

**Andrei D. Polyanin
Alexander V. Manzhirov**



**HANDBOOK OF
INTEGRAL
EQUATIONS**

SECOND EDITION

 **Chapman & Hall/CRC**
Taylor & Francis Group

**HANDBOOK OF
INTEGRAL
EQUATIONS**

SECOND EDITION

Handbooks of Mathematical Equations

Handbook of Linear Partial Differential Equations for Engineers and Scientists

A. D. Polyanin, 2002

Handbook of First Order Partial Differential Equations

A. D. Polyanin, V. F. Zaitsev, and A. Moussiaux, 2002

Handbook of Exact Solutions for Ordinary Differential Equations, 2nd Edition

A. D. Polyanin and V. F. Zaitsev, 2003

Handbook of Nonlinear Partial Differential Equations

A. D. Polyanin and V. F. Zaitsev, 2004

Handbook of Integral Equations, 2nd Edition

A. D. Polyanin and A. V. Manzhirov, 2008

See also:

Handbook of Mathematics for Engineers and Scientists

A. D. Polyanin and A. V. Manzhirov, 2007

HANDBOOK OF INTEGRAL EQUATIONS

SECOND EDITION

Andrei D. Polyanin
Alexander V. Manzhirov

 **Chapman & Hall/CRC**
Taylor & Francis Group

Boca Raton London New York

Chapman & Hall/CRC is an imprint of the
Taylor & Francis Group, an informa business

Chapman & Hall/CRC
Taylor & Francis Group
6000 Broken Sound Parkway NW, Suite 300
Boca Raton, FL 33487-2742

© 2008 by Taylor & Francis Group, LLC
Chapman & Hall/CRC is an imprint of Taylor & Francis Group, an Informa business

No claim to original U.S. Government works
Printed in the United States of America on acid-free paper
10 9 8 7 6 5 4 3 2 1

International Standard Book Number-13: 978-1-58488-507-8 (Hardcover)

This book contains information obtained from authentic and highly regarded sources. Reprinted material is quoted with permission, and sources are indicated. A wide variety of references are listed. Reasonable efforts have been made to publish reliable data and information, but the author and the publisher cannot assume responsibility for the validity of all materials or for the consequences of their use.

Except as permitted under U.S. Copyright Law, no part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access www.copyright.com (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC) 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

Trademark Notice: Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

Library of Congress Cataloging-in-Publication Data

Polianin, A. D. (Andrei Dmitrievich)
Handbook of integral equations / Andrei D. Polyanin and Alexander V. Manzhirov. -- 2nd ed.
p. cm.
Includes bibliographical references and index.
ISBN-13: 978-1-58488-507-8 (hardcover : alk. paper)
ISBN-10: 1-58488-507-6 (hardcover : alk. paper)
1. Integral equations--Handbooks, manuals, etc. I. Manzhirov, A. V. (Aleksandr Vladimirovich) II.
Title.

QA431.P65 2008
515'.45--dc22

2007035725

Visit the Taylor & Francis Web site at
<http://www.taylorandfrancis.com>

and the CRC Press Web site at
<http://www.crcpress.com>

CONTENTS

Authors	xxix
Preface	xxxii
Some Remarks and Notation	xxxiii

Part I. Exact Solutions of Integral Equations

1. Linear Equations of the First Kind with Variable Limit of Integration	3
1.1. Equations Whose Kernels Contain Power-Law Functions	4
1.1-1. Kernels Linear in the Arguments x and t	4
1.1-2. Kernels Quadratic in the Arguments x and t	4
1.1-3. Kernels Cubic in the Arguments x and t	5
1.1-4. Kernels Containing Higher-Order Polynomials in x and t	6
1.1-5. Kernels Containing Rational Functions	7
1.1-6. Kernels Containing Square Roots	9
1.1-7. Kernels Containing Arbitrary Powers	12
1.1-8. Two-Dimensional Equation of the Abel Type	15
1.2. Equations Whose Kernels Contain Exponential Functions	15
1.2-1. Kernels Containing Exponential Functions	15
1.2-2. Kernels Containing Power-Law and Exponential Functions	19
1.3. Equations Whose Kernels Contain Hyperbolic Functions	22
1.3-1. Kernels Containing Hyperbolic Cosine	22
1.3-2. Kernels Containing Hyperbolic Sine	28
1.3-3. Kernels Containing Hyperbolic Tangent	36
1.3-4. Kernels Containing Hyperbolic Cotangent	38
1.3-5. Kernels Containing Combinations of Hyperbolic Functions	39
1.4. Equations Whose Kernels Contain Logarithmic Functions	42
1.4-1. Kernels Containing Logarithmic Functions	42
1.4-2. Kernels Containing Power-Law and Logarithmic Functions	45
1.5. Equations Whose Kernels Contain Trigonometric Functions	46
1.5-1. Kernels Containing Cosine	46
1.5-2. Kernels Containing Sine	52
1.5-3. Kernels Containing Tangent	60
1.5-4. Kernels Containing Cotangent	62
1.5-5. Kernels Containing Combinations of Trigonometric Functions	63
1.6. Equations Whose Kernels Contain Inverse Trigonometric Functions	66
1.6-1. Kernels Containing Arccosine	66
1.6-2. Kernels Containing Arcsine	68
1.6-3. Kernels Containing Arctangent	70
1.6-4. Kernels Containing Arccotangent	71

1.7. Equations Whose Kernels Contain Combinations of Elementary Functions	73
1.7-1. Kernels Containing Exponential and Hyperbolic Functions	73
1.7-2. Kernels Containing Exponential and Logarithmic Functions	77
1.7-3. Kernels Containing Exponential and Trigonometric Functions	78
1.7-4. Kernels Containing Hyperbolic and Logarithmic Functions	83
1.7-5. Kernels Containing Hyperbolic and Trigonometric Functions	84
1.7-6. Kernels Containing Logarithmic and Trigonometric Functions	85
1.8. Equations Whose Kernels Contain Special Functions	86
1.8-1. Kernels Containing Error Function or Exponential Integral	86
1.8-2. Kernels Containing Sine and Cosine Integrals	87
1.8-3. Kernels Containing Fresnel Integrals	87
1.8-4. Kernels Containing Incomplete Gamma Functions	88
1.8-5. Kernels Containing Bessel Functions	88
1.8-6. Kernels Containing Modified Bessel Functions	97
1.8-7. Kernels Containing Legendre Polynomials	105
1.8-8. Kernels Containing Associated Legendre Functions	107
1.8-9. Kernels Containing Confluent Hypergeometric Functions	107
1.8-10. Kernels Containing Hermite Polynomials	108
1.8-11. Kernels Containing Chebyshev Polynomials	109
1.8-12. Kernels Containing Laguerre Polynomials	110
1.8-13. Kernels Containing Jacobi Theta Functions	110
1.8-14. Kernels Containing Other Special Functions	111
1.9. Equations Whose Kernels Contain Arbitrary Functions	111
1.9-1. Equations with Degenerate Kernel: $K(x, t) = g_1(x)h_1(t) + g_2(x)h_2(t)$	111
1.9-2. Equations with Difference Kernel: $K(x, t) = K(x - t)$	114
1.9-3. Other Equations	122
1.10. Some Formulas and Transformations	124
2. Linear Equations of the Second Kind with Variable Limit of Integration	127
2.1. Equations Whose Kernels Contain Power-Law Functions	127
2.1-1. Kernels Linear in the Arguments x and t	127
2.1-2. Kernels Quadratic in the Arguments x and t	129
2.1-3. Kernels Cubic in the Arguments x and t	132
2.1-4. Kernels Containing Higher-Order Polynomials in x and t	133
2.1-5. Kernels Containing Rational Functions	136
2.1-6. Kernels Containing Square Roots and Fractional Powers	138
2.1-7. Kernels Containing Arbitrary Powers	139
2.2. Equations Whose Kernels Contain Exponential Functions	144
2.2-1. Kernels Containing Exponential Functions	144
2.2-2. Kernels Containing Power-Law and Exponential Functions	151
2.3. Equations Whose Kernels Contain Hyperbolic Functions	154
2.3-1. Kernels Containing Hyperbolic Cosine	154
2.3-2. Kernels Containing Hyperbolic Sine	156
2.3-3. Kernels Containing Hyperbolic Tangent	161
2.3-4. Kernels Containing Hyperbolic Cotangent	162
2.3-5. Kernels Containing Combinations of Hyperbolic Functions	164
2.4. Equations Whose Kernels Contain Logarithmic Functions	164
2.4-1. Kernels Containing Logarithmic Functions	164
2.4-2. Kernels Containing Power-Law and Logarithmic Functions	165

2.5. Equations Whose Kernels Contain Trigonometric Functions	166
2.5-1. Kernels Containing Cosine	166
2.5-2. Kernels Containing Sine	169
2.5-3. Kernels Containing Tangent	174
2.5-4. Kernels Containing Cotangent	175
2.5-5. Kernels Containing Combinations of Trigonometric Functions	176
2.6. Equations Whose Kernels Contain Inverse Trigonometric Functions	176
2.6-1. Kernels Containing Arccosine	176
2.6-2. Kernels Containing Arcsine	177
2.6-3. Kernels Containing Arctangent	178
2.6-4. Kernels Containing Arccotangent	178
2.7. Equations Whose Kernels Contain Combinations of Elementary Functions	179
2.7-1. Kernels Containing Exponential and Hyperbolic Functions	179
2.7-2. Kernels Containing Exponential and Logarithmic Functions	180
2.7-3. Kernels Containing Exponential and Trigonometric Functions	181
2.7-4. Kernels Containing Hyperbolic and Logarithmic Functions	185
2.7-5. Kernels Containing Hyperbolic and Trigonometric Functions	186
2.7-6. Kernels Containing Logarithmic and Trigonometric Functions	187
2.8. Equations Whose Kernels Contain Special Functions	187
2.8-1. Kernels Containing Bessel Functions	187
2.8-2. Kernels Containing Modified Bessel Functions	189
2.9. Equations Whose Kernels Contain Arbitrary Functions	191
2.9-1. Equations with Degenerate Kernel: $K(x, t) = g_1(x)h_1(t) + \cdots + g_n(x)h_n(t)$	191
2.9-2. Equations with Difference Kernel: $K(x, t) = K(x - t)$	203
2.9-3. Other Equations	212
2.10. Some Formulas and Transformations	215
3. Linear Equations of the First Kind with Constant Limits of Integration	217
3.1. Equations Whose Kernels Contain Power-Law Functions	217
3.1-1. Kernels Linear in the Arguments x and t	217
3.1-2. Kernels Quadratic in the Arguments x and t	219
3.1-3. Kernels Containing Integer Powers of x and t or Rational Functions	220
3.1-4. Kernels Containing Square Roots	222
3.1-5. Kernels Containing Arbitrary Powers	223
3.1-6. Equations Containing the Unknown Function of a Complicated Argument	227
3.1-7. Singular Equations	228
3.2. Equations Whose Kernels Contain Exponential Functions	231
3.2-1. Kernels Containing Exponential Functions of the Form $e^{\lambda x-t }$	231
3.2-2. Kernels Containing Exponential Functions of the Forms $e^{\lambda x}$ and $e^{\mu t}$	234
3.2-3. Kernels Containing Exponential Functions of the Form $e^{\lambda xt}$	234
3.2-4. Kernels Containing Power-Law and Exponential Functions	236
3.2-5. Kernels Containing Exponential Functions of the Form $e^{\lambda(x \pm t)^2}$	236
3.2-6. Other Kernels	237
3.3. Equations Whose Kernels Contain Hyperbolic Functions	238
3.3-1. Kernels Containing Hyperbolic Cosine	238
3.3-2. Kernels Containing Hyperbolic Sine	238
3.3-3. Kernels Containing Hyperbolic Tangent	241
3.3-4. Kernels Containing Hyperbolic Cotangent	242

3.4. Equations Whose Kernels Contain Logarithmic Functions	242
3.4-1. Kernels Containing Logarithmic Functions	242
3.4-2. Kernels Containing Power-Law and Logarithmic Functions	244
3.4-3. Equation Containing the Unknown Function of a Complicated Argument	246
3.5. Equations Whose Kernels Contain Trigonometric Functions	246
3.5-1. Kernels Containing Cosine	246
3.5-2. Kernels Containing Sine	247
3.5-3. Kernels Containing Tangent	251
3.5-4. Kernels Containing Cotangent	252
3.5-5. Kernels Containing a Combination of Trigonometric Functions	252
3.5-6. Equations Containing the Unknown Function of a Complicated Argument	254
3.5-7. Singular Equations	255
3.6. Equations Whose Kernels Contain Combinations of Elementary Functions	255
3.6-1. Kernels Containing Hyperbolic and Logarithmic Functions	255
3.6-2. Kernels Containing Logarithmic and Trigonometric Functions	256
3.6-3. Kernels Containing Combinations of Exponential and Other Elementary Functions	257
3.7. Equations Whose Kernels Contain Special Functions	258
3.7-1. Kernels Containing Error Function, Exponential Integral or Logarithmic Integral	258
3.7-2. Kernels Containing Sine Integrals, Cosine Integrals, or Fresnel Integrals	258
3.7-3. Kernels Containing Gamma Functions	260
3.7-4. Kernels Containing Incomplete Gamma Functions	260
3.7-5. Kernels Containing Bessel Functions of the First Kind	261
3.7-6. Kernels Containing Bessel Functions of the Second Kind	264
3.7-7. Kernels Containing Combinations of the Bessel Functions	265
3.7-8. Kernels Containing Modified Bessel Functions of the First Kind	266
3.7-9. Kernels Containing Modified Bessel Functions of the Second Kind	266
3.7-10. Kernels Containing a Combination of Bessel and Modified Bessel Functions	269
3.7-11. Kernels Containing Legendre Functions	270
3.7-12. Kernels Containing Associated Legendre Functions	271
3.7-13. Kernels Containing Kummer Confluent Hypergeometric Functions	272
3.7-14. Kernels Containing Tricomi Confluent Hypergeometric Functions	274
3.7-15. Kernels Containing Whittaker Confluent Hypergeometric Functions	274
3.7-16. Kernels Containing Gauss Hypergeometric Functions	276
3.7-17. Kernels Containing Parabolic Cylinder Functions	276
3.7-18. Kernels Containing Other Special Functions	277
3.8. Equations Whose Kernels Contain Arbitrary Functions	278
3.8-1. Equations with Degenerate Kernel	278
3.8-2. Equations Containing Modulus	279
3.8-3. Equations with Difference Kernel: $K(x, t) = K(x - t)$	284
3.8-4. Other Equations of the Form $\int_a^b K(x, t)y(t) dt = F(x)$	285
3.8-5. Equations of the Form $\int_a^b K(x, t)y(\cdot) dt = F(x)$	289
3.9. Dual Integral Equations of the First Kind	295
3.9-1. Kernels Containing Trigonometric Functions	295
3.9-2. Kernels Containing Bessel Functions of the First Kind	297
3.9-3. Kernels Containing Bessel Functions of the Second Kind	299
3.9-4. Kernels Containing Legendre Spherical Functions of the First Kind, $i^2 = -1$	299

4. Linear Equations of the Second Kind with Constant Limits of Integration	301
4.1. Equations Whose Kernels Contain Power-Law Functions	301
4.1-1. Kernels Linear in the Arguments x and t	301
4.1-2. Kernels Quadratic in the Arguments x and t	304
4.1-3. Kernels Cubic in the Arguments x and t	307
4.1-4. Kernels Containing Higher-Order Polynomials in x and t	311
4.1-5. Kernels Containing Rational Functions	314
4.1-6. Kernels Containing Arbitrary Powers	317
4.1-7. Singular Equations	319
4.2. Equations Whose Kernels Contain Exponential Functions	320
4.2-1. Kernels Containing Exponential Functions	320
4.2-2. Kernels Containing Power-Law and Exponential Functions	326
4.3. Equations Whose Kernels Contain Hyperbolic Functions	327
4.3-1. Kernels Containing Hyperbolic Cosine	327
4.3-2. Kernels Containing Hyperbolic Sine	329
4.3-3. Kernels Containing Hyperbolic Tangent	332
4.3-4. Kernels Containing Hyperbolic Cotangent	333
4.3-5. Kernels Containing Combination of Hyperbolic Functions	334
4.4. Equations Whose Kernels Contain Logarithmic Functions	334
4.4-1. Kernels Containing Logarithmic Functions	334
4.4-2. Kernels Containing Power-Law and Logarithmic Functions	335
4.5. Equations Whose Kernels Contain Trigonometric Functions	335
4.5-1. Kernels Containing Cosine	335
4.5-2. Kernels Containing Sine	337
4.5-3. Kernels Containing Tangent	342
4.5-4. Kernels Containing Cotangent	343
4.5-5. Kernels Containing Combinations of Trigonometric Functions	344
4.5-6. Singular Equation	344
4.6. Equations Whose Kernels Contain Inverse Trigonometric Functions	344
4.6-1. Kernels Containing Arccosine	344
4.6-2. Kernels Containing Arcsine	345
4.6-3. Kernels Containing Arctangent	346
4.6-4. Kernels Containing Arccotangent	347
4.7. Equations Whose Kernels Contain Combinations of Elementary Functions	348
4.7-1. Kernels Containing Exponential and Hyperbolic Functions	348
4.7-2. Kernels Containing Exponential and Logarithmic Functions	349
4.7-3. Kernels Containing Exponential and Trigonometric Functions	349
4.7-4. Kernels Containing Hyperbolic and Logarithmic Functions	351
4.7-5. Kernels Containing Hyperbolic and Trigonometric Functions	352
4.7-6. Kernels Containing Logarithmic and Trigonometric Functions	353
4.8. Equations Whose Kernels Contain Special Functions	353
4.8-1. Kernels Containing Bessel Functions	353
4.8-2. Kernels Containing Modified Bessel Functions	355
4.9. Equations Whose Kernels Contain Arbitrary Functions	357
4.9-1. Equations with Degenerate Kernel: $K(x, t) = g_1(x)h_1(t) + \dots + g_n(x)h_n(t)$	357
4.9-2. Equations with Difference Kernel: $K(x, t) = K(x - t)$	372
4.9-3. Other Equations of the Form $y(x) + \int_a^b K(x, t)y(t) dt = F(x)$	374
4.9-4. Equations of the Form $y(x) + \int_a^b K(x, t)y(\dots) dt = F(x)$	381
4.10. Some Formulas and Transformations	390

5. Nonlinear Equations of the First Kind with Variable Limit of Integration	393
5.1. Equations with Quadratic Nonlinearity That Contain Arbitrary Parameters	393
5.1-1. Equations of the Form $\int_0^x y(t)y(x-t) dt = f(x)$	393
5.1-2. Equations of the Form $\int_0^x K(x,t)y(t)y(x-t) dt = f(x)$	395
5.1-3. Equations of the Form $\int_0^x y(t)y(\dots) dt = f(x)$	396
5.2. Equations with Quadratic Nonlinearity That Contain Arbitrary Functions	397
5.2-1. Equations of the Form $\int_a^x K(x,t)[Ay(t) + By^2(t)] dt = f(x)$	397
5.2-2. Equations of the Form $\int_a^x K(x,t)y(t)y(ax+bt) dt = f(x)$	398
5.3. Equations with Nonlinearity of General Form	399
5.3-1. Equations of the Form $\int_a^x K(x,t)f(t,y(t)) dt = g(x)$	399
5.3-2. Other Equations	401
6. Nonlinear Equations of the Second Kind with Variable Limit of Integration	403
6.1. Equations with Quadratic Nonlinearity That Contain Arbitrary Parameters	403
6.1-1. Equations of the Form $y(x) + \int_a^x K(x,t)y^2(t) dt = F(x)$	403
6.1-2. Equations of the Form $y(x) + \int_a^x K(x,t)y(t)y(x-t) dt = F(x)$	406
6.2. Equations with Quadratic Nonlinearity That Contain Arbitrary Functions	406
6.2-1. Equations of the Form $y(x) + \int_a^x K(x,t)y^2(t) dt = F(x)$	406
6.2-2. Other Equations	407
6.3. Equations with Power-Law Nonlinearity	408
6.3-1. Equations Containing Arbitrary Parameters	408
6.3-2. Equations Containing Arbitrary Functions	410
6.4. Equations with Exponential Nonlinearity	411
6.4-1. Equations Containing Arbitrary Parameters	411
6.4-2. Equations Containing Arbitrary Functions	413
6.5. Equations with Hyperbolic Nonlinearity	414
6.5-1. Integrands with Nonlinearity of the Form $\cosh[\beta y(t)]$	414
6.5-2. Integrands with Nonlinearity of the Form $\sinh[\beta y(t)]$	415
6.5-3. Integrands with Nonlinearity of the Form $\tanh[\beta y(t)]$	416
6.5-4. Integrands with Nonlinearity of the Form $\coth[\beta y(t)]$	418
6.6. Equations with Logarithmic Nonlinearity	419
6.6-1. Integrands Containing Power-Law Functions of x and t	419
6.6-2. Integrands Containing Exponential Functions of x and t	419
6.6-3. Other Integrands	420
6.7. Equations with Trigonometric Nonlinearity	420
6.7-1. Integrands with Nonlinearity of the Form $\cos[\beta y(t)]$	420
6.7-2. Integrands with Nonlinearity of the Form $\sin[\beta y(t)]$	422
6.7-3. Integrands with Nonlinearity of the Form $\tan[\beta y(t)]$	423
6.7-4. Integrands with Nonlinearity of the Form $\cot[\beta y(t)]$	424
6.8. Equations with Nonlinearity of General Form	425
6.8-1. Equations of the Form $y(x) + \int_a^x K(x,t)G(y(t)) dt = F(x)$	425
6.8-2. Equations of the Form $y(x) + \int_a^x K(x-t)G(t,y(t)) dt = F(x)$	428
6.8-3. Other Equations	431
7. Nonlinear Equations of the First Kind with Constant Limits of Integration	433
7.1. Equations with Quadratic Nonlinearity That Contain Arbitrary Parameters	433
7.1-1. Equations of the Form $\int_a^b K(t)y(x)y(t) dt = F(x)$	433
7.1-2. Equations of the Form $\int_a^b K(t)y(t)y(xt) dt = F(x)$	435
7.1-3. Other Equations	436

7.2. Equations with Quadratic Nonlinearity That Contain Arbitrary Functions	437
7.2-1. Equations of the Form $\int_a^b K(t)y(t)y(\cdot \cdot) dt = F(x)$	437
7.2-2. Equations of the Form $\int_a^b [K(x, t)y(t) + M(x, t)y^2(t)] dt = F(x)$	443
7.3. Equations with Power-Law Nonlinearity That Contain Arbitrary Functions	444
7.3-1. Equations of the Form $\int_a^b K(t)y^\mu(x)y^\gamma(t) dt = F(x)$	444
7.3-2. Equations of the Form $\int_a^b K(t)y^\gamma(t)y(xt) dt = F(x)$	444
7.3-3. Equations of the Form $\int_a^b K(t)y^\gamma(t)y(x + \beta t) dt = F(x)$	445
7.3-4. Equations of the Form $\int_a^b [K(x, t)y(t) + M(x, t)y^\gamma(t)] dt = f(x)$	446
7.3-5. Other Equations	446
7.4. Equations with Nonlinearity of General Form	447
7.4-1. Equations of the Form $\int_a^b \varphi(y(x))K(t, y(t)) dt = F(x)$	447
7.4-2. Equations of the Form $\int_a^b y(xt)K(t, y(t)) dt = F(x)$	447
7.4-3. Equations of the Form $\int_a^b y(x + \beta t)K(t, y(t)) dt = F(x)$	449
7.4-4. Equations of the Form $\int_a^b [K(x, t)y(t) + \varphi(x)\Psi(t, y(t))] dt = F(x)$	450
7.4-5. Other Equations	451
8. Nonlinear Equations of the Second Kind with Constant Limits of Integration	453
8.1. Equations with Quadratic Nonlinearity That Contain Arbitrary Parameters	453
8.1-1. Equations of the Form $y(x) + \int_a^b K(x, t)y^2(t) dt = F(x)$	453
8.1-2. Equations of the Form $y(x) + \int_a^b K(x, t)y(x)y(t) dt = F(x)$	454
8.1-3. Equations of the Form $y(x) + \int_a^b K(t)y(t)y(\cdot \cdot) dt = F(x)$	455
8.2. Equations with Quadratic Nonlinearity That Contain Arbitrary Functions	456
8.2-1. Equations of the Form $y(x) + \int_a^b K(x, t)y^2(t) dt = F(x)$	456
8.2-2. Equations of the Form $y(x) + \int_a^b \sum K_{nm}(x, t)y^n(x)y^m(t) dt = F(x)$, $n + m \leq 2$	457
8.2-3. Equations of the Form $y(x) + \int_a^b K(t)y(t)y(\cdot \cdot) dt = F(x)$	460
8.3. Equations with Power-Law Nonlinearity	464
8.3-1. Equations of the Form $y(x) + \int_a^b K(x, t)y^\beta(t) dt = F(x)$	464
8.3-2. Other Equations	465
8.4. Equations with Exponential Nonlinearity	467
8.4-1. Integrands with Nonlinearity of the Form $\exp[\beta y(t)]$	467
8.4-2. Other Integrands	468
8.5. Equations with Hyperbolic Nonlinearity	468
8.5-1. Integrands with Nonlinearity of the Form $\cosh[\beta y(t)]$	468
8.5-2. Integrands with Nonlinearity of the Form $\sinh[\beta y(t)]$	469
8.5-3. Integrands with Nonlinearity of the Form $\tanh[\beta y(t)]$	469
8.5-4. Integrands with Nonlinearity of the Form $\coth[\beta y(t)]$	470
8.5-5. Other Integrands	471
8.6. Equations with Logarithmic Nonlinearity	472
8.6-1. Integrands with Nonlinearity of the Form $\ln[\beta y(t)]$	472
8.6-2. Other Integrands	473
8.7. Equations with Trigonometric Nonlinearity	473
8.7-1. Integrands with Nonlinearity of the Form $\cos[\beta y(t)]$	473
8.7-2. Integrands with Nonlinearity of the Form $\sin[\beta y(t)]$	474
8.7-3. Integrands with Nonlinearity of the Form $\tan[\beta y(t)]$	475
8.7-4. Integrands with Nonlinearity of the Form $\cot[\beta y(t)]$	475
8.7-5. Other Integrands	476

8.8. Equations with Nonlinearity of General Form	477
8.8-1. Equations of the Form $y(x) + \int_a^b K(x-t)G(y(t)) dt = F(x)$	477
8.8-2. Equations of the Form $y(x) + \int_a^b K(x,t)G(t, y(t)) dt = F(x)$	479
8.8-3. Equations of the Form $y(x) + \int_a^b G(x, t, y(t)) dt = F(x)$	483
8.8-4. Equations of the Form $y(x) + \int_a^b y(xt)G(t, y(t)) dt = F(x)$	485
8.8-5. Equations of the Form $y(x) + \int_a^b y(x + \beta t)G(t, y(t)) dt = F(x)$	487
8.8-6. Other Equations	494

Part II. Methods for Solving Integral Equations

9. Main Definitions and Formulas. Integral Transforms	501
9.1. Some Definitions, Remarks, and Formulas	501
9.1-1. Some Definitions	501
9.1-2. Structure of Solutions to Linear Integral Equations	502
9.1-3. Integral Transforms	503
9.1-4. Residues. Calculation Formulas. Cauchy's Residue Theorem	504
9.1-5. Jordan Lemma	505
9.2. Laplace Transform	505
9.2-1. Definition. Inversion Formula	505
9.2-2. Inverse Transforms of Rational Functions	506
9.2-3. Inversion of Functions with Finitely Many Singular Points	507
9.2-4. Convolution Theorem. Main Properties of the Laplace Transform	507
9.2-5. Limit Theorems	507
9.2-6. Representation of Inverse Transforms as Convergent Series	509
9.2-7. Representation of Inverse Transforms as Asymptotic Expansions as $x \rightarrow \infty$	509
9.2-8. Post-Widder Formula	510
9.3. Mellin Transform	510
9.3-1. Definition. Inversion Formula	510
9.3-2. Main Properties of the Mellin Transform	511
9.3-3. Relation Among the Mellin, Laplace, and Fourier Transforms	511
9.4. Fourier Transform	512
9.4-1. Definition. Inversion Formula	512
9.4-2. Asymmetric Form of the Transform	512
9.4-3. Alternative Fourier Transform	512
9.4-4. Convolution Theorem. Main Properties of the Fourier Transforms	513
9.5. Fourier Cosine and Sine Transforms	514
9.5-1. Fourier Cosine Transform	514
9.5-2. Fourier Sine Transform	514
9.6. Other Integral Transforms	515
9.6-1. Hankel Transform	515
9.6-2. Meijer Transform	516
9.6-3. Kontorovich-Lebedev Transform	516
9.6-4. Y -transform	516
9.6-5. Summary Table of Integral Transforms	517
10. Methods for Solving Linear Equations of the Form $\int_a^x K(x,t)y(t) dt = f(x)$	519
10.1. Volterra Equations of the First Kind	519
10.1-1. Equations of the First Kind. Function and Kernel Classes	519
10.1-2. Existence and Uniqueness of a Solution	520
10.1-3. Some Problems Leading to Volterra Integral Equations of the First Kind	520

10.2. Equations with Degenerate Kernel: $K(x, t) = g_1(x)h_1(t) + \cdots + g_n(x)h_n(t)$	522
10.2-1. Equations with Kernel of the Form $K(x, t) = g_1(x)h_1(t) + g_2(x)h_2(t)$	522
10.2-2. Equations with General Degenerate Kernel	523
10.3. Reduction of Volterra Equations of the First Kind to Volterra Equations of the Second Kind	524
10.3-1. First Method	524
10.3-2. Second Method	524
10.4. Equations with Difference Kernel: $K(x, t) = K(x - t)$	524
10.4-1. Solution Method Based on the Laplace Transform	524
10.4-2. Case in Which the Transform of the Solution is a Rational Function	525
10.4-3. Convolution Representation of a Solution	526
10.4-4. Application of an Auxiliary Equation	527
10.4-5. Reduction to Ordinary Differential Equations	527
10.4-6. Reduction of a Volterra Equation to a Wiener–Hopf Equation	528
10.5. Method of Fractional Differentiation	529
10.5-1. Definition of Fractional Integrals	529
10.5-2. Definition of Fractional Derivatives	529
10.5-3. Main Properties	530
10.5-4. Solution of the Generalized Abel Equation	531
10.5-5. Erdélyi–Kober Operators	532
10.6. Equations with Weakly Singular Kernel	532
10.6-1. Method of Transformation of the Kernel	532
10.6-2. Kernel with Logarithmic Singularity	533
10.7. Method of Quadratures	534
10.7-1. Quadrature Formulas	534
10.7-2. General Scheme of the Method	535
10.7-3. Algorithm Based on the Trapezoidal Rule	536
10.7-4. Algorithm for an Equation with Degenerate Kernel	536
10.8. Equations with Infinite Integration Limit	537
10.8-1. Equation of the First Kind with Variable Lower Limit of Integration	537
10.8-2. Reduction to a Wiener–Hopf Equation of the First Kind	538
11. Methods for Solving Linear Equations of the Form $y(x) - \int_a^x K(x, t)y(t) dt = f(x)$ 539	
11.1. Volterra Integral Equations of the Second Kind	539
11.1-1. Preliminary Remarks. Equations for the Resolvent	539
11.1-2. Relationship Between Solutions of Some Integral Equations	540
11.2. Equations with Degenerate Kernel: $K(x, t) = g_1(x)h_1(t) + \cdots + g_n(x)h_n(t)$	540
11.2-1. Equations with Kernel of the Form $K(x, t) = \varphi(x) + \psi(x)(x - t)$	540
11.2-2. Equations with Kernel of the Form $K(x, t) = \varphi(t) + \psi(t)(t - x)$	541
11.2-3. Equations with Kernel of the Form $K(x, t) = \sum_{m=1}^n \varphi_m(x)(x - t)^{m-1}$	542
11.2-4. Equations with Kernel of the Form $K(x, t) = \sum_{m=1}^n \varphi_m(t)(t - x)^{m-1}$	543
11.2-5. Equations with Degenerate Kernel of the General Form	543
11.3. Equations with Difference Kernel: $K(x, t) = K(x - t)$	544
11.3-1. Solution Method Based on the Laplace Transform	544
11.3-2. Method Based on the Solution of an Auxiliary Equation	546
11.3-3. Reduction to Ordinary Differential Equations	547
11.3-4. Reduction to a Wiener–Hopf Equation of the Second Kind	547
11.3-5. Method of Fractional Integration for the Generalized Abel Equation	548
11.3-6. Systems of Volterra Integral Equations	549

11.4. Operator Methods for Solving Linear Integral Equations	549
11.4-1. Application of a Solution of a “Truncated” Equation of the First Kind	549
11.4-2. Application of the Auxiliary Equation of the Second Kind	551
11.4-3. Method for Solving “Quadratic” Operator Equations	552
11.4-4. Solution of Operator Equations of Polynomial Form	553
11.4-5. Some Generalizations	554
11.5. Construction of Solutions of Integral Equations with Special Right-Hand Side	555
11.5-1. General Scheme	555
11.5-2. Generating Function of Exponential Form	555
11.5-3. Power-Law Generating Function	557
11.5-4. Generating Function Containing Sines and Cosines	558
11.6. Method of Model Solutions	559
11.6-1. Preliminary Remarks	559
11.6-2. Description of the Method	560
11.6-3. Model Solution in the Case of an Exponential Right-Hand Side	561
11.6-4. Model Solution in the Case of a Power-Law Right-Hand Side	562
11.6-5. Model Solution in the Case of a Sine-Shaped Right-Hand Side	562
11.6-6. Model Solution in the Case of a Cosine-Shaped Right-Hand Side	563
11.6-7. Some Generalizations	563
11.7. Method of Differentiation for Integral Equations	564
11.7-1. Equations with Kernel Containing a Sum of Exponential Functions	564
11.7-2. Equations with Kernel Containing a Sum of Hyperbolic Functions	564
11.7-3. Equations with Kernel Containing a Sum of Trigonometric Functions	564
11.7-4. Equations Whose Kernels Contain Combinations of Various Functions	565
11.8. Reduction of Volterra Equations of the Second Kind to Volterra Equations of the First Kind	565
11.8-1. First Method	565
11.8-2. Second Method	566
11.9. Successive Approximation Method	566
11.9-1. General Scheme	566
11.9-2. Formula for the Resolvent	567
11.10. Method of Quadratures	568
11.10-1. General Scheme of the Method	568
11.10-2. Application of the Trapezoidal Rule	568
11.10-3. Case of a Degenerate Kernel	569
11.11. Equations with Infinite Integration Limit	569
11.11-1. Equation of the Second Kind with Variable Lower Integration Limit	570
11.11-2. Reduction to a Wiener–Hopf Equation of the Second Kind	571
12. Methods for Solving Linear Equations of the Form $\int_a^b K(x, t)y(t) dt = f(x)$	573
12.1. Some Definition and Remarks	573
12.1-1. Fredholm Integral Equations of the First Kind	573
12.1-2. Integral Equations of the First Kind with Weak Singularity	574
12.1-3. Integral Equations of Convolution Type	574
12.1-4. Dual Integral Equations of the First Kind	575
12.1-5. Some Problems Leading to Integral Equations of the First Kind	575
12.2. Integral Equations of the First Kind with Symmetric Kernel	577
12.2-1. Solution of an Integral Equation in Terms of Series in Eigenfunctions of Its Kernel	577
12.2-2. Method of Successive Approximations	579

12.3. Integral Equations of the First Kind with Nonsymmetric Kernel	580
12.3-1. Representation of a Solution in the Form of Series. General Description	580
12.3-2. Special Case of a Kernel That is a Generating Function	580
12.3-3. Special Case of the Right-Hand Side Represented in Terms of Orthogonal Functions	582
12.3-4. General Case. Galerkin's Method	582
12.3-5. Utilization of the Schmidt Kernels for the Construction of Solutions of Equations	582
12.4. Method of Differentiation for Integral Equations	583
12.4-1. Equations with Modulus	583
12.4-2. Other Equations. Some Generalizations	585
12.5. Method of Integral Transforms	586
12.5-1. Equation with Difference Kernel on the Entire Axis	586
12.5-2. Equations with Kernel $K(x, t) = K(x/t)$ on the Semiaxis	587
12.5-3. Equation with Kernel $K(x, t) = K(xt)$ and Some Generalizations	587
12.6. Krein's Method and Some Other Exact Methods for Integral Equations of Special Types	588
12.6-1. Krein's Method for an Equation with Difference Kernel with a Weak Singularity	588
12.6-2. Kernel is the Sum of a Nondegenerate Kernel and an Arbitrary Degenerate Kernel	589
12.6-3. Reduction of Integral Equations of the First Kind to Equations of the Second Kind	591
12.7. Riemann Problem for the Real Axis	592
12.7-1. Relationships Between the Fourier Integral and the Cauchy Type Integral	592
12.7-2. One-Sided Fourier Integrals	593
12.7-3. Analytic Continuation Theorem and the Generalized Liouville Theorem	595
12.7-4. Riemann Boundary Value Problem	595
12.7-5. Problems with Rational Coefficients	601
12.7-6. Exceptional Cases. The Homogeneous Problem	602
12.7-7. Exceptional Cases. The Nonhomogeneous Problem	604
12.8. Carleman Method for Equations of the Convolution Type of the First Kind	606
12.8-1. Wiener-Hopf Equation of the First Kind	606
12.8-2. Integral Equations of the First Kind with Two Kernels	607
12.9. Dual Integral Equations of the First Kind	610
12.9-1. Carleman Method for Equations with Difference Kernels	610
12.9-2. General Scheme of Finding Solutions of Dual Integral Equations	611
12.9-3. Exact Solutions of Some Dual Equations of the First Kind	613
12.9-4. Reduction of Dual Equations to a Fredholm Equation	615
12.10. Asymptotic Methods for Solving Equations with Logarithmic Singularity	618
12.10-1. Preliminary Remarks	618
12.10-2. Solution for Large λ	619
12.10-3. Solution for Small λ	620
12.10-4. Integral Equation of Elasticity	621
12.11. Regularization Methods	621
12.11-1. Lavrentiev Regularization Method	621
12.11-2. Tikhonov Regularization Method	622
12.12. Fredholm Integral Equation of the First Kind as an Ill-Posed Problem	623
12.12-1. General Notions of Well-Posed and Ill-Posed Problems	623
12.12-2. Integral Equation of the First Kind is an Ill-Posed Problem	624

13. Methods for Solving Linear Equations of the Form $y(x) - \int_a^b K(x, t)y(t) dt = f(x)$	625
13.1. Some Definition and Remarks	625
13.1-1. Fredholm Equations and Equations with Weak Singularity of the Second Kind	625
13.1-2. Structure of the Solution	626
13.1-3. Integral Equations of Convolution Type of the Second Kind	626
13.1-4. Dual Integral Equations of the Second Kind	627
13.2. Fredholm Equations of the Second Kind with Degenerate Kernel. Some Generalizations	627
13.2-1. Simplest Degenerate Kernel	627
13.2-2. Degenerate Kernel in the General Case	628
13.2-3. Kernel is the Sum of a Nondegenerate Kernel and an Arbitrary Degenerate Kernel	631
13.3. Solution as a Power Series in the Parameter. Method of Successive Approximations	632
13.3-1. Iterated Kernels	632
13.3-2. Method of Successive Approximations	633
13.3-3. Construction of the Resolvent	633
13.3-4. Orthogonal Kernels	634
13.4. Method of Fredholm Determinants	635
13.4-1. Formula for the Resolvent	635
13.4-2. Recurrent Relations	636
13.5. Fredholm Theorems and the Fredholm Alternative	637
13.5-1. Fredholm Theorems	637
13.5-2. Fredholm Alternative	638
13.6. Fredholm Integral Equations of the Second Kind with Symmetric Kernel	639
13.6-1. Characteristic Values and Eigenfunctions	639
13.6-2. Bilinear Series	640
13.6-3. Hilbert–Schmidt Theorem	641
13.6-4. Bilinear Series of Iterated Kernels	642
13.6-5. Solution of the Nonhomogeneous Equation	642
13.6-6. Fredholm Alternative for Symmetric Equations	643
13.6-7. Resolvent of a Symmetric Kernel	644
13.6-8. Extremal Properties of Characteristic Values and Eigenfunctions	644
13.6-9. Kellog’s Method for Finding Characteristic Values in the Case of Symmetric Kernel	645
13.6-10. Trace Method for the Approximation of Characteristic Values	646
13.6-11. Integral Equations Reducible to Symmetric Equations	647
13.6-12. Skew-Symmetric Integral Equations	647
13.6-13. Remark on Nonsymmetric Kernels	647
13.7. Integral Equations with Nonnegative Kernels	648
13.7-1. Positive Principal Eigenvalues. Generalized Jentzch Theorem	648
13.7-2. Positive Solutions of a Nonhomogeneous Integral Equation	649
13.7-3. Estimates for the Spectral Radius	649
13.7-4. Basic Definition and Theorems for Oscillating Kernels	651
13.7-5. Stochastic Kernels	654
13.8. Operator Method for Solving Integral Equations of the Second Kind	655
13.8-1. Simplest Scheme	655
13.8-2. Solution of Equations of the Second Kind on the Semiaxis	655

13.9. Methods of Integral Transforms and Model Solutions	656
13.9-1. Equation with Difference Kernel on the Entire Axis	656
13.9-2. Equation with the Kernel $K(x, t) = t^{-1}Q(x/t)$ on the Semiaxis	657
13.9-3. Equation with the Kernel $K(x, t) = t^{\beta}Q(xt)$ on the Semiaxis	658
13.9-4. Method of Model Solutions for Equations on the Entire Axis	659
13.10. Carleman Method for Integral Equations of Convolution Type of the Second Kind	660
13.10-1. Wiener–Hopf Equation of the Second Kind	660
13.10-2. Integral Equation of the Second Kind with Two Kernels	664
13.10-3. Equations of Convolution Type with Variable Integration Limit	668
13.10-4. Dual Equation of Convolution Type of the Second Kind	670
13.11. Wiener–Hopf Method	671
13.11-1. Some Remarks	671
13.11-2. Homogeneous Wiener–Hopf Equation of the Second Kind	673
13.11-3. General Scheme of the Method. The Factorization Problem	676
13.11-4. Nonhomogeneous Wiener–Hopf Equation of the Second Kind	677
13.11-5. Exceptional Case of a Wiener–Hopf Equation of the Second Kind	678
13.12. Krein’s Method for Wiener–Hopf Equations	679
13.12-1. Some Remarks. The Factorization Problem	679
13.12-2. Solution of the Wiener–Hopf Equations of the Second Kind	681
13.12-3. Hopf–Fock Formula	683
13.13. Methods for Solving Equations with Difference Kernels on a Finite Interval	683
13.13-1. Krein’s Method	683
13.13-2. Kernels with Rational Fourier Transforms	685
13.13-3. Reduction to Ordinary Differential Equations	686
13.14. Method of Approximating a Kernel by a Degenerate One	687
13.14-1. Approximation of the Kernel	687
13.14-2. Approximate Solution	688
13.15. Bateman Method	689
13.15-1. General Scheme of the Method	689
13.15-2. Some Special Cases	690
13.16. Collocation Method	692
13.16-1. General Remarks	692
13.16-2. Approximate Solution	693
13.16-3. Eigenfunctions of the Equation	694
13.17. Method of Least Squares	695
13.17-1. Description of the Method	695
13.17-2. Construction of Eigenfunctions	696
13.18. Bubnov–Galerkin Method	697
13.18-1. Description of the Method	697
13.18-2. Characteristic Values	697
13.19. Quadrature Method	698
13.19-1. General Scheme for Fredholm Equations of the Second Kind	698
13.19-2. Construction of the Eigenfunctions	699
13.19-3. Specific Features of the Application of Quadrature Formulas	700
13.20. Systems of Fredholm Integral Equations of the Second Kind	701
13.20-1. Some Remarks	701
13.20-2. Method of Reducing a System of Equations to a Single Equation	701

13.21. Regularization Method for Equations with Infinite Limits of Integration	702
13.21-1. Basic Equation and Fredholm Theorems	702
13.21-2. Regularizing Operators	703
13.21-3. Regularization Method	704
14. Methods for Solving Singular Integral Equations of the First Kind	707
14.1. Some Definitions and Remarks	707
14.1-1. Integral Equations of the First Kind with Cauchy Kernel	707
14.1-2. Integral Equations of the First Kind with Hilbert Kernel	707
14.2. Cauchy Type Integral	708
14.2-1. Definition of the Cauchy Type Integral	708
14.2-2. Hölder Condition	709
14.2-3. Principal Value of a Singular Integral	709
14.2-4. Multivalued Functions	711
14.2-5. Principal Value of a Singular Curvilinear Integral	712
14.2-6. Poincaré–Bertrand Formula	714
14.3. Riemann Boundary Value Problem	714
14.3-1. Principle of Argument. The Generalized Liouville Theorem	714
14.3-2. Hermite Interpolation Polynomial	716
14.3-3. Notion of the Index	716
14.3-4. Statement of the Riemann Problem	718
14.3-5. Solution of the Homogeneous Problem	720
14.3-6. Solution of the Nonhomogeneous Problem	721
14.3-7. Riemann Problem with Rational Coefficients	723
14.3-8. Riemann Problem for a Half-Plane	725
14.3-9. Exceptional Cases of the Riemann Problem	727
14.3-10. Riemann Problem for a Multiply Connected Domain	731
14.3-11. Riemann Problem for Open Curves	734
14.3-12. Riemann Problem with a Discontinuous Coefficient	739
14.3-13. Riemann Problem in the General Case	741
14.3-14. Hilbert Boundary Value Problem	742
14.4. Singular Integral Equations of the First Kind	743
14.4-1. Simplest Equation with Cauchy Kernel	743
14.4-2. Equation with Cauchy Kernel on the Real Axis	743
14.4-3. Equation of the First Kind on a Finite Interval	744
14.4-4. General Equation of the First Kind with Cauchy Kernel	745
14.4-5. Equations of the First Kind with Hilbert Kernel	746
14.5. Multhopp–Kalandiya Method	747
14.5-1. Solution That is Unbounded at the Endpoints of the Interval	747
14.5-2. Solution Bounded at One Endpoint of the Interval	749
14.5-3. Solution Bounded at Both Endpoints of the Interval	750
14.6. Hypersingular Integral Equations	751
14.6-1. Hypersingular Integral Equations with Cauchy- and Hilbert-Type Kernels	751
14.6-2. Definition of Hypersingular Integrals	751
14.6-3. Exact Solution of the Simplest Hypersingular Equation with Cauchy-Type Kernel	753
14.6-4. Exact Solution of the Simplest Hypersingular Equation with Hilbert-Type Kernel	754
14.6-5. Numerical Methods for Hypersingular Equations	754

15. Methods for Solving Complete Singular Integral Equations	757
15.1. Some Definitions and Remarks	757
15.1-1. Integral Equations with Cauchy Kernel	757
15.1-2. Integral Equations with Hilbert Kernel	759
15.1-3. Fredholm Equations of the Second Kind on a Contour	759
15.2. Carleman Method for Characteristic Equations	761
15.2-1. Characteristic Equation with Cauchy Kernel	761
15.2-2. Transposed Equation of a Characteristic Equation	764
15.2-3. Characteristic Equation on the Real Axis	765
15.2-4. Exceptional Case of a Characteristic Equation	767
15.2-5. Characteristic Equation with Hilbert Kernel	769
15.2-6. Tricomi Equation	769
15.3. Complete Singular Integral Equations Solvable in a Closed Form	770
15.3-1. Closed-Form Solutions in the Case of Constant Coefficients	770
15.3-2. Closed-Form Solutions in the General Case	771
15.4. Regularization Method for Complete Singular Integral Equations	772
15.4-1. Certain Properties of Singular Operators	772
15.4-2. Regularizer	774
15.4-3. Methods of Left and Right Regularization	775
15.4-4. Problem of Equivalent Regularization	776
15.4-5. Fredholm Theorems	777
15.4-6. Carleman–Vekua Approach to the Regularization	778
15.4-7. Regularization in Exceptional Cases	779
15.4-8. Complete Equation with Hilbert Kernel	780
15.5. Analysis of Solutions Singularities for Complete Integral Equations with Generalized Cauchy Kernels	783
15.5-1. Statement of the Problem and Preliminary Remarks	783
15.5-2. Auxiliary Results	784
15.5-3. Equations for the Exponents of Singularity of a Solution	787
15.5-4. Analysis of Equations for Singularity Exponents	789
15.5-5. Application to an Equation Arising in Fracture Mechanics	791
15.6. Direct Numerical Solution of Singular Integral Equations with Generalized Kernels ..	792
15.6-1. Preliminary Remarks	792
15.6-2. Quadrature Formulas for Integrals with the Jacobi Weight Function	793
15.6-3. Approximation of Solutions in Terms of a System of Orthogonal Polynomials	795
15.6-4. Some Special Functions and Their Calculations	797
15.6-5. Numerical Solution of Singular Integral Equations	799
15.6-6. Numerical Solutions of Singular Integral Equations of Bueckner Type	801
16. Methods for Solving Nonlinear Integral Equations	805
16.1. Some Definitions and Remarks	805
16.1-1. Nonlinear Equations with Variable Limit of Integration (Volterra Equations) ..	805
16.1-2. Nonlinear Equations with Constant Integration Limits (Urysohn Equations) ..	806
16.1-3. Some Special Features of Nonlinear Integral Equations	807
16.2. Exact Methods for Nonlinear Equations with Variable Limit of Integration	809
16.2-1. Method of Integral Transforms	809
16.2-2. Method of Differentiation for Nonlinear Equations with Degenerate Kernel ..	810

16.3. Approximate and Numerical Methods for Nonlinear Equations with Variable Limit of Integration	811
16.3-1. Successive Approximation Method	811
16.3-2. Newton–Kantorovich Method	813
16.3-3. Collocation Method	815
16.3-4. Quadrature Method	816
16.4. Exact Methods for Nonlinear Equations with Constant Integration Limits	817
16.4-1. Nonlinear Equations with Degenerate Kernels	817
16.4-2. Method of Integral Transforms	819
16.4-3. Method of Differentiating for Integral Equations	820
16.4-4. Method for Special Urysohn Equations of the First Kind	821
16.4-5. Method for Special Urysohn Equations of the Second Kind	822
16.4-6. Some Generalizations	824
16.5. Approximate and Numerical Methods for Nonlinear Equations with Constant Integration Limits	826
16.5-1. Successive Approximation Method	826
16.5-2. Newton–Kantorovich Method	827
16.5-3. Quadrature Method	829
16.5-4. Tikhonov Regularization Method	829
16.6. Existence and Uniqueness Theorems for Nonlinear Equations	830
16.6-1. Hammerstein Equations	830
16.6-2. Urysohn Equations	832
16.7. Nonlinear Equations with a Parameter: Eigenfunctions, Eigenvalues, Bifurcation Points	834
16.7-1. Eigenfunctions and Eigenvalues of Nonlinear Integral Equations	834
16.7-2. Local Solutions of a Nonlinear Integral Equation with a Parameter	835
16.7-3. Bifurcation Points of Nonlinear Integral Equations	835
17. Methods for Solving Multidimensional Mixed Integral Equations	839
17.1. Some Definition and Remarks	839
17.1-1. Basic Classes of Functions	839
17.1-2. Mixed Equations on a Finite Interval	840
17.1-3. Mixed Equation on a Ring-Shaped (Circular) Domain	841
17.1-4. Mixed Equations on a Closed Bounded Set	842
17.2. Methods of Solution of Mixed Integral Equations on a Finite Interval	843
17.2-1. Equation with a Hilbert–Schmidt Kernel and a Given Right-Hand Side	843
17.2-2. Equation with Hilbert–Schmidt Kernel and Auxiliary Conditions	845
17.2-3. Equation with a Schmidt Kernel and a Given Right-Hand Side on an Interval	848
17.2-4. Equation with a Schmidt Kernel and Auxiliary Conditions	851
17.3. Methods of Solving Mixed Integral Equations on a Ring-Shaped Domain	855
17.3-1. Equation with a Hilbert–Schmidt Kernel and a Given Right-Hand Side	855
17.3-2. Equation with a Hilbert–Schmidt Kernel and Auxiliary Conditions	856
17.3-3. Equation with a Schmidt Kernel and a Given Right-Hand Side	859
17.3-4. Equation with a Schmidt Kernel and Auxiliary Conditions on Ring-Shaped Domain	862
17.4. Projection Method for Solving Mixed Equations on a Bounded Set	866
17.4-1. Mixed Operator Equation with a Given Right-Hand Side	866
17.4-2. Mixed Operator Equations with Auxiliary Conditions	869
17.4-3. General Projection Problem for Operator Equation	873

18. Application of Integral Equations for the Investigation of Differential Equations . . .	875
18.1. Reduction of the Cauchy Problem for ODEs to Integral Equations	875
18.1-1. Cauchy Problem for First-Order ODEs. Uniqueness and Existence Theorems	875
18.1-2. Cauchy Problem for First-Order ODEs. Method of Successive Approximations	876
18.1-3. Cauchy Problem for Second-Order ODEs. Method of Successive Approximations	876
18.1-4. Cauchy Problem for a Special n -Order Linear ODE	876
18.2. Reduction of Boundary Value Problems for ODEs to Volterra Integral Equations. Calculation of Eigenvalues	877
18.2-1. Reduction of Differential Equations to Volterra Integral Equations	877
18.2-2. Application of Volterra Equations to the Calculation of Eigenvalues	879
18.3. Reduction of Boundary Value Problems for ODEs to Fredholm Integral Equations with the Help of the Green's Function	881
18.3-1. Linear Ordinary Differential Equations. Fundamental Solutions	881
18.3-2. Boundary Value Problems for n th Order Differential Equations. Green's Function	882
18.3-3. Boundary Value Problems for Second-Order Differential Equations. Green's Function	883
18.3-4. Nonlinear Problem of Nonisothermal Flow in Plane Channel	884
18.4. Reduction of PDEs with Boundary Conditions of the Third Kind to Integral Equations	887
18.4-1. Usage of Particular Solutions of PDEs for the Construction of Other Solutions	887
18.4-2. Mass Transfer to a Particle in Fluid Flow Complicated by a Surface Reaction	888
18.4-3. Integral Equations for Surface Concentration and Diffusion Flux	890
18.4-4. Method of Numerical Integration of the Equation for Surface Concentration .	891
18.5. Representation of Linear Boundary Value Problems in Terms of Potentials	892
18.5-1. Basic Types of Potentials for the Laplace Equation and Their Properties	892
18.5-2. Integral Identities. Green's Formula	895
18.5-3. Reduction of Interior Dirichlet and Neumann Problems to Integral Equations .	895
18.5-4. Reduction of Exterior Dirichlet and Neumann Problems to Integral Equations	896
18.6. Representation of Solutions of Nonlinear PDEs in Terms of Solutions of Linear Integral Equations (Inverse Scattering)	898
18.6-1. Description of the Zakharov–Shabat Method	898
18.6-2. Korteweg–de Vries Equation and Other Nonlinear Equations	899

Supplements

Supplement 1. Elementary Functions and Their Properties	905
1.1. Power, Exponential, and Logarithmic Functions	905
1.1-1. Properties of the Power Function	905
1.1-2. Properties of the Exponential Function	905
1.1-3. Properties of the Logarithmic Function	906
1.2. Trigonometric Functions	907
1.2-1. Simplest Relations	907
1.2-2. Reduction Formulas	907
1.2-3. Relations Between Trigonometric Functions of Single Argument	908
1.2-4. Addition and Subtraction of Trigonometric Functions	908
1.2-5. Products of Trigonometric Functions	908
1.2-6. Powers of Trigonometric Functions	908
1.2-7. Addition Formulas	909

1.2-8. Trigonometric Functions of Multiple Arguments	909
1.2-9. Trigonometric Functions of Half Argument	909
1.2-10. Differentiation Formulas	910
1.2-11. Integration Formulas	910
1.2-12. Expansion in Power Series	910
1.2-13. Representation in the Form of Infinite Products	910
1.2-14. Euler and de Moivre Formulas. Relationship with Hyperbolic Functions	911
1.3. Inverse Trigonometric Functions	911
1.3-1. Definitions of Inverse Trigonometric Functions	911
1.3-2. Simplest Formulas	912
1.3-3. Some Properties	912
1.3-4. Relations Between Inverse Trigonometric Functions	912
1.3-5. Addition and Subtraction of Inverse Trigonometric Functions	912
1.3-6. Differentiation Formulas	913
1.3-7. Integration Formulas	913
1.3-8. Expansion in Power Series	913
1.4. Hyperbolic Functions	913
1.4-1. Definitions of Hyperbolic Functions	913
1.4-2. Simplest Relations	913
1.4-3. Relations Between Hyperbolic Functions of Single Argument ($x \geq 0$)	914
1.4-4. Addition and Subtraction of Hyperbolic Functions	914
1.4-5. Products of Hyperbolic Functions	914
1.4-6. Powers of Hyperbolic Functions	914
1.4-7. Addition Formulas	915
1.4-8. Hyperbolic Functions of Multiple Argument	915
1.4-9. Hyperbolic Functions of Half Argument	915
1.4-10. Differentiation Formulas	916
1.4-11. Integration Formulas	916
1.4-12. Expansion in Power Series	916
1.4-13. Representation in the Form of Infinite Products	916
1.4-14. Relationship with Trigonometric Functions	916
1.5. Inverse Hyperbolic Functions	917
1.5-1. Definitions of Inverse Hyperbolic Functions	917
1.5-2. Simplest Relations	917
1.5-3. Relations Between Inverse Hyperbolic Functions	917
1.5-4. Addition and Subtraction of Inverse Hyperbolic Functions	917
1.5-5. Differentiation Formulas	917
1.5-6. Integration Formulas	918
1.5-7. Expansion in Power Series	918
Supplement 2. Finite Sums and Infinite Series	919
2.1. Finite Numerical Sums	919
2.1-1. Progressions	919
2.1-2. Sums of Powers of Natural Numbers Having the Form $\sum k^m$	919
2.1-3. Alternating Sums of Powers of Natural Numbers, $\sum (-1)^k k^m$	920
2.1-4. Other Sums Containing Integers	920
2.1-5. Sums Containing Binomial Coefficients	920
2.1-6. Other Numerical Sums	921

2.2. Finite Functional Sums	922
2.2-1. Sums Involving Hyperbolic Functions	922
2.2-2. Sums Involving Trigonometric Functions	922
2.3. Infinite Numerical Series	924
2.3-1. Progressions	924
2.3-2. Other Numerical Series	924
2.4. Infinite Functional Series	925
2.4-1. Power Series	925
2.4-2. Trigonometric Series in One Variable Involving Sine	927
2.4-3. Trigonometric Series in One Variable Involving Cosine	928
2.4-4. Trigonometric Series in Two Variables	930
Supplement 3. Tables of Indefinite Integrals	933
3.1. Integrals Involving Rational Functions	933
3.1-1. Integrals Involving $a + bx$	933
3.1-2. Integrals Involving $a + x$ and $b + x$	933
3.1-3. Integrals Involving $a^2 + x^2$	934
3.1-4. Integrals Involving $a^2 - x^2$	935
3.1-5. Integrals Involving $a^3 + x^3$	936
3.1-6. Integrals Involving $a^3 - x^3$	936
3.1-7. Integrals Involving $a^4 \pm x^4$	937
3.2. Integrals Involving Irrational Functions	937
3.2-1. Integrals Involving $x^{1/2}$	937
3.2-2. Integrals Involving $(a + bx)^{p/2}$	938
3.2-3. Integrals Involving $(x^2 + a^2)^{1/2}$	938
3.2-4. Integrals Involving $(x^2 - a^2)^{1/2}$	938
3.2-5. Integrals Involving $(a^2 - x^2)^{1/2}$	939
3.2-6. Integrals Involving Arbitrary Powers. Reduction Formulas	939
3.3. Integrals Involving Exponential Functions	940
3.4. Integrals Involving Hyperbolic Functions	940
3.4-1. Integrals Involving $\cosh x$	940
3.4-2. Integrals Involving $\sinh x$	941
3.4-3. Integrals Involving $\tanh x$ or $\coth x$	942
3.5. Integrals Involving Logarithmic Functions	943
3.6. Integrals Involving Trigonometric Functions	944
3.6-1. Integrals Involving $\cos x$ ($n = 1, 2, \dots$)	944
3.6-2. Integrals Involving $\sin x$ ($n = 1, 2, \dots$)	945
3.6-3. Integrals Involving $\sin x$ and $\cos x$	947
3.6-4. Reduction Formulas	947
3.6-5. Integrals Involving $\tan x$ and $\cot x$	947
3.7. Integrals Involving Inverse Trigonometric Functions	948
Supplement 4. Tables of Definite Integrals	951
4.1. Integrals Involving Power-Law Functions	951
4.1-1. Integrals Over a Finite Interval	951
4.1-2. Integrals Over an Infinite Interval	952
4.2. Integrals Involving Exponential Functions	954
4.3. Integrals Involving Hyperbolic Functions	955
4.4. Integrals Involving Logarithmic Functions	955


4.5. Integrals Involving Trigonometric Functions	956
4.5-1. Integrals Over a Finite Interval	956
4.5-2. Integrals Over an Infinite Interval	957
4.6. Integrals Involving Bessel Functions	958
4.6-1. Integrals Over an Infinite Interval	958
4.6-2. Other Integrals	959
Supplement 5. Tables of Laplace Transforms	961
5.1. General Formulas	961
5.2. Expressions with Power-Law Functions	963
5.3. Expressions with Exponential Functions	963
5.4. Expressions with Hyperbolic Functions	964
5.5. Expressions with Logarithmic Functions	965
5.6. Expressions with Trigonometric Functions	966
5.7. Expressions with Special Functions	967
Supplement 6. Tables of Inverse Laplace Transforms	969
6.1. General Formulas	969
6.2. Expressions with Rational Functions	971
6.3. Expressions with Square Roots	975
6.4. Expressions with Arbitrary Powers	977
6.5. Expressions with Exponential Functions	978
6.6. Expressions with Hyperbolic Functions	979
6.7. Expressions with Logarithmic Functions	980
6.8. Expressions with Trigonometric Functions	981
6.9. Expressions with Special Functions	981
Supplement 7. Tables of Fourier Cosine Transforms	983
7.1. General Formulas	983
7.2. Expressions with Power-Law Functions	983
7.3. Expressions with Exponential Functions	984
7.4. Expressions with Hyperbolic Functions	985
7.5. Expressions with Logarithmic Functions	985
7.6. Expressions with Trigonometric Functions	986
7.7. Expressions with Special Functions	987
Supplement 8. Tables of Fourier Sine Transforms	989
8.1. General Formulas	989
8.2. Expressions with Power-Law Functions	989
8.3. Expressions with Exponential Functions	990
8.4. Expressions with Hyperbolic Functions	991
8.5. Expressions with Logarithmic Functions	992
8.6. Expressions with Trigonometric Functions	992
8.7. Expressions with Special Functions	993

Supplement 9. Tables of Mellin Transforms	997
9.1. General Formulas	997
9.2. Expressions with Power-Law Functions	998
9.3. Expressions with Exponential Functions	998
9.4. Expressions with Logarithmic Functions	999
9.5. Expressions with Trigonometric Functions	999
9.6. Expressions with Special Functions	1000
Supplement 10. Tables of Inverse Mellin Transforms	1001
10.1. Expressions with Power-Law Functions	1001
10.2. Expressions with Exponential and Logarithmic Functions	1002
10.3. Expressions with Trigonometric Functions	1003
10.4. Expressions with Special Functions	1004
Supplement 11. Special Functions and Their Properties	1007
11.1. Some Coefficients, Symbols, and Numbers	1007
11.1-1. Binomial Coefficients	1007
11.1-2. Pochhammer Symbol	1007
11.1-3. Bernoulli Numbers	1008
11.1-4. Euler Numbers	1008
11.2. Error Functions. Exponential and Logarithmic Integrals	1009
11.2-1. Error Function and Complementary Error Function	1009
11.2-2. Exponential Integral	1010
11.2-3. Logarithmic Integral	1010
11.3. Sine Integral and Cosine Integral. Fresnel Integrals	1011
11.3-1. Sine Integral	1011
11.3-2. Cosine Integral	1011
11.3-3. Fresnel Integrals and Generalized Fresnel Integrals	1012
11.4. Gamma Function, Psi Function, and Beta Function	1012
11.4-1. Gamma Function	1012
11.4-2. Psi Function (Digamma Function)	1013
11.4-3. Beta Function	1014
11.5. Incomplete Gamma and Beta Functions	1014
11.5-1. Incomplete Gamma Function	1014
11.5-2. Incomplete Beta Function	1015
11.6. Bessel Functions (Cylindrical Functions)	1016
11.6-1. Definitions and Basic Formulas	1016
11.6-2. Integral Representations and Asymptotic Expansions	1017
11.6-3. Zeros of Bessel Functions	1019
11.6-4. Orthogonality Properties of Bessel Functions	1019
11.6-5. Hankel Functions (Bessel Functions of the Third Kind)	1020
11.7. Modified Bessel Functions	1021
11.7-1. Definitions. Basic Formulas	1021
11.7-2. Integral Representations and Asymptotic Expansions	1022
11.8. Airy Functions	1023
11.8-1. Definition and Basic Formulas	1023
11.8-2. Power Series and Asymptotic Expansions	1023

11.9. Confluent Hypergeometric Functions	1024
11.9-1. Kummer and Tricomi Confluent Hypergeometric Functions	1024
11.9-2. Integral Representations and Asymptotic Expansions	1027
11.9-3. Whittaker Confluent Hypergeometric Functions	1027
11.10. Gauss Hypergeometric Functions	1028
11.10-1. Various Representations of the Gauss Hypergeometric Function	1028
11.10-2. Basic Properties	1028
11.11. Legendre Polynomials, Legendre Functions, and Associated Legendre Functions	1030
11.11-1. Legendre Polynomials and Legendre Functions	1030
11.11-2. Associated Legendre Functions with Integer Indices and Real Argument	1031
11.11-3. Associated Legendre Functions. General Case	1032
11.12. Parabolic Cylinder Functions	1034
11.12-1. Definitions. Basic Formulas	1034
11.12-2. Integral Representations, Asymptotic Expansions, and Linear Relations	1035
11.13. Elliptic Integrals	1035
11.13-1. Complete Elliptic Integrals	1035
11.13-2. Incomplete Elliptic Integrals (Elliptic Integrals)	1037
11.14. Elliptic Functions	1038
11.14-1. Jacobi Elliptic Functions	1039
11.14-2. Weierstrass Elliptic Function	1042
11.15. Jacobi Theta Functions	1043
11.15-1. Series Representation of the Jacobi Theta Functions. Simplest Properties	1043
11.15-2. Various Relations and Formulas. Connection with Jacobi Elliptic Functions	1044
11.16. Mathieu Functions and Modified Mathieu Functions	1045
11.16-1. Mathieu Functions	1045
11.16-2. Modified Mathieu Functions	1046
11.17. Orthogonal Polynomials	1047
11.17-1. Laguerre Polynomials and Generalized Laguerre Polynomials	1047
11.17-2. Chebyshev Polynomials and Functions	1048
11.17-3. Hermite Polynomials and Functions	1050
11.17-4. Jacobi Polynomials	1051
11.17-5. Gegenbauer Polynomials	1051
11.18. Nonorthogonal Polynomials	1052
11.18-1. Bernoulli Polynomials	1052
11.18-2. Euler Polynomials	1053
Supplement 12. Some Notions of Functional Analysis	1055
12.1. Functions of Bounded Variation	1055
12.1-1. Definition of a Function of Bounded Variation	1055
12.1-2. Classes of Functions of Bounded Variation	1056
12.1-3. Properties of Functions of Bounded Variation	1056
12.1-4. Criteria for Functions to Have Bounded Variation	1057
12.1-5. Properties of Continuous Functions of Bounded Variation	1057
12.2. Stieltjes Integral	1057
12.2-1. Basic Definitions	1057
12.2-2. Properties of the Stieltjes Integral	1058
12.2-3. Existence Theorems for the Stieltjes Integral	1058

12.3. Lebesgue Integral	1059
12.3-1. Riemann Integral and the Lebesgue Integral	1059
12.3-2. Sets of Zero Measure. Notion of “Almost Everywhere”	1060
12.3-3. Step Functions and Measurable Functions	1060
12.3-4. Definition and Properties of the Lebesgue Integral	1061
12.3-5. Measurable Sets	1062
12.3-6. Integration Over Measurable Sets	1063
12.3-7. Case of an Infinite Interval	1063
12.3-8. Case of Several Variables	1064
12.3-9. Spaces L_p	1064
12.4. Linear Normed Spaces	1065
12.4-1. Linear Spaces	1065
12.4-2. Linear Normed Spaces	1065
12.4-3. Space of Continuous Functions $C(a, b)$	1066
12.4-4. Lebesgue Space $L_p(a, b)$	1066
12.4-5. Hölder Space $C_\alpha(0, 1)$	1066
12.4-6. Space of Functions of Bounded Variation $V(0, 1)$	1066
12.5. Euclidean and Hilbert Spaces. Linear Operators in Hilbert Spaces	1067
12.5-1. Preliminary Remarks	1067
12.5-2. Euclidean and Hilbert Spaces	1067
12.5-3. Linear Operators in Hilbert Spaces	1068
References	1071
Index	1081

AUTHORS



Andrei D. Polyinin, D.Sc., Ph.D., is a well-known scientist of broad interests and is active in various areas of mathematics, mechanics, and chemical engineering sciences. He is one of the most prominent authors in the field of reference literature on mathematics and physics.

Professor Polyinin graduated with honors from the Department of Mechanics and Mathematics of Moscow State University in 1974. He received his Ph.D. degree in 1981 and D.Sc. degree in 1986 at the Institute for Problems in Mechanics of the Russian (former USSR) Academy of Sciences. Since 1975, Professor Polyinin has been working at the Institute for Problems in Mechanics of the Russian Academy of Sciences; he is also Professor of Mathematics at Bauman Moscow State Technical University. He is a member of the Russian National Committee on Theoretical and Applied Mechanics and of the Mathematics and Mechanics Expert Council of

the Higher Certification Committee of the Russian Federation.

Professor Polyinin has made important contributions to exact and approximate analytical methods in the theory of differential equations, mathematical physics, integral equations, engineering mathematics, theory of heat and mass transfer, and chemical hydrodynamics. He obtained exact solutions for several thousand ordinary differential, partial differential, and integral equations.

Professor Polyinin is an author of more than 30 books in English, Russian, German, and Bulgarian as well as over 120 research papers and three patents. He has written a number of fundamental handbooks, including A. D. Polyinin and V. F. Zaitsev, *Handbook of Exact Solutions for Ordinary Differential Equations*, CRC Press, 1995 and 2003; A. D. Polyinin and A. V. Manzhirov, *Handbook of Integral Equations*, CRC Press, 1998; A. D. Polyinin, *Handbook of Linear Partial Differential Equations for Engineers and Scientists*, Chapman & Hall/CRC Press, 2002; A. D. Polyinin, V. F. Zaitsev, and A. Moussiaux, *Handbook of First Order Partial Differential Equations*, Taylor & Francis, 2002; A. D. Polyinin and V. F. Zaitsev, *Handbook of Nonlinear Partial Differential Equations*, Chapman & Hall/CRC Press, 2004, and A. D. Polyinin and A. V. Manzhirov, *Handbook of Mathematics for Engineers and Scientists*, Chapman & Hall/CRC Press, 2007.

Professor Polyinin is editor of the book series *Differential and Integral Equations and Their Applications*, Chapman & Hall/CRC Press, London/Boca Raton, and *Physical and Mathematical Reference Literature*, Fizmatlit, Moscow. He is also Editor-in-Chief of the international scientific-educational Website *EqWorld—The World of Mathematical Equations* (<http://eqworld.ipmnet.ru>), which is visited by over 1700 users a day worldwide. Professor Polyinin is a member of the Editorial Board of the journal *Theoretical Foundations of Chemical Engineering*.

In 1991, Professor Polyinin was awarded a Chaplygin Prize of the Russian Academy of Sciences for his research in mechanics. In 2001, he received an award from the Ministry of Education of the Russian Federation.

Address: Institute for Problems in Mechanics, Vernadsky Ave. 101 Bldg 1, 119526 Moscow, Russia

Home page: <http://eqworld.ipmnet.ru/polyinin-ew.htm>



Alexander V. Manzhirov, D.Sc., Ph.D., is a noted scientist in the fields of mechanics and applied mathematics, integral equations, and their applications.

After graduating with honors from the Department of Mechanics and Mathematics of Rostov State University in 1979, Alexander Manzhirov attended postgraduate courses at Moscow Institute of Civil Engineering. He received his Ph.D. degree in 1983 at Moscow Institute of Electronic Engineering Industry and D.Sc. degree in 1993 at the Institute for Problems in Mechanics of the Russian (former USSR) Academy of Sciences. Since 1983, Alexander Manzhirov has been working at the Institute for Problems in Mechanics of the Russian Academy of Sciences. Currently, he is head of the Laboratory for Modeling in Solid Mechanics at the same institute.

Professor Manzhirov is also head of a branch of the Department of Applied Mathematics at Bauman Moscow State Technical University, professor of mathematics at Moscow State University of Engineering and Computer Science, vice-chairman of Mathematics and Mechanics Expert Council of the Higher Certification Committee of the Russian Federation, executive secretary of Solid Mechanics Scientific Council of the Russian Academy of Sciences, and expert in mathematics, mechanics, and computer science of the Russian Foundation for Basic Research. He is a member of the Russian National Committee on Theoretical and Applied Mechanics and the European Mechanics Society (EUROMECH), and member of the editorial board of the journal *Mechanics of Solids* and the international scientific-educational Website *EqWorld—The World of Mathematical Equations* (<http://eqworld.ipmnet.ru>).

Professor Manzhirov has made important contributions to new mathematical methods for solving problems in the fields of integral equations and their applications, mechanics of growing solids, contact mechanics, tribology, viscoelasticity, and creep theory. He is an author of more than ten books (including *Contact Problems in Mechanics of Growing Solids* [in Russian], Nauka, Moscow, 1991; *Handbook of Integral Equations*, CRC Press, Boca Raton, 1998; *Handbuch der Integralgleichungen: Exakte Lösungen*, Spektrum Akad. Verlag, Heidelberg, 1999; *Contact Problems in the Theory of Creep* [in Russian], National Academy of Sciences of Armenia, Yerevan, 1999; A. D. Polyanin and A. V. Manzhirov, *Handbook of Mathematics for Engineers and Scientists*, Chapman & Hall/CRC Press, Boca Raton, 2007), more than 70 research papers, and two patents.

Professor Manzhirov is a winner of the First Competition of the Science Support Foundation 2001, Moscow.

Address: Institute for Problems in Mechanics, Vernadsky Ave. 101 Bldg 1, 119526 Moscow, Russia.

Home page: <http://eqworld.ipmnet.ru/en/board/manzhirov.htm>.

PREFACE TO THE NEW EDITION

Handbook of Integral Equations, Second Edition, a unique reference for engineers and scientists, contains over 2,500 integral equations with solutions, as well as analytical and numerical methods for solving linear and nonlinear equations. It considers Volterra, Fredholm, Wiener–Hopf, Hammerstein, Urysohn, and other equations, which arise in mathematics, physics, engineering sciences, economics, etc. In total, the number of equations described is an order of magnitude greater than in any other book available.

The second edition has been substantially updated, revised, and extended. It includes new chapters on mixed multidimensional equations, methods of integral equations for ODEs and PDEs, and about 400 new equations with exact solutions. It presents a considerable amount of new material on Volterra, Fredholm, singular, hypersingular, dual, and nonlinear integral equations, integral transforms, and special functions. Many examples were added for illustrative purposes. The new edition has been increased by a total of over 300 pages.

Note that the first part of the book can be used as a database of test problems for numerical and approximate methods for solving linear and nonlinear integral equations.

We would like to express our deep gratitude to Alexei Zhurov and Vasilii Silvestrov for fruitful discussions. We also appreciate the help of Grigory Yosifian in translating new sections of this book and valuable remarks.

The authors hope that the handbook will prove helpful for a wide audience of researchers, college and university teachers, engineers, and students in various fields of applied mathematics, mechanics, physics, chemistry, biology, economics, and engineering sciences.

A. D. Polyinin

A. V. Manzhirov

PREFACE TO THE FIRST EDITION

Integral equations are encountered in various fields of science and numerous applications (in elasticity, plasticity, heat and mass transfer, oscillation theory, fluid dynamics, filtration theory, electrostatics, electrodynamics, biomechanics, game theory, control, queuing theory, electrical engineering, economics, medicine, etc.).

Exact (closed-form) solutions of integral equations play an important role in the proper understanding of qualitative features of many phenomena and processes in various areas of natural science. Lots of equations of physics, chemistry, and biology contain functions or parameters which are obtained from experiments and hence are not strictly fixed. Therefore, it is expedient to choose the structure of these functions so that it would be easier to analyze and solve the equation. As a possible selection criterion, one may adopt the requirement that the model integral equation admits a solution in a closed form. Exact solutions can be used to verify the consistency and estimate errors of various numerical, asymptotic, and approximate methods.

More than 2,100 integral equations and their solutions are given in the first part of the book (Chapters 1–6). A lot of new exact solutions to linear and nonlinear equations are included. Special attention is paid to equations of general form, which depend on arbitrary functions. The other equations contain one or more free parameters (the book actually deals with families of integral

equations); it is the reader's option to fix these parameters. In total, the number of equations described in this handbook is an order of magnitude greater than in any other book currently available.

The second part of the book (Chapters 7–14) presents exact, approximate analytical, and numerical methods for solving linear and nonlinear integral equations. Apart from the classical methods, some new methods are also described. When selecting the material, the authors have given a pronounced preference to practical aspects of the matter; that is, to methods that allow effectively “constructing” the solution. For the reader's better understanding of the methods, each section is supplied with examples of specific equations. Some sections may be used by lecturers of colleges and universities as a basis for courses on integral equations and mathematical physics equations for graduate and postgraduate students.

For the convenience of a wide audience with different mathematical backgrounds, the authors tried to do their best, wherever possible, to avoid special terminology. Therefore, some of the methods are outlined in a schematic and somewhat simplified manner, with necessary references made to books where these methods are considered in more detail. For some nonlinear equations, only solutions of the simplest form are given. The book does not cover two-, three-, and multidimensional integral equations.

The handbook consists of chapters, sections, and subsections. Equations and formulas are numbered separately in each section. The equations within a section are arranged in increasing order of complexity. The extensive table of contents provides rapid access to the desired equations.

For the reader's convenience, the main material is followed by a number of supplements, where some properties of elementary and special functions are described, tables of indefinite and definite integrals are given, as well as tables of Laplace, Mellin, and other transforms, which are used in the book.

The first and second parts of the book, just as many sections, were written so that they could be read independently from each other. This allows the reader to quickly get to the heart of the matter.

We would like to express our deep gratitude to Rolf Sulanke and Alexei Zhurov for fruitful discussions and valuable remarks. We also appreciate the help of Vladimir Nazaikinskii and Alexander Shtern in translating the second part of this book, and are thankful to Inna Shingareva for her assistance in preparing the camera-ready copy of the book.

The authors hope that the handbook will prove helpful for a wide audience of researchers, college and university teachers, engineers, and students in various fields of mathematics, mechanics, physics, chemistry, biology, economics, and engineering sciences.

A. D. Polyinin
A. V. Manzhirov

SOME REMARKS AND NOTATION

1. In Chapters 1–11, 14, and 18 in the original integral equations, the independent variable is denoted by x , the integration variable by t , and the unknown function by $y = y(x)$.

2. For a function of one variable $f = f(x)$, we use the following notation for the derivatives:

$$f'_x = \frac{df}{dx}, \quad f''_{xx} = \frac{d^2f}{dx^2}, \quad f'''_{xxx} = \frac{d^3f}{dx^3}, \quad f''''_{xxxx} = \frac{d^4f}{dx^4}, \quad \text{and} \quad f^{(n)}_x = \frac{d^n f}{dx^n} \quad \text{for } n \geq 5.$$

Occasionally, we use the similar notation for partial derivatives of a function of two variables, for example, $K'_x(x, t) = \frac{\partial}{\partial x} K(x, t)$.

3. In some cases, we use the operator notation $\left[f(x) \frac{d}{dx} \right]^n g(x)$, which is defined recursively by

$$\left[f(x) \frac{d}{dx} \right]^n g(x) = f(x) \frac{d}{dx} \left\{ \left[f(x) \frac{d}{dx} \right]^{n-1} g(x) \right\}.$$

4. It is indicated in the beginning of Chapters 1–8 that $f = f(x)$, $g = g(x)$, $K = K(x)$, etc. are arbitrary functions, and A , B , etc. are free parameters. This means that:

- (a) $f = f(x)$, $g = g(x)$, $K = K(x)$, etc. are assumed to be continuous real-valued functions of real arguments;*
- (b) if the solution contains derivatives of these functions, then the functions are assumed to be sufficiently differentiable;**
- (c) if the solution contains integrals with these functions (in combination with other functions), then the integrals are supposed to converge;
- (d) the free parameters A , B , etc. may assume any real values for which the expressions occurring in the equation and the solution make sense (for example, if a solution contains a factor $\frac{A}{1-A}$, then it is implied that $A \neq 1$; as a rule, this is not specified in the text).

5. The notations $\text{Re } z$ and $\text{Im } z$ stand, respectively, for the real and the imaginary part of a complex quantity z .

6. In the first part of the book (Chapters 1–8) when referencing a particular equation, we use a notation like 2.3.15, which implies equation 15 from Section 2.3.

7. To highlight portions of the text, the following symbols are used in the book:

- ▶ indicates important information pertaining to a group of equations (Chapters 1–8);
- ⊙ indicates the literature used in the preparation of the text in specific equations (Chapters 1–8) or sections (Chapters 9–18).

* Less severe restrictions on these functions are presented in the second part of the book.

** Restrictions (b) and (c) imposed on $f = f(x)$, $g = g(x)$, $K = K(x)$, etc. are not mentioned in the text.

Part I

**Exact Solutions of
Integral Equations**

Chapter 1

Linear Equations of the First Kind with Variable Limit of Integration

► **Notation:** $f = f(x)$, $g = g(x)$, $h = h(x)$, $K = K(x)$, and $M = M(x)$ are arbitrary functions (these may be composite functions of the argument depending on two variables x and t); $A, B, C, D, E, a, b, c, \alpha, \beta, \gamma, \lambda$, and μ are free parameters; and m and n are nonnegative integers.

► **Preliminary remarks.** For equations of the form

$$\int_a^x K(x, t)y(t) dt = f(x), \quad a \leq x \leq b,$$

where the functions $K(x, t)$ and $f(x)$ are continuous, the right-hand side must satisfy the following conditions:

1°. If $K(a, a) \neq 0$, then we must have $f(a) = 0$ (for example, the right-hand sides of equations 1.1.1 and 1.2.1 must satisfy this condition).

2°. If $K(a, a) = K'_x(a, a) = \dots = K_x^{(n-1)}(a, a) = 0$, $0 < |K_x^{(n)}(a, a)| < \infty$, then the right-hand side of the equation must satisfy the conditions

$$f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0.$$

For example, with $n = 1$, these are constraints for the right-hand side of equation 1.1.2.

3°. If $K(a, a) = K'_x(a, a) = \dots = K_x^{(n-1)}(a, a) = 0$, $K_x^{(n)}(a, a) = \infty$, then the right-hand side of the equation must satisfy the conditions

$$f(a) = f'_x(a) = \dots = f_x^{(n-1)}(a) = 0.$$

For example, with $n = 1$, this is a constraint for the right-hand side of equation 1.1.30.

4°. For unbounded $K(x, t)$ with integrable power-law or logarithmic singularity at $x = t$ and continuous $f(x)$, no additional conditions are imposed on the right-hand side of the integral equation (e.g., see Abel's equation 1.1.36).

In the case of a difference kernel, $K(x, t) = K(x - t)$, that can be represented as $x \rightarrow t$ in the form

$$K(x - t) = A(x - t)^\lambda + o((x - t)^\lambda) \quad (0 < |A| < \infty),$$

the right-hand side of the integral equation, for $\lambda \geq 0$, must satisfy the conditions

$$f(a) = f'_x(a) = \dots = f_x^{([\lambda])}(a) = 0,$$

where $[\lambda]$ is the integer part of λ . For $-1 < \lambda < 0$, there are no additional conditions imposed on the function $f(x)$.

In Chapter 1, conditions 1°–3° are as a rule not specified.

1.1. Equations Whose Kernels Contain Power-Law Functions

1.1-1. Kernels Linear in the Arguments x and t .

$$1. \quad \int_a^x y(t) dt = f(x).$$

Solution: $y(x) = f'_x(x)$.

$$2. \quad \int_a^x (x-t)y(t) dt = f(x).$$

Solution: $y(x) = f''_{xx}(x)$.

$$3. \quad \int_a^x (Ax + Bt + C)y(t) dt = f(x).$$

This is a special case of equation 1.9.5 with $g(x) = x$.

1°. Solution with $B \neq -A$:

$$y(x) = \frac{d}{dx} \left\{ [(A+B)x + C]^{-\frac{A}{A+B}} \int_a^x [(A+B)t + C]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

2°. Solution with $B = -A$:

$$y(x) = \frac{1}{C} \frac{d}{dx} \left[\exp\left(-\frac{A}{C}x\right) \int_a^x \exp\left(\frac{A}{C}t\right) f'_t(t) dt \right].$$

1.1-2. Kernels Quadratic in the Arguments x and t .

$$4. \quad \int_a^x (x-t)^2 y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Solution: $y(x) = \frac{1}{2} f'''_{xxx}(x)$.

$$5. \quad \int_a^x (x^2 - t^2)y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This is a special case of equation 1.9.2 with $g(x) = x^2$.

$$\text{Solution: } y(x) = \frac{1}{2x^2} [x f''_{xx}(x) - f'_x(x)].$$

$$6. \quad \int_a^x (Ax^2 + Bt^2)y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = x^2$. For $B = -A$, see equation 1.1.5.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left[x^{-\frac{2A}{A+B}} \int_a^x t^{-\frac{2B}{A+B}} f'_t(t) dt \right].$$

$$7. \quad \int_a^x (Ax^2 + Bt^2 + C)y(t) dt = f(x).$$

This is a special case of equation 1.9.5 with $g(x) = x^2$.

Solution:

$$y(x) = \text{sign } \varphi(x) \frac{d}{dx} \left\{ |\varphi(x)|^{-\frac{A}{A+B}} \int_a^x |\varphi(t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}, \quad \varphi(x) = (A+B)x^2 + C.$$

$$8. \int_a^x [Ax^2 + (B - A)xt - Bt^2]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Differentiating with respect to x yields an equation of the form 1.1.3:

$$\int_a^x [2Ax + (B - A)t]y(t) dt = f'_x(x).$$

Solution:

$$y(x) = \frac{1}{A + B} \frac{d}{dx} \left[x^{-\frac{2A}{A+B}} \int_a^x t^{\frac{A-B}{A+B}} f''_{tt}(t) dt \right].$$

$$9. \int_a^x (Ax^2 + Bt^2 + Cx + Dt + E)y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^2 + Cx$ and $h(t) = Bt^2 + Dt + E$.

$$10. \int_a^x (Axt + Bt^2 + Cx + Dt + E)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = x$, $h_1(t) = At + C$, $g_2(x) = 1$, and $h_2(t) = Bt^2 + Dt + E$.

$$11. \int_a^x (Ax^2 + Bxt + Cx + Dt + E)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Bx + D$, $h_1(t) = t$, $g_2(x) = Ax^2 + Cx + E$, and $h_2(t) = 1$.

1.1-3. Kernels Cubic in the Arguments x and t .

$$12. \int_a^x (x - t)^3 y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0.$$

Solution: $y(x) = \frac{1}{6} f'''_{xxx}(x)$.

$$13. \int_a^x (x^3 - t^3)y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This is a special case of equation 1.9.2 with $g(x) = x^3$.

Solution: $y(x) = \frac{1}{3x^3} [x f'''_{xxx}(x) - 2f'_x(x)]$.

$$14. \int_a^x (Ax^3 + Bt^3)y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = x^3$. For $B = -A$, see equation 1.1.13.

Solution with $0 \leq a \leq x$: $y(x) = \frac{1}{A + B} \frac{d}{dx} \left[x^{-\frac{3A}{A+B}} \int_a^x t^{-\frac{3B}{A+B}} f'_t(t) dt \right]$.

$$15. \int_a^x (Ax^3 + Bt^3 + C)y(t) dt = f(x).$$

This is a special case of equation 1.9.5 with $g(x) = x^3$.

$$16. \int_a^x (x^2t - xt^2)y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This is a special case of equation 1.9.11 with $g(x) = x^2$ and $h(x) = x$.

$$\text{Solution: } y(x) = \frac{1}{x} \frac{d^2}{dx^2} \left[\frac{1}{x} f(x) \right].$$

$$17. \int_a^x (Ax^2t + Bxt^2)y(t) dt = f(x).$$

This is a special case of equation 1.9.12 with $g(x) = x^2$ and $h(x) = x$. For $B = -A$, see equation 1.1.16.

Solution:

$$y(x) = \frac{1}{(A+B)x} \frac{d}{dx} \left\{ x^{-\frac{A}{A+B}} \int_a^x t^{-\frac{B}{A+B}} \frac{d}{dt} \left[\frac{1}{t} f(t) \right] dt \right\}.$$

$$18. \int_a^x (Ax^3 + Bxt^2)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^3$, $h_1(t) = 1$, $g_2(x) = Bx$, and $h_2(t) = t^2$.

$$19. \int_a^x (Ax^3 + Bx^2t)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^3$, $h_1(t) = 1$, $g_2(x) = Bx^2$, and $h_2(t) = t$.

$$20. \int_a^x (Ax^2t + Bt^3)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^2$, $h_1(t) = t$, $g_2(x) = B$, and $h_2(t) = t^3$.

$$21. \int_a^x (Axt^2 + Bt^3)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax$, $h_1(t) = t^2$, $g_2(x) = B$, and $h_2(t) = t^3$.

$$22. \int_a^x (A_3x^3 + B_3t^3 + A_2x^2 + B_2t^2 + A_1x + B_1t + C)y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A_3x^3 + A_2x^2 + A_1x + C$ and $h(t) = B_3t^3 + B_2t^2 + B_1t$.

1.1-4. Kernels Containing Higher-Order Polynomials in x and t .

$$23. \int_a^x (x-t)^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

It is assumed that the right-hand of the equation satisfies the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{1}{n!} f_x^{(n+1)}(x).$$

Example. For $f(x) = Ax^m$, where m is a positive integer, $m > n$, the solution has the form

$$y(x) = \frac{Am!}{n!(m-n-1)!} x^{m-n-1}.$$

$$24. \int_a^x (x^n - t^n)y(t) dt = f(x), \quad f(a) = f'_x(a) = 0, \quad n = 1, 2, \dots$$

$$\text{Solution: } y(x) = \frac{1}{n} \frac{d}{dx} \left[\frac{f'_x(x)}{x^{n-1}} \right].$$

$$25. \int_a^x (t^n x^{n+1} - x^n t^{n+1})y(t) dt = f(x), \quad n = 2, 3, \dots$$

This is a special case of equation 1.9.11 with $g(x) = x^{n+1}$ and $h(x) = x^n$.

$$\text{Solution: } y(x) = \frac{1}{x^n} \frac{d^2}{dx^2} \left[\frac{f(x)}{x^n} \right].$$

1.1-5. Kernels Containing Rational Functions.

$$26. \int_0^x \frac{y(t) dt}{x+t} = f(x).$$

1°. For a polynomial right-hand side, $f(x) = \sum_{n=0}^N A_n x^n$, the solution has the form

$$y(x) = \sum_{n=0}^N \frac{A_n}{B_n} x^n, \quad B_n = (-1)^n \left[\ln 2 + \sum_{k=1}^n \frac{(-1)^k}{k} \right].$$

2°. For $f(x) = x^\lambda \sum_{n=0}^N A_n x^n$, where λ is an arbitrary number ($\lambda > -1$), the solution has the form

$$y(x) = x^\lambda \sum_{n=0}^N \frac{A_n}{B_n} x^n, \quad B_n = \int_0^1 \frac{t^{\lambda+n} dt}{1+t}.$$

3°. For $f(x) = \ln x \left(\sum_{n=0}^N A_n x^n \right)$, the solution has the form

$$y(x) = \ln x \sum_{n=0}^N \frac{A_n}{B_n} x^n + \sum_{n=0}^N \frac{A_n I_n}{B_n^2} x^n,$$

$$B_n = (-1)^n \left[\ln 2 + \sum_{k=1}^n \frac{(-1)^k}{k} \right], \quad I_n = (-1)^n \left[\frac{\pi^2}{12} + \sum_{k=1}^n \frac{(-1)^k}{k^2} \right].$$

4°. For $f(x) = \sum_{n=0}^N A_n (\ln x)^n$, the solution of the equation has the form

$$y(x) = \sum_{n=0}^N A_n Y_n(x),$$

where the functions $Y_n = Y_n(x)$ are given by

$$Y_n(x) = \left\{ \frac{d^n}{d\lambda^n} \left[\frac{x^\lambda}{I(\lambda)} \right] \right\}_{\lambda=0}, \quad I(\lambda) = \int_0^1 \frac{z^\lambda dz}{1+z}.$$

5°. For $f(x) = \sum_{n=1}^N A_n \cos(\lambda_n \ln x) + \sum_{n=1}^N B_n \sin(\lambda_n \ln x)$, the solution of the equation has the form

$$y(x) = \sum_{n=1}^N C_n \cos(\lambda_n \ln x) + \sum_{n=1}^N D_n \sin(\lambda_n \ln x),$$

where the constants C_n and D_n are found by the method of undetermined coefficients.

6°. For arbitrary $f(x)$, the transformation

$$x = \frac{1}{2}e^{2z}, \quad t = \frac{1}{2}e^{2\tau}, \quad y(t) = e^{-\tau} w(\tau), \quad f(x) = e^{-z} g(z)$$

leads to an integral equation with difference kernel of the form 1.9.27:

$$\int_{-\infty}^z \frac{w(\tau) d\tau}{\cosh(z - \tau)} = g(z).$$

27. $\int_0^x \frac{y(t) dt}{ax + bt} = f(x), \quad a > 0, \quad a + b > 0.$

1°. For a polynomial right-hand side, $f(x) = \sum_{n=0}^N A_n x^n$, the solution has the form

$$y(x) = \sum_{n=0}^N \frac{A_n}{B_n} x^n, \quad B_n = \int_0^1 \frac{t^n dt}{a + bt}.$$

2°. For $f(x) = x^\lambda \sum_{n=0}^N A_n x^n$, where λ is an arbitrary number ($\lambda > -1$), the solution has the form

$$y(x) = x^\lambda \sum_{n=0}^N \frac{A_n}{B_n} x^n, \quad B_n = \int_0^1 \frac{t^{\lambda+n} dt}{a + bt}.$$

3°. For $f(x) = \ln x \left(\sum_{n=0}^N A_n x^n \right)$, the solution has the form

$$y(x) = \ln x \sum_{n=0}^N \frac{A_n}{B_n} x^n - \sum_{n=0}^N \frac{A_n C_n}{B_n^2} x^n, \quad B_n = \int_0^1 \frac{t^n dt}{a + bt}, \quad C_n = \int_0^1 \frac{t^n \ln t}{a + bt} dt.$$

4°. For some other special forms of the right-hand side (see items 4 and 5, equation 1.1.26), the solution may be found by the method of undetermined coefficients.

28. $\int_0^x \frac{y(t) dt}{ax^2 + bt^2} = f(x), \quad a > 0, \quad a + b > 0.$

1°. For a polynomial right-hand side, $f(x) = \sum_{n=0}^N A_n x^n$, the solution has the form

$$y(x) = \sum_{n=0}^N \frac{A_n}{B_n} x^{n+1}, \quad B_n = \int_0^1 \frac{t^{n+1} dt}{a + bt^2}.$$

Example. For $a = b = 1$ and $f(x) = Ax^2 + Bx + C$, the solution of the integral equation is:

$$y(x) = \frac{2A}{1 - \ln 2}x^3 + \frac{4B}{4 - \pi}x^2 + \frac{2C}{\ln 2}x.$$

2°. For $f(x) = x^\lambda \sum_{n=0}^N A_n x^n$, where λ is an arbitrary number ($\lambda > -1$), the solution has the form

$$y(x) = x^\lambda \sum_{n=0}^N \frac{A_n}{B_n} x^{n+1}, \quad B_n = \int_0^1 \frac{t^{\lambda+n+1} dt}{a + bt^2}.$$

3°. For $f(x) = \ln x \left(\sum_{n=0}^N A_n x^n \right)$, the solution has the form

$$y(x) = \ln x \sum_{n=0}^N \frac{A_n}{B_n} x^{n+1} - \sum_{n=0}^N \frac{A_n C_n}{B_n^2} x^{n+1}, \quad B_n = \int_0^1 \frac{t^{n+1} dt}{a + bt^2}, \quad C_n = \int_0^1 \frac{t^{n+1} \ln t}{a + bt^2} dt.$$

29.
$$\int_0^x \frac{y(t) dt}{ax^m + bt^m} = f(x), \quad a > 0, \quad a + b > 0, \quad m = 1, 2, \dots$$

1°. For a polynomial right-hand side, $f(x) = \sum_{n=0}^N A_n x^n$, the solution has the form

$$y(x) = \sum_{n=0}^N \frac{A_n}{B_n} x^{m+n-1}, \quad B_n = \int_0^1 \frac{t^{m+n-1} dt}{a + bt^m}.$$

2°. For $f(x) = x^\lambda \sum_{n=0}^N A_n x^n$, where λ is an arbitrary number ($\lambda > -1$), the solution has the form

$$y(x) = x^\lambda \sum_{n=0}^N \frac{A_n}{B_n} x^{m+n-1}, \quad B_n = \int_0^1 \frac{t^{\lambda+m+n-1} dt}{a + bt^m}.$$

3°. For $f(x) = \ln x \left(\sum_{n=0}^N A_n x^n \right)$, the solution has the form

$$y(x) = \ln x \sum_{n=0}^N \frac{A_n}{B_n} x^{m+n-1} - \sum_{n=0}^N \frac{A_n C_n}{B_n^2} x^{m+n-1},$$

$$B_n = \int_0^1 \frac{t^{m+n-1} dt}{a + bt^m}, \quad C_n = \int_0^1 \frac{t^{m+n-1} \ln t}{a + bt^m} dt.$$

1.1-6. Kernels Containing Square Roots.

30.
$$\int_a^x \sqrt{x-t} y(t) dt = f(x).$$

Differentiating with respect to x , we arrive at Abel's equation 1.1.36:

$$\int_a^x \frac{y(t) dt}{\sqrt{x-t}} = 2f'_x(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \frac{d^2}{dx^2} \int_a^x \frac{f(t) dt}{\sqrt{x-t}}.$$

$$31. \int_a^x (\sqrt{x} - \sqrt{t}) y(t) dt = f(x).$$

This is a special case of equation 1.1.45 with $\mu = \frac{1}{2}$.

$$\text{Solution: } y(x) = 2 \frac{d}{dx} [\sqrt{x} f'_x(x)].$$

$$32. \int_a^x (A\sqrt{x} + B\sqrt{t}) y(t) dt = f(x).$$

This is a special case of equation 1.1.46 with $\mu = \frac{1}{2}$.

$$33. \int_a^x (1 + b\sqrt{x-t}) y(t) dt = f(x).$$

Differentiating with respect to x , we arrive at Abel's equation of the second kind 2.1.46:

$$y(x) + \frac{b}{2} \int_a^x \frac{y(t) dt}{\sqrt{x-t}} = f'_x(x).$$

$$34. \int_a^x (t\sqrt{x} - x\sqrt{t}) y(t) dt = f(x).$$

This is a special case of equation 1.9.11 with $g(x) = \sqrt{x}$ and $h(x) = x$.

$$35. \int_a^x (At\sqrt{x} + Bx\sqrt{t}) y(t) dt = f(x).$$

This is a special case of equation 1.9.12 with $g(x) = \sqrt{x}$ and $h(t) = t$.

$$36. \int_a^x \frac{y(t) dt}{\sqrt{x-t}} = f(x).$$

Abel's equation.

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{\sqrt{x-t}} = \frac{f(a)}{\pi\sqrt{x-a}} + \frac{1}{\pi} \int_a^x \frac{f'_t(t) dt}{\sqrt{x-t}}.$$

⊙ Reference: E. T. Whittaker and G. N. Watson (1958).

$$37. \int_a^x \left(b + \frac{1}{\sqrt{x-t}} \right) y(t) dt = f(x).$$

Let us rewrite the equation in the form

$$\int_a^x \frac{y(t) dt}{\sqrt{x-t}} = f(x) - b \int_a^x y(t) dt.$$

Assuming the right-hand side to be known, we solve this equation as Abel's equation 1.1.36. After some manipulations, we arrive at Abel's equation of the second kind 2.1.46:

$$y(x) + \frac{b}{\pi} \int_a^x \frac{y(t) dt}{\sqrt{x-t}} = F(x), \quad \text{where } F(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{\sqrt{x-t}}.$$

$$38. \int_a^x \left(\frac{1}{\sqrt{x}} - \frac{1}{\sqrt{t}} \right) y(t) dt = f(x).$$

This is a special case of equation 1.1.45 with $\mu = -\frac{1}{2}$.

$$\text{Solution: } y(x) = -2 \left[x^{3/2} f'_x(x) \right]'_x, \quad a > 0.$$

$$39. \int_a^x \left(\frac{A}{\sqrt{x}} + \frac{B}{\sqrt{t}} \right) y(t) dt = f(x).$$

This is a special case of equation 1.1.46 with $\mu = -\frac{1}{2}$.

$$40. \int_{-x}^x \sqrt{\frac{x-t}{x+t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{\text{sign } x}{2\pi} \left[\frac{d}{dx} \int_0^{|x|} \frac{f(t) - f(-t)}{\sqrt{x^2 - t^2}} dt - \frac{1}{x} \frac{d}{dx} \int_0^{|x|} \frac{t[f(t) - f(-t)]}{\sqrt{x^2 - t^2}} dt \right].$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992).

$$41. \int_a^x \frac{y(t) dt}{\sqrt{x^2 - t^2}} = f(x).$$

$$\text{Solution: } y = \frac{2}{\pi} \frac{d}{dx} \int_a^x \frac{t f(t) dt}{\sqrt{x^2 - t^2}}.$$

⊙ Reference: P. P. Zabreyko, A. I. Koshelev, et al. (1975).

$$42. \int_0^x \frac{y(t) dt}{\sqrt{ax^2 + bt^2}} = f(x), \quad a > 0, \quad a + b > 0.$$

1°. For a polynomial right-hand side, $f(x) = \sum_{n=0}^N A_n x^n$, the solution has the form

$$y(x) = \sum_{n=0}^N \frac{A_n}{B_n} x^n, \quad B_n = \int_0^1 \frac{t^n dt}{\sqrt{a + bt^2}}.$$

2°. For $f(x) = x^\lambda \sum_{n=0}^N A_n x^n$, where λ is an arbitrary number ($\lambda > -1$), the solution has the form

$$y(x) = x^\lambda \sum_{n=0}^N \frac{A_n}{B_n} x^n, \quad B_n = \int_0^1 \frac{t^{\lambda+n} dt}{\sqrt{a + bt^2}}.$$

3°. For $f(x) = \ln x \left(\sum_{n=0}^N A_n x^n \right)$, the solution has the form

$$y(x) = \ln x \sum_{n=0}^N \frac{A_n}{B_n} x^n - \sum_{n=0}^N \frac{A_n C_n}{B_n^2} x^n, \quad B_n = \int_0^1 \frac{t^n dt}{\sqrt{a + bt^2}}, \quad C_n = \int_0^1 \frac{t^n \ln t}{\sqrt{a + bt^2}} dt.$$

4°. For $f(x) = \sum_{n=0}^N A_n (\ln x)^n$, the solution of the equation has the form

$$y(x) = \sum_{n=0}^N A_n Y_n(x),$$

where the functions $Y_n = Y_n(x)$ are given by

$$Y_n(x) = \left\{ \frac{d^n}{d\lambda^n} \left[\frac{x^\lambda}{I(\lambda)} \right] \right\}_{\lambda=0}, \quad I(\lambda) = \int_0^1 \frac{z^\lambda dz}{\sqrt{a+bz^2}}.$$

5°. For $f(x) = \sum_{n=1}^N A_n \cos(\lambda_n \ln x) + \sum_{n=1}^N B_n \sin(\lambda_n \ln x)$, the solution of the equation has the form

$$y(x) = \sum_{n=1}^N C_n \cos(\lambda_n \ln x) + \sum_{n=1}^N D_n \sin(\lambda_n \ln x),$$

where the constants C_n and D_n are found by the method of undetermined coefficients.

1.1-7. Kernels Containing Arbitrary Powers.

43. $\int_a^x (x-t)^\lambda y(t) dt = f(x), \quad f(a) = 0, \quad 0 < \lambda < 1.$

Differentiating with respect to x , we arrive at the generalized Abel equation 1.1.47:

$$\int_a^x \frac{y(t) dt}{(x-t)^{1-\lambda}} = \frac{1}{\lambda} f'_x(x).$$

Solution:

$$y(x) = k \frac{d^2}{dx^2} \int_a^x \frac{f(t) dt}{(x-t)^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

⊙ Reference: F. D. Gakhov (1977).

44. $\int_a^x (x-t)^\mu y(t) dt = f(x).$

For $\mu = 0, 1, 2, \dots$, see equations 1.1.1, 1.1.2, 1.1.4, 1.1.12, and 1.1.23. For $0 < \mu < 1$, see equation 1.1.43.

Set $\mu = n - \lambda$, where $n = 1, 2, \dots$ and $0 \leq \lambda < 1$, and $f(a) = f'_x(a) = \dots = f_x^{(n-1)}(a) = 0$.

On differentiating the equation n times, we arrive at an equation of the form 1.1.47:

$$\int_a^x \frac{y(t) d\tau}{(x-t)^\lambda} = \frac{\Gamma(\mu - n + 1)}{\Gamma(\mu + 1)} f_x^{(n)}(x),$$

where $\Gamma(\mu)$ is the gamma function.

Example. Set $f(x) = Ax^\beta$, where $\beta \geq 0$, and let $\mu > -1$ and $\mu - \beta \neq 0, 1, 2, \dots$. In this case, the solution has the form $y(x) = \frac{A\Gamma(\beta+1)}{\Gamma(\mu+1)\Gamma(\beta-\mu)} x^{\beta-\mu-1}$.

⊙ Reference: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971).

45. $\int_a^x (x^\mu - t^\mu) y(t) dt = f(x).$

This is a special case of equation 1.9.2 with $g(x) = x^\mu$.

Solution: $y(x) = \frac{1}{\mu} [x^{1-\mu} f'_x(x)]'_x.$

$$46. \int_a^x (Ax^\mu + Bt^\mu)y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = x^\mu$. For $B = -A$, see equation 1.1.44.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left[x^{-\frac{A\mu}{A+B}} \int_a^x t^{-\frac{B\mu}{A+B}} f'_t(t) dt \right].$$

$$47. \int_a^x \frac{y(t) dt}{(x-t)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

The generalized Abel equation.

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{(x-t)^{1-\lambda}} = \frac{\sin(\pi\lambda)}{\pi} \left[\frac{f(a)}{(x-a)^{1-\lambda}} + \int_a^x \frac{f'_t(t) dt}{(x-t)^{1-\lambda}} \right].$$

⊙ Reference: E. T. Whittaker and G. N. Watson (1958).

$$48. \int_a^x \left[b + \frac{1}{(x-t)^\lambda} \right] y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Rewrite the equation in the form

$$\int_a^x \frac{y(t) dt}{(x-t)^\lambda} = f(x) - b \int_a^x y(t) dt.$$

Assuming the right-hand side to be known, we solve this equation as the generalized Abel equation 1.1.47. After some manipulations, we arrive at Abel's equation of the second kind 2.1.60:

$$y(x) + \frac{b \sin(\pi\lambda)}{\pi} \int_a^x \frac{y(t) dt}{(x-t)^{1-\lambda}} = F(x), \quad \text{where } F(x) = \frac{\sin(\pi\lambda)}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{(x-t)^{1-\lambda}}.$$

$$49. \int_a^x (\sqrt{x} - \sqrt{t})^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{k}{\sqrt{x}} \left(\sqrt{x} \frac{d}{dx} \right)^2 \int_a^x \frac{f(t) dt}{\sqrt{t} (\sqrt{x} - \sqrt{t})^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$50. \int_a^x \frac{y(t) dt}{(\sqrt{x} - \sqrt{t})^\lambda} = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{2\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{\sqrt{t} (\sqrt{x} - \sqrt{t})^{1-\lambda}}.$$

$$51. \int_a^x (Ax^\lambda + Bt^\mu)y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\lambda$ and $h(t) = Bt^\mu$.

$$52. \int_a^x [1 + A(x^\lambda t^\mu - x^{\lambda+\mu})] y(t) dt = f(x).$$

This is a special case of equation 1.9.13 with $g(x) = Ax^\mu$ and $h(x) = x^\lambda$.

Solution:

$$y(x) = \frac{d}{dx} \left\{ \frac{x^\lambda}{\Phi(x)} \int_a^x [t^{-\lambda} f(t)]'_t \Phi(t) dt \right\}, \quad \Phi(x) = \exp\left(-\frac{A\mu}{\mu + \lambda} x^{\mu+\lambda}\right).$$

$$53. \int_a^x (Ax^\beta t^\gamma + Bx^\delta t^\lambda) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\beta$, $h_1(t) = t^\gamma$, $g_2(x) = Bx^\delta$, and $h_2(t) = t^\lambda$.

$$54. \int_a^x [Ax^\lambda(t^\mu - x^\mu) + Bx^\beta(t^\gamma - x^\gamma)] y(t) dt = f(x).$$

This is a special case of equation 1.9.47 with $g_1(x) = Ax^\lambda$, $h_1(x) = x^\mu$, $g_2(x) = Bx^\beta$, and $h_2(x) = x^\gamma$.

$$55. \int_a^x [Ax^\lambda t^\mu + Bx^{\lambda+\beta} t^{\mu-\beta} - (A+B)x^{\lambda+\gamma} t^{\mu-\gamma}] y(t) dt = f(x).$$

This is a special case of equation 1.9.49 with $g(x) = x$.

$$56. \int_a^x t^\sigma (x^\mu - t^\mu)^\lambda y(t) dt = f(x), \quad \sigma > -1, \quad \mu > 0, \quad \lambda > -1.$$

The transformation $\tau = t^\mu$, $z = x^\mu$, $w(\tau) = t^{\sigma-\mu+1} y(t)$ leads to an equation of the form 1.1.43:

$$\int_A^z (z - \tau)^\lambda w(\tau) d\tau = F(z),$$

where $A = a^\mu$ and $F(z) = \mu f(z^{1/\mu})$.

Solution with $-1 < \lambda < 0$:

$$y(x) = -\frac{\mu \sin(\pi\lambda)}{\pi x^\sigma} \frac{d}{dx} \left[\int_a^x t^{\mu-1} (x^\mu - t^\mu)^{-1-\lambda} f(t) dt \right].$$

$$57. \int_0^x \frac{y(t) dt}{(x+t)^\mu} = f(x).$$

This is a special case of equation 1.1.58 with $\lambda = 1$ and $a = b = 1$.

The transformation

$$x = \frac{1}{2} e^{2z}, \quad t = \frac{1}{2} e^{2\tau}, \quad y(t) = e^{(\mu-2)\tau} w(\tau), \quad f(x) = e^{-\mu z} g(z)$$

leads to an equation with difference kernel of the form 1.9.27:

$$\int_{-\infty}^z \frac{w(\tau) d\tau}{\cosh^\mu(z - \tau)} = g(z).$$

$$58. \int_0^x \frac{y(t) dt}{(ax^\lambda + bt^\lambda)^\mu} = f(x), \quad a > 0, \quad a + b > 0.$$

1°. The substitution $t = xz$ leads to a special case of equation 3.8.45:

$$\int_0^1 \frac{y(xz) dz}{(a + bz^\lambda)^\mu} = x^{\lambda\mu-1} f(x). \quad (1)$$

2°. For a polynomial right-hand side, $f(x) = \sum_{m=0}^n A_m x^m$, the solution has the form

$$y(x) = x^{\lambda\mu-1} \sum_{m=0}^n \frac{A_m}{I_m} x^m, \quad I_m = \int_0^1 \frac{z^{m+\lambda\mu-1} dz}{(a + bz^\lambda)^\mu}.$$

The integrals I_m are supposed to be convergent.

3°. The solution structure for some other right-hand sides of the integral equation may be obtained using (1) and the results presented for the more general equation 3.8.53 (see also equations 3.8.34–3.8.40).

4°. For $a = b$, the equation can be reduced, just as equation 1.1.57, to an integral equation with difference kernel of the form 1.9.27.

$$59. \int_a^x \frac{(\sqrt{x} + \sqrt{x-t})^{2\lambda} + (\sqrt{x} - \sqrt{x-t})^{2\lambda}}{2t^\lambda \sqrt{x-t}} y(t) dt = f(x).$$

The equation can be rewritten in terms of the Gaussian hypergeometric functions in the form

$$\int_a^x (x-t)^{\gamma-1} F\left(\lambda, -\lambda, \gamma; 1 - \frac{x}{t}\right) y(t) dt = f(x), \quad \text{where } \gamma = \frac{1}{2}.$$

See 1.8.135 for the solution of this equation.

1.1-8. Two-Dimensional Equation of the Abel Type.

$$60. \iint_{\Delta} \frac{u(x, y) dx dy}{\sqrt{(y_0 - y)^2 - (x_0 - x)^2}} = f(x_0, y_0).$$

Here Δ is an isosceles right triangle with apex at the point (x_0, y_0) and base on the x -axis.

Solution:

$$u(x_0, y_0) = \frac{1}{2\pi^2} \left(\frac{\partial^2 g}{\partial x_0^2} - \frac{\partial^2 g}{\partial y_0^2} \right), \quad g(x_0, y_0) = \iint_{\Delta} \frac{f(x, y) dx dy}{\sqrt{(y_0 - y)^2 - (x_0 - x)^2}}.$$

⊙ Reference: P. P. Zabreyko, A. I. Koshelev, et al. (1975).

1.2. Equations Whose Kernels Contain Exponential Functions

1.2-1. Kernels Containing Exponential Functions.

$$1. \int_a^x e^{\lambda(x-t)} y(t) dt = f(x).$$

Solution: $y(x) = f'_x(x) - \lambda f(x)$.

Example. In the special case $a = 0$ and $f(x) = Ax$, the solution has the form $y(x) = A(1 - \lambda x)$.

$$2. \quad \int_a^x e^{\lambda x + \beta t} y(t) dt = f(x).$$

$$\text{Solution: } y(x) = e^{-(\lambda + \beta)x} [f'_x(x) - \lambda f(x)].$$

Example. In the special case $a = 0$ and $f(x) = A \sin(\gamma x)$, the solution has the form $y(x) = Ae^{-(\lambda + \beta)x} \times [\gamma \cos(\gamma x) - \lambda \sin(\gamma x)]$.

$$3. \quad \int_a^x [e^{\lambda(x-t)} - 1] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = \frac{1}{\lambda} f''_{xx}(x) - f'_x(x).$$

$$4. \quad \int_a^x [e^{\lambda(x-t)} + b] y(t) dt = f(x).$$

For $b = -1$, see equation 1.2.3. Differentiating with respect to x yields an equation of the form 2.2.1:

$$y(x) + \frac{\lambda}{b+1} \int_a^x e^{\lambda(x-t)} y(t) dt = \frac{f'_x(x)}{b+1}.$$

Solution:

$$y(x) = \frac{f'_x(x)}{b+1} - \frac{\lambda}{(b+1)^2} \int_a^x \exp\left[\frac{\lambda b}{b+1}(x-t)\right] f'_t(t) dt.$$

$$5. \quad \int_a^x (e^{\lambda x + \beta t} + b) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = e^{\lambda x}$, $h_1(t) = e^{\beta t}$, $g_2(x) = 1$, and $h_2(t) = b$. For $\beta = -\lambda$, see equation 1.2.4.

$$6. \quad \int_a^x (e^{\lambda x} - e^{\lambda t}) y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This is a special case of equation 1.9.2 with $g(x) = e^{\lambda x}$.

$$\text{Solution: } y(x) = e^{-\lambda x} \left[\frac{1}{\lambda} f''_{xx}(x) - f'_x(x) \right].$$

$$7. \quad \int_a^x (e^{\lambda x} - e^{\lambda t} + b) y(t) dt = f(x).$$

This is a special case of equation 1.9.3 with $g(x) = e^{\lambda x}$. For $b = 0$, see equation 1.2.6.

Solution:

$$y(x) = \frac{1}{b} f'_x(x) - \frac{\lambda}{b^2} e^{\lambda x} \int_a^x \exp\left(\frac{e^{\lambda t} - e^{\lambda x}}{b}\right) f'_t(t) dt.$$

$$8. \quad \int_a^x (Ae^{\lambda x} + Be^{\lambda t}) y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = e^{\lambda x}$. For $B = -A$, see equation 1.2.6.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left[\exp\left(-\frac{A\lambda}{A+B}x\right) \int_a^x \exp\left(-\frac{B\lambda}{A+B}t\right) f'_t(t) dt \right].$$

$$9. \quad \int_a^x (Ae^{\lambda x} + Be^{\lambda t} + C) y(t) dt = f(x).$$

This is a special case of equation 1.9.5 with $g(x) = e^{\lambda x}$.

$$10. \int_a^x (Ae^{\lambda x} + Be^{\mu t})y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ae^{\lambda x}$ and $h(t) = Be^{\mu t}$. For $\lambda = \mu$, see equation 1.2.8.

$$11. \int_a^x [e^{\lambda(x-t)} - e^{\mu(x-t)}]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{1}{\lambda - \mu} [f''_{xx} - (\lambda + \mu)f'_x + \lambda\mu f], \quad f = f(x).$$

$$12. \int_a^x [Ae^{\lambda(x-t)} + Be^{\mu(x-t)}]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\lambda x}$, $h_1(t) = e^{-\lambda t}$, $g_2(x) = Be^{\mu x}$, and $h_2(t) = e^{-\mu t}$. For $B = -A$, see equation 1.2.11.

Solution:

$$y(x) = \frac{e^{\lambda x}}{A+B} \frac{d}{dx} \left\{ e^{(\mu-\lambda)x} \Phi(x) \int_a^x \left[\frac{f(t)}{e^{\mu t}} \right]'_t \frac{dt}{\Phi(t)} \right\}, \quad \Phi(x) = \exp \left[\frac{B(\lambda - \mu)}{A+B} x \right].$$

$$13. \int_a^x [Ae^{\lambda(x-t)} + Be^{\mu(x-t)} + C]y(t) dt = f(x).$$

This is a special case of equation 1.2.14 with $\beta = 0$.

$$14. \int_a^x [Ae^{\lambda(x-t)} + Be^{\mu(x-t)} + Ce^{\beta(x-t)}]y(t) dt = f(x).$$

Differentiating the equation with respect to x yields

$$(A+B+C)y(x) + \int_a^x [A\lambda e^{\lambda(x-t)} + B\mu e^{\mu(x-t)} + C\beta e^{\beta(x-t)}]y(t) dt = f'_x(x).$$

Eliminating the term with $e^{\beta(x-t)}$ with the aid of the original equation, we arrive at an equation of the form 2.2.10:

$$(A+B+C)y(x) + \int_a^x [A(\lambda - \beta)e^{\lambda(x-t)} + B(\mu - \beta)e^{\mu(x-t)}]y(t) dt = f'_x(x) - \beta f(x).$$

In the special case $A+B+C=0$, this is an equation of the form 1.2.12.

$$15. \int_a^x [Ae^{\lambda(x-t)} + Be^{\mu(x-t)} + Ce^{\beta(x-t)} - A - B - C]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Differentiating with respect to x , we arrive at an equation of the form 1.2.14:

$$\int_a^x [A\lambda e^{\lambda(x-t)} + B\mu e^{\mu(x-t)} + C\beta e^{\beta(x-t)}]y(t) dt = f'_x(x).$$

$$16. \int_a^x (e^{\lambda x + \mu t} - e^{\mu x + \lambda t})y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This is a special case of equation 1.9.11 with $g(x) = e^{\lambda x}$ and $h(t) = e^{\mu t}$.

Solution:

$$y(x) = \frac{f''_{xx} - (\lambda + \mu)f'_x(x) + \lambda\mu f(x)}{(\lambda - \mu) \exp[(\lambda + \mu)x]}.$$

$$17. \int_a^x (Ae^{\lambda x + \mu t} + Be^{\mu x + \lambda t})y(t) dt = f(x).$$

This is a special case of equation 1.9.12 with $g(x) = e^{\lambda x}$ and $h(t) = e^{\mu t}$. For $B = -A$, see equation 1.2.16.

Solution:

$$y(x) = \frac{1}{(A+B)e^{\mu x}} \frac{d}{dx} \left\{ \Phi^A(x) \int_a^x \Phi^B(t) \frac{d}{dt} \left[\frac{f(t)}{e^{\mu t}} \right] dt \right\}, \quad \Phi(x) = \exp\left(\frac{\mu - \lambda}{A+B}x\right).$$

$$18. \int_a^x (Ae^{\lambda x + \mu t} + Be^{\beta x + \gamma t})y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\lambda x}$, $h_1(t) = e^{\mu t}$, $g_2(x) = Be^{\beta x}$, and $h_2(t) = e^{\gamma t}$.

$$19. \int_a^x (Ae^{2\lambda x} + Be^{2\beta t} + Ce^{\lambda x} + De^{\beta t} + E)y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ae^{2\lambda x} + Ce^{\lambda x}$ and $h(t) = Be^{2\beta t} + De^{\beta t} + E$.

$$20. \int_a^x (Ae^{\lambda x + \beta t} + Be^{2\beta t} + Ce^{\lambda x} + De^{\beta t} + E)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = e^{\lambda x}$, $h_1(t) = Ae^{\beta t} + C$, and $g_2(x) = 1$, $h_2(t) = Be^{2\beta t} + De^{\beta t} + E$.

$$21. \int_a^x (Ae^{2\lambda x} + Be^{\lambda x + \beta t} + Ce^{\lambda x} + De^{\beta t} + E)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Be^{\lambda x} + D$, $h_1(t) = e^{\beta t}$, and $g_2(x) = Ae^{2\lambda x} + Ce^{\lambda x} + E$, $h_2(t) = 1$.

$$22. \int_a^x [1 + Ae^{\lambda x}(e^{\mu t} - e^{\mu x})]y(t) dt = f(x).$$

This is a special case of equation 1.9.13 with $g(x) = e^{\mu x}$ and $h(x) = Ae^{\lambda x}$.

Solution:

$$y(x) = \frac{d}{dx} \left\{ e^{\lambda x} \Phi(x) \int_a^x \left[\frac{f(t)}{e^{\lambda t}} \right]' \frac{dt}{\Phi(t)} \right\}, \quad \Phi(x) = \exp\left[\frac{A\mu}{\lambda + \mu}e^{(\lambda + \mu)x}\right].$$

$$23. \int_a^x [Ae^{\lambda x}(e^{\mu x} - e^{\mu t}) + Be^{\beta x}(e^{\gamma x} - e^{\gamma t})]y(t) dt = f(x).$$

This is a special case of equation 1.9.47 with $g_1(x) = Ae^{\lambda x}$, $h_1(t) = -e^{\mu t}$, $g_2(x) = Be^{\beta x}$, and $h_2(t) = -e^{\gamma t}$.

$$24. \int_a^x \{A \exp(\lambda x + \mu t) + B \exp[(\lambda + \beta)x + (\mu - \beta)t] - (A + B) \exp[(\lambda + \gamma)x + (\mu - \gamma)t]\}y(t) dt = f(x).$$

This is a special case of equation 1.9.49 with $g_1(x) = e^x$.

$$25. \int_a^x (e^{\lambda x} - e^{\lambda t})^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

Solution:

$$y(x) = \frac{1}{\lambda^n n!} e^{\lambda x} \left(\frac{1}{e^{\lambda x}} \frac{d}{dx} \right)^{n+1} f(x).$$

$$26. \int_a^x \sqrt{e^{\lambda x} - e^{\lambda t}} y(t) dt = f(x), \quad \lambda > 0.$$

Solution:

$$y(x) = \frac{2}{\pi} e^{\lambda x} \left(e^{-\lambda x} \frac{d}{dx} \right)^2 \int_a^x \frac{e^{\lambda t} f(t) dt}{\sqrt{e^{\lambda x} - e^{\lambda t}}}.$$

$$27. \int_a^x \frac{y(t) dt}{\sqrt{e^{\lambda x} - e^{\lambda t}}} = f(x), \quad \lambda > 0.$$

Solution:

$$y(x) = \frac{\lambda}{\pi} \frac{d}{dx} \int_a^x \frac{e^{\lambda t} f(t) dt}{\sqrt{e^{\lambda x} - e^{\lambda t}}}.$$

$$28. \int_a^x (e^{\lambda x} - e^{\lambda t})^\mu y(t) dt = f(x), \quad \lambda > 0, \quad 0 < \mu < 1.$$

Solution:

$$y(x) = k e^{\lambda x} \left(e^{-\lambda x} \frac{d}{dx} \right)^2 \int_a^x \frac{e^{\lambda t} f(t) dt}{(e^{\lambda x} - e^{\lambda t})^\mu}, \quad k = \frac{\sin(\pi\mu)}{\pi\mu}.$$

$$29. \int_a^x \frac{y(t) dt}{(e^{\lambda x} - e^{\lambda t})^\mu} = f(x), \quad \lambda > 0, \quad 0 < \mu < 1.$$

Solution:

$$y(x) = \frac{\lambda \sin(\pi\mu)}{\pi} \frac{d}{dx} \int_a^x \frac{e^{\lambda t} f(t) dt}{(e^{\lambda x} - e^{\lambda t})^{1-\mu}}.$$

1.2-2. Kernels Containing Power-Law and Exponential Functions.

$$30. \int_a^x [A(x-t) + B e^{\lambda(x-t)}] y(t) dt = f(x).$$

Differentiating with respect to x , we arrive at an equation of the form 2.2.4:

$$B y(x) + \int_a^x [A + B \lambda e^{\lambda(x-t)}] y(t) dt = f'_x(x).$$

$$31. \int_a^x (x-t) e^{\lambda(x-t)} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution: $y(x) = f''_{xx}(x) - 2\lambda f'_x(x) + \lambda^2 f(x)$.

$$32. \int_a^x (Ax + Bt + C) e^{\lambda(x-t)} y(t) dt = f(x).$$

The substitution $u(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 1.1.3:

$$\int_a^x (Ax + Bt + C) u(t) dt = e^{-\lambda x} f(x).$$

$$33. \int_a^x (Axe^{\lambda t} + Bte^{\mu x})y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax$, $h_1(t) = e^{\lambda t}$, and $g_2(x) = Be^{\mu x}$, $h_2(t) = t$.

$$34. \int_a^x [Axe^{\lambda(x-t)} + Bte^{\mu(x-t)}]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Axe^{\lambda x}$, $h_1(t) = e^{-\lambda t}$, $g_2(x) = Be^{\mu x}$, and $h_2(t) = te^{-\mu t}$.

$$35. \int_a^x (x-t)^2 e^{\lambda(x-t)}y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Solution: $y(x) = \frac{1}{2} [f'''_{xxx}(x) - 3\lambda f''_{xx}(x) + 3\lambda^2 f'_x(x) - \lambda^3 f(x)]$.

$$36. \int_a^x (x-t)^n e^{\lambda(x-t)}y(t) dt = f(x), \quad n = 1, 2, \dots$$

It is assumed that $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

Solution: $y(x) = \frac{1}{n!} e^{\lambda x} \frac{d^{n+1}}{dx^{n+1}} [e^{-\lambda x} f(x)]$.

$$37. \int_a^x (Ax^\beta + Be^{\lambda t})y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = Be^{\lambda t}$.

$$38. \int_a^x (Ae^{\lambda x} + Bt^\beta)y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ae^{\lambda x}$ and $h(t) = Bt^\beta$.

$$39. \int_a^x (Ax^\beta e^{\lambda t} + Bt^\gamma e^{\mu x})y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\beta$, $h_1(t) = e^{\lambda t}$, $g_2(x) = Be^{\mu x}$, and $h_2(t) = t^\gamma$.

$$40. \int_a^x e^{\lambda(x-t)} \sqrt{x-t} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} e^{\lambda x} \frac{d^2}{dx^2} \int_a^x \frac{e^{-\lambda t} f(t) dt}{\sqrt{x-t}}.$$

$$41. \int_a^x \frac{e^{\lambda(x-t)}}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} e^{\lambda x} \frac{d}{dx} \int_a^x \frac{e^{-\lambda t} f(t) dt}{\sqrt{x-t}}.$$

$$42. \int_a^x (x-t)^\lambda e^{\mu(x-t)} y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k e^{\mu x} \frac{d^2}{dx^2} \int_a^x \frac{e^{-\mu t} f(t) dt}{(x-t)^\lambda}, \quad k = \frac{\sin(\pi \lambda)}{\pi \lambda}.$$

$$43. \int_a^x \frac{e^{\lambda(x-t)}}{(x-t)^\mu} y(t) dt = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi \mu)}{\pi} e^{\lambda x} \frac{d}{dx} \int_a^x \frac{e^{-\lambda t} f(t)}{(x-t)^{1-\mu}} dt.$$

$$44. \int_a^x (\sqrt{x} - \sqrt{t})^\lambda e^{\mu(x-t)} y(t) dt = f(x), \quad 0 < \lambda < 1.$$

The substitution $u(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.1.49:

$$\int_a^x (\sqrt{x} - \sqrt{t})^\lambda u(t) dt = e^{-\mu x} f(x).$$

$$45. \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{(\sqrt{x} - \sqrt{t})^\lambda} = f(x), \quad 0 < \lambda < 1.$$

The substitution $u(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.1.50:

$$\int_a^x \frac{u(t) dt}{(\sqrt{x} - \sqrt{t})^\lambda} = e^{-\mu x} f(x).$$

$$46. \int_a^x \frac{e^{\lambda(x-t)}}{\sqrt{x^2 - t^2}} y(t) dt = f(x).$$

$$\text{Solution: } y = \frac{2}{\pi} e^{\lambda x} \frac{d}{dx} \int_a^x \frac{t e^{-\lambda t}}{\sqrt{x^2 - t^2}} f(t) dt.$$

$$47. \int_a^x \exp[\lambda(x^2 - t^2)] y(t) dt = f(x).$$

$$\text{Solution: } y(x) = f'_x(x) - 2\lambda x f(x).$$

$$48. \int_a^x [\exp(\lambda x^2) - \exp(\lambda t^2)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \exp(\lambda x^2)$.

$$\text{Solution: } y(x) = \frac{1}{2\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{x \exp(\lambda x^2)} \right].$$

$$49. \int_a^x [A \exp(\lambda x^2) + B \exp(\lambda t^2) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.5 with $g(x) = \exp(\lambda x^2)$.

$$50. \int_a^x [A \exp(\lambda x^2) + B \exp(\mu t^2)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \exp(\lambda x^2)$ and $h(t) = B \exp(\mu t^2)$.

$$51. \int_a^x \sqrt{x-t} \exp[\lambda(x^2 - t^2)] y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \exp(\lambda x^2) \frac{d^2}{dx^2} \int_a^x \frac{\exp(-\lambda t^2)}{\sqrt{x-t}} f(t) dt.$$

$$52. \int_a^x \frac{\exp[\lambda(x^2 - t^2)]}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \exp(\lambda x^2) \frac{d}{dx} \int_a^x \frac{\exp(-\lambda t^2)}{\sqrt{x-t}} f(t) dt.$$

$$53. \int_a^x (x-t)^\lambda \exp[\mu(x^2 - t^2)] y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k \exp(\mu x^2) \frac{d^2}{dx^2} \int_a^x \frac{\exp(-\mu t^2)}{(x-t)^\lambda} f(t) dt, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$54. \int_a^x \exp[\lambda(x^\beta - t^\beta)] y(t) dt = f(x).$$

Solution: $y(x) = f'_x(x) - \lambda\beta x^{\beta-1} f(x)$.

$$55. \int_0^x (-1)^{[(x-t)/b]} y(t) dt = f(x), \quad f(0) = f'_x(0) = 0.$$

Here $b = \text{const}$ and $[A]$ stands for the integer part of the number A .

Solution:

$$y(x) = \frac{1}{2} \int_0^x \left(2 \left[\frac{x-t}{b} \right] + 1 \right) f''_{tt}(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 434).

1.3. Equations Whose Kernels Contain Hyperbolic Functions

1.3-1. Kernels Containing Hyperbolic Cosine.

$$1. \int_a^x \cosh[\lambda(x-t)] y(t) dt = f(x).$$

Solution: $y(x) = f'_x(x) - \lambda^2 \int_a^x f(x) dx$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 435).

$$2. \int_a^x \{ \cosh[\lambda(x-t)] - 1 \} y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Solution: $y(x) = \frac{1}{\lambda^2} f'''_{xxx}(x) - f'_x(x)$.

$$3. \int_a^x \{\cosh[\lambda(x-t)] + b\}y(t) dt = f(x).$$

For $b = 0$, see equation 1.3.1. For $b = -1$, see equation 1.3.2. For $\lambda = 0$, see equation 1.1.1. Differentiating the equation with respect to x , we arrive at an equation of the form 2.3.16:

$$y(x) + \frac{\lambda}{b+1} \int_a^x \sinh[\lambda(x-t)]y(t) dt = \frac{f'_x(x)}{b+1}.$$

1°. Solution with $b(b+1) < 0$:

$$y(x) = \frac{f'_x(x)}{b+1} - \frac{\lambda^2}{k(b+1)^2} \int_a^x \sin[k(x-t)]f'_t(t) dt, \quad \text{where } k = \lambda\sqrt{\frac{-b}{b+1}}.$$

2°. Solution with $b(b+1) > 0$:

$$y(x) = \frac{f'_x(x)}{b+1} - \frac{\lambda^2}{k(b+1)^2} \int_a^x \sinh[k(x-t)]f'_t(t) dt, \quad \text{where } k = \lambda\sqrt{\frac{b}{b+1}}.$$

$$4. \int_a^x \cosh(\lambda x + \beta t)y(t) dt = f(x).$$

For $\beta = -\lambda$, see equation 1.3.1.

Differentiating the equation with respect to x twice, we obtain

$$\cosh[(\lambda+\beta)x]y(x) + \lambda \int_a^x \sinh(\lambda x + \beta t)y(t) dt = f'_x(x), \quad (1)$$

$$\{\cosh[(\lambda+\beta)x]y(x)\}'_x + \lambda \sinh[(\lambda+\beta)x]y(x) + \lambda^2 \int_a^x \cosh(\lambda x + \beta t)y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the first-order linear ordinary differential equation

$$w'_x + \lambda \tanh[(\lambda+\beta)x]w = f''_{xx}(x) - \lambda^2 f(x), \quad w = \cosh[(\lambda+\beta)x]y(x). \quad (3)$$

Setting $x = a$ in (1) yields the initial condition $w(a) = f'_x(a)$. On solving equation (3) with this condition, after some manipulations we obtain the solution of the original integral equation in the form

$$y(x) = \frac{1}{\cosh[(\lambda+\beta)x]} f'_x(x) - \frac{\lambda \sinh[(\lambda+\beta)x]}{\cosh^2[(\lambda+\beta)x]} f(x) + \frac{\lambda\beta}{\cosh^{k+1}[(\lambda+\beta)x]} \int_a^x f(t) \cosh^{k-2}[(\lambda+\beta)t] dt, \quad k = \frac{\lambda}{\lambda+\beta}.$$

$$5. \int_a^x [\cosh(\lambda x) - \cosh(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \cosh(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sinh(\lambda x)} \right].$$

$$6. \int_a^x [A \cosh(\lambda x) + B \cosh(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cosh(\lambda x)$. For $B = -A$, see equation 1.3.5.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\cosh(\lambda x)]^{-\frac{A}{A+B}} \int_a^x [\cosh(\lambda t)]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$7. \int_a^x [A \cosh(\lambda x) + B \cosh(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh(\lambda x)$ and $h(t) = B \cosh(\mu t) + C$.

$$8. \int_a^x \{A_1 \cosh[\lambda_1(x-t)] + A_2 \cosh[\lambda_2(x-t)]\} y(t) dt = f(x).$$

The equation is equivalent to the equation

$$\int_a^x \{B_1 \sinh[\lambda_1(x-t)] + B_2 \sinh[\lambda_2(x-t)]\} y(t) dt = F(x),$$

$$B_1 = \frac{A_1}{\lambda_1}, \quad B_2 = \frac{A_2}{\lambda_2}, \quad F(x) = \int_a^x f(t) dt,$$

of the form 1.3.49. (Differentiating this equation yields the original equation.)

$$9. \int_a^x \cosh^2[\lambda(x-t)] y(t) dt = f(x).$$

Differentiation yields an equation of the form 2.3.16:

$$y(x) + \lambda \int_a^x \sinh[2\lambda(x-t)] y(t) dt = f'_x(x).$$

Solution:

$$y(x) = f'_x(x) - \frac{2\lambda^2}{k} \int_a^x \sinh[k(x-t)] f'_t(t) dt, \quad \text{where } k = \lambda\sqrt{2}.$$

$$10. \int_a^x [\cosh^2(\lambda x) - \cosh^2(\lambda t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sinh(2\lambda x)} \right].$$

$$11. \int_a^x [A \cosh^2(\lambda x) + B \cosh^2(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cosh^2(\lambda x)$. For $B = -A$, see equation 1.3.10.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\cosh(\lambda x)]^{-\frac{2A}{A+B}} \int_a^x [\cosh(\lambda t)]^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$12. \int_a^x [A \cosh^2(\lambda x) + B \cosh^2(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh^2(\lambda x)$, and $h(t) = B \cosh^2(\mu t) + C$.

$$13. \int_a^x \cosh[\lambda(x-t)] \cosh[\lambda(x+t)] y(t) dt = f(x).$$

Using the formula

$$\cosh(\alpha - \beta) \cosh(\alpha + \beta) = \frac{1}{2} [\cosh(2\alpha) + \cosh(2\beta)], \quad \alpha = \lambda x, \quad \beta = \lambda t,$$

we transform the original equation to an equation of the form 1.3.6 with $A = B = 1$:

$$\int_a^x [\cosh(2\lambda x) + \cosh(2\lambda t)] y(t) dt = 2f(x).$$

Solution:

$$y(x) = \frac{d}{dx} \left[\frac{1}{\sqrt{\cosh(2\lambda x)}} \int_a^x \frac{f'_t(t) dt}{\sqrt{\cosh(2\lambda t)}} \right].$$

$$14. \int_a^x [\cosh(\lambda x) \cosh(\mu t) + \cosh(\beta x) \cosh(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \cosh(\lambda x)$, $h_1(t) = \cosh(\mu t)$, $g_2(x) = \cosh(\beta x)$, and $h_2(t) = \cosh(\gamma t)$.

$$15. \int_a^x \cosh^3[\lambda(x-t)] y(t) dt = f(x).$$

Using the formula $\cosh^3 \beta = \frac{1}{4} \cosh 3\beta + \frac{3}{4} \cosh \beta$, we arrive at an equation of the form 1.3.8:

$$\int_a^x \left\{ \frac{1}{4} \cosh[3\lambda(x-t)] + \frac{3}{4} \cosh[\lambda(x-t)] \right\} y(t) dt = f(x).$$

$$16. \int_a^x [\cosh^3(\lambda x) - \cosh^3(\lambda t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = \frac{1}{3\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sinh(\lambda x) \cosh^2(\lambda x)} \right].$$

$$17. \int_a^x [A \cosh^3(\lambda x) + B \cosh^3(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cosh^3(\lambda x)$. For $B = -A$, see equation 1.3.16.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\cosh(\lambda x)]^{-\frac{3A}{A+B}} \int_a^x [\cosh(\lambda t)]^{-\frac{3B}{A+B}} f'_t(t) dt \right\}.$$

$$18. \int_a^x [A \cosh^2(\lambda x) \cosh(\mu t) + B \cosh(\beta x) \cosh^2(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \cosh^2(\lambda x)$, $h_1(t) = \cosh(\mu t)$, $g_2(x) = B \cosh(\beta x)$, and $h_2(t) = \cosh^2(\gamma t)$.

$$19. \int_a^x \cosh^4[\lambda(x-t)] y(t) dt = f(x).$$

Let us transform the kernel of the integral equation using the formula

$$\cosh^4 \beta = \frac{1}{8} \cosh 4\beta + \frac{1}{2} \cosh 2\beta + \frac{3}{8}, \quad \text{where } \beta = \lambda(x-t),$$

and differentiate the resulting equation with respect to x . Then we obtain an equation of the form 2.3.18:

$$y(x) + \lambda \int_a^x \left\{ \frac{1}{2} \sinh[4\lambda(x-t)] + \sinh[2\lambda(x-t)] \right\} y(t) dt = f'_x(x).$$

$$20. \int_a^x [\cosh(\lambda x) - \cosh(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{\sinh(\lambda x)}{\lambda^n n!} \left[\frac{1}{\sinh(\lambda x)} \frac{d}{dx} \right]^{n+1} f(x).$$

$$21. \int_a^x \sqrt{\cosh x - \cosh t} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \sinh x \left(\frac{1}{\sinh x} \frac{d}{dx} \right)^2 \int_a^x \frac{\sinh t f(t) dt}{\sqrt{\cosh x - \cosh t}}.$$

$$22. \int_a^x \frac{y(t) dt}{\sqrt{\cosh x - \cosh t}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{\sinh t f(t) dt}{\sqrt{\cosh x - \cosh t}}.$$

$$23. \int_a^x (\cosh x - \cosh t)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k \sinh x \left(\frac{1}{\sinh x} \frac{d}{dx} \right)^2 \int_a^x \frac{\sinh t f(t) dt}{(\cosh x - \cosh t)^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$24. \int_a^x (\cosh^\mu x - \cosh^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \cosh^\mu x$.

$$\text{Solution: } y(x) = \frac{1}{\mu} \frac{d}{dx} \left[\frac{f'_x(x)}{\sinh x \cosh^{\mu-1} x} \right].$$

$$25. \int_a^x (A \cosh^\mu x + B \cosh^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cosh^\mu x$. For $B = -A$, see equation 1.3.24.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\cosh(\lambda x)]^{-\frac{A\mu}{A+B}} \int_a^x [\cosh(\lambda t)]^{-\frac{B\mu}{A+B}} f'_t(t) dt \right\}.$$

$$26. \int_a^x \frac{y(t) dt}{(\cosh x - \cosh t)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi} \frac{d}{dx} \int_a^x \frac{\sinh t f(t) dt}{(\cosh x - \cosh t)^{1-\lambda}}.$$

$$27. \int_a^x (x-t) \cosh[\lambda(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Differentiating the equation twice yields

$$y(x) + 2\lambda \int_a^x \sinh[\lambda(x-t)] y(t) dt + \lambda^2 \int_a^x (x-t) \cosh[\lambda(x-t)] y(t) dt = f''_{xx}(x).$$

Eliminating the third term on the right-hand side with the aid of the original equation, we arrive at an equation of the form 2.3.16:

$$y(x) + 2\lambda \int_a^x \sinh[\lambda(x-t)] y(t) dt = f''_{xx}(x) - \lambda^2 f(x).$$

$$28. \int_a^x \frac{\cosh[\lambda(x-t)]}{\sqrt{x-t}} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \int_a^x \frac{\cosh[\lambda(x-t)]}{\sqrt{x-t}} [f''_{tt}(t) - \lambda^2 f(t)] dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 436).

$$29. \int_a^x \sqrt{x-t} \cosh(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} f'_t(t) dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 437), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$30. \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 437), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$31. \int_x^\infty \frac{\cosh(\lambda\sqrt{t-x})}{\sqrt{t-x}} y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{1}{\pi} \frac{d}{dx} \int_x^\infty \frac{\cos(\lambda\sqrt{t-x})}{\sqrt{t-x}} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 439), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$32. \int_0^x \frac{\cosh(\lambda\sqrt{x^2-t^2})}{\sqrt{x^2-t^2}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \frac{d}{dx} \int_0^x t \frac{\cos(\lambda\sqrt{x^2-t^2})}{\sqrt{x^2-t^2}} f(t) dt.$$

$$33. \int_x^\infty \frac{\cosh(\lambda\sqrt{t^2-x^2})}{\sqrt{t^2-x^2}} y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{2}{\pi} \frac{d}{dx} \int_x^\infty t \frac{\cos(\lambda\sqrt{t^2-x^2})}{\sqrt{t^2-x^2}} f(t) dt.$$

$$34. \int_0^x \frac{\cosh(\lambda\sqrt{xt-t^2})}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi x} \int_0^x \frac{\cos(\lambda\sqrt{x^2-xt})}{\sqrt{x-t}} [f(t)/2 + t f'_t(t)] dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 438), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$35. \int_0^x \frac{\cosh(\lambda\sqrt{x^2-xt})}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{\sqrt{x}}{\pi} \frac{d}{dx} \left[\sqrt{x} \int_0^x \frac{\cos(\lambda\sqrt{xt-t^2})}{\sqrt{x-t}} f(t) dt \right].$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 438), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$36. \int_a^x \frac{\cosh[\lambda\sqrt{(x-t)(x-t+\gamma)}]}{\sqrt{x-t}} y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{2}{\pi\lambda^2} \int_a^x \frac{\sinh[\lambda\sqrt{(x-t)(x-t-\gamma)}]}{\sqrt{x-t-\gamma}} \int_a^t \sinh[\lambda(t-s)] \left(\frac{d^2}{ds^2} - \lambda^2 \right)^2 f(s) ds dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 438), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$37. \int_a^x [Ax^\beta + B \cosh^\gamma(\lambda t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = B \cosh^\gamma(\lambda t) + C$.

$$38. \int_a^x [A \cosh^\gamma(\lambda x) + Bt^\beta + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh^\gamma(\lambda x)$ and $h(t) = Bt^\beta + C$.

$$39. \int_a^x (Ax^\lambda \cosh^\mu t + Bt^\beta \cosh^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \cosh^\mu t$, $g_2(x) = B \cosh^\gamma x$, and $h_2(t) = t^\beta$.

1.3-2. Kernels Containing Hyperbolic Sine.

$$40. \int_a^x \sinh[\lambda(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution: $y(x) = \frac{1}{\lambda} f''_{xx}(x) - \lambda f(x)$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 435).

$$41. \int_a^x \frac{\sinh[\lambda(x-t)]}{\sqrt{x-t}} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \int_a^x \frac{\sinh[\lambda(x-t)]}{\sqrt{x-t}} [f''_{tt}(t) - \lambda^2 f(t)] dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 436).

$$42. \int_a^x \frac{\sinh[\lambda(x-t)]}{(x-t)^{3/2}} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \int_a^x \frac{\sinh[\lambda(x-t)]}{\sqrt{x-t}} \left[f''_{tt}(t) - \lambda^2 f(t) + \frac{f'(t)}{x-t} \right] dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 437).

$$43. \int_a^x \{ \sinh[\lambda(x-t)] + b \} y(t) dt = f(x).$$

For $b = 0$, see equation 1.3.40. Assume that $b \neq 0$.

Differentiating the equation with respect to x , we arrive at an equation of the form 2.3.3:

$$y(x) + \frac{\lambda}{b} \int_a^x \cosh[\lambda(x-t)] y(t) dt = \frac{1}{b} f'_x(x).$$

Solution:

$$y(x) = \frac{1}{b} f'_x(x) + \int_a^x R(x-t) f'_t(t) dt,$$

$$R(x) = \frac{\lambda}{b^2} \exp\left(-\frac{\lambda x}{2b}\right) \left[\frac{\lambda}{2bk} \sinh(kx) - \cosh(kx) \right], \quad k = \frac{\lambda\sqrt{1+4b^2}}{2b}.$$

$$44. \int_a^x \sinh(\lambda x + \beta t) y(t) dt = f(x).$$

For $\beta = -\lambda$, see equation 1.3.40. Assume that $\beta \neq -\lambda$.

Differentiating the equation with respect to x twice yields

$$\sinh[(\lambda + \beta)x] y(x) + \lambda \int_a^x \cosh(\lambda x + \beta t) y(t) dt = f'_x(x), \quad (1)$$

$$\{ \sinh[(\lambda + \beta)x] y(x) \}'_x + \lambda \cosh[(\lambda + \beta)x] y(x) + \lambda^2 \int_a^x \sinh(\lambda x + \beta t) y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the first-order linear ordinary differential equation

$$w'_x + \lambda \coth[(\lambda + \beta)x] w = f''_{xx}(x) - \lambda^2 f(x), \quad w = \sinh[(\lambda + \beta)x] y(x). \quad (3)$$

Setting $x = a$ in (1) yields the initial condition $w(a) = f'_x(a)$. On solving equation (3) with this condition, after some manipulations we obtain the solution of the original integral equation in the form

$$y(x) = \frac{1}{\sinh[(\lambda + \beta)x]} f'_x(x) - \frac{\lambda \cosh[(\lambda + \beta)x]}{\sinh^2[(\lambda + \beta)x]} f(x)$$

$$- \frac{\lambda\beta}{\sinh^{k+1}[(\lambda + \beta)x]} \int_a^x f(t) \sinh^{k-2}[(\lambda + \beta)t] dt, \quad k = \frac{\lambda}{\lambda + \beta}.$$

$$45. \int_a^x [\sinh(\lambda x) - \sinh(\lambda t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This is a special case of equation 1.9.2 with $g(x) = \sinh(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\cosh(\lambda x)} \right].$$

$$46. \int_a^x [A \sinh(\lambda x) + B \sinh(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \sinh(\lambda x)$. For $B = -A$, see equation 1.3.45.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\sinh(\lambda x)]^{-\frac{A}{A+B}} \int_a^x [\sinh(\lambda t)]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$47. \int_a^x [A \sinh(\lambda x) + B \sinh(\mu t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sinh(\lambda x)$ and $h(t) = B \sinh(\mu t)$.

$$48. \int_a^x \{ \mu \sinh[\lambda(x-t)] - \lambda \sinh[\mu(x-t)] \} y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{f'''_{xxx} - (\lambda^2 + \mu^2)f''_{xx} + \lambda^2\mu^2 f}{\mu\lambda^3 - \lambda\mu^3}, \quad f = f(x).$$

$$49. \int_a^x \{ A_1 \sinh[\lambda_1(x-t)] + A_2 \sinh[\lambda_2(x-t)] \} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

1°. Introduce the notation

$$I_1 = \int_a^x \sinh[\lambda_1(x-t)]y(t) dt, \quad I_2 = \int_a^x \sinh[\lambda_2(x-t)]y(t) dt,$$

$$J_1 = \int_a^x \cosh[\lambda_1(x-t)]y(t) dt, \quad J_2 = \int_a^x \cosh[\lambda_2(x-t)]y(t) dt.$$

Let us successively differentiate the integral equation four times. As a result, we have (the first line is the original equation):

$$A_1 I_1 + A_2 I_2 = f, \quad f = f(x), \tag{1}$$

$$A_1 \lambda_1 J_1 + A_2 \lambda_2 J_2 = f'_x, \tag{2}$$

$$(A_1 \lambda_1 + A_2 \lambda_2)y + A_1 \lambda_1^2 I_1 + A_2 \lambda_2^2 I_2 = f''_{xx}, \tag{3}$$

$$(A_1 \lambda_1 + A_2 \lambda_2)y'_x + A_1 \lambda_1^3 J_1 + A_2 \lambda_2^3 J_2 = f'''_{xxx}, \tag{4}$$

$$(A_1 \lambda_1 + A_2 \lambda_2)y''_{xx} + (A_1 \lambda_1^3 + A_2 \lambda_2^3)y + A_1 \lambda_1^4 I_1 + A_2 \lambda_2^4 I_2 = f''''_{xxxx}. \tag{5}$$

Eliminating I_1 and I_2 from (1), (3), and (5), we arrive at the following second-order linear ordinary differential equation with constant coefficients:

$$(A_1 \lambda_1 + A_2 \lambda_2)y''_{xx} - \lambda_1 \lambda_2 (A_1 \lambda_2 + A_2 \lambda_1)y = f''''_{xxxx} - (\lambda_1^2 + \lambda_2^2)f''_{xx} + \lambda_1^2 \lambda_2^2 f. \tag{6}$$

The initial conditions can be obtained by substituting $x = a$ into (3) and (4):

$$(A_1 \lambda_1 + A_2 \lambda_2)y(a) = f''_{xx}(a), \quad (A_1 \lambda_1 + A_2 \lambda_2)y'_x(a) = f'''_{xxx}(a). \tag{7}$$

Solving the differential equation (6) under conditions (7) allows us to find the solution of the integral equation.

2°. Denote

$$\Delta = \lambda_1 \lambda_2 \frac{A_1 \lambda_2 + A_2 \lambda_1}{A_1 \lambda_1 + A_2 \lambda_2}.$$

2.1. Solution for $\Delta > 0$:

$$(A_1 \lambda_1 + A_2 \lambda_2)y(x) = f''_{xx}(x) + Bf(x) + C \int_a^x \sinh[k(x-t)]f(t) dt,$$

$$k = \sqrt{\Delta}, \quad B = \Delta - \lambda_1^2 - \lambda_2^2, \quad C = \frac{1}{\sqrt{\Delta}} [\Delta^2 - (\lambda_1^2 + \lambda_2^2)\Delta + \lambda_1^2 \lambda_2^2].$$

2.2. Solution for $\Delta < 0$:

$$(A_1 \lambda_1 + A_2 \lambda_2)y(x) = f''_{xx}(x) + Bf(x) + C \int_a^x \sin[k(x-t)]f(t) dt,$$

$$k = \sqrt{-\Delta}, \quad B = \Delta - \lambda_1^2 - \lambda_2^2, \quad C = \frac{1}{\sqrt{-\Delta}} [\Delta^2 - (\lambda_1^2 + \lambda_2^2)\Delta + \lambda_1^2 \lambda_2^2].$$

2.3. Solution for $\Delta = 0$:

$$(A_1 \lambda_1 + A_2 \lambda_2)y(x) = f''_{xx}(x) - (\lambda_1^2 + \lambda_2^2)f(x) + \lambda_1^2 \lambda_2^2 \int_a^x (x-t)f(t) dt.$$

2.4. Solution for $\Delta = \infty$:

$$y(x) = \frac{f''''_{xxxx} - (\lambda_1^2 + \lambda_2^2)f''_{xx} + \lambda_1^2 \lambda_2^2 f}{A_1 \lambda_1^3 + A_2 \lambda_2^3}, \quad f = f(x).$$

In the last case, the relation $A_1 \lambda_1 + A_2 \lambda_2 = 0$ is valid, and the right-hand side of the integral equation is assumed to satisfy the conditions $f(a) = f'_x(a) = f''_{xx}(a) = f''''_{xxxx}(a) = 0$.

50.
$$\int_a^x \{ A \sinh[\lambda(x-t)] + B \sinh[\mu(x-t)] + C \sinh[\beta(x-t)] \} y(t) dt = f(x).$$

It assumed that $f(a) = f'_x(a) = 0$. Differentiating the integral equation twice yields

$$(A\lambda + B\mu + C\beta)y(x) + \int_a^x \{ A\lambda^2 \sinh[\lambda(x-t)] + B\mu^2 \sinh[\mu(x-t)] \} y(t) dt$$

$$+ C\beta^2 \int_a^x \sinh[\beta(x-t)]y(t) dt = f''_{xx}(x).$$

Eliminating the last integral with the aid of the original equation, we arrive at an equation of the form 2.3.18:

$$(A\lambda + B\mu + C\beta)y(x)$$

$$+ \int_a^x \{ A(\lambda^2 - \beta^2) \sinh[\lambda(x-t)] + B(\mu^2 - \beta^2) \sinh[\mu(x-t)] \} y(t) dt = f''_{xx}(x) - \beta^2 f(x).$$

In the special case $A\lambda + B\mu + C\beta = 0$, this is an equation of the form 1.3.49.

51.
$$\int_a^x \sinh^2[\lambda(x-t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Differentiating yields an equation of the form 1.3.40:

$$\int_a^x \sinh[2\lambda(x-t)]y(t) dt = \frac{1}{\lambda} f'_x(x).$$

Solution:
$$y(x) = \frac{1}{2} \lambda^{-2} f''''_{xxxx}(x) - 2 f'_x(x).$$

$$52. \int_a^x [\sinh^2(\lambda x) - \sinh^2(\lambda t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sinh(2\lambda x)} \right].$$

$$53. \int_a^x [A \sinh^2(\lambda x) + B \sinh^2(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \sinh^2(\lambda x)$. For $B = -A$, see equation 1.3.52.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\sinh(\lambda x)]^{-\frac{2A}{A+B}} \int_a^x [\sinh(\lambda t)]^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$54. \int_a^x [A \sinh^2(\lambda x) + B \sinh^2(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sinh^2(\lambda x)$ and $h(t) = B \sinh^2(\mu t)$.

$$55. \int_a^x \sinh[\lambda(x-t)] \sinh[\lambda(x+t)] y(t) dt = f(x).$$

Using the formula

$$\sinh(\alpha - \beta) \sinh(\alpha + \beta) = \frac{1}{2} [\cosh(2\alpha) - \cosh(2\beta)], \quad \alpha = \lambda x, \quad \beta = \lambda t,$$

we reduce the original equation to an equation of the form 1.3.5:

$$\int_a^x [\cosh(2\lambda x) - \cosh(2\lambda t)] y(t) dt = 2f(x).$$

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sinh(2\lambda x)} \right].$$

$$56. \int_a^x [A \sinh(\lambda x) \sinh(\mu t) + B \sinh(\beta x) \sinh(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \sinh(\lambda x)$, $h_1(t) = \sinh(\mu t)$, $g_2(x) = B \sinh(\beta x)$, and $h_2(t) = \sinh(\gamma t)$.

$$57. \int_a^x \sinh^3[\lambda(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0.$$

Using the formula $\sinh^3 \beta = \frac{1}{4} \sinh 3\beta - \frac{3}{4} \sinh \beta$, we arrive at an equation of the form 1.3.49:

$$\int_a^x \left\{ \frac{1}{4} \sinh[3\lambda(x-t)] - \frac{3}{4} \sinh[\lambda(x-t)] \right\} y(t) dt = f(x).$$

$$58. \int_a^x [\sinh^3(\lambda x) - \sinh^3(\lambda t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This is a special case of equation 1.9.2 with $g(x) = \sinh^3(\lambda x)$.

$$59. \int_a^x [A \sinh^3(\lambda x) + B \sinh^3(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \sinh^3(\lambda x)$.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\sinh(\lambda x)]^{-\frac{3A}{A+B}} \int_a^x [\sinh(\lambda t)]^{-\frac{3B}{A+B}} f'_t(t) dt \right\}.$$

$$60. \int_a^x [A \sinh^2(\lambda x) \sinh(\mu t) + B \sinh(\beta x) \sinh^2(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \sinh^2(\lambda x)$, $h_1(t) = \sinh(\mu t)$, $g_2(x) = B \sinh(\beta x)$, and $h_2(t) = \sinh^2(\gamma t)$.

$$61. \int_a^x \sinh^4[\lambda(x-t)] y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = \dots = f''''_{xxxx}(a) = 0$.

Let us transform the kernel of the integral equation using the formula

$$\sinh^4 \beta = \frac{1}{8} \cosh 4\beta - \frac{1}{2} \cosh 2\beta + \frac{3}{8}, \quad \text{where } \beta = \lambda(x-t),$$

and differentiate the resulting equation with respect to x . Then we arrive at an equation of the form 1.3.49:

$$\lambda \int_a^x \left\{ \frac{1}{2} \sinh[4\lambda(x-t)] - \sinh[2\lambda(x-t)] \right\} y(t) dt = f'_x(x).$$

$$62. \int_a^x \sinh^n[\lambda(x-t)] y(t) dt = f(x), \quad n = 2, 3, \dots$$

It is assumed that $f(a) = f'_x(a) = \dots = f^{(n)}_x(a) = 0$.

1°. Let us differentiate the equation with respect to x twice and transform the kernel of the resulting integral equation using the formula $\cosh^2 \beta = 1 + \sinh^2 \beta$, where $\beta = \lambda(x-t)$. Then we have

$$\lambda^2 n^2 \int_a^x \sinh^n[\lambda(x-t)] y(t) dt + \lambda^2 n(n-1) \int_a^x \sinh^{n-2}[\lambda(x-t)] y(t) dt = f''_{xx}(x).$$

Eliminating the first term on the left-hand side with the aid of the original equation, we obtain

$$\int_a^x \sinh^{n-2}[\lambda(x-t)] y(t) dt = \frac{1}{\lambda^2 n(n-1)} [f''_{xx}(x) - \lambda^2 n^2 f(x)].$$

This equation has the same form as the original equation, but the exponent of the kernel has been reduced by two.

By applying this technique sufficiently many times, we finally arrive at simple integral equations of the form 1.1.1 (for even n) or 1.3.40 (for odd n).

2°. Solution:

$$y(x) = \frac{1}{\lambda^n n!} \left(\frac{d}{dx} + n\lambda \right) \left(\frac{d}{dx} + (n-2)\lambda \right) \dots \left(\frac{d}{dx} - n\lambda \right) f(x).$$

$$63. \int_a^x \sinh(\lambda\sqrt{x-t})y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \frac{d^2}{dx^2} \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 437), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$64. \int_x^\infty \sinh(\lambda\sqrt{t-x})y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \frac{d^2}{dx^2} \int_x^\infty \frac{\cos(\lambda\sqrt{t-x})}{\sqrt{t-x}} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 439), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$65. \int_a^x \sqrt{\sinh x - \sinh t} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \cosh x \left(\frac{1}{\cosh x} \frac{d}{dx} \right)^2 \int_a^x \frac{\cosh t f(t) dt}{\sqrt{\sinh x - \sinh t}}.$$

$$66. \int_a^x \frac{y(t) dt}{\sqrt{\sinh x - \sinh t}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{\cosh t f(t) dt}{\sqrt{\sinh x - \sinh t}}.$$

$$67. \int_a^x (\sinh x - \sinh t)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k \cosh x \left(\frac{1}{\cosh x} \frac{d}{dx} \right)^2 \int_a^x \frac{\cosh t f(t) dt}{(\sinh x - \sinh t)^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$68. \int_a^x (\sinh^\mu x - \sinh^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \sinh^\mu x$.

$$\text{Solution: } y(x) = \frac{1}{\mu} \frac{d}{dx} \left[\frac{f'_x(x)}{\cosh x \sinh^{\mu-1} x} \right].$$

$$69. \int_a^x [A \sinh^\mu(\lambda x) + B \sinh^\mu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \sinh^\mu(\lambda x)$.

Solution with $B \neq -A$:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\sinh(\lambda x)]^{-\frac{A\mu}{A+B}} \int_a^x [\sinh(\lambda t)]^{-\frac{B\mu}{A+B}} f'_t(t) dt \right\}.$$

$$70. \int_a^x \frac{y(t) dt}{(\sinh x - \sinh t)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi} \frac{d}{dx} \int_a^x \frac{\cosh t f(t) dt}{(\sinh x - \sinh t)^{1-\lambda}}.$$

$$71. \int_a^x (x-t) \sinh[\lambda(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Double differentiation yields

$$2\lambda \int_a^x \cosh[\lambda(x-t)] y(t) dt + \lambda^2 \int_a^x (x-t) \sinh[\lambda(x-t)] y(t) dt = f''_{xx}(x).$$

Eliminating the second term on the left-hand side with the aid of the original equation, we arrive at an equation of the form 1.3.1:

$$\int_a^x \cosh[\lambda(x-t)] y(t) dt = \frac{1}{2\lambda} [f''_{xx}(x) - \lambda^2 f(x)].$$

Solution:

$$y(x) = \frac{1}{2\lambda} f'''_{xxx}(x) - \lambda f'_x(x) + \frac{1}{2} \lambda^3 \int_a^x f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 436).

$$72. \int_a^x \frac{\sinh[\lambda(x-t)]}{\sqrt{x-t}} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \int_a^x \frac{\sinh[\lambda(x-t)]}{\sqrt{x-t}} [f''_{tt}(t) - \lambda^2 f(t)] dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 436).

$$73. \int_a^x \frac{\sinh[\lambda(x-t)]}{(x-t)^{3/2}} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \int_a^x \frac{\sinh[\lambda(x-t)]}{\sqrt{x-t}} \left[f''_{tt}(t) - \lambda^2 f(t) + \frac{f'(t)}{x-t} \right] dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 437).

$$74. \int_a^x \frac{\sinh[\lambda\sqrt{(x-t)(x-t+\gamma)}]}{\sqrt{x-t+\gamma}} y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{2}{\pi\lambda^2} \int_a^x \frac{\cosh[\lambda\sqrt{(x-t)(x-t+\gamma)}]}{\sqrt{x-t}} \int_a^t \sinh[\lambda(t-s)] \left(\frac{d^2}{ds^2} + \lambda^2 \right)^2 f(s) ds dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 438), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$75. \int_a^x [Ax^\beta + B \sinh^\gamma(\lambda t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = B \sinh^\gamma(\lambda t) + C$.

$$76. \int_a^x [A \sinh^\gamma(\lambda x) + Bt^\beta + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sinh^\gamma(\lambda x)$ and $h(t) = Bt^\beta + C$.

$$77. \int_a^x (Ax^\lambda \sinh^\mu t + Bt^\beta \sinh^\gamma x)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \sinh^\mu t$, $g_2(x) = B \sinh^\gamma x$, and $h_2(t) = t^\beta$.

1.3-3. Kernels Containing Hyperbolic Tangent.

$$78. \int_a^x [\tanh(\lambda x) - \tanh(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \tanh(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda} [\cosh^2(\lambda x) f'_x(x)]'_x.$$

$$79. \int_a^x [A \tanh(\lambda x) + B \tanh(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \tanh(\lambda x)$. For $B = -A$, see equation 1.3.78.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\tanh(\lambda x)]^{-\frac{A}{A+B}} \int_a^x [\tanh(\lambda t)]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$80. \int_a^x [A \tanh(\lambda x) + B \tanh(\mu t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh(\lambda x)$ and $h(t) = B \tanh(\mu t) + C$.

$$81. \int_a^x [\tanh^2(\lambda x) - \tanh^2(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \tanh^2(\lambda x)$.

$$\text{Solution: } y(x) = \frac{d}{dx} \left[\frac{\cosh^3(\lambda x) f'_x(x)}{2\lambda \sinh(\lambda x)} \right].$$

$$82. \int_a^x [A \tanh^2(\lambda x) + B \tanh^2(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \tanh^2(\lambda x)$. For $B = -A$, see equation 1.3.81.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\tanh(\lambda x)]^{-\frac{2A}{A+B}} \int_a^x [\tanh(\lambda t)]^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$83. \int_a^x [A \tanh^2(\lambda x) + B \tanh^2(\mu t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh^2(\lambda x)$ and $h(t) = B \tanh^2(\mu t) + C$.

$$84. \int_a^x [\tanh(\lambda x) - \tanh(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{1}{\lambda^n n! \cosh^2(\lambda x)} \left[\cosh^2(\lambda x) \frac{d}{dx} \right]^{n+1} f(x).$$

$$85. \int_a^x \sqrt{\tanh x - \tanh t} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi \cosh^2 x} \left(\cosh^2 x \frac{d}{dx} \right)^2 \int_a^x \frac{f(t) dt}{\cosh^2 t \sqrt{\tanh x - \tanh t}}.$$

$$86. \int_a^x \frac{y(t) dt}{\sqrt{\tanh x - \tanh t}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{\cosh^2 t \sqrt{\tanh x - \tanh t}}.$$

$$87. \int_a^x (\tanh x - \tanh t)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi\lambda \cosh^2 x} \left(\cosh^2 x \frac{d}{dx} \right)^2 \int_a^x \frac{f(t) dt}{\cosh^2 t (\tanh x - \tanh t)^\lambda}.$$

$$88. \int_a^x (\tanh^\mu x - \tanh^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \tanh^\mu x$.

$$\text{Solution: } y(x) = \frac{1}{\mu} \frac{d}{dx} \left[\frac{\cosh^{\mu+1} x f'_x(x)}{\sinh^{\mu-1} x} \right].$$

$$89. \int_a^x (A \tanh^\mu x + B \tanh^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \tanh^\mu x$. For $B = -A$, see equation 1.3.88.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\tanh(\lambda x)]^{-\frac{A\mu}{A+B}} \int_a^x [\tanh(\lambda t)]^{-\frac{B\mu}{A+B}} f'_t(t) dt \right\}.$$

$$90. \int_a^x \frac{y(t) dt}{[\tanh(\lambda x) - \tanh(\lambda t)]^\mu} = f(x), \quad 0 < \mu < 1.$$

This is a special case of equation 1.9.44 with $g(x) = \tanh(\lambda x)$ and $h(x) \equiv 1$.

Solution:

$$y(x) = \frac{\lambda \sin(\pi\mu)}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{\cosh^2(\lambda t) [\tanh(\lambda x) - \tanh(\lambda t)]^{1-\mu}}.$$

$$91. \int_a^x [Ax^\beta + B \tanh^\gamma(\lambda t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = B \tanh^\gamma(\lambda t) + C$.

$$92. \int_a^x [A \tanh^\gamma(\lambda x) + Bt^\beta + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh^\gamma(\lambda x)$ and $h(t) = Bt^\beta + C$.

$$93. \int_a^x (Ax^\lambda \tanh^\mu t + Bt^\beta \tanh^\gamma x)y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \tanh^\mu t$, $g_2(x) = B \tanh^\gamma x$, and $h_2(t) = t^\beta$.

1.3-4. Kernels Containing Hyperbolic Cotangent.

$$94. \int_a^x [\operatorname{coth}(\lambda x) - \operatorname{coth}(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \operatorname{coth}(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{1}{\lambda} \frac{d}{dx} [\sinh^2(\lambda x) f'_x(x)].$$

$$95. \int_a^x [A \operatorname{coth}(\lambda x) + B \operatorname{coth}(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \operatorname{coth}(\lambda x)$. For $B = -A$, see equation 1.3.94.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\tanh(\lambda x)]^{\frac{A}{A+B}} \int_a^x [\tanh(\lambda t)]^{\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$96. \int_a^x [A \operatorname{coth}(\lambda x) + B \operatorname{coth}(\mu t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \operatorname{coth}(\lambda x)$ and $h(t) = B \operatorname{coth}(\mu t) + C$.

$$97. \int_a^x [\operatorname{coth}^2(\lambda x) - \operatorname{coth}^2(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \operatorname{coth}^2(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{d}{dx} \left[\frac{\sinh^3(\lambda x) f'_x(x)}{2\lambda \cosh(\lambda x)} \right].$$

$$98. \int_a^x [A \operatorname{coth}^2(\lambda x) + B \operatorname{coth}^2(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \operatorname{coth}^2(\lambda x)$. For $B = -A$, see equation 1.3.97.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\tanh(\lambda x)]^{\frac{2A}{A+B}} \int_a^x [\tanh(\lambda t)]^{\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$99. \int_a^x [A \operatorname{coth}^2(\lambda x) + B \operatorname{coth}^2(\mu t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \operatorname{coth}^2(\lambda x)$ and $h(t) = B \operatorname{coth}^2(\mu t) + C$.

$$100. \int_a^x [\coth(\lambda x) - \coth(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{(-1)^n}{\lambda^n n! \sinh^2(\lambda x)} \left[\sinh^2(\lambda x) \frac{d}{dx} \right]^{n+1} f(x).$$

$$101. \int_a^x (\coth^\mu x - \coth^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \coth^\mu x$.

$$\text{Solution: } y(x) = -\frac{1}{\mu} \frac{d}{dx} \left[\frac{\sinh^{\mu+1} x f'_x(x)}{\cosh^{\mu-1} x} \right].$$

$$102. \int_a^x (A \coth^\mu x + B \coth^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \coth^\mu x$. For $B = -A$, see equation 1.3.101.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ \left| \tanh x \right|^{\frac{A\mu}{A+B}} \int_a^x \left| \tanh t \right|^{\frac{B\mu}{A+B}} f'_t(t) dt \right\}.$$

$$103. \int_a^x [Ax^\beta + B \coth^\gamma(\lambda t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = B \coth^\gamma(\lambda t) + C$.

$$104. \int_a^x [A \coth^\gamma(\lambda x) + Bt^\beta + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \coth^\gamma(\lambda x)$ and $h(t) = Bt^\beta + C$.

$$105. \int_a^x (Ax^\lambda \coth^\mu t + Bt^\beta \coth^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \coth^\mu t$, $g_2(x) = B \coth^\gamma x$, and $h_2(t) = t^\beta$.

1.3-5. Kernels Containing Combinations of Hyperbolic Functions.

$$106. \int_a^x \{ \cosh[\lambda(x-t)] + A \sinh[\mu(x-t)] \} y(t) dt = f(x).$$

Let us differentiate the equation with respect to x and then eliminate the integral with the hyperbolic cosine. As a result, we arrive at an equation of the form 2.3.16:

$$y(x) + (\lambda - A^2\mu) \int_a^x \sinh[\mu(x-t)] y(t) dt = f'_x(x) - A\mu f(x).$$

$$107. \int_a^x [A \cosh(\lambda x) + B \sinh(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh(\lambda x)$ and $h(t) = B \sinh(\mu t) + C$.

$$108. \int_a^x [A \cosh^2(\lambda x) + B \sinh^2(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh^2(\lambda x)$ and $h(t) = B \sinh^2(\mu t) + C$.

$$109. \int_a^x \sinh[\lambda(x-t)] \cosh[\lambda(x+t)] y(t) dt = f(x).$$

Using the formula

$$\sinh(\alpha - \beta) \cosh(\alpha + \beta) = \frac{1}{2} [\sinh(2\alpha) - \sinh(2\beta)], \quad \alpha = \lambda x, \quad \beta = \lambda t,$$

we reduce the original equation to an equation of the form 1.3.45:

$$\int_a^x [\sinh(2\lambda x) - \sinh(2\lambda t)] y(t) dt = 2f(x).$$

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\cosh(2\lambda x)} \right].$$

$$110. \int_a^x \cosh[\lambda(x-t)] \sinh[\lambda(x+t)] y(t) dt = f(x).$$

Using the formula

$$\cosh(\alpha - \beta) \sinh(\alpha + \beta) = \frac{1}{2} [\sinh(2\alpha) + \sinh(2\beta)], \quad \alpha = \lambda x, \quad \beta = \lambda t,$$

we reduce the original equation to an equation of the form 1.3.46 with $A = B = 1$:

$$\int_a^x [\sinh(2\lambda x) + \sinh(2\lambda t)] y(t) dt = 2f(x).$$

$$111. \int_a^x [A \cosh(\lambda x) \sinh(\mu t) + B \cosh(\beta x) \sinh(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \cosh(\lambda x)$, $h_1(t) = \sinh(\mu t)$, $g_2(x) = B \cosh(\beta x)$, and $h_2(t) = \sinh(\gamma t)$.

$$112. \int_a^x [\sinh(\lambda x) \cosh(\mu t) + \sinh(\beta x) \cosh(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \sinh(\lambda x)$, $h_1(t) = \cosh(\mu t)$, $g_2(x) = \sinh(\beta x)$, and $h_2(t) = \cosh(\gamma t)$.

$$113. \int_a^x [\cosh(\lambda x) \cosh(\mu t) + \sinh(\beta x) \sinh(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \cosh(\lambda x)$, $h_1(t) = \cosh(\mu t)$, $g_2(x) = \sinh(\beta x)$, and $h_2(t) = \sinh(\gamma t)$.

$$114. \int_a^x [A \cosh^\beta(\lambda x) + B \sinh^\gamma(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh^\beta(\lambda x)$ and $h(t) = B \sinh^\gamma(\mu t)$.

$$115. \int_a^x [A \sinh^\beta(\lambda x) + B \cosh^\gamma(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sinh^\beta(\lambda x)$ and $h(t) = B \cosh^\gamma(\mu t)$.

$$116. \int_a^x (Ax^\lambda \cosh^\mu t + Bt^\beta \sinh^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \cosh^\mu t$, $g_2(x) = B \sinh^\gamma x$, and $h_2(t) = t^\beta$.

$$117. \int_a^x \{ (x-t) \sinh[\lambda(x-t)] - \lambda(x-t)^2 \cosh[\lambda(x-t)] \} y(t) dt = f(x).$$

Solution:

$$y(x) = \int_a^x g(t) dt,$$

where

$$g(t) = \sqrt{\frac{\pi}{2\lambda}} \frac{1}{64\lambda^5} \left(\frac{d^2}{dt^2} - \lambda^2 \right)^6 \int_a^t (t-\tau)^{\frac{5}{2}} I_{\frac{5}{2}}[\lambda(t-\tau)] f(\tau) d\tau.$$

$$118. \int_a^x \left\{ \frac{\sinh[\lambda(x-t)]}{x-t} - \lambda \cosh[\lambda(x-t)] \right\} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{2\lambda^4} \left(\frac{d^2}{dx^2} - \lambda^2 \right)^3 \int_a^x \sinh[\lambda(x-t)] f(t) dt.$$

$$119. \int_a^x [\sinh(\lambda\sqrt{x-t}) - \lambda\sqrt{x-t} \cosh(\lambda\sqrt{x-t})] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = -\frac{4}{\pi\lambda^3} \frac{d^3}{dx^3} \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$120. \int_a^x (Ax^\lambda \sinh^\mu t + Bt^\beta \cosh^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \sinh^\mu t$, $g_2(x) = B \cosh^\gamma x$, and $h_2(t) = t^\beta$.

$$121. \int_a^x [A \tanh(\lambda x) + B \coth(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh(\lambda x)$ and $h(t) = B \coth(\mu t) + C$.

$$122. \int_a^x [A \tanh^2(\lambda x) + B \coth^2(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh^2(\lambda x)$ and $h(t) = B \coth^2(\mu t)$.

$$123. \int_a^x [\tanh(\lambda x) \coth(\mu t) + \tanh(\beta x) \coth(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \tanh(\lambda x)$, $h_1(t) = \coth(\mu t)$, $g_2(x) = \tanh(\beta x)$, and $h_2(t) = \coth(\gamma t)$.

$$124. \int_a^x [\coth(\lambda x) \tanh(\mu t) + \coth(\beta x) \tanh(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \coth(\lambda x)$, $h_1(t) = \tanh(\mu t)$, $g_2(x) = \coth(\beta x)$, and $h_2(t) = \tanh(\gamma t)$.

$$125. \int_a^x [\tanh(\lambda x) \tanh(\mu t) + \coth(\beta x) \coth(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \tanh(\lambda x)$, $h_1(t) = \tanh(\mu t)$, $g_2(x) = \coth(\beta x)$, and $h_2(t) = \coth(\gamma t)$.

$$126. \int_a^x [A \tanh^\beta(\lambda x) + B \coth^\gamma(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh^\beta(\lambda x)$ and $h(t) = B \coth^\gamma(\mu t)$.

$$127. \int_a^x [A \coth^\beta(\lambda x) + B \tanh^\gamma(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \coth^\beta(\lambda x)$ and $h(t) = B \tanh^\gamma(\mu t)$.

$$128. \int_a^x (Ax^\lambda \tanh^\mu t + Bt^\beta \coth^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \tanh^\mu t$, $g_2(x) = B \coth^\gamma x$, and $h_2(t) = t^\beta$.

$$129. \int_a^x (Ax^\lambda \coth^\mu t + Bt^\beta \tanh^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \coth^\mu t$, $g_2(x) = B \tanh^\gamma x$, and $h_2(t) = t^\beta$.

1.4. Equations Whose Kernels Contain Logarithmic Functions

1.4-1. Kernels Containing Logarithmic Functions.

$$1. \int_a^x (\ln x - \ln t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \ln x$.

Solution: $y(x) = x f''_{xx}(x) + f'_x(x)$.

$$2. \int_0^x \ln(x-t) y(t) dt = f(x).$$

Solution:

$$y(x) = - \int_0^x f''_{tt}(t) dt \int_0^\infty \frac{(x-t)^z e^{-Cz}}{\Gamma(z+1)} dz - f'_x(0) \int_0^\infty \frac{x^z e^{-Cz}}{\Gamma(z+1)} dz,$$

where $C = \lim_{k \rightarrow \infty} \left(1 + \frac{1}{2} + \dots + \frac{1}{k+1} - \ln k \right) = 0.5772 \dots$ is the Euler constant and $\Gamma(z)$ is the gamma function.

⊙ References: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971), A. G. Butkovskii (1979).

$$3. \int_a^x [\ln(x-t) + A]y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{d}{dx} \int_a^x \nu_A(x-t)f(t) dt, \quad \nu_A(x) = \frac{d}{dx} \int_0^\infty \frac{x^z e^{(A-C)z}}{\Gamma(z+1)} dz,$$

where $C = 0.5772 \dots$ is the Euler constant and $\Gamma(z)$ is the gamma function.

For $a = 0$, the solution can be written in the form

$$y(x) = -\int_0^x f''_{tt}(t) dt \int_0^\infty \frac{(x-t)^z e^{(A-C)z}}{\Gamma(z+1)} dz - f'_x(0) \int_0^\infty \frac{x^z e^{(A-C)z}}{\Gamma(z+1)} dz.$$

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$4. \int_a^x (A \ln x + B \ln t)y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \ln x$. For $B = -A$, see equation 1.4.1.

Solution:

$$y(x) = \frac{\text{sign}(\ln x)}{A+B} \frac{d}{dx} \left\{ |\ln x|^{-\frac{A}{A+B}} \int_a^x |\ln t|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$5. \int_a^x (A \ln x + B \ln t + C)y(t) dt = f(x).$$

This is a special case of equation 1.9.5 with $g(x) = x$.

$$6. \int_a^x [\ln^2(\lambda x) - \ln^2(\lambda t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = \frac{d}{dx} \left[\frac{x f'_x(x)}{2 \ln(\lambda x)} \right].$$

$$7. \int_a^x [A \ln^2(\lambda x) + B \ln^2(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \ln^2(\lambda x)$. For $B = -A$, see equation 1.4.6.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |\ln(\lambda x)|^{-\frac{2A}{A+B}} \int_a^x |\ln(\lambda t)|^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$8. \int_a^x [A \ln^2(\lambda x) + B \ln^2(\mu t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \ln^2(\lambda x)$ and $h(t) = B \ln^2(\mu t) + C$.

$$9. \int_a^x [\ln(x/t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{1}{n! x} \left(x \frac{d}{dx} \right)^{n+1} f(x).$$

$$10. \int_a^x (\ln^2 x - \ln^2 t)^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{\ln x}{2^n n! x} \left(x \frac{d}{dx} \right)^{n+1} f(x).$$

$$11. \int_a^x \ln \left(\frac{x+b}{t+b} \right) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \ln(x+b)$.

$$\text{Solution: } y(x) = (x+b) f''_{xx}(x) + f'_x(x).$$

$$12. \int_a^x \sqrt{\ln(x/t)} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi x} \left(x \frac{d}{dx} \right)^2 \int_a^x \frac{f(t) dt}{t \sqrt{\ln(x/t)}}.$$

$$13. \int_a^x \frac{y(t) dt}{\sqrt{\ln(x/t)}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{t \sqrt{\ln(x/t)}}.$$

$$14. \int_a^x [\ln^\mu(\lambda x) - \ln^\mu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \ln^\mu(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\mu} \frac{d}{dx} [x \ln^{1-\mu}(\lambda x) f'_x(x)].$$

$$15. \int_a^x [A \ln^\beta(\lambda x) + B \ln^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \ln^\beta(\lambda x)$ and $h(t) = B \ln^\gamma(\mu t) + C$.

$$16. \int_a^x [\ln(x/t)]^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{k}{x} \left(x \frac{d}{dx} \right)^2 \int_a^x \frac{f(t) dt}{t [\ln(x/t)]^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$17. \int_a^x \frac{y(t) dt}{[\ln(x/t)]^\lambda} = f(x), \quad 0 < \lambda < 1.$$

This is a special case of equation 1.9.44 with $g(x) = \ln x$ and $h(x) \equiv 1$.

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{t [\ln(x/t)]^{1-\lambda}}.$$

$$18. \int_0^x \ln \frac{\sqrt{x} + \sqrt{x-t}}{\sqrt{x} - \sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_0^x \frac{\sqrt{t}}{\sqrt{x-t}} \frac{d}{dt} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 451).

$$19. \int_x^\infty \ln \frac{\sqrt{t} + \sqrt{t-x}}{\sqrt{t} - \sqrt{t-x}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{1}{\sqrt{x}} \frac{d}{dx} \int_x^\infty \frac{t}{\sqrt{t-x}} \frac{d}{dt} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 452).

1.4-2. Kernels Containing Power-Law and Logarithmic Functions.

$$20. \int_a^x (x-t) [\ln(x-t) + A] y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{d^2}{dx^2} \int_a^x \nu_A(x-t) f(t) dt, \quad \nu_A(x) = \frac{d}{dx} \int_0^\infty \frac{x^z e^{(A-C)z}}{\Gamma(z+1)} dz,$$

where $C = 0.5772 \dots$ is the Euler constant and $\Gamma(z)$ is the gamma function.

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$21. \int_a^x \frac{\ln(x-t) + A}{(x-t)^\lambda} y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = -\frac{\sin(\pi\lambda)}{\pi} \frac{d}{dx} \int_a^x \frac{F(t) dt}{(x-t)^{1-\lambda}}, \quad F(x) = \int_a^x \nu_h(x-t) f(t) dt,$$

$$\nu_h(x) = \frac{d}{dx} \int_0^\infty \frac{x^z e^{hz}}{\Gamma(z+1)} dz, \quad h = A + \psi(1-\lambda),$$

where $\Gamma(z)$ is the gamma function and $\psi(z) = [\Gamma(z)]'_z$ is the logarithmic derivative of the gamma function.

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$22. \int_a^x \frac{(x-t)^{\alpha-1}}{\Gamma(\alpha)} [\ln(x-t) + A] y(t) dt = f(x), \quad \alpha > 0.$$

Solution:

$$y(x) = -\frac{1}{\Gamma([\alpha] - \alpha + 1)} \left(\frac{d}{dx} \right)^{[\alpha]+1} \int_a^x \frac{F(t) dt}{(x-t)^{\alpha-[\alpha]}}, \quad F(x) = \int_a^x \nu_h(x-t) f(t) dt,$$

$$\nu_h(x) = \frac{d}{dx} \int_0^\infty \frac{x^z e^{hz}}{\Gamma(z+1)} dz, \quad h = A + \psi(\alpha),$$

where $\Gamma(z)$ is the gamma function and $\psi(z) = [\Gamma(z)]'_z$ is the logarithmic derivative of the gamma function.

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993, p. 483).

$$23. \int_a^x (t^\beta \ln^\lambda x - x^\beta \ln^\lambda t) y(t) dt = f(x).$$

This is a special case of equation 1.9.11 with $g(x) = \ln^\lambda x$ and $h(t) = t^\beta$.

$$24. \int_a^x (At^\beta \ln^\lambda x + Bx^\mu \ln^\gamma t) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \ln^\lambda x$, $h_1(t) = t^\beta$, $g_2(x) = Bx^\mu$, and $h_2(t) = \ln^\gamma t$.

$$25. \int_a^x \ln \left(\frac{x^\mu + b}{ct^\lambda + s} \right) y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = \ln(x^\mu + b)$ and $h(t) = -\ln(ct^\lambda + s)$.

1.5. Equations Whose Kernels Contain Trigonometric Functions

1.5-1. Kernels Containing Cosine.

$$1. \int_a^x \cos[\lambda(x-t)] y(t) dt = f(x).$$

$$\text{Solution: } y(x) = f'_x(x) + \lambda^2 \int_a^x f(x) dx.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 442).

$$2. \int_a^x \{ \cos[\lambda(x-t)] - 1 \} y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

$$\text{Solution: } y(x) = -\frac{1}{\lambda^2} f'''_{xxx}(x) - f'_x(x).$$

$$3. \int_a^x \{ \cos[\lambda(x-t)] + b \} y(t) dt = f(x).$$

For $b = 0$, see equation 1.5.1. For $b = -1$, see equation 1.5.2. For $\lambda = 0$, see equation 1.1.1. Differentiating the equation with respect to x , we arrive at an equation of the form 2.5.16:

$$y(x) - \frac{\lambda}{b+1} \int_a^x \sin[\lambda(x-t)] y(t) dt = \frac{f'_x(x)}{b+1}.$$

1°. Solution with $b(b+1) > 0$:

$$y(x) = \frac{f'_x(x)}{b+1} + \frac{\lambda^2}{k(b+1)^2} \int_a^x \sin[k(x-t)] f'_t(t) dt, \quad \text{where } k = \lambda \sqrt{\frac{b}{b+1}}.$$

2°. Solution with $b(b+1) < 0$:

$$y(x) = \frac{f'_x(x)}{b+1} + \frac{\lambda^2}{k(b+1)^2} \int_a^x \sinh[k(x-t)] f'_t(t) dt, \quad \text{where } k = \lambda \sqrt{\frac{-b}{b+1}}.$$

$$4. \int_a^x \cos(\lambda x + \beta t) y(t) dt = f(x).$$

Differentiating the equation with respect to x twice yields

$$\cos[(\lambda + \beta)x]y(x) - \lambda \int_a^x \sin(\lambda x + \beta t)y(t) dt = f'_x(x), \quad (1)$$

$$\left\{ \cos[(\lambda + \beta)x]y(x) \right\}'_x - \lambda \sin[(\lambda + \beta)x]y(x) - \lambda^2 \int_a^x \cos(\lambda x + \beta t)y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the first-order linear ordinary differential equation

$$w'_x - \lambda \tan[(\lambda + \beta)x]w = f''_{xx}(x) + \lambda^2 f(x), \quad w = \cos[(\lambda + \beta)x]y(x). \quad (3)$$

Setting $x = a$ in (1) yields the initial condition $w(a) = f'_x(a)$. On solving equation (3) under this condition, after some transformations we obtain the solution of the original integral equation in the form

$$y(x) = \frac{1}{\cos[(\lambda + \beta)x]} f'_x(x) + \frac{\lambda \sin[(\lambda + \beta)x]}{\cos^2[(\lambda + \beta)x]} f(x) - \frac{\lambda \beta}{\cos^{k+1}[(\lambda + \beta)x]} \int_a^x f(t) \cos^{k-2}[(\lambda + \beta)t] dt, \quad k = \frac{\lambda}{\lambda + \beta}.$$

$$5. \int_a^x [\cos(\lambda x) - \cos(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \cos(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sin(\lambda x)} \right].$$

$$6. \int_a^x [A \cos(\lambda x) + B \cos(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cos(\lambda x)$. For $B = -A$, see equation 1.5.5.

Solution with $B \neq -A$:

$$y(x) = \frac{\text{sign } \cos(\lambda x)}{A + B} \frac{d}{dx} \left\{ |\cos(\lambda x)|^{-\frac{A}{A+B}} \int_a^x |\cos(\lambda t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$7. \int_a^x [A \cos(\lambda x) + B \cos(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cos(\lambda x)$ and $h(t) = B \cos(\mu t) + C$.

$$8. \int_a^x \{ A_1 \cos[\lambda_1(x - t)] + A_2 \cos[\lambda_2(x - t)] \} y(t) dt = f(x).$$

The equation is equivalent to the equation

$$\int_a^x \{ B_1 \sin[\lambda_1(x - t)] + B_2 \sin[\lambda_2(x - t)] \} y(t) dt = F(x),$$

$$B_1 = \frac{A_1}{\lambda_1}, \quad B_2 = \frac{A_2}{\lambda_2}, \quad F(x) = \int_a^x f(t) dt,$$

which has the form 1.5.41. (Differentiation of this equation yields the original integral equation.)

$$9. \int_a^x \cos^2[\lambda(x-t)]y(t) dt = f(x).$$

Differentiating yields an equation of the form 2.5.16:

$$y(x) - \lambda \int_a^x \sin[2\lambda(x-t)]y(t) dt = f'_x(x).$$

Solution:

$$y(x) = f'_x(x) + \frac{2\lambda^2}{k} \int_a^x \sin[k(x-t)]f'_t(t) dt, \quad \text{where } k = \lambda\sqrt{2}.$$

$$10. \int_a^x [\cos^2(\lambda x) - \cos^2(\lambda t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = -\frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sin(2\lambda x)} \right].$$

$$11. \int_a^x [A \cos^2(\lambda x) + B \cos^2(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cos^2(\lambda x)$. For $B = -A$, see equation 1.5.10.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\cos(\lambda x)]^{-\frac{2A}{A+B}} \int_a^x [\cos(\lambda t)]^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$12. \int_a^x [A \cos^2(\lambda x) + B \cos^2(\mu t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cos^2(\lambda x)$ and $h(t) = B \cos^2(\mu t) + C$.

$$13. \int_a^x \cos[\lambda(x-t)] \cos[\lambda(x+t)]y(t) dt = f(x).$$

Using the trigonometric formula

$$\cos(\alpha - \beta) \cos(\alpha + \beta) = \frac{1}{2} [\cos(2\alpha) + \cos(2\beta)], \quad \alpha = \lambda x, \quad \beta = \lambda t,$$

we reduce the original equation to an equation of the form 1.5.6 with $A = B = 1$:

$$\int_a^x [\cos(2\lambda x) + \cos(2\lambda t)]y(t) dt = 2f(x).$$

Solution with $\cos(2\lambda x) > 0$:

$$y(x) = \frac{d}{dx} \left[\frac{1}{\sqrt{\cos(2\lambda x)}} \int_a^x \frac{f'_t(t) dt}{\sqrt{\cos(2\lambda t)}} \right].$$

$$14. \int_a^x \cos[\lambda(x-t)] \cos[\mu(x-t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{1}{\sqrt{\lambda^2 - \mu^2}} \left[\frac{d^2}{dx^2} + (\lambda + \mu)^2 \right] \left[\frac{d^2}{dx^2} + (\lambda - \mu)^2 \right] \int_a^x \int_a^t \sin[\sqrt{\lambda^2 + \mu^2} (t-s)] f(s) ds dt.$$

$$15. \int_a^x [A \cos(\lambda x) \cos(\mu t) + B \cos(\beta x) \cos(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \cos(\lambda x)$, $h_1(t) = \cos(\mu t)$, $g_2(x) = B \cos(\beta x)$, and $h_2(t) = \cos(\gamma t)$.

$$16. \int_a^x \cos^3[\lambda(x-t)] y(t) dt = f(x).$$

Using the formula $\cos^3 \beta = \frac{1}{4} \cos 3\beta + \frac{3}{4} \cos \beta$, we arrive at an equation of the form 1.5.8:

$$\int_a^x \left\{ \frac{1}{4} \cos[3\lambda(x-t)] + \frac{3}{4} \cos[\lambda(x-t)] \right\} y(t) dt = f(x).$$

$$17. \int_a^x [\cos^3(\lambda x) - \cos^3(\lambda t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = -\frac{1}{3\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sin(\lambda x) \cos^2(\lambda x)} \right].$$

$$18. \int_a^x [A \cos^3(\lambda x) + B \cos^3(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cos^3(\lambda x)$. For $B = -A$, see equation 1.5.17.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\cos(\lambda x)]^{-\frac{3A}{A+B}} \int_a^x [\cos(\lambda t)]^{-\frac{3B}{A+B}} f'_t(t) dt \right\}.$$

$$19. \int_a^x [\cos^2(\lambda x) \cos(\mu t) + \cos(\beta x) \cos^2(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \cos^2(\lambda x)$, $h_1(t) = \cos(\mu t)$, $g_2(x) = \cos(\beta x)$, and $h_2(t) = \cos^2(\gamma t)$.

$$20. \int_a^x \cos^4[\lambda(x-t)] y(t) dt = f(x).$$

Let us transform the kernel of the integral equation using the trigonometric formula $\cos^4 \beta = \frac{1}{8} \cos 4\beta + \frac{1}{2} \cos 2\beta + \frac{3}{8}$, where $\beta = \lambda(x-t)$, and differentiate the resulting equation with respect to x . Then we arrive at an equation of the form 2.5.18:

$$y(x) - \lambda \int_a^x \left\{ \frac{1}{2} \sin[4\lambda(x-t)] + \sin[2\lambda(x-t)] \right\} y(t) dt = f'_x(x).$$

$$21. \int_a^x [\cos(\lambda x) - \cos(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{(-1)^n}{\lambda^n n!} \sin(\lambda x) \left[\frac{1}{\sin(\lambda x)} \frac{d}{dx} \right]^{n+1} f(x).$$

$$22. \int_a^x \sqrt{\cos t - \cos x} y(t) dt = f(x).$$

This is a special case of equation 1.9.40 with $g(x) = 1 - \cos x$.

Solution:

$$y(x) = \frac{2}{\pi} \sin x \left(\frac{1}{\sin x} \frac{d}{dx} \right)^2 \int_a^x \frac{\sin t f(t) dt}{\sqrt{\cos t - \cos x}}.$$

$$23. \int_a^x \frac{y(t) dt}{\sqrt{\cos t - \cos x}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{\sin t f(t) dt}{\sqrt{\cos t - \cos x}}.$$

$$24. \int_a^x (\cos t - \cos x)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k \sin x \left(\frac{1}{\sin x} \frac{d}{dx} \right)^2 \int_a^x \frac{\sin t f(t) dt}{(\cos t - \cos x)^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$25. \int_a^x (\cos^\mu x - \cos^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \cos^\mu x$.

$$\text{Solution: } y(x) = -\frac{1}{\mu} \frac{d}{dx} \left[\frac{f'_x(x)}{\sin x \cos^{\mu-1} x} \right].$$

$$26. \int_a^x (A \cos^\mu x + B \cos^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cos^\mu x$. For $B = -A$, see equation 1.5.25.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |\cos x|^{-\frac{A\mu}{A+B}} \int_a^x |\cos t|^{-\frac{B\mu}{A+B}} f'_t(t) dt \right\}.$$

$$27. \int_a^x \frac{y(t) dt}{(\cos t - \cos x)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi} \frac{d}{dx} \int_a^x \frac{\sin t f(t) dt}{(\cos t - \cos x)^{1-\lambda}}.$$

$$28. \int_a^x (x-t) \cos[\lambda(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Differentiating the equation twice yields

$$y(x) - 2\lambda \int_a^x \sin[\lambda(x-t)] y(t) dt - \lambda^2 \int_a^x (x-t) \cos[\lambda(x-t)] y(t) dt = f''_{xx}(x).$$

Eliminating the third term on the left-hand side with the aid of the original equation, we arrive at an equation of the form 2.5.16:

$$y(x) - 2\lambda \int_a^x \sin[\lambda(x-t)] y(t) dt = f''_{xx}(x) + \lambda^2 f(x).$$

$$29. \int_a^x \frac{\cos[\lambda(x-t)]}{\sqrt{x-t}} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \int_a^x \frac{\sin[\lambda(x-t)]}{\sqrt{x-t}} [f''_{tt}(t) + \lambda^2 f(t)] dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 445).

$$30. \int_a^x \sqrt{x-t} \cos(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f'_t(t) dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 445–446), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$31. \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 446), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$32. \int_x^\infty \frac{\cos(\lambda\sqrt{t-x})}{\sqrt{t-x}} y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{1}{\pi} \frac{d}{dx} \int_x^\infty \frac{\cosh(\lambda\sqrt{t-x})}{\sqrt{t-x}} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 448), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$33. \int_0^x \frac{\cos(\lambda\sqrt{x^2-t^2})}{\sqrt{x^2-t^2}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \frac{d}{dx} \int_0^x t \frac{\cosh(\lambda\sqrt{x^2-t^2})}{\sqrt{x^2-t^2}} f(t) dt.$$

$$34. \int_x^\infty \frac{\cos(\lambda\sqrt{t^2-x^2})}{\sqrt{t^2-x^2}} y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{2}{\pi} \frac{d}{dx} \int_x^\infty t \frac{\cosh(\lambda\sqrt{t^2-x^2})}{\sqrt{t^2-x^2}} f(t) dt.$$

$$35. \int_0^x \frac{\cos(\lambda\sqrt{xt-t^2})}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi x} \int_0^x \frac{\cosh(\lambda\sqrt{x^2-xt})}{\sqrt{x-t}} [f(t)/2 + t f'_t(t)] dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 446).

$$36. \int_0^x \frac{\cos(\lambda\sqrt{x^2 - xt})}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{\sqrt{x}}{\pi} \frac{d}{dx} \left[\sqrt{x} \int_0^x \frac{\cosh(\lambda\sqrt{xt-t^2})}{\sqrt{x-t}} f(t) dt \right].$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 446).

$$37. \int_a^x \frac{\cos[\lambda\sqrt{(x-t)(x-t+\gamma)}]}{\sqrt{x-t}} y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{2}{\pi\lambda^2} \int_a^x \frac{\sin[\lambda\sqrt{(x-t)(x-t-\gamma)}]}{\sqrt{x-t-\gamma}} \int_a^t \sin[\lambda(t-s)] \left(\frac{d^2}{ds^2} + \lambda^2 \right)^2 f(s) ds dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 447).

$$38. \int_a^x [Ax^\beta + B \cos^\gamma(\lambda t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = B \cos^\gamma(\lambda t) + C$.

$$39. \int_a^x [A \cos^\gamma(\lambda x) + Bt^\beta + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cos^\gamma(\lambda x)$ and $h(t) = Bt^\beta + C$.

$$40. \int_a^x (Ax^\lambda \cos^\mu t + Bt^\beta \cos^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \cos^\mu t$, $g_2(x) = B \cos^\gamma x$, and $h_2(t) = t^\beta$.

1.5-2. Kernels Containing Sine.

$$41. \int_a^x \sin[\lambda(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution: $y(x) = \frac{1}{\lambda} f''_{xx}(x) + \lambda f(x)$.

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 442).

$$42. \int_a^x \{ \sin[\lambda(x-t)] + b \} y(t) dt = f(x).$$

For $b = 0$, see equation 1.5.41. Assume that $b \neq 0$.

Differentiating the equation with respect to x yields an equation of the form 2.5.3:

$$y(x) + \frac{\lambda}{b} \int_a^x \cos[\lambda(x-t)] y(t) dt = \frac{1}{b} f'_x(x).$$

$$43. \int_a^x \sin(\lambda x + \beta t) y(t) dt = f(x).$$

For $\beta = -\lambda$, see equation 1.5.41. Assume that $\beta \neq -\lambda$.

Differentiating the equation with respect to x twice yields

$$\sin[(\lambda + \beta)x]y(x) + \lambda \int_a^x \cos(\lambda x + \beta t)y(t) dt = f'_x(x), \quad (1)$$

$$\{\sin[(\lambda + \beta)x]y(x)\}'_x + \lambda \cos[(\lambda + \beta)x]y(x) - \lambda^2 \int_a^x \sin(\lambda x + \beta t)y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the first-order linear ordinary differential equation

$$w'_x + \lambda \cot[(\lambda + \beta)x]w = f''_{xx}(x) + \lambda^2 f(x), \quad w = \sin[(\lambda + \beta)x]y(x). \quad (3)$$

Setting $x = a$ in (1) yields the initial condition $w(a) = f'_x(a)$. On solving equation (3) under this condition, after some transformation we obtain the solution of the original integral equation in the form

$$y(x) = \frac{1}{\sin[(\lambda + \beta)x]} f'_x(x) - \frac{\lambda \cos[(\lambda + \beta)x]}{\sin^2[(\lambda + \beta)x]} f(x) - \frac{\lambda \beta}{\sin^{k+1}[(\lambda + \beta)x]} \int_a^x f(t) \sin^{k-2}[(\lambda + \beta)t] dt, \quad k = \frac{\lambda}{\lambda + \beta}.$$

$$44. \int_a^x [\sin(\lambda x) - \sin(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \sin(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\cos(\lambda x)} \right].$$

$$45. \int_a^x [A \sin(\lambda x) + B \sin(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \sin(\lambda x)$. For $B = -A$, see equation 1.5.44.

Solution with $B \neq -A$:

$$y(x) = \frac{\text{sign} \sin(\lambda x)}{A + B} \frac{d}{dx} \left\{ |\sin(\lambda x)|^{-\frac{A}{A+B}} \int_a^x |\sin(\lambda t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$46. \int_a^x [A \sin(\lambda x) + B \sin(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sin(\lambda x)$ and $h(t) = B \sin(\mu t) + C$.

$$47. \int_a^x \{\mu \sin[\lambda(x-t)] - \lambda \sin[\mu(x-t)]\} y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{f''''_{xxx} + (\lambda^2 + \mu^2) f''_{xx} + \lambda^2 \mu^2 f}{\lambda \mu^3 - \lambda^3 \mu}, \quad f = f(x).$$

$$48. \int_a^x \{A_1 \sin[\lambda_1(x-t)] + A_2 \sin[\lambda_2(x-t)]\} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This equation can be solved in the same manner as equation 1.3.49, i.e., by reducing it to a second-order linear ordinary differential equation with constant coefficients.

Let

$$\Delta = -\lambda_1 \lambda_2 \frac{A_1 \lambda_2 + A_2 \lambda_1}{A_1 \lambda_1 + A_2 \lambda_2}.$$

1°. Solution for $\Delta > 0$:

$$(A_1 \lambda_1 + A_2 \lambda_2) y(x) = f''_{xx}(x) + B f(x) + C \int_a^x \sinh[k(x-t)] f(t) dt,$$

$$k = \sqrt{\Delta}, \quad B = \Delta + \lambda_1^2 + \lambda_2^2, \quad C = \frac{1}{\sqrt{\Delta}} [\Delta^2 + (\lambda_1^2 + \lambda_2^2) \Delta + \lambda_1^2 \lambda_2^2].$$

2°. Solution for $\Delta < 0$:

$$(A_1 \lambda_1 + A_2 \lambda_2) y(x) = f''_{xx}(x) + B f(x) + C \int_a^x \sin[k(x-t)] f(t) dt,$$

$$k = \sqrt{-\Delta}, \quad B = \Delta + \lambda_1^2 + \lambda_2^2, \quad C = \frac{1}{\sqrt{-\Delta}} [\Delta^2 + (\lambda_1^2 + \lambda_2^2) \Delta + \lambda_1^2 \lambda_2^2].$$

3°. Solution for $\Delta = 0$:

$$(A_1 \lambda_1 + A_2 \lambda_2) y(x) = f''_{xx}(x) + (\lambda_1^2 + \lambda_2^2) f(x) + \lambda_1^2 \lambda_2^2 \int_a^x (x-t) f(t) dt.$$

4°. Solution for $\Delta = \infty$:

$$y(x) = -\frac{f''''_{xxxx} + (\lambda_1^2 + \lambda_2^2) f''_{xx} + \lambda_1^2 \lambda_2^2 f}{A_1 \lambda_1^3 + A_2 \lambda_2^3}, \quad f = f(x).$$

In the last case, the relation $A_1 \lambda_1 + A_2 \lambda_2 = 0$ holds and the right-hand side of the integral equation is assumed to satisfy the conditions $f(a) = f'_x(a) = f''_{xx}(a) = f''''_{xxxx}(a) = 0$.

Remark. The solution can be obtained from the solution of equation 1.3.49 in which the change of variables $\lambda_k \rightarrow i \lambda_k$, $A_k \rightarrow -i A_k$, $i^2 = -1$ ($k = 1, 2$), should be made.

$$49. \int_a^x \{A \sin[\lambda(x-t)] + B \sin[\mu(x-t)] + C \sin[\beta(x-t)]\} y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = 0$. Differentiating the integral equation twice yields

$$(A\lambda + B\mu + C\beta)y(x) - \int_a^x \{A\lambda^2 \sin[\lambda(x-t)] + B\mu^2 \sin[\mu(x-t)]\} y(t) dt$$

$$- C\beta^2 \int_a^x \sin[\beta(x-t)] y(t) dt = f''_{xx}(x).$$

Eliminating the last integral with the aid of the original equation, we arrive at an equation of the form 2.5.18:

$$(A\lambda + B\mu + C\beta)y(x) + \int_a^x \{A(\beta^2 - \lambda^2) \sin[\lambda(x-t)]$$

$$+ B(\beta^2 - \mu^2) \sin[\mu(x-t)]\} y(t) dt = f''_{xx}(x) + \beta^2 f(x).$$

In the special case $A\lambda + B\mu + C\beta = 0$, this is an equation of the form 1.5.41.

$$50. \int_a^x \sin^2[\lambda(x-t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Differentiation yields an equation of the form 1.5.41:

$$\int_a^x \sin[2\lambda(x-t)]y(t) dt = \frac{1}{\lambda} f'_x(x).$$

$$\text{Solution: } y(x) = \frac{1}{2}\lambda^{-2} f'''_{xxx}(x) + 2f'_x(x).$$

$$51. \int_a^x [\sin^2(\lambda x) - \sin^2(\lambda t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sin(2\lambda x)} \right].$$

$$52. \int_a^x [A \sin^2(\lambda x) + B \sin^2(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \sin^2(\lambda x)$. For $B = -A$, see equation 1.5.51.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |\sin(\lambda x)|^{-\frac{2A}{A+B}} \int_a^x |\sin(\lambda t)|^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$53. \int_a^x [A \sin^2(\lambda x) + B \sin^2(\mu t) + C]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sin^2(\lambda x)$ and $h(t) = B \sin^2(\mu t) + C$.

$$54. \int_a^x \sin[\lambda(x-t)] \sin[\lambda(x+t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Using the trigonometric formula

$$\sin(\alpha - \beta) \sin(\alpha + \beta) = \frac{1}{2} [\cos(2\beta) - \cos(2\alpha)], \quad \alpha = \lambda x, \quad \beta = \lambda t,$$

we reduce the original equation to an equation of the form 1.5.5:

$$\int_a^x [\cos(2\lambda x) - \cos(2\lambda t)]y(t) dt = -2f(x).$$

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sin(2\lambda x)} \right].$$

$$55. \int_a^x \sin[\lambda(x-t)] \sin[\mu(x-t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Solution:

$$y(x) = \left[\frac{d^2}{dx^2} + (\lambda + \mu)^2 \right] \left[\frac{d^2}{dx^2} + (\lambda - \mu)^2 \right] \frac{1}{2\lambda\mu} \int_a^x f(t) dt.$$

$$56. \int_a^x [\sin(\lambda x) \sin(\mu t) + \sin(\beta x) \sin(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \sin(\lambda x)$, $h_1(t) = \sin(\mu t)$, $g_2(x) = \sin(\beta x)$, and $h_2(t) = \sin(\gamma t)$.

$$57. \int_a^x \sin^3[\lambda(x-t)] y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Using the formula $\sin^3 \beta = -\frac{1}{4} \sin 3\beta + \frac{3}{4} \sin \beta$, we arrive at an equation of the form 1.5.48:

$$\int_a^x \left\{ -\frac{1}{4} \sin[3\lambda(x-t)] + \frac{3}{4} \sin[\lambda(x-t)] \right\} y(t) dt = f(x).$$

$$58. \int_a^x [\sin^3(\lambda x) - \sin^3(\lambda t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

This is a special case of equation 1.9.2 with $g(x) = \sin^3(\lambda x)$.

$$59. \int_a^x [A \sin^3(\lambda x) + B \sin^3(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \sin^3(\lambda x)$. For $B = -A$, see equation 1.5.58.

Solution:

$$y(x) = \frac{\text{sign} \sin(\lambda x)}{A+B} \frac{d}{dx} \left\{ |\sin(\lambda x)|^{-\frac{3A}{A+B}} \int_a^x |\sin(\lambda t)|^{-\frac{3B}{A+B}} f'_t(t) dt \right\}.$$

$$60. \int_a^x [\sin^2(\lambda x) \sin(\mu t) + \sin(\beta x) \sin^2(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \sin^2(\lambda x)$, $h_1(t) = \sin(\mu t)$, $g_2(x) = \sin(\beta x)$, and $h_2(t) = \sin^2(\gamma t)$.

$$61. \int_a^x \sin^4[\lambda(x-t)] y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = \dots = f''''_{xxxx}(a) = 0$.

Let us transform the kernel of the integral equation using the trigonometric formula $\sin^4 \beta = \frac{1}{8} \cos 4\beta - \frac{1}{2} \cos 2\beta + \frac{3}{8}$, where $\beta = \lambda(x-t)$, and differentiate the resulting equation with respect to x . Then we obtain an equation of the form 1.5.48:

$$\lambda \int_a^x \left\{ -\frac{1}{2} \sin[4\lambda(x-t)] + \sin[2\lambda(x-t)] \right\} y(t) dt = f'_x(x).$$

$$62. \int_a^x \sin^n[\lambda(x-t)] y(t) dt = f(x), \quad n = 2, 3, \dots$$

It is assumed that $f(a) = f'_x(a) = \dots = f^{(n)}_x(a) = 0$.

1°. Let us differentiate the equation with respect to x twice and transform the kernel of the resulting integral equation using the formula $\cos^2 \beta = 1 - \sin^2 \beta$, where $\beta = \lambda(x-t)$. We have

$$-\lambda^2 n^2 \int_a^x \sin^n[\lambda(x-t)] y(t) dt + \lambda^2 n(n-1) \int_a^x \sin^{n-2}[\lambda(x-t)] y(t) dt = f''_{xx}(x).$$

Eliminating the first term on the left-hand side with the aid of the original equation, we obtain

$$\int_a^x \sin^{n-2}[\lambda(x-t)]y(t) dt = \frac{1}{\lambda^2 n(n-1)} [f''_{xx}(x) + \lambda^2 n^2 f(x)].$$

This equation has the same form as the original equation, but the degree characterizing the kernel has been reduced by two.

By applying this technique sufficiently many times, we finally arrive at simple integral equations of the form 1.1.1 (for even n) or 1.5.41 (for odd n).

2°. Solution:

$$y(x) = \frac{1}{\lambda^n n!} \left(\frac{d}{dx} \right)^{1-\alpha} \prod_{k=1}^{\beta} \left[\frac{d^2}{dx^2} + (2k + \alpha)\lambda^2 \right] f(x),$$

where $\alpha = n - 2[n/2]$, $\beta = [(n+1)/2]$, $[A]$ denotes the integer part of number A .

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 443).

$$63. \int_a^x (x-t) \sin[\lambda(x-t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Solution:

$$y(x) = \frac{1}{2\lambda} \left(\frac{d^2}{dx^2} + \lambda^2 \right)^2 \int_a^x f(t) dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 444), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$64. \int_a^x \sin(\lambda\sqrt{x-t})y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \frac{d^2}{dx^2} \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

See also Example 2 in Section 10.4.

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 445), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$65. \int_x^\infty \sin(\lambda\sqrt{t-x})y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \frac{d^2}{dx^2} \int_x^\infty \frac{\cos(\lambda\sqrt{t-x})}{\sqrt{t-x}} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 447).

$$66. \int_a^x \frac{\sin[\lambda(x-t)]}{\sqrt{x-t}} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi\lambda} \int_a^x \frac{\cos[\lambda(x-t)]}{\sqrt{x-t}} [f''_{tt}(t) + \lambda^2 f(t)] dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 445), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$67. \int_a^x \frac{\sin[\lambda(x-t)]}{(x-t)^{3/2}} y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi\lambda^2} \int_a^x \frac{\sin[\lambda(x-t)]}{\sqrt{x-t}} \left[f''_{tt}(t) + \lambda^2 f(t) + \frac{f'(t)}{x-t} \right] dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 445), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$68. \int_a^x \frac{\sin [\lambda\sqrt{(x-t)(x-t+\gamma)}]}{\sqrt{x-t+\gamma}} y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{2}{\pi\lambda^2} \int_a^x \frac{\cos[\lambda\sqrt{(x-t)(x-t-\gamma)}]}{\sqrt{x-t}} \int_a^t \sin[\lambda(t-s)] \left(\frac{d^2}{ds^2} + \lambda^2 \right) f(s) ds dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 447).

$$69. \int_a^x \sqrt{\sin x - \sin t} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \cos x \left(\frac{1}{\cos x} \frac{d}{dx} \right)^2 \int_a^x \frac{\cos t f(t) dt}{\sqrt{\sin x - \sin t}}.$$

$$70. \int_a^x \frac{y(t) dt}{\sqrt{\sin x - \sin t}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{\cos t f(t) dt}{\sqrt{\sin x - \sin t}}.$$

$$71. \int_a^x (\sin x - \sin t)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k \cos x \left(\frac{1}{\cos x} \frac{d}{dx} \right)^2 \int_a^x \frac{\cos t f(t) dt}{(\sin x - \sin t)^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$72. \int_a^x (\sin^\mu x - \sin^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \sin^\mu x$.

$$\text{Solution: } y(x) = \frac{1}{\mu} \frac{d}{dx} \left[\frac{f'_x(x)}{\cos x \sin^{\mu-1} x} \right].$$

$$73. \int_a^x \{ A |\sin(\lambda x)|^\mu + B |\sin(\lambda t)|^\mu \} y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = |\sin(\lambda x)|^\mu$.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |\sin(\lambda x)|^{-\frac{A\mu}{A+B}} \int_a^x |\sin(\lambda t)|^{-\frac{B\mu}{A+B}} f'_t(t) dt \right\}.$$

$$74. \int_a^x \frac{y(t) dt}{[\sin(\lambda x) - \sin(\lambda t)]^\mu} = f(x), \quad 0 < \mu < 1.$$

This is a special case of equation 1.9.44 with $g(x) = \sin(\lambda x)$ and $h(x) \equiv 1$.

Solution:

$$y(x) = \frac{\lambda \sin(\pi\mu)}{\pi} \frac{d}{dx} \int_a^x \frac{\cos(\lambda t) f(t) dt}{[\sin(\lambda x) - \sin(\lambda t)]^{1-\mu}}.$$

$$75. \int_a^x (x-t) \sin[\lambda(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Double differentiation yields

$$2\lambda \int_a^x \cos[\lambda(x-t)] y(t) dt - \lambda^2 \int_a^x (x-t) \sin[\lambda(x-t)] y(t) dt = f''_{xx}(x).$$

Eliminating the second integral on the left-hand side of this equation with the aid of the original equation, we arrive at an equation of the form 1.5.1:

$$\int_a^x \cos[\lambda(x-t)] y(t) dt = \frac{1}{2\lambda} [f''_{xx}(x) + \lambda^2 f(x)].$$

Solution:

$$y(x) = \frac{1}{2\lambda} f'''_{xxx}(x) + \lambda f'_x(x) + \frac{1}{2} \lambda^3 \int_a^x f(t) dt.$$

$$76. \int_a^x |\sin(\lambda(x-t))| y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Solution:

$$y(x) = \frac{1}{\lambda} \int_a^x (-1)^{[\lambda(x-t)/\pi]} (f'''_{ttt}(t) + \lambda^2 f'_t(t)) dt,$$

where $[A]$ denotes the integer part of number A .

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 443).

$$77. \int_a^x [Ax^\beta + B \sin^\gamma(\lambda t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = B \sin^\gamma(\lambda t) + C$.

$$78. \int_a^x [A \sin^\gamma(\lambda x) + Bt^\beta + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sin^\gamma(\lambda x)$ and $h(t) = Bt^\beta + C$.

$$79. \int_a^x (Ax^\lambda \sin^\mu t + Bt^\beta \sin^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \sin^\mu t$, $g_2(x) = B \sin^\gamma x$, and $h_2(t) = t^\beta$.

1.5-3. Kernels Containing Tangent.

$$80. \int_a^x [\tan(\lambda x) - \tan(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \tan(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} [\cos^2(\lambda x) f'_x(x)].$$

$$81. \int_a^x [A \tan(\lambda x) + B \tan(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \tan(\lambda x)$. For $B = -A$, see equation 1.5.80.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\tan(\lambda x)]^{-\frac{A}{A+B}} \int_a^x [\tan(\lambda t)]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$82. \int_a^x [A \tan(\lambda x) + B \tan(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tan(\lambda x)$ and $h(t) = B \tan(\mu t) + C$.

$$83. \int_a^x [\tan^2(\lambda x) - \tan^2(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \tan^2(\lambda x)$.

$$\text{Solution: } y(x) = \frac{d}{dx} \left[\frac{\cos^3(\lambda x) f'_x(x)}{2\lambda \sin(\lambda x)} \right].$$

$$84. \int_a^x [A \tan^2(\lambda x) + B \tan^2(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \tan^2(\lambda x)$. For $B = -A$, see equation 1.5.83.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |\tan(\lambda x)|^{-\frac{2A}{A+B}} \int_a^x |\tan(\lambda t)|^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$85. \int_a^x [A \tan^2(\lambda x) + B \tan^2(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tan^2(\lambda x)$ and $h(t) = B \tan^2(\mu t) + C$.

$$86. \int_a^x [\tan(\lambda x) - \tan(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{1}{\lambda^n n! \cos^2(\lambda x)} \left[\cos^2(\lambda x) \frac{d}{dx} \right]^{n+1} f(x).$$

$$87. \int_a^x \sqrt{\tan x - \tan t} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi \cos^2 x} \left(\cos^2 x \frac{d}{dx} \right)^2 \int_a^x \frac{f(t) dt}{\cos^2 t \sqrt{\tan x - \tan t}}.$$

$$88. \int_a^x \frac{y(t) dt}{\sqrt{\tan x - \tan t}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{\cos^2 t \sqrt{\tan x - \tan t}}.$$

$$89. \int_a^x (\tan x - \tan t)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi\lambda \cos^2 x} \left(\cos^2 x \frac{d}{dx} \right)^2 \int_a^x \frac{f(t) dt}{\cos^2 t (\tan x - \tan t)^\lambda}.$$

$$90. \int_a^x (\tan^\mu x - \tan^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \tan^\mu x$.

$$\text{Solution: } y(x) = \frac{1}{\mu} \frac{d}{dx} \left[\frac{\cos^{\mu+1} x f'_x(x)}{\sin^{\mu-1} x} \right].$$

$$91. \int_a^x (A \tan^\mu x + B \tan^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \tan^\mu x$. For $B = -A$, see equation 1.5.90.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\tan(\lambda x)]^{-\frac{A\mu}{A+B}} \int_a^x [\tan(\lambda t)]^{-\frac{B\mu}{A+B}} f'_t(t) dt \right\}.$$

$$92. \int_a^x \frac{y(t) dt}{[\tan(\lambda x) - \tan(\lambda t)]^\mu} = f(x), \quad 0 < \mu < 1.$$

This is a special case of equation 1.9.44 with $g(x) = \tan(\lambda x)$ and $h(x) \equiv 1$.

Solution:

$$y(x) = \frac{\lambda \sin(\pi\mu)}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{\cos^2(\lambda t) [\tan(\lambda x) - \tan(\lambda t)]^{1-\mu}}.$$

$$93. \int_a^x [Ax^\beta + B \tan^\gamma(\lambda t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = B \tan^\gamma(\lambda t) + C$.

$$94. \int_a^x [A \tan^\gamma(\lambda x) + Bt^\beta + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tan^\gamma(\lambda x)$ and $h(t) = Bt^\beta + C$.

$$95. \int_a^x (Ax^\lambda \tan^\mu t + Bt^\beta \tan^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \tan^\mu t$, $g_2(x) = B \tan^\gamma x$, and $h_2(t) = t^\beta$.

1.5-4. Kernels Containing Cotangent.

$$96. \int_a^x [\cot(\lambda x) - \cot(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \cot(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{1}{\lambda} \frac{d}{dx} [\sin^2(\lambda x) f'_x(x)].$$

$$97. \int_a^x [A \cot(\lambda x) + B \cot(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cot(\lambda x)$. For $B = -A$, see equation 1.5.96.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\tan(\lambda x)]^{\frac{A}{A+B}} \int_a^x [\tan(\lambda t)]^{\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$98. \int_a^x [A \cot(\lambda x) + B \cot(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cot(\lambda x)$ and $h(t) = B \cot(\mu t) + C$.

$$99. \int_a^x [\cot^2(\lambda x) - \cot^2(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \cot^2(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{d}{dx} \left[\frac{\sin^3(\lambda x) f'_x(x)}{2\lambda \cos(\lambda x)} \right].$$

$$100. \int_a^x [A \cot^2(\lambda x) + B \cot^2(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cot^2(\lambda x)$. For $B = -A$, see equation 1.5.99.

$$\text{Solution: } y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |\tan(\lambda x)|^{\frac{2A}{A+B}} \int_a^x |\tan(\lambda t)|^{\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$101. \int_a^x [A \cot^2(\lambda x) + B \cot^2(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cot^2(\lambda x)$ and $h(t) = B \cot^2(\mu t) + C$.

$$102. \int_a^x [\cot(\lambda x) - \cot(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{(-1)^n}{\lambda^n n! \sin^2(\lambda x)} \left[\sin^2(\lambda x) \frac{d}{dx} \right]^{n+1} f(x).$$

$$103. \int_a^x (\cot^\mu x - \cot^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \cot^\mu x$.

$$\text{Solution: } y(x) = -\frac{1}{\mu} \frac{d}{dx} \left[\frac{\sin^{\mu+1} x f'_x(x)}{\cos^{\mu-1} x} \right].$$

$$104. \int_a^x (A \cot^\mu x + B \cot^\mu t) y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \cot^\mu x$. For $B = -A$, see equation 1.5.103.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |\tan x|^{\frac{A\mu}{A+B}} \int_a^x |\tan t|^{\frac{B\mu}{A+B}} f'_t(t) dt \right\}.$$

$$105. \int_a^x [Ax^\beta + B \cot^\gamma(\lambda t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = Ax^\beta$ and $h(t) = B \cot^\gamma(\lambda t) + C$.

$$106. \int_a^x [A \cot^\gamma(\lambda x) + Bt^\beta + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cot^\gamma(\lambda x)$ and $h(t) = Bt^\beta + C$.

$$107. \int_a^x (Ax^\lambda \cot^\mu t + Bt^\beta \cot^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \cot^\mu t$, $g_2(x) = B \cot^\gamma x$, and $h_2(t) = t^\beta$.

1.5-5. Kernels Containing Combinations of Trigonometric Functions.

$$108. \int_a^x \{ \cos[\lambda(x-t)] + A \sin[\mu(x-t)] \} y(t) dt = f(x).$$

Differentiating the equation with respect to x followed by eliminating the integral with the cosine yields an equation of the form 2.3.16:

$$y(x) - (\lambda + A^2\mu) \int_a^x \sin[\mu(x-t)] y(t) dt = f'_x(x) - A\mu f(x).$$

$$109. \int_a^x [A \cos(\lambda x) + B \sin(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cos(\lambda x)$ and $h(t) = B \sin(\mu t) + C$.

$$110. \int_a^x [A \sin(\lambda x) + B \cos(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sin(\lambda x)$ and $h(t) = B \cos(\mu t) + C$.

$$111. \int_a^x [A \cos^2(\lambda x) + B \sin^2(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cos^2(\lambda x)$ and $h(t) = B \sin^2(\mu t)$.

$$112. \int_a^x \sin[\lambda(x-t)] \cos[\lambda(x+t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Using the trigonometric formula

$$\sin(\alpha - \beta) \cos(\alpha + \beta) = \frac{1}{2} [\sin(2\alpha) - \sin(2\beta)], \quad \alpha = \lambda x, \quad \beta = \lambda t,$$

we reduce the original equation to an equation of the form 1.5.44:

$$\int_a^x [\sin(2\lambda x) - \sin(2\lambda t)] y(t) dt = 2f(x).$$

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\cos(2\lambda x)} \right].$$

$$113. \int_a^x \cos[\lambda(x-t)] \sin[\lambda(x+t)] y(t) dt = f(x).$$

Using the trigonometric formula

$$\cos(\alpha - \beta) \sin(\alpha + \beta) = \frac{1}{2} [\sin(2\alpha) + \sin(2\beta)], \quad \alpha = \lambda x, \quad \beta = \lambda t,$$

we reduce the original equation to an equation of the form 1.5.45 with $A = B = 1$:

$$\int_a^x [\sin(2\lambda x) + \sin(2\lambda t)] y(t) dt = 2f(x).$$

Solution with $\sin(2\lambda x) > 0$:

$$y(x) = \frac{d}{dx} \left[\frac{1}{\sqrt{\sin(2\lambda x)}} \int_a^x \frac{f'_t(t) dt}{\sqrt{\sin(2\lambda t)}} \right].$$

$$114. \int_a^x \sin[\lambda(x-t)] \cos[\mu(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Solution with $\mu < \lambda$:

$$y(x) = \frac{1}{\lambda \sqrt{\lambda^2 - \mu^2}} \left[\frac{d^2}{dx^2} + (\lambda + \mu)^2 \right] \left[\frac{d^2}{dx^2} + (\lambda - \mu)^2 \right] \int_a^x \sin[\sqrt{\lambda^2 - \mu^2} (x-t)] f(t) dt.$$

Solution with $\mu > \lambda$:

$$y(x) = \frac{1}{\lambda \sqrt{\lambda^2 - \mu^2}} \left[\frac{d^2}{dx^2} + (\lambda + \mu)^2 \right] \left[\frac{d^2}{dx^2} + (\lambda - \mu)^2 \right] \int_a^x \sinh[\sqrt{\mu^2 - \lambda^2} (x-t)] f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 444).

$$115. \int_a^x [A \cos(\lambda x) \sin(\mu t) + B \cos(\beta x) \sin(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \cos(\lambda x)$, $h_1(t) = \sin(\mu t)$, $g_2(x) = B \cos(\beta x)$, and $h_2(t) = \sin(\gamma t)$.

$$116. \int_a^x [A \sin(\lambda x) \cos(\mu t) + B \sin(\beta x) \cos(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \sin(\lambda x)$, $h_1(t) = \cos(\mu t)$, $g_2(x) = B \sin(\beta x)$, and $h_2(t) = \cos(\gamma t)$.

$$117. \int_a^x [A \cos(\lambda x) \cos(\mu t) + B \sin(\beta x) \sin(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A \cos(\lambda x)$, $h_1(t) = \cos(\mu t)$, $g_2(x) = B \sin(\beta x)$, and $h_2(t) = \sin(\gamma t)$.

$$118. \int_a^x [A \cos^\beta(\lambda x) + B \sin^\gamma(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cos^\beta(\lambda x)$ and $h(t) = B \sin^\gamma(\mu t)$.

$$119. \int_a^x [A \sin^\beta(\lambda x) + B \cos^\gamma(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sin^\beta(\lambda x)$ and $h(t) = B \cos^\gamma(\mu t)$.

$$120. \int_a^x (Ax^\lambda \cos^\mu t + Bt^\beta \sin^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \cos^\mu t$, $g_2(x) = B \sin^\gamma x$, and $h_2(t) = t^\beta$.

$$121. \int_a^x (Ax^\lambda \sin^\mu t + Bt^\beta \cos^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \sin^\mu t$, $g_2(x) = B \cos^\gamma x$, and $h_2(t) = t^\beta$.

$$122. \int_a^x \{ (x-t) \sin[\lambda(x-t)] - \lambda(x-t)^2 \cos[\lambda(x-t)] \} y(t) dt = f(x).$$

Solution:

$$y(x) = \int_a^x g(t) dt,$$

where

$$g(t) = \sqrt{\frac{\pi}{2\lambda}} \frac{1}{64\lambda^5} \left(\frac{d^2}{dt^2} + \lambda^2 \right)^6 \int_a^t (t-\tau)^{5/2} J_{5/2}[\lambda(t-\tau)] f(\tau) d\tau.$$

$$123. \int_a^x \left\{ \frac{\sin[\lambda(x-t)]}{x-t} - \lambda \cos[\lambda(x-t)] \right\} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{2\lambda^4} \left(\frac{d^2}{dx^2} + \lambda^2 \right)^3 \int_a^x \sin[\lambda(x-t)] f(t) dt.$$

$$124. \int_a^x [\sin(\lambda\sqrt{x-t}) - \lambda\sqrt{x-t} \cos(\lambda\sqrt{x-t})] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution:

$$y(x) = \frac{4}{\pi\lambda^3} \frac{d^3}{dx^3} \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$125. \int_a^x [A \tan(\lambda x) + B \cot(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tan(\lambda x)$ and $h(t) = B \cot(\mu t) + C$.

$$126. \int_a^x [A \tan^2(\lambda x) + B \cot^2(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tan^2(\lambda x)$ and $h(t) = B \cot^2(\mu t)$.

$$127. \int_a^x [\tan(\lambda x) \cot(\mu t) + \tan(\beta x) \cot(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \tan(\lambda x)$, $h_1(t) = \cot(\mu t)$, $g_2(x) = \tan(\beta x)$, and $h_2(t) = \cot(\gamma t)$.

$$128. \int_a^x [\cot(\lambda x) \tan(\mu t) + \cot(\beta x) \tan(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \cot(\lambda x)$, $h_1(t) = \tan(\mu t)$, $g_2(x) = \cot(\beta x)$, and $h_2(t) = \tan(\gamma t)$.

$$129. \int_a^x [\tan(\lambda x) \tan(\mu t) + \cot(\beta x) \cot(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = \tan(\lambda x)$, $h_1(t) = \tan(\mu t)$, $g_2(x) = \cot(\beta x)$, and $h_2(t) = \cot(\gamma t)$.

$$130. \int_a^x [A \tan^\beta(\lambda x) + B \cot^\gamma(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tan^\beta(\lambda x)$ and $h(t) = B \cot^\gamma(\mu t)$.

$$131. \int_a^x [A \cot^\beta(\lambda x) + B \tan^\gamma(\mu t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cot^\beta(\lambda x)$ and $h(t) = B \tan^\gamma(\mu t)$.

$$132. \int_a^x (Ax^\lambda \tan^\mu t + Bt^\beta \cot^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \tan^\mu t$, $g_2(x) = B \cot^\gamma x$, and $h_2(t) = t^\beta$.

$$133. \int_a^x (Ax^\lambda \cot^\mu t + Bt^\beta \tan^\gamma x) y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = \cot^\mu t$, $g_2(x) = B \tan^\gamma x$, and $h_2(t) = t^\beta$.

1.6. Equations Whose Kernels Contain Inverse Trigonometric Functions

1.6-1. Kernels Containing Arccosine.

$$1. \int_a^x [\arccos(\lambda x) - \arccos(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \arccos(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{1}{\lambda} \frac{d}{dx} \left[\sqrt{1 - \lambda^2 x^2} f'_x(x) \right].$$

$$2. \int_a^x [A \arccos(\lambda x) + B \arccos(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \arccos(\lambda x)$. For $B = -A$, see equation 1.6.1.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\arccos(\lambda x)]^{-\frac{A}{A+B}} \int_a^x [\arccos(\lambda t)]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$3. \int_a^x [A \arccos(\lambda x) + B \arccos(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \arccos(\lambda x)$ and $h(t) = B \arccos(\mu t) + C$.

$$4. \int_a^x [\arccos(\lambda x) - \arccos(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

Solution:

$$y(x) = \frac{(-1)^n}{\lambda^n n! \sqrt{1-\lambda^2 x^2}} \left(\sqrt{1-\lambda^2 x^2} \frac{d}{dx} \right)^{n+1} f(x).$$

$$5. \int_a^x \sqrt{\arccos(\lambda t) - \arccos(\lambda x)} y(t) dt = f(x).$$

This is a special case of equation 1.9.40 with $g(x) = 1 - \arccos(\lambda x)$.

Solution:

$$y(x) = \frac{2}{\pi} \varphi(x) \left(\frac{1}{\varphi(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{\varphi(t) f(t) dt}{\sqrt{\arccos(\lambda t) - \arccos(\lambda x)}}, \quad \varphi(x) = \frac{1}{\sqrt{1-\lambda^2 x^2}}.$$

$$6. \int_a^x \frac{y(t) dt}{\sqrt{\arccos(\lambda t) - \arccos(\lambda x)}} = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{\pi} \frac{d}{dx} \int_a^x \frac{\varphi(t) f(t) dt}{\sqrt{\arccos(\lambda t) - \arccos(\lambda x)}}, \quad \varphi(x) = \frac{1}{\sqrt{1-\lambda^2 x^2}}.$$

$$7. \int_a^x [\arccos(\lambda t) - \arccos(\lambda x)]^\mu y(t) dt = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = k \varphi(x) \left(\frac{1}{\varphi(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{\varphi(t) f(t) dt}{[\arccos(\lambda t) - \arccos(\lambda x)]^\mu},$$

$$\varphi(x) = \frac{1}{\sqrt{1-\lambda^2 x^2}}, \quad k = \frac{\sin(\pi \mu)}{\pi \mu}.$$

$$8. \int_a^x [\arccos^\mu(\lambda x) - \arccos^\mu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \arccos^\mu(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{1}{\lambda \mu} \frac{d}{dx} \left[\frac{f'_x(x) \sqrt{1-\lambda^2 x^2}}{\arccos^{\mu-1}(\lambda x)} \right].$$

$$9. \int_a^x \frac{y(t) dt}{[\arccos(\lambda t) - \arccos(\lambda x)]^\mu} = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = \frac{\lambda \sin(\pi\mu)}{\pi} \frac{d}{dx} \int_a^x \frac{\varphi(t)f(t) dt}{[\arccos(\lambda t) - \arccos(\lambda x)]^{1-\mu}}, \quad \varphi(x) = \frac{1}{\sqrt{1-\lambda^2x^2}}.$$

$$10. \int_a^x [A \arccos^\beta(\lambda x) + B \arccos^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \arccos^\beta(\lambda x)$ and $h(t) = B \arccos^\gamma(\mu t) + C$.

1.6-2. Kernels Containing Arcsine.

$$11. \int_a^x [\arcsin(\lambda x) - \arcsin(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \arcsin(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} \left[\sqrt{1-\lambda^2x^2} f'_x(x) \right].$$

$$12. \int_a^x [A \arcsin(\lambda x) + B \arcsin(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \arcsin(\lambda x)$. For $B = -A$, see equation 1.6.11.

Solution:

$$y(x) = \frac{\text{sign } x}{A+B} \frac{d}{dx} \left\{ |\arcsin(\lambda x)|^{-\frac{A}{A+B}} \int_a^x |\arcsin(\lambda t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$13. \int_a^x [A \arcsin(\lambda x) + B \arcsin(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \arcsin(\lambda x)$ and $h(t) = B \arcsin(\mu t) + C$.

$$14. \int_a^x [\arcsin(\lambda x) - \arcsin(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

Solution:

$$y(x) = \frac{1}{\lambda^n n! \sqrt{1-\lambda^2x^2}} \left(\sqrt{1-\lambda^2x^2} \frac{d}{dx} \right)^{n+1} f(x).$$

$$15. \int_a^x \sqrt{\arcsin(\lambda x) - \arcsin(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \varphi(x) \left(\frac{1}{\varphi(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{\varphi(t)f(t) dt}{\sqrt{\arcsin(\lambda x) - \arcsin(\lambda t)}}, \quad \varphi(x) = \frac{1}{\sqrt{1-\lambda^2x^2}}.$$

$$16. \int_a^x \frac{y(t) dt}{\sqrt{\arcsin(\lambda x) - \arcsin(\lambda t)}} = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{\pi} \frac{d}{dx} \int_a^x \frac{\varphi(t)f(t) dt}{\sqrt{\arcsin(\lambda x) - \arcsin(\lambda t)}}, \quad \varphi(x) = \frac{1}{\sqrt{1 - \lambda^2 x^2}}.$$

$$17. \int_a^x [\arcsin(\lambda x) - \arcsin(\lambda t)]^\mu y(t) dt = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = k\varphi(x) \left(\frac{1}{\varphi(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{\varphi(t)f(t) dt}{[\arcsin(\lambda x) - \arcsin(\lambda t)]^\mu},$$

$$\varphi(x) = \frac{1}{\sqrt{1 - \lambda^2 x^2}}, \quad k = \frac{\sin(\pi\mu)}{\pi\mu}.$$

$$18. \int_a^x [\arcsin^\mu(\lambda x) - \arcsin^\mu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \arcsin^\mu(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda\mu} \frac{d}{dx} \left[\frac{f'_x(x)\sqrt{1 - \lambda^2 x^2}}{\arcsin^{\mu-1}(\lambda x)} \right].$$

$$19. \int_a^x \frac{y(t) dt}{[\arcsin(\lambda x) - \arcsin(\lambda t)]^\mu} = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = \frac{\lambda \sin(\pi\mu)}{\pi} \frac{d}{dx} \int_a^x \frac{\varphi(t)f(t) dt}{[\arcsin(\lambda x) - \arcsin(\lambda t)]^{1-\mu}}, \quad \varphi(x) = \frac{1}{\sqrt{1 - \lambda^2 x^2}}.$$

$$20. \int_a^x [A \arcsin^\beta(\lambda x) + B \arcsin^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \arcsin^\beta(\lambda x)$ and $h(t) = B \arcsin^\gamma(\mu t) + C$.

$$21. \int_0^x \arcsin \sqrt{1 - \frac{t}{x}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \frac{1}{\sqrt{x}} \frac{d}{dx} \int_0^x \frac{t}{\sqrt{x-t}} \frac{d}{dt} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 452).

$$22. \int_x^\infty \arcsin \sqrt{1 - \frac{x}{t}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \frac{d}{dx} \int_x^\infty \frac{\sqrt{t}}{\sqrt{t-x}} \frac{d}{dt} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 453).

1.6-3. Kernels Containing Arctangent.

$$23. \int_a^x [\arctan(\lambda x) - \arctan(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \arctan(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda} \frac{d}{dx} [(1 + \lambda^2 x^2) f'_x(x)].$$

$$24. \int_a^x [A \arctan(\lambda x) + B \arctan(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \arctan(\lambda x)$. For $B = -A$, see equation 1.6.21.

Solution:

$$y(x) = \frac{\text{sign } x}{A + B} \frac{d}{dx} \left\{ |\arctan(\lambda x)|^{-\frac{A}{A+B}} \int_a^x |\arctan(\lambda t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$25. \int_a^x [A \arctan(\lambda x) + B \arctan(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \arctan(\lambda x)$ and $h(t) = B \arctan(\mu t) + C$.

$$26. \int_a^x [\arctan(\lambda x) - \arctan(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

Solution:

$$y(x) = \frac{1}{\lambda^n n! (1 + \lambda^2 x^2)} \left((1 + \lambda^2 x^2) \frac{d}{dx} \right)^{n+1} f(x).$$

$$27. \int_a^x \sqrt{\arctan(\lambda x) - \arctan(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \varphi(x) \left(\frac{1}{\varphi(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{\varphi(t) f(t) dt}{\sqrt{\arctan(\lambda x) - \arctan(\lambda t)}}, \quad \varphi(x) = \frac{1}{1 + \lambda^2 x^2}.$$

$$28. \int_a^x \frac{y(t) dt}{\sqrt{\arctan(\lambda x) - \arctan(\lambda t)}} = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{\pi} \frac{d}{dx} \int_a^x \frac{\varphi(t) f(t) dt}{\sqrt{\arctan(\lambda x) - \arctan(\lambda t)}}, \quad \varphi(x) = \frac{1}{1 + \lambda^2 x^2}.$$

$$29. \int_a^x \sqrt{t} \arctan \left(\sqrt{\frac{x-t}{t}} \right) y(t) dt = f(x).$$

The equation can be rewritten in terms of the Gaussian hypergeometric function in the form

$$\int_a^x (x-t)^{\gamma-1} F \left(\alpha, \beta, \gamma; 1 - \frac{x}{t} \right) y(t) dt = f(x), \quad \text{where } \alpha = \frac{1}{2}, \quad \beta = 1, \quad \gamma = \frac{3}{2}.$$

See 1.8.135 for the solution of this equation.

$$30. \int_a^x [\arctan(\lambda x) - \arctan(\lambda t)]^\mu y(t) dt = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = k\varphi(x) \left(\frac{1}{\varphi(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{\varphi(t)f(t) dt}{[\arctan(\lambda x) - \arctan(\lambda t)]^\mu},$$

$$\varphi(x) = \frac{1}{1 + \lambda^2 x^2}, \quad k = \frac{\sin(\pi\mu)}{\pi\mu}.$$

$$31. \int_a^x [\arctan^\mu(\lambda x) - \arctan^\mu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \arctan^\mu(\lambda x)$.

$$\text{Solution: } y(x) = \frac{1}{\lambda\mu} \frac{d}{dx} \left[\frac{(1 + \lambda^2 x^2) f'_x(x)}{\arctan^{\mu-1}(\lambda x)} \right].$$

$$32. \int_a^x \frac{y(t) dt}{[\arctan(\lambda x) - \arctan(\lambda t)]^\mu} = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = \frac{\lambda \sin(\pi\mu)}{\pi} \frac{d}{dx} \int_a^x \frac{\varphi(t)f(t) dt}{[\arctan(\lambda x) - \arctan(\lambda t)]^{1-\mu}}, \quad \varphi(x) = \frac{1}{1 + \lambda^2 x^2}.$$

$$33. \int_a^x [A \arctan^\beta(\lambda x) + B \arctan^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \arctan^\beta(\lambda x)$ and $h(t) = B \arctan^\gamma(\mu t) + C$.

1.6-4. Kernels Containing Arccotangent.

$$34. \int_a^x [\operatorname{arccot}(\lambda x) - \operatorname{arccot}(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \operatorname{arccot}(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{1}{\lambda} \frac{d}{dx} [(1 + \lambda^2 x^2) f'_x(x)].$$

$$35. \int_a^x [A \operatorname{arccot}(\lambda x) + B \operatorname{arccot}(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.4 with $g(x) = \operatorname{arccot}(\lambda x)$. For $B = -A$, see equation 1.6.34.

Solution:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [\operatorname{arccot}(\lambda x)]^{-\frac{A}{A+B}} \int_a^x [\operatorname{arccot}(\lambda t)]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$36. \int_a^x [A \operatorname{arccot}(\lambda x) + B \operatorname{arccot}(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \operatorname{arccot}(\lambda x)$ and $h(t) = B \operatorname{arccot}(\mu t) + C$.

$$37. \int_a^x [\operatorname{arccot}(\lambda x) - \operatorname{arccot}(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

Solution:

$$y(x) = \frac{(-1)^n}{\lambda^n n! (1 + \lambda^2 x^2)} \left((1 + \lambda^2 x^2) \frac{d}{dx} \right)^{n+1} f(x).$$

$$38. \int_a^x \sqrt{\operatorname{arccot}(\lambda t) - \operatorname{arccot}(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \varphi(x) \left(\frac{1}{\varphi(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{\varphi(t) f(t) dt}{\sqrt{\operatorname{arccot}(\lambda t) - \operatorname{arccot}(\lambda x)}}, \quad \varphi(x) = \frac{1}{1 + \lambda^2 x^2}.$$

$$39. \int_a^x \frac{y(t) dt}{\sqrt{\operatorname{arccot}(\lambda t) - \operatorname{arccot}(\lambda x)}} = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{\pi} \frac{d}{dx} \int_a^x \frac{\varphi(t) f(t) dt}{\sqrt{\operatorname{arccot}(\lambda t) - \operatorname{arccot}(\lambda x)}}, \quad \varphi(x) = \frac{1}{1 + \lambda^2 x^2}.$$

$$40. \int_a^x [\operatorname{arccot}(\lambda t) - \operatorname{arccot}(\lambda x)]^\mu y(t) dt = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = k \varphi(x) \left(\frac{1}{\varphi(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{\varphi(t) f(t) dt}{[\operatorname{arccot}(\lambda t) - \operatorname{arccot}(\lambda x)]^\mu},$$

$$\varphi(x) = \frac{1}{1 + \lambda^2 x^2}, \quad k = \frac{\sin(\pi \mu)}{\pi \mu}.$$

$$41. \int_a^x [\operatorname{arccot}^\mu(\lambda x) - \operatorname{arccot}^\mu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = \operatorname{arccot}^\mu(\lambda x)$.

$$\text{Solution: } y(x) = -\frac{1}{\lambda \mu} \frac{d}{dx} \left[\frac{(1 + \lambda^2 x^2) f'_x(x)}{\operatorname{arccot}^{\mu-1}(\lambda x)} \right].$$

$$42. \int_a^x \frac{y(t) dt}{[\operatorname{arccot}(\lambda t) - \operatorname{arccot}(\lambda x)]^\mu} = f(x), \quad 0 < \mu < 1.$$

Solution:

$$y(x) = \frac{\lambda \sin(\pi \mu)}{\pi} \frac{d}{dx} \int_a^x \frac{\varphi(t) f(t) dt}{[\operatorname{arccot}(\lambda t) - \operatorname{arccot}(\lambda x)]^{1-\mu}}, \quad \varphi(x) = \frac{1}{1 + \lambda^2 x^2}.$$

$$43. \int_a^x [A \operatorname{arccot}^\beta(\lambda x) + B \operatorname{arccot}^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \operatorname{arccot}^\beta(\lambda x)$ and $h(t) = B \operatorname{arccot}^\gamma(\mu t) + C$.

1.7. Equations Whose Kernels Contain Combinations of Elementary Functions

1.7-1. Kernels Containing Exponential and Hyperbolic Functions.

$$1. \quad \int_a^x e^{\mu(x-t)} \{A_1 \cosh[\lambda_1(x-t)] + A_2 \cosh[\lambda_2(x-t)]\} y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.8:

$$\int_a^x \{A_1 \cosh[\lambda_1(x-t)] + A_2 \cosh[\lambda_2(x-t)]\} w(t) dt = e^{-\mu x} f(x).$$

$$2. \quad \int_a^x e^{\mu(x-t)} \cosh^2[\lambda(x-t)] y(t) dt = f(x).$$

Solution:

$$y(x) = \varphi(x) - \frac{2\lambda^2}{k} \int_a^x e^{\mu(x-t)} \sinh[k(x-t)] \varphi(x) dt, \quad k = \lambda\sqrt{2}, \quad \varphi(x) = f'_x(x) - \mu f(x).$$

$$3. \quad \int_a^x e^{\mu(x-t)} \cosh^3[\lambda(x-t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.15:

$$\int_a^x \cosh^3[\lambda(x-t)] w(t) dt = e^{-\mu x} f(x).$$

$$4. \quad \int_a^x e^{\mu(x-t)} \cosh^4[\lambda(x-t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.19:

$$\int_a^x \cosh^4[\lambda(x-t)] w(t) dt = e^{-\mu x} f(x).$$

$$5. \quad \int_a^x e^{\mu(x-t)} [\cosh(\lambda x) - \cosh(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

Solution:

$$y(x) = \frac{1}{\lambda^n n!} e^{\mu x} \sinh(\lambda x) \left[\frac{1}{\sinh(\lambda x)} \frac{d}{dx} \right]^{n+1} F_\mu(x), \quad F_\mu(x) = e^{-\mu x} f(x).$$

$$6. \quad \int_a^x e^{\mu(x-t)} \sqrt{\cosh x - \cosh t} y(t) dt = f(x), \quad f(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi} e^{\mu x} \sinh x \left(\frac{1}{\sinh x} \frac{d}{dx} \right)^2 \int_a^x \frac{e^{-\mu t} \sinh t f(t) dt}{\sqrt{\cosh x - \cosh t}}.$$

$$7. \quad \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{\sqrt{\cosh x - \cosh t}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} e^{\mu x} \frac{d}{dx} \int_a^x \frac{e^{-\mu t} \sinh t f(t) dt}{\sqrt{\cosh x - \cosh t}}.$$

$$8. \int_a^x e^{\mu(x-t)} (\cosh x - \cosh t)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.23:

$$\int_a^x (\cosh x - \cosh t)^\lambda w(t) dt = e^{-\mu x} f(x).$$

$$9. \int_a^x [Ae^{\mu(x-t)} + B \cosh^\lambda x] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \cosh^\lambda x$, and $h_2(t) = 1$.

$$10. \int_a^x [Ae^{\mu(x-t)} + B \cosh^\lambda t] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \cosh^\lambda t$.

$$11. \int_a^x e^{\mu(x-t)} (\cosh^\lambda x - \cosh^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.24:

$$\int_a^x (\cosh^\lambda x - \cosh^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$12. \int_a^x e^{\mu(x-t)} (A \cosh^\lambda x + B \cosh^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.25:

$$\int_a^x (A \cosh^\lambda x + B \cosh^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$13. \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{(\cosh x - \cosh t)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi} e^{\mu x} \frac{d}{dx} \int_a^x \frac{e^{-\mu t} \sinh t f(t) dt}{(\cosh x - \cosh t)^{1-\lambda}}.$$

$$14. \int_a^x e^{\mu(x-t)} \{A_1 \sinh[\lambda_1(x-t)] + A_2 \sinh[\lambda_2(x-t)]\} y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.49:

$$\int_a^x \{A_1 \sinh[\lambda_1(x-t)] + A_2 \sinh[\lambda_2(x-t)]\} w(t) dt = e^{-\mu x} f(x).$$

$$15. \int_a^x e^{\mu(x-t)} \sinh^2[\lambda(x-t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.51:

$$\int_a^x \sinh^2[\lambda(x-t)] w(t) dt = e^{-\mu x} f(x).$$

$$16. \int_a^x e^{\mu(x-t)} \sinh^3[\lambda(x-t)]y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.3.57:

$$\int_a^x \sinh^3[\lambda(x-t)]w(t) dt = e^{-\mu x} f(x).$$

$$17. \int_a^x e^{\mu(x-t)} \sinh^n[\lambda(x-t)]y(t) dt = f(x), \quad n = 2, 3, \dots$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.3.62:

$$\int_a^x \sinh^n[\lambda(x-t)]w(t) dt = e^{-\mu x} f(x).$$

$$18. \int_a^x e^{\mu(x-t)} \sinh(k\sqrt{x-t})y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi k} e^{\mu x} \frac{d^2}{dx^2} \int_a^x \frac{e^{-\mu t} \cos(k\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$19. \int_a^x e^{\mu(x-t)} \sqrt{\sinh x - \sinh t} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} e^{\mu x} \cosh x \left(\frac{1}{\cosh x} \frac{d}{dx} \right)^2 \int_a^x \frac{e^{-\mu t} \cosh t f(t) dt}{\sqrt{\sinh x - \sinh t}}.$$

$$20. \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{\sqrt{\sinh x - \sinh t}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} e^{\mu x} \frac{d}{dx} \int_a^x \frac{e^{-\mu t} \cosh t f(t) dt}{\sqrt{\sinh x - \sinh t}}.$$

$$21. \int_a^x e^{\mu(x-t)} (\sinh x - \sinh t)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.3.67:

$$\int_a^x (\sinh x - \sinh t)^\lambda w(t) dt = e^{-\mu x} f(x).$$

$$22. \int_a^x e^{\mu(x-t)} (\sinh^\lambda x - \sinh^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.3.68:

$$\int_a^x (\sinh^\lambda x - \sinh^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$23. \int_a^x e^{\mu(x-t)} (A \sinh^\lambda x + B \sinh^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.3.69:

$$\int_a^x (A \sinh^\lambda x + B \sinh^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$24. \int_a^x [Ae^{\mu(x-t)} + B \sinh^\lambda x] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \sinh^\lambda x$, and $h_2(t) = 1$.

$$25. \int_a^x [Ae^{\mu(x-t)} + B \sinh^\lambda t] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \sinh^\lambda t$.

$$26. \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{(\sinh x - \sinh t)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi} e^{\mu x} \frac{d}{dx} \int_a^x \frac{e^{-\mu t} \cosh t f(t) dt}{(\sinh x - \sinh t)^{1-\lambda}}.$$

$$27. \int_a^x e^{\mu(x-t)} (A \tanh^\lambda x + B \tanh^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.89:

$$\int_a^x (A \tanh^\lambda x + B \tanh^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$28. \int_a^x e^{\mu(x-t)} (A \tanh^\lambda x + B \tanh^\beta t + C) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.9.6 with $g(x) = A \tanh^\lambda x$, $h(t) = B \tanh^\beta t + C$:

$$\int_a^x (A \tanh^\lambda x + B \tanh^\beta t + C) w(t) dt = e^{-\mu x} f(x).$$

$$29. \int_a^x [Ae^{\mu(x-t)} + B \tanh^\lambda x] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \tanh^\lambda x$, and $h_2(t) = 1$.

$$30. \int_a^x [Ae^{\mu(x-t)} + B \tanh^\lambda t] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \tanh^\lambda t$.

$$31. \int_a^x e^{\mu(x-t)} (A \coth^\lambda x + B \coth^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.3.102:

$$\int_a^x (A \coth^\lambda x + B \coth^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$32. \int_a^x e^{\mu(x-t)} (A \coth^\lambda x + B \coth^\beta t + C) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.9.6 with $g(x) = A \coth^\lambda x$, $h(t) = B \coth^\beta t + C$:

$$\int_a^x (A \coth^\lambda x + B \coth^\beta t + C) w(t) dt = e^{-\mu x} f(x).$$

$$33. \int_a^x [A e^{\mu(x-t)} + B \coth^\lambda x] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A e^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \coth^\lambda x$, and $h_2(t) = 1$.

$$34. \int_a^x [A e^{\mu(x-t)} + B \coth^\lambda t] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A e^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \coth^\lambda t$.

1.7-2. Kernels Containing Exponential and Logarithmic Functions.

$$35. \int_a^x e^{\lambda(x-t)} (\ln x - \ln t) y(t) dt = f(x).$$

Solution:

$$y(x) = e^{\lambda x} [x \varphi''_{xx}(x) + \varphi'_x(x)], \quad \varphi(x) = e^{-\lambda x} f(x).$$

$$36. \int_0^x e^{\lambda(x-t)} \ln(x-t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 1.4.2:

$$\int_0^x \ln(x-t) w(t) dt = e^{-\lambda x} f(x).$$

$$37. \int_a^x e^{\lambda(x-t)} (A \ln x + B \ln t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 1.4.4:

$$\int_a^x (A \ln x + B \ln t) w(t) dt = e^{-\lambda x} f(x).$$

$$38. \int_a^x e^{\mu(x-t)} [A \ln^2(\lambda x) + B \ln^2(\lambda t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 1.4.7:

$$\int_a^x [A \ln^2(\lambda x) + B \ln^2(\lambda t)] w(t) dt = e^{-\lambda x} f(x).$$

$$39. \int_a^x e^{\lambda(x-t)} [\ln(x/t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

Solution:

$$y(x) = \frac{1}{n!} e^{\lambda x} \left(x \frac{d}{dx} \right)^{n+1} F_\lambda(x), \quad F_\lambda(x) = e^{-\lambda x} f(x).$$

$$40. \int_a^x e^{\lambda(x-t)} \sqrt{\ln(x/t)} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2e^{\lambda x}}{\pi x} \left(x \frac{d}{dx} \right)^2 \int_a^x \frac{e^{-\lambda t} f(t) dt}{t \sqrt{\ln(x/t)}}.$$

$$41. \int_a^x \frac{e^{\lambda(x-t)}}{\sqrt{\ln(x/t)}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} e^{\lambda x} \frac{d}{dx} \int_a^x \frac{e^{-\lambda t} f(t) dt}{t \sqrt{\ln(x/t)}}.$$

$$42. \int_a^x [Ae^{\mu(x-t)} + B \ln^\nu(\lambda x)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \ln^\nu(\lambda x)$, and $h_2(t) = 1$.

$$43. \int_a^x [Ae^{\mu(x-t)} + B \ln^\nu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \ln^\nu(\lambda t)$.

$$44. \int_a^x e^{\mu(x-t)} [\ln(x/t)]^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.4.16:

$$\int_a^x [\ln(x/t)]^\lambda w(t) dt = e^{-\mu x} f(x).$$

$$45. \int_a^x \frac{e^{\mu(x-t)}}{[\ln(x/t)]^\lambda} y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi} e^{\mu x} \frac{d}{dx} \int_a^x \frac{f(t) dt}{t e^{\mu t} [\ln(x/t)]^{1-\lambda}}.$$

1.7-3. Kernels Containing Exponential and Trigonometric Functions.

$$46. \int_a^x e^{\mu(x-t)} \cos[\lambda(x-t)] y(t) dt = f(x).$$

Solution: $y(x) = f'_x(x) - \mu f(x) + \lambda^2 \int_a^x e^{\mu(x-t)} f(t) dt.$

$$47. \int_a^x e^{\mu(x-t)} \{A_1 \cos[\lambda_1(x-t)] + A_2 \cos[\lambda_2(x-t)]\} y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.8:

$$\int_a^x \{A_1 \cos[\lambda_1(x-t)] + A_2 \cos[\lambda_2(x-t)]\} w(t) dt = e^{-\mu x} f(x).$$

$$48. \int_a^x e^{\mu(x-t)} \cos^2[\lambda(x-t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.9.

Solution:

$$y(x) = \varphi(x) + \frac{2\lambda^2}{k} \int_a^x e^{\mu(x-t)} \sin[k(x-t)] \varphi(t) dt, \quad k = \lambda\sqrt{2}, \quad \varphi(x) = f'_x(x) - \mu f(x).$$

$$49. \int_a^x e^{\mu(x-t)} \cos^3[\lambda(x-t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.16:

$$\int_a^x \cos^3[\lambda(x-t)] w(t) dt = e^{-\mu x} f(x).$$

$$50. \int_a^x e^{\mu(x-t)} \cos^4[\lambda(x-t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.20:

$$\int_a^x \cos^4[\lambda(x-t)] w(t) dt = e^{-\mu x} f(x).$$

$$51. \int_a^x e^{\mu(x-t)} [\cos(\lambda x) - \cos(\lambda t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

Solution:

$$y(x) = \frac{(-1)^n}{\lambda^n n!} e^{\mu x} \sin(\lambda x) \left[\frac{1}{\sin(\lambda x)} \frac{d}{dx} \right]^{n+1} F_\mu(x), \quad F_\mu(x) = e^{-\mu x} f(x).$$

$$52. \int_a^x e^{\mu(x-t)} \sqrt{\cos t - \cos x} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} e^{\mu x} \sin x \left(\frac{1}{\sin x} \frac{d}{dx} \right)^2 \int_a^x \frac{e^{-\mu t} \sin t f(t) dt}{\sqrt{\cos t - \cos x}}.$$

$$53. \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{\sqrt{\cos t - \cos x}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} e^{\mu x} \frac{d}{dx} \int_a^x \frac{e^{-\mu t} \sin t f(t) dt}{\sqrt{\cos t - \cos x}}.$$

$$54. \int_a^x e^{\mu(x-t)} (\cos t - \cos x)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k e^{\mu x} \sin x \left(\frac{1}{\sin x} \frac{d}{dx} \right)^2 \int_a^x \frac{e^{-\mu t} \sin t f(t) dt}{(\cos t - \cos x)^\lambda}, \quad k = \frac{\sin(\pi \lambda)}{\pi \lambda}.$$

$$55. \int_a^x e^{\mu(x-t)} (\cos^\lambda x - \cos^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.25:

$$\int_a^x (\cos^\lambda x - \cos^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$56. \int_a^x e^{\mu(x-t)} (A \cos^\lambda x + B \cos^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.26:

$$\int_a^x (A \cos^\lambda x + B \cos^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$57. \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{(\cos t - \cos x)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.27:

$$\int_a^x \frac{w(t) dt}{(\cos t - \cos x)^\lambda} = e^{-\mu x} f(x).$$

$$58. \int_a^x [A e^{\mu(x-t)} + B \cos^\nu(\lambda x)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A e^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \cos^\nu(\lambda x)$, and $h_2(t) = 1$.

$$59. \int_a^x [A e^{\mu(x-t)} + B \cos^\nu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A e^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \cos^\nu(\lambda t)$.

$$60. \int_a^x e^{\mu(x-t)} \sin[\lambda(x-t)] y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

Solution: $y(x) = \frac{1}{\lambda} [f''_{xx}(x) - 2\mu f'_x(x) + (\lambda^2 + \mu^2) f(x)]$.

$$61. \int_a^x e^{\mu(x-t)} \{A_1 \sin[\lambda_1(x-t)] + A_2 \sin[\lambda_2(x-t)]\} y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.48:

$$\int_a^x \{A_1 \sin[\lambda_1(x-t)] + A_2 \sin[\lambda_2(x-t)]\} w(t) dt = e^{-\mu x} f(x).$$

$$62. \int_a^x e^{\mu(x-t)} \sin^2[\lambda(x-t)]y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.5.50:

$$\int_a^x \sin^2[\lambda(x-t)]w(t) dt = e^{-\mu x} f(x).$$

$$63. \int_a^x e^{\mu(x-t)} \sin^3[\lambda(x-t)]y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.5.57:

$$\int_a^x \sin^3[\lambda(x-t)]w(t) dt = e^{-\mu x} f(x).$$

$$64. \int_a^x e^{\mu(x-t)} \sin^n[\lambda(x-t)]y(t) dt = f(x), \quad n = 2, 3, \dots$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.5.62:

$$\int_a^x \sin^n[\lambda(x-t)]w(t) dt = e^{-\mu x} f(x).$$

$$65. \int_a^x e^{\mu(x-t)} \sin(k\sqrt{x-t})y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi k} e^{\mu x} \frac{d^2}{dx^2} \int_a^x \frac{e^{-\mu t} \cosh(k\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$66. \int_a^x e^{\mu(x-t)} \sqrt{\sin x - \sin t} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} e^{\mu x} \cos x \left(\frac{1}{\cos x} \frac{d}{dx} \right)^2 \int_a^x \frac{e^{-\mu t} \cos t f(t) dt}{\sqrt{\sin x - \sin t}}.$$

$$67. \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{\sqrt{\sin x - \sin t}} = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} e^{\mu x} \frac{d}{dx} \int_a^x \frac{e^{-\mu t} \cos t f(t) dt}{\sqrt{\sin x - \sin t}}.$$

$$68. \int_a^x e^{\mu(x-t)} (\sin x - \sin t)^\lambda y(t) dt = f(x), \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k e^{\mu x} \cos x \left(\frac{1}{\cos x} \frac{d}{dx} \right)^2 \int_a^x \frac{e^{-\mu t} \cos t f(t) dt}{(\sin x - \sin t)^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$69. \int_a^x e^{\mu(x-t)} (\sin^\lambda x - \sin^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x}y(x)$ leads to an equation of the form 1.5.72:

$$\int_a^x (\sin^\lambda x - \sin^\lambda t)w(t) dt = e^{-\mu x} f(x).$$

$$70. \int_a^x e^{\mu(x-t)} (A \sin^\lambda x + B \sin^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.9.4 with $g(x) = \sin^\lambda x$:

$$\int_a^x (A \sin^\lambda x + B \sin^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$71. \int_a^x \frac{e^{\mu(x-t)} y(t) dt}{(\sin x - \sin t)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.74:

$$\int_a^x \frac{w(t) dt}{(\sin x - \sin t)^\lambda} = e^{-\mu x} f(x).$$

$$72. \int_a^x [Ae^{\mu(x-t)} + B \sin^\nu(\lambda x)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \sin^\nu(\lambda x)$, and $h_2(t) = 1$.

$$73. \int_a^x [Ae^{\mu(x-t)} + B \sin^\nu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \sin^\nu(\lambda t)$.

$$74. \int_a^x e^{\mu(x-t)} (A \tan^\lambda x + B \tan^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.91:

$$\int_a^x (A \tan^\lambda x + B \tan^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$75. \int_a^x e^{\mu(x-t)} (A \tan^\lambda x + B \tan^\beta t + C) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.9.6:

$$\int_a^x (A \tan^\lambda x + B \tan^\beta t + C) w(t) dt = e^{-\mu x} f(x).$$

$$76. \int_a^x [Ae^{\mu(x-t)} + B \tan^\nu(\lambda x)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \tan^\nu(\lambda x)$, and $h_2(t) = 1$.

$$77. \int_a^x [Ae^{\mu(x-t)} + B \tan^\nu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ae^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \tan^\nu(\lambda t)$.

$$78. \int_a^x e^{\mu(x-t)} (A \cot^\lambda x + B \cot^\lambda t) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.5.104:

$$\int_a^x (A \cot^\lambda x + B \cot^\lambda t) w(t) dt = e^{-\mu x} f(x).$$

$$79. \int_a^x e^{\mu(x-t)} (A \cot^\lambda x + B \cot^\beta t + C) y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 1.9.6:

$$\int_a^x (A \cot^\lambda x + B \cot^\beta t + C) w(t) dt = e^{-\mu x} f(x).$$

$$80. \int_a^x [A e^{\mu(x-t)} + B \cot^\nu(\lambda x)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A e^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B \cot^\nu(\lambda x)$, and $h_2(t) = 1$.

$$81. \int_a^x [A e^{\mu(x-t)} + B \cot^\nu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = A e^{\mu x}$, $h_1(t) = e^{-\mu t}$, $g_2(x) = B$, and $h_2(t) = \cot^\nu(\lambda t)$.

1.7-4. Kernels Containing Hyperbolic and Logarithmic Functions.

$$82. \int_a^x [A \cosh^\beta(\lambda x) + B \ln^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh^\beta(\lambda x)$ and $h(t) = B \ln^\gamma(\mu t) + C$.

$$83. \int_a^x [A \cosh^\beta(\lambda t) + B \ln^\gamma(\mu x) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = B \ln^\gamma(\mu x) + C$ and $h(t) = A \cosh^\beta(\lambda t)$.

$$84. \int_a^x [A \sinh^\beta(\lambda x) + B \ln^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sinh^\beta(\lambda x)$ and $h(t) = B \ln^\gamma(\mu t) + C$.

$$85. \int_a^x [A \sinh^\beta(\lambda t) + B \ln^\gamma(\mu x) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = B \ln^\gamma(\mu x)$ and $h(t) = A \sinh^\beta(\lambda t) + C$.

$$86. \int_a^x [A \tanh^\beta(\lambda x) + B \ln^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh^\beta(\lambda x)$ and $h(t) = B \ln^\gamma(\mu t) + C$.

$$87. \int_a^x [A \tanh^\beta(\lambda t) + B \ln^\gamma(\mu x) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = B \ln^\gamma(\mu x)$ and $h(t) = A \tanh^\beta(\lambda t) + C$.

$$88. \int_a^x [A \coth^\beta(\lambda x) + B \ln^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \coth^\beta(\lambda x)$ and $h(t) = B \ln^\gamma(\mu t) + C$.

$$89. \int_a^x [A \coth^\beta(\lambda t) + B \ln^\gamma(\mu x) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = B \ln^\gamma(\mu x)$ and $h(t) = A \coth^\beta(\lambda t) + C$.

1.7-5. Kernels Containing Hyperbolic and Trigonometric Functions.

$$90. \int_a^x [A \cosh^\beta(\lambda x) + B \cos^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh^\beta(\lambda x)$ and $h(t) = B \cos^\gamma(\mu t) + C$.

$$91. \int_a^x [A \cosh^\beta(\lambda t) + B \sin^\gamma(\mu x) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = B \sin^\gamma(\mu x) + C$ and $h(t) = A \cosh^\beta(\lambda t)$.

$$92. \int_a^x [A \cosh^\beta(\lambda x) + B \tan^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cosh^\beta(\lambda x)$ and $h(t) = B \tan^\gamma(\mu t) + C$.

$$93. \int_a^x [A \sinh^\beta(\lambda x) + B \cos^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sinh^\beta(\lambda x)$ and $h(t) = B \cos^\gamma(\mu t) + C$.

$$94. \int_a^x [A \sinh^\beta(\lambda t) + B \sin^\gamma(\mu x) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = B \sin^\gamma(\mu x)$ and $h(t) = A \sinh^\beta(\lambda t) + C$.

$$95. \int_a^x [A \sinh^\beta(\lambda x) + B \tan^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sinh^\beta(\lambda x)$ and $h(t) = B \tan^\gamma(\mu t) + C$.

$$96. \int_a^x [A \tanh^\beta(\lambda x) + B \cos^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh^\beta(\lambda x)$ and $h(t) = B \cos^\gamma(\mu t) + C$.

$$97. \int_a^x [A \tanh^\beta(\lambda x) + B \sin^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \tanh^\beta(\lambda x)$ and $h(t) = B \sin^\gamma(\mu t) + C$.

$$98. \int_a^x \sinh[\lambda(x-t)] - \sin[\lambda(x-t)]y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{1}{2\lambda^3} \left(\frac{d^4}{dx^4} - \lambda^4 \right) f(x).$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 449).

$$99. \int_a^x \sinh[\lambda(x-t)] \sin[\lambda(x-t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = f''_{xx}(a) = 0.$$

Solution:

$$y(x) = \frac{1}{2\lambda^2} \left(\frac{d^4}{dx^4} + 4\lambda^4 \right) \int_a^x f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 449).

$$100. \int_a^x \sinh[\lambda(x-t)] \cos[\lambda(x-t)]y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{1}{\sqrt{2}\lambda^2} \left(\frac{d^4}{dx^4} + 4\lambda^4 \right) \int_a^x \sinh[\sqrt{2}\lambda(x-t)]f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 449).

$$101. \int_a^x \cosh[\lambda(x-t)] \sin[\lambda(x-t)]y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$.

Solution:

$$y(x) = \frac{1}{\sqrt{2}\lambda^2} \left(\frac{d^4}{dx^4} + 4\lambda^4 \right) \int_a^x \sin[\sqrt{2}\lambda(x-t)]f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 450).

$$102. \int_a^x \cosh[\lambda(x-t)] \cos[\lambda(x-t)]y(t) dt = f(x), \quad f(a) = 0.$$

Solution:

$$y(x) = \frac{1}{2} \left(\frac{d^4}{dx^4} + 4\lambda^4 \right) \int_a^x (x-t)^2 f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 450).

1.7-6. Kernels Containing Logarithmic and Trigonometric Functions.

$$103. \int_a^x [A \cos^\beta(\lambda x) + B \ln^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \cos^\beta(\lambda x)$ and $h(t) = B \ln^\gamma(\mu t) + C$.

$$104. \int_a^x [A \cos^\beta(\lambda t) + B \ln^\gamma(\mu x) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = B \ln^\gamma(\mu x) + C$ and $h(t) = A \cos^\beta(\lambda t)$.

$$105. \int_a^x [A \sin^\beta(\lambda x) + B \ln^\gamma(\mu t) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A \sin^\beta(\lambda x)$ and $h(t) = B \ln^\gamma(\mu t) + C$.

$$106. \int_a^x [A \sin^\beta(\lambda t) + B \ln^\gamma(\mu x) + C] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = B \ln^\gamma(\mu x)$ and $h(t) = A \sin^\beta(\lambda t) + C$.

1.8. Equations Whose Kernels Contain Special Functions*

1.8-1. Kernels Containing Error Function or Exponential Integral.

$$1. \int_0^x \operatorname{erf}(\lambda \sqrt{x-t}) y(t) dt = f(x), \quad f(0) = f'_x(0) = 0.$$

Here $\operatorname{erf} z$ is the error function (see Supplement 11.2-1).

Solution:

$$y(x) = \frac{1}{\sqrt{\pi} \lambda} e^{-\lambda^2 x} \frac{d}{dx} \int_0^x \frac{e^{\lambda^2 t}}{\sqrt{x-t}} f'_t(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 458).

$$2. \int_x^\infty \operatorname{erf}(\lambda \sqrt{t-x}) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\sqrt{\pi} \lambda} e^{\lambda^2 x} \frac{d}{dx} \int_x^\infty \frac{1}{e^{\lambda^2 t} \sqrt{t-x}} f'_t(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 459).

$$3. \int_0^x \operatorname{Ei}(\lambda(t-x)) y(t) dt = f(x), \quad f(0) = f'_x(0) = 0.$$

Here $\operatorname{Ei}(z)$ is the exponential integral (see Supplement 11.2-2).

Solution:

$$y(x) = -\frac{1}{\lambda} \int_0^x e^{\lambda(t-x)} \nu(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda \frac{d}{dt} \right) f(t) dt,$$

$$\text{where } \nu(z) = \int_0^\infty \frac{z^\xi d\xi}{\Gamma(\xi+1)}.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 455).

* For notation and properties of special functions, see Supplement 11.

1.8-2. Kernels Containing Sine and Cosine Integrals.

$$4. \quad \int_0^x [\sin(x-t) \text{Si}(x-t) - \cos(x-t) \text{ci}(x-t)]y(t) dt = f(x), \quad f(0) = f'_x(0) = 0.$$

Here $\text{Si}(z)$ is the sine integral and $\text{ci}(z)$ is the cosine integral (see Supplements 11.3-1 and 11.3-2).

Solution:

$$y(x) = \int_0^x \nu(x-t) \left(\frac{d^2}{dt^2} + 1 \right) f(t) dt,$$

$$\text{where } \nu(z) = \int_0^\infty \frac{z^\xi d\xi}{\Gamma(\xi+1)}.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 458).

$$5. \quad \int_0^x [\cos(x-t) \text{Si}(x-t) - \sin(x-t) \text{ci}(x-t)]y(t) dt = f(x), \quad f(0) = f'_x(0) = f''_{xx}(0) = 0.$$

Solution:

$$y(x) = \int_0^x \nu(x-t) \left(\frac{d^3}{dt^3} + \frac{d}{dt} \right) f(t) dt,$$

$$\text{where } \nu(z) = \int_0^\infty \frac{z^\xi d\xi}{\Gamma(\xi+1)}.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 458).

1.8-3. Kernels Containing Fresnel Integrals.

$$6. \quad \int_0^x S(x-t)y(t) dt = f(x), \quad f(0) = f'_x(0) = f''_{xx}(0) = f'''_{xxx}(0) = 0.$$

Here $S(z)$ is the Fresnel sine integral (see Supplement 11.3-3).

Solution:

$$y(x) = 4 \int_0^x C(x-t) \left(\frac{d^4}{dt^4} + \frac{d^2}{dt^2} \right) y(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 460).

$$7. \quad \int_0^x C(x-t)y(t) dt = f(x), \quad f(0) = f'_x(0) = f''_{xx}(0) = f'''_{xxx}(0) = 0.$$

Here $C(z)$ is the Fresnel cosine integral (see Supplement 11.3-3).

Solution:

$$y(x) = 4 \int_0^x S(x-t) \left(\frac{d^4}{dt^4} + \frac{d^2}{dt^2} \right) y(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 460).

1.8-4. Kernels Containing Incomplete Gamma Functions.

$$8. \int_0^x \gamma(\nu, \lambda(x-t))y(t) dt = f(x).$$

Here $\gamma(\nu, z)$ is the incomplete gamma function (see Supplement 11.5-1).

1°. Let $\operatorname{Re} \nu > 0$, $m = [\operatorname{Re} \nu] + 1$, where $[\operatorname{Re} \nu]$ denotes the integer part of the number $\operatorname{Re} \nu$, and $f(0) = f'_x(0) = \dots = f_x^{(m)}(0) = 0$. Then the solution is

$$y(x) = \frac{\lambda^{-\nu}}{\Gamma(\nu)\Gamma(m-\nu)} e^{-\lambda x} \left(\frac{d}{dx}\right)^m \int_0^x \frac{e^{-\lambda t}}{(x-t)^{\nu-m+1}} f'_t(t) dt.$$

2°. Let $\nu = n/2$, where n is a positive integer, and $f(0) = f'(0) = \dots = f^{(n+1)}(0) = 0$. Then the solution is

$$y(x) = \frac{\lambda^{-n}}{\Gamma^2(n/2)} \int_0^x \gamma\left(\frac{n}{2}, \lambda(x-t)\right) \frac{d^2}{dt^2} \left(\frac{d}{dt} + \lambda\right)^n f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 461).

$$9. \int_x^\infty \gamma(\nu, \lambda(t-x))y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{\lambda^{-\nu}}{\Gamma(\nu)\Gamma(m-\nu)} e^{\lambda x} \left(-\frac{d}{dx}\right)^m \int_x^\infty \frac{e^{-\lambda t}}{(t-x)^{\nu-m+1}} f'_t(t) dt,$$

where $\operatorname{Re} \nu > 0$, $m = [\operatorname{Re} \nu] + 1$, and $[\operatorname{Re} \nu]$ denotes the integer part of the number $\operatorname{Re} \nu$.

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 462).

$$10. \int_0^x \Gamma(\nu, \lambda(x-t))y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{e^{-\lambda x}}{\Gamma(\nu)} \int_0^x E_\nu([\lambda(x-t)]^\nu) \left(\frac{d^2}{dt^2} - \lambda \frac{d}{dt}\right) (e^{\lambda t} f(t)) dt,$$

where $\operatorname{Re} \nu > 0$ and $E_\nu(z)$ are the Weber function,

$$E_\nu(z) = \frac{1}{\pi} \int_0^\pi \sin(\nu t - z \sin t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 462).

1.8-5. Kernels Containing Bessel Functions.

$$11. \int_a^x J_0(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.17 with $n = 0$ and $J_0(z)$ is the Bessel function (see Supplement 11.6-1). If $f(a) = f'_x(a) = 0$ then the solution is

$$y(x) = \int_a^x J_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2\right) f(t) dt.$$

Example. In the special case $\lambda = 1$ and $f(x) = A \sin x$, the solution has the form $y(x) = A J_0(x)$.

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 470).

* For notation and properties of special functions, see Supplement 11.

$$12. \int_a^x [J_0(\lambda x) - J_0(\lambda t)]y(t) dt = f(x).$$

$$\text{Solution: } y(x) = -\frac{d}{dx} \left[\frac{f'_x(x)}{\lambda J_1(\lambda x)} \right].$$

$$13. \int_a^x [AJ_0(\lambda x) + BJ_0(\lambda t)]y(t) dt = f(x).$$

For $B = -A$, see equation 1.8.12. We consider the interval $[a, x]$ in which $J_0(\lambda x)$ does not change its sign.

Solution with $B \neq -A$:

$$y(x) = \pm \frac{1}{A+B} \frac{d}{dx} \left\{ |J_0(\lambda x)|^{-\frac{A}{A+B}} \int_a^x |J_0(\lambda t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

Here the sign of $J_0(\lambda x)$ should be taken.

$$14. \int_a^x (x-t) J_0(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.18 with $n = 0$. If $f(a) = f'_x(a) = f''_{xx}(a) = 0$ then the solution is

$$y(x) = \int_a^x J_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2 \right)^2 F(t) dt, \quad F(t) = \int_a^t f(s) ds.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$15. \int_a^x (x-t) J_1(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.17 with $n = 1$. If $f(a) = f'_x(a) = 0$ then the solution is

$$y(x) = \frac{f'_x(x)}{\lambda} + \frac{1}{\lambda} \int_a^x J_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2 \right) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 471).

$$16. \int_a^x (x-t)^2 J_1(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.18 with $n = 1$. If $f(a) = f'_x(a) = \dots = f''''_{xxxx}(a) = 0$ then the solution is

$$y(x) = \frac{1}{3\lambda} \int_a^x J_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2 \right)^3 F(t) dt, \quad F(t) = \int_a^t f(s) ds.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$17. \int_a^x (x-t)^n J_n(\lambda(x-t))y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

If $f(a) = f'_x(a) = \dots = f^{(2n+1)}_x(a) = 0$ then the solution is

$$y(x) = \frac{2^n n!}{(2n)! \lambda^n} \int_a^x J_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2 \right)^{n+1} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 471–472).

$$18. \int_a^x (x-t)^{n+1} J_n(\lambda(x-t))y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

If $f(a) = f'_x(a) = \dots = f_x^{(2n+2)}(a) = 0$ then the solution is

$$y(x) = \frac{2^{n+1}(n+1)!}{(2n+2)!\lambda^n} \int_a^x J_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2 \right)^{n+2} F(t) dt, \quad F(t) = \int_a^t f(s) ds.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$19. \int_a^x (x-t)^{1/2} J_{1/2}(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.23 with $n = 1$. If $f(a) = f'_x(a) = 0$ then the solution is

$$y(x) = \sqrt{\frac{\pi}{2\lambda}} [f''_{xx}(x) + \lambda^2 f(x)].$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 471).

$$20. \int_a^x (x-t)^{3/2} J_{1/2}(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.24 with $n = 1$. Let $f(a) = f'_x(a) = f''_{xx}(a) = 0$. Then the solution is

$$y(x) = \frac{\sqrt{\pi}}{2\sqrt{2\lambda}} \left(\frac{d^2}{dx^2} + \lambda^2 \right)^2 \int_a^x f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$21. \int_a^x (x-t)^{3/2} J_{3/2}(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.23 with $n = 2$. If $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$ then the solution is

$$y(x) = \frac{\sqrt{\pi}}{(2\lambda)^{3/2}} \left(\frac{d^2}{dx^2} + \lambda^2 \right)^2 f(x).$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 471).

$$22. \int_a^x (x-t)^{5/2} J_{3/2}(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.24 with $n = 2$. Let $f(a) = f'_x(a) = \dots = f''''_{xxxx}(a) = 0$. Then the solution is

$$y(x) = \frac{\sqrt{\pi}}{4(2\lambda)^{3/2}} \left(\frac{d^2}{dx^2} + \lambda^2 \right)^3 \int_a^x f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$23. \int_a^x (x-t)^{\frac{2n-1}{2}} J_{\frac{2n-1}{2}}(\lambda(x-t))y(t) dt = f(x), \quad n = 1, 2, 3, \dots$$

Let $f(a) = f'_x(a) = \dots = f_x^{(2n-1)}(a) = 0$. Then the solution is

$$y(x) = \frac{\sqrt{\pi}}{(2\lambda)^{\frac{2n-1}{2}} (n-1)!} \left(\frac{d^2}{dx^2} + \lambda^2 \right)^n f(x).$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 471).

$$24. \int_a^x (x-t)^{\frac{2n+1}{2}} J_{\frac{2n-1}{2}}(\lambda(x-t))y(t) dt = f(x), \quad n = 1, 2, 3, \dots$$

Let $f(a) = f'_x(a) = \dots = f^{(2n)}(a) = 0$. Then the solution is

$$y(x) = \frac{\sqrt{\pi}}{2(2\lambda)^{n-1/2}n!} \left(\frac{d^2}{dx^2} + \lambda^2 \right)^{n+1} \int_a^x f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$25. \int_a^x [J_\nu(\lambda x) - J_\nu(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = J_\nu(\lambda x)$, where $J_\nu(z)$ is the Bessel function (see Supplement 11.6-1).

$$\text{Solution: } y(x) = \frac{d}{dx} \left[\frac{x f'_x(x)}{\nu J_\nu(\lambda x) - \lambda x J_{\nu+1}(\lambda x)} \right].$$

$$26. \int_a^x [A J_\nu(\lambda x) + B J_\nu(\lambda t)]y(t) dt = f(x).$$

For $B = -A$, see equation 1.8.25. We consider the interval $[a, x]$ in which $J_\nu(\lambda x)$ does not change its sign.

Solution with $B \neq -A$:

$$y(x) = \pm \frac{1}{A+B} \frac{d}{dx} \left\{ |J_\nu(\lambda x)|^{-\frac{A}{A+B}} \int_a^x |J_\nu(\lambda t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

Here the sign of $J_\nu(\lambda x)$ should be taken.

$$27. \int_a^x [A J_\nu(\lambda x) + B J_\mu(\beta t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = A J_\nu(\lambda x)$ and $h(t) = B J_\mu(\beta t)$.

$$28. \int_a^x (x-t)^\nu J_\nu(\lambda(x-t))y(t) dt = f(x).$$

1°. Let $\text{Re } \nu > -1/2$ and $f(a) = f'_x(a) = \dots = f_x^{(2n-1)}(a) = 0$, where $n = [\text{Re } \nu + 1/2] + 1$ and $[A]$ stands for the integer part of the number A . Then the solution is

$$y(x) = \frac{\pi(2\lambda)^{1-n}}{\Gamma(\nu+1/2)\Gamma(n-\nu-1/2)} \int_a^x (x-t)^{n-\nu-1} J_{n-\nu-1}(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2 \right)^n f(t) dt.$$

2°. For $\nu = n$ and $\nu = n - 1/2$ (n is an integer) see equations 1.8.17 and 1.8.23.

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 471), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$29. \int_a^x (x-t)^{\nu+1} J_\nu(\lambda(x-t))y(t) dt = f(x).$$

1°. Let $\text{Re } \nu > -1$ and $f(a) = f'_x(a) = \dots = f^{(2n-2)}(a) = 0$, where $n = [\text{Re } \nu + 3/2] + 1$ and $[A]$ stands for the integer part of the number A . Then the solution is

$$y(x) = \frac{2^{1-n} \lambda^{2-n} \pi}{\Gamma(\nu+3/2)\Gamma(n-\nu-3/2)} \int_a^x (x-t)^{n-\nu-2} J_{n-\nu-2}(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2 \right)^n F(t) dt,$$

where $F(t) = \int_a^t f(s) ds$.

2°. For $\nu = n$ and $\nu = n - 1/2$ (n is an integer) see equations 1.8.18 and 1.8.24.

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$30. \int_0^x \frac{J_\nu(\lambda(x-t))}{x-t} y(t) dt = f(x), \quad \operatorname{Re} \nu > 0.$$

1°. If $\nu = n$ is a positive integer number and $f(0) = f'_x(0) = \dots = f_x^{(n)}(0) = 0$ then

$$y(x) = \frac{n}{\lambda^n} \sum_{k=0}^{[n/2]} C_n^{2k} \left(\frac{d}{dx}\right)^{n-2k} \left(\frac{d^2}{dx^2} + \lambda^2\right)^k f(x) \\ + \frac{n}{\lambda^n} \int_0^x J_0(\lambda(x-t)) \sum_{k=0}^{[(n-1)/2]} C_n^{2k+1} \left(\frac{d}{dt}\right)^{n-2k-1} \left(\frac{d^2}{dt^2} + \lambda^2\right)^{k+1} f(t) dt,$$

where $[A]$ stands for the integer part of the number A and $C_n^k = \frac{n!}{k!(n-k)!}$ are binomial coefficients ($0! = 1$).

2°. If ν is not an integer, $[\operatorname{Re} \nu] + 1 = m > 1$, and $f(0) = f'_x(0) = \dots = f_x^{(m)}(0) = 0$ then

$$y(x) = \frac{\nu}{\lambda^m} \int_0^x J_{m-\nu}(\lambda(x-t)) \sum_{k=0}^{[(m-1)/2]} C_m^{2k+1} \left(\frac{d}{dt}\right)^{m-2k-1} \left(\frac{d^2}{dt^2} + \lambda^2\right)^{k+1} f(t) dt \\ + \frac{\nu(m-\nu)}{\lambda^m} \int_0^x \frac{J_{m-\nu}(\lambda(x-t))}{x-t} \sum_{k=0}^{[m/2]} C_m^{2k} \left(\frac{d}{dt}\right)^{m-2k} \left(\frac{d^2}{dt^2} + \lambda^2\right)^k f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 470–471).

$$31. \int_a^x J_0(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

This is a special case of equation 1.8.38 with $n = 0$. If $f(a) = f'_x(a) = 0$ then the solution is

$$y(x) = \frac{d^2}{dx^2} \int_a^x I_0(\lambda\sqrt{x-t}) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$32. \int_a^x [AJ_\nu(\lambda\sqrt{x}) + BJ_\nu(\lambda\sqrt{t})] y(t) dt = f(x).$$

We consider the interval $[a, x]$ in which $J_\nu(\lambda\sqrt{x})$ does not change its sign.

Solution with $B \neq -A$:

$$y(x) = \pm \frac{1}{A+B} \frac{d}{dx} \left\{ |J_\nu(\lambda\sqrt{x})|^{-\frac{A}{A+B}} \int_a^x |J_\nu(\lambda\sqrt{t})|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

Here the sign $J_\nu(\lambda\sqrt{x})$ should be taken.

$$33. \int_a^x [AJ_\nu(\lambda\sqrt{x}) + BJ_\mu(\beta\sqrt{t})] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = AJ_\nu(\lambda\sqrt{x})$ and $h(t) = BJ_\mu(\beta\sqrt{t})$.

$$34. \int_a^x \sqrt{x-t} J_1(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

This is a special case of equation 1.8.38 with $n = 1$. If the conditions $f(a) = f'_x(a) = f''_{xx}(a) = 0$ are satisfied, then the solution is

$$y(x) = \frac{2}{\lambda} \frac{d^3}{dx^3} \int_a^x I_0(\lambda\sqrt{x-t}) f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$35. \int_a^x (x-t)^{1/4} J_{1/2}(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

This is a special case of equation 1.8.39 with $n = 1$. If the conditions $f(a) = f'_x(a) = 0$ are satisfied, then the solution is

$$y(x) = \sqrt{\frac{2}{\pi\lambda}} \frac{d^2}{dx^2} \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$36. \int_a^x (x-t)^{3/4} J_{3/2}(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

This is a special case of equation 1.8.39 with $n = 2$. If the conditions $f(a) = f'_x(a) = f''_{xx}(a) = 0$ are satisfied, then the solution is

$$y(x) = \frac{2^{3/2}}{\sqrt{\pi}\lambda^{3/2}} \frac{d^3}{dx^3} \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$37. \int_a^x (x-t)^{-1/4} J_{-1/2}(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

This is a special case of equation 1.8.39 with $n = 0$. If the condition $f(a) = 0$ is satisfied, then the solution is

$$y(x) = \sqrt{\frac{\lambda}{2\pi}} \frac{d}{dx} \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$38. \int_a^x (x-t)^{n/2} J_n(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

This is a special case of equation 1.8.40 with $\nu = n$ and $m = n + 2$. If the conditions $f(a) = f'_x(a) = \dots = f^{(n+1)}_x(a) = 0$ are satisfied, then the solution is

$$y(x) = \left(\frac{2}{\lambda}\right)^n \frac{d^{n+2}}{dx^{n+2}} \int_a^x I_0(\lambda\sqrt{x-t}) f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$39. \int_a^x (x-t)^{\frac{2n-1}{4}} J_{\frac{2n-1}{2}}(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

This is a special case of equation 1.8.40 with $\nu = n - 1/2$ and $m = n + 1$. If the conditions $f(a) = f'_x(a) = \dots = f^{(n)}_x(a) = 0$ are satisfied, then the solution is

$$y(x) = \frac{1}{\sqrt{\pi}} \left(\frac{2}{\lambda}\right)^{\frac{2n-1}{2}} \frac{d^{n+1}}{dx^{n+1}} \int_a^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$40. \int_a^x (x-t)^{\nu/2} J_\nu(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad \operatorname{Re} \nu > -1.$$

1°. Let $f(a) = f'_x(a) = \dots = f_x^{(m-1)}(a) = 0$, where $m = [\operatorname{Re} \nu + 1] + 1$ and $[A]$ stands for the integer part of the number A . Then the solution is

$$y(x) = \left(\frac{2}{\lambda}\right)^{m-2} \frac{d^m}{dx^m} \int_a^x (x-t)^{\frac{m-\nu-2}{2}} I_{m-\nu-2}(\lambda\sqrt{x-t}) f(t) dt.$$

2°. For $\nu = n$ and $\nu = n - 1/2$ (n is an integer) see equations 1.8.38 and 1.8.39.

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 472).

$$41. \int_x^\infty (t-x)^{\nu/2} J_\nu(\lambda\sqrt{t-x}) y(t) dt = f(x), \quad \operatorname{Re} \nu > -1.$$

Solution:

$$y(x) = \left(\frac{\lambda}{2}\right)^{2-m} \left(-\frac{d}{dx}\right)^m \int_x^\infty (t-x)^{(m-\nu)/2-1} I_{m-\nu-2}(\lambda\sqrt{t-x}) f(t) dt,$$

where $m = [\operatorname{Re} \nu + 1] + 1$ and $[A]$ stands for the integer part of the number A .

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 474), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$42. \int_0^x (x-t)^{\nu/2} J_\nu(\lambda\sqrt{t(x-t)}) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{2} x^{-1/2} \int_0^x (x-t)^{-(\nu+1)/2} J_{-\nu-1}(\lambda\sqrt{x(x-t)}) t^{\nu+1} d(t^{-\nu} f(t)),$$

where $-1 < \operatorname{Re} \nu < 0$.

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 473), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$43. \int_0^x (x-t)^{\nu/2} J_\nu(\lambda\sqrt{x(x-t)}) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{2} x^{-\nu} \frac{d}{dx} \left(x^{\nu+1} \int_0^x t^{\nu/2} (x-t)^{-(\nu+1)/2} I_{-\nu-1}(\lambda\sqrt{t(x-t)}) f(t) dt \right),$$

where $-1 < \operatorname{Re} \nu < 0$.

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 473), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$44. \int_0^x J_0(\lambda\sqrt{x^2-t^2}) y(t) dt = f(x).$$

Solution:

$$y(x) = f'_x(x) + \lambda \frac{d}{dx} \int_0^x \frac{t}{\sqrt{x^2-t^2}} I_1(\lambda\sqrt{x^2-t^2}) f(t) dt.$$

⊙ Reference: S. Fenyö and H. W. Stolle (1984, p. 328).

$$45. \int_0^x (x^2 - t^2)^{-1/4} J_{-1/2}(\lambda\sqrt{x^2 - t^2}) y(t) dt = f(x).$$

Solution:

$$y(x) = \sqrt{\frac{2\lambda}{\pi}} \frac{d}{dx} \int_0^x t \frac{\cosh(\lambda\sqrt{x^2 - t^2})}{\sqrt{x^2 - t^2}} f(t) dt.$$

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$46. \int_x^\infty (t^2 - x^2)^{-1/4} J_{-1/2}(\lambda\sqrt{t^2 - x^2}) y(t) dt = f(x).$$

Solution:

$$y(x) = -\sqrt{\frac{2\lambda}{\pi}} \frac{d}{dx} \int_x^\infty t \frac{\cosh(\lambda\sqrt{t^2 - x^2})}{\sqrt{t^2 - x^2}} f(t) dt.$$

$$47. \int_0^x (x^2 - t^2)^{\nu/2} J_\nu(\lambda\sqrt{x^2 - t^2}) y(t) dt = f(x), \quad -1 < \nu < 0.$$

Solution:

$$y(x) = \lambda \frac{d}{dx} \int_0^x t (x^2 - t^2)^{-(\nu+1)/2} I_{-\nu-1}(\lambda\sqrt{x^2 - t^2}) f(t) dt.$$

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$48. \int_x^\infty (t^2 - x^2)^{\nu/2} J_\nu(\lambda\sqrt{t^2 - x^2}) y(t) dt = f(x), \quad -1 < \nu < 0.$$

Solution:

$$y(x) = -\lambda \frac{d}{dx} \int_x^\infty t (t^2 - x^2)^{-(\nu+1)/2} I_{-\nu-1}(\lambda\sqrt{t^2 - x^2}) f(t) dt.$$

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$49. \int_a^x [At^k J_\nu(\lambda x) + Bx^m J_\mu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = AJ_\nu(\lambda x)$, $h_1(t) = t^k$, $g_2(x) = Bx^m$, and $h_2(t) = J_\mu(\lambda t)$.

$$50. \int_a^x [AJ_\nu^2(\lambda x) + BJ_\nu^2(\lambda t)] y(t) dt = f(x).$$

Solution with $B \neq -A$:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |J_\nu(\lambda x)|^{-\frac{2A}{A+B}} \int_a^x |J_\nu(\lambda t)|^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$51. \int_a^x [AJ_\nu^k(\lambda x) + BJ_\mu^m(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = AJ_\nu^k(\lambda x)$ and $h(t) = BJ_\mu^m(\beta t)$.

$$52. \int_0^x \left(\frac{x-t}{x-t+\gamma} \right)^{\nu/2} J_\nu(\lambda\sqrt{(x-t)(x-t+\gamma)})y(t) dt = f(x).$$

Let $-1 < \operatorname{Re} \nu < m+1 < 2n+1$ (n and m are the minimal integer numbers), and $f(0) = f'_x(0) = \dots = f_x^{(2n+m+1)}(0) = 0$. Then

$$y(x) = \sqrt{\pi} \lambda^{-m} \int_0^x \left(\frac{x-t}{x-t-\gamma} \right)^{(m-\nu)/2} J_{m-\nu}(\lambda\sqrt{(x-t)(x-t-\gamma)}) \\ \times \int_0^t \sum_{j=0}^m \frac{C_m^j}{\Gamma(n-j/2)} \left(\frac{t-s}{2\lambda} \right)^{n-(j+1)/2} J_{n-(j+1)/2}(\lambda(t-s)) \left(\frac{d^2}{ds^2} + \lambda^2 \right)^{n+1} \left(\frac{d}{ds} \right)^{m-j} f(s) ds dt,$$

where C_n^k are binomial coefficients.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 473), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$53. \int_a^x [Y_0(\lambda x) - Y_0(\lambda t)]y(t) dt = f(x).$$

$$\text{Solution: } y(x) = -\frac{d}{dx} \left[\frac{f'_x(x)}{\lambda Y_1(\lambda x)} \right].$$

$$54. \int_a^x [Y_\nu(\lambda x) - Y_\nu(\lambda t)]y(t) dt = f(x).$$

$$\text{Solution: } y(x) = \frac{d}{dx} \left[\frac{x f'_x(x)}{\nu Y_\nu(\lambda x) - \lambda x Y_{\nu+1}(\lambda x)} \right].$$

$$55. \int_a^x [AY_\nu(\lambda x) + BY_\nu(\lambda t)]y(t) dt = f(x).$$

For $B = -A$, see equation 1.8.54. We consider the interval $[a, x]$ in which $Y_\nu(\lambda x)$ does not change its sign.

Solution with $B \neq -A$:

$$y(x) = \pm \frac{1}{A+B} \frac{d}{dx} \left\{ |Y_\nu(\lambda x)|^{-\frac{A}{A+B}} \int_a^x |Y_\nu(\lambda t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

Here the sign of $Y_\nu(\lambda x)$ should be taken.

$$56. \int_a^x [At^k Y_\nu(\lambda x) + Bx^m Y_\mu(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = AY_\nu(\lambda x)$, $h_1(t) = t^k$, $g_2(x) = Bx^m$, and $h_2(t) = Y_\mu(\lambda t)$.

$$57. \int_a^x [AJ_\nu(\lambda x)Y_\mu(\beta t) + BJ_\nu(\lambda t)Y_\mu(\beta x)]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = AJ_\nu(\lambda x)$, $h_1(t) = Y_\mu(\beta t)$, $g_2(x) = BY_\mu(\beta x)$, and $h_2(t) = J_\nu(\lambda t)$.

1.8-6. Kernels Containing Modified Bessel Functions.

$$58. \int_a^x I_0(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.64 with $n = 0$ and $I_0(z)$ is the modified Bessel function (see Supplement 11.7-1). If $f(a) = f'_x(a) = 0$ then the solution is

$$y(x) = \int_a^x I_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2 \right) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 481).

$$59. \int_a^x [I_0(\lambda x) - I_0(\lambda t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

$$\text{Solution: } y(x) = \frac{d}{dx} \left[\frac{f'_x(x)}{\lambda I_1(\lambda x)} \right].$$

$$60. \int_a^x [AI_0(\lambda x) + BI_0(\lambda t)]y(t) dt = f(x).$$

For $B = -A$, see equation 1.8.59. Solution with $B \neq -A$:

$$y(x) = \pm \frac{1}{A+B} \frac{d}{dx} \left\{ |I_0(\lambda x)|^{-\frac{A}{A+B}} \int_a^x |I_0(\lambda t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

Here the sign of $I_\nu(\lambda x)$ should be taken.

$$61. \int_a^x (x-t)I_0(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.65 with $n = 0$. If $f(a) = f'_x(a) = f''_{xx}(a) = 0$ then the solution is

$$y(x) = \int_a^x I_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2 \right)^2 F(t) dt, \quad F(t) = \int_a^t f(s) ds.$$

$$62. \int_a^x (x-t)I_1(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.64 with $n = 1$. If $f(a) = f'_x(a) = 0$ then the solution is

$$y(x) = \lambda^{-1} f'_x(x) + \lambda^{-1} \int_a^x I_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2 \right) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 481).

$$63. \int_a^x (x-t)^2 I_1(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.65 with $n = 1$. If $f(a) = f'_x(a) = \dots = f''''_{xxxx}(a) = 0$ then the solution is

$$y(x) = \frac{1}{3\lambda} \int_a^x I_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2 \right)^3 F(t) dt, \quad F(t) = \int_a^t f(s) ds.$$

$$64. \int_a^x (x-t)^n I_n(\lambda(x-t))y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

If $f(a) = f'_x(a) = \dots = f_x^{(2n+1)}(a) = 0$ then the solution is

$$y(x) = \frac{2^n n!}{(2n)! \lambda^n} \int_a^x I_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2 \right)^{n+1} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 481), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$65. \int_a^x (x-t)^{n+1} I_n(\lambda(x-t))y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

This is a special case of equation 1.8.78 with $\nu = n$ and $m = n + 2$. If $f(a) = f'_x(a) = \dots = f_x^{(2n+2)}(a) = 0$ then the solution is

$$y(x) = \frac{2^{n+1}(n+1)!}{(2n+2)! \lambda^n} \int_a^x I_0(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2 \right)^{n+2} F(t) dt, \quad F(t) = \int_a^t f(s) ds.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 482), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$66. \int_a^x (x-t)^{1/2} I_{1/2}(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.70 with $n = 1$. If $f(a) = f'_x(a) = 0$ then the solution is

$$y(x) = \sqrt{\frac{\pi}{2\lambda}} [f''_{xx}(x) - \lambda^2 f(x)].$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 481).

$$67. \int_a^x (x-t)^{3/2} I_{1/2}(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.71 with $n = 1$. If $f(a) = f'_x(a) = f''_{xx}(a) = 0$ then the solution is

$$y(x) = \frac{\sqrt{\pi}}{2(2\lambda)^{1/2}} \left(\frac{d^2}{dx^2} - \lambda^2 \right)^2 \int_a^x f(t) dt.$$

$$68. \int_a^x (x-t)^{3/2} I_{3/2}(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.70 with $n = 2$. If $f(a) = f'_x(a) = f''_{xx}(a) = f'''_{xxx}(a) = 0$ then the solution is

$$y(x) = \frac{\sqrt{\pi}}{(2\lambda)^{3/2}} \left(\frac{d^2}{dx^2} - \lambda^2 \right)^2 f(x).$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 481).

$$69. \int_a^x (x-t)^{5/2} I_{3/2}(\lambda(x-t))y(t) dt = f(x).$$

This is a special case of equation 1.8.71 with $n = 2$. If $f(a) = f'_x(a) = \dots = f_{xxxx}(a) = 0$ then the solution is

$$y(x) = \frac{\sqrt{\pi}}{4(2\lambda)^{3/2}} \left(\frac{d^2}{dx^2} - \lambda^2 \right)^3 \int_a^x f(t) dt.$$

$$70. \int_a^x (x-t)^{\frac{2n-1}{2}} I_{\frac{2n-1}{2}}(\lambda(x-t))y(t) dt = f(x), \quad n = 1, 2, 3, \dots$$

If $f(a) = f'_x(a) = \dots = f_x^{(2n-1)}(a) = 0$ then the solution is

$$y(x) = \frac{\sqrt{\pi}}{(2\lambda)^{n-1/2}(n-1)!} \left(\frac{d^2}{dx^2} - \lambda^2 \right)^n f(x).$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 481), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$71. \int_a^x (x-t)^{\frac{2n+1}{2}} I_{\frac{2n+1}{2}}(\lambda(x-t))y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

If $f(a) = f'_x(a) = \dots = f_x^{(2n)}(a) = 0$ then the solution is

$$y(x) = \frac{\sqrt{\pi}}{2(2\lambda)^{n-1/2}n!} \left(\frac{d^2}{dx^2} - \lambda^2 \right)^{n+1} \int_a^x f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 482).

$$72. \int_a^x [I_\nu(\lambda x) - I_\nu(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = I_\nu(\lambda x)$, where $I_\nu(z)$ is the modified Bessel function (see Supplement 11.7-1).

$$73. \int_a^x [AI_\nu(\lambda x) + BI_\nu(\lambda t)]y(t) dt = f(x).$$

Solution with $B \neq -A$:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [I_\nu(\lambda x)]^{-\frac{A}{A+B}} \int_a^x [I_\nu(\lambda t)]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$74. \int_a^x [AI_\nu(\lambda x) + BI_\mu(\beta t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = AI_\nu(\lambda x)$ and $h(t) = BI_\mu(\beta t)$.

$$75. \int_0^x I_\nu(\lambda(x-t))y(t) dt = f(x).$$

1°. Let $-1 < \operatorname{Re} \nu < 1$ and $f(0) = f'_x(0) = 0$. Then the solution is

$$y(x) = \int_0^x I_{-\nu}(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2 \right) f(t) dt.$$

2°. Let $\nu = n \geq 0$ (n is an integer number) and $f(0) = f'_x(0) = \dots = f_x^{(n+1)}(0) = 0$. Then the solution is

$$y(x) = \lambda^{-n} \sum_{k=0}^{[(n-1)/2]} C_n^{2k+1} \left(\frac{d}{dx} \right)^{n-2k-1} \left(\frac{d^2}{dx^2} - \lambda^2 \right)^{k+1} f(x) \\ + \lambda^{-n} \int_0^x I_0(\lambda(x-t)) \sum_{k=0}^{[n/2]} C_n^{2k} \left(\frac{d}{dt} \right)^{n-2k} \left(\frac{d^2}{dt^2} - \lambda^2 \right)^{k+1} f(t) dt,$$

where $[A]$ stands for the integer part of the number A and C_n^k are binomial coefficients.

3°. Let $\operatorname{Re} \nu > -1$ and $f(0) = f'_x(0) = \dots = f_x^{(m+1)}(0) = 0$, where $m = [\operatorname{Re} \nu + 1]$. Then the solution is

$$y(x) = \frac{m - \nu}{\lambda^m} \int_0^x \frac{I_{m-\nu}(\lambda(x-t))}{x-t} \sum_{k=0}^{[(m-1)/2]} C_m^{2k+1} \left(\frac{d}{dt}\right)^{m-2k-1} \left(\frac{d^2}{dt^2} - \lambda^2\right)^{k+1} f(t) dt \\ + \lambda^{-m} \int_0^x I_{m-\nu}(\lambda(x-t)) \sum_{k=0}^{[m/2]} C_m^{2k} \left(\frac{d}{dt}\right)^{m-2k} \left(\frac{d^2}{dt^2} - \lambda^2\right)^{k+1} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 479–480), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

76. $\int_0^x (x-t)^{-1} I_\nu(\lambda(x-t)) y(t) dt = f(x).$

1°. Let $\operatorname{Re} \nu > 0$ and $f(0) = f'_x(0) = \dots = f_x^{(m)}(0) = 0$, where $m = [\operatorname{Re} \nu] + 1$ and $[A]$ stands for the integer part of the number A . Then the solution is

$$y(x) = \nu \lambda^{-m} \int_0^x I_{m-\nu}(\lambda(x-t)) \sum_{k=0}^{[(m-1)/2]} C_m^{2k+1} \left(\frac{d}{dt}\right)^{m-2k-1} \left(\frac{d^2}{dt^2} - \lambda^2\right)^{k+1} f(t) dt \\ + \nu(m-\nu) \lambda^{-m} \int_0^x (x-t)^{-1} I_{m-\nu}(\lambda(x-t)) \sum_{k=0}^{[m/2]} C_m^{2k} \left(\frac{d}{dt}\right)^{m-2k} \left(\frac{d^2}{dt^2} - \lambda^2\right)^k f(t) dt,$$

where C_n^k are binomial coefficients.

2°. If $\nu = n > 0$ (n is an integer number) and $f(0) = f'_x(0) = \dots = f_x^{(n)}(0) = 0$ then

$$y(x) = n \lambda^{-n} \sum_{k=0}^{[n/2]} C_n^{2k} \left(\frac{d}{dx}\right)^{n-2k} \left(\frac{d^2}{dx^2} - \lambda^2\right)^k f(x) \\ + n \lambda^{-n} \int_0^x I_0(\lambda(x-t)) \sum_{k=0}^{[(n-1)/2]} C_n^{2k+1} \left(\frac{d}{dt}\right)^{n-2k-1} \left(\frac{d^2}{dt^2} - \lambda^2\right)^{k+1} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 480–481).

77. $\int_a^x (x-t)^\nu I_\nu(\lambda(x-t)) y(t) dt = f(x), \quad \operatorname{Re} \nu > -1/2.$

1°. Let $f(a) = f'_x(a) = \dots = f_x^{(2m-1)}(a) = 0$, where $m = [\operatorname{Re} \nu + 1/2] + 1$ and $[A]$ stands for the integer part of the number A . Then the solution is

$$y(x) = \frac{(2\lambda)^{1-m} \pi}{\Gamma(\nu + 1/2) \Gamma(m - \nu - 1/2)} \int_a^x (x-t)^{m-\nu-1} I_{m-\nu-1}(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2\right)^m f(t) dt.$$

2°. Let $f(a) = f'_x(a) = \dots = f_x^{(m-1)}(a) = 0$, where $m = [2 \operatorname{Re} \nu + 1] + 1$. Then the solution is

$$y(x) = \frac{\sqrt{\pi} \lambda \Gamma(-\nu - 1)}{2^{2\nu+1} \Gamma(\nu + 1/2)} e^{-\lambda x} \frac{d^m}{dx^m} \left[e^{\lambda x} \int_a^x (x-t)^{m-\nu-1} \right. \\ \left. \times \sum_{k=0}^m \frac{(-m)_k (-2\nu - 2)_k}{\Gamma(m+k-2\nu-1) k!} (k-\nu-1) I_{k-\nu-1}(\lambda(x-t)) f(t) dt \right],$$

where $(a)_k = a(a+1) \dots (a+k-1)$ is the Pochhammer symbol.

3°. For $\nu = n$ and $\nu = n - 1/2$ (n is an integer) see equations 1.8.64 and 1.8.70.

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 481), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$78. \int_a^x (x-t)^{\nu+1} I_\nu(\lambda(x-t))y(t) dt = f(x).$$

1°. Let $\operatorname{Re} \nu > -1$ and $f(a) = f'_x(a) = \dots = f^{(2m-2)}(a) = 0$, where $m = [\operatorname{Re} \nu + 3/2] + 1$ and $[A]$ stands for the integer part of the number A . Then the solution is

$$y(x) = \frac{2^{1-m} \lambda^{2-m} \pi}{\Gamma(\nu + 3/2) \Gamma(m - \nu - 3/2)} \int_a^x (x-t)^{m-\nu-2} I_{m-\nu-2}(\lambda(x-t)) \left(\frac{d^2}{dt^2} - \lambda^2 \right)^m F(t) dt,$$

where $F(t) = \int_a^t f(s) ds$.

2°. For $\nu = n$ and $\nu = n - 1/2$ (n is an integer) see equations 1.8.65 and 1.8.71.

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 482), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$79. \int_a^x I_0(\lambda\sqrt{x-t})y(t) dt = f(x).$$

This is a special case of equation 1.8.86 with $n = 0$. If the conditions $f(a) = f'_x(a) = 0$ are satisfied, then the solution is

$$y(x) = \frac{d^2}{dx^2} \int_a^x J_0(\lambda\sqrt{x-t}) f(t) dt.$$

$$80. \int_a^x [AI_\nu(\lambda\sqrt{x}) + BI_\nu(\lambda\sqrt{t})]y(t) dt = f(x).$$

Solution with $B \neq -A$:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [I_\nu(\lambda\sqrt{x})]^{-\frac{A}{A+B}} \int_a^x [I_\nu(\lambda\sqrt{t})]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$81. \int_a^x [AI_\nu(\lambda\sqrt{x}) + BI_\mu(\beta\sqrt{t})]y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = AI_\nu(\lambda\sqrt{x})$ and $h(t) = BI_\mu(\beta\sqrt{t})$.

$$82. \int_a^x \sqrt{x-t} I_1(\lambda\sqrt{x-t})y(t) dt = f(x).$$

This is a special case of equation 1.8.86 with $n = 1$. If the conditions $f(a) = f'_x(a) = f''_{xx}(a) = 0$ are satisfied, then the solution is

$$y(x) = \frac{2}{\lambda} \frac{d^3}{dx^3} \int_a^x J_0(\lambda\sqrt{x-t}) f(t) dt.$$

$$83. \int_a^x (x-t)^{1/4} I_{1/2}(\lambda\sqrt{x-t})y(t) dt = f(x).$$

This is a special case of equation 1.8.87 with $n = 1$. If the conditions $f(a) = f'_x(a) = 0$ are satisfied, then the solution is

$$y(x) = \sqrt{\frac{2}{\pi\lambda}} \frac{d^2}{dx^2} \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$84. \int_a^x (x-t)^{3/4} I_{3/2}(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

This is a special case of equation 1.8.87 with $n = 2$. If the conditions $f(a) = f'_x(a) = f''_{xx}(a) = 0$ are satisfied, then the solution is

$$y(x) = \frac{2^{3/2}}{\sqrt{\pi} \lambda^{3/2}} \frac{d^3}{dx^3} \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$85. \int_a^x (x-t)^{-1/4} I_{-1/2}(\lambda\sqrt{x-t}) y(t) dt = f(x).$$

This is a special case of equation 1.8.87 with $n = 0$. If the condition $f(a) = 0$ is satisfied, then the solution is

$$y(x) = \sqrt{\frac{\lambda}{2\pi}} \frac{d}{dx} \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$86. \int_a^x (x-t)^{n/2} I_n(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

This is a special case of equation 1.8.88 with $\nu = n$ and $m = n + 2$. If the conditions $f(a) = f'_x(a) = \dots = f_x^{(n+1)}(a) = 0$ are satisfied, then the solution is

$$y(x) = \left(\frac{2}{\lambda}\right)^n \frac{d^{n+2}}{dx^{n+2}} \int_a^x J_0(\lambda\sqrt{x-t}) f(t) dt.$$

$$87. \int_a^x (x-t)^{\frac{2n-1}{4}} I_{\frac{2n-1}{2}}(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

This is a special case of equation 1.8.88 with $\nu = n - 1/2$ and $m = n + 1$. If the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$ are satisfied, then the solution is

$$y(x) = \frac{1}{\sqrt{\pi}} \left(\frac{2}{\lambda}\right)^{\frac{2n-1}{2}} \frac{d^{n+1}}{dx^{n+1}} \int_a^x \frac{\cos(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

$$88. \int_a^x (x-t)^{\nu/2} I_\nu(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad \operatorname{Re} \nu > -1.$$

1°. Let $f(a) = f'_x(a) = \dots = f_x^{(m-1)}(a) = 0$, where $m = [\operatorname{Re} \nu + 1] + 1$ and $[A]$ stands for the integer part of the number A . Then the solution is

$$y(x) = \left(\frac{2}{\lambda}\right)^{m-2} \frac{d^m}{dx^m} \int_a^x (x-t)^{\frac{m-\nu-2}{2}} J_{m-\nu-2}(\lambda\sqrt{x-t}) f(t) dt.$$

2°. For $\nu = n$ and $\nu = n - 1/2$ (n is an integer) see equations 1.8.86 and 1.8.87.

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 482), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$89. \int_x^\infty (t-x)^{\nu/2} I_\nu(\lambda\sqrt{t-x}) y(t) dt = f(x), \quad \operatorname{Re} \nu > -1.$$

Solution:

$$y(x) = \left(\frac{\lambda}{2}\right)^{2-m} \left(-\frac{d}{dx}\right)^m \int_x^\infty (t-x)^{(m-\nu)/2-1} J_{m-\nu-2}(\lambda\sqrt{t-x}) f(t) dt,$$

where $m = [\operatorname{Re} \nu + 1] + 1$ and $[A]$ stands for the integer part of the number A .

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 484), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$90. \int_0^x (x-t)^{\nu/2} I_\nu(\lambda\sqrt{t(x-t)}) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{2} x^{-1/2} \int_0^x (x-t)^{-(\nu+1)/2} J_{-\nu-1}(\lambda\sqrt{x(x-t)}) t^{\nu+1} d(t^{-\nu} f(t)),$$

where $-1 < \operatorname{Re} \nu < 0$.

⊙ References: K. Soni (1968), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 483), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$91. \int_0^x (x-t)^{\nu/2} I_\nu(\lambda\sqrt{x(x-t)}) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{2} x^{-\nu} \frac{d}{dx} \left[x^{\nu+1} \int_0^x t^{\nu/2} (x-t)^{-(\nu+1)/2} J_{-\nu-1}(\lambda\sqrt{t(x-t)}) f(t) dt \right],$$

where $-1 < \operatorname{Re} \nu < 0$.

⊙ References: K. Soni (1968), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 483), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$92. \int_0^x (x^2-t^2)^{-1/4} I_{-1/2}(\lambda\sqrt{x^2-t^2}) y(t) dt = f(x).$$

Solution:

$$y(x) = \sqrt{\frac{2\lambda}{\pi}} \frac{d}{dx} \int_0^x t \frac{\cos(\lambda\sqrt{x^2-t^2})}{\sqrt{x^2-t^2}} f(t) dt.$$

$$93. \int_x^\infty (t^2-x^2)^{-1/4} I_{-1/2}(\lambda\sqrt{t^2-x^2}) y(t) dt = f(x).$$

Solution:

$$y(x) = -\sqrt{\frac{2\lambda}{\pi}} \frac{d}{dx} \int_x^\infty t \frac{\cos(\lambda\sqrt{t^2-x^2})}{\sqrt{t^2-x^2}} f(t) dt.$$

$$94. \int_0^x (x^2-t^2)^{\nu/2} I_\nu(\lambda\sqrt{x^2-t^2}) y(t) dt = f(x), \quad -1 < \nu < 0.$$

Solution:

$$y(x) = \lambda \frac{d}{dx} \int_0^x t (x^2-t^2)^{-(\nu+1)/2} J_{-\nu-1}(\lambda\sqrt{x^2-t^2}) f(t) dt.$$

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$95. \int_x^\infty (t^2-x^2)^{\nu/2} I_\nu(\lambda\sqrt{t^2-x^2}) y(t) dt = f(x), \quad -1 < \nu < 0.$$

Solution:

$$y(x) = -\lambda \frac{d}{dx} \int_x^\infty t (t^2-x^2)^{-(\nu+1)/2} J_{-\nu-1}(\lambda\sqrt{t^2-x^2}) f(t) dt.$$

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$96. \int_0^x \left(\frac{x-t}{x-t+\gamma} \right)^{\nu/2} I_\nu(\lambda\sqrt{(x-t)(x-t+\gamma)}) y(t) dt = f(x).$$

Let $-1 < \operatorname{Re} \nu < m+1 < 2n+1$ (n and m are the minimal integer numbers), and $f(0) = f'_x(0) = \dots = f_x^{(2n+m+1)}(0) = 0$. Then

$$y(x) = \sqrt{\pi} \lambda^{-m} \int_0^x \left(\frac{x-t}{x-t+\gamma} \right)^{(m-\nu)/2} I_{m-\nu}(\lambda\sqrt{(x-t)(x-t+\gamma)}) \\ \times \int_0^t \sum_{j=0}^m \frac{C_m^j}{\Gamma(n-j/2)} \left(\frac{t-s}{2\lambda} \right)^{n-(j+1)/2} I_{n-(j+1)/2}(\lambda(t-s)) \left(\frac{d^2}{ds^2} - \lambda^2 \right)^{n+1} \left(\frac{d}{ds} \right)^{m-j} f(s) ds dt,$$

where C_n^k are binomial coefficients.

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 483–484), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$97. \int_a^x [At^k I_\nu(\lambda x) + Bx^s I_\mu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = AI_\nu(\lambda x)$, $h_1(t) = t^k$, $g_2(x) = Bx^s$, and $h_2(t) = I_\mu(\lambda t)$.

$$98. \int_a^x [AI_\nu^2(\lambda x) + BI_\nu^2(\lambda t)] y(t) dt = f(x).$$

Solution with $B \neq -A$:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ |I_\nu(\lambda x)|^{-\frac{2A}{A+B}} \int_a^x |I_\nu(\lambda t)|^{-\frac{2B}{A+B}} f'_t(t) dt \right\}.$$

$$99. \int_a^x [AI_\nu^k(\lambda x) + BI_\mu^s(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.6 with $g(x) = AI_\nu^k(\lambda x)$ and $h(t) = BI_\mu^s(\beta t)$.

$$100. \int_a^x [K_0(\lambda x) - K_0(\lambda t)] y(t) dt = f(x).$$

$$\text{Solution: } y(x) = -\frac{d}{dx} \left[\frac{f'_x(x)}{\lambda K_1(\lambda x)} \right].$$

$$101. \int_a^x [K_\nu(\lambda x) - K_\nu(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.2 with $g(x) = K_\nu(\lambda x)$.

$$102. \int_a^x [AK_\nu(\lambda x) + BK_\nu(\lambda t)] y(t) dt = f(x).$$

Solution with $B \neq -A$:

$$y(x) = \frac{1}{A+B} \frac{d}{dx} \left\{ [K_\nu(\lambda x)]^{-\frac{A}{A+B}} \int_a^x [K_\nu(\lambda t)]^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$103. \int_a^x [At^k K_\nu(\lambda x) + Bx^s K_\mu(\lambda t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = AK_\nu(\lambda x)$, $h_1(t) = t^k$, $g_2(x) = Bx^s$, and $h_2(t) = K_\mu(\lambda t)$.

$$104. \int_a^x [AI_\nu(\lambda x)K_\mu(\beta t) + BI_\nu(\lambda t)K_\mu(\beta x)]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = AI_\nu(\lambda x)$, $h_1(t) = K_\mu(\beta t)$, $g_2(x) = BK_\mu(\beta x)$, and $h_2(t) = I_\nu(\lambda t)$.

1.8-7. Kernels Containing Legendre Polynomials.

$$105. \int_1^x P_n\left(\frac{x}{t}\right)y(t) dt = f(x), \quad f(1) = 0, \quad x \geq 1.$$

Here $P_n(x)$ is the Legendre polynomial (see Supplement 11.11-1).

Solution:

$$y(x) = \frac{x^{n+1}}{(n-1)!} \left(\frac{1}{x} \frac{d}{dx}\right)^{n+1} \int_1^x (x-t)^{n-1} f(t) dt,$$

where $n = 1, 2, 3, \dots$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 495–496).

$$106. \int_1^x P_n\left(\frac{t}{x}\right)y(t) dt = f(x), \quad f(1) = f'_x(1) = 0, \quad x \geq 1.$$

Solution:

$$y(x) = \int_1^x t^{2-n} P_{n-2}\left(\frac{x}{t}\right) \left(\frac{1}{t} \frac{d}{dt}\right)^2 [t^n f(t)] dt,$$

where $n = 2, 3, 4, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 496).

$$107. \int_x^1 P_n\left(\frac{x}{t}\right)y(t) dt = f(x), \quad f(1) = f'_x(1) = 0, \quad 0 < x \leq 1.$$

Solution:

$$y(x) = x^{-2} \int_x^1 t^{n+2} P_{n-2}\left(\frac{t}{x}\right) \left(\frac{1}{t} \frac{d}{dt}\right)^2 [t^{2-n} f(t)] dt,$$

where $n = 2, 3, 4, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 496).

$$108. \int_x^1 P_n\left(\frac{t}{x}\right)y(t) dt = f(x), \quad f(1) = f'_x(1) = 0, \quad 0 < x \leq 1.$$

Solution:

$$y(x) = \int_x^1 t^{2-n} P_{n-2}\left(\frac{x}{t}\right) \left(\frac{1}{t} \frac{d}{dt}\right)^2 [t^n f(t)] dt,$$

where $n = 2, 3, 4, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 496).

$$109. \int_0^x P_n \left(2 \frac{x}{t} - 1 \right) y(t) dt = f(x), \quad f(0) = 0, \quad x > 0.$$

Solution:

$$y(x) = \frac{x^n}{(n-1)!} \frac{d^{n+1}}{dx^{n+1}} \left[x^{-n} \int_0^x (x-t)^{n-1} f(t) dt \right],$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 497).

$$110. \int_x^1 P_n \left(2 \frac{x}{t} - 1 \right) y(t) dt = f(x), \quad f(1) = 0, \quad x \leq 1.$$

Solution:

$$y(x) = x^n \left(-\frac{d}{dx} \right)^{n+1} \left[x^{-n} \int_x^1 \frac{(t-x)^{n-1}}{(n-1)!} f(t) dt \right],$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 497).

$$111. \int_x^1 P_n \left(2 \frac{t}{x} - 1 \right) y(t) dt = f(x), \quad f(1) = 0, \quad x \leq 1.$$

Solution:

$$y(x) = \left(-\frac{d}{dx} \right)^{n+1} \left[x^{n+1} \int_x^1 \frac{(t-x)^{n-1}}{(n-1)!} t^{-n-1} f(t) dt \right],$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 498).

$$112. \int_0^x P_n(\cosh(x-t)) y(t) dt = f(x), \quad f(0) = f'_x(0) = 0.$$

Solution:

$$y(x) = \left(\frac{d^2}{dx^2} - (n+1)^2 \right) \int_0^x P_{n+1}(\cosh(x-t)) f(t) dt,$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 498).

$$113. \int_0^x P_n(\cos(x-t)) y(t) dt = f(x), \quad f(0) = f'_x(0) = 0.$$

Solution:

$$y(x) = \left(\frac{d^2}{dx^2} + (n+1)^2 \right) \int_0^x P_{n+1}(\cos(x-t)) f(t) dt,$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 498).

1.8-8. Kernels Containing Associated Legendre Functions.

$$114. \int_a^x (x^2 - t^2)^{-\mu/2} P_\nu^\mu \left(\frac{x}{t} \right) y(t) dt = f(x), \quad 0 \leq a < x.$$

Here $P_\nu^\mu(x)$ is the modified associated Legendre function (see Supplement 11.11-3).

Let $1 - n < \operatorname{Re} \mu < 1$ ($n = 1, 2, \dots$) and $f(a) = f'_x(a) = \dots = f_x^{(n-1)}(a) = 0$. Then the solution is

$$y(x) = x^{n+\mu-1} \frac{d^n}{dx^n} \left[x^{1-\mu} \int_a^x (x^2 - t^2)^{\frac{n+\mu-2}{2}} t^{-n} P_\nu^{2-n-\mu} \left(\frac{t}{x} \right) f(t) dt \right].$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 515), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$115. \int_a^x (x^2 - t^2)^{-\mu/2} P_\nu^\mu \left(\frac{t}{x} \right) y(t) dt = f(x), \quad 0 \leq a < x.$$

Let $1 - n < \operatorname{Re} \mu < 1$ ($n = 1, 2, \dots$) and $f(a) = f'_x(a) = \dots = f_x^{(n-1)}(a) = 0$. Then the solution is

$$y(x) = \frac{d^n}{dx^n} \int_a^x (x^2 - t^2)^{\frac{n+\mu-2}{2}} P_\nu^{2-n-\mu} \left(\frac{x}{t} \right) f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 515), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$116. \int_x^\infty (t^2 - x^2)^{-\mu/2} P_\nu^\mu \left(\frac{x}{t} \right) y(t) dt = f(x).$$

Let $1 - n < \operatorname{Re} \mu < 1$ ($n = 1, 2, \dots$). Then the solution is

$$y(x) = (-1)^n x^{n+\mu-1} \frac{d^n}{dx^n} \left[x^{1-\mu} \int_x^b (t^2 - x^2)^{\frac{n+\mu-2}{2}} t^{-n} P_\nu^{2-n-\mu} \left(\frac{t}{x} \right) f(t) dt \right].$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 516), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$117. \int_x^\infty (t^2 - x^2)^{-\mu/2} P_\nu^\mu \left(\frac{t}{x} \right) y(t) dt = f(x).$$

Let $1 - n < \operatorname{Re} \mu < 1$ ($n = 1, 2, \dots$). Then the solution is

$$y(x) = (-1)^n \frac{d^n}{dx^n} \int_x^b (t^2 - x^2)^{\frac{n+\mu-2}{2}} P_\nu^{2-n-\mu} \left(\frac{x}{t} \right) f(t) dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 516), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

1.8-9. Kernels Containing Confluent Hypergeometric Functions.

$$118. \int_s^x (x-t)^{b-1} \Phi(a, b; \lambda(x-t)) y(t) dt = f(x).$$

Here $\Phi(a, b; z)$ is the Kummer confluent hypergeometric function (see Supplement 11.9-1).

Let $0 < \operatorname{Re} b < n$ ($n = 1, 2, \dots$) and $f(s) = f'_x(s) = \dots = f_x^{(n-1)}(s) = 0$. Then the solution is

$$y(x) = \frac{d^n}{dx^n} \int_s^x \frac{(x-t)^{n-b-1}}{\Gamma(b)\Gamma(n-b)} \Phi(-a, n-b; \lambda(x-t)) f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 530), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$119. \int_x^\infty (t-x)^{b-1} \Phi(a, b; \lambda(x-t)) y(t) dt = f(x).$$

Here $\Phi(a, b; z)$ is the Kummer confluent hypergeometric function (see Supplement 11.9-1).
If $0 < \operatorname{Re} b < n$ ($n = 1, 2, \dots$) then the solution is

$$y(x) = \int_x^\infty \frac{(t-x)^{n-b-1}}{\Gamma(b)\Gamma(n-b)} \Phi(-a, n-b; \lambda(x-t)) f_t^{(n)}(t) dt.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 530), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$120. \int_0^x (x-t)^{\nu-1/2} M_{\mu,\nu}(\lambda(x-t)) y(t) dt = f(x).$$

Here $M_{\mu,\nu}(z)$ is the Whittaker confluent hypergeometric function (see Supplement 11.9-3).
Let $-1/2 < \operatorname{Re} \nu < (n-1)/2$ and $f(0) = f'_x(0) = \dots = f_x^{(n-1)}(0) = 0$. Then solution is

$$y(x) = \frac{\lambda^{-n/2}}{\Gamma(2\nu+1)} e^{-\lambda x/2} \frac{d^n}{dx^n} \left[e^{\lambda x/2} \int_0^x \frac{(x-t)^{(\nu-3)/2-\nu}}{\Gamma((\nu-1)/2-\nu)} M_{n/2-\mu, n/2-\nu-1}(\lambda(x-t)) f(t) dt \right].$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 522).

$$121. \int_x^\infty (t-x)^{\nu-1/2} M_{\mu,\nu}(\lambda(t-x)) y(t) dt = f(x).$$

Here $M_{\mu,\nu}(z)$ is the Whittaker confluent hypergeometric function (see Supplement 11.9-3).
Let $-1/2 < \operatorname{Re} \nu < (n-1)/2$. Then solution is

$$y(x) = \frac{\lambda^{-n/2}}{\Gamma(2\nu+1)} \int_x^\infty \frac{(t-x)^{(\nu-3)/2-\nu}}{\Gamma((\nu-1)/2-\nu)} e^{\lambda t/2} M_{n/2-\mu, n/2-\nu-1}(\lambda(t-x)) \frac{d^n}{dt^n} [e^{-\lambda t/2} f(t)] dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 522).

1.8-10. Kernels Containing Hermite Polynomials.

$$122. \int_0^x (x-t)^{-1/2} H_{2n}(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad f(0) = 0.$$

Here $H_m(x)$ is the Hermite polynomial (see Supplement 11.17-3).

Solution:

$$y(x) = \frac{(-1)^n n!}{\sqrt{\pi} (2n)!} \left(\frac{d}{dx} \right)^m \int_0^x \frac{(x-t)^{m-3/2}}{\Gamma(m-1/2)} F\left(n, m-\frac{1}{2}; \lambda^2(x-t)\right) f(t) dt,$$

where $m \geq 1$ and $F(a, b; x)$ is the Kummer confluent hypergeometric function (see Supplement 11.9-1).

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 556).

$$123. \int_0^x H_{2n+1}(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad f(0) = f'_x(0) = 0.$$

Solution:

$$y(x) = \frac{(-1)^n n!}{\lambda\sqrt{\pi} (2n+1)!} \left(\frac{d}{dx} \right)^m \int_0^x \frac{(x-t)^{m-5/2}}{\Gamma(m-3/2)} F\left(n, m-\frac{3}{2}; \lambda^2(x-t)\right) f(t) dt,$$

where $m \geq 2$ and $F(a, b; x)$ is the Kummer confluent hypergeometric function (see Supplement 11.9-1).

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 556).

1.8-11. Kernels Containing Chebyshev Polynomials.

$$124. \int_1^x (x^2 - t^2)^{-1/2} T_n \left(\frac{x}{t} \right) y(t) dt = f(x), \quad f(1) = 0, \quad x \geq 1.$$

Here $T_n(x)$ is the Chebyshev polynomials of the first kind (see Supplement 11.17-2).

Solution:

$$y(x) = \frac{2}{\pi} \int_1^x t^n (x^2 - t^2)^{-1/2} T_{n-1} \left(\frac{t}{x} \right) \frac{d}{dt} [t^{1-n} f(t)] dt,$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 499).

$$125. \int_1^x (x^2 - t^2)^{-1/2} T_n \left(\frac{t}{x} \right) y(t) dt = f(x), \quad f(1) = 0, \quad x \geq 1.$$

Solution:

$$y(x) = \frac{2}{\pi} x^{n+1} \frac{d}{dx} \left[x^{-n} \int_1^x (x^2 - t^2)^{-1/2} T_{n+1} \left(\frac{x}{t} \right) f(t) dt \right],$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 499).

$$126. \int_x^1 (t^2 - x^2)^{-1/2} T_n \left(\frac{x}{t} \right) y(t) dt = f(x), \quad f(1) = 0, \quad 0 < x \leq 1.$$

Solution:

$$y(x) = -\frac{2}{\pi} x^{-n} \frac{d}{dx} \left[x^{n+1} \int_x^1 (t^2 - x^2)^{-1/2} T_{n+1} \left(\frac{t}{x} \right) f(t) dt \right],$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 499).

$$127. \int_x^1 (t^2 - x^2)^{-1/2} T_n \left(\frac{t}{x} \right) y(t) dt = f(x), \quad f(1) = 0, \quad 0 < x \leq 1.$$

Solution:

$$y(x) = -\frac{2}{\pi} \int_x^1 t^{1-n} (t^2 - x^2)^{-1/2} T_{n-1} \left(\frac{x}{t} \right) \frac{d}{dt} [t^n f(t)] dt,$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 500).

$$128. \int_0^x (x-t)^{-1/2} T_n \left(2 \frac{x}{t} - 1 \right) y(t) dt = f(x), \quad f(0) = 0, \quad x > 0.$$

Solution:

$$y(x) = \frac{x^n}{\sqrt{\pi} \Gamma(n-1/2)} \frac{d^n}{dx^n} \left[x^{-n} \int_0^x (x-t)^{n-3/2} f(t) dt \right],$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 500).

$$129. \int_x^1 (t-x)^{-1/2} T_n \left(2 \frac{x}{t} - 1 \right) y(t) dt = f(x), \quad f(1) = 0, \quad x \leq 1.$$

Solution:

$$y(x) = \frac{x^n}{\sqrt{\pi}} \left(-\frac{d}{dx} \right)^n \left[x^{-n} \int_x^1 \frac{(t-x)^{n-3/2}}{\Gamma(n-1/2)} f(t) dt \right],$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 501).

$$130. \int_x^1 (t-x)^{-1/2} T_n \left(2 \frac{t}{x} - 1 \right) y(t) dt = f(x), \quad f(1) = 0, \quad x \leq 1.$$

Solution:

$$y(x) = \frac{1}{\sqrt{\pi x}} \left(-\frac{d}{dx} \right)^n \left[x^{n+1/2} \int_x^1 \frac{(t-x)^{n-3/2}}{\Gamma(n-1/2)} t^{-n} f(t) dt \right],$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 501).

1.8-12. Kernels Containing Laguerre Polynomials.

$$131. \int_0^x L_n(\lambda(x-t)) y(t) dt = f(x), \quad f(0) = f'_x(0) = 0, \quad x > 0.$$

Here $L_n(x)$ is the Laguerre polynomial (see Supplement 11.17-1).

Solution:

$$y(x) = e^{\lambda x} \int_0^x L_{n-1}(\lambda(t-x)) e^{-\lambda t} f''_{tt}(t) dt,$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 504).

$$132. \int_x^\infty L_n(\lambda(t-x)) y(t) dt = f(x), \quad f(0) = f'_x(0) = 0, \quad x > 0.$$

Here $L_n(x)$ is the Laguerre polynomial (see Supplement 11.17-1).

Solution:

$$y(x) = e^{-\lambda x} \int_x^\infty L_{n+1}(\lambda(x-t)) e^{\lambda t} f''_{tt}(t) dt,$$

where $n = 1, 2, 3, \dots$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 505).

1.8-13. Kernels Containing Jacobi Theta Functions.

$$133. \int_0^x \vartheta_2(0, x-t) y(t) dt = f(x), \quad f(0) = 0.$$

Here $\vartheta_2(v, q)$ is the Jacobi theta function (see Supplement 11.15-1).

Solution:

$$y(x) = \frac{1}{\pi} \int_0^x \vartheta_3(0, x-t) f'_t(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 551).

$$134. \int_0^x \vartheta_3(0, x-t)y(t) dt = f(x), \quad f(0) = 0.$$

Here $\vartheta_3(v, q)$ is the Jacobi theta function (see Supplement 11.15-1).

Solution:

$$y(x) = \frac{1}{\pi} \int_0^x \vartheta_2(0, x-t)f'_t(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 551).

1.8-14. Kernels Containing Other Special Functions.

$$135. \int_s^x (x-t)^{c-1} F\left(a, b, c; 1 - \frac{x}{t}\right) y(t) dt = f(x).$$

Here $\Phi(a, b, c; z)$ is the Gaussian hypergeometric function (see Supplement 11.10-1).

Solution:

$$y(x) = x^{-a} \frac{d^n}{dx^n} \left\{ x^a \int_s^x \frac{(x-t)^{n-c-1}}{\Gamma(c)\Gamma(n-c)} F\left(-a, n-b, n-c; 1 - \frac{t}{x}\right) f(t) dt \right\},$$

where $0 < c < n$ and $n = 1, 2, \dots$

If the right-hand side of the equation is differentiable sufficiently many times and the conditions $f(s) = f'_x(s) = \dots = f_x^{(n-1)}(s) = 0$ are satisfied, then the solution of the integral equation can be written in the form

$$y(x) = \int_s^x \frac{(x-t)^{n-c-1}}{\Gamma(c)\Gamma(n-c)} F\left(-a, -b, n-c; 1 - \frac{t}{x}\right) f_t^{(n)}(t) dt.$$

⊙ Reference: S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$136. \int_0^x (x-t)^{-(\nu+1)/2} D_\nu(\lambda\sqrt{x-t})y(t) dt = f(x).$$

Here $D_\nu(z)$ is the parabolic cylinder function (see Supplement 11.12-1) and $-1 < \operatorname{Re} \nu < 1$.

Solution:

$$y(x) = \frac{1}{\pi} \int_0^x (x-t)^{(\nu-1)/2} e^{\lambda^2 t/4} D_\nu(\lambda\sqrt{t-x}) \left(\frac{d}{dt} + \frac{\lambda^2}{2} \right) \left(e^{-\lambda^2 t/4} f(t) \right) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 464).

1.9. Equations Whose Kernels Contain Arbitrary Functions

1.9-1. Equations with Degenerate Kernel: $K(x, t) = g_1(x)h_1(t) + g_2(x)h_2(t)$.

$$1. \int_a^x g(x)h(t)y(t) dt = f(x).$$

$$\text{Solution: } y = \frac{1}{h(x)} \frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{1}{g(x)h(x)} f'_x(x) - \frac{g'_x(x)}{g^2(x)h(x)} f(x).$$

$$2. \quad \int_a^x [g(x) - g(t)]y(t) dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = 0$ and $f'_x/g'_x \neq \text{const.}$

$$\text{Solution: } y(x) = \frac{d}{dx} \left[\frac{f'_x(x)}{g'_x(x)} \right].$$

$$3. \quad \int_a^x [g(x) - g(t) + b]y(t) dt = f(x).$$

For $b = 0$, see equation 1.9.2. Assume that $b \neq 0$.

Differentiation with respect to x yields an equation of the form 2.9.2:

$$y(x) + \frac{1}{b}g'_x(x) \int_a^x y(t) dt = \frac{1}{b}f'_x(x).$$

Solution:

$$y(x) = \frac{1}{b}f'_x(x) - \frac{1}{b^2}g'_x(x) \int_a^x \exp\left[\frac{g(t) - g(x)}{b}\right] f'_t(t) dt.$$

$$4. \quad \int_a^x [Ag(x) + Bg(t)]y(t) dt = f(x).$$

For $B = -A$, see equation 1.9.2. Assume that $B \neq -A$.

Solution with $B \neq -A$:

$$y(x) = \frac{\text{sign } g(x)}{A+B} \frac{d}{dx} \left\{ |g(x)|^{-\frac{A}{A+B}} \int_a^x |g(t)|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$5. \quad \int_a^x [Ag(x) + Bg(t) + C]y(t) dt = f(x).$$

For $B = -A$, see equation 1.9.3. Assume that $B \neq -A$ and $(A+B)g(x) + C > 0$.

Solution:

$$y(x) = \frac{d}{dx} \left\{ |(A+B)g(x) + C|^{-\frac{A}{A+B}} \int_a^x |(A+B)g(t) + C|^{-\frac{B}{A+B}} f'_t(t) dt \right\}.$$

$$6. \quad \int_a^x [g(x) + h(t)]y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{d}{dx} \left[\frac{\Phi(x)}{g(x) + h(x)} \int_a^x \frac{f'_t(t) dt}{\Phi(t)} \right], \quad \Phi(x) = \exp \left[\int_a^x \frac{h'_t(t) dt}{g(t) + h(t)} \right].$$

$$7. \quad \int_a^x [g(x) + (x-t)h(x)]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = g(x) + xh(x)$, $h_1(t) = 1$, $g_2(x) = h(x)$, and $h_2(t) = -t$.

Solution:

$$y(x) = \frac{d}{dx} \left\{ \Phi(x) \frac{h(x)}{g(x)} \int_a^x \left[\frac{f(t)}{h(t)} \right]'_t \frac{dt}{\Phi(t)} \right\}, \quad \Phi(t) = \exp \left[- \int_a^x \frac{h(t)}{g(t)} dt \right].$$

$$8. \int_a^x [g(t) + (x-t)h(t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = x$, $h_1(t) = h(t)$, $g_2(x) = 1$, and $h_2(t) = g(t) - th(t)$.

$$9. \int_a^x [g(x) + (Ax^\lambda + Bt^\mu)h(x)]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = g(x) + Ax^\lambda h(x)$, $h_1(t) = 1$, $g_2(x) = h(x)$, and $h_2(t) = Bt^\mu$.

$$10. \int_a^x [g(t) + (Ax^\lambda + Bt^\mu)h(t)]y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = Ax^\lambda$, $h_1(t) = h(t)$, $g_2(x) = 1$, and $h_2(t) = g(t) + Bt^\mu h(t)$.

$$11. \int_a^x [g(x)h(t) - h(x)g(t)]y(t) dt = f(x), \quad f(a) = f'_x(a) = 0.$$

For $g = \text{const}$ or $h = \text{const}$, see equation 1.9.2.

Solution:

$$y(x) = \frac{1}{h} \frac{d}{dx} \left[\frac{(f/h)'_x}{(g/h)'_x} \right], \quad \text{where } f = f(x), \quad g = g(x), \quad h = h(x).$$

Here $Af + Bg + Ch \neq 0$, with A , B , and C being some constants.

$$12. \int_a^x [Ag(x)h(t) + Bg(t)h(x)]y(t) dt = f(x).$$

For $B = -A$, see equation 1.9.11.

Solution with $B \neq -A$:

$$y(x) = \frac{1}{(A+B)h(x)} \frac{d}{dx} \left\{ \left[\frac{h(x)}{g(x)} \right]^{\frac{A}{A+B}} \int_a^x \left[\frac{h(t)}{g(t)} \right]^{\frac{B}{A+B}} \frac{d}{dt} \left[\frac{f(t)}{h(t)} \right] dt \right\}.$$

$$13. \int_a^x \{1 + [g(t) - g(x)]h(x)\}y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = 1 - g(x)h(x)$, $h_1(t) = 1$, $g_2(x) = h(x)$, and $h_2(t) = g(t)$.

Solution:

$$y(x) = \frac{d}{dx} \left\{ h(x)\Phi(x) \int_a^x \left[\frac{f(t)}{h(t)} \right]'_t \frac{dt}{\Phi(t)} \right\}, \quad \Phi(x) = \exp \left[\int_a^x g'_t(t)h(t) dt \right].$$

$$14. \int_a^x \{e^{-\lambda(x-t)} + [e^{\lambda x}g(t) - e^{\lambda t}g(x)]h(x)\}y(t) dt = f(x).$$

This is a special case of equation 1.9.15 with $g_1(x) = e^{\lambda x}h(x)$, $h_1(t) = g(t)$, $g_2(x) = e^{-\lambda x} - g(x)h(x)$, and $h_2(t) = e^{\lambda t}$.

$$15. \int_a^x [g_1(x)h_1(t) + g_2(x)h_2(t)]y(t) dt = f(x).$$

For $g_2/g_1 = \text{const}$ or $h_2/h_1 = \text{const}$, see equation 1.9.1.

1°. Solution with $g_1(x)h_1(x) + g_2(x)h_2(x) \neq 0$ and $f(x) \neq \text{const } g_2(x)$:

$$y(x) = \frac{1}{h_1(x)} \frac{d}{dx} \left\{ \frac{g_2(x)h_1(x)\Phi(x)}{g_1(x)h_1(x) + g_2(x)h_2(x)} \int_a^x \left[\frac{f(t)}{g_2(t)} \right]'_t \frac{dt}{\Phi(t)} \right\}, \quad (1)$$

where

$$\Phi(x) = \exp \left\{ \int_a^x \left[\frac{h_2(t)}{h_1(t)} \right]'_t \frac{g_2(t)h_1(t) dt}{g_1(t)h_1(t) + g_2(t)h_2(t)} \right\}. \quad (2)$$

If $f(x) \equiv \text{const } g_2(x)$, the solution is given by formulas (1) and (2) in which the subscript 1 must be changed by 2 and vice versa.

2°. Solution with $g_1(x)h_1(x) + g_2(x)h_2(x) \equiv 0$:

$$y(x) = \frac{1}{h_1} \frac{d}{dx} \left[\frac{(f/g_2)'_x}{(g_1/g_2)'_x} \right] = -\frac{1}{h_1} \frac{d}{dx} \left[\frac{(f/g_2)'_x}{(h_2/h_1)'_x} \right],$$

where $f = f(x)$, $g_2 = g_2(x)$, $h_1 = h_1(x)$, and $h_2 = h_2(x)$.

1.9-2. Equations with Difference Kernel: $K(x, t) = K(x - t)$.

$$16. \int_a^x K(x - t)y(t) dt = f(x).$$

1°. Let $K(0) = 1$ and $f(a) = 0$. Differentiating the equation with respect to x yields a Volterra equation of the second kind:

$$y(x) + \int_a^x K'_x(x - t)y(t) dt = f'_x(x).$$

The solution of this equation can be represented in the form

$$y(x) = f'_x(x) + \int_a^x R(x - t)f'_t(t) dt.$$

Here the resolvent $R(x)$ is related to the kernel $K(x)$ of the original equation by

$$R(x) = \mathfrak{L}^{-1} \left[\frac{1}{p\tilde{K}(p)} - 1 \right], \quad \tilde{K}(p) = \mathfrak{L}[K(x)],$$

where \mathfrak{L} and \mathfrak{L}^{-1} are the operators of the direct and inverse Laplace transforms, respectively.

$$\tilde{K}(p) = \mathfrak{L}[K(x)] = \int_0^\infty e^{-px} K(x) dx, \quad R(x) = \mathfrak{L}^{-1}[\tilde{R}(p)] = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{R}(p) dp.$$

2°. Let $K(x)$ have an integrable power-law singularity at $x = 0$. Denote by $w = w(x)$ the solution of the simpler auxiliary equation (compared with the original equation) with $a = 0$ and constant right-hand side $f \equiv 1$,

$$\int_0^x K(x - t)w(t) dt = 1. \quad (1)$$

Then the solution of the original integral equation with arbitrary right-hand side is expressed in terms of w as follows:

$$y(x) = \frac{d}{dx} \int_a^x w(x - t)f(t) dt = f(a)w(x - a) + \int_a^x w(x - t)f'_t(t) dt. \quad (2)$$

Remark. The integral equation and its solution (2) form the *Sonine transform pair*.

⊙ References: E. C. Titchmarsh (1986), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 426), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$17. \int_x^\infty K(x-t)y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{d}{dx} \int_x^\infty H(t-x)f(t) dt,$$

where

$$\int_0^x K(t)H(x-t) dt = 1.$$

$$18. \int_{-\infty}^x K(x-t)y(t) dt = Ax^n, \quad n = 0, 1, 2, \dots$$

This is a special case of equation 1.9.20 with $\lambda = 0$.

1°. Solution with $n = 0$:

$$y(x) = \frac{A}{B}, \quad B = \int_0^\infty K(z) dz.$$

2°. Solution with $n = 1$:

$$y(x) = \frac{A}{B}x + \frac{AC}{B^2}, \quad B = \int_0^\infty K(z) dz, \quad C = \int_0^\infty zK(z) dz.$$

3°. Solution with $n = 2$:

$$y_2(x) = \frac{A}{B}x^2 + 2\frac{AC}{B^2}x + 2\frac{AC^2}{B^3} - \frac{AD}{B^2},$$

$$B = \int_0^\infty K(z) dz, \quad C = \int_0^\infty zK(z) dz, \quad D = \int_0^\infty z^2K(z) dz.$$

4°. Solution with $n = 3, 4, \dots$ is given by:

$$y_n(x) = A \left\{ \frac{\partial^n}{\partial \lambda^n} \left[\frac{e^{\lambda x}}{B(\lambda)} \right] \right\}_{\lambda=0}, \quad B(\lambda) = \int_0^\infty K(z)e^{-\lambda z} dz.$$

$$19. \int_{-\infty}^x K(x-t)y(t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = \frac{A}{B}e^{\lambda x}, \quad B = \int_0^\infty K(z)e^{-\lambda z} dz = \mathfrak{L}\{K(z), \lambda\}.$$

$$20. \int_{-\infty}^x K(x-t)y(t) dt = Ax^n e^{\lambda x}, \quad n = 1, 2, \dots$$

1°. Solution with $n = 1$:

$$y_1(x) = \frac{A}{B}xe^{\lambda x} + \frac{AC}{B^2}e^{\lambda x},$$

$$B = \int_0^\infty K(z)e^{-\lambda z} dz, \quad C = \int_0^\infty zK(z)e^{-\lambda z} dz.$$

It is convenient to calculate the coefficients B and C using tables of Laplace transforms according to the formulas $B = \mathfrak{L}\{K(z), \lambda\}$ and $C = \mathfrak{L}\{zK(z), \lambda\}$.

2°. Solution with $n = 2$:

$$y_2(x) = \frac{A}{B}x^2e^{\lambda x} + 2\frac{AC}{B^2}xe^{\lambda x} + \left(2\frac{AC^2}{B^3} - \frac{AD}{B^2}\right)e^{\lambda x},$$

$$B = \int_0^\infty K(z)e^{-\lambda z} dz, \quad C = \int_0^\infty zK(z)e^{-\lambda z} dz, \quad D = \int_0^\infty z^2K(z)e^{-\lambda z} dz.$$

3°. Solution with $n = 3, 4, \dots$ is given by:

$$y_n(x) = \frac{\partial}{\partial \lambda} y_{n-1}(x) = A \frac{\partial^n}{\partial \lambda^n} \left[\frac{e^{\lambda x}}{B(\lambda)} \right], \quad B(\lambda) = \int_0^\infty K(z)e^{-\lambda z} dz.$$

21. $\int_{-\infty}^x K(x-t)y(t) dt = A \cosh(\lambda x).$

Solution:

$$y(x) = \frac{A}{2B_-}e^{\lambda x} + \frac{A}{2B_+}e^{-\lambda x} = \frac{1}{2}\left(\frac{A}{B_-} + \frac{A}{B_+}\right) \cosh(\lambda x) + \frac{1}{2}\left(\frac{A}{B_-} - \frac{A}{B_+}\right) \sinh(\lambda x),$$

$$B_- = \int_0^\infty K(z)e^{-\lambda z} dz, \quad B_+ = \int_0^\infty K(z)e^{\lambda z} dz.$$

22. $\int_{-\infty}^x K(x-t)y(t) dt = A \sinh(\lambda x).$

Solution:

$$y(x) = \frac{A}{2B_-}e^{\lambda x} - \frac{A}{2B_+}e^{-\lambda x} = \frac{1}{2}\left(\frac{A}{B_-} - \frac{A}{B_+}\right) \cosh(\lambda x) + \frac{1}{2}\left(\frac{A}{B_-} + \frac{A}{B_+}\right) \sinh(\lambda x),$$

$$B_- = \int_0^\infty K(z)e^{-\lambda z} dz, \quad B_+ = \int_0^\infty K(z)e^{\lambda z} dz.$$

23. $\int_{-\infty}^x K(x-t)y(t) dt = A \cos(\lambda x).$

Solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \cos(\lambda x) - B_s \sin(\lambda x)],$$

$$B_c = \int_0^\infty K(z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(z) \sin(\lambda z) dz.$$

24. $\int_{-\infty}^x K(x-t)y(t) dt = A \sin(\lambda x).$

Solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \sin(\lambda x) + B_s \cos(\lambda x)],$$

$$B_c = \int_0^\infty K(z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(z) \sin(\lambda z) dz.$$

$$25. \int_{-\infty}^x K(x-t)y(t) dt = Ae^{\mu x} \cos(\lambda x).$$

Solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} e^{\mu x} [B_c \cos(\lambda x) - B_s \sin(\lambda x)],$$

$$B_c = \int_0^\infty K(z)e^{-\mu z} \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(z)e^{-\mu z} \sin(\lambda z) dz.$$

$$26. \int_{-\infty}^x K(x-t)y(t) dt = Ae^{\mu x} \sin(\lambda x).$$

Solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} e^{\mu x} [B_c \sin(\lambda x) + B_s \cos(\lambda x)],$$

$$B_c = \int_0^\infty K(z)e^{-\mu z} \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(z)e^{-\mu z} \sin(\lambda z) dz.$$

$$27. \int_{-\infty}^x K(x-t)y(t) dt = f(x).$$

1°. For a polynomial right-hand side of the equation, $f(x) = \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients. The solution can also be obtained by the formula given in 1.9.18 (item 4°).

2°. For $f(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients. The solution can also be obtained by the formula given in 1.9.20 (item 3°).

3°. For $f(x) = \sum_{k=0}^n A_k \exp(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_k} \exp(\lambda_k x), \quad B_k = \int_0^\infty K(z) \exp(-\lambda_k z) dz.$$

4°. For $f(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

5°. For $f(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

6°. For $f(x) = \sum_{k=0}^n A_k \cos(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_{ck}^2 + B_{sk}^2} [B_{ck} \cos(\lambda_k x) - B_{sk} \sin(\lambda_k x)],$$

$$B_{ck} = \int_0^\infty K(z) \cos(\lambda_k z) dz, \quad B_{sk} = \int_0^\infty K(z) \sin(\lambda_k z) dz.$$

7°. For $f(x) = \sum_{k=0}^n A_k \sin(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_{ck}^2 + B_{sk}^2} [B_{ck} \sin(\lambda_k x) + B_{sk} \cos(\lambda_k x)],$$

$$B_{ck} = \int_0^\infty K(z) \cos(\lambda_k z) dz, \quad B_{sk} = \int_0^\infty K(z) \sin(\lambda_k z) dz.$$

28. $\int_x^\infty K(x-t)y(t) dt = f(x).$

Solution:

$$y(x) = -\frac{d}{dx} \int_x^\infty H(t-x)f(t) dt,$$

where

$$\int_0^x K(t)H(x-t) dt = 1.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 426), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

29. $\int_x^\infty K(x-t)y(t) dt = Ax^n, \quad n = 0, 1, 2, \dots$

This is a special case of equation 1.9.31 with $\lambda = 0$.

1°. Solution with $n = 0$:

$$y(x) = \frac{A}{B}, \quad B = \int_0^\infty K(-z) dz.$$

2°. Solution with $n = 1$:

$$y(x) = \frac{A}{B}x - \frac{AC}{B^2}, \quad B = \int_0^\infty K(-z) dz, \quad C = \int_0^\infty zK(-z) dz.$$

3°. Solution with $n = 2$:

$$y_2(x) = \frac{A}{B}x^2 - 2\frac{AC}{B^2}x + 2\frac{AC^2}{B^3} - \frac{AD}{B^2},$$

$$B = \int_0^\infty K(-z) dz, \quad C = \int_0^\infty zK(-z) dz, \quad D = \int_0^\infty z^2K(-z) dz.$$

4°. Solution with $n = 3, 4, \dots$ is given by

$$y_n(x) = A \left\{ \frac{\partial^n}{\partial \lambda^n} \left[\frac{e^{\lambda x}}{B(\lambda)} \right] \right\}_{\lambda=0}, \quad B(\lambda) = \int_0^\infty K(-z)e^{\lambda z} dz.$$

$$30. \int_x^\infty K(x-t)y(t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = \frac{A}{B}e^{\lambda x}, \quad B = \int_0^\infty K(-z)e^{\lambda z} dz.$$

The expression for B is the Laplace transform of the function $K(-z)$ with parameter $p = -\lambda$ and can be calculated with the aid of tables of Laplace transforms given (e.g., see Supplement 5).

$$31. \int_x^\infty K(x-t)y(t) dt = Ax^n e^{\lambda x}, \quad n = 1, 2, \dots$$

1°. Solution with $n = 1$:

$$y_1(x) = \frac{A}{B}xe^{\lambda x} - \frac{AC}{B^2}e^{\lambda x},$$

$$B = \int_0^\infty K(-z)e^{\lambda z} dz, \quad C = \int_0^\infty zK(-z)e^{\lambda z} dz.$$

It is convenient to calculate the coefficients B and C using tables of Laplace transforms with parameter $p = -\lambda$.

2°. Solution with $n = 2$:

$$y_2(x) = \frac{A}{B}x^2e^{\lambda x} - 2\frac{AC}{B^2}xe^{\lambda x} + \left(2\frac{AC^2}{B^3} - \frac{AD}{B^2}\right)e^{\lambda x},$$

$$B = \int_0^\infty K(-z)e^{\lambda z} dz, \quad C = \int_0^\infty zK(-z)e^{\lambda z} dz, \quad D = \int_0^\infty z^2K(-z)e^{\lambda z} dz.$$

3°. Solution with $n = 3, 4, \dots$ is given by:

$$y_n(x) = \frac{\partial}{\partial \lambda} y_{n-1}(x) = A \frac{\partial^n}{\partial \lambda^n} \left[\frac{e^{\lambda x}}{B(\lambda)} \right], \quad B(\lambda) = \int_0^\infty K(-z)e^{\lambda z} dz.$$

$$32. \int_x^\infty K(x-t)y(t) dt = A \cosh(\lambda x).$$

Solution:

$$y(x) = \frac{A}{2B_+}e^{\lambda x} + \frac{A}{2B_-}e^{-\lambda x} = \frac{1}{2} \left(\frac{A}{B_+} + \frac{A}{B_-} \right) \cosh(\lambda x) + \frac{1}{2} \left(\frac{A}{B_+} - \frac{A}{B_-} \right) \sinh(\lambda x),$$

$$B_+ = \int_0^\infty K(-z)e^{\lambda z} dz, \quad B_- = \int_0^\infty K(-z)e^{-\lambda z} dz.$$

$$33. \int_x^\infty K(x-t)y(t) dt = A \sinh(\lambda x).$$

Solution:

$$y(x) = \frac{A}{2B_+}e^{\lambda x} - \frac{A}{2B_-}e^{-\lambda x} = \frac{1}{2} \left(\frac{A}{B_+} - \frac{A}{B_-} \right) \cosh(\lambda x) + \frac{1}{2} \left(\frac{A}{B_+} + \frac{A}{B_-} \right) \sinh(\lambda x),$$

$$B_+ = \int_0^\infty K(-z)e^{\lambda z} dz, \quad B_- = \int_0^\infty K(-z)e^{-\lambda z} dz.$$

$$34. \int_x^\infty K(x-t)y(t) dt = A \cos(\lambda x).$$

Solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \cos(\lambda x) + B_s \sin(\lambda x)],$$

$$B_c = \int_0^\infty K(-z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z) \sin(\lambda z) dz.$$

$$35. \int_x^\infty K(x-t)y(t) dt = A \sin(\lambda x).$$

Solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \sin(\lambda x) - B_s \cos(\lambda x)],$$

$$B_c = \int_0^\infty K(-z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z) \sin(\lambda z) dz.$$

$$36. \int_x^\infty K(x-t)y(t) dt = A e^{\mu x} \cos(\lambda x).$$

Solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} e^{\mu x} [B_c \cos(\lambda x) + B_s \sin(\lambda x)],$$

$$B_c = \int_0^\infty K(-z) e^{\mu z} \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z) e^{\mu z} \sin(\lambda z) dz.$$

$$37. \int_x^\infty K(x-t)y(t) dt = A e^{\mu x} \sin(\lambda x).$$

Solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} e^{\mu x} [B_c \sin(\lambda x) - B_s \cos(\lambda x)],$$

$$B_c = \int_0^\infty K(-z) e^{\mu z} \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z) e^{\mu z} \sin(\lambda z) dz.$$

$$38. \int_x^\infty K(x-t)y(t) dt = f(x).$$

1°. For a polynomial right-hand side of the equation, $f(x) = \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients. The solution can also be obtained by the formula given in 1.9.29 (item 4°).

2°. For $f(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients. The solution can also be obtained by the formula given in 1.9.31 (item 3°).

3°. For $f(x) = \sum_{k=0}^n A_k \exp(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_k} \exp(\lambda_k x), \quad B_k = \int_0^{\infty} K(-z) \exp(\lambda_k z) dz.$$

4°. For $f(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

5°. For $f(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

6°. For $f(x) = \sum_{k=0}^n A_k \cos(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_{ck}^2 + B_{sk}^2} [B_{ck} \cos(\lambda_k x) + B_{sk} \sin(\lambda_k x)],$$

$$B_{ck} = \int_0^{\infty} K(-z) \cos(\lambda_k z) dz, \quad B_{sk} = \int_0^{\infty} K(-z) \sin(\lambda_k z) dz.$$

7°. For $f(x) = \sum_{k=0}^n A_k \sin(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_{ck}^2 + B_{sk}^2} [B_{ck} \sin(\lambda_k x) - B_{sk} \cos(\lambda_k x)],$$

$$B_{ck} = \int_0^{\infty} K(-z) \cos(\lambda_k z) dz, \quad B_{sk} = \int_0^{\infty} K(-z) \sin(\lambda_k z) dz.$$

8°. For arbitrary right-hand side $f = f(x)$, the solution of the integral equation can be calculated by the formula

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(p)}{\tilde{k}(-p)} e^{px} dp,$$

$$\tilde{f}(p) = \int_0^{\infty} f(x) e^{-px} dx, \quad \tilde{k}(-p) = \int_0^{\infty} K(-z) e^{pz} dz.$$

To calculate $\tilde{f}(p)$ and $\tilde{k}(-p)$, it is convenient to use tables of Laplace transforms, and to determine $y(x)$, tables of inverse Laplace transforms.

1.9-3. Other Equations.

$$39. \int_a^x [g(x) - g(t)]^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

The right-hand side of the equation is assumed to satisfy the conditions $f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0$.

$$\text{Solution: } y(x) = \frac{1}{n!} g'_x(x) \left(\frac{1}{g'_x(x)} \frac{d}{dx} \right)^{n+1} f(x).$$

$$40. \int_a^x \sqrt{g(x) - g(t)} y(t) dt = f(x), \quad f(a) = 0.$$

Solution:

$$y(x) = \frac{2}{\pi} g'_x(x) \left(\frac{1}{g'_x(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{f(t) g'_t(t) dt}{\sqrt{g(x) - g(t)}}.$$

$$41. \int_a^x \frac{y(t) dt}{\sqrt{g(x) - g(t)}} = f(x), \quad g'_x > 0.$$

Solution:

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) g'_t(t) dt}{\sqrt{g(x) - g(t)}}.$$

$$42. \int_a^x \frac{e^{\lambda(x-t)} y(t) dt}{\sqrt{g(x) - g(t)}} = f(x), \quad g'_x > 0.$$

Solution:

$$y(x) = \frac{1}{\pi} e^{\lambda x} \frac{d}{dx} \int_a^x \frac{e^{-\lambda t} f(t) g'_t(t) dt}{\sqrt{g(x) - g(t)}}.$$

$$43. \int_a^x [g(x) - g(t)]^\lambda y(t) dt = f(x), \quad f(a) = 0, \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = k g'_x(x) \left(\frac{1}{g'_x(x)} \frac{d}{dx} \right)^2 \int_a^x \frac{g'_t(t) f(t) dt}{[g(x) - g(t)]^\lambda}, \quad k = \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

$$44. \int_a^x \frac{h(t) y(t) dt}{[g(x) - g(t)]^\lambda} = f(x), \quad g'_x > 0, \quad 0 < \lambda < 1.$$

Solution:

$$y(x) = \frac{\sin(\pi\lambda)}{\pi h(x)} \frac{d}{dx} \int_a^x \frac{f(t) g'_t(t) dt}{[g(x) - g(t)]^{1-\lambda}}.$$

$$45. \int_0^x K\left(\frac{t}{x}\right) y(t) dt = Ax^\lambda + Bx^\mu.$$

Solution:

$$y(x) = \frac{A}{I_\lambda} x^{\lambda-1} + \frac{B}{I_\mu} x^{\mu-1}, \quad I_\lambda = \int_0^1 K(z) z^{\lambda-1} dz, \quad I_\mu = \int_0^1 K(z) z^{\mu-1} dz.$$

$$46. \int_0^x K\left(\frac{t}{x}\right)y(t) dt = P_n(x), \quad P_n(x) = x^\lambda \sum_{m=0}^n A_m x^m.$$

Solution:

$$y(x) = x^\lambda \sum_{m=0}^n \frac{A_m}{I_m} x^{m-1}, \quad I_m = \int_0^1 K(z)z^{\lambda+m-1} dz.$$

The integral I_0 is supposed to converge.

$$47. \int_a^x \{g_1(x)[h_1(t) - h_1(x)] + g_2(x)[h_2(t) - h_2(x)]\} y(t) dt = f(x).$$

This is a special case of equation 1.9.52 with $g_3(x) = -g_1(x)h_1(x) - g_2(x)h_2(x)$ and $h_3(t) = 1$.

The substitution $Y(x) = \int_a^x y(t) dt$ followed by integration by parts leads to an integral equation of the form 1.9.15:

$$\int_a^x \{g_1(x)[h_1(t)]'_t + g_2(x)[h_2(t)]'_t\} Y(t) dt = -f(x).$$

$$48. \int_a^x \{g_1(x)[h_1(t) - e^{\lambda(x-t)}h_1(x)] + g_2(x)[h_2(t) - e^{\lambda(x-t)}h_2(x)]\} y(t) dt = f(x).$$

This is a special case of equation 1.9.52 with $g_3(x) = -e^{\lambda x}[g_1(x)h_1(x) + g_2(x)h_2(x)]$, and $h_3(t) = e^{-\lambda t}$.

The substitution $Y(x) = \int_a^x e^{-\lambda t} y(t) dt$ followed by integration by parts leads to an integral equation of the form 1.9.15:

$$\int_a^x \{g_1(x)[e^{\lambda t}h_1(t)]'_t + g_2(x)[e^{\lambda t}h_2(t)]'_t\} Y(t) dt = -f(x).$$

$$49. \int_a^x [Ag^\lambda(x)g^\mu(t) + Bg^{\lambda+\beta}(x)g^{\mu-\beta}(t) - (A+B)g^{\lambda+\gamma}(x)g^{\mu-\gamma}(t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.52 with $g_1(x) = Ag^\lambda(x)$, $h_1(t) = g^\mu(t)$, $g_2(x) = Bg^{\lambda+\beta}(x)$, $h_2(t) = g^{\mu-\beta}(t)$, $g_3(x) = -(A+B)g^{\lambda+\gamma}(x)$, and $h_3(t) = g^{\mu-\gamma}(t)$.

$$50. \int_a^x [Ag^\lambda(x)h(x)g^\mu(t) + Bg^{\lambda+\beta}(x)h(x)g^{\mu-\beta}(t) - (A+B)g^{\lambda+\gamma}(x)g^{\mu-\gamma}(t)h(t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.52 with $g_1(x) = Ag^\lambda(x)h(x)$, $h_1(t) = g^\mu(t)$, $g_2(x) = Bg^{\lambda+\beta}(x)h(x)$, $h_2(t) = g^{\mu-\beta}(t)h(t)$, $g_3(x) = -(A+B)g^{\lambda+\gamma}(x)$, and $h_3(t) = g^{\mu-\gamma}(t)h(t)$.

$$51. \int_a^x [Ag^\lambda(x)h(x)g^\mu(t) + Bg^{\lambda+\beta}(x)h(t)g^{\mu-\beta}(t) - (A+B)g^{\lambda+\gamma}(x)g^{\mu-\gamma}(t)h(t)] y(t) dt = f(x).$$

This is a special case of equation 1.9.52 with $g_1(x) = Ag^\lambda(x)h(x)$, $h_1(t) = g^\mu(t)$, $g_2(x) = Bg^{\lambda+\beta}(x)$, $h_2(t) = g^{\mu-\beta}(t)h(t)$, $g_3(x) = -(A+B)g^{\lambda+\gamma}(x)$, and $h_3(t) = g^{\mu-\gamma}(t)h(t)$.

$$52. \int_a^x [g_1(x)h_1(t) + g_2(x)h_2(t) + g_3(x)h_3(t)]y(t) dt = f(x),$$

$$\text{where } g_1(x)h_1(x) + g_2(x)h_2(x) + g_3(x)h_3(x) \equiv 0.$$

The substitution $Y(x) = \int_a^x h_3(t)y(t) dt$ followed by integration by parts leads to an integral equation of the form 1.9.15:

$$\int_a^x \left\{ g_1(x) \left[\frac{h_1(t)}{h_3(t)} \right]' + g_2(x) \left[\frac{h_2(t)}{h_3(t)} \right]' \right\} Y(t) dt = -f(x).$$

$$53. \int_{-\infty}^x Q(x-t)e^{\alpha t}y(\xi) dt = Ae^{px}, \quad \xi = e^{\beta t}g(x-t).$$

Solution:

$$y(\xi) = \frac{A}{q} \xi^{\frac{p-\alpha}{\beta}}, \quad q = \int_0^\infty Q(z)[g(z)]^{\frac{p-\alpha}{\beta}} e^{-pz} dz.$$

1.10. Some Formulas and Transformations

1. Let the solution of the integral equation

$$\int_a^x K(x,t)y(t) dt = f(x) \tag{1}$$

have the form

$$y(x) = \mathcal{F}[f(x)], \tag{2}$$

where \mathcal{F} is some linear integro-differential operator. Then the solution of the more complicated integral equation

$$\int_a^x K(x,t)g(x)h(t)y(t) dt = f(x) \tag{3}$$

has the form

$$y(x) = \frac{1}{h(x)} \mathcal{F} \left[\frac{f(x)}{g(x)} \right]. \tag{4}$$

Below are formulas for the solutions of integral equations of the form (3) for some specific functions $g(x)$ and $h(t)$. In all cases, it is assumed that the solution of equation (1) is known and is determined by formula (2).

(a) The solution of the equation

$$\int_a^x K(x,t)(x/t)^\lambda y(t) dt = f(x)$$

has the form

$$y(x) = x^\lambda \mathcal{F}[x^{-\lambda} f(x)].$$

(b) The solution of the equation

$$\int_a^x K(x,t)e^{\lambda(x-t)}y(t) dt = f(x)$$

has the form

$$y(x) = e^{\lambda x} \mathcal{F}[e^{-\lambda x} f(x)].$$

2. Let the solution of the integral equation (1) have the form

$$y(x) = L_1\left(x, \frac{d}{dx}\right)f(x) + L_2\left(x, \frac{d}{dx}\right) \int_a^x R(x, t)f(t) dt, \quad (5)$$

where L_1 and L_2 are some linear differential operators.

The solution of the more complicated integral equation

$$\int_a^x K(\varphi(x), \varphi(t))y(t) dt = f(x), \quad (6)$$

where $\varphi(x)$ is an arbitrary monotone function (differentiable sufficiently many times, $\varphi'_x > 0$), is determined by the formula

$$\begin{aligned} y(x) = & \varphi'_x(x)L_1\left(\varphi(x), \frac{1}{\varphi'_x(x)}\frac{d}{dx}\right)f(x) \\ & + \varphi'_x(x)L_2\left(\varphi(x), \frac{1}{\varphi'_x(x)}\frac{d}{dx}\right) \int_a^x R(\varphi(x), \varphi(t))\varphi'_t(t)f(t) dt. \end{aligned} \quad (7)$$

Below are formulas for the solutions of integral equations of the form (6) for some specific functions $\varphi(x)$. In all cases, it is assumed that the solution of equation (1) is known and is determined by formula (5).

(a) For $\varphi(x) = x^\lambda$,

$$y(x) = \lambda x^{\lambda-1}L_1\left(x^\lambda, \frac{1}{\lambda x^{\lambda-1}}\frac{d}{dx}\right)f(x) + \lambda^2 x^{\lambda-1}L_2\left(x^\lambda, \frac{1}{\lambda x^{\lambda-1}}\frac{d}{dx}\right) \int_a^x R(x^\lambda, t^\lambda)t^{\lambda-1}f(t) dt.$$

(b) For $\varphi(x) = e^{\lambda x}$,

$$y(x) = \lambda e^{\lambda x}L_1\left(e^{\lambda x}, \frac{1}{\lambda e^{\lambda x}}\frac{d}{dx}\right)f(x) + \lambda^2 e^{\lambda x}L_2\left(e^{\lambda x}, \frac{1}{\lambda e^{\lambda x}}\frac{d}{dx}\right) \int_a^x R(e^{\lambda x}, e^{\lambda t})e^{\lambda t}f(t) dt.$$

(c) For $\varphi(x) = \ln(\lambda x)$,

$$y(x) = \frac{1}{x}L_1\left(\ln(\lambda x), x\frac{d}{dx}\right)f(x) + \frac{1}{x}L_2\left(\ln(\lambda x), x\frac{d}{dx}\right) \int_a^x \frac{1}{t}R(\ln(\lambda x), \ln(\lambda t))f(t) dt.$$

(d) For $\varphi(x) = \cos(\lambda x)$,

$$\begin{aligned} y(x) = & -\lambda \sin(\lambda x)L_1\left(\cos(\lambda x), \frac{-1}{\lambda \sin(\lambda x)}\frac{d}{dx}\right)f(x) \\ & + \lambda^2 \sin(\lambda x)L_2\left(\cos(\lambda x), \frac{-1}{\lambda \sin(\lambda x)}\frac{d}{dx}\right) \int_a^x R(\cos(\lambda x), \cos(\lambda t))\sin(\lambda t)f(t) dt. \end{aligned}$$

(e) For $\varphi(x) = \sin(\lambda x)$,

$$\begin{aligned} y(x) = & \lambda \cos(\lambda x)L_1\left(\sin(\lambda x), \frac{1}{\lambda \cos(\lambda x)}\frac{d}{dx}\right)f(x) \\ & + \lambda^2 \cos(\lambda x)L_2\left(\sin(\lambda x), \frac{1}{\lambda \cos(\lambda x)}\frac{d}{dx}\right) \int_a^x R(\sin(\lambda x), \sin(\lambda t))\cos(\lambda t)f(t) dt. \end{aligned}$$

Chapter 2

Linear Equations of the Second Kind with Variable Limit of Integration

► **Notation:** $f = f(x)$, $g = g(x)$, $h = h(x)$, $K = K(x)$, and $M = M(x)$ are arbitrary functions (these may be composite functions of the argument depending on two variables x and t); $A, B, C, D, a, b, c, \alpha, \beta, \gamma, \lambda$, and μ are free parameters; and m and n are nonnegative integers.

2.1. Equations Whose Kernels Contain Power-Law Functions

2.1-1. Kernels Linear in the Arguments x and t .

1. $y(x) - \lambda \int_a^x y(t) dt = f(x).$

Solution:

$$y(x) = f(x) + \lambda \int_a^x e^{\lambda(x-t)} f(t) dt.$$

2. $y(x) + \lambda x \int_a^x y(t) dt = f(x).$

Solution:

$$y(x) = f(x) - \lambda \int_a^x x \exp\left[\frac{1}{2}\lambda(t^2 - x^2)\right] f(t) dt.$$

3. $y(x) + \lambda \int_a^x ty(t) dt = f(x).$

Solution:

$$y(x) = f(x) - \lambda \int_a^x t \exp\left[\frac{1}{2}\lambda(t^2 - x^2)\right] f(t) dt.$$

4. $y(x) + \lambda \int_a^x (x-t)y(t) dt = f(x).$

This is a special case of equation 2.1.34 with $n = 1$.

1°. Solution with $\lambda > 0$:

$$y(x) = f(x) - k \int_a^x \sin[k(x-t)]f(t) dt, \quad k = \sqrt{\lambda}.$$

2°. Solution with $\lambda < 0$:

$$y(x) = f(x) + k \int_a^x \sinh[k(x-t)]f(t) dt, \quad k = \sqrt{-\lambda}.$$

5. $y(x) + \int_a^x [A + B(x-t)] y(t) dt = f(x).$

1°. Solution with $A^2 > 4B$:

$$y(x) = f(x) - \int_a^x R(x-t)f(t) dt,$$

$$R(x) = \exp(-\frac{1}{2}Ax) \left[A \cosh(\beta x) + \frac{2B - A^2}{2\beta} \sinh(\beta x) \right], \quad \beta = \sqrt{\frac{1}{4}A^2 - B}.$$

2°. Solution with $A^2 < 4B$:

$$y(x) = f(x) - \int_a^x R(x-t)f(t) dt,$$

$$R(x) = \exp(-\frac{1}{2}Ax) \left[A \cos(\beta x) + \frac{2B - A^2}{2\beta} \sin(\beta x) \right], \quad \beta = \sqrt{B - \frac{1}{4}A^2}.$$

3°. Solution with $A^2 = 4B$:

$$y(x) = f(x) - \int_a^x R(x-t)f(t) dt, \quad R(x) = \exp(-\frac{1}{2}Ax) (A - \frac{1}{4}A^2x).$$

6. $y(x) - \int_a^x (Ax + Bt + C)y(t) dt = f(x).$

This is a special case of equation 2.9.6 with $g(x) = -Ax$ and $h(t) = -Bt - C$. For $B = -A$ see equation 2.1.5.

By differentiation followed by the substitution $Y(x) = \int_a^x y(t) dt$, the original equation can be reduced to the second-order linear ordinary differential equation

$$Y''_{xx} - [(A+B)x + C]Y'_x - AY = f'_x(x) \quad (1)$$

under the initial conditions

$$Y(a) = 0, \quad Y'_x(a) = f(a). \quad (2)$$

A fundamental system of solutions of the homogeneous equation (1) with $f \equiv 0$ has the form

$$Y_1(x) = \Phi(\alpha, \frac{1}{2}; kz^2), \quad Y_2(x) = \Psi(\alpha, \frac{1}{2}; kz^2),$$

$$\alpha = \frac{A}{2(A+B)}, \quad k = \frac{A+B}{2}, \quad z = x + \frac{C}{A+B},$$

where $\Phi(\alpha, \beta; x)$ and $\Psi(\alpha, \beta; x)$ are degenerate hypergeometric functions.

Solving the homogeneous equation (1) under conditions (2) for an arbitrary function $f = f(x)$ and taking into account the relation $y(x) = Y'_x(x)$, we thus obtain the solution of the integral equation in the form

$$y(x) = f(x) - \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = \frac{\partial^2}{\partial x \partial t} \left[\frac{Y_1(x)Y_2(t) - Y_2(x)Y_1(t)}{W(t)} \right], \quad W(t) = \frac{2\sqrt{\pi k}}{\Gamma(\alpha)} \exp \left[k \left(t + \frac{C}{A+B} \right)^2 \right].$$

2.1-2. Kernels Quadratic in the Arguments x and t .

$$7. \quad y(x) + A \int_a^x x^2 y(t) dt = f(x).$$

This is a special case of equation 2.1.50 with $\lambda = 2$ and $\mu = 0$.

Solution:

$$y(x) = f(x) - A \int_a^x x^2 \exp\left[\frac{1}{3}A(t^3 - x^3)\right] f(t) dt.$$

$$8. \quad y(x) + A \int_a^x x t y(t) dt = f(x).$$

This is a special case of equation 2.1.50 with $\lambda = 1$ and $\mu = 1$.

Solution:

$$y(x) = f(x) - A \int_a^x x t \exp\left[\frac{1}{3}A(t^3 - x^3)\right] f(t) dt.$$

$$9. \quad y(x) + A \int_a^x t^2 y(t) dt = f(x).$$

This is a special case of equation 2.1.50 with $\lambda = 0$ and $\mu = 2$.

Solution:

$$y(x) = f(x) - A \int_a^x t^2 \exp\left[\frac{1}{3}A(t^3 - x^3)\right] f(t) dt.$$

$$10. \quad y(x) + \lambda \int_a^x (x-t)^2 y(t) dt = f(x).$$

This is a special case of equation 2.1.34 with $n = 2$.

Solution:

$$y(x) = f(x) - \int_a^x R(x-t) f(t) dt,$$

$$R(x) = \frac{2}{3} k e^{-2kx} - \frac{2}{3} k e^{kx} \left[\cos(\sqrt{3} kx) - \sqrt{3} \sin(\sqrt{3} kx) \right], \quad k = \left(\frac{1}{4}\lambda\right)^{1/3}.$$

$$11. \quad y(x) + A \int_a^x (x^2 - t^2) y(t) dt = f(x).$$

This is a special case of equation 2.9.5 with $g(x) = Ax^2$.

Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x [u_1'(x)u_2'(t) - u_2'(x)u_1'(t)] f(t) dt,$$

where the primes denote differentiation with respect to the argument specified in the parentheses; $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $u''_{xx} + 2Axu = 0$; and the functions $u_1(x)$ and $u_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of the parameter A :

For $A > 0$,

$$W = 3/\pi, \quad u_1(x) = \sqrt{x} J_{1/3}\left(\sqrt{\frac{8}{9}Ax^{3/2}}\right), \quad u_2(x) = \sqrt{x} Y_{1/3}\left(\sqrt{\frac{8}{9}Ax^{3/2}}\right).$$

For $A < 0$,

$$W = -\frac{3}{2}, \quad u_1(x) = \sqrt{x} I_{1/3}\left(\sqrt{\frac{8}{9}|A|x^{3/2}}\right), \quad u_2(x) = \sqrt{x} K_{1/3}\left(\sqrt{\frac{8}{9}|A|x^{3/2}}\right).$$

$$12. \quad y(x) + A \int_a^x (xt - t^2)y(t) dt = f(x).$$

This is a special case of equation 2.9.4 with $g(t) = At$. Solution:

$$y(x) = f(x) + \frac{A}{W} \int_a^x t [y_1(x)y_2(t) - y_2(x)y_1(t)] f(t) dt,$$

where $y_1(x), y_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $y''_{xx} + Axy = 0$; the functions $y_1(x)$ and $y_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of the parameter A :

For $A > 0$,

$$W = 3/\pi, \quad y_1(x) = \sqrt{x} J_{1/3}(\frac{2}{3}\sqrt{A}x^{3/2}), \quad y_2(x) = \sqrt{x} Y_{1/3}(\frac{2}{3}\sqrt{A}x^{3/2}).$$

For $A < 0$,

$$W = -\frac{3}{2}, \quad y_1(x) = \sqrt{x} I_{1/3}(\frac{2}{3}\sqrt{|A|}x^{3/2}), \quad y_2(x) = \sqrt{x} K_{1/3}(\frac{2}{3}\sqrt{|A|}x^{3/2}).$$

$$13. \quad y(x) + A \int_a^x (x^2 - xt)y(t) dt = f(x).$$

This is a special case of equation 2.9.3 with $g(x) = Ax$. Solution:

$$y(x) = f(x) + \frac{A}{W} \int_a^x x [y_1(x)y_2(t) - y_2(x)y_1(t)] f(t) dt,$$

where $y_1(x), y_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $y''_{xx} + Axy = 0$; the functions $y_1(x)$ and $y_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of the parameter A :

For $A > 0$,

$$W = 3/\pi, \quad y_1(x) = \sqrt{x} J_{1/3}(\frac{2}{3}\sqrt{A}x^{3/2}), \quad y_2(x) = \sqrt{x} Y_{1/3}(\frac{2}{3}\sqrt{A}x^{3/2}).$$

For $A < 0$,

$$W = -\frac{3}{2}, \quad y_1(x) = \sqrt{x} I_{1/3}(\frac{2}{3}\sqrt{|A|}x^{3/2}), \quad y_2(x) = \sqrt{x} K_{1/3}(\frac{2}{3}\sqrt{|A|}x^{3/2}).$$

$$14. \quad y(x) + A \int_a^x (t^2 - 3x^2)y(t) dt = f(x).$$

This is a special case of equation 2.1.55 with $\lambda = 1$ and $\mu = 2$.

$$15. \quad y(x) + A \int_a^x (2xt - 3x^2)y(t) dt = f(x).$$

This is a special case of equation 2.1.55 with $\lambda = 2$ and $\mu = 1$.

$$16. \quad y(x) - \int_a^x (ABxt - ABx^2 + Ax + B)y(t) dt = f(x).$$

This is a special case of equation 2.9.16 with $g(x) = Ax$ and $h(x) = B$.

Solution:

$$y(x) = f(x) + \int_a^x R(x, t)f(t) dt,$$

$$R(x, t) = (Ax + B) \exp\left[\frac{1}{2}A(x^2 - t^2)\right] + B^2 \int_t^x \exp\left[\frac{1}{2}A(s^2 - t^2) + B(x - s)\right] ds.$$

$$17. \quad y(x) + \int_a^x (Ax^2 - At^2 + Bx - Ct + D)y(t) dt = f(x).$$

This is a special case of equation 2.9.6 with $g(x) = Ax^2 + Bx + D$ and $h(t) = -At^2 - Ct$.

Solution:

$$y(x) = f(x) + \int_a^x \frac{\partial^2}{\partial x \partial t} \left[\frac{Y_1(x)Y_2(t) - Y_2(x)Y_1(t)}{W(t)} \right] f(t) dt.$$

Here $Y_1(x), Y_2(x)$ is a fundamental system of solutions of the second-order homogeneous ordinary differential equation $Y''_{xx} + [(B-C)x + D]Y'_x + (2Ax + B)Y = 0$ (see A. D. Polyanin and V. F. Zaitsev (2003) for details about this equation):

$$Y_1(x) = \exp(-kx)\Phi\left(\alpha, \frac{1}{2}; \frac{1}{2}(C-B)z^2\right), \quad Y_2(x) = \exp(-kx)\Psi\left(\alpha, \frac{1}{2}; \frac{1}{2}(C-B)z^2\right),$$

$$W(x) = -\frac{\sqrt{2\pi(C-B)}}{\Gamma(\alpha)} \exp\left[\frac{1}{2}(C-B)z^2 - 2kx\right], \quad k = \frac{2A}{B-C},$$

$$\alpha = -\frac{4A^2 + 2AD(C-B) + B(C-B)^2}{2(C-B)^3}, \quad z = x - \frac{4A + (C-B)D}{(C-B)^2},$$

where $\Phi(\alpha, \beta; x)$ and $\Psi(\alpha, \beta; x)$ are degenerate hypergeometric functions and $\Gamma(\alpha)$ is the gamma function.

$$18. \quad y(x) - \int_a^x [Ax + B + (Cx + D)(x - t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.11 with $g(x) = Ax + B$ and $h(x) = Cx + D$.

Solution with $A \neq 0$:

$$y(x) = f(x) + \int_a^x [Y_2''(x)Y_1(t) - Y_1''(x)Y_2(t)] \frac{f(t)}{W(t)} dt.$$

Here $Y_1(x), Y_2(x)$ is a fundamental system of solutions of the second-order homogeneous ordinary differential equation $Y''_{xx} - (Ax + B)Y'_x - (Cx + D)Y = 0$ (see A. D. Polyanin and V. F. Zaitsev (2003) for details about this equation):

$$Y_1(x) = \exp(-kx)\Phi\left(\alpha, \frac{1}{2}; \frac{1}{2}Az^2\right), \quad Y_2(x) = \exp(-kx)\Psi\left(\alpha, \frac{1}{2}; \frac{1}{2}Az^2\right),$$

$$W(x) = -\sqrt{2\pi A} [\Gamma(\alpha)]^{-1} \exp\left(\frac{1}{2}Az^2 - 2kx\right), \quad k = C/A,$$

$$\alpha = \frac{1}{2}(A^2D - ABC - C^2)A^{-3}, \quad z = x + (AB + 2C)A^{-2},$$

where $\Phi(\alpha, \beta; x)$ and $\Psi(\alpha, \beta; x)$ are degenerate hypergeometric functions, $\Gamma(\alpha)$ is the gamma function.

$$19. \quad y(x) + \int_a^x [At + B + (Ct + D)(t - x)] y(t) dt = f(x).$$

This is a special case of equation 2.9.12 with $g(t) = -At - B$ and $h(t) = -Ct - D$.

Solution with $A \neq 0$:

$$y(x) = f(x) - \int_a^x [Y_1(x)Y_2''(t) - Y_1''(t)Y_2(x)] \frac{f(t)}{W(x)} dt.$$

Here $Y_1(x), Y_2(x)$ is a fundamental system of solutions of the second-order homogeneous ordinary differential equation $Y''_{xx} - (Ax + B)Y'_x - (Cx + D)Y = 0$ (see A. D. Polyanin and V. F. Zaitsev (2003) for details about this equation):

$$Y_1(x) = \exp(-kx)\Phi\left(\alpha, \frac{1}{2}; \frac{1}{2}Az^2\right), \quad Y_2(x) = \exp(-kx)\Psi\left(\alpha, \frac{1}{2}; \frac{1}{2}Az^2\right),$$

$$W(x) = -\sqrt{2\pi A} [\Gamma(\alpha)]^{-1} \exp\left(\frac{1}{2}Az^2 - 2kx\right), \quad k = C/A,$$

$$\alpha = \frac{1}{2}(A^2D - ABC - C^2)A^{-3}, \quad z = x + (AB + 2C)A^{-2},$$

where $\Phi(\alpha, \beta; x)$ and $\Psi(\alpha, \beta; x)$ are degenerate hypergeometric functions and $\Gamma(\alpha)$ is the gamma function.

2.1-3. Kernels Cubic in the Arguments x and t .

20. $y(x) + A \int_a^x x^3 y(t) dt = f(x).$

Solution:

$$y(x) = f(x) - A \int_a^x x^3 \exp\left[\frac{1}{4}A(t^4 - x^4)\right] f(t) dt.$$

21. $y(x) + A \int_a^x x^2 t y(t) dt = f(x).$

Solution:

$$y(x) = f(x) - A \int_a^x x^2 t \exp\left[\frac{1}{4}A(t^4 - x^4)\right] f(t) dt.$$

22. $y(x) + A \int_a^x x t^2 y(t) dt = f(x).$

Solution:

$$y(x) = f(x) - A \int_a^x x t^2 \exp\left[\frac{1}{4}A(t^4 - x^4)\right] f(t) dt.$$

23. $y(x) + A \int_a^x t^3 y(t) dt = f(x).$

Solution:

$$y(x) = f(x) - A \int_a^x t^3 \exp\left[\frac{1}{4}A(t^4 - x^4)\right] f(t) dt.$$

24. $y(x) + \lambda \int_a^x (x-t)^3 y(t) dt = f(x).$

This is a special case of equation 2.1.34 with $n = 3$.

Solution:

$$y(x) = f(x) - \int_a^x R(x-t) f(t) dt,$$

where

$$R(x) = \begin{cases} k [\cosh(kx) \sin(kx) - \sinh(kx) \cos(kx)], & k = \left(\frac{3}{2}\lambda\right)^{1/4} \text{ for } \lambda > 0, \\ \frac{1}{2}s [\sin(sx) - \sinh(sx)], & s = (-6\lambda)^{1/4} \text{ for } \lambda < 0. \end{cases}$$

25. $y(x) + A \int_a^x (x^3 - t^3) y(t) dt = f(x).$

This is a special case of equation 2.1.52 with $\lambda = 3$.

26. $y(x) - A \int_a^x (4x^3 - t^3) y(t) dt = f(x).$

This is a special case of equation 2.1.55 with $\lambda = 1$ and $\mu = 3$.

27. $y(x) + A \int_a^x (x t^2 - t^3) y(t) dt = f(x).$

This is a special case of equation 2.1.49 with $\lambda = 2$.

$$28. \quad y(x) + A \int_a^x (x^2 t - t^3) y(t) dt = f(x).$$

The transformation $z = x^2$, $\tau = t^2$, $y(x) = w(z)$ leads to an equation of the form 2.1.4:

$$w(z) + \frac{1}{2} A \int_{a^2}^z (z - \tau) w(\tau) d\tau = F(z), \quad F(z) = f(x).$$

$$29. \quad y(x) + \int_a^x (Ax^2 t + Bt^3) y(t) dt = f(x).$$

The transformation $z = x^2$, $\tau = t^2$, $y(x) = w(z)$ leads to an equation of the form 2.1.6:

$$w(z) + \int_{a^2}^z \left(\frac{1}{2} Az + \frac{1}{2} B\tau \right) w(\tau) d\tau = F(z), \quad F(z) = f(x).$$

$$30. \quad y(x) + B \int_a^x (2x^3 - xt^2) y(t) dt = f(x).$$

This is a special case of equation 2.1.55 with $\lambda = 2$, $\mu = 2$, and $B = -2A$.

$$31. \quad y(x) - A \int_a^x (4x^3 - 3x^2 t) y(t) dt = f(x).$$

This is a special case of equation 2.1.55 with $\lambda = 3$ and $\mu = 1$.

$$32. \quad y(x) + \int_a^x (ABx^3 - ABx^2 t - Ax^2 - B) y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $g(x) = Ax^2$ and $\lambda = B$.

Solution:

$$y(x) = f(x) + \int_a^x R(x-t) f(t) dt,$$

$$R(x, t) = (Ax^2 + B) \exp\left[\frac{1}{3} A(x^3 - t^3)\right] + B^2 \int_t^x \exp\left[\frac{1}{3} A(s^3 - t^3) + B(x-s)\right] ds.$$

$$33. \quad y(x) + \int_a^x (ABxt^2 - ABt^3 + At^2 + B) y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $g(t) = At^2$ and $\lambda = B$.

Solution:

$$y(x) = f(x) + \int_a^x R(x-t) f(t) dt,$$

$$R(x, t) = -(At^2 + B) \exp\left[\frac{1}{3} A(t^3 - x^3)\right] + B^2 \int_t^x \exp\left[\frac{1}{3} A(s^3 - x^3) + B(t-s)\right] ds.$$

2.1-4. Kernels Containing Higher-Order Polynomials in x and t .

$$34. \quad y(x) + A \int_a^x (x-t)^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

1°. Differentiating the equation $n + 1$ times with respect to x yields an $(n + 1)$ st-order linear ordinary differential equation with constant coefficients for $y = y(x)$:

$$y_x^{(n+1)} + An! y = f_x^{(n+1)}(x).$$

This equation under the initial conditions $y(a) = f(a)$, $y'_x(a) = f'_x(a)$, \dots , $y_x^{(n)}(a) = f_x^{(n)}(a)$ determines the solution of the original integral equation.

2°. Solution:

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt,$$

$$R(x) = \frac{1}{n+1} \sum_{k=0}^n \exp(\sigma_k x) [\sigma_k \cos(\beta_k x) - \beta_k \sin(\beta_k x)],$$

where the coefficients σ_k and β_k are given by

$$\begin{aligned} \sigma_k &= |An!|^{\frac{1}{n+1}} \cos\left(\frac{2\pi k}{n+1}\right), & \beta_k &= |An!|^{\frac{1}{n+1}} \sin\left(\frac{2\pi k}{n+1}\right) & \text{for } A < 0, \\ \sigma_k &= |An!|^{\frac{1}{n+1}} \cos\left(\frac{2\pi k + \pi}{n+1}\right), & \beta_k &= |An!|^{\frac{1}{n+1}} \sin\left(\frac{2\pi k + \pi}{n+1}\right) & \text{for } A > 0. \end{aligned}$$

35. $y(x) + A \int_x^\infty (t-x)^n y(t) dt = f(x), \quad n = 1, 2, \dots$

The Picard–Goursat equation. This is a special case of equation 2.9.62 with $K(z) = A(-z)^n$.

1°. A solution of the homogeneous equation ($f \equiv 0$) is

$$y(x) = C e^{-\lambda x}, \quad \lambda = (-An!)^{\frac{1}{n+1}},$$

where C is an arbitrary constant and $A < 0$. This is a unique solution for $n = 0, 1, 2, 3$.

The general solution of the homogeneous equation for any sign of A has the form

$$y(x) = \sum_{k=1}^s C_k \exp(-\lambda_k x). \quad (1)$$

Here C_k are arbitrary constants and λ_k are the roots of the algebraic equation $\lambda^{n+1} + An! = 0$ that satisfy the condition $\operatorname{Re} \lambda_k > 0$. The number of terms in (1) is determined by the inequality $s \leq 2\left[\frac{n}{4}\right] + 1$, where $[a]$ stands for the integral part of a number a . For more details about the solution of the homogeneous Picard–Goursat equation, see Subsection 11.11-1 (Example 1).

2°. For $f(x) = \sum_{k=1}^m a_k \exp(-\beta_k x)$, where $\beta_k > 0$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^m \frac{a_k \beta_k^{n+1}}{\beta_k^{n+1} + An!} \exp(-\beta_k x), \quad (2)$$

where $\beta_k^{n+1} + An! \neq 0$. For $A > 0$, this formula can also be used for arbitrary $f(x)$ expandable into a convergent exponential series (which corresponds to $m = \infty$).

3°. For $f(x) = e^{-\beta x} \sum_{k=1}^m a_k x^k$, where $\beta > 0$, a solution of the equation has the form

$$y(x) = e^{-\beta x} \sum_{k=0}^m B_k x^k, \quad (3)$$

where the constants B_k are found by the method of undetermined coefficients. The solution can also be constructed using the formulas given in item 3°, equation 2.9.55.

4°. For $f(x) = \cos(\beta x) \sum_{k=1}^m a_k \exp(-\mu_k x)$, a solution of the equation has the form

$$y(x) = \cos(\beta x) \sum_{k=1}^m B_k \exp(-\mu_k x) + \sin(\beta x) \sum_{k=1}^m C_k \exp(-\mu_k x), \quad (4)$$

where the constants B_k and C_k are found by the method of undetermined coefficients. The solution can also be constructed using the formulas given in 2.9.60.

5°. For $f(x) = \sin(\beta x) \sum_{k=1}^m a_k \exp(-\mu_k x)$, a solution of the equation has the form

$$y(x) = \cos(\beta x) \sum_{k=1}^m B_k \exp(-\mu_k x) + \sin(\beta x) \sum_{k=1}^m C_k \exp(-\mu_k x), \quad (5)$$

where the constants B_k and C_k are found by the method of undetermined coefficients. The solution can also be constructed using the formulas given in 2.9.61.

6°. To obtain the general solution in item 2°–5°, the solution (1) of the homogeneous equation must be added to each right-hand side of (2)–(5).

$$36. \quad y(x) + A \int_a^x (x-t)t^n y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 2.1.49 with $\lambda = n$.

$$37. \quad y(x) + A \int_a^x (x^n - t^n) y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 2.1.52 with $\lambda = n$.

$$38. \quad y(x) + \int_a^x (ABx^{n+1} - ABx^n t - Ax^n - B) y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 2.9.7 with $g(x) = Ax^n$ and $\lambda = B$.

Solution:

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt,$$

$$R(x, t) = (Ax^n + B) \exp\left[\frac{A}{n+1}(x^{n+1} - t^{n+1})\right] + B^2 \int_t^x \exp\left[\frac{A}{n+1}(s^{n+1} - t^{n+1}) + B(x-s)\right] ds.$$

$$39. \quad y(x) + \int_a^x (ABxt^n - ABt^{n+1} + At^n + B) y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 2.9.8 with $g(t) = At^n$ and $\lambda = B$.

Solution:

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt,$$

$$R(x, t) = -(At^n + B) \exp\left[\frac{A}{n+1}(t^{n+1} - x^{n+1})\right] + B^2 \int_t^x \exp\left[\frac{A}{n+1}(s^{n+1} - x^{n+1}) + B(t-s)\right] ds.$$

2.1-5. Kernels Containing Rational Functions.

$$40. \quad y(x) + x^{-3} \int_a^x t [2Ax + (1-A)t] y(t) dt = f(x).$$

This equation can be obtained by differentiating the equation

$$\int_a^x [Ax^2 t + (1-A)xt^2] y(t) dt = F(x), \quad F(x) = \int_a^x t^3 f(t) dt,$$

which has the form 1.1.17:

Solution:

$$y(x) = \frac{1}{x} \frac{d}{dx} \left[x^{-A} \int_a^x t^{A-1} \varphi'_t(t) dt \right], \quad \varphi(x) = \frac{1}{x} \int_a^x t^3 f(t) dt.$$

$$41. \quad y(x) - \lambda \int_0^x \frac{y(t) dt}{x+t} = f(x).$$

Dixon's equation. This is a special case of equation 2.1.62 with $a = b = 1$ and $\mu = 0$.

1°. The solution of the homogeneous equation ($f \equiv 0$) is

$$y(x) = Cx^\beta \quad (\beta > -1, \lambda > 0). \quad (1)$$

Here C is an arbitrary constant, and $\beta = \beta(\lambda)$ is determined by the transcendental equation

$$\lambda I(\beta) = 1, \quad \text{where} \quad I(\beta) = \int_0^1 \frac{z^\beta dz}{1+z}. \quad (2)$$

2°. For a polynomial right-hand side,

$$f(x) = \sum_{n=0}^N A_n x^n$$

the solution bounded at zero is given by

$$y(x) = \begin{cases} \sum_{n=0}^N \frac{A_n}{1-(\lambda/\lambda_n)} x^n & \text{for } \lambda < \lambda_0, \\ \sum_{n=0}^N \frac{A_n}{1-(\lambda/\lambda_n)} x^n + Cx^\beta & \text{for } \lambda > \lambda_0 \text{ and } \lambda \neq \lambda_n, \end{cases}$$

$$\lambda_n = \frac{1}{I(n)}, \quad I(n) = (-1)^n \left[\ln 2 + \sum_{m=1}^n \frac{(-1)^m}{m} \right],$$

where C is an arbitrary constant, and $\beta = \beta(\lambda)$ is determined by the transcendental equation (2).

For special $\lambda = \lambda_n$ ($n = 1, 2, \dots$), the solution differs in one term and has the form

$$y(x) = \sum_{m=0}^{n-1} \frac{A_m}{1-(\lambda_n/\lambda_m)} x^m + \sum_{m=n+1}^N \frac{A_m}{1-(\lambda_n/\lambda_m)} x^m - A_n \frac{\bar{\lambda}_n}{\lambda_n} x^n \ln x + Cx^n,$$

where $\bar{\lambda}_n = (-1)^{n+1} \left[\frac{\pi^2}{12} + \sum_{k=1}^n \frac{(-1)^k}{k^2} \right]^{-1}$.

Remark. For arbitrary $f(x)$, expandable into power series, the formulas of item 2° can be used, in which one should set $N = \infty$. In this case, the radius of convergence of the solution $y(x)$ is equal to the radius of convergence of $f(x)$.

3°. For logarithmic-polynomial right-hand side,

$$f(x) = \ln x \left(\sum_{n=0}^N A_n x^n \right),$$

the solution with logarithmic singularity at zero is given by

$$y(x) = \begin{cases} \ln x \sum_{n=0}^N \frac{A_n}{1 - (\lambda/\lambda_n)} x^n + \sum_{n=0}^N \frac{A_n D_n \lambda}{[1 - (\lambda/\lambda_n)]^2} x^n & \text{for } \lambda < \lambda_0, \\ \ln x \sum_{n=0}^N \frac{A_n}{1 - (\lambda/\lambda_n)} x^n + \sum_{n=0}^N \frac{A_n D_n \lambda}{[1 - (\lambda/\lambda_n)]^2} x^n + Cx^\beta & \text{for } \lambda > \lambda_0 \text{ and } \lambda \neq \lambda_n, \end{cases}$$

$$\lambda_n = \frac{1}{I(n)}, \quad I(n) = (-1)^n \left[\ln 2 + \sum_{k=1}^n \frac{(-1)^k}{k} \right], \quad D_n = (-1)^{n+1} \left[\frac{\pi^2}{12} + \sum_{k=1}^n \frac{(-1)^k}{k^2} \right].$$

4°. For arbitrary $f(x)$, the transformation

$$x = \frac{1}{2}e^{2z}, \quad t = \frac{1}{2}e^{2\tau}, \quad y(x) = e^{-z}w(z), \quad f(x) = e^{-z}g(z)$$

leads to an integral equation with difference kernel of the form 2.9.51:

$$w(z) - \lambda \int_{-\infty}^z \frac{w(\tau) d\tau}{\cosh(z - \tau)} = g(z).$$

42. $y(x) - \lambda \int_a^x \frac{x+b}{t+b} y(t) dt = f(x).$

This is a special case of equation 2.9.1 with $g(x) = x + b$.

Solution:

$$y(x) = f(x) + \lambda \int_a^x \frac{x+b}{t+b} e^{\lambda(x-t)} f(t) dt.$$

43. $y(x) = \frac{2}{(1-\lambda^2)x^2} \int_{\lambda x}^x \frac{t}{1+t} y(t) dt.$

This equation is encountered in nuclear physics and describes deceleration of neutrons in matter.

1°. Solution with $\lambda = 0$:

$$y(x) = \frac{C}{(1+x)^2},$$

where C is an arbitrary constant.

2°. For $\lambda \neq 0$, the solution can be found in the series form

$$y(x) = \sum_{n=0}^{\infty} A_n x^n.$$

2.1-6. Kernels Containing Square Roots and Fractional Powers.

$$44. \quad y(x) + A \int_a^x (x-t)\sqrt{t} y(t) dt = f(x).$$

This is a special case of equation 2.1.49 with $\lambda = \frac{1}{2}$.

$$45. \quad y(x) + A \int_a^x (\sqrt{x} - \sqrt{t}) y(t) dt = f(x).$$

This is a special case of equation 2.1.52 with $\lambda = \frac{1}{2}$.

$$46. \quad y(x) + \lambda \int_a^x \frac{y(t) dt}{\sqrt{x-t}} = f(x).$$

Abel's equation of the second kind. This equation is encountered in problems of heat and mass transfer.

Solution:

$$y(x) = F(x) + \pi \lambda^2 \int_a^x \exp[\pi \lambda^2 (x-t)] F(t) dt,$$

where

$$F(x) = f(x) - \lambda \int_a^x \frac{f(t) dt}{\sqrt{x-t}}.$$

⊙ References: H. Brakhage, K. Nickel, and P. Rieder (1965), Yu. I. Babenko (1986).

$$47. \quad y(x) - \lambda \int_0^x \frac{y(t) dt}{\sqrt{ax^2 + bt^2}} = f(x), \quad a > 0, \quad b > 0.$$

1°. The solution of the homogeneous equation ($f \equiv 0$) is

$$y(x) = Cx^\beta \quad (\beta > -1, \quad \lambda > 0). \quad (1)$$

Here C is an arbitrary constant, and $\beta = \beta(\lambda)$ is determined by the transcendental equation

$$\lambda I(\beta) = 1, \quad \text{where} \quad I(\beta) = \int_0^1 \frac{z^\beta dz}{\sqrt{a + bz^2}}. \quad (2)$$

2°. For a polynomial right-hand side,

$$f(x) = \sum_{n=0}^N A_n x^n$$

the solution bounded at zero is given by

$$y(x) = \begin{cases} \sum_{n=0}^N \frac{A_n}{1 - (\lambda/\lambda_n)} x^n & \text{for } \lambda < \lambda_0, \\ \sum_{n=0}^N \frac{A_n}{1 - (\lambda/\lambda_n)} x^n + Cx^\beta & \text{for } \lambda > \lambda_0 \text{ and } \lambda \neq \lambda_n, \end{cases}$$

$$\lambda_0 = \frac{\sqrt{b}}{\text{Arsinh}(\sqrt{b/a})}, \quad \lambda_n = \frac{1}{I(n)}, \quad I(n) = \int_0^1 \frac{z^n dz}{\sqrt{a + bz^2}}.$$

Here C is an arbitrary constant, and $\beta = \beta(\lambda)$ is determined by the transcendental equation (2).

3°. For special $\lambda = \lambda_n$ ($n = 1, 2, \dots$), the solution differs in one term and has the form

$$y(x) = \sum_{m=0}^{n-1} \frac{A_m}{1 - (\lambda_n/\lambda_m)} x^m + \sum_{m=n+1}^N \frac{A_m}{1 - (\lambda_n/\lambda_m)} x^m - A_n \frac{\bar{\lambda}_n}{\lambda_n} x^n \ln x + Cx^n,$$

where $\bar{\lambda}_n = \left[\int_0^1 \frac{z^n \ln z dz}{\sqrt{a + bz^2}} \right]^{-1}$.

4°. For arbitrary $f(x)$, expandable into power series, the formulas of item 2° can be used, in which one should set $N = \infty$. In this case, the radius of convergence of the solution $y(x)$ is equal to the radius of convergence of $f(x)$.

48. $y(x) + \lambda \int_a^x \frac{y(t) dt}{(x-t)^{3/4}} = f(x).$

This equation admits solution by quadratures (see equation 2.1.60 and Example 2 in Subsection 11.4-2).

2.1-7. Kernels Containing Arbitrary Powers.

49. $y(x) + A \int_a^x (x-t)t^\lambda y(t) dt = f(x).$

This is a special case of equation 2.9.4 with $g(t) = At^\lambda$.

Solution:

$$y(x) = f(x) + \frac{A}{W} \int_a^x [y_1(x)y_2(t) - y_2(x)y_1(t)] t^\lambda f(t) dt,$$

where $y_1(x), y_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $y''_{xx} + Ax^\lambda y = 0$; the functions $y_1(x)$ and $y_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of A :

For $A > 0$,

$$W = \frac{2q}{\pi}, \quad y_1(x) = \sqrt{x} J_{\frac{1}{2q}} \left(\frac{\sqrt{A}}{q} x^q \right), \quad y_2(x) = \sqrt{x} Y_{\frac{1}{2q}} \left(\frac{\sqrt{A}}{q} x^q \right), \quad q = \frac{\lambda+2}{2},$$

For $A < 0$,

$$W = -q, \quad y_1(x) = \sqrt{x} I_{\frac{1}{2q}} \left(\frac{\sqrt{|A|}}{q} x^q \right), \quad y_2(x) = \sqrt{x} K_{\frac{1}{2q}} \left(\frac{\sqrt{|A|}}{q} x^q \right), \quad q = \frac{\lambda+2}{2}.$$

50. $y(x) + A \int_a^x x^\lambda t^\mu y(t) dt = f(x).$

This is a special case of equation 2.9.2 with $g(x) = -Ax^\lambda$ and $h(t) = t^\mu$ (λ and μ are arbitrary numbers).

Solution:

$$y(x) = f(x) - \int_a^x R(x,t) f(t) dt,$$

$$R(x,t) = \begin{cases} Ax^\lambda t^\mu \exp \left[\frac{A}{\lambda + \mu + 1} (t^{\lambda+\mu+1} - x^{\lambda+\mu+1}) \right] & \text{for } \lambda + \mu + 1 \neq 0, \\ Ax^{\lambda-A} t^{\mu+A} & \text{for } \lambda + \mu + 1 = 0. \end{cases}$$

$$51. \quad y(x) + A \int_a^x (x-t)x^\lambda t^\mu y(t) dt = f(x).$$

The substitution $u(x) = x^{-\lambda}y(x)$ leads to an equation of the form 2.1.49:

$$u(x) + A \int_a^x (x-t)t^{\lambda+\mu}u(t) dt = f(x)x^{-\lambda}.$$

$$52. \quad y(x) + A \int_a^x (x^\lambda - t^\lambda)y(t) dt = f(x).$$

This is a special case of equation 2.9.5 with $g(x) = Ax^\lambda$.

Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x [u'_1(x)u'_2(t) - u'_2(x)u'_1(t)] f(t) dt,$$

where the primes denote differentiation with respect to the argument specified in the parentheses, and $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $u''_{xx} + A\lambda x^{\lambda-1}u = 0$; the functions $u_1(x)$ and $u_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of A :

For $A\lambda > 0$,

$$W = \frac{2q}{\pi}, \quad u_1(x) = \sqrt{x} J_{\frac{1}{2q}} \left(\frac{\sqrt{A\lambda}}{q} x^q \right), \quad u_2(x) = \sqrt{x} Y_{\frac{1}{2q}} \left(\frac{\sqrt{A\lambda}}{q} x^q \right), \quad q = \frac{\lambda+1}{2},$$

For $A\lambda < 0$,

$$W = -q, \quad u_1(x) = \sqrt{x} I_{\frac{1}{2q}} \left(\frac{\sqrt{|A\lambda|}}{q} x^q \right), \quad u_2(x) = \sqrt{x} \lambda K_{\frac{1}{2q}} \left(\frac{\sqrt{|A\lambda|}}{q} x^q \right), \quad q = \frac{\lambda+1}{2}.$$

$$53. \quad y(x) - \int_a^x (Ax^\lambda t^{\lambda-1} + Bt^{2\lambda-1})y(t) dt = f(x).$$

The transformation

$$z = x^\lambda, \quad \tau = t^\lambda, \quad y(x) = Y(z)$$

leads to an equation of the form 2.1.6:

$$Y(z) - \int_b^z \left(\frac{A}{\lambda} z + \frac{B}{\lambda} \tau \right) Y(\tau) d\tau = F(z), \quad F(z) = f(x), \quad b = a^\lambda.$$

$$54. \quad y(x) - \int_a^x (Ax^{\lambda+\mu} t^{\lambda-\mu-1} + Bx^\mu t^{2\lambda-\mu-1})y(t) dt = f(x).$$

The substitution $y(x) = x^\mu w(x)$ leads to an equation of the form 2.1.53:

$$w(x) - \int_a^x (Ax^\lambda t^{\lambda-1} + Bt^{2\lambda-1})w(t) dt = x^{-\mu} f(x).$$

$$55. \quad y(x) + A \int_a^x [\lambda x^{\lambda-1} t^\mu - (\lambda + \mu)x^{\lambda+\mu-1}] y(t) dt = f(x).$$

This equation can be obtained by differentiating equation 1.1.52:

$$\int_a^x [1 + A(x^\lambda t^\mu - x^{\lambda+\mu})] y(t) dt = F(x), \quad F(x) = \int_a^x f(x) dx.$$

Solution:

$$y(x) = \frac{d}{dx} \left\{ \frac{x^\lambda}{\Phi(x)} \int_a^x [t^{-\lambda} F(t)]'_t \Phi(t) dt \right\}, \quad \Phi(x) = \exp \left(-\frac{A\mu}{\mu + \lambda} x^{\mu+\lambda} \right).$$

$$56. \quad y(x) + \int_a^x (ABx^{\lambda+1} - ABx^\lambda t - Ax^\lambda - B)y(t) dt = f(x).$$

This is a special case of equation 2.9.7.

Solution:

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt,$$

$$R(x, t) = (Ax^\lambda + B) \exp\left[\frac{A}{\lambda+1}(x^{\lambda+1} - t^{\lambda+1})\right] + B^2 \int_t^x \exp\left[\frac{A}{\lambda+1}(s^{\lambda+1} - t^{\lambda+1}) + B(x-s)\right] ds.$$

$$57. \quad y(x) + \int_a^x (ABxt^\lambda - ABt^{\lambda+1} + At^\lambda + B)y(t) dt = f(x).$$

This is a special case of equation 2.9.8.

Solution:

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt,$$

$$R(x, t) = -(At^\lambda + B) \exp\left[\frac{A}{\lambda+1}(t^{\lambda+1} - x^{\lambda+1})\right] + B^2 \int_t^x \exp\left[\frac{A}{\lambda+1}(s^{\lambda+1} - x^{\lambda+1}) + B(t-s)\right] ds.$$

$$58. \quad y(x) - \lambda \int_a^x \left(\frac{x+b}{t+b}\right)^\mu y(t) dt = f(x).$$

This is a special case of equation 2.9.1 with $g(x) = (x+b)^\mu$.

Solution:

$$y(x) = f(x) + \lambda \int_a^x \left(\frac{x+b}{t+b}\right)^\mu e^{\lambda(x-t)} f(t) dt.$$

$$59. \quad y(x) - \lambda \int_a^x \frac{x^\mu + b}{t^\mu + b} y(t) dt = f(x).$$

This is a special case of equation 2.9.1 with $g(x) = x^\mu + b$.

Solution:

$$y(x) = f(x) + \lambda \int_a^x \frac{x^\mu + b}{t^\mu + b} e^{\lambda(x-t)} f(t) dt.$$

$$60. \quad y(x) - \lambda \int_0^x \frac{y(t) dt}{(x-t)^\alpha} = f(x), \quad 0 < \alpha < 1.$$

Generalized Abel equation of the second kind.

1°. Assume that the number α can be represented in the form

$$\alpha = 1 - \frac{m}{n}, \quad \text{where } m = 1, 2, \dots, \quad n = 2, 3, \dots \quad (m < n).$$

In this case, the solution of the generalized Abel equation of the second kind can be written in closed form (in quadratures):

$$y(x) = f(x) + \int_0^x R(x-t)f(t) dt,$$

where

$$R(x) = \sum_{\nu=1}^{n-1} \frac{\lambda^\nu \Gamma^\nu(m/n)}{\Gamma(\nu m/n)} x^{(\nu m/n)-1} + \frac{b}{m} \sum_{\mu=0}^{m-1} \varepsilon_\mu \exp(\varepsilon_\mu b x) + \frac{b}{m} \sum_{\nu=1}^{n-1} \frac{\lambda^\nu \Gamma^\nu(m/n)}{\Gamma(\nu m/n)} \left[\sum_{\mu=0}^{m-1} \varepsilon_\mu \exp(\varepsilon_\mu b x) \int_0^x t^{(\nu m/n)-1} \exp(-\varepsilon_\mu b t) dt \right],$$

$$b = \lambda^{n/m} \Gamma^{n/m}(m/n), \quad \varepsilon_\mu = \exp\left(\frac{2\pi\mu i}{m}\right), \quad i^2 = -1, \quad \mu = 0, 1, \dots, m-1.$$

2°. Solution with any α from $0 < \alpha < 1$:

$$y(x) = f(x) + \int_0^x R(x-t)f(t) dt, \quad \text{where} \quad R(x) = \sum_{n=1}^{\infty} \frac{[\lambda \Gamma(1-\alpha)x^{1-\alpha}]^n}{x \Gamma[n(1-\alpha)]}.$$

⊙ References: H. Brakhage, K. Nickel, and P. Rieder (1965), V. I. Smirnov (1974).

61. $y(x) - \frac{\lambda}{x^\alpha} \int_0^x \frac{y(t) dt}{(x-t)^{1-\alpha}} = f(x), \quad 0 < \alpha \leq 1.$

1°. The solution of the homogeneous equation ($f \equiv 0$) is

$$y(x) = Cx^\beta \quad (\beta > -1, \lambda > 0). \quad (1)$$

Here C is an arbitrary constant, and $\beta = \beta(\lambda)$ is determined by the transcendental equation

$$\lambda B(\alpha, \beta + 1) = 1, \quad (2)$$

where $B(p, q) = \int_0^1 z^{p-1}(1-z)^{q-1} dz$ is the beta function.

2°. For a polynomial right-hand side,

$$f(x) = \sum_{n=0}^N A_n x^n$$

the solution bounded at zero is given by

$$y(x) = \begin{cases} \sum_{n=0}^N \frac{A_n}{1 - (\lambda/\lambda_n)} x^n & \text{for } \lambda < \alpha, \\ \sum_{n=0}^N \frac{A_n}{1 - (\lambda/\lambda_n)} x^n + Cx^\beta & \text{for } \lambda > \alpha \text{ and } \lambda \neq \lambda_n, \end{cases}$$

$$\lambda_n = \frac{(\alpha)_{n+1}}{n!}, \quad (\alpha)_{n+1} = \alpha(\alpha+1)\dots(\alpha+n).$$

Here C is an arbitrary constant, and $\beta = \beta(\lambda)$ is determined by the transcendental equation (2).

For special $\lambda = \lambda_n$ ($n = 1, 2, \dots$), the solution differs in one term and has the form

$$y(x) = \sum_{m=0}^{n-1} \frac{A_m}{1 - (\lambda_n/\lambda_m)} x^m + \sum_{m=n+1}^N \frac{A_m}{1 - (\lambda_n/\lambda_m)} x^m - A_n \frac{\bar{\lambda}_n}{\lambda_n} x^n \ln x + Cx^n,$$

where $\bar{\lambda}_n = \left[\int_0^1 (1-z)^{\alpha-1} z^n \ln z dz \right]^{-1}.$

3°. For arbitrary $f(x)$, expandable into power series, the formulas of item 2° can be used, in which one should set $N = \infty$. In this case, the radius of convergence of the solution $y(x)$ is equal to the radius of convergence of $f(x)$.

4°. For

$$f(x) = \ln(kx) \sum_{n=0}^N A_n x^n,$$

a solution has the form

$$y(x) = \ln(kx) \sum_{n=0}^N B_n x^n + \sum_{n=0}^N D_n x^n,$$

where the constants B_n and D_n are found by the method of undetermined coefficients. To obtain the general solution we must add the solution (1) of the homogeneous equation.

In Mikhailov (1966), solvability conditions for the integral equation in question were investigated for various classes of $f(x)$.

62.
$$y(x) - \frac{\lambda}{x^\mu} \int_0^x \frac{y(t) dt}{(ax + bt)^{1-\mu}} = f(x).$$

Here $a > 0$, $b > 0$, and μ is an arbitrary number.

1°. The solution of the homogeneous equation ($f \equiv 0$) is

$$y(x) = Cx^\beta \quad (\beta > -1, \lambda > 0). \quad (1)$$

Here C is an arbitrary constant, and $\beta = \beta(\lambda)$ is determined by the transcendental equation

$$\lambda I(\beta) = 1, \quad \text{where} \quad I(\beta) = \int_0^1 z^\beta (a + bz)^{\mu-1} dz. \quad (2)$$

2°. For a polynomial right-hand side,

$$f(x) = \sum_{n=0}^N A_n x^n$$

the solution bounded at zero is given by

$$y(x) = \begin{cases} \sum_{n=0}^N \frac{A_n}{1 - (\lambda/\lambda_n)} x^n & \text{for } \lambda < \lambda_0, \\ \sum_{n=0}^N \frac{A_n}{1 - (\lambda/\lambda_n)} x^n + Cx^\beta & \text{for } \lambda > \lambda_0 \text{ and } \lambda \neq \lambda_n, \end{cases}$$

$$\lambda_n = \frac{1}{I(n)}, \quad I(n) = \int_0^1 z^n (a + bz)^{\mu-1} dz.$$

Here C is an arbitrary constant, and $\beta = \beta(\lambda)$ is determined by the transcendental equation (2).

3°. For special $\lambda = \lambda_n$ ($n = 1, 2, \dots$), the solution differs in one term and has the form

$$y(x) = \sum_{m=0}^{n-1} \frac{A_m}{1 - (\lambda_n/\lambda_m)} x^m + \sum_{m=n+1}^N \frac{A_m}{1 - (\lambda_n/\lambda_m)} x^m - A_n \frac{\bar{\lambda}_n}{\lambda_n} x^n \ln x + Cx^n,$$

where $\bar{\lambda}_n = \left[\int_0^1 z^n (a + bz)^{\mu-1} \ln z dz \right]^{-1}$.

4°. For arbitrary $f(x)$ expandable into power series, the formulas of item 2° can be used, in which one should set $N = \infty$. In this case, the radius of convergence of the solution $y(x)$ is equal to the radius of convergence of $f(x)$.

2.2. Equations Whose Kernels Contain Exponential Functions

2.2-1. Kernels Containing Exponential Functions.

$$1. \quad y(x) + A \int_a^x e^{\lambda(x-t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) - A \int_a^x e^{(\lambda-A)(x-t)} f(t) dt.$$

$$2. \quad y(x) + A \int_a^x e^{\lambda x + \beta t} y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ae^{\lambda x}$ and $h(t) = e^{\beta t}$. For $\beta = -\lambda$, see equation 2.2.1.

Solution:

$$y(x) = f(x) - \int_a^x R(x, t) f(t) dt, \quad R(x, t) = Ae^{\lambda x + \beta t} \exp\left\{\frac{A}{\lambda + \beta} [e^{(\lambda + \beta)t} - e^{(\lambda + \beta)x}]\right\}.$$

$$3. \quad y(x) + A \int_a^x [e^{\lambda(x-t)} - 1] y(t) dt = f(x).$$

1°. Solution with $D \equiv \lambda(\lambda - 4A) > 0$:

$$y(x) = f(x) - \frac{2A\lambda}{\sqrt{D}} \int_a^x R(x-t) f(t) dt, \quad R(x) = \exp\left(\frac{1}{2}\lambda x\right) \sinh\left(\frac{1}{2}\sqrt{D}x\right).$$

2°. Solution with $D \equiv \lambda(\lambda - 4A) < 0$:

$$y(x) = f(x) - \frac{2A\lambda}{\sqrt{|D|}} \int_a^x R(x-t) f(t) dt, \quad R(x) = \exp\left(\frac{1}{2}\lambda x\right) \sin\left(\frac{1}{2}\sqrt{|D|}x\right).$$

3°. Solution with $\lambda = 4A$:

$$y(x) = f(x) - 4A^2 \int_a^x (x-t) \exp[2A(x-t)] f(t) dt.$$

$$4. \quad y(x) + \int_a^x [Ae^{\lambda(x-t)} + B] y(t) dt = f(x).$$

This is a special case of equation 2.2.10 with $A_1 = A$, $A_2 = B$, $\lambda_1 = \lambda$, and $\lambda_2 = 0$.

1°. The structure of the solution depends on the sign of the discriminant

$$D \equiv (A - B - \lambda)^2 + 4AB \tag{1}$$

of the square equation

$$\mu^2 + (A + B - \lambda)\mu - B\lambda = 0. \tag{2}$$

2°. If $D > 0$, then equation (2) has the real different roots

$$\mu_1 = \frac{1}{2}(\lambda - A - B) + \frac{1}{2}\sqrt{D}, \quad \mu_2 = \frac{1}{2}(\lambda - A - B) - \frac{1}{2}\sqrt{D}.$$

In this case, the original integral equation has the solution

$$y(x) = f(x) + \int_a^x [E_1 e^{\mu_1(x-t)} + E_2 e^{\mu_2(x-t)}] f(t) dt,$$

where

$$E_1 = A \frac{\mu_1}{\mu_2 - \mu_1} + B \frac{\mu_1 - \lambda}{\mu_2 - \mu_1}, \quad E_2 = A \frac{\mu_2}{\mu_1 - \mu_2} + B \frac{\mu_2 - \lambda}{\mu_1 - \mu_2}.$$

3°. If $D < 0$, then equation (2) has the complex conjugate roots

$$\mu_1 = \sigma + i\beta, \quad \mu_2 = \sigma - i\beta, \quad \sigma = \frac{1}{2}(\lambda - A - B), \quad \beta = \frac{1}{2}\sqrt{-D}.$$

In this case, the original integral equation has the solution

$$y(x) = f(x) + \int_a^x \left\{ E_1 e^{\sigma(x-t)} \cos[\beta(x-t)] + E_2 e^{\sigma(x-t)} \sin[\beta(x-t)] \right\} f(t) dt,$$

where

$$E_1 = -A - B, \quad E_2 = \frac{1}{\beta}(-A\sigma - B\sigma + B\lambda).$$

5. $y(x) + A \int_a^x (e^{\lambda x} - e^{\lambda t}) y(t) dt = f(x).$

This is a special case of equation 2.9.5 with $g(x) = Ae^{\lambda x}$.

Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x [u'_1(x)u'_2(t) - u'_2(x)u'_1(t)] f(t) dt,$$

where the primes denote differentiation with respect to the argument specified in the parentheses, and $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $u''_{xx} + A\lambda e^{\lambda x}u = 0$; the functions $u_1(x)$ and $u_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of A :

For $A\lambda > 0$,

$$W = \frac{\lambda}{\pi}, \quad u_1(x) = J_0\left(\frac{2\sqrt{A\lambda}}{\lambda}e^{\lambda x/2}\right), \quad u_2(x) = Y_0\left(\frac{2\sqrt{A\lambda}}{\lambda}e^{\lambda x/2}\right),$$

For $A\lambda < 0$,

$$W = -\frac{\lambda}{2}, \quad u_1(x) = I_0\left(\frac{2\sqrt{|A\lambda|}}{\lambda}e^{\lambda x/2}\right), \quad u_2(x) = K_0\left(\frac{2\sqrt{|A\lambda|}}{\lambda}e^{\lambda x/2}\right).$$

6. $y(x) + \int_a^x (Ae^{\lambda x} + Be^{\lambda t}) y(t) dt = f(x).$

This is a special case of equation 2.9.6 with $g(x) = Ae^{\lambda x}$ and $h(t) = Be^{\lambda t}$. For $B = -A$, see equation 2.2.5.

Differentiating the original integral equation followed by substituting $Y(x) = \int_a^x y(t) dt$ yields the second-order linear ordinary differential equation

$$Y''_{xx} + (A + B)e^{\lambda x}Y'_x + A\lambda e^{\lambda x}Y = f'_x(x) \quad (1)$$

under the initial conditions

$$Y(a) = 0, \quad Y'_x(a) = f(a). \quad (2)$$

A fundamental system of solutions of the homogeneous equation (1) with $f \equiv 0$ has the form

$$Y_1(x) = \Phi\left(\frac{A}{m}, 1; -\frac{m}{\lambda}e^{\lambda x}\right), \quad Y_2(x) = \Psi\left(\frac{A}{m}, 1; -\frac{m}{\lambda}e^{\lambda x}\right), \quad m = A + B,$$

where $\Phi(\alpha, \beta; x)$ and $\Psi(\alpha, \beta; x)$ are degenerate hypergeometric functions.

Solving the homogeneous equation (1) under conditions (2) for an arbitrary function $f = f(x)$ and taking into account the relation $y(x) = Y'_x(x)$, we thus obtain the solution of the integral equation in the form

$$y(x) = f(x) - \int_a^x R(x, t)f(t) dt,$$

$$R(x, t) = \frac{\Gamma(A/m)}{\lambda} \frac{\partial^2}{\partial x \partial t} \left\{ \exp\left(\frac{m}{\lambda}e^{\lambda t}\right) [Y_1(x)Y_2(t) - Y_2(x)Y_1(t)] \right\}.$$

7. $y(x) + A \int_a^x [e^{\lambda(x+t)} - e^{2\lambda t}] y(t) dt = f(x).$

The transformation $z = e^{\lambda x}$, $\tau = e^{\lambda t}$ leads to an equation of the form 2.1.4.

1°. Solution with $A\lambda > 0$:

$$y(x) = f(x) - \lambda k \int_a^x e^{\lambda t} \sin[k(e^{\lambda x} - e^{\lambda t})] f(t) dt, \quad k = \sqrt{A/\lambda}.$$

2°. Solution with $A\lambda < 0$:

$$y(x) = f(x) + \lambda k \int_a^x e^{\lambda t} \sinh[k(e^{\lambda x} - e^{\lambda t})] f(t) dt, \quad k = \sqrt{|A/\lambda|}.$$

8. $y(x) + A \int_a^x [e^{\lambda x + \mu t} - e^{(\lambda + \mu)t}] y(t) dt = f(x).$

The transformation $z = e^{\mu x}$, $\tau = e^{\mu t}$, $Y(z) = y(x)$ leads to an equation of the form 2.1.52:

$$Y(z) + \frac{A}{\mu} \int_b^z (z^k - \tau^k) Y(\tau) d\tau = F(z), \quad F(z) = f(x),$$

where $k = \lambda/\mu$, $b = e^{\mu a}$.

9. $y(x) + A \int_a^x [\lambda e^{\lambda x + \mu t} - (\lambda + \mu)e^{(\lambda + \mu)x}] y(t) dt = f(x).$

This equation can be obtained by differentiating an equation of the form 1.2.22:

$$\int_a^x [1 + Ae^{\lambda x}(e^{\mu t} - e^{\mu x})] y(t) dt = F(x), \quad F(x) = \int_a^x f(t) dt.$$

Solution:

$$y(x) = \frac{d}{dx} \left\{ e^{\lambda x} \Phi(x) \int_a^x \left[\frac{F(t)}{e^{\lambda t}} \right]'_t \frac{dt}{\Phi(t)} \right\}, \quad \Phi(x) = \exp \left[\frac{A\mu}{\lambda + \mu} e^{(\lambda + \mu)x} \right].$$

$$10. \quad y(x) + \int_a^x [A_1 e^{\lambda_1(x-t)} + A_2 e^{\lambda_2(x-t)}] y(t) dt = f(x).$$

1°. Introduce the notation

$$I_1 = \int_a^x e^{\lambda_1(x-t)} y(t) dt, \quad I_2 = \int_a^x e^{\lambda_2(x-t)} y(t) dt.$$

Differentiating the integral equation twice yields (the first line is the original equation)

$$y + A_1 I_1 + A_2 I_2 = f, \quad f = f(x), \quad (1)$$

$$y'_x + (A_1 + A_2)y + A_1 \lambda_1 I_1 + A_2 \lambda_2 I_2 = f'_x, \quad (2)$$

$$y''_{xx} + (A_1 + A_2)y'_x + (A_1 \lambda_1 + A_2 \lambda_2)y + A_1 \lambda_1^2 I_1 + A_2 \lambda_2^2 I_2 = f''_{xx}. \quad (3)$$

Eliminating I_1 and I_2 , we arrive at the second-order linear ordinary differential equation with constant coefficients

$$y''_{xx} + (A_1 + A_2 - \lambda_1 - \lambda_2)y'_x + (\lambda_1 \lambda_2 - A_1 \lambda_2 - A_2 \lambda_1)y = f''_{xx} - (\lambda_1 + \lambda_2)f'_x + \lambda_1 \lambda_2 f. \quad (4)$$

Substituting $x = a$ into (1) and (2) yields the initial conditions

$$y(a) = f(a), \quad y'_x(a) = f'_x(a) - (A_1 + A_2)f(a). \quad (5)$$

Solving the differential equation (4) under conditions (5), we can find the solution of the integral equation.

2°. Consider the characteristic equation

$$\mu^2 + (A_1 + A_2 - \lambda_1 - \lambda_2)\mu + \lambda_1 \lambda_2 - A_1 \lambda_2 - A_2 \lambda_1 = 0 \quad (6)$$

which corresponds to the homogeneous differential equation (4) (with $f(x) \equiv 0$). The structure of the solution of the integral equation depends on the sign of the discriminant

$$D \equiv (A_1 - A_2 - \lambda_1 + \lambda_2)^2 + 4A_1 A_2$$

of the quadratic equation (6).

If $D > 0$, the quadratic equation (6) has the real different roots

$$\mu_1 = \frac{1}{2}(\lambda_1 + \lambda_2 - A_1 - A_2) + \frac{1}{2}\sqrt{D}, \quad \mu_2 = \frac{1}{2}(\lambda_1 + \lambda_2 - A_1 - A_2) - \frac{1}{2}\sqrt{D}.$$

In this case, the solution of the original integral equation has the form

$$y(x) = f(x) + \int_a^x [B_1 e^{\mu_1(x-t)} + B_2 e^{\mu_2(x-t)}] f(t) dt,$$

where

$$B_1 = A_1 \frac{\mu_1 - \lambda_2}{\mu_2 - \mu_1} + A_2 \frac{\mu_1 - \lambda_1}{\mu_2 - \mu_1}, \quad B_2 = A_1 \frac{\mu_2 - \lambda_2}{\mu_1 - \mu_2} + A_2 \frac{\mu_2 - \lambda_1}{\mu_1 - \mu_2}.$$

If $D < 0$, the quadratic equation (6) has the complex conjugate roots

$$\mu_1 = \sigma + i\beta, \quad \mu_2 = \sigma - i\beta, \quad \sigma = \frac{1}{2}(\lambda_1 + \lambda_2 - A_1 - A_2), \quad \beta = \frac{1}{2}\sqrt{-D}.$$

In this case, the solution of the original integral equation has the form

$$y(x) = f(x) + \int_a^x \{B_1 e^{\sigma(x-t)} \cos[\beta(x-t)] + B_2 e^{\sigma(x-t)} \sin[\beta(x-t)]\} f(t) dt,$$

where

$$B_1 = -A_1 - A_2, \quad B_2 = \frac{1}{\beta} [A_1(\lambda_2 - \sigma) + A_2(\lambda_1 - \sigma)].$$

$$11. \quad y(x) + \int_a^x [Ae^{\lambda(x+t)} - Ae^{2\lambda t} + Be^{\lambda t}] y(t) dt = f(x).$$

The transformation $z = e^{\lambda x}$, $\tau = e^{\lambda t}$, $Y(z) = y(x)$ leads to an equation of the form 2.1.5:

$$Y(z) + \int_b^z [B_1(z - \tau) + A_1] Y(\tau) d\tau = F(z), \quad F(z) = f(x),$$

where $A_1 = B/\lambda$, $B_1 = A/\lambda$, $b = e^{\lambda a}$.

$$12. \quad y(x) + \int_a^x [Ae^{\lambda(x+t)} + Be^{2\lambda t} + Ce^{\lambda t}] y(t) dt = f(x).$$

The transformation $z = e^{\lambda x}$, $\tau = e^{\lambda t}$, $Y(z) = y(x)$ leads to an equation of the form 2.1.6:

$$Y(z) - \int_b^z (A_1 z + B_1 \tau + C_1) Y(\tau) d\tau = F(z), \quad F(z) = f(x),$$

where $A_1 = -A/\lambda$, $B_1 = -B/\lambda$, $C_1 = -C/\lambda$, $b = e^{\lambda a}$.

$$13. \quad y(x) + \int_a^x [\lambda e^{\lambda(x-t)} + A(\mu e^{\mu x + \lambda t} - \lambda e^{\lambda x + \mu t})] y(t) dt = f(x).$$

This is a special case of equation 2.9.23 with $h(t) = A$.

Solution:

$$y(x) = \frac{1}{e^{\lambda x}} \frac{d}{dx} \left\{ \Phi(x) \int_a^x \left[\frac{F(t)}{e^{\lambda t}} \right]'_t \frac{e^{2\lambda t}}{\Phi(t)} dt \right\},$$

$$\Phi(x) = \exp \left[A \frac{\lambda - \mu}{\lambda + \mu} e^{(\lambda + \mu)x} \right], \quad F(x) = \int_a^x f(t) dt.$$

$$14. \quad y(x) - \int_a^x [\lambda e^{-\lambda(x-t)} + A(\mu e^{\lambda x + \mu t} - \lambda e^{\mu x + \lambda t})] y(t) dt = f(x).$$

This is a special case of equation 2.9.24 with $h(x) = A$.

Assume that $f(a) = 0$. Solution:

$$y(x) = \int_a^x w(t) dt, \quad w(x) = e^{-\lambda x} \frac{d}{dx} \left\{ \frac{e^{2\lambda x}}{\Phi(x)} \int_a^x \left[\frac{f(t)}{e^{\lambda t}} \right]'_t \Phi(t) dt \right\},$$

$$\Phi(x) = \exp \left[A \frac{\lambda - \mu}{\lambda + \mu} e^{(\lambda + \mu)x} \right].$$

$$15. \quad y(x) + \int_a^x [\lambda e^{\lambda(x-t)} + Ae^{\beta t} (\mu e^{\mu x + \lambda t} - \lambda e^{\lambda x + \mu t})] y(t) dt = f(x).$$

This is a special case of equation 2.9.23 with $h(t) = Ae^{\beta t}$.

Solution:

$$y(x) = e^{-(\lambda + \beta)x} \frac{d}{dx} \left\{ \Phi(x) \int_a^x \left[\frac{F(t)}{e^{\lambda t}} \right]'_t \frac{e^{(2\lambda + \beta)t}}{\Phi(t)} dt \right\},$$

$$\Phi(x) = \exp \left[A \frac{\lambda - \mu}{\lambda + \mu + \beta} e^{(\lambda + \mu + \beta)x} \right], \quad F(x) = \int_a^x f(t) dt.$$

$$16. \quad y(x) - \int_a^x [\lambda e^{-\lambda(x-t)} + A e^{\beta x} (\mu e^{\lambda x + \mu t} - \lambda e^{\mu x + \lambda t})] y(t) dt = f(x).$$

This is a special case of equation 2.9.24 with $h(x) = A e^{\beta x}$.

Assume that $f(a) = 0$. Solution:

$$y(x) = \int_a^x w(t) dt, \quad w(x) = e^{-\lambda x} \frac{d}{dx} \left\{ \frac{e^{(2\lambda+\beta)x}}{\Phi(x)} \int_a^x \left[\frac{f(t)}{e^{(\lambda+\beta)t}} \right]' \Phi(t) dt \right\},$$

$$\Phi(x) = \exp \left[A \frac{\lambda - \mu}{\lambda + \mu + \beta} e^{(\lambda+\mu+\beta)x} \right].$$

$$17. \quad y(x) + \int_a^x [A B e^{(\lambda+1)x+t} - A B e^{\lambda x+2t} - A e^{\lambda x+t} - B e^t] y(t) dt = f(x).$$

The transformation $z = e^x$, $\tau = e^t$, $Y(z) = y(x)$ leads to an equation of the form 2.1.56:

$$Y(z) + \int_b^z (A B z^{\lambda+1} - A B z^{\lambda} \tau - A z^{\lambda} - B) Y(\tau) d\tau = F(z),$$

where $F(z) = f(x)$ and $b = e^a$.

$$18. \quad y(x) + \int_a^x [A B e^{x+\lambda t} - A B e^{(\lambda+1)t} + A e^{\lambda t} + B e^t] y(t) dt = f(x).$$

The transformation $z = e^x$, $\tau = e^t$, $Y(z) = y(x)$ leads to an equation of the form 2.1.57 (in which λ is substituted by $\lambda - 1$):

$$Y(z) + \int_b^z (A B z \tau^{\lambda-1} - A B \tau^{\lambda} + A \tau^{\lambda-1} + B) Y(\tau) d\tau = F(z),$$

where $F(z) = f(x)$ and $b = e^a$.

$$19. \quad y(x) + \int_a^x \left[\sum_{k=1}^n A_k e^{\lambda_k(x-t)} \right] y(t) dt = f(x).$$

1°. This integral equation can be reduced to an n th-order linear nonhomogeneous ordinary differential equation with constant coefficients. Set

$$I_k(x) = \int_a^x e^{\lambda_k(x-t)} y(t) dt. \quad (1)$$

Differentiating (1) with respect to x yields

$$I_k' = y(x) + \lambda_k \int_a^x e^{\lambda_k(x-t)} y(t) dt, \quad (2)$$

where the prime stands for differentiation with respect to x . From the comparison of (1) with (2) we see that

$$I_k' = y(x) + \lambda_k I_k, \quad I_k = I_k(x). \quad (3)$$

The integral equation can be written in terms of $I_k(x)$ as follows:

$$y(x) + \sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x and taking account of (3), we obtain

$$y'_x(x) + \sigma_n y(x) + \sum_{k=1}^n A_k \lambda_k I_k = f'_x(x), \quad \sigma_n = \sum_{k=1}^n A_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5), we find that

$$y'_x(x) + (\sigma_n - \lambda_n)y(x) + \sum_{k=1}^{n-1} A_k(\lambda_k - \lambda_n)I_k = f'_x(x) - \lambda_n f(x). \quad (6)$$

Differentiating (6) with respect to x and eliminating I_{n-1} from the resulting equation with the aid of (6), we obtain a similar equation whose left-hand side is a second-order linear differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} A_k^1 I_k$. If we proceed with successively eliminating $I_{n-2}, I_{n-3}, \dots, I_1$ with the aid of differentiation and formula (3), then we will finally arrive at an n th-order linear nonhomogeneous ordinary differential equation with constant coefficients.

The initial conditions for $y(x)$ can be obtained by setting $x = a$ in the integral equation and all its derivative equations.

2°. The solution of the equation can be represented in the form

$$y(x) = f(x) + \int_a^x \left[\sum_{k=1}^n B_k e^{\mu_k(x-t)} \right] f(t) dt. \quad (7)$$

The unknown constants μ_k are the roots of the algebraic equation

$$\sum_{k=1}^n \frac{A_k}{z - \lambda_k} + 1 = 0, \quad (8)$$

which is reduced (by separating the numerator) to the problem of finding the roots of an n th-order characteristic polynomial.

After the μ_k have been calculated, the coefficients B_k can be found from the following linear system of algebraic equations:

$$\sum_{k=1}^n \frac{B_k}{\lambda_m - \mu_k} + 1 = 0, \quad m = 1, \dots, n. \quad (9)$$

Another way of determining the B_k is presented in item 3° below.

If all the roots μ_k of equation (8) are real and different, then the solution of the original integral equation can be calculated by formula (7).

To a pair of complex conjugate roots $\mu_{k,k+1} = \alpha \pm i\beta$ of the characteristic polynomial (8) there corresponds a pair of complex conjugate coefficients $B_{k,k+1}$ in equation (9). In this case, the corresponding terms $B_k e^{\mu_k(x-t)} + B_{k+1} e^{\mu_{k+1}(x-t)}$ in solution (7) can be written in the form $\overline{B}_k e^{\alpha(x-t)} [\cos \beta(x-t)] + \overline{B}_{k+1} e^{\alpha(x-t)} [\sin \beta(x-t)]$, where \overline{B}_k and \overline{B}_{k+1} are real coefficients.

3°. For $a = 0$, the solution of the original integral equation is given by

$$y(x) = f(x) - \int_0^x R(x-t)f(t) dt, \quad R(x) = \mathfrak{L}^{-1}[\overline{R}(p)], \quad (10)$$

where $\mathfrak{L}^{-1}[\overline{R}(p)]$ is the inverse Laplace transform of the function

$$\overline{R}(p) = \frac{\overline{K}(p)}{1 + \overline{K}(p)}, \quad \overline{K}(p) = \sum_{k=1}^n \frac{A_k}{p - \lambda_k}. \quad (11)$$

The transform $\overline{R}(p)$ of the resolvent $R(x)$ can be represented as a regular fractional function:

$$\overline{R}(p) = \frac{Q(p)}{P(p)}, \quad P(p) = (p - \mu_1)(p - \mu_2) \dots (p - \mu_n),$$

where $Q(p)$ is a polynomial in p of degree $< n$. The roots μ_k of the polynomial $P(p)$ coincide with the roots of equation (8). If all μ_k are real and different, then the resolvent can be determined by the formula

$$R(x) = \sum_{k=1}^n B_k e^{\mu_k x}, \quad B_k = \frac{Q(\mu_k)}{P'(\mu_k)},$$

where the prime stands for differentiation.

2.2-2. Kernels Containing Power-Law and Exponential Functions.

20. $y(x) + A \int_a^x x e^{\lambda(x-t)} y(t) dt = f(x).$

Solution:

$$y(x) = f(x) - A \int_a^x x \exp\left[\frac{1}{2}A(t^2 - x^2) + \lambda(x-t)\right] f(t) dt.$$

21. $y(x) + A \int_a^x t e^{\lambda(x-t)} y(t) dt = f(x).$

Solution:

$$y(x) = f(x) - A \int_a^x t \exp\left[\frac{1}{2}A(t^2 - x^2) + \lambda(x-t)\right] f(t) dt.$$

22. $y(x) + A \int_a^x (x-t) e^{\lambda t} y(t) dt = f(x).$

This is a special case of equation 2.9.4 with $g(t) = A e^{\lambda t}$.

Solution:

$$y(x) = f(x) + \frac{A}{W} \int_a^x [u_1(x)u_2(t) - u_2(x)u_1(t)] e^{\lambda t} f(t) dt,$$

where $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $u''_{xx} + A e^{\lambda x} u = 0$; the functions $u_1(x)$ and $u_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions, depending on sign A :

$$\begin{aligned} W = \frac{\lambda}{\pi}, \quad u_1(x) = J_0\left(\frac{2\sqrt{A}}{\lambda} e^{\lambda x/2}\right), \quad u_2(x) = Y_0\left(\frac{2\sqrt{A}}{\lambda} e^{\lambda x/2}\right) & \text{ for } A > 0, \\ W = -\frac{\lambda}{2}, \quad u_1(x) = I_0\left(\frac{2\sqrt{|A|}}{\lambda} e^{\lambda x/2}\right), \quad u_2(x) = K_0\left(\frac{2\sqrt{|A|}}{\lambda} e^{\lambda x/2}\right) & \text{ for } A < 0. \end{aligned}$$

$$23. \quad y(x) + A \int_a^x (x-t)e^{\lambda(x-t)} y(t) dt = f(x).$$

1°. Solution with $A > 0$:

$$y(x) = f(x) - k \int_a^x e^{\lambda(x-t)} \sin[k(x-t)] f(t) dt, \quad k = \sqrt{A}.$$

2°. Solution with $A < 0$:

$$y(x) = f(x) + k \int_a^x e^{\lambda(x-t)} \sinh[k(x-t)] f(t) dt, \quad k = \sqrt{-A}.$$

$$24. \quad y(x) + A \int_a^x (x-t)e^{\lambda x + \mu t} y(t) dt = f(x).$$

The substitution $u(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 2.2.22:

$$u(x) + A \int_a^x (x-t)e^{(\lambda+\mu)t} u(t) dt = f(x)e^{-\lambda x}.$$

$$25. \quad y(x) - \int_a^x (Ax + Bt + C)e^{\lambda(x-t)} y(t) dt = f(x).$$

The substitution $u(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 2.1.6:

$$u(x) - \int_a^x (Ax + Bt + C)u(t) dt = f(x)e^{-\lambda x}.$$

$$26. \quad y(x) + A \int_a^x x^2 e^{\lambda(x-t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) - A \int_a^x x^2 \exp\left[\frac{1}{3}A(t^3 - x^3) + \lambda(x-t)\right] f(t) dt.$$

$$27. \quad y(x) + A \int_a^x xte^{\lambda(x-t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) - A \int_a^x xt \exp\left[\frac{1}{3}A(t^3 - x^3) + \lambda(x-t)\right] f(t) dt.$$

$$28. \quad y(x) + A \int_a^x t^2 e^{\lambda(x-t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) - A \int_a^x t^2 \exp\left[\frac{1}{3}A(t^3 - x^3) + \lambda(x-t)\right] f(t) dt.$$

$$29. \quad y(x) + A \int_a^x (x-t)^2 e^{\lambda(x-t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) - \int_a^x R(x-t)f(t) dt,$$

$$R(x) = \frac{2}{3}ke^{(\lambda-2k)x} - \frac{2}{3}ke^{(\lambda+k)x} [\cos(\sqrt{3}kx) - \sqrt{3}\sin(\sqrt{3}kx)], \quad k = \left(\frac{1}{4}A\right)^{1/3}.$$

$$30. \quad y(x) + A \int_0^x (x^2 - t^2) e^{\lambda(x-t)} y(t) dt = f(x).$$

The substitution $u(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 2.1.11:

$$u(x) + A \int_0^x (x^2 - t^2) u(t) dt = f(x) e^{-\lambda x}.$$

$$31. \quad y(x) + A \int_a^x (x-t)^n e^{\lambda(x-t)} y(t) dt = f(x), \quad n = 1, 2, \dots$$

Solution:

$$y(x) = f(x) + \int_a^x R(x-t) f(t) dt,$$

$$R(x) = \frac{1}{n+1} e^{\lambda x} \sum_{k=0}^n \exp(\sigma_k x) [\sigma_k \cos(\beta_k x) - \beta_k \sin(\beta_k x)],$$

where

$$\sigma_k = |An!|^{\frac{1}{n+1}} \cos\left(\frac{2\pi k}{n+1}\right), \quad \beta_k = |An!|^{\frac{1}{n+1}} \sin\left(\frac{2\pi k}{n+1}\right) \quad \text{for } A < 0,$$

$$\sigma_k = |An!|^{\frac{1}{n+1}} \cos\left(\frac{2\pi k + \pi}{n+1}\right), \quad \beta_k = |An!|^{\frac{1}{n+1}} \sin\left(\frac{2\pi k + \pi}{n+1}\right) \quad \text{for } A > 0.$$

$$32. \quad y(x) + b \int_a^x \frac{\exp[\lambda(x-t)]}{\sqrt{x-t}} y(t) dt = f(x).$$

Solution:

$$y(x) = e^{\lambda x} \left\{ F(x) + \pi b^2 \int_a^x \exp[\pi b^2(x-t)] F(t) dt \right\},$$

where

$$F(x) = e^{-\lambda x} f(x) - b \int_a^x \frac{e^{-\lambda t} f(t)}{\sqrt{x-t}} dt.$$

$$33. \quad y(x) + A \int_a^x (x-t)t^k e^{\lambda(x-t)} y(t) dt = f(x).$$

The substitution $u(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 2.1.49:

$$u(x) + A \int_a^x (x-t)t^k u(t) dt = f(x) e^{-\lambda x}.$$

$$34. \quad y(x) + A \int_a^x (x^k - t^k) e^{\lambda(x-t)} y(t) dt = f(x).$$

The substitution $u(x) = e^{-\lambda x} y(x)$ leads to an equation of the form 2.1.52:

$$u(x) + A \int_a^x (x^k - t^k) u(t) dt = f(x) e^{-\lambda x}.$$

$$35. \quad y(x) - \lambda \int_0^x \frac{e^{\mu(x-t)}}{(x-t)^\alpha} y(t) dt = f(x), \quad 0 < \alpha < 1.$$

Solution:

$$y(x) = f(x) + \int_0^x R(x-t) f(t) dt, \quad \text{where } R(x) = e^{\mu x} \sum_{n=1}^{\infty} \frac{[\lambda \Gamma(1-\alpha) x^{1-\alpha}]^n}{x \Gamma[n(1-\alpha)]}.$$

$$36. \quad y(x) + A \int_a^x \exp[\lambda(x^2 - t^2)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) - A \int_a^x \exp[\lambda(x^2 - t^2) - A(x - t)] f(t) dt.$$

$$37. \quad y(x) + A \int_a^x \exp(\lambda x^2 + \beta t^2) y(t) dt = f(x).$$

In the case $\beta = -\lambda$, see equation 2.2.36. This is a special case of equation 2.9.2 with $g(x) = -A \exp(\lambda x^2)$ and $h(t) = \exp(\beta t^2)$.

$$38. \quad y(x) + A \int_x^\infty \exp(-\lambda\sqrt{t-x}) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = A \exp(-\lambda\sqrt{-x})$.

$$39. \quad y(x) + A \int_a^x \exp[\lambda(x^\mu - t^\mu)] y(t) dt = f(x), \quad \mu > 0.$$

This is a special case of equation 2.9.2 with $g(x) = -A \exp(\lambda x^\mu)$ and $h(t) = \exp(-\lambda t^\mu)$.

Solution:

$$y(x) = f(x) - A \int_a^x \exp[\lambda(x^\mu - t^\mu) - A(x - t)] f(t) dt.$$

$$40. \quad y(x) + k \int_0^x \frac{1}{x} \exp\left(-\lambda \frac{t}{x}\right) y(t) dt = g(x).$$

This is a special case of equation 2.9.71 with $f(z) = k e^{-\lambda z}$.

For a polynomial right-hand side, $g(x) = \sum_{n=0}^N A_n x^n$, a solution is given by

$$y(x) = \sum_{n=0}^N \frac{A_n}{1 + k B_n} x^n, \quad B_n = \frac{n!}{\lambda^{n+1}} - e^{-\lambda} \sum_{k=0}^n \frac{n!}{k!} \frac{1}{\lambda^{n-k+1}}.$$

2.3. Equations Whose Kernels Contain Hyperbolic Functions

2.3-1. Kernels Containing Hyperbolic Cosine.

$$1. \quad y(x) - A \int_a^x \cosh(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cosh(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \cosh(\lambda x) \exp\left\{\frac{A}{\lambda} [\sinh(\lambda x) - \sinh(\lambda t)]\right\} f(t) dt.$$

$$2. \quad y(x) - A \int_a^x \cosh(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \cosh(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \cosh(\lambda t) \exp\left\{\frac{A}{\lambda} [\sinh(\lambda x) - \sinh(\lambda t)]\right\} f(t) dt.$$

$$3. \quad y(x) + A \int_a^x \cosh[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.28 with $g(t) = A$. Therefore, solving the original integral equation is reduced to solving the second-order linear nonhomogeneous ordinary differential equation with constant coefficients

$$y''_{xx} + Ay'_x - \lambda^2 y = f''_{xx} - \lambda^2 f, \quad f = f(x),$$

under the initial conditions

$$y(a) = f(a), \quad y'_x(a) = f'_x(a) - Af(a).$$

Solution:

$$y(x) = f(x) + \int_a^x R(x-t) f(t) dt,$$

$$R(x) = \exp\left(-\frac{1}{2}Ax\right) \left[\frac{A^2}{2k} \sinh(kx) - A \cosh(kx) \right], \quad k = \sqrt{\lambda^2 + \frac{1}{4}A^2}.$$

$$4. \quad y(x) + \int_a^x \left\{ \sum_{k=1}^n A_k \cosh[\lambda_k(x-t)] \right\} y(t) dt = f(x).$$

This equation can be reduced to an equation of the form 2.2.19 by using the identity $\cosh z \equiv \frac{1}{2}(e^z + e^{-z})$. Therefore, the integral equation in question can be reduced to a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients.

$$5. \quad y(x) - A \int_a^x \frac{\cosh(\lambda x)}{\cosh(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\cosh(\lambda x)}{\cosh(\lambda t)} f(t) dt.$$

$$6. \quad y(x) - A \int_a^x \frac{\cosh(\lambda t)}{\cosh(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\cosh(\lambda t)}{\cosh(\lambda x)} f(t) dt.$$

$$7. \quad y(x) - A \int_a^x \cosh^k(\lambda x) \cosh^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cosh^k(\lambda x)$ and $h(t) = \cosh^m(\mu t)$.

$$8. \quad y(x) + A \int_a^x t \cosh[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.28 with $g(t) = At$.

$$9. \quad y(x) + A \int_a^x t^k \cosh^m(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -A \cosh^m(\lambda x)$ and $h(t) = t^k$.

$$10. \quad y(x) + A \int_a^x x^k \cosh^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ax^k$ and $h(t) = \cosh^m(\lambda t)$.

$$11. \quad y(x) - \int_a^x [A \cosh(kx) + B - AB(x-t) \cosh(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \cosh(kx)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t) f(t) dt,$$

$$R(x,t) = [A \cosh(kx) + B] \frac{G(x)}{G(t)} + \frac{B^2}{G(t)} \int_t^x e^{B(x-s)} G(s) ds, \quad G(x) = \exp \left[\frac{A}{k} \sinh(kx) \right].$$

$$12. \quad y(x) + \int_a^x [A \cosh(kt) + B + AB(x-t) \cosh(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \cosh(kt)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t) f(t) dt,$$

$$R(x,t) = -[A \cosh(kt) + B] \frac{G(t)}{G(x)} + \frac{B^2}{G(x)} \int_t^x e^{B(t-s)} G(s) ds, \quad G(x) = \exp \left[\frac{A}{k} \sinh(kx) \right].$$

$$13. \quad y(x) + A \int_x^\infty \cosh(\lambda \sqrt{t-x}) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = A \cosh(\lambda \sqrt{-x})$.

2.3-2. Kernels Containing Hyperbolic Sine.

$$14. \quad y(x) - A \int_a^x \sinh(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sinh(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \sinh(\lambda x) \exp \left\{ \frac{A}{\lambda} [\cosh(\lambda x) - \cosh(\lambda t)] \right\} f(t) dt.$$

$$15. \quad y(x) - A \int_a^x \sinh(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \sinh(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \sinh(\lambda t) \exp\left\{\frac{A}{\lambda} [\cosh(\lambda x) - \cosh(\lambda t)]\right\} f(t) dt.$$

$$16. \quad y(x) + A \int_a^x \sinh[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.30 with $g(x) = A$.

1°. Solution with $\lambda(A - \lambda) > 0$:

$$y(x) = f(x) - \frac{A\lambda}{k} \int_a^x \sin[k(x-t)] f(t) dt, \quad \text{where } k = \sqrt{\lambda(A - \lambda)}.$$

2°. Solution with $\lambda(A - \lambda) < 0$:

$$y(x) = f(x) - \frac{A\lambda}{k} \int_a^x \sinh[k(x-t)] f(t) dt, \quad \text{where } k = \sqrt{\lambda(\lambda - A)}.$$

3°. Solution with $A = \lambda$:

$$y(x) = f(x) - \lambda^2 \int_a^x (x-t) f(t) dt.$$

$$17. \quad y(x) + A \int_a^x \sinh^3[\lambda(x-t)] y(t) dt = f(x).$$

Using the formula $\sinh^3 \beta = \frac{1}{4} \sinh 3\beta - \frac{3}{4} \sinh \beta$, we arrive at an equation of the form 2.3.18:

$$y(x) + \int_a^x \left\{ \frac{1}{4} A \sinh[3\lambda(x-t)] - \frac{3}{4} A \sinh[\lambda(x-t)] \right\} y(t) dt = f(x).$$

$$18. \quad y(x) + \int_a^x \{ A_1 \sinh[\lambda_1(x-t)] + A_2 \sinh[\lambda_2(x-t)] \} y(t) dt = f(x).$$

1°. Introduce the notation

$$I_1 = \int_a^x \sinh[\lambda_1(x-t)] y(t) dt, \quad I_2 = \int_a^x \sinh[\lambda_2(x-t)] y(t) dt,$$

$$J_1 = \int_a^x \cosh[\lambda_1(x-t)] y(t) dt, \quad J_2 = \int_a^x \cosh[\lambda_2(x-t)] y(t) dt.$$

Successively differentiating the integral equation four times yields (the first line is the original equation)

$$y + A_1 I_1 + A_2 I_2 = f, \quad f = f(x), \quad (1)$$

$$y'_x + A_1 \lambda_1 J_1 + A_2 \lambda_2 J_2 = f'_x, \quad (2)$$

$$y''_{xx} + (A_1 \lambda_1 + A_2 \lambda_2) y + A_1 \lambda_1^2 I_1 + A_2 \lambda_2^2 I_2 = f''_{xx}, \quad (3)$$

$$y'''_{xxx} + (A_1 \lambda_1 + A_2 \lambda_2) y'_x + A_1 \lambda_1^3 J_1 + A_2 \lambda_2^3 J_2 = f'''_{xxx}, \quad (4)$$

$$y''''_{xxxx} + (A_1 \lambda_1 + A_2 \lambda_2) y''_{xx} + (A_1 \lambda_1^3 + A_2 \lambda_2^3) y + A_1 \lambda_1^4 I_1 + A_2 \lambda_2^4 I_2 = f''''_{xxxx}. \quad (5)$$

Eliminating I_1 and I_2 from (1), (3), and (5), we arrive at a fourth-order linear ordinary differential equation with constant coefficients:

$$\begin{aligned} y_{xxxx}''' - (\lambda_1^2 + \lambda_2^2 - A_1\lambda_1 - A_2\lambda_2)y_{xx}'' + (\lambda_1^2\lambda_2^2 - A_1\lambda_1\lambda_2^2 - A_2\lambda_1^2\lambda_2)y = \\ f_{xxxx}''' - (\lambda_1^2 + \lambda_2^2)f_{xx}'' + \lambda_1^2\lambda_2^2 f. \end{aligned} \quad (6)$$

The initial conditions can be obtained by setting $x = a$ in (1)–(4):

$$\begin{aligned} y(a) = f(a), \quad y_x'(a) = f_x'(a), \\ y_{xx}''(a) = f_{xx}''(a) - (A_1\lambda_1 + A_2\lambda_2)f(a), \\ y_{xxx}'''(a) = f_{xxx}'''(a) - (A_1\lambda_1 + A_2\lambda_2)f_x'(a). \end{aligned} \quad (7)$$

On solving the differential equation (6) under conditions (7), we thus find the solution of the integral equation.

2°. Consider the characteristic equation

$$z^2 - (\lambda_1^2 + \lambda_2^2 - A_1\lambda_1 - A_2\lambda_2)z + \lambda_1^2\lambda_2^2 - A_1\lambda_1\lambda_2^2 - A_2\lambda_1^2\lambda_2 = 0, \quad (8)$$

whose roots, z_1 and z_2 , determine the solution structure of the integral equation.

Assume that the discriminant of equation (8) is positive:

$$D \equiv (A_1\lambda_1 - A_2\lambda_2 - \lambda_1^2 + \lambda_2^2)^2 + 4A_1A_2\lambda_1\lambda_2 > 0.$$

In this case, the quadratic equation (8) has the real (different) roots

$$z_1 = \frac{1}{2}(\lambda_1^2 + \lambda_2^2 - A_1\lambda_1 - A_2\lambda_2) + \frac{1}{2}\sqrt{D}, \quad z_2 = \frac{1}{2}(\lambda_1^2 + \lambda_2^2 - A_1\lambda_1 - A_2\lambda_2) - \frac{1}{2}\sqrt{D}.$$

Depending on the signs of z_1 and z_2 the following three cases are possible.

Case 1. If $z_1 > 0$ and $z_2 > 0$, then the solution of the integral equation has the form ($i = 1, 2$):

$$y(x) = f(x) + \int_a^x \{B_1 \sinh[\mu_1(x-t)] + B_2 \sinh[\mu_2(x-t)]\} f(t) dt, \quad \mu_i = \sqrt{z_i},$$

where

$$B_1 = A_1 \frac{\lambda_1(\mu_1^2 - \lambda_2^2)}{\mu_1(\mu_2^2 - \mu_1^2)} + A_2 \frac{\lambda_2(\mu_1^2 - \lambda_1^2)}{\mu_1(\mu_2^2 - \mu_1^2)}, \quad B_2 = A_1 \frac{\lambda_1(\mu_2^2 - \lambda_2^2)}{\mu_2(\mu_1^2 - \mu_2^2)} + A_2 \frac{\lambda_2(\mu_2^2 - \lambda_1^2)}{\mu_2(\mu_1^2 - \mu_2^2)}.$$

Case 2. If $z_1 < 0$ and $z_2 < 0$, then the solution of the integral equation has the form

$$y(x) = f(x) + \int_a^x \{B_1 \sin[\mu_1(x-t)] + B_2 \sin[\mu_2(x-t)]\} f(t) dt, \quad \mu_i = \sqrt{|z_i|},$$

where the coefficients B_1 and B_2 are found by solving the following system of linear algebraic equations:

$$\frac{B_1\mu_1}{\lambda_1^2 + \mu_1^2} + \frac{B_2\mu_2}{\lambda_1^2 + \mu_2^2} + 1 = 0, \quad \frac{B_1\mu_1}{\lambda_2^2 + \mu_1^2} + \frac{B_2\mu_2}{\lambda_2^2 + \mu_2^2} + 1 = 0.$$

Case 3. If $z_1 > 0$ and $z_2 < 0$, then the solution of the integral equation has the form

$$y(x) = f(x) + \int_a^x \{B_1 \sinh[\mu_1(x-t)] + B_2 \sin[\mu_2(x-t)]\} f(t) dt, \quad \mu_i = \sqrt{|z_i|},$$

where B_1 and B_2 are determined from the following system of linear algebraic equations:

$$\frac{B_1\mu_1}{\lambda_1^2 - \mu_1^2} + \frac{B_2\mu_2}{\lambda_1^2 + \mu_2^2} + 1 = 0, \quad \frac{B_1\mu_1}{\lambda_2^2 - \mu_1^2} + \frac{B_2\mu_2}{\lambda_2^2 + \mu_2^2} + 1 = 0.$$

$$19. \quad y(x) + \int_a^x \left\{ \sum_{k=1}^n A_k \sinh[\lambda_k(x-t)] \right\} y(t) dt = f(x).$$

1°. This equation can be reduced to an equation of the form 2.2.19 with the aid of the formula $\sinh z = \frac{1}{2}(e^z - e^{-z})$. Therefore, the original integral equation can be reduced to a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients.

2°. Let us find the roots z_k of the algebraic equation

$$\sum_{k=1}^n \frac{\lambda_k A_k}{z - \lambda_k^2} + 1 = 0. \quad (1)$$

By reducing it to a common denominator, we arrive at the problem of determining the roots of an n th-degree characteristic polynomial.

Assume that all z_k are real, different, and nonzero. Let us divide the roots into two groups

$$\begin{aligned} z_1 > 0, \quad z_2 > 0, \quad \dots, \quad z_s > 0 & \quad (\text{positive roots}); \\ z_{s+1} < 0, \quad z_{s+2} < 0, \quad \dots, \quad z_n < 0 & \quad (\text{negative roots}). \end{aligned}$$

Then the solution of the integral equation can be written in the form

$$y(x) = f(x) + \int_a^x \left\{ \sum_{k=1}^s B_k \sinh[\mu_k(x-t)] + \sum_{k=s+1}^n C_k \sin[\mu_k(x-t)] \right\} f(t) dt, \quad \mu_k = \sqrt{|z_k|}. \quad (2)$$

The coefficients B_k and C_k are determined from the following system of linear algebraic equations:

$$\sum_{k=0}^s \frac{B_k \mu_k}{\lambda_m^2 - \mu_k^2} + \sum_{k=s+1}^n \frac{C_k \mu_k}{\lambda_m^2 + \mu_k^2} + 1 = 0, \quad \mu_k = \sqrt{|z_k|}, \quad m = 1, \dots, n. \quad (3)$$

In the case of a nonzero root $z_s = 0$, we can introduce the new constant $D = B_s \mu_s$ and proceed to the limit $\mu_s \rightarrow 0$. As a result, the term $D(x-t)$ appears in solution (2) instead of $B_s \sinh[\mu_s(x-t)]$ and the corresponding terms $D\lambda_m^{-2}$ appear in system (3).

$$20. \quad y(x) - A \int_a^x \frac{\sinh(\lambda x)}{\sinh(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\sinh(\lambda x)}{\sinh(\lambda t)} f(t) dt.$$

$$21. \quad y(x) - A \int_a^x \frac{\sinh(\lambda t)}{\sinh(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\sinh(\lambda t)}{\sinh(\lambda x)} f(t) dt.$$

$$22. \quad y(x) - A \int_a^x \sinh^k(\lambda x) \sinh^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sinh^k(\lambda x)$ and $h(t) = \sinh^m(\mu t)$.

$$23. \quad y(x) + A \int_a^x t \sinh[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.30 with $g(t) = At$.

Solution:

$$y(x) = f(x) + \frac{A\lambda}{W} \int_a^x t [u_1(x)u_2(t) - u_2(x)u_1(t)] f(t) dt,$$

where $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear ordinary differential equation $u''_{xx} + \lambda(Ax - \lambda)u = 0$, and W is the Wronskian.

The functions $u_1(x)$ and $u_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of $A\lambda$, as follows:

if $A\lambda > 0$, then

$$u_1(x) = \xi^{1/2} J_{1/3}(\frac{2}{3}\sqrt{A\lambda}\xi^{3/2}), \quad u_2(x) = \xi^{1/2} Y_{1/3}(\frac{2}{3}\sqrt{A\lambda}\xi^{3/2}), \\ W = 3/\pi, \quad \xi = x - (\lambda/A);$$

if $A\lambda < 0$, then

$$u_1(x) = \xi^{1/2} I_{1/3}(\frac{2}{3}\sqrt{-A\lambda}\xi^{3/2}), \quad u_2(x) = \xi^{1/2} K_{1/3}(\frac{2}{3}\sqrt{-A\lambda}\xi^{3/2}), \\ W = -\frac{3}{2}, \quad \xi = x - (\lambda/A).$$

$$24. \quad y(x) + A \int_a^x x \sinh[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.31 with $g(x) = Ax$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + \frac{A\lambda}{W} \int_a^x x [u_1(x)u_2(t) - u_2(x)u_1(t)] f(t) dt,$$

where $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear ordinary differential equation $u''_{xx} + \lambda(Ax - \lambda)u = 0$, and W is the Wronskian.

The functions $u_1(x), u_2(x)$, and W are specified in 2.3.23.

$$25. \quad y(x) + A \int_a^x t^k \sinh^m(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -A \sinh^m(\lambda x)$ and $h(t) = t^k$.

$$26. \quad y(x) + A \int_a^x x^k \sinh^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ax^k$ and $h(t) = \sinh^m(\lambda t)$.

$$27. \quad y(x) - \int_a^x [A \sinh(kx) + B - AB(x-t) \sinh(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \sinh(kx)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t) f(t) dt, \\ R(x,t) = [A \sinh(kx) + B] \frac{G(x)}{G(t)} + \frac{B^2}{G(t)} \int_t^x e^{B(x-s)} G(s) ds, \quad G(x) = \exp\left[\frac{A}{k} \cosh(kx)\right].$$

$$28. \quad y(x) + \int_a^x [A \sinh(kt) + B + AB(x-t) \sinh(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \sinh(kt)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t) f(t) dt,$$

$$R(x,t) = -[\sinh(kt) + B] \frac{G(t)}{G(x)} + \frac{B^2}{G(x)} \int_t^x e^{B(t-s)} G(s) ds, \quad G(x) = \exp \left[\frac{A}{k} \cosh(kx) \right].$$

$$29. \quad y(x) + A \int_x^\infty \sinh(\lambda \sqrt{t-x}) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = A \sinh(\lambda \sqrt{-x})$.

2.3-3. Kernels Containing Hyperbolic Tangent.

$$30. \quad y(x) - A \int_a^x \tanh(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tanh(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \tanh(\lambda x) \left[\frac{\cosh(\lambda x)}{\cosh(\lambda t)} \right]^{A/\lambda} f(t) dt.$$

$$31. \quad y(x) - A \int_a^x \tanh(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \tanh(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \tanh(\lambda t) \left[\frac{\cosh(\lambda x)}{\cosh(\lambda t)} \right]^{A/\lambda} f(t) dt.$$

$$32. \quad y(x) + A \int_a^x [\tanh(\lambda x) - \tanh(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.5 with $g(x) = A \tanh(\lambda x)$.

Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x [Y_1'(x) Y_2'(t) - Y_2'(x) Y_1'(t)] f(t) dt,$$

where $Y_1(x), Y_2(x)$ is a fundamental system of solutions of the second-order linear ordinary differential equation $\cosh^2(\lambda x) Y_{xx} + A \lambda Y = 0$, W is the Wronskian, and the primes stand for the differentiation with respect to the argument specified in the parentheses.

As shown in A. D. Polyanin and V. F. Zaitsev (2003), the functions $Y_1(x)$ and $Y_2(x)$ can be represented in the form

$$Y_1(x) = F\left(\alpha, \beta, 1; \frac{e^{\lambda x}}{1 + e^{\lambda x}}\right), \quad Y_2(x) = Y_1(x) \int_a^x \frac{d\xi}{Y_1^2(\xi)}, \quad W = 1,$$

where $F(\alpha, \beta, \gamma; z)$ is the hypergeometric function, in which α and β are determined from the algebraic system $\alpha + \beta = 1$, $\alpha\beta = -A/\lambda$.

$$33. \quad y(x) - A \int_a^x \frac{\tanh(\lambda x)}{\tanh(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\tanh(\lambda x)}{\tanh(\lambda t)} f(t) dt.$$

$$34. \quad y(x) - A \int_a^x \frac{\tanh(\lambda t)}{\tanh(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\tanh(\lambda t)}{\tanh(\lambda x)} f(t) dt.$$

$$35. \quad y(x) - A \int_a^x \tanh^k(\lambda x) \tanh^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tanh^k(\lambda x)$ and $h(t) = \tanh^m(\mu t)$.

$$36. \quad y(x) + A \int_a^x t^k \tanh^m(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -A \tanh^m(\lambda x)$ and $h(t) = t^k$.

$$37. \quad y(x) + A \int_a^x x^k \tanh^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ax^k$ and $h(t) = \tanh^m(\lambda t)$.

$$38. \quad y(x) + A \int_x^\infty \tanh[\lambda(t-x)] y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(z) = A \tanh(-\lambda z)$.

$$39. \quad y(x) + A \int_x^\infty \tanh(\lambda\sqrt{t-x}) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(z) = A \tanh(\lambda\sqrt{-z})$.

$$40. \quad y(x) - \int_a^x [A \tanh(kx) + B - AB(x-t) \tanh(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \tanh(kx)$.

$$41. \quad y(x) + \int_a^x [A \tanh(kt) + B + AB(x-t) \tanh(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \tanh(kt)$.

2.3-4. Kernels Containing Hyperbolic Cotangent.

$$42. \quad y(x) - A \int_a^x \coth(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \coth(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \coth(\lambda x) \left[\frac{\sinh(\lambda x)}{\sinh(\lambda t)} \right]^{A/\lambda} f(t) dt.$$

$$43. \quad y(x) - A \int_a^x \coth(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \coth(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \coth(\lambda t) \left[\frac{\sinh(\lambda x)}{\sinh(\lambda t)} \right]^{A/\lambda} f(t) dt.$$

$$44. \quad y(x) - A \int_a^x \frac{\coth(\lambda t)}{\coth(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\coth(\lambda t)}{\coth(\lambda x)} f(t) dt.$$

$$45. \quad y(x) - A \int_a^x \frac{\coth(\lambda x)}{\coth(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\coth(\lambda x)}{\coth(\lambda t)} f(t) dt.$$

$$46. \quad y(x) - A \int_a^x \coth^k(\lambda x) \coth^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \coth^k(\lambda x)$ and $h(t) = \coth^m(\mu t)$.

$$47. \quad y(x) + A \int_a^x t^k \coth^m(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -A \coth^m(\lambda x)$ and $h(t) = t^k$.

$$48. \quad y(x) + A \int_a^x x^k \coth^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ax^k$ and $h(t) = \coth^m(\lambda t)$.

$$49. \quad y(x) + A \int_x^\infty \coth[\lambda(t-x)] y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(z) = A \coth(-\lambda z)$.

$$50. \quad y(x) + A \int_x^\infty \coth(\lambda \sqrt{t-x}) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(z) = A \coth(\lambda \sqrt{-z})$.

$$51. \quad y(x) - \int_a^x [A \coth(kx) + B - AB(x-t) \coth(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \coth(kx)$.

$$52. \quad y(x) + \int_a^x [A \coth(kt) + B + AB(x-t) \coth(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \coth(kt)$.

2.3-5. Kernels Containing Combinations of Hyperbolic Functions.

53. $y(x) - A \int_a^x \cosh^k(\lambda x) \sinh^m(\mu t) y(t) dt = f(x).$

This is a special case of equation 2.9.2 with $g(x) = A \cosh^k(\lambda x)$ and $h(t) = \sinh^m(\mu t)$.

54. $y(x) - \int_a^x \{A + B \cosh(\lambda x) + B(x-t)[\lambda \sinh(\lambda x) - A \cosh(\lambda x)]\} y(t) dt = f(x).$

This is a special case of equation 2.9.32 with $b = B$ and $g(x) = A$.

55. $y(x) - \int_a^x \{A + B \sinh(\lambda x) + B(x-t)[\lambda \cosh(\lambda x) - A \sinh(\lambda x)]\} y(t) dt = f(x).$

This is a special case of equation 2.9.33 with $b = B$ and $g(x) = A$.

56. $y(x) - A \int_a^x \tanh^k(\lambda x) \coth^m(\mu t) y(t) dt = f(x).$

This is a special case of equation 2.9.2 with $g(x) = A \tanh^k(\lambda x)$ and $h(t) = \coth^m(\mu t)$.

2.4. Equations Whose Kernels Contain Logarithmic Functions

2.4-1. Kernels Containing Logarithmic Functions.
--

1. $y(x) - A \int_a^x \ln(\lambda x) y(t) dt = f(x).$

This is a special case of equation 2.9.2 with $g(x) = A \ln(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \ln(\lambda x) e^{-A(x-t)} \frac{(\lambda x)^{Ax}}{(\lambda t)^{At}} f(t) dt.$$

2. $y(x) - A \int_a^x \ln(\lambda t) y(t) dt = f(x).$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \ln(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \ln(\lambda t) e^{-A(x-t)} \frac{(\lambda x)^{Ax}}{(\lambda t)^{At}} f(t) dt.$$

3. $y(x) + A \int_a^x (\ln x - \ln t) y(t) dt = f(x).$

This is a special case of equation 2.9.5 with $g(x) = A \ln x$.

Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x [u_1'(x) u_2'(t) - u_2'(x) u_1'(t)] f(t) dt,$$

where the primes denote differentiation with respect to the argument specified in the parentheses; and $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $u''_{xx} + Ax^{-1}u = 0$, with $u_1(x)$ and $u_2(x)$ expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of A :

$$\begin{aligned} W = \frac{1}{\pi}, \quad u_1(x) &= \sqrt{x} J_1(2\sqrt{Ax}), \quad u_2(x) = \sqrt{x} Y_1(2\sqrt{Ax}) & \text{for } A > 0, \\ W = -\frac{1}{2}, \quad u_1(x) &= \sqrt{x} I_1(2\sqrt{-Ax}), \quad u_2(x) = \sqrt{x} K_1(2\sqrt{-Ax}) & \text{for } A < 0. \end{aligned}$$

$$4. \quad y(x) - A \int_a^x \frac{\ln(\lambda x)}{\ln(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\ln(\lambda x)}{\ln(\lambda t)} f(t) dt.$$

$$5. \quad y(x) - A \int_a^x \frac{\ln(\lambda t)}{\ln(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\ln(\lambda t)}{\ln(\lambda x)} f(t) dt.$$

$$6. \quad y(x) - A \int_a^x \ln^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$7. \quad y(x) + a \int_x^\infty \ln(t-x) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = a \ln(-x)$.

For $f(x) = \sum_{k=1}^m A_k \exp(-\lambda_k x)$, where $\lambda_k > 0$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^m \frac{A_k}{B_k} \exp(-\lambda_k x), \quad B_k = 1 - \frac{a}{\lambda_k} (\ln \lambda_k + \mathcal{C}),$$

where $\mathcal{C} = 0.5772 \dots$ is the Euler constant.

$$8. \quad y(x) + a \int_x^\infty \ln^2(t-x) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = a \ln^2(-x)$.

For $f(x) = \sum_{k=1}^m A_k \exp(-\lambda_k x)$, where $\lambda_k > 0$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^m \frac{A_k}{B_k} \exp(-\lambda_k x), \quad B_k = 1 + \frac{a}{\lambda_k} \left[\frac{1}{6} \pi^2 + (\ln \lambda_k + \mathcal{C})^2 \right],$$

where $\mathcal{C} = 0.5772 \dots$ is the Euler constant.

2.4-2. Kernels Containing Power-Law and Logarithmic Functions.

$$9. \quad y(x) - A \int_a^x x^k \ln^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = Ax^k$ and $h(t) = \ln^m(\lambda t)$.

$$10. \quad y(x) - A \int_a^x t^k \ln^m(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\lambda x)$ and $h(t) = t^k$.

$$11. \quad y(x) - \int_a^x [A \ln(kx) + B - AB(x-t) \ln(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \ln(kx)$.

$$12. \quad y(x) + \int_a^x [A \ln(kt) + B + AB(x-t) \ln(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \ln(kt)$.

$$13. \quad y(x) + a \int_x^\infty (t-x)^n \ln(t-x) y(t) dt = f(x), \quad n = 1, 2, \dots$$

For $f(x) = \sum_{k=1}^m A_k \exp(-\lambda_k x)$, where $\lambda_k > 0$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^m \frac{A_k}{B_k} \exp(-\lambda_k x), \quad B_k = 1 + \frac{an!}{\lambda_k^{n+1}} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} - \ln \lambda_k - \mathcal{C}\right),$$

where $\mathcal{C} = 0.5772 \dots$ is the Euler constant.

$$14. \quad y(x) + a \int_x^\infty \frac{\ln(t-x)}{\sqrt{t-x}} y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(-x) = ax^{-1/2} \ln x$.

For $f(x) = \sum_{k=1}^m A_k \exp(-\lambda_k x)$, where $\lambda_k > 0$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^m \frac{A_k}{B_k} \exp(-\lambda_k x), \quad B_k = 1 - a \sqrt{\frac{\pi}{\lambda_k}} [\ln(4\lambda_k) + \mathcal{C}],$$

where $\mathcal{C} = 0.5772 \dots$ is the Euler constant.

2.5. Equations Whose Kernels Contain Trigonometric Functions

2.5-1. Kernels Containing Cosine.

$$1. \quad y(x) - A \int_a^x \cos(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cos(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \cos(\lambda x) \exp\left\{\frac{A}{\lambda} [\sin(\lambda x) - \sin(\lambda t)]\right\} f(t) dt.$$

$$2. \quad y(x) - A \int_a^x \cos(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \cos(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \cos(\lambda t) \exp\left\{\frac{A}{\lambda} [\sin(\lambda x) - \sin(\lambda t)]\right\} f(t) dt.$$

$$3. \quad y(x) + A \int_a^x \cos[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.34 with $g(t) = A$. Therefore, solving this integral equation is reduced to solving the following second-order linear nonhomogeneous ordinary differential equation with constant coefficients:

$$y''_{xx} + Ay'_x + \lambda^2 y = f''_{xx} + \lambda^2 f, \quad f = f(x),$$

with the initial conditions

$$y(a) = f(a), \quad y'_x(a) = f'_x(a) - Af(a).$$

1°. Solution with $|A| > 2|\lambda|$:

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt,$$

$$R(x) = \exp(-\frac{1}{2}Ax) \left[\frac{A^2}{2k} \sinh(kx) - A \cosh(kx) \right], \quad k = \sqrt{\frac{1}{4}A^2 - \lambda^2}.$$

2°. Solution with $|A| < 2|\lambda|$:

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt,$$

$$R(x) = \exp(-\frac{1}{2}Ax) \left[\frac{A^2}{2k} \sin(kx) - A \cos(kx) \right], \quad k = \sqrt{\lambda^2 - \frac{1}{4}A^2}.$$

3°. Solution with $\lambda = \pm \frac{1}{2}A$:

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt, \quad R(x) = \exp(-\frac{1}{2}Ax) \left(\frac{1}{2}A^2 x - A \right).$$

$$4. \quad y(x) + \int_a^x \left\{ \sum_{k=1}^n A_k \cos[\lambda_k(x-t)] \right\} y(t) dt = f(x).$$

This integral equation is reduced to a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients. Set

$$I_k(x) = \int_a^x \cos[\lambda_k(x-t)] y(t) dt. \quad (1)$$

Differentiating (1) with respect to x twice yields

$$I'_k = y(x) - \lambda_k \int_a^x \sin[\lambda_k(x-t)] y(t) dt, \quad (2)$$

$$I''_k = y'_x(x) - \lambda_k^2 \int_a^x \cos[\lambda_k(x-t)] y(t) dt,$$

where the primes stand for differentiation with respect to x . Comparing (1) and (2), we see that

$$I''_k = y'_x(x) - \lambda_k^2 I_k, \quad I_k = I_k(x). \quad (3)$$

With the aid of (1), the integral equation can be rewritten in the form

$$y(x) + \sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x twice taking into account (3) yields

$$y''_{xx}(x) + \sigma_n y'_x(x) - \sum_{k=1}^n A_k \lambda_k^2 I_k = f''_{xx}(x), \quad \sigma_n = \sum_{k=1}^n A_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5), we obtain

$$y''_{xx}(x) + \sigma_n y'_x(x) + \lambda_n^2 y(x) + \sum_{k=1}^{n-1} A_k (\lambda_n^2 - \lambda_k^2) I_k = f''_{xx}(x) + \lambda_n^2 f(x). \quad (6)$$

Differentiating (6) with respect to x twice followed by eliminating I_{n-1} from the resulting expression with the aid of (6) yields a similar equation whose left-hand side is a fourth-order differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} B_k I_k$. Successively eliminating the terms I_{n-2}, I_{n-3}, \dots using double differentiation and formula (3), we finally arrive at a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients.

The initial conditions for $y(x)$ can be obtained by setting $x = a$ in the integral equation and all its derivative equations.

5. $y(x) - A \int_a^x \frac{\cos(\lambda x)}{\cos(\lambda t)} y(t) dt = f(x).$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\cos(\lambda x)}{\cos(\lambda t)} f(t) dt.$$

6. $y(x) - A \int_a^x \frac{\cos(\lambda t)}{\cos(\lambda x)} y(t) dt = f(x).$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\cos(\lambda t)}{\cos(\lambda x)} f(t) dt.$$

7. $y(x) - A \int_a^x \cos^k(\lambda x) \cos^m(\mu t) y(t) dt = f(x).$

This is a special case of equation 2.9.2 with $g(x) = A \cos^k(\lambda x)$ and $h(t) = \cos^m(\mu t)$.

8. $y(x) + A \int_a^x t \cos[\lambda(x-t)] y(t) dt = f(x).$

This is a special case of equation 2.9.34 with $g(t) = At$.

9. $y(x) + A \int_a^x t^k \cos^m(\lambda x) y(t) dt = f(x).$

This is a special case of equation 2.9.2 with $g(x) = -A \cos^m(\lambda x)$ and $h(t) = t^k$.

$$10. \quad y(x) + A \int_a^x x^k \cos^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ax^k$ and $h(t) = \cos^m(\lambda t)$.

$$11. \quad y(x) - \int_a^x [A \cos(kx) + B - AB(x-t) \cos(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \cos(kx)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x, t) f(t) dt,$$

$$R(x, t) = [A \cos(kx) + B] \frac{G(x)}{G(t)} + \frac{B^2}{G(t)} \int_t^x e^{B(x-s)} G(s) ds, \quad G(x) = \exp \left[\frac{A}{k} \sin(kx) \right].$$

$$12. \quad y(x) + \int_a^x [A \cos(kt) + B + AB(x-t) \cos(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \cos(kt)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x, t) f(t) dt,$$

$$R(x, t) = -[A \cos(kt) + B] \frac{G(t)}{G(x)} + \frac{B^2}{G(x)} \int_t^x e^{B(t-s)} G(s) ds, \quad G(x) = \exp \left[\frac{A}{k} \sin(kx) \right].$$

$$13. \quad y(x) + A \int_x^\infty \cos(\lambda \sqrt{t-x}) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = A \cos(\lambda \sqrt{-x})$.

2.5-2. Kernels Containing Sine.

$$14. \quad y(x) - A \int_a^x \sin(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sin(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \sin(\lambda x) \exp \left\{ \frac{A}{\lambda} [\cos(\lambda t) - \cos(\lambda x)] \right\} f(t) dt.$$

$$15. \quad y(x) - A \int_a^x \sin(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \sin(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \sin(\lambda t) \exp \left\{ \frac{A}{\lambda} [\cos(\lambda t) - \cos(\lambda x)] \right\} f(t) dt.$$

$$16. \quad y(x) + A \int_a^x \sin[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.36 with $g(t) = A$.

1°. Solution with $\lambda(A + \lambda) > 0$:

$$y(x) = f(x) - \frac{A\lambda}{k} \int_a^x \sin[k(x-t)] f(t) dt, \quad \text{where } k = \sqrt{\lambda(A + \lambda)}.$$

2°. Solution with $\lambda(A + \lambda) < 0$:

$$y(x) = f(x) - \frac{A\lambda}{k} \int_a^x \sinh[k(x-t)] f(t) dt, \quad \text{where } k = \sqrt{-\lambda(\lambda + A)}.$$

3°. Solution with $A = -\lambda$:

$$y(x) = f(x) + \lambda^2 \int_a^x (x-t) f(t) dt.$$

$$17. \quad y(x) + A \int_a^x \sin^3[\lambda(x-t)] y(t) dt = f(x).$$

Using the formula $\sin^3 \beta = -\frac{1}{4} \sin 3\beta + \frac{3}{4} \sin \beta$, we arrive at an equation of the form 2.5.18:

$$y(x) + \int_a^x \left\{ -\frac{1}{4} A \sin[3\lambda(x-t)] + \frac{3}{4} A \sin[\lambda(x-t)] \right\} y(t) dt = f(x).$$

$$18. \quad y(x) + \int_a^x \{ A_1 \sin[\lambda_1(x-t)] + A_2 \sin[\lambda_2(x-t)] \} y(t) dt = f(x).$$

This equation can be solved by the same method as equation 2.3.18, by reducing it to a fourth-order linear ordinary differential equation with constant coefficients.

Consider the characteristic equation

$$z^2 + (\lambda_1^2 + \lambda_2^2 + A_1 \lambda_1 + A_2 \lambda_2) z + \lambda_1^2 \lambda_2^2 + A_1 \lambda_1 \lambda_2^2 + A_2 \lambda_1^2 \lambda_2 = 0, \quad (1)$$

whose roots, z_1 and z_2 , determine the solution structure of the integral equation.

Assume that the discriminant of equation (1) is positive:

$$D \equiv (A_1 \lambda_1 - A_2 \lambda_2 + \lambda_1^2 - \lambda_2^2)^2 + 4 A_1 A_2 \lambda_1 \lambda_2 > 0.$$

In this case, the quadratic equation (1) has the real (different) roots

$$z_1 = -\frac{1}{2}(\lambda_1^2 + \lambda_2^2 + A_1 \lambda_1 + A_2 \lambda_2) + \frac{1}{2}\sqrt{D}, \quad z_2 = -\frac{1}{2}(\lambda_1^2 + \lambda_2^2 + A_1 \lambda_1 + A_2 \lambda_2) - \frac{1}{2}\sqrt{D}.$$

Depending on the signs of z_1 and z_2 the following three cases are possible.

Case 1. If $z_1 > 0$ and $z_2 > 0$, then the solution of the integral equation has the form ($i = 1, 2$):

$$y(x) = f(x) + \int_a^x \{ B_1 \sinh[\mu_1(x-t)] + B_2 \sinh[\mu_2(x-t)] \} f(t) dt, \quad \mu_i = \sqrt{z_i},$$

where the coefficients B_1 and B_2 are determined from the following system of linear algebraic equations:

$$\frac{B_1 \mu_1}{\lambda_1^2 + \mu_1^2} + \frac{B_2 \mu_2}{\lambda_1^2 + \mu_2^2} - 1 = 0, \quad \frac{B_1 \mu_1}{\lambda_2^2 + \mu_1^2} + \frac{B_2 \mu_2}{\lambda_2^2 + \mu_2^2} - 1 = 0.$$

Case 2. If $z_1 < 0$ and $z_2 < 0$, then the solution of the integral equation has the form

$$y(x) = f(x) + \int_a^x \{B_1 \sin[\mu_1(x-t)] + B_2 \sin[\mu_2(x-t)]\} f(t) dt, \quad \mu_i = \sqrt{|z_i|},$$

where B_1 and B_2 are determined from the system

$$\frac{B_1\mu_1}{\lambda_1^2 - \mu_1^2} + \frac{B_2\mu_2}{\lambda_1^2 - \mu_2^2} - 1 = 0, \quad \frac{B_1\mu_1}{\lambda_2^2 - \mu_1^2} + \frac{B_2\mu_2}{\lambda_2^2 - \mu_2^2} - 1 = 0.$$

Case 3. If $z_1 > 0$ and $z_2 < 0$, then the solution of the integral equation has the form

$$y(x) = f(x) + \int_a^x \{B_1 \sinh[\mu_1(x-t)] + B_2 \sin[\mu_2(x-t)]\} f(t) dt, \quad \mu_i = \sqrt{|z_i|},$$

where B_1 and B_2 are determined from the system

$$\frac{B_1\mu_1}{\lambda_1^2 + \mu_1^2} + \frac{B_2\mu_2}{\lambda_1^2 - \mu_2^2} - 1 = 0, \quad \frac{B_1\mu_1}{\lambda_2^2 + \mu_1^2} + \frac{B_2\mu_2}{\lambda_2^2 - \mu_2^2} - 1 = 0.$$

Remark. The solution of the original integral equation can be obtained from the solution of equation 2.3.18 by performing the following change of parameters:

$$\lambda_k \rightarrow i\lambda_k, \quad \mu_k \rightarrow i\mu_k, \quad A_k \rightarrow -iA_k, \quad B_k \rightarrow -iB_k, \quad i^2 = -1 \quad (k = 1, 2).$$

19.
$$y(x) + \int_a^x \left\{ \sum_{k=1}^n A_k \sin[\lambda_k(x-t)] \right\} y(t) dt = f(x).$$

1°. This integral equation can be reduced to a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients. Set

$$I_k(x) = \int_a^x \sin[\lambda_k(x-t)] y(t) dt. \quad (1)$$

Differentiating (1) with respect to x twice yields

$$I_k' = \lambda_k \int_a^x \cos[\lambda_k(x-t)] y(t) dt, \quad I_k'' = \lambda_k y(x) - \lambda_k^2 \int_a^x \sin[\lambda_k(x-t)] y(t) dt, \quad (2)$$

where the primes stand for differentiation with respect to x . Comparing (1) and (2), we see that

$$I_k'' = \lambda_k y(x) - \lambda_k^2 I_k, \quad I_k = I_k(x). \quad (3)$$

With aid of (1), the integral equation can be rewritten in the form

$$y(x) + \sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x twice taking into account (3) yields

$$y''_{xx}(x) + \sigma_n y(x) - \sum_{k=1}^n A_k \lambda_k^2 I_k = f''_{xx}(x), \quad \sigma_n = \sum_{k=1}^n A_k \lambda_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5), we obtain

$$y''_{xx}(x) + (\sigma_n + \lambda_n^2)y(x) + \sum_{k=1}^{n-1} A_k(\lambda_n^2 - \lambda_k^2)I_k = f''_{xx}(x) + \lambda_n^2 f(x). \quad (6)$$

Differentiating (6) with respect to x twice followed by eliminating I_{n-1} from the resulting expression with the aid of (6) yields a similar equation whose left-hand side is a fourth-order differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} B_k I_k$. Successively eliminating the terms I_{n-2}, I_{n-3}, \dots using double differentiation and formula (3), we finally arrive at a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients.

The initial conditions for $y(x)$ can be obtained by setting $x = a$ in the integral equation and all its derivative equations.

2°. Let us find the roots z_k of the algebraic equation

$$\sum_{k=1}^n \frac{\lambda_k A_k}{z + \lambda_k^2} + 1 = 0. \quad (7)$$

By reducing it to a common denominator, we arrive at the problem of determining the roots of an n th-degree characteristic polynomial.

Assume that all z_k are real, different, and nonzero. Let us divide the roots into two groups

$$\begin{aligned} z_1 > 0, \quad z_2 > 0, \quad \dots, \quad z_s > 0 & \quad (\text{positive roots}); \\ z_{s+1} < 0, \quad z_{s+2} < 0, \quad \dots, \quad z_n < 0 & \quad (\text{negative roots}). \end{aligned}$$

Then the solution of the integral equation can be written in the form

$$y(x) = f(x) + \int_a^x \left\{ \sum_{k=1}^s B_k \sinh[\mu_k(x-t)] + \sum_{k=s+1}^n C_k \sin[\mu_k(x-t)] \right\} f(t) dt, \quad \mu_k = \sqrt{|z_k|}. \quad (8)$$

The coefficients B_k and C_k are determined from the following system of linear algebraic equations:

$$\sum_{k=0}^s \frac{B_k \mu_k}{\lambda_m^2 + \mu_k^2} + \sum_{k=s+1}^n \frac{C_k \mu_k}{\lambda_m^2 - \mu_k^2} - 1 = 0, \quad \mu_k = \sqrt{|z_k|} \quad m = 1, 2, \dots, n. \quad (9)$$

In the case of a nonzero root $z_s = 0$, we can introduce the new constant $D = B_s \mu_s$ and proceed to the limit $\mu_s \rightarrow 0$. As a result, the term $D(x-t)$ appears in solution (8) instead of $B_s \sinh[\mu_s(x-t)]$ and the corresponding terms $D\lambda_m^{-2}$ appear in system (9).

20. $y(x) - A \int_a^x \frac{\sin(\lambda x)}{\sin(\lambda t)} y(t) dt = f(x).$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\sin(\lambda x)}{\sin(\lambda t)} f(t) dt.$$

21. $y(x) - A \int_a^x \frac{\sin(\lambda t)}{\sin(\lambda x)} y(t) dt = f(x).$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\sin(\lambda t)}{\sin(\lambda x)} f(t) dt.$$

$$22. \quad y(x) - A \int_a^x \sin^k(\lambda x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sin^k(\lambda x)$ and $h(t) = \sin^m(\mu t)$.

$$23. \quad y(x) + A \int_a^x t \sin[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.36 with $g(t) = At$.

Solution:

$$y(x) = f(x) + \frac{A\lambda}{W} \int_a^x t [u_1(x)u_2(t) - u_2(x)u_1(t)] f(t) dt,$$

where $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear ordinary differential equation $u''_{xx} + \lambda(Ax + \lambda)u = 0$, and W is the Wronskian.

Depending on the sign of $A\lambda$, the functions $u_1(x)$ and $u_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions as follows:

if $A\lambda > 0$, then

$$u_1(x) = \xi^{1/2} J_{1/3} \left(\frac{2}{3} \sqrt{A\lambda} \xi^{3/2} \right), \quad u_2(x) = \xi^{1/2} Y_{1/3} \left(\frac{2}{3} \sqrt{A\lambda} \xi^{3/2} \right), \\ W = 3/\pi, \quad \xi = x + (\lambda/A);$$

if $A\lambda < 0$, then

$$u_1(x) = \xi^{1/2} I_{1/3} \left(\frac{2}{3} \sqrt{-A\lambda} \xi^{3/2} \right), \quad u_2(x) = \xi^{1/2} K_{1/3} \left(\frac{2}{3} \sqrt{-A\lambda} \xi^{3/2} \right), \\ W = -\frac{3}{2}, \quad \xi = x + (\lambda/A).$$

$$24. \quad y(x) + A \int_a^x x \sin[\lambda(x-t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.37 with $g(x) = Ax$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + \frac{A\lambda}{W} \int_a^x x [u_1(x)u_2(t) - u_2(x)u_1(t)] f(t) dt,$$

where $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear ordinary differential equation $u''_{xx} + \lambda(Ax + \lambda)u = 0$, and W is the Wronskian.

The functions $u_1(x), u_2(x)$, and W are specified in 2.5.23.

$$25. \quad y(x) + A \int_a^x t^k \sin^m(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -A \sin^m(\lambda x)$ and $h(t) = t^k$.

$$26. \quad y(x) + A \int_a^x x^k \sin^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ax^k$ and $h(t) = \sin^m(\lambda t)$.

$$27. \quad y(x) - \int_a^x [A \sin(kx) + B - AB(x-t) \sin(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \sin(kx)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t) f(t) dt, \\ R(x,t) = [A \sin(kx) + B] \frac{G(x)}{G(t)} + \frac{B^2}{G(t)} \int_t^x e^{B(x-s)} G(s) ds, \quad G(x) = \exp \left[-\frac{A}{k} \cos(kx) \right].$$

$$28. \quad y(x) + \int_a^x [A \sin(kt) + B + AB(x-t) \sin(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \sin(kt)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t) f(t) dt,$$

$$R(x,t) = -[A \sin(kt) + B] \frac{G(t)}{G(x)} + \frac{B^2}{G(x)} \int_t^x e^{B(t-s)} G(s) ds, \quad G(x) = \exp\left[-\frac{A}{k} \cos(kx)\right].$$

$$29. \quad y(x) + A \int_x^\infty \sin(\lambda\sqrt{t-x}) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = A \sin(\lambda\sqrt{-x})$.

2.5-3. Kernels Containing Tangent.

$$30. \quad y(x) - A \int_a^x \tan(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tan(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \tan(\lambda x) \left| \frac{\cos(\lambda t)}{\cos(\lambda x)} \right|^{A/\lambda} f(t) dt.$$

$$31. \quad y(x) - A \int_a^x \tan(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \tan(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \tanh(\lambda t) \left| \frac{\cos(\lambda t)}{\cos(\lambda x)} \right|^{A/\lambda} f(t) dt.$$

$$32. \quad y(x) + A \int_a^x [\tan(\lambda x) - \tan(\lambda t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.5 with $g(x) = A \tan(\lambda x)$.

Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x [Y_1'(x)Y_2'(t) - Y_2'(x)Y_1'(t)] f(t) dt,$$

where $Y_1(x), Y_2(x)$ is a fundamental system of solutions of the second-order linear ordinary differential equation $\cos^2(\lambda x)Y''_{xx} + A\lambda Y = 0$, W is the Wronskian, and the primes stand for the differentiation with respect to the argument specified in the parentheses.

As shown in A. D. Polyanin and V. F. Zaitsev (2003), the functions $Y_1(x)$ and $Y_2(x)$ can be expressed via the hypergeometric function.

$$33. \quad y(x) - A \int_a^x \frac{\tan(\lambda x)}{\tan(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\tan(\lambda x)}{\tan(\lambda t)} f(t) dt.$$

$$34. \quad y(x) - A \int_a^x \frac{\tan(\lambda t)}{\tan(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\tan(\lambda t)}{\tan(\lambda x)} f(t) dt.$$

$$35. \quad y(x) - A \int_a^x \tan^k(\lambda x) \tan^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tan^k(\lambda x)$ and $h(t) = \tan^m(\mu t)$.

$$36. \quad y(x) + A \int_a^x t^k \tan^m(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -A \tan^m(\lambda x)$ and $h(t) = t^k$.

$$37. \quad y(x) + A \int_a^x x^k \tan^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ax^k$ and $h(t) = \tan^m(\lambda t)$.

$$38. \quad y(x) - \int_a^x [A \tan(kx) + B - AB(x-t) \tan(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \tan(kx)$.

$$39. \quad y(x) + \int_a^x [A \tan(kt) + B + AB(x-t) \tan(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \tan(kt)$.

2.5-4. Kernels Containing Cotangent.

$$40. \quad y(x) - A \int_a^x \cot(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cot(\lambda x)$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + A \int_a^x \cot(\lambda x) \left| \frac{\sin(\lambda x)}{\sin(\lambda t)} \right|^{A/\lambda} f(t) dt.$$

$$41. \quad y(x) - A \int_a^x \cot(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \cot(\lambda t)$.

Solution:

$$y(x) = f(x) + A \int_a^x \coth(\lambda t) \left| \frac{\sin(\lambda x)}{\sin(\lambda t)} \right|^{A/\lambda} f(t) dt.$$

$$42. \quad y(x) - A \int_a^x \frac{\cot(\lambda x)}{\cot(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\cot(\lambda x)}{\cot(\lambda t)} f(t) dt.$$

$$43. \quad y(x) - A \int_a^x \frac{\cot(\lambda t)}{\cot(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\cot(\lambda t)}{\cot(\lambda x)} f(t) dt.$$

$$44. \quad y(x) + A \int_a^x t^k \cot^m(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -A \cot^m(\lambda x)$ and $h(t) = t^k$.

$$45. \quad y(x) + A \int_a^x x^k \cot^m(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = -Ax^k$ and $h(t) = \cot^m(\lambda t)$.

$$46. \quad y(x) - \int_a^x [A \cot(kx) + B - AB(x-t) \cot(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \cot(kx)$.

$$47. \quad y(x) + \int_a^x [A \cot(kt) + B + AB(x-t) \cot(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \cot(kt)$.

2.5-5. Kernels Containing Combinations of Trigonometric Functions.

$$48. \quad y(x) - A \int_a^x \cos^k(\lambda x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cos^k(\lambda x)$ and $h(t) = \sin^m(\mu t)$.

$$49. \quad y(x) - \int_a^x \{A + B \cos(\lambda x) - B(x-t)[\lambda \sin(\lambda x) + A \cos(\lambda x)]\} y(t) dt = f(x).$$

This is a special case of equation 2.9.38 with $b = B$ and $g(x) = A$.

$$50. \quad y(x) - \int_a^x \{A + B \sin(\lambda x) + B(x-t)[\lambda \cos(\lambda x) - A \sin(\lambda x)]\} y(t) dt = f(x).$$

This is a special case of equation 2.9.39 with $b = B$ and $g(x) = A$.

$$51. \quad y(x) - A \int_a^x \tan^k(\lambda x) \cot^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tan^k(\lambda x)$ and $h(t) = \cot^m(\mu t)$.

2.6. Equations Whose Kernels Contain Inverse Trigonometric Functions

2.6-1. Kernels Containing Arccosine.

$$1. \quad y(x) - A \int_a^x \arccos(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \arccos(\lambda x)$ and $h(t) = 1$.

$$2. \quad y(x) - A \int_a^x \arccos(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \arccos(\lambda t)$.

$$3. \quad y(x) - A \int_a^x \frac{\arccos(\lambda x)}{\arccos(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\arccos(\lambda x)}{\arccos(\lambda t)} f(t) dt.$$

$$4. \quad y(x) - A \int_a^x \frac{\arccos(\lambda t)}{\arccos(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\arccos(\lambda t)}{\arccos(\lambda x)} f(t) dt.$$

$$5. \quad y(x) - \int_a^x [A \arccos(kx) + B - AB(x-t) \arccos(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \arccos(kx)$.

$$6. \quad y(x) + \int_a^x [A \arccos(kt) + B + AB(x-t) \arccos(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \arccos(kt)$.

2.6-2. Kernels Containing Arcsine.

$$7. \quad y(x) - A \int_a^x \arcsin(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \arcsin(\lambda x)$ and $h(t) = 1$.

$$8. \quad y(x) - A \int_a^x \arcsin(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \arcsin(\lambda t)$.

$$9. \quad y(x) - A \int_a^x \frac{\arcsin(\lambda x)}{\arcsin(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\arcsin(\lambda x)}{\arcsin(\lambda t)} f(t) dt.$$

$$10. \quad y(x) - A \int_a^x \frac{\arcsin(\lambda t)}{\arcsin(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\arcsin(\lambda t)}{\arcsin(\lambda x)} f(t) dt.$$

$$11. \quad y(x) - \int_a^x [A \arcsin(kx) + B - AB(x-t) \arcsin(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \arcsin(kx)$.

$$12. \quad y(x) + \int_a^x [A \arcsin(kt) + B + AB(x-t) \arcsin(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \arcsin(kt)$.

2.6-3. Kernels Containing Arctangent.

$$13. \quad y(x) - A \int_a^x \arctan(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \arctan(\lambda x)$ and $h(t) = 1$.

$$14. \quad y(x) - A \int_a^x \arctan(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \arctan(\lambda t)$.

$$15. \quad y(x) - A \int_a^x \frac{\arctan(\lambda x)}{\arctan(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\arctan(\lambda x)}{\arctan(\lambda t)} f(t) dt.$$

$$16. \quad y(x) - A \int_a^x \frac{\arctan(\lambda t)}{\arctan(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\arctan(\lambda t)}{\arctan(\lambda x)} f(t) dt.$$

$$17. \quad y(x) + A \int_x^\infty \arctan[\lambda(t-x)] y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = A \arctan(-\lambda x)$.

$$18. \quad y(x) - \int_a^x [A \arctan(kx) + B - AB(x-t) \arctan(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \arctan(kx)$.

$$19. \quad y(x) + \int_a^x [A \arctan(kt) + B + AB(x-t) \arctan(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \arctan(kt)$.

2.6-4. Kernels Containing Arccotangent.

$$20. \quad y(x) - A \int_a^x \operatorname{arccot}(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \operatorname{arccot}(\lambda x)$ and $h(t) = 1$.

$$21. \quad y(x) - A \int_a^x \operatorname{arccot}(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = \operatorname{arccot}(\lambda t)$.

$$22. \quad y(x) - A \int_a^x \frac{\operatorname{arccot}(\lambda x)}{\operatorname{arccot}(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\operatorname{arccot}(\lambda x)}{\operatorname{arccot}(\lambda t)} f(t) dt.$$

$$23. \quad y(x) - A \int_a^x \frac{\operatorname{arccot}(\lambda t)}{\operatorname{arccot}(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{\operatorname{arccot}(\lambda t)}{\operatorname{arccot}(\lambda x)} f(t) dt.$$

$$24. \quad y(x) + A \int_x^\infty \operatorname{arccot}[\lambda(t-x)] y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = A \operatorname{arccot}(-\lambda x)$.

$$25. \quad y(x) - \int_a^x [A \operatorname{arccot}(kx) + B - AB(x-t) \operatorname{arccot}(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = A \operatorname{arccot}(kx)$.

$$26. \quad y(x) + \int_a^x [A \operatorname{arccot}(kt) + B + AB(x-t) \operatorname{arccot}(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = A \operatorname{arccot}(kt)$.

2.7. Equations Whose Kernels Contain Combinations of Elementary Functions

2.7-1. Kernels Containing Exponential and Hyperbolic Functions.

$$1. \quad y(x) + A \int_a^x e^{\mu(x-t)} \cosh[\lambda(x-t)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_a^x R(x-t) f(t) dt,$$

$$R(x) = \exp\left[\left(\mu - \frac{1}{2}A\right)x\right] \left[\frac{A^2}{2k} \sinh(kx) - A \cosh(kx) \right], \quad k = \sqrt{\lambda^2 + \frac{1}{4}A^2}.$$

$$2. \quad y(x) + A \int_a^x e^{\mu(x-t)} \sinh[\lambda(x-t)] y(t) dt = f(x).$$

1°. Solution with $\lambda(A - \lambda) > 0$:

$$y(x) = f(x) - \frac{A\lambda}{k} \int_a^x e^{\mu(x-t)} \sin[k(x-t)] f(t) dt, \quad \text{where } k = \sqrt{\lambda(A - \lambda)}.$$

2°. Solution with $\lambda(A - \lambda) < 0$:

$$y(x) = f(x) - \frac{A\lambda}{k} \int_a^x e^{\mu(x-t)} \sinh[k(x-t)] f(t) dt, \quad \text{where } k = \sqrt{\lambda(\lambda - A)}.$$

3°. Solution with $A = \lambda$:

$$y(x) = f(x) - \lambda^2 \int_a^x (x-t) e^{\mu(x-t)} f(t) dt.$$

$$3. \quad y(x) + \int_a^x e^{\mu(x-t)} \{A_1 \sinh[\lambda_1(x-t)] + A_2 \sinh[\lambda_2(x-t)]\} y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 2.3.18:

$$w(x) + \int_a^x \{A_1 \sinh[\lambda_1(x-t)] + A_2 \sinh[\lambda_2(x-t)]\} w(t) dt = e^{-\mu x} f(x).$$

$$4. \quad y(x) + A \int_a^x t e^{\mu(x-t)} \sinh[\lambda(x-t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 2.3.23:

$$w(x) + A \int_a^x t \sinh[\lambda(x-t)] w(t) dt = e^{-\mu x} f(x).$$

2.7-2. Kernels Containing Exponential and Logarithmic Functions.

$$5. \quad y(x) - A \int_a^x e^{\mu t} \ln(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln(\lambda x)$ and $h(t) = e^{\mu t}$.

$$6. \quad y(x) - A \int_a^x e^{\mu x} \ln(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A e^{\mu x}$ and $h(t) = \ln(\lambda t)$.

$$7. \quad y(x) - A \int_a^x e^{\mu(x-t)} \ln(\lambda x) y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{(\mu-A)(x-t)} \ln(\lambda x) \frac{(\lambda x)^{Ax}}{(\lambda t)^{At}} f(t) dt.$$

$$8. \quad y(x) - A \int_a^x e^{\mu(x-t)} \ln(\lambda t) y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{(\mu-A)(x-t)} \ln(\lambda t) \frac{(\lambda x)^{Ax}}{(\lambda t)^{At}} f(t) dt.$$

$$9. \quad y(x) + A \int_a^x e^{\mu(x-t)} (\ln x - \ln t) y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x e^{\mu(x-t)} [u_1'(x)u_2'(t) - u_2'(x)u_1'(t)] f(t) dt,$$

where the primes stand for the differentiation with respect to the argument specified in the parentheses, and $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear homogeneous ordinary differential equation $u''_{xx} + Ax^{-1}u = 0$, with $u_1(x)$ and $u_2(x)$ expressed in terms of Bessel functions or modified Bessel functions, depending on the sign of A :

$$\begin{aligned} W = \frac{1}{\pi}, \quad u_1(x) = \sqrt{x} J_1(2\sqrt{Ax}), \quad u_2(x) = \sqrt{x} Y_1(2\sqrt{Ax}) & \quad \text{for } A > 0, \\ W = -\frac{1}{2}, \quad u_1(x) = \sqrt{x} I_1(2\sqrt{-Ax}), \quad u_2(x) = \sqrt{x} K_1(2\sqrt{-Ax}) & \quad \text{for } A < 0. \end{aligned}$$

$$10. \quad y(x) + a \int_x^\infty e^{\lambda(x-t)} \ln(t-x) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = ae^{\lambda x} \ln(-x)$.

2.7-3. Kernels Containing Exponential and Trigonometric Functions.

$$11. \quad y(x) - A \int_a^x e^{\mu t} \cos(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cos(\lambda x)$ and $h(t) = e^{\mu t}$.

$$12. \quad y(x) - A \int_a^x e^{\mu x} \cos(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = Ae^{\mu x}$ and $h(t) = \cos(\lambda t)$.

$$13. \quad y(x) + A \int_a^x e^{\mu(x-t)} \cos[\lambda(x-t)] y(t) dt = f(x).$$

1°. Solution with $|A| > 2|\lambda|$:

$$\begin{aligned} y(x) &= f(x) + \int_a^x R(x-t) f(t) dt, \\ R(x) &= \exp\left[(\mu - \frac{1}{2}A)x\right] \left[\frac{A^2}{2k} \sinh(kx) - A \cosh(kx) \right], \quad k = \sqrt{\frac{1}{4}A^2 - \lambda^2}. \end{aligned}$$

2°. Solution with $|A| < 2|\lambda|$:

$$\begin{aligned} y(x) &= f(x) + \int_a^x R(x-t) f(t) dt, \\ R(x) &= \exp\left[(\mu - \frac{1}{2}A)x\right] \left[\frac{A^2}{2k} \sin(kx) - A \cos(kx) \right], \quad k = \sqrt{\lambda^2 - \frac{1}{4}A^2}. \end{aligned}$$

3°. Solution with $\lambda = \pm \frac{1}{2}A$:

$$y(x) = f(x) + \int_a^x R(x-t) f(t) dt, \quad R(x) = \left(\frac{1}{2}A^2 x - A\right) \exp\left[(\mu - \frac{1}{2}A)x\right].$$

$$14. \quad y(x) - \int_a^x e^{\mu(x-t)} [A \cos(kx) + B - AB(x-t) \cos(kx)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_a^x e^{\mu(x-t)} M(x, t) f(t) dt,$$

$$M(x, t) = [A \cos(kx) + B] \frac{G(x)}{G(t)} + \frac{B^2}{G(t)} \int_t^x e^{B(x-s)} G(s) ds, \quad G(x) = \exp \left[\frac{A}{k} \sin(kx) \right].$$

$$15. \quad y(x) + \int_a^x e^{\mu(x-t)} [A \cos(kt) + B + AB(x-t) \cos(kt)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_a^x e^{\mu(x-t)} M(x, t) f(t) dt,$$

$$M(x, t) = -[A \cos(kt) + B] \frac{G(t)}{G(x)} + \frac{B^2}{G(x)} \int_t^x e^{B(t-s)} G(s) ds, \quad G(x) = \exp \left[\frac{A}{k} \sin(kx) \right].$$

$$16. \quad y(x) - A \int_a^x e^{\mu t} \sin(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sin(\lambda x)$ and $h(t) = e^{\mu t}$.

$$17. \quad y(x) - A \int_a^x e^{\mu x} \sin(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = Ae^{\mu x}$ and $h(t) = \sin(\lambda t)$.

$$18. \quad y(x) + A \int_a^x e^{\mu(x-t)} \sin[\lambda(x-t)] y(t) dt = f(x).$$

1°. Solution with $\lambda(A + \lambda) > 0$:

$$y(x) = f(x) - \frac{A\lambda}{k} \int_a^x e^{\mu(x-t)} \sin[k(x-t)] f(t) dt, \quad \text{where } k = \sqrt{\lambda(A + \lambda)}.$$

2°. Solution with $\lambda(A + \lambda) < 0$:

$$y(x) = f(x) - \frac{A\lambda}{k} \int_a^x e^{\mu(x-t)} \sinh[k(x-t)] f(t) dt, \quad \text{where } k = \sqrt{-\lambda(\lambda + A)}.$$

3°. Solution with $A = -\lambda$:

$$y(x) = f(x) + \lambda^2 \int_a^x (x-t) e^{\mu(x-t)} f(t) dt.$$

$$19. \quad y(x) + A \int_a^x e^{\mu(x-t)} \sin^3[\lambda(x-t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 2.5.17:

$$w(x) + A \int_a^x \sin^3[\lambda(x-t)] w(t) dt = e^{-\mu x} f(x).$$

$$20. \quad y(x) + \int_a^x e^{\mu(x-t)} \{A_1 \sin[\lambda_1(x-t)] + A_2 \sin[\lambda_2(x-t)]\} y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 2.5.18:

$$w(x) + \int_a^x \{A_1 \sin[\lambda_1(x-t)] + A_2 \sin[\lambda_2(x-t)]\} w(t) dt = e^{-\mu x} f(x).$$

$$21. \quad y(x) + \int_a^x e^{\mu(x-t)} \left\{ \sum_{k=1}^n A_k \sin[\lambda_k(x-t)] \right\} y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 2.5.19:

$$w(x) + \int_a^x \left\{ \sum_{k=1}^n A_k \sin[\lambda_k(x-t)] \right\} w(t) dt = e^{-\mu x} f(x).$$

$$22. \quad y(x) + A \int_a^x t e^{\mu(x-t)} \sin[\lambda(x-t)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \frac{A\lambda}{W} \int_a^x t e^{\mu(x-t)} [u_1(x)u_2(t) - u_2(x)u_1(t)] f(t) dt,$$

where $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear ordinary differential equation $u''_{xx} + \lambda(Ax + \lambda)u = 0$, and W is the Wronskian.

Depending on the sign of $A\lambda$, the functions $u_1(x)$ and $u_2(x)$ are expressed in terms of Bessel functions or modified Bessel functions as follows:

if $A\lambda > 0$, then

$$u_1(x) = \xi^{1/2} J_{1/3} \left(\frac{2}{3} \sqrt{A\lambda} \xi^{3/2} \right), \quad u_2(x) = \xi^{1/2} Y_{1/3} \left(\frac{2}{3} \sqrt{A\lambda} \xi^{3/2} \right), \\ W = 3/\pi, \quad \xi = x + (\lambda/A);$$

if $A\lambda < 0$, then

$$u_1(x) = \xi^{1/2} I_{1/3} \left(\frac{2}{3} \sqrt{-A\lambda} \xi^{3/2} \right), \quad u_2(x) = \xi^{1/2} K_{1/3} \left(\frac{2}{3} \sqrt{-A\lambda} \xi^{3/2} \right), \\ W = -\frac{3}{2}, \quad \xi = x + (\lambda/A).$$

$$23. \quad y(x) + A \int_a^x x e^{\mu(x-t)} \sin[\lambda(x-t)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \frac{A\lambda}{W} \int_a^x x e^{\mu(x-t)} [u_1(x)u_2(t) - u_2(x)u_1(t)] f(t) dt,$$

where $u_1(x), u_2(x)$ is a fundamental system of solutions of the second-order linear ordinary differential equation $u''_{xx} + \lambda(Ax + \lambda)u = 0$, and W is the Wronskian.

The functions $u_1(x), u_2(x)$, and W are specified in 2.7.22.

$$24. \quad y(x) + A \int_x^\infty e^{\mu(t-x)} \sin(\lambda\sqrt{t-x}) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = Ae^{-\mu x} \sin(\lambda\sqrt{-x})$.

$$25. \quad y(x) - \int_a^x e^{\mu(x-t)} [A \sin(kx) + B - AB(x-t) \sin(kx)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_a^x e^{\mu(x-t)} M(x, t) f(t) dt,$$

$$M(x, t) = [A \sin(kx) + B] \frac{G(x)}{G(t)} + \frac{B^2}{G(t)} \int_t^x e^{B(x-s)} G(s) ds, \quad G(x) = \exp \left[-\frac{A}{k} \cos(kx) \right].$$

$$26. \quad y(x) + \int_a^x e^{\mu(x-t)} [A \sin(kt) + B + AB(x-t) \sin(kt)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_a^x e^{\mu(x-t)} M(x, t) f(t) dt,$$

$$M(x, t) = -[A \sin(kt) + B] \frac{G(t)}{G(x)} + \frac{B^2}{G(x)} \int_t^x e^{B(t-s)} G(s) ds, \quad G(x) = \exp \left[-\frac{A}{k} \cos(kx) \right].$$

$$27. \quad y(x) - A \int_a^x e^{\mu t} \tan(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tan(\lambda x)$ and $h(t) = e^{\mu t}$.

$$28. \quad y(x) - A \int_a^x e^{\mu x} \tan(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = Ae^{\mu x}$ and $h(t) = \tan(\lambda t)$.

$$29. \quad y(x) + A \int_a^x e^{\mu(x-t)} [\tan(\lambda x) - \tan(\lambda t)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 2.5.32:

$$w(x) + A \int_a^x [\tan(\lambda x) - \tan(\lambda t)] w(t) dt = e^{-\mu x} f(x).$$

$$30. \quad y(x) - \int_a^x e^{\mu(x-t)} [A \tan(kx) + B - AB(x-t) \tan(kx)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 2.9.7 with $\lambda = B$ and $g(x) = A \tan(kx)$:

$$w(x) - \int_a^x [A \tan(kx) + B - AB(x-t) \tan(kx)] w(t) dt = e^{-\mu x} f(x).$$

$$31. \quad y(x) + \int_a^x e^{\mu(x-t)} [A \tan(kt) + B + AB(x-t) \tan(kt)] y(t) dt = f(x).$$

The substitution $w(x) = e^{-\mu x} y(x)$ leads to an equation of the form 2.9.8 with $\lambda = B$ and $g(t) = A \tan(kt)$:

$$w(x) + \int_a^x [A \tan(kt) + B + AB(x-t) \tan(kt)] w(t) dt = e^{-\mu x} f(x).$$

$$32. \quad y(x) - A \int_a^x e^{\mu t} \cot(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cot(\lambda x)$ and $h(t) = e^{\mu t}$.

$$33. \quad y(x) - A \int_a^x e^{\mu x} \cot(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = Ae^{\mu x}$ and $h(t) = \cot(\lambda t)$.

2.7-4. Kernels Containing Hyperbolic and Logarithmic Functions.

$$34. \quad y(x) - A \int_a^x \cosh^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cosh^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$35. \quad y(x) - A \int_a^x \cosh^k(\lambda t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\mu x)$ and $h(t) = \cosh^k(\lambda t)$.

$$36. \quad y(x) - A \int_a^x \sinh^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sinh^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$37. \quad y(x) - A \int_a^x \sinh^k(\lambda t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\mu x)$ and $h(t) = \sinh^k(\lambda t)$.

$$38. \quad y(x) - A \int_a^x \tanh^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tanh^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$39. \quad y(x) - A \int_a^x \tanh^k(\lambda t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\mu x)$ and $h(t) = \tanh^k(\lambda t)$.

$$40. \quad y(x) - A \int_a^x \coth^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \coth^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$41. \quad y(x) - A \int_a^x \coth^k(\lambda t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\mu x)$ and $h(t) = \coth^k(\lambda t)$.

2.7-5. Kernels Containing Hyperbolic and Trigonometric Functions.

$$42. \quad y(x) - A \int_a^x \cosh^k(\lambda x) \cos^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cosh^k(\lambda x)$ and $h(t) = \cos^m(\mu t)$.

$$43. \quad y(x) - A \int_a^x \cosh^k(\lambda t) \cos^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cos^m(\mu x)$ and $h(t) = \cosh^k(\lambda t)$.

$$44. \quad y(x) - A \int_a^x \cosh^k(\lambda x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cosh^k(\lambda x)$ and $h(t) = \sin^m(\mu t)$.

$$45. \quad y(x) - A \int_a^x \cosh^k(\lambda t) \sin^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sin^m(\mu x)$ and $h(t) = \cosh^k(\lambda t)$.

$$46. \quad y(x) - A \int_a^x \sinh^k(\lambda x) \cos^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sinh^k(\lambda x)$ and $h(t) = \cos^m(\mu t)$.

$$47. \quad y(x) - A \int_a^x \sinh^k(\lambda t) \cos^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cos^m(\mu x)$ and $h(t) = \sinh^k(\lambda t)$.

$$48. \quad y(x) - A \int_a^x \sinh^k(\lambda x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sinh^k(\lambda x)$ and $h(t) = \sin^m(\mu t)$.

$$49. \quad y(x) - A \int_a^x \sinh^k(\lambda t) \sin^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sin^m(\mu x)$ and $h(t) = \sinh^k(\lambda t)$.

$$50. \quad y(x) - A \int_a^x \tanh^k(\lambda x) \cos^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tanh^k(\lambda x)$ and $h(t) = \cos^m(\mu t)$.

$$51. \quad y(x) - A \int_a^x \tanh^k(\lambda t) \cos^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cos^m(\mu x)$ and $h(t) = \tanh^k(\lambda t)$.

$$52. \quad y(x) - A \int_a^x \tanh^k(\lambda x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tanh^k(\lambda x)$ and $h(t) = \sin^m(\mu t)$.

$$53. \quad y(x) - A \int_a^x \tanh^k(\lambda t) \sin^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sin^m(\mu x)$ and $h(t) = \tanh^k(\lambda t)$.

2.7-6. Kernels Containing Logarithmic and Trigonometric Functions.

$$54. \quad y(x) - A \int_a^x \cos^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cos^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$55. \quad y(x) - A \int_a^x \cos^k(\lambda t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\mu x)$ and $h(t) = \cos^k(\lambda t)$.

$$56. \quad y(x) - A \int_a^x \sin^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \sin^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$57. \quad y(x) - A \int_a^x \sin^k(\lambda t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\mu x)$ and $h(t) = \sin^k(\lambda t)$.

$$58. \quad y(x) - A \int_a^x \tan^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \tan^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$59. \quad y(x) - A \int_a^x \tan^k(\lambda t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\mu x)$ and $h(t) = \tan^k(\lambda t)$.

$$60. \quad y(x) - A \int_a^x \cot^k(\lambda x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \cot^k(\lambda x)$ and $h(t) = \ln^m(\mu t)$.

$$61. \quad y(x) - A \int_a^x \cot^k(\lambda t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A \ln^m(\mu x)$ and $h(t) = \cot^k(\lambda t)$.

2.8. Equations Whose Kernels Contain Special Functions

2.8-1. Kernels Containing Bessel Functions.

$$1. \quad y(x) - \lambda \int_0^x J_0(x-t) y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_0^x R(x-t) f(t) dt,$$

where

$$R(x) = \lambda \cos(\sqrt{1-\lambda^2} x) + \frac{\lambda^2}{\sqrt{1-\lambda^2}} \sin(\sqrt{1-\lambda^2} x) + \frac{\lambda}{\sqrt{1-\lambda^2}} \int_0^x \sin[\sqrt{1-\lambda^2}(x-t)] \frac{J_1(t)}{t} dt.$$

⊙ Reference: V. I. Smirnov (1974).

$$2. \quad y(x) - A \int_a^x J_\nu(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = AJ_\nu(\lambda x)$ and $h(t) = 1$.

$$3. \quad y(x) - A \int_a^x J_\nu(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = J_\nu(\lambda t)$.

$$4. \quad y(x) - A \int_a^x \frac{J_\nu(\lambda x)}{J_\nu(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{J_\nu(\lambda x)}{J_\nu(\lambda t)} f(t) dt.$$

$$5. \quad y(x) - A \int_a^x \frac{J_\nu(\lambda t)}{J_\nu(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{J_\nu(\lambda t)}{J_\nu(\lambda x)} f(t) dt.$$

$$6. \quad y(x) + A \int_x^\infty J_\nu(\lambda(t-x)) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = AJ_\nu(-\lambda x)$.

$$7. \quad y(x) - \int_a^x [AJ_\nu(kx) + B - AB(x-t)J_\nu(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = AJ_\nu(kx)$.

$$8. \quad y(x) + \int_a^x [AJ_\nu(kt) + B + AB(x-t)J_\nu(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = AJ_\nu(kt)$.

$$9. \quad y(x) - \lambda \int_0^x e^{\mu(x-t)} J_0(x-t) y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_0^x R(x-t) f(t) dt,$$

where

$$R(x) = e^{\mu x} \left\{ \lambda \cos(\sqrt{1-\lambda^2} x) + \frac{\lambda^2}{\sqrt{1-\lambda^2}} \sin(\sqrt{1-\lambda^2} x) + \frac{\lambda}{\sqrt{1-\lambda^2}} \int_0^x \sin[\sqrt{1-\lambda^2}(x-t)] \frac{J_1(t)}{t} dt \right\}.$$

$$10. \quad y(x) - A \int_a^x Y_\nu(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = AY_\nu(\lambda x)$ and $h(t) = 1$.

$$11. \quad y(x) - A \int_a^x Y_\nu(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = Y_\nu(\lambda t)$.

$$12. \quad y(x) - A \int_a^x \frac{Y_\nu(\lambda x)}{Y_\nu(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{Y_\nu(\lambda x)}{Y_\nu(\lambda t)} f(t) dt.$$

$$13. \quad y(x) - A \int_a^x \frac{Y_\nu(\lambda t)}{Y_\nu(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{Y_\nu(\lambda t)}{Y_\nu(\lambda x)} f(t) dt.$$

$$14. \quad y(x) + A \int_x^\infty Y_\nu(\lambda(t-x)) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = AY_\nu(-\lambda x)$.

$$15. \quad y(x) - \int_a^x [AY_\nu(kx) + B - AB(x-t)Y_\nu(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = AY_\nu(kx)$.

$$16. \quad y(x) + \int_a^x [AY_\nu(kt) + B + AB(x-t)Y_\nu(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = AY_\nu(kt)$.

2.8-2. Kernels Containing Modified Bessel Functions.

$$17. \quad y(x) - A \int_a^x I_\nu(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = AI_\nu(\lambda x)$ and $h(t) = 1$.

$$18. \quad y(x) - A \int_a^x I_\nu(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = I_\nu(\lambda t)$.

$$19. \quad y(x) - A \int_a^x \frac{I_\nu(\lambda x)}{I_\nu(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{I_\nu(\lambda x)}{I_\nu(\lambda t)} f(t) dt.$$

$$20. \quad y(x) - A \int_a^x \frac{I_\nu(\lambda t)}{I_\nu(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{I_\nu(\lambda t)}{I_\nu(\lambda x)} f(t) dt.$$

$$21. \quad y(x) + A \int_x^\infty I_\nu(\lambda(t-x)) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = AI_\nu(-\lambda x)$.

$$22. \quad y(x) - \int_a^x [AI_\nu(kx) + B - AB(x-t)I_\nu(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = AI_\nu(kx)$.

$$23. \quad y(x) + \int_a^x [AI_\nu(kt) + B + AB(x-t)I_\nu(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = AI_\nu(kt)$.

$$24. \quad y(x) - A \int_a^x K_\nu(\lambda x) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = AK_\nu(\lambda x)$ and $h(t) = 1$.

$$25. \quad y(x) - A \int_a^x K_\nu(\lambda t) y(t) dt = f(x).$$

This is a special case of equation 2.9.2 with $g(x) = A$ and $h(t) = K_\nu(\lambda t)$.

$$26. \quad y(x) - A \int_a^x \frac{K_\nu(\lambda x)}{K_\nu(\lambda t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{K_\nu(\lambda x)}{K_\nu(\lambda t)} f(t) dt.$$

$$27. \quad y(x) - A \int_a^x \frac{K_\nu(\lambda t)}{K_\nu(\lambda x)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + A \int_a^x e^{A(x-t)} \frac{K_\nu(\lambda t)}{K_\nu(\lambda x)} f(t) dt.$$

$$28. \quad y(x) + A \int_x^\infty K_\nu(\lambda(t-x)) y(t) dt = f(x).$$

This is a special case of equation 2.9.62 with $K(x) = AK_\nu(-\lambda x)$.

$$29. \quad y(x) - \int_a^x [AK_\nu(kx) + B - AB(x-t)K_\nu(kx)] y(t) dt = f(x).$$

This is a special case of equation 2.9.7 with $\lambda = B$ and $g(x) = AK_\nu(kx)$.

$$30. \quad y(x) + \int_a^x [AK_\nu(kt) + B + AB(x-t)K_\nu(kt)] y(t) dt = f(x).$$

This is a special case of equation 2.9.8 with $\lambda = B$ and $g(t) = AK_\nu(kt)$.

2.9. Equations Whose Kernels Contain Arbitrary Functions

2.9-1. Equations with Degenerate Kernel: $K(x, t) = g_1(x)h_1(t) + \dots + g_n(x)h_n(t)$.

1.
$$y(x) - \lambda \int_a^x \frac{g(x)}{g(t)} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \lambda \int_a^x e^{\lambda(x-t)} \frac{g(x)}{g(t)} f(t) dt.$$

2.
$$y(x) - \int_a^x g(x)h(t)y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_a^x R(x, t)f(t) dt, \quad \text{where } R(x, t) = g(x)h(t) \exp \left[\int_t^x g(s)h(s) ds \right].$$

3.
$$y(x) + \int_a^x (x-t)g(x)y(t) dt = f(x).$$

This is a special case of equation 2.9.11.

1°. Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x [Y_1(x)Y_2(t) - Y_2(x)Y_1(t)] g(x)f(t) dt, \quad (1)$$

where $Y_1 = Y_1(x)$ and $Y_2 = Y_2(x)$ are two linearly independent solutions ($Y_1/Y_2 \neq \text{const}$) of the second-order linear homogeneous differential equation $Y''_{xx} + g(x)Y = 0$. In this case, the Wronskian is a constant: $W = Y_1(Y_2)'_x - Y_2(Y_1)'_x \equiv \text{const}$.

2°. Given only one nontrivial solution $Y_1 = Y_1(x)$ of the linear homogeneous differential equation $Y''_{xx} + g(x)Y = 0$, one can obtain the solution of the integral equation by formula (1) with

$$W = 1, \quad Y_2(x) = Y_1(x) \int_b^x \frac{d\xi}{Y_1^2(\xi)},$$

where b is an arbitrary number.

4.
$$y(x) + \int_a^x (x-t)g(t)y(t) dt = f(x).$$

This is a special case of equation 2.9.12.

1°. Solution:

$$y(x) = f(x) + \frac{1}{W} \int_a^x [Y_1(x)Y_2(t) - Y_2(x)Y_1(t)] g(t)f(t) dt, \quad (1)$$

where $Y_1 = Y_1(x)$ and $Y_2 = Y_2(x)$ are two linearly independent solutions ($Y_1/Y_2 \neq \text{const}$) of the second-order linear homogeneous differential equation $Y''_{xx} + g(x)Y = 0$. In this case, the Wronskian is a constant: $W = Y_1(Y_2)'_x - Y_2(Y_1)'_x \equiv \text{const}$.

2°. Given only one nontrivial solution $Y_1 = Y_1(x)$ of the linear homogeneous differential equation $Y''_{xx} + g(x)Y = 0$, one can obtain the solution of the integral equation by formula (1) with

$$W = 1, \quad Y_2(x) = Y_1(x) \int_b^x \frac{d\xi}{Y_1^2(\xi)},$$

where b is an arbitrary number.

$$5. \quad y(x) + \int_a^x [g(x) - g(t)] y(t) dt = f(x).$$

1°. Differentiating the equation with respect to x yields

$$y'_x(x) + g'_x(x) \int_a^x y(t) dt = f'_x(x). \quad (1)$$

Introducing the new variable $Y(x) = \int_a^x y(t) dt$, we obtain the second-order linear ordinary differential equation

$$Y''_{xx} + g'_x(x)Y = f'_x(x), \quad (2)$$

which must be supplemented by the initial conditions

$$Y(a) = 0, \quad Y'_x(a) = f(a). \quad (3)$$

Conditions (3) follow from the original equation and the definition of $Y(x)$.

For exact solutions of second-order linear ordinary differential equations (2) with various $f(x)$, see E. Kamke (1977), G. M. Murphy (1960), and A. D. Polyanin and V. F. Zaitsev (2003).

2°. Let $Y_1 = Y_1(x)$ and $Y_2 = Y_2(x)$ be two linearly independent solutions ($Y_1/Y_2 \neq \text{const}$) of the second-order linear homogeneous differential equation $Y''_{xx} + g'_x(x)Y = 0$, which follows from (2) for $f(x) \equiv 0$. In this case, the Wronskian is a constant:

$$W = Y_1(Y_2)'_x - Y_2(Y_1)'_x \equiv \text{const}.$$

Solving the nonhomogeneous equation (2) under the initial conditions (3) with arbitrary $f = f(x)$ and taking into account $y(x) = Y'_x(x)$, we obtain the solution of the original integral equation in the form

$$y(x) = f(x) + \frac{1}{W} \int_a^x [Y'_1(x)Y'_2(t) - Y'_2(x)Y'_1(t)] f(t) dt, \quad (4)$$

where the primes stand for the differentiation with respect to the argument specified in the parentheses.

3°. Given only one nontrivial solution $Y_1 = Y_1(x)$ of the linear homogeneous differential equation $Y''_{xx} + g'_x(x)Y = 0$, one can obtain the solution of the nonhomogeneous equation (2) under the initial conditions (3) by formula (4) with

$$W = 1, \quad Y_2(x) = Y_1(x) \int_b^x \frac{d\xi}{Y_1^2(\xi)},$$

where b is an arbitrary number.

$$6. \quad y(x) + \int_a^x [g(x) + h(t)] y(t) dt = f(x).$$

1°. Differentiating the equation with respect to x yields

$$y'_x(x) + [g(x) + h(x)]y(x) + g'_x(x) \int_a^x y(t) dt = f'_x(x).$$

Introducing the new variable $Y(x) = \int_a^x y(t) dt$, we obtain the second-order linear ordinary differential equation

$$Y''_{xx} + [g(x) + h(x)]Y'_x + g'_x(x)Y = f'_x(x), \quad (1)$$

which must be supplemented by the initial conditions

$$Y(a) = 0, \quad Y'_x(a) = f(a). \quad (2)$$

Conditions (3) follow from the original equation and the definition of $Y(x)$.

For exact solutions of second-order linear ordinary differential equations (1) with various $f(x)$, see E. Kamke (1977), G. M. Murphy (1960), and A. D. Polyanin and V. F. Zaitsev (2003).

2°. Let $Y_1 = Y_1(x)$ and $Y_2 = Y_2(x)$ be two linearly independent solutions ($Y_1/Y_2 \neq \text{const}$) of the second-order linear homogeneous differential equation $Y''_{xx} + [g(x) + h(x)]Y'_x + g'_x(x)Y = 0$, which follows from (1) for $f(x) \equiv 0$.

Solving the nonhomogeneous equation (1) under the initial conditions (2) with arbitrary $f = f(x)$ and taking into account $y(x) = Y'_x(x)$, we obtain the solution of the original integral equation in the form

$$y(x) = f(x) + \int_a^x R(x, t)f(t) dt,$$

$$R(x, t) = \frac{\partial^2}{\partial x \partial t} \left[\frac{Y_1(x)Y_2(t) - Y_2(x)Y_1(t)}{W(t)} \right], \quad W(x) = Y_1(x)Y'_2(x) - Y_2(x)Y'_1(x),$$

where $W(x)$ is the Wronskian and the primes stand for the differentiation with respect to the argument specified in the parentheses.

7. $y(x) - \int_a^x [g(x) + \lambda - \lambda(x-t)g(x)] y(t) dt = f(x).$

This is a special case of equation 2.9.16 with $h(x) = \lambda$.

Solution:

$$y(x) = f(x) + \int_a^x R(x, t)f(t) dt,$$

$$R(x, t) = [g(x) + \lambda] \frac{G(x)}{G(t)} + \frac{\lambda^2}{G(t)} \int_t^x e^{\lambda(x-s)} G(s) ds, \quad G(x) = \exp \left[\int_a^x g(s) ds \right].$$

8. $y(x) + \int_a^x [g(t) + \lambda + \lambda(x-t)g(t)] y(t) dt = f(x).$

Solution:

$$y(x) = f(x) + \int_a^x R(x, t)f(t) dt,$$

$$R(x, t) = -[g(t) + \lambda] \frac{G(t)}{G(x)} + \frac{\lambda^2}{G(x)} \int_t^x e^{\lambda(t-s)} G(s) ds, \quad G(x) = \exp \left[\int_a^x g(s) ds \right].$$

9. $y(x) - \int_a^x [g_1(x) + g_2(x)t] y(t) dt = f(x).$

This equation can be rewritten in the form of equation 2.9.11 with $g_1(x) = g(x) + xh(x)$ and $g_2(x) = -h(x)$.

$$10. \quad y(x) - \int_a^x [g_1(t) + g_2(t)x] y(t) dt = f(x).$$

This equation can be rewritten in the form of equation 2.9.12 with $g_1(t) = g(t) + th(t)$ and $g_2(t) = -h(t)$.

$$11. \quad y(x) - \int_a^x [g(x) + h(x)(x-t)] y(t) dt = f(x).$$

1°. The solution of the integral equation can be represented in the form $y(x) = Y''_{xx}$, where $Y = Y(x)$ is the solution of the second-order linear nonhomogeneous ordinary differential equation

$$Y''_{xx} - g(x)Y'_x - h(x)Y = f(x), \quad (1)$$

under the initial conditions

$$Y(a) = Y'_x(a) = 0. \quad (2)$$

2°. Let $Y_1 = Y_1(x)$ and $Y_2 = Y_2(x)$ be two nontrivial linearly independent solutions of the second-order linear homogeneous differential equation $Y''_{xx} - g(x)Y'_x - h(x)Y = 0$, which follows from (1) for $f(x) \equiv 0$. Then the solution of the nonhomogeneous differential equation (1) under conditions (2) is given by

$$Y(x) = \int_a^x [Y_2(x)Y_1(t) - Y_1(x)Y_2(t)] \frac{f(t)}{W(t)} dt, \quad W(t) = Y_1(t)Y'_2(t) - Y_2(t)Y'_1(t), \quad (3)$$

where $W(t)$ is the Wronskian and the primes denote the derivatives.

Substituting (3) into (1), we obtain the solution of the original integral equation in the form

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt, \quad R(x,t) = \frac{1}{W(t)} [Y''_2(x)Y_1(t) - Y''_1(x)Y_2(t)]. \quad (4)$$

3°. Let $Y_1 = Y_1(x)$ be a nontrivial particular solution of the homogeneous differential equation (1) (with $f \equiv 0$) satisfying the initial condition $Y_1(a) \neq 0$. Then the function

$$Y_2(x) = Y_1(x) \int_a^x \frac{W(t)}{[Y_1(t)]^2} dt, \quad W(x) = \exp \left[\int_a^x g(s) ds \right] \quad (5)$$

is another nontrivial solution of the homogeneous equation. Substituting (5) into (4) yields the solution of the original integral equation in the form

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = g(x) \frac{W(x)}{Y_1(x)} \frac{Y_1(t)}{W(t)} + [g(x)Y'_1(x) + h(x)Y_1(x)] \frac{Y_1(t)}{W(t)} \int_t^x \frac{W(s)}{[Y_1(s)]^2} ds,$$

where $W(x) = \exp \left[\int_a^x g(s) ds \right]$.

$$12. \quad y(x) - \int_a^x [g(t) + h(t)(t-x)] y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = g(t) \frac{Y(x)W(x)}{Y(t)W(t)} + Y(x)W(x)[g(t)Y'_t(t) + h(t)Y(t)] \int_x^t \frac{ds}{W(s)[Y(s)]^2},$$

$$W(t) = \exp \left[\int_b^t g(t) dt \right],$$

where $Y = Y(x)$ is an arbitrary nontrivial solution of the second-order homogeneous differential equation

$$Y''_{xx} + g(x)Y'_x + h(x)Y = 0$$

satisfying the condition $Y(a) \neq 0$.

13. $y(x) + \int_a^x (x-t)g(x)h(t)y(t) dt = f(x).$

The substitution $y(x) = g(x)u(x)$ leads to an equation of the form 2.9.4:

$$u(x) + \int_a^x (x-t)g(t)h(t)u(t) dt = f(x)/g(x).$$

14. $y(x) - \int_a^x \{g(x) + \lambda x^n + \lambda(x-t)x^{n-1}[n - xg(x)]\}y(t) dt = f(x).$

This is a special case of equation 2.9.16 with $h(x) = \lambda x^n$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = [g(x) + \lambda x^n] \frac{G(x)}{G(t)} + \lambda(\lambda x^{2n} + nx^{n-1}) \frac{H(x)}{G(t)} \int_t^x \frac{G(s)}{H(s)} ds,$$

where $G(x) = \exp \left[\int_a^x g(s) ds \right]$ and $H(x) = \exp \left(\frac{\lambda}{n+1} x^{n+1} \right)$.

15. $y(x) - \int_a^x \{g(x) + \lambda + (x-t)[g'_x(x) - \lambda g(x)]\}y(t) dt = f(x).$

This is a special case of equation 2.9.16.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = [g(x) + \lambda]e^{\lambda(x-t)} + \{[g(x)]^2 + g'_x(x)\}G(x) \int_t^x \frac{e^{\lambda(s-t)}}{G(s)} ds,$$

where $G(x) = \exp \left[\int_a^x g(s) ds \right]$.

16. $y(x) - \int_a^x \{g(x) + h(x) + (x-t)[h'_x(x) - g(x)h(x)]\}y(t) dt = f(x).$

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = [g(x) + h(x)] \frac{G(x)}{G(t)} + \{[h(x)]^2 + h'_x(x)\} \frac{H(x)}{G(t)} \int_t^x \frac{G(s)}{H(s)} ds,$$

where $G(x) = \exp \left[\int_a^x g(s) ds \right]$ and $H(x) = \exp \left[\int_a^x h(s) ds \right]$.

$$17. \quad y(x) + \int_a^x \left\{ \frac{\varphi'_x(x)}{\varphi(t)} + [\varphi(t)g'_x(x) - \varphi'_x(x)g(t)]h(t) \right\} y(t) dt = f(x).$$

1°. This equation is equivalent to the equation

$$\int_a^x \left\{ \frac{\varphi(x)}{\varphi(t)} + [\varphi(t)g(x) - \varphi(x)g(t)]h(t) \right\} y(t) dt = F(x), \quad F(x) = \int_a^x f(x) dx, \quad (1)$$

obtained by differentiating the original equation with respect to x . Equation (1) is a special case of equation 1.9.15 with

$$g_1(x) = g(x), \quad h_1(t) = \varphi(t)h(t), \quad g_2(x) = \varphi(x), \quad h_2(t) = \frac{1}{\varphi(t)} - g(t)h(t).$$

2°. Solution:

$$y(x) = \frac{1}{\varphi(x)h(x)} \frac{d}{dx} \left\{ \Xi(x) \int_a^x \left[\frac{F(t)}{\varphi(t)} \right]'_t \frac{\varphi^2(t)h(t)}{\Xi(t)} dt \right\},$$

$$F(x) = \int_a^x f(x) dx, \quad \Xi(x) = \exp \left\{ - \int_a^x \left[\frac{g(t)}{\varphi(t)} \right]'_t \varphi^2(t)h(t) dt \right\}.$$

$$18. \quad y(x) - \int_a^x \left\{ \frac{\varphi'_t(t)}{\varphi(x)} + [\varphi(x)g'_t(t) - \varphi'_t(t)g(x)]h(x) \right\} y(t) dt = f(x).$$

1°. Let $f(a) = 0$. The change

$$y(x) = \int_a^x w(t) dt \quad (1)$$

followed by the integration by parts leads to the equation

$$\int_a^x \left\{ \frac{\varphi(t)}{\varphi(x)} + [\varphi(x)g(t) - \varphi(t)g(x)]h(x) \right\} w(t) dt = f(x), \quad (2)$$

which is a special case of equation 1.9.15 with

$$g_1(x) = \frac{1}{\varphi(x)} - g(x)h(x), \quad h_1(t) = \varphi(t), \quad g_2(x) = \varphi(x)h(x), \quad h_2(t) = g(t).$$

The solution of equation (2) is given by

$$y(x) = \frac{1}{\varphi(x)} \frac{d}{dx} \left\{ \varphi^2(x)h(x)\Phi(x) \int_a^x \left[\frac{f(t)}{\varphi(t)h(t)} \right]'_t \frac{dt}{\Phi(t)} \right\},$$

$$\Phi(x) = \exp \left\{ \int_a^x \left[\frac{g(t)}{\varphi(t)} \right]'_t \varphi^2(t)h(t) dt \right\}.$$

2°. Let $f(a) \neq 0$. The substitution $y(x) = \bar{y}(x) + f(a)$ leads to the integral equation $\bar{y}(x)$ with the right-hand side $\bar{f}(x)$ satisfying the condition $\bar{f}(a) = 0$. Thus we obtain case 1°.

$$19. \quad y(x) - \int_a^x \left[\sum_{k=1}^n g_k(x)(x-t)^{k-1} \right] y(t) dt = f(x).$$

The solution can be represented in the form

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt. \quad (1)$$

Here the resolvent $R(x, t)$ is given by

$$R(x, t) = w_x^{(n)}, \quad w_x^{(n)} = \frac{d^n w}{dx^n}, \quad (2)$$

where w is the solution of the n th-order linear homogeneous ordinary differential equation

$$w_x^{(n)} - g_1(x)w_x^{(n-1)} - g_2(x)w_x^{(n-2)} - 2g_3(x)w_x^{(n-3)} - \dots - (n-1)!g_n(x)w = 0 \quad (3)$$

satisfying the following initial conditions at $x = t$:

$$w|_{x=t} = w'_x|_{x=t} = \dots = w_x^{(n-2)}|_{x=t} = 0, \quad w_x^{(n-1)}|_{x=t} = 1. \quad (4)$$

Note that the differential equation (3) implicitly depends on t via the initial conditions (4).

⊙ References: E. Goursat (1923), A. F. Verlan' and V. S. Sizikov (1987).

$$20. \quad y(x) - \int_a^x \left[\sum_{k=1}^n g_k(t)(t-x)^{k-1} \right] y(t) dt = f(x).$$

The solution can be represented in the form

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt. \quad (1)$$

Here the resolvent $R(x, t)$ is given by

$$R(x, t) = -u_t^{(n)}, \quad u_t^{(n)} = \frac{d^n u}{dt^n}, \quad (2)$$

where u is the solution of the n th-order linear homogeneous ordinary differential equation

$$u_t^{(n)} + g_1(t)u_t^{(n-1)} + g_2(t)u_t^{(n-2)} + 2g_3(t)u_t^{(n-3)} + \dots + (n-1)!g_n(t)u = 0, \quad (3)$$

satisfying the following initial conditions at $t = x$:

$$u|_{t=x} = u'_t|_{t=x} = \dots = u_t^{(n-2)}|_{t=x} = 0, \quad u_t^{(n-1)}|_{t=x} = 1. \quad (4)$$

Note that the differential equation (3) implicitly depends on x via the initial conditions (4).

⊙ References: E. Goursat (1923), A. F. Verlan' and V. S. Sizikov (1987).

$$21. \quad y(x) + \int_a^x (e^{\lambda x + \mu t} - e^{\mu x + \lambda t})g(t)y(t) dt = f(x).$$

Let us differentiate the equation twice and then eliminate the integral terms from the resulting relations and the original equation. As a result, we arrive at the second-order linear ordinary differential equation

$$y''_{xx} - (\lambda + \mu)y'_x + [(\lambda - \mu)e^{(\lambda + \mu)x}g(x) + \lambda\mu]y = f''_{xx}(x) - (\lambda + \mu)f'_x(x) + \lambda\mu f(x),$$

which must be supplemented by the initial conditions $y(a) = f(a)$, $y'_x(a) = f'_x(a)$.

$$22. \quad y(x) + \int_a^x [e^{\lambda x} g(t) + e^{\mu x} h(t)] y(t) dt = f(x).$$

Let us differentiate the equation twice and then eliminate the integral terms from the resulting relations and the original equation. As a result, we arrive at the second-order linear ordinary differential equation

$$y''_{xx} + [e^{\lambda x} g(x) + e^{\mu x} h(x) - \lambda - \mu] y'_x + [e^{\lambda x} g'_x(x) + e^{\mu x} h'_x(x) + (\lambda - \mu)e^{\lambda x} g(x) + (\mu - \lambda)e^{\mu x} h(x) + \lambda\mu] y = f''_{xx}(x) - (\lambda + \mu)f'_x(x) + \lambda\mu f(x),$$

which must be supplemented by the initial conditions

$$y(a) = f(a), \quad y'_x(a) = f'_x(a) - [e^{\lambda a} g(a) + e^{\mu a} h(a)] f(a).$$

Example. The Arutyunyan equation,

$$y(x) - \int_a^x \varphi(t) \frac{\partial}{\partial t} \left\{ \frac{1}{\varphi(t)} + \psi(t) [1 - e^{-\lambda(x-t)}] \right\} y(t) dt = f(x),$$

can be reduced to the above equation. The former is encountered in the theory of viscoelasticity for aging solids. The solution of the Arutyunyan equation is given by

$$y(x) = f(x) - \int_a^x \frac{1}{\varphi(t)} \frac{\partial}{\partial t} \left[\varphi(t) - \lambda \psi(t) \varphi^2(t) e^{\eta(t)} \int_t^x e^{-\eta(s)} ds \right] f(t) dt,$$

where

$$\eta(x) = \int_a^x \left\{ \lambda [1 + \psi(t) \varphi(t)] - \frac{\varphi'(t)}{\varphi(t)} \right\} dt.$$

⊙ Reference: N. Kh. Arutyunyan (1966).

$$23. \quad y(x) + \int_a^x [\lambda e^{\lambda(x-t)} + (\mu e^{\mu x + \lambda t} - \lambda e^{\lambda x + \mu t}) h(t)] y(t) dt = f(x).$$

This is a special case of equation 2.9.17 with $\varphi(x) = e^{\lambda x}$ and $g(x) = e^{\mu x}$.

Solution:

$$y(x) = \frac{1}{e^{\lambda x} h(x)} \frac{d}{dx} \left\{ \Phi(x) \int_a^x \left[\frac{F(t)}{e^{\lambda t}} \right]'_t \frac{e^{2\lambda t} h(t)}{\Phi(t)} dt \right\},$$

$$F(x) = \int_a^x f(t) dt, \quad \Phi(x) = \exp \left[(\lambda - \mu) \int_a^x e^{(\lambda + \mu)t} h(t) dt \right].$$

$$24. \quad y(x) - \int_a^x [\lambda e^{-\lambda(x-t)} + (\mu e^{\lambda x + \mu t} - \lambda e^{\mu x + \lambda t}) h(x)] y(t) dt = f(x).$$

This is a special case of equation 2.9.18 with $\varphi(x) = e^{\lambda x}$ and $g(x) = e^{\mu x}$.

Assume that $f(a) = 0$. Solution:

$$y(x) = \int_a^x w(t) dt, \quad w(x) = e^{-\lambda x} \frac{d}{dx} \left\{ \frac{e^{2\lambda x} h(x)}{\Phi(x)} \int_a^x \left[\frac{f(t)}{e^{\lambda t} h(t)} \right]'_t \Phi(t) dt \right\},$$

$$\Phi(x) = \exp \left[(\lambda - \mu) \int_a^x e^{(\lambda + \mu)t} h(t) dt \right].$$

$$25. \quad y(x) - \int_a^x \{g(x) + be^{\lambda x} + b(x-t)e^{\lambda x}[\lambda - g(x)]\}y(t) dt = f(x).$$

This is a special case of equation 2.9.16 with $h(x) = be^{\lambda x}$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = [g(x) + be^{\lambda x}] \frac{G(x)}{G(t)} + (b^2 e^{2\lambda x} + b\lambda e^{\lambda x}) \frac{H(x)}{G(t)} \int_t^x \frac{G(s)}{H(s)} ds,$$

where $G(x) = \exp\left[\int_a^x g(s) ds\right]$ and $H(x) = \exp\left(\frac{b}{\lambda} e^{\lambda x}\right)$.

$$26. \quad y(x) + \int_a^x \{\lambda e^{\lambda(x-t)} + [e^{\lambda t} g'_x(x) - \lambda e^{\lambda x} g(t)]h(t)\}y(t) dt = f(x).$$

This is a special case of equation 2.9.17 with $\varphi(x) = e^{\lambda x}$.

$$27. \quad y(x) - \int_a^x \{\lambda e^{-\lambda(x-t)} + [e^{\lambda x} g'_t(t) - \lambda e^{\lambda t} g(x)]h(x)\}y(t) dt = f(x).$$

This is a special case of equation 2.9.18 with $\varphi(x) = e^{\lambda x}$.

$$28. \quad y(x) + \int_a^x \cosh[\lambda(x-t)]g(t)y(t) dt = f(x).$$

Differentiating the equation with respect to x twice yields

$$y'_x(x) + g(x)y(x) + \lambda \int_a^x \sinh[\lambda(x-t)]g(t)y(t) dt = f'_x(x), \quad (1)$$

$$y''_{xx}(x) + [g(x)y(x)]'_x + \lambda^2 \int_a^x \cosh[\lambda(x-t)]g(t)y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the second-order linear ordinary differential equation

$$y''_{xx} + [g(x)y]'_x - \lambda^2 y = f''_{xx}(x) - \lambda^2 f(x). \quad (3)$$

By setting $x = a$ in the original equation and (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = f(a), \quad y'_x(a) = f'_x(a) - f(a)g(a). \quad (4)$$

Equation (3) under conditions (4) determines the solution of the original integral equation.

$$29. \quad y(x) + \int_a^x \cosh[\lambda(x-t)]g(x)h(t)y(t) dt = f(x).$$

The substitution $y(x) = g(x)u(x)$ leads to an equation of the form 2.9.28:

$$u(x) + \int_a^x \cosh[\lambda(x-t)]g(t)h(t)u(t) dt = f(x)/g(x).$$

$$30. \quad y(x) + \int_a^x \sinh[\lambda(x-t)]g(t)y(t) dt = f(x).$$

1°. Differentiating the equation with respect to x twice yields

$$y'_x(x) + \lambda \int_a^x \cosh[\lambda(x-t)]g(t)y(t) dt = f'_x(x), \quad (1)$$

$$y''_{xx}(x) + \lambda g(x)y(x) + \lambda^2 \int_a^x \sinh[\lambda(x-t)]g(t)y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the second-order linear ordinary differential equation

$$y''_{xx} + \lambda[g(x) - \lambda]y = f''_{xx}(x) - \lambda^2 f(x). \quad (3)$$

By setting $x = a$ in the original equation and (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = f(a), \quad y'_x(a) = f'_x(a). \quad (4)$$

For exact solutions of second-order linear ordinary differential equations (3) with various $g(x)$, see E. Kamke (1977), G. M. Murphy (1960), and A. D. Polyanin and V. F. Zaitsev (2003).

2°. Let $y_1 = y_1(x)$ and $y_2 = y_2(x)$ be two linearly independent solutions ($y_1/y_2 \neq \text{const}$) of the homogeneous differential equation $y''_{xx} + \lambda[g(x) - \lambda]y = 0$, which follows from (3) for $f(x) \equiv 0$. In this case, the Wronskian is a constant:

$$W = y_1(y_2)'_x - y_2(y_1)'_x \equiv \text{const}.$$

The solution of the nonhomogeneous equation (3) under conditions (4) with arbitrary $f = f(x)$ has the form

$$y(x) = f(x) + \frac{\lambda}{W} \int_a^x [y_1(x)y_2(t) - y_2(x)y_1(t)]g(t)f(t) dt \quad (5)$$

and determines the solution of the original integral equation.

3°. Given only one nontrivial solution $y_1 = y_1(x)$ of the linear homogeneous differential equation $y''_{xx} + \lambda[g(x) - \lambda]y = 0$, one can obtain the solution of the nonhomogeneous equation (3) under the initial conditions (4) by formula (5) with

$$W = 1, \quad y_2(x) = y_1(x) \int_b^x \frac{d\xi}{y_1^2(\xi)},$$

where b is an arbitrary number.

$$31. \quad y(x) + \int_a^x \sinh[\lambda(x-t)]g(x)h(t)y(t) dt = f(x).$$

The substitution $y(x) = g(x)u(x)$ leads to an equation of the form 2.9.30:

$$u(x) + \int_a^x \sinh[\lambda(x-t)]g(t)h(t)u(t) dt = f(x)/g(x).$$

$$32. \quad y(x) - \int_a^x \{g(x) + b \cosh(\lambda x) + b(x-t)[\lambda \sinh(\lambda x) - \cosh(\lambda x)g(x)]\} y(t) dt = f(x).$$

This is a special case of equation 2.9.16 with $h(x) = b \cosh(\lambda x)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = [g(x) + b \cosh(\lambda x)] \frac{G(x)}{G(t)} + [b^2 \cosh^2(\lambda x) + b\lambda \sinh(\lambda x)] \frac{H(x)}{G(t)} \int_t^x \frac{G(s)}{H(s)} ds,$$

where $G(x) = \exp \left[\int_a^x g(s) ds \right]$ and $H(x) = \exp \left[\frac{b}{\lambda} \sinh(\lambda x) \right]$.

$$33. \quad y(x) - \int_a^x \{g(x) + b \sinh(\lambda x) + b(x-t)[\lambda \cosh(\lambda x) - \sinh(\lambda x)g(x)]\} y(t) dt = f(x).$$

This is a special case of equation 2.9.16 with $h(x) = b \sinh(\lambda x)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = [g(x) + b \sinh(\lambda x)] \frac{G(x)}{G(t)} + [b^2 \sinh^2(\lambda x) + b\lambda \cosh(\lambda x)] \frac{H(x)}{G(t)} \int_t^x \frac{G(s)}{H(s)} ds,$$

where $G(x) = \exp \left[\int_a^x g(s) ds \right]$ and $H(x) = \exp \left[\frac{b}{\lambda} \cosh(\lambda x) \right]$.

$$34. \quad y(x) + \int_a^x \cos[\lambda(x-t)]g(t)y(t) dt = f(x).$$

Differentiating the equation with respect to x twice yields

$$y'_x(x) + g(x)y(x) - \lambda \int_a^x \sin[\lambda(x-t)]g(t)y(t) dt = f'_x(x), \quad (1)$$

$$y''_{xx}(x) + [g(x)y(x)]'_x - \lambda^2 \int_a^x \cos[\lambda(x-t)]g(t)y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the second-order linear ordinary differential equation

$$y''_{xx} + [g(x)y]'_x + \lambda^2 y = f''_{xx}(x) + \lambda^2 f(x). \quad (3)$$

By setting $x = a$ in the original equation and (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = f(a), \quad y'_x(a) = f'_x(a) - f(a)g(a). \quad (4)$$

$$35. \quad y(x) + \int_a^x \cos[\lambda(x-t)]g(x)h(t)y(t) dt = f(x).$$

The substitution $y(x) = g(x)u(x)$ leads to an equation of the form 2.9.34:

$$u(x) + \int_a^x \cos[\lambda(x-t)]g(t)h(t)u(t) dt = f(x)/g(x).$$

$$36. \quad y(x) + \int_a^x \sin[\lambda(x-t)]g(t)y(t) dt = f(x).$$

1°. Differentiating the equation with respect to x twice yields

$$y'_x(x) + \lambda \int_a^x \cos[\lambda(x-t)]g(t)y(t) dt = f'_x(x), \quad (1)$$

$$y''_{xx}(x) + \lambda g(x)y(x) - \lambda^2 \int_a^x \sin[\lambda(x-t)]g(t)y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the second-order linear ordinary differential equation

$$y''_{xx} + \lambda[g(x) + \lambda]y = f''_{xx}(x) + \lambda^2 f(x). \quad (3)$$

By setting $x = a$ in the original equation and (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = f(a), \quad y'_x(a) = f'_x(a). \quad (4)$$

For exact solutions of second-order linear ordinary differential equations (3) with various $f(x)$, see E. Kamke (1977) and A. D. Polyanin and V. F. Zaitsev (2003).

2°. Let $y_1 = y_1(x)$ and $y_2 = y_2(x)$ be two linearly independent solutions ($y_1/y_2 \neq \text{const}$) of the homogeneous differential equation $y''_{xx} + \lambda[g(x) - \lambda]y = 0$, which follows from (3) for $f(x) \equiv 0$. In this case, the Wronskian is a constant:

$$W = y_1(y_2)'_x - y_2(y_1)'_x \equiv \text{const}.$$

The solution of the nonhomogeneous equation (3) under conditions (4) with arbitrary $f = f(x)$ has the form

$$y(x) = f(x) + \frac{\lambda}{W} \int_a^x [y_1(x)y_2(t) - y_2(x)y_1(t)]g(t)f(t) dt \quad (5)$$

and determines the solution of the original integral equation.

3°. Given only one nontrivial solution $y_1 = y_1(x)$ of the linear homogeneous differential equation $y''_{xx} + \lambda[g(x) + \lambda]y = 0$, one can obtain the solution of the nonhomogeneous equation (3) under the initial conditions (4) by formula (5) with

$$W = 1, \quad y_2(x) = y_1(x) \int_b^x \frac{d\xi}{y_1^2(\xi)},$$

where b is an arbitrary number.

$$37. \quad y(x) + \int_a^x \sin[\lambda(x-t)]g(x)h(t)y(t) dt = f(x).$$

The substitution $y(x) = g(x)u(x)$ leads to an equation of the form 2.9.36:

$$u(x) + \int_a^x \sin[\lambda(x-t)]g(t)h(t)u(t) dt = f(x)/g(x).$$

$$38. \quad y(x) - \int_a^x \{g(x) + b \cos(\lambda x) - b(x-t)[\lambda \sin(\lambda x) + \cos(\lambda x)g(x)]\} y(t) dt = f(x).$$

This is a special case of equation 2.9.16 with $h(x) = b \cos(\lambda x)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = [g(x) + b \cos(\lambda x)] \frac{G(x)}{G(t)} + [b^2 \cos^2(\lambda x) - b\lambda \sin(\lambda x)] \frac{H(x)}{G(t)} \int_t^x \frac{G(s)}{H(s)} ds,$$

where $G(x) = \exp \left[\int_a^x g(s) ds \right]$ and $H(x) = \exp \left[\frac{b}{\lambda} \sin(\lambda x) \right]$.

$$39. \quad y(x) - \int_a^x \{g(x) + b \sin(\lambda x) + b(x-t)[\lambda \cos(\lambda x) - \sin(\lambda x)g(x)]\} y(t) dt = f(x).$$

This is a special case of equation 2.9.16 with $h(x) = b \sin(\lambda x)$.

Solution:

$$y(x) = f(x) + \int_a^x R(x,t)f(t) dt,$$

$$R(x,t) = [g(x) + b \sin(\lambda x)] \frac{G(x)}{G(t)} + [b^2 \sin^2(\lambda x) + b\lambda \cos(\lambda x)] \frac{H(x)}{G(t)} \int_t^x \frac{G(s)}{H(s)} ds,$$

where $G(x) = \exp \left[\int_a^x g(s) ds \right]$ and $H(x) = \exp \left[-\frac{b}{\lambda} \cos(\lambda x) \right]$.

2.9-2. Equations with Difference Kernel: $K(x,t) = K(x-t)$.

$$40. \quad y(x) + \int_a^x K(x-t)y(t) dt = f(x).$$

Renewal equation.

1°. To solve this integral equation, direct and inverse Laplace transforms are used.

The solution can be represented in the form

$$y(x) = f(x) - \int_a^x R(x-t)f(t) dt. \quad (1)$$

Here the resolvent $R(x)$ is expressed via the kernel $K(x)$ of the original equation as follows:

$$R(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \tilde{R}(p)e^{px} dp,$$

$$\tilde{R}(p) = \frac{\tilde{K}(p)}{1 + \tilde{K}(p)}, \quad \tilde{K}(p) = \int_0^\infty K(x)e^{-px} dx.$$

⊙ References: R. Bellman and K. L. Cooke (1963), M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971), V. I. Smirnov (1974).

2°. Let $w = w(x)$ be the solution of the simpler auxiliary equation with $a = 0$ and $f \equiv 1$:

$$w(x) + \int_0^x K(x-t)w(t) dt = 1. \quad (2)$$

Then the solution of the original integral equation with arbitrary $f = f(x)$ is expressed via the solution of the auxiliary equation (2) as

$$y(x) = \frac{d}{dx} \int_a^x w(x-t)f(t) dt = f(a)w(x-a) + \int_a^x w(x-t)f'_t(t) dt.$$

⊙ Reference: R. Bellman and K. L. Cooke (1963).

41. $y(x) + \int_{-\infty}^x K(x-t)y(t) dt = 0.$

Eigenfunctions of this integral equation are determined by the roots of the following transcendental (algebraic) equation for the parameter λ :

$$\int_0^\infty K(z)e^{-\lambda z} dz = -1. \tag{1}$$

The left-hand side of this equation is the Laplace transform of the kernel of the integral equation.

1°. For a real simple root λ_k of equation (1) there is a corresponding eigenfunction

$$y_k(x) = \exp(\lambda_k x).$$

2°. For a real root λ_k of multiplicity r there are corresponding r eigenfunctions

$$y_{k1}(x) = \exp(\lambda_k x), \quad y_{k2}(x) = x \exp(\lambda_k x), \quad \dots, \quad y_{kr}(x) = x^{r-1} \exp(\lambda_k x).$$

3°. For a complex simple root $\lambda_k = \alpha_k + i\beta_k$ of equation (1) there is a corresponding eigenfunction pair

$$y_k^{(1)}(x) = \exp(\alpha_k x) \cos(\beta_k x), \quad y_k^{(2)}(x) = \exp(\alpha_k x) \sin(\beta_k x).$$

4°. For a complex root $\lambda_k = \alpha_k + i\beta_k$ of multiplicity r there are corresponding r eigenfunction pairs

$$\begin{aligned} y_{k1}^{(1)}(x) &= \exp(\alpha_k x) \cos(\beta_k x), & y_{k1}^{(2)}(x) &= \exp(\alpha_k x) \sin(\beta_k x), \\ y_{k2}^{(1)}(x) &= x \exp(\alpha_k x) \cos(\beta_k x), & y_{k2}^{(2)}(x) &= x \exp(\alpha_k x) \sin(\beta_k x), \\ &\dots & & \dots \\ y_{kr}^{(1)}(x) &= x^{r-1} \exp(\alpha_k x) \cos(\beta_k x), & y_{kr}^{(2)}(x) &= x^{r-1} \exp(\alpha_k x) \sin(\beta_k x). \end{aligned}$$

The general solution is the combination (with arbitrary constants) of the eigenfunctions of the homogeneous integral equation.

► For equations 2.9.42–2.9.51, only particular solutions are given. To obtain the general solution, one must add the general solution of the corresponding homogeneous equation 2.9.41 to the particular solution.

42. $y(x) + \int_{-\infty}^x K(x-t)y(t) dt = Ax^n, \quad n = 0, 1, 2, \dots$

This is a special case of equation 2.9.44 with $\lambda = 0$.

1°. A solution with $n = 0$:

$$y(x) = \frac{A}{B}, \quad B = 1 + \int_0^\infty K(z) dz.$$

2°. A solution with $n = 1$:

$$y(x) = \frac{A}{B}x + \frac{AC}{B^2}, \quad B = 1 + \int_0^\infty K(z) dz, \quad C = \int_0^\infty zK(z) dz.$$

3°. A solution with $n = 2$:

$$\begin{aligned} y_2(x) &= \frac{A}{B}x^2 + 2\frac{AC}{B^2}x + 2\frac{AC^2}{B^3} - \frac{AD}{B^2}, \\ B &= 1 + \int_0^\infty K(z) dz, \quad C = \int_0^\infty zK(z) dz, \quad D = \int_0^\infty z^2K(z) dz. \end{aligned}$$

4°. A solution with $n = 3, 4, \dots$ is given by:

$$y_n(x) = A \left\{ \frac{\partial^n}{\partial \lambda^n} \left[\frac{e^{\lambda x}}{B(\lambda)} \right] \right\}_{\lambda=0}, \quad B(\lambda) = 1 + \int_0^\infty K(z)e^{-\lambda z} dz.$$

$$43. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = Ae^{\lambda x}.$$

A solution:

$$y(x) = \frac{A}{B}e^{\lambda x}, \quad B = 1 + \int_0^{\infty} K(z)e^{-\lambda z} dz.$$

The integral term in the expression for B is the Laplace transform of $K(z)$, which may be calculated using tables of Laplace transforms (e.g., see Supplement 5).

$$44. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = Ax^n e^{\lambda x}, \quad n = 1, 2, \dots$$

1°. A solution with $n = 1$:

$$y_1(x) = \frac{A}{B}xe^{\lambda x} + \frac{AC}{B^2}e^{\lambda x},$$

$$B = 1 + \int_0^{\infty} K(z)e^{-\lambda z} dz, \quad C = \int_0^{\infty} zK(z)e^{-\lambda z} dz.$$

It is convenient to calculate B and C using tables of Laplace transforms.

2°. A solution with $n = 2$:

$$y_2(x) = \frac{A}{B}x^2e^{\lambda x} + 2\frac{AC}{B^2}xe^{\lambda x} + \left(2\frac{AC^2}{B^3} - \frac{AD}{B^2}\right)e^{\lambda x},$$

$$B = 1 + \int_0^{\infty} K(z)e^{-\lambda z} dz, \quad C = \int_0^{\infty} zK(z)e^{-\lambda z} dz, \quad D = \int_0^{\infty} z^2K(z)e^{-\lambda z} dz.$$

3°. A solution with $n = 3, 4, \dots$ is given by:

$$y_n(x) = \frac{\partial}{\partial \lambda} y_{n-1}(x) = A \frac{\partial^n}{\partial \lambda^n} \left[\frac{e^{\lambda x}}{B(\lambda)} \right], \quad B(\lambda) = 1 + \int_0^{\infty} K(z)e^{-\lambda z} dz.$$

$$45. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = A \cosh(\lambda x).$$

A solution:

$$y(x) = \frac{A}{2B_-}e^{\lambda x} + \frac{A}{2B_+}e^{-\lambda x} = \frac{1}{2} \left(\frac{A}{B_-} + \frac{A}{B_+} \right) \cosh(\lambda x) + \frac{1}{2} \left(\frac{A}{B_-} - \frac{A}{B_+} \right) \sinh(\lambda x),$$

$$B_- = 1 + \int_0^{\infty} K(z)e^{-\lambda z} dz, \quad B_+ = 1 + \int_0^{\infty} K(z)e^{\lambda z} dz.$$

$$46. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = A \sinh(\lambda x).$$

A solution:

$$y(x) = \frac{A}{2B_-}e^{\lambda x} - \frac{A}{2B_+}e^{-\lambda x} = \frac{1}{2} \left(\frac{A}{B_-} - \frac{A}{B_+} \right) \cosh(\lambda x) + \frac{1}{2} \left(\frac{A}{B_-} + \frac{A}{B_+} \right) \sinh(\lambda x),$$

$$B_- = 1 + \int_0^{\infty} K(z)e^{-\lambda z} dz, \quad B_+ = 1 + \int_0^{\infty} K(z)e^{\lambda z} dz.$$

$$47. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = A \cos(\lambda x).$$

A solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \cos(\lambda x) - B_s \sin(\lambda x)],$$

$$B_c = 1 + \int_0^\infty K(z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(z) \sin(\lambda z) dz.$$

$$48. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = A \sin(\lambda x).$$

A solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \sin(\lambda x) + B_s \cos(\lambda x)],$$

$$B_c = 1 + \int_0^\infty K(z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(z) \sin(\lambda z) dz.$$

$$49. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = A e^{\mu x} \cos(\lambda x).$$

A solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} e^{\mu x} [B_c \cos(\lambda x) - B_s \sin(\lambda x)],$$

$$B_c = 1 + \int_0^\infty K(z) e^{-\mu z} \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(z) e^{-\mu z} \sin(\lambda z) dz.$$

$$50. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = A e^{\mu x} \sin(\lambda x).$$

A solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} e^{\mu x} [B_c \sin(\lambda x) + B_s \cos(\lambda x)],$$

$$B_c = 1 + \int_0^\infty K(z) e^{-\mu z} \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(z) e^{-\mu z} \sin(\lambda z) dz.$$

$$51. \quad y(x) + \int_{-\infty}^x K(x-t)y(t) dt = f(x).$$

1°. For a polynomial right-hand side, $f(x) = \sum_{k=0}^n A_k x^k$, a solution has the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients. One can also make use of the formula given in item 4° of equation 2.9.42 to construct the solution.

2°. For $f(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the B_k are found by the method of undetermined coefficients. One can also make use of the formula given in item 3° of equation 2.9.44 to construct the solution.

3°. For $f(x) = \sum_{k=0}^n A_k \exp(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_k} \exp(\lambda_k x), \quad B_k = 1 + \int_0^{\infty} K(z) \exp(-\lambda_k z) dz.$$

4°. For $f(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

5°. For $f(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

6°. For $f(x) = \sum_{k=0}^n A_k \cos(\lambda_k x)$, the solution of a equation has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_{ck}^2 + B_{sk}^2} [B_{ck} \cos(\lambda_k x) - B_{sk} \sin(\lambda_k x)],$$

$$B_{ck} = 1 + \int_0^{\infty} K(z) \cos(\lambda_k z) dz, \quad B_{sk} = \int_0^{\infty} K(z) \sin(\lambda_k z) dz.$$

7°. For $f(x) = \sum_{k=0}^n A_k \sin(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_{ck}^2 + B_{sk}^2} [B_{ck} \sin(\lambda_k x) + B_{sk} \cos(\lambda_k x)],$$

$$B_{ck} = 1 + \int_0^{\infty} K(z) \cos(\lambda_k z) dz, \quad B_{sk} = \int_0^{\infty} K(z) \sin(\lambda_k z) dz.$$

8°. For $f(x) = \cos(\lambda x) \sum_{k=0}^n A_k \exp(\mu_k x)$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n \frac{A_k B_{ck}}{B_{ck}^2 + B_{sk}^2} \exp(\mu_k x) - \sin(\lambda x) \sum_{k=0}^n \frac{A_k B_{sk}}{B_{ck}^2 + B_{sk}^2} \exp(\mu_k x),$$

$$B_{ck} = 1 + \int_0^{\infty} K(z) \exp(-\mu_k z) \cos(\lambda z) dz, \quad B_{sk} = \int_0^{\infty} K(z) \exp(-\mu_k z) \sin(\lambda z) dz.$$

9°. For $f(x) = \sin(\lambda x) \sum_{k=0}^n A_k \exp(\mu_k x)$, a solution of the equation has the form

$$y(x) = \sin(\lambda x) \sum_{k=0}^n \frac{A_k B_{ck}}{B_{ck}^2 + B_{sk}^2} \exp(\mu_k x) + \cos(\lambda x) \sum_{k=0}^n \frac{A_k B_{sk}}{B_{ck}^2 + B_{sk}^2} \exp(\mu_k x),$$

$$B_{ck} = 1 + \int_0^\infty K(z) \exp(-\mu_k z) \cos(\lambda z) dz, \quad B_{sk} = \int_0^\infty K(z) \exp(-\mu_k z) \sin(\lambda z) dz.$$

52. $y(x) + \int_x^\infty K(x-t)y(t) dt = 0.$

Eigenfunctions of this integral equation are determined by the roots of the following transcendental (algebraic) equation for the parameter λ :

$$\int_0^\infty K(-z)e^{\lambda z} dz = -1. \tag{1}$$

The left-hand side of this equation is the Laplace transform of the function $K(-z)$ with parameter $-\lambda$.

1°. For a real simple root λ_k of equation (1) there is a corresponding eigenfunction

$$y_k(x) = \exp(\lambda_k x).$$

2°. For a real root λ_k of multiplicity r there are corresponding r eigenfunctions

$$y_{k1}(x) = \exp(\lambda_k x), \quad y_{k2}(x) = x \exp(\lambda_k x), \quad \dots, \quad y_{kr}(x) = x^{r-1} \exp(\lambda_k x).$$

3°. For a complex simple root $\lambda_k = \alpha_k + i\beta_k$ of equation (1) there is a corresponding eigenfunction pair

$$y_k^{(1)}(x) = \exp(\alpha_k x) \cos(\beta_k x), \quad y_k^{(2)}(x) = \exp(\alpha_k x) \sin(\beta_k x).$$

4°. For a complex root $\lambda_k = \alpha_k + i\beta_k$ of multiplicity r there are corresponding r eigenfunction pairs

$$\begin{array}{ll} y_{k1}^{(1)}(x) = \exp(\alpha_k x) \cos(\beta_k x), & y_{k1}^{(2)}(x) = \exp(\alpha_k x) \sin(\beta_k x), \\ y_{k2}^{(1)}(x) = x \exp(\alpha_k x) \cos(\beta_k x), & y_{k2}^{(2)}(x) = x \exp(\alpha_k x) \sin(\beta_k x), \\ \dots & \dots \\ y_{kr}^{(1)}(x) = x^{r-1} \exp(\alpha_k x) \cos(\beta_k x), & y_{kr}^{(2)}(x) = x^{r-1} \exp(\alpha_k x) \sin(\beta_k x). \end{array}$$

The general solution is the combination (with arbitrary constants) of the eigenfunctions of the homogeneous integral equation.

► For equations 2.9.53–2.9.62, only particular solutions are given. To obtain the general solution, one must add the general solution of the corresponding homogeneous equation 2.9.52 to the particular solution.

53. $y(x) + \int_x^\infty K(x-t)y(t) dt = Ax^n, \quad n = 0, 1, 2, \dots$

This is a special case of equation 2.9.55 with $\lambda = 0$.

1°. A solution with $n = 0$:

$$y(x) = \frac{A}{B}, \quad B = 1 + \int_0^\infty K(-z) dz.$$

2°. A solution with $n = 1$:

$$y(x) = \frac{A}{B}x - \frac{AC}{B^2}, \quad B = 1 + \int_0^\infty K(-z) dz, \quad C = \int_0^\infty zK(-z) dz.$$

3°. A solution with $n = 2$:

$$y_2(x) = \frac{A}{B}x^2 - 2\frac{AC}{B^2}x + 2\frac{AC^2}{B^3} - \frac{AD}{B^2},$$

$$B = 1 + \int_0^\infty K(-z) dz, \quad C = \int_0^\infty zK(-z) dz, \quad D = \int_0^\infty z^2K(-z) dz.$$

4°. A solution with $n = 3, 4, \dots$ is given by:

$$y_n(x) = A \left\{ \frac{\partial^n}{\partial \lambda^n} \left[\frac{e^{\lambda x}}{B(\lambda)} \right] \right\}_{\lambda=0}, \quad B(\lambda) = 1 + \int_0^\infty K(-z)e^{\lambda z} dz.$$

54. $y(x) + \int_x^\infty K(x-t)y(t) dt = Ae^{\lambda x}.$

A solution:

$$y(x) = \frac{A}{B}e^{\lambda x}, \quad B = 1 + \int_0^\infty K(-z)e^{\lambda z} dz = 1 + \mathfrak{L}\{K(-z), -\lambda\}.$$

The integral term in the expression for B is the Laplace transform of $K(-z)$ with parameter $-\lambda$, which may be calculated using tables of Laplace transforms (e.g., see H. Bateman and A. Erdélyi (vol. 1, 1954), V. A. Ditkin and A. P. Prudnikov (1965), and Supplement 5).

55. $y(x) + \int_x^\infty K(x-t)y(t) dt = Ax^n e^{\lambda x}, \quad n = 1, 2, \dots$

1°. A solution with $n = 1$:

$$y_1(x) = \frac{A}{B}xe^{\lambda x} - \frac{AC}{B^2}e^{\lambda x},$$

$$B = 1 + \int_0^\infty K(-z)e^{\lambda z} dz, \quad C = \int_0^\infty zK(-z)e^{\lambda z} dz.$$

It is convenient to calculate B and C using tables of Laplace transforms (with parameter $-\lambda$).

2°. A solution with $n = 2$:

$$y_2(x) = \frac{A}{B}x^2e^{\lambda x} - 2\frac{AC}{B^2}xe^{\lambda x} + \left(2\frac{AC^2}{B^3} - \frac{AD}{B^2}\right)e^{\lambda x},$$

$$B = 1 + \int_0^\infty K(-z)e^{\lambda z} dz, \quad C = \int_0^\infty zK(-z)e^{\lambda z} dz, \quad D = \int_0^\infty z^2K(-z)e^{\lambda z} dz.$$

3°. A solution with $n = 3, 4, \dots$ is given by

$$y_n(x) = \frac{\partial}{\partial \lambda} y_{n-1}(x) = A \frac{\partial^n}{\partial \lambda^n} \left[\frac{e^{\lambda x}}{B(\lambda)} \right], \quad B(\lambda) = 1 + \int_0^\infty K(-z)e^{\lambda z} dz.$$

$$56. \quad y(x) + \int_x^\infty K(x-t)y(t) dt = A \cosh(\lambda x).$$

A solution:

$$y(x) = \frac{A}{2B_+} e^{\lambda x} + \frac{A}{2B_-} e^{-\lambda x} = \frac{1}{2} \left(\frac{A}{B_+} + \frac{A}{B_-} \right) \cosh(\lambda x) + \frac{1}{2} \left(\frac{A}{B_+} - \frac{A}{B_-} \right) \sinh(\lambda x),$$

$$B_+ = 1 + \int_0^\infty K(-z)e^{\lambda z} dz, \quad B_- = 1 + \int_0^\infty K(-z)e^{-\lambda z} dz.$$

$$57. \quad y(x) + \int_x^\infty K(x-t)y(t) dt = A \sinh(\lambda x).$$

A solution:

$$y(x) = \frac{A}{2B_+} e^{\lambda x} - \frac{A}{2B_-} e^{-\lambda x} = \frac{1}{2} \left(\frac{A}{B_+} - \frac{A}{B_-} \right) \cosh(\lambda x) + \frac{1}{2} \left(\frac{A}{B_+} + \frac{A}{B_-} \right) \sinh(\lambda x),$$

$$B_+ = 1 + \int_0^\infty K(-z)e^{\lambda z} dz, \quad B_- = 1 + \int_0^\infty K(-z)e^{-\lambda z} dz.$$

$$58. \quad y(x) + \int_x^\infty K(x-t)y(t) dt = A \cos(\lambda x).$$

A solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \cos(\lambda x) + B_s \sin(\lambda x)],$$

$$B_c = 1 + \int_0^\infty K(-z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z) \sin(\lambda z) dz.$$

$$59. \quad y(x) + \int_x^\infty K(x-t)y(t) dt = A \sin(\lambda x).$$

A solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \sin(\lambda x) - B_s \cos(\lambda x)],$$

$$B_c = 1 + \int_0^\infty K(-z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z) \sin(\lambda z) dz.$$

$$60. \quad y(x) + \int_x^\infty K(x-t)y(t) dt = Ae^{\mu x} \cos(\lambda x).$$

A solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} e^{\mu x} [B_c \cos(\lambda x) + B_s \sin(\lambda x)],$$

$$B_c = 1 + \int_0^\infty K(-z)e^{\mu z} \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z)e^{\mu z} \sin(\lambda z) dz.$$

$$61. \quad y(x) + \int_x^\infty K(x-t)y(t) dt = Ae^{\mu x} \sin(\lambda x).$$

A solution:

$$y(x) = \frac{A}{B_c^2 + B_s^2} e^{\mu x} [B_c \sin(\lambda x) - B_s \cos(\lambda x)],$$

$$B_c = 1 + \int_0^\infty K(-z)e^{\mu z} \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z)e^{\mu z} \sin(\lambda z) dz.$$

$$62. \quad y(x) + \int_x^\infty K(x-t)y(t) dt = f(x).$$

1°. For a polynomial right-hand side, $f(x) = \sum_{k=0}^n A_k x^k$, a solution has the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients. One can also make use of the formula given in item 4° of equation 2.9.53 to construct the solution.

2°. For $f(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients. One can also make use of the formula given in item 3° of equation 2.9.55 to construct the solution.

3°. For $f(x) = \sum_{k=0}^n A_k \exp(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_k} \exp(\lambda_k x), \quad B_k = 1 + \int_0^\infty K(-z) \exp(\lambda_k z) dz.$$

4°. For $f(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$ a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

5°. For $f(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the B_k and C_k are found by the method of undetermined coefficients.

6°. For $f(x) = \sum_{k=0}^n A_k \cos(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_{ck}^2 + B_{sk}^2} [B_{ck} \cos(\lambda_k x) + B_{sk} \sin(\lambda_k x)],$$

$$B_{ck} = 1 + \int_0^\infty K(-z) \cos(\lambda_k z) dz, \quad B_{sk} = \int_0^\infty K(-z) \sin(\lambda_k z) dz.$$

7°. For $f(x) = \sum_{k=0}^n A_k \sin(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_{ck}^2 + B_{sk}^2} [B_{ck} \sin(\lambda_k x) - B_{sk} \cos(\lambda_k x)],$$

$$B_{ck} = 1 + \int_0^\infty K(-z) \cos(\lambda_k z) dz, \quad B_{sk} = \int_0^\infty K(-z) \sin(\lambda_k z) dz.$$

8°. For $f(x) = \cos(\lambda x) \sum_{k=0}^n A_k \exp(\mu_k x)$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n \frac{A_k B_{ck}}{B_{ck}^2 + B_{sk}^2} \exp(\mu_k x) + \sin(\lambda x) \sum_{k=0}^n \frac{A_k B_{sk}}{B_{ck}^2 + B_{sk}^2} \exp(\mu_k x),$$

$$B_{ck} = 1 + \int_0^\infty K(-z) \exp(\mu_k z) \cos(\lambda z) dz, \quad B_{sk} = \int_0^\infty K(-z) \exp(\mu_k z) \sin(\lambda z) dz.$$

9°. For $f(x) = \sin(\lambda x) \sum_{k=0}^n A_k \exp(\mu_k x)$, a solution of the equation has the form

$$y(x) = \sin(\lambda x) \sum_{k=0}^n \frac{A_k B_{ck}}{B_{ck}^2 + B_{sk}^2} \exp(\mu_k x) - \cos(\lambda x) \sum_{k=0}^n \frac{A_k B_{sk}}{B_{ck}^2 + B_{sk}^2} \exp(\mu_k x),$$

$$B_{ck} = 1 + \int_0^\infty K(-z) \exp(\mu_k z) \cos(\lambda z) dz, \quad B_{sk} = \int_0^\infty K(-z) \exp(\mu_k z) \sin(\lambda z) dz.$$

10°. In the general case of arbitrary right-hand side $f = f(x)$, the solution of the integral equation can be represented in the form

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(p)}{1 + \tilde{k}(-p)} e^{px} dp,$$

$$\tilde{f}(p) = \int_0^\infty f(x) e^{-px} dx, \quad \tilde{k}(-p) = \int_0^\infty K(-z) e^{pz} dz.$$

To calculate $\tilde{f}(p)$ and $\tilde{k}(-p)$, it is convenient to use tables of Laplace transforms, and to determine $y(x)$, tables of inverse Laplace transforms.

2.9-3. Other Equations.

63. $y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = 0.$

Eigenfunctions of this integral equation are determined by the roots of the following transcendental (algebraic) equation for the parameter λ :

$$\int_0^1 f(z) z^\lambda dz = -1. \quad (1)$$

1°. For a real simple root λ_k of equation (1) there is a corresponding eigenfunction

$$y_k(x) = x^{\lambda_k}.$$

2°. For a real root λ_k of multiplicity r there are corresponding r eigenfunctions

$$y_{k1}(x) = x^{\lambda_k}, \quad y_{k2}(x) = x^{\lambda_k} \ln x, \quad \dots, \quad y_{kr}(x) = x^{\lambda_k} \ln^{r-1} x.$$

3°. For a complex simple root $\lambda_k = \alpha_k + i\beta_k$ of equation (1) there is a corresponding eigenfunction pair

$$y_k^{(1)}(x) = x^{\alpha_k} \cos(\beta_k \ln x), \quad y_k^{(2)}(x) = x^{\alpha_k} \sin(\beta_k \ln x).$$

4°. For a complex root $\lambda_k = \alpha_k + i\beta_k$ of multiplicity r there are corresponding r eigenfunction pairs

$$\begin{aligned} y_{k1}^{(1)}(x) &= x^{\alpha_k} \cos(\beta_k \ln x), & y_{k1}^{(2)}(x) &= x^{\alpha_k} \sin(\beta_k \ln x), \\ y_{k2}^{(1)}(x) &= x^{\alpha_k} \ln x \cos(\beta_k \ln x), & y_{k2}^{(2)}(x) &= x^{\alpha_k} \ln x \sin(\beta_k \ln x), \\ &\dots\dots\dots & & \dots\dots\dots \\ y_{kr}^{(1)}(x) &= x^{\alpha_k} \ln^{r-1} x \cos(\beta_k \ln x), & y_{kr}^{(2)}(x) &= x^{\alpha_k} \ln^{r-1} x \sin(\beta_k \ln x). \end{aligned}$$

The general solution is the combination (with arbitrary constants) of the eigenfunctions of the homogeneous integral equation.

► For equations 2.9.64–2.9.71, only particular solutions are given. To obtain the general solution, one must add the general solution of the corresponding homogeneous equation 2.9.63 to the particular solution.

64. $y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = Ax + B.$

A solution:

$$y(x) = \frac{A}{1 + I_1} x + \frac{B}{1 + I_0}, \quad I_0 = \int_0^1 f(t) dt, \quad I_1 = \int_0^1 t f(t) dt.$$

65. $y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = Ax^\beta.$

A solution:

$$y(x) = \frac{A}{B} x^\beta, \quad B = 1 + \int_0^1 f(t) t^\beta dt.$$

66. $y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = A \ln x + B.$

A solution:

$$y(x) = p \ln x + q,$$

where

$$p = \frac{A}{1 + I_0}, \quad q = \frac{B}{1 + I_0} - \frac{AI_1}{(1 + I_0)^2}, \quad I_0 = \int_0^1 f(t) dt, \quad I_1 = \int_0^1 f(t) \ln t dt.$$

67. $y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = Ax^\beta \ln x.$

A solution:

$$y(x) = px^\beta \ln x + qx^\beta,$$

where

$$p = \frac{A}{1 + I_1}, \quad q = -\frac{AI_2}{(1 + I_1)^2}, \quad I_1 = \int_0^1 f(t) t^\beta dt, \quad I_2 = \int_0^1 f(t) t^\beta \ln t dt.$$

$$68. \quad y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = A \cos(\ln x).$$

A solution:

$$y(x) = \frac{AI_c}{I_c^2 + I_s^2} \cos(\ln x) + \frac{AI_s}{I_c^2 + I_s^2} \sin(\ln x),$$

$$I_c = 1 + \int_0^1 f(t) \cos(\ln t) dt, \quad I_s = \int_0^1 f(t) \sin(\ln t) dt.$$

$$69. \quad y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = A \sin(\ln x).$$

A solution:

$$y(x) = -\frac{AI_s}{I_c^2 + I_s^2} \cos(\ln x) + \frac{AI_c}{I_c^2 + I_s^2} \sin(\ln x),$$

$$I_c = 1 + \int_0^1 f(t) \cos(\ln t) dt, \quad I_s = \int_0^1 f(t) \sin(\ln t) dt.$$

$$70. \quad y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = Ax^\beta \cos(\ln x) + Bx^\beta \sin(\ln x).$$

A solution:

$$y(x) = px^\beta \cos(\ln x) + qx^\beta \sin(\ln x),$$

where

$$p = \frac{AI_c - BI_s}{I_c^2 + I_s^2}, \quad q = \frac{AI_s + BI_c}{I_c^2 + I_s^2},$$

$$I_c = 1 + \int_0^1 f(t)t^\beta \cos(\ln t) dt, \quad I_s = \int_0^1 f(t)t^\beta \sin(\ln t) dt.$$

$$71. \quad y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t) dt = g(x).$$

1°. For a polynomial right-hand side,

$$g(x) = \sum_{n=0}^N A_n x^n$$

a solution bounded at zero is given by

$$y(x) = \sum_{n=0}^N \frac{A_n}{1 + f_n} x^n, \quad f_n = \int_0^1 f(z) z^n dz.$$

Here it is assumed that $f_0 < \infty$ and $f_n \neq -1$ ($n = 0, 1, 2, \dots$).

If for some n the relation $f_n = -1$ holds, then a solution differs from the above case in one term and has the form

$$y(x) = \sum_{m=0}^{n-1} \frac{A_m}{1 + f_m} x^m + \sum_{m=n+1}^N \frac{A_m}{1 + f_m} x^m + \frac{A_n}{\bar{f}_n} x^n \ln x, \quad \bar{f}_n = \int_0^1 f(z) z^n \ln z dz.$$

For arbitrary $g(x)$ expandable into power series, the formulas of item 1° can be used, in which one should set $N = \infty$. In this case, the convergence radius of the obtained solution $y(x)$ is equal to that of the function $g(x)$.

2°. For $g(x) = \ln x \sum_{k=0}^n A_k x^k$, a solution has the form

$$y(x) = \ln x \sum_{k=0}^n B_k x^k + \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

3°. For $g(x) = \sum_{k=0}^n A_k (\ln x)^k$, a solution of the equation has the form

$$y(x) = \sum_{k=0}^n B_k (\ln x)^k,$$

where the B_k are found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k \ln x)$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k \ln x) + \sum_{k=1}^n C_k \sin(\lambda_k \ln x),$$

where the B_k and C_k are found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k \ln x)$ a solution of the equation has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k \ln x) + \sum_{k=1}^n C_k \sin(\lambda_k \ln x),$$

where the B_k and C_k are found by the method of undetermined coefficients.

6°. For arbitrary right-hand side $g(x)$, the transformation

$$x = e^{-z}, \quad t = e^{-\tau}, \quad y(x) = e^z w(z), \quad f(\xi) = F(\ln \xi), \quad g(x) = e^z G(z)$$

leads to an equation with difference kernel of the form 2.9.62:

$$w(z) + \int_z^\infty F(z - \tau)w(\tau) d\tau = G(z).$$

7°. For arbitrary right-hand side $g(x)$, the solution of the integral equation can be expressed via the inverse Mellin transform (see Example 2 in Subsection 11.6-4).

2.10. Some Formulas and Transformations

Let the solution of the integral equation

$$y(x) + \int_a^x K(x, t)y(t) dt = f(x) \quad (1)$$

have the form

$$y(x) = f(x) + \int_a^x R(x, t)f(t) dt. \quad (2)$$

Then the solution of the more complicated integral equation

$$y(x) + \int_a^x K(x, t) \frac{g(x)}{g(t)} y(t) dt = f(x) \quad (3)$$

has the form

$$y(x) = f(x) + \int_a^x R(x, t) \frac{g(x)}{g(t)} f(t) dt. \quad (4)$$

Below are formulas for the solutions of integral equations of the form (3) for some specific functions $g(x)$. In all cases, it is assumed that the solution of equation (1) is known and is given by (2).

1°. The solution of the equation

$$y(x) + \int_a^x K(x, t)(x/t)^\lambda y(t) dt = f(x)$$

has the form

$$y(x) = f(x) + \int_a^x R(x, t)(x/t)^\lambda f(t) dt.$$

2°. The solution of the equation

$$y(x) + \int_a^x K(x, t)e^{\lambda(x-t)} y(t) dt = f(x)$$

has the form

$$y(x) = f(x) + \int_a^x R(x, t)e^{\lambda(x-t)} f(t) dt.$$

Chapter 3

Linear Equations of the First Kind with Constant Limits of Integration

► **Notation:** $f = f(x)$, $g = g(x)$, $h = h(x)$, $K = K(x)$, and $M = M(x)$ are arbitrary functions (these may be composite functions of the argument depending on two variables x and t); $A, B, C, a, b, c, k, \alpha, \beta, \gamma, \lambda$, and μ are free parameters; and n is a nonnegative integer.

3.1. Equations Whose Kernels Contain Power-Law Functions

3.1-1. Kernels Linear in the Arguments x and t .

1.
$$\int_0^1 |x - t| y(t) dt = f(x).$$

1°. Let us remove the modulus in the integrand:

$$\int_0^x (x - t)y(t) dt + \int_x^1 (t - x)y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x yields

$$\int_0^x y(t) dt - \int_x^1 y(t) dt = f'_x(x). \quad (2)$$

Differentiating (2) yields the solution

$$y(x) = \frac{1}{2} f''_{xx}(x). \quad (3)$$

2°. Let us demonstrate that the right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = 0$ and $x = 1$ in (1), we obtain two corollaries $\int_0^1 ty(t) dt = f(0)$ and $\int_0^1 (1 - t)y(t) dt = f(1)$, which can be rewritten in the form

$$\int_0^1 ty(t) dt = f(0), \quad \int_0^1 y(t) dt = f(0) + f(1). \quad (4)$$

In Section 3.1, we mean that kernels of the integral equations discussed may contain power-law functions or modulus of power-law functions.

Substitute $y(x)$ of (3) into (4). Integration by parts yields $f'_x(1) = f(1) + f(0)$ and $f'_x(1) - f'_x(0) = 2f(1) + 2f(0)$. Hence, we obtain the desired constraints for $f(x)$:

$$f'_x(1) = f(0) + f(1), \quad f'_x(0) + f'_x(1) = 0. \quad (5)$$

Conditions (5) make it possible to find the admissible general form of the right-hand side of the integral equation:

$$f(x) = F(x) + Ax + B, \\ A = -\frac{1}{2}[F'_x(1) + F'_x(0)], \quad B = \frac{1}{2}[F'_x(1) - F(1) - F(0)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function with bounded first derivative.

$$2. \quad \int_a^b |x-t| y(t) dt = f(x), \quad 0 \leq a < b < \infty.$$

This is a special case of equation 3.8.3 with $g(x) = x$.

Solution:

$$y(x) = \frac{1}{2} f''_{xx}(x).$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations. The general form of $f(x)$ is as follows:

$$f(x) = F(x) + Ax + B, \\ A = -\frac{1}{2}[F'_x(a) + F'_x(b)], \quad B = \frac{1}{2}[aF'_x(a) + bF'_x(b) - F(a) - F(b)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

$$3. \quad \int_0^a |\lambda x - t| y(t) dt = f(x), \quad \lambda > 0.$$

Here $0 \leq x \leq a$ and $0 \leq t \leq a$.

1°. Let us remove the modulus in the integrand:

$$\int_0^{\lambda x} (\lambda x - t)y(t) dt + \int_{\lambda x}^a (t - \lambda x)y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x , we find that

$$\lambda \int_0^{\lambda x} y(t) dt - \lambda \int_{\lambda x}^a y(t) dt = f'_x(x). \quad (2)$$

Differentiating (2) yields $2\lambda^2 y(\lambda x) = f''_{xx}(x)$. Hence, we obtain the solution

$$y(x) = \frac{1}{2\lambda^2} f''_{xx}\left(\frac{x}{\lambda}\right). \quad (3)$$

2°. Let us demonstrate that the right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = 0$ in (1) and (2), we obtain two corollaries

$$\int_0^a t y(t) dt = f(0), \quad \lambda \int_0^a y(t) dt = -f'_x(0). \quad (4)$$

Substitute $y(x)$ from (3) into (4). Integrating by parts yields the desired constraints for $f(x)$:

$$(a/\lambda)f'_x(a/\lambda) = f(0) + f(a/\lambda), \quad f'_x(0) + f'_x(a/\lambda) = 0. \quad (5)$$

Conditions (5) make it possible to establish the admissible general form of the right-hand side of the integral equation:

$$f(x) = F(z) + Az + B, \quad z = \lambda x; \\ A = -\frac{1}{2}[F'_z(a) + F'_z(0)], \quad B = \frac{1}{2}[aF'_z(a) - F(a) - F(0)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

$$4. \quad \int_0^a |x - \lambda t| y(t) dt = f(x), \quad \lambda > 0.$$

Here $0 \leq x \leq a$ and $0 \leq t \leq a$.

Solution:

$$y(x) = \frac{1}{2} \lambda f''_{xx}(\lambda x).$$

The right-hand side $f(x)$ of the integral equation must satisfy the relations

$$a \lambda f'_x(a \lambda) = f(0) + f(a \lambda), \quad f'_x(0) + f'_x(a \lambda) = 0.$$

Hence, it follows the general form of the right-hand side:

$$f(x) = F(x) + Ax + B, \quad A = -\frac{1}{2} [F'_x(\lambda a) + F'_x(0)], \quad B = \frac{1}{2} [a \lambda F'_x(a \lambda) - F(\lambda a) - F(0)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

3.1-2. Kernels Quadratic in the Arguments x and t .

$$5. \quad \int_0^a |Ax + Bx^2 - t| y(t) dt = f(x), \quad A > 0, \quad B > 0.$$

This is a special case of equation 3.8.5 with $g(x) = Ax + Bx^2$.

$$6. \quad \int_0^a |x - At - Bt^2| y(t) dt = f(x), \quad A > 0, \quad B > 0.$$

This is a special case of equation 3.8.6 with $g(x) = At + Bt^2$.

$$7. \quad \int_a^b |xt - t^2| y(t) dt = f(x), \quad 0 \leq a < b < \infty.$$

The substitution $w(t) = ty(t)$ leads to an equation of the form 3.1.2:

$$\int_a^b |x - t| w(t) dt = f(x).$$

$$8. \quad \int_a^b |x^2 - t^2| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = x^2$.

Solution: $y(x) = \frac{d}{dx} \left[\frac{f'_x(x)}{4x} \right]$. The right-hand side $f(x)$ of the equation must satisfy certain constraints, given in 3.8.3.

$$9. \quad \int_0^a |x^2 - \beta t^2| y(t) dt = f(x), \quad \beta > 0.$$

This is a special case of equation 3.8.4 with $g(x) = x^2$ and $\beta = \lambda^2$.

$$10. \quad \int_0^a |Ax + Bx^2 - A\lambda t - B\lambda^2 t^2| y(t) dt = f(x), \quad \lambda > 0.$$

This is a special case of equation 3.8.4 with $g(x) = Ax + Bx^2$.

3.1-3. Kernels Containing Integer Powers of x and t or Rational Functions.

$$11. \int_a^b |x-t|^3 y(t) dt = f(x).$$

Let us remove the modulus in the integrand:

$$\int_a^x (x-t)^3 y(t) dt + \int_x^b (t-x)^3 y(t) dt = f(x). \quad (1)$$

Differentiating (1) twice yields

$$6 \int_a^x (x-t)y(t) dt + 6 \int_x^b (t-x)y(t) dt = f''_{xx}(x).$$

This equation can be rewritten in the form 3.1.2:

$$\int_a^b |x-t| y(t) dt = \frac{1}{6} f''_{xx}(x). \quad (2)$$

Therefore the solution of the integral equation is given by

$$y(x) = \frac{1}{12} y''''_{xxxx}(x). \quad (3)$$

The right-hand side $f(x)$ of the equation must satisfy certain conditions. To obtain these conditions, one must substitute solution (3) into (1) with $x = a$ and $x = b$ and into (2) with $x = a$ and $x = b$, and then integrate the four resulting relations by parts.

$$12. \int_a^b |x^3 - t^3| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = x^3$.

$$13. \int_a^b |xt^2 - t^3| y(t) dt = f(x) \quad 0 \leq a < b < \infty.$$

The substitution $w(t) = t^2 y(t)$ leads to an equation of the form 3.1.2:

$$\int_a^b |x-t| w(t) dt = f(x).$$

$$14. \int_a^b |x^2 t - t^3| y(t) dt = f(x).$$

The substitution $w(t) = |t| y(t)$ leads to an equation of the form 3.1.8:

$$\int_a^b |x^2 - t^2| w(t) dt = f(x).$$

$$15. \int_0^a |x^3 - \beta t^3| y(t) dt = f(x), \quad \beta > 0.$$

This is a special case of equation 3.8.4 with $g(x) = x^3$ and $\beta = \lambda^3$.

$$16. \int_a^b |x-t|^{2n+1} y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

Solution:

$$y(x) = \frac{1}{2(2n+1)!} f_x^{(2n+2)}(x). \quad (1)$$

The right-hand side $f(x)$ of the equation must satisfy certain conditions. To obtain these conditions, one must substitute solution (1) into the relations

$$\int_a^b (t-a)^{2n+1} y(t) dt = f(a), \quad \int_a^b (t-a)^{2n-k} y(t) dt = \frac{(-1)^{k+1}}{A_k} f_x^{(k+1)}(a),$$

$$A_k = (2n+1)(2n) \dots (2n+1-k); \quad k = 0, 1, \dots, 2n,$$

and then integrate the resulting equations by parts.

$$17. \int_0^\infty \frac{y(t) dt}{x+t} = f(x).$$

The left-hand side of this equation is the *Stieltjes transform*.

1°. By setting

$$x = e^z, \quad t = e^\tau, \quad y(t) = e^{-\tau/2} w(\tau), \quad f(x) = e^{-z/2} g(z),$$

we obtain an integral equation with difference kernel of the form 3.8.15:

$$\int_{-\infty}^\infty \frac{w(\tau) d\tau}{2 \cosh\left[\frac{1}{2}(z-\tau)\right]} = g(z),$$

whose solution is given by

$$w(z) = \frac{1}{\sqrt{2\pi^3}} \int_{-\infty}^\infty \cosh(\pi u) \tilde{g}(u) e^{iux} du, \quad \tilde{g}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty g(z) e^{-iuz} dz, \quad i^2 = -1.$$

2°. Solution:

$$y(x) = \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow +0} [f(-x - i\varepsilon) - f(-x + i\varepsilon)]$$

$$= \frac{1}{\pi\sqrt{x}} \sum_{k=0}^\infty \frac{(-1)^k}{(2k)!} \left(\frac{\pi}{x} \frac{d}{dx}\right)^{2k} [\sqrt{x} f(x)].$$

3°. Under some assumptions, the solution of the original equation can be represented in the form

$$y(x) = \lim_{n \rightarrow \infty} \frac{(-1)^n}{(n+1)!(n-1)!} [x^{2n+1} f_x^{(n)}(x)]_x^{(n+1)}, \quad (1)$$

which is the real inversion of the Stieltjes transform.

An alternative form of the solution is

$$y(x) = \lim_{n \rightarrow \infty} \frac{(-1)^n}{2\pi} \left(\frac{e}{n}\right)^{2n} [x^{2n} f_x^{(n)}(x)]_x^{(n)}. \quad (2)$$

To obtain an approximate solution of the integral equation, one restricts oneself to a specific value of n in (1) or (2) instead of taking the limit.

© References: E. A. C. Paley and N. Wiener (1934), D. V. Widder (1939, 1971), I. I. Hirschman and D. V. Widder (1955), P. P. Zabreyko, A. I. Koshelev, et al. (1975), E. C. Titchmarsh (1986), Yu. A. Brychkov and A. P. Prudnikov (1989), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 428).

3.1-4. Kernels Containing Square Roots.

$$18. \int_0^a |\sqrt{x} - \sqrt{t}| y(t) dt = f(x), \quad 0 < a < \infty.$$

This is a special case of equation 3.8.3 with $g(x) = \sqrt{x}$.

Solution:

$$y(x) = \frac{d}{dx} [\sqrt{x} f'_x(x)].$$

The right-hand side $f(x)$ of the equation must satisfy certain conditions. The general form of the right-hand side is

$$f(x) = F(x) + Ax + B, \quad A = -F'_x(a), \quad B = \frac{1}{2} [aF'_x(a) - F(a) - F(0)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

$$19. \int_0^a |\sqrt{x} - \beta\sqrt{t}| y(t) dt = f(x), \quad \beta > 0.$$

This is a special case of equation 3.8.4 with $g(x) = \sqrt{x}$ and $\beta = \sqrt{\lambda}$.

$$20. \int_0^a |\sqrt{x} - t| y(t) dt = f(x).$$

This is a special case of equation 3.8.5 with $g(x) = \sqrt{x}$ (see item 3° of 3.8.5).

$$21. \int_0^a |x - \sqrt{t}| y(t) dt = f(x).$$

This is a special case of equation 3.8.6 with $g(t) = \sqrt{t}$ (see item 3° of 3.8.6).

$$22. \int_0^a \frac{y(t)}{\sqrt{|x-t|}} dt = f(x), \quad 0 < a \leq \infty.$$

This is a special case of equation 3.1.30 with $k = \frac{1}{2}$.

Solution:

$$y(x) = -\frac{A}{x^{1/4}} \frac{d}{dx} \left[\int_x^a \frac{dt}{(t-x)^{1/4}} \int_0^t \frac{f(s) ds}{s^{1/4}(t-s)^{1/4}} \right], \quad A = \frac{1}{\sqrt{8\pi} \Gamma^2(3/4)}.$$

$$23. \int_{-\infty}^{\infty} \frac{y(t)}{\sqrt{|x-t|}} dt = f(x).$$

This is a special case of equation 3.1.35 with $\lambda = \frac{1}{2}$.

Solution:

$$y(x) = \frac{1}{4\pi} \int_{-\infty}^{\infty} \frac{f(x) - f(t)}{|x-t|^{3/2}} dt.$$

$$24. \int_{-1}^1 \frac{y(t) dt}{\sqrt{1+x^2-2xt}} = f(x).$$

Solution:

$$y(x) = \frac{1}{2} \sum_{n=0}^{\infty} \frac{2n+1}{n!} f_x^{(n)}(0) P_n(x),$$

where $P_n(x)$ are the Legendre polynomials (see Supplement 11.11-1)

$$P_n(x) = \frac{1}{n! 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n.$$

3.1-5. Kernels Containing Arbitrary Powers.

$$25. \int_0^a |x^k - t^k| y(t) dt = f(x), \quad 0 < k < 1, \quad 0 < a < \infty.$$

1°. Let us remove the modulus in the integrand:

$$\int_0^x (x^k - t^k)y(t) dt + \int_x^a (t^k - x^k)y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x yields

$$kx^{k-1} \int_0^x y(t) dt - kx^{k-1} \int_x^a y(t) dt = f'_x(x). \quad (2)$$

Let us divide both sides of (2) by kx^{k-1} and differentiate the resulting equation. As a result, we obtain the solution

$$y(x) = \frac{1}{2k} \frac{d}{dx} [x^{1-k} f'_x(x)]. \quad (3)$$

2°. Let us demonstrate that the right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x=0$ and $x=a$, in (1), we obtain two corollaries $\int_0^a t^k y(t) dt = f(0)$ and $\int_0^a (a^k - t^k)y(t) dt = f(a)$, which can be rewritten in the form

$$\int_0^a t^k y(t) dt = f(0), \quad a^k \int_0^a y(t) dt = f(0) + f(a). \quad (4)$$

Substitute $y(x)$ of (3) into (4). Integrating by parts yields the relations $a f'_x(a) = k f(a) + k f(0)$ and $a f'_x(a) = 2k f(a) + 2k f(0)$. Hence, the desired constraints for $f(x)$ have the form

$$f(0) + f(a) = 0, \quad f'_x(a) = 0. \quad (5)$$

Conditions (5) make it possible to find the admissible general form of the right-hand side of the integral equation:

$$f(x) = F(x) + Ax + B, \quad A = -F'_x(a), \quad B = \frac{1}{2} [aF'_x(a) - F(a) - F(0)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function with bounded first derivative. The first derivative may be unbounded at $x = 0$, in which case the conditions $[x^{1-k} F'_x]_{x=0} = 0$ must hold.

$$26. \int_0^a |x^k - \beta t^k| y(t) dt = f(x), \quad 0 < k < 1, \quad \beta > 0.$$

This is a special case of equation 3.8.4 with $g(x) = x^k$ and $\beta = \lambda^k$.

$$27. \int_0^a |x^k t^m - t^{k+m}| y(t) dt = f(x), \quad 0 < k < 1, \quad 0 < a < \infty.$$

The substitution $w(t) = t^m y(t)$ leads to an equation of the form 3.1.25:

$$\int_0^a |x^k - t^k| w(t) dt = f(x).$$

$$28. \int_0^1 |x^k - t^m| y(t) dt = f(x), \quad k > 0, \quad m > 0.$$

The transformation

$$z = x^k, \quad \tau = t^m, \quad w(\tau) = \tau^{\frac{1-m}{m}} y(t)$$

leads to an equation of the form 3.1.1:

$$\int_0^1 |z - \tau| w(\tau) d\tau = F(z), \quad F(z) = m f(z^{1/k}).$$

$$29. \int_a^b |x - t|^{1+\lambda} y(t) dt = f(x), \quad 0 \leq \lambda < 1.$$

For $\lambda = 0$, see equation 3.1.2. Assume that $0 < \lambda < 1$.

1°. Let us remove the modulus in the integrand:

$$\int_a^x (x-t)^{1+\lambda} y(t) dt + \int_x^b (t-x)^{1+\lambda} y(t) dt = f(x). \quad (1)$$

Let us differentiate (1) with respect to x twice and then divide both the sides by $\lambda(\lambda + 1)$. As a result, we obtain

$$\int_a^x (x-t)^{\lambda-1} y(t) dt + \int_x^b (t-x)^{\lambda-1} y(t) dt = \frac{1}{\lambda(\lambda+1)} f''_{xx}(x). \quad (2)$$

Rewrite equation (2) in the form

$$\int_a^b \frac{y(t) dt}{|x-t|^k} = \frac{1}{\lambda(\lambda+1)} f''_{xx}(x), \quad k = 1 - \lambda. \quad (3)$$

See 3.1.30 and 3.1.31 for the solutions of equation (3) for various a and b .

2°. The right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = a$ and $x = b$ in (1), we obtain two corollaries

$$\int_a^b (t-a)^{1+\lambda} y(t) dt = f(a), \quad \int_a^b (b-t)^{1+\lambda} y(t) dt = f(b). \quad (4)$$

On substituting the solution $y(x)$ of (3) into (4) and then integrating by parts, we obtain the desired constraints for $f(x)$.

$$30. \int_0^a \frac{y(t)}{|x-t|^k} dt = f(x), \quad 0 < k < 1, \quad 0 < a \leq \infty.$$

1°. Solution:

$$y(x) = -Ax^{\frac{k-1}{2}} \frac{d}{dx} \left[\int_x^a \frac{t^{\frac{1-2k}{2}} dt}{(t-x)^{\frac{1-k}{2}}} \int_0^t \frac{f(s) ds}{s^{\frac{1-k}{2}} (t-s)^{\frac{1-k}{2}}} \right],$$

$$A = \frac{1}{2\pi} \cos\left(\frac{\pi k}{2}\right) \Gamma(k) \left[\Gamma\left(\frac{1+k}{2}\right) \right]^{-2},$$

where $\Gamma(k)$ is the gamma function.

2°. The transformation $x = z^2$, $t = \xi^2$, $w(\xi) = 2\xi y(t)$ leads to an equation of the form 3.1.32:

$$\int_0^{\sqrt{a}} \frac{w(\xi)}{|z^2 - \xi^2|^k} d\xi = f(z^2).$$

$$31. \int_a^b \frac{y(t)}{|x-t|^k} dt = f(x), \quad 0 < k < 1.$$

It is assumed that $|a| + |b| < \infty$. Solution:

$$y(x) = \frac{1}{2\pi} \cot\left(\frac{1}{2}\pi k\right) \frac{d}{dx} \int_a^x \frac{f(t) dt}{(x-t)^{1-k}} - \frac{1}{\pi^2} \cos^2\left(\frac{1}{2}\pi k\right) \int_a^x \frac{Z(t)F(t)}{(x-t)^{1-k}} dt,$$

where

$$Z(t) = (t-a)^{\frac{1+k}{2}} (b-t)^{\frac{1-k}{2}}, \quad F(t) = \frac{d}{dt} \left[\int_a^t \frac{d\tau}{(t-\tau)^k} \int_\tau^b \frac{f(s) ds}{Z(s)(s-\tau)^{1-k}} \right].$$

⊙ Reference: F. D. Gakhov (1977).

$$32. \int_0^a \frac{y(t)}{|x^2-t^2|^k} dt = f(x), \quad 0 < k < 1, \quad 0 < a \leq \infty.$$

Solution:

$$y(x) = -\frac{2\Gamma(k)\cos\left(\frac{1}{2}\pi k\right)}{\pi\left[\Gamma\left(\frac{1+k}{2}\right)\right]^2} x^{k-1} \frac{d}{dx} \int_x^a \frac{t^{2-2k} F(t) dt}{(t^2-x^2)^{\frac{1-k}{2}}}, \quad F(t) = \int_0^t \frac{s^k f(s) ds}{(t^2-s^2)^{\frac{1-k}{2}}}.$$

⊙ Reference: P. P. Zabreyko, A. I. Koshelev, et al. (1975).

$$33. \int_a^b \frac{y(t)}{|x^\lambda-t^\lambda|^k} dt = f(x), \quad 0 < k < 1, \quad \lambda > 0.$$

1°. The transformation

$$z = x^\lambda, \quad \tau = t^\lambda, \quad w(\tau) = \tau^{\frac{1-\lambda}{\lambda}} y(t)$$

leads to an equation of the form 3.1.31:

$$\int_A^B \frac{w(\tau)}{|z-\tau|^k} d\tau = F(z),$$

where $A = a^\lambda$, $B = b^\lambda$, $F(z) = \lambda f(z^{1/\lambda})$.

2°. Solution with $a = 0$:

$$y(x) = -Ax^{\frac{\lambda(k-1)}{2}} \frac{d}{dx} \left[\int_x^b \frac{t^{\frac{\lambda(3-2k)-2}{2}} dt}{(t^\lambda-x^\lambda)^{\frac{1-k}{2}}} \int_0^t \frac{s^{\frac{\lambda(k+1)-2}{2}} f(s) ds}{(t^\lambda-s^\lambda)^{\frac{1-k}{2}}} \right],$$

$$A = \frac{\lambda^2}{2\pi} \cos\left(\frac{\pi k}{2}\right) \Gamma(k) \left[\Gamma\left(\frac{1+k}{2}\right) \right]^{-2},$$

where $\Gamma(k)$ is the gamma function.

$$34. \int_0^1 \frac{y(t)}{|x^\lambda-t^m|^k} dt = f(x), \quad 0 < k < 1, \quad \lambda > 0, \quad m > 0.$$

The transformation

$$z = x^\lambda, \quad \tau = t^m, \quad w(\tau) = \tau^{\frac{1-m}{m}} y(t)$$

leads to an equation of the form 3.1.31:

$$\int_0^1 \frac{w(\tau)}{|z-\tau|^k} d\tau = F(z), \quad F(z) = m f(z^{1/\lambda}).$$

$$35. \int_{-\infty}^{\infty} \frac{y(t)}{|x-t|^{1-\lambda}} dt = f(x), \quad 0 < \operatorname{Re} \lambda < 1.$$

Solution:

$$\begin{aligned} y(x) &= \frac{\lambda}{2\pi} \tan\left(\frac{\pi\lambda}{2}\right) \int_{-\infty}^{\infty} \frac{f(x) - f(t)}{|x-t|^{1+\lambda}} dt \\ &= \frac{\lambda}{2\pi} \tan\left(\frac{\pi\lambda}{2}\right) \int_0^{\infty} \frac{2f(x) - f(x+t) - f(x-t)}{t^{1+\lambda}} dt. \end{aligned}$$

It is assumed that the condition $\int_{-\infty}^{\infty} |f(x)|^p dx < \infty$ is satisfied for some p , $1 < p < 1/\lambda$.

The integral equation and its solution form the *Riesz transform pair* (the *Riesz potential*).

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 428), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$36. \int_{-\infty}^{\infty} \frac{y(t)}{|x^3-t|^{1-\lambda}} dt = f(x), \quad 0 < \lambda < 1.$$

The substitution $z = x^3$ leads to an equation of the form 3.1.35:

$$\int_{-\infty}^{\infty} \frac{y(t)}{|z-t|^{1-\lambda}} dt = f(z^{1/3}).$$

$$37. \int_{-\infty}^{\infty} \frac{y(t)}{|x^3-t^3|^{1-\lambda}} dt = f(x), \quad 0 < \lambda < 1.$$

The transformation

$$z = x^3, \quad \tau = t^3, \quad w(\tau) = \tau^{-2/3} y(t)$$

leads to an equation of the form 3.1.35:

$$\int_{-\infty}^{\infty} \frac{w(\tau)}{|z-\tau|^{1-\lambda}} d\tau = F(z), \quad F(z) = 3f(z^{1/3}).$$

$$38. \int_{-\infty}^{\infty} \frac{\operatorname{sign}(x-t)}{|x-t|^{1-\lambda}} y(t) dt = f(x), \quad 0 < \operatorname{Re} \lambda < 1.$$

Solution:

$$\begin{aligned} y(x) &= \frac{\lambda}{2\pi} \cot\left(\frac{\pi\lambda}{2}\right) \int_{-\infty}^{\infty} \frac{f(x) - f(t)}{|x-t|^{1+\lambda}} \operatorname{sign}(x-t) dt \\ &= \frac{\lambda}{2\pi} \cot\left(\frac{\pi\lambda}{2}\right) \int_0^{\infty} \frac{f(x+t) - f(x-t)}{t^{1+\lambda}} dt \\ &= \frac{\lambda}{2\pi} \cot\left(\frac{\pi\lambda}{2}\right) \frac{d}{dx} \int_{-\infty}^{\infty} \frac{f(t)}{|x-t|^\lambda} dt. \end{aligned}$$

The integral equation and its solution form the *Feller transform pair* (the *Feller potential*).

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 428), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$39. \int_{-\infty}^{\infty} \frac{a + b \operatorname{sign}(x-t)}{|x-t|^{1-\lambda}} y(t) dt = f(x), \quad 0 < \operatorname{Re} \lambda < 1.$$

Solution:

$$\begin{aligned} y(x) &= C\lambda \int_{-\infty}^{\infty} \frac{a + b \operatorname{sign}(x-t)}{|x-t|^{1+\lambda}} [f(x) - f(t)] dt \\ &= C\lambda \int_0^{\infty} t^{-1-\lambda} [2af(x) - (a+b)f(x-t) - (a-b)f(x+t)] dt \\ &= C \frac{d}{dx} \int_{-\infty}^{\infty} \frac{b + a \operatorname{sign}(x-t)}{|x-t|^\lambda} f(t) dt, \end{aligned}$$

where

$$C = \frac{\sin(\pi\lambda)}{4\pi \left[a^2 \cos^2\left(\frac{1}{2}\pi\lambda\right) + b^2 \sin^2\left(\frac{1}{2}\pi\lambda\right) \right]}.$$

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 431), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

$$40. \int_0^\infty \frac{y(t) dt}{(ax + bt)^k} = f(x), \quad a > 0, \quad b > 0, \quad k > 0.$$

By setting

$$x = \frac{1}{2a}e^{2z}, \quad t = \frac{1}{2b}e^{2\tau}, \quad y(t) = be^{(k-2)\tau}w(\tau), \quad f(x) = e^{-kz}g(z),$$

we obtain an integral equation with the difference kernel of the form 3.8.15:

$$\int_{-\infty}^\infty \frac{w(\tau) d\tau}{\cosh^k(z - \tau)} = g(z).$$

$$41. \int_0^\infty t^{z-1}y(t) dt = f(z).$$

The left-hand side of this equation is the Mellin transform of $y(t)$ (z is treated as a complex variable).

Solution:

$$y(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} t^{-z} f(z) dz, \quad i^2 = -1.$$

For specific $f(z)$, one can use tables of Mellin and Laplace integral transforms to calculate the integral.

⊙ References: H. Bateman and A. Erdélyi (vol. 2, 1954), V. A. Ditkin and A. P. Prudnikov (1965).

3.1-6. Equations Containing the Unknown Function of a Complicated Argument.

$$42. \int_0^1 y(xt) dt = f(x).$$

Solution:

$$y(x) = x f'_x(x) + f(x).$$

The function $f(x)$ is assumed to satisfy the condition $[x f(x)]_{x=0} = 0$.

$$43. \int_0^1 t^\lambda y(xt) dt = f(x).$$

The substitution $\xi = xt$ leads to equation $\int_0^x \xi^\lambda y(\xi) d\xi = x^{\lambda+1} f(x)$. Differentiating with respect to x yields the solution

$$y(x) = x f'_x(x) + (\lambda + 1) f(x).$$

The function $f(x)$ is assumed to satisfy the condition $[x^{\lambda+1} f(x)]_{x=0} = 0$.

$$44. \int_0^1 (Ax^k + Bt^m)y(xt) dt = f(x).$$

The substitution $\xi = xt$ leads to an equation of the form 1.1.51:

$$\int_0^x (Ax^{k+m} + B\xi^m)y(\xi) d\xi = x^{m+1}f(x).$$

$$45. \int_0^1 \frac{y(xt) dt}{\sqrt{1-t}} = f(x).$$

The substitution $\xi = xt$ leads to Abel's equation 1.1.36:

$$\int_0^x \frac{y(\xi) d\xi}{\sqrt{x-\xi}} = \sqrt{x} f(x).$$

$$46. \int_0^1 \frac{y(xt) dt}{(1-t)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

The substitution $\xi = xt$ leads to the generalized Abel equation 1.1.47:

$$\int_0^x \frac{y(\xi) d\xi}{(x-\xi)^\lambda} = x^{1-\lambda} f(x).$$

$$47. \int_0^1 \frac{t^\mu y(xt) dt}{(1-t)^\lambda} = f(x), \quad 0 < \lambda < 1.$$

The transformation $\xi = xt$, $w(\xi) = \xi^\mu y(\xi)$ leads to the generalized Abel equation 1.1.47:

$$\int_0^x \frac{w(\xi) d\xi}{(x-\xi)^\lambda} = x^{1+\mu-\lambda} f(x).$$

$$48. \int_0^\infty \frac{y(x+t) - y(x-t)}{t} dt = f(x).$$

Solution:

$$y(x) = -\frac{1}{\pi^2} \int_0^\infty \frac{f(x+t) - f(x-t)}{t} dt.$$

⊙ References: V. A. Ditkin and A. P. Prudnikov (1965), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 427).

3.1-7. Singular Equations.

In this subsection, all singular integrals are understood in the sense of the Cauchy principal value.

$$49. \int_{-\infty}^\infty \frac{y(t) dt}{t-x} = f(x).$$

Solution:

$$y(x) = -\frac{1}{\pi^2} \int_{-\infty}^\infty \frac{f(t) dt}{t-x}.$$

The integral equation and its solution form a *Hilbert transform pair* (in the asymmetric form).

⊙ References: V. A. Ditkin and A. P. Prudnikov (1965), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 427).

$$50. \int_0^{\infty} \frac{y(t) dt}{t-x} = f(x).$$

Solution:

$$y(x) = -\frac{\sqrt{x}}{\pi^2} \int_0^{\infty} \frac{f(t)}{\sqrt{t}(t-x)} dt.$$

The integral equation and its solution form a *Hilbert transform pair on the semiaxis* (in the asymmetric form).

⊙ References: D. Hilbert (1953), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 427), I. K. Lifanov, L. N. Poltavskii, and G. M. Vainikko (2004, p. 8).

$$51. \int_a^b \frac{y(t) dt}{t-x} = f(x).$$

This equation is encountered in hydrodynamics in solving the problem on the flow of an ideal inviscid fluid around a thin profile ($a \leq x \leq b$). It is assumed that $|a| + |b| < \infty$.

1°. The solution bounded at the endpoints is

$$y(x) = -\frac{1}{\pi^2} \sqrt{(x-a)(b-x)} \int_a^b \frac{f(t)}{\sqrt{(t-a)(b-t)} t-x} dt,$$

provided that

$$\int_a^b \frac{f(t) dt}{\sqrt{(t-a)(b-t)}} = 0.$$

2°. The solution bounded at the endpoint $x = a$ and unbounded at the endpoint $x = b$ is

$$y(x) = -\frac{1}{\pi^2} \sqrt{\frac{x-a}{b-x}} \int_a^b \sqrt{\frac{b-t}{t-a}} \frac{f(t)}{t-x} dt.$$

3°. The solution unbounded at the endpoints is

$$y(x) = -\frac{1}{\pi^2 \sqrt{(x-a)(b-x)}} \left[\int_a^b \frac{\sqrt{(t-a)(b-t)}}{t-x} f(t) dt + C \right],$$

where C is an arbitrary constant. The formula $\int_a^b y(t) dt = C/\pi$ holds.

Solutions that have a singularity point $x = s$ inside the interval $[a, b]$ can be found in Subsection 14.4-3.

⊙ Reference: F. D. Gakhov (1977).

$$52. \int_{-1}^1 \left(\frac{1}{t-x} + \frac{1}{x+t+2} \right) y(t) dt = f(x), \quad -1 < x < 1.$$

Solution for $f(x) = \pi q = \text{const}$:

$$y(t) = q \frac{1+t}{\sqrt{(1-t)(3+t)}}.$$

⊙ Reference: H. F. Bueckner (1966).

$$53. \int_0^1 \left(\frac{1}{t-x} + \frac{\lambda}{t+x} \right) y(t) dt = f(x), \quad 0 < x < 1.$$

Solution for $f(x) = \pi q = \text{const}$:

$$y(x) = \frac{q}{2 \sin(\frac{1}{2}\pi\beta)} \left[\left(\frac{x}{1 + \sqrt{1-x^2}} \right)^\beta \left(\frac{\beta}{\sqrt{1-x^2}} + 1 \right) + \left(\frac{x}{1 + \sqrt{1-x^2}} \right)^{-\beta} \left(\frac{\beta}{\sqrt{1-x^2}} - 1 \right) \right],$$

where β is given by

$$\cos(\pi\beta) = -\lambda, \quad 0 < \beta < 1.$$

We assume that the following necessary condition holds

$$\int_0^1 y(t) dt = 0.$$

⊙ References: H. F. Bueckner (1966), P. S. Theocaris and N. I. Ioakimidis (1977).

$$54. \frac{1}{\pi i} \int_{-a}^a \left(\frac{1}{t-x} - \frac{\lambda x}{xt-a^2} \right) y(t) dt = f(x), \quad -a < x < a \quad (i^2 = -1).$$

1°. Solution:

$$y(x) = \left(\frac{a-x}{-a-x} \right)^\beta \frac{1}{2\pi i} \int_{-a}^a \left(\frac{a-t}{-a-t} \right)^{-\beta} \left(\frac{1}{t-x} - \frac{x}{xt-a^2} \right) f(t) dt \\ + \left(\frac{a-x}{-a-x} \right)^{-\beta} \frac{1}{2\pi i} \int_{-a}^a \left(\frac{a-t}{-a-t} \right)^\beta \left(\frac{1}{t-x} - \frac{x}{xt-a^2} \right) f(t) dt,$$

where $\lambda = \cos \theta$ and $\beta = 1 - \frac{\theta}{\pi}$. We assume that the following necessary condition holds

$$\frac{1}{2\pi i} \int_{-a}^a \left[e^{-\pi i \beta} \left(\frac{a-t}{-a-t} \right)^\beta - e^{\pi i \beta} \left(\frac{a-t}{-a-t} \right)^{-\beta} \right] \frac{f(t)}{t} dt = 0.$$

2°. Solution for $f(x) \equiv 0$:

$$y(x) = C_1 \Lambda_1(x) + C_2 \Lambda_2(x) + C_3 \Lambda_3(x),$$

where C_1 , C_2 , and C_3 are arbitrary constants, and

$$\Lambda_1(x) = (1 + \lambda) e^{i\pi\beta} \left(\frac{a-t}{-a-t} \right)^{1-\beta} + (1 - \lambda) e^{-i\pi\beta} \left(\frac{a-t}{-a-t} \right)^\beta, \\ \Lambda_2(x) = (1 + \lambda) e^{-i\pi\beta} \left(\frac{a-t}{-a-t} \right)^{-1+\beta} + (1 - \lambda) e^{i\pi\beta} \left(\frac{a-t}{-a-t} \right)^{-\beta}, \\ \Lambda_3(x) = e^{i\pi\beta} \left(\frac{a-t}{-a-t} \right)^{1-\beta} + e^{-i\pi\beta} \left(\frac{a-t}{-a-t} \right)^{-1+\beta}.$$

⊙ Reference: D. I. Sherman (1969).

$$55. \int_a^b \frac{y(t)}{(x-t)^2} dt = f(x), \quad a \leq x \leq b.$$

The simple hypersingular equation of the first kind with Cauchy-type kernel. This equation governs circulation-free flow of an ideal incompressible fluid past the segment $[a, b]$.

Let the conditions $y(a) = y(b) = 0$ be satisfied. Then the solution is

$$y(x) = \frac{1}{\pi^2} \int_a^b \ln \left| \frac{\sqrt{(b-t)(x-a)} - \sqrt{(b-x)(t-a)}}{\sqrt{(b-t)(x-a)} + \sqrt{(b-x)(t-a)}} \right| f'_t(t) dt.$$

This equation is discussed in Subsection 14.6-3 in detail.

⊙ Reference: I. K. Lifanov, L. N. Poltavskii, and G. M. Vainikko (2004, p. 7).

$$56. \frac{1}{\pi^2} \int_{-1}^1 \int_{-1}^1 \frac{u(x, y) dx dy}{(x_0 - x)(y_0 - y)} = f(x_0, y_0).$$

A two-dimensional singular equation.

A solution, which is bounded on the lines $x = \pm 1$ and $y = \pm 1$ but which is unbounded on the line $x = q$ ($-1 < q < 1$), is given by the formula

$$u(x_0, y_0) = \frac{\sqrt{(1-x_0^2)(1-y_0^2)}}{\pi^2} \int_{-1}^1 \int_{-1}^1 \frac{f(x, y) dx dy}{\sqrt{(1-x^2)(1-y^2)}(x-x_0)(y-y_0)} \\ - \frac{\sqrt{(1-x_0^2)(1-y_0^2)}}{\pi^2(q-x_0)} \int_{-1}^1 \frac{dx}{\sqrt{1-x^2}} \left(\frac{1}{\pi^2} \int_{-1}^1 \frac{f(x, y) dy}{\sqrt{1-y^2}(y-y_0)} \right),$$

provided that

$$\int_{-1}^1 \frac{f(x_0, y) dy}{\sqrt{1-y^2}} = 0, \quad -1 \leq x_0 \leq 1.$$

⊙ Reference: I. K. Lifanov, L. N. Poltavskii, and G. M. Vainikko (2004, pp. 16–20).

3.2. Equations Whose Kernels Contain Exponential Functions

3.2-1. Kernels Containing Exponential Functions of the Form $e^{\lambda|x-t|}$.

$$1. \int_{-\infty}^{\infty} e^{-\lambda|x-t|} y(t) dt = f(x), \quad f(\pm\infty) = 0.$$

Solution:

$$y(x) = \frac{1}{2\lambda} [\lambda^2 f(x) - f''_{xx}(x)].$$

⊙ References: I. I. Hirschman and D. V. Widder (1955), F. D. Gakhov and Yu. I. Cherskii (1978), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 433).

$$2. \int_0^{\infty} e^{-\lambda|x-t|} y(t) dt = f(x), \quad f(\infty) = 0.$$

1°. Solution:

$$y(x) = \frac{1}{2\lambda} e^{-\lambda x} \frac{d}{dx} e^{2\lambda x} \frac{d}{dx} e^{-\lambda x} f(x).$$

2°. If $f'_x(0) - \lambda f(0) = 0$ then

$$y(x) = \frac{1}{2\lambda} [\lambda^2 f(x) - f''_{xx}(x)].$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 433).

$$3. \int_a^b e^{\lambda|x-t|} y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

1°. Let us remove the modulus in the integrand:

$$\int_a^x e^{\lambda(x-t)} y(t) dt + \int_x^b e^{\lambda(t-x)} y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x twice yields

$$2\lambda y(x) + \lambda^2 \int_a^x e^{\lambda(x-t)} y(t) dt + \lambda^2 \int_x^b e^{\lambda(t-x)} y(t) dt = f''_{xx}(x). \quad (2)$$

By eliminating the integral terms from (1) and (2), we obtain the solution

$$y(x) = \frac{1}{2\lambda} [f''_{xx}(x) - \lambda^2 f(x)]. \quad (3)$$

2°. The right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = a$ and $x = b$ in (1), we obtain two corollaries

$$\int_a^b e^{\lambda t} y(t) dt = e^{\lambda a} f(a), \quad \int_a^b e^{-\lambda t} y(t) dt = e^{-\lambda b} f(b). \quad (4)$$

On substituting the solution $y(x)$ of (3) into (4) and then integrating by parts, we see that

$$\begin{aligned} e^{\lambda b} f'_x(b) - e^{\lambda a} f'_x(a) &= \lambda e^{\lambda a} f(a) + \lambda e^{\lambda b} f(b), \\ e^{-\lambda b} f'_x(b) - e^{-\lambda a} f'_x(a) &= \lambda e^{-\lambda a} f(a) + \lambda e^{-\lambda b} f(b). \end{aligned}$$

Hence, we obtain the desired constraints for $f(x)$:

$$f'_x(a) + \lambda f(a) = 0, \quad f'_x(b) - \lambda f(b) = 0. \quad (5)$$

The general form of the right-hand side satisfying conditions (5) is given by

$$f(x) = F(x) + Ax + B,$$

$$A = \frac{1}{b\lambda - a\lambda - 2} [F'_x(a) + F'_x(b) + \lambda F(a) - \lambda F(b)], \quad B = -\frac{1}{\lambda} [F'_x(a) + \lambda F(a) + Aa\lambda + A],$$

where $F(x)$ is an arbitrary bounded, twice differentiable function.

$$4. \int_a^b (Ae^{\lambda|x-t|} + Be^{\mu|x-t|}) y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

Let us remove the modulus in the integrand and differentiate the resulting equation with respect to x twice to obtain

$$2(A\lambda + B\mu)y(x) + \int_a^b (A\lambda^2 e^{\lambda|x-t|} + B\mu^2 e^{\mu|x-t|}) y(t) dt = f''_{xx}(x). \quad (1)$$

Eliminating the integral term with $e^{\mu|x-t|}$ from (1) with the aid of the original integral equation, we find that

$$2(A\lambda + B\mu)y(x) + A(\lambda^2 - \mu^2) \int_a^b e^{\lambda|x-t|} y(t) dt = f''_{xx}(x) - \mu^2 f(x). \quad (2)$$

For $A\lambda + B\mu = 0$, this is an equation of the form 3.2.3, and for $A\lambda + B\mu \neq 0$, this is an equation of the form 4.2.15.

The right-hand side $f(x)$ must satisfy certain relations, which can be obtained by setting $x = a$ and $x = b$ in the original equation (a similar procedure is used in 3.2.3).

$$5. \int_a^b \left[\sum_{k=1}^n A_k \exp(\lambda_k |x-t|) \right] y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

1°. Let us remove the modulus in the k th summand of the integrand:

$$I_k(x) = \int_a^b \exp(\lambda_k |x-t|) y(t) dt = \int_a^x \exp[\lambda_k(x-t)] y(t) dt + \int_x^b \exp[\lambda_k(t-x)] y(t) dt. \quad (1)$$

Differentiating (1) with respect to x twice yields

$$\begin{aligned} I_k' &= \lambda_k \int_a^x \exp[\lambda_k(x-t)] y(t) dt - \lambda_k \int_x^b \exp[\lambda_k(t-x)] y(t) dt, \\ I_k'' &= 2\lambda_k y(x) + \lambda_k^2 \int_a^x \exp[\lambda_k(x-t)] y(t) dt + \lambda_k^2 \int_x^b \exp[\lambda_k(t-x)] y(t) dt, \end{aligned} \quad (2)$$

where the primes denote the derivatives with respect to x . By comparing formulas (1) and (2), we find the relation between I_k'' and I_k :

$$I_k'' = 2\lambda_k y(x) + \lambda_k^2 I_k, \quad I_k = I_k(x). \quad (3)$$

2°. With the aid of (1), the integral equation can be rewritten in the form

$$\sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x twice and taking into account (3), we obtain

$$\sigma_1 y(x) + \sum_{k=1}^n A_k \lambda_k^2 I_k = f''_{xx}(x), \quad \sigma_1 = 2 \sum_{k=1}^n A_k \lambda_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5) yields

$$\sigma_1 y(x) + \sum_{k=1}^{n-1} A_k (\lambda_k^2 - \lambda_n^2) I_k = f''_{xx}(x) - \lambda_n^2 f(x). \quad (6)$$

Differentiating (6) with respect to x twice and eliminating I_{n-1} from the resulting equation with the aid of (6), we obtain a similar equation whose right-hand side is a second-order linear differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} B_k I_k$. If we successively eliminate $I_{n-2}, I_{n-3}, \dots, I_1$ with the aid of double differentiation, then we finally arrive at a linear nonhomogeneous ordinary differential equation of order $2(n-1)$ with constant coefficients.

3°. The right-hand side $f(x)$ must satisfy certain conditions. To find these conditions, one must set $x = a$ in the integral equation and its derivatives. (Alternatively, these conditions can be found by setting $x = a$ and $x = b$ in the integral equation and all its derivatives obtained by means of double differentiation.)

3.2-2. Kernels Containing Exponential Functions of the Forms $e^{\lambda x}$ and $e^{\mu t}$.

$$6. \quad \int_a^b |e^{\lambda x} - e^{\lambda t}| y(t) dt = f(x), \quad \lambda > 0.$$

This is a special case of equation 3.8.3 with $g(x) = e^{\lambda x}$.

Solution:

$$y(x) = \frac{1}{2\lambda} \frac{d}{dx} [e^{-\lambda x} f'_x(x)].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$7. \quad \int_0^a |e^{\beta x} - e^{\mu t}| y(t) dt = f(x), \quad \beta > 0, \quad \mu > 0.$$

This is a special case of equation 3.8.4 with $g(x) = e^{\beta x}$ and $\lambda = \mu/\beta$.

$$8. \quad \int_a^b \frac{y(t) dt}{|e^{\lambda x} - e^{\lambda t}|^k} = f(x), \quad 0 < k < 1.$$

The transformation $z = e^{\lambda x}$, $\tau = e^{\lambda t}$, $w(\tau) = e^{-\lambda t} y(t)$ leads to an equation of the form 3.1.31:

$$\int_A^B \frac{w(\tau) d\tau}{|z - \tau|^k} = F(z),$$

where $A = e^{\lambda a}$, $B = e^{\lambda b}$, $F(z) = \lambda f\left(\frac{1}{\lambda} \ln z\right)$.

$$9. \quad \int_0^\infty \frac{y(t) dt}{(e^{\lambda x} + e^{\lambda t})^k} = f(x), \quad \lambda > 0, \quad k > 0.$$

This equation can be rewritten as an equation with difference kernel in the form 3.8.16:

$$\int_0^\infty \frac{w(t) dt}{\cosh^k\left[\frac{1}{2}\lambda(x-t)\right]} = g(x),$$

where $w(t) = 2^{-k} \exp(-\frac{1}{2}\lambda kt) y(t)$ and $g(x) = \exp(\frac{1}{2}\lambda kx) f(x)$.

3.2-3. Kernels Containing Exponential Functions of the Form $e^{\lambda xt}$.

$$10. \quad \int_{-\infty}^\infty e^{-xt} y(t) dt = f(x).$$

Solution:

$$\begin{aligned} y(t) &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{st} f(s) ds \\ &= \frac{1}{\sqrt{2\pi^3}} \int_0^\infty e^{-\xi^2/2} d\xi \int_{-\infty}^\infty e^{-x^2/2} \cos(\xi(x+t)) f(x) dx. \end{aligned}$$

The integral equation and its solution form a *two-side Laplace transform pair*.

⊙ References: B. Van der Pol and H. Bremmer (1955), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 433).

$$11. \int_{-\infty}^{\infty} e^{\lambda xt} y(t) dt = f(x), \quad \lambda \neq 0.$$

1°. The transformation

$$x = -\frac{1}{\lambda}z, \quad f(x) = F(z)$$

leads to an equation of the form 3.2.10:

$$\int_{-\infty}^{\infty} e^{-zt} y(t) dt = F(z).$$

2°. The transformation

$$y(t) = \exp(-t^2)Y(t), \quad x = \frac{2}{\lambda}\zeta, \quad f(x) = \exp(\zeta^2)\Phi(\zeta)$$

leads to an equation of the form 3.2.17:

$$\int_{-\infty}^{\infty} e^{-(\zeta-t)^2} Y(t) dt = \Phi(\zeta).$$

$$12. \int_{-\infty}^{\infty} e^{-ixt} y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ixt} f(x) dx.$$

Up to constant factors, the function $f(x)$ and the solution $y(t)$ are the *Fourier transform pair*.

⊙ References: V. A. Ditkin and A. P. Prudnikov (1965), J. W. Miles (1971), B. Davis (1978), F. Oberhettinger (1980), Yu. A. Brychkov and A. P. Prudnikov (1989), W. H. Beyer (1991), I. Sneddon (1995), A. Pinkus and S. Zafrany (1997), R. Bracewell (1999), A. D. Poularikas (2000), R. J. Beerends, H. G. ter Morschem, and J. C. van den Berg (2003), L. Debnath and D. Bhatta (2007).

$$13. \int_0^{\infty} e^{-zt} y(t) dt = f(z).$$

The left-hand side of the equation is the *Laplace transform* of $y(t)$ (z is treated as a complex variable).

1°. Solution:

$$y(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{zt} f(z) dz, \quad i^2 = -1.$$

For specific functions $f(z)$, one may use tables of inverse Laplace transforms to calculate the integral (e.g., see Supplement 6).

2°. For real $z = x$, under some assumptions the solution of the original equation can be represented in the form

$$y(x) = \lim_{n \rightarrow \infty} \frac{(-1)^n}{n!} \left(\frac{n}{x}\right)^{n+1} f_x^{(n)}\left(\frac{n}{x}\right),$$

which is the real inversion of the Laplace transform. To calculate the solution approximately, one should restrict oneself to a specific value of n in this formula instead of taking the limit.

⊙ References: G. Doetsch (1950, 1956, 1958, 1974), H. Bateman and A. Erdélyi (vol. 1, 1954), I. I. Hirschman and D. V. Widder (1955), V. A. Ditkin and A. P. Prudnikov (1965), J. W. Miles (1971), F. Oberhettinger (1973), B. Davis (1978), W. R. LePage (1980), R. Bellman and R. Roth (1984), Yu. A. Brychkov and A. P. Prudnikov (1989), W. H. Beyer (1991), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, Vols 4 and 5), R. J. Beerends, H. G. ter Morschem, and J. C. van den Berg (2003).

3.2-4. Kernels Containing Power-Law and Exponential Functions.
--

14. $\int_0^a |ke^{\lambda x} - k - t| y(t) dt = f(x).$

This is a special case of equation 3.8.5 with $g(x) = ke^{\lambda x} - k$.

15. $\int_0^a |x - ke^{\lambda t} - k| y(t) dt = f(x).$

This is a special case of equation 3.8.6 with $g(t) = ke^{\lambda t} + k$.

16. $\int_{-\infty}^{\infty} t^{-ix-1/2} \exp\left(\frac{2x-i}{4}\pi\right) y(t) dt = f(x), \quad i^2 = -1.$

Solution:

$$y(x) = \frac{1}{4\pi} \int_{-\infty}^{\infty} x^{it-1/2} \exp\left(\frac{2t+i}{4}\pi\right) \frac{f(t)}{\cosh(\pi t)} dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 463).

3.2-5. Kernels Containing Exponential Functions of the Form $e^{\lambda(x \pm t)^2}$.
--

17. $\int_{-\infty}^{\infty} e^{-(x-t)^2} y(t) dt = f(x).$

1°. The transformation

$$Y(t) = \exp(-t^2)y(t), \quad z = -2x, \quad F(z) = \exp(x^2)f(x)$$

leads to an equation of the form 3.2.10:

$$\int_{-\infty}^{\infty} e^{-zt}Y(t) dt = F(z).$$

2°. Solution:

$$\begin{aligned} y(t) &= \frac{1}{\pi^{3/2}} \int_0^{\infty} e^{s^2/4} ds \int_{-\infty}^{\infty} \cos(s(t-x)) f(x) dx \\ &= \exp\left[-\frac{1}{4\sqrt{\pi}} \frac{d^2}{dt^2} f(t)\right] \equiv \sum_{k=0}^{\infty} \frac{1}{k!} \left(-\frac{1}{4\sqrt{\pi}}\right)^k \frac{d^{2k} f(t)}{dt^{2k}}. \end{aligned}$$

(See equation 3.2.18 for $\lambda = 1$.)

3°. Solution:

$$y(x) = \frac{1}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{f_x^{(n)}(0)}{2^n n!} H_n(x),$$

where $H_n(x)$ are the Hermite polynomials (see Supplement 11.17-3)

$$H_m(x) = (-1)^m \exp(x^2) \frac{d^m}{dx^m} \exp(-x^2).$$

⊙ References: P. M. Morse and H. Feshbach (1953), I. I. Hirschman and D. V. Widder (1955), P. G. Rooney (1963), M. L. Krasnov (1975).

$$18. \quad \frac{1}{\sqrt{\pi\lambda}} \int_{-\infty}^{\infty} \exp\left[-\frac{(x-t)^2}{\lambda}\right] y(t) dt = f(x).$$

It is the *Gauss transform* (the *Weierstrass transform* for $\lambda = 4$).

Solution:

$$\begin{aligned} y(t) &= \frac{1}{\pi} \int_0^{\infty} e^{\lambda s^2/4} ds \int_{-\infty}^{\infty} \cos(s(t-x)) f(x) dx \\ &= \exp\left[-\frac{\lambda}{4} \frac{d^2}{dt^2} f(t)\right] \equiv \sum_{k=0}^{\infty} \frac{1}{k!} \left(-\frac{\lambda}{4}\right)^k \frac{d^{2k} f(t)}{dt^{2k}}. \end{aligned}$$

⊙ References: I. I. Hirschman and D. V. Widder (1955), P. G. Rooney (1963), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 435).

$$19. \quad \int_{-\infty}^{\infty} e^{i(x+t)^2} y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(x) = \frac{1}{\pi} \int_{-\infty}^{\infty} e^{-i(x+t)^2} f(t) dt.$$

⊙ References: E. A. C. Paley and N. Wiener (1934), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 435).

3.2-6. Other Kernels.

$$20. \quad \int_a^b |\exp(\lambda x^2) - \exp(\lambda t^2)| y(t) dt = f(x), \quad \lambda > 0.$$

This is a special case of equation 3.8.3 with $g(x) = \exp(\lambda x^2)$.

Solution:

$$y(x) = \frac{1}{4\lambda} \frac{d}{dx} \left[\frac{1}{x} \exp(-\lambda x^2) f'_x(x) \right].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$21. \quad \frac{1}{\sqrt{\pi x}} \int_0^{\infty} \exp\left(-\frac{t^2}{4x}\right) y(t) dt = f(x).$$

Applying the Laplace transformation to the equation, we obtain

$$\frac{\tilde{y}(\sqrt{p})}{\sqrt{p}} = \tilde{f}(p), \quad \tilde{f}(p) = \int_0^{\infty} e^{-pt} f(t) dt.$$

Substituting p by p^2 and solving for the transform \tilde{y} , we find that $\tilde{y}(p) = p\tilde{f}(p^2)$. The inverse Laplace transform provides the solution of the original integral equation:

$$y(t) = \mathfrak{L}^{-1}\{p\tilde{f}(p^2)\}, \quad \mathfrak{L}^{-1}\{g(p)\} \equiv \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{pt} g(p) dp.$$

3.3. Equations Whose Kernels Contain Hyperbolic Functions

3.3-1. Kernels Containing Hyperbolic Cosine.

$$1. \int_a^b |\cosh(\lambda x) - \cosh(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \cosh(\lambda x)$.

Solution:

$$y(x) = \frac{1}{2\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sinh(\lambda x)} \right].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$2. \int_0^a |\cosh(\beta x) - \cosh(\mu t)| y(t) dt = f(x), \quad \beta > 0, \quad \mu > 0.$$

This is a special case of equation 3.8.4 with $g(x) = \cosh(\beta x)$ and $\lambda = \mu/\beta$.

$$3. \int_a^b |\cosh^k x - \cosh^k t| y(t) dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \cosh^k x$.

Solution:

$$y(x) = \frac{1}{2k} \frac{d}{dx} \left[\frac{f'_x(x)}{\sinh x \cosh^{k-1} x} \right].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$4. \int_a^b \frac{y(t)}{|\cosh(\lambda x) - \cosh(\lambda t)|^k} dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.7 with $g(x) = \cosh(\lambda x) + \beta$, where β is an arbitrary number.

3.3-2. Kernels Containing Hyperbolic Sine.

$$5. \int_a^b \sinh(\lambda|x-t|) y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

1°. Let us remove the modulus in the integrand:

$$\int_a^x \sinh[\lambda(x-t)] y(t) dt + \int_x^b \sinh[\lambda(t-x)] y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x twice yields

$$2\lambda y(x) + \lambda^2 \int_a^x \sinh[\lambda(x-t)] y(t) dt + \lambda^2 \int_x^b \sinh[\lambda(t-x)] y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral terms from (1) and (2), we obtain the solution

$$y(x) = \frac{1}{2\lambda} [f''_{xx}(x) - \lambda^2 f(x)]. \quad (3)$$

2°. The right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = a$ and $x = b$ in (1), we obtain two corollaries

$$\int_a^b \sinh[\lambda(t-a)]y(t) dt = f(a), \quad \int_a^b \sinh[\lambda(b-t)]y(t) dt = f(b). \quad (4)$$

Substituting solution (3) into (4) and integrating by parts yields the desired conditions for $f(x)$:

$$\begin{aligned} \sinh[\lambda(b-a)]f'_x(b) - \lambda \cosh[\lambda(b-a)]f(b) &= \lambda f(a), \\ \sinh[\lambda(b-a)]f'_x(a) + \lambda \cosh[\lambda(b-a)]f(a) &= -\lambda f(b). \end{aligned} \quad (5)$$

The general form of the right-hand side is given by

$$f(x) = F(x) + Ax + B, \quad (6)$$

where $F(x)$ is an arbitrary bounded twice differentiable function, and the coefficients A and B are expressed in terms of $F(a)$, $F(b)$, $F'_x(a)$, and $F'_x(b)$ and can be determined by substituting formula (6) into conditions (5).

$$6. \quad \int_a^b \left\{ A \sinh(\lambda|x-t|) + B \sinh(\mu|x-t|) \right\} y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

Let us remove the modulus in the integrand and differentiate the equation with respect to x twice to obtain

$$2(A\lambda + B\mu)y(x) + \int_a^b \left\{ A\lambda^2 \sinh(\lambda|x-t|) + B\mu^2 \sinh(\mu|x-t|) \right\} y(t) dt = f''_{xx}(x). \quad (1)$$

Eliminating the integral term with $\sinh(\mu|x-t|)$ from (1) yields

$$2(A\lambda + B\mu)y(x) + A(\lambda^2 - \mu^2) \int_a^b \sinh(\lambda|x-t|)y(t) dt = f''_{xx}(x) - \mu^2 f(x). \quad (2)$$

For $A\lambda + B\mu = 0$, this is an equation of the form 3.3.5, and for $A\lambda + B\mu \neq 0$, this is an equation of the form 4.3.26.

The right-hand side $f(x)$ must satisfy certain relations, which can be obtained by setting $x = a$ and $x = b$ in the original equation (a similar procedure is used in 3.3.5).

$$7. \quad \int_a^b |\sinh(\lambda x) - \sinh(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \sinh(\lambda x)$.

Solution:

$$y(x) = \frac{1}{2\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\cosh(\lambda x)} \right].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$8. \int_0^a |\sinh(\beta x) - \sinh(\mu t)| y(t) dt = f(x), \quad \beta > 0, \quad \mu > 0.$$

This is a special case of equation 3.8.4 with $g(x) = \sinh(\beta x)$ and $\lambda = \mu/\beta$.

$$9. \int_a^b \sinh^3(\lambda|x-t|) y(t) dt = f(x).$$

Using the formula $\sinh^3 \beta = \frac{1}{4} \sinh 3\beta - \frac{3}{4} \sinh \beta$, we arrive at an equation of the form 3.3.6:

$$\int_a^b \left[\frac{1}{4} A \sinh(3\lambda|x-t|) - \frac{3}{4} A \sinh(\lambda|x-t|) \right] y(t) dt = f(x).$$

$$10. \int_a^b \left[\sum_{k=1}^n A_k \sinh(\lambda_k|x-t|) \right] y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

1°. Let us remove the modulus in the k th summand of the integrand:

$$I_k(x) = \int_a^b \sinh(\lambda_k|x-t|) y(t) dt = \int_a^x \sinh[\lambda_k(x-t)] y(t) dt + \int_x^b \sinh[\lambda_k(t-x)] y(t) dt. \quad (1)$$

Differentiating (1) with respect to x twice yields

$$I_k' = \lambda_k \int_a^x \cosh[\lambda_k(x-t)] y(t) dt - \lambda_k \int_x^b \cosh[\lambda_k(t-x)] y(t) dt, \quad (2)$$

$$I_k'' = 2\lambda_k y(x) + \lambda_k^2 \int_a^x \sinh[\lambda_k(x-t)] y(t) dt + \lambda_k^2 \int_x^b \sinh[\lambda_k(t-x)] y(t) dt,$$

where the primes denote the derivatives with respect to x . By comparing formulas (1) and (2), we find the relation between I_k'' and I_k :

$$I_k'' = 2\lambda_k y(x) + \lambda_k^2 I_k, \quad I_k = I_k(x). \quad (3)$$

2°. With the aid of (1), the integral equation can be rewritten in the form

$$\sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x twice and taking into account (3), we find that

$$\sigma_1 y(x) + \sum_{k=1}^n A_k \lambda_k^2 I_k = f_{xx}(x), \quad \sigma_1 = 2 \sum_{k=1}^n A_k \lambda_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5) yields

$$\sigma_1 y(x) + \sum_{k=1}^{n-1} A_k (\lambda_k^2 - \lambda_n^2) I_k = f_{xx}(x) - \lambda_n^2 f(x). \quad (6)$$

Differentiating (6) with respect to x twice and eliminating I_{n-1} from the resulting equation with the aid of (6), we obtain a similar equation whose right-hand side is a second-order linear differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} B_k I_k$.

If we successively eliminate I_{n-2}, I_{n-3}, \dots , with the aid of double differentiation, then we finally arrive at a linear nonhomogeneous ordinary differential equation of order $2(n-1)$ with constant coefficients.

3°. The right-hand side $f(x)$ must satisfy certain conditions. To find these conditions, one should set $x = a$ in the integral equation and its derivatives. (Alternatively, these conditions can be found by setting $x = a$ and $x = b$ in the integral equation and all its derivatives obtained by means of double differentiation.)

$$11. \int_0^b |\sinh^k x - \sinh^k t| y(t) dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \sinh^k x$.

Solution:

$$y(x) = \frac{1}{2k} \frac{d}{dx} \left[\frac{f'_x(x)}{\cosh x \sinh^{k-1} x} \right].$$

The right-hand side $f(x)$ must satisfy certain conditions. As follows from item 3° of equation 3.8.3, the admissible general form of the right-hand side is given by

$$f(x) = F(x) + Ax + B, \quad A = -F'_x(b), \quad B = \frac{1}{2} [bF'_x(b) - F(0) - F(b)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

$$12. \int_a^b \frac{y(t)}{|\sinh(\lambda x) - \sinh(\lambda t)|^k} dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.7 with $g(x) = \sinh(\lambda x) + \beta$, where β is an arbitrary number.

$$13. \int_0^a |k \sinh(\lambda x) - t| y(t) dt = f(x).$$

This is a special case of equation 3.8.5 with $g(x) = k \sinh(\lambda x)$.

$$14. \int_0^a |x - k \sinh(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.6 with $g(x) = k \sinh(\lambda t)$.

3.3-3. Kernels Containing Hyperbolic Tangent.

$$15. \int_a^b |\tanh(\lambda x) - \tanh(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \tanh(\lambda x)$.

Solution:

$$y(x) = \frac{1}{2\lambda} \frac{d}{dx} [\cosh^2(\lambda x) f'_x(x)].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$16. \int_0^a |\tanh(\beta x) - \tanh(\mu t)| y(t) dt = f(x), \quad \beta > 0, \quad \mu > 0.$$

This is a special case of equation 3.8.4 with $g(x) = \tanh(\beta x)$ and $\lambda = \mu/\beta$.

$$17. \int_0^b |\tanh^k x - \tanh^k t| y(t) dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \tanh^k x$.

Solution:

$$y(x) = \frac{1}{2k} \frac{d}{dx} [\cosh^2 x \coth^{k-1} x f'_x(x)].$$

The right-hand side $f(x)$ must satisfy certain conditions. As follows from item 3° of equation 3.8.3, the admissible general form of the right-hand side is given by

$$f(x) = F(x) + Ax + B, \quad A = -F'_x(b), \quad B = \frac{1}{2} [bF'_x(b) - F(0) - F(b)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

$$18. \int_a^b \frac{y(t)}{|\tanh(\lambda x) - \tanh(\lambda t)|^k} dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.7 with $g(x) = \tanh(\lambda x) + \beta$, where β is an arbitrary number.

$$19. \int_0^a |k \tanh(\lambda x) - t| y(t) dt = f(x).$$

This is a special case of equation 3.8.5 with $g(x) = k \tanh(\lambda x)$.

$$20. \int_0^a |x - k \tanh(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.6 with $g(x) = k \tanh(\lambda t)$.

3.3-4. Kernels Containing Hyperbolic Cotangent.

$$21. \int_a^b |\coth(\lambda x) - \coth(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \coth(\lambda x)$.

$$22. \int_0^b |\coth^k x - \coth^k t| y(t) dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \coth^k x$.

3.4. Equations Whose Kernels Contain Logarithmic Functions

3.4-1. Kernels Containing Logarithmic Functions.

$$1. \int_a^b |\ln(x/t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \ln x$.

Solution:

$$y(x) = \frac{1}{2} \frac{d}{dx} [x f'_x(x)].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$2. \quad \int_a^b \ln|x-t| y(t) dt = f(x).$$

Carleman's equation.

1°. Solution with $b-a \neq 4$:

$$y(x) = \frac{1}{\pi^2 \sqrt{(x-a)(b-x)}} \left[\int_a^b \frac{\sqrt{(t-a)(b-t)} f'_t(t) dt}{t-x} + \frac{1}{\ln\left[\frac{1}{4}(b-a)\right]} \int_a^b \frac{f(t) dt}{\sqrt{(t-a)(b-t)}} \right].$$

2°. If $b-a = 4$, then for the equation to be solvable, the condition

$$\int_a^b f(t)(t-a)^{-1/2}(b-t)^{-1/2} dt = 0$$

must be satisfied. In this case, the solution has the form

$$y(x) = \frac{1}{\pi^2 \sqrt{(x-a)(b-x)}} \left[\int_a^b \frac{\sqrt{(t-a)(b-t)} f'_t(t) dt}{t-x} + C \right],$$

where C is an arbitrary constant.

⊙ Reference: F. D. Gakhov (1977).

$$3. \quad \int_a^b (\ln|x-t| + \beta) y(t) dt = f(x).$$

By setting

$$x = e^{-\beta} z, \quad t = e^{-\beta} \tau, \quad y(t) = Y(\tau), \quad f(x) = e^{-\beta} g(z),$$

we arrive at an equation of the form 3.4.2:

$$\int_A^B \ln|z-\tau| Y(\tau) d\tau = g(z), \quad A = ae^\beta, \quad B = be^\beta.$$

$$4. \quad \int_{-a}^a \left(\ln \frac{A}{|x-t|} \right) y(t) dt = f(x), \quad -a \leq x \leq a.$$

This is a special case of equation 3.4.3 with $b = -a$. Solution with $0 < a < 2A$:

$$\begin{aligned} y(x) = & \frac{1}{2M'(a)} \left[\frac{d}{da} \int_{-a}^a w(t, a) f(t) dt \right] w(x, a) \\ & - \frac{1}{2} \int_{|x|}^a w(x, \xi) \frac{d}{d\xi} \left[\frac{1}{M'(\xi)} \frac{d}{d\xi} \int_{-\xi}^{\xi} w(t, \xi) f(t) dt \right] d\xi \\ & - \frac{1}{2} \frac{d}{dx} \int_{|x|}^a \frac{w(x, \xi)}{M'(\xi)} \left[\int_{-\xi}^{\xi} w(t, \xi) df(t) \right] d\xi, \end{aligned}$$

where

$$M(\xi) = \left(\ln \frac{2A}{\xi} \right)^{-1}, \quad w(x, \xi) = \frac{M(\xi)}{\pi \sqrt{\xi^2 - x^2}},$$

and the prime stands for the derivative.

⊙ Reference: I. C. Gohberg and M. G. Krein (1967).

$$5. \int_0^a \ln \left| \frac{x+t}{x-t} \right| y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{2}{\pi^2} \frac{d}{dx} \int_x^a \frac{F(t) dt}{\sqrt{t^2 - x^2}}, \quad F(t) = \frac{d}{dt} \int_0^t \frac{sf(s) ds}{\sqrt{t^2 - s^2}}.$$

⊙ Reference: P. P. Zabreyko, A. I. Koshelev, et al. (1975).

$$6. \int_a^b \left| \ln \frac{1+\lambda x}{1+\lambda t} \right| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \ln(1 + \lambda x)$.

Solution:

$$y(x) = \frac{1}{2\lambda} \frac{d}{dx} [(1 + \lambda x) f'_x(x)].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2^o of equation 3.8.3).

$$7. \int_a^b |\ln^\beta x - \ln^\beta t| y(t) dt = f(x), \quad 0 < \beta < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \ln^\beta x$.

$$8. \int_a^b \frac{y(t)}{|\ln(x/t)|^\beta} dt = f(x), \quad 0 < \beta < 1.$$

This is a special case of equation 3.8.7 with $g(x) = \ln x + A$, where A is an arbitrary number.

3.4-2. Kernels Containing Power-Law and Logarithmic Functions.

$$9. \int_0^1 (\ln |x-t| + \beta t^k) y(t) dt = f(x).$$

See Example 3 in Subsection 12.6-2 with $\psi(t) = \beta t^k$.

$$10. \int_0^a |k \ln(1 + \lambda x) - t| y(t) dt = f(x).$$

This is a special case of equation 3.8.5 with $g(x) = k \ln(1 + \lambda x)$.

$$11. \int_0^a |x - k \ln(1 + \lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.6 with $g(x) = k \ln(1 + \lambda t)$.

$$12. \int_0^\infty \frac{1}{t} \ln \left| \frac{x+t}{x-t} \right| y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{x}{\pi^2} \frac{d}{dx} \int_0^\infty \frac{df(t)}{dt} \ln \left| 1 - \frac{x^2}{t^2} \right| dt.$$

⊙ Reference: P. P. Zabreyko, A. I. Koshelev, et al. (1975).

$$13. \int_0^{\infty} \frac{\ln x - \ln t}{x - t} y(t) dt = f(x).$$

The left-hand side of this equation is the iterated Stieltjes transform.

Under some assumptions, the solution of the integral equation can be represented in the form

$$y(x) = \frac{1}{4\pi^2} \lim_{n \rightarrow \infty} \left(\frac{e}{n}\right)^{4n} D^n x^{2n} D^{2n} x^{2n} D^n f(x), \quad D = \frac{d}{dx}.$$

To calculate the solution approximately, one should restrict oneself to a specific value of n in this formula instead of taking the limit.

⊙ Reference: I. I. Hirschman and D. V. Widder (1955).

$$14. \int_a^b \ln |x^\beta - t^\beta| y(t) dt = f(x), \quad \beta > 0.$$

The transformation

$$z = x^\beta, \quad \tau = t^\beta, \quad w(\tau) = t^{1-\beta} y(t)$$

leads to Carleman's equation 3.4.2:

$$\int_A^B \ln |z - \tau| w(\tau) d\tau = F(z), \quad A = a^\beta, \quad B = b^\beta,$$

where $F(z) = \beta f(z^{1/\beta})$.

$$15. \int_0^1 \ln |x^\beta - t^\mu| y(t) dt = f(x), \quad \beta > 0, \quad \mu > 0.$$

The transformation

$$z = x^\beta, \quad \tau = t^\mu, \quad w(\tau) = t^{1-\mu} y(t)$$

leads to an equation of the form 3.4.2:

$$\int_0^1 \ln |z - \tau| w(\tau) d\tau = F(z), \quad F(z) = \mu f(z^{1/\beta}).$$

$$16. \int_0^{\infty} \frac{1}{\sqrt{xt} \ln(xt)} y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{1}{\pi^2} \int_0^{\infty} \frac{1}{\sqrt{xt} \ln(xt)} f(t) dt.$$

⊙ References: E. C. Titchmarsh (1986), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 450).

$$17. \frac{d}{dx} \int_{-\infty}^{\infty} \ln \left| 1 - \frac{x}{t} \right| y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{1}{\pi^2} \frac{d}{dx} \int_{-\infty}^{\infty} \ln \left| 1 - \frac{x}{t} \right| f(t) dt.$$

⊙ References: E. C. Titchmarsh (1986), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 450).

$$18. \int_0^{\infty} (xt)^{-[1+i \ln(xt)]/2} y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(x) = \frac{1}{2\pi} \int_0^{\infty} (xt)^{-[1-i \ln(xt)]/2} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 452).

3.4-3. Equation Containing the Unknown Function of a Complicated Argument.

$$19. \int_0^1 (A \ln t + B)y(xt) dt = f(x).$$

The substitution $\xi = xt$ leads to an equation of the form 1.9.3 with $g(x) = -A \ln x$:

$$\int_0^x (A \ln \xi - A \ln x + B)y(\xi) d\xi = xf(x).$$

3.5. Equations Whose Kernels Contain Trigonometric Functions

3.5-1. Kernels Containing Cosine.

$$1. \int_0^\infty \cos(xt)y(t) dt = f(x).$$

Solution: $y(x) = \frac{2}{\pi} \int_0^\infty \cos(xt)f(t) dt.$

Up to constant factors, the function $f(x)$ and the solution $y(t)$ are the *Fourier cosine transform pair*.

⊙ References: E. A. C. Paley and N. Wiener (1934), S. Bochner and K. C. Chandrasekharan (1949), G. N. Watson (1952), H. Bateman and A. Erdélyi (Vol. 1, 1954), S. Bochner (1959), V. A. Ditkin and A. P. Prudnikov (1965), B. Davis (1978), F. Oberhettinger (1980), E. C. Titchmarsh (1986), Ya. A. Brychkov and A. P. Prudnikov (1989), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 440), I. Sneddon (1995), A. D. Poularikas (2000).

$$2. \int_a^b \cos(xt)y(t) dt = f(x), \quad 0 \leq x < \infty.$$

Solution:

$$y(t) = \begin{cases} \frac{2}{\pi} \int_0^\infty \cos(xt)f(x) dx & \text{if } a < t < b, \\ 0 & \text{if } 0 < t < a \text{ or } t > b, \end{cases}$$

where $0 \leq a \leq b \leq \infty$.

$$3. \int_a^b |\cos(\lambda x) - \cos(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \cos(\lambda x)$.

Solution:

$$y(x) = -\frac{1}{2\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\sin(\lambda x)} \right].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$4. \int_0^a |\cos(\beta x) - \cos(\mu t)| y(t) dt = f(x), \quad \beta > 0, \quad \mu > 0.$$

This is a special case of equation 3.8.4 with $g(x) = \cos(\beta x)$ and $\lambda = \mu/\beta$.

$$5. \int_a^b |\cos^k x - \cos^k t| y(t) dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \cos^k x$.

Solution:

$$y(x) = -\frac{1}{2k} \frac{d}{dx} \left[\frac{f'_x(x)}{\sin x \cos^{k-1} x} \right].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$6. \int_a^b \frac{y(t)}{|\cos(\lambda x) - \cos(\lambda t)|^k} dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.7 with $g(x) = \cos(\lambda x) + \beta$, where β is an arbitrary number.

$$7. \int_0^\infty t^{-ix-1/2} \cos\left(\frac{1+2ix}{4}\pi\right) y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(t) = \frac{1}{\pi} \int_{-\infty}^\infty t^{ix-1/2} \cos\left(\frac{1-2ix}{4}\pi\right) \frac{f(x)}{\cosh(\pi x)} dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 463).

3.5-2. Kernels Containing Sine.

$$8. \int_0^\infty \sin(xt)y(t) dt = f(x).$$

$$\text{Solution: } y(x) = \frac{2}{\pi} \int_0^\infty \sin(xt)f(t) dt.$$

Up to constant factors, the function $f(x)$ and the solution $y(t)$ are the *Fourier sine transform pair*.

⊙ References: E. A. C. Paley and N. Wiener (1934), S. Bochner and K. C. Chandrasekharan (1949), G. N. Watson (1952), H. Bateman and A. Erdélyi (Vol. 1, 1954), S. Bochner (1959), V. A. Ditkin and A. P. Prudnikov (1965), B. Davis (1978), F. Oberhettinger (1980), E. C. Titchmarsh (1986), Ya. A. Brychkov and A. P. Prudnikov (1989), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 440), I. Sneddon (1995), A. D. Poularikas (2000).

$$9. \int_a^b \sin(xt)y(t) dt = f(x), \quad 0 \leq x < \infty.$$

Solution:

$$y(t) = \begin{cases} \frac{2}{\pi} \int_0^\infty \sin(xt)f(x) dx & \text{if } a < t < b, \\ 0 & \text{if } 0 < t < a \text{ or } t > b, \end{cases}$$

where $0 \leq a \leq b \leq \infty$.

$$10. \int_{-\infty}^\infty \sin(\lambda|x-t|)y(t) dt = f(x), \quad f(\pm\infty) = 0.$$

Solution:

$$y(x) = \frac{1}{2\lambda} [f''_{xx}(x) + \lambda^2 f(x)].$$

$$11. \int_a^b \sin(\lambda|x-t|)y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

1°. Let us remove the modulus in the integrand:

$$\int_a^x \sin[\lambda(x-t)]y(t) dt + \int_x^b \sin[\lambda(t-x)]y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x twice yields

$$2\lambda y(x) - \lambda^2 \int_a^x \sin[\lambda(x-t)]y(t) dt - \lambda^2 \int_x^b \sin[\lambda(t-x)]y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral terms from (1) and (2), we obtain the solution

$$y(x) = \frac{1}{2\lambda} [f''_{xx}(x) + \lambda^2 f(x)]. \quad (3)$$

2°. The right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = a$ and $x = b$ in (1), we obtain two corollaries

$$\int_a^b \sin[\lambda(t-a)]y(t) dt = f(a), \quad \int_a^b \sin[\lambda(b-t)]y(t) dt = f(b). \quad (4)$$

Substituting solution (3) into (4) followed by integrating by parts yields the desired conditions for $f(x)$:

$$\begin{aligned} \sin[\lambda(b-a)]f'_x(b) - \lambda \cos[\lambda(b-a)]f(b) &= \lambda f(a), \\ \sin[\lambda(b-a)]f'_x(a) + \lambda \cos[\lambda(b-a)]f(a) &= -\lambda f(b). \end{aligned} \quad (5)$$

The general form of the right-hand side of the integral equation is given by

$$f(x) = F(x) + Ax + B, \quad (6)$$

where $F(x)$ is an arbitrary bounded twice differentiable function, and the coefficients A and B are expressed in terms of $F(a)$, $F(b)$, $F'_x(a)$, and $F'_x(b)$ and can be determined by substituting formula (6) into conditions (5).

$$12. \int_a^b \{A \sin(\lambda|x-t|) + B \sin(\mu|x-t|)\}y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

Let us remove the modulus in the integrand and differentiate the equation with respect to x twice to obtain

$$2(A\lambda + B\mu)y(x) - \int_a^b \{A\lambda^2 \sin(\lambda|x-t|) + B\mu^2 \sin(\mu|x-t|)\}y(t) dt = f''_{xx}(x). \quad (1)$$

Eliminating the integral term with $\sin(\mu|x-t|)$ from (1) with the aid of the original equation, we find that

$$2(A\lambda + B\mu)y(x) + A(\mu^2 - \lambda^2) \int_a^b \sin(\lambda|x-t|)y(t) dt = f''_{xx}(x) + \mu^2 f(x). \quad (2)$$

For $A\lambda + B\mu = 0$, this is an equation of the form 3.5.11 and for $A\lambda + B\mu \neq 0$, this is an equation of the form 4.5.29.

The right-hand side $f(x)$ must satisfy certain relations, which can be obtained by setting $x = a$ and $x = b$ in the original equation (a similar procedure is used in 3.5.11).

$$13. \int_a^b |\sin(\lambda x) - \sin(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \sin(\lambda x)$.

Solution:

$$y(x) = \frac{1}{2\lambda} \frac{d}{dx} \left[\frac{f'_x(x)}{\cos(\lambda x)} \right].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$14. \int_0^a |\sin(\beta x) - \sin(\mu t)| y(t) dt = f(x), \quad \beta > 0, \quad \mu > 0.$$

This is a special case of equation 3.8.4 with $g(x) = \sin(\beta x)$ and $\lambda = \mu/\beta$.

$$15. \int_a^b \sin^3(\lambda|x-t|) y(t) dt = f(x).$$

Using the formula $\sin^3 \beta = -\frac{1}{4} \sin 3\beta + \frac{3}{4} \sin \beta$, we arrive at an equation of the form 3.5.12:

$$\int_a^b \left[-\frac{1}{4} A \sin(3\lambda|x-t|) + \frac{3}{4} A \sin(\lambda|x-t|) \right] y(t) dt = f(x).$$

$$16. \int_a^b \left[\sum_{k=1}^n A_k \sin(\lambda_k|x-t|) \right] y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

1°. Let us remove the modulus in the k th summand of the integrand:

$$I_k(x) = \int_a^b \sin(\lambda_k|x-t|) y(t) dt = \int_a^x \sin[\lambda_k(x-t)] y(t) dt + \int_x^b \sin[\lambda_k(t-x)] y(t) dt. \quad (1)$$

Differentiating (1) with respect to x yields

$$\begin{aligned} I'_k &= \lambda_k \int_a^x \cos[\lambda_k(x-t)] y(t) dt - \lambda_k \int_x^b \cos[\lambda_k(t-x)] y(t) dt, \\ I''_k &= 2\lambda_k y(x) - \lambda_k^2 \int_a^x \sin[\lambda_k(x-t)] y(t) dt - \lambda_k^2 \int_x^b \sin[\lambda_k(t-x)] y(t) dt, \end{aligned} \quad (2)$$

where the primes denote the derivatives with respect to x . By comparing formulas (1) and (2), we find the relation between I''_k and I_k :

$$I''_k = 2\lambda_k y(x) - \lambda_k^2 I_k, \quad I_k = I_k(x). \quad (3)$$

2°. With the aid of (1), the integral equation can be rewritten in the form

$$\sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x twice and taking into account (3), we find that

$$\sigma_1 y(x) - \sum_{k=1}^n A_k \lambda_k^2 I_k = f''_{xx}(x), \quad \sigma_1 = 2 \sum_{k=1}^n A_k \lambda_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5) yields

$$\sigma_1 y(x) + \sum_{k=1}^{n-1} A_k (\lambda_n^2 - \lambda_k^2) I_k = f''_{xx}(x) + \lambda_n^2 f(x). \quad (6)$$

Differentiating (6) with respect to x twice and eliminating I_{n-1} from the resulting equation with the aid of (6), we obtain a similar equation whose left-hand side is a second-order linear differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} B_k I_k$.

If we successively eliminate I_{n-2}, I_{n-3}, \dots , with the aid of double differentiation, then we finally arrive at a linear nonhomogeneous ordinary differential equation of order $2(n-1)$ with constant coefficients.

3°. The right-hand side $f(x)$ must satisfy certain conditions. To find these conditions, one should set $x = a$ in the integral equation and its derivatives. (Alternatively, these conditions can be found by setting $x = a$ and $x = b$ in the integral equation and all its derivatives obtained by means of double differentiation.)

$$17. \int_0^b |\sin^k x - \sin^k t| y(t) dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \sin^k x$.

Solution:

$$y(x) = \frac{1}{2k} \frac{d}{dx} \left[\frac{f'_x(x)}{\cos x \sin^{k-1} x} \right].$$

The right-hand side $f(x)$ must satisfy certain conditions. As follows from item 3° of equation 3.8.3, the admissible general form of the right-hand side is given by

$$f(x) = F(x) + Ax + B, \quad A = -F'_x(b), \quad B = \frac{1}{2} [bF'_x(b) - F(0) - F(b)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

$$18. \int_a^b \frac{y(t)}{|\sin(\lambda x) - \sin(\lambda t)|^k} dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.7 with $g(x) = \sin(\lambda x) + \beta$, where β is an arbitrary number.

$$19. \int_0^a |k \sin(\lambda x) - t| y(t) dt = f(x).$$

This is a special case of equation 3.8.5 with $g(x) = k \sin(\lambda x)$.

$$20. \int_0^a |x - k \sin(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.6 with $g(t) = k \sin(\lambda t)$.

$$21. \int_0^\infty \frac{\sin t}{t^2} [y(x+t) - y(x-t)] dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi} \int_0^\infty \left[\frac{\cos t}{t} + \text{Si}(t) \right] [f(x-t) - f(x+t)] dt,$$

where $\text{Si}(t)$ is sine integral (see Supplement 11.3-1).

The integral equation and its solution form the *Boas transform pair*.

$$22. \int_0^{\infty} t^{-ix-1/2} \sin\left(\frac{1+2ix}{4}\pi\right) y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(t) = \frac{1}{\pi} \int_{-\infty}^{\infty} t^{ix-1/2} \sin\left(\frac{1-2ix}{4}\pi\right) \frac{f(x)}{\cosh(\pi x)} dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 463).

3.5-3. Kernels Containing Tangent.

$$23. \int_a^b |\tan(\lambda x) - \tan(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \tan(\lambda x)$.

Solution:

$$y(x) = \frac{1}{2\lambda} \frac{d}{dx} \left[\cos^2(\lambda x) f'_x(x) \right].$$

The right-hand side $f(x)$ of the integral equation must satisfy certain relations (see item 2° of equation 3.8.3).

$$24. \int_0^a |\tan(\beta x) - \tan(\mu t)| y(t) dt = f(x), \quad \beta > 0, \quad \mu > 0.$$

This is a special case of equation 3.8.4 with $g(x) = \tan(\beta x)$ and $\lambda = \mu/\beta$.

$$25. \int_0^b |\tan^k x - \tan^k t| y(t) dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \tan^k x$.

Solution:

$$y(x) = \frac{1}{2k} \frac{d}{dx} \left[\cos^2 x \cot^{k-1} x f'_x(x) \right].$$

The right-hand side $f(x)$ must satisfy certain conditions. As follows from item 3° of equation 3.8.3, the admissible general form of the right-hand side is given by

$$f(x) = F(x) + Ax + B, \quad A = -F'_x(b), \quad B = \frac{1}{2} [bF'_x(b) - F(0) - F(b)],$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

$$26. \int_a^b \frac{y(t)}{|\tan(\lambda x) - \tan(\lambda t)|^k} dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.7 with $g(x) = \tan(\lambda x) + \beta$, where β is an arbitrary number.

$$27. \int_0^a |k \tan(\lambda x) - t| y(t) dt = f(x).$$

This is a special case of equation 3.8.5 with $g(x) = k \tan(\lambda x)$.

$$28. \int_0^a |x - k \tan(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.6 with $g(t) = k \tan(\lambda t)$.

3.5-4. Kernels Containing Cotangent.

$$29. \int_a^b |\cot(\lambda x) - \cot(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = \cot(\lambda x)$.

$$30. \int_a^b |\cot^k x - \cot^k t| y(t) dt = f(x), \quad 0 < k < 1.$$

This is a special case of equation 3.8.3 with $g(x) = \cot^k x$.

3.5-5. Kernels Containing a Combination of Trigonometric Functions.

$$31. \int_{-\infty}^{\infty} [\cos(xt) + \sin(xt)] y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} [\cos(xt) + \sin(xt)] f(t) dt.$$

Up to constant factors, the function $f(x)$ and the solution $y(t)$ are the *Hartley transform pair*.

⊙ Reference: D. Zwillinger (1989).

$$32. \int_0^{\infty} [\sin(xt) - xt \cos(xt)] y(t) dt = f(x).$$

This equation can be reduced to a special case of equation 3.7.17 with $\nu = \frac{3}{2}$.

Solution:

$$y(x) = \frac{2}{\pi} \int_0^{\infty} \frac{\sin(xt) - xt \cos(xt)}{x^2 t^2} f(t) dt.$$

$$33. \int_0^{\infty} [\sin(xt) + xt \cos(xt)] y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{2}{\pi} \int_0^{\infty} \text{si}(xt) y(t) dt,$$

where $\text{si}(z)$ is the sine integral (see Supplement 11.3-1).

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 457).

$$34. \int_0^{\infty} [1 - \cos(xt) + xt \sin(xt)] y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^{\infty} \text{ci}(xt) f(t) dt,$$

where $\text{ci}(z)$ is the cosine integral (see Supplement 11.3-2).

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 457).

$$35. \int_0^{\infty} (xt)^{1/2} \left[\frac{\sin(xt)}{xt} + 2 \cos(xt) \right] y(t) dt = f(x).$$

Solution:

$$y(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \left[\frac{1}{2} - S(xt) \right] f(t) dt.$$

where $S(z)$ is the Fresnel sine integral (see Supplement 11.3-3).

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 459).

$$36. \int_0^{\infty} (xt)^{1/2} \left[\frac{\cos(xt) - 1}{xt} - 2 \sin(xt) \right] y(t) dt = f(x).$$

Solution:

$$y(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \left[\frac{1}{2} - C(xt) \right] f(t) dt,$$

where $C(z)$ is the Fresnel cosine integral (see Supplement 11.3-3).

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 460).

$$37. \int_0^{\infty} \frac{(1-\nu) \sin(xt) + xt \cos(xt)}{(xt)^{\nu}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^{\infty} S(xt, \nu) f(t) dt,$$

where $S(z, \nu)$ is the generalized Fresnel sine integral (see Supplement 11.3-3).

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 461).

$$38. \int_0^{\infty} \frac{(1-\nu) \cos(xt) - xt \sin(xt)}{(xt)^{\nu}} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^{\infty} C(xt, \nu) y(t) dt,$$

where $C(z, \nu)$ is the generalized Fresnel cosine integral (see Supplement 11.3-3).

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 461).

$$39. \int_0^{\pi} \left[\frac{a \sin(x+t)}{1-2a \cos(x+t)+a^2} + \frac{a \sin(x-t)}{1-2a \cos(x-t)+a^2} \right] y(t) dt = f(x), \quad 0 < a < 1.$$

Solution:

$$y(x) = C + \frac{2}{\pi^2} \sum_{n=1}^{\infty} \frac{f_n}{a^n} \cos(nx), \quad f_n = \int_0^{\pi} f(x) \sin(nx) dx,$$

where C is an arbitrary constant.

Remark. The kernel of the integral equation can be represented as a series in powers of a :

$$K(x, t) = \frac{a \sin(x+t)}{1-2a \cos(x+t)+a^2} + \frac{a \sin(x-t)}{1-2a \cos(x-t)+a^2} = 2 \sum_{n=1}^{\infty} a^n \sin(nx) \cos(nt).$$

⊙ References: W. Schmeidler (1950, p. 169), S. Fenyö and H. W. Stolle (1984, pp. 18–19).

3.5-6. Equations Containing the Unknown Function of a Complicated Argument.

$$40. \int_0^{\pi/2} y(\xi) dt = f(x), \quad \xi = x \sin t.$$

Schlömilch equation.

Solution:

$$y(x) = \frac{2}{\pi} \left[f(0) + x \int_0^{\pi/2} f'_\xi(\xi) dt \right], \quad \xi = x \sin t.$$

⊙ References: E. T. Whittaker and G. N. Watson (1958), F. D. Gakhov (1977).

$$41. \int_0^{\pi/2} y(\xi) dt = f(x), \quad \xi = x \sin^k t.$$

Generalized Schlömilch equation.

This is a special case of equation 3.5.43 for $\lambda = 0$ and $m = 0$.

Solution:

$$y(x) = \frac{2k}{\pi} x^{\frac{k-1}{k}} \frac{d}{dx} \left[x^{\frac{1}{k}} \int_0^x \sin t f(\xi) dt \right], \quad \xi = x \sin^k t.$$

$$42. \int_0^{\pi/2} \sin^\lambda t y(\xi) dt = f(x), \quad \xi = x \sin^k t.$$

This is a special case of equation 3.5.43 for $m = 0$.

Solution:

$$y(x) = \frac{2k}{\pi} x^{\frac{k-\lambda-1}{k}} \frac{d}{dx} \left[x^{\frac{\lambda+1}{k}} \int_0^x \sin^{\lambda+1} t f(\xi) dt \right], \quad \xi = x \sin^k t.$$

$$43. \int_0^{\pi/2} \sin^\lambda t \cos^m t y(\xi) dt = f(x), \quad \xi = x \sin^k t.$$

1°. Let $\lambda > -1$, $m > -1$, and $k > 0$. The transformation

$$z = x^{\frac{2}{k}}, \quad \zeta = z \sin^2 t, \quad w(\zeta) = \zeta^{\frac{\lambda-1}{2}} y\left(\zeta^{\frac{k}{2}}\right)$$

leads to an equation of the form 1.1.44:

$$\int_0^z (z - \zeta)^{\frac{m-1}{2}} w(\zeta) d\zeta = F(z), \quad F(z) = 2z^{\frac{\lambda+m}{2}} f\left(z^{\frac{k}{2}}\right).$$

2°. Solution with $-1 < m < 1$:

$$y(x) = \frac{2k}{\pi} \sin\left[\frac{\pi(1-m)}{2}\right] x^{\frac{k-\lambda-1}{k}} \frac{d}{dx} \left[x^{\frac{\lambda+1}{k}} \int_0^{\pi/2} \sin^{\lambda+1} t \tan^m t f(\xi) dt \right],$$

where $\xi = x \sin^k t$.

3.5-7. Singular Equations.

$$44. \int_0^{2\pi} \cot\left(\frac{t-x}{2}\right) y(t) dt = f(x), \quad 0 \leq x \leq 2\pi.$$

Here the integral is understood in the sense of the Cauchy principal value and the right-hand side is assumed to satisfy the condition $\int_0^{2\pi} f(t) dt = 0$.

Solution:

$$y(x) = -\frac{1}{4\pi^2} \int_0^{2\pi} \cot\left(\frac{t-x}{2}\right) f(t) dt + C,$$

where C is an arbitrary constant.

It follows from the solution that $\int_0^{2\pi} y(t) dt = 2\pi C$.

The equation and its solution form a *Hilbert transform pair* (in the asymmetric form).

⊙ Reference: F. D. Gakhov (1977).

$$45. \int_{-\pi}^{\pi} \left[1 + \cot\left(\frac{x-t}{2}\right) \right] y(t) dt = f(x), \quad -\pi \leq x \leq \pi.$$

Hilbert–Plessner equation.

Solution:

$$y(x) = \frac{1}{4\pi^2} \int_{-\pi}^{\pi} \left[1 + \cot\left(\frac{x-t}{2}\right) \right] f(t) dt.$$

⊙ Reference: S. Fenyő and H. W. Stolle (1984, pp. 36–38).

$$46. \int_0^{2\pi} \left[\sin\left(\frac{\xi-x}{2}\right) \right]^{-2} y(\xi) d\xi = f(x), \quad 0 \leq x \leq 2\pi.$$

The simple hypersingular equation of the first kind with Hilbert-type kernel.

Let the periodic conditions $y(0) = y(2\pi)$ be satisfied. Then the solution is

$$y(x) = -\frac{1}{4\pi^2} \int_0^{2\pi} f(\xi) \ln \left| \sin\left(\frac{\xi-x}{2}\right) \right| d\xi + C,$$

where C is an arbitrary constant.

This equation is discussed in Subsection 14.6-4 in detail.

⊙ Reference: I. K. Lifanov, L. N. Poltavskii, and G. M. Vainikko (2004, p. 8).

3.6. Equations Whose Kernels Contain Combinations of Elementary Functions

3.6-1. Kernels Containing Hyperbolic and Logarithmic Functions.

$$1. \int_a^b \ln |\cosh(\lambda x) - \cosh(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.9 with $g(x) = \cosh(\lambda x)$.

$$2. \int_a^b \ln |\sinh(\lambda x) - \sinh(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.9 with $g(x) = \sinh(\lambda x)$.

$$3. \int_{-a}^a \ln \left[\frac{\sinh(\frac{1}{2}A)}{2 \sinh(\frac{1}{2}|x-t|)} \right] y(t) dt = f(x), \quad -a \leq x \leq a.$$

Solution with $0 < a < A$:

$$\begin{aligned} y(x) &= \frac{1}{2M'(a)} \left[\frac{d}{da} \int_{-a}^a w(t, a) f(t) dt \right] w(x, a) \\ &\quad - \frac{1}{2} \int_{|x|}^a w(x, \xi) \frac{d}{d\xi} \left[\frac{1}{M'(\xi)} \frac{d}{d\xi} \int_{-\xi}^{\xi} w(t, \xi) f(t) dt \right] d\xi \\ &\quad - \frac{1}{2} \frac{d}{dx} \int_{|x|}^a \frac{w(x, \xi)}{M'(\xi)} \left[\int_{-\xi}^{\xi} w(t, \xi) df(t) \right] d\xi, \end{aligned}$$

where the prime stands for the derivative with respect to the argument and

$$M(\xi) = \left[\ln \left(\frac{\sinh(\frac{1}{2}A)}{\sinh(\frac{1}{2}\xi)} \right) \right]^{-1}, \quad w(x, \xi) = \frac{\cosh(\frac{1}{2}x) M(\xi)}{\pi \sqrt{2 \cosh \xi - 2 \cosh x}}.$$

⊙ Reference: I. C. Gohberg and M. G. Krein (1967).

$$4. \int_a^b \ln |\tanh(\lambda x) - \tanh(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.9 with $g(x) = \tanh(\lambda x)$.

$$5. \int_{-a}^a \ln [\coth(\frac{1}{4}|x-t|)] y(t) dt = f(x), \quad -a \leq x \leq a.$$

Solution:

$$\begin{aligned} y(x) &= \frac{1}{2M'(a)} \left[\frac{d}{da} \int_{-a}^a w(t, a) f(t) dt \right] w(x, a) \\ &\quad - \frac{1}{2} \int_{|x|}^a w(x, \xi) \frac{d}{d\xi} \left[\frac{1}{M'(\xi)} \frac{d}{d\xi} \int_{-\xi}^{\xi} w(t, \xi) f(t) dt \right] d\xi \\ &\quad - \frac{1}{2} \frac{d}{dx} \int_{|x|}^a \frac{w(x, \xi)}{M'(\xi)} \left[\int_{-\xi}^{\xi} w(t, \xi) df(t) \right] d\xi, \end{aligned}$$

where the prime stands for the derivative with respect to the argument and

$$M(\xi) = \frac{P_{-1/2}(\cosh \xi)}{Q_{-1/2}(\cosh \xi)}, \quad w(x, \xi) = \frac{1}{\pi Q_{-1/2}(\cosh \xi) \sqrt{2 \cosh \xi - 2 \cosh x}},$$

and $P_{-1/2}(\cosh \xi)$ and $Q_{-1/2}(\cosh \xi)$ are the Legendre functions of the first and second kind, respectively.

⊙ Reference: I. C. Gohberg and M. G. Krein (1967).

3.6-2. Kernels Containing Logarithmic and Trigonometric Functions.

$$6. \int_a^b \ln |\cos(\lambda x) - \cos(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.9 with $g(x) = \cos(\lambda x)$.

$$7. \int_a^b \ln |\sin(\lambda x) - \sin(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.9 with $g(x) = \sin(\lambda x)$.

$$8. \int_0^\pi \ln \frac{1 - \cos(x+t)}{1 - \cos(x-t)} y(t) dt = f(x), \quad 0 \leq x \leq \pi.$$

Solution:

$$y(x) = \frac{2}{\pi^2} \sum_{n=1}^{\infty} n f_n \sin(nx), \quad f_n = \int_0^\pi f(x) \sin(nx) dx.$$

⊙ Reference: S. Fenyő and H. W. Stolle (1984, p. 44).

$$9. \int_{-a}^a \ln \left[\frac{\sin(\frac{1}{2}A)}{2 \sin(\frac{1}{2}|x-t|)} \right] y(t) dt = f(x), \quad -a \leq x \leq a.$$

Solution with $0 < a < A$:

$$\begin{aligned} y(x) &= \frac{1}{2M'(a)} \left[\frac{d}{da} \int_{-a}^a w(t, a) f(t) dt \right] w(x, a) \\ &\quad - \frac{1}{2} \int_{|x|}^a w(x, \xi) \frac{d}{d\xi} \left[\frac{1}{M'(\xi)} \frac{d}{d\xi} \int_{-\xi}^{\xi} w(t, \xi) f(t) dt \right] d\xi \\ &\quad - \frac{1}{2} \frac{d}{dx} \int_{|x|}^a \frac{w(x, \xi)}{M'(\xi)} \left[\int_{-\xi}^{\xi} w(t, \xi) df(t) \right] d\xi, \end{aligned}$$

where the prime stands for the derivative with respect to the argument and

$$M(\xi) = \left[\ln \left(\frac{\sin(\frac{1}{2}A)}{\sin(\frac{1}{2}\xi)} \right) \right]^{-1}, \quad w(x, \xi) = \frac{\cos(\frac{1}{2}\xi) M(\xi)}{\pi \sqrt{2 \cos x - 2 \cos \xi}}.$$

⊙ Reference: I. C. Gohberg and M. G. Krein (1967).

$$10. \frac{d}{dx} \int_{-\pi}^{\pi} \ln \left(2 \left| \sin \frac{x-t}{2} \right| \right) y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{1}{\pi^2} \frac{d}{dx} \int_{-\pi}^{\pi} \ln \left(2 \left| \sin \frac{x-t}{2} \right| \right) f(t) dt, \quad \int_{-\pi}^{\pi} y(t) dt = 0.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 452).

3.6-3. Kernels Containing Combinations of Exponential and Other Elementary Functions.

$$11. \int_a^b (\ln |x-t| + Ae^{-\alpha x - \beta t}) y(t) dt = f(x).$$

This is a special case of equation 3.8.28 with $\varphi(x) = Ae^{-\alpha x}$ and $\psi(t) = e^{-\beta t}$.

$$12. \int_0^\infty [\sin(xt) + Ae^{-\alpha x - \beta t}] y(t) dt = f(x).$$

This is a special case of equation 3.8.29 with $\varphi(x) = Ae^{-\alpha x}$ and $\psi(t) = e^{-\beta t}$.

$$13. \int_0^\infty [\cos(xt) + Ae^{-\alpha x - \beta t}] y(t) dt = f(x).$$

This is a special case of equation 3.8.30 with $\varphi(x) = Ae^{-\alpha x}$ and $\psi(t) = e^{-\beta t}$.

3.7. Equations Whose Kernels Contain Special Functions*

3.7-1. Kernels Containing Error Function, Exponential Integral or Logarithmic Integral.

$$1. \int_0^{\infty} [\exp(i(x+t)^2) \operatorname{erf}(e^{\pi i/4}(x+t)) + \exp(i(x-t)^2) \operatorname{erf}(e^{\pi i/4}(x-t))] y(t) dt = f(x).$$

Here $\operatorname{erf} z$ is the error function (see Supplement 11.2-1) and $i^2 = -1$.

Solution:

$$y(x) = -\frac{1}{\pi} \int_0^{\infty} [\exp(-i(t+x)^2) \operatorname{erf}(e^{3\pi i/4}(t+x)) + \exp(-i(t-x)^2) \operatorname{erf}(e^{3\pi i/4}(t-x))] f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 459).

$$2. \int_0^{\infty} e^{-ixt} \operatorname{Ei}(ixt) y(t) dt = f(x), \quad i^2 = -1.$$

Here $\operatorname{Ei}(z)$ is the exponential integral (see Supplement 11.2-2).

Solution:

$$y(t) = \frac{1}{2\pi^2} \int_{-\infty}^{\infty} \left[e^{ixt} \operatorname{erf}(e^{\pi i/4}\sqrt{xt}) - \frac{1+i}{\sqrt{2\pi xt}} \right] f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 456).

$$3. \int_1^{\infty} \operatorname{li}\left(\frac{x}{t}\right) y(t) dt = f(x), \quad f(1) = f'(1) = 0.$$

Here $\operatorname{li}(z)$ is the logarithmic integral (see Supplement 11.2-3).

Solution:

$$y(t) = -\int_1^x t^{-2} \nu\left(\ln \frac{t}{x}\right) \left[\left(t \frac{d}{dt}\right)^2 - t \frac{d}{dt} \right] f(t) dt,$$

$$\text{where } \nu(z) = \int_0^{\infty} \frac{z^{\xi} d\xi}{\Gamma(\xi+1)}.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 457).

3.7-2. Kernels Containing Sine Integrals, Cosine Integrals, or Fresnel Integrals.

$$4. \int_0^{\infty} \operatorname{si}(xt) y(t) dt = f(x).$$

Here $\operatorname{si}(z)$ is the sine integral (see Supplement 11.3-1).

Solution:

$$y(x) = -\frac{2}{\pi} \int_0^{\infty} [\sin(xt) + xt \cos(xt)] f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 457).

* For notation and properties of special functions, see Supplement 11.

$$5. \int_0^{\infty} \text{ci}(xt)y(t) dt = f(x).$$

Here $\text{ci}(z)$ is the cosine integral (see Supplement 11.3-2).

Solution:

$$y(x) = \frac{2}{\pi} \int_0^{\infty} [1 - \cos(xt) + xt \sin(xt)] f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 457).

$$6. \int_0^{\infty} \left[\frac{1}{2} - S(xt) \right] y(t) dt = f(x).$$

Here $S(z)$ is the Fresnel sine integral (see Supplement 11.3-3).

Solution:

$$y(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} (xt)^{1/2} \left[\frac{\sin(xt)}{xt} + 2 \cos(xt) \right] f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 459).

$$7. \int_0^{\infty} \left[\frac{1}{2} - C(xt) \right] y(t) dt = f(x).$$

Here $C(z)$ is the Fresnel cosine integral (see Supplement 11.3-3).

Solution:

$$y(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} (xt)^{1/2} \left[\frac{\cos(xt) - 1}{xt} - 2 \sin(xt) \right] f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 460).

$$8. \int_0^{\infty} S(xt, \nu) y(t) dt = f(x).$$

Here $S(z, \nu)$ is the generalized Fresnel sine integral (see Supplement 11.3-3).

Solution:

$$y(x) = \frac{2}{\pi} \int_0^{\infty} \frac{(1 - \nu) \sin(xt) + xt \cos(xt)}{(xt)^{\nu}} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 461).

$$9. \int_0^{\infty} C(xt, \nu) y(t) dt = f(x).$$

Here $C(z, \nu)$ is the generalized Fresnel cosine integral (see Supplement 11.3-3).

Solution:

$$y(x) = \frac{2}{\pi} \int_0^{\infty} \frac{(1 - \nu) \cos(xt) - xt \sin(xt)}{(xt)^{\nu}} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 461).

3.7-3. Kernels Containing Gamma Functions.
--

10.
$$\int_0^{\infty} (xt)^{-(\pi+1)/2} \Gamma(\pm i \ln(xt)) y(t) dt = f(x), \quad i^2 = -1.$$

Here $\Gamma(z)$ is the incomplete gamma function (see Supplement 11.4-1).

Solution:

$$y(x) = \frac{1}{4\pi^2} \int_0^{\infty} (xt)^{-(\pi+1)/2} \Gamma(\mp i \ln(xt)) f(t) dt.$$

The integral equation and its solution form a *Paley–Wiener transform pair* (in the asymmetric form).

⊙ References: E. C. Titchmarsh (1986), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 453).

11.
$$\int_{-\infty}^{\infty} e^{-\pi(x+t)/2} \Gamma(\pm i(x+t)) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{4\pi^2} \int_{-\infty}^{\infty} e^{-\pi(x+t)/2} \Gamma(\mp i(x+t)) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 453).

12.
$$\int_{-\infty}^{\infty} \Gamma(\alpha + i(x+t)) \Gamma(\alpha - i(x+t)) y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{\alpha \sin(2\pi\alpha)}{2\pi^3} \int_{-\infty}^{\infty} \Gamma(-\alpha + i(x+t)) \Gamma(-\alpha - i(x+t)) f(t) dt,$$

where $\operatorname{Re} \alpha < 0$ ($2\alpha \neq -1, -2, \dots$).

⊙ References: J. Wimp (1971), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 453).

3.7-4. Kernels Containing Incomplete Gamma Functions.

13.
$$\int_{-\infty}^{\infty} (t-x)^{\alpha-1} \gamma(1-\alpha, 2i(t-x)) y(t) dt = f(x), \quad i^2 = -1.$$

Here $\gamma(\nu, z)$ is the incomplete gamma function (see Supplement 11.5-1).

Solution:

$$y(x) = -\frac{1}{4\pi^2} \int_{-\infty}^{\infty} (t-x)^{-\alpha-1} \gamma(1+\alpha, 2i(t-x)) f(t) dt,$$

where $-1/2 < \operatorname{Re} \alpha \leq 0$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 462).

14.
$$\int_{-\infty}^{\infty} \left[\exp\left(\frac{2x-i}{4}\pi\right) t^{-ix-1/2} + (b-a)a^{ix-1/2} e^{iat} \Gamma\left(\frac{1}{2} - ix, iat\right) \right] y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{4\pi} \int_{-\infty}^{\infty} \left[\exp\left(\frac{2t+i}{4}\pi\right) x^{it-1/2} + (a-b)b^{-it-1/2} e^{-ibx} \Gamma\left(\frac{1}{2} + it, -ibx\right) \right] \frac{f(t)}{\cosh(\pi t)} dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 463).

$$15. \int_0^{\infty} \left\{ t^{-ix-1/2} \sin\left(\frac{1+2ix}{4}\pi\right) + \frac{i}{2}(b-a)a^{ix-1/2} \left[e^{-iat}\Gamma\left(\frac{1}{2}-ix, -iat\right) - e^{iat}\Gamma\left(\frac{1}{2}-ix, iat\right) \right] \right\} y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{1}{\pi} \int_{-\infty}^{\infty} \left\{ t^{ix-1/2} \sin\left(\frac{1-2ix}{4}\pi\right) + \frac{i}{2}(a-b)b^{-ix-1/2} \left[e^{-ibt}\Gamma\left(\frac{1}{2}+ix, -ibt\right) - e^{ibt}\Gamma\left(\frac{1}{2}+ix, ibt\right) \right] \right\} \frac{f(x)}{\cosh(\pi x)} dx,$$

where $a, b \notin (-\infty, 0)$ are complex numbers.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 463).

$$16. \int_0^{\infty} \left\{ t^{-ix-1/2} \cos\left(\frac{1+2ix}{4}\pi\right) + \frac{1}{2}(b-a)a^{ix-1/2} \left[e^{-iat}\Gamma\left(\frac{1}{2}-ix, -iat\right) + e^{iat}\Gamma\left(\frac{1}{2}-ix, iat\right) \right] \right\} y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{1}{\pi} \int_{-\infty}^{\infty} \left\{ t^{ix-1/2} \cos\left(\frac{1-2ix}{4}\pi\right) + \frac{1}{2}(a-b)b^{-ix-1/2} \left[e^{-ibt}\Gamma\left(\frac{1}{2}+ix, -ibt\right) + e^{ibt}\Gamma\left(\frac{1}{2}+ix, ibt\right) \right] \right\} \frac{f(x)}{\cosh(\pi x)} dx,$$

where $a, b \notin (-\infty, 0)$ are complex numbers.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 463).

3.7-5. Kernels Containing Bessel Functions of the First Kind.

$$17. \int_0^{\infty} t J_{\nu}(xt) y(t) dt = f(x).$$

Here $J_{\nu}(z)$ is the Bessel function of the first kind (see Supplement 11.6-1).

Solution:

$$y(x) = \begin{cases} \int_0^{\infty} t J_{\nu}(xt) f(t) dt & \text{if } \operatorname{Re} \nu \geq -1 \text{ or } \nu = -2, -3, \dots, \\ \int_0^{\infty} t \left[J_{\nu}(xt) - \sum_{k=0}^{n-1} \frac{(-1)^k (xt/2)^{2k+\nu}}{k! \Gamma(\nu+k+1)} \right] f(t) dt & \text{if } \operatorname{Re} \nu < -1 \text{ and } \nu \neq -2, -3, \dots, \\ \text{where } -n-1 < \operatorname{Re} \nu < -n, \quad n = 1, 2, \dots \end{cases}$$

The functions $f(x)$ and $y(x)$ are the *Hankel transform pair*.

⊙ References: E. C. Titchmarsh (1923), J. L. Griffith (1958), V. A. Ditkin and A. P. Prudnikov (1965), F. Oberhettinger (1972), I. Sneddon (1972), H. M. Srivastava and R. G. Buschman (1977), B. Davis (1978), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 468), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993), I. Sneddon (1995).

$$18. \int_a^b t J_\nu(xt) y(t) dt = f(x), \quad 0 \leq x < \infty.$$

Solution:

$$y(t) = \begin{cases} \int_0^\infty x J_\nu(xt) f(x) dx & \text{if } a < t < b, \\ 0 & \text{if } 0 < t < a \text{ or } t > b, \end{cases}$$

where $0 \leq a \leq b \leq \infty$ and $\operatorname{Re} \nu > -1$.

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 468), I. N. Sneddon (1995).

$$19. \int_0^\infty t J_0(xt) y(t) dt = 0, \quad a \leq x < \infty.$$

Homogeneous integral equation of the first kind.

Solution:

$$y(t) = \int_0^a \cos(xt) \varphi(x) dx,$$

where $\varphi(x)$ is an arbitrary continuously differentiable function.

⊙ Reference: Ya. S. Uflyand (1977).

$$20. \int_0^\infty t J_\nu(xt) y(t) dt = 0, \quad a \leq x < \infty.$$

Homogeneous integral equation of the first kind, $\operatorname{Re} \nu > -1/2$.

Solution:

$$y(t) = \sqrt{\frac{\pi t}{2}} \int_0^a \sqrt{x} J_{\nu-1/2}(xt) \varphi(x) dx,$$

where $\varphi(x)$ is an arbitrary continuously differentiable function.

⊙ Reference: Ya. S. Uflyand (1977).

$$21. \int_a^b |J_\nu(\lambda x) - J_\nu(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = J_\nu(\lambda x)$, where $J_\nu(z)$ is the Bessel function of the first kind.

$$22. \int_0^\infty J_\nu(\lambda(x-t)) y(t) dt = f(x).$$

1°. If $|\operatorname{Re} \nu| < 1$ and $f(0) = f'(0) = 0$ then

$$y(x) = \int_0^x J_{-\nu}(\lambda(x-t)) \left(\frac{d^2}{dt^2} + \lambda^2 \right) f(t) dt.$$

2°. If $\nu = n$ is a positive integer number and $f(0) = f'(0) = \dots = f^{(n-1)}(0) = 0$ then

$$y(x) = \frac{1}{\lambda^n} \sum_{k=0}^{[(n-1)/2]} C_n^{2k+1} \left(\frac{d}{dx} \right)^{n-2k-1} \left(\frac{d^2}{dx^2} + \lambda^2 \right)^{k+1} f(x) \\ + \frac{1}{\lambda^n} \int_0^x J_0(\lambda(x-t)) \sum_{k=0}^{[n/2]} C_n^{2k} \left(\frac{d}{dt} \right)^{n-2k} \left(\frac{d^2}{dt^2} + \lambda^2 \right)^{k+1} f(t) dt,$$

where $[A]$ stands for the integer part of the number A and $C_n^k = \frac{n!}{k!(n-k)!}$ are binomial coefficients ($0! = 1$).

3°. If ν is not an integer, $m - 1 < \operatorname{Re} \nu < m$ ($m = 0, 1, 2, \dots$), and $f(0) = f'(0) = \dots = f^{(m+1)}(0) = 0$ then

$$y(x) = \frac{m - \nu}{\lambda^m} \int_0^x \frac{J_{m-\nu}(\lambda(x-t))}{x-t} \sum_{k=0}^{[(m-1)/2]} C_m^{2k+1} \left(\frac{d}{dt}\right)^{m-2k-1} \left(\frac{d^2}{dt^2} + \lambda^2\right)^{k+1} f(t) dt \\ + \frac{1}{\lambda^m} \int_0^x J_{m-\nu}(\lambda(x-t)) \sum_{k=0}^{[m/2]} C_m^{2k} \left(\frac{d}{dt}\right)^{m-2k} \left(\frac{d^2}{dt^2} + \lambda^2\right)^{k+1} f(t) dt.$$

⊙ References: H. M. Srivastava and R. G. Buschman (1977), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 470), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

23. $\int_{-\infty}^{\infty} |x-t|^\nu J_\nu(\lambda|x-t|)y(t) dt = f(x).$

Solution:

$$y(x) = -\frac{\lambda \cos(\nu\pi)}{4 \sin^2(\nu\pi)} \int_{-\infty}^{\infty} \frac{\operatorname{sign}(t-x)}{|t-x|^{2\nu+1}} \frac{d}{dt} \left[|t-x|^{\nu+1} J_{-\nu-1}(\lambda|t-x|)f(t) \right] dt,$$

where $0 < \operatorname{Re} \nu < 1/2$.

⊙ References: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 469), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

24. $\int_0^\infty J_{n/2-1}(2\pi xt)G(x,t)y(t) dt = f(x), \quad G(x,t) = 2\pi x(t/x)^{n/2}, \quad n = 1, 2, \dots$

Solution:

$$y(x) = \int_0^\infty J_{n/2-1}(2\pi xt)G(x,t)f(t) dt.$$

The functions $f(x)$ and $y(t)$ are the *Bochner transform pair*.

⊙ Reference: Yu. A. Brychkov and A. P. Prudnikov (1979).

25. $\int_0^\infty \frac{d}{dx} \left[x J_\nu^2(xt) \right] t y(t) dt = f(x).$

Solution:

$$y(x) = -2\pi \int_0^\infty t J_\nu(x,t) Y_\nu(xt) f(t) dt \\ = \pi \int_0^\infty t \left\{ \sin(2\nu\pi) [J_{-\nu}^2(xt) - Y_\nu^2(xt)] - 2 \cos(2\nu\pi) J_{-\nu}(xt) Y_{-\nu}(xt) \right\} f(t) dt.$$

⊙ References: I. I. Hirschman and D. V. Widder (1955), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 474).

26. $\int_0^\infty t [J_{-\mu}(xt)J_{-\nu}(xt) \pm J_\mu(xt)J_\nu(xt)] y(t) dt = f(x).$

Solution:

$$y(x) = \frac{\pi}{2 \cos\left[\frac{\pi}{2}(\nu \pm \mu)\right] \sin\left[\frac{\pi}{2}(\nu \mp \mu)\right]} \int_0^\infty t \frac{d}{dt} \left[t (J_\mu(xt)J_{-\nu}(xt) \mp J_{-\mu}(xt)J_\nu(xt)) \right] f(t) dt,$$

where $\operatorname{Re}(\mu + \nu) < 3/2$.

⊙ References: I. I. Hirschman and D. V. Widder (1955), E. C. Titchmarsh (1986), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 475).

$$27. \int_0^{\infty} [J_{ix}(t) + J_{-ix}(t)]y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(x) = \frac{1}{2x} \int_0^{\infty} \frac{t[J_{it}(x) + J_{-it}(x)]}{\sinh(\pi t)} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 469).

$$28. \int_0^{\infty} [J_{it}(x) + J_{-it}(x)]y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(x) = \frac{x}{2 \sinh(\pi x)} \int_0^{\infty} \frac{J_{ix}(t) + J_{-ix}(t)}{t} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 469).

3.7-6. Kernels Containing Bessel Functions of the Second Kind.

$$29. \int_0^{\infty} tY_{\nu}(xt)y(t) dt = f(x).$$

Here $Y_{\nu}(z)$ is the Bessel function of the second kind (see Supplement 11.6-1).

1°. If $|\operatorname{Re} \nu| < 1$ then

$$y(x) = \int_0^{\infty} t\mathbf{H}_{\nu}(xt)f(t) dt,$$

where $\mathbf{H}_{\nu}(x)$ is the Struve function, which is defined as

$$\mathbf{H}_{\nu}(x) = \sum_{j=0}^{\infty} \frac{(-1)^j (x/2)^{\nu+2j+1}}{\Gamma(j + \frac{3}{2})\Gamma(\nu + j + \frac{3}{2})}.$$

The function $f(x)$ and the solution $y(x)$ are the Y_{ν} -transform pair.

2°. If $1 < |\operatorname{Re} \nu| < 3$ then

$$y(x) = \int_0^{\infty} t \left[\mathbf{H}_{\nu}(xt) - \frac{(xt)^{\nu-1}}{2^{\nu-1} \sqrt{\pi} \Gamma(\nu + 1/2)} \right] f(t) dt.$$

⊙ References: E. C. Titchmarsh (1948), G. N. Watson (1952), J. L. Griffith (1958), F. Oberhettinger (1972), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 475).

$$30. \int_a^b |Y_{\nu}(\lambda x) - Y_{\nu}(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = Y_{\nu}(\lambda x)$, where $Y_{\nu}(z)$ is the Bessel function of the second kind.

3.7-7. Kernels Containing Combinations of the Bessel Functions.

$$31. \int_0^{\infty} [\cos(p\pi)J_{\nu}(xt) + \sin(p\pi)Y_{\nu}(xt)]ty(t) dt = f(x).$$

Solution:

$$y(x) = \int_0^{\infty} \Phi(xt)tf(t) dt, \quad \Phi(z) = \sum_{n=0}^{\infty} \frac{(-1)^n(z/2)^{\nu+2p+2n}}{\Gamma(p+n+1)\Gamma(\nu+p+n+1)}.$$

The functions $f(x)$ and $y(x)$ are the *Hardy transform pair*.

⊙ Reference: Yu. A. Brychkov and A. P. Prudnikov (1989).

$$32. \int_0^{\infty} tJ_{\nu}(xt)Y_{\nu}(xt)y(t) dt = f(x).$$

Solution:

$$y(x) = 2\pi \int_0^{\infty} t \frac{d}{dt} [tJ_{\nu}^2(xt)] f(t) dt,$$

where $\operatorname{Re} \nu > -1/4$.

⊙ References: E. C. Titchmarsh (1948), I. I. Hirschman and D. V. Widder (1955), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 476).

$$33. \int_a^{\infty} t[J_{\nu}(ax)Y_{\nu}(xt) - Y_{\nu}(ax)J_{\nu}(xt)]y(t) dt = f(x).$$

Solution:

$$y(x) = \int_0^{\infty} \frac{t[J_{\nu}(at)Y_{\nu}(xt) - Y_{\nu}(at)J_{\nu}(xt)]}{J_{\nu}^2(at) + Y_{\nu}^2(at)} f(t) dt.$$

The function $f(x)$ and the solution $y(x)$ are the *Weber transform pair*.

⊙ References: G. N. Watson (1952), Yu. A. Brychkov and A. P. Prudnikov (1979, 1989), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 477).

$$34. \int_0^{\infty} t[J_{\nu}(at)Y_{\nu}(xt) - Y_{\nu}(at)J_{\nu}(xt)]y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{x}{J_{\nu}^2(ax) + Y_{\nu}^2(ax)} \int_0^{\infty} t[J_{\nu}(ax)Y_{\nu}(xt) - Y_{\nu}(ax)J_{\nu}(xt)]f(t) dt.$$

⊙ References: G. N. Watson (1952), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 477).

$$35. \int_{-\infty}^{\infty} e^{\pm\pi(x-t)/2} H_{i(t-x)}^{(1)}(a)y(t) dt = f(x), \quad i^2 = -1.$$

Here $H_{\nu}^{(1)}(z) = J_{\nu}(z) + iY_{\nu}(z)$ is the Hankel function of the first kind (see Supplement 11.6-5).

Solution:

$$y(x) = \frac{1}{4} \int_{-\infty}^{\infty} e^{\pm\pi(t-x)/2} H_{i(t-x)}^{(1)}(a)f(t) dt,$$

where $a > 0$.

⊙ References: Vu Kim Tuan (1988), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 479).

$$36. \int_{-\infty}^{\infty} e^{\pm\pi(x-t)/2} H_{i(t-x)}^{(2)}(a)y(t) dt = f(x).$$

Here $H_{\nu}^{(2)}(z) = J_{\nu}(z) - iY_{\nu}(z)$ is the Hankel function of the second kind (see Supplement 11.6-5).

Solution:

$$y(x) = \frac{1}{4} \int_{-\infty}^{\infty} e^{\pm\pi(t-x)/2} H_{i(t-x)}^{(2)}(a)f(t) dt,$$

where $a > 0$.

⊙ References: Vu Kim Tuan (1988), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 479).

3.7-8. Kernels Containing Modified Bessel Functions of the First Kind.

$$37. \int_a^b |I_{\nu}(\lambda x) - I_{\nu}(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = I_{\nu}(\lambda x)$, where $I_{\nu}(z)$ is the modified Bessel function of the first kind (see Supplement 11.7-1).

$$38. \int_0^{\infty} \frac{d}{dx} I_{it}^2(x)y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(x) = \frac{2i}{\pi} x \int_0^{\infty} K_{ix}^2(t)f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 485).

$$39. \int_{-\infty}^{\infty} \text{Ai}(x+t)y(t) dt = f(x).$$

Here $\text{Ai}(x) = \frac{1}{3}\sqrt{x} [L_{-1/3}(z) - I_{1/3}(z)]$ is the Airy function (see Supplement 11.8-1).

Solution:

$$y(x) = \int_{-\infty}^{\infty} \text{Ai}(x+t)f(t) dt.$$

⊙ References: Vu Kim Tuan (1988), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 485).

3.7-9. Kernels Containing Modified Bessel Functions of the Second Kind.

$$40. \int_{-\infty}^{\infty} K_0(|x-t|)y(t) dt = f(x).$$

Here $K_0(z)$ is the modified Bessel function of the second kind (the MacDonald function), see Supplement 11.7-1.

Solution:

$$y(x) = -\frac{1}{\pi^2} \left(\frac{d^2}{dx^2} - 1 \right) \int_{-\infty}^{\infty} K_0(|x-t|)f(t) dt.$$

⊙ Reference: D. Naylor (1986).

$$41. \int_a^b |K_{\nu}(\lambda x) - K_{\nu}(\lambda t)| y(t) dt = f(x).$$

This is a special case of equation 3.8.3 with $g(x) = K_{\nu}(\lambda x)$.

$$42. \int_0^{\infty} \sqrt{zt} K_{\nu}(zt) y(t) dt = f(z).$$

Here $K_{\nu}(z)$ is the modified Bessel function of the second kind.

Up to a constant factor, the left-hand side of this equation is the Meijer transform of $y(t)$ (z is treated as a complex variable).

Solution:

$$y(t) = \frac{1}{\pi i} \int_{c-i\infty}^{c+i\infty} \sqrt{zt} I_{\nu}(zt) f(z) dz.$$

For specific $f(z)$, one may use tables of Meijer integral transforms to calculate the integral.

⊙ Reference: V. A. Ditkin and A. P. Prudnikov (1965).

$$43. \int_0^{\infty} K_{ix}(t) y(t) dt = f(x), \quad i^2 = -1.$$

Solution:

$$y(x) = \frac{2}{\pi^2 x} \int_0^{\infty} t \sinh(\pi t) K_{it}(x) f(t) dt.$$

The function $f(x)$ and the solution $y(x)$ are the *Kontorovich-Lebedev transform pair*.

⊙ References: V. A. Ditkin and A. P. Prudnikov (1965), F. Oberhettinger (1972), Yu. A. Brychkov and A. P. Prudnikov (1989), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 487).

$$44. \int_0^{\infty} K_{it}(x) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{2x \sinh(\pi x)}{\pi^2} \int_0^{\infty} \frac{K_{ix}(t)}{t} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 487).

$$45. \int_0^{\infty} K_{it}^2(x) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{4x \sinh(\pi x)}{\pi^2} \int_0^{\infty} \frac{d}{dt} \{ [I_{ix}(t) + I_{-ix}(t)] K_{ix}(t) \} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 492).

$$46. \int_0^{\infty} \operatorname{Re} K_{ix+1/2}(t) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{4}{\pi^2} \int_0^{\infty} \cosh(\pi t) \operatorname{Re} K_{it+1/2}(x) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 488).

$$47. \int_0^{\infty} \operatorname{Im} K_{ix+1/2}(t) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{4}{\pi^2} \int_0^{\infty} \cosh(\pi t) \operatorname{Im} K_{it+1/2}(x) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 488).

$$48. \int_0^{\infty} \operatorname{Re} K_{it+1/2}(x)y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{4}{\pi^2} \cosh(\pi x) \int_0^{\infty} \operatorname{Re} K_{ix+1/2}(t)f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 488).

$$49. \int_0^{\infty} \operatorname{Im} K_{it+1/2}(x)y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{4}{\pi^2} \cosh(\pi x) \int_0^{\infty} \operatorname{Im} K_{ix+1/2}(t)f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 488).

$$50. \int_{-\infty}^{\infty} e^{\pi(x+t)/2} K_{i(x+t)}(a)y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi^2} \int_{-\infty}^{\infty} e^{\pi(x+t)/2} K_{i(x+t)}(a)f(t) dt,$$

where $a > 0$. The function $f(x)$ and the solution $y(x)$ are a *Crum transform pair* (in the asymmetric form).

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 488).

$$51. \int_{-\infty}^{\infty} K_{i(x+t)}(\pm ia)y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi^2} \int_{-\infty}^{\infty} K_{i(x+t)}(\mp ia)f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 488–489).

$$52. \int_{-\infty}^{\infty} t^{-\frac{1}{4}(2ix+1)} K_{\frac{1}{2}+ix}(2i\lambda\sqrt{t})y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{\lambda}{\pi^2} \int_{-\infty}^{\infty} x^{\frac{1}{4}(2it-1)} K_{\frac{1}{2}-it}(2i\lambda\sqrt{x})f(t) dt,$$

where $\lambda > 0$ and $\sqrt{x} = -i\sqrt{|x|}$ for $x < 0$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 489).

$$53. \int_0^{\infty} \left[(a+t)^{-\frac{1}{4}(2ix+1)} K_{\frac{1}{2}+ix}(2i\lambda\sqrt{a+t}) \right. \\ \left. + (a-t)^{-\frac{1}{4}(2ix+1)} K_{\frac{1}{2}+ix}(2i\lambda\sqrt{a-t}) \right] y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{\lambda}{\pi^2} \int_{-\infty}^{\infty} \left[(a+t)^{\frac{1}{4}(2ix-1)} K_{\frac{1}{2}-ix}(-2i\lambda\sqrt{a+t}) + (a-t)^{\frac{1}{4}(2ix-1)} K_{\frac{1}{2}-ix}(-2i\lambda\sqrt{a-t}) \right] f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 489).

$$54. \int_{-\infty}^{\infty} x^{\frac{1}{4}(2it-1)} K_{\frac{1}{2}-it}(2i\lambda\sqrt{x}) y(t) dt = f(x), \quad \lambda > 0.$$

Solution:

$$y(x) = \frac{\lambda}{\pi^2} \int_{-\infty}^{\infty} t^{-\frac{1}{4}(2ix+1)} K_{\frac{1}{2}+ix}(2i\lambda\sqrt{t}) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 489).

$$55. \int_{-\infty}^{\infty} \left[(a+t)^{\frac{1}{4}(2it-1)} K_{\frac{1}{2}-it}(-2i\lambda\sqrt{a+x}) \right. \\ \left. + (a-t)^{\frac{1}{4}(2it-1)} K_{\frac{1}{2}-it}(-2i\lambda\sqrt{a-x}) \right] y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{\lambda}{\pi^2} \int_0^{\infty} \left[(a+x)^{-\frac{1}{4}(2it+1)} K_{\frac{1}{2}+it}(2i\lambda\sqrt{a+x}) + (a-x)^{-\frac{1}{4}(2it+1)} K_{\frac{1}{2}+it}(2i\lambda\sqrt{a-x}) \right] f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 490).

$$56. \int_{-\infty}^{\infty} \exp\left(\frac{\pi x}{2} \operatorname{sign} t\right) K_{ix}(|t|) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{1}{\pi^2 x} \int_{-\infty}^{\infty} t \exp\left(\frac{\pi t}{2} \operatorname{sign} x\right) K_{it}(|x|) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 490).

3.7-10. Kernels Containing a Combination of Bessel and Modified Bessel Functions.

$$57. \int_0^{\infty} [I_{ix}(t) + I_{-ix}(t)] K_{ix}(t) y(t) dt = f(x).$$

Solution:

$$y(x) = -\frac{4}{\pi^2} \frac{d}{dx} \int_0^{\infty} t \sinh(\pi t) K_{it}^2(x) f(t) dt.$$

The integral equation and its solution form the *Lebedev transform pair*.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 493).

$$58. \int_0^{\infty} [K_{it}(a) I_{it}(x) - I_{it}(a) K_{it}(x)] y(t) dt = f(x), \quad 0 < x < a.$$

Solution:

$$y(t) = \frac{2t \sinh(\pi t)}{\pi^2 [I_{ia}(a)]^2} \int_0^a x^{-1} [K_{it}(a) I_{it}(x) - I_{it}(a) K_{it}(x)] f(x) dx, \quad t > 0.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 494).

$$59. \int_0^{\infty} t \left[Y_0(xt) - \frac{2}{\pi} K_0(xt) \right] y(t) dt = f(x).$$

Solution:

$$y(x) = \int_0^{\infty} t \left[Y_0(xt) - \frac{2}{\pi} K_0(xt) \right] f(t) dt.$$

The integral equation and its solution form the *divisor transform pair*.

⊙ References: F. Oberhettinger (1973), E. C. Titchmarsh (1986), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 492).

$$60. \int_0^{\infty} t \left[Y_{2n+1}(xt) \pm \frac{2}{\pi} K_{2n+1}(xt) \right] y(t) dt = f(x), \quad n = 1, 2, \dots$$

Solution:

$$y(x) = \int_0^{\infty} t \left[Y_{2n+1}(xt) \mp \frac{2}{\pi} K_{2n+1}(xt) \right] f(t) dt.$$

⊙ References: E. C. Titchmarsh (1986), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 493).

$$61. \int_0^{\infty} t \left[Y_{2n}(xt) + \frac{2}{\pi} K_{2n}(xt) \right] y(t) dt = f(x), \quad n = 1, 2, \dots$$

Solution:

$$y(x) = \int_0^{\infty} t \left[Y_{2n}(xt) + \frac{2}{\pi} K_{2n}(xt) \right] f(t) dt.$$

⊙ References: E. C. Titchmarsh (1986), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 493).

3.7-11. Kernels Containing Legendre Functions.

$$62. \int_1^{\infty} P_{-\frac{1}{2}+ix}(t)y(t) dt = f(x), \quad 0 \leq x < \infty.$$

Here $P_\nu(x)$ is the Legendre function of the first kind (see Supplement 11.11-3) and $i^2 = -1$.

Solution:

$$y(t) = \int_0^{\infty} x \tanh(\pi x) P_{ix-1/2}(t) f(x) dx.$$

The functions $f(x)$ and $y(t)$ are the *Mehler–Fock transform pair*.

Remark. The Legendre function of the first kind can be represented in the form

$$P_{-\frac{1}{2}+ix}(t) = \frac{2}{\pi} \cosh(\pi x) \int_0^{\infty} \frac{\cos(xs) ds}{\sqrt{2(t + \cosh s)}}, \quad 1 \leq t < \infty.$$

⊙ References: N. N. Lebedev (1965), V. A. Ditkin and A. P. Prudnikov (1965), Yu. A. Brychkov and A. P. Prudnikov (1989), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 512).

$$63. \int_0^{\infty} P_{-\frac{1}{2}+it}(x)y(t) dt = f(x), \quad 1 \leq x < \infty.$$

Solution:

$$y(t) = t \tanh(\pi t) \int_1^{\infty} P_{-\frac{1}{2}+it}(x) f(x) dx.$$

⊙ References: N. N. Lebedev (1965), V. A. Ditkin and A. P. Prudnikov (1965), Yu. A. Brychkov and A. P. Prudnikov (1989), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 513).

$$64. \int_0^{\infty} [P_{-\frac{1}{2}+ix}(it) \pm P_{-\frac{1}{2}+ix}(-it)]y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{1}{2} \int_0^{\infty} \frac{\sinh(\pi x)}{\cosh^2(\pi x)} [P_{-\frac{1}{2}+ix}(-it) \pm P_{-\frac{1}{2}+ix}(it)] f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 513).

$$65. \int_0^{\infty} [P_{-\frac{1}{2}+it}(ix) \pm P_{-\frac{1}{2}+it}(-ix)]y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{t \sinh(\pi t)}{2 \cosh^2(\pi t)} \int_0^{\infty} [P_{-\frac{1}{2}+it}(-ix) \pm P_{-\frac{1}{2}+it}(ix)]f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 514).

$$66. \int_{-\infty}^{\infty} [ie^{-i\pi x} P_{-\frac{1}{2}+x}(\cos t) + P_{-\frac{1}{2}+x}(-\cos t)]y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{1}{2} \sin t \int_{-\infty}^{\infty} \frac{x}{\sinh(2\pi x)} [ie^{i\pi x} P_{-\frac{1}{2}+x}(\cos t) + P_{-\frac{1}{2}+x}(-\cos t)]f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 513).

$$67. \int_0^{\infty} [P_{-\frac{1}{2}+it}(x)]^2 y(t) dt = f(x), \quad 1 \leq x < \infty.$$

Solution:

$$y(t) = t \tanh(\pi t) \int_1^{\infty} P_{-\frac{1}{2}+it}(x) [Q_{-\frac{1}{2}+it}(x) + Q_{-\frac{1}{2}-it}(x)] (x^2 - 1)^{1/2} \frac{d}{dx} [(x^2 - 1)^{1/2} f(x)] dx,$$

where $Q_{\nu}(x)$ is the Legendre function of the second kind.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 514).

$$68. \int_1^{\infty} P_{-\frac{1}{2}+ix}(t) [Q_{-\frac{1}{2}+ix}(t) + Q_{-\frac{1}{2}-ix}(t)] y(t) dt = f(x), \quad 0 \leq x < \infty.$$

Here $Q_{\nu}(x)$ is the Legendre function of the second kind.

Solution:

$$y(t) = (t^2 - 1)^{1/2} \frac{d}{dt} \left[(t^2 - 1)^{1/2} \int_0^{\infty} x \tanh(\pi x) [P_{-\frac{1}{2}+ix}(t)]^2 f(x) dx \right].$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 519).

3.7-12. Kernels Containing Associated Legendre Functions.

$$69. \int_1^{\infty} P_{-\frac{1}{2}+ix}^{\mu}(t) y(t) dt = f(x), \quad 0 \leq x < \infty.$$

Here $P_{\nu}^{\mu}(x)$ is the associated Legendre function of the first kind (see Supplement 11.11-3) and $i^2 = -1$.

Solution:

$$y(t) = \frac{1}{\pi} \int_0^{\infty} x \sinh(\pi x) \Gamma\left(\frac{1}{2} - \mu + ix\right) \Gamma\left(\frac{1}{2} - \mu - ix\right) P_{ix-1/2}^{\mu}(t) f(x) dx.$$

The functions $f(x)$ and $y(t)$ are the *generalized Mehler–Fock transform pair*.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 518).

$$70. \int_0^{\infty} P_{-\frac{1}{2}+it}^{\mu}(x)y(t) dt = f(x), \quad 1 \leq x < \infty.$$

Solution:

$$y(t) = \frac{1}{\pi} t \sinh(\pi t) \Gamma\left(\frac{1}{2} - \mu + it\right) \Gamma\left(\frac{1}{2} - \mu - it\right) \int_1^{\infty} P_{it-1/2}^{\mu}(x) f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 519).

$$71. \int_{-1}^1 P_{-\frac{1}{2}+ia}^{ix}(\pm t)y(t) dt = f(x), \quad -\infty < x < \infty.$$

Solution:

$$y(t) = \frac{1}{2\pi i(1-t)} \int_{-\infty}^{\infty} x \Gamma\left(\frac{1}{2} + ia - ix\right) \Gamma\left(\frac{1}{2} - ia - ix\right) P_{-\frac{1}{2}+ia}^{ix}(\mp t) f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 518).

$$72. \int_0^{\infty} [(x+t-1)^2 - 4xt]^{-1/2} Q_{\nu-\frac{1}{2}}^1\left(\frac{x+t-1}{2\sqrt{xt}}\right) y(t) dt = f(x), \quad \operatorname{Re} \nu > -1.$$

Here $Q_{\nu}^{\mu}(x)$ is the associated Legendre function of the second kind (see Supplement 11.11-3).

Solution:

$$y(t) = \frac{1}{4\pi^2} \int_0^{\infty} (xt)^{-1/2} [(x+t-1)^2 - 4xt]^{-1/2} Q_{\nu-\frac{1}{2}}^1\left(\frac{x+t-1}{2\sqrt{xt}}\right) f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 520).

3.7-13. Kernels Containing Kummer Confluent Hypergeometric Functions.

$$73. \int_0^{\infty} F(a, b; ixt)y(t) dt = f(x).$$

Here $F(a, b; x)$ is the Kummer confluent hypergeometric function (see Supplement 11.9-1) and $i^2 = -1$.

Let $\operatorname{Re}(b-a) < n < \operatorname{Re} b - 1/2$. Then the solution is

$$y(t) = \frac{\Gamma(a)}{2\pi\Gamma(b)} t^{b-1} \left(\frac{d}{dt}\right)^n \left[t^{n-b+1} \int_{-\infty}^{\infty} e^{-ixt} \Psi(n+a-b, n-b+2; ixt) f(x) dx \right].$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 530).

$$74. \int_0^{\infty} F\left(\frac{1}{2}b \pm ix, b; -it\right) y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{t^{b-1}}{2\pi\Gamma^2(b)} \int_{-\infty}^{\infty} e^{\mp\pi x} \Gamma\left(\frac{1}{2}b + ix\right) \Gamma\left(\frac{1}{2}b - ix\right) F\left(\frac{1}{2}b \mp ix, b; it\right) f(x) dx,$$

where $\operatorname{Re} b > 0$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 531).

$$75. \int_0^{\infty} t^{ix} F\left(\frac{1}{2} + ix, b + ix; i\alpha t\right) y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{t^{b-1}}{2\pi} \left(-\frac{d}{dt}\right)^n \int_{-\infty}^{\infty} t^{n-b-ix} e^{-i\alpha t} \frac{\Gamma\left(\frac{1}{2} + ix\right)}{\Gamma(b + ix)} \Psi\left(n - b + \frac{1}{2}, n - b + 1 - ix; i\alpha t\right) f(x) dx,$$

where $\text{Im } \alpha = 0$ and $0 < \text{Re } b - 1/2 < n < \text{Re } b$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 531).

$$76. \int_{-\infty}^{\infty} F(a, b; i\beta(x-t)) y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{\beta^2(a-1)(a-b+1) \sin(\pi b)}{4\pi(b-1)(b-2)(b-3) \sin(\pi a) \sin[\pi(b-a)]} \int_{-\infty}^{\infty} F(2-a, b-a-1; i\beta(x-t)) f(x) dx,$$

where $1 < \text{Re } a < 3/2$ and $-1 < \text{Re}(b-a) < -1/2$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 531).

$$77. \int_{-\infty}^{\infty} F\left(\frac{1}{2} \pm ia, \frac{1}{2}; \pm i(x-t)^2\right) y(t) dt = f(x), \quad a > 0.$$

Solution:

$$y(t) = \frac{e^{\pi a}}{\pi \cosh(\pi a)} \int_{-\infty}^{\infty} F\left(\frac{1}{2} \mp ia, \frac{1}{2}; \mp i(x-t)^2\right) f(x) dx.$$

⊙ References: Vu Kim Tuan (1988), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 532).

$$78. \int_{-\infty}^{\infty} F\left(\frac{1}{2}b \pm it, b; ix\right) y(t) dt = f(x), \quad \text{Re } b > 0.$$

Solution:

$$y(t) = \frac{e^{\pm\pi t}}{2\pi\Gamma^2(b)} \Gamma\left(\frac{1}{2}b + it\right) \Gamma\left(\frac{1}{2}b - it\right) \int_0^{\infty} x^{b-1} F\left(\frac{1}{2}b \mp it, b; -ix\right) f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 532).

$$79. \int_{-\infty}^{\infty} F\left(\frac{1}{2}b \pm it, b; -ix\right) y(t) dt = f(x), \quad \text{Re } b > 0.$$

Solution:

$$y(t) = \frac{e^{\mp\pi t}}{2\pi\Gamma^2(b)} \Gamma\left(\frac{1}{2}b + it\right) \Gamma\left(\frac{1}{2}b - it\right) \int_0^{\infty} x^{b-1} F\left(\frac{1}{2}b \mp it, b; ix\right) f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 532).

$$80. \int_{-\infty}^{\infty} x^{-it} F\left(\frac{1}{2} - it, b - it; i\beta x\right) y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{\Gamma\left(\frac{1}{2}(1-it)\right)}{2\pi\Gamma\left(b - \frac{1}{2}it\right)} \int_0^{\infty} x^{n-b+it} e^{-i\beta x} \Psi\left(n + \frac{1}{2} - b, n + 1 - b + it; i\beta x\right) \left(\frac{d}{dx}\right)^n [x^{b-1} f(x)] dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 533).

3.7-14. Kernels Containing Tricomi Confluent Hypergeometric Functions.

$$81. \int_0^{\infty} t^{ix} \Psi(a + ix, 2ix + 1; t) y(t) dt = f(x).$$

Here $\Psi(a, b; x)$ is the Tricomi confluent hypergeometric function (see Supplement 11.9-1) and $i^2 = -1$.

Solution:

$$y(t) = \frac{e^{-t}}{\pi^2 t} \int_0^{\infty} x \sinh(2\pi x) \Gamma(a - ix) \Gamma(a + ix) t^{ix} \Psi(a + ix, 2ix + 1; t) f(x) dx.$$

⊙ References: J. Wimp (1971), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 534).

$$82. \int_0^{\infty} x^{it} \Psi(a + it, 2it + 1; t) y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{t}{\pi^2} \sinh(2\pi t) \Gamma(a - it) \Gamma(a + it) \int_0^{\infty} x^{-1+it} e^{-x} \Psi(a + it, 2it + 1; x) f(x) dx.$$

⊙ References: J. Wimp (1971), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 535).

$$83. \int_{-\infty}^{\infty} \Psi\left(\frac{1}{2} + ix, \frac{3}{2} - i\beta + ix; \pm it\right) y(t) dt = f(x), \quad \mathbf{Im} \beta = 0.$$

Solution:

$$y(t) = \frac{1}{4\pi} \int_{-\infty}^{\infty} \frac{1}{\cosh(\pi x)} \Psi\left(\frac{1}{2} - ix, \frac{3}{2} + i\beta - ix; \mp it\right) f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 536).

3.7-15. Kernels Containing Whittaker Confluent Hypergeometric Functions.

$$84. \int_0^{\infty} M_{\pm ix, \nu}(it) y(t) dt = f(x), \quad \mathbf{Re} \nu > -\frac{1}{2}.$$

Here $M_{\mu, \nu}(z)$ is the Whittaker confluent hypergeometric function (see Supplement 11.9-3) and $i^2 = -1$.

Solution:

$$y(t) = \frac{1}{2\pi \Gamma^2(2\nu + 1) t} \int_{-\infty}^{\infty} e^{\mp \pi x} \Gamma\left(\frac{1}{2} + \nu + ix\right) \Gamma\left(\frac{1}{2} + \nu - ix\right) M_{\pm ix, \nu}(-it) f(x) dx.$$

The integral equation and its solution form the *Buchholz transform pair*.

⊙ References: H. Buchholz (1969), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 523).

$$85. \int_0^{\infty} M_{\pm ix, \nu}(-it) y(t) dt = f(x), \quad \mathbf{Re} \nu > -\frac{1}{2}.$$

Solution:

$$y(t) = \frac{1}{2\pi \Gamma^2(2\nu + 1) t} \int_{-\infty}^{\infty} e^{\pm \pi x} \Gamma\left(\frac{1}{2} + \nu + ix\right) \Gamma\left(\frac{1}{2} + \nu - ix\right) M_{\mp ix, \nu}(it) f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 523–524).

$$86. \int_{-\infty}^{\infty} M_{\pm it, \nu}(ix)y(t) dt = f(x), \quad \operatorname{Re} \nu > -\frac{1}{2}.$$

Solution:

$$y(t) = \frac{e^{\mp \pi t}}{2\pi\Gamma^2(2\nu+1)}\Gamma\left(\frac{1}{2} + \nu + it\right)\Gamma\left(\frac{1}{2} + \nu - it\right) \int_0^{\infty} x^{-1} M_{\mp it, \nu}(-ix)f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 524).

$$87. \int_{-\infty}^{\infty} M_{\pm it, \nu}(-ix)y(t) dt = f(x), \quad \operatorname{Re} \nu > -\frac{1}{2}.$$

Solution:

$$y(t) = \frac{e^{\pm \pi t}}{2\pi\Gamma^2(2\nu+1)}\Gamma\left(\frac{1}{2} + \nu + it\right)\Gamma\left(\frac{1}{2} + \nu - it\right) \int_0^{\infty} x^{-1} M_{\mp it, \nu}(ix)f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 524–525).

$$88. \int_{-\infty}^{\infty} \Gamma\left(\frac{1}{2} + \nu + ix - it\right)\Gamma\left(\frac{1}{2} + \nu - ix + it\right) M_{it-ix, \nu}(a)y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{(2\nu+1)\sin(2\pi\nu)}{4\pi^3} \int_{-\infty}^{\infty} \Gamma\left(-\frac{1}{2} - \nu + ix - it\right)\Gamma\left(-\frac{1}{2} - \nu - ix + it\right) M_{it-ix, -\nu-1}(a)f(x) dx.$$

⊙ References: J. Wimp (1971), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 526).

$$89. \int_0^{\infty} W_{\mu, ix}(t)y(t) dt = f(x).$$

Here $W_{\mu, \nu}(z)$ is the Whittaker confluent hypergeometric function (see Supplement 11.9-3).

Solution:

$$y(t) = \frac{1}{\pi^2 t^2} \int_0^{\infty} x \sinh(2\pi x)\Gamma\left(\frac{1}{2} - \mu - ix\right)\Gamma\left(\frac{1}{2} - \mu + ix\right) W_{\mu, ix}(t)f(x) dx.$$

⊙ References: J. Wimp (1971), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 527).

$$90. \int_0^{\infty} W_{\mu, it}(x)y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{t}{\pi^2} \sinh(2\pi t)\Gamma\left(\frac{1}{2} - \mu - it\right)\Gamma\left(\frac{1}{2} - \mu + it\right) \int_0^{\infty} x^{-2} W_{\mu, it}(x)f(x) dx.$$

⊙ References: J. Wimp (1971), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 527).

$$91. \int_{-\infty}^{\infty} e^{-ixt/2} W_{\mu, \nu}(ixt)y(t) dt = f(x).$$

Solution:

$$y(t) = \frac{\Gamma\left(\frac{3}{2} - \mu - \nu\right)}{2\pi\Gamma(1+n-2\nu)}(it)^{-n/2-1} \\ \times \int_0^{\infty} x^{(n-1)/2-\nu} e^{ixt/2} W_{\mu+n/2-1, n/2-\nu}(ixt) \left(\frac{d}{dx}\right)^n [x^{\nu-1/2} f(x)] dx,$$

where $\operatorname{Re} \mu < \operatorname{Re} \nu + 1/2 < 3/4$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 528).

3.7-16. Kernels Containing Gauss Hypergeometric Functions.

$$92. \int_0^a F\left(\frac{\beta}{2}, \frac{\beta+1}{2}, \mu; \frac{4x^2t^2}{(x^2+t^2)^2}\right) \frac{y(t) dt}{(x^2+t^2)^\beta} = f(x).$$

Here $0 < a \leq \infty$, $0 < \beta < \mu < \beta + 1$, and $F(a, b, c; z)$ is the Gauss hypergeometric function (see Supplement 11.10-1).

1°. Solution:

$$y(x) = \frac{x^{2\mu-2}}{\Gamma(1+\beta-\mu)} \frac{d}{dx} \int_x^a \frac{tg(t) dt}{(t^2-x^2)^{\mu-\beta}},$$

$$g(t) = \frac{2\Gamma(\beta) \sin[(\beta-\mu)\pi]}{\pi\Gamma(\mu)} t^{1-2\beta} \frac{d}{dt} \int_0^t \frac{s^{2\mu-1} f(s) ds}{(t^2-s^2)^{\mu-\beta}}.$$

2°. If $a = \infty$ and $f(x)$ is a differentiable function, then the solution can be represented in the form

$$y(x) = A \frac{d}{dt} \int_0^\infty \frac{(xt)^{2\mu} f'_t(t)}{(x^2+t^2)^{2\mu-\beta}} F\left(\mu - \frac{\beta}{2}, \mu + \frac{1-\beta}{2}, \mu + 1; \frac{4x^2t^2}{(x^2+t^2)^2}\right) dt,$$

where $A = \frac{\Gamma(\beta) \Gamma(2\mu - \beta) \sin[(\beta - \mu)\pi]}{\pi\Gamma(\mu) \Gamma(1 + \mu)}$.

⊙ Reference: P. P. Zabreyko, A. I. Koshelev, et al. (1975).

$$93. \int_0^\infty F(a+ix, a-ix, c; -t)y(t) dt = f(x), \quad a, c > 0.$$

Solution:

$$y(t) = \frac{t^{c-1}(1+t)^{2a-c}}{\pi^2\Gamma^2(c)} \int_0^\infty x \sinh(2\pi x) |\Gamma(a+ix)\Gamma(c-a+ix)|^2 F(a+ix, a-ix, c; -t) f(x) dx.$$

The integral equation and its solution form the *Olevskii transform pair*.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 538).

3.7-17. Kernels Containing Parabolic Cylinder Functions.

$$94. \int_{-\infty}^\infty D_{-ix-1/2}(\pm e^{-\pi i/4}t)y(t) dt = f(x), \quad i^2 = -1.$$

Here $D_\nu(z)$ is the parabolic cylinder function (see Supplement 11.12-1).

Solution:

$$y(x) = \frac{1}{4\pi} \int_{-\infty}^\infty \frac{e^{-\pi t/2}}{\cosh(\pi t)} D_{it-1/2}(\pm e^{\pi i/4}x) f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 467).

$$95. \int_{-\infty}^\infty \exp\left[\pm \frac{i(x-t)^2}{4}\right] [D_{\pm i\alpha}(e^{\mp \pi i/4}(t-x)) - D_{\pm i\alpha}(e^{\mp \pi i/4}(x-t))] y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{e^{\pi\alpha/2}}{8\pi \cosh^2(\pi\alpha/2)} \int_{-\infty}^\infty \exp\left[\mp \frac{i(x-t)^2}{4}\right] [D_{\mp i\alpha}(e^{\pm \pi i/4}(t-x)) + D_{\mp i\alpha}(e^{\pm \pi i/4}(x-t))] f(t) dt,$$

where $\alpha > 0$.

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 466).

$$96. \int_0^\infty \left\{ \exp \left[\frac{i(x+t)^2}{4} \right] [D_{2i\alpha}(e^{3\pi i/4}(x+t)) - D_{2i\alpha}(e^{-\pi i/4}(x+t))] \right. \\ \left. + \exp \left[\frac{i(x-t)^2}{4} \right] [D_{2i\alpha}(e^{3\pi i/4}(x-t)) - D_{2i\alpha}(e^{-\pi i/4}(x-t))] \right\} y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{e^{\pi\alpha}}{8\pi \sinh^2(\pi\alpha)} \int_0^\infty \left\{ \exp \left[-\frac{i(x+t)^2}{4} \right] [D_{-2i\alpha}(-e^{\pi i/4}(x+t)) - D_{-2i\alpha}(e^{\pi i/4}(x+t))] \right. \\ \left. + \exp \left[-\frac{i(t-x)^2}{4} \right] [D_{-2i\alpha}(-e^{\pi i/4}(t-x)) - D_{-2i\alpha}(e^{\pi i/4}(t-x))] \right\} f(t) dt.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, pp. 465–466).

3.7-18. Kernels Containing Other Special Functions.

$$97. \int_0^a K \left(\frac{2\sqrt{xt}}{x+t} \right) \frac{y(t) dt}{x+t} = f(x).$$

Here $K(z) = \int_0^1 \frac{dt}{\sqrt{(1-t^2)(1-z^2t^2)}}$ is the complete elliptic integral of the first kind (see Supplement 11.13-1).

Solution:

$$y(x) = -\frac{4}{\pi^2} \frac{d}{dx} \int_x^a \frac{tF(t) dt}{\sqrt{t^2-x^2}}, \quad F(t) = \frac{d}{dt} \int_0^t \frac{sf(s) ds}{\sqrt{t^2-s^2}}.$$

⊙ Reference: P. P. Zabreyko, A. I. Koshelev, et al. (1975).

$$98. \int_0^\infty \left[\zeta \left(\frac{1}{2} + ix, it \right) - \zeta \left(\frac{1}{2} + ix, \frac{1}{2} + it \right) \right] y(t) dt = f(x).$$

Here $\zeta(z, v) = \sum_{k=0}^\infty \frac{1}{(v+k)^z}$ is the generalized Riemann zeta function ($\operatorname{Re} z > 1$; $v \neq 0, -1, -2, \dots$).

Solution:

$$y(t) = \frac{e^{\pi i/4}}{4\pi\sqrt{t}} \int_{-\infty}^\infty \frac{e^{\pi x/2}}{\cosh(\pi x)} \left[1 + \left(1 + \frac{i}{2t} \right)^{ix-1/2} \right] t^{ix} f(x) dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 454).

$$99. \int_0^\infty \left\{ t^{-ix-1/2} \sin \frac{(1+2ix)\pi}{4} + 2^{-ix-3/2} e^{\pi x} \left[\zeta \left(\frac{1}{2} + ix, \frac{1-it}{2} \right) - \zeta \left(\frac{1}{2} + ix, \frac{1+it}{2} \right) \right. \right. \\ \left. \left. - \zeta \left(\frac{1}{2} + ix, -\frac{it}{2} \right) + \zeta \left(\frac{1}{2} + ix, \frac{it}{2} \right) \right] \right\} y(t) dt = f(x).$$

Here $\zeta(z, v)$ is the generalized Riemann zeta function (see Eq. 3.7.98).

Solution:

$$y(t) = \frac{1}{\pi} \int_{-\infty}^\infty \left\{ t^{ix-1/2} \sin \frac{(1-2ix)\pi}{4} + \sin \left[\left(\frac{1}{2} - ix \right) \arctan t \right] (t^2+1)^{ix/2-1/4} \right\} \frac{f(x)}{\cosh(\pi x)} dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 454).

$$100. \int_0^{\infty} \left\{ t^{-ix-1/2} \cos \frac{(1+2ix)\pi}{4} - i2^{-ix-3/2} e^{\pi x} \left[\zeta \left(\frac{1}{2} + ix, \frac{1-it}{2} \right) + \zeta \left(\frac{1}{2} + ix, \frac{1+it}{2} \right) - \zeta \left(\frac{1}{2} + ix, -\frac{it}{2} \right) - \zeta \left(\frac{1}{2} + ix, \frac{it}{2} \right) \right] \right\} y(t) dt = f(x).$$

Here $\zeta(z, v)$ is the generalized Riemann zeta function (see Eq. 3.7.98).

Solution:

$$y(t) = \frac{1}{\pi} \int_{-\infty}^{\infty} \left\{ t^{ix-1/2} \cos \frac{(1-2ix)\pi}{4} + \cos \left[\left(\frac{1}{2} - ix \right) \arctan t \right] (t^2+1)^{ix/2-1/4} \right\} \frac{f(x)}{\cosh(\pi x)} dx.$$

⊙ Reference: A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 455).

3.8. Equations Whose Kernels Contain Arbitrary Functions

3.8-1. Equations with Degenerate Kernel.

$$1. \int_a^b [g_1(x)h_1(t) + g_2(x)h_2(t)] y(t) dt = f(x).$$

This integral equation has solutions only if its right-hand side is representable in the form

$$f(x) = A_1 g_1(x) + A_2 g_2(x), \quad A_1 = \text{const}, \quad A_2 = \text{const}. \quad (1)$$

In this case, any function $y = y(x)$ satisfying the normalization type conditions

$$\int_a^b h_1(t)y(t) dt = A_1, \quad \int_a^b h_2(t)y(t) dt = A_2 \quad (2)$$

is a solution of the integral equation. Otherwise, the equation has no solutions.

$$2. \int_a^b \left[\sum_{k=0}^n g_k(x)h_k(t) \right] y(t) dt = f(x).$$

This integral equation has solutions only if its right-hand side is representable in the form

$$f(x) = \sum_{k=0}^n A_k g_k(x), \quad (1)$$

where the A_k are some constants. In this case, any function $y = y(x)$ satisfying the normalization type conditions

$$\int_a^b h_k(t)y(t) dt = A_k \quad (k = 1, \dots, n) \quad (2)$$

is a solution of the integral equation. Otherwise, the equation has no solutions.

3.8-2. Equations Containing Modulus.

$$3. \quad \int_a^b |g(x) - g(t)| y(t) dt = f(x).$$

Let $a \leq x \leq b$ and $a \leq t \leq b$; it is assumed in items 1° and 2° that $0 < g'_x(x) < \infty$.

1°. Let us remove the modulus in the integrand:

$$\int_a^x [g(x) - g(t)] y(t) dt + \int_x^b [g(t) - g(x)] y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x yields

$$g'_x(x) \int_a^x y(t) dt - g'_x(x) \int_x^b y(t) dt = f'_x(x). \quad (2)$$

Divide both sides of (2) by $g'_x(x)$ and differentiate the resulting equation to obtain the solution

$$y(x) = \frac{1}{2} \frac{d}{dx} \left[\frac{f'_x(x)}{g'_x(x)} \right]. \quad (3)$$

2°. Let us demonstrate that the right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = a$ and $x = b$, in (1), we obtain two corollaries

$$\int_a^b [g(t) - g(a)] y(t) dt = f(a), \quad \int_a^b [g(b) - g(t)] y(t) dt = f(b). \quad (4)$$

Substitute $y(x)$ of (3) into (4). Integrating by parts yields the desired constraints for $f(x)$:

$$\begin{aligned} [g(b) - g(a)] \frac{f'_x(b)}{g'_x(b)} &= f(a) + f(b), \\ [g(a) - g(b)] \frac{f'_x(a)}{g'_x(a)} &= f(a) + f(b). \end{aligned} \quad (5)$$

Let us point out a useful property of these constraints: $f'_x(b)g'_x(a) + f'_x(a)g'_x(b) = 0$.

Conditions (5) make it possible to find the admissible general form of the right-hand side of the integral equation:

$$f(x) = F(x) + Ax + B, \quad (6)$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative), and the coefficients A and B are given by

$$\begin{aligned} A &= -\frac{g'_x(a)F'_x(b) + g'_x(b)F'_x(a)}{g'_x(a) + g'_x(b)}, \\ B &= -\frac{1}{2}A(a+b) - \frac{1}{2}[F(a) + F(b)] - \frac{g(b) - g(a)}{2g'_x(a)}[A + F'_x(a)]. \end{aligned}$$

3°. If $g(x)$ is representable in the form $g(x) = O(x-a)^k$ with $0 < k < 1$ in the vicinity of the point $x = a$ (in particular, the derivative g'_x is unbounded as $x \rightarrow a$), then the solution of the integral equation is given by formula (3) as well. In this case, the right-hand side of the integral equation must satisfy the conditions

$$f(a) + f(b) = 0, \quad f'_x(b) = 0. \quad (7)$$

As before, the right-hand side of the integral equation is given by (6), with

$$A = -F'_x(b), \quad B = \frac{1}{2} [(a+b)F'_x(b) - F(a) - F(b)].$$

4°. For $g'_x(a) = 0$, the right-hand side of the integral equation must satisfy the conditions

$$f'_x(a) = 0, \quad [g(b) - g(a)]f'_x(b) = [f(a) + f(b)]g'_x(b).$$

As before, the right-hand side of the integral equation is given by (6), with

$$A = -F'_x(a), \quad B = \frac{1}{2} [(a+b)F'_x(a) - F(a) - F(b)] + \frac{g(b) - g(a)}{2g'_x(b)} [F'_x(b) - F'_x(a)].$$

4.
$$\int_0^a |g(x) - g(\lambda t)| y(t) dt = f(x), \quad \lambda > 0.$$

Assume that $0 \leq x \leq a$, $0 \leq t \leq a$, and $0 < g'_x(x) < \infty$.

1°. Let us remove the modulus in the integrand:

$$\int_0^{x/\lambda} [g(x) - g(\lambda t)] y(t) dt + \int_{x/\lambda}^a [g(\lambda t) - g(x)] y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x yields

$$g'_x(x) \int_0^{x/\lambda} y(t) dt - g'_x(x) \int_{x/\lambda}^a y(t) dt = f'_x(x). \quad (2)$$

Let us divide both sides of (2) by $g'_x(x)$ and differentiate the resulting equation to obtain $y(x/\lambda) = \frac{1}{2} \lambda [f'_x(x)/g'_x(x)]'_x$. Substituting x by λx yields the solution

$$y(x) = \frac{\lambda}{2} \frac{d}{dz} \left[\frac{f'_z(z)}{g'_z(z)} \right], \quad z = \lambda x. \quad (3)$$

2°. Let us demonstrate that the right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = 0$ in (1) and (2), we obtain two corollaries

$$\int_0^a [g(\lambda t) - g(0)] y(t) dt = f(0), \quad g'_x(0) \int_0^a y(t) dt = -f'_x(0). \quad (4)$$

Substitute $y(x)$ of (3) into (4). Integrating by parts yields the desired constraints for $f(x)$:

$$\begin{aligned} f'_x(0)g'_x(\lambda a) + f'_x(\lambda a)g'_x(0) &= 0, \\ [g(\lambda a) - g(0)] \frac{f'_x(\lambda a)}{g'_x(\lambda a)} &= f(0) + f(\lambda a). \end{aligned} \quad (5)$$

Conditions (5) make it possible to find the admissible general form of the right-hand side of the integral equation:

$$f(x) = F(x) + Ax + B, \quad (6)$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative), and the coefficients A and B are given by

$$\begin{aligned} A &= -\frac{g'_x(0)F'_x(\lambda a) + g'_x(\lambda a)F'_x(0)}{g'_x(0) + g'_x(\lambda a)}, \\ B &= -\frac{1}{2} Aa\lambda - \frac{1}{2} [F(0) + F(\lambda a)] - \frac{g(\lambda a) - g(0)}{2g'_x(0)} [A + F'_x(0)]. \end{aligned}$$

3°. If $g(x)$ is representable in the form $g(x) = O(x)^k$ with $0 < k < 1$ in the vicinity of the point $x = 0$ (in particular, the derivative g'_x is unbounded as $x \rightarrow 0$), then the solution of the integral equation is given by formula (3) as well. In this case, the right-hand side of the integral equation must satisfy the conditions

$$f(0) + f(\lambda a) = 0, \quad f'_x(\lambda a) = 0. \quad (7)$$

As before, the right-hand side of the integral equation is given by (6), with

$$A = -F'_x(\lambda a), \quad B = \frac{1}{2} [a\lambda F'_x(\lambda a) - F(0) - F(\lambda a)].$$

5.
$$\int_0^a |g(x) - t| y(t) dt = f(x).$$

Assume that $0 \leq x \leq a$, $0 \leq t \leq a$; $g(0) = 0$, and $0 < g'_x(x) < \infty$.

1°. Let us remove the modulus in the integrand:

$$\int_0^{g(x)} [g(x) - t] y(t) dt + \int_{g(x)}^a [t - g(x)] y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x yields

$$g'_x(x) \int_0^{g(x)} y(t) dt - g'_x(x) \int_{g(x)}^a y(t) dt = f'_x(x). \quad (2)$$

Let us divide both sides of (2) by $g'_x(x)$ and differentiate the resulting equation to obtain $2g'_x(x)y(g(x)) = [f'_x(x)/g'_x(x)]'_x$. Hence, we find the solution:

$$y(x) = \frac{1}{2g'_z(z)} \frac{d}{dz} \left[\frac{f'_z(z)}{g'_z(z)} \right], \quad z = g^{-1}(x), \quad (3)$$

where g^{-1} is the inverse of g .

2°. Let us demonstrate that the right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = 0$ in (1) and (2), we obtain two corollaries

$$\int_0^a ty(t) dt = f(0), \quad g'_x(0) \int_0^a y(t) dt = -f'_x(0). \quad (4)$$

Substitute $y(x)$ of (3) into (4). Integrating by parts yields the desired constraints for $f(x)$:

$$\begin{aligned} f'_x(0)g'_x(x_a) + f'_x(x_a)g'_x(0) &= 0, \quad x_a = g^{-1}(a); \\ g(x_a) \frac{f'_x(x_a)}{g'_x(x_a)} &= f(0) + f(x_a). \end{aligned} \quad (5)$$

Conditions (5) make it possible to find the admissible general form of the right-hand side of the integral equation in question:

$$f(x) = F(x) + Ax + B, \quad (6)$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative), and the coefficients A and B are given by

$$A = -\frac{g'_x(0)F'_x(x_a) + g'_x(x_a)F'_x(0)}{g'_x(0) + g'_x(x_a)}, \quad x_a = g^{-1}(a),$$

$$B = -\frac{1}{2}Ax_a - \frac{1}{2}[F(0) + F(x_a)] - \frac{g(x_a)}{2g'_x(0)}[A + F'_x(0)].$$

3°. If $g(x)$ is representable in the vicinity of the point $x = 0$ in the form $g(x) = O(x)^k$ with $0 < k < 1$ (i.e., the derivative g'_x is unbounded as $x \rightarrow 0$), then the solution of the integral equation is given by formula (3) as well. In this case, the right-hand side of the integral equation must satisfy the conditions

$$f(0) + f(x_a) = 0, \quad f'_x(x_a) = 0. \quad (7)$$

As before, the right-hand side of the integral equation is given by (6), with

$$A = -F'_x(x_a), \quad B = \frac{1}{2}[x_a F'_x(x_a) - F(0) - F(x_a)].$$

6.
$$\int_0^a |x - g(t)| y(t) dt = f(x).$$

Assume that $0 \leq x \leq a$, $0 \leq t \leq a$; $g(0) = 0$, and $0 < g'_x(x) < \infty$.

1°. Let us remove the modulus in the integrand:

$$\int_0^{g^{-1}(x)} [x - g(t)]y(t) dt + \int_{g^{-1}(x)}^a [g(t) - x]y(t) dt = f(x), \quad (1)$$

where g^{-1} is the inverse of g . Differentiating (1) with respect to x yields

$$\int_0^{g^{-1}(x)} y(t) dt - \int_{g^{-1}(x)}^a y(t) dt = f'_x(x). \quad (2)$$

Differentiating the resulting equation yields $2y(g^{-1}(x)) = g'_x(x)f''_{xx}(x)$. Hence, we obtain the solution

$$y(x) = \frac{1}{2}g'_z(z)f''_{zz}(z), \quad z = g(x). \quad (3)$$

2°. Let us demonstrate that the right-hand side $f(x)$ of the integral equation must satisfy certain relations. By setting $x = 0$ in (1) and (2), we obtain two corollaries

$$\int_0^a g(t)y(t) dt = f(0), \quad \int_0^a y(t) dt = -f'_x(0). \quad (4)$$

Substitute $y(x)$ of (3) into (4). Integrating by parts yields the desired constraints for $f(x)$:

$$x_a f'_x(x_a) = f(0) + f(x_a), \quad f'_x(0) + f'_x(x_a) = 0, \quad x_a = g(a). \quad (5)$$

Conditions (5) make it possible to find the admissible general form of the right-hand side of the integral equation:

$$f(x) = F(x) + Ax + B,$$

$$A = -\frac{1}{2}[F'_x(0) + F'_x(x_a)], \quad B = \frac{1}{2}[x_a F'_x(0) - F(x_a) - F(0)], \quad x_a = g(a),$$

where $F(x)$ is an arbitrary bounded twice differentiable function (with bounded first derivative).

$$7. \int_a^b \frac{y(t)}{|g(x) - g(t)|^k} dt = f(x), \quad 0 < k < 1.$$

Let $g'_x \neq 0$. The transformation

$$z = g(x), \quad \tau = g(t), \quad w(\tau) = \frac{1}{g'_t(t)} y(t)$$

leads to an equation of the form 3.1.31:

$$\int_A^B \frac{w(\tau)}{|z - \tau|^k} d\tau = F(z), \quad A = g(a), \quad B = g(b),$$

where $F = F(z)$ is the function which is obtained from $z = g(x)$ and $F = f(x)$ by eliminating x .

$$8. \int_0^1 \frac{y(t)}{|g(x) - h(t)|^k} dt = f(x), \quad 0 < k < 1.$$

Let $g(0) = 0$, $g(1) = 1$, $g'_x > 0$; $h(0) = 0$, $h(1) = 1$, and $h'_t > 0$.

The transformation

$$z = g(x), \quad \tau = h(t), \quad w(\tau) = \frac{1}{h'_t(t)} y(t)$$

leads to an equation of the form 3.1.30:

$$\int_0^1 \frac{w(\tau)}{|z - \tau|^k} d\tau = F(z),$$

where $F = F(z)$ is the function which is obtained from $z = g(x)$ and $F = f(x)$ by eliminating x .

$$9. \int_a^b y(t) \ln |g(x) - g(t)| dt = f(x).$$

Let $g'_x \neq 0$. The transformation

$$z = g(x), \quad \tau = g(t), \quad w(\tau) = \frac{1}{g'_t(t)} y(t)$$

leads to Carleman's equation 3.4.2:

$$\int_A^B \ln |z - \tau| w(\tau) d\tau = F(z), \quad A = g(a), \quad B = g(b),$$

where $F = F(z)$ is the function which is obtained from $z = g(x)$ and $F = f(x)$ by eliminating x .

$$10. \int_0^1 y(t) \ln |g(x) - h(t)| dt = f(x).$$

Let $g(0) = 0$, $g(1) = 1$, $g'_x > 0$; $h(0) = 0$, $h(1) = 1$, and $h'_t > 0$.

The transformation

$$z = g(x), \quad \tau = h(t), \quad w(\tau) = \frac{1}{h'_t(t)} y(t)$$

leads to an equation of the form 3.4.2:

$$\int_0^1 \ln |z - \tau| w(\tau) d\tau = F(z),$$

where $F = F(z)$ is the function which is obtained from $z = g(x)$ and $F = f(x)$ by eliminating x .

3.8-3. Equations with Difference Kernel: $K(x, t) = K(x - t)$.

$$11. \int_{-\infty}^{\infty} K(x-t)y(t) dt = Ax^n, \quad n = 0, 1, 2, \dots$$

1°. Solution with $n = 0$:

$$y(x) = \frac{A}{B}, \quad B = \int_{-\infty}^{\infty} K(x) dx.$$

2°. Solution with $n = 1$:

$$y(x) = \frac{A}{B}x + \frac{AC}{B^2}, \quad B = \int_{-\infty}^{\infty} K(x) dx, \quad C = \int_{-\infty}^{\infty} xK(x) dx.$$

3°. Solution with $n \geq 2$:

$$y(x) = \left\{ \frac{d^n}{d\lambda^n} \left[\frac{Ae^{\lambda x}}{B(\lambda)} \right] \right\}_{\lambda=0}, \quad B(\lambda) = \int_{-\infty}^{\infty} K(x)e^{-\lambda x} dx.$$

$$12. \int_{-\infty}^{\infty} K(x-t)y(t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = \frac{A}{B}e^{\lambda x}, \quad B = \int_{-\infty}^{\infty} K(x)e^{-\lambda x} dx.$$

$$13. \int_{-\infty}^{\infty} K(x-t)y(t) dt = Ax^n e^{\lambda x}, \quad n = 1, 2, \dots$$

1°. Solution with $n = 1$:

$$y(x) = \frac{A}{B}xe^{\lambda x} + \frac{AC}{B^2}e^{\lambda x},$$

$$B = \int_{-\infty}^{\infty} K(x)e^{-\lambda x} dx, \quad C = \int_{-\infty}^{\infty} xK(x)e^{-\lambda x} dx.$$

2°. Solution with $n \geq 2$:

$$y(x) = \frac{d^n}{d\lambda^n} \left[\frac{Ae^{\lambda x}}{B(\lambda)} \right], \quad B(\lambda) = \int_{-\infty}^{\infty} K(x)e^{-\lambda x} dx.$$

$$14. \int_{-\infty}^{\infty} K(x-t)y(t) dt = A \cos(\lambda x) + B \sin(\lambda x).$$

Solution:

$$y(x) = \frac{AI_c + BI_s}{I_c^2 + I_s^2} \cos(\lambda x) + \frac{BI_c - AI_s}{I_c^2 + I_s^2} \sin(\lambda x),$$

$$I_c = \int_{-\infty}^{\infty} K(z) \cos(\lambda z) dz, \quad I_s = \int_{-\infty}^{\infty} K(z) \sin(\lambda z) dz.$$

$$15. \int_{-\infty}^{\infty} K(x-t)y(t) dt = f(x).$$

The Fourier transform is used to solve this equation.

1°. Solution:

$$y(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\tilde{f}(u)}{\tilde{K}(u)} e^{iux} du,$$

$$\tilde{f}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-iux} dx, \quad \tilde{K}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K(x) e^{-iux} dx.$$

The following statement is valid. Let $f(x) \in L_2(-\infty, \infty)$ and $K(x) \in L_1(-\infty, \infty)$. Then for a solution $y(x) \in L_2(-\infty, \infty)$ of the integral equation to exist, it is necessary and sufficient that $\tilde{f}(u)/\tilde{K}(u) \in L_2(-\infty, \infty)$.

2°. Let the function $P(s)$ defined by the formula

$$\frac{1}{P(s)} = \int_{-\infty}^{\infty} e^{-st} K(t) dt$$

be a polynomial of degree n with real roots of the form

$$P(s) = \left(1 - \frac{s}{a_1}\right) \left(1 - \frac{s}{a_2}\right) \dots \left(1 - \frac{s}{a_n}\right).$$

Then the solution of the integral equation is given by

$$y(x) = P(D)f(x), \quad D = \frac{d}{dx}.$$

⊙ References: I. I. Hirschman and D. V. Widder (1955), V. A. Ditkin and A. P. Prudnikov (1965).

$$16. \int_0^{\infty} K(x-t)y(t) dt = f(x).$$

The Wiener-Hopf equation of the first kind. This equation is discussed in Subsection 12.8-1 in detail.

3.8-4. Other Equations of the Form $\int_a^b K(x,t)y(t) dt = F(x)$.

$$17. \int_{-\infty}^{\infty} K(ax-t)y(t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = \frac{A}{B} \exp\left(\frac{\lambda}{a}x\right), \quad B = \int_{-\infty}^{\infty} K(z) \exp\left(-\frac{\lambda}{a}z\right) dz.$$

$$18. \int_{-\infty}^{\infty} K(ax-t)y(t) dt = f(x).$$

The substitution $z = ax$ leads to an equation of the form 3.8.15:

$$\int_{-\infty}^{\infty} K(z-t)y(t) dt = f(z/a).$$

$$19. \int_{-\infty}^{\infty} K(ax+t)y(t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = \frac{A}{B} \exp\left(-\frac{\lambda}{a}x\right), \quad B = \int_{-\infty}^{\infty} K(z) \exp\left(-\frac{\lambda}{a}z\right) dz.$$

$$20. \int_{-\infty}^{\infty} K(ax+t)y(t) dt = f(x).$$

The transformation $\tau = -t$, $z = ax$, $y(t) = Y(\tau)$ leads to an equation of the form 3.8.15:

$$\int_{-\infty}^{\infty} K(z-\tau)Y(\tau) dt = f(z/a).$$

$$21. \int_{-\infty}^{\infty} [e^{\beta t} K(ax+t) + e^{\mu t} M(ax-t)]y(t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = A \frac{I_k(q)e^{px} - I_m(p)e^{qx}}{I_k(p)I_k(q) - I_m(p)I_m(q)}, \quad p = -\frac{\lambda}{a} - \beta, \quad q = \frac{\lambda}{a} - \mu,$$

where

$$I_k(q) = \int_{-\infty}^{\infty} K(z)e^{(\beta+q)z} dz, \quad I_m(q) = \int_{-\infty}^{\infty} M(z)e^{-(\mu+q)z} dz.$$

$$22. \int_0^{\infty} g(xt)y(t) dt = f(x).$$

By setting

$$x = e^z, \quad t = e^{-\tau}, \quad y(t) = e^{\tau}w(\tau), \quad g(\xi) = G(\ln \xi), \quad f(\xi) = F(\ln \xi),$$

we arrive at an integral equation with difference kernel of the form 3.8.15:

$$\int_{-\infty}^{\infty} G(z-\tau)w(\tau) d\tau = F(z).$$

$$23. \int_0^{\infty} g\left(\frac{x}{t}\right)y(t) dt = f(x).$$

By setting

$$x = e^z, \quad t = e^{\tau}, \quad y(t) = e^{-\tau}w(\tau), \quad g(\xi) = G(\ln \xi), \quad f(\xi) = F(\ln \xi),$$

we arrive at an integral equation with difference kernel of the form 3.8.15:

$$\int_{-\infty}^{\infty} G(z-\tau)w(\tau) d\tau = F(z).$$

$$24. \int_0^{\infty} g(x^{\beta}t^{\lambda})y(t) dt = f(x), \quad \beta > 0, \quad \lambda > 0.$$

By setting

$$x = e^{z/\beta}, \quad t = e^{-\tau/\lambda}, \quad y(t) = e^{\tau/\lambda}w(\tau), \quad g(\xi) = G(\ln \xi), \quad f(\xi) = \frac{1}{\lambda}F(\beta \ln \xi),$$

we arrive at an integral equation with difference kernel of the form 3.8.15:

$$\int_{-\infty}^{\infty} G(z-\tau)w(\tau) d\tau = F(z).$$

$$25. \int_0^{\infty} g\left(\frac{x^\beta}{t^\lambda}\right) y(t) dt = f(x), \quad \beta > 0, \quad \lambda > 0.$$

By setting

$$x = e^{z/\beta}, \quad t = e^{\tau/\lambda}, \quad y(t) = e^{-\tau/\lambda} w(\tau), \quad g(\xi) = G(\ln \xi), \quad f(\xi) = \frac{1}{\lambda} F(\beta \ln \xi),$$

we arrive at an integral equation with difference kernel of the form 3.8.15:

$$\int_{-\infty}^{\infty} G(z - \tau) w(\tau) d\tau = F(z).$$

$$26. \int_0^a \left[\frac{1}{|x-t|^k} + \varphi(x)\psi(t) \right] y(t) dt = f(x), \quad 0 < k < 1.$$

The solution can be obtained by the methods described in Subsection 12.6-2; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.1.30.

$$27. \int_0^{\infty} \exp[-g(x)t^2] y(t) dt = f(x).$$

Assume that $g(0) = \infty$, $g(\infty) = 0$, and $g'_x < 0$.

The substitution $z = \frac{1}{4g(x)}$ leads to equation 3.2.21:

$$\frac{1}{\sqrt{\pi z}} \int_0^{\infty} \exp\left(-\frac{t^2}{4z}\right) y(t) dt = F(z),$$

where the function $F(z)$ is determined by the relations $F = \frac{2}{\sqrt{\pi}} f(x) \sqrt{g(x)}$ and $z = \frac{1}{4g(x)}$ by means of eliminating x .

$$28. \int_a^b \left[\ln|x-t| + \varphi(x)\psi(t) \right] y(t) dt = f(x).$$

The solution can be obtained by the methods described in Subsection 12.6-2; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.4.2. See also Example 3 in Subsection 12.6-2.

$$29. \int_0^{\infty} [\sin(xt) + \varphi(x)\psi(t)] y(t) dt = f(x).$$

The solution can be obtained by the methods described in Subsection 12.6-2; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.5.8.

Solution:

$$y(t) = y_f(t) + Ay_\varphi(t),$$

where

$$y_f(t) = \frac{2}{\pi} \int_0^{\infty} \sin(xt) f(x) dx, \quad y_\varphi(t) = \frac{2}{\pi} \int_0^{\infty} \sin(xt) \varphi(x) dx, \quad A = -\frac{\int_0^{\infty} \psi(t) y_f(t) dt}{1 + \int_0^{\infty} \psi(t) y_\varphi(t) dt}.$$

$$30. \int_0^{\infty} [\cos(xt) + \varphi(x)\psi(t)]y(t) dt = f(x).$$

The solution can be obtained by the methods described in Subsection 12.6-2; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.5.1.

Solution:

$$y(t) = y_f(t) + Ay_{\varphi}(t),$$

where

$$y_f(t) = \frac{2}{\pi} \int_0^{\infty} \cos(xt)f(x) dx, \quad y_{\varphi}(t) = \frac{2}{\pi} \int_0^{\infty} \cos(xt)\varphi(x) dx, \quad A = -\frac{\int_0^{\infty} \psi(t)y_f(t) dt}{1 + \int_0^{\infty} \psi(t)y_{\varphi}(t) dt}.$$

$$31. \int_0^{\infty} t^{a-1} \cos [\varphi(x)t^a] y(t) dt = f(x), \quad a > 0.$$

Transformation

$$z = \varphi(x), \quad \tau = t^a, \quad Y(\tau) = y(t), \quad F(z) = af(x)$$

leads to an equation of the form 3.5.1:

$$\int_0^{\infty} \cos(z\tau)Y(\tau) d\tau = F(z).$$

$$32. \int_0^{\infty} t^{a-1} \sin [\varphi(x)t^a] y(t) dt = f(x), \quad a > 0.$$

Transformation

$$z = \varphi(x), \quad \tau = t^a, \quad Y(\tau) = y(t), \quad F(z) = af(x)$$

leads to an equation of the form 3.5.8:

$$\int_0^{\infty} \sin(z\tau)Y(\tau) d\tau = F(z).$$

$$33. \int_0^{\infty} [tJ_{\nu}(xt) + \varphi(x)\psi(t)]y(t) dt = f(x), \quad \nu > -1.$$

Here $J_{\nu}(z)$ is the Bessel function of the first kind. The solution can be obtained by the methods described in Subsection 12.6-2; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.7.17.

Solution:

$$y(t) = y_f(t) + Ay_{\varphi}(t),$$

where

$$y_f(t) = \int_0^{\infty} xJ_{\nu}(xt)f(x) dx, \quad y_{\varphi}(t) = \int_0^{\infty} xJ_{\nu}(xt)\varphi(x) dx, \quad A = -\frac{\int_0^{\infty} \psi(t)y_f(t) dt}{1 + \int_0^{\infty} \psi(t)y_{\varphi}(t) dt}.$$

3.8-5. Equations of the Form $\int_a^b K(x, t)y(\cdot \cdot \cdot) dt = F(x)$.

34. $\int_a^b f(t)y(xt) dt = Ax + B.$

Solution:

$$y(x) = \frac{A}{I_1}x + \frac{B}{I_0}, \quad I_0 = \int_a^b f(t) dt, \quad I_1 = \int_a^b tf(t) dt.$$

35. $\int_a^b f(t)y(xt) dt = Ax^\beta.$

Solution:

$$y(x) = \frac{A}{B}x^\beta, \quad B = \int_a^b f(t)t^\beta dt.$$

36. $\int_a^b f(t)y(xt) dt = A \ln x + B.$

Solution:

$$y(x) = p \ln x + q,$$

where

$$p = \frac{A}{I_0}, \quad q = \frac{B}{I_0} - \frac{AI_1}{I_0^2}, \quad I_0 = \int_a^b f(t) dt, \quad I_1 = \int_a^b f(t) \ln t dt.$$

37. $\int_a^b f(t)y(xt) dt = Ax^\beta \ln x.$

Solution:

$$y(x) = px^\beta \ln x + qx^\beta,$$

where

$$p = \frac{A}{I_1}, \quad q = -\frac{AI_2}{I_1^2}, \quad I_1 = \int_a^b f(t)t^\beta dt, \quad I_2 = \int_a^b f(t)t^\beta \ln t dt.$$

38. $\int_a^b f(t)y(xt) dt = A \cos(\ln x).$

Solution:

$$y(x) = \frac{AI_c}{I_c^2 + I_s^2} \cos(\ln x) + \frac{AI_s}{I_c^2 + I_s^2} \sin(\ln x),$$

$$I_c = \int_a^b f(t) \cos(\ln t) dt, \quad I_s = \int_a^b f(t) \sin(\ln t) dt.$$

39. $\int_a^b f(t)y(xt) dt = A \sin(\ln x).$

Solution:

$$y(x) = -\frac{AI_s}{I_c^2 + I_s^2} \cos(\ln x) + \frac{AI_c}{I_c^2 + I_s^2} \sin(\ln x),$$

$$I_c = \int_a^b f(t) \cos(\ln t) dt, \quad I_s = \int_a^b f(t) \sin(\ln t) dt.$$

$$40. \int_a^b f(t)y(xt) dt = Ax^\beta \cos(\ln x) + Bx^\beta \sin(\ln x).$$

Solution:

$$y(x) = px^\beta \cos(\ln x) + qx^\beta \sin(\ln x),$$

where

$$p = \frac{AI_c - BI_s}{I_c^2 + I_s^2}, \quad q = \frac{AI_s + BI_c}{I_c^2 + I_s^2},$$

$$I_c = \int_a^b f(t)t^\beta \cos(\ln t) dt, \quad I_s = \int_a^b f(t)t^\beta \sin(\ln t) dt.$$

$$41. \int_a^b f(t)y(x-t) dt = Ax + B.$$

Solution:

$$y(x) = px + q,$$

where

$$p = \frac{A}{I_0}, \quad q = \frac{AI_1}{I_0^2} + \frac{B}{I_0}, \quad I_0 = \int_a^b f(t) dt, \quad I_1 = \int_a^b tf(t) dt.$$

$$42. \int_a^b f(t)y(x-t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = \frac{A}{B}e^{\lambda x}, \quad B = \int_a^b f(t) \exp(-\lambda t) dt.$$

$$43. \int_a^b f(t)y(x-t) dt = A \cos(\lambda x).$$

Solution:

$$y(x) = -\frac{AI_s}{I_c^2 + I_s^2} \sin(\lambda x) + \frac{AI_c}{I_c^2 + I_s^2} \cos(\lambda x),$$

$$I_c = \int_a^b f(t) \cos(\lambda t) dt, \quad I_s = \int_a^b f(t) \sin(\lambda t) dt.$$

$$44. \int_a^b f(t)y(x-t) dt = A \sin(\lambda x).$$

Solution:

$$y(x) = \frac{AI_c}{I_c^2 + I_s^2} \sin(\lambda x) + \frac{AI_s}{I_c^2 + I_s^2} \cos(\lambda x),$$

$$I_c = \int_a^b f(t) \cos(\lambda t) dt, \quad I_s = \int_a^b f(t) \sin(\lambda t) dt.$$

$$45. \int_a^b f(t)y(x-t) dt = e^{\mu x}(A \sin \lambda x + B \cos \lambda x).$$

Solution:

$$y(x) = e^{\mu x}(p \sin \lambda x + q \cos \lambda x),$$

where

$$p = \frac{AI_c - BI_s}{I_c^2 + I_s^2}, \quad q = \frac{AI_s + BI_c}{I_c^2 + I_s^2},$$

$$I_c = \int_a^b f(t)e^{-\mu t} \cos(\lambda t) dt, \quad I_s = \int_a^b f(t)e^{-\mu t} \sin(\lambda t) dt.$$

$$46. \int_a^b f(t)y(x-t) dt = g(x).$$

1°. For $g(x) = \sum_{k=1}^n A_k \exp(\lambda_k x)$, the solution of the equation has the form

$$y(x) = \sum_{k=1}^n \frac{A_k}{B_k} \exp(\lambda_k x), \quad B_k = \int_a^b f(t) \exp(-\lambda_k t) dt.$$

2°. For a polynomial right-hand side, $g(x) = \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients.

3°. For $g(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

6°. For $g(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

7°. For $g(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

8°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \cos(\lambda_k x)$, the solution has the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

9°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \sin(\lambda_k x)$, the solution has the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

10°. For $g(x) = \cos(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

11°. For $g(x) = \sin(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

47. $\int_a^b f(t)y(x + \beta t) dt = Ax + B.$

Solution:

$$y(x) = px + q,$$

where

$$p = \frac{A}{I_0}, \quad q = \frac{B}{I_0} - \frac{AI_1\beta}{I_0^2}, \quad I_0 = \int_a^b f(t) dt, \quad I_1 = \int_a^b t f(t) dt.$$

$$48. \int_a^b f(t)y(x + \beta t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = \frac{A}{B}e^{\lambda x}, \quad B = \int_a^b f(t) \exp(\lambda \beta t) dt.$$

$$49. \int_a^b f(t)y(x + \beta t) dt = A \sin \lambda x + B \cos \lambda x.$$

Solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x,$$

where

$$p = \frac{AI_c + BI_s}{I_c^2 + I_s^2}, \quad q = \frac{BI_c - AI_s}{I_c^2 + I_s^2},$$

$$I_c = \int_a^b f(t) \cos(\lambda \beta t) dt, \quad I_s = \int_a^b f(t) \sin(\lambda \beta t) dt.$$

$$50. \int_0^1 y(\xi) dt = f(x), \quad \xi = g(x)t.$$

Assume that $g(0) = 0$, $g(1) = 1$, and $g'_x \geq 0$.

1°. The substitution $z = g(x)$ leads to an equation of the form 3.1.42: $\int_0^1 y(zt) dt = F(z)$, where the function $F(z)$ is obtained from $z = g(x)$ and $F = f(x)$ by eliminating x .

2°. Solution $y = y(z)$ in the parametric form:

$$y(z) = \frac{g(x)}{g'_x(x)} f'_x(x) + f(x), \quad z = g(x).$$

$$51. \int_0^1 t^\lambda y(\xi) dt = f(x), \quad \xi = g(x)t.$$

Assume that $g(0) = 0$, $g(1) = 1$, and $g'_x \geq 0$.

1°. The substitution $z = g(x)$ leads to an equation of the form 3.1.43: $\int_0^1 t^\lambda y(zt) dt = F(z)$, where the function $F(z)$ is obtained from $z = g(x)$ and $F = f(x)$ by eliminating x .

2°. Solution $y = y(z)$ in the parametric form:

$$y(z) = \frac{g(x)}{g'_x(x)} f'_x(x) + (\lambda + 1)f(x), \quad z = g(x).$$

$$52. \int_a^b f(t)y(\xi) dt = Ax^\beta, \quad \xi = x\varphi(t).$$

Solution:

$$y(x) = \frac{A}{B}x^\beta, \quad B = \int_a^b f(t) [\varphi(t)]^\beta dt. \quad (1)$$

$$53. \int_a^b f(t)y(\xi) dt = g(x), \quad \xi = x\varphi(t).$$

1°. For $g(x) = \sum_{k=0}^n A_k x^k$, the solution of the equation has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_k} x^k, \quad B_k = \int_a^b f(t) [\varphi(t)]^k dt.$$

2°. For $g(x) = \sum_{k=0}^n A_k x^{\lambda_k}$, the solution has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_k} x^{\lambda_k}, \quad B_k = \int_a^b f(t) [\varphi(t)]^{\lambda_k} dt.$$

3°. For $g(x) = \ln x \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \ln x \sum_{k=0}^n B_k x^k + \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=0}^n A_k (\ln x)^k$, the solution has the form

$$y(x) = \sum_{k=0}^n B_k (\ln x)^k,$$

where the constants B_k are found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k \ln x)$, the solution has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k \ln x) + \sum_{k=1}^n C_k \sin(\lambda_k \ln x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

6°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k \ln x)$, the solution has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k \ln x) + \sum_{k=1}^n C_k \sin(\lambda_k \ln x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

$$54. \int_a^b f(t)y(\xi) dt = g(x), \quad \xi = x + \varphi(t).$$

1°. For $g(x) = \sum_{k=1}^n A_k \exp(\lambda_k x)$, the solution of the equation has the form

$$y(x) = \sum_{k=1}^n \frac{A_k}{B_k} \exp(\lambda_k x), \quad B_k = \int_a^b f(t) \exp[\lambda_k \varphi(t)] dt.$$

2°. For a polynomial right-hand side, $g(x) = \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients.

3°. For $g(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k are found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k x)$ the solution has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k x)$, the solution has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

6°. For $g(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

7°. For $g(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

8°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \cos(\lambda_k x)$, the solution has the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

9°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \sin(\lambda_k x)$, the solution has the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

10°. For $g(x) = \cos(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

11°. For $g(x) = \sin(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, the solution has the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k are found by the method of undetermined coefficients.

3.9. Dual Integral Equations of the First Kind

3.9-1. Kernels Containing Trigonometric Functions.

$$1. \quad \int_0^{\infty} \cos(xt)y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} \sin(xt)y(t) dt = 0 \quad \text{for } 1 < x < \infty.$$

Solution:

$$y(x) = \frac{2x}{\pi} \int_0^1 t J_0(xt) \left[\int_0^t \frac{f(s) ds}{\sqrt{t^2 - s^2}} \right] dt.$$

⊙ References: C. Nasim and B. D. Aggarwala (1984), B. N. Mandal and N. Mandal (1999, pp. 134–136).

$$2. \quad \int_0^{\infty} \cos(xt)y(t) dt = 0 \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} \sin(xt)y(t) dt = f(x) \quad \text{for } 1 < x < \infty.$$

Solution:

$$y(x) = \frac{2x}{\pi} \int_1^{\infty} t J_0(xt) \left[\int_t^{\infty} \frac{f(s) ds}{\sqrt{s^2 - t^2}} \right] dt.$$

⊙ References: C. Nasim and B. D. Aggarwala (1984), B. N. Mandal and N. Mandal (1999, pp. 136–137).

$$3. \quad \int_0^{\infty} \cos(xt)y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} t \cos(xt)y(t) dt = 0 \quad \text{for } 1 < x < \infty.$$

Solution:

$$y(x) = \frac{2x}{\pi} \int_0^1 t J_0(xt) \left[\int_0^t \frac{f(s) ds}{\sqrt{t^2 - s^2}} \right] dt - \frac{2}{\pi} J_1(x) \int_0^1 \frac{f(s) ds}{\sqrt{1 - s^2}}.$$

⊙ References: I. W. Busbridge (1938), B. N. Mandal and N. Mandal (1999, pp. 138–139).

$$4. \quad \int_0^{\infty} t \cos(xt)y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} \cos(xt)y(t) dt = 0 \quad \text{for } 1 < x < \infty.$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^1 t J_0(xt) \left[\int_0^t \frac{f(s) ds}{\sqrt{t^2 - s^2}} \right] dt.$$

⊙ References: I. W. (1937, p. 339), B. N. Mandal and N. Mandal (1999, pp. 139–140).

$$5. \quad \int_0^{\infty} \sin(xt)y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} t \sin(xt)y(t) dt = 0 \quad \text{for } 1 < x < \infty.$$

It is assumed that $f(0) = 0$.

Solution:

$$y(x) = \frac{2}{\pi} \int_0^1 t J_0(xt) \left[\int_0^t \frac{f'_s(s) ds}{\sqrt{t^2 - s^2}} \right] dt.$$

⊙ References: I. W. Busbridge (1938), B. N. Mandal and N. Mandal (1999, pp. 140–141).

$$6. \quad \int_0^{\infty} t \sin(xt)y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} \sin(xt)y(t) dt = 0 \quad \text{for } 1 < x < \infty.$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^1 J_1(xt) \left[\int_0^t \frac{s f(s) ds}{\sqrt{t^2 - s^2}} \right] dt.$$

⊙ References: B. Noble (1963), B. N. Mandal and N. Mandal (1999, pp. 141–142).

$$7. \quad \int_0^{\infty} [a \sin(xt) + t \cos(xt)]y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} t[a \sin(xt) + t \cos(xt)]y(t) dt = 0 \quad \text{for } 1 < x < \infty.$$

Solution:

$$y(x) = \int_0^1 t J_0(xt) F(t) dt + \frac{F(1)}{K_0(a)} \int_1^{\infty} t J_0(xt) \left[\int_t^{\infty} \frac{e^{-as} ds}{\sqrt{s^2 - t^2}} \right] dt,$$

where

$$F(t) = \frac{2}{\pi} \frac{d}{dt} \int_0^t \frac{z\varphi(z) dz}{\sqrt{t^2 - z^2}} = \frac{2}{\pi} \int_0^t \frac{\varphi'_z(z) dz}{\sqrt{t^2 - z^2}}, \quad \varphi(z) = e^{-az} \int_0^z e^{as} f(s) ds \quad (0 < z < 1),$$

and $K_0(x)$ is the modified Bessel functions of the second kind.

⊙ References: B. D. Aggarwala and C. Nasim (1996), B. N. Mandal and N. Mandal (1999, pp. 143–145).

$$8. \quad \int_0^\infty t[a \sin(xt) + t \cos(xt)]y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^\infty [a \sin(xt) + t \cos(xt)]y(t) dt = g(x) \quad \text{for } 1 < x < \infty.$$

Solution for $a \neq 0$:

$$y(x) = \int_0^1 J_1(xt)F(t) dt - \int_1^\infty tJ_1(xt)G(t) dt + \frac{2Da}{\pi} \int_t^\infty J_1(xt)K_1(at) dt, \quad D = \frac{F(1) + G(1)}{aK_1(a)},$$

where

$$F(t) = \frac{2}{\pi} \int_0^t \frac{z\varphi(z) dz}{\sqrt{t^2 - z^2}}, \quad \varphi(z) = e^{-az} \int_0^z e^{as} f(s) ds \quad (0 < z < 1),$$

$$G(t) = \frac{2}{\pi} \frac{d}{dt} \int_t^\infty \frac{\psi(z) dz}{\sqrt{z^2 - t^2}}, \quad \psi(z) = e^{-az} \int_1^z e^{as} g(s) ds \quad (1 < z < \infty),$$

and $K_0(x)$ is the modified Bessel functions of the second kind.

⊙ References: B. D. Aggarwala and C. Nasim (1996), B. N. Mandal and N. Mandal (1999, pp. 143, 147–148).

3.9-2. Kernels Containing Bessel Functions of the First Kind.

$$9. \quad \int_0^\infty J_0(xt)y(t) dt = f(x) \quad \text{for } 0 < x < a,$$

$$\int_0^\infty tJ_0(xt)y(t) dt = 0 \quad \text{for } a < x < \infty.$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^a \cos(xt) \left[\frac{d}{dt} \int_0^t \frac{sf(s) ds}{\sqrt{t^2 - s^2}} \right] dt.$$

$$10. \quad \int_0^\infty tJ_0(xt)y(t) dt = f(x) \quad \text{for } 0 < x < a,$$

$$\int_0^\infty J_0(xt)y(t) dt = 0 \quad \text{for } a < x < \infty.$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^a \sin(xt) \left[\frac{d}{dt} \int_0^t \frac{sf(s) ds}{\sqrt{t^2 - s^2}} \right] dt.$$

$$11. \quad \int_0^\infty tJ_\mu(xt)y(t) dt = f(x) \quad \text{for } 0 < x < a,$$

$$\int_0^\infty J_\mu(xt)y(t) dt = 0 \quad \text{for } a < x < \infty.$$

Solution:

$$y(x) = \sqrt{\frac{2x}{\pi}} \int_0^a t^{3/2} J_{\mu+\frac{1}{2}}(xt) \left[\int_0^{\pi/2} \sin^{\mu+1} \theta f(t \sin \theta) d\theta \right] dt.$$

$$12. \int_0^{\infty} J_{\mu}(xt)y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} t^2 J_{\mu}(xt)y(t) dt = 0 \quad \text{for } 1 < x < \infty.$$

Solution:

$$y(x) = f(1)J_{\mu-1}(x) + x \int_0^1 t J_{\mu}(xt)f(t) dt.$$

⊙ Reference: B. N. Mandal and N. Mandal (1999, p. 31).

$$13. \int_0^{\infty} t^{2\beta} J_{\mu}(xt)y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} J_{\mu}(xt)y(t) dt = 0 \quad \text{for } 1 < x < \infty.$$

1°. Solution for $\beta > 0$:

$$y(x) = \frac{(2x)^{1-\beta}}{\Gamma(\beta)} \int_0^1 t^{1+\beta} J_{\mu+\beta}(xt)F(t) dt, \quad F(t) = \int_0^1 f(t\zeta)\zeta^{\mu+1}(1-\zeta^2)^{\beta-1} d\zeta. \quad (1)$$

2°. Solution for $\beta > -1$:

$$y(x) = \frac{(2x)^{-\beta}}{\Gamma(1+\beta)} \left[x^{1+\beta} J_{\mu+\beta}(x) \int_0^1 t^{\mu+1}(1-t^2)^{\beta} f(t) dt + \int_0^1 t^{\mu+1}(1-t^2)^{\beta} \Phi(x, t) dt \right], \quad (2)$$

$$\Phi(x, t) = \int_0^1 (x\xi)^{2+\beta} J_{\mu+\beta+1}(x\xi)f(\xi) d\xi.$$

Formula (2) holds for $\beta > -1$ and for $-\mu - \frac{1}{2} < 2\beta < \mu + \frac{3}{2}$. It can be shown that for $\beta > 0$ the solution of Eq. (2) can be reduced to the form (1).

$$14. \int_0^{\infty} t^{-2\alpha} J_{\mu}(xt)y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} t^{-2\beta} J_{\mu}(xt)y(t) dt = g(x) \quad \text{for } 1 < x < \infty.$$

Solution for $0 < \beta - \alpha < 1$:

$$y(x) = \frac{2^{1+\alpha-\beta} x^{1+\alpha+\beta}}{\Gamma(\beta-\alpha)} \int_0^1 t^{1+\alpha-\beta-\mu} J_{\mu+\beta-\alpha}(xt) \left[\int_0^t s^{1+\mu}(t^2-s^2)^{\beta-\alpha-1} f(s) ds \right] dt$$

$$- \frac{2^{\beta-\alpha}}{\Gamma(1+\alpha-\beta)} x^{1+\alpha+\beta} \int_1^{\infty} t^{\mu+\beta-\alpha} J_{\mu+\beta-\alpha}(xt) \left[\frac{d}{dt} \int_t^{\infty} s^{1-\mu}(s^2-t^2)^{\alpha-\beta} g(s) ds \right] dt.$$

⊙ References: C. Nasim and B. D. Aggarwala (1984), B. N. Mandal and N. Mandal (1999, pp. 40–44).

$$15. \int_0^{\infty} J_0(xt)y(t) dt = f(x) \quad \text{for } 0 < x < a,$$

$$\int_0^{\infty} \cos(xt)y(t) dt = g(x) \quad \text{for } a < x < \infty.$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^a \cos(xt) \left[\frac{d}{dt} \int_0^t \frac{s f(s) ds}{\sqrt{t^2-s^2}} \right] dt + \frac{2}{\pi} \int_a^{\infty} \cos(xt)g(t) dt.$$

⊙ Reference: B. N. Mandal and N. Mandal (1999, pp. 194–195).

$$16. \int_0^{\infty} t J_0(xt) y(t) dt = f(x) \quad \text{for } 0 < x < a,$$

$$\int_0^{\infty} \sin(xt) y(t) dt = g(x) \quad \text{for } a < x < \infty.$$

Solution:

$$y(x) = \frac{2}{\pi} \int_0^a \sin(xt) \left[\int_0^t \frac{s f(s) ds}{\sqrt{t^2 - s^2}} \right] dt + \frac{2}{\pi} \int_a^{\infty} \sin(xt) g(t) dt.$$

⊙ Reference: B. N. Mandal and N. Mandal (1999, pp. 195–196).

3.9-3. Kernels Containing Bessel Functions of the Second Kind.

$$17. \int_0^{\infty} t^{-2\alpha} Y_{\mu}(xt) y(t) dt = f(x) \quad \text{for } 0 < x < 1,$$

$$\int_0^{\infty} t^{-2\beta} Y_{\nu}(xt) y(t) dt = g(x) \quad \text{for } 1 < x < \infty.$$

Let $2(\alpha - \beta) = \nu - \mu > 0$, $|\mu| < \frac{1}{2}$, $|\nu| < \frac{1}{2}$.

1°. Solution for $0 < \nu - \mu < 1$:

$$y(x) = -\frac{2^{\nu-\mu}}{\Gamma(1+\mu-\nu)} x^{2\beta+1} \int_0^1 t^{\nu} \mathbf{H}_{\nu}(xt) \left[\frac{d}{dt} \int_t^1 s^{1-\mu} (s^2 - t^2)^{\mu-\nu} f(s) ds \right] dt$$

$$+ \frac{2^{\mu-\nu}}{\Gamma(\mu-\nu)} \int_1^{\infty} t^{1+\mu} \mathbf{H}_{\mu}(xt) \left[\int_t^{\infty} s^{1-\nu} (s^2 - t^2)^{\nu-\mu-1} g(s) ds \right] dt,$$

where $\mathbf{H}_{\mu}(x)$ is the Struve function, which is defined as

$$\mathbf{H}_{\mu}(x) = \sum_{j=0}^{\infty} \frac{(-1)^j (x/2)^{\mu+2j+1}}{\Gamma(j + \frac{3}{2}) \Gamma(\mu + j + \frac{3}{2})}.$$

2°. Solution for $-1 < \nu - \mu < 0$:

$$y(x) = \frac{2^{1-\nu-\mu}}{\Gamma(\mu-\nu)} x^{2\beta+1} \int_0^1 t^{\nu+1} \mathbf{H}_{\nu}(xt) \left[\int_t^1 s^{1-\mu} (s^2 - t^2)^{\mu-\nu-1} f(s) ds \right] dt$$

$$+ \frac{2^{\mu-\nu}}{\Gamma(1-\mu+\nu)} x^{2\alpha+1} \int_1^{\infty} t^{\mu} \mathbf{H}_{\mu}(xt) \left[\int_t^{\infty} s^{1-\nu} (s^2 - t^2)^{\nu-\mu} g(s) ds \right] dt.$$

⊙ References: C. Nasim and B. D. Aggarwala (1984), B. N. Mandal and N. Mandal (1999, pp. 58–59).

3.9-4. Kernels Containing Legendre Spherical Functions of the First Kind, $i^2 = -1$.

$$18. \int_0^{\infty} t P_{-\frac{1}{2}+it}(\cosh x) y(t) dt = f(x) \quad \text{for } 0 < x < a,$$

$$\int_0^{\infty} \tanh(\pi t) P_{-\frac{1}{2}+it}(\cosh x) y(t) dt = 0 \quad \text{for } a < x < \infty.$$

Solution:

$$y(x) = \frac{\sqrt{2}}{\pi} \int_0^a \sin(xt) \left[\int_0^t \frac{f(s) \sinh s}{\sqrt{\cosh t - \cosh s}} ds \right] dt.$$

Note that

$$P_{-\frac{1}{2}+it}(\cosh x) = \frac{\sqrt{2}}{\pi} \int_0^x \frac{\cos(ts)}{\sqrt{\cosh x - \cosh s}} ds, \quad x > 0,$$

where the integral on the right-hand side is called the *Meler integral*.

Chapter 4

Linear Equations of the Second Kind with Constant Limits of Integration

► **Notation:** $f = f(x)$, $g = g(x)$, $h = h(x)$, $v = v(x)$, $w = w(x)$, $K = K(x)$ are arbitrary functions; $A, B, C, D, E, a, b, c, l, \alpha, \beta, \gamma, \delta, \mu$, and ν are arbitrary parameters; n is a nonnegative integer; and i is the imaginary unit.

► **Preliminary remarks.** A number λ is called a *characteristic value* of the integral equation

$$y(x) - \lambda \int_a^b K(x, t)y(t) dt = f(x)$$

if there exist nontrivial solutions of the corresponding homogeneous equation (with $f(x) \equiv 0$). The nontrivial solutions themselves are called the *eigenfunctions* of the integral equation corresponding to the characteristic value λ . If λ is a characteristic value, the number $1/\lambda$ is called an *eigenvalue* of the integral equation. A value of the parameter λ is said to be *regular* if for this value the homogeneous equation has only the trivial solution. Sometimes the characteristic values and the eigenfunctions of a Fredholm integral equation are called the *characteristic values* and the *eigenfunctions of the kernel* $K(x, t)$. In the above equation, it is usually assumed that $a \leq x \leq b$.

4.1. Equations Whose Kernels Contain Power-Law Functions

4.1-1. Kernels Linear in the Arguments x and t .

1. $y(x) - \lambda \int_a^b (x - t)y(t) dt = f(x).$

Solution:

$$y(x) = f(x) + \lambda(A_1x + A_2),$$

where

$$A_1 = \frac{12f_1 + 6\lambda(f_1\Delta_2 - 2f_2\Delta_1)}{\lambda^2\Delta_1^4 + 12}, \quad A_2 = \frac{-12f_2 + 2\lambda(3f_2\Delta_2 - 2f_1\Delta_3)}{\lambda^2\Delta_1^4 + 12},$$
$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b xf(x) dx, \quad \Delta_n = b^n - a^n.$$

2. $y(x) - \lambda \int_a^b (x+t)y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_1 = \frac{6(b+a) + 4\sqrt{3(a^2+ab+b^2)}}{(a-b)^3}, \quad \lambda_2 = \frac{6(b+a) - 4\sqrt{3(a^2+ab+b^2)}}{(a-b)^3}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1x + A_2),$$

where

$$A_1 = \frac{12f_1 - 6\lambda(f_1\Delta_2 - 2f_2\Delta_1)}{12 - 12\lambda\Delta_2 - \lambda^2\Delta_1^4}, \quad A_2 = \frac{12f_2 - 2\lambda(3f_2\Delta_2 - 2f_1\Delta_3)}{12 - 12\lambda\Delta_2 - \lambda^2\Delta_1^4},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b xf(x) dx, \quad \Delta_n = b^n - a^n.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 :

$$y_1(x) = x + \frac{1}{\lambda_1(b-a)} - \frac{b+a}{2}.$$

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. The equation has no multiple characteristic values.

3. $y(x) - \lambda \int_a^b (Ax + Bt)y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{3(A+B)(b+a) \pm \sqrt{9(A-B)^2(b+a)^2 + 48AB(a^2+ab+b^2)}}{AB(a-b)^3}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1x + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{12Af_1 - 6AB\lambda(f_1\Delta_2 - 2f_2\Delta_1)}{12 - 6(A+B)\lambda\Delta_2 - AB\lambda^2\Delta_1^4}, \quad A_2 = \frac{12Bf_2 - 2AB\lambda(3f_2\Delta_2 - 2f_1\Delta_3)}{12 - 6(A+B)\lambda\Delta_2 - AB\lambda^2\Delta_1^4},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b xf(x) dx, \quad \Delta_n = b^n - a^n.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 :

$$y_1(x) = x + \frac{1}{\lambda_1 A(b-a)} - \frac{b+a}{2}.$$

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq \pm B$, where the characteristic value $\lambda_* = \frac{4}{(A+B)(b^2-a^2)}$ is double:

$$y(x) = f(x) + C y_*(x),$$

where C is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* :

$$y_*(x) = x - \frac{(A-B)(b+a)}{4A}.$$

The equation has no multiple characteristic values if $A = \pm B$.

4. $y(x) - \lambda \int_a^b [A + B(x-t)]y(t) dt = f(x).$

This is a special case of equation 4.9.8 with $h(t) = 1$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.8.

5. $y(x) - \lambda \int_a^b (Ax + Bt + C)y(t) dt = f(x).$

This is a special case of equation 4.9.7 with $g(x) = x$ and $h(t) = 1$.

Solution:

$$y(x) = f(x) + \lambda(A_1x + A_2),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.7.

6. $y(x) + A \int_a^b |x-t| y(t) dt = f(x).$

This is a special case of equation 4.9.36 with $g(t) = A$.

1°. The function $y = y(x)$ obeys the following second-order linear nonhomogeneous ordinary differential equation with constant coefficients:

$$y''_{xx} + 2Ay = f''_{xx}(x). \quad (1)$$

The boundary conditions for (1) have the form (see 4.9.36)

$$\begin{aligned} y'_x(a) + y'_x(b) &= f'_x(a) + f'_x(b), \\ y(a) + y(b) + (b-a)y'_x(a) &= f(a) + f(b) + (b-a)f'_x(a). \end{aligned} \quad (2)$$

Equation (1) under the boundary conditions (2) determines the solution of the original integral equation.

2°. For $A < 0$, the general solution of equation (1) is given by

$$y(x) = C_1 \cosh(kx) + C_2 \sinh(kx) + f(x) + k \int_a^x \sinh[k(x-t)]f(t) dt, \quad k = \sqrt{-2A}, \quad (3)$$

where C_1 and C_2 are arbitrary constants.

For $A > 0$, the general solution of equation (1) is given by

$$y(x) = C_1 \cos(kx) + C_2 \sin(kx) + f(x) - k \int_a^x \sin[k(x-t)]f(t) dt, \quad k = \sqrt{2A}. \quad (4)$$

The constants C_1 and C_2 in solutions (3) and (4) are determined by conditions (2).

3°. In the special case $a = 0$ and $A > 0$, the solution of the integral equation is given by formula (4) with

$$C_1 = k \frac{I_s(1 + \cos \lambda) - I_c(\lambda + \sin \lambda)}{2 + 2 \cos \lambda + \lambda \sin \lambda}, \quad C_2 = k \frac{I_s \sin \lambda + I_c(1 + \cos \lambda)}{2 + 2 \cos \lambda + \lambda \sin \lambda},$$

$$k = \sqrt{2A}, \quad \lambda = bk, \quad I_s = \int_0^b \sin[k(b-t)]f(t) dt, \quad I_c = \int_0^b \cos[k(b-t)]f(t) dt.$$

4.1-2. Kernels Quadratic in the Arguments x and t .

7. $y(x) - \lambda \int_a^b (x^2 + t^2)y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_1 = \frac{1}{\frac{1}{3}(b^3 - a^3) + \sqrt{\frac{1}{5}(b^5 - a^5)(b - a)}}, \quad \lambda_2 = \frac{1}{\frac{1}{3}(b^3 - a^3) - \sqrt{\frac{1}{5}(b^5 - a^5)(b - a)}}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 x^2 + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda(\frac{1}{3}f_1\Delta_3 - f_2\Delta_1)}{\lambda^2(\frac{1}{9}\Delta_3^2 - \frac{1}{5}\Delta_1\Delta_5) - \frac{2}{3}\lambda\Delta_3 + 1}, \quad A_2 = \frac{f_2 - \lambda(\frac{1}{3}f_2\Delta_3 - \frac{1}{5}f_1\Delta_5)}{\lambda^2(\frac{1}{9}\Delta_3^2 - \frac{1}{5}\Delta_1\Delta_5) - \frac{2}{3}\lambda\Delta_3 + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^2 f(x) dx, \quad \Delta_n = b^n - a^n.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = x^2 + \sqrt{\frac{b^5 - a^5}{5(b - a)}},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_2(x), \quad y_2(x) = x^2 - \sqrt{\frac{b^5 - a^5}{5(b - a)}},$$

where C is an arbitrary constant and $y_2(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_2 .

4°. The equation has no multiple characteristic values.

$$8. \quad y(x) - \lambda \int_a^b (x^2 - t^2)y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \pm \frac{1}{\sqrt{\frac{1}{9}(b^3 - a^3)^2 - \frac{1}{5}(b^5 - a^5)(b - a)}}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1x^2 + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 + \lambda \left(\frac{1}{3}f_1\Delta_3 - f_2\Delta_1 \right)}{\lambda^2 \left(\frac{1}{5}\Delta_1\Delta_5 - \frac{1}{9}\Delta_2^2 \right) + 1}, \quad A_2 = \frac{-f_2 + \lambda \left(\frac{1}{3}f_2\Delta_3 - \frac{1}{5}f_1\Delta_5 \right)}{\lambda^2 \left(\frac{1}{5}\Delta_1\Delta_5 - \frac{1}{9}\Delta_2^2 \right) + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^2 f(x) dx, \quad \Delta_n = b^n - a^n.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = x^2 + \frac{3 - \lambda_1(b^3 - a^3)}{3\lambda_1(b - a)},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. The equation has no multiple characteristic values.

$$9. \quad y(x) - \lambda \int_a^b (Ax^2 + Bt^2)y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{\frac{1}{3}(A + B)\Delta_3 \pm \sqrt{\frac{1}{9}(A - B)^2\Delta_3^2 + \frac{4}{5}AB\Delta_1\Delta_5}}{2AB\left(\frac{1}{9}\Delta_3^2 - \frac{1}{5}\Delta_1\Delta_5\right)}, \quad \Delta_n = b^n - a^n.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1x^2 + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{Af_1 - AB\lambda \left(\frac{1}{3}f_1\Delta_3 - f_2\Delta_1 \right)}{AB\lambda^2 \left(\frac{1}{9}\Delta_3^2 - \frac{1}{5}\Delta_1\Delta_5 \right) - \frac{1}{3}(A + B)\lambda\Delta_3 + 1},$$

$$A_2 = \frac{Bf_2 - AB\lambda \left(\frac{1}{3}f_2\Delta_3 - \frac{1}{5}f_1\Delta_5 \right)}{AB\lambda^2 \left(\frac{1}{9}\Delta_3^2 - \frac{1}{5}\Delta_1\Delta_5 \right) - \frac{1}{3}(A + B)\lambda\Delta_3 + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^2 f(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = x^2 + \frac{3 - \lambda_1 A(b^3 - a^3)}{3\lambda_1 A(b - a)},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq \pm B$, where $\lambda_* = \frac{6}{(A+B)(b^3 - a^3)}$ is the double characteristic value:

$$y(x) = f(x) + C_1 y_*(x),$$

where C_1 is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* :

$$y_*(x) = x^2 - \frac{(A-B)(b^3 - a^3)}{6A(b-a)}.$$

The equation has no multiple characteristic values if $A = \pm B$.

10. $y(x) - \lambda \int_a^b (xt - t^2)y(t) dt = f(x).$

This is a special case of equation 4.9.8 with $A = 0$, $B = 1$, and $h(t) = t$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.8.

11. $y(x) - \lambda \int_a^b (x^2 - xt)y(t) dt = f(x).$

This is a special case of equation 4.9.10 with $A = 0$, $B = 1$, and $h(x) = x$.

Solution:

$$y(x) = f(x) + \lambda(E_1x^2 + E_2x),$$

where E_1 and E_2 are the constants determined by the formulas presented in 4.9.10.

12. $y(x) - \lambda \int_a^b (Bxt + Ct^2)y(t) dt = f(x).$

This is a special case of equation 4.9.9 with $A = 0$ and $h(t) = t$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.9.

13. $y(x) - \lambda \int_a^b (Bx^2 + Cxt)y(t) dt = f(x).$

This is a special case of equation 4.9.11 with $A = 0$ and $h(x) = x$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^2 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.11.

14. $y(x) - \lambda \int_a^b (Axt + Bx^2 + Cx + D)y(t) dt = f(x).$

This is a special case of equation 4.9.18 with $g_1(x) = Bx^2 + Cx + D$, $h_1(t) = 1$, $g_2(x) = x$, and $h_2(t) = At$.

Solution:

$$y(x) = f(x) + \lambda[A_1(Bx^2 + Cx + D) + A_2x],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$15. \quad y(x) - \lambda \int_a^b (Ax^2 + Bt^2 + Cx + Dt + E)y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = Ax^2 + Cx$, $h_1(t) = 1$, $g_2(x) = 1$, and $h_2(t) = Bt^2 + Dt + E$.

Solution:

$$y(x) = f(x) + \lambda[A_1(Ax^2 + Cx) + A_2],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$16. \quad y(x) - \lambda \int_a^b [Ax + B + (Cx + D)(x - t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = Cx^2 + (A + D)x + B$, $h_1(t) = 1$, $g_2(x) = Cx + D$, and $h_2(t) = -t$.

Solution:

$$y(x) = f(x) + \lambda[A_1(Cx^2 + Ax + Dx + B) + A_2(Cx + D)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$17. \quad y(x) - \lambda \int_a^b [At + B + (Ct + D)(t - x)]y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = 1$, $h_1(t) = Ct^2 + (A + D)t + B$, $g_2(x) = x$, and $h_2(t) = -(Ct + D)$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$18. \quad y(x) - \lambda \int_a^b (x - t)^2 y(t) dt = f(x).$$

This is a special case of equation 4.9.19 with $g(x) = x$, $h(t) = -t$, and $m = 2$.

$$19. \quad y(x) - \lambda \int_a^b (Ax + Bt)^2 y(t) dt = f(x).$$

This is a special case of equation 4.9.19 with $g(x) = Ax$, $h(t) = Bt$, and $m = 2$.

4.1-3. Kernels Cubic in the Arguments x and t .

$$20. \quad y(x) - \lambda \int_a^b (x^3 + t^3)y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_1 = \frac{1}{\frac{1}{4}(b^4 - a^4) + \sqrt{\frac{1}{7}(b^7 - a^7)(b - a)}}, \quad \lambda_2 = \frac{1}{\frac{1}{4}(b^4 - a^4) - \sqrt{\frac{1}{7}(b^7 - a^7)(b - a)}}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1x^3 + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda \left(\frac{1}{4} f_1 \Delta_4 - f_2 \Delta_1 \right)}{\lambda^2 \left(\frac{1}{16} \Delta_4^2 - \frac{1}{7} \Delta_1 \Delta_7 \right) - \frac{1}{2} \lambda \Delta_4 + 1}, \quad A_2 = \frac{f_2 - \lambda \left(\frac{1}{4} f_2 \Delta_4 - \frac{1}{7} f_1 \Delta_7 \right)}{\lambda^2 \left(\frac{1}{16} \Delta_4^2 - \frac{1}{7} \Delta_1 \Delta_7 \right) - \frac{1}{2} \lambda \Delta_4 + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^3 f(x) dx, \quad \Delta_n = b^n - a^n.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = x^3 + \sqrt{\frac{b^7 - a^7}{7(b-a)}},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_2(x), \quad y_2(x) = x^3 - \sqrt{\frac{b^7 - a^7}{7(b-a)}},$$

where C is an arbitrary constant and $y_2(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_2 .

4°. The equation has no multiple characteristic values.

21. $y(x) - \lambda \int_a^b (x^3 - t^3) y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_{1,2} = \pm \frac{1}{\sqrt{\frac{1}{4}(a^4 - b^4)^2 - \frac{1}{7}(a^7 - b^7)(b-a)}}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 x^3 + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 + \lambda \left(\frac{1}{4} f_1 \Delta_4 - f_2 \Delta_1 \right)}{\lambda^2 \left(\frac{1}{7} \Delta_1 \Delta_7 - \frac{1}{16} \Delta_4^2 \right) + 1}, \quad A_2 = \frac{-f_2 + \lambda \left(\frac{1}{4} f_2 \Delta_4 - \frac{1}{7} f_1 \Delta_7 \right)}{\lambda^2 \left(\frac{1}{7} \Delta_1 \Delta_7 - \frac{1}{16} \Delta_4^2 \right) + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^3 f(x) dx, \quad \Delta_n = b^n - a^n.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = x^3 + \frac{4 - \lambda_1(b^4 - a^4)}{4\lambda_1(b-a)},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. The equation has no multiple characteristic values.

$$22. \quad y(x) - \lambda \int_a^b (Ax^3 + Bt^3)y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{\frac{1}{4}(A+B)\Delta_4 \pm \sqrt{\frac{1}{16}(A-B)^2\Delta_4^2 + \frac{4}{7}AB\Delta_1\Delta_7}}{2AB\left(\frac{1}{16}\Delta_4^2 - \frac{1}{7}\Delta_1\Delta_7\right)}, \quad \Delta_n = b^n - a^n.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1x^3 + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{Af_1 - AB\lambda\left(\frac{1}{4}f_1\Delta_4 - f_2\Delta_1\right)}{AB\lambda^2\left(\frac{1}{16}\Delta_4^2 - \frac{1}{7}\Delta_1\Delta_7\right) - \frac{1}{4}\lambda(A+B)\Delta_4 + 1},$$

$$A_2 = \frac{Bf_2 - AB\lambda\left(\frac{1}{4}f_2\Delta_4 - \frac{1}{7}f_1\Delta_7\right)}{AB\lambda^2\left(\frac{1}{16}\Delta_4^2 - \frac{1}{7}\Delta_1\Delta_7\right) - \frac{1}{4}\lambda(A+B)\Delta_4 + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^3 f(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = x^3 + \frac{4 - \lambda_1 A(b^4 - a^4)}{4\lambda_1 A(b - a)},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq \pm B$, where $\lambda_* = \frac{8}{(A+B)(b^4 - a^4)}$ is the double characteristic value:

$$y(x) = f(x) + Cy_*(x), \quad y_*(x) = x^3 - \frac{(A-B)(b^4 - a^4)}{8A(b-a)},$$

where C is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A = \pm B$.

$$23. \quad y(x) - \lambda \int_a^b (xt^2 - t^3)y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $A = 0$, $B = 1$, and $h(t) = t^2$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.8.

$$24. \quad y(x) - \lambda \int_a^b (Bxt^2 + Ct^3)y(t) dt = f(x).$$

This is a special case of equation 4.9.9 with $A = 0$ and $h(t) = t^2$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.9.

$$25. \quad y(x) - \lambda \int_a^b (Ax^2t + Bxt^2)y(t) dt = f(x).$$

This is a special case of equation 4.9.17 with $g(x) = x^2$ and $h(x) = x$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^2 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.17.

$$26. \quad y(x) - \lambda \int_a^b (Ax^3 + Bxt^2)y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = x^3$, $h_1(t) = A$, $g_2(x) = x$, and $h_2(t) = Bt^2$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^3 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$27. \quad y(x) - \lambda \int_a^b (Ax^3 + Bx^2t + Cx^2 + D)y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = Ax^3 + Cx^2 + D$, $h_1(t) = 1$, $g_2(x) = x^2$, and $h_2(t) = Bt$.

Solution:

$$y(x) = f(x) + \lambda[A_1(Ax^3 + Cx^2 + D) + A_2x^2],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$28. \quad y(x) - \lambda \int_a^b (Axt^2 + Bt^3 + Ct^2 + D)y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = x$, $h_1(t) = At^2$, $g_2(x) = 1$, and $h_2(t) = Bt^3 + Ct^2 + D$.

Solution:

$$y(x) = f(x) + \lambda(A_1x + A_2),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$29. \quad y(x) - \lambda \int_a^b (x-t)^3 y(t) dt = f(x).$$

This is a special case of equation 4.9.19 with $g(x) = x$, $h(t) = -t$, and $m = 3$.

$$30. \quad y(x) - \lambda \int_a^b (Ax + Bt)^3 y(t) dt = f(x).$$

This is a special case of equation 4.9.19 with $g(x) = Ax$, $h(t) = Bt$, and $m = 3$.

4.1-4. Kernels Containing Higher-Order Polynomials in x and t .

$$31. \quad y(x) - \lambda \int_a^b (x^n + t^n)y(t) dt = f(x), \quad n = 1, 2, \dots$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{1}{\Delta_n \pm \sqrt{\Delta_0 \Delta_{2n}}}, \quad \text{where } \Delta_n = \frac{1}{n+1}(b^{n+1} - a^{n+1}).$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 x^n + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda(f_1 \Delta_n - f_2 \Delta_0)}{\lambda^2(\Delta_n^2 - \Delta_0 \Delta_{2n}) - 2\lambda \Delta_n + 1}, \quad A_2 = \frac{f_2 - \lambda(f_2 \Delta_n - f_1 \Delta_{2n})}{\lambda^2(\Delta_n^2 - \Delta_0 \Delta_{2n}) - 2\lambda \Delta_n + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^n f(x) dx, \quad \Delta_n = \frac{1}{n+1}(b^{n+1} - a^{n+1}).$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = x^n + \sqrt{\Delta_{2n}/\Delta_0},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_2(x), \quad y_2(x) = x^n - \sqrt{\Delta_{2n}/\Delta_0},$$

where C is an arbitrary constant and $y_2(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_2 .

4°. The equation has no multiple characteristic values.

$$32. \quad y(x) - \lambda \int_a^b (x^n - t^n)y(t) dt = f(x), \quad n = 1, 2, \dots$$

The characteristic values of the equation:

$$\lambda_{1,2} = \pm \left[\frac{1}{(n+1)^2} (b^{n+1} - a^{n+1})^2 - \frac{1}{2n+1} (b^{2n+1} - a^{2n+1})(b-a) \right]^{-1/2}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 x^n + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 + \lambda(f_1 \Delta_n - f_2 \Delta_0)}{\lambda^2(\Delta_0 \Delta_{2n} - \Delta_n^2) + 1}, \quad A_2 = \frac{-f_2 + \lambda(f_2 \Delta_n - f_1 \Delta_{2n})}{\lambda^2(\Delta_0 \Delta_{2n} - \Delta_n^2) + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^n f(x) dx, \quad \Delta_n = \frac{1}{n+1}(b^{n+1} - a^{n+1}).$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = x^n + \frac{1 - \lambda_1 \Delta_n}{\lambda_1 \Delta_0},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. The equation has no multiple characteristic values.

$$33. \quad y(x) - \lambda \int_a^b (Ax^n + Bt^n)y(t) dt = f(x), \quad n = 1, 2, \dots$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{(A+B)\Delta_n \pm \sqrt{(A-B)^2\Delta_n^2 + 4AB\Delta_0\Delta_{2n}}}{2AB(\Delta_n^2 - \Delta_0\Delta_{2n})}, \quad \Delta_n = \frac{1}{n+1}(b^{n+1} - a^{n+1}).$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1x^n + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{Af_1 - AB\lambda(f_1\Delta_n - f_2\Delta_0)}{AB\lambda^2(\Delta_n^2 - \Delta_0\Delta_{2n}) - (A+B)\lambda\Delta_n + 1},$$

$$A_2 = \frac{Bf_2 - AB\lambda(f_2\Delta_n - f_1\Delta_{2n})}{AB\lambda^2(\Delta_n^2 - \Delta_0\Delta_{2n}) - (A+B)\lambda\Delta_n + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b x^n f(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = x^n + \frac{1 - A\lambda_1\Delta_n}{A\lambda_1\Delta_0},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq \pm B$, where the characteristic value $\lambda_* = 2/[(A+B)\Delta_n]$ is double:

$$y(x) = f(x) + Cy_*(x), \quad y_*(x) = x^n - \frac{(A-B)\Delta_n}{2A\Delta_0}.$$

Here C is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A = \pm B$.

$$34. \quad y(x) - \lambda \int_a^b (x-t)t^m y(t) dt = f(x), \quad m = 1, 2, \dots$$

This is a special case of equation 4.9.8 with $A = 0$, $B = 1$, and $h(t) = t^m$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.8.

$$35. \quad y(x) - \lambda \int_a^b (x-t)x^m y(t) dt = f(x), \quad m = 1, 2, \dots$$

This is a special case of equation 4.9.10 with $A = 0$, $B = 1$, and $h(x) = x^m$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^{m+1} + A_2x^m),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.10.

$$36. \quad y(x) - \lambda \int_a^b (Ax^{m+1} + Bx^m t + Cx^m + D)y(t) dt = f(x), \quad m = 1, 2, \dots$$

This is a special case of equation 4.9.18 with $g_1(x) = Ax^{m+1} + Cx^m + D$, $h_1(t) = 1$, $g_2(x) = x^m$, and $h_2(t) = Bt$.

Solution:

$$y(x) = f(x) + \lambda[A_1(Ax^{m+1} + Cx^m + D) + A_2x^m],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$37. \quad y(x) - \lambda \int_a^b (Axt^m + Bt^{m+1} + Ct^m + D)y(t) dt = f(x), \quad m = 1, 2, \dots$$

This is a special case of equation 4.9.18 with $g_1(x) = x$, $h_1(t) = At^m$, $g_2(x) = 1$, and $h_2(t) = Bt^{m+1} + Ct^m + D$.

Solution:

$$y(x) = f(x) + \lambda(A_1x + A_2),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$38. \quad y(x) - \lambda \int_a^b (Ax^n t^n + Bx^m t^m)y(t) dt = f(x), \quad n, m = 1, 2, \dots, \quad n \neq m.$$

This is a special case of equation 4.9.14 with $g(x) = x^n$ and $h(t) = t^m$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^n + A_2x^m),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.14.

$$39. \quad y(x) - \lambda \int_a^b (Ax^n t^m + Bx^m t^n)y(t) dt = f(x), \quad n, m = 1, 2, \dots, \quad n \neq m.$$

This is a special case of equation 4.9.17 with $g(x) = x^n$ and $h(t) = t^m$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^n + A_2x^m),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.17.

$$40. \quad y(x) - \lambda \int_a^b (x-t)^m y(t) dt = f(x), \quad m = 1, 2, \dots$$

This is a special case of equation 4.9.19 with $g(x) = x$ and $h(t) = -t$.

$$41. \quad y(x) - \lambda \int_a^b (Ax + Bt)^m y(t) dt = f(x), \quad m = 1, 2, \dots$$

This is a special case of equation 4.9.19 with $g(x) = Ax$ and $h(t) = Bt$.

$$42. \quad y(x) + A \int_a^b |x-t|t^k y(t) dt = f(x).$$

This is a special case of equation 4.9.36 with $g(t) = At^k$. Solving the integral equation is reduced to solving the ordinary differential equation $y''_{xx} + 2Ax^k y = f''_{xx}(x)$, the general solution of which can be expressed via Bessel functions or modified Bessel functions (the boundary conditions are given in 4.9.36).

$$43. \quad y(x) + A \int_a^b |x-t|^{2n+1} y(t) dt = f(x), \quad n = 0, 1, 2, \dots$$

Let us remove the modulus in the integrand:

$$y(x) + A \int_a^x (x-t)^{2n+1} y(t) dt + A \int_x^b (t-x)^{2n+1} y(t) dt = f(x). \quad (1)$$

The k -fold differentiation of (1) with respect to x yields

$$y_x^{(k)}(x) + AB_k \int_a^x (x-t)^{2n+1-k} y(t) dt + (-1)^k AB_k \int_x^b (t-x)^{2n+1-k} y(t) dt = f_x^{(k)}(x), \quad (2)$$

$$B_k = (2n+1)(2n) \dots (2n+2-k), \quad k = 1, 2, \dots, 2n+1.$$

Differentiating (2) with $k = 2n+1$, we arrive at the following linear nonhomogeneous differential equation with constant coefficients for $y = y(x)$:

$$y_x^{(2n+2)} + 2(2n+1)! Ay = f_x^{(2n+2)}(x). \quad (3)$$

Equation (3) must satisfy the initial conditions which can be obtained by setting $x = a$ in (1) and (2):

$$y(a) + A \int_a^b (t-a)^{2n+1} y(t) dt = f(a), \quad (4)$$

$$y_x^{(k)}(a) + (-1)^k AB_k \int_a^b (t-a)^{2n+1-k} y(t) dt = f_x^{(k)}(a), \quad k = 1, 2, \dots, 2n+1.$$

These conditions can be reduced to a more habitual form containing no integrals. To this end, y must be expressed from equation (3) in terms of $y_x^{(2n+2)}$ and $f_x^{(2n+2)}$ and substituted into (4), and then one must integrate the resulting expressions by parts (sufficiently many times).

4.1-5. Kernels Containing Rational Functions.

$$44. \quad y(x) - \lambda \int_a^b \left(\frac{1}{x} + \frac{1}{t} \right) y(t) dt = f(x).$$

This is a special case of equation 4.9.2 with $g(x) = 1/x$.

Solution:

$$y(x) = f(x) + \lambda \left(\frac{A_1}{x} + A_2 \right),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.2.

$$45. \quad y(x) - \lambda \int_a^b \left(\frac{1}{x} - \frac{1}{t} \right) y(t) dt = f(x).$$

This is a special case of equation 4.9.3 with $g(x) = 1/x$.

Solution:

$$y(x) = f(x) + \lambda \left(\frac{A_1}{x} + A_2 \right),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.3.

$$46. \quad y(x) - \lambda \int_a^b \left(\frac{A}{x} + \frac{B}{t} \right) y(t) dt = f(x).$$

This is a special case of equation 4.9.4 with $g(x) = 1/x$.

Solution:

$$y(x) = f(x) + \lambda \left(\frac{A_1}{x} + A_2 \right),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.4.

$$47. \quad y(x) - \lambda \int_a^b \left(\frac{A}{x + \alpha} + \frac{B}{t + \beta} \right) y(t) dt = f(x).$$

This is a special case of equation 4.9.5 with $g(x) = \frac{A}{x + \alpha}$ and $h(t) = \frac{B}{t + \beta}$.

Solution:

$$y(x) = f(x) + \lambda \left(A_1 \frac{A}{x + \alpha} + A_2 \right),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.5.

$$48. \quad y(x) - \lambda \int_a^b \left(\frac{x}{t} - \frac{t}{x} \right) y(t) dt = f(x).$$

This is a special case of equation 4.9.16 with $g(x) = x$ and $h(t) = 1/t$.

Solution:

$$y(x) = f(x) + \lambda \left(A_1 x + \frac{A_2}{x} \right),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.16.

$$49. \quad y(x) - \lambda \int_a^b \left(\frac{Ax}{t} + \frac{Bt}{x} \right) y(t) dt = f(x).$$

This is a special case of equation 4.9.17 with $g(x) = x$ and $h(t) = 1/t$.

Solution:

$$y(x) = f(x) + \lambda \left(A_1 x + \frac{A_2}{x} \right),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.17.

$$50. \quad y(x) - \lambda \int_a^b \left(A \frac{x + \alpha}{t + \beta} + B \frac{t + \alpha}{x + \beta} \right) y(t) dt = f(x).$$

This is a special case of equation 4.9.17 with $g(x) = x + \alpha$ and $h(t) = \frac{1}{t + \beta}$.

Solution:

$$y(x) = f(x) + \lambda \left[A_1(x + \alpha) + \frac{A_2}{x + \beta} \right],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.17.

$$51. \quad y(x) - \lambda \int_a^b \left[A \frac{(x + \alpha)^n}{(t + \beta)^m} + B \frac{(t + \alpha)^n}{(x + \beta)^m} \right] y(t) dt = f(x), \quad n, m = 0, 1, 2, \dots$$

This is a special case of equation 4.9.17 with $g(x) = (x + \alpha)^n$ and $h(t) = (t + \beta)^{-m}$.

Solution:

$$y(x) = f(x) + \lambda \left[A_1(x + \alpha)^n + \frac{A_2}{(x + \beta)^m} \right],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.17.

$$52. \quad y(x) - \lambda \int_1^{\infty} \frac{y(t)}{x+t} dt = f(x), \quad 1 \leq x < \infty, \quad -\infty < \pi\lambda < 1.$$

Solution:

$$y(x) = \int_0^{\infty} \frac{\tau \sinh(\pi\tau) F(\tau)}{\cosh(\pi\tau) - \pi\lambda} P_{-\frac{1}{2}+i\tau}(x) d\tau,$$

$$F(\tau) = \int_1^{\infty} f(x) P_{-\frac{1}{2}+i\tau}(x) dx,$$

where $P_{\nu}(x) = F(-\nu, \nu + 1, 1; \frac{1}{2}(1-x))$ is the Legendre spherical function of the first kind, for which the integral representation

$$P_{-\frac{1}{2}+i\tau}(\cosh \alpha) = \frac{2}{\pi} \int_0^{\alpha} \frac{\cos(\tau s) ds}{\sqrt{2(\cosh \alpha - \cosh s)}} \quad (\alpha \geq 0)$$

can be used.

⊙ Reference: V. A. Ditkin and A. P. Prudnikov (1965).

$$53. \quad (x^2 + b^2)y(x) = \frac{\lambda}{\pi} \int_{-\infty}^{\infty} \frac{a^3 y(t)}{a^2 + (x-t)^2} dt.$$

This equation is encountered in atomic and nuclear physics.

We seek the solution in the form

$$y(x) = \sum_{m=0}^{\infty} \frac{A_m x}{x^2 + (am + b)^2}. \quad (1)$$

The coefficients A_m obey the equations

$$mA_m \left(\frac{m+2b}{a} \right) + \lambda A_{m-1} = 0, \quad \sum_{m=0}^{\infty} A_m = 0. \quad (2)$$

Using the first equation of (2) to express all A_m via A_0 (A_0 can be chosen arbitrarily), substituting the result into the second equation of (2), and dividing by A_0 , we obtain

$$1 + \sum_{m=1}^{\infty} \frac{(-\lambda)^m}{m!} \frac{1}{(1+2b/a)(2+2b/a)\dots(m+2b/a)} = 0. \quad (3)$$

It follows from the definitions of the Bessel functions of the first kind that equation (3) can be rewritten in the form

$$\lambda^{-b/a} J_{2b/a}(2\sqrt{\lambda}) = 0. \quad (4)$$

In this sort of problem, a and λ are usually assumed to be given and b , which is proportional to the system energy, to be unknown. The quantity b can be determined by tables of zeros of Bessel functions. In some cases, b and a are given and λ is unknown.

⊙ Reference: I. Sneddon (1995).

$$54. \quad \int_{-1}^1 \frac{y(x) - y(t)}{|x-t|} dt = \lambda y(x).$$

The characteristic values of the equation:

$$\lambda_n = 2 \left(1 + \frac{1}{2} + \dots + \frac{1}{n} \right), \quad \text{where } n = 1, 2, \dots$$

The eigenfunctions of the equation:

$$y_n(x) = P_n(x), \quad \text{where } n = 1, 2, \dots$$

Here $P_n(x) = \frac{1}{n! 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n$ are the Legendre polynomials.

⊙ Reference: A. G. Petrov (1986).

4.1-6. Kernels Containing Arbitrary Powers.

$$55. \quad y(x) - \lambda \int_a^b (x-t)t^\mu y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $A = 0$, $B = 1$, and $h(t) = t^\mu$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.8.

$$56. \quad y(x) - \lambda \int_a^b (x-t)x^\nu y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $A = 0$, $B = 1$, and $h(x) = x^\nu$.

Solution:

$$y(x) = f(x) + \lambda(E_1x^{\nu+1} + E_2x^\nu),$$

where E_1 and E_2 are the constants determined by the formulas presented in 4.9.10.

$$57. \quad y(x) - \lambda \int_a^b (x^\mu - t^\mu)y(t) dt = f(x).$$

This is a special case of equation 4.9.3 with $g(x) = x^\mu$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^\mu + A_2),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.3.

$$58. \quad y(x) - \lambda \int_a^b (Ax^\nu + Bt^\nu)t^\mu y(t) dt = f(x).$$

This is a special case of equation 4.9.6 with $g(x) = x^\nu$ and $h(t) = t^\mu$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^\nu + A_2),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.6.

$$59. \quad y(x) - \lambda \int_a^b (Dx^\nu + Et^\mu)x^\gamma y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = x^{\nu+\gamma}$, $h_1(t) = D$, $g_2(x) = x^\gamma$, and $h_2(t) = Et^\mu$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^{\nu+\gamma} + A_2x^\gamma),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$60. \quad y(x) - \lambda \int_a^b (Ax^\nu t^\mu + Bx^\gamma t^\delta)y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = x^\nu$, $h_1(t) = At^\mu$, $g_2(x) = x^\gamma$, and $h_2(t) = Bt^\delta$.

Solution:

$$y(x) = f(x) + \lambda(A_1x^\nu + A_2x^\gamma),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$61. \quad y(x) - \lambda \int_a^b (A + Bxt^\mu + Ct^{\mu+1})y(t) dt = f(x).$$

This is a special case of equation 4.9.9 with $h(t) = t^\mu$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.9.

$$62. \quad y(x) - \lambda \int_a^b (At^\alpha + Bx^\beta t^\mu + Ct^{\mu+\gamma})y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = 1$, $h_1(t) = At^\alpha + Ct^{\mu+\gamma}$, $g_2(x) = x^\beta$, and $h_2(t) = Bt^\mu$.

Solution:

$$y(x) = f(x) + \lambda(A_1 + A_2x^\beta),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$63. \quad y(x) - \lambda \int_a^b (Ax^\alpha t^\gamma + Bx^\beta t^\gamma + Cx^\mu t^\nu)y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = Ax^\alpha + Bx^\beta$, $h_1(t) = t^\gamma$, $g_2(x) = x^\mu$, and $h_2(t) = Ct^\nu$.

Solution:

$$y(x) = f(x) + \lambda[A_1(Ax^\alpha + Bx^\beta) + A_2x^\mu],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$64. \quad y(x) - \lambda \int_a^b \left[A \frac{(x+p_1)^\beta}{(t+q_1)^\gamma} + B \frac{(x+p_2)^\mu}{(t+q_2)^\delta} \right] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = (x+p_1)^\beta$, $h_1(t) = A(t+q_1)^{-\gamma}$, $g_2(x) = (x+p_2)^\mu$, and $h_2(t) = B(t+q_2)^{-\delta}$.

Solution:

$$y(x) = f(x) + \lambda[A_1(x+p_1)^\beta + A_2(x+p_2)^\mu],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$65. \quad y(x) - \lambda \int_a^b \left(A \frac{x^\mu + a}{t^\nu + b} + B \frac{x^\gamma + c}{t^\delta + d} \right) y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = x^\mu + a$, $h_1(t) = \frac{A}{t^\nu + b}$, $g_2(x) = x^\gamma + c$, and $h_2(t) = \frac{B}{t^\delta + d}$.

Solution:

$$y(x) = f(x) + \lambda[A_1(x^\mu + a) + A_2(x^\gamma + c)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

4.1-7. Singular Equations.

In this subsection, all singular integrals are understood in the sense of the Cauchy principal value.

$$66. \quad Ay(x) + \frac{B}{\pi} \int_{-1}^1 \frac{y(t) dt}{t-x} = f(x), \quad -1 < x < 1.$$

Here A and B are real numbers such that $B \neq 0$, $A \pm B \neq 0$, and $A^2 + B^2 = 1$.

1°. The solution bounded at the endpoints:

$$y(x) = Af(x) - \frac{B}{\pi} \int_{-1}^1 \frac{g(t) f(t) dt}{g(t) t-x}, \quad g(x) = (1+x)^\alpha (1-x)^{1-\alpha}, \quad (1)$$

where α is the solution of the trigonometric equation

$$A + B \cot(\pi\alpha) = 0 \quad (2)$$

on the interval $0 < \alpha < 1$. This solution $y(x)$ exists if and only if $\int_{-1}^1 \frac{f(t)}{g(t)} dt = 0$.

2°. The solution bounded at the endpoint $x = 1$ and unbounded at the endpoint $x = -1$:

$$y(x) = Af(x) - \frac{B}{\pi} \int_{-1}^1 \frac{g(t) f(t) dt}{g(t) t-x}, \quad g(x) = (1+x)^\alpha (1-x)^{-\alpha}, \quad (3)$$

where α is the solution of the trigonometric equation (2) on the interval $-1 < \alpha < 0$.

3°. The solution unbounded at the endpoints:

$$y(x) = Af(x) - \frac{B}{\pi} \int_{-1}^1 \frac{g(t) f(t) dt}{g(t) t-x} + Cg(x), \quad g(x) = (1+x)^\alpha (1-x)^{-1-\alpha}, \quad (4)$$

where C is an arbitrary constant and α is the solution of the trigonometric equation (2) on the interval $-1 < \alpha < 0$.

⊙ References: N. I. Muskhelishvili (1992), I. K. Lifanov, L. N. Poltavskii, and G. M. Vainikko (2004, pp. 6–7).

$$67. \quad y(x) - \lambda \int_{-\infty}^{\infty} \frac{y(t) dt}{t-x} = f(x).$$

Solution:

$$y(x) = \frac{1}{1 + \pi^2 \lambda^2} \left[f(x) + \lambda \int_{-\infty}^{\infty} \frac{f(t) dt}{t-x} \right].$$

⊙ Reference: M. L. Krasnov (1975).

$$68. \quad y(x) - \lambda \int_0^1 \left(\frac{1}{t-x} - \frac{1}{x+t-2xt} \right) y(t) dt = f(x), \quad 0 < x < 1.$$

Tricomi's equation.

Solution:

$$y(x) = \frac{1}{1 + \lambda^2 \pi^2} \left[f(x) + \int_0^1 \frac{t^\alpha (1-x)^\alpha}{x^\alpha (1-t)^\alpha} \left(\frac{1}{t-x} - \frac{1}{x+t-2xt} \right) f(t) dt \right] + \frac{C(1-x)^\beta}{x^{1+\beta}},$$

$$\alpha = \frac{2}{\pi} \arctan(\lambda\pi) \quad (-1 < \alpha < 1), \quad \tan \frac{\beta\pi}{2} = \lambda\pi \quad (-2 < \beta < 0),$$

where C is an arbitrary constant.

⊙ References: P. P. Zabreyko, A. I. Koshelev, et al. (1975), F. G. Tricomi (1985).

$$69. \quad y(x) + \lambda \int_{-1}^1 \left(\frac{1+t}{1+x} \right)^{\frac{2}{n+2}} \left(\frac{1}{t-x} - \frac{1}{1-xt} \right) y(t) dt = f(x), \quad 0 < x < 1.$$

Tricomi–Gellerstedt equation.

In the class of functions $y(x)$ for which integrals $\int_{-1}^1 |y(x)| \ln|x \pm 1| dx$ are finite the unique solution of the equation has the form

$$y(x) = \frac{1}{1 + \lambda^2 \pi^2} \left[f(x) - \lambda \int_{-1}^1 \left(\frac{1-t^2}{1-x^2} \right)^{\frac{1}{n+2}} \left(\frac{1}{t-x} - \frac{1}{1-xt} \right) f(t) dt \right].$$

⊙ Reference: S. G. Mikhlin (1967).

4.2. Equations Whose Kernels Contain Exponential Functions

4.2-1. Kernels Containing Exponential Functions.

$$1. \quad y(x) - \lambda \int_a^b (e^{\beta x} + e^{\beta t}) y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{\beta}{e^{\beta b} - e^{\beta a} \pm \sqrt{\frac{1}{2}\beta(b-a)(e^{2\beta b} - e^{2\beta a})}}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 e^{\beta x} + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda[f_1 \Delta_\beta - (b-a)f_2]}{\lambda^2[\Delta_\beta^2 - (b-a)\Delta_{2\beta}] - 2\lambda\Delta_\beta + 1}, \quad A_2 = \frac{f_2 - \lambda(f_2 \Delta_\beta - f_1 \Delta_{2\beta})}{\lambda^2[\Delta_\beta^2 - (b-a)\Delta_{2\beta}] - 2\lambda\Delta_\beta + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b f(x) e^{\beta x} dx, \quad \Delta_\beta = \frac{1}{\beta}(e^{\beta b} - e^{\beta a}).$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = e^{\beta x} + \sqrt{\frac{e^{2\beta b} - e^{2\beta a}}{2\beta(b-a)}},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_2(x), \quad y_2(x) = e^{\beta x} - \sqrt{\frac{e^{2\beta b} - e^{2\beta a}}{2\beta(b-a)}},$$

where C is an arbitrary constant and $y_2(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_2 .

4°. The equation has no multiple characteristic values.

$$2. \quad y(x) - \lambda \int_a^b (e^{\beta x} - e^{\beta t})y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \pm \frac{\beta}{\sqrt{(e^{\beta b} - e^{\beta a})^2 - \frac{1}{2}\beta(b-a)(e^{2\beta b} - e^{2\beta a})}}.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 e^{\beta x} + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 + \lambda[f_1 \Delta_\beta - (b-a)f_2]}{\lambda^2[(b-a)\Delta_{2\beta} - \Delta_\beta^2] + 1}, \quad A_2 = \frac{-f_2 + \lambda(f_2 \Delta_\beta - f_1 \Delta_{2\beta})}{\lambda^2[(b-a)\Delta_{2\beta} - \Delta_\beta^2] + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b f(x)e^{\beta x} dx, \quad \Delta_\beta = \frac{1}{\beta}(e^{\beta b} - e^{\beta a}).$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = e^{\beta x} + \frac{1 - \lambda_1 \Delta_\beta}{\lambda_1(b-a)},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. The equation has no multiple characteristic values.

$$3. \quad y(x) - \lambda \int_a^b (Ae^{\beta x} + Be^{\beta t})y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{(A+B)\Delta_\beta \pm \sqrt{(A-B)^2 \Delta_\beta^2 + 4AB(b-a)\Delta_{2\beta}}}{2AB[\Delta_\beta^2 - (b-a)\Delta_{2\beta}]}, \quad \Delta_\beta = \frac{1}{\beta}(e^{\beta b} - e^{\beta a}).$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 e^{\beta x} + A_2),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{A f_1 - AB\lambda[f_1 \Delta_\beta - (b-a)f_2]}{AB\lambda^2[\Delta_\beta^2 - (b-a)\Delta_{2\beta}] - (A+B)\lambda\Delta_\beta + 1},$$

$$A_2 = \frac{B f_2 - AB\lambda(f_2 \Delta_\beta - f_1 \Delta_{2\beta})}{AB\lambda^2[\Delta_\beta^2 - (b-a)\Delta_{2\beta}] - (A+B)\lambda\Delta_\beta + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b f(x)e^{\beta x} dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = e^{\beta x} + \frac{1 - A\lambda_1 \Delta_\beta}{A(b-a)\lambda_1},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq \pm B$, where the characteristic value $\lambda_* = \frac{2}{(A+B)\Delta_\beta}$ is double:

$$y(x) = f(x) + Cy_*(x), \quad y_*(x) = e^{\beta x} - \frac{(A-B)\Delta_\beta}{2A(b-a)},$$

where C is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A = \pm B$.

4.
$$y(x) - \lambda \int_a^b [Ae^{\beta(x-t)} + B]y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\beta x}$, $h_1(t) = Ae^{-\beta t}$, $g_2(x) = 1$, and $h_2(t) = B$.

Solution:

$$y(x) = f(x) + \lambda(A_1 e^{\beta x} + A_2),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

5.
$$y(x) - \lambda \int_a^b [Ae^{\beta x + \mu t} + Be^{(\beta + \mu)t}]y(t) dt = f(x).$$

This is a special case of equation 4.9.6 with $g(x) = e^{\beta x}$ and $h(t) = e^{\mu t}$.

Solution:

$$y(x) = f(x) + \lambda(A_1 e^{\beta x} + A_2),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.6.

6.
$$y(x) - \lambda \int_a^b [Ae^{\alpha(x+t)} + Be^{\beta(x+t)}]y(t) dt = f(x).$$

This is a special case of equation 4.9.14 with $g(x) = e^{\alpha x}$ and $h(t) = e^{\beta t}$.

Solution:

$$y(x) = f(x) + \lambda(A_1 e^{\alpha x} + A_2 e^{\beta x}),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.14.

7.
$$y(x) - \lambda \int_a^b (Ae^{\alpha x + \beta t} + Be^{\beta x + \alpha t})y(t) dt = f(x).$$

This is a special case of equation 4.9.17 with $g(x) = e^{\alpha x}$ and $h(t) = e^{\beta t}$.

Solution:

$$y(x) = f(x) + \lambda(A_1 e^{\alpha x} + A_2 e^{\beta x}),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.17.

8.
$$y(x) - \lambda \int_a^b [De^{(\gamma + \mu)x} + Ee^{\nu t + \mu x}]y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{(\gamma + \mu)x}$, $h_1(t) = D$, $g_2(x) = e^{\mu x}$, and $h_2(t) = Ee^{\nu t}$.

Solution:

$$y(x) = f(x) + \lambda[A_1 e^{(\gamma + \mu)x} + A_2 e^{\mu x}],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$9. \quad y(x) - \lambda \int_a^b (Ae^{\alpha x + \beta t} + Be^{\gamma x + \delta t})y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\alpha x}$, $h_1(t) = Ae^{\beta t}$, $g_2(x) = e^{\gamma x}$, and $h_2(t) = Be^{\delta t}$.

Solution:

$$y(x) = f(x) + \lambda(A_1 e^{\alpha x} + A_2 e^{\gamma x}),$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.18.

$$10. \quad y(x) - \lambda \int_a^b \left[\sum_{k=1}^n A_k e^{\gamma_k(x-t)} \right] y(t) dt = f(x).$$

This is a special case of equation 4.9.20 with $g_k(x) = e^{\gamma_k x}$ and $h_k(t) = A_k e^{-\gamma_k t}$.

$$11. \quad y(x) - \frac{1}{2} \int_0^\infty e^{-|x-t|} y(t) dt = Ae^{\mu x}, \quad 0 < \mu < 1.$$

Solution:

$$y(x) = C(1+x) + A\mu^{-2}[(\mu^2 - 1)e^{\mu x} - \mu + 1],$$

where C is an arbitrary constant.

⊙ Reference: P. P. Zabreyko, A. I. Koshelev, et al. (1975).

$$12. \quad y(x) + \lambda \int_0^\infty e^{-|x-t|} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_0^\infty \exp(-\sqrt{1+2\lambda}|x-t|) f(t) dt \\ + \left(1 - \frac{\lambda+1}{\sqrt{1+2\lambda}} \right) \int_0^\infty \exp[-\sqrt{1+2\lambda}(x+t)] f(t) dt,$$

where $\lambda > -\frac{1}{2}$.

⊙ Reference: F. D. Gakhov and Yu. I. Cherskii (1978).

$$13. \quad y(x) - \lambda \int_{-\infty}^\infty e^{-|x-t|} y(t) dt = 0, \quad \lambda > 0.$$

The Lalesco–Picard equation.

Solution:

$$y(x) = \begin{cases} C_1 \exp(x\sqrt{1-2\lambda}) + C_2 \exp(-x\sqrt{1-2\lambda}) & \text{for } 0 < \lambda < \frac{1}{2}, \\ C_1 + C_2 x & \text{for } \lambda = \frac{1}{2}, \\ C_1 \cos(x\sqrt{2\lambda-1}) + C_2 \sin(x\sqrt{2\lambda-1}) & \text{for } \lambda > \frac{1}{2}, \end{cases}$$

where C_1 and C_2 are arbitrary constants.

⊙ Reference: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971).

$$14. \quad y(x) + \lambda \int_{-\infty}^{\infty} e^{-|x-t|} y(t) dt = f(x).$$

1°. Solution with $\lambda > -\frac{1}{2}$:

$$y(x) = f(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) f(t) dt.$$

2°. If $\lambda \leq -\frac{1}{2}$, for the equation to be solvable the conditions

$$\int_{-\infty}^{\infty} f(x) \cos(ax) dx = 0, \quad \int_{-\infty}^{\infty} f(x) \sin(ax) dx = 0,$$

where $a = \sqrt{-1-2\lambda}$, must be satisfied. In this case, the solution has the form

$$y(x) = f(x) - \frac{a^2 + 1}{2a} \int_0^{\infty} \sin(at) f(x+t) dt, \quad (-\infty < x < \infty).$$

In the class of solutions not belonging to $L_2(-\infty, \infty)$, the homogeneous equation (with $f(x) \equiv 0$) has a nontrivial solution. In this case, the general solution of the corresponding nonhomogeneous equation with $\lambda \leq -\frac{1}{2}$ has the form

$$y(x) = C_1 \sin(ax) + C_2 \cos(ax) + f(x) - \frac{a^2 + 1}{4a} \int_{-\infty}^{\infty} \sin(a|x-t|) f(t) dt.$$

⊙ Reference: F. D. Gakhov and Yu. I. Cherskii (1978).

$$15. \quad y(x) + A \int_a^b e^{\lambda|x-t|} y(t) dt = f(x).$$

This is a special case of equation 4.9.37 with $g(t) = A$.

1°. The function $y = y(x)$ obeys the following second-order linear nonhomogeneous ordinary differential equation with constant coefficients:

$$y''_{xx} + \lambda(2A - \lambda)y = f''_{xx}(x) - \lambda^2 f(x). \quad (1)$$

The boundary conditions for (1) have the form (see 4.9.37)

$$\begin{aligned} y'_x(a) + \lambda y(a) &= f'_x(a) + \lambda f(a), \\ y'_x(b) - \lambda y(b) &= f'_x(b) - \lambda f(b). \end{aligned} \quad (2)$$

Equation (1) under the boundary conditions (2) determines the solution of the original integral equation.

2°. For $\lambda(2A - \lambda) < 0$, the general solution of equation (1) is given by

$$\begin{aligned} y(x) &= C_1 \cosh(kx) + C_2 \sinh(kx) + f(x) - \frac{2A\lambda}{k} \int_a^x \sinh[k(x-t)] f(t) dt, \\ k &= \sqrt{\lambda(\lambda - 2A)}, \end{aligned} \quad (3)$$

where C_1 and C_2 are arbitrary constants.

For $\lambda(2A - \lambda) > 0$, the general solution of equation (1) is given by

$$\begin{aligned} y(x) &= C_1 \cos(kx) + C_2 \sin(kx) + f(x) - \frac{2A\lambda}{k} \int_a^x \sin[k(x-t)] f(t) dt, \\ k &= \sqrt{\lambda(2A - \lambda)}. \end{aligned} \quad (4)$$

For $\lambda = 2A$, the general solution of equation (1) is given by

$$y(x) = C_1 + C_2 x + f(x) - 4A^2 \int_a^x (x-t) f(t) dt. \quad (5)$$

The constants C_1 and C_2 in solutions (3)–(5) are determined by conditions (2).

3°. In the special case $a = 0$ and $\lambda(2A - \lambda) > 0$, the solution of the integral equation is given by formula (4) with

$$C_1 = \frac{A(kI_c - \lambda I_s)}{(\lambda - A) \sin \mu - k \cos \mu}, \quad C_2 = -\frac{\lambda}{k} \frac{A(kI_c - \lambda I_s)}{(\lambda - A) \sin \mu - k \cos \mu},$$

$$k = \sqrt{\lambda(2A - \lambda)}, \quad \mu = bk, \quad I_s = \int_0^b \sin[k(b-t)]f(t) dt, \quad I_c = \int_0^b \cos[k(b-t)]f(t) dt.$$

16. $y(x) + \int_a^b \left[\sum_{k=1}^n A_k \exp(\lambda_k |x-t|) \right] y(t) dt = f(x), \quad -\infty < a < b < \infty.$

1°. Let us remove the modulus in the k th summand of the integrand:

$$I_k(x) = \int_a^b \exp(\lambda_k |x-t|)y(t) dt = \int_a^x \exp[\lambda_k(x-t)]y(t) dt + \int_x^b \exp[\lambda_k(t-x)]y(t) dt. \quad (1)$$

Differentiating (1) with respect to x twice yields

$$I'_k = \lambda_k \int_a^x \exp[\lambda_k(x-t)]y(t) dt - \lambda_k \int_x^b \exp[\lambda_k(t-x)]y(t) dt,$$

$$I''_k = 2\lambda_k y(x) + \lambda_k^2 \int_a^x \exp[\lambda_k(x-t)]y(t) dt + \lambda_k^2 \int_x^b \exp[\lambda_k(t-x)]y(t) dt, \quad (2)$$

where the primes denote the derivatives with respect to x . By comparing formulas (1) and (2), we find the relation between I''_k and I_k :

$$I''_k = 2\lambda_k y(x) + \lambda_k^2 I_k, \quad I_k = I_k(x). \quad (3)$$

2°. With the aid of (1), the integral equation can be rewritten in the form

$$y(x) + \sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x twice and taking into account (3), we find that

$$y''_{xx}(x) + \sigma_n y(x) + \sum_{k=1}^n A_k \lambda_k^2 I_k = f''_{xx}(x), \quad \sigma_n = 2 \sum_{k=1}^n A_k \lambda_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5) yields

$$y''_{xx}(x) + (\sigma_n - \lambda_n^2)y(x) + \sum_{k=1}^{n-1} A_k (\lambda_k^2 - \lambda_n^2) I_k = f''_{xx}(x) - \lambda_n^2 f(x). \quad (6)$$

Differentiating (6) with respect to x twice and eliminating I_{n-1} from the resulting equation with the aid of (6), we obtain a similar equation whose left-hand side is a second-order linear differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} B_k I_k$. If we successively eliminate I_{n-2}, I_{n-3}, \dots , with the aid of double differentiation, then we finally arrive at a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients.

3°. The boundary conditions for $y(x)$ can be found by setting $x = a$ in the integral equation and all its derivatives. (Alternatively, these conditions can be found by setting $x = a$ and $x = b$ in the integral equation and all its derivatives obtained by means of double differentiation.)

4.2-2. Kernels Containing Power-Law and Exponential Functions.

$$17. \quad y(x) - \lambda \int_a^b (x-t)e^{\gamma t} y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $A = 0$, $B = 1$, and $h(t) = e^{\gamma t}$.

$$18. \quad y(x) - \lambda \int_a^b (x-t)e^{\gamma x} y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $A = 0$, $B = 1$, and $h(x) = e^{\gamma x}$.

$$19. \quad y(x) - \lambda \int_a^b (x-t)e^{\gamma x + \mu t} y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = xe^{\gamma x}$, $h_1(t) = e^{\mu t}$, $g_2(x) = e^{\gamma x}$, and $h_2(t) = -te^{\mu t}$.

$$20. \quad y(x) - \lambda \int_a^b [A + (Bx + Ct)e^{\gamma x}] y(t) dt = f(x).$$

This is a special case of equation 4.9.11 with $h(x) = e^{\gamma x}$.

$$21. \quad y(x) - \lambda \int_0^b (x^2 + t^2)e^{\gamma(x+t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.15 with $g(x) = x^2 e^{\gamma x}$ and $h(t) = e^{\gamma t}$.

$$22. \quad y(x) - \lambda \int_0^b (x^2 - t^2)e^{\gamma(x-t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = x^2 e^{\gamma x}$, $h_1(t) = e^{-\gamma t}$, $g_2(x) = e^{\gamma x}$, and $h_2(t) = -t^2 e^{-\gamma t}$.

$$23. \quad y(x) - \lambda \int_0^b (Ax^n + Bt^n)e^{\alpha x + \beta t} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.18 with $g_1(x) = x^n e^{\alpha x}$, $h_1(t) = Ae^{\beta t}$, $g_2(x) = e^{\alpha x}$, and $h_2(t) = Bt^n e^{\beta t}$.

$$24. \quad y(x) - \lambda \int_a^b \left[\sum_{k=1}^n A_k t^{\nu_k} e^{\alpha_k x + \beta_k t} \right] y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.20 with $g_k(x) = e^{\alpha_k x}$ and $h_k(t) = A_k t^{\nu_k} e^{\beta_k t}$.

$$25. \quad y(x) - \lambda \int_a^b \left[\sum_{k=1}^n A_k x^{\nu_k} e^{\alpha_k x + \beta_k t} \right] y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.20 with $g_k(x) = A_k x^{\nu_k} e^{\alpha_k x}$ and $h_k(t) = e^{\beta_k t}$.

$$26. \quad y(x) - \lambda \int_a^b (x-t)^n e^{\gamma(x-t)} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.20.

$$27. \quad y(x) - \lambda \int_a^b (x-t)^n e^{\alpha x + \beta t} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.20.

$$28. \quad y(x) - \lambda \int_a^b (Ax + Bt)^n e^{\alpha x + \beta t} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.20.

$$29. \quad y(x) + A \int_a^b t e^{\lambda|x-t|} y(t) dt = f(x).$$

This is a special case of equation 4.9.37 with $g(t) = At$. The solution of the integral equation can be written via the Bessel functions (or modified Bessel functions) of order $1/3$.

$$30. \quad y(x) + \int_0^\infty (a + b|x-t|) \exp(-|x-t|) y(t) dt = f(x).$$

Let the biquadratic polynomial $P(k) = k^4 + 2(a-b+1)k^2 + 2a+2b+1$ have no real roots and let $k = \alpha + i\beta$ be a root of the equation $P(k) = 0$ such that $\alpha > 0$ and $\beta > 0$. In this case, the solution has the form

$$\begin{aligned} y(x) = & f(x) + \rho \int_0^\infty \exp(-\beta|x-t|) \cos(\theta + \alpha|x-t|) f(t) dt \\ & + \frac{[\alpha + (\beta-1)^2]^2}{4\alpha^2\beta} \int_0^\infty \exp[-\beta(x+t)] \cos[\alpha(x-t)] f(t) dt \\ & + \frac{R}{4\alpha^2} \int_0^\infty \exp[-\beta(x+t)] \cos[\psi + \alpha(x+t)] f(t) dt, \end{aligned}$$

where the parameters ρ , θ , R , and ψ are determined from the system of algebraic equations obtained by separating real and imaginary parts in the relations

$$\rho e^{i\theta} = \frac{\mu}{\beta - i\alpha}, \quad R e^{i\psi} = \frac{(\beta - 1 - i\alpha)^4}{8\alpha^2(\beta - i\alpha)}.$$

⊙ Reference: F. D. Gakhov and Yu. I. Cherskii (1978).

4.3. Equations Whose Kernels Contain Hyperbolic Functions

4.3-1. Kernels Containing Hyperbolic Cosine.

$$1. \quad y(x) - \lambda \int_a^b \cosh(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cosh(\beta x)$ and $h(t) = 1$.

$$2. \quad y(x) - \lambda \int_a^b \cosh(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \cosh(\beta t)$.

$$3. \quad y(x) - \lambda \int_a^b \cosh[\beta(x-t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.13 with $g(x) = \cosh(\beta x)$ and $h(t) = \sinh(\beta t)$.

Solution:

$$y(x) = f(x) + \lambda [A_1 \cosh(\beta x) + A_2 \sinh(\beta x)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.13.

$$4. \quad y(x) - \lambda \int_a^b \cosh[\beta(x+t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.12 with $g(x) = \cosh(\beta x)$ and $h(t) = \sinh(\beta t)$.

Solution:

$$y(x) = f(x) + \lambda [A_1 \cosh(\beta x) + A_2 \sinh(\beta x)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.12.

$$5. \quad y(x) - \lambda \int_a^b \left\{ \sum_{k=1}^n A_k \cosh[\beta_k(x-t)] \right\} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.20.

$$6. \quad y(x) - \lambda \int_a^b \frac{\cosh(\beta x)}{\cosh(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cosh(\beta x)$ and $h(t) = \frac{1}{\cosh(\beta t)}$.

$$7. \quad y(x) - \lambda \int_a^b \frac{\cosh(\beta t)}{\cosh(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\cosh(\beta x)}$ and $h(t) = \cosh(\beta t)$.

$$8. \quad y(x) - \lambda \int_a^b \cosh^k(\beta x) \cosh^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cosh^k(\beta x)$ and $h(t) = \cosh^m(\mu t)$.

$$9. \quad y(x) - \lambda \int_a^b t^k \cosh^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cosh^m(\beta x)$ and $h(t) = t^k$.

$$10. \quad y(x) - \lambda \int_a^b x^k \cosh^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \cosh^m(\beta t)$.

$$11. \quad y(x) - \lambda \int_a^b [A + B(x-t) \cosh(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \cosh(\beta x)$.

$$12. \quad y(x) - \lambda \int_a^b [A + B(x-t) \cosh(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \cosh(\beta t)$.

$$13. \quad y(x) + \lambda \int_{-\infty}^{\infty} \frac{y(t) dt}{\cosh[b(x-t)]} = f(x).$$

Solution with $b > \pi|\lambda|$:

$$y(x) = f(x) - \frac{2\lambda b}{\sqrt{b^2 - \pi^2 \lambda^2}} \int_{-\infty}^{\infty} \frac{\sinh[2k(x-t)]}{\sinh[2b(x-t)]} f(t) dt, \quad k = \frac{b}{\pi} \arccos\left(\frac{\pi \lambda}{b}\right).$$

⊙ Reference: F. D. Gakhov and Yu. I. Cherskii (1978).

4.3-2. Kernels Containing Hyperbolic Sine.

$$14. \quad y(x) - \lambda \int_a^b \sinh(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sinh(\beta x)$ and $h(t) = 1$.

$$15. \quad y(x) - \lambda \int_a^b \sinh(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \sinh(\beta t)$.

$$16. \quad y(x) - \lambda \int_a^b \sinh[\beta(x-t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.16 with $g(x) = \sinh(\beta x)$ and $h(t) = \cosh(\beta t)$.

Solution:

$$y(x) = f(x) + \lambda [A_1 \sinh(\beta x) + A_2 \cosh(\beta x)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.16.

$$17. \quad y(x) - \lambda \int_a^b \sinh[\beta(x+t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.15 with $g(x) = \sinh(\beta x)$ and $h(t) = \cosh(\beta t)$.

Solution:

$$y(x) = f(x) + \lambda [A_1 \sinh(\beta x) + A_2 \cosh(\beta x)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.15.

$$18. \quad y(x) - \lambda \int_a^b \left\{ \sum_{k=1}^n A_k \sinh[\beta_k(x-t)] \right\} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.20.

$$19. \quad y(x) - \lambda \int_a^b \frac{\sinh(\beta x)}{\sinh(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sinh(\beta x)$ and $h(t) = \frac{1}{\sinh(\beta t)}$.

$$20. \quad y(x) - \lambda \int_a^b \frac{\sinh(\beta t)}{\sinh(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\sinh(\beta x)}$ and $h(t) = \sinh(\beta t)$.

$$21. \quad y(x) - \lambda \int_a^b \sinh^k(\beta x) \sinh^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sinh^k(\beta x)$ and $h(t) = \sinh^m(\mu t)$.

$$22. \quad y(x) - \lambda \int_a^b t^k \sinh^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sinh^m(\beta x)$ and $h(t) = t^k$.

$$23. \quad y(x) - \lambda \int_a^b x^k \sinh^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \sinh^m(\beta t)$.

$$24. \quad y(x) - \lambda \int_a^b [A + B(x - t) \sinh(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \sinh(\beta t)$.

$$25. \quad y(x) - \lambda \int_a^b [A + B(x - t) \sinh(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \sinh(\beta x)$.

$$26. \quad y(x) + A \int_a^b \sinh(\lambda|x - t|) y(t) dt = f(x).$$

This is a special case of equation 4.9.38 with $g(t) = A$.

1°. The function $y = y(x)$ obeys the following second-order linear nonhomogeneous ordinary differential equation with constant coefficients:

$$y''_{xx} + \lambda(2A - \lambda)y = f''_{xx}(x) - \lambda^2 f(x). \quad (1)$$

The boundary conditions for (1) have the form (see 4.9.38)

$$\begin{aligned} \sinh[\lambda(b - a)]\varphi'_x(b) - \lambda \cosh[\lambda(b - a)]\varphi(b) &= \lambda\varphi(a), \\ \sinh[\lambda(b - a)]\varphi'_x(a) + \lambda \cosh[\lambda(b - a)]\varphi(a) &= -\lambda\varphi(b), \end{aligned} \quad \varphi(x) = y(x) - f(x). \quad (2)$$

Equation (1) under the boundary conditions (2) determines the solution of the original integral equation.

2°. For $\lambda(2A - \lambda) = -k^2 < 0$, the general solution of equation (1) is given by

$$y(x) = C_1 \cosh(kx) + C_2 \sinh(kx) + f(x) - \frac{2A\lambda}{k} \int_a^x \sinh[k(x - t)]f(t) dt, \quad (3)$$

where C_1 and C_2 are arbitrary constants.

For $\lambda(2A - \lambda) = k^2 > 0$, the general solution of equation (1) is given by

$$y(x) = C_1 \cos(kx) + C_2 \sin(kx) + f(x) - \frac{2A\lambda}{k} \int_a^x \sin[k(x - t)]f(t) dt. \quad (4)$$

For $\lambda = 2A$, the general solution of equation (1) is given by

$$y(x) = C_1 + C_2 x + f(x) - 4A^2 \int_a^x (x - t)f(t) dt. \quad (5)$$

The constants C_1 and C_2 in solutions (3)–(5) are determined by conditions (2).

$$27. \quad y(x) + A \int_a^b t \sinh(\lambda|x-t|)y(t) dt = f(x).$$

This is a special case of equation 4.9.38 with $g(t) = At$. The solution of the integral equation can be written via the Bessel functions (or modified Bessel functions) of order $1/3$.

$$28. \quad y(x) + A \int_a^b \sinh^3(\lambda|x-t|)y(t) dt = f(x).$$

Using the formula $\sinh^3 \beta = \frac{1}{4} \sinh 3\beta - \frac{3}{4} \sinh \beta$, we arrive at an equation of the form 4.3.29 with $n = 2$:

$$y(x) + \int_a^b \left[\frac{1}{4} A \sinh(3\lambda|x-t|) - \frac{3}{4} A \sinh(\lambda|x-t|) \right] y(t) dt = f(x).$$

$$29. \quad y(x) + \int_a^b \left[\sum_{k=1}^n A_k \sinh(\lambda_k|x-t|) \right] y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

1°. Let us remove the modulus in the k th summand of the integrand:

$$I_k(x) = \int_a^b \sinh(\lambda_k|x-t|)y(t) dt = \int_a^x \sinh[\lambda_k(x-t)]y(t) dt + \int_x^b \sinh[\lambda_k(t-x)]y(t) dt. \quad (1)$$

Differentiating (1) with respect to x twice yields

$$I'_k = \lambda_k \int_a^x \cosh[\lambda_k(x-t)]y(t) dt - \lambda_k \int_x^b \cosh[\lambda_k(t-x)]y(t) dt, \quad (2)$$

$$I''_k = 2\lambda_k y(x) + \lambda_k^2 \int_a^x \sinh[\lambda_k(x-t)]y(t) dt + \lambda_k^2 \int_x^b \sinh[\lambda_k(t-x)]y(t) dt,$$

where the primes denote the derivatives with respect to x . By comparing formulas (1) and (2), we find the relation between I''_k and I_k :

$$I''_k = 2\lambda_k y(x) + \lambda_k^2 I_k, \quad I_k = I_k(x). \quad (3)$$

2°. With the aid of (1), the integral equation can be rewritten in the form

$$y(x) + \sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x twice and taking into account (3), we find that

$$y''_{xx}(x) + \sigma_n y(x) + \sum_{k=1}^n A_k \lambda_k^2 I_k = f''_{xx}(x), \quad \sigma_n = 2 \sum_{k=1}^n A_k \lambda_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5) yields

$$y''_{xx}(x) + (\sigma_n - \lambda_n^2)y(x) + \sum_{k=1}^{n-1} A_k (\lambda_k^2 - \lambda_n^2) I_k = f''_{xx}(x) - \lambda_n^2 f(x). \quad (6)$$

Differentiating (6) with respect to x twice and eliminating I_{n-1} from the resulting equation with the aid of (6), we obtain a similar equation whose left-hand side is a second-order linear differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} B_k I_k$. If we successively eliminate I_{n-2}, I_{n-3}, \dots , with the aid of double differentiation, then we finally arrive at a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients.

3°. The boundary conditions for $y(x)$ can be found by setting $x = a$ in the integral equation and its derivatives. (Alternatively, these conditions can be found by setting $x = a$ and $x = b$ in the integral equation and all its derivatives obtained by means of double differentiation.)

4.3-3. Kernels Containing Hyperbolic Tangent.

$$30. \quad y(x) - \lambda \int_a^b \tanh(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tanh(\beta x)$ and $h(t) = 1$.

$$31. \quad y(x) - \lambda \int_a^b \tanh(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \tanh(\beta t)$.

$$32. \quad y(x) - \lambda \int_a^b [A \tanh(\beta x) + B \tanh(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.4 with $g(x) = \tanh(\beta x)$.

$$33. \quad y(x) - \lambda \int_a^b \frac{\tanh(\beta x)}{\tanh(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tanh(\beta x)$ and $h(t) = \frac{1}{\tanh(\beta t)}$.

$$34. \quad y(x) - \lambda \int_a^b \frac{\tanh(\beta t)}{\tanh(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\tanh(\beta x)}$ and $h(t) = \tanh(\beta t)$.

$$35. \quad y(x) - \lambda \int_a^b \tanh^k(\beta x) \tanh^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tanh^k(\beta x)$ and $h(t) = \tanh^m(\mu t)$.

$$36. \quad y(x) - \lambda \int_a^b t^k \tanh^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tanh^m(\beta x)$ and $h(t) = t^k$.

$$37. \quad y(x) - \lambda \int_a^b x^k \tanh^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \tanh^m(\beta t)$.

$$38. \quad y(x) - \lambda \int_a^b [A + B(x - t) \tanh(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \tanh(\beta t)$.

$$39. \quad y(x) - \lambda \int_a^b [A + B(x - t) \tanh(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \tanh(\beta x)$.

4.3-4. Kernels Containing Hyperbolic Cotangent.

$$40. \quad y(x) - \lambda \int_a^b \coth(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \coth(\beta x)$ and $h(t) = 1$.

$$41. \quad y(x) - \lambda \int_a^b \coth(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \coth(\beta t)$.

$$42. \quad y(x) - \lambda \int_a^b [A \coth(\beta x) + B \coth(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.4 with $g(x) = \coth(\beta x)$.

$$43. \quad y(x) - \lambda \int_a^b \frac{\coth(\beta x)}{\coth(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \coth(\beta x)$ and $h(t) = \frac{1}{\coth(\beta t)}$.

$$44. \quad y(x) - \lambda \int_a^b \frac{\coth(\beta t)}{\coth(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\coth(\beta x)}$ and $h(t) = \coth(\beta t)$.

$$45. \quad y(x) - \lambda \int_a^b \coth^k(\beta x) \coth^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \coth^k(\beta x)$ and $h(t) = \coth^m(\mu t)$.

$$46. \quad y(x) - \lambda \int_a^b t^k \coth^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \coth^m(\beta x)$ and $h(t) = t^k$.

$$47. \quad y(x) - \lambda \int_a^b x^k \coth^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \coth^m(\beta t)$.

$$48. \quad y(x) - \lambda \int_a^b [A + B(x-t) \coth(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \coth(\beta t)$.

$$49. \quad y(x) - \lambda \int_a^b [A + B(x-t) \coth(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \coth(\beta x)$.

4.3-5. Kernels Containing Combination of Hyperbolic Functions.

$$50. \quad y(x) - \lambda \int_a^b \cosh^k(\beta x) \sinh^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cosh^k(\beta x)$ and $h(t) = \sinh^m(\mu t)$.

$$51. \quad y(x) - \lambda \int_a^b [A \sinh(\alpha x) \cosh(\beta t) + B \sinh(\gamma x) \cosh(\delta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = \sinh(\alpha x)$, $h_1(t) = A \cosh(\beta t)$, $g_2(x) = \sinh(\gamma x)$, and $h_2(t) = B \cosh(\delta t)$.

$$52. \quad y(x) - \lambda \int_a^b \tanh^k(\gamma x) \coth^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tanh^k(\gamma x)$ and $h(t) = \coth^m(\mu t)$.

$$53. \quad y(x) - \lambda \int_a^b [A \tanh(\alpha x) \coth(\beta t) + B \tanh(\gamma x) \coth(\delta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = \tanh(\alpha x)$, $h_1(t) = A \coth(\beta t)$, $g_2(x) = \tanh(\gamma x)$, and $h_2(t) = B \coth(\delta t)$.

4.4. Equations Whose Kernels Contain Logarithmic Functions

4.4-1. Kernels Containing Logarithmic Functions.

$$1. \quad y(x) - \lambda \int_a^b \ln(\gamma x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln(\gamma x)$ and $h(t) = 1$.

$$2. \quad y(x) - \lambda \int_a^b \ln(\gamma t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \ln(\gamma t)$.

$$3. \quad y(x) - \lambda \int_a^b (\ln x - \ln t) y(t) dt = f(x).$$

This is a special case of equation 4.9.3 with $g(x) = \ln x$.

$$4. \quad y(x) - \lambda \int_a^b \frac{\ln(\gamma x)}{\ln(\gamma t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln(\gamma x)$ and $h(t) = \frac{1}{\ln(\gamma t)}$.

$$5. \quad y(x) - \lambda \int_a^b \frac{\ln(\gamma t)}{\ln(\gamma x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\ln(\gamma x)}$ and $h(t) = \ln(\gamma t)$.

$$6. \quad y(x) - \lambda \int_a^b \ln^k(\gamma x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^k(\gamma x)$ and $h(t) = \ln^m(\mu t)$.

4.4-2. Kernels Containing Power-Law and Logarithmic Functions.

$$7. \quad y(x) - \lambda \int_a^b t^k \ln^m(\gamma x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\gamma x)$ and $h(t) = t^k$.

$$8. \quad y(x) - \lambda \int_a^b x^k \ln^m(\gamma t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \ln^m(\gamma t)$.

$$9. \quad y(x) - \lambda \int_a^b [A + B(x-t) \ln(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \ln(\gamma t)$.

$$10. \quad y(x) - \lambda \int_a^b [A + B(x-t) \ln(\gamma x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \ln(\gamma x)$.

$$11. \quad y(x) - \lambda \int_a^b [A + (Bx + Ct) \ln(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.9 with $h(t) = \ln(\gamma t)$.

$$12. \quad y(x) - \lambda \int_a^b [A + (Bx + Ct) \ln(\gamma x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.11 with $h(x) = \ln(\gamma x)$.

$$13. \quad y(x) - \lambda \int_a^b [At^n \ln^m(\beta x) + Bx^k \ln^l(\gamma t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = \ln^m(\beta x)$, $h_1(t) = At^n$, $g_2(x) = x^k$, and $h_2(t) = B \ln^l(\gamma t)$.

4.5. Equations Whose Kernels Contain Trigonometric Functions

4.5-1. Kernels Containing Cosine.

$$1. \quad y(x) - \lambda \int_a^b \cos(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos(\beta x)$ and $h(t) = 1$.

$$2. \quad y(x) - \lambda \int_a^b \cos(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \cos(\beta t)$.

$$3. \quad y(x) - \lambda \int_a^b \cos[\beta(x-t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.12 with $g(x) = \cos(\beta x)$ and $h(t) = \sin(\beta t)$.

Solution:

$$y(x) = f(x) + \lambda [A_1 \cos(\beta x) + A_2 \sin(\beta x)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.12.

$$4. \quad y(x) - \lambda \int_a^b \cos[\beta(x+t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.13 with $g(x) = \cos(\beta x)$ and $h(t) = \sin(\beta t)$.

Solution:

$$y(x) = f(x) + \lambda [A_1 \cos(\beta x) + A_2 \sin(\beta x)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.13.

$$5. \quad y(x) - \lambda \int_0^\infty \cos(xt) y(t) dt = 0.$$

Characteristic values: $\lambda = \pm \sqrt{2/\pi}$. For the characteristic values, the integral equation has infinitely many linearly independent eigenfunctions.

Eigenfunctions for $\lambda = +\sqrt{2/\pi}$ have the form

$$y_+(x) = f(x) + \sqrt{\frac{2}{\pi}} \int_0^\infty f(t) \cos(xt) dt, \quad (1)$$

where $f = f(x)$ is any continuous function absolutely integrable on the interval $[0, \infty)$.

Eigenfunctions for $\lambda = -\sqrt{2/\pi}$ have the form

$$y_-(x) = f(x) - \sqrt{\frac{2}{\pi}} \int_0^\infty f(t) \cos(xt) dt, \quad (2)$$

where $f = f(x)$ is any continuous function absolutely integrable on the interval $[0, \infty)$.

In particular, from (1) and (2) with $f(x) = e^{-ax}$ we obtain

$$\begin{aligned} y_+(x) &= e^{-ax} + \sqrt{\frac{2}{\pi}} \frac{a}{a^2 + x^2} & \text{for } \lambda = +\sqrt{\frac{2}{\pi}}, \\ y_-(x) &= e^{-ax} - \sqrt{\frac{2}{\pi}} \frac{a}{a^2 + x^2} & \text{for } \lambda = -\sqrt{\frac{2}{\pi}}, \end{aligned}$$

where a is any positive number.

⊙ Reference: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971).

$$6. \quad y(x) - \lambda \int_0^\infty \cos(xt) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^\infty \cos(xt) f(t) dt,$$

where $\lambda \neq \pm \sqrt{2/\pi}$.

⊙ Reference: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971).

$$7. \quad y(x) - \lambda \int_a^b \left\{ \sum_{k=1}^n A_k \cos[\beta_k(x-t)] \right\} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This equation can be reduced to a special case of equation 4.9.20; the formula $\cos[\beta(x-t)] = \cos(\beta x) \cos(\beta t) + \sin(\beta x) \sin(\beta t)$ must be used.

$$8. \quad y(x) - \lambda \int_a^b \frac{\cos(\beta x)}{\cos(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos(\beta x)$ and $h(t) = \frac{1}{\cos(\beta t)}$.

$$9. \quad y(x) - \lambda \int_a^b \frac{\cos(\beta t)}{\cos(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\cos(\beta x)}$ and $h(t) = \cos(\beta t)$.

$$10. \quad y(x) - \lambda \int_a^b \cos^k(\beta x) \cos^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos^k(\beta x)$ and $h(t) = \cos^m(\mu t)$.

$$11. \quad y(x) - \lambda \int_a^b t^k \cos^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos^m(\beta x)$ and $h(t) = t^k$.

$$12. \quad y(x) - \lambda \int_a^b x^k \cos^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \cos^m(\beta t)$.

$$13. \quad y(x) - \lambda \int_a^b [A + B(x-t) \cos(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \cos(\beta x)$.

$$14. \quad y(x) - \lambda \int_a^b [A + B(x-t) \cos(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \cos(\beta t)$.

4.5-2. Kernels Containing Sine.

$$15. \quad y(x) - \lambda \int_a^b \sin(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin(\beta x)$ and $h(t) = 1$.

$$16. \quad y(x) - \lambda \int_a^b \sin(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \sin(\beta t)$.

$$17. \quad y(x) - \lambda \int_a^b \sin[\beta(x-t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.16 with $g(x) = \sin(\beta x)$ and $h(t) = \cos(\beta t)$.

Solution:

$$y(x) = f(x) + \lambda [A_1 \sin(\beta x) + A_2 \cos(\beta x)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.16.

$$18. \quad y(x) - \lambda \int_a^b \sin[\beta(x+t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.15 with $g(x) = \sin(\beta x)$ and $h(t) = \cos(\beta t)$.

Solution:

$$y(x) = f(x) + \lambda [A_1 \sin(\beta x) + A_2 \cos(\beta x)],$$

where A_1 and A_2 are the constants determined by the formulas presented in 4.9.15.

Example. Let us consider the case of $a = 0$, $b = \pi$, $\beta = 1$ in detail.

1°. Solution for $\lambda \neq \pm 2/\pi$:

$$y(x) = f(x) + \lambda A \sin x + \lambda B \cos x, \quad (1)$$

where

$$A = \frac{f_1 + \frac{1}{2}\pi\lambda f_2}{1 - \frac{1}{4}\pi^2\lambda^2}, \quad B = \frac{\frac{1}{2}\pi\lambda f_1 + f_2}{1 - \frac{1}{4}\pi^2\lambda^2}, \quad f_1 = \int_0^\pi f(t) \cos t dt, \quad f_2 = \int_0^\pi f(t) \sin t dt. \quad (2)$$

2°. Characteristic values and normed eigenfunctions of the homogeneous equation for $f(x) \equiv 0$ are given by the formulas

$$\lambda_1 = -\frac{2}{\pi}, \quad y_1(x) = \frac{1}{\sqrt{\pi}}(\sin x - \cos x);$$

$$\lambda_2 = \frac{2}{\pi}, \quad y_2(x) = \frac{1}{\sqrt{\pi}}(\sin x + \cos x).$$

3°. If $\lambda = -2/\pi$ and $f_1 = f_2$ (values of f_1 and f_2 can be found using formulas of Item 1°). In this case the solution can be obtained with the help of formula (1) in which $B = f_1 - A$ where A is an arbitrary constant.

If $\lambda = 2/\pi$ and $f_1 = -f_2$ then the solution can be found using formula (1) in which $B = A - f_1$ where A is an arbitrary constant.

4°. If $\lambda = -2/\pi$ and $f_1 \neq f_2$ or $\lambda = 2/\pi$ and $f_1 \neq -f_2$, then the equation under consideration has no solutions.

$$19. \quad y(x) - \lambda \int_0^\infty \sin(xt)y(t) dt = 0.$$

Characteristic values: $\lambda = \pm\sqrt{2/\pi}$. For the characteristic values, the integral equation has infinitely many linearly independent eigenfunctions.

Eigenfunctions for $\lambda = +\sqrt{2/\pi}$ have the form

$$y_+(x) = f(x) + \sqrt{\frac{2}{\pi}} \int_0^\infty f(t) \sin(xt) dt,$$

where $f = f(x)$ is any continuous function absolutely integrable on the interval $[0, \infty)$.

Eigenfunctions for $\lambda = -\sqrt{2/\pi}$ have the form

$$y_-(x) = f(x) - \sqrt{\frac{2}{\pi}} \int_0^\infty f(t) \sin(xt) dt,$$

where $f = f(x)$ is any continuous function absolutely integrable on the interval $[0, \infty)$.

$$20. \quad y(x) - \lambda \int_0^{\infty} \sin(xt)y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt)f(t) dt,$$

where $\lambda \neq \pm \sqrt{2/\pi}$.

⊙ References: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971), F. D. Gakhov and Yu. I. Cherskii (1978).

$$21. \quad y(x) - \lambda \int_a^b \left\{ \sum_{k=1}^n A_k \sin[\beta_k(x-t)] \right\} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This equation can be reduced to a special case of equation 4.9.20; the formula $\sin[\beta(x-t)] = \sin(\beta x) \cos(\beta t) - \sin(\beta t) \cos(\beta x)$ must be used.

$$22. \quad y(x) - \lambda \int_a^b \frac{\sin(\beta x)}{\sin(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin(\beta x)$ and $h(t) = \frac{1}{\sin(\beta t)}$.

$$23. \quad y(x) - \lambda \int_a^b \frac{\sin(\beta t)}{\sin(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\sin(\beta x)}$ and $h(t) = \sin(\beta t)$.

$$24. \quad y(x) - \lambda \int_a^b \sin^k(\beta x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin^k(\beta x)$ and $h(t) = \sin^m(\mu t)$.

$$25. \quad y(x) - \lambda \int_a^b t^k \sin^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin^m(\beta x)$ and $h(t) = t^k$.

$$26. \quad y(x) - \lambda \int_a^b x^k \sin^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \sin^m(\beta t)$.

$$27. \quad y(x) - \lambda \int_a^b [A + B(x-t) \sin(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \sin(\beta t)$.

$$28. \quad y(x) - \lambda \int_a^b [A + B(x-t) \sin(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \sin(\beta x)$.

$$29. \quad y(x) + A \int_a^b \sin(\lambda|x-t|)y(t) dt = f(x).$$

This is a special case of equation 4.9.39 with $g(t) = A$.

1°. The function $y = y(x)$ obeys the following second-order linear nonhomogeneous ordinary differential equation with constant coefficients:

$$y''_{xx} + \lambda(2A + \lambda)y = f''_{xx}(x) + \lambda^2 f(x). \quad (1)$$

The boundary conditions for (1) have the form (see 4.9.39)

$$\begin{aligned} \sin[\lambda(b-a)]\varphi'_x(b) - \lambda \cos[\lambda(b-a)]\varphi(b) &= \lambda\varphi(a), \\ \sin[\lambda(b-a)]\varphi'_x(a) + \lambda \cos[\lambda(b-a)]\varphi(a) &= -\lambda\varphi(b), \end{aligned} \quad \varphi(x) = y(x) - f(x). \quad (2)$$

Equation (1) under the boundary conditions (2) determines the solution of the original integral equation.

2°. For $\lambda(2A + \lambda) = -k^2 < 0$, the general solution of equation (1) is given by

$$y(x) = C_1 \cosh(kx) + C_2 \sinh(kx) + f(x) - \frac{2A\lambda}{k} \int_a^x \sinh[k(x-t)] f(t) dt, \quad (3)$$

where C_1 and C_2 are arbitrary constants.

For $\lambda(2A + \lambda) = k^2 > 0$, the general solution of equation (1) is given by

$$y(x) = C_1 \cos(kx) + C_2 \sin(kx) + f(x) - \frac{2A\lambda}{k} \int_a^x \sin[k(x-t)] f(t) dt. \quad (4)$$

For $\lambda = 2A$, the general solution of equation (1) is given by

$$y(x) = C_1 + C_2 x + f(x) + 4A^2 \int_a^x (x-t)f(t) dt. \quad (5)$$

The constants C_1 and C_2 in solutions (3)–(5) are determined by conditions (2).

$$30. \quad y(x) + A \int_a^b t \sin(\lambda|x-t|)y(t) dt = f(x).$$

This is a special case of equation 4.9.39 with $g(t) = At$. The solution of the integral equation can be written via the Bessel functions (or modified Bessel functions) of order $1/3$.

$$31. \quad y(x) + A \int_a^b \sin^3(\lambda|x-t|)y(t) dt = f(x).$$

Using the formula $\sin^3 \beta = -\frac{1}{4} \sin 3\beta + \frac{3}{4} \sin \beta$, we arrive at an equation of the form 4.5.32 with $n = 2$:

$$y(x) + \int_a^b \left[-\frac{1}{4} A \sin(3\lambda|x-t|) + \frac{3}{4} A \sin(\lambda|x-t|) \right] y(t) dt = f(x).$$

$$32. \quad y(x) + \int_a^b \left[\sum_{k=1}^n A_k \sin(\lambda_k |x-t|) \right] y(t) dt = f(x), \quad -\infty < a < b < \infty.$$

1°. Let us remove the modulus in the k th summand of the integrand:

$$I_k(x) = \int_a^b \sin(\lambda_k |x-t|) y(t) dt = \int_a^x \sin[\lambda_k(x-t)] y(t) dt + \int_x^b \sin[\lambda_k(t-x)] y(t) dt. \quad (1)$$

Differentiating (1) with respect to x twice yields

$$\begin{aligned} I_k' &= \lambda_k \int_a^x \cos[\lambda_k(x-t)] y(t) dt - \lambda_k \int_x^b \cos[\lambda_k(t-x)] y(t) dt, \\ I_k'' &= 2\lambda_k y(x) - \lambda_k^2 \int_a^x \sin[\lambda_k(x-t)] y(t) dt - \lambda_k^2 \int_x^b \sin[\lambda_k(t-x)] y(t) dt, \end{aligned} \quad (2)$$

where the primes denote the derivatives with respect to x . By comparing formulas (1) and (2), we find the relation between I_k'' and I_k :

$$I_k'' = 2\lambda_k y(x) - \lambda_k^2 I_k, \quad I_k = I_k(x). \quad (3)$$

2°. With the aid of (1), the integral equation can be rewritten in the form

$$y(x) + \sum_{k=1}^n A_k I_k = f(x). \quad (4)$$

Differentiating (4) with respect to x twice and taking into account (3), we find that

$$y''_{xx}(x) + \sigma_n y(x) - \sum_{k=1}^n A_k \lambda_k^2 I_k = f''_{xx}(x), \quad \sigma_n = 2 \sum_{k=1}^n A_k \lambda_k. \quad (5)$$

Eliminating the integral I_n from (4) and (5) yields

$$y''_{xx}(x) + (\sigma_n + \lambda_n^2) y(x) + \sum_{k=1}^{n-1} A_k (\lambda_n^2 - \lambda_k^2) I_k = f''_{xx}(x) + \lambda_n^2 f(x). \quad (6)$$

Differentiating (6) with respect to x twice and eliminating I_{n-1} from the resulting equation with the aid of (6), we obtain a similar equation whose left-hand side is a second-order linear differential operator (acting on y) with constant coefficients plus the sum $\sum_{k=1}^{n-2} B_k I_k$. If we successively eliminate I_{n-2}, I_{n-3}, \dots , with the aid of double differentiation, then we finally arrive at a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients.

3°. The boundary conditions for $y(x)$ can be found by setting $x = a$ in the integral equation and all its derivatives. (Alternatively, these conditions can be found by setting $x = a$ and $x = b$ in the integral equation and all its derivatives obtained by means of double differentiation.)

$$33. \quad y(x) - \lambda \int_{-\infty}^{\infty} \frac{\sin(x-t)}{x-t} y(t) dt = f(x).$$

Solution:

$$y(x) = f(x) + \frac{\lambda}{\sqrt{2\pi - \pi\lambda}} \int_{-\infty}^{\infty} \frac{\sin(x-t)}{x-t} f(t) dt, \quad \lambda \neq \sqrt{\frac{2}{\pi}}.$$

4.5-3. Kernels Containing Tangent.

$$34. \quad y(x) - \lambda \int_a^b \tan(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tan(\beta x)$ and $h(t) = 1$.

$$35. \quad y(x) - \lambda \int_a^b \tan(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \tan(\beta t)$.

$$36. \quad y(x) - \lambda \int_a^b [A \tan(\beta x) + B \tan(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.4 with $g(x) = \tan(\beta x)$.

$$37. \quad y(x) - \lambda \int_a^b \frac{\tan(\beta x)}{\tan(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tan(\beta x)$ and $h(t) = \frac{1}{\tan(\beta t)}$.

$$38. \quad y(x) - \lambda \int_a^b \frac{\tan(\beta t)}{\tan(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\tan(\beta x)}$ and $h(t) = \tan(\beta t)$.

$$39. \quad y(x) - \lambda \int_a^b \tan^k(\beta x) \tan^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tan^k(\beta x)$ and $h(t) = \tan^m(\mu t)$.

$$40. \quad y(x) - \lambda \int_a^b t^k \tan^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tan^m(\beta x)$ and $h(t) = t^k$.

$$41. \quad y(x) - \lambda \int_a^b x^k \tan^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \tan^m(\beta t)$.

$$42. \quad y(x) - \lambda \int_a^b [A + B(x - t) \tan(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \tan(\beta t)$.

$$43. \quad y(x) - \lambda \int_a^b [A + B(x - t) \tan(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \tan(\beta x)$.

4.5-4. Kernels Containing Cotangent.

$$44. \quad y(x) - \lambda \int_a^b \cot(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cot(\beta x)$ and $h(t) = 1$.

$$45. \quad y(x) - \lambda \int_a^b \cot(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \cot(\beta t)$.

$$46. \quad y(x) - \lambda \int_a^b [A \cot(\beta x) + B \cot(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.4 with $g(x) = \cot(\beta x)$.

$$47. \quad y(x) - \lambda \int_a^b \frac{\cot(\beta x)}{\cot(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cot(\beta x)$ and $h(t) = \frac{1}{\cot(\beta t)}$.

$$48. \quad y(x) - \lambda \int_a^b \frac{\cot(\beta t)}{\cot(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\cot(\beta x)}$ and $h(t) = \cot(\beta t)$.

$$49. \quad y(x) - \lambda \int_a^b \cot^k(\beta x) \cot^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cot^k(\beta x)$ and $h(t) = \cot^m(\mu t)$.

$$50. \quad y(x) - \lambda \int_a^b t^k \cot^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cot^m(\beta x)$ and $h(t) = t^k$.

$$51. \quad y(x) - \lambda \int_a^b x^k \cot^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \cot^m(\beta t)$.

$$52. \quad y(x) - \lambda \int_a^b [A + B(x-t) \cot(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \cot(\beta t)$.

$$53. \quad y(x) - \lambda \int_a^b [A + B(x-t) \cot(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \cot(\beta x)$.

4.5-5. Kernels Containing Combinations of Trigonometric Functions.

$$54. \quad y(x) - \lambda \int_a^b \cos^k(\beta x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos^k(\beta x)$ and $h(t) = \sin^m(\mu t)$.

$$55. \quad y(x) - \lambda \int_a^b [A \sin(\alpha x) \cos(\beta t) + B \sin(\gamma x) \cos(\delta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = \sin(\alpha x)$, $h_1(t) = A \cos(\beta t)$, $g_2(x) = \sin(\gamma x)$, and $h_2(t) = B \cos(\delta t)$.

$$56. \quad y(x) - \lambda \int_a^b \tan^k(\gamma x) \cot^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tan^k(\gamma x)$ and $h(t) = \cot^m(\mu t)$.

$$57. \quad y(x) - \lambda \int_a^b [A \tan(\alpha x) \cot(\beta t) + B \tan(\gamma x) \cot(\delta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = \tan(\alpha x)$, $h_1(t) = A \cot(\beta t)$, $g_2(x) = \tan(\gamma x)$, and $h_2(t) = B \cot(\delta t)$.

4.5-6. Singular Equation.

$$58. \quad Ay(x) - \frac{B}{2\pi} \int_0^{2\pi} \cot\left(\frac{t-x}{2}\right) y(t) dt = f(x), \quad 0 \leq x \leq 2\pi.$$

Here the integral is understood in the sense of the Cauchy principal value. Without loss of generality we may assume that $A^2 + B^2 = 1$.

Solution:

$$y(x) = Af(x) + \frac{B}{2\pi} \int_0^{2\pi} \cot\left(\frac{t-x}{2}\right) f(t) dt + \frac{B^2}{2\pi A} \int_0^{2\pi} f(t) dt.$$

⊙ Reference: I. K. Lifanov (1996).

4.6. Equations Whose Kernels Contain Inverse Trigonometric Functions

4.6-1. Kernels Containing Arccosine.

$$1. \quad y(x) - \lambda \int_a^b \arccos(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arccos(\beta x)$ and $h(t) = 1$.

$$2. \quad y(x) - \lambda \int_a^b \arccos(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \arccos(\beta t)$.

$$3. \quad y(x) - \lambda \int_a^b \frac{\arccos(\beta x)}{\arccos(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arccos(\beta x)$ and $h(t) = \frac{1}{\arccos(\beta t)}$.

$$4. \quad y(x) - \lambda \int_a^b \frac{\arccos(\beta t)}{\arccos(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\arccos(\beta x)}$ and $h(t) = \arccos(\beta t)$.

$$5. \quad y(x) - \lambda \int_a^b \arccos^k(\beta x) \arccos^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arccos^k(\beta x)$ and $h(t) = \arccos^m(\mu t)$.

$$6. \quad y(x) - \lambda \int_a^b t^k \arccos^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arccos^m(\beta x)$ and $h(t) = t^k$.

$$7. \quad y(x) - \lambda \int_a^b x^k \arccos^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \arccos^m(\beta t)$.

$$8. \quad y(x) - \lambda \int_a^b [A + B(x - t) \arccos(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \arccos(\beta x)$.

$$9. \quad y(x) - \lambda \int_a^b [A + B(x - t) \arccos(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \arccos(\beta t)$.

4.6-2. Kernels Containing Arcsine.

$$10. \quad y(x) - \lambda \int_a^b \arcsin(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arcsin(\beta x)$ and $h(t) = 1$.

$$11. \quad y(x) - \lambda \int_a^b \arcsin(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \arcsin(\beta t)$.

$$12. \quad y(x) - \lambda \int_a^b \frac{\arcsin(\beta x)}{\arcsin(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arcsin(\beta x)$ and $h(t) = \frac{1}{\arcsin(\beta t)}$.

$$13. \quad y(x) - \lambda \int_a^b \frac{\arcsin(\beta t)}{\arcsin(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\arcsin(\beta x)}$ and $h(t) = \arcsin(\beta t)$.

$$14. \quad y(x) - \lambda \int_a^b \arcsin^k(\beta x) \arcsin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arcsin^k(\beta x)$ and $h(t) = \arcsin^m(\mu t)$.

$$15. \quad y(x) - \lambda \int_a^b t^k \arcsin^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arcsin^m(\beta x)$ and $h(t) = t^k$.

$$16. \quad y(x) - \lambda \int_a^b x^k \arcsin^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \arcsin^m(\beta t)$.

$$17. \quad y(x) - \lambda \int_a^b [A + B(x - t) \arcsin(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \arcsin(\beta t)$.

$$18. \quad y(x) - \lambda \int_a^b [A + B(x - t) \arcsin(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \arcsin(\beta x)$.

4.6-3. Kernels Containing Arctangent.

$$19. \quad y(x) - \lambda \int_a^b \arctan(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arctan(\beta x)$ and $h(t) = 1$.

$$20. \quad y(x) - \lambda \int_a^b \arctan(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \arctan(\beta t)$.

$$21. \quad y(x) - \lambda \int_a^b [A \arctan(\beta x) + B \arctan(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.4 with $g(x) = \arctan(\beta x)$.

$$22. \quad y(x) - \lambda \int_a^b \frac{\arctan(\beta x)}{\arctan(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arctan(\beta x)$ and $h(t) = \frac{1}{\arctan(\beta t)}$.

$$23. \quad y(x) - \lambda \int_a^b \frac{\arctan(\beta t)}{\arctan(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\arctan(\beta x)}$ and $h(t) = \arctan(\beta t)$.

$$24. \quad y(x) - \lambda \int_a^b \arctan^k(\beta x) \arctan^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arctan^k(\beta x)$ and $h(t) = \arctan^m(\mu t)$.

$$25. \quad y(x) - \lambda \int_a^b t^k \arctan^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \arctan^m(\beta x)$ and $h(t) = t^k$.

$$26. \quad y(x) - \lambda \int_a^b x^k \arctan^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \arctan^m(\beta t)$.

$$27. \quad y(x) - \lambda \int_a^b [A + B(x-t) \arctan(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \arctan(\beta t)$.

$$28. \quad y(x) - \lambda \int_a^b [A + B(x-t) \arctan(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \arctan(\beta x)$.

4.6-4. Kernels Containing Arccotangent.

$$29. \quad y(x) - \lambda \int_a^b \operatorname{arccot}(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \operatorname{arccot}(\beta x)$ and $h(t) = 1$.

$$30. \quad y(x) - \lambda \int_a^b \operatorname{arccot}(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = \operatorname{arccot}(\beta t)$.

$$31. \quad y(x) - \lambda \int_a^b [A \operatorname{arccot}(\beta x) + B \operatorname{arccot}(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.4 with $g(x) = \operatorname{arccot}(\beta x)$.

$$32. \quad y(x) - \lambda \int_a^b \frac{\operatorname{arccot}(\beta x)}{\operatorname{arccot}(\beta t)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \operatorname{arccot}(\beta x)$ and $h(t) = \frac{1}{\operatorname{arccot}(\beta t)}$.

$$33. \quad y(x) - \lambda \int_a^b \frac{\operatorname{arccot}(\beta t)}{\operatorname{arccot}(\beta x)} y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \frac{1}{\operatorname{arccot}(\beta x)}$ and $h(t) = \operatorname{arccot}(\beta t)$.

$$34. \quad y(x) - \lambda \int_a^b \operatorname{arccot}^k(\beta x) \operatorname{arccot}^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \operatorname{arccot}^k(\beta x)$ and $h(t) = \operatorname{arccot}^m(\mu t)$.

$$35. \quad y(x) - \lambda \int_a^b t^k \operatorname{arccot}^m(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \operatorname{arccot}^m(\beta x)$ and $h(t) = t^k$.

$$36. \quad y(x) - \lambda \int_a^b x^k \operatorname{arccot}^m(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = x^k$ and $h(t) = \operatorname{arccot}^m(\beta t)$.

$$37. \quad y(x) - \lambda \int_a^b [A + B(x - t) \operatorname{arccot}(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = \operatorname{arccot}(\beta t)$.

$$38. \quad y(x) - \lambda \int_a^b [A + B(x - t) \operatorname{arccot}(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = \operatorname{arccot}(\beta x)$.

4.7. Equations Whose Kernels Contain Combinations of Elementary Functions

4.7-1. Kernels Containing Exponential and Hyperbolic Functions.

$$1. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} \cosh[\beta(x-t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\mu x} \cosh(\beta x)$, $h_1(t) = e^{-\mu t} \cosh(\beta t)$, $g_2(x) = e^{\mu x} \sinh(\beta x)$, and $h_2(t) = -e^{-\mu t} \sinh(\beta t)$.

$$2. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} \sinh[\beta(x-t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\mu x} \sinh(\beta x)$, $h_1(t) = e^{-\mu t} \cosh(\beta t)$, $g_2(x) = e^{\mu x} \cosh(\beta x)$, and $h_2(t) = -e^{-\mu t} \sinh(\beta t)$.

$$3. \quad y(x) - \lambda \int_a^b t e^{\mu(x-t)} \sinh[\beta(x-t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\mu x} \sinh(\beta x)$, $h_1(t) = t e^{-\mu t} \cosh(\beta t)$, $g_2(x) = e^{\mu x} \cosh(\beta x)$, and $h_2(t) = -t e^{-\mu t} \sinh(\beta t)$.

4.7-2. Kernels Containing Exponential and Logarithmic Functions.

$$4. \quad y(x) - \lambda \int_a^b e^{\mu t} \ln(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln(\beta x)$ and $h(t) = e^{\mu t}$.

$$5. \quad y(x) - \lambda \int_a^b e^{\mu x} \ln(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = e^{\mu x}$ and $h(t) = \ln(\beta t)$.

$$6. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} \ln(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = e^{\mu x} \ln(\beta x)$ and $h(t) = e^{-\mu t}$.

$$7. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} \ln(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = e^{\mu x}$ and $h(t) = e^{-\mu t} \ln(\beta t)$.

$$8. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} (\ln x - \ln t) y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\mu x} \ln x$, $h_1(t) = e^{-\mu t}$, $g_2(x) = e^{\mu x}$, and $h_2(t) = -e^{-\mu t} \ln t$.

$$9. \quad y(x) + \frac{b^2 - a^2}{2a} \int_0^\infty \frac{1}{t} \exp\left(-a \left| \ln \frac{x}{t} \right| \right) y(t) dt = f(x).$$

Solution with $a > 0$, $b > 0$, and $x > 0$:

$$y(x) = f(x) + \frac{a^2 - b^2}{2b} \int_0^\infty \frac{1}{t} \exp\left(-b \left| \ln \frac{x}{t} \right| \right) f(t) dt.$$

⊙ Reference: F. D. Gakhov and Yu. I. Cherskii (1978).

4.7-3. Kernels Containing Exponential and Trigonometric Functions.

$$10. \quad y(x) - \lambda \int_a^b e^{\mu t} \cos(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos(\beta x)$ and $h(t) = e^{\mu t}$.

$$11. \quad y(x) - \lambda \int_a^b e^{\mu x} \cos(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = e^{\mu x}$ and $h(t) = \cos(\beta t)$.

$$12. \quad y(x) - \lambda \int_0^\infty e^{\mu(x-t)} \cos(xt) y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^\infty e^{\mu(x-t)} \cos(xt) f(t) dt, \quad \lambda \neq \pm \sqrt{2/\pi}.$$

$$13. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} \cos[\beta(x-t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\mu x} \cos(\beta x)$, $h_1(t) = e^{-\mu t} \cos(\beta t)$, $g_2(x) = e^{\mu x} \sin(\beta x)$, and $h_2(t) = e^{-\mu t} \sin(\beta t)$.

$$14. \quad y(x) - \lambda \int_a^b e^{\mu t} \sin(\beta x)y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin(\beta x)$ and $h(t) = e^{\mu t}$.

$$15. \quad y(x) - \lambda \int_a^b e^{\mu x} \sin(\beta t)y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = e^{\mu x}$ and $h(t) = \sin(\beta t)$.

$$16. \quad y(x) - \lambda \int_0^\infty e^{\mu(x-t)} \sin(xt)y(t) dt = f(x).$$

Solution:

$$y(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^\infty e^{\mu(x-t)} \sin(xt)f(t) dt, \quad \lambda \neq \pm\sqrt{2/\pi}.$$

$$17. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} \sin[\beta(x-t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\mu x} \sin(\beta x)$, $h_1(t) = e^{-\mu t} \cos(\beta t)$, $g_2(x) = e^{\mu x} \cos(\beta x)$, and $h_2(t) = -e^{-\mu t} \sin(\beta t)$.

$$18. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} \left\{ \sum_{k=1}^n A_k \sin[\beta_k(x-t)] \right\} y(t) dt = f(x), \quad n = 1, 2, \dots$$

This is a special case of equation 4.9.20.

$$19. \quad y(x) - \lambda \int_a^b t e^{\mu(x-t)} \sin[\beta(x-t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\mu x} \sin(\beta x)$, $h_1(t) = t e^{-\mu t} \cos(\beta t)$, $g_2(x) = e^{\mu x} \cos(\beta x)$, and $h_2(t) = -t e^{-\mu t} \sin(\beta t)$.

$$20. \quad y(x) - \lambda \int_a^b x e^{\mu(x-t)} \sin[\beta(x-t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = x e^{\mu x} \sin(\beta x)$, $h_1(t) = e^{-\mu t} \cos(\beta t)$, $g_2(x) = x e^{\mu x} \cos(\beta x)$, and $h_2(t) = -e^{-\mu t} \sin(\beta t)$.

$$21. \quad y(x) - \lambda \int_a^b e^{\mu t} \tan(\beta x)y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tan(\beta x)$ and $h(t) = e^{\mu t}$.

$$22. \quad y(x) - \lambda \int_a^b e^{\mu x} \tan(\beta t)y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = e^{\mu x}$ and $h(t) = \tan(\beta t)$.

$$23. \quad y(x) - \lambda \int_a^b e^{\mu(x-t)} [\tan(\beta x) - \tan(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.18 with $g_1(x) = e^{\mu x} \tan(\beta x)$, $h_1(t) = e^{-\mu t}$, $g_2(x) = e^{\mu x}$, and $h_2(t) = -e^{-\mu t} \tan(\beta t)$.

$$24. \quad y(x) - \lambda \int_a^b e^{\mu t} \cot(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cot(\beta x)$ and $h(t) = e^{\mu t}$.

$$25. \quad y(x) - \lambda \int_a^b e^{\mu x} \cot(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = e^{\mu x}$ and $h(t) = \cot(\beta t)$.

4.7-4. Kernels Containing Hyperbolic and Logarithmic Functions.

$$26. \quad y(x) - \lambda \int_a^b \cosh^k(\beta x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cosh^k(\beta x)$ and $h(t) = \ln^m(\mu t)$.

$$27. \quad y(x) - \lambda \int_a^b \cosh^k(\beta t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\mu x)$ and $h(t) = \cosh^k(\beta t)$.

$$28. \quad y(x) - \lambda \int_a^b \sinh^k(\beta x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sinh^k(\beta x)$ and $h(t) = \ln^m(\mu t)$.

$$29. \quad y(x) - \lambda \int_a^b \sinh^k(\beta t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\mu x)$ and $h(t) = \sinh^k(\beta t)$.

$$30. \quad y(x) - \lambda \int_a^b \tanh^k(\beta x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tanh^k(\beta x)$ and $h(t) = \ln^m(\mu t)$.

$$31. \quad y(x) - \lambda \int_a^b \tanh^k(\beta t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\mu x)$ and $h(t) = \tanh^k(\beta t)$.

$$32. \quad y(x) - \lambda \int_a^b \coth^k(\beta x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \coth^k(\beta x)$ and $h(t) = \ln^m(\mu t)$.

$$33. \quad y(x) - \lambda \int_a^b \coth^k(\beta t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\mu x)$ and $h(t) = \coth^k(\beta t)$.

4.7-5. Kernels Containing Hyperbolic and Trigonometric Functions.

$$34. \quad y(x) - \lambda \int_a^b \cosh^k(\beta x) \cos^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cosh^k(\beta x)$ and $h(t) = \cos^m(\mu t)$.

$$35. \quad y(x) - \lambda \int_a^b \cosh^k(\beta t) \cos^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos^m(\mu x)$ and $h(t) = \cosh^k(\beta t)$.

$$36. \quad y(x) - \lambda \int_a^b \cosh^k(\beta x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cosh^k(\beta x)$ and $h(t) = \sin^m(\mu t)$.

$$37. \quad y(x) - \lambda \int_a^b \cosh^k(\beta t) \sin^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin^m(\mu x)$ and $h(t) = \cosh^k(\beta t)$.

$$38. \quad y(x) - \lambda \int_a^b \sinh^k(\beta x) \cos^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sinh^k(\beta x)$ and $h(t) = \cos^m(\mu t)$.

$$39. \quad y(x) - \lambda \int_a^b \sinh^k(\beta t) \cos^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos^m(\mu x)$ and $h(t) = \sinh^k(\beta t)$.

$$40. \quad y(x) - \lambda \int_a^b \sinh^k(\beta x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sinh^k(\beta x)$ and $h(t) = \sin^m(\mu t)$.

$$41. \quad y(x) - \lambda \int_a^b \sinh^k(\beta t) \sin^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin^m(\mu x)$ and $h(t) = \sinh^k(\beta t)$.

$$42. \quad y(x) - \lambda \int_a^b \tanh^k(\beta x) \cos^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tanh^k(\beta x)$ and $h(t) = \cos^m(\mu t)$.

$$43. \quad y(x) - \lambda \int_a^b \tanh^k(\beta t) \cos^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos^m(\mu x)$ and $h(t) = \tanh^k(\beta t)$.

$$44. \quad y(x) - \lambda \int_a^b \tanh^k(\beta x) \sin^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tanh^k(\beta x)$ and $h(t) = \sin^m(\mu t)$.

$$45. \quad y(x) - \lambda \int_a^b \tanh^k(\beta t) \sin^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin^m(\mu x)$ and $h(t) = \tanh^k(\beta t)$.

4.7-6. Kernels Containing Logarithmic and Trigonometric Functions.

$$46. \quad y(x) - \lambda \int_a^b \cos^k(\beta x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cos^k(\beta x)$ and $h(t) = \ln^m(\mu t)$.

$$47. \quad y(x) - \lambda \int_a^b \cos^k(\beta t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\mu x)$ and $h(t) = \cos^k(\beta t)$.

$$48. \quad y(x) - \lambda \int_a^b \sin^k(\beta x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \sin^k(\beta x)$ and $h(t) = \ln^m(\mu t)$.

$$49. \quad y(x) - \lambda \int_a^b \sin^k(\beta t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\mu x)$ and $h(t) = \sin^k(\beta t)$.

$$50. \quad y(x) - \lambda \int_a^b \tan^k(\beta x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \tan^k(\beta x)$ and $h(t) = \ln^m(\mu t)$.

$$51. \quad y(x) - \lambda \int_a^b \tan^k(\beta t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\mu x)$ and $h(t) = \tan^k(\beta t)$.

$$52. \quad y(x) - \lambda \int_a^b \cot^k(\beta x) \ln^m(\mu t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \cot^k(\beta x)$ and $h(t) = \ln^m(\mu t)$.

$$53. \quad y(x) - \lambda \int_a^b \cot^k(\beta t) \ln^m(\mu x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = \ln^m(\mu x)$ and $h(t) = \cot^k(\beta t)$.

4.8. Equations Whose Kernels Contain Special Functions

4.8-1. Kernels Containing Bessel Functions.

$$1. \quad y(x) - \lambda \int_a^b J_\nu(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = J_\nu(\beta x)$ and $h(t) = 1$.

$$2. \quad y(x) - \lambda \int_a^b J_\nu(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = J_\nu(\beta t)$.

$$3. \quad y(x) + \lambda \int_0^\infty t J_\nu(xt) y(t) dt = 0, \quad \nu > -1.$$

Characteristic values: $\lambda = \pm 1$. For the characteristic values, the integral equation has infinitely many linearly independent eigenfunctions.

Eigenfunctions for $\lambda = +1$ have the form

$$y_+(x) = f(x) - \int_0^\infty t J_\nu(xt) f(t) dt,$$

where $f = f(x)$ is an arbitrary function.

Eigenfunctions for $\lambda = -1$ have the form

$$y_-(x) = f(x) + \int_0^\infty t J_\nu(xt) f(t) dt,$$

where $f = f(x)$ is an arbitrary function.

$$4. \quad y(x) + \lambda \int_0^\infty t J_\nu(xt) y(t) dt = f(x), \quad \nu > -1.$$

Solution:

$$y(x) = \frac{f(x)}{1 - \lambda^2} - \frac{\lambda}{1 - \lambda^2} \int_0^\infty t J_\nu(xt) f(t) dt, \quad \lambda \neq \pm 1.$$

$$5. \quad y(x) + \lambda \int_0^\infty J_\nu(2\sqrt{xt}) y(t) dt = f(x).$$

By setting $x = \frac{1}{2}z^2$, $t = \frac{1}{2}\tau^2$, $y(x) = Y(z)$, and $f(x) = F(z)$, we arrive at an equation of the form 4.8.4:

$$Y(z) + \lambda \int_0^\infty \tau J_\nu(z\tau) Y(\tau) d\tau = F(z).$$

$$6. \quad y(x) - \lambda \int_a^b [A + B(x-t)J_\nu(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = J_\nu(\beta t)$.

$$7. \quad y(x) - \lambda \int_a^b [A + B(x-t)J_\nu(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = J_\nu(\beta x)$.

$$8. \quad y(x) - \lambda \int_a^b [AJ_\mu(\alpha x) + BJ_\nu(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.5 with $g(x) = AJ_\mu(\alpha x)$ and $h(t) = BJ_\nu(\beta t)$.

$$9. \quad y(x) - \lambda \int_a^b [AJ_\mu(x)J_\nu(t) + BJ_\nu(x)J_\mu(t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.17 with $g(x) = J_\mu(x)$ and $h(t) = J_\nu(t)$.

$$10. \quad y(x) - \lambda \int_a^b Y_\nu(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = Y_\nu(\beta x)$ and $h(t) = 1$.

$$11. \quad y(x) - \lambda \int_a^b Y_\nu(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = Y_\nu(\beta t)$.

$$12. \quad y(x) - \lambda \int_a^b [A + B(x-t)Y_\nu(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = Y_\nu(\beta t)$.

$$13. \quad y(x) - \lambda \int_a^b [A + B(x-t)Y_\nu(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = Y_\nu(\beta x)$.

$$14. \quad y(x) - \lambda \int_a^b [AY_\mu(\alpha x) + BY_\nu(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.5 with $g(x) = AY_\mu(\alpha x)$ and $h(t) = BY_\nu(\beta t)$.

$$15. \quad y(x) - \lambda \int_a^b [AY_\mu(x)Y_\nu(t) + BY_\nu(x)Y_\mu(t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.14 with $g(x) = Y_\mu(x)$ and $h(t) = Y_\nu(t)$.

$$16. \quad y(x) - \lambda \int_a^b [AY_\mu(x)Y_\nu(t) + BY_\nu(x)Y_\mu(t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.17 with $g(x) = Y_\mu(x)$ and $h(t) = Y_\nu(t)$.

4.8-2. Kernels Containing Modified Bessel Functions.
--

$$17. \quad y(x) - \lambda \int_a^b I_\nu(\beta x) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = I_\nu(\beta x)$ and $h(t) = 1$.

$$18. \quad y(x) - \lambda \int_a^b I_\nu(\beta t) y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = I_\nu(\beta t)$.

$$19. \quad y(x) - \lambda \int_a^b [A + B(x-t)I_\nu(\beta t)] y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = I_\nu(\beta t)$.

$$20. \quad y(x) - \lambda \int_a^b [A + B(x-t)I_\nu(\beta x)] y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = I_\nu(\beta x)$.

$$21. \quad y(x) - \lambda \int_a^b [AI_\mu(\alpha x) + BI_\nu(\beta t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.5 with $g(x) = AI_\mu(\alpha x)$ and $h(t) = BI_\nu(\beta t)$.

$$22. \quad y(x) - \lambda \int_a^b [AI_\mu(x)I_\mu(t) + BI_\nu(x)I_\nu(t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.14 with $g(x) = I_\mu(x)$ and $h(t) = I_\nu(t)$.

$$23. \quad y(x) - \lambda \int_a^b [AI_\mu(x)I_\nu(t) + BI_\nu(x)I_\mu(t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.17 with $g(x) = I_\mu(x)$ and $h(t) = I_\nu(t)$.

$$24. \quad y(x) - \lambda \int_a^b K_\nu(\beta x)y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = K_\nu(\beta x)$ and $h(t) = 1$.

$$25. \quad y(x) - \lambda \int_a^b K_\nu(\beta t)y(t) dt = f(x).$$

This is a special case of equation 4.9.1 with $g(x) = 1$ and $h(t) = K_\nu(\beta t)$.

$$26. \quad y(x) - \lambda \int_a^b [A + B(x - t)K_\nu(\beta t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.8 with $h(t) = K_\nu(\beta t)$.

$$27. \quad y(x) - \lambda \int_a^b [A + B(x - t)K_\nu(\beta x)]y(t) dt = f(x).$$

This is a special case of equation 4.9.10 with $h(x) = K_\nu(\beta x)$.

$$28. \quad y(x) - \lambda \int_a^b [AK_\mu(\alpha x) + BK_\nu(\beta t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.5 with $g(x) = AK_\mu(\alpha x)$ and $h(t) = BK_\nu(\beta t)$.

$$29. \quad y(x) - \lambda \int_a^b [AK_\mu(x)K_\mu(t) + BK_\nu(x)K_\nu(t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.14 with $g(x) = K_\mu(x)$ and $h(t) = K_\nu(t)$.

$$30. \quad y(x) - \lambda \int_a^b [AK_\mu(x)K_\nu(t) + BK_\nu(x)K_\mu(t)]y(t) dt = f(x).$$

This is a special case of equation 4.9.17 with $g(x) = K_\mu(x)$ and $h(t) = K_\nu(t)$.

4.9. Equations Whose Kernels Contain Arbitrary Functions

4.9-1. Equations with Degenerate Kernel: $K(x, t) = g_1(x)h_1(t) + \dots + g_n(x)h_n(t)$.

1. $y(x) - \lambda \int_a^b g(x)h(t)y(t) dt = f(x).$

1°. Assume that $\lambda \neq \left(\int_a^b g(t)h(t) dt \right)^{-1}$.

Solution:

$$y(x) = f(x) + \lambda k g(x), \quad \text{where } k = \left(1 - \lambda \int_a^b g(t)h(t) dt \right)^{-1} \int_a^b h(t)f(t) dt.$$

2°. Assume that $\lambda = \left(\int_a^b g(t)h(t) dt \right)^{-1}$.

For $\int_a^b h(t)f(t) dt = 0$, the solution has the form

$$y = f(x) + Cg(x),$$

where C is an arbitrary constant.

For $\int_a^b h(t)f(t) dt \neq 0$, there is no solution.

The limits of integration may take the values $a = -\infty$ and/or $b = \infty$, provided that the corresponding improper integral converges.

2. $y(x) - \lambda \int_a^b [g(x) + g(t)]y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_1 = \frac{1}{g_1 + \sqrt{(b-a)g_2}}, \quad \lambda_2 = \frac{1}{g_1 - \sqrt{(b-a)g_2}},$$

where

$$g_1 = \int_a^b g(x) dx, \quad g_2 = \int_a^b g^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda[f_1g_1 - (b-a)f_2]}{[g_1^2 - (b-a)g_2]\lambda^2 - 2g_1\lambda + 1}, \quad A_2 = \frac{f_2 - \lambda(f_2g_1 - f_1g_2)}{[g_1^2 - (b-a)g_2]\lambda^2 - 2g_1\lambda + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b f(x)g(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \sqrt{\frac{g_2}{b-a}},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_2(x), \quad y_2(x) = g(x) - \sqrt{\frac{g_2}{b-a}},$$

where C is an arbitrary constant and $y_2(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_2 .

4°. The equation has no multiple characteristic values.

3. $y(x) - \lambda \int_a^b [g(x) - g(t)]y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_1 = \frac{1}{\sqrt{g_1^2 - (b-a)g_2}}, \quad \lambda_2 = -\frac{1}{\sqrt{g_1^2 - (b-a)g_2}},$$

where

$$g_1 = \int_a^b g(x) dx, \quad g_2 = \int_a^b g^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 + \lambda[f_1g_1 - (b-a)f_2]}{[(b-a)g_2 - g_1^2]\lambda^2 + 1}, \quad A_2 = \frac{-f_2 + \lambda(f_2g_1 - f_1g_2)}{[(b-a)g_2 - g_1^2]\lambda^2 + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b f(x)g(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \frac{1 - \lambda_1g_1}{\lambda_1(b-a)},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. The equation has no multiple characteristic values.

4. $y(x) - \lambda \int_a^b [Ag(x) + Bg(t)]y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{(A+B)g_1 \pm \sqrt{(A-B)^2g_1^2 + 4AB(b-a)g_2}}{2AB[g_1^2 - (b-a)g_2]},$$

where

$$g_1 = \int_a^b g(x) dx, \quad g_2 = \int_a^b g^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{Af_1 - \lambda AB[f_1g_1 - (b-a)f_2]}{AB[g_1^2 - (b-a)g_2]\lambda^2 - (A+B)g_1\lambda + 1}, \quad A_2 = \frac{Bf_2 - \lambda AB(f_2g_1 - f_1g_2)}{AB[g_1^2 - (b-a)g_2]\lambda^2 - (A+B)g_1\lambda + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b f(x)g(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \frac{1 - \lambda_1 Ag_1}{\lambda_1 A(b-a)},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq \pm B$, where the characteristic value $\lambda_* = \frac{2}{(A+B)g_1}$ is double:

$$y(x) = f(x) + Cy_*(x), \quad y_*(x) = g(x) - \frac{(A-B)g_1}{2A(b-a)}.$$

Here C is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A = \pm B$.

5. $y(x) - \lambda \int_a^b [g(x) + h(t)]y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{s_1 + s_3 \pm \sqrt{(s_1 - s_3)^2 + 4(b-a)s_2}}{2[s_1s_3 - (b-a)s_2]},$$

where

$$s_1 = \int_a^b g(x) dx, \quad s_2 = \int_a^b g(x)h(x) dx, \quad s_3 = \int_a^b h(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda[f_1s_3 - (b-a)f_2]}{[s_1s_3 - (b-a)s_2]\lambda^2 - (s_1 + s_3)\lambda + 1}, \quad A_2 = \frac{f_2 - \lambda(f_2s_1 - f_1s_2)}{[s_1s_3 - (b-a)s_2]\lambda^2 - (s_1 + s_3)\lambda + 1},$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b f(x)h(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \frac{1 - \lambda_1 s_1}{\lambda_1 (b-a)},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $s_1 \neq \pm s_3$, where the characteristic value $\lambda_* = \frac{2}{s_1 + s_3}$ is double:

$$y(x) = f(x) + C y_*(x), \quad y_*(x) = g(x) - \frac{s_1 - s_3}{2(b-a)}.$$

Here C is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $s_1 = \pm s_3$.

6.
$$y(x) - \lambda \int_a^b [A g(x) + B g(t)] h(t) y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{(A+B)s_1 \pm \sqrt{(A-B)^2 s_1^2 + 4ABs_0 s_2}}{2AB(s_1^2 - s_0 s_2)},$$

where

$$s_0 = \int_a^b h(x) dx, \quad s_1 = \int_a^b g(x) h(x) dx, \quad s_2 = \int_a^b g^2(x) h(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda [A_1 g(x) + A_2],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{A f_1 - AB \lambda (f_1 s_1 - f_2 s_0)}{AB(s_1^2 - s_0 s_2) \lambda^2 - (A+B)s_1 \lambda + 1}, \quad A_2 = \frac{B f_2 - AB \lambda (f_2 s_1 - f_1 s_2)}{AB(s_1^2 - s_0 s_2) \lambda^2 - (A+B)s_1 \lambda + 1},$$

$$f_1 = \int_a^b f(x) h(x) dx, \quad f_2 = \int_a^b f(x) g(x) h(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + C y_1(x), \quad y_1(x) = g(x) + \frac{1 - \lambda_1 A s_1}{\lambda_1 A s_0},$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq \pm B$, where the characteristic value $\lambda_* = \frac{2}{(A+B)s_1}$ is double:

$$y(x) = f(x) + C y_*(x),$$

where C is an arbitrary constant and

$$y_*(x) = g(x) - \frac{(A-B)s_1}{2A s_0}$$

is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A = \pm B$.

$$7. \quad y(x) - \lambda \int_a^b [Ag(x) + Bg(t) + C]h(t) y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{(A+B)s_1 + Cs_0 \pm \sqrt{(A-B)^2s_1^2 + 2(A+B)Cs_1s_0 + C^2s_0^2 + 4ABs_0s_2}}{2AB(s_1^2 - s_0s_2)},$$

where

$$s_0 = \int_a^b h(x) dx, \quad s_1 = \int_a^b g(x)h(x) dx, \quad s_2 = \int_a^b g^2(x)h(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{Af_1 - AB\lambda(f_1s_1 - f_2s_0)}{AB(s_1^2 - s_0s_2)\lambda^2 - [(A+B)s_1 + Cs_0]\lambda + 1},$$

$$A_2 = \frac{C_1f_1 + Bf_2 - AB\lambda(f_2s_1 - f_1s_2)}{AB(s_1^2 - s_0s_2)\lambda^2 - [(A+B)s_1 + Cs_0]\lambda + 1},$$

$$f_1 = \int_a^b f(x)h(x) dx, \quad f_2 = \int_a^b f(x)g(x)h(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + \tilde{C}y_1(x), \quad y_1(x) = g(x) + \frac{1 - \lambda_1As_1}{\lambda_1As_0},$$

where \tilde{C} is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $(A \pm B)s_1 \pm Cs_0 \neq 0$, where the characteristic value $\lambda_* = \frac{2}{(A+B)s_1 + Cs_0}$ is double:

$$y(x) = f(x) + \tilde{C}y_*(x),$$

where \tilde{C} is an arbitrary constant and

$$y_*(x) = g(x) - \frac{(A-B)s_1 - Cs_0}{2As_0}$$

is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $(A \pm B)s_1 \pm Cs_0 = 0$.

$$8. \quad y(x) - \lambda \int_a^b [A + B(x-t)h(t)]y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{A(b-a) \pm \sqrt{[A(b-a) - 2Bh_1]^2 + 2Bh_0[A(b^2 - a^2) - 2Bh_2]}}{B\{A(b-a)[2h_1 - (b+a)h_0] - 2B(h_1^2 - h_0h_2)\}},$$

where

$$h_0 = \int_a^b h(x) dx, \quad h_1 = \int_a^b xh(x) dx, \quad h_2 = \int_a^b x^2h(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda [B(f_1h_1 + f_2h_2) - \frac{1}{2}Af_2(b^2 - a^2)]}{B\{A(b-a)[h_1 - \frac{1}{2}(b+a)h_0] - B(h_1^2 - h_0h_2)\} \lambda^2 + A(b-a)\lambda + 1},$$

$$A_2 = \frac{f_2 - \lambda[A(b-a)f_2 - B(f_1h_0 + f_2h_1)]}{B\{A(b-a)[h_1 - \frac{1}{2}(b+a)h_0] - B(h_1^2 - h_0h_2)\} \lambda^2 + A(b-a)\lambda + 1},$$

$$f_1 = A \int_a^b f(x) dx - B \int_a^b xf(x)h(x) dx, \quad f_2 = B \int_a^b f(x)h(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = 1 + \frac{2 - 2\lambda_1[A(b-a) - Bh_1]}{\lambda_1[A(b^2 - a^2) - 2Bh_2]} x,$$

where C is an arbitrary constant, and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq 0$ or $2Bh_1 - A(b-a) \neq 0$, where the characteristic value $\lambda_* = \frac{2}{A(b-a)}$ is double:

$$y(x) = f(x) + Cy_*(x),$$

where C is an arbitrary constant, and

$$y_*(x) = 1 - \frac{A(b-a) - 2Bh_1}{A(b^2 - a^2) - 2Bh_2} x$$

is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A = 0$ or $2Bh_1 - A(b-a) = 0$.

$$9. \quad y(x) - \lambda \int_a^b [A + (Bx + Ct)h(t)]y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{A(b-a) + (C+B)h_1 \pm \sqrt{D}}{B\{A(b-a)[2h_1 - (b+a)h_0] + 2C(h_1^2 - h_0h_2)\}},$$

$$D = [A(b-a) + (C-B)h_1]^2 + 2Bh_0[A(b^2 - a^2) + 2Ch_2],$$

where

$$h_0 = \int_a^b h(x) dx, \quad h_1 = \int_a^b xh(x) dx, \quad h_2 = \int_a^b x^2 h(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda(A_1 + A_2x),$$

where the constants A_1 and A_2 are given by

$$A_1 = \Delta^{-1} \left\{ f_1 - \lambda \left[Bf_1h_1 - Cf_2h_2 - \frac{1}{2}A(b^2 - a^2)f_2 \right] \right\},$$

$$A_2 = \Delta^{-1} \left\{ f_2 - \lambda \left[A(b-a)f_2 - Bf_1h_0 + Cf_2h_1 \right] \right\},$$

$$\Delta = B \left\{ A(b-a) \left[h_1 - \frac{1}{2}(b+a)h_0 \right] + C(h_1^2 - h_0h_2) \right\} \lambda^2 + [A(b-a) + (B+C)h_1] \lambda + 1,$$

$$f_1 = A \int_a^b f(x) dx + C \int_a^b xf(x)h(x) dx, \quad f_2 = B \int_a^b f(x)h(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + \tilde{C}y_1(x), \quad y_1(x) = 1 + \frac{2 - 2\lambda_1[A(b-a) + Ch_1]}{\lambda_1[A(b^2 - a^2) + 2Ch_2]} x,$$

where \tilde{C} is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $\pm A(b-a) + (B \pm C)h_1 \neq 0$, where the characteristic value $\lambda_* = \frac{2}{A(b-a) + (B+C)h_1}$ is double:

$$y(x) = f(x) + \tilde{C}y_*(x),$$

where \tilde{C} is an arbitrary constant and

$$y_*(x) = 1 - \frac{A(b-a) + (C-B)h_1}{A(b^2 - a^2) + 2Ch_2} x$$

is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $\pm A(b-a) + (B \pm C)h_1 = 0$.

$$10. \quad y(x) - \lambda \int_a^b [A + B(x-t)h(x)]y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{A(b-a) \pm \sqrt{[A(b-a) + 2Bh_1]^2 - 4Bh_0[A(b-a) + Bh_2]}}{2B\{h_0[A(b-a) + Bh_2] - h_1[A(b-a) + Bh_1]\}},$$

where

$$h_0 = \int_a^b h(x) dx, \quad h_1 = \int_a^b xh(x) dx, \quad h_2 = \int_a^b x^2 h(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda [AE_1 + (BE_1x + E_2)h(x)],$$

where the constants E_1 and E_2 are given by

$$\begin{aligned} E_1 &= \Delta^{-1} [f_1 + \lambda B(f_1 h_1 - f_2 h_0)], \\ E_2 &= \Delta^{-1} \{f_2 - \lambda f_2 [A(b-a) + Bh_1] - \lambda f_1 [A(b-a) + Bh_2]\}, \\ \Delta &= B \{h_0 [A(b-a) + Bh_2] - h_1 [A(b-a) + Bh_1]\} \lambda^2 - A(b-a)\lambda + 1, \\ f_1 &= \int_a^b f(x) dx, \quad f_2 = \int_a^b xf(x) dx. \end{aligned}$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = A + Bxh(x) + \frac{1 - \lambda_1 [A(b-a) + Bh_1]}{\lambda_1 h_0} h(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq 0$ or $A(b-a) + 4Bh_1 \neq 0$, where the characteristic value $\lambda_* = \frac{2}{A(b-a)}$ is double:

$$y(x) = f(x) + Cy_*(x),$$

where C is an arbitrary constant and

$$y_*(x) = A + Bxh(x) - \frac{A(b-a) + 2Bh_1}{2h_0} h(x)$$

is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A = 0$ or $A(b-a) + 4Bh_1 = 0$.

$$11. \quad y(x) - \lambda \int_a^b [A + (Bx + Ct)h(x)]y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{A(b-a) + (B+C)h_1 \pm \sqrt{D}}{2C\{h_1[A(b-a) + Bh_1] - h_0[A(b-a) + Bh_2]\}},$$

$$D = [A(b-a) + (B-C)h_1]^2 + 4Ch_0[A(b-a) + Bh_2],$$

where

$$h_0 = \int_a^b h(x) dx, \quad h_1 = \int_a^b xh(x) dx, \quad h_2 = \int_a^b x^2 h(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda [AE_1 + (BE_1x + E_2)h(x)],$$

where the constants E_1 and E_2 are given by

$$E_1 = \Delta^{-1}[f_1 - \lambda C(f_1 h_1 - f_2 h_0)],$$

$$E_2 = C\Delta^{-1}\{f_2 - \lambda f_2[A(b-a) + Bh_1] - \lambda f_1[A(b-a) + Bh_2]\},$$

$$\Delta = C\{h_1[A(b-a) + Bh_1] - h_0[A(b-a) + Bh_2]\}\lambda^2 - [A(b-a) + (B+C)h_1]\lambda + 1,$$

$$f_1 = \int_a^b f(x) dx, \quad f_2 = \int_a^b xf(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + \tilde{C}y_1(x), \quad y_1(x) = A + Bxh(x) + \frac{1 - \lambda_1[A(b-a) + Bh_1]}{\lambda_1 h_0} h(x),$$

where \tilde{C} is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A(b-a) + (B \pm C)h_1 \neq 0$, where the characteristic value $\lambda_* = \frac{2}{A(b-a) + (B+C)h_1}$ is double:

$$y(x) = f(x) + \tilde{C}y_*(x),$$

where \tilde{C} is an arbitrary constant and

$$y_*(x) = A + Bxh(x) - \frac{A(b-a) + (B-C)h_1}{2h_0} h(x)$$

is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A(b-a) + (B \pm C)h_1 = 0$.

$$12. \quad y(x) - \lambda \int_a^b [g(x)g(t) + h(x)h(t)]y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_1 = \frac{s_1 + s_3 + \sqrt{(s_1 - s_3)^2 + 4s_2^2}}{2(s_1s_3 - s_2^2)}, \quad \lambda_2 = \frac{s_1 + s_3 - \sqrt{(s_1 - s_3)^2 + 4s_2^2}}{2(s_1s_3 - s_2^2)},$$

where

$$s_1 = \int_a^b g^2(x) dx, \quad s_2 = \int_a^b g(x)h(x) dx, \quad s_3 = \int_a^b h^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2h(x)],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda(f_1s_3 - f_2s_2)}{(s_1s_3 - s_2^2)\lambda^2 - (s_1 + s_3)\lambda + 1}, \quad A_2 = \frac{f_2 - \lambda(f_2s_1 - f_1s_2)}{(s_1s_3 - s_2^2)\lambda^2 - (s_1 + s_3)\lambda + 1},$$

$$f_1 = \int_a^b f(x)g(x) dx, \quad f_2 = \int_a^b f(x)h(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + (1 - \lambda_1s_1)h(x)/(\lambda_1s_2),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $s_1 \neq \pm s_3$, where the characteristic value $\lambda_* = 1/s_1$ is double:

$$y(x) = f(x) + \tilde{C}_1g(x) + \tilde{C}_2h(x),$$

where \tilde{C}_1 and \tilde{C}_2 are arbitrary constants.

The equation has no multiple characteristic values if $s_1 = \pm s_3$.

$$13. \quad y(x) - \lambda \int_a^b [g(x)g(t) - h(x)h(t)]y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_1 = \frac{s_1 - s_3 + \sqrt{(s_1 + s_3)^2 - 4s_2^2}}{2(s_2^2 - s_1s_3)}, \quad \lambda_2 = \frac{s_1 - s_3 - \sqrt{(s_1 + s_3)^2 - 4s_2^2}}{2(s_2^2 - s_1s_3)},$$

where

$$s_1 = \int_a^b g^2(x) dx, \quad s_2 = \int_a^b g(x)h(x) dx, \quad s_3 = \int_a^b h^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2h(x)],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 + \lambda(f_1s_3 - f_2s_2)}{(s_2^2 - s_1s_3)\lambda^2 - (s_1 - s_3)\lambda + 1}, \quad A_2 = \frac{-f_2 + \lambda(f_2s_1 - f_1s_2)}{(s_2^2 - s_1s_3)\lambda^2 - (s_1 - s_3)\lambda + 1},$$

$$f_1 = \int_a^b f(x)g(x) dx, \quad f_2 = \int_a^b f(x)h(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \frac{1 - \lambda_1s_1}{\lambda_1s_2}h(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $s_1 \neq \pm s_3$, where the characteristic value $\lambda_* = \frac{2}{s_1 - s_3}$ is double:

$$y(x) = f(x) + Cy_*(x), \quad y_*(x) = g(x) - \frac{s_1 + s_3}{2s_2}h(x),$$

where C is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $s_1 = \pm s_3$.

14. $y(x) - \lambda \int_a^b [Ag(x)g(t) + Bh(x)h(t)]y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{As_1 + Bs_3 \pm \sqrt{(As_1 - Bs_3)^2 + 4ABs_2^2}}{2AB(s_1s_3 - s_2^2)},$$

where

$$s_1 = \int_a^b g^2(x) dx, \quad s_2 = \int_a^b g(x)h(x) dx, \quad s_3 = \int_a^b h^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2h(x)],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{Af_1 - \lambda AB(f_1s_3 - f_2s_2)}{AB(s_1s_3 - s_2^2)\lambda^2 - (As_1 + Bs_3)\lambda + 1}, \quad A_2 = \frac{Bf_2 - \lambda AB(f_2s_1 - f_1s_2)}{AB(s_1s_3 - s_2^2)\lambda^2 - (As_1 + Bs_3)\lambda + 1},$$

$$f_1 = \int_a^b f(x)g(x) dx, \quad f_2 = \int_a^b f(x)h(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \frac{1 - \lambda_1As_1}{\lambda_1As_2}h(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $As_1 \neq \pm Bs_3$, where the characteristic value $\lambda_* = \frac{2}{As_1 + Bs_3}$ is double:

$$y(x) = f(x) + Cy_*(x),$$

where C is an arbitrary constant and

$$y_*(x) = g(x) - \frac{As_1 - Bs_3}{2As_2} h(x)$$

is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $As_1 = \pm Bs_3$.

15. $y(x) - \lambda \int_a^b [g(x)h(t) + h(x)g(t)]y(t) dt = f(x).$

The characteristic values of the equation:

$$\lambda_1 = \frac{1}{s_1 + \sqrt{s_2 s_3}}, \quad \lambda_2 = \frac{1}{s_1 - \sqrt{s_2 s_3}},$$

where

$$s_1 = \int_a^b h(x)g(x) dx, \quad s_2 = \int_a^b h^2(x) dx, \quad s_3 = \int_a^b g^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1 g(x) + A_2 h(x)],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda(f_1 s_1 - f_2 s_2)}{(s_1^2 - s_2 s_3)\lambda^2 - 2s_1 \lambda + 1}, \quad A_2 = \frac{f_2 - \lambda(f_2 s_1 - f_1 s_3)}{(s_1^2 - s_2 s_3)\lambda^2 - 2s_1 \lambda + 1},$$

$$f_1 = \int_a^b f(x)h(x) dx, \quad f_2 = \int_a^b f(x)g(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \sqrt{\frac{s_3}{s_2}} h(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_2(x), \quad y_2(x) = g(x) - \sqrt{\frac{s_3}{s_2}} h(x),$$

where C is an arbitrary constant and $y_2(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_2 .

4°. The equation has no multiple characteristic values.

$$16. \quad y(x) - \lambda \int_a^b [g(x)h(t) - h(x)g(t)]y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_1 = \frac{1}{\sqrt{s_1^2 - s_2s_3}}, \quad \lambda_2 = -\frac{1}{\sqrt{s_1^2 - s_2s_3}},$$

where

$$s_1 = \int_a^b h(x)g(x) dx, \quad s_2 = \int_a^b h^2(x) dx, \quad s_3 = \int_a^b g^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2h(x)],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 + \lambda(f_1s_1 - f_2s_2)}{(s_2s_3 - s_1^2)\lambda^2 + 1}, \quad A_2 = \frac{-f_2 + \lambda(f_2s_1 - f_1s_3)}{(s_2s_3 - s_1^2)\lambda^2 + 1},$$

$$f_1 = \int_a^b f(x)h(x) dx, \quad f_2 = \int_a^b f(x)g(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \frac{\sqrt{s_1^2 - s_2s_3} - s_1}{s_2} h(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. Solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_2(x), \quad y_2(x) = g(x) - \frac{\sqrt{s_1^2 - s_2s_3} + s_1}{s_2} h(x),$$

where C is an arbitrary constant and $y_2(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_2 .

4°. The equation has no multiple characteristic values.

$$17. \quad y(x) - \lambda \int_a^b [Ag(x)h(t) + Bh(x)g(t)]y(t) dt = f(x).$$

The characteristic values of the equation:

$$\lambda_{1,2} = \frac{(A+B)s_1 \pm \sqrt{(A-B)^2s_1^2 + 4ABs_2s_3}}{2AB(s_1^2 - s_2s_3)},$$

where

$$s_1 = \int_a^b h(x)g(x) dx, \quad s_2 = \int_a^b h^2(x) dx, \quad s_3 = \int_a^b g^2(x) dx.$$

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g(x) + A_2h(x)],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{Af_1 - \lambda AB(f_1s_1 - f_2s_2)}{AB(s_1^2 - s_2s_3)\lambda^2 - (A+B)s_1\lambda + 1}, \quad A_2 = \frac{Bf_2 - \lambda AB(f_2s_1 - f_1s_3)}{AB(s_1^2 - s_2s_3)\lambda^2 - (A+B)s_1\lambda + 1},$$

$$f_1 = \int_a^b f(x)h(x) dx, \quad f_2 = \int_a^b f(x)g(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x), \quad y_1(x) = g(x) + \frac{1 - \lambda_1 A s_1}{\lambda_1 A s_2} h(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 .

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $A \neq \pm B$, where the characteristic value $\lambda_* = \frac{2}{(A+B)s_1}$ is double:

$$y(x) = f(x) + Cy_*(x), \quad y_*(x) = g(x) - \frac{(A-B)s_1}{2As_2} h(x).$$

Here C is an arbitrary constant and $y_*(x)$ is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $A = \pm B$.

18. $y(x) - \lambda \int_a^b [g_1(x)h_1(t) + g_2(x)h_2(t)]y(t) dt = f(x).$

The characteristic values of the equation λ_1 and λ_2 are given by

$$\lambda_{1,2} = \frac{s_{11} + s_{22} \pm \sqrt{(s_{11} - s_{22})^2 + 4s_{12}s_{21}}}{2(s_{11}s_{22} - s_{12}s_{21})},$$

provided that the integrals

$$s_{11} = \int_a^b h_1(x)g_1(x) dx, \quad s_{12} = \int_a^b h_1(x)g_2(x) dx, \quad s_{21} = \int_a^b h_2(x)g_1(x) dx, \quad s_{22} = \int_a^b h_2(x)g_2(x) dx$$

are convergent.

1°. Solution with $\lambda \neq \lambda_{1,2}$:

$$y(x) = f(x) + \lambda[A_1g_1(x) + A_2g_2(x)],$$

where the constants A_1 and A_2 are given by

$$A_1 = \frac{f_1 - \lambda(f_1s_{22} - f_2s_{12})}{(s_{11}s_{22} - s_{12}s_{21})\lambda^2 - (s_{11} + s_{22})\lambda + 1}, \quad A_2 = \frac{f_2 - \lambda(f_2s_{11} - f_1s_{21})}{(s_{11}s_{22} - s_{12}s_{21})\lambda^2 - (s_{11} + s_{22})\lambda + 1},$$

$$f_1 = \int_a^b f(x)h_1(x) dx, \quad f_2 = \int_a^b f(x)h_2(x) dx.$$

2°. Solution with $\lambda = \lambda_1 \neq \lambda_2$ and $f_1 = f_2 = 0$:

$$y(x) = f(x) + Cy_1(x),$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction of the equation corresponding to the characteristic value λ_1 :

$$y_1(x) = g_1(x) + \frac{1 - \lambda_1 s_{11}}{\lambda_1 s_{12}} g_2(x) = g_1(x) + \frac{\lambda_1 s_{21}}{1 - \lambda_1 s_{22}} g_2(x).$$

3°. The solution with $\lambda = \lambda_2 \neq \lambda_1$ and $f_1 = f_2 = 0$ is given by the formulas of item 2° in which one must replace λ_1 and $y_1(x)$ by λ_2 and $y_2(x)$, respectively.

4°. Solution with $\lambda = \lambda_{1,2} = \lambda_*$ and $f_1 = f_2 = 0$ provided that $s_{11} \neq \pm s_{22}$, where the characteristic value $\lambda_* = \frac{2}{s_{11} + s_{22}}$ is double:

$$y(x) = f(x) + Cy_*(x),$$

where C is an arbitrary constant and

$$y_*(x) = g_1(x) - \frac{s_{11} - s_{22}}{2s_{12}} g_2(x)$$

is an eigenfunction of the equation corresponding to λ_* .

The equation has no multiple characteristic values if $s_{11} = \pm s_{22}$.

$$19. \quad y(x) - \lambda \int_a^b [g(x) + h(t)]^m y(t) dt = f(x), \quad m = 1, 2, \dots$$

This is a special case of equation 4.9.20 with $g_k(x) = g^k(x)$, $h_k(t) = C_m^k h^{m-k}(t)$, and $k = 1, \dots, m$.

Solution:

$$y(x) = f(x) + \lambda \sum_{k=0}^m A_k g^k(x),$$

where the A_k are constants that can be determined from 4.9.20.

$$20. \quad y(x) - \lambda \int_a^b \left[\sum_{k=1}^n g_k(x) h_k(t) \right] y(t) dt = f(x), \quad n = 2, 3, \dots$$

The characteristic values of the integral equation (counting the multiplicity, we have exactly n of them) are the roots of the algebraic equation

$$\Delta(\lambda) = 0,$$

where

$$\Delta(\lambda) = \begin{vmatrix} 1 - \lambda s_{11} & -\lambda s_{12} & -\lambda s_{13} & \cdots & -\lambda s_{1n} \\ -\lambda s_{21} & 1 - \lambda s_{22} & -\lambda s_{23} & \cdots & -\lambda s_{2n} \\ -\lambda s_{31} & -\lambda s_{32} & 1 - \lambda s_{33} & \cdots & -\lambda s_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\lambda s_{n1} & -\lambda s_{n2} & -\lambda s_{n3} & \cdots & 1 - \lambda s_{nn} \end{vmatrix} \\ = (-\lambda)^n \begin{vmatrix} s_{11} - \lambda^{-1} & s_{12} & s_{13} & \cdots & s_{1n} \\ s_{21} & s_{22} - \lambda^{-1} & s_{23} & \cdots & s_{2n} \\ s_{31} & s_{32} & s_{33} - \lambda^{-1} & \cdots & s_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ s_{n1} & s_{n2} & s_{n3} & \cdots & s_{nn} - \lambda^{-1} \end{vmatrix},$$

and the integrals

$$s_{mk} = \int_a^b h_m(x) g_k(x) dx; \quad m, k = 1, \dots, n,$$

are assumed to be convergent.

Solution with regular λ :

$$y(x) = f(x) + \lambda \sum_{k=1}^n A_k g_k(x),$$

where the constants A_k form the solution of the following system of algebraic equations:

$$A_m - \lambda \sum_{k=1}^n s_{mk} A_k = f_m, \quad f_m = \int_a^b f(x) h_m(x) dx, \quad m = 1, \dots, n.$$

The A_k can be calculated by Cramer's rule:

$$A_k = \Delta_k(\lambda) / \Delta(\lambda),$$

where

$$\Delta_k(\lambda) = \begin{vmatrix} 1 - \lambda s_{11} & \cdots & -\lambda s_{1k-1} & f_1 & -\lambda s_{1k+1} & \cdots & -\lambda s_{1n} \\ -\lambda s_{21} & \cdots & -\lambda s_{2k-1} & f_2 & -\lambda s_{2k+1} & \cdots & -\lambda s_{2n} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -\lambda s_{n1} & \cdots & -\lambda s_{nk-1} & f_n & -\lambda s_{nk+1} & \cdots & 1 - \lambda s_{nn} \end{vmatrix}.$$

For solutions of the equation in the case in which λ is a characteristic value, see Subsection 13.2-2.

⊙ Reference: S. G. Mikhlin (1960).

4.9-2. Equations with Difference Kernel: $K(x, t) = K(x - t)$.

21. $y(x) = \lambda \int_{-\pi}^{\pi} K(x-t)y(t) dt, \quad K(x) = K(-x).$

Characteristic values:

$$\lambda_n = \frac{1}{\pi a_n}, \quad a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} K(x) \cos(nx) dx \quad (n = 0, 1, 2, \dots).$$

The corresponding eigenfunctions are

$$y_0(x) = 1, \quad y_n^{(1)}(x) = \cos(nx), \quad y_n^{(2)}(x) = \sin(nx) \quad (n = 1, 2, \dots).$$

For each value λ_n with $n \neq 0$, there are two corresponding linearly independent eigenfunctions $y_n^{(1)}(x)$ and $y_n^{(2)}(x)$.

⊙ Reference: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971).

22. $y(x) + \int_{-\infty}^{\infty} K(x-t)y(t) dt = Ae^{\lambda x}.$

Solution:

$$y(x) = \frac{A}{1+q} e^{\lambda x}, \quad q = \int_{-\infty}^{\infty} K(x) e^{-\lambda x} dx.$$

23. $y(x) + \int_{-\infty}^{\infty} K(x-t)y(t) dt = A \cos(\lambda x) + B \sin(\lambda x).$

Solution:

$$y(x) = \frac{AI_c + BI_s}{I_c^2 + I_s^2} \cos(\lambda x) + \frac{BI_c - AI_s}{I_c^2 + I_s^2} \sin(\lambda x),$$

$$I_c = 1 + \int_{-\infty}^{\infty} K(z) \cos(\lambda z) dz, \quad I_s = \int_{-\infty}^{\infty} K(z) \sin(\lambda z) dz.$$

$$24. \quad y(x) - \int_{-\infty}^{\infty} K(x-t)y(t) dt = f(x).$$

Here $-\infty < x < \infty$, $f(x) \in L_1(-\infty, \infty)$, and $K(x) \in L_1(-\infty, \infty)$.

For the integral equation to be solvable (in L_1), it is necessary and sufficient that

$$1 - \sqrt{2\pi} \tilde{K}(u) \neq 0, \quad -\infty < u < \infty, \quad (1)$$

where $\tilde{K}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K(x)e^{-iux} dx$ is the Fourier transform of $K(x)$. In this case, the equation has a unique solution, which is given by

$$y(x) = f(x) + \int_{-\infty}^{\infty} R(x-t)f(t) dt,$$

$$R(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \tilde{R}(u)e^{iux} du, \quad \tilde{R}(u) = \frac{\tilde{K}(u)}{1 - \sqrt{2\pi} \tilde{K}(u)}.$$

⊙ Reference: V. A. Ditkin and A. P. Prudnikov (1965).

$$25. \quad y(x) - \int_0^{\infty} K(x-t)y(t) dt = f(x).$$

*The Wiener-Hopf equation of the second kind.**

Here $0 \leq x < \infty$, $K(x) \in L_1(-\infty, \infty)$, $f(x) \in L_1(0, \infty)$, and $y(x) \in L_1(0, \infty)$.

For the integral equation to be solvable, it is necessary and sufficient that

$$\Omega(u) = 1 - \check{K}(u) \neq 0, \quad -\infty < u < \infty, \quad (1)$$

where $\check{K}(u) = \int_{-\infty}^{\infty} K(x)e^{iux} dx$ is the Fourier transform (in the asymmetric form) of $K(x)$. In this case, the index of the equation can be introduced,

$$\nu = -\text{ind } \Omega(u) = -\frac{1}{2\pi} [\arg \Omega(u)]_{-\infty}^{\infty}.$$

1°. Solution with $\nu = 0$:

$$y(x) = f(x) + \int_0^{\infty} R(x,t)f(t) dt,$$

where

$$R(x,t) = R_+(x-t) + R_-(t-x) + \int_0^{\infty} R_+(x-s)R_-(t-s) ds,$$

and the functions $R_+(x)$ and $R_-(x)$ satisfy the conditions $R_+(x) = 0$ and $R_-(x) = 0$ for $x < 0$ and are uniquely defined by their Fourier transforms as follows:

$$1 + \int_0^{\infty} R_{\pm}(t)e^{\pm iut} dt = \exp \left[-\frac{1}{2} \ln \Omega(u) \mp \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\ln \Omega(t)}{t-u} dt \right].$$

Alternatively, $R_+(x)$ and $R_-(x)$ can be obtained by constructing the solutions of the equations

$$R_+(x) + \int_0^{\infty} K(x-t)R_+(t) dt = K(x), \quad 0 \leq x \leq \infty,$$

$$R_-(x) + \int_0^{\infty} K(t-x)R_-(t) dt = K(-x), \quad 0 \leq x \leq \infty.$$

* A comprehensive discussion of this equation is given in Subsection 13.10-1, Section 13.11, and Section 13.12.

2°. Solution with $\nu > 0$:

$$y(x) = f(x) + \sum_{m=1}^{\nu} C_m x^{m-1} e^{-x} + \int_0^{\infty} R^{\circ}(x, t) \left[f(t) + \sum_{m=1}^{\nu} C_m t^{m-1} e^{-t} \right] dt,$$

where the C_m are arbitrary constants,

$$R^{\circ}(x, t) = R_+^{(0)}(x-t) + R_-^{(1)}(t-x) + \int_0^{\infty} R_+^{(0)}(x-s) R_-^{(1)}(t-s) ds,$$

and the functions $R_+^{(0)}(x)$ and $R_-^{(1)}(x)$ are uniquely defined by their Fourier transforms:

$$1 + \int_0^{\infty} R_{\pm}^{(1)}(t) e^{\pm iut} dt = \left(\frac{u-i}{u+i} \right)^{\nu} \left[1 + \int_0^{\infty} R_{\pm}^{(0)}(t) e^{\pm iut} dt \right],$$

$$1 + \int_0^{\infty} R_{\pm}^{(0)}(t) e^{\pm iut} dt = \exp \left[-\frac{1}{2} \ln \Omega^{\circ}(u) \mp \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\ln \Omega^{\circ}(t)}{t-u} dt \right],$$

$$\Omega^{\circ}(u)(u+i)^{\nu} = \Omega(u)(u-i)^{\nu}.$$

3°. For $\nu < 0$, the solution exists only if the conditions

$$\int_0^{\infty} f(x) \psi_m(x) dx = 0, \quad m = 1, 2, \dots, -\nu,$$

are satisfied. Here $\psi_1(x), \dots, \psi_{-\nu}(x)$ is the system of linearly independent solutions of the transposed homogeneous equation

$$\psi(x) - \int_0^{\infty} K(t-x) \psi(t) dt = 0.$$

Then

$$y(x) = f(x) + \int_0^{\infty} R^*(x, t) f(t) dt,$$

where

$$R^*(x, t) = R_+^{(1)}(x-t) + R_-^{(0)}(t-x) + \int_0^{\infty} R_+^{(1)}(x-s) R_-^{(0)}(t-s) ds,$$

and the functions $R_+^{(1)}(x)$ and $R_-^{(0)}(x)$ are uniquely defined in item 2° by their Fourier transforms.

⊙ References: V. I. Smirnov (1974), F. D. Gakhov and Yu. I. Cherskii (1978), I. M. Vinogradov (1979).

4.9-3. Other Equations of the Form $y(x) + \int_a^b K(x, t)y(t) dt = F(x)$.

26. $y(x) - \int_{-\infty}^{\infty} K(x+t)y(t) dt = f(x).$

The Fourier transform is used to solve this equation.

Solution:

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\tilde{f}(u) + \sqrt{2\pi} \tilde{f}(-u) \tilde{K}(u)}{1 - \sqrt{2\pi} \tilde{K}(u) \tilde{K}(-u)} e^{iux} du,$$

where

$$\tilde{f}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-iux} dx, \quad \tilde{K}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K(x) e^{-iux} dx.$$

⊙ Reference: V. A. Ditkin and A. P. Prudnikov (1965).

$$27. \quad y(x) + \int_{-\infty}^{\infty} e^{\beta t} K(x+t)y(t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = \frac{e^{\lambda x} - k(\lambda)e^{-(\beta+\lambda)x}}{1 - k(\lambda)k(-\beta-\lambda)}, \quad k(\lambda) = \int_{-\infty}^{\infty} K(x)e^{(\lambda+\beta)x} dx.$$

$$28. \quad y(x) + \int_{-\infty}^{\infty} [e^{\beta t} K(x+t) + M(x-t)]y(t) dt = Ae^{\lambda x}.$$

Solution:

$$y(x) = A \frac{I_k(\lambda)e^{px} - [1 + I_m(p)]e^{\lambda x}}{I_k(\lambda)I_k(p) - [1 + I_m(\lambda)][1 + I_m(p)]}, \quad p = -\lambda - \beta,$$

where

$$I_k(\lambda) = \int_{-\infty}^{\infty} K(z)e^{(\beta+\lambda)z} dz, \quad I_m(\lambda) = \int_{-\infty}^{\infty} M(z)e^{-\lambda z} dz.$$

$$29. \quad y(x) - \int_0^{\infty} K(xt)y(t) dt = f(x).$$

The solution can be obtained with the aid of the inverse Mellin transform:

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(s) + \tilde{K}(s)\tilde{f}(1-s)}{1 - \tilde{K}(s)\tilde{K}(1-s)} x^{-s} ds,$$

where \tilde{f} and \tilde{K} stand for the Mellin transforms of the right-hand side and of the kernel of the integral equation,

$$\tilde{f}(s) = \int_0^{\infty} f(x)x^{s-1} dx, \quad \tilde{K}(s) = \int_0^{\infty} K(x)x^{s-1} dx.$$

⊙ Reference: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971).

$$30. \quad y(x) - \int_0^{\infty} K(xt)t^{\beta}y(t) dt = Ax^{\lambda}.$$

Solution:

$$y(x) = A \frac{x^{\lambda} + I_{\beta+\lambda}x^{-\beta-\lambda-1}}{1 - I_{\beta+\lambda}I_{-\lambda-1}}, \quad I_{\mu} = \int_0^{\infty} K(\xi)\xi^{\mu} d\xi.$$

It is assumed that all improper integrals are convergent.

$$31. \quad y(x) - \int_0^{\infty} K(xt)t^{\beta}y(t) dt = f(x).$$

The solution can be obtained with the aid of the inverse Mellin transform as follows:

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(s) + \tilde{K}(s)\tilde{f}(1+\beta-s)}{1 - \tilde{K}(s)\tilde{K}(1+\beta-s)} x^{-s} ds,$$

where \tilde{f} and \tilde{K} stand for the Mellin transforms of the right-hand side and of the kernel of the integral equation,

$$\tilde{f}(s) = \int_0^{\infty} f(x)x^{s-1} dx, \quad \tilde{K}(s) = \int_0^{\infty} K(x)x^{s-1} dx.$$

32. $y(x) - \int_0^\infty g(xt)x^\lambda t^\mu y(t) dt = f(x).$

This equation can be rewritten in the form of equation 4.9.31 by setting $K(z) = z^\lambda g(z)$ and $\beta = \mu - \lambda.$

33. $y(x) - \int_0^\infty \frac{1}{t} K\left(\frac{x}{t}\right) y(t) dt = 0.$

Eigenfunctions of this integral equation are determined by the roots of the following transcendental (algebraic) equation for the parameter $\lambda:$

$$\int_0^\infty K\left(\frac{1}{z}\right) z^{\lambda-1} dz = 1. \tag{1}$$

1°. For a real simple root λ_n of equation (1), there is a corresponding eigenfunction

$$y_n(x) = x^{\lambda_n}.$$

2°. For a real root λ_n of multiplicity $r,$ there are corresponding r eigenfunctions

$$y_{n1}(x) = x^{\lambda_n}, \quad y_{n2}(x) = x^{\lambda_n} \ln x, \quad \dots, \quad y_{nr}(x) = x^{\lambda_n} \ln^{r-1} x.$$

3°. For a complex simple root $\lambda_n = \alpha_n + i\beta_n$ of equation (1), there is a corresponding pair of eigenfunctions

$$y_n^{(1)}(x) = x^{\alpha_n} \cos(\beta_n \ln x), \quad y_n^{(2)}(x) = x^{\alpha_n} \sin(\beta_n \ln x).$$

4°. For a complex root $\lambda_n = \alpha_n + i\beta_n$ of multiplicity $r,$ there are corresponding r eigenfunction pairs

$$\begin{array}{ll} y_{n1}^{(1)}(x) = x^{\alpha_n} \cos(\beta_n \ln x), & y_{n1}^{(2)}(x) = x^{\alpha_n} \sin(\beta_n \ln x), \\ y_{n2}^{(1)}(x) = x^{\alpha_n} \ln x \cos(\beta_n \ln x), & y_{n2}^{(2)}(x) = x^{\alpha_n} \ln x \sin(\beta_n \ln x), \\ \dots & \dots \\ y_{nr}^{(1)}(x) = x^{\alpha_n} \ln^{r-1} x \cos(\beta_n \ln x), & y_{nr}^{(2)}(x) = x^{\alpha_n} \ln^{r-1} x \sin(\beta_n \ln x). \end{array}$$

The general solution is the linear combination (with arbitrary constants) of the eigenfunctions of the homogeneous integral equation.

34. $y(x) - \int_0^\infty \frac{1}{t} K\left(\frac{x}{t}\right) y(t) dt = Ax^b.$

A solution:

$$y(x) = \frac{A}{B} x^b, \quad B = 1 - \int_0^\infty K\left(\frac{1}{\xi}\right) \xi^{b-1} d\xi.$$

It is assumed that the improper integral is convergent and $B \neq 0.$ The general solution of the integral equations is the sum of the above solution and the solution of the homogeneous equation 4.9.33.

$$35. \quad y(x) - \int_0^{\infty} \frac{1}{t} K\left(\frac{x}{t}\right) y(t) dt = f(x).$$

The solution can be obtained with the aid of the inverse Mellin transform:

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(s)}{1 - \tilde{K}(s)} x^{-s} ds,$$

where \tilde{f} and \tilde{K} stand for the Mellin transforms of the right-hand side and the kernel of the integral equation,

$$\tilde{f}(s) = \int_0^{\infty} f(x)x^{s-1} dx, \quad \tilde{K}(s) = \int_0^{\infty} K(x)x^{s-1} dx.$$

Example. For $f(x) = Ae^{-\lambda x}$ and $K(x) = \frac{1}{2}e^{-x}$, the solution of the integral equation has the form

$$y(x) = \begin{cases} \frac{4A}{(3-2C)(\lambda x)^3} & \text{for } \lambda x > 1, \\ -2A \sum_{k=1}^{\infty} \frac{1}{(\lambda x)^{s_k} \psi(s_k)} & \text{for } \lambda x < 1. \end{cases}$$

Here $C = 0.5772\dots$ is the Euler constant, $\psi(z) = [\ln \Gamma(z)]'_z$ is the logarithmic derivative of the gamma function, and the s_k are the negative roots of the transcendental equation $\Gamma(s_k) = 2$, where $\Gamma(z)$ is the gamma function.

⊙ Reference: M. L. Krasnov, A. I. Kisilev, and G. I. Makarenko (1971).

$$36. \quad y(x) + \int_a^b |x-t|g(t)y(t) dt = f(x), \quad a \leq x \leq b.$$

1°. Let us remove the modulus in the integrand,

$$y(x) + \int_a^x (x-t)g(t)y(t) dt + \int_x^b (t-x)g(t)y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x yields

$$y'_x(x) + \int_a^x g(t)y(t) dt - \int_x^b g(t)y(t) dt = f'_x(x). \quad (2)$$

Differentiating (2), we arrive at a second-order ordinary differential equation for $y = y(x)$,

$$y''_{xx} + 2g(x)y = f''_{xx}(x). \quad (3)$$

2°. Let us derive the boundary conditions for equation (3). We assume that the limits of integration satisfy the conditions $-\infty < a < b < \infty$. By setting $x = a$ and $x = b$ in (1), we obtain two consequences

$$\begin{aligned} y(a) + \int_a^b (t-a)g(t)y(t) dt &= f(a), \\ y(b) + \int_a^b (b-t)g(t)y(t) dt &= f(b). \end{aligned} \quad (4)$$

Let us express $g(x)y$ from (3) via y''_{xx} and f''_{xx} and substitute the result into (4). Integrating by parts yields the desired boundary conditions for $y(x)$,

$$\begin{aligned} y(a) + y(b) + (b-a)[f'_x(b) - y'_x(b)] &= f(a) + f(b), \\ y(a) + y(b) + (a-b)[f'_x(a) - y'_x(a)] &= f(a) + f(b). \end{aligned} \quad (5)$$

Note a useful consequence of (5),

$$y'_x(a) + y'_x(b) = f'_x(a) + f'_x(b), \quad (6)$$

which can be used together with one of conditions (5).

Equation (3) under the boundary conditions (5) determines the solution of the original integral equation. Conditions (5) make it possible to calculate the constants of integration that occur in the solution of the differential equation (3).

$$37. \quad y(x) + \int_a^b e^{\lambda|x-t|} g(t)y(t) dt = f(x), \quad a \leq x \leq b.$$

1°. Let us remove the modulus in the integrand:

$$y(x) + \int_a^x e^{\lambda(x-t)} g(t)y(t) dt + \int_x^b e^{\lambda(t-x)} g(t)y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x twice yields

$$y''_{xx}(x) + 2\lambda g(x)y(x) + \lambda^2 \int_a^x e^{\lambda(x-t)} g(t)y(t) dt + \lambda^2 \int_x^b e^{\lambda(t-x)} g(t)y(t) dt = f''_{xx}(x). \quad (2)$$

Eliminating the integral terms from (1) and (2), we arrive at a second-order ordinary differential equation for $y = y(x)$,

$$y''_{xx} + 2\lambda g(x)y - \lambda^2 y = f''_{xx}(x) - \lambda^2 f(x). \quad (3)$$

2°. Let us derive the boundary conditions for equation (3). We assume that the limits of integration satisfy the conditions $-\infty < a < b < \infty$. By setting $x = a$ and $x = b$ in (1), we obtain two consequences

$$\begin{aligned} y(a) + e^{-\lambda a} \int_a^b e^{\lambda t} g(t)y(t) dt &= f(a), \\ y(b) + e^{\lambda b} \int_a^b e^{-\lambda t} g(t)y(t) dt &= f(b). \end{aligned} \quad (4)$$

Let us express $g(x)y$ from (3) via y''_{xx} and f''_{xx} and substitute the result into (4). Integrating by parts yields the conditions

$$\begin{aligned} e^{\lambda b} \varphi'_x(b) - e^{\lambda a} \varphi'_x(a) &= \lambda e^{\lambda a} \varphi(a) + \lambda e^{\lambda b} \varphi(b), \\ e^{-\lambda b} \varphi'_x(b) - e^{-\lambda a} \varphi'_x(a) &= \lambda e^{-\lambda a} \varphi(a) + \lambda e^{-\lambda b} \varphi(b), \end{aligned} \quad \varphi(x) = y(x) - f(x).$$

Finally, after some manipulations, we arrive at the desired boundary conditions for $y(x)$:

$$\varphi'_x(a) + \lambda \varphi(a) = 0, \quad \varphi'_x(b) - \lambda \varphi(b) = 0; \quad \varphi(x) = y(x) - f(x). \quad (5)$$

Equation (3) under the boundary conditions (5) determines the solution of the original integral equation. Conditions (5) make it possible to calculate the constants of integration that occur in solving the differential equation (3).

$$38. \quad y(x) + \int_a^b \sinh(\lambda|x-t|)g(t)y(t) dt = f(x), \quad a \leq x \leq b.$$

1°. Let us remove the modulus in the integrand:

$$y(x) + \int_a^x \sinh[\lambda(x-t)]g(t)y(t) dt + \int_x^b \sinh[\lambda(t-x)]g(t)y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x twice yields

$$\begin{aligned} y''_{xx}(x) + 2\lambda g(x)y(x) + \lambda^2 \int_a^x \sinh[\lambda(x-t)]g(t)y(t) dt \\ + \lambda^2 \int_x^b \sinh[\lambda(t-x)]g(t)y(t) dt = f''_{xx}(x). \end{aligned} \quad (2)$$

Eliminating the integral terms from (1) and (2), we arrive at a second-order ordinary differential equation for $y = y(x)$,

$$y''_{xx} + 2\lambda g(x)y - \lambda^2 y = f''_{xx}(x) - \lambda^2 f(x). \quad (3)$$

2°. Let us derive the boundary conditions for equation (3). We assume that the limits of integration satisfy the conditions $-\infty < a < b < \infty$. By setting $x = a$ and $x = b$ in (1), we obtain two corollaries

$$\begin{aligned} y(a) + \int_a^b \sinh[\lambda(t-a)]g(t)y(t) dt &= f(a), \\ y(b) + \int_a^b \sinh[\lambda(b-t)]g(t)y(t) dt &= f(b). \end{aligned} \quad (4)$$

Let us express $g(x)y$ from (3) via y''_{xx} and f''_{xx} and substitute the result into (4). Integrating by parts yields the desired boundary conditions for $y(x)$,

$$\begin{aligned} \sinh[\lambda(b-a)]\varphi'_x(b) - \lambda \cosh[\lambda(b-a)]\varphi(b) &= \lambda\varphi(a), \\ \sinh[\lambda(b-a)]\varphi'_x(a) + \lambda \cosh[\lambda(b-a)]\varphi(a) &= -\lambda\varphi(b); \quad \varphi(x) = y(x) - f(x). \end{aligned} \quad (5)$$

Equation (3) under the boundary conditions (5) determines the solution of the original integral equation. Conditions (5) make it possible to calculate the constants of integration that occur in solving the differential equation (3).

39. $y(x) + \int_a^b \sin[\lambda|x-t|]g(t)y(t) dt = f(x), \quad a \leq x \leq b.$

1°. Let us remove the modulus in the integrand:

$$y(x) + \int_a^x \sin[\lambda(x-t)]g(t)y(t) dt + \int_x^b \sin[\lambda(t-x)]g(t)y(t) dt = f(x). \quad (1)$$

Differentiating (1) with respect to x twice yields

$$\begin{aligned} y''_{xx}(x) + 2\lambda g(x)y(x) - \lambda^2 \int_a^x \sin[\lambda(x-t)]g(t)y(t) dt \\ - \lambda^2 \int_x^b \sin[\lambda(t-x)]g(t)y(t) dt = f''_{xx}(x). \end{aligned} \quad (2)$$

Eliminating the integral terms from (1) and (2), we arrive at a second-order ordinary differential equation for $y = y(x)$,

$$y''_{xx} + 2\lambda g(x)y + \lambda^2 y = f''_{xx}(x) + \lambda^2 f(x). \quad (3)$$

2°. Let us derive the boundary conditions for equation (3). We assume that the limits of integration satisfy the conditions $-\infty < a < b < \infty$. By setting $x = a$ and $x = b$ in (1), we obtain two consequences

$$\begin{aligned} y(a) + \int_a^b \sin[\lambda(t-a)]g(t)y(t) dt &= f(a), \\ y(b) + \int_a^b \sin[\lambda(b-t)]g(t)y(t) dt &= f(b). \end{aligned} \quad (4)$$

Let us express $g(x)y$ from (3) via y''_{xx} and f''_{xx} and substitute the result into (4). Integrating by parts yields the desired boundary conditions for $y(x)$,

$$\begin{aligned} \sin[\lambda(b-a)]\varphi'_x(b) - \lambda \cos[\lambda(b-a)]\varphi(b) &= \lambda\varphi(a), \\ \sin[\lambda(b-a)]\varphi'_x(a) + \lambda \cos[\lambda(b-a)]\varphi(a) &= -\lambda\varphi(b); \quad \varphi(x) = y(x) - f(x). \end{aligned} \quad (5)$$

Equation (3) under the boundary conditions (5) determines the solution of the original integral equation. Conditions (5) make it possible to calculate the constants of integration that occur in solving the differential equation (3).

$$40. \quad y(x) + \int_{-\infty}^{\infty} [\lambda e^{-|x-t|} + \varphi(x)\psi(t)]y(t) dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 13.2-3; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.2.14.

Solution:

$$y(x) = Y_f(x) + AY_\varphi(x),$$

where

$$Y_f(x) = f(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) f(t) dt,$$

$$Y_\varphi(x) = \varphi(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) \varphi(t) dt,$$

$$A = -\frac{\int_0^{\infty} \psi(t)Y_f(t) dt}{1 + \int_0^{\infty} \psi(t)Y_\varphi(t) dt}, \quad \lambda > -\frac{1}{2}.$$

$$41. \quad y(x) - \int_0^{\infty} [\lambda \sin(xt) + \varphi(x)\psi(t)]y(t) dt = f(x).$$

The solution can be obtained by the methods described in Subsection 13.2-3; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.5.20.

Solution:

$$y(x) = Y_f(x) + AY_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt)f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt)\varphi(t) dt,$$

$$A = \frac{\int_0^{\infty} \psi(t)Y_f(t) dt}{1 - \int_0^{\infty} \psi(t)Y_\varphi(t) dt}, \quad \lambda \neq \pm\sqrt{\frac{2}{\pi}}.$$

$$42. \quad y(x) - \int_0^{\infty} [\lambda \cos(xt) + \varphi(x)\psi(t)]y(t) dt = f(x).$$

The solution can be obtained by the methods described in Subsection 13.2-3; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.5.6.

Solution:

$$y(x) = Y_f(x) + AY_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \cos(xt)f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \cos(xt)\varphi(t) dt,$$

$$A = \frac{\int_0^{\infty} \psi(t)Y_f(t) dt}{1 - \int_0^{\infty} \psi(t)Y_\varphi(t) dt}, \quad \lambda \neq \pm\sqrt{\frac{2}{\pi}}.$$

43. $y(x) + \int_0^\infty [\lambda t J_\nu(xt) + \varphi(x)\psi(t)]y(t) dt = f(x), \quad \nu > -1.$

Here $J_\nu(z)$ is the Bessel function of the first kind. The solution can be obtained by the methods described in Subsection 13.2-3; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.8.4.

Solution:

$$y(x) = Y_f(x) + AY_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1-\lambda^2} - \frac{\lambda}{1-\lambda^2} \int_0^\infty t J_\nu(xt) f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1-\lambda^2} - \frac{\lambda}{1-\lambda^2} \int_0^\infty t J_\nu(xt) \varphi(t) dt,$$

$$A = -\frac{\int_0^\infty \psi(t) Y_f(t) dt}{1 + \int_0^\infty \psi(t) Y_\varphi(t) dt}, \quad \lambda \neq \pm 1.$$

4.9-4. Equations of the Form $y(x) + \int_a^b K(x, t)y(\cdot \cdot) dt = F(x).$

44. $y(x) + \int_a^b f(t)y(x-t) dt = 0.$

Eigenfunctions of this integral equation* are determined by the roots of the following characteristic (transcendental or algebraic) equation for μ :

$$\int_a^b f(t) \exp(-\mu t) dt = -1. \tag{1}$$

1°. For a real (simple) root μ_k of equation (1), there is a corresponding eigenfunction

$$y_k(x) = \exp(\mu_k x).$$

2°. For a real root μ_k of multiplicity r , there are corresponding r eigenfunctions

$$y_{k1}(x) = \exp(\mu_k x), \quad y_{k2}(x) = x \exp(\mu_k x), \quad \dots, \quad y_{kr}(x) = x^{r-1} \exp(\mu_k x).$$

3°. For a complex (simple) root $\mu_k = \alpha_k + i\beta_k$ of equation (1), there is a corresponding pair of eigenfunctions

$$y_k^{(1)}(x) = \exp(\alpha_k x) \cos(\beta_k x), \quad y_k^{(2)}(x) = \exp(\alpha_k x) \sin(\beta_k x).$$

4°. For a complex root $\mu_k = \alpha_k + i\beta_k$ of multiplicity r , there are corresponding r pairs of eigenfunctions

$$\begin{aligned} y_{k1}^{(1)}(x) &= \exp(\alpha_k x) \cos(\beta_k x), & y_{k1}^{(2)}(x) &= \exp(\alpha_k x) \sin(\beta_k x), \\ y_{k2}^{(1)}(x) &= x \exp(\alpha_k x) \cos(\beta_k x), & y_{k2}^{(2)}(x) &= x \exp(\alpha_k x) \sin(\beta_k x), \\ &\dots & \dots & \\ y_{kr}^{(1)}(x) &= x^{r-1} \exp(\alpha_k x) \cos(\beta_k x), & y_{kr}^{(2)}(x) &= x^{r-1} \exp(\alpha_k x) \sin(\beta_k x). \end{aligned}$$

The general solution is the linear combination (with arbitrary constants) of the eigenfunctions of the homogeneous integral equation.

* In the equations below that contain $y(x-t)$ in the integrand, the arguments can have, for example, the domain (a) $-\infty < x < \infty, -\infty < t < \infty$ for $a = -\infty$ and $b = \infty$ or (b) $a \leq t \leq b, -\infty \leq x < \infty$, for a and b such that $-\infty < a < b < \infty$. Case (b) is a special case of (a) if $f(t)$ is nonzero only on the interval $a \leq t \leq b$.

► For equations 4.9.45–4.9.50, only particular solutions are given. To obtain the general solution, one must add the particular solution to the general solution of the corresponding homogeneous equation 4.9.44.

$$45. \quad y(x) + \int_a^b f(t)y(x-t) dt = Ax + B.$$

A solution:

$$y(x) = px + q,$$

where the coefficients p and q are given by

$$p = \frac{A}{1 + I_0}, \quad q = \frac{AI_1}{(1 + I_0)^2} + \frac{B}{1 + I_0}, \quad I_0 = \int_a^b f(t) dt, \quad I_1 = \int_a^b t f(t) dt.$$

$$46. \quad y(x) + \int_a^b f(t)y(x-t) dt = Ae^{\lambda x}.$$

A solution:

$$y(x) = \frac{A}{B} e^{\lambda x}, \quad B = 1 + \int_a^b f(t) \exp(-\lambda t) dt.$$

The general solution of the integral equation is the sum of the specified particular solution and the general solution of the homogeneous equation 4.9.44.

$$47. \quad y(x) + \int_a^b f(t)y(x-t) dt = A \sin(\lambda x).$$

A solution:

$$y(x) = \frac{AI_c}{I_c^2 + I_s^2} \sin(\lambda x) + \frac{AI_s}{I_c^2 + I_s^2} \cos(\lambda x),$$

where the coefficients I_c and I_s are given by

$$I_c = 1 + \int_a^b f(t) \cos(\lambda t) dt, \quad I_s = \int_a^b f(t) \sin(\lambda t) dt.$$

$$48. \quad y(x) + \int_a^b f(t)y(x-t) dt = A \cos(\lambda x).$$

A solution:

$$y(x) = -\frac{AI_s}{I_c^2 + I_s^2} \sin(\lambda x) + \frac{AI_c}{I_c^2 + I_s^2} \cos(\lambda x),$$

where the coefficients I_c and I_s are given by

$$I_c = 1 + \int_a^b f(t) \cos(\lambda t) dt, \quad I_s = \int_a^b f(t) \sin(\lambda t) dt.$$

$$49. \quad y(x) + \int_a^b f(t)y(x-t) dt = e^{\mu x}(A \sin \lambda x + B \cos \lambda x).$$

A solution:

$$y(x) = e^{\mu x}(p \sin \lambda x + q \cos \lambda x),$$

where the coefficients p and q are given by

$$p = \frac{AI_c - BI_s}{I_c^2 + I_s^2}, \quad q = \frac{AI_s + BI_c}{I_c^2 + I_s^2},$$

$$I_c = 1 + \int_a^b f(t)e^{-\mu t} \cos(\lambda t) dt, \quad I_s = \int_a^b f(t)e^{-\mu t} \sin(\lambda t) dt.$$

50. $y(x) + \int_a^b f(t)y(x-t) dt = g(x).$

1°. For $g(x) = \sum_{k=1}^n A_k \exp(\lambda_k x)$, the equation has a solution

$$y(x) = \sum_{k=1}^n \frac{A_k}{B_k} \exp(\lambda_k x), \quad B_k = 1 + \int_a^b f(t) \exp(-\lambda_k t) dt.$$

2°. For polynomial right-hand side of the equation, $g(x) = \sum_{k=0}^n A_k x^k$, a solution has the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

3°. For $g(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

6°. For $g(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

7°. For $g(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

8°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \cos(\lambda_k x)$, a solution of the equation has the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

9°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \sin(\lambda_k x)$, a solution of the equation has the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

10°. For $g(x) = \cos(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

11°. For $g(x) = \sin(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

51. $y(x) + \int_a^b f(t)y(x + \beta t) dt = Ax + B.$

A solution:*

$$y(x) = px + q,$$

where

$$p = \frac{A}{1 + I_0}, \quad q = \frac{B}{1 + I_0} - \frac{AI_1\beta}{(1 + I_0)^2}, \quad I_0 = \int_a^b f(t) dt, \quad I_1 = \int_a^b tf(t) dt.$$

52. $y(x) + \int_a^b f(t)y(x + \beta t) dt = Ae^{\lambda x}.$

A solution:

$$y(x) = \frac{A}{B} e^{\lambda x}, \quad B = 1 + \int_a^b f(t) \exp(\lambda \beta t) dt.$$

* In the equations below that contain $y(x + \beta t)$, $\beta > 0$, in the integrand, the arguments can have, for example, the domain (a) $0 \leq x < \infty, 0 \leq t < \infty$ for $a = 0$ and $b = \infty$ or (b) $a \leq t \leq b, 0 \leq x < \infty$ for a and b such that $0 \leq a < b < \infty$. Case (b) is a special case of (a) if $f(t)$ is nonzero only on the interval $a \leq t \leq b$.

$$53. \quad y(x) + \int_a^b f(t)y(x + \beta t) dt = A \sin \lambda x + B \cos \lambda x.$$

A solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x,$$

where the coefficients p and q are given by

$$p = \frac{AI_c + BI_s}{I_c^2 + I_s^2}, \quad q = \frac{BI_c - AI_s}{I_c^2 + I_s^2},$$

$$I_c = 1 + \int_a^b f(t) \cos(\lambda \beta t) dt, \quad I_s = \int_a^b f(t) \sin(\lambda \beta t) dt.$$

$$54. \quad y(x) + \int_a^b f(t)y(x + \beta t) dt = g(x).$$

1°. For $g(x) = \sum_{k=1}^n A_k \exp(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^n \frac{A_k}{B_k} \exp(\lambda_k x), \quad B_k = 1 + \int_a^b f(t) \exp(\beta \lambda_k t) dt.$$

2°. For polynomial right-hand side of the equation, $g(x) = \sum_{k=0}^n A_k x^k$, a solution has the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

3°. For $g(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k x)$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

6°. For $g(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

7°. For $g(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

8°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \cos(\lambda_k x)$, a solution of the equation has the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

9°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \sin(\lambda_k x)$, a solution of the equation has the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

10°. For $g(x) = \cos(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

11°. For $g(x) = \sin(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, a solution of the equation has the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

55. $y(x) + \int_a^b f(t)y(xt) dt = 0.$

Eigenfunctions of this integral equation* are determined by the roots of the following transcendental (or algebraic) equation for λ :

$$\int_a^b f(t)t^\lambda dt = -1. \tag{1}$$

1°. For a real (simple) root λ_k of equation (1), there is a corresponding eigenfunction

$$y_k(x) = x^{\lambda_k}.$$

2°. For a real root λ_k of multiplicity r , there are corresponding r eigenfunctions

$$y_{k1}(x) = x^{\lambda_k}, \quad y_{k2}(x) = x^{\lambda_k} \ln x, \quad \dots, \quad y_{kr}(x) = x^{\lambda_k} \ln^{r-1} x.$$

3°. For a complex (simple) root $\lambda_k = \alpha_k + i\beta_k$ of equation (1), there is a corresponding pair of eigenfunctions

$$y_k^{(1)}(x) = x^{\alpha_k} \cos(\beta_k \ln x), \quad y_k^{(2)}(x) = x^{\alpha_k} \sin(\beta_k \ln x).$$

4°. For a complex root $\lambda_k = \alpha_k + i\beta_k$ of multiplicity r , there are corresponding r pairs of eigenfunctions

$$\begin{aligned} y_{k1}^{(1)}(x) &= x^{\alpha_k} \cos(\beta_k \ln x), & y_{k1}^{(2)}(x) &= x^{\alpha_k} \sin(\beta_k \ln x), \\ y_{k2}^{(1)}(x) &= x^{\alpha_k} \ln x \cos(\beta_k \ln x), & y_{k2}^{(2)}(x) &= x^{\alpha_k} \ln x \sin(\beta_k \ln x), \\ &\dots & &\dots \\ y_{kr}^{(1)}(x) &= x^{\alpha_k} \ln^{r-1} x \cos(\beta_k \ln x), & y_{kr}^{(2)}(x) &= x^{\alpha_k} \ln^{r-1} x \sin(\beta_k \ln x). \end{aligned}$$

The general solution is the linear combination (with arbitrary constants) of the eigenfunctions of the homogeneous integral equation.

► For equations 4.9.56–4.9.62, only particular solutions are given. To obtain the general solution, one must add the particular solution to the general solution of the corresponding homogeneous equation 4.9.55.

56. $y(x) + \int_a^b f(t)y(xt) dt = Ax + B.$

A solution:

$$y(x) = \frac{A}{1 + I_1}x + \frac{B}{1 + I_0}, \quad I_0 = \int_a^b f(t) dt, \quad I_1 = \int_a^b tf(t) dt.$$

57. $y(x) + \int_a^b f(t)y(xt) dt = Ax^\beta.$

A solution:

$$y(x) = \frac{A}{B}x^\beta, \quad B = 1 + \int_a^b f(t)t^\beta dt.$$

* In the equations below that contain $y(xt)$ in the integrand, the arguments can have, for example, the domain (a) $0 \leq x \leq 1$, $0 \leq t \leq 1$ for $a = 0$ and $b = 1$, (b) $1 \leq x < \infty$, $1 \leq t < \infty$ for $a = 1$ and $b = \infty$, (c) $0 \leq x < \infty$, $0 \leq t < \infty$ for $a = 0$ and $b = \infty$, or (d) $a \leq t \leq b$, $0 \leq x < \infty$ for a and b such that $0 \leq a < b \leq \infty$. Case (d) is a special case of (c) if $f(t)$ is nonzero only on the interval $a \leq t \leq b$.

$$58. \quad y(x) + \int_a^b f(t)y(xt) dt = A \ln x + B.$$

A solution:

$$y(x) = p \ln x + q,$$

where

$$p = \frac{A}{1 + I_0}, \quad q = \frac{B}{1 + I_0} - \frac{AI_1}{(1 + I_0)^2}, \quad I_0 = \int_a^b f(t) dt, \quad I_1 = \int_a^b f(t) \ln t dt.$$

$$59. \quad y(x) + \int_a^b f(t)y(xt) dt = Ax^\beta \ln x.$$

A solution:

$$y(x) = px^\beta \ln x + qx^\beta,$$

where

$$p = \frac{A}{1 + I_1}, \quad q = -\frac{AI_2}{(1 + I_1)^2}, \quad I_1 = \int_a^b f(t)t^\beta dt, \quad I_2 = \int_a^b f(t)t^\beta \ln t dt.$$

$$60. \quad y(x) + \int_a^b f(t)y(xt) dt = A \cos(\ln x).$$

A solution:

$$y(x) = \frac{AI_c}{I_c^2 + I_s^2} \cos(\ln x) + \frac{AI_s}{I_c^2 + I_s^2} \sin(\ln x),$$

$$I_c = 1 + \int_a^b f(t) \cos(\ln t) dt, \quad I_s = \int_a^b f(t) \sin(\ln t) dt.$$

$$61. \quad y(x) + \int_a^b f(t)y(xt) dt = A \sin(\ln x).$$

A solution:

$$y(x) = -\frac{AI_s}{I_c^2 + I_s^2} \cos(\ln x) + \frac{AI_c}{I_c^2 + I_s^2} \sin(\ln x),$$

$$I_c = 1 + \int_a^b f(t) \cos(\ln t) dt, \quad I_s = \int_a^b f(t) \sin(\ln t) dt.$$

$$62. \quad y(x) + \int_a^b f(t)y(xt) dt = Ax^\beta \cos(\lambda \ln x) + Bx^\beta \sin(\lambda \ln x).$$

A solution:

$$y(x) = px^\beta \cos(\lambda \ln x) + qx^\beta \sin(\lambda \ln x),$$

where

$$p = \frac{AI_c - BI_s}{I_c^2 + I_s^2}, \quad q = \frac{AI_s + BI_c}{I_c^2 + I_s^2},$$

$$I_c = 1 + \int_a^b f(t)t^\beta \cos(\lambda \ln t) dt, \quad I_s = \int_a^b f(t)t^\beta \sin(\lambda \ln t) dt.$$

63. $y(x) + \int_a^b f(t)y(\xi) dt = 0, \quad \xi = x\varphi(t).$

Eigenfunctions of this integral equation are determined by the roots of the following transcendental (or algebraic) equation for λ :

$$\int_a^b f(t)[\varphi(t)]^\lambda dt = -1. \tag{1}$$

1°. For a real (simple) root λ_k of equation (1), there is a corresponding eigenfunction

$$y_k(x) = x^{\lambda_k}.$$

2°. For a real root λ_k of multiplicity r , there are corresponding r eigenfunctions

$$y_{k1}(x) = x^{\lambda_k}, \quad y_{k2}(x) = x^{\lambda_k} \ln x, \quad \dots, \quad y_{kr}(x) = x^{\lambda_k} \ln^{r-1} x.$$

3°. For a complex (simple) root $\lambda_k = \alpha_k + i\beta_k$ of equation (1), there is a corresponding pair of eigenfunctions

$$y_k^{(1)}(x) = x^{\alpha_k} \cos(\beta_k \ln x), \quad y_k^{(2)}(x) = x^{\alpha_k} \sin(\beta_k \ln x).$$

4°. For a complex root $\lambda_k = \alpha_k + i\beta_k$ of multiplicity r , there are corresponding r pairs of eigenfunctions

$$\begin{aligned} y_{k1}^{(1)}(x) &= x^{\alpha_k} \cos(\beta_k \ln x), & y_{k1}^{(2)}(x) &= x^{\alpha_k} \sin(\beta_k \ln x), \\ y_{k2}^{(1)}(x) &= x^{\alpha_k} \ln x \cos(\beta_k \ln x), & y_{k2}^{(2)}(x) &= x^{\alpha_k} \ln x \sin(\beta_k \ln x), \\ &\dots & &\dots \\ y_{kr}^{(1)}(x) &= x^{\alpha_k} \ln^{r-1} x \cos(\beta_k \ln x), & y_{kr}^{(2)}(x) &= x^{\alpha_k} \ln^{r-1} x \sin(\beta_k \ln x). \end{aligned}$$

The general solution is the linear combination (with arbitrary constants) of the eigenfunctions of the homogeneous integral equation.

64. $y(x) + \int_a^b f(t)y(\xi) dt = Ax^\beta, \quad \xi = x\varphi(t).$

A solution:

$$y(x) = \frac{A}{B}x^\beta, \quad B = 1 + \int_a^b f(t)[\varphi(t)]^\beta dt.$$

It is assumed that $B \neq 0$. A linear combination of eigenfunctions of the corresponding homogeneous equation (see 4.9.63) can be added to this solution.

65. $y(x) + \int_a^b f(t)y(\xi) dt = g(x), \quad \xi = x\varphi(t).$

1°. For $g(x) = \sum_{k=0}^n A_k x^k$, a solution of the equation has the form

$$y(x) = \sum_{k=0}^n \frac{A_k}{B_k} x^k, \quad B_k = 1 + \int_a^b f(t)[\varphi(t)]^k dt. \tag{1}$$

2°. For $g(x) = \ln x \sum_{k=0}^n A_k x^k$, a solution has the form

$$y(x) = \ln x \sum_{k=0}^n B_k x^k + \sum_{k=0}^n C_k x^k, \tag{2}$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

3°. For $g(x) = \sum_{k=0}^n A_k (\ln x)^k$, a solution of the equation has the form

$$y(x) = \sum_{k=0}^n B_k (\ln x)^k, \quad (3)$$

where the constants B_k can be found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k \ln x)$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k \ln x) + \sum_{k=1}^n C_k \sin(\lambda_k \ln x), \quad (4)$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k \ln x)$, a solution of the equation has the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k \ln x) + \sum_{k=1}^n C_k \sin(\lambda_k \ln x), \quad (5)$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

Remark. A linear combination of eigenfunctions of the corresponding homogeneous equation (see 4.9.63) can be added to solutions (1)–(5).

4.10. Some Formulas and Transformations

Let the solution of the integral equation

$$y(x) + \int_a^b K(x, t)y(t) dt = f(x) \quad (1)$$

have the form

$$y(x) = f(x) + \int_a^b R(x, t)f(t) dt. \quad (2)$$

Then the solution of the more complicated integral equation

$$y(x) + \int_a^b K(x, t) \frac{g(x)}{g(t)} y(t) dt = f(x) \quad (3)$$

has the form

$$y(x) = f(x) + \int_a^b R(x, t) \frac{g(x)}{g(t)} f(t) dt. \quad (4)$$

Below are formulas for the solutions of integral equations of the form (3) for some specific functions $g(x)$. In all cases, it is assumed that the solution of equation (1) is known and is given by (2).

1°. The solution of the equation

$$y(x) + \int_a^b K(x, t)(x/t)^\lambda y(t) dt = f(x)$$

has the form

$$y(x) = f(x) + \int_a^b R(x, t)(x/t)^\lambda f(t) dt.$$

2°. The solution of the equation

$$y(x) + \int_a^b K(x, t)e^{\lambda(x-t)}y(t) dt = f(x)$$

has the form

$$y(x) = f(x) + \int_a^b R(x, t)e^{\lambda(x-t)}f(t) dt.$$

Chapter 5

Nonlinear Equations of the First Kind with Variable Limit of Integration

► **Notation:** $f, g, h,$ and K are arbitrary functions of an argument specified in the parentheses (the argument can depend on $t, x,$ and y); $A, B, a, b, k, \beta, \lambda,$ and μ are arbitrary parameters.

5.1. Equations with Quadratic Nonlinearity That Contain Arbitrary Parameters

5.1-1. Equations of the Form $\int_0^x y(t)y(x-t) dt = f(x).$

1.
$$\int_0^x y(t)y(x-t) dt = Ax + B, \quad A, B > 0.$$

Solutions:

$$y(x) = \pm\sqrt{B} \left[\frac{1}{\sqrt{\pi x}} \exp\left(-\frac{A}{B}x\right) + \sqrt{\frac{A}{B}} \operatorname{erf}\left(\sqrt{\frac{A}{B}}x\right) \right],$$

where $\operatorname{erf} z = \frac{2}{\sqrt{\pi}} \int_0^z \exp(-t^2) dt$ is the error function.

2.
$$\int_0^x y(t)y(x-t) dt = A^2 x^\lambda.$$

Solutions:

$$y(x) = \pm A \frac{\sqrt{\Gamma(\lambda+1)}}{\Gamma\left(\frac{\lambda+1}{2}\right)} x^{\frac{\lambda-1}{2}},$$

where $\Gamma(z)$ is the gamma function.

3.
$$\int_0^x y(t)y(x-t) dt = Ax^{\lambda-1} + Bx^\lambda, \quad \lambda > 0.$$

Solutions:

$$y(x) = \pm \frac{\sqrt{A\Gamma(\lambda)}}{\Gamma(\lambda/2)} x^{\frac{\lambda-2}{2}} \exp\left(-\lambda \frac{B}{A}x\right) \Phi\left(\frac{\lambda+1}{2}, \frac{\lambda}{2}; \lambda \frac{B}{A}x\right),$$

where $\Phi(a, c; x)$ is the degenerate hypergeometric function (Kummer's function).

4.
$$\int_0^x y(t)y(x-t) dt = A^2 e^{\lambda x}.$$

Solutions: $y(x) = \pm \frac{A}{\sqrt{\pi x}} e^{\lambda x}.$

$$5. \quad \int_0^x y(t)y(x-t) dt = (Ax + B)e^{\lambda x}, \quad A, B > 0.$$

Solutions:

$$y(x) = \pm \sqrt{B} e^{\lambda x} \left[\frac{1}{\sqrt{\pi x}} \exp\left(-\frac{A}{B}x\right) + \sqrt{\frac{A}{B}} \operatorname{erf}\left(\sqrt{\frac{A}{B}}x\right) \right],$$

where $\operatorname{erf} z = \frac{2}{\sqrt{\pi}} \int_0^z \exp(-t^2) dt$ is the error function.

$$6. \quad \int_0^x y(t)y(x-t) dt = A^2 x^\mu e^{\lambda x}.$$

Solutions:

$$y(x) = \pm \frac{A\sqrt{\Gamma(\mu+1)}}{\Gamma(\frac{\mu+1}{2})} x^{\frac{\mu-1}{2}} e^{\lambda x}.$$

$$7. \quad \int_0^x y(t)y(x-t) dt = (Ax^{\mu-1} + Bx^\mu)e^{\lambda x}.$$

Solutions:

$$y(x) = \pm \frac{\sqrt{A\Gamma(\mu)}}{\Gamma(\mu/2)} x^{\frac{\mu-2}{2}} \exp\left[\left(\lambda - \mu \frac{B}{A}\right)x\right] \Phi\left(\frac{\mu+1}{2}, \frac{\mu}{2}; \mu \frac{B}{A}x\right),$$

where $\Phi(a, c; x)$ is the degenerate hypergeometric function (Kummer's function).

$$8. \quad \int_0^x y(t)y(x-t) dt = A^2 \cosh(\lambda x).$$

Solutions: $y(x) = \pm \frac{A}{\sqrt{\pi}} \frac{d}{dx} \int_0^x \frac{I_0(\lambda t) dt}{\sqrt{x-t}}$, where $I_0(z)$ is the modified Bessel function.

$$9. \quad \int_0^x y(t)y(x-t) dt = A \sinh(\lambda x).$$

Solutions: $y = \pm \sqrt{A\lambda} I_0(\lambda x)$, where $I_0(z)$ is the modified Bessel function.

$$10. \quad \int_0^x y(t)y(x-t) dt = A \sinh(\lambda \sqrt{x}).$$

Solutions: $y = \pm \sqrt{A} \pi^{1/4} 2^{-7/8} \lambda^{3/4} x^{-1/8} I_{-1/4}\left(\lambda \sqrt{\frac{1}{2}x}\right)$, where $I_{-1/4}(z)$ is the modified Bessel function.

$$11. \quad \int_0^x y(t)y(x-t) dt = A^2 \cos(\lambda x).$$

Solutions: $y(x) = \pm \frac{A}{\sqrt{\pi}} \frac{d}{dx} \int_0^x \frac{J_0(\lambda t) dt}{\sqrt{x-t}}$, where $J_0(z)$ is the Bessel function.

$$12. \quad \int_0^x y(t)y(x-t) dt = A \sin(\lambda x).$$

Solutions: $y = \pm \sqrt{A\lambda} J_0(\lambda x)$, where $J_0(z)$ is the Bessel function.

$$13. \int_0^x y(t)y(x-t) dt = A \sin(\lambda\sqrt{x}).$$

Solutions: $y = \pm\sqrt{A} \pi^{1/4} 2^{-7/8} \lambda^{3/4} x^{-1/8} J_{-1/4}\left(\lambda\sqrt{\frac{1}{2}x}\right)$, where $J_{-1/4}(z)$ is the Bessel function.

$$14. \int_0^x y(t)y(x-t) dt = A^2 e^{\mu x} \cosh(\lambda x).$$

Solutions: $y(x) = \pm \frac{A}{\sqrt{\pi}} e^{\mu x} \frac{d}{dx} \int_0^x \frac{I_0(\lambda t) dt}{\sqrt{x-t}}$, where $I_0(z)$ is the modified Bessel function.

$$15. \int_0^x y(t)y(x-t) dt = A e^{\mu x} \sinh(\lambda x).$$

Solutions: $y = \pm\sqrt{A\lambda} e^{\mu x} I_0(\lambda x)$, where $I_0(z)$ is the modified Bessel function.

$$16. \int_0^x y(t)y(x-t) dt = A^2 e^{\mu x} \cos(\lambda x).$$

Solutions: $y(x) = \pm \frac{A}{\sqrt{\pi}} e^{\mu x} \frac{d}{dx} \int_0^x \frac{J_0(\lambda t) dt}{\sqrt{x-t}}$, where $J_0(z)$ is the Bessel function.

$$17. \int_0^x y(t)y(x-t) dt = A e^{\mu x} \sin(\lambda x).$$

Solutions: $y = \pm\sqrt{A\lambda} e^{\mu x} J_0(\lambda x)$, where $J_0(z)$ is the Bessel function.

5.1-2. Equations of the Form $\int_0^x K(x,t)y(t)y(x-t) dt = f(x)$.

$$18. \int_0^x t^k y(t)y(x-t) dt = A x^\lambda, \quad A > 0.$$

Solutions:

$$y(x) = \pm \left[\frac{A\Gamma(\lambda+1)}{\Gamma\left(\frac{\lambda+1+k}{2}\right)\Gamma\left(\frac{\lambda+1-k}{2}\right)} \right]^{1/2} x^{\frac{\lambda-k-1}{2}},$$

where $\Gamma(z)$ is the gamma function.

$$19. \int_0^x t^k y(t)y(x-t) dt = A e^{\lambda x}.$$

Solutions:

$$y(x) = \pm \left[\frac{A}{\Gamma\left(\frac{k+1}{2}\right)\Gamma\left(\frac{1-k}{2}\right)} \right]^{1/2} x^{-\frac{k+1}{2}} e^{\lambda x},$$

where $\Gamma(z)$ is the gamma function.

$$20. \int_0^x t^k y(t)y(x-t) dt = A x^\mu e^{\lambda x}.$$

Solutions:

$$y(x) = \pm \left[\frac{A\Gamma(\mu+1)}{\Gamma\left(\frac{\mu+k+1}{2}\right)\Gamma\left(\frac{\mu-k+1}{2}\right)} \right]^{1/2} x^{\frac{\mu-k-1}{2}} e^{\lambda x},$$

where $\Gamma(z)$ is the gamma function.

$$21. \int_0^x \frac{y(t)y(x-t)}{ax+bt} dt = Ax^\lambda.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\lambda/2}, \quad I = \int_0^1 z^{\lambda/2}(1-z)^{\lambda/2} \frac{dz}{a+bz}.$$

$$22. \int_0^x \frac{y(t)y(x-t)}{ax+bt} dt = Ae^{\lambda x}.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} e^{\lambda x}, \quad I = \frac{1}{b} \ln\left(1 + \frac{b}{a}\right).$$

$$23. \int_0^x \frac{y(t)y(x-t)}{ax+bt} dt = Ax^\mu e^{\lambda x}.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\mu/2} e^{\lambda x}, \quad I = \int_0^1 z^{\mu/2}(1-z)^{\mu/2} \frac{dz}{a+bz}.$$

$$24. \int_0^x \frac{y(t)y(x-t)}{\sqrt{ax^2+bt^2}} dt = Ax^\lambda.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\lambda/2}, \quad I = \int_0^1 z^{\lambda/2}(1-z)^{\lambda/2} \frac{dz}{\sqrt{a+bz^2}}.$$

$$25. \int_0^x \frac{y(t)y(x-t)}{\sqrt{ax^2+bt^2}} dt = Ae^{\lambda x}.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} e^{\lambda x}, \quad I = \int_0^1 \frac{dz}{\sqrt{a+bz^2}}.$$

$$26. \int_0^x \frac{y(t)y(x-t)}{\sqrt{ax^2+bt^2}} dt = Ax^\mu e^{\lambda x}.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\mu/2} e^{\lambda x}, \quad I = \int_0^1 z^{\mu/2}(1-z)^{\mu/2} \frac{dz}{\sqrt{a+bz^2}}.$$

5.1-3. Equations of the Form $\int_0^x y(t)y(\dots) dt = f(x)$.

$$27. \int_0^x y(t)y(ax+bt) dt = Ax^\lambda.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\frac{\lambda-1}{2}}, \quad I = \int_0^1 z^{\frac{\lambda-1}{2}} (a+bz)^{\frac{\lambda-1}{2}} dz.$$

$$28. \int_0^x y(t)y(ax-t) dt = Ae^{\lambda x}, \quad a \geq 1.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} \frac{\exp(\lambda x/a)}{\sqrt{x}}, \quad I = \int_0^1 \frac{dz}{\sqrt{z(a-z)}}.$$

$$29. \int_0^x y(t)y(ax-t) dt = Ax^\mu e^{\lambda x}, \quad a \geq 1.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\frac{\mu-1}{2}} \exp(\lambda x/a), \quad I = \int_0^1 z^{\frac{\mu-1}{2}} (a-z)^{\frac{\mu-1}{2}} dz.$$

$$30. \int_0^x y(t)y(xt) dt = Ax^\mu.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{1}{3} A(2\mu+1)} x^{\frac{\mu-1}{3}} \quad (A > 0, \mu \geq 0).$$

5.2. Equations with Quadratic Nonlinearity That Contain Arbitrary Functions

5.2-1. Equations of the Form $\int_a^x K(x,t)[Ay(t) + By^2(t)] dt = f(x)$.

$$1. \int_a^x (x-t)[Ay(t) + By^2(t)] dt = f(x), \quad f(a) = f'(a) = 0.$$

Solution in implicit form:

$$Ay + By^2 - f''_{xx}(x) = 0.$$

$$2. \int_a^x (x-t)^n [Ay(t) + By^2(t)] dt = f(x), \quad f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0.$$

Here $n = 1, 2, \dots$. Solution in implicit form:

$$n! (Ay + By^2) - f_x^{(n+1)}(x) = 0.$$

$$3. \int_a^x e^{\lambda(x-t)} [Ay(t) + By^2(t)] dt = f(x), \quad f(a) = 0.$$

Solution in implicit form:

$$Ay + By^2 + \lambda f(x) - f'_x(x) = 0.$$

$$4. \int_a^x \sinh[\lambda(x-t)] [Ay(t) + By^2(t)] dt = f(x), \quad f(a) = f'(a) = 0.$$

Solution in implicit form:

$$\lambda(Ay + By^2) + \lambda^2 f(x) - f''_{xx}(x) = 0.$$

$$5. \int_a^x \cosh[\lambda(x-t)][Ay(t) + By^2(t)] dt = f(x), \quad f(a) = 0.$$

Solution in implicit form:

$$Ay + By^2 + \lambda^2 \int_a^x f(t) dt - f'_x(x) = 0.$$

$$6. \int_a^x \sin[\lambda(x-t)][Ay(t) + By^2(t)] dt = f(x), \quad f(a) = f'(a) = 0.$$

Solution in implicit form:

$$\lambda(Ay + By^2) - \lambda^2 f(x) - f''_{xx}(x) = 0.$$

$$7. \int_a^x \cos[\lambda(x-t)][Ay(t) + By^2(t)] dt = f(x).$$

Solution in implicit form:

$$Ay + By^2 - \lambda^2 \int_a^x f(t) dt - f'_x(x) = 0.$$

$$8. \int_a^x [g(x) - g(t)][Ay(t) + By^2(t)] dt = f(x).$$

It is assumed that $f(a) = f'_x(a) = 0$ and $f'_x/g'_x \neq \text{const.}$

Solution in implicit form:

$$Ay + By^2 = \frac{d}{dx} \left[\frac{f'_x(x)}{g'_x(x)} \right].$$

$$9. \int_a^x K(x, t)[Ay(t) + By^2(t)] dt = f(x).$$

The substitution $w(t) = Ay(t) + By^2(t)$ leads to the linear integral equation of the first kind

$$\int_a^x K(x, t)w(t) dt = f(x).$$

For the exact solutions of the equation with various $K(x, t)$ and $f(x)$, see Chapter 1.

5.2-2. Equations of the Form $\int_a^x K(x, t)y(t)y(ax + bt) dt = f(x)$.

$$10. \int_a^x K(t)y(x)y(t) dt = f(x).$$

Solutions:

$$y(x) = \pm f(x) \left[2 \int_a^x K(t)f(t) dt \right]^{-1/2}.$$

$$11. \int_0^x f\left(\frac{t}{x}\right)y(t)y(x-t) dt = Ax^\lambda.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\frac{\lambda-1}{2}}, \quad I = \int_0^1 f(z)z^{\frac{\lambda-1}{2}}(1-z)^{\frac{\lambda-1}{2}} dz.$$

$$12. \int_0^x f\left(\frac{t}{x}\right) y(t)y(x-t) dt = Ae^{\lambda x}.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} \frac{e^{\lambda x}}{\sqrt{x}}, \quad I = \int_0^1 \frac{f(z) dz}{\sqrt{z(1-z)}}.$$

$$13. \int_0^x f\left(\frac{t}{x}\right) y(t)y(x-t) dt = Ax^\mu e^{\lambda x}.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\frac{\mu-1}{2}} e^{\lambda x}, \quad I = \int_0^1 f(z) z^{\frac{\mu-1}{2}} (1-z)^{\frac{\mu-1}{2}} dz.$$

$$14. \int_0^x f\left(\frac{t}{x}\right) y(t)y(ax+bt) dt = Ax^\lambda.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\frac{\lambda-1}{2}}, \quad I = \int_0^1 f(z) z^{\frac{\lambda-1}{2}} (a+bz)^{\frac{\lambda-1}{2}} dz.$$

$$15. \int_0^x f\left(\frac{t}{x}\right) y(t)y(ax-t) dt = Ae^{\lambda x}, \quad a \geq 1.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} \frac{\exp(\lambda x/a)}{\sqrt{x}}, \quad I = \int_0^1 \frac{f(z) dz}{\sqrt{z(a-z)}}.$$

$$16. \int_0^x f\left(\frac{t}{x}\right) y(t)y(ax-t) dt = Ax^\mu e^{\lambda x}, \quad a \geq 1.$$

Solutions:

$$y(x) = \pm \sqrt{\frac{A}{I}} x^{\frac{\mu-1}{2}} \exp(\lambda x/a), \quad I = \int_0^1 f(z) z^{\frac{\mu-1}{2}} (a-z)^{\frac{\mu-1}{2}} dz.$$

5.3. Equations with Nonlinearity of General Form

5.3-1. Equations of the Form $\int_a^x K(x,t)f(t,y(t)) dt = g(x)$.

$$1. \int_a^x f(t,y(t)) dt = g(x), \quad g(a) = 0.$$

Solution in implicit form:

$$f(x,y) - g'_x(x) = 0.$$

$$2. \int_a^x (x-t)f(t,y(t)) dt = g(x), \quad g(a) = g'(a) = 0.$$

Solution in implicit form:

$$f(x,y) - g''_{xx}(x) = 0.$$

$$3. \quad \int_a^x (x-t)^n f(t, y(t)) dt = g(x), \quad g(a) = g'_x(a) = \cdots = g_x^{(n)}(a) = 0.$$

Here $n = 1, 2, \dots$. Solution in implicit form:

$$n! f(x, y) - g_x^{(n+1)}(x) = 0.$$

$$4. \quad \int_a^x e^{\lambda(x-t)} f(t, y(t)) dt = g(x), \quad g(a) = 0.$$

Solution in implicit form:

$$f(x, y) + \lambda g(x) - g'_x(x) = 0.$$

$$5. \quad \int_a^x \sinh[\lambda(x-t)] f(t, y(t)) dt = g(x), \quad g(a) = g'(a) = 0.$$

Solution in implicit form:

$$\lambda f(x, y) + \lambda^2 g(x) - g''_{xx}(x) = 0.$$

$$6. \quad \int_a^x \cosh[\lambda(x-t)] f(t, y(t)) dt = g(x), \quad g(a) = 0.$$

Solution in implicit form:

$$f(x, y) + \lambda^2 \int_a^x g(t) dt - g'_x(x) = 0.$$

$$7. \quad \int_a^x \sin[\lambda(x-t)] f(t, y(t)) dt = g(x), \quad g(a) = g'(a) = 0.$$

Solution in implicit form:

$$\lambda f(x, y) - \lambda^2 g(x) - g''_{xx}(x) = 0.$$

$$8. \quad \int_a^x \cos[\lambda(x-t)] f(t, y(t)) dt = g(x).$$

Solution in implicit form:

$$f(x, y) - \lambda^2 \int_a^x g(t) dt - g'_x(x) = 0.$$

$$9. \quad \int_a^x [h(x) - h(t)] f(t, y(t)) dt = g(x).$$

It is assumed that $g(a) = g'_x(a) = 0$ and $g'_x/h'_x \neq \text{const.}$

Solution in implicit form:

$$f(x, y) = \frac{d}{dx} \left[\frac{g'_x(x)}{h'_x(x)} \right].$$

$$10. \quad \int_a^x K(x, t) f(t, y(t)) dt = g(x).$$

The substitution $w(t) = f(t, y(t))$ leads to the linear integral equation of the first kind

$$\int_a^x K(x, t) w(t) dt = g(x).$$

For the exact solutions of the equation with various $K(x, t)$ and $g(x)$, see Chapter 1.

5.3-2. Other Equations.

$$11. \int_0^x f\left(\frac{t}{x}, y(t), y(x)\right) dt = Ax.$$

A solution: $y(x) = \lambda$, where λ is a root of the algebraic (or transcendental) equation

$$F(\lambda) - A = 0, \quad F(\lambda) = \int_0^1 f(z, \lambda, \lambda) dz.$$

$$12. \int_0^x f\left(\frac{t}{x}, \frac{y(t)}{y(x)}\right) dt = Ax.$$

A solution: $y(x) = Cx^\lambda$, where C is an arbitrary constant and λ is a root of the algebraic (or transcendental) equation

$$F(\lambda) - A = 0, \quad F(\lambda) = \int_0^1 f(z, z^\lambda) dz.$$

$$13. \int_0^x f\left(\frac{t}{x}, \frac{y(t)}{y(x)}\right) y^\alpha(t) dt = Ax^\beta, \quad \alpha \neq 0.$$

A solution:

$$y(x) = A^{1/\alpha} x^\lambda, \quad \lambda = \frac{\beta - 1}{\alpha},$$

where λ is a root of the algebraic (or transcendental) equation

$$F(\lambda) - 1 = 0, \quad F(\lambda) = \int_0^1 f(z, z^\lambda) z^{\alpha\lambda} dz.$$

$$14. \int_0^x f\left(\frac{t}{x}, \frac{y(t)}{t}, \frac{y(x)}{x}\right) dt = Ax.$$

A solution: $y(x) = \lambda x$, where λ is a root of the algebraic (or transcendental) equation

$$F(\lambda) - A = 0, \quad F(\lambda) = \int_0^1 f(z, \lambda, \lambda) dz.$$

$$15. \int_x^\infty f(t-x, y(t-x)) y(t) dt = Ae^{-\lambda x}.$$

Solutions: $y(x) = b_k e^{-\lambda x}$, where b_k are roots of the algebraic (or transcendental) equation

$$bI(b) = A, \quad I(b) = \int_0^\infty f(z, be^{-\lambda z}) e^{-\lambda z} dz.$$

Chapter 6

Nonlinear Equations of the Second Kind with Variable Limit of Integration

► **Notation:** f , g , and h are arbitrary functions of an argument specified in the parentheses (the argument can depend on t , x , and y); A , B , C , a , b , k , β , λ , and μ are arbitrary parameters.

6.1. Equations with Quadratic Nonlinearity That Contain Arbitrary Parameters

6.1-1. Equations of the Form $y(x) + \int_a^x K(x, t)y^2(t) dt = F(x)$.

1. $y(x) + A \int_a^x y^2(t) dt = Bx + C.$

By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

1°. Solution with $AB > 0$:

$$y(x) = k \frac{(k + y_a) \exp[2Ak(x - a)] + y_a - k}{(k + y_a) \exp[2Ak(x - a)] - y_a + k}, \quad k = \sqrt{\frac{B}{A}}, \quad y_a = aB + C.$$

2°. Solution with $AB < 0$:

$$y(x) = k \tan \left[Ak(a - x) + \arctan \frac{y_a}{k} \right], \quad k = \sqrt{-\frac{B}{A}}, \quad y_a = aB + C.$$

3°. Solution with $B = 0$:

$$y(x) = \frac{C}{AC(x - a) + 1}.$$

2. $y(x) + k \int_a^x (x - t)y^2(t) dt = Ax^2 + Bx + C.$

This is a special case of equation 6.8.3 with $f(y) = ky^2$.

Solution in an implicit form:

$$\int_{y_0}^y [4Au - 2kF(u) + B^2 - 4AC]^{-1/2} du = \pm(x - a),$$
$$F(u) = \frac{1}{3}(u^3 - y_0^3), \quad y_0 = Aa^2 + Ba + C.$$

$$3. \quad y(x) + A \int_a^x t^\lambda y^2(t) dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = Ay^2$. By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

Solution in an implicit form:

$$(\lambda + 1) \int_{y_a}^y \frac{du}{Au^2 - B(\lambda + 1)} + x^{\lambda+1} - a^{\lambda+1} = 0, \quad y_a = Ba^{\lambda+1} + C.$$

$$4. \quad y(x) + A \int_0^x x^{-\lambda-1} y^2(t) dt = Bx^\lambda, \quad \lambda > -\frac{1}{2}.$$

Solutions: $y_1(x) = \beta_1 x^\lambda$ and $y_2(x) = \beta_2 x^\lambda$, where $\beta_{1,2}$ are the roots of the quadratic equation $A\beta^2 + (2\lambda + 1)\beta - B(2\lambda + 1) = 0$.

$$5. \quad y(x) + \int_0^x \frac{y^2(t) dt}{ax + bt} = A.$$

Solutions: $y_1(x) = \lambda_1$ and $y_2(x) = \lambda_2$, where $\lambda_{1,2}$ are the roots of the quadratic equation $\ln\left(1 + \frac{b}{a}\right)\lambda^2 + b\lambda - Ab = 0$.

$$6. \quad y(x) + A \int_0^x \frac{y^2(t) dt}{x^2 + t^2} = Bx.$$

Solutions: $y_1(x) = \lambda_1 x$ and $y_2(x) = \lambda_2 x$, where $\lambda_{1,2}$ are the roots of the quadratic equation $(1 - \frac{1}{4}\pi)A\lambda^2 + \lambda - B = 0$.

$$7. \quad y(x) + \int_0^x \frac{y^2(t) dt}{\sqrt{ax^2 + bt^2}} = A.$$

Solutions: $y_1(x) = \lambda_1$ and $y_2(x) = \lambda_2$, where $\lambda_{1,2}$ are the roots of the quadratic equation

$$I\lambda^2 + \lambda - A = 0, \quad I = \int_0^1 \frac{dz}{\sqrt{a + bz^2}}.$$

$$8. \quad y(x) + A \int_0^x (ax^n + bt^n)^{-\frac{\lambda+1}{n}} y^2(t) dt = Bx^\lambda.$$

Solutions: $y_1(x) = \beta_1 x^\lambda$ and $y_2(x) = \beta_2 x^\lambda$, where $\beta_{1,2}$ are the roots of the quadratic equation

$$AI\beta^2 + \beta - B = 0, \quad I = \int_0^1 z^{2\lambda} (a + bz^n)^{-\frac{\lambda+1}{n}} dz.$$

$$9. \quad y(x) + A \int_a^x e^{\lambda t} y^2(t) dt = Be^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = Ay^2$. By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

Solution in an implicit form:

$$\lambda \int_{y_0}^y \frac{du}{Au^2 - B\lambda} + e^{\lambda x} - e^{\lambda a} = 0, \quad y_0 = Be^{\lambda a} + C.$$

$$10. \quad y(x) + A \int_a^x e^{\lambda(x-t)} y^2(t) dt = B.$$

This is a special case of equation 6.8.10. By differentiation, this integral equation can be reduced to the separable ordinary differential equation

$$y'_x + Ay^2 - \lambda y + \lambda B = 0, \quad y(a) = B.$$

Solution in an implicit form:

$$\int_B^y \frac{du}{Au^2 - \lambda u + \lambda B} + x - a = 0.$$

$$11. \quad y(x) + k \int_a^x e^{\lambda(x-t)} y^2(t) dt = Ae^{\lambda x} + B.$$

Solution in an implicit form:

$$\int_{y_0}^y \frac{du}{\lambda u - ku^2 - \lambda B} = x - a, \quad y_0 = Ae^{\lambda a} + B.$$

$$12. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] y^2(t) dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = ky^2$.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 Cu - 2k\lambda F(u) + \lambda^2(C^2 - 4AB)]^{-1/2} du = \pm(x - a),$$

$$F(u) = \frac{1}{3}(u^3 - y_0^3), \quad y_0 = Ae^{\lambda a} + Be^{-\lambda a} + C.$$

$$13. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] y^2(t) dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = ky^2$.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 Bu - 2k\lambda F(u) + \lambda^2(B^2 - A^2)]^{-1/2} du = \pm(x - a),$$

$$F(u) = \frac{1}{3}(u^3 - y_0^3), \quad y_0 = A \cosh(\lambda a) + B.$$

$$14. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] y^2(t) dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = ky^2$.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 Bu - 2k\lambda F(u) + \lambda^2(A^2 + B^2)]^{-1/2} du = \pm(x - a),$$

$$F(u) = \frac{1}{3}(u^3 - y_0^3), \quad y_0 = A \sinh(\lambda a) + B.$$

$$15. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] y^2(t) dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = ky^2$.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 D - \lambda^2 u^2 + 2\lambda^2 Cu - 2k\lambda F(u)]^{-1/2} du = \pm(x - a),$$

$$y_0 = A \sin(\lambda a) + B \cos(\lambda a) + C, \quad D = A^2 + B^2 - C^2, \quad F(u) = \frac{1}{3}(u^3 - y_0^3).$$

6.1-2. Equations of the Form $y(x) + \int_a^x K(x, t)y(t)y(x-t) dt = F(x)$.

16. $y(x) + A \int_0^x y(t)y(x-t) dt = AB^2x + B.$

A solution: $y(x) = B.$

17. $y(x) + A \int_0^x y(t)y(x-t) dt = (AB^2x + B)e^{\lambda x}.$

A solution: $y(x) = Be^{\lambda x}.$

18. $y(x) + \frac{\lambda}{2\beta} \int_0^x y(t)y(x-t) dt = \frac{1}{2}\beta \sinh(\lambda x).$

A solution: $y(x) = \beta I_1(\lambda x)$, where $I_1(x)$ is the modified Bessel function.

19. $y(x) - \frac{\lambda}{2\beta} \int_0^x y(t)y(x-t) dt = \frac{1}{2}\beta \sin(\lambda x).$

A solution: $y(x) = \beta J_1(\lambda x)$, where $J_1(x)$ is the Bessel function.

20. $y(x) + A \int_0^x x^{-\lambda-1}y(t)y(x-t) dt = Bx^\lambda.$

Solutions: $y_1(x) = \beta_1 x^\lambda$ and $y_2(x) = \beta_2 x^\lambda$, where $\beta_{1,2}$ are the roots of the quadratic equation

$$AI\beta^2 + \beta - B = 0, \quad I = \int_0^1 z^\lambda(1-z)^\lambda dz = \frac{\Gamma^2(\lambda+1)}{\Gamma(2\lambda+2)}.$$

6.2. Equations with Quadratic Nonlinearity That Contain Arbitrary Functions

6.2-1. Equations of the Form $y(x) + \int_a^x K(x, t)y^2(t) dt = F(x)$.

1. $y(x) + \int_a^x f(t)y^2(t) dt = A.$

Solution:

$$y(x) = A \left[1 + A \int_a^x f(t) dt \right]^{-1}.$$

2. $y(x) + \int_a^x e^{\lambda(x-t)}g(t)y^2(t) dt = f(x).$

Differentiating the equation with respect to x yields

$$y'_x + g(x)y^2 + \lambda \int_a^x e^{\lambda(x-t)}g(t)y^2(t) dt = f'_x(x). \quad (1)$$

Eliminating the integral term from (1) with the aid of the original equation, we arrive at a Riccati ordinary differential equation,

$$y'_x + g(x)y^2 - \lambda y + \lambda f(x) - f'_x(x) = 0, \quad (2)$$

under the initial condition $y(a) = f(a)$. Equation (2) can be reduced to a second-order linear ordinary differential equation. For the exact solutions of equation (2) with various specific functions f and g , see, for example, E. Kamke (1977) and A. D. Polyanin and V. F. Zaitsev (2003).

$$3. \quad y(x) + \int_a^x g(x)h(t)y^2(t) dt = f(x).$$

Differentiating the equation with respect to x yields

$$y'_x + g(x)h(x)y^2 + g'_x(x) \int_a^x h(t)y^2(t) dt = f'_x(x). \quad (1)$$

Eliminating the integral term from (1) with the aid of the original equation, we arrive at a Riccati ordinary differential equation,

$$y'_x + g(x)h(x)y^2 - \frac{g'_x(x)}{g(x)}y = f'_x(x) - \frac{g'_x(x)}{g(x)}f(x), \quad (2)$$

under the initial condition $y(a) = f(a)$. Equation (2) can be reduced to a second-order linear ordinary differential equation. For the exact solutions of equation (2) with various specific functions f , g , and h , see, for example, E. Kamke (1977) and A. D. Polyanin and V. F. Zaitsev (2003).

$$4. \quad y(x) + \int_0^x x^{-\lambda-1} f\left(\frac{t}{x}\right) y^2(t) dt = Ax^\lambda.$$

Solutions: $y_1(x) = \beta_1 x^\lambda$ and $y_2(x) = \beta_2 x^\lambda$, where $\beta_{1,2}$ are the roots of the quadratic equation

$$I\beta^2 + \beta - A = 0, \quad I = \int_0^1 f(z)z^{2\lambda} dz.$$

$$5. \quad y(x) - \int_{-\infty}^x e^{\lambda t + \beta x} f(x-t)y^2(t) dt = 0.$$

This is a special case of equation 6.3.19 with $k = 2$.

$$6. \quad y(x) - \int_x^\infty e^{\lambda t + \beta x} f(x-t)y^2(t) dt = 0.$$

A solution:

$$y(x) = \frac{1}{A} e^{-(\lambda+\beta)x}, \quad A = \int_0^\infty e^{-(\lambda+2\beta)z} f(-z) dz.$$

6.2-2. Other Equations.

$$7. \quad y(x) + \int_0^x \frac{1}{x} f\left(\frac{t}{x}\right) y(t)y(x-t) dt = Ae^{\lambda x}.$$

Solutions:

$$y_1(x) = B_1 e^{\lambda x}, \quad y_2(x) = B_2 e^{\lambda x},$$

where B_1 and B_2 are the roots of the quadratic equation

$$IB^2 + B - A = 0, \quad I = \int_0^1 f(z) dz.$$

$$8. \quad y(x) + A \int_0^x x^{-\lambda-1} f\left(\frac{t}{x}\right) y(t) y(x-t) dt = Bx^\lambda.$$

Solutions: $y_1(x) = \beta_1 x^\lambda$ and $y_2(x) = \beta_2 x^\lambda$, where $\beta_{1,2}$ are the roots of the quadratic equation

$$AI\beta^2 + \beta - B = 0, \quad I = \int_0^1 f(z) z^\lambda (1-z)^\lambda dz.$$

$$9. \quad y(x) + \int_x^\infty f(t-x) y(t-x) y(t) dt = a e^{-\lambda x}.$$

Solutions: $y(x) = b_k e^{-\lambda x}$, where b_k ($k = 1, 2$) are the roots of the quadratic equation

$$b^2 I + b - a = 0, \quad I = \int_0^\infty f(z) e^{-2\lambda z} dz.$$

To calculate the integral I , it is convenient to use tables of Laplace transforms (with parameter $p = 2\lambda$).

6.3. Equations with Power-Law Nonlinearity

6.3-1. Equations Containing Arbitrary Parameters.

$$1. \quad y(x) = a \int_0^x y^k(t) dt + b, \quad a > 0, b > 0, k > 0.$$

Solution:

$$y(x) = \begin{cases} [b^{1-k} + a(1-k)x]^{\frac{1}{1-k}} & \text{if } 0 < k < 1, \\ be^{ax} & \text{if } k = 1, \\ [b^{1-k} - a(k-1)x]^{\frac{1}{1-k}} & \text{if } k > 1. \end{cases}$$

If $0 < k \leq 1$, the solution exists for all $x \geq 0$. If $k > 1$, the continuous solution exists only in a limited interval of argument variation

$$0 \leq x < x_* = \frac{b^{1-k}}{a(k-1)}.$$

$$2. \quad y(x) + A \int_a^x t^\lambda y^k(t) dt = Bx^{\lambda+1} + C.$$

By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

Solution in an implicit form:

$$(\lambda + 1) \int_{y_0}^y \frac{du}{Au^k - B(\lambda + 1)} + x^{\lambda+1} - a^{\lambda+1} = 0, \quad y_0 = Ba^{\lambda+1} + C.$$

$$3. \quad y(x) + \int_0^x \frac{y^k(t)}{ax + bt} dt = A.$$

A solution: $y(x) = \lambda$, where λ is a root of the algebraic (or transcendental) equation

$$\ln\left(1 + \frac{b}{a}\right) \lambda^k + b\lambda - Ab = 0.$$

$$4. \quad y(x) + Ax \int_0^x \frac{y^k(t) dt}{x^2 + t^2} = B.$$

A solution: $y(x) = \lambda$, where λ is a root of the algebraic (or transcendental) equation

$$\lambda + \frac{1}{4}A\pi\lambda^k = B.$$

$$5. \quad y(x) + \int_0^x \frac{y^k(t) dt}{\sqrt{ax^2 + bt^2}} = A.$$

A solution: $y(x) = \lambda$, where λ is a root of the algebraic (or transcendental) equation

$$I\lambda^k + \lambda - A = 0, \quad I = \int_0^1 \frac{dz}{\sqrt{a + bz^2}}.$$

$$6. \quad y(x) + A \int_a^x (ax^n + bt^n)^{\frac{\lambda-k\lambda-1}{n}} y^k(t) dt = Bx^\lambda.$$

A solution: $y = \beta x^\lambda$, where β is a root of the algebraic (or transcendental) equation

$$AI\beta^k + \beta - B = 0, \quad I = \int_0^1 z^{k\lambda} (a + bz^n)^{\frac{\lambda-k\lambda-1}{n}} dz.$$

$$7. \quad y(x) + A \int_a^x e^{\lambda t} y^\mu(t) dt = Be^{\lambda x} + C.$$

By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

Solution in an implicit form:

$$\lambda \int_{y_0}^y \frac{du}{Au^\mu - B\lambda} + e^{\lambda x} - e^{\lambda a} = 0, \quad y_0 = Be^{\lambda a} + C.$$

$$8. \quad y(x) + k \int_a^x e^{\lambda(x-t)} y^\mu(t) dt = Ae^{\lambda x} + B.$$

Solution in an implicit form:

$$\int_{y_0}^y \frac{dt}{\lambda t - kt^\mu - \lambda B} = x - a, \quad y_0 = Ae^{\lambda a} + B.$$

$$9. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] y^\mu(t) dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = ky^\mu$.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 C u - 2k\lambda F(u) + \lambda^2 (C^2 - 4AB)]^{-1/2} du = \pm(x - a),$$

$$F(u) = \frac{1}{\mu + 1} (u^{\mu+1} - y_0^{\mu+1}), \quad y_0 = Ae^{\lambda a} + Be^{-\lambda a} + C.$$

$$10. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)]y^\mu(t) dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = ky^\mu$.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 B u - 2k\lambda F(u) + \lambda^2(B^2 - A^2)]^{-1/2} du = \pm(x-a),$$

$$F(u) = \frac{1}{\mu+1} (u^{\mu+1} - y_0^{\mu+1}), \quad y_0 = A \cosh(\lambda a) + B.$$

$$11. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)]y^\mu(t) dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = ky^\mu$.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 B u - 2k\lambda F(u) + \lambda^2(A^2 + B^2)]^{-1/2} du = \pm(x-a),$$

$$F(u) = \frac{1}{\mu+1} (u^{\mu+1} - y_0^{\mu+1}), \quad y_0 = A \sinh(\lambda a) + B.$$

$$12. \quad y(x) + k \int_a^x \sin[\lambda(x-t)]y^\mu(t) dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = ky^\mu$.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 D - \lambda^2 u^2 + 2\lambda^2 C u - 2k\lambda F(u)]^{-1/2} du = \pm(x-a),$$

$$y_0 = A \sin(\lambda a) + B \cos(\lambda a) + C, \quad D = A^2 + B^2 - C^2, \quad F(u) = \frac{1}{\mu+1} (u^{\mu+1} - y_0^{\mu+1}).$$

6.3-2. Equations Containing Arbitrary Functions.

$$13. \quad y(x) + \int_a^x f(t)y^k(t) dt = A.$$

Solution:

$$y(x) = \left[A^{1-k} + (k-1) \int_a^x f(t) dt \right]^{\frac{1}{1-k}}.$$

$$14. \quad y(x) - \int_a^x f(x)g(t)y^k(t) dt = 0.$$

1°. Differentiating the equation with respect to x and eliminating the integral term (using the original equation), we obtain the Bernoulli ordinary differential equation

$$y'_x - f(x)g(x)y^k - \frac{f'_x(x)}{f(x)}y = 0, \quad y(a) = 0.$$

2°. Solution with $k < 1$:

$$y(x) = f(x) \left[(1-k) \int_a^x f^k(t)g(t) dt \right]^{\frac{1}{1-k}}.$$

Additionally, for $k > 0$, there is the trivial solution $y(x) \equiv 0$.

$$15. \quad y(x) + \int_0^x x^{\lambda-k\lambda-1} f\left(\frac{t}{x}\right) y^k(t) dt = Ax^\lambda.$$

A solution: $y(x) = \beta x^\lambda$, where β is a root of the algebraic equation

$$I\beta^k + \beta - A = 0, \quad I = \int_0^1 f(z)z^{k\lambda} dz.$$

$$16. \quad y(x) + \int_0^x f\left(\frac{t}{x}\right) \sqrt{y(t)} dt = Ax^2.$$

Solutions: $y_k(x) = B_k^2 x^2$, where B_k ($k = 1, 2$) are the roots of the quadratic equations

$$B^2 \pm IB - A = 0, \quad I = \int_0^1 z f(z) dz.$$

$$17. \quad y(x) - \int_0^x t^\alpha f\left(\frac{t}{x}\right) y^k(t) dt = 0, \quad k \neq 1.$$

A solution:

$$y(x) = Ax^{\frac{1+\alpha}{1-k}}, \quad A^{1-k} = \int_0^1 z^{\frac{\alpha+k}{1-k}} f(z) dz.$$

$$18. \quad y(x) - \int_x^\infty e^{\lambda t + \beta x} f(x-t) y^k(t) dt = 0, \quad k \neq 1.$$

A solution:

$$y(x) = A \exp\left(\frac{\lambda + \beta}{1-k} x\right), \quad A^{1-k} = \int_0^\infty \exp\left(\frac{\lambda + \beta k}{1-k} z\right) f(-z) dz.$$

$$19. \quad y(x) - \int_{-\infty}^x e^{\lambda t + \beta x} f(x-t) y^k(t) dt = 0, \quad k \neq 1.$$

A solution:

$$y(x) = A \exp\left(\frac{\lambda + \beta}{1-k} x\right), \quad A^{1-k} = \int_0^\infty \exp\left(-\frac{\lambda + \beta k}{1-k} z\right) f(z) dz.$$

6.4. Equations with Exponential Nonlinearity

6.4-1. Equations Containing Arbitrary Parameters.

$$1. \quad y(x) + A \int_a^x \exp[\lambda y(t)] dt = B.$$

Solution:

$$y(x) = -\frac{1}{\lambda} \ln [A\lambda(x-a) + e^{-B\lambda}].$$

$$2. \quad y(x) + A \int_a^x \exp[\lambda y(t)] dt = Bx + C.$$

For $B = 0$, see equation 6.4.1.

Solution with $B \neq 0$:

$$y(x) = -\frac{1}{\lambda} \ln \left[\frac{A}{B} + \left(e^{-\lambda y_0} - \frac{A}{B} \right) e^{\lambda B(a-x)} \right], \quad y_0 = aB + C.$$

$$3. \quad y(x) + k \int_a^x (x-t) \exp[\lambda y(t)] dt = Ax^2 + Bx + C.$$

1°. This is a special case of equation 6.8.3 with $f(y) = ke^{\lambda y}$. The solution of this integral equation is determined by the solution of the second-order autonomous ordinary differential equation

$$y''_{xx} + ke^{\lambda y} - 2A = 0$$

under the initial conditions

$$y(a) = Aa^2 + Ba + C, \quad y'_x(a) = 2Aa + B.$$

2°. Solution in an implicit form:

$$\int_{y_0}^y [4Au - 2F(u) + B^2 - 4AC]^{-1/2} du = \pm(x-a),$$

$$F(u) = \frac{k}{\lambda} (e^{\lambda u} - e^{\lambda y_0}), \quad y_0 = Aa^2 + Ba + C.$$

$$4. \quad y(x) + A \int_a^x t^\lambda \exp[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

Solution in an implicit form:

$$(\lambda + 1) \int_{y_0}^y \frac{du}{Ae^{\beta u} - B(\lambda + 1)} + x^{\lambda+1} - a^{\lambda+1} = 0, \quad y_0 = Ba^{\lambda+1} + C.$$

$$5. \quad y(x) + \int_0^x \frac{\exp[\lambda y(t)]}{ax + bt} dt = A.$$

A solution: $y(x) = \beta$, where β is a root of the transcendental equation

$$\ln\left(1 + \frac{b}{a}\right) e^{\lambda\beta} + b\beta - Ab = 0.$$

$$6. \quad y(x) + \int_0^x \frac{\exp[\lambda y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

A solution: $y(x) = \beta$, where β is a root of the transcendental equation

$$ke^{\lambda\beta} + \beta - A = 0, \quad k = \int_0^1 \frac{dz}{\sqrt{a + bz^2}}.$$

$$7. \quad y(x) + A \int_a^x \exp[\lambda t + \beta y(t)] dt = Be^{\lambda x} + C.$$

By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

Solution in an implicit form:

$$\lambda \int_{y_0}^y \frac{du}{Ae^{\beta u} - B\lambda} + e^{\lambda x} - e^{\lambda a} = 0, \quad y_0 = Be^{\lambda a} + C.$$

$$8. \quad y(x) + k \int_a^x \exp[\lambda(x-t) + \beta y(t)] dt = A.$$

Solution in an implicit form:

$$\int_A^y \frac{dt}{\lambda t - k e^{\beta t} - \lambda A} = x - a.$$

$$9. \quad y(x) + k \int_a^x \exp[\lambda(x-t) + \beta y(t)] dt = A e^{\lambda x} + B.$$

Solution in an implicit form:

$$\int_{y_0}^y \frac{dt}{\lambda t - k e^{\beta t} - \lambda B} = x - a, \quad y_0 = A e^{\lambda a} + B.$$

$$10. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \exp[\beta y(t)] dt = A e^{\lambda x} + B e^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k e^{\beta y}$.

$$11. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \exp[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k e^{\beta y}$.

$$12. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \exp[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k e^{\beta y}$.

$$13. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \exp[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k e^{\beta y}$.

6.4-2. Equations Containing Arbitrary Functions.

$$14. \quad y(x) + \int_a^x f(t) \exp[\lambda y(t)] dt = A.$$

Solution:

$$y(x) = -\frac{1}{\lambda} \ln \left[\lambda \int_a^x f(t) dt + e^{-A\lambda} \right].$$

$$15. \quad y(x) + \int_a^x g(t) \exp[\lambda y(t)] dt = f(x).$$

1°. By differentiation, this integral equation can be reduced to the first-order ordinary differential equation

$$y'_x + g(x) e^{\lambda y} = f'_x(x) \tag{1}$$

under the initial condition $y(a) = f(a)$. The substitution $w = e^{-\lambda y}$ reduces (1) to the linear equation

$$w'_x + \lambda f'_x(x) w - \lambda g(x) = 0, \quad w(a) = \exp[-\lambda f(a)].$$

2°. Solution:

$$y(x) = f(x) - \frac{1}{\lambda} \ln \left\{ 1 + \lambda \int_a^x g(t) \exp[\lambda f(t)] dt \right\}.$$

$$16. \quad y(x) + \frac{1}{x} \int_0^x f\left(\frac{t}{x}\right) \exp[\lambda y(t)] dt = A.$$

A solution: $y(x) = \beta$, where β is a root of the transcendental equation

$$\beta + Ie^{\lambda\beta} - A = 0, \quad I = \int_0^1 f(z) dz.$$

6.5. Equations with Hyperbolic Nonlinearity

6.5-1. Integrands with Nonlinearity of the Form $\cosh[\beta y(t)]$.

$$1. \quad y(x) + k \int_a^x \cosh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \cosh(\beta y)$.

$$2. \quad y(x) + k \int_a^x \cosh[\beta y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \cosh(\beta y)$.

$$3. \quad y(x) + k \int_a^x (x-t) \cosh[\beta y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \cosh(\beta y)$.

$$4. \quad y(x) + k \int_a^x t^\lambda \cosh[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \cosh(\beta y)$.

$$5. \quad y(x) + \int_a^x g(t) \cosh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \cosh(\beta y)$.

$$6. \quad y(x) + \int_0^x \frac{\cosh[\beta y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \cosh(\beta y)$.

$$7. \quad y(x) + \int_0^x \frac{\cosh[\beta y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \cosh(\beta y)$.

$$8. \quad y(x) + k \int_a^x e^{\lambda t} \cosh[\beta y(t)] dt = Be^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \cosh(\beta y)$.

$$9. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \cosh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \cosh(\beta y)$.

$$10. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \cosh[\beta y(t)] dt = Ae^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \cosh(\beta y)$.

$$11. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cosh[\beta y(t)] dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \cosh(\beta y)$.

$$12. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cosh[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \cosh(\beta y)$.

$$13. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cosh[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \cosh(\beta y)$.

$$14. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \cosh[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \cosh(\beta y)$.

6.5-2. Integrands with Nonlinearity of the Form $\sinh[\beta y(t)]$.

$$15. \quad y(x) + k \int_a^x \sinh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \sinh(\beta y)$.

$$16. \quad y(x) + k \int_a^x \sinh[\beta y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \sinh(\beta y)$.

$$17. \quad y(x) + k \int_a^x (x-t) \sinh[\beta y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \sinh(\beta y)$.

$$18. \quad y(x) + k \int_a^x t^\lambda \sinh[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \sinh(\beta y)$.

$$19. \quad y(x) + \int_a^x g(t) \sinh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \sinh(\beta y)$.

$$20. \quad y(x) + \int_0^x \frac{\sinh[\beta y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \sinh(\beta y)$.

$$21. \quad y(x) + \int_0^x \frac{\sinh[\beta y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \sinh(\beta y)$.

$$22. \quad y(x) + k \int_a^x e^{\lambda t} \sinh[\beta y(t)] dt = B e^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \sinh(\beta y)$.

$$23. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \sinh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \sinh(\beta y)$.

$$24. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \sinh[\beta y(t)] dt = A e^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \sinh(\beta y)$.

$$25. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \sinh[\beta y(t)] dt = A e^{\lambda x} + B e^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \sinh(\beta y)$.

$$26. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \sinh[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \sinh(\beta y)$.

$$27. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \sinh[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \sinh(\beta y)$.

$$28. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \sinh[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \sinh(\beta y)$.

6.5-3. Integrands with Nonlinearity of the Form $\tanh[\beta y(t)]$.

$$29. \quad y(x) + k \int_a^x \tanh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \tanh(\beta y)$.

$$30. \quad y(x) + k \int_a^x \tanh[\beta y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \tanh(\beta y)$.

$$31. \quad y(x) + k \int_a^x (x-t) \tanh[\beta y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \tanh(\beta y)$.

$$32. \quad y(x) + k \int_a^x t^\lambda \tanh[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \tanh(\beta y)$.

$$33. \quad y(x) + \int_a^x g(t) \tanh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \tanh(\beta y)$.

$$34. \quad y(x) + \int_0^x \frac{\tanh[\beta y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \tanh(\beta y)$.

$$35. \quad y(x) + \int_0^x \frac{\tanh[\beta y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \tanh(\beta y)$.

$$36. \quad y(x) + k \int_a^x e^{\lambda t} \tanh[\beta y(t)] dt = Be^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \tanh(\beta y)$.

$$37. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \tanh[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \tanh(\beta y)$.

$$38. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \tanh[\beta y(t)] dt = Ae^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \tanh(\beta y)$.

$$39. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \tanh[\beta y(t)] dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \tanh(\beta y)$.

$$40. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \tanh[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \tanh(\beta y)$.

$$41. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \tanh[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \tanh(\beta y)$.

$$42. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \tanh[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \tanh(\beta y)$.

6.5-4. Integrands with Nonlinearity of the Form $\coth[\beta y(t)]$.

$$43. \quad y(x) + k \int_a^x \coth[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \coth(\beta y)$.

$$44. \quad y(x) + k \int_a^x \coth[\beta y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \coth(\beta y)$.

$$45. \quad y(x) + k \int_a^x (x-t) \coth[\beta y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \coth(\beta y)$.

$$46. \quad y(x) + k \int_a^x t^\lambda \coth[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \coth(\beta y)$.

$$47. \quad y(x) + \int_a^x g(t) \coth[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \coth(\beta y)$.

$$48. \quad y(x) + \int_0^x \frac{\coth[\beta y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \coth(\beta y)$.

$$49. \quad y(x) + \int_0^x \frac{\coth[\beta y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \coth(\beta y)$.

$$50. \quad y(x) + k \int_a^x e^{\lambda t} \coth[\beta y(t)] dt = Be^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \coth(\beta y)$.

$$51. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \coth[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \coth(\beta y)$.

$$52. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \coth[\beta y(t)] dt = Ae^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \coth(\beta y)$.

$$53. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \coth[\beta y(t)] dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \coth(\beta y)$.

$$54. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \coth[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \coth(\beta y)$.

$$55. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \coth[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \coth(\beta y)$.

$$56. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \coth[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \coth(\beta y)$.

6.6. Equations with Logarithmic Nonlinearity

6.6-1. Integrands Containing Power-Law Functions of x and t .

$$1. \quad y(x) + k \int_a^x \ln[\lambda y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \ln(\lambda y)$.

$$2. \quad y(x) + k \int_a^x \ln[\lambda y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \ln(\lambda y)$.

$$3. \quad y(x) + k \int_a^x (x-t) \ln[\lambda y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \ln(\lambda y)$.

$$4. \quad y(x) + k \int_a^x t^\lambda \ln[\mu y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \ln(\mu y)$.

$$5. \quad y(x) + \int_0^x \frac{\ln[\lambda y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \ln(\lambda y)$.

$$6. \quad y(x) + \int_0^x \frac{\ln[\lambda y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \ln(\lambda y)$.

6.6-2. Integrands Containing Exponential Functions of x and t .

$$7. \quad y(x) + k \int_a^x e^{\lambda t} \ln[\mu y(t)] dt = Be^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \ln(\mu y)$.

$$8. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \ln[\mu y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \ln(\mu y)$.

$$9. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \ln[\mu y(t)] dt = Ae^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \ln(\mu y)$.

6.6-3. Other Integrands.

$$10. \quad y(x) + \int_a^x g(t) \ln[\lambda y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \ln(\lambda y)$.

$$11. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \ln[\mu y(t)] dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \ln(\mu y)$.

$$12. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \ln[\mu y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \ln(\mu y)$.

$$13. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \ln[\mu y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \ln(\mu y)$.

$$14. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \ln[\mu y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \ln(\mu y)$.

6.7. Equations with Trigonometric Nonlinearity

6.7-1. Integrands with Nonlinearity of the Form $\cos[\beta y(t)]$.
--

$$1. \quad y(x) + k \int_a^x \cos[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \cos(\beta y)$.

$$2. \quad y(x) + k \int_a^x \cos[\beta y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \cos(\beta y)$.

$$3. \quad y(x) + k \int_a^x (x-t) \cos[\beta y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \cos(\beta y)$.

$$4. \quad y(x) + k \int_a^x t^\lambda \cos[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \cos(\beta y)$.

$$5. \quad y(x) + \int_a^x g(t) \cos[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \cos(\beta y)$.

$$6. \quad y(x) + \int_0^x \frac{\cos[\beta y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \cos(\beta y)$.

$$7. \quad y(x) + \int_0^x \frac{\cos[\beta y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \cos(\beta y)$.

$$8. \quad y(x) + k \int_a^x e^{\lambda t} \cos[\beta y(t)] dt = Be^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \cos(\beta y)$.

$$9. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \cos[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \cos(\beta y)$.

$$10. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \cos[\beta y(t)] dt = Ae^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \cos(\beta y)$.

$$11. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cos[\beta y(t)] dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \cos(\beta y)$.

$$12. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cos[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \cos(\beta y)$.

$$13. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cos[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \cos(\beta y)$.

$$14. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \cos[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \cos(\beta y)$.

6.7-2. Integrands with Nonlinearity of the Form $\sin[\beta y(t)]$.

$$15. \quad y(x) + k \int_a^x \sin[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \sin(\beta y)$.

$$16. \quad y(x) + k \int_a^x \sin[\beta y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \sin(\beta y)$.

$$17. \quad y(x) + k \int_a^x (x-t) \sin[\beta y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \sin(\beta y)$.

$$18. \quad y(x) + k \int_a^x t^\lambda \sin[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \sin(\beta y)$.

$$19. \quad y(x) + \int_a^x g(t) \sin[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \sin(\beta y)$.

$$20. \quad y(x) + \int_0^x \frac{\sin[\beta y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \sin(\beta y)$.

$$21. \quad y(x) + \int_0^x \frac{\sin[\beta y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \sin(\beta y)$.

$$22. \quad y(x) + k \int_a^x e^{\lambda t} \sin[\beta y(t)] dt = Be^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \sin(\beta y)$.

$$23. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \sin[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \sin(\beta y)$.

$$24. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \sin[\beta y(t)] dt = Ae^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \sin(\beta y)$.

$$25. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \sin[\beta y(t)] dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \sin(\beta y)$.

$$26. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \sin[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \sin(\beta y)$.

$$27. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \sin[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \sin(\beta y)$.

$$28. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \sin[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \sin(\beta y)$.

6.7-3. Integrands with Nonlinearity of the Form $\tan[\beta y(t)]$.

$$29. \quad y(x) + k \int_a^x \tan[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \tan(\beta y)$.

$$30. \quad y(x) + k \int_a^x \tan[\beta y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \tan(\beta y)$.

$$31. \quad y(x) + k \int_a^x (x-t) \tan[\beta y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \tan(\beta y)$.

$$32. \quad y(x) + k \int_a^x t^\lambda \tan[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \tan(\beta y)$.

$$33. \quad y(x) + \int_a^x g(t) \tan[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \tan(\beta y)$.

$$34. \quad y(x) + \int_0^x \frac{\tan[\beta y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \tan(\beta y)$.

$$35. \quad y(x) + \int_0^x \frac{\tan[\beta y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \tan(\beta y)$.

$$36. \quad y(x) + k \int_a^x e^{\lambda t} \tan[\beta y(t)] dt = Be^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \tan(\beta y)$.

$$37. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \tan[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \tan(\beta y)$.

$$38. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \tan[\beta y(t)] dt = Ae^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \tan(\beta y)$.

$$39. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \tan[\beta y(t)] dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \tan(\beta y)$.

$$40. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \tan[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \tan(\beta y)$.

$$41. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \tan[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \tan(\beta y)$.

$$42. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \tan[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \tan(\beta y)$.

6.7-4. Integrands with Nonlinearity of the Form $\cot[\beta y(t)]$.

$$43. \quad y(x) + k \int_a^x \cot[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.1 with $f(y) = k \cot(\beta y)$.

$$44. \quad y(x) + k \int_a^x \cot[\beta y(t)] dt = Ax + B.$$

This is a special case of equation 6.8.2 with $f(y) = k \cot(\beta y)$.

$$45. \quad y(x) + k \int_a^x (x-t) \cot[\beta y(t)] dt = Ax^2 + Bx + C.$$

This is a special case of equation 6.8.3 with $f(y) = k \cot(\beta y)$.

$$46. \quad y(x) + k \int_a^x t^\lambda \cot[\beta y(t)] dt = Bx^{\lambda+1} + C.$$

This is a special case of equation 6.8.4 with $f(y) = k \cot(\beta y)$.

$$47. \quad y(x) + \int_a^x g(t) \cot[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.5 with $f(y) = \cot(\beta y)$.

$$48. \quad y(x) + \int_0^x \frac{\cot[\beta y(t)]}{ax + bt} dt = A.$$

This is a special case of equation 6.8.6 with $f(y) = \cot(\beta y)$.

$$49. \quad y(x) + \int_0^x \frac{\cot[\beta y(t)]}{\sqrt{ax^2 + bt^2}} dt = A.$$

This is a special case of equation 6.8.7 with $f(y) = \cot(\beta y)$.

$$50. \quad y(x) + k \int_a^x e^{\lambda t} \cot[\beta y(t)] dt = B e^{\lambda x} + C.$$

This is a special case of equation 6.8.9 with $f(y) = k \cot(\beta y)$.

$$51. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \cot[\beta y(t)] dt = A.$$

This is a special case of equation 6.8.10 with $f(y) = k \cot(\beta y)$.

$$52. \quad y(x) + k \int_a^x e^{\lambda(x-t)} \cot[\beta y(t)] dt = A e^{\lambda x} + B.$$

This is a special case of equation 6.8.11 with $f(y) = k \cot(\beta y)$.

$$53. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cot[\beta y(t)] dt = A e^{\lambda x} + B e^{-\lambda x} + C.$$

This is a special case of equation 6.8.12 with $f(y) = k \cot(\beta y)$.

$$54. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cot[\beta y(t)] dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.13 with $f(y) = k \cot(\beta y)$.

$$55. \quad y(x) + k \int_a^x \sinh[\lambda(x-t)] \cot[\beta y(t)] dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.14 with $f(y) = k \cot(\beta y)$.

$$56. \quad y(x) + k \int_a^x \sin[\lambda(x-t)] \cot[\beta y(t)] dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

This is a special case of equation 6.8.15 with $f(y) = k \cot(\beta y)$.

6.8. Equations with Nonlinearity of General Form

6.8-1. Equations of the Form $y(x) + \int_a^x K(x,t)G(y(t)) dt = F(x)$.

$$1. \quad y(x) + \int_a^x f(y(t)) dt = A.$$

This is a special case of equation 6.8.16.

Solution in an implicit form:

$$\int_A^y \frac{du}{f(u)} + x - a = 0.$$

$$2. \quad y(x) + \int_a^x f(y(t)) dt = Ax + B.$$

Solution in an implicit form:

$$\int_{y_0}^y \frac{du}{A - f(u)} = x - a, \quad y_0 = Aa + B.$$

$$3. \quad y(x) + \int_a^x (x - t)f(y(t)) dt = Ax^2 + Bx + C.$$

1°. This is a special case of equation 6.8.17. The solution of this integral equation is determined by the solution of the second-order autonomous ordinary differential equation

$$y''_{xx} + f(y) - 2A = 0$$

under the initial conditions

$$y(a) = Aa^2 + Ba + C, \quad y'_x(a) = 2Aa + B.$$

2°. Solutions in an implicit form:

$$\int_{y_0}^y [4Au - 2F(u) + B^2 - 4AC]^{-1/2} du = \pm(x - a),$$

$$F(u) = \int_{y_0}^u f(t) dt, \quad y_0 = Aa^2 + Ba + C.$$

$$4. \quad y(x) + \int_a^x t^\lambda f(y(t)) dt = Bx^{\lambda+1} + C.$$

By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

Solution in an implicit form:

$$(\lambda + 1) \int_{y_a}^y \frac{du}{f(u) - B(\lambda + 1)} + x^{\lambda+1} - a^{\lambda+1} = 0, \quad y_a = Ba^{\lambda+1} + C.$$

$$5. \quad y(x) + \int_a^x g(t)f(y(t)) dt = A.$$

Solution in an implicit form:

$$\int_A^y \frac{du}{f(u)} + \int_a^x g(t) dt = 0.$$

$$6. \quad y(x) + \int_0^x \frac{f(y(t))}{ax + bt} dt = A.$$

A solution: $y(x) = \lambda$, where λ is a root of the algebraic (or transcendental) equation

$$\ln\left(1 + \frac{b}{a}\right)f(\lambda) + b\lambda - Ab = 0.$$

$$7. \quad y(x) + \int_0^x \frac{f(y(t))}{\sqrt{ax^2 + bt^2}} dt = A.$$

A solution: $y(x) = \lambda$, where λ is a root of the algebraic (or transcendental) equation

$$kf(\lambda) + \lambda - A = 0, \quad k = \int_0^1 \frac{dz}{\sqrt{a + bz^2}}.$$

$$8. \quad y(x) + x \int_0^x f(y(t)) \frac{dt}{x^2 + t^2} = A.$$

A solution: $y(x) = \lambda$, where λ is a root of the algebraic (or transcendental) equation

$$\lambda + \frac{1}{4}\pi f(\lambda) = A.$$

$$9. \quad y(x) + \int_a^x e^{\lambda t} f(y(t)) dt = Be^{\lambda x} + C.$$

By differentiation, this integral equation can be reduced to a separable ordinary differential equation.

Solution in an implicit form:

$$\lambda \int_{y_0}^y \frac{du}{f(u) - B\lambda} + e^{\lambda x} - e^{\lambda a} = 0, \quad y_0 = Be^{\lambda a} + C.$$

$$10. \quad y(x) + \int_a^x e^{\lambda(x-t)} f(y(t)) dt = A.$$

This is a special case of equation 6.8.19.

Solution in an implicit form:

$$\int_A^y \frac{du}{\lambda u - f(u) - \lambda A} = x - a.$$

$$11. \quad y(x) + \int_a^x e^{\lambda(x-t)} f(y(t)) dt = Ae^{\lambda x} + B.$$

This is a special case of equation 6.8.19.

Solution in an implicit form:

$$\int_{y_0}^y \frac{du}{\lambda u - f(u) - \lambda B} = x - a, \quad y_0 = Ae^{\lambda a} + B.$$

$$12. \quad y(x) + \int_a^x \sinh[\lambda(x-t)] f(y(t)) dt = Ae^{\lambda x} + Be^{-\lambda x} + C.$$

1°. This is a special case of equation 6.8.21. The solution of this integral equation is determined by the solution of the second-order autonomous ordinary differential equation

$$y''_{xx} + \lambda f(y) - \lambda^2 y + \lambda^2 C = 0$$

under the initial conditions

$$y(a) = Ae^{\lambda a} + Be^{-\lambda a} + C, \quad y'_x(a) = \lambda Ae^{\lambda a} - \lambda Be^{-\lambda a}.$$

2°. Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 C u - 2\lambda F(u) + \lambda^2(C^2 - 4AB)]^{-1/2} du = \pm(x - a),$$

$$F(u) = \int_{y_0}^u f(t) dt, \quad y_0 = Ae^{\lambda a} + Be^{-\lambda a} + C.$$

$$13. \quad y(x) + \int_a^x \sinh[\lambda(x-t)]f(y(t)) dt = A \cosh(\lambda x) + B.$$

This is a special case of equation 6.8.12.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 B u - 2\lambda F(u) + \lambda^2 (B^2 - A^2)]^{-1/2} du = \pm(x-a),$$

$$F(u) = \int_{y_0}^u f(t) dt, \quad y_0 = A \cosh(\lambda a) + B.$$

$$14. \quad y(x) + \int_a^x \sinh[\lambda(x-t)]f(y(t)) dt = A \sinh(\lambda x) + B.$$

This is a special case of equation 6.8.21.

Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 u^2 - 2\lambda^2 B u - 2\lambda F(u) + \lambda^2 (A^2 + B^2)]^{-1/2} du = \pm(x-a),$$

$$F(u) = \int_{y_0}^u f(t) dt, \quad y_0 = A \sinh(\lambda a) + B.$$

$$15. \quad y(x) + \int_a^x \sin[\lambda(x-t)]f(y(t)) dt = A \sin(\lambda x) + B \cos(\lambda x) + C.$$

1°. This is a special case of equation 6.8.23. The solution of this integral equation is determined by the solution of the second-order autonomous ordinary differential equation

$$y''_{xx} + \lambda f(y) + \lambda^2 y - \lambda^2 C = 0$$

under the initial conditions

$$y(a) = A \sin(\lambda a) + B \cos(\lambda a) + C, \quad y'_x(a) = A \lambda \cos(\lambda a) - B \lambda \sin(\lambda a).$$

2°. Solution in an implicit form:

$$\int_{y_0}^y [\lambda^2 D - \lambda^2 u^2 + 2\lambda^2 C u - 2\lambda F(u)]^{-1/2} du = \pm(x-a),$$

$$y_0 = A \sin(\lambda a) + B \cos(\lambda a) + C, \quad D = A^2 + B^2 - C^2, \quad F(u) = \int_{y_0}^u f(t) dt.$$

6.8-2. Equations of the Form $y(x) + \int_a^x K(x-t)G(t, y(t)) dt = F(x)$.

$$16. \quad y(x) + \int_a^x f(t, y(t)) dt = g(x).$$

The solution of this integral equation is determined by the solution of the first-order ordinary differential equation

$$y'_x + f(x, y) - g'_x(x) = 0$$

under the initial condition $y(a) = g(a)$. For the exact solutions of the first-order differential equations with various $f(x, y)$ and $g(x)$, see E. Kamke (1977) and A. D. Polyanin and V. F. Zaitsev (2003).

$$17. \quad y(x) + \int_a^x (x-t)f(t, y(t)) dt = g(x).$$

Differentiating the equation with respect to x yields

$$y'_x + \int_a^x f(t, y(t)) dt = g'_x(x). \quad (1)$$

In turn, differentiating this equation with respect to x yields the second-order nonlinear ordinary differential equation

$$y''_{xx} + f(x, y) - g''_{xx}(x) = 0. \quad (2)$$

By setting $x = a$ in the original equation and equation (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = g(a), \quad y'_x(a) = g'_x(a). \quad (3)$$

Equation (2) under conditions (3) defines the solution of the original integral equation. For the exact solutions of the second-order differential equation (2) with various $f(x, y)$ and $g(x)$, see A. D. Polyanin and V. F. Zaitsev (2003).

$$18. \quad y(x) + \int_a^x (x-t)^n f(t, y(t)) dt = g(x), \quad n = 1, 2, \dots$$

Differentiating the equation $n+1$ times with respect to x , we obtain an $(n+1)$ st-order nonlinear ordinary differential equation for $y = y(x)$:

$$y_x^{(n+1)} + n! f(x, y) - g_x^{(n+1)}(x) = 0.$$

This equation under the initial conditions

$$y(a) = g(a), \quad y'_x(a) = g'_x(a), \quad \dots, \quad y_x^{(n)}(a) = g_x^{(n)}(a),$$

defines the solution of the original integral equation.

$$19. \quad y(x) + \int_a^x e^{\lambda(x-t)} f(t, y(t)) dt = g(x).$$

Differentiating the equation with respect to x yields

$$y'_x + f(x, y(x)) + \lambda \int_a^x e^{\lambda(x-t)} f(t, y(t)) dt = g'_x(x).$$

Eliminating the integral term with the aid of the original equation, we obtain the first-order nonlinear ordinary differential equation

$$y'_x + f(x, y) - \lambda y + \lambda g(x) - g'_x(x) = 0.$$

The unknown function $y = y(x)$ must satisfy the initial condition $y(a) = g(a)$. For the exact solutions of the first-order differential equations with various $f(x, y)$ and $g(x)$, see E. Kamke (1977) and A. D. Polyanin and V. F. Zaitsev (2003).

$$20. \quad y(x) + \int_a^x \cosh[\lambda(x-t)]f(t, y(t)) dt = g(x).$$

Differentiating the equation with respect to x twice yields

$$y'_x(x) + f(x, y(x)) + \lambda \int_a^x \sinh[\lambda(x-t)]f(t, y(t)) dt = g'_x(x), \quad (1)$$

$$y''_{xx}(x) + [f(x, y(x))]'_x + \lambda^2 \int_a^x \cosh[\lambda(x-t)]f(t, y(t)) dt = g''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the second-order nonlinear ordinary differential equation

$$y''_{xx} + [f(x, y)]'_x - \lambda^2 y + \lambda^2 g(x) - g''_{xx}(x) = 0. \quad (3)$$

By setting $x = a$ in the original equation and in (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = g(a), \quad y'_x(a) = g'_x(a) - f(a, g(a)). \quad (4)$$

Equation (3) under conditions (4) defines the solution of the original integral equation.

$$21. \quad y(x) + \int_a^x \sinh[\lambda(x-t)]f(t, y(t)) dt = g(x).$$

Differentiating the equation with respect to x twice yields

$$y'_x(x) + \lambda \int_a^x \cosh[\lambda(x-t)]f(t, y(t)) dt = g'_x(x), \quad (1)$$

$$y''_{xx}(x) + \lambda f(x, y(x)) + \lambda^2 \int_a^x \sinh[\lambda(x-t)]f(t, y(t)) dt = g''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the second-order nonlinear ordinary differential equation

$$y''_{xx} + \lambda f(x, y) - \lambda^2 y + \lambda^2 g(x) - g''_{xx}(x) = 0. \quad (3)$$

By setting $x = a$ in the original equation and in (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = g(a), \quad y'_x(a) = g'_x(a). \quad (4)$$

Equation (3) under conditions (4) defines the solution of the original integral equation. For the exact solutions of the second-order differential equation (3) with various $f(x, y)$ and $g(x)$, see A. D. Polyanin and V. F. Zaitsev (2003).

$$22. \quad y(x) + \int_a^x \cos[\lambda(x-t)]f(t, y(t)) dt = g(x).$$

Differentiating the equation with respect to x twice yields

$$y'_x(x) + f(x, y(x)) - \lambda \int_a^x \sin[\lambda(x-t)]f(t, y(t)) dt = g'_x(x), \quad (1)$$

$$y''_{xx}(x) + [f(x, y(x))]'_x - \lambda^2 \int_a^x \cos[\lambda(x-t)]f(t, y(t)) dt = g''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the second-order nonlinear ordinary differential equation

$$y''_{xx} + [f(x, y)]'_x + \lambda^2 y - \lambda^2 g(x) - g''_{xx}(x) = 0. \quad (3)$$

By setting $x = a$ in the original equation and in (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = g(a), \quad y'_x(a) = g'_x(a) - f(a, g(a)). \quad (4)$$

Equation (3) under conditions (4) defines the solution of the original integral equation.

$$23. \quad y(x) + \int_a^x \sin[\lambda(x-t)]f(t, y(t)) dt = g(x).$$

Differentiating the equation with respect to x twice yields

$$y'_x(x) + \lambda \int_a^x \cos[\lambda(x-t)]f(t, y(t)) dt = g'_x(x), \quad (1)$$

$$y''_{xx}(x) + \lambda f(x, y(x)) - \lambda^2 \int_a^x \sin[\lambda(x-t)]f(t, y(t)) dt = g''_{xx}(x). \quad (2)$$

Eliminating the integral term from (2) with the aid of the original equation, we arrive at the second-order nonlinear ordinary differential equation

$$y''_{xx} + \lambda f(x, y) + \lambda^2 y - \lambda^2 g(x) - g''_{xx}(x) = 0. \quad (3)$$

By setting $x = a$ in the original equation and in (1), we obtain the initial conditions for $y = y(x)$:

$$y(a) = g(a), \quad y'_x(a) = g'_x(a). \quad (4)$$

Equation (3) under conditions (4) defines the solution of the original integral equation. For the exact solutions of the second-order differential equation (3) with various $f(x, y)$ and $g(x)$, see A. D. Polyanin and V. F. Zaitsev (2003).

6.8-3. Other Equations.

$$24. \quad y(x) + \frac{1}{x} \int_0^x f\left(\frac{t}{x}, y(t), y(x)\right) dt = A.$$

A solution: $y(x) = \lambda$, where λ is a root of the algebraic (or transcendental) equation

$$\lambda + F(\lambda) - A = 0, \quad F(\lambda) = \int_0^1 f(z, \lambda, \lambda) dz.$$

$$25. \quad y(x) + \int_0^x f\left(\frac{t}{x}, \frac{y(t)}{t}, \frac{y(x)}{x}\right) dt = Ax.$$

A solution: $y(x) = \lambda x$, where λ is a root of the algebraic (or transcendental) equation

$$\lambda + F(\lambda) - A = 0, \quad F(\lambda) = \int_0^1 f(z, \lambda, \lambda) dz.$$

$$26. \quad y(x) + \int_x^\infty f(t-x, y(t-x))y(t) dt = ae^{-\lambda x}.$$

Solutions: $y(x) = b_k e^{-\lambda x}$, where b_k are roots of the algebraic (or transcendental) equation

$$b + bI(b) = a, \quad I(b) = \int_0^\infty f(z, be^{-\lambda z})e^{-\lambda z} dz.$$

Chapter 7

Nonlinear Equations of the First Kind with Constant Limits of Integration

► **Notation:** $f, g, \varphi,$ and ψ are arbitrary functions of an argument specified in the parentheses (the argument can depend on $t, x,$ and y); and $A, B, a, b, c, \beta, \gamma, \lambda,$ and μ are arbitrary parameters.

7.1. Equations with Quadratic Nonlinearity That Contain Arbitrary Parameters

7.1-1. Equations of the Form $\int_a^b K(t)y(x)y(t) dt = F(x).$

1.
$$\int_0^1 y(x)y(t) dt = Ax^\lambda, \quad A > 0, \quad \lambda > -1.$$

This is a special case of equation 7.2.1 with $f(t) = 1, g(x) = Ax^\lambda, a = 0,$ and $b = 1.$

Solutions: $y(x) = \pm \sqrt{A(\lambda + 1)} x^\lambda.$

2.
$$\int_0^1 y(x)y(t) dt = Ae^{\beta x}, \quad A > 0.$$

This is a special case of equation 7.2.1 with $f(t) = 1, g(x) = Ae^{\beta x}, a = 0,$ and $b = 1.$

Solutions: $y(x) = \pm \sqrt{\frac{A\beta}{e^\beta - 1}} e^{\beta x}.$

3.
$$\int_0^1 y(x)y(t) dt = A \cosh(\beta x), \quad A > 0.$$

This is a special case of equation 7.2.1 with $f(t) = 1, g(x) = A \cosh(\beta x), a = 0,$ and $b = 1.$

Solutions: $y(x) = \pm \sqrt{\frac{A\beta}{\sinh \beta}} \cosh(\beta x).$

4.
$$\int_0^1 y(x)y(t) dt = A \sinh(\beta x), \quad A\beta > 0.$$

This is a special case of equation 7.2.1 with $f(t) = 1, g(x) = A \sinh(\beta x), a = 0,$ and $b = 1.$

Solutions: $y(x) = \pm \sqrt{\frac{A\beta}{\cosh \beta - 1}} \sinh(\beta x).$

$$5. \quad \int_0^1 y(x)y(t) dt = A \tanh(\beta x), \quad A\beta > 0.$$

This is a special case of equation 7.2.1 with $f(t) = 1$, $g(x) = A \tanh(\beta x)$, $a = 0$, and $b = 1$.

$$\text{Solutions: } y(x) = \pm \sqrt{\frac{A\beta}{\ln \cosh \beta}} \tanh(\beta x).$$

$$6. \quad \int_0^1 y(x)y(t) dt = A \ln(\beta x), \quad A(\ln \beta - 1) > 0.$$

This is a special case of equation 7.2.1 with $f(t) = 1$, $g(x) = A \ln(\beta x)$, $a = 0$, and $b = 1$.

$$\text{Solutions: } y(x) = \pm \sqrt{\frac{A}{\ln \beta - 1}} \ln(\beta x).$$

$$7. \quad \int_0^1 y(x)y(t) dt = A \cos(\beta x), \quad A > 0.$$

This is a special case of equation 7.2.1 with $f(t) = 1$, $g(x) = A \cos(\beta x)$, $a = 0$, and $b = 1$.

$$\text{Solutions: } y(x) = \pm \sqrt{\frac{A\beta}{\sin \beta}} \cos(\beta x).$$

$$8. \quad \int_0^1 y(x)y(t) dt = A \sin(\beta x), \quad A\beta > 0.$$

This is a special case of equation 7.2.1 with $f(t) = 1$, $g(x) = A \sin(\beta x)$, $a = 0$, and $b = 1$.

$$\text{Solutions: } y(x) = \pm \sqrt{\frac{A\beta}{1 - \cos \beta}} \sin(\beta x).$$

$$9. \quad \int_0^1 y(x)y(t) dt = A \tan(\beta x), \quad A\beta > 0.$$

This is a special case of equation 7.2.1 with $f(t) = 1$, $g(x) = A \tan(\beta x)$, $a = 0$, and $b = 1$.

$$\text{Solutions: } y(x) = \pm \sqrt{\frac{-A\beta}{\ln |\cos \beta|}} \tan(\beta x).$$

$$10. \quad \int_0^1 t^\mu y(x)y(t) dt = Ax^\lambda, \quad A > 0, \quad \mu + \lambda > -1.$$

This is a special case of equation 7.2.1 with $f(t) = t^\mu$, $g(x) = Ax^\lambda$, $a = 0$, and $b = 1$.

$$\text{Solutions: } y(x) = \pm \sqrt{A(\mu + \lambda + 1)} x^\lambda.$$

$$11. \quad \int_0^1 e^{\mu t} y(x)y(t) dt = Ae^{\beta x}, \quad A > 0.$$

This is a special case of equation 7.2.1 with $f(t) = e^{\mu t}$, $g(x) = Ae^{\beta x}$, $a = 0$, and $b = 1$.

$$\text{Solutions: } y(x) = \pm \sqrt{\frac{A(\mu + \beta)}{e^{\mu + \beta} - 1}} e^{\beta x}.$$

7.1-2. Equations of the Form $\int_a^b K(t)y(t)y(xt) dt = F(x)$.

12. $\int_0^1 y(t)y(xt) dt = A, \quad 0 \leq x \leq 1.$

This is a special case of equation 7.2.2 with $f(t) = 1$, $a = 0$, and $b = 1$.

1°. Solutions:

$$\begin{aligned} y_1(x) &= \sqrt{A}, & y_2(x) &= -\sqrt{A}, \\ y_3(x) &= \sqrt{A}(3x-2), & y_4(x) &= -\sqrt{A}(3x-2), \\ y_5(x) &= \sqrt{A}(10x^2-12x+3), & y_6(x) &= -\sqrt{A}(10x^2-12x+3). \end{aligned}$$

2°. The integral equation has some other solutions; for example,

$$\begin{aligned} y_7(x) &= \frac{\sqrt{A}}{C} [(2C+1)x^C - C - 1], & y_8(x) &= -\frac{\sqrt{A}}{C} [(2C+1)x^C - C - 1], \\ y_9(x) &= \sqrt{A}(\ln x + 1), & y_{10}(x) &= -\sqrt{A}(\ln x + 1), \end{aligned}$$

where C is an arbitrary constant.

3°. See 7.2.2 for some other solutions.

13. $\int_0^1 y(t)y(xt^\beta) dt = A, \quad \beta > 0.$

1°. Solutions:

$$\begin{aligned} y_1(x) &= \sqrt{A}, & y_2(x) &= -\sqrt{A}, \\ y_3(x) &= \sqrt{B}[(\beta+2)x - \beta - 1], & y_4(x) &= -\sqrt{B}[(\beta+2)x - \beta - 1], \end{aligned}$$

where $B = \sqrt{\frac{2A}{\beta(\beta+1)}}$.

2°. The integral equation has some other (more complicated solutions) of the polynomial form $y(x) = \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic equations.

14. $\int_1^\infty y(t)y(xt) dt = Ax^{-\lambda}, \quad \lambda > 0, \quad 1 \leq x < \infty.$

This is a special case of equation 7.2.3 with $f(t) = 1$, $a = 1$, and $b = \infty$.

1°. Solutions:

$$\begin{aligned} y_1(x) &= Bx^{-\lambda}, & y_2(x) &= -Bx^{-\lambda}, & \lambda &> \frac{1}{2}; \\ y_3(x) &= B[(2\lambda-3)x - 2\lambda + 2]x^{-\lambda}, & y_4(x) &= -B[(2\lambda-3)x - 2\lambda + 2]x^{-\lambda}, & \lambda &> \frac{3}{2}; \end{aligned}$$

where $B = \sqrt{A(2\lambda-1)}$.

2°. For sufficiently large λ , the integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic equations.

$$15. \int_0^{\infty} e^{-\lambda t} y(t)y(xt) dt = A, \quad \lambda > 0, \quad 0 \leq x < \infty.$$

This is a special case of equation 7.2.2 with $f(t) = e^{-\lambda t}$, $a = 0$, and $b = \infty$.

1°. Solutions:

$$\begin{aligned} y_1(x) &= \sqrt{A\lambda}, & y_2(x) &= -\sqrt{A\lambda}, \\ y_3(x) &= \sqrt{\frac{1}{2}A\lambda}(\lambda x - 2), & y_4(x) &= -\sqrt{\frac{1}{2}A\lambda}(\lambda x - 2). \end{aligned}$$

2°. The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic equations. See 7.2.2 for some other solutions.

7.1-3. Other Equations.

$$16. \int_0^1 y(t)y(x + \lambda t) dt = A, \quad 0 \leq x < \infty.$$

This is a special case of equation 7.2.7 with $f(t) \equiv 1$, $a = 0$, and $b = 1$.

Solutions:

$$\begin{aligned} y_1(x) &= \sqrt{A}, & y_2(x) &= -\sqrt{A}, \\ y_3(x) &= \sqrt{3A/\lambda}(1 - 2x), & y_4(x) &= -\sqrt{3A/\lambda}(1 - 2x). \end{aligned}$$

$$17. \int_0^{\infty} y(t)y(x + \lambda t) dt = A e^{-\beta x}, \quad A, \lambda, \beta > 0, \quad 0 \leq x < \infty.$$

This is a special case of equation 7.2.9 with $f(t) \equiv 1$, $a = 0$, and $b = \infty$.

Solutions:

$$\begin{aligned} y_1(x) &= \sqrt{A\beta(\lambda + 1)} e^{-\beta x}, & y_2(x) &= -\sqrt{A\beta(\lambda + 1)} e^{-\beta x}, \\ y_3(x) &= B[\beta(\lambda + 1)x - 1] e^{-\beta x}, & y_4(x) &= -B[\beta(\lambda + 1)x - 1] e^{-\beta x}, \end{aligned}$$

where $B = \sqrt{A\beta(\lambda + 1)/\lambda}$.

$$18. \int_0^1 y(t)y(x - t) dt = A, \quad -\infty < x < \infty.$$

This is a special case of equation 7.2.10 with $f(t) \equiv 1$, $a = 0$, and $b = 1$.

1°. Solutions with $A > 0$:

$$\begin{aligned} y_1(x) &= \sqrt{A}, & y_2(x) &= -\sqrt{A}, \\ y_3(x) &= \sqrt{5A}(6x^2 - 6x + 1), & y_4(x) &= -\sqrt{5A}(6x^2 - 6x + 1). \end{aligned}$$

2°. Solutions with $A < 0$:

$$y_1(x) = \sqrt{-3A}(1 - 2x), \quad y_2(x) = -\sqrt{-3A}(1 - 2x).$$

The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic equations.

$$19. \int_0^{\infty} e^{-\lambda t} y\left(\frac{x}{t}\right) y(t) dt = Ax^b, \quad \lambda > 0.$$

Solutions: $y(x) = \pm \sqrt{A\lambda} x^b$.

7.2. Equations with Quadratic Nonlinearity That Contain Arbitrary Functions

7.2-1. Equations of the Form $\int_a^b K(t)y(t)y(\cdot\cdot\cdot) dt = F(x)$.

1.
$$\int_a^b f(t)y(x)y(t) dt = g(x).$$

Solutions:

$$y(x) = \pm \lambda g(x), \quad \lambda = \left[\int_a^b f(t)g(t) dt \right]^{-1/2}.$$

2.
$$\int_a^b f(t)y(t)y(xt) dt = A.$$

1°. Solutions:*

$$\begin{aligned} y_1(x) &= \sqrt{A/I_0}, & y_2(x) &= -\sqrt{A/I_0}, \\ y_3(x) &= q(I_1x - I_2), & y_4(x) &= -q(I_1x - I_2), \end{aligned}$$

where

$$I_m = \int_a^b t^m f(t) dt, \quad q = \left(\frac{A}{I_0 I_2^2 - I_1^2 I_2} \right)^{1/2}, \quad m = 0, 1, 2.$$

The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic equations.

2°. Solutions:

$$\begin{aligned} y_5(x) &= q(I_1 x^C - I_2), & y_6(x) &= -q(I_1 x^C - I_2), \\ q &= \left(\frac{A}{I_0 I_2^2 - I_1^2 I_2} \right)^{1/2}, & I_m &= \int_a^b t^{mC} f(t) dt, \quad m = 0, 1, 2, \end{aligned}$$

where C is an arbitrary constant.

The equation has more complicated solutions of the form $y(x) = \sum_{k=0}^n B_k x^{kC}$, where C is an arbitrary constant and the coefficients B_k can be found from the corresponding system of algebraic equations.

3°. Solutions:

$$\begin{aligned} y_7(x) &= p(J_0 \ln x - J_1), & y_8(x) &= -p(J_0 \ln x - J_1), \\ p &= \left(\frac{A}{J_0^2 J_2 - J_0 J_1^2} \right)^{1/2}, & J_m &= \int_a^b (\ln t)^m f(t) dt. \end{aligned}$$

The equation has more complicated solutions of the form $y(x) = \sum_{k=0}^n E_k (\ln x)^k$, where the constants E_k can be found from the corresponding system of algebraic equations.

* The arguments of the equations containing $y(xt)$ in the integrand can vary, for example, within the following intervals: (a) $0 \leq t \leq 1$, $0 \leq x \leq 1$ for $a = 0$ and $b = 1$; (b) $1 \leq t < \infty$, $1 \leq x < \infty$ for $a = 1$ and $b = \infty$; (c) $0 \leq t < \infty$, $0 \leq x < \infty$ for $a = 0$ and $b = \infty$; or (d) $a \leq t \leq b$, $0 \leq x < \infty$ for arbitrary a and b such that $0 \leq a < b \leq \infty$. Case (d) is a special case of (c) if $f(t)$ is nonzero only on the interval $a \leq t \leq b$.

$$3. \int_a^b f(t)y(t)y(xt) dt = Ax^\beta.$$

1°. Solutions:

$$\begin{aligned} y_1(x) &= \sqrt{A/I_0} x^\beta, & y_2(x) &= -\sqrt{A/I_0} x^\beta, \\ y_3(x) &= q(I_1x - I_2) x^\beta, & y_4(x) &= -q(I_1x - I_2) x^\beta, \end{aligned}$$

where

$$I_m = \int_a^b t^{2\beta+m} f(t) dt, \quad q = \sqrt{\frac{A}{I_2(I_0I_2 - I_1^2)}}, \quad m = 0, 1, 2.$$

2°. The substitution $y(x) = x^\beta w(x)$ leads to an equation of the form 7.2.2:

$$\int_a^b g(t)w(t)w(xt) dt = A, \quad g(x) = f(x)x^{2\beta}.$$

Therefore, the integral equation in question has more complicated solutions.

$$4. \int_a^b f(t)y(t)y(xt) dt = A \ln x + B.$$

This equation has solutions of the form $y(x) = p \ln x + q$. The constants p and q are determined from the following system of two second-order algebraic equations:

$$I_1p^2 + I_0pq = A, \quad I_2p^2 + 2I_1pq + I_0q^2 = B,$$

where

$$I_m = \int_a^b f(t)(\ln t)^m dt, \quad m = 0, 1, 2.$$

$$5. \int_a^b f(t)y(t)y(xt) dt = Ax^\lambda \ln x + Bx^\lambda.$$

The substitution $y(x) = x^\lambda w(x)$ leads to an equation of the form 7.2.4:

$$\int_a^b g(t)w(t)w(xt) dt = A \ln x + B, \quad g(t) = f(t)t^{2\lambda}.$$

$$6. \int_0^\infty f(t)y(t)y\left(\frac{x}{t}\right) dt = Ax^\lambda.$$

Solutions:

$$y_1(x) = \sqrt{\frac{A}{I}} x^\lambda, \quad y_2(x) = -\sqrt{\frac{A}{I}} x^\lambda, \quad I = \int_0^\infty f(t) dt.$$

$$7. \int_a^b f(t)y(t)y(x + \lambda t) dt = A, \quad \lambda > 0.$$

1°. Solutions*

$$\begin{aligned} y_1(x) &= \sqrt{A/I_0}, & y_2(x) &= -\sqrt{A/I_0}, \\ y_3(x) &= q(I_0x - I_1), & y_4(x) &= -q(I_0x - I_1), \end{aligned}$$

* The arguments of the equations containing $y(x+\lambda t)$ in the integrand can vary within the following intervals: (a) $0 \leq t < \infty$, $0 \leq x < \infty$ for $a = 0$ and $b = \infty$ or (b) $a \leq t \leq b$, $0 \leq x < \infty$ for arbitrary a and b such that $0 \leq a < b < \infty$. Case (b) is a special case of (a) if $f(t)$ is nonzero only on the interval $a \leq t \leq b$.

where

$$I_m = \int_a^b t^m f(t) dt, \quad q = \sqrt{\frac{A}{\lambda(I_0^2 I_2 - I_0 I_1^2)}}, \quad m = 0, 1, 2.$$

2°. The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic equations.

$$8. \quad \int_a^b f(t)y(t)y(x + \lambda t) dt = Ax + B, \quad \lambda > 0.$$

A solution: $y(x) = \beta x + \mu$, where the constants β and μ are determined from the following system of two second-order algebraic equations:

$$I_0 \beta \mu + I_1 \beta^2 = A, \quad I_0 \mu^2 + (\lambda + 1) I_1 \beta \mu + \lambda I_2 \beta^2 = B, \quad I_m = \int_a^b t^m f(t) dt. \quad (1)$$

Multiplying the first equation by B and the second by $-A$ and adding the resulting equations, we obtain the quadratic equation

$$AI_0 z^2 + [(\lambda + 1)AI_1 - BI_0]z + \lambda AI_2 - BI_1 = 0, \quad z = \mu/\beta. \quad (2)$$

In general, to each root of equation (2) two solutions of system (1) correspond. Therefore, the original integral equation can have at most four solutions of this form. If the discriminant of equation (2) is negative, then the integral equation has no such solutions.

The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n \beta_k x^k$, where the constants β_k can be found from the corresponding system of algebraic equations.

$$9. \quad \int_a^b f(t)y(t)y(x + \lambda t) dt = Ae^{-\beta x}, \quad \lambda > 0.$$

1°. Solutions:

$$\begin{aligned} y_1(x) &= \sqrt{A/I_0} e^{-\beta x}, & y_2(x) &= -\sqrt{A/I_0} e^{-\beta x}, \\ y_3(x) &= q(I_0 x - I_1) e^{-\beta x}, & y_4(x) &= -q(I_0 x - I_1) e^{-\beta x}, \end{aligned}$$

where

$$I_m = \int_a^b t^m e^{-\beta(\lambda+1)t} f(t) dt, \quad q = \sqrt{\frac{A}{\lambda(I_0^2 I_2 - I_0 I_1^2)}}, \quad m = 0, 1, 2.$$

2°. The equation has more complicated solutions of the form $y(x) = e^{-\beta x} \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic equations.

3°. The substitution $y(x) = e^{-\beta x} w(x)$ leads to an equation of the form 7.2.7:

$$\int_a^b e^{-\beta(\lambda+1)t} f(t)w(t)w(x + \lambda t) dt = A.$$

$$10. \int_a^b f(t)y(t)y(x-t) dt = A.$$

1°. Solutions*

$$\begin{aligned} y_1(x) &= \sqrt{A/I_0}, & y_2(x) &= -\sqrt{A/I_0}, \\ y_3(x) &= q(I_0x - I_1), & y_4(x) &= -q(I_0x - I_1), \end{aligned}$$

where

$$I_m = \int_a^b t^m f(t) dt, \quad q = \sqrt{\frac{A}{I_0I_1^2 - I_0^2I_2}}, \quad m = 0, 1, 2.$$

2°. The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n \lambda_k x^k$, where the constants λ_k can be found from the corresponding system of algebraic equations. For $n = 3$, such a solution is presented in 7.1.18.

$$11. \int_a^b f(t)y(t)y(x-t) dt = Ax + B.$$

A solution: $y(x) = \lambda x + \mu$, where the constants λ and μ are determined from the following system of two second-order algebraic equations:

$$I_0\lambda\mu + I_1\lambda^2 = A, \quad I_0\mu^2 - I_2\lambda^2 = B, \quad I_m = \int_a^b t^m f(t) dt, \quad m = 0, 1, 2. \quad (1)$$

Multiplying the first equation by B and the second by $-A$ and adding the results, we obtain the quadratic equation

$$AI_0z^2 - BI_0z - AI_2 - BI_1 = 0, \quad z = \mu/\lambda. \quad (2)$$

In general, to each root of equation (2) two solutions of system (1) correspond. Therefore, the original integral equation can have at most four solutions of this form. If the discriminant of equation (2) is negative, then the integral equation has no such solutions.

The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n \lambda_k x^k$, where the constants λ_k can be found from the corresponding system of algebraic equations.

$$12. \int_a^b f(t)y(t)y(x-t) dt = \sum_{k=0}^n A_k x^k.$$

This equation has solutions of the form

$$y(x) = \sum_{k=0}^n \lambda_k x^k, \quad (1)$$

where the constants λ_k are determined from the system of algebraic equations obtained by substituting solution (1) into the original integral equation and matching the coefficients of like powers of x .

* The arguments of the equations containing $y(x-t)$ in the integrand can vary within the following intervals: (a) $-\infty < t < \infty$, $-\infty < x < \infty$ for $a = -\infty$ and $b = \infty$ or (b) $a \leq t \leq b$, $-\infty \leq x < \infty$, for arbitrary a and b such that $-\infty < a < b < \infty$. Case (b) is a special case of (a) if $f(t)$ is nonzero only on the interval $a \leq t \leq b$.

$$13. \int_a^b f(t)y(x-t)y(t) dt = Ae^{\lambda x}.$$

Solutions:

$$\begin{aligned} y_1(x) &= \sqrt{A/I_0} e^{\lambda x}, & y_2(x) &= -\sqrt{A/I_0} e^{\lambda x}, \\ y_3(x) &= q(I_0x - I_1)e^{\lambda x}, & y_4(x) &= -q(I_0x - I_1)e^{\lambda x}, \end{aligned}$$

where

$$I_m = \int_a^b t^m f(t) dt, \quad q = \sqrt{\frac{A}{I_0I_1^2 - I_0^2I_2}}, \quad m = 0, 1, 2.$$

The integral equation has more complicated solutions of the form $y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic equations.

$$14. \int_a^b f(t)y(t)y(x-t) dt = A \sinh \lambda x.$$

A solution:

$$y(x) = p \sinh \lambda x + q \cosh \lambda x. \quad (1)$$

Here p and q are roots of the algebraic system

$$I_0pq + I_{cs}(p^2 - q^2) = A, \quad I_{cc}q^2 - I_{ss}p^2 = 0, \quad (2)$$

where the notation

$$\begin{aligned} I_0 &= \int_a^b f(t) dt, & I_{cs} &= \int_a^b f(t) \cosh(\lambda t) \sinh(\lambda t) dt, \\ I_{cc} &= \int_a^b f(t) \cosh^2(\lambda t) dt, & I_{ss} &= \int_a^b f(t) \sinh^2(\lambda t) dt \end{aligned}$$

is used. Different solutions of system (2) generate different solutions (1) of the integral equation.

It follows from the second equation of (2) that $q = \pm \sqrt{I_{ss}/I_{cc}} p$. Using this expression to eliminate q from the first equation of (2), we obtain the following four solutions:

$$\begin{aligned} y_{1,2}(x) &= p(\sinh \lambda x \pm k \cosh \lambda x), & y_{3,4}(x) &= -p(\sinh \lambda x \pm k \cosh \lambda x), \\ k &= \sqrt{\frac{I_{ss}}{I_{cc}}}, & p &= \sqrt{\frac{A}{(1-k^2)I_{cs} \pm kI_0}}. \end{aligned}$$

$$15. \int_a^b f(t)y(t)y(x-t) dt = A \cosh \lambda x.$$

A solution:

$$y(x) = p \sinh \lambda x + q \cosh \lambda x. \quad (1)$$

Here p and q are roots of the algebraic system

$$I_0pq + I_{cs}(p^2 - q^2) = 0, \quad I_{cc}q^2 - I_{ss}p^2 = A, \quad (2)$$

where we use the notation introduced in 7.2.14. Different solutions of system (2) generate different solutions (1) of the integral equation.

$$16. \int_a^b f(t)y(t)y(x-t) dt = A \sin \lambda x.$$

A solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x. \quad (1)$$

Here p and q are roots of the algebraic system

$$I_0 p q + I_{cs}(p^2 + q^2) = A, \quad I_{cc} q^2 - I_{ss} p^2 = 0, \quad (2)$$

where

$$I_0 = \int_a^b f(t) dt, \quad I_{cs} = \int_a^b f(t) \cos(\lambda t) \sin(\lambda t) dt,$$

$$I_{cc} = \int_a^b f(t) \cos^2(\lambda t) dt, \quad I_{ss} = \int_a^b f(t) \sin^2(\lambda t) dt.$$

It follows from the second equation of (2) that $q = \pm \sqrt{I_{ss}/I_{cc}} p$. Using this expression to eliminate q from the first equation of (2), we obtain the following four solutions:

$$y_{1,2}(x) = p(\sin \lambda x \pm k \cos \lambda x), \quad y_{3,4}(x) = -p(\sin \lambda x \pm k \cos \lambda x),$$

$$k = \sqrt{\frac{I_{ss}}{I_{cc}}}, \quad p = \sqrt{\frac{A}{(1+k^2)I_{cs} \pm kI_0}}.$$

$$17. \int_a^b f(t)y(t)y(x-t) dt = A \cos \lambda x.$$

A solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x. \quad (1)$$

Here p and q are roots of the algebraic system

$$I_0 p q + I_{cs}(p^2 + q^2) = 0, \quad I_{cc} q^2 - I_{ss} p^2 = A, \quad (2)$$

where we use the notation introduced in 7.2.16. Different solutions of system (2) generate different solutions (1) of the integral equation.

$$18. \int_0^1 y(t)y(\xi) dt = A, \quad \xi = f(x)t.$$

1°. Solutions:

$$y_1(t) = \sqrt{A}, \quad y_2(t) = -\sqrt{A},$$

$$y_3(t) = \sqrt{A}(3t-2), \quad y_4(t) = -\sqrt{A}(3t-2),$$

$$y_5(t) = \sqrt{A}(10t^2-12t+3), \quad y_6(t) = -\sqrt{A}(10t^2-12t+3).$$

2°. The integral equation has some other (more complicated) solutions of the polynomial form $y(t) = \sum_{k=0}^n B_k t^k$, where the constants B_k can be found from the corresponding system of algebraic equations.

3°. The substitution $z = f(x)$ leads to an equation of the form 7.1.12.

7.2-2. Equations of the Form $\int_a^b [K(x, t)y(t) + M(x, t)y^2(t)] dt = F(x)$.

$$19. \int_0^a \left[\frac{1}{|x-t|^k} y(t) + \varphi(x)\psi(t)y^2(t) \right] dt = f(x), \quad 0 < k < 1.$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.1.30.

$$20. \int_a^b \left[\ln|x-t|y(t) + \varphi(x)\psi(t)y^2(t) \right] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.4.2.

$$21. \int_0^\infty [\sin(xt)y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.5.8.

Solutions:

$$y_{1,2}(t) = Y_f(t) + A_{1,2}Y_\varphi(t),$$

where

$$Y_f(t) = \frac{2}{\pi} \int_0^\infty \sin(xt)f(x) dx, \quad Y_\varphi(t) = \frac{2}{\pi} \int_0^\infty \sin(xt)\varphi(x) dx,$$

and $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0,$$

$$p = \int_0^\infty \psi(t)Y_\varphi^2(t) dt, \quad q = 1 + 2 \int_0^\infty \psi(t)Y_f(t)Y_\varphi(t) dt, \quad r = \int_0^\infty \psi(t)Y_f^2(t) dt.$$

⊙ Reference: A. D. Polyanin and A. I. Zhurov (2007).

$$22. \int_0^\infty [\cos(xt)y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.5.1.

Solutions:

$$y_{1,2}(t) = Y_f(t) + A_{1,2}Y_\varphi(t),$$

where

$$Y_f(t) = \frac{2}{\pi} \int_0^\infty \cos(xt)f(x) dx, \quad Y_\varphi(t) = \frac{2}{\pi} \int_0^\infty \cos(xt)\varphi(x) dx,$$

and $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0,$$

$$p = \int_0^\infty \psi(t)Y_\varphi^2(t) dt, \quad q = 1 + 2 \int_0^\infty \psi(t)Y_f(t)Y_\varphi(t) dt, \quad r = \int_0^\infty \psi(t)Y_f^2(t) dt.$$

$$23. \int_0^{\infty} [tJ_{\nu}(xt)y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x), \quad \nu > -1.$$

Here $J_{\nu}(z)$ is the Bessel function of the first kind. The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.7.17.

Solutions:

$$y_{1,2}(t) = Y_f(t) + A_{1,2}Y_{\varphi}(t),$$

where

$$Y_f(t) = \int_0^{\infty} xJ_{\nu}(xt)f(x) dx, \quad Y_{\varphi}(t) = \int_0^{\infty} xJ_{\nu}(xt)\varphi(x) dx,$$

and $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0,$$

$$p = \int_0^{\infty} \psi(t)Y_{\varphi}^2(t) dt, \quad q = 1 + 2 \int_0^{\infty} \psi(t)Y_f(t)Y_{\varphi}(t) dt, \quad r = \int_0^{\infty} \psi(t)Y_f^2(t) dt.$$

7.3. Equations with Power-Law Nonlinearity That Contain Arbitrary Functions

7.3-1. Equations of the Form $\int_a^b K(t)y^{\mu}(x)y^{\gamma}(t) dt = F(x)$.

$$1. \int_a^b t^{\lambda}y^{\mu}(x)y^{\gamma}(t) dt = g(x).$$

A solution:

$$y(x) = A[g(x)]^{\frac{1}{\mu}}, \quad A = \left\{ \int_a^b t^{\lambda} [g(t)]^{\frac{\gamma}{\mu}} dt \right\}^{-\frac{1}{\mu+\gamma}}.$$

$$2. \int_a^b e^{\lambda t}y^{\mu}(x)y^{\gamma}(t) dt = g(x).$$

A solution:

$$y(x) = A[g(x)]^{\frac{1}{\mu}}, \quad A = \left\{ \int_a^b e^{\lambda t} [g(t)]^{\frac{\gamma}{\mu}} dt \right\}^{-\frac{1}{\mu+\gamma}}.$$

7.3-2. Equations of the Form $\int_a^b K(t)y^{\gamma}(t)y(xt) dt = F(x)$.

$$3. \int_a^b f(t)y^{\gamma}(t)y(xt) dt = A.$$

This is a special case of equation 7.4.4.

$$4. \int_a^b f(t)y^{\gamma}(t)y(xt) dt = Ax + B.$$

This is a special case of equation 7.4.5.

$$5. \int_a^b f(t)y^\gamma(t)y(xt) dt = Ax^\beta.$$

This equation has solutions of the form $y(x) = kx^\beta$, where k is a constant.

$$6. \int_a^b f(t)y^\gamma(t)y(xt) dt = A \ln x + B.$$

This equation has solutions of the form $y(x) = p \ln x + q$, where p and q are some constants.

$$7. \int_a^b f(t)y^\gamma(t)y(xt) dt = Ax^\beta \ln x.$$

This equation has solutions of the form $y(x) = px^\beta \ln x + qx^\beta$, where p and q are some constants.

$$8. \int_a^b f(t)y^\gamma(t)y(xt) dt = A \cos(\beta \ln x).$$

This equation has solutions of the form $y(x) = p \cos(\beta \ln x) + q \sin(\beta \ln x)$, where p and q are some constants.

$$9. \int_a^b f(t)y^\gamma(t)y(xt) dt = A \sin(\beta \ln x).$$

This equation has solutions of the form $y(x) = p \cos(\beta \ln x) + q \sin(\beta \ln x)$, where p and q are some constants.

7.3-3. Equations of the Form $\int_a^b K(t)y^\gamma(t)y(x + \beta t) dt = F(x)$.

$$10. \int_a^b f(t)y^\gamma(t)y(x + \beta t) dt = Ax + B.$$

This is a special case of equation 7.4.16.

$$11. \int_a^b f(t)y^\gamma(t)y(x + \beta t) dt = Ae^{-\lambda x}.$$

This is a special case of equation 7.4.17.

$$12. \int_a^b f(t)y^\gamma(t)y(x + \beta t) dt = A \cos \lambda x.$$

This equation has solutions of the form $y(x) = p \sin \lambda x + q \cos \lambda x$, where p and q are some constants.

$$13. \int_a^b f(t)y^\gamma(t)y(x + \beta t) dt = A \sin \lambda x.$$

This equation has solutions of the form $y(x) = p \sin \lambda x + q \cos \lambda x$, where p and q are some constants.

$$14. \int_a^b f(t)y^\gamma(t)y(x + \beta t) dt = e^{-\mu x}(A \cos \lambda x + B \sin \lambda x).$$

This equation has solutions of the form $y(x) = e^{-\mu x}(p \sin \lambda x + q \cos \lambda x)$, where p and q are some constants.

7.3-4. Equations of the Form $\int_a^b [K(x, t)y(t) + M(x, t)y^\gamma(t)] dt = f(x)$.

$$15. \int_0^a \left[\frac{1}{\sqrt{|x-t|}} y(t) + \varphi(x)\psi(t)y^\gamma(t) \right] dt = f(x), \quad 0 < a \leq \infty.$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.1.22.

$$16. \int_0^a \left[\frac{1}{|x-t|^k} y(t) + \varphi(x)\psi(t)y^\gamma(t) \right] dt = f(x), \quad 0 < k < 1.$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.1.30.

$$17. \int_a^b \left[\ln|x-t|y(t) + \varphi(x)\psi(t)y^\gamma(t) \right] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.4.2.

$$18. \int_0^\infty [\sin(xt)y(t) + \varphi(x)\psi(t)y^\gamma(t)] dt = f(x).$$

This is a special case of equation 7.4.24.

$$19. \int_0^\infty [\cos(xt)y(t) + \varphi(x)\psi(t)y^\gamma(t)] dt = f(x).$$

This is a special case of equation 7.4.25.

7.3-5. Other Equations.

$$20. \int_0^\infty f(x^a t)t^b y^\gamma(t) y(x^k t) dt = Ax^c.$$

A solution:

$$y(x) = \left(\frac{A}{I} \right)^{\frac{1}{\gamma+1}} x^\lambda, \quad \lambda = \frac{a+c+ab}{k-a-a\gamma},$$

$$I = \int_0^\infty f(t)t^\beta dt, \quad \beta = \frac{a+c+a\gamma+bk+c\gamma}{k-a-a\gamma}.$$

$$21. \int_0^1 [y(xt) + \varphi(x)\psi(t)y^\gamma(t)] dt = f(x).$$

This is a special case of equation 7.4.27.

$$22. \int_0^{\pi/2} [y(x \sin t) + \varphi(x)\psi(t)y^\gamma(t)] dt = f(x).$$

This is a special case of equation 7.4.28.

7.4. Equations with Nonlinearity of General Form

7.4-1. Equations of the Form $\int_a^b \varphi(y(x))K(t, y(t)) dt = F(x)$.

1.
$$\int_a^b y(x)f(t, y(t)) dt = g(x).$$

A solution: $y(x) = \lambda g(x)$, where λ is determined by the algebraic (or transcendental) equation

$$\lambda \int_a^b f(t, \lambda g(t)) dt = 1.$$

2.
$$\int_a^b y^k(x)f(t, y(t)) dt = g(x).$$

A solution: $y(x) = \lambda[g(x)]^{1/k}$, where λ is determined from the algebraic (or transcendental)

equation $\lambda^k \int_a^b f(t, \lambda g^{1/k}(t)) dt = 1$.

3.
$$\int_a^b \varphi(y(x))f(t, y(t)) dt = g(x).$$

A solution in an implicit form:

$$\lambda \varphi(y(x)) - g(x) = 0, \tag{1}$$

where λ is determined by the algebraic (or transcendental) equation

$$\lambda - F(\lambda) = 0, \quad F(\lambda) = \int_a^b f(t, y(t)) dt. \tag{2}$$

Here the function $y(x) = y(x, \lambda)$ obtained by solving (1) must be substituted into (2).

The number of solutions of the integral equation is determined by the number of the solutions obtained from (1) and (2).

7.4-2. Equations of the Form $\int_a^b y(xt)K(t, y(t)) dt = F(x)$.

4.
$$\int_a^b y(xt)f(t, y(t)) dt = A.$$

1°. Solutions: $y(x) = \lambda_k$, where λ_k are roots of the algebraic (or transcendental) equation

$$\lambda \int_a^b f(t, \lambda) dt = A.$$

2°. Solutions: $y(x) = px + q$, where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\int_a^b tf(t, pt + q) dt = 0, \quad q \int_a^b f(t, pt + q) dt = A.$$

In the case $f(t, y(t)) = \bar{f}(t)y(t)$, see 7.2.2 for solutions of this system.

2°. The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic (or transcendental) equations.

4°. The integral equation can have logarithmic solutions similar to those presented in item 3° of equation 7.2.2.

$$5. \quad \int_a^b y(xt) f(t, y(t)) dt = Ax + B.$$

1°. A solution:

$$y(x) = px + q, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$p \int_a^b t f(t, pt + q) dt - A = 0, \quad q \int_a^b f(t, pt + q) dt - B = 0. \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

2°. The integral equation has some other (more complicated) solutions of the polynomial form $y(x) = \sum_{k=0}^n B_k x^k$, where the constants B_k can be found from the corresponding system of algebraic (or transcendental) equations.

$$6. \quad \int_a^b y(xt) f(t, y(t)) dt = Ax^\beta.$$

A solution:

$$y(x) = kx^\beta, \quad (1)$$

where k is a root of the algebraic (or transcendental) equation

$$kF(k) - A = 0, \quad F(k) = \int_a^b t^\beta f(t, kt^\beta) dt. \quad (2)$$

Each root of equation (2) generates a solution of the integral equation which has the form (1).

$$7. \quad \int_a^b y(xt) f(t, y(t)) dt = A \ln x + B.$$

A solution:

$$y(x) = p \ln x + q, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$p \int_a^b f(t, p \ln t + q) dt - A = 0, \quad \int_a^b (p \ln t + q) f(t, p \ln t + q) dt - B = 0. \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$8. \quad \int_a^b y(xt) f(t, y(t)) dt = Ax^\beta \ln x.$$

This equation has solutions of the form $y(x) = px^\beta \ln x + qx^\beta$, where p and q are some constants.

$$9. \quad \int_a^b y(xt) f(t, y(t)) dt = A \cos(\beta \ln x).$$

This equation has solutions of the form $y(x) = p \cos(\beta \ln x) + q \sin(\beta \ln x)$, where p and q are some constants.

$$10. \int_a^b y(xt) f(t, y(t)) dt = A \sin(\beta \ln x).$$

This equation has solutions of the form $y(x) = p \cos(\beta \ln x) + q \sin(\beta \ln x)$, where p and q are some constants.

$$11. \int_a^b y(xt) f(t, y(t)) dt = Ax^\beta \cos(\beta \ln x) + Bx^\beta \sin(\beta \ln x).$$

This equation has solutions of the form $y(x) = px^\beta \cos(\beta \ln x) + qx^\beta \sin(\beta \ln x)$, where p and q are some constants.

7.4-3. Equations of the Form $\int_a^b y(x + \beta t) K(t, y(t)) dt = F(x)$.

$$12. \int_a^b y(x - t) f(t, y(t)) dt = Ax + B.$$

This equation has solutions of the form $y(x) = px + q$, where p and q are some constants.

$$13. \int_a^b y(x - t) f(t, y(t)) dt = Ae^{\lambda x}.$$

This equation has solutions of the form $y(x) = pe^{\lambda x}$, where p is some constant.

$$14. \int_a^b y(x - t) f(t, y(t)) dt = A \cos \lambda x.$$

This equation has solutions of the form $y(x) = p \sin \lambda x + q \cos \lambda x$, where p and q are some constants.

$$15. \int_a^b y(x - t) f(t, y(t)) dt = e^{-\mu x} (A \cos \lambda x + B \sin \lambda x).$$

This equation has solutions of the form $y(x) = e^{-\mu x} (p \sin \lambda x + q \cos \lambda x)$, where p and q are some constants.

$$16. \int_a^b y(x + \beta t) f(t, y(t)) dt = Ax + B, \quad \beta > 0.$$

A solution:

$$y(x) = px + q, \tag{1}$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$p \int_a^b f(t, pt + q) dt - A = 0, \quad \int_a^b (\beta pt + q) f(t, pt + q) dt - B = 0. \tag{2}$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$17. \int_a^b y(x + \beta t) f(t, y(t)) dt = Ae^{-\lambda x}, \quad \beta > 0.$$

Solutions:

$$y(x) = k_n e^{-\lambda x},$$

where k_n are roots of the algebraic (or transcendental) equation

$$kF(k) - A = 0, \quad F(k) = \int_a^b f(t, ke^{-\lambda t}) e^{-\beta \lambda t} dt.$$

$$18. \int_a^b y(x + \beta t) f(t, y(t)) dt = A \cos \lambda x, \quad \beta > 0.$$

This equation has solutions of the form $y(x) = p \sin \lambda x + q \cos \lambda x$, where p and q are some constants.

$$19. \int_a^b y(x + \beta t) f(t, y(t)) dt = A \sin \lambda x, \quad \beta > 0.$$

This equation has solutions of the form $y(x) = p \sin \lambda x + q \cos \lambda x$, where p and q are some constants.

$$20. \int_a^b y(x + \beta t) f(t, y(t)) dt = e^{-\mu x} (A \cos \lambda x + B \sin \lambda x), \quad \beta > 0.$$

This equation has solutions of the form $y(x) = e^{-\mu x} (p \sin \lambda x + q \cos \lambda x)$, where p and q are some constants.

7.4-4. Equations of the Form $\int_a^b [K(x, t)y(t) + \varphi(x)\Psi(t, y(t))] dt = F(x)$.

$$21. \int_0^a \left[\frac{1}{\sqrt{|x-t|}} y(t) + \varphi(x)\Psi(t, y(t)) \right] dt = f(x), \quad 0 < a \leq \infty.$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.1.22.

$$22. \int_0^a \left[\frac{1}{|x-t|^k} y(t) + \varphi(x)\Psi(t, y(t)) \right] dt = f(x), \quad 0 < k < 1.$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.1.30.

$$23. \int_a^b \left[\ln|x-t| y(t) + \varphi(x)\Psi(t, y(t)) \right] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.4.2.

$$24. \int_0^\infty [\sin(xt)y(t) + \varphi(x)\Psi(t, y(t))] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.5.8.

Solutions:

$$y_m(t) = Y_f(t) + A_m Y_\varphi(t),$$

where

$$Y_f(t) = \frac{2}{\pi} \int_0^\infty \sin(xt) f(x) dx, \quad Y_\varphi(t) = \frac{2}{\pi} \int_0^\infty \sin(xt) \varphi(x) dx,$$

and A_m are roots of the algebraic (transcendental) equation

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

$$25. \int_0^{\infty} [\cos(xt)y(t) + \varphi(x)\Psi(t, y(t))] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.5.1.

Solutions:

$$y_m(t) = Y_f(t) + A_m Y_\varphi(t),$$

where

$$Y_f(t) = \frac{2}{\pi} \int_0^{\infty} \cos(xt)f(x) dx, \quad Y_\varphi(t) = \frac{2}{\pi} \int_0^{\infty} \cos(xt)\varphi(x) dx,$$

and A_m are roots of the algebraic (transcendental) equation

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

$$26. \int_0^{\infty} [tJ_\nu(xt)y(t) + \varphi(x)\Psi(t, y(t))] dt = f(x), \quad \nu > -1.$$

Here $J_\nu(z)$ is the Bessel function of the first kind. The solutions can be obtained by the methods described in Subsection 16.4-4; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.7.17.

Solutions:

$$y_m(t) = Y_f(t) + A_m Y_\varphi(t),$$

where

$$Y_f(t) = \int_0^{\infty} xJ_\nu(xt)f(x) dx, \quad Y_\varphi(t) = \int_0^{\infty} xJ_\nu(xt)\varphi(x) dx,$$

and A_m are roots of the algebraic (transcendental) equation

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

7.4-5. Other Equations.

$$27. \int_0^1 [y(xt) + \varphi(x)\Psi(t, y(t))] dt = f(x).$$

Solutions:

$$y_m(t) = Y_f(t) + A_m Y_\varphi(t),$$

where

$$Y_f(t) = t f'_t(t) + f(t), \quad Y_\varphi(t) = t \varphi'_t(t) + \varphi(t),$$

and A_m are roots of the algebraic (transcendental) equation

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

The functions $f(x)$ and $\varphi(x)$ are assumed to satisfy the conditions $[xf(x)]_{x=0} = [x\varphi(x)]_{x=0} = 0$.

$$28. \int_0^{\pi/2} [y(x \sin t) + \varphi(x)\Psi(t, y(t))] dt = f(x).$$

For $\varphi(x) = 0$, it is the Schlömilch equation, see Eq. 3.5.40.

Solutions:

$$y_m(z) = Y_f(z) + A_m Y_\varphi(z),$$

where

$$Y_f(z) = \frac{2}{\pi} \left[f(0) + z \int_0^{\pi/2} f'_\xi(\xi) d\tau \right], \quad Y_\varphi(z) = \frac{2}{\pi} \left[\varphi(0) + z \int_0^{\pi/2} \varphi'_\xi(\xi) d\tau \right], \quad \xi = z \sin \tau,$$

and A_m are roots of the algebraic (transcendental) equation

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

⊙ Reference: A. D. Polyanin and A. I. Zhurov (2007).

Chapter 8

Nonlinear Equations of the Second Kind with Constant Limits of Integration

► **Notation:** $f, g, h, \varphi, \Psi,$ and ψ are arbitrary functions of an argument specified in the parentheses (the argument can depend on $t, x,$ and y); and $A, B, C, a, b, \beta, \gamma, \lambda,$ and μ are arbitrary parameters.

8.1. Equations with Quadratic Nonlinearity That Contain Arbitrary Parameters

8.1-1. Equations of the Form $y(x) + \int_a^b K(x, t)y^2(t) dt = F(x)$.

1. $y(x) + A \int_a^b x^\lambda y^2(t) dt = 0.$

Solutions:

$$y_1(x) = 0, \quad y_2(x) = -\frac{2\lambda + 1}{A(b^{2\lambda+1} - a^{2\lambda+1})} x^\lambda.$$

2. $y(x) + A \int_a^b x^\lambda t^\mu y^2(t) dt = 0.$

Solutions:

$$y_1(x) = 0, \quad y_2(x) = -\frac{2\lambda + \mu + 1}{A(b^{2\lambda+\mu+1} - a^{2\lambda+\mu+1})} x^\lambda.$$

3. $y(x) + A \int_a^b e^{-\lambda x} y^2(t) dt = 0.$

Solutions:

$$y_1(x) = 0, \quad y_2(x) = \frac{2\lambda}{A(e^{-2\lambda b} - e^{-2\lambda a})} e^{-\lambda x}.$$

4. $y(x) + A \int_a^b e^{-\lambda x - \mu t} y^2(t) dt = 0.$

Solutions:

$$y_1(x) = 0, \quad y_2(x) = \frac{2\lambda + \mu}{A[e^{-(2\lambda+\mu)b} - e^{-(2\lambda+\mu)a}]} e^{-\lambda x}.$$

$$5. \quad y(x) + A \int_a^b x^\lambda e^{-\mu t} y^2(t) dt = 0.$$

This is a special case of equation 8.2.2 with $f(x) = Ax^\lambda$ and $g(t) = e^{-\mu t}$.

$$6. \quad y(x) + A \int_a^b e^{-\mu x} t^\lambda y^2(t) dt = 0.$$

This is a special case of equation 8.2.2 with $f(x) = Ae^{-\mu x}$ and $g(t) = t^\lambda$.

$$7. \quad y(x) + A \int_0^1 y^2(t) dt = Bx^\mu, \quad \mu > -1.$$

This is a special case of equation 8.2.4 with $g(t) = A$, $f(x) = Bx^\mu$, $a = 0$, and $b = 1$.

A solution: $y(x) = Bx^\mu + \lambda$, where λ is determined by the quadratic equation

$$\lambda^2 + \frac{1}{A} \left(1 + \frac{2AB}{\mu + 1} \right) \lambda + \frac{B^2}{2\mu + 1} = 0.$$

$$8. \quad y(x) + A \int_a^b t^\beta y^2(t) dt = Bx^\mu.$$

This is a special case of equation 8.2.4 with $g(t) = At^\beta$ and $f(x) = Bx^\mu$.

$$9. \quad y(x) + A \int_a^b e^{\beta t} y^2(t) dt = Be^{\mu x}.$$

This is a special case of equation 8.2.4 with $g(t) = Ae^{\beta t}$ and $f(x) = Be^{\mu x}$.

$$10. \quad y(x) + A \int_a^b x^\beta y^2(t) dt = Bx^\mu.$$

This is a special case of equation 8.2.5 with $g(x) = Ax^\beta$ and $f(x) = Bx^\mu$.

$$11. \quad y(x) + A \int_a^b e^{\beta x} y^2(t) dt = Be^{\mu x}.$$

This is a special case of equation 8.2.5 with $g(x) = Ae^{\beta x}$ and $f(x) = Be^{\mu x}$.

8.1-2. Equations of the Form $y(x) + \int_a^b K(x, t)y(x)y(t) dt = F(x)$.

$$12. \quad y(x) + A \int_a^b t^\beta y(x)y(t) dt = Bx^\mu.$$

This is a special case of equation 8.2.7 with $g(t) = At^\beta$ and $f(x) = Bx^\mu$.

$$13. \quad y(x) + A \int_a^b e^{\beta t} y(x)y(t) dt = Be^{\mu x}.$$

This is a special case of equation 8.2.7 with $g(t) = Ae^{\beta t}$ and $f(x) = Be^{\mu x}$.

$$14. \quad y(x) + A \int_a^b x^\beta y(x)y(t) dt = Bx^\mu.$$

This is a special case of equation 8.2.8 with $g(x) = Ax^\beta$ and $f(x) = Bx^\mu$.

$$15. \quad y(x) + A \int_a^b e^{\beta x} y(x)y(t) dt = Be^{\mu x}.$$

This is a special case of equation 8.2.8 with $g(x) = Ae^{\beta x}$ and $f(x) = Be^{\mu x}$.

8.1-3. Equations of the Form $y(x) + \int_a^b K(t)y(t)y(\cdot \cdot \cdot) dt = F(x)$.

16. $y(x) + A \int_0^1 y(t)y(xt) dt = 0$.

This is a special case of equation 8.2.16 with $f(t) = A$, $a = 0$, and $b = 1$.

1°. Solutions:

$$y_1(x) = -\frac{1}{A}(2C+1)x^C, \quad y_2(x) = \frac{(I_1 - I_0)x + I_1 - I_2}{I_0I_2 - I_1^2}x^C,$$

$$I_m = \frac{A}{2C + m + 1}, \quad m = 0, 1, 2,$$

where C is an arbitrary nonnegative constant.

There are more complicated solutions of the form $y(x) = x^C \sum_{k=0}^n B_k x^k$, where C is an arbitrary constant and the coefficients B_k can be found from the corresponding system of algebraic equations.

2°. A solution:

$$y_3(x) = \frac{(I_1 - I_0)x^\beta + I_1 - I_2}{I_0I_2 - I_1^2}x^C, \quad I_m = \frac{A}{2C + m\beta + 1}, \quad m = 0, 1, 2,$$

where C and β are arbitrary constants.

There are more complicated solutions of the form $y(x) = x^C \sum_{k=0}^n D_k x^{k\beta}$, where C and β are arbitrary constants and the coefficients D_k can be found from the corresponding system of algebraic equations.

3°. A solution:

$$y_4(x) = \frac{x^C(J_1 \ln x - J_2)}{J_0J_2 - J_1^2}, \quad J_m = \int_0^1 t^{2C} (\ln t)^m dt, \quad m = 0, 1, 2,$$

where C is an arbitrary constant.

There are more complicated solutions of the form $y(x) = x^C \sum_{k=0}^n E_k (\ln x)^k$, where C is an arbitrary constant and the coefficients E_k can be found from the corresponding system of algebraic equations.

17. $y(x) + A \int_1^\infty y(t)y(xt) dt = 0$.

This is a special case of equation 8.2.16 with $f(t) = A$, $a = 1$, and $b = \infty$.

18. $y(x) + \lambda \int_1^\infty y(t)y(xt) dt = Ax^\beta$.

This is a special case of equation 8.2.17 with $f(t) = \lambda$, $a = 0$, and $b = 1$.

$$19. \quad y(x) + A \int_0^1 y(t)y(x + \lambda t) dt = 0.$$

This is a special case of equation 8.2.21 with $f(t) \equiv A$, $a = 0$, and $b = 1$.

1°. A solution:

$$y(x) = \frac{C(\lambda + 1)}{A[1 - e^{C(\lambda+1)}]} e^{Cx},$$

where C is an arbitrary constant.

2°. There are more complicated solutions of the form $y(x) = e^{Cx} \sum_{m=0}^n B_m x^m$, where C is an arbitrary constant and the coefficients B_m can be found from the corresponding system of algebraic equations.

$$20. \quad y(x) + A \int_0^\infty y(t)y(x + \lambda t) dt = 0, \quad \lambda > 0, \quad 0 \leq x < \infty.$$

This is a special case of equation 8.2.21 with $f(t) \equiv A$, $a = 0$, and $b = \infty$.

A solution:

$$y(x) = -\frac{C(\lambda + 1)}{A} e^{-Cx},$$

where C is an arbitrary positive constant.

$$21. \quad y(x) + A \int_0^\infty e^{-\lambda t} y\left(\frac{x}{t}\right) y(t) dt = 0, \quad \lambda > 0.$$

A solution: $y(x) = -\frac{\lambda}{A} x^C$, where C is an arbitrary constant.

$$22. \quad y(x) + A \int_0^\infty e^{-\lambda t} y\left(\frac{x}{t}\right) y(t) dt = Bx^b, \quad \lambda > 0.$$

Solutions:

$$y_1(x) = \beta_1 x^b, \quad y_2(x) = \beta_2 x^b,$$

where β_1 and β_2 are the roots of the quadratic equation $A\beta^2 + \lambda\beta - B\lambda = 0$.

8.2. Equations with Quadratic Nonlinearity That Contain Arbitrary Functions

8.2-1. Equations of the Form $y(x) + \int_a^b K(x, t)y^2(t) dt = F(x)$.

$$1. \quad y(x) + \int_a^b f(x)y^2(t) dt = 0.$$

Solutions: $y_1(x) = 0$ and $y_2(x) = \lambda f(x)$, where $\lambda = -\left[\int_a^b f^2(t) dt\right]^{-1}$.

$$2. \quad y(x) + \int_a^b f(x)g(t)y^2(t) dt = 0.$$

This is a special case of equation 8.8.9.

Solutions: $y_1(x) = 0$ and $y_2(x) = \lambda f(x)$, where $\lambda = -\left[\int_a^b f^2(t)g(t) dt\right]^{-1}$.

$$3. \quad y(x) + A \int_a^b y^2(t) dt = f(x).$$

This is a special case of equation 8.8.7.

A solution: $y(x) = f(x) + \lambda$, where λ is determined by the quadratic equation

$$A(b-a)\lambda^2 + (1 + 2AI_1)\lambda + AI_2 = 0, \quad \text{where } I_1 = \int_a^b f(t) dt, \quad I_2 = \int_a^b f^2(t) dt.$$

$$4. \quad y(x) + \int_a^b g(t)y^2(t) dt = f(x).$$

This is a special case of equation 8.8.9.

A solution: $y(x) = f(x) + \lambda$, where λ is determined by the quadratic equation

$$I_0\lambda^2 + (1 + 2I_1)\lambda + I_2 = 0, \quad \text{where } I_m = \int_a^b f^m(t)g(t) dt, \quad m = 0, 1, 2.$$

$$5. \quad y(x) + \int_a^b g(x)y^2(t) dt = f(x).$$

Solution: $y(x) = \lambda g(x) + f(x)$, where λ is determined by the quadratic equation

$$I_{gg}\lambda^2 + (1 + 2I_{fg})\lambda + I_{ff} = 0,$$

$$I_{gg} = \int_a^b g^2(t) dt, \quad I_{fg} = \int_a^b f(t)g(t) dt, \quad I_{ff} = \int_a^b f^2(t) dt.$$

$$6. \quad y(x) + \int_a^b [g_1(x)h_1(t) + g_2(x)h_2(t)] y^2(t) dt = f(x).$$

A solution: $y(x) = \lambda_1 g_1(x) + \lambda_2 g_2(x) + f(x)$, where the constants λ_1 and λ_2 can be found from a system of two second-order algebraic equations (this system can be obtained from the more general system presented in 8.8.19).

8.2-2. Equations of the Form $y(x) + \int_a^b \sum K_{nm}(x, t)y^n(x)y^m(t) dt = F(x)$, $n+m \leq 2$.

$$7. \quad y(x) + \int_a^b g(t)y(x)y(t) dt = f(x).$$

Solutions:

$$y_1(x) = \lambda_1 f(x), \quad y_2(x) = \lambda_2 f(x),$$

where λ_1 and λ_2 are the roots of the quadratic equation

$$I\lambda^2 + \lambda - 1 = 0, \quad I = \int_a^b f(t)g(t) dt.$$

$$8. \quad y(x) + \int_a^b g(x)y(x)y(t) dt = f(x).$$

A solution:

$$y(x) = \frac{f(x)}{1 + \lambda g(x)},$$

where λ is a root of the algebraic (or transcendental) equation

$$\lambda - \int_a^b \frac{f(t) dt}{1 + \lambda g(t)} = 0.$$

Different roots generate different solutions of the integral equation.

$$9. \quad y(x) + \int_a^b [g_1(t)y^2(x) + g_2(x)y(t)] dt = f(x).$$

Solution in an implicit form:

$$y(x) + Iy^2(x) + \lambda g_2(x) - f(x) = 0, \quad I = \int_a^b g_1(t) dt, \quad (1)$$

where λ is determined by the algebraic equation

$$\lambda = \int_a^b y(t) dt. \quad (2)$$

Here the function $y(x) = y(x, \lambda)$ obtained by solving the quadratic equation (1) must be substituted in the integrand of (2).

$$10. \quad y(x) + \int_a^b [g_1(t)y^2(x) + g_2(x)y^2(t)] dt = f(x).$$

Solution in an implicit form:

$$y(x) + Iy^2(x) + \lambda g_2(x) - f(x) = 0, \quad I = \int_a^b g_1(t) dt, \quad (1)$$

where λ is determined by the algebraic equation

$$\lambda = \int_a^b y^2(t) dt. \quad (2)$$

Here the function $y(x) = y(x, \lambda)$ obtained by solving the quadratic equation (1) must be substituted into the integrand of (2).

$$11. \quad y(x) + \int_a^b [g_{11}(x)h_{11}(t)y^2(x) + g_{12}(x)h_{12}(t)y(x)y(t) + g_{22}(x)h_{22}(t)y^2(t) \\ + g_1(x)h_1(t)y(x) + g_2(x)h_2(t)y(t)] dt = f(x).$$

This is a special case of equation 8.8.49.

$$12. \quad y(x) + \int_{-\infty}^{\infty} [\lambda e^{-|x-t|}y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.2.14.

Solutions for $\lambda > -\frac{1}{2}$:

$$y_{1,2}(x) = Y_f(x) + A_{1,2}Y_\varphi(x),$$

where

$$Y_f(x) = f(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) f(t) dt, \\ Y_\varphi(x) = \varphi(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) \varphi(t) dt,$$

and $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0, \\ p = \int_0^{\infty} \psi(t)Y_\varphi^2(t) dt, \quad q = 1 + 2 \int_0^{\infty} \psi(t)Y_f(t)Y_\varphi(t) dt, \quad r = \int_0^{\infty} \psi(t)Y_f^2(t) dt.$$

$$13. \quad y(x) - \int_0^{\infty} [\lambda \sin(xt)y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.5.20.

Solutions for $\lambda \neq \pm\sqrt{\frac{2}{\pi}}$:

$$y_{1,2}(x) = Y_f(x) + A_{1,2}Y_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt)f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt)\varphi(t) dt,$$

and $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0,$$

$$p = \int_0^{\infty} \psi(t)Y_\varphi^2(t) dt, \quad q = 2 \int_0^{\infty} \psi(t)Y_f(t)Y_\varphi(t) dt - 1, \quad r = \int_0^{\infty} \psi(t)Y_f^2(t) dt.$$

$$14. \quad y(x) - \int_0^{\infty} [\lambda \cos(xt)y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.5.6.

Solutions for $\lambda \neq \pm\sqrt{\frac{2}{\pi}}$:

$$y_{1,2}(x) = Y_f(x) + A_{1,2}Y_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \cos(xt)f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \cos(xt)\varphi(t) dt,$$

and $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0,$$

$$p = \int_0^{\infty} \psi(t)Y_\varphi^2(t) dt, \quad q = 2 \int_0^{\infty} \psi(t)Y_f(t)Y_\varphi(t) dt - 1, \quad r = \int_0^{\infty} \psi(t)Y_f^2(t) dt.$$

$$15. \quad y(x) + \int_0^{\infty} [\lambda t J_\nu(xt)y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x), \quad \nu > -1.$$

Here $J_\nu(z)$ is the Bessel function of the first kind. The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.8.4.

Solutions for $\lambda \neq \pm 1$:

$$y_{1,2}(x) = Y_f(x) + A_{1,2}Y_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1 - \lambda^2} - \frac{\lambda}{1 - \lambda^2} \int_0^{\infty} t J_\nu(xt)f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1 - \lambda^2} - \frac{\lambda}{1 - \lambda^2} \int_0^{\infty} t J_\nu(xt)\varphi(t) dt,$$

and $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0,$$

$$p = \int_0^\infty \psi(t)Y_\varphi^2(t) dt, \quad q = 1 + 2 \int_0^\infty \psi(t)Y_f(t)Y_\varphi(t) dt, \quad r = \int_0^\infty \psi(t)Y_f^2(t) dt.$$

8.2-3. Equations of the Form $y(x) + \int_a^b K(t)y(t)y(\cdot \cdot) dt = F(x)$.

16. $y(x) + \int_a^b f(t)y(t)y(xt) dt = 0.$

1°. Solutions:

$$y_1(x) = -\frac{1}{I_0}x^C, \quad y_2(x) = \frac{(I_1 - I_0)x + I_1 - I_2}{I_0I_2 - I_1^2}x^C,$$

$$I_m = \int_a^b f(t)t^{2C+m} dt, \quad m = 0, 1, 2,$$

where C is an arbitrary constant.

There are more complicated solutions of the form $y(x) = x^C \sum_{k=0}^n B_k x^k$, where C is an arbitrary constant and the coefficients B_k can be found from the corresponding system of algebraic equations.

2°. A solution:

$$y_3(x) = \frac{(I_1 - I_0)x^\beta + I_1 - I_2}{I_0I_2 - I_1^2}x^C,$$

$$I_m = \int_a^b f(t)t^{2C+m\beta} dt, \quad m = 0, 1, 2,$$

where C and β are arbitrary constants.

There are more complicated solutions of the form $y(x) = x^C \sum_{k=0}^n D_k x^{k\beta}$, where C and β are arbitrary constants and the coefficients D_k can be found from the corresponding system of algebraic equations.

3°. A solution:

$$y_4(x) = \frac{x^C(J_1 \ln x - J_2)}{J_0J_2 - J_1^2},$$

$$J_m = \int_a^b f(t)t^{2C}(\ln t)^m dt, \quad m = 0, 1, 2,$$

where C is an arbitrary constant.

There are more complicated solutions of the form $y(x) = x^C \sum_{k=0}^n E_k (\ln x)^k$, where C is an arbitrary constant and the coefficients E_k can be found from the corresponding system of algebraic equations.

4°. The equation also has the trivial solution $y(x) \equiv 0$.

5°. The substitution $y(x) = x^\beta w(x)$ leads to an equation of the same form,

$$w(x) + \int_a^b g(t)w(t)w(xt) dt = 0, \quad g(x) = f(x)x^{2\beta}.$$

$$17. \quad y(x) + \int_a^b f(t)y(t)y(xt) dt = Ax^\beta.$$

1°. Solutions:

$$y_1(x) = k_1x^\beta, \quad y_2(x) = k_2x^\beta,$$

where k_1 and k_2 are the roots of the quadratic equation

$$Ik^2 + k - A = 0, \quad I = \int_a^b f(t)t^{2\beta} dt.$$

2°. Solutions:

$$y(x) = x^\beta(\lambda x + \mu),$$

where λ and μ are determined from the following system of two algebraic equations (this system can be reduced to a quadratic equation):

$$I_2\lambda + I_1\mu + 1 = 0, \quad I_1\lambda\mu + I_0\mu^2 + \mu - A = 0$$

where $I_m = \int_a^b f(t)t^{2\beta+m} dt$, $m = 0, 1, 2$.

3°. There are more complicated solutions of the form $y(x) = x^\beta \sum_{m=0}^n B_m x^m$, where the B_m can be found from the corresponding system of algebraic equations.

$$18. \quad y(x) + \int_a^b f(t)y(t)y(xt) dt = A \ln x + B.$$

This equation has solutions of the form $y(x) = p \ln x + q$, where the constants p and q can be found from a system of two second-order algebraic equations.

$$19. \quad y(x) + \int_0^\infty f(t)y(t)y\left(\frac{x}{t}\right) dt = 0.$$

1°. A solution:

$$y(x) = -kx^C, \quad k = \left[\int_0^\infty f(t) dt \right]^{-1},$$

where C is an arbitrary constant.

2°. The equation has the trivial solution $y(x) \equiv 0$.

3°. The substitution $y(x) = x^\beta w(x)$ leads to an equation of the same form,

$$w(x) + \int_0^\infty f(t)w(t)w\left(\frac{x}{t}\right) dt = 0.$$

$$20. \quad y(x) + \int_0^\infty f(t)y(t)y\left(\frac{x}{t}\right) dt = Ax^b.$$

Solutions:

$$y_1(x) = \lambda_1 x^b, \quad y_2(x) = \lambda_2 x^b,$$

where λ_1 and λ_2 are the roots of the quadratic equation

$$I\lambda^2 + \lambda - A = 0, \quad I = \int_0^\infty f(t) dt.$$

$$21. \quad y(x) + \int_a^b f(t)y(t)y(x + \lambda t) dt = 0, \quad \lambda > 0.$$

1°. Solutions:

$$y_1(x) = -\frac{1}{I_0} \exp(-Cx), \quad y_2(x) = \frac{I_2 - I_1 x}{I_1^2 - I_0 I_2} \exp(-Cx),$$

$$I_m = \int_a^b t^m \exp[-C(\lambda + 1)t] f(t) dt, \quad m = 0, 1, 2,$$

where C is an arbitrary constant.

2°. There are more complicated solutions of the form $y(x) = \exp(-Cx) \sum_{k=0}^n A_k x^k$, where C is an arbitrary constant and the coefficients A_k can be found from the corresponding system of algebraic equations.

3°. The equation also has the trivial solution $y(x) \equiv 0$.

4°. The substitution $y(x) = e^{\beta x} w(x)$ leads to a similar equation:

$$w(x) + \int_a^b g(t)w(t)w(x + \lambda t) dt = 0, \quad g(t) = e^{\beta(\lambda+1)t} f(t).$$

$$22. \quad y(x) + \int_a^b f(t)y(x + \lambda t)y(t) dt = A e^{-\mu x}, \quad \lambda > 0.$$

1°. Solutions:

$$y_1(x) = k_1 e^{-\mu x}, \quad y_2(x) = k_2 e^{-\mu x},$$

where k_1 and k_2 are the roots of the quadratic equation

$$Ik^2 + k - A = 0, \quad I = \int_a^b e^{-\mu(\lambda+1)t} f(t) dt.$$

2°. There are more complicated solutions of the form $y(x) = e^{-\mu x} \sum_{m=0}^n B_m x^m$, where the B_m can be found from the corresponding system of algebraic equations.

3°. The substitution $y(x) = e^{\beta x} w(x)$ leads to an equation of the same form,

$$w(x) + \int_a^b g(t)w(t)w(x - t) dt = A e^{(\lambda-\beta)x}, \quad g(t) = f(t) e^{\beta(\lambda+1)t}.$$

$$23. \quad y(x) + \int_a^b f(t)y(t)y(x - t) dt = 0.$$

1°. Solutions:

$$y_1(x) = -\frac{1}{I_0} \exp(Cx), \quad y_2(x) = \frac{I_2 - I_1 x}{I_1^2 - I_0 I_2} \exp(Cx), \quad I_m = \int_a^b t^m f(t) dt,$$

where C is an arbitrary constant and $m = 0, 1, 2$.

2°. There are more complicated solutions of the form $y(x) = \exp(Cx) \sum_{k=0}^n A_k x^k$, where C is an arbitrary constant and the coefficients A_k can be found from the corresponding system of algebraic equations.

3°. The equation also has the trivial solution $y(x) \equiv 0$.

4°. The substitution $y(x) = \exp(Cx)w(x)$ leads to an equation of the same form:

$$w(x) + \int_a^b f(t)w(t)w(x - t) dt = 0.$$

$$24. \quad y(x) + \int_a^b f(t)y(x-t)y(t) dt = Ae^{\lambda x}.$$

1°. Solutions:

$$y_1(x) = k_1 e^{\lambda x}, \quad y_2(x) = k_2 e^{\lambda x},$$

where k_1 and k_2 are the roots of the quadratic equation

$$Ik^2 + k - A = 0, \quad I = \int_a^b f(t) dt.$$

2°. The substitution $y(x) = e^{\beta x} w(x)$ leads to an equation of the same form,

$$w(x) + \int_a^b f(t)w(t)w(x-t) dt = Ae^{(\lambda-\beta)x}.$$

$$25. \quad y(x) + \int_a^b f(t)y(t)y(x-t) dt = A \sinh \lambda x.$$

A solution:

$$y(x) = p \sinh \lambda x + q \cosh \lambda x. \quad (1)$$

Here p and q are roots of the algebraic system

$$p + I_0 p q + I_{cs}(p^2 - q^2) = A, \quad q + I_{cc} q^2 - I_{ss} p^2 = 0, \quad (2)$$

where

$$I_0 = \int_a^b f(t) dt, \quad I_{cs} = \int_a^b f(t) \cosh(\lambda t) \sinh(\lambda t) dt,$$

$$I_{cc} = \int_a^b f(t) \cosh^2(\lambda t) dt, \quad I_{ss} = \int_a^b f(t) \sinh^2(\lambda t) dt.$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$26. \quad y(x) + \int_a^b f(t)y(t)y(x-t) dt = A \cosh \lambda x.$$

A solution:

$$y(x) = p \sinh \lambda x + q \cosh \lambda x. \quad (1)$$

Here p and q are roots of the algebraic system

$$p + I_0 p q + I_{cs}(p^2 - q^2) = 0, \quad q + I_{cc} q^2 - I_{ss} p^2 = A, \quad (2)$$

where we use the notation introduced in 8.2.25. Different solutions of system (2) generate different solutions (1) of the integral equation.

$$27. \quad y(x) + \int_a^b f(t)y(t)y(x-t) dt = A \sin \lambda x.$$

A solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x. \quad (1)$$

Here p and q are roots of the algebraic system

$$p + I_0 p q + I_{cs}(p^2 + q^2) = A, \quad q + I_{cc} q^2 - I_{ss} p^2 = 0, \quad (2)$$

where

$$I_0 = \int_a^b f(t) dt, \quad I_{cs} = \int_a^b f(t) \cos(\lambda t) \sin(\lambda t) dt,$$

$$I_{cc} = \int_a^b f(t) \cos^2(\lambda t) dt, \quad I_{ss} = \int_a^b f(t) \sin^2(\lambda t) dt.$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$28. \quad y(x) + \int_a^b f(t)y(t)y(x-t) dt = A \cos \lambda x.$$

A solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x. \quad (1)$$

Here p and q are roots of the algebraic system

$$p + I_0 p q + I_{cs}(p^2 + q^2) = 0, \quad q + I_{cc} q^2 - I_{ss} p^2 = A, \quad (2)$$

where we use the notation introduced in 8.2.27. Different solutions of system (2) generate different solutions (1) of the integral equation.

8.3. Equations with Power-Law Nonlinearity

8.3-1. Equations of the Form $y(x) + \int_a^b K(x,t)y^\beta(t) dt = F(x)$.

$$1. \quad y(x) + A \int_a^b t^\lambda y^\beta(t) dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\lambda y^\beta$.

$$2. \quad y(x) + A \int_a^b e^{\mu t} y^\beta(t) dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = Ae^{\mu t} y^\beta$.

$$3. \quad y(x) + A \int_a^b e^{\lambda(x-t)} y^\beta(t) dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = Ay^\beta$.

$$4. \quad y(x) - \int_a^b g(x)y^\beta(t) dt = 0.$$

A solution:

$$y(x) = \lambda g(x), \quad \lambda = \left[\int_a^b g^\beta(t) dt \right]^{\frac{1}{1-\beta}}.$$

For $\beta > 0$, the equation also has the trivial solution $y(x) \equiv 0$.

$$5. \quad y(x) - \int_a^b g(x)y^\beta(t) dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = -y^\beta$.

$$6. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) y^\beta(t) dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = Ay^\beta$.

$$7. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) y^\beta(t) dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = Ay^\beta$.

$$8. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) y^\beta(t) dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = Ay^\beta$.

$$9. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) y^\beta(t) dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = Ay^\beta$.

$$10. \quad y(x) + \int_0^\infty f\left(\frac{t}{x}\right) \sqrt{y(t)} dt = Ax^2.$$

Solutions: $y_k(x) = \beta_k^2 x^2$, where β_k ($k = 1, 2$) are the roots of the quadratic equations

$$\beta^2 \pm I\beta - A = 0, \quad I = \int_0^\infty z f(z) dz.$$

$$11. \quad y(x) - \int_0^\infty t^\lambda f\left(\frac{t}{x}\right) y^\beta(t) dt = 0, \quad \beta \neq 1.$$

A solution:

$$y(x) = Ax^{\frac{1+\lambda}{1-\beta}}, \quad A^{1-\beta} = \int_0^\infty z^{\frac{\lambda+\beta}{1-\beta}} f(z) dz.$$

$$12. \quad y(x) - \int_{-\infty}^\infty e^{\lambda t} f(ax + bt) y^\beta(t) dt = 0, \quad b \neq 0, \quad a\beta \neq -b.$$

A solution:

$$y(x) = A \exp\left(-\frac{a\lambda}{a\beta + b} x\right), \quad A^{1-\beta} = \int_{-\infty}^\infty \exp\left(\frac{\lambda b}{a\beta + b} z\right) f(bz) dz.$$

8.3-2. Other Equations.

$$13. \quad y(x) + A \int_a^b y^\beta(x) y^\mu(t) dt = f(x).$$

Solution in an implicit form:

$$y(x) + A\lambda y^\beta(x) - f(x) = 0, \tag{1}$$

where λ is determined by the algebraic (or transcendental) equation

$$\lambda = \int_a^b y^\mu(t) dt. \tag{2}$$

Here the function $y(x) = y(x, \lambda)$ obtained by solving the quadratic equation (1) must be substituted in the integrand of (2).

$$14. \quad y(x) + \int_a^b g(t) y(x) y^\mu(t) dt = f(x).$$

A solution: $y(x) = \lambda f(x)$, where λ is determined from the algebraic (or transcendental) equation

$$I\lambda^{\mu+1} + \lambda - 1 = 0, \quad I = \int_a^b g(t) f^\mu(t) dt.$$

$$15. \quad y(x) + \int_a^b g(x)y(x)y^\mu(t) dt = f(x).$$

A solution:

$$y(x) = \frac{f(x)}{1 + \lambda g(x)},$$

where λ is a root of the algebraic (or transcendental) equation

$$\lambda - \int_a^b \frac{f^\mu(t) dt}{[1 + \lambda g(t)]^\mu} = 0.$$

Different roots generate different solutions of the integral equation.

$$16. \quad y(x) + \int_a^b [g_1(t)y^2(x) + g_2(x)y^\mu(t)] dt = f(x).$$

Solution in an implicit form:

$$y(x) + Iy^2(x) + \lambda g_2(x) - f(x) = 0, \quad I = \int_a^b g_1(t) dt, \quad (1)$$

where λ is determined by the algebraic (or transcendental) equation

$$\lambda = \int_a^b y^\mu(t) dt. \quad (2)$$

Here the function $y(x) = y(x, \lambda)$ obtained by solving the quadratic equation (1) must be substituted in the integrand of (2).

$$17. \quad y(x) + \int_a^b [g_1(x)h_1(t)y^k(x)y^s(t) + g_2(x)h_2(t)y^p(x)y^q(t)] dt = f(x).$$

This is a special case of equation 8.8.49.

$$18. \quad y(x) + A \int_a^b f(t)y(xt)y^\beta(t) dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = Af(t)y^\beta$.

$$19. \quad y(x) + \int_{-\infty}^{\infty} [\lambda e^{-|x-t|}y(t) + \varphi(x)\psi(t)y^\beta(t)] dt = f(x).$$

This is a special case of equation 8.8.21. The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.2.14.

$$20. \quad y(x) - \int_0^{\infty} [\lambda \sin(xt)y(t) + \varphi(x)\psi(t)y^\beta(t)] dt = f(x).$$

This is a special case of equation 8.8.22. The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.5.20.

$$21. \quad y(x) - \int_0^{\infty} [\lambda \cos(xt)y(t) + \varphi(x)\psi(t)y^\beta(t)] dt = f(x).$$

This is a special case of equation 8.8.23. The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.5.6.

$$22. \quad y(x) + \int_0^{\infty} [\lambda t J_\nu(xt)y(t) + \varphi(x)\psi(t)y^\beta(t)] dt = f(x).$$

Here $J_\nu(z)$ is the Bessel function of the first kind. This is a special case of equation 8.8.24.

8.4. Equations with Exponential Nonlinearity

8.4-1. Integrands with Nonlinearity of the Form $\exp[\beta y(t)]$.

$$1. \quad y(x) + A \int_a^b \exp[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \exp(\beta y)$.

$$2. \quad y(x) + A \int_a^b t^\mu \exp[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \exp(\beta y)$.

$$3. \quad y(x) + A \int_a^b \exp[\mu t + \beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \exp(\mu t) \exp(\beta y)$.

$$4. \quad y(x) + A \int_a^b \exp[\lambda(x-t) + \beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \exp(\beta y)$.

$$5. \quad y(x) + \int_a^b g(x) \exp[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \exp(\beta y)$.

$$6. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \exp[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \exp(\beta y)$.

$$7. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \exp[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \exp(\beta y)$.

$$8. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \exp[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \exp(\beta y)$.

$$9. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \exp[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \exp(\beta y)$.

8.4-2. Other Integrands.

$$10. \quad y(x) + A \int_a^b \exp[\beta y(x) + \gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \exp(\beta y)$ and $f(t, y) = \exp(\gamma y)$.

$$11. \quad y(x) + A \int_a^b y(xt) \exp[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \exp(\beta y)$.

8.5. Equations with Hyperbolic Nonlinearity

8.5-1. Integrands with Nonlinearity of the Form $\cosh[\beta y(t)]$.

$$1. \quad y(x) + A \int_a^b \cosh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \cosh(\beta y)$.

$$2. \quad y(x) + A \int_a^b t^\mu \cosh^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A t^\mu \cosh^k(\beta y)$.

$$3. \quad y(x) + A \int_a^b \cosh(\mu t) \cosh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \cosh(\mu t) \cosh(\beta y)$.

$$4. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \cosh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \cosh(\beta y)$.

$$5. \quad y(x) + \int_a^b g(x) \cosh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \cosh(\beta y)$.

$$6. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \cosh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \cosh(\beta y)$.

$$7. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \cosh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \cosh(\beta y)$.

$$8. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \cosh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \cosh(\beta y)$.

$$9. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \cosh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \cosh(\beta y)$.

8.5-2. Integrands with Nonlinearity of the Form $\sinh[\beta y(t)]$.

$$10. \quad y(x) + A \int_a^b \sinh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \sinh(\beta y)$.

$$11. \quad y(x) + A \int_a^b t^\mu \sinh^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \sinh^k(\beta y)$.

$$12. \quad y(x) + A \int_a^b \sinh(\mu t) \sinh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \sinh(\mu t) \sinh(\beta y)$.

$$13. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \sinh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \sinh(\beta y)$.

$$14. \quad y(x) + \int_a^b g(x) \sinh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \sinh(\beta y)$.

$$15. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \sinh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \sinh(\beta y)$.

$$16. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \sinh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \sinh(\beta y)$.

$$17. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \sinh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \sinh(\beta y)$.

$$18. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \sinh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \sinh(\beta y)$.

8.5-3. Integrands with Nonlinearity of the Form $\tanh[\beta y(t)]$.

$$19. \quad y(x) + A \int_a^b \tanh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \tanh(\beta y)$.

$$20. \quad y(x) + A \int_a^b t^\mu \tanh^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \tanh^k(\beta y)$.

$$21. \quad y(x) + A \int_a^b \tanh(\mu t) \tanh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \tanh(\mu t) \tanh(\beta y)$.

$$22. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \tanh[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \tanh(\beta y)$.

$$23. \quad y(x) + \int_a^b g(x) \tanh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \tanh(\beta y)$.

$$24. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \tanh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \tanh(\beta y)$.

$$25. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \tanh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \tanh(\beta y)$.

$$26. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \tanh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \tanh(\beta y)$.

$$27. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \tanh[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \tanh(\beta y)$.

8.5-4. Integrands with Nonlinearity of the Form $\coth[\beta y(t)]$.

$$28. \quad y(x) + A \int_a^b \coth[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \coth(\beta y)$.

$$29. \quad y(x) + A \int_a^b t^\mu \coth^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \coth^k(\beta y)$.

$$30. \quad y(x) + A \int_a^b \coth(\mu t) \coth[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \coth(\mu t) \coth(\beta y)$.

$$31. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \coth[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \coth(\beta y)$.

$$32. \quad y(x) + \int_a^b g(x) \coth[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \coth(\beta y)$.

$$33. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \coth[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \coth(\beta y)$.

$$34. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \coth[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \coth(\beta y)$.

$$35. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \coth[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \coth(\beta y)$.

$$36. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \coth[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \coth(\beta y)$.

8.5-5. Other Integrands.

$$37. \quad y(x) + A \int_a^b \cosh[\beta y(x)] \cosh[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \cosh(\beta y)$ and $f(t, y) = \cosh(\gamma y)$.

$$38. \quad y(x) + A \int_a^b y(xt) \cosh[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \cosh(\beta y)$.

$$39. \quad y(x) + A \int_a^b \sinh[\beta y(x)] \sinh[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \sinh(\beta y)$ and $f(t, y) = \sinh(\gamma y)$.

$$40. \quad y(x) + A \int_a^b y(xt) \sinh[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \sinh(\beta y)$.

$$41. \quad y(x) + A \int_a^b \tanh[\beta y(x)] \tanh[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \tanh(\beta y)$ and $f(t, y) = \tanh(\gamma y)$.

$$42. \quad y(x) + A \int_a^b y(xt) \tanh[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \tanh(\beta y)$.

$$43. \quad y(x) + A \int_a^b \coth[\beta y(x)] \coth[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \coth(\beta y)$ and $f(t, y) = \coth(\gamma y)$.

$$44. \quad y(x) + A \int_a^b y(xt) \coth[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \coth(\beta y)$.

8.6. Equations with Logarithmic Nonlinearity

8.6-1. Integrands with Nonlinearity of the Form $\ln[\beta y(t)]$.

$$1. \quad y(x) + A \int_a^b \ln[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \ln(\beta y)$.

$$2. \quad y(x) + A \int_a^b t^\mu \ln^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \ln^k(\beta y)$.

$$3. \quad y(x) + A \int_a^b \ln(\mu t) \ln[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \ln(\mu t) \ln(\beta y)$.

$$4. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \ln[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \ln(\beta y)$.

$$5. \quad y(x) + \int_a^b g(x) \ln[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \ln(\beta y)$.

$$6. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \ln[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \ln(\beta y)$.

$$7. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \ln[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \ln(\beta y)$.

$$8. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \ln[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \ln(\beta y)$.

$$9. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \ln[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \ln(\beta y)$.

8.6-2. Other Integrands.

$$10. \quad y(x) + A \int_a^b \ln[\beta y(x)] \ln[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \ln(\beta y)$ and $f(t, y) = \ln(\gamma y)$.

$$11. \quad y(x) + A \int_a^b y(xt) \ln[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \ln(\beta y)$.

8.7. Equations with Trigonometric Nonlinearity

8.7-1. Integrands with Nonlinearity of the Form $\cos[\beta y(t)]$.

$$1. \quad y(x) + A \int_a^b \cos[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \cos(\beta y)$.

$$2. \quad y(x) + A \int_a^b t^\mu \cos^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \cos^k(\beta y)$.

$$3. \quad y(x) + A \int_a^b \cos(\mu t) \cos[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \cos(\mu t) \cos(\beta y)$.

$$4. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \cos[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \cos(\beta y)$.

$$5. \quad y(x) + \int_a^b g(x) \cos[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \cos(\beta y)$.

$$6. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \cos[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \cos(\beta y)$.

$$7. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \cos[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \cos(\beta y)$.

$$8. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \cos[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \cos(\beta y)$.

$$9. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \cos[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \cos(\beta y)$.

8.7-2. Integrands with Nonlinearity of the Form $\sin[\beta y(t)]$.

$$10. \quad y(x) + A \int_a^b \sin[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \sin(\beta y)$.

$$11. \quad y(x) + A \int_a^b t^\mu \sin^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \sin^k(\beta y)$.

$$12. \quad y(x) + A \int_a^b \sin(\mu t) \sin[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \sin(\mu t) \sin(\beta y)$.

$$13. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \sin[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \sin(\beta y)$.

$$14. \quad y(x) + \int_a^b g(x) \sin[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \sin(\beta y)$.

$$15. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \sin[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \sin(\beta y)$.

$$16. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \sin[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \sin(\beta y)$.

$$17. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \sin[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \sin(\beta y)$.

$$18. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \sin[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \sin(\beta y)$.

8.7-3. Integrands with Nonlinearity of the Form $\tan[\beta y(t)]$.

$$19. \quad y(x) + A \int_a^b \tan[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \tan(\beta y)$.

$$20. \quad y(x) + A \int_a^b t^\mu \tan^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \tan^k(\beta y)$.

$$21. \quad y(x) + A \int_a^b \tan(\mu t) \tan[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \tan(\mu t) \tan(\beta y)$.

$$22. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \tan[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \tan(\beta y)$.

$$23. \quad y(x) + \int_a^b g(x) \tan[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \tan(\beta y)$.

$$24. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \tan[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \tan(\beta y)$.

$$25. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \tan[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \tan(\beta y)$.

$$26. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \tan[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \tan(\beta y)$.

$$27. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \tan[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \tan(\beta y)$.

8.7-4. Integrands with Nonlinearity of the Form $\cot[\beta y(t)]$.

$$28. \quad y(x) + A \int_a^b \cot[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \cot(\beta y)$.

$$29. \quad y(x) + A \int_a^b t^\mu \cot^k[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = At^\mu \cot^k(\beta y)$.

$$30. \quad y(x) + A \int_a^b \cot(\mu t) \cot[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.7 with $f(t, y) = A \cot(\mu t) \cot(\beta y)$.

$$31. \quad y(x) + A \int_a^b e^{\lambda(x-t)} \cot[\beta y(t)] dt = g(x).$$

This is a special case of equation 8.8.8 with $f(t, y) = A \cot(\beta y)$.

$$32. \quad y(x) + \int_a^b g(x) \cot[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.9 with $f(t, y) = \cot(\beta y)$.

$$33. \quad y(x) + A \int_a^b \cosh(\lambda x + \mu t) \cot[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.11 with $f(t, y) = A \cot(\beta y)$.

$$34. \quad y(x) + A \int_a^b \sinh(\lambda x + \mu t) \cot[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.12 with $f(t, y) = A \cot(\beta y)$.

$$35. \quad y(x) + A \int_a^b \cos(\lambda x + \mu t) \cot[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.13 with $f(t, y) = A \cot(\beta y)$.

$$36. \quad y(x) + A \int_a^b \sin(\lambda x + \mu t) \cot[\beta y(t)] dt = h(x).$$

This is a special case of equation 8.8.14 with $f(t, y) = A \cot(\beta y)$.

8.7-5. Other Integrands.

$$37. \quad y(x) + A \int_a^b \cos[\beta y(x)] \cos[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \cos(\beta y)$ and $f(t, y) = \cos(\gamma y)$.

$$38. \quad y(x) + A \int_a^b y(xt) \cos[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \cos(\beta y)$.

$$39. \quad y(x) + A \int_a^b \sin[\beta y(x)] \sin[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \sin(\beta y)$ and $f(t, y) = \sin(\gamma y)$.

$$40. \quad y(x) + A \int_a^b y(xt) \sin[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \sin(\beta y)$.

$$41. \quad y(x) = \lambda \int_0^1 f(x)g(t) \sin\left(\frac{y(t)}{f(t)}\right) y(t) dt.$$

Solutions are sought in the form $y(x) = Af(x)$, where the constant A is determined from the transcendental equation (the trivial solution corresponding to $A = 0$ is not taken into account):

$$1 = \lambda \sigma \sin A, \quad \sigma = \int_0^1 f(t)g(t) dt.$$

For $|\lambda| < 1/|\sigma|$, the integral equation has no real solutions (the case $\sigma = 0$ is included).

For any λ satisfying the inequality $|\lambda| > 1/|\sigma|$, the integral equation has infinitely many real solutions.

$$42. \quad y(x) + A \int_a^b \tan[\beta y(x)] \tan[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \tan(\beta y)$ and $f(t, y) = \tan(\gamma y)$.

$$43. \quad y(x) + A \int_a^b y(xt) \tan[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \tan(\beta y)$.

$$44. \quad y(x) + A \int_a^b \cot[\beta y(x)] \cot[\gamma y(t)] dt = h(x).$$

This is a special case of equation 8.8.48 with $g(x, y) = A \cot(\beta y)$ and $f(t, y) = \cot(\gamma y)$.

$$45. \quad y(x) + A \int_a^b y(xt) \cot[\beta y(t)] dt = 0.$$

This is a special case of equation 8.8.25 with $f(t, y) = A \cot(\beta y)$.

8.8. Equations with Nonlinearity of General Form

8.8-1. Equations of the Form $y(x) + \int_a^b K(|x-t|)G(y(t)) dt = F(x)$.

$$1. \quad y(x) + \int_a^b |x-t|f(y(t)) dt = Ax^2 + Bx + C.$$

This is a special case of equation 8.8.15 with $f(t, y) = f(y)$ and $g(x) = Ax^2 + Bx + C$.

The function $y = y(x)$ obeys the second-order autonomous differential equation

$$y''_{xx} + 2f(y) = 2A,$$

whose solution can be represented in an implicit form:

$$\int_{y_a}^y \frac{du}{\sqrt{w_a^2 + 4A(u - y_a) - 4F(u, y_a)}} = \pm(x - a), \quad F(u, v) = \int_v^u f(t) dt, \quad (1)$$

where $y_a = y(a)$ and $w_a = y'_x(a)$ are constants of integration. These constants, as well as the unknowns $y_b = y(b)$ and $w_b = y'_x(b)$, are determined by the algebraic (or transcendental) system

$$\begin{aligned} y_a + y_b - (a - b)w_a &= (b^2 + 2ab - a^2)A + 2bB + 2C, \\ w_a + w_b &= 2(a + b)A + 2B, \\ w_b^2 &= w_a^2 + 4A(y_b - y_a) - 4F(y_b, y_a), \\ \int_{y_a}^{y_b} \frac{du}{\sqrt{w_a^2 + 4A(u - y_a) - 4F(u, y_a)}} &= \pm(b - a). \end{aligned} \quad (2)$$

Here the first equation is obtained from the second condition of (5) in 8.8.15, the second equation is obtained from condition (6) in 8.8.15, and the third and fourth equations are consequences of (1).

Each solution of system (2) generates a solution of the integral equation.

$$2. \quad y(x) + \int_a^b e^{\lambda|x-t|} f(y(t)) dt = A + Be^{\lambda x} + Ce^{-\lambda x}.$$

This is a special case of equation 8.8.16 with $f(t, y) = f(y)$ and $g(x) = A + Be^{\lambda x} + Ce^{-\lambda x}$.

The function $y = y(x)$ satisfies the second-order autonomous differential equation

$$y''_{xx} + 2\lambda f(y) - \lambda^2 y = -\lambda^2 A, \quad (1)$$

whose solution can be written in an implicit form:

$$\int_{y_a}^y \frac{du}{\sqrt{w_a^2 + \lambda^2(u^2 - y_a^2) - 2A\lambda^2(u - y_a) - 4\lambda F(u, y_a)}} = \pm(x - a), \quad F(u, v) = \int_v^u f(t) dt, \quad (2)$$

where $y_a = y(a)$ and $w_a = y'_x(a)$ are constants of integration. These constants, as well as the unknowns $y_b = y(b)$ and $w_b = y'_x(b)$, are determined by the algebraic (or transcendental) system

$$\begin{aligned} w_a + \lambda y_a &= A\lambda + 2B\lambda e^{\lambda a}, \\ w_b - \lambda y_b &= -A\lambda - 2C\lambda e^{-\lambda b}, \\ w_b^2 &= w_a^2 + \lambda^2(y_b^2 - y_a^2) - 2A\lambda^2(y_b - y_a) - 4\lambda F(y_b, y_a), \\ \int_{y_a}^{y_b} \frac{du}{\sqrt{w_a^2 + \lambda^2(u^2 - y_a^2) - 2A\lambda^2(u - y_a) - 4\lambda F(u, y_a)}} &= \pm(b - a). \end{aligned} \quad (3)$$

Here the first and second equations are obtained from conditions (5) in 8.8.16, and the third and fourth equations are consequences of (2).

Each solution of system (3) generates a solution of the integral equation.

$$3. \quad y(x) + \int_a^b e^{\lambda|x-t|} f(y(t)) dt = \beta \cosh(\lambda x).$$

This is a special case of equation 8.8.2 with $A = 0$ and $B = C = \frac{1}{2}\beta$.

$$4. \quad y(x) + \int_a^b e^{\lambda|x-t|} f(y(t)) dt = \beta \sinh(\lambda x).$$

This is a special case of equation 8.8.2 with $A = 0$, $B = \frac{1}{2}\beta$, and $C = -\frac{1}{2}\beta$.

$$5. \quad y(x) + \int_a^b \sinh(\lambda|x-t|) f(y(t)) dt = A + B \cosh(\lambda x) + C \sinh(\lambda x).$$

This is a special case of equation 8.8.17 with $f(t, y) = f(y)$ and $g(x) = A + B \cosh(\lambda x) + C \sinh(\lambda x)$.

The function $y = y(x)$ satisfies the second-order autonomous differential equation

$$y''_{xx} + 2\lambda f(y) - \lambda^2 y = -\lambda^2 A,$$

whose solution can be represented in an implicit form:

$$\int_{y_a}^y \frac{du}{\sqrt{w_a^2 + \lambda^2(u^2 - y_a^2) - 2A\lambda^2(u - y_a) - 4\lambda F(u, y_a)}} = \pm(x - a), \quad F(u, v) = \int_v^u f(t) dt,$$

where $y_a = y(a)$ and $w_a = y'_x(a)$ are constants of integration, which can be determined from the boundary conditions (5) in 8.8.17.

$$6. \quad y(x) + \int_a^b \sin(\lambda|x-t|) f(y(t)) dt = A + B \cos(\lambda x) + C \sin(\lambda x).$$

This is a special case of equation 8.8.18 with $f(t, y) = f(y)$ and $g(x) = A + B \cos(\lambda x) + C \sin(\lambda x)$.

The function $y = y(x)$ satisfies the second-order autonomous differential equation

$$y''_{xx} + 2\lambda f(y) + \lambda^2 y = \lambda^2 A,$$

whose solution can be represented in an implicit form:

$$\int_{y_a}^y \frac{du}{\sqrt{w_a^2 - \lambda^2(u^2 - y_a^2) + 2A\lambda^2(u - y_a) - 4\lambda F(u, y_a)}} = \pm(x - a), \quad F(u, v) = \int_v^u f(t) dt,$$

where $y_a = y(a)$ and $w_a = y'_x(a)$ are constants of integration, which can be determined from the boundary conditions (5) in 8.8.18.

8.8-2. Equations of the Form $y(x) + \int_a^b K(x, t)G(t, y(t)) dt = F(x)$.

$$7. \quad y(x) + \int_a^b f(t, y(t)) dt = g(x).$$

A solution: $y(x) = g(x) + \lambda$, where λ is determined by the algebraic (or transcendental) equation

$$\lambda + F(\lambda) = 0, \quad F(\lambda) = \int_a^b f(t, g(t) + \lambda) dt.$$

$$8. \quad y(x) + \int_a^b e^{\lambda(x-t)} f(t, y(t)) dt = g(x).$$

A solution: $y(x) = \beta e^{\lambda x} + g(x)$, where λ is determined by the algebraic (or transcendental) equation

$$\beta + F(\beta) = 0, \quad F(\beta) = \int_a^b e^{-\lambda t} f(t, \beta e^{\lambda t} + g(t)) dt.$$

9. $y(x) + \int_a^b g(x)f(t, y(t)) dt = h(x).$

A solution: $y(x) = \lambda g(x) + h(x)$, where λ is determined by the algebraic (or transcendental) equation

$$\lambda + F(\lambda) = 0, \quad F(\lambda) = \int_a^b f(t, \lambda g(t) + h(t)) dt.$$

10. $y(x) + \int_a^b (Ax + Bt)f(t, y(t)) dt = g(x).$

A solution: $y(x) = g(x) + \lambda x + \mu$, where the constants λ and μ are determined from the algebraic (or transcendental) system

$$\lambda + A \int_a^b f(t, g(t) + \lambda t + \mu) dt = 0, \quad \mu + B \int_a^b t f(t, g(t) + \lambda t + \mu) dt = 0.$$

11. $y(x) + \int_a^b \cosh(\lambda x + \mu t)f(t, y(t)) dt = h(x).$

Using the formula $\cosh(\lambda x + \mu t) = \cosh(\lambda x) \cosh(\mu t) + \sinh(\mu t) \sinh(\lambda x)$, we arrive at an equation of the form 8.8.19:

$$y(x) + \int_a^b [\cosh(\lambda x)f_1(t, y(t)) + \sinh(\lambda x)f_2(t, y(t))] dt = h(x),$$

$$f_1(t, y(t)) = \cosh(\mu t)f(t, y(t)), \quad f_2(t, y(t)) = \sinh(\mu t)f(t, y(t)).$$

12. $y(x) + \int_a^b \sinh(\lambda x + \mu t)f(t, y(t)) dt = h(x).$

Using the formula $\sinh(\lambda x + \mu t) = \cosh(\lambda x) \sinh(\mu t) + \cosh(\mu t) \sinh(\lambda x)$, we arrive at an equation of the form 8.8.19:

$$y(x) + \int_a^b [\cosh(\lambda x)f_1(t, y(t)) + \sinh(\lambda x)f_2(t, y(t))] dt = h(x),$$

$$f_1(t, y(t)) = \sinh(\mu t)f(t, y(t)), \quad f_2(t, y(t)) = \cosh(\mu t)f(t, y(t)).$$

13. $y(x) + \int_a^b \cos(\lambda x + \mu t)f(t, y(t)) dt = h(x).$

Using the formula $\cos(\lambda x + \mu t) = \cos(\lambda x) \cos(\mu t) - \sin(\mu t) \sin(\lambda x)$, we arrive at an equation of the form 8.8.19:

$$y(x) + \int_a^b [\cos(\lambda x)f_1(t, y(t)) + \sin(\lambda x)f_2(t, y(t))] dt = h(x),$$

$$f_1(t, y(t)) = \cos(\mu t)f(t, y(t)), \quad f_2(t, y(t)) = -\sin(\mu t)f(t, y(t)).$$

14. $y(x) + \int_a^b \sin(\lambda x + \mu t)f(t, y(t)) dt = h(x).$

Using the formula $\sin(\lambda x + \mu t) = \cos(\lambda x) \sin(\mu t) + \cos(\mu t) \sin(\lambda x)$, we arrive at an equation of the form 8.8.19:

$$y(x) + \int_a^b [\cos(\lambda x)f_1(t, y(t)) + \sin(\lambda x)f_2(t, y(t))] dt = h(x),$$

$$f_1(t, y(t)) = \sin(\mu t)f(t, y(t)), \quad f_2(t, y(t)) = \cos(\mu t)f(t, y(t)).$$

$$15. \quad y(x) + \int_a^b |x-t|f(t, y(t)) dt = g(x), \quad a \leq x \leq b.$$

1°. Let us remove the modulus in the integrand:

$$y(x) + \int_a^x (x-t)f(t, y(t)) dt + \int_x^b (t-x)f(t, y(t)) dt = g(x). \quad (1)$$

Differentiating (1) with respect to x yields

$$y'_x(x) + \int_a^x f(t, y(t)) dt - \int_x^b f(t, y(t)) dt = g'_x(x). \quad (2)$$

Differentiating (2), we arrive at a second-order ordinary differential equation for $y = y(x)$:

$$y''_{xx} + 2f(x, y) = g''_{xx}(x). \quad (3)$$

2°. Let us derive the boundary conditions for equation (3). We assume that $-\infty < a < b < \infty$. By setting $x = a$ and $x = b$ in (1), we obtain the relations

$$\begin{aligned} y(a) + \int_a^b (t-a)f(t, y(t)) dt &= g(a), \\ y(b) + \int_a^b (b-t)f(t, y(t)) dt &= g(b). \end{aligned} \quad (4)$$

Let us solve equation (3) for $f(x, y)$ and substitute the result into (4). Integrating by parts yields the desired boundary conditions for $y(x)$:

$$\begin{aligned} y(a) + y(b) + (b-a)[g'_x(b) - y'_x(b)] &= g(a) + g(b), \\ y(a) + y(b) + (a-b)[g'_x(a) - y'_x(a)] &= g(a) + g(b). \end{aligned} \quad (5)$$

Let us point out a useful consequence of (5):

$$y'_x(a) + y'_x(b) = g'_x(a) + g'_x(b), \quad (6)$$

which can be used together with one of conditions (5).

Equation (3) under the boundary conditions (5) determines the solution of the original integral equation (there may be several solutions). Conditions (5) make it possible to calculate the constants of integration that occur in solving the differential equation (3).

$$16. \quad y(x) + \int_a^b e^{\lambda|x-t|}f(t, y(t)) dt = g(x), \quad a \leq x \leq b.$$

1°. Let us remove the modulus in the integrand:

$$y(x) + \int_a^x e^{\lambda(x-t)}f(t, y(t)) dt + \int_x^b e^{\lambda(t-x)}f(t, y(t)) dt = g(x). \quad (1)$$

Differentiating (1) with respect to x twice yields

$$y''_{xx}(x) + 2\lambda f(x, y(x)) + \lambda^2 \int_a^x e^{\lambda(x-t)}f(t, y(t)) dt + \lambda^2 \int_x^b e^{\lambda(t-x)}f(t, y(t)) dt = g''_{xx}(x). \quad (2)$$

Eliminating the integral terms from (1) and (2), we arrive at a second-order ordinary differential equation for $y = y(x)$:

$$y''_{xx} + 2\lambda f(x, y) - \lambda^2 y = g''_{xx}(x) - \lambda^2 g(x). \tag{3}$$

2°. Let us derive the boundary conditions for equation (3). We assume that $-\infty < a < b < \infty$. By setting $x = a$ and $x = b$ in (1), we obtain the relations

$$\begin{aligned} y(a) + e^{-\lambda a} \int_a^b e^{\lambda t} f(t, y(t)) dt &= g(a), \\ y(b) + e^{\lambda b} \int_a^b e^{-\lambda t} f(t, y(t)) dt &= g(b). \end{aligned} \tag{4}$$

Let us solve equation (3) for $f(x, y)$ and substitute the result into (4). Integrating by parts yields

$$\begin{aligned} e^{\lambda b} \varphi'_x(b) - e^{\lambda a} \varphi'_x(a) &= \lambda e^{\lambda a} \varphi(a) + \lambda e^{\lambda b} \varphi(b), \quad \varphi(x) = y(x) - g(x); \\ e^{-\lambda b} \varphi'_x(b) - e^{-\lambda a} \varphi'_x(a) &= \lambda e^{-\lambda a} \varphi(a) + \lambda e^{-\lambda b} \varphi(b). \end{aligned}$$

Hence, we obtain the boundary conditions for $y(x)$:

$$\varphi'_x(a) + \lambda \varphi(a) = 0, \quad \varphi'_x(b) - \lambda \varphi(b) = 0; \quad \varphi(x) = y(x) - g(x). \tag{5}$$

Equation (3) under the boundary conditions (5) determines the solution of the original integral equation (there may be several solutions). Conditions (5) make it possible to calculate the constants of integration that occur in solving the differential equation (3).

17. $y(x) + \int_a^b \sinh(\lambda|x-t|) f(t, y(t)) dt = g(x), \quad a \leq x \leq b.$

1°. Let us remove the modulus in the integrand:

$$y(x) + \int_a^x \sinh[\lambda(x-t)] f(t, y(t)) dt + \int_x^b \sinh[\lambda(t-x)] f(t, y(t)) dt = g(x). \tag{1}$$

Differentiating (1) with respect to x twice yields

$$\begin{aligned} y''_{xx}(x) + 2\lambda f(x, y(x)) + \lambda^2 \int_a^x \sinh[\lambda(x-t)] f(t, y(t)) dt \\ + \lambda^2 \int_x^b \sinh[\lambda(t-x)] f(t, y(t)) dt = g''_{xx}(x). \end{aligned} \tag{2}$$

Eliminating the integral terms from (1) and (2), we arrive at a second-order ordinary differential equation for $y = y(x)$:

$$y''_{xx} + 2\lambda f(x, y) - \lambda^2 y = g''_{xx}(x) - \lambda^2 g(x). \tag{3}$$

2°. Let us derive the boundary conditions for equation (3). We assume that $-\infty < a < b < \infty$. By setting $x = a$ and $x = b$ in (1), we obtain the relations

$$\begin{aligned} y(a) + \int_a^b \sinh[\lambda(t-a)] f(t, y(t)) dt &= g(a), \\ y(b) + \int_a^b \sinh[\lambda(b-t)] f(t, y(t)) dt &= g(b). \end{aligned} \tag{4}$$

Let us solve equation (3) for $f(x, y)$ and substitute the result into (4). Integrating by parts yields

$$\begin{aligned} \sinh[\lambda(b-a)] \varphi'_x(b) - \lambda \cosh[\lambda(b-a)] \varphi(b) &= \lambda \varphi(a), \quad \varphi(x) = y(x) - g(x); \\ \sinh[\lambda(b-a)] \varphi'_x(a) + \lambda \cosh[\lambda(b-a)] \varphi(a) &= -\lambda \varphi(b). \end{aligned} \tag{5}$$

Equation (3) under the boundary conditions (5) determines the solution of the original integral equation (there may be several solutions). Conditions (5) make it possible to calculate the constants of integration that occur in solving the differential equation (3).

$$18. \quad y(x) + \int_a^b \sin[\lambda|x-t|] f(t, y(t)) dt = g(x), \quad a \leq x \leq b.$$

1°. Let us remove the modulus in the integrand:

$$y(x) + \int_a^x \sin[\lambda(x-t)] f(t, y(t)) dt + \int_x^b \sin[\lambda(t-x)] f(t, y(t)) dt = g(x). \quad (1)$$

Differentiating (1) with respect to x twice yields

$$\begin{aligned} y''_{xx}(x) + 2\lambda f(x, y(x)) - \lambda^2 \int_a^x \sin[\lambda(x-t)] f(t, y(t)) dt \\ - \lambda^2 \int_x^b \sin[\lambda(t-x)] f(t, y(t)) dt = g''_{xx}(x). \end{aligned} \quad (2)$$

Eliminating the integral terms from (1) and (2), we arrive at a second-order ordinary differential equation for $y = y(x)$:

$$y''_{xx} + 2\lambda f(x, y) + \lambda^2 y = g''_{xx}(x) + \lambda^2 g(x). \quad (3)$$

2°. Let us derive the boundary conditions for equation (3). We assume that $-\infty < a < b < \infty$. By setting $x = a$ and $x = b$ in (1), we obtain the relations

$$\begin{aligned} y(a) + \int_a^b \sin[\lambda(t-a)] f(t, y(t)) dt = g(a), \\ y(b) + \int_a^b \sin[\lambda(b-t)] f(t, y(t)) dt = g(b). \end{aligned} \quad (4)$$

Let us solve equation (3) for $f(x, y)$ and substitute the result into (4). Integrating by parts yields

$$\begin{aligned} \sin[\lambda(b-a)] \varphi'_x(b) - \lambda \cos[\lambda(b-a)] \varphi(b) = \lambda \varphi(a), \quad \varphi(x) = y(x) - g(x); \\ \sin[\lambda(b-a)] \varphi'_x(a) + \lambda \cos[\lambda(b-a)] \varphi(a) = -\lambda \varphi(b). \end{aligned} \quad (5)$$

Equation (3) under the boundary conditions (5) determines the solution of the original integral equation (there may be several solutions). Conditions (5) make it possible to calculate the constants of integration that occur in solving the differential equation (3).

8.8-3. Equations of the Form $y(x) + \int_a^b G(x, t, y(t)) dt = F(x)$.

$$19. \quad y(x) + \int_a^b [g_1(x)f_1(t, y(t)) + g_2(x)f_2(t, y(t))] dt = h(x).$$

A solution:

$$y(x) = h(x) + \lambda_1 g_1(x) + \lambda_2 g_2(x),$$

where the constants λ_1 and λ_2 are determined from the algebraic (or transcendental) system

$$\begin{aligned} \lambda_1 + \int_a^b f_1(t, h(t) + \lambda_1 g_1(t) + \lambda_2 g_2(t)) dt = 0, \\ \lambda_2 + \int_a^b f_2(t, h(t) + \lambda_1 g_1(t) + \lambda_2 g_2(t)) dt = 0. \end{aligned}$$

$$20. \quad y(x) + \int_a^b \left[\sum_{k=1}^n g_k(x) f_k(t, y(t)) \right] dt = h(x).$$

A solution:

$$y(x) = h(x) + \sum_{k=1}^n \lambda_k g_k(x),$$

where the coefficients λ_k are determined from the algebraic (or transcendental) system

$$\lambda_m + \int_a^b f_m \left(t, h(t) + \sum_{k=1}^n \lambda_k g_k(t) \right) dt = 0; \quad m = 1, \dots, n.$$

Different roots of this system generate different solutions of the integral equation.

⊙ Reference: A. F. Verlan' and V. S. Sizikov (1986).

$$21. \quad y(x) + \int_{-\infty}^{\infty} [\lambda e^{-|x-t|} y(t) + \varphi(x) \Psi(t, y(t))] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.2.14.

Solutions for $\lambda > -\frac{1}{2}$:

$$y_m(x) = Y_f(x) + A_m Y_\varphi(x),$$

where

$$Y_f(x) = f(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) f(t) dt,$$

$$Y_\varphi(x) = \varphi(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) \varphi(t) dt,$$

and A_m are roots of the algebraic (transcendental) equation

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

⊙ Reference: A. D. Polyanin and A. I. Zhurov (2007).

$$22. \quad y(x) - \int_0^{\infty} [\lambda \sin(xt) y(t) + \varphi(x) \Psi(t, y(t))] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.5.20.

Solutions for $\lambda \neq \pm \sqrt{\frac{2}{\pi}}$:

$$y_m(x) = Y_f(x) + A_m Y_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt) f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt) \varphi(t) dt,$$

and A_m are roots of the algebraic (transcendental) equation

$$A - \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

$$23. \quad y(x) - \int_0^{\infty} [\lambda \cos(xt)y(t) + \varphi(x)\Psi(t, y(t))] dt = f(x).$$

The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.5.6.

Solutions for $\lambda \neq \pm\sqrt{\frac{2}{\pi}}$:

$$y_m(x) = Y_f(x) + A_m Y_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \cos(xt)f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \cos(xt)\varphi(t) dt,$$

and A_m are roots of the algebraic (transcendental) equation

$$A - \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

$$24. \quad y(x) + \int_0^{\infty} [\lambda t J_\nu(xt)y(t) + \varphi(x)\Psi(t, y(t))] dt = f(x), \quad \nu > -1.$$

Here $J_\nu(z)$ is the Bessel function of the first kind. The solutions can be obtained by the methods described in Subsection 16.4-5; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 4.8.4.

Solutions for $\lambda \neq \pm 1$:

$$y_m(x) = Y_f(x) + A_m Y_\varphi(x),$$

where

$$Y_f(x) = \frac{f(x)}{1 - \lambda^2} - \frac{\lambda}{1 - \lambda^2} \int_0^{\infty} t J_\nu(xt)f(t) dt,$$

$$Y_\varphi(x) = \frac{\varphi(x)}{1 - \lambda^2} - \frac{\lambda}{1 - \lambda^2} \int_0^{\infty} t J_\nu(xt)\varphi(t) dt,$$

and A_m are roots of the algebraic (transcendental) equation

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0.$$

8.8-4. Equations of the Form $y(x) + \int_a^b y(xt)G(t, y(t)) dt = F(x)$.

$$25. \quad y(x) + \int_a^b y(xt)f(t, y(t)) dt = 0.$$

1°. A solution:

$$y(x) = kx^C, \tag{1}$$

where C is an arbitrary constant and the dependence $k = k(C)$ is determined by the algebraic (or transcendental) equation

$$1 + \int_a^b t^C f(t, kt^C) dt = 0. \tag{2}$$

Each root of equation (2) generates a solution of the integral equation which has the form (1).

2°. The integral equation can have some other solutions similar to those indicated in items 1°–3° of equation 8.2.16.

$$26. \quad y(x) + \int_a^b y(xt) f(t, y(t)) dt = Ax + B.$$

A solution:

$$y(x) = px + q, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + p \int_a^b t f(t, pt + q) dt - A &= 0, \\ q + q \int_a^b f(t, pt + q) dt - B &= 0. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$27. \quad y(x) + \int_a^b y(xt) f(t, y(t)) dt = Ax^\beta.$$

A solution:

$$y(x) = kx^\beta, \quad (1)$$

where k is a root of the algebraic (or transcendental) equation

$$k + kF(k) - A = 0, \quad F(k) = \int_a^b t^\beta f(t, kt^\beta) dt. \quad (2)$$

Each root of equation (2) generates a solution of the integral equation which has the form (1).

$$28. \quad y(x) + \int_a^b y(xt) f(t, y(t)) dt = A \ln x + B.$$

A solution:

$$y(x) = p \ln x + q, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + p \int_a^b f(t, p \ln t + q) dt - A &= 0, \\ q + \int_a^b (p \ln t + q) f(t, p \ln t + q) dt - B &= 0. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$29. \quad y(x) + \int_a^b y(xt) f(t, y(t)) dt = Ax^\beta \ln x.$$

A solution:

$$y(x) = px^\beta \ln x + qx^\beta, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + p \int_a^b t^\beta f(t, pt^\beta \ln t + qt^\beta) dt &= A, \\ q + \int_a^b (pt^\beta \ln t + qt^\beta) f(t, pt^\beta \ln t + qt^\beta) dt &= 0. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$30. \quad y(x) + \int_a^b y(xt) f(t, y(t)) dt = A \cos(\ln x).$$

A solution:

$$y(x) = p \cos(\ln x) + q \sin(\ln x),$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b [p \cos(\ln t) + q \sin(\ln t)] f(t, p \cos(\ln t) + q \sin(\ln t)) dt &= A, \\ q + \int_a^b [q \cos(\ln t) - p \sin(\ln t)] f(t, p \cos(\ln t) + q \sin(\ln t)) dt &= 0. \end{aligned}$$

$$31. \quad y(x) + \int_a^b y(xt) f(t, y(t)) dt = A \sin(\ln x).$$

A solution:

$$y(x) = p \cos(\ln x) + q \sin(\ln x),$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b [p \cos(\ln t) + q \sin(\ln t)] f(t, p \cos(\ln t) + q \sin(\ln t)) dt &= 0, \\ q + \int_a^b [q \cos(\ln t) - p \sin(\ln t)] f(t, p \cos(\ln t) + q \sin(\ln t)) dt &= A. \end{aligned}$$

$$32. \quad y(x) + \int_a^b y(xt) f(t, y(t)) dt = Ax^\beta \cos(\ln x) + Bx^\beta \sin(\ln x).$$

A solution:

$$y(x) = px^\beta \cos(\ln x) + qx^\beta \sin(\ln x), \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b t^\beta [p \cos(\ln t) + q \sin(\ln t)] f(t, pt^\beta \cos(\ln t) + qt^\beta \sin(\ln t)) dt &= A, \\ q + \int_a^b t^\beta [q \cos(\ln t) - p \sin(\ln t)] f(t, pt^\beta \cos(\ln t) + qt^\beta \sin(\ln t)) dt &= B. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

8.8-5. Equations of the Form $y(x) + \int_a^b y(x + \beta t) G(t, y(t)) dt = F(x)$.

$$33. \quad y(x) + \int_a^b y(x-t) f(t, y(t)) dt = 0.$$

1°. A solution:

$$y(x) = ke^{Cx}, \quad (1)$$

where C is an arbitrary constant and the dependence $k = k(C)$ is determined by the algebraic (or transcendental) equation

$$1 + \int_a^b f(t, ke^{Ct}) e^{-Ct} dt = 0. \quad (2)$$

Each root of equation (2) generates a solution of the integral equation which has the form (1).

2°. The equation has solutions of the form $y(x) = \sum_{m=0}^n E_m x^m$, where the constants E_m can be found by the method of undetermined coefficients.

$$34. \quad y(x) + \int_a^b y(x-t)f(t, y(t)) dt = Ax + B.$$

A solution:

$$y(x) = px + q, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + p \int_a^b f(t, pt + q) dt - A &= 0, \\ q + \int_a^b (q - pt)f(t, pt + q) dt - B &= 0. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$35. \quad y(x) + \int_a^b y(x-t)f(t, y(t)) dt = Ae^{\lambda x}.$$

Solutions:

$$y(x) = k_n e^{\lambda x},$$

where k_n are roots of the algebraic (or transcendental) equation

$$k + kF(k) - A = 0, \quad F(k) = \int_a^b f(t, ke^{\lambda t})e^{-\lambda t} dt.$$

$$36. \quad y(x) + \int_a^b y(x-t)f(t, y(t)) dt = A \sinh \lambda x.$$

A solution:

$$y(x) = p \sinh \lambda x + q \cosh \lambda x, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b (p \cosh \lambda t - q \sinh \lambda t)f(t, p \sinh \lambda t + q \cosh \lambda t) dt &= A, \\ q + \int_a^b (q \cosh \lambda t - p \sinh \lambda t)f(t, p \sinh \lambda t + q \cosh \lambda t) dt &= 0. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$37. \quad y(x) + \int_a^b y(x-t)f(t, y(t)) dt = A \cosh \lambda x.$$

A solution:

$$y(x) = p \sinh \lambda x + q \cosh \lambda x,$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b (p \cosh \lambda t - q \sinh \lambda t)f(t, p \sinh \lambda t + q \cosh \lambda t) dt &= 0, \\ q + \int_a^b (q \cosh \lambda t - p \sinh \lambda t)f(t, p \sinh \lambda t + q \cosh \lambda t) dt &= A. \end{aligned}$$

$$38. \quad y(x) + \int_a^b y(x-t)f(t, y(t)) dt = A \sin \lambda x.$$

A solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b (p \cos \lambda t + q \sin \lambda t) f(t, p \sin \lambda t + q \cos \lambda t) dt &= A, \\ q + \int_a^b (q \cos \lambda t - p \sin \lambda t) f(t, p \sin \lambda t + q \cos \lambda t) dt &= 0. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$39. \quad y(x) + \int_a^b y(x-t)f(t, y(t)) dt = A \cos \lambda x.$$

A solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x,$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b (p \cos \lambda t + q \sin \lambda t) f(t, p \sin \lambda t + q \cos \lambda t) dt &= 0, \\ q + \int_a^b (q \cos \lambda t - p \sin \lambda t) f(t, p \sin \lambda t + q \cos \lambda t) dt &= A. \end{aligned}$$

$$40. \quad y(x) + \int_a^b y(x-t)f(t, y(t)) dt = e^{\mu x} (A \sin \lambda x + B \cos \lambda x).$$

A solution:

$$y(x) = e^{\mu x} (p \sin \lambda x + q \cos \lambda x), \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b (p \cos \lambda t + q \sin \lambda t) e^{-\mu t} f(t, p e^{\mu t} \sin \lambda t + q e^{\mu t} \cos \lambda t) dt &= A, \\ q + \int_a^b (q \cos \lambda t - p \sin \lambda t) e^{-\mu t} f(t, p e^{\mu t} \sin \lambda t + q e^{\mu t} \cos \lambda t) dt &= B. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$41. \quad y(x) + \int_a^b y(x-t)f(t, y(t)) dt = g(x).$$

1°. For $g(x) = \sum_{k=1}^n A_k \exp(\lambda_k x)$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \exp(\lambda_k x),$$

where the constants B_k are determined from the nonlinear algebraic (or transcendental) system

$$\begin{aligned} B_k + B_k F_k(\vec{B}) - A_k &= 0, \quad k = 1, \dots, n, \\ \vec{B} = \{B_1, \dots, B_n\}, \quad F_k(\vec{B}) &= \int_a^b f\left(t, \sum_{m=1}^n B_m \exp(\lambda_m t)\right) \exp(-\lambda_k t) dt. \end{aligned}$$

Different solutions of this system generate different solutions of the integral equation.

2°. For a polynomial right-hand side, $g(x) = \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

3°. For $g(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k x)$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k x)$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

6°. For $g(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

7°. For $g(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

8°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \cos(\lambda_k x)$, the equation has a solution of the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

9°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \sin(\lambda_k x)$, the equation has a solution of the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

10°. For $g(x) = \cos(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, the equation has a solution of the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

11°. For $g(x) = \sin(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, the equation has a solution of the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

42. $y(x) + \int_a^b y(x + \beta t) f(t, y(t)) dt = Ax + B.$

A solution:

$$y(x) = px + q, \tag{1}$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + p \int_a^b f(t, pt + q) dt - A &= 0, \\ q + \int_a^b (\beta pt + q) f(t, pt + q) dt - B &= 0. \end{aligned} \tag{2}$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

43. $y(x) + \int_a^b y(x + \beta t) f(t, y(t)) dt = Ae^{\lambda x}.$

Solutions:

$$y(x) = k_n e^{\lambda x},$$

where k_n are roots of the algebraic (or transcendental) equation

$$k + kF(k) - A = 0, \quad F(k) = \int_a^b f(t, ke^{\lambda t}) e^{\beta \lambda t} dt.$$

$$44. \quad y(x) + \int_a^b y(x + \beta t) f(t, y(t)) dt = A \sin \lambda x + B \cos \lambda x.$$

A solution:

$$y(x) = p \sin \lambda x + q \cos \lambda x, \quad (1)$$

where p and q are roots of the following system of algebraic (or transcendental) equations:

$$\begin{aligned} p + \int_a^b [p \cos(\lambda \beta t) - q \sin(\lambda \beta t)] f(t, p \sin \lambda t + q \cos \lambda t) dt &= A, \\ q + \int_a^b [q \cos(\lambda \beta t) + p \sin(\lambda \beta t)] f(t, p \sin \lambda t + q \cos \lambda t) dt &= B. \end{aligned} \quad (2)$$

Different solutions of system (2) generate different solutions (1) of the integral equation.

$$45. \quad y(x) + \int_a^b y(x + \beta t) f(t, y(t)) dt = g(x).$$

1°. For $g(x) = \sum_{k=1}^n A_k \exp(\lambda_k x)$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \exp(\lambda_k x),$$

where the constants B_k are determined from the nonlinear algebraic (or transcendental) system

$$\begin{aligned} B_k + B_k F_k(\vec{B}) - A_k &= 0, \quad k = 1, \dots, n, \\ \vec{B} = \{B_1, \dots, B_n\}, \quad F_k(\vec{B}) &= \int_a^b f\left(t, \sum_{m=1}^n B_m \exp(\lambda_m t)\right) \exp(\lambda_k \beta t) dt. \end{aligned}$$

Different solutions of this system generate different solutions of the integral equation.

2°. For a polynomial right-hand side, $g(x) = \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \sum_{k=0}^n B_k x^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

3°. For $g(x) = e^{\lambda x} \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = e^{\lambda x} \sum_{k=0}^n B_k x^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k x)$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k x)$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k x) + \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

6°. For $g(x) = \cos(\lambda x) \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

7°. For $g(x) = \sin(\lambda x) \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \cos(\lambda x) \sum_{k=0}^n B_k x^k + \sin(\lambda x) \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

8°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \cos(\lambda_k x)$, the equation has a solution of the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

9°. For $g(x) = e^{\mu x} \sum_{k=1}^n A_k \sin(\lambda_k x)$, the equation has a solution of the form

$$y(x) = e^{\mu x} \sum_{k=1}^n B_k \cos(\lambda_k x) + e^{\mu x} \sum_{k=1}^n C_k \sin(\lambda_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

10°. For $g(x) = \cos(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, the equation has a solution of the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

11°. For $g(x) = \sin(\lambda x) \sum_{k=1}^n A_k \exp(\mu_k x)$, the equation has a solution of the form

$$y(x) = \cos(\lambda x) \sum_{k=1}^n B_k \exp(\mu_k x) + \sin(\lambda x) \sum_{k=1}^n C_k \exp(\mu_k x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

8.8-6. Other Equations.

$$46. \quad y(x) + \int_a^b y(x) f(t, y(t)) dt = g(x).$$

A solution: $y(x) = \lambda g(x)$, where λ is determined by the algebraic (or transcendental) equation

$$\lambda + \lambda F(\lambda) - 1 = 0, \quad F(\lambda) = \int_a^b f(t, \lambda g(t)) dt.$$

$$47. \quad y(x) + \int_a^b g(x) y(x) f(t, y(t)) dt = h(x).$$

A solution: $y(x) = \frac{h(x)}{1 + \lambda g(x)}$, where λ is determined from the algebraic (or transcendental) equation

$$\lambda - F(\lambda) = 0, \quad F(\lambda) = \int_a^b f\left(t, \frac{h(t)}{1 + \lambda g(t)}\right) dt.$$

$$48. \quad y(x) + \int_a^b g(x, y(x)) f(t, y(t)) dt = h(x).$$

Solution in an implicit form:

$$y(x) + \lambda g(x, y(x)) - h(x) = 0, \quad (1)$$

where λ is determined from the algebraic (or transcendental) equation

$$\lambda - F(\lambda) = 0, \quad F(\lambda) = \int_a^b f(t, y(t)) dt. \quad (2)$$

Here the function $y(x) = y(x, \lambda)$ obtained by solving (1) must be substituted into (2).

The number of solutions of the integral equation is determined by the number of the solutions obtained from (1) and (2).

$$49. \quad f(x, y(x)) + \int_a^b \left[\sum_{k=1}^n g_k(x, y(x)) h_k(t, y(t)) \right] dt = 0.$$

Solution in an implicit form:

$$f(x, y(x)) + \sum_{k=1}^n \lambda_k g_k(x, y(x)) = 0, \quad (1)$$

where the λ_k are determined from the algebraic (or transcendental) system

$$\begin{aligned} \lambda_k - H_k(\vec{\lambda}) &= 0, \quad k = 1, \dots, n; \\ H_k(\vec{\lambda}) &= \int_a^b h_k(t, y(t)) dt, \quad \vec{\lambda} = \{\lambda_1, \dots, \lambda_n\}. \end{aligned} \quad (2)$$

Here the function $y(x) = y(x, \vec{\lambda})$ obtained by solving (1) must be substituted into (2).

The number of solutions of the integral equation is determined by the number of the solutions obtained from (1) and (2).

$$50. \quad y(x) + \int_a^b y(xt^\beta) f(t, y(t)) dt = g(x), \quad \beta > 0.$$

1°. For $g(x) = \sum_{k=1}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k x^k,$$

where B_k are roots of the algebraic (or transcendental) equations

$$B_k + B_k F_k(\vec{B}) - A_k = 0, \quad F_k(\vec{B}) = \int_a^b t^{k\beta} f\left(t, \sum_{m=1}^n B_m t^m\right) dt.$$

Different roots of this system generate different solutions of the integral equation.

2°. For $g(x) = \ln x \sum_{k=0}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \ln x \sum_{k=0}^n B_k x^k + \sum_{k=0}^n C_k x^k,$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

3°. For $g(x) = \sum_{k=0}^n A_k (\ln x)^k$, the equation has a solution of the form

$$y(x) = \sum_{k=0}^n B_k (\ln x)^k,$$

where the constants B_k can be found by the method of undetermined coefficients.

4°. For $g(x) = \sum_{k=1}^n A_k \cos(\lambda_k \ln x)$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k \ln x) + \sum_{k=1}^n C_k \sin(\lambda_k \ln x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

5°. For $g(x) = \sum_{k=1}^n A_k \sin(\lambda_k \ln x)$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \cos(\lambda_k \ln x) + \sum_{k=1}^n C_k \sin(\lambda_k \ln x),$$

where the constants B_k and C_k can be found by the method of undetermined coefficients.

$$51. \quad y(x) + \int_a^b y(\xi) f(t, y(t)) dt = 0, \quad \xi = x\varphi(t).$$

1°. A solution:

$$y(x) = kx^C, \quad (1)$$

where C is an arbitrary constant and the dependence $k = k(C)$ is determined by the algebraic (or transcendental) equation

$$1 + \int_a^b [\varphi(t)]^C f(t, kt^C) dt = 0. \quad (2)$$

Each root of equation (2) generates a solution of the integral equation which has the form (1).

2°. The equation has solutions of the form $y(x) = \sum_{m=0}^n E_m x^m$, where the constants E_m can be found by the method of undetermined coefficients.

$$52. \quad y(x) + \int_a^b y(\xi) f(t, y(t)) dt = g(x), \quad \xi = x\varphi(t).$$

1°. For $g(x) = \sum_{k=1}^n A_k x^k$, the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k x^k,$$

where B_k are roots of the algebraic (or transcendental) equations

$$B_k + B_k F_k(\vec{B}) - A_k = 0, \quad k = 1, \dots, n,$$

$$\vec{B} = \{B_1, \dots, B_n\}, \quad F_k(\vec{B}) = \int_a^b [\varphi(t)]^k f\left(t, \sum_{m=1}^n B_m t^m\right) dt.$$

Different roots generate different solutions of the integral equation.

2°. A form of solutions for some other functions $g(x)$ can be found in items 2°–5° of equation 8.8.50.

$$53. \quad y(x) + \int_a^b y(\xi) f(t, y(t)) dt = 0, \quad \xi = x + \varphi(t).$$

1°. A solution:

$$y(x) = ke^{Cx}, \quad (1)$$

where C is an arbitrary constant and the dependence $k = k(C)$ is determined by the algebraic (or transcendental) equation

$$1 + \int_a^b e^{C\varphi(t)} f(t, ke^{Ct}) dt = 0. \quad (2)$$

Each root of equation (2) generates a solution of the integral equation which has the form (1).

2°. The equation has a solution of the form $y(x) = \sum_{m=0}^n E_m x^m$, where the constants E_m can be found by the method of undetermined coefficients.

$$54. \quad y(x) + \int_a^b y(\xi) f(t, y(t)) dt = g(x), \quad \xi = x + \varphi(t).$$

1°. For $g(x) = \sum_{k=1}^n A_k \exp(\lambda_k x)$ the equation has a solution of the form

$$y(x) = \sum_{k=1}^n B_k \exp(\lambda_k x),$$

where the constants B_k are determined from the nonlinear algebraic (or transcendental) system

$$B_k + B_k F_k(\vec{B}) - A_k = 0, \quad k = 1, \dots, n,$$

$$\vec{B} = \{B_1, \dots, B_n\}, \quad F_k(\vec{B}) = \int_a^b f\left(t, \sum_{m=1}^n B_m \exp(\lambda_m t)\right) \exp[\lambda_k \varphi(t)] dt.$$

2°. A form of solutions for some other functions $g(x)$ can be found in items 2°–11° of equation 8.8.45.

Part II

**Methods for Solving
Integral Equations**

Chapter 9

Main Definitions and Formulas. Integral Transforms

9.1. Some Definitions, Remarks, and Formulas

9.1-1. Some Definitions.

A function $f(x)$ is said to be *square integrable* on an interval $[a, b]$ if $f^2(x)$ is integrable on $[a, b]$. The set of all square integrable functions is denoted by $L_2(a, b)$ or, briefly, L_2 .^{*} Likewise, the set of all integrable functions on $[a, b]$ is denoted by $L_1(a, b)$ or, briefly, L_1 .

Let us list the main properties of functions from L_2 .

- 1°. The sum of two square integrable functions is a square integrable function.
- 2°. The product of a square integrable function by a constant is a square integrable function.
- 3°. The product of two square integrable functions is an integrable function.
- 4°. If $f(x) \in L_2$ and $g(x) \in L_2$, then the following Cauchy–Schwarz–Bunyakovsky inequality holds:

$$(f, g)^2 \leq \|f\|^2 \|g\|^2,$$
$$(f, g) = \int_a^b f(x)g(x) dx, \quad \|f\|^2 = (f, f) = \int_a^b f^2(x) dx.$$

The number (f, g) is called the *inner product* of the functions $f(x)$ and $g(x)$ and the number $\|f\|$ is called the L_2 -norm of $f(x)$.

- 5°. For $f(x) \in L_2$ and $g(x) \in L_2$, the following *triangle inequality* holds:

$$\|f + g\| \leq \|f\| + \|g\|.$$

- 6°. Let functions $f(x)$ and $f_1(x), f_2(x), \dots, f_n(x), \dots$ be square integrable on an interval $[a, b]$. If

$$\lim_{n \rightarrow \infty} \int_a^b [f_n(x) - f(x)]^2 dx = 0,$$

then the sequence $f_1(x), f_2(x), \dots$ is said to be *mean-square convergent* to $f(x)$.

Note that if a sequence of functions $\{f_n(x)\}$ from L_2 converges uniformly to $f(x)$, then $f(x) \in L_2$ and $\{f_n(x)\}$ is mean-square convergent to $f(x)$.

^{*} In the most general case the integral is understood as the Lebesgue integral of measurable functions (see Supplement 12.3). As usual, two equivalent functions (i.e., equal everywhere, or distinct on a negligible set (of zero measure)) are regarded as one and the same element of L_2 .

The notion of an integrable function of several variables is similar. For instance, a function $f(x, t)$ is said to be *square integrable* in a domain $S = \{a \leq x \leq b, a \leq t \leq b\}$ if $f(x, t)$ is measurable and

$$\|f\|^2 \equiv \int_a^b \int_a^b f^2(x, t) dx dt < \infty.$$

Here $\|f\|$ denotes the norm of the function $f(x, t)$, as above.

9.1-2. Structure of Solutions to Linear Integral Equations.

A linear integral equation with variable integration limit has the form

$$\beta y(x) + \int_a^x K(x, t)y(t) dt = f(x), \quad (1)$$

where $y(x)$ is the unknown function.

A linear integral equation with constant integration limits has the form

$$\beta y(x) + \int_a^b K(x, t)y(t) dt = f(x). \quad (2)$$

For $\beta = 0$, Eqs. (1) and (2) are called *linear integral equations of the first kind*, and for $\beta \neq 0$, *linear integral equations of the second kind*.*

Equations of the form (1) and (2) with specific conditions imposed on the kernels and the right-hand sides form various classes of integral equations (Volterra equations, Fredholm equations, convolution equations, etc.), which are considered in detail in Chapters 10–14.

For brevity, we shall sometimes represent the linear equations (1) and (2) in the operator form

$$\mathbf{L}[y] = f(x). \quad (3)$$

A linear operator \mathbf{L} possesses the properties

$$\begin{aligned} \mathbf{L}[y_1 + y_2] &= \mathbf{L}[y_1] + \mathbf{L}[y_2], \\ \mathbf{L}[\sigma y] &= \sigma \mathbf{L}[y], \quad \sigma = \text{const.} \end{aligned}$$

A linear equation is called *homogeneous* if $f(x) \equiv 0$ and *nonhomogeneous* otherwise.

An arbitrary homogeneous linear integral equation has the trivial solution $y \equiv 0$.

If $y_1 = y_1(x)$ and $y_2 = y_2(x)$ are particular solutions of a linear homogeneous integral equation, then the linear combination $C_1 y_1 + C_2 y_2$ with arbitrary constants C_1 and C_2 is also a solution (in physical problems, this property is called the *linear superposition principle*).

The general solution of a linear nonhomogeneous integral equation (3) is the sum of the general solution $Y = Y(x)$ of the corresponding homogeneous equation $\mathbf{L}[Y] = 0$ and an arbitrary particular solution $\bar{y} = \bar{y}(x)$ of the nonhomogeneous equation $\mathbf{L}[\bar{y}] = f(x)$, that is,

$$y = Y + \bar{y}. \quad (4)$$

If the homogeneous integral equation has only the trivial solution $Y \equiv 0$, then the solution of the corresponding nonhomogeneous equation is unique (if it exists).

Let \bar{y}_1 and \bar{y}_2 be solutions of nonhomogeneous linear integral equations with the same left-hand sides and different right-hand sides, $\mathbf{L}[\bar{y}_1] = f_1(x)$ and $\mathbf{L}[\bar{y}_2] = f_2(x)$. Then the function $\bar{y} = \bar{y}_1 + \bar{y}_2$ is a solution of the equation $\mathbf{L}[\bar{y}] = f_1(x) + f_2(x)$.

The transformation

$$x = g(z), \quad t = g(\tau), \quad y(x) = \varphi(z)w(z) + \psi(z), \quad (5)$$

where $g(z)$, $\varphi(z)$, and $\psi(z)$ are arbitrary continuous functions ($g'_z \neq 0$), reduces Eqs. (1) and (2) to linear equations of the same form for the unknown function $w = w(z)$. Such transformations are frequently used for constructing exact solutions of linear integral equations.

* In Chapters 1–4, which deal with equations with variable and constant limits of integration, we sometimes consider more general equations in which the integrand contains the unknown function $y(z)$, where $z = z(x, t)$, instead of $y(t)$.

9.1-3. Integral Transforms.

Integral transforms have the form

$$\tilde{f}(\lambda) = \int_a^b \varphi(x, \lambda) f(x) dx.$$

The function $\tilde{f}(\lambda)$ is called the *transform* of the function $f(x)$ and $\varphi(x, \lambda)$ is called the *kernel* of the integral transform. The function $f(x)$ is called the *inverse transform* of $\tilde{f}(\lambda)$. The limits of integration a and b are real numbers (usually, $a = 0$, $b = \infty$ or $a = -\infty$, $b = \infty$).

In Subsections 9.2–9.6, the most popular (Laplace, Mellin, Fourier, etc.) integral transforms, applied in this book to the solution of specific integral equations, are described. These subsections also describe the corresponding inversion formulas, which have the form

$$f(x) = \int_{\mathcal{L}} \psi(x, \lambda) \tilde{f}(\lambda) d\lambda$$

and make it possible to recover $f(x)$ if $\tilde{f}(\lambda)$ is given. The integration path \mathcal{L} can lie either on the real axis or in the complex plane.

Integral transforms are used in the solution of various differential and integral equations (see, for example, Sections 10.4, 11.3, 11.6, 12.5, and 13.9). Figure 1 outlines the overall scheme of solving some special classes of linear integral equations by means of integral transforms (by applying appropriate integral transforms to this sort of integral equations, one obtains first-order linear algebraic equations for $\tilde{f}(\lambda)$).




Figure 1. Principal scheme of applying integral transforms for solving integral equations.

In many cases, to calculate definite integrals, in particular, to find the inverse Laplace, Mellin, and Fourier transforms, methods of the theory of functions of a complex variable can be applied, including the residue theorem and the Jordan lemma, which are presented below in Subsections 9.1-4 and 9.1-5.

9.1-4. Residues. Calculation Formulas. Cauchy's Residue Theorem.

1°. The *residue* of a function $f(z)$ holomorphic in a deleted neighborhood of a point $z = a$ (thus, a is an isolated singularity of f) of the complex plane z is the number

$$\operatorname{res}_{z=a} f(z) = \frac{1}{2\pi i} \int_{c_\varepsilon} f(z) dz, \quad i^2 = -1,$$

where c_ε is a circle of sufficiently small radius ε described by the equation $|z - a| = \varepsilon$.

If the point $z = a$ is a pole of order n^* of the function $f(z)$, then we have

$$\operatorname{res}_{z=a} f(z) = \frac{1}{(n-1)!} \lim_{z \rightarrow a} \frac{d^{n-1}}{dz^{n-1}} [(z-a)^n f(z)].$$

For a simple pole, which corresponds to $n = 1$, this implies

$$\operatorname{res}_{z=a} f(z) = \lim_{z \rightarrow a} [(z-a)f(z)].$$

If $f(z) = \frac{\varphi(z)}{\psi(z)}$, where $\varphi(a) \neq 0$ and $\psi(z)$ has a simple zero at the point $z = a$, i.e., $\psi(a) = 0$ and $\psi'_z(a) \neq 0$, then

$$\operatorname{res}_{z=a} f(z) = \frac{\varphi(a)}{\psi'_z(a)}.$$

2°. A function $f(z)$ is said to be continuous on the boundary C of the domain D if for each boundary point z_0 there exists a limit $\lim_{z \rightarrow z_0} f(z) = f(z_0)$ as $z \rightarrow z_0$, $z \in D$.

CAUCHY'S RESIDUE THEOREM. Let $f(z)$ be a function continuous on the boundary C of a domain D and analytic in the interior of D everywhere except for finitely many points a_1, \dots, a_n . Then

$$\int_C f(z) dz = 2\pi i \sum_{k=1}^n \operatorname{res} f(a_k),$$

where the integral is taken in the positive sense of C .

The residue of a function $f(z)$ at infinity is defined as

$$\operatorname{res} f(\infty) = \frac{1}{2\pi i} \oint_{\Gamma} f(z) dz,$$

where Γ is a circle of sufficiently large radius $|z| = \rho$ and the integral is taken in the clockwise sense (so that the neighborhood of the point $z = \infty$ remains to the left of the contour, just as in the case of a finite point).

Note that

$$\operatorname{res} f(\infty) = \lim_{z \rightarrow \infty} [-zf(z)],$$

provided that this limit exists.

THEOREM. If a function $f(z)$ has finitely many singular points a_1, \dots, a_n in the extended complex plane, then the sum of all its residues, including the residue at infinity, is zero:

$$\operatorname{res} f(\infty) + \sum_{k=1}^n \operatorname{res} f(a_k) = 0.$$

* In a neighborhood of this point we have $f(z) \approx \text{const} (z-a)^{-n}$.

9.1-5. Jordan Lemma.

JORDAN LEMMA. If a function $f(z)$ is continuous in the domain $|z| \geq R_0$, $\text{Im } z \geq \alpha$, where α is a chosen real number, and if $\lim_{z \rightarrow \infty} f(z) = 0$, then

$$\lim_{R \rightarrow \infty} \int_{C_R} e^{i\lambda z} f(z) dz = 0$$

for any $\lambda > 0$, where C_R is the arc of the circle $|z| = R$ that lies in this domain.

If a function $f(z)$ is analytic for $|z| > R_0$ and $zf(z) \rightarrow 0$ as $|z| \rightarrow \infty$ for $y \geq 0$ (or $x \geq 0$), then

$$\lim_{R \rightarrow \infty} \int_{C_R} f(z) dz = 0,$$

where C_R is the arc of the circle $|z| = R$ in the upper half-plane (or right half-plane).

⊙ References for Section 9.1: A. G. Sveshnikov and A. N. Tikhonov (1970), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), W. R. LePage (1980), A. D. Polyanin and A. V. Manzhirov (1998), A. N. Kolmogorov and S. V. Fomin (1999), S. G. Krantz (1999).

9.2. Laplace Transform

9.2-1. Definition. Inversion Formula.

The Laplace transform of an arbitrary (complex-valued) function $f(x)$ of a real variable x ($x \geq 0$) is defined by

$$\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx, \quad (1)$$

where $p = s + i\sigma$ is a complex variable.

The Laplace transform exists for any continuous or piecewise-continuous function satisfying the condition $|f(x)| < Me^{\sigma_0 x}$ with some $M > 0$ and $\sigma_0 \geq 0$. In the following, σ_0 often means the greatest lower bound of the possible values of σ_0 in this estimate; this value is called the *growth exponent* of the function $f(x)$.

For any $f(x)$, the transform $\tilde{f}(p)$ is defined in the half-plane $\text{Re } p > \sigma_0$ and is analytic there.

For brevity, we shall write formula (1) as follows:

$$\tilde{f}(p) = \mathfrak{L}\{f(x)\}, \quad \text{or} \quad \tilde{f}(p) = \mathfrak{L}\{f(x), p\}.$$

Given the transform $\tilde{f}(p)$, the function can be found by means of the inverse Laplace transform

$$f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \tilde{f}(p)e^{px} dp, \quad i^2 = -1, \quad (2)$$

where the integration path is parallel to the imaginary axis and lies to the right of all singularities of $\tilde{f}(p)$, which corresponds to $c > \sigma_0$.

The integral in (2) is understood in the sense of the Cauchy principal value:

$$\int_{c-i\infty}^{c+i\infty} \tilde{f}(p)e^{px} dp = \lim_{\omega \rightarrow \infty} \int_{c-i\omega}^{c+i\omega} \tilde{f}(p)e^{px} dp.$$

In the domain $x < 0$, formula (2) gives $f(x) \equiv 0$.

Formula (2) holds for continuous functions. If $f(x)$ has a (finite) jump discontinuity at a point $x = x_0 > 0$, then the left-hand side of (2) is equal to $\frac{1}{2}[f(x_0 - 0) + f(x_0 + 0)]$ at this point (for $x_0 = 0$, the first term in the square brackets must be omitted).

For brevity, we write the Laplace inversion formula (2) as follows:

$$f(x) = \mathfrak{L}^{-1}\{\tilde{f}(p)\}, \quad \text{or} \quad f(x) = \mathfrak{L}^{-1}\{\tilde{f}(p), x\}.$$

9.2-2. Inverse Transforms of Rational Functions.

Consider the important case in which the transform is a rational function of the form

$$\tilde{f}(p) = \frac{R(p)}{Q(p)}, \quad (3)$$

where $Q(p)$ and $R(p)$ are polynomials in the variable p and the degree of $Q(p)$ exceeds that of $R(p)$.

Assume that the zeros of the denominator are simple, i.e.,

$$Q(p) \equiv \text{const}(p - \lambda_1)(p - \lambda_2) \dots (p - \lambda_n).$$

Then the inverse transform can be determined by the formula

$$f(x) = \sum_{k=1}^n \frac{R(\lambda_k)}{Q'(\lambda_k)} \exp(\lambda_k x), \quad (4)$$

where the primes denote the derivatives.

If $Q(p)$ has multiple zeros, i.e.,

$$Q(p) \equiv \text{const}(p - \lambda_1)^{s_1}(p - \lambda_2)^{s_2} \dots (p - \lambda_m)^{s_m},$$

then

$$f(x) = \sum_{k=1}^m \frac{1}{(s_k - 1)!} \lim_{p \rightarrow s_k} \frac{d^{s_k-1}}{dp^{s_k-1}} [(p - \lambda_k)^{s_k} \tilde{f}(p) e^{px}]. \quad (5)$$

Example 1. The transform

$$\tilde{f}(p) = \frac{b}{p^2 - a^2} \quad (a, b \text{ real numbers})$$

can be represented as the fraction (3) with $R(p) = b$ and $Q(p) = (p - a)(p + a)$. The denominator $Q(p)$ has two simple roots, $\lambda_1 = a$ and $\lambda_2 = -a$. Using formula (4) with $n = 2$ and $Q'(p) = 2p$, we obtain the inverse transform in the form

$$f(x) = \frac{b}{2a} e^{ax} - \frac{b}{2a} e^{-ax} = \frac{b}{a} \sinh(ax).$$

Example 2. The transform

$$\tilde{f}(p) = \frac{b}{p^2 + a^2} \quad (a, b \text{ real numbers})$$

can be written as the fraction (3) with $R(p) = b$ and $Q(p) = (p - ia)(p + ia)$, $i^2 = -1$. The denominator $Q(p)$ has two simple pure imaginary roots, $\lambda_1 = ia$ and $\lambda_2 = -ia$. Using formula (4) with $n = 2$, we find the inverse transform:

$$f(x) = \frac{b}{2ia} e^{iax} - \frac{b}{2ia} e^{-iax} = -\frac{bi}{2a} [\cos(ax) + i \sin(ax)] + \frac{bi}{2a} [\cos(ax) - i \sin(ax)] = \frac{b}{a} \sin(ax).$$

Example 3. The transform

$$\tilde{f}(p) = ap^{-n},$$

where n is a positive integer, can be written as the fraction (3) with $R(p) = a$ and $Q(p) = p^n$. The denominator $Q(p)$ has one root of multiplicity n , $\lambda_1 = 0$. By formula (5) with $m = 1$ and $s_1 = n$, we find the inverse transform:

$$f(x) = \frac{a}{(n-1)!} x^{n-1}.$$

► Fairly detailed tables of inverse Laplace transforms can be found in Supplement 6.

9.2-3. Inversion of Functions with Finitely Many Singular Points.

If the function $\tilde{f}(p)$ has finitely many singular points, p_1, p_2, \dots, p_n , and tends to zero as $p \rightarrow \infty$, then the integral in the Laplace inversion formula (2) may be evaluated using the residue theory by applying the Jordan lemma (see Subsection 9.1-5). In this case

$$f(x) = \sum_{k=1}^n \operatorname{res}_{p=p_k} [\tilde{f}(p)e^{px}]. \quad (6)$$

Formula (6) can be extended to the case where $\tilde{f}(p)$ has infinitely many singular points. In this case, $f(x)$ is represented as an infinite series.

9.2-4. Convolution Theorem. Main Properties of the Laplace Transform.

1°. The *convolution* of two functions $f(x)$ and $g(x)$ is defined as an integral $\int_0^x f(t)g(x-t) dt$, and is usually denoted by $f(x) * g(x)$,

$$f(x) * g(x) = \int_0^x f(t)g(x-t) dt.$$

By performing substitution $x - t = u$, we see that the convolution is symmetric with respect to the convolved functions: $f(x) * g(x) = g(x) * f(x)$.

The *convolution theorem* states that

$$\mathfrak{L} \{ f(x) * g(x) \} = \mathfrak{L} \{ f(x) \} \mathfrak{L} \{ g(x) \}$$

and is frequently applied to solve Volterra equations with kernels depending on the difference of the arguments.

2°. The main properties of the correspondence between functions and their Laplace transforms are gathered in Table 1.

3°. The Laplace transforms of some functions are listed in Table 2; for more detailed tables of direct and inverse Laplace transforms, see Supplements 5–6 and the list of references at the end of this section.

9.2-5. Limit Theorems.

THEOREM 1. Let $0 \leq x < \infty$ and $\tilde{f}(p) = \mathfrak{L} \{ f(x) \}$ be the Laplace transform of $f(x)$. If a limit of $f(x)$ as $x \rightarrow 0$ exists, then

$$\lim_{x \rightarrow 0} f(x) = \lim_{p \rightarrow \infty} [p\tilde{f}(p)].$$

THEOREM 2. If a limit of $f(x)$ as $x \rightarrow \infty$ exists, then

$$\lim_{x \rightarrow \infty} f(x) = \lim_{p \rightarrow 0} [p\tilde{f}(p)].$$

TABLE 1
Main properties of the Laplace transform

No.	Function	Laplace transform	Operation
1	$af_1(x) + bf_2(x)$	$a\tilde{f}_1(p) + b\tilde{f}_2(p)$	Linearity
2	$f(x/a), a > 0$	$a\tilde{f}(ap)$	Scaling
3	$f(x-a),$ $f(\xi) \equiv 0$ for $\xi < 0$	$e^{-ap}\tilde{f}(p)$	Shift of the argument
4	$x^n f(x); n = 1, 2, \dots$	$(-1)^n \tilde{f}_p^{(n)}(p)$	Differentiation of the transform
5	$\frac{1}{x} f(x)$	$\int_p^\infty \tilde{f}(q) dq$	Integration of the transform
6	$e^{ax} f(x)$	$\tilde{f}(p-a)$	Shift in the complex plane
7	$f'_x(x)$	$p\tilde{f}(p) - f(+0)$	Differentiation
8	$f_x^{(n)}(x)$	$p^n \tilde{f}(p) - \sum_{k=1}^n p^{n-k} f_x^{(k-1)}(+0)$	Differentiation
9	$x^m f_x^{(n)}(x), m = 1, 2, \dots$	$(-1)^m \frac{d^m}{dp^m} \left[p^n \tilde{f}(p) - \sum_{k=1}^n p^{n-k} f_x^{(k-1)}(+0) \right]$	Differentiation
10	$\frac{d^n}{dx^n} [x^m f(x)], m \geq n$	$(-1)^m p^n \frac{d^m}{dp^m} \tilde{f}(p)$	Differentiation
11	$\int_0^x f(t) dt$	$\frac{\tilde{f}(p)}{p}$	Integration
12	$\int_0^x f_1(t)f_2(x-t) dt$	$\tilde{f}_1(p)\tilde{f}_2(p)$	Convolution

TABLE 2
The Laplace transforms of some functions

No.	Function, $f(x)$	Laplace transform, $\tilde{f}(p)$	Remarks
1	1	$1/p$	
2	x^n	$\frac{n!}{p^{n+1}}$	$n = 1, 2, \dots$
3	x^a	$\Gamma(a+1)p^{-a-1}$	$a > -1$
4	e^{-ax}	$(p+a)^{-1}$	
5	$x^a e^{-bx}$	$\Gamma(a+1)(p+b)^{-a-1}$	$a > -1$
6	$\sinh(ax)$	$\frac{a}{p^2 - a^2}$	
7	$\cosh(ax)$	$\frac{p}{p^2 - a^2}$	
8	$\ln x$	$-\frac{1}{p}(\ln p + C)$	$C = 0.5772 \dots$ is the Euler constant
9	$\sin(ax)$	$\frac{a}{p^2 + a^2}$	
10	$\cos(ax)$	$\frac{p}{p^2 + a^2}$	
11	$\operatorname{erfc}\left(\frac{a}{2\sqrt{x}}\right)$	$\frac{1}{p} \exp(-a\sqrt{p})$	$a \geq 0$
12	$J_0(ax)$	$\frac{1}{\sqrt{p^2 + a^2}}$	$J_0(x)$ is the Bessel function

9.2-6. Representation of Inverse Transforms as Convergent Series.

THEOREM 1. Suppose the transform $\tilde{f}(p)$ can be expanded into series in negative powers of p ,

$$\tilde{f}(p) = \sum_{n=1}^{\infty} \frac{a_n}{p^n},$$

convergent for $|p| > R$, where R is an arbitrary positive number; note that the transform tends to zero as $|p| \rightarrow \infty$. Then the inverse transform can be obtained by the formula

$$f(x) = \sum_{n=1}^{\infty} \frac{a_n}{(n-1)!} x^{n-1},$$

where the series on the right-hand side is convergent for all x .

THEOREM 2. Suppose the transform $\tilde{f}(p)$, $|p| > R$, is represented by an absolutely convergent series,

$$\tilde{f}(p) = \sum_{n=0}^{\infty} \frac{a_n}{p^{\lambda_n}}, \quad (7)$$

where $\{\lambda_n\}$ is any positive increasing sequence, $0 < \lambda_0 < \lambda_1 < \dots \rightarrow \infty$. Then it is possible to proceed termwise from series (7) to the following inverse transform series:

$$f(x) = \sum_{n=0}^{\infty} \frac{a_n}{\Gamma(\lambda_n)} x^{\lambda_n-1}, \quad (8)$$

where $\Gamma(\lambda)$ is the Gamma function. Series (8) is convergent for all real and complex values of x other than zero (if $\lambda_0 \geq 1$, the series is convergent for all x).

9.2-7. Representation of Inverse Transforms as Asymptotic Expansions as $x \rightarrow \infty$.

1°. Let $p = p_0$ be a singular point of the Laplace transform $\tilde{f}(p)$ with the greatest real part (it is assumed there is only one such point). If $\tilde{f}(p)$ can be expanded near $p = p_0$ into an absolutely convergent series,

$$\tilde{f}(p) = \sum_{n=0}^{\infty} c_n (p - p_0)^{\lambda_n} \quad (\lambda_0 < \lambda_1 < \dots \rightarrow \infty) \quad (9)$$

with arbitrary λ_n , then the inverse transform $f(x)$ can be expressed in the form of the asymptotic expansion

$$f(x) \sim e^{p_0 x} \sum_{n=0}^{\infty} \frac{c_n}{\Gamma(-\lambda_n)} x^{-\lambda_n-1} \quad \text{as } x \rightarrow \infty.$$

The terms corresponding to nonnegative integer λ_n must be omitted from the summation, since $\Gamma(0) = \Gamma(-1) = \Gamma(-2) = \dots = \infty$.

2°. If the transform $\tilde{f}(p)$ has several singular points, p_1, \dots, p_m , with the same greatest real part, $\operatorname{Re} p_1 = \dots = \operatorname{Re} p_m$, then expansions of the form (9) should be obtained for each of these points and the resulting expressions must be added together.

9.2-8. Post–Widder Formula.

In applications, one can find $f(x)$ if the Laplace transform $\tilde{f}(t)$ on the real semiaxis is known for $t = p \geq 0$. To this end, one uses the Post–Widder formula

$$f(x) = \lim_{n \rightarrow \infty} \left[\frac{(-1)^n}{n!} \left(\frac{n}{x}\right)^{n+1} \tilde{f}_t^{(n)}\left(\frac{n}{x}\right) \right]. \quad (10)$$

Approximate inversion formulas are obtained by taking sufficiently large positive integer n in (10) instead of passing to the limit.

⊙ References for Section 9.2: G. Doetsch (1950, 1956, 1958, 1974), H. Bateman and A. Erdélyi (1954), I. I. Hirschman and D. V. Widder (1955), V. A. Ditkin and A. P. Prudnikov (1965), J. W. Miles (1971), F. Oberhettinger (1973), B. Davis (1978), W. R. LePage (1980), R. Bellman and R. Roth (1984), Yu. A. Brychkov and A. P. Prudnikov (1989), W. H. Beyer (1991), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, Vols 4 and 5), R. J. Beerends, H. G. ter Morschem, and J. C. van den Berg (2003).

9.3. Mellin Transform

9.3-1. Definition. Inversion Formula.

Suppose that a function $f(x)$ is defined for positive x and satisfies the conditions

$$\int_0^1 |f(x)|x^{\sigma_1-1} dx < \infty, \quad \int_1^\infty |f(x)|x^{\sigma_2-1} dx < \infty$$

for some real numbers σ_1 and σ_2 , $\sigma_1 < \sigma_2$.

The *Mellin transform* of $f(x)$ is defined by

$$\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx, \quad (1)$$

where $s = \sigma + i\tau$ is a complex variable ($\sigma_1 < \sigma < \sigma_2$).

For brevity, we rewrite formula (1) as follows:

$$\hat{f}(s) = \mathfrak{M}\{f(x)\}, \quad \text{or} \quad \hat{f}(s) = \mathfrak{M}\{f(x), s\}.$$

Given $\hat{f}(s)$, the function can be found by means of the *inverse Mellin transform*

$$f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds, \quad (\sigma_1 < \sigma < \sigma_2) \quad (2)$$

where the integration path is parallel to the imaginary axis of the complex plane s and the integral is understood in the sense of the Cauchy principal value.

Formula (2) holds for continuous functions. If $f(x)$ has a (finite) jump discontinuity at a point $x = x_0 > 0$, then the left-hand side of (2) is equal to $\frac{1}{2} [f(x_0-0) + f(x_0+0)]$ at this point (for $x_0 = 0$, the first term in the square brackets must be omitted).

For brevity, we rewrite formula (2) in the form

$$f(x) = \mathfrak{M}^{-1}\{\hat{f}(s)\}, \quad \text{or} \quad f(x) = \mathfrak{M}^{-1}\{\hat{f}(s), x\}.$$

9.3-2. Main Properties of the Mellin Transform.

1°. The integral relations

$$\int_0^{\infty} f(xt)g(t) dt = \mathfrak{M}^{-1}\{\hat{f}(s)\hat{g}(1-s)\}, \quad (3)$$

$$\int_0^{\infty} f\left(\frac{x}{t}\right)g(t)\frac{dt}{t} = \mathfrak{M}^{-1}\{\hat{f}(s)\hat{g}(s)\} \quad (4)$$

hold for fairly general assumptions about the integrability of the functions involved (see Ditkin and Prudnikov, 1965).

2°. The main properties of the correspondence between the functions and their Mellin transforms are gathered in Table 3.

TABLE 3
Main properties of the Mellin transform

No	Function	Mellin Transform	Operation
1	$af_1(x) + bf_2(x)$	$a\hat{f}_1(s) + b\hat{f}_2(s)$	Linearity
2	$f(ax), a > 0$	$a^{-s}\hat{f}(s)$	Scaling
3	$x^a f(x)$	$\hat{f}(s+a)$	Shift of the argument of the transform
4	$f(x^2)$	$\frac{1}{2}\hat{f}\left(\frac{1}{2}s\right)$	Squared argument
5	$f(1/x)$	$\hat{f}(-s)$	Inversion of the argument of the transform
6	$x^\lambda f(ax^\beta), a > 0, \beta \neq 0$	$\frac{1}{\beta}a^{-\frac{s+\lambda}{\beta}}\hat{f}\left(\frac{s+\lambda}{\beta}\right)$	Power law transform
7	$f'_x(x)$	$-(s-1)\hat{f}(s-1)$	Differentiation
8	$xf'_x(x)$	$-s\hat{f}(s)$	Differentiation
9	$f_x^{(n)}(x)$	$(-1)^n \frac{\Gamma(s)}{\Gamma(s-n)}\hat{f}(s-n)$	Multiple differentiation
10	$\left(x\frac{d}{dx}\right)^n f(x)$	$(-1)^n s^n \hat{f}(s)$	Multiple differentiation
11	$x^\alpha \int_0^\infty t^\beta f_1(xt)f_2(t) dt$	$\hat{f}_1(s+\alpha)\hat{f}_2(1-s-\alpha+\beta)$	Complicated integration
12	$x^\alpha \int_0^\infty t^\beta f_1\left(\frac{x}{t}\right)f_2(t) dt$	$\hat{f}_1(s+\alpha)\hat{f}_2(s+\alpha+\beta+1)$	Complicated integration

9.3-3. Relation Among the Mellin, Laplace, and Fourier Transforms.

There are tables of direct and inverse Mellin transforms (see Supplements 9 and 10), which are useful in solving specific integral and differential equations. The Mellin transform is related to the Laplace and Fourier transforms as follows:

$$\mathfrak{M}\{f(x), s\} = \mathfrak{L}\{f(e^x), -s\} + \mathfrak{L}\{f(e^{-x}), s\} = \mathfrak{F}\{f(e^x), is\},$$

which makes it possible to apply much more common tables of direct and inverse Laplace and Fourier transforms.

⊙ References for Section 9.3: V. A. Ditkin and A. P. Prudnikov (1965), F. Oberhettinger (1974), Yu. A. Brychkov and A. P. Prudnikov (1989).

9.4. Fourier Transform

9.4-1. Definition. Inversion Formula.

The *Fourier transform* is defined as follows:

$$\tilde{f}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x)e^{-iux} dx. \quad (1)$$

For brevity, we rewrite formula (1) as follows:

$$\tilde{f}(u) = \mathfrak{F}\{f(x)\}, \quad \text{or} \quad \tilde{f}(u) = \mathfrak{F}\{f(x), u\}.$$

Given $\tilde{f}(u)$, the function $f(x)$ can be found by means of the *inverse Fourier transform*

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \tilde{f}(u)e^{iux} du. \quad (2)$$

Formula (2) holds for continuous functions. If $f(x)$ has a (finite) jump discontinuity at a point $x = x_0$, then the left-hand side of (2) is equal to $\frac{1}{2}[f(x_0 - 0) + f(x_0 + 0)]$ at this point.

For brevity, we rewrite formula (2) as follows:

$$f(x) = \mathfrak{F}^{-1}\{\tilde{f}(u)\}, \quad \text{or} \quad f(x) = \mathfrak{F}^{-1}\{\tilde{f}(u), x\}.$$

9.4-2. Asymmetric Form of the Transform.

Sometimes it is more convenient to define the Fourier transform by

$$\check{f}(u) = \int_{-\infty}^{\infty} f(x)e^{-iux} dx. \quad (3)$$

For brevity, we rewrite formula (3) as follows: $\check{f}(u) = \mathcal{F}\{f(x)\}$ or $\check{f}(u) = \mathcal{F}\{f(x), u\}$.

In this case, the *Fourier inversion formula* reads

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \check{f}(u)e^{iux} du, \quad (4)$$

and we use the following symbolic notation for relation (4): $f(x) = \mathcal{F}^{-1}\{\check{f}(u)\}$, or $f(x) = \mathcal{F}^{-1}\{\check{f}(u), x\}$.

9.4-3. Alternative Fourier Transform.

Sometimes, for instance, in the theory of boundary value problems, the alternative Fourier transform is used (and called merely the *Fourier transform*) in the form

$$\mathcal{F}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x)e^{iux} dx. \quad (5)$$

For brevity, we rewrite formula (5) as follows:

$$\mathcal{F}(u) = \mathbf{F}\{f(x)\}, \quad \text{or} \quad \mathcal{F}(u) = \mathbf{F}\{f(x), u\}.$$

For given $\mathcal{F}(u)$, the function $f(x)$ can be found by means of the inverse transform

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{F}(u)e^{-iux} du. \quad (6)$$

TABLE 4
Main properties of the Fourier transform

No.	Function	Fourier transform	Operation
1	$af_1(x) + bf_2(x)$	$a\tilde{f}_1(u) + b\tilde{f}_2(u)$	Linearity
2	$f(x/a), a > 0$	$a\tilde{f}(au)$	Scaling
3	$x^n f(x); n = 1, 2, \dots$	$i^n \tilde{f}_u^{(n)}(u)$	Differentiation of the transform
4	$f''_{xx}(x)$	$-u^2 \tilde{f}(u)$	Differentiation
5	$f_x^{(n)}(x)$	$(iu)^n \tilde{f}(u)$	Differentiation
6	$\int_{-\infty}^{\infty} f_1(\xi)f_2(x-\xi) d\xi$	$\tilde{f}_1(u)\tilde{f}_2(u)$	Convolution

For brevity, we rewrite formula (6) as follows:

$$f(x) = \mathbf{F}^{-1}\{\mathcal{F}(u)\}, \quad \text{or} \quad f(x) = \mathbf{F}^{-1}\{\mathcal{F}(u), x\}.$$

The function $\mathcal{F}(u)$ is also called the *Fourier integral* of $f(x)$.

We can introduce an asymmetric form for the alternative Fourier transform similarly to that of the Fourier transform:

$$\check{\mathcal{F}}(u) = \int_{-\infty}^{\infty} f(x)e^{iux} dx, \quad f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \check{\mathcal{F}}(u)e^{-iux} du, \quad (7)$$

where the direct and the inverse transforms (7) are briefly denoted by $\check{\mathcal{F}}(u) = \check{\mathbf{F}}\{f(x)\}$ and $f(x) = \check{\mathbf{F}}^{-1}\{\check{\mathcal{F}}(u)\}$, or by $\check{\mathcal{F}}(u) = \check{\mathbf{F}}\{f(x), u\}$ and $f(x) = \check{\mathbf{F}}^{-1}\{\check{\mathcal{F}}(u), x\}$.

9.4-4. Convolution Theorem. Main Properties of the Fourier Transforms.

1°. The *convolution* of two functions $f(x)$ and $g(x)$ is defined as

$$f(x) * g(x) \equiv \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x-t)g(t) dt.$$

By performing substitution $x - t = u$, we see that the convolution is symmetric with respect to the convolved functions: $f(x) * g(x) = g(x) * f(x)$.

The *convolution theorem* states that

$$\mathfrak{F}\{f(x) * g(x)\} = \mathfrak{F}\{f(x)\} \mathfrak{F}\{g(x)\}. \quad (8)$$

For the alternative Fourier transform, the convolution theorem reads

$$\mathbf{F}\{f(x) * g(x)\} = \mathbf{F}\{f(x)\} \mathbf{F}\{g(x)\}. \quad (9)$$

Formulas (8) and (9) will be used in Chapters 12 and 13 for solving linear integral equations with difference kernel.

2°. The main properties of the correspondence between functions and their Fourier transforms are gathered in Table 4.

© References for Section 9.4: V. A. Ditkin and A. P. Prudnikov (1965), J. W. Miles (1971), B. Davis (1978), F. Oberhettinger (1980), Yu. A. Brychkov and A. P. Prudnikov (1989), W. H. Beyer (1991), I. Sneddon (1995), A. Pinkus and S. Zafrany (1997), R. Bracewell (1999), A. D. Poularikas (2000), R. J. Beerends, H. G. ter Morschem, J. C. van den Berg (2003), L. Debnath and D. Bhatta (2007).

9.5. Fourier Cosine and Sine Transforms

9.5-1. Fourier Cosine Transform.

1°. Let a function $f(x)$ be integrable on the semiaxis $0 \leq x < \infty$. The *Fourier cosine transform* is defined by

$$\tilde{f}_c(u) = \sqrt{\frac{2}{\pi}} \int_0^\infty f(x) \cos(xu) dx, \quad 0 < u < \infty. \quad (1)$$

For given $\tilde{f}_c(u)$, the function can be found by means of the Fourier cosine inversion formula

$$f(x) = \sqrt{\frac{2}{\pi}} \int_0^\infty \tilde{f}_c(u) \cos(xu) du, \quad 0 < x < \infty. \quad (2)$$

The Fourier cosine transform (1) is denoted for brevity by $\tilde{f}_c(u) = \mathfrak{F}_c\{f(x)\}$.

2°. It follows from formula (2) that the Fourier cosine transform has the property $\mathfrak{F}_c^2 = 1$.

Some other properties of the Fourier cosine transform:

$$\begin{aligned} \mathfrak{F}_c\{x^{2n} f(x)\} &= (-1)^n \frac{d^{2n}}{du^{2n}} \mathfrak{F}_c\{f(x)\}, \quad n = 1, 2, \dots; \\ \mathfrak{F}_c\{f''(x)\} &= -u^2 \mathfrak{F}_c\{f(x)\}. \end{aligned}$$

Here $f(x)$ is assumed to vanish sufficiently rapidly (exponentially) as $x \rightarrow \infty$. For the second formula, the condition $f'(0) = 0$ is assumed to hold.

Parseval's relation for the Fourier cosine transform:

$$\int_0^\infty \mathfrak{F}_c\{f(x)\} \mathfrak{F}_c\{g(x)\} du = \int_0^\infty f(x)g(x) dx.$$

There are tables of the Fourier cosine transform (see Supplement 7 and the references listed at the end of the current section) which prove useful in the solution of specific integral equations.

3°. Sometimes the asymmetric form of the Fourier cosine transform is applied, which is given by the pair of formulas

$$\check{f}_c(u) = \int_0^\infty f(x) \cos(xu) dx, \quad f(x) = \frac{2}{\pi} \int_0^\infty \check{f}_c(u) \cos(xu) du. \quad (3)$$

The direct and inverse Fourier cosine transforms (3) are denoted by $\check{f}_c(u) = \mathcal{F}_c\{f(x)\}$ and $f(x) = \mathcal{F}_c^{-1}\{\check{f}_c(u)\}$, respectively.

9.5-2. Fourier Sine Transform.

1°. Let a function $f(x)$ be integrable on the semiaxis $0 \leq x < \infty$. The *Fourier sine transform* is defined by

$$\tilde{f}_s(u) = \sqrt{\frac{2}{\pi}} \int_0^\infty f(x) \sin(xu) dx, \quad 0 < u < \infty. \quad (4)$$

For given $\tilde{f}_s(u)$, the function $f(x)$ can be found by means of the inverse Fourier sine transform

$$f(x) = \sqrt{\frac{2}{\pi}} \int_0^\infty \tilde{f}_s(u) \sin(xu) du, \quad 0 < x < \infty. \quad (5)$$

The Fourier sine transform (4) is briefly denoted by $\tilde{f}_s(u) = \mathfrak{F}_s\{f(x)\}$.

2°. It follows from formula (5) that the Fourier sine transform has the property $\mathfrak{F}_s^2 = 1$.

Some other properties of the Fourier sine transform:

$$\begin{aligned}\mathfrak{F}_s\{x^{2n}f(x)\} &= (-1)^n \frac{d^{2n}}{du^{2n}} \mathfrak{F}_s\{f(x)\}, \quad n = 1, 2, \dots; \\ \mathfrak{F}_s\{f''(x)\} &= -u^2 \mathfrak{F}_s\{f(x)\}.\end{aligned}$$

Here $f(x)$ is assumed to vanish sufficiently rapidly (exponentially) as $x \rightarrow \infty$. For the second formula, the condition $f(0) = 0$ is assumed to hold.

Parseval's relation for the Fourier sine transform:

$$\int_0^\infty \mathfrak{F}_s\{f(x)\} \mathfrak{F}_s\{g(x)\} du = \int_0^\infty f(x)g(x) dx.$$

There are tables of the Fourier sine transform (see Supplement 8 and the references listed at the end of the current section), which are useful in solving specific integral equations.

3°. Sometimes it is more convenient to apply the asymmetric form of the Fourier sine transform defined by the following two formulas:

$$\check{f}_s(u) = \int_0^\infty f(x) \sin(xu) dx, \quad f(x) = \frac{2}{\pi} \int_0^\infty \check{f}_s(u) \sin(xu) du. \quad (6)$$

The direct and inverse Fourier sine transforms (6) are denoted by $\check{f}_s(u) = \mathcal{F}_s\{f(x)\}$ and $f(x) = \mathcal{F}_s^{-1}\{\check{f}_s(u)\}$, respectively.

⊙ References for Section 9.5: E. A. C. Paley and N. Wiener (1934), S. Bochner and K. C. Chandrasekharan (1949), G. N. Watson (1952), H. Bateman and A. Erdélyi (Vol. 1, 1954), S. Bochner (1959), V. A. Ditkin and A. P. Prudnikov (1965), J. W. Miles (1971), B. Davis (1978), F. Oberhettinger (1980), E. C. Titchmarsh (1986), Ya. A. Brychkov and A. P. Prudnikov (1989), W. H. Beyer (1991), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, p. 440), I. Sneddon (1995), A. D. Poularikas (2000).

9.6. Other Integral Transforms

9.6-1. Hankel Transform.

The *Hankel transform* is defined as follows:

$$\check{f}_\nu(u) = \int_0^\infty x J_\nu(ux) f(x) dx, \quad 0 < u < \infty, \quad (1)$$

where $\nu > -1$ and $J_\nu(x)$ is the Bessel function of the first kind of order ν (see Supplement 11.6).

For given $\check{f}_\nu(u)$, the function $f(x)$ can be found by means of the Hankel inversion formula

$$f(x) = \int_0^\infty u J_\nu(ux) \check{f}_\nu(u) du, \quad 0 < x < \infty. \quad (2)$$

Note that if $f(x) = O(x^\alpha)$ as $x \rightarrow 0$, where $\alpha + \nu + 2 > 0$, and $f(x) = O(x^\beta)$ as $x \rightarrow \infty$, where $\beta + \frac{3}{2} < 0$, then the integral (1) is convergent.

The inversion formula (2) holds for continuous functions. If $f(x)$ has a (finite) jump discontinuity at a point $x = x_0$, then the left-hand side of (2) is equal to $\frac{1}{2}[f(x_0 - 0) + f(x_0 + 0)]$ at this point.

For brevity, we denote the Hankel transform (1) by $\check{f}_\nu(u) = \mathcal{H}_\nu\{f(x)\}$. It follows from formula (2) that the Hankel transform has the property $\mathcal{H}_\nu^2 = 1$.

Parseval's relation for the Hankel transform:

$$\int_0^\infty u \mathcal{H}_\nu\{f(x)\} \mathcal{H}_\nu\{g(x)\} du = \int_0^\infty x f(x)g(x) dx, \quad \nu > -\frac{1}{2}.$$

9.6-2. Meijer Transform.

The Meijer transform is defined as follows:

$$\hat{f}_\mu(s) = \sqrt{\frac{2}{\pi}} \int_0^\infty \sqrt{sx} K_\mu(sx) f(x) dx, \quad 0 < s < \infty,$$

where $K_\mu(x)$ is the modified Bessel function of the second kind (the Macdonald function) of order μ (see Supplement 11.7).

For given $\hat{f}_\mu(s)$, the function $f(x)$ can be found by means of the *Meijer inversion formula*

$$f(x) = \frac{1}{i\sqrt{2\pi}} \int_{c-i\infty}^{c+i\infty} \sqrt{sx} I_\mu(sx) \hat{f}_\mu(s) ds, \quad 0 < x < \infty,$$

where $I_\mu(x)$ is the modified Bessel function of the first kind of order μ (see Supplement 11.7). For the Meijer transform, a convolution is defined and an operational calculus is developed.

9.6-3. Kontorovich–Lebedev Transform.

The Kontorovich–Lebedev transform is introduced as follows:

$$F(\tau) = \int_0^\infty K_{i\tau}(x) f(x) dx, \quad 0 < \tau < \infty,$$

where $K_\mu(x)$ is the modified Bessel function of the second kind (the Macdonald function) of order μ (see Supplement 11.7) and $i = \sqrt{-1}$.

For given $F(\tau)$, the function can be found by means of the *Kontorovich–Lebedev inversion formula*

$$f(x) = \frac{2}{\pi^2 x} \int_0^\infty \tau \sinh(\pi\tau) K_{i\tau}(x) F(\tau) d\tau, \quad 0 < x < \infty.$$

Parseval's relation for the Kontorovich–Lebedev transform:

$$\int_0^\infty F_1(\tau) F_2(\tau) d\tau = \int_0^\infty f_1(x) f_2(x) dx.$$

9.6-4. Y-transform.

The *Y-transform* is defined by

$$F_\nu(u) = \int_0^\infty \sqrt{ux} Y_\nu(ux) f(x) dx,$$

where $Y_\nu(x)$ is the Bessel function of the second kind of order ν .

Given a transform $F_\nu(u)$, the inverse *Y-transform* $f(x)$ is found by the inversion formula

$$f(x) = \int_0^\infty \sqrt{ux} \mathbf{H}_\nu(ux) F_\nu(u) du,$$

where $\mathbf{H}_\nu(x)$ is the Struve function, which is defined as

$$\mathbf{H}_\nu(x) = \sum_{j=0}^{\infty} \frac{(-1)^j (x/2)^{\nu+2j+1}}{\Gamma(j + \frac{3}{2}) \Gamma(\nu + j + \frac{3}{2})}.$$

9.6-5. Summary Table of Integral Transforms.

Table 5 summarizes the integral transforms considered above and also lists some other integral transforms; for the constraints imposed on the functions and parameters occurring in the integrand, see the references given at the end of this section.

TABLE 5
Main integral transforms

Integral transform	Definition	Inversion formula
Laplace transform	$\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$	$f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
Laplace-Carlson transform	$\tilde{f}(p) = p \int_0^{\infty} e^{-px} f(x) dx$	$f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \frac{\tilde{f}(p)}{p} dp$
Two-sided Laplace transform	$\tilde{f}_*(p) = \int_{-\infty}^{\infty} e^{-px} f(x) dx$	$f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}_*(p) dp$
Fourier transform	$\tilde{f}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-iux} f(x) dx$	$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{iux} \tilde{f}(u) du$
Fourier sine transform	$\tilde{f}_s(u) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \sin(xu) f(x) dx$	$f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \sin(xu) \tilde{f}_s(u) du$
Fourier cosine transform	$\tilde{f}_c(u) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \cos(xu) f(x) dx$	$f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \cos(xu) \tilde{f}_c(u) du$
Hartley transform	$\tilde{f}_h(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (\cos xu + \sin xu) f(x) dx$	$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (\cos xu + \sin xu) \tilde{f}_h(u) du$
Mellin transform	$\hat{f}(s) = \int_0^{\infty} x^{s-1} f(x) dx$	$f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} x^{-s} \hat{f}(s) ds$
Hankel transform	$\hat{f}_\nu(w) = \int_0^{\infty} x J_\nu(xw) f(x) dx$	$f(x) = \int_0^{\infty} w J_\nu(xw) \hat{f}_\nu(w) dw$
Y-transform	$F_\nu(u) = \int_0^{\infty} \sqrt{ux} Y_\nu(ux) f(x) dx$	$f(x) = \int_0^{\infty} \sqrt{ux} \mathbf{H}_\nu(ux) F_\nu(u) du$
Meijer transform (K -transform)	$\hat{f}(s) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \sqrt{sx} K_\nu(sx) f(x) dx$	$f(x) = \frac{1}{i\sqrt{2\pi}} \int_{c-i\infty}^{c+i\infty} \sqrt{sx} I_\nu(sx) \hat{f}(s) ds$
Bochner transform	$\tilde{f}(r) = \int_0^{\infty} J_{n/2-1}(2\pi xr) G(x, r) f(x) dx,$ $G(x, r) = 2\pi r(x/r)^{n/2}, \quad n = 1, 2, \dots$	$f(x) = \int_0^{\infty} J_{n/2-1}(2\pi rx) G(r, x) \tilde{f}(r) dr$
Weber transform	$F_a(u) = \int_0^{\infty} W_\nu(xu, au) x f(x) dx,$ $W_\nu(\beta, \mu) \equiv J_\nu(\beta) Y_\nu(\mu) - J_\nu(\mu) Y_\nu(\beta)$	$f(x) = \int_0^{\infty} \frac{W_\nu(xu, au)}{J_\nu^2(au) + Y_\nu^2(au)} u F_a(u) du$
Hardy transform	$F(u) = \int_0^{\infty} C_\nu(xu) x f(x) dx,$ $C_\nu(z) \equiv \cos(\pi p) J_\nu(z) + \sin(\pi p) Y_\nu(z)$	$f(x) = \int_0^{\infty} \Phi(xu) u F(u) du,$ $\Phi(z) = \sum_{n=0}^{\infty} \frac{(-1)^n (z/2)^{\nu+2p+2n}}{\Gamma(p+n+1) \Gamma(\nu+p+n+1)}$
Kontorovich-Lebedev transform	$F(\tau) = \int_0^{\infty} K_{i\tau}(x) f(x) dx$	$f(x) = \frac{2}{\pi^2 x} \int_0^{\infty} \tau \sinh(\pi\tau) K_{i\tau}(x) F(\tau) d\tau$

TABLE 5 (continued)
Main integral transforms

Integral transform	Definition	Inversion formula
Mehler–Fock transform	$F(x) = \int_0^\infty P_{-\frac{1}{2}+i\tau}(x)f(\tau) d\tau, \quad 1 \leq x < \infty$	$f(\tau) = \tau \tanh(\pi\tau) \int_1^\infty P_{-\frac{1}{2}+i\tau}(x)F(x) dx$
Euler transform of the 1st kind	$F(x) = \frac{1}{\Gamma(\mu)} \int_a^x \frac{f(t) dt}{(x-t)^{1-\mu}}, \quad 0 < \mu < 1, \quad x > a$	$f(x) = \frac{1}{\Gamma(1-\mu)} \frac{d}{dx} \int_a^x \frac{F(t) dt}{(x-t)^\mu}$
Euler transform of the 2nd kind	$F(x) = \frac{1}{\Gamma(\mu)} \int_x^a \frac{f(t) dt}{(t-x)^{1-\mu}}, \quad 0 < \mu < 1, \quad x < a$	$f(x) = -\frac{1}{\Gamma(1-\mu)} \frac{d}{dx} \int_x^a \frac{F(t) dt}{(t-x)^\mu}$
Gauss transform	$F(x) = \frac{1}{\sqrt{\pi a}} \int_{-\infty}^\infty \exp\left[-\frac{(x-t)^2}{a}\right] f(t) dt$	$f(x) = \exp\left(-\frac{a}{4} \frac{d^2}{dx^2}\right) F(x)$
Hilbert transform	$\widehat{F}(s) = \frac{1}{\pi} \int_{-\infty}^\infty \frac{f(x)}{x-s} dx$	$f(x) = -\frac{1}{\pi} \int_{-\infty}^\infty \frac{\widehat{F}(s)}{s-x} ds$

Notations for Table 5: $i = \sqrt{-1}$, $J_\mu(x)$ and $Y_\mu(x)$ are the Bessel functions of the first and the second kind, respectively; $I_\mu(x)$ and $K_\mu(x)$ are the modified Bessel functions of the first and the second kind, respectively; $P_\mu(x)$ is the Legendre spherical function of the first kind; and $\mathbf{H}_\mu(x)$ is the Struve function (see Subsection 9.6-4).

Remark 1. The Euler transform of the first kind is also known as Riemann–Liouville integral (the left fractional integral of order μ or, for short, the fractional integral), see Section 10.5. The Euler transform of the second kind is also called the right fractional integral of order μ .

Remark 2. If $a = 4$, the Gauss transform is called the Weierstrass transform. In the inversion formula, the exponential is represented by an operator series: $\exp\left(k \frac{d^2}{dx^2}\right) \equiv 1 + \sum_{n=1}^\infty \frac{k^n}{n!} \frac{d^{2n}}{dx^{2n}}$.

Remark 3. In the direct and inverse Hilbert transforms, the integrals are understood in the sense of the Cauchy principal value.

Remark 4. Some other integral transforms are described in Chapter 3.

⊙ References for Section 9.6: H. Bateman and A. Erdélyi (1954, Vols 1 and 2), J. L. Griffith (1958), V. A. Ditkin and A. P. Prudnikov (1965), J. W. Miles (1971), F. Oberhettinger (1972), I. Sneddon (1972), H. M. Srivastava and R. G. Buschman (1977), B. Davis (1978), D. Zwillinger (1989), Yu. A. Brychkov and A. P. Prudnikov (1989), W. H. Beyer (1991), M. Ya. Antimirov (1993), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993), A. D. Polyaniin and A. V. Manzhirev (1998, 2007), A. D. Poularikas (2000), L. Debnath and D. Bhatta (2007).

Chapter 10

Methods for Solving Linear Equations of the Form $\int_a^x K(x, t)y(t) dt = f(x)$

10.1. Volterra Equations of the First Kind

10.1-1. Equations of the First Kind. Function and Kernel Classes.

In this chapter we present methods for solving Volterra linear equations of the first kind. These equations have the form

$$\int_a^x K(x, t)y(t) dt = f(x), \quad (1)$$

where $y(x)$ is the unknown function ($a \leq x \leq b$), $K(x, t)$ is the kernel of the integral equation, and $f(x)$ is a given function, the *right-hand side* of Eq. (1). The functions $y(x)$ and $f(x)$ are usually assumed to be continuous or square integrable on $[a, b]$. The kernel $K(x, t)$ is usually assumed either to be continuous on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ or to satisfy the condition

$$\int_a^b \int_a^b K^2(x, t) dx dt = B^2 < \infty, \quad (2)$$

where B is a constant, that is, to be square integrable on this square. It is assumed in (2) that $K(x, t) \equiv 0$ for $t > x$.

The kernel $K(x, t)$ is said to be *degenerate* if it can be represented in the form

$$K(x, t) = g_1(x)h_1(t) + \cdots + g_n(x)h_n(t).$$

The kernel $K(x, t)$ of an integral equation is called *difference kernel* if it depends only on the difference of the arguments, $K(x, t) = K(x - t)$.

Polar kernels

$$K(x, t) = \frac{L(x, t)}{(x - t)^\beta} + M(x, t), \quad 0 < \beta < 1, \quad (3)$$

and logarithmic kernels (kernels with logarithmic singularity)

$$K(x, t) = L(x, t) \ln(x - t) + M(x, t), \quad (4)$$

where $L(x, t)$ and $M(x, t)$ are continuous on S and $L(x, x) \neq 0$, are often considered as well.

Polar and logarithmic kernels form a class of kernels with weak singularity. Equations containing such kernels are called *equations with weak singularity*.

The following *generalized Abel equation* is a special case of Eq. (1) with the kernel of the form (3):

$$\int_a^x \frac{y(t)}{(x - t)^\beta} dt = f(x), \quad 0 < \beta < 1.$$

In case the functions $K(x, t)$ and $f(x)$ are continuous, the right-hand side of Eq. (1) must satisfy the following conditions:

1°. If $K(a, a) \neq 0$, then $f(x)$ must be constrained by $f(a) = 0$.

2°. If $K(a, a) = K'_x(a, a) = \dots = K_x^{(n-1)}(a, a) = 0$, $0 < |K_x^{(n)}(a, a)| < \infty$, then the right-hand side of the equation must satisfy the conditions

$$f(a) = f'_x(a) = \dots = f_x^{(n)}(a) = 0.$$

3°. If $K(a, a) = K'_x(a, a) = \dots = K_x^{(n-1)}(a, a) = 0$, $K_x^{(n)}(a, a) = \infty$, then the right-hand side of the equation must satisfy the conditions

$$f(a) = f'_x(a) = \dots = f_x^{(n-1)}(a) = 0.$$

For polar kernels of the form (3) or (4) and continuous $f(x)$, no additional conditions are imposed on the right-hand side of the integral equation.

Remark 1. Generally, the case in which the integration limit a is infinite is not excluded.

10.1-2. Existence and Uniqueness of a Solution.

Assume that in Eq. (1) the functions $f(x)$ and $K(x, t)$ are continuous together with their first derivatives on $[a, b]$ and on S , respectively. If $K(x, x) \neq 0$ ($x \in [a, b]$) and $f(a) = 0$, then there exists a unique continuous solution $y(x)$ of Eq. (1).

Remark 2. The problem of existence and uniqueness of a solution to a Volterra equation of the first kind is closely related to conditions under which this equation can be reduced to Volterra equations of the second kind (see Section 10.3).

Remark 3. A Volterra equation of the first kind can be treated as a Fredholm equation of the first kind whose kernel $K(x, t)$ vanishes for $t > x$ (see Chapter 12).

10.1-3. Some Problems Leading to Volterra Integral Equations of the First Kind.

1°. **Abel problem** (generalization of the tautochrone problem*).

Statement of the problem. Suppose a point mass (a bead) can move along a curve in the vertical plane (ξ, η) under the gravitational force. Determine the curve if the bead, initially having an ordinate x and zero velocity, must reach the $O\xi$ axis in a time $t = f_1(x)$, where $f_1(x)$ is a given function.

Derivation of the integral equation. The absolute value of the bead velocity is expressed as

$$v = \sqrt{2g(x - \eta)}.$$

Let $\beta = \beta(\eta)$ denote the angle between the tangent to the curve and the $O\xi$ axis, as shown in Fig. 2. Then the η -component of the velocity is found as

$$\frac{d\eta}{dt} = -\sqrt{2g(x - \eta)} \sin \beta.$$

It follows that

$$dt = -\frac{d\eta}{\sqrt{2g(x - \eta)} \sin \beta}.$$

* Find the curve down which a heavy bead having zero initial velocity and placed anywhere will fall to the bottom in the same amount of time.

Integrating over η from 0 to x and setting $\frac{1}{\sin \beta} = y(\eta)$, one arrives at the Abel equation

$$\int_0^x \frac{y(\eta)}{\sqrt{x-\eta}} d\eta = -\sqrt{2g} f_1(x).$$

Denoting $-\sqrt{2g} f_1(x) = f(x)$ yields

$$\int_0^x \frac{y(\eta)}{\sqrt{x-\eta}} d\eta = f(x).$$

Here $y(x)$ is the unknown function and $f(x)$ is a given function.




Figure 2. Curve along which the bead moves in the Abel problem.

Having found $y(\eta)$, one readily obtains the equation of the desired curve. Indeed, since $y(\eta) = 1/\sin \beta$, we have $\eta = \Phi(\beta)$. Further,

$$d\xi = \frac{d\eta}{\tan \beta} = \frac{\Phi'(\beta) d\beta}{\tan \beta},$$

and therefore

$$\xi = \int \frac{\Phi'(\beta)}{\tan \beta} d\beta = \Psi(\beta).$$

Hence, the desired curve is determined parametrically by the equations

$$\xi = \Psi(\beta), \quad \eta = \Phi(\beta).$$

In particular, if $f(x) = C = \text{const}$, the desired curve is a cycloid.

2°. A model problem on buying and selling goods.

Statement of the problem. There is a shop that buys and sells various types of goods. It is assumed that:

- 1) buying and selling are continuous processes and the goods bought are put on sale immediately;
- 2) any type of goods is purchased by the shop in consignments, the quantity of goods in each consignment equal to the quantity sold by the shop for a time T , the same for all types of goods;
- 3) each new consignment is sold uniformly over the time T .

The shop starts selling a new consignment the cost of which is equal to unity. Find the law $y(t)$ according to which the goods should be bought, in order that the cost of the goods present in the shop remains constant.

Derivation of the integral equation. The cost of the initially bought goods remaining in the shop by an instant t is equal to

$$K(t) = \begin{cases} 1 - \frac{t}{T} & \text{if } t \leq T, \\ 0 & \text{if } t > T. \end{cases}$$

Suppose that on the time interval from τ to $\tau + d\tau$ the cost of the goods bought is equal to $y(\tau) d\tau$. This stock of the goods is decreased through selling, so that by the instant $t > \tau$ the cost of the remainder is $K(t - \tau)y(\tau) d\tau$. Therefore, by the time t , the cost of the unsold portion of the goods purchased by the shop will be equal to

$$\int_0^t K(t - \tau)y(\tau) d\tau.$$

On the other hand, the cost of the unsold portion of the goods bought by the shop is equal to $1 - K(t)$. Equating these two expressions gives

$$1 - K(t) = \int_0^t K(t - \tau)y(\tau) d\tau.$$

This is a convolution integral equation of the first kind for the unknown function $y(t)$.

© References for Section 10.1: E. Goursat (1923), H. M. Müntz (1934), F. G. Tricomi (1957), V. Volterra (1959), S. G. Mikhlín (1960), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), J. A. Cochran (1972), C. Corduneanu (1973), V. I. Smirnov (1974), P. P. Zabreyko, A. I. Koshelev, et al. (1975), A. J. Jerry (1985), A. F. Verlan' and V. S. Sizikov (1986), P. Linz (1987).

10.2. Equations with Degenerate Kernel:

$$K(x, t) = g_1(x)h_1(t) + \dots + g_n(x)h_n(t)$$

10.2-1. Equations with Kernel of the Form $K(x, t) = g_1(x)h_1(t) + g_2(x)h_2(t)$.

Any equation of this type can be rewritten in the form

$$g_1(x) \int_a^x h_1(t)y(t) dt + g_2(x) \int_a^x h_2(t)y(t) dt = f(x). \quad (1)$$

It is assumed that $g_1(x) \neq \text{const } g_2(x)$, $h_1(t) \neq \text{const } h_2(t)$, $0 < g_1^2(a) + g_2^2(a) < \infty$, and $f(a) = 0$.

The change of variables

$$u(x) = \int_a^x h_1(t)y(t) dt \quad (2)$$

followed by the integration by parts in the second integral in (1) with regard to the relation $u(a) = 0$ yields the following Volterra equation of the second kind:

$$[g_1(x)h_1(x) + g_2(x)h_2(x)]u(x) - g_2(x)h_1(x) \int_a^x \left[\frac{h_2(t)}{h_1(t)} \right]'_t u(t) dt = h_1(x)f(x). \quad (3)$$

The substitution

$$w(x) = \int_a^x \left[\frac{h_2(t)}{h_1(t)} \right]'_t u(t) dt \quad (4)$$

reduces Eq. (3) to the first-order linear ordinary differential equation

$$[g_1(x)h_1(x) + g_2(x)h_2(x)]w'_x - g_2(x)h_1(x) \left[\frac{h_2(x)}{h_1(x)} \right]'_x w = f(x)h_1(x) \left[\frac{h_2(x)}{h_1(x)} \right]'_x. \quad (5)$$

1°. In the case $g_1(x)h_1(x) + g_2(x)h_2(x) \neq 0$, the solution of equation (5) satisfying the condition $w(a) = 0$ (this condition is a consequence of the substitution (4)) has the form

$$w(x) = \Phi(x) \int_a^x \left[\frac{h_2(t)}{h_1(t)} \right]'_t \frac{f(t)h_1(t) dt}{\Phi(t)[g_1(t)h_1(t) + g_2(t)h_2(t)]}, \tag{6}$$

$$\Phi(x) = \exp \left\{ \int_a^x \left[\frac{h_2(t)}{h_1(t)} \right]'_t \frac{g_2(t)h_1(t) dt}{g_1(t)h_1(t) + g_2(t)h_2(t)} \right\}. \tag{7}$$

Let us differentiate relation (4) and substitute the function (6) into the resulting expression. After integrating by parts with regard to the relations $f(a) = 0$ and $w(a) = 0$, for $f \neq \text{const } g_2$ we obtain

$$u(x) = \frac{g_2(x)h_1(x)\Phi(x)}{g_1(x)h_1(x) + g_2(x)h_2(x)} \int_a^x \left[\frac{f(t)}{g_2(t)} \right]'_t \frac{dt}{\Phi(t)}.$$

Using formula (2), we find a solution of the original equation in the form

$$y(x) = \frac{1}{h_1(x)} \frac{d}{dx} \left\{ \frac{g_2(x)h_1(x)\Phi(x)}{g_1(x)h_1(x) + g_2(x)h_2(x)} \int_a^x \left[\frac{f(t)}{g_2(t)} \right]'_t \frac{dt}{\Phi(t)} \right\}, \tag{8}$$

where the function $\Phi(x)$ is given by (7).

If $f(x) \equiv \text{const } g_2(x)$, the solution is given by formulas (8) and (7) in which the subscript 1 must be changed by 2 and vice versa.

2°. In the case $g_1(x)h_1(x) + g_2(x)h_2(x) \equiv 0$, the solution has the form

$$y(x) = \frac{1}{h_1} \frac{d}{dx} \left[\frac{(f/g_2)'_x}{(g_1/g_2)'_x} \right] = -\frac{1}{h_1} \frac{d}{dx} \left[\frac{(f/g_2)'_x}{(h_2/h_1)'_x} \right].$$

10.2-2. Equations with General Degenerate Kernel.

A Volterra equation of the first kind with general degenerate kernel has the form

$$\sum_{m=1}^n g_m(x) \int_a^x h_m(t)y(t) dt = f(x). \tag{9}$$

Using the notation

$$w_m(x) = \int_a^x h_m(t)y(t) dt, \quad m = 1, \dots, n, \tag{10}$$

we can rewrite Eq. (9) as follows:

$$\sum_{m=1}^n g_m(x)w_m(x) = f(x). \tag{11}$$

On differentiating formulas (10) and eliminating $y(x)$ from the resulting equations, we arrive at the following linear differential equations for the functions $w_m = w_m(x)$:

$$h_1(x)w'_m = h_m(x)w'_1, \quad m = 2, \dots, n, \tag{12}$$

(the prime stands for the derivative with respect to x) with the initial conditions

$$w_m(a) = 0, \quad m = 1, \dots, n.$$

Any solution of system (11), (12) determines a solution of the original integral equation (9) by each of the expressions

$$y(x) = \frac{w'_m(x)}{h_m(x)}, \quad m = 1, \dots, n,$$

which can be obtained by differentiating formula (10).

System (11), (12) can be reduced to a linear differential equation of order $n - 1$ for any function $w_m(x)$ ($m = 1, \dots, n$) by multiple differentiation of Eq. (11) with regard to (12).

⊙ References for Section 10.2: E. Goursat (1923), A. F. Verlan' and V. S. Sizikov (1986).

10.3. Reduction of Volterra Equations of the First Kind to Volterra Equations of the Second Kind

10.3-1. First Method.

Suppose that the kernel and the right-hand side of the equation

$$\int_a^x K(x, t)y(t) dt = f(x), \quad (1)$$

have continuous derivatives with respect to x and that the condition $K(x, x) \neq 0$ holds. In this case, after differentiating relation (1) and dividing the resulting expression by $K(x, x)$ we arrive at the following Volterra equation of the second kind:

$$y(x) + \int_a^x \frac{K'_x(x, t)}{K(x, x)} y(t) dt = \frac{f'_x(x)}{K(x, x)}. \quad (2)$$

Equations of this type are considered in Chapter 11. If $K(x, x) \equiv 0$, then, on differentiating Eq. (1) with respect to x twice and assuming that $K'_x(x, t)|_{t=x} \neq 0$, we obtain the Volterra equation of the second kind

$$y(x) + \int_a^x \frac{K''_{xx}(x, t)}{K'_x(x, t)|_{t=x}} y(t) dt = \frac{f''_{xx}(x)}{K'_x(x, t)|_{t=x}}.$$

If $K'_x(x, x) \equiv 0$, we can again apply differentiation, and so on. If the first $m - 2$ partial derivatives of the kernel with respect to x are identically zero and the $(m - 1)$ st derivative is nonzero, then the m -fold differentiation of the original equation gives the following Volterra equation of the second kind:

$$y(x) + \int_a^x \frac{K_x^{(m)}(x, t)}{K_x^{(m-1)}(x, t)|_{t=x}} y(t) dt = \frac{f_x^{(m)}(x)}{K_x^{(m-1)}(x, t)|_{t=x}}.$$

10.3-2. Second Method.

Let us introduce the new variable

$$Y(x) = \int_a^x y(t) dt$$

and integrate the right-hand side of Eq. (1) by parts taking into account the relation $f(a) = 0$. After dividing the resulting expression by $K(x, x)$, we arrive at the Volterra equation of the second kind

$$Y(x) - \int_a^x \frac{K'_t(x, t)}{K(x, x)} Y(t) dt = \frac{f(x)}{K(x, x)},$$

for which the condition $K(x, x) \neq 0$ must hold.

⊙ References for Section 10.3: E. Goursat (1923), V. Volterra (1959).

10.4. Equations with Difference Kernel: $K(x, t) = K(x - t)$

10.4-1. Solution Method Based on the Laplace Transform.

Volterra equations of the first kind with kernel depending on the difference of the arguments have the form

$$\int_0^x K(x - t)y(t) dt = f(x). \quad (1)$$

To solve these equations, the Laplace transform can be used (see Section 9.2). In what follows we need the transforms of the kernel and the right-hand side; they are given by the formulas

$$\tilde{K}(p) = \int_0^{\infty} K(x)e^{-px} dx, \quad \tilde{f}(p) = \int_0^{\infty} f(x)e^{-px} dx. \quad (2)$$

Applying the Laplace transform \mathfrak{L} to Eq. (1) and taking into account the fact that an integral with kernel depending on the difference of the arguments is transformed to the product by the rule (see Subsection 9.2-4)

$$\mathfrak{L} \left\{ \int_0^x K(x-t)y(t) dt \right\} = \tilde{K}(p)\tilde{y}(p),$$

we obtain the following equation for the transform $\tilde{y}(p)$:

$$\tilde{K}(p)\tilde{y}(p) = \tilde{f}(p). \quad (3)$$

The solution of Eq. (3) is given by the formula

$$\tilde{y}(p) = \frac{\tilde{f}(p)}{\tilde{K}(p)}. \quad (4)$$

On applying the Laplace inversion formula (if it is applicable) to (4), we obtain a solution of Eq. (1) in the form

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(p)}{\tilde{K}(p)} e^{px} dp. \quad (5)$$

When applying formula (5) in practice, the following two technical problems occur:

1°. Finding the transform $\tilde{K}(p) = \int_0^{\infty} K(x)e^{-px} dx$ for a given kernel $K(x)$.

2°. Finding the resolvent (5) whose transform $\tilde{R}(p)$ is given by formula (4).

To calculate the corresponding integrals, tables of direct and inverse Laplace transforms can be applied (see Supplements 5 and 6), and, in many cases, to find the inverse transform, methods of the theory of functions of a complex variable are applied, including the Cauchy residue theorem (see Subsection 9.1-4).

Remark. If the lower limit in the integral of a Volterra equation with difference kernel is a , then this equation can be reduced to Eq. (1) by means of the change of variables $x = \bar{x} - a$, $t = \bar{t} - a$.

10.4-2. Case in Which the Transform of the Solution is a Rational Function.

Consider the important special case in which the transform (4) of the solution is a rational function of the form

$$\tilde{y}(p) = \frac{\tilde{f}(p)}{\tilde{K}(p)} \equiv \frac{R(p)}{Q(p)},$$

where $Q(p)$ and $R(p)$ are polynomials in the variable p and the degree of $Q(p)$ exceeds that of $R(p)$.

If the zeros of the denominator $Q(p)$ are simple, i.e.,

$$Q(p) \equiv \text{const} (p - \lambda_1)(p - \lambda_2) \dots (p - \lambda_n),$$

and $\lambda_i \neq \lambda_j$ for $i \neq j$, then the solution has the form

$$y(x) = \sum_{k=1}^n \frac{R(\lambda_k)}{Q'(\lambda_k)} \exp(\lambda_k x),$$

where the prime stands for the derivatives.

Example 1. Consider the Volterra integral equation of the first kind

$$\int_0^x e^{-a(x-t)} y(t) dt = A \sinh(bx).$$

We apply the Laplace transform to this equation and obtain (see Supplement 5)

$$\frac{1}{p+a} \tilde{y}(p) = \frac{Ab}{p^2 - b^2}.$$

This implies

$$\tilde{y}(p) = \frac{Ab(p+a)}{p^2 - b^2} = \frac{Ab(p+a)}{(p-b)(p+b)}.$$

We have $Q(p) = (p-b)(p+b)$, $R(p) = Ab(p+a)$, $\lambda_1 = b$, and $\lambda_2 = -b$. Therefore, the solution of the integral equation has the form

$$y(x) = \frac{1}{2} A(b+a)e^{bx} + \frac{1}{2} A(b-a)e^{-bx} = Aa \sinh(bx) + Ab \cosh(bx).$$

10.4-3. Convolution Representation of a Solution.

In solving Volterra integral equations of the first kind with difference kernel $K(x-t)$ by means of the Laplace transform, it is sometimes useful to apply the following approach.

Let us represent the transform (4) of a solution in the form

$$\tilde{y}(p) = \tilde{N}(p)\tilde{M}(p)\tilde{f}(p), \quad \tilde{N}(p) \equiv \frac{1}{\tilde{K}(p)\tilde{M}(p)}. \quad (6)$$

If we can find a function $\tilde{M}(p)$ for which the inverse transforms

$$\mathfrak{L}^{-1}\{\tilde{M}(p)\} = M(x), \quad \mathfrak{L}^{-1}\{\tilde{N}(p)\} = N(x) \quad (7)$$

exist and can be found in a closed form, then the solution can be written as the convolution

$$y(x) = \int_0^x N(x-t)F(t) dt, \quad F(t) = \int_0^t M(t-s)f(s) ds. \quad (8)$$

Example 2. Consider the equation

$$\int_0^x \sin(\lambda\sqrt{x-t}) y(t) dt = f(x), \quad f(0) = 0. \quad (9)$$

Applying the Laplace transform, we obtain (see Supplement 5)

$$\tilde{y}(p) = \frac{2}{\sqrt{\pi}\lambda} p^{3/2} \exp(\alpha/p) \tilde{f}(p), \quad \alpha = \frac{1}{4}\lambda^2. \quad (10)$$

Let us rewrite the right-hand side of (10) in the equivalent form

$$\tilde{y}(p) = \frac{2}{\sqrt{\pi}\lambda} p^2 [p^{-1/2} \exp(\alpha/p)] \tilde{f}(p), \quad \alpha = \frac{1}{4}\lambda^2, \quad (11)$$

where the factor in the square brackets corresponds to $\tilde{M}(p)$ in formula (6) and $\tilde{N}(p) = \text{const } p^2$.

By applying the Laplace inversion formula according to the above scheme to formula (11) with regard to the relation (see Supplement 6)

$$\mathfrak{L}^{-1}\{p^2 \tilde{\varphi}(p)\} = \frac{d^2}{dx^2} \varphi(x), \quad \mathfrak{L}^{-1}\{p^{-1/2} \exp(\alpha/p)\} = \frac{1}{\sqrt{\pi x}} \cosh(\lambda\sqrt{x}),$$

we find the solution

$$y(x) = \frac{2}{\pi\lambda} \frac{d^2}{dx^2} \int_0^x \frac{\cosh(\lambda\sqrt{x-t})}{\sqrt{x-t}} f(t) dt.$$

10.4-4. Application of an Auxiliary Equation.

Consider the equation

$$\int_a^x K(x-t)y(t) dt = f(x), \quad (12)$$

where the kernel $K(x)$ has an integrable singularity at $x = 0$.

Let $w = w(x)$ be the solution of the simpler auxiliary equation with $f(x) \equiv 1$ and $a = 0$,

$$\int_0^x K(x-t)w(t) dt = 1. \quad (13)$$

Then the solution of the original equation (12) with arbitrary right-hand side can be expressed as follows via the solution of the auxiliary equation (13):

$$y(x) = \frac{d}{dx} \int_a^x w(x-t)f(t) dt = f(a)w(x-a) + \int_a^x w(x-t)f'_t(t) dt. \quad (14)$$

Example 3. Consider the generalized Abel equation

$$\int_a^x \frac{y(t) dt}{(x-t)^\mu} = f(x), \quad 0 < \mu < 1. \quad (15)$$

We seek a solution of the corresponding auxiliary equation

$$\int_0^x \frac{w(t) dt}{(x-t)^\mu} = 1, \quad 0 < \mu < 1, \quad (16)$$

by the method of indeterminate coefficients in the form

$$w(x) = Ax^\beta. \quad (17)$$

Let us substitute (17) into (16) and then perform the change of variable $t = x\xi$ in the integral. Taking into account the relationship

$$B(p, q) = \int_0^1 \xi^{p-1}(1-\xi)^{q-1} d\xi = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

between the beta and gamma functions, we obtain

$$A \frac{\Gamma(\beta+1)\Gamma(1-\mu)}{\Gamma(2+\beta-\mu)} x^{\beta+1-\mu} = 1.$$

From this relation we find the coefficients A and β :

$$\beta = \mu - 1, \quad A = \frac{1}{\Gamma(\mu)\Gamma(1-\mu)} = \frac{\sin(\pi\mu)}{\pi}. \quad (18)$$

Formulas (17) and (18) define the solution of the auxiliary equation (16) and make it possible to find the solution of the generalized Abel equation (15) by means of formula (14) as follows:

$$y(x) = \frac{\sin(\pi\mu)}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) dt}{(x-t)^{1-\mu}} = \frac{\sin(\pi\mu)}{\pi} \left[\frac{f(a)}{(x-a)^{1-\mu}} + \int_a^x \frac{f'_t(t) dt}{(x-t)^{1-\mu}} \right]. \quad (19)$$

10.4-5. Reduction to Ordinary Differential Equations.

Consider the special case in which the transform of the kernel of the integral equation (1) can be represented in the form

$$\tilde{K}(p) = \frac{M(p)}{N(p)}, \quad (20)$$

where $M(p)$ and $N(p)$ are some polynomials of degrees m and n , respectively:

$$M(p) = \sum_{k=0}^m A_k p^k, \quad N(p) = \sum_{k=0}^n B_k p^k. \quad (21)$$

In this case, the solution of the integral equation (1) (if it exists) satisfies the following linear nonhomogeneous ordinary differential equation of order m with constant coefficients:

$$\sum_{k=0}^m A_k y_x^{(k)}(x) = \sum_{k=0}^n B_k f_x^{(k)}(x). \quad (22)$$

We can rewrite Eq. (22) in the operator form

$$M(D)y(x) = N(D)f(x), \quad D \equiv \frac{d}{dx}.$$

The initial data for the differential equation (22), as well as the conditions that must be imposed on the right-hand side of the integral equation (1), can be obtained from the relation

$$\sum_{k=0}^m A_k \sum_{s=0}^{k-1} p^{k-1-s} y_x^{(s)}(0) - \sum_{k=0}^n B_k \sum_{s=0}^{k-1} p^{k-1-s} f_x^{(s)}(0) = 0 \quad (23)$$

by matching the coefficients of like powers of the parameter p .

The proof of this assertion can be given by applying the Laplace transform to the differential equation (22) followed by comparing the resulting expression with Eq. (3) with regard to (20).

10.4-6. Reduction of a Volterra Equation to a Wiener–Hopf Equation.

A Volterra equation of the first kind with difference kernel of the form

$$\int_0^x K(x-t)y(t) dt = f(x), \quad 0 < x < \infty, \quad (24)$$

can be reduced to the following Wiener–Hopf equation of the first kind:

$$\int_0^\infty K_+(x-t)y(t) dt = f(x), \quad 0 < x < \infty, \quad (25)$$

where the kernel $K_+(x-t)$ is given by

$$K_+(s) = \begin{cases} K(s) & \text{for } s > 0, \\ 0 & \text{for } s < 0. \end{cases}$$

Methods for solving Eq. (25) are presented in Section 12.8.

© References for Section 10.4: G. Doetsch (1956), V. A. Ditkin and A. P. Prudnikov (1965), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), V. I. Smirnov (1974), P. P. Zabreyko, A. I. Koshelev, et al. (1975), F. D. Gakhov and Yu. I. Cherskii (1978).

10.5. Method of Fractional Differentiation

10.5-1. Definition of Fractional Integrals.

A function $f(x)$ is said to be *absolutely continuous* on a closed interval $[a, b]$ if for each $\varepsilon > 0$ there exists a $\delta > 0$ such that for any finite system of disjoint intervals $[a_k, b_k] \subset [a, b]$, $k = 1, \dots, n$, such that $\sum_{k=1}^n (b_k - a_k) < \delta$ the inequality $\sum_{k=1}^n |f(b_k) - f(a_k)| < \varepsilon$ holds. The class of all these functions is denoted by AC .

Let AC^n , $n = 1, 2, \dots$, be the class of functions $f(x)$ that are continuously differentiable on $[a, b]$ up to the order $n - 1$ and for which $f^{(n-1)}(x) \in AC$.

Let $\varphi(x) \in L_1(a, b)$. The integrals

$$\mathbf{I}_{a+}^{\mu} \varphi(x) \equiv \frac{1}{\Gamma(\mu)} \int_a^x \frac{\varphi(t)}{(x-t)^{1-\mu}} dt, \quad x > a, \quad (1)$$

$$\mathbf{I}_{b-}^{\mu} \varphi(x) \equiv \frac{1}{\Gamma(\mu)} \int_x^b \frac{\varphi(t)}{(t-x)^{1-\mu}} dt, \quad x < b, \quad (2)$$

where $\mu > 0$, are called the *integrals of fractional order μ* . Sometimes the integral (1) is called *left-sided* and the integral (2) is called *right-sided*. The operators \mathbf{I}_{a+}^{μ} and \mathbf{I}_{b-}^{μ} are called the *operators of fractional integration*.

The integrals (1) and (2) are usually called the *Riemann–Liouville fractional integrals*.

The following formula holds:

$$\int_a^b \varphi(x) \mathbf{I}_{a+}^{\mu} \psi(x) dx = \int_a^b \psi(x) \mathbf{I}_{b-}^{\mu} \varphi(x) dx, \quad (3)$$

which is sometimes called the *formula of fractional integration by parts*.

Fractional integration has the property

$$\mathbf{I}_{a+}^{\mu} \mathbf{I}_{a+}^{\beta} \varphi(x) = \mathbf{I}_{a+}^{\mu+\beta} \varphi(x), \quad \mathbf{I}_{b-}^{\mu} \mathbf{I}_{b-}^{\beta} \varphi(x) = \mathbf{I}_{b-}^{\mu+\beta} \varphi(x), \quad \mu > 0, \quad \beta > 0. \quad (4)$$

Property (4) is called the *semigroup property of fractional integration*.

10.5-2. Definition of Fractional Derivatives.

It is natural to introduce fractional differentiation as the operation inverse to fractional integration. For a function $f(x)$ defined on a closed interval $[a, b]$, the expressions

$$\mathbf{D}_{a+}^{\mu} f(x) = \frac{1}{\Gamma(1-\mu)} \frac{d}{dx} \int_a^x \frac{f(t)}{(x-t)^{\mu}} dt, \quad (5)$$

$$\mathbf{D}_{b-}^{\mu} f(x) = -\frac{1}{\Gamma(1-\mu)} \frac{d}{dx} \int_x^b \frac{f(t)}{(t-x)^{\mu}} dt \quad (6)$$

are called the *left* and the *right fractional derivative of order μ* , respectively. It is assumed here that $0 < \mu < 1$.

The fractional derivatives (5) and (6) are usually called the *Riemann–Liouville derivatives*.

Note that the fractional integrals are defined for any order $\mu > 0$, but the fractional derivatives are so far defined only for $0 < \mu < 1$.

If $f(x) \in AC$, then the derivatives $\mathbf{D}_{a+}^\mu f(x)$ and $\mathbf{D}_{b-}^\mu f(x)$, $0 < \mu < 1$, exist almost everywhere, and we have $\mathbf{D}_{a+}^\mu f(x) \in L_r(a, b)$ and $\mathbf{D}_{b-}^\mu f(x) \in L_r(a, b)$, $1 \leq r < 1/\mu$. These derivatives have the representations

$$\mathbf{D}_{a+}^\mu f(x) = \frac{1}{\Gamma(1-\mu)} \left[\frac{f(a)}{(x-a)^\mu} + \int_a^x \frac{f'_t(t)}{(x-t)^\mu} dt \right], \quad (7)$$

$$\mathbf{D}_{b-}^\mu f(x) = \frac{1}{\Gamma(1-\mu)} \left[\frac{f(b)}{(b-x)^\mu} - \int_x^b \frac{f'_t(t)}{(t-x)^\mu} dt \right]. \quad (8)$$

Finally, let us pass to the fractional derivatives of order $\mu \geq 1$. We shall use the following notation: $[\mu]$ stands for the integral part of a real number μ and $\{\mu\}$ is the fractional part of μ , $0 \leq \{\mu\} < 1$, so that

$$\mu = [\mu] + \{\mu\}. \quad (9)$$

If μ is an integer, then by the fractional derivative of order μ we mean the ordinary derivative

$$\mathbf{D}_{a+}^\mu = \left(\frac{d}{dx} \right)^\mu, \quad \mathbf{D}_{b-}^\mu = \left(-\frac{d}{dx} \right)^\mu, \quad \mu = 1, 2, \dots \quad (10)$$

However, if μ is not integral, then $\mathbf{D}_{a+}^\mu f$ and $\mathbf{D}_{b-}^\mu f$ are introduced by the formulas

$$\mathbf{D}_{a+}^\mu f(x) \equiv \left(\frac{d}{dx} \right)^{[\mu]} \mathbf{D}_{a+}^{\{\mu\}} f(x) = \left(\frac{d}{dx} \right)^{[\mu]+1} \mathbf{I}_{a+}^{1-\{\mu\}} f(x), \quad (11)$$

$$\mathbf{D}_{b-}^\mu f(x) \equiv \left(-\frac{d}{dx} \right)^{[\mu]} \mathbf{D}_{b-}^{\{\mu\}} f(x) = \left(-\frac{d}{dx} \right)^{[\mu]+1} \mathbf{I}_{b-}^{1-\{\mu\}} f(x). \quad (12)$$

Thus,

$$\mathbf{D}_{a+}^\mu f(x) = \frac{1}{\Gamma(n-\mu)} \left(\frac{d}{dx} \right)^n \int_a^x \frac{f(t)}{(x-t)^{\mu-n+1}} dt, \quad n = [\mu] + 1, \quad (13)$$

$$\mathbf{D}_{b-}^\mu f(x) = \frac{(-1)^n}{\Gamma(n-\mu)} \left(\frac{d}{dx} \right)^n \int_x^b \frac{f(t)}{(t-x)^{\mu-n+1}} dt, \quad n = [\mu] + 1. \quad (14)$$

A sufficient condition for the existence of the derivatives (13) and (14) is as follows:

$$\int_a^x \frac{f(t) dt}{(x-t)^{\{\mu\}}} \in AC^{[\mu]}.$$

This sufficient condition holds whenever $f(x) \in AC^{[\mu]}$.

Remark. The definitions of the fractional integrals and fractional derivatives can be extended to the case of complex μ (e.g., see S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993)).

10.5-3. Main Properties.

Let $\mathbf{I}_{a+}^\mu(L_1)$, $\mu > 0$, be the class of functions $f(x)$ that can be represented by the left fractional integral of order μ of an integrable function: $f(x) = \mathbf{I}_{a+}^\mu \varphi(x)$, $\varphi(x) \in L_1(a, b)$, $1 \leq p < \infty$.

For the relation $f(x) \in \mathbf{I}_{a+}^\mu(L_1)$, $\mu > 0$, to hold, it is necessary and sufficient that

$$f_{n-\mu}(x) \equiv \mathbf{I}_{a+}^{n-\mu} f \in AC^n, \quad (15)$$

where $n = [\mu] + 1$, and*

$$f_{n-\mu}^{(k)}(a) = 0, \quad k = 0, 1, \dots, n-1. \quad (16)$$

Let $\mu > 0$. We say that a function $f(x) \in L_1$ has an *integrable fractional derivative* $\mathbf{D}_{a+}^\mu f$ if $\mathbf{I}_{a+}^{n-\mu} f(x) \in AC^n$, where $n = [\mu] + 1$.

In other words, this definition introduces a notion involving only the first of the two conditions (15) and (16) describing the class $\mathbf{I}_{a+}^\mu(L_1)$.

Let $\mu > 0$. In this case the relation

$$\mathbf{D}_{a+}^\mu \mathbf{I}_{a+}^\mu \varphi(x) = \varphi(x) \quad (17)$$

holds for any integrable function $\varphi(x)$, and the relation

$$\mathbf{I}_{a+}^\mu \mathbf{D}_{a+}^\mu f(x) = f(x) \quad (18)$$

holds for any function $f(x)$ such that

$$f(x) \in \mathbf{I}_{a+}^\mu(L_1). \quad (19)$$

If we replace (19) by the condition that the function $f(x) \in L_1(a, b)$ has an integrable derivative $\mathbf{D}_{a+}^\mu f(x)$, then relation (18) fails in general and must be replaced by the formula

$$\mathbf{I}_{a+}^\mu \mathbf{D}_{a+}^\mu f(x) = f(x) - \sum_{k=0}^{n-1} \frac{(x-a)^{\mu-k-1}}{\Gamma(\mu-k)} f_{n-\mu}^{(n-k-1)}(a), \quad (20)$$

where $n = [\mu] + 1$ and $f_{n-\mu}(x) = \mathbf{I}_{a+}^{n-\mu} f(x)$. In particular, for $0 < \mu < 1$ we have

$$\mathbf{I}_{a+}^\mu \mathbf{D}_{a+}^\mu f(x) = f(x) - \frac{f_{1-\mu}(a)}{\Gamma(\mu)} (x-a)^{\mu-1}. \quad (21)$$

10.5-4. Solution of the Generalized Abel Equation.

Consider the Abel integral equation

$$\int_a^x \frac{y(t)}{(x-t)^\mu} dt = f(x), \quad (22)$$

where $0 < \mu < 1$. Suppose that $x \in [a, b]$, $f(x) \in AC$, and $y(t) \in L_1$, and apply the technique of fractional differentiation. We divide Eq. (22) by $\Gamma(1-\mu)$, and, by virtue of (1), rewrite this equation as follows:

$$\mathbf{I}_{a+}^{1-\mu} y(x) = \frac{f(x)}{\Gamma(1-\mu)}, \quad x > a. \quad (23)$$

Let us apply the operator of fractional differentiation $\mathbf{D}_{a+}^{1-\mu}$ to (23). Using the properties of the operators of fractional integration and differentiation, we obtain

$$y(x) = \frac{\mathbf{D}_{a+}^{1-\mu} f(x)}{\Gamma(1-\mu)}, \quad (24)$$

or, in the detailed notation,

$$y(x) = \frac{1}{\Gamma(\mu)\Gamma(1-\mu)} \left[\frac{f(a)}{(x-a)^{1-\mu}} + \int_a^x \frac{f'_t(t)}{(x-t)^{1-\mu}} dt \right]. \quad (25)$$

Taking into account the relation

$$\frac{1}{\Gamma(\mu)\Gamma(1-\mu)} = \frac{\sin(\pi\mu)}{\pi},$$

we now arrive at the solution of the generalized Abel equation in the form

$$y(x) = \frac{\sin(\pi\mu)}{\pi} \left[\frac{f(a)}{(x-a)^{1-\mu}} + \int_a^x \frac{f'_t(t) dt}{(x-t)^{1-\mu}} \right], \quad (26)$$

which coincides with that obtained above in Subsection 10.4-4.

* From now on in Section 10.5, by $f^{(n)}(x)$ we mean the n th derivative of $f(x)$ with respect to x and $f^{(n)}(a) \equiv f^{(n)}(x)|_{x=a}$.

10.5-5. Erdélyi–Kober Operators.

Generalized Erdélyi–Kober operators are defined by the relations

$$I_{\beta, \alpha}[f] \equiv \frac{2}{\Gamma(\alpha)} x^{-2(\alpha+\beta)} \int_a^x t^{2\beta+1} (x^2 - t^2)^{\alpha-1} f(t) dt \quad \text{if } 0 < \alpha < \infty \quad (27)$$

and

$$K_{\beta, \alpha}[f] \equiv \frac{2}{\Gamma(\alpha)} x^{2\beta} \int_x^b t^{1-2\beta-2\alpha} (t^2 - x^2)^{\alpha-1} f(t) dt \quad \text{if } 0 < \alpha < \infty.$$

The following identities hold:

$$\begin{aligned} I_{\beta, \alpha} I_{\alpha+\beta, \gamma} &= I_{\beta, \alpha+\gamma}, \\ I_{\beta, \alpha}[t^{2\gamma} f(t)] &= x^{2\gamma} I_{\beta+\gamma, \alpha}[f(t)], \\ K_{\beta, \alpha} I_{\alpha+\beta, \gamma} &= K_{\beta, \alpha+\gamma}, \\ K_{\beta, \alpha}[t^{2\gamma} f(t)] &= x^{2\gamma} K_{\beta-\gamma, \alpha}[f(t)]. \end{aligned}$$

Defining the inverse operators, one can show that

$$\begin{aligned} I_{\beta, \alpha}^{-1} &= I_{\alpha+\beta, -\alpha}, \\ K_{\beta, \alpha}^{-1} &= K_{\alpha+\beta, -\alpha}. \end{aligned} \quad (28)$$

Generalized Erdélyi–Kober operators (27) and inversion formulas (28) are used for solving some dual integral equations.

⊙ References for Section 10.5: K. B. Oldham and J. Spanier (1974), C. Nasim and B. D. Aggarwala (1984), Yu. I. Babenko (1986), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

10.6. Equations with Weakly Singular Kernel

10.6-1. Method of Transformation of the Kernel.

Consider the Volterra integral equation of the first kind with polar kernel

$$K(x, t) = \frac{L(x, t)}{(x-t)^\alpha}, \quad 0 < \alpha < 1. \quad (1)$$

The integral equation in question can be represented in the form

$$\int_0^x \frac{L(x, t)}{(x-t)^\alpha} y(t) dt = f(x), \quad (2)$$

where we assume that the functions $L(x, t)$ and $\partial L(x, t)/\partial x$ are continuous and bounded. To solve Eq. (2), we multiply it by $dx/(\xi-x)^{1-\alpha}$ and integrate from 0 to ξ , thus obtaining

$$\int_0^\xi \left[\int_0^x \frac{L(x, t)}{(x-t)^\alpha} y(t) dt \right] \frac{dx}{(\xi-x)^{1-\alpha}} = \int_0^\xi \frac{f(x) dx}{(\xi-x)^{1-\alpha}}.$$

By setting

$$\begin{aligned} K^*(\xi, t) &= \int_t^\xi \frac{L(x, t) dx}{(\xi-x)^{1-\alpha} (x-t)^\alpha}, \\ \varphi(\xi) &= \int_0^\xi \frac{f(x) dx}{(\xi-x)^{1-\alpha}}, \quad \varphi(0) = 0, \end{aligned}$$

we obtain another integral equation of the first kind with the unknown function $y(t)$:

$$\int_0^\xi K^*(\xi, t)y(t) dt = \varphi(\xi), \quad (3)$$

in which the kernel $K^*(\xi, t)$ has no singularities.

It can be shown that any solution of Eq. (3) is a solution of Eq. (2). Thus, after transforming Eq. (2) to the form (3), we can apply any methods available for continuous kernels to the latter equation.

10.6-2. Kernel with Logarithmic Singularity.

Consider the equation

$$\int_0^x \ln(x-t)y(t) dt = f(x), \quad f(0) = 0. \quad (4)$$

Let us apply the Laplace transform to solve this equation. Note that

$$\mathfrak{L}\{x^\nu\} = \int_0^\infty e^{-px} x^\nu dx = \frac{\Gamma(\nu+1)}{p^{\nu+1}}, \quad \nu > -1. \quad (5)$$

Let us differentiate relation (5) with respect to ν . We obtain

$$\mathfrak{L}\{x^\nu \ln x\} = \frac{\Gamma(\nu+1)}{p^{\nu+1}} \left[\frac{\Gamma'_z(\nu+1)}{\Gamma(\nu+1)} + \ln \frac{1}{p} \right]. \quad (6)$$

From Supplement 11.4-2, it follows that

$$\frac{\Gamma'_z(1)}{\Gamma(1)} = -\mathcal{C},$$

where $\mathcal{C} = 0.5772\dots$ is the Euler constant. With regard to the last relation, formula (6) with $\nu = 0$ becomes

$$\mathfrak{L}\{\ln x\} = -\frac{\ln p + \mathcal{C}}{p}. \quad (7)$$

Applying the Laplace transform to Eq. (4) and taking into account (7), we obtain

$$-\frac{\ln p + \mathcal{C}}{p} \tilde{y}(p) = \tilde{f}(p),$$

and hence

$$\tilde{y}(p) = -\frac{p\tilde{f}(p)}{\ln p + \mathcal{C}}. \quad (8)$$

Now let us express $\tilde{y}(p)$ in the form

$$\tilde{y}(p) = -\frac{p^2\tilde{f}(p) - f'_x(0)}{p(\ln p + \mathcal{C})} - \frac{f'_x(0)}{p(\ln p + \mathcal{C})}. \quad (9)$$

Since $f(0) = 0$, it follows that

$$\mathfrak{L}\{f''_{xx}(x)\} = p^2\tilde{f}(p) - f'_x(0). \quad (10)$$

Let us rewrite formula (5) as

$$\mathfrak{L}\left\{\frac{x^\nu}{\Gamma(\nu+1)}\right\} = \frac{1}{p^{\nu+1}} \quad (11)$$

and integrate (11) with respect to ν from 0 to ∞ . We obtain

$$\mathfrak{L}\left\{\int_0^\infty \frac{x^\nu}{\Gamma(\nu+1)} d\nu\right\} = \int_0^\infty \frac{d\nu}{p^{\nu+1}} = \frac{1}{p \ln p}.$$

Applying the scaling formula for the Laplace transform (see Table 1 in Subsection 9.2-5) we see that

$$\mathfrak{L}\left\{\int_0^\infty \frac{(x/a)^\nu}{\Gamma(\nu+1)} d\nu\right\} = \frac{1}{p \ln ap} = \frac{1}{p(\ln p + \ln a)}.$$

We set $a = e^C$ and obtain

$$\mathfrak{L} \left\{ \int_0^\infty \frac{x^\nu e^{-C\nu}}{\Gamma(\nu+1)} d\nu \right\} = \frac{1}{p(\ln p + C)}. \quad (12)$$

Let us proceed with relation (9). By (12), we have

$$\frac{f'_x(0)}{p(\ln p + C)} = \mathfrak{L} \left\{ f'_x(0) \int_0^\infty \frac{x^\nu e^{-C\nu}}{\Gamma(\nu+1)} d\nu \right\}. \quad (13)$$

Taking into account (10) and (12), we can regard the first summand on the right-hand side in (9) as a product of transforms. To find this summand itself we apply the convolution theorem:

$$\frac{p^2 \tilde{f}(p) - f'_x(0)}{p(\ln p + C)} = \mathfrak{L} \left\{ \int_0^x f''_{tt}(t) \int_0^\infty \frac{(x-t)^\nu e^{-C\nu}}{\Gamma(\nu+1)} d\nu dt \right\}. \quad (14)$$

On the basis of relations (9), (13), and (14) we obtain the solution of the integral equation (4) in the form

$$y(x) = - \int_0^x f''_{tt}(t) \int_0^\infty \frac{(x-t)^\nu e^{-C\nu}}{\Gamma(\nu+1)} d\nu dt - f'_x(0) \int_0^\infty \frac{x^\nu e^{-C\nu}}{\Gamma(\nu+1)} d\nu. \quad (15)$$

⊙ References for Section 10.6: V. Volterra (1959), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971).

10.7. Method of Quadratures

10.7-1. Quadrature Formulas.

The *method of quadratures* is a method for constructing an approximate solution of an integral equation based on the replacement of integrals by finite sums according to some formula. Such formulas are called *quadrature formulas* and, in general, have the form

$$\int_a^b \psi(x) dx = \sum_{i=1}^n A_i \psi(x_i) + \varepsilon_n[\psi], \quad (1)$$

where x_i ($i = 1, \dots, n$) are the abscissas of the partition points of the integration interval $[a, b]$, or *quadrature (interpolation) nodes*, A_i ($i = 1, \dots, n$) are numerical coefficients independent of the choice of the function $\psi(x)$, and $\varepsilon_n[\psi]$ is the remainder (the truncation error) of formula (1). As a rule, $A_i \geq 0$ and $\sum_{i=1}^n A_i = b - a$.

There are quite a few quadrature formulas of the form (1). The following formulas are the simplest and most frequently used in practice.

Rectangle rule:

$$\begin{aligned} A_1 = A_2 = \dots = A_{n-1} = h, \quad A_n = 0, \\ h = \frac{b-a}{n-1}, \quad x_i = a + h(i-1) \quad (i = 1, \dots, n). \end{aligned} \quad (2)$$

Trapezoidal rule:

$$\begin{aligned} A_1 = A_n = \frac{1}{2}h, \quad A_2 = A_3 = \dots = A_{n-1} = h, \\ h = \frac{b-a}{n-1}, \quad x_i = a + h(i-1) \quad (i = 1, \dots, n). \end{aligned} \quad (3)$$

Simpson's rule (or prismoidal formula):

$$\begin{aligned} A_1 = A_{2m+1} = \frac{1}{3}h, \quad A_2 = \dots = A_{2m} = \frac{4}{3}h, \quad A_3 = \dots = A_{2m-1} = \frac{2}{3}h, \\ h = \frac{b-a}{n-1}, \quad x_i = a + h(i-1) \quad (n = 2m+1, \quad i = 1, \dots, n), \end{aligned} \quad (4)$$

where m is a positive integer.

In formulas (2)–(4), h is a constant integration step.

The quadrature formulas due to Chebyshev and Gauss with various numbers of interpolation nodes are also widely applied. Let us illustrate these formulas by an example.

Example. For the interval $[-1, 1]$, the parameters in formula (1) acquire the following values:
Chebyshev's formula ($n = 6$):

$$\begin{aligned} A_1 = A_2 = \dots = \frac{2}{n} = \frac{1}{3}, & \quad x_1 = -x_6 = -0.8662468181, \\ x_2 = -x_5 = -0.4225186538, & \quad x_3 = -x_4 = -0.2666354015. \end{aligned} \quad (5)$$

Gauss's formula ($n = 7$):

$$\begin{aligned} A_1 = A_7 = 0.1294849662, & \quad A_2 = A_6 = 0.2797053915, \\ A_3 = A_5 = 0.3818300505, & \quad A_4 = 0.4179591837, \\ x_1 = -x_7 = -0.9491079123, & \quad x_2 = -x_6 = -0.7415311856, \\ x_3 = -x_5 = -0.4058451514, & \quad x_4 = 0. \end{aligned} \quad (6)$$

Note that a vast literature is devoted to quadrature formulas, and the reader can find books of interest (e.g., see G. A. Korn and T. M. Korn (1968), N. S. Bakhvalov (1973), S. M. Nikol'skii (1979)).

10.7-2. General Scheme of the Method.

Let us solve the Volterra integral equation of the first kind

$$\int_a^x K(x, t)y(t) dt = f(x), \quad f(a) = 0, \quad (7)$$

on an interval $a \leq x \leq b$ by the method of quadratures. The procedure of constructing the solution involves two stages:

1°. First, we determine the initial value $y(a)$. To this end, we differentiate Eq. (7) with respect to x , thus obtaining

$$K(x, x)y(x) + \int_a^x K'_x(x, t)y(t) dt = f'_x(x).$$

By setting $x = a$, we find that

$$y_1 = y(a) = \frac{f'_x(a)}{K(a, a)} = \frac{f'_x(a)}{K_{11}}.$$

2°. Let us choose a constant integration step h and consider the discrete set of points $x_i = a + h(i-1)$, $i = 1, \dots, n$. For $x = x_i$, Eq. (7) acquires the form

$$\int_a^{x_i} K(x_i, t)y(t) dt = f(x_i), \quad i = 2, \dots, n, \quad (8)$$

Applying the quadrature formula (1) to the integral in (8) and choosing x_j ($j = 1, \dots, i$) to be the nodes in t , we arrive at the system of equations

$$\sum_{j=1}^i A_{ij} K(x_i, x_j)y(x_j) = f(x_i) + \varepsilon_i[y], \quad i = 2, \dots, n, \quad (9)$$

where the A_{ij} are the coefficients of the quadrature formula on the interval $[a, x_i]$ and $\varepsilon_i[y]$ is the truncation error. Assume that the $\varepsilon_i[y]$ are small and neglect them; then we obtain a system of linear algebraic equations in the form

$$\sum_{j=1}^i A_{ij} K_{ij} y_j = f_i, \quad i = 2, \dots, n, \quad (10)$$

where $K_{ij} = K(x_i, x_j)$ ($j = 1, \dots, i$), $f_i = f(x_i)$, and y_j are approximate values of the unknown function at the nodes x_i .

Now system (10) permits one, provided that $A_{ii}K_{ii} \neq 0$ ($i = 2, \dots, n$), to successively find the desired approximate values by the formulas

$$y_1 = \frac{f'_x(a)}{K_{11}}, \quad y_2 = \frac{f_2 - A_{21}K_{21}y_1}{A_{22}K_{22}}, \quad \dots, \quad y_n = \frac{f_n - \sum_{j=1}^{n-1} A_{nj}K_{nj}y_j}{A_{nn}K_{nn}},$$

whose specific form depends on the choice of the quadrature formula.

10.7-3. Algorithm Based on the Trapezoidal Rule.

According to the trapezoidal rule (3), we have

$$A_{i1} = A_{ii} = \frac{1}{2}h, \quad A_{i2} = \dots = A_{i,i-1} = h, \quad i = 2, \dots, n.$$

The application of the trapezoidal rule in the general scheme leads to the following step algorithm:

$$y_1 = \frac{f'_x(a)}{K_{11}}, \quad f'_x(a) = \frac{-3f_1 + 4f_2 - f_3}{2h},$$

$$y_i = \frac{2}{K_{ii}} \left(\frac{f_i}{h} - \sum_{j=1}^{i-1} \beta_j K_{ij} y_j \right), \quad \beta_j = \begin{cases} \frac{1}{2} & \text{for } j = 1, \\ 1 & \text{for } j > 1, \end{cases} \quad i = 2, \dots, n,$$

where the notation coincides with that introduced in Subsection 10.7-2. The trapezoidal rule is quite simple and effective and frequently used in practice for solving integral equations with variable limit of integration.

On the basis of Subsections 10.7-1 and 10.7-2, one can write out similar expressions for other quadrature formulas. However, they must be used with care. For example, the application of Simpson's rule must be alternated, for odd nodes, with some other rule, e.g., the rectangle rule or the trapezoidal rule. For equations with variable integration limit, the use of Chebyshev's formula or Gauss's formula also has some difficulties as well.

10.7-4. Algorithm for an Equation with Degenerate Kernel.

A general property of the algorithms of the method of quadratures in the solution of the Volterra equations of the first kind with arbitrary kernel is that the amount of computational work at each step is proportional to the number of the step: all operations of the previous step are repeated with new data and another term in the sum is added.

However, if the kernel in Eq. (7) is degenerate, i.e.,

$$K(x, t) = \sum_{k=1}^m p_k(x)q_k(t), \quad (11)$$

or if the kernel under consideration can be approximated by a degenerate kernel, then an algorithm can be constructed for which the number of operations does not depend on the index of the digitalization node. With regard to (11), Eq. (7) becomes

$$\sum_{k=1}^m p_k(x) \int_a^x q_k(t)y(t) dt = f(x). \quad (12)$$

By applying the trapezoidal rule to (12), we obtain recurrent expressions for the solution of the equation (see formulas in Subsection 10.7-3):

$$y(a) = \frac{f'_x(a)}{\sum_{k=1}^m p_k(a)q_k(a)}, \quad y_i = \frac{2}{\sum_{k=1}^m p_{ki}q_{ki}} \left[\frac{f_i}{h} - \sum_{k=1}^m p_{ki} \sum_{j=1}^{i-1} \beta_j q_{kj} y_j \right],$$

where y_i are approximate values of $y(x)$ at x_i , $f_i = f(x_i)$, $p_{ki} = p_k(x_i)$, and $q_{ki} = q_k(x_i)$.

© References for Section 10.7: G. A. Korn and T. M. Korn (1968), N. S. Bakhvalov (1973), V. I. Krylov, V. V. Bobkov, and P. I. Monastyryi (1984), A. F. Verlan' and V. S. Sizikov (1986).

10.8. Equations with Infinite Integration Limit

Integral equations of the first kind with difference kernel in which one of the limits of integration is variable and the other is infinite are of interest. Sometimes the kernels and the functions of these equations do not belong to the classes described in the beginning of the chapter. The investigation of these equations can be performed by the method of model solutions (see Section 11.6) or by the method of reducing to equations of the convolution type. Let us consider these methods for an example of an equation of the first kind with variable lower limit of integration.

10.8-1. Equation of the First Kind with Variable Lower Limit of Integration.

Consider the equation of the first kind with difference kernel

$$\int_x^\infty K(x-t)y(t) dt = f(x). \quad (1)$$

Equation (1) cannot be solved by direct application of the Laplace transform, because the convolution theorem cannot be used here. According to the method of model solutions whose detailed exposition can be found in Section 11.6, we consider the auxiliary equation with exponential right-hand side

$$\int_x^\infty K(x-t)y(t) dt = e^{px}. \quad (2)$$

The solution of (2) has the form

$$Y(x, p) = \frac{1}{\tilde{K}(-p)} e^{px}, \quad \tilde{K}(-p) = \int_0^\infty K(-z)e^{pz} dz. \quad (3)$$

On the basis of these formulas and formula (11) from Section 11.6, we obtain the solution of Eq. (1) for an arbitrary right-hand side $f(x)$ in the form

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(p)}{\tilde{K}(-p)} e^{px} dp, \quad (4)$$

where $\tilde{f}(p)$ is the Laplace transform of the function $f(x)$.

Example. Consider the following integral equation of the first kind with variable lower limit of integration:

$$\int_x^\infty e^{a(x-t)}y(t) dt = A \sin(bx), \quad a > 0. \quad (5)$$

According to (3) and (4), we can write out the expressions for $\tilde{f}(p)$ (see Supplement 5) and $\tilde{K}(-p)$,

$$\tilde{f}(p) = \frac{Ab}{p^2 + b^2}, \quad \tilde{K}(-p) = \int_0^\infty e^{(p-a)z} dz = \frac{1}{a-p}, \quad (6)$$

and the solution of Eq. (5) in the form

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{Ab(a-p)}{p^2 + b^2} e^{px} dp. \quad (7)$$

Now using the tables of inverse Laplace transforms (see Supplement 6), we obtain the exact solution

$$y(x) = Aa \sin(bx) - Ab \cos(bx), \quad a > 0, \quad (8)$$

which can readily be verified by substituting (8) into (5) and using the tables of integrals in Supplement 3.

10.8-2. Reduction to a Wiener–Hopf Equation of the First Kind.
--

Equation (1) can be reduced to a first-kind one-sided equation

$$\int_0^{\infty} K_-(x-t)y(t) dt = -f(x), \quad 0 < x < \infty, \quad (9)$$

where the kernel $K_-(x-t)$ has the following form:

$$K_-(s) = \begin{cases} 0 & \text{for } s > 0, \\ -K(s) & \text{for } s < 0. \end{cases}$$

Methods for studying Eq. (9) are described in Chapter 12.

- ⊙ References for Section 10.8: F. D. Gakhov and Yu. I. Cherskii (1978), A. D. Polyanin and A. V. Manzhirov (1997).

Chapter 11

Methods for Solving Linear Equations of the Form $y(x) - \int_a^x K(x, t)y(t) dt = f(x)$

11.1. Volterra Integral Equations of the Second Kind

11.1-1. Preliminary Remarks. Equations for the Resolvent.

In this chapter we present methods for solving Volterra integral equations of the second kind, which have the form

$$y(x) - \int_a^x K(x, t)y(t) dt = f(x), \quad (1)$$

where $y(x)$ is the unknown function ($a \leq x \leq b$), $K(x, t)$ is the kernel of the integral equation, and $f(x)$ is the *right-hand side* of the integral equation. The function classes to which $y(x)$, $f(x)$, and $K(x, t)$ can belong are defined in Subsection 10.1-1. In these function classes, there exists a unique solution of the Volterra integral equation of the second kind.

Equation (1) is said to be *homogeneous* if $f(x) \equiv 0$ and *nonhomogeneous* otherwise.

The kernel $K(x, t)$ is said to be *degenerate* if it can be represented in the form

$$K(x, t) = g_1(x)h_1(t) + \cdots + g_n(x)h_n(t).$$

The kernel $K(x, t)$ of an integral equation is called *difference kernel* if it depends only on the difference of the arguments,

$$K(x, t) = K(x - t).$$

Remark 1. A homogeneous Volterra integral equation of the second kind has only the trivial solution.

Remark 2. The existence and uniqueness of the solution of a Volterra integral equation of the second kind hold for a much wider class of kernels and functions.

Remark 3. A Volterra equation of the second kind can be regarded as a Fredholm equation of the second kind whose kernel $K(x, t)$ vanishes for $t > x$ (see Chapter 13).

Remark 4. The case in which $a = -\infty$ and/or $b = \infty$ is not excluded, but in this case the square integrability of the kernel $K(x, t)$ on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ is especially significant.

The solution of Eq. (1) can be presented in the form

$$y(x) = f(x) + \int_a^x R(x, t)f(t) dt, \quad (2)$$

where the resolvent $R(x, t)$ is independent of $f(x)$ and the lower limit of integration a and is determined by the kernel of the integral equation alone.

The resolvent of the Volterra equation (1) satisfies the following two integral equations:

$$R(x, t) = K(x, t) + \int_t^x K(x, s)R(s, t) ds, \quad (3)$$

$$R(x, t) = K(x, t) + \int_t^x K(s, t)R(x, s) ds, \quad (4)$$

in which the integration is performed with respect to different pairs of variables of the kernel and the resolvent.

11.1-2. Relationship Between Solutions of Some Integral Equations.

Let us present two useful formulas that express the solution of one integral equation via the solutions of other integral equations.

1°. Assume that the Volterra equation of the second kind with kernel $K(x, t)$ has a resolvent $R(x, t)$. Then the Volterra equation of the second kind with kernel $K^*(x, t) = -K(t, x)$ has the resolvent $R^*(x, t) = -R(t, x)$.

2°. Assume that two Volterra equations of the second kind with kernels $K_1(x, t)$ and $K_2(x, t)$ are given and that resolvents $R_1(x, t)$ and $R_2(x, t)$ correspond to these equations. In this case the Volterra equation with kernel

$$K(x, t) = K_1(x, t) + K_2(x, t) - \int_t^x K_1(x, s)K_2(s, t) ds \quad (5)$$

has the resolvent

$$R(x, t) = R_1(x, t) + R_2(x, t) + \int_t^x R_1(s, t)R_2(x, s) ds. \quad (6)$$

Note that in formulas (5) and (6), the integration is performed with respect to different pairs of variables.

⊙ References for Section 11.1: E. Goursat (1923), H. M. Müntz (1934), V. Volterra (1959), S. G. Mikhailin (1960), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), J. A. Cochran (1972), V. I. Smirnov (1974), P. P. Zabreyko, A. I. Koshelev, et al. (1975), A. J. Jerry (1985), F. G. Tricomi (1985), A. F. Verlan' and V. S. Sizikov (1986), P. Linz (1987), G. Gripenberg, S.-O. Londen, and O. Staffans (1990), C. Corduneanu (1991), R. Gorenflo and S. Vessella (1991), A. C. Pipkin (1991).

11.2. Equations with Degenerate Kernel:

$$K(x, t) = g_1(x)h_1(t) + \cdots + g_n(x)h_n(t)$$

11.2-1. Equations with Kernel of the Form $K(x, t) = \varphi(x) + \psi(x)(x-t)$.

The solution of a Volterra equation (see Subsection 11.1-1) with kernel of this type can be expressed by the formula

$$y = w''_{xx}, \quad (1)$$

where $w = w(x)$ is the solution of the second-order linear nonhomogeneous ordinary differential equation

$$w''_{xx} - \varphi(x)w'_x - \psi(x)w = f(x), \quad (2)$$

with the initial conditions

$$w(a) = w'_x(a) = 0. \quad (3)$$

Let $w_1 = w_1(x)$ be a nontrivial particular solution of the corresponding homogeneous linear differential equation (2) for $f(x) \equiv 0$. Assume that $w_1(a) \neq 0$. In this case, the other nontrivial particular solution $w_2 = w_2(x)$ of this homogeneous linear differential equation has the form

$$w_2(x) = w_1(x) \int_a^x \frac{\Phi(t)}{[w_1(t)]^2} dt, \quad \Phi(x) = \exp \left[\int_a^x \varphi(s) ds \right].$$

The solution of the nonhomogeneous equation (2) with the initial conditions (3) is given by the formula

$$w(x) = w_2(x) \int_a^x \frac{w_1(t)}{\Phi(t)} f(t) dt - w_1(x) \int_a^x \frac{w_2(t)}{\Phi(t)} f(t) dt. \quad (4)$$

On substituting expression (4) into formula (1) we obtain the solution of the original integral equation in the form

$$y(x) = f(x) + \int_a^x R(x, t) f(t) dt,$$

where

$$\begin{aligned} R(x, t) &= [w_2''(x)w_1(t) - w_1''(x)w_2(t)] \frac{1}{\Phi(t)} \\ &= \varphi(x) \frac{\Phi(x)}{w_1(x)} \frac{w_1(t)}{\Phi(t)} + [\varphi(x)w_1'(x) + \psi(x)w_1(x)] \frac{w_1(t)}{\Phi(t)} \int_t^x \frac{\Phi(s)}{[w_1(s)]^2} ds. \end{aligned}$$

Here $\Phi(x) = \exp \left[\int_a^x \varphi(s) ds \right]$ and the primes stand for x -derivatives.

For a degenerate kernel of the above form, the resolvent can be defined by the formula

$$R(x, t) = u''_{xx},$$

where the auxiliary function u is the solution of the homogeneous linear second-order ordinary differential equation

$$u''_{xx} - \varphi(x)u'_x - \psi(x)u = 0 \quad (5)$$

with the following initial conditions at $x = t$:

$$u|_{x=t} = 0, \quad u'_x|_{x=t} = 1. \quad (6)$$

The parameter t occurs only in the initial conditions (6), and Eq. (5) itself is independent of t .

Remark 1. The kernel of the integral equation in question can be rewritten in the form $K(x, t) = G_1(x) + tG_2(x)$, where $G_1(x) = \varphi(x) + x\psi(x)$ and $G_2(x) = -\varphi(x)$.

11.2-2. Equations with Kernel of the Form $K(x, t) = \varphi(t) + \psi(t)(t - x)$.

For a degenerate kernel of the above form, the resolvent is determined by the expression

$$R(x, t) = -v''_{tt}, \quad (7)$$

where the auxiliary function v is the solution of the homogeneous linear second-order ordinary differential equation

$$v''_{tt} + \varphi(t)v'_t + \psi(t)v = 0 \quad (8)$$

with the following initial conditions at $t = x$:

$$v|_{t=x} = 0, \quad v'_t|_{t=x} = 1. \quad (9)$$

The point x occurs only in the initial data (9) as a parameter, and Eq. (8) itself is independent of x .

Assume that $v_1 = v_1(t)$ is a nontrivial particular solution of Eq. (8). In this case, the general solution of this differential equation is given by the formula

$$v(t) = C_1 v_1(t) + C_2 v_1(t) \int_a^t \frac{ds}{\Phi(s)[v_1(s)]^2}, \quad \Phi(t) = \exp\left[\int_a^t \varphi(s) ds\right].$$

Taking into account the initial data (9), we find the dependence of the integration constants C_1 and C_2 on the parameter x . As a result, we obtain the solution of problem (8), (9):

$$v = v_1(x)\Phi(x) \int_x^t \frac{ds}{\Phi(s)[v_1(s)]^2}. \tag{10}$$

On substituting the expression (10) into formula (7) and eliminating the second derivative by means of Eq. (8) we find the resolvent:

$$R(x, t) = \varphi(t) \frac{v_1(x)\Phi(x)}{v_1(t)\Phi(t)} + v_1(x)\Phi(x)[\varphi(t)v'_t(t) + \psi(t)v_1(t)] \int_x^t \frac{ds}{\Phi(s)[v_1(s)]^2}.$$

Remark 2. The kernel of the integral equation under consideration can be rewritten in the form $K(x, t) = G_1(t) + xG_2(t)$, where $G_1(t) = \varphi(t) + t\psi(t)$ and $G_2(t) = -\varphi(t)$.

11.2-3. Equations with Kernel of the Form $K(x, t) = \sum_{m=1}^n \varphi_m(x)(x-t)^{m-1}$.

To find the resolvent, we introduce an auxiliary function as follows:

$$u(x, t) = \frac{1}{(n-1)!} \int_t^x R(s, t)(x-s)^{n-1} ds + \frac{(x-t)^{n-1}}{(n-1)!};$$

at $x = t$, this function vanishes together with the first $n - 2$ derivatives with respect to x , and the $(n - 1)$ st derivative at $x = t$ is equal to 1. Moreover,

$$R(x, t) = u_x^{(n)}(x, t), \quad u_x^{(n)} = \frac{d^n u(x, t)}{dx^n}. \tag{11}$$

On substituting relation (11) into the resolvent equation (3) of Subsection 11.1-1, we see that

$$u_x^{(n)}(x, t) = K(x, t) + \int_t^x K(x, s)u_s^{(n)}(s, t) ds. \tag{12}$$

Integrating by parts the right-hand side in (12), we obtain

$$u_x^{(n)}(x, t) = K(x, t) + \sum_{m=0}^{n-1} (-1)^m K_s^{(m)}(x, s)u_s^{(n-m-1)}(s, t)\Big|_{s=t}^{s=x}. \tag{13}$$

On substituting the expressions for $K(x, t)$ and $u(x, t)$ into (13), we arrive at a linear homogeneous ordinary differential equation of order n for the function $u(x, t)$.

Thus, the resolvent $R(x, t)$ of the Volterra integral equation with degenerate kernel of the above form can be obtained by means of (11), where $u(x, t)$ satisfies the following differential equation and initial conditions:

$$u_x^{(n)} - \varphi_1(x)u_x^{(n-1)} - \varphi_2(x)u_x^{(n-2)} - 2\varphi_3(x)u_x^{(n-3)} - \dots - (n-1)! \varphi_n(x)u = 0, \\ u|_{x=t} = u'_x|_{x=t} = \dots = u_x^{(n-2)}|_{x=t} = 0, \quad u_x^{(n-1)}|_{x=t} = 1.$$

The parameter t occurs only in the initial conditions, and the equation itself is independent of t explicitly.

Remark 3. A kernel of the form $K(x, t) = \sum_{m=1}^n \phi_m(x)t^{m-1}$ can be reduced to a kernel of the above type by elementary transformations.

11.2-4. Equations with Kernel of the Form $K(x, t) = \sum_{m=1}^n \varphi_m(t)(t-x)^{m-1}$.

Let us represent the resolvent of this degenerate kernel in the form

$$R(x, t) = -v_t^{(n)}(x, t), \quad v_t^{(n)} = \frac{d^n v(x, t)}{dt^n},$$

where the auxiliary function $v(x, t)$ vanishes at $t = x$ together with $n - 2$ derivatives with respect to t , and the $(n - 1)$ st derivative with respect to t at $t = x$ is equal to 1. On substituting the expression for the resolvent into Eq. (3) of Subsection 11.1-1, we obtain

$$v_t^{(n)}(x, t) = \int_t^x K(s, t)v_s^{(n)}(x, s) ds - K(x, t).$$

Let us apply integration by parts to the integral on the right-hand side. Taking into account the properties of the auxiliary function $v(x, t)$, we arrive at the following Cauchy problem for an n th-order ordinary differential equation:

$$v_t^{(n)} + \varphi_1(t)v_t^{(n-1)} + \varphi_2(t)v_t^{(n-2)} + 2\varphi_3(t)v_t^{(n-3)} + \dots + (n - 1)! \varphi_n(t)v = 0, \\ v|_{t=x} = v'_t|_{t=x} = \dots = v_t^{(n-2)}|_{t=x} = 0, \quad v_t^{(n-1)}|_{t=x} = 1.$$

The parameter x occurs only in the initial conditions, and the equation itself is independent of x explicitly.

Remark 4. A kernel of the form $K(x, t) = \sum_{m=1}^n \phi_m(t)x^{m-1}$ can be reduced to a kernel of the above type by elementary transformations.

11.2-5. Equations with Degenerate Kernel of the General Form.

In this case, the Volterra equation of the second kind can be represented in the form

$$y(x) - \sum_{m=1}^n g_m(x) \int_a^x h_m(t)y(t) dt = f(x). \tag{14}$$

Let us introduce the notation

$$w_j(x) = \int_a^x h_j(t)y(t) dt, \quad j = 1, \dots, n, \tag{15}$$

and rewrite Eq. (14) as follows:

$$y(x) = \sum_{m=1}^n g_m(x)w_m(x) + f(x). \tag{16}$$

On differentiating the expressions (15) with regard to formula (16), we arrive at the following system of linear differential equations for the functions $w_j = w_j(x)$:

$$w'_j = h_j(x) \left[\sum_{m=1}^n g_m(x)w_m + f(x) \right], \quad j = 1, \dots, n,$$

with the initial conditions

$$w_j(a) = 0, \quad j = 1, \dots, n.$$

Once the solution of this system is found, the solution of the original integral equation (14) is defined by formula (16) or any of the expressions

$$y(x) = \frac{w'_j(x)}{h_j(x)}, \quad j = 1, \dots, n,$$

which can be obtained from formula (15) by differentiation.

⊙ References for Section 11.2: E. Goursat (1923), H. M. Müntz (1934), A. F. Verlan' and V. S. Sizikov (1986), A. D. Polyaniin and A. V. Manzhirrov (1998).

11.3. Equations with Difference Kernel: $K(x, t) = K(x - t)$

11.3-1. Solution Method Based on the Laplace Transform.

Volterra equations of the second kind with kernel depending on the difference of the arguments have the form

$$y(x) - \int_0^x K(x-t)y(t) dt = f(x). \quad (1)$$

Applying the Laplace transform \mathfrak{L} to Eq. (1) and taking into account the fact that by the convolution theorem (see Subsection 9.2-4) the integral with kernel depending on the difference of the arguments is transformed into the product $\tilde{K}(p)\tilde{y}(p)$, we arrive at the following equation for the transform of the unknown function:

$$\tilde{y}(p) - \tilde{K}(p)\tilde{y}(p) = \tilde{f}(p). \quad (2)$$

The solution of Eq. (2) is given by the formula

$$\tilde{y}(p) = \frac{\tilde{f}(p)}{1 - \tilde{K}(p)}, \quad (3)$$

which can be written equivalently in the form

$$\tilde{y}(p) = \tilde{f}(p) + \tilde{R}(p)\tilde{f}(p), \quad \tilde{R}(p) = \frac{\tilde{K}(p)}{1 - \tilde{K}(p)}. \quad (4)$$

On applying the Laplace inversion formula to (4), we obtain the solution of Eq. (1) in the form

$$y(x) = f(x) + \int_0^x R(x-t)f(t) dt, \quad (5)$$

$$R(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \tilde{R}(p)e^{px} dp.$$

When applying formula (5) in practice, the following two technical problems occur:

1°. Finding the transform $\tilde{K}(p) = \int_0^\infty K(x)e^{-px} dx$ for a given kernel $K(x)$.

2°. Finding the resolvent (5) whose transform $\tilde{R}(p)$ is given by formula (4).

To calculate the corresponding integrals, tables of direct and inverse Laplace transforms can be applied (see Supplements 5 and 6), and, in many cases, to find the inverse transform, methods of the theory of functions of a complex variable are applied, including the Cauchy residue theorem (see Subsection 9.1-4).

Remark. If the lower limit of the integral in the Volterra equation with kernel depending on the difference of the arguments is equal to a , then this equation can be reduced to Eq. (1) by the change of variables $x = \bar{x} - a$, $t = \bar{t} - a$.

Figure 3 depicts the principal scheme of solving Volterra integral equations of the second kind with difference kernel by means of the Laplace integral transform.




Figure 3. Scheme of solving Volterra integral equations of the second kind with difference kernel by means of the Laplace integral transform, $R(x)$ is the inverse transform of the function $\tilde{R}(p) = \frac{\tilde{K}(p)}{1 - \tilde{K}(p)}$.

Example 1. Consider the equation

$$y(x) + A \int_0^x \sin[\lambda(x-t)]y(t) dt = f(x), \quad (6)$$

which is a special case of Eq. (1) for $K(x) = -A \sin(\lambda x)$.

We first apply the table of Laplace transforms (see Supplement 5) and obtain the transform of the kernel of the integral equation in the form

$$\tilde{K}(p) = -\frac{A\lambda}{p^2 + \lambda^2}.$$

Next, by formula (4) we find the transform of the resolvent:

$$\tilde{R}(p) = -\frac{A\lambda}{p^2 + \lambda(A + \lambda)}.$$

Furthermore, applying the table of inverse Laplace transforms (see Supplement 6) we obtain the resolvent:

$$R(x) = \begin{cases} -\frac{A\lambda}{k} \sin(kx) & \text{for } \lambda(A + \lambda) > 0, \\ -\frac{A\lambda}{k} \sinh(kx) & \text{for } \lambda(A + \lambda) < 0, \end{cases} \quad \text{where } k = |\lambda(A + \lambda)|^{1/2}.$$

Moreover, in the special case $\lambda = -A$, we have $R(x) = A^2 x$. On substituting the expressions for the resolvent into formula (5), we find the solution of the integral equation (6). In particular, for $\lambda(A + \lambda) > 0$, this solution has the form

$$y(x) = f(x) - \frac{A\lambda}{k} \int_0^x \sin[k(x-t)]f(t) dt, \quad k = \sqrt{\lambda(A + \lambda)}. \quad (7)$$

The Laplace transformation can also be used for finding solutions of integro-differential equations with difference kernel.

Example 2. Consider the Cauchy problem for the integro-differential equation

$$\frac{dy}{dx} + \int_0^x K(x-t)y(t) dt = f(x) \quad (0 \leq x < \infty) \quad (8)$$

with the initial condition

$$y = a \quad \text{at} \quad x = 0. \quad (9)$$

Let us multiply equation (8) by e^{-px} and integrate the result with respect to x from zero to infinity. Using properties 7 and 12 of the Laplace transform (Table 1, Subsection 9.2-4) and taking into account the initial condition (9), we obtain a linear algebraic equation for the transform $\tilde{y}(p)$:

$$p\tilde{y}(p) - a + \tilde{K}(p)\tilde{y}(p) = \tilde{f}(p).$$

It follows that

$$\tilde{y}(p) = \frac{\tilde{f}(p) + a}{p + \tilde{K}(p)}.$$

By the inversion formula (see formula (2) of Subsection 9.2-1), the solution to the original problem (8)–(9) is found in the form

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(p) + a}{p + \tilde{K}(p)} e^{px} dp, \quad i^2 = -1. \quad (10)$$

Consider the special case of $a = 0$ and $K(x) = \cos(bx)$. From row 10 of Table 2 it follows that $\tilde{K}(p) = \frac{p}{p^2 + b^2}$. Rearranging the integrand in (10), we get

$$\frac{\tilde{f}(p)}{p + \tilde{K}(p)} = \frac{p^2 + b^2}{p(p^2 + b^2 + 1)} \tilde{f}(p) = \left(\frac{1}{p} - \frac{1}{p(p^2 + b^2 + 1)} \right) \tilde{f}(p).$$

In order to invert this expression, let us use the convolution theorem (see row 12 of Table 1) as well as formulas 1 and 28 for the inversion of rational functions, Supplement 6.2. As a result, we arrive at the solution in the form

$$y(x) = \int_0^x \frac{b^2 + \cos(t\sqrt{b^2 + 1})}{b^2 + 1} f(x-t) dt.$$

11.3-2. Method Based on the Solution of an Auxiliary Equation.

Consider the integral equation

$$Ay(x) + B \int_a^x K(x-t)y(t) dt = f(x). \quad (11)$$

Let $w = w(x)$ be a solution of the simpler auxiliary equation with $f(x) \equiv 1$ and $a = 0$,

$$Aw(x) + B \int_0^x K(x-t)w(t) dt = 1. \quad (12)$$

In this case, the solution of the original equation (11) with an arbitrary right-hand side can be expressed via the solution of the auxiliary equation (12) by the formula

$$y(x) = \frac{d}{dx} \int_a^x w(x-t)f(t) dt = f(a)w(x-a) + \int_a^x w(x-t)f'_t(t) dt. \quad (13)$$

Let us prove this assertion. We rewrite expression (13) (in which we first redenote the integration parameter t by s) in the form

$$y(x) = \frac{d}{dx} I(x), \quad I(x) = \int_a^x w(x-s)f(s) ds \quad (14)$$

and substitute it into the left-hand side of Eq. (11). After some algebraic manipulations and after changing the order of integration in the double integral with regard to (12), we obtain

$$\begin{aligned} \frac{d}{dx} AI(x) + B \int_a^x K(x-t) \frac{d}{dt} I(t) dt &= \frac{d}{dx} AI(x) + \frac{d}{dx} B \int_a^x K(x-t)I(t) dt \\ &= \frac{d}{dx} \left[A \int_a^x w(x-s)f(s) ds + B \int_a^x \int_a^t K(x-t)w(t-s)f(s) ds dt \right] \\ &= \frac{d}{dx} \left\{ \int_a^x f(s) \left[Aw(x-s) + B \int_s^x K(x-t)w(t-s) dt \right] ds \right\} \\ &= \frac{d}{dx} \left\{ \int_a^x f(s) \left[Aw(x-s) + B \int_0^{x-s} K(x-s-\lambda)w(\lambda) d\lambda \right] ds \right\} = \frac{d}{dx} \int_a^x f(s) ds = f(x), \end{aligned}$$

which proves the desired assertion.

11.3-3. Reduction to Ordinary Differential Equations.

Consider the special case in which the transform of the kernel of the integral equation (1) can be expressed in the form

$$1 - \tilde{K}(p) = \frac{Q(p)}{R(p)}, \quad (15)$$

where $Q(p)$ and $R(p)$ are polynomials of degree n :

$$Q(p) = \sum_{k=0}^n A_k p^k, \quad R(p) = \sum_{k=0}^n B_k p^k. \quad (16)$$

In this case, the solution of the integral equation (1) satisfies the following linear nonhomogeneous ordinary differential equation of order n with constant coefficients:

$$\sum_{k=0}^n A_k y_x^{(k)}(x) = \sum_{k=0}^n B_k f_x^{(k)}(x). \quad (17)$$

Equation (17) can be rewritten in the operator form

$$Q(D)y(x) = R(D)f(x), \quad D \equiv \frac{d}{dx}.$$

The initial conditions for Eq. (17) can be found from the relation

$$\sum_{k=0}^n A_k \sum_{s=0}^{k-1} p^{k-1-s} y_x^{(s)}(0) - \sum_{k=0}^n B_k \sum_{s=0}^{k-1} p^{k-1-s} f_x^{(s)}(0) = 0 \quad (18)$$

by matching the coefficients of like powers of the parameter p .

The proof of this assertion can be performed by applying the Laplace transform to the differential equation (17) and by the subsequent comparison of the resulting expression with Eq. (2) with regard to (15).

Another method of reducing an integral equation to an ordinary differential equation is described in Section 11.7.

11.3-4. Reduction to a Wiener–Hopf Equation of the Second Kind.

A Volterra equation of the second kind with the difference kernel of the form

$$y(x) + \int_0^x K(x-t)y(t) dt = f(x), \quad 0 < x < \infty, \quad (19)$$

can be reduced to the Wiener–Hopf equation

$$y(x) + \int_0^\infty K_+(x-t)y(t) dt = f(x), \quad 0 < x < \infty, \quad (20)$$

where the kernel $K_+(x-t)$ is given by

$$K_+(s) = \begin{cases} K(s) & \text{for } s > 0, \\ 0 & \text{for } s < 0. \end{cases}$$

Methods for studying Eq. (20) are described in Chapter 13, where an example of constructing a solution of a Volterra equation of the second kind with difference kernel by means of constructing a solution of the corresponding Wiener–Hopf equation of the second kind is presented (see Subsection 13.10-3).

11.3-5. Method of Fractional Integration for the Generalized Abel Equation.

Consider the generalized Abel equation of the second kind

$$y(x) - \lambda \int_a^x \frac{y(t)}{(x-t)^\mu} dt = f(x), \quad x > a, \quad (21)$$

where $0 < \mu < 1$. Let us assume that $x \in [a, b]$, $f(x) \in AC$, and $y(t) \in L_1$, and apply the technique of the fractional integration (see Section 10.5). We set

$$\mu = 1 - \beta, \quad 0 < \beta < 1, \quad \lambda = \frac{\nu}{\Gamma(\beta)}, \quad (22)$$

and use formula (8) from Subsection 10.5-1 to rewrite Eq. (21) in the form

$$(1 - \nu \mathbf{I}_{a+}^\beta) y(x) = f(x), \quad x > a. \quad (23)$$

Now the solution of the generalized Abel equation of the second kind can be symbolically written as follows:

$$y(x) = (1 - \nu \mathbf{I}_{a+}^\beta)^{-1} f(x), \quad x > a. \quad (24)$$

On expanding the operator expression in the parentheses in a series in powers of the operator by means of the formula for a geometric progression, we obtain

$$y(x) = \left[1 + \sum_{n=1}^{\infty} (\nu \mathbf{I}_{a+}^\beta)^n \right] f(x), \quad x > a. \quad (25)$$

Taking into account the relation $(\mathbf{I}_{a+}^\beta)^n = \mathbf{I}_{a+}^{\beta n}$, we can rewrite formula (25) in the expanded form

$$y(x) = f(x) + \sum_{n=1}^{\infty} \frac{\nu^n}{\Gamma(\beta n)} \int_a^x (x-t)^{\beta n - 1} f(t) dt, \quad x > a. \quad (26)$$

Let us transpose the integration and summation in the expression (26). Note that

$$\sum_{n=1}^{\infty} \frac{\nu^n (x-t)^{\beta n - 1}}{\Gamma(\beta n)} = \frac{d}{dx} \sum_{n=1}^{\infty} \frac{\nu^n (x-t)^{\beta n}}{\Gamma(1 + \beta n)}.$$

In this case, taking into account the change of variables (22), we see that a solution of the generalized Abel equation of the second kind becomes

$$y(x) = f(x) + \int_a^x R(x-t) f(t) dt, \quad x > a, \quad (27)$$

where the resolvent $R(x-t)$ is given by the formula

$$R(x-t) = \frac{d}{dx} \sum_{n=1}^{\infty} \frac{[\lambda \Gamma(1-\mu)(x-t)^{(1-\mu)}]^n}{\Gamma[1 + (1-\mu)n]}. \quad (28)$$

In some cases, the sum of the series in the representation (28) of the resolvent can be found, and a closed-form expression for this sum can be obtained.

Example 3. Consider the Abel equation of the second kind (we set $\mu = \frac{1}{2}$ in Eq. (21))

$$y(x) - \lambda \int_a^x \frac{y(t)}{\sqrt{x-t}} dt = f(x), \quad x > a. \quad (29)$$

By virtue of formula (28), the resolvent for Eq. (29) is given by the expression

$$R(x-t) = \frac{d}{dx} \sum_{n=1}^{\infty} \frac{[\lambda \sqrt{\pi(x-t)}]^n}{\Gamma(1 + \frac{1}{2}n)}. \quad (30)$$

We have

$$\sum_{n=1}^{\infty} \frac{x^{n/2}}{\Gamma(1 + \frac{1}{2}n)} = e^x \operatorname{erf} \sqrt{x}, \quad \operatorname{erf} x \equiv \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt, \quad (31)$$

where $\operatorname{erf} x$ is the error function. By (30) and (31), in this case the expression for the resolvent can be rewritten in the form

$$R(x-t) = \frac{d}{dx} \left\{ \exp[\lambda^2 \pi(x-t)] \operatorname{erf} [\lambda \sqrt{\pi(x-t)}] \right\}. \quad (32)$$

Applying relations (27) and (32), we obtain the solution of the Abel integral equation of the second kind (29) in the form

$$y(x) = f(x) + \frac{d}{dx} \int_a^x \left\{ \exp[\lambda^2 \pi(x-t)] \operatorname{erf} [\lambda \sqrt{\pi(x-t)}] \right\} f(t) dt, \quad x > a. \quad (33)$$

Note that in the case under consideration, the solution is constructed in the closed form.

11.3-6. Systems of Volterra Integral Equations.

The Laplace transform can be applied to solve systems of Volterra integral equations of the form

$$y_m(x) - \sum_{k=1}^n \int_0^x K_{mk}(x-t)y_k(t) dt = f_m(x), \quad m = 1, \dots, n. \quad (34)$$

Let us apply the Laplace transform to system (34). We obtain the relations

$$\tilde{y}_m(p) - \sum_{k=1}^n \tilde{K}_{mk}(p)\tilde{y}_k(p) = \tilde{f}_m(p), \quad m = 1, \dots, n. \quad (35)$$

On solving this system of linear algebraic equations, we find $\tilde{y}_m(p)$, and the solution of the system under consideration becomes

$$y_m(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \tilde{y}_m(p)e^{px} dp. \quad (36)$$

The Laplace transform can be applied to construct a solution of systems of Volterra equations of the first kind and of integro-differential equations as well.

⊙ References for Section 11.3: V. A. Ditkin and A. P. Prudnikov (1965), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), V. I. Smirnov (1974), K. B. Oldham and J. Spanier (1974), P. P. Zabreyko, A. I. Koshelev, et al. (1975), F. D. Gakhov and Yu. I. Cherskii (1978), Yu. I. Babenko (1986), R. Gorenflo and S. Vessella (1991), S. G. Samko, A. A. Kilbas, and O. I. Marichev (1993).

11.4. Operator Methods for Solving Linear Integral Equations

11.4-1. Application of a Solution of a "Truncated" Equation of the First Kind.

Consider the linear equation of the second kind

$$y(x) + \mathbf{L}[y] = f(x), \quad (1)$$

where \mathbf{L} is a linear (integral) operator.

Assume that the solution of the auxiliary “truncated” equation of the first kind,

$$\mathbf{L}[u] = g(x), \tag{2}$$

can be represented in the form

$$u(x) = \mathbf{M}[\mathbf{L}[g]], \tag{3}$$

where \mathbf{M} is a known linear operator. Formula (3) means that

$$\mathbf{L}^{-1} = \mathbf{M}\mathbf{L}.$$

Let us apply the operator \mathbf{L}^{-1} to Eq. (1). The resulting relation has the form

$$\mathbf{M}[\mathbf{L}[y]] + y(x) = \mathbf{M}[\mathbf{L}[f]]. \tag{4}$$

On eliminating $y(x)$ from (1) and (4) we obtain the equation

$$\mathbf{M}[w] - w(x) = F(x), \tag{5}$$

in which the following notation is used:

$$w = \mathbf{L}[y], \quad F(x) = \mathbf{M}[\mathbf{L}[f]] - f(x).$$

In some cases, Eq. (5) is simpler than the original equation (1). For example, this is the case if the operator \mathbf{M} is a constant (see Section 13.8) or a differential operator:

$$\mathbf{M} = a_n D^n + a_{n-1} D^{n-1} + \dots + a_1 D + a_0, \quad D \equiv \frac{d}{dx}.$$

In the latter case, Eq. (5) is an ordinary linear differential equation for the function w .

If a solution $w = w(x)$ of Eq. (5) is obtained, then a solution of Eq. (1) is given by the formula $y(x) = \mathbf{M}[\mathbf{L}[w]]$.

Example 1. Consider the Abel equation of the second kind

$$y(x) + \lambda \int_a^x \frac{y(t) dt}{\sqrt{x-t}} = f(x). \tag{6}$$

To solve this equation, we apply a slight modification of the above scheme, which corresponds to the case $\mathbf{M} \equiv \text{const } \frac{d}{dx}$.

Let us rewrite Eq. (6) as follows:

$$\int_a^x \frac{y(t) dt}{\sqrt{x-t}} = \frac{f(x) - y(x)}{\lambda}. \tag{7}$$

Let us assume that the right-hand side of Eq. (7) is known and treat Eq. (7) as an Abel equation of the first kind. Its solution can be written in the following form (see Example 3 in Subsection 10.4-4):

$$y(x) = \frac{1}{\pi} \frac{d}{dx} \int_a^x \frac{f(t) - y(t)}{\lambda \sqrt{x-t}} dt$$

or

$$y(x) + \frac{1}{\pi\lambda} \frac{d}{dx} \int_a^x \frac{y(t) dt}{\sqrt{x-t}} = \frac{1}{\pi\lambda} \frac{d}{dx} \int_a^x \frac{f(t) dt}{\sqrt{x-t}}. \tag{8}$$

Let us differentiate both sides of Eq. (6) with respect to x , multiply Eq. (8) by $-\pi\lambda^2$, and add the resulting expressions term by term. We eventually arrive at the following first-order linear ordinary differential equation for the function $y = y(x)$:

$$y'_x - \pi\lambda^2 y = F'_x(x), \tag{9}$$

where

$$F(x) = f(x) - \lambda \int_a^x \frac{f(t) dt}{\sqrt{x-t}}. \tag{10}$$

We must supplement Eq. (9) with initial condition

$$y(a) = f(a), \tag{11}$$

which is a consequence of (6).

The solution of problem (9)–(11) has the form

$$y(x) = F(x) + \pi\lambda^2 \int_a^x \exp[\pi\lambda^2(x-t)]F(t) dt, \tag{12}$$

and defines the solution of the Abel equation of the second kind (6).

11.4-2. Application of the Auxiliary Equation of the Second Kind.

The solution of the Abel equation of the second kind (6) can also be obtained by another method, presented below.

Consider the linear equation

$$y(x) - \mathbf{L}[y] = f(x), \quad (13)$$

where \mathbf{L} is a linear operator. Assume that the solution of the auxiliary equation

$$w(x) - \mathbf{L}^n[w] = \Phi(x), \quad \mathbf{L}^n[w] \equiv \mathbf{L}[\mathbf{L}^{n-1}[w]], \quad (14)$$

which involves the n th power of the operator \mathbf{L} , is known and is defined by the formula

$$w(x) = \mathbf{M}[\Phi(x)]. \quad (15)$$

In this case, the solution of the original equation (13) has the form

$$y(x) = \mathbf{M}[\Phi(x)], \quad \Phi(x) = \mathbf{L}^{n-1}[f] + \mathbf{L}^{n-2}[f] + \cdots + \mathbf{L}[f] + f(x). \quad (16)$$

This assertion can be proved by applying the operator $\mathbf{L}^{n-1} + \mathbf{L}^{n-2} + \cdots + \mathbf{L} + 1$ to Eq. (13), with regard to the operator relation

$$(1 - \mathbf{L})(\mathbf{L}^{n-1} + \mathbf{L}^{n-2} + \cdots + \mathbf{L} + 1) = 1 - \mathbf{L}^n$$

together with formula (16) for $\Phi(x)$. In Eq. (14) we may write $y(x)$ instead of $w(x)$.

Example 2. Let us apply the operator method (for $n = 2$) to solve the generalized Abel equation with exponent $3/4$:

$$y(x) - b \int_0^x \frac{y(t) dt}{(x-t)^{3/4}} = f(x). \quad (17)$$

We first consider the integral operator with difference kernel

$$\mathbf{L}[y(x)] \equiv \int_0^x K(x-t)y(t) dt.$$

Let us find \mathbf{L}^2 :

$$\begin{aligned} \mathbf{L}^2[y] &\equiv \mathbf{L}[\mathbf{L}[y]] = \int_0^x \int_0^t K(x-t)K(t-s)y(s) ds dt \\ &= \int_0^x y(s) ds \int_s^x K(x-t)K(t-s) dt = \int_0^x K_2(x-s)y(s) ds, \\ K_2(z) &= \int_0^z K(\xi)K(z-\xi) d\xi. \end{aligned} \quad (18)$$

In the proof of this formula, we have reversed the order of integration and performed the change of variables $\xi = t - s$.

For the power-law kernel

$$K(\xi) = b\xi^\mu,$$

we have

$$K_2(z) = b^2 \frac{\Gamma^2(1 + \mu)}{\Gamma(2 + 2\mu)} z^{1+2\mu}. \quad (19)$$

For Eq. (17) we obtain

$$\mu = -\frac{3}{4}, \quad K_2(z) = A \frac{1}{\sqrt{z}}, \quad A = \frac{b^2}{\sqrt{\pi}} \Gamma^2\left(\frac{1}{4}\right).$$

Therefore, the auxiliary equation (14) corresponding to $n = 2$ has the form

$$y(x) - A \int_0^x \frac{y(t) dt}{\sqrt{x-t}} = \Phi(x), \quad (20)$$

where

$$\Phi(x) = f(x) + b \int_0^x \frac{f(t) dt}{(x-t)^{3/4}}.$$

After the substitution $A \rightarrow -\lambda$ and $\Phi \rightarrow f$, relation (20) coincides with Eq. (6), and the solution of Eq. (20) can be obtained by formula (12).

Remark. It follows from (19) that the solution of the generalized Abel equation with exponent β

$$y(x) + \lambda \int_0^x \frac{y(t) dt}{(x-t)^\beta} = f(x)$$

can be reduced to the solution of a similar equation with the different exponent $\beta_1 = 2\beta - 1$. In particular, the Abel equation (6), which corresponds to $\beta = \frac{1}{2}$, is reduced to the solution of an equation with degenerate kernel for $\beta_1 = 0$.

11.4-3. Method for Solving "Quadratic" Operator Equations.

Suppose that the solution of the linear (integral, differential, etc.) equation

$$y(x) - \lambda \mathbf{L}[y] = f(x) \quad (21)$$

is known for an arbitrary right-hand side $f(x)$ and for any λ from the interval $(\lambda_{\min}, \lambda_{\max})$. We denote this solution by

$$y = Y(f, \lambda). \quad (22)$$

Let us construct the solution of the more complicated equation

$$y(x) - a\mathbf{L}[y] - b\mathbf{L}^2[y] = f(x), \quad (23)$$

where a and b are some numbers and $f(x)$ is an arbitrary function. To this end, we represent the left-hand side of Eq. (23) by the product of operators

$$(1 - a\mathbf{L} - b\mathbf{L}^2)[y] \equiv (1 - \lambda_1\mathbf{L})(1 - \lambda_2\mathbf{L})[y], \quad (24)$$

where λ_1 and λ_2 are the roots of the quadratic equation

$$\lambda^2 - a\lambda - b = 0. \quad (25)$$

We assume that $\lambda_{\min} < \lambda_1, \lambda_2 < \lambda_{\max}$.

Let us solve the auxiliary equation

$$w(x) - \lambda_2\mathbf{L}[w] = f(x), \quad (26)$$

which is the special case of Eq. (21) for $\lambda = \lambda_2$. The solution of this equation is given by the formula

$$w(x) = Y(f, \lambda_2). \quad (27)$$

Taking into account (24) and (26), we can rewrite Eq. (23) in the form

$$(1 - \lambda_1\mathbf{L})(1 - \lambda_2\mathbf{L})[y] = (1 - \lambda_2\mathbf{L})[w],$$

or, in view of the identity $(1 - \lambda_1\mathbf{L})(1 - \lambda_2\mathbf{L}) \equiv (1 - \lambda_2\mathbf{L})(1 - \lambda_1\mathbf{L})$, in the form

$$(1 - \lambda_2\mathbf{L}) \left\{ (1 - \lambda_1\mathbf{L})[y] - w(x) \right\} = 0.$$

This relation holds if the unknown function $y(x)$ satisfies the equation

$$y(x) - \lambda_1\mathbf{L}[y] = w(x). \quad (28)$$

The solution of this equation is given by the formula

$$y(x) = Y(w, \lambda_1), \quad \text{where } w = Y(f, \lambda_2). \quad (29)$$

If the homogeneous equation $y(x) - \lambda_2 \mathbf{L}[y] = 0$ has only the trivial* solution $y \equiv 0$, then formula (29) defines the unique solution of the original equation (23).

Example 3. Consider the integral equation

$$y(x) - \int_0^x \left(\frac{A}{\sqrt{x-t}} + B \right) y(t) dt = f(x).$$

It follows from the results of Example 2 that this equation can be written in the form of Eq. (23):

$$y(x) - A\mathbf{L}[y] - \frac{1}{\pi} B\mathbf{L}^2[y] = f(x), \quad \mathbf{L}[y] \equiv \int_0^x \frac{y(t) dt}{\sqrt{x-t}}.$$

Therefore, the solution (in the form of antiderivatives) of the integral equation can be given by the formulas

$$y(x) = Y(w, \lambda_1), \quad w = Y(f, \lambda_2), \\ Y(f, \lambda) = F(x) + \pi\lambda^2 \int_0^x \exp[\pi\lambda^2(x-t)] F(t) dt, \quad F(x) = f(x) + \lambda \int_0^x \frac{f(t) dt}{\sqrt{x-t}},$$

where λ_1 and λ_2 are the roots of the quadratic equation $\lambda^2 - A\lambda - \frac{1}{\pi}B = 0$.

This method can also be applied to solve (in the form of antiderivatives) more general equations of the form

$$y(x) - \int_0^x \left[\frac{A}{(x-t)^\beta} + \frac{B}{(x-t)^{2\beta-1}} \right] y(t) dt = f(x),$$

where β is a rational number satisfying the condition $0 < \beta < 1$ (see Example 2 and Eq. 2.1.60 from the first part of the book).

11.4-4. Solution of Operator Equations of Polynomial Form.

The method described in Subsection 11.4-3 can be generalized to the case of operator equations of polynomial form. Suppose that the solution of the linear nonhomogeneous equation (21) is given by formula (22) and that the corresponding homogeneous equation has only the trivial solution.

Let us construct the solution of the more complicated equation with polynomial left-hand side with respect to the operator \mathbf{L} :

$$y(x) - \sum_{k=1}^n A_k \mathbf{L}^k [y] = f(x), \quad \mathbf{L}^k \equiv \mathbf{L}(\mathbf{L}^{k-1}), \quad (30)$$

where A_k are some numbers and $f(x)$ is an arbitrary function.

We denote by $\lambda_1, \dots, \lambda_n$ the roots of the characteristic equation

$$\lambda^n - \sum_{k=1}^n A_k \lambda^{n-k} = 0. \quad (31)$$

The left-hand side of Eq. (30) can be expressed in the form of a product of operators:

$$y(x) - \sum_{k=1}^n A_k \mathbf{L}^k [y] \equiv \prod_{k=1}^n (1 - \lambda_k \mathbf{L}) [y]. \quad (32)$$

The solution of the auxiliary equation (26), in which we use the substitution $w \rightarrow y_{n-1}$ and $\lambda_2 \rightarrow \lambda_n$, is given by the formula $y_{n-1}(x) = Y(f, \lambda_n)$. Reasoning similar to that in Subsection 11.4-3 shows that the solution of Eq. (30) is reduced to the solution of the simpler equation

$$\prod_{k=1}^{n-1} (1 - \lambda_k \mathbf{L}) [y] = y_{n-1}(x), \quad (33)$$

* If the homogeneous equation $y(x) - \lambda_2 \mathbf{L}[y] = 0$ has nontrivial solutions, then the right-hand side of Eq. (28) must contain the function $w(x) + y_0(x)$ instead of $w(x)$, where y_0 is the general solution of the homogeneous equation.

whose degree is less by one than that of the original equation with respect to the operator \mathbf{L} . We can show in a similar way that Eq. (33) can be reduced to the solution of the simpler equation

$$\prod_{k=1}^{n-2} (1 - \lambda_k \mathbf{L})[y] = y_{n-2}(x), \quad y_{n-2}(x) = Y(y_{n-1}, \lambda_{n-1}).$$

Successively reducing the order of the equation, we eventually arrive at an equation of the form (28) whose right-hand side contains the function $y_1(x) = Y(y_2, \lambda_2)$. The solution of this equation is given by the formula $y(x) = Y(y_1, \lambda_1)$.

The solution of the original equation (30) is defined recursively by the following formulas:

$$y_{k-1}(x) = Y(y_k, \lambda_k); \quad k = n, \dots, 1, \quad \text{where } y_n(x) \equiv f(x), \quad y_0(x) \equiv y(x).$$

Note that here the decreasing sequence $k = n, \dots, 1$ is used.

11.4-5. Some Generalizations.

Suppose that the left-hand side of a linear (integral) equation

$$y(x) - \mathbf{Q}[y] = f(x) \tag{34}$$

can be represented in the form of a product

$$y(x) - \mathbf{Q}[y] \equiv \prod_{k=1}^n (1 - \mathbf{L}_k)[y], \tag{35}$$

where the \mathbf{L}_k are linear operators. Suppose that the solutions of the auxiliary equations

$$y(x) - \mathbf{L}_k[y] = f(x), \quad k = 1, \dots, n \tag{36}$$

are known and are given by the formulas

$$y(x) = Y_k[f(x)], \quad k = 1, \dots, n. \tag{37}$$

The solution of the auxiliary equation (36) for $k = n$, in which we apply the substitution $y \rightarrow y_{n-1}$, is given by the formula $y_{n-1}(x) = Y_n[f(x)]$. Reasoning similar to that used in Subsection 11.4-3 shows that the solution of Eq. (34) can be reduced to the solution of the simpler equation

$$\prod_{k=1}^{n-1} (1 - \mathbf{L}_k)[y] = y_{n-1}(x).$$

Successively reducing the order of the equation, we eventually arrive at an equation of the form (36) for $k = 1$, whose right-hand side contains the function $y_1(x) = Y_2[y_2(x)]$. The solution of this equation is given by the formula $y(x) = Y_1[y_1(x)]$.

The solution of the original equation (35) can be defined recursively by the following formulas:

$$y_{k-1}(x) = Y_k[y_k(x)]; \quad k = n, \dots, 1, \quad \text{where } y_n(x) \equiv f(x), \quad y_0(x) \equiv y(x).$$

Note that here the decreasing sequence $k = n, \dots, 1$ is used.

11.5. Construction of Solutions of Integral Equations with Special Right-Hand Side

In this section we describe some approaches to the construction of solutions of integral equations with special right-hand side. These approaches are based on the application of auxiliary solutions that depend on a free parameter.

11.5-1. General Scheme.

Consider a linear equation, which we shall write in the following brief form:

$$\mathbf{L}[y] = f_g(x, \lambda), \quad (1)$$

where \mathbf{L} is a linear operator (integral, differential, etc.) that acts with respect to the variable x and is independent of the parameter λ , and $f_g(x, \lambda)$ is a given function that depends on the variable x and the parameter λ .

Suppose that the solution of Eq. (1) is known:

$$y = y(x, \lambda). \quad (2)$$

Let \mathbf{M} be a linear operator (integral, differential, etc.) that acts with respect to the parameter λ and is independent of the variable x . Consider the (usual) case in which \mathbf{M} commutes with \mathbf{L} . We apply the operator \mathbf{M} to Eq. (1) and find that the equation

$$\mathbf{L}[w] = f_M(x), \quad f_M(x) = \mathbf{M}[f_g(x, \lambda)], \quad (3)$$

has the solution

$$w = \mathbf{M}[y(x, \lambda)]. \quad (4)$$

By choosing the operator \mathbf{M} in a different way, we can obtain solutions for other right-hand sides of Eq. (1). The original function $f_g(x, \lambda)$ is called the *generating function* for the operator \mathbf{L} .

11.5-2. Generating Function of Exponential Form.

Consider a linear equation with exponential right-hand side

$$\mathbf{L}[y] = e^{\lambda x}. \quad (5)$$

Suppose that the solution is known and is given by formula (2). In Table 6 we present solutions of the equation $\mathbf{L}[y] = f(x)$ with various right-hand sides; these solutions are expressed via the solution of Eq. (5).

Remark 1. When applying the formulas indicated in the table, we need not know the left-hand side of the linear equation (5) (the equation can be integral, differential, etc.) provided that a particular solution of this equation for exponential right-hand side is known. It is only of importance that the left-hand side of the equation is independent of the parameter λ .

Remark 2. When applying formulas indicated in the table, the convergence of the integrals occurring in the resulting solution must be verified.

TABLE 6
Solutions of the equation $\mathbf{L}[y] = f(x)$ with generating function of the exponential form

No	Right-Hand Side $f(x)$	Solution y	Solution Method
1	$e^{\lambda x}$	$y(x, \lambda)$	Original Equation
2	$A_1 e^{\lambda_1 x} + \dots + A_n e^{\lambda_n x}$	$A_1 y(x, \lambda_1) + \dots + A_n y(x, \lambda_n)$	Follows from linearity
3	$Ax + B$	$A \frac{\partial}{\partial \lambda} [y(x, \lambda)]_{\lambda=0} + B y(x, 0)$	Follows from linearity and the results of row No 4
4	$Ax^n,$ $n = 0, 1, 2, \dots$	$A \left\{ \frac{\partial^n}{\partial \lambda^n} [y(x, \lambda)] \right\}_{\lambda=0}$	Follows from the results of row No 6 for $\lambda = 0$
5	$\frac{A}{x+a}, \quad a > 0$	$A \int_0^\infty e^{-a\lambda} y(x, -\lambda) d\lambda$	Integration with respect to the parameter λ
6	$Ax^n e^{\lambda x},$ $n = 0, 1, 2, \dots$	$A \frac{\partial^n}{\partial \lambda^n} [y(x, \lambda)]$	Differentiation with respect to the parameter λ
7	a^x	$y(x, \ln a)$	Follows from row No 1
8	$A \cosh(\lambda x)$	$\frac{1}{2} A [y(x, \lambda) + y(x, -\lambda)]$	Linearity and relations to the exponential
9	$A \sinh(\lambda x)$	$\frac{1}{2} A [y(x, \lambda) - y(x, -\lambda)]$	Linearity and relations to the exponential
10	$Ax^m \cosh(\lambda x),$ $m = 1, 3, 5, \dots$	$\frac{1}{2} A \frac{\partial^m}{\partial \lambda^m} [y(x, \lambda) - y(x, -\lambda)]$	Differentiation with respect to λ and relation to the exponential
11	$Ax^m \cosh(\lambda x),$ $m = 2, 4, 6, \dots$	$\frac{1}{2} A \frac{\partial^m}{\partial \lambda^m} [y(x, \lambda) + y(x, -\lambda)]$	Differentiation with respect to λ and relation to the exponential
12	$Ax^m \sinh(\lambda x),$ $m = 1, 3, 5, \dots$	$\frac{1}{2} A \frac{\partial^m}{\partial \lambda^m} [y(x, \lambda) + y(x, -\lambda)]$	Differentiation with respect to λ and relation to the exponential
13	$Ax^m \sinh(\lambda x),$ $m = 2, 4, 6, \dots$	$\frac{1}{2} A \frac{\partial^m}{\partial \lambda^m} [y(x, \lambda) - y(x, -\lambda)]$	Differentiation with respect to λ and relation to the exponential
14	$A \cos(\beta x)$	$A \operatorname{Re} [y(x, i\beta)]$	Selection of the real part for $\lambda = i\beta$
15	$A \sin(\beta x)$	$A \operatorname{Im} [y(x, i\beta)]$	Selection of the imaginary part for $\lambda = i\beta$
16	$Ax^n \cos(\beta x),$ $n = 1, 2, 3, \dots$	$A \operatorname{Re} \left\{ \frac{\partial^n}{\partial \lambda^n} [y(x, \lambda)] \right\}_{\lambda=i\beta}$	Differentiation with respect to λ and selection of the real part for $\lambda = i\beta$
17	$Ax^n \sin(\beta x),$ $n = 1, 2, 3, \dots$	$A \operatorname{Im} \left\{ \frac{\partial^n}{\partial \lambda^n} [y(x, \lambda)] \right\}_{\lambda=i\beta}$	Differentiation with respect to λ and selection of the imaginary part for $\lambda = i\beta$
18	$Ae^{\mu x} \cos(\beta x)$	$A \operatorname{Re} [y(x, \mu + i\beta)]$	Selection of the real part for $\lambda = \mu + i\beta$
19	$Ae^{\mu x} \sin(\beta x)$	$A \operatorname{Im} [y(x, \mu + i\beta)]$	Selection of the imaginary part for $\lambda = \mu + i\beta$
20	$Ax^n e^{\mu x} \cos(\beta x),$ $n = 1, 2, 3, \dots$	$A \operatorname{Re} \left\{ \frac{\partial^n}{\partial \lambda^n} [y(x, \lambda)] \right\}_{\lambda=\mu+i\beta}$	Differentiation with respect to λ and selection of the real part for $\lambda = \mu + i\beta$
21	$Ax^n e^{\mu x} \sin(\beta x),$ $n = 1, 2, 3, \dots$	$A \operatorname{Im} \left\{ \frac{\partial^n}{\partial \lambda^n} [y(x, \lambda)] \right\}_{\lambda=\mu+i\beta}$	Differentiation with respect to λ and selection of the imaginary part for $\lambda = \mu + i\beta$

Example 1. We seek a solution of the equation with exponential right-hand side

$$y(x) + \int_x^\infty K(x-t)y(t) dt = e^{\lambda x} \quad (6)$$

in the form $y(x, \lambda) = ke^{\lambda x}$ by the method of indeterminate coefficients. Then we obtain

$$y(x, \lambda) = \frac{1}{B(\lambda)} e^{\lambda x}, \quad B(\lambda) = 1 + \int_0^\infty K(-z)e^{\lambda z} dz. \quad (7)$$

It follows from row 3 of Table 6 that the solution of the equation

$$y(x) + \int_x^\infty K(x-t)y(t) dt = Ax \quad (8)$$

has the form

$$y(x) = \frac{A}{D}x - \frac{AC}{D^2},$$

where

$$D = 1 + \int_0^\infty K(-z) dz, \quad C = \int_0^\infty zK(-z) dz.$$

For such a solution to exist, it is necessary that the improper integrals of the functions $K(-z)$ and $zK(-z)$ exist. This holds if the function $K(-z)$ decreases more rapidly than z^{-2} as $z \rightarrow \infty$. Otherwise a solution can be nonexistent. It is of interest that for functions $K(-z)$ with power-law growth as $z \rightarrow \infty$ in the case $\lambda < 0$, the solution of Eq. (6) exists and is given by formula (7), whereas Eq. (8) does not have a solution. Therefore, we must be careful when using formulas from Table 6 and verify the convergence of the integrals occurring in the solution.

It follows from row 15 of Table 6 that the solution of the equation

$$y(x) + \int_x^\infty K(x-t)y(t) dt = A \sin(\lambda x) \quad (9)$$

is given by the formula

$$y(x) = \frac{A}{B_c^2 + B_s^2} [B_c \sin(\lambda x) - B_s \cos(\lambda x)],$$

where

$$B_c = 1 + \int_0^\infty K(-z) \cos(\lambda z) dz, \quad B_s = \int_0^\infty K(-z) \sin(\lambda z) dz.$$

11.5-3. Power-Law Generating Function.

Consider the linear equation with power-law right-hand side

$$\mathbf{L}[y] = x^\lambda. \quad (10)$$

Suppose that the solution is known and is given by formula (2). In Table 7, solutions of the equation $\mathbf{L}[y] = f(x)$ with various right-hand sides are presented which can be expressed via the solution of Eq. (10).

Example 2. We seek a solution of the equation with power-law right-hand side

$$y(x) + \int_0^x \frac{1}{x} K\left(\frac{t}{x}\right) y(t) dt = x^\lambda$$

in the form $y(x, \lambda) = kx^\lambda$ by the method of indeterminate coefficients. We finally obtain

$$y(x, \lambda) = \frac{1}{1 + B(\lambda)} x^\lambda, \quad B(\lambda) = \int_0^1 K(t)t^\lambda dt.$$

It follows from row 3 of Table 7 that the solution of the equation with logarithmic right-hand side

$$y(x) + \int_0^x \frac{1}{x} K\left(\frac{t}{x}\right) y(t) dt = A \ln x$$

has the form

$$y(x) = \frac{A}{1 + I_0} \ln x - \frac{AI_1}{(1 + I_0)^2},$$

where

$$I_0 = \int_0^1 K(t) dt, \quad I_1 = \int_0^1 K(t) \ln t dt.$$

TABLE 7
Solutions of the equation $\mathbf{L}[y] = f(x)$ with generating function of power-law form

No	Right-Hand Side $f(x)$	Solution y	Solution Method
1	x^λ	$y(x, \lambda)$	Original Equation
2	$\sum_{k=0}^n A_k x^k$	$\sum_{k=0}^n A_k y(x, k)$	Follows from linearity
3	$A \ln x + B$	$A \frac{\partial}{\partial \lambda} \left[y(x, \lambda) \right]_{\lambda=0} + B y(x, 0)$	Follows from linearity and from the results of row No 4
4	$A \ln^n x,$ $n = 0, 1, 2, \dots$	$A \left\{ \frac{\partial^n}{\partial \lambda^n} \left[y(x, \lambda) \right] \right\}_{\lambda=0}$	Follows from the results of row No 5 for $\lambda = 0$
5	$A x^\lambda \ln^n x,$ $n = 0, 1, 2, \dots$	$A \frac{\partial^n}{\partial \lambda^n} \left[y(x, \lambda) \right]$	Differentiation with respect to the parameter λ
6	$A \cos(\beta \ln x)$	$A \operatorname{Re} \left[y(x, i\beta) \right]$	Selection of the real part for $\lambda = i\beta$
7	$A \sin(\beta \ln x)$	$A \operatorname{Im} \left[y(x, i\beta) \right]$	Selection of the imaginary part for $\lambda = i\beta$
8	$A x^\mu \cos(\beta \ln x)$	$A \operatorname{Re} \left[y(x, \mu + i\beta) \right]$	Selection of the real part for $\lambda = \mu + i\beta$
9	$A x^\mu \sin(\beta \ln x)$	$A \operatorname{Im} \left[y(x, \mu + i\beta) \right]$	Selection of the imaginary part for $\lambda = \mu + i\beta$

11.5-4. Generating Function Containing Sines and Cosines.

Consider the linear equation

$$\mathbf{L}[y] = \sin(\lambda x). \quad (11)$$

We assume that the solution of this equation is known and is given by formula (2). In Table 8, solutions of the equation $\mathbf{L}[y] = f(x)$ with various right-hand sides are given, which are expressed via the solution of Eq. (11).

Consider the linear equation

$$\mathbf{L}[y] = \cos(\lambda x). \quad (12)$$

We assume that the solution of this equation is known and is given by formula (2). In Table 9, solutions of the equation $\mathbf{L}[y] = f(x)$ with various right-hand sides are given, which are expressed via the solution of Eq. (12).

TABLE 8
Solutions of the equation $\mathbf{L}[y] = f(x)$ with sine-shaped generating function

No	Right-Hand Side $f(x)$	Solution y	Solution Method
1	$\sin(\lambda x)$	$y(x, \lambda)$	Original Equation
2	$\sum_{k=1}^n A_k \sin(\lambda_k x)$	$\sum_{k=1}^n A_k y(x, \lambda_k)$	Follows from linearity
3	$Ax^m,$ $m = 1, 3, 5, \dots$	$A(-1)^{\frac{m-1}{2}} \left[\frac{\partial^m}{\partial \lambda^m} y(x, \lambda) \right]_{\lambda=0}$	Follows from the results of row 5 for $\lambda = 0$
4	$Ax^m \sin(\lambda x),$ $m = 2, 4, 6, \dots$	$A(-1)^{\frac{m}{2}} \frac{\partial^m}{\partial \lambda^m} y(x, \lambda)$	Differentiation with respect to the parameter λ
5	$Ax^m \cos(\lambda x),$ $m = 1, 3, 5, \dots$	$A(-1)^{\frac{m-1}{2}} \frac{\partial^m}{\partial \lambda^m} y(x, \lambda)$	Differentiation with respect to the parameter λ
6	$\sinh(\beta x)$	$-iy(x, i\beta)$	Relation to the hyperbolic sine, $\lambda = i\beta$
7	$x^m \sinh(\beta x),$ $m = 2, 4, 6, \dots$	$i(-1)^{\frac{m+2}{2}} \left[\frac{\partial^m}{\partial \lambda^m} y(x, \lambda) \right]_{\lambda=i\beta}$	Differentiation with respect to λ and relation to the hyperbolic sine, $\lambda = i\beta$

TABLE 9
Solutions of the equation $\mathbf{L}[y] = f(x)$ with cosine-shaped generating function

No	Right-Hand Side $f(x)$	Solution y	Solution Method
1	$\cos(\lambda x)$	$y(x, \lambda)$	Original Equation
2	$\sum_{k=1}^n A_k \cos(\lambda_k x)$	$\sum_{k=1}^n A_k y(x, \lambda_k)$	Follows from linearity
3	$Ax^m,$ $m = 0, 2, 4, \dots$	$A(-1)^{\frac{m}{2}} \left[\frac{\partial^m}{\partial \lambda^m} y(x, \lambda) \right]_{\lambda=0}$	Follows from the results of row 4 for $\lambda = 0$
4	$Ax^m \cos(\lambda x),$ $m = 2, 4, 6, \dots$	$A(-1)^{\frac{m}{2}} \frac{\partial^m}{\partial \lambda^m} y(x, \lambda)$	Differentiation with respect to the parameter λ
5	$Ax^m \sin(\lambda x),$ $m = 1, 3, 5, \dots$	$A(-1)^{\frac{m+1}{2}} \frac{\partial^m}{\partial \lambda^m} y(x, \lambda)$	Differentiation with respect to the parameter λ
6	$\cosh(\beta x)$	$y(x, i\beta)$	Relation to the hyperbolic cosine, $\lambda = i\beta$
7	$x^m \cosh(\beta x),$ $m = 2, 4, 6, \dots$	$(-1)^{\frac{m}{2}} \left[\frac{\partial^m}{\partial \lambda^m} y(x, \lambda) \right]_{\lambda=i\beta}$	Differentiation with respect to λ and relation to the hyperbolic cosine, $\lambda = i\beta$

11.6. Method of Model Solutions

11.6-1. Preliminary Remarks*.

Consider a linear equation, which we briefly write out in the form

$$\mathbf{L}[y(x)] = f(x), \tag{1}$$

where \mathbf{L} is a linear (integral) operator, $y(x)$ is an unknown function, and $f(x)$ is a known function.

We first define arbitrarily a test solution

$$y_0 = y_0(x, \lambda), \tag{2}$$

* Before reading this section, it is useful to look over Section 11.5.

which depends on an auxiliary parameter λ (it is assumed that the operator \mathbf{L} is independent of λ and $y_0 \neq \text{const}$). By means of Eq. (1) we define the right-hand side that corresponds to the test solution (2):

$$f_0(x, \lambda) = \mathbf{L}[y_0(x, \lambda)].$$

Let us multiply Eq. (1), for $y = y_0$ and $f = f_0$, by some function $\varphi(\lambda)$ and integrate the resulting relation with respect to λ over an interval $[a, b]$. We finally obtain

$$\mathbf{L}[y_\varphi(x)] = f_\varphi(x), \quad (3)$$

where

$$y_\varphi(x) = \int_a^b y_0(x, \lambda)\varphi(\lambda) d\lambda, \quad f_\varphi(x) = \int_a^b f_0(x, \lambda)\varphi(\lambda) d\lambda. \quad (4)$$

It follows from formulas (3) and (4) that, for the right-hand side $f = f_\varphi(x)$, the function $y = y_\varphi(x)$ is a solution of the original equation (1). Since the choice of the function $\varphi(\lambda)$ (as well as of the integration interval) is arbitrary, the function $f_\varphi(x)$ can be arbitrary in principle. Here the main problem is how to choose a function $\varphi(\lambda)$ to obtain a given function $f_\varphi(x)$. This problem can be solved if we can find a test solution such that the right-hand side of Eq. (1) is the kernel of a known inverse integral transform (we denote such a test solution by $Y(x, \lambda)$ and call it a *model solution*).

11.6-2. Description of the Method.

Indeed, let \mathfrak{P} be an invertible integral transform that takes each function $f(x)$ to the corresponding transform $F(\lambda)$ by the rule

$$F(\lambda) = \mathfrak{P}\{f(x)\}. \quad (5)$$

Assume that the inverse transform \mathfrak{P}^{-1} has the kernel $\psi(x, \lambda)$ and acts as follows:

$$\mathfrak{P}^{-1}\{F(\lambda)\} = f(x), \quad \mathfrak{P}^{-1}\{F(\lambda)\} \equiv \int_a^b F(\lambda)\psi(x, \lambda) d\lambda. \quad (6)$$

The limits of integration a and b and the integration path in (6) may well lie in the complex plane.

Suppose that we succeeded in finding a model solution $Y(x, \lambda)$ of the auxiliary problem for Eq. (1) whose right-hand side is the kernel of the inverse transform \mathfrak{P}^{-1} :

$$\mathbf{L}[Y(x, \lambda)] = \psi(x, \lambda). \quad (7)$$

Let us multiply Eq. (7) by $F(\lambda)$ and integrate with respect to λ within the same limits that stand in the inverse transform (6). Taking into account the fact that the operator \mathbf{L} is independent of λ and applying the relation $\mathfrak{P}^{-1}\{F(\lambda)\} = f(x)$, we obtain

$$\mathbf{L}\left[\int_a^b Y(x, \lambda)F(\lambda) d\lambda\right] = f(x).$$

Therefore, the solution of Eq. (1) for an arbitrary function $f(x)$ on the right-hand side is expressed via a solution of the simpler auxiliary equation (7) by the formula

$$y(x) = \int_a^b Y(x, \lambda)F(\lambda) d\lambda, \quad (8)$$

where $F(\lambda)$ is the transform (5) of the function $f(x)$.

For the right-hand side of the auxiliary equation (7) we can take, for instance, exponential, power-law, and trigonometric function, which are the kernels of the Laplace, Mellin, and sine and cosine Fourier transforms (up to a constant factor). Sometimes it is rather easy to find a model solution by means of the method of indeterminate coefficients (by prescribing its structure). Afterwards, to construct a solution of the equation with arbitrary right-hand side, we can apply formulas written out below in Subsections 11.6-3–11.6-6.

11.6-3. Model Solution in the Case of an Exponential Right-Hand Side.

Assume that we have found a model solution $Y = Y(x, \lambda)$ that corresponds to the exponential right-hand side:

$$\mathbf{L}[Y(x, \lambda)] = e^{\lambda x}. \quad (9)$$

Consider two cases:

1°. *Equations on the semiaxis*, $0 \leq x < \infty$. Let $\tilde{f}(p)$ be the Laplace transform of the function $f(x)$:

$$\tilde{f}(p) = \mathfrak{L}\{f(x)\}, \quad \mathfrak{L}\{f(x)\} \equiv \int_0^{\infty} f(x)e^{-px} dx. \quad (10)$$

The solution of Eq. (1) for an arbitrary right-hand side $f(x)$ can be expressed via the solution of the simpler auxiliary equation with exponential right-hand side (9) for $\lambda = p$ by the formula

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} Y(x, p)\tilde{f}(p) dp. \quad (11)$$

2°. *Equations on the entire axis*, $-\infty < x < \infty$. Let $\tilde{f}(u)$ denote the Fourier transform of the function $f(x)$:

$$\tilde{f}(u) = \mathfrak{F}\{f(x)\}, \quad \mathfrak{F}\{f(x)\} \equiv \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x)e^{-iux} dx. \quad (12)$$

The solution of Eq. (1) for an arbitrary right-hand side $f(x)$ can be expressed via the solution of the simpler auxiliary equation with exponential right-hand side (9) for $\lambda = iu$ by the formula

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} Y(x, iu)\tilde{f}(u) du. \quad (13)$$

In the calculation of the integrals on the right-hand sides in (11) and (13), methods of the theory of functions of a complex variable are applied, including the Cauchy residue theorem and the Jordan lemma (see Subsections 9.1-4 and 9.1-5).

Remark 1. The structure of a model solution $Y(x, \lambda)$ can differ from that of the kernel of the Laplace or Fourier inversion formula.

Remark 2. When applying the method under consideration, the left-hand side of Eq. (1) need not be known (the equation can be integral, differential, functional, etc.) if a particular solution of this equation is known for the exponential right-hand side. Here only the most general information is important, namely, that the equation is linear, and its left-hand side is independent of the parameter λ .

Remark 3. The above method can be used in the solution of linear integral (differential, integro-differential, and functional) equations with composed argument of the unknown function.

Example 1. Consider the following Volterra equation of the second kind with difference kernel:

$$y(x) + \int_x^{\infty} K(x-t)y(t) dt = f(x). \quad (14)$$

This equation cannot be solved by direct application of the Laplace transform because the convolution theorem cannot be used here.

In accordance with the method of model solutions, we consider the auxiliary equation with exponential right-hand side

$$y(x) + \int_x^{\infty} K(x-t)y(t) dt = e^{px}. \quad (15)$$

Its solution has the form (see Example 1 of Section 11.5)

$$Y(x, p) = \frac{1}{1 + \tilde{K}(-p)} e^{px}, \quad \tilde{K}(-p) = \int_0^{\infty} K(-z)e^{pz} dz. \quad (16)$$

This, by means of formula (11), yields a solution of Eq. (14) for an arbitrary right-hand side,

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\tilde{f}(p)}{1 + \tilde{K}(-p)} e^{px} dp, \quad (17)$$

where $\tilde{f}(p)$ is the Laplace transform (10) of the function $f(x)$ (see also Section 11.11).

Note that a solution to Eq. (12) was obtained in the book of M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971) in a more complicated way.

11.6-4. Model Solution in the Case of a Power-Law Right-Hand Side.
--

Suppose that we have succeeded in finding a model solution $Y = Y(x, s)$ that corresponds to a power-law right-hand side of the equation:

$$\mathbf{L}[Y(x, s)] = x^{-s}, \quad \lambda = -s. \quad (18)$$

Let $\hat{f}(s)$ be the Mellin transform of the function $f(x)$:

$$\hat{f}(s) = \mathfrak{M}\{f(x)\}, \quad \mathfrak{M}\{f(x)\} \equiv \int_0^\infty f(x)x^{s-1} dx. \quad (19)$$

The solution of Eq. (1) for an arbitrary right-hand side $f(x)$ can be expressed via the solution of the simpler auxiliary equation with power-law right-hand side (18) by the formula

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} Y(x, s)\hat{f}(s) ds. \quad (20)$$

In the calculation of the corresponding integrals on the right-hand side of formula (20), one can use tables of inverse Mellin transforms (e.g., see Supplement 10), as well as methods of the theory of functions of a complex variable, including the Cauchy residue theorem and the Jordan lemma (see Subsections 9.1-4 and 9.1-5).

Example 2. Consider the equation

$$y(x) + \int_0^x \frac{1}{x} K\left(\frac{t}{x}\right)y(t) dt = f(x). \quad (21)$$

In accordance with the method of model solutions, we consider the following auxiliary equation with power-law right-hand side:

$$y(x) + \int_0^x \frac{1}{x} K\left(\frac{t}{x}\right)y(t) dt = x^{-s}. \quad (22)$$

Its solution has the form (see Example 2 for $\lambda = -s$ in Section 11.5)

$$Y(x, s) = \frac{1}{1+B(s)}x^{-s}, \quad B(s) = \int_0^1 K(t)t^{-s} dt. \quad (23)$$

This, by means of formula (20), yields the solution of Eq. (21) for an arbitrary right-hand side:

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\hat{f}(s)}{1+B(s)}x^{-s} ds, \quad (24)$$

where $\hat{f}(s)$ is the Mellin transform (19) of the function $f(x)$.

11.6-5. Model Solution in the Case of a Sine-Shaped Right-Hand Side.
--

Suppose that we have succeeded in finding a model solution $Y = Y(x, u)$ that corresponds to the sine on the right-hand side:

$$\mathbf{L}[Y(x, u)] = \sin(ux), \quad \lambda = u. \quad (25)$$

Let $\check{f}_s(u)$ be the asymmetric sine Fourier transform of the function $f(x)$:

$$\check{f}_s(u) = \mathcal{F}_s\{f(x)\}, \quad \mathcal{F}_s\{f(x)\} \equiv \int_0^\infty f(x) \sin(ux) dx. \quad (26)$$

The solution of Eq. (1) for an arbitrary right-hand side $f(x)$ can be expressed via the solution of the simpler auxiliary equation with sine-shape right-hand side (25) by the formula

$$y(x) = \frac{2}{\pi} \int_0^\infty Y(x, u)\check{f}_s(u) du. \quad (27)$$

11.6-6. Model Solution in the Case of a Cosine-Shaped Right-Hand Side.

Suppose that we have succeeded in finding a model solution $Y = Y(x, u)$ that corresponds to the cosine on the right-hand side:

$$\mathbf{L}[Y(x, u)] = \cos(ux), \quad \lambda = u. \quad (28)$$

Let $\check{f}_c(u)$ be the asymmetric Fourier cosine transform of the function $f(x)$:

$$\check{f}_c(u) = \mathcal{F}_c\{f(x)\}, \quad \mathcal{F}_c\{f(x)\} \equiv \int_0^\infty f(x) \cos(ux) dx. \quad (29)$$

The solution of Eq. (1) for an arbitrary right-hand side $f(x)$ can be expressed via the solution of the simpler auxiliary equation with cosine right-hand side (28) by the formula

$$y(x) = \frac{2}{\pi} \int_0^\infty Y(x, u) \check{f}_c(u) du. \quad (30)$$

11.6-7. Some Generalizations.

Just as above we assume that \mathfrak{B} is an invertible transform taking each function $f(x)$ to the corresponding transform $F(\lambda)$ by the rule (5) and that the inverse transform is defined by formula (6).

Suppose that we have succeeded in finding a model solution $Y(x, \lambda)$ of the following auxiliary problem for Eq. (1):

$$\mathbf{L}_x [Y(x, \lambda)] = \mathbf{H}_\lambda [\psi(x, \lambda)]. \quad (31)$$

The right-hand side of Eq. (31) contains an invertible linear operator (which is integral, differential, or functional) that is independent of the variable x and acts with respect to the parameter λ on the kernel $\psi(x, \lambda)$ of the inverse transform, see formula (6). For clarity, the operator on the left-hand side of Eq. (31) is labeled by the subscript x (it acts with respect to the variable x and is independent of λ).

Let us apply the inverse operator \mathbf{H}_λ^{-1} to Eq. (31). As a result, we obtain the kernel $\psi(x, \lambda)$ on the right-hand side. On the left-hand side we intertwine the operators by the rule $\mathbf{H}_\lambda^{-1} \mathbf{L}_x = \mathbf{L}_x \mathbf{H}_\lambda^{-1}$ (this is as a rule possible because the operators act with respect to different variables). Furthermore, let us multiply the resulting relation by $F(\lambda)$ and integrate with respect to λ within the limits that stand in the inverse transform (6). Taking into account the relation $\mathfrak{B}^{-1}\{F(\lambda)\} = f(x)$, we finally obtain

$$\mathbf{L}_x \left[\int_a^b F(\lambda) \mathbf{H}_\lambda^{-1} [Y(x, \lambda)] d\lambda \right] = f(x). \quad (32)$$

Hence, a solution of Eq. (1) with an arbitrary function $f(x)$ on the right-hand side can be expressed via the solution of the simpler auxiliary equation (31) by the formula

$$y(x) = \int_a^b F(\lambda) \mathbf{H}_\lambda^{-1} [Y(x, \lambda)] d\lambda, \quad (33)$$

where $F(\lambda)$ is the transform of the function $f(x)$ obtained by means of the transform \mathfrak{B} (5).

Since the choice of the operator \mathbf{H}_λ is arbitrary, this approach extends the abilities of the method of model solutions.

11.7. Method of Differentiation for Integral Equations

In some cases, the differentiation of integral equations (once, twice, and so on) with the subsequent elimination of integral terms by means of the original equation makes it possible to reduce a given equation to an ordinary differential equation. Sometimes by differentiating we can reduce a given equation to a simpler integral equation whose solution is known. Below we list some classes of integral equations that can be reduced to ordinary differential equations with constant coefficients.

11.7-1. Equations with Kernel Containing a Sum of Exponential Functions.

Consider the equation

$$y(x) + \int_a^x \left[\sum_{k=1}^n A_k e^{\lambda_k(x-t)} \right] y(t) dt = f(x). \quad (1)$$

In the general case, this equation can be reduced to a linear nonhomogeneous ordinary differential equation of n th order with constant coefficients (see equation 2.2.19 of the first part of the book).

In a wide range of the parameters A_k and λ_k , the solution can be represented as follows:

$$y(x) = f(x) + \int_a^x \left[\sum_{k=1}^n B_k e^{\mu_k(x-t)} \right] f(t) dt, \quad (2)$$

where the parameters B_k and μ_k of the solution are related to the parameters A_k and λ_k of the equation by algebraic relations.

For the solution of Eq. (1) with $n = 2$, see Section 2.2 of the first part of the book (equation 2.2.10).

11.7-2. Equations with Kernel Containing a Sum of Hyperbolic Functions.

By means of the formulas $\cosh \beta = \frac{1}{2}(e^\beta + e^{-\beta})$ and $\sinh \beta = \frac{1}{2}(e^\beta - e^{-\beta})$, any equation with difference kernel of the form

$$y(x) + \int_a^x K(x-t)y(t) dt = f(x), \quad (3)$$

$$K(x) = \sum_{k=1}^m A_k \cosh(\lambda_k x) + \sum_{k=1}^s B_k \sinh(\mu_k x),$$

can be represented in the form of Eq. (1) with $n = 2m + 2s$, and hence these equations can be reduced to linear nonhomogeneous ordinary differential equations with constant coefficients.

11.7-3. Equations with Kernel Containing a Sum of Trigonometric Functions.

Equations with difference kernel of the form

$$y(x) + \int_a^x K(x-t)y(t) dt = f(x), \quad K(x) = \sum_{k=1}^m A_k \cos(\lambda_k x), \quad (4)$$

$$y(x) + \int_a^x K(x-t)y(t) dt = f(x), \quad K(x) = \sum_{k=1}^m A_k \sin(\lambda_k x), \quad (5)$$

can also be reduced to linear nonhomogeneous ordinary differential equations of order $2m$ with constant coefficients (see equations 2.5.4 and 2.5.19 in the first part of the book).

In a wide range of the parameters A_k and λ_k , the solution of Eq. (5) can be represented in the form

$$y(x) = f(x) + \int_a^x R(x-t)f(t) dt, \quad R(x) = \sum_{k=1}^m B_k \sin(\mu_k x), \quad (6)$$

where the parameters B_k and μ_k of the solution are related to the parameters A_k and λ_k of the equation by algebraic relations.

Equations with difference kernels containing both cosines and sines can also be reduced to linear nonhomogeneous ordinary differential equations with constant coefficients.

11.7-4. Equations Whose Kernels Contain Combinations of Various Functions.

Any equation with difference kernel that contains a linear combination of summands of the form

$$(x-t)^m \quad (m = 0, 1, 2, \dots), \quad \exp[\alpha(x-t)], \quad (7)$$

$$\cosh[\beta(x-t)], \quad \sinh[\gamma(x-t)], \quad \cos[\lambda(x-t)], \quad \sin[\mu(x-t)],$$

can also be reduced by differentiation to a linear nonhomogeneous ordinary differential equation with constant coefficients, where exponential, hyperbolic, and trigonometric functions can also be multiplied by $(x-t)^n$ ($n = 1, 2, \dots$).

Remark. The method of differentiation can be successfully used to solve more complicated equations with nondifference kernel to which the Laplace transform cannot be applied (see, for instance, Eqs. 2.9.5, 2.9.28, 2.9.30, 2.9.34, and 2.9.36 in the first part of the book).

11.8. Reduction of Volterra Equations of the Second Kind to Volterra Equations of the First Kind

The Volterra equation of the second kind

$$y(x) - \int_a^x K(x,t)y(t) dt = f(x) \quad (1)$$

can be reduced to a Volterra equation of the first kind in two ways.

11.8-1. First Method.

We integrate Eq. (1) with respect to x from a to x and then reverse the order of integration in the double integral. We finally obtain the Volterra equation of the first kind

$$\int_a^x M(x,t)y(t) dt = F(x), \quad (2)$$

where $M(x,t)$ and $F(x)$ are defined as follows:

$$M(x,t) = 1 - \int_t^x K(s,t) ds, \quad F(x) = \int_a^x f(t) dt. \quad (3)$$

11.8-2. Second Method.

Assume that the condition $f(a) = 0$ is satisfied. In this case Eq. (1) can be reduced to a Volterra equation of the first kind for the derivative of the unknown function,

$$\int_a^x N(x, t)y'_t(t) dt = f(x), \quad y(a) = 0, \quad (4)$$

where

$$N(x, t) = 1 - \int_t^x K(x, s) ds. \quad (5)$$

Indeed, on integrating by parts the right-hand side of formula (4) with regard to formula (5), we arrive at Eq. (1).

Remark. For $f(a) \neq 0$, Eq. (1) implies the relation $y(a) = f(a)$. In this case the substitution $z(x) = y(x) - f(a)$ yields the Volterra equation of the second kind

$$z(x) - \int_a^x K(x, t)z(t) dt = \Phi(x),$$

$$\Phi(x) = f(x) - f(a) + f(a) \int_a^x K(x, t) dt,$$

whose right-hand side satisfies the condition $\Phi(a) = 0$, and hence this equation can be reduced by the second method to a Volterra equation of the first kind.

⊙ References for Section 11.8: V. Volterra (1959), A. F. Verlan' and V. S. Sizikov (1986).

11.9. Successive Approximation Method

11.9-1. General Scheme.

1°. Consider a Volterra integral equation of the second kind

$$y(x) - \int_a^x K(x, t)y(t) dt = f(x). \quad (1)$$

Assume that $f(x)$ is continuous on the interval $[a, b]$ and the kernel $K(x, t)$ is continuous for $a \leq x \leq b$ and $a \leq t \leq x$.

Let us seek the solution by the successive approximation method. To this end, we set

$$y(x) = f(x) + \sum_{n=1}^{\infty} \varphi_n(x), \quad (2)$$

where the $\varphi_n(x)$ are determined by the formulas

$$\varphi_1(x) = \int_a^x K(x, t)f(t) dt,$$

$$\varphi_2(x) = \int_a^x K(x, t)\varphi_1(t) dt = \int_a^x K_2(x, t)f(t) dt,$$

$$\varphi_3(x) = \int_a^x K(x, t)\varphi_2(t) dt = \int_a^x K_3(x, t)f(t) dt, \quad \text{etc.}$$

Here

$$K_n(x, t) = \int_a^x K(x, z)K_{n-1}(z, t) dz, \quad (3)$$

where $n = 2, 3, \dots$, and we have the relations $K_1(x, t) \equiv K(x, t)$ and $K_n(x, t) = 0$ for $t > x$. The functions $K_n(x, t)$ given by formulas (3) are called *iterated kernels*. These kernels satisfy the relation

$$K_n(x, t) = \int_a^x K_m(x, s)K_{n-m}(s, t) ds, \quad (4)$$

where m is an arbitrary positive integer less than n .

2°. The successive approximations can be implemented in a more general scheme:

$$y_n(x) = f(x) + \int_a^x K(x, t)y_{n-1}(t) dt, \quad n = 1, 2, \dots, \quad (5)$$

where the function $y_0(x)$ is continuous on the interval $[a, b]$. The functions $y_1(x), y_2(x), \dots$ which are obtained from (5) are also continuous on $[a, b]$.

Under the assumptions adopted in item 1° for $f(x)$ and $K(x, t)$, the sequence $\{y_n(x)\}$ converges, as $n \rightarrow \infty$, to the continuous solution $y(x)$ of the integral equation. A successful choice of the “zeroth” approximation $y_0(x)$ can result in a rapid convergence of the procedure.

Note that in the special case $y_0(x) = f(x)$, this method becomes that described in item 1°.

Remark 1. If the kernel $K(x, t)$ is square integrable on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ and $f(x) \in L_2(a, b)$, then the successive approximations are mean-square convergent to the solution $y(x) \in L_2(a, b)$ of the integral equation (1) for any initial approximation $y_0(x) \in L_2(a, b)$.

Example. Consider the integral equation

$$y(x) + \int_0^x (x-t)y(t) dt = 1$$

and use the method of successive approximations for finding its solution. To that end, we take $f(x) = 1$, $K(x, t) = -(x-t)$ in (5) and choose the initial function $y_0(x) = 0$. As a result, we get

$$\begin{aligned} y_1(x) &= 1, \\ y_2(x) &= 1 - \frac{x^2}{2!}, \\ &\dots\dots\dots, \\ y_n(x) &= 1 - \frac{x^2}{2!} + \dots + (-1)^{n-1} \frac{x^{2n-2}}{(2n-2)!}. \end{aligned}$$

It follows that

$$y(x) = \lim_{n \rightarrow \infty} y_n(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots = \cos x.$$

It is easy to check that $y(x) = \cos x$ is an exact solution of the integral equation under consideration.

11.9-2. Formula for the Resolvent.

The resolvent of the integral equation (1) is determined via the iterated kernels by the formula

$$R(x, t) = \sum_{n=1}^{\infty} K_n(x, t), \quad (6)$$

where the convergent series on the right-hand side is called the *Neumann series* of the kernel $K(x, t)$. Now the solution of the Volterra equation of the second kind (1) can be rewritten in the traditional form

$$y(x) = f(x) + \int_a^x R(x, t)f(t) dt. \quad (7)$$

Remark 2. In the case of a kernel with weak singularity, the solution of Eq. (1) can be obtained by the successive approximation method. In this case the kernels $K_n(x, t)$ are continuous starting from some n . For $\alpha < \frac{1}{2}$, even the kernel $K_2(x, t)$ is continuous.

© References for Section 11.9: W. V. Lovitt (1950), V. Volterra (1959), S. G. Mikhlin (1960), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), V. I. Smirnov (1974).

11.10. Method of Quadratures

11.10-1. General Scheme of the Method.

Let us consider the linear Volterra integral equation of the second kind

$$y(x) - \int_a^x K(x, t)y(t) dt = f(x), \quad (1)$$

on an interval $a \leq x \leq b$. Assume that the kernel and the right-hand side of the equation are continuous functions.

From Eq. (1) we find that $y(a) = f(a)$. Let us choose a constant integration step h and consider the discrete set of points $x_i = a + h(i - 1)$, $i = 1, \dots, n$. For $x = x_i$, Eq. (1) acquires the form

$$y(x_i) - \int_a^{x_i} K(x_i, t)y(t) dt = f(x_i), \quad i = 1, \dots, n. \quad (2)$$

Applying the quadrature formula (see Subsection 10.7-1) to the integral in (2) and choosing x_j ($j = 1, \dots, i$) to be the nodes in t , we arrive at the system of equations

$$y(x_i) - \sum_{j=1}^i A_{ij}K(x_i, x_j)y(x_j) = f(x_i) + \varepsilon_i[y], \quad i = 2, \dots, n, \quad (3)$$

where $\varepsilon_i[y]$ is the truncation error and A_{ij} are the coefficients of the quadrature formula on the interval $[a, x_i]$ (see Subsection 10.7-1). Suppose that $\varepsilon_i[y]$ are small and neglect them; then we obtain a system of linear algebraic equations in the form

$$y_1 = f_1, \quad y_i - \sum_{j=1}^i A_{ij}K_{ij}y_j = f_i, \quad i = 2, \dots, n, \quad (4)$$

where $K_{ij} = K(x_i, x_j)$, $f_i = f(x_i)$, and y_i are approximate values of the unknown function $y(x)$ at the nodes x_i .

From (4) we obtain the recurrent formula

$$y_1 = f_1, \quad y_i = \frac{f_i + \sum_{j=1}^{i-1} A_{ij}K_{ij}y_j}{1 - A_{ii}K_{ii}}, \quad i = 2, \dots, n, \quad (5)$$

valid under the condition

$$1 - A_{ii}K_{ii} \neq 0, \quad (6)$$

which can always be ensured by an appropriate choice of the nodes and by guaranteeing that the coefficients A_{ii} are sufficiently small.

11.10-2. Application of the Trapezoidal Rule.

According to the trapezoidal rule (see Subsection 10.7-1), we have

$$A_{i1} = A_{ii} = \frac{1}{2}h, \quad A_{i2} = \dots = A_{i,i-1} = h, \quad i = 2, \dots, n.$$

The application of the trapezoidal rule in the general scheme leads to the following step algorithm:

$$y_1 = f_1, \quad y_i = \frac{f_i + h \sum_{j=1}^{i-1} \beta_j K_{ij} y_j}{1 - \frac{1}{2}h K_{ii}}, \quad i = 2, \dots, n,$$

$$x_i = a + (i-1)h, \quad n = \frac{b-a}{h} + 1, \quad \beta_j = \begin{cases} \frac{1}{2} & \text{for } j = 1, \\ 1 & \text{for } j > 1, \end{cases}$$

where the notation coincides with that introduced in Subsection 11.10-1. The trapezoidal rule is quite simple and effective, and frequently used in practice. Some peculiarities of using the quadrature method for solving integral equations with variable limits of integration are indicated in Subsection 10.7-3.

11.10-3. Case of a Degenerate Kernel.

When solving a Volterra integral equation of the second kind with arbitrary kernel, the amount of calculations increases as the index of the integration step increases. However, if the kernel is degenerate, then it is possible to construct algorithms with a constant amount of calculations at each step. Indeed, for a degenerate kernel

$$K(x, t) = \sum_{k=1}^m p_k(x) q_k(t),$$

we can rewrite Eq. (1) in the form

$$y(x) = \sum_{k=1}^m p_k(x) \int_a^x q_k(t) y(t) dt + f(x).$$

The application of the trapezoidal rule makes it possible to obtain the following recurrent expression (see Subsection 11.10-2):

$$y_1 = f_1, \quad y_i = \frac{f_i + h \sum_{k=1}^m p_{ki} \sum_{j=1}^{i-1} \beta_j q_{kj} y_j}{1 - \frac{1}{2}h \sum_{k=1}^m p_{ki} q_{ki}},$$

where y_i are approximate values of the unknown function $y(x)$ at the nodes x_i , $f_i = f(x_i)$, $p_{ki} = p_k(x_i)$, and $q_{ki} = q_k(x_i)$, and this expression shows that the amount of calculations is the same at each step.

⊙ References for Section 11.10: S. G. Mikhailin and K. L. Smolitskiy (1967), G. A. Korn and T. M. Korn (1968), V. I. Krylov, V. V. Bobkov, and P. I. Monastyrnyi (1984), A. F. Verlan' and V. S. Sizikov (1986), H. Brunner (2004).

11.11. Equations with Infinite Integration Limit

Integral equations of the second kind with difference kernel and with a variable limit of integration for which the other limit is infinite are also of interest. Kernels and functions in such equations need not belong to the classes described in the beginning of the chapter. In this case their investigation can be performed by the method of model solutions (see Section 11.6) or by the reduction to equations of convolution type. We consider the latter method by an example of an equation of the second kind with variable lower limit.

11.11-1. Equation of the Second Kind with Variable Lower Integration Limit.

Integral equations of the second kind with variable lower limit, in the case of a difference kernel, have the form

$$y(x) + \int_x^\infty K(x-t)y(t) dt = f(x), \quad 0 < x < \infty. \quad (1)$$

This equation substantially differs from Volterra equations of the second kind studied above for which a solution exists and is unique. A solution of the corresponding homogeneous equation

$$y(x) + \int_x^\infty K(x-t)y(t) dt = 0 \quad (2)$$

can be nontrivial.

The eigenfunctions of the integral equation (2) are determined by the roots of the following transcendental (or algebraic) equation for the parameter λ :

$$\int_0^\infty K(-z)e^{-\lambda z} dz = -1. \quad (3)$$

The left-hand side of this equation is the Laplace transform of the function $K(-z)$ with parameter λ .

To a real simple root λ_k of Eq. (3) there corresponds an eigenfunction

$$y_k(x) = \exp(-\lambda_k x).$$

The general solution is the linear combination (with arbitrary constants) of the eigenfunctions of the homogeneous integral equation (2).

For solutions of Eq. (2) in the case of multiple or complex roots, see equation 52 in Section 2.9 (see also Example 1 below).

The general solution of the integral equation (1) is the sum of the general solution of the homogeneous equation (2) and a particular solution of the nonhomogeneous equation (1).

Example 1. Consider the homogeneous Picard–Goursat equation

$$y(x) + A \int_x^\infty (t-x)^n y(t) dt = 0, \quad n = 0, 1, 2, \dots, \quad (4)$$

which is a special case of Eq. (1) with $K(z) = A(-z)^n$.

The general solution of the homogeneous equation has the form

$$y(x) = \sum_{k=1}^m C_k \exp(-\lambda_k x), \quad (5)$$

where C_k are arbitrary constants and λ_k are the roots of the algebraic equation

$$\lambda^{n+1} + An! = 0 \quad (6)$$

that satisfy the condition $\operatorname{Re} \lambda_k > 0$ (m is the number of the roots of Eq. (6) that satisfy this condition). Equation (6) is a special case of Eq. (3) with $K(z) = A(-z)^n$. The roots of Eq. (6) such that $\operatorname{Re} \lambda_k \leq 0$ must be dropped out, since for them the integral in (3) is divergent.

Equation (6) has complex roots. Consider two cases that correspond to different signs of A .

1°. Let $A < 0$. A solution of the Eq. (4) is

$$y(x) = Ce^{-\lambda x}, \quad \lambda = (-An!)^{\frac{1}{n+1}}, \quad (7)$$

where C is an arbitrary constant. This solution is unique for $n = 0, 1, 2, 3$.

For $n \geq 4$, taking the real and the imaginary part in (5), one arrives at the general solution of the homogeneous Picard–Goursat equation in the form

$$y(x) = C e^{-\lambda x} + \sum_{k=1}^{[n/4]} \exp(-\alpha_k x) [C_k^{(1)} \cos(\beta_k x) + C_k^{(2)} \sin(\beta_k x)], \quad (8)$$

where $C_k^{(1)}$ and $C_k^{(2)}$ are arbitrary constants, $[a]$ stands for the integral part of a number a , λ is defined in (7), and the coefficients α_k and β_k are given by

$$\alpha_k = |An!|^{\frac{1}{n+1}} \cos\left(\frac{2\pi k}{n+1}\right), \quad \beta_k = |An!|^{\frac{1}{n+1}} \sin\left(\frac{2\pi k}{n+1}\right).$$

Note that Eq. (8) contains an odd number of terms.

2°. Let $A > 0$. By taking the real and the imaginary part in (5), one obtains the general solution of the homogeneous Picard–Goursat equation in the form

$$y(x) = \sum_{k=0}^{\left[\frac{n+2}{4}\right]} \exp(-\alpha_k x) [C_k^{(1)} \cos(\beta_k x) + C_k^{(2)} \sin(\beta_k x)], \quad (9)$$

where $C_k^{(1)}$ and $C_k^{(2)}$ are arbitrary constants, and the coefficients α_k and β_k are given by

$$\alpha_k = (An!)^{\frac{1}{n+1}} \cos\left(\frac{2\pi k + \pi}{n+1}\right), \quad \beta_k = (An!)^{\frac{1}{n+1}} \sin\left(\frac{2\pi k + \pi}{n+1}\right).$$

Note that Eq. (9) contains an even number of terms. In the special cases of $n = 0$ and $n = 1$, Eq. (9) gives the trivial solution $y(x) \equiv 0$.

Example 2. Consider the nonhomogeneous Picard–Goursat equation

$$y(x) + A \int_x^\infty (t-x)^n y(t) dt = B e^{-\mu x}, \quad n = 0, 1, 2, \dots, \quad (10)$$

which is a special case of Eq. (1) with $K(z) = A(-z)^n$ and $f(x) = B e^{-\mu x}$.

Let $\mu > 0$. Consider two cases.

1°. Let $\mu^{n+1} + An! \neq 0$. A particular solution of the nonhomogeneous equation is

$$\bar{y}(x) = D e^{-\mu x}, \quad D = \frac{B \mu^{n+1}}{\mu^{n+1} + An!}. \quad (11)$$

For $A < 0$, the general solution of the nonhomogeneous Picard–Goursat equation is the sum of solutions (8) and (11). For $A > 0$, the general solution of the Eq. (10) is the sum of solutions (9) and (11).

2°. Let $\mu^{n+1} + An! = 0$. Since μ is positive, it follows that A must be negative. A particular solution of the nonhomogeneous equation is

$$\bar{y}(x) = E x e^{-\mu x}, \quad E = \frac{B \mu^{n+2}}{A(n+1)!}. \quad (12)$$

The general solution of the nonhomogeneous Picard–Goursat equation is the sum of solutions (8) and (12).

11.11-2. Reduction to a Wiener–Hopf Equation of the Second Kind.

Equation (1) can be reduced to a one-sided equation of the second kind of the form

$$y(x) - \int_0^\infty K_-(x-t)y(t) dt = f(x), \quad 0 < x < \infty, \quad (13)$$

where the kernel $K_-(x-t)$ has the form

$$K_-(s) = \begin{cases} 0 & \text{for } s > 0, \\ -K(s) & \text{for } s < 0. \end{cases}$$

Methods for studying Eq. (13) are described in Chapter 13, where equations of the second kind with constant limits are considered. In the same chapter, in Subsection 13.10-3, an equation of the second kind with difference kernel and variable lower limit is studied by means of reduction to a Wiener–Hopf equation of the second kind.

☉ Reference for Section 11.11: F. D. Gakhov and Yu. I. Cherskii (1978), A. D. Polyaniin and A. V. Manzhurov (1998).

Chapter 12

Methods for Solving Linear Equations of the Form $\int_a^b K(x, t)y(t) dt = f(x)$

12.1. Some Definition and Remarks

12.1-1. Fredholm Integral Equations of the First Kind.

Linear integral equations of the first kind with constant limits of integration have the form

$$\int_a^b K(x, t)y(t) dt = f(x), \quad (1)$$

where $y(x)$ is the unknown function ($a \leq x \leq b$), $K(x, t)$ is the *kernel* of the integral equation, and $f(x)$ is a given function, which is called the *right-hand side* of Eq. (1). The functions $y(x)$ and $f(x)$ are usually assumed to be continuous or square integrable on $[a, b]$. If the kernel of the integral equation (1) is continuous on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ or at least square integrable on this square, i.e.,

$$\int_a^b \int_a^b K^2(x, t) dx dt = B^2 < \infty, \quad (2)$$

where B is a constant, then this kernel is called a *Fredholm kernel*. Equations of the form (1) with constant integration limits and Fredholm kernel are called *Fredholm equations of the first kind*.

The kernel $K(x, t)$ of an integral equation is said to be *degenerate* if it can be represented in the form $K(x, t) = g_1(x)h_1(t) + \cdots + g_n(x)h_n(t)$.

The kernel $K(x, t)$ of an integral equation is called a *difference kernel* if it depends only on the difference of the arguments: $K(x, t) = K(x - t)$.

The kernel $K(x, t)$ of an integral equation is said to be *symmetric* if it satisfies the condition $K(x, t) = K(t, x)$.

The integral equation obtained from (1) by replacing the kernel $K(x, t)$ by $K(t, x)$ is said to be *transposed* to (1).

Remark 1. The variables t and x in Eq. (1) may vary within different intervals (e.g., $a \leq t \leq b$ and $c \leq x \leq d$).

It is important to observe that integral equations of the first kind (1), even with very smooth kernels and right-hand sides, may have no solutions at all or have several (infinitely many) solutions.

Example 1. The equation

$$\int_0^1 y(t) dt = 1 + t$$

has no solutions.

Example 2. The equation

$$\int_0^1 y(t) dt = 1$$

has the solutions $y(x) = 1$ and $y(x) = 1 + C(2x - 1)$, where C is an arbitrary constant. Moreover, this equation has the solution

$$y(x) = \begin{cases} \varphi(x)/A & \text{if } A \neq 0, \\ 1 + C\varphi(x) & \text{if } A = 0, \end{cases} \quad A = \int_0^1 \varphi(x) dx,$$

where $\varphi(x)$ is an arbitrary function.

It should also be mentioned that Fredholm integral equations of the first kind belong to the class of ill-posed problems (for details see Section 12.12).

12.1-2. Integral Equations of the First Kind with Weak Singularity.

If the kernel of the integral equation (1) is polar, i.e., if

$$K(x, t) = \frac{L(x, t)}{|x - t|^\alpha} + M(x, t), \quad 0 < \alpha < 1, \quad (3)$$

or logarithmic, i.e.,

$$K(x, t) = L(x, t) \ln|x - t| + M(x, t), \quad (4)$$

where $L(x, t)$ and $M(x, t)$ are continuous on S and $L(x, x) \neq 0$, then $K(x, t)$ is called a *kernel with weak singularity*, and the equation itself is called an *equation with weak singularity*.

Remark 2. Kernels with logarithmic singularity and polar kernels with $0 < \alpha < \frac{1}{2}$ are Fredholm kernels.

Remark 3. In general, the case in which the limits of integration a and/or b can be infinite is not excluded, but in this case the validity of condition (2) must be verified with special care.

12.1-3. Integral Equations of Convolution Type.

The integral equation of the first kind with difference kernel on the entire axis (this equation is sometimes called an *equation of convolution type of the first kind with a single kernel*) has the form

$$\int_{-\infty}^{\infty} K(x - t)y(t) dt = f(x), \quad -\infty < x < \infty, \quad (5)$$

where $f(x)$ and $K(x)$ are the right-hand side and the kernel of the integral equation and $y(x)$ is the unknown function (in what follows we use the above notation).

An integral equation of the first kind with difference kernel on the semiaxis has the form

$$\int_0^{\infty} K(x - t)y(t) dt = f(x), \quad 0 < x < \infty. \quad (6)$$

Equation (6) is also called a *one-sided equation of the first kind* or a *Wiener-Hopf integral equation of the first kind*.

An integral equation of convolution type with two kernels of the first kind has the form

$$\int_0^{\infty} K_1(x - t)y(t) dt + \int_{-\infty}^0 K_2(x - t)y(t) dt = f(x), \quad -\infty < x < \infty, \quad (7)$$

where $K_1(x)$ and $K_2(x)$ are the kernels of the integral equation (7).

Recall that a function $g(x)$ satisfies the *Hölder condition on the real axis* if for any real x_1 and x_2 we have the inequality

$$|g(x_2) - g(x_1)| \leq A|x_2 - x_1|^\lambda, \quad 0 < \lambda \leq 1,$$

and for any x_1 and x_2 sufficiently large in absolute value we have

$$|g(x_2) - g(x_1)| \leq A \left| \frac{1}{x_2} - \frac{1}{x_1} \right|^\lambda, \quad 0 < \lambda \leq 1,$$

where A and λ are positive (the latter inequality is the Hölder condition in the vicinity of the point at infinity).

Assume that the functions $y(x)$ and $f(x)$ and the kernels $K(x)$, $K_1(x)$, and $K_2(x)$ are such that their Fourier transforms belong to $L_2(-\infty, \infty)$ and, moreover, satisfy the Hölder condition.

For a function $y(x)$ to belong to the above function class it suffices to require $y(x)$ to belong to $L_2(-\infty, \infty)$ and $xy(x)$ to be absolutely integrable on $(-\infty, \infty)$.

12.1-4. Dual Integral Equations of the First Kind.

A dual integral equation of the first kind with difference kernels (of convolution type) has the form

$$\begin{aligned} \int_{-\infty}^{\infty} K_1(x-t)y(t) dt &= f(x), & 0 < x < \infty, \\ \int_{-\infty}^{\infty} K_2(x-t)y(t) dt &= f(x), & -\infty < x < 0, \end{aligned} \quad (8)$$

where the notation and the classes of functions and kernels coincide with those introduced above for equations of convolution type.

In the general case, a dual integral equation of the first kind has the form

$$\begin{aligned} \int_a^{\infty} K_1(x,t)y(t) dt &= f_1(x), & a < x < b, \\ \int_a^{\infty} K_2(x,t)y(t) dt &= f_2(x), & b < x < \infty, \end{aligned}$$

where $f_1(x)$ and $f_2(x)$ are the right-hand sides, $K_1(x,t)$ and $K_2(x,t)$ are the kernels of Eq. (8), and $y(x)$ is the unknown function. Various forms of this equation are considered in Subsections 12.9-3 and 12.9-4.

The integral equations obtained from (5)–(8) by replacing the kernel $K(x-t)$ with $K(t-x)$ are called *transposed* equations.

Remark 3. Some equations whose kernels contain the product or the ratio of the variables x and t can be reduced to equations of the form (5)–(8).

Remark 4. Equations (5)–(8) of the convolution type are sometimes written in the form in which the integrals are multiplied by the coefficient $1/\sqrt{2\pi}$.

12.1-5. Some Problems Leading to Integral Equations of the First Kind.

1°. Historically, one of the first problems that can be associated with integral equations was that of inverting the integral

$$g(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x)e^{ixt} dx,$$

i.e., finding a function $f(x)$ from a given $g(t)$. This problem was solved in 1811 by Fourier, who obtained its solution in the form

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(t)e^{-ixt} dt.$$

2°. Making an elastic string acquire a given shape under the action of a distributed force.

Suppose there is a weightless elastic string of length l that resists tension but does not resist changing its shape. Assume that the string obeys Hooke's law in tension, so that the force required to extend the string by Δl is equal to $\gamma \Delta l$, where γ is some constant.

Let the ends of the string be fixed at points A and B (Fig. 4) and let the string position coincide with the segment AB of the Ox axis when acted upon by only a horizontal tensile force T_0 , very large compared to any other force under consideration.




Figure 4. Shape of an elastic string fixed at points A and B and acted upon by a force P at a point C_0 .

Suppose that a force P is applied to the string at a point C_0 with $x = \xi$. Then the string will take the shape of a broken line ACB . Assume that the displacement $CC_0 = \delta$ is small compared to AC_0 and C_0B , which results from the assumption that P is small compared to T_0 . Also assume that the tension of the string remains equal to T_0 . Projecting the tensile forces at C and the force P onto the vertical, we write down the equilibrium condition to obtain

$$T_0 \sin \alpha + T_0 \sin \beta = P.$$

Since δ is considered to be small, we have

$$\sin \alpha \approx \frac{\delta}{\xi}, \quad \sin \beta \approx \frac{\delta}{l - \xi}.$$

Then the equilibrium condition can be rewritten as

$$T_0 \frac{\delta}{\xi} + T_0 \frac{\delta}{l - \xi} = P.$$

It follows that

$$\delta(\xi) = P \frac{(l - \xi)\xi}{T_0 l}.$$

Let $y(x)$ denote the amount of sag of the string at the point with abscissa x . Then

$$y(x) = PG(x, \xi),$$

where

$$G(x, \xi) = \begin{cases} \frac{x(l - \xi)}{T_0 l} & \text{if } 0 \leq x \leq \xi, \\ \frac{(l - \xi)\xi}{T_0 l} & \text{if } \xi \leq x \leq l. \end{cases}$$

Indeed, for $x < \xi$, from the similarity of the triangles AC_0C and AA_1C_1 (Fig. 4) it follows that

$$\frac{y(x)}{\delta(\xi)} = \frac{x}{\xi}, \quad \text{or} \quad \frac{PG(x, \xi)}{\delta(\xi)} = \frac{x}{\xi}.$$

Hence,

$$G(x, \xi) = \frac{x\delta(\xi)}{P\xi} = \frac{x(l-\xi)}{T_0l}.$$

The case of $\xi < x$ is treated similarly. It is apparent that $G(x, \xi) = G(\xi, x)$.

If the string is acted upon by a continuously distributed force with line density $p(\xi)$, then the small segment between ξ and $\xi + \Delta\xi$ is subjected to the force approximately equal to $p(\xi) \Delta\xi$ and is displaced by the distance $G(x, \xi) p(\xi) \Delta\xi$. Since the displacements caused by the elementary forces $p(\xi) \Delta\xi$ are summed (according to the principle of superposition), the total amount of sag $y(x)$ is approximately equal to

$$\sum_{(\xi)} G(x, \xi) p(\xi) \Delta\xi.$$

On passing to the limit as $\Delta\xi \rightarrow 0$, one arrives at a Fredholm integral equation of the first kind:

$$y(x) = \int_0^l G(x, \xi) p(\xi) d\xi.$$

This equation serves to determine the force density $p(x)$ under the action of which the string will take the given shape $y = y(x)$. The function $G(x, \xi)$ is called an *influence function*.

© References for Section 12.1: B. Noble (1958), S. G. Mikhlin (1960), I. C. Gohberg and M. G. Krein (1967), L. Ya. Tslaf (1970), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), P. P. Zabreyko, A. I. Koshelev, et al. (1975), Ya. S. Uflyand (1977), F. D. Gakhov and Yu. I. Cherskii (1978), A. J. Jerry (1985), A. F. Verlan' and V. S. Sizikov (1986), I. Sneddon (1995), A. V. Bitsadze (1995), L. A. Sakhnovich (1996).

12.2. Integral Equations of the First Kind with Symmetric Kernel

12.2-1. Solution of an Integral Equation in Terms of Series in Eigenfunctions of Its Kernel.

Suppose $K(x, t)$ is a real symmetric kernel defined on a segment $[a, b]$. Let us write out the system of characteristic values and eigenfunctions of this kernel* as the sequences

$$\begin{aligned} \lambda_1, \quad \lambda_2, \quad \dots, \quad \lambda_n, \quad \dots; \\ y_1(x), \quad y_2(x), \quad \dots, \quad y_n(x), \quad \dots, \end{aligned} \tag{1}$$

where

$$y_n(x) - \lambda_n \int_a^b K(x, t) y_n(t) dt = 0.$$

It is assumed that the following conditions hold:

- 1) The values λ_n are ordered so that their moduli form a nondecreasing sequence, i.e., $|\lambda_{n-1}| \leq |\lambda_n|$.
- 2) Each characteristic value appears as many times as its multiplicity (rank), so that one and the same value λ in (1) may occur several times, each corresponding to only one eigenfunction.
- 3) Eigenfunctions $y_n(x)$ are normalized and mutually orthogonal in $L_2[a, b]$ (for details, see Subsection 13.6-1).

* For definitions of characteristic values and eigenfunctions of a kernel $K(x, t)$, see Subsection 13.1-1.

A symmetric kernel $K(x, t)$ defined on $[a, b]$ is called *complete* (or *closed*), if the system of the corresponding eigenfunctions is complete in $L_2[a, b]$; otherwise, the kernel is called *incomplete*.

Consider a nonhomogeneous integral equation of the first kind

$$\int_a^b K(x, t)y(t) dt = f(x) \quad (2)$$

with a real symmetric kernel and $f \in L_2[a, b]$.

PICARD THEOREM. Equation (2) has a solution if and only if $f(x)$ can be expanded into a mean-square convergent series with respect to eigenfunctions of the kernel $K(x, t)$:

$$f(x) = \sum_{k=1}^{\infty} f_k y_k(x), \quad f_k = \int_a^b f(x)y_k(x) dx, \quad (3)$$

and the series

$$\sum_{k=1}^{\infty} \lambda_k^2 |f_k|^2 \quad (4)$$

is convergent. In this case, the general solution of equation (2) has the form

$$y(x) = y_0(x) + \sum_{k=1}^{\infty} \lambda_k f_k y_k(x),$$

where $y_0(x)$ is an arbitrary solution of the homogeneous equation (2) for $f(x) \equiv 0$. If the kernel $K(x, t)$ is complete, then $y_0(x) \equiv 0$, and equation (2) has only one solution,

$$y(x) = \sum_{k=1}^{\infty} \lambda_k f_k y_k(x). \quad (5)$$

Example. Consider the integral equation

$$\int_0^1 K(x, t)y(t) dt = \sin^3(\pi x) \quad (6)$$

with the real symmetric kernel

$$K(x, t) = \begin{cases} (1-x)t & \text{if } 0 \leq t \leq x, \\ (1-t)x & \text{if } x \leq t \leq 1. \end{cases} \quad (7)$$

Let us use the Picard theorem to find its solution. First, we find the characteristic values and the corresponding normalized eigenfunctions of the kernel (7):

$$\begin{aligned} \lambda_1 &= \pi^2, & \lambda_2 &= (2\pi)^2, & \dots, & \lambda_n &= (n\pi)^2, & \dots; \\ y_1(x) &= \sqrt{2} \sin(\pi x), & y_2(x) &= \sqrt{2} \sin(2\pi x), & \dots, & y_n(x) &= \sqrt{2} \sin(n\pi x), & \dots \end{aligned} \quad (8)$$

Then we express the right-hand side of (6) in terms of the eigenfunctions:

$$f(x) \equiv \sin^3(\pi x) = \frac{3}{4} \sin(\pi x) - \frac{1}{4} \sin(3\pi x) = \frac{3}{4\sqrt{2}} y_1(x) - \frac{1}{4\sqrt{2}} y_3(x)$$

and write out the corresponding coefficients in the expansion of $f(x)$:

$$f_1 = \frac{3}{4\sqrt{2}}, \quad f_2 = 0, \quad f_3 = -\frac{1}{4\sqrt{2}}, \quad f_m = 0 \quad \text{for } m = 4, 5, \dots$$

The series (4) in this case reduces to the finite sum

$$\sum_{k=1}^{\infty} \lambda_k^2 |f_k|^2 = (\pi^2)^2 \left(\frac{3}{4\sqrt{2}}\right)^2 + (9\pi^2)^2 \left(-\frac{1}{4\sqrt{2}}\right)^2 = \frac{45}{16} \pi^4$$

and is therefore convergent. The system of eigenfunctions (8) is a complete orthonormal system on $[0, 1]$, i.e., the kernel is complete. By the Picard theorem, equation (6)–(7) has the unique solution

$$y(x) = \lambda_1 f_1 y_1(x) + \lambda_3 f_3 y_3(x),$$

which can be written in the form

$$y(x) = \frac{3}{4} \pi^2 [\sin(\pi x) - 3 \sin(3\pi x)].$$

12.2-2. Method of Successive Approximations.

THEOREM. Let $K(x, t)$ be a symmetric positive kernel and suppose that the equation

$$\int_a^b K(x, t)y(t) dt = f(x), \quad f(x) \in L_2[a, b], \tag{9}$$

admits one and only one solution. Then the sequence of functions $\{y_n(x)\}$ defined by the recurrent relation

$$y_n(x) = y_{n-1}(x) + \lambda \left[f(x) - \int_a^b K(x, t)y_{n-1}(t) dt \right], \quad n = 1, 2, \dots, \tag{10}$$

where

$$y_0(x) \in L_2[a, b], \quad 0 < \lambda < 2\lambda_1, \tag{11}$$

λ_1 is the smallest characteristic value of the kernel $K(x, t)$, is mean-square convergent to the solution of equation (9).

Remark. If there is no information about the solution of equation (9), one can take $y_0(x) = 0$ as the zero approximation. If the smallest characteristic value λ_1 is unknown, then λ should be chosen sufficiently small and one should check (control) the convergence of the process (10).

Example. Consider the integral equation

$$\int_0^1 K(x, t)y(t) dt = \sin(\pi x), \tag{12}$$

where

$$K(x, t) = \begin{cases} (1-x)t & \text{if } 0 \leq t \leq x, \\ (1-t)x & \text{if } x \leq t \leq 1. \end{cases} \tag{13}$$

Let us construct successive approximations by formulas (10), taking $y_0(x) = 0$ and imposing no constraints on λ so far. We have

$$\begin{aligned} y_1(x) &= \lambda \sin(\pi x), \\ y_2(x) &= \lambda \sin(\pi x) + \lambda \left(1 - \frac{\lambda}{\pi^2} \right) \sin(\pi x), \\ y_3(x) &= \lambda \sin(\pi x) + \lambda \left(1 - \frac{\lambda}{\pi^2} \right) \sin(\pi x) + \lambda \left(1 - \frac{\lambda}{\pi^2} \right)^2 \sin(\pi x), \\ &\dots \\ y_n(x) &= \lambda \left[1 + \left(1 - \frac{\lambda}{\pi^2} \right) + \left(1 - \frac{\lambda}{\pi^2} \right)^2 + \dots + \left(1 - \frac{\lambda}{\pi^2} \right)^{n-1} \right] \sin(\pi x), \end{aligned} \tag{14}$$

The square brackets contain the finite sum of a geometrical progression with ratio $q = 1 - \frac{\lambda}{\pi^2}$. This sum is calculated by the formula

$$\sum_{m=0}^{n-1} q^m = \frac{1-q^n}{1-q}, \quad q = 1 - \frac{\lambda}{\pi^2}.$$

For $n \rightarrow \infty$, this sum has a finite limit equal to $\frac{1}{1-q}$, provided that $|q| < 1$, which yields the following constraint on the coefficient λ :

$$0 < \lambda < 2\pi^2. \tag{15}$$

Passing to the limit in (14) as $n \rightarrow \infty$, under the condition (15), we find that

$$\lim_{n \rightarrow \infty} y_n(x) = \frac{\lambda}{1-q} \sin(\pi x) = \pi^2 \sin(\pi x). \tag{16}$$

It is easy to check that the limit solution (16) coincides with the exact solution of the integral equation (12)–(13). It can be shown that the smallest characteristic value of the kernel (13) is $\lambda_1 = \pi^2$. Therefore, condition (11), in this case, turns into (15).

© References for Section 12.2: M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), M. L. Krasnov (1975), P. P. Zabreyko, A. I. Koshelev et al. (1975).

12.3. Integral Equations of the First Kind with Nonsymmetric Kernel

12.3-1. Representation of a Solution in the Form of Series. General Description.

Consider an integral equation

$$\int_a^b K(x, t)y(t) dt = f(x) \quad (1)$$

with an arbitrary (symmetric or nonsymmetric) kernel. Let us seek its solution in the form of a sum

$$y(t) = \sum_{n=1}^N A_n \varphi_n(x), \quad (2)$$

where $\varphi_n(x)$ is a (complete) system of functions on the interval (a, b) , the upper limit of the sum, N , can be either finite or infinite. It is important to mention that in some cases it is possible to obtain an exact solution of the integral equation (1) in the form of series (2) for $N = \infty$ (see Examples 1 and 2 in Subsection 12.3-2).

Substituting (2) into (1), we get

$$f(x) = \sum_{n=1}^N A_n g_n(x), \quad (3)$$

where $g_n(x)$ are known functions,

$$g_n(x) = \int_a^b K(x, t)\varphi_n(t) dt. \quad (4)$$

In order to find the coefficients A_n in the right-hand side of (3), different methods can be used, depending on the structure of the functions $g_n(x)$. Some basic methods are described below.

12.3-2. Special Case of a Kernel That is a Generating Function.

For power-type functions $g_n(x) = b_n x^n$, the right hand side of (3) is a polynomial or a power series (for $N = \infty$). The coefficients A_n of that series can be found by way of comparison with the corresponding coefficients in the expansion of $f(x)$ in powers of x . This case takes place if the kernel of the integral equation is a generating function for a system of orthogonal polynomials. Recall that $G(x, t)$ is called a *generating function* for a system of functions

$$h_0(t), h_1(t), \dots, h_m(t), \dots$$

if $G(x, t)$ admits the following expansion in powers of x :

$$G(x, t) = \sum_{m=0}^{\infty} c_m h_m(t) x^m \quad (c_m \neq 0).$$

Example 1. Consider the equation

$$\int_{-1}^1 \frac{y(t) dt}{\sqrt{1+x^2-2xt}} = f(x). \quad (5)$$

Its kernel is a generating function for the Legendre polynomials (see Supplement 11.11-1):

$$\frac{1}{\sqrt{1+x^2-2xt}} = \sum_{m=0}^{\infty} P_m(t)x^m, \quad P_m(x) = \frac{1}{m! 2^m} \frac{d^m}{dx^m} (x^2-1)^m. \quad (6)$$

Let us seek a solution of equation (5) in the form

$$y(x) = \sum_{n=0}^{\infty} A_n P_n(x). \quad (7)$$

Substituting (6) and (7) into equation (5) and taking into account the orthogonality conditions for the Legendre polynomials,

$$\int_{-1}^1 P_n(x) P_m(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{2}{2n+1} & \text{if } n = m, \end{cases}$$

we find that

$$2 \sum_{n=0}^{\infty} \frac{A_n}{2n+1} x^n = f(x).$$

Expanding the right-hand side into a Maclaurin series and equating the coefficients of equal powers of x , we obtain

$$A_n = \frac{2n+1}{2n!} f_x^{(n)}(0).$$

Inserting these coefficients into (7), we obtain a solution of the integral equation (5) in the form

$$y(x) = \frac{1}{2} \sum_{n=0}^{\infty} \frac{2n+1}{n!} f_x^{(n)}(0) P_n(x). \quad (8)$$

It is easy to see that if the right-hand side of equation (5) is a polynomial, then its solution (8) is a polynomial of the same degree.

Example 2. Consider the equation

$$\int_{-\infty}^{\infty} e^{-(x-t)^2} y(t) dt = f(x) \quad (9)$$

whose kernel is a generating function for the Hermitian polynomials (see Supplement 11.17-3)

$$e^{-(x-t)^2} = \sum_{m=0}^{\infty} \frac{1}{m!} e^{-t^2} H_m(t) x^m, \quad H_m(x) = (-1)^m \exp(x^2) \frac{d^m}{dx^m} \exp(-x^2). \quad (10)$$

Let us seek a solution of equation (9) in the form of expansion

$$y(x) = \sum_{n=0}^{\infty} A_n H_n(x). \quad (11)$$

Substituting (10) and (11) into (9) and taking into account the orthogonality of the Hermitian polynomials, together with the relations

$$\int_{-\infty}^{\infty} e^{-t^2} H_n^2(t) dt = 2^n n! \sqrt{\pi},$$

we obtain

$$f(x) = \sqrt{\pi} \sum_{n=0}^{\infty} A_n 2^n x^n.$$

Hence, we find the coefficients A_n :

$$A_n = \frac{f_x^{(n)}(0)}{2^n n! \sqrt{\pi}}.$$

Substituting these into (11), we obtain a solution of the original integral equation (9):

$$y(x) = \frac{1}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{f_x^{(n)}(0)}{2^n n!} H_n(x).$$

12.3-3. Special Case of the Right-Hand Side Represented in Terms of Orthogonal Functions.

Suppose that the functions $g_n(x)$ in the right-hand side of (3) form an orthogonal, with some weight $\rho(x)$, system on the interval (a, b) :

$$\int_a^b \rho(x)g_n(x)g_m(x) dx = 0 \quad \text{if } n \neq m.$$

Then the coefficients A_n are obtained by multiplying (3) by $\rho(x)g_m(x)$ with subsequent integration in x over the segment $[a, b]$. As a result, we get

$$A_n = \frac{1}{\beta_n} \int_a^b f(x)\rho(x)g_n(x) dx, \quad \beta_n = \int_a^b \rho(x)g_n^2(x) dx.$$

12.3-4. General Case. Galerkin's Method.
--

In the general case, one chooses a sequence of functions $\psi_m(x)$, $m = 1, \dots, n$, multiplies the relation (3) by these in successive order, and then integrates in x over the segment $[a, b]$. The result is a system of linear algebraic equations for the coefficients A_n :

$$\sum_{n=1}^N \sigma_{nm}A_n = B_m, \quad m = 1, \dots, n;$$

$$\sigma_{nm} = \int_a^b g_n(x)\psi_m(x) dx, \quad B_m = \int_a^b f(x)\psi_m(x) dx.$$

Finding the coefficients A_n from this system and substituting these in (3), one obtains an approximate solution of the integral equation (1).

Remark. The Galerkin method and its modifications, when applied to the solution of integral equations of the second kind, may result in large errors (connected with the instability of solutions with respect to small perturbations of the right-hand side of the equation; see Section 12.12). For this reason the said methods are rarely used in practice.

12.3-5. Utilization of the Schmidt Kernels for the Construction of Solutions of Equations.
--

Let $K(x, t)$ be a real (or complex) nonsymmetric kernel, $K(x, t) \neq K(t, x)$, such that

$$\int_a^b \int_a^b |K(x, t)|^2 dx dt < \infty.$$

The kernels $K(x, t)$ and $K^*(x, t) = \overline{K(t, s)}$ are called *conjugate*.

Consider auxiliary functions

$$K_1(x, t) = \int_a^b K^*(x, s)K(s, t) ds = \int_a^b \overline{K(s, x)}K(s, t) ds, \quad (12)$$

$$K_2(x, t) = \int_a^b K(x, s)K^*(s, t) ds = \int_a^b K(x, s)\overline{K(t, s)} ds, \quad (13)$$

representing symmetric positive kernels called the *Schmidt kernels* corresponding to $K(x, t)$. It can be shown that the system of characteristic values of the kernels (12) and (13) coincide.

Denote by μ_n ($n = 1, 2, \dots$) the characteristic values of the Schmidt kernels, by $u_n(x)$ the orthonormalized eigenfunctions corresponding to $K_2(x, t)$, and by $v_n(x)$ orthonormalized eigenfunctions corresponding to $K_1(x, t)$. Each $u_n(x)$ and $v_n(x)$ can be multiplied by an arbitrary constant coefficient whose absolute value is equal to unity. These coefficients can be chosen such that the following formulas hold:

$$K(x, t) = \sum_{n=1}^{\infty} \frac{u_n(x)\overline{v_n(t)}}{\sqrt{\mu_n}}, \quad K^*(x, t) = \sum_{n=1}^{\infty} \frac{v_n(x)\overline{u_n(t)}}{\sqrt{\mu_n}}. \quad (14)$$

These series are mean-square convergent on $[a, b]$ (with respect to the variables x and t jointly). The following inequality holds:

$$\sum_{n=1}^{\infty} \frac{1}{\mu_n} \leq \int_a^b \int_a^b |K(x, t)|^2 dx dt.$$

For a nonhomogeneous equation of the first kind

$$\int_a^b K(x, t)y(t) dt = f(x), \quad f \in L_2[a, b], \quad (15)$$

with nonsymmetric kernel to have a solution it is necessary and sufficient that the free term $f(x)$ could be expanded into a mean-square convergent series in terms of eigenfunctions u_n :

$$f(x) = \sum_{n=1}^{\infty} f_n u_n(x), \quad f_n = \int_a^b f(x) u_n(x) dx,$$

and that the series

$$\sum_{n=1}^{\infty} \mu_n |f_n|^2$$

be convergent. Under these conditions, the general solution of equation (15) has the form

$$y(x) = y_0(x) + \sum_{n=1}^{\infty} \sqrt{\mu_n} f_n v_n(x),$$

where $y_0(x)$ is any solution of the homogeneous equation (15) with $f(x) \equiv 0$. If the Schmidt kernel $K_1(x, t)$ is complete, then $y_0(x) \equiv 0$, and equation (15) has only one solution,

$$y(x) = \sum_{n=1}^{\infty} \sqrt{\mu_n} f_n v_n(x).$$

⊙ References for Section 12.3: P. M. Morse and H. Feshbach (1953), L. Ya. Tslaf (1970), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), V. I. Smirnov (1974), P. P. Zabreyko, A. I. Koshelev et al. (1975).

12.4. Method of Differentiation for Integral Equations

12.4-1. Equations with Modulus.

In some cases, differentiation of integral equations (once, twice, etc.) with subsequent elimination of integral terms by means of the original equation makes it possible to find solutions of the latter.

The class of integral equations whose solutions can be obtained by the method of differentiation includes integral equations of the first kind with difference kernel,

$$\int_a^b K(|x-t|)y(t) dt = f(x), \quad (1)$$

for the following types of $K(z)$:

$$K(z) = \sum_{m=1}^n A_m z^m, \quad (2)$$

$$K(z) = \sum_{m=1}^n A_m \exp(\lambda_m z), \quad (3)$$

$$K(z) = \sum_{m=1}^n A_m \sinh(\lambda_m z), \quad (4)$$

$$K(z) = \sum_{m=1}^n A_m \sin(\lambda_m z). \quad (5)$$

Example 1. Consider the equation

$$\int_0^1 |x-t| y(t) dt = f(x), \quad (6)$$

which is a special case of (1), (2) for $n = 1$.

1°. Let us remove the modulus in the integrand:

$$\int_0^x (x-t)y(t) dt + \int_x^1 (t-x)y(t) dt = f(x). \quad (7)$$

Differentiating (7) in x yields

$$\int_0^x y(t) dt - \int_x^1 y(t) dt = f'_x(x). \quad (8)$$

Differentiating (8) in x yields the solution

$$y(x) = \frac{1}{2} f''_{xx}(x). \quad (9)$$

2°. The right-hand side $f(x)$ of the integral equation (6) must satisfy certain additional relations. In order to obtain these, let us substitute the solution (9) into the transformed original equation (7). Integrating by parts, we get

$$\begin{aligned} \int_0^x (x-t)y(t) dt &= \frac{1}{2} \int_0^x (x-t)f''_{tt}(t) dt = \frac{1}{2} [-xf'(0) + f(x) - f(0)], \\ \int_x^1 (t-x)y(t) dt &= \frac{1}{2} \int_x^1 (t-x)f''_{tt}(t) dt = \frac{1}{2} [-xf'(1) + f(x) + f'(1) - f(1)]. \end{aligned}$$

Substituting these integrals into the left-hand side of equation (7) and reducing the result by $f(x)$, we obtain

$$-\frac{1}{2}x[f'(0) + f'(1)] + \frac{1}{2}[f'(1) - f(1) - f(0)] = 0.$$

Since this relation must hold for all x , we obtain the following two conditions:

$$\begin{aligned} f'(0) + f'(1) &= 0, \\ f'(1) - f(1) - f(0) &= 0, \end{aligned}$$

which should be satisfied by the right-hand side of the integral equation (6).

Example 2. Consider the equation

$$\int_a^b e^{\lambda|x-t|}y(t) dt = f(x), \quad (10)$$

which is a special case of (1) with kernel (3) for $n = 1$.

Let us remove the modulus in the integrand:

$$\int_a^x e^{\lambda(x-t)}y(t) dt + \int_x^b e^{\lambda(t-x)}y(t) dt = f(x). \quad (11)$$

Differentiating (11) with respect to x twice yields

$$2\lambda y(x) + \lambda^2 \int_a^x e^{\lambda(x-t)}y(t) dt + \lambda^2 \int_x^b e^{\lambda(t-x)}y(t) dt = f''_{xx}(x). \quad (12)$$

Eliminating the integral terms from (11) and (12), we obtain the solution

$$y(x) = \frac{1}{2\lambda} [f''_{xx}(x) - \lambda^2 f(x)]. \quad (13)$$

The right-hand side $f(x)$ of the integral equation (10) must satisfy certain additional relations. In order to obtain these, one should substitute the solution (13) into the original equation (10) or its corollary (11). Another method of finding additional conditions on $f(x)$ is described in Section 3.2 (see Eq. 3, Item 2°).

Other examples of solutions of such equations can be found in Section 3.1 (equations 2, 8, 11, and 16), Section 3.2 (equations 3, 4, and 6), Section 3.3 (equations 5, 6, and 10), Section 3.5 (equations 10, 11, 12, and 16).

In a similar way, one can find solutions of the equation

$$\int_a^b |g(x) - h(t)|y(t) dt = f(x).$$

Some equations of this type are considered in 3.8 (see equations 4–6).

12.4-2. Other Equations. Some Generalizations.

1°. Sometimes, differentiation helps to reduce a given equation

$$\int_a^b K(x, t)y(t) dt = f(x) \quad (14)$$

to a simpler integral equation

$$\int_a^b K'_x(x, t)y(t) dt = f'_x(x) \quad (15)$$

whose solution is known.

Note that equations (14) and (15) may be nonequivalent. Thus, if $y(t)$ is a solution of (14), it is also a solution of (15) (provided that the integral on the left-hand side of the equation exists). On the other hand, if $y(t)$ is a solution of (15), it will satisfy equation (14) only under the additional condition

$$\int_a^b K(c, t)y(t) dt = f(c) \quad (a < c < b), \quad (16)$$

which is obtained by taking $x = c$ in the original equation (14).

Example 3. Consider the equation

$$\int_{-1}^1 \ln \left| \tanh \frac{\lambda(t-x)}{2} \right| y(t) dt = -f(x), \quad -1 \leq x \leq 1. \quad (17)$$

Differentiating this equation in x , we obtain the following singular equation:

$$\lambda \int_{-1}^1 \frac{y(t) dt}{\sinh[\lambda(t-x)]} = f'_x(x). \quad (18)$$

Equations (18) and (17) are equivalent under the additional condition

$$\int_{-1}^1 \ln \left| \tanh \frac{\lambda t}{2} \right| y(t) dt = -f(0). \quad (19)$$

Let us rewrite (18) as follows:

$$\lambda \int_{-1}^1 \frac{y(t) dt}{\cosh(\lambda t)[\tanh(\lambda t) - \tanh(\lambda x)]} = \cosh(\lambda x) f'_x(x),$$

and make the transformation

$$z = \tanh(\lambda x), \quad \tau = \tanh(\lambda t), \quad Y(\tau) = \cosh(\lambda t)y(t), \quad h(z) = \cosh(\lambda x) f'_x(x). \quad (20)$$

As a result, we obtain a much simpler equation,

$$\int_{-a}^a \frac{Y(\tau) d\tau}{\tau - z} = h(z) \quad (a = \tanh \lambda, |z| \leq a),$$

whose solution is given in Section 3.1 (see equation 51). Using this solution and going back to the original variables by formulas (20), one can find a solution of the original equation (17).

2°. As a preliminary step, one can multiply equation (14) by a function $\varphi(x)$, and then integrate the result in x . Thus, one obtains the equation

$$\int_a^b \frac{\partial}{\partial x} [\varphi(x)K(x, t)]y(t) dt = [\varphi(x)f(x)]'_x.$$

If one can find a solution of the last equation, its solution should be inserted into the original equation (14) or into (16) for verification.

Remark 1. Instead of differentiation, one can multiply equation (14) by a function $\varphi(x)$, integrate the result in x from a to x , and try to find a solution of the equation thus obtained.

Remark 2. If equation (14) does not depend on the parameter μ , it can be multiplied by a function $\varphi(x, \mu)$ and then integrated with respect to μ from α to β .

⊙ References for Section 12.4: I. I. Vorovich, V. M. Aleksandrov, and V. A. Babeshko (1974), A. D. Polyanin and A. V. Manzhirov (1998).

12.5. Method of Integral Transforms

The method of integral transforms enables one to reduce some integral equations on the entire axis and on the semiaxis to algebraic equations for transforms. These algebraic equations can readily be solved for the transform of the desired function. The solution of the original integral equation is then obtained by applying the inverse integral transform.

12.5-1. Equation with Difference Kernel on the Entire Axis.

Consider the integral equation

$$\int_{-\infty}^{\infty} K(x-t)y(t) dt = f(x), \quad -\infty < x < \infty, \quad (1)$$

where $f(x), y(x) \in L_2(-\infty, \infty)$ and $K(x) \in L_1(-\infty, \infty)$.

Let us apply the Fourier transform to Eq. (1). In this case, taking into account the convolution theorem (see Subsection 9.4-4), we obtain

$$\sqrt{2\pi} \tilde{K}(u)\tilde{y}(u) = \tilde{f}(u). \quad (2)$$

Thus, by means of the Fourier transform we have reduced the solution of the original integral equation (1) to the solution of the algebraic equation (2) for the Fourier transform of the desired solution. The solution of the latter equation has the form

$$\tilde{y}(u) = \frac{1}{\sqrt{2\pi}} \frac{\tilde{f}(u)}{\tilde{K}(u)}, \quad (3)$$

where the function $\tilde{f}(u)/\tilde{K}(u)$ must belong to the space $L_2(-\infty, \infty)$.

Thus, the Fourier transform of the solution of the original integral equation is expressed via the Fourier transforms of known functions, namely, the kernel and the right-hand side of the equation. The solution itself can be expressed via its Fourier transform by means of the Fourier inversion formula:

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \tilde{y}(u) e^{iux} du = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\tilde{f}(u)}{\tilde{K}(u)} e^{iux} du. \quad (4)$$

12.5-2. Equations with Kernel $K(x, t) = K(x/t)$ on the Semiaxis.

The integral equation of the first kind

$$\int_0^{\infty} K(x/t)y(t) dt = f(x), \quad 0 \leq x < \infty, \quad (5)$$

can be reduced to the form (1) by the change of variables $x = e^\xi$, $t = e^\tau$, $w(\tau) = ty(t)$. The solution to this equation can also be obtained by straightforward application of the Mellin transform, and this method is applied in a similar situation in the next section.

12.5-3. Equation with Kernel $K(x, t) = K(xt)$ and Some Generalizations.

1°. We first consider the equation

$$\int_0^{\infty} K(xt)y(t) dt = f(x), \quad 0 \leq x < \infty. \quad (6)$$

By changing variables $x = e^\xi$ and $t = e^{-\tau}$ this equation can be reduced to the form (1), but it is more convenient here to apply the Mellin transform (see Section 9.3). On multiplying Eq. (6) by x^{s-1} and integrating with respect to x from 0 to ∞ , we obtain

$$\int_0^{\infty} y(t) dt \int_0^{\infty} K(xt)x^{s-1} dx = \int_0^{\infty} f(x)x^{s-1} dx.$$

We make the change of variables $z = xt$ in the inner integral of the double integral. This implies the relation

$$\hat{K}(s) \int_0^{\infty} y(t)t^{-s} dt = \hat{f}(s). \quad (7)$$

Taking into account the formula

$$\int_0^{\infty} y(t)t^{-s} dt = \hat{y}(1-s),$$

we can rewrite Eq. (7) in the form

$$\hat{K}(s)\hat{y}(1-s) = \hat{f}(s). \quad (8)$$

Replacing $1 - s$ by s in (8) and solving the resulting relation for $\hat{y}(s)$, we obtain the transform

$$\hat{y}(s) = \frac{\hat{f}(1-s)}{\hat{K}(1-s)} \quad (9)$$

of the desired solution.

Applying the Mellin inversion formula, we obtain the solution of the integral equation (6) in the form

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\hat{f}(1-s)}{\hat{K}(1-s)} x^{-s} ds.$$

2°. Now we consider the more complicated equation

$$\int_0^\infty K(\varphi(x)\psi(t))g(t)y(t) dt = f(x). \quad (10)$$

Assume that the conditions $\varphi(0) = 0$, $\varphi(\infty) = \infty$, $\varphi'_x > 0$, $\psi(0) = 0$, $\psi(\infty) = \infty$, and $\psi'_x > 0$ are satisfied.

The transform

$$z = \varphi(x), \quad \tau = \psi(t), \quad y(t) = \frac{g(t)}{\psi'_t(t)} w(\tau)$$

takes (10) to the following equation of the form (6):

$$\int_0^\infty K(z\tau)w(\tau) d\tau = F(z),$$

where the function $F(z)$ is defined parametrically by $F = f(x)$, $z = \varphi(x)$. In many cases, on eliminating x from these relations, we obtain the dependence $F = F(z)$ in an explicit form.

⊙ References for Section 12.5: V. A. Ditkin and A. P. Prudnikov (1965), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971).

12.6. Krein's Method and Some Other Exact Methods for Integral Equations of Special Types

12.6-1. Krein's Method for an Equation with Difference Kernel with a Weak Singularity.

1°. Here we describe a method for constructing exact closed-form solutions of linear integral equations of the first kind with weak singularity and with arbitrary right-hand side. The method is based on the construction of the auxiliary solution of the simpler equation whose right-hand side is equal to one. The auxiliary solution is then used to construct the solution of the original equation for an arbitrary right-hand side.

Consider the equation

$$\int_{-a}^a K(x-t)y(t) dt = f(x), \quad -a \leq x \leq a. \quad (1)$$

Suppose that the kernel of the integral equation (1) is polar or logarithmic and that $K(x)$ is an even positive definite function that can be expressed in the form

$$K(x) = \beta|x|^{-\mu} + M(x), \quad 0 < \mu < 1,$$

$$K(x) = \beta \ln \frac{1}{|x|} + M(x),$$

respectively, where $\beta > 0$, $-2a \leq x \leq 2a$, and $M(x)$ is a sufficiently smooth function.

Along with (1), we consider the following auxiliary equation containing a parameter ξ ($0 \leq \xi \leq a$):

$$\int_{-\xi}^{\xi} K(x-t)w(t, \xi) dt = 1, \quad -\xi \leq x \leq \xi. \quad (2)$$

2°. For any continuous function $f(x)$, the solution of the original equation (1) can be expressed via the solution of the auxiliary equation (2) by the formula

$$\begin{aligned} y(x) = & \frac{1}{2M'(a)} \left[\frac{d}{da} \int_{-a}^a w(t, a)f(t) dt \right] w(x, a) \\ & - \frac{1}{2} \int_{|x|}^a w(x, \xi) \frac{d}{d\xi} \left[\frac{1}{M'(\xi)} \frac{d}{d\xi} \int_{-\xi}^{\xi} w(t, \xi)f(t) dt \right] d\xi \\ & - \frac{1}{2} \frac{d}{dx} \int_{|x|}^a \frac{w(x, \xi)}{M'(\xi)} \left[\int_{-\xi}^{\xi} w(t, \xi) df(t) \right] d\xi, \end{aligned} \quad (3)$$

where $M(\xi) = \int_0^{\xi} w(x, \xi) dx$, the prime stands for the derivative, and the last inner integral is treated as a Stieltjes integral.

Formula (3) permits one to obtain some exact solutions of integral equations of the form (1) with arbitrary right-hand side, see Section 3.6 of the first part of the book.

Example 1. The solution of the integral equation

$$\int_{-a}^a \ln \left(\frac{A}{|x-t|} \right) y(t) dt = f(x),$$

which arises in elasticity, is given by formula (3), where

$$M(\xi) = \left(\ln \frac{2A}{\xi} \right)^{-1}, \quad w(t, \xi) = \frac{M(\xi)}{\pi \sqrt{\xi^2 - t^2}}.$$

Example 2. Consider the integral equation

$$\int_{-a}^a \frac{y(t) dt}{|x-t|^\mu} = f(x), \quad 0 < \mu < 1,$$

which arises in the theory of elasticity. The solution is given by formula (3), where

$$M(\xi) = \frac{2\sqrt{\pi}}{\mu \Gamma\left(\frac{\mu}{2}\right) \Gamma\left(\frac{1-\mu}{2}\right)} \xi^\mu, \quad w(t, \xi) = \frac{1}{\pi} \cos\left(\frac{\pi\mu}{2}\right) (\xi^2 - t^2)^{\frac{\mu-1}{2}}.$$

12.6-2. Kernel is the Sum of a Nondegenerate Kernel and an Arbitrary Degenerate Kernel.

1°. Consider the Fredholm equation of the first kind

$$\int_a^b K(x, t)y(t) dt = f(x). \quad (4)$$

Suppose equation (4) can be solved for any $f(x)$ from some class of functions $L_{\mathcal{F}}$. Let $y_f(x)$ denote the corresponding solution.

Now consider the more complex integral equation

$$\int_a^b [K(x, t) + \varphi(x)\psi(t)]u(t) dt = f(x) \quad (5)$$

with its kernel containing an additional term $\varphi(x)\psi(t)$. A solution to equation (5) will be sought in the form

$$u(x) = y_f(x) + Ay_\varphi(x), \quad (6)$$

where $y_\varphi(x)$ is the solution to equation (4) in which $f(x)$ must be replaced with $\varphi(x)$. Substituting (6) into (5) results in the coefficient A :

$$A = -\frac{\int_a^b \psi(t)y_f(t) dt}{1 + \int_a^b \psi(t)y_\varphi(t) dt}. \quad (7)$$

Formulas (6)–(7) define a solution to equation (5), provided the integrals in the numerator and denominator exist, with $\int_a^b \psi(t)y_\varphi(t) dt \neq -1$. In addition, the condition $\varphi(x) \in L_{\mathcal{F}}$ must be satisfied.

Example 3. The solution of Carleman's equation

$$\int_0^1 \ln|x-t|y(t) dt = f(x) \quad (8)$$

is expressed as

$$y_f(x) = \frac{1}{\pi^2 \sqrt{x(1-x)}} \left[\int_0^1 \frac{\sqrt{t(1-t)}f'_t(t) dt}{t-x} - \frac{1}{\ln 4} \int_0^1 \frac{f(t) dt}{\sqrt{t(1-t)}} \right]. \quad (9)$$

Now consider the more complex integral equation

$$\int_0^1 [\ln|x-t| + \psi(t)] u(t) dt = f(x) \quad (10)$$

with its kernel containing an arbitrary additive function $\psi(t)$. In terms of equation (5), we have $\varphi(x) = 1$ in (10). The corresponding solution (9) to equation (8) with $f(x) = 1$ is written as

$$y_1(x) = -\frac{1}{\pi \ln 4 \sqrt{x(1-x)}}. \quad (11)$$

Hence, equation (10) has the solution

$$u(x) = y_f(x) + Ay_1(x), \quad A = -\frac{\int_0^1 \psi(t)y_f(t) dt}{1 + \int_0^1 \psi(t)y_1(t) dt}.$$

Example 4. Consider the integral equation

$$\int_0^\infty [\cos(xt) + \varphi(x)\psi(t)]y(t) dt = f(x).$$

Its solution can be obtained by the methods described in Subsection 12.6-2; it must be taken into account that the truncated equation, with $\varphi(x) = 0$, coincides with equation 3.5.1 from Section 3.5. Therefore the solution is

$$y(t) = y_f(t) + Ay_\varphi(t),$$

where

$$y_f(t) = \frac{2}{\pi} \int_0^\infty \cos(xt)f(x) dx, \quad y_\varphi(t) = \frac{2}{\pi} \int_0^\infty \cos(xt)\varphi(x) dx, \quad A = -\frac{\int_0^\infty \psi(t)y_f(t) dt}{1 + \int_0^\infty \psi(t)y_\varphi(t) dt}.$$

2°. The integral equation

$$\int_a^b \left[K(x, t) + \sum_{m=1}^n \varphi_m(x)\psi_m(t) \right] u(t) dt = f(x) \quad (12)$$

whose kernel is the sum of the kernel of equation (4) and an arbitrary degenerate kernel can be solved in a similar manner. The solution is sought in the additive form

$$u(x) = y_f(x) + \sum_{m=1}^n A_m y_{\varphi_m}(x), \quad (13)$$

where $y_{\varphi_m}(x)$ is the solution to equation (4) in which $f(x)$ must be replaced with $\varphi_m(x)$. Substituting (13) into (12) results in the following linear algebraic system of equations for the coefficients A_m :

$$A_m + \sum_{j=1}^n A_j \sigma_{mj} = -\sigma_{m0}, \quad m = 1, \dots, n; \quad (14)$$

$$\sigma_{mj} = \int_a^b \psi_m(t) y_{\varphi_j}(t) dt, \quad \sigma_{m0} = \int_a^b \psi_m(t) y_f(t) dt.$$

Corollary. Given a solution to the integral equation (4) with a difference kernel $K(x, t) = K(x-t)$, one can obtain a solution to the integral equation with a difference kernel of the form

$$K(x-t) + P_n(x-t),$$

where $P_n(x)$ is an arbitrary polynomial of any (finite) degree n .

3°. Let a function $y(x)$ solve equation (4) and let the condition

$$\int_a^b K(x, t) dt = 0$$

be satisfied. Then the function

$$y(x) + C,$$

with C an arbitrary constant, also solves equation (4).

12.6-3. Reduction of Integral Equations of the First Kind to Equations of the Second Kind.

In some cases it is possible to reduce integral equations of the first kind with constant limits of integration to integral equations of the second kind.

In order to be definite, let us consider an integral equation of the first kind on semiaxis

$$\int_0^{\infty} [K(x, t)g(t) + L(x, t)]y(t) dt = f(x). \quad (15)$$

Suppose the truncated linear equation

$$\int_0^{\infty} K(x, t)u(t) dt = f(x) \quad (16)$$

obtained from (15) by setting $L(x, t) = 0$ and $g(t) = 1$, to be an integral transform (see Subsections 9.1-3 and 9.6-5) with the following inverse formula:

$$u(x) = \int_0^{\infty} M(x, s)f(s) ds. \quad (17)$$

Let us rewrite (15) in such a way, that its left-hand side coincides with (16):

$$\int_0^{\infty} K(x, t)u(t) dt = f(x) - \int_0^{\infty} L(x, t)y(t) dt, \quad u(t) = g(t)y(t). \quad (18)$$

Applying the inverse formula (17) to (18), provided that the function $f(x)$ must be substituted for $f(x) - \int_0^{\infty} L(x, t)y(t) dt$, and changing the integration order, we obtain an integral equation of the second kind with constant limits of integration

$$y(x) + \int_0^{\infty} N(x, t)y(t) dt = F(x), \quad (19)$$

where

$$N(x, t) = \frac{1}{g(x)} \int_0^{\infty} M(x, s)L(s, t) ds, \quad F(x) = \frac{1}{g(x)} \int_0^{\infty} M(x, s)f(s) ds.$$

Here, all integrals are supposed to converge.

Example 5. Consider the integral equation of the first kind

$$\int_0^\infty [\sin(xt) + L(x, t)]y(t) dt = f(x). \quad (20)$$

The solution of the truncated linear integral equation

$$\int_0^\infty \sin(xt)u(t) dt = f(x) \quad (21)$$

is expressed as (see equation 3.5.8 in Section 3.5)

$$u(x) = \frac{2}{\pi} \int_0^\infty \sin(xs)f(s) ds. \quad (22)$$

Up to constant factors, the function $f(x)$ and the solution $u(x)$ in (21)–(22) are the Fourier sine transform pair.

In accordance with the method described, the integral equation of the first kind with constant limits of integration (20) can be reduced to the integral equation of the second kind with constant limits of integration (19) where

$$N(x, t) = \frac{2}{\pi} \int_0^\infty \sin(xs)L(s, t) ds, \quad F(x) = \frac{2}{\pi} \int_0^\infty \sin(xs)f(s) ds.$$

☉ References for Section 12.6: N. Kh. Arutyunyan (1959), I. C. Gohberg and M. G. Krein (1967), F. D. Gakhov (1977, 1990), S. Fenyö and H. W. Stolle (1984, pp. 236–237), A. D. Polyaniin and A. I. Zhurov (2007).

12.7. Riemann Problem for the Real Axis

The Riemann boundary value problem is one of the main tools for constructing solutions of integral equations provided that various integral transforms can be applied to a given equation and the corresponding convolution-type theorems can be applied. This problem is investigated by an example of the Fourier integral transform.

12.7-1. Relationships Between the Fourier Integral and the Cauchy Type Integral.

Let $\mathcal{Y}(\tau)$ be a function integrable on a closed or nonclosed contour L on the complex plane of the variable $z = u + iv$ (τ is the complex coordinate of the contour points). Consider the integral of the Cauchy type (see Section 14.2):

$$\frac{1}{2\pi i} \int_L \frac{\mathcal{Y}(\tau)}{\tau - z} d\tau.$$

This integral defines a function that is analytic on the complex plane with a cut along the contour L . If L is a closed curve, then the integral is a function that is analytic on each of the connected parts of the plane bounded by L . If the contour L is the real axis, then we have

$$\frac{1}{2\pi i} \int_{-\infty}^\infty \frac{\mathcal{Y}(\tau)}{\tau - z} d\tau = \begin{cases} \mathcal{Y}^+(z) & \text{if } \text{Im } z > 0, \\ \mathcal{Y}^-(z) & \text{if } \text{Im } z < 0. \end{cases} \quad (1)$$

Moreover, there exist limit values of the functions $\mathcal{Y}^\pm(z)$ on the real axis, and these values are related to the density \mathcal{Y} of the integral by the *Sokhotski–Plemelj formulas*

$$\begin{aligned} \mathcal{Y}^+(u) &= \frac{1}{2}\mathcal{Y}(u) + \frac{1}{2\pi i} \int_{-\infty}^\infty \frac{\mathcal{Y}(\tau)}{\tau - u} d\tau, \\ \mathcal{Y}^-(u) &= -\frac{1}{2}\mathcal{Y}(u) + \frac{1}{2\pi i} \int_{-\infty}^\infty \frac{\mathcal{Y}(\tau)}{\tau - u} d\tau, \end{aligned} \quad (2)$$

or

$$\mathcal{Y}^+(u) - \mathcal{Y}^-(u) = \mathcal{Y}(u), \quad \mathcal{Y}^+(u) + \mathcal{Y}^-(u) = \frac{1}{\pi i} \int_{-\infty}^\infty \frac{\mathcal{Y}(\tau)}{\tau - u} d\tau. \quad (3)$$

In the latter formulas, the integral is understood as a singular integral in the sense of the Cauchy principal value.

In the Fourier integral*

$$\mathcal{Y}(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} y(x)e^{iux} dx,$$

the real parameter u occurs in an analytic function, and therefore we can replace u in this integral by a complex variable z . The function $\mathcal{Y}(z)$ defined by the integral

$$\mathcal{Y}(z) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} y(x)e^{izx} dx \quad (4)$$

is analytic in the part of the complex plane of the variable $z = u + iv$ in which the integral (4) is absolutely convergent. If this is a domain indeed, i.e., if it is not reduced to the real axis, then the integral (4) gives an analytic continuation of the Fourier integral into the complex plane. The integral (4) will also be called the *Fourier integral*.

Let us establish a relationship between this integral and the integral of the Cauchy type with density $\mathcal{Y}(u)$ taken along the entire axis. We have

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\mathcal{Y}(\tau)}{\tau - z} d\tau = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} y(x)e^{izx} dx, \quad \text{Im } z > 0, \quad (5)$$

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\mathcal{Y}(\tau)}{\tau - z} d\tau = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 y(x)e^{izx} dx, \quad \text{Im } z < 0. \quad (6)$$

12.7-2. One-Sided Fourier Integrals.

If $\mathcal{Y}(z) = \mathcal{Y}^+(z)$ is an analytic function in the upper half-plane whose limit value on the real axis is given by the function $\mathcal{Y}(u) = \mathcal{Y}^+(u) \in L_2(-\infty, \infty)$, then the function $\mathcal{Y}^+(z)$ can be expressed by means of the Cauchy integral. Hence, by virtue of (5) we have

$$\mathcal{Y}^+(z) = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} y(x)e^{izx} dx,$$

and, since the integral defines a continuous function, the limit values on the axis can be obtained from the last relation merely by setting $z = u$:

$$\mathcal{Y}^+(u) = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} y(x)e^{iux} dx,$$

where, according to (5), $y(x)$ the inverse transform of $\mathcal{Y}(u)$. The right-hand side can be regarded as the Fourier integral of a function that is identically zero for negative x . Hence, by the uniqueness of the representation of the function $\mathcal{Y}^+(u)$ by a Fourier integral, it follows that $y(x) \equiv 0$ on the negative semiaxis.

Conversely, if $y \equiv 0$ for $x < 0$, then the Fourier integral of this function becomes

$$\mathcal{Y}(u) = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} y(x)e^{iux} dx.$$

* In Sections 12.7–12.9, the alternative Fourier transform is used (see Subsection 9.4-3).

If we replace the parameter u by a complex number z belonging to the upper half-plane, then the integral will converge even better. This implies the analyticity of the function

$$\mathcal{Y}(z) = \frac{1}{\sqrt{2\pi}} \int_0^\infty y(x)e^{izx} dx$$

in the upper half-plane.

The case of the lower half-plane can be treated in a similar way.

The integrals

$$\mathcal{Y}^+(z) = \frac{1}{\sqrt{2\pi}} \int_0^\infty y(x)e^{izx} dx, \quad \mathcal{Y}^-(z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 y(x)e^{izx} dx \quad (7)$$

are called *one-sided Fourier integrals*, namely, the *right* and the *left* Fourier integral, respectively. As well as in formula (1), the symbols \pm over symbols of functions mean that the corresponding function is analytic in the upper or lower half-plane, respectively.

Let us introduce the functions

$$y_+(x) = \begin{cases} y(x) & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad y_-(x) = \begin{cases} 0 & \text{for } x > 0, \\ -y(x) & \text{for } x < 0. \end{cases} \quad (8)$$

These functions are said to be *one-sided functions* for $y(x)$, namely, the *right function* and the *left function*, respectively. Obviously, the following relation holds:

$$y(x) = y_+(x) - y_-(x). \quad (9)$$

Applying the well-known function $\text{sign } x$ defined by

$$\text{sign } x = \begin{cases} 1 & \text{for } x > 0, \\ -1 & \text{for } x < 0, \end{cases} \quad (10)$$

we can express y_\pm in terms of y as follows:

$$y_\pm(x) = \frac{1}{2}(\pm 1 + \text{sign } x)y(x). \quad (11)$$

The symbols \pm on symbols of one-sided functions will be always subscripts.

The Fourier integrals of the right and left one-sided functions are the boundary values of functions that are analytic on the upper and lower half-planes, respectively.

Let us indicate the following analogs of the Sokhotski–Plemelj formulas (3) in the Fourier integrals:

$$\begin{aligned} \mathcal{Y}(u) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty y(x)e^{iux} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_0^\infty y(x)e^{iux} dx + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 y(x)e^{iux} dx = \mathcal{Y}^+(u) - \mathcal{Y}^-(u), \\ \frac{1}{\pi i} \int_{-\infty}^\infty \frac{\mathcal{Y}(\tau)}{\tau - u} d\tau &= \mathcal{Y}^+(u) + \mathcal{Y}^-(u) \\ &= \frac{1}{\sqrt{2\pi}} \int_0^\infty y(x)e^{iux} dx - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 y(x)e^{iux} dx = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty y(x) \text{sign } x e^{iux} dx. \end{aligned} \quad (12)$$

Thus, in this setting, the first Sokhotski–Plemelj formula (a representation of an arbitrary function in the form of the difference of boundary values of analytic functions) is an obvious consequence of the decomposition of a Fourier integral into the right and the left integral. The second formula can also be rewritten as follows:

$$\mathbf{F}\{y(x) \text{sign } x\} = \frac{1}{\pi i} \int_{-\infty}^\infty \frac{\mathcal{Y}(\tau)}{\tau - u} d\tau, \quad \mathbf{F}^{-1}\left\{\frac{1}{\pi i} \int_{-\infty}^\infty \frac{\mathcal{Y}(\tau)}{\tau - u} d\tau\right\} = y(x) \text{sign } x. \quad (13)$$

12.7-3. Analytic Continuation Theorem and the Generalized Liouville Theorem.

Below is the analytic continuation theorem and the generalized Liouville theorem combined into a single statement, which will be used in Chapters 12 and 13.

Let functions $\mathcal{Y}_1(z)$ and $\mathcal{Y}_2(z)$ be analytic in the upper and lower half-planes, respectively, possibly except for a point $z_* \neq \infty$, at which these functions have a pole. If $\mathcal{Y}_1(z)$ and $\mathcal{Y}_2(z)$ are bounded at infinity, the principal parts of their expansions in a neighborhood of z_* have the form

$$\frac{c_1}{z - z_*} + \frac{c_2}{(z - z_*)^2} + \cdots + \frac{c_m}{(z - z_*)^m} \equiv \frac{\mathcal{P}_{m-1}(z)}{(z - z_*)^m},$$

and if the functions themselves coincide on the real axis, then these functions represent a single rational function on the entire plane:

$$\mathcal{Y}(z) = c_0 + \frac{\mathcal{P}_{m-1}(z)}{(z - z_*)^m},$$

where c_0 is a constant. The pole z_* can belong either to the open half-planes or to the real axis.

Let us also give a more general version of the above statement.

If functions $\mathcal{Y}_1(z)$ and $\mathcal{Y}_2(z)$ are analytic in the upper and lower half-planes, respectively, possibly except for finitely many points $z_0 = \infty, z_k$ ($k = 1, \dots, n$), at which these functions can have poles if the principal parts of the expansions of these functions in a neighborhood of a pole have the form

$$\begin{aligned} c_1^0 z + c_2^0 z^2 + \cdots + c_{m_0}^0 z^{m_0} &\equiv \mathcal{P}_0(z) && \text{at the point } z_0, \\ \frac{c_1^k}{z - z_k} + \frac{c_2^k}{(z - z_k)^2} + \cdots + \frac{c_{m_k}^k}{(z - z_k)^{m_k}} &\equiv \frac{\mathcal{P}_{m_k-1}(z)}{(z - z_k)^{m_k}} && \text{at the points } z_k, \end{aligned}$$

and if the functions themselves coincide on the real axis, then these functions represent a single rational function on the entire plane:

$$\mathcal{Y}(z) = C + \mathcal{P}_0(z) + \sum_{k=1}^n \frac{\mathcal{P}_{m_k-1}(z)}{(z - z_k)^{m_k}}$$

where C is a constant. The poles z_k can belong either to the open half-planes or to the real axis.

12.7-4. Riemann Boundary Value Problem.

The solution of the Riemann problem in this section differs from the traditional one, because it is expressed not by means of integrals of the Cauchy type (see Subsection 14.3-8) but by means of Fourier integrals. To solve equations of convolution type under consideration, the Fourier integral technique is more convenient.

By the *index* of a continuous complex-valued nonvanishing function $\mathcal{M}(u)$ ($\mathcal{M}(u) = \mathcal{M}_1(u) + i\mathcal{M}_2(u)$, $-\infty < u < \infty$, $\mathcal{M}(-\infty) = \mathcal{M}(\infty)$) we mean the variation of the argument of this function on the real axis expressed in the number of full rotations:

$$\text{Ind } \mathcal{M}(u) = \frac{1}{2\pi} [\arg \mathcal{M}(u)]_{-\infty}^{\infty} = \frac{1}{2\pi i} [\ln \mathcal{M}(u)]_{-\infty}^{\infty} = \frac{1}{2\pi i} \int_{-\infty}^{\infty} d \ln \mathcal{M}(u).$$

If $\mathcal{M}(u)$ is not differentiable but is of bounded variation, then the last integral must be understood as the Stieltjes integral.

If an analytic function $\mathcal{Y}(z)$ has a representation of the form

$$\mathcal{Y}(z) = (z - z_0)^m \mathcal{Y}_1(z)$$

in a neighborhood of some point z_0 , where $\mathcal{Y}_1(z)$ is analytic and $\mathcal{Y}_1(z_0) \neq 0$, then the integer m (which can be positive, negative, or zero) is called the *order* of the function $\mathcal{Y}(z)$ at the point z_0 .

If $m > 0$, then the order of the function is the order of its zero, and if $m < 0$, then the order of the function is minus the order of its pole. If the order of the function at z_0 is zero, then at this point the function takes a finite nonzero value. When considering the point at infinity we must replace the difference $z - z_0$ by $1/z$.

Let us pose the Riemann problem. Let two functions be given on the real axis, namely, $\mathcal{D}(u)$, the *coefficient of the problem*, and $\mathcal{H}(u)$, the *right-hand side*, and let the following normality condition hold: $\mathcal{D}(u) \neq 0$. The functions $\mathcal{H}(u)$ and $\mathcal{D}(u) - 1$ belong to $L_2(-\infty, \infty)$ and simultaneously satisfy the Hölder condition. The problem is to find two functions $\mathcal{Y}^\pm(z)$ that are analytic in the upper and the lower half-plane, respectively,* whose limit values on the real axis satisfy the following boundary condition:

$$\mathcal{Y}^+(u) = \mathcal{D}(u)\mathcal{Y}^-(u) + \mathcal{H}(u). \quad (14)$$

It follows from the representation of $\mathcal{D}(u)$ that $\mathcal{D}(\infty) = 1$. The last condition implies no loss of generality of subsequent reasoning because by dividing the boundary condition (14) by $\mathcal{D}(\infty)$ we can always obtain the necessary form of the problem.**

If $\mathcal{D}(u) \equiv 1$, then the Riemann problem is called the *jump problem*. For $\mathcal{H}(u) \equiv 0$, the Riemann problem is said to be *homogeneous*. The index ν of the coefficient $\mathcal{D}(u)$ of the boundary value problem is called the *index of the Riemann problem*.

Consider the *jump problem*, i.e., the problem of finding $\mathcal{Y}^\pm(z)$ from the boundary condition

$$\mathcal{Y}^+(u) - \mathcal{Y}^-(u) = \mathcal{H}(u). \quad (15)$$

The solution of this problem is given by the first formula in (12):

$$\mathcal{Y}^+(z) = \frac{1}{\sqrt{2\pi}} \int_0^\infty H(x)e^{izx} dx, \quad \mathcal{Y}^-(z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 H(x)e^{izx} dx, \quad (16)$$

where

$$H(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \mathcal{H}(u)e^{-iux} du. \quad (17)$$

Let us construct a particular solution $\mathcal{X}(z)$ of the homogeneous Riemann problem (14), which we need in what follows:

$$\mathcal{X}^+(u) = \mathcal{D}(u)\mathcal{X}^-(u), \quad \mathcal{D}(\infty) = 1, \quad (18)$$

where $\mathcal{X}(z)$ is assumed to be nonzero on the real axis with the additional condition $\mathcal{X}^\pm(\infty) = 1$. Denote by N_+ and N_- the numbers of zeros of the functions $\mathcal{X}^+(z)$ and $\mathcal{X}^-(z)$ in the upper and lower half-planes, respectively. On calculating the index of both sides of the boundary condition (18) and applying the properties of the index, we obtain

$$N_+ + N_- = \text{Ind } \mathcal{D}(u) = \nu. \quad (19)$$

We first assume that $\nu = 0$. In this case, $\ln \mathcal{D}(u)$ is a single-valued function. It follows from relation (19) that $N_+ = N_- = 0$, i.e., the solution has no zeros on the entire plane. Therefore, the functions $\ln \mathcal{X}^+(z)$ and $\ln \mathcal{X}^-(z)$ are analytic in the corresponding half-planes, and hence are single-valued together with their boundary values $\ln \mathcal{X}^+(u)$ and $\ln \mathcal{X}^-(u)$. Taking the logarithm of the boundary condition (18), we obtain

$$\ln \mathcal{X}^+(u) - \ln \mathcal{X}^-(u) = \ln \mathcal{D}(u). \quad (20)$$

* A couple of functions $\mathcal{Y}^\pm(z)$ can be treated as a single function $\mathcal{Y}(z)$ piecewise analytic in the entire complex plane. In some cases, we use the latter notation.

** Since the boundary condition is the main analytic expression of the Riemann problem, in references to the corresponding problem we shall often indicate its boundary condition only and write, for instance, "Riemann problem (14)."

On choosing a branch of $\ln \mathcal{D}(u)$ such that $\ln \mathcal{D}(\infty) = 0$ (it can be shown that the final result does not depend on the choice of the branch) we arrive at a jump problem. In this case, on the basis of (15)–(17) and (20), the solution of problem (18) can be represented in the form

$$\begin{aligned} \mathcal{X}^+(z) &= e^{\mathcal{G}^+(z)}, \quad \mathcal{X}^-(z) = e^{\mathcal{G}^-(z)}, \\ \mathcal{G}^+(z) &= \frac{1}{\sqrt{2\pi}} \int_0^\infty g(x)e^{izx} dx, \quad \mathcal{G}^-(z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 g(x)e^{izx} dx, \\ g(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \ln \mathcal{D}(u) e^{-iux} du. \end{aligned} \quad (21)$$

Relations (21) imply the following important fact: a function $\mathcal{D}(u)$ of zero index that is nonvanishing on the real axis and satisfies the condition $\mathcal{D}(\infty) = 1$ can be represented as the ratio of functions that are the boundary values of nonzero analytic functions in the upper and the lower half-plane, respectively.

Let us pass to the case in which the index of the homogeneous Riemann problem (18) is arbitrary. By a *canonical function* $\mathcal{X}(z)$ (of the homogeneous Riemann problem) we mean a function that satisfies the boundary condition (18) and the condition $\mathcal{X}^\pm(\infty) = 1$ and has zero order everywhere possibly except for the point $-i$, at which the order of $\mathcal{X}(z)$ is equal to the index ν of the Riemann problem. Such a function can be constructed by reducing the homogeneous Riemann problem to the above case of zero index. Indeed, let us write out the boundary condition of the homogeneous Riemann problem (18) in the form

$$\mathcal{X}^+(u) = \left[\left(\frac{u-i}{u+i} \right)^{-\nu} \mathcal{D}(u) \right] \left[\left(\frac{u-i}{u+i} \right)^\nu \mathcal{X}^-(u) \right]. \quad (22)$$

In this case, the function in the first square brackets has zero index and can be represented as the ratio of the boundary values of functions that are analytic in the upper and the lower half-plane. This, together with the boundary condition (22), gives the following expression for the canonical function:

$$\begin{aligned} \mathcal{X}^+(z) &= e^{\mathcal{G}^+(z)}, \quad \mathcal{X}^-(z) = \left(\frac{z-i}{z+i} \right)^{-\nu} e^{\mathcal{G}^-(z)}, \\ \mathcal{G}^+(z) &= \frac{1}{\sqrt{2\pi}} \int_0^\infty g(x)e^{izx} dx, \quad \mathcal{G}^-(z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 g(x)e^{izx} dx, \\ g(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \ln \left[\left(\frac{u-i}{u+i} \right)^{-\nu} \mathcal{D}(u) \right] e^{-iux} du, \end{aligned} \quad (23)$$

where, at the point $-i$, $\mathcal{X}^-(z)$ has a zero of order ν for $\nu > 0$ and a pole of order $|\nu|$ for the case $\nu < 0$.

The coefficient $\mathcal{D}(u)$ of the Riemann boundary value problem can be represented as the ratio of the boundary values of the canonical function (see (22) and (23)):

$$\mathcal{D}(u) = \frac{\mathcal{X}^+(u)}{\mathcal{X}^-(u)}. \quad (24)$$

Such a representation of $\mathcal{D}(u)$ in the form of the ratio of boundary values of the canonical function is often called a *factorization*.

Now we consider the homogeneous Riemann problem with the boundary condition

$$\mathcal{Y}^+(u) = \mathcal{D}(u)\mathcal{Y}^-(u), \quad \mathcal{D}(\infty) = 1. \quad (25)$$

On substituting the expression (24) for $\mathcal{D}(u)$ into (25) we reduce the boundary condition to the form

$$\frac{\mathcal{Y}^+(u)}{\mathcal{X}^+(u)} = \frac{\mathcal{Y}^-(u)}{\mathcal{X}^-(u)}. \quad (26)$$

According to formulas (23) for $\mathcal{X}(z)$, the left- and the right-hand sides of Eq. (26) contain the boundary values of functions that are analytic on the upper and lower half-planes, respectively, possibly except for the point $-i$ at which the order is equal to ν . In the chosen function class, each function vanishes at infinity. In this case, it follows from the analytic continuation theorem and the generalized Liouville theorem (see Subsection 12.7-3) that for $\nu > 0$ we have

$$\frac{\mathcal{Y}^+(z)}{\mathcal{X}^+(z)} = \frac{\mathcal{Y}^-(z)}{\mathcal{X}^-(z)} = \frac{\mathcal{P}_{\nu-1}(z)}{(z+i)^\nu}, \quad (27)$$

where $\mathcal{P}_{\nu-1}(z)$ is an arbitrary polynomial of degree $\nu - 1$ (the degree of the numerator is less than that of the denominator because $\mathcal{Y}(\infty) = 0$). Hence,

$$\mathcal{Y}(z) = \mathcal{X}(z) \frac{\mathcal{P}_{\nu-1}(z)}{(z+i)^\nu}. \quad (28)$$

For $\nu \leq 0$, it follows from $\mathcal{Y}(\infty) = 0$ that $\mathcal{Y}(z) \equiv 0$ by the generalized Liouville theorem.

Hence, for $\nu > 0$, the homogeneous Riemann boundary value problem has precisely ν linearly independent solutions of the form

$$\frac{z^{k-1} \mathcal{X}(z)}{(z+i)^\nu}, \quad k = 1, 2, \dots, \nu,$$

and for $\nu \leq 0$, there are no nontrivial solutions.

The right-hand side of Eq. (28) has exactly ν zeros on the entire plane, including the zero at infinity. These zeros can lie at arbitrary points of the upper and lower half-plane or on the real axis. Denote the number of zeros on the real axis by N_0 . In the general case (without the requirement that there are no zeros on the real axis), formula (19) is replaced by the relation

$$N_+ + N_- + N_0 = \text{Ind } \mathcal{D}(u) = \nu. \quad (29)$$

Let us pass to the solution of the nonhomogeneous Riemann problem with the boundary condition (14). We apply relation (24) and reduce the boundary condition to the form

$$\frac{\mathcal{Y}^+(u)}{\mathcal{X}^+(u)} = \frac{\mathcal{Y}^-(u)}{\mathcal{X}^-(u)} + \frac{\mathcal{H}(u)}{\mathcal{X}^+(u)}. \quad (30)$$

Let us express the last summand as the difference of the boundary values of functions that are analytic in the upper and the lower half-plane (see the jump problem), that is,

$$\mathcal{W}^+(u) - \mathcal{W}^-(u) = \frac{\mathcal{H}(u)}{\mathcal{X}^+(u)}, \quad (31)$$

where

$$\begin{aligned} \mathcal{W}^+(z) &= \frac{1}{\sqrt{2\pi}} \int_0^\infty w(x) e^{izx} dx, & \mathcal{W}^-(z) &= -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 w(x) e^{izx} dx, \\ w(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \frac{\mathcal{H}(u)}{\mathcal{X}^+(u)} e^{-iux} du. \end{aligned} \quad (32)$$

On substituting (31) into (30), we obtain

$$\frac{\mathcal{Y}^+(u)}{\mathcal{X}^+(u)} - \mathcal{W}^+(u) = \frac{\mathcal{Y}^-(u)}{\mathcal{X}^-(u)} - \mathcal{W}^-(u). \quad (33)$$

For $\nu > 0$, it follows from the analytic continuation theorem and the generalized Liouville theorem that

$$\frac{\mathcal{Y}^+(z)}{\mathcal{X}^+(z)} - \mathcal{W}^+(z) = \frac{\mathcal{Y}^-(z)}{\mathcal{X}^-(z)} - \mathcal{W}^-(z) = \frac{\mathcal{P}_{\nu-1}(z)}{(z+i)^\nu}.$$

Hence, for $\nu > 0$ we have

$$\mathcal{Y}(z) = \mathcal{X}(z) \left[\mathcal{W}(z) + \frac{\mathcal{P}_{\nu-1}(z)}{(z+i)^\nu} \right]. \tag{34}$$

The right-hand side of formula (34) contains the general solution (28) of the homogeneous problem as a summand, and hence the general solution of the nonhomogeneous problem is obtained.

For $\nu \leq 0$ we must set $\mathcal{P}_{\nu-1}(z) \equiv 0$, and the desired solution becomes

$$\mathcal{Y}(z) = \mathcal{X}(z)\mathcal{W}(z). \tag{35}$$

However, formula (35) gives a solution that satisfies all conditions for $\nu = 0$ only. For $\nu < 0$, the function $\mathcal{X}(z)$ has a pole of order $|\nu|$ at the point $-i$. In this case, for the existence of a solution in the chosen class of functions it is necessary that the second factor have a zero of the corresponding order at the point $-i$. On the basis of relations (6) and (32), we represent the function $\mathcal{W}^-(z)$ in the form

$$\mathcal{W}^-(z) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\mathcal{H}(\tau)}{\mathcal{X}^+(\tau)} \frac{d\tau}{\tau - z}.$$

On expanding the last integral in series in powers of $z + i$ and equating the coefficients of $(z + i)^{k-1}$ ($k = 1, 2, \dots, |\nu|$) with zero, we obtain the solvability conditions for the problem in the form

$$\int_{-\infty}^{\infty} \frac{\mathcal{H}(u)}{\mathcal{X}^+(u)} \frac{du}{(u+i)^k} = 0, \quad k = 1, 2, \dots, |\nu|. \tag{36}$$

Figure 5 depicts a scheme of the above method for solving the Riemann problem on the real axis.

Let us state the results concerning the solution of the Riemann problem in the final form. If the index ν of the problem satisfies the condition $\nu > 0$, then the homogeneous and the nonhomogeneous Riemann problems are unconditionally solvable, and their solutions

$$\mathcal{Y}^\pm(z) = \mathcal{X}^\pm(z) \frac{\mathcal{P}_{\nu-1}(z)}{(z+i)^\nu} \quad (\text{the homogeneous problem}), \tag{37}$$

$$\mathcal{Y}^\pm(z) = \mathcal{X}^\pm(z) \left[\mathcal{W}^\pm(z) + \frac{\mathcal{P}_{\nu-1}(z)}{(z+i)^\nu} \right] \quad (\text{the nonhomogeneous problem}) \tag{38}$$

depend on ν arbitrary complex constants, where $\mathcal{P}_{\nu-1}(z)$ is a polynomial of degree $\nu - 1$. If $\nu \leq 0$, then the homogeneous problem has only the trivial zero solution, and the nonhomogeneous problem has the unique solution

$$\mathcal{Y}^\pm(z) = \mathcal{X}^\pm(z)\mathcal{W}^\pm(z) \tag{39}$$

provided that $|\nu|$ conditions (36) hold. Here we have

$$g(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \ln \left[\left(\frac{u-i}{u+i} \right)^{-\nu} \mathcal{D}(u) \right] e^{-iux} du, \tag{40}$$

$$\mathcal{G}^+(z) = \frac{1}{\sqrt{2\pi}} \int_0^\infty g(x)e^{izx} dx, \quad \mathcal{G}^-(z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 g(x)e^{izx} dx, \tag{41}$$

$$\mathcal{X}^+(z) = e^{\mathcal{G}^+(z)}, \quad \mathcal{X}^-(z) = \left(\frac{z-i}{z+i} \right)^{-\nu} e^{\mathcal{G}^-(z)}, \tag{42}$$

$$w(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\mathcal{H}(u)}{\mathcal{X}^+(u)} e^{-iux} du, \tag{43}$$

$$\mathcal{W}^+(z) = \frac{1}{\sqrt{2\pi}} \int_0^\infty w(x)e^{izx} dx, \quad \mathcal{W}^-(z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 w(x)e^{izx} dx. \tag{44}$$




Figure 5. Scheme of solving the Riemann boundary value problem for the functions $\mathcal{Y}^+(z)$ and $\mathcal{Y}^-(z)$ that are analytic, respectively, in the upper and the lower half-plane of the complex plane $z = u + iv$. It is assumed that $\mathcal{D}(u) \neq 0$ and $\mathcal{P}_{\nu-1}(z) \equiv 0$ for $\nu \leq 0$.

The sequence of operations to construct a solution can be described as follows.

1°. By virtue of formula (40) we find $g(x)$, and then, with the help of (41), for the given $g(x)$ we find $\mathcal{G}^\pm(z)$.

2°. By formulas (42) the canonical function $\mathcal{X}^\pm(z)$ is determined.

3°. By formula (43) we determine $w(x)$, and then apply formula (44) to find $\mathcal{W}^\pm(z)$.

After this, solutions of the homogeneous and nonhomogeneous problems can be found by formulas (37)–(39) and (42). For the case $\nu < 0$, it is also necessary to verify the solvability conditions (36).

12.7-5. Problems with Rational Coefficients.

The solution of the Riemann problem thus obtained requires evaluation of several Fourier integrals. This can also be readily expressed by means of integrals of the Cauchy type. As a rule, the integrals cannot be evaluated in the closed form and are calculated by various approximate methods. This process is rather cumbersome, and therefore it is of interest to select cases in which the solution can be obtained directly from the boundary condition by applying the method of analytic continuation without using the antiderivatives.

Assume that in the boundary condition (14) we have

$$\mathcal{D}(u) = \frac{\mathcal{R}_+(u) \mathcal{R}_-(u)}{\mathcal{Q}_+(u) \mathcal{Q}_-(u)}.$$

Here $\mathcal{R}_+(u)$ and $\mathcal{Q}_+(u)$ ($\mathcal{R}_-(u)$ and $\mathcal{Q}_-(u)$) are polynomials whose zeros belong to the upper (lower) half-plane (we must avoid confusing these polynomials with the one-sided functions introduced above, which have similar notation). Denote the degrees of the polynomials \mathcal{P}_+ , \mathcal{R}_+ , \mathcal{Q}_+ , and \mathcal{Q}_- by m_+ , m_- , n_+ , and n_- , respectively. Since, by the assumption of the problem, the value $\mathcal{D}(\infty)$ can be neither zero nor infinity, it follows that the relation $m_+ + m_- = n_+ + n_-$ holds. The index of the problem can be expressed by the formula

$$\nu = \text{Ind } \mathcal{D}(u) = m_+ - n_+ = -(m_- - n_-).$$

On multiplying the boundary condition by $\mathcal{Q}_-(u)/\mathcal{P}_-(u)$ we obtain

$$\frac{\mathcal{Q}_-(u)}{\mathcal{R}_-(u)} \mathcal{Y}^+(u) - \frac{\mathcal{R}_+(u)}{\mathcal{Q}_+(u)} \mathcal{Y}^-(u) = \frac{\mathcal{Q}_-(u)}{\mathcal{R}_-(u)} \mathcal{H}(u).$$

If $\mathcal{H}(u)$ is a rational function as well, then the jump problem can readily be solved:

$$\mathcal{W}^+(u) - \mathcal{W}^-(u) = \frac{\mathcal{Q}_-(u)}{\mathcal{R}_-(u)} \mathcal{H}(u). \quad (45)$$

To this end, it suffices to decompose the right-hand side into the sum of partial fractions. Then $\mathcal{W}^+(u)$ and $\mathcal{W}^-(u)$ are the sums of the partial fractions with poles in the lower and the upper half-planes, respectively. We can directly apply the continuity principle (the analytic continuation theorem) and the generalized Liouville theorem to the resulting relation

$$\frac{\mathcal{Q}_-(u)}{\mathcal{R}_-(u)} \mathcal{Y}^+(u) - \mathcal{W}^+(u) = \frac{\mathcal{R}_+(u)}{\mathcal{Q}_+(u)} \mathcal{Y}^-(u) - \mathcal{W}^-(u).$$

The only exceptional point at which the analytic function, which is the same on the entire complex plane, can have a nonzero order is the point at infinity, at which the order of the function is equal to $\nu - 1 = m_+ - n_+ - 1 = n_- - m_- - 1$.

For $\nu > 0$, the solution can be written in the form

$$\mathcal{Y}^+(z) = \frac{\mathcal{R}_-(z)}{\mathcal{Q}_-(z)} [\mathcal{W}^+(z) + \mathcal{P}_{\nu-1}(z)], \quad \mathcal{Y}^-(z) = \frac{\mathcal{Q}_+(z)}{\mathcal{R}_+(z)} [\mathcal{W}^-(z) + \mathcal{P}_{\nu-1}(z)].$$

For $\nu \leq 0$ we must set $\mathcal{P}_{\nu-1} \equiv 0$; moreover, for $\nu < 0$ we must also write out the solvability conditions that can be obtained by equating with zero the first $|\nu|$ terms of the expansion of the rational function $\mathcal{W}(z)$ in a series (in powers of $1/z$) in a neighborhood of the point at infinity.

The solution of the jump problem (45) can be obtained either by applying the method of indeterminate coefficients, as is usually performed in the integration of rational functions, or using the theory of residuals of analytic functions. Let z_k be a pole, of multiplicity m , of the function $[\mathcal{Q}_-(z)/\mathcal{R}_-(z)]\mathcal{H}(z)$. Then the coefficients of the principal part of the decomposition of this function in a neighborhood of the point z_k , which has the form

$$\frac{c_1^k}{z - z_k} + \cdots + \frac{c_m^k}{(z - z_k)^m},$$

can be found by the formula

$$c_j^k = \frac{1}{(j-1)!} \frac{d^{j-1}}{dz^{j-1}} \left[\frac{\mathcal{Q}_-(z)}{\mathcal{R}_-(z)} \mathcal{H}(z) \right]_{z=z_k}.$$

The above case is not only of independent interest, as it frequently occurs in practice, but also of importance as a possible way of solving the problem under general assumptions. The approximation of arbitrary coefficients of the class under consideration by rational functions is a widespread method of approximate solution of the Riemann boundary value problem.

12.7-6. Exceptional Cases. The Homogeneous Problem.

Assume that the coefficient $\mathcal{D}(u)$ of a Riemann boundary value problem has zeros of orders $\alpha_1, \dots, \alpha_r$ at points a_1, \dots, a_r , respectively, and poles* of the orders β_1, \dots, β_s at points b_1, \dots, b_s ($\alpha_1, \dots, \alpha_r$ and β_1, \dots, β_s are positive integers). Thus, the coefficient can be represented in the form

$$\mathcal{D}(u) = \frac{\prod_{i=1}^r (u - a_i)^{\alpha_i}}{\prod_{j=1}^s (u - b_j)^{\beta_j}} \mathcal{D}_1(u), \quad \mathcal{D}_1(u) \neq 0, \quad -\infty < u < \infty, \quad \sum_{i=1}^r \alpha_i = m, \quad \sum_{j=1}^s \beta_j = n. \quad (46)$$

In turn, we represent the function $\mathcal{D}_1(u)$ (see Subsection 12.7-5) in the form

$$\mathcal{D}_1(u) = \frac{\mathcal{R}_+(u)\mathcal{R}_-(u)}{\mathcal{Q}_+(u)\mathcal{Q}_-(u)} \mathcal{D}_2(u), \quad (47)$$

where, as above, $\mathcal{R}_+(u)$ and $\mathcal{Q}_+(u)$ ($\mathcal{R}_-(u)$ and $\mathcal{Q}_-(u)$) are polynomials of degrees m_+ and n_+ (m_- and n_-) whose zeros belong to the upper (lower) half-plane. The function $\mathcal{D}_2(u)$ satisfies the Hölder condition, has zero index, and nowhere vanishes on the real axis. Moreover, this function can be subjected to some differentiability conditions in neighborhoods of the points a_i and b_j and possibly in a neighborhood of the point at infinity.

The boundary condition of the homogeneous Riemann problem can be rewritten in the form

$$\mathcal{Y}^+(u) = \frac{\prod_{i=1}^r (u - a_i)^{\alpha_i} \mathcal{R}_+(u)\mathcal{R}_-(u)}{\prod_{j=1}^s (u - b_j)^{\beta_j} \mathcal{Q}_+(u)\mathcal{Q}_-(u)} \mathcal{D}_2(u)\mathcal{Y}^-(u). \quad (48)$$

* For the case in which the function $\mathcal{D}(u)$ is not analytic, the term "pole" will be used for points at which the function tends to infinity with integer order.

We seek a solution in the class of functions that are bounded on the real axis and vanish at infinity:

$$\mathcal{Y}(\infty) = 0. \quad (49)$$

The coefficient $\mathcal{D}(u)$ has the order

$$\eta = n + n_+ + n_- - m - m_+ - m_- \quad (50)$$

at infinity. The number

$$\nu = m_+ - n_+ \quad (51)$$

is called the *index* of the problem. Let us introduce the notation

$$h = n_- - m_-. \quad (52)$$

Then the order at infinity is expressed by the formula

$$\eta = h - \nu + n - m. \quad (53)$$

Now let us proceed with the solution of problem (48). Applying general methods, we set

$$\begin{aligned} \mathcal{D}_2(u) &= \frac{e^{\mathcal{G}^+(u)}}{e^{\mathcal{G}^-(u)}}, \quad g(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \ln \mathcal{D}_2(u) e^{-iux} du, \\ \mathcal{G}^+(z) &= \frac{1}{\sqrt{2\pi}} \int_0^{\infty} g(x) e^{izx} dx, \quad \mathcal{G}^-(z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 g(x) e^{izx} dx \end{aligned} \quad (54)$$

and rewrite the boundary condition in the form

$$\frac{\prod_{i=1}^r (u - a_i)^{\alpha_i} \mathcal{R}_-(u) e^{\mathcal{G}^+(u)}}{\prod_{j=1}^s (u - b_j)^{\beta_j} \mathcal{Q}_+(u) e^{\mathcal{G}^-(u)}} = \frac{\mathcal{R}_+(u) \mathcal{Y}^-(u)}{\mathcal{Q}_-(u) \mathcal{Y}^+(u)}. \quad (55)$$

As above, we can apply the analytic continuation and the generalized Liouville theorem and obtain a pole at infinity as the only possible singularity.

Two cases are possible:

1°. Let the order η of the coefficient of the boundary value problem at infinity satisfy the condition $\eta \geq 0$, i.e., let $\mathcal{D}(u)$ have a zero of order η at infinity. It follows from (53) that $n - \nu \geq m - h$. On equating the left- and right-hand sides of relation (55) with a polynomial $\mathcal{P}_{\nu-n-1}(z)$, we obtain the solution of the boundary value problem in the form

$$\begin{aligned} \mathcal{Y}^+(z) &= \prod_{i=1}^r (z - a_i)^{\alpha_i} \frac{\mathcal{R}_-(z)}{\mathcal{Q}_-(z)} e^{\mathcal{G}^+(z)} \mathcal{P}_{\nu-n-1}(z), \\ \mathcal{Y}^-(z) &= \prod_{j=1}^s (z - b_j)^{\beta_j} \frac{\mathcal{Q}_+(z)}{\mathcal{R}_+(z)} e^{\mathcal{G}^-(z)} \mathcal{P}_{\nu-n-1}(z). \end{aligned} \quad (56)$$

This problem has $\nu - n$ linearly independent solutions for $\nu - n > 0$ and only the trivial zero solution for $\nu - n \leq 0$.

2°. Let $\eta < 0$, i.e., let $\mathcal{D}(u)$ have a pole of order $-\eta$ at infinity. In this case, $m - h > n - \nu$, and we can obtain the general solution from (56) by replacing $\mathcal{P}_{\nu-n-1}(z)$ by $\mathcal{P}_{h-m-1}(z)$ in this expression. In this case, the problem has $h - m$ solutions for $h - m > 0$ and only the trivial zero solution for $h - m \leq 0$.

According to (53), we have

$$h - m = \nu - n + \eta. \quad (57)$$

Thus, in both cases under consideration, the number of linearly independent solutions is equal to the index minus the total number of the poles (including the pole at infinity) of the coefficient $\mathcal{D}(u)$. Hence, we have the following law: the number of linearly independent solutions of a homogeneous Riemann problem is not affected by the number of zeros of the coefficient and is reduced by the total number of its poles.

12.7-7. Exceptional Cases. The Nonhomogeneous Problem.

Assume that the right-hand side has the same poles as the coefficient. The boundary condition can be rewritten as follows:

$$\mathcal{Y}^+(u) = \frac{\prod_{i=1}^r (u - a_i)^{\alpha_i} \mathcal{R}_+(u) \mathcal{R}_-(u)}{\prod_{j=1}^s (u - b_j)^{\beta_j} \mathcal{Q}_+(u) \mathcal{Q}_-(u)} \mathcal{D}_2(u) \mathcal{Y}^-(u) + \frac{\mathcal{H}_1(u)}{\prod_{j=1}^s (u - b_j)^{\beta_j}}, \quad (58)$$

where $\mathcal{D}_2(u)$ and $\mathcal{H}_1(u)$ satisfy the Hölder condition and some additional differentiability conditions near the points a_i , b_j , and ∞ .

1°. Assume that the order η at infinity of the coefficient of the boundary value problem satisfies the condition $\eta \geq 0$. Since the first two terms of relation (58) vanish at infinity, it follows that the minimal possible order of $\mathcal{H}_1(u)$ at infinity is equal to $1 - n$. Just as in the homogeneous problem, we replace $\mathcal{D}_2(u)$ by the ratio of two functions (54) and write out the boundary condition in the following form (under the braces, the orders of the functions at infinity are indicated):

$$\underbrace{\frac{\prod_{j=1}^s (u - b_j)^{\beta_j} \mathcal{Q}_-(u) \mathcal{Y}^+(u)}{\mathcal{R}_-(u) e^{\mathcal{G}^+(u)}}}_{1-n-h} = \underbrace{\frac{\prod_{i=1}^r (u - a_i)^{\alpha_i} \mathcal{R}_+(u) \mathcal{Y}^-(u)}{\mathcal{Q}_+(u) e^{\mathcal{G}^-(u)}}}_{1-m-\nu} + \underbrace{\frac{\mathcal{H}_1(u) \mathcal{Q}_-(u)}{\mathcal{R}_-(u) e^{\mathcal{G}^+(u)}}}_{1-n-h}.$$

Assume that a polynomial $\mathcal{S}(u)$ of the degree $n + h - 1$ represents the principal part of the decomposition of the last term in a neighborhood of the point at infinity (for the case in which $n + h - 1 \geq 0$):

$$\frac{\mathcal{H}_1(u) \mathcal{Q}_-(u)}{\mathcal{R}_-(u) e^{\mathcal{G}^+(u)}} = \mathcal{S}(u) + \mathcal{W}(u), \quad \mathcal{W}(\infty) = 0.$$

On replacing the function $\mathcal{W}(u)$ by the difference of boundary values of analytic functions

$$\mathcal{W}(u) = \mathcal{W}^+(u) - \mathcal{W}^-(u), \quad (59)$$

where

$$w(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{W}(u) e^{-iux} du, \quad (60)$$

$$\mathcal{W}^+(u) = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} w(x) e^{iux} dx, \quad \mathcal{W}^-(u) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 w(x) e^{iux} dx,$$

we reduce the boundary condition to the form

$$\frac{\prod_{j=1}^s (u - b_j)^{\beta_j} \mathcal{Q}_-(u) \mathcal{Y}^+(u)}{\mathcal{R}_-(u) e^{\mathcal{G}^+(u)}} - \mathcal{S}(u) - \mathcal{W}^+(u) = \frac{\prod_{i=1}^r (u - a_i)^{\alpha_i} \mathcal{R}_+(u) \mathcal{Y}^-(u)}{\mathcal{Q}_+(u) e^{\mathcal{G}^-(u)}} - \mathcal{W}^-(u).$$

On applying the analytic continuation theorem and the generalized Liouville theorem and taking into account the fact that the only possible singular point of the function under consideration is the point at infinity, while we have the relation $-n - h \leq -m - \nu$ ($\eta \geq 0$), we obtain the expressions

$$\mathcal{Y}^+(z) = \frac{\mathcal{R}_-(z) e^{\mathcal{G}^+(z)}}{\prod_{j=1}^s (z - b_j)^{\beta_j} \mathcal{Q}_-(z)} [\mathcal{W}^+(z) + \mathcal{S}(z) + \mathcal{P}_{\nu+m-1}(z)],$$

$$\mathcal{Y}^-(z) = \frac{\mathcal{Q}_+(z) e^{\mathcal{G}^-(z)}}{\prod_{i=1}^r (z - a_i)^{\alpha_i} \mathcal{R}_+(z)} [\mathcal{W}^-(z) + \mathcal{P}_{\nu+m-1}(z)]. \quad (61)$$

The last formulas define a solution that has pole singularities at the points a_i and b_j . To obtain a bounded solution, we apply the canonical function of the nonhomogeneous problem.

By a *canonical function* $\mathcal{V}(z)$ of the nonhomogeneous Riemann problem in the exceptional case we mean a piecewise analytic function that satisfies the boundary condition (58), has the zero order on the entire finite part of the complex plane, including the points a_i and b_j , and has the least possible order at infinity.

Let $\mathcal{U}_p(z)$ be the Hermite interpolation polynomial with interpolation nodes of orders α_i and β_j at the points a_i and b_j , respectively. Such a polynomial of degree $p = m + n - 1$ exists and is determined uniquely (see Subsection 14.3-2). The functions $\mathcal{D}_1(u)$ and $\mathcal{H}_1(u)$ must be subjected to the additional condition that in neighborhoods of the points a_i and b_j these functions have derivatives of the orders α_i and β_j , respectively, and these derivatives satisfy the Hölder condition. Then the canonical function of the nonhomogeneous problem can be represented in the form

$$\begin{aligned}\mathcal{V}^+(z) &= \frac{\mathcal{R}_-(z)e^{\mathcal{G}^+(z)}}{\prod_{j=1}^s (z-b_j)^{\beta_j} \mathcal{Q}_-(z)} [\mathcal{W}^+(z) + \mathcal{S}(z) - \mathcal{U}_p(z)], \\ \mathcal{V}^-(z) &= \frac{\mathcal{Q}_+(z)e^{\mathcal{G}^-(z)}}{\prod_{i=1}^r (z-a_i)^{\alpha_i} \mathcal{R}_+(z)} [\mathcal{W}^-(z) - \mathcal{U}_p(z)].\end{aligned}\tag{62}$$

Adding $\mathcal{V}(z)$ to the above general solution of the homogeneous problem, we find the general solution of the nonhomogeneous problem under consideration:

$$\begin{aligned}\mathcal{Y}^+(z) &= \mathcal{V}^+(z) + \prod_{i=1}^r (z-a_i)^{\alpha_i} \frac{\mathcal{R}_-(z)}{\mathcal{Q}_-(z)} e^{\mathcal{G}^+(z)} \mathcal{P}_{\nu-n-1}(z), \\ \mathcal{Y}^-(z) &= \mathcal{V}^-(z) + \prod_{j=1}^s (z-b_j)^{\beta_j} \frac{\mathcal{Q}_+(z)}{\mathcal{R}_+(z)} e^{\mathcal{G}^-(z)} \mathcal{P}_{\nu-n-1}(z).\end{aligned}\tag{63}$$

For $\nu - n > 0$, the problem has $\nu - n$ linearly independent solutions. In the case $\nu - n \leq 0$ we must set $\mathcal{P}_{\nu-n-1}(z) \equiv 0$. For $\nu - n < 0$, the canonical function $\mathcal{V}(z)$ has the order $\nu - n < 0$ at infinity and hence is no longer a solution of the nonhomogeneous problem. However, on subjecting the right-hand side to $n - \nu$ conditions we can increase the order of the function $\mathcal{V}(u)$ at infinity by $n - \nu$ and thus make the canonical function $\mathcal{V}(z)$ be a solution of the nonhomogeneous problem again.

To make the above operations possible, it suffices to require that the functions $u^k \mathcal{H}_1(u)$ and $\mathcal{D}_2(u)$ have derivatives of order $\leq n - \nu$ at infinity, and these derivatives satisfy the Hölder condition.

2°. Let $\eta < 0$. The least possible order at infinity of $\mathcal{H}_1(u)$ is $h - \nu - m + 1$. In this case, the function $[\mathcal{H}_1(u)\mathcal{Q}_-(u)]/[\mathcal{R}_-(u)e^{\mathcal{G}^+(u)}]$ in the boundary condition (58) has the order $1 - m - \nu$ at infinity. After selecting the principal part of the expansion of $[\mathcal{H}_1(u)\mathcal{Q}_-(u)]/[\mathcal{R}_-(u)e^{\mathcal{G}^+(u)}]$ in a neighborhood of the point at infinity for $m + \nu - 1 > 0$, the boundary condition can be rewritten in the form

$$\frac{\prod_{j=1}^s (u-b_j)^{\beta_j} \mathcal{Q}_-(u) \mathcal{Y}^+(u)}{\mathcal{R}_-(u)e^{\mathcal{G}^+(u)}} - \mathcal{W}^+(u) = \frac{\prod_{i=1}^r (u-a_i)^{\alpha_i} \mathcal{R}_+(u) \mathcal{Y}^-(u)}{\mathcal{Q}_+(u)e^{\mathcal{G}^-(u)}} - \mathcal{W}^-(u) + \mathcal{S}(u).$$

The canonical function of the nonhomogeneous problem can be expressed via the interpolation

polynomial as follows:

$$\begin{aligned}\mathcal{V}_1^+(z) &= \frac{\mathcal{R}_-(z)e^{\mathcal{G}^+(z)}}{\prod_{j=1}^s (z - b_j)^{\beta_j} \mathcal{Q}_-(z)} [\mathcal{W}^+(z) - \mathcal{U}_p(z)], \\ \mathcal{V}_1^-(z) &= \frac{\mathcal{Q}_+(z)e^{\mathcal{G}^-(z)}}{\prod_{i=1}^r (z - a_i)^{\alpha_i} \mathcal{R}_+(z)} [\mathcal{W}^-(z) - \mathcal{S}(z) - \mathcal{U}_p(z)].\end{aligned}\tag{64}$$

The general solution of problem (58) becomes

$$\begin{aligned}\mathcal{Y}^+(z) &= \mathcal{V}_1^+(z) + \prod_{i=1}^r (z - a_i)^{\alpha_i} \frac{\mathcal{R}_-(z)}{\mathcal{Q}_-(z)} e^{\mathcal{G}^+(z)} \mathcal{P}_{h-m-1}(z), \\ \mathcal{Y}^-(z) &= \mathcal{V}_1^-(z) + \prod_{j=1}^s (z - b_j)^{\beta_j} \frac{\mathcal{Q}_+(z)}{\mathcal{R}_+(z)} e^{\mathcal{G}^-(z)} \mathcal{P}_{h-m-1}(z).\end{aligned}\tag{65}$$

For $h - m > 0$, the problem has $h - m$ linearly independent solutions. In the case $h - m \leq 0$, we must set the polynomial $\mathcal{P}_{h-m-1}(z)$ to be identically zero and, for the case in which $h - m < 0$, impose $m - h$ conditions of the same type as in the previous case on the right-hand side. Under these conditions, the nonhomogeneous problem (58) has a unique solution.

Remark. In Section 12.8 we consider equations that can be reduced to the problem by applying the convolution theorem for the Fourier transform. Equations to which the convolution theorems for other integral transforms can be applied, for instance, for the Mellin transform, can be investigated in a similar way.

⊙ References for Section 12.7: F. D. Gakhov and Yu. I. Cherskii (1978), S. G. Mikhlin and S. Prössdorf (1986), A. V. Bitsadze (1995), N. I. Muskhelishvili (1992).

12.8. Carleman Method for Equations of the Convolution Type of the First Kind

By the *Carleman method* we mean the method of reducing an integral equation to a boundary value problem of the theory of analytic functions, in particular, to the Riemann problem. For equations of convolution type, this reduction can be performed by means of the integral transforms. After solving the boundary value problem, the desired function can be obtained by applying the inverse integral transform.

12.8-1. Wiener–Hopf Equation of the First Kind.

Consider the Wiener–Hopf equation of the first kind

$$\frac{1}{\sqrt{2\pi}} \int_0^\infty K(x-t)y(t) dt = f(x), \quad 0 < x < \infty,\tag{1}$$

which is frequently encountered in applications. Let us extend its domain to the negative semiaxis by introducing one-sided functions,

$$y_+(x) = \begin{cases} y(x) & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad f_+(x) = \begin{cases} f(x) & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad y_-(x) = 0 \quad \text{for } x > 0.$$

Using these one-sided functions, we can rewrite Eq. (1) in the form

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K(x-t)y_+(t) dt = f_+(x) + y_-(x), \quad -\infty < x < \infty. \quad (2)$$

The auxiliary function $y_-(x)$ is introduced to compensate for the left-hand side of Eq. (2) for $x < 0$. Note that $y_-(x)$ is unknown in the domain $x < 0$ and is to be found in solving the problem.

Let us now apply the alternative Fourier transform to Eq. (2). Then we obtain the boundary value problem

$$\mathcal{Y}^+(u) = \frac{1}{\mathcal{K}(u)} \mathcal{Y}^-(u) + \frac{\mathcal{F}^+(u)}{\mathcal{K}(u)}. \quad (3)$$

If σ is the order of $\mathcal{K}(u)$ at infinity, then the order of the coefficient of the boundary value problem at infinity is $\eta = -\sigma < 0$. The general solution of problem (3) can be obtained on the basis of relations (65) from Subsection 12.7-7 by replacing $\mathcal{P}_{h-m-1}(z)$ with $\mathcal{P}_{\nu-n+\eta-1}(z)$ there. The solution of the original equation (1) can be obtained from the solution of problem (3) by means of the inversion formula

$$y(x) = y_+(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{Y}^+(u) e^{-iux} du, \quad x > 0. \quad (4)$$

Note that in formula (4), only the function $\mathcal{Y}^+(u)$ occurs explicitly, which is related to the function $\mathcal{Y}^-(u)$ by (3).

12.8-2. Integral Equations of the First Kind with Two Kernels.

Consider the integral equation of the first kind

$$\frac{1}{\sqrt{2\pi}} \int_0^{\infty} K_1(x-t)y(t) dt + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 K_2(x-t)y(t) dt = f(x), \quad -\infty < x < \infty. \quad (5)$$

The Fourier transform of Eq. (5) results in the following boundary value problem:

$$\mathcal{Y}^+(u) = \frac{\mathcal{K}_2(u)}{\mathcal{K}_1(u)} \mathcal{Y}^-(u) + \frac{\mathcal{F}(u)}{\mathcal{K}_1(u)}, \quad -\infty < u < \infty. \quad (6)$$

The coefficient of this problem is the ratio of functions that vanish at infinity, and hence, in contrast to the preceding case, it can have a zero or a pole of some order at infinity.

Let $\mathcal{K}_1(u) = \mathcal{T}_1(u)/u^\lambda$ and $\mathcal{K}_2(u) = \mathcal{T}_2(u)/u^\mu$, where the functions $\mathcal{T}_1(u)$ and $\mathcal{T}_2(u)$ have zero order at infinity. In the dependence of the sign of the difference $\eta = \mu - \lambda$, two cases can occur. For generality, we assume that there are exceptional points at finite distances as well. Let the functions $\mathcal{K}_1(u)$ and $\mathcal{K}_2(u)$ have the representations

$$\mathcal{K}_1(u) = \prod_{j=1}^s (u - b_j)^{\beta_j} \prod_{k=1}^p (u - c_k)^{\gamma_k} \mathcal{K}_{11}(u),$$

$$\mathcal{K}_2(u) = \prod_{i=1}^r (u - a_i)^{\alpha_i} \prod_{k=1}^p (u - c_k)^{\gamma_k} \mathcal{K}_{12}(u).$$

Along with the common zeros at points c_k of multiplicity γ_k , the functions $\mathcal{K}_1(u)$ and $\mathcal{K}_2(u)$ have a common zero of order $\min(\lambda, \mu)$ at infinity.

The coefficient of the Riemann problem can be represented in the form

$$\mathcal{D}(u) = \frac{\prod_{i=1}^r (u - a_i)^{\alpha_i} \mathcal{R}_+(u) \mathcal{R}_-(u)}{\prod_{j=1}^s (u - b_j)^{\beta_j} \mathcal{Q}_+(u) \mathcal{Q}_-(u)} \mathcal{D}_2(u).$$

It follows from (6) that this problem and the integral equation (5) are solvable if at any point c_k that is a common zero of the functions $\mathcal{K}_1(u)$ and $\mathcal{K}_2(u)$, the function $\mathcal{F}(u)$ has zero of order γ_k , i.e., $\mathcal{F}(u)$ has the form

$$\mathcal{F}(u) = \prod_{k=1}^p (u - c_k)^{\gamma_k} \mathcal{F}_1(u).$$

To this end, the following $\gamma_1 + \dots + \gamma_p = l$ conditions must hold:

$$[\mathcal{F}_u^{(j_k)}(u)]_{u=c_k} = 0, \quad j_k = 0, 1, \dots, \gamma_k - 1, \quad (7)$$

or, which is the same, the conditions

$$\int_{-\infty}^{\infty} f(x) x^{j_k} e^{ic_k x} dx = 0. \quad (8)$$

For the case under consideration in which the equation is of the first kind, we must add other d conditions, where

$$d = \min(\lambda, \mu) + 1, \quad (9)$$

that are imposed on the behavior of $\mathcal{F}(u)$ at infinity because the functions $\mathcal{K}_1(u)$ and $\mathcal{K}_2(u)$ have a common zero of order $\min(\lambda, \mu)$ at infinity. Hence, $\mathcal{F}(u)$ must satisfy the conditions (8) and have at least the order d at infinity.

If these conditions are satisfied, then the boundary value problem (6) becomes

$$\mathcal{Y}^+(u) = \frac{\prod_{i=1}^r (u - a_i)^{\alpha_i} \mathcal{R}_+(u) \mathcal{R}_-(u)}{\prod_{j=1}^s (u - b_j)^{\beta_j} \mathcal{Q}_+(u) \mathcal{Q}_-(u)} \mathcal{D}_2(u) \mathcal{Y}^-(u) + \frac{\mathcal{H}_1(u)}{\prod_{j=1}^s (u - b_j)^{\beta_j}}.$$

The solution was given above in Subsection 12.7-7. For the case in which $\eta \geq 0$ ($\mu \geq \lambda$), this solution can be rewritten in the form

$$\begin{aligned} \mathcal{Y}^+(z) &= \mathcal{V}^+(z) + \prod_{i=1}^r (z - a_i)^{\alpha_i} \frac{\mathcal{R}_-(z)}{\mathcal{Q}_-(z)} e^{\mathcal{G}^+(z)} \mathcal{P}_{\nu-n+1}(z), \\ \mathcal{Y}^-(z) &= \mathcal{V}^-(z) + \prod_{j=1}^s (z - b_j)^{\beta_j} \frac{\mathcal{Q}_+(z)}{\mathcal{R}_+(z)} e^{\mathcal{G}^-(z)} \mathcal{P}_{\nu-n+1}(z). \end{aligned} \quad (10)$$

For the case in which $\eta < 0$ ($\mu < \lambda$), this solution becomes

$$\begin{aligned} \mathcal{Y}^+(z) &= \mathcal{V}_1^+(z) + \prod_{i=1}^r (z - a_i)^{\alpha_i} \frac{\mathcal{R}_-(z)}{\mathcal{Q}_-(z)} e^{\mathcal{G}^+(z)} \mathcal{P}_{h-m-1}(z), \\ \mathcal{Y}^-(z) &= \mathcal{V}_1^-(z) + \prod_{j=1}^s (z - b_j)^{\beta_j} \frac{\mathcal{Q}_+(z)}{\mathcal{R}_+(z)} e^{\mathcal{G}^-(z)} \mathcal{P}_{h-m-1}(z). \end{aligned} \quad (11)$$

In both cases, the solution of the original integral equation can be obtained by substituting the expressions (10) and (11) into the formula

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} [\mathcal{Y}^+(u) - \mathcal{Y}^-(u)] e^{-iux} du. \tag{12}$$

Example. Consider the following equation of the first kind:

$$\frac{1}{\sqrt{2\pi}} \int_0^{\infty} K_1(x-t)y(t) dt + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 K_2(x-t)y(t) dt = f(x),$$

where

$$K_1(x) = \begin{cases} 0 & \text{for } x > 0, \\ \sqrt{2\pi} (e^{3x} - e^{2x}) & \text{for } x < 0, \end{cases} \quad K_2(x) = \begin{cases} -\sqrt{2\pi} i e^{-2x} & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad f(x) = \begin{cases} 0 & \text{for } x > 0, \\ \sqrt{2\pi} (e^{3x} - e^{2x}) & \text{for } x < 0. \end{cases} \tag{13}$$

Applying the Fourier transform to the functions in (13), we obtain

$$\mathcal{K}_1(u) = \frac{1}{(u-2i)(u-3i)}, \quad \mathcal{K}_2(u) = \frac{1}{u+2i}, \quad \mathcal{F}(u) = \frac{1}{(u-2i)(u-3i)}.$$

Here the boundary value problem (6) becomes

$$\mathcal{Y}^+(u) = \frac{(u-2i)(u-3i)}{u+2i} \mathcal{Y}^-(u) + 1.$$

The coefficient $\mathcal{D}(u)$ has a first-order pole at infinity ($\nu = -1$). In this case

$$m_+ = 2, \quad n_+ = 0, \quad \nu = m_+ - n_+ = 2, \quad \min(\lambda, \mu) = 1, \quad d = 2.$$

The function $\mathcal{F}(u)$ has second-order zero at infinity, and hence the necessary condition for the solvability is satisfied.

In the class of functions that vanish at infinity, the homogeneous problem

$$\mathcal{Y}^+(u) = \frac{(u-2i)(u-3i)}{u+2i} \mathcal{Y}^-(u)$$

has the following solution:

$$\mathcal{Y}^+(z) = \frac{C}{z+2i}, \quad \mathcal{Y}^-(z) = \frac{C}{(z-2i)(z-3i)},$$

where C is an arbitrary constant.

The number of linearly independent solutions of problem (13) is less by one than the index, because $\mathcal{D}(u)$ has a first-order pole at infinity.

The solution of the nonhomogeneous problem in the class of functions vanishing at infinity has the form

$$\mathcal{Y}^+(z) = \frac{C}{z+2i}, \quad \mathcal{Y}^-(z) = \frac{C-2i-z}{(z-2i)(z-3i)},$$

$$y(x) = \begin{cases} -\sqrt{2\pi} i C e^{-2x} & \text{for } x > 0, \\ \sqrt{2\pi} C (e^{2x} - e^{3x}) - 4i\sqrt{2\pi} e^{2x} + 5i\sqrt{2\pi} e^{3x} & \text{for } x < 0. \end{cases}$$

For the chosen right-hand side, the equation turns out to be solvable. However, if we take, for instance,

$$f(x) = \begin{cases} 0 & \text{for } x > 0, \\ \sqrt{2\pi} i (5e^{3x} - 4e^{2x}) & \text{for } x < 0, \end{cases} \tag{14}$$

then we have $\mathcal{F}(u) = (u+2i)/[(u-2i)(u-3i)]$. The corresponding Riemann boundary value problem has the form

$$\mathcal{Y}^+(u) = \frac{(u-2i)(u-3i)}{u+2i} \mathcal{Y}^-(u) + u + 2i.$$

In the class of functions bounded at infinity, its solution can be represented in the form

$$\mathcal{Y}^+(z) = \frac{C-z}{z+2i}, \quad \mathcal{Y}^-(z) = \frac{C-z-(z+2i)^2}{(z-2i)(z-3i)}. \tag{15}$$

For no choice of the constant C the solution vanishes at infinity, and hence the equation with the right-hand side defined by (14) has no solutions integrable on the real axis.

⊙ References for Section 12.8: F. D. Gakhov and Yu. I. Cherskii (1978), S. G. Mikhlín and S. Prössdorf (1986), N. I. Muskhelishvili (1992).

12.9. Dual Integral Equations of the First Kind

12.9-1. Carleman Method for Equations with Difference Kernels.

Consider the following dual integral equation of convolution type:

$$\begin{aligned} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K_1(x-t)y(t) dt &= f(x), & 0 < x < \infty, \\ \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K_2(x-t)y(t) dt &= f(x), & -\infty < x < 0, \end{aligned} \quad (1)$$

in which the function $y(x)$ is to be found.

In order to apply the Fourier transform technique (see Subsections 9.4-3, 12.7-1, and 12.7-2), we extend the domain of both conditions in Eq. (1) by formally rewriting them for all real values of x . This can be achieved by introducing new unknown functions into the right-hand sides. These functions must be chosen so that the conditions given on the semiaxis are not violated. Hence, the first condition in (1) must be complemented by a summand that vanishes on the positive semiaxis and the second by a summand that vanishes on the negative semiaxis. Thus, the dual equation can be written in the form

$$\begin{aligned} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K_1(x-t)y(t) dt &= f(x) + \xi_-(x), & -\infty < x < \infty, \\ \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K_2(x-t)y(t) dt &= f(x) + \xi_+(x), \end{aligned}$$

where the $\xi_{\pm}(x)$ are some right and left one-sided functions so far unknown.

On applying the Fourier integral transform, we have

$$\mathcal{K}_1(u)\mathcal{Y}(u) = \mathcal{F}(u) + \Xi^-(u), \quad \mathcal{K}_2(u)\mathcal{Y}(u) = \mathcal{F}(u) + \Xi^+(u). \quad (2)$$

Here the three functions $\mathcal{Y}(u)$, $\Xi^+(u)$, and $\Xi^-(u)$ are unknown.

Let us eliminate $\mathcal{Y}(u)$ from relations (2). We obtain the Riemann boundary value problem in the form

$$\Xi^+(u) = \frac{\mathcal{K}_2(u)}{\mathcal{K}_1(u)}\Xi^-(u) + \frac{\mathcal{K}_2(u) - \mathcal{K}_1(u)}{\mathcal{K}_1(u)}\mathcal{F}(u), \quad -\infty < u < \infty.$$

In the present case, the coefficient of the boundary condition is the ratio of functions that vanish at infinity, and hence this coefficient can have a zero or a pole of some order at infinity. The solution of the Riemann boundary value problem can be constructed on the basis of Subsections 12.7-6 and 12.7-7, and the solution of the integral equation (1) can be defined by the formula

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\Xi^+(u) + \mathcal{F}(u)}{\mathcal{K}_2(u)} e^{-iux} du = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\Xi^-(u) + \mathcal{F}(u)}{\mathcal{K}_1(u)} e^{-iux} du. \quad (3)$$

Example 1. Let us solve the dual equation (1), where

$$K_1(x) = \begin{cases} \sqrt{2\pi}(e^{3x} - e^{2x}) & \text{for } x < 0, \\ 0 & \text{for } x > 0, \end{cases} \quad K_2(x) = \begin{cases} 0 & \text{for } x < 0, \\ -\sqrt{2\pi}ie^{-2x} & \text{for } x > 0, \end{cases} \quad f(x) = \begin{cases} \frac{1}{4}\sqrt{2\pi}e^{2x} & \text{for } x < 0, \\ -\frac{1}{4}\sqrt{2\pi}e^{-2x} & \text{for } x > 0. \end{cases}$$

We find the Fourier integrals

$$\mathcal{K}_1(u) = \frac{1}{(u-2i)(u-3i)}, \quad \mathcal{K}_2(u) = \frac{1}{u+2i}, \quad \mathcal{F}(u) = \frac{1}{u^2+4}.$$

In this case, the boundary value problem (2) corresponding to this equation becomes

$$\Xi^+(u) = \frac{(u-2i)(u-3i)}{u+2i} \Xi^-(u) + \frac{u-3i}{(u+2i)^2} - \frac{1}{u^2+4}. \quad (4)$$

The coefficient $\mathcal{D}(u)$ has a first-order pole at infinity (with index $\nu = -1$). The functions $\mathcal{K}_1(u)$ and $\mathcal{K}_2(u)$ have a common zero of the first order at infinity. We find that

$$m_+ = 2, \quad n_+ = 0, \quad \nu = m_+ - n_+ = 2.$$

On representing the boundary condition in the form

$$(u+2i)\Xi^+(u) - \frac{u-3i}{u+2i} = (u-2i)(u-3i)\Xi^-(u) - \frac{1}{u-2i}$$

and applying the analytic continuation and the generalized Liouville theorem, we see that the general solution of problem (4) in the class of functions vanishing at infinity is given by

$$\Xi^+(z) = \frac{1}{z+2i} \left(\frac{z-3i}{z+2i} + C \right), \quad \Xi^-(z) = \frac{1}{(z-2i)(z-3i)} \left(\frac{1}{z-2i} + C \right), \quad (5)$$

where C is an arbitrary constant.

The solution of the integral equation in question is given by the expression

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\Xi^+(u) + \mathcal{F}(u)}{\mathcal{K}_2(u)} e^{-iux} du.$$

Since the function $\mathcal{K}_2(u)$ has a first-order zero at infinity, it follows that the function $\Xi^+(u) + \mathcal{F}(u)$ must have a zero at infinity whose order is at least two. This condition implies the relation $C = -1$.

For $C = -1$, formulas (5) become

$$\Xi^+(z) = \frac{-5i}{(z+2i)^2}, \quad \Xi^-(z) = \frac{1+2i-z}{(z-2i)^2(z-3i)}, \quad y(x) = \begin{cases} i\sqrt{2\pi} e^{2x} & \text{for } x < 0, \\ 5\sqrt{2\pi} e^{-2x} & \text{for } x > 0. \end{cases}$$

Thus, we have succeeded in satisfying the solvability condition, which follows from the existence of a common zero of the functions $\mathcal{K}_1(u)$ and $\mathcal{K}_2(u)$, by choosing an appropriate constant that enters the general solution, and the integral equation turns out to be unconditionally and uniquely solvable.

12.9-2. General Scheme of Finding Solutions of Dual Integral Equations.

In applications (for example, in elasticity, thermal conduction, and electrostatics), one encounters dual integral equations of the form

$$\begin{aligned} \int_0^{\infty} K(x,t)y(t) dt &= f(x) & \text{if } 0 \leq x \leq a, \\ \int_0^{\infty} M(x,t)y(t) dt &= g(x) & \text{if } a < x < \infty, \end{aligned} \quad (6)$$

where $K(x,t)$, $M(x,t)$, $f(x)$, and $g(x)$ are known functions and $y(x)$ is the function to be found.

Methods for solving various types of these equations are described, for instance, in the books mentioned in the references at the end of this section. Below we outline the general scheme of finding solutions of such equations.

A solution of equation (6) can be represented as the sum

$$y(x) = y_1(x) + y_2(x),$$

where $y_1(x)$ and $y_2(x)$ are solutions of simpler auxiliary dual equations

$$\begin{aligned} \int_0^{\infty} K(x,t)y_1(t) dt &= f(x) & \text{if } 0 \leq x \leq a, \\ \int_0^{\infty} M(x,t)y_1(t) dt &= 0 & \text{if } a < x < \infty \end{aligned} \quad (7)$$

and

$$\begin{aligned} \int_0^\infty K(x, t)y_2(t) dt &= 0 & \text{if } 0 \leq x \leq a, \\ \int_0^\infty M(x, t)y_2(t) dt &= g(x) & \text{if } a < x < \infty. \end{aligned} \quad (8)$$

For instance, consider dual equation (7) [dual equation (8) can be considered in a similar manner]. Let us supplement the second equation by the relation

$$\int_0^\infty M(x, t)y_1(t) dt = \varphi(x) \quad \text{if } 0 \leq x \leq a, \quad (9)$$

where $\varphi(x)$ is an auxiliary function to be determined. Suppose that the integral transformation

$$\int_0^\infty M(x, t)y(t) dt = z(x) \quad (0 \leq x < \infty)$$

can be inverted in the form

$$\int_0^\infty M_1(x, t)z(t) dt = y(x) \quad (0 \leq x < \infty). \quad (10)$$

Then, from (7), (9), and (10), using the relation $z(x) = \begin{cases} \varphi(x) & \text{if } 0 \leq x \leq a, \\ 0 & \text{if } a < x < \infty, \end{cases}$ we obtain

$$y_1(x) = \int_0^a M_1(x, t)\varphi(t) dt \quad (0 \leq x < \infty). \quad (11)$$

Substituting this expression into the first equation in (7) and changing the integration order, we obtain Fredholm integral equations of the first kind for the auxiliary function $\varphi(x)$:

$$\int_0^a N(x, s)\varphi(s) ds = f(x), \quad N(x, s) = \int_0^\infty K(x, t)M_1(t, s) dt. \quad (12)$$

After finding a solution of equation (12), one can use formula (11) to obtain a solution of the dual integral equation. In some cases, it is possible to find a solution of equation (12) in closed form (see Example 2, Subsection 12.9-3, and Section 3.9).

In a number of cases the kernel of integral equation (12) can be presented in the form of sum of the kernel of integral transform and some function. Then, using method described in Subsection 12.6-3, one can reduce integral equation of the first kind with constant limits of integration (12) to an integral equation of the second kind (see Subsection 12.9-4).

Example 2. Consider the dual integral equations

$$\begin{aligned} \int_0^\infty \cos(xt)y(t) dt &= f(x) & \text{if } 0 < x < 1, \\ \int_0^\infty \sin(xt)y(t) dt &= 0 & \text{if } 1 < x < \infty, \end{aligned} \quad (13)$$

which arises in crack problems in the classical theory of elasticity.

The second equation in (13) can be written as

$$\sqrt{\frac{2}{\pi}} \int_0^\infty \sin(xt)y(t) dt = \begin{cases} \varphi(x) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x < \infty, \end{cases} \quad (14)$$

where $\varphi(x)$ is an auxiliary function. The right-hand side of equation (14) is the Fourier sine transform. Applying the Fourier sine inversion formula (see Subsection 9.5-2) to (14), we get

$$y(t) = \sqrt{\frac{2}{\pi}} \int_0^1 \sin(xt)\varphi(x) dx. \quad (15)$$

Using the integral representation (see Supplement 4.6-2, integral 3)

$$\sin(xt) = t \int_0^x \frac{uJ_0(ut)}{\sqrt{x^2 - u^2}} du,$$

one can transform equation (15) (after changing the integration order) to

$$\begin{aligned} y(t) &= \sqrt{\frac{2}{\pi}} t \int_0^1 \varphi(x) \left[\int_0^x \frac{uJ_0(ut)}{\sqrt{x^2 - u^2}} du \right] dx \\ &= \sqrt{\frac{2}{\pi}} t \int_0^1 uJ_0(ut) \left[\int_u^1 \frac{\varphi(x) dx}{\sqrt{x^2 - u^2}} \right] du = t \int_0^1 J_0(ut) u\psi(u) du, \end{aligned} \quad (16)$$

where

$$\psi(u) = \sqrt{\frac{2}{\pi}} \int_u^1 \frac{\varphi(x) dx}{\sqrt{x^2 - u^2}}. \quad (17)$$

Let us rewrite the first equation in (13) as follows:

$$f(x) = \frac{d}{dx} \int_0^\infty \frac{1}{t} \sin(xt) y(t) dt \quad (0 < x < 1). \quad (18)$$

Substituting $y(t)$ from (16) into (18), we have

$$\begin{aligned} f(x) &= \frac{d}{dx} \int_0^\infty \sin(xt) \left[\int_0^1 J_0(ut) u\psi(u) du \right] dt \\ &= \frac{d}{dx} \int_0^1 u\psi(u) \left[\int_0^\infty \sin(xt) J_0(ut) dt \right] du. \end{aligned} \quad (19)$$

The last integral in square brackets can be calculated (see Supplement 4.6-1, integral 3). As a result, equation (19) becomes

$$f(x) = \frac{d}{dx} \int_0^x \frac{u\psi(u) du}{\sqrt{x^2 - u^2}} \quad (0 < x < 1). \quad (20)$$

To within obvious changes of notation, the right-hand side of equation (20) coincides with the inverse Abel-type integral equation 41 from Subsection 1.1-6. Therefore, the solution of equation (20) has the form

$$\psi(u) = \frac{2}{\pi} \int_0^u \frac{f(x) dx}{\sqrt{u^2 - x^2}}.$$

Substituting this function into (16), we find a solution of the original dual integral equation (13):

$$y(t) = \frac{2}{\pi} t \int_0^1 uJ_0(ut) \left[\int_0^u \frac{f(x) dx}{\sqrt{u^2 - x^2}} \right] du. \quad (21)$$

Below we give solutions for some classes of dual integral equations that occur most frequently in applications.

12.9-3. Exact Solutions of Some Dual Equations of the First Kind.

Below we present solutions of some classes of dual integral equations that occur most frequently in applications.

1°. Consider the following dual integral equation:

$$\begin{aligned} \int_0^\infty J_0(xt) y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty t J_0(xt) y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (22)$$

where $J_0(x)$ is the Bessel function of zero order. We can obtain the solution of Eqs. (22) by applying the Hankel transform. This solution is given by

$$y(x) = \frac{2}{\pi} \int_0^a \cos(xt) \left[\frac{d}{dt} \int_0^t \frac{sf(s) ds}{\sqrt{t^2 - s^2}} \right] dt. \quad (23)$$

2°. The exact solution of the dual integral equation

$$\begin{aligned} \int_0^\infty t J_0(xt)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty J_0(xt)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (24)$$

where $J_0(x)$ is the Bessel function of zero order, can be constructed by means of the Hankel transform,

$$y(x) = \frac{2}{\pi} \int_0^a \sin(xt) \left[\frac{d}{dt} \int_0^t \frac{sf(s) ds}{\sqrt{t^2 - s^2}} \right] dt. \quad (25)$$

3°. The exact solution of the dual integral equation

$$\begin{aligned} \int_0^\infty t J_\mu(xt)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty J_\mu(xt)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (26)$$

where $J_\mu(x)$ is the Bessel function of order μ , can be defined by the following expression (here the calculation also involves the Hankel transform):

$$y(x) = \sqrt{\frac{2x}{\pi}} \int_0^a t^{3/2} J_{\mu+\frac{1}{2}}(xt) \left[\int_0^{\pi/2} \sin^{\mu+1} \theta f(t \sin \theta) d\theta \right] dt. \quad (27)$$

4°. Consider the dual integral equation

$$\begin{aligned} \int_0^\infty t^{2\beta} J_\mu(xt)y(t) dt &= f(x) & \text{for } 0 < x < 1, \\ \int_0^\infty J_\mu(xt)y(t) dt &= 0 & \text{for } 1 < x < \infty, \end{aligned} \quad (28)$$

where $J_\mu(x)$ is the Bessel function of order μ .

The solution of Eq. (28) can be obtained by applying the Mellin transform. For $\beta > 0$, this solution is defined by the formulas

$$y(x) = \frac{(2x)^{1-\beta}}{\Gamma(\beta)} \int_0^1 t^{1+\beta} J_{\mu+\beta}(xt) F(t) dt, \quad F(t) = \int_0^1 f(t\zeta) \zeta^{\mu+1} (1-\zeta^2)^{\beta-1} d\zeta. \quad (29)$$

For $\beta > -1$, the solution of the dual equation (28) has the form

$$y(x) = \frac{(2x)^{-\beta}}{\Gamma(1+\beta)} \left[x^{1+\beta} J_{\mu+\beta}(x) \int_0^1 t^{\mu+1} (1-t^2)^\beta f(t) dt + \int_0^1 t^{\mu+1} (1-t^2)^\beta \Phi(x, t) dt \right], \quad (30)$$

$$\Phi(x, t) = \int_0^1 (x\xi)^{2+\beta} J_{\mu+\beta+1}(x\xi) f(\xi) d\xi.$$

Formula (30) holds for $\beta > -1$ and for $-\mu - \frac{1}{2} < 2\beta < \mu + \frac{3}{2}$. It can be shown that for $\beta > 0$ the solution of Eq. (30) can be reduced to the form (29).

5°. The exact solution of the dual integral equation

$$\begin{aligned} \int_0^\infty t P_{-\frac{1}{2}+it}(\cosh x) y(t) dt &= f(x) && \text{for } 0 < x < a, \\ \int_0^\infty \tanh(\pi t) P_{-\frac{1}{2}+it}(\cosh x) y(t) dt &= 0 && \text{for } a < x < \infty, \end{aligned} \quad (31)$$

where $P_\mu(x)$ is the Legendre spherical function of the first kind (see Supplement 11.11) and $i^2 = -1$, can be constructed by means of the Meler–Fock integral transform (see Section 9.6) and is given by the formula

$$y(x) = \frac{\sqrt{2}}{\pi} \int_0^a \sin(xt) \left[\int_0^t \frac{f(s) \sinh s}{\sqrt{\cosh t - \cosh s}} ds \right] dt. \quad (32)$$

Note that

$$P_{-\frac{1}{2}+it}(\cosh x) = \frac{\sqrt{2}}{\pi} \int_0^x \frac{\cos(ts)}{\sqrt{\cosh x - \cosh s}} ds, \quad x > 0,$$

where the integral on the right-hand side is called the *Meler integral*.

12.9-4. Reduction of Dual Equations to a Fredholm Equation.

One of the most effective methods for the approximate solution of dual integral equations of the first kind is the method of reducing these equations to Fredholm integral equations of the second kind (see Chapter 13). In what follows, we present some dual equations encountered in problems of mechanics and physics and related Fredholm equations of the second kind.

1°. The solution of the dual integral equation of the first kind

$$\begin{aligned} \int_0^\infty g(t) J_0(xt) y(t) dt &= f(x) && \text{for } 0 < x < a, \\ \int_0^\infty t J_0(xt) y(t) dt &= 0 && \text{for } a < x < \infty, \end{aligned} \quad (33)$$

where $g(x)$ is a given function and $J_0(x)$ is the Bessel function of zero order, has the form

$$y(x) = \int_0^a \varphi(t) \cos(xt) dt, \quad (34)$$

where the function $\varphi(x)$ to be found from the following Fredholm equation of the second kind:

$$\varphi(x) - \frac{1}{\pi} \int_0^a K(x, t) \varphi(t) dt = \psi(x), \quad 0 < x < a, \quad (35)$$

where the symmetric kernel $K(x, t)$ and the right-hand side $\psi(x)$ are given by

$$K(x, t) = 2 \int_0^\infty [1 - g(s)] \cos(xs) \cos(ts) ds, \quad \psi(x) = \frac{2}{\pi} \frac{d}{dx} \int_0^x \frac{tf(t)}{\sqrt{x^2 - t^2}} dt. \quad (36)$$

Methods for the investigation of these equations are presented in Chapter 13.

2°. The solution of the dual integral equation of the first kind

$$\begin{aligned} \int_0^\infty tg(t)J_0(xt)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty J_0(xt)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (37)$$

where $g(x)$ is a given function and $J_0(x)$ is the Bessel function of zero order, has the form

$$y(x) = \int_0^a \varphi(t) \sin(xt) dt, \quad (38)$$

where the function $\varphi(x)$ is to be found from the Fredholm equation (35) of the second kind with

$$K(x, t) = 2 \int_0^\infty [1 - g(s)] \sin(xs) \sin(ts) ds, \quad \psi(x) = \frac{2}{\pi} \int_0^x \frac{tf(t)}{\sqrt{x^2 - t^2}} dt.$$

Note that the kernel $K(x, t)$ is symmetric.

3°. The solution of the dual integral equation of the first kind

$$\begin{aligned} \int_0^\infty g(t)J_\mu(xt)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty tJ_\mu(xt)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (39)$$

where $g(x)$ is a given function and $J_\mu(x)$ is the Bessel function of order μ , has the form

$$y(x) = \sqrt{\frac{\pi x}{2}} \int_0^a \sqrt{t} J_{\mu-\frac{1}{2}}(xt) \varphi(t) dt, \quad (40)$$

where the function $\varphi(x)$ is to be found from the Fredholm equation (35) of the second kind with

$$\begin{aligned} K(x, t) &= \pi \sqrt{xt} \int_0^\infty [1 - g(s)] s J_{\mu-\frac{1}{2}}(xs) J_{\mu-\frac{1}{2}}(ts) ds, \\ \psi(x) &= \frac{2}{\pi} \left\{ f(0) + \int_0^{\pi/2} [\mu(\sin \theta)^{\mu-1} f(x \sin \theta) + x(\sin \theta)^\mu f'(x \sin \theta)] d\theta \right\}. \end{aligned}$$

Note that $f'(x \sin \theta) = f'_\xi(\xi)|_{\xi=x \sin \theta}$, and the kernel $K(x, t)$ is symmetric.

4°. The solution of the integral equation of the first kind

$$\begin{aligned} \int_0^\infty tg(t)J_\mu(xt)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty J_\mu(xt)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (41)$$

where $g(x)$ is a given function and $J_\mu(x)$ is the Bessel function of order μ , has the form

$$y(x) = \sqrt{\frac{\pi x}{2}} \int_0^a \sqrt{t} J_{\mu+\frac{1}{2}}(xt) \varphi(t) dt, \quad (42)$$

where the function $\varphi(x)$ is to be found by solving the Fredholm equation (35) of the second kind with

$$K(x, t) = \pi \sqrt{xt} \int_0^\infty [1 - g(s)] s J_{\mu+\frac{1}{2}}(xs) J_{\mu+\frac{1}{2}}(ts) ds, \quad \psi(x) = \frac{2x}{\pi} \int_0^{\pi/2} f(x \sin \theta) (\sin \theta)^{\mu+1} d\theta,$$

and the kernel $K(x, t)$ is symmetric.

5°. The solution of the dual integral equation of the first kind

$$\begin{aligned} \int_0^\infty g(t)J_\mu(xt)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty J_\mu(xt)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (43)$$

where $g(x)$ is a given function and $J_\mu(x)$ is the Bessel function of order μ , has the form

$$y(x) = x\sqrt{\frac{\pi x}{2}} \int_0^a \sqrt{t} J_{\mu-\frac{1}{2}}(xt)\varphi(t) dt, \quad (44)$$

and the function $\varphi(x)$ is to be found from the Fredholm equation (35) of the second kind with

$$\begin{aligned} K(x, t) &= x^\mu \sqrt{2\pi t} \int_x^a \frac{\rho^{1-\mu}}{\sqrt{\rho^2 - x^2}} \int_0^\infty [1 - g(s)]s^{3/2} J_\mu(\rho s) J_{\mu-\frac{1}{2}}(ts) ds d\rho, \\ \psi(x) &= \frac{2}{\pi} x^\mu \int_x^a \frac{\rho^{1-\mu}}{\sqrt{\rho^2 - x^2}} d\rho. \end{aligned}$$

6°. The solution of the dual integral equation of the first kind

$$\begin{aligned} \int_0^\infty t^{2\beta} g(t)J_\mu(xt)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty J_\mu(xt)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (45)$$

where $0 < \beta < 1$, $g(x)$ is a given function, and $J_\mu(x)$ is the Bessel function of order μ , has the form

$$y(x) = \sqrt{\frac{\pi}{2}} x^{1-\beta} \int_0^a \sqrt{t} J_{\mu+\beta}(xt)\varphi(t) dt, \quad (46)$$

and the function $\varphi(x)$ is to be found from the Fredholm equation (35) of the second kind with

$$\begin{aligned} K(x, t) &= \pi\sqrt{xt} \int_0^\infty [1 - g(s)]s J_{\mu+\beta}(xs) J_{\mu+\beta}(ts) ds, \\ \psi(x) &= \frac{2^{1-\beta}}{\Gamma(\beta)} \sqrt{\frac{2x}{\pi}} x^\beta \int_0^{\pi/2} f(x \sin \theta) (\sin \theta)^{\mu+1} (\cos \theta)^{2\beta-1} d\theta, \end{aligned}$$

and the kernel $K(x, t)$ is symmetric.

7°. The solution of the dual integral equation of the first kind

$$\begin{aligned} \int_0^\infty g(t)P_{-\frac{1}{2}+it}(\cosh x)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty t \tanh(\pi t)P_{-\frac{1}{2}+it}(\cosh x)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (47)$$

where $P_\mu(x)$ is the Legendre spherical function of the first kind (see Supplement 11.11), $i^2 = -1$, and $g(x)$ is a given function, is determined by the formula

$$y(x) = \int_0^a \cos(xt)\varphi(t) dt, \quad (48)$$

and the function $\varphi(x)$ is to be found from the Fredholm equation (35) of the second kind in which

$$\begin{aligned} K(x, t) &= \int_0^\infty [1 - g(s)]\{\cos[(x+t)s] + \cos[(x-t)s]\} ds, \\ \psi(x) &= \frac{\sqrt{2}}{\pi} \frac{d}{dx} \int_0^x \frac{f(s) \sinh s}{\sqrt{\cosh x - \cosh s}} ds. \end{aligned} \quad (49)$$

On the basis of relations (49), we can readily see that the kernel $K(x, t)$ is symmetric.

8°. The solution of the dual integral equation of the first kind

$$\begin{aligned} \int_0^\infty tg(t)P_{-\frac{1}{2}+it}(\cosh x)y(t) dt &= f(x) & \text{for } 0 < x < a, \\ \int_0^\infty \tanh(\pi t)P_{-\frac{1}{2}+it}(\cosh x)y(t) dt &= 0 & \text{for } a < x < \infty, \end{aligned} \quad (50)$$

where $P_\mu(x)$ is the spherical Legendre function of the first kind (see Supplement 11.11), $i^2 = -1$, and $g(x)$ is a given function, is determined by the formula

$$y(x) = \int_0^a \sin(xt)\varphi(t) dt, \quad (51)$$

and the function $\varphi(x)$ is to be found from the Fredholm equation (35) of the second kind in which

$$\begin{aligned} K(x, t) &= \int_0^\infty [1 - g(s)]\{\cos[(x-t)s] - \cos[(x+t)s]\} ds, \\ \psi(x) &= \frac{\sqrt{2}}{\pi} \int_0^x \frac{f(s) \sinh s}{\sqrt{\cosh x - \cosh s}} ds. \end{aligned} \quad (52)$$

On the basis of relations (52), we can readily see that the kernel $K(x, t)$ is symmetric.

⊙ References for Section 12.9: Ya. S. Uflyand (1977), F. D. Gakhov and Yu. I. Cherskii (1978), C. Nasim and B. D. Aggarwala (1984), E. C. Titchmarsh (1986), I. Sneddon (1995), B. N. Mandal and N. Mandal (1999, pp. 134–136).

12.10. Asymptotic Methods for Solving Equations with Logarithmic Singularity

12.10-1. Preliminary Remarks.

Consider the Fredholm integral equation of the first kind of the form

$$\int_{-1}^1 K\left(\frac{x-t}{\lambda}\right)y(t) dt = f(x), \quad -1 \leq x \leq 1, \quad (1)$$

with parameter λ ($0 < \lambda < \infty$).

We assume that the kernel $K = K(x)$ is an even function continuous for $x \neq 0$ which has a logarithmic singularity as $x \rightarrow 0$ and exponentially decays as $x \rightarrow \infty$. Equations with such a kernel arise in solving various problems of continuum mechanics with mixed boundary conditions.

Let $f(x)$ belong to the space of functions whose first derivatives satisfy the Hölder condition with exponent $\alpha > \frac{1}{2}$ on $[-1, 1]$. In this case, the solution of the integral equation (1) in the class of functions satisfying the Hölder condition exists and is unique for any $\lambda \in (0, \infty)$ and has the structure

$$y(x) = \frac{\omega(x)}{\sqrt{1-x^2}}, \quad (2)$$

where $\omega(x)$ is a continuous function that does not vanish at $x = \pm 1$.*

It follows from formula (2) that the solution of Eq. (1) is unbounded as $x \rightarrow \pm 1$. This important circumstance will be taken into account in Subsection 12.10-3 in constructing the asymptotic solution in the case $\lambda \rightarrow 0$.

Note that more general equations with difference kernel and arbitrary finite limits of integration can always be reduced to Eq. (1) by a change of variables. The form (1) is taken here for further convenience.

* The situation $\omega(\pm 1) = 0$ is only possible in exceptional cases for special values of λ .

12.10-2. Solution for Large λ .

Let the representation

$$K(x) = \ln|x| \sum_{n=0}^{\infty} a_n |x|^n + \sum_{n=0}^{\infty} b_n |x|^n, \quad (3)$$

where $a_0 \neq 0$, be valid for the kernel of the integral equation (1) as $x \rightarrow 0$.

It is obvious from (3) that two different-scale large parameters λ and $\ln \lambda$ occur in Eq. (1) as $\lambda \rightarrow \infty$. The latter, “quasiconstant” parameter grows much slower than the former (for instance, for $\lambda = 100$ and $\lambda = 1000$ we have $\ln \lambda \approx 4.6$ and $\ln \lambda \approx 6.9$, respectively).

Let us drop out all terms decaying as $\lambda \rightarrow \infty$ in Eq. (1). In view of (3), for the main (zeroth) approximation we have

$$\int_{-1}^1 (a_0 \ln|x-t| - a_0 \ln \lambda + b_0) y_0(t) dt = f(x), \quad -1 \leq x \leq 1. \quad (4)$$

It should be noted that one cannot retain in the integrand only one term proportional to $\ln \lambda$ (since the corresponding “truncated” equation is unsolvable). The constant b_0 must also be included in (4) for the main-approximation equation to be invariant with respect to the scaling parameter λ in Eq. (1).

The exact closed-form solution of Eq. (4) is given in Section 3.4 (see equations 3 and 4).

To construct an asymptotic solution of Eq. (1) as $\lambda \rightarrow \infty$, it is convenient to do the following. First, we consider the auxiliary integral equation

$$\int_{-1}^1 \mathcal{K}(x-t, \beta, \lambda) y(t) dt = f(x), \quad -1 \leq x \leq 1, \quad (5)$$

$$\mathcal{K}(x, \beta, \lambda) = (\ln|x| - \beta) \sum_{n=0}^{\infty} \frac{a_n}{\lambda^n} |x|^n + \sum_{n=0}^{\infty} \frac{b_n}{\lambda^n} |x|^n,$$

with two parameters λ and β . We seek its solution in the form of a regular asymptotic expansion in negative powers of λ (for fixed β). That is, we have

$$y(x, \beta, \lambda) = \sum_{n=0}^N \lambda^{-n} y_n(x, \beta) + o(\lambda^{-N}). \quad (6)$$

Substituting (6) into (5) yields a recurrent chain of integral equations of the form (4):

$$\int_{-1}^1 (a_0 \ln|x-t| - a_0 \beta + b_0) y_n(t, \beta) dt = g_n(x, \beta), \quad -1 \leq x \leq 1, \quad (7)$$

from which the functions $y_n(x, \beta)$ can be successively calculated. The right-hand sides $g_n(x, \beta)$ depend only on the previously determined functions y_0, y_1, \dots, y_{n-1} .

Note that for $\beta = \ln \lambda$ the auxiliary equation (5) coincides with the original equation (1) into which the expansion (3) is substituted. Therefore, the asymptotic solution of Eq. (1) can be obtained with the aid of (6) and (7) with $\beta = \ln \lambda$.

Some contact problems of elasticity can be reduced to Eq. (1), in which the kernel can be represented in the form (3) with $a_n = 0$ for all $n > 0$ and $b_{2m+1} = 0$ for $m = 0, 1, 2, \dots$. In this case, one must set $y_n(x, \beta) \equiv 0$ ($n = 1, 3, 5, \dots$) in the solution (6). In practice, it usually suffices to retain the terms up to λ^{-4} .

12.10-3. Solution for Small λ .

In analyzing the limit case $\lambda \rightarrow 0$, we take into account the singularities of the solution at the endpoints of the interval $-1 \leq x \leq 1$ (see formula (2)). Consider the following auxiliary system of two integral equations:

$$\begin{aligned} \int_{-1}^{\infty} K\left(\frac{x-t}{\lambda}\right)y_1(t) dt &= f_1(x) + \int_{-\infty}^{-1} K\left(\frac{x-t}{\lambda}\right)y_2(t) dt, & -1 \leq x < \infty, \\ \int_{-\infty}^1 K\left(\frac{x-t}{\lambda}\right)y_2(t) dt &= f_2(x) + \int_1^{\infty} K\left(\frac{x-t}{\lambda}\right)y_1(t) dt, & -\infty < x \leq 1. \end{aligned} \quad (8)$$

The former equation provides for selecting the singularity at $x = -1$ and the latter for selecting the singularity at $x = +1$.

The functions $f_1(x)$ and $f_2(x)$ are such that

$$\begin{aligned} f_1(x) + f_2(x) &= f(x), & -1 \leq x \leq 1, \\ f_1(x) &= O(e^{-\alpha_1 x}) & \text{as } x \rightarrow \infty, \\ f_2(x) &= O(e^{\alpha_2 x}) & \text{as } x \rightarrow -\infty, \end{aligned} \quad (9)$$

where $\alpha_1 > 0$ and $\alpha_2 > 0$.

The first condition in (9) makes it possible to seek the solution of the integral equation (1) as the sum of the solutions of the integral equations (8), that is,

$$y(x) = y_1(x) + y_2(x), \quad -1 \leq x \leq 1. \quad (10)$$

Note that by virtue of the last two conditions in (9), the relations

$$\begin{aligned} y_1(x) &= O(e^{-\beta_1 x}) & \text{as } x \rightarrow \infty, \\ y_2(x) &= O(e^{\beta_2 x}) & \text{as } x \rightarrow -\infty, \end{aligned} \quad (11)$$

where $\beta_1 > 0$ and $\beta_2 > 0$, are valid.

Recall that the kernel $K(x)$ is an even function. Therefore, if $f(x)$ in Eq. (1) is an even or odd function, then one must set

$$f_1(x) = \pm f_2(-x), \quad y_1(x) = \pm y_2(-x) \quad (12)$$

in system (8).*

In both cases, system (8) can be reduced by changes of variables to the same integral equation

$$\int_0^{\infty} K(z-\tau)w(\tau) d\tau = F(z) \pm \int_{2/\lambda}^{\infty} K(2/\lambda - z - \tau)w(\tau) d\tau, \quad 0 \leq z < \infty, \quad (13)$$

in which the following notation is used:

$$z = \frac{x+1}{\lambda}, \quad \tau = \frac{t+1}{\lambda}, \quad w(\tau) = y(t), \quad F(z) = \frac{1}{\lambda} f_1(x). \quad (14)$$

In view of the properties of the kernel $K(x)$ (see Subsection 12.10-1) and the first relation in (11), the asymptotic estimate

$$I(w) \equiv \int_{2/\lambda}^{\infty} K(2/\lambda - z - \tau)w(\tau) d\tau = O(e^{-2\beta_1/\lambda}) \quad (15)$$

can be obtained, which is uniform with respect to τ .

* In formulas (12), (13), (16), and (17), the plus sign corresponds to even $f(x)$ and the minus sign to odd $f(x)$.

According to (15), for small λ the iterative scheme

$$\int_0^\infty K(z - \tau)w_n(\tau) d\tau = F(z) \pm I(w_{n-1}), \quad n = 1, 2, \dots, \quad (16)$$

can be used to solve the integral equation (13) by the method of successive approximations. In the main approximation, the integral $I(w_0)$ can be omitted on the right-hand side. Equations (16) are Wiener–Hopf integral equations of the first kind, which can be solved in a closed form (see Subsection 12.8-1).

It follows from formulas (10), (12), and (14) that, as $\lambda \rightarrow 0$, the leading term of the asymptotic expansion of the solution of the integral equation (1) has the form

$$y(x) = w_1 \left(\frac{1+x}{\lambda} \right) \pm w_1 \left(\frac{1-x}{\lambda} \right), \quad (17)$$

where $w_1 = w_1(\tau)$ is the solution of Eq. (16) with $n = 1$ and $w_0 \equiv 0$.

For practical purposes, formula (17) is usually sufficient.

12.10-4. Integral Equation of Elasticity.

The integral equation (1) whose kernel is given via the Fourier cosine transform,

$$K(x) = \int_0^\infty \frac{L(u)}{u} \cos(ux) du, \quad (18)$$

frequently occurs in contact problems of elasticity. The function $L(u)$ in (18) is continuous and positive for $0 < u < \infty$ and satisfies the asymptotic relations

$$\begin{aligned} L(u) &= Au + O(u^3) && \text{as } u \rightarrow 0, \\ L(u) &= \sum_{n=0}^{N-1} B_n u^{-n} + O(u^{-N}) && \text{as } u \rightarrow \infty, \end{aligned} \quad (19)$$

where $A > 0$ and $B_0 > 0$.

Formula (18) implies that the kernel is an even function: $K(x) = K(-x)$.

It is usually assumed that $L(u)u^{-1}$ and $u[L(u)]^{-1}$, treated as functions of the complex variable $w = u + iv$, are regular at the pole $|v| \leq \gamma_1$ and the pole $|v| \leq \gamma_2$, respectively. It follows in particular that the kernel $K(x)$ decays at least as $\exp(-\gamma_1|t|)$ at infinity.

Formulas (18) and (19) imply that $K(x)$ has a logarithmic singularity at $x = 0$. Moreover, the representation (3) is valid with $a_n = 0$ for $n = 1, 3, 5, \dots$

Thus, the kernel given by (18) has the same characteristic features as those inherent by assumption in the kernel of the integral equation (1). Therefore, the results of Subsections 12.10-2 and 12.10-3 can be used for the asymptotic analysis of Eq. (1) with kernel (18) as $\lambda \rightarrow \infty$ and $\lambda \rightarrow 0$.

⊙ References for Section 12.10: I. I. Vorovich, V. M. Aleksandrov, and V. A. Babeshko (1974), V. M. Aleksandrov and E. V. Kovalenko (1986), V. M. Aleksandrov (1993).

12.11. Regularization Methods

12.11-1. Lavrentiev Regularization Method.

Consider the Fredholm equation of the first kind

$$\int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b, \quad (1)$$

where $f(x) \in L_2(a, b)$ and $y(x) \in L_2(a, b)$. The kernel $K(x, t)$ is square integrable, symmetric, and positive definite (see Subsection 13.6-2), that is, for all $\varphi(x) \in L_2(a, b)$, we have

$$\int_a^b \int_a^b K(x, t)\varphi(x)\varphi(t) dx dt \geq 0,$$

where the equality is attained only for $\varphi(x) \equiv 0$.

In the above classes of functions and kernels, the problem of finding a solution of Eq. (1) is ill-posed, i.e., unstable with respect to small variations in the right-hand side of the integral equation.

Following the Lavrentiev regularization method, along with Eq. (1) we consider the regularized equation

$$\varepsilon y_\varepsilon(x) + \int_a^b K(x, t)y_\varepsilon(t) dt = f(x), \quad a \leq x \leq b, \quad (2)$$

where $\varepsilon > 0$ is the regularization parameter. This equation is a Fredholm equation of the second kind, so it can be solved by the methods presented in Chapter 13, whence the solution exists and is unique.

On taking a sufficiently small ε in Eq. (2), we find a solution $y_\varepsilon(x)$ of the equation and substitute this solution into Eq. (1), thus obtaining

$$\int_a^b K(x, t)y_\varepsilon(t) dt = f_\varepsilon(x), \quad a \leq x \leq b. \quad (3)$$

If the function $f_\varepsilon(x)$ thus obtained differs only slightly from $f(x)$, that is,

$$\|f(x) - f_\varepsilon(x)\| \leq \delta, \quad (4)$$

where δ is a prescribed small positive number, then the solution $y_\varepsilon(x)$ is regarded as a sufficiently good approximate solution of Eq. (1).

The parameter δ usually defines the error of the initial data provided that the right-hand side of Eq. (1) is defined or determined by an experiment with some accuracy.

For the case in which, for a given ε , condition (4) fails, we must choose another value of the regularization parameter and repeat the above procedure.

The next subsection describes the regularization method suitable for equations of the first kind with arbitrary square-integrable kernels.

12.11-2. Tikhonov Regularization Method.

Consider the Fredholm integral equation of the first kind

$$\int_a^b K(x, t)y(t) dt = f(x), \quad c \leq x \leq d. \quad (5)$$

Assume that $K(x, t)$ is any function square-integrable in the domain $\{a \leq t \leq b, c \leq x \leq d\}$, $f(x) \in L_2(c, d)$, and $y(x) \in L_2(a, b)$. The problem of finding the solution of Eq. (5) is also ill-posed in the above sense.

Following the Tikhonov (zero-order) regularization method, along with (5) we consider the following Fredholm integral equation of the second kind (see Chapter 13):

$$\varepsilon y_\varepsilon(x) + \int_a^b K^*(x, t)y_\varepsilon(t) dt = f^*(x), \quad a \leq x \leq b, \quad (6)$$

where

$$K^*(x, t) = K^*(t, x) = \int_c^d K(s, x)K(s, t) ds, \quad f^*(x) = \int_c^d K(s, x)f(s) ds, \quad (7)$$

and the positive number ε is the regularization parameter. Equation (6) is said to be a *regularized integral equation*, and its solution exists and is unique.

Taking a sufficiently small ε in Eq. (6), we find a solution $y_\varepsilon(x)$ of the equation and substitute this solution into Eq. (5), thus obtaining

$$\int_a^b K(x, t)y_\varepsilon(t) dt = f_\varepsilon(x), \quad c \leq x \leq d. \quad (8)$$

By comparing the right-hand side with the given $f(x)$ using formula (4), we either regard $f_\varepsilon(x)$ as a satisfactory approximate solution obtained in accordance with the above simple algorithm, or continue the procedure for a new value of the regularization parameter.

Presented above are the simplest principles of finding an approximate solution of the Fredholm equation of the first kind. More perfect and complex algorithms can be found in the references cited below.

© References for Section 12.11: M. M. Lavrentiev (1967), A. N. Tikhonov and V. Ya. Arsenin (1979), M. M. Lavrentiev, V. G. Romanov, and S. P. Shishatskii (1980), A. F. Verlan' and V. S. Sizikov (1986), R. Kress (1999).

12.12. Fredholm Integral Equation of the First Kind as an Ill-Posed Problem

12.12-1. General Notions of Well-Posed and Ill-Posed Problems.

To solve a quantitative mathematical problem usually means to find an element y , called a “solution of the problem”, from a given element f , called “data of the problem”. Assume that y and f are elements of some metric spaces Y (space of solutions) and F (space of data) with the respective distances between their elements $\rho_Y(y_1, y_2)$ and $\rho_F(f_1, f_2)$.

A solution y corresponding to f is called *stable*, if for any $\varepsilon > 0$ there is $\delta(\varepsilon) > 0$ such that for any $f_1 \in F$ such that $\rho_F(f, f_1) \leq \delta(\varepsilon)$ we have $\rho_Y(y, y_1) \leq \varepsilon$, where y_1 is a solution corresponding to f_1 . In other words, if small variations of data cause a small variation of solutions.

Such a problem is called *well-posed* on a pair of metrics spaces (Y, F) , if the following conditions hold:

- 1) for each $f \in F$, there is a solution $y \in Y$;
- 2) the solution is unique;
- 3) the solution is stable.

Problems that do not satisfy one of these requirements are called *ill-posed*.

Remark. The metrics in the spaces Y and F determine in what sense small variations of y and f are understood. The choice of these metrics determines whether a solution y is stable or not under the variation of f , and therefore, a particular problem may be well-posed or ill-posed, depending on the metrics.

Now, for definiteness, assume that y and f are elements of the space of continuous functions on an interval $[a, b]$ with the metrics

$$\rho_Y(y_1, y_2) = \sup_{a \leq x \leq b} |y_1(x) - y_2(x)|, \quad \rho_F(f_1, f_2) = \sup_{a \leq x \leq b} |f_1(x) - f_2(x)|. \quad (1)$$

12.12-2. Integral Equation of the First Kind is an Ill-Posed Problem.

Consider the Fredholm equation of the first kind

$$\int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b, \quad (2)$$

with continuous kernel $K(x, t)$, and assume that y and f are elements of the space of continuous functions with the metrics (1).

It is easy to see that the continuity of $f(x)$, in general, does not guarantee the existence of a continuous solution. Indeed, suppose that the function $f(x)$ is continuous, but its derivative is discontinuous at some points $x \in (a, b)$, and the kernel is continuously differentiable in x . Then for any continuous $y(x)$ the left-hand side of (2) has a continuous derivative at all points of (a, b) , while the derivative of the right-hand side has discontinuities on (a, b) . Therefore, relation (2) can hold for no continuous function $y(x)$, which means that equation (2) has no solutions.

Consider the problem of stability of a solution. Assume that the kernel $K(x, t)$ is continuous, together with its derivative in t . Let $y(x)$ be a solution of equation (2). Take $z(x) = y(x) + \cos(\omega x)$, where ω is a parameter. We have

$$\int_a^b K(x, t)[z(t) - \cos(\omega t)] dt = f(x).$$

After elementary transformations, we get

$$\begin{aligned} \int_a^b K(x, t)z(t) dt &= g(x) \equiv f(x) + \int_a^b K(x, t) \cos(\omega t) dt \\ &= f(x) + K(x, t) \frac{\sin(\omega t)}{\omega} \Big|_a^b - \int_a^b K'_t(x, t) \cos(\omega t) \frac{\sin(\omega t)}{\omega} dt. \end{aligned}$$

This relation, for a finite interval $[a, b]$, implies the estimate

$$\sup_{a \leq x \leq b} |f(x) - g(x)| \leq \frac{C}{\omega},$$

where C is a constant that does not depend on ω . Therefore, for sufficiently large ω , the value

$$\rho_F(f, g) = \sup_{a \leq x \leq b} |f(x) - g(x)|$$

becomes arbitrarily small, $\rho_F(f, g) \rightarrow 0$. On the other hand, we have

$$\rho_Y(y, z) = \sup_{a \leq x \leq b} |y(x) - z(x)| = \sup_{a \leq x \leq b} |\cos(\omega x)| = 1,$$

and this quantity is not small. Hence, an important conclusion can be made: the Fredholm integral equation of the first kind (2) admits a solution which is unstable with respect to perturbations of the right-hand side $f(x)$. Thus, equation (2) belongs to the class of ill-posed problems.

This instability of solutions of integral equations of the first kind causes great difficulties when using such equations for practical purposes, since small errors in input data may cause large variations of a solution. For this reason, there existed a widespread opinion that Fredholm equations of the first kind (as well as other ill-posed problems) are unsuitable for the description of physical processes. At present this view has changed drastically due to the development of the general theory of ill-posed problems and the corresponding regularization methods (see Section 12.11 and the references below).

© References for Section 12.12: M. M. Lavrentiev (1967), A. N. Tikhonov and V. Ya. Arsenin (1979), M. M. Lavrentiev, V. G. Romanov, and S. P. Shishatskii (1980), A. B. Vasilieva and A. N. Tikhonov (1989), R. Kress (1999).

Chapter 13

Methods for Solving Linear Equations of the Form $y(x) - \int_a^b K(x, t)y(t) dt = f(x)$

13.1. Some Definition and Remarks

13.1-1. Fredholm Equations and Equations with Weak Singularity of the Second Kind.
--

Linear integral equations of the second kind with constant limits of integration have the form

$$y(x) - \lambda \int_a^b K(x, t)y(t) dt = f(x), \quad (1)$$

where $y(x)$ is the unknown function ($a \leq x \leq b$), $K(x, t)$ is the *kernel* of the integral equation, and $f(x)$ is a given function, which is called the *right-hand side* of Eq. (1). For convenience of analysis, a number λ is traditionally singled out in Eq. (1), which is called the *parameter of integral equation*. The classes of functions and kernels under consideration were defined above in Subsections 12.1-1 and 12.1-2. Note that equations of the form (1) with constant limits of integration and with Fredholm kernels or kernels with weak singularity are called *Fredholm equations of the second kind* and *equations with weak singularity of the second kind*, respectively.

A number λ is called a *characteristic value* of the integral equation (1) if there exist nontrivial solutions of the corresponding homogeneous equation (with $f(x) \equiv 0$). The nontrivial solutions themselves are called the *eigenfunctions* of the integral equation corresponding to the characteristic value λ . If λ is a characteristic value, the number $1/\lambda$ is called an *eigenvalue* of the integral equation (1). A value of the parameter λ is said to be *regular* if for this value the above homogeneous equation has only the trivial solution. Sometimes the characteristic values and the eigenfunctions of a Fredholm integral equation are called the *characteristic values* and the *eigenfunctions of the kernel* $K(x, t)$.

The kernel $K(x, t)$ of the integral equation (1) is called a *degenerate kernel* if it has the form $K(x, t) = g_1(x)h_1(t) + \dots + g_n(x)h_n(t)$, a *difference kernel* if it depends on the difference of the arguments ($K(x, t) = K(x-t)$), and a *symmetric kernel* if it satisfies the condition $K(x, t) = K(t, x)$.

The *transposed* integral equation is obtained from (1) by replacing the kernel $K(x, t)$ by $K(t, x)$.

Remark 1. The variables t and x may vary in different ranges (e.g., $a \leq t \leq b$ and $c \leq x \leq d$). To be specific, from now on we assume that $c = a$ and $d = b$ (this can be achieved by the linear substitution $x = \alpha\bar{x} + \beta$ with the aid of an appropriate choice of the constants α and β).

Remark 2. In general, the case in which the limits of integration a and/or b can be infinite is not excluded; however, in this case, the validity of the condition that the kernel $K(x, t)$ is square integrable on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ is especially significant.

13.1-2. Structure of the Solution.

The solution of Eq. (1) can be presented in the form

$$y(x) = f(x) + \lambda \int_a^b R(x, t; \lambda) f(t) dt,$$

where the resolvent $R(x, t; \lambda)$ is independent of $f(x)$ and is determined by the kernel of the integral equation.

The resolvent of the Fredholm equation (1) satisfies the following two integral equations:

$$R(x, t; \lambda) = K(x, t) + \int_a^b K(x, s)R(s, t; \lambda) ds,$$

$$R(x, t; \lambda) = K(x, t) + \int_a^b K(s, t)R(x, s; \lambda) ds,$$

in which the integration is performed with respect to different pairs of arguments of the kernel and the resolvent.

13.1-3. Integral Equations of Convolution Type of the Second Kind.

By the *integral equations of convolution type* (see also Subsection 12.1-3) we mean the integral equations that can be reduced, by applying some integral transform and the convolution theorem for this transform, to an algebraic equation for the transforms or to boundary value problems of the theory of analytic functions. Consider equations of convolution type of the second kind related to the Fourier transform.

An integral equation of the second kind with difference kernel on the entire axis (this equation is sometimes called an *equation of convolution type of the second kind with a single kernel*) has the form

$$y(x) + \int_{-\infty}^{\infty} K(x-t)y(t) dt = f(x), \quad -\infty < x < \infty, \quad (2)$$

where $f(x)$ and $K(x)$ are the right-hand side and the kernel of the integral equation and $y(x)$ is the function to be found.

An integral equation of the second kind with difference kernel on the semiaxis has the form

$$y(x) + \int_0^{\infty} K(x-t)y(t) dt = f(x), \quad 0 < x < \infty. \quad (3)$$

Equation (3) is also called a *one-sided equation of the second kind* or a *Wiener–Hopf integral equation of the second kind*.

An integral equation of convolution type of the second kind with two kernels has the form

$$y(x) + \int_0^{\infty} K_1(x-t)y(t) dt + \int_{-\infty}^0 K_2(x-t)y(t) dt = f(x), \quad -\infty < x < \infty, \quad (4)$$

where $K_1(x)$ and $K_2(x)$ are the *kernels* of the integral equation (4). The class of functions and kernels for equations of convolution type was introduced above in Subsection 12.1-3.

13.1-4. Dual Integral Equations of the Second Kind.

A dual integral equation of the second kind with difference kernels (of convolution type) has the form

$$\begin{aligned} y(x) + \int_{-\infty}^{\infty} K_1(x-t)y(t) dt &= f(x), & 0 < x < \infty, \\ y(x) + \int_{-\infty}^{\infty} K_2(x-t)y(t) dt &= f(x), & -\infty < x < 0, \end{aligned} \quad (5)$$

where the notation and the class of the functions and kernels coincide with those introduced for the equations of convolution type in Subsection 12.1-3.

In a sufficiently general case, a dual integral equation of the second kind has the form

$$\begin{aligned} y(x) + \int_a^{\infty} K_1(x,t)y(t) dt &= f_1(x), & a < x < b, \\ y(x) + \int_a^{\infty} K_2(x,t)y(t) dt &= f_2(x), & b < x < \infty, \end{aligned} \quad (6)$$

where $f_1(x)$ and $f_2(x)$ (and $K_1(x,t)$ and $K_2(x,t)$) are the known right-hand sides (and the kernels) of Eq. (6) and $y(x)$ is the function to be found. These equations can be studied by the methods of various integral transforms with reduction to boundary value problems of the theory of analytic functions and also by other methods developed for dual integral equations of the first kind (e.g., see I. Sneddon (1995) and Ya. S. Uflyand (1977)).

The integral equations obtained from (2)–(5) by replacing the kernel $K(x-t)$ by $K(t-x)$ are said to be *transposed* to the original equations.

If the right-hand sides of Eqs. (1)–(6) are identically zero, then these equations are said to be *homogeneous*. For the case in which the right-hand side of an equation of the type (1)–(6) does not vanish on the entire domain, the corresponding equation is said to be *nonhomogeneous*.

Remark 3. Some equations whose kernel contains the product or the ratio of the variables x and t can be reduced to Eqs. (2)–(5).

Remark 4. Sometimes equations of convolution type of the form (2)–(5) are written in the form in which the integrals are multiplied by the coefficient $1/\sqrt{2\pi}$.

Remark 5. The cases in which the class of functions and kernels for equations of convolution type (in particular, for Wiener–Hopf equations) differs from those introduced in Subsections 12.1-3 are always mentioned explicitly (see Sections 13.11 and 13.12).

⊙ References for Section 13.1: E. Goursat (1923), F. Riesz and B. Sz.-Nagy (1955), I. G. Petrovskii (1957), B. Noble (1958), M. G. Krein (1958), S. G. Mikhlin (1960), L. V. Kantorovich and G. P. Akilov (1964), A. N. Kolmogorov and S. V. Fomin (1970), L. Ya. Tslaf (1970), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), J. A. Cochran (1972), V. I. Smirnov (1974), P. P. Zabreyko, A. I. Koshelev, et al. (1975), F. D. Gakhov and Yu. I. Cherskii (1978), A. G. Butkovskii (1979), L. M. Delves and J. L. Mohamed (1985), F. G. Tricomi (1985), A. J. Jerry (1985), A. F. Verlan' and V. S. Sizikov (1986), A. Golberg (1990), D. Porter and D. S. G. Stirling (1990), C. Corduneanu (1991), J. Kondo (1991), S. Prössdorf and B. Silbermann (1991), W. Hackbusch (1995), R. P. Kanwal (1996).

13.2. Fredholm Equations of the Second Kind with Degenerate Kernel. Some Generalizations

13.2-1. Simplest Degenerate Kernel.

Consider Fredholm integral equations of the second kind with the simplest degenerate kernel:

$$y(x) - \lambda \int_a^b g(x)h(t)y(t) dt = f(x), \quad a \leq x \leq b. \quad (1)$$

We seek a solution of Eq. (1) in the form

$$y(x) = f(x) + \lambda Ag(x). \quad (2)$$

On substituting the expressions (2) into Eq. (1), after simple algebraic manipulations we obtain

$$A \left[1 - \lambda \int_a^b h(t)g(t) dt \right] = \int_a^b f(t)h(t) dt. \quad (3)$$

Both integrals occurring in Eq. (3) are supposed to exist. On the basis of (1)–(3) and taking into account the fact that the unique characteristic value λ_1 of Eq. (1) is given by the expression

$$\lambda_1 = \left[\int_a^b h(t)g(t) dt \right]^{-1}, \quad (4)$$

we obtain the following results.

1°. If $\lambda \neq \lambda_1$, then for an arbitrary right-hand side there exists a unique solution of Eq. (1), which can be written in the form

$$y(x) = f(x) + \frac{\lambda \lambda_1 f_1}{\lambda_1 - \lambda} g(x), \quad f_1 = \int_a^b f(t)h(t) dt. \quad (5)$$

2°. If $\lambda = \lambda_1$ and $f_1 = 0$, then any solution of Eq. (1) can be represented in the form

$$y = f(x) + Cy_1(x), \quad y_1(x) = g(x), \quad (6)$$

where C is an arbitrary constant and $y_1(x)$ is an eigenfunction that corresponds to the characteristic value λ_1 .

3°. If $\lambda = \lambda_1$ and $f_1 \neq 0$, then there are no solutions.

13.2-2. Degenerate Kernel in the General Case.

In the general case, a Fredholm integral equation of the second kind with degenerate kernel has the form

$$y(x) - \lambda \int_a^b \left[\sum_{k=1}^n g_k(x)h_k(t) \right] y(t) dt = f(x), \quad n = 2, 3, \dots \quad (7)$$

Let us rewrite Eq. (7) in the form

$$y(x) = f(x) + \lambda \sum_{k=1}^n g_k(x) \int_a^b h_k(t)y(t) dt, \quad n = 2, 3, \dots \quad (8)$$

We assume that Eq. (8) has a solution and introduce the notation

$$A_k = \int_a^b h_k(t)y(t) dt. \quad (9)$$

In this case we have

$$y(x) = f(x) + \lambda \sum_{k=1}^n A_k g_k(x), \quad (10)$$

and hence the solution of the integral equation with degenerate kernel is reduced to the definition of the constants A_k .

Let us multiply Eq. (10) by $h_m(x)$ and integrate with respect to x from a to b . We obtain the following system of linear algebraic equations for the coefficients A_k :

$$A_m - \lambda \sum_{k=1}^n s_{mk} A_k = f_m, \quad m = 1, \dots, n, \quad (11)$$

where

$$s_{mk} = \int_a^b h_m(x) g_k(x) dx, \quad f_m = \int_a^b f(x) h_m(x) dx; \quad m, k = 1, \dots, n. \quad (12)$$

In the calculation of the coefficients s_{mk} and f_m for specific degenerate kernels, the tables of integrals can be applied; see Supplements 3 and 4, as well as I. S. Gradshteyn and I. M. Ryzhik (1980), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1986).

Once we construct a solution of system (11), we obtain a solution of the integral equation with degenerate kernel (7) as well. The values of the parameter λ at which the determinant of system (11) vanishes are characteristic values of the integral equation (7), and it is clear that there are just n such values counted according to their multiplicities.

Now we can state the main results on the solution of Eq. (7).

1°. If λ is a regular value, then for an arbitrary right-hand side $f(x)$, there exists a unique solution of the Fredholm integral equation with degenerate kernel and this solution can be represented in the form (10), in which the coefficients A_k make up a solution of system (11). The constants A_k can be determined, for instance, by Cramer's rule (see equation 4.9.20, Chapter 4).

2°. If λ is a characteristic value and $f(x) \equiv 0$, then every solution of the homogeneous equation with degenerate kernel has the form

$$y(x) = \sum_{i=1}^p C_i y_i(x), \quad (13)$$

where the C_i are arbitrary constants and the $y_i(x)$ are linearly independent eigenfunctions of the kernel corresponding to the characteristic value λ :

$$y_i(x) = \sum_{k=1}^n A_{k(i)} g_k(x). \quad (14)$$

Here the constants $A_{k(i)}$ form p ($p \leq n$) linearly independent solutions of the following homogeneous system of algebraic equations:

$$A_{m(i)} - \lambda \sum_{k=1}^n s_{mk} A_{k(i)} = 0; \quad m = 1, \dots, n, \quad i = 1, \dots, p. \quad (15)$$

3°. If λ is a characteristic value and $f(x) \neq 0$, then for the nonhomogeneous integral equation (7) to be solvable, it is necessary and sufficient that the right-hand side $f(x)$ is such that the p conditions

$$\sum_{k=1}^n B_{k(i)} f_k = 0, \quad i = 1, \dots, p, \quad p \leq n, \quad (16)$$

are satisfied. Here the constants $B_{k(i)}$ form p linearly independent solutions of the homogeneous system of algebraic equations which is the transpose of system (15). In this case, every solution of Eq. (7) has the form

$$y(x) = y_0(x) + \sum_{i=1}^p C_i y_i(x), \quad (17)$$

where $y_0(x)$ is a particular solution of the nonhomogeneous equation (7) and the sum represents the general solution of the corresponding homogeneous equation (see item 2°). In particular, if $f(x) \neq 0$ but all f_k are zero, we have

$$y(x) = f(x) + \sum_{i=1}^p C_i y_i(x). \quad (18)$$

Remark. When studying Fredholm equations of the second kind with degenerate kernel, it is useful for the reader to be acquainted with equations 4.9.18 and 4.9.20 of the first part of the book.

Example 1. Let us solve the integral equation

$$y(x) - \lambda \int_{-\pi}^{\pi} (x \cos t + t^2 \sin x + \cos x \sin t)y(t) dt = x, \quad -\pi \leq x \leq \pi. \quad (19)$$

Let us denote

$$A_1 = \int_{-\pi}^{\pi} y(t) \cos t dt, \quad A_2 = \int_{-\pi}^{\pi} t^2 y(t) dt, \quad A_3 = \int_{-\pi}^{\pi} y(t) \sin t dt, \quad (20)$$

where A_1 , A_2 , and A_3 are unknown constants. Then Eq. (19) can be rewritten in the form

$$y(x) = A_1 \lambda x + A_2 \lambda \sin x + A_3 \lambda \cos x + x. \quad (21)$$

On substituting the expression (21) into relations (20), we obtain

$$\begin{aligned} A_1 &= \int_{-\pi}^{\pi} (A_1 \lambda t + A_2 \lambda \sin t + A_3 \lambda \cos t + t) \cos t dt, \\ A_2 &= \int_{-\pi}^{\pi} (A_1 \lambda t + A_2 \lambda \sin t + A_3 \lambda \cos t + t) t^2 dt, \\ A_3 &= \int_{-\pi}^{\pi} (A_1 \lambda t + A_2 \lambda \sin t + A_3 \lambda \cos t + t) \sin t dt. \end{aligned}$$

On calculating the integrals occurring in these equations, we obtain the following system of algebraic equations for the unknowns A_1 , A_2 , and A_3 :

$$\begin{aligned} A_1 - \lambda \pi A_3 &= 0, \\ A_2 + 4\lambda \pi A_3 &= 0, \\ -2\lambda \pi A_1 - \lambda \pi A_2 + A_3 &= 2\pi. \end{aligned} \quad (22)$$

The determinant of this system is

$$\Delta(\lambda) = \begin{vmatrix} 1 & 0 & -\lambda \pi \\ 0 & 1 & 4\lambda \pi \\ -2\lambda \pi & -\lambda \pi & 1 \end{vmatrix} = 1 + 2\lambda^2 \pi^2 \neq 0.$$

Thus, system (22) has the unique solution

$$A_1 = \frac{2\lambda \pi^2}{1 + 2\lambda^2 \pi^2}, \quad A_2 = -\frac{8\lambda \pi^2}{1 + 2\lambda^2 \pi^2}, \quad A_3 = \frac{2\pi}{1 + 2\lambda^2 \pi^2}.$$

On substituting the above values of A_1 , A_2 , and A_3 into (21), we obtain the solution of the original integral equation:

$$y(x) = \frac{2\lambda \pi}{1 + 2\lambda^2 \pi^2} (\lambda \pi x - 4\lambda \pi \sin x + \cos x) + x.$$

13.2-3. Kernel is the Sum of a Nondegenerate Kernel and an Arbitrary Degenerate Kernel.

1°. Consider a linear integral equation of the second kind

$$y(x) + \int_a^b K(x, t)y(t) dt = f(x). \quad (23)$$

Suppose equation (23) can be solved for any $f(x)$ from some class of functions $L_{\mathcal{F}}$. Let $y_f(x)$ denote the corresponding solution.

Now consider the more complex integral equation

$$u(x) + \int_a^b [K(x, t) + \varphi(x)\psi(t)]u(t) dt = f(x), \quad (24)$$

with its kernel containing an additional term $\varphi(x)\psi(t)$. A solution to equation (24) will be sought in the form

$$u(x) = y_f(x) + Ay_{\varphi}(x), \quad (25)$$

where $y_{\varphi}(x)$ is the solution to equation (23) in which $f(x)$ must be replaced with $\varphi(x)$. Substituting (25) into (24) results in the coefficient A :

$$A = -\frac{\int_a^b \psi(t)y_f(t) dt}{1 + \int_a^b \psi(t)y_{\varphi}(t) dt}. \quad (26)$$

Formulas (25)–(26) define a solution to equation (24), provided the integrals in the numerator and denominator exist, with $\int_a^b \psi(t)y_{\varphi}(t) dt \neq -1$. In addition, the condition $\varphi(x) \in L_{\mathcal{F}}$ must be satisfied.

Example 2. The solution of the integral equation

$$y(x) - \lambda \int_0^{\infty} \sin(xt)y(t) dt = f(x) \quad (27)$$

is expressed as (see Eq. 4.5.20)

$$y_f(x) = \frac{f(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt)f(t) dt, \quad (28)$$

where $\lambda \neq \pm\sqrt{2/\pi}$.

Now consider the more complex integral equation

$$u(x) - \lambda \int_0^{\infty} [\sin(xt) + \varphi(x)\psi(t)]u(t) dt = f(x) \quad (29)$$

with its kernel containing an arbitrary additive function $\varphi(x)\psi(t)$. The corresponding solution (28) to equation (27) with $f(x) = \varphi(x)$ is written as

$$y_{\varphi}(x) = \frac{\varphi(x)}{1 - \frac{\pi}{2}\lambda^2} + \frac{\lambda}{1 - \frac{\pi}{2}\lambda^2} \int_0^{\infty} \sin(xt)\varphi(t) dt.$$

Hence, equation (29) has the solution

$$u(x) = y_f(x) + Ay_{\varphi}(x), \quad A = -\frac{\int_0^1 \psi(t)y_f(t) dt}{1 + \int_0^1 \psi(t)y_{\varphi}(t) dt}.$$

2°. The integral equation

$$u(x) + \int_a^b \left[K(x, t) + \sum_{m=1}^n \varphi_m(x)\psi_m(t) \right] u(t) dt = f(x), \quad (30)$$

with its kernel being the sum of the kernel of equation (23) and an arbitrary degenerate kernel, can be solved in a similar manner. The solution is sought in the additive form

$$u(x) = y_f(x) + \sum_{m=1}^n A_m y_{\varphi_m}(x), \quad (31)$$

where $y_{\varphi_m}(x)$ is the solution to equation (23) in which $f(x)$ must be replaced with $\varphi_m(x)$. Substituting (31) into (30) results in the following linear algebraic system of equations for the coefficients A_m :

$$A_m + \sum_{j=1}^n A_j \sigma_{mj} = -\sigma_{m0}, \quad m = 1, \dots, n;$$

$$\sigma_{mj} = \int_a^b \psi_m(t) y_{\varphi_j}(t) dt, \quad \sigma_{m0} = \int_a^b \psi_m(t) y_f(t) dt.$$

© References for Section 13.2: S. G. Mikhlin (1960), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), P. P. Zabreyko, A. I. Koshelev, et al. (1975), A. J. Jerry (1985), A. F. Verlan' and V. S. Sizikov (1986), A. D. Polyanin and A. I. Zhurov (2007).

13.3. Solution as a Power Series in the Parameter. Method of Successive Approximations

13.3-1. Iterated Kernels.

Consider the Fredholm integral equation of the second kind:

$$y(x) - \lambda \int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b. \quad (1)$$

We seek the solution in the form of a series in powers of the parameter λ :

$$y(x) = f(x) + \sum_{n=1}^{\infty} \lambda^n \psi_n(x). \quad (2)$$

Substitute series (2) into Eq. (1). On matching the coefficients of like powers of λ , we obtain a recurrent system of equations for the functions $\psi_n(x)$. The solution of this system yields

$$\psi_1(x) = \int_a^b K(x, t)f(t) dt,$$

$$\psi_2(x) = \int_a^b K(x, t)\psi_1(t) dt = \int_a^b K_2(x, t)f(t) dt,$$

$$\psi_3(x) = \int_a^b K(x, t)\psi_2(t) dt = \int_a^b K_3(x, t)f(t) dt, \quad \text{etc.}$$

Here

$$K_n(x, t) = \int_a^b K(x, z)K_{n-1}(z, t) dz, \quad (3)$$

where $n = 2, 3, \dots$, and we have $K_1(x, t) \equiv K(x, t)$. The functions $K_n(x, t)$ defined by formulas (3) are called *iterated kernels*. These kernels satisfy the relation

$$K_n(x, t) = \int_a^b K_m(x, s)K_{n-m}(s, t) ds, \quad (4)$$

where m is an arbitrary positive integer less than n .

The iterated kernels $K_n(x, t)$ can be directly expressed via $K(x, t)$ by the formula

$$K_n(x, t) = \underbrace{\int_a^b \int_a^b \cdots \int_a^b}_{n-1} K(x, s_1) K(s_1, s_2) \cdots K(s_{n-1}, t) ds_1 ds_2 \cdots ds_{n-1}.$$

All iterated kernels $K_n(x, t)$, beginning with $K_2(x, t)$, are continuous functions on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ if the original kernel $K(x, t)$ is square integrable on S .

If $K(x, t)$ is symmetric, then all iterated kernels $K_n(x, t)$ are also symmetric.

13.3-2. Method of Successive Approximations.

The results of Subsection 13.3-1 can also be obtained by means of the method of successive approximations. To this end, one should use the recurrent formula

$$y_n(x) = f(x) + \lambda \int_a^b K(x, t) y_{n-1}(t) dt, \quad n = 1, 2, \dots,$$

with the zeroth approximation $y_0(x) = f(x)$.

13.3-3. Construction of the Resolvent.

The resolvent of the integral equation (1) is defined via the iterated kernels by the formula

$$R(x, t; \lambda) = \sum_{n=1}^{\infty} \lambda^{n-1} K_n(x, t), \quad (5)$$

where the series on the right-hand side is called the *Neumann series of the kernel* $K(x, t)$. It converges to a unique square integrable solution of Eq. (1) provided that

$$|\lambda| < \frac{1}{B}, \quad B = \sqrt{\int_a^b \int_a^b K^2(x, t) dx dt}. \quad (6)$$

If, in addition, we have

$$\int_a^b K^2(x, t) dt \leq A, \quad a \leq x \leq b,$$

where A is a constant, then the Neumann series converges absolutely and uniformly on $[a, b]$.

A solution of a Fredholm equation of the second kind of the form (1) is expressed by the formula

$$y(x) = f(x) + \lambda \int_a^b R(x, t; \lambda) f(t) dt, \quad a \leq x \leq b. \quad (7)$$

Inequality (6) is essential for the convergence of the series (5). However, a solution of Eq. (1) can exist for values $|\lambda| > 1/B$ as well.

Remark 1. A solution of the equation

$$y(x) - \lambda \int_a^b K(x, t) y(t) dt = f(x), \quad a \leq x \leq b,$$

with weak singularity, where the kernel $K(x, t)$ has the form

$$K(x, t) = \frac{L(x, t)}{|x - t|^\alpha}, \quad 0 < \alpha < 1,$$

and $L(x, t)$ is a function continuous on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$, can be obtained by the successive approximation method provided that

$$|\lambda| < \frac{1 - \alpha}{2B^*(b - a)^{1-\alpha}}, \quad B^* = \sup |L(x, t)|.$$

The equation itself can be reduced to a Fredholm equation of the form

$$y(x) - \lambda^n \int_a^b K_n(x, t)y(t) dt = F(x), \quad a \leq x \leq b,$$

$$F(x) = f(x) + \sum_{p=1}^{n-1} \lambda^p \int_a^b K_p(x, t)f(t) dt,$$

where $K_p(x, t)$ ($p = 1, \dots, n$) is the p th iterated kernel, with $K_n(x, t)$ being a Fredholm kernel for $n > \frac{1}{2}(1 - \alpha)^{-1}$ and bounded for $n > (1 - \alpha)^{-1}$.

Example 1. Let us solve the integral equation

$$y(x) - \lambda \int_0^1 xty(t) dt = f(x), \quad 0 \leq x \leq 1,$$

by the method of successive approximations. Here we have $K(x, t) = xt$, $a = 0$, and $b = 1$. We successively define

$$K_1(x, t) = xt, \quad K_2(x, t) = \int_0^1 (xz)(zt) dz = \frac{xt}{3}, \quad K_3(x, t) = \frac{1}{3} \int_0^1 (xz)(zt) dz = \frac{xt}{3^2}, \quad \dots, \quad K_n(x, t) = \frac{xt}{3^{n-1}}.$$

According to formula (5) for the resolvent, we obtain

$$R(x, t; \lambda) = \sum_{n=1}^{\infty} \lambda^{n-1} K_n(x, t) = xt \sum_{n=1}^{\infty} \left(\frac{\lambda}{3}\right)^{n-1} = \frac{3xt}{3 - \lambda},$$

where $|\lambda| < 3$, and it follows from formula (7) that the solution of the integral equation can be rewritten in the form

$$y(x) = f(x) + \lambda \int_0^1 \frac{3xt}{3 - \lambda} f(t) dt, \quad 0 \leq x \leq 1, \quad \lambda \neq 3.$$

In particular, for $f(x) = x$ we obtain

$$y(x) = \frac{3x}{3 - \lambda}, \quad 0 \leq x \leq 1, \quad \lambda \neq 3.$$

13.3-4. Orthogonal Kernels.

For some Fredholm equations, the Neumann series (5) for the resolvent is convergent for all values of λ . Let us establish this fact.

Assume that two kernels $K(x, t)$ and $L(x, t)$ are given. These kernels are said to be *orthogonal* if the following two conditions hold:

$$\int_a^b K(x, z)L(z, t) dz = 0, \quad \int_a^b L(x, z)K(z, t) dz = 0 \quad (8)$$

for all admissible values of x and t .

There exist kernels that are orthogonal to themselves. For these kernels we have $K_2(x, t) \equiv 0$, where $K_2(x, t)$ is the second iterated kernel. It is clear that in this case all the subsequent iterated kernels also vanish, and the resolvent coincides with the kernel $K(x, t)$.

Example 2. Let us find the resolvent of the kernel $K(x, t) = \sin(x - 2t)$, $0 \leq x \leq 2\pi$, $0 \leq t \leq 2\pi$. We have

$$\begin{aligned} \int_0^{2\pi} \sin(x - 2z) \sin(z - 2t) dz &= \frac{1}{2} \int_0^{2\pi} [\cos(x + 2t - 3z) - \cos(x - 2t - z)] dz = \\ &= \frac{1}{2} \left[-\frac{1}{3} \sin(x + 2t - 3z) + \sin(x - 2t - z) \right]_{z=0}^{z=2\pi} = 0. \end{aligned}$$

Thus, in this case the resolvent of the kernel is equal to the kernel itself:

$$R(x, t; \lambda) \equiv \sin(x - 2t),$$

so that the Neumann series (5) consists of a single term and clearly converges for any λ .

Example 3. The kernel

$$K(x, t) = \sum_{n=1}^{\infty} a_n \sin(nx) \cos(nt), \quad 0 \leq x, t \leq 2\pi,$$

with a convergent series $\sum_{n=1}^{\infty} |a_n|$ is orthogonal to itself.

Remark 2. If the kernels $M^{(1)}(x, t), \dots, M^{(n)}(x, t)$ are pairwise orthogonal, then the resolvent corresponding to the sum

$$K(x, t) = \sum_{m=1}^n M^{(m)}(x, t)$$

is equal to the sum of the resolvents corresponding to each of the summands.

⊙ References for Section 13.3: S. G. Mikhlin (1960), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), J. A. Cochran (1972), V. I. Smirnov (1974), A. J. Jerry (1985).

13.4. Method of Fredholm Determinants

13.4-1. Formula for the Resolvent.

A solution of the Fredholm equation of the second kind

$$y(x) - \lambda \int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b, \quad (1)$$

is given by the formula

$$y(x) = f(x) + \lambda \int_a^b R(x, t; \lambda)f(t) dt, \quad a \leq x \leq b, \quad (2)$$

where the resolvent $R(x, t; \lambda)$ is defined by the relation

$$R(x, t; \lambda) = \frac{D(x, t; \lambda)}{D(\lambda)}, \quad D(\lambda) \neq 0. \quad (3)$$

Here $D(x, t; \lambda)$ and $D(\lambda)$ are power series in λ ,

$$D(x, t; \lambda) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} A_n(x, t) \lambda^n, \quad D(\lambda) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} B_n \lambda^n, \quad (4)$$

with coefficients defined by the formulas

$$A_0(x, t) = K(x, t), \quad A_n(x, t) = \underbrace{\int_a^b \cdots \int_a^b}_n \begin{vmatrix} K(x, t) & K(x, t_1) & \cdots & K(x, t_n) \\ K(t_1, t) & K(t_1, t_1) & \cdots & K(t_1, t_n) \\ \vdots & \vdots & \ddots & \vdots \\ K(t_n, t) & K(t_n, t_1) & \cdots & K(t_n, t_n) \end{vmatrix} dt_1 \cdots dt_n, \quad (5)$$

$$B_0 = 1, \quad B_n = \underbrace{\int_a^b \cdots \int_a^b}_n \begin{vmatrix} K(t_1, t_1) & K(t_1, t_2) & \cdots & K(t_1, t_n) \\ K(t_2, t_1) & K(t_2, t_2) & \cdots & K(t_2, t_n) \\ \vdots & \vdots & \ddots & \vdots \\ K(t_n, t_1) & K(t_n, t_2) & \cdots & K(t_n, t_n) \end{vmatrix} dt_1 \cdots dt_n; \quad n = 0, 1, 2, \dots \quad (6)$$

The function $D(x, t; \lambda)$ is called the *Fredholm minor* and $D(\lambda)$ the *Fredholm determinant*. The series (4) converge for all values of λ and hence define entire analytic functions of λ . The resolvent $R(x, t; \lambda)$ is an analytic function of λ everywhere except for the values of λ that are roots of $D(\lambda)$. These roots coincide with the characteristic values of the equation and are poles of the resolvent $R(x, t; \lambda)$.

Example 1. Consider the integral equation

$$y(x) - \lambda \int_0^1 x e^t y(t) dt = f(x), \quad 0 \leq x \leq 1, \quad \lambda \neq 1.$$

We have

$$A_0(x, t) = x e^t, \quad A_1(x, t) = \int_0^1 \begin{vmatrix} x e^t & x e^{t_1} \\ t_1 e^t & t_1 e^{t_1} \end{vmatrix} dt_1 = 0, \quad A_2(x, t) = \int_0^1 \int_0^1 \begin{vmatrix} x e^t & x e^{t_1} & x e^{t_2} \\ t_1 e^t & t_1 e^{t_1} & t_1 e^{t_2} \\ t_2 e^t & t_2 e^{t_1} & t_2 e^{t_2} \end{vmatrix} dt_1 dt_2 = 0,$$

since the determinants in the integrand are zero. It is clear that the relation $A_n(x, t) = 0$ holds for the subsequent coefficients. Let us find the coefficients B_n :

$$B_1 = \int_0^1 K(t_1, t_1) dt_1 = \int_0^1 t_1 e^{t_1} dt_1 = 1, \quad B_2 = \int_0^1 \int_0^1 \begin{vmatrix} t_1 e^{t_1} & t_1 e^{t_2} \\ t_2 e^{t_1} & t_2 e^{t_2} \end{vmatrix} dt_1 dt_2 = 0.$$

It is clear that $B_n = 0$ for all subsequent coefficients as well.

According to formulas (4), we have

$$D(x, t; \lambda) = K(x, t) = x e^t; \quad D(\lambda) = 1 - \lambda.$$

Thus,

$$R(x, t; \lambda) = \frac{D(x, t; \lambda)}{D(\lambda)} = \frac{x e^t}{1 - \lambda},$$

and the solution of the equation can be represented in the form

$$y(x) = f(x) + \lambda \int_0^1 \frac{x e^t}{1 - \lambda} f(t) dt, \quad 0 \leq x \leq 1, \quad \lambda \neq 1.$$

In particular, for $f(x) = e^{-x}$ we obtain

$$y(x) = e^{-x} + \frac{\lambda}{1 - \lambda} x, \quad 0 \leq x \leq 1, \quad \lambda \neq 1.$$

13.4-2. Recurrent Relations.

In practice, the calculation of the coefficients $A_n(x, t)$ and B_n of the series (4) by means of formulas (5) and (6) is seldom possible. However, formulas (5) and (6) imply the following recurrent relations:

$$A_n(x, t) = B_n K(x, t) - n \int_a^b K(x, s) A_{n-1}(s, t) ds, \quad (7)$$

$$B_n = \int_a^b A_{n-1}(s, s) ds. \quad (8)$$

Example 2. Let us use formulas (7) and (8) to find the resolvent of the kernel $K(x, t) = x - 2t$, where $0 \leq x \leq 1$ and $0 \leq t \leq 1$.

Indeed, we have $B_0 = 1$ and $A_0(x, t) = x - 2t$. Applying formula (8), we see that

$$B_1 = \int_0^1 (-s) ds = -\frac{1}{2}.$$

Formula (7) implies the relation

$$A_1(x, t) = -\frac{x-2t}{2} - \int_0^1 (x-2s)(s-2t) ds = -x-t+2xt+\frac{2}{3}.$$

Furthermore, we have

$$\begin{aligned} B_2 &= \int_0^1 (-2s+2s^2+\frac{2}{3}) ds = \frac{1}{3}, \\ A_2(x, t) &= \frac{x-2t}{3} - 2 \int_0^1 (x-2s)(-s-t+2st+\frac{2}{3}) ds = 0, \\ B_3 = B_4 = \dots &= 0, \quad A_3(x, t) = A_4(x, t) = \dots = 0. \end{aligned}$$

Hence,

$$D(\lambda) = 1 + \frac{1}{2}\lambda + \frac{1}{6}\lambda^2; \quad D(x, t; \lambda) = x - 2t + \lambda(x + t - 2xt - \frac{2}{3}).$$

The resolvent has the form

$$R(x, t; \lambda) = \frac{x-2t + \lambda(x+t-2xt-\frac{2}{3})}{1 + \frac{1}{2}\lambda + \frac{1}{6}\lambda^2}.$$

⊙ References for Section 13.4: S. G. Mikhlin (1960), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), V. I. Smirnov (1974).

13.5. Fredholm Theorems and the Fredholm Alternative

13.5-1. Fredholm Theorems.

THEOREM 1. *If λ is a regular value, then both the Fredholm integral equation of the second kind and the transposed equation are solvable for any right-hand side, and both the equations have unique solutions. The corresponding homogeneous equations have only the trivial solutions.*

THEOREM 2. *For the nonhomogeneous integral equation to be solvable, it is necessary and sufficient that the right-hand side $f(x)$ satisfies the conditions*

$$\int_a^b f(x)\psi_k(x) dx = 0, \quad k = 1, \dots, n,$$

where $\psi_k(x)$ is a complete set of linearly independent solutions of the corresponding transposed homogeneous equation.

THEOREM 3. *If λ is a characteristic value, then both the homogeneous integral equation and the transposed homogeneous equation have nontrivial solutions. The number of linearly independent solutions of the homogeneous integral equation is finite and is equal to the number of linearly independent solutions of the transposed homogeneous equation.*

THEOREM 4. *A Fredholm equation of the second kind has at most countably many characteristic values, whose only possible accumulation point is the point at infinity.*

Example. To illustrate the Fredholm theorems, consider the degenerate integral equation

$$y(x) - \lambda \int_0^\pi \sin(x+t)y(t) dt = f(x). \quad (1)$$

Using the trigonometric formula

$$\sin(x + t) = \sin x \cos t + \cos x \sin t,$$

we transform equation (1) to

$$y(x) = \lambda A \sin x + \lambda B \cos x + f(x), \quad (2)$$

$$A = \int_0^\pi \cos t y(t) dt, \quad B = \int_0^\pi \sin t y(t) dt. \quad (3)$$

Substituting (2) into (3), we come to the system of linear algebraic equations for the coefficients A and B :

$$\begin{aligned} A - \frac{1}{2}\pi\lambda B &= f_1, \\ -\frac{1}{2}\pi\lambda A + B &= f_2, \end{aligned} \quad (4)$$

where

$$f_1 = \int_0^\pi f(t) \cos t dt, \quad f_2 = \int_0^\pi f(t) \sin t dt.$$

The determinant of this system

$$\begin{vmatrix} 1 & -\frac{1}{2}\pi\lambda \\ -\frac{1}{2}\pi\lambda & 1 \end{vmatrix} = 1 - \frac{1}{4}\pi^2\lambda^2$$

has the roots

$$\lambda_1 = -2/\pi, \quad \lambda_2 = 2/\pi, \quad (5)$$

which coincide with the characteristic values of equation (1); the other values $\lambda \neq \pm 2/\pi$ are regular.

If λ differs from the characteristic values (5), then the determinant of system (4) differs from zero and the coefficients A and B are uniquely defined by

$$A = \frac{f_1 + \frac{1}{2}\pi\lambda f_2}{1 - \frac{1}{4}\pi^2\lambda^2}, \quad B = \frac{\frac{1}{2}\pi\lambda f_1 + f_2}{1 - \frac{1}{4}\pi^2\lambda^2}$$

and yield the unique solution (2) of the nonhomogeneous integral equation (1). For these values $\lambda \neq \pm 2/\pi$, the corresponding homogeneous integral equation (1) with $f(x) \equiv 0$ has only the trivial solution $y(x) \equiv 0$. (This illustrates Theorem 1.)

Now, suppose that $\lambda = \lambda_{1,2}$ is one of the characteristic values (5). In this case, both equations of the homogeneous system (4) with $f(x) \equiv 0$ are proportional and one can take $A = \frac{1}{2}\pi\lambda B$, where B is an arbitrary constant. The corresponding eigenfunctions have the form

$$y_{1,2}(x) = \frac{2}{\pi} B(\sin x \mp \cos x). \quad (6)$$

The constant B can be chosen, for instance, from the following normalization condition for eigenfunctions:

$$\|y_{1,2}\|^2 = \int_0^\pi |y_{1,2}(x)|^2 dx = 1,$$

which yields $B = \frac{1}{2}\sqrt{\pi}$.

If we take $\lambda = \lambda_1 = -2/\pi$ in the nonhomogeneous equation (1), then the algebraic system (4) takes the form

$$A + B = f_1, \quad A + B = f_2;$$

and for its solvability it is necessary and sufficient that $f_1 = f_2$. This condition means that the right-hand side $f(x)$ is orthogonal to the eigenfunction $y_1(x)$. Similarly, for $\lambda = \lambda_2 = 2/\pi$, system (4) has a solution if and only if $f_1 = -f_2$, i.e., the functions $f(x)$ and $y_2(x)$ are orthogonal. The eigenfunctions obtained in this example and the orthogonality conditions illustrate the statements of Theorems 2 and 3 (in the case under consideration, the kernel $K(x, t)$ coincides with its conjugate).

13.5-2. Fredholm Alternative.

The Fredholm theorems imply the so-called Fredholm alternative, which is most frequently used in the investigation of integral equations.

THE FREDHOLM ALTERNATIVE. *Either the nonhomogeneous equation is solvable for any right-hand side or the corresponding homogeneous equation has nontrivial solutions.*

The first part of the alternative holds if the given value of the parameter is regular and the second if it is characteristic.

Remark. The Fredholm theory is also valid for integral equations of the second kind with weak singularity.

© References for Section 13.5: S. G. Mikhlin (1960), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), J. A. Cochran (1972), V. I. Smirnov (1974), A. J. Jerry (1985), D. Porter and D. S. G. Stirling (1990), C. Corduneanu (1991), J. Kondo (1991), W. Hackbusch (1995), R. P. Kanwal (1996), R. Kress (1999).

13.6. Fredholm Integral Equations of the Second Kind with Symmetric Kernel

13.6-1. Characteristic Values and Eigenfunctions.

Integral equations whose kernels are *symmetric*, that is, satisfy the condition $K(x, t) = K(t, x)$, are called *symmetric integral equations*.

Each symmetric kernel that is not identically zero has at least one characteristic value.

For any n , the set of characteristic values of the n th iterated kernel coincides with the set of n th powers of the characteristic values of the first kernel.

The eigenfunctions of a symmetric kernel corresponding to distinct characteristic values are orthogonal, i.e., if

$$\varphi_1(x) = \lambda_1 \int_a^b K(x, t)\varphi_1(t) dt, \quad \varphi_2(x) = \lambda_2 \int_a^b K(x, t)\varphi_2(t) dt, \quad \lambda_1 \neq \lambda_2,$$

then

$$(\varphi_1, \varphi_2) = 0, \quad (\varphi, \psi) \equiv \int_a^b \varphi(x)\psi(x) dx.$$

The characteristic values of a symmetric kernel are real.

The eigenfunctions can be normalized; namely, we can divide each characteristic function by its norm. If several linearly independent eigenfunctions correspond to the same characteristic value, say, $\varphi_1(x), \dots, \varphi_n(x)$, then each linear combination of these functions is an eigenfunction as well, and these linear combinations can be chosen so that the corresponding eigenfunctions are orthonormal.

Indeed, the function

$$\psi_1(x) = \frac{\varphi_1(x)}{\|\varphi_1\|}, \quad \|\varphi_1\| = \sqrt{(\varphi_1, \varphi_1)},$$

has the norm equal to one, i.e., $\|\psi_1\| = 1$. Let us form a linear combination $\alpha\psi_1 + \varphi_2$ and choose α so that

$$(\alpha\psi_1 + \varphi_2, \psi_1) = 0,$$

i.e.,

$$\alpha = -\frac{(\varphi_2, \psi_1)}{(\psi_1, \psi_1)} = -(\varphi_2, \psi_1).$$

The function

$$\psi_2(x) = \frac{\alpha\psi_1 + \varphi_2}{\|\alpha\psi_1 + \varphi_2\|}$$

is orthogonal to $\psi_1(x)$ and has the unit norm. Next, we choose a linear combination $\alpha\psi_1 + \beta\psi_2 + \varphi_3$, where the constants α and β can be found from the orthogonality relations

$$(\alpha\psi_1 + \beta\varphi_2 + \varphi_3, \psi_1) = 0, \quad (\alpha\psi_1 + \beta\psi_2 + \varphi_3, \psi_2) = 0.$$

For the coefficients α and β thus defined, the function

$$\psi_3 = \frac{\alpha\psi_1 + \beta\psi_2 + \varphi_3}{\|\alpha\psi_1 + \beta\varphi_2 + \varphi_3\|}$$

is orthogonal to ψ_1 and ψ_2 and has the unit norm, and so on.

As was noted above, the eigenfunctions corresponding to distinct characteristic values are orthogonal. Hence, the sequence of eigenfunctions of a symmetric kernel can be made orthonormal.

In what follows we assume that the sequence of eigenfunctions of a symmetric kernel is orthonormal.

We also assume that the characteristic values are always numbered in the increasing order of their absolute values. Thus, if

$$\lambda_1, \lambda_2, \dots, \lambda_n, \dots \quad (1)$$

is the sequence of characteristic values of a symmetric kernel, and if a sequence of eigenfunctions

$$\varphi_1, \varphi_2, \dots, \varphi_n, \dots \quad (2)$$

corresponds to the sequence (1) so that

$$\varphi_n(x) - \lambda_n \int_a^b K(x,t)\varphi_n(t) dt = 0, \quad (3)$$

then

$$\int_a^b \varphi_i(x)\varphi_j(x) dx = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j, \end{cases} \quad (4)$$

and

$$|\lambda_1| \leq |\lambda_2| \leq \dots \leq |\lambda_n| \leq \dots \quad (5)$$

If there are infinitely many characteristic values, then it follows from the fourth Fredholm theorem that their only accumulation point is the point at infinity, and hence $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$.

The set of all characteristic values and the corresponding normalized eigenfunctions of a symmetric kernel is called the *system of characteristic values and eigenfunctions* of the kernel. The system of eigenfunctions is said to be *incomplete* if there exists a nonzero square integrable function that is orthogonal to all functions of the system. Otherwise, the system of eigenfunctions is said to be *complete*.

13.6-2. Bilinear Series.

Assume that a kernel $K(x,t)$ admits an expansion in a uniformly convergent series with respect to the orthonormal system of its eigenfunctions:

$$K(x,t) = \sum_{k=1}^{\infty} a_k(x)\varphi_k(t) \quad (6)$$

for all x in the case of a continuous kernel or for almost all x in the case of a square integrable kernel.

We have

$$a_k(x) = \int_a^b K(x,t)\varphi_k(t) dt = \frac{\varphi_k(x)}{\lambda_k}, \quad (7)$$

and hence

$$K(x,t) = \sum_{k=1}^{\infty} \frac{\varphi_k(x)\varphi_k(t)}{\lambda_k}. \quad (8)$$

Conversely, if the series

$$\sum_{k=1}^{\infty} \frac{\varphi_k(x)\varphi_k(t)}{\lambda_k} \quad (9)$$

is uniformly convergent, then

$$K(x, t) = \sum_{k=1}^{\infty} \frac{\varphi_k(x)\varphi_k(t)}{\lambda_k}.$$

The following assertion holds: the bilinear series (9) converges in mean-square to the kernel $K(x, t)$.

If a symmetric kernel $K(x, t)$ has finitely many characteristic values, then it is degenerate, because in this case we have

$$K(x, t) = \sum_{k=1}^n \frac{\varphi_k(x)\varphi_k(t)}{\lambda_k}. \quad (10)$$

A kernel $K(x, t)$ is said to be *positive definite* if for all functions $\varphi(x)$ that are not identically zero we have

$$\int_a^b \int_a^b K(x, t)\varphi(x)\varphi(t) dx dt > 0,$$

and the above quadratic functional vanishes for $\varphi(x)=0$ only. Such a kernel has positive characteristic values only. A *negative definite* kernel is defined similarly.

Each symmetric positive definite (or negative definite) continuous kernel can be decomposed in a bilinear series in eigenfunctions that is absolutely and uniformly convergent with respect to the variables x, t .

The assertion remains valid if we assume that the kernel has finitely many negative (positive, respectively) characteristic values.

If a kernel $K(x, t)$ is symmetric, continuous on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$, and has uniformly bounded partial derivatives on this square, then this kernel can be expanded in a uniformly convergent bilinear series in eigenfunctions.

13.6-3. Hilbert–Schmidt Theorem.

If a function $f(x)$ can be represented in the form

$$f(x) = \int_a^b K(x, t)g(t) dt, \quad (11)$$

where the symmetric kernel $K(x, t)$ is square integrable and $g(t)$ is a square integrable function, then $f(x)$ can be represented by its *Fourier series* with respect to the orthonormal system of eigenfunctions of the kernel $K(x, t)$:

$$f(x) = \sum_{k=1}^{\infty} a_k \varphi_k(x), \quad (12)$$

where

$$a_k = \int_a^b f(x)\varphi_k(x) dx, \quad k = 1, 2, \dots$$

Moreover, if

$$\int_a^b K^2(x, t) dt \leq A < \infty, \quad (13)$$

then the series (12) is absolutely and uniformly convergent for any function $f(x)$ of the form (11).

Remark 1. In the Hilbert–Schmidt theorem, the completeness of the system of eigenfunctions is not assumed.

13.6-4. Bilinear Series of Iterated Kernels.

By the definition of the iterated kernels, we have

$$K_m(x, t) = \int_a^b K(x, z)K_{m-1}(z, t) dz, \quad m = 2, 3, \dots \quad (14)$$

The Fourier coefficients $a_k(t)$ of the kernel $K_m(x, t)$, regarded as a function of the variable x , with respect to the orthonormal system of eigenfunctions of the kernel $K(x, t)$ are equal to

$$a_k(t) = \int_a^b K_m(x, t)\varphi_k(x) dx = \frac{\varphi_k(t)}{\lambda_k^m}. \quad (15)$$

On applying the Hilbert–Schmidt theorem to (14), we obtain

$$K_m(x, t) = \sum_{k=1}^{\infty} \frac{\varphi_k(x)\varphi_k(t)}{\lambda_k^m}, \quad m = 2, 3, \dots \quad (16)$$

In formula (16), the sum of the series is understood as the limit in mean-square. If in addition to the above assumptions, inequality (13) is satisfied, then the series in (16) is uniformly convergent.

13.6-5. Solution of the Nonhomogeneous Equation.

Let us represent an integral equation

$$y(x) - \lambda \int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b, \quad (17)$$

where the parameter λ is not a characteristic value, in the form

$$y(x) - f(x) = \lambda \int_a^b K(x, t)y(t) dt \quad (18)$$

and apply the Hilbert–Schmidt theorem to the function $y(x) - f(x)$:

$$y(x) - f(x) = \sum_{k=1}^{\infty} A_k \varphi_k(x),$$

$$A_k = \int_a^b [y(x) - f(x)]\varphi_k(x) dx = \int_a^b y(x)\varphi_k(x) dx - \int_a^b f(x)\varphi_k(x) dx = y_k - f_k.$$

Taking into account the expansion (8), we obtain

$$\lambda \int_a^b K(x, t)y(t) dt = \lambda \sum_{k=1}^{\infty} \frac{y_k}{\lambda_k} \varphi_k(x),$$

and thus

$$\lambda \frac{y_k}{\lambda_k} = y_k - f_k, \quad y_k = \frac{\lambda_k f_k}{\lambda_k - \lambda}, \quad A_k = \frac{\lambda f_k}{\lambda_k - \lambda}. \quad (19)$$

Hence,

$$y(x) = f(x) + \lambda \sum_{k=1}^{\infty} \frac{f_k}{\lambda_k - \lambda} \varphi_k(x). \quad (20)$$

However, if λ is a characteristic value, i.e.,

$$\lambda = \lambda_p = \lambda_{p+1} = \cdots = \lambda_q, \quad (21)$$

then, for $k \neq p, p+1, \dots, q$, the terms (20) preserve their form. For $k = p, p+1, \dots, q$, formula (19) implies the relation $f_k = A_k(\lambda - \lambda_k)/\lambda$, and by (21) we obtain $f_p = f_{p+1} = \cdots = f_q = 0$. The last relation means that

$$\int_a^b f(x)\varphi_k(x) dx = 0$$

for $k = p, p+1, \dots, q$, i.e., the right-hand side of the equation must be orthogonal to the eigenfunctions that correspond to the characteristic value λ .

In this case, the solutions of Eqs. (17) have the form

$$y(x) = f(x) + \lambda \sum_{k=1}^{\infty} \frac{f_k}{\lambda_k - \lambda} \varphi_k(x) + \sum_{k=p}^q C_k \varphi_k(x), \quad (22)$$

where the terms in the first of the sums (22) with indices $k = p, p+1, \dots, q$ must be omitted (for these indices, f_k and $\lambda - \lambda_k$ vanish in this sum simultaneously). The coefficients C_k in the second sum are arbitrary constants.

Remark 2. On the basis of the bilinear expansion (8) and the Hilbert–Schmidt theorem, the solution of the symmetric Fredholm integral equation of the first kind

$$\int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b,$$

can be constructed in a similar way in the form

$$y(x) = \sum_{k=1}^{\infty} f_k \lambda_k \varphi_k(x),$$

and the necessary and sufficient condition for the existence and uniqueness of such a solution in $L_2(a, b)$ is the completeness of the system of the eigenfunctions $\varphi_k(x)$ of the kernel $K(x, t)$ together with the convergence of the series $\sum_{k=1}^{\infty} f_k^2 \lambda_k^2$, where the λ_k are the corresponding characteristic values.

It should be noted that the verification of the last condition for specific equations is quite complicated. In the solution of Fredholm equations of the first kind, the methods presented in Chapter 12 are usually applied.

13.6-6. Fredholm Alternative for Symmetric Equations.

The above results can be unified in the following alternative form.

A symmetric integral equation

$$y(x) - \lambda \int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b, \quad (23)$$

for a given λ , either has a unique square integrable solution for an arbitrarily given function $f(x) \in L_2(a, b)$, in particular, $y = 0$ for $f = 0$, or the corresponding homogeneous equation has finitely many linearly independent solutions $Y_1(x), \dots, Y_r(x)$, $r > 0$.

For the second case, the nonhomogeneous equation has a solution if and only if the right-hand side $f(x)$ is orthogonal to all the functions $Y_1(x), \dots, Y_r(x)$ on the interval $[a, b]$. Here the solution is defined only up to an arbitrary additive linear combination $A_1 Y_1(x) + \cdots + A_r Y_r(x)$.

13.6-7. Resolvent of a Symmetric Kernel.

The solution of a Fredholm equation of the second kind (23) can be written in the form

$$y(x) = f(x) + \lambda \int_a^b R(x, t; \lambda) f(t) dt, \quad (24)$$

where the resolvent $R(x, t; \lambda)$ is given by the series

$$R(x, t; \lambda) = \sum_{k=1}^{\infty} \frac{\varphi_k(x)\varphi_k(t)}{\lambda_k - \lambda}. \quad (25)$$

Here the collections $\varphi_k(x)$ and λ_k form the system of eigenfunctions and characteristic values of Eqs. (23). It follows from formula (25) that the resolvent of a symmetric kernel has only simple poles.

13.6-8. Extremal Properties of Characteristic Values and Eigenfunctions.

Let us introduce the notation

$$(u, w) = \int_a^b u(x)w(x) dx, \quad \|u\|^2 = (u, u),$$

$$(Ku, u) = \int_a^b \int_a^b K(x, t)u(x)u(t) dx dt,$$

where (u, w) is the *inner product* of functions $u(x)$ and $w(x)$, $\|u\|$ is the *norm* of a function $u(x)$, and (Ku, u) is the *quadratic form* generated by the kernel $K(x, t)$.

Let λ_1 be the characteristic value of the symmetric kernel $K(x, t)$ with minimum absolute value and let $y_1(x)$ be the eigenfunction corresponding to this value. Then

$$\frac{1}{|\lambda_1|} = \max_{y \neq 0} \frac{|(Ky, y)|}{\|y\|^2}, \quad (26)$$

in particular, the maximum is attained, and $y = y_1$ is a maximum point.

Let $\lambda_1, \dots, \lambda_n$ be the first n characteristic values of a symmetric kernel $K(x, t)$ (in the ascending order of their absolute values) and let $y_1(x), \dots, y_n(x)$ be orthonormal eigenfunctions corresponding to $\lambda_1, \dots, \lambda_n$, respectively. Then the formula

$$\frac{1}{|\lambda_{n+1}|} = \max \frac{|(Ky, y)|}{\|y\|^2} \quad (27)$$

is valid for the characteristic value λ_{n+1} following λ_n . The maximum is taken over the set of functions y which are orthogonal to all y_1, \dots, y_n and are not identically zero, that is, $y \neq 0$

$$(y, y_j) = 0, \quad j = 1, \dots, n; \quad (28)$$

in particular, the maximum in (27) is attained, and $y = y_{n+1}$ is a maximum point, where y_{n+1} is any eigenfunction corresponding to the characteristic value λ_{n+1} which is orthogonal to y_1, \dots, y_n .

Remark 3. For a positive definite kernel $K(x, t)$, the symbol of modulus on the right-hand sides of (27) and (28) can be omitted.

13.6-9. Kellogg's Method for Finding Characteristic Values in the Case of Symmetric Kernel.

Let $K(x, t)$ be a symmetric positive kernel ($a \leq x, t \leq b$). For an arbitrary function $\varphi_0(x) \in L_2(a, b)$, let us construct a sequence of functions by the recurrent formula

$$\varphi_n(x) = \int_a^b K(x, t)\varphi_{n-1}(t) dt, \quad n = 1, 2, \dots$$

and consider the numerical sequence

$$\left\{ \frac{\|\varphi_{n-1}\|}{\|\varphi_n\|} \right\}, \tag{29}$$

where $\|\varphi_n\| = \sqrt{\int_a^b |\varphi_n(x)|^2 dx}$. Let $y_1(x), y_2(x), \dots$ be orthonormal eigenfunctions of the kernel $K(x, t)$, and $\lambda_1 \leq \lambda_2 \leq \dots$ the corresponding characteristic values. Suppose that the initial function $\varphi_0(x)$ has been chosen orthogonal to the functions $y_1(x), \dots, y_{k-1}(x)$, but nonorthogonal to the eigenfunction $y_k(x)$. Then the limit of the sequence (29) is equal to the k th characteristic value λ_k . The sequence

$$\left\{ \frac{1}{\sqrt[n]{\|\varphi_n\|}} \right\} \tag{30}$$

has the same limit as (29). In this case, the sequence of functions

$$\left\{ \frac{\varphi_n(x)}{\|\varphi_n\|} \right\}$$

converges to a function which is a linear combination of eigenfunctions corresponding to the characteristic value λ_k .

Suppose that the functions $y_1(x)$ and $\varphi_0(x)$ are nonorthogonal, $\int_a^b y_1(x)\varphi_0(x) dx \neq 0$. Then, from (29) and (30) we obtain the following two approximation formulas for the smallest characteristic value:

$$\lambda_1 \approx \|\varphi_{n-1}\|/\|\varphi_n\|, \tag{31}$$

$$\lambda_1 \approx (\|\varphi_n\|)^{-1/n}. \tag{32}$$

Formula (31) yields an upper bound for λ_1 . For a suitably chosen initial function $\varphi_0(x)$, the Kellogg method is relatively simple with regard to calculations.

Example 1. Let us apply the Kellogg method for the calculation of the smallest characteristic value of the kernel $K(x, t) = x^2t^2, 0 \leq x, t \leq 1$.

Taking $\varphi_0(x) = x$ as the initial function, we find that

$$\begin{aligned} \varphi_1(x) &= \int_0^1 x^2t^2t dt = \int_0^1 x^2t^3 dt = \frac{1}{4}x^2, \\ \varphi_2(x) &= \int_0^1 x^2t^2 \frac{1}{4}t^2 dt = \frac{1}{4} \int_0^1 x^2t^4 dt = \frac{1}{4 \times 5}x^2, \\ \varphi_3(x) &= \int_0^1 x^2t^2 \frac{1}{4 \times 5}t^2 dt = \frac{1}{4 \times 5} \int_0^1 x^2t^4 dt = \frac{1}{4 \times 5^2}x^2, \\ &\dots\dots\dots \\ \varphi_n(x) &= \int_0^1 x^2t^2 \frac{1}{4 \times 5^{n-2}}t^2 dt = \frac{1}{4 \times 5^{n-2}} \int_0^1 x^2t^4 dt = \frac{1}{4 \times 5^{n-1}}x^2. \end{aligned}$$

Now, we define the norm

$$\|\varphi_n\| = \frac{1}{4 \times 5^{n-1}} \sqrt{\int_0^1 |x^4| dx} = \frac{1}{4 \times 5^{n-1} \sqrt{5}}.$$

According to (31), we find the first characteristic value:

$$\lambda_1 \approx \frac{\|\varphi_{n-1}\|}{\|\varphi_n\|} = 5.$$

It is easy to check that $\lambda_1 = 5$ is the exact characteristic value.

If the kernel $K(x, t)$ is not positive definite, then formulas (31) and (32) yield approximations for the smallest absolute value of the corresponding characteristic values.

13.6-10. Trace Method for the Approximation of Characteristic Values.

The m -trace of the kernel $K(x, t)$ is defined by

$$A_m = \int_a^b K_m(t, t) dt,$$

where $K_m(x, t)$ is the m th iterated kernel (see Subsection 13.3-1).

Formula (16) for a symmetric kernel implies that

$$A_m = \sum_{n=1}^{\infty} \frac{1}{\lambda_n^m} \quad (m = 2, 3, \dots).$$

For sufficiently large m , the leading term in this expression is $1/\lambda_1^m$, and therefore, we obtain the approximate relations

$$A_{2m} \approx \frac{1}{\lambda_1^{2m}}, \quad A_{2m+2} \approx \frac{1}{\lambda_1^{2m+2}}.$$

It follows that for the smallest characteristic value λ_1 , for large enough m , the following approximation formula holds:

$$|\lambda_1| \approx \sqrt{\frac{A_{2m}}{A_{2m+2}}}, \quad (33)$$

which is an upper bound for $|\lambda_1|$.

In order to calculate the second characteristic value, one can use the approximation formulas

$$|\lambda_2| \approx \frac{1}{|\lambda_1|} \sqrt{\frac{B_{2m}}{B_{2m+2}}}, \quad |\lambda_2| \approx \frac{1}{|\lambda_1|} \left(\frac{2}{B_{2m}} \right)^{1/(2m)},$$

where $B_{2m} = A_{2m}^2 - A_{4m}$.

Traces of even orders for a symmetric kernel are calculated by the formula

$$A_{2m} = \int_a^b \int_a^b K_m^2(x, t) dx dt = 2 \int_a^b \int_a^x K_m^2(x, t) dt dx. \quad (34)$$

Example 2. Let us use the trace method to find the first characteristic value of the kernel

$$K(x, t) = \begin{cases} x & \text{if } 0 \leq x \leq t \leq 1, \\ t & \text{if } 0 \leq t \leq x \leq 1. \end{cases}$$

Since $K(x, t)$ is symmetric, it suffices to find $K_2(x, t)$ for $t < x$. We have

$$K_2(x, t) = \int_0^1 K(x, z)K(z, t) dz = \int_0^t z^2 dz + \int_t^x zt dz + \int_x^1 xt dt = xt - \frac{1}{2}x^2t - \frac{1}{6}t^3.$$

Now, using (34) for $m = 1$ and $m = 2$, we find that

$$\begin{aligned} A_2 &= 2 \int_0^1 dx \int_0^x K_1^2(x, t) dt = 2 \int_0^1 dx \int_0^x t^2 dt = 2 \int_0^1 \frac{x^3}{3} dx = \frac{1}{6}, \\ A_4 &= 2 \int_0^1 dx \int_0^x K_2^2(x, t) dt \\ &= 2 \int_0^1 dx \int_0^x \left(x^2 t^2 + \frac{x^4 t^2}{4} + \frac{t^6}{36} - x^3 t^2 - \frac{x t^4}{3} + \frac{x^2 t^4}{6} \right) dt \\ &= 2 \int_0^1 \left(\frac{x^2 t^3}{3} + \frac{x^4 t^3}{12} + \frac{t^7}{7 \times 36} - \frac{x^3 t^3}{3} - \frac{x t^5}{15} + \frac{x^2 t^5}{30} \right) \Big|_{t=0}^{t=x} dx \\ &= 2 \int_0^1 \left(\frac{x^5}{3} + \frac{x^7}{12} + \frac{x^7}{7 \times 36} - \frac{x^6}{3} - \frac{x^6}{15} + \frac{x^7}{30} \right) dx = \frac{17}{630}. \end{aligned}$$

By (33) we obtain an approximation for the smallest characteristic value,

$$\lambda_1 \approx \sqrt{\frac{A_2}{A_4}} = \sqrt{\frac{\frac{1}{6}}{\frac{17}{630}}} \approx 2.485.$$

This is in good agreement with the exact value $\lambda_1 = \frac{1}{4}\pi^2 \approx 2.467$ (the error is less than 1%).

13.6-11. Integral Equations Reducible to Symmetric Equations.

An equation of the form

$$y(x) - \lambda \int_a^b K(x, t) \rho(t) y(t) dt = f(x), \quad (35)$$

where $K(x, t)$ is a symmetric kernel and $\rho(t) > 0$ is a continuous function on $[a, b]$, can be reduced to a symmetric equation. Indeed, on multiplying Eq. (35) by $\sqrt{\rho(x)}$ and introducing the new unknown function $z(x) = \sqrt{\rho(x)} y(x)$, we arrive at the integral equation

$$z(x) - \lambda \int_a^b L(x, t) z(t) dt = f(x) \sqrt{\rho(x)}, \quad L(x, t) = K(x, t) \sqrt{\rho(x) \rho(t)}, \quad (36)$$

where $L(x, t)$ is a symmetric kernel.

13.6-12. Skew-Symmetric Integral Equations.

By a *skew-symmetric integral equation* we mean an equation whose kernel is skew-symmetric, i.e., an equation of the form

$$y(x) - \lambda \int_a^b K(x, t) y(t) dt = f(x) \quad (37)$$

whose kernel $K(x, t)$ has the property

$$K(t, x) = -K(x, t). \quad (38)$$

Equation (37) with the skew-symmetric kernel (38) has at least one characteristic value, and all its characteristic values are purely imaginary.

13.6-13. Remark on Nonsymmetric Kernels.

An integral equation with a *nonsymmetric kernel* (i.e., such that $K(x, t) \neq K(t, x)$ for some x, t) may happen to have no characteristic values.

Example. Consider the homogeneous integral equation with nonsymmetric degenerate kernel

$$y(x) = \lambda \int_0^\pi \cos x \sin t y(t) dt, \quad (1)$$

which can be written in the form

$$y(x) = A \cos x, \quad A = \lambda \int_0^\pi \sin t y(t) dt. \quad (2)$$

Substituting (2) into (1) and dividing the result by $\cos x$, we get

$$A = \lambda \int_0^\pi \sin t A \cos t dt = 0.$$

Therefore, equation (1) has only the trivial solution for any λ .

⊙ References for Section 13.6: E. Goursat (1923), G. Wiarda (1930), R. Courant and D. Hilbert (1931), S. G. Mikhlin (1960), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), J. A. Cochran (1972), V. I. Smirnov (1974), P. P. Zabreyko, A. I. Koshelev, et al. (1975), A. J. Jerry (1985), F. G. Tricomi (1985), D. Porter and D. S. G. Stirling (1990), C. Corduneanu (1991), J. Kondo (1991), W. Hackbusch (1995), R. P. Kanwal (1996).

13.7. Integral Equations with Nonnegative Kernels

13.7-1. Positive Principal Eigenvalues. Generalized Jentzch Theorem.

In this section, we consider nonnegative kernels $K(x, t) \geq 0$ that are either continuous or square-summable in the domain $a \leq x, t \leq b$.

Such kernels, under minimal additional assumptions, admit nonnegative eigenfunctions. The associated eigenvalues μ (resp., characteristic values $\lambda = 1/\mu$) will be called *positive principal eigenvalues* (resp., *positive principal characteristic values*). Each positive principal eigenvalue is obviously nonzero. Under fairly general conditions, a nonnegative eigenfunction is unique (up to a constant coefficient) and the corresponding positive principal eigenvalue is an upper bound for the modulus of any other eigenvalue.

THEOREM 1 (GENERALIZED JENTZCH THEOREM). *If a continuous or polar kernel $K(x, t)$ is positive, then its characteristic values λ_0 with the smallest modulus is positive and simple, and the corresponding eigenfunction $y_0(x)$ does not change sign on the interval $a \leq x \leq b$.*

Remark 1. The generalized Jentzch theorem holds for a symmetric, as well as a nonsymmetric, polar positive kernel. It is allowed that the kernel may vanish at isolated points (on a set of zero measure) of the domain $a \leq x, t \leq b$.

THEOREM 2. *Suppose a nonnegative kernel $K(x, t)$ has at least one (real or complex) eigenvalue. Then it has a nonnegative eigenvalue μ_0 .*

Remark 2. Not every nonnegative kernel has a nonnegative eigenfunction.

Example. Any nonnegative kernel $K(x, t) \geq 0$ ($a \leq x, t \leq b$) satisfying the condition $K(x, t) \equiv 0$ for $t \geq x$ has no eigenfunctions corresponding to nonzero eigenvalues.

THEOREM 3. *Let $K(x, t)$ be a nonnegative kernel. Suppose that there is a function $u_0(x)$ which is positive on a set of nonzero measure and satisfies the inequality*

$$\int_a^b K_n(x, t)u_0(t) dt \geq \beta u_0(x) \quad (a \leq x \leq b),$$

where $\beta > 0$ and $K_n(x, t)$ is an iterated kernel of some order n . Then the kernel $K(x, t)$ has at least one positive principal eigenvalue μ_0 . This eigenvalue satisfies the inequality $\mu_0 \geq \beta^{1/n}$.

THEOREM 4. *All (real and complex) eigenvalues μ of the nonnegative kernel $K(x, t)$ satisfy the inequality $|\mu| < \Delta$, where Δ is the largest positive principal eigenvalue.*

13.7-2. Positive Solutions of a Nonhomogeneous Integral Equation.

Consider a nonhomogeneous integral equation with a parameter μ :

$$\mu y(x) = \int_a^b K(x, t)y(t) dt + f(x) \quad (a \leq x \leq b), \quad (1)$$

where the kernel $K(x, t) \geq 0$ is either continuous or square-summable. The functions $y(t)$ and $f(x)$ are also assumed either continuous or square-summable.

THEOREM 1. *Let $\mu > \Delta$, where Δ is the largest positive principal eigenvalue of the kernel $K(x, t)$. Then, for any nonnegative function $f(x)$, equation (1) has one and only one nonnegative solution $y(x)$, which can be obtained by the method of successive approximations based on the formula*

$$\mu y_{n+1}(x) = \int_a^b K(x, t)y_n(t) dt + f(x) \quad (n = 0, 1, \dots) \quad (2)$$

with any initial approximation $y_0(x)$.

For $y_0(x) = 0$, the solution can be represented as the series

$$y(x) = \sum_{n=0}^{\infty} \frac{\mathbf{K}^{(n)}[f(x)]}{\mu^{n+1}}, \quad \mathbf{K}[f(x)] = \int_a^b K(x, t)f(t) dt, \quad \mathbf{K}^{(n)}[f(x)] = \mathbf{K}[\mathbf{K}^{(n-1)}[f(x)]].$$

Under the assumptions of Theorem 1, the rate of convergence of the successive approximations to the solution of equation (1) is characterized by the inequality

$$\|y - y_n\| \leq C(\mu) \left(\frac{\Delta}{\mu} \right)^n \quad (n = 1, 2, \dots),$$

where $C(\mu)$ is a constant. If the kernel $K(x, t)$ and the function $f(x)$ are continuous, then the norm is introduced by $\|y\| = \max_{a \leq x \leq b} |y(x)|$. If $K(x, t)$ and $f(x)$ are square-summable, then one takes the

$$\text{norm } \|y\| = \left[\int_a^b y^2(x) dx \right]^{1/2}.$$

THEOREM 2. *If equation (1) admits a positive solution for at least one positive function $f_0(x)$, then $\mu > \Delta$, and therefore, equation (1) has a nonnegative solution for any nonnegative function $f(x)$.*

13.7-3. Estimates for the Spectral Radius.

1°. The greatest among the moduli of the eigenvalues of the kernel $K(x, t)$ is called the *spectral radius of the kernel* or *spectral radius of the integral operator*

$$\mathbf{K}[y(x)] = \int_a^b K(x, t)y(t) dt$$

and is denoted $\rho(\mathbf{K})$.

The role of the spectral radius can be characterized, for instance, by the fact that the integral equation (1) with a continuous kernel and a continuous free term has a continuous solution that can be obtained by the method of successive approximations (2) if and only if $|\mu| > \rho(\mathbf{K})$. Theorem 4 of Subsection 13.7-1 implies that the spectral radius of the nonnegative kernel $K(x, t) \geq 0$ is either equal to zero or coincides with its largest positive principal eigenvalue. Therefore, estimates for the

spectral radius in the case of a nonnegative kernel coincide with estimates for the largest positive principal eigenvalue.

Estimates for spectral radii of nonnegative kernels can be used for studying kernels of alternating sign, since the spectral radius of a nonnegative kernel $K(x, t)$ is an upper bound for the spectral radius of any kernel $M(x, t)$ such that

$$|M(x, t)| \leq K(x, t) \quad (a \leq x \leq b).$$

In the following statements, it is assumed that all functions are either continuous or square-summable and $K(x, t) \geq 0$.

2°. The simplest upper bounds for the spectral radius have the form

$$\begin{aligned} \rho(\mathbf{K}) &\leq \max_{a \leq x \leq b} \int_a^b K(x, t) dt, \\ \rho(\mathbf{K}) &\leq \left[\int_a^b \int_a^b K^2(x, t) dx dt \right]^{1/2}. \end{aligned}$$

More precise estimates are obtained in terms of iterations of the kernel:

$$\begin{aligned} \rho(\mathbf{K}) &\leq \left[\max_{a \leq x \leq b} \int_a^b K_n(x, t) dt \right]^{1/n}, \\ \rho(\mathbf{K}) &\leq \left[\int_a^b \int_a^b K_n^2(x, t) dx dt \right]^{1/(2n)}. \end{aligned}$$

Let us give two more estimates.

Suppose that for some $\beta_1 > 0$, the following inequality holds:

$$\int_a^b K(x, t) \left[\beta_1 - \int_a^b K_n(t, \tau) d\tau \right] dt \geq 0.$$

Then $\rho(\mathbf{K}) \leq \beta_1^{1/n}$.

Suppose that for some $\beta_2 > 0$, we have

$$\int_a^b K_n(x, t) \left[\beta_2 - \int_a^b K(t, \tau) d\tau \right] dt \geq 0.$$

Then $\rho(\mathbf{K}) \leq \beta_2$.

THEOREM 1. Suppose that for some $\beta_1 > 0$ and some nonnegative function $u_1(x)$ taking positive values on a set of nonzero measure, the following inequality holds:

$$\int_a^b K(x, t)u_1(t) dt \geq \beta_1 u_1(x) \quad (a \leq x \leq b).$$

Then $\rho(\mathbf{K}) \geq \beta_1$.

THEOREM 2. Suppose that for some $\beta_2 > 0$ and some nonnegative function $u_2(x)$ taking zero values only on a set of zero measure (say, at finitely many points), the following inequality holds:

$$\int_a^b K_n(x, t)u_2(t) dt \leq \beta_2 u_2(x) \quad (a \leq x \leq b).$$

Then $\rho(\mathbf{K}) \leq \beta_2^{1/n}$.

3°. Consider a continuous kernel $K(x, t)$ defined on the square $a \leq x, t \leq b$. Let us split the segment $[a, b]$ into n parts:

$$a = x_0 < x_1 < \dots < x_{n-1} < x_n = b.$$

Setting

$$m_{ij} = \max_{x_{i-1} \leq x \leq x_i} \int_{x_{j-1}}^{x_j} |K(x, t)| dt \quad (i, j = 1, \dots, n), \quad (3)$$

let us construct the matrix

$$S \equiv \begin{pmatrix} m_{11} & m_{12} & \cdots & m_{1n} \\ m_{21} & m_{22} & \cdots & m_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ m_{n1} & m_{n2} & \cdots & m_{nn} \end{pmatrix}. \quad (4)$$

THEOREM 3. *The spectral radius $\rho(\mathbf{K})$ does not exceed the largest eigenvalue of the matrix S .*

The likewise is true if, instead of (3), the elements of the matrix (4) are defined by

$$m_{ij} = \left(\int_{x_{i-1}}^{x_i} \int_{x_{j-1}}^{x_j} K^2(x, t) dx dt \right)^{1/2}. \quad (5)$$

Example. Consider the kernel $K(x, t)$ that coincides with the Green function $G(x, t)$ for the equation of vibrations of a string with fixed ends,

$$K(x, t) = G(x, t) = \begin{cases} x(1-t) & \text{if } 0 \leq x \leq t \leq 1, \\ t(1-x) & \text{if } 0 \leq t \leq x \leq 1. \end{cases}$$

Let us construct the matrix (4), taking $n = 5$, $x_i = \frac{1}{5}i$ ($i = 0, 1, \dots, 5$). The elements of this matrix are calculated as in (5).

The largest eigenvalue of the matrix S in this case is equal to 0.10216. This gives the estimate $\rho(\mathbf{K}) \leq 0.10216$. The exact largest eigenvalue is $1/\pi^2 \approx 0.10132$.

13.7-4. Basic Definition and Theorems for Oscillating Kernels.

1°. A continuous function of two variables $K(x, t)$ ($a \leq x, t \leq b$) is called an *oscillation kernel*, if the following inequalities hold:

- (a) $K(x, t) > 0$, $a < x < b$, $a < t < b$;
- (b) $\det K(x_i, t_j) \geq 0$, $a < x_1 < x_2 < \dots < x_n < b$, $a < t_1 < t_2 < \dots < t_n < b$;
- (c) $\det K(x_i, x_j) > 0$, $a < x_1 < x_2 < \dots < x_n < b$,

where n is an arbitrary positive integer and the points x_i, t_j that satisfy the above inequalities are otherwise selected arbitrarily.

It can be shown that the product of two (or finitely many) oscillation kernels is an oscillating kernel.

THEOREM 1. *Consider an integral equation of the form*

$$y(x) = \lambda \int_a^b K(x, t)\sigma(t)y(t) dt, \quad (6)$$

where $K(x, t)$ is an oscillation kernel and $\sigma(t) > 0$ is a continuous function. Then the following statements hold:

1. All characteristic values (6) are positive and simple; $0 < \lambda_0 < \lambda_1 < \dots$.
2. The eigenfunction $y_0(x)$ corresponding to λ_0 has no zeros on the interval $a < x < b$.

3. The eigenfunction $y_k(x)$ corresponding to λ_k has precisely k nodes ($y_k(x)$ changes sign at each node) and has no other zeros.

4. For arbitrary integers k and m ($0 \leq k \leq m$) and real c_k, c_{k+1}, \dots, c_m ($\sum_{i=k}^m c_i^2 > 0$), the linear combination

$$y(x) = \sum_{i=k}^m c_i y_i(x)$$

has at least k nodes and at most m zeros.

5. The nodes of neighboring eigenfunctions alternate.

In order to apply the theory of integral equations with oscillation kernels to the investigation of ordinary differential equations, one has to reduce the latter to the former with the help of the Green's function, and it is necessary to answer the question whether the Green's function represents an oscillation kernel. Next, we give some results that suggest an answer to this question.

2°. Consider a differential operator

$$L[y] = \sum_{s=0}^n \gamma_s(x) \frac{d^s y}{dx^s}, \quad n \geq 2, \tag{7}$$

on the interval $a \leq x \leq b$ with positive coefficients

$$\gamma_s(x) > 0$$

and the homogeneous boundary conditions

$$\begin{aligned} \sum_{m=0}^{n-1} \alpha_{im} y_x^{(m)} &= 0 \quad \text{for } x = a \quad (i = 1, \dots, p), \\ \sum_{m=0}^{n-1} \beta_{im} y_x^{(m)} &= 0 \quad \text{for } x = b \quad (i = 1, \dots, q), \end{aligned} \tag{8}$$

where $p + q = 1$.

THEOREM 2. Suppose that the system of boundary conditions (8) corresponds to the Green's function $G(x, t)$ of the differential operator (7) such that $(-1)^q G(x, t)$ is an oscillation kernel. Then the same property holds for the following simpler system of boundary conditions:

$$\begin{aligned} y(a) = y'_x(a) = \dots = y_x^{(p-1)}(a) &= 0, \\ y(b) = y'_x(b) = \dots = y_x^{(q-1)}(b) &= 0, \end{aligned} \tag{9}$$

where $p + q = n$, i.e., $(-1)^q G_{p,q}(x, t)$ is an oscillation kernel, where $G_{p,q}(x, t)$ is the Green's function of the operator (7) with the boundary conditions (9).

THEOREM 3. The system of boundary conditions (9) ($1 \leq p < n$) of the operator (7) corresponds to an oscillation kernel $(-1)^q G_{p,q}(x, t)$ if and only if the following two conditions hold:

1. The differential equation with the truncated system of boundary conditions

$$L[y] = 0; \quad y(b) = y'_x(b) = \dots = y_x^{(q-1)}(b) = 0$$

has p solutions $y_1 = y_1(x), \dots, y_p = y_p(x)$ such that

$$y_1 > 0, \quad W(y_1, y_2) > 0, \quad \dots, \quad W(y_1, \dots, y_p) > 0 \quad \text{for } a < x < b,$$

where $W(y_1, \dots, y_k)$ is the Wronskian determinant

$$W(y_1, \dots, y_k) = \begin{vmatrix} y_1(x) & \cdots & y_k(x) \\ y_1'(x) & \cdots & y_k'(x) \\ \cdots & \cdots & \cdots \\ y_1^{(k-1)}(x) & \cdots & y_k^{(k-1)}(x) \end{vmatrix}.$$

2. The differential equation with the truncated system of boundary conditions

$$L[y] = 0; \quad y(a) = y_x'(a) = \cdots = y_x^{(p-1)}(a) = 0$$

has $q = n - p$ solutions $y_{p+1} = y_1(x), \dots, y_n = y_n(x)$ such that

$$W(y_1, \dots, y_p, y_{p+1}) > 0, \quad \dots, \quad W(y_1, \dots, y_p, \dots, y_n) > 0 \quad \text{for } a < x < b.$$

THEOREM 4. *Conditions 1 and 2 of Theorem 3, under the assumption that the Green's function exists, are equivalent to the condition that the differential operator $L[y]$ admits a representation of the form*

$$L[y] = \mu_0(x) \frac{d}{dx} \mu_1(x) \frac{d}{dx} \mu_2(x) \cdots \frac{d}{dx} \mu_n(x) y, \quad (10)$$

where $\mu_k(x)$ are positive weight functions with k continuous derivatives on (a, b) .

If the differential operator $L[y]$ admits the representation (10), then the equation $L[y] = 0$ has a particular solution $y = \text{const} / \mu_n(x)$.

THEOREM 5 (KREIN'S CRITERION). *The condition that for each p ($1 \leq p < n$), the differential operator (7) with the boundary conditions (9) admits a Green's function $G_{p,q}(x, t)$ such that $(-1)^q G_{p,q}(x, t)$ is an oscillation kernel, is equivalent to the condition that the operator $L[y]$ on the interval (a, b) admits the representation (10) with strictly positive functions $\mu_k(x)$ having k continuous derivatives on (a, b) .*

Remark. Suppose that the operator (10) with the boundary conditions (8) admits a Green's function $G(x, t)$ (it is assumed that $\mu_k(x) > 0$ and have k continuous derivatives). Then the function $(-1)^q G(x, t)$ is an oscillation kernel.

Example 1. Consider the second-order linear differential operator

$$L[y] = f(x)y''_{xx} + g(x)y'_x, \quad (11)$$

where $f(x) > 0$ and $g(x) > 0$ for $x \in [a, b]$, with the homogeneous boundary conditions of the first kind

$$y(a) = 0, \quad y(b) = 0. \quad (12)$$

The differential operator (11) can be represented as an iterated operator (10) with positive weights:

$$L[y] = \mu_0(x) \frac{d}{dx} \mu_1(x) \frac{d}{dx} \mu_2(x) y, \\ \mu_0(x) = f(x) \exp \left[\int \frac{g(x)}{f(x)} dx \right], \quad \mu_1(x) = \exp \left[\int \frac{g(x)}{f(x)} dx \right], \quad \mu_2(x) = 1.$$

Boundary conditions (12) represent a special case of (9) for $p = q = 1$. It is not difficult to show that the operator (11) with the conditions (12) has the following Green's function [constructed with the help of formulas (12) from Subsection 18.3-3]:

$$G(x, t) = - \begin{cases} \frac{Y(a, x)Y(t, b)}{f(t)\Phi(t)Y(a, b)} & \text{if } a \leq x \leq t, \\ \frac{Y(a, t)Y(x, b)}{f(t)\Phi(t)Y(a, b)} & \text{if } t \leq x \leq b, \end{cases}$$

where

$$\Phi(t) = \exp\left[-\int_0^t \frac{g(\tau)}{f(\tau)} d\tau\right], \quad Y(a, b) = \int_a^b \Phi(z) dz.$$

Krein's criterion ensures that the function $K(x, t) = -G(x, t)$ is an oscillation kernel.

Example 2. Consider the third-order differential operator

$$L[y] = y_{xxx} \tag{13}$$

with the boundary conditions

$$\begin{aligned} y'_x &= 0 & \text{for } x &= 0, \\ y &= y''_{xx} = 0 & \text{for } x &= 1. \end{aligned} \tag{14}$$

It is easy to check that the operator (13) with the boundary conditions (14) has the Green's function

$$G(x, t) = \begin{cases} t - \frac{1}{2}(x^2 + t^2) & \text{for } 0 \leq x \leq t \leq 1, \\ t - xt & \text{for } 0 \leq t \leq x \leq 1. \end{cases} \tag{15}$$

Here $p = 1$ and $q = 2$. Therefore, it follows from Theorem 5 (see the remark) that $G(x, t)$ is an oscillation kernel.

Now, let us examine the eigenvalue problem for the third-order equation

$$y_{xxx} = \lambda \sigma(x)y, \quad \sigma(x) > 0,$$

with boundary conditions (14). With the help of the Green's function (15), this problem is reduced to the integral equation

$$y(x) = \lambda \int_0^1 G(x, t)\sigma(t)y(t) dt. \tag{16}$$

Since $G(x, t)$ is an oscillation kernel and $\sigma(x) > 0$, the results of Theorem 1 can be applied to equation (16).

13.7-5. Stochastic Kernels.

1°. A nonnegative continuous kernel $K(x, t)$ in the domain $a \leq x, t \leq b$ is called a *stochastic kernel*, if

$$\int_a^b K(x, t) dt \equiv 1 \quad (a \leq x \leq b).$$

Obviously, for any integral operator with a stochastic kernel $K(x, t)$,

$$y_0(x) \equiv 1 \quad (a \leq x \leq b)$$

is an eigenfunction corresponding to the characteristic value $\lambda_0 = 1$. The other characteristic values λ satisfy the inequality $|\lambda| \geq 1$. Integral operators with stochastic kernels may have characteristic values $\lambda \neq 1$ such that $|\lambda| = 1$. The corresponding eigenvalues $\mu = 1/\lambda$ are called *permutators*.

2°. Properties of stochastic kernels:

1. All eigenvalues μ of an integral operator with stochastic kernel such that $|\mu| = 1$ are integer roots of unity.

2. The set of all eigenfunctions of an integral operator with stochastic kernel corresponding to an eigenvalue $\mu = 1/\lambda = 1$ contains a basis that consists of nonnegative functions $y_1(x), \dots, y_m(x)$ with the following properties:

(a) for every $y_j(x)$ ($j = 1, \dots, m$), there is at least one point at which this function is positive and all other functions of the basis are equal to zero;

(b) for each $x \in [a, b]$, there is at least one function of the basis that is positive at x .

© References for Section 13.7: M. G. Krein (1939), F. P. Gantmakher and M. G. Krein (1950), S. Karlin (1968), J. M. Karon (1969), P. P. Zabreyko, A. I. Koshelev, et al. (1975), D. D. Joseph (1976), R. P. Agarwal, D. O'Regan, and P. J. Y. Wong (1998).

13.8. Operator Method for Solving Integral Equations of the Second Kind

13.8-1. Simplest Scheme.

Consider a linear equation of the second kind of the special form

$$y(x) - \lambda \mathbf{L}[y] = f(x), \quad (1)$$

where \mathbf{L} is a linear (integral) operator such that $\mathbf{L}^2 = k$, $k = \text{const}$.

Let us apply the operator \mathbf{L} to Eq. (1). We obtain

$$\mathbf{L}[y] - k\lambda y(x) = \mathbf{L}[f(x)]. \quad (2)$$

On eliminating the term $\mathbf{L}[y]$ from (1) and (2), we find the solution

$$y(x) = \frac{1}{1 - k\lambda^2} \{f(x) + \lambda \mathbf{L}[f]\}. \quad (3)$$

Remark. In Section 11.4, various generalizations of the above method are described.

13.8-2. Solution of Equations of the Second Kind on the Semiaxis.

1°. Consider the equation

$$y(x) - \lambda \int_0^\infty \cos(xt)y(t) dt = f(x). \quad (4)$$

In this case, the operator \mathbf{L} coincides, up to a constant factor, with the Fourier cosine transform:

$$\mathbf{L}[y] = \int_0^\infty \cos(xt)y(t) dt = \sqrt{\frac{\pi}{2}} \mathfrak{F}_c[y] \quad (5)$$

and acts by the rule $\mathbf{L}^2 = k$, where $k = \frac{\pi}{2}$ (see Subsection 9.5-1).

We obtain the solution by formula (3) taking into account Eq. (5):

$$y(x) = \frac{2}{2 - \pi\lambda^2} \left[f(x) + \lambda \int_0^\infty \cos(xt)f(t) dt \right], \quad \lambda \neq \pm \sqrt{\frac{2}{\pi}}. \quad (6)$$

2°. Consider the equation

$$y(x) - \lambda \int_0^\infty t J_\nu(xt)y(t) dt = f(x), \quad (7)$$

where $J_\nu(x)$ is the Bessel function, $\text{Re } \nu > -1$.

Here the operator \mathbf{L} coincides, up to a constant factor, with the Hankel transform:

$$\mathbf{L}[y] = \int_0^\infty t J_\nu(xt)y(t) dt \quad (8)$$

and acts by the rule $\mathbf{L}^2 = 1$ (see Subsection 9.6-1).

We obtain the solution by formula (3), for $k = 1$, taking into account Eq. (8):

$$y(x) = \frac{1}{1 - \lambda^2} \left[f(x) + \lambda \int_0^\infty t J_\nu(xt)f(t) dt \right], \quad \lambda \neq \pm 1. \quad (9)$$

13.9. Methods of Integral Transforms and Model Solutions

13.9-1. Equation with Difference Kernel on the Entire Axis.

Consider an integral equation of convolution type of the second kind with one kernel

$$y(x) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K(x-t)y(t) dt = f(x), \quad -\infty < x < \infty, \quad (1)$$

where $f(x)$ and $K(x)$ are the known right-hand side and the kernel of the integral equation and $y(x)$ is the unknown function. Let us apply the (alternative) Fourier transform to Eq. (1). In this case, taking into account the convolution theorem (see Subsection 9.4-4), we obtain

$$\mathcal{Y}(u)[1 + \mathcal{K}(u)] = \mathcal{F}(u). \quad (2)$$

Thus, on applying the Fourier transform we reduce the solution of the original integral equation (1) to the solution of the algebraic equation (2) for the transform of the unknown function. The solution of Eq. (2) has the form

$$\mathcal{Y}(u) = \frac{\mathcal{F}(u)}{1 + \mathcal{K}(u)}. \quad (3)$$

Formula (3) gives the transform of the solution of the original integral equation in terms of the transforms of the known functions, namely, the kernel and the right-hand side of the equation. The solution itself can be obtained by applying the Fourier inversion formula:

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{Y}(u)e^{-iux} du = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\mathcal{F}(u)}{1 + \mathcal{K}(u)} e^{-iux} du. \quad (4)$$

In fact, formula (4) solves the problem; however, sometimes it is not convenient because it requires the calculation of the transform $F(u)$ for each right-hand side $f(x)$. In many cases, the representation of the solution of the nonhomogeneous integral equation via the resolvent of the original equation is more convenient. To obtain the desired representation, we note that formula (3) can be transformed to the expression

$$\mathcal{Y}(u) = [1 - \mathcal{R}(u)]\mathcal{F}(u), \quad \mathcal{R}(u) = \frac{\mathcal{K}(u)}{1 + \mathcal{K}(u)}. \quad (5)$$

On the basis of (5), by applying the Fourier inversion formula and the convolution theorem (for transforms) we obtain

$$y(x) = f(x) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} R(x-t)f(t) dt, \quad (6)$$

where the resolvent $R(x-t)$ of the integral equation (1) is given by the relation

$$R(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\mathcal{K}(u)}{1 + \mathcal{K}(u)} e^{-iux} du, \quad (7)$$

Thus, to determine the solution of the original integral equation (1), it suffices to find the function $R(x)$ by formula (7).

The function $R(x)$ is a solution of Eq. (1) for a special form of the function $f(x)$. Indeed, it follows from formulas (3) and (5) that for $\mathcal{Y}(u) = \mathcal{R}(u)$ the function $\mathcal{F}(u)$ is equal to $\mathcal{K}(u)$. This means that, for $f(x) \equiv K(x)$, the function $y(x) \equiv R(x)$ is a solution of Eq. (1), i.e., the resolvent of Eq. (1) satisfies the integral equation

$$R(x) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K(x-t)R(t) dt = K(x), \quad -\infty < x < \infty. \quad (8)$$

Note that to calculate direct and inverse Fourier transforms, one can use the corresponding tables from Supplements 7 and 8 and the books by H. Bateman and A. Erdélyi (1954) and by V. A. Ditkin and A. P. Prudnikov (1965).

Example. Let us solve the integral equation

$$y(x) - \lambda \int_{-\infty}^{\infty} \exp(\alpha|x-t|)y(t) dt = f(x), \quad -\infty < x < \infty, \quad (9)$$

which is a special case of Eq. (1) with kernel $K(x-t)$ given by the expression

$$K(x) = -\sqrt{2\pi} \lambda e^{-\alpha|x|}, \quad \alpha > 0. \quad (10)$$

Let us find the function $R(x)$. To this end, we calculate the integral

$$\mathcal{K}(u) = - \int_{-\infty}^{\infty} \lambda e^{-\alpha|x|} e^{iux} dx = -\frac{2\alpha\lambda}{u^2 + \alpha^2}. \quad (11)$$

In this case, formula (5) implies

$$\mathcal{R}(u) = \frac{\mathcal{K}(u)}{1 + \mathcal{K}(u)} = -\frac{2\alpha\lambda}{u^2 + \alpha^2 - 2\alpha\lambda}, \quad (12)$$

and hence

$$R(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{R}(u) e^{-iux} du = -\sqrt{\frac{2}{\pi}} \int_{-\infty}^{\infty} \frac{\alpha\lambda}{u^2 + \alpha^2 - 2\alpha\lambda} e^{-iux} du. \quad (13)$$

Assume that $\lambda < \frac{1}{2}\alpha$. In this case the integral (13) makes sense and can be calculated by means of the theory of residues on applying the Jordan lemma (see Subsections 9.1-4 and 9.1-5). After some algebraic manipulations, we obtain

$$R(x) = -\sqrt{2\pi} \frac{\alpha\lambda}{\sqrt{\alpha^2 - 2\alpha\lambda}} \exp(-|x|\sqrt{\alpha^2 - 2\alpha\lambda}) \quad (14)$$

and finally, in accordance with (6), we obtain

$$y(x) = f(x) + \frac{\alpha\lambda}{\sqrt{\alpha^2 - 2\alpha\lambda}} \int_{-\infty}^{\infty} \exp(-|x-t|\sqrt{\alpha^2 - 2\alpha\lambda}) f(t) dt, \quad -\infty < x < \infty. \quad (15)$$

13.9-2. Equation with the Kernel $K(x, t) = t^{-1}Q(x/t)$ on the Semiaxis.

Here we consider the following equation on the semiaxis:

$$y(x) - \int_0^{\infty} \frac{1}{t} Q\left(\frac{x}{t}\right) y(t) dt = f(x). \quad (16)$$

To solve this equation we apply the Mellin transform which is defined as follows (see also Section 9.3):

$$\hat{f}(s) = \mathfrak{M}\{f(x), s\} \equiv \int_0^{\infty} f(x)x^{s-1} dx, \quad (17)$$

where $s = \sigma + i\tau$ is a complex variable ($\sigma_1 < \sigma < \sigma_2$) and $\hat{f}(s)$ is the transform of the function $f(x)$. In what follows, we briefly denote the Mellin transform by $\mathfrak{M}\{f(x)\} \equiv \mathfrak{M}\{f(x), s\}$.

For known $\hat{f}(s)$, the original function can be found by means of the Mellin inversion formula

$$f(x) = \mathfrak{M}^{-1}\{\hat{f}(s)\} \equiv \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \hat{f}(s)x^{-s} ds, \quad \sigma_1 < c < \sigma_2, \quad (18)$$

where the integration path is parallel to the imaginary axis of the complex plane s and the integral is understood in the sense of the Cauchy principal value.

On applying the Mellin transform to Eq. (16) and taking into account the fact that the integral with such a kernel is transformed into the product by the rule (see Subsection 9.3-2)

$$\mathfrak{M}\left\{\int_0^{\infty} \frac{1}{t} Q\left(\frac{x}{t}\right) y(t) dt\right\} = \hat{Q}(s)\hat{y}(s),$$

we obtain the following equation for the transform $\hat{y}(s)$:

$$\hat{y}(s) - \hat{Q}(s)\hat{y}(s) = \hat{f}(s).$$

The solution of this equation is given by the formula

$$\hat{y}(s) = \frac{\hat{f}(s)}{1 - \hat{Q}(s)}. \tag{19}$$

On applying the Mellin inversion formula to Eq. (19) we obtain the solution of the original integral equation

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\hat{f}(s)}{1 - \hat{Q}(s)} x^{-s} ds. \tag{20}$$

This solution can also be represented via the resolvent in the form

$$y(x) = f(x) + \int_0^\infty \frac{1}{t} N\left(\frac{x}{t}\right) f(t) dt, \tag{21}$$

where we have used the notation

$$N(x) = \mathfrak{M}^{-1}\{\hat{N}(s)\}, \quad \hat{N}(s) = \frac{\hat{Q}(s)}{1 - \hat{Q}(s)}. \tag{22}$$

Under the application of this analytical method of solution, the following technical difficulties can occur: (a) in the calculation of the transform for a given kernel $K(x)$ and (b) in the calculation of the solution for the known transform $\hat{y}(s)$. To find the corresponding integrals, tables of direct and inverse Mellin transforms are applied (e.g., see Supplements 9 and 10). In many cases, the relationship between the Mellin transform and the Fourier and Laplace transforms is first used:

$$\mathfrak{M}\{f(x), s\} = \mathfrak{F}\{f(e^x), is\} = \mathfrak{L}\{f(e^x), -s\} + \mathfrak{L}\{f(e^{-x}), s\}, \tag{23}$$

and then tables of direct and inverse Fourier transforms and Laplace transforms are applied (see Supplements 5–8).

Remark 1. The equation

$$y(x) - \int_0^\infty H\left(\frac{x}{t}\right) x^\alpha t^{-\alpha-1} y(t) dt = f(x) \tag{24}$$

can be rewritten in the form of Eq. (16) under the notation $K(z) = z^\alpha H(z)$.

13.9-3. Equation with the Kernel $K(x, t) = t^\beta Q(xt)$ on the Semiaxis.

Consider the following equation on the semiaxis:

$$y(x) - \int_0^\infty t^\beta Q(xt)y(t) dt = f(x). \tag{25}$$

To solve this equation, we apply the Mellin transform. On multiplying Eq. (25) by x^{s-1} and integrating with respect to x from zero to infinity, we obtain

$$\int_0^\infty y(x)x^{s-1} dx - \int_0^\infty y(t)t^\beta dt \int_0^\infty Q(xt)x^{s-1} dx = \int_0^\infty f(x)x^{s-1} dx. \tag{26}$$

Let us make the change of variables $z = xt$. We finally obtain

$$\hat{y}(s) - \hat{Q}(s) \int_0^{\infty} y(t)t^{\beta-s} dt = \hat{f}(s). \quad (27)$$

Taking into account the relation

$$\int_0^{\infty} y(t)t^{\beta-s} dt = \hat{y}(1 + \beta - s),$$

we rewrite Eq. (27) in the form

$$\hat{y}(s) - \hat{Q}(s)\hat{y}(1 + \beta - s) = \hat{f}(s). \quad (28)$$

On replacing s by $1 + \beta - s$ in Eq. (28), we obtain

$$\hat{y}(1 + \beta - s) - \hat{Q}(1 + \beta - s)\hat{y}(s) = \hat{f}(1 + \beta - s). \quad (29)$$

Let us eliminate $\hat{y}(1 + \beta - s)$ and solve the resulting equation for $\hat{y}(s)$. We thus find the transform of the solution:

$$\hat{y}(s) = \frac{\hat{f}(s) + \hat{Q}(s)\hat{f}(1 + \beta - s)}{1 - \hat{Q}(s)\hat{Q}(1 + \beta - s)}. \quad (30)$$

On applying the Mellin inversion formula, we obtain the solution of the integral equation (25) in the form

$$y(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\hat{f}(s) + \hat{Q}(s)\hat{f}(1 + \beta - s)}{1 - \hat{Q}(s)\hat{Q}(1 + \beta - s)} x^{-s} ds. \quad (31)$$

Remark 2. The equation

$$y(x) - \int_0^{\infty} H(xt)x^p t^q y(t) dt = f(x)$$

can be rewritten in the form of Eq. (25) under the notation $Q(z) = z^p H(z)$, where $\beta = q - p$.

13.9-4. Method of Model Solutions for Equations on the Entire Axis.

Let us illustrate the capability of a generalized modification of the method of model solutions (see Subsection 11.6) by an example of the equation

$$Ay(x) + \int_{-\infty}^{\infty} Q(x+t)e^{\beta t} y(t) dt = f(x), \quad (32)$$

where $Q = Q(z)$ and $f(x)$ are arbitrary functions and A and β are arbitrary constants satisfying some constraints.

For clarity, instead of the original equation (32) we write

$$\mathbf{L}[y(x)] = f(x). \quad (33)$$

For a test solution, we take the exponential function

$$y_0 = e^{px}. \quad (34)$$

On substituting (34) into the left-hand side of Eq. (33), after some algebraic manipulations we obtain

$$\mathbf{L}[e^{px}] = Ae^{px} + q(p)e^{-(p+\beta)x}, \quad \text{where } q(p) = \int_{-\infty}^{\infty} Q(z)e^{(p+\beta)z} dz. \quad (35)$$

The right-hand side of (35) can be regarded as a functional equation for the kernel e^{px} of the inverse Laplace transform. To solve it, we replace p by $-p - \beta$ in Eq. (35). We finally obtain

$$\mathbf{L}[e^{-(p+\beta)x}] = Ae^{-(p+\beta)x} + q(-p - \beta)e^{px}. \quad (36)$$

Let us multiply Eq. (35) by A and Eq. (36) by $-q(p)$ and add the resulting relations. This yields

$$\mathbf{L}[Ae^{px} - q(p)e^{-(p+\beta)x}] = [A^2 - q(p)q(-p - \beta)]e^{px}. \quad (37)$$

On dividing Eq. (37) by the constant $A^2 - q(p)q(-p - \beta)$, we obtain the original model solution

$$Y(x, p) = \frac{Ae^{px} - q(p)e^{-(p+\beta)x}}{A^2 - q(p)q(-p - \beta)}, \quad \mathbf{L}[Y(x, p)] = e^{px}. \quad (38)$$

Since here $-\infty < x < \infty$, one must set $p = iu$ and use the formulas from Subsection 11.6-3. Then the solution of Eq. (32) for an arbitrary function $f(x)$ can be represented in the form

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} Y(x, iu)\tilde{f}(u) du, \quad \tilde{f}(u) = \int_{-\infty}^{\infty} f(x)e^{-iux} dx. \quad (39)$$

⊙ References for Section 13.9: M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), V. I. Smirnov (1974), P. P. Zabreyko, A. I. Koshelev, et al. (1975), F. D. Gakhov and Yu. I. Cherskii (1978), A. D. Polyanin and A. V. Manzhirov (1997, 1998).

13.10. Carleman Method for Integral Equations of Convolution Type of the Second Kind

13.10-1. Wiener-Hopf Equation of the Second Kind.

Equations of convolution type of the second kind of the form*

$$y(x) + \frac{1}{\sqrt{2\pi}} \int_0^{\infty} K(x-t)y(t) dt = f(x), \quad 0 < x < \infty, \quad (1)$$

frequently occur in applications. Here the domain of the kernel $K(x)$ is the entire real axis.

Let us extend the equation domain to the negative semiaxis by introducing one-sided functions,

$$y_+(x) = \begin{cases} y(x) & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad f_+(x) = \begin{cases} f(x) & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad y_-(x) = 0 \quad \text{for } x > 0.$$

Then we obtain an equation,

$$y_+(x) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K(x-t)y_+(t) dt = y_-(x) + f_+(x), \quad -\infty < x < \infty, \quad (2)$$

which coincides with (1) for $x > 0$.

* Prior to reading this section looking through Sections 12.7 and 12.8 is recommended.

The auxiliary function $y_-(x)$ is introduced to compensate for the left-hand side of Eq. (2) for $x < 0$. Note that $y_-(x)$ is unknown for $x < 0$ and is to be found in solving the problem.

Let us pass to the Fourier integrals in Eq. (2) (see Subsections 9.4-3, 12.7-1, and 12.7-2). We obtain a Riemann problem in the form

$$\mathcal{Y}^+(u) = \frac{\mathcal{Y}^-(u)}{1 + \mathcal{K}(u)} + \frac{\mathcal{F}^+(u)}{1 + \mathcal{K}(u)}, \quad -\infty < u < \infty. \quad (3)$$

1°. Assume that the normality condition is satisfied, i.e.,

$$1 + \mathcal{K}(u) \neq 0,$$

then we rewrite the Riemann problem in the usual form

$$\mathcal{Y}^+(u) = \mathcal{D}(u)\mathcal{Y}^-(u) + \mathcal{H}(u), \quad -\infty < u < \infty, \quad (4)$$

where

$$\mathcal{D}(u) = \frac{1}{1 + \mathcal{K}(u)}, \quad \mathcal{H}(u) = \frac{\mathcal{F}(u)}{1 + \mathcal{K}(u)}. \quad (5)$$

The Riemann problem (4) is equivalent to Eq. (1); in particular, these equations are simultaneously solvable or unsolvable and have an equal number of arbitrary constants in their general solutions. If the *index* ν of the Riemann problem, which is given by the relation

$$\nu = \text{Ind} \frac{1}{1 + \mathcal{K}(u)} \quad (6)$$

(which is also sometimes called the *index of the Wiener–Hopf equation of the second kind*), is positive, then the homogeneous equation (1) ($f(x) \equiv 0$) has exactly ν linearly independent solutions, and the nonhomogeneous equation is unconditionally solvable and its solution depends on ν arbitrary complex constants.

In the case $\nu \leq 0$, the homogeneous equation has no nonzero solutions. For $\nu = 0$, the nonhomogeneous equation is unconditionally solvable, and the solution is unique. If the index ν is negative, then the conditions

$$\int_{-\infty}^{\infty} \frac{\mathcal{F}(u) du}{\mathcal{X}^+(u)[1 + \mathcal{K}(u)](u + i)^k} = 0, \quad k = 1, 2, \dots, -\nu, \quad (7)$$

are necessary and sufficient for the solvability of the nonhomogeneous equation (see Subsection 12.7-4).

For all cases in which the solution of Eq. (1) exists, it can be found by the formula

$$y(x) = y_+(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{Y}^+(u) e^{-iux} du, \quad x > 0, \quad (8)$$

where $\mathcal{Y}^+(u)$ is the solution of the Riemann problem (4) and (5) that is constructed by the scheme of Subsection 12.7-4 (see Fig. 5). The last formula shows that the solution does not depend on $\mathcal{Y}^-(u)$, i.e., is independent of the choice of the extension of the equation to the negative semiaxis.

2°. Now let us study the exceptional case of the integral equation (1) in which the normality condition for the Riemann problem (3) (see Subsections 12.7-6 and 12.7-7) is violated. In this case, the coefficient $D(u) = [1 + \mathcal{K}(u)]^{-1}$ has no zeros, and its order at infinity is $\eta = 0$. The general solution to the boundary value problem (3) can be obtained by formulas (63) of Subsection 12.7-7




Figure 6. Scheme of solving the Wiener–Hopf integral equations. For $\beta = 0$, we have the equation of the first kind, and for $\beta = 1$, we have the equation of the second kind.

for $\alpha_i = 0$. The solution of the original integral equation (1) can be determined from the solution of the boundary value problem on applying formula (8).

Figure 6 depicts a scheme of solving the Wiener–Hopf equations (see also Subsection 12.8-1).

Example. Consider the equation

$$y(x) + \int_0^\infty (a + b|x-t|)e^{-|x-t|}y(t) dt = f(x), \quad x > 0, \tag{9}$$

where the constants a and b are real, and $b \neq 0$. The kernel $K(x-t)$ of Eq. (1) is given by the expression

$$K(x) = \sqrt{2\pi} (a + b|x|)e^{-|x|}.$$

Let us find the transform of the kernel,

$$\mathcal{K}(u) = \int_{-\infty}^\infty (a + b|x|)e^{-|x|+iux} dx = 2 \frac{u^2(a-b) + a+b}{(u^2+1)^2}.$$

Hence,

$$1 + \mathcal{K}(u) = \frac{P(u)}{(u^2+1)^2}, \quad P(z) = z^4 + 2(a-b+1)z^2 + 2a+2b+1.$$

On the basis of the normality condition, we assume that the constants a and b are such that the polynomial $P(z)$ has no real roots. Let $\alpha + i\beta$ be a root of the biquadratic equation $P(z) = 0$ such that $\alpha > 0$ and $\beta > 0$. Since the coefficients of the equation are real, it is clear that $(\alpha - i\beta)$, $(-\alpha + i\beta)$, and $(-\alpha - i\beta)$ are the other three roots. Since the function $1 + \mathcal{K}(u)$ is real as well, it follows that it has zero index, and hence Eq. (9) is uniquely solvable.

On factorizing, we obtain the relation $1 + \mathcal{K}(u) = \mathcal{X}^-(u)/\mathcal{X}^+(u)$, where

$$\mathcal{X}^+(u) = \frac{(u+i)^2}{(u+\alpha+i\beta)(u-\alpha+i\beta)}, \quad \mathcal{X}^-(u) = \frac{(u-\alpha-i\beta)(u+\alpha-i\beta)}{(u-i)^2}.$$

Applying this result, we represent the boundary condition (4), (5) in the form

$$\frac{\mathcal{Y}^+(u)}{\mathcal{X}^+(u)} - \frac{(u-i)^2 \mathcal{F}^+(u)}{(u-\alpha-i\beta)(u+\alpha-i\beta)} = \frac{\mathcal{Y}^-(u)}{\mathcal{X}^-(u)}, \quad -\infty < u < \infty. \tag{10}$$

It follows from the theorem on the analytic continuation and the generalized Liouville theorem (see Subsection 12.7-3) that both sides of the above relation are equal to

$$\frac{C_1}{u-\alpha-i\beta} + \frac{C_2}{u+\alpha-i\beta},$$

where the constants C_1 and C_2 must be defined. Hence,

$$\mathcal{Y}^+(u) = \mathcal{X}^+(u) \left(\frac{(u-i)^2 \mathcal{F}^+(u)}{(u-\alpha-i\beta)(u+\alpha-i\beta)} + \frac{C_1}{u-\alpha-i\beta} + \frac{C_2}{u+\alpha-i\beta} \right). \tag{11}$$

For the poles $(\alpha + i\beta)$ and $(-\alpha + i\beta)$ to be deleted, it is necessary and sufficient that

$$C_1 = -\frac{(\alpha+i\beta-i)^2 \mathcal{F}^+(\alpha+i\beta)}{2\alpha}, \quad C_2 = -\frac{(-\alpha+i\beta-i)^2 \mathcal{F}^+(-\alpha+i\beta)}{-2\alpha}. \tag{12}$$

Since the problem is more or less cumbersome, we pass from the transform (11) to the corresponding original function in two stages. We first find the inverse transform of the summand

$$\mathcal{Y}_1(u) = \mathcal{X}^+(u) \frac{(u-i)^2 \mathcal{F}^+(u)}{(u-\alpha-i\beta)(u+\alpha-i\beta)} = \frac{1}{1+\mathcal{K}(u)} \mathcal{F}^+(u) = \mathcal{F}^+(u) + \mathcal{R}(u)\mathcal{F}^+(u).$$

Here

$$\mathcal{R}(u) = -\frac{2u^2(a-b) + 2a + 2b}{[u^2 - (\alpha+i\beta)^2][u^2 - (\alpha-i\beta)^2]} = \frac{\mu}{u^2 - (\alpha+i\beta)^2} + \frac{\bar{\mu}}{u^2 - (\alpha-i\beta)^2}, \quad \mu = i \frac{(\alpha+i\beta)^2(a-b) + a+b}{2\alpha\beta}.$$

Let us find the inverse transform of the first fraction:

$$\mathbf{F}^{-1} \left\{ \frac{\mu}{u^2 - (\alpha+i\beta)^2} \right\} = \sqrt{\frac{\pi}{2}} \frac{\mu}{\beta-i\alpha} e^{-(\beta-i\alpha)|x|}.$$

The inverse transform of the second fraction can be found in the form

$$\mathbf{F}^{-1} \left\{ \frac{\bar{\mu}}{u^2 - (\alpha-i\beta)^2} \right\} = \sqrt{\frac{\pi}{2}} \frac{\bar{\mu}}{\beta+i\alpha} e^{-(\beta+i\alpha)|x|}. \tag{13}$$

Thus,

$$R(x) = \sqrt{\frac{\pi}{2}} \rho (e^{i\theta+i\alpha|x|} + e^{-i\theta-i\alpha|x|}) e^{-\beta|x|} = \sqrt{2\pi} \rho e^{-\beta|x|} \cos(\theta + \alpha|x|)$$

and

$$y_1(x) = f(x) + \rho \int_0^\infty e^{-\beta|x-t|} \cos(\theta + \alpha|x-t|) f(t) dt, \quad x > 0, \quad \rho e^{i\theta} = \frac{\mu}{\beta-i\alpha}. \tag{14}$$

Note that, as a by-product, we have found the resolvent $R(x-t)$ of the following integral equation on the entire axis:

$$y_0(x) + \int_{-\infty}^\infty (a+b|x-t|) e^{-|x-t|} y_0(t) dt = f_0(x), \quad -\infty < x < \infty.$$

Now consider the remaining part of the transform (11):

$$\mathcal{Y}_2(u) = \mathcal{X}^+(u) \left(\frac{C_1}{u-\alpha-i\beta} + \frac{C_2}{u+\alpha-i\beta} \right).$$

We can calculate the integrals

$$\mathbf{F}^{-1} \{ \mathcal{Y}_2(u) \} = \frac{C_1}{\sqrt{2\pi}} \int_{-\infty}^\infty \frac{(u+i)^2 e^{-iux} du}{(u+i\beta-\alpha)(u+i\beta+\alpha)(u-\alpha-i\beta)} + \frac{C_2}{\sqrt{2\pi}} \int_{-\infty}^\infty \frac{(u+i)^2 e^{-iux} du}{(u+i\beta-\alpha)(u+i\beta+\alpha)(u+\alpha-i\beta)}$$

by means of the residue theory (see Subsections 9.1-4 and 9.1-5) and substitute the values (12) into the constants C_1 and C_2 . For $x > 0$, we obtain

$$y_2(x) = \frac{[\alpha + (\beta - 1)^2]^2}{4\alpha^2\beta} \int_0^\infty e^{-\beta(x+t)} \cos[\alpha(x-t)]f(t) dt + \frac{\rho^*}{4\alpha^2} \int_0^\infty e^{-\beta(x+t)} \cos[\psi + \alpha(x+t)]f(t) dt, \quad \rho^* e^{i\psi} = \frac{(\beta - 1 - i\alpha)^4}{8\alpha^2(\beta - i\alpha)}. \quad (15)$$

Since $\mathcal{Y}^+(u) = \mathcal{Y}_1(u) + \mathcal{Y}_2(u)$, it follows that the desired solution is the sum of the functions (14) and (15).

13.10-2. Integral Equation of the Second Kind with Two Kernels.

Consider an integral equation of convolution type of the second kind with two kernels of the form

$$y(x) + \frac{1}{\sqrt{2\pi}} \int_0^\infty K_1(x-t)y(t) dt + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 K_2(x-t)y(t) dt = f(x), \quad -\infty < x < \infty. \quad (16)$$

Note that each of the kernels $K_1(x)$ and $K_2(x)$ is defined on the entire real axis. On representing the desired function as the difference of one-sided functions,

$$y(x) = y_+(x) - y_-(x), \quad (17)$$

we rewrite the equation in the form

$$y_+(x) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty K_1(x-t)y_+(t) dt - y_-(x) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty K_2(x-t)y_-(t) dt = f(x). \quad (18)$$

Applying the Fourier integral transform (see Subsection 9.4-3), we obtain

$$[1 + \mathcal{K}_1(u)]\mathcal{Y}^+(u) - [1 + \mathcal{K}_2(u)]\mathcal{Y}^-(u) = \mathcal{F}(u). \quad (19)$$

This implies the relation

$$\mathcal{Y}^+(u) = \frac{1 + \mathcal{K}_2(u)}{1 + \mathcal{K}_1(u)}\mathcal{Y}^-(u) + \frac{\mathcal{F}(u)}{1 + \mathcal{K}_1(u)}. \quad (20)$$

Here $\mathcal{K}_1(u)$, $\mathcal{K}_2(u)$, and $\mathcal{F}(u)$ stand for the Fourier integrals of known functions. The unknown transforms $\mathcal{Y}^+(u)$ and $\mathcal{Y}^-(u)$ are the boundary values of functions that are analytic on the upper and lower half-planes, respectively. Thus, we have obtained a Riemann boundary value problem.

1°. Assume that the normality conditions are satisfied, i.e.,

$$1 + \mathcal{K}_1(u) \neq 0, \quad 1 + \mathcal{K}_2(u) \neq 0,$$

then we can rewrite the Riemann problem in the usual form (see Subsection 12.7-4):

$$\mathcal{Y}^+(u) = \mathcal{D}(u)\mathcal{Y}^-(u) + \mathcal{H}(u), \quad -\infty < u < \infty, \quad (21)$$

where

$$\mathcal{D}(u) = \frac{1 + \mathcal{K}_2(u)}{1 + \mathcal{K}_1(u)}, \quad \mathcal{H}(u) = \frac{\mathcal{F}(u)}{1 + \mathcal{K}_1(u)}. \quad (22)$$

The Riemann problem (21), (22) is equivalent to Eq. (16): these problems are solvable or unsolvable simultaneously, and have the same number of arbitrary constants in their general solutions.

If the index

$$\nu = \text{Ind} \frac{1 + \mathcal{K}_2(u)}{1 + \mathcal{K}_1(u)} \quad (23)$$

is positive, then the homogeneous equation (16) ($f(x) \equiv 0$) has precisely ν linearly independent solutions, and the nonhomogeneous equation is unconditionally solvable; moreover, the solution of this equation depends on ν arbitrary complex constants.

In the case $\nu \leq 0$, the homogeneous equation has no nonzero solutions. The nonhomogeneous equation is unconditionally solvable for $\nu = 0$, and the solution is unique. For the case in which the index ν is negative, the conditions

$$\int_{-\infty}^{\infty} \frac{\mathcal{F}(u) du}{\mathcal{X}^+(u)[1 + \mathcal{K}_1(u)](u+i)^k} = 0, \quad k = 1, 2, \dots, -\nu, \quad (24)$$

are necessary and sufficient for the solvability of the nonhomogeneous equation.

In all cases for which the solution of Eq. (16) exists, this solution can be found by the formula

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} [\mathcal{Y}^+(u) - \mathcal{Y}^-(u)] e^{-iux} du, \quad -\infty < x < \infty, \quad (25)$$

where $\mathcal{Y}^+(u)$, $\mathcal{Y}^-(u)$ is the solution of the Riemann problem (21), (22) constructed with respect to the scheme of Subsection 12.7-4 (see Fig. 5).

Thus, the solution of Eq. (16) is equivalent to the solution of a Riemann boundary value problem and is reduced to the calculation of finitely many Fourier integrals.

2°. Now let us study the exceptional case of an integral equation of the form (16). Assume that the functions $1 + \mathcal{K}_1(u)$ and $1 + \mathcal{K}_2(u)$ can have zeros, and these zeros can be both different and coinciding points of the contour. Let us write out the expansion of these functions on selecting the coinciding zeros:

$$\begin{aligned} 1 + \mathcal{K}_1(u) &= \prod_{j=1}^s (u - b_j)^{\beta_j} \prod_{k=1}^p (u - d_k)^{\gamma_k} \mathcal{K}_{11}(u), \\ 1 + \mathcal{K}_2(u) &= \prod_{i=1}^r (u - a_i)^{\alpha_i} \prod_{k=1}^p (u - d_k)^{\gamma_k} \mathcal{K}_{12}(u), \quad \sum_{k=1}^p \gamma_k = l. \end{aligned} \quad (26)$$

Here $a_i \neq b_j$, but it is possible that some points d_k ($k = 1, \dots, p$) coincide with either a_i or b_j . This corresponds to the case in which the functions $1 + \mathcal{K}_1(u)$ and $1 + \mathcal{K}_2(u)$ have a common zero of different multiplicity. We do not select these points especially because their presence does not affect the solvability conditions and the number of solutions of the problem.

It follows from Eq. (19) and from the condition that a solution must be finite on the contour that, for the solvability of the problem, and all the more for the solvability of Eq. (16), it is necessary that the function $\mathcal{F}(u)$ have zero of order γ_k at any point d_k , i.e., $\mathcal{F}(u)$ must have the form

$$\mathcal{F}(u) = \prod_{k=1}^p (u - d_k)^{\gamma_k} \mathcal{F}_1(u).$$

To this end, the following $\gamma_1 + \dots + \gamma_p = l$ conditions must be satisfied:

$$\mathcal{F}_1^{(j_k)}(d_k) = 0, \quad j_k = 0, 1, \dots, \gamma_k - 1, \quad (27)$$

or, which is the same,

$$\int_{-\infty}^{\infty} f(x) x^{j_k} e^{id_k x} dx = 0. \quad (28)$$

Since the functions $\mathcal{K}_1(u)$ and $\mathcal{K}_2(u)$ vanish at infinity, it follows that the point at infinity is a regular point of $\mathcal{D}(u)$.

Assume that conditions (28) are satisfied. In this case the Riemann boundary value problem (20) can be rewritten in the form (see Subsections 12.7-6 and 12.7-7)

$$\mathcal{Y}^+(u) = \frac{\prod_{i=1}^r (u - a_i)^{\alpha_i} \mathcal{R}_+(u) \mathcal{R}_-(u)}{\prod_{j=1}^s (u - b_j)^{\beta_j} \mathcal{Q}_+(u) \mathcal{Q}_-(u)} \mathcal{D}_2(u) \mathcal{Y}^-(u) + \frac{\mathcal{H}_1(u)}{\prod_{j=1}^s (u - b_j)^{\beta_j}}. \tag{29}$$

On finding its general solution in the exceptional case under consideration, we obtain the general solution of the original equation by means of formula (25).

Let us state the conclusions on the solvability conditions and on the number of solutions of Eq. (16). For the solvability of Eq. (16), it is necessary that the Fourier transform of the right-hand side of the equation satisfies l conditions of the form (27). If these conditions are satisfied, then, for $\nu - n > 0$, problem (20) and the integral equation (16) have exactly $\nu - n$ linearly independent solutions. For $\nu - n \leq 0$, we must take the polynomial $\mathcal{P}_{\nu-n-1}(z)$ to be identically zero, and, for the case in which $\nu - n < 0$, the right-hand side must satisfy another $n - \nu$ conditions. If the latter conditions are satisfied, then the integral equation has a unique solution.

Example. Consider Eq. (16) for which

$$K_1(x) = \begin{cases} -(1 + \alpha)\sqrt{2\pi} e^{-x} & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad K_2(x) = \begin{cases} -(1 + \beta)\sqrt{2\pi} e^{-x} & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad f(x) = \begin{cases} 0 & \text{for } x > 0, \\ -\sqrt{2\pi} e^x & \text{for } x < 0, \end{cases}$$

where α and β are real constants. In this case, $K_1(x - t) = 0$ for $x < t$ and $K_2(x - t) = 0$ for $x < t$. Hence, the equation under consideration has the form

$$\begin{aligned} y(x) - (1 + \alpha) \int_0^x e^{-(x-t)} y(t) dt - (1 + \beta) \int_{-\infty}^0 e^{-(x-t)} y(t) dt &= 0, & x > 0, \\ y(x) - (1 + \beta) \int_{-\infty}^x e^{-(x-t)} y(t) dt &= -\sqrt{2\pi} e^x, & x < 0. \end{aligned}$$

Let us calculate the Fourier integrals

$$\mathcal{K}_1(u) = -(1 + \alpha) \int_0^\infty e^{-x} e^{iux} dx = -\frac{i(1 + \alpha)}{u + i}, \quad \mathcal{K}_2(u) = -\frac{i(1 + \beta)}{u + i}, \quad \mathcal{F}(u) = \frac{i}{u - i}, \quad \mathcal{D}(u) = \frac{u - i\beta}{u - i\alpha}.$$

The boundary condition can be rewritten in the form

$$\mathcal{Y}^+(u) = \frac{u - i\beta}{u - i\alpha} \mathcal{Y}^-(u) + \frac{i(u + i)}{(u - i)(u - i\alpha)}. \tag{30}$$

The solution of the Riemann problem depends on the signs of α and β .

1°. Let $\alpha > 0$ and $\beta > 0$. In this case we have $\nu = \text{Ind } \mathcal{D}(u) = 0$. The left-hand side and the right-hand side of the boundary condition contain functions that have analytic continuations to the upper and the lower half-plane, respectively. On applying the theorem on the analytic continuation directly and the generalized Liouville theorem (Subsection 12.7-3), we see that

$$\mathcal{Y}^+(z) = 0, \quad \frac{z - i\beta}{z - i\alpha} \mathcal{Y}^-(z) + \frac{i(z + i)}{(z - i)(z - i\alpha)} = 0.$$

Hence,

$$y_+(x) = 0, \quad y(x) = -y_-(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \frac{i(u + i)}{(u - i)(u - i\beta)} e^{-iux} du.$$

On calculating the last integral, under the assumption that $\beta \neq 1$, by the Cauchy residue theorem (see Subsections 9.1-4 and 9.1-5) we obtain

$$y(x) = \begin{cases} 0 & \text{for } x > 0, \\ -\frac{\sqrt{2\pi}}{1 - \beta} [2e^x - (1 + \beta)e^{\beta x}] & \text{for } x < 0. \end{cases}$$

In the case $\beta = 1$, we have

$$y(x) = \begin{cases} 0 & \text{for } x > 0, \\ -\sqrt{2\pi} e^x (1 + 2x) & \text{for } x < 0. \end{cases}$$

2°. Let $\alpha < 0$ and $\beta < 0$. Here we again have $\nu = 0$, $\mathcal{X}^+(z) = (z - i\beta)(z - i\alpha)^{-1}$, and $\mathcal{X}^-(z) = 1$. On grouping the terms containing the boundary values of functions that are analytic in each of the half-planes and then applying the analytic continuation theorem and the generalized Liouville theorem (Subsection 12.7-3), we see that

$$\frac{\mathcal{Y}^+(z)}{\mathcal{X}^+(z)} + \frac{\beta + 1}{i(\beta - 1)} \frac{1}{z - i\beta} = \frac{\mathcal{Y}^-(z)}{\mathcal{X}^-(z)} + \frac{2}{i(\beta - 1)} \frac{1}{z - i} = 0.$$

Hence,

$$\mathcal{Y}^+(z) = \frac{\beta + 1}{\beta - 1} \frac{i}{z - i\alpha}, \quad \mathcal{Y}^-(z) = \frac{2i}{\beta - 1} \frac{1}{z - i},$$

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} [\mathcal{Y}^+(u) - \mathcal{Y}^-(u)] e^{-iux} du = \begin{cases} \sqrt{2\pi} \frac{\beta + 1}{\beta - 1} e^{\alpha x} & \text{for } x > 0, \\ \frac{2\sqrt{2\pi}}{\beta - 1} e^x & \text{for } x < 0. \end{cases}$$

3°. Let $\alpha < 0$ and $\beta > 0$. In this case we have $\nu = 1$. Let us rewrite the boundary condition (30) in the form

$$\mathcal{Y}^+(u) + \frac{i(1 + \alpha)}{1 - \alpha} \frac{1}{u - i\alpha} = \frac{u - i\beta}{u - i\alpha} \mathcal{Y}^-(u) - \frac{2i}{1 - \alpha} \frac{1}{u - i}.$$

On applying the analytic continuation theorem and the generalized Liouville theorem (Subsection 12.7-3), we see that

$$\mathcal{Y}^+(z) + \frac{i(1 + \alpha)}{1 - \alpha} \frac{1}{z - i\alpha} = \frac{z - i\beta}{z - i\alpha} \mathcal{Y}^-(z) - \frac{2i}{1 - \alpha} \frac{1}{z - i} = \frac{C}{z - i\alpha}.$$

Therefore,

$$\mathcal{Y}^+(z) = \left(C - i \frac{1 + \alpha}{1 - \alpha} \right) \frac{1}{z - i\alpha}, \quad \mathcal{Y}^-(z) = \frac{C}{z - i\beta} - \frac{2i}{1 - \alpha} \frac{z - i\alpha}{(z - i)(z - i\beta)},$$

where C is an arbitrary constant. Now, by means of the Fourier inversion formula, we obtain the general solution of the integral equation in the form

$$y(x) = \begin{cases} -\sqrt{2\pi} \left(iC + \frac{1 + \alpha}{1 - \alpha} \right) e^{\alpha x} & \text{for } x > 0, \\ -\sqrt{2\pi} \left[iC + \frac{2(\alpha - \beta)}{(1 - \alpha)(1 - \beta)} \right] e^{\beta x} - \frac{2\sqrt{2\pi}}{1 - \beta} e^x & \text{for } x < 0. \end{cases}$$

4°. Let $\alpha > 0$ and $\beta < 0$. In this case we have $\nu = -1$. By the Liouville theorem (see Subsection 12.7-3), we obtain

$$\mathcal{Y}^+(z) = \frac{z - i\beta}{z - i\alpha} \mathcal{Y}^-(z) + \frac{i(z + i)}{(z - i)(z - i\alpha)} = 0,$$

and hence

$$\mathcal{Y}^+(z) = 0, \quad \mathcal{Y}^-(z) = -\frac{i(z + i)}{(z - i)(z - i\beta)}.$$

It can be seen from the expression for $\mathcal{Y}^-(z)$ that the singularity of the function $\mathcal{Y}^-(z)$ at the point $i\beta$ disappears if we set $\beta = -1$. The last condition is exactly the solvability condition of the Riemann problem. In this case we have the unique solution

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{i}{u - i} e^{-iux} du = \begin{cases} 0 & \text{for } x > 0, \\ -\sqrt{2\pi} e^x & \text{for } x < 0. \end{cases}$$

Remark 1. Some equations whose kernels contain not the difference but certain other combinations of arguments, namely, the product or, more frequently, the ratio, can be reduced to equations considered in Subsection 13.10-2. For instance, the equation

$$Y(\xi) + \int_0^1 \frac{1}{\tau} N_1 \left(\frac{\xi}{\tau} \right) Y(\tau) d\tau + \int_1^\infty \frac{1}{\tau} N_2 \left(\frac{\xi}{\tau} \right) Y(\tau) d\tau = g(\xi), \quad \xi > 0, \quad (31)$$

becomes a usual equation with two kernels after the following changes of the functions and their arguments: $\xi = e^x$, $\tau = e^t$, $N_1(\xi) = K_1(x)$, $N_2(\xi) = K_2(x)$, $g(\xi) = f(x)$, and $Y(\xi) = y(x)$.

13.10-3. Equations of Convolution Type with Variable Integration Limit.

1°. Consider the Volterra integral equation of the second kind

$$y(x) + \frac{1}{\sqrt{2\pi}} \int_0^x K(x-t)y(t) dt = f(x), \quad 0 \leq x < T, \quad (32)$$

where the interval $[0, T]$ can be either finite or infinite. In contrast with Eq. (1), where the kernel is defined on the entire real axis, here the kernel is defined on the positive semiaxis.

Equation (32) can be regarded as a special case of the one-sided equation (1) of Subsection 13.10-1. To see this, we can rewrite Eq. (32) in the form

$$y(x) + \frac{1}{\sqrt{2\pi}} \int_0^\infty K_+(x-t)y(t) dt = f(x), \quad 0 < x < \infty,$$

which can be reduced to the following boundary value problem:

$$\mathcal{Y}^+(u) = \frac{\mathcal{Y}^-(u)}{1 + \mathcal{K}^+(u)} + \frac{\mathcal{F}^+(u)}{1 + \mathcal{K}^+(u)}.$$

Here the coefficient $[1 + \mathcal{K}^+(u)]^{-1}$ of the problem is a function that has an analytic continuation to the upper half-plane, possibly except for finitely many poles that are zeros of the function $1 + \mathcal{K}^+(z)$ (we assume that $1 + \mathcal{K}^+(z) \neq 0$ on the real axis). Therefore, the index ν of the problem is always nonpositive, $\nu \leq 0$. On rewriting the problem in the form $[1 + \mathcal{K}^+(u)]\mathcal{Y}^+(u) = \mathcal{Y}^-(u) + \mathcal{F}^+(u)$, we see that $\mathcal{Y}^-(u) \equiv 0$, which implies

$$\mathcal{Y}^+(u) = \frac{\mathcal{F}^+(u)}{1 + \mathcal{K}^+(u)}. \quad (33)$$

Consider the following cases.

1.1. The function $1 + \mathcal{K}^+(z)$ has no zeros on the upper half-plane (this means that $\nu = 0$). In this case, Eq. (32) has a unique solution for an arbitrary right-hand side $f(x)$, and this solution can be expressed via the resolvent:

$$y(x) = f(x) + \frac{1}{\sqrt{2\pi}} \int_0^x R(x-t)f(t) dt, \quad x > 0, \quad (34)$$

where

$$R(x) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \frac{\mathcal{K}^+(u)}{1 + \mathcal{K}^+(u)} e^{-iux} du.$$

1.2. The function $1 + \mathcal{K}^+(z)$ has zeros at the points $z = a_1, \dots, a_m$ of the upper half-plane (in this case we have $\nu < 0$, and ν is equal to the minus total order of the zeros). The following two possibilities can occur.

(a) The function $\mathcal{F}^+(z)$ vanishes at the points a_1, \dots, a_m , and the orders of these zeros are not less than the orders of the corresponding zeros of the function $1 + \mathcal{K}^+(z)$. In this case, the function $\mathcal{F}^+(z)[1 + \mathcal{K}^+(z)]^{-1}$ has no poles again, and thus the equation has the unique solution (34).

The assumption $d^k \mathcal{F}^+(a_j)/dz^k = 0$ on the zeros of the function $\mathcal{F}^+(z)$ is equivalent to the conditions

$$\int_{-\infty}^\infty f(t)e^{-ia_j t} t^k dt = 0, \quad k = 0, \dots, \eta_j - 1, \quad j = 1, \dots, m, \quad (35)$$

where η_j is the multiplicity of the zero of the function $1 + \mathcal{K}^+(z)$ at the point a_j . In this case, conditions (35) are imposed directly on the right-hand side of the equation.

(b) The function $\mathcal{F}^+(z)$ does not vanish at the points a_1, \dots, a_m (or vanishes with less multiplicity than $1 + \mathcal{K}^+(z)$). In this case, the function $\mathcal{F}^+(z)[1 + \mathcal{K}^+(z)]^{-1}$ has poles, and therefore the function (33) does not belong to the class under consideration. Equation (32) has no solutions in the chosen class of functions. In this case, conditions (35) fail.

The last result does not contradict the well-known fact that a Volterra equation always has a unique solution. Equation (32) belongs to the class of Volterra type equations, and therefore is also solvable in case (b), but in a broader space of functions with exponential growth.

2°. Another simple special case of Eq. (1) in Subsection 13.10-1 is the following equation with variable lower limit:

$$y(x) + \frac{1}{\sqrt{2\pi}} \int_x^\infty K(x-t)y(t) dt = f(x), \quad 0 < x < \infty. \tag{36}$$

This corresponds to the case in which the function $K(x)$ in Eq. (1) is left one-sided: $K(x) = K_-(x)$. Under the assumption $1 + \mathcal{K}^-(u) \neq 0$, the Riemann problem becomes

$$\mathcal{Y}^+(u) = \frac{\mathcal{Y}^-(u)}{1 + \mathcal{K}^-(u)} + \frac{\mathcal{F}^+(u)}{1 + \mathcal{K}^-(u)}. \tag{37}$$

2.1. The function $1 + \mathcal{K}^-(z)$ has no zeros on the lower half-plane. This means that the inverse transform of the function $\mathcal{Y}^-(u)[1 + \mathcal{K}^-(u)]^{-1}$ is left one-sided, and such a function does not influence the relation between the inverse transforms of (37) for $x > 0$. Thus, if we introduce the function

$$\mathcal{R}^-(u) = -\frac{\mathcal{K}^-(u)}{1 + \mathcal{K}^-(u)}$$

(for convenience of the final formula), then by applying the Fourier inversion formula to Eq. (37) and by setting $x > 0$ we obtain the unique solution to Eq. (36),

$$y(x) = f(x) + \frac{1}{\sqrt{2\pi}} \int_x^\infty R_-(x-t)f(t) dt, \quad x > 0.$$

2.2. The function $1 + \mathcal{K}^-(z)$ has zeros in the lower half-plane. Since this function is nonzero both on the entire real axis and at infinity, it follows that the number of zeros is finite. The Riemann problem (37) has a positive index which is just equal to the number of zeros in the lower half-plane (the zeros are counted according to their multiplicities):

$$\nu = \text{Ind} \frac{1}{1 + \mathcal{K}^-(u)} = -\text{Ind}[1 + \mathcal{K}^-(u)] = \eta_1 + \dots + \eta_n > 0.$$

Here η_k are the multiplicities of the zeros z_k of the function $1 + \mathcal{K}^-(z)$, $k = 1, \dots, n$.

Let

$$\frac{C_{1k}}{z - z_k} + \frac{C_{2k}}{(z - z_k)^2} + \dots + \frac{C_{\eta_k k}}{(z - z_k)^{\eta_k}}$$

be the principal part of the Laurent series expansion of the function $\mathcal{Y}^-(z)[1 + \mathcal{K}^-(z)]^{-1}$ in powers of $(z - z_k)$, $k = 1, \dots, n$. In this case, Eq. (37) becomes

$$\mathcal{Y}^+(u) = \frac{\mathcal{F}^+(u)}{1 + \mathcal{K}^-(u)} + \sum_{k=1}^n \sum_{j=1}^{\eta_k} \frac{C_{jk}}{(z - z_k)^j} + \dots, \tag{38}$$

where the dots denote a function whose inverse transform vanishes for $x > 0$. Under the passage to the inverse transforms in Eq. (38), for $x > 0$ we obtain

$$y(x) = f(x) + \frac{1}{\sqrt{2\pi}} \int_x^\infty R_-(x-t)f(t) dt + \sum_{k=1}^n P_k(x)e^{-iz_k x}, \quad x > 0. \tag{39}$$

Here the $P_k(x)$ are polynomials of degree $\eta_k - 1$. We can verify that the function (39) is a solution of Eq. (36) for arbitrary coefficients of the polynomials. Since the number of linearly independent solutions of the homogeneous equation (36) is equal to the index, it follows that the above solution (39) is the general solution of the nonhomogeneous equation.

13.10-4. Dual Equation of Convolution Type of the Second Kind.

Consider the dual integral equation of the second kind

$$\begin{aligned} y(x) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K_1(x-t)y(t) dt &= f(x), & 0 < x < \infty, \\ y(x) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K_2(x-t)y(t) dt &= f(x), & -\infty < x < 0, \end{aligned} \quad (40)$$

in which the function $y(x)$ is to be found.

In order to apply the Fourier transform technique (see Subsections 11.4-3, 12.7-1, and 12.7-2), we extend the domain of both conditions in Eq. (40) by formally rewriting them for all real values of x . This can be achieved by introducing new unknown functions into the right-hand sides. These functions must be chosen so that the conditions given on the semiaxis are not violated. Hence, the first condition in (40) must be complemented by a summand that vanishes on the positive semiaxis and the second by a summand that vanishes on the negative semiaxis. Thus, the dual equation can be written in the form

$$\begin{aligned} y(x) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K_1(x-t)y(t) dt &= f(x) + \xi_-(x), \\ y(x) + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K_2(x-t)y(t) dt &= f(x) + \xi_+(x), \end{aligned} \quad -\infty < x < \infty, \quad (41)$$

where the $\xi_{\pm}(x)$ are some right and left one-sided functions so far unknown.

On applying the Fourier integral transform, we arrive at the relations

$$[1 + \mathcal{K}_1(u)]\mathcal{Y}(u) = \mathcal{F}(u) + \Xi^-(u), \quad [1 + \mathcal{K}_2(u)]\mathcal{Y}(u) = \mathcal{F}(u) + \Xi^+(u). \quad (42)$$

Here the three functions $\mathcal{Y}(u)$, $\Xi^+(u)$, and $\Xi^-(u)$ are unknown.

Now on the basis of (42) we can find

$$\mathcal{Y}(u) = \frac{\mathcal{F}(u) + \Xi^-(u)}{1 + \mathcal{K}_1(u)} = \frac{\mathcal{F}(u) + \Xi^+(u)}{1 + \mathcal{K}_2(u)} \quad (43)$$

and eliminate the function $\mathcal{Y}(u)$ from relations (42) by applying formula (43). We obtain the Riemann boundary value problem in the form

$$\Xi^+(u) = \frac{1 + \mathcal{K}_2(u)}{1 + \mathcal{K}_1(u)} \Xi^-(u) + \frac{\mathcal{K}_2(u) - \mathcal{K}_1(u)}{1 + \mathcal{K}_1(u)} \mathcal{F}(u), \quad -\infty < u < \infty. \quad (44)$$

1°. Assume that the normality conditions are satisfied, i.e.,

$$1 + \mathcal{K}_1(u) \neq 0, \quad 1 + \mathcal{K}_2(u) \neq 0;$$

then we can rewrite the Riemann problem (44) in the usual form (see Subsection 12.7-4)

$$\Xi^+(u) = \mathcal{D}(u)\Xi^-(u) + \mathcal{H}(u), \quad -\infty < u < \infty, \quad (45)$$

where

$$\mathcal{D}(u) = \frac{1 + \mathcal{K}_2(u)}{1 + \mathcal{K}_1(u)}, \quad \mathcal{H}(u) = \frac{\mathcal{K}_2(u) - \mathcal{K}_1(u)}{1 + \mathcal{K}_1(u)} \mathcal{F}(u). \quad (46)$$

The Riemann problem (45), (46) is equivalent to Eq. (40); in particular, they are solvable and unsolvable simultaneously and have the same number of arbitrary constants in the general solutions.

If the index

$$\nu = \text{Ind} \frac{1 + \mathcal{K}_2(u)}{1 + \mathcal{K}_1(u)} \quad (47)$$

is positive, then the homogeneous equation (40) ($f(x) \equiv 0$) has exactly ν linearly independent solutions, and the nonhomogeneous equation is unconditionally solvable and the solution depends on ν arbitrary complex constants.

For the case $\nu \leq 0$, the homogeneous equation has no nonzero solutions. For $\nu = 0$, the nonhomogeneous equation is unconditionally solvable, and a solution is unique. If the index ν is negative, then the conditions

$$\int_{-\infty}^{\infty} \frac{\mathcal{K}_2(u) - \mathcal{K}_1(u)}{\mathcal{X}^+(u)[1 + \mathcal{K}_1(u)]} \mathcal{F}(u) \frac{du}{(u+i)^k} = 0, \quad k = 1, 2, \dots, -\nu \quad (48)$$

are necessary and sufficient for the solvability of the nonhomogeneous equation.

For all cases in which a solution of Eq. (40) exists, it can be found by the formula

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\mathcal{F}(u) + \Xi^-(u)}{1 + \mathcal{K}_1(u)} e^{-iux} du = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{\mathcal{F}(u) + \Xi^+(u)}{1 + \mathcal{K}_2(u)} e^{-iux} du, \quad (49)$$

where $\Xi^+(u)$, $\Xi^-(u)$ is a solution of the Riemann problem (45), (46) that is constructed by the scheme of Subsection 12.7-4 (see Fig. 5).

2°. Let us investigate the exceptional case of the integral equation (40). Assume that the functions $1 + \mathcal{K}_1(u)$ and $1 + \mathcal{K}_2(u)$ can have zeros that can be either different or coinciding points of the contour. Take the expansions of these functions on selecting the coinciding zeros in the form of (26) and further repeat the reasoning performed for the equations of convolution type of the second kind with two kernels. After finding the general solution of the Riemann boundary value problem (44) in this exceptional case (see Subsection 12.7-7), we obtain the general solution of the original equation (40) by formula (49).

The conclusions on the solvability conditions and on the number of solutions of Eq. (40) are similar to those made above for the equations with two kernels in Subsection 13.10-2.

Remark 2. Equations treated in Section 13.10 are sometimes called *characteristic equations of convolution type*.

⊙ References for Section 13.10: F. D. Gakhov and Yu. I. Cherskii (1978), F. D. Gakhov (1990).

13.11. Wiener–Hopf Method

13.11-1. Some Remarks.

Suppose that the Fourier transform of the function $y(x)$ exists (see Subsection 9.4-3):

$$\mathcal{Y}(z) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} y(x) e^{izx} dx. \quad (1)$$

Assume that the parameter z that enters the transform (1) can take complex values as well. Let us study the properties of the function $\mathcal{Y}(z)$ regarded as a function of the complex variable z . To this end, we represent the function $y(x)$ in the form*

$$y(x) = y^+(x) + y^-(x), \quad (2)$$

* Do not confuse the functions $y^\pm(x)$ and $\mathcal{Y}_\pm(x)$ introduced in this section with the functions $y_\pm(x)$ and $\mathcal{Y}^\pm(x)$ introduced in Subsection 12.7-2 and used in solving the Riemann boundary value problem on the real axis.

where the functions $y^+(x)$ and $y^-(x)$ are given by the relations

$$y^+(x) = \begin{cases} y(x) & \text{for } x > 0, \\ 0 & \text{for } x < 0, \end{cases} \quad y^-(x) = \begin{cases} 0 & \text{for } x > 0, \\ y(x) & \text{for } x < 0. \end{cases} \quad (3)$$

In this case the transform $\mathcal{Y}(z)$ of the function $y(x)$ is clearly equal to the sum of the transforms $\mathcal{Y}_+(z)$ and $\mathcal{Y}_-(z)$ of the functions $y^+(x)$ and $y^-(x)$, respectively. Let us clarify the analytic properties of the function $\mathcal{Y}(z)$ by establishing the analytic properties of the functions $\mathcal{Y}_+(z)$ and $\mathcal{Y}_-(z)$. Consider the function $y^+(x)$ given by relations (3). Its transform is equal to

$$\mathcal{Y}_+(z) = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} y^+(x)e^{izx} dx. \quad (4)$$

It can be shown that if the function $y^+(x)$ satisfies the condition

$$|y^+(x)| < Me^{v_-x} \quad \text{as } x \rightarrow \infty, \quad (5)$$

where M is a constant, then the function $\mathcal{Y}_+(z)$ given by formula (4) is an analytic function of the complex variable $z = u + iv$ in the domain $\text{Im } z > v_-$, and in this domain we have $\mathcal{Y}_+(z) \rightarrow 0$ as $|z| \rightarrow \infty$. We can also show that the functions $y^+(x)$ and $\mathcal{Y}_+(z)$ are related as follows:

$$y^+(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty+iv}^{\infty+iv} \mathcal{Y}_+(z)e^{-izx} dz, \quad (6)$$

where the integration is performed over any line $\text{Im } z = v > v_-$ in the complex plane z , which is parallel to the real axis.

For $v_- < 0$ (i.e., for functions $y(x)$ with exponential decay at infinity), the real axis belongs to the domain in which the function $\mathcal{Y}_+(z)$ is analytic, and we can integrate over the real axis in formula (6). However, if the only possible values of v_- are positive (for instance, if the function $y^+(x)$ has nontrivial growth at infinity, which does not exceed the exponential growth with linear exponent), then the analyticity domain of the function $\mathcal{Y}_+(z)$ is strictly above the real axis of the complex plane z (and in this case, the integral (4) can be divergent on the real axis). Similarly, if the function $y^-(x)$ in relations (3) satisfies the condition

$$|y^-(x)| < Me^{v_+x} \quad \text{as } x \rightarrow -\infty, \quad (7)$$

then its transform, i.e., the function

$$\mathcal{Y}_-(z) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 y^-(x)e^{izx} dx, \quad (8)$$

is an analytic function of the complex variable z in the domain $\text{Im } z < v_+$. The function $y^-(x)$ can be expressed via $\mathcal{Y}_-(z)$ by means of the relation

$$y^-(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty+iv}^{\infty+iv} \mathcal{Y}_-(z)e^{-izx} dz, \quad \text{Im } z = v < v_+. \quad (9)$$

For $v_+ > 0$, the analyticity domain of the function $\mathcal{Y}_-(z)$ contains the real axis.

It is clear that for $v_- < v_+$, the function $\mathcal{Y}(z)$ defined by formula (1) is an analytic function of the complex variable z in the strip $v_- < \text{Im } z < v_+$. In this case, the functions $y(x)$ and $\mathcal{Y}(z)$ are related by the Fourier inversion formula

$$y(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty+iv}^{\infty+iv} \mathcal{Y}(z)e^{-izx} dz, \quad (10)$$

where the integration is performed over an arbitrary line in the complex plane z belonging to the strip $v_- < \text{Im } z < v_+$. In particular, for $v_- < 0$ and $v_+ > 0$, the function $\mathcal{Y}(z)$ is analytic in the strip containing the real axis of the complex plane z .

Example 1. For $\alpha > 0$, the function $K(x) = e^{-\alpha|x|}$ has the transform

$$\mathcal{K}(z) = \frac{1}{\sqrt{2\pi}} \frac{2\alpha}{\alpha^2 + z^2},$$

which is an analytic function of the complex variable z in the strip $-\alpha < \text{Im } z < \alpha$, which contains the real axis.

13.11-2. Homogeneous Wiener-Hopf Equation of the Second Kind.

Consider a homogeneous integral Wiener-Hopf equation of the second kind in the form

$$y(x) = \int_0^{\infty} K(x-t)y(t) dt, \quad (11)$$

whose solution can obviously be determined up to an arbitrary constant factor only. Here the domain of the function $K(x)$ is the entire real axis. This factor can be found from additional conditions of the problem, for instance, from normalization conditions.

We assume that Eq. (11) defines a function $y(x)$ for all values of the variable x , positive and negative. Let us introduce the functions $y^-(x)$ and $y^+(x)$ by formulas (3). Obviously, we have $y(x) = y^+(x) + y^-(x)$, and Eq. (11) can be rewritten in the form

$$y^+(x) = \int_0^{\infty} K(x-t)y^+(t) dt, \quad x > 0 \quad (12)$$

$$y^-(x) = \int_0^{\infty} K(x-t)y^+(t) dt, \quad x < 0. \quad (13)$$

That is, the function $y^+(x)$ can be determined by the solution of the integral equation (12) and the function $y^-(x)$ can be expressed via the functions $y^+(x)$ and $K(x)$ by means of formulas (13). In this case, we have the relation

$$y^+(x) + y^-(x) = \int_{-\infty}^{\infty} K(x-t)y^+(t) dt, \quad (14)$$

which is equivalent to the original equation (11).

Let the function $K(x)$ satisfy the condition

$$\begin{aligned} |K(x)| < M e^{v_- x} & \text{ as } x \rightarrow \infty, \\ |K(x)| < M e^{v_+ x} & \text{ as } x \rightarrow -\infty, \end{aligned} \quad (15)$$

where $v_- < 0$ and $v_+ > 0$. In this case, the function

$$\mathcal{K}(z) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} K(x)e^{izx} dx \quad (16)$$

is analytic in the strip $v_- < \text{Im } z < v_+$.

Let us seek the solution of Eq. (11) satisfying the condition

$$|y^+(x)| < M_1 e^{\mu x} \quad \text{as } x \rightarrow \infty, \quad (17)$$

where $\mu < v_+$ (such a solution exists). In this case we can readily verify that the integrals on the right-hand sides in (12) and (13) are convergent, and the function $y^-(x)$ satisfies the estimate

$$|y^-(x)| < M_2 e^{v_+ x} \quad \text{as } x \rightarrow -\infty. \quad (18)$$

It follows from conditions (17) and (18) that the transforms $\mathcal{Y}_+(z)$ and $\mathcal{Y}_-(z)$ of the functions $y^+(x)$ and $y^-(x)$ are analytic functions of the complex variable z for $\text{Im } z > \mu$ and $\text{Im } z < v_+$, respectively.

Let us pass to the solution of the integral equation (11) or of Eq. (14), which is equivalent to (11). To this end, we apply the (alternative) Fourier transform. By the convolution theorem (see Subsection 9.4-4), it follows from (14) that

$$\mathcal{Y}_+(z) + \mathcal{Y}_-(z) = \sqrt{2\pi} \mathcal{K}(z) \mathcal{Y}_+(z),$$

or

$$\mathcal{W}(z)\mathcal{Y}_+(z) + \mathcal{Y}_-(z) = 0, \tag{19}$$

where

$$\mathcal{W}(z) = 1 - \sqrt{2\pi} \mathcal{K}(z) \neq 0. \tag{20}$$

Thus, by means of the Fourier transform, we succeeded in the passage from the original integral equation to an algebraic equation for the transforms. However, in this case Eq. (19) involves two unknown functions. In general, a single algebraic equation cannot uniquely determine two unknown functions. The Wiener–Hopf method makes it possible to solve this problem for a certain class of functions. This method is mainly related to the study of the analyticity domains of the functions that enter the equation and to a special representation of this equation. The main idea of the Wiener–Hopf method is as follows.

Let Eq. (19) be representable in the form

$$\mathcal{W}_+(z)\mathcal{Y}_+(z) = -\mathcal{W}_-(z)\mathcal{Y}_-(z), \tag{21}$$

where the left-hand side is analytic in the upper half-plane $\text{Im } z > \mu$ and the right-hand side is analytic in the lower half-plane $\text{Im } z < v_+$, where $\mu < v_+$, so that there exists a common analyticity strip of these functions: $\mu < \text{Im } z < v_+$. Since the analytic continuation is unique, it follows that there exists a unique entire function of the complex variable that coincides with the left-hand side of (21) in the upper half-plane and with the right-hand side of (21) in the lower half-plane, respectively. If, in addition, the functions that enter Eq. (21) have at most power-law growth with respect to z at infinity, then it follows from the generalized Liouville theorem (see Subsection 12.7-3) that the entire function under consideration is a polynomial. In particular, for the case of a function that is bounded at infinity we obtain

$$\mathcal{W}_+(z)\mathcal{Y}_+(z) = -\mathcal{W}_-(z)\mathcal{Y}_-(z) = \text{const}. \tag{22}$$

These relations uniquely determine the functions $\mathcal{Y}_+(z)$ and $\mathcal{Y}_-(z)$.

Thus, let us apply the above scheme to the solution of Eq. (19). It follows from the above reasoning that the analyticity domains of the functions $\mathcal{Y}_+(z)$, $\mathcal{Y}_-(z)$, and $\mathcal{W}(z) = 1 - \sqrt{2\pi} \mathcal{K}(z)$, respectively, are the upper half-plane $\text{Im } z > \mu$, the lower half-plane $\text{Im } z < v_+$, and the strip $v_- < \text{Im } z < v_+$. Therefore, this equation holds in the strip* $\mu < \text{Im } z < v_+$, which is the common analyticity domain for all functions that enter the equation. In order to transform Eq. (19) to the form (21), we assume that it is possible to decompose the function $\mathcal{W}(z)$ as follows:

$$\mathcal{W}(z) = \frac{\mathcal{W}_+(z)}{\mathcal{W}_-(z)}, \tag{23}$$

where the functions $\mathcal{W}_+(z)$ and $\mathcal{W}_-(z)$ are analytic for $\text{Im } z > \mu$ and $\text{Im } z < v_+$, respectively. Moreover, we assume that, in the corresponding analyticity domains, these functions grow at infinity no faster than z^n , where n is a positive integer. A representation of an analytic function $\mathcal{W}(z)$ in the form (23) is often called a *factorization* of $\mathcal{W}(z)$.

Thus, as the result of factorization, the original equation is reduced to the form (21). It follows from the above reasoning that this equation determines an entire function of the complex variable z .

Since $\mathcal{Y}_\pm(z) \rightarrow 0$ as $|z| \rightarrow \infty$ and the growth of the functions $\mathcal{W}_\pm(z)$ does not exceed that of a power function z^n , it follows that the entire function under consideration can be only a polynomial $\mathcal{P}_{n-1}(z)$ of degree at most $n - 1$.

If the growth of the functions $\mathcal{W}_\pm(z)$ at infinity is only linear with respect to the variable z , then it follows from relations (22), by virtue of the Liouville theorem (see Subsection 12.7-3), that the

* To be definite, we set $\mu > v_-$. Otherwise, the common domain of analyticity is the strip $v_- < \text{Im } z < v_+$.

corresponding entire function is a constant C . In this case we obtain the following relations for the unknown functions $\mathcal{Y}_+(z)$ and $\mathcal{Y}_-(z)$:

$$\mathcal{Y}_+(z) = \frac{C}{\mathcal{W}_+(z)}, \quad \mathcal{Y}_-(z) = -\frac{C}{\mathcal{W}_-(z)}, \quad (24)$$

which define the transform of the solution up to a constant factor, which can be found at least from the normalization conditions. In the general case, the expressions

$$\mathcal{Y}_+(z) = \frac{\mathcal{P}_{n-1}(z)}{\mathcal{W}_+(z)}, \quad \mathcal{Y}_-(z) = -\frac{\mathcal{P}_{n-1}(z)}{\mathcal{W}_-(z)}, \quad (25)$$

define the transform of the desired solution of the integral equation (11) up to indeterminate constants, which can be found from the additional conditions of the problem. The solution itself is defined by means of the Fourier inversion formula (6), (9), and (10).

Example 2. Consider the equation

$$y(x) = \lambda \int_0^\infty e^{-|x-t|} y(t) dt, \quad 0 < \lambda < \infty, \quad (26)$$

whose kernel has the form $K(x) = \lambda e^{-|x|}$.

Let us find the transform of the function $K(x)$:

$$\mathcal{K}(z) = \frac{\lambda}{\sqrt{2\pi}} \int_{-\infty}^\infty K(x) e^{izx} dx = \sqrt{\frac{2}{\pi}} \frac{\lambda}{z^2 + 1}. \quad (27)$$

The function $\mathcal{K}(z)$ is analytic with respect to the complex variable z in the strip $-1 < \text{Im } z < 1$. Let us represent the expression

$$\mathcal{W}(z) = 1 - \sqrt{2\pi} \mathcal{K}(z) = \frac{z^2 - 2\lambda + 1}{z^2 + 1} \quad (28)$$

in the form (23), where

$$\mathcal{W}_+(z) = \frac{z^2 - 2\lambda + 1}{z + i}, \quad \mathcal{W}_-(z) = z - i. \quad (29)$$

The function $\mathcal{W}_+(z)$ in Eq. (29) is analytic with respect to z and nonzero in the domain $\text{Im } z > \text{Im } \sqrt{2\lambda - 1}$. For $0 < \lambda < \frac{1}{2}$, this domain is defined by the condition $\text{Im } z > \sqrt{1 - 2\lambda}$, and $\sqrt{1 - 2\lambda} \leq \mu < 1$. For $\lambda > \frac{1}{2}$, the function $\mathcal{W}_+(z)$ is analytic and nonzero in the domain $\text{Im } z > 0$. It is clear that the function $\mathcal{W}_-(z)$ is a nonzero analytic function in the domain $\text{Im } z < 1$. Therefore, for $0 < \lambda < \frac{1}{2}$ both functions satisfy the required conditions in the domain $\mu < \text{Im } z < 1$.

For $\lambda > \frac{1}{2}$, the strip $0 < \text{Im } z < 1$ is the common domain of analyticity of the functions $\mathcal{W}_+(z)$ and $\mathcal{W}_-(z)$. Thus, we have obtained the desired factorization of the function (28).

Consider the expressions $\mathcal{Y}_\pm(z)\mathcal{W}_\pm(z)$. Since $\mathcal{Y}_\pm(z) \rightarrow 0$ as $|z| \rightarrow \infty$, and, according to (29), the growth of the functions $\mathcal{W}_\pm(z)$ at infinity is linear with respect to z , it follows that the entire function $\mathcal{P}_{n-1}(z)$ that coincides with $\mathcal{Y}_+(z)\mathcal{W}_+(z)$ for $\text{Im } z > \mu$ and with $\mathcal{Y}_-(z)\mathcal{W}_-(z)$ for $\text{Im } z < 1$ can be a polynomial of zero degree only. Therefore,

$$\mathcal{Y}_+(z)\mathcal{W}_+(z) = C. \quad (30)$$

Hence,

$$\mathcal{Y}_+(z) = C \frac{z + i}{z^2 - 2\lambda + 1}, \quad (31)$$

and it follows from (6) that

$$y^+(x) = \frac{C}{\sqrt{2\pi}} \int_{-\infty+iv}^{\infty+iv} \frac{z + i}{z^2 - 2\lambda + 1} e^{-izx} dz, \quad (32)$$

where $\mu < v < 1$.

On closing the integration contour for $x > 0$ by a semicircle in the lower half-plane and estimating the integral over this semicircle by means of the Jordan lemma (see Subsections 9.1-4 and 9.1-5), after some calculations we obtain

$$y^+(x) = C \left[\cos(\sqrt{2\lambda - 1} x) + \frac{\sin(\sqrt{2\lambda - 1} x)}{\sqrt{2\lambda - 1}} \right], \quad (33)$$

where C is a constant. For $0 < \lambda < \frac{1}{2}$, this solution has exponential growth with respect to x , and for $\frac{1}{2} < \lambda < \infty$, it is bounded at infinity.

13.11-3. General Scheme of the Method. The Factorization Problem.

In the general case, the problem which is solved by the Wiener–Hopf method can be reduced to the following problem. It is required to find functions $\mathcal{Y}_+(z)$ and $\mathcal{Y}_-(z)$ of the complex variable z that are analytic in the half-planes $\text{Im } z > v_-$ and $\text{Im } z < v_+$, respectively ($v_- < v_+$), vanish as $|z| \rightarrow \infty$ in their analyticity domains, and satisfy the following functional equation in the strip ($v_- < \text{Im } z < v_+$):

$$A(z)\mathcal{Y}_+(z) + B(z)\mathcal{Y}_-(z) + C(z) = 0. \quad (34)$$

Here $A(z)$, $B(z)$, and $C(z)$ are given functions of the complex variable z that are analytic in the strip $v_- < \text{Im } z < v_+$, and the functions $A(z)$ and $B(z)$ are nonzero in this strip.

The main idea of the solution of this problem is based on the possibility of a factorization of the expression $A(z)/B(z)$, i.e., of a representation in the form

$$\frac{A(z)}{B(z)} = \frac{\mathcal{W}_+(z)}{\mathcal{W}_-(z)}, \quad (35)$$

where the functions $\mathcal{W}_+(z)$ and $\mathcal{W}_-(z)$ are analytic and nonzero in the half-planes $\text{Im } z > v'_-$ and $\text{Im } z < v'_+$, and the strips $v_- < \text{Im } z < v_+$ and $v'_- < \text{Im } z < v'_+$ have a nonempty common part. In this case Eq. (34), with regard to Eq. (35), can be rewritten in the form

$$\mathcal{W}_+(z)\mathcal{Y}_+(z) + \mathcal{W}_-(z)\mathcal{Y}_-(z) + \mathcal{W}_-(z)\frac{C(z)}{B(z)} = 0. \quad (36)$$

If the last summand in Eq. (36) can be represented as the sum

$$\mathcal{W}_-(z)\frac{C(z)}{B(z)} = \mathcal{D}_+(z) + \mathcal{D}_-(z), \quad (37)$$

where the functions $\mathcal{D}_+(z)$ and $\mathcal{D}_-(z)$ are analytic in the half-planes $\text{Im } z > v''_-$ and $\text{Im } z < v''_+$, respectively, and all three strips $v_- < \text{Im } z < v_+$, $v'_- < \text{Im } z < v'_+$, and $v''_- < \text{Im } z < v''_+$ have a nonempty common part, for a strip $v_-^0 < \text{Im } z < v_+^0$, then, in this common strip, the following functional equation holds:

$$\mathcal{W}_+(z)\mathcal{Y}_+(z) + \mathcal{D}_+(z) = -\mathcal{W}_-(z)\mathcal{Y}_-(z) - \mathcal{D}_-(z). \quad (38)$$

The left-hand side of Eq. (38) is a function analytic in the half-plane $v_-^0 < \text{Im } z$, and the right-hand side is a function analytic in the domain $\text{Im } z < v_+^0$. Since these functions coincide in the strip $v_-^0 < \text{Im } z < v_+^0$, it follows that there exists a unique entire function that coincides with the left-hand side and the right-hand side of (38) in their analyticity domains, respectively. If the growth at infinity of all functions that enter the right-hand sides of Eqs. (35) and (37), in their analyticity domains, is at most that of z^n , then it follows from the limit relation $\mathcal{Y}_\pm(z) \rightarrow 0$ as $|z| \rightarrow \infty$ that this entire function is a polynomial $\mathcal{P}_{n-1}(z)$ of degree at most $n - 1$. Thus, the relations

$$\mathcal{Y}_+(z) = \frac{\mathcal{P}_{n-1}(z) - \mathcal{D}_+(z)}{\mathcal{W}_+(z)}, \quad \mathcal{Y}_-(z) = \frac{-\mathcal{P}_{n-1}(z) - \mathcal{D}_-(z)}{\mathcal{W}_-(z)} \quad (39)$$

determine the desired functions up to constants. These constants can be found from the additional conditions of the problem.

The application of the Wiener–Hopf method is based on the representations (35) and (37). If a function $\mathcal{G}(z)$ is analytic in the strip $v_- < \text{Im } z < v_+$ and if in this strip the function $\mathcal{G}(z)$ uniformly tends to zero as $|z| \rightarrow \infty$, then in this strip the following representation is possible:

$$\mathcal{G}(z) = \mathcal{G}_+(z) + \mathcal{G}_-(z), \quad (40)$$

where the function $\mathcal{G}_+(z)$ is analytic in the half-plane $\text{Im } z > v_-$, the function $\mathcal{G}_-(z)$ is analytic in the half-plane $\text{Im } z < v_+$, and

$$\mathcal{G}_+(z) = \frac{1}{2\pi i} \int_{-\infty+iv'_-}^{\infty+iv'_-} \frac{\mathcal{G}(\tau)}{\tau-z} d\tau, \quad v_- < v'_- < \text{Im } z < v_+, \quad (41)$$

$$\mathcal{G}_-(z) = -\frac{1}{2\pi i} \int_{-\infty+iv'_+}^{\infty+iv'_+} \frac{\mathcal{G}(\tau)}{\tau-z} d\tau, \quad v_- < \text{Im } z < v'_+ < v_+. \quad (42)$$

The integrals (41) and (42), being regarded as integrals depending on a parameter, define analytic functions of the complex variable z under the assumption that the point z does not belong to the integration contour.

In particular, $\mathcal{G}_+(z)$ is an analytic function in the half-plane $\text{Im } z > v'_-$ and $\mathcal{G}_-(z)$ in the half-plane $\text{Im } z > v'_+$.

Moreover, if a function $\mathcal{H}(z)$ is analytic and nonzero in the strip $v_- < \text{Im } z < v_+$ and if $\mathcal{H}(z) \rightarrow 1$ uniformly in this strip as $|z| \rightarrow \infty$, then the following representation holds in the strip:

$$\mathcal{H}(z) = \mathcal{H}_+(z)\mathcal{H}_-(z), \quad (43)$$

$$\mathcal{H}_+(z) = \exp \left[\frac{1}{2\pi i} \int_{-\infty+iv'_-}^{\infty+iv'_-} \frac{\ln \mathcal{H}(\tau)}{\tau-z} d\tau \right], \quad v_- < v'_- < \text{Im } z < v_+, \quad (44)$$

$$\mathcal{H}_-(z) = \exp \left[-\frac{1}{2\pi i} \int_{-\infty+iv'_+}^{\infty+iv'_+} \frac{\ln \mathcal{H}(\tau)}{\tau-z} d\tau \right], \quad v_- < \text{Im } z < v'_+ < v_+, \quad (45)$$

where the functions $\mathcal{H}_+(z)$ and $\mathcal{H}_-(z)$ are analytic and nonzero in the half-planes $\text{Im } z > v_-$ and $\text{Im } z < v_+$, respectively. The representation (43) is called a *factorization* of the function $\mathcal{H}(z)$.

13.11-4. Nonhomogeneous Wiener–Hopf Equation of the Second Kind.

Consider the Wiener–Hopf equation of the second kind

$$y(x) - \int_0^\infty K(x-t)y(t) dt = f(x). \quad (46)$$

Suppose that the kernel $K(x)$ of the equation and the right-hand side $f(x)$ satisfy conditions (15). Let us seek the solution $y^+(x)$ to Eq. (46) for which condition (17) is satisfied.

In this case, reasoning similar to that in the derivation of the functional equation (19) for a homogeneous integral equation shows that, in the case of Eq. (46), the following functional equation must hold on the strip $\mu < \text{Im } z < v_+$:

$$\mathcal{Y}_+(z) + \mathcal{Y}_-(z) = \sqrt{2\pi} \mathcal{K}(z)\mathcal{Y}_+(z) + \mathcal{F}_+(z) + \mathcal{F}_-(z), \quad (47)$$

or

$$\mathcal{W}(z)\mathcal{Y}_+(z) + \mathcal{Y}_-(z) - \mathcal{F}(z) = 0, \quad (48)$$

where $\mathcal{W}(z)$ is subjected to condition (20), as well as in the case of a homogeneous equation.

We now note that Eq. (48) is a special case of Eq. (34). In the strip $v_- < \text{Im } z < v_+$, the function $\mathcal{W}(z)$ is analytic and uniformly tends to 1 as $|z| \rightarrow \infty$ because $|\mathcal{K}(z)| \rightarrow 0$ as $|z| \rightarrow \infty$. In this case, this function has the representation (see (43)–(45))

$$\mathcal{W}(z) = \frac{\mathcal{W}_+(z)}{\mathcal{W}_-(z)}, \quad (49)$$

where the function $\mathcal{W}_+(z)$ is analytic in the upper half-plane $\text{Im } z > v_-$ and $\mathcal{W}_-(z)$ is analytic in the lower half-plane $\text{Im } z < v_+$, and the growth at infinity of the functions $\mathcal{W}_\pm(z)$ does not exceed that of z^n .

On the basis of the representation (49), Eq. (48) becomes

$$\mathcal{W}_+(z)\mathcal{Y}_+(z) + \mathcal{W}_-(z)\mathcal{Y}_-(z) - \mathcal{W}_-(z)\mathcal{F}_-(z) - \mathcal{W}_-(z)\mathcal{F}_+(z) = 0. \quad (50)$$

To reduce Eq. (50) to the form (38), it suffices to decompose the last summand

$$\mathcal{F}_+(z)\mathcal{W}_-(z) = \mathcal{D}_+(z) + \mathcal{D}_-(z) \quad (51)$$

into the sum of functions $\mathcal{D}_+(z)$ and $\mathcal{D}_-(z)$ that are analytic in the half-planes $\text{Im } z > \mu$ and $\text{Im } z < v_+$, respectively.

To establish the possibility of a representation (51), we note that the function $\mathcal{F}_+(z)$ is analytic in the upper half-plane $\text{Im } z > v_-$ and uniformly tends to zero as $|z| \rightarrow \infty$. The function $\mathcal{W}_-(z)$ is analytic in the lower half-plane $\text{Im } z < v_+$, and, according to the method of its construction, we can perform the factorization (49) so that the function $\mathcal{W}_-(z)$ remains bounded in the strip $v_- < \text{Im } z < v_+$ as $|z| \rightarrow \infty$. Hence (see (40)–(42)), the functions $\mathcal{F}_+(z)\mathcal{W}_-(z)$ in the strip $v_- < \text{Im } z < v_+$ satisfy all conditions that are sufficient for the validity of the representation (51).

The above reasoning makes it possible to take into account the fact that the growth at infinity of the functions $\mathcal{W}_\pm(z)$ does not exceed that of z^n , and thus to present the transform of the solution of the nonhomogeneous integral equation (46) in the form

$$\mathcal{Y}_+(z) = \frac{\mathcal{P}_{n-1}(z) + \mathcal{D}_+(z)}{\mathcal{W}_+(z)}, \quad \mathcal{Y}_-(z) = \frac{-\mathcal{P}_{n-1}(z) + \mathcal{W}_-(z)\mathcal{F}_-(z) + \mathcal{D}_-(z)}{\mathcal{W}_-(z)}. \quad (52)$$

The solution itself can be obtained from (52) by means of the Fourier inversion formula (6), (9), and (10).

13.11-5. Exceptional Case of a Wiener–Hopf Equation of the Second Kind.

Consider the exceptional case of a Wiener–Hopf equation of the second kind in which the function $\mathcal{W}(z) = 1 - \sqrt{2\pi} \mathcal{K}(z)$ has finitely many zeros N (counted according to their multiplicities) in the strip $v_- < \text{Im } z < v_+$. In this case, the factorization is also possible. To this end, it suffices to introduce the auxiliary function

$$\mathcal{W}_1(z) = \ln \left[(z^2 + b^2)^{N/2} \mathcal{W}(z) \prod_i (z - z_i)^{-\alpha_i} \right], \quad (53)$$

where α_i is the multiplicity of the zero z_i and a positive constant $b > \{|v_-|, |v_+|\}$ is chosen so that the function in the square brackets has no additional zeros in the strip $v_- < \text{Im } z < v_+$.

However, in the exceptional case, the Wiener–Hopf method gives the answer only if the number of zeros of the function $\mathcal{W}(z)$ is even. This restriction is due to the fact that only for the case in which the number of zeros is even is it possible to achieve the necessary behavior at infinity (for the application of the Wiener–Hopf method) of the function $(z^2 + b^2)^{N/2}$ (see F. D. Gakhov and Yu. I. Cherskii (1978)). The last restriction makes no real obstacle to the broad use of the Wiener–Hopf method in solving applied problems in which the kernel $K(x)$ of the corresponding integral equation is frequently an even function, and thus the reasoning below can be applied completely.

Remark 1. The Wiener–Hopf equation of the second kind for functions vanishing at infinity can be reduced to a Riemann boundary value problem on the real axis (see Subsection 13.10-1). In this case, the assumption that the number of zeros of the function $\mathcal{W}(z)$ is even, as well as the assumption that the kernel $K(x)$ is even in the exceptional case, are unessential.

Remark 2. For functions with nontrivial growth at infinity, the complete solution of Wiener-Hopf equations of the second kind is presented in the cited book by F. D. Gakhov and Yu. I. Cherskii (1978).

Remark 3. The Wiener-Hopf method can be applied to solve Wiener-Hopf integral equations of the first kind under the assumption that the kernels of these equations are even.

⊙ References for Section 13.11: B. Noble (1958), A. G. Sveshnikov and A. N. Tikhonov (1970), V. I. Smirnov (1974), F. D. Gakhov (1977, 1990), F. D. Gakhov and Yu. I. Cherskii (1978).

13.12. Krein's Method for Wiener-Hopf Equations

13.12-1. Some Remarks. The Factorization Problem.

Consider the Wiener-Hopf equation of the second kind

$$y(x) - \int_0^{\infty} K(x-t)y(t) dt = f(x), \quad 0 \leq x < \infty, \quad (1)$$

where $f(x)$, $y(x) \in L_1(0, \infty)$ and $K(x) \in L_1(-\infty, \infty)$. Let us use the classes of functions that can be represented as Fourier transforms (alternative Fourier transform in the asymmetric form, see Subsection 9.4-3), of functions from $L_1(-\infty, \infty)$, $L_1(0, \infty)$, and $L_1(-\infty, 0)$. For brevity, instead of these symbols we simply write L , L_+ , and L_- . Let functions $h(x)$, $h_1(x)$, and $h_2(x)$ belong to L , L_+ , and L_- , respectively; in this case, their transforms can be represented in the form

$$\check{H}(u) = \int_{-\infty}^{\infty} h(x)e^{iux} dx, \quad \check{H}_1(u) = \int_0^{\infty} h_1(x)e^{iux} dx, \quad \check{H}_2(u) = \int_{-\infty}^0 h_2(x)e^{iux} dx.$$

Let Q , Q_+ , and Q_- be the classes of functions representable in the form

$$\check{W}(u) = 1 + \check{H}(u), \quad \check{W}_1(u) = 1 + \check{H}_1(u), \quad \check{W}_2(u) = 1 + \check{H}_2(u), \quad (2)$$

respectively, where the functions from the classes Q_+ and Q_- , treated as functions of the complex variable $z = u + iv$, are analytic for $\text{Im } z > 0$ and $\text{Im } z < 0$, respectively, and are continuous up to the real axis.

Let $T(x)$ belong to L and let $\check{T}(u)$ be its transform. Assume that

$$1 - \check{T}(u) \neq 0, \quad \text{Ind}[1 - \check{T}(u)] = \frac{1}{2\pi} \left\{ \arg[1 - \check{T}(u)] \right\}_{-\infty}^{\infty} = 0, \quad -\infty < u < \infty. \quad (3)$$

In this case there exists a $q(x) \in L$ such that

$$\ln[1 - \check{T}(u)] = \int_{-\infty}^{\infty} q(x)e^{iux} dx. \quad (4)$$

This formula readily implies the relation $\ln[1 - \check{T}(u)] \rightarrow 0$ as $u \rightarrow \pm\infty$.

In what follows, we apply the *factorization* of functions $\check{M}(u)$ of the class Q that are continuous on the interval $-\infty \leq u \leq \infty$. Here the factorization means a representation of the function $\check{M}(u)$ in the form of a product

$$\check{M}(u) = \check{M}_+(u) \left(\frac{u-i}{u+i} \right)^k \check{M}_-(u), \quad (5)$$

where $\check{M}_-(z)$ and $\check{M}_+(z)$ are analytic functions in the corresponding half-planes $\text{Im } z > 0$ and $\text{Im } z < 0$ continuous up to the real axis. Moreover,

$$\check{M}_+(z) \neq 0 \quad \text{for} \quad \text{Im } z \geq 0 \quad \text{and} \quad \check{M}_-(z) \neq 0 \quad \text{for} \quad \text{Im } z \leq 0. \quad (6)$$

Relation (5) implies the formula

$$k = \text{Ind } \check{M}(u).$$

The factorization (5) is said to be *canonical* provided that $k = 0$.

In what follows we consider only functions of the form

$$\check{M}(u) = 1 - \check{T}(u) \quad (7)$$

such that $\check{M}(\pm\infty) = 1$. We can also assume that

$$\check{M}_+(\pm\infty) = \check{M}_-(\pm\infty) = 1. \quad (8)$$

Let us state the main results concerning the factorization problem.

A function (7) admits a canonical factorization if and only if the following two conditions hold:

$$\check{M}(u) \neq 0, \quad \text{Ind } \check{M}(u) = 0. \quad (9)$$

In this case, the canonical factorization is unique. Moreover, if conditions (9) hold, then there exists a function $M(x)$ in the class L such that

$$\check{M}(u) = \exp \left[\int_{-\infty}^{\infty} M(x)e^{iux} dx \right], \quad (10)$$

$$\check{M}_+(u) = \exp \left[\int_0^{\infty} M(x)e^{iux} dx \right], \quad \check{M}_-(u) = \exp \left[\int_{-\infty}^0 M(x)e^{iux} dx \right]. \quad (11)$$

Hence, we have $\check{M}(u) \in Q$ and $\check{M}_{\pm}(u) \in Q_{\pm}$. The factors in the canonical factorization are also described by the following formulas:

$$\ln \check{M}_+(z) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\ln \check{M}(\tau)}{\tau - z} d\tau, \quad \text{Im } z > 0, \quad (12)$$

$$\ln \check{M}_-(z) = -\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\ln \check{M}(\tau)}{\tau - z} d\tau, \quad \text{Im } z < 0. \quad (13)$$

In the general case of the factorization, the following assertion holds. A function (7) admits a factorization (5) if and only if the following condition is satisfied:

$$\check{M}(u) \neq 0, \quad -\infty < u < \infty.$$

In this case, relation (5) can be rewritten in the form

$$\left(\frac{u-i}{u+i} \right)^{-k} \check{M}(u) = \check{M}_-(u)\check{M}_+(u), \quad -\infty < u < \infty.$$

The last relation implies the canonical factorization for the function

$$\check{M}_1(u) = \left(\frac{u-i}{u+i} \right)^{-k} \check{M}(u).$$

Hence, the factors $\check{M}_{\pm}(u)$ satisfy formulas (10)–(13) if we replace $\check{M}(u)$ in these formulas by $\check{M}_1(u)$.

Now we return to Eq. (1) for which

$$\check{K}(u) = \int_{-\infty}^{\infty} K(x)e^{iux} dx. \quad (14)$$

13.12-2. Solution of the Wiener-Hopf Equations of the Second Kind.

THEOREM 1. For Eq. (1) to have a unique solution of the class L_+ for an arbitrary $f(x) \in L_+$, it is necessary and sufficient that the following conditions hold:

$$1 - \check{K}(u) \neq 0, \quad -\infty < u < \infty, \quad (15)$$

$$\nu = -\text{Ind}[1 - \check{K}(u)] = 0. \quad (16)$$

THEOREM 2. If condition (15) holds, then the inequality $\nu > 0$ is necessary and sufficient for the existence of nonzero solutions in the class L_+ of the homogeneous equation

$$y(x) - \int_0^\infty K(x-t)y(t) dt = 0. \quad (17)$$

The set of these solutions has a basis formed by ν functions $\varphi_k(x)$ ($k = 1, \dots, \nu$) that tend to zero as $x \rightarrow \infty$ and that are related as follows:

$$\varphi_k(x) = \int_0^x \varphi_{k+1}(t) dt, \quad k = 1, 2, \dots, \nu - 1, \quad \varphi_\nu(x) = \int_0^x \psi(t) dt + C, \quad (18)$$

where C is a nonzero constant and the functions $\varphi_k(t)$ and $\psi(t)$ belong to L_+ .

THEOREM 3. If condition (15) holds and if $\nu > 0$, then for any $f(x) \in L_+$ Eq. (1) has infinitely many solutions in L_+ .

However, if $\nu < 0$, then, for a given $f(x) \in L_+$, Eq. (1) has either no solutions from L_+ or a unique solution. For the latter case to hold, it is necessary and sufficient that the following conditions be satisfied:

$$\int_0^\infty f(x)\psi_k(x) dx = 0, \quad k = 1, 2, \dots, |\nu|, \quad (19)$$

where $\psi_k(x)$ is a basis of the linear space of all solutions of the transposed homogeneous equation

$$\psi(x) - \int_0^\infty K(t-x)\psi(t) dt = 0. \quad (20)$$

1°. If conditions (15) and (16) hold, then there exists a unique factorization

$$[1 - \check{K}(u)]^{-1} = \check{M}_+(u)\check{M}_-(u), \quad (21)$$

and

$$\check{M}_+(u) = 1 + \int_0^\infty R_+(t)e^{iut} dt, \quad \check{M}_-(u) = 1 + \int_0^\infty R_-(t)e^{-iut} dt. \quad (22)$$

The resolvent is defined by the formula

$$R(x, t) = R_+(x-t) + R_-(t-x) + \int_0^\infty R_+(x-s)R_-(t-s) ds \quad (23)$$

where $0 \leq x < \infty$, $0 \leq t < \infty$, $R_+(x) = 0$, and $R_-(x) = 0$ for $x < 0$, so that, for $f(x)$ from L_+ , the solution of the equation is determined by the expression

$$y(x) = f(x) + \int_0^\infty R(x, t)f(t) dt. \quad (24)$$

Formula (23) can be rewritten as follows:

$$R(x, t) = R(x - t, 0) + R(0, t - x) + \int_0^\infty R(x - s, 0)R(0, t - s) ds. \quad (25)$$

If $K(x - t) = K(t - x)$, then formula (25) becomes

$$R(x, t) = R(|x - t|, 0) + \int_0^{\min(x, t)} R(x - s, 0)R(t - s, 0) ds. \quad (26)$$

Note that $R_+(x) = R(x, 0)$ and $R_-(x) = R(0, x)$ are unique solutions, in the class L_+ , of the following equations ($0 \leq x < \infty$):

$$\begin{aligned} R_+(x) + \int_0^\infty K(x - t)R_+(t) dt &= K(x), \\ R_-(x) + \int_0^\infty K(t - x)R_-(t) dt &= K(-x). \end{aligned} \quad (27)$$

2°. Suppose that condition (15) holds, but

$$\nu = -\text{Ind}[1 - \check{K}(u)] > 0.$$

In this case, the function $[1 - \check{K}(u)]^{-1}$ admits the factorization

$$[1 - \check{K}(u)]^{-1} = \check{G}_-(u) \left(\frac{u - i}{u + i} \right)^\nu \check{G}_+(u), \quad -\infty < u < \infty. \quad (28)$$

For the functions $\check{M}_-(u)$ and $\check{M}_+(u)$ defined by the relations

$$\check{M}_-(u) = \check{G}_-(u) \quad \text{and} \quad \check{M}_+(u) = \left(\frac{u - i}{u + i} \right)^\nu \check{G}_+(u), \quad (29)$$

we have the representation (22) and formula (23) for the resolvent.

Moreover, for $k = 1, \dots, \nu$, the following representations hold:

$$\frac{i^k \check{M}_+(u)}{(u - i)^k} = \int_0^\infty g_k(x) e^{iux} dx, \quad (30)$$

where $g_k(x)$ is the solution of the homogeneous equation (17). The solutions $\varphi_k(x)$ mentioned in Theorem 2 can also naturally be expressed via the functions $g_k(x)$.

3°. If $\nu = -\text{Ind}[1 - \check{K}(u)] < 0$, then the transposed equation

$$y(x) - \int_0^\infty K(t - x)y(t) dt = f(x) \quad (31)$$

has the index $-\nu > 0$. If formula (28) defines a factorization for Eq. (1), then the transposed equation admits a factorization of the form

$$[1 - \check{K}(u)]^{-1} = \check{M}_-(-u)\check{M}_+(-u),$$

and $\check{M}_-(-u)$ plays the role of $\check{M}_+(u)$, and $\check{M}_+(-u)$ plays the role of $\check{M}_-(u)$.

13.12-3. Hopf–Fock Formula.

Let us give a useful formula that allows one to express the solution of Eq. (1) with an arbitrary right-hand side $f(x)$ via the solution to a simpler auxiliary integral equation with an exponential right-hand side.

Assume that in Eq. (1) we have

$$f(x) = e^{i\zeta x}, \quad \text{Im } \zeta > 0, \quad y(x) = y_\zeta(x), \quad (32)$$

and moreover, conditions (15) and (16) hold. In this case

$$y_\zeta(x) = e^{i\zeta x} + \int_0^\infty R(x, t)e^{i\zeta t} dt, \quad (33)$$

where $R(x, t)$ has the form (25). After some manipulations, we can see that

$$y_\zeta(x) = \check{\mathcal{M}}_-(-\zeta) \left[1 + \int_0^x R(t, 0)e^{-i\zeta t} dt \right] e^{i\zeta x}. \quad (34)$$

On setting $x = 0$ in (34), we have

$$y_\zeta(0) = \check{\mathcal{M}}_-(-\zeta), \quad (35)$$

and if the function $K(x)$ describing the kernel of the integral equation is even, then

$$y_\zeta(0) = \check{\mathcal{M}}_+(\zeta). \quad (36)$$

On the basis of formula (34), we can obtain the solution of Eq. (1) for a general $f(x)$ as well (see also Section 11.6):

$$y(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \check{\mathcal{F}}_+(-\zeta)y_\zeta(x) d\zeta, \quad \check{\mathcal{F}}_+(u) = \int_0^\infty f(x)e^{iux} dx. \quad (37)$$

Remark 1. All results obtained in Section 13.12 concerning Wiener–Hopf equations of the second kind remain valid for continuous, square integrable, and some other classes of functions, which are discussed in detail in the paper by M. G. Krein (1958) and in the book by C. Corduneanu (1973).

Remark 2. The solution of the Wiener–Hopf equation can be also obtained in other classes of functions for the exceptional case in which $1 - \check{\mathcal{K}}(u) = 0$ (see Subsections 13.10-1 and 13.11-5).

© References for Section 13.12: V. A. Fock (1942), M. G. Krein (1958), C. Corduneanu (1973), V. I. Smirnov (1974), P. P. Zabreyko, A. I. Koshelev, et al. (1975).

13.13. Methods for Solving Equations with Difference Kernels on a Finite Interval

13.13-1. Krein’s Method.

Consider a method for constructing exact analytic solutions of linear integral equations with an arbitrary right-hand side. The method is based on the construction of two auxiliary solutions of simpler equations with the right-hand side equal to 1. The auxiliary solutions are used to construct a solution of the original equation for an arbitrary right-hand side.

1°. Let the equation

$$y(x) - \int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b, \tag{1}$$

be given. Along with (1), we consider two auxiliary equations depending on a parameter ξ ($a \leq \xi \leq b$):

$$\begin{aligned} w(x, \xi) - \int_a^\xi K(x, t)w(t, \xi) dt &= 1, \\ w^*(x, \xi) - \int_a^\xi K(t, x)w^*(t, \xi) dt &= 1, \end{aligned} \tag{2}$$

where $a \leq x \leq \xi$. Assume that for any ξ the auxiliary equations (2) have unique continuous solutions $w(x, \xi)$ and $w^*(x, \xi)$, respectively, which satisfy the condition $w(\xi, \xi)w^*(\xi, \xi) \neq 0$ ($a \leq \xi \leq b$). In this case, for any continuous function $f(x)$, the unique continuous solution of Eq. (1) can be obtained by the formula

$$y(x) = F(b)w(x, b) - \int_x^b w(x, \xi)F'_\xi(\xi) d\xi, \quad F(\xi) = \frac{1}{m(\xi)} \frac{d}{d\xi} \int_a^\xi w^*(t, \xi)f(t) dt, \tag{3}$$

where

$$m(\xi) = w(\xi, \xi)w^*(\xi, \xi).$$

Formula (3) permits one to construct a solution of Eq. (1) with an arbitrary right-hand side $f(x)$ by means of solutions to the two simpler auxiliary equations (2) (depending on the parameter ξ) with a constant right-hand side equal to 1.

2°. Consider now an equation with the kernel depending on the difference of the arguments:

$$y(x) + \int_a^b K(x - t)y(t) dt = f(x), \quad a \leq x \leq b. \tag{4}$$

It is assumed that $K(x)$ is an even function integrable on $[a - b, b - a]$. Along with (4) we consider the following auxiliary equation depending on a parameter ξ ($a \leq \xi \leq b$):

$$w(x, \xi) + \int_a^\xi K(x - t)w(t, \xi) dt = 1, \quad a \leq x \leq \xi. \tag{5}$$

Assume that for an arbitrary ξ the auxiliary equation (5) has a unique continuous solution $w(x, \xi)$. In this case, for any continuous function $f(x)$, a solution of Eq. (4) can be obtained from formula (3) by setting $w^*(x, t) = w(x, t)$ in this formula.

Now let us indicate another useful formula for equations whose kernel depends on the difference of the arguments:

$$y(x) + \int_{-a}^a K(x - t)y(t) dt = f(x), \quad -a \leq x \leq a. \tag{6}$$

It is assumed that $K(x)$ is an even function that is integrable on the segment $[-2a, 2a]$. Along with (6) we consider an auxiliary equation depending on a parameter ξ ($0 < \xi \leq a$):

$$w(x, \xi) + \int_{-\xi}^\xi K(x - t)w(t, \xi) dt = 1, \quad -\xi \leq x \leq \xi. \tag{7}$$

Let the auxiliary equation (7) have a unique continuous solution $w(x, \xi)$ for any ξ . In this case, for an arbitrary continuous function $f(x)$, the solution of Eq. (6) can be obtained by the following formula:

$$\begin{aligned}
 y(x) = & \frac{1}{2M(a)} \left[\frac{d}{da} \int_{-a}^a w(t, a) f(t) dt \right] w(x, a) \\
 & - \frac{1}{2} \int_{|x|}^a w(x, \xi) \frac{d}{d\xi} \left[\frac{1}{M(\xi)} \frac{d}{d\xi} \int_{-\xi}^{\xi} w(t, \xi) f(t) dt \right] d\xi \\
 & - \frac{1}{2} \frac{d}{dx} \int_{|x|}^a \frac{w(x, \xi)}{M(\xi)} \left[\int_{-\xi}^{\xi} w(t, \xi) df(t) \right] d\xi,
 \end{aligned} \tag{8}$$

where $M(\xi) = w^2(\xi, \xi)$, and the last inner integral is treated as a Stieltjes integral.

13.13-2. Kernels with Rational Fourier Transforms.

Consider an equation of the form

$$y(x) - \int_0^T K(x-t)y(t) dt = f(x), \tag{9}$$

where $0 \leq x \leq T < \infty$. If the kernel $K(x)$ is integrable on $[-T, T]$, then the Fredholm theory can be applied to this equation.

Since the equation involves the values of the kernel $K(x)$ for the points of $[-T, T]$ only, it follows that we can extend the kernel outside this interval in an arbitrary way. Assume that the kernel is extended to the entire axis so that the extended function is integrable. In the general case, Eq. (9) in the space $L_2(0, T)$ can be reduced to a boundary value problem of the theory of analytic functions (Riemann problem) for two pairs of unknown functions.

If the Fourier transform of the kernel

$$\check{K}(u) = \int_{-\infty}^{\infty} K(x)e^{iux} dx$$

is rational, then Eq. (9) can be solved in the closed form. Assume that $1 - \check{K}(u) \neq 0$ ($-\infty < u < \infty$). In this case, the transform of the solution of the integral equation (9) is given by the formula

$$\check{Y}(u) = \frac{1}{1 - \check{K}(u)} [\check{F}(u) - \check{W}^+(u) - e^{-iT u} \check{W}^-(u)] \tag{10}$$

in which

$$\check{W}^{\pm}(u) = \sum_n \sum_{k=1}^{p_n^{\pm}} \frac{M_{nk}^{\pm}}{(u - b_n^{\pm})^k},$$

where the b_n^+ and the b_n^- are poles of the functions $1 - \check{K}(u)$ that belong to the upper and lower half-planes, respectively, and the p_n^{\pm} are their multiplicities. The constants M_{nk}^{\pm} can be determined from the conditions

$$\begin{aligned}
 \frac{d^s}{du^s} [\check{W}^+(u) + e^{-iT u} \check{W}^-(u) - \check{F}(u)]_{u=a_n^+} &= 0; & s = 0, 1, \dots, q_n^+ - 1; \\
 \frac{d^s}{du^s} [\check{W}^+(u) + e^{-iT u} - \check{F}(u)]_{u=a_n^-} &= 0; & s = 0, 1, \dots, q_n^- - 1;
 \end{aligned}$$

where a_n^+ and a_n^- are the zeros of the functions $1 - \check{K}(u)$ that belong to the upper and lower half-planes, respectively, and q_n^{\pm} are their multiplicities. The constants M_{nk}^{\pm} can also be determined by substituting the solution into the original equation. The solution of the integral equation (9) can be obtained by inverting formula (10).

13.13-3. Reduction to Ordinary Differential Equations.

1°. Consider the special case in which the Fourier transform of the kernel of the integral equation (9) can be represented in the form

$$\check{K}(u) = \frac{\check{M}(u)}{\check{N}(u)}, \quad (11)$$

where $\check{M}(u)$ and $\check{N}(u)$ are some polynomials of degrees m and n , respectively:

$$\check{M}(u) = \sum_{k=0}^m A_k u^k, \quad \check{N}(u) = \sum_{k=0}^n B_k u^k. \quad (12)$$

In this case, the solution of the integral equation (9) (if it exists) satisfies the following linear nonhomogeneous ordinary differential equation of the order m with constant coefficients:

$$\check{M}\left(i\frac{d}{dx}\right)y(x) = \check{N}\left(i\frac{d}{dx}\right)f(x), \quad 0 < x < T. \quad (13)$$

The solution of Eq. (13) contains m arbitrary constants that are defined by substituting the solution into the original equation (9). Here a system of linear algebraic equations is obtained for these constants.

2°. Consider the Fredholm equation of the second kind with a difference kernel that contains a sum of the exponential functions:

$$y(x) + \int_a^b \left(\sum_{k=1}^n A_k e^{\lambda_k |x-t|} \right) y(t) dt = f(x). \quad (14)$$

In the general case, this equation can be reduced to a linear nonhomogeneous ordinary differential equation of order $2n$ with constant coefficients (see equation 4.2.16 in the first part of the book).

For the solution of Eq. (14) with $n = 1$, see equation 4.2.15 in the first part of the book.

3°. Equations with a difference kernel that contains a sum of hyperbolic functions,

$$y(x) + \int_a^b K(x-t)y(t) dt = f(x), \quad K(x) = \sum_{k=1}^n A_k \sinh(\lambda_k |x|), \quad (15)$$

can be also reduced by differentiation to linear nonhomogeneous ordinary differential equations of order $2n$ with constant coefficients (see equation 4.3.29 in the first part of the book).

For the solution of Eq. (15) with $n = 1$, see equation 4.3.26 in the first part of the book.

4°. Equations with a difference kernel containing a sum of trigonometric functions

$$y(x) + \int_a^b K(x-t)y(t) dt = f(x), \quad K(x) = \sum_{k=1}^n A_k \sin(\lambda_k |x|), \quad (16)$$

can be also reduced to linear nonhomogeneous ordinary differential equations of order $2n$ with constant coefficients (see equations 4.5.29 and 4.5.32 in the first part of the book).

13.14. Method of Approximating a Kernel by a Degenerate One

13.14-1. Approximation of the Kernel.

For the approximate solution of the Fredholm integral equation of the second kind

$$y(x) - \int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b, \quad (1)$$

where, for simplicity, the functions $f(x)$ and $K(x, t)$ are assumed to be continuous, it is useful to replace the kernel $K(x, t)$ by a close degenerate kernel

$$K_{(n)}(x, t) = \sum_{k=0}^n g_k(x)h_k(t). \quad (2)$$

Let us indicate several ways to perform such a change. If the kernel $K(x, t)$ is differentiable with respect to x on $[a, b]$ sufficiently many times, then, for a degenerate kernel $K_{(n)}(x, t)$, we can take a finite segment of the Taylor series:

$$K_{(n)}(x, t) = \sum_{m=0}^n \frac{(x - x_0)^m}{m!} K_x^{(m)}(x_0, t), \quad (3)$$

where $x_0 \in [a, b]$. A similar trick can be applied for the case in which $K(x, t)$ is differentiable with respect to t on $[a, b]$ sufficiently many times.

To construct a degenerate kernel, a finite segment of the double Fourier series can be used:

$$K_{(n)}(x, t) = \sum_{p=0}^n \sum_{q=0}^n a_{pq} (x - x_0)^p (t - t_0)^q, \quad (4)$$

where

$$a_{pq} = \frac{1}{p! q!} \frac{\partial^{p+q}}{\partial x^p \partial t^q} K(x, t) \Big|_{\substack{x=x_0 \\ t=t_0}}, \quad a \leq x_0 \leq b, \quad a \leq t_0 \leq b.$$

A continuous kernel $K(x, t)$ admits an approximation by a trigonometric polynomial of period $2l$, where $l = b - a$.

For instance, we can set

$$K_{(n)}(x, t) = \frac{1}{2} a_0(t) + \sum_{k=1}^n a_k(t) \cos\left(\frac{k\pi x}{l}\right), \quad (5)$$

where the $a_k(t)$ ($k = 0, 1, 2, \dots$) are the Fourier coefficients

$$a_k(t) = \frac{2}{l} \int_a^b K(x, t) \cos\left(\frac{p\pi x}{l}\right) dx. \quad (6)$$

A similar decomposition can be obtained by interchanging the roles of the variables x and t . A finite segment of the double Fourier series can also be applied by setting, for instance,

$$a_k(t) \approx \frac{1}{2} a_{k0} + \sum_{m=1}^n a_{km} \cos\left(\frac{m\pi t}{l}\right), \quad k = 0, 1, \dots, n, \quad (7)$$

and it follows from formulas (5)–(7) that

$$K_{(n)}(x, t) = \frac{1}{4}a_{00} + \frac{1}{2} \sum_{k=1}^n a_{k0} \cos\left(\frac{k\pi x}{l}\right) + \frac{1}{2} \sum_{m=1}^n a_{0m} \cos\left(\frac{m\pi t}{l}\right) + \sum_{k=1}^n \sum_{m=1}^n a_{km} \cos\left(\frac{k\pi x}{l}\right) \cos\left(\frac{m\pi t}{l}\right),$$

where

$$a_{km} = \frac{4}{l^2} \int_a^b \int_a^b K(x, t) \cos\left(\frac{k\pi x}{l}\right) \cos\left(\frac{m\pi t}{l}\right) dx dt. \tag{8}$$

One can also use other methods of interpolating and approximating the kernel $K(x, t)$.

13.14-2. Approximate Solution.

If $K_{(n)}(x, t)$ is an approximate degenerate kernel for a given exact kernel $K(x, t)$ and if a function $f_n(x)$ is close to $f(x)$, then the solution $y_n(x)$ of the integral equation

$$y_n(x) - \int_a^b K_{(n)}(x, t)y_n(t) dt = f_n(x) \tag{9}$$

can be regarded as an approximation to the solution $y(x)$ of Eq. (1).

Assume that the following error estimates hold:

$$\int_a^b |K(x, t) - K_{(n)}(x, t)| dt \leq \varepsilon, \quad |f(x) - f_n(x)| \leq \delta.$$

Next, let the resolvent $R_n(x, t)$ of Eq. (9) satisfy the relation

$$\int_a^b |R_n(x, t)| dt \leq M_n$$

for $a \leq x \leq b$. Finally, assume that the following inequality holds:

$$q = \varepsilon(1 + M_n) < 1.$$

In this case, Eq. (1) has a unique solution $y(x)$ and

$$|y(x) - y_n(x)| \leq \varepsilon \frac{N(1 + M_n)^2}{1 - q} + \delta, \quad N = \max_{a \leq x \leq b} |f(x)|. \tag{10}$$

Example. Let us find an approximate solution of the equation

$$y(x) - \int_0^{1/2} e^{-x^2 t^2} y(t) dt = 1. \tag{11}$$

Applying the expansion in a double Taylor series, we replace the kernel

$$K(x, t) = e^{-x^2 t^2}$$

by the degenerate kernel

$$K_{(2)}(x, t) = 1 - x^2 t^2 + \frac{1}{2} x^4 t^4.$$

Hence, instead of Eq. (11) we obtain

$$y_2(x) = 1 + \int_0^{1/2} (1 - x^2 t^2 + \frac{1}{2} x^4 t^4) y_2(t) dt. \quad (12)$$

Therefore,

$$y_2(x) = 1 + A_1 + A_2 x^2 + A_3 x^4, \quad (13)$$

where

$$A_1 = \int_0^{1/2} y_2(x) dx, \quad A_2 = - \int_0^{1/2} x^2 y_2(x) dx, \quad A_3 = \frac{1}{2} \int_0^{1/2} x^4 y_2(x) dx. \quad (14)$$

From (13) and (14) we obtain a system of three equations with three unknowns; to the fourth decimal place, the solution is

$$A_1 = 0.9930, \quad A_2 = -0.0833, \quad A_3 = 0.0007.$$

Hence,

$$y(x) \approx y_2(x) = 1.9930 - 0.0833 x^2 + 0.0007 x^4, \quad 0 \leq x \leq \frac{1}{2}. \quad (15)$$

An error estimate for the approximate solution (15) can be performed by formula (10).

⊙ References for Section 13.14: L. V. Kantorovich and V. I. Krylov (1958), S. G. Mikhlin (1960), B. P. Demidovich, I. A. Maron, and E. Z. Shuvalova (1963), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), K. E. Atkinson (1997).

13.15. Bateman Method

13.15-1. General Scheme of the Method.

In some cases it is useful, instead of replacing a given kernel by a degenerate kernel, to represent the given kernel approximately as the sum of a kernel whose resolvent is known and a degenerate kernel. For the latter, the resolvent can be written out in a closed form.

Consider the Fredholm integral equation of the second kind

$$y(x) - \lambda \int_a^b k(x, t) y(t) dt = f(x) \quad (1)$$

with kernel $k(x, t)$ whose resolvent $r(x, t; \lambda)$ is known; thus, the solution of (1) can be represented in the form

$$y(x) = f(x) + \lambda \int_a^b r(x, t; \lambda) f(t) dt. \quad (2)$$

Then, for the integral equation with kernel

$$K(x, t) = \frac{1}{\Delta(a_{ij})} \begin{vmatrix} k(x, t) & g_1(x) & \cdots & g_n(x) \\ h_1(t) & a_{11} & \cdots & a_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ h_n(t) & a_{n1} & \cdots & a_{nn} \end{vmatrix}, \quad \Delta(a_{ij}) = \begin{vmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{vmatrix}, \quad (3)$$

where $g_k(x)$ and $h_k(t)$ ($k = 1, \dots, n$) are arbitrary functions and a_{ij} ($i, j = 1, \dots, n$) are arbitrary numbers, the resolvent has the form

$$R(x, t; \lambda) = \frac{1}{\Delta(a_{ij} + \lambda b_{ij})} \begin{vmatrix} r(x, t; \lambda) & \varphi_1(x) & \cdots & \varphi_n(x) \\ \psi_1(t) & a_{11} + \lambda b_{11} & \cdots & a_{1n} + \lambda b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ \psi_n(t) & a_{n1} + \lambda b_{n1} & \cdots & a_{nn} + \lambda b_{nn} \end{vmatrix}, \quad (4)$$

where

$$\begin{aligned} \varphi_k(x) &= g_k(x) + \lambda \int_a^b r(x, t; \lambda) g_k(t) dt, & \psi_k(x) &= h_k(x) + \lambda \int_a^b r(x, t; \lambda) h_k(t) dt, \\ b_{ij} &= \int_a^b g_j(x) h_i(x) dx, & k, i, j &= 1, \dots, n. \end{aligned} \quad (5)$$

13.15-2. Some Special Cases.

Assume that

$$K(x, t) = k(x, t) - \sum_{k=1}^n g_k(x)h_k(t), \tag{6}$$

i.e., in formula (3) we have $a_{ij} = 0$ for $i \neq j$ and $a_{ii} = 1$. For this case, the resolvent is equal to

$$R(x, t; \lambda) = \frac{1}{\Delta_*} \begin{vmatrix} r(x, t; \lambda) & \varphi_1(x) & \cdots & \varphi_n(x) \\ \psi_1(t) & 1 + \lambda b_{11} & \cdots & \lambda b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ \psi_n(t) & \lambda b_{n1} & \cdots & 1 + \lambda b_{nn} \end{vmatrix}, \quad \Delta_* = \begin{vmatrix} 1 + \lambda b_{11} & \lambda b_{12} & \cdots & \lambda b_{1n} \\ \lambda b_{21} & 1 + \lambda b_{22} & \cdots & \lambda b_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \lambda b_{n1} & \lambda b_{n2} & \cdots & 1 + \lambda b_{nn} \end{vmatrix}. \tag{7}$$

Moreover, assume that $k(x, t) = 0$, i.e., the kernel $K(x, t)$ is degenerate:

$$K(x, t) = - \sum_{k=1}^n g_k(x)h_k(t). \tag{8}$$

In this case it is clear that $r(x, t; \lambda) = 0$ and, by virtue of (7),

$$\varphi_k(x) = g_k(x), \quad \psi_k(x) = h_k(x), \quad b_{ij} = \int_a^b g_j(x)h_i(x) dx.$$

Therefore, the resolvent becomes

$$R(x, t; \lambda) = \frac{1}{\Delta_*} \begin{vmatrix} 0 & g_1(x) & \cdots & g_n(x) \\ h_1(t) & 1 + \lambda b_{11} & \cdots & \lambda b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ h_n(t) & \lambda b_{n1} & \cdots & 1 + \lambda b_{nn} \end{vmatrix}. \tag{9}$$

Now we consider an integral equation with some kernel $Q(x, t)$. On the interval (a, b) we arbitrarily choose points x_1, \dots, x_n and t_1, \dots, t_n , and in relation (3) we set

$$k(x, t) = 0, \quad g_k(x) = Q(x, t_k), \quad h_k(t) = -Q(x_k, t), \quad a_{ij} = Q(x_i, t_j).$$

In this case it is clear that $r(x, t; \lambda) = 0$, and the kernel $K(x, t)$ acquires the form

$$K(x, t) = \frac{1}{D} \begin{vmatrix} 0 & Q(x, t_1) & \cdots & Q(x, t_n) \\ Q(x_1, t) & Q(x_1, t_1) & \cdots & Q(x_1, t_n) \\ \vdots & \vdots & \ddots & \vdots \\ Q(x_n, t) & Q(x_n, t_1) & \cdots & Q(x_n, t_n) \end{vmatrix}, \quad D = \begin{vmatrix} Q(x_1, t_1) & \cdots & Q(x_1, t_n) \\ \vdots & \ddots & \vdots \\ Q(x_n, t_1) & \cdots & Q(x_n, t_n) \end{vmatrix}.$$

It is convenient to rewrite this formula in the form

$$K(x, t) = Q(x, t) - \frac{1}{D} \begin{vmatrix} Q(x, t) & Q(x, t_1) & \cdots & Q(x, t_n) \\ Q(x_1, t) & Q(x_1, t_1) & \cdots & Q(x_1, t_n) \\ \vdots & \vdots & \ddots & \vdots \\ Q(x_n, t) & Q(x_n, t_1) & \cdots & Q(x_n, t_n) \end{vmatrix}. \tag{10}$$

The kernel $K(x, t)$ is degenerate and, moreover, it coincides with the kernel $Q(x, t)$ on the straight lines $x = x_i, t = t_j$ ($i, j = 1, \dots, n$). Indeed, if we set $x = x_i$ or $t = t_j$, then the determinant in the numerator of the second term has two equal rows or columns and hence vanishes, and therefore,

$$K(x_i, t) = Q(x_i, t), \quad K(x, t_j) = Q(x, t_j).$$

This coincidence on $2n$ straight lines permits us to expect that $K(x, t)$ is close to $Q(x, t)$ and the solution of the equation with kernel $K(x, t)$ is close to the solution of the equation with kernel $Q(x, t)$. It should be noted that if $Q(x, t)$ is degenerate, i.e., has the form

$$Q(x, t) = \sum_{k=1}^n g_k(x)h_k(t), \quad (11)$$

then the determinant in the numerator is identically zero, and hence in this case we have

$$K(x, t) \equiv Q(x, t). \quad (12)$$

For the kernel $K(x, t)$, the resolvent can be evaluated on the basis of the following relations:

$$\begin{aligned} r(x, t; \lambda) &= 0, \quad \varphi_i(x) = g_i(x) = Q(x, t_i), \quad \psi_j(t) = h_j(t) = -Q(x_j, t), \\ b_{ij} &= -\int_a^b Q(x, t_j)Q(x_i, x) dx = -Q_2(x_i, t_j), \quad i, j = 1, \dots, n, \end{aligned} \quad (13)$$

where $Q_2(x, t)$ is the second iterated kernel for $Q(x, t)$:

$$Q_2(x, y) = \int_a^b Q(x, s)Q(s, t) ds,$$

and hence

$$R(x, t; \lambda) = \frac{1}{D - \lambda D_2} \begin{vmatrix} 0 & Q(x, t_1) & \cdots & Q(x, t_n) \\ Q(x_1, t) & Q(x_1, t_1) - \lambda Q_2(x_1, t_1) & \cdots & Q(x_1, t_n) - \lambda Q_2(x_1, t_n) \\ \vdots & \vdots & \ddots & \vdots \\ Q(x_n, t) & Q(x_n, t_1) - \lambda Q_2(x_n, t_1) & \cdots & Q(x_n, t_n) - \lambda Q_2(x_n, t_n) \end{vmatrix}, \quad (14)$$

where

$$D_2 = \begin{vmatrix} Q_2(x_1, t_1) & \cdots & Q_2(x_1, t_n) \\ \vdots & \ddots & \vdots \\ Q_2(x_n, t_1) & \cdots & Q_2(x_n, t_n) \end{vmatrix}.$$

By using the resolvent $R(x, t; \lambda)$, we can obtain an approximate solution of the equation with kernel $Q(x, t)$. In particular, approximate characteristic values $\tilde{\lambda}$ of this kernel can be found by equating the determinant in the denominator of (14) with zero.

Example. Consider the equation

$$\begin{aligned} y(x) - \lambda \int_0^1 Q(x, t)y(t) dt &= 0, \quad 0 \leq x \leq 1, \\ Q(x, t) &= \begin{cases} x(t-1) & \text{for } x \leq t, \\ t(x-1) & \text{for } x \geq t. \end{cases} \end{aligned} \quad (15)$$

Let us find its characteristic values. To this end, we apply formula (14), where for the second iterated kernel we have

$$Q_2(x, t) = \int_0^1 Q(x, s)Q(s, t) ds = \begin{cases} \frac{1}{6}x(1-t)(2t-x^2-t^2) & \text{for } x \leq t, \\ \frac{1}{6}t(1-x)(2x-x^2-t^2) & \text{for } x \geq t. \end{cases}$$

We choose equidistant points x_i and t_j and take $n = 5$. This implies

$$x_1 = t_1 = \frac{1}{6}, \quad x_2 = t_2 = \frac{2}{6}, \quad x_3 = t_3 = \frac{3}{6}, \quad x_4 = t_4 = \frac{4}{6}, \quad x_5 = t_5 = \frac{5}{6}.$$

Let us equate the determinant in the denominator of (14) with zero. After some algebraic manipulations, we obtain the following equation:

$$130\mu^5 - 441\mu^4 + 488\mu^3 - 206\mu^2 + 30\mu - 1 = 0 \quad (\tilde{\lambda} = 216\mu),$$

which can be rewritten in the form

$$(\mu - 1)(2\mu - 1)(5\mu - 1)(13\mu^2 - 22\mu + 1) = 0. \quad (16)$$

On solving (16), we obtain

$$\bar{\lambda}_1 = 10.02, \quad \bar{\lambda}_2 = 43.2, \quad \bar{\lambda}_3 = 108, \quad \bar{\lambda}_4 = 216, \quad \bar{\lambda}_5 = 355.2.$$

The exact values of the characteristic values of the equation under consideration are known:

$$\lambda_1 = \pi^2 = 9.869 \dots, \quad \lambda_2 = (2\pi)^2 = 39.478 \dots, \quad \lambda_3 = (3\pi)^2 = 88.826 \dots,$$

and hence the calculation error is 2% for the first characteristic value, 9% for the second characteristic value, and 20% for the third characteristic value.

The result can be improved by choosing another collection of points x_i and t_i ($i = 1, \dots, 5$). However, for this number of ordinates we cannot have very high precision, because the kernel $Q(x, t)$ itself has a singularity, namely, its derivative is discontinuous for $x = t$, and thus the kernels under consideration cannot provide a good approximation of the given kernel.

⊙ References for Section 13.15: H. Bateman (1922), E. Goursat (1923), L. V. Kantorovich and V. I. Krylov (1958), P. K. Kythe and P. Puri (2002).

13.16. Collocation Method

13.16-1. General Remarks.

Let us rewrite the Fredholm integral equation of the second kind in the form

$$\varepsilon[y(x)] \equiv y(x) - \lambda \int_a^b K(x, t)y(t) dt - f(x) = 0. \quad (1)$$

Let us seek an approximate solution of Eq. (1) in the special form

$$Y_n(x) = \Phi(x, A_1, \dots, A_n) \quad (2)$$

with free parameters A_1, \dots, A_n (undetermined coefficients). On substituting the expression (2) into Eq. (1), we obtain the residual

$$\varepsilon[Y_n(x)] = Y_n(x) - \lambda \int_a^b K(x, t)Y_n(t) dt - f(x). \quad (3)$$

If $y(x)$ is an exact solution, then, clearly, the residual $\varepsilon[y(x)]$ is zero. Therefore, one tries to choose the parameters A_1, \dots, A_n so that, in a sense, the residual $\varepsilon[Y_n(x)]$ is as small as possible. The residual $\varepsilon[Y_n(x)]$ can be minimized in several ways. Usually, to simplify the calculations, a function $Y_n(x)$ linearly depending on the parameters A_1, \dots, A_n is taken. On finding the parameters A_1, \dots, A_n , we obtain an approximate solution (2). If

$$\lim_{n \rightarrow \infty} Y_n(x) = y(x), \quad (4)$$

then, by taking a sufficiently large number of parameters A_1, \dots, A_n , we find that the solution $y(x)$ can be found with an arbitrary prescribed precision.

Now let us go to the description of a concrete method of construction of an approximate solution $Y_n(x)$.

13.16-2. Approximate Solution.

We set

$$Y_n(x) = \varphi_0(x) + \sum_{i=1}^n A_i \varphi_i(x), \quad (5)$$

where $\varphi_0(x), \varphi_1(x), \dots, \varphi_n(x)$ are given functions (*coordinate functions*) and A_1, \dots, A_n are indeterminate coefficients, and assume that the functions $\varphi_i(x)$ ($i = 1, \dots, n$) are linearly independent. Note that, in particular, we can take $\varphi_0(x) = f(x)$ or $\varphi_0(x) \equiv 0$. On substituting the expression (5) into the left-hand side of Eq. (1), we obtain the residual

$$\varepsilon[Y_n(x)] = \varphi_0(x) + \sum_{i=1}^n A_i \varphi_i(x) - f(x) - \lambda \int_a^b K(x, t) \left[\varphi_0(t) + \sum_{i=1}^n A_i \varphi_i(t) \right] dt,$$

or

$$\varepsilon[Y_n(x)] = \psi_0(x, \lambda) + \sum_{i=1}^n A_i \psi_i(x, \lambda), \quad (6)$$

where

$$\begin{aligned} \psi_0(x, \lambda) &= \varphi_0(x) - f(x) - \lambda \int_a^b K(x, t) \varphi_0(t) dt, \\ \psi_i(x, \lambda) &= \varphi_i(x) - \lambda \int_a^b K(x, t) \varphi_i(t) dt, \quad i = 1, \dots, n. \end{aligned} \quad (7)$$

According to the collocation method, we require that the residual $\varepsilon[Y_n(x)]$ be zero at the given system of *the collocation points* x_1, \dots, x_n on the interval $[a, b]$, i.e., we set

$$\varepsilon[Y_n(x_j)] = 0, \quad j = 1, \dots, n,$$

where

$$a \leq x_1 < x_2 < \dots < x_{n-1} < x_n \leq b.$$

It is common practice to set $x_1 = a$ and $x_n = b$.

This, together with formula (6), implies the linear algebraic system

$$\sum_{i=1}^n A_i \psi_i(x_j, \lambda) = -\psi_0(x_j, \lambda), \quad j = 1, \dots, n, \quad (8)$$

for the coefficients A_1, \dots, A_n . If the determinant of system (8) is nonzero,

$$\det[\psi_i(x_j, \lambda)] = \begin{vmatrix} \psi_1(x_1, \lambda) & \psi_1(x_2, \lambda) & \cdots & \psi_1(x_n, \lambda) \\ \psi_2(x_1, \lambda) & \psi_2(x_2, \lambda) & \cdots & \psi_2(x_n, \lambda) \\ \vdots & \vdots & \ddots & \vdots \\ \psi_n(x_1, \lambda) & \psi_n(x_2, \lambda) & \cdots & \psi_n(x_n, \lambda) \end{vmatrix} \neq 0,$$

then system (8) uniquely determines the numbers A_1, \dots, A_n , and hence makes it possible to find the approximate solution $Y_n(x)$ by formula (5).

13.16-3. Eigenfunctions of the Equation.

On equating the determinant with zero, we obtain the relation

$$\det[\psi_i(x_j, \lambda)] = 0,$$

which, in general, enables us to find approximate values $\tilde{\lambda}_k$ ($k = 1, \dots, n$) for the characteristic values of the kernel $K(x, t)$.

If we set

$$f(x) \equiv 0, \quad \varphi_0(x) \equiv 0, \quad \lambda = \tilde{\lambda}_k,$$

then, instead of system (8), we obtain the homogeneous system

$$\sum_{i=1}^n \tilde{A}_i^{(k)} \psi_i(x_j, \tilde{\lambda}_k) = 0, \quad j = 1, \dots, n. \quad (9)$$

On finding nonzero solutions $\tilde{A}_i^{(k)}$ ($i = 1, \dots, n$) of system (9), we obtain approximate eigenfunctions for the kernel $K(x, t)$:

$$\tilde{Y}_n^{(k)}(x) = \sum_{i=1}^n \tilde{A}_i^{(k)} \varphi_i(x),$$

that correspond to its characteristic value $\lambda_k \approx \tilde{\lambda}_k$.

Example. Let us solve the equation

$$y(x) - \int_0^1 \frac{t^2 y(t)}{x^2 + t^2} dt = x \arctan \frac{1}{x} \quad (10)$$

by the collocation method.

We set

$$Y_2(x) = A_1 + A_2 x.$$

On substituting this expression into Eq. (10), we obtain the residual

$$\varepsilon[Y_2(x)] = -A_1 x \arctan \frac{1}{x} + A_2 \left[x - \frac{1}{2} + \frac{x^2}{2} \ln \left(1 + \frac{1}{x^2} \right) \right] - x \arctan \frac{1}{x}.$$

On choosing the collocation points $x_1 = 0$ and $x_2 = 1$ and taking into account the relations

$$\lim_{x \rightarrow 0} x \arctan \frac{1}{x} = 0, \quad \lim_{x \rightarrow 0} x^2 \ln \left(1 + \frac{1}{x^2} \right) = 0,$$

we obtain the following system for the coefficients A_1 and A_2 :

$$\begin{aligned} 0 \times A_1 - \frac{1}{2} A_2 &= 0, \\ -\frac{\pi}{4} A_1 + \frac{1}{2} (1 + \ln 2) A_2 &= \frac{\pi}{4}. \end{aligned}$$

This implies $A_2 = 0$ and $A_1 = -1$. Thus,

$$Y_2(x) = -1. \quad (11)$$

We can readily verify that the approximate solution (11) thus obtained is exact.

13.17. Method of Least Squares

13.17-1. Description of the Method.

By analogy with the collocation method, for the equation

$$\varepsilon[y(x)] \equiv y(x) - \lambda \int_a^b K(x, t)y(t) dt - f(x) = 0 \quad (1)$$

we set

$$Y_n(x) = \varphi_0(x) + \sum_{i=1}^n A_i \varphi_i(x), \quad (2)$$

where $\varphi_0(x), \varphi_1(x), \dots, \varphi_n(x)$ are given functions, A_1, \dots, A_n are indeterminate coefficients, and the $\varphi_i(x)$ ($i = 1, \dots, n$) are linearly independent.

On substituting (2) into the left-hand side of Eq. (1), we obtain the residual

$$\varepsilon[Y_n(x)] = \psi_0(x, \lambda) + \sum_{i=1}^n A_i \psi_i(x, \lambda), \quad (3)$$

where $\psi_0(x, \lambda)$ and the $\psi_i(x, \lambda)$ ($i = 1, \dots, n$) are defined by formulas (7) of Subsection 13.16-2.

According to the method of least squares, the coefficients A_i ($i = 1, \dots, n$) can be found from the condition for the minimum of the integral

$$I = \int_a^b \{\varepsilon[Y_n(x)]\}^2 dx = \int_a^b \left[\psi_0(x, \lambda) + \sum_{i=1}^n A_i \psi_i(x, \lambda) \right]^2 dx. \quad (4)$$

This requirement leads to the algebraic system of equations

$$\frac{\partial I}{\partial A_j} = 0, \quad j = 1, \dots, n, \quad (5)$$

and hence, on the basis of (4), by differentiating with respect to the parameters A_1, \dots, A_n under the integral sign, we obtain

$$\frac{1}{2} \frac{\partial I}{\partial A_j} = \int_a^b \psi_j(x, \lambda) \left[\psi_0(x, \lambda) + \sum_{i=1}^n A_i \psi_i(x, \lambda) \right] dx = 0, \quad j = 1, \dots, n. \quad (6)$$

Using the notation

$$c_{ij}(\lambda) = \int_a^b \psi_i(x, \lambda) \psi_j(x, \lambda) dx, \quad (7)$$

we can rewrite system (6) in the form of the *normal system of the method of least squares*:

$$\begin{aligned} c_{11}(\lambda)A_1 + c_{12}(\lambda)A_2 + \dots + c_{1n}(\lambda)A_n &= -c_{10}(\lambda), \\ c_{21}(\lambda)A_1 + c_{22}(\lambda)A_2 + \dots + c_{2n}(\lambda)A_n &= -c_{20}(\lambda), \\ \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \\ c_{n1}(\lambda)A_1 + c_{n2}(\lambda)A_2 + \dots + c_{nn}(\lambda)A_n &= -c_{n0}(\lambda). \end{aligned} \quad (8)$$

Note that if $\varphi_0(x) \equiv 0$, then $\psi_0(x) = -f(x)$. Moreover, since $c_{ij}(\lambda) = c_{ji}(\lambda)$, the matrix of system (8) is symmetric.

13.17-2. Construction of Eigenfunctions.

The method of least squares can also be applied for the approximate construction of characteristic values and eigenfunctions of the kernel $K(x, s)$, similarly to the way in which it can be done in the collocation method. Namely, by setting $f(x) \equiv 0$ and $\varphi_0(x) \equiv 0$, which implies $\psi_0(x) \equiv 0$, we determine approximate values of the characteristic values from the algebraic equation

$$\det[c_{ij}(\lambda)] = 0. \quad (9)$$

After this, approximate eigenfunctions can be found from the homogeneous system of the form (8), where, instead of λ , the corresponding approximate value is substituted.

Example. Let us find an approximate solution of the equation

$$y(x) = x^2 + \int_{-1}^1 \sinh(x+t)y(t) dt \quad (10)$$

by the method of least squares.

For the form of an approximate solution we take $Y_2(x) = x^2 + A_2x + A_1$. This implies

$$\varphi_1(x) = 1, \quad \varphi_2(x) = x, \quad \varphi_0(x) = x^2.$$

Taking into account the relations

$$\int_{-1}^1 \sinh(x+t) dt = a \sinh x, \quad \int_{-1}^1 t \sinh(x+t) dt = b \sinh x, \quad \int_{-1}^1 t^2 \sinh(x+t) dt = c \sinh x,$$

$$a = 2 \sinh 1 = 2.3504, \quad b = 2e^{-1} = 0.7358, \quad c = 6 \sinh 1 - 4 \cosh 1 = 0.8788,$$

on the basis of formulas (7) of Subsection 13.16-2 we have

$$\psi_1 = 1 - a \sinh x, \quad \psi_2 = x - b \cosh x, \quad \psi_0 = -c \sinh x.$$

Furthermore, we see that (to the fourth decimal place)

$$c_{11} = 2 + a^2 \left(\frac{1}{2} \sinh 2 - 1\right) = 6.4935, \quad c_{22} = \frac{2}{3} + b^2 \left(\frac{1}{2} \sinh 2 + 1\right) = 2.1896,$$

$$c_{12} = -4(ae^{-1} + b \sinh 1) = -8e^{-1} \sinh 1 = -3.4586, \quad c_{10} = ac \left(\frac{1}{2} \sinh 2 - 1\right) = 1.6800, \quad c_{20} = -2ce^{-1} = -0.6466,$$

and obtain the following system for the coefficients A_1 and A_2 :

$$6.4935A_1 - 3.4586A_2 = -1.6800,$$

$$-3.4586A_1 + 2.1896A_2 = 0.6466.$$

Hence, we have $A_1 = -0.5423$ and $A_2 = -0.5613$. Thus,

$$Y_2(x) = x^2 - 0.5613x - 0.5423. \quad (11)$$

Since the kernel

$$K(x, t) = \sinh(x+t) = \sinh x \cosh t + \cosh x \sinh t$$

of Eq. (10) is degenerate, we can readily obtain the exact solution

$$y(x) = x^2 + \alpha \sinh x + \beta \cosh x, \quad (12)$$

$$\alpha = \frac{6 \sinh 1 - 4 \cosh 1}{2 - \left(\frac{1}{2} \sinh 2\right)^2} = -0.6821, \quad \beta = \alpha \left(\frac{1}{2} \sinh 2 - 1\right) = -0.5548.$$

On comparing formulas (11) and (12) we conclude that the approximate solution $Y_2(x)$ is close to the exact solution $y(x)$ if $|x|$ is small. At the endpoints $x = \pm 1$, the discrepancy $|y(x) - Y_2(x)|$ is rather significant.

13.18. Bubnov–Galerkin Method

13.18-1. Description of the Method.

Let

$$\varepsilon[y(x)] \equiv y(x) - \lambda \int_a^b K(x, t)y(t) dt - f(x) = 0. \quad (1)$$

Similarly to the above reasoning, we seek an approximate solution of Eq. (1) in the form of a finite sum

$$Y_n(x) = f(x) + \sum_{i=1}^n A_i \varphi_i(x), \quad i = 1, \dots, n, \quad (2)$$

where the $\varphi_i(x)$ ($i = 1, \dots, n$) are some given linearly independent functions (*coordinate functions*) and A_1, \dots, A_n are indeterminate coefficients. On substituting the expression (2) into the left-hand side of Eq. (1), we obtain the residual

$$\varepsilon[Y_n(x)] = \sum_{j=1}^n A_j \left[\varphi_j(x) - \lambda \int_a^b K(x, t)\varphi_j(t) dt \right] - \lambda \int_a^b K(x, t)f(t) dt. \quad (3)$$

According to the Bubnov–Galerkin method, the coefficients A_i ($i = 1, \dots, n$) are defined from the condition that the residual is orthogonal to all coordinate functions $\varphi_1(x), \dots, \varphi_n(x)$. This gives the system of equations

$$\int_a^b \varepsilon[Y_n(x)]\varphi_i(x) dx = 0, \quad i = 1, \dots, n,$$

or, by virtue of (3),

$$\sum_{j=1}^n (\alpha_{ij} - \lambda\beta_{ij})A_j = \lambda\gamma_i, \quad i = 1, \dots, n, \quad (4)$$

where

$$\alpha_{ij} = \int_a^b \varphi_i(x)\varphi_j(x) dx, \quad \beta_{ij} = \int_a^b \int_a^b K(x, t)\varphi_i(x)\varphi_j(t) dt dx, \quad \gamma_i = \int_a^b \int_a^b K(x, t)\varphi_i(x)f(t) dt dx.$$

If the determinant of system (4)

$$D(\lambda) = \det[\alpha_{ij} - \lambda\beta_{ij}]$$

is nonzero, then this system uniquely determines the coefficients A_1, \dots, A_n . In this case, formula (2) gives an approximate solution of the integral equation (1).

13.18-2. Characteristic Values.

The equation $D(\lambda) = 0$ gives approximate characteristic values $\tilde{\lambda}_1, \dots, \tilde{\lambda}_n$ of the integral equation. On finding nonzero solutions of the homogeneous linear system

$$\sum_{j=1}^n (\alpha_{ij} - \tilde{\lambda}_k\beta_{ij})\tilde{A}_j^{(k)} = 0, \quad i = 1, \dots, n,$$

we can construct approximate eigenfunctions $\tilde{Y}_n^{(k)}(x)$ corresponding to characteristic values $\tilde{\lambda}_k$:

$$\tilde{Y}_n^{(k)}(x) = \sum_{i=1}^n \tilde{A}_i^{(k)} \varphi_i(x).$$

It can be shown that the Bubnov–Galerkin method is equivalent to the replacement of the kernel $K(x, t)$ by some degenerate kernel $K_{(n)}(x, t)$. Therefore, for the approximate solution $Y_n(x)$ we have an error estimate similar to that presented in Subsection 13.14-2.

Example. Let us find the first two characteristic values of the integral equation

$$\varepsilon[y(x)] \equiv y(x) - \lambda \int_0^1 K(x, t)y(t) dt = 0,$$

where

$$K(x, t) = \begin{cases} t & \text{for } t \leq x, \\ x & \text{for } t > x. \end{cases} \quad (5)$$

On the basis of (5), we have

$$\varepsilon[y(x)] = y(x) - \lambda \left\{ \int_0^x ty(t) dt + \int_x^1 xy(t) dt \right\}.$$

We set $Y_2(x) = A_1x + A_2x^2$. In this case

$$\begin{aligned} \varepsilon[Y_2(x)] &= A_1x + A_2x^2 - \lambda \left[\frac{1}{3}A_1x^3 + \frac{1}{4}A_2x^4 + x \left(\frac{1}{2}A_1 + \frac{1}{3}A_2 \right) - \left(\frac{1}{2}A_1x^3 + \frac{1}{3}A_2x^4 \right) \right] = \\ &= A_1 \left[\left(1 - \frac{1}{2}\lambda \right) x + \frac{1}{6}\lambda x^3 \right] + A_2 \left[-\frac{1}{3}\lambda x + x^2 + \frac{1}{12}\lambda x^4 \right]. \end{aligned}$$

On orthogonalizing the residual $\varepsilon[Y_2(x)]$, we obtain the system

$$\begin{aligned} \int_0^1 \varepsilon[Y_2(x)]x dx &= 0, \\ \int_0^1 \varepsilon[Y_2(x)]x^2 dx &= 0, \end{aligned}$$

or the following homogeneous system of two algebraic equations with two unknowns:

$$\begin{aligned} A_1(120 - 48\lambda) + A_2(90 - 35\lambda) &= 0 \\ A_1(630 - 245\lambda) + A_2(504 - 180\lambda) &= 0. \end{aligned} \quad (6)$$

On equating the determinant of system (6) with zero, we obtain the following equation for the characteristic values:

$$D(\lambda) \equiv \begin{vmatrix} 120 - 48\lambda & 90 - 35\lambda \\ 630 - 245\lambda & 504 - 180\lambda \end{vmatrix} = 0.$$

Hence,

$$\lambda^2 - 26.03\lambda + 58.15 = 0. \quad (7)$$

Equations (7) imply

$$\tilde{\lambda}_1 = 2.462 \dots \quad \text{and} \quad \tilde{\lambda}_2 = 23.568 \dots$$

For comparison we present the exact characteristic values:

$$\lambda_1 = \frac{1}{4}\pi^2 = 2.467 \dots \quad \text{and} \quad \lambda_2 = \frac{9}{4}\pi^2 = 22.206 \dots,$$

which can be obtained from the solution of the following boundary value problem equivalent to the original equation:

$$y''_{xx}(x) + \lambda y(x) = 0; \quad y(0) = 0, \quad y'_x(1) = 0.$$

Thus, the error of $\tilde{\lambda}_1$ is approximately equal to 0.2% and that of $\tilde{\lambda}_2$, to 6%.

© References for Section 13.18: L. V. Kantorovich and V. I. Krylov (1958), B. P. Demidovich, I. A. Maron, and E. Z. Shulvalova (1963), A. F. Verlan' and V. S. Sizikov (1986), K. E. Atkinson (1997), R. Kress (1999).

13.19. Quadrature Method

13.19-1. General Scheme for Fredholm Equations of the Second Kind.

In the solution of an integral equation, the reduction to the solution of systems of algebraic equations obtained by replacing the integrals with finite sums is one of the most effective tools. The method of quadratures is related to the approximation methods. It is widespread in practice because it is rather universal with respect to the principle of constructing algorithms for solving both linear and nonlinear equations.

Just as in the case of Volterra equations, the method is based on a quadrature formula (see Subsection 10.7-1):

$$\int_a^b \varphi(x) dx = \sum_{j=1}^n A_j \varphi(x_j) + \varepsilon_n[\varphi], \quad (1)$$

where the x_j are the nodes of the quadrature formula, the A_j are given coefficients that do not depend on the function $\varphi(x)$, and $\varepsilon_n[\varphi]$ is the error of replacement of the integral by the sum (the truncation error).

If in the Fredholm integral equation of the second kind,

$$y(x) - \lambda \int_a^b K(x, t)y(t) dt = f(x), \quad a \leq x \leq b, \quad (2)$$

we set $x = x_i$ ($i = 1, \dots, n$), then we obtain the following relation that is the basic formula for the method under consideration:

$$y(x_i) - \lambda \int_a^b K(x_i, t)y(t) dt = f(x_i), \quad i = 1, \dots, n. \quad (3)$$

Applying the quadrature formula (1) to the integral in (3), we arrive at the following system of equations:

$$y(x_i) - \lambda \sum_{j=1}^n A_j K(x_i, x_j)y(x_j) = f(x_i) + \lambda \varepsilon_n[y]. \quad (4)$$

By neglecting the small term $\lambda \varepsilon_n[y]$ in this formula, we obtain the system of linear algebraic equations for approximate values y_i of the solution $y(x)$ at the nodes x_1, \dots, x_n :

$$y_i - \lambda \sum_{j=1}^n A_j K_{ij}y_j = f_i, \quad i = 1, \dots, n, \quad (5)$$

where $K_{ij} = K(x_i, x_j)$, $f_i = f(x_i)$.

The solution of system (5) gives the values y_1, \dots, y_n , which determine an approximate solution of the integral equation (2) on the entire interval $[a, b]$ by interpolation. Here for the approximate solution we can take the function obtained by linear interpolation, i.e., the function that coincides with y_i at the points x_i and is linear on each of the intervals $[x_i, x_{i+1}]$. Moreover, for an analytic expression of the approximate solution to the equation, a function

$$\tilde{y}(x) = f(x) + \lambda \sum_{j=1}^n A_j K(x, x_j)y_j \quad (6)$$

can be chosen, which also takes the values y_1, \dots, y_n at the points x_1, \dots, x_n .

13.19-2. Construction of the Eigenfunctions.

The method of quadratures can also be applied for solutions of homogeneous Fredholm equations of the second kind. In this case, system (5) becomes homogeneous ($f_i = 0$) and has a nontrivial solution only if its determinant $D(\lambda)$ is equal to zero. The algebraic equation $D(\lambda) = 0$ of degree n for λ makes it possible to find the roots $\tilde{\lambda}_1, \dots, \tilde{\lambda}_n$, which are approximate values of n characteristic values of the equation. The substitution of each value $\tilde{\lambda}_k$ ($k = 1, \dots, n$) into (5) for $f_i \equiv 0$ leads to the system of equations

$$y_i^{(k)} - \tilde{\lambda}_k \sum_{j=1}^n A_j K_{ij}y_j^{(k)} = 0, \quad i = 1, \dots, n,$$

whose nonzero solutions $y_i^{(k)}$ make it possible to obtain approximate expressions for the eigenfunctions of the integral equation:

$$\tilde{y}_k(x) = \tilde{\lambda}_k \sum_{j=1}^n A_j K(x, x_j) y_j^{(k)}.$$

If λ differs from each of the roots $\tilde{\lambda}_k$, then the nonhomogeneous system of linear algebraic equations (5) has a unique solution. In the same case, the homogeneous system of equations (5) has only the trivial solution.

13.19-3. Specific Features of the Application of Quadrature Formulas.

The accuracy of the resulting solutions essentially depends on the smoothness of the kernel and the constant term. When choosing the quadrature formula, it is necessary to take into account that the more accurate an applied formula is, the more serious requirements must be imposed on the smoothness of the kernel, the solution, and the right-hand side.

If the right-hand side or the kernel have singularities, then it is reasonable to perform a preliminary transform of the original equation to obtain a more accurate approximate solution. Here the following methods can be applied.

If the right-hand side $f(x)$ has singularities and the kernel is smooth, then we can introduce the new unknown function $z(x) = y(x) - f(x)$ instead of $y(x)$, and the substitution of $z(x)$ in the original equation leads to the equation

$$z(x) - \lambda \int_a^b K(x, t)z(t) dt = \lambda \int_a^b K(x, t)f(t) dt,$$

in which the right-hand side is smoothed, and hence a solution $z(x)$ is smoother. From the function $z(x)$ thus obtained we can readily find the desired solution $y(x)$.

For the cases in which the kernel $K(x, t)$ or its derivatives with respect to t have discontinuities on the diagonal $x = t$, it is useful to rewrite the equation under consideration in the equivalent form

$$y(x) \left[1 - \lambda \int_a^b K(x, t) dt \right] - \lambda \int_a^b K(x, t)[y(t) - y(x)] dt = f(x),$$

where the integrand in the second integral has no singularities because the difference $y(t) - y(x)$ vanishes on the diagonal $x = t$, and the calculation of the integral $\int_a^b K(x, t) dt$ is performed without unknown functions and is possible in the explicit form.

Example. Consider the equation

$$y(x) - \frac{1}{2} \int_0^1 xty(t) dt = \frac{5}{6}x.$$

Let us choose the nodes $x_1 = 0$, $x_2 = \frac{1}{2}$, $x_3 = 1$ and calculate the values of the right-hand side $f(x) = \frac{5}{6}x$ and of the kernel $K(x, t) = xt$ at these nodes:

$$\begin{aligned} f(0) &= 0, & f\left(\frac{1}{2}\right) &= \frac{5}{12}, & f(1) &= \frac{5}{6}, \\ K(0, 0) &= 0, & K\left(0, \frac{1}{2}\right) &= 0, & K(0, 1) &= 0, & K\left(\frac{1}{2}, 0\right) &= 0, & K\left(\frac{1}{2}, \frac{1}{2}\right) &= \frac{1}{4}, \\ K\left(\frac{1}{2}, 1\right) &= \frac{1}{2}, & K(1, 0) &= 0, & K\left(1, \frac{1}{2}\right) &= \frac{1}{2}, & K(1, 1) &= 1. \end{aligned}$$

On applying Simpson's rule (see Subsection 10.7-1)

$$\int_0^1 F(x) dx \approx \frac{1}{6} \left[F(0) + 4F\left(\frac{1}{2}\right) + F(1) \right]$$

to determine the approximate values y_i ($i = 1, 2, 3$) of the solution $y(x)$ at the nodes x_i we obtain the system

$$\begin{aligned} y_1 &= 0, \\ \frac{11}{12}y_2 - \frac{1}{24}y_3 &= \frac{5}{12}, \\ -\frac{2}{12}y_2 + \frac{11}{12}y_3 &= \frac{5}{6}, \end{aligned}$$

whose solution is $y_1 = 0$, $y_2 = \frac{1}{2}$, $y_3 = 1$. In accordance with the expression (6), the approximate solution can be presented in the form

$$\tilde{y}(x) = \frac{5}{6}x + \frac{1}{2} \times \frac{1}{6} \left(0 + 4 \times \frac{1}{2} \times \frac{1}{2}x + 1 \times 1 \times x \right) = x.$$

We can readily verify that it coincides with the exact solution.

⊙ References for Section 13.19: N. S. Bakhvalov (1973), V. I. Krylov, V. V. Bobkov, and P. I. Monastyrnyi (1984), A. F. Verlan' and V. S. Sizikov (1986).

13.20. Systems of Fredholm Integral Equations of the Second Kind

13.20-1. Some Remarks.

A system of Fredholm integral equations of the second kind has the form

$$y_i(x) - \lambda \sum_{j=1}^n \int_a^b K_{ij}(x, t) y_j(t) dt = f_i(x), \quad a \leq x \leq b, \quad i = 1, \dots, n. \quad (1)$$

Assume that the kernels $K_{ij}(x, t)$ are continuous or square integrable on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ and the right-hand sides $f_i(x)$ are continuous or square integrable on $[a, b]$. We also assume that the functions $y_i(x)$ to be defined are continuous or square integrable on $[a, b]$ as well. The theory developed above for Fredholm equations of the second kind can be completely extended to such systems. In particular, it can be shown that for systems (1), the successive approximations converge in mean-square to the solution of the system if λ satisfies the inequality

$$|\lambda| < \frac{1}{B_*}, \quad (2)$$

where

$$\sum_{i=1}^n \sum_{j=1}^n \int_a^b \int_a^b |K_{ij}(x, t)|^2 dx dt = B_*^2 < \infty. \quad (3)$$

If the kernel $K_{ij}(x, t)$ satisfies the additional condition

$$\int_a^b K_{ij}^2(x, t) dt \leq A_{ij}, \quad a \leq x \leq b, \quad (4)$$

where A_{ij} are some constants, then the successive approximations converge absolutely and uniformly.

If all kernels $K_{ij}(x, t)$ are degenerate, then system (1) can be reduced to a linear algebraic system. It can be established that for a system of Fredholm integral equations, all Fredholm theorems are satisfied.

13.20-2. Method of Reducing a System of Equations to a Single Equation.

System (1) can be transformed into a single Fredholm integral equation of the second kind. Indeed, let us introduce the functions $Y(x)$ and $F(x)$ on $[a, nb - (n-1)a]$ by setting

$$Y(x) = y_i(x - (i-1)(b-a)), \quad F(x) = f_i(x - (i-1)(b-a)),$$

for

$$(i-1)b - (i-2)a \leq x \leq ib - (i-1)a.$$

Let us define a kernel $K(x, t)$ on the square $\{a \leq x \leq nb - (n-1)a, a \leq t \leq nb - (n-1)a\}$ as follows:

$$K(x, t) = K_{ij}(x - (i-1)(b-a), t - (j-1)(b-a))$$

for

$$(i-1)b - (i-2)a \leq x \leq ib - (i-1)a, \quad (j-1)b - (j-2)a \leq t \leq jb - (j-1)a.$$

Now system (1) can be rewritten as the single Fredholm equation

$$Y(x) - \lambda \int_a^{nb-(n-1)a} K(x, t)Y(t) dt = F(x), \quad a \leq x \leq nb - (n-1)a.$$

If the kernels $K_{ij}(x, t)$ are square integrable on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ and the right-hand sides $f_i(x)$ are square integrable on $[a, b]$, then the kernel $K(x, t)$ is square integrable on the new square

$$S_n = \{a < x < nb - (n-1)a, a < t < nb - (n-1)a\},$$

and the right-hand side $F(x)$ is square integrable on $[a, nb - (n-1)a]$.

If condition (4) is satisfied, then the kernel $K(x, t)$ satisfies the inequality

$$\int_a^b K^2(x, t) dt \leq A_*, \quad a < x < nb - (n-1)a,$$

where A_* is a constant.

⊙ Reference for Section 13.20: S. G. Mikhlin (1960).

13.21. Regularization Method for Equations with Infinite Limits of Integration

13.21-1. Basic Equation and Fredholm Theorems.

Consider an integral equation of the second kind in the form

$$y(x) + \frac{1}{\sqrt{2\pi}} \int_0^\infty K_1(x-t)y(t) dt + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 K_2(x-t)y(t) dt + \int_{-\infty}^\infty M(x, t)y(t) dt = f(x), \quad (1)$$

where $-\infty < x < \infty$. We assume that the functions $y(x)$ and $f(x)$ and the kernels $K_1(x)$ and $K_2(x)$ are such that their Fourier transforms belong to $L_2(-\infty, \infty)$ and satisfy the Hölder condition. We also assume that the Fourier transforms of the kernel $M(x, t)$ with respect to each variable belong to $L_2(-\infty, \infty)$ and satisfy the Hölder condition and, in addition,

$$\int_{-\infty}^\infty \int_{-\infty}^\infty |M(x, t)|^2 dx dt < \infty.$$

It should be noted that Eq. (1) with $M(x, t) \equiv 0$ is the convolution-type integral equation with two kernels which was discussed in Subsection 13.10-2.

The transposed homogeneous equation has the form

$$\varphi(x) + \frac{1}{\sqrt{2\pi}} \int_0^\infty K_1(t-x)\varphi(t) dt + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 K_2(t-x)\varphi(t) dt + \int_{-\infty}^\infty M(t, x)\varphi(t) dt = 0, \quad (2)$$

where $-\infty < x < \infty$.

Assume that the normality conditions (see Subsection 13.10-2) hold, that is,

$$1 + \mathcal{K}_1(u) \neq 0, \quad 1 + \mathcal{K}_2(u) \neq 0, \quad -\infty < u < \infty. \quad (3)$$

THEOREM 1. *The number of linearly independent solutions of the homogeneous ($f(x) \equiv 0$) equation (1) and that of the transposed homogeneous ($g(x) \equiv 0$) equation (2) are finite.*

THEOREM 2. *For the nonhomogeneous equation (1) to be solvable, it is necessary and sufficient that*

$$\int_{-\infty}^\infty f(t)\varphi_k(t) dt = 0, \quad k = 1, \dots, N, \quad (4)$$

where $\varphi_k(x)$ is a complete finite set of linearly independent solutions to the transposed homogeneous equation (2).

THEOREM 3. *The difference between the number of linearly independent solutions to the homogeneous equation (1) and the number of linearly independent solutions to the homogeneous transposed equation (2) is equal to the index*

$$\nu = \text{Ind} \frac{1 + \mathcal{K}_2(u)}{1 + \mathcal{K}_1(u)} = \frac{1}{2\pi} \left[\arg \frac{1 + \mathcal{K}_2(u)}{1 + \mathcal{K}_1(u)} \right]_{-\infty}^\infty. \quad (5)$$

13.21-2. Regularizing Operators.

An important method for the theoretical investigation and practical solution of the integral equations in question is a regularization of these equations, i.e., their reduction to a Fredholm equation of the second kind.

Let us denote by \mathbf{K} the operator determined by the left-hand side of Eq. (1):

$$\mathbf{K}[y(x)] \equiv y(x) + \frac{1}{\sqrt{2\pi}} \int_0^\infty K_1(x-t)y(t) dt + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 K_2(x-t)y(t) dt + \int_{-\infty}^\infty M(x, t)y(t) dt \quad (6)$$

and introduce the similar operator

$$\mathbf{L}[\omega(x)] \equiv \omega(x) + \frac{1}{\sqrt{2\pi}} \int_0^\infty L_1(x-t)\omega(t) dt + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 L_2(x-t)\omega(t) dt + \int_{-\infty}^\infty Q(x, t)\omega(t) dt. \quad (7)$$

Let us find an operator \mathbf{L} such that the product \mathbf{LK} is determined by the left-hand side of a Fredholm equation of the second kind with a kernel $K(x, t)$:

$$\mathbf{LK}[y(x)] \equiv y(x) + \int_{-\infty}^\infty K(x, t)y(t) dt, \quad \int_{-\infty}^\infty \int_{-\infty}^\infty |K(x, t)|^2 dx dt < \infty. \quad (8)$$

The operator \mathbf{L} is called a *left regularizer*.

For the operator \mathbf{K} of the integral equation (1) to have a left regularizer \mathbf{L} of the form (7), it is necessary and sufficient that the normality conditions (3) hold.

If conditions (3) are satisfied, then the left regularizer \mathbf{L} has the form

$$\mathbf{L}\omega(x) \equiv \omega(x) - \frac{1}{\sqrt{2\pi}} \int_0^\infty R_1(x-t)\omega(t) dt - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 R_2(x-t)\omega(t) dt + \int_{-\infty}^\infty Q(x, t)\omega(t) dt, \quad (9)$$

where the resolvents $R_1(x-t)$ and $R_2(x-t)$ of the kernels $K_1(x-t)$ and $K_2(x-t)$ are given by (see Subsection 13.9-1)

$$R_j(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \frac{\mathcal{K}_j(u)}{1 + \mathcal{K}_j(u)} e^{-iux} du, \quad \mathcal{K}_j(u) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty K_j(x) e^{iux} dx, \quad j = 1, 2,$$

and $Q(x, t)$ is any function such that

$$\int_{-\infty}^\infty \int_{-\infty}^\infty |Q(x, t)|^2 dx dt < \infty.$$

If condition (3) is satisfied, then the operator \mathbf{L} given by formula (9) is simultaneously a *right regularizer* of the operator \mathbf{K} :

$$\mathbf{KL}[y(x)] \equiv y(x) + \int_{-\infty}^\infty K_*(x, t)y(t) dt, \quad (10)$$

where the function $K_*(x, t)$ satisfies the condition

$$\int_{-\infty}^\infty \int_{-\infty}^\infty |K_*(x, t)|^2 dx dt < \infty. \quad (11)$$

13.21-3. Regularization Method.

Consider the equation of the form

$$\mathbf{K}[y(x)] = f(x), \quad -\infty < x < \infty, \quad (12)$$

where the operator \mathbf{K} is defined by (6).

There are several ways of regularizing this equation, i.e., of its reduction to a Fredholm equation. First, this equation can be reduced to an equation with a Cauchy kernel. On regularizing the last equation by a method presented in Section 15.4, we can achieve our aim. This approach can be applied if we can find, for given functions $K_1(x)$, $K_2(x)$, $M(x, t)$, and $f(x)$, simple expressions for their Fourier integrals. Otherwise it is natural to perform the regularization of Eq. (12) directly, without passing to the inverse transforms.

A left regularization of Eq. (12) involves the application of the regularizer \mathbf{L} constructed in the previous subsection to both its sides:

$$\mathbf{LK}[y(x)] = \mathbf{L}[f(x)]. \quad (13)$$

It follows from (8) that Eq. (13) is a Fredholm equation

$$y(x) + \int_{-\infty}^\infty K(x, t)y(t) dt = \mathbf{L}[f(x)]. \quad (14)$$

Thus, Eq. (12) can be transformed by left regularization to a Fredholm equation with the same unknown function $y(x)$ and the known right-hand side $\mathbf{L}[f(x)]$. Left regularization is known to imply no loss of solutions: all solutions of the original equation (12) are solutions of the regularized equation. However, in the general case, a solution of the regularized equation need not be a solution of the original equation.

The right regularization consists in the substitution of the expression

$$y(x) = \mathbf{L}[\omega(x)] \quad (15)$$

for the desired function into Eq. (12), where $\omega(x)$ is a new unknown function. We finally arrive at the following integral equation:

$$\mathbf{KL}[\omega(x)] = f(x), \quad (16)$$

which is a Fredholm equation as well by virtue of (10):

$$\mathbf{KL}[\omega(x)] \equiv \omega(x) + \int_{-\infty}^{\infty} K_*(x, t)\omega(t) dt = f(x), \quad -\infty < x < \infty. \quad (17)$$

Thus, we have passed from Eq. (12) for the unknown function $y(x)$ to a Fredholm integral equation for a new unknown function $\omega(x)$. On solving the Fredholm equation (17), we find a solution of the original equation (12) by formula (15). Right regularization can give no irrelevant solutions, but it is known that it can lead to a loss of a solution.

A solution of the problem on an equivalent regularization, for which neither the loss of solutions nor the appearance of irrelevant “solutions” occur, is of significant theoretical and practical interest.

For Eq. (12) with an arbitrary right-hand side $f(x)$ to admit an equivalent left regularization, it is necessary and sufficient that the index ν given by formula (5) be nonnegative. For an equivalently regularizing operator we can take the operator

$$\mathbf{L}^\circ[\omega(x)] \equiv \omega(x) - \frac{1}{\sqrt{2\pi}} \int_0^\infty R_1(x-t)\omega(t) dt - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 R_2(x-t)\omega(t) dt.$$

Thus, the Fredholm equation

$$\mathbf{L}^\circ \mathbf{K}[y(x)] = \mathbf{L}^\circ[f(x)], \quad (18)$$

for the case $\nu \geq 0$, has those and only those solutions that are solutions to Eq. (12).

For the case in which the index ν is nonpositive, the operator \mathbf{L}° performs an equivalent right regularization of Eq. (12) for an arbitrary right-hand side $f(x)$. In other words, for $\nu \leq 0$, on finding the solution to the Fredholm equation

$$\mathbf{KL}^\circ[\omega(x)] = f(x),$$

we can obtain all solutions of the original equation (12) by the formula $y(x) = \mathbf{L}^\circ[\omega(x)]$.

Another method of regularization is known, the so-called Carleman–Vekua regularization, which is based on the solution of the corresponding characteristic equation. Equation (12) can formally be rewritten as a convolution type equation with two kernels:

$$y(x) + \frac{1}{\sqrt{2\pi}} \int_0^\infty K_1(x-t)y(t) dt + \frac{1}{\sqrt{2\pi}} \int_{-\infty}^0 K_2(x-t)y(t) dt = f_1(x), \quad (19)$$

where

$$f_1(x) = f(x) - \int_{-\infty}^{\infty} M(x, t)y(t) dt.$$

Next, the function $f_1(x)$ is provisionally assumed to be known, and Eq. (19) is solved (see Subsection 13.10-2). The analysis of the resulting formula for the function $y(x)$ shows that, for $\nu = 0$, this is a Fredholm integral equation with the unknown function $y(x)$. For the case in which $\nu > 0$, the resulting equation contains ν arbitrary constants. For a negative index ν , solvability conditions must be added to the equation.

Chapter 14

Methods for Solving Singular Integral Equations of the First Kind

14.1. Some Definitions and Remarks

14.1-1. Integral Equations of the First Kind with Cauchy Kernel.

A *singular integral equation of the first kind with Cauchy kernel* has the form

$$\frac{1}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau = f(t), \quad i^2 = -1, \quad (1)$$

where L is a smooth closed or nonclosed contour in the complex plane of the variable $z = x + iy$, t and τ are the complex coordinates on L , $\varphi(t)$ is the unknown function, $\frac{1}{\tau - t}$ is the Cauchy kernel, and $f(t)$ is a given function, which is called the right-hand side of Eq. (1). The integral on the left-hand side only exists in the sense of the Cauchy principal value (see Subsection 14.2-5).

A singular integral equation in which L is a smooth closed contour, as well as an equation of the form

$$\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\varphi(t)}{t - x} dt = f(x), \quad -\infty < x < \infty, \quad (2)$$

on the real axis and an equation with Cauchy kernel

$$\frac{1}{\pi} \int_a^b \frac{\varphi(t)}{t - x} dt = f(x), \quad a \leq x \leq b, \quad (3)$$

on a finite interval, are special cases of Eq. (1).

A general singular integral equation of the first kind with Cauchy kernel has the form

$$\frac{1}{\pi i} \int_L \frac{M(t, \tau)}{\tau - t} \varphi(\tau) d\tau = f(t), \quad (4)$$

where $M(t, \tau)$ is a given function. This equation can also be rewritten in a different (equivalent) form, which is given in Subsection 14.4-4.

Assume that all functions in Eqs. (1)–(4) satisfy the Hölder condition (Subsection 14.2-2) and the function $M(t, \tau)$ satisfies this condition with respect to both variables.

14.1-2. Integral Equations of the First Kind with Hilbert Kernel.

The simplest *singular integral equation of the first kind with Hilbert kernel* has the form

$$\frac{1}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi - x}{2}\right) \varphi(\xi) d\xi = f(x), \quad (5)$$

where $\varphi(x)$ is the unknown function ($0 \leq x \leq 2\pi$), $\cot[\frac{1}{2}(\xi - x)]$ is the Hilbert kernel, and $f(x)$ is the given right-hand side of the equation ($0 \leq x \leq 2\pi$).

A general singular integral equation of the first kind with Hilbert kernel has the form

$$-\frac{1}{2\pi} \int_0^{2\pi} N(x, \xi) \cot\left(\frac{\xi - x}{2}\right) \varphi(\xi) d\xi = f(x), \quad (6)$$

where $N(x, \xi)$ is a given function. Equation (6) can often be rewritten in an equivalent form, which is presented in Subsection 14.4-5.

Assume that all functions in Eqs. (5) and (6) also satisfy the Hölder condition (see Subsection 14.2-2) and the function $N(x, \xi)$ satisfies this condition with respect to both variables.

If the right-hand sides of Eqs. (1)–(6) are identically zero, then the equations are said to be *homogeneous*, otherwise they are said to be *nonhomogeneous*.

© References for Section 14.1: F. D. Gakhov (1977), S. G. Mikhlín and S. Prössdorf (1986), S. Prössdorf and B. Silbermann (1991), A. Dzuraev (1992), N. I. Muskhelishvili (1992), I. K. Lifanov (1996), R. Estrada and R. P. Kanwal (1999), E. G. Ladopoulos (2000).

14.2. Cauchy Type Integral

14.2-1. Definition of the Cauchy Type Integral.

Let L be a smooth closed contour* on the plane of a complex variable $z = x + iy$. The domain inside the contour L is called the *interior domain* and is denoted by Ω^+ , and the complement of $\Omega^+ \cup L$, which contains the point at infinity, is called the *exterior domain* and is denoted by Ω^- .

If a function $f(z)$ is analytic in Ω^+ and continuous in $\Omega^+ \cup L$, then according to the familiar Cauchy formula in the theory of functions of a complex variable we have

$$\frac{1}{2\pi i} \int_L \frac{f(\tau)}{\tau - z} d\tau = \begin{cases} f(z) & \text{for } z \in \Omega^+, \\ 0 & \text{for } z \in \Omega^-. \end{cases} \quad (1)$$

If a function $f(z)$ is analytic in Ω^- and continuous in $\Omega^- \cup L$, then

$$\frac{1}{2\pi i} \int_L \frac{f(\tau)}{\tau - z} d\tau = \begin{cases} f(\infty) & \text{for } z \in \Omega^+, \\ -f(z) + f(\infty) & \text{for } z \in \Omega^-. \end{cases} \quad (2)$$

As usual, the positive direction on L is defined as the direction for which the domain Ω^+ remains to the left of the contour.

The Cauchy formula permits one to calculate the values of a function at any point of the domain provided that the values on the boundary of the domain are known, i.e., the Cauchy formula solves the boundary value problem for analytic functions. The integral on the left-hand side in (1) and (2) is called the *Cauchy integral*.

Assume that L is a smooth closed or nonclosed contour that entirely belongs to the finite part of the complex plane. Let τ be the complex coordinate on L , and let $\varphi(\tau)$ be a continuous function of a point of the contour. In this case the integral

$$\Phi(z) = \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - z} d\tau, \quad (3)$$

which is constructed in the same way as the Cauchy integral, is called a *Cauchy type integral*. The function $\varphi(\tau)$ is called its *density* and $1/(\tau - z)$ its *kernel*.

* By a *smooth contour* we mean a simple curve (i.e., a curve without points of self-intersection) that is either closed or nonclosed, has a continuous tangent, and has no cuspidal points.

For a Cauchy type integral with continuous density $\varphi(\tau)$, the only points at which the integrand is not analytic with respect to z are the points of the integration curve L . This curve is singular for the function $\Phi(z)$.

If L is a nonclosed contour, then $\Phi(z)$ is an analytic function on the entire plane with the singularity curve L . Assume that L is a closed contour. In this case, $\Phi(z)$ splits into two independent functions: a function $\Phi^+(z)$ defined on the domain Ω^+ and a function $\Phi^-(z)$ defined on the domain Ω^- . In general, these functions are not analytic continuations of each other.

By a *piecewise analytic function* we mean an analytic function $\Phi(z)$ defined by two independent expressions $\Phi^+(z)$ and $\Phi^-(z)$ on two complementary domains Ω^+ and Ω^- of the complex plane.

We note an important property of a Cauchy type integral. The function $\Phi(z)$ expressed by a Cauchy type integral of the form (3) vanishes at infinity, i.e., $\Phi^-(\infty) = 0$. This condition is also sufficient for the representability of a piecewise analytic function by a Cauchy type integral.

14.2-2. Hölder Condition.

Let L be a smooth curve in the complex plane $z = x + iy$, and let $\varphi(t)$ be a function on this curve. We say that $\varphi(t)$ satisfies the *Hölder condition* on L if for any two points $t_1, t_2 \in L$ we have

$$|\varphi(t_2) - \varphi(t_1)| < A|t_2 - t_1|^\lambda, \quad (4)$$

where A and λ are positive constants. The number A is called the *Hölder constant* and λ is called the *Hölder exponent*. If $\lambda > 1$, then by condition (4) the derivative $\varphi'_t(t)$ vanishes everywhere, and $\varphi(t)$ must be constant. Therefore, we assume that $0 < \lambda \leq 1$. For $\lambda = 1$, the Hölder condition is often called the *Lipschitz condition*. Sometimes the Hölder condition is called the *Lipschitz condition of order λ* .

If t_1 and t_2 are sufficiently close to each other and if the Hölder condition holds for some exponent λ_1 , then this condition certainly holds for each exponent $\lambda < \lambda_1$. In general, the converse assertion fails. The smaller λ , the broader the class of Hölder continuous functions is. The narrowest class is that of functions satisfying the Lipschitz condition.

It follows from the last property that if functions $\varphi_1(t)$ and $\varphi_2(t)$ satisfy the Hölder condition with exponents λ_1 and λ_2 , respectively, then their sum and the product, as well as their ratio provided that the denominator is nonzero, satisfy the Hölder condition with exponent $\lambda = \min(\lambda_1, \lambda_2)$.

If $\varphi(t)$ is differentiable and has a bounded derivative, then $\varphi(t)$ satisfies the Lipschitz condition. In general, the converse assertion fails.

14.2-3. Principal Value of a Singular Integral.

Consider the integral

$$\int_a^b \frac{dx}{x-c}, \quad a < c < b.$$

Evaluating this integral as an improper integral, we obtain

$$\int_a^b \frac{dx}{x-c} = \lim_{\substack{\varepsilon_1 \rightarrow 0 \\ \varepsilon_2 \rightarrow 0}} \left(-\int_a^{c-\varepsilon_1} \frac{dx}{c-x} + \int_{c+\varepsilon_2}^b \frac{dx}{x-c} \right) = \ln \frac{b-c}{c-a} + \lim_{\substack{\varepsilon_1 \rightarrow 0 \\ \varepsilon_2 \rightarrow 0}} \ln \frac{\varepsilon_1}{\varepsilon_2}. \quad (5)$$

The limit of the last expression obviously depends on the way in which ε_1 and ε_2 tend to zero. Hence, the improper integral does not exist. This integral is called a *singular integral*. However, this integral can be assigned a meaning if we assume that there is some relationship between ε_1 and ε_2 . For example, if the deleted interval is symmetric with respect to the point c , i.e.,

$$\varepsilon_1 = \varepsilon_2 = \varepsilon, \quad (6)$$

we arrive at the notion of the Cauchy principal value of a singular integral.

The *Cauchy principal value* of the singular integral

$$\int_a^b \frac{dx}{x-c}, \quad a < c < b$$

is the number

$$\lim_{\varepsilon \rightarrow 0} \left(\int_a^{c-\varepsilon} \frac{dx}{x-c} + \int_{c+\varepsilon}^b \frac{dx}{x-c} \right).$$

With regard to formula (5), we have

$$\int_a^b \frac{dx}{x-c} = \ln \frac{b-c}{c-a}. \quad (7)$$

Consider the more general integral

$$\int_a^b \frac{\varphi(x)}{x-c} dx, \quad (8)$$

where $\varphi(x) \in [a, b]$ is a function satisfying the Hölder condition. Let us understand this integral in the sense of the Cauchy principal value, which we define as follows:

$$\int_a^b \frac{\varphi(x)}{x-c} dx = \lim_{\varepsilon \rightarrow 0} \left(\int_a^{c-\varepsilon} \frac{\varphi(x)}{x-c} dx + \int_{c+\varepsilon}^b \frac{\varphi(x)}{x-c} dx \right).$$

We have the identity

$$\int_a^b \frac{\varphi(x)}{x-c} dx = \int_a^b \frac{\varphi(x) - \varphi(c)}{x-c} dx + \varphi(c) \int_a^b \frac{dx}{x-c};$$

moreover, the first integral on the right-hand side is convergent as an improper integral, because it follows from the Hölder condition that

$$\left| \frac{\varphi(x) - \varphi(c)}{x-c} \right| < \frac{A}{|x-c|^{1-\lambda}}, \quad 0 < \lambda \leq 1,$$

and the second integral coincides with (7).

Thus, we see that the singular integral (8), where $\varphi(x)$ satisfies the Hölder condition, exists in the sense of the Cauchy principal value and is equal to

$$\int_a^b \frac{\varphi(x)}{x-c} dx = \int_a^b \frac{\varphi(x) - \varphi(c)}{x-c} dx + \varphi(c) \ln \frac{b-c}{c-a}.$$

Some authors denote singular integrals by special symbols like v.p. \int (valeur principale). However, this is not necessary because, on one hand, if an integral of the form (8) exists as a proper or an improper integral, then it exists in the sense of the Cauchy principal value, and their values coincide; on the other hand, we shall always understand a singular integral in the sense of the Cauchy principal value. For this reason, we denote a singular integral by the usual integral sign.

14.2-4. Multivalued Functions.

In the representation $z = \rho e^{i\theta}$ of a complex number, the modulus ρ is determined uniquely, whereas the argument θ is only defined modulo 2π . This does not make the representation of a number ambiguous, because the argument enters this representation via the function $e^{i\theta}$, which is 2π -periodic. However, if the dependence of an analytic function on the argument θ is not 2π -periodic, then this function turns out to be multivalued. Of the elementary functions, the logarithm and the power function with noninteger exponent have this property:

$$\ln(z - z_0) = \ln|z - z_0| + i \arg(z - z_0) = \ln \rho + i\Theta, \quad (9)$$

$$(z - z_0)^\gamma = \rho^\gamma e^{i\gamma\Theta} = \rho^\alpha [\cos(\beta \ln \rho) + i \sin(\beta \ln \rho)] e^{i\gamma\theta} e^{i2\pi k\gamma}, \quad \gamma = \alpha + i\beta. \quad (10)$$

In our reasoning, the logarithm of the modulus of a complex number is always understood as a real number, according to the usual definition. The general representation of the argument Θ has the form

$$\Theta = \theta + 2\pi k,$$

where k ranges over all integers ($k = 0, \pm 1, \pm 2, \dots$) and θ is the argument with the least absolute value.

To any k , there corresponds a branch of the multivalued function. The logarithmic function has infinitely many branches. The same holds for the power function with an irrational or nonreal exponent. However, if the exponent is rational, $\gamma = p/q$, with $\gcd(p, q) = 1$, then the power function has q branches. The branches of the logarithm differ by a constant of the form $i2\pi m$, and the branches of a power function differ by a factor of the form $e^{i2\pi m\gamma}$ (m is an integer). Obviously, to define a multivalued function, it is necessary to indicate which branch is chosen. However, in contrast to the case of functions of a real variable, this is not sufficient for the complete definition of a multivalued function of a complex variable. For the latter functions, there are points on the plane with the following property: as the independent variable goes along a closed contour surrounding this point and returns to the initial value, the chosen branch of the function changes to some other branch. Such points are called the *branching points* of the multivalued function. For the functions (9) and (10), the branching points are z_0 and the point at infinity. If the variable is going along a contour surrounding the point z_0 counterclockwise or clockwise, then the argument Θ is changed by 2π or by -2π , respectively.

Accordingly, the logarithm is increased or decreased by $i2\pi$, and the power function is multiplied by $e^{i2\pi\gamma}$ or $e^{-i2\pi\gamma}$. Hence, the branch corresponding to the value $k = n$ passes to the neighboring branch corresponding to $k = n + 1$ or $k = n - 1$. As usual, the study of the point at infinity is performed by the substitution $z = 1/\zeta$ with the subsequent investigation at the point $\zeta = 0$.

We can preserve a chosen branch of a function only if we forbid going around an arbitrary branching point. To this end, we may use *cuts* joining the branching points. In the above cases of the logarithmic and the power function, we can make a cut along a curve issuing from the point z_0 and passing to infinity. A multivalued function is defined uniquely if the branch is chosen and the cut is given.

The range of Θ is determined by the position of the cut. For example, if the cut passes along the ray that forms an angle θ_0 with the real axis, then for the principal branch ($k = 0$) we have $\theta_0 \leq \Theta \leq \theta_0 + 2\pi$. In particular, for the cut that passes along the positive real axis, we have $0 \leq \Theta \leq 2\pi$; and for cut along the negative real axis, we obtain $-\pi \leq \Theta \leq \pi$. If the cut is curvilinear, then the range of the argument depends on the functions of a point. The initial value of the argument corresponds to the left edge of the cut (with respect to z_0) and the final value corresponds to the right edge. Let us denote the value of the argument on the left and on the right edge of the cut by Θ^+ and Θ^- , respectively. Then we have

$$\Theta^- - \Theta^+ = 2\pi.$$

For the chosen branch, the cut is a curve of discontinuity. On the edges of the cut we have

$$\begin{aligned}\ln(z^- - z_0) &= \ln(z^+ - z_0) + i2\pi, \\ (z^- - z_0)^\gamma &= e^{i2\pi\gamma}(z^+ - z_0)^\gamma.\end{aligned}$$

This discontinuity property of branches of multivalued functions on the edges of a cut is widely used in the solution of boundary value problems with discontinuous boundary conditions. The logarithm is applied for the case in which a discontinuous function enters the boundary condition as a summand, and the power function corresponds to the case of a discontinuous factor in the boundary conditions.

14.2-5. Principal Value of a Singular Curvilinear Integral.

Let L be a smooth contour and let τ and t be complex coordinates of its points. Consider the singular curvilinear integral

$$\int_L \frac{\varphi(\tau)}{\tau - t} d\tau. \quad (11)$$

Let us take a circle of some radius ρ centered at the point t on the contour. Let t_1 and t_2 be the points of intersection of this circle with the curve. Assume that the radius is so small that the circle has no other points of intersection with L . Let l be the part of the contour L cut out by the circle. Consider the integral over the remaining arc,

$$\int_{L-l} \frac{\varphi(\tau)}{\tau - t} d\tau. \quad (12)$$

The limit of the integral (12) as $\rho \rightarrow 0$ is called the *principal value* of the singular integral (11). Using the representation

$$\int_L \frac{\varphi(\tau)}{\tau - t} d\tau = \int_L \frac{\varphi(\tau) - \varphi(t)}{\tau - t} d\tau + \varphi(t) \int_L \frac{d\tau}{\tau - t}$$

and the same reasoning as above, we see that the singular integral (11) exists in the sense of the Cauchy principal value for any function $\varphi(\tau)$ satisfying the Hölder condition.

At any point of smoothness, this integral can be presented in two forms:

$$\begin{aligned}\int_L \frac{\varphi(\tau)}{\tau - t} d\tau &= \int_L \frac{\varphi(\tau) - \varphi(t)}{\tau - t} d\tau + \varphi(t) \left(\ln \frac{b-t}{a-t} + i\pi \right) \\ \int_L \frac{\varphi(\tau)}{\tau - t} d\tau &= \int_L \frac{\varphi(\tau) - \varphi(t)}{\tau - t} d\tau + \varphi(t) \ln \frac{b-t}{t-a},\end{aligned}$$

where a and b are the endpoints of L .

In particular, if the contour is closed, then by setting $a = b$ we obtain

$$\int_L \frac{\varphi(\tau)}{\tau - t} d\tau = \int_L \frac{\varphi(\tau) - \varphi(t)}{\tau - t} d\tau + i\pi\varphi(t).$$

Throughout the following, any singular integral will be understood in the sense of the Cauchy principal value.

Let L be a smooth contour (closed or nonclosed) and let $\varphi(\tau)$ be a Hölder function of a point on the contour. Then the Cauchy type integral

$$\Phi(z) = \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - z} d\tau \quad (13)$$

has limit values $\Phi^+(t)$ and $\Phi^-(t)$ at any point of $t \in L$ other than the endpoints of the contour, as $z \rightarrow t$ from the left or from the right along any path; and these limit values can be expressed via the density $\varphi(t)$ of the integral and via the singular integral (13) by the *Sokhotski–Plemelj formulas*

$$\Phi^+(t) = \frac{1}{2}\varphi(t) + \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau, \quad \Phi^-(t) = -\frac{1}{2}\varphi(t) + \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau. \quad (14)$$

The sum and the difference of formulas (14) give the equivalent formulas

$$\Phi^+(t) - \Phi^-(t) = \varphi(t), \quad (15)$$

$$\Phi^+(t) + \Phi^-(t) = \frac{1}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau, \quad (16)$$

which are often used instead of (14).

The *Sokhotski–Plemelj formulas for the real axis* have the form

$$\Phi^+(x) = \frac{1}{2}\varphi(x) + \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\varphi(\tau)}{\tau - x} d\tau, \quad \Phi^-(x) = -\frac{1}{2}\varphi(x) + \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\varphi(\tau)}{\tau - x} d\tau. \quad (17)$$

Moreover, we have

$$\Phi^+(\infty) = \frac{1}{2}\varphi(\infty), \quad \Phi^-(\infty) = -\frac{1}{2}\varphi(\infty).$$

This, together with (17), implies

$$\Phi^+(\infty) + \Phi^-(\infty) = 0, \quad (18)$$

$$\lim_{x \rightarrow \infty} \int_{-\infty}^{\infty} \frac{\varphi(\tau)}{\tau - x} d\tau = 0. \quad (19)$$

Any function representable by a Cauchy type integral on the real axis necessarily satisfies condition (18). This condition is also sufficient for the representability of a piecewise analytic function in the upper and the lower half-plane by an integral over the real axis.

Consider a Cauchy type integral over the real axis and assume that z is not real:

$$\Phi(z) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\varphi(x)}{x - z} dx, \quad (20)$$

where $\varphi(x)$ is a complex function of a real variable x satisfying the Hölder condition on the real axis.

If a function $\varphi(z)$ is analytic in the upper half-plane, is continuous in the closed upper half-plane, and satisfies the Hölder condition on the real axis, then

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\varphi(x)}{x - z} dx = \begin{cases} \varphi(z) - \frac{1}{2}\varphi(\infty) & \text{for } \operatorname{Im} z > 0, \\ -\frac{1}{2}\varphi(\infty) & \text{for } \operatorname{Im} z < 0. \end{cases} \quad (21)$$

We also have the formula

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\varphi(x) - \varphi(\infty)}{x - z} dx = \begin{cases} \frac{1}{2}\varphi(\infty) & \text{for } \operatorname{Im} z > 0, \\ -\varphi(z) + \frac{1}{2}\varphi(\infty) & \text{for } \operatorname{Im} z < 0 \end{cases} \quad (22)$$

provided that $\varphi(z)$ is analytic in the lower half-plane, continuous in the closed lower half-plane, and satisfies the Hölder condition on the real axis.

14.2-6. Poincaré–Bertrand Formula.

Consider the following pair of iterated singular integrals:

$$N(t) = \frac{1}{\pi i} \int_L \frac{d\tau}{\tau - t} \frac{1}{\pi i} \int_L \frac{K(\tau, \tau_1)}{\tau_1 - \tau} d\tau_1, \quad (23)$$

$$M(t) = \frac{1}{\pi i} \int_L d\tau_1 \frac{1}{\pi i} \int_L \frac{K(\tau, \tau_1)}{(\tau - t)(\tau_1 - \tau)} d\tau, \quad (24)$$

where L is a smooth contour and the function $K(\tau, \tau_1)$ satisfies the Hölder condition with respect to both variables.

Both integrals make sense, and although N differs from M only by the order of integration, they are not equal, as shown by the following *Poincaré–Bertrand formula*

$$\frac{1}{\pi i} \int_L \frac{d\tau}{\tau - t} \frac{1}{\pi i} \int_L \frac{K(\tau, \tau_1)}{\tau_1 - \tau} d\tau_1 = K(t, t) + \frac{1}{\pi i} \int_L d\tau_1 \frac{1}{\pi i} \int_L \frac{K(\tau, \tau_1)}{(\tau - t)(\tau_1 - \tau)} d\tau, \quad (25)$$

which can also be rewritten in the form

$$\int_L \frac{d\tau}{\tau - t} \int_L \frac{K(\tau, \tau_1)}{\tau_1 - \tau} d\tau_1 = -\pi^2 K(t, t) + \int_L d\tau_1 \int_L \frac{K(\tau, \tau_1)}{(\tau - t)(\tau_1 - \tau)} d\tau. \quad (26)$$

Example. Let us evaluate the Cauchy type integral over the unit circle $|z| = 1$ with density $\varphi(\tau) = 2/[\tau(\tau - 2)]$, i.e.,

$$\Phi(z) = \frac{1}{2\pi i} \int_L \frac{1}{\tau - 2} \frac{d\tau}{\tau - z} - \frac{1}{2\pi i} \int_L \frac{1}{\tau} \frac{d\tau}{\tau - z}.$$

The function $1/(z - 2)$ is analytic in Ω^+ , and $1/z$ is analytic in Ω^- and vanishes at infinity. By formula (1), the first integral is equal to $1/(z - 2)$ for $z \in \Omega^+$ and is zero for $z \in \Omega^-$. By formula (2), the second integral is equal to $-1/z$ for $z \in \Omega^-$ and is zero for $z \in \Omega^+$. Hence,

$$\Phi^+(z) = \frac{1}{z - 2}, \quad \Phi^-(z) = \frac{1}{z}.$$

☉ References for Section 14.2: F. D. Gakhov (1977), S. G. Mikhlin and S. Prössdorf (1986), N. I. Muskhelishvili (1992).

14.3. Riemann Boundary Value Problem

14.3-1. Principle of Argument. The Generalized Liouville Theorem.

THE THEOREM ON THE ANALYTIC CONTINUATION (THE PRINCIPLE OF CONTINUITY). Assume that a domain Ω_1 borders a domain Ω_2 along a smooth curve L . Let analytic functions $f_1(z)$ and $f_2(z)$ be given in Ω_1 and Ω_2 . Assume that, as the point z tends to L , both functions tend to the same continuous limit function on the curve L . Under these assumptions, the functions $f_1(z)$ and $f_2(z)$ are analytic continuations of each other.

Assume that a function $f(z)$ is analytic in a domain Ω bounded by a contour L except for finitely many points, where it may have poles. Let us write out the power series expansion of $f(z)$ around some point z_0 :

$$f(z) = c_n(z - z_0)^n + c_{n+1}(z - z_0)^{n+1} + \dots = (z - z_0)^n f_1(z), \quad f_1(z_0) = c_n \neq 0.$$

The number n is called the *order of the function $f(z)$ at the point z_0* . If $n > 0$, then the order of the function is the order of zero; if $n < 0$, then the order of the function is minus the order of the pole. If the order of a function at z_0 is zero, then at z_0 the function has a finite nonzero value at z_0 . When considering the point at infinity, we must replace the difference $z - z_0$ by $1/z$. If $z_0 \in L$, then we define the order of the function to be equal to $\frac{1}{2}n$.

Let N_Ω and P_Ω (N_L and P_L) be the numbers of zeros and poles on the domain (on the contour, respectively), where each zero and pole is taken according to its multiplicity. Let $[\delta]_L$ denote the increment of the variable δ when going around the contour in the positive direction. As usual, by the positive direction we mean the direction the domain under consideration remains to the left of the contour.

THE PRINCIPLE OF ARGUMENT. Let $f(z)$ be a single-valued analytic function in a multiply connected domain Ω bounded by a smooth contour $L = L_0 + L_1 + \dots + L_m$ except for finitely many points at which $f(z)$ may have poles, and let $f(z)$ be continuous in the closed domain $\Omega \cup L$ (except for these poles) and have at most finitely many zeros of integer order on the contour. In this case, the following formula holds:

$$N_\Omega - P_\Omega + \frac{1}{2}(N_L - P_L) = \frac{1}{2\pi}[\arg f(z)]_L.$$

THE GENERALIZED LIOUVILLE THEOREM. Assume that a function $f(z)$ is analytic on the entire complex plane except for points $a_0 = \infty, a_k$ ($k = 1, \dots, n$), where it has poles, and that the principal parts of the Laurent series expansions of $f(z)$ at the poles have the form

$$Q_0(z) = c_1^0 z + c_2^0 z^2 + \dots + c_{m_0}^0 z^{m_0} \quad \text{at the point } a_0,$$

$$Q_k\left(\frac{1}{z - a_k}\right) = \frac{c_1^k}{z - a_k} + \frac{c_2^k}{(z - a_k)^2} + \dots + \frac{c_{m_k}^k}{(z - a_k)^{m_k}} \quad \text{at the points } a_k.$$

Then $f(z)$ is a rational function, and can be represented by the formula

$$f(z) = C + Q_0(z) + \sum_{k=1}^n Q_k\left(\frac{1}{z - a_k}\right),$$

where C is a constant. In particular, if the only singularity of $f(z)$ is a pole of order m at infinity, then $f(z)$ is a polynomial of degree m ,

$$f(z) = c_0 + c_1 z + \dots + c_m z^m.$$

The following notation is customary:

- (a) $\overline{f(z)}$ is the function conjugate to a given function $f(z)$;
- (b) $f(\bar{z})$ is the function obtained from $f(z)$ by replacing z by \bar{z} , i.e., y by $-y$ in $f(z)$;
- (c) $\bar{f}(z)$ is the function defined by the condition $f(z) = \bar{f}(\bar{z})$.

If $z = x + iy$ and $f(z) = u(x, y) + iv(x, y)$, then

$$\overline{f(z)} = u(x, y) - iv(x, y), \quad f(\bar{z}) = u(x, -y) + iv(x, -y), \quad \bar{f}(z) = u(x, -y) - iv(x, -y).$$

In particular, if $f(z)$ is given by a series $f(z) = \sum_{k=0}^n c_k z^k$, then

$$\overline{f(z)} = \sum_{k=0}^n \bar{c}_k \bar{z}^k, \quad f(\bar{z}) = \sum_{k=0}^n c_k \bar{z}^k, \quad \bar{f}(z) = \sum_{k=0}^n \bar{c}_k z^k.$$

For a function represented by a Cauchy type integral

$$f(z) = \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - z} d\tau,$$

we have

$$\overline{f(z)} = -\frac{1}{2\pi i} \int_L \frac{\overline{\varphi(\tau)}}{\bar{\tau} - \bar{z}} d\bar{\tau}, \quad f(\bar{z}) = \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - \bar{z}} d\tau, \quad \bar{f}(z) = -\frac{1}{2\pi i} \int_L \frac{\overline{\varphi(\tau)}}{\bar{\tau} - z} d\bar{\tau}.$$

Note that if a function satisfies the condition $\bar{f}(z) = f(z)$, then it takes real values for all real values of z . The converse assertion also holds.

14.3-2. Hermite Interpolation Polynomial.

The Hermite interpolation polynomial is used for the construction of the canonical function of the nonhomogeneous Riemann problem in Subsections 12.4-7 and 14.3-9.

Let distinct points z_k ($k = 1, \dots, m$) be given, and a number $\Delta_k^{(j)}$ ($j = 0, 1, \dots, n_k - 1$) be assigned to each point z_k , where the n_k are given positive integers. It is required to construct a polynomial $\mathcal{U}_p(z)$ of the least possible degree such that

$$\mathcal{U}_p^{(j)}(z_k) = \Delta_k^{(j)}, \quad k = 1, \dots, m, \quad j = 0, 1, \dots, n_k - 1,$$

where the $\mathcal{U}_p^{(j)}(z_k)$ are the values of the j th-order derivatives of the polynomial at the points z_k . The numbers z_k are called the *interpolation nodes* and n_k the *interpolation multiplicities at the nodes* z_k .

There exists a unique polynomial with these properties. It has the form (e.g., see V. I. Smirnov and N. A. Lebedev (1964))

$$\mathcal{U}_p(z) = \sum_{k=1}^m \frac{\zeta(z)}{(z - z_k)^{n_k}} \sum_{r=0}^{n_k-1} A_{k,r} (z - z_k)^r, \quad p = \sum_{k=1}^m n_k - 1,$$

$$\zeta(z) = \prod_{k=1}^m (z - z_k)^{n_k}, \quad A_{k,r} = \sum_{j=0}^r \frac{\Delta_k^{(j)}}{j! (r-j)!} \left[\frac{d^{r-j}}{dz^{r-j}} \frac{(z - z_k)^{n_k}}{\zeta(z)} \right]_{z=z_k},$$

$$k = 1, \dots, m, \quad r = 0, 1, \dots, n_k - 1;$$

and this polynomial is unique.

The interpolation polynomial $\mathcal{U}_p(z)$ constructed for some function $f(z)$ must satisfy the following conditions at the points z_k :

$$\mathcal{U}_p^{(j)}(z_k) = \Delta_k^{(j)} = f^{(j)}(z_k), \quad k = 1, \dots, m, \quad j = 0, 1, \dots, n_k - 1,$$

where $f^{(j)}(z_k)$ is the value of the j th-order derivative of $f(z)$ at the point z_k .

14.3-3. Notion of the Index.

Let L be a smooth closed contour, and let $D(t)$ be a continuous nowhere vanishing function on this contour.

The *index* ν of the function $D(t)$ with respect to the contour L is the increment of the argument of $D(t)$ along L (traversed in the positive direction) divided by 2π :

$$\nu = \text{Ind } D(t) = \frac{1}{2\pi} [\arg D(t)]_L. \quad (1)$$

Since $\ln D(t) = \ln |D(t)| + i \arg D(t)$ and since after the traverse the function $|D(t)|$ returns to its original value, it follows that $[\ln D(t)]_L = i [\arg D(t)]_L$, and hence

$$\nu = \frac{1}{2\pi i} [\ln D(t)]_L. \quad (2)$$

The index can be expressed in the form of an integral as follows:

$$\nu = \text{Ind } D(t) = \frac{1}{2\pi i} \int_L d \ln D(t) = \frac{1}{2\pi} \int_L d \arg D(t). \quad (3)$$

If the function $D(t)$ is not differentiable but has bounded variation, then the integral is regarded as the Stieltjes integral. Since $D(t)$ is continuous, the image $\dot{\Gamma}$ of the closed contour L is a closed contour as well, and the increment of the argument $D(t)$ along L is a multiple of 2π . Hence, the following assertions hold.

1°. The index of a function that is continuous on a closed contour and vanishes nowhere is an integer (possibly zero).

2°. The index of the product of two functions is equal to the sum of the indexes of the factors. The index of a ratio is equal to the difference of the indexes of the numerator and the denominator.

We now assume that $D(t)$ is differentiable and is the boundary value of a function analytic in the interior or exterior of L . In this case, the number

$$\nu = \frac{1}{2\pi i} \int_L d \ln D(t) = \frac{1}{2\pi i} \int_L \frac{D'_t(t)}{D(t)} dt \quad (4)$$

is equal to the logarithmic residue of the function $D(t)$. The principle of argument (see Subsection 14.3-1) implies the following properties of the index:

3°. If $D(t)$ is the boundary value of a function analytic in the interior or exterior of the contour, then its index is equal to the number of zeros inside the contour or minus the number of zeros outside the contour, respectively.

4°. If a function $D(z)$ is analytic in the interior of the contour except for finitely many points at which it may have poles, then the number of zeros must be replaced by the difference of the number of zeros and the number of poles.

Here the zeros and the poles are counted according to their multiplicities. We also note that the indexes of complex conjugate functions have opposite signs.

Let

$$t = t_1(s) + it_2(s) \quad (0 \leq s \leq l)$$

be the equation of the contour L . On substituting the expression of the complex coordinate t into the function $D(t)$, we obtain

$$D(t) = D(t_1(s) + it_2(s)) = \xi(s) + i\eta(s). \quad (5)$$

Let us regard ξ and η as Cartesian coordinates. Then

$$\xi = \xi(s), \quad \eta = \eta(s)$$

is a parametric equation of some curve Γ . Since the function $D(t)$ is continuous and the contour L is closed, it follows that the curve Γ is closed as well.

The number of turns of the curve Γ around the origin, i.e., the number of full rotations of the radius vector as the variable s varies from 0 to l , is obviously the index of the function $D(t)$. This number is often called the *winding number of the curve* Γ with respect to the origin.

If the curve Γ is successfully constructed, then the winding number can be observed directly. There are many examples for which the index can be found by analyzing the shape of the curve Γ . For instance, if $D(t)$ is a real or a pure imaginary function that does not vanish, then Γ is a line segment (traversed an even number of times), and the index $D(t)$ is equal to zero. If the real part $\xi(s)$ or the imaginary part $\eta(s)$ preserves its sign, then the index is obviously zero, and so on. If the function $D(t)$ can be represented as the product or the ratio of functions that are limit values of functions analytic in the interior or exterior of the contour, then the index can be calculated on the basis of properties 2°, 3°, and 4°.

In the general case, the calculation of the index can be performed by formula (3). On the basis of formula (5) we substitute the expression

$$d \arg D(t) = d \arctan \frac{\eta(s)}{\xi(s)}$$

into (3) and assume that ξ and η are differentiable. Then we obtain

$$\nu = \frac{1}{2\pi} \int_{\Gamma} \frac{\xi d\eta - \eta d\xi}{\xi^2 + \eta^2} = \frac{1}{2\pi} \int_0^l \frac{\xi(s)\eta'_s(s) - \eta(s)\xi'_s(s)}{\xi^2(s) + \eta^2(s)} ds. \quad (6)$$

Example 1. Let us calculate the index of $D(t) = t^n$ with respect to an arbitrary contour L surrounding the origin.

First method. The function t^n is the boundary value of the function z^n , which has precisely one zero of order n inside the contour. Hence

$$\nu = \text{Ind } t^n = n.$$

Second method. If the argument of t is φ , then the argument of t^n is $n\varphi$. As the point t traverses the contour L and returns to the original value, the argument φ obtains the increment 2π . Hence,

$$\text{Ind } t^n = n.$$

The index can also be found numerically. Since the index is integer-valued, an approximate value whose error is less than $\frac{1}{2}$ can be rounded off to the nearest integer to obtain the exact value.

14.3-4. Statement of the Riemann Problem.

Let L be a simple smooth closed contour which divides the complex plane into the interior domain Ω^+ and the exterior domain Ω^- , and let two functions of points of the contours $D(t)$ and $H(t)$ satisfying the Hölder condition (see Subsection 14.2-2) be given; moreover, suppose that $D(t)$ does not vanish.

The Riemann Problem. Find two functions (or a single piecewise analytic function), namely, a function $\Phi^+(z)$ analytic in Ω^+ and a function $\Phi^-(z)$ analytic in the domain Ω^- including $z = \infty$, so that the following linear relation is satisfied on the contour L :

$$\Phi^+(t) = D(t)\Phi^-(t) \quad (\text{the homogeneous problem}) \quad (7)$$

or

$$\Phi^+(t) = D(t)\Phi^-(t) + H(t) \quad (\text{the nonhomogeneous problem}). \quad (8)$$

The function $D(t)$ is called the *coefficient* of the Riemann problem, and the function $H(t)$ is called the *right-hand side*.

We first consider a Riemann problem of special form that is called the *jump problem*. Let a function $\varphi(t)$ defined on a closed contour L satisfy the Hölder condition. The problem is to find a piecewise analytic function $\Phi(z)$ ($\Phi(z) = \Phi^+(z)$ for $z \in \Omega^+$ and $\Phi(z) = \Phi^-(z)$ for $z \in \Omega^-$) that vanishes at infinity and has a jump of magnitude $\varphi(t)$ on L , i.e., such that

$$\Phi^+(t) - \Phi^-(t) = \varphi(t).$$

It follows from the Sokhotski–Plemelj formulas (see Subsection 14.2-5) that the function

$$\Phi(z) = \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - z} d\tau$$

is the unique solution to the above problem.

Thus, an arbitrary function $\varphi(t)$ given on the closed contour and satisfying the Hölder condition can be uniquely represented as the difference of functions $\Phi^+(t)$ and $\Phi^-(t)$ that are the boundary values of analytic functions $\Phi^+(z)$ and $\Phi^-(z)$ under the additional condition $\Phi^-(\infty) = 0$.

If we neglect the additional condition $\Phi^-(\infty) = 0$, then the solution will be given by the formula

$$\Phi(z) = \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - z} d\tau + \text{const}. \quad (9)$$

Let us seek a particular solution of the homogeneous problem (7) in the class of functions that do not vanish on the contour. Let N_+ and N_- be the numbers of zeros of the desired functions in the domains Ω^+ and Ω^- , respectively. Taking the index of both parts of Eq. (7), on the basis of properties 2° and 3° we obtain

$$N_+ + N_- = \text{Ind } D(t) = \nu. \quad (10)$$

We call the index ν of the coefficient $D(t)$ the *index of the Riemann problem*.

Let $\nu = 0$. Under this condition, $\ln D(t)$ is a single-valued function. It follows from (10) that $N_+ = N_- = 0$, i.e., the solution has no zeros on the entire plane. Therefore, the functions $\ln \Phi^\pm(z)$ are analytic in their domains and hence single-valued together with the boundary values $\ln \Phi^\pm(t)$.

Taking the logarithm of the boundary condition (7), we obtain

$$\ln \Phi^+(t) - \ln \Phi^-(t) = \ln D(t). \quad (11)$$

We can choose an arbitrary branch of $\ln D(t)$ because the final result is independent of the choice of this branch. Thus, we must find a piecewise analytic function $\ln \Phi(z)$ with a prescribed jump on L . The solution of this problem under the additional condition $\ln \Phi^-(\infty) = 0$ is given by the formula

$$\ln \Phi(z) = \frac{1}{2\pi i} \int_L \frac{\ln D(\tau)}{\tau - z} d\tau. \quad (12)$$

For brevity, we write

$$\frac{1}{2\pi i} \int_L \frac{\ln D(\tau)}{\tau - z} d\tau = G(z). \quad (13)$$

It readily follows from the Sokhotski–Plemelj formulas that the functions

$$\Phi^+(z) = e^{G^+(z)} \quad \text{and} \quad \Phi^-(z) = e^{G^-(z)} \quad (14)$$

are the solution of the boundary value problem (7) with the condition $\Phi^-(\infty) = 1$.

If we neglect the additional condition $\Phi^-(\infty) = 1$, then in formula (12) we must add an arbitrary constant, and the solution becomes

$$\Phi^+(z) = C e^{G^+(z)}, \quad \Phi^-(z) = C e^{G^-(z)}, \quad (15)$$

where C is an arbitrary constant. Since $G^-(\infty) = 0$, it follows that C is the value of $\Phi^-(z)$ at infinity.

Thus, in the case $\nu = 0$ and for arbitrary $\Phi^-(\infty) \neq 0$, the solution contains a single arbitrary constant, and hence there is a unique linearly independent solution. If $\Phi^-(\infty) = 0$, then $C = 0$, and the problem has only the trivial solution (which is identically zero), which is natural because $N_- = 0$.

This gives an important corollary. An arbitrary function $D(t) \neq 0$ on L that satisfies the Hölder condition and has zero index can be represented as the ratio of the boundary values $\Phi^+(t)$ and $\Phi^-(t)$ of functions that are analytic in Ω^+ and Ω^- and have no zeros in these domains. These functions are determined modulo an arbitrary constant factor and are given by formulas (15).

On passing to the general case, we seek a piecewise analytic function satisfying the homogeneous boundary condition (7) and having zero order on the entire plane except for the point at infinity, where the order of the function is equal to the index of the problem.

By the *canonical function* (of the homogeneous Riemann problem) $X(z)$ we mean the function satisfying the boundary condition (7) and piecewise analytic on the entire plane except for the point at infinity, where the order of this function is equal to the index of the problem.

This function can be constructed by reducing the problem to the case of zero index. Indeed, let us rewrite the boundary condition (7) in the form

$$\Phi^+(t) = t^{-\nu} D(t) t^\nu \Phi^-(t).$$

On representing the function $t^{-\nu}D(t)$ with zero index as the ratio of boundary values of analytic functions,

$$t^{-\nu}D(t) = \frac{e^{G^+(t)}}{e^{G^-(t)}}, \quad G(z) = \frac{1}{2\pi i} \int_L \frac{\ln[\tau^{-\nu}D(\tau)]}{\tau - z} d\tau, \quad (16)$$

we obtain the following expression for the canonical function:

$$X^+(z) = e^{G^+(z)}, \quad X^-(z) = z^{-\nu} e^{G^-(z)}. \quad (17)$$

Since $X^+(t) = D(t)X^-(t)$, it follows that the coefficient of the Riemann problem can be represented as the ratio of canonical functions:

$$D(t) = \frac{X^+(t)}{X^-(t)}. \quad (18)$$

The representation (18) is often called a *factorization*.

For $\nu \geq 0$, the canonical function, which has a zero of order ν at infinity, is a particular solution of the boundary value problem (7). For $\nu < 0$, the canonical function has a pole of order $|\nu|$ at infinity and is not a solution, but in this case it is still used as an auxiliary function in the solution of the nonhomogeneous problem.

14.3-5. Solution of the Homogeneous Problem.

Let $\nu = \text{Ind } D(t)$ be an arbitrary integer. On representing $D(t)$ by formula (18), we reduce the boundary condition (7) to the form

$$\frac{\Phi^+(t)}{X^+(t)} = \frac{\Phi^-(t)}{X^-(t)}.$$

The left-hand side of the last relation contains the boundary value of a function that is analytic in Ω^+ , and the right-hand side contains the boundary value of a function that has at least the order $-\nu$ at infinity. By the principle of continuity (see Subsection 14.3-1), the functions on the left-hand side and on the right-hand side are analytic continuations of each other to the entire plane possibly except for the point at infinity at which, in the case $\nu > 0$, a pole of order $\leq \nu$ can occur. Hence, for $\nu > 0$, by the generalized Liouville theorem (see Subsection 14.3-1), this single analytic function is a polynomial of degree $\leq \nu$ with arbitrary coefficients. For $\nu < 0$, it follows from the Liouville theorem that this function is constant. However, since this function must vanish at infinity, it follows that it is identically zero. Hence, for $\nu < 0$, the homogeneous problem has only the trivial solution (which is identically zero). A problem that has no nontrivial solutions is said to be *unsolvable*. Thus, for a negative index, the homogeneous problem (7) is unsolvable.

Let $\nu > 0$. Let $P_\nu(z)$ stand for a polynomial of degree ν with arbitrary coefficients. In this case, we obtain a solution in the form

$$\Phi(z) = P_\nu(z)X(z),$$

or

$$\Phi^+(z) = P_\nu(z)e^{G^+(z)}, \quad \Phi^-(z) = z^{-\nu}P_\nu(z)e^{G^-(z)}, \quad (19)$$

where $G(z)$ is determined by formula (16).

Thus, if the index ν of the Riemann boundary value problem is nonnegative, then the homogeneous problem (7) has $\nu + 1$ linearly independent solutions

$$\Phi_k^+(z) = z^k e^{G^+(z)}, \quad \Phi_k^-(z) = z^{k-\nu} e^{G^-(z)} \quad (k = 0, 1, \dots, \nu). \quad (20)$$

The general solution contains $\nu + 1$ arbitrary constants and is given by formula (19). For a negative index, problem (7) is unsolvable.

The polynomial $P_\nu(z)$ has exactly ν zeros in the complex plane. It follows from formulas (19) that the number of all zeros of a solution to the homogeneous Riemann boundary value problem is equal to the index ν . Depending on the choice of the coefficients of the polynomial, these zeros can occur in each of the domains Ω^\pm and also on the contour itself. Just as above, we denote by N_\pm the number of zeros in the domains Ω^\pm and by N_0 the number of zeros on the contour L . We can see that in the general case (without the condition that there are no zeros on the contour), formula (10) becomes

$$N_+ + N_- + N_0 = \nu. \quad (21)$$

14.3-6. Solution of the Nonhomogeneous Problem.

On replacing the coefficient $D(t)$ in the boundary condition (8) by the ratio of the boundary values of the canonical functions by formula (18), we reduce (8) to the form

$$\frac{\Phi^+(t)}{X^+(t)} = \frac{\Phi^-(t)}{X^-(t)} + \frac{H(t)}{X^+(t)}. \quad (22)$$

The function $H(t)/X^+(t)$ satisfies the Hölder condition. Let us replace it by the difference of the boundary values of analytic functions (see the jump problem in Subsection 14.3-4):

$$\frac{H(t)}{X^+(t)} = \Psi^+(t) - \Psi^-(t),$$

where

$$\Psi(z) = \frac{1}{2\pi i} \int_L \frac{H(\tau)}{X^+(\tau)} \frac{d\tau}{\tau - z}. \quad (23)$$

Then the boundary condition (22) can be rewritten in the form

$$\frac{\Phi^+(t)}{X^+(t)} - \Psi^+(t) = \frac{\Phi^-(t)}{X^-(t)} - \Psi^-(t).$$

Note that for $\nu \geq 0$ the function $\Phi^-(z)/X^-(z)$ has a pole at infinity, and for $\nu < 0$ it has a zero of order ν .

By the same reasoning as in the solution of the homogeneous problem, we obtain the following results.

Let $\nu \geq 0$. In this case,

$$\frac{\Phi^+(t)}{X^+(t)} - \Psi^+(t) = \frac{\Phi^-(t)}{X^-(t)} - \Psi^-(t) = P_\nu(t).$$

This gives the solution

$$\Phi(z) = X(z)[\Psi(z) + P_\nu(z)], \quad (24)$$

where the functions $X(z)$ and $\Psi(z)$ are expressed by formulas (17) and (23) and P_ν is a polynomial of degree ν with arbitrary coefficients.

We can readily see that formula (24) gives the general solution of the nonhomogeneous problem since it contains the general solution $X(z)P_\nu(z)$ of the homogeneous problem as a summand.

Let $\nu < 0$. In this case, $\Phi^-(z)/X^-(z)$ vanishes at infinity and

$$\frac{\Phi^+(t)}{X^+(t)} - \Psi^+(t) = \frac{\Phi^-(t)}{X^-(t)} - \Psi^-(t) = 0,$$

so that

$$\Phi(z) = X(z)\Psi(z). \quad (25)$$

In the expression for the function $\Phi^-(z)$, the first factor has a pole of order $-\nu$ at infinity by virtue of formula (17), and the second factor is the Cauchy type integral (23) and, in general, has a first-order zero at infinity. Hence, $\Phi^-(z)$ has a pole of order $\leq -\nu - 1$ at infinity. Thus, if $\nu < -1$, then the nonhomogeneous problem is unsolvable in general. It is solvable only if the constant term satisfies some additional conditions. To find these conditions, we expand the Cauchy type integral (23) in a series in a neighborhood of the point at infinity:

$$\Psi^-(z) = \sum_{k=1}^{\infty} c_k z^{-k}, \quad \text{where} \quad c_k = -\frac{1}{2\pi i} \int_L \frac{H(\tau)}{X^+(\tau)} \tau^{k-1} d\tau.$$

For $\Phi^-(z)$ to be analytic at the point at infinity, it is necessary that the first $-\nu - 1$ coefficients of the expansion of $\Psi^-(z)$ be zero. This means that for the solvability of the nonhomogeneous problem in the case of negative index ($\nu < -1$), it is necessary and sufficient that the following $-\nu - 1$ conditions hold:

$$\int_L \frac{H(\tau)}{X^+(\tau)} \tau^{k-1} d\tau = 0, \quad k = 1, 2, \dots, -\nu - 1. \tag{26}$$

Thus, in the case $\nu \geq 0$, the nonhomogeneous Riemann problem is solvable for an arbitrary right-hand side, and the general solution is given by the formula

$$\Phi(z) = \frac{X(z)}{2\pi i} \int_L \frac{H(\tau)}{X^+(\tau)} \frac{d\tau}{\tau - z} + X(z)P_\nu(z), \tag{27}$$

where the canonical function $X(z)$ is given by (17) and $P_\nu(z)$ is a polynomial of degree ν with arbitrary complex coefficients. If $\nu = -1$, then the nonhomogeneous problem is also solvable and has a unique solution.

In the case $\nu < -1$, the nonhomogeneous problem is unsolvable in general. For this problem to be solvable, it is necessary and sufficient that the right-hand side of the problem satisfy $-\nu - 1$ conditions (26). If these conditions are satisfied, then the solution of the problem is unique and is given by formula (27), where we must set $P_\nu(z) \equiv 0$.

The solution with the additional condition of vanishing at infinity has important applications. In this case, instead of a polynomial of degree ν , we must take a polynomial of degree $\nu - 1$. For the solvability of the problem in the case of negative index, it is necessary that the coefficient $c_{-\nu}$ be zero as well.

Hence, under the assumption that $\Phi^-(\infty) = 0$, the solution is given for $\nu \geq 0$ by the formula

$$\Phi(z) = X(z)[\Psi(z) + P_{\nu-1}(z)], \tag{28}$$

where, for $\nu = 0$, we must set $P_{\nu-1}(z) \equiv 0$.

If $\nu < 0$, then the solution is still given by formula (28) with $P_{\nu-1}(z) \equiv 0$ under the following $-\nu$ solvability conditions:

$$\int_L \frac{H(\tau)}{X^+(\tau)} \tau^{k-1} d\tau = 0, \quad k = 1, 2, \dots, -\nu. \tag{29}$$

In this case, the assertion on the solvability of the nonhomogeneous problem acquires a more symmetric form. For $\nu \geq 0$, the general solution of the nonhomogeneous problem linearly depends on ν arbitrary constants. For $\nu < 0$, the number of the solvability conditions is equal to $-\nu$. Note that for $\nu = 0$ the nonhomogeneous problem is unconditionally solvable, and the solution is unique.

On the basis of the above reasoning, the solution of the Riemann boundary value problem is mainly reduced to the following two operations:

1°. A representation of an arbitrary function given on the contour in the form of the difference of boundary values of analytic functions in the domains Ω^+ and Ω^- (the jump problem).

2°. A representation of a nonvanishing function in the form of the ratio of boundary values of analytic functions (factorization).

Here the second operation can be reduced to the first by taking the logarithm. Some complications related to the case of a nonzero index are due to the multivaluedness of the logarithm only. The first operation for arbitrary functions is equivalent to the calculation of a Cauchy type integral. In this connection, the solution to the problem by formulas (17) and (23)–(25) is explicitly expressed (in the closed form) via Cauchy type integrals.

14.3-7. Riemann Problem with Rational Coefficients.

Consider the Riemann boundary value problem with a contour that consists of finitely many simple curves and with coefficient $D(t)$ a rational function that has neither zeros nor poles on the contour. Note that an arbitrary continuous function (and all the functions satisfying the Hölder condition) can be approximated with arbitrary accuracy by rational functions, and the solution of problems with rational coefficients can serve as a basis for the approximate solution in the general case. Assume that the Riemann problem has the form

$$\Phi^+(t) = \frac{p(t)}{q(t)}\Phi^-(t) + H(t), \quad (30)$$

and the polynomials $p(z)$ and $q(z)$ can be factorized as follows:

$$p(z) = p_+(z)p_-(z), \quad q(z) = q_+(z)q_-(z), \quad (31)$$

where $p_+(z)$ and $q_+(z)$ are polynomials whose roots belong to Ω^+ and $p_-(z)$ and $q_-(z)$ are polynomials with roots in Ω^- . It readily follows from property 4° of the index (Subsection 14.3-3) that $\nu = m_+ - n_+$, where m_+ and n_+ are the numbers of zeros of the polynomials $p_+(z)$ and $q_+(z)$.

Since the coefficient of the problem is a function that can be analytically continued to the domain Ω^\pm , it follows that in this case it is reasonable to avoid using the general formulas and obtain a solution directly by analytic continuation; here the role of the standard function of the type t^ν that is used in the reduction of the index to zero can be played by the product $\prod_{j=1}^{\nu} (t - a_j)$, where a_1, \dots, a_ν are arbitrary points of the domain Ω^+ . On representing the boundary condition in the form

$$\frac{q_-(t)}{p_-(t)}\Phi^+(t) - \frac{p_+(t)}{q_+(t)}\Phi^-(t) = \frac{q_-(t)}{p_-(t)}H(t),$$

where the canonical function is determined by the expressions

$$X^+(z) = \frac{p_-(z)}{q_-(z)}, \quad X^-(z) = \frac{q_+(z)}{p_+(z)}, \quad (32)$$

we obtain the solution by the same reasoning as in Subsection 14.3-6 in the following form:

$$\Phi^+(z) = \frac{p_-(z)}{q_-(z)}[\Psi(z) + P_{\nu-1}(z)], \quad \Phi^-(z) = \frac{q_+(z)}{p_+(z)}[\Psi(z) + P_{\nu-1}(z)], \quad (33)$$

where

$$\Psi(z) = \frac{1}{2\pi i} \int_L \frac{q_-(\tau)}{p_-(\tau)} \frac{H(\tau) d\tau}{\tau - z}, \quad \Phi^-(\infty) = 0.$$

If the index is negative, then we must set $P_{\nu-1}(z) \equiv 0$ and add the solvability conditions

$$\int_L \frac{q_-(\tau)}{p_-(\tau)} H(\tau) \tau^{k-1} d\tau = 0, \quad k = 1, 2, \dots, -\nu, \quad (34)$$

which agree with the general formula (29), because the canonical function has the form (32).

Note that for the general case, in the practical solution of the Riemann problem, it can also be convenient to express the coefficient in the form

$$D(t) = \frac{p_+(t)p_-(t)}{q_+(t)q_-(t)}D_1(t),$$

where $D_1(t)$ is a function with zero index and the polynomials $p_{\pm}(t)$ and $q_{\pm}(t)$ are chosen for a given coefficient in a special way. For an appropriate choice of such polynomials, the solution can be obtained in the simplest possible way.

Example 2. Consider the Riemann problem

$$\Phi^+(t) = \frac{t}{t^2 - 1}\Phi^-(t) + \frac{t^3 - t^2 + 1}{t^3 - t}$$

under the assumption that $\Phi^-(\infty) = 0$ and L is an arbitrary smooth closed contour of one of the following forms:

- 1°. The interior of the contour L contains the point $z_1 = 0$ and does not contain the points $z_2 = 1$ and $z_3 = -1$.
- 2°. The interior of the contour L contains the points $z_1 = 0$ and $z_2 = 1$ and does not contain the point $z_3 = -1$.
- 3°. The interior of the contour L contains the points $z_1 = 0, z_2 = 1,$ and $z_3 = -1$.
- 4°. The interior of the contour L contains the points $z_2 = 1$ and $z_3 = -1$ and does not contain the point $z_1 = 0$.

Consider cases 1°–4° in order. In the solution we apply the method of Subsection 14.3-7.

1°. We have

$$p_+(t) = t, \quad p_-(t) = 1, \quad q_+(t) = 1, \quad q_-(t) = t^2 - 1; \quad m_+ = 1, \quad n_+ = 0, \quad \nu = m_+ - n_+ = 1.$$

Let us rewrite the boundary condition in the form

$$(t^2 - 1)\Phi^+(t) - t\Phi^-(t) = \frac{1}{t}(t^3 - t^2 + 1)(t + 1).$$

Hence,

$$\Psi(z) = \frac{1}{2\pi i} \int_L \frac{q_-(\tau)}{p_-(\tau)} H(\tau) \frac{d\tau}{\tau - z} = \frac{1}{2\pi i} \int_L \frac{\tau^3 - \tau + 1}{\tau - z} d\tau + \frac{1}{2\pi i} \int_L \frac{1/\tau}{\tau - z} d\tau,$$

and the formulas for the Cauchy integral (see Subsection 14.2-1) imply

$$\Psi^+(z) = z^3 - z + 1, \quad \Psi^-(z) = -\frac{1}{z}.$$

The general solution of the problem contains a single (arbitrary) constant. By formula (33), we obtain

$$\Phi^+(z) = \frac{1}{z^2 - 1}(z^3 - z + 1 + C) = \frac{z^3 - z + 1}{z^2 - 1} + \frac{C}{z^2 - 1}, \quad \Phi^-(z) = \frac{1}{z} \left(-\frac{1}{z} + C \right) = -\frac{1}{z^2} + \frac{C}{z},$$

where C is an arbitrary constant. On replacing C by $C - 1$ we can rewrite the solution in the form

$$\Phi^+(z) = z + \frac{C}{z^2 - 1}, \quad \Phi^-(z) = -\frac{z + 1}{z^2} + \frac{C}{z}.$$

2°. We have

$$p_+(t) = t, \quad p_-(t) = 1, \quad q_+(t) = t - 1, \quad q_-(t) = t + 1, \quad m_+ = n_+ = 1, \quad \nu = 0,$$

$$(t + 1)\Phi^+(t) - \frac{t}{t - 1}\Phi^-(t) = \frac{(t + 1)(t^3 - t^2 + 1)}{t(t - 1)},$$

$$\Psi(z) = \frac{1}{2\pi i} \int_L \frac{\tau^2 + \tau}{\tau - z} d\tau + \frac{1}{2\pi i} \int_L \frac{(\tau + 1)/[\tau(\tau - 1)]}{\tau - z} d\tau = \begin{cases} z^2 + z & \text{for } z \in \Omega^+, \\ -\frac{z + 1}{z(z - 1)} & \text{for } z \in \Omega^-. \end{cases}$$

The problem has the unique solution

$$\begin{aligned} \Phi^+(z) &= \frac{p_-(z)}{q_-(z)}\Phi^+(z) = \frac{1}{z + 1}(z^2 + z) = z, \\ \Phi^-(z) &= \frac{q_+(z)}{p_+(z)}\Phi^-(z) = \frac{z - 1}{z} \left(-\frac{z + 1}{z(z - 1)} \right) = -\frac{z + 1}{z^2}. \end{aligned}$$

3°. We have

$$p_+(t) = t, \quad p_-(t) = 1, \quad q_+(t) = t^2 - 1, \quad q_-(t) = 1, \quad m_+ = 1, \quad n_+ = 2, \quad \nu = -1,$$

$$\Psi(z) = \frac{1}{2\pi i} \int_L \frac{\tau}{\tau - z} d\tau + \frac{1}{2\pi i} \int_L \frac{1/[\tau(\tau - 1)]}{\tau - z} d\tau = \begin{cases} z & \text{for } z \in \Omega^+, \\ -\frac{1}{z(z-1)} & \text{for } z \in \Omega^-. \end{cases}$$

The solution of the problem exists only under the solvability conditions (34) or, for the case in question, under the single condition

$$\int_L \frac{q_-(\tau)}{p_-(\tau)} H(\tau) d\tau = 0.$$

On calculating this integral, we obtain

$$\int_L \frac{\tau^3 - \tau^2 + 1}{\tau^2 - \tau} d\tau = \int_L \tau d\tau + \int_L \frac{d\tau}{\tau - 1} - \int_L \frac{d\tau}{\tau} = 0 + 2\pi i - 2\pi i = 0.$$

Thus, the solvability condition holds, and the unique solution of the problem is

$$\Phi^+(z) = z, \quad \Phi^-(z) = -\frac{z+1}{z^2}.$$

4°. We have

$$p_+(t) = 1, \quad p_-(t) = t, \quad q_+(t) = t^2 - 1, \quad q_-(t) = 1, \quad \nu = m_+ - n_+ = -2 < 0.$$

For the solvability of the problem, the following two conditions are necessary:

$$\int_L \frac{q_-(\tau)}{p_-(\tau)} H(\tau) \tau^{k-1} d\tau = 0, \quad k = 1, 2.$$

On calculating the last integral for $k = 1$, we obtain

$$\int_L \frac{\tau^3 - \tau^2 + 1}{\tau(\tau^2 - \tau)} d\tau = \int_L \left(1 - \frac{1}{\tau} - \frac{1}{\tau^2} + \frac{1}{\tau - 1} \right) d\tau = 2\pi i \neq 0.$$

Thus, the solvability condition fails, and hence the problem has no solution.

Note that if we formally calculate the function $\Phi(z)$, then it has a pole at infinity, and hence cannot be a solution of the problem.

14.3-8. Riemann Problem for a Half-Plane.

Let the contour L be the real axis. Just as above, the Riemann problem is to find two bounded analytic functions $\Phi^+(z)$ and $\Phi^-(z)$ in the upper and the lower half-plane, respectively (or a single piecewise analytic function $\Phi(z)$ on the plane), whose limit values on the contour satisfy the boundary condition

$$\Phi^+(x) = D(x)\Phi^-(x) + H(x), \quad -\infty < x < \infty. \quad (35)$$

The given functions $D(x)$ and $H(x)$ satisfy the Hölder condition both at the endpoints and in a neighborhood of the point at infinity on the contour. We also assume that $D(x) \neq 0$.

The main difference from the above case of a finite curve is that here the point at infinity and the origin belong to the contour itself, and therefore cannot be taken as exceptional points at which the canonical function can have a nonzero order. Instead of the auxiliary function t which was used in the above discussion (and has the unit index with respect to L), we use the linear-fractional function on the real axis with the same property:

$$\frac{x - i}{x + i}.$$

The argument of this function

$$\arg \frac{x - i}{x + i} = \arg \frac{(x - i)^2}{x^2 + i} = 2 \arg(x - i)$$

increases by 2π as x ranges over the real axis in the positive direction. Thus,

$$\text{Ind } \frac{x-i}{x+i} = 1.$$

If $\text{Ind } D(x) = \nu$, then the function

$$\left(\frac{x-i}{x+i} \right)^{-\nu} D(x)$$

has zero index. Its logarithm is single-valued on the real axis.

We construct the canonical function for which the point $-i$ is the exceptional point as follows:

$$X^+(z) = e^{G^+(z)}, \quad X^-(z) = \left(\frac{z-i}{z+i} \right)^{-\nu} e^{G^-(z)}, \quad (36)$$

where

$$G(z) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \ln \left[\left(\frac{\tau-i}{\tau+i} \right)^{-\nu} D(\tau) \right] \frac{d\tau}{\tau-z}.$$

Using the limit values of this function, we transform the boundary condition (35) to the form

$$\frac{\Phi^+(x)}{X^+(x)} = \frac{\Phi^-(x)}{X^-(x)} + \frac{H(x)}{X^+(x)}.$$

Next, introducing the analytic function

$$\Psi(z) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{H(\tau)}{X^+(\tau)} \frac{d\tau}{\tau-z}, \quad (37)$$

we represent the boundary condition in the form

$$\frac{\Phi^+(x)}{X^+(x)} - \Psi^+(x) = \frac{\Phi^-(x)}{X^-(x)} - \Psi^-(x).$$

Note that, in contrast with the case of a finite contour, here we have $\Psi^-(\infty) \neq 0$ in general. On applying the theorem on analytic continuation and taking into account the fact that the only possible singularity of the function under consideration is a pole at the point $z = -i$ of order $\leq \nu$ (for $\nu > 0$), on the basis of the generalized Liouville theorem we obtain (see Subsection 14.3-1)

$$\frac{\Phi^+(z)}{X^+(z)} - \Psi^+(z) = \frac{\Phi^-(z)}{X^-(z)} - \Psi^-(z) = \frac{P_\nu(z)}{(z+i)^\nu}, \quad \nu \geq 0,$$

where $P_\nu(z)$ is a polynomial of degree $\leq \nu$ with arbitrary coefficients. This gives the general solution of the problem:

$$\Phi(z) = X(z) \left[\Psi(z) + \frac{P_\nu(z)}{(z+i)^\nu} \right] \quad \text{for } \nu \geq 0, \quad (38)$$

$$\Phi(z) = X(z) [\Psi(z) + C] \quad \text{for } \nu < 0, \quad (39)$$

where C is an arbitrary constant. For $\nu < 0$, the function $X(z)$ has a pole of order $-\nu$ at the point $z = -i$, and therefore for the solvability of the problem we must set $C = -\Psi(-i)$. For $\nu < -1$, the following conditions must additionally hold:

$$\int_{-\infty}^{\infty} \frac{H(x)}{X^+(x)} \frac{dx}{(x+i)^k} = 0, \quad k = 2, 3, \dots, -\nu. \quad (40)$$

Thus, we obtained results similar to those for a finite contour.

Indeed, for $\nu \geq 0$, the homogeneous and nonhomogeneous Riemann boundary value problems for the half-plane are unconditionally solvable, and their solution linearly depends on $\nu + 1$ arbitrary constants. For $\nu < 0$, the homogeneous problem is unsolvable. For $\nu < 0$, the nonhomogeneous problem is uniquely solvable; moreover, in the case $\nu = -1$ the problem is unconditionally solvable, and in the case $\nu < -1$, it is solvable under $-\nu - 1$ solvability conditions (40) only.

Let us also discuss the case of solutions vanishing at infinity. On substituting the relation $\Phi^+(\infty) = \Phi^-(\infty) = 0$ into the boundary condition, we obtain $H(\infty) = 0$. Hence, for a Riemann problem to have a solution that vanishes at infinity, the right-hand side of the boundary condition must vanish at infinity. Assume that this condition is satisfied. To obtain a solution for the case under consideration, we must replace the expression $P_\nu(z)$ in (38) by $P_{\nu-1}(z)$ and equate the constant C in (39) with zero. Thus,

$$\Phi(z) = X(z) \left[\Psi(z) + \frac{P_{\nu-1}(z)}{(z+i)^\nu} \right]. \quad (41)$$

For $\nu \leq 0$, we must set $P_{\nu-1}(z) \equiv 0$ in this formula. We must add another condition to the solvability conditions (40), namely, $\Psi(-i) = 0$, and finally we obtain the following solvability conditions:

$$\int_{-\infty}^{\infty} \frac{H(x)}{X^+(x)} \frac{dx}{(x+i)^k} = 0, \quad k = 1, 2, \dots, -\nu. \quad (42)$$

Now, for $\nu > 0$ we have a solution that depends on ν arbitrary constants. For $\nu \leq 0$, a solution is unique, and for $\nu < 0$, a solution exists if and only if $-\nu$ conditions hold.

14.3-9. Exceptional Cases of the Riemann Problem

In the statement of the Riemann boundary value problem it was required that the coefficient $D(t)$ satisfies the Hölder condition (this prevents infinite values of this coefficient) and vanishes nowhere. As can be observed from the solution (the use of $\ln D(t)$), these restrictions are essential. Now we assume that $D(t)$ vanishes or tends to infinity, with an integer order, at some points of the contour. We assume that the contour L consists of a single closed curve.

Consider the homogeneous problem. We rewrite the boundary condition of the homogeneous Riemann problem in the form

$$\Phi^+(t) = \frac{\prod_{k=1}^{\mu} (t - \alpha_k)^{m_k}}{\prod_{j=1}^{\kappa} (t - \beta_j)^{p_j}} D_1(t) \Phi^-(t). \quad (43)$$

Here α_k ($k = 1, \dots, \mu$) and β_j ($j = 1, \dots, \kappa$) are some points of the contour, m_k and p_j are positive integers, and $D_1(t)$ is a function that is everywhere nonzero and satisfies the Hölder condition. The points α_k are zeros of the function $D(t)$. The points β_j will be called the *poles* of this function. The use of the term “pole” is not completely rigorous because the function $D(t)$ is not analytic. We shall use this term for brevity for a point at which a function (not analytic) tends to infinity with some integer order. We write

$$\text{Ind } D_1(t) = \nu, \quad \sum_{j=1}^{\kappa} p_j = p, \quad \sum_{k=1}^{\mu} m_k = m.$$

We seek the solution in the class of functions bounded on the contour.

Let $X(z)$ be the canonical function of the Riemann problem with coefficient $D_1(t)$. Let us substitute the expression $D_1(t) = X^+(t)/X^-(t)$ into (43) and rewrite the boundary condition in the form

$$\frac{\Phi^+(t)}{X^+(t) \prod_{k=1}^{\mu} (t - \alpha_k)^{m_k}} = \frac{\Phi^-(t)}{X^-(t) \prod_{j=1}^{\kappa} (t - \beta_j)^{p_j}}. \tag{44}$$

To the last relation we apply the theorem on analytic continuation and the generalized Liouville theorem (see Subsection 14.3-1). The points α_k and β_j cannot be singular points of the same analytic function because this would contradict the assumption that $\Phi^+(t)$ or $\Phi^-(t)$ be bounded. Hence, the only possible singularity is the point at infinity. The order at infinity of $X^-(z)$ is ν , and the order of $\prod_{j=1}^{\kappa} (z - \beta_j)^{p_j}$ is equal to $-p$. Hence, the order at infinity of the function $\Phi^-(z)/[X^-(z) \prod_{j=1}^{\kappa} (z - \beta_j)^{p_j}]$ is $-\nu + p$. For $\nu - p \geq 0$ it follows from the generalized Liouville theorem that

$$\frac{\Phi^+(z)}{X^+(z) \prod_{k=1}^{\mu} (z - \alpha_k)^{m_k}} = \frac{\Phi^-(z)}{X^-(z) \prod_{j=1}^{\kappa} (z - \beta_j)^{p_j}} = P_{\nu-p}(z),$$

and hence

$$\Phi^+(z) = X^+(z) \prod_{k=1}^{\mu} (z - \alpha_k)^{m_k} P_{\nu-p}(z), \quad \Phi^-(z) = X^-(z) \prod_{j=1}^{\kappa} (z - \beta_j)^{p_j} P_{\nu-p}(z). \tag{45}$$

If $\nu - p < 0$, then we must set $P_{\nu-p}(z) \equiv 0$, and hence the problem has no solutions.

The boundary value problem with coefficient $D_1(t)$ is called the *reduced problem*. The index ν of the reduced problem will be called the *index* of the original problem. Formulas (45) show that the degree of the occurring polynomial is less by p than the index ν of the problem.

Hence, the number of solutions of problem (43) in the class of functions bounded on the contour is independent of the number of zeros of the coefficient and is diminished by the total number of all poles. In particular, if the index is less than the total order of the poles, then the problem is unsolvable. If the problem is solvable, then its solution can be expressed by formulas (45) in which the canonical function $X(z)$ of the reduced problem can be found by formulas (16) and (17) after replacing $D(t)$ by $D_1(t)$ in these formulas. Under the additional condition $\Phi^-(\infty) = 0$, the number of solutions is diminished by one, and the degree of the polynomial in (45) must be at most $\nu - p - 1$.

Now let us extend the class of solutions by assuming that one of the desired functions $\Phi^+(z)$ and $\Phi^-(z)$ can tend to infinity with integral order at some points of the contour, and at the same time another function remains bounded at these points. We can readily see that this assumption implies no modifications at nonexceptional points. Here the boundedness of one of the functions automatically implies the boundedness of the other. This is not the case for the exceptional points. Let us rewrite the boundary condition (43) in the form

$$\frac{\prod_{j=1}^{\kappa} (t - \beta_j)^{p_j} \Phi^+(t)}{X^+(t)} = \frac{\prod_{k=1}^{\mu} (t - \alpha_k)^{m_k} \Phi^-(t)}{X^-(t)}. \tag{46}$$

Applying the above reasoning and taking into account the fact that the right-hand side has a pole of order $\nu + m$ at infinity, we obtain the general solution in the form

$$\Phi^+(z) = X^+(z) \prod_{k=1}^{\mu} (z - \alpha_k)^{-m_k} P_{\nu+m}(z), \quad \Phi^-(z) = X^-(z) \prod_{j=1}^{\kappa} (z - \beta_j)^{-p_j} P_{\nu+m}(z). \tag{47}$$

Formulas (47) show that in the class of solutions with admissible polar singularity for one of the functions, the number of solutions is greater than that in the class of functions bounded on the contour (for $\nu > 0$) by the total order of all zeros and poles of the coefficient.

We now consider the nonhomogeneous problem. Let us write out the boundary condition in the form

$$\Phi^+(t) = \frac{\prod_{k=1}^{\mu} (t - \alpha_k)^{m_k}}{\prod_{j=1}^{\kappa} (t - \beta_j)^{p_j}} D_1(t) \Phi^-(t) + H(t). \quad (48)$$

We can readily see that the boundary condition cannot be satisfied by finite functions $\Phi^+(t)$ and $\Phi^-(t)$ if we assume that $H(t)$ has poles at points that differ from β_j or if at these points, the orders of the poles of $H(t)$ exceed p_j . Hence, we assume that $H(t)$ can have poles at the points β_j only and that their orders do not exceed p_j . To perform the subsequent reasoning, we must also assume that the functions $D_1(t)$ and $\prod_{j=1}^{\kappa} (t - \beta_j)^{p_j} H(t)$ at the exceptional points are differentiable sufficiently many times.

Just as in the homogeneous problem, we replace $D_1(t)$ by the ratio of the canonical functions $X^+(t)/X^-(t)$ and rewrite the boundary condition (48) in the form

$$\prod_{j=1}^{\kappa} (t - \beta_j)^{p_j} \frac{\Phi^+(t)}{X^+(t)} = \prod_{k=1}^{\mu} (t - \alpha_k)^{m_k} \frac{\Phi^-(t)}{X^-(t)} + \prod_{j=1}^{\kappa} (t - \beta_j)^{p_j} \frac{H(t)}{X^+(t)}. \quad (49)$$

On replacing the function defined by the second summand on the right-hand side in (49) by the difference of the boundary values of analytic functions

$$\prod_{j=1}^{\kappa} (t - \beta_j)^{p_j} \frac{H(t)}{X^+(t)} = \Psi^+(t) - \Psi^-(t),$$

where

$$\Psi(z) = \frac{1}{2\pi i} \int_L \prod_{j=1}^{\kappa} (\tau - \beta_j)^{p_j} \frac{H(\tau)}{X^+(\tau)} \frac{d\tau}{\tau - z}, \quad (50)$$

we reduce the boundary condition to the form

$$\prod_{j=1}^{\kappa} (t - \beta_j)^{p_j} \frac{\Phi^+(t)}{X^+(t)} - \Psi^+(t) = \prod_{k=1}^{\mu} (t - \alpha_k)^{m_k} \frac{\Phi^-(t)}{X^-(t)} - \Psi^-(t).$$

On applying the theorem on analytic continuation and the generalized Liouville theorem (see Subsection 14.3-1), we obtain

$$\Phi^+(z) = \frac{X^+(z)}{\prod_{j=1}^{\kappa} (z - \beta_j)^{p_j}} [\Psi^+(z) + P_{\nu+m}(z)], \quad \Phi^-(z) = \frac{X^-(z)}{\prod_{k=1}^{\mu} (z - \alpha_k)^{m_k}} [\Psi^-(z) + P_{\nu+m}(z)]. \quad (51)$$

In general, the last formulas give solutions that can tend to infinity at the points α_k and β_k . For a solution to be bounded it is necessary that the function $\Psi^+(z) + P_{\nu+m}(z)$ have zeros of orders p_j at the points β_j and the function $\Psi^-(z) + P_{\nu+m}(z)$ have zeros of orders m_k at the points α_k . These requirements form $m+p$ conditions for the coefficients of the polynomial $P_{\nu+m}(z)$. If the coefficients

of the polynomial $P_{\nu+m}(z)$ are chosen in accordance with the above conditions, then formulas (51) give a solution of the nonhomogeneous problem (48) in the class of bounded functions.

Consider another way of constructing a solution, which is more convenient and based on the construction of a special particular solution.

By the *canonical function* $Y(z)$ of the nonhomogeneous problem we mean a piecewise analytic function that satisfies the boundary condition (48), has zero order everywhere in the finite part of the domain (including the points α_k and β_j), and has the least possible order at infinity.

In the construction of the canonical function, we start from the solution given by formulas (51). Let us construct a polynomial $\mathcal{U}_n(z)$ that satisfies the following conditions:

$$\begin{aligned} \mathcal{U}_n^{(i)}(\beta_j) &= \Psi^{+(i)}(\beta_j), & i &= 0, 1, \dots, p_j - 1, & j &= 1, \dots, \kappa, \\ \mathcal{U}_n^{(l)}(\alpha_k) &= \Psi^{-(l)}(\alpha_k), & l &= 0, 1, \dots, m_k - 1, & k &= 1, \dots, \mu, \end{aligned}$$

where $\Psi^{+(i)}(\beta_j)$ and $\Psi^{-(l)}(\alpha_k)$ are the values of the i th and the l th derivatives at the corresponding points. Thus, $\mathcal{U}_n(z)$ is the Hermite interpolation polynomial for the functions

$$\Psi(z) = \begin{cases} \Psi^+(z) & \text{at the points } \beta_j, \\ \Psi^-(z) & \text{at the points } \alpha_k \end{cases}$$

with interpolation nodes β_j and α_k of multiplicities p_j and m_k , respectively (see Subsection 14.3-2). Such a polynomial is uniquely determined, and its degree is at most $n = m + p - 1$.

The canonical function of the nonhomogeneous problem can be expressed via the interpolation polynomial as follows:

$$Y^+(z) = X^+(z) \frac{\Psi^+(z) - \mathcal{U}_n(z)}{\prod_{j=1}^{\kappa} (z - \beta_j)^{p_j}}, \quad Y^-(z) = X^-(z) \frac{\Psi^-(z) - \mathcal{U}_n(z)}{\prod_{k=1}^{\mu} (z - \alpha_k)^{m_k}}. \quad (52)$$

To construct the general solution of the nonhomogeneous problem (48), we use the fact that this general solution is the sum of a particular solution of the nonhomogeneous problem and of the general solution of the homogeneous problem. Applying formulas (47) and (52), we obtain

$$\begin{aligned} \Phi^+(z) &= Y^+(z) + X^+(z) \prod_{k=1}^{\mu} (z - \alpha_k)^{m_k} P_{\nu-p}(z), \\ \Phi^-(z) &= Y^-(z) + X^-(z) \prod_{j=1}^{\kappa} (z - \beta_j)^{p_j} P_{\nu-p}(z). \end{aligned} \quad (53)$$

For the case in which $\nu - p < 0$, we must set $P_{\nu-p}(z) \equiv 0$. Applying formula (52), we readily find that the order of $Y^-(z)$ at infinity is equal to $\nu - p + 1$. If $\nu < p - 1$, then $Y^-(z)$ has a pole at infinity, and the canonical function is no longer a solution of the nonhomogeneous problem.

However, on subjecting the constant term $H(t)$ to $p - \nu - 1$ conditions, we can increase the order of the functions $Y(z)$ at infinity by $p - \nu - 1$ and thus again make the canonical function $Y(z)$ a solution of the nonhomogeneous problem. Obviously, to this end it is necessary and sufficient that in the expansion of the function $\Psi(z) - \mathcal{U}_n(z)$ in a neighborhood of the point at infinity, the first $p - \nu - 1$ coefficients be zero. This gives just $p - \nu - 1$ solvability conditions of the problem for the case under consideration. Let us clarify the character of these conditions. The expansion of $\Psi(z) - \mathcal{U}_n(z)$ can be represented in the form

$$\Psi(z) - \mathcal{U}_n(z) = -a_n z^n - a_{n-1} z^{n-1} - \dots - a_0 + a_{-1} z^{-1} + a_{-2} z^{-2} + \dots + a_{-k} z^{-k} + \dots,$$

where a_0, a_1, \dots, a_n are the coefficients of the polynomial $\mathcal{U}_n(z)$, and the a_{-k} are the coefficients of the expansion of the function $\Psi(z)$, which are given by the obvious formula

$$a_{-k} = -\frac{1}{2\pi i} \int_L \prod_{j=1}^k (\tau - \beta_j)^{p_j} \frac{H(\tau)\tau^{k-1}}{X^+(\tau)} d\tau.$$

The solvability conditions acquire the form

$$a_n = a_{n-1} = \dots = a_{n-p+\nu+2} = 0.$$

If a solution must satisfy the additional condition $\Phi^-(\infty) = 0$, then, for $\nu - p > 0$, in formulas (53) we must take the polynomial $P_{\nu-p-1}(z)$, and for $\nu - p < 0$, $p - \nu$ conditions must be satisfied.

14.3-10. Riemann Problem for a Multiply Connected Domain.

Let $L = L_0 + L_1 + \dots + L_m$ be a collection of $m + 1$ disjoint contours, and let the interior of the contour L_0 contain the other contours. By Ω^+ we denote the $(m + 1)$ -connected domain interior for L_0 and exterior for L_1, \dots, L_m . By Ω^- we denote the complement of $\Omega^+ + L$ in the entire complex plane. To be definite, we assume that the origin lies in Ω^+ . The positive direction of the contour L is that for which the domain Ω^+ remains to the left, i.e., the contour L_0 must be traversed counterclockwise and the contours L_1, \dots, L_m , clockwise.

We first note that the jump problem

$$\Phi^+(t) - \Phi^-(t) = H(t)$$

is solved by the same formula

$$\Phi(z) = \frac{1}{2\pi i} \int_L \frac{H(\tau) d\tau}{\tau - z}$$

as in the case of a simply connected domain. This follows from the Sokhotski–Plemelj formulas, which have the same form for a multiply connected domain as for a simply connected domain.

The Riemann problem (homogeneous and nonhomogeneous) can be posed in the same way as for a simply connected domain.

We write $\nu_k = \frac{1}{2\pi} [\arg D(t)]_{L_k}$ (all contours are passed in the positive direction). By the *index of the problem* we mean the number

$$\nu = \sum_{k=0}^m \nu_k. \tag{54}$$

If ν_k ($k = 1, \dots, m$) are zero for the inner contours, then the solution of the problem has just the same form as for a simply connected domain.

To reduce the general case to the simple one, we introduce the function

$$\prod_{k=1}^m (t - z_k)^{\nu_k},$$

where the z_k are some points inside the contours L_k ($k = 1, \dots, m$). Taking into account the fact that $[\arg(t - z_k)]_{L_j} = 0$ for $k \neq j$ and $[\arg(t - z_j)]_{L_j} = -2\pi$, we obtain

$$\frac{1}{2\pi} \left[\arg \prod_{k=1}^m (t - z_k)^{\nu_k} \right]_{L_j} = \frac{1}{2\pi} [\arg(t - z_j)^{\nu_j}]_{L_j} = -\nu_j, \quad j = 1, \dots, m.$$

Hence,

$$\left[\arg \left(D(t) \prod_{k=1}^m (t - z_k)^{\nu_k} \right) \right]_{L_j} = 0, \quad j = 1, \dots, m.$$

Let us calculate the increment of the argument of the function $D(t) \prod_{k=1}^m (t - z_k)^{\nu_k}$ with respect to the contour L_0 :

$$\frac{1}{2\pi} \left[\arg \left(D(t) \prod_{k=1}^m (t - z_k)^{\nu_k} \right) \right]_{L_0} = \frac{1}{2\pi} [\arg D(t)]_{L_0} + \frac{1}{2\pi} \sum_{k=1}^m [\nu_k \arg(t - z_k)]_{L_0} = \nu_0 + \sum_{k=1}^m \nu_k = \nu.$$

Since the origin belongs to the domain Ω^+ , it follows that

$$[\arg t]_{L_k} = 0, \quad k = 1, \dots, m, \quad [\arg t]_{L_0} = 2\pi.$$

Therefore,

$$\left[\arg \left(t^{-\nu} \prod_{k=1}^m (t - z_k)^{\nu_k} D(t) \right) \right]_L = 0. \quad (55)$$

1°. *The Homogeneous Problem.* Let us rewrite the boundary condition

$$\Phi^+(t) = D(t)\Phi^-(t) \quad (56)$$

in the form

$$\Phi^+(t) = \frac{t^\nu}{\prod_{k=1}^m (t - z_k)^{\nu_k}} \left(t^{-\nu} \prod_{k=1}^m (t - z_k)^{\nu_k} D(t) \right) \Phi^-(t). \quad (57)$$

The function $t^{-\nu} \prod_{k=1}^m (t - z_k)^{\nu_k} D(t)$ has zero index on each of the contours L_k ($k = 1, \dots, m$), and hence it can be expressed as the ratio

$$t^{-\nu} \prod_{k=1}^m (t - z_k)^{\nu_k} D(t) = \frac{e^{G^+(t)}}{e^{G^-(t)}}, \quad (58)$$

where

$$G(z) = \frac{1}{2\pi i} \int_L \ln \left(\tau^{-\nu} \prod_{k=1}^m (\tau - z_k)^{\nu_k} D(\tau) \right) \frac{d\tau}{\tau - z}. \quad (59)$$

The canonical function of the problem is given by the formulas

$$X^+(z) = \prod_{k=1}^m (z - z_k)^{-\nu_k} e^{G^+(z)}, \quad X^-(z) = z^{-\nu} e^{G^-(z)}. \quad (60)$$

Now the boundary condition (57) can be rewritten in the form

$$\frac{\Phi^+(t)}{X^+(t)} = \frac{\Phi^-(t)}{X^-(t)}.$$

As usual, by applying the theorem on analytic continuation and the generalized Liouville theorem (see Subsection 14.3-1), we obtain

$$\Phi^+(z) = \prod_{k=1}^m (z - z_k)^{-\nu_k} e^{G^+(z)} P_\nu(z), \quad \Phi^-(z) = z^{-\nu} e^{G^-(z)} P_\nu(z). \tag{61}$$

We can see that this solution differs from the above solution of the problem for a simply connected domain only in that the function $\Phi^+(z)$ has the factor $\prod_{k=1}^m (z - z_k)^{-\nu_k}$. Under the additional condition $\Phi^-(\infty) = 0$, in formulas (61) we must take the polynomial $P_{\nu-1}(z)$.

Applying the Sokhotski–Plemelj formulas, we obtain

$$G^\pm(t) = \pm \frac{1}{2} \ln[t^{-\nu} \Pi(t) D(t)] + G(t),$$

where $G(t)$ is the Cauchy principal value of the integral (59) and

$$\Pi(t) = \prod_{k=1}^m (t - z_k)^{\nu_k}.$$

On passing to the limit as $z \rightarrow t$ in formulas (60) we obtain

$$X^+(t) = \sqrt{\frac{D(t)}{t^\nu \Pi(t)}} e^{G(t)}, \quad X^-(t) = \frac{1}{\sqrt{t^\nu \Pi(t) D(t)}} e^{G(t)}. \tag{62}$$

The sign of the root is determined by the (arbitrary) choice of a branch of the function $\ln[t^{-\nu} \Pi(t) D(t)]$.

2°. *The Nonhomogeneous Problem.* By the same reasoning as above, we represent the boundary condition

$$\Phi^+(t) = D(t) \Phi^-(t) + H(t) \tag{63}$$

in the form

$$\frac{\Phi^+(t)}{X^+(t)} - \Psi^+(t) = \frac{\Phi^-(t)}{X^-(t)} - \Psi^-(t),$$

where $\Psi(z)$ is defined by the formula

$$\Psi(z) = \frac{1}{2\pi i} \int_L \frac{H(\tau)}{X^+(\tau)} \frac{d\tau}{\tau - z}.$$

This gives the general solution

$$\Phi(z) = X(z)[\Psi(z) + P_\nu(z)] \tag{64}$$

or

$$\Phi(z) = X(z)[\Psi(z) + P_{\nu-1}(z)], \tag{65}$$

if the solution satisfies the condition $\Phi^-(\infty) = 0$.

For $\nu < 0$, the nonhomogeneous problem is solvable if and only if the following conditions are satisfied:

$$\int_L \frac{H(t)}{X^+(t)} t^{k-1} dt = 0, \tag{66}$$

where k ranges from 1 to $-\nu - 1$ if we seek solutions bounded at infinity and from 1 to $-\nu$ if we assume that $\Phi^-(\infty) = 0$.

Under conditions (66), the solution can also be found from formulas (64) or (65) by setting $P_\nu \equiv 0$.

If the external contour L_0 is absent and the domain Ω^+ is the plane with holes, then the main difference from the preceding case is that here the zero index with respect to all contours L_k ($k = 1, \dots, m$) is attained by the function $\prod_{k=1}^m (t - z_k)^{\nu_k} D(t)$ that does not involve the factor $t^{-\nu}$.

Therefore, to obtain a solution to the problem, it suffices to repeat the above reasoning on omitting this factor.

14.3-11. Riemann Problem for Open Curves.

Let L be a curve that consists of m simple smooth open curves (arcs) $L_k = a_k b_k$, $k = 1, 2, \dots, m$, without common interior or endpoints. Let $D(t), H(t)$ be two functions defined on L and satisfying the Hölder condition on each arc L_k , and suppose that $D(t) \neq 0$ for all t . On different arcs L_k the functions $D(t), H(t)$ can be defined by the analytical formulas:

$$D(t) = D_k(t), \quad H(t) = H_k(t), \quad t \in L_k, \quad k = 1, 2, \dots, m.$$

The arc L_k is directed from the point a_k to the point b_k .

For the points a_k, b_k we will use a unified notation c_j , so that the set of all c -points consists of $2m$ ordered points c_1, c_2, \dots, c_{2m} , each coinciding with some a_k or b_k , but their order may be different from that of the sequences a_k and b_k . For instance, one can take

$$\{c_1; c_2; c_3; c_4; \dots; c_{2m-1}; c_{2m}\} = \{a_1; b_1; a_2; b_2; \dots; a_m; b_m\},$$

or

$$\{c_1; c_2; \dots; c_m; c_{m+1}; c_{m+2}; \dots; c_{2m}\} = \{a_1; a_2; \dots; a_m; b_1; b_2; \dots; b_m\},$$

or some other combination of a_k, b_k .

On each arc $L_k = a_k b_k$, we fix some continuous branch of the function $\ln D(t) = \ln D_k(t)$, $t \in L_k$ by the condition

$$0 \leq \operatorname{Im} \ln D(a_k) < 2\pi \quad \iff \quad 0 \leq \arg D(a_k) < 2\pi. \quad (67)$$

Then

$$\ln D(b_k) = \ln D(a_k) + i [\arg D(t)]_{L_k} = \ln D(a_k) + i \int_{L_k} d \arg D(t). \quad (68)$$

Note that a branch $\ln D(t)$ on L_k may be fixed by other conditions. For instance, instead of (67), one can take $-\pi < \arg D(a_k) \leq \pi$ or $0 \leq \arg D(b_k) < 2\pi$, or some other condition.

For each point c_k , we calculate the numbers

$$\alpha_k + i\beta_k = \mp \frac{\ln D(c_k)}{2\pi i}, \quad k = 1, 2, \dots, 2m, \quad (69)$$

where the upper minus corresponds to c_k coinciding with some a_j and the lower plus corresponds to c_k coinciding with some b_j . The points c_k , as well as the corresponding endpoints a_j, b_j of the curve L , for which α_k are integer numbers, are called *singular*, while the other c_k and the corresponding endpoints of the curve L are called *nonsingular*. Clearly, c_k is a singular point if and only if $D(c_k)$ is real and positive.

Let us renumber the points c_1, c_2, \dots, c_{2m} so that the first and the second groups of subscripts would respectively designate nonsingular and singular points. Let c_1, c_2, \dots, c_n ($0 \leq n \leq 2m$) be all nonsingular endpoints of the curve L . From these points, we choose p ($0 \leq p \leq n$) points and move them to the first p places; we may assume these to be c_1, c_2, \dots, c_p (after renumbering, if necessary).

THE RIEMANN PROBLEM. Find a function $\Phi(z)$ which is analytic on the entire plane outside the curve L , on which it has continuous boundary values

$$\Phi^+(t), \quad \Phi^-(t), \quad t \in L \setminus \{\text{endpoints}\},$$

satisfying the boundary condition (7) or (8), bounded near the nonsingular endpoints c_1, c_2, \dots, c_p , and admitting integrable singularities near the other nonsingular points, i.e.,

$$|\Phi(z)| \leq \frac{M_k}{|z - c_k|^{\lambda_k}}, \quad M_k = \text{const}, \quad \lambda_k = \text{const}, \quad 0 \leq \lambda_k < 1 \quad \text{near } c_k, \quad k = p+1, \dots, n.$$

In this case, in contrast to the Riemann problem for closed curves, the boundary condition (7) or (8) should hold only at the points other than the endpoints, near which the sought function should have a prescribed behavior. Moreover, the behavior of the sought function is prescribed only near nonsingular endpoints and is left unspecified near singular endpoints, since any solution of the homogeneous problem (7) near a singular endpoint c_k is always bounded and a solution of the nonhomogeneous problem for $\beta_k \neq 0$ is bounded and for $\beta_k = 0$ has a logarithmic singularity. If c_k is a nonsingular endpoint, then any solution of the homogeneous Riemann problem always vanishes at the point c_k , while a solution of the nonhomogeneous problem will only be bounded.

A solution of the above Riemann problem is called a *solution of class h_p or class $h(c_1, c_2, \dots, c_p)$* if nonsingular endpoints c_1, c_2, \dots, c_p are fixed a priori. The class h_0 consists of all solutions of the problem that admit integrable singularities near all n nonsingular points of the line L . This class contains all other classes $h_p, 1 \leq p \leq n$. The class h_n belongs to all other classes $h_p, 0 \leq p \leq n-1$, and consists of all solutions of the Riemann problem that are bounded near all nonsingular endpoints of the line L .

As in the case of one or several closed curves, let us construct a particular solution $X(z)$ of the homogeneous problem (7), which, in addition, does not vanish on the entire plane including the edges of the cuts $L_k = a_k b_k$, except at the endpoints of the arcs near which its behavior is determined by the class $h_p = h(c_1, c_2, \dots, c_p)$. Consider the Cauchy integral

$$\Gamma(z) = \frac{1}{2\pi i} \int_L \frac{\ln D(\tau) d\tau}{\tau - z} = \sum_{k=1}^m \frac{1}{2\pi i} \int_{a_k}^{b_k} \frac{\ln D_k(\tau) d\tau}{\tau - z}, \tag{70}$$

where $\ln D_k(t)$ are the logarithmic branches fixed above. This integral has a discontinuity on the curve L with the jump

$$\Gamma^+(t) - \Gamma^-(t) = \ln D(t), \quad t \in L \setminus \{c_1, c_2, \dots, c_{2m}\},$$

and near the endpoints c_k admits the representation

$$\Gamma(z) = (\alpha_k + i\beta_k) \ln(z - c_k) + \Gamma_*(z).$$

Here, $\ln(z - c_k)$ is a branch which is single-valued on the plane with the cut joining the points c_k and ∞ and going along the arc L_j with an endpoint at c_k ; the function $\Gamma_*(z)$ is analytic in a small neighborhood of c_k with the cut along L_j and tends to a certain limit as $z \rightarrow c_k$ along any path. Therefore, the function

$$X(z) = e^{\Gamma(z)} \prod_{k=1}^{2m} (z - c_k)^{-\nu_k}, \tag{71}$$

where ν_k are integers such that

$$\begin{aligned} 0 < \alpha_k - \nu_k < 1 &\iff \nu_k = [\alpha_k], & k = 1, 2, \dots, p, \\ -1 < \alpha_k - \nu_k < 0 &\iff \nu_k = 1 + [\alpha_k], & k = p + 1, \dots, n, \\ \alpha_k - \nu_k = 0 &\iff \nu_k = \alpha_k, & k = n + 1, \dots, 2m \end{aligned} \tag{72}$$

($[\alpha_k]$ is the integer part of α_k), has all the above properties of a particular solution of the homogeneous problem (7): both functions $X(z)$ and $1/X(z)$ are analytic on the plane with the cut along L , on which $X^\pm(t) \neq 0, X^+(t) = D(t)X^-(t), t \in L \setminus \{c_k\}$, and

$$\begin{aligned} X(z) &\sim A_k(z - c_k)^{\lambda_k} \text{ as } z \rightarrow c_k, & \lambda_k = \alpha_k - \nu_k, & k = 1, 2, \dots, n, \\ X(z) &\sim A_k \text{ as } z \rightarrow c_k, & k = n + 1, \dots, 2m, \end{aligned}$$

where $A_k = \text{const} \neq 0$ for all k , and $0 < \lambda_k < 1$ for $k = 1, 2, \dots, p$, and $-1 < \lambda_k < 0$ for $k = p+1, \dots, n$. At ∞ , this function is of the order

$$\nu = \nu_1 + \nu_2 + \dots + \nu_{2m},$$

i.e., $X(z) \sim Az^{-\nu}$, $A = \text{const} \neq 0$ as $z \rightarrow \infty$.

The function $X(z)$ is called a *canonical function of a problem of class $h_p = h(c_1, c_2, \dots, c_p)$* , and the integer ν is called the *index of a problem of class h_p* .

With the help of the canonical function $X(z)$, the Riemann problem for open curves is solved along the same lines as in the case of a simply-connected domain.

1°. *The jump problem*

$$\Phi^+(t) - \Phi^-(t) = H(t), \quad t \in L \setminus \{\text{endpoints}\}$$

in the class of functions vanishing at ∞ (independently of the class h_p) has a unique solution,

$$\Phi(z) = \frac{1}{2\pi i} \int_L \frac{H(\tau) d\tau}{\tau - z} = \sum_{k=1}^m \frac{1}{2\pi i} \int_{a_k}^{b_k} \frac{H_k(\tau) d\tau}{\tau - z}.$$

In order to obtain a solution bounded at ∞ or with a pole at ∞ of an order $\leq \mu$, one should take the sum of $\Phi(z)$ and an arbitrary constant C or an arbitrary polynomial of degree μ , respectively.

2°. *The homogeneous problem* (7), with the help of the representation (18), is reduced to the construction of a function $\Phi(z)/X(z)$ which is analytic on the entire plane and has removable singularities at all the endpoints c_k . A solution of this problem that vanishes at ∞ and belongs to the class $h_p = h(c_1, c_2, \dots, c_p)$, for $\nu \geq 0$, is given by the formula

$$\Phi(z) = P_\nu(z)X(z),$$

where $P_\nu(z)$ is an arbitrary polynomial of degree ν . For $\nu < 0$, the homogeneous problem has no nontrivial solutions.

3°. *The nonhomogeneous problem* (8), with the help of the transformation (18), is reduced to the jump problem (22), and its solution of class $h_p = h(c_1, c_2, \dots, c_p)$ decaying at ∞ for $\nu \geq 0$ is again given by formulas (23), (28). For $\nu < 0$, a nontrivial solution of class h_p exists and is unique, provided that $-\nu$ solvability conditions (29) are satisfied; the solution has the form $\Phi(z) = X(z)\Psi(z)$, where $X(z)$ is a canonical function of class h_p , and $\Psi(z)$ is the integral (23).

In order to obtain a solution of the homogeneous or the nonhomogeneous problem bounded at ∞ or with a pole at ∞ of order $< \mu$, one should replace ν by $\nu + 1$ or $\nu + \mu$, respectively.

Remark. If $X_0(z)$ is a canonical function of the widest class h_0 , then

$$X(z) = (z - c_1)(z - c_2) \dots (z - c_p)X_0(z)$$

is a canonical function of class $h_p = h(c_1, c_2, \dots, c_p)$. A similar relation holds for canonical functions of any two classes h_p and h_q . Thus, for the construction of a canonical function of class h_p , it suffices to construct a canonical function of any other class h_q , in particular, h_0 .

In order to obtain the canonical function $X_n(z)$ of the narrowest class h_n , one should take $\nu_k = [\alpha_k]$ for all $k = 1, 2, \dots, 2m$ in (71). This function is bounded near all endpoints of the line L , both singular and nonsingular. In terms of $X_n(z)$, the canonical function of class $h_p = h(c_1, c_2, \dots, c_p)$ is found by the formula

$$X(z) = (z - c_{p+1})^{-1} \dots (z - c_n)^{-1} X_n(z).$$

Example 3. Let the line L consist of a segment $L_1 = [a; b]$ ($a > 0$) of the real axis and the segment $L_2 = [2\pi i; 3\pi i]$ of the imaginary axis, and let

$$D(t) = \begin{cases} it & \text{if } t \in L_1, \\ e^t & \text{if } t \in L_2. \end{cases}$$

Let us find possible classes h_p of solutions of the Riemann problem and construct the canonical function in these classes.

1) Let us fix the branches

$$\ln D(t) = \ln(it) = \ln(t) + \frac{\pi i}{2}, \quad t \in [a; b],$$

$$\ln D(t) = \ln(e^t) = t - 2\pi i, \quad t \in [2\pi i; 3\pi i],$$

so that the values

$$\ln D(a) = \ln(a) + \frac{\pi i}{2}, \quad \ln D(2\pi i) = 0$$

satisfy condition (67). We have

$$\ln D(b) = \ln(b) + \frac{\pi i}{2}, \quad \ln D(3\pi i) = \pi i.$$

2) Taking $c_1 = a, c_2 = b, c_3 = 2\pi i, c_4 = 3\pi i$, let us calculate the numbers

$$\alpha_1 + i\beta_1 = -\frac{\ln D(a)}{2\pi i} = -\frac{1}{4} + i\frac{\ln a}{2\pi}, \quad \alpha_2 + i\beta_2 = \frac{\ln D(b)}{2\pi i} = \frac{1}{4} - i\frac{\ln b}{2\pi},$$

$$\alpha_3 + i\beta_3 = -\frac{\ln D(2\pi i)}{2\pi i} = 0, \quad \alpha_4 + i\beta_4 = \frac{\ln D(3\pi i)}{2\pi i} = \frac{1}{2}.$$

Since $\alpha_3 = 0$ is integer and all the other α_k are noninteger, the endpoint $c_3 = 2\pi i$ is singular and the rest of the endpoints c_1, c_2, c_4 are nonsingular. In this connection, let us renumber the points c_k as follows: $c_1 = a, c_2 = b, c_3 = 3\pi i, c_4 = 2\pi i$, and for these we have the new

$$\alpha_1 = -\frac{1}{4}, \quad \alpha_2 = \frac{1}{4}, \quad \alpha_3 = \frac{1}{2}, \quad \alpha_4 = 0.$$

3) All possible classes of solutions of the Riemann problem (and therefore, the classes of the canonical function) are determined by the points c_1, c_2, c_3 . These classes are the following: $h_0, h(c_1), h(c_2), h(c_3), h(c_1, c_2), h(c_1, c_3), h(c_2, c_3), h_3 = h(c_1, c_2, c_3)$, with h_0 being the widest class and $h_3 = h(a, b, 3\pi i)$ the narrowest class.

4) Let us construct the canonical function $X_0(z)$ of class h_0 . In view of (72), we have

$$\nu_1 = 1 + [\alpha_1] = 0, \quad \nu_2 = 1 + [\alpha_2] = 1, \quad \nu_3 = 1 + [\alpha_3] = 1, \quad \nu_4 = \alpha_4 = 0,$$

and by (70) and (71),

$$X_0(z) = e^{\Gamma(z)}(z - b)^{-1}(z - 3\pi i)^{-1},$$

$$\Gamma(z) = \frac{1}{2\pi i} \int_a^b \left(\ln \tau + \frac{\pi i}{2} \right) \frac{d\tau}{\tau - z} + \frac{1}{2\pi i} \int_{2\pi i}^{3\pi i} \frac{\tau - 2\pi i}{\tau - z} d\tau,$$

and therefore,

$$X_0(z) = \frac{\sqrt{e}(\tau - 2\pi i)}{(z - b)(\tau - 3\pi i)^2} \left(\frac{z - b}{z - a} \right)^{\frac{1}{4}} \left(\frac{z - 3\pi i}{z - 2\pi i} \right)^{\frac{z}{2\pi i}} \exp \left(\frac{1}{2\pi i} \int_a^b \frac{\ln \tau}{\tau - z} d\tau \right),$$

where we have chosen branches (of the multiple-valued functions involved) which are single-valued on the plane with cuts along the segments $L_1 = [a; b]$ and $L_2 = [2\pi i; 3\pi i]$, respectively, and take the value 1 at ∞ .

5) According to the above remark, the canonical function of class $h(a)$ is obtained from $X_0(z)$ by its multiplication by $z - a$, and the canonical function of class $h(b)$ is obtained by multiplying $X_0(z)$ by $z - b$, etc. These functions can be found directly with the help of formulas (70)–(72).

4°. *The case of a piecewise constant coefficient of the problem.* The canonical function of the Riemann problem for open curves can be found explicitly, provided that its coefficient $D(t)$ takes constant values on the arcs $L_k = a_k b_k$ (the values may be different on different arcs). Let

$$D(t) = D_k, \quad D_k = \text{const} \neq 0, \quad t \in a_k b_k, \quad k = 1, 2, \dots, m.$$

Then, according to (67)–(69), we have

$$\ln D(a_k) = \ln D(b_k) = \ln D_k, \quad \alpha_k + i\beta_k = \mp(\gamma_k - i\delta_k),$$

$$\gamma_k = \frac{\arg D_k}{2\pi}, \quad \delta_k = \frac{\ln |D_k|}{2\pi}, \quad 0 \leq \arg D_k < 2\pi, \tag{73}$$

where the upper sign corresponds to the initial points a_k , and the lower sign corresponds to the endpoints b_k of the arcs L_k . Thus for each specific arc L_k , its endpoints a_k, b_k are singular or nonsingular simultaneously, depending on whether γ_k is integer or noninteger, and if γ_k is integer, it must be equal to zero. Therefore, the number of nonsingular endpoints, as well as of singular ones, is always even.

Suppose that $a_1, b_1, \dots, a_n, b_n$ ($0 \leq n \leq m$) are nonsingular endpoints and $a_{n+1}, b_{n+1}, \dots, a_m, b_m$ are singular endpoints of the line L . For this distribution of nonsingular and singular endpoints, it might be necessary to renumber the arcs L_k . According to (70)–(72), the canonical function $X_0(z)$ of class h_0 (i.e., for this function, all nonsingular endpoints are infinity points of order < 1 , and this function is bounded near all singular endpoints) is given by the formula

$$X_0(z) = X_n(z) \prod_{k=1}^n \frac{1}{z - b_k}, \quad X_n(z) = \prod_{k=1}^n \left(\frac{z - b_k}{z - a_k} \right)^{\gamma_k + i\delta_k} \prod_{k=n+1}^m \left(\frac{z - b_k}{z - a_k} \right)^{i\delta_k}, \quad (74)$$

where the multiple-valued functions in the last two products are replaced by their branches that take the value 1 at ∞ and are single-valued on the plane along the arcs $L_k = a_k b_k$. Since $0 \leq \gamma_k < 1$ for all k , it can be seen that $X_n(z)$ is a canonical function of class $h(b_1, b_2, \dots, b_n)$. This function is bounded near all endpoints b_k (both singular and nonsingular); the points a_1, a_2, \dots, a_n are its infinity points of order < 1 , and near the points a_{n+1}, \dots, a_m (if these exist) it is also bounded.

In order to obtain a canonical function of class $h(c_1, c_2, \dots, c_p)$, where c_k is the general notation for a_j, b_j , one should multiply $X_0(z)$ by $(z - c_1)(z - c_2) \dots (z - c_p)$. Taking different systems c_k , one obtains, in particular, the following canonical functions:

$$\begin{aligned} X_{n*}(z) &= X_n(z) \prod_{k=1}^n \frac{z - a_k}{z - b_k} = \\ &= \prod_{k=1}^m \left(\frac{z - a_k}{z - b_k} \right)^{1 - \gamma_k - i\delta_k} \prod_{k=m+1}^n \left(\frac{z - b_k}{z - a_k} \right)^{i\delta_k}, \quad X_{n*}(z) \in h(a_1, a_2, \dots, a_n), \end{aligned} \quad (75)$$

$$X_{2n}(z) = X_n(z) \prod_{k=1}^n (z - a_k), \quad X_{2n}(z) \in h_{2n}, \quad h_{2n} = h(a_1, b_1, \dots, a_n, b_n)$$

etc. The last function in (75) is bounded near all endpoints of the curve L .

5°. *The case of a constant coefficient of the problem.* Let

$$D(t) = D_0, \quad D_0 = \text{const}, \quad t \in a_k b_k, \quad k = 1, 2, \dots, m.$$

Then, for nonreal or negative real D_0 , formulas (73)–(75) yield

$$\begin{aligned} X_m(z) &= \left(\prod_{k=1}^m \frac{z - b_k}{z - a_k} \right)^{\gamma + i\delta}, \quad \gamma = \frac{\arg D_0}{2\pi}, \quad \delta = \frac{\ln |D_0|}{2\pi}, \\ 0 &< \arg D_0 < 2\pi, \quad X_m(z) \in h(b_1, b_2, \dots, b_n); \\ X_{m*}(z) &= \left(\prod_{k=1}^m \frac{z - a_k}{z - b_k} \right)^{\gamma + i\delta}, \quad X_{m*}(z) \in h(a_1, a_2, \dots, a_n); \\ X_0(z) &= X_m(z) \prod_{k=1}^m \frac{1}{z - b_k}, \quad X_0(z) \in h_0; \\ X_{2m}(z) &= X_m(z) \prod_{k=1}^m (z - a_k), \quad X_{2m}(z) \in h_{2m}, \quad h_{2m} = h(a_1, b_1, \dots, a_m, b_m). \end{aligned} \quad (76)$$

The function $X_m(z)$ is bounded near all points b_k , and a_k are its infinity points of an order < 1 . Conversely, the function $X_{m^*}(z)$ is bounded near a_k , and at the points b_k has integrable singularities. For the function $X_0(z)$, the endpoints a_k, b_k are infinity points of integrable character. The function $X_{2m}(z)$ is bounded near all endpoints a_k, b_k .

If $D_0 \neq 1$ is a real positive number, then all endpoints of the curve L are singular and different classes h_p cannot be defined for the Riemann problem. In this case, there is a single (to within a nonzero constant coefficient) canonical function

$$X(z) = \left(\prod_{k=1}^m \frac{z - b_k}{z - a_k} \right)^{i\delta}, \quad \delta = \frac{1}{2\pi} \ln |D_0|,$$

which is bounded near all endpoints of the curve, although for $z \rightarrow a_k$ and $z \rightarrow b_k$ it has no limits.

In applications, one often encounters the Riemann problem with

$$\Phi^+(t) - \Phi^-(t) = H(t), \quad t \in L,$$

the coefficient $D(t) \equiv -1$, and $\gamma = \frac{1}{2}$, $\delta = 0$ in (76). In this situation,

$$\begin{aligned} X_0(z) &= \prod_{k=1}^m \frac{1}{\sqrt{(z - a_k)(z - b_k)}}, & X_0(z) &\in h_0; \\ X_m(z) &= \prod_{k=1}^m \sqrt{\frac{z - b_k}{z - a_k}}, & X_m(z) &\in h(b_1, b_2, \dots, b_m); \\ X_{m^*}(z) &= \prod_{k=1}^m \sqrt{\frac{z - a_k}{z - b_k}}, & X_{m^*}(z) &\in h(a_1, a_2, \dots, a_m); \\ X_{2m}(z) &= \prod_{k=1}^m \sqrt{(z - a_k)(z - b_k)}, & X_{2m}(z) &\in h_{2m}. \end{aligned}$$

14.3-12. Riemann Problem with a Discontinuous Coefficient.

Let L be a smooth closed curve and suppose that the coefficient $D(t)$ of the Riemann problem is continuous on L except at finitely many points t_1, t_2, \dots, t_m in which it has jumps of the first kind. On each arc $L_k = t_k t_{k+1}$, $k = 1, 2, \dots, m$ (it is assumed that $t_{m+1} = t_1$), the functions $D(t), H(t)$ satisfy the Hölder condition and $D(t) \neq 0$ for all t .

On an arc L_k , we fix a continuous branch of the logarithmic function $\ln D(t)$. This can also be done as in the case of an open curve L by fixing the values of $\ln D(t)$ at the initial points of the arcs:

$$\begin{aligned} \ln D(t_k + 0) &= |\ln D(t_k + 0)| + i \arg D(t_k + 0), \\ 0 \leq \arg D(t_k + 0) &< 2\pi, \quad k = 1, 2, \dots, m, \end{aligned} \tag{77}$$

where $D(t_k + 0) = \lim_{t \rightarrow t_k, t \in L_k} D(t)$ is the value of the function at the point t_k regarded as the initial point of the arc L_k . Then, at the finite point t_{k+1} of this arc, we have

$$\ln D(t_{k+1} - 0) = \ln D(t_k + 0) + i [\arg D(t)]_{L_k}, \quad k = 1, 2, \dots, m.$$

Let us calculate the numbers

$$\gamma_k + i\delta_k = \frac{1}{2\pi i} [\ln D(t_k - 0) - \ln D(t_k + 0)], \quad k = 1, 2, \dots, m, \tag{78}$$

and determine nonsingular discontinuity points t_k for which γ_k are noninteger, and singular points t_k for which γ_k are integer. Then we denote the points t_k by c_j , so that all the nonsingular points c_j occupy the first places and all singular points appear after these. Moreover, for $c_j = t_k$, the point c_j is associated with the number $\alpha_j + i\beta_j = \gamma_k + i\delta_k$.

Let c_1, c_2, \dots, c_n ($0 \leq n \leq m$) be all the nonsingular discontinuity points of the coefficient $D(t)$. The Riemann problem consists in finding two functions $\Phi^+(t)$ and $\Phi^-(t)$ that are analytic in the interior Ω^+ and the exterior Ω^- of the curve L , respectively, have boundary values $\Phi^+(t)$, $\Phi^-(t)$ continuous on L except, possibly, the points t_1, t_2, \dots, t_m , satisfying the boundary condition (8), bounded near the nonsingular points c_1, c_2, \dots, c_p ($0 \leq p \leq n$), having infinity of an order < 1 at the other nonsingular points c_{p+1}, \dots, c_n , and possibly, having logarithmic singularities at the points c_{n+1}, \dots, c_m .

Often, the condition of logarithmic singularity of the sought functions is replaced by the more general condition of almost boundedness:

$$\lim_{z \rightarrow c_j} |z - c_j|^\varepsilon \Phi^\pm(z) = 0 \quad \text{for any } \varepsilon > 0.$$

Similarly to the case of open curves, a canonical function $X(z)$ of class $h_p = h(c_1, c_2, \dots, c_p)$ for this problem can be constructed in the form

$$X(z) = e^{\Gamma(z)} \prod_{k=1}^m (z - c_j)^{-\nu_j}, \quad \Gamma(z) = \frac{1}{2\pi i} \int_L \frac{\ln D(\tau) d\tau}{\tau - z},$$

where $\nu_j = [\alpha_j]$ for the nonsingular points c_j , $j = 1, 2, \dots, p$, that determine the class h_p ; $\nu_j = 1 + [\alpha_j]$ for the other nonsingular points c_j , $j = p + 1, \dots, n$; $\nu_j = \alpha_j$ for singular points c_j , $j = n + 1, \dots, m$. In particular, in order to obtain a canonical function in the narrowest class $h_n = h(c_1, c_2, \dots, c_n)$ of functions bounded near all discontinuity points (both singular and nonsingular), one should take $\nu_j = [\alpha_j]$, $j = 1, \dots, m$.

With the help of the canonical function $X(z)$, the Riemann problem with a discontinuous coefficient is solved in the same way as in Subsections 12.3-4, 12.3-10, and 12.3-11. All the results of these subsections are valid for this problem, provided that one takes into account that the index of the problem is equal to $\nu = \nu_1 + \nu_2 + \dots + \nu_m$.

In the case of a piecewise-constant coefficient

$$D(t) = D_k = \text{const}, \quad t \in t_k t_{k+1}, \quad k = 1, 2, \dots, m,$$

the canonical function of the narrowest class h_n , which is bounded at the discontinuity points t_k , $k = 1, 2, \dots, m$, has the form

$$X_n(z) = \prod_{k=1}^m (z - t_k)^{\{\gamma_k\} + i\delta_k}, \tag{79}$$

where $\{\gamma_k\} = \gamma_k - [\gamma_k]$ is the fractional part of γ_k , which is the real part of the complex number

$$\begin{aligned} \gamma_k + i\delta_k &= \frac{1}{2\pi i} (\ln D_{k-1} - \ln D_k), \\ \ln D_k &= \ln |D_k| + i \arg D_k, \quad 0 \leq \arg D_k < 2\pi, \quad k = 1, 2, \dots, m \\ &\text{(it is assumed that } D_0 = D_m). \end{aligned} \tag{80}$$

Note that in general the difference of logarithms in (78), (80) cannot be replaced by the logarithm of fraction. For instance, if $D_{k-1} = 2$ and $D_k = 2i$, then for the logarithmic branch fixed by the condition $0 \leq \arg D_j < 2\pi$, we have

$$\ln D_{k-1} - \ln D_k = \ln 2 - \left(\ln 2 + \frac{\pi i}{2}\right) = -\frac{\pi i}{2} \quad \text{and} \quad \ln \frac{D_{k-1}}{D_k} = \ln(-i) = \frac{3\pi i}{2}.$$

Example 4. Let L be the unit circle $t = e^{i\varphi}$, $0 \leq \varphi \leq 2\pi$, and let the coefficient of the Riemann problem have the form

$$D(t) = D(e^{i\varphi}) = \begin{cases} -i, & 0 < \varphi < \frac{\pi}{2}, \\ 1 + i, & \frac{\pi}{2} < \varphi < \pi, \\ -1, & \pi < \varphi < 2\pi. \end{cases}$$

The function $D(t)$ is piecewise constant with discontinuities of the first kind at the points $t_1 = e^{i0} = 1$, $t_2 = e^{i\pi/2} = i$, $t_3 = e^{i\pi} = -1$. By (78), we find that

$$\begin{aligned} \ln D_1 &= \ln(-i) = \frac{3\pi i}{2}, & \ln D_2 &= \ln(1 + i) = \ln \sqrt{2} + \frac{\pi i}{4}, \\ \ln D_3 &= \ln(-1) = \pi i, & \ln D_0 &= \ln D_3 = \pi i, \\ \gamma_1 + i\delta_1 &= -\frac{1}{4}, & \gamma_2 + i\delta_2 &= \frac{5}{8} + i\frac{\ln 2}{4\pi}, & \gamma_3 + i\delta_3 &= -\frac{3}{8} - i\frac{\ln 2}{4\pi}. \end{aligned}$$

Since all γ_k are noninteger, all three discontinuity points are nonsingular. Then, according to (79), the canonical function $X_3(z)$ of class $h_3 = h(1, i, -1)$ (i.e., the function bounded near all discontinuity points) has the form

$$X_3(z) = (z - 1)^{3/4} (z - i)^{5/8 + i\delta} (z + 1)^{5/8 - i\delta}, \quad \delta = \frac{\ln 2}{4\pi}.$$

In order to obtain the canonical function $X_0(z)$ of the widest class h_0 , one should divide $X_3(z)$ by $(z - 1)(z - i)(z + 1)$; and to obtain the function $X(z) \in h(1)$, one should divide $X_3(z)$ by $(z - i)(z + 1)$, etc.

14.3-13. Riemann Problem in the General Case.

Let L be the union of finitely many smooth closed and open oriented curves with finitely many common points (L is a piecewise smooth line), and let $D(t)$, $H(t)$ be two functions on L that satisfy the Hölder condition everywhere except for finitely many first kind discontinuity points, $D(t) \neq 0$ everywhere on L . Denote by t_k the endpoints, the nodes, the angular points of the line L , and the discontinuity points of the function $D(t)$. On the closed curves belonging to L , we chose arbitrary points regarded as the initial and the ending points of these curves and include these points into the set of t_k .

Let t_1, t_2, \dots, t_m be all the above-specified points of the line L , which is split into finitely many oriented open arcs L_j by these points. On each arc, we fix a certain continuous branch of the logarithmic function $\ln D(t)$, so that if $t_k + 0$ is the initial point of some arc L_j , then the value $\ln D(t)$ at that point is found by the formula (77). Then, at the ending point $t_l - 0$ of that arc, we have

$$\ln D(t_l - 0) = \ln D(t_k + 0) + i[\arg D(t)]_{L_j}.$$

Note that one and the same t_k may happen to be the initial point of some arcs L_j (one or more) and the ending point of other arcs.

For each t_k , we calculate the number

$$\gamma_k + i\delta_k = \frac{1}{2\pi i} \left[\sum \ln D(t_k - 0) - \sum \ln D(t_k + 0) \right], \quad k = 1, 2, \dots, m,$$

where the first sum is over all $t_k - 0$ that are the endpoints of the arcs ending at the point t_k , and the second sum is over all $t_k + 0$ that are the initial points of the arcs issuing from t_k . For instance, if t_k is the initial point of the arc L_1, L_2, \dots, L_{m_1} , then the second sum has the form

$$\sum \ln D(t_k + 0) = \sum_{j=1}^{m_1} \ln D(t_k) \Big|_{t_k \in L_j} = \sum_{j=1}^{m_1} \lim_{t \rightarrow t_k, t \in L_k} D(t).$$

Further, as in the previous subsection, the condition that γ_k is integer or noninteger determines singular and nonsingular nodes c_j of the line L , after which the Riemann problem is formulated and solved as in Subsections 12.3-11 and 12.3-12.

Remark. One of the crucial steps when solving the Riemann problem is the construction of a canonical function of a given class. This function can be constructed in a simpler way.

Suppose that the line L is split by the points t_1, t_2, \dots, t_m into (closed and open) curves L_1, L_2, \dots, L_n . For each L_j , we can construct a canonical function $X_j(z)$ of the homogeneous Riemann problem $X_j^+(t) = D(t)X_j^-(t)$, $t \in L_j$, without taking care of its specific class. Then the function $X_0(z) = X_1(z)X_2(z) \dots X_n(z)$ satisfies the homogeneous boundary condition (7) and near the points t_k admits the representation

$$X_0(z) \sim A_k(z - t_k)^{\lambda_k + i\mu_k}, \quad A_k = \text{const} \neq 0, \quad k = 1, 2, \dots, m,$$

where $\lambda_k + i\mu_k$ are certain numbers found on the basis of the behavior of the functions $X_j(z)$ in a neighborhood of t_k . Knowing λ_k , it is easy to determine singular and nonsingular points t_k and find the canonical function $X(z)$ of a given class. This function has the form

$$X(z) = \prod_{j=1}^n X_j(z) \prod_{k=1}^m (z - t_k)^{-\omega_k},$$

where ω_k are integers to be chosen such that $X(z)$ should belong to the given class.

Example 5. Let L consist of the segment $L_1 = [-1; 1]$ on the real axis and the segment $L_2 = [0; i]$ on the imaginary axis, and

$$D(t) = \begin{cases} 2i, & t \in [-1; 0), \\ 2, & t \in (0; 1], \\ -1, & t \in (0; i]. \end{cases}$$

Let us construct the canonical function of the homogeneous Riemann problem with the coefficient $D(t)$, requiring that this function is bounded near all endpoints of the line L and near the node $t = 0$. For the points $t_1 = -1$ (the initial point of the segment $[-1; 0]$), $t_2 = 0$ (the ending point of the segment $[-1; 0]$ and the initial point of the segments $[0; 1]$, $[0; i]$), $t_3 = 1$ (the ending point of the segment $[0; 1]$), $t_4 = i$ (the ending point of the segment $[0; i]$), we find the numbers

$$\begin{aligned} \gamma_1 + i\delta_1 &= -\frac{1}{2\pi i} \ln(2i) = -\frac{1}{4} + i\frac{\ln 2}{2\pi}, & \gamma_2 + i\delta_2 &= \frac{1}{2\pi i} [\ln(2i) - \ln 2 - \ln(-1)] = -\frac{1}{4}, \\ \gamma_3 + i\delta_3 &= \frac{1}{2\pi i} \ln 2 = -i\frac{\ln 2}{2\pi}, & \gamma_4 + i\delta_4 &= \frac{1}{2\pi i} \ln(-1) = \frac{1}{2}. \end{aligned}$$

Since $\gamma_3 = 0$, the point $t_3 = 1$ is singular, while the rest of t_k are nonsingular. The canonical function that is bounded near all endpoints is found by (79) and has the form

$$X_3(z) = (z + 1)^{3/4 + i\delta} z^{3/4} (z - 1)^{-i\delta} (z - i)^{1/2}, \quad \delta = \frac{\ln 2}{2\pi}.$$

The canonical functions of the other classes are obtained from $X_3(z)$ by its division by $z + 1$, z , $z - i$, all or some of these, depending on the class.

14.3-14. Hilbert Boundary Value Problem.

Let a simple smooth closed contour L and real Hölder functions $a(s)$, $b(s)$, and $c(s)$ of the arc length s on the contour be given.

By the *Hilbert boundary value problem* we mean the following problem. Find a function

$$f(z) = u(x, y) + iv(x, y)$$

that is analytic on the domain Ω^+ and continuous on the contour for which the limit values of the real and the imaginary part on the contour satisfy the linear relation

$$a(s)u(s) + b(s)v(s) = c(s). \tag{81}$$

For $c(s) \equiv 0$ we obtain the *homogeneous* problem and, for nonzero $c(s)$, a *nonhomogeneous*. The Hilbert boundary value problem can be reduced to the Riemann boundary value problem. The methods of this reduction can be found in the references cited at the end of the section.

⊙ References for Section 14.3: F. D. Gakhov (1977), N. I. Muskhelishvili (1992).

14.4. Singular Integral Equations of the First Kind

14.4-1. Simplest Equation with Cauchy Kernel.

Consider the singular integral equation of the first kind

$$\frac{1}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau = f(t), \quad (1)$$

where L is a closed contour. Let us construct the solution. In this relation we replace the variable t by τ_1 , multiply by $\frac{1}{\pi i} \frac{d\tau_1}{\tau_1 - t}$, integrate along the contour L , and change the order of integration according to the Poincaré–Bertrand formula (see Subsection 14.2-6). Then we obtain

$$\frac{1}{\pi i} \int_L \frac{f(\tau_1)}{\tau_1 - t} d\tau_1 = \varphi(t) + \frac{1}{\pi i} \int_L \varphi(\tau) d\tau \frac{1}{\pi i} \int_L \frac{d\tau_1}{(\tau_1 - t)(\tau - \tau_1)}. \quad (2)$$

Let us calculate the second integral on the right-hand side of (2):

$$\int_L \frac{d\tau_1}{(\tau_1 - t)(\tau - \tau_1)} = \frac{1}{\tau - t} \left(\int_L \frac{d\tau_1}{\tau_1 - t} - \int_L \frac{d\tau_1}{\tau_1 - \tau} \right) = \frac{1}{\tau - t} (i\pi - i\pi) = 0.$$

Thus,

$$\varphi(t) = \frac{1}{\pi i} \int_L \frac{f(\tau)}{\tau - t} d\tau. \quad (3)$$

The last formula gives the solution of the singular integral equation of the first kind (1) for a closed contour L .

14.4-2. Equation with Cauchy Kernel on the Real Axis.

Consider the following singular integral equation of the first kind on the real axis:

$$\frac{1}{\pi i} \int_{-\infty}^{\infty} \frac{\varphi(t)}{t - x} dt = f(x), \quad -\infty < x < \infty. \quad (4)$$

Equation (4) is a special case of the characteristic integral equation on the real axis (see Subsection 15.2-3). In the class of functions vanishing at infinity, Eq. (4) has the solution

$$\varphi(x) = \frac{1}{\pi i} \int_{-\infty}^{\infty} \frac{f(t)}{t - x} dt, \quad -\infty < x < \infty. \quad (5)$$

Denoting $f(x) = F(x)i^{-1}$, we rewrite Eqs. (4) and (5) in the form

$$\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\varphi(t)}{t - x} dt = F(x), \quad \varphi(x) = -\frac{1}{\pi} \int_{-\infty}^{\infty} \frac{F(t)}{t - x} dt, \quad -\infty < x < \infty. \quad (6)$$

The two formulas (6) are called the Hilbert transform pair (see Subsection 9.6-5).

14.4-3. Equation of the First Kind on a Finite Interval.

Consider the singular integral equation of the first kind

$$\frac{1}{\pi} \int_a^b \frac{\varphi(t)}{t-x} dt = f(x), \quad a \leq x \leq b, \quad (7)$$

on a finite interval. Its solutions can be constructed by using the theory of the Riemann boundary value problem for a nonclosed contour (see Subsection 14.3-11). Let us present the final results.

1°. A solution that is unbounded at both endpoints:

$$\varphi(x) = -\frac{1}{\pi} \frac{1}{\sqrt{(x-a)(b-x)}} \left(\int_a^b \frac{\sqrt{(t-a)(b-t)}}{t-x} f(t) dt + C \right), \quad (8)$$

where C is an arbitrary constant and

$$\int_a^b \varphi(t) dt = C. \quad (9)$$

2°. A solution bounded at the endpoint a and unbounded at the endpoint b :

$$\varphi(x) = -\frac{1}{\pi} \sqrt{\frac{x-a}{b-x}} \int_a^b \sqrt{\frac{b-t}{t-a}} \frac{f(t)}{t-x} dt. \quad (10)$$

3°. A solution bounded at both endpoints:

$$\varphi(x) = -\frac{1}{\pi} \sqrt{(x-a)(b-x)} \int_a^b \frac{f(t)}{\sqrt{(t-a)(b-t)}} \frac{dt}{t-x}, \quad (11)$$

under the condition that

$$\int_a^b \frac{f(t) dt}{\sqrt{(t-a)(b-t)}} = 0. \quad (12)$$

Solutions that have a singularity point s inside the interval $[a, b]$ can also be constructed. These solutions have the following form:

4°. A singular solution that is unbounded at both endpoints:

$$\varphi(x) = -\frac{1}{\pi} \frac{1}{\sqrt{(x-a)(b-x)}} \left(\int_a^b \frac{\sqrt{(t-a)(b-t)}}{t-x} f(t) dt + C_1 + \frac{C_2}{x-s} \right), \quad (13)$$

where C_1 and C_2 are arbitrary constants.

5°. A singular solution bounded at one endpoint:

$$\varphi(x) = -\frac{1}{\pi} \sqrt{(x-a)(b-x)} \left(\int_a^b \sqrt{\frac{b-t}{t-a}} \frac{f(t)}{t-x} dt + \frac{C}{x-s} \right), \quad (14)$$

where C is an arbitrary constant.

6°. A singular solution bounded at both endpoints:

$$\varphi(x) = -\frac{1}{\pi} \sqrt{(x-a)(b-x)} \left(\int_a^b \frac{f(t)}{\sqrt{(t-a)(b-t)}} \frac{dt}{t-x} + \frac{A}{x-s} \right), \quad A = \int_{-1}^1 \frac{f(t) dt}{\sqrt{(t-a)(b-t)}}. \quad (15)$$

14.4-4. General Equation of the First Kind with Cauchy Kernel.

Consider the general equation of the first kind with Cauchy kernel

$$\frac{1}{\pi i} \int_L \frac{M(t, \tau)}{\tau - t} \varphi(\tau) d\tau = f(t), \quad (16)$$

where the integral is understood in the sense of the Cauchy principal value and is taken over a closed or nonclosed contour L . As usual, the functions $a(t)$, $f(t)$, and $M(t, \tau)$ on L are assumed to satisfy the Hölder condition, where the last function satisfies this condition with respect to both variables.

We perform the following manipulation with the kernel:

$$\frac{M(t, \tau)}{\tau - t} = \frac{M(t, \tau) - M(t, t)}{\tau - t} + \frac{M(t, t)}{\tau - t}$$

and write

$$M(t, t) = b(t), \quad \frac{1}{\pi i} \frac{M(t, \tau) - M(t, t)}{\tau - t} = K(t, \tau). \quad (17)$$

We can rewrite Eq. (16) in the form

$$\frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L K(t, \tau) \varphi(\tau) d\tau = f(t). \quad (18)$$

It follows from formulas (17) that the function $b(t)$ satisfies the Hölder condition on the entire contour L and $K(t, \tau)$ satisfies this condition everywhere except for the points with $\tau = t$ at which this function satisfies the estimate

$$|K(t, \tau)| < \frac{A}{|\tau - t|^\lambda}, \quad 0 \leq \lambda < 1.$$

The general singular integral equation of the first kind with Cauchy kernel is frequently written in the form (18).

The general singular integral equation of the first kind is a special case of the complete singular integral equation whose theory is treated in Chapter 15. In general, it cannot be solved in a closed form. However, there are some cases in which such a solution is possible.

Let the function $M(t, \tau)$ in Eq. (16), which satisfies the Hölder condition with respect to both variables on the smooth closed contour L by assumption, have an analytic continuation to the domain Ω^+ with respect to each of the variables. If $M(t, t) \equiv 1$, then the solution of Eq. (16) can be obtained by means of the Poincaré–Bertrand formula (see Subsection 14.2-6). This solution is given by the relation

$$\varphi(t) = \frac{1}{\pi i} \int_L \frac{M(t, \tau)}{\tau - t} f(\tau) d\tau. \quad (19)$$

Equation (16) can be solved without the assumption that the function $M(t, \tau)$ satisfies the condition $M(t, t) \equiv 1$. Namely, assume that the function $M(t, \tau)$ has the analytic continuation to Ω^+ with respect to each of the variables and that $M(z, z) \neq 0$ for $z \in \overline{\Omega^+}$. In this case, the solution of Eq. (16) has the form

$$\varphi(t) = \frac{1}{\pi i} \frac{1}{M(t, t)} \int_L \frac{M(t, \tau)}{M(\tau, \tau)} \frac{f(\tau)}{\tau - t} d\tau. \quad (20)$$

In Section 14.5, a numerical method for solving a special case of the general equation of the first kind is given, which is of independent interest from the viewpoint of applications.

Remark 1. The solutions of complete singular integral equations that are constructed in Subsection 14.4-4 can also be applied for the case in which the contour L is a collection of finitely many disjoint smooth closed contours.

14.4-5. Equations of the First Kind with Hilbert Kernel.

1°. Consider the simplest singular integral equation of the first kind with Hilbert kernel

$$\frac{1}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) \varphi(\xi) d\xi = f(x), \quad 0 \leq x \leq 2\pi, \quad (21)$$

under the additional assumption

$$\int_0^{2\pi} \varphi(x) dx = 0. \quad (22)$$

Equation (21) can have a solution only if a solvability condition is satisfied. This condition is obtained by integrating Eq. (21) with respect to x from zero to 2π and, with regard for the relation

$$\int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) dx = 0,$$

becomes

$$\int_0^{2\pi} f(x) dx = 0. \quad (23)$$

To construct a solution of Eq. (21), we apply the solution of the simplest singular integral equation of the first kind with Cauchy kernel by assuming that the contour L is the circle of unit radius centered at the origin (see Subsection 14.4-1). We rewrite the equation with Cauchy kernel and its solution in the form

$$\frac{1}{\pi} \int_L \frac{\varphi_1(\tau)}{\tau-t} d\tau = f_1(t), \quad (24)$$

$$\varphi_1(t) = -\frac{1}{\pi} \int_L \frac{f_1(\tau)}{\tau-t} d\tau, \quad (25)$$

which is obtained by substituting the function $\varphi_1(t)$ instead of $\varphi(t)$ and the function $f_1(t)i^{-1}$ instead of $f(t)$ into the relations of 14.4-1.

We set $t = e^{ix}$ and $\tau = e^{i\xi}$ and find the relationship between the Cauchy kernel and the Hilbert kernel:

$$\frac{d\tau}{\tau-t} = \frac{1}{2} \cot\left(\frac{\xi-x}{2}\right) d\xi + \frac{i}{2} d\xi. \quad (26)$$

On substituting relation (26) into Eq. (24) and into solution (25), with regard to the change of variables $\varphi(x) = \varphi_1(t)$ and $f(x) = f_1(t)$ we obtain

$$\frac{1}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) \varphi(\xi) d\xi + \frac{i}{2\pi} \int_0^{2\pi} \varphi(\xi) d\xi = f(x), \quad (27)$$

$$\varphi(x) = -\frac{1}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) f(\xi) d\xi - \frac{i}{2\pi} \int_0^{2\pi} f(\xi) d\xi. \quad (28)$$

Equation (21), under the additional assumption (22), coincides with Eq. (27), and hence its solution is given by the expression (28). Taking into account the solvability conditions (23), on the basis of (28) we rewrite a solution of Eq. (21) in the form

$$\varphi(x) = -\frac{1}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) f(\xi) d\xi. \quad (29)$$

Formulas (21) and (29), together with conditions (22) and (23), are called the *Hilbert inversion formula*.

Remark 2. Equation (21) is a special case of the characteristic singular integral equation with Hilbert kernel (see Subsections 15.1-2 and 15.2-5).

2°. Consider the general singular integral equation of the first kind with Hilbert kernel

$$\frac{1}{2\pi} \int_0^{2\pi} N(x, \xi) \cot\left(\frac{\xi-x}{2}\right) \varphi(\xi) d\xi = f(x). \quad (30)$$

Let us represent its kernel in the form

$$N(x, \xi) \cot\frac{\xi-x}{2} = [N(x, \xi) - N(x, x)] \cot\frac{\xi-x}{2} + N(x, x) \cot\frac{\xi-x}{2}.$$

We introduce the notation

$$N(x, x) = -b(x), \quad \frac{1}{2\pi} [N(x, \xi) - N(x, x)] \cot\frac{\xi-x}{2} = K(x, \xi), \quad (31)$$

and rewrite Eq. (30) as follows:

$$-\frac{b(x)}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) \varphi(\xi) d\xi + \int_0^{2\pi} K(x, \xi) \varphi(\xi) d\xi = f(x). \quad (32)$$

It follows from formulas (31) that the function $b(x)$ satisfies the Hölder condition, whereas the kernel $K(x, \xi)$ satisfies the Hölder condition everywhere except possibly for the points $x = \xi$, at which the following estimate holds:

$$|K(x, \xi)| < \frac{A}{|\xi-x|^\lambda}, \quad A = \text{const} < \infty, \quad 0 \leq \lambda < 1.$$

The general singular integral equation of the first kind with Hilbert kernel is frequently written in the form (32). It is a special case of the complete singular integral equation with Hilbert kernel, which is treated in Subsections 15.1-2 and 15.4-8.

⊙ References for Section 14.4: F. D. Gakhov (1977), F. D. Gakhov and Yu. I. Cherskii (1978), S. G. Mikhlin and S. Prössdorf (1986), N. I. Muskhelishvili (1992), I. K. Lifanov (1996).

14.5. Multhopp–Kalandiya Method

Consider a general singular integral equation of the first kind with Cauchy kernel on the finite interval $[-1, 1]$ of the form

$$\frac{1}{\pi} \int_{-1}^1 \frac{\varphi(t) dt}{t-x} + \frac{1}{\pi} \int_{-1}^1 K(x, t) \varphi(t) dt = f(x). \quad (1)$$

This equation frequently occurs in applications, especially in aerodynamics and 2D elasticity.

We present here a method of approximate solution of Eq. (1) under the assumption that this equation has a solution in the classes indicated below.

14.5-1. Solution That is Unbounded at the Endpoints of the Interval.

According to the general theory of singular integral equations (e.g., see N. I. Muskhelishvili (1992)), such a solution can be represented in the form

$$\varphi(x) = \frac{\psi(x)}{\sqrt{1-x^2}}, \quad (2)$$

where $\psi(x)$ is a bounded function on $[-1, 1]$. Let us substitute the expression (2) into Eq. (1) and introduce new variables θ and τ by the relations $x = \cos \theta$ and $t = \cos \tau$, $0 \leq \theta \leq \pi$, $0 \leq \tau \leq \pi$. In this case, Eq. (1) becomes

$$\frac{1}{\pi} \int_0^\pi \frac{\psi(\cos \tau) d\tau}{\cos \tau - \cos \theta} + \frac{1}{\pi} \int_0^\pi K(\cos \theta, \cos \tau) \psi(\cos \tau) d\tau = f(\cos x). \quad (3)$$

Let us construct the *Lagrange interpolation polynomial* for the desired function $\psi(x)$ with the *Chebyshev nodes*

$$x_m = \cos \theta_m, \quad \theta_m = \frac{2m-1}{2n} \pi, \quad m = 1, \dots, n.$$

This polynomial is known to have the form

$$L_n(\psi; \cos \theta) = \frac{1}{n} \sum_{l=1}^n (-1)^{l+1} \psi(\cos \theta_l) \frac{\cos n\theta \sin \theta_l}{\cos \theta - \cos \theta_l}. \quad (4)$$

Note that for each l the fraction on the right-hand side in (4) is an even trigonometric polynomial of degree $\leq n-1$. We define the coefficients of this polynomial by means of the known relations

$$\frac{1}{\pi} \int_0^\pi \frac{\cos n\tau d\tau}{\cos \tau - \cos \theta} = \frac{\sin n\theta}{\sin \theta}, \quad 0 \leq \theta \leq \pi, \quad n = 0, 1, 2, \dots \quad (5)$$

and rewrite (4) in the form

$$L_n(\psi; \cos \theta) = \frac{2}{n} \sum_{l=1}^n \psi(\cos \theta_l) \sum_{m=0}^{n-1} \cos m\theta_l \cos m\theta - \frac{1}{n} \sum_{l=1}^n \psi(\cos \theta_l). \quad (6)$$

On the basis of the above two relations we write out the following quadrature formula for the singular integral:

$$\frac{1}{\pi} \int_{-1}^1 \frac{\varphi(t) dt}{t-x} = \frac{2}{n \sin \theta} \sum_{l=1}^n \psi(\cos \theta_l) \sum_{m=1}^{n-1} \cos m\theta_l \sin m\theta. \quad (7)$$

This formula is exact for the case in which $\psi(t)$ is a polynomial of order $\leq n-1$ in t .

To the second integral on the left-hand side of Eq. (1), we apply the formula

$$\frac{1}{\pi} \int_{-1}^1 \frac{P(x) dx}{\sqrt{1-x^2}} = \frac{1}{n} \sum_{l=1}^n P(\cos \theta_l), \quad (8)$$

which holds for any polynomial $P(x)$ of degree $\leq 2n-1$. In this case, by (8) we have

$$\frac{1}{\pi} \int_{-1}^1 K(x, t) \varphi(t) dt = \frac{1}{n} \sum_{l=1}^n K(\cos \theta, \cos \theta_l) \psi(\cos \theta_l). \quad (9)$$

On substituting relations (7) and (9) into Eq. (1), we obtain

$$\frac{2}{n \sin \theta} \sum_{l=1}^n \psi(\cos \theta_l) \sum_{m=1}^{n-1} \cos m\theta_l \sin m\theta + \frac{1}{n} \sum_{l=1}^n K(\cos \theta, \cos \theta_l) \psi(\cos \theta_l) = f(\cos \theta). \quad (10)$$

By setting $\theta = \theta_k$ ($k = 1, \dots, n$) and with regard to the formula

$$\sum_{m=1}^{n-1} \cos m\theta_l \sin m\theta_k = \frac{1}{2} \cot \frac{\theta_k \pm \theta_l}{2}, \quad (11)$$

where the sign “plus” is taken for the case in which $|k - l|$ is even and “minus” if $|k - l|$ odd, we obtain the following system of linear algebraic equations for the approximate values ψ_l of the desired function $\psi(x)$ at the nodes:

$$\begin{aligned} \sum_{l=1}^n a_{kl} \psi_l &= f_k, \quad f_k = f(\cos \theta_k), \quad k = 1, \dots, n, \\ a_{kl} &= \frac{1}{n} \left[\frac{1}{\sin \theta_k} \cot \frac{\theta_k \pm \theta_l}{2} + K(\cos \theta_k, \cos \theta_l) \right]. \end{aligned} \quad (12)$$

After solving the system (12), the corresponding approximate solution to Eq. (1) can be found by formulas (2) and (4).

14.5-2. Solution Bounded at One Endpoint of the Interval.

In this case we set

$$\varphi(x) = \sqrt{\frac{1-x}{1+x}} \zeta(x), \quad (13)$$

where $\zeta(x)$ is a bounded function on $[-1, 1]$.

We take the same interpolation nodes as in Subsection 14.5-1, replace $\zeta(x)$ by the polynomial

$$L_n(\zeta; \cos \theta) = \frac{1}{n} \sum_{l=1}^n (-1)^{l+1} \zeta(\cos \theta_l) \frac{\cos n\theta \sin \theta_l}{\cos \theta - \cos \theta_l}, \quad (14)$$

and substitute the result into the singular integral that enters the expression (1). Just as above, we obtain the following quadrature formula:

$$\frac{1}{\pi} \int_{-1}^1 \frac{\varphi(t) dt}{t-x} = 2 \frac{1-\cos \theta}{n \sin \theta} \sum_{l=1}^n \zeta(\cos \theta_l) \sum_{m=1}^{n-1} \cos m\theta_l \sin m\theta - \frac{1}{n} \sum_{l=1}^n \zeta(\cos \theta_l). \quad (15)$$

This formula is exact for the case in which $\zeta(t)$ is a polynomial of order $\leq n-1$ in t .

The formula for the second summand on the left-hand side of the equation becomes

$$\frac{1}{\pi} \int_{-1}^1 K(x, t) \varphi(t) dt = \frac{1}{n} \sum_{l=1}^n (1 - \cos \theta_l) K(\cos \theta, \cos \theta_l) \zeta(\cos \theta_l). \quad (16)$$

This formula is exact if the integrand is a polynomial in t of degree $\leq 2n-2$.

On substituting relations (15) and (16) into Eq. (1) and on setting $\theta = \theta_k$ ($k = 1, \dots, n$), with regard to formula (11), we obtain a system of linear algebraic equations for the approximate values ζ_l of the desired function $\zeta(x)$ at the nodes:

$$\begin{aligned} \sum_{l=1}^n b_{kl} \zeta_l &= f_k, \quad f_k = f(\cos \theta_k), \quad k = 1, \dots, n, \\ b_{kl} &= \frac{1}{n} \left[\tan \frac{\theta_k}{2} \cot \frac{\theta_k \pm \theta_l}{2} - 1 + 2 \sin^2 \frac{\theta_l}{2} K(\cos \theta_k, \cos \theta_l) \right]. \end{aligned} \quad (17)$$

After solving system (17), the corresponding approximate solution to Eq. (1) can be found by formulas (13) and (14).

14.5-3. Solution Bounded at Both Endpoints of the Interval.

A solution of Eq. (1) that is bounded at the endpoints of the interval vanishes at the endpoints,

$$\varphi(1) = \varphi(-1) = 0. \quad (18)$$

Let us approximate the function $\varphi(x)$ by an even trigonometric polynomial of θ constructed for the interpolation nodes that are the roots of the corresponding *Chebyshev polynomial of the second kind*:

$$x_k = \cos \theta_k, \quad \theta_k = \frac{k\pi}{n+1}, \quad k = 1, \dots, n. \quad (19)$$

This polynomial has the form

$$M_n(\varphi; \cos \theta) = \frac{2}{n+1} \sum_{l=1}^n \varphi(\cos \theta_l) \sum_{m=1}^n \sin m\theta_l \sin m\theta. \quad (20)$$

We thus obtain the following quadrature formula:

$$\frac{1}{\pi} \int_{-1}^1 \frac{\varphi(t) dt}{t-x} = -\frac{2}{n+1} \sum_{l=1}^n \varphi(\cos \theta_l) \sum_{m=1}^n \sin m\theta_l \cos m\theta. \quad (21)$$

This formula holds for any odd trigonometric polynomial $\varphi(x)$ of degree $\leq n$.

To the regular integral in Eq. (1) we apply the formula

$$\int_{-1}^1 \sqrt{1-x^2} P(x) dx = \frac{\pi}{n+1} \sum_{l=1}^n \sin^2 \theta_l P(\cos \theta_l), \quad (22)$$

whose accuracy coincides with that of formula (8). On the basis of (22), we have

$$\frac{1}{\pi} \int_{-1}^1 K(x, t) \varphi(t) dt = \frac{1}{n+1} \sum_{l=1}^n \sin \theta_l K(\cos \theta, \cos \theta_l) \varphi(\cos \theta_l). \quad (23)$$

On substituting relations (21) and (23) into Eq. (1) and on setting $\theta = \theta_k$ ($k = 1, \dots, n$), we obtain a system of linear algebraic equations in the form

$$\sum_{l=1}^n c_{kl} \varphi_l = f_k, \quad k = 1, \dots, n, \quad (24)$$

$$c_{kl} = \frac{\sin \theta_l}{n+1} \left[\frac{2\varepsilon_{kl}}{\cos \theta_l - \cos \theta_k} + K(\cos \theta_k, \cos \theta_l) \right], \quad \varepsilon_{kl} = \begin{cases} 0 & \text{for even } |k-l|, \\ 1 & \text{for odd } |k-l|, \end{cases}$$

where $f_k = f(\cos \theta_k)$ and φ_l are approximate values of the unknown function $\varphi(x)$ at the nodes.

After solving system (24), the corresponding approximate solution is defined by formula (20).

When solving a singular integral equation by the Multhopp–Kalandiya method, it is important that the desired solutions have a representation

$$\varphi(x) = (1-x)^\alpha (1+x)^\beta \chi(x), \quad (25)$$

where $\alpha = \pm \frac{1}{2}$, $\beta = \pm \frac{1}{2}$, and $\chi(x)$ is a bounded function on the interval with well-defined values at the endpoints. If the representation (25) holds, then the method can be applied to the complete singular integral equation, which is treated in Chapter 15.

In the literature cited below, some other methods of numerical solution of singular integral equations are discussed as well.

⊙ References for Section 14.5: A. I. Kalandiya (1973), N. I. Muskhelishvili (1992), S. M. Belotserkovskii and I. K. Lifanov (1993), and I. K. Lifanov (1996).

14.6. Hypersingular Integral Equations

14.6-1. Hypersingular Integral Equations with Cauchy- and Hilbert-Type Kernels.

The simplest *hypersingular integral equation of the first kind with Cauchy-type kernel* on a finite interval has the form

$$\frac{1}{\pi} \int_a^b \frac{\varphi(t)}{(x-t)^2} dt = f'_x(x), \quad a \leq x \leq b, \quad (1)$$

where $\varphi(t)$ is the unknown function, $\frac{1}{(x-t)^2}$ is *Cauchy-type kernel*, $f'_x(x)$ is a function called the *free term* or the *right-hand side of equation* (1). The integral on the left-hand side exists only in the sense of Hadamard principal value (see Subsection 14.6-2).

The *general hypersingular equation of the first kind with Cauchy-type kernel* on a finite interval has the form

$$\frac{1}{\pi} \int_a^b \frac{\varphi(t)}{(x-t)^2} dt + \frac{1}{\pi} \int_a^b K'_x(x, t)\varphi(t) dt = f'_x(x), \quad a \leq x \leq b. \quad (2)$$

Assume that the functions $\varphi(x)$, $f(x)$ in equations (1), (2) are differentiable and $K(x, t)$ is differentiable in both variables everywhere except at the points $x = t$, near which it satisfies the estimate

$$|K(x, t)| \leq \frac{A}{|x-t|^\lambda}, \quad A = \text{const} < \infty, \quad 0 \leq \lambda < 1.$$

Remark 1. The notation in (1) and (2) is meant to emphasize the fact that these equations are obtained from equation (3) of Subsection 14.1-1 and equation (1) of Section 14.5 by their differentiation in x .

The simplest *hypersingular equation of the first kind with Hilbert-type kernel* has the form

$$\frac{1}{4\pi} \int_0^{2\pi} \left[\sin\left(\frac{\xi-x}{2}\right) \right]^{-2} \varphi(\xi) d\xi = f'_x(x), \quad 0 \leq x \leq 2\pi, \quad (3)$$

where $\varphi(x)$ is the unknown function, $1/\sin^2[\frac{1}{2}(\xi-x)]$ is *Hilbert-type kernel*, $f(x)$ is a given right-hand side of the equation.

The *general hypersingular equation of the first kind with Hilbert-type kernel* has the form

$$\frac{1}{4\pi} \int_0^{2\pi} \left[\sin\left(\frac{\xi-x}{2}\right) \right]^{-2} \varphi(\xi) d\xi + \frac{1}{4\pi} \int_a^b K'_x(x, \xi)\varphi(\xi) d\xi = f'_x(x), \quad 0 \leq x \leq 2\pi, \quad (4)$$

where $\varphi(x)$, $f(x)$, and $K(x, t)$ are functions with the properties specified above.

If the right-hand sides of equations (1)–(4) are identically equal to zero, the equations are called *homogeneous*; otherwise, they are called *nonhomogeneous*.

Remark 2. Note that there is a relation between hypersingular integral equations (3), (4) and singular integral equations (5) from Subsection 14.1-2 and (32) from Subsection 14.4-5: the latter are obtained from the former by the integration in ξ .

14.6-2. Definition of Hypersingular Integrals.

Hypersingular integrals in equations (1)–(4) exist neither in the sense of improper integrals nor in the sense of the Cauchy principal value. Taking as an example hypersingular integrals with the Cauchy-type kernel, consider some definitions of such integrals.

1°. Hypersingular integral as the derivative of an integral in the sense of the Cauchy principal value:

$$\int_a^b \frac{\varphi(t)}{(x-t)^2} dt = \frac{d}{dx} \int_a^b \frac{\varphi(t)}{t-x} dt. \quad (5)$$

2°. Hypersingular integral in the sense of *Hadamard principal value*:

$$\int_a^b \frac{\varphi(t)}{(x-t)^2} dt = \lim_{\varepsilon \rightarrow +0} \left[\left(\int_{\alpha}^{x-\varepsilon} + \int_{x+\varepsilon}^b \right) \frac{\varphi(t) dt}{(x-t)^2} - \frac{2\varphi(x)}{\varepsilon} \right]. \quad (6)$$

3°. Hypersingular integral as an analytic continuation of the integral

$$\int_a^b |x-t|^\alpha \varphi(t) dt \quad (7)$$

understood in the sense of distributions, where $\alpha = -2$.

Example. Let us calculate the values of hypersingular integrals using formulas (5)–(7) for $\varphi(x) \equiv 1$.

1°. Using the first definition, we have

$$\int_a^b \frac{dt}{(x-t)^2} = \frac{d}{dx} \int_a^b \frac{dt}{t-x} = \frac{d}{dx} \ln \left(\frac{b-x}{x-a} \right) = \frac{a-b}{(x-a)(b-x)}.$$

2°. For the Hadamard principal value, we have

$$\int_a^b \frac{dt}{(x-t)^2} = \lim_{\varepsilon \rightarrow +0} \left[\left(-\frac{1}{x-a} + \frac{1}{\varepsilon} + \frac{1}{\varepsilon} + \frac{1}{x-b} \right) - \frac{2}{\varepsilon} \right] = \frac{a-b}{(x-a)(b-x)}.$$

3°. Using formula (3), for $\operatorname{Re}(\alpha) > -1$ we get

$$\int_a^b |x-t|^\alpha dt = \int_a^x (x-t)^\alpha dt + \int_x^b (t-x)^\alpha dt = \frac{(x-a)^{\alpha+1}}{\alpha+1} + \frac{(b-x)^{\alpha+1}}{\alpha+1}.$$

Analytic continuation of this function from the half-plane $\operatorname{Re}(\alpha) > -1$ for $\alpha = -2$ yields

$$\int_a^b \frac{dt}{(x-t)^2} = \lim_{\alpha \rightarrow -2} \left[\frac{(x-a)^{\alpha+1}}{\alpha+1} + \frac{(b-x)^{\alpha+1}}{\alpha+1} \right] = \frac{a-b}{(x-a)(b-x)}.$$

For a differentiable function $\varphi(x)$ on the segment (a, b) the above three definitions of hypersingular integrals are equivalent.

The expression

$$\left(\int_{\alpha}^{x-\varepsilon} + \int_{x+\varepsilon}^b \right) \frac{\varphi(t) - \varphi(x) - \varphi'_x(x)(t-x)}{(x-t)^2} dt + \varphi(x) \frac{a-b}{(x-a)(b-x)} + \varphi'_x(x) \ln \frac{b-x}{x-a},$$

which is equivalent to the right-hand side of (2), shows that for a differentiable $\varphi(x)$, $x \in (a, b)$, a finite value of the hypersingular integral $\int_a^b \varphi(t)(x-t)^{-2} dt$ exists always, since this expression has a finite limit as $\varepsilon \rightarrow +0$.

Remark 3. Hypersingular integrals with the Hilbert-type kernel can be defined by analogy with the above definitions in the case of integrals with the Cauchy-type kernel. Note also that equation (26) of Section 14.4 establishes a relation between the Cauchy and the Hilbert kernels.

Remark 4. From definition of hypersingular integral in the sense of Hadamard principal value (6) we can see that

$$\int_a^b \frac{\varphi(t)}{(x-t)^2} dt = \frac{\varphi(a)}{a-x} - \frac{\varphi(b)}{b-x} + \int_a^b \frac{\varphi'_t(t) dt}{t-x},$$

which means that the right-hand side of this equation can be understood as a result of formal integration by parts.

14.6-3. Exact Solution of the Simplest Hypersingular Equation with Cauchy-Type Kernel.
--

Consider the simplest hypersingular equation of the first kind with Cauchy-type kernel on a finite interval

$$\frac{1}{\pi} \int_a^b \frac{\varphi(t)}{(x-t)^2} dt = f'_x(x), \quad a \leq x \leq b, \quad (8)$$

where $\varphi(a) = \varphi(b) = 0$. Let us construct its solution by two methods.

1°. According to definition (5) from Subsection 14.6-2, this simplest equation can be written in the form

$$\frac{1}{\pi} \frac{d}{dx} \int_a^b \frac{\varphi(t) dt}{t-x} = f'_x(x), \quad a \leq x \leq b.$$

Integrating the last equation with respect to x , we obtain

$$\frac{1}{\pi} \int_a^b \frac{\varphi(t) dt}{t-x} = f(x) + C, \quad a \leq x \leq b, \quad (9)$$

where C is an arbitrary constant. A bounded solution of equation (9) has been obtained in Subsection 14.4-3. This solution has the form

$$\varphi(x) = -\frac{1}{\pi} \sqrt{(x-a)(b-x)} \int_a^b \frac{f(t)}{\sqrt{(t-a)(b-t)} t-x} dt, \quad C = \frac{1}{\pi} \int_a^b \frac{f(t)}{\sqrt{(t-a)(b-t)}} dt. \quad (10)$$

2°. Integrating by parts equation (8) (see Remark 4), we get

$$\frac{1}{\pi} \left[\frac{\varphi(a)}{a-x} - \frac{\varphi(b)}{b-x} + \int_a^b \frac{\varphi'_t(t) dt}{t-x} \right] = f'_x(x).$$

Using this relation and the conditions $\varphi(a) = \varphi(b) = 0$, we finally come to the equation

$$\frac{1}{\pi} \int_a^b \frac{\varphi'_t(t) dt}{t-x} = f'_x(x). \quad (11)$$

Consider the solution of equation (11) given in Subsection 14.4-3:

$$\varphi'_x(x) = -\frac{1}{\pi} \frac{1}{\sqrt{(x-a)(b-x)}} \int_a^b \frac{\sqrt{(t-a)(b-t)}}{t-x} f'_t(t) dt, \quad \int_a^b \varphi'_t(t) dt = 0. \quad (12)$$

Integrating (12) from a to x , we get

$$\varphi(x) = -\frac{1}{\pi} \int_a^x \frac{1}{\sqrt{(\tau-a)(b-\tau)}} \int_a^b \frac{\sqrt{(t-a)(b-t)}}{t-\tau} f'_t(t) dt d\tau.$$

Hence, changing the order of integration, we obtain

$$\varphi(x) = \frac{1}{\pi} \int_a^b \int_a^x \frac{1}{\sqrt{(\tau-a)(b-\tau)}} \frac{d\tau}{\tau-t} f'_t(t) \sqrt{(t-a)(b-t)} dt. \quad (13)$$

The internal integral in (6) can be calculated by the formula

$$\begin{aligned} & \int \frac{1}{\sqrt{(\tau-a)(b-\tau)}} \frac{d\tau}{\tau-t} \\ &= -\frac{1}{2\sqrt{(t-a)(b-t)}} \ln \frac{\frac{1}{2}(a+b)(t+\tau) - ab - t\tau + \sqrt{(\tau-a)(b-\tau)(t-a)(b-t)}}{\frac{1}{2}(a+b)(t+\tau) - ab - t\tau - \sqrt{(\tau-a)(b-\tau)(t-a)(b-t)}} \\ &= \frac{1}{\sqrt{(t-a)(b-t)}} \ln \left| \frac{\sqrt{(b-t)(\tau-a)} - \sqrt{(b-\tau)(t-a)}}{\sqrt{(b-t)(\tau-a)} + \sqrt{(b-\tau)(t-a)}} \right|. \end{aligned}$$

Thus, a solution of the simplest hypersingular equation with Cauchy-type kernel (8) can be obtained in the form

$$\varphi(x) = \frac{1}{\pi} \int_a^b \ln \left| \frac{\sqrt{(b-t)(x-a)} - \sqrt{(b-x)(t-a)}}{\sqrt{(b-t)(x-a)} + \sqrt{(b-x)(t-a)}} \right| f'_t(t) dt, \quad (14)$$

which, in contrast to (10), contains no singular integrals.

14.6-4. Exact Solution of the Simplest Hypersingular Equation with Hilbert-Type Kernel.

Consider the simplest hypersingular integral equation of the first kind with Hilbert-type kernel on the finite interval

$$\frac{1}{4\pi} \int_0^{2\pi} \left[\sin\left(\frac{\xi-x}{2}\right) \right]^{-2} \varphi(\xi) d\xi = f'_x(x), \quad 0 \leq x \leq 2\pi \quad (15)$$

with the periodic conditions $\varphi(0) = \varphi(2\pi)$. Let us construct its solution by two methods.

1°. According to definition (1) from Subsection 14.2-1, this equation can be written in the form

$$\frac{1}{2\pi} \frac{d}{dx} \int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) \varphi(\xi) d\xi = f'_x(x), \quad 0 \leq x \leq 2\pi. \quad (16)$$

Integrating (16), we reduce the problem of finding a solution of the hypersingular equation under consideration to that of finding a solution of the following singular integral equation with the Hilbert kernel:

$$\frac{1}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) \varphi(\xi) d\xi = f(x) + C, \quad 0 \leq x \leq 2\pi, \quad (17)$$

where C is an arbitrary constant. This equation is considered in Subsection 14.4-5.

2°. Integrating (15) by parts, we obtain

$$\frac{1}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi-x}{2}\right) \varphi'_\xi(\xi) d\xi = f'_x(x), \quad 0 \leq x \leq 2\pi. \quad (18)$$

To find a solution of equation (18), let us use the result obtained in Subsection 14.4-5 for a singular integral equation of the first kind with the Hilbert kernel. We finally get

$$\varphi(x) = -\frac{1}{\pi} \int_0^{2\pi} f'_\xi(\xi) \ln \left| \sin\left(\frac{\xi-x}{2}\right) \right| d\xi + C,$$

where C is an arbitrary constant.

14.6-5. Numerical Methods for Hypersingular Equations.

1°. Consider collocation method for the simplest equation (1). Let us partition the interval $[a, b]$ into n equal segments of length $h = (b-a)/n$ with endpoints at the nodes $a = t_0, t_1, t_2, \dots, t_{n-1}, t_n = b$, $t = a + jh$, $j = 0, 1, \dots, n$. Denote the midpoints of the segments $[t_{i-1}, t_i]$ by x_i . It is easy to see that $x_i = a + (i-1/2)h$ for $i = 1, \dots, n$.

Let us represent an approximate value of the integral from (1) as a finite sum. Then, for $x = x_i$, we have

$$\begin{aligned} \frac{1}{\pi} \int_a^b \frac{\varphi(t) dt}{(x_i-t)^2} &\approx \frac{1}{\pi} \sum_{j=0}^n \varphi(t_j) \int_{t_{j-1}}^{t_j} \frac{dt}{(x_i-t)^2} = \frac{1}{\pi} \varphi(t_i) \int_{-h/2}^{h/2} \frac{dt}{t^2} \\ &+ \frac{1}{\pi} \sum_{j \neq i} \varphi(t_j) \left(\frac{1}{x_i-t_j} - \frac{1}{x_i-t_{j-1}} \right) = \frac{1}{\pi} \sum_{j=1}^n \varphi(t_j) \left(\frac{1}{x_i-t_j} - \frac{1}{x_i-t_{j-1}} \right). \end{aligned} \quad (19)$$

Now, let us replace the hypersingular integral equation under consideration by an approximate expression in the form of a system of linear algebraic equations:

$$\frac{1}{\pi} \sum_{j=1}^n \varphi(t_j) \left(\frac{1}{x_i-t_j} - \frac{1}{x_i-t_{j-1}} \right) = f'(x_i), \quad i = 1, \dots, n. \quad (20)$$

It can be shown that for a fixed $x = x_l \in (a, b)$, the difference of the solutions $\varphi(x_l)$ of system (20) and equation (19) tends to zero as $n \rightarrow \infty$, i.e.,

$$\begin{aligned}\varphi(x_l) &\sim -\frac{h}{\pi} \sqrt{(x_l - a)(b - x_l)} \sum_{m=1}^n \frac{f(t_m)}{\sqrt{(t_m - a)(b - t_m)} (x_l - t_m)} \\ &\sim -\frac{\sqrt{(x_l - a)(b - x_l)}}{\pi} \int_a^b \frac{f(t)}{\sqrt{(t - a)(b - t)} t - x_l} dt.\end{aligned}$$

2°. By analogy with the above considerations, one can obtain an approximate solution of the general hypersingular integral equation (2) by a collocation method solving the following system of algebraic equations:

$$\sum_{j=1}^n \left[\frac{1}{x_i - t_j} - \frac{1}{x_i - t_{j-1}} + hK'_x(x_i, t_j) \right] \varphi(t_j) = f'_x(x_i), \quad i = 1, \dots, n.$$

3°. Consider the general hypersingular integral equation (2) of the first kind with the Cauchy-type kernel on a finite interval and write this equation in the form

$$\frac{1}{\pi} \int_a^b \frac{\varphi(t)}{(x - t)^2} dt = f'_x(x) - \frac{1}{\pi} \int_a^b K'_x(x, t) \varphi(t) dt, \quad a \leq x \leq b. \quad (21)$$

A bounded solution of equation (21) can be constructed by resolving this equation with respect to the right-hand side, and thereby reducing it to a Fredholm equation of the second kind. Indeed,

$$\varphi(x) - \int_a^b N(x, t) \varphi(t) dt = F(x), \quad a \leq x \leq b,$$

where

$$\begin{aligned}N(x, t) &= -\frac{\sqrt{(x - a)(b - x)}}{\pi^2} \int_a^b \frac{K(\tau, t)}{\sqrt{(\tau - a)(b - \tau)} \tau - x} d\tau, \\ F(x) &= -\frac{\sqrt{(x - a)(b - x)}}{\pi} \int_a^b \frac{f(\tau)}{\sqrt{(\tau - a)(b - \tau)} \tau - x} d\tau.\end{aligned}$$

The problem of solving Fredholm equations of the second kind is considered in detail in Chapter 13.

© References for Section 14.6: N. I. Muskhelishvili (1968), A. I. Kalandia (1973), F. D. Gakhov (1977, 1990), F. D. Gakhov and Yu. I. Cherskii (1978), S. G. Mikhlín and S. Prössdorf (1986), S. Prössdorf and B. Silbermann (1991), I. K. Lifanov (1996), S. G. Samko (2000), G. Iovane, I. K. Lifanov, and M. A. Sumbatyan (2003), M. A. I. K. Lifanov, L. N. Poltavskii, and G. M. Vainikko (2004).

Chapter 15

Methods for Solving Complete Singular Integral Equations

15.1. Some Definitions and Remarks

15.1-1. Integral Equations with Cauchy Kernel.

A complete singular integral equation with Cauchy kernel has the form

$$a(t)\varphi(t) + \frac{1}{\pi i} \int_L \frac{M(t, \tau)}{\tau - t} \varphi(\tau) d\tau = f(t), \quad i^2 = -1, \quad (1)$$

where the integral, which is understood in the sense of the Cauchy principal value, is taken over a closed or nonclosed contour L and t and τ are the complex coordinates of points of the contour. It is assumed that the functions $a(t)$, $f(t)$, and $M(t, \tau)$ given on L and the unknown function $\varphi(t)$ satisfy the Hölder condition (see Subsection 14.2-2), and $M(t, \tau)$ satisfies this condition with respect to both variables.

The integral in Eq. (1) can also be written in a frequently used equivalent form. To this end, we consider the following transformation of the kernel:

$$\frac{M(t, \tau)}{\tau - t} = \frac{M(t, \tau) - M(t, t)}{\tau - t} + \frac{M(t, t)}{\tau - t}, \quad (2)$$

where we set

$$M(t, t) = b(t), \quad \frac{1}{\pi i} \frac{M(t, \tau) - M(t, t)}{\tau - t} = K(t, \tau). \quad (3)$$

In this case Eq. (1), with regard to (2) and (3), becomes

$$a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L K(t, \tau)\varphi(\tau) d\tau = f(t). \quad (4)$$

It follows from formulas (3) that the function $b(t)$ satisfies the Hölder condition on the entire contour L and $K(t, \tau)$ satisfies the Hölder condition everywhere except for the points $\tau = t$, at which one has the estimate

$$|K(t, \tau)| < \frac{A}{|\tau - t|^\lambda}, \quad A = \text{const} < \infty, \quad 0 \leq \lambda < 1.$$

Naturally, Eq. (4) is also called a *complete singular integral equation with Cauchy kernel*. The functions $a(t)$ and $b(t)$ are called the *coefficients* of Eq. (4), $\frac{1}{\tau - t}$ is called the *Cauchy kernel*, and the known function $f(t)$ is called the *right-hand side* of the equation. The first and the second terms

on the left-hand side of Eq. (4) form the *characteristic part* or the *characteristic* of the complete singular equation and the third summand is called the *regular part*, and the function $K(t, \tau)$ is called the *kernel of the regular part*. It follows from the above estimate for the kernel of the regular part that $K(t, \tau)$ is a Fredholm kernel.

For Eqs. (1) and (4) we shall use the operator notation

$$\mathbf{K}[\varphi(t)] = f(t), \quad (5)$$

where the operator \mathbf{K} is called a *singular operator*.

The equation

$$\mathbf{K}^\circ[\varphi(t)] \equiv a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau = f(t) \quad (6)$$

is called the *characteristic equation* corresponding to the complete equation (4), and the operator \mathbf{K}° is called the *characteristic operator*.

For the regular part of the equation we introduce the notation

$$\mathbf{K}_r[\varphi(t)] \equiv \int_L K(t, \tau)\varphi(\tau) d\tau,$$

where the operator \mathbf{K}_r is called a *regular (Fredholm) operator*, and we rewrite the complete singular equation in another operator form:

$$\mathbf{K}[\varphi(t)] \equiv \mathbf{K}^\circ[\varphi(t)] + \mathbf{K}_r[\varphi(t)] = f(t), \quad (7)$$

which will be used in what follows.

The equation

$$\mathbf{K}^*[\psi(t)] \equiv a(t)\psi(t) - \frac{1}{\pi i} \int_L \frac{b(\tau)\psi(\tau)}{\tau - t} d\tau + \int_L K(\tau, t)\psi(\tau) d\tau = g(t), \quad (8)$$

obtained from Eq. (4) by transposing the variables in the kernel, is said to be *transposed* to (4). The operator \mathbf{K}^* is said to be *transposed* to the operator \mathbf{K} .

In particular, the equation

$$\mathbf{K}^{\circ*}[\psi(t)] \equiv a(t)\psi(t) - \frac{1}{\pi i} \int_L \frac{b(\tau)}{\tau - t} \psi(\tau) d\tau = g(t) \quad (9)$$

is the equation transposed to the characteristic equation (6). It should be noted that the operator $\mathbf{K}^{\circ*}$ transposed to the characteristic operator \mathbf{K}° differs from the operator $\mathbf{K}^{*\circ}$ that is characteristic for the transposed equation (9). The latter is defined by the formula

$$\mathbf{K}^{*\circ}[\psi(t)] \equiv a(t)\psi(t) - \frac{b(t)}{\pi i} \int_L \frac{\psi(\tau)}{\tau - t} d\tau. \quad (10)$$

Throughout the following we assume that in the general case the contour L consists of $m + 1$ closed smooth curves $L = L_0 + L_1 + \dots + L_m$. For equations with nonclosed contours, see, for example, the books by F. D. Gakhov (1977, 1990) and N. I. Muskhelishvili (1992).

Remark 1. The above relationship between Eqs. (1) and (4) that involves the properties of these equations is violated if we modify the condition and assume that in Eq. (1) the function $M(t, \tau)$ satisfies the Hölder condition everywhere on the contour except for finitely many points at which M has jump discontinuities. In this case, the complete singular integral equation must be represented in the form (4) with separated characteristic and regular parts in some way that differs from the transformation (2) and (3) because the above transformation of Eq. (1) does not lead to the desired decomposition. For equations with discontinuous coefficients, see the cited books.

15.1-2. Integral Equations with Hilbert Kernel.

A complete singular integral equation with Hilbert kernel has the form

$$a(x)\varphi(x) + \frac{1}{2\pi} \int_0^{2\pi} N(x, \xi) \cot \frac{\xi - x}{2} \varphi(\xi) d\xi = f(x), \quad (11)$$

where the real functions $a(x)$, $f(x)$, and $N(x, \xi)$ and the unknown function $\varphi(x)$ satisfy the Hölder condition (see Subsection 14.2-2), with the function $N(x, \xi)$ satisfying the condition with respect to both variables.

The integral equation (11) can also be written in the following equivalent form, which is frequently used. We transform the kernel as follows:

$$N(x, \xi) \cot \frac{\xi - x}{2} = [N(x, \xi) - N(x, x)] \cot \frac{\xi - x}{2} + N(x, x) \cot \frac{\xi - x}{2}, \quad (12)$$

where we write

$$N(x, x) = -b(x), \quad \frac{1}{2\pi} [N(x, \xi) - N(x, x)] \cot \frac{\xi - x}{2} = K(x, \xi). \quad (13)$$

In this case, Eq. (11) with regard to (12) and (13) becomes

$$a(x)\varphi(x) - \frac{b(x)}{2\pi} \int_0^{2\pi} \cot \frac{\xi - x}{2} \varphi(\xi) d\xi + \int_0^{2\pi} K(x, \xi)\varphi(\xi) d\xi = f(x). \quad (14)$$

It follows from formulas (13) that the function $b(x)$ satisfies the Hölder condition, and the kernel $K(x, \xi)$ satisfies the Hölder condition everywhere except possibly for the points $x = \xi$ at which the following estimate holds:

$$|K(x, \xi)| < \frac{A}{|\xi - x|^\lambda}, \quad A = \text{const} < \infty, \quad 0 \leq \lambda < 1.$$

The equation in the form (14) is also called a complete singular integral equation with Hilbert kernel. The functions $a(x)$ and $b(x)$ are called the *coefficients* of Eq. (14), $\cot \left[\frac{1}{2}(\xi - x) \right]$ is called the *Hilbert kernel*, and the known function $f(x)$ is called the *right-hand side* of the equation. The first and second summands in Eq. (14) form the so-called *characteristic part* or the *characteristic* of the complete singular equation, and the third summand is called its *regular part*; the function $K(x, \xi)$ is called the *kernel of the regular part*.

The equation

$$a(x)\varphi(x) - \frac{b(x)}{2\pi} \int_0^{2\pi} \cot \frac{\xi - x}{2} \varphi(\xi) d\xi = f(x) \quad (15)$$

is called the *characteristic equation* corresponding to the complete equation (14).

As usual, the above and the forthcoming equations whose right-hand sides are zero everywhere on their domains are said to be *homogeneous*, and otherwise they are said to be *nonhomogeneous*.

15.1-3. Fredholm Equations of the Second Kind on a Contour.

Fredholm theory and methods for solving Fredholm integral equations of the second kind presented in Chapter 13 remain valid if all functions and parameters in the equations are treated as complex ones and an interval of the real axis is replaced by a contour L . Here we present only some information and write the Fredholm integral equation of the second kind in the form that is convenient for the purposes of this chapter.

Consider the Fredholm integral equation

$$\varphi(t) + \lambda \int_L K(t, \tau) \varphi(\tau) d\tau = f(t), \quad (16)$$

where L is a smooth contour, t and τ are complex coordinates of its points, $\varphi(t)$ is the desired function, $f(t)$ is the right-hand side of the equation, and $K(t, \tau)$ is the kernel.

If for some λ , the homogeneous Fredholm equation has a nontrivial solution (or nontrivial solutions), then λ is called a *characteristic value*, and the nontrivial solutions themselves are called *eigenfunctions* of the kernel $K(t, \tau)$ or of Eq. (16).

The set of characteristic values of Eq. (16) is at most countable. If this set is infinite, then its only limit point is the point at infinity. To each characteristic value, there are corresponding finitely many linearly independent eigenfunctions. The set of characteristic values of an integral equation is called its *spectrum*. The spectrum of a Fredholm integral equation is a discrete set.

If λ does not coincide with any characteristic value (in this case the value λ is said to be regular), i.e., the homogeneous equation has only the trivial solution, then the nonhomogeneous equation (16) is solvable for any right-hand side $f(t)$.

The general solution is given by the formula

$$\varphi(t) = f(t) - \int_L R(t, \tau; \lambda) f(\tau) d\tau, \quad (17)$$

where the function $R(t, \tau; \lambda)$ is called the *resolvent of the equation* or the *resolvent of the kernel* $K(t, \tau)$ and can be expressed via $K(t, \tau)$.

If a value of the parameter λ is characteristic for Eq. (16), then the homogeneous integral equation

$$\varphi(t) + \lambda \int_L K(t, \tau) \varphi(\tau) d\tau = 0, \quad (18)$$

as well as the transposed homogeneous equation

$$\psi(t) + \lambda \int_L K(\tau, t) \varphi(\tau) d\tau = 0, \quad (19)$$

has nontrivial solutions, and the number of solutions of Eq. (18) is finite and is equal to the number of linearly independent solutions of Eq. (19).

The general solution of the homogeneous equation can be represented in the form

$$\varphi(t) = \sum_{k=1}^n C_k \varphi_k(t), \quad (20)$$

where $\varphi_1(t), \dots, \varphi_n(t)$ is a (complete) finite set of linearly independent eigenfunctions that correspond to the characteristic value λ , and C_k are arbitrary constants.

If the homogeneous equation (18) is solvable, then the nonhomogeneous equation (16) is, in general, unsolvable. This equation is solvable if and only if the following conditions hold:

$$\int_L f(t) \psi_k(t) dt = 0, \quad (21)$$

where $\{\psi_k(t)\}$ ($k = 1, \dots, n$) is a (complete) finite set of linearly independent eigenfunctions of the transposed equation that correspond to the characteristic value λ .

If conditions (21) are satisfied, then the general solution of the nonhomogeneous equation (16) can be given by the formula (e.g., see Subsection 15.6-5)

$$\varphi(t) = f(t) - \int_L R_g(t, \tau; \lambda) f(\tau) d\tau + \sum_{k=1}^n C_k \varphi_k(t), \quad (22)$$

where $R_g(t, \tau; \lambda)$ is called the *generalized resolvent* and the sum on the right-hand side of (22) is the general solution of the corresponding homogeneous equation.

Now we consider an equation of the second kind with weak singularity on the contour:

$$\varphi(t) + \int_L \frac{M(t, \tau)}{|\tau - t|^\alpha} \varphi(\tau) d\tau = f(t), \quad (23)$$

where $M(t, \tau)$ is a continuous function and $0 < \alpha < 1$. By iterating we can reduce this equation to a Fredholm integral equation of the second kind (e.g., see Remark 1 in Section 13.3). It has all properties of a Fredholm equation.

For the above reasons, in the theory of singular integral equations it is customary to make no difference between Fredholm equations and equations with weak singularity and use for them the same notation

$$\varphi(t) + \lambda \int_L K(t, \tau) \varphi(\tau) d\tau = 0, \quad K(t, \tau) = \frac{M(t, \tau)}{|\tau - t|^\alpha}, \quad 0 \leq \alpha < 1. \quad (24)$$

The integral equation (24) is called simply a *Fredholm equation*, and its kernel is called a *Fredholm kernel*.

If in Eq. (24) the known functions satisfy the Hölder condition, and $M(t, \tau)$ satisfies this condition with respect to both variables, then each bounded integrable solution of Eq. (24) also satisfies the Hölder condition.

Remark 2. By the above estimates, the kernels of the regular parts of the above singular integral equations are Fredholm kernels.

Remark 3. The complete and characteristic singular integral equations are sometimes called singular integral equations of the second kind.

⊙ References for Section 15.1: F. D. Gakhov (1977, 1990), F. G. Tricomi (1985), S. G. Mikhlin and S. Prössdorf (1986), A. Dzhuraev (1992), N. I. Muskhelishvili (1992), I. K. Lifanov (1996), R. Estrada and R. P. Kanwal (1999), E. G. Ladopoulos (2000).

15.2. Carleman Method for Characteristic Equations

15.2-1. Characteristic Equation with Cauchy Kernel.

Consider a characteristic equation with Cauchy kernel:

$$\mathbf{K}^\circ[\varphi(t)] \equiv a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau = f(t), \quad (1)$$

where the contour L consists of $m + 1$ closed smooth curves $L = L_0 + L_1 + \cdots + L_m$.

Solving Eq. (1) can be reduced to solving a Riemann boundary value problem (see Subsection 14.3-10), and the solution of the equation can be presented in a closed form.

Let us introduce the piecewise analytic function given by the Cauchy integral whose density is the desired solution of the characteristic equation:

$$\Phi(z) = \frac{1}{2\pi i} \int_L \frac{\varphi(\tau)}{\tau - z} d\tau. \quad (2)$$

According to the Sokhotski–Plemelj formulas (see Subsection 14.2-5), we have

$$\begin{aligned}\varphi(t) &= \Phi^+(t) - \Phi^-(t), \\ \frac{1}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - z} d\tau &= \Phi^+(t) + \Phi^-(t).\end{aligned}\quad (3)$$

On substituting (3) into (1) and solving the resultant equation for $\Phi^+(t)$, we see that the piecewise analytic function $\Phi(z)$ must be a solution of the Riemann boundary value problem

$$\Phi^+(t) = D(t)\Phi^-(t) + H(t), \quad (4)$$

where

$$D(t) = \frac{a(t) - b(t)}{a(t) + b(t)}, \quad H(t) = \frac{f(t)}{a(t) + b(t)}. \quad (5)$$

Since the function $\Phi(z)$ is represented by a Cauchy type integral, it follows that this function must satisfy the additional condition

$$\Phi^-(\infty) = 0. \quad (6)$$

The index ν of the coefficient $D(t)$ of the Riemann problem (4) is called the *index of the integral equation* (1). On solving the boundary value problem (4), we find the solution of Eq. (1) by the first formula in (3).

Thus, the integral equation (1) is reduced to the Riemann boundary value problem (4). To establish the equivalence of the equation to the boundary value problem we note that, conversely, the function $\varphi(t)$ that is found by the above-mentioned method from the solution of the boundary value problem necessarily satisfies Eq. (1).

We first consider the following normal (nonexceptional) case in which the coefficient $D(t)$ of the Riemann problem (4) admits no zero or infinite values, which amounts to the condition

$$a(t) \pm b(t) \neq 0 \quad (7)$$

for Eq. (1). To simplify the subsequent formulas, we assume that the coefficients of Eq. (1) satisfy the condition

$$a^2(t) - b^2(t) = 1. \quad (8)$$

This can always be achieved by dividing the equation by $\sqrt{a^2(t) - b^2(t)}$.

Let us write out the solution of the Riemann boundary value problem (4) under the assumption $\nu \geq 0$ and then use the Sokhotski–Plemelj formulas to find the limit values of the corresponding functions (see Subsections 14.2-5, 14.3-6, and 14.3-10):

$$\Phi^+(t) = X^+(t) \left[\frac{1}{2} \frac{H(t)}{X^+(t)} + \Psi(t) - \frac{1}{2} P_{\nu-1}(t) \right], \quad \Phi^-(t) = X^-(t) \left[-\frac{1}{2} \frac{H(t)}{X^+(t)} + \Psi(t) - \frac{1}{2} P_{\nu-1}(t) \right], \quad (9)$$

where

$$\Psi(t) = \frac{1}{2\pi i} \int_L \frac{H(\tau)}{X^+(\tau)} \frac{d\tau}{\tau - t}. \quad (10)$$

The arbitrary polynomial is taken in the form $-\frac{1}{2}P_{\nu-1}(t)$, which is convenient for the subsequent notation.

Hence, by formula (3) we have

$$\varphi(t) = \frac{1}{2} \left[1 + \frac{X^-(t)}{X^+(t)} \right] H(t) + X^+(t) \left[1 - \frac{X^-(t)}{X^+(t)} \right] \left[\Psi(t) - \frac{1}{2} P_{\nu-1}(t) \right].$$

Representing the coefficient of the Riemann problem in the form $D(t) = X^+(t)/X^-(t)$ and replacing the function $\Psi(t)$ by the expression on the right-hand side in (10), we obtain

$$\varphi(t) = \frac{1}{2} \left[1 + \frac{1}{D(t)} \right] H(t) + X^+(t) \left[1 - \frac{1}{D(t)} \right] \left[\frac{1}{2\pi i} \int_L \frac{H(\tau)}{X^+(\tau)} \frac{d\tau}{\tau - t} - \frac{1}{2} P_{\nu-1}(t) \right].$$

Finally, on replacing $X^+(t)$ by the expression (62) in Subsection 14.3-10 and substituting the expressions for $D(t)$ and $H(t)$ given in (5), we obtain

$$\varphi(t) = a(t)f(t) - \frac{b(t)Z(t)}{\pi i} \int_L \frac{f(\tau)}{Z(\tau)} \frac{d\tau}{\tau - t} + b(t)Z(t)P_{\nu-1}(t), \quad (11)$$

where

$$Z(t) = [a(t) + b(t)]X^+(t) = [a(t) - b(t)]X^-(t) = \frac{e^{G(t)}}{\sqrt{t^\nu \Pi(t)}}, \quad (12)$$

$$G(t) = \frac{1}{2\pi i} \int_L \ln \left[\tau^{-\nu} \Pi(\tau) \frac{a(\tau) - b(\tau)}{a(\tau) + b(\tau)} \right] \frac{d\tau}{\tau - t}, \quad \Pi(t) = \sum_{k=1}^m (t - z_k)^{\nu_k},$$

and the coefficients $a(t)$ and $b(t)$ satisfy condition (7). Here $\Pi(t) \equiv 1$ for the case in which L is a simple contour enclosing a simply connected domain. Since the functions $a(t)$, $b(t)$, and $f(t)$ satisfy the Hölder condition, it follows from the properties of the limit values of the Cauchy type integral that the function $\varphi(t)$ also satisfies the Hölder condition.

The last term in formula (11) is the general solution of the homogeneous equation ($f(t) \equiv 0$), and the first two terms form a particular solution of the nonhomogeneous equation.

The particular solution of Eq. (1) can be represented in the form $\mathbf{R}[f(t)]$, where \mathbf{R} is the operator defined by

$$\mathbf{R}[f(t)] = a(t)f(t) - \frac{b(t)Z(t)}{\pi i} \int_L \frac{f(\tau)}{Z(\tau)} \frac{d\tau}{\tau - t}.$$

In this case, the general solution of Eq. (1) becomes

$$\varphi(t) = \mathbf{R}[f(t)] + \sum_{k=1}^{\nu} c_k \varphi_k(t), \quad (13)$$

where $\varphi_k(t) = b(t)Z(t)t^{k-1}$ ($k = 1, 2, \dots, \nu$) are the linearly independent eigenfunctions of the characteristic equation.

If $\nu < 0$, then the Riemann problem (4) is in general unsolvable. The solvability conditions

$$\int_L \frac{H(\tau)}{X^+(\tau)} \tau^{k-1} d\tau = 0, \quad k = 1, 2, \dots, -\nu, \quad (14)$$

for problem (4) are the solvability conditions for Eq. (1) as well.

Replacing $H(\tau)$ and $X^+(\tau)$ by their expressions from (5) and (12), we can rewrite the solvability conditions in the form

$$\int_L \frac{f(\tau)}{Z(\tau)} \tau^{k-1} d\tau = 0, \quad k = 1, 2, \dots, -\nu. \quad (15)$$

If the solvability conditions hold, then the solution of the nonhomogeneous equation (4) is given by formula (11) for $P_{\nu-1} \equiv 0$.

1.° If $\nu > 0$, then the homogeneous equation $\mathbf{K}^\circ[\varphi(t)] = 0$ has ν linearly independent solutions

$$\varphi_k(t) = b(t)Z(t)t^{k-1}, \quad k = 1, 2, \dots, \nu.$$

2.° If $\nu \leq 0$, then the homogeneous equation is unsolvable (has only the trivial solution).

3.° If $\nu \geq 0$, then the nonhomogeneous equation is solvable for an arbitrary right-hand side $f(t)$, and its general solution linearly depends on ν arbitrary constants.

4.° If $\nu < 0$, then the nonhomogeneous equation is solvable if and only if its right-hand side f satisfies the $-\nu$ conditions,

$$\int_L \psi_k(t) f(t) dt = 0, \quad \psi_k(t) = \frac{t^{k-1}}{Z(t)}. \quad (16)$$

The above properties of characteristic singular integral equations are essentially different from the properties of Fredholm integral equations (see Subsection 15.1-3). With Fredholm equations, if the homogeneous equation is solvable, then the nonhomogeneous equation is in general unsolvable, and conversely, if the homogeneous equation is unsolvable, then the nonhomogeneous equation is solvable. However, for a singular equation, if the homogeneous equation is solvable, then the nonhomogeneous equation is unconditionally solvable, and if the homogeneous equation is unsolvable, then the nonhomogeneous equation is in general unsolvable as well.

By analogy with the case of Fredholm equations, we introduce a parameter λ into the kernel of the characteristic equation and consider the equation

$$a(t)\varphi(t) + \frac{\lambda b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau = 0.$$

As shown above, the last equation is solvable if

$$\nu = \text{Ind} \frac{a(t) - \lambda b(t)}{a(t) + \lambda b(t)} > 0.$$

The index of a continuous function changes by jumps and only for the values of λ such that $a(t) \mp \lambda b(t) = 0$. If in the complex plane $\lambda = \lambda_1 + i\lambda_2$ we draw the curves $\lambda = \pm a(t)/b(t)$, then these curves divide the plane into domains in each of which the index is constant. Thus, the characteristic values of the characteristic integral equation occupy entire domains, and hence the spectrum is continuous, in contrast with the spectrum of a Fredholm equation.

15.2-2. Transposed Equation of a Characteristic Equation.

The equation

$$\mathbf{K}^{\circ*}[\psi(t)] \equiv a(t)\psi(t) - \frac{1}{\pi i} \int_L \frac{b(\tau)\psi(\tau)}{\tau - t} d\tau = g(t), \quad (17)$$

which is transposed to the characteristic equation $\mathbf{K}^{\circ}[\varphi(t)] = f(t)$, is not characteristic. However, the substitution

$$b(t)\psi(t) = \omega(t) \quad (18)$$

reduces it to a characteristic equation for the function $\omega(t)$:

$$a(t)\omega(t) - \frac{b(t)}{\pi i} \int_L \frac{\omega(\tau)}{\tau - t} d\tau = b(t)g(t). \quad (19)$$

From the last equation we find $\omega(t)$, by the formula obtained by adding (17) to (18), and determine the desired function $\psi(t)$:

$$\psi(t) = \frac{1}{a(t) + b(t)} \left[\omega(t) + \frac{1}{\pi i} \int_L \frac{\omega(\tau)}{\tau - t} d\tau + g(t) \right].$$

Introducing the piecewise analytic function

$$\Phi_*(z) = \frac{1}{2\pi i} \int_L \frac{\omega(\tau)}{\tau - z} d\tau, \quad (20)$$

we arrive at the Riemann boundary value problem

$$\Phi_*^+(t) = \frac{a(t) + b(t)}{a(t) - b(t)} \Phi_*^-(t) + \frac{b(t)g(t)}{a(t) - b(t)}. \quad (21)$$

The coefficient of the boundary value problem (21) is the inverse of the coefficient of the Riemann problem (4) corresponding to the equation $\mathbf{K}^\circ[\varphi(t)] = f(t)$. Hence,

$$\nu^* = \text{Ind} \frac{a(t) + b(t)}{a(t) - b(t)} = -\text{Ind} \frac{a(t) - b(t)}{a(t) + b(t)} = -\nu. \quad (22)$$

Note that it follows from formulas (17) in Subsection 14.3-4 that the canonical function $X^*(z)$ for Eq. (21) and the canonical function $X(z)$ for (4) are reciprocal:

$$X^*(z) = \frac{1}{X(z)}.$$

By analogy with the reasoning in Subsection 15.2-1, we obtain a solution of the singular integral equation (17) for $\nu^* = -\nu \geq 0$ in the form

$$\psi(t) = a(t)g(t) + \frac{1}{\pi i Z(t)} \int_L \frac{b(\tau)Z(\tau)g(\tau)}{\tau - t} d\tau + \frac{1}{Z(t)} Q_{\nu^*-1}(t), \quad (23)$$

where $Z(t)$ is given by formula (12) and $Q_{\nu^*-1}(t)$ is a polynomial of degree at most $\nu^* - 1$ with arbitrary coefficients. If $\nu^* = 0$, then we must set $Q_{\nu^*-1}(t) \equiv 0$.

If $\nu^* = -\nu < 0$, then for the solvability of Eq. (17) it is necessary and sufficient that

$$\int_L b(t)Z(t)g(t)t^{k-1} dt = 0, \quad k = 1, 2, \dots, -\nu^*, \quad (24)$$

and if these conditions hold, then the solution is given by formula (23), where we must set $Q_{\nu^*-1}(t) \equiv 0$.

The results of simultaneous investigation of a characteristic equation and the transposed equation show another essential difference from the properties of Fredholm equations (see Subsection 15.1-3). Transposed homogeneous characteristic equations cannot be solvable simultaneously. Either they are both unsolvable ($\nu = 0$), or, for a nonzero index, only the equation with a positive index is solvable.

We point out that the difference between the numbers of solutions of a characteristic homogeneous equation and the transposed equation is equal to the index ν .

Assertions 1^o and 2^o and assertions 3^o and 4^o in Subsection 15.2-1 are called, respectively, the *first Fredholm theorem* and the *second Fredholm theorem* for a characteristic equation, and the relationship between the index of an equation and the number of solutions of the homogeneous equations $\mathbf{K}^\circ[\varphi(t)] = 0$ and $\mathbf{K}^{\circ*}[\psi(t)] = 0$ is called the *third Fredholm theorem*.

15.2-3. Characteristic Equation on the Real Axis.

The theory of the Cauchy type integral (see Section 14.2) shows that if the density of the Cauchy type integral taken over an infinite curve vanishes at infinity, then the properties of the integral for the cases in which the contour is finite and infinite are essentially the same. Therefore, the theory of singular integral equations on an infinite contour in the class of functions that vanish at infinity coincides with the theory of equations on a finite contour.

Just as for the case of a finite contour, the characteristic integral equation

$$a(x)\varphi(x) + \frac{b(x)}{\pi i} \int_{-\infty}^{\infty} \frac{\varphi(\tau)}{\tau - x} d\tau = f(x) \quad (25)$$

can be reduced by means of the Cauchy type integral

$$\Phi(z) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\varphi(\tau)}{\tau - z} d\tau \quad (26)$$

and the Sokhotski–Plemelj formulas (see Subsection 14.2-5), to the following Riemann boundary value problem for the real axis (see Subsection 14.3-8):

$$\Phi^+(x) = \frac{a(x) - b(x)}{a(x) + b(x)} \Phi^-(x) + \frac{f(x)}{a(x) + b(x)}, \quad -\infty < x < \infty. \quad (27)$$

We assume that

$$a^2(x) - b^2(x) = 1, \quad (28)$$

because Eq. (25) can always be reduced to case (28) by the division by $\sqrt{a^2(t) - b^2(t)}$. Note that the index ν of the integral equation (25) is given by the formula

$$\nu = \text{Ind} \frac{a(x) - b(x)}{a(x) + b(x)}. \quad (29)$$

In this case for $\nu \geq 0$ we obtain

$$\varphi(x) = a(x)f(x) - \frac{b(x)Z(x)}{\pi i} \int_{-\infty}^{\infty} \frac{f(\tau)}{Z(\tau)} \frac{d\tau}{\tau - x} + b(x)Z(x) \frac{P_{\nu-1}(x)}{(x+i)^\nu}, \quad (30)$$

where

$$Z(x) = [a(x) + b(x)]X^+(x) = [a(x) - b(x)]X^-(x) = \left(\frac{x-i}{x+i}\right)^{-\nu/2} e^{G(x)},$$

$$G(x) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \ln \left[\left(\frac{\tau-i}{\tau+i}\right)^{-\nu} \frac{a(\tau) - b(\tau)}{a(\tau) + b(\tau)} \right] \frac{d\tau}{\tau - x}.$$

For the case in which $\nu \leq 0$ we must set $P_{\nu-1}(x) \equiv 0$. For $\nu < 0$, we must also impose the solvability conditions

$$\int_{-\infty}^{\infty} \frac{f(x)}{Z(x)} \frac{dx}{(x+i)^k} = 0, \quad k = 1, 2, \dots, -\nu. \quad (31)$$

For the solution of Eq. (25) in the class of functions bounded at infinity, see F. D. Gakhov (1977, 1990).

The analog of the characteristic equation on the real axis is the equation of the form

$$a(x)\varphi(x) + \frac{b(x)}{\pi i} \int_{-\infty}^{\infty} \frac{x - z_0}{\tau - z_0} \frac{\varphi(\tau)}{\tau - x} d\tau = f(x), \quad (32)$$

where z_0 is a point that does not belong to the contour. For this equation, all qualitative results obtained for the characteristic equation with finite contour are still valid together with the formulas. In particular, the following inversion formulas for the Cauchy type integral hold:

$$\psi(x) = \frac{1}{\pi i} \int_{-\infty}^{\infty} \frac{x - z_0}{\tau - z_0} \frac{\varphi(\tau)}{\tau - x} d\tau, \quad \varphi(x) = \frac{1}{\pi i} \int_{-\infty}^{\infty} \frac{x - z_0}{\tau - z_0} \frac{\psi(\tau)}{\tau - x} d\tau. \quad (33)$$

15.2-4. Exceptional Case of a Characteristic Equation.

In the study of the characteristic equation in Subsection 15.2-1, the case in which the functions $a(t) \pm b(t)$ can vanish on the contour L was excluded. The reason was that the coefficient $D(t)$ of the Riemann problem to which the characteristic equation can be reduced has in the exceptional case zeros and poles on the contour, and hence this problem is outside the framework of the general theory. Let us perform an investigation of the above exceptional case.

We assume that the coefficients of the singular equations under consideration have properties that provide the additional differentiability requirements that were introduced in the consideration of exceptional cases of the Riemann problem (see 14.3-9).

Consider a characteristic equation with Cauchy kernel (1) under the assumption that the functions $a(t)-b(t)$ and $a(t)+b(t)$ have zeros on the contour at the points $\alpha_1, \dots, \alpha_\mu$ and $\beta_1, \dots, \beta_\eta$, respectively, of integral orders, and hence are representable in the form

$$a(t) - b(t) = \prod_{k=1}^{\mu} (t - \alpha_k)^{m_k} r(t), \quad a(t) + b(t) = \prod_{j=1}^{\eta} (t - \beta_j)^{p_j} s(t),$$

where $r(t)$ and $s(t)$ vanish nowhere. We assume that all points α_k and β_j are different.

Assume that the coefficients of Eq. (1) satisfy the relation

$$a^2(t) - b^2(t) = \prod_{k=1}^{\mu} (t - \alpha_k)^{m_k} \prod_{j=1}^{\eta} (t - \beta_j)^{p_j} = A_0(t). \tag{34}$$

The equation under consideration can be reduced to the above case by dividing it by $\sqrt{s(t)r(t)}$.

In the exceptional case, by analogy with the case studied in Subsection 15.2-1, Eq. (1) can be reduced to the Riemann problem

$$\Phi^+(t) = \frac{\prod_{k=1}^{\mu} (t - \alpha_k)^{m_k}}{\prod_{j=1}^{\eta} (t - \beta_j)^{p_j}} D_1(t) \Phi^-(t) + \frac{f(t)}{\prod_{j=1}^{\eta} (t - \beta_j)^{p_j} s(t)}, \tag{35}$$

where $D_1(t) = r(t)/s(t)$. The solution of this problem in the class of functions that satisfy the condition $\Phi(\infty) = 0$ is given by the formulas

$$\begin{aligned} \Phi^+(z) &= \frac{X^+(z)}{\eta} [\Psi^+(z) - \mathcal{U}_\rho(z) + A_0(z)P_{\nu-p-1}(z)], \\ &\quad \prod_{j=1}^{\eta} (z - \beta_j)^{p_j} \\ \Phi^-(z) &= \frac{X^-(z)}{\mu} [\Psi^-(z) - \mathcal{U}_\rho(z) + A_0(z)P_{\nu-p-1}(z)], \\ &\quad \prod_{k=1}^{\mu} (z - \alpha_k)^{m_k} \end{aligned} \tag{36}$$

where

$$\Psi(z) = \frac{1}{2\pi i} \int_L \frac{f(\tau)}{s(\tau)X^+(\tau)} \frac{d\tau}{\tau - z}, \tag{37}$$

and $\mathcal{U}_\rho(z)$ is the Hermite interpolation polynomial (see Subsection 14.3-2) for the function $\Psi(z)$ of degree $\rho = m + p - 1$ with nodes at the points α_k and β_j , respectively, and of the multiplicities m_k and p_j , respectively, where $m = \sum m_k$ and $p = \sum p_j$.

We regard the polynomial $\mathcal{U}_\rho(z)$ as an operator that maps the right-hand side $f(t)$ of Eq. (1) to the polynomial that interpolates the Cauchy type integral (37) as above. Let us denote this operator by

$$\frac{1}{2}\mathbf{T}[f(t)] = \mathcal{U}_\rho(z). \tag{38}$$

Here the coefficient $\frac{1}{2}$ is taken for the convenience of the subsequent manipulations.

Furthermore, by analogy with the normal case, from (36) we can find

$$\begin{aligned} \Phi^+(t) &= \frac{X^+(t)}{\prod_{j=1}^{\eta}(t-\beta_j)^{p_j}} \left[\frac{1}{2} \frac{f(t)}{s(t)X^+(t)} + \frac{1}{2\pi i} \int_L \frac{f(\tau)}{s(\tau)X^+(\tau)} \frac{d\tau}{\tau-t} - \frac{1}{2}\mathbf{T}[f(t)] - \frac{1}{2}A_0(t)P_{\nu-p-1}(t) \right], \\ \Phi^-(t) &= \frac{X^-(t)}{\prod_{k=1}^{\mu}(t-\alpha_k)^{m_k}} \left[-\frac{1}{2} \frac{f(t)}{s(t)X^+(t)} + \frac{1}{2\pi i} \int_L \frac{f(\tau)}{s(\tau)X^+(\tau)} \frac{d\tau}{\tau-t} - \frac{1}{2}\mathbf{T}[f(t)] - \frac{1}{2}A_0(t)P_{\nu-p-1}(t) \right]. \end{aligned}$$

We introduced the coefficient $-\frac{1}{2}$ in the last summands of these formulas using the fact that the coefficients of the polynomial $P_{\nu-p-1}(t)$ are arbitrary. Hence,

$$\varphi(t) = \Phi^+(t) - \Phi^-(t) = \frac{\Delta_1(t)f(t)}{s(t)X^+(t)} + \Delta_2(t) \left[\frac{1}{\pi i} \int_L \frac{f(\tau) d\tau}{s(\tau)X^+(\tau)(\tau-t)} - \mathbf{T}[f(t)] - A_0(t)P_{\nu-p-1}(t) \right], \tag{39}$$

where

$$\Delta_1(t) = \frac{X^+(t)}{2 \prod_{j=1}^{\eta}(t-\beta_j)^{p_j}} + \frac{X^-(t)}{2 \prod_{k=1}^{\mu}(t-\alpha_k)^{m_k}}, \quad \Delta_2(t) = \frac{X^+(t)}{2 \prod_{j=1}^{\eta}(t-\beta_j)^{p_j}} - \frac{X^-(t)}{2 \prod_{k=1}^{\mu}(t-\alpha_k)^{m_k}}.$$

We write

$$Z(t) = s(t)X^+(t) = r(t)X^-(t), \tag{40}$$

and, applying relation (34), represent formula (39) as follows:

$$\varphi(t) = \frac{1}{A_0(t)} \left[a(t)f(t) - \frac{b(t)Z(t)}{\pi i} \int_L \frac{f(\tau)}{Z(\tau)} \frac{d\tau}{\tau-t} + b(t)Z(t)\mathbf{T}[f(t)] \right] + b(t)Z(t)P_{\nu-p-1}(t).$$

Let us introduce the operator $\mathbf{R}_1[f(t)]$ by the formula

$$\mathbf{R}_1[f(t)] \equiv \frac{1}{A_0(t)} \left[a(t)f(t) - \frac{b(t)Z(t)}{\pi i} \int_L \frac{f(\tau)}{Z(\tau)} \frac{d\tau}{\tau-t} + b(t)Z(t)\mathbf{T}[f(t)] \right], \tag{41}$$

and finally obtain

$$\varphi(t) = \mathbf{R}_1[f(t)] + b(t)Z(t)P_{\nu-p-1}(t). \tag{42}$$

Formula (42) gives a solution of Eq. (1) for the exceptional case in which $\nu - p > 0$. This solution linearly depends on $\nu - p$ arbitrary constants. If $\nu - p < 0$, then the solution exists only under $p - \nu$ special solvability conditions imposed on $f(t)$, which follow from the solvability conditions for the Riemann problem (35) corresponding to this case.

15.2-5. Characteristic Equation with Hilbert Kernel.

Consider the characteristic equation with Hilbert kernel

$$a(x)\varphi(x) - \frac{b(x)}{2\pi} \int_0^{2\pi} \cot \frac{\xi - x}{2} \varphi(\xi) d\xi = f(x). \quad (43)$$

Just as the characteristic integral equation with Cauchy kernel is related to the Riemann boundary value problem, so the characteristic equation (43) with Hilbert kernel can be analytically reduced to a Hilbert problem in a straightforward manner. In turn, the Hilbert problem can be reduced to the Riemann problem (see Subsection 14.3-12), and hence the solution of Eq. (43) can be constructed in a closed form.

For $\nu > 0$, the homogeneous equation (43) ($f(x) \equiv 0$) has 2ν linearly independent solutions, and the nonhomogeneous problem is unconditionally solvable and linearly depends on 2ν real constants.

For $\nu < 0$, the homogeneous equation is unsolvable, and the nonhomogeneous equation is solvable only under -2ν real solvability conditions.

Taking into account the fact that any complex parameter contains two real parameters, and a complex solvability condition is equivalent to two real conditions, we see that, for $\nu \neq 0$, the qualitative results of investigating the characteristic equation with Hilbert kernel completely agree with the corresponding results for the characteristic equation with Cauchy kernel.

15.2-6. Tricomi Equation.

The singular integral Tricomi equation has the form

$$\varphi(x) - \lambda \int_0^1 \left(\frac{1}{\xi - x} - \frac{1}{x + \xi - 2x\xi} \right) \varphi(\xi) d\xi = f(x), \quad 0 \leq x \leq 1. \quad (44)$$

The kernel of this equation consists of two terms. The first term is the Cauchy kernel. The second term is continuous if at least one of the variables x and ξ varies strictly inside the interval $[0, 1]$; however, for $x = \xi = 0$ and for $x = \xi = 1$, this kernel becomes infinite and is nonintegrable in the square $\{0 \leq x \leq 1, 0 \leq \xi \leq 1\}$.

By using the function

$$\Phi(z) = \frac{1}{2\pi i} \int_0^1 \left(\frac{1}{\xi - z} - \frac{1}{z + \xi - 2z\xi} \right) \varphi(\xi) d\xi,$$

which is piecewise analytic in the upper and the lower half-plane, we can reduce Eq. (44) to the Riemann problem with boundary condition on the real axis. The solution of the Tricomi equation has the form

$$y(x) = \frac{1}{1 + \lambda^2 \pi^2} \left[f(x) + \int_0^1 \frac{\xi^\alpha (1-x)^\alpha}{x^\alpha (1-\xi)^\alpha} \left(\frac{1}{\xi - x} - \frac{1}{x + \xi - 2x\xi} \right) f(\xi) d\xi \right] + \frac{C(1-x)^\beta}{x^{1+\beta}},$$

$$\alpha = \frac{2}{\pi} \arctan(\lambda\pi) \quad (-1 < \alpha < 1), \quad \tan \frac{\beta\pi}{2} = \lambda\pi \quad (-2 < \beta < 0),$$

where C is an arbitrary constant.

15.3. Complete Singular Integral Equations Solvable in a Closed Form

In contrast with characteristic equations and their transposed equations, complete singular integral equations cannot be solved in the closed form in general. However, there are some cases in which complete equations can be solved in a closed form.

15.3-1. Closed-Form Solutions in the Case of Constant Coefficients.

Consider the complete singular integral equation with Cauchy kernel in the form (see Subsection 15.1-1)

$$a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L K(t, \tau)\varphi(\tau) d\tau = f(t), \quad (1)$$

where L is an arbitrary closed contour. Let us show that Eq. (1) can be solved in a closed form if $a(t) = a$ and $b(t) = b$ are constants and $K(t, \tau)$ is an arbitrary function that has an analytic continuation to the domain Ω^+ with respect to each variable.

Under the above assumptions, Eq. (1) has the form

$$a\varphi(t) + \frac{1}{\pi i} \int_L \frac{M(t, \tau)}{\tau - t} \varphi(\tau) d\tau = f(t), \quad (2)$$

where $M(t, \tau) = b + \pi i(t - \tau)K(t, \tau)$, so that $M(t, t) = b = \text{const}$. Let $b \neq 0$. We write

$$\psi(t) = \frac{1}{b\pi i} \int_L \frac{M(t, \tau)}{\tau - t} \varphi(\tau) d\tau. \quad (3)$$

According to Subsection 14.4-4, the function $\varphi(t)$ can be expressed via $\psi(t)$ and $\psi(t)$ can be expressed via $\varphi(t)$. Then we rewrite Eq. (2) as follows:

$$a\varphi(t) + b\psi(t) = f(t). \quad (4)$$

On applying the operation (3) to this equation, we obtain

$$a\psi(t) + b\varphi(t) = w(t), \quad (5)$$

where

$$w(t) = \frac{1}{b\pi i} \int_L \frac{M(t, \tau)}{\tau - t} f(\tau) d\tau.$$

By solving system (4), (5) we find $\varphi(t)$:

$$\varphi(t) = \frac{1}{a^2 - b^2} \left[af(t) - \frac{1}{\pi i} \int_L \frac{M(t, \tau)}{\tau - t} f(\tau) d\tau \right] \quad (6)$$

under the assumption that $a \neq \pm b$.

Thus, for $a \neq \pm b$ and for a kernel $K(t, \tau)$ that can be analytically continued, Eq. (1) or (2) is solvable and has the unique solution given by formula (6).

Equation (1) was studied above for $b \neq 0$. This assumption is natural because, for $b \equiv 0$, Eq. (1) is no longer singular. However, the Fredholm equation obtained for $b = 0$, that is,

$$a\varphi(t) + \int_L K(t, \tau)\varphi(\tau) d\tau = f(t), \quad a = \text{const}, \quad (7)$$

is solvable in a closed form for a kernel $K(t, \tau)$ that has analytic continuation.

Let a function $K(t, \tau)$ have an analytic continuation to the domain Ω^+ with respect to each of the variables and continuous for $t, \tau \in L$. In this case, the following assertions hold.

1°. The function

$$\Phi^+(t) = \int_L K(t, \tau) \varphi(\tau) d\tau$$

has an analytic continuation to the domain Ω^+ for any function $\varphi(t)$ satisfying the Hölder condition.

2°. If a function $\varphi^+(t)$ satisfying the Hölder condition has an analytic continuation to the domain Ω^+ , then

$$\int_L K(t, \tau) \varphi^+(\tau) d\tau = 0. \quad (8)$$

This implies the relation

$$\int_L K(t, \tau) \int_L K(\tau, \tau_1) \varphi(\tau_1) d\tau_1 d\tau = 0 \quad (9)$$

for each function $\varphi(t)$ (satisfying the Hölder condition). Therefore, it follows from (7) that

$$a \int_L K(t, \tau) \varphi(\tau) d\tau = \int_L K(t, \tau) f(\tau) d\tau,$$

and hence

$$\varphi(t) = \frac{1}{a^2} \left[a f(t) - \int_L K(t, \tau) f(\tau) d\tau \right]. \quad (10)$$

Therefore, if a kernel $K(t, \tau)$ is analytic in the domain Ω^+ with respect to each of the variables and continuous for $t, \tau \in L$, then Eq. (7) is solvable for each right-hand side, and the solution is given by formula (10).

15.3-2. Closed-Form Solutions in the General Case.

Let us pass to the general case of the solvability of Eq. (1) in a closed form under the condition that a function $K(t, \tau)[a(t) + b(t)]^{-1}$ is analytic with respect to τ and meromorphic with respect to t in the domain Ω^+ .

For brevity, we write

$$\mathbf{K}_r[\varphi(t)] = \int_L K(t, \tau) \varphi(\tau) d\tau$$

and note that

$$\mathbf{K}_r[\varphi^+(t)] = 0 \quad (11)$$

for each function $\varphi^+(t)$ that has an analytic continuation to the domain Ω^+ . By setting $\varphi(t) = \varphi^+(t) - \varphi^-(t)$ and with regard to (11), we reduce Eq. (1) to a relation similar to that of the Riemann problem:

$$\varphi^+(t) - \frac{1}{a(t) + b(t)} \mathbf{K}_r[\varphi^-(t)] = D(t) \varphi^-(t) + H(t), \quad (12)$$

where

$$D(t) = \frac{a(t) - b(t)}{a(t) + b(t)}, \quad H(t) = \frac{f(t)}{a(t) + b(t)}.$$

By assumption, we have

$$\frac{K(t, \tau)}{a(t) + b(t)} = \frac{A^+(t, \tau)}{\Pi^+(t)}, \quad \Pi^+(t) = \prod_{k=1}^n (t - z_k)^{m_k}, \quad (13)$$

where $z_k \in \Omega^+$ and m_k are positive integers and the function $A^+(t, \tau)$ is analytic with respect to t and with respect to τ on Ω^+ .

Relation (12) becomes

$$\Pi^+(t)\varphi^+(t) + \mathbf{A}^+[\varphi^-(t)] = \Pi^+(t)[D(t)\varphi^-(t) + H(t)], \tag{14}$$

where \mathbf{A}^+ is the integral operator with kernel $A^+(t, \tau)$. Since the function $\mathbf{A}^+[\varphi^-(t)]$ is analytic on Ω^+ , it follows that the last relation is an ordinary Riemann problem for which the functions $\Pi^+(t)\varphi^+(t) + \mathbf{A}^+[\varphi^-(t)]$ and $\varphi^-(t)$ can be defined in a closed form, and hence the same holds for $\varphi(t)$. Namely, let us rewrite the function $D(t)$ in the form $D(t) = X^+(t)/X^-(t)$, where $X^\pm(z)$ is the canonical function of the Riemann problem, and reduce relation (14) to the form in which the generalized Liouville theorem can be applied (see Subsection 14.3-1). We arrive at a polynomial of degree at most $\nu - 1 + \sum_{k=1}^n m_k$ with arbitrary coefficients (for the case in which $\nu + \sum_{k=1}^n m_k > 0$). However, the presence of the factor $\Pi^+(t)$ (on $\varphi^+(t)$), which vanishes in Ω^+ with total order of zeros $\sum_{k=1}^n m_k$, clearly reduces the number of arbitrary constants in the general solution.

Remark 1. Following the lines of the discussion in Subsection 15.3-2 we can treat the case in which the kernel $K(t, \tau)$ is meromorphic with respect to τ as well. In this case, Eq. (1) can be reduced to a Riemann problem of the type (12) and a linear algebraic system.

Remark 2. The solutions of a complete singular integral equation that are constructed in Section 15.3 can be applied for the case in which the contour L is a collection of finitely many disjoint smooth closed contours.

Example 1. Consider the equation

$$\lambda\varphi(t) + \frac{1}{\pi i} \int_L \frac{\cos(\tau - t)}{\tau - t} \varphi(\tau) d\tau = f(t), \tag{15}$$

where L is an arbitrary closed contour.

Note that the function $M(t, \tau) = \cos(\tau - t)$ has the property $M(t, t) \equiv 1$. Therefore, it remains to apply formula (6), and thus for (15) we have

$$\varphi(t) = \frac{1}{\lambda^2 - 1} \left[\lambda f(t) - \frac{1}{\pi i} \int_L \frac{\cos(\tau - t)}{\tau - t} f(\tau) d\tau \right], \quad \lambda \neq \pm 1.$$

Example 2. Consider the equation

$$\lambda\varphi(t) + \frac{1}{\pi i} \int_L \frac{\sin(\tau - t)}{(\tau - t)^2} \varphi(\tau) d\tau = f(t), \tag{16}$$

where L is an arbitrary closed contour.

The function $M(t, \tau) = \sin(\tau - t)/(\tau - t)$ has the property $M(t, t) \equiv 1$. Therefore, applying formula (6), for (16) we obtain

$$\varphi(t) = \frac{1}{\lambda^2 - 1} \left[\lambda f(t) - \frac{1}{\pi i} \int_L \frac{\sin(\tau - t)}{(\tau - t)^2} f(\tau) d\tau \right], \quad \lambda \neq \pm 1.$$

⊙ Reference for Section 15.3: F. D. Gakhov (1977, 1990).

15.4. Regularization Method for Complete Singular Integral Equations

15.4-1. Certain Properties of Singular Operators.

Let \mathbf{K}_1 and \mathbf{K}_2 be singular operators,

$$\mathbf{K}_1[\varphi(t)] \equiv a_1(t)\varphi(t) + \frac{1}{\pi i} \int_L \frac{M_1(t, \tau)}{\tau - t} \varphi(\tau) d\tau, \tag{1}$$

$$\mathbf{K}_2[\omega(t)] \equiv a_2(t)\omega(t) + \frac{1}{\pi i} \int_L \frac{M_2(t, \tau)}{\tau - t} \omega(\tau) d\tau. \tag{2}$$

The operator $\mathbf{K} = \mathbf{K}_2\mathbf{K}_1$ defined by the formula $\mathbf{K}[\varphi(t)] = \mathbf{K}_2[\mathbf{K}_1[\varphi(t)]]$ is called the *composition* or the *product* of the operators \mathbf{K}_1 and \mathbf{K}_2 .

Let us form the expression for the operator \mathbf{K} ,

$$\begin{aligned} \mathbf{K}[\varphi(t)] = \mathbf{K}_2\mathbf{K}_1[\varphi(t)] \equiv a_2(t) & \left[a_1(t)\varphi(t) + \frac{1}{\pi i} \int_L \frac{M_1(t, \tau)}{\tau - t} \varphi(\tau) d\tau \right] \\ & + \frac{1}{\pi i} \int_L \frac{M_2(t, \tau)}{\tau - t} \left[a_1(\tau)\varphi(\tau) + \frac{1}{\pi i} \int_L \frac{M_1(\tau, \tau_1)}{\tau_1 - \tau} \varphi(\tau_1) d\tau_1 \right] d\tau, \end{aligned} \quad (3)$$

and select its characteristic part. To this end, we perform the following manipulations:

$$\begin{aligned} \int_L \frac{M_1(t, \tau)}{\tau - t} \varphi(\tau) d\tau &= M_1(t, t) \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L \frac{M_1(t, \tau) - M_1(t, t)}{\tau - t} \varphi(\tau) d\tau, \\ \int_L \frac{a_1(\tau)M_2(t, \tau)}{\tau - t} \varphi(\tau) d\tau &= a_1(t)M_2(t, t) \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L \frac{a_1(\tau)M_2(t, \tau) - a_1(t)M_2(t, t)}{\tau - t} \varphi(\tau) d\tau, \\ \int_L \frac{M_2(t, \tau)}{\tau - t} d\tau \int_L \frac{M_1(\tau, \tau_1)}{\tau_1 - \tau} \varphi(\tau_1) d\tau_1 &= -\pi^2 M_2(t, t)M_1(t, t)\varphi(t) + \int_L \varphi(\tau_1) d\tau_1 \int_L \frac{M_2(t, \tau)M_1(\tau, \tau_1)}{(\tau_1 - \tau)(\tau - t)} d\tau. \end{aligned} \quad (4)$$

Here we applied the Poincaré–Bertrand formula (see Subsection 14.2-6). We can see that all kernels of the integrals of the last summands on the right-hand sides in (4) are Fredholm kernels.

We write

$$M_1(t, t) = b_1(t), \quad M_2(t, t) = b_2(t) \quad (5)$$

and see that the characteristic operator \mathbf{K}° of the composition (product) \mathbf{K} of two singular operators \mathbf{K}_1 and \mathbf{K}_2 can be expressed by the formula

$$\mathbf{K}^\circ[\varphi(t)] = (\mathbf{K}_2\mathbf{K}_1)^\circ[\varphi(t)] = [a_2(t)a_1(t) + b_2(t)b_1(t)]\varphi(t) + \frac{a_2(t)b_1(t) + b_2(t)a_1(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau. \quad (6)$$

Let us write out the operator \mathbf{K}_1 and \mathbf{K}_2 in the form (3) with explicitly expressed characteristic parts:

$$\mathbf{K}_1[\varphi(t)] \equiv a_1(t)\varphi(t) + \frac{b_1(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L K_1(t, \tau)\varphi(\tau) d\tau, \quad (7)$$

$$\mathbf{K}_2[\omega(t)] \equiv a_2(t)\omega(t) + \frac{b_2(t)}{\pi i} \int_L \frac{\omega(\tau)}{\tau - t} d\tau + \int_L K_2(t, \tau)\omega(\tau) d\tau. \quad (8)$$

Thus, the coefficients $a(t)$ and $b(t)$ of the characteristic part of the product of the operators \mathbf{K}_1 and \mathbf{K}_2 can be expressed by the formulas

$$a(t) = a_2(t)a_1(t) + b_2(t)b_1(t), \quad b(t) = a_2(t)b_1(t) + b_2(t)a_1(t). \quad (9)$$

These formulas do not contain regular kernels k_1 and k_2 and are symmetric with respect to the indices 1 and 2. This means that the characteristic part of the product of singular operators depends neither on their regular parts nor on the order of these operators in the product.

Thus, any change of order of the factors, as well as a change of the regular parts of the factors, influences the regular part of the product of the operators only and preserves the characteristic part of the product.

Let us calculate the coefficient of the Riemann problem that corresponds to the characteristic operator $(\mathbf{K}_2\mathbf{K}_1)^\circ$:

$$D(t) = \frac{a(t) - b(t)}{a(t) + b(t)} = \frac{[a_2(t) - b_2(t)][a_1(t) - b_1(t)]}{[a_2(t) + b_2(t)][a_1(t) + b_1(t)]} = D_2(t)D_1(t), \quad (10)$$

where we denote by

$$D_1(t) = \frac{a_1(t) - b_1(t)}{a_1(t) + b_1(t)}, \quad D_2(t) = \frac{a_2(t) - b_2(t)}{a_2(t) + b_2(t)} \quad (11)$$

the coefficients of the Riemann problems that correspond to the operators \mathbf{K}_1° and \mathbf{K}_2° . This means that the coefficient of the Riemann problem for the operator $(\mathbf{K}_2\mathbf{K}_1)^\circ$ is equal to the product of the coefficients of the Riemann problems for the operators \mathbf{K}_1° and \mathbf{K}_2° , and hence the index of the product of singular operators is equal to the sum of indices of the factors:

$$\nu = \nu_1 + \nu_2. \tag{12}$$

In its complete form, the operator $\mathbf{K}_2\mathbf{K}_1$ is defined by the expression

$$\mathbf{K}_2\mathbf{K}_1[\varphi(t)] \equiv a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L K(t, \tau)\varphi(\tau) d\tau,$$

where $a(t)$ and $b(t)$ are defined by formulas (9). For a regular kernel $K(t, \tau)$, on the basis of formulas (4) we can write out the explicit expression.

For a singular operator \mathbf{K} and its transposed operator \mathbf{K}^* (see Subsection 15.1-1), the following relations hold:

$$\int_L \psi(t)\mathbf{K}[\varphi(t)] dt = \int_L \varphi\mathbf{K}^*[\psi(t)] dt$$

for any functions $\varphi(t)$ and $\psi(t)$ that satisfy the Hölder condition, and

$$(\mathbf{K}_2\mathbf{K}_1)^* = \mathbf{K}_1^*\mathbf{K}_2^*.$$

15.4-2. Regularizer.

The regularization method is a reduction of a singular integral equation to a Fredholm equation. The reduction process itself is known as *regularization*.

If a singular operator \mathbf{K}_2 is such that the operator $\mathbf{K}_2\mathbf{K}_1$ is regular (Fredholm), i.e., contains no singular integral ($b(t) \equiv 0$), then \mathbf{K}_2 is called a *regularizing operator* with respect to the singular operator \mathbf{K}_1 or, briefly, a *regularizer*. Note that if \mathbf{K}_2 is a regularizer, then the operator $\mathbf{K}_1\mathbf{K}_2$ is regular as well.

Let us find the general form of a regularizer. By definition, the following relation must hold:

$$b(t) = a_2(t)b_1(t) + b_2(t)a_1(t) = 0, \tag{13}$$

which implies that

$$a_2(t) = g(t)a_1(t), \quad b_2(t) = -g(t)b_1(t), \tag{14}$$

where $g(t)$ is an arbitrary function that vanishes nowhere and satisfies the Hölder condition.

Hence, if \mathbf{K} is a singular operator,

$$\mathbf{K}[\varphi(t)] \equiv a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L K(t, \tau)\varphi(\tau) d\tau, \tag{15}$$

then, in general, the regularizer $\tilde{\mathbf{K}}$ can be expressed as follows:

$$\tilde{\mathbf{K}}[\omega(t)] \equiv g(t)a(t)\omega(t) - \frac{g(t)b(t)}{\pi i} \int_L \frac{\omega(\tau)}{\tau - t} d\tau + \int_L \tilde{K}(t, \tau)\omega(\tau) d\tau, \tag{16}$$

where $\tilde{K}(t, \tau)$ is an arbitrary Fredholm kernel and $g(t)$ is an arbitrary function satisfying the Hölder condition.

Since the index of a regular operator ($b(t) \equiv 0$) is clearly equal to zero, it follows from the property of the product of operators that the index of the regularizer has the same modulus as the

index of the original operator and the opposite sign. The same fact can be established directly by the form of a regularizer (16) from the formula

$$\tilde{D}(t) = \frac{\tilde{a}(t) - \tilde{b}(t)}{\tilde{a}(t) + \tilde{b}(t)} = \frac{a(t) + b(t)}{a(t) - b(t)} = \frac{1}{D(t)}.$$

Thus, for any singular operator with Cauchy kernel (15) of the normal type ($a(t) \pm b(t) \neq 0$), there exist infinitely many regularizers (16) whose characteristic part depends on an arbitrary function $g(t)$ that contains an arbitrary regular kernel $\tilde{K}(t, \tau)$.

Since the elements $g(t)$ and $\tilde{K}(t, \tau)$ are arbitrary, we can choose them so that the regularizer will satisfy some additional conditions. For instance, we can make the coefficient of $\varphi(t)$ in the regularized equation be normalized, i.e., equal to one. To this end we must set $g(t) = [a^2(t) - b^2(t)]^{-1}$. If no conditions are imposed, then it is natural to apply the simplest regularizers. These can be obtained by setting $g(t) \equiv 1$ and $\tilde{K}(t, \tau) \equiv 0$ in formula (16), which gives the regularizer

$$\tilde{\mathbf{K}}[\omega(t)] = \mathbf{K}^{*\circ}[\omega(t)] \equiv a(t)\omega(t) - \frac{b(t)}{\pi i} \int_L \frac{\omega(\tau)}{\tau - t} d\tau, \quad (17)$$

or we can set $g(t) \equiv 1$ and $\tilde{K}(t, \tau) = -\frac{1}{\pi i} \frac{b(\tau) - b(t)}{\tau - t}$ and obtain

$$\tilde{\mathbf{K}}[\omega(t)] = \mathbf{K}^{\circ*}[\omega(t)] \equiv a(t)\omega(t) - \frac{1}{\pi i} \int_L \frac{b(\tau)\omega(\tau)}{\tau - t} d\tau. \quad (18)$$

The simplest operators $\mathbf{K}^{*\circ}$ and $\mathbf{K}^{\circ*}$ are most frequently used as regularizers.

Since the multiplication of operators is not generally commutative, one should distinguish two forms of regularization: left regularization, which gives the operator $\tilde{\mathbf{K}}\mathbf{K}$, and right regularization which leads to the operator $\mathbf{K}\tilde{\mathbf{K}}$. On the basis of the above remark we can claim that a right regularizer is simultaneously a left regularizer, and vice versa. Thus, the operation of regularization is commutative.

If an operator $\tilde{\mathbf{K}}$ is a regularizer for an operator \mathbf{K} , then, in turn, the operator \mathbf{K} is a regularizer for the operator $\tilde{\mathbf{K}}$. The operators $\mathbf{K}_1\mathbf{K}_2$ and $\mathbf{K}_2\mathbf{K}_1$ can differ by a regular part only.

15.4-3. Methods of Left and Right Regularization.

Let a complete singular integral equation be given:

$$\mathbf{K}[\varphi(t)] \equiv a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L K(t, \tau)\varphi(\tau) d\tau = f(t). \quad (19)$$

Three methods of regularization are used. The first two methods are based on the composition of a given singular operator and its regularizer (left and right regularization). The third method differs essentially from the first two, namely, the elimination of the singular integral is performed by solving the corresponding characteristic equation.

1°. *Left regularization.* Let us take the regularizer (16):

$$\tilde{\mathbf{K}}[\omega(t)] \equiv g(t)a(t)\omega(t) - \frac{g(t)b(t)}{\pi i} \int_L \frac{\omega(\tau)}{\tau - t} d\tau + \int_L \tilde{K}(t, \tau)\omega(\tau) d\tau. \quad (20)$$

On replacing the function $\omega(t)$ in $\tilde{\mathbf{K}}[\omega(t)]$ with the expression $\mathbf{K}[\varphi(t)] - f(t)$ we arrive at the integral equation

$$\tilde{\mathbf{K}}\mathbf{K}[\varphi(t)] = \tilde{\mathbf{K}}[f(t)]. \quad (21)$$

By definition, $\tilde{\mathbf{K}}\mathbf{K}$ is a Fredholm operator, because $\tilde{\mathbf{K}}$ is a regularizer. Hence, Eq. (21) is a Fredholm equation. Thus, we have transformed the singular integral equation (19) into the Fredholm integral equation (21) for the same unknown function $\varphi(t)$.

This is the first regularization method, which is called *left regularization*.

2°. *Right Regularization.* On replacing in Eq. (19) the desired function by the expression (20),

$$\varphi(t) = \tilde{\mathbf{K}}[\omega(t)], \quad (22)$$

where $\omega(t)$ is a new unknown function, we arrive at the integral equation

$$\mathbf{K}\tilde{\mathbf{K}}[\omega(t)] = f(t), \quad (23)$$

which is a Fredholm equation as well. Thus, from the singular integral equation (19) for the unknown function $\varphi(t)$ we passed to the Fredholm integral equation for the new unknown function $\omega(t)$.

On solving the Fredholm equation (23), we find a solution of the original equation (19) by formula (22). The application of formula (22) requires integration only (a proper integral and a singular integral must be found).

This is the second method of the regularization, which is called *right regularization*.

15.4-4. Problem of Equivalent Regularization.

In the reduction of a singular integral equation to a regular one we perform a functional transformation over the corresponding equation. In general, this transformation can either introduce new irrelevant solutions that do not satisfy the original equation or imply a loss of some solutions. Therefore, in general, the resultant equation is not equivalent to the original equation. Consider the relationship between the solutions of these equations and find out in what cases these equations are equivalent.

1°. *Left Regularization.* Consider a singular equation

$$\mathbf{K}[\varphi(t)] = f(t) \quad (24)$$

and the corresponding regular equation

$$\tilde{\mathbf{K}}\mathbf{K}[\varphi(t)] = \tilde{\mathbf{K}}[f(t)]. \quad (25)$$

Let us write out Eq. (25) in the form

$$\tilde{\mathbf{K}}[\mathbf{K}[\varphi(t)] - f(t)] = 0. \quad (26)$$

Since the operator $\tilde{\mathbf{K}}$ is homogeneous, it follows that each solution of the original equation (24) (a function that vanishes the expression $\mathbf{K}[\varphi(t)] - f(t)$) satisfies Eq. (26) as well. Hence, the left regularization implies no loss of solutions. However, a solution of the regularized equation need not be a solution of the original equation.

Consider the singular integral equation corresponding to the regularizer

$$\tilde{\mathbf{K}}[\omega(t)] = 0. \quad (27)$$

Let $\omega_1(t), \dots, \omega_p(t)$ be a complete system of its solutions, i.e., a maximal collection of linearly independent eigenfunctions of the regularizer $\tilde{\mathbf{K}}$.

We regard Eq. (26) as a singular equation of the form (27) with the unknown function $\omega(t) = \mathbf{K}[\varphi(t)] - f(t)$. We obtain

$$\mathbf{K}[\varphi(t)] - f(t) = \sum_{j=1}^p \alpha_j \omega_j(t), \quad (28)$$

where the α_j are some constants.

We see that the regularized equation is equivalent to Eq. (28) rather than the original equation (24).

Thus, Eq. (25) is equivalent to Eq. (28) in which α_j are arbitrary or definite constants. It may occur that Eq. (28) is solvable only under the assumption that all α_j satisfy the condition $\alpha_j = 0$. In this case, Eq. (25) is equivalent to the original equation (24), and the regularizer defines an equivalent transformation. In particular, if the regularizer has no eigenfunctions, then the right-hand side of Eq. (28) is identically zero, and it must be equivalent. This operator certainly exists for $\nu \geq 0$. For instance, we can take the regularizer $\mathbf{K}^{*\circ}$, which has no eigenfunctions for the case under consideration because the index of the regularizer $\mathbf{K}^{*\circ}$ is equal to $-\nu \leq 0$.

2°. *Right Regularization.* Consider Eq. (24) and the corresponding regularized equation

$$\mathbf{K}\tilde{\mathbf{K}}[\omega(t)] = f(t), \quad (29)$$

which is obtained by substitution

$$\tilde{\mathbf{K}}[\omega(t)] = \varphi(t). \quad (30)$$

If $\omega_j(t)$ is a solution of Eq. (29), then formula (30) gives the corresponding solution of the original equation

$$\varphi_j(t) = \tilde{\mathbf{K}}[\omega_j(t)].$$

Hence, the right regularization cannot lead to irrelevant solutions.

Conversely, assume that $\varphi_k(t)$ is a solution of the original equation. In this case a solution of the regularized equation (29) can be obtained as a solution of the nonhomogeneous singular equation

$$\tilde{\mathbf{K}}[\omega(t)] = \varphi_k(t);$$

however, this solution may be unsolvable. Thus, the right regularization can lead to loss of solutions. We have no loss of solutions if Eq. (30) is solvable for each right-hand side. In this case the operator $\tilde{\mathbf{K}}$ will be an equivalent right regularizer.

3°. *The Equivalent Regularization.* The operator $\tilde{\mathbf{K}} = \mathbf{K}^{*\circ}$ is an equivalent regularizer for any index; for $\nu \geq 0$, we must apply left regularization, while for $\nu \leq 0$ we must use right regularization.

In the latter case we obtain an equation for a new function $\omega(t)$, and if it is determined, then we can construct all solutions to the original equation in antiderivatives, and it follows from the properties of the right regularization that no irrelevant solutions can occur.

For the other methods of equivalent regularization, see the references at the end of this section.

15.4-5. Fredholm Theorems.

Let a complete singular integral equation be given:

$$\mathbf{K}[\varphi(t)] = f(t). \quad (31)$$

THEOREM 1. *The number of solutions of the singular integral equation (31) is finite.*

THEOREM 2. *A necessary and sufficient solvability condition for the singular equation (31) is*

$$\int_L f(t)\psi_j(t) dt = 0, \quad j = 1, \dots, m, \quad (32)$$

where $\psi_1(t), \dots, \psi_m(t)$ is a maximal finite set of linearly independent solutions of the transposed homogeneous equation $\mathbf{K}^*[\psi(t)] = 0$. (Since the functions under consideration are complex, it follows that condition (32) is not the orthogonality condition for the functions $f(t)$ and $\psi_j(t)$.)

THEOREM 3. *The difference between the number n of linearly independent solutions of the singular equation $\mathbf{K}[\varphi(t)] = 0$ and the number m of linearly independent solutions of the transposed equation $\mathbf{K}^*[\psi(t)] = 0$ depends on the characteristic part of the operator \mathbf{K} only and is equal to its index, i.e.,*

$$n - m = \nu. \quad (33)$$

Corollary. The number of linearly independent solutions of characteristic equations is minimal among all singular equations with given index ν .

15.4-6. Carleman–Vekua Approach to the Regularization.

Let us transfer the regular part of a singular equation to the right-hand side and rewrite the equation as follows:

$$a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau = f(t) - \int_L K(t, \tau)\varphi(\tau) d\tau, \quad (34)$$

or, in the operator form,

$$\mathbf{K}^\circ[\varphi(t)] = f(t) - \mathbf{K}_r[\varphi(t)]. \quad (35)$$

We regard the last equation as a characteristic one and solve it by temporarily assuming that the right-hand side is a known function. In this case (see Subsection 15.2-1)

$$\begin{aligned} \varphi(t) = & \left[a(t)f(t) - \frac{b(t)Z(t)}{\pi i} \int_L \frac{f(\tau)}{Z(\tau)} \frac{d\tau}{\tau - t} + b(t)Z(t)P_{\nu-1}(t) \right] \\ & - \left[a(t) \int_L K(t, \tau)\varphi(\tau) d\tau - \frac{b(t)Z(t)}{\pi i} \int_L \frac{d\tau_1}{Z(\tau_1)(\tau_1 - t)} \int_L K(\tau_1, \tau)\varphi(\tau) d\tau \right], \end{aligned} \quad (36)$$

where for $\nu \leq 0$ we must set $P_{\nu-1}(t) \equiv 0$. Let us reverse the order of integration in the iterated integral and rewrite the expression in the last parentheses as follows:

$$\int_L \left[a(t)K(t, \tau) - \frac{b(t)Z(t)}{\pi i} \int_L \frac{K(\tau_1, \tau)}{Z(\tau_1)(\tau_1 - t)} d\tau_1 \right] \varphi(\tau) d\tau.$$

Since $Z(t)$ satisfies the Hölder condition (and hence is bounded) and does not vanish and since $K(\tau_1, \tau)$ satisfies the estimate $|K(\tau_1, \tau)| < A|\tau_1 - \tau|^{-\lambda}$ (with $0 \leq \lambda < 1$) near the point $\tau_1 = \tau$, we can see that the entire integral

$$\int_L \frac{K(\tau_1, \tau)}{Z(\tau_1)(\tau_1 - t)} d\tau_1$$

satisfies an estimate similar to that for $K(\tau_1, \tau)$. Hence, the kernel

$$N(t, \tau) = a(t)K(t, \tau) - \frac{b(t)Z(t)}{\pi i} \int_L \frac{K(\tau_1, \tau)}{Z(\tau_1)(\tau_1 - t)} d\tau_1 \quad (37)$$

is a Fredholm kernel. On transferring the terms with $\varphi(t)$ to the right-hand side, we obtain

$$\varphi(t) + \int_L N(t, \tau)\varphi(\tau) d\tau = f_1(t), \quad (38)$$

where $N(t, \tau)$ is the Fredholm kernel defined by formula (37) and $f_1(t)$ has the form

$$f_1(t) = a(t)f(t) - \frac{b(t)Z(t)}{\pi i} \int_L \frac{f(\tau)}{Z(\tau)} \frac{d\tau}{\tau - t} + b(t)Z(t)P_{\nu-1}(t). \quad (39)$$

If the index of Eq. (34) ν is negative, then the function must satisfy not only the Fredholm equation (38) but also the relations

$$\int_L \left[\int_L \frac{K(t, \tau)}{Z(t)} t^{k-1} dt \right] \varphi(\tau) d\tau = \int_L \frac{f(t)}{Z(t)} t^{k-1} dt, \quad k = 1, 2, \dots, -\nu. \quad (40)$$

Thus, if $\nu \geq 0$, then the solution of a complete singular integral equation (34) is reduced to the solution of the Fredholm integral equation (38). If $\nu < 0$, then Eq. (34) can be reduced to Eq. (38) (where we must set $P_{\nu-1}(t) \equiv 0$) together with conditions (40), which can be rewritten in the form

$$\begin{aligned} \int_L \rho_k(\tau)\varphi(\tau) d\tau &= f_k, \quad k = 1, 2, \dots, -\nu, \\ \rho_k(\tau) &= \int_L \frac{K(t, \tau)}{Z(t)} t^{k-1} dt, \quad f_k = \int_L \frac{f(t)}{Z(t)} t^{k-1} dt, \end{aligned} \quad (41)$$

where the $\rho_k(\tau)$ are known functions and the f_k are known constants.

Relations (41) are the solvability conditions for the regularized equation (38). However, they need not be the solvability conditions for the original singular integral equation (34). Some of them can be the equivalence conditions for these two equations. Let us select the conditions of these two types.

Assume that among the functions $\rho_k(t)$ there are precisely h linearly independent functions. We can choose the numbering so that these are the functions $\rho_1(t), \dots, \rho_h(t)$. In this case we have

$$\int_L \rho_k(t)\varphi(t) dt = f_k, \quad k = 1, 2, \dots, h. \tag{42}$$

Moreover, the following $\eta = |\nu| - h$ linearly independent relations must hold:

$$\alpha_{j1}\rho_1(t) + \dots + \alpha_{j|\nu|}\rho_{|\nu|}(t) = 0, \quad j = 1, 2, \dots, \eta.$$

Let us multiply the relations in (40) successively by $\alpha_{j1}, \dots, \alpha_{j|\nu|}$ and sum the products. Taking into account the last relations, we have

$$\int_L f(t)\psi_j(t) dt = 0, \quad \psi_j(t) = \frac{1}{Z(t)} \sum_{k=1}^{|\nu|} \alpha_{jk}t^{k-1}; \quad j = 1, 2, \dots, \eta. \tag{43}$$

These relations, which do not involve the desired function $\varphi(t)$, are the necessary solvability conditions on the right-hand side $f(t)$ for the original singular equation and the regularized equation to be solvable. Relations (42) are the equivalence conditions for the original singular equation and the regularized equation. The solution of the Fredholm equation (38) satisfies the original singular equation (34) if and only if it satisfies conditions (42).

Thus, for $\nu \geq 0$, the regularized equation (38) is equivalent to the original singular equation. For $\nu < 0$, the original equation is equivalent to the regularized equation (with common solvability conditions (43)) together with conditions (42).

Remark 1. If the kernel of the regular part of a complete singular integral equation with Cauchy kernel is degenerate, then by the Carleman–Vekua regularization this equation can be reduced to the investigation of a system of linear algebraic equations (see, e.g., S. G. Mikhlin and K. L. Smolitskiy (1967)).

Remark 2. The Carleman–Vekua regularization is sometimes called the regularization by solving the characteristic equation.

15.4-7. Regularization in Exceptional Cases.

Consider the complete singular equation with Cauchy kernel

$$\mathbf{K}[\varphi(t)] \equiv a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau + \int_L K(t, \tau)\varphi(\tau) d\tau = f(t) \tag{44}$$

under the same conditions on the functions $a(t) \pm b(t)$ as above in Subsection 15.2-4.

We represent this equation in the form

$$\mathbf{K}^\circ[\varphi(t)] = f(t) - \int_L K(t, \tau)\varphi(\tau) d\tau,$$

and apply the Carleman–Vekua regularization. In this case by formula (42) of Subsection 15.2-4 we obtain the equation

$$\varphi(t) + \mathbf{R}_1 \left[\int_L K(t, \tau)\varphi(\tau) d\tau \right] = \mathbf{R}_1[f(t)] + b(t)Z(t)P_{\nu-p-1}(t), \tag{45}$$

where the operator \mathbf{R}_1 is defined by formula (41) of Subsection 15.2-4.

In the expression for the second summand on the left-hand side in (45), the operation \mathbf{R}_1 with respect to the variable t commutes with the operation of integration with respect to τ . Therefore, Eq. (45) can be rewritten in the form

$$\varphi(t) + \int_L \mathbf{R}_1^t [K(t, \tau)] \varphi(\tau) d\tau = \mathbf{R}_1[f(t)] + b(t)Z(t)P_{\nu-p-1}(t), \quad (46)$$

where the superscript t at the symbol of the operator \mathbf{R}_1^t means that the operation is performed with respect to the variable t .

Since the operator \mathbf{R}_1 is bounded, it follows that the resulting integral equation (46) is a Fredholm equation, and hence the regularization problem for the singular equation (44) is solved.

It follows from the general theory of the regularization that Eq. (44) is equivalent to Eq. (46) for $\nu - p \geq 0$ and to Eq. (46) and a system of functional equations for $\nu - p < 0$.

In conclusion we note that for the above cases of singular integral equations, the Fredholm theorems fail in general.

Remark 3. Exceptional cases of singular integral equations with Cauchy kernel can be reduced to equations of the normal type.

15.4-8. Complete Equation with Hilbert Kernel.

Consider the complete singular integral equation with Hilbert kernel (see Subsection 15.1-2)

$$a(x)\varphi(x) - \frac{b(x)}{2\pi} \int_0^{2\pi} \cot\left(\frac{\xi - x}{2}\right) \varphi(\xi) d\xi + \int_0^{2\pi} K(x, \xi) \varphi(\xi) d\xi = f(x). \quad (47)$$

Let us show that Eq. (47) can be reduced to a complete singular integral equation with a kernel of the Cauchy type, and in this connection, the theory of the latter equation can be directly extended to Eq. (47). Since the regular parts of these two types of equations have the same character, it follows that it suffices to apply the relationship between the Hilbert kernel and the Cauchy kernel (see Subsection 14.4-5):

$$\frac{d\tau}{\tau - t} = \frac{1}{2} \cot\left(\frac{\xi - x}{2}\right) d\xi + \frac{i}{2} d\xi. \quad (48)$$

Hence,

$$\frac{1}{2} \cot\left(\frac{\xi - x}{2}\right) d\xi = \frac{d\tau}{\tau - t} - \frac{1}{2} \frac{d\tau}{\tau}, \quad (49)$$

where $t = e^{ix}$ and $\tau = e^{i\xi}$ are the complex coordinates of points of the contour L , that is, the unit circle.

On replacing the Hilbert kernel in Eq. (47) with the expression (49) and on substituting $x = -i \ln t$, $\xi = -i \ln \tau$, and $d\xi = -i\tau^{-1} d\tau$, after obvious manipulations we reduce Eq. (47) to a complete singular integral equation with Cauchy kernel of the form

$$a_1(t)\varphi_1(t) - \frac{ib_1(t)}{\pi i} \int_L \frac{\varphi_1(\tau)}{\tau - t} d\tau + \int_L K_1(t, \tau) d\tau = f_1(t). \quad (50)$$

The coefficient of the Riemann problem corresponding to Eq. (50) is

$$D(t) = \frac{a_1(t) + ib_1(t)}{a_1(t) - ib_1(t)} = \frac{a(x) + ib(x)}{a(x) - ib(x)}, \quad (51)$$

and the index is expressed by the formula

$$\text{Ind } D(t) = 2 \text{Ind}[a(x) + ib(x)]. \quad (52)$$

Example. Let us perform the regularization of the following singular integral equations in different ways:

$$\mathbf{K}[\varphi(t)] \equiv (t + t^{-1})\varphi(t) + \frac{t - t^{-1}}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau - \frac{1}{2\pi i} \int_L (t + t^{-1})(\tau + \tau^{-1})\varphi(\tau) d\tau = 2t^2, \quad (53)$$

where L is the unit circle.

The regular part of the kernel is degenerate. Therefore, in the same way as was applied in the solution of Fredholm equations with degenerate kernel (see Section 13.2), the equation can be reduced to the investigation of the characteristic equation and a linear algebraic equation, and hence it can be solved in a closed form. Thus, we need no regularization. However, the equation under consideration is useful in the illustration of general methods because all calculations can be performed to the very end.

For convenience of the subsequent discussion, we first solve this equation. We write

$$\frac{1}{2\pi i} \int_L (\tau + \tau^{-1})\varphi(\tau) d\tau = A, \quad (54)$$

and write out the equation in the characteristic form:

$$(t + t^{-1})\varphi(t) + \frac{t - t^{-1}}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau = 2t^2 + A(t + t^{-1}).$$

For the corresponding Riemann boundary value problem

$$\Phi^+(t) = t^{-2}\Phi^-(t) + t + \frac{1}{2}A(1 + t^{-2}), \quad (55)$$

we have the index $\nu = -2$, and the solvability conditions (see Subsection 15.2-1) hold for $A = 0$ only. In this case, $\Phi^+(z) = z$ and $\Phi^-(z) = 0$. This gives a solution to Eq. (53) in the form $\varphi(t) = \Phi^+(t) - \Phi^-(t) = t$. On substituting the last expression into Eq. (54) we see that this relation holds for $A = 0$. Hence, the given equation is solvable and has a unique solution of the form

$$\varphi(t) = t.$$

1°. *Left Regularization.* Since the equation index $\nu = -2$ is negative, any regularizer of the equation has eigenfunctions (at least two linearly independent), and hence the left regularization leads, in general, to an equation that is not equivalent to the original one.

We first consider the left regularization by means of the simplest regularizer $\mathbf{K}^{*\circ}$. Let us find the linearly independent eigenfunctions of the equation

$$\mathbf{K}^{*\circ}[\omega(t)] \equiv (t + t^{-1})\omega(t) - \frac{t - t^{-1}}{\pi i} \int_L \frac{\omega(\tau)}{\tau - t} d\tau = 0.$$

The corresponding Riemann boundary value problem

$$\Phi^+(t) = t^2\Phi^-(t)$$

now has the index $\nu = 2$. We can find the eigenfunctions of the operator $\mathbf{K}^{*\circ}$ by the formulas of Subsection 15.2-1 and obtain

$$\omega_1(t) = 1 - t^{-2}, \quad \omega_2(t) = t - t^{-1}.$$

On the basis of the general theory (see Subsection 15.4-4), the regular equation $\mathbf{K}^{*\circ}\mathbf{K}[\varphi(t)] = \mathbf{K}^{*\circ}[f(t)]$ is equivalent to the singular equation:

$$\mathbf{K}[\varphi(t)] = f(t) + \alpha_1\omega_1(t) + \alpha_2\omega_2(t), \quad (56)$$

where α_1 and α_2 are constants that can be either arbitrary or definite. Taking into account Eq. (54), we write out Eq. (56) in the form of a characteristic equation:

$$(t + t^{-1})\varphi(t) + \frac{t - t^{-1}}{\pi i} \int_L \frac{\varphi(\tau)}{\tau - t} d\tau = 2t^2 + A(t + t^{-1}) + \alpha_1(1 - t^{-2}) + \alpha_2(t - t^{-1}).$$

The corresponding Riemann boundary value problem has the form

$$\Phi^+(t) = t^{-2}\Phi^-(t) + t + \frac{1}{2}A(1 + t^{-2}) + \frac{1}{2}\alpha_1(t^{-1} + t^{-3}) + \frac{1}{2}\alpha_2(1 - t^{-2}).$$

Its solution can be represented as follows:

$$\Phi^+(z) = z + \frac{1}{2}A + \frac{1}{2}\alpha_2, \quad \Phi^-(z) = \frac{1}{2}z^2[\alpha_1 z^{-3} + (\alpha_2 - A)z^{-2} - \alpha_1 z^{-1}].$$

The solvability conditions give $\alpha_1 = 0$ and $\alpha_2 = A$. In this case, the solution of Eq. (56) is defined by the formula

$$\varphi(t) = \Phi^+(t) - \Phi^-(t) = t + A.$$

On substituting the above expression for $\varphi(t)$ into Eq. (54) we obtain the identity $A = A$. Hence, the constant $\alpha_2 = A$ remains arbitrary, and the regularized equation is equivalent not to the original equation but to the equation

$$\mathbf{K}[\varphi(t)] = f(t) + \alpha_2\omega_2(t),$$

which has the solution $\varphi(t) = t + A$, where A is an arbitrary constant. The last function φ satisfies the original equation only for $A = 0$.

2°. *Right Regularization.* For a right regularizer we take the simplest operator $\mathbf{K}^{*\circ}$. By setting

$$\varphi(t) = \mathbf{K}^{*\circ}[\omega(t)] \equiv (t + t^{-1})\omega(t) - \frac{t - t^{-1}}{\pi i} \int_L \frac{\omega(\tau)}{\tau - t} d\tau, \tag{57}$$

we obtain the following Fredholm equation with respect to the function $\omega(t)$:

$$\mathbf{K}\mathbf{K}^{*\circ}[\omega(t)] \equiv \omega(t) - \frac{1}{4\pi i} \int_L [t(\tau^2 - 1 + \tau^{-2}) + 2\tau^{-1} + t^{-1}(\tau^2 + 3 + \tau^{-2}) - 2\tau^{-2}\tau^{-1}]\omega(\tau) d\tau = \frac{1}{2}t^2. \tag{58}$$

The last equation is degenerate. On solving it we obtain

$$\omega(t) = \frac{1}{2}t^2 + \alpha(t - t^{-1}) + \beta(1 - t^{-2}),$$

where α and β are arbitrary constants.

Thus, the regularized equation for $\omega(t)$ has two linearly independent solutions, while the original equation (53) has a unique solution. On substituting the above expression for $\omega(t)$ into formula (57) we obtain

$$\varphi(t) = \mathbf{K}^{*\circ} \left[\frac{1}{2}t^2 + \alpha(t - t^{-1}) + \beta(1 - t^{-2}) \right] = t,$$

where $\varphi(t)$ is the (unique) solution of the original singular equation. The result agrees with the general theory because, for a negative index, the right regularization by means of the operator $\mathbf{K}^{*\circ}$ is an equivalent regularization.

3°. *The Carleman–Vekua Regularization.* This method of regularization is performed by formulas (36)–(39). However, we must recall that these formulas can be applied only for an equation such that $a^2(t) - b^2(t) = 1$. Therefore, we must first divide Eq. (53) by two. In this case, we have

$$\begin{aligned} a &= \frac{1}{2}(t + t^{-1}), \quad b = \frac{1}{2}(t - t^{-1}), \quad f(t) = t^2, \\ K(t, \tau) &= -\frac{1}{4\pi i}(t + t^{-1})(\tau + \tau^{-1}), \quad X^+(z) = 1, \quad Z(t) = (a + b)X^+ = t, \\ f_1(t) &= \frac{1}{2}(t + t^{-1})t^2 - \frac{(t - t^{-1})t}{2\pi i} \int_L \frac{\tau^2}{\tau} \frac{d\tau}{\tau - t} = t, \\ N(t, \tau) &= -\frac{1}{2}(t + t^{-1})\frac{1}{4\pi i}(t + t^{-1})(\tau + \tau^{-1}) + \frac{(t - t^{-1})t(\tau + \tau^{-1})}{2\pi i \cdot 4\pi i} \int_L \frac{\tau_1 + \tau_1^{-1}}{\tau_1} \frac{d\tau_1}{\tau_1 - t} = -\frac{1}{2\pi i}(\tau + \tau^{-1}). \end{aligned}$$

The regularized equation has the form

$$\varphi(t) - \frac{1}{2\pi i} \int_L (\tau + \tau^{-1})\varphi(\tau) d\tau = t. \tag{59}$$

To this equation we must add conditions (41) for $k = 1, 2$. This equation is degenerate, and on solving it we find the general solution $\varphi(t) = t + A$, where A is an arbitrary constant. Let us write out conditions (42) and (43). Here we have

$$\begin{aligned} \rho_k(\tau) &= \int_L \frac{K(t, \tau)}{Z(t)} t^{k-1} dt = -\frac{\tau + \tau^{-1}}{4\pi i} \int_L (1 + t^{-2})t^{k-1} dt, \quad k = 1, 2, \\ \rho_1(\tau) &= 0, \quad \rho_2(\tau) = -\frac{1}{2}(\tau + \tau^{-1}), \quad f_k = \int_L \frac{f(t)}{Z(t)} t^{k-1} dt = \int_L t^k dt, \quad f_1 = f_2 = 0. \end{aligned}$$

The functions $\rho_1(t)$ and $\rho_2(t)$ are linearly dependent. The dependence $\alpha_{j1}\rho_1(t) + \dots + \alpha_{j|\nu|}\rho_{|\nu|}(t) = 0$ (see Subsection 15.4-6) has the form

$$\alpha_1 \rho_1(t) + 0 \cdot \rho_2(t) = 0.$$

Hence, the solvability condition (43) holds identically. The equivalence condition (42)

$$\int_L \rho_2(\tau)\varphi(\tau) d\tau = -\frac{1}{2} \int_L (\tau + \tau^{-1})(\tau + A) d\tau = 0$$

holds for $A = 0$ only. Hence, among the solutions to the regularized equation, $\varphi(t) = t + A$, only the function $\varphi(t) = t$ satisfies the original equation.

⊙ References for Section 15.4: F. D. Gakhov (1977, 1990), S. G. Mikhlin and S. Prössdorf (1986), N. I. Muskhelishvili (1992).

15.5. Analysis of Solutions Singularities for Complete Integral Equations with Generalized Cauchy Kernels*

15.5-1. Statement of the Problem and Preliminary Remarks.

Consider a complete integral equation of the second kind in the form

$$a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_{-1}^1 \frac{\varphi(\tau) d\tau}{\tau - t} + \int_{-1}^1 K_g(t, \tau)\varphi(\tau) d\tau + \int_{-1}^1 L_g(t, \tau)\overline{\varphi(\tau)} d\tau + \int_{-1}^1 K(t, \tau)\varphi(\tau) d\tau + \int_{-1}^1 L(t, \tau)\overline{\varphi(\tau)} d\tau = f(t), \quad -1 < t < 1, \quad (1)$$

where $\varphi(\tau)$ is an unknown function and $\overline{\varphi(\tau)}$ is its complex conjugate; $f(t)$ is a given continuous function on the closed interval $[-1, 1]$; the functions $a(t)$, $b(t)$ and the kernels $K(t, \tau)$, $L(t, \tau)$ are bounded and continuous (or satisfy the Hölder condition in all their arguments), and the *generalized kernels* $K_g(t, \tau)$, $L_g(t, \tau)$ have fixed singularities that are first-order poles at the endpoints of the integration interval, as the (real) parameters τ and t simultaneously tend to either endpoint of the interval $[-1, 1]$ ($\tau = t \rightarrow \pm 1$). The kernels of equation (1), as well as the functions $a(t)$, $b(t)$, $f(t)$, may be either real- or complex-valued. Note that the method described below is suitable for the asymptotic analysis of equations of the second as well as first kind for $a(t) \equiv 0$.

Assume that the solution of equation (1) belongs to the class of functions that have, at the endpoints of the integration interval, integrable singularities (generally complex) of power type due to both a “movable” singularity of the integral in the sense of the principal value (the first integral in (1)) and fixed singularities of the kernels $K_g(t, \tau)$ and $L_g(t, \tau)$. This type of asymptotic behavior of the unknown function can be taken into account by the introduction of a special weight function

$$w(\tau) = (1 - \tau)^\alpha(1 + \tau)^\beta, \quad -1 \leq \tau \leq 1, \quad -1 < \operatorname{Re} \alpha, \operatorname{Re} \beta < 0, \quad (2)$$

which is present as a coefficient in the unknown function, i.e.,

$$\varphi(\tau) = u(\tau)w(\tau). \quad (3)$$

Here, $u(\tau)$ is a new unknown function satisfying the Hölder condition and different from zero at the endpoints of the interval. The last requirement is connected with the fact that the sought weight function (2) should reflect the leading singular asymptotics of the unknown function (3). Note that the presence of fixed singularities in the kernels $K_g(t, \tau)$ and $L_g(t, \tau)$ significantly effects the asymptotic behavior of the solution near the endpoints of the integration interval, which in this situation usually has the form $(1 \mp \tau)^\lambda$, $-1 < \operatorname{Re} \lambda < 0$, $\tau \rightarrow \pm 1$ ($\lambda = \alpha, \beta$), and $\operatorname{Re} \lambda \neq -1/2$. (Sometimes the weight function may be bounded on one end of the integration interval, which corresponds to $\operatorname{Re} \lambda \geq 0$.)

Assume that the generalized kernels can be represented in the form

$$K_g(t, \tau) = \sum_{p,j,k,r} A_k(t) \frac{(1 + \tau)^p(1 + t)^j}{(\tau - z_r^*)^{p+j+1}} + \sum_{l,m,n,s} B_l(t) \frac{(1 - \tau)^m(1 - t)^n}{(\tau - z_s^{**})^{m+n+1}}, \quad (4)$$

$$L_g(t, \tau) = \sum_{p,j,k,r} C_k(t) \frac{(1 + \tau)^p(1 + t)^j}{(\tau - z_r^*)^{p+j+1}} + \sum_{l,m,n,s} D_l(t) \frac{(1 - \tau)^m(1 - t)^n}{(\tau - z_s^{**})^{m+n+1}}. \quad (5)$$

* Section 15.5 was written by A. V. Andreev.

Here, the indices and the power exponents p, j, k, l, m, n, r, s take the values $0, 1, 2, \dots$ and vary independently in each sum; the quantities z_r^* and z_s^{**} depend on the variable t as follows ($i^2 = -1$):

$$\begin{aligned} z_r^*(t) &= -1 + (1+t)e^{i\theta_r}, & 0 < \theta_r < 2\pi; \\ z_s^{**}(t) &= 1 + (1-t)e^{i\theta_s}, & -\pi < \theta_s < \pi. \end{aligned} \tag{6}$$

It is also assumed that the functions $A_k(t), C_k(t)$ and $B_l(t), D_l(t)$ in (4)–(5) are finite and nonzero at $t = -1$ and $t = +1$.

The geometrical meaning of expressions (6) is that for $t \rightarrow -1$ (resp., $t \rightarrow +1$) the points z_r^* (resp., z_s^{**}), on the complex plane, tend to -1 (resp., $+1$) along the ray obtained from the integration line by its rotation by the angle θ_r (resp., θ_s) about the point -1 (resp., $+1$).

Remark. Note that in mathematical statements of some applied problems, the representations of $z_r^*(t)$ and $z_s^{**}(t)$ in the form (6) may involve $\theta_r < 0$ or $\theta_r > 2\pi$. In view of the inequality $0 < \theta_r < 2\pi$ indicating that the values of the angle θ_r in (6) are counted counterclockwise from the positive direction of the axis Ox , one should replace θ_r in each such case by a suitable equivalent value obtained by that rule. Similarly, if $\theta_s > \pi$ or $\theta_s < -\pi$, this value of θ_s should be replaced by a suitable acute angle. Such replacements ensure fixed signs in the formulas for $z_r^*(t)$ and $z_s^{**}(t)$, which appear in the generalized kernels (4) and (5).

15.5-2. Auxiliary Results.

Using (2) and (3), let us rewrite equation (1) in the equivalent form

$$\begin{aligned} u(t) &\left[a(t)w(t) + \frac{b(t)}{\pi i} \int_{-1}^1 \frac{w(\tau) d\tau}{\tau - t} + \int_{-1}^1 K_g(t, \tau)w(\tau) d\tau \right] + \overline{u(t)} \int_{-1}^1 L_g(t, \tau)\overline{w(\tau)} d\tau \\ &+ \int_{-1}^1 \left[\frac{b(t)}{\pi i} \frac{1}{\tau - t} + K_g(t, \tau) \right] \left[u(\tau) - u(t) \right] w(\tau) d\tau + \int_{-1}^1 L_g(t, \tau)[\overline{u(\tau)} - \overline{u(t)}]\overline{w(\tau)} d\tau \\ &+ \int_{-1}^1 K(t, \tau)u(\tau)w(\tau) d\tau + \int_{-1}^1 L(t, \tau)\overline{u(\tau)w(\tau)} d\tau = f(t), \quad -1 < t < 1. \end{aligned} \tag{7}$$

It is easy to see from (7) that for Hölder continuous $u(\tau)$, the characteristic part of equation (1), which goes to infinity as $t \rightarrow \pm 1$, has the form

$$I_s(t) = u(t) \left[a(t)w(t) + \frac{b(t)}{\pi i} \int_{-1}^1 \frac{w(\tau) d\tau}{\tau - t} + \int_{-1}^1 K_g(t, \tau)w(\tau) d\tau \right] + \overline{u(t)} \int_{-1}^1 L_g(t, \tau)\overline{w(\tau)} d\tau. \tag{8}$$

The most general approach to solving an integral equation with conjugate unknown functions consists in regarding this equation as a system of equations for two unknown functions $\varphi(\tau)$ and $\overline{\varphi(\tau)}$, where the second equation of the system is obtained by passing from (1) to conjugate values. Let us write out the characteristic part of the equation conjugate to (1) (to be used in the sequel):

$$\overline{I_s(t)} = u(t) \int_{-1}^1 \overline{L_g(t, \tau)w(\tau)} d\tau + \overline{u(t)} \left[\overline{a(t)w(t)} - \frac{\overline{b(t)}}{\pi i} \int_{-1}^1 \frac{\overline{w(\tau)} d\tau}{\tau - t} + \int_{-1}^1 \overline{K_g(t, \tau)w(\tau)} d\tau \right]. \tag{9}$$

Let us examine the asymptotic behavior of the characteristic part (8) and its conjugate (9) as $t \rightarrow \pm 1$ ($z_r^* \rightarrow -1, z_s^{**} \rightarrow +1$). To that end, we obtain expressions for the leading terms of the integrals in the sense of the principal value and the integrals containing generalized kernels (4), (5) as $t \rightarrow \pm 1$.

In order to calculate the integrals in (8), we use the integral representation of the zero-order Jacobi function of the second kind $Q_0^{(\alpha, \beta)}(z)$:

$$Q_0^{(\alpha, \beta)}(z) = \frac{1}{w(z)} \int_{-1}^1 \frac{w(\tau)}{\tau - z} d\tau, \quad z \notin [-1, 1]. \tag{10}$$

Using the representations of the Jacobi functions of the second kind in terms of the hypergeometric function $F(a, b, c, \vartheta)$, functional relations for this function, and the formula for the gamma function $\Gamma(\zeta)$, we obtain from (10) the following formal expressions of the Cauchy integral:

$$\begin{aligned} \int_{-1}^1 \frac{w(\tau)}{\tau - z} d\tau &= -\frac{\pi w(z)}{(-1)^\beta \sin(\pi\beta)} + 2^{\alpha+\beta} \frac{\Gamma(\alpha + 1)\Gamma(\beta)}{\Gamma(\alpha + \beta + 1)} F\left(1, -\alpha - \beta, 1 - \beta, \frac{1+z}{2}\right) \\ &= \frac{\pi w(z)}{(-1)^\alpha \sin(\pi\alpha)} - 2^{\alpha+\beta} \frac{\Gamma(\alpha)\Gamma(\beta + 1)}{\Gamma(\alpha + \beta + 1)} F\left(1, -\alpha - \beta, 1 - \alpha, \frac{1-z}{2}\right), \end{aligned} \tag{11}$$

where $z \notin [-1, 1]$. Since $F(a, b, c, 0) = 1$, the singularities of the Cauchy integral for $z \rightarrow \pm 1$ are completely determined by the first terms in the expressions (11).

We fix the multivaluedness of $(-1)^\lambda$ ($\lambda = \alpha, \beta$) in (11) so that the resulting expressions correctly reflect the behavior of the Cauchy integral as $z \rightarrow \pm 1$ from the complex plane cut along the segment $[-1, 1]$. The leading terms of the asymptotic expansion of the Cauchy integral near the endpoints of the integration interval are obtained from (11):

$$\left\{ \int_{-1}^1 \frac{w(\tau)}{\tau - z} d\tau \right\}_{z \rightarrow -1} = -\frac{2^\alpha \pi e^{-i\pi\beta}}{\sin(\pi\beta)} \{(1+z)^\beta\}_{z \rightarrow -1}, \quad z \notin [-1, 1]; \tag{12}$$

$$\left\{ \int_{-1}^1 \frac{w(\tau)}{\tau - z} d\tau \right\}_{z \rightarrow +1} = \frac{2^\beta \pi e^{i\pi\alpha}}{\sin(\pi\alpha)} \{(1-z)^\alpha\}_{z \rightarrow +1}, \quad z \notin [-1, 1]. \tag{13}$$

Here and in subsequent asymptotic formulas, we use the notation $\{F(x)\}_{x \rightarrow a} = F(x)|_{x \rightarrow a}$, and only the leading term of the expansion is kept in the right-hand side.

From (12) and (13), using the Sokhotski–Plemelj formula $2\Phi(x) = \Phi^+(x) + \Phi^-(x)$, we obtain the following asymptotic formulas for the leading part of the integral in the sense of the principal value:

$$\left\{ \int_{-1}^1 \frac{w(\tau)}{\tau - t} d\tau \right\}_{t \rightarrow -1} = -2^\alpha \pi \cot(\pi\beta) \{(1+t)^\beta\}_{t \rightarrow -1}, \tag{14}$$

$$\left\{ \int_{-1}^1 \frac{w(\tau)}{\tau - t} d\tau \right\}_{t \rightarrow +1} = 2^\beta \pi \cot(\pi\alpha) \{(1-t)^\alpha\}_{t \rightarrow +1}. \tag{15}$$

Here, it has been taken into account that the power functions $(1+t)^\beta$ and $(1-t)^\alpha$ acquire the coefficients $e^{2i\pi\beta}$ and $e^{-2i\pi\alpha}$, respectively, as one goes around the points -1 and $+1$.

Taking into account the explicit formulas (6) and using (12), (13), we get

$$\left\{ \int_{-1}^1 \frac{w(\tau)}{\tau - z_r^*(t)} d\tau \right\}_{t \rightarrow -1} = -\frac{2^\alpha \pi e^{-i\pi\beta} e^{i\theta_r\beta}}{\sin(\pi\beta)} \{(1+t)^\beta\}_{t \rightarrow -1}, \tag{16}$$

$$\left\{ \int_{-1}^1 \frac{w(\tau)}{\tau - z_s^{**}(t)} d\tau \right\}_{t \rightarrow +1} = \frac{2^\beta \pi e^{i\theta_s\alpha}}{\sin(\pi\alpha)} \{(1-t)^\alpha\}_{t \rightarrow +1}. \tag{17}$$

Note that when deriving the last expression, we have chosen the value $e^{-i\pi\alpha}$ of the multi-valued quantity $(-1)^\alpha$ like for (13) (see also (11)).

The representation of the integrals in (8) of the terms of the kernels (4) and (5) with denominators of degree > 1 are obtained by differentiating the Cauchy integral in the parameter z :

$$\int_{-1}^1 \frac{w(\tau) d\tau}{(\tau - z)^s} = \frac{1}{(s-1)!} \frac{d^{s-1}}{dz^{s-1}} \int_{-1}^1 \frac{w(\tau) d\tau}{\tau - z}, \quad s = 2, 3, \dots \tag{18}$$

For the weight function (2) we introduce the notation

$$w(\tau) = (1 - \tau)^\alpha(1 + \tau)^\beta \equiv w^{(\alpha, \beta)}(\tau), \quad \overline{w(\tau)} = w^{(\bar{\alpha}, \bar{\beta})}(\tau). \tag{19}$$

Consecutively differentiating relations (12), (13) and using (18), we obtain the following expressions for the leading terms of the expansions of the corresponding integrals ($s = 2, 3, \dots$):

$$\left\{ \int_{-1}^1 \frac{w^{(\alpha, \beta)}(\tau) d\tau}{(\tau - z)^s} \right\}_{z \rightarrow -1} = -\frac{2^\alpha \pi e^{-i\pi\beta}}{\sin(\pi\beta)} \frac{\beta(\beta - 1)\dots(\beta - s + 2)}{(s - 1)!} \{(1 + z)^{\beta - s + 1}\}_{z \rightarrow -1}, \tag{20}$$

$$\left\{ \int_{-1}^1 \frac{w^{(\alpha, \beta)}(\tau) d\tau}{(\tau - z)^s} \right\}_{z \rightarrow +1} = \frac{2^\beta \pi e^{i\pi\alpha}}{\sin(\pi\alpha)} \frac{\alpha(\alpha - 1)\dots(\alpha - s + 2)}{(s - 1)!} \{(1 - z)^{\alpha - s + 1}\}_{z \rightarrow +1}. \tag{21}$$

In view of (20) and (6), for the generic term of the first series in (4) (the second series is bounded for $t \rightarrow -1$), we have

$$\begin{aligned} & \left\{ \int_{-1}^1 A_k(t) \frac{(1 + \tau)^p(1 + t)^j}{(\tau - z_r^*(t))^{p+j+1}} w^{(\alpha, \beta)}(\tau) d\tau \right\}_{t \rightarrow -1} = \left\{ A_k(t)(1 + t)^j \int_{-1}^1 \frac{w^{(\alpha, \beta+p)}(\tau)}{(\tau - z_r^*(t))^{p+j+1}} d\tau \right\}_{t \rightarrow -1} \\ & = -A_k(-1) \frac{2^\alpha \pi e^{-i\pi(\beta+p)}}{\sin[\pi(\beta + p)]} \frac{(\beta + p)(\beta + p - 1)\dots(\beta - j + 1)}{(p + j)!} \{(1 + t)^j(1 + z_r^*(t))^{\beta - j}\}_{t \rightarrow -1} \\ & = -2^\alpha A_k(-1) \frac{\pi e^{-i\pi\beta} e^{i\theta_r(\beta - j)}}{\sin(\pi\beta)} \frac{(\beta + p)(\beta + p - 1)\dots(\beta - j + 1)}{(p + j)!} \{(1 + t)^\beta\}_{t \rightarrow -1}. \end{aligned} \tag{22}$$

In a similar way, using (21), we obtain an expression for the leading term of the integral of the generic term of the second series in (4) that goes to infinity as $t \rightarrow +1$:

$$\begin{aligned} & \left\{ \int_{-1}^1 B_l(t) \frac{(1 - \tau)^m(1 - t)^n}{(\tau - z_s^{**}(t))^{m+n+1}} w^{(\alpha, \beta)}(\tau) d\tau \right\}_{t \rightarrow +1} \\ & = 2^\beta B_l(+1) \frac{(-1)^n \pi e^{i\theta_s(\alpha - n)}}{\sin(\pi\alpha)} \frac{(\alpha + m)(\alpha + m - 1)\dots(\alpha - n + 1)}{(m + n)!} \{(1 - t)^\alpha\}_{t \rightarrow +1}. \end{aligned} \tag{23}$$

The expressions for the leading parts of the integrals of the generic terms of the series in (5) with the weight $w(\tau)$ (see (8)) can be obtained by replacing α (β) by $\bar{\alpha}$ ($\bar{\beta}$) and A_k (B_l) by C_k (D_l) in (22) and (23):

$$\begin{aligned} & \left\{ \int_{-1}^1 C_k(t) \frac{(1 + \tau)^p(1 + t)^j}{(\tau - z_r^*(t))^{p+j+1}} w^{(\bar{\alpha}, \bar{\beta})}(\tau) d\tau \right\}_{t \rightarrow -1} \\ & = -2^{\bar{\alpha}} C_k(-1) \frac{\pi e^{-i\pi\bar{\beta}} e^{i\theta_r(\bar{\beta} - j)}}{\sin(\pi\bar{\beta})} \frac{(\bar{\beta} + p)(\bar{\beta} + p - 1)\dots(\bar{\beta} - j + 1)}{(p + j)!} \{(1 + t)^{\bar{\beta}}\}_{t \rightarrow -1}, \end{aligned} \tag{24}$$

$$\begin{aligned} & \left\{ \int_{-1}^1 D_l(t) \frac{(1 - \tau)^m(1 - t)^n}{(\tau - z_s^{**}(t))^{m+n+1}} w^{(\bar{\alpha}, \bar{\beta})}(\tau) d\tau \right\}_{t \rightarrow +1} \\ & = 2^{\bar{\beta}} D_l(+1) \frac{(-1)^n \pi e^{i\theta_s(\bar{\alpha} - n)}}{\sin(\pi\bar{\alpha})} \frac{(\bar{\alpha} + m)(\bar{\alpha} + m - 1)\dots(\bar{\alpha} - n + 1)}{(m + n)!} \{(1 - t)^{\bar{\alpha}}\}_{t \rightarrow +1}. \end{aligned} \tag{25}$$

It can be seen that representations (22)–(25) of the integrals of the generic terms of the series in (4) and (5) also cover the cases $p = j = 0$ and $m = n = 0$ (see (16) and (17)). Summation of these representations with respect to the parameters of the corresponding series in (4) and (5), together

with (14) and (15), allows us to obtain analytic formulas for the leading parts of all integrals in (8) and separate the factors that go to infinity as $t \rightarrow \pm 1$ ($-1 < \text{Re } \alpha, \text{Re } \beta < 0$).

Let us examine the representation of the conjugate characteristic part (9) of equation (1). The expression for the integral in the sense of principal value in (9) is easily obtained by the replacement of α with $\bar{\alpha}$ and β with $\bar{\beta}$ in (14) and (15). In order to obtain representations for the integrals of the generalized kernels (4) and (5) in (9), one should perform similar transformations $\alpha \leftrightarrow \bar{\alpha}$ and $\beta \leftrightarrow \bar{\beta}$ in (22)–(25). Moreover, it should be taken into account that passing to conjugates in these kernels is accompanied by the replacement of the coefficients A_k, B_l, C_k, D_l and the functions $z_r^*(t), z_s^{**}(t)$ by their conjugates (see (4) and (5)), the last operation, in view of (6), being equivalent to the replacement of θ_r by $2\pi - \theta_r$ and θ_s by $-\theta_s$. The final expressions will not be written out here, since the comparison of the representations obtained in the above way with (22)–(25) shows that the former can be obtained from the latter by formal passage to the conjugate quantities.

15.5-3. Equations for the Exponents of Singularity of a Solution.

In order to obtain an equation for the exponent β , we write the expressions of the characteristic part (8) of equation (1) and its complex conjugate (9) for $t \rightarrow -1$:

$$\begin{aligned} I_s(t) &= 2^\alpha \Delta_{11}^-(\beta) \left\{ (1+t)^\beta \right\}_{t \rightarrow -1} u(-1) + 2^{\bar{\alpha}} \Delta_{12}^-(\bar{\beta}) \left\{ (1+t)^{\bar{\beta}} \right\}_{t \rightarrow -1} \overline{u(-1)}, \\ \bar{I}_s(t) &= 2^\alpha \Delta_{21}^-(\beta) \left\{ (1+t)^\beta \right\}_{t \rightarrow -1} u(-1) + 2^{\bar{\alpha}} \Delta_{22}^-(\bar{\beta}) \left\{ (1+t)^{\bar{\beta}} \right\}_{t \rightarrow -1} \overline{u(-1)}. \end{aligned} \tag{26}$$

Here,

$$\begin{aligned} \Delta_{11}^-(\beta) &= a(-1) + i \cot(\pi\beta)b(-1) + 2^{-\alpha} \{K_g, w\}_{t \rightarrow -1}, \\ \Delta_{12}^-(\bar{\beta}) &= 2^{-\bar{\alpha}} \{L_g, \bar{w}\}_{t \rightarrow -1}, \\ \Delta_{21}^-(\beta) &= \overline{\Delta_{12}^-(\bar{\beta})} = 2^{-\alpha} \{\bar{L}_g, w\}_{t \rightarrow -1}, \\ \Delta_{22}^-(\bar{\beta}) &= \overline{\Delta_{11}^-(\beta)} = \overline{a(-1)} - i \cot(\pi\bar{\beta})\bar{b}(-1) + 2^{-\bar{\alpha}} \{\bar{K}_g, \bar{w}\}_{t \rightarrow -1}, \end{aligned} \tag{27}$$

where $\{K_g, w\}_{t \rightarrow -1}$ stands for the coefficient of the leading term of the asymptotic expansion of the integral of kernel $K_g(t, \tau)$ with weight $w(\tau)$ as $t \rightarrow -1$. In view of (4), this coefficient is a sum of bounded factors of expressions calculated on the basis of (22), i.e.,

$$\{K_g, w\}_{t \rightarrow -1} = -\frac{2^\alpha \pi e^{-i\pi\beta}}{\sin(\pi\beta)} \sum_{p,j,k,r} \left[A_k(-1) e^{i\theta_r(\beta-j)} \frac{(\beta+p)(\beta+p-1)\dots(\beta-j+1)}{(p+j)!} \right]. \tag{28}$$

Using (24), we obtain a similar expression for the integral of the function $L_g(t, \tau)\overline{w(\tau)}$ in (8):

$$\{L_g, \bar{w}\}_{t \rightarrow -1} = -\frac{2^{\bar{\alpha}} \pi e^{-i\pi\bar{\beta}}}{\sin(\pi\bar{\beta})} \sum_{p,j,k,r} \left[C_k(-1) e^{i\theta_r(\bar{\beta}-j)} \frac{(\bar{\beta}+p)(\bar{\beta}+p-1)\dots(\bar{\beta}-j+1)}{(p+j)!} \right]. \tag{29}$$

There is no need to write out the coefficients of the leading asymptotic terms of the integrals of generalized kernels in (9), because of the above-mentioned fact that these coefficients are the complex conjugates of the coefficients (28) and (29). This fact is reflected in the relation between the functions $\Delta_{hq}^-(\beta)$ ($h, q = 1, 2$) in (27).

Let us rewrite the expression (26) in the form

$$\begin{aligned} I_s(t) &= \left\{ (1+t)^{\text{Re } \beta} \right\}_{t \rightarrow -1} \left[2^\alpha \Delta_{11}^-(\beta) \left\{ (1+t)^{i \text{Im } \beta} \right\}_{t \rightarrow -1} u(-1) \right. \\ &\quad \left. + 2^{\bar{\alpha}} \Delta_{12}^-(\bar{\beta}) \left\{ (1+t)^{-i \text{Im } \beta} \right\}_{t \rightarrow -1} \overline{u(-1)} \right], \\ \bar{I}_s(t) &= \left\{ (1+t)^{\text{Re } \beta} \right\}_{t \rightarrow -1} \left[2^\alpha \Delta_{21}^-(\beta) \left\{ (1+t)^{i \text{Im } \beta} \right\}_{t \rightarrow -1} u(-1) \right. \\ &\quad \left. + 2^{\bar{\alpha}} \Delta_{22}^-(\bar{\beta}) \left\{ (1+t)^{-i \text{Im } \beta} \right\}_{t \rightarrow -1} \overline{u(-1)} \right]. \end{aligned}$$

It is easy to see that the first factors in the right-hand sides go to infinity as $t \rightarrow -1$. Since the other terms of equations (7), not involved in (8), and the right-hand side of (7) are bounded for $t \rightarrow -1$, it has to be required that the second factors (those in square brackets) be equal to zero. This brings us to a system of two homogeneous algebraic equations for the values $u(-1)$ and $\overline{u(-1)}$:

$$\begin{aligned} 2^\alpha \Delta_{11}^-(\beta) \{(1+t)^{i \operatorname{Im} \beta}\}_{t \rightarrow -1} u(-1) + 2^{\bar{\alpha}} \Delta_{12}^-(\bar{\beta}) \{(1+t)^{-i \operatorname{Im} \beta}\}_{t \rightarrow -1} \overline{u(-1)} &= 0, \\ 2^\alpha \Delta_{21}^-(\beta) \{(1+t)^{i \operatorname{Im} \beta}\}_{t \rightarrow -1} u(-1) + 2^{\bar{\alpha}} \Delta_{22}^-(\bar{\beta}) \{(1+t)^{-i \operatorname{Im} \beta}\}_{t \rightarrow -1} \overline{u(-1)} &= 0. \end{aligned} \tag{30}$$

According to the statement of the problem, the unknown function $u(\tau)$ does not vanish at the ends of the interval $[-1, 1]$ (see (3)), and therefore, in order to satisfy this system it is necessary to require that its determinant be equal to zero. Interpreting an ambiguity of the form $x^{ix} (x \rightarrow 0)$, dividing by equal factors, and taking into account the relation between functions (27), we finally obtain the following transcendental equation for the singularity exponent of the solution of equation (1) at the left endpoint of the integration interval:

$$\Delta_{11}^-(\beta) \overline{\Delta_{11}^-(\beta)} - \Delta_{12}^-(\bar{\beta}) \overline{\Delta_{12}^-(\bar{\beta})} = 0, \tag{31}$$

where

$$\begin{aligned} \Delta_{11}^-(\beta) &= a(-1) + i \cot(\pi\beta)b(-1) \\ &\quad - \frac{\pi e^{-i\pi\beta}}{\sin(\pi\beta)} \sum_{p,j,k,r} \left[A_k(-1) e^{i\theta_r(\beta-j)} \frac{(\beta+p)(\beta+p-1)\dots(\beta-j+1)}{(p+j)!} \right], \end{aligned} \tag{32}$$

$$\Delta_{12}^-(\bar{\beta}) = -\frac{\pi e^{-i\pi\bar{\beta}}}{\sin(\pi\bar{\beta})} \sum_{p,j,k,r} \left[C_k(-1) e^{i\theta_r(\bar{\beta}-j)} \frac{(\bar{\beta}+p)(\bar{\beta}+p-1)\dots(\bar{\beta}-j+1)}{(p+j)!} \right]. \tag{33}$$

Note that the relation between $u(-1)$ and its conjugate $\overline{u(-1)}$ represented by either equation (30) is actually only seeming, since for complex β ($\operatorname{Im} \beta \neq 0$) the limit $\{(1+\eta)^{\pm i \operatorname{Im} \beta}\}_{\eta \rightarrow -1}$ does not exist.

To obtain an equation for the singularity exponent α , one should write the expressions for the singular part (8) of equation (1) and its conjugate (9) for $t \rightarrow +1$, and then argue as above. Omitting intermediate calculations, we obtain the following transcendental equation for α :

$$\Delta_{11}^+(\alpha) \overline{\Delta_{11}^+(\alpha)} - \Delta_{12}^+(\bar{\alpha}) \overline{\Delta_{12}^+(\bar{\alpha})} = 0, \tag{34}$$

where

$$\begin{aligned} \Delta_{11}^+(\alpha) &= a(+1) - i \cot(\pi\alpha)b(+1) \\ &\quad + \frac{\pi}{\sin(\pi\alpha)} \sum_{l,m,n,s} \left[B_l(+1)(-1)^n e^{i\theta_s(\alpha-n)} \frac{(\alpha+m)(\alpha+m-1)\dots(\alpha-n+1)}{(m+n)!} \right], \end{aligned} \tag{35}$$

$$\Delta_{12}^+(\bar{\alpha}) = \frac{\pi}{\sin(\pi\bar{\alpha})} \sum_{l,m,n,s} \left[D_l(+1)(-1)^n e^{i\theta_s(\bar{\alpha}-n)} \frac{(\bar{\alpha}+m)(\bar{\alpha}+m-1)\dots(\bar{\alpha}-n+1)}{(m+n)!} \right]. \tag{36}$$

Thus, the problem of finding the exponents of the asymptotic solution of equation (1) at the endpoints of the integration interval has been reduced to two independent transcendental equations (31) and (34) for these exponents. The roots of these equations lying in the strip $-1 < \operatorname{Re} \alpha, \operatorname{Re} \beta < 0$, are the desired singularity exponents in the weight function (2), and the corresponding root with the minimal real part is the leading exponent of the singularity in the solution of equation (1) at a given endpoint of the integration interval.

15.5-4. Analysis of Equations for Singularity Exponents.

Let us give a theoretical analysis of possible solutions of equations (31) and (34). For definiteness, consider equation (31). Using the relations between the terms involved in that equation, we can transform it to

$$\left(|\Delta_{11}^-(\beta)| - |\Delta_{12}^-(\bar{\beta})|\right) \left(|\Delta_{11}^-(\beta)| + |\Delta_{12}^-(\bar{\beta})|\right) = 0. \quad (37)$$

It is easy to see that the left-hand side of the equation obtained is a real-valued function of the complex singularity exponent β , and the zeroes of this function can be found by equating to zero its first and its second factors, which are also real-valued functions.

Equating to zero the first factor in (37), we obtain the equation

$$|\Delta_{11}^-(\beta)| - |\Delta_{12}^-(\bar{\beta})| = 0, \quad (38)$$

which can be considered as an (implicit) equation $g(x, y) = 0$ of some curve on the plane xy , where x and y are, respectively, the real and the imaginary parts of the exponent β ($x = \text{Re } \beta$, $y = \text{Im } \beta$). Therefore, if at least some part of this curve lies in the strip $-1 < x < 0$, then equation (1) allows for the existence of infinitely many singularity exponents of its solution. In this situation, the quantity β can be fixed only if the unique solvability of equation takes place only under an additional condition, and this condition, in its turn, imposes certain constraints on the singularity exponents. The solvability conditions occurring in applications impose no constraints of that kind (see the references at the end of this section), and the theory of equation (1) with generalized kernels, which might give a definite answer in regard to such a condition, has not been developed to a sufficient extent,* in spite of the fact that equations of type (1) quite often occur in problems of mechanics and mathematical physics. It is apparent from (33) (see also (5) and (7)), that it is the integral of the function $L_g(t, \tau)\varphi(\tau)$ in (1) that is responsible for the appearance of the term $|\Delta_{12}^-(\bar{\beta})|$ in (38). If $L_g(t, \tau) \equiv 0$, this term is absent and (38) reduces to the equation

$$\Delta_{11}^-(\beta) = 0, \quad (39)$$

whose left-hand side is a complex-valued function. This means that in this case there is a system of two real equations

$$\begin{aligned} \text{Re } [\Delta_{11}^-(x + iy)] &\equiv h(x, y) = 0, \\ \text{Im } [\Delta_{11}^-(x + iy)] &\equiv p(x, y) = 0 \end{aligned} \quad (40)$$

for the real and the imaginary parts of the singularity exponent. Of course, in some special cases the curves $h(x, y) = 0$ and $p(x, y) = 0$ may have infinitely many common points (i.e., coincide on a finite arc L). However in actual applied problems as a rule, there are finitely many points of intersection of these curves, and therefore, finitely many solutions of system (40), which are admissible singularity exponents $\beta = x + iy$ for solutions of equation (1).

Thus, analysis of equation (38) shows that the integral (with generalized kernel) of the conjugate of the unknown function in equation (1) leads to a qualitatively new behavior of the singularity exponent: equation (31) defines infinitely many singularity exponents admissible for solutions of equation (1) (provided that a finite part of the curve $g(x, y) = 0$ associated with equation (38) belongs to the strip $-1 < x < 0$).

Equating to zero the second factor in (37) brings us to the system of equations

$$\Delta_{11}^-(\beta) = 0, \quad \Delta_{12}^-(\bar{\beta}) = 0, \quad (41)$$

whose left-hand sides are complex-valued functions. This system is overdetermined, since it imposes four real conditions on two real unknown variables x and y ($\beta = x + iy$). Although (41) is

* One of the rare publications in this area is the monograph by Duduchava (1979) that dealt with singular equations with generalized kernels not containing integrals of the conjugate of the unknown function.

an overdetermined system, it cannot be excluded that there exist β in the strip $-1 < x < 0$ satisfying (41), and this shows that the second factor on the left-hand side of (37) should be taken into account when solving specific problems.

Going back to equation (31), we note that a more narrow class of its solutions can be obtained a priori by taking a real β ($\text{Im } \beta \equiv 0$). In this case, equation (31) becomes

$$\Delta_{11}^-(\beta)\overline{\Delta_{11}^-(\beta)} - \Delta_{12}^-(\beta)\overline{\Delta_{12}^-(\beta)} = 0, \tag{42}$$

where the conjugation applies only to the functions (32) and (33), not to the parameter β ($\bar{\beta}$ in (33) must be replaced by β). This real equation serves to determine real singularity exponents admissible for equation (1). As a rule, there are finitely many such exponents.

There is an important point that should be mentioned in connection with the a priori assumption of β being real. By similarity with an algebraic equation with real coefficients, equation (42) may admit (mutually conjugate) complex roots, in particular. Of course, such roots should be excluded from consideration, since neither they nor their real parts satisfy the original equation (31), and, therefore, are inadmissible for equation (1).

Selecting a root with the minimal real part ($-1 < \text{Re } \beta < 0$) in the solutions of equations (37) and (42) allows us to determine the leading singularity exponent of the solution of equation (1) at the left endpoint of the integration interval.

Prior to solving equation (31), it is convenient to perform a regularization by extracting the factors $1/\sin(\pi\beta)$ and $1/\sin(\pi\bar{\beta})$ in (32) and (33), respectively. After the division of equation (31) by the factor $1/|\sin(\pi\beta)|$, which does not vanish in the strip $-1 < \text{Re } \beta < 0$, the left-hand side of the equation becomes an analytic function in a finite region of the complex plane $x + iy = \beta$. This allows us to use the methods of the theory of analytic functions for solving the equations constructed above.

Similar arguments and remarks are valid for equation (34) for the singularity exponent α .

Table 10 summarizes the above analysis and other results known about exponents of singularity of solutions of singular integral equations of the form (1).

TABLE 10
Singularity exponents for solutions of different cases of integral equation (1)

Functions involved in equation (1)	Singularity exponents	Qualitative character of singularities
$a(t)=0,$ $K_g(t, \tau)=0,$ $L_g(t, \tau)=0$	$\alpha=\beta=-1/2$ or $\alpha=-\beta=\pm 1/2$	Real singularity, unique in the interval $-1 < \alpha, \beta < 0$
$K_g(t, \tau)=0,$ $L_g(t, \tau)=0$	$\alpha=-1/2 + i\omega, \beta=-1/2 - i\omega$ or $\alpha=-\beta=\pm 1/2 + i\omega,$ where $\omega = -\frac{1}{2\pi} \ln \frac{b(\pm 1) - a(\pm 1)}{b(\pm 1) + a(\pm 1)}$ (upper sign corresponds to the exponent α , lower corresponds to β)	Complex singularity, unique in the region $-1 < \text{Re } \alpha, \text{Re } \beta < 0$
$L_g(t, \tau)=0$	are determined by the equations: $\Delta_{11}^+(\alpha)=0, \Delta_{11}^-(\beta)=0$	Singularities are complex (in general) and form a discrete set, $\text{Re } \alpha, \text{Re } \beta \neq -1/2$
Complete equation (1)	are determined by the equations: $\Delta_{11}^+(\alpha)\overline{\Delta_{11}^+(\alpha)} - \Delta_{12}^+(\alpha)\overline{\Delta_{12}^+(\alpha)}=0,$ $\Delta_{11}^-(\beta)\overline{\Delta_{11}^-(\beta)} - \Delta_{12}^-(\beta)\overline{\Delta_{12}^-(\beta)}=0$	Complex singularities have continuous distribution. Under the a priori assumption $\text{Im } \alpha = \text{Im } \beta = 0$, real singularities form a discrete set
<i>Remark.</i> Singularity exponents α and β are independent of the Fredholm kernels $K(t, \tau)$ and $L(t, \tau)$ in the integral equation (1).		

15.5-5. Application to an Equation Arising in Fracture Mechanics.

As an application, we use the above approach to determine singularity exponents for an equation that arises in a two-dimensional elasticity problem for a rectilinear crack of unit half-length with a vertex on the interface between two materials with different elastic properties (Linkov, 1999). This problem can be reduced to the integral equation

$$\int_{-1}^1 \frac{\varphi(\tau)}{\tau - t} d\tau + \int_{-1}^1 K_g(t, \tau)\varphi(\tau) d\tau + \int_{-1}^1 L_g(t, \tau)\overline{\varphi(\tau)} d\tau = f(t), \quad -1 < t < 1, \quad (43)$$

where

$$K_g(t, \tau) = \frac{A_0}{\tau - z_0^*} + \frac{A_1}{\tau - z_1^*} + A_2 \frac{1 + \tau}{(\tau - z_0^*)^2} + A_3 \frac{(1 + \tau)(1 + t)}{(\tau - z_0^*)^3},$$

$$A_0 = -\frac{\chi_1}{2}e^{2i\gamma}, \quad A_1 = \frac{\chi_2}{2}e^{-2i\gamma}, \quad A_2 = \frac{\chi_2}{2}(1 - e^{-2i\gamma})^2e^{4i\gamma}, \quad A_3 = 2a_2e^{2i\gamma};$$

$$L_g(t, \tau) = C_0 \frac{1 + \tau}{(\tau - z_0^*)^2} + C_1 \frac{1 + \tau}{(\tau - z_1^*)^2}, \quad C_0 = -\frac{\chi_2}{2}(1 - e^{-2i\gamma})e^{4i\gamma}, \quad C_1 = -\frac{\chi_2}{2}(1 - e^{2i\gamma})e^{-2i\gamma};$$

$$z_0^* = -1 + (1 + t)e^{i\theta_0}, \quad 0 < \theta_0 = 2\gamma < 2\pi; \quad z_1^* = -1 + (1 + t)e^{i\theta_1}, \quad 0 < \theta_1 = 2(\pi - \gamma) < 2\pi;$$

$$\chi_1 \equiv \frac{\kappa_2\mu_1 - \kappa_1\mu_2}{\mu_2 + \kappa_2\mu_1}, \quad \chi_2 \equiv \frac{\mu_2 - \mu_1}{\mu_1 + \kappa_1\mu_2},$$

$p(t) = f(t)/\pi$ is a self-balanced load on the crack surface, $\kappa_r = 3 - 4\nu_r$ for the plane-strain state, and $\kappa_r = (3 - \nu_r)/(1 + \nu_r)$ for the plane-stress state; ν_r is the Poisson ratio, μ_r is the shear modulus ($r = 1, 2$). The index 2 in the last expressions refers to the upper half-plane (i.e., μ_2 and ν_2 are its elastic constants), and the index 1 refers to the lower half-plane with the crack whose line forms angle γ with the positive direction of the axis Ox associated with the interface ($0 < \gamma < \pi$).

Equation (43) has a unique solution in the class of functions that may go to infinity at the endpoints of the integration interval, provided that an additional condition is satisfied. The condition is that the displacement jump at the endpoints of the crack is zero:

$$\int_{-1}^1 \varphi(\tau) d\tau = 0.$$

Since the kernels in (43) are bounded for $\tau = t \rightarrow +1$, the expression (36) and the last term in (35) are equal to zero, and from (34) we obtain the equation $\cot(\pi\alpha) = 0$ ($a(t) \equiv 0$). The solution of this equation with the minimal real part ($\text{Re } \alpha > -1$) is the root $\alpha_1 = -1/2$ corresponding to the common root singularity of the unknown function.

Calculating the expressions (32) and (33) for the kernels in (43), we have

$$\Delta_{11}^-(\beta) = F(\beta) \left\{ \cos(\pi\beta) + e^{-i\pi\beta} \left[A_0e^{i\beta\theta_0} + A_1e^{i\beta\theta_1} + A_2e^{i\beta\theta_0}(\beta + 1) + \frac{1}{2}A_3e^{i(\beta-1)\theta_0}(\beta + 1)\beta \right] \right\}, \quad (44)$$

$$\Delta_{12}^-(\bar{\beta}) = F(\bar{\beta})e^{-i\pi\bar{\beta}} \left[C_0e^{i\bar{\beta}\theta_0}(\bar{\beta} + 1) + C_1e^{i\bar{\beta}\theta_1}(\bar{\beta} + 1) \right],$$

where $F(\beta) = -\pi / \sin(\pi\beta)$.

The complex solution of equation (31) with the minimal real part was obtained with the help of graphical analysis and numerical methods (the Müller method, the chord method, and the golden section method).

Figure 7 shows the dependence of the leading complex singularity exponent β on the angle γ within the range $0 < \gamma < \pi/2$ ($\beta(\pi - \gamma) = \bar{\beta}(\gamma)$) for $\nu_1 = \nu_2 = 0.3$ (plane strain) for two cases




Figure 7. Dependence of the leading complex singularity exponent β on the angle γ resulting from Eq. (38).




Figure 8. Dependence of the leading singularity exponent β on the angle γ resulting from Eq. (42).

$\mu_1/\mu_2 = 0.1$ and $\mu_1/\mu_2 = 10$ (the respective exponents are labelled by $\beta_{0,1}$ and β_{10}). Note that the function $g(x, y) = 0$ (see the interpretation of equation (38) in Section 15.5-4) has a kink at the point corresponding to the root with the minimal real part. Note also that for the parameters of the problem under consideration, numerical experiments have shown that the equation $\Delta_{11}^-(\beta) = 0$, the first equation in system (41), has suitable roots (with $-1 < \text{Re } \beta < 0$) and the equation $\Delta_{12}^-(\beta) = 0$ has no roots.

Figure 8 gives calculation results for the singularity exponent which were obtained using simplified equation (42) with the parameters of the problem being the same. The set of roots of this equation consists of two real values and, for some parameters of the problem, two complex-conjugate ones. The minimal real root is always greater than the corresponding real part of the complex root (Fig. 7), except for the points at which its imaginary part changes sign.

References for Section 15.5: F. E. Erdogan (1975), F. D. Gakhov (1977, 1990), R. Duduchava (1979), A. F. Nikiforov and V. B. Uvarov (1988), N. I. Muskhelishvili (1992), W. H. Press et al. (1992), M. P. Savruk et al. (1999), A. M. Linkov (2002), A. V. Andreev (2007).

15.6. Direct Numerical Solution of Singular Integral Equations with Generalized Kernels*

15.6-1. Preliminary Remarks.

Below, we describe some approaches to the direct numerical solution of integral equations with generalized kernels of the Cauchy type (see Section 15.5). These approaches are based on the method of collocation and are more or less traditional, but due to the class of equations examined here have some specific features which require some special considerations.

The first characteristic feature of the class of equations considered here is the presence of nontrivial (generally complex) singularities of the solution at the endpoints of the integration interval.

In order to obtain integral (nonlocal) characteristics of solutions of equations with generalized kernels, one can adopt well-known numerical approaches that do not take into account the asymptotic behavior of a solution near its singular points at the ends of the integration interval. On the other hand, numerical experiments show that in some situations such methods (for instance, the method of discrete vortices) applied to integral equations with generalized kernels give inadequate results, even if used to find integral characteristics of a solution (see the next paragraph). Moreover, it is of special interest to obtain a fairly accurate local distribution of solution within the framework of the process of its numerical construction, which requires utilization of methods explicitly taking

* Section 15.6 was written by A. V. Andreev.

into account asymptotic behavior of solutions. In particular, it is very important for construction of correct solution asymptotics near the endpoints of the integration interval. Thus, direct numerical solution of singular integral equations with generalized kernels presumes that one has to find a bounded function $u(\tau)^*$, while the Jacobi weight function $w(\tau)$ with singularities is supposed to be known from preliminary analysis, and its asymptotic behavior at the ends of the integration interval is explicitly taken into account in numerical approximations of integrals and other calculations.

The second characteristic feature of equations with generalized kernels is that, as a rule, the analytic continuation of integral kernels (in the integration variable) has singularities outside the integration line, and this fact necessitates the application of high-precision quadrature methods for the numerical approximation of integrals with such kernels. In this connection, quadrature formulas of the highest algebraic accuracy (like the Gauss method) are used below, and quadrature formulas of interpolation type are used to ensure greater flexibility of the collocation method.

The material presented below can be divided into two parts: first, in Sections 15.6-2 to 15.6-4 we describe auxiliary numerical-analytical results, and then, in Sections 15.6-5, 15.6-6 we apply them to the construction of solutions to singular integral equations; in particular, we give examples of their numerical realization and compare its results with exact analytical solutions.

15.6-2. Quadrature Formulas for Integrals with the Jacobi Weight Function.

For the numerical approximation of a nonsingular integral with the weight function $w(\tau)$ in the form of a sum we use the Gauss–Jacobi quadrature formula (of the highest algebraic precision):

$$\int_{-1}^1 u(\tau)w(\tau) d\tau = \sum_{k=1}^n W_k u(\tau_k), \quad \text{Re } \alpha, \text{Re } \beta > -1. \tag{1}$$

Here

$$q_n^{(\alpha,\beta)}(t) = \int_{-1}^1 \frac{w(\tau)P_n^{(\alpha,\beta)}(\tau)}{\tau - t} d\tau, \quad W_k = \frac{q_n^{(\alpha,\beta)}(\tau_k)}{[P_n^{(\alpha,\beta)}(\tau_k)]'}, \tag{2}$$

and $P_n^{(\alpha,\beta)}(\tau)$ is the Jacobi polynomial defined by

$$\begin{aligned} P_n^{(\alpha,\beta)}(\tau) &= \frac{(-1)^n}{2^n n!} (1 - \tau)^{-\alpha} (1 + \tau)^{-\beta} \frac{d^n}{d\tau^n} \left[(1 - \tau)^{\alpha+n} (1 + \tau)^{\beta+n} \right] \\ &= 2^{-n} \sum_{m=0}^n C_{n+\alpha}^m C_{n+\beta}^{n-m} (\tau - 1)^{n-m} (\tau + 1)^m, \end{aligned} \tag{3}$$

C_b^a are binomial coefficients, and the nodes τ_k of the quadrature formulas form the set of roots of this polynomial,

$$P_n^{(\alpha,\beta)}(\tau_k) = 0, \quad k = 1, 2, \dots, n. \tag{4}$$

Formula (1) is exact if $u(\tau)$ is a polynomial of a degree $\leq 2n - 1$ (or briefly, $u(\tau) \in \mathcal{P}_{2n-1}$). In formula (2) and below we use the notation $[F(\tau_k)]' = \left. \frac{dF}{d\tau} \right|_{\tau=\tau_k}$.

For $\text{Re } \alpha > -1$, $\text{Re } \beta > -1$, and $\text{Im } \alpha = \text{Im } \beta = 0$, the roots of the Jacobi polynomial are simple and belong to the interval $\tau \in (-1, 1)$. The quadrature formula (1) remains valid in the case of complex values of α and β , but in this case the roots of the Jacobi polynomial also turn out to be complex ($\text{Im } \tau_k \neq 0$) and lie near the interval $\tau \in (-1, 1)$.

* Recall that a solution of a singular integral equation is sought in the form of the product

$$\varphi(\tau) = u(\tau)w(\tau), \quad \text{where } w(\tau) = (1 - \tau)^\alpha (1 + \tau)^\beta, \quad -1 \leq \tau \leq 1, \quad \text{Re } \alpha, \text{Re } \beta > -1$$

(see formulas (2)–(3) in Subsection 15.5-1).

Note that the function $q_n^{(\alpha,\beta)}(t)$ can be easily expressed through the Jacobi function of the second kind $Q_n^{(\alpha,\beta)}(t)$ and the weight function $w(\tau)$:

$$q_n^{(\alpha,\beta)}(t) = w(t)Q_n^{(\alpha,\beta)}(t), \quad (5)$$

and the zeroes of these functions coincide on the interval $t \in (-1, 1)$.

For a singular integral with the Cauchy kernel, the following modification of the Gauss–Jacobi quadrature formula holds:

$$\int_{-1}^1 \frac{u(\tau)w(\tau)}{\tau - t} d\tau = \sum_{k=1}^n W_k \frac{u(\tau_k)}{\tau_k - t} + u(t) \frac{q_n^{(\alpha,\beta)}(t)}{P_n^{(\alpha,\beta)}(t)}, \quad -1 < t \neq \tau_k < 1, \quad \operatorname{Re} \alpha, \operatorname{Re} \beta > -1. \quad (6)$$

This formula is exact if $u(\tau) \in \mathcal{P}_{2n}$ (i.e., $u(\tau)$ is a polynomial of degree $\leq 2n$).

For a discrete set of points t_m such that (see (5))

$$Q_n^{(\alpha,\beta)}(t_m) = 0, \quad t_m \neq \tau_k, \quad (7)$$

the quadrature formula (6) becomes similar to (1):

$$\int_{-1}^1 \frac{u(\tau)w(\tau)}{\tau - t_m} d\tau = \sum_{k=1}^n W_k \frac{u(\tau_k)}{\tau_k - t_m}. \quad (8)$$

For the restoration of the values of the unknown function $u(\tau)$ on the entire interval $\tau \in [-1, 1]$ from its values on the discrete set τ_k ($k = 1, 2, \dots, n$), one can use the Lagrange interpolation polynomial, which it is convenient to write in the following form (since the interpolation is with respect to the zeroes of the Jacobi polynomial):

$$u(\tau) = P_n^{(\alpha,\beta)}(\tau) \sum_{k=1}^n \frac{u(\tau_k)}{(\tau - \tau_k)[P_n^{(\alpha,\beta)}(\tau_k)]'}. \quad (9)$$

This interpolation representation is exact if $u(\tau)$ is a polynomial of a degree $\leq n - 1$. Note that the representation (9) may be useful for the approximation of the term outside the integral in a singular equation of the second kind.

Moreover, on the basis of the approximation (9), one can construct quadrature formulas of interpolation type for a singular integral. Substituting (9) into (6), we obtain the following quadrature formula for the singular integral:

$$\int_{-1}^1 \frac{u(\tau)w(\tau)}{\tau - t} d\tau = \sum_{k=1}^n W_k^{(s)}(t)u(\tau_k), \quad -1 < t < 1, \quad \operatorname{Re} \alpha, \operatorname{Re} \beta > -1, \quad (10)$$

which is precise for $u(\tau) \in \mathcal{S}_{n-1}$. Here,

$$W_k^{(s)}(t) = \begin{cases} \frac{q_n^{(\alpha,\beta)}(\tau_k) - q_n^{(\alpha,\beta)}(t)}{[P_n^{(\alpha,\beta)}(\tau_k)]'(\tau_k - t)} & \text{if } t \neq \tau_k, \\ \frac{[q_n^{(\alpha,\beta)}(\tau_k)]'}{[P_n^{(\alpha,\beta)}(\tau_k)]'} & \text{if } t = \tau_k. \end{cases} \quad (11)$$

Note that the lower expression for the weight in (11) is obtained from the upper one by passing to the limit as $t \rightarrow \tau_k$. It can be seen that the quadrature formula (10) for the singular integral (unlike the similar formula (6)) holds also at $t = \tau_k$. Moreover, formula (10) yields an expression which, in contrast to (6), is an approximation based only on the density values at the nodes of the quadrature formula (see also (8)).

15.6-3. Approximation of Solutions in Terms of a System of Orthogonal Polynomials.

As mentioned in the previous subsection, for $\text{Im } \alpha \neq 0$ or $\text{Im } \beta \neq 0$, the roots of the Jacobi polynomial are complex ($\text{Im } \tau_k \neq 0$). This is an obstacle to the utilization of the above quadrature formulas for the approximation of integrals, since it becomes necessary to find the unknown function of the integral equation outside its domain $\tau \in [-1, 1]$.

To construct a solution of an equation with complex asymptotics at the endpoints of the integration interval, let us represent the unknown function $u(\tau)$ in the form of expansion in terms of a finite system of Jacobi polynomials which are orthogonal on the segment $[-1, 1]$ with weight function $w(\tau)$:

$$u(\tau) = \sum_{k=0}^n c_k P_k^{(\alpha, \beta)}(\tau). \quad (12)$$

Here, c_k are complex constants to be determined, and $P_k^{(\alpha, \beta)}(\tau)$ is a Jacobi polynomial of real argument τ with complex α and β . Such a representation allows us to perform analytical integration of the singular integral in terms of special functions on the basis of the following integral representation of the Jacobi function of the second kind $Q_k^{(\alpha, \beta)}(t)$ (see (2), (5)):

$$\int_{-1}^1 \frac{w(\tau) P_k^{(\alpha, \beta)}(\tau)}{\tau - t} d\tau = w(t) Q_k^{(\alpha, \beta)}(t), \quad -1 < t < 1. \quad (13)$$

When using (12) for the approximation of a solution of an integral equation, one has to deal with integrals of the form

$$h_k(t) \equiv \frac{1}{w(t)} \int_{-1}^1 k(t, \tau) w(\tau) P_k^{(\alpha, \beta)}(\tau) d\tau, \quad -1 < t < 1, \quad (14)$$

where $k(t, \tau)$ is a generalized kernel. In general, such a kernel (see, for instance, (4) and (5) in Section 15.5) is nonanalytic for τ in a neighborhood of the segment $\tau \in [-1, 1]$ on the complex plane. At the same time, if for a specific kernel one can separate its poles from the region of the complex roots of the Jacobi polynomial, then a direct and fairly precise approach to the calculation of integrals (14) can be realized by the method of mechanical Gauss–Jacobi quadratures (1). If such an operation is impossible or entails very difficult calculations, the following technique can be used.

Let us approximate the kernel $k(t, \tau)$ by a degenerate kernel in the form of a polynomial of degree N with respect to τ :

$$k(t, \tau) = \sum_{s=0}^N c_s(t) \tau^s, \quad t \in (-1, 1). \quad (15)$$

Obviously, the Jacobi polynomials can be represented in a similar form

$$P_k^{(\alpha, \beta)}(\tau) = \sum_{l=0}^k g_l^{(k)} \tau^l. \quad (16)$$

Here, the superscript in the coefficients $g_l^{(k)}$ refers to the highest degree of the polynomial.

Using (15) and (16), we obtain the following expression for the integral (14):

$$h_k(t) = \frac{1}{w(t)} \sum_{j=0}^{N+k} d_j^{(k)}(t) \int_{-1}^1 \tau^j w(\tau) d\tau. \quad (17)$$

Here, the coefficients $d_j^{(k)}(t)$ can be written in the form

$$d_j^{(k)}(t) = \sum_{s=0}^j c_s(t)g_{j-s}^{(k)}, \tag{18}$$

which is obtained on the basis of multiplication of the series (15) and (16). Note that in the sum (18), one should take $c_s(t) = 0$ for $s > N$ and $g_{j-s}^{(k)} = 0$ for $j - s > k$.

To calculate the integrals involved in (17), we use the identity

$$\int_{-1}^1 \tau^j w(\tau) d\tau = \int_{-1}^1 \frac{\tau^{j+1} w(\tau)}{\tau - t_*} d\tau - t_* \int_{-1}^1 \frac{\tau^j w(\tau)}{\tau - t_*} d\tau. \tag{19}$$

Here and in what follows, $-1 < t_* < 1$ is a fixed auxiliary parameter. The integrals in the right-hand side of the last expression can be calculated with the help of (13), which, in view of (16), can be represented in the form

$$\begin{aligned} Q_m^{(\alpha, \beta)}(t_*) &= \frac{1}{w(t_*)} \sum_{l=0}^m g_l^{(m)} \int_{-1}^1 \frac{\tau^l w(\tau)}{\tau - t_*} d\tau \\ &= \frac{1}{w(t_*)} \left[g_0^{(m)} \int_{-1}^1 \frac{w(\tau)}{\tau - t_*} d\tau + g_1^{(m)} \int_{-1}^1 \frac{\tau w(\tau)}{\tau - t_*} d\tau + \dots + g_m^{(m)} \int_{-1}^1 \frac{\tau^m w(\tau)}{\tau - t_*} d\tau \right]. \end{aligned}$$

The last expression implies that the integrals of the form

$$\begin{aligned} I_m(t_*) &\equiv \int_{-1}^1 \frac{\tau^m w(\tau)}{\tau - t_*} d\tau = \frac{1}{g_m^{(m)}} \left[w(t_*) Q_m^{(\alpha, \beta)}(t_*) - \sum_{p=0}^{m-1} g_p^{(m)} I_p(t_*) \right], \tag{20} \\ I_0(t_*) &= w(t_*) Q_0^{(\alpha, \beta)}(t_*). \end{aligned}$$

can be calculated on the basis of the above recurrent relation (note that $g_0^{(0)} = 1$).

Let us introduce the function

$$\begin{aligned} S_m(t_*) &\equiv \frac{I_m(t_*)}{w(t_*)} = \frac{1}{g_m^{(m)}} \left[Q_m^{(\alpha, \beta)}(t_*) - \sum_{p=0}^{m-1} g_p^{(m)} S_p(t_*) \right], \tag{21} \\ S_0(t_*) &= Q_0^{(\alpha, \beta)}(t_*), \end{aligned}$$

calculated on the basis of a similar recurrent relation. Substituting (19) into (17) and using (20), (21), we finally obtain

$$h_k(t) = \frac{w(t_*)}{w(t)} \sum_{j=0}^{N+k} d_j^{(k)}(t) [S_{j+1}(t_*) - t_* S_j(t_*)], \quad -1 < t < 1. \tag{22}$$

Note that the above approach to the calculation of integrals (14) based on the expansion of the generalized kernel in power series (15) might be especially convenient if the kernel $k(t, \tau)$ cannot be expressed explicitly and one has to use its representation as an integral. In such a situation, the representation (15) can be obtained with the help of the expansion of the integrand in power series with respect to τ and subsequent analytical integration of that series.

15.6-4. Some Special Functions and Their Calculations.

In order to implement the above methods for the approximation of integrals with the Jacobi weight function, it is necessary to calculate special functions and their roots with great precision and efficiency. Next, we sum up the results necessary for the implementation of the methods and approaches proposed above (see also the references at the end of this section).

To calculate Jacobi polynomials and their set, as well as the corresponding functions of the second kind, it is convenient to use a single recurrent procedure based on the relation

$$\begin{aligned} a_n \Psi_{n+1}^{(\alpha, \beta)}(\tau) &= (b_n + c_n \tau) \Psi_n^{(\alpha, \beta)}(\tau) - d_n \Psi_{n-1}^{(\alpha, \beta)}(\tau), \\ a_n &= 2(n+1)(n+\alpha+\beta+1)(2n+\alpha+\beta), \\ b_n &= (2n+\alpha+\beta+1)(\alpha^2 - \beta^2), \\ c_n &= (2n+\alpha+\beta)(2n+\alpha+\beta+1)(2n+\alpha+\beta+2), \\ d_n &= 2(n+\alpha)(n+\beta)(2n+\alpha+\beta+2). \end{aligned} \quad (23)$$

Here and henceforth in this subsection, we use the symbol Ψ to denote the polynomial P and the function of the second kind Q if they satisfy identical relations.

For the coefficients $b_l^{(k)}$ of a Jacobi polynomial of the form (16), one can construct recurrent relations that can be used for the determination of the coefficients of a polynomial of degree $n+1$ in terms of the coefficients of polynomials of smaller degrees n and $n-1$. Thus, substituting (16) into (23) and equating the coefficients of equal powers of τ , we obtain

$$a_n b_l^{(n+1)} = b_n b_l^{(n)} + c_n b_{l-1}^{(n)} - d_n b_l^{(n-1)}, \quad a_n b_{n+1}^{(n+1)} = c_n b_n^{(n)}, \quad b_n^{(n-1)} = b_{-1}^{(n)} = 0, \quad l = 0, 1, \dots, n. \quad (24)$$

Since the derivative of a Jacobi polynomial (function of the second kind) is expressed through two consecutive polynomials (functions) of the corresponding orders with the same parameters α , β and argument τ :

$$(1 - \tau^2)[\Psi_n^{(\alpha, \beta)}(\tau)]' = (\tilde{a}_n + \tilde{b}_n \tau) \Psi_n^{(\alpha, \beta)}(\tau) + \tilde{c}_n \Psi_{n-1}^{(\alpha, \beta)}(\tau); \quad (25)$$

$$\tilde{a}_n = \frac{n(\alpha - \beta)}{2n + \alpha + \beta}, \quad \tilde{b}_n = -n, \quad \tilde{c}_n = \frac{2(n + \alpha)(n + \beta)}{(2n + \alpha + \beta)}, \quad (26)$$

its calculation reduces to the calculation of coefficients (26) and their substitution into (25) on the final stage of the recurrent procedure (23).

As the initial values in (23), one can use the Jacobi polynomials of the zero and the first orders,

$$P_0^{(\alpha, \beta)}(\tau) = 1, \quad P_1^{(\alpha, \beta)}(\tau) = \frac{1}{2}(\alpha - \beta) + \frac{1}{2}(2 + \alpha + \beta)\tau,$$

and for a function of the second kind, use the initial values obtained from the explicit expression

$$\begin{aligned} Q_n^{(\alpha, \beta)}(\tau) &= -\frac{\pi P_n^{(\alpha, \beta)}(\tau)}{\tan(\pi\beta)} + \frac{(-1)^n 2^{\alpha+\beta}}{w(\tau)} B(n + \alpha + 1, \beta) F\left(n + 1, -n - \alpha - \beta, 1 - \beta; \frac{1 + \tau}{2}\right) \\ &= \frac{\pi P_n^{(\alpha, \beta)}(\tau)}{\tan(\pi\alpha)} - \frac{2^{\alpha+\beta}}{w(\tau)} B(n + \beta + 1, \alpha) F\left(n + 1, -n - \alpha - \beta, 1 - \alpha; \frac{1 - \tau}{2}\right), \end{aligned} \quad (27)$$

where $B(x, y)$ is the beta function, $F(a, b, c; z)$ is the hypergeometric function, and $-1 < \tau < 1$.

For the calculation of the hypergeometric function for $-1 < z = (1 \pm \tau)/2 < 1$, one can use its representation as the Gauss series (see Supplement 11.10)

$$F(a, b, c; z) = 1 + \sum_{m=1}^{\infty} \frac{(a)_m (b)_m}{(c)_m} \frac{z^m}{m!}, \quad (a)_m = a(a+1) \dots (a+m-1). \quad (28)$$

The beta function is related to the gamma function $\Gamma(x)$ by $B(x, y) = \Gamma(x)\Gamma(y)/\Gamma(x+y)$, and the latter can be calculated quite accurately by the Lanczos approximation

$$\Gamma(z) = \sqrt{2\pi} \frac{(z + C_1 - 1/2)^{z-1/2}}{e^{z+C_1-1/2}} \left(s_1 + \sum_{k=2}^m \frac{s_k}{z+k-2} \right), \quad z \neq 0, -1, -2, \dots \quad (29)$$

For $m = 15$, the coefficients of the approximation (29) have the form

$$\begin{aligned} C_1 &= 607/128, & s_1 &= 0.999999999999997, \\ s_2 &= 57.15623566586292, & s_3 &= -59.59796035547549, \\ s_4 &= 14.13609797474174, & s_5 &= -0.491913816097620, \\ s_6 &= 0.339946499848118 \times 10^{-4}, & s_7 &= 0.465236289270485 \times 10^{-4}, \\ s_8 &= -0.983744753048795 \times 10^{-4}, & s_9 &= 0.158088703224912 \times 10^{-3}, \\ s_{10} &= -0.210264441724104 \times 10^{-3}, & s_{11} &= 0.217439618115212 \times 10^{-3}, \\ s_{12} &= -0.164318106536763 \times 10^{-3}, & s_{13} &= 0.844182239838527 \times 10^{-4}, \\ s_{14} &= -0.261908384015814 \times 10^{-4}, & s_{15} &= 0.368991826595316 \times 10^{-5}. \end{aligned}$$

Note that the above methods for the calculation of special functions are applicable for both real and complex parameters and arguments of these functions.

Moreover, for real values one can obtain explicit expressions for the nodes and the weights in quadrature formulas. These expressions were obtained for large values of the discretization parameter, $n \gg 1$. In this case, the Jacobi polynomials and the integral (2) can be written in terms of elementary functions:

$$P_n^{(\alpha, \beta)}(\cos \theta) = \frac{\cos \{ [n + (\alpha + \beta + 1)/2] \theta - (2\alpha + 1)\pi/4 \}}{\sqrt{\pi n} (\sin(\theta/2))^{\alpha+1/2} (\cos(\theta/2))^{\beta+1/2}} + O(n^{-3/2}), \quad 0 < \theta < \pi, \quad (30)$$

$$q_n^{(\alpha, \beta)}(\cos \theta) = 2^{\alpha+\beta} \sqrt{\frac{\pi}{n}} \frac{\sin \{ [n + (\alpha + \beta + 1)/2] \theta - (2\alpha + 1)\pi/4 \}}{(\sin(\theta/2))^{-\alpha+1/2} (\cos(\theta/2))^{-\beta+1/2}} + O(n^{-3/2}). \quad (31)$$

On the basis of these results, one obtains the following approximate expressions for the nodes and the weights in the quadrature formulas:

$$\tau_k \approx \cos \theta_k, \quad \theta_k = \frac{2\alpha - 1 + 4k}{2n + \alpha + \beta + 1} \frac{\pi}{2}, \quad k = 1, 2, \dots, n; \quad (32)$$

$$W_k \approx \frac{2\pi}{2n + \alpha + \beta + 1} \sqrt{1 - \tau_k^2} (1 - \tau_k)^\alpha (1 + \tau_k)^\beta. \quad (33)$$

Note that these expressions for the nodes and weights are precise for $\alpha = \pm 1/2$ and $\beta = \pm 1/2$ for any n .

In the general case, the real roots τ_k of a Jacobi polynomial (or function of the second kind) can be calculated by means of the following algorithm. Choosing a suitable initial approximation $\tau_k^{(1)}$ for the k th root, its value can be found, quickly enough and with given accuracy, in an iteration

process based on the Newton method (of tangential lines), by consecutively refining the position of a root with the help of the expression (i is the number of the iteration)

$$\tau_k^{(i+1)} = \tau_k^{(i)} - \frac{\Psi_n^{(\alpha, \beta)}(\tau_k^{(i)})}{[\Psi_n^{(\alpha, \beta)}(\tau_k^{(i)})]'}, \quad i = 1, 2, \dots \quad (34)$$

It is possible to choose initial approximations for (real) roots of Jacobi polynomials in the form

$$\begin{aligned} \tau_1^{(1)} &= 1 - \frac{(1 + \alpha)[2.78/(4 + n^2) + 0.768\alpha/n^2]}{1 + 1.48\alpha/n + 0.96\beta/n + 0.452\alpha^2/n^2 + 0.83\alpha\beta/n^2}, \\ \tau_2^{(1)} &= \tilde{\tau}_1 - \frac{(1 - \tilde{\tau}_1)(4.1 + \alpha)}{(1 + \alpha)(1 + 0.156\alpha)} \left[1 + \frac{0.06(n - 8)(1 + 0.12\alpha)}{n} \right] \left[1 + \frac{0.012\beta(1 + 0.25|\alpha|)}{n} \right], \\ \tau_3^{(1)} &= \tilde{\tau}_2 - (\tilde{\tau}_1 - \tilde{\tau}_2) \frac{1.67 + 0.28\alpha}{1 + 0.37\alpha} \left[1 + \frac{0.22(n - 8)}{n} \right] \left[1 + \frac{8\beta}{(6.28 + \beta)n^2} \right], \\ \tau_k^{(1)} &= 3\tilde{\tau}_{k-1} - 3\tilde{\tau}_{k-2} + \tilde{\tau}_{k-3}, \quad 3 < k < n - 1, \\ \tau_{n-1}^{(1)} &= \tilde{\tau}_{n-2} + (\tilde{\tau}_{n-2} - \tilde{\tau}_{n-3}) \frac{1 + 0.235\beta}{0.766 + 0.119\beta} \left[1 + \frac{1 + 0.639(n - 4)}{1 + 0.71(n - 4)} \right]^{-1} \left[1 + \frac{20\alpha}{(7.5 + \alpha)n^2} \right]^{-1}, \\ \tau_n^{(1)} &= \tilde{\tau}_{n-1} + (\tilde{\tau}_{n-1} - \tilde{\tau}_{n-2}) \frac{1 + 0.37\beta}{1.67 + 0.28\beta} \left[1 + \frac{0.22(n - 8)}{n} \right]^{-1} \left[1 + \frac{8\alpha}{(6.28 + \alpha)n^2} \right]^{-1}. \end{aligned}$$

Here, the quantities marked with tilde denote approximate values of the roots of the polynomial which were obtained as a result of previous iteration processes (as regards this process).

Initial approximations for (real) roots of a function of the second kind may be chosen in the form

$$\begin{aligned} \tilde{t}_1^{(1)} &= 1 - n^{-5/2}(\alpha + 1/2)^2(1 + \tilde{\tau}_1), \quad \alpha > -1/2; \\ \tilde{t}_k^{(1)} &= (\tilde{\tau}_k + \tilde{\tau}_{k-1})/2, \quad k = 2, 3, \dots, n; \\ \tilde{t}_{n+1}^{(1)} &= -1 - n^{-5/2}(\beta + 1/2)^2(-1 + \tilde{\tau}_n), \quad \beta > -1/2. \end{aligned} \quad (35)$$

When calculating complex roots of Jacobi polynomials ($\text{Im } \alpha \neq 0$ or $\text{Im } \beta \neq 0$), one can take as the initial approximation $\tau_k^{(1)}$ in (34) the roots of the real polynomial $P_n^{(\text{Re } \alpha, \text{Re } \beta)}(\tau)$, which can be found from the equation $P_n^{(\text{Re } \alpha, \text{Re } \beta)}(\tau_k^{(1)}) = 0$, $k = 1, 2, \dots, n$.

15.6-5. Numerical Solution of Singular Integral Equations.

Consider a complete singular integral equation of the first kind

$$\int_{-1}^1 \frac{\varphi(\tau) d\tau}{\tau - t} + \int_{-1}^1 k(t, \tau)\varphi(\tau) d\tau = f(t), \quad -1 < t < 1. \quad (36)$$

Using approximations of the integrals (1) and (6), we write it in the form

$$u(t) \frac{q_n^{(\alpha, \beta)}(t)}{P_n^{(\alpha, \beta)}(t)} + \sum_{k=1}^n W_k u(\tau_k) \left[\frac{1}{\tau_k - t} + k(t, \tau_k) \right] = f(t), \quad -1 < t < 1, \quad t \neq \tau_k. \quad (37)$$

Next, one can realize several versions of the construction of a complete system of algebraic equations for the values $u(\tau_k)$ ($k = 1, 2, \dots, n$) on the basis of the collocation method.

If the function of the second kind $Q_n^{(\alpha,\beta)}(t)$ (see (5) and (7)) has $\geq n$ zeroes on the interval $t \in (-1, 1)$, then, using these zeroes as collocation points (see also (8)), one can construct the complete system of algebraic equations

$$\sum_{k=1}^n W_k u(\tau_k) \left[\frac{1}{\tau_k - t_m} + k(t_m, \tau_k) \right] = f(t_m), \quad m = 1, 2, \dots, n. \tag{38}$$

This situation takes place in the case of $\alpha > -1/2$ or $\beta > -1/2$ (see (35)). If neither of these conditions holds, then the function $Q_n^{(\alpha,\beta)}(t)$ has $n - 1$ zeroes on the interval $t \in (-1, 1)$, and one should use the following approaches.

First of all, we note that in actual applied problems, equation (36) whose solution has singularities in the region $-1 < \alpha, \beta \leq -1/2$ as a rule is accompanied by the condition

$$\int_{-1}^1 \varphi(\tau) d\tau = A \tag{39}$$

(A being a known constant), whose quadrature analogue (see (1))

$$\sum_{k=1}^n W_k u(\tau_k) = A \tag{40}$$

allows us to complete the algebraic system of equations.

At the same time, in cases not covered by this rule, one can use (9) for the interpolation of the unknown function in the first term in (37) and the construction of a complete system of linear equations on an arbitrary set of collocation points $-1 < t_m \neq \tau_k < 1$ ($m = 1, 2, \dots, n$). An equivalent approach is to use the quadrature formula (10) for the approximation of the singular integral in (36), and in the latter case, collocation points can be chosen coincident with the nodes of the quadrature formulas $t_m = \tau_k$ ($m, k = 1, 2, \dots, n$):

$$\sum_{k=1}^n u(\tau_k) \left[W_k^{(s)}(t_m) + W_k k(t_m, \tau_k) \right] = f(t_m), \quad k, m = 1, 2, \dots, n. \tag{41}$$

The last approach is especially convenient in that the special functions necessary for its realization are calculated only for a single system of points, namely, for the nodes of the quadrature formulas.

Note that an important feature of all approaches described above and realized in the framework of the collocation method is the utilization of a quadrature formula of the highest algebraic precision for the approximation of an integral containing a generalized kernel $k(t, \tau)$.

Consider the complete singular integral equation of the second kind with generalized kernel $k(t, \tau)$:

$$a(t)\varphi(t) + \frac{b(t)}{\pi i} \int_{-1}^1 \frac{\varphi(\tau) d\tau}{\tau - t} + \int_{-1}^1 k(t, \tau)\varphi(\tau) d\tau = f(t), \quad -1 < t < 1. \tag{42}$$

If the solution of this equation has real singularities at the ends of the integration interval, then a numerical approximate solution can be constructed by quadrature-collocation methods similar to those described above, with the interpolation polynomial (9) used for the term outside the integral. However, in many actual applied problems, singularities of a solution of equation (42) are complex, and this requires the approach described below.

Using the approximation (12), from (13) and (42) we get:

$$w(t) \sum_{k=0}^n c_k \left[a(t)P_k^{(\alpha,\beta)}(t) + \frac{b(t)}{\pi i} Q_k^{(\alpha,\beta)}(t) + h_k(t) \right] = f(t), \quad -1 < t < 1. \tag{43}$$

To obtain a system of linear algebraic equations for the unknown coefficients c_k ($k = 0, 1, \dots, n$), the expression (43) can be written for the corresponding number of collocation points t_r :

$$w(t_r) \sum_{k=0}^n c_k \left[a(t_r) P_k^{(\alpha, \beta)}(t_r) + \frac{b(t_r)}{\pi i} Q_k^{(\alpha, \beta)}(t_r) + h_k(t_r) \right] = f(t_r), \quad (44)$$

$$r = 0, 1, \dots, n, \quad -1 < t_r < 1.$$

If equation (42) is accompanied by a condition of the form (39), it is necessary to use a slightly modified approach.

Using (12), let us rewrite the additional condition (39) in the form

$$\sum_{k=0}^n c_k \int_{-1}^1 w(\tau) P_k^{(\alpha, \beta)}(\tau) d\tau = A. \quad (45)$$

From the orthogonality condition of the Jacobi polynomials on the interval $\tau \in [-1, 1]$ with the weight function $w(\tau)$, we have

$$\int_{-1}^1 w(\tau) P_k^{(\alpha, \beta)}(\tau) d\tau = \begin{cases} 2^{\alpha+\beta+1} B(\alpha+1, \beta+1) & \text{if } k=0, \\ 0 & \text{if } k>0. \end{cases}$$

Thus, condition (45) immediately allows us to find one of the unknown constants:

$$c_0 = \frac{2^{-1-\alpha-\beta}}{B(\alpha+1, \beta+1)} A. \quad (46)$$

Therefore, in this situation it is necessary to take $k = 1$ in the lower limit of the sum (44), decrease the number of collocation points t_r by 1, and determine the unknown constant c_0 from (46).

Note that in calculating expressions (14) according to (22), when constructing system (44), it is convenient to choose the auxiliary point $t_* \in (-1, 1)$, introduced in (19), to be coincident with one of the points t_r . This allows us to reduce calculations by using in (21) the values $Q_k^{(\alpha, \beta)}(t_r)$ ($r = 0, 1, \dots, n$) obtained on the stage of calculations of the second term in the sum (44). Numerical experiments show that the accuracy of a solution is little affected by which point t_r is chosen as the auxiliary point.

Thus, solving integral equations on the basis of the collocation method amounts to solving of systems of linear algebraic equations, which allows us to determine the coefficients of the approximation of the unknown function or its values on a discrete set of points. Note that the approaches described above can be directly extended to the case of an equation also containing an integral (with regular or generalized kernel) of the complex-conjugate of the unknown function (see (1) in Section 15.1).

15.6-6. Numerical Solutions of Singular Integral Equations of Bueckner Type.

Example 1. Consider a singular integral equation of *Bueckner type* (1966):

$$\int_{-1}^1 \frac{\varphi(\tau) d\tau}{\tau-t} + \int_{-1}^1 \frac{\varphi(\tau) d\tau}{\tau+t+2} = \pi h(t), \quad -1 < t < 1. \quad (47)$$

This equation with generalized kernel $k(t, \tau) = 1/(\tau + t + 2)$ has a unique solution, which, for $h(t) = q = \text{const}$, can be expressed in terms of elementary functions,

$$\varphi(\tau) = q \frac{1+\tau}{\sqrt{1-\tau}\sqrt{3+\tau}}. \quad (48)$$

It can be seen that in this case the weight function, which reflects the asymptotic behavior of the solution at the ends of the integration interval, has the form

$$w(\tau) = (1 - \tau)^{-1/2}(1 + \tau). \tag{49}$$

Table 11 gives the results of numerical solution of equation (47) for $h(t) = q = 1$. This numerical solution was obtained in the framework of the quadrature-collocation approach (38) on the set of ten points τ_k listed in the second column. The third column lists the values of the exact analytical solution (48) calculated at these points for $q = 1$.

TABLE 11
Comparison of exact and numerical solutions of the Bueckner equation (47)

k	Points τ_k	Exact solution (48)	Numerical solution
1	0.9893260	9.64037	9.64044
2	0.9052947	3.13291	3.13293
3	0.7443578	1.78291	1.78293
4	0.5201627	1.16966	1.16968
5	0.2517209	0.80245	0.80246
6	-0.0382037	0.54848	0.54850
7	-0.3250257	0.35852	0.35854
8	-0.5844217	0.21242	0.21246
9	-0.7943919	0.10335	0.10342
10	-0.9371120	0.03145	0.03172

As one can see from the table, the numerical solution is accurate to four or five significant digits for the first nine τ_k even for only $n = 10$. Note that the relative error increases towards the left endpoint of the interval $(-1, 1)$, where it reaches $\epsilon_{\max} = 0.83\%$. This increase is due to the fact that as t decreases, the pole of the function $g(z) = 1/(z + t + 2)$ approaches the left endpoint of the interval, which worsens the quadrature approximation of the integrals.

Example 2. Now consider a more general Bueckner equation

$$\int_{-1}^1 \frac{\varphi(\tau) d\tau}{\tau - t} + D \int_{-1}^1 \frac{\varphi(\tau) d\tau}{\tau + t + 2} = \pi h(t), \quad -1 < t < 1, \tag{50}$$

where D is a complex constant.

Numerical experiments show that, as a rule, the roots of the Jacobi polynomial (4) for complex α and β lie in the strip $|\operatorname{Re} \tau_k| < 1$ in a small neighborhood of the segment $\tau \in [-1, 1]$ ($|\operatorname{Im} \tau_k| \ll 1$), and these roots approach this segment with the growth of n . Only if the imaginary parts of the singularity exponents are sufficiently large, the roots can lie outside the strip, but in this case the following estimate holds: $|\operatorname{Re} \tau_k| < 1 + \epsilon, 0 < \epsilon \ll 1$. This justifies the utilization of the first method proposed in Subsection 15.6-2 for the calculation of integrals (14), since the (real) pole of the generalized kernel satisfies the inequality $z_* < -1$.

Note that in order to obtain a unique solution of equation (50) for $D \neq 1$, an additional condition should be introduced:

$$\int_{-1}^1 \varphi(\tau) d\tau = 0. \tag{51}$$

For $h(t) \equiv q = \text{const}$, the solution of equation (50) can be written in closed form,

$$\varphi(\tau) = \frac{q}{\pi \sqrt{2(1+D)}} \left[\left(\frac{\theta}{\sqrt{1-x^2} + 1} \right) \left(\frac{x}{1 + \sqrt{1-x^2}} \right)^\theta + \left(\frac{\theta}{\sqrt{1-x^2}} - 1 \right) \left(\frac{x}{1 + \sqrt{1-x^2}} \right)^{-\theta} \right], \tag{52}$$

where $x = (1 + \tau)/2$ and $\theta = \arccos(-D)/\pi$ is complex.

We see that the asymptotic behavior of the solution near the left endpoint of the integration interval has the form $\rho^{-\theta}$ as $\rho \rightarrow 0$ and at the right endpoint the solution has a root singularity. This corresponds to the weight function

$$w(\tau) = (1 - \tau)^{-1/2}(1 + \tau)^{-\theta}. \tag{53}$$

For the construction of a numerical solution of equation (50) we use two methods: (i) integrals (14) are calculated according to the Gauss–Jacobi quadrature formulas (1):

$$h_k(t) = \frac{1}{w(t)} \sum_{k=1}^s W_k k(t, \tau_k) P_k^{(\alpha, \beta)}(\tau_k), \quad k(t, \tau) = \frac{1}{\tau + t + 2}; \quad (54)$$

(ii) the method based on the expansion of this kernel into series (15). In case (ii), we use two types of polynomial approximation: Maclaurin series

$$k(t, \tau) = \sum_{s=0}^N a_s(t) \tau^s, \quad a_s(t) = \frac{(-1)^s}{(t+2)^{s+1}}, \quad (55)$$

and the expansion with respect to Chebyshev polynomials of the first kind $T_n(\tau)$:

$$k(t, \tau) = -\frac{g_0(t)}{2} + \sum_{m=0}^N g_m(t) T_m(\tau), \quad g_j(t) = \frac{2}{N+1} \sum_{k=1}^{N+1} k(t, \tau_k) T_j(\tau_k), \quad (56)$$

$$\tau_k = \cos\left(\frac{2k-1}{N+1} \frac{\pi}{2}\right), \quad T_j(\tau_k) = \cos\left(\frac{2k-1}{N+1} \frac{\pi j}{2}\right).$$

Note that the coefficients $a_s(t)$ in the expansion (15) can be obtained from the coefficients $g_m(t)$ (56) by the method used above for the derivation of (18).

As t_r ($r = 1, \dots, n$) we use the uniform grid

$$t_r = (1 - \delta)[2(r-1)/(n-1) - 1],$$

where $0 < \delta < 1$ is a small parameter that fixes the position of the minimal and the maximal collocation points ($\min t_r = -1 + \delta$, $\max t_r = 1 - \delta$).

Table 12 gives calculation results for equation (50) (with the additional condition (51)) obtained for the following parameter values: $D = 0.5 + 0.5i$ in (72), $q = 1$ in (52), $\delta = 0.2$, $n = 10$ in (12), $s = 10$ in (54), $N = 25$ in (55) and (56). For the given D , we have $\theta = 0.644 + 0.169i$, i.e., the solution has a sufficiently strong singularity near the endpoint $\tau = -1$ ($\operatorname{Re} \beta < -1/2$; see (53)).

TABLE 12
Comparison of exact and numerical solutions of the Bueckner equation (50). N.s. is shorthand notation for “numerical solution”

τ	Exact solution	N.s. based on (54)	N.s. based on (55)	N.s. based on (56)
-0.875	-0.62953 - 0.00548 i	-0.62870 - 0.00542 i	-0.62785 - 0.00758 i	-0.62870 - 0.00541 i
-0.750	-0.36772 + 0.03491 i	-0.36747 + 0.03498 i	-0.36711 + 0.03354 i	-0.36747 + 0.03498 i
-0.625	-0.24804 + 0.04166 i	-0.24791 + 0.04172 i	-0.24765 + 0.04101 i	-0.24791 + 0.04172 i
-0.500	-0.17148 + 0.04234 i	-0.17139 + 0.04239 i	-0.17118 + 0.04198 i	-0.17139 + 0.04239 i
-0.375	-0.11389 + 0.04151 i	-0.11383 + 0.04155 i	-0.11365 + 0.04125 i	-0.11383 + 0.04155 i
-0.250	-0.06598 + 0.04038 i	-0.06592 + 0.04042 i	-0.06577 + 0.04018 i	-0.06592 + 0.04042 i
-0.125	-0.02311 + 0.03937 i	-0.02306 + 0.03940 i	-0.02292 + 0.03921 i	-0.02306 + 0.03940 i
0.000	0.01755 + 0.03865 i	0.01759 + 0.03868 i	0.01771 + 0.03851 i	0.01759 + 0.03868 i
0.125	0.05818 + 0.03832 i	0.05821 + 0.03834 i	0.05833 + 0.03819 i	0.05821 + 0.03834 i
0.250	0.10088 + 0.03848 i	0.10091 + 0.03851 i	0.10102 + 0.03837 i	0.10091 + 0.03851 i
0.375	0.14831 + 0.03930 i	0.14834 + 0.03933 i	0.14845 + 0.03920 i	0.14834 + 0.03933 i
0.500	0.20461 + 0.04108 i	0.20464 + 0.04111 i	0.20475 + 0.04098 i	0.20464 + 0.04111 i
0.625	0.27778 + 0.04447 i	0.27781 + 0.04450 i	0.27792 + 0.04436 i	0.27781 + 0.04450 i
0.750	0.38768 + 0.05118 i	0.38771 + 0.05122 i	0.38783 + 0.05104 i	0.38771 + 0.05122 i
0.875	0.61138 + 0.06818 i	0.61140 + 0.06824 i	0.61151 + 0.06794 i	0.61140 + 0.06824 i

As one can see, all three calculation techniques provide quite good agreement between the numerical and exact analytical solution.

A slightly lower accuracy (to 3 or 4 significant digits) is attained using the approximation (55), while the approximation (56), as well as the solution based on (54), provides a considerably higher accuracy (to 4 or 5 significant digits) for the same N . This is due to a higher accuracy of the approximation (56)—it is closer to the polynomial of best uniform approximation. In particular, the maximum relative error of the approximation (56) in the rectangle $\{\min t_r \leq t \leq \max t_r, -1 \leq \tau \leq 1\}$ for the above calculation parameters is $\max \varepsilon_{\text{chebyshev}} = 0.09\%$, while that of the approximation (55) is $\max \varepsilon_{\text{maclaurin}} = 0.87\%$. The accuracy $\max \varepsilon_{\text{maclaurin}} \approx 0.1\%$ can be attained using the calculations based on (55) by increasing N to $N = 37$, while the approximation (56) provides the same accuracy for $N = 25$. This means that it is not the technique but the accuracy of approximation of the kernel that makes the main effect on the error of calculation of (14) using (22).

It is noteworthy also that the calculation error slightly increases towards the left endpoint of the integration interval. This is due to the reason mentioned in Example 1 and, possibly, to the presence of the second asymptotic term in the expansion near that endpoint.

As one could expect, the quantity δ has a considerable effect on the calculation error. This is because it is δ that controls the position of the minimum and maximum points of collocation, and their positions determine the maximum error of the approximations (55) and (56) for $t_r \in [\min t_r, \max t_r]$. However, a large increase in δ may result in ill-conditioning in the generated algebraic system.

☉ References for Section 15.6: H. F. Bueckner (1966), F. E. Erdogan, G. D. Gupta and T. S. Cook (1973), P. S. Theocaris and N. I. Ioakimidis (1979), M. P. Savruk et al. (1989, 1999), W. H. Press, S. A. Teukolsky et al. (1992), N. G. Moiseyev and G. Ya. Popov (1994), S. M. Belotserkovskii and I. K. Lifanov (1993), A. M. Linkov (2002), A. V. Andreev (2005, 2006).

Chapter 16

Methods for Solving Nonlinear Integral Equations

16.1. Some Definitions and Remarks

16.1-1. Nonlinear Equations with Variable Limit of Integration (Volterra Equations).

Nonlinear Volterra integral equations can be represented in the form

$$\int_a^x K(x, t, y(t)) dt = F(x, y(x)), \quad (1)$$

where $K(x, t, y(t))$ is the kernel of the integral equation and $y(x)$ is the unknown function ($a \leq x \leq b$). All functions in (1) are usually assumed to be continuous.

The form (1) does not cover all possible forms of nonlinear Volterra integral equations; however, it includes the types of nonlinear equations which are most frequently used and studied. A nonlinear integral equation (1) is called a *Volterra integral equation in the Urysohn form*.

In some cases, Eq. (1) can be rewritten in the form

$$\int_a^x K(x, t, y(t)) dt = f(x). \quad (2)$$

Equation (2) is called a *Volterra equation of the first kind in the Urysohn form*. Similarly, the equation

$$y(x) - \int_a^x K(x, t, y(t)) dt = f(x) \quad (3)$$

is called a *Volterra equation of the second kind in the Urysohn form*.

By the substitution $u(x) = y(x) - f(x)$, Eq. (3) can be reduced to the canonical form

$$u(x) = \int_a^x \mathcal{K}(x, t, u(t)) dt, \quad (4)$$

where $\mathcal{K}(x, t, u(t))$ is the kernel* of the canonical integral equation.

The kernel $K(x, t, y(t))$ is said to be *degenerate* if

$$K(x, t, y(t)) = \sum_{k=1}^n g_k(x) h_k(t, y(t)).$$

* There are other ways of reducing Eq. (3) to the form (4) for which the form of the function \mathcal{K} may be different.

If in Eq. (1) the kernel is $K(x, t, y(t)) = Q(x, t)\Phi(t, y(t))$, where $Q(x, t)$ and $\Phi(t, y)$ are known functions, then we obtain the Volterra integral equation in the Hammerstein form:

$$\int_a^x Q(x, t)\Phi(t, y(t)) dt = F(x, y(x)). \quad (5)$$

In some cases Eq. (5) can be rewritten in the form

$$\int_a^x Q(x, t)\Phi(t, y(t)) dt = f(x). \quad (6)$$

Equation (6) is called a *Volterra equation of the first kind in the Hammerstein form*. Similarly, an equation of the form

$$y(x) - \int_a^x Q(x, t)\Phi(t, y(t)) dt = f(x), \quad (7)$$

is called a *Volterra equation of the second kind in the Hammerstein form*.

It is possible to reduce Eq. (7) to the canonical form

$$u(x) = \int_a^x Q(x, t)\Phi_*(t, u(t)) dt, \quad (8)$$

where $u(x) = y(x) - f(x)$.

Remark 1. Since a Volterra equation in the Hammerstein form is a special case of a Volterra equation in the Urysohn form, the methods discussed below for the latter are certainly applicable to the former.

Remark 2. Some other types of nonlinear integral equations with variable limits of integration are considered in Chapters 5–6.

16.1-2. Nonlinear Equations with Constant Integration Limits (Urysohn Equations).

Nonlinear integral equations with constant integration limits can be represented in the form

$$\int_a^b K(x, t, y(t)) dt = F(x, y(x)), \quad \alpha \leq x \leq \beta, \quad (9)$$

where $K(x, t, y(t))$ is the kernel of the integral equation and $y(x)$ is the unknown function. Usually, all functions in (9) are assumed to be continuous and the case of $\alpha = a$ and $\beta = b$ is considered.

The form (9) does not cover all possible forms of nonlinear integral equations with constant integration limits; however, just as the form (1) for the Volterra equations, it includes the most frequently used and most studied types of these equations. A nonlinear integral equation (9) with constant limits of integration is called an *integral equation of the Urysohn type*.

If Eq. (9) can be rewritten in the form

$$\int_a^b K(x, t, y(t)) dt = f(x), \quad (10)$$

then (10) is called an *Urysohn equation of the first kind*. Similarly, the equation

$$y(x) - \int_a^b K(x, t, y(t)) dt = f(x) \quad (11)$$

is called an *Urysohn equation of the second kind*.

An Urysohn equation of the second kind can be rewritten in the canonical form

$$u(x) = \int_a^b \mathcal{K}(x, t, u(t)) dt. \quad (12)$$

Remark 3. Conditions for existence and uniqueness of the solution of an Urysohn equation are discussed below in Section 16.6.

If in Eq. (9) the kernel is $K(x, t, y(t)) = Q(x, t)\Phi(t, y(t))$, and $Q(x, t)$ and $\Phi(t, y)$ are given functions, then we obtain an *integral equation of the Hammerstein type*:

$$\int_a^b Q(x, t) \Phi(t, y(t)) dt = F(x, y(x)), \quad (13)$$

where, as usual, all functions in the equation are assumed to be continuous.

If Eq. (13) can be rewritten in the form

$$\int_a^b Q(x, t) \Phi(t, y(t)) dt = f(x), \quad (14)$$

then (14) is called a *Hammerstein equation of the first kind*. Similarly, an equation of the form

$$y(x) - \int_a^b Q(x, t) \Phi(t, y(t)) dt = f(x) \quad (15)$$

is called a *Hammerstein equation of the second kind*.

A Hammerstein equation of the second kind can be rewritten in the *canonical form*

$$u(x) = \int_a^b Q(x, t) \Phi_*(t, u(t)) dt. \quad (16)$$

The existence of the canonical forms (4), (8), (12), and (16) means that the distinction between the inhomogeneous and homogeneous nonlinear integral equations is unessential, unlike the case of linear equations. Another specific feature of a nonlinear equation is that it frequently has several solutions.

Remark 4. Since a Hammerstein equation is a special case of an Urysohn equation, the methods discussed below for the latter are certainly applicable to the former.

Remark 5. Some other types of nonlinear integral equations with constant limits of integration are considered in Chapters 7–8.

16.1-3. Some Special Features of Nonlinear Integral Equations.

Even simplest nonlinear equations, such as those of Volterra or Hammerstein, exhibit some new phenomena characteristic only of nonlinear equations and having no analogues in the theory of linear integral equations.

Example 1. Consider the Volterra integral equation with power nonlinearity

$$y(x) = a \int_0^x y^n(t) dt + b, \quad a > 0, b \geq 0, n > 0. \quad (17)$$

By the differentiation in x , this equation is reduced to the Cauchy problem for the first-order ODE:

$$y'_x = ay^n, \quad y(0) = b. \quad (18)$$

The solution of problem (18) depends on the parameter n and, for $b > 0$, is defined by the formulas

$$y(x) = \begin{cases} [b^{1-n} + a(1-n)x]^{\frac{1}{1-n}} & \text{if } 0 < n < 1, \\ be^{ax} & \text{if } n = 1, \\ [b^{1-n} - a(n-1)x]^{\frac{1}{1-n}} & \text{if } n > 1. \end{cases} \quad (19)$$

It is easy to see that for $0 < n \leq 1$ the solution exists for all $x \geq 0$, and for $0 < n < 1$ and large x , the function y has power growth, while for $n = 1$ it has exponential growth. For $n > 1$, a continuous solution exists only on the finite interval

$$0 \leq x < x_* = \frac{b^{1-n}}{a(n-1)}.$$

Such a situation is not observed for linear Volterra equations.

Now consider the limit case $b = 0$. Then for any $0 < n < \infty$, $0 < a < \infty$, equation (17) has the trivial solution $y(x) \equiv 0$. Moreover, for $0 < n < 1$, $0 < a < \infty$, equation (17) admits another real solution, $y(x) = [a(1-n)x]^{\frac{1}{1-n}}$.

Example 2. Consider the Hammerstein integral equation with a quadratic nonlinearity

$$y(x) = \lambda \int_0^1 x^2 t y^2(t) dt, \quad (20)$$

where λ is a free parameter.

Setting

$$A = \int_0^1 t y^2(t) dt, \quad (21)$$

let us represent equation (20) in the form

$$y(x) = A\lambda x^2.$$

Substituting this expression into (21), we obtain a quadratic equation for the determination of the constant A :

$$A = \frac{1}{6} A^2 \lambda^2. \quad (22)$$

Its solutions are $A_1 = 0$ and $A_2 = 6\lambda^{-2}$. Therefore, the original integral equation (20) has two solutions for any $\lambda \neq 0$:

$$y_1(x) \equiv 0, \quad y_2(x) = \frac{6}{\lambda} x^2.$$

Note that the linear homogeneous integral equation

$$y(x) = \lambda \int_0^1 x^2 t y(t) dt \quad (23)$$

with the same kernel $K(x, t) = x^2 t$ has a nontrivial solution only for a single value of λ , namely, $\lambda = 4$, which is a characteristic value of the kernel $K(x, t)$. Therefore, if we follow the terminology of linear equations and say that λ is a characteristic value of a nonlinear equation if this equation has a nontrivial solution for that λ , it turns out that equation (23) has infinite intervals of characteristic values $(-\infty, 0)$ and $(0, \infty)$.

Example 3. Consider another integral equation of Hammerstein's type with a quadratic nonlinearity

$$y(x) = \lambda \int_0^1 y^2(t) dt + 1. \quad (24)$$

This equation can be written as

$$y(x) = A\lambda + 1, \quad (25)$$

where

$$A = \int_0^1 y^2(t) dt. \quad (26)$$

Substituting (25) into (26), we obtain the quadratic equation

$$\lambda^2 A^2 + (2\lambda - 1)A + 1 = 0$$

with the roots

$$A = \frac{1 - 2\lambda \pm \sqrt{1 - 4\lambda}}{2\lambda^2}.$$

Thus, equation (24) has real solutions only for $\lambda \leq 1/4$. It has two solutions for $\lambda < 1/4$ and one solution for $\lambda = 1/4$ (for $\lambda = 0$ there is one bounded solution $y(x) = 1$).

The corresponding equation with no free term

$$y(x) = \lambda \int_0^1 ty^2(t) dt,$$

for any $\lambda \neq 0$, admits the nontrivial solution $y(x) = 1/\lambda$. Obviously, this does not mean that equation (24) with a free term had infinitely many solutions.

Example 4. Now consider an integral equation of Hammerstein's type with a transcendental nonlinearity

$$y(x) = \lambda \int_0^1 f(x)g(t) \sin\left(\frac{y(t)}{f(t)}\right) y(t) dt. \quad (27)$$

Its solutions are sought in the form $y(x) = Af(x)$, where the constant A is determined from the transcendental equation*

$$1 = \lambda\sigma \sin A, \quad \sigma = \int_0^1 f(t)g(t) dt. \quad (28)$$

For $|\lambda| < 1/|\sigma|$, equation (28), and therefore equation (27), has no real solutions (the case $\sigma = 0$ is included).

For any λ satisfying the inequality $|\lambda| > 1/|\sigma|$, equation (28), and therefore equation (27), has infinitely many real solutions.

⊙ References for Section 16.1: N. S. Smirnov (1951), M. A. Krasnosel'skii (1964), M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), M. L. Krasnov (1975), P. P. Zabreyko, A. I. Koshelev, et al. (1975), F. G. Tricomi (1985), A. F. Verlan' and V. S. Sizikov (1986).

16.2. Exact Methods for Nonlinear Equations with Variable Limit of Integration

16.2-1. Method of Integral Transforms.

Consider a Volterra integral equation with quadratic nonlinearity

$$\mu y(x) - \lambda \int_0^x y(x-t)y(t) dt = f(x). \quad (1)$$

This equation can be solved using the Laplace transform. In doing so, one applies the convolution theorem (see Section 9.2) to obtain a quadratic equation for the transform $\tilde{y}(p) = \mathfrak{L}\{y(x)\}$:

$$\mu\tilde{y}(p) - \lambda\tilde{y}^2(p) = \tilde{f}(p).$$

This implies

$$\tilde{y}(p) = \frac{\mu \pm \sqrt{\mu^2 - 4\lambda\tilde{f}(p)}}{2\lambda}. \quad (2)$$

The inverse Laplace transform $y(x) = \mathfrak{L}^{-1}\{\tilde{y}(p)\}$, if it exists, is a solution to Eq. (1). Note that for the two different signs in formula (2), there are two corresponding solutions of the original equation.

Example. Consider the integral equation

$$\int_0^x y(x-t)y(t) dt = Ax^m, \quad m > -1.$$

Applying the Laplace transform to this equation and taking into account the relation $\mathfrak{L}\{x^m\} = \Gamma(m+1)p^{-m-1}$, we obtain

$$\tilde{y}^2(p) = A\Gamma(m+1)p^{-m-1},$$

where $\Gamma(m)$ is the Gamma function. Taking the square root of both sides of the equation, we obtain

$$\tilde{y}(p) = \pm \sqrt{A\Gamma(m+1)} p^{-\frac{m+1}{2}}.$$

Applying the Laplace inversion formula, we obtain two solutions to the original integral equation

$$y_1(x) = -\frac{\sqrt{A\Gamma(m+1)}}{\Gamma\left(\frac{m+1}{2}\right)} x^{\frac{m-1}{2}}, \quad y_2(x) = \frac{\sqrt{A\Gamma(m+1)}}{\Gamma\left(\frac{m+1}{2}\right)} x^{\frac{m-1}{2}}.$$

* The trivial solution corresponding to $A = 0$ is not taken into account.

16.2-2. Method of Differentiation for Nonlinear Equations with Degenerate Kernel.

Sometimes, differentiation (possibly multiple) of a nonlinear integral equation with subsequent elimination of the integral term using the original equation makes it possible to reduce this equation to a nonlinear ordinary differential equation. Listed below are some equations of this type.

1°. The equation

$$y(x) + \int_a^x f(t, y(t)) dt = g(x) \quad (3)$$

can be reduced by differentiation to the nonlinear first-order equation

$$y'_x + f(x, y) - g'_x(x) = 0 \quad (4)$$

with the initial condition $y(a) = g(a)$.

2°. The equation

$$y(x) + \int_a^x (x-t)f(t, y(t)) dt = g(x) \quad (5)$$

can be reduced by double differentiation (with the subsequent elimination of the integral term using the original equation) to the nonlinear second-order equation:

$$y''_{xx} + f(x, y) - g''_{xx}(x) = 0. \quad (6)$$

The initial conditions for the function $y = y(x)$ have the form

$$y(a) = g(a), \quad y'_x(a) = g'_x(a).$$

3°. The equation

$$y(x) + \int_a^x e^{\lambda(x-t)} f(t, y(t)) dt = g(x) \quad (7)$$

can be reduced by differentiation to the nonlinear first-order equation

$$y'_x + f(x, y) - \lambda y + \lambda g(x) - g'_x(x) = 0. \quad (8)$$

The desired function $y = y(x)$ must satisfy the initial condition $y(a) = g(a)$.

Remark 1. A considerable number of exact solutions to the ordinary differential equations (4), (6), and (8) for various functions $f(x, y)$ and $g(x)$ can be found in the book by Polyanin and Zaitsev (2003).

4°. Equations of the form

$$\begin{aligned} y(x) + \int_a^x \cosh[\lambda(x-t)] f(t, y(t)) dt &= g(x), \\ y(x) + \int_a^x \sinh[\lambda(x-t)] f(t, y(t)) dt &= g(x), \\ y(x) + \int_a^x \cos[\lambda(x-t)] f(t, y(t)) dt &= g(x), \\ y(x) + \int_a^x \sin[\lambda(x-t)] f(t, y(t)) dt &= g(x) \end{aligned}$$

can also be reduced to second-order ordinary differential equations by double differentiation. For these equations, see Section 6.8 in the first part of the book (Eqs. 20, 21, 22, and 23, respectively).

5°. Consider the nonlinear Volterra equation of the second kind with the general degenerate kernel

$$y(x) - \sum_{m=1}^n \varphi_m(x) \int_a^x f_m(t, y(t)) dt = g(x). \quad (9)$$

Let us introduce the notation

$$w_j(x) = \int_a^x f_j(t, y(t)) dt, \quad j = 1, \dots, n, \quad (10)$$

and rewrite Eq. (9) as follows:

$$y(x) = g(x) + \sum_{m=1}^n \varphi_m(x) w_m(x). \quad (11)$$

On differentiating the expressions (10) with regard to formula (11), we arrive at the following system of nonlinear differential equations for the functions $w_j = w_j(x)$:

$$w'_j = f_j \left(x, g(x) + \sum_{m=1}^n \varphi_m(x) w_m \right), \quad j = 1, \dots, n,$$

with the initial conditions

$$w_j(a) = 0, \quad j = 1, \dots, n.$$

Once a solution of this system is found, the corresponding solution of the original integral equation (9) is defined by formula (11).

Remark 2. Equations (3), (5), and (7) are special cases of equation (9). The equations of Item 4° can be reduced to (9) using hyperbolic and trigonometric formulas (see the addition formulas in Supplements 1.4-7 and 1.2-7, respectively).

© References for Section 16.2: M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), A. F. Verlan' and V. S. Sizikov (1986), A. D. Polyaniin and A. V. Manzhirov (1998).

16.3. Approximate and Numerical Methods for Nonlinear Equations with Variable Limit of Integration

16.3-1. Successive Approximation Method.

1°. In many cases, the successive approximation method can be successfully applied to solve various types of integral equations. The principles of constructing the iteration process are the same as in the case of linear equations. For Volterra equations of the second kind in the Urysohn form

$$y(x) - \int_a^x K(x, t, y(t)) dt = f(x), \quad a \leq x \leq b, \quad (1)$$

the corresponding recursive expression has the form

$$y_{n+1}(x) = f(x) + \int_a^x K(x, t, y_n(t)) dt, \quad n = 0, 1, 2, \dots \quad (2)$$

It is customary to take the initial approximation either in the form $y_0(x) \equiv 0$ or in the form $y_0(x) = f(x)$.

In contrast to the case of linear equations, the successive approximation method has a smaller domain of convergence. Let us present the convergence conditions for the iteration process (2), which are simultaneously the existence conditions for a solution of Eq. (1). To be specific, we assume that $y_0(x) = f(x)$.

If for any z_1 and z_2 the relations

$$|K(x, t, z_1) - K(x, t, z_2)| \leq \varphi(x, t)|z_1 - z_2|$$

and

$$\left| \int_a^x K(x, t, f(t)) dt \right| \leq \psi(x)$$

hold, with

$$\int_a^x \psi^2(t) dt \leq N^2, \quad \int_a^b \int_a^x \varphi^2(x, t) dt dx \leq M^2,$$

where N and M are some constants, then the successive approximations converge to a unique solution of Eq. (1) almost everywhere absolutely and uniformly.

Example 1. Let us apply the successive approximation method to solve the equation

$$y(x) = \int_0^x \frac{1 + y^2(t)}{1 + t^2} dt.$$

If $y_0(x) \equiv 0$, then

$$y_1(x) = \int_0^x \frac{dt}{1 + t^2} = \arctan x,$$

$$y_2(x) = \int_0^x \frac{1 + \arctan^2 t}{1 + t^2} dt = \arctan x + \frac{1}{3} \arctan^3 x,$$

$$y_3(x) = \int_0^x \frac{1 + \arctan t + \frac{1}{3} \arctan^3 t}{1 + t^2} dt = \arctan x + \frac{1}{3} \arctan^3 x + \frac{2}{3 \cdot 5} \arctan^5 x + \frac{1}{7 \cdot 9} \arctan^7 x.$$

On continuing this process, we can observe that $y_n(x) \rightarrow \tan(\arctan x) = x$ as $n \rightarrow \infty$, i.e., $y(x) = x$. This result is validated by substituting it into the original equation.

Example 2. For the nonlinear equation

$$y(x) = \int_0^x [ty^2(t) - 1] dt,$$

we wish to obtain the first three approximations. If we set $y_0(x) = 0$, then

$$y_1(x) = \int_0^x (-1) dt = -x,$$

$$y_2(x) = \int_0^x (t^3 - 1) dt = -x + \frac{1}{4}x^4,$$

$$y_3(x) = \int_0^x \left[t \left(\frac{1}{16}t^8 - \frac{1}{2}t^5 + t^2 \right) - 1 \right] dt = -x + \frac{1}{4}x^4 - \frac{1}{14}x^7 + \frac{1}{160}x^{10}.$$

2°. Suppose that in the nonlinear Volterra equation

$$y(x) = \int_0^x K(x, t, y(t)) dt,$$

the function $K(x, t, y)$ and its partial derivative $K'_y(x, t, y)$ are continuous in the domain $x, t \geq 0$, $-\infty < y < \infty$, and the following inequality holds:

$$|K(x, t, y)| \leq \varphi(y),$$

where $\varphi(y)$ is a nondecreasing function on the half-line $[0, \infty)$. If the Cauchy problem for the differential equation

$$u'_x = \varphi(|u|), \quad u(0) = 0$$

has a solution on the interval $[0, \omega]$, then the above Volterra equation has a solution on $[0, \omega]$. For $y_0(x) \equiv 0$ as the initial function, the successive approximations

$$y_n(x) = \int_0^x K(x, t, y_{n-1}(t)) dt \quad (n = 1, 2, \dots)$$

are uniformly convergent on $[0, \omega]$ to a solution of the Volterra equation.

Note that all approximations do not abandon the domain $-u(x) \leq y_n(x) \leq u(x)$ and satisfy the inequality

$$|y_n(x) - y_{n-1}(x)| \leq \frac{M}{L} \frac{(Lt)^n}{n!} \quad (n = 1, 2, \dots),$$

where L and M are constants such that

$$\begin{aligned} |K(x, t, 0)| &\leq M \quad \text{for } 0 \leq x, t \leq \omega, \\ |K(x, t, y_1) - K(x, t, y_2)| &\leq L|y_1 - y_2| \quad \text{for } 0 \leq x, t \leq \omega, \quad -u(x) \leq y_1, y_2 \leq u(x). \end{aligned}$$

3°. The successive approximation method can be applied to solve other forms of nonlinear equations, for instance, equations of the form

$$y(x) = F\left(x, \int_a^x K(x, t)y(t) dt\right)$$

solved for $y(x)$ in which the integral has x as the upper integration limit. This makes it possible to obtain a numerical solution by applying small steps with respect to x and by linearization at each step, which usually provides the uniqueness of the result of the iterations for an arbitrary initial approximation.

4°. The initial approximation has a substantial effect on the number of iterations required to obtain the result with a prescribed accuracy. Therefore, when choosing this approximation, some additional arguments are usually applied. Namely, for the equation

$$Ay(x) - \int_0^x Q(x-t)\Phi(y(t)) dt = f(x),$$

where A is a constant, a good initial approximation $y_0(x)$ can sometimes be found from the solution of the following (in general, transcendental) equation for $\tilde{y}_0(p)$:

$$A\tilde{y}_0(p) - \tilde{Q}(p)\Phi(\tilde{y}_0(p)) = \tilde{f}(p),$$

where $\tilde{y}_0(p)$, $\tilde{Q}(p)$, and $\tilde{f}(p)$ are the Laplace transforms of the respective functions. If $\tilde{y}_0(p)$ is defined, then the initial approximation can be found by applying the Laplace inversion formula: $y_0(x) = \mathfrak{L}^{-1}\{\tilde{y}_0(p)\}$.

16.3-2. Newton–Kantorovich Method.

A merit of the iteration methods when applied to Volterra linear equations of the second kind is their unconditional convergence under weak restrictions on the kernel and the right-hand side. When solving nonlinear equations, the applicability domain of the method of simple iterations is smaller, and if the process is still convergent, then, in many cases, the rate of convergence can be

very low. An effective method that makes it possible to overcome the indicated complications is the Newton–Kantorovich method. The main objective of this method is the solution of nonlinear integral equations of the second kind with constant limits of integration. Nevertheless, this method is useful in the solution of many problems for the Volterra equations and makes it possible to significantly increase the rate of convergence compared with the successive approximation method.

Let us apply the Newton–Kantorovich method to solve a Volterra equation of the second kind in the Urysohn form

$$y(x) = f(x) + \int_a^x K(x, t, y(t)) dt. \quad (3)$$

We obtain the following iteration process:

$$y_n(x) = y_{n-1}(x) + \varphi_{n-1}(x), \quad n = 1, 2, \dots, \quad (4)$$

$$\varphi_{n-1}(x) = \varepsilon_{n-1}(x) + \int_a^x K'_y(x, t, y_{n-1}(t)) \varphi_{n-1}(t) dt, \quad (5)$$

$$\varepsilon_{n-1}(x) = f(x) + \int_a^x K(x, t, y_{n-1}(t)) dt - y_{n-1}(x). \quad (6)$$

The algorithm is based on the solution of the linear integral equation (5) for the correction $\varphi_{n-1}(x)$ with the kernel and right-hand side that vary from step to step. This process has a high rate of convergence, but it is rather complicated because we must solve a new equation at each step of iteration. To simplify the problem, we can replace Eq. (5) by the equation

$$\varphi_{n-1}(x) = \varepsilon_{n-1}(x) + \int_a^x K'_y(x, t, y_0(t)) \varphi_{n-1}(t) dt \quad (7)$$

or by the equation

$$\varphi_{n-1}(x) = \varepsilon_{n-1}(x) + \int_a^x K'_y(x, t, y_m(t)) \varphi_{n-1}(t) dt, \quad (8)$$

whose kernels do not vary. In Eq. (8), m is fixed and satisfies the condition $m < n - 1$.

It is reasonable to apply Eq. (7) with an appropriately chosen initial approximation. Otherwise we can stop at some m th approximation and, beginning with this approximation, apply the simplified equation (8). The iteration process thus obtained is the modified Newton–Kantorovich method. In principle, it converges somewhat slower than the original process (4)–(6); however, it is not so cumbersome in the calculations.

Example 3. Let us apply the Newton–Kantorovich method to solve the equation

$$y(x) = \int_0^x [ty^2(t) - 1] dt.$$

The derivative of the integrand with respect to y has the form

$$K'_y(t, y(t)) = 2ty(t).$$

For the zero approximation we take $y_0(x) \equiv 0$. According to (5) and (6) we obtain $\varphi_0(x) = -x$ and $y_1(x) = -x$. Furthermore, $y_2(x) = y_1(x) + \varphi_1(x)$. By (6) we have

$$\varepsilon_1(x) = \int_0^x [t(-t)^2 - 1] dt + x = \frac{1}{4}x^4.$$

The equation for the correction has the form

$$\varphi_1(x) = -2 \int_0^x t^2 \varphi_1(t) dt + \frac{1}{4}x^4$$

and can be solved by any of the known methods for Volterra linear equations of the second kind. In the case under consideration, we apply the successive approximation method, which leads to the following results (the number of the step is indicated in the superscript):

$$\begin{aligned}\varphi_1^{(0)} &= \frac{1}{4}x^4, \\ \varphi_1^{(1)} &= \frac{1}{4}x^4 - 2 \int_0^x \frac{1}{4}t^6 dt = \frac{1}{4}x^4 - \frac{1}{14}x^7, \\ \varphi_1^{(2)} &= \frac{1}{4}x^4 - 2 \int_0^x t^2 \left(\frac{1}{4}t^4 - \frac{1}{14}x^7 \right) dt = \frac{1}{4}x^4 - \frac{1}{14}x^7 + \frac{1}{70}x^{10}.\end{aligned}$$

We restrict ourselves to the second approximation and obtain

$$y_2(x) = -x + \frac{1}{4}x^4 - \frac{1}{14}x^7 + \frac{1}{70}x^{10}$$

and then pass to the third iteration step of the Newton–Kantorovich method:

$$\begin{aligned}y_3(x) &= y_2(x) + \varphi_2(x), \\ \varepsilon_2(x) &= \frac{1}{160}x^{10} - \frac{1}{1820}x^{13} - \frac{1}{7840}x^{16} + \frac{1}{9340}x^{19} + \frac{1}{107800}x^{22}, \\ \varphi_2(x) &= \varepsilon_2(x) + 2 \int_0^x t \left(-t + \frac{1}{4}t^4 - \frac{1}{14}t^7 + \frac{1}{70}t^{10} \right) \varphi_2(t) dt.\end{aligned}$$

When solving the last equation, we restrict ourselves to the zero approximation and obtain

$$y_3(x) = -x + \frac{1}{4}x^4 - \frac{1}{14}x^7 + \frac{23}{112}x^{10} - \frac{1}{1820}x^{13} - \frac{1}{7840}x^{16} + \frac{1}{9340}x^{19} + \frac{1}{107800}x^{22}.$$

The application of the successive approximation method to the original equation leads to the same result at the fourth step.

As usual, in the numerical solution the integral is replaced by a quadrature formula. The main difficulty of the implementation of the method in this case is in evaluating the derivative of the kernel. The problem can be simplified if the kernel is given as an analytic expression that can be differentiated in the analytic form. However, if the kernel is given by a table, then the evaluation must be performed numerically.

16.3-3. Collocation Method.

When applied to the solution of a Volterra equation of the first kind in the Urysohn form

$$\int_a^x K(x, t, y(t)) dt = f(x), \quad a \leq x \leq b, \quad (9)$$

the *collocation method* is as follows. The interval $[a, b]$ is divided into N parts on each of which the desired solution can be presented by a function of a certain form

$$\tilde{y}(x) = \Phi(x, A_1, \dots, A_m), \quad (10)$$

involving free parameters $A_i, i = 1, \dots, m$.

On the $(k+1)$ st part $x_k \leq x \leq x_{k+1}$, where $k = 0, 1, \dots, N-1$, the solution can be written in the form

$$\int_{x_k}^x K(x, t, \tilde{y}(t)) dt = f(x) - \Psi_k(x), \quad (11)$$

where the integral

$$\Psi_k(x) = \int_a^{x_k} K(x, t, \tilde{y}(t)) dt \quad (12)$$

can always be calculated for the approximate solution $\tilde{y}(x)$, which is known on the interval $a \leq x \leq x_k$ and was previously obtained for $k-1$ parts. The initial value $y(a)$ of the desired solution can be found by an auxiliary method or is assumed to be given.

To solve Eq. (11), representation (10) is applied, and the free parameters A_i ($i = 1, \dots, m$) can be defined from the condition that the residuals vanish:

$$\varepsilon(A_i, x_{k,j}) = \int_{x_k}^{x_{k,j}} K(x_{k,j}, t, \Phi(t, A_1, \dots, A_m)) dt - f(x_{k,j}) - \Psi_k(x_{k,j}), \quad (13)$$

where the $x_{k,j}$ ($j = 1, \dots, m$) are the nodes that correspond to the partition of the interval $[x_k, x_{k+1}]$ into m parts (subintervals). System (13) is a system of m equations for A_1, \dots, A_m .

For convenience of the calculations, it is reasonable to present the desired solution on any part as a polynomial

$$\tilde{y}(x) = \sum_{i=1}^m A_i \varphi_i(x), \quad (14)$$

where the $\varphi_i(x)$ are linearly independent coordinate functions. For the functions $\varphi_i(x)$, power and trigonometric polynomials are frequently used; for instance, $\varphi_i(x) = x^{i-1}$.

In applications, the concrete form of the functions $\varphi_i(x)$ in formula (14), as well as the form of the functions Φ in (10), can sometimes be given on the basis of physical reasoning or defined by the structure of the solution of a simpler model equation.

16.3-4. Quadrature Method.

To solve a nonlinear Volterra equation, we can apply the method based on the use of quadrature formulas. The procedure of constructing the approximate system of equations is the same as in the linear case (see Subsection 11.10-1).

1°. We consider the nonlinear Volterra equation of the second kind in the Urysohn form

$$y(x) - \int_a^x K(x, t, y(t)) dt = f(x) \quad (15)$$

on an interval $a \leq x \leq b$. Assume that $K(x, t, y(t))$ and $f(x)$ are continuous functions.

From Eq. (15) we find that $y(a) = f(a)$. Let us choose a constant integration step h and consider the discrete set of points $x_i = a + h(i-1)$, where $i = 1, \dots, n$. For $x = x_i$, Eq. (15) becomes

$$y(x_i) - \int_a^{x_i} K(x_i, t, y(t)) dt = f(x_i). \quad (16)$$

Applying the quadrature formula (see Subsection 10.7-1) to the integral in (16), choosing x_j ($j = 1, \dots, i$) to be the nodes in t , and neglecting the truncation error, we arrive at the following system of nonlinear algebraic (or transcendental) equations:

$$y_1 = f_1, \quad y_i - \sum_{j=1}^i A_{ij} K_{ij}(y_j) = f_i, \quad i = 2, \dots, n, \quad (17)$$

where the A_{ij} are the coefficients of the quadrature formula on the interval $[a, x_i]$, the y_i are the approximate values of the solution $y(x)$ at the nodes x_i , $f_i = f(x_i)$, and $K_{ij}(y_j) = K(x_i, t_j, y_j)$.

Relations (17) can be rewritten as a sequence of recursive nonlinear equations,

$$y_1 = f_1, \quad y_i - A_{ii} K_{ii}(y_i) = f_i + \sum_{j=1}^{i-1} A_{ij} K_{ij}(y_j), \quad i = 2, \dots, n, \quad (18)$$

for the approximate values of the desired solution at the nodes.

2°. When applied to the Volterra equation of the second kind in the Hammerstein form

$$y(x) - \int_a^x Q(x, t) \Phi(t, y(t)) dt = f(x), \quad (19)$$

the main relations of the quadrature method have the form ($x_1 = a$)

$$y_1 = f_1, \quad y_i - \sum_{j=1}^i A_{ij} Q_{ij} \Phi_j(y_j) = f_i, \quad i = 2, \dots, n, \quad (20)$$

where $Q_{ij} = Q(x_i, t_j)$ and $\Phi_j(y_j) = \Phi(t_j, y_j)$. These relations lead to the sequence of nonlinear recursive equations

$$y_1 = f_1, \quad y_i - A_{ii} Q_{ii} \Phi_i(y_i) = f_i + \sum_{j=1}^{i-1} A_{ij} Q_{ij} \Phi_j(y_j), \quad i = 2, \dots, n, \quad (21)$$

whose solutions give approximate values of the desired function.

Example 4. In the solution of the equation

$$y(x) - \int_0^x e^{-(x-t)} y^2(t) dt = e^{-x}, \quad 0 \leq x \leq 0.1,$$

where $Q(x, t) = e^{-(x-t)}$, $\Phi(t, y(t)) = y^2(t)$, and $f(x) = e^{-x}$, the approximate expression has the form

$$y(x_i) - \int_0^{x_i} e^{-(x_i-t)} y^2(t) dt = e^{-x_i}.$$

On applying the trapezoidal rule to evaluate the integral (with step $h = 0.02$) and finding the solution at the nodes $x_i = 0, 0.02, 0.04, 0.06, 0.08, 0.1$, we obtain, according to (21), the following system of computational relations:

$$y_1 = f_1, \quad y_i - 0.01 Q_{ii} y_i^2 = f_i + \sum_{j=1}^{i-1} 0.02 Q_{ij} y_j^2, \quad i = 2, \dots, 6.$$

Thus, to find an approximate solution, we must solve a quadratic equation for each value y_i , which makes it possible to write out the answer

$$y_i = 50 \pm 50 \left[1 - 0.04 \left(f_i + \sum_{j=1}^{i-1} 0.02 Q_{ij} y_j^2 \right) \right]^{1/2}, \quad i = 2, \dots, 6.$$

⊙ References for Section 16.3: M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), P. P. Zabreyko, A. I. Koshelev, et al. (1975), A. F. Verlan' and V. S. Sizikov (1986).

16.4. Exact Methods for Nonlinear Equations with Constant Integration Limits

16.4-1. Nonlinear Equations with Degenerate Kernels.

1°. Consider a Hammerstein equation of the second kind in the canonical form

$$y(x) = \int_a^b Q(x, t) \Phi(t, y(t)) dt, \quad (1)$$

where $Q(x, t)$ and $\Phi(t, y)$ are given functions and $y(x)$ is the unknown function.

Let the kernel $Q(x, t)$ be degenerate, i.e.,

$$Q(x, t) = \sum_{k=1}^m g_k(x)h_k(t). \quad (2)$$

In this case Eq. (1) becomes

$$y(x) = \sum_{k=1}^m g_k(x) \int_a^b h_k(t) \Phi(t, y(t)) dt. \quad (3)$$

We write

$$A_k = \int_a^b h_k(t) \Phi(t, y(t)) dt, \quad k = 1, \dots, m, \quad (4)$$

where the constants A_k are yet unknown. Then it follows from (3) that

$$y(x) = \sum_{k=1}^m A_k g_k(x). \quad (5)$$

On substituting the expression (5) for $y(x)$ into relations (4), we obtain (in the general case) m transcendental equations of the form

$$A_k = \Psi_k(A_1, \dots, A_m), \quad k = 1, \dots, m, \quad (6)$$

which contain m unknown numbers A_1, \dots, A_m .

For the case in which $\Phi(t, y)$ is a polynomial in y , i.e.,

$$\Phi(t, y) = p_0(t) + p_1(t)y + \dots + p_n(t)y^n, \quad (7)$$

where $p_0(t), \dots, p_n(t)$ are, for instance, continuous functions of t on the interval $[a, b]$, system (6) becomes a system of nonlinear algebraic equations for A_1, \dots, A_m .

The number of solutions of the integral equation (3) is equal to the number of solutions of system (6). Each solution of system (6) generates a solution (5) of the integral equation.

2°. Consider the Urysohn equation of the second kind with the simplified degenerate kernel of the following form:

$$y(x) + \int_a^b \left\{ \sum_{k=1}^n g_k(x) f_k(t, y(t)) \right\} dt = h(x). \quad (8)$$

Its solution has the form

$$y(x) = h(x) + \sum_{k=1}^n \lambda_k g_k(x), \quad (9)$$

where the constants λ_k can be defined by solving the algebraic (or transcendental) system of equations

$$\lambda_m + \int_a^b f_m \left(t, h(t) + \sum_{k=1}^n \lambda_k g_k(t) \right) dt = 0, \quad m = 1, \dots, n. \quad (10)$$

To different roots of this system, there are different corresponding solutions of the nonlinear integral equation. It may happen that (real) solutions are absent.

A solution of an Urysohn equation of the second kind with degenerate kernel in the general form

$$f(x, y(x)) + \int_a^b \left\{ \sum_{k=1}^n g_k(x, y(x)) h_k(t, y(t)) \right\} dt = 0 \quad (11)$$

can be represented in the implicit form

$$f(x, y(x)) + \sum_{k=1}^n \lambda_k g_k(x, y(x)) = 0, \quad (12)$$

where the parameters λ_k are determined from the system of algebraic (or transcendental) equations:

$$\begin{aligned} \lambda_k - H_k(\vec{\lambda}) &= 0, \quad k = 1, \dots, n, \\ H_k(\vec{\lambda}) &= \int_a^b h_k(t, y(t)) dt, \quad \vec{\lambda} = \{\lambda_1, \dots, \lambda_n\}. \end{aligned} \quad (13)$$

Into system (13), we must substitute the function $y(x) = y(x, \vec{\lambda})$, which can be obtained by solving Eq. (12).

The number of solutions of the integral equation is defined by the number of solutions obtained from (12) and (13). It can occur that there is no solution.

Example 1. Let us solve the integral equation

$$y(x) = \lambda \int_0^1 xty^3(t) dt \quad (14)$$

with parameter λ . We write

$$A = \int_0^1 ty^3(t) dt. \quad (15)$$

In this case, it follows from (14) that

$$y(x) = \lambda Ax. \quad (16)$$

On substituting $y(x)$ in the form (16) into relation (15), we obtain

$$A = \int_0^1 t\lambda^3 A^3 t^3 dt.$$

Hence,

$$A = \frac{1}{5} \lambda^3 A^3. \quad (17)$$

For $\lambda > 0$, Eq. (17) has three solutions:

$$A_1 = 0, \quad A_2 = \left(\frac{5}{\lambda^3}\right)^{1/2}, \quad A_3 = -\left(\frac{5}{\lambda^3}\right)^{1/2}.$$

Hence, the integral equation (14) also has three solutions for any $\lambda > 0$:

$$y_1(x) \equiv 0, \quad y_2(x) = \left(\frac{5}{\lambda^3}\right)^{1/2} x, \quad y_3(x) = -\left(\frac{5}{\lambda^3}\right)^{1/2} x.$$

For $\lambda \leq 0$, Eq. (17) has only the trivial solution $y(x) \equiv 0$.

16.4-2. Method of Integral Transforms.

1°. Consider the following nonlinear integral equation with quadratic nonlinearity on a semi-axis:

$$\mu y(x) - \lambda \int_0^\infty \frac{1}{t} y\left(\frac{x}{t}\right) y(t) dt = f(x). \quad (18)$$

To solve this equation, the Mellin transform can be applied, which, with regard to the convolution theorem (see Section 9.3), leads to a quadratic equation for the transform $\hat{y}(s) = \mathfrak{M}\{y(x)\}$:

$$\mu\hat{y}(s) - \lambda\hat{y}^2(s) = \hat{f}(s).$$

This implies

$$\hat{y}(s) = \frac{\mu \pm \sqrt{\mu^2 - 4\lambda\hat{f}(s)}}{2\lambda}. \quad (19)$$

The inverse transform $y(x) = \mathfrak{M}^{-1}\{\hat{y}(s)\}$ obtained by means of the Mellin inversion formula (if it exists) is a solution of Eq. (18). To different signs in the formula for the images (19), there are two corresponding solutions of the original equation.

2°. By applying the Mellin transform, one can solve nonlinear integral equations of the form

$$y(x) - \lambda \int_0^\infty t^\beta y(xt)y(t) dt = f(x). \quad (20)$$

The Mellin transform (see Table 3 in Section 9.3) reduces (20) to the following functional equation for the transform $\hat{y}(s) = \mathfrak{M}\{y(x)\}$:

$$\hat{y}(s) - \lambda\hat{y}(s)\hat{y}(1-s+\beta) = \hat{f}(s). \quad (21)$$

On replacing s by $1-s+\beta$ in (21), we obtain the relationship

$$\hat{y}(1-s+\beta) - \lambda\hat{y}(s)\hat{y}(1-s+\beta) = \hat{f}(1-s+\beta). \quad (22)$$

On eliminating the quadratic term from (21) and (22), we obtain

$$\hat{y}(s) - \hat{f}(s) = \hat{y}(1-s+\beta) - \hat{f}(1-s+\beta).$$

We express $\hat{y}(1-s+\beta)$ from this relation and substitute it into (21). We arrive at the quadratic equation

$$\lambda\hat{y}^2(s) - [1 + \hat{f}(s) - \hat{f}(1-s+\beta)]\hat{y}(s) + \hat{f}(s) = 0.$$

On solving this equation for $\hat{y}(s)$, by means of the Mellin inversion formula we can find a solution of the original integral equation (20).

16.4-3. Method of Differentiating for Integral Equations.

1°. The nonlinear integral equation

$$y(x) + \int_a^b |x-t|f(t, y(t)) dt = g(x), \quad a \leq x \leq b \quad (23)$$

can be reduced to a nonlinear second-order equation by double differentiation.

Let us remove the modulus in the integrand:

$$y(x) + \int_a^x (x-t)f(t, y(t)) dt + \int_x^b (t-x)f(t, y(t)) dt = g(x). \quad (24)$$

Differentiating (24) with respect to x yields

$$y'_x(x) + \int_a^x f(t, y(t)) dt - \int_x^b f(t, y(t)) dt = g'_x(x). \quad (25)$$

Differentiating (25), we arrive at a second-order ordinary differential equation for $y = y(x)$:

$$y''_{xx} + 2f(x, y) = g''_{xx}(x). \quad (26)$$

For the boundary conditions for this equation, see Section 8.8 in the first part of the book (Eq. 8.8.15).

2°. The equation

$$y(x) + \int_a^b e^{\lambda|x-t|} f(t, y(t)) dt = g(x) \quad (27)$$

can also be reduced to a nonlinear second-order equation by double differentiation (with subsequent elimination of the integral term by using the original equation):

$$y''_{xx} + 2\lambda f(x, y) - \lambda^2 y = g''_{xx}(x) - \lambda^2 g(x). \quad (28)$$

For the boundary conditions for this equation, see Section 8.8 of the first part of the book (Eq. 8.8.16).

Remark. A considerable number of exact solutions to ordinary differential equations (26) and (28) for various functions $f(x, y)$ and $g(x)$ can be found in the book by Polyanin and Zaitsev (2003).

3°. The equations

$$y(x) + \int_a^b \sinh(\lambda|x-t|) f(t, y(t)) dt = g(x),$$

$$y(x) + \int_a^b \sin(\lambda|x-t|) f(t, y(t)) dt = g(x),$$

can also be reduced to second-order ordinary differential equations by means of the differentiation. For these equations, see Section 8.8 of the first part of the book (Eqs. 8.8.17 and 8.8.18).

16.4-4. Method for Special Urysohn Equations of the First Kind.

1°. Consider the linear integral equation of the first kind

$$\int_a^b K(x, t)Y(t) dt = f(x). \quad (29)$$

Suppose equation (29) can be solved for any $f(x)$ from some class of functions $L_{\mathcal{F}}$. Let $Y_f(t)$ denote the corresponding solution.

Now consider the more complex nonlinear Urysohn equation of the first kind

$$\int_a^b [K(x, t)y(t) + \varphi(x)\Psi(t, y(t))] dt = f(x) \quad (30)$$

with its kernel containing an additional nonlinear term $\varphi(x)\Psi(t, y(t))$. A solution to equation (30) will be sought in the form

$$y(t) = Y_f(t) + AY_{\varphi}(t), \quad (31)$$

where $Y_{\varphi}(t)$ is the solution to equation (29) in which $f(x)$ must be replaced with $\varphi(x)$. Substituting (31) into (30) we have the following algebraic (transcendental) equation for the coefficient A :

$$A + \int_a^b \Psi(t, Y_f(t) + AY_{\varphi}(t)) dt = 0. \quad (32)$$

Formulas (31)–(32) can define one, several, or infinitely many solutions (or even none) to equation (30). In addition, the condition $\varphi(x) \in L_{\mathcal{F}}$ must be satisfied.

Example 2. The solution of the linear integral equation of the first kind

$$\int_0^{\infty} \sin(xt)Y(t) dt = f(x) \quad (33)$$

is expressed as (see equation 3.5.8 in Section 3.5)

$$Y_f(t) = \frac{2}{\pi} \int_0^{\infty} \sin(xt)f(x) dx. \quad (34)$$

Up to constant factors, the function $f(x)$ and the solution $Y_f(t)$ in (33)–(34) are the Fourier sine transform pair.

Now consider the more complex integral equation with quadratic nonlinearity

$$\int_0^{\infty} [\sin(xt)y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x). \quad (35)$$

In terms of equation (30), we have $\Psi(t, y(t)) = \psi(t)y^2(t)$ in (35). The corresponding solution (34) to equation (33) with $\varphi(x)$ is written as

$$Y_{\varphi}(t) = \frac{2}{\pi} \int_0^{\infty} \sin(xt)\varphi(x) dx. \quad (36)$$

Hence, equation (35) has the two solutions

$$y(t) = Y_f(t) + A_{1,2}Y_{\varphi}(t),$$

where $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0, \\ p = \int_0^{\infty} \psi(t)Y_{\varphi}^2(t) dt, \quad q = 1 + 2 \int_0^{\infty} \psi(t)Y_f(t)Y_{\varphi}(t) dt, \quad r = \int_0^{\infty} \psi(t)Y_f^2(t) dt.$$

Here all integrals are supposed to converge.

2°. The integral equation

$$\int_a^b \left[K(x, t)y(t) + \sum_{m=1}^n \varphi_m(x)\Psi_m(t, y(t)) \right] dt = f(x) \quad (37)$$

whose kernel is the sum of the kernel of equation (29) and an arbitrary degenerate nonlinear kernel can be solved in a similar manner. The solution is sought in the additive form

$$y(t) = Y_f(t) + \sum_{m=1}^n A_m Y_{\varphi_m}(t), \quad (38)$$

where $Y_{\varphi_m}(x)$ is the solution to equation (29) in which $f(x)$ must be replaced with $\varphi_m(x)$. Substituting (38) into (37) results in the following algebraic (transcendental) system of equations for the coefficients A_m :

$$A_m + \int_a^b \Psi_m \left(t, Y_f(t) + \sum_{j=1}^n A_j Y_{\varphi_j}(t) \right) dt = 0, \quad m = 1, \dots, n. \quad (39)$$

16.4-5. Method for Special Urysohn Equations of the Second Kind.

1°. Consider the linear equation of the second kind

$$Y(x) + \int_a^b K(x, t)Y(t) dt = f(x). \quad (40)$$

Suppose equation (40) can be solved for any $f(x)$ from some class of functions $L_{\mathcal{F}}$. Let $Y_f(x)$ denote the corresponding solution.

Now consider the more complex nonlinear Urysohn equation of the second kind

$$y(x) + \int_a^b [K(x, t)y(t) + \varphi(x)\Psi(t, y(t))] dt = f(x), \tag{41}$$

with its kernel containing an additional term $\varphi(x)\Psi(t, y(t))$. A solution to equation (41) will be sought in the form

$$y(x) = Y_f(x) + AY_\varphi(x), \tag{42}$$

where $Y_\varphi(x)$ is the solution to equation (40) in which $f(x)$ must be replaced with $\varphi(x)$. Substituting (42) into (41) we have the following algebraic (transcendental) equation for the coefficient A :

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0. \tag{43}$$

Formulas (42)–(43) can define one, several, or infinitely many solutions (or even none) to equation (41). In addition, the condition $\varphi(x) \in L_{\mathcal{F}}$ must be satisfied.

Example 3. The solution of the linear integral equation of the second kind

$$Y(x) + \lambda \int_{-\infty}^{\infty} e^{-|x-t|} Y(t) dt = f(x), \quad \lambda > -\frac{1}{2}, \tag{44}$$

is expressed as (see equation 4.2.14 in Section 4.2)

$$Y_f(x) = f(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) f(t) dt. \tag{45}$$

Now consider the more complex integral equation with quadratic nonlinearity

$$y(x) + \int_{-\infty}^{\infty} [\lambda e^{-|x-t|} y(t) + \varphi(x)\psi(t)y^2(t)] dt = f(x). \tag{46}$$

In terms of equation (41), we have $\Psi(t, y(t)) = \psi(t)y^2(t)$ in (46). The corresponding solution (45) to equation (44) with $\varphi(x)$ is written as

$$Y_\varphi(x) = \varphi(x) - \frac{\lambda}{\sqrt{1+2\lambda}} \int_{-\infty}^{\infty} \exp(-\sqrt{1+2\lambda}|x-t|) \varphi(t) dt.$$

Hence, equation (46) has the two solutions

$$y(t) = Y_f(t) + A_{1,2}Y_\varphi(t),$$

where $A_{1,2}$ are roots of the quadratic equation

$$pA^2 + qA + r = 0, \\ p = \int_0^\infty \psi(t)Y_\varphi^2(t) dt, \quad q = 1 + 2 \int_0^\infty \psi(t)Y_f(t)Y_\varphi(t) dt, \quad r = \int_0^\infty \psi(t)Y_f^2(t) dt.$$

2°. The integral equation

$$y(x) + \int_a^b \left[K(x, t)y(t) + \sum_{m=1}^n \varphi_m(x)\Psi_m(t, y(t)) \right] dt = f(x), \tag{47}$$

with its kernel being the sum of the kernel of equation (40) and an arbitrary degenerate nonlinear kernel, can be solved in a similar manner. The solution is sought in the additive form

$$y(x) = Y_f(x) + \sum_{m=1}^n A_m Y_{\varphi_m}(x), \tag{48}$$

where $Y_{\varphi_m}(x)$ is the solution to equation (40) in which $f(x)$ must be replaced with $\varphi_m(x)$. Substituting (48) into (47) results in the following algebraic (transcendental) system of equations for the coefficients A_m :

$$A_m + \int_a^b \Psi_m \left(t, Y_f(t) + \sum_{j=1}^n A_j Y_{\varphi_j}(t) \right) dt = 0, \quad m = 1, \dots, n. \tag{49}$$

Remark 1. Formulas (38)–(39) and (48)–(49), which define solutions to the special Urysohn equations of the first and second kind (37) and (47), respectively, are coincident (but the functions $Y_f(x)$ and $Y_{\varphi_m}(x)$ are different).

Remark 2. The method outlined may be used for approximate solution of nonlinear integral equations of the form

$$y(x) + \int_a^b [K(x, t)y(t) + \Psi(x, t, y(t))] dt = f(x)$$

by appropriately selecting an approximation of the nonlinear part of the kernel, $\Psi(x, t, y(t)) \approx \sum_{m=1}^n \varphi_m(x)\Psi_m(t, y(t))$.

16.4-6. Some Generalizations.

The method presented in Subsections 16.4-4 and 16.4-5 for the special Urysohn equations of the first and second kind admits generalizations.

Consider an abstract nonlinear equation for the function $y = y(x)$:

$$\mathbf{L}[y] + \sum_{m=1}^n \varphi_m(x)I_m[y] = f(x), \quad (50)$$

where $\mathbf{L}[y]$ is a linear operator (it can be integral, functional, differential,* or other) and $I_m[y]$ are some nonlinear functionals (i.e., numbers for any given $y(x)$).

Examples of nonlinear functionals:

$$I_1[y] = ay^2(0) + by(1), \quad I_2[y] = \max_{0 \leq x \leq 1} |y(x)|, \quad I_3[y] = \int_a^b K(t, y(t), y'_t(t), y''_{tt}(t)) dt.$$

Suppose the truncated linear equation

$$\mathbf{L}[Y] = f(x), \quad (51)$$

obtained from (50) by setting $\varphi_m(x) = 0$ ($m = 1, \dots, n$), can be solved for any $f(x)$ from some class of functions $L_{\mathcal{F}}$. Let $Y_f(x)$ denote the corresponding solution.

Let the conditions $\varphi_m(x) \in L_{\mathcal{F}}$ ($m = 1, \dots, n$) be satisfied.

Solutions to the nonlinear equation (50) are sought in the form

$$y(x) = Y_f(x) + \sum_{m=1}^n A_m Y_{\varphi_m}(x), \quad (52)$$

where $Y_{\varphi_m}(x)$ is the solution to equation (51) in which $f(x)$ must be replaced with $\varphi_m(x)$. Substituting (52) into (50) results in the following algebraic (transcendental) system of equations for the coefficients A_m :

$$A_m + I_m \left[Y_f(x) + \sum_{j=1}^n A_j Y_{\varphi_j}(x) \right], \quad m = 1, \dots, n. \quad (53)$$

Formulas (52)–(53) can define one, several, or infinitely many solutions (or even none) to equation (50).

* In this case, the equation must be supplemented with appropriate homogeneous boundary conditions.

Example 4. Consider the nonlinear functional integral equation

$$y(x) + \lambda y(a-x) + \int_0^a \varphi(x) \Psi(t, y(t), y(a-t)) dt = f(x), \quad \lambda \neq \pm 1, \quad (54)$$

where $0 \leq x \leq a$, $0 \leq t \leq a$.

The truncated linear functional equations (54), with $\varphi(x) = 0$, has the solution

$$Y_f(x) = \frac{f(x) - \lambda f(a-x)}{1 - \lambda^2}.$$

Therefore, solutions to the nonlinear equation (54) are sought in the form

$$y(x) = Y_f(x) + AY_\varphi(x), \quad Y_\varphi(x) = \frac{\varphi(x) - \lambda\varphi(a-x)}{1 - \lambda^2},$$

where the constant A is determined from the algebraic (transcendental) equation

$$A + \int_0^a \Psi(t, Y_f(t) + AY_\varphi(t), Y_f(a-t) + AY_\varphi(a-t)) dt = 0.$$

Example 5. Consider the nonlinear integro-functional-differential equation

$$\int_0^{\pi/2} [y(x \sin t) + \varphi(x) \Psi(t, y(t), y'_t(t))] dt = f(x). \quad (55)$$

For $\varphi(x) = 0$, it is the Schlömilch equation. Its solution is given in Subsection 3.5 (see Eq. 3.5.40). It should be noted that equation (50) contains the unknown function with different arguments, $y(x \sin t)$ and $y(t)$.

Following the method described above, we look for solutions to equation (55) in the form

$$y_m(z) = Y_f(z) + A_m Y_\varphi(z),$$

where

$$Y_f(z) = \frac{2}{\pi} \left[f(0) + z \int_0^{\pi/2} f'_\xi(\xi) d\tau \right], \quad Y_\varphi(z) = \frac{2}{\pi} \left[\varphi(0) + z \int_0^{\pi/2} \varphi'_\xi(\xi) d\tau \right], \quad \xi = z \sin \tau,$$

and A_m are roots of the algebraic (transcendental) equation

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t), Y'_f(t) + AY'_\varphi(t)) dt = 0.$$

Example 6. Consider a boundary value problem for the nonlinear integro-differential equation

$$y''_{xx} + \varphi(x) \int_0^1 \Psi(t, y(t)) dt = f(x) \quad (56)$$

with homogeneous boundary conditions

$$y(0) = y(1) = 0. \quad (57)$$

The solution of an auxiliary linear boundary value problem

$$Y''_{xx} = f(x); \quad Y(0) = Y(1) = 0$$

has the form

$$Y_f(x) = \int_0^1 G(x, \xi) f(\xi) d\xi, \quad (58)$$

$$G(x, \xi) = \begin{cases} (\xi - 1)x & \text{for } 0 \leq x \leq \xi \leq 1, \\ (x - 1)\xi & \text{for } 0 \leq \xi \leq x \leq 1. \end{cases}$$

Therefore solutions of the original boundary value problem for nonlinear integro-differential equation (56) with boundary conditions (57) can be constructed in the form

$$y(x) = Y_f(x) + AY_\varphi(x), \quad (59)$$

where $Y_\varphi(x)$ is determined by the right-hand side of formula (58), in which function $f(x)$ is changed by function $\varphi(x)$. Substitution of (59) to (56) leads to the following algebraic (transcendental) equation for determining of A :

$$A + \int_a^b \Psi(t, Y_f(t) + AY_\varphi(t)) dt = 0. \quad (60)$$

In particular case of $f(x) = 0$ and $\varphi(x) = 1$ in formulas (59)–(60) it is necessary to set $Y_f(x) = 0$, $Y_\varphi(x) = \frac{1}{2}x(x-1)$.

Remark. In the case of inhomogeneous boundary conditions for integro-differential equation it is necessary to change variables using relation $y(x) = \bar{y}(x) + g(x)$, where $g(x)$ is an arbitrary sufficiently smooth function that satisfies boundary conditions. Finally we obtain the problem with homogeneous boundary conditions. For example, for integro-differential equation (56) with inhomogeneous boundary conditions $y(0) = a$, $y(1) = b$ one can take $g(x) = a + (b - a)x$.

⊙ References for Section 16.4: M. L. Krasnov, A. I. Kiselev, and G. I. Makarenko (1971), A. D. Polyanin and A. V. Manzhirov (1998), A. D. Polyanin and A. I. Zhurov (2007).

16.5. Approximate and Numerical Methods for Nonlinear Equations with Constant Integration Limits

16.5-1. Successive Approximation Method.

Consider the nonlinear Urysohn integral equation in the canonical form:

$$y(x) = \int_a^b \mathcal{K}(x, t, y(t)) dt, \quad a \leq x \leq b. \quad (1)$$

The iteration process for this equation is constructed by the formula

$$y_k(x) = \int_a^b \mathcal{K}(x, t, y_{k-1}(t)) dt, \quad k = 1, 2, \dots \quad (2)$$

If the function $\mathcal{K}(x, t, y)$ is jointly continuous together with the derivative $\mathcal{K}'_y(x, t, y)$ (with respect to the variables x, t , and y), ρ , $a \leq x \leq b$, $a \leq t \leq b$, and $|y| \leq \rho$ and if

$$\int_a^b \sup_y |\mathcal{K}(x, t, y)| dt \leq \rho, \quad \int_a^b \sup_y |\mathcal{K}'_y(x, t, y)| dt \leq \beta < 1, \quad (3)$$

then for any continuous function $y_0(x)$ of the initial approximation from the domain $\{|y| \leq \rho, a \leq x \leq b\}$, the successive approximations (2) converge to a continuous solution $y^*(x)$, which lies in the same domain and is unique in this domain. The rate of convergence is defined by the inequality

$$|y^*(x) - y_k(x)| \leq \frac{\beta^k}{1 - \beta} \sup_x |y_1(x) - y_0(x)|, \quad a \leq x \leq b, \quad (4)$$

which gives an *a priori* estimate for the error of the k th approximation. The *a posteriori* estimate (which is, in general, more precise) has the form

$$|y^*(x) - y_k(x)| \leq \frac{\beta}{1 - \beta} \sup_x |y_k(x) - y_{k-1}(x)|, \quad a \leq x \leq b. \quad (5)$$

A solution of an equation of the form (1) with an additional term $f(x)$ on the right-hand side can be constructed in a similar manner.

Example 1. Let us apply the successive approximation method to solve the equation

$$y(x) = \int_0^1 xty^2(t) dt - \frac{5}{12}x + 1.$$

The recursive formula has the form

$$y_k(x) = \int_0^1 xty_{k-1}^2(t) dt - \frac{5}{12}x + 1, \quad k = 1, 2, \dots$$

For the initial approximation we take $y_0(x) = 1$. The calculation yields

$$\begin{aligned} y_1(x) &= 1 + 0.083x, & y_2(x) &= 1 + 0.14x, & y_3(x) &= 1 + 0.18x, & \dots, \\ y_8(x) &= 1 + 0.27x, & y_9(x) &= 1 + 0.26x, & y_{10}(x) &= 1 + 0.29x, & \dots, \\ y_{16}(x) &= 1 + 0.318x, & y_{17}(x) &= 1 + 0.321x, & y_{18}(x) &= 1 + 0.323x, & \dots \end{aligned}$$

Thus, the approximations tend to the exact solution $y(x) = 1 + \frac{1}{3}x$. We see that the rate of convergence of the iteration process is fairly small.

Note that in Subsection 16.5-2, the equation in question is solved by a more efficient method.

16.5-2. Newton–Kantorovich Method.

The solution of nonlinear integral equations is a complicated problem of computational mathematics, which is related to difficulties of both a principal and computational character. In this connection, methods are developed that are especially designed for solving nonlinear equations, including the Newton–Kantorovich method, which makes it possible to provide and accelerate the convergence of iteration processes in many cases.

We consider this method in connection with the Urysohn equation in the canonical form (1). The iteration process is constructed as follows:

$$y_k(x) = y_{k-1}(x) + \varphi_{k-1}(x), \quad k = 1, 2, \dots, \quad (6)$$

$$\varphi_{k-1}(x) = \varepsilon_{k-1}(x) + \int_a^b \mathcal{K}'_y(x, t, y_{k-1}(t)) \varphi_{k-1}(t) dt, \quad (7)$$

$$\varepsilon_{k-1}(x) = \int_a^b \mathcal{K}(x, t, y_{k-1}(t)) dt - y_{k-1}(x). \quad (8)$$

At each step of the algorithm, a linear integral equation for the correction $\varphi_{k-1}(x)$ is solved. Under some conditions, the process (6) has high rate of convergence; however, it is rather complicated, because at each iteration we must obtain the new kernel $\mathcal{K}'_y(x, t, y_{k-1}(t))$ for Eqs. (7).

The algorithm can be simplified by using the equation

$$\varphi_{k-1}(x) = \varepsilon_{k-1}(x) + \int_a^b \mathcal{K}'_y(x, t, y_0(t)) \varphi_{k-1}(t) dt \quad (9)$$

instead of (7). If the initial approximation is chosen successfully, then the difference between the integral operators in (7) and (9) is small, and the kernel in (9) remains the same in the course of the solution.

The successive approximation method that consists in the application of formulas (6), (8), and (9) is called the *modified Newton–Kantorovich method*. In principle, its rate of convergence is less than that of the original (unmodified) method; however, this version of the method is less complicated in calculations, and therefore it is frequently preferable.

Let the function $\mathcal{K}(x, t, y)$ be jointly continuous together with the derivatives $\mathcal{K}'_y(x, t, y)$ and $\mathcal{K}''_{yy}(x, t, y)$ with respect to the variables x, t, y , where $a \leq x \leq b$ and $a \leq t \leq b$, and let the following conditions hold:

1°. For the initial approximation $y_0(x)$, the resolvent $\mathcal{R}(x, t)$ of the linear integral equation (7) with the kernel $\mathcal{K}'_y(x, t, y_0(t))$ satisfies the condition

$$\int_a^b |\mathcal{R}(x, t)| dt \leq A < \infty, \quad a \leq x \leq b.$$

2°. The residual $\varepsilon_0(x)$ of Eq. (8) for the approximation $y_0(x)$ satisfies the inequality

$$|\varepsilon_0(x)| = \left| \int_a^b \mathcal{K}(x, t, y_0(t)) dt - y_0(x) \right| \leq B < \infty.$$

3°. In the domain $|y(x) - y_0(x)| \leq 2(1 + A)B$, the following relation holds:

$$\int_a^b \sup_y |\mathcal{K}''_{yy}(x, t, y)| dt \leq D < \infty.$$

4°. The constants A, B , and D satisfy the condition

$$H = (1 + A)^2 BD \leq \frac{1}{2}.$$

In this case, under assumptions $1^\circ-4^\circ$, the process (6) converges to a solution $y^*(x)$ of Eq. (1) in the domain

$$|y(x) - y_0(x)| \leq (1 - \sqrt{1 - 2H})H^{-1}(1 - A)B, \quad a \leq x \leq b.$$

This solution is unique in the domain

$$|y(x) - y_0(x)| \leq 2(1 + A)B, \quad a \leq x \leq b.$$

The rate of convergence is determined by the estimate

$$|y^*(x) - y_k(x)| \leq 2^{1-k}(2H)^{2^k-1}(1 - A)B, \quad a \leq x \leq b.$$

Thus, the above conditions establish the convergence of the algorithm and the existence, the position, and the uniqueness domain of a solution of the nonlinear equation (1). These conditions impose certain restrictions on the initial approximation $y_0(x)$ whose choice is an important independent problem that has no unified approach. As usual, the initial approximation is determined either by more detailed *a priori* analysis of the equation under consideration or by physical reasoning implied by the essence of the problem described by this equation. Under a successful choice of the initial approximation, the Newton–Kantorovich method provides a high rate of convergence of the iteration process to obtain an approximate solution with given accuracy.

Remark. Let the right-hand side of Eq. (1) contain an additional term $f(x)$. Then such an equation can be represented in the form (1), where the integrand is $K(x, t, y(t)) + (b - a)^{-1}f(x)$.

Example 2. Let us apply the Newton–Kantorovich method to solve the equation

$$y(x) = \int_0^1 xty^2(t) dt - \frac{5}{12}x + 1. \quad (10)$$

For the initial approximation we take $y_0(x) = 1$. According to (8), we find the residual

$$\varepsilon_0(x) = \int_0^1 xty_0^2(t) dt - \frac{5}{12}x + 1 - y_0(x) = x \int_0^1 t dt - \frac{5}{12}x + 1 - 1 = \frac{1}{12}x.$$

The y -derivative of the kernel $\mathcal{K}(x, t, y) = xty^2(t)$, which is needed in the calculations, has the form $\mathcal{K}'_y(x, t, y) = 2xty(t)$. According to (7), we form the following equation for $\varphi_0(x)$:

$$\varphi_0(x) = \frac{1}{12}x + 2x \int_0^1 ty_0(t)\varphi_0(t) dt,$$

where the kernel turns out to be degenerate, which makes it possible to obtain the solution $\varphi_0(x) = \frac{1}{4}x$ directly.

Now we define the first approximation to the desired function:

$$y_1(x) = y_0(x) + \varphi_0(x) = 1 + \frac{1}{4}x.$$

We continue the iteration process and obtain

$$\varepsilon_1(x) = \int_0^1 xt(1 + \frac{1}{4}t) dt + (1 - \frac{5}{12}x) - (1 + \frac{1}{4}x) = \frac{1}{64}x.$$

The equation for $\varphi_1(x)$ has the form

$$\varphi_1(x) = \frac{1}{64}x + 2x \int_0^1 t(1 + \frac{1}{4}t) dt + (1 - \frac{5}{12}x) - (1 + \frac{1}{4}x),$$

and the solution is $\varphi_1(x) = \frac{3}{40}x$. Hence, $y_2(x) = 1 + \frac{1}{4}x + \frac{3}{40}x = 1 + 0.325x$. The maximal difference between the exact solution $y(x) = 1 + \frac{1}{3}x$ and the approximate solution $y_2(x)$ is observed at $x = 1$ and is less than 0.5%.

This solution is not unique. The other solution can be obtained by taking the function $y_0(x) = 1 + 0.8x$ for the initial approximation. In this case we can repeat the above sequence of approximations and obtain the following results (the numerical coefficient of x is rounded):

$$y_1(x) = 1 + 0.82x, \quad y_2(x) = 1 + 1.13x, \quad y_3(x) = 1 + 0.98x, \quad \dots,$$

and the subsequent approximations tend to the exact solution $y(x) = 1 + x$.

We see that the rate of convergence of the iteration process performed by the Newton–Kantorovich method is significantly higher than that performed by the method of successive approximations (see Example 1 in Subsection 16.5-1).

To estimate the rate of convergence of the performed iteration process, we can compare the above results with the realization of the modified Newton–Kantorovich method. In connection with the latter, for the above versions of the approximations we can obtain

$$y_n(x) = 1 + k_n x;$$

k_0	k_1	k_2	k_3	k_4	k_5	k_6	k_7	k_8	\dots
0	0.25	0.69	0.60	0.51	0.44	0.38	0.36	0.345	\dots

The iteration process converges to the exact solution $y(x) = 1 + \frac{1}{3}x$.

We see that the modified Newton–Kantorovich method is less efficient than the Newton–Kantorovich method, but more efficient than the method of successive approximations (see Example 1 in Subsection 16.5-1).

16.5-3. Quadrature Method.

To solve an arbitrary nonlinear equation, we can apply the method based on the application of quadrature formulas. The procedure of composing the approximating system of equations is the same as in the linear case (see Subsection 13.19-1). We consider this procedure for an example of the Urysohn equation of the second kind:

$$y(x) - \int_a^b K(x, t, y(t)) dt = f(x), \quad a \leq x \leq b. \quad (11)$$

We set $x = x_i$ ($i = 1, \dots, n$). Then we obtain

$$y(x_i) - \int_a^b K(x_i, t, y(t)) dt = f(x_i), \quad i = 1, \dots, n, \quad (12)$$

On applying the quadrature formula from Subsection 13.19-1 and neglecting the approximation error, we transform relations (12) into the system of nonlinear equations

$$y_i - \sum_{j=1}^n A_j K_{ij}(y_j) = f_i, \quad i = 1, \dots, n, \quad (13)$$

for the approximate values y_i of the solution $y(x)$ at the nodes x_1, \dots, x_n , where $f_i = f(x_i)$ and $K_{ij}(y_j) = K(x_i, t_j, y_j)$, and A_j are the coefficients of the quadrature formula.

The solution of the nonlinear system (13) gives values y_1, \dots, y_n for which by interpolation we find an approximate solution of the integral equation (11) on the entire interval $[a, b]$. For the analytic expression of an approximate solution, we can take the function

$$\tilde{y}(x) = f(x) + \sum_{j=1}^n A_j K(x, x_j, y_j). \quad (14)$$

16.5-4. Tikhonov Regularization Method.

In connection with the nonlinear Urysohn integral equation of the first kind

$$\int_a^b K(x, t, y(t)) dt = f(x), \quad c \leq x \leq d, \quad (15)$$

where $f(x) \in L_2(c, d)$ and $y(t) \in L_2(a, b)$, the Tikhonov regularization method leads to a regularized nonlinear integral equation in the form

$$\alpha y_\alpha(x) + \int_a^b M(t, x, y_\alpha(t), y_\alpha(x)) dt = F(x, y_\alpha(x)), \quad a \leq x \leq b, \quad (16)$$

$$M(t, x, y(t), y(x)) = \int_c^d K(s, t, y(t)) K'_y(s, x, y(x)) ds, \quad (17)$$

$$F(x, y(x)) = \int_c^d K'_y(t, x, y(x)) f(t) dt, \quad (18)$$

where α is a regularization parameter.

For instance, by applying the quadrature method on the basis of the trapezoidal rule, we can reduce Eq. (16) to a system of nonlinear algebraic equations. An approximate solution of (15) is constructed by the principle described above for linear equations (see Section 12.11).

⊙ References for Section 16.5: N. S. Smirnov (1951), P. P. Zabreyko, A. I. Koshelev, et al. (1975), F. G. Tricomi (1985), A. F. Verlan' and V. S. Sizikov (1986).

16.6 Existence and Uniqueness Theorems for Nonlinear Equations

16.6-1. Hammerstein Equations.

1°. Consider the Hammerstein equation

$$y(x) = \int_a^b K(x, t) \Phi(t, y(t)) dt, \quad a \leq x \leq b. \quad (1)$$

Assume that the function $\Phi(t, y)$ is continuous, and the kernel $K(x, t)$ is positive definite, continuous, and symmetric, $K(x, t) = K(t, x)$.

THEOREM 1. *Suppose that the inequality*

$$|\Phi(t, y)| \leq C_1 |y| + C_2$$

holds with some positive constants C_1 and C_2 such that $C_1 < \lambda_1$, and λ_1 is the smallest characteristic value of the kernel $K(x, t)$. Then the nonlinear integral equation (1) has at least one continuous solution.

THEOREM 2. *If for any fixed $t \in [a, b]$, the function $\Phi(t, y)$ is nondecreasing with respect to y , then the nonlinear integral equation (1) has at most one solution.*

THEOREM 3. *The nonlinear integral equation (1) has at most one solution if the function $\Phi(t, y)$ satisfies the uniform Lipschitz condition*

$$|\Phi(t, y_2) - \Phi(t, y_1)| \leq \sigma |y_2 - y_1|,$$

where $0 < \sigma < \lambda_1$, λ_1 is the smallest characteristic value of the kernel $K(x, t)$.

THEOREM 4 (ON NONEXISTENCE OF SOLUTIONS). *Suppose that $K(x, t) \geq 0$, $K(x, t) \not\equiv 0$, and the eigenfunction $y_1(x)$ of the kernel $K(x, t)$ corresponding to the smallest characteristic value λ_1 does not change sign in the domain $a \leq x, t \leq b$. Then the condition*

$$\Phi(t, y(t)) > \lambda_1 y(t) \quad (\text{for all } t \in [a, b])$$

ensures that equation (1) has no solutions.

2°. Assume now that the kernel $K(x, t)$ of equation (1) is continuous and positive definite (possibly, nonsymmetric, $K(x, t) \neq K(t, x)$), and the function $\Phi(t, y)$ is continuous.

THEOREM 5. *Suppose that the inequality*

$$\int_0^y \Phi(t, y) dy \leq \frac{1}{2}Ay^2 + B \quad (t \in \Omega, |y| < \infty) \quad (2)$$

holds with a constant $A < \lambda_1$, where λ_1 is the smallest characteristic value of the kernel $K(x, t)$. Then equation (1) has at least one continuous solution.

Now consider the case of an unbounded positive-definite kernel $K(x, t)$. Then the following result holds.

THEOREM 6. *Suppose that the kernel $K(x, t)$ satisfies the condition*

$$\int_a^b \int_a^b |K(x, t)|^p dx dt < \infty, \quad p \geq 2,$$

and the function $\Phi(t, y)$ satisfies the inequality (2) and the condition

$$|\Phi(t, y)| \leq a + b|u|^{p-1} \quad (a \leq x, t \leq b, |y| < \infty).$$

Then equation (1) has at least one solution.

THEOREM 7. *Let $K(x, t)$ be positive and continuous in the domain $a \leq x, t \leq b$. Suppose that the function $\Phi(t, y)$ is continuous in the domain $a \leq t \leq b, y > 0$, nonnegative for $y \geq 0$ and strictly positive for $y > 0$ and almost all t . Suppose also that one of the following conditions holds:*

1) $\Phi(t, y)$ does not decrease in y , and $y^{-\beta}\Phi(t, y)$ does not increase in y , where β is a point from the interval $(0, 1)$;

2) $\Phi(t, y)$ does not increase in y , and $y^\beta\Phi(t, y)$ increases in y , where β is a point from the interval.

Then equation (1) has one and only one positive solution. This solution is the uniform limit of the successive approximations

$$y_n(x) = \int_a^b K(x, t) \Phi(t, y_{n-1}(t)) dt \quad (n = 1, 2, \dots),$$

where $y_0(x)$ is an arbitrary nonzero nonnegative initial function.

3°. Consider a system of integral equations of Hammerstein's type

$$y_i(x) = \int_a^b K_i(x, t) \Phi_i(t, y_1(t), \dots, y_n(t)) dt \quad (i = 1, \dots, n) \quad (3)$$

with continuous symmetric positive-definite kernels $K_i(x, t)$, where $\Phi_i(t, y_1, \dots, y_n)$ are continuous functions in all their arguments.

THEOREM 8. *Suppose that the functions $\Phi_i(t, y_1, \dots, y_n)$ satisfy the inequality*

$$\sum_{i=1}^n y_i \Phi_i(t, y_1, \dots, y_n) \leq A \sum_{i=1}^n y_i^2 + B,$$

where $A < \lambda_0$ and λ_0 is the smallest characteristic value of the kernels $K_i(x, t)$. Then system (3) has at least one continuous solution.

16.6-2. Urysohn Equations.

Consider the nonlinear Urysohn equation

$$y(x) = \lambda \int_a^b K(x, t, y(t)) dt + f(x). \quad (4)$$

THEOREM 1. Let $K(x, t, y)$ and $f(x)$ be continuous functions of their arguments $a \leq x, t \leq b$, $-\infty < y < \infty$, and suppose that $K(x, t, y)$ satisfies the Lipschitz condition in y ,

$$|K(x, t, y_2) - K(x, t, y_1)| \leq L|y_2 - y_1|,$$

where L is a constant independent of y_1 and y_2 . Then the condition

$$\lambda < \frac{b-a}{L}$$

ensures that equation (4) has one and only one continuous solution. This solution can be found by the method of successive approximations

$$y_{n+1}(x) = \lambda \int_a^b K(x, t, y_n(t)) dt + f(x), \quad n = 0, 1, \dots,$$

with an arbitrary continuous function $y_0(x)$.

Remark. If the function $K(x, t, y)$ has a bounded partial derivative in y :

$$\left| \frac{\partial K}{\partial y} \right| \leq \Lambda \quad (a \leq x, t \leq b, -\infty < y < \infty),$$

then $K(x, y, t)$ satisfies the Lipschitz condition in y with a constant $L \leq \Lambda$.

THEOREM 2. Let $f(x) \in L_2(a, b)$ and suppose that

$$|K(x, t, y_1) - K(x, t, y_2)| \leq M(x, t)|y_1 - y_2| \quad (a \leq x, t \leq b, -\infty < y < \infty),$$

where

$$\int_a^b \int_a^b |M(x, t)|^2 dx dt = B^2 < \infty.$$

Then, for $\lambda < \frac{1}{B}$, equation (4) has one and only one solution in $L_2(a, b)$.

THEOREM 3. Suppose that the function $K(x, t, y)$ is continuous in y and satisfies the inequality

$$|K(x, t, y)| \leq M(x, t)(A + B|y|^p) \quad (a \leq x, t \leq b, -\infty < y < \infty),$$

where $A, B, p > 0$, and

$$\int_a^b \int_a^b |M(x, t)|^{p+1} dx dt < \infty.$$

Then, for any sufficiently small $|\lambda|$ and any $f(x) \in L_{p+1}(a, b)$, equation (4) has a solution $y(x) \in L_{p+1}(a, b)$. If $p < 1$, then a solution exists for any λ .

THEOREM 4. Suppose that the function $K(x, t, y)$ is continuous in the domain $\Omega = \{a \leq x \leq b, a \leq t \leq b, |y| \leq \rho\}$ and its partial derivative in y is bounded,

$$\left| \frac{\partial K}{\partial y} \right| \leq C \quad (x, t, y \in \Omega).$$

Then the equation

$$y(x) = \lambda \int_a^b K(x, t, y(t)) dt, \quad (5)$$

where

$$|\lambda|C|b - a| < 1,$$

$$|\lambda| \max_{a \leq x \leq b} \int_a^b \max_{|y| \leq \rho} |K(x, t, y)| dt \leq \rho,$$

has one and only one continuous solution $y(x)$ ($a \leq x \leq b$) satisfying the inequality $|y(x)| \leq \rho$.

If $y_0(x)$ is an arbitrary continuous function satisfying the inequality $|y_0(x)| \leq \rho$ ($a \leq x \leq b$), then the successive approximations

$$y_{n+1}(x) = \lambda \int_a^b K(x, t, y_n(t)) dt \quad (n = 0, 1, \dots)$$

are uniformly convergent on $[a, b]$ to this solution.

THEOREM 5. Suppose that the function $K(x, t, y)$ is continuous in the domain $\Omega = \{a \leq x \leq b, a \leq t \leq b, |y| \leq \rho\}$. Then the condition

$$|\lambda| \leq \frac{\rho}{(b-a) \max_{x,t,y \in \Omega} |K(x, t, y)|}$$

ensures that the integral equation (5) has at least one continuous solution satisfying the inequality $|y(x)| \leq \rho$.

THEOREM 6. Let $K(x, t, y)$ be continuous in the domain $\Omega = \{a \leq x \leq b, a \leq t \leq b, -\infty < y < \infty\}$ and let

$$\varphi(r) = \max_{a \leq x, t \leq b, y \leq r} |K(x, t, y)|,$$

$$\Lambda = \sup_{0 < r < \infty} \frac{r}{(b-a)\varphi(r)}.$$

Then, for $|\lambda| < \Lambda$, the integral equation (5) has at least one continuous solution.

In particular, if

$$\lim_{r \rightarrow \infty} \frac{\varphi(r)}{r} = 0,$$

then the integral equation (5) has a solution for any λ . In this case, equation (4), too, has a solution for any λ and any continuous $f(x)$.

THEOREM 7. Suppose a function $K(x, t, y)$ is continuous in x, t, y and satisfies the inequalities

$$0 \leq K(x, t, y) \leq a + L(x, t)y \quad (a \leq x, t \leq b, y \geq 0),$$

where $L(x, t)$ is a nonnegative kernel the smallest characteristic value of which satisfies the condition $\lambda > 1$. Then the equation

$$y(x) = \int_a^b K(x, t, y(t)) dt$$

has at least one continuous nonnegative solution.

16.7. Nonlinear Equations with a Parameter: Eigenfunctions, Eigenvalues, Bifurcation Points

16.7-1. Eigenfunctions and Eigenvalues of Nonlinear Integral Equations.

Consider a nonlinear integral equation

$$y = \lambda \mathbf{A}[y] \quad (1)$$

with a parameter λ . An example of such an equation is provided by the Urysohn equation (5) from Subsection 16.6-2, which has a solution for sufficiently small λ (this solution tends to zero as $\lambda \rightarrow 0$). Of most importance for applications are nontrivial solutions corresponding to finite values of λ . By analogy with the theory of linear integral equations, nontrivial solutions of the nonlinear equation (1) are called *eigenfunctions* of the corresponding nonlinear operator \mathbf{A} (or *eigenfunctions* of the nonlinear equation (1)), and λ for which such solutions exist are called *characteristic values* (and the numbers $\mu = 1/\lambda$ are called *eigenvalues*).

For nonlinear integral equations (1) with a parameter (1), a typical situation is the existence of nontrivial solutions for all λ from some interval (α, β) (thus, there is a continuous spectrum of characteristic values corresponding to a continuum of eigenfunctions; see Section 16.1-3 for relevant examples), which is in harsh contrast with the situation typical for linear integral equations with a discrete spectrum of characteristic values.

THEOREM 1. *Suppose that $\Phi(t, y)$ is continuous and the kernel $K(x, t)$ is continuous, symmetric, and positive definite. Then the Hammerstein equation*

$$y(x) = \lambda \int_a^b K(x, t) \Phi(t, y(t)) dt, \quad a \leq x \leq b, \quad (2)$$

has a continuum of eigenfunctions.

THEOREM 2. *Let $K(x, t)$ be symmetric, positive definite, and satisfying the inequality*

$$\int_a^b \int_a^b |K(x, t)|^p dx dt < \infty, \quad p \geq 2,$$

and suppose that $\Phi(t, y)$ is continuous and satisfies the inequality

$$|\Phi(t, y)| \leq C_1 + C_2 |y|^{p-1}.$$

Then the Hammerstein equation (2) has a continuum of eigenfunctions.

THEOREM 3. *Suppose that the kernel $K(x, t)$ is continuous, symmetric, and has finitely many negative characteristic values. Suppose also that $\Phi(t, y)$ is continuous and satisfies the inequality*

$$y\Phi(t, y) \geq Ay^2 - B \quad (a \leq x, t \leq b, -\infty < y < \infty),$$

for $A > 0$. Then the Hammerstein equation (2) has a continuum of eigenfunctions.

THEOREM 4. *Suppose that $K(x, t, y)$ is continuous and satisfies the inequality*

$$K(x, t, y) \geq L(x, t)y \quad (a \leq x, t \leq b, y > 0),$$

where $L(x, t)$ is a positive continuous kernel. Then the Urysohn equation

$$y(x) = \lambda \int_a^b K(x, t, y(t)) dt$$

has a continuum of eigenfunctions.

16.7-2. Local Solutions of a Nonlinear Integral Equation with a Parameter.

Consider the Urysohn equation

$$y(x) = \int_a^b K(x, t, y(t); \lambda) dt \quad (a \leq x \leq b) \quad (3)$$

with the integrand depending on the parameter λ in an arbitrary manner. Assume that a function $y_0(x)$ is a solution of equation (3) for $\lambda = \lambda_0$. It is important to know the conditions under which equation (3) has solutions $y(x)$ close to $y_0(x)$ for λ close to λ_0 .

THEOREM. *Suppose that the function $K(x, t, y; \lambda)$ and its partial derivative $K'_y(x, t, y; \lambda)$ are continuous in all their arguments and a continuous function $y_0(x)$ is a solution of equation (3) for $\lambda = \lambda_0$,*

$$y_0(x) = \int_a^b K(x, t, y_0(t); \lambda_0) dt \quad (a \leq x \leq b). \quad (4)$$

If $\Lambda = 1$ is not a characteristic value of the kernel $K'_y(x, t, y_0(t); \lambda_0)$, then equation (3) has one and only one continuous solution $y = y(x, \lambda)$ close to $y_0(x)$ for λ close to λ_0 .

Remark. The solution of equation (3) can be sought in the form of expansion in powers of $(\lambda - \lambda_0)$:

$$y(x, \lambda) = y_0(x) + (\lambda - \lambda_0)y_1(x) + (\lambda - \lambda_0)^2y_2(x) + \dots$$

For $y_k(x)$ one obtains the triangular system of linear integral equations

$$y_k(x) = \int_a^b K'_y(x, t, y_0(t); \lambda_0)y_k(t) dt + \int_a^b F_k(x, t, y_0(t), \dots, y_{k-1}(t); \lambda_0) dt, \quad k = 1, 2, \dots,$$

which can be solved in consecutive manner with $y_0(x)$ being the solution of equation (4).

16.7-3. Bifurcation Points of Nonlinear Integral Equations.

Here it is assumed that for all values of the parameter λ , the integral equation (3) admits the trivial solution $y(x) \equiv 0$, i.e., $K(x, t, 0; \lambda) = 0$.

A value λ_* is called a *bifurcation point* for equation (3) if for any $\varepsilon > 0$ there is $\lambda \in (\lambda_* - \varepsilon, \lambda_* + \varepsilon)$ for which the equation has a nontrivial solution $y(x) = y(x, \lambda)$ that satisfies the inequality $\|y(x)\| < \varepsilon$.

In simple words, the meaning of a bifurcation point λ_* is that the number of solutions changes as λ crosses that point.

Example 1. Consider the linear integral equation with a continuous kernel

$$y(x) = \lambda \int_a^b K(x, t)y(t) dt. \quad (5)$$

For any λ , this equation admits the trivial solution $y(x) \equiv 0$.

Let $\lambda = \lambda_*$ be a characteristic value of equation (1) corresponding to a nontrivial solution $y_*(x)$. Since this solution is defined to within a constant coefficient, it can be made arbitrarily small in the norm of $C(a, b)$, i.e., for any $\varepsilon > 0$, there is a solution $y_*(x)$ such that

$$\|y_*(x)\| = \max_{a \leq x \leq b} |y_*(x)| < \varepsilon.$$

Thus, for any $\varepsilon > 0$ there is $\lambda \in (\lambda_* - \varepsilon, \lambda_* + \varepsilon)$ (in this case $\lambda = \lambda_*$) for which equation (5) has a nontrivial solution $y_*(x)$ satisfying the condition $\|y_*(x)\| < \varepsilon$. By definition, the characteristic value λ_* of the kernel $K(x, t)$ is a bifurcation point of equation (5).

Example 2. Consider the Hammerstein equation with a degenerate kernel containing a quadratic nonlinearity:

$$y(x) = \lambda \int_0^1 xt[y(t) + y^2(t)] dt. \tag{6}$$

For any value of the parameter λ , equation (6) the trivial solution $y(x) \equiv 0$.

Denote

$$A_1 = \int_0^1 ty(t) dt, \quad A_2 = \int_0^1 ty^2(t) dt. \tag{7}$$

With this notation, equation (6) can be rewritten as

$$y(t) = \lambda(A_1 + A_2)x. \tag{8}$$

Substituting (8) into (7), one obtains a second-order algebraic system for the determination of the coefficients A_1 and A_2 :

$$\begin{aligned} A_1 &= \frac{1}{3}\lambda(A_1 + A_2), \\ A_2 &= \frac{1}{4}\lambda^2(A_1 + A_2)^2. \end{aligned}$$

The solution of this system leads us to two solutions of the integral equation (6), one of which is trivial, $y(x) \equiv 0$, and the other has the form

$$y(x) = \frac{4(3 - \lambda)}{3\lambda}x. \tag{9}$$

Let us show that $\lambda = 3$ is a bifurcation point for equation (6). Indeed, for any $\varepsilon > 0$ there is $\lambda \in (3 - \varepsilon, 3 + \varepsilon)$ (for instance, any $\lambda \neq 3$ from this interval) for which equation (6) has the nontrivial solution (9) satisfying the condition

$$\|y_*(x)\| = \max_{0 \leq x \leq 1} \left| \frac{4(3 - \lambda)}{3\lambda} \right| < \varepsilon.$$

THEOREM 1. Let λ_* be a bifurcation point for the nonlinear integral equation (3). Then 1 is a characteristic value of the kernel $L(x, t) = K'_y(x, t, 0; \lambda_*)$. (It is assumed that the integrand $K(x, t, y; \lambda)$ and its partial derivative $K'_y(x, t, y; \lambda)$ are continuous in their arguments.)

In other words, a necessary condition for the existence of a bifurcation point $\lambda = \lambda_*$ for equation (3) is the existence of a nontrivial solution of the homogeneous linear integral equation

$$u(x) = \int_a^b K'_y(x, t, 0; \lambda_*)u(t) dt \quad (a \leq x \leq b). \tag{10}$$

Theorem 1 suggests which values $\lambda = \lambda_*$ can be expected to be bifurcation points.

THEOREM 2. Let 1 be a characteristic value of the kernel $L(x, t) = K'_y(x, t, 0; \lambda_*)$ and let the multiplicity of this value be equal to 1 (is odd). Then λ_* is a bifurcation point of the nonlinear integral equation (3).

Example 3. Consider the Nekrasov equation

$$y(x) = \lambda \int_0^{2\pi} \frac{K(x, t) \sin y(t)}{1 + \int_0^t \sin y(s) ds} dt, \quad K(x, t) = \frac{1}{3} \sum_{n=1}^{\infty} \frac{\sin(nx) \sin(nt)}{n},$$

that describes waves on the surface of ideal incompressible fluid. For all values of the parameter λ , this equation admits the trivial solution $y(x) \equiv 0$. Its bifurcation points correspond to the values $\lambda = \lambda_*$ for which waves are produced.

The linearized equation (10) in this case can be written as

$$u(x) = \lambda_* \int_0^{2\pi} K(x, t)u(t) dt.$$

Its characteristic values and the corresponding eigenfunctions have the form

$$\lambda_n = 3n, \quad u_n(x) = \sin(nx).$$

All characteristic values are simple, and therefore, these and only these are bifurcation points of the Nekrasov equation.

THEOREM 3. *Let the kernel $K(x, t)$ be continuous and positive definite and let the function $\Phi(t, y)$ and its derivative $\Phi'_y(t, y)$ be continuous, with $\Phi(t, 0) \equiv 0$. Then the bifurcation points of the Hammerstein equation (2) with parameter λ coincide with the characteristic values of the kernel $L(x, t) = K(x, t)\Phi'_y(t, 0)$.*

Example 4. For the Hammerstein equation with degenerate kernel (6), we have

$$K(x, t) = xt, \quad \Phi(t, y) = y + y^2, \quad \Phi'_y(t, y) = 1 + 2y, \quad \Phi(t, 0) \equiv 0.$$

Therefore, the assumptions of Theorem 3 hold. Since $\Phi'_y(t, 0) = 1$, we see that the bifurcation points of equation (6) coincide with the characteristic values of the kernel $K(x, t) = xt$,

$$u(x) = \lambda_* \int_0^1 xtu(t) dt.$$

Substituting $u(x) = A\lambda_*x$ into this equation, we obtain the unique characteristic value $\lambda_* = 3$, which is a bifurcation point of the Hammerstein equation.

Remark. The bifurcation point $\lambda_* = 3$ in Example 2 was obtained in a more difficult way, by direct examination of equation (6).

© References for Section 16.7: M. A. Krasnosel'skii (1964), M. G. Krein (1972), M. L. Krasnov (1975), P. P. Zabreyko, A. I. Koshelev, et al. (1975), R. Precup (2006).

Chapter 17

Methods for Solving Multidimensional Mixed Integral Equations

17.1. Some Definition and Remarks

17.1-1. Basic Classes of Functions.

Integral equations containing both the Volterra kernels (see Subsection 10.1-1) and the Fredholm kernels (see Subsections 12.1-1 and 12.1-2) are called *mixed integral equations*. Such integral equations arise in applications and are a fairly new object of mathematical studies. So far, no definite classification of such equations has been given, and such a classification is likely to be vast and ramified. Here, we consider some integral equations and related problems that have been studied in more detail.

Mixed integral equations are multidimensional (at least two-dimensional). In the integral terms of such equations with Volterra kernels, the unknown function of several variables is integrated in the variable that has the meaning of the time; and in the integral terms with Fredholm kernels, the integration of the same unknown function is over some (one- or multi-dimensional) domain.

Let us describe the main classes of multidimensional real-valued functions that appear in mixed integral equations.

For a bounded closed domain Ω in \mathbb{R}^n , the set $L_2(\Omega)$ consists of all real-valued functions $f(\vec{x})$ defined in Ω and having their squared absolute value $|f(\vec{x})|^2$ integrable in Ω . The set $L_2(\Omega)$ is a Hilbert space with the following scalar product and the norm (see Supplement 12.5-2):

$$(f, g) = \int_{\Omega} f(\vec{x})g(\vec{x}) d\Omega_x, \quad \|f\| = (f, f)^{1/2} = \sqrt{\int_{\Omega} |f(\vec{x})|^2 d\Omega_x},$$

where $\vec{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$.

Example.

1°. If $n = 2$ and Ω is the ring $\omega = \{a \leq r \leq b, 0 \leq \varphi \leq 2\pi\}$, then

$$L_2(\Omega) \equiv L_2(\omega), \quad f(\vec{x}) \equiv f(r, \varphi), \quad \int_{\Omega} \equiv \int_{\omega} \equiv \int_a^b \int_0^{2\pi}, \quad d\Omega \equiv r dr d\varphi.$$

2°. The subspace of $L_2(\omega)$ consisting of functions that depend only on the radial coordinate r is denoted by $\widehat{L}_2(\omega)$. In this case,

$$f(\vec{x}) \equiv f(r), \quad \int_{\Omega} f(\vec{x}) d\Omega = \int_{\omega} f(r) r dr d\varphi = 2\pi \int_a^b r f(r) dr,$$

The space $\widehat{L}_2(\omega)$ is a Hilbert space with the scalar product

$$(f, g) = 2\pi \int_a^b f(r)g(r)r dr.$$

Frequently the space $\widehat{L}_2(\omega)$ is considered as the space of functions of the form $\widehat{f}(r) = \sqrt{2\pi}f(r)$ for which the scalar product has the like form but without factor 2π ,

$$(\widehat{f}, \widehat{g}) = \int_a^b \widehat{f}(r)\widehat{g}(r)r dr.$$

In what follows we will use the latter definition of the scalar product. To simplify notation we will omit the hat symbol over functions.

Let $F(t)$ be a function of the real argument $t \in [\tau_0, T]$ with values in a Banach space B , i.e., $F: [\tau_0, T] \rightarrow B$. The function $F(t)$ is called *continuous* on $[\tau_0, T]$ if for any $t \in [\tau_0, T]$, we have

$$\|F(t) - F(t_1)\|_B \rightarrow 0 \quad \text{as } t_1 \rightarrow t,$$

where $\|\cdot\|_B$ is the norm in B . The space of such continuous functions is denoted by $C([\tau_0, T], B)$.

For example, if $B = L_2(\Omega)$ with the above norm $\|\cdot\|$, we can consider a function $y(\vec{x}, t)$ such that for each $t \in [\tau_0, T]$ its value belongs to $L_2(\Omega)$. Regarded as a function of t with values in $L_2(\Omega)$, such a function is called *continuous* in t on the interval $[\tau_0, T]$ if for any $t \in [\tau_0, T]$, we have

$$\|y(\vec{x}, t) - y(\vec{x}, t_1)\| \rightarrow 0 \quad \text{as } t_1 \rightarrow t.$$

Accordingly, the space of such functions is denoted by $C([\tau_0, T], L_2(\Omega))$.

For a function $y(\vec{x}, t) \in C([\tau_0, T], L_2(\Omega))$, the following properties hold:

- 1) the norm $\|y(\vec{x}, t)\|$ is continuous in $t \in [\tau_0, T]$;
- 2) for any $f(\vec{x}) \in L_2(\Omega)$, the scalar product $(y(\vec{x}, t), f(\vec{x}))$ is continuous in $t \in [\tau_0, T]$;
- 3) $y(\vec{x}, t) \in L_2(\Omega \times (\tau_0, T))$ if the interval (τ_0, T) is finite.

In what follows, we consider the cases of Ω being a finite interval, a circle, or an arbitrary closed bounded set.

17.1-2. Mixed Equations on a Finite Interval.

1°. For continuous functions of $t \in [\tau_0, T]$ with values in $L_2[a, b]$, the mixed two-dimensional integral equation with symmetric Fredholm kernel has the form

$$\begin{aligned} \sigma(t) \left[y(x, t) - \int_{\tau_0}^t V_1(t, \tau) y(x, \tau) d\tau \right] + \int_a^b F(x, \xi) y(\xi, t) d\xi \\ - \int_{\tau_0}^t V_2(t, \tau) \int_a^b F(x, \xi) y(\xi, \tau) d\xi d\tau = f(x, t), \quad a \leq x \leq b, \quad \tau_0 \leq t \leq T, \end{aligned} \quad (1)$$

where $\sigma(t)$ is a known continuous positive function; $y(x, t)$ is the unknown function of class $C([\tau_0, T], L_2[a, b])$; $f(x, t) \in C([\tau_0, T], L_2[a, b])$ is a given function; $V_1(t, \tau)$ and $V_2(t, \tau)$ are Volterra kernels (see Subsection 10.1-1); and $F(x, \xi)$ is a Fredholm kernel, so that

$$\int_a^b \int_a^b F^2(x, \xi) dx d\xi = B^2 < \infty.$$

Assume, in addition, that the kernel $F(x, \xi)$ is symmetric and positive definite. Such a kernel is also called a *Hilbert-Schmidt kernel* (see also Subsection 13.6-2).

2°. For functions of class $C([\tau_0, T], L_2[a, b])$, a mixed two-dimensional integral equation with a Schmidt kernel has the form

$$\begin{aligned} \sigma(t) \left[y(x, t) - \int_{\tau_0}^t V_1(t, \tau) y(x, \tau) d\tau \right] + \int_a^b S(x, \xi) y(\xi, t) d\xi \\ - \int_{\tau_0}^t V_2(t, \tau) \int_a^b S(x, \xi) y(\xi, \tau) d\xi d\tau = \frac{f(x, t)}{h(x)}, \end{aligned} \quad (2)$$

$$S(x, \xi) = \frac{F(x, \xi)}{h(x)}, \quad a \leq x \leq b, \quad \tau_0 \leq t \leq T,$$

where $h(x) > 0$ is a given function of class $L_2[a, b]$, and the other quantities are similar to those introduced for equation (1). The kernel $S(x, \xi)$ is called a *Schmidt kernel*. This kernel is nonsymmetric, but has all the properties of symmetric kernels. Equation (2) is often written in the following equivalent form:

$$\begin{aligned} \sigma(t)h(x) \left[y(x, t) - \int_{\tau_0}^t V_1(t, \tau)y(x, \tau) d\tau \right] + \int_a^b F(x, \xi)y(\xi, t) d\xi \\ - \int_{\tau_0}^t V_2(t, \tau) \int_a^b F(x, \xi)y(\xi, \tau) d\xi d\tau = f(x, t), \quad a \leq x \leq b, \quad \tau_0 \leq t \leq T. \end{aligned} \quad (3)$$

Together with equations (1), (2), we consider some problems that contain, in addition to (1) or (2), some auxiliary integral conditions on the unknown function. Such conditions are introduced in some cases if the right hand-side of the equation is not determined completely. Often, such conditions have the form

$$\int_a^b y(\xi, t) d\xi = M_1(t), \quad \int_a^b \left(\xi - \frac{a+b}{2} \right) y(\xi, t) d\xi = M_2(t). \quad (4)$$

17.1-3. Mixed Equation on a Ring-Shaped (Circular) Domain.

1°. A mixed two-dimensional integral equation with Fredholm kernel for functions of class $C([\tau_0, T], \widehat{L}_2(\omega))$ has the form

$$\begin{aligned} \sigma(t) \left[y(r, t) - \int_{\tau_0}^t V_1(t, \tau)y(r, \tau) d\tau \right] + \int_a^b F_\omega(r, \rho)y(\rho, t)\rho d\rho \\ - \int_{\tau_0}^t V_2(t, \tau) \int_a^b F_\omega(r, \rho)y(\rho, \tau)\rho d\rho d\tau = f(r, t), \quad a \leq r \leq b, \quad \tau_0 \leq t \leq T, \end{aligned} \quad (5)$$

where ω is the ring of internal radius a and external radius b (for $a = 0$, the domain ω is the circle of radius b); $\sigma(t)$ is a given positive continuous function; $y(r, t)$ is the unknown function of class $C([\tau_0, T], \widehat{L}_2(\omega))$; $f(r, t)$ is a given right-hand side of class $C([\tau_0, T], \widehat{L}_2(\omega))$; $V_1(t, \tau)$ and $V_2(t, \tau)$ are Volterra kernels (see Subsection 10.1-1); and $F_\omega(r, \rho)$ is a Fredholm kernel, so that

$$\int_a^b \int_a^b F_\omega^2(r, \rho)r\rho dr d\rho = B_\omega^2 < \infty.$$

Assume, in addition, that the kernel $F_\omega(r, \rho)$ is symmetric and positive definite (see also Subsection 13.6-2), i.e.,

$$F_\omega(r, \rho) = F_\omega(\rho, r), \quad \int_a^b \int_a^b F_\omega(r, \rho)\varphi(r)\varphi(\rho) dr d\rho \geq 0,$$

and the second relation holds as equality only for $\varphi(r) = 0$. As above, a symmetric positive Fredholm kernel will be called a Hilbert–Schmidt kernel.

2°. A mixed two-dimensional integral equation with a Schmidt kernel for functions of class $C([\tau_0, T], \widehat{L}_2(\omega))$ has the form

$$\begin{aligned} \sigma(t) \left[y(r, t) - \int_{\tau_0}^t V_1(t, \tau)y(r, \tau) d\tau \right] + \int_a^b S_\omega(r, \rho)y(\rho, t)\rho d\rho \\ - \int_{\tau_0}^t V_2(t, \tau) \int_a^b S_\omega(r, \rho)y(\rho, \tau)\rho d\rho d\tau = \frac{f(r, t)}{h(r)}, \end{aligned} \quad (6)$$

$$S_\omega(r, \rho) = \frac{F_\omega(r, \rho)}{h(r)}, \quad a \leq r \leq b, \quad \tau_0 \leq t \leq T,$$

where $h(r) > 0$ is a given function in $\widehat{L}_2(\omega)$, and the other notations are similar to those introduced for equation (5). The kernel $S_\omega(r, \rho)$ is a Schmidt kernel, which possesses all the properties of Hilbert–Schmidt kernels. Equation (6) is often written in the following equivalent form:

$$\begin{aligned} \sigma(t)h(r) \left[y(r, t) - \int_{\tau_0}^t V_1(t, \tau)y(r, \tau) d\tau \right] + \int_a^b F_\omega(r, \rho)y(\rho, t)\rho d\rho \\ - \int_{\tau_0}^t V_2(t, \tau) \int_a^b F_\omega(r, \rho)y(\rho, \tau)\rho d\rho d\tau = f(r, t), \quad a \leq r \leq b, \quad \tau_0 \leq t \leq T. \end{aligned} \tag{7}$$

Together with equations (5)–(7), we consider some problems with an auxiliary integral condition on the unknown function. Such conditions are introduced if there is not enough information about the right-hand side of the equation. For equations (5)–(7), this condition often has the form

$$\int_a^b y(\rho, t)\rho d\rho = M(t). \tag{8}$$

17.1-4. Mixed Equations on a Closed Bounded Set.

1°. A mixed multi-dimensional integral equation with a symmetric Fredholm kernel for functions of class $C([\tau_0, T], L_2(\Omega))$ has the form

$$\begin{aligned} \sigma(t)(\mathbf{I} - \mathbf{V}_1)y(\vec{x}, t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{F}y(\vec{x}, t) = f(\vec{x}, t), \quad \vec{x} \in \Omega, \quad \tau_0 \leq t \leq T, \\ \mathbf{F}y(\vec{x}, t) = \int_\Omega F_\Omega(\vec{x}, \vec{\xi})y(\vec{\xi}, t) d\Omega_\xi, \quad \mathbf{V}_p y(\vec{x}, t) = \int_{\tau_0}^t V_p(t, \tau)y(\vec{x}, \tau) d\tau, \end{aligned} \tag{9}$$

where $\vec{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$; Ω is a closed bounded set in \mathbb{R}^n ; $\sigma(t)$ is a continuous function of t on $[\tau_0, T]$; $y(\vec{x}, t) \in C([\tau_0, T], L_2(\Omega))$ is the unknown function; $f(\vec{x}, t) \in C([\tau_0, T], L_2(\Omega))$ is a given right-hand side of the equation; \mathbf{I} is the identity operator; \mathbf{V}_p ($p = 1, 2$) are Volterra integral operators with continuous or polar kernels $V_p(t, \tau)$; and \mathbf{F} is a Fredholm integral operator, which is a compact operator from $L_2(\Omega)$ to $L_2(\Omega)$ (see Supplement 12.5-3). Its properties are determined by the kernel $F_\Omega(\vec{x}, \vec{\xi})$, which is assumed to satisfy the condition

$$\int_\Omega \int_\Omega F_\Omega^2(\vec{x}, \vec{\xi}) d\Omega_x d\Omega_\xi = B_\Omega^2 < \infty. \tag{10}$$

Relation (10) is a sufficient condition for the compactness of the integral operator \mathbf{F} .

If the kernel of an integral operator satisfies the relation

$$F_\Omega(\vec{x}, \vec{\xi}) = F_\Omega(\vec{\xi}, \vec{x}), \tag{11}$$

then this operator is self-adjoint.

If, moreover,

$$\int_\Omega \int_\Omega F_\Omega(\vec{x}, \vec{\xi})\varphi(\vec{x})\varphi(\vec{\xi}) d\Omega_x d\Omega_\xi \geq 0, \tag{12}$$

and (12) holds as equality only for $\varphi(\vec{x}) = 0$, then the integral operator is called positive definite.

Compact self-adjoint positive definite operators are called *Hilbert–Schmidt operators*.

2°. A mixed multi-dimensional integral equation with a Schmidt operator for functions of class $C([\tau_0, T], L_2(\Omega))$ has the form

$$\begin{aligned} \sigma(t)(\mathbf{I} - \mathbf{V}_1)y(\vec{x}, t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{S}y(\vec{x}, t) &= \frac{f(\vec{x}, t)}{h(\vec{x})}, \\ \mathbf{S}y(\vec{x}, t) &= \int_{\Omega} S_{\Omega}(\vec{x}, \vec{\xi})y(\vec{\xi}, t) d\Omega_{\xi}, \quad S_{\Omega}(\vec{x}, \vec{\xi}) = \frac{F_{\Omega}(\vec{x}, \vec{\xi})}{h(\vec{x})}, \\ \mathbf{V}_p y(\vec{x}, t) &= \int_{\tau_0}^t V_p(t, \tau)y(\vec{x}, \tau) d\tau, \quad \vec{x} \in \Omega, \quad \tau_0 \leq t \leq T, \end{aligned} \quad (13)$$

where $h(\vec{x}) > 0$ is a given function in $L_2(\Omega)$; \mathbf{S} is a Schmidt integral operator; and the other notations are similar to those introduced for equation (12). Equation (13) is often written in the following equivalent form:

$$\begin{aligned} \sigma(t)h(\vec{x})(\mathbf{I} - \mathbf{V}_1)y(\vec{x}, t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{F}y(\vec{x}, t) &= f(\vec{x}, t), \quad \vec{x} \in \Omega, \quad \tau_0 \leq t \leq T, \\ \mathbf{F}y(\vec{x}, t) &= \int_{\Omega} F_{\Omega}(\vec{x}, \vec{\xi})y(\vec{\xi}, t) d\Omega_{\xi}, \quad \mathbf{V}_p y(\vec{x}, t) = \int_{\tau_0}^t V_p(t, \tau)y(\vec{x}, \tau) d\tau. \end{aligned} \quad (14)$$

Together with equations (9), (13), and (14), we consider some problems with auxiliary conditions on the unknown function. Such conditions are introduced if there is not enough information about the right-hand side of the equation. Such conditions usually have the form

$$\int_{\Omega} y(\vec{x}, t)f_i(\vec{x}) d\Omega_x = M_i(t), \quad i = 1, \dots, N, \quad (15)$$

where $f_i(\vec{x})$ is a system of N linearly independent functions of class $L_2(\Omega)$.

Remark 1. Any equation with a Schmidt kernel (integral operator) can always be reduced (by changing the variables) to an equation with a symmetric Hilbert–Schmidt kernel (self-adjoint integral operator).

Remark 2. A compact operator is a generalization of a Fredholm integral operator. Equations with compact operators are studied in the framework of the Riesz–Schauder theory.

Remark 3. A compact self-adjoint operator is a generalization of a Fredholm integral operator with a symmetric kernel. If its kernel is positive definite, then the corresponding operator is also positive definite (see Supplement 12.5-3). Equations with compact self-adjoint and positive definite operators are studied in the framework of the Hilbert–Schmidt theory.

© References for Section 17.1: E. Goursat (1923), F. Riesz and B. Sz.-Nagy (1955), V. S. Vladimirov (1981), V. M. Aleksandrov and S. M. Mkhitarian (1983), N. Kh. Arutyunyan, A. V. Manzhurov, and V.E. Naumov (1991), A. N. Kolmogorov and S. V. Fomin (1999), A. V. Manzhurov (2001, 2005).

17.2. Methods of Solution of Mixed Integral Equations on a Finite Interval

17.2-1. Equation with a Hilbert–Schmidt Kernel and a Given Right-Hand Side.

Consider the mixed integral equation (1) of Subsection 17.1-2 with a Hilbert–Schmidt kernel. By changing the variables, this equation can always be reduced to a similar equation with the parameters $a = -1$, $b = 1$, $\tau_0 = 1$:

$$\begin{aligned} \sigma(t) \left[y(x, t) - \int_1^t V_1(t, \tau)y(x, \tau) d\tau \right] + \int_{-1}^1 F(x, \xi)y(\xi, t) d\xi \\ - \int_{\tau_0}^t V_2(t, \tau) \int_{-1}^1 F(x, \xi)y(\xi, \tau) d\xi d\tau = f(x, t), \quad -1 \leq x \leq 1, \quad 1 \leq t \leq T. \end{aligned} \quad (1)$$

Suppose that the right-hand side $f(x, t)$ of (1) is known and we have to find the function $y(x, t)$. Here $f(x, t)$ and $y(x, t)$ are functions of class $C([1, T], L_2(-1, 1))$.

Let us seek a solution of equation (1) in the form of a series

$$y(x, t) = \sum_{k=1}^{\infty} y_k(t) \varphi_k(x), \quad (2)$$

where $\varphi_k(x)$ are eigenfunctions of the kernel $F(x, \xi)$ corresponding to the eigenvalues $\mu_k > 0$, i.e.,

$$\int_{-1}^1 F(x, \xi) \varphi_k(\xi) d\xi = \mu_k \varphi_k(x), \quad k = 1, 2, \dots \quad (3)$$

The representation of a solution in the form (2) is possible, since eigenfunctions of the kernel $F(x, \xi)$ form a complete orthonormal system of functions in $L_2[-1, 1]$ (a basis in $L_2[-1, 1]$; see Subsection 13.6-1 and Supplement 12.5-3). For the same reason, the right-hand side of the equation can be represented in the form

$$f(x, t) = \sum_{k=1}^{\infty} f_k(t) \varphi_k(x), \quad f_k(t) = \int_{-1}^1 f(x, t) \varphi_k(x) dx. \quad (4)$$

Substituting (2) into (1) and taking into account (3) and (4), we obtain the following sequence of Volterra equations for the unknown functions $y_k(t)$:

$$y_k(t) - \int_1^t V_k(t, \tau) y_k(\tau) d\tau = \delta_k(t), \quad \delta_k(t) = \frac{f_k(t)}{\sigma(t) + \mu_k}, \quad (5)$$

$$V_k(t, \tau) = \frac{\sigma(t) V_1(t, \tau) + \mu_k V_2(t, \tau)}{\sigma(t) + \mu_k}, \quad k = 1, 2, \dots, \quad (6)$$

where $V_k(t, \tau)$ are Volterra kernels which belong to the same class of functions as the kernels $V_1(t, \tau)$, $V_2(t, \tau)$, since $\mu_k \rightarrow 0$ as $k \rightarrow \infty$.

A solution of the infinite sequence of Volterra equations (5) can be constructed by analytical and numerical methods of Chapter 11. This solution can be written in the form

$$y_k(t) = \delta_k(t) + \int_1^t R_k(t, \tau) \delta_k(\tau) d\tau, \quad (7)$$

where $R_k(t, \tau)$ is the resolvent of the kernel $V_k(t, \tau)$.

Thus, the desired solution has been constructed. The series (2) converges in $L_2[-1, 1]$ uniformly in $t \in [1, T]$, and its sum is a continuous function of $t \in [1, T]$ with values in $L_2[-1, 1]$.

In order to justify the above method of constructing a solution, it remains to construct the eigenfunctions and calculate the eigenvalues of the Hilbert–Schmidt integral operator. Let us represent the k th eigenfunction as a series in terms of any basis $p_i(x)$ of $L_2[-1, 1]$. For definiteness, we take the orthonormal Legendre polynomials $P_{i-1}^*(x)$ as the basis. Then

$$\varphi_k(x) = \sum_{i=1}^{\infty} \varphi_{i(k)} p_i(x), \quad p_i(x) = P_{i-1}^*(x). \quad (8)$$

Let us expand the Hilbert–Schmidt kernel in double series with respect to the chosen basis:

$$F(x, \xi) = \sum_{m=1}^{\infty} \sum_{i=1}^{\infty} F_{mn} p_m(x) p_n(\xi), \quad (9)$$

$$F_{mn} = \int_{-1}^1 \int_{-1}^1 F(x, \xi) p_m(x) p_n(\xi) dx d\xi, \quad F_{mn} = F_{nm}.$$

Substituting (8) and (9) into (3), we obtain a linear system of algebraic equations for the determination of eigenvalues and eigenfunction expansion coefficients. This algebraic system has a symmetric matrix and can be written as follows:

$$\sum_{n=1}^{\infty} F_{mn} \varphi_{n(k)} = \mu_k \varphi_{m(k)}, \quad m = 1, 2, \dots \quad (10)$$

Of course, for practical calculations, one has to limit the number of expansion terms. For instance, taking N orthonormal Legendre polynomials, we obtain the N th approximation of the solution. In this case, to construct eigenfunctions and eigenvalues of the Hilbert–Schmidt kernel, it is necessary to find eigenvalues and orthonormal eigenvectors of the matrix

$$[F_{NN}] = \begin{pmatrix} F_{11} & F_{12} & F_{13} & \cdots & F_{1N} \\ F_{12} & F_{22} & F_{23} & \cdots & F_{2N} \\ F_{13} & F_{23} & F_{33} & \cdots & F_{3N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ F_{1N} & F_{2N} & F_{3N} & \cdots & F_{NN} \end{pmatrix}. \quad (11)$$

Eigenvalues of the matrix (11) give approximate values of the first N eigenvalues of the Hilbert–Schmidt kernel, and the components of orthonormal eigenvectors of the matrix give expansion coefficients of the first N eigenfunctions of the Hilbert–Schmidt kernel with respect to N orthonormal Legendre polynomials.

17.2-2. Equation with Hilbert–Schmidt Kernel and Auxiliary Conditions.

Consider equation (1) with the right-hand side of the form $f(x, t) = \alpha_1(t) + \alpha_2(t)x - g(x, t)$ and two auxiliary integral conditions of the form (4) of Subsection 17.1-2. The problem is to find a solution of the mixed integral equation

$$\begin{aligned} \sigma(t) \left[y(x, t) - \int_1^t V_1(t, \tau) y(x, \tau) d\tau \right] + \int_{-1}^1 F(x, \xi) y(\xi, t) d\xi \\ - \int_1^t V_2(t, \tau) \int_{-1}^1 F(x, \xi) y(\xi, \tau) d\xi d\tau = \alpha_1(t) + \alpha_2(t)x - g(x, t), \end{aligned} \quad (12)$$

$$-1 \leq x \leq 1, \quad 1 \leq t \leq T$$

with the auxiliary conditions

$$\int_{-1}^1 y(\xi, t) d\xi = M_1(t), \quad \int_{-1}^1 \xi y(\xi, t) d\xi = M_2(t), \quad (13)$$

regarding $y(x, t)$, $\alpha_1(t)$, and $\alpha_2(t)$ as unknown. All other functions in (12) are assumed given and $g(x, t)$ is of class $C([1, T], L_2[-1, 1])$.

Note that the Hilbert space $L_2[-1, 1]$ can be represented as the direct sum of its orthogonal subspaces, $L_2[-1, 1] = L_2^{\circ}[-1, 1] \oplus L_2^*[-1, 1]$ (see Supplement 12.5-3.), where $L_2^{\circ}[-1, 1]$ is the Euclidean space with the basis $p_1(x) = P_0^*(x) = 1/\sqrt{2}$, $p_2(x) = P_1^*(x) = \sqrt{3}/2 x$, and $L_2^*[-1, 1]$ is the Hilbert space with the basis $p_k(x) = P_{k-1}^*(x)$ ($k = 3, 4, \dots$).

Note also that the integrand and the right-hand side can be represented as a sum of functions continuous in $t \in [1, T]$ with values in $L_2^{\circ}[-1, 1]$ and $L_2^*[-1, 1]$, respectively, i.e.,

$$y(x, t) = y^{\circ}(x, t) + y^*(x, t), \quad f(x, t) = f^{\circ}(x, t) + f^*(x, t), \quad (14)$$

where

$$\begin{aligned}
 y^\circ(x, t) &= y_1^\circ(t)p_1(x) + y_2^\circ(t)p_2(x), \quad y_1^\circ(t) = \frac{1}{\sqrt{2}}M_1(t), \quad y_2^\circ(t) = \sqrt{\frac{3}{2}}M_2(t) \\
 f^\circ(x, t) &= [\sqrt{2}\alpha_1(t) - g_1^\circ(t)]p_1(x) + \left[\sqrt{\frac{2}{3}}\alpha_2(t) - g_2^\circ(t)\right]p_2(x), \\
 f^*(x, t) &= -g^*(x, t), \quad g(x, t) = g^\circ(x, t) + g^*(x, t), \quad g^\circ(x, t) = g_1^\circ(t)p_1(x) + g_2^\circ(t)p_2(x), \\
 g_1^\circ(t) &= \frac{1}{\sqrt{2}} \int_{-1}^1 g(x, t) dx, \quad g_2^\circ(t) = \sqrt{\frac{3}{2}} \int_{-1}^1 g(x, t) dx.
 \end{aligned}
 \tag{15}$$

Note that in the representation (14) of $y(x, t)$, the term $y^\circ(x, t)$ is known and is determined by the auxiliary conditions; and the term $y^*(x, t)$ is to be found. Conversely, for the right-hand side, $f^\circ(x, t)$ is the unknown and $f^*(x, t)$ is determined by the function $g(x, t)$. These features allow us to classify the resulting problem as a special case of the general projection problem formulated and solved in Subsection 17.4-3.

Applying the general method to the present case, we introduce an operator of orthogonal projection that maps $L_2[-1, 1]$ onto $L_2^\circ[-1, 1]$:

$$\mathbf{P}^\circ f(x, t) = \int_{-1}^1 f(\xi, t)[p_1(x)p_1(\xi) + p_2(x)p_2(\xi)] d\xi.
 \tag{16}$$

Obviously, $\mathbf{P}^* = \mathbf{I} - \mathbf{P}^\circ$ is the orthogonal projector of $L_2[-1, 1]$ onto $L_2^*[-1, 1]$. Moreover, the following relations hold:

$$\begin{aligned}
 \mathbf{P}^\circ y(x, t) &= y^\circ(x, t), \quad \mathbf{P}^* y(x, t) = y^*(x, t), \\
 \mathbf{P}^\circ f(x, t) &= f^\circ(x, t), \quad \mathbf{P}^* f(x, t) = f^*(x, t).
 \end{aligned}
 \tag{17}$$

Following the method of Section 17.4, let us apply the projection operator \mathbf{P}^* to equation (12). As a result, for $y^*(x, t)$ we obtain an integral equation on the space $L_2^*[-1, 1]$ with a known right-hand side:

$$\begin{aligned}
 \sigma(t) &\left[y^*(x, t) - \int_1^t V_1(t, \tau)y^*(x, \tau) d\tau \right] \\
 &+ \int_{-1}^1 F^*(x, \xi)y^*(\xi, t) d\xi - \int_1^t V_2(t, \tau) \int_{-1}^1 F^*(x, \xi)y^*(\xi, \tau) d\xi d\tau \\
 &= -g^*(x, t) - \int_{-1}^1 F^*(x, \xi)y^\circ(\xi, t) d\xi + \int_1^t V_2(t, \tau) \int_{-1}^1 F^*(x, \xi)y^\circ(\xi, \tau) d\xi d\tau, \\
 &-1 \leq x \leq 1, \quad 1 \leq t \leq T,
 \end{aligned}
 \tag{18}$$

where the kernel of the integral equation

$$F^*(x, \xi) = F(x, \xi) - \int_{-1}^1 F(s, \xi)[p_1(x)p_1(s) + p_2(x)p_2(s)] ds
 \tag{19}$$

is of Fredholm type and, moreover, is symmetric and positive definite. Let us construct a solution of equation (18) in the form of a series with eigenfunctions of the kernel (19). These eigenfunction form a basis in the Hilbert space $L_2^*[-1, 1]$. We start, however, with the construction of the said eigenfunctions.

Let $\varphi_k^*(x)$ be eigenfunctions and μ_k^* the corresponding eigenvalues of the kernel $F^*(x, \xi)$. Then

$$\int_{-1}^1 F^*(x, \xi)\varphi_k^*(\xi) d\xi = \mu_k^*\varphi_k^*(x), \quad k = 3, 4, \dots
 \tag{20}$$

Let us represent the eigenfunction $\varphi_k^*(x)$ in the form of a series with respect to the basis $p_i(x)$ ($i \geq 3$):

$$\varphi_k^*(x) = \sum_{i=3}^{\infty} \varphi_{i(k)}^* p_i(x), \quad p_i(x) = P_{i-1}^*(x), \quad k = 3, 4, \dots \tag{21}$$

Using (9) and (19), we obtain a double series expansion for the kernel $F^*(x, \xi)$:

$$F^*(x, \xi) = \sum_{m=3}^{\infty} \sum_{n=3}^{\infty} F_{mn} p_m(x) p_n(\xi) + \sum_{n=3}^{\infty} F_{1n} p_n(x) p_1(\xi) + \sum_{n=3}^{\infty} F_{2n} p_n(x) p_2(\xi). \tag{22}$$

Note that the coefficients of the expansion of the kernel $F^*(x, \xi)$ in (22) coincide with the coefficients of the expansion of the kernel $F(x, \xi)$, which allows us to avoid recalculation of the coefficients of the new problem and use the already available data.

Substituting (21) and (22) into (20), we obtain the following infinite system of algebraic equations (with a symmetric matrix) for the determination of eigenvalues and eigenfunction expansion coefficients:

$$\sum_{n=3}^{\infty} F_{mn} \varphi_{n(k)}^* = \mu_k^* \varphi_{m(k)}^*, \quad m = 3, 4, \dots \tag{23}$$

Now, let us construct a solution of equation (18). For this purpose, we represent the functions $y^*(x, t)$ and $g^*(x, t)$ in the form of series with respect to eigenfunctions of the kernel $F^*(x, \xi)$:

$$y^*(x, t) = \sum_{k=3}^{\infty} y_k^*(t) \varphi_k^*(x), \quad g^*(x, t) = \sum_{k=3}^{\infty} g_k^*(t) \varphi_k^*(x), \quad g_k^*(t) = \int_{-1}^1 g^*(x, t) \varphi_k^*(x) dx, \tag{24}$$

and substitute these into (18). Then, taking into account (15), (19)–(22), we obtain the following sequence of independent Volterra equations:

$$\begin{aligned} y_k^*(t) - \int_1^t V_k^*(t, \tau) y_k^*(\tau) d\tau &= f_k^*(t), \quad V_k^*(t, \tau) = \frac{\sigma(t) V_1(t, \tau) + \mu_k^* V_2(t, \tau)}{\sigma(t) + \mu_k^*}, \\ f_k^*(t) &= -\frac{1}{\sigma(t) + \mu_k^*} \left[g_k^*(t) + \sum_{i=1}^2 F_{k(i)} y_i^{\circ}(t) - \int_1^t V_2(t, \tau) \sum_{i=1}^2 F_{k(i)} y_i^{\circ}(\tau) d\tau \right], \\ F_{k(i)} &= \sum_{n=3}^{\infty} F_{in} \varphi_{n(k)}^*, \quad i = 1, 2, \quad k = 3, 4, \dots \end{aligned} \tag{25}$$

Resolving (25) with respect to $y_k^*(t)$ by the methods of Chapter 11, we obtain

$$y_k^*(t) = f_k^*(t) + \int_1^t R_k^*(t, \tau) f_k^*(\tau) d\tau, \tag{26}$$

where $R_k^*(t, \tau)$ is the resolvent of the kernel $V_k^*(t, \tau)$.

Now, in view of (24)–(26), the function $y^*(x, t)$ has been determined, and therefore, the function $y(x, t)$ has also been found, since $y^{\circ}(x, t)$ is known by assumption (see (14) and (15)).

Before passing to the determination of the other unknown quantities of the problem, we make some remarks that may be useful in practice.

For practical calculations one should restrict the number of expansion terms. For instance, taking the orthonormal Legendre polynomials from the third to the N th, we obtain the N th approximation of the desired solution. In this case, for the construction of eigenvalues and eigenfunctions of the

Hilbert–Schmidt kernel $F^{*}(x, \xi)$ one should find the eigenvalues and orthonormal eigenvectors of the matrix

$$[F_{NN}^{*}] = \begin{pmatrix} F_{33} & F_{34} & F_{35} & \cdots & F_{3N} \\ F_{34} & F_{44} & F_{45} & \cdots & F_{4N} \\ F_{35} & F_{45} & F_{55} & \cdots & F_{5N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ F_{3N} & F_{4N} & F_{5N} & \cdots & F_{NN} \end{pmatrix}. \tag{27}$$

The eigenvalues of the matrix (27) give approximate values of the first $N - 2$ eigenvalues of the Fredholm operator, and the components of the orthonormal eigenvectors of this matrix give expansion coefficients for the first $N - 2$ eigenfunctions of the Hilgert–Schmidt operator with respect to the chosen orthonormal Legendre polynomials. Recall that the first two terms (i.e., two terms of $y^{\circ}(x, t)$) of the function $y(x, t)$ are known by assumption. Therefore, constructing the next $N - 2$ terms of the expansion, we obtain the N th approximation.

It is important to emphasize the relation between the matrices (11) and (27), which, in general, correspond to two different problems. The matrix (27) can be obtained from the matrix (11) by deleting its first two rows and columns. This allows us to construct the expansion of the original kernel only once, and then use that data for the examination of the new kernel arising in the problem with auxiliary conditions.

Now, we find the functions $\alpha_1(t)$ and $\alpha_2(t)$. To that end, we apply the orthogonal projection operators \mathbf{P}^{*} to equation (12). As a result, we obtain the following formulas:

$$\alpha_1(t) = \frac{1}{\sqrt{2}} \left\{ g_1^{\circ}(t) + \sigma(t) \left[y_1^{\circ}(t) - \int_1^t V_1(t, \tau) y_1^{\circ}(\tau) d\tau \right] + F_{11} y_1^{\circ}(t) + F_{12} y_2^{\circ}(t) + \sum_{k=3}^{\infty} F_{k(1)} y_k^{*}(t) - \int_1^t V_1(t, \tau) \left[F_{11} y_1^{\circ}(\tau) + F_{12} y_2^{\circ}(\tau) + \sum_{k=3}^{\infty} F_{k(1)} y_k^{*}(\tau) \right] d\tau \right\}, \tag{28}$$

$$\alpha_2(t) = \sqrt{\frac{3}{2}} \left\{ g_2^{\circ}(t) + \sigma(t) \left[y_2^{\circ}(t) - \int_1^t V_1(t, \tau) y_2^{\circ}(\tau) d\tau \right] + F_{12} y_1^{\circ}(t) + F_{22} y_2^{\circ}(t) + \sum_{k=3}^{\infty} F_{k(2)} y_k^{*}(t) - \int_1^t V_1(t, \tau) \left[F_{12} y_1^{\circ}(\tau) + F_{22} y_2^{\circ}(\tau) + \sum_{k=3}^{\infty} F_{k(2)} y_k^{*}(\tau) \right] d\tau \right\}. \tag{29}$$

Thus, we have obtained a complete solution of the integral equation (12) with the auxiliary conditions (13).

17.2-3. Equation with a Schmidt Kernel and a Given Right-Hand Side on an Interval.

Consider a mixed integral equation of the form (17.1.3) with a Schmidt kernel. Changing the variables, we can always reduce this equation to the following equation with the parameters $a = -1$, $b = 1$, $\tau_0 = 1$:

$$\begin{aligned} \sigma(t) \left[y(x, t) - \int_1^t V_1(t, \tau) y(x, \tau) d\tau \right] + \int_{-1}^1 S(x, \xi) y(\xi, t) d\xi \\ - \int_1^t V_2(t, \tau) \int_{-1}^1 S(x, \xi) y(\xi, \tau) d\xi d\tau = \frac{f(x, t)}{h(x)}, \tag{30} \\ S(x, \xi) = \frac{F(x, \xi)}{h(x)}, \quad -1 \leq x \leq 1, \quad 1 \leq t \leq T. \end{aligned}$$

Suppose that the right-hand side $f(x, t)$ in (30) is known, and it is required to find the function $y(x, t)$. Here, $f(x, t)$ and $y(x, t)$ are functions of class $C([1, T], L_2[-1, 1])$; $\sigma(t)$ is a given positive continuous function; $h(x) > 0$ is a given function in $L_2[a, b]$; $V_1(t, \tau)$ and $V_2(t, \tau)$ are

Volterra kernels; $S(x, \xi)$ is a Schmidt kernel; $F(x, \xi)$ is a symmetric positive definite Fredholm kernel.

Let us transform the equation with the Schmidt kernel to a Hilbert–Schmidt equation. To that end, we multiply (30) by $\sqrt{h(x)}$ and change the variables as follows:

$$q(x, t) = \sqrt{h(x)}y(x, t), \quad F^h(x, \xi) = \frac{S(x, \xi)\sqrt{h(x)}}{\sqrt{h(\xi)}} = \frac{F(x, \xi)}{\sqrt{h(x)}\sqrt{h(\xi)}}. \tag{31}$$

Then we have

$$\begin{aligned} \sigma(t) \left[q(x, t) - \int_1^t V_1(t, \tau)q(x, \tau) d\tau \right] + \int_{-1}^1 F^h(x, \xi)q(\xi, t) d\xi \\ - \int_{\tau_0}^t V_2(t, \tau) \int_{-1}^1 F^h(x, \xi)q(\xi, \tau) d\xi d\tau = \frac{f(x, t)}{\sqrt{h(x)}}, \quad -1 \leq x \leq 1, \quad 1 \leq t \leq T, \end{aligned} \tag{32}$$

where $q(x, t)$ and $f(x, t)/\sqrt{h(x)}$ are functions of class $C([1, T], L_2[-1, 1])$; $F^h(x, \xi)$ is a symmetric positive definite kernel of Hilbert–Schmidt type (due to the properties of Schmidt kernels); and the other functions have been specified above.

Suppose that the right-hand side of equation (32) is known, and it is required to find the function $q(x, t)$.

Let us seek a solution of the mixed equation (32) in the form or a series

$$q(x, t) = \sum_{k=1}^{\infty} q_k(t)\varphi_k^h(x), \tag{33}$$

where $\varphi_k^h(x)$ are eigenfunctions of the kernel $F^h(x, \xi)$ corresponding to eigenvalues $\mu_k^h > 0$, i.e.,

$$\int_{-1}^1 F^h(x, \xi)\varphi_k^h(\xi) d\xi = \mu_k^h\varphi_k^h(x), \quad k = 1, 2, \dots \tag{34}$$

The representation of a solution in the form (33) is possible, since the system of eigenfunctions of the kernel $F^h(x, \xi)$ forms a basis in $L_2[-1, 1]$.

Here, in contrast to the above case, we construct the basis functions in the form

$$\varphi_k^h(x) = \frac{\Phi_k^h(x)}{\sqrt{h(x)}} \quad k = 1, 2, \dots \tag{35}$$

with an explicit function $h(x)$, where

$$\int_{-1}^1 \varphi_i^h(\xi)\varphi_j^h(\xi) d\xi = \int_{-1}^1 \frac{\Phi_i^h(\xi)\Phi_j^h(\xi)}{h(\xi)} d\xi = \delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases} \tag{36}$$

In order to construct such eigenfunctions, we first construct a certain basis $p_n(x)$ in $L_2[-1, 1]$ for which

$$\int_{-1}^1 p_i^h(\xi)p_j^h(\xi) d\xi = \delta_{ij}, \quad p_n^h(x) = \frac{P_{n-1}^h(x)}{\sqrt{h(x)}}, \quad n = 1, 2, \dots \tag{37}$$

Such a basis can be constructed with the help of the formulas

$$P_0^h(x) = \frac{1}{\sqrt{J_0}}, \quad P_n^h(x) = \frac{1}{\sqrt{\Delta_{n-1}\Delta_n}} \begin{vmatrix} J_0 & J_1 & \cdots & J_n \\ J_1 & J_2 & \cdots & J_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ J_{n-1} & J_n & \cdots & J_{2n-1} \\ 1 & x & \cdots & x^n \end{vmatrix}, \tag{38}$$

$$\Delta_{-1} = 1, \quad \Delta_0 = J_0, \quad \Delta_n = \begin{vmatrix} J_0 & J_1 & \cdots & J_n \\ J_1 & J_2 & \cdots & J_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ J_n & J_{n+1} & \cdots & J_{2n} \end{vmatrix}, \quad J_n = \int_{-1}^1 \frac{\xi^n}{h(\xi)} d\xi.$$

Note that for $h(x) = 1$, the basis functions of the Hilbert space $L_2[-1, 1]$ obtained from (37) and (38) coincide with orthonormal Legendre polynomials.

Let us represent the k th eigenfunction as a series in terms of the basis $p_i^h(x)$ of $L_2[-1, 1]$. Then

$$\varphi_k^h(x) = \sum_{i=1}^{\infty} \varphi_{i(k)}^h p_i^h(x), \quad p_i^h(x) = \frac{P_{i-1}^h(x)}{\sqrt{h(x)}}, \quad \Phi_k^h(x) = \sum_{i=1}^{\infty} \varphi_{i(k)}^h P_{i-1}^h(x). \tag{39}$$

Expanding the Hilbert–Schmidt kernel $F^h(x, \xi)$ as a double series in terms of the chosen basis, we can write

$$F^h(x, \xi) = \sum_{m=1}^{\infty} \sum_{i=1}^{\infty} F_{mn}^h p_m^h(x) p_n^h(\xi), \tag{40}$$

$$F_{mn}^h = \int_{-1}^1 \int_{-1}^1 F^h(x, \xi) p_m^h(x) p_n^h(\xi) dx d\xi, \quad F_{mn}^h = F_{nm}^h.$$

Substituting (39) and (40) into (34), we obtain the following infinite system of linear algebraic equations (with a symmetric matrix) for the determination of the eigenvalues and the eigenfunction expansion coefficients:

$$\sum_{n=1}^{\infty} F_{mn}^h \varphi_{n(k)}^h = \mu_k^h \varphi_{m(k)}^h, \quad m = 1, 2, \dots \tag{41}$$

To calculate approximations of N eigenvalues and eigenfunctions of the Hilbert–Schmidt kernel, one should find the eigenvalues and orthonormal eigenvectors of the matrix

$$[F_{NN}^h] = \begin{pmatrix} F_{11}^h & F_{12}^h & F_{13}^h & \dots & F_{1N}^h \\ F_{12}^h & F_{22}^h & F_{23}^h & \dots & F_{2N}^h \\ F_{13}^h & F_{23}^h & F_{33}^h & \dots & F_{3N}^h \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ F_{1N}^h & F_{2N}^h & F_{3N}^h & \dots & F_{NN}^h \end{pmatrix}. \tag{42}$$

The eigenvalues of the matrix (42) give approximate values of the first N eigenvalues of the Hilbert–Schmidt kernel, and the components of its orthonormal eigenvectors give the coefficients in the expansion of the first N eigenfunctions of that kernel in terms of N orthonormal functions of the basis.

Now, let us expand the right-hand side of equation (32) into the following series:

$$\frac{f(x, t)}{\sqrt{h(x)}} = \sum_{k=1}^{\infty} f_k^h(t) \varphi_k^h(x) = \sum_{k=1}^{\infty} f_k^h(t) \frac{\Phi_k^h(x)}{\sqrt{h(x)}}, \tag{43}$$

$$f_k^h(t) = \int_{-1}^1 \frac{f(x, t)}{\sqrt{h(x)}} \varphi_k^h(x) dx = \int_{-1}^1 \frac{f(x, t)}{h(x)} \Phi_k^h(x) dx.$$

Substituting (33) and (43) into (32) and taking into account (34), we obtain the following sequence of Volterra equations for the unknown functions $q_k(t)$:

$$q_k(t) - \int_1^t V_k^h(t, \tau) q_k(\tau) d\tau = \delta_k^h(t), \quad \delta_k^h(t) = \frac{f_k(t)}{\sigma(t) + \mu_k^h}, \tag{44}$$

$$V_k^h(t, \tau) = \frac{\sigma(\tau) V_1(t, \tau) + \mu_k^h V_2(t, \tau)}{\sigma(t) + \mu_k^h}, \quad k = 1, 2, \dots, \tag{45}$$

where $V_k^h(t, \tau)$ are Volterra kernels belonging to the same class as the kernels $V_1(t, \tau)$ and $V_2(t, \tau)$, since $\mu_k^h \rightarrow 0$ as $k \rightarrow \infty$.

A solution of the infinite system of Volterra equations (44) can be constructed by analytical and numerical methods of Chapter 11. This solution can be written in the form

$$q_k(t) = \delta_k^h(t) + \int_1^t R_k^h(t, \tau) \delta_k^h(\tau) d\tau, \tag{46}$$

where $R_k^h(t, \tau)$ is the resolvent of the kernel $V_k^h(t, \tau)$.

The series (33) converges in $L_2[-1, 1]$ uniformly in $t \in [1, T]$, and its sum is a continuous function of $t \in [1, T]$ with values in $L_2[-1, 1]$.

Inserting (45) into (33) and taking into account (43), one can also represent the solution in the form

$$q(x, t) = \frac{f(x)}{\sqrt{h(x)}} + \sum_{k=1}^{\infty} \int_1^t R_k^h(t, \tau) f_k^h(\tau) d\tau \varphi_k^h(x).$$

Finally, in view of the transformation of the variable (31) and the formula for eigenfunctions (35), we have

$$y(x, t) = \frac{1}{h(x)} \sum_{k=1}^{\infty} q_k(t) \Phi_k^h(x). \tag{47}$$

Note that the solution (47) involves the function $h(x)$ in an explicit manner, which allows us to solve equation (30) with great accuracy by preserving a small number of terms. In the case of a strongly oscillating function $h(x)$, the other known methods can hardly be used for the construction of solutions.

17.2-4. Equation with a Schmidt Kernel and Auxiliary Conditions.

Consider equation (30) with the right-hand side of the form $f(x, t) = \alpha_1(t) + \alpha_2(t)x - g(x, t)$ and two auxiliary integral conditions of the form (17.1.4) on the unknown function $y(x, t)$. The problem is to find a solution of the mixed integral equation

$$\begin{aligned} \sigma(t) \left[y(x, t) - \int_1^t V_1(t, \tau) y(x, \tau) d\tau \right] + \int_{-1}^1 S(x, \xi) y(\xi, t) d\xi \\ - \int_1^t V_2(t, \tau) \int_{-1}^1 S(x, \xi) y(\xi, \tau) d\xi d\tau = \frac{\alpha_1(t)}{h(x)} + \frac{\alpha_2(t)x}{h(x)} - \frac{g(x, t)}{h(x)}, \tag{48} \\ S(x, \xi) = \frac{F(x, \xi)}{h(x)}, \quad -1 \leq x \leq 1, \quad 1 \leq t \leq T. \end{aligned}$$

with the auxiliary conditions

$$\int_{-1}^1 y(\xi, t) d\xi = M_1(t), \quad \int_{-1}^1 \xi y(\xi, t) d\xi = M_2(t), \tag{49}$$

the unknown functions being $y(x, t)$, $\alpha_1(t)$, and $\alpha_2(t)$. The other functions in (48) are assumed given, and $g(x, t)$ is a continuous function of $t \in [1, T]$ with values in $L_2[-1, 1]$.

Let us transform the equation with the Schmidt kernel to an equation with a Hilbert–Schmidt kernel by changing the variables as in (31). As a result, equation (48) and the auxiliary conditions (49) become

$$\begin{aligned} \sigma(t) \left[q(x, t) - \int_1^t V_1(t, \tau) q(x, \tau) d\tau \right] + \int_{-1}^1 F^h(x, \xi) q(\xi, t) d\xi \\ - \int_{\tau_0}^t V_2(t, \tau) \int_{-1}^1 F^h(x, \xi) q(\xi, \tau) d\xi d\tau = \frac{\alpha_1(t)}{\sqrt{h(x)}} + \frac{\alpha_2(t)x}{\sqrt{h(x)}} - \frac{g(x, t)}{\sqrt{h(x)}}, \tag{50} \end{aligned}$$

$$\int_{-1}^1 \frac{q(\xi, t)}{\sqrt{h(\xi)}} d\xi = M_1(t), \quad \int_{-1}^1 \frac{q(\xi, t)}{\sqrt{h(\xi)}} \xi d\xi = M_2(t), \quad -1 \leq x \leq 1, \quad 1 \leq t \leq T. \tag{51}$$

In order to construct a solution of the mixed integral equation (48) with the auxiliary conditions (49), we use the basis $p_n^h(x)$ in $L_2[-1, 1]$ (see (37) and (38)) and note that the space $L_2[-1, 1]$ can be represented as a direct sum of its orthogonal subspaces, $L_2[-1, 1] = L_2^{h^\circ}[-1, 1] \oplus L_2^{h^*}[-1, 1]$ (see Supplement 12.5-3), where $L_2^{h^\circ}[-1, 1]$ is the Euclidean space with the basis $p_1^h(x)$ and $p_2^h(x)$, and $L_2^{h^*}[-1, 1]$ is the Hilbert space with the basis $p_k^h(x)$ ($k = 3, 4, \dots$). It can be seen that the integrand and the right-hand side can be represented as a sum of continuous functions of $t \in [1, T]$ with values in $L_2^{h^\circ}[-1, 1]$ and $L_2^{h^*}[-1, 1]$, respectively, i.e.,

$$q(x, t) = q^\circ(x, t) + q^*(x, t), \quad \frac{f(x, t)}{\sqrt{h(x)}} = f_h^\circ(x, t) + f_h^*(x, t), \tag{52}$$

and the following representations hold:

$$\begin{aligned} q^\circ(x, t) &= q_1^\circ(t)p_1^h(x) + q_2^\circ(t)p_2^h(x), & q_1^\circ(t) &= \frac{M_1(t)}{\sqrt{J_0}}, & q_2^\circ(t) &= \frac{J_0M_2(t) - J_1M_1(t)}{\sqrt{J_0(J_0J_2 - J_1^2)}}, \\ \frac{f(x, t)}{\sqrt{h(x)}} &= \frac{\alpha_1(t)}{\sqrt{h(x)}} + \frac{\alpha_2(t)x}{\sqrt{h(x)}} - \frac{g(x, t)}{\sqrt{h(x)}}, & \frac{g(x, t)}{\sqrt{h(x)}} &= g_h^\circ(x, t) + g_h^*(x, t), \\ f_h^\circ(x, t) &= \left[\sqrt{J_0}\alpha_1(t) + \frac{J_1}{\sqrt{J_0}}\alpha_2(t) - g_1(t) \right] p_1^h(x) + \left[\frac{\sqrt{J_0J_2 - J_1^2}}{\sqrt{J_0}}\alpha(t) - g_1 \right] p_2^h(x), \\ f_h^*(x, t) &= -g_h^*(x), & g_h^\circ(x, t) &= g_1^{h^\circ}(t)p_1^h(x) + g_2^{h^\circ}(t)p_2^h(x), \\ g_1^\circ(t) &= \int_{-1}^1 \frac{g(x, t)}{\sqrt{h(x)}} p_1^h(x) dx, & g_2^\circ(t) &= \int_{-1}^1 \frac{g(x, t)}{\sqrt{h(x)}} p_2^h(x) dx. \end{aligned} \tag{53}$$

Note that in the representation (52) for $q(x, t)$, the function $q^\circ(x, t)$ is known as determined by the auxiliary conditions, and the term $q^*(x, t)$ should be found. Conversely, for the right-hand side, $f_h^\circ(x, t)$ should be found and $f_h^*(x, t)$ is determined by the function $g(x, t)/\sqrt{h(x)}$. The facts mentioned above allow us to classify the resulting problem as a special case of the general projection problem whose solution is given in Subsection 17.4-3.

According to the general method, in the present case one can introduce an operator of orthogonal projection that maps the space $L_2[-1, 1]$ onto $L_2^{h^\circ}[-1, 1]$:

$$\mathbf{P}_h^\circ \phi(x, t) = \int_{-1}^1 \phi(\xi, t) [p_1^h(x)p_1^h(\xi) + p_2^h(x)p_2^h(\xi)] d\xi. \tag{54}$$

Obviously, the projector $\mathbf{P}_h^* = \mathbf{I} - \mathbf{P}_h^\circ$ maps $L_2[-1, 1]$ onto $L_2^{h^*}[-1, 1]$. Moreover, the following relations hold:

$$\begin{aligned} \mathbf{P}_h^\circ q(x, t) &= q^\circ(x, t), & \mathbf{P}_h^* q(x, t) &= q^*(x, t), \\ \mathbf{P}_h^\circ \frac{f(x, t)}{\sqrt{h(x)}} &= f_h^\circ(x, t), & \mathbf{P}_h^* \frac{f(x, t)}{\sqrt{h(x)}} &= f_h^*(x, t). \end{aligned} \tag{55}$$

According to Subsections 17.4-2 and 17.4-3, let us apply the projection operator \mathbf{P}_h^* to equation (50). Then, for the determination of $q^*(x, t)$, we obtain the following integral equation in $L_2^{h^*}[-1, 1]$ with a known right-hand side:

$$\begin{aligned} \sigma(t) &\left[q^*(x, t) - \int_1^t V_1(t, \tau) q^*(x, \tau) d\tau \right] \\ &+ \int_{-1}^1 F_h^*(x, \xi) q^*(\xi, t) d\xi - \int_{\tau_0}^t V_2(t, \tau) \int_{-1}^1 F_h^*(x, \xi) q^*(\xi, \tau) d\xi d\tau \\ &= -g_h^*(x, t) - \int_{-1}^1 F_h^*(x, \xi) q^\circ(\xi, t) d\xi + \int_1^t V_2(t, \tau) \int_{-1}^1 F_h^*(x, \xi) q^\circ(\xi, \tau) d\xi d\tau, \\ &-1 \leq x \leq 1, \quad 1 \leq t \leq T, \end{aligned} \tag{56}$$

where the kernel of the integral equation

$$F_h^*(x, \xi) = F^h(x, \xi) - \int_{-1}^1 F^h(s, \xi)[p_1^h(x)p_1^h(s) + p_2^h(x)p_2^h(s)] ds \tag{57}$$

is a Hilbert–Schmidt kernel. A solution of equation (56) can be constructed in the form of a series in terms of eigenfunctions of the kernel (57). These form a basis in the Hilbert space $L_2^{h*}[-1, 1]$. Let us construct a system of these eigenfunctions.

For an eigenfunction $\varphi_k^{h*}(x)$, let μ_k^{h*} be the corresponding eigenvalue of the kernel $F_p^*(x, \xi)$. Then

$$\int_{-1}^1 F_h^*(x, \xi)\varphi_k^{h*}(\xi) d\xi = \mu_k^{h*}\varphi_k^{h*}(x), \quad k = 3, 4, \dots \tag{58}$$

Let us represent the eigenfunction $\varphi_i^{h*}(x)$ as a series in terms of the basis functions $p_i^h(x)$ ($i \geq 3$):

$$\varphi_k^{h*}(x) = \sum_{i=3}^{\infty} \varphi_{i(k)}^{h*} p_i^h(x), \quad k = 3, 4, \dots \tag{59}$$

The double series expansion of the kernel $F_h^*(x, \xi)$ is obtained with the help of (40) and (57):

$$F_h^*(x, \xi) = \sum_{m=3}^{\infty} \sum_{n=3}^{\infty} F_{mn}^h p_m^h(x)p_n^h(\xi) + \sum_{n=3}^{\infty} F_{1n}^h p_n^h(x)p_1^h(\xi) + \sum_{n=3}^{\infty} F_{2n}^h p_n^h(x)p_2^h(\xi). \tag{60}$$

Note that the coefficients of the expansion of the kernel $F_h^*(x, \xi)$ in (60) coincide with coefficients of the expansion of the kernel $F^h(x, \xi)$, which allows us to avoid recalculation of the coefficients of the new problem and use the existing data.

Substituting (59) and (60) into (58), we obtain the following infinite system of linear algebraic equations (with a symmetric matrix) for the eigenvalues and the eigenfunction expansion coefficients:

$$\sum_{n=3}^{\infty} F_{mn}^h \varphi_n^{h*} = \mu_k^{h*} \varphi_m^{h*}, \quad m = 3, 4, \dots \tag{61}$$

Now, let us construct a solution of equation (56). To that end, we represent the functions $q^*(x, t)$ and $g_h^*(x, t)$ as series in terms of eigenfunctions of the kernel $F_h^*(x, \xi)$:

$$q^*(x, t) = \sum_{k=3}^{\infty} q_k^*(t)\varphi_k^{h*}(x), \quad g_h^*(x, t) = \sum_{k=3}^{\infty} g_k^{h*}(t)\varphi_k^{h*}(x), \quad g_k^{h*}(t) = \int_{-1}^1 g_k^*(x, t)\varphi_k^{h*}(x) dx. \tag{62}$$

Substituting these into (56) and taking into account (53), (57)–(60), we obtain the following sequence of independent Volterra equations of the second kind:

$$\begin{aligned} q_k^*(t) - \int_1^t V_k^{h*}(t, \tau)q_k^*(\tau) d\tau &= \delta_k^{h*}(t), \quad V_k^{h*}(t, \tau) = \frac{\sigma(t)V_1(t, \tau) + \mu_k^{h*}V_2(t, \tau)}{\sigma(t) + \mu_k^{h*}}, \\ \delta_k^{h*}(t) &= -\frac{1}{\sigma(t) + \mu_k^{h*}} \left[g_k^{h*}(t) + \sum_{i=1}^2 F_{k(i)}^h q_i^{\circ}(t) - \int_1^t V_2(t, \tau) \sum_{i=1}^2 F_{k(i)}^h q_i^{\circ}(\tau) d\tau \right], \\ F_{k(i)}^h &= \sum_{n=3}^{\infty} F_{in}^h \varphi_n^{h*}, \quad i = 1, 2, \quad k = 3, 4, \dots \end{aligned} \tag{63}$$

Resolving (63) with respect to $q_k^*(t)$ by the methods of Chapter 11, we obtain

$$q_k^*(t) = f_k^{h*}(t) + \int_1^t R_k^{h*}(t, \tau) f_k^{h*}(\tau) d\tau, \tag{64}$$

where $R_k^{h*}(t, \tau)$ is the resolvent of the kernel $V_k^{h*}(t, \tau)$.

Now, in view of (62)–(64), the function $q^*(x, t)$ has been determined, and therefore, we easily find $q(x, t)$, since $q^\circ(x, t)$ is known by assumption (see (52) and (53)).

In practical calculations, the number of expansion terms is naturally limited. For instance, taking the basis functions $p_k^h(x)$ for $k = 3, \dots, N$, we obtain the N th approximation of the desired solution. In this case, for the construction of eigenvalues and eigenfunctions of the Hilbert–Schmidt kernel $F_h^*(x, \xi)$, one should find the eigenvalues and orthonormal eigenvectors of the matrix

$$[F_{NN}^h] = \begin{pmatrix} F_{33}^h & F_{34}^h & F_{35}^h & \dots & F_{3N}^h \\ F_{34}^h & F_{44}^h & F_{45}^h & \dots & F_{4N}^h \\ F_{35}^h & F_{45}^h & F_{55}^h & \dots & F_{5N}^h \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ F_{3N}^h & F_{4N}^h & F_{5N}^h & \dots & F_{NN}^h \end{pmatrix}. \tag{65}$$

The eigenvalues of the matrix (65) give approximations for the first $N - 2$ eigenvalues if the Hilbert–Schmidt operator and the components of the orthonormal eigenvectors of the matrix give expansion coefficients of the first $N - 2$ eigenfunctions of this operator in terms of the chosen orthonormal Legendre polynomials. Recall that the first two terms of the expansion (i.e., two terms of $q^\circ(x, t)$) of the function $q(x, t)$ are known by assumption. Therefore, constructing the next $N - 2$ terms of the expansion, we obtain the N th approximation of the solution.

It is important to keep in mind the relation between the matrices (42) and (65). The matrix (65) can be obtained from (42) by deleting its first two rows and columns. This allows us to construct the expansion of the original series only once, and then use this information for studying the new kernel in the problem with auxiliary conditions.

Now, let us find the functions $\alpha_1(t)$, and $\alpha_2(t)$. To this end, we apply the projection operator \mathbf{P}_h° to equation (56). As a result, we obtain the following formulas:

$$\begin{aligned} \alpha_1(t) = & \frac{1}{\sqrt{J_0}} \left\{ g_1^{h^\circ}(t) - \frac{J_1}{\sqrt{J_0}} \alpha_2(t) + \sigma(t) \left[q_1^\circ(t) - \int_1^t V_1(t, \tau) q_1^\circ(\tau) d\tau \right] + F_{11}^h q_1^\circ(t) \right. \\ & \left. + F_{12}^h q_2^\circ(t) + \sum_{k=3}^\infty F_{k(1)}^h q_k^*(t) - \int_1^t V_1(t, \tau) \left[F_{11}^h q_1^\circ(\tau) + F_{12}^h q_2^\circ(\tau) + \sum_{k=3}^\infty F_{k(1)}^h q_k^*(\tau) \right] d\tau \right\}, \tag{66} \end{aligned}$$

$$\begin{aligned} \alpha_2(t) = & \sqrt{\frac{J_0}{J_0 J_2 - J_1^2}} \left\{ g_2^{h^\circ}(t) + \sigma(t) \left[q_2^\circ(t) - \int_1^t V_1(t, \tau) q_2^\circ(\tau) d\tau \right] + F_{12}^h q_1^\circ(t) \right. \\ & \left. + F_{22}^h q_2^\circ(t) + \sum_{k=3}^\infty F_{k(2)}^h q_k^*(t) - \int_1^t V_1(t, \tau) \left[F_{12}^h q_1^\circ(\tau) + F_{22}^h q_2^\circ(\tau) + \sum_{k=3}^\infty F_{k(2)}^h q_k^*(\tau) \right] d\tau \right\}. \tag{67} \end{aligned}$$

Note that relations (66) and (67) form a system of two linear algebraic equations (with a triangular matrix) for the determination of the unknown quantities $\alpha_1(t)$ and $\alpha_2(t)$.

Thus, we have constructed a solution of the integral equation (48) with the auxiliary conditions (49).

© References for Section 17.2: E. Goursat (1923), G. Szegö (1975), V. M. Aleksandrov and S. M. Mkhitarian (1983), V. M. Aleksandrov and A. V. Manzhirou (1987), A. V. Manzhirou (2001, 2005), A. V. Manzhirou and K. E. Kazakov (2006).

17.3. Methods of Solving Mixed Integral Equations on a Ring-Shaped Domain

17.3-1. Equation with a Hilbert–Schmidt Kernel and a Given Right-Hand Side.

Consider a mixed integral equation with a Hilbert–Schmidt kernel on a ring-shape domain (see Subsection 17.1-3). By a suitable transformation of the variables, this equation can be reduced to a similar equation with the parameters $a = 0$, $b = 1$, $\tau_0 = 1$: approximation of the solution

$$\sigma(t) \left[y(r, t) - \int_1^t V_1(t, \tau) y(r, \tau) d\tau \right] + \int_0^1 G(r, \rho) y(\rho, t) \rho d\rho - \int_1^t V_2(t, \tau) \int_0^1 G(r, \rho) y(\rho, \tau) \rho d\rho d\tau = f(r, t), \quad 0 \leq r \leq 1, \quad 1 \leq t \leq T. \quad (1)$$

Suppose that the right-hand side $f(r, t)$ of equation (1) is known and it is required to find the function $y(r, t)$. Here, $f(r, t)$, $y(r, t)$ are supposed to be continuous functions of $t \in [1, T]$ with values in $\widehat{L}_2(\omega)$.

Let us seek a solution of equation (1) in the form of a series

$$y(x, t) = \sum_{k=1}^{\infty} y_k(t) \psi_k(r), \quad (2)$$

where $\psi_k(r)$ are eigenfunctions of the kernel $F(r, \rho)$ corresponding to eigenvalues $\nu_k > 0$, i.e.,

$$\int_0^1 G(r, \rho) \psi_k(\rho) \rho d\rho = \nu_k \psi_k(r), \quad k = 1, 2, \dots \quad (3)$$

The representation (2) is justified by the fact that the system of eigenfunctions of the kernel $F(r, \rho)$ forms a complete orthonormal system in $\widehat{L}_2(\omega)$, in other words, an orthonormal basis in $\widehat{L}_2(\omega)$ (see Subsection 13.6-1 and Supplement 12.5-3). This fact also allows us to represent the right-hand side of the equation in the form

$$f(r, t) = \sum_{k=1}^{\infty} f_k(t) \psi_k(r), \quad f_k(t) = \int_0^1 f(\rho, t) \psi_k(\rho) \rho d\rho. \quad (4)$$

Substituting (2) into (1) and taking into account (3) and (4), we obtain the following sequence of Volterra equations of the second kind for the unknown functions $y_k(t)$:

$$y_k(t) - \int_1^t V_{k(\nu)}(t, \tau) y_k(\tau) d\tau = \gamma_k(t), \quad \gamma_k(t) = \frac{f_k(t)}{\sigma(t) + \nu_k}, \quad (5)$$

$$V_{k(\nu)}(t, \tau) = \frac{\sigma(t) V_1(t, \tau) + \nu_k V_2(t, \tau)}{\sigma(t) + \nu_k}, \quad k = 1, 2, \dots, \quad (6)$$

where $V_k^\nu(t, \tau)$ are Volterra kernels of the same class as $V_1(t, \tau)$ and $V_2(t, \tau)$, since $\nu_k \rightarrow 0$ as $k \rightarrow \infty$.

A solution of the sequence of Volterra equations (5) can be constructed by analytical and numerical methods of Chapter 11. This solution can be written in the form

$$y_k(t) = \gamma_k(t) + \int_1^t R_{k(\nu)}(t, \tau) \gamma_k(\tau) d\tau, \quad (6)$$

where $R_k^\nu(t, \tau)$ is the resolvent of the kernel $V_k(t, \tau)$.

Thus, we have found a solution of equation (1). The series (2) converges in $\widehat{L}_2(\omega)$ uniformly in $t \in [1, T]$, and its sum is a continuous function of t with values in $\widehat{L}_2(\omega)$.

Now, let us construct eigenfunctions and eigenvalues of the Hilbert–Schmidt kernel. We represent the k th eigenfunction as a series in terms of any orthonormal basis $z_i(r)$ in $\widehat{L}_2(\omega)$. For definiteness, we take

$$z_k(r) = \sqrt{4k + 2} P_{k-1}(1 - 2r^2), \quad k = 1, 2, \dots, \tag{7}$$

where $P_n(x)$ are the Legendre polynomials. Now we can represent the eigenfunctions of the kernel $G(r, \rho)$ in the form of a series

$$\psi_k(r) = \sum_{i=1}^{\infty} \psi_{i(k)} z_i(r), \quad k = 1, 2, \dots \tag{8}$$

We write a double series expansion (in terms of the chosen basis) for the kernel of the equation:

$$G(r, \rho) = \sum_{m=1}^{\infty} \sum_{i=1}^{\infty} G_{mn} z_m(r) z_n(\rho), \tag{9}$$

$$G_{mn} = \int_0^1 \int_0^1 G(r, \rho) z_m(r) z_n(\rho) r \rho \, dr \, d\rho, \quad G_{mn} = G_{nm}.$$

Substituting (8) and (9) into (3), we obtain an infinite system of linear algebraic equations (with a symmetric matrix) for the determination of the eigenfunctions. This system has the form

$$\sum_{n=1}^{\infty} G_{mn} \psi_{n(k)} = \nu_k \psi_{m(k)}, \quad m = 1, 2, \dots \tag{10}$$

Naturally, in practical calculations one has to limit the number of expansion terms. For instance, taking N orthonormal Legendre polynomials, we obtain the N th approximation of the solution. And in order to construct the eigenvalues and eigenfunctions of the Hilbert–Schmidt kernel, in this case, one should find the eigenvalues and orthonormal eigenvectors of the matrix

$$[G_{NN}] = \begin{pmatrix} G_{11} & G_{12} & G_{13} & \cdots & G_{1N} \\ G_{12} & G_{22} & G_{23} & \cdots & G_{2N} \\ G_{13} & G_{23} & G_{33} & \cdots & G_{3N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ G_{1N} & G_{2N} & G_{3N} & \cdots & G_{NN} \end{pmatrix}. \tag{11}$$

Eigenvalues of the matrix (11) give approximations for the first N eigenvalues of the Hilbert–Schmidt kernel, and the components of its orthonormal eigenvectors give the coefficients in the expansion of the first N eigenfunctions of the Hilbert–Schmidt kernel in terms of N orthonormal Legendre polynomials.

17.3-2. Equation with a Hilbert–Schmidt Kernel and Auxiliary Conditions.

Consider equation (1) with the right-hand side of the form $f(r, t) = \beta(t) - w(r, t)$ and an auxiliary condition of the form (8) of Subsection 17.1-3 for the unknown function $y(r, t)$. The problem is to find a solution of the mixed integral equation

$$\sigma(t) \left[y(r, t) - \int_1^t V_1(t, \tau) y(r, \tau) \, d\tau \right] + \int_0^1 G(r, \rho) y(\rho, t) \rho \, d\rho - \int_1^t V_2(t, \tau) \int_0^1 G(r, \rho) y(\rho, \tau) \rho \, d\rho \, d\tau = \beta(t) - w(r, t), \tag{12}$$

$$0 \leq r \leq 1, \quad 1 \leq t \leq T$$

with the auxiliary condition

$$\int_0^1 y(\rho, t)\rho d\rho = M(t), \tag{13}$$

the unknown functions being $y(r, t)$ and $\beta(t)$. All other functions in (12) are assumed known, and $w(r, t)$ is a continuous function of $t \in [1, T]$ with values in $\widehat{L}_2(\omega)$.

Note that the Hilbert space $\widehat{L}_2(\omega)$ can be represented as the direct sum of its orthogonal subspaces: $\widehat{L}_2(\omega) = L_2^\circ(\omega) \oplus L_2^*(\omega)$ (see Supplement 12.5-3), where $\widehat{L}_2^\circ(\omega)$ is the Euclidean space with the basis $z_1(r) = \sqrt{2}P_0(r) = \sqrt{2}$ and $\widehat{L}_2^*(\omega)$ is the Hilbert space with the basis $z_k(r)$ ($k = 2, 3, \dots$). Note that the integrand and the right-hand side can be represented as a sum of functions that are continuous in $t \in [1, T]$ and take values in $\widehat{L}_2^\circ(\omega)$ and $\widehat{L}_2^*(\omega)$, respectively. Thus, we can write

$$y(r, t) = y^\circ(r, t) + y^*(r, t), \quad f(r, t) = f^\circ(r, t) + f^*(r, t), \tag{14}$$

and the following expansions hold:

$$\begin{aligned} y^\circ(r, t) &= y_1^\circ(t)z_1(r), & y_1^\circ(t) &= \sqrt{2}M(t), \\ f^\circ(r, t) &= \left[\frac{\beta(t)}{\sqrt{2}} - w_1^\circ(t) \right] z_1(r), & f^*(r, t) &= -w^*(r, t), \\ w(x, t) &= w^\circ(x, t) + w^*(x, t), & w^\circ(x, t) &= w_1^\circ(t)z_1(r), & w_1^\circ(t) &= \sqrt{2} \int_0^1 w(\rho, t)\rho d\rho. \end{aligned} \tag{15}$$

Note that in the representation (14) for $y(r, t)$, the term $y^\circ(r, t)$ is known (as defined by the auxiliary conditions), and the term $y^*(r, t)$ should be found. Conversely, for the right-hand side, one should find $f^\circ(r, t)$ and the term $f^*(r, t)$ is determined by the function $w(r, t)$. These considerations allow us to classify the problem as a special case of the general projection problem examined in Subsection 17.4-3.

According to the general method, in the present case we can introduce an operator of orthogonal projection that maps the space $\widehat{L}_2(\omega)$ onto $\widehat{L}_2^\circ(\omega)$:

$$\mathbf{Q}^\circ f(r, t) = \int_0^1 f(\rho, t)z_1(r)z_1(\rho)\rho d\rho = 2 \int_0^1 f(\rho, t)\rho d\rho. \tag{16}$$

Obviously, the orthogonal projector $\mathbf{Q}^* = \mathbf{I} - \mathbf{Q}^\circ$ maps the space $\widehat{L}_2(\omega)$ onto $\widehat{L}_2^*(\omega)$. Moreover, the following relations hold:

$$\begin{aligned} \mathbf{Q}^\circ y(r, t) &= y^\circ(r, t), & \mathbf{Q}^* y(r, t) &= y^*(r, t), \\ \mathbf{Q}^\circ f(r, t) &= f^\circ(r, t), & \mathbf{Q}^* f(r, t) &= f^*(r, t). \end{aligned} \tag{17}$$

Following Section 17.4, we apply the projection operator \mathbf{Q}^* to equation (12). As a result, we obtain an integral equation for $y^*(r, t)$ in the space $\widehat{L}_2^*(\omega)$ with a known right-hand side:

$$\begin{aligned} \sigma(t) &\left[y^*(r, t) - \int_1^t V_1(t, \tau)y^*(r, \tau) d\tau \right] \\ &+ \int_0^1 G^*(r, \rho)y^*(\rho, t)\rho d\rho - \int_1^t V_2(t, \tau) \int_0^1 G^*(r, \rho)y^*(\rho, \tau)\rho d\rho d\tau \\ &= -w^*(r, t) - \int_0^1 G^*(r, \rho)y^\circ(\rho, t)\rho d\rho + \int_1^t V_2(t, \tau) \int_0^1 G^*(r, \rho)y^\circ(\rho, \tau)\rho d\rho d\tau, \\ &0 \leq r \leq 1, \quad 1 \leq t \leq T, \end{aligned} \tag{18}$$

where the kernel of the integral equation

$$G^*(r, \rho) = G(r, \rho) - 2 \int_0^1 G(r, \rho)r \, dr \tag{19}$$

is of Hilbert–Schmidt type.

Let us construct a solution of equation (18) in the form of a series in terms of eigenfunctions of the kernel (19). These eigenfunctions form a basis in the Hilbert space $\widehat{L}_2^*(\omega)$. Let us construct the system of these eigenfunctions.

For an eigenfunction $\psi_k^*(r)$ of the kernel $G^*(r, \rho)$, let ν_k^* be the corresponding eigenvalue. Then,

$$\int_0^1 G^*(r, \rho)\psi_k^*(\rho)\rho \, d\rho = \nu_k^*\psi_k^*(r), \quad k = 2, 3, \dots \tag{20}$$

Let us represent the eigenfunction $\psi_k^*(r)$ as a series with respect to the basis $z_i(r)$ ($i \geq 2$):

$$\psi_k^*(r) = \sum_{i=2}^{\infty} \psi_{i(k)}^* z_i(r), \quad k = 2, 3, \dots \tag{21}$$

For the kernel $G^*(r, \rho)$, we construct a double series expansion with the help of (9) and (19):

$$G^*(r, \rho) = \sum_{m=2}^{\infty} \sum_{n=2}^{\infty} G_{mn} z_m(r) z_n(\rho) + \sum_{n=2}^{\infty} \sqrt{2} G_{1n} z_n(r). \tag{22}$$

Note that the coefficients in the expansion of the kernel $G^*(r, \rho)$ in (22) coincide with those in the expansion of $G(r, \rho)$, and this allows us to avoid recalculating the coefficients of the new problem and use the available information.

Substituting (21) and (22) into (20), we obtain an infinite system of linear algebraic equations for the eigenvalues and the eigenfunction expansion coefficients. This system has a symmetric matrix and can be written as follows:

$$\sum_{n=2}^{\infty} G_{mn} \psi_{n(k)}^* = \nu_k^* \psi_{m(k)}^*, \quad m = 2, 3, \dots \tag{23}$$

Now let us construct a solution of equation (18). For this purpose, we represent the functions $y^*(r, t)$ and $w^*(r, t)$ in the form of series in terms of eigenfunctions of the kernel $G^*(r, \rho)$:

$$y^*(x, t) = \sum_{k=2}^{\infty} y_k^*(t) \psi_k^*(r), \quad w^*(r, t) = \sum_{k=2}^{\infty} w_k^*(t) \psi_k^*(r), \quad w_k^*(t) = \int_0^1 w^*(r, t) \psi_k^*(\rho) \rho \, d\rho. \tag{24}$$

Substituting these into (18) and taking into account (15), (19)–(22), we obtain the following sequence of independent Volterra equations:

$$\begin{aligned} y_k^*(t) - \int_1^t V_{k(\nu)}^*(t, \tau) y_k^*(\tau) \, d\tau &= \gamma_k^*(t), \quad V_{k(\nu)}^*(t, \tau) = \frac{\sigma(t)V_1(t, \tau) + \nu_k^* V_2(t, \tau)}{\sigma(t) + \nu_k^*}, \\ \gamma_k^*(t) &= -\frac{1}{\sigma(t) + \nu_k^*} \left[w_k^*(t) + G_k y_1^\circ(t) - \int_1^t V_2(t, \tau) G_k y_1^\circ(\tau) \, d\tau \right], \\ G_k &= \sum_{n=1}^{\infty} G_{1n} \psi_{n(k)}^*, \quad k = 2, 3, \dots \end{aligned} \tag{25}$$

Resolving (25) with respect to $y_k^*(t)$ by the methods of Chapter 11, we get

$$y_k^*(t) = \gamma_k^*(t) + \int_1^t R_{k(\nu)}^*(t, \tau) \gamma_k^*(\tau) d\tau, \tag{26}$$

where $R_{k(\nu)}^*(t, \tau)$ is the resolvent of the kernel $V_{k(\nu)}^*(t, \tau)$.

Thus, in view of (24)–(26), the function $y^*(r, t)$ has been determined, and we easily find $y(r, t)$, since $y^\circ(r, t)$ is known by assumption (see (14) and (15)).

Before we go on to find the other unknown quantities of the problem, we make some practical recommendations. Naturally, in practical calculations the number of expansion terms should be limited. For instance, taking the Legendre polynomials from the second to the N th, we obtain the N th approximation of the desired solution. In this case, for the construction of eigenvalues and eigenfunctions of the Hilbert–Schmidt kernel $G^*(r, \rho)$, one should find the eigenvalues and the orthonormal eigenvectors of the matrix

$$[G_{NN}^*] = \begin{pmatrix} G_{22} & G_{23} & G_{24} & \cdots & G_{2N} \\ G_{23} & G_{33} & G_{34} & \cdots & G_{3N} \\ G_{24} & G_{34} & G_{44} & \cdots & G_{4N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ G_{2N} & G_{3N} & G_{4N} & \cdots & G_{NN} \end{pmatrix}. \tag{27}$$

The eigenvalues of the matrix (27) give approximations of the first N eigenvalues of the Hilbert–Schmidt kernel, and the components of its eigenvectors give the coefficients in the expansion of the first $N - 1$ eigenfunctions of that kernel in terms of the chosen Legendre polynomials. Recall that the first term $y^\circ(x, t)$ of the expansion of $y(x, t)$ is known by assumption. Therefore, constructing the next $N - 1$ terms of the expansion, we obtain the N th approximation.

Note that the matrix (27) can be obtained from the matrix (11) by deleting its first two rows and columns. This allows us to construct the expansion of the original kernel only once and then use that data for the examination of the new kernel arising in the problem with auxiliary conditions.

Now, let us find the function $\beta(t)$. To that end, we apply the orthogonal projection operator \mathbf{Q}^* to equation (12). As a result we obtain the following formula:

$$\begin{aligned} \beta(t) = \sqrt{2} \left\{ w_1^\circ(t) + \sigma(t) \left[y_1^\circ(t) - \int_1^t V_1(t, \tau) y_1^\circ(\tau) d\tau \right] + G_{11} y_1^\circ(t) \right. \\ \left. + \sum_{k=2}^\infty G_{k(1)} y_k^*(t) - \int_1^t V_1(t, \tau) \left[G_{11} y_1^\circ(\tau) + \sum_{k=2}^\infty G_k y_k^*(\tau) \right] d\tau \right\}. \end{aligned} \tag{28}$$

Thus, we have obtained a complete solution of the integral equation (12) with the auxiliary conditions (13).

17.3-3. Equation with a Schmidt Kernel and a Given Right-Hand Side.

Consider a mixed integral equation of the form (6) from Subsection 17.1-3 with a Schmidt kernel. Changing the variables, we can easily transform this equation to a similar equation with the parameters $a = 0, b = 1, \tau_0 = 1$:

$$\begin{aligned} \sigma(t) \left[y(r, t) - \int_1^t V_1(t, \tau) y(\rho, \tau) d\tau \right] + \int_0^1 S_\omega(r, \rho) y(\rho, t) \rho d\rho \\ - \int_1^t V_2(t, \tau) \int_0^1 S_\omega(r, \rho) y(\rho, \tau) \rho d\rho d\tau = \frac{f(r, t)}{h(r)}, \end{aligned} \tag{29}$$

$$S_\omega(r, \rho) = \frac{G(r, \rho)}{h(r)}, \quad 0 \leq r \leq 1, \quad 1 \leq t \leq T.$$

Suppose that the right-hand side $f(r, t)/h(r)$ of (29) is known, and it is required to find the function $y(r, t)$. Here, $f(r, t)$ and $y(r, t)$ are continuous functions of $t \in [1, T]$ with values in $\widehat{L}_2(\omega)$; $\sigma(t)$ is a given positive continuous function, $h(r) > 0$ is a given function in $\widehat{L}_2(\omega)$; $V_1(t, \tau)$ and $V_2(t, \tau)$ are Volterra kernels; $S_\omega(r, \rho)$ is a Schmidt kernel; $G(r, \rho)$ is a symmetric positive definite Fredholm kernel.

Let us transform the equation with the Schmidt kernel to an equation with Hilbert–Schmidt kernel. To this end, we multiply equation (28) by $\sqrt{h(r)}$ and change the variables as follows:

$$q(r, t) = \sqrt{h(r)}y(r, t), \quad G^h(r, \rho) = \frac{S(r, \rho)\sqrt{h(r)}}{\sqrt{h(\rho)}} = \frac{G(r, \rho)}{\sqrt{h(r)h(\rho)}}. \quad (30)$$

Then, we have

$$\begin{aligned} \sigma(t) \left[q(r, t) - \int_1^t V_1(t, \tau)q(r, \tau) d\tau \right] + \int_0^1 G^h(r, \rho)q(\rho, t)\rho d\rho \\ - \int_1^t V_2(t, \tau) \int_0^1 G^h(r, \rho)q(\rho, \tau)\rho d\rho d\tau = \frac{f(r, t)}{\sqrt{h(r)}}, \quad 0 \leq r \leq 1, \quad 1 \leq t \leq T, \end{aligned} \quad (31)$$

where $q(r, t)$ and $f(r, t)/\sqrt{h(r)}$ are continuous functions of $t \in [1, T]$ with values in $\widehat{L}_2(\omega)$; $G^h(r, \rho)$ is a symmetric positive definite Hilbert–Schmidt kernel; and the other functions are the same as above.

Suppose that the right hand side of equation (31) is known and it is required to find the function $q(r, t)$.

Let us seek a solution of the mixed equation (31) in the form of a series

$$q(r, t) = \sum_{k=1}^{\infty} q_k(t)\varphi_k^h(r), \quad (32)$$

where $\varphi_k^h(r)$ are eigenfunctions of the kernel $G^h(r, \rho)$ corresponding to eigenvalues $\nu_k^h > 0$, i.e.,

$$\int_0^1 G^h(r, \rho)\varphi_k^h(\rho)\rho d\rho = \nu_k^h\varphi_k^h(r), \quad k = 1, 2, \dots \quad (33)$$

The representation (33) is possible, since the system of eigenfunctions of the kernel $G^h(r, \rho)$ forms a basis in $\widehat{L}_2(\omega)$.

Here, in contrast to the above cases, we construct the basis in the form

$$\varphi_k^h(r) = \frac{\Psi_k^h(r)}{\sqrt{h(r)}} \quad k = 1, 2, \dots \quad (34)$$

with explicit dependence on the function $h(r)$, where

$$\int_0^1 \varphi_i^h(\rho)\varphi_j^h(\rho)\rho d\rho = \int_0^1 \frac{\Psi_i^h(\rho)\Psi_j^h(\rho)}{h(\rho)}\rho d\rho = \delta_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases} \quad (35)$$

In order to construct such eigenfunctions, we first construct a basis $z_n^h(r)$ in $\widehat{L}_2(\omega)$ for which

$$\int_0^1 z_i^h(\rho)z_j^h(\rho)\rho d\rho = \delta_{ij}, \quad z_n^h(r) = \frac{Q_{n-1}^h(r)}{\sqrt{h(r)}}, \quad n = 1, 2, \dots \quad (36)$$

Such a basis can be constructed by the formulas

$$Q_0^h(r) = \frac{1}{\sqrt{I_0}}, \quad Q_n^h(r) = \frac{1}{\sqrt{D_{n-1}D_n}} \begin{vmatrix} I_0 & I_1 & \cdots & I_n \\ I_1 & I_2 & \cdots & I_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ I_{n-1} & I_n & \cdots & I_{2n-1} \\ 1 & r^2 & \cdots & r^{2n} \end{vmatrix}, \tag{37}$$

$$D_{-1} = 1, \quad D_0 = I_0, \quad D_n = \begin{vmatrix} I_0 & I_1 & \cdots & I_n \\ I_1 & I_2 & \cdots & I_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ I_n & I_{n+1} & \cdots & I_{2n} \end{vmatrix}, \quad I_n = \int_0^1 \frac{\rho^{2n+1}}{h(\rho)} d\rho.$$

Let us represent the k th eigenfunction as a series in terms of the basis $z_i^h(r)$ in $\widehat{L}_2(\omega)$. Then

$$\psi_k^h(r) = \sum_{i=1}^{\infty} \psi_{i(k)}^h z_i^h(r), \quad z_i^h(r) = \frac{Q_{i-1}^h(r)}{\sqrt{h(r)}}, \quad \Psi_k^h(r) = \sum_{i=1}^{\infty} \psi_{i(k)}^h Q_{i-1}^h(r). \tag{38}$$

For the Hilbert–Schmidt kernel $Q^h(r, \rho)$ we use the double series expansion with respect to the chosen basis:

$$Q^h(r, \rho) = \sum_{m=1}^{\infty} \sum_{i=1}^{\infty} Q_{mn}^h z_m^h(r) z_n^h(\rho), \tag{39}$$

$$Q_{mn}^h = \int_0^1 \int_0^1 Q^h(r, \rho) z_m^h(r) z_n^h(\rho) r \rho dr d\rho, \quad Q_{mn}^h = Q_{nm}^h.$$

Substituting (38) and (39) into (33), we obtain an infinite system of linear algebraic equations for the determination of the eigenvalues and the eigenfunction expansion coefficients. This system has a symmetric matrix and can be written in the form

$$\sum_{n=1}^{\infty} Q_{mn}^h \psi_{n(k)}^h = \nu_k^h \psi_{m(k)}^h, \quad m = 1, 2, \dots \tag{40}$$

In order to calculate approximations for N eigenvalues and eigenfunctions of the Hilbert–Schmidt kernel, it is necessary to find the eigenvalues and orthonormal eigenvectors of the matrix

$$[G_{NN}^h] = \begin{pmatrix} G_{11}^h & G_{12}^h & G_{13}^h & \cdots & G_{1N}^h \\ G_{12}^h & G_{22}^h & G_{23}^h & \cdots & G_{2N}^h \\ G_{13}^h & G_{23}^h & G_{33}^h & \cdots & G_{3N}^h \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ G_{1N}^h & G_{2N}^h & G_{3N}^h & \cdots & G_{NN}^h \end{pmatrix}. \tag{41}$$

The eigenvalues of the matrix (41) give approximations of the first N eigenvalues of the Hilbert–Schmidt kernel, and the components of its orthonormal eigenvectors give the coefficients in the expansion of the first N eigenfunctions of this kernel in terms of N orthonormal basis functions of the space $\widehat{L}_2(\omega)$.

Now, consider the expansion of the right-hand side of equation (31) into the following series:

$$\frac{f(r, t)}{\sqrt{h(r)}} = \sum_{k=1}^{\infty} f_k^h(t) \psi_k^h(r) = \sum_{k=1}^{\infty} f_k^h(t) \frac{\Psi_k^h(r)}{\sqrt{h(r)}}, \tag{42}$$

$$f_k^h(t) = \int_0^1 \frac{f(\rho, t)}{\sqrt{h(\rho)}} \psi_k^h(\rho) \rho d\rho = \int_0^1 \frac{f(\rho, t)}{h(\rho)} \Psi_k^h(\rho) \rho d\rho.$$

Substituting (32), (42) into (31) and taking into account (33), we obtain the following sequence of Volterra equations for the unknown functions $q_k(t)$:

$$q_k(t) - \int_1^t V_{k(\nu)}^h(t, \tau) q_k(\tau) d\tau = \gamma_k^h(t), \quad \gamma_k^h(t) = \frac{f_k^h(t)}{\sigma(t) + \nu_k^h} \quad (43)$$

$$V_{k(\nu)}^h(t, \tau) = \frac{\sigma(t)V_1(t, \tau) + \nu_k^h V_2(t, \tau)}{\sigma(t) + \nu_k^h}, \quad k = 1, 2, \dots, \quad (44)$$

where $V_{k(\nu)}^h(t, \tau)$ are Volterra kernels of the same class as $V_1(t, \tau)$ and $V_2(t, \tau)$, since $\nu_k^h \rightarrow 0$ as $k \rightarrow \infty$.

A solution of the sequence of Volterra equations (44) can be constructed by analytical and numerical methods of Chapter 11. This solution can be written in the form

$$q_k(t) = \gamma_k^h(t) + \int_1^t R_{k(\nu)}^h(t, \tau) \gamma_k^h(\tau) d\tau, \quad (45)$$

where $R_{k(\nu)}^h(t, \tau)$ is the resolvent of the kernel $V_{k(\nu)}^h(t, \tau)$.

The series (32) converges in $\widehat{L}_2(\omega)$ uniformly with respect to $t \in [1, T]$, and its sum is a continuous function of $t \in [1, T]$ with values in $\widehat{L}_2(\omega)$.

Finally, in view of the transformation of the variables (30) and formula (34) for the eigenfunctions, we have

$$y(r, t) = \frac{1}{h(r)} \sum_{k=1}^{\infty} q_k(t) \Psi_k^h(r). \quad (46)$$

Note that the solution (46) explicitly depends on the function $h(r)$, which allows us to solve equation (29) with great accuracy by keeping a small number of terms of the series. In the case of a strongly oscillating function $h(r)$, it is hardly possible to construct a solution by other known methods.

17.3-4. Equation with a Schmidt Kernel and Auxiliary Conditions on Ring-Shaped Domain.

Consider equation (29) with the right-hand side of the form $f(r, t) = \beta(t) - w(r, t)$ and an integral condition of the form (8) from Subsection 17.1-3 on the unknown function $y(r, t)$. The problem is to find a solution of the mixed integral equation

$$\begin{aligned} \sigma(t) \left[y(r, t) - \int_1^t V_1(t, \tau) y(r, \tau) d\tau \right] + \int_0^1 S_\omega(r, \rho) y(\rho, t) \rho d\rho \\ - \int_1^t V_2(t, \tau) \int_0^1 S_\omega(r, \rho) y(\rho, \tau) \rho d\rho d\tau = \frac{\beta(t)}{h(r)} - \frac{w(r, t)}{h(r)}, \quad (47) \\ S_\omega(r, \rho) = \frac{G(r, \rho)}{h(r)}, \quad 0 \leq r \leq 1, \quad 1 \leq t \leq T \end{aligned}$$

with the auxiliary condition

$$\int_0^1 y(\rho, t) \rho d\rho = M(t), \quad (48)$$

where the unknown functions are the following: $y(r, t)$, $\beta(t)$. All the other functions in (47) and (48) are assumed known, and $w(r, t)$ is a continuous function of $t \in [1, T]$ with values in $\widehat{L}_2(\omega)$.

Let us transform the equation with the Schmidt kernel to an equation with a Hilbert–Schmidt kernel by changing the variables according to (30). Then, equation (47) and the auxiliary conditions (48) become

$$\sigma(t) \left[q(r, t) - \int_1^t V_1(t, \tau) q(r, \tau) d\tau \right] + \int_0^1 G^h(r, \rho) q(\rho, t) \rho d\rho - \int_{\tau_0}^t V_2(t, \tau) \int_0^1 G^h(r, \rho) q(\rho, \tau) \rho d\rho d\tau = \frac{\beta(t)}{\sqrt{h(r)}} - \frac{w(r, t)}{\sqrt{h(r)}}, \tag{49}$$

$$\int_0^1 \frac{q(\rho, t)}{\sqrt{h(\rho)}} \rho d\rho = M(t), \quad 0 \leq r \leq 1, \quad 1 \leq t \leq T. \tag{50}$$

To construct a solution of the mixed integral equation (47) with the auxiliary conditions (48), we use the basis $z_n^h(r)$ of $\widehat{L}_2(\omega)$ (see (36) and (37)) and note that the space $\widehat{L}_2(\omega)$ can be represented as a direct sum of its orthogonal subspaces: $\widehat{L}_2(\omega) = L_2^{h^o}(\omega) \oplus L_2^{h^*}(\omega)$ (see Supplement 12.5-3), where $\widehat{L}_2^{h^o}(\omega)$ is the Euclidean space with the basis $z_1^h(r)$, and $\widehat{L}_2^{h^*}(\omega)$ is the Hilbert space with the basis $p_k^h(r)$ ($k = 2, 3, \dots$). Note also that the integrand and the right-hand side can be represented as a sum of continuous functions of $t \in [1, T]$ with values in $\widehat{L}_2^{h^o}(\omega)$ and $\widehat{L}_2^{h^*}(\omega)$, respectively, i.e.,

$$q(r, t) = q^\circ(r, t) + q^*(r, t), \quad \frac{f(r, t)}{\sqrt{h(r)}} = f_h^\circ(r, t) + f_h^*(r, t), \tag{51}$$

where the following representations hold:

$$\begin{aligned} q^\circ(r, t) &= q_1^\circ(t) z_1^h(r), & q_1^\circ(t) &= \frac{M(t)}{\sqrt{I_0}}, \\ \frac{f(r, t)}{\sqrt{h(r)}} &= \frac{\beta(t)}{\sqrt{h(r)}} - \frac{w(r, t)}{\sqrt{h(r)}}, & \frac{w(r, t)}{\sqrt{h(r)}} &= w_h^\circ(r, t) + w_h^*(r, t), \\ f_h^\circ(r, t) &= [\sqrt{I_0} \beta(t) - w_1(t)] z_1^h(r), & f_h^*(r, t) &= -w_h^*(r, t), \\ w_h^\circ(r, t) &= w_1^{h^o}(t) z_1^h(r), & w_1^{h^o}(t) &= \int_0^1 \frac{w(\rho, t)}{\sqrt{h(\rho)}} z_1^h(\rho) \rho d\rho. \end{aligned} \tag{52}$$

Note that in the representation (52) for $q(r, t)$, the term $q^\circ(r, t)$ is known (as determined by the auxiliary conditions), and the term $q^*(r, t)$ is to be found. For the right-hand side, the term $f_h^\circ(r, t)$ should be found and $f_h^*(r, t)$ is given by $g(r, t)/\sqrt{h(r)}$. Thus, we have come to a special case of the general projection problem whose solution is constructed in Subsection 17.4-3.

According to the general method, in this case, one can introduce an operator of orthogonal projection that maps $\widehat{L}_2(\omega)$ onto $\widehat{L}_2^{h^o}(\omega)$:

$$\mathbf{Q}_h^\circ \phi(r, t) = \int_0^1 \phi(\rho, t) z_1^h(r) z_1^h(\rho) \rho d\rho. \tag{53}$$

Obviously, the orthogonal projector $\mathbf{Q}_h^* = \mathbf{I} - \mathbf{Q}_h^\circ$ maps $\widehat{L}_2(\omega)$ onto $\widehat{L}_2^{h^*}(\omega)$. Moreover, the following relations hold:

$$\begin{aligned} \mathbf{Q}_h^\circ q(r, t) &= q^\circ(r, t), & \mathbf{Q}_h^* q(r, t) &= q^*(r, t), \\ \mathbf{Q}_h^\circ \frac{f(r, t)}{\sqrt{h(r)}} &= f_h^\circ(r, t), & \mathbf{Q}_h^* \frac{f(r, t)}{\sqrt{h(r)}} &= f_h^*(r, t). \end{aligned} \tag{54}$$

Following Section 17.4, we apply the projection operator \mathbf{Q}_h^* to equation (49) and obtain an integral equation in the space $\widehat{L}_2^{h*}(\omega)$ with a known right-hand side. This is the equation for the determination of $q^*(x, t)$:

$$\begin{aligned} &\sigma(t) \left[q^*(r, t) - \int_1^t V_1(t, \tau) q^*(r, \tau) d\tau \right] \\ &\quad + \int_0^1 G_h^*(r, \rho) q^*(\rho, t) \rho d\rho - \int_1^t V_2(t, \tau) \int_0^1 G_h^*(r, \rho) q^*(\rho, \tau) \rho d\rho d\tau \\ &= -g_h^*(r, t) - \int_0^1 G_h^*(r, \rho) q^\circ(\rho, t) \rho d\rho + \int_1^t V_2(t, \tau) \int_0^1 G_h^*(r, \rho) q^\circ(\rho, \tau) \rho d\rho d\tau, \\ &\quad 0 \leq r \leq 1, \quad 1 \leq t \leq T, \end{aligned} \tag{55}$$

where the kernel of the integral equation

$$G_h^*(r, \rho) = G^h(r, \rho) - \int_0^1 G^h(s, \rho) z_1^h(r) z_1^h(s) ds \tag{56}$$

is of Hilbert–Schmidt type.

Let us construct a solution of equation (55) in the form of a series with respect to eigenfunctions of the kernel (56). These eigenfunctions form a basis in the Hilbert space $\widehat{L}_2^{h*}(\omega)$. Let us construct a system of these functions.

Let $\psi_k^{h*}(r)$ be eigenfunctions and ν_k^{h*} the corresponding eigenvalues of the kernel $G_h^*(r, \rho)$, i.e.,

$$\int_0^1 G_h^*(r, \rho) \psi_k^{h*}(\rho) \rho d\rho = \nu_k^{h*} \psi_k^{h*}(r), \quad k = 2, 3, \dots \tag{57}$$

Let us represent the eigenfunction $\psi_i^{h*}(r)$ in the form of a series with respect to the basis $z_i^h(r)$ ($i \geq 2$):

$$\psi_k^{h*}(r) = \sum_{i=2}^\infty \psi_{i(k)}^{h*} z_i^h(r), \quad z_i^h(r) = \frac{Q_{i-1}^h(r)}{\sqrt{h(r)}}, \quad \Psi_k^{h*}(r) = \sum_{i=2}^\infty \psi_{i(k)}^{h*} Q_{i-1}^h(r). \tag{58}$$

Using (39) and (56), we obtain a double series expansion for the kernel $G_h^*(r, \rho)$:

$$G_h^*(x, \xi) = \sum_{m=2}^\infty \sum_{n=2}^\infty G_{mn}^h z_m^h(r) z_n^h(\rho) + \sum_{n=2}^\infty G_{1n}^h z_n^h(r) z_1^h(\rho). \tag{59}$$

Note that the coefficients of the expansion of $G_h^*(r, \rho)$ in (59) coincide with those of the expansion of $G^h(r, \rho)$, which allows us to avoid recalculation of the coefficients of the new problem and use the available data.

Substituting (58) and (59) into (57), we obtain an infinite system of linear algebraic equations for the determination of the eigenvalues and eigenfunction expansion coefficients. This system has a symmetric matrix and can be written as follows:

$$\sum_{n=2}^\infty G_{mn}^h \psi_{n(k)}^{h*} = \nu_k^{h*} \psi_{m(k)}^{h*}, \quad m = 2, 3, \dots \tag{60}$$

Now, let us construct a solution of equation (55). For this purpose, we represent the functions $q^*(r, t)$ and $w_h^*(r, t)$ in the form of series with eigenfunctions of the kernel $G_h^*(r, \rho)$:

$$q^*(r, t) = \sum_{k=2}^{\infty} q_k^*(t)\psi_k^{h*}(r), \quad w_h^*(r, t) = \sum_{k=2}^{\infty} w_k^{h*}(t)\psi_k^{h*}(r), \quad w_k^{h*}(t) = \int_0^1 w_h^*(\rho, t)\psi_k^{h*}(\rho)\rho \, d\rho, \quad (61)$$

and substitute these into (55). Then, taking into account (52), (56)–(59), we obtain a sequence of independent Volterra equations of the second kind:

$$\begin{aligned} q_k^*(t) - \int_1^t V_{k(\nu)}^{h*}(t, \tau)q_k^*(\tau) \, d\tau &= \gamma_k^{h*}(t), \quad V_{k(\nu)}^{h*}(t, \tau) = \frac{\sigma(t)V_1(t, \tau) + \nu_k^{h*}V_2(t, \tau)}{\sigma(t) + \nu_k^{h*}}, \\ \gamma_k^{h*}(t) &= -\frac{1}{\sigma(t) + \mu_k^{h*}} \left[g_k^{h*}(t) + G_k^h q_1^\circ(t) - \int_1^t V_2(t, \tau)G_k^h q_1^\circ(\tau) \, d\tau \right], \\ G_k^h &= \sum_{n=2}^{\infty} G_n^h \psi_n^{h*}, \quad k = 2, 3, \dots \end{aligned} \quad (62)$$

Resolving (62) with respect to $q_k^*(t)$ by the methods of Chapter 11, we get

$$q_k^*(t) = \gamma_k^{h*}(t) + \int_1^t R_{k(\nu)}^{h*}(t, \tau)\gamma_k^{h*}(\tau) \, d\tau, \quad (63)$$

where $R_{k(\nu)}^{h*}(t, \tau)$ is the resolvent of the kernel $V_{k(\nu)}^{h*}(t, \tau)$.

Now, in view of (61)–(63), the function $q^*(r, t)$ has been found, as well as the function $q(r, t)$, since $q^\circ(r, t)$ is known by assumption (see (51) and (52)). Hence, using (30), we finally obtain

$$y(r, t) = \frac{1}{h(r)} \left[\frac{M(t)}{\sqrt{I_0}} Q_0^h(r) + \sum_{k=2}^{\infty} q_k^*(t)\Psi_k^{h*}(r) \right]. \quad (64)$$

Note that the function $h(r)$ enters solution (64) explicitly, which allows us to solve equation (47) with high accuracy by keeping a relatively small number of terms of the series even in the case of a rapidly oscillating $h(r)$.

In practical calculations, the number of terms in the series is taken finite. For instance, taking the functions $z_k^h(r)$ of the basis with $k = 2, \dots, N$, we obtain the N th approximation of the desired solution. In this case, for the construction of eigenvalues and eigenfunctions of the Hilbert–Schmidt kernel $G_h^*(r, \rho)$ one should find the eigenvalues and the orthonormal eigenfunctions of the matrix

$$[G_{NN}^h] = \begin{pmatrix} G_{33}^h & G_{34}^h & G_{35}^h & \dots & G_{3N}^h \\ G_{34}^h & G_{44}^h & G_{45}^h & \dots & G_{4N}^h \\ G_{35}^h & G_{45}^h & G_{55}^h & \dots & G_{5N}^h \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ G_{3N}^h & G_{4N}^h & G_{5N}^h & \dots & G_{NN}^h \end{pmatrix}. \quad (65)$$

The eigenvalues of the matrix (65) give approximations for the first $N - 1$ eigenvalues of the Hilbert–Schmidt operator, and the components of its orthonormal eigenvectors give the expansion coefficients for the first $N - 1$ eigenfunctions of that operator. Recall that the first term $q^\circ(r, t)$ in the expansion of $q(r, t)$ is known by assumption. Therefore, constructing the next $N - 1$ terms of the expansion, we obtain the N th approximation of the solution.

It is important to keep in mind the relation between the matrices (41) and (65). The matrix (64) is obtained from the matrix (41) by deleting its first row and column. This allows us to construct

an expansion of the original kernel only once and then use this data for the examination of the new kernel arising in the problem with auxiliary conditions.

Now, let us find the function $\beta(t)$. To this end, we apply the projection operator \mathbf{Q}_h° to equation (55). As a result, we obtain the following formulas:

$$\beta(t) = \frac{1}{\sqrt{I_0}} \left\{ w^{h_0}(t) + \sigma(t) \left[q_1^\circ(t) - \int_1^t V_1(t, \tau) q_1^\circ(\tau) d\tau \right] + G_{11}^h q_1^\circ(t) + \sum_{k=3}^\infty G_k^h q_k^*(t) - \int_1^t V_2(t, \tau) \left[G_{11}^h q_1^\circ(\tau) + \sum_{k=2}^\infty G_k^h q_k^*(\tau) \right] d\tau \right\}. \tag{66}$$

Thus, we have constructed a complete solution of the integral equation (47) with the auxiliary conditions (48).

⊙ References for Section 17.3: E. Goursat (1923), G. Szegö (1975), A. V. Manzhirov (1985, 2005), N. Kh. Arutynyan, A. V. Manzhirov, and V.E. Naumov (1991), N. Kh. Arutynyan, A. V. Manzhirov (1999), A. V. Manzhirov and K. E. Kazakov (2006).

17.4. Projection Method for Solving Mixed Equations on a Bounded Set

17.4-1. Mixed Operator Equation with a Given Right-Hand Side.

Consider a mixed multi-dimensional equation of the form (10) of Subsection 17.1-4 with integral operators of Volterra and Schmidt types:

$$\begin{aligned} \sigma(t)(\mathbf{I} - \mathbf{V}_1)y(\vec{x}, t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{S}y(\vec{x}, t) &= \frac{f(\vec{x}, t)}{h(\vec{x})}, \\ \mathbf{S}y(\vec{x}, t) &= \int_\Omega S(\vec{x}, \vec{\xi})y(\vec{\xi}, t) d\Omega_\xi, \quad S(\vec{x}, \vec{\xi}) = \frac{F(\vec{x}, \vec{\xi})}{h(\vec{x})}, \\ \mathbf{V}_p y(\vec{x}, t) &= \int_{\tau_0}^t V_p(t, \tau)y(\vec{x}, \tau) d\tau, \quad \vec{x} \in \Omega, \quad \tau_0 \leq t \leq T. \end{aligned} \tag{1}$$

In this section, we consider some general questions of the theory of mixed equations. For this reason we do not single out equations with the Hilbert–Schmidt integral operator of the form (9) from Subsection 17.1-4 and only mention that this equation is a special case of equation (1) with the Schmidt integral operator with $h(\vec{x}) = 1$.

Let the right-hand side $f(\vec{x}, t)/h(\vec{x})$ of equation (1) be known. It is required to find the function $y(\vec{x}, t)$. Here, $f(\vec{x}, t)$ and $y(\vec{x}, t)$ are continuous functions of $t \in [1, T]$ with values in $L_2(\Omega)$; $\sigma(t)$ is a given positive continuous function; $h(\vec{x}) > 0$ is a given function of class $L_2(\Omega)$; $F(\vec{x}, \vec{\xi})$ is a symmetric positive definite Fredholm kernel; \mathbf{V}_1 and \mathbf{V}_2 are Volterra operators; and \mathbf{S} is a Schmidt operator.

Let us transform the equation with the Schmidt operator to an equation with a Hilbert–Schmidt operator. To this end, we multiply (1) by $\sqrt{h(\vec{x})}$ and change the variables as follows:

$$q(\vec{x}, t) = \sqrt{h(\vec{x})} y(\vec{x}, t), \quad F^h(\vec{x}, \vec{\xi}) = \frac{S(\vec{x}, \vec{\xi})\sqrt{h(\vec{x})}}{\sqrt{h(\vec{\xi})}} = \frac{F(\vec{x}, \vec{\xi})}{\sqrt{h(\vec{x})h(\vec{\xi})}}. \tag{2}$$

Then

$$\begin{aligned} \sigma(t)(\mathbf{I} - \mathbf{V}_1)q(\vec{x}, t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{F}^h q(\vec{x}, t) &= \frac{f(\vec{x}, t)}{\sqrt{h(\vec{x})}}, \\ \mathbf{F}^h q(\vec{x}, t) &= \int_\Omega F^h(\vec{x}, \vec{\xi})q(\vec{\xi}, t) d\Omega_\xi, \quad \vec{x} \in \Omega, \quad \tau_0 \leq t \leq T. \end{aligned} \tag{3}$$

where $q(\vec{x}, t)$ and $f(\vec{x}, t)/\sqrt{h(\vec{x})}$ are continuous functions of $t \in [\tau_0, T]$ with values in the Hilbert space $L_2(\Omega)$; \mathbf{F}^h is a Hilbert–Schmidt operator; and the other functions have been specified above.

Suppose that the right-hand side of equation (3) is known and we have to find the function $q(\vec{x}, t)$. Let us seek a solution of the mixed equation (3) in the form of a series

$$q(\vec{x}, t) = \sum_{k=1}^{\infty} q_k(t)\varphi_k^h(\vec{x}), \tag{4}$$

where $\varphi_k^h(\vec{x})$ are eigenfunctions of the operator \mathbf{F}^h corresponding to eigenvalues $\mu_k^h > 0$, i.e.,

$$\mathbf{F}^h \varphi_k^h(\vec{x}) d\xi = \mu_k^h \varphi_k^h(\vec{x}), \quad k = 1, 2, \dots \tag{5}$$

The representation (4) is possible, since the system of eigenfunctions of the operator \mathbf{F}^h forms a basis in $L_2(\Omega)$.

Let us construct the functions of the basis in the form

$$\varphi_k^h(\vec{x}) = \frac{\Phi_k^h(\vec{x})}{\sqrt{h(\vec{x})}}, \quad k = 1, 2, \dots \tag{6}$$

with explicit dependence on the function $h(\vec{x})$, where

$$\int_{\Omega} \varphi_i^h(\vec{\xi})\varphi_j^h(\vec{\xi}) d\Omega_{\xi} = \int_{\Omega} \frac{\Phi_i^h(\vec{\xi})\Phi_j^h(\vec{\xi})}{h(\vec{\xi})} d\Omega_{\xi} = \delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases} \tag{7}$$

In order to construct such eigenfunctions, we first construct a basis $p_n^h(\vec{x})$ in $L_2(\Omega)$ for which

$$\int_{\Omega} p_i^h(\vec{\xi})p_j^h(\vec{\xi}) d\Omega_{\xi} = \delta_{ij}, \quad p_n^h(\vec{x}) = \frac{P_n^h(\vec{x})}{\sqrt{h(\vec{x})}}, \quad n = 1, 2, \dots \tag{8}$$

Such a basis can be constructed by the formulas

$$P_1^h(\vec{x}) = \frac{f_1(\vec{x})}{\sqrt{H_{11}}}, \quad P_n^h(\vec{x}) = \frac{1}{\sqrt{\Delta_{n-1}\Delta_n}} = \begin{vmatrix} H_{11} & H_{12} & \dots & H_{1n} \\ H_{21} & H_{22} & \dots & H_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ f_1(\vec{x}), & f_2(\vec{x}) & \dots & f_n(\vec{x}) \end{vmatrix}, \tag{9}$$

$$\Delta_0 = 1, \quad \Delta_1 = H_{11}, \quad \Delta_n = \begin{vmatrix} H_{11} & H_{12} & \dots & H_{1n} \\ H_{21} & H_{22} & \dots & H_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ H_{n1} & H_{n2} & \dots & H_{nn} \end{vmatrix}, \quad H_{ij} = \int_{\Omega} \frac{f_i(\vec{\xi})f_j(\vec{\xi})}{h(\vec{\xi})} d\Omega_{\xi},$$

where $f_i(\vec{x})$ is an arbitrary complete system of linearly independent function in $L_2(\Omega)$.

Let us represent the k th eigenfunction in the form of a series with respect to the basis $p_i^h(\vec{x})$ of $L_2(\Omega)$. We have

$$\varphi_k^h(\vec{x}) = \sum_{i=1}^{\infty} \varphi_{i(k)}^h p_i^h(\vec{x}), \quad p_i^h(\vec{x}) = \frac{P_i^h(\vec{x})}{\sqrt{h(\vec{x})}}, \quad \Phi_k^h(\vec{x}) = \sum_{i=1}^{\infty} \varphi_{i(k)}^h P_i^h(\vec{x}). \tag{10}$$

The Hilbert–Schmidt kernel $F^h(\vec{x}, \vec{\xi})$ can be expanded into double series with respect to the chosen basis:

$$F^h(\vec{x}, \vec{\xi}) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} F_{mn}^h p_m^h(\vec{x}) p_n^h(\vec{\xi}), \tag{11}$$

$$F_{mn}^h = \int_{\Omega} \int_{\Omega} F^h(\vec{x}, \vec{\xi}) p_m^h(\vec{x}) p_n^h(\vec{\xi}) d\Omega_x d\Omega_{\xi}, \quad F_{mn}^h = F_{nm}^h.$$

Substituting (10) and (11) into (5), we obtain an infinite system of linear algebraic equations for the determination of the eigenvalues and the eigenfunction expansion coefficients. This system has a symmetric matrix and can be written as follows:

$$\sum_{n=1}^{\infty} F_{mn}^h \varphi_{n(k)}^h = \mu_k^h \varphi_{m(k)}^h, \quad m = 1, 2, \dots \tag{12}$$

In order to calculate approximations for N eigenvalues and eigenfunctions of the Hilbert–Schmidt operator \mathbf{F}^h , it is necessary to find the eigenvalues and orthonormal eigenvectors of the matrix

$$[F_{NN}^h] = \begin{pmatrix} F_{11}^h & F_{12}^h & F_{13}^h & \dots & F_{1N}^h \\ F_{12}^h & F_{22}^h & F_{23}^h & \dots & F_{2N}^h \\ F_{13}^h & F_{23}^h & F_{33}^h & \dots & F_{3N}^h \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ F_{1N}^h & F_{2N}^h & F_{3N}^h & \dots & F_{NN}^h \end{pmatrix}. \tag{13}$$

The eigenvalues of the matrix (13) give approximations for the first N eigenvalues of the Hilbert–Schmidt operator, and the components of its orthonormal eigenvectors give approximate values of the expansion coefficients for the first N eigenfunctions of that series with N orthonormal functions of the basis.

Let us write the right-hand side of equation (3) in the form

$$\begin{aligned} \frac{f(\vec{x}, t)}{\sqrt{h(\vec{x})}} &= \sum_{k=1}^{\infty} f_k^h(t) \varphi_k^h(\vec{x}) = \sum_{k=1}^{\infty} f_k^h(t) \frac{\Phi_k^h(\vec{x})}{\sqrt{h(\vec{x})}}, \\ f_k^h(t) &= \int_{\Omega} \frac{f(\vec{\xi}, t)}{\sqrt{h(\vec{\xi})}} \varphi_k^h(\vec{\xi}) d\Omega_{\xi} = \int_{-1}^1 \frac{f(\vec{x}, t)}{h(\vec{x})} \Phi_k^h(\vec{x}) dx. \end{aligned} \tag{14}$$

Substituting (4), (14) into (3) and taking into account (5), we obtain the following sequence of Volterra equations for the unknown functions $q_k(t)$:

$$\begin{aligned} (\mathbf{I} - \mathbf{V}_k^h) q_k(t) &= \delta_k^h(t), \quad \delta_k^h(t) = \frac{f_k^h(t)}{\sigma(t) + \mu_k^h}, \\ \mathbf{V}_k^h &= \frac{\sigma(t) \mathbf{V}_1 + \mu_k^h \mathbf{V}_2}{\sigma(t) + \mu_k^h}, \quad \mathbf{V}_k^h f(t) = \int_{\tau_0}^t V_k^h(t, \tau) f(\tau) d\tau, \\ V_k^h &= \frac{\sigma(t) V_1(t, \tau) + \mu_k^h V_2(t, \tau)}{\sigma(t) + \mu_k^h}, \quad k = 1, 2, \dots, \end{aligned} \tag{15}$$

where all operators \mathbf{V}_k^h are of Volterra type, just as the operators \mathbf{V}_1 and \mathbf{V}_2 , since $\mu_k^h \rightarrow 0$ as $k \rightarrow \infty$.

A solution of the sequence of Volterra equations (15) can be constructed by analytical and numerical methods of Chapter 11. This solution can be written in the form

$$q_k(t) = (\mathbf{I} + \mathbf{R}_k^h) \delta_k^h(t), \quad (\mathbf{I} - \mathbf{V}_k^h)^{-1} = (\mathbf{I} + \mathbf{R}_k^h), \quad \mathbf{R}_k^h f(t) = \int_{\tau_0}^t R_k^h(t, \tau) f(\tau) d\tau, \tag{16}$$

where \mathbf{R}_k^h is the resolvent operator for \mathbf{V}_k^h , and $R_k(t, \tau)$ is the resolvent of the kernel $V_k^h(t, \tau)$. The series (13) converges in $L_2(\Omega)$ uniformly in $t \in [\tau_0, t]$, and its sum is a continuous function of t with values in $L_2(\Omega)$.

Finally, taking into account (2), (4), and (6), (16), we find that

$$y(\vec{x}, t) = \frac{1}{h(\vec{x})} \sum_{k=1}^{\infty} (\mathbf{I} + \mathbf{R}_k^h) \delta_k^h(t) \Phi_k^h(\vec{x}). \tag{17}$$

Note that the function $h(\vec{x})$ enters the solution (17) in explicit form, which allows us to solve equation (1) with high accuracy, even for a rapidly oscillating function $h(\vec{x})$.

17.4-2. Mixed Operator Equations with Auxiliary Conditions.

Consider equation (1) with the right-hand side $f(\vec{x}, t) = \sum_{i=1}^N \alpha_i(t) f_i(\vec{x}) - g(\vec{x}, t)$ and N auxiliary integral conditions (of the form (12) from Subsection 17.1-4) on the unknown function $y(x, t)$. The problem is to find a solution of the operator equation

$$\sigma(t)(\mathbf{I} - \mathbf{V}_1)y(\vec{x}, t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{S}y(\vec{x}, t) = \sum_{i=1}^N \alpha_i(t) \frac{f_i(\vec{x})}{h(\vec{x})} - \frac{g(\vec{x}, t)}{h(\vec{x})}, \quad \vec{x} \in \Omega, \quad \tau_0 \leq t \leq T \quad (18)$$

with the auxiliary conditions

$$\int_{\Omega} y(\vec{\xi}, t) f_i(\vec{\xi}) d\Omega_{\xi} = M_i(t), \quad i = 1, \dots, N, \quad (19)$$

regarding $y(\vec{x}, t)$ and $\alpha_1(t), \dots, \alpha_N(t)$ as unknown functions. All other functions in (18) are assumed given, and $g(\vec{x}, t)$ is a continuous function of t with values in $L_2(\Omega)$; $f_i(\vec{x})$ is a system of N linearly independent functions in $L_2(\Omega)$.

Let us transform the equation with the Schmidt operator to an equation with a Hilbert–Schmidt operator by changing the variables as in (2). Then, equation (18) and the auxiliary conditions (19) become

$$\sigma(t)(\mathbf{I} - \mathbf{V}_1)q(\vec{x}, t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{F}^h q(\vec{x}, t) = \sum_{i=1}^N \alpha_i(t) \frac{f_i(\vec{x})}{\sqrt{h(\vec{x})}} - \frac{g(\vec{x}, t)}{\sqrt{h(\vec{x})}}, \quad (20)$$

$$\vec{x} \in \Omega, \quad \tau_0 \leq t \leq T,$$

$$\int_{\Omega} q(\vec{\xi}, t) \frac{f_i(\vec{\xi})}{\sqrt{h(\vec{\xi})}} d\Omega_{\xi} = M_i(t), \quad i = 1, \dots, N. \quad (21)$$

In order to construct a solution of the mixed integral equation (18) with the auxiliary conditions (19), we construct a special basis in $L_2(\Omega)$ with explicit dependence on the function $1/\sqrt{h(\vec{x})}$. To this end, we complement the system of N linearly independent functions $f_i(\vec{x})$, so as to obtain a complete system in $L_2(\Omega)$, and then use formulas (8) and (9). As a result, we obtain a basis $p_n^h(\vec{x})$ in $L_2(\Omega)$ for which (in view of (8) and (9)) the following expansion holds:

$$p_i^h(\vec{x}) = \frac{P_i^h(\vec{x})}{\sqrt{h(\vec{x})}} = \sum_{k=1}^i a_{ik} \frac{f_k(\vec{x})}{\sqrt{h(\vec{x})}}, \quad i = 1, \dots, N. \quad (22)$$

Resolving the system of algebraic equations (22), we obtain

$$\frac{f_i(\vec{x})}{\sqrt{h(\vec{x})}} = \sum_{k=1}^i b_{ik} p_k^h(\vec{x}), \quad i = 1, \dots, N, \quad (23)$$

the matrix of system (23) being the inverse of the matrix corresponding to system (22).

Let us represent the Hilbert space $L_2(\Omega)$ as the direct sum of its orthogonal subspaces:

$$L_2(\Omega) = L_2^{\circ}(\Omega) \oplus L_2^*(\Omega), \quad (24)$$

where $L_2^{\circ}(\Omega)$ is the Euclidean space with the basis $p_1(\vec{x}), \dots, p_N(\vec{x})$, and $L_2^*(\Omega)$ is the Hilbert space with the basis $\{p_k(\vec{x})\} (k = N + 1, N + 2, \dots)$.

Note that any continuous function of t with values in $L_2(\Omega)$ can be represented as a sum of continuous functions of t with values in $L_2^{\circ}(\Omega)$ and $L_2^*(\Omega)$. Let us write such a representation for the integrand:

$$q(\vec{x}, t) = q^{\circ}(\vec{x}, t) + q^*(\vec{x}, t), \quad q^{\circ}(\vec{x}, t) = \sum_{n=1}^N q_n^{\circ}(t) p_n^h(\vec{x}). \tag{25}$$

Using the auxiliary conditions (21), together with (23) and (25), we obtain the following system of equations:

$$\sum_{k=1}^i b_{ik} q_n^{\circ}(t) = M_i(t), \quad i = 1, \dots, N. \tag{26}$$

The solution of this system determines the coefficients of the first term in the expansion (25) of $q^{\circ}(\vec{x}, t)$:

$$q_i^{\circ}(t) = \sum_{k=1}^i a_{ik} M_k(t), \quad i = 1, \dots, N. \tag{27}$$

In view of (23), the right-hand side of the equation can be written in the form

$$\begin{aligned} \frac{f(\vec{x}, t)}{\sqrt{h(\vec{x})}} &= \sum_{i=1}^N \alpha_i(t) \frac{f_i(\vec{x})}{\sqrt{h(\vec{x})}} - \frac{g(\vec{x}, t)}{\sqrt{h(\vec{x})}} = f_h^{\circ}(\vec{x}, t) + f_h^*(\vec{x}, t), \\ f_h^{\circ}(\vec{x}, t) &= \sum_{k=1}^N \sum_{i=k}^N [\alpha_i(t) b_{ik} - g_k^{h^{\circ}}(t)] p_k^h(\vec{x}), \quad f_h^*(\vec{x}, t) = -g_h^*(\vec{x}, t), \\ \frac{g(\vec{x}, t)}{\sqrt{h(\vec{x})}} &= g_h^{\circ}(\vec{x}, t) + g_h^*(\vec{x}, t), \quad g_h^{\circ}(\vec{x}, t) = \sum_{k=1}^N g_k^{h^{\circ}}(t) p_k^h(\vec{x}), \\ g_k^{h^{\circ}}(t) &= \int_{\Omega} \frac{g(\vec{\xi}, t)}{\sqrt{h(\vec{\xi})}} p_k^h(\vec{\xi}) d\Omega_{\xi}, \quad k = 1, \dots, N. \end{aligned} \tag{28}$$

Note that in the representations (25)–(28) for $q(\vec{x}, t)$, the function $q^{\circ}(\vec{x}, t)$ is known (as determined by the auxiliary conditions), and the term $q^*(\vec{x}, t)$ is to be found. Conversely, for the right-hand side, we should find $f_h^{\circ}(\vec{x}, t)$, and $f_h^*(\vec{x}, t)$ is given by $g(\vec{x}, t)/\sqrt{h(\vec{x})}$. The facts mentioned above allow us to classify the resulting problem as a special case of the general projection problem considered in Subsection 17.4-3.

According to the general method, in the present case, one can introduce an operator of orthogonal projection that maps the space $L_2(\Omega)$ onto $L_2^{h^{\circ}}(\Omega)$:

$$\mathbf{P}_h^{\circ} f(\vec{x}) = \int_{\Omega} f(\vec{\xi}) \sum_{i=1}^N p_i^h(\vec{x}) p_i^h(\vec{\xi}) d\Omega_{\xi}. \tag{29}$$

Obviously, the orthogonal projector $\mathbf{P}_h^* = \mathbf{I} - \mathbf{P}_h^{\circ}$ maps $L_2(\Omega)$ onto $L_2^{h^*}(\Omega)$. Moreover, the following relations hold:

$$\begin{aligned} \mathbf{P}_h^{\circ} q(\vec{x}, t) &= q^{\circ}(\vec{x}, t), & \mathbf{P}_h^* q(\vec{x}, t) &= q^*(\vec{x}, t), \\ \mathbf{P}_h^{\circ} \frac{f(\vec{x}, t)}{\sqrt{h(\vec{x})}} &= f_h^{\circ}(\vec{x}, t), & \mathbf{P}_h^* \frac{f(\vec{x}, t)}{\sqrt{h(\vec{x})}} &= f_h^*(\vec{x}, t). \end{aligned} \tag{30}$$

Following Section 17.4, we apply the projection operator \mathbf{P}_h^* to equation (20) and obtain an integral equation in $L_2^{h*}(\Omega)$ (with a known right-hand side) for the determination of $q^*(\vec{x}, t)$:

$$\begin{aligned} \sigma(t)(\mathbf{I} - \mathbf{V}_1)q^*(\vec{x}, t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{P}_h^*\mathbf{F}^h q^*(\vec{x}, t) &= -g^*(\vec{x}, t) - (\mathbf{I} - \mathbf{V}_2)\mathbf{P}_h^*\mathbf{F}^h q^o(\vec{x}, t), \\ \mathbf{P}_h^*\mathbf{F}^h \phi(\vec{x}, t) &= \int_{\Omega} F_h^*(\vec{x}, \vec{\xi})\phi(\vec{\xi}, t) d\Omega_{\xi}, \quad \vec{x} \in \Omega, \quad \tau_0 \leq t \leq T, \\ F_h^*(\vec{x}, \vec{\xi}) &= F^h(\vec{x}, \vec{\xi}) - \int_{\Omega} F^h(\vec{s}, \vec{\xi}) \sum_{i=1}^N p_i^h(\vec{x})p_i^h(\vec{s}) d\Omega_s. \end{aligned} \tag{31}$$

The operator $\mathbf{P}_h^*\mathbf{F}^h$ is a Hilbert–Schmidt operator from $L_2^{h*}(\Omega)$ to $L_2^{h*}(\Omega)$. Let us construct a solution of equation (31) in the form of a series with respect to its eigenfunctions that form a basis in $L_2^{h*}(\Omega)$. Let us construct the system of these functions.

Let $\varphi_k^{h*}(\vec{x})$ be eigenfunctions of the operator $\mathbf{P}_h^*\mathbf{F}^h$ and μ_k^{h*} the corresponding eigenvalues. We have

$$\mathbf{P}_h^*\mathbf{F}^h \varphi_k^{h*}(\vec{x}) = \mu_k^{h*} \varphi_k^{h*}(\vec{x}), \quad k = N + 1, N + 2, \dots \tag{32}$$

Let us represent the eigenfunction $\varphi_i^{h*}(\vec{x})$ as a series with respect to the basis $p_i^h(\vec{x})$ ($i \geq N + 1$):

$$\varphi_k^{h*}(\vec{x}) = \sum_{i=N+1}^{\infty} \varphi_{i(k)}^{h*} p_i^h(\vec{x}), \quad p_i^h(\vec{x}) = \frac{P_i^h(\vec{x})}{\sqrt{h(\vec{x})}}, \quad \Phi_k^{h*}(\vec{x}) = \sum_{i=N+1}^{\infty} \varphi_{i(k)}^{h*} P_i^h(\vec{x}). \tag{33}$$

Using (11) and (31), we obtain the following double series expansion for the kernel $F_h^*(\vec{x}, \vec{\xi})$:

$$F_h^*(\vec{x}, \vec{\xi}) = \sum_{m=N+1}^{\infty} \sum_{n=N+1}^{\infty} F_{mn}^h p_m^h(\vec{x})p_n^h(\vec{\xi}) + \sum_{i=1}^N \sum_{n=N+1}^{\infty} F_{in}^h p_n^h(\vec{x})p_i^h(\vec{\xi}). \tag{34}$$

Note that the coefficients in the expansion of the kernel $F_h^*(\vec{x}, \vec{\xi})$ in (34) coincide with those in the expansion of the kernel $F^h(\vec{x}, \vec{\xi})$, and this allows us to use the available data instead of recalculating the coefficients of the new problem.

Substituting (33) and (34) into (32), we obtain an infinite system of linear algebraic equations for the determination of the eigenvalues and the eigenfunction expansion coefficients. This system has a symmetric matrix and can be written in the form

$$\sum_{n=N+1}^{\infty} F_{mn}^h \varphi_{n(k)}^{h*} = \mu_k^{h*} \varphi_{m(k)}^{h*}, \quad m = N + 1, N + 2, \dots \tag{35}$$

Now, let us construct a solution of equation (31). To this end, we represent the functions $q^*(\vec{x}, t)$ and $g_h^*(\vec{x}, t)$ in the form of series with eigenfunctions of the operator $\mathbf{P}_h^*\mathbf{F}^h$:

$$\begin{aligned} q^*(\vec{x}, t) &= \sum_{k=N+1}^{\infty} q_k^*(t)\varphi_k^{h*}(\vec{x}), \\ g_h^*(\vec{x}, t) &= \sum_{k=N+1}^{\infty} g_k^{h*}(t)\varphi_k^{h*}(\vec{x}), \quad g_k^{h*}(t) = \int_{-1}^1 g_h^*(\vec{\xi}, t)\varphi_k^{h*}(\vec{\xi}) d\Omega_{\xi}, \end{aligned} \tag{36}$$

and substitute these into (31). Then, taking into account (25)–(28) and (30)–(34), we obtain the following sequence of independent Volterra equations of the second kind:

$$\begin{aligned}
 (\mathbf{I} - \mathbf{V}_k^{h*})q_k^*(t) &= \delta_k^{h*}(t), \quad \mathbf{V}_k^{h*} = \frac{\sigma(t)\mathbf{V}_1 + \mu_k^{h*}\mathbf{V}_2}{\sigma(t) + \mu_k^{h*}}, \\
 \mathbf{V}_k^{h*} f(t) &= \int_{\tau_0}^t V_k^{h*}(t, \tau)f(\tau) d\tau, \quad V_k^{h*}(t, \tau) = \frac{\sigma(t)V_1(t, \tau) + \mu_k^{h*}V_2(t, \tau)}{\sigma(t) + \mu_k^{h*}}, \\
 \delta_k^{h*}(t) &= -\frac{1}{\sigma(t) + \mu_k^{h*}} \left[g_k^{h*}(t) + (\mathbf{I} - \mathbf{V}_2) \sum_{i=1}^N F_{k(i)}^h q_i^\circ(t) \right], \\
 F_{k(i)}^h &= \sum_{n=N+1}^\infty F_{in}^h \varphi_n^{h*}, \quad i = 1, \dots, N, \quad k = N + 1, N + 2, \dots
 \end{aligned} \tag{37}$$

Resolving (37) with respect to $q_k^*(t)$ by the methods of Chapter 11, we get

$$q_k^*(t) = (\mathbf{I} + \mathbf{R}_k^{h*}) \delta_k^{h*}(t), \quad (\mathbf{I} - \mathbf{V}_k^{h*})^{-1} = (\mathbf{I} + \mathbf{R}_k^{h*}), \quad \mathbf{R}_k^{h*} f(t) = \int_{\tau_0}^t R_k^{h*}(t, \tau)f(\tau) d\tau, \tag{38}$$

where $R_k^{h*}(t, \tau)$ is the resolvent of the kernel $V_k^{h*}(t, \tau)$.

We see that in view of (35)–(38) the function $q^*(\vec{x}, t)$ has been determined, and it is easy to find $q(\vec{x}, t)$, since $q^\circ(\vec{x}, t)$ is known by assumption (see (25)–(27)). Hence, taking into account the transformation of the variables (2), we finally obtain

$$y(\vec{x}, t) = \frac{1}{h(\vec{x})} \left[\sum_{n=1}^N \sum_{k=1}^n a_{nk} M_k(t) P_n^h(\vec{x}) + \sum_{k=N+1}^\infty q_k^*(t) \Phi_k^{h*}(\vec{x}) \right]. \tag{39}$$

The solution (39) depends on the function $h(x)$ in explicit manner, and this allows us to solve equation (48) with high accuracy by keeping a relatively small number of terms in the series even for a rapidly oscillating $h(\vec{x})$.

In practical calculations, the number of terms in the expansions has to be limited. For instance, taking the basis functions $p_k^h(\vec{x})$ with $k = N + 1, \dots, M$, we obtain the M th approximation of the desired solution. In this case, for the construction of eigenvalues and eigenfunctions of the Hilbert–Schmidt operator $\mathbf{P}_h^* \mathbf{F}^h$ one should find the eigenvalues and orthonormal eigenvectors of the matrix

$$[F_{MM}^h] = \begin{pmatrix} F_{N+1N+1}^h & F_{N+1N+2}^h & F_{35}^h & \cdots & F_{N+1M}^h \\ F_{N+1N+2}^h & F_{N+2N+2}^h & F_{N+2N+3}^h & \cdots & F_{N+2M}^h \\ F_{N+1N+3}^h & F_{N+2N+3}^h & F_{N+3N+3}^h & \cdots & F_{N+3M}^h \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ F_{N+1M}^h & F_{N+2M}^h & F_{N+3M}^h & \cdots & F_{MM}^h \end{pmatrix}. \tag{40}$$

The eigenvalues of the matrix (65) give approximations of the first $M - N$ eigenvalues of the Hilbert–Schmidt operator, and the components of its orthonormal eigenvectors approximate the coefficients in the expansion of the first $M - N$ eigenfunctions of this operator. Recall that the first N terms of the expansion (25) of $q^\circ(\vec{x}, t)$ of the function $q(\vec{x}, t)$ are known by assumption. Therefore, constructing the next $M - N$ terms of the expansion (36) of $q^*(\vec{x}, t)$, we obtain the M th approximation of the solution $q(\vec{x}, t)$ (see (25)).

It is important to observe that the matrix (40) can be obtained from the matrix (13) by deleting its first N rows and columns. This allows us to construct an expansion of the original kernel only once and then use these data for the examination of the new kernel arising in the problem with auxiliary conditions.

Now, in order to find the functions $\alpha_i(t)$ ($i = 1, \dots, N$), we apply the projection operator \mathbf{P}_h° to equation (31). As a result, we get

$$\alpha_k(t) = \sum_{i=k}^N a_{ik} \left\{ g_i^{h^\circ}(t) + \sigma(t)(\mathbf{I} - \mathbf{V}_1) \sum_{m=1}^i a_{im} M_m(t) + (\mathbf{I} - \mathbf{V}_2) \left[\sum_{j=1}^N F_{ji}^h \sum_{m=1}^j a_{jm} M_m(t) + \sum_{j=N+1}^\infty F_{j(i)}^h q_j^*(t) \right] \right\}. \tag{41}$$

Note that relations (41) form a system of N linear algebraic equations (with a triangular matrix) for the determination of the unknown quantities $\alpha_1(t), \dots, \alpha_N(t)$.

Thus, we have constructed a complete solution of the integral equation (18) with the auxiliary conditions (19).

17.4-3. General Projection Problem for Operator Equation.

Consider the equation

$$c(t)(\mathbf{I} - \mathbf{V}_1)y(t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{F}y(t) = f(t), \tag{42}$$

where $y(t)$ and $f(t)$ are continuous functions of t with values in an abstract Hilbert space H ; $c(t) > 0$ is a continuous scalar function of t ; \mathbf{I} is the identity operator; \mathbf{F} is a compact self-adjoint positive operator from H to H ; \mathbf{V}_1 and \mathbf{V}_2 are Volterra operators (with respect to t) such that the operators $(\mathbf{I} - \mathbf{V}_1)$, $(\mathbf{I} - \mathbf{V}_2)$, and $(\mathbf{I} - (\omega_1(t)\mathbf{V}_1 + \omega_2(t)\mathbf{V}_2))$ and their inverse operators preserve the class of continuous functions, provided that $\omega_1(t)$ and $\omega_2(t)$ are continuous in t .

Let us represent the Hilbert space H as a sum of its orthogonal subspaces $H = H^\circ \oplus H^*$. For continuous functions of t with values in H the following representations hold:

$$f(t) = f^\circ(t) + f^*(t), \quad y(t) = y^\circ(t) + y^*(t), \tag{43}$$

where $f(t)^\circ, y(t)^\circ$ are continuous functions of t with values in H° , and $f(t)^*, y(t)^*$ are continuous functions of t with values in H^* .

Consider the operator \mathbf{P}° of orthogonal projection from H onto H° . The operator $\mathbf{P}^* = \mathbf{I} - \mathbf{P}^\circ$ projects H onto H^* . Obviously,

$$\mathbf{P}^\circ f(t) = f(t)^\circ, \quad \mathbf{P}^* f(t) = f(t)^*, \quad \mathbf{P}^\circ y(t) = y(t)^\circ, \quad \mathbf{P}^* y(t) = y(t)^*. \tag{44}$$

General projection problem. Let $y(t)$ and $f(t)$ satisfy equation (42). For given $y^\circ(t)$ and $f^*(t)$, it is required to find the unknown $y^*(t)$ and $f^\circ(t)$.

Let us apply the operator \mathbf{P}^* to equation (42). As a result, we obtain a new equation which, after simple transformations, can be written in the form

$$c(t)(\mathbf{I} - \mathbf{V}_1)y^*(t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{P}^*\mathbf{F}y^*(t) = f^*(t) - (\mathbf{I} - \mathbf{V}_2)\mathbf{P}^*\mathbf{F}y^\circ(t). \tag{45}$$

THEOREM 1. *The operator $\mathbf{P}^*\mathbf{F}$ is compact, self-adjoint, and positive definite as an operator from H^* to H^* .*

Let φ_i be eigenfunctions of the operator $\mathbf{P}^*\mathbf{F}$ corresponding to its eigenvalues μ_i , i.e.,

$$\mathbf{P}^*\mathbf{F}\varphi_i = \mu_i\varphi_i. \tag{46}$$

All these eigenfunctions form a basis in H^* . Then, for continuous functions of t with values in H^* the following representations hold:

$$\begin{aligned} y^*(t) &= \sum_i a_i(t)\varphi_i, & f^*(t) &= \sum_i f_i(t)\varphi_i, \\ g^*(t) &= (\mathbf{I} - \mathbf{V}_2)\mathbf{P}^*\mathbf{F}y^*(t) = \sum_i g_i(t)\varphi_i, \end{aligned} \quad (47)$$

where $a_i(t)$, $f_i(t)$, $g_i(t)$ are continuous in t .

Substituting (47) into (45) and taking into account (46), we get

$$\begin{aligned} (\mathbf{I} - \mathbf{V}^i)a_i(t) &= \Phi_i(t), & \mathbf{V}^i &= \frac{c(t)\mathbf{V}_1 + \alpha_i\mathbf{V}_2}{c(t) + \alpha_i}, \\ \Phi_i(t) &= \frac{f_i(t) - g_i(t)}{c(t) + \alpha_i}, & a_i(t) &= (\mathbf{I} + \mathbf{R}^i)\Phi_i(t), \end{aligned} \quad (48)$$

where \mathbf{R}^i is the resolvent Volterra operator for \mathbf{V}^i .

Note that \mathbf{V}^i and $\Phi_i(t)$ are always defined since $c(t) > 0$, $\alpha_i > 0$, and $\alpha_i \rightarrow 0$, $\mathbf{V}^i \rightarrow \mathbf{V}_1$ as $i \rightarrow \infty$ (see Supplement 12.5-3). Due to the conditions imposed on the functions and the operators, the series (47) for $y^*(t)$ converges in H uniformly with respect to t , and its sum is a continuous function of t with values in H^* . Equation (45) is linear and for $f^*(t) = 0$, $y^\circ(t) = 0$ has the trivial solutions.

THEOREM 2. *In the above classes of continuous functions, equation (45) has one and only one solution.*

Thus, we have found $y^*(t)$. In order to find $f^\circ(t)$, let us apply the operator \mathbf{P}° to equation (42). Then

$$f^\circ(t) = c(t)(\mathbf{I} - \mathbf{V}_1)y^\circ(t) + (\mathbf{I} - \mathbf{V}_2)\mathbf{P}^\circ\mathbf{F}(y^\circ(t) + y^*(t)), \quad (49)$$

which immediately yields an expression for $f^\circ(t)$, since $y^\circ(t)$ is given and $y^*(t)$ has been found. The question about the existence and the uniqueness of the solution $f^\circ(t)$ is solved simultaneously with that of the existence and the uniqueness of the solution $y^*(t)$.

For the justification of this method the following theorem is needed.

THEOREM 3. *Functions $y(t)$ and $f(t)$ satisfy equation (42) for given projections $\mathbf{P}^\circ y(t)$ and $\mathbf{P}^* f(t)$ if and only if relations (45) and (49) hold.*

THEOREM 4. *A solution of equation (42) for given $y^\circ(t)$ and $f^*(t)$ exists and is unique if and only if equation (45) has one and only one solution.*

Remark 1. For $\mathbf{P}^* = \mathbf{I}$, the above projection problem reduces to the classical problem for an equation with a given right-hand side. Thus, the problem considered here is a generalization of the classical approach to more complex cases of equations with auxiliary conditions.

Remark 2. The projection method considered here can be regarded as an extension of the Hilbert–Schmidt method to multidimensional equations with auxiliary conditions.

Remark 3. For given auxiliary conditions, the basic operator of the problem is $\mathbf{P}^*\mathbf{F}$ and not \mathbf{F} , which makes the problem considered here essentially different from the problem with a given right-hand side.

Remark 4. Eigenfunctions and eigenvalues of the kernels and operators can be found by various methods described in literature, and not only those represented in Chapter 17.

Chapter 18

Application of Integral Equations for the Investigation of Differential Equations

► **Preliminary remarks.** Integral equations play an important role in the theory of ordinary and partial differential equations and boundary value problems. The reduction of boundary value problems to integral equations allows for the application of iteration and finite-difference methods of solving integral equations. These methods are, as a rule, substantially simpler than those used for solving differential equations. Moreover, many delicate proofs and qualitative results of the theory of differential equations have been obtained by the investigation of the corresponding integral equations.

18.1. Reduction of the Cauchy Problem for ODEs to Integral Equations

18.1-1. Cauchy Problem for First-Order ODEs. Uniqueness and Existence Theorems.

The *Cauchy problem*: find a solution of the equation

$$y'_x = f(x, y) \tag{1}$$

that satisfies the *initial condition*

$$y(x_0) = y_0 \tag{2}$$

for given y_0 and x_0 .

Geometrical meaning of the Cauchy problem: find an integral curve of equation (1) passing through the point (x_0, y_0) .

THEOREM (EXISTENCE, PEANO). *Let the function $f(x, y)$ be continuous in an open domain D of the xy -plane. Then there is at least one integral curve of equation (1) that passes through each point $(x_0, y_0) \in D$; each of these curves can be extended at both ends up to the boundary of any closed domain $D_0 \subset D$ such that (x_0, y_0) belongs to the interior of D_0 .*

THEOREM (UNIQUENESS). *Let the function $f(x, y)$ be continuous in an open domain D and have a bounded partial derivative in D with respect to y (or satisfy the Lipschitz condition: $|f(x, y) - f(x, z)| \leq M|y - z|$, where $M > 0$ is a constant). Then there is a unique solution of equation (1) satisfying condition (2).*

18.1-2. Cauchy Problem for First-Order ODEs. Method of Successive Approximations.

The method of successive approximations (the Picard method) consists of two stages. On the first stage, the Cauchy problem (1)–(2) is reduced to the equivalent integral equation:

$$y(x) = y_0 + \int_{x_0}^x f(t, y(t)) dt. \tag{3}$$

Then a solution of equation (3) is sought using the formula of successive approximations:

$$y_{n+1}(x) = y_0 + \int_{x_0}^x f(t, y_n(t)) dt; \quad n = 0, 1, 2, \dots$$

The initial approximation $y_0(x)$ can be chosen arbitrarily; the simplest way is to take y_0 a constant. The iterative process converges as $n \rightarrow \infty$, provided the assumptions of the theorems in Subsection 18.1-1 are satisfied.

18.1-3. Cauchy Problem for Second-Order ODEs. Method of Successive Approximations.

The method of successive approximations is implemented in two steps. First, the Cauchy problem

$$\begin{aligned} y''_{xx} &= f(x, y, y'_x) && \text{(equation),} \\ y(x_0) &= y_0, \quad y'_x(x_0) = y'_0 && \text{(initial conditions)} \end{aligned}$$

is reduced to an equivalent system of integral equations by the introduction of the new variable $u(x) = y'_x$. These integral equations have the form

$$u(x) = y'_0 + \int_{x_0}^x f(t, y(t), u(t)) dt, \quad y(x) = y_0 + \int_{x_0}^x u(t) dt. \tag{4}$$

Then the solution of system (4) is sought by means of successive approximations defined by the following recurrence formulas:

$$u_{n+1}(x) = y'_0 + \int_{x_0}^x f(t, y_n(t), u_n(t)) dt, \quad y_{n+1}(x) = y_0 + \int_{x_0}^x u_n(t) dt; \quad n = 0, 1, 2, \dots$$

As the initial approximation, one can take $y_0(x) = y_0$ and $u_0(x) = y'_0$. The iterative process converges as $n \rightarrow \infty$, under assumptions similar to those formulated in the theorems of Subsection 18.1-1.

Remark. In a similar way, the Cauchy problem for an n th order ODE can be reduced to a system of integral equations.

18.1-4. Cauchy Problem for a Special n -Order Linear ODE.

Consider the Cauchy problem for the following linear n th order ODE:

$$y_x^{(n)} + f_{n-1}(x)y_x^{(n-1)} + \dots + f_1(x)y'_x + f_0(x)y = g(x) \tag{5}$$

with the homogeneous initial conditions at the point $x = a$:

$$y(a) = y'_x(a) = \dots = y_x^{(n-1)}(a) = 0. \tag{6}$$

Introducing a new unknown function by

$$y(x) = \frac{1}{(n-1)!} \int_a^x (x-t)^{n-1} u(t) dt \tag{7}$$

and differentiating (7) n times, we get

$$\begin{aligned} y_x^{(k)}(x) &= \frac{1}{(n-k-1)!} \int_a^x (x-t)^{n-k-1} u(t) dt, \quad k = 1, \dots, n-1; \\ y_x^{(n)}(x) &= u(x). \end{aligned} \quad (8)$$

Obviously, the function (7) satisfies the initial conditions (6). Substituting (8) into the left-hand side of equation (5), we obtain

$$u(x) + \int_a^x K(x,t)u(t) dt = g(x), \quad (9)$$

where

$$K(x,t) = f_{n-1}(x) + f_{n-2}(x) \frac{x-t}{1!} + \dots + f_0(x) \frac{(x-t)^{n-1}}{(n-1)!}. \quad (10)$$

Thus, the Cauchy problem (5)–(6) has been reduced to the integral equation (9)–(10), which is a Volterra equation of the second kind. Finding the function $u(x)$ from (9) and using formula (7) we obtain the desired solution $y(x)$.

Remark. The Cauchy problem for equation (5) with nonhomogeneous boundary conditions

$$y(a) = b_0, \quad y'_x(a) = b_1, \quad \dots, \quad y_x^{(n-1)}(a) = b_{n-1}$$

can be reduced to a Cauchy problem with homogeneous boundary conditions for another function $w(x)$ with the help of the substitution

$$y(x) = w(x) + \sum_{k=1}^{n-1} b_k \frac{(x-a)^k}{k!}.$$

© References for Section 18.1: W. V. Lovitt (1950), E. Kamke (1977), R. P. Kanwal (1996), A. D. Polyanin and A. V. Manzhirov (2007).

18.2. Reduction of Boundary Value Problems for ODEs to Volterra Integral Equations. Calculation of Eigenvalues

18.2-1. Reduction of Differential Equations to Volterra Integral Equations.

1°. Consider a linear nonhomogeneous ODE for the function $y = y(x)$:

$$L_n[y] = h(x) \quad (a < x < b), \quad (1)$$

where

$$L_n[y] = \sum_{k=0}^n f_k(x)y_x^{(k)}, \quad f_n(x) \neq 0. \quad (2)$$

Let $\varphi_1(x), \dots, \varphi_n(x)$ be a fundamental system of solutions of the truncated homogeneous equation

$$L_n[\varphi] = 0. \quad (3)$$

Denote by $W(x)$ its Wronskian determinant

$$W(x) = \begin{vmatrix} \varphi_1(x) & \cdots & \varphi_n(x) \\ \varphi_1'(x) & \cdots & \varphi_n'(x) \\ \vdots & \ddots & \vdots \\ \varphi_1^{(n-2)}(x) & \cdots & \varphi_n^{(n-2)}(x) \\ \varphi_1^{(n-1)}(x) & \cdots & \varphi_n^{(n-1)}(x) \end{vmatrix},$$

and by $W_\nu(x)$ the determinant obtained from $W(x)$ by replacing the ν th column by $0, \dots, 0, h(x)$. The general solution of equation (1) can be written in the form

$$y(x) = \sum_{\nu=1}^n \varphi_\nu(x) \int_a^x \frac{W_\nu(\xi)}{f_n(\xi)W(\xi)} d\xi + \sum_{\nu=1}^n C_\nu \varphi_\nu(x), \tag{4}$$

where the first sum is a particular solution of equation (1), the second sum is the general solution of the homogeneous equation (3), and C_ν are arbitrary constants.

For boundary value problems, the constants C_ν are found from the corresponding boundary conditions, and for the Cauchy problem, C_ν are obtained from the initial conditions.

2°. Consider the linear ODE for the function $y = y(x)$ with a parameter λ :

$$L_n[y] = h(x) - \lambda g(x)y \quad (a < x < b), \tag{5}$$

where L_n is the differential operator (2). Equation (5) differs from (1) only by an additional term in the right-hand side. Therefore, replacing the function $h(x)$ by $h(x) - \lambda g(x)y(x)$ in the solution (4) and performing simple transformations, we come to the Volterra integral equation

$$y(x) + \lambda \int_a^x K(x, \xi)y(\xi) d\xi = F(x), \tag{6}$$

where

$$K(x, \xi) = \frac{g(\xi)}{f_n(\xi)W(\xi)} D(x, \xi), \quad F(x) = \int_a^x \frac{h(\xi)}{f_n(\xi)W(\xi)} D(x, \xi) d\xi + \sum_{\nu=1}^n C_\nu \varphi_\nu(x), \tag{7}$$

and

$$D(x, \xi) = \begin{vmatrix} \varphi_1(\xi) & \cdots & \varphi_n(\xi) \\ \varphi_1'(\xi) & \cdots & \varphi_n'(\xi) \\ \vdots & \ddots & \vdots \\ \varphi_1^{(n-2)}(\xi) & \cdots & \varphi_n^{(n-2)}(\xi) \\ \varphi_1(x) & \cdots & \varphi_n(x) \end{vmatrix}. \tag{8}$$

The Volterra integral equation (6)–(8) is equivalent to the differential equation (5).

3°. In a similar way, one can approach nonlinear ODEs of the form

$$L_n[y] = h(x, y) \quad (a < x < b) \tag{9}$$

with the same differential operator (2). This equation can be reduced to the nonlinear Volterra integral equation

$$y(x) = \int_a^x \frac{h(\xi, y(\xi))}{f_n(\xi)W(\xi)} D(x, \xi) d\xi + \sum_{\nu=1}^n C_\nu \varphi_\nu(x), \tag{10}$$

where the function $D(x, \xi)$ is defined by (8).

Note that both sides of equation (9) may depend on the spectral parameter λ . The linear equation corresponds to the right-hand side $h(x, y) = h_1(x)y + h_0(x)$.

18.2-2. Application of Volterra Equations to the Calculation of Eigenvalues.

1°. The Volterra integral equation (6) can be used for the calculation of the smallest eigenvalue and the corresponding eigenfunction of various boundary value problems for the ODE (5). For this purpose, one utilizes the method of successive approximations: the first term $y(x)$ of the integral equation is replaced by $y_n(x)$, and $y(\xi)$ in the integrand is replaced by $y_{n-1}(\xi)$. On each step, the parameter λ is chosen such that the function $y_n(x)$ would satisfy the boundary conditions. This procedure can be illustrated by the following example.

Example 1. Consider the equation

$$y''_{xx} + \lambda g(x)y = 0 \quad (0 < x < 1) \quad (11)$$

with the homogeneous boundary conditions of the first kind

$$y(0) = y(1) = 0. \quad (12)$$

Equation (11) is a special case of (5) for $n = 2$, $L_2[y] = y''_{xx}$, $h(x) \equiv 0$, $a = 0$, $b = 0$.

The fundamental system of solutions of the truncated equation $L_2[\varphi] = 0$ has the form

$$\varphi_1(x) = 1, \quad \varphi_2(x) = x. \quad (13)$$

Simple transformations with the help of (8) yield

$$\begin{aligned} W(x) &= \varphi_1(x)[\varphi_2(x)]'_x - \varphi_2(x)[\varphi_1(x)]'_x = 1, \\ D(x, \xi) &= \varphi_1(\xi)\varphi_2(x) - \varphi_1(x)\varphi_2(\xi) = x - \xi. \end{aligned} \quad (14)$$

Substituting (13)–(14) into (6)–(7), we come to the Volterra equation

$$y(x) = C_1 + C_2x - \lambda \int_0^x (x - \xi)g(\xi)y(\xi) d\xi. \quad (15)$$

From the first boundary condition in (12), we get $C_1 = 0$. Since eigenfunctions are defined to within a constant coefficient, we can take $C_2 = 1$ in (15). As a result we get

$$y(x) = x - \lambda \int_0^x (x - \xi)g(\xi)y(\xi) d\xi. \quad (16)$$

This equation can be solved by the method of successive approximations based on the formula

$$y_n(x) = x - \lambda \int_0^x (x - \xi)g(\xi)y_{n-1}(\xi) d\xi, \quad n = 1, 2, \dots \quad (17)$$

Next, consider more closely the simplest case $g(x) = 1$. As the zero approximation, we take $y_0 = 1$ and find that

$$y_1(x) = x - \lambda \int_0^x (x - \xi) d\xi = x - \frac{1}{2}\lambda x^2.$$

From the second boundary condition in (12), we get $y_1(1) = 0$, and therefore, $\lambda = \lambda_1 = 2$. It follows that

$$y_1(x) = x - x^2.$$

Let us insert this function into the right-hand side of (17), where $g(x) = 1$. We have

$$y_2(x) = x - \lambda \int_0^x (x - \xi)(\xi - \xi^2) d\xi = x - \lambda \left(\frac{1}{6}x^3 - \frac{1}{12}x^4 \right).$$

Satisfying the second boundary condition in (12), i.e., $y_2(1) = 0$, we obtain

$$\lambda_2 = 12, \quad y_2(x) = x - 2x^3 + x^4.$$

In a similar way, we find that

$$\lambda_3 = 10, \quad y_2(x) = x - \frac{5}{3}x^3 + 3x^5 - \frac{1}{3}x^6. \quad (18)$$

The exact smallest eigenvalue for $g(x) = 1$ is equal to $\lambda = \pi^2 \approx 9.87$, and the corresponding eigenfunction has the form

$$y(x) = \frac{1}{\pi} \sin(\pi x) \approx x - 1.6x^3 + 0.8x^5.$$

In the case under consideration, the choice of the initial approximation, $y_0(x) = 1$, was not quite good, since both boundary conditions in (12) do not hold for this function. The convergence rate may be increased by taking the initial approximation of the form $y_0(x) = x - x^2$, in which case both boundary conditions in (12) are satisfied.

For an arbitrary continuous function $g(x)$, in the absence of information about eigenfunctions, it is convenient to take the initial approximation in (17) of the form

$$y_0(x) = x - x^2 \quad \text{or} \quad y_0(x) = \frac{1}{\pi} \sin(\pi x),$$

since these functions satisfy the boundary conditions (12). In the special case of $g(x) = 1$, the first initial function ensures fast convergence of the expression (17) to the exact result, and the second yields the exact result immediately.

2°. The Volterra integral equation (6) can be used for obtaining asymptotic expansions of eigenvalues and eigenfunctions of the corresponding boundary value problems for the ODE (5) for large λ . Note that in this situation, various cases are possible; for instance, the operator $L_n[y]$ may involve the spectral parameter λ , while the right-hand side of the differential equation is independent on λ .

Example 2. Consider the equation

$$y''_{xx} + [f(x) + \lambda^2]y = 0 \quad (a < x < b) \tag{19}$$

with the homogeneous boundary condition of the first kind

$$y(a) = y(b) = 0. \tag{20}$$

The function $f(x)$ is assumed continuous on the finite segment $[a, b]$.

Let us write equation (19), using the notation from (9). As the differential operator and the right-hand side of the equation we take

$$\begin{aligned} L_2[y] &= y''_{xx} + \lambda^2 y, \\ h(x, y) &= -f(x)y(x). \end{aligned} \tag{21}$$

The fundamental system of solutions of the truncated equation $L_2[\varphi] = 0$ has the form

$$\varphi_1 = \cos(\lambda x), \quad \varphi_2 = \sin(\lambda x). \tag{22}$$

After elementary calculations with the help of (8), we get

$$\begin{aligned} W(x) &= \varphi_1(x)[\varphi_2(x)]'_x - \varphi_2(x)[\varphi_1(x)]'_x = \lambda, \\ D(x, \xi) &= \varphi_1(\xi)\varphi_2(x) - \varphi_1(x)\varphi_2(\xi) = \sin[\lambda(x - \xi)]. \end{aligned} \tag{23}$$

Substituting the second expression from (21), as well as (22) and (23), into (10), we come to the Volterra integral equation

$$y(x) = -\frac{1}{\lambda} \int_a^x \sin[\lambda(x - \xi)]f(\xi)y(\xi) d\xi + C_1 \cos(\lambda x) + C_2 \sin(\lambda x).$$

The first boundary condition in (20) yields $C_1 \cos(\lambda a) + C_2 \sin(\lambda a) = 0$. Therefore,

$$C_1 \cos(\lambda x) + C_2 \sin(\lambda x) = C \sin[\lambda(x - a)].$$

Since eigenfunctions are defined to within an arbitrary constant coefficient, we come to the integral equation

$$y(x) = \sin[\lambda(x - a)] - \frac{1}{\lambda} \int_a^x \sin[\lambda(x - \xi)]f(\xi)y(\xi) d\xi. \tag{24}$$

It is easy to see that the functions $y(x)$ are uniformly bounded for sufficiently small $\lambda > 0$. From the second boundary condition in (20), using (24), we find that

$$\sin[\lambda(b - a)] = \frac{1}{\lambda} \int_a^b \sin[\lambda(x - \xi)]f(\xi)y(\xi) d\xi = O\left(\frac{1}{\lambda}\right). \tag{25}$$

Hence, we obtain the following asymptotic formula for the eigenvalues $\lambda = \lambda_n$:

$$\lambda_n = \frac{\pi n}{b - a} + O\left(\frac{1}{\lambda}\right) = \frac{\pi n}{b - a} + O\left(\frac{1}{n}\right), \tag{26}$$

where n is a large positive integer. The corresponding eigenfunctions are obtained by substituting the values (26) into (24):

$$y_n(x) = \sin[\lambda_n(x - a)] + O\left(\frac{1}{\lambda}\right) = \sin[\lambda_n(x - a)] + O\left(\frac{1}{n}\right).$$

Inserting this function into (25), one can refine the asymptotic formula (26), etc.

A similar approach can be taken with regard to other boundary conditions for equation (19).

⊙ References for Section 18.2: E. Kamke (1977), A. D. Polyanin and V. F. Zaitsev (2003).

18.3. Reduction of Boundary Value Problems for ODEs to Fredholm Integral Equations with the Help of the Green's Function

18.3-1. Linear Ordinary Differential Equations. Fundamental Solutions.

Consider a homogeneous linear ordinary differential equation

$$L[y] \equiv \sum_{k=0}^n f_k(x)y_x^{(k)} = 0 \quad (a < x < b), \tag{1}$$

where $f_k(x)$ are continuous functions on the segment $a \leq x \leq b$ and $f_n(x) \neq 0$.

A *fundamental solution* of the differential equation (1) is a function of two variables $g(x, \xi)$ defined on the square $a \leq x, \xi \leq b$ and having the following properties:

- (a) in each of the triangles $a \leq x \leq \xi \leq b$ and $a \leq \xi \leq x \leq b$, the function $g(x, \xi)$ has partial derivatives in x of the orders $\leq n$, and these derivatives are continuous in x and ξ in each triangle;
- (b) $g(x, \xi)$, as a function of x , satisfies equation (1) in each of the triangles;
- (c) on the entire square $a \leq x, \xi \leq b$, the function $g(x, \xi)$ is continuous and has partial derivative in x up to the order $(n - 2)$, and these derivatives are continuous in x and ξ on that square;
- (d) for $a < \xi < b$, the following relation holds:

$$\frac{\partial^{n-1}g}{\partial x^{n-1}} \Big|_{x=\xi+0} - \frac{\partial^{n-1}g}{\partial x^{n-1}} \Big|_{x=\xi-0} = \frac{1}{f_n(\xi)}. \tag{2}$$

Fundamental solutions exist always. For instance, one can take

$$g(x, \xi) = \frac{\text{sign}(x - \xi)}{2f_n(\xi)W(\xi)} \begin{vmatrix} y_1(\xi) & \cdots & y_n(\xi) \\ y_1'(\xi) & \cdots & y_n'(\xi) \\ \vdots & \ddots & \vdots \\ y_1^{(n-2)}(\xi) & \cdots & y_n^{(n-2)}(\xi) \\ y_1(x) & \cdots & y_n(x) \end{vmatrix}, \quad W(x) = \begin{vmatrix} y_1(x) & \cdots & y_n(x) \\ y_1'(x) & \cdots & y_n'(x) \\ \vdots & \ddots & \vdots \\ y_1^{(n-2)}(x) & \cdots & y_n^{(n-2)}(x) \\ y_1^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}, \tag{3}$$

where $y_1(x), \dots, y_n(x)$ is a fundamental system of solutions of equation (1), and $W(x)$ is its Wronskian determinant. This special fundamental solution has the following property:

$$g(\xi, \xi) = g'_\xi(\xi, \xi) = \cdots = g_x^{(n-2)}(\xi, \xi) = 0.$$

The set of all fundamental solutions can be described by the sum

$$g(x, \xi) + C_1(\xi)y_1(x) + \cdots + C_n(\xi)y_n(x),$$

where $C_k(\xi)$ are arbitrary continuous functions.

Fundamental solutions play an important role in the theory of linear differential equations, since the function

$$y(x) = \int_a^b g(x, \xi)\varphi(\xi) d\xi$$

is a particular solution of the nonhomogeneous linear ODE

$$L[y] = \varphi(x).$$

18.3-2. Boundary Value Problems for n th Order Differential Equations. Green's Function.

Consider a homogeneous linear boundary value problem for equation (1) with the boundary condition

$$\Gamma_m[y] = 0, \quad m = 1, \dots, n. \tag{4}$$

Assume that the functions $f_k(x)$ are continuous on the segment $[a, b]$ and the left-hand sides of the boundary conditions have the form

$$\Gamma_m[y] = \Gamma_{m,a}[y] + \Gamma_{m,b}[y], \tag{5}$$

where $\Gamma_{m,a}[y]$ and $\Gamma_{m,b}[y]$ are linear differential forms of an order $\leq n-1$ calculated at the endpoints $x = a$ and $x = b$.

A function $G(x, \xi)$ defined on the square $a \leq x, \xi \leq b$ is called the *Green's function* or the *influence function* for problem (1), (4) if it is a fundamental solution of equation (1) and for any fixed ξ ($a < \xi < b$) satisfies boundary conditions (4) as a function of x .

If the boundary value problem (1), (4) admits only the trivial solution $y \equiv 0$, then there is only one Green's function for this problem. Knowing a fundamental system of solutions $y_1(x), \dots, y_n(x)$ of equation (1), one can construct the Green's function as follows. For each ξ ($a \leq \xi \leq b$), we find a solution $c_1 = c_1(\xi), \dots, c_n = c_n(\xi)$ of the system of linear algebraic equations

$$\begin{aligned} \sum_{\nu=1}^n c_\nu y_\nu^{(\sigma)}(\xi) &= 0, \quad \sigma = 0, \dots, n-2, \\ \sum_{\nu=1}^n c_\nu y_\nu^{(n-1)}(\xi) &= \frac{1}{f_n(\xi)}, \end{aligned}$$

and then a solution $b_1 = b_1(\xi), \dots, b_n = b_n(\xi)$ of another system

$$\sum_{\nu=1}^n b_\nu \Gamma_m[y_\nu] = \sum_{\nu=1}^n c_\nu \Gamma_{m,a}[y_\nu], \quad m = 1, \dots, n.$$

The Green's function can be defined by the formula

$$G(x, \xi) = \begin{cases} \sum_{\nu=1}^n a_\nu(\xi) y_\nu(x) & \text{for } a \leq x \leq \xi \leq b, \\ \sum_{\nu=1}^n b_\nu(\xi) y_\nu(x) & \text{for } a \leq \xi \leq x \leq b, \end{cases} \tag{6}$$

where $a_\nu(\xi) = b_\nu(\xi) - c_\nu(\xi)$.

The Green's function can be expressed in terms of a fundamental system of solutions $y_1(x), \dots, y_n(x)$ of equation (1), the fundamental solution $g(x, \xi)$, and the differential forms (4):

$$G(x, \xi) = \frac{Z(x, \xi)}{\Delta}, \quad Z(x, \xi) = \begin{vmatrix} g(x, \xi) & y_1(x) & \cdots & y_n(x) \\ \Gamma_1[g] & \Gamma_1[y_1] & \cdots & \Gamma_1[y_n] \\ \vdots & \vdots & \ddots & \vdots \\ \Gamma_n[g] & \Gamma_n[y_1] & \cdots & \Gamma_n[y_n] \end{vmatrix}, \tag{7}$$

where Δ stands for the determinant

$$\Delta = \det[\Gamma_i[y_j]], \quad i, j = 1, \dots, n.$$

Remark. For $\xi = a$ and $\xi = b$, the representation (7), is not valid, in general. However, in this case the following limit relations hold:

$$G(x, a) = \lim_{\xi \rightarrow a} G(x, \xi), \quad G(x, b) = \lim_{\xi \rightarrow b} G(x, \xi).$$

The Green's function plays an important role in the theory of linear boundary value problems, since the function

$$y(x) = \int_a^b G(x, \xi) \varphi(\xi) d\xi$$

is a solution of the linear nonhomogeneous differential equation $L[y] = \varphi(x)$ with the homogeneous boundary conditions (4).

A nonlinear boundary value problem for a nonlinear ODE of the form

$$\sum_{k=0}^n f_k(x) y_x^{(k)} = \Phi(x, y) \quad (a < x < b), \quad (8)$$

with the boundary conditions (4) can be reduced, with the help of the Green's function, to the nonlinear integral equation

$$y(x) = \int_a^b G(x, \xi) \Phi(\xi, y(\xi)) d\xi, \quad (9)$$

whose investigation is, as a rule, much simpler than that of the original boundary value problem (8), (4).

Remark. In applications, one often has to deal with linear eigenvalue problems in which equation (8) is considered with

$$\Phi(x, y) = \lambda p(x)y,$$

where λ is a spectral parameter.

18.3-3. Boundary Value Problems for Second-Order Differential Equations. Green's Function.

The Green's function for the boundary value problem for the linear second-order equation

$$f_2(x)y''_{xx} + f_1(x)y'_{xx} + f_0(x)y = 0 \quad (10)$$

with the homogeneous boundary conditions

$$\begin{aligned} k_1 y'_x + s_1 y &= 0 & \text{at } x = a, \\ k_2 y'_x + s_2 y &= 0 & \text{at } x = b, \end{aligned} \quad (11)$$

can be written as

$$G(x, \xi) = \begin{cases} \frac{y_1(x)y_2(\xi)}{f_2(\xi)W(\xi)} & \text{if } a \leq x \leq \xi \leq b, \\ \frac{y_1(\xi)y_2(x)}{f_2(\xi)W(\xi)} & \text{if } a \leq \xi \leq x \leq b, \end{cases} \quad (12)$$

where $y_1(x)$ is any nontrivial solution of equation (10) satisfying the first boundary condition in (11), and $y_2(x)$ is any nontrivial solution of equation (10) satisfying the second boundary condition in (11); $W(x) = y_1(x)y'_2(x) - y'_1(x)y_2(x)$ is the Wronskian determinant.

The Green's function (12) can be used for constructing solutions of nonhomogeneous linear or nonlinear boundary value problems for second-order ODEs.

Example. Consider the boundary value problem for the nonlinear second-order equation

$$y''_{xx} = \Phi(x, y(x)) \tag{13}$$

with the homogeneous boundary conditions of the first kind

$$y(0) = 0, \quad y(1) = 0. \tag{14}$$

Let us construct the Green's function for the linear equation

$$y''_{xx} = 0 \tag{15}$$

with boundary conditions (14). Taking into account that the general solution of equation (15) has the form $y = C_1 + C_2x$, we take in (12) the solutions $y_1(x) = x$, $y_2(x) = 1 - x$ [each of these satisfies one of the boundary conditions from (14)] and $f_2(x) = 1$, $W(x) = -1$, $a = 0$, $b = 1$. As a result, we get

$$G(x, \xi) = \begin{cases} (\xi - 1)x & \text{if } 0 \leq x \leq \xi \leq 1, \\ (x - 1)\xi & \text{if } 0 \leq \xi \leq x \leq 1. \end{cases}$$

Regarding the right-hand side of equation (13) as known, we obtain

$$y(x) = \int_0^1 G(x, \xi)\Phi(\xi, y(\xi)) d\xi.$$

Thus, solving the boundary value problem (13)–(14) amounts to solving a nonlinear integral equation of Hammerstein type with the kernel being the Green's function for problem (15), (14).

Table 13 contains simplest examples of Green's functions $G(x, \xi)$ for some linear boundary value problems for ODEs. In all these examples, $G(x, \xi) = G(\xi, x)$, and therefore the Green's function is specified only in the domain $x \leq \xi$. For equations with the operator $L[y] = -[f(x)y'_x]'_x$, it is assumed that $f(x) > 0$ and $q(x) = \int_0^x \frac{dt}{f(t)}$.

18.3-4. Nonlinear Problem of Nonisothermal Flow in Plane Channel.

It is known that the dynamic viscosity of a fluid μ essentially depends on temperature T (μ decreases as T increases) and the other physical parameters of the fluid have small variation. For high-viscosity fluids (like glycerol, liquid oil, or petroleum) it is common to assume the exponential dependence

$$\mu = \mu_0 \exp[-\beta(T - T_0)], \tag{16}$$

where μ_0 , β , and T_0 are empirical constants.

Stationary nonisothermal flows of viscous incompressible fluid are described by the following system of equations:

$$\sum_{j=1}^3 \left(u_j \frac{\partial u_i}{\partial X_j} - \frac{1}{\rho} \frac{\partial p_{ij}}{\partial X_j} \right) = 0, \quad p_{ij} = -p\delta_{ij} + \mu \left(\frac{\partial u_i}{\partial X_j} + \frac{\partial u_j}{\partial X_i} \right), \quad i = 1, 2, 3; \tag{17}$$

$$\frac{\partial u_1}{\partial X_1} + \frac{\partial u_2}{\partial X_2} + \frac{\partial u_3}{\partial X_3} = 0, \tag{18}$$

$$u_1 \frac{\partial T}{\partial X_1} + u_2 \frac{\partial T}{\partial X_2} + u_3 \frac{\partial T}{\partial X_3} = \sigma \left(\frac{\partial^2 T}{\partial X_1^2} + \frac{\partial^2 T}{\partial X_2^2} + \frac{\partial^2 T}{\partial X_3^2} \right). \tag{19}$$

Here u_j are fluid velocity components, X_j are Cartesian coordinates, ρ is density, p is pressure, σ is the heat transfer coefficient, and $\delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases}$

Stationary rectilinear flows in a plane channel correspond to solutions of the form

$$u_1 = u_2 = 0, \quad u_3 = u(X), \quad p = p(X, Z), \quad T = T(X, Z), \tag{20}$$

where $Z = X_3$ is the longitudinal coordinate in the channel and $X = X_1$ is the transverse coordinate.

TABLE 13
Green's function for some boundary value problems for linear ODEs $L[y] = 0$

Differential operator, $L[y]$	Boundary conditions	Green's function, $G(x, \xi)$
$-y''_{xx}$	$y(0) = y(a) = 0$	$x\left(1 - \frac{\xi}{a}\right)$
$-y''_{xx}$	$y(0) = y'_x(a) = 0$	ξ
$-y''_{xx}$	$y'_x(0) = y(a) = 0$	$a - \xi$
$-y''_{xx} - k^2y$	$y(0) = y(1) = 0$	$\frac{\sin(kx)\sin[k(1-\xi)]}{k\sin k}$
$-y''_{xx} + k^2y$	$y(0) = y(1) = 0$	$\frac{\sinh(kx)\sinh[k(1-\xi)]}{k\sinh k}$
$-xy''_{xx} - y'_x$	$y(0) \neq \infty, y(a) = 0$	$-\ln \frac{\xi}{a}$
$-(xy'_x)'_x + \frac{n^2}{x}y$	$y(0) \neq \infty, y(a) = 0$	$\frac{1}{2n}\left(\frac{x}{a}\right)^n - \frac{(x\xi)^n}{2na^{2n}} \quad (n = 1, 2, \dots)$
$-[f(x)y'_x]'_x$	$y(0) = y(a) = 0$	$q(x) - \frac{q(x)q(\xi)}{q(a)}$
$-[f(x)y'_x]'_x$	$y(0) = y'_x(a) = 0$	$q(x)$
$-[f(x)y'_x]'_x$	$y(0) = 0, ky'_x(a) + y(a) = 0$	$q(x) - \frac{f(a)q(x)q(\xi)}{f(a)q(a) + k} \quad (k > 0)$
y''''_{xxxx}	$y(0) = y'_x(0) = 0,$ $y(1) = y'_x(1) = 0$	$\left(\frac{1}{2}\xi - \xi^2 + \frac{1}{2}\xi^3\right)x^2 - \left(\frac{1}{6}\xi - \frac{1}{2}\xi^2 + \frac{1}{3}\xi^3\right)x^3$
y''''_{xxxx}	$y(0) = y'_x(0) = 0,$ $y''_{xx}(1) = y''''_{xxxx}(1) = 0$	$\frac{1}{6}x^2(3\xi - x)$

Substituting expressions (20) into equations (17)–(19) and letting $u'_X = du/dX$, we obtain the following three equations:

$$\frac{\partial p}{\partial X} = u'_X \frac{\partial \mu}{\partial Z}, \quad \frac{\partial p}{\partial Z} = \frac{\partial}{\partial X}(\mu u'_X), \quad (21)$$

$$u(X) \frac{\partial T}{\partial Z} = \sigma \left(\frac{\partial^2 T}{\partial X^2} + \frac{\partial^2 T}{\partial Z^2} \right). \quad (22)$$

Using differentiation, we eliminate the pressure p from (21) and obtain

$$\left(\frac{\partial^2}{\partial Z^2} - \frac{\partial^2}{\partial X^2} \right) (\mu u'_X) = 0. \quad (23)$$

The general solution of equation (23) can be written in the form

$$\mu u'_X = \Phi(Z + X) + \Psi(Z - X), \quad (24)$$

where Φ and Ψ are arbitrary functions.

Let $X = 0$ correspond to the middle line of the plane channel of width $2h$, i.e., $-h \leq X \leq h$. On the walls we have the conditions of adhesion

$$u = 0 \quad \text{for} \quad X = \pm h. \tag{25}$$

Moreover, assume that the temperature has linear variation on the channel walls,

$$T = T_0 - EZ \quad \text{for} \quad X = \pm h. \tag{26}$$

Instead of the domain $-h \leq X \leq h$, we can consider its half $0 \leq X \leq h$ with the symmetry condition on the middle line:

$$u'_X = T_X = 0 \quad \text{for} \quad X = 0. \tag{27}$$

Equations (22) and (24) (with the viscosity defined by (16)) and the boundary conditions (25), (26) (for $x = h$), and (27) can be satisfied if one seeks a solution in the form

$$\Phi(\zeta) = -Ae^{\beta E\zeta}, \quad \Psi(\zeta) = Ae^{\beta E\zeta}, \quad u = u(X), \quad T = T_0 - EZ + \frac{E}{\sigma}\theta(X). \tag{28}$$

As a result, we obtain a system of ODEs,

$$\begin{aligned} u'_X &= -\frac{A}{\mu_0} \exp\left(\frac{\beta E}{\sigma}\theta\right) [e^{\beta EX} - e^{-\beta EX}], \\ u &= -\theta''_{XX}. \end{aligned} \tag{29}$$

Eliminating u from these equations and taking into account (25)–(26), (28), we come to the following nonlinear boundary value problem for the excessive temperature:

$$w'''_{xxx} = \gamma e^w \sinh(\varepsilon x), \tag{30}$$

$$w'_x = 0 \quad \text{for} \quad x = 0, \quad w = 0 \quad \text{for} \quad x = 1, \quad w''_{xx} = 0 \quad \text{for} \quad x = 1, \tag{31}$$

where we have set

$$x = \frac{X}{h}, \quad w = \frac{\beta E\theta}{\sigma}, \quad \gamma = \frac{2Ah^3\beta E}{\sigma\mu_0}, \quad \varepsilon = \beta Eh.$$

The last boundary condition in (31) has been derived by passing to the limit for $X \rightarrow h$ in the second equation in (29) with the adhesion condition (25) on the walls taken into account.

Note that the volume rate of flow Q is calculated in terms of the heat flow on the walls by the formula

$$Q = -2\theta'_X(h) = -2\sigma(\beta Eh)^{-1}w'_x(1).$$

Let us prove that for sufficiently large $\gamma > 0$, the boundary value problem (30)–(31) has no solutions.

It is not difficult to show that the Green's function for the linear boundary value problem (30)–(31) with $\gamma = 0$ (see Subsection 18.3-2) has the form

$$G(x, \xi) = \begin{cases} \xi - \frac{1}{2}(x^2 + \xi^2) & \text{for } 0 \leq x \leq \xi \leq 1, \\ \xi - x\xi & \text{for } 0 \leq \xi \leq x \leq 1. \end{cases} \tag{32}$$

Therefore, the nonlinear boundary value problem (30)–(31) is equivalent to the nonlinear integral equation

$$w(x) = \gamma \int_0^1 G(x, \xi) e^{w(\xi)} \sinh(\varepsilon\xi) d\xi. \tag{33}$$

Since $G(x, \xi) \geq 0$, it follows that for $\gamma > 0$ we have $w(x) > 0$. This inequality has a clear physical meaning: if the walls are cooled by the environment, the temperature in the channel is larger than that of the walls.

Consider an auxiliary linear boundary value problem for eigenvalues:

$$y''''_{xxx} = -\lambda \sinh(\varepsilon x)y, \quad (34)$$

$$y = 0 \quad \text{for } x = 0, \quad y''_{xx} = 0 \quad \text{for } x = 0, \quad y'_x = 0 \quad \text{for } x = 1. \quad (35)$$

This problem is equivalent to the linear Fredholm integral equation

$$y(x) = \lambda \int_0^1 G(\xi, x)y(\xi) \sinh(\varepsilon\xi) d\xi. \quad (36)$$

Here the Green's function $G(\xi, x)$ corresponds to the transposition of the variables x and ξ in (32).

Since the kernel of the integral operator (33) is positive, the generalized Jentzch theorem implies that the smallest eigenvalue is positive, $\lambda_0 > 0$, and the corresponding eigenfunction $y_0(x)$ does not change sign on the interval $[0, 1]$. Let us multiply both sides of equation (30) by $y_0(x)$ and integrate the resulting expression in x from 0 to 1. Taking into account the relations

$$\begin{aligned} y_0 w''''_{xxx} &= (y_0 w''_{xx})'_x - (y'_0 w'_x)'_x + (y''_{0xx} w)'_x - y''''_{0xxx} w, \\ y''''_{0xxx} &= -\lambda_0 \sinh(\varepsilon x)y_0, \end{aligned}$$

and the boundary conditions (31) and (35) for the functions w and y_0 , we come to the relation

$$\frac{\lambda_0}{\gamma} = \frac{\int_0^1 y_0(\xi)e^{w(\xi)} \sinh(\varepsilon\xi) d\xi}{\int_0^1 y_0(\xi)w(\xi) \sinh(\varepsilon\xi) d\xi}. \quad (37)$$

Since $w \geq 0$, we have $e^w \geq ew$. This inequality, together with (37), implies the estimate $\lambda_0/\gamma \geq e$. Therefore, for

$$\gamma > \lambda_0/e,$$

the boundary value problem (30)–(31) has no solutions, and for the critical value γ_* we have $\gamma_* < \lambda_0/e$.

Remark. It can be shown that for $0 < \gamma < \gamma_*$, the boundary value problem (30)–(31) has two solutions (one stable and another unstable). For $\gamma = \gamma_*$, there is only one solution.

⊙ References for Section 18.3: P. P. Zabreyko, A. I. Koshelev, et al. (1975), E. Kamke (1977), V. I. Naidenov and A. D. Polyaniin (1990), R. P. Agarwal, D. O'Regan, and P. J. Y. Wong (1998), A. D. Polyaniin and V. F. Zaitsev (2003).

18.4. Reduction of PDEs with Boundary Conditions of the Third Kind to Integral Equations

18.4-1. Usage of Particular Solutions of PDEs for the Construction of Other Solutions.

Let

$$L[w] = 0 \quad (1)$$

be an arbitrary homogeneous linear partial differential equation of any order in the variables x, t with sufficiently smooth coefficients (t may stand for the time or a spatial variable).

There is an effective way to construct solutions of this equation. Suppose that equation (1) has a particular solution

$$\tilde{w}(x, t; \mu) \quad (2)$$

depending on a parameter μ , and the coefficients of the linear differential operator L are independent of μ . Multiplying the particular solution (2) by an arbitrary function $\varphi(\mu)$ and integrating the result with respect to μ over some interval $[\alpha, \beta]$, we obtain the new function

$$w = \int_{\alpha}^{\beta} \tilde{w}(x, t; \mu) \varphi(\mu) d\mu, \quad (3)$$

which is also a solution of the original homogeneous linear equation (1).

Let us mention some useful facts applied for the construction of solutions of boundary value problems with the help of integral representations like (3).

1. The domain of integration in (3) usually coincides with the domain of one of the independent variables of the PDEs under consideration (in particular, if t is the time, one often takes $\alpha = 0$ and $\beta = \infty$).

2. Let $\tilde{w}(x, t)$ be a particular solution of equation (1) with the coefficients independent of t . Then, for any constant μ , the function $\tilde{w}(x, t - \mu)$ is also a solution of equation (1).

3. Suppose that the particular solution (2) satisfies one or several initial or boundary conditions of the form

$$\begin{aligned} w = 0 \quad \text{for} \quad t = 0, & \quad w_t = 0 \quad \text{for} \quad t = 0, \\ w = 0 \quad \text{for} \quad x = a, & \quad w_x = 0 \quad \text{for} \quad x = b. \end{aligned} \quad (4)$$

Then the function (3) also satisfies equation (1) with the same initial or boundary conditions.

Consider a boundary value problem for equation (1). Suppose that one of the boundary conditions has the form (generalized boundary conditions of the third kind)

$$w_x = F(t, w) \quad \text{for} \quad x = 0, \quad (5)$$

and the other boundary and initial conditions are homogeneous and have the form (4).

Suppose that the special solution (2) of equation (1) satisfies all the homogeneous initial and boundary conditions except (5). Let us seek a solution of the corresponding boundary value problem in the form of the integral (3). Substituting this integral into the boundary condition (5), we obtain an integral equation for the function $\varphi(\mu)$. It is important to make a proper choice of the particular solution (2). As an illustration, consider the following example.

18.4-2. Mass Transfer to a Particle in Fluid Flow Complicated by a Surface Reaction.

Consider steady-state diffusion to a particle in laminar viscous incompressible fluid flow. Assume that on the surface of the particle a chemical reaction occurs with rate $F_*(C)$, where C is the mass concentration of a reactant. In particular, for a reaction of order n we have

$$F_*(C) = KC^n, \quad (6)$$

where K is the reaction rate coefficient.

It is assumed that the velocity field in the fluid is known from the solution of the corresponding hydrodynamic problem and can be specified in terms of a flow function ψ (a flow function can be introduced, for instance, for plane and axisymmetric flows). In the diffusion boundary layer approximation, the dimensionless equation of stationary convective diffusion and the boundary

conditions in curvilinear orthogonal coordinates ξ, η, ζ associated with the body surface $\xi = \xi_s$ and the lines of flow have the form

$$\frac{1}{\sqrt{g^s}} \left(\frac{\partial \psi}{\partial \xi} \frac{\partial w}{\partial \eta} - \frac{\partial \psi}{\partial \eta} \frac{\partial w}{\partial \xi} \right) = \frac{1}{\text{Pe}} \frac{\partial^2 w}{\partial \xi^2}, \tag{7}$$

$$-\frac{\partial w}{\partial \xi} = F(w) \quad \text{for } \xi = \xi_s, \quad w \rightarrow 0 \quad \text{for } \xi \rightarrow \infty, \tag{8}$$

where

$$\begin{aligned} \psi &= (\xi - \xi_s)^m f(\eta), \quad g^s = g^s(\eta) = (g_{\xi\xi}g_{\eta\eta}g_{\zeta\zeta})|_{\xi=\xi_s}, \\ w &= \frac{C_\infty - C}{C_\infty}, \quad \text{Pe} = \frac{aU}{D}, \quad F(w) = \frac{aF_*(C)}{DC_\infty}, \end{aligned}$$

a is the characteristic size of the particle (radius), C_∞ is concentration far away from the particle, U is the characteristic flow velocity (far away from the particle), D is the diffusion coefficient, Pe is the Peclet number, and $g_{\xi\xi}, g_{\eta\eta}, g_{\zeta\zeta}$ are the metric tensor components; the value $m = 1$ corresponds to drops or bubbles and $m = 2$ corresponds to solid particles. In the problem stated in terms of (7)–(8), the boundary condition for $\eta = 0$ has been dropped [for $f(0) = 0$, one imposes the condition that the solution is bounded for $\eta = 0$]. When writing equation (7) and the first boundary condition in (8), it has been assumed that the coordinate ξ near the surface $\xi = \xi_s$ is chosen such that the difference $\xi - \xi_s$ determines the distance between the point (ξ_s, η) on the surface of the body and the point (ξ, η) in the flow (i.e., it is assumed that $g_{\xi\xi}|_{\xi=\xi_s} = 1$).

For a reaction of order n (6), the dimensionless rate of surface chemical reaction is described by the expression

$$F(w) = k(1 - w)^n, \quad k = aKC_\infty^{n-1}/D. \tag{9}$$

Further, it is assumed that the domain under consideration is specified by the inequalities $\xi_s \leq \xi < \infty, 0 \leq \eta \leq \eta_0$, and also that the inequality $f(\eta) > 0$ holds for $0 < \eta < \eta_0$, and $f(0) \geq 0$.

Introducing the new variables

$$t = t(\eta) = \frac{1}{n} \int_0^\eta f^{1/n}(\eta)[g^s(\eta)]^{1/2} d\eta, \quad x = \frac{2}{n+1} \text{Pe}^{1/2} \psi^{(n+1)/(2n)}, \tag{10}$$

we reduce (7)–(8) to the following boundary value problem for the unknown function $w(x, t)$:

$$\frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2} + \frac{1-2\nu}{x} \frac{\partial w}{\partial x}, \tag{11}$$

$$w = 0 \quad \text{for } t = 0, \quad w \rightarrow 0 \quad \text{as } x \rightarrow \infty, \tag{12}$$

$$x^{1-2\nu} \frac{\partial w}{\partial x} + \text{Pe}^{-\nu} (2\nu)^{1-2\nu} h_\nu(t) F(w) = 0 \quad \text{for } x = 0, \tag{13}$$

where

$$\nu = (n + 1)^{-1}, \quad h_\nu(t) = f^{-1/n}(\eta(t)).$$

The function $h_\nu(t)$ in the boundary condition on the particle surface (13) is found from the parametric relations $h_\nu = f^{-1/n}(\eta), t = t(\eta)$ (see the first formula in (10)).

Simple verification shows that equation (11) admits the particular solution

$$\tilde{w}(x, t; \mu) = \begin{cases} A(t - \mu)^{\nu-1} \exp\left[-\frac{x^2}{4(t - \mu)}\right] & \text{for } t > \mu, \\ 0 & \text{for } t \leq \mu. \end{cases} \tag{14}$$

Note that for $\nu = \frac{1}{2}$ equation (11) turns into the classical heat transfer equation. In this case, the function (14) for $\mu = 0$ and $A = \frac{1}{2}\pi^{-1/2}$ coincides with the fundamental solution of the heat equation.

Let us seek a solution of problem (11)–(13) in the form of the integral (3) (for $\alpha = 0$ and $\beta = \infty$), into which the function (14) should be inserted. As a result, we obtain

$$w(x, t) = \frac{2^{2\nu-1}}{\Gamma(1-\nu)} \int_0^t (t-\mu)^{\nu-1} \exp\left[-\frac{x^2}{4(t-\mu)}\right] \varphi(\mu) d\mu, \tag{15}$$

where $\Gamma(\nu)$ is the gamma function (here, for the sake of definiteness, we have taken a specific value of the constant A). Obviously, the function (15) satisfies the initial and the boundary conditions (12).

As shown by Sutton (1943), the function (15) has the following limit properties:

$$\lim_{x \rightarrow 0} w = \frac{2^{2\nu-1}}{\Gamma(1-\nu)} \int_0^t (t-\mu)^{\nu-1} \varphi(\mu) d\mu, \quad \lim_{x \rightarrow 0} \left(x^{1-2\nu} \frac{\partial w}{\partial x}\right) = -\varphi(t). \tag{16}$$

Substituting (15) into the boundary condition (13) and taking into account (16), we come to the integral equation for the function $\varphi(t)$:

$$\varphi(t) = \text{Pe}^{-\nu} (2\nu)^{1-2\nu} h_\nu(t) F\left(\frac{2^{2\nu-1}}{\Gamma(1-\nu)} \int_0^t (t-\mu)^{\nu-1} \varphi(\mu) d\mu\right). \tag{17}$$

The replacement

$$\varphi(t) = \text{Pe}^{-\nu} (2\nu)^{1-2\nu} h_\nu(t) F(\psi(t))$$

reduces equation (17) to a more common form (it is assumed that the function F is invertible)

$$\psi(t) = \frac{\nu^{2\nu-1} \text{Pe}^{-\nu}}{\Gamma(1-\nu)} \int_0^t (t-\mu)^{\nu-1} h_\nu(\mu) F(\psi(\mu)) d\mu. \tag{18}$$

After solving equation (17) or (18), formulas (10), (15) can be used to obtain the distribution of concentration in the diffusive boundary layer of the particle.

18.4-3. Integral Equations for Surface Concentration and Diffusion Flux.

Instead of equation (17) for the function φ , it is convenient to consider directly the equations for surface concentration or local diffusion flux—the quantities with a clear physical meaning (in most practical problems these two are the desired quantities).

In view of (10) and (15), surface concentration is determined by the expression

$$w_s = w_s(t) \equiv w(0, t) = \frac{2^{2\nu-1}}{\Gamma(1-\nu)} \int_0^t (t-\mu)^{\nu-1} \varphi(\mu) d\mu = \Lambda_\nu(0, t) * \varphi(t). \tag{19}$$

Note that the operator $\Lambda_\nu(0, t)$ coincides, to within a constant coefficient, with an integral of fractional order ν (see Subsection 10.5-1). Applying the operator $\Lambda_\nu(0, t)$ to both terms of equation (17) and using (17), we obtain the equation for surface concentration:

$$w_s = \text{Pe}^{-\nu} (2\nu)^{1-2\nu} \Lambda_\nu(0, t) * [h_n(t) F(w_s)], \quad w_s = w_s(t). \tag{20}$$

Applying the inverse operator $\Lambda_\nu^{-1}(0, t)$ to both sides of this equation, we come to the following equivalent equation:

$$j_\infty(t) t^\nu \Xi_\nu(t) * w_s(t) = F(w_s(t)), \quad \Xi_\nu(t) \equiv \Gamma(\nu) 2^{2\nu-1} \Lambda_\nu^{-1}(0, t); \tag{21}$$

$$\Xi_\nu(t) * z(t) \equiv \frac{d}{dt} \int_0^t z(\lambda) (t-\lambda)^{-\nu} d\lambda = \frac{z(0)}{t^\nu} + \int_0^t \frac{dz(\lambda)}{d\lambda} (t-\lambda)^{-\nu} d\lambda,$$

where $j_\infty = j_\infty(t)$ is the local diffusion flux corresponding to the diffusion mode of the reaction on the surface (i.e., the boundary condition $w = 1$ for $\xi = \xi_s$):

$$j_\infty(t) = \nu^{2\nu-1} [\Gamma(\nu)]^{-1} \text{Pe}^\nu h_n^{-1}(t) t^{-\nu} = \nu^{2\nu-1} [\Gamma(\nu)]^{-1} \text{Pe}^\nu f^{1/n}(\eta) t^{-\nu}(\eta). \quad (22)$$

From equation (20), combined with the formula

$$j = F(w_s) \quad (23)$$

and the identity $\Gamma(\nu)\Gamma(1-\nu) = \pi / \sin(\pi\nu)$, we obtain a relation between surface concentration and local diffusion flux:

$$w_s(t) = \frac{\sin(\pi\nu)}{\pi} \int_0^t \frac{j(\lambda)}{j_\infty(\lambda)} \lambda^{-\nu} (t-\lambda)^{\nu-1} d\lambda. \quad (24)$$

Substituting (24) into the right-hand side of (23), we obtain an integral equation for local diffusion flux on the particle surface:

$$j = F\left(\frac{\sin(\pi\nu)}{\pi} \int_0^t \frac{j(\lambda)}{j_\infty(\lambda)} \lambda^{-\nu} (t-\lambda)^{\nu-1} d\lambda\right). \quad (25)$$

It is not difficult to show that if the limit local diffusion flux on a part of the body is constant,

$$j_\infty(t) = j_\infty = \text{const} \quad (0 \leq t \leq t_0), \quad (26)$$

then the solution of the nonlinear integral equation (25) reduces to the solution of the algebraic (transcendental) equation

$$j = F(j/j_\infty) \quad (0 \leq t \leq t_0). \quad (27)$$

In view of (23), (27), surface concentration, under the condition (26), is also determined by solving an algebraic equation,

$$w_s = j_\infty^{-1} F(w_s). \quad (28)$$

In the general case, for $j_\infty = j_\infty(t) \neq 0$, it is impossible to obtain an exact analytical solution of integral equations for surface concentration and local diffusion flux (21)–(22) and (25). Therefore, one has to resort to the methods of numerical or approximate integration of these equations.

In engineering, approximations of surface concentration and local flux are sometimes constructed by the *method of equidistant surface*. The essence of this method can be described as follows. First, formula (22) is used to determine the limit local diffusion flux $j = j_\infty(t)$, and then this expression is inserted into equations (27) and (28), i.e., instead of the original integral equations (21) and (25), one solves algebraic (transcendental) equations. Comparison of the approximate results obtained by this method with those of numerical analysis for many typical cases shows that the method of equidistant surface is fairly accurate (for relatively simple reactions, the error does not exceed 20%; see the references at the end of this section). Therefore, when using iteration methods for solving integral equations, it is reasonable to take a solution obtained by the said method as the initial approximation.

18.4-4. Method of Numerical Integration of the Equation for Surface Concentration.

Consider more closely a method of numerical integration of the equation for surface concentration (21)–(22); local diffusion flux in this case is found with the help of (23). Let us represent equation (21) in the form

$$j_\infty(t) \left[w_s(0) + t^\nu \int_0^t \frac{dw_s(\lambda)}{d\lambda} (t-\lambda)^{-\nu} d\lambda \right] = F(w_s(t)). \quad (29)$$

Numerical integration of equation (29) is carried out as follows. First, the segment $[0, t_0]$, $t_0 = t(\eta_0)$, is split into M equal parts

$$[(i - 1)\Delta t, i\Delta t] \quad (i = 1, \dots, M)$$

of length $\Delta t = t_0/M$ and equation (29) is transformed to

$$w_s(0) + (m\Delta t)^\nu \sum_{i=1}^m \int_{(i-1)\Delta t}^{i\Delta t} \frac{dw_s(\lambda)}{d\lambda} (m\Delta t - \lambda)^{-\nu} d\lambda = \frac{F(w_s)}{j_\infty(m\Delta t)}. \tag{30}$$

Then, the derivative $dw_s/d\lambda$ should be approximated on every segment $(i - 1)\Delta t \leq \lambda \leq i\Delta t$ by the expression

$$\frac{dw_s(\lambda)}{d\lambda} = \frac{1}{\Delta t} [w_s(i\Delta t) - w_s((i - 1)\Delta t)],$$

and then one integrates equation (30). After suitable transformations, one obtains

$$\begin{aligned} & \frac{1 - \nu}{m^\nu} w_s(0) - w_s(0) \{ m^{1-\nu} - (m - 1)^{1-\nu} \} + \sum_{i=1}^{m-1} w_s(i\Delta t) A_{m-i} \\ & = -w_s(m\Delta t) + \frac{1 - \nu}{m^\nu} \frac{F(w_s(m\Delta t))}{j_\infty(m\Delta t)}, \quad m = 1, \dots, M - 1, \tag{31} \\ & \nu = \frac{1}{n + 1}, \quad A_m = -2m^{1-\nu} + (m + 1)^{1-\nu} + (m - 1)^{1-\nu}. \end{aligned}$$

The algebraic equation (31) is solved for $m = 1, 2, 3, \dots$ with respect to surface concentration $w_s(m\Delta t)$, which corresponds to $t = m\Delta t$. The left-hand side of (31) contains the values of $w_s(i\Delta t)$ for $1 \leq i \leq m - 1$, which have already been calculated, while the right-hand side of (31) contains the unknown quantity $w_s(m\Delta t)$. Numerical integration of equation (31) starts with preliminary determination of surface concentration $w_s(0)$ at the point of diffusion boundary layer initiation (for $t = 0$) by solving the auxiliary equation

$$w_s(0) = j_\infty^{-1}(0)F(w_s(0)), \tag{32}$$

which coincides with (28) for $t = 0$. If $j_\infty(0) = \infty$, then $w_s(0) = 0$ (this situation occurs at the front critical point of a plate streamlined by fluid). Then the solution procedure goes on in successive order for $m = 1, 2, 3, \dots$, and this process is direct in the sense that no repeated calculations are needed. Naturally, the precision of these calculations depends on the value of Δt .

Remark. The algebraic (transcendental) equation (32) may have several roots, depending on the structure of the function $F(w)$ (thus, there may exist several stationary regimes of reaction on the particle surface). In this case, one has to examine the stability of the solutions.

© References for Section 18.4: W. G. L. Sutton (1943), A. Acrivos and P. L. Shambire (1957), A. D. Polyanin and Yu. A. Sergeev (1980), D. A. Frank-Kamenetskii (1987), Yu. P. Gupalo, A. D. Polyanin, and Yu. S. Ryazantsev (1985).

18.5. Representation of Linear Boundary Value Problems in Terms of Potentials

18.5-1. Basic Types of Potentials for the Laplace Equation and Their Properties.

1°. Let S be a smooth closed surface in the n -dimensional Euclidean space \mathbb{R}^n ($n \geq 2$) that coincides with the boundary of a finite domain $G = G^+$, and let G^- be the exterior infinite domain ($G^+ \cup S \cup G^- = \mathbb{R}^n$).

Consider the n -dimensional Laplace equation

$$\Delta w \equiv \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} = 0. \quad (1)$$

The fundamental solution of equation (1) has the form

$$\mathcal{E}(x, y) = \mathcal{E}(|x - y|) = \begin{cases} \frac{1}{\Omega_n(n-2)} \frac{1}{|x - y|^{n-1}} & \text{if } n \geq 3, \\ \frac{1}{2\pi} \ln \frac{1}{|x - y|} & \text{if } n = 2, \end{cases} \quad (2)$$

where

$$|x - y| = \left[\sum_{k=1}^n (x_k - y_k)^2 \right]^{1/2}, \quad \Omega_n = \frac{2\pi^{n/2}}{\Gamma(n/2)},$$

$|x - y|$ is the distance between points $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$, Ω_n is the area of the unit sphere in \mathbb{R}^n , and $\Gamma(z)$ is the gamma function.

Three integrals depending on x as a parameter define different potentials:

$$\begin{aligned} V(x) &= \int_S \mu(y) \mathcal{E}(x, y) dS_y && \text{(single layer potential),} \\ W(x) &= \int_S \nu(y) \frac{\partial}{\partial n_y} \mathcal{E}(x, y) dS_y && \text{(double layer potential),} \\ Z(x) &= \int_G \rho(y) \mathcal{E}(x, y) dy && \text{(volume potential).} \end{aligned} \quad (3)$$

Here n_y is the direction of the outward (with respect to G^+) normal to the surface S at the point $y \in S$. The functions $\mu(y)$, $\nu(y)$, and $\rho(y)$ are called *densities* of the respective potentials. In what follows, these densities are always assumed absolutely integrable on S or G .

2°. Let $\mu(y) \in C^1(S)$. The single layer potential $V(x)$ is a *harmonic function* [i.e., a function satisfying the Laplace equation (1)] for $x \notin S$, and

$$\lim_{|x| \rightarrow \infty} \frac{V(x)}{\mathcal{E}(x, 0)} = M_1, \quad M_1 = \int_S \mu(y) dS_y;$$

in particular, $\lim_{|x| \rightarrow \infty} V(x) = 0$ for $n \geq 3$, but $\lim_{|x| \rightarrow \infty} V(x) = 0$ for $n = 2$, if and only if $\int_S \mu(y) dS_y = 0$.

The single layer potential is continuous everywhere in \mathbb{R}^n . Moreover, $V(x)$ and its tangential derivatives are continuous across the surface S . The normal derivative of the single layer potential has a jump across the surface S :

$$\left(\frac{\partial V}{\partial n_x} \right)^+ = \frac{1}{2} \mu(x) + \frac{\partial V}{\partial n_x}, \quad \left(\frac{\partial V}{\partial n_x} \right)^- = -\frac{1}{2} \mu(x) + \frac{\partial V}{\partial n_x}. \quad (4)$$

Here the superscripts $+$ and $-$ in the left-hand sides mark the limit values of the normal derivatives from the direction of G^+ and G^- , respectively, i.e.,

$$\left(\frac{\partial V}{\partial n_x} \right)^+ = \lim_{x' \rightarrow x, x' \in G^+} \frac{\partial V}{\partial n_x}, \quad \left(\frac{\partial V}{\partial n_x} \right)^- = \lim_{x' \rightarrow x, x' \in G^-} \frac{\partial V}{\partial n_x}.$$

And in the right-hand sides of (4), the normal derivative is calculated directly on the surface S , i.e.,

$$\frac{\partial V}{\partial n_x} = \int_S \mu(y) \frac{\partial}{\partial n_x} \mathcal{E}(x, y) dS_y, \quad x \in S,$$

which is a continuous function of $x \in S$, and the kernel has a weak singularity on S :

$$\left| \frac{\partial}{\partial n_x} \mathcal{E}(x, y) \right| \leq \frac{\text{const}}{|x - y|^{n-2}}, \quad x, y \in S.$$

3°. Let $\nu(y) \in C^1(S)$. The double layer potential $W(x)$ is a harmonic function of $x \notin S$ and

$$\lim_{|x| \rightarrow \infty} \Omega_n |x|^{n-1} W(x) = M_2, \quad M_2 = \int_S \nu(y) dS_y.$$

Across the surface S , the double layer potential has a jump:

$$W^+(x) = -\frac{1}{2} \nu(x) + W(x), \quad W^-(x) = \frac{1}{2} \nu(x) + W(x), \quad x \in S, \tag{5}$$

where $W^+(x)$ and $W^-(x)$ are the limit values of the double layer potential in the directions from G^+ and G^- , i.e.,

$$W^+(x) = \lim_{x' \rightarrow x, x' \in G^+} W(x'), \quad W^-(x) = \lim_{x' \rightarrow x, x' \in G^-} W(x').$$

The right-hand sides of (5) involve the direct value of the double layer potential on the surface S ,

$$W(x) = \int_S \nu(y) \frac{\partial}{\partial n_y} \mathcal{E}(x, y) dS_y, \quad x \in S,$$

which is a continuous function of $x \in S$, and the kernel has a weak singularity on S :

$$\left| \frac{\partial}{\partial n_y} \mathcal{E}(x, y) \right| \leq \frac{\text{const}}{|x - y|^{n-2}}, \quad x, y \in S.$$

The tangential derivatives of the double layer potential also have a jump across the surface S , but its normal derivative preserves its value across S :

$$\left(\frac{\partial W}{\partial n_x} \right)^+ = \left(\frac{\partial W}{\partial n_x} \right)^-, \quad x \in S.$$

In the case of constant density $\nu = 1$, the *Gauss formula*

$$-\int_S \frac{\partial}{\partial n_y} \mathcal{E}(x, y) dS_y = q(x) \equiv \begin{cases} 1 & \text{if } x \in G^+, \\ 1/2 & \text{if } x \in S, \\ 0 & \text{if } x \in G^- \end{cases} \tag{6}$$

holds for the double layer potential. The integral on the left is interpreted as the solid angle, divided by $\Omega_n(n - 2)$, under which the surface S is seen from the point x .

4°. For $\rho(y) \in C^1(G \cup S)$, the volume potential and its first-order derivatives are continuous everywhere in \mathbb{R}^n and can be calculated by the differentiation under the sign of the integral. Thus, $Z \in C^1(\mathbb{R}^n)$. Moreover,

$$\lim_{|x| \rightarrow \infty} \frac{Z(x)}{\mathcal{E}(x, 0)} = M_3, \quad M_3 = \int_G \rho(y) dy.$$

Its second-order derivatives are continuous outside S , but have a jump on S . In the interior domain G^+ the *Poisson equation* holds

$$\Delta Z = \rho(x), \quad x \in G^+,$$

and in the exterior domain G^- the volume potential satisfies the Laplace equation

$$\Delta Z = 0, \quad x \in G^-.$$

For a finite domain G_1 in \mathbb{R}^n with the boundary $S_1 = \partial G_1$ of class C^1 , the *Gauss formula* for the volume potential holds:

$$\int_{S_1} \frac{\partial Z}{\partial n_x} dS_{1x} = - \int_{G \cap G_1} \rho(y) dy.$$

The integration in the first integral is over the variable x .

18.5-2. Integral Identities. Green's Formula.

Let $\Phi(x)$ be a function of class $C^2(G \cup S)$, where S is a surface of class C^2 . Then the following integral identity, called the *Green's formula*, holds:

$$-\int_G \Delta \Phi(y) \mathcal{E}(x, y) dy + \int_S \left[\frac{\partial \Phi(y)}{\partial n_y} \mathcal{E}(x, y) - \Phi(y) \frac{\partial}{\partial n_y} \mathcal{E}(x, y) \right] dS_y = q(x) \Phi(x). \quad (7)$$

Here $q(x)$ is the function defined by (6). Formula (7) implies that in the domain G the function $\Phi(x)$ can be represented as the sum of a single layer potential, a double layer potential, and a volume potential with the respective densities

$$\mu(y) = \frac{\partial \Phi(y)}{\partial n_y}, \quad \nu(y) = -\Phi(y), \quad \rho(y) = -\Delta \Phi(y).$$

For a function $u(x)$ which is harmonic in the domain G and belongs to the class $C^1(G \cup S)$, the following identity holds:

$$\int_S \left[\frac{\partial w(y)}{\partial n_y} \mathcal{E}(x, y) - w(y) \frac{\partial}{\partial n_y} \mathcal{E}(x, y) \right] dS_y = q(x) w(x), \quad (8)$$

and thus, $w(x)$ can be represented in G as the sum of a single layer potential and a double layer potential with the respective densities

$$\mu(y) = \frac{\partial w(y)}{\partial n_y}, \quad \nu(y) = -w(y).$$

However, the densities in (8) cannot be chosen arbitrary on S , because they are related by the integral identity obtained from (8) for $x \in G^+$.

18.5-3. Reduction of Interior Dirichlet and Neumann Problems to Integral Equations.

1°. *Interior Dirichlet problem (first boundary value problem)*: find a function $w(x)$ that satisfies equation (1) in G^+ and the boundary condition

$$w(x) = \varphi^+(x) \quad \text{for } x \in S, \quad (9)$$

where $\varphi^+(x)$ is a given continuous function on S .

Problem (1), (9) has a solution.* This solution is unique and can be represented in the form of the double layer potential

$$w(x) = \int_S \nu(y) \frac{\partial}{\partial n_y} \mathcal{E}(x, y) dS_y$$

with density $\nu(y)$ which is found as the unique solution of the following Fredholm integral equation of the second kind:

$$-\frac{1}{2}\nu(x) + \int_S \nu(y) \frac{\partial}{\partial n_y} \mathcal{E}(x, y) dS_y = \varphi^+(x), \quad x \in S.$$

2°. *Interior Neumann problem (second boundary value problem)*: find a function $w(x)$ that satisfies equation (1) in G^+ and the boundary condition

$$\frac{\partial w(x)}{\partial n_x} = \psi^+(x) \quad \text{for } x \in S, \quad (10)$$

* In Subsections 18.5-3 and 18.5-4 it is assumed that the surface S is sufficiently smooth.

where $\psi^+(x)$ is a given continuous function on S . This problem has a solution if and only if the function $\psi^+(x)$ satisfies the *compatibility condition*

$$\int_S \psi^+(x) dS_x = 0. \tag{11}$$

A solution of problem (1), (10) with the condition (11) is defined to within an additive constant,

$$w(x) = V(x) + C,$$

where

$$V(x) = \int_S \mu(y) \mathcal{E}(x, y) dS_y$$

is the single layer potential with density μ found by solving the Fredholm integral equation of the second kind

$$\frac{1}{2}\mu(x) + \int_S \mu(y) \frac{\partial}{\partial n_x} \mathcal{E}(x, y) dS_y = \psi^+(x), \quad x \in S. \tag{12}$$

The corresponding homogeneous equation (with $\psi^+(x) = 0$) has a nontrivial solution $\mu_0(x)$, and the nonhomogeneous equation (12) has a solution if the condition (11) is satisfied. The general solution of equation (12) has the form $\mu(x) + A\mu_0(x)$, where A is an arbitrary constant.

18.5-4. Reduction of Exterior Dirichlet and Neumann Problems to Integral Equations.

1°. *Exterior Dirichlet problem (first boundary value problem)*: find a function $w(x)$ that satisfies equation (1) in G^- ($0 \notin G^-$) and the boundary condition

$$w(x) = \varphi^-(x) \quad \text{for } x \in S, \tag{13}$$

where $\varphi^-(x)$ is a given continuous function on S , and it is also required that the following *regularity condition* holds at infinity:

$$\lim_{|x| \rightarrow \infty} |x|^{n-2} w(x) = \text{const}. \tag{14}$$

This problem has a solution; this solution is unique and can be represented in the form

$$w(x) = W(x) + \frac{A}{|x|^{n-2}}, \quad W(x) = \int_S \nu(y) \frac{\partial}{\partial n_y} \mathcal{E}(x, y) dS_y,$$

where A is a constant, $W(x)$ is a double layer potential, $\nu(y)$ its density, which is found by solving the Fredholm integral equation of the second kind:

$$\frac{1}{2}\nu(x) + \int_S \nu(y) \frac{\partial}{\partial n_y} \mathcal{E}(x, y) dS_y = \varphi^-(x) - \frac{A}{|x|^{n-2}}, \quad x \in S. \tag{15}$$

The corresponding homogeneous equation has the nontrivial solution $\tilde{\nu}_0 = 1$. For suitable A , the solution of the nonhomogeneous equation (15) has the form

$$\nu(y) = \nu^-(y) + C,$$

where C is an arbitrary constant, $\nu^-(y)$ is a particular solution of equation (15). The constant A is chosen of the form

$$A = - \int_S \varphi^-(x) \nu_0(x) dS_x,$$

where the auxiliary density $\nu_0(x)$ should satisfy the normalization condition

$$\int_S \frac{\nu_0(y)}{|y|^{n-2}} dS_y = 1. \quad (16)$$

The density $\nu_0(x)$ is a nontrivial solution of the integral equation (12) for the interior Neumann problem with the Neumann boundary values $\psi^+(x) = 0$, $x \in S$, and this density satisfies the following normalization condition equivalent to (16) for $n \geq 3$:

$$V_0(x) \equiv \int_S \nu_0(y) \mathcal{E}(x, y) dS_y = 1, \quad x \in G^+ \cup S.$$

The single layer potential $V_0(x)$ with density $\nu_0(x)$ is called the *equilibrium potential* or the *Roben potential*. The density $\nu_0(x)$ yields the solution of the *Roben electrostatic problem* for charge distribution in a conductor S that produces an equilibrium potential which is constant in the domain G^+ . A certain complexity of the solution of the external Dirichlet problem is due to the fact that a harmonic function $w(x)$ satisfying the regularity condition at infinity generally has a slower decay rate (as $|x| \rightarrow \infty$) than the double layer potential. Therefore, in the general case, $w(x)$ cannot be represented merely in terms of the double layer potential.

2°. *External Neumann problem (second boundary value problem)*: find a function $w(x)$ that satisfies equation (1) in G^- ($0 \notin G^-$) and the boundary condition

$$\frac{\partial w(x)}{\partial n_x} = \psi^-(x) \quad \text{for } x \in S,$$

where $\psi^-(x)$ is a given continuous function on S , and it is also required that the regularity condition (14) hold at infinity.

For $n \geq 3$, a solution of this problem exists and is unique. For $n = 2$, a solution exists if and only if the function $\psi^-(x)$ satisfies the *compatibility condition*

$$\int_S \psi^-(x) dS_x = 0; \quad (17)$$

and the solution is defined to within an arbitrary additive constant. The solution of the external Neumann problem can be represented as the single layer potential

$$w(x) = \int_S \mu(y) \mathcal{E}(x, y) dS_y$$

whose density $\mu(y)$ is determined by solving the Fredholm integral equation of the second kind:

$$-\frac{1}{2}\mu(x) + \int_S \mu(y) \frac{\partial}{\partial n_x} \mathcal{E}(x, y) dS_y = \psi^-(x), \quad x \in S. \quad (18)$$

For $n \geq 3$, this equation has one and only one solution. For $n = 2$, the corresponding homogeneous integral equation (with $\psi^-(x) = 0$) admits the nontrivial solution $\mu_0(x)$, and therefore, the nonhomogeneous equation (18), with the solvability condition (17), has a unique solution $\tilde{\mu}(x)$ such that

$$\int_S \tilde{\mu}(x) dS_x = 0,$$

and its general solution has the form

$$\mu(x) = \tilde{\mu}(x) + c\mu_0(x),$$

where c is an arbitrary constant.

Remark. In a similar way, potentials can be introduced for the heat equation and other equations of mathematical physics. These potentials can also be used for the reduction of the corresponding stationary and nonstationary linear problems to integral equations.

© References for Section 18.5: S. G. Mikhlin (1967), P. P. Zabreyko, A. I. Koshelev et al. (1975), R. Courant and D. Hilbert (1989), A. N. Tikhonov and A. A. Samarskii (1990), I. G. Petrovsky (1991), R. B. Guenther and J. W. Lee (1996), W. McLean (2000).

18.6. Representation of Solutions of Nonlinear PDEs in Terms of Solutions of Linear Integral Equations (Inverse Scattering)

18.6-1. Description of the Zakharov–Shabat Method.

Solutions of some nonlinear PDEs can be expressed through solutions of linear integral equations. Below we outline an approach based on the application of linear integral equations of the form*

$$K(x, y) = F(x, y) + \int_x^\infty K(x, z)N(x, z, y) dz, \quad y \geq x, \quad (1)$$

where the functions F , N , and K may depend on some additional parameters other than the specified arguments. In each specific case, the function N is explicitly expressed through F , and both functions F and N are solutions of some linear PDEs.

Define an operator A_x by

$$A_x f(y) = \begin{cases} \int_x^\infty f(z)N(x, z, y) dz & \text{if } y \geq x, \\ 0 & \text{if } y < x \end{cases}$$

and assume that for each chosen N , it is possible to prove that the operator $I - A_x$ is invertible and its inverse, $(I - A_x)^{-1}$, is continuous, where I is the identity operator. The following three steps represent an algorithm for finding a nonlinear equation that can then be solved by the inverse scattering method.

1°. A specific structure is chosen for the integral equation (1). To that end, one prescribes a relation between the functions N and F (N is expressed through F).

2°. Two suitable linear differential (ordinary or partial) equations are introduced for the function F :

$$L_m F = 0, \quad m = 1, 2. \quad (2)$$

3°. The function K is related to F by equation (1), which can be rewritten as

$$(I - A_x)K = F. \quad (3)$$

Applying the operators L_m involved in (2) to equation (3), we obtain

$$L_m(I - A_x)K = 0, \quad m = 1, 2.$$

This equation can be rewritten in the form

$$(I - A_x)(L_m K) = R_m, \quad m = 1, 2,$$

where R_m contains all nonzero terms of the commutator $[L_m, (I - A_x)]$. Moreover, (1) and (2) should be chosen so that R_m could be represented in the form

$$R_m = (I - A_x)M_m(K), \quad m = 1, 2,$$

where $M_m(K)$ is a nonlinear functional of K . But the operator $I - A_x$ is invertible, and therefore, the function K satisfies the nonlinear differential equations

$$L_m K - M_m(K) = 0, \quad m = 1, 2. \quad (4)$$

It follows that each solution of the linear integral equation (1) is a solution of nonlinear differential equations (4). Of most interest, as a rule, are special cases of one of the equations in (4) or equations derived from (4).

Remark. The first two steps of the algorithm are fundamental and most difficult. Linear differential equations (2) usually correspond to a linear eigenvalue problem (for $m = 1$) and a problem of time-evolution of eigenfunctions (for $m = 2$).

* Such equations are called integral equations of the Gel'fand–Levitan–Marchenko type.

18.6-2. Korteweg–de Vries Equation and Other Nonlinear Equations.

To clarify basic features of the above algorithm, consider some examples.

Example 1. Let us consider the integral equation

$$K(x, y) = F(x, y) + \int_x^\infty K(x, z)F(z, y) dz \tag{5}$$

and write out some identities to be used in the sequel,

$$\partial_x^n \int_x^\infty K(x, z)F(z, y) dz = \int_x^\infty F(z, y)\partial_x^n K(x, z) dz + A_n, \tag{6}$$

$$\int_x^\infty K(x, z)\partial_x^n F(z, y) dz = (-1)^n \int_x^\infty F(z, y)\partial_z^n K(x, z) dz + B_n, \tag{7}$$

where A_n are defined by the recurrence relations

$$A_1 = -K(x, x)F(x, y), \quad A_n = (A_{n-1})_x - F(x, y)[\partial_x^{n-1} K(x, z)]_{z=x},$$

and

$$B_1 = -K(x, x)F(x, y), \quad B_2 = -K(x, x)\partial_x F(x, y) + [\partial_z K(x, z)]_{z=x}F(x, y), \quad \dots$$

Let us introduce an operator L_1 and require that F satisfy the linear equation

$$L_1 F \equiv (\partial_x^2 - \partial_y^2)F(x, y) = 0. \tag{8}$$

Applying the operator L_1 to (5) and taking into account (6), (7), we obtain

$$(\partial_x^2 - \partial_y^2)K(x, y) = \int_x^\infty F(x, z)(\partial_x^2 - \partial_y^2)K(x, z) dz - 2F(x, y)\frac{d}{dx}K(x, x).$$

Using the equation $F = (I - A_x)K$ and taking into account that the operator $I - A_x$ is invertible, we finally get

$$(\partial_x^2 - \partial_y^2)K(x, y) + u(x)K(x, y) = 0, \tag{9}$$

where the function $u(x)$ is defined by

$$u(x) = 2\frac{d}{dx}K(x, x). \tag{10}$$

Let us require that F satisfy the linear equation

$$L_2 F = (\partial_t + (\partial_x + \partial_y)^3)F = 0 \tag{11}$$

and apply the operator L_2 to (5). We thus obtain

$$(\partial_t + (\partial_x + \partial_y)^3)K(x, y) = (\partial_t + (\partial_x + \partial_y)^3) \int_x^\infty K(x, z)F(z, y) dz.$$

A procedure similar to the above calculations for the operator L_1 yields

$$K_t + (\partial_x + \partial_y)^3 K + 3u(\partial_x + \partial_y)K = 0. \tag{12}$$

For the characteristic $y = x$, equation (12) can be rewritten in terms of $u = 2(d/dx)K(x, x)$. Differentiating (12) with respect to x and rearranging terms, we arrive at the *Korteweg–de Vries equation*

$$u_t + 6uu_x + u_{xxx} = 0.$$

Any function F satisfying the linear equations (8), (11) and rapidly decaying as $x \rightarrow +\infty$ generates a solution of the Korteweg–de Vries equation. To this end, one should solve the linear integral equation (5) for the function K and express u through K by (10).

Example 2. Consider the integral equation

$$K(x, y) = F(x, y) + \frac{\sigma}{4} \int_x^\infty \int_x^\infty K(x, z)F(z, u)F(u, y) dz du, \tag{13}$$

where $\sigma = \pm 1$. Here and in what follows, the coefficients are chosen with a view to simplifying the calculations. Let the operator L_1 have the form

$$L_1 F = (\partial_x - \partial_y)F = 0, \tag{14}$$

which implies that

$$F(x, y) = F\left(\frac{x+y}{2}\right).$$

Shifting the lower limit of integration to zero, we rewrite equation (13) in the form

$$K(x, y) = F\left(\frac{x+y}{2}\right) + \frac{\sigma}{4} \int_0^\infty \int_0^\infty K(x, x+\zeta) F\left(\frac{2x+\zeta+\eta}{2}\right) F\left(\frac{x+\eta+y}{2}\right) d\zeta d\eta, \tag{15}$$

or, equivalently,

$$[(I - \sigma A_x)K](x, y) = F\left(\frac{x+y}{2}\right),$$

where the operator A_x is defined by

$$A_x f(y) = \frac{1}{4} \int_0^\infty \int_0^\infty f(\zeta) F\left(\frac{2x+\zeta+\eta}{2}\right) F\left(\frac{x+\eta+y}{2}\right) d\zeta d\eta.$$

Introducing the function

$$K_2(x, z) = \int_0^\infty K(x, x+\zeta) F\left(\frac{x+\zeta+z}{2}\right) d\zeta, \tag{16}$$

we can rewrite equation (13) as

$$K(x, y) = F\left(\frac{x+y}{2}\right) + \frac{\sigma}{4} \int_0^\infty K_2(x, x+\eta) F\left(\frac{x+\eta+y}{2}\right) d\eta. \tag{17}$$

Applying the operator L_1 of (14) to equation (17), and the operator $\partial_x + \partial_z$ to (16), and taking into account the invertibility of $I - \sigma A_x$, we find, after appropriate calculations, that

$$(\partial_x + \partial_y)K_2(x, y) = -2K(x, x)K(x, y), \tag{18}$$

$$(\partial_x - \partial_y)K(x, y) = -\frac{\sigma}{2} K(x, x)K_2(x, y). \tag{19}$$

Applying the operator $\partial_x + \partial_y$ to (15), we get

$$F'\left(\frac{x+y}{2}\right) = (I - \sigma A_x) \left[(\partial_x + \partial_y)K(x, y) + \frac{\sigma}{2} K_2(x, x)K(x, y) \right]. \tag{20}$$

Let us require that the function F satisfy the second linear equation

$$L_2 F = (\partial_t + (\partial_x + \partial_y)^3)F = 0. \tag{21}$$

Applying the operator L_2 to equation (15) and taking into account the above auxiliary relations (18)–(20), we ultimately find that

$$[\partial_t + (\partial_x + \partial_y)^3]K(x, y) = 3\sigma K(x, x)K(x, y)\partial_x K(x, x) + 3\sigma K^2(x, x)(\partial_x + \partial_y)K(x, y) \tag{22}$$

for $y \geq x$. Now, by setting $q(x, t) = K(x, x; t)$, we rewrite equation (22), for $y = x$, in terms of the dependent variable q to obtain the *modified Korteweg–de Vries equation*

$$q_t + q_{xxx} = 6\sigma q^2 q_x. \tag{23}$$

Thus, each solution of the equations $L_i F = 0$, $i = 1, 2$, with a sufficiently fast decay rate as $x \rightarrow \infty$ determines a solution of equation (23). Note that we have to solve the linear integral equation (13) at an intermediate step.

Example 3. Consider the *Boussinesq equation*

$$w_{tt} + (w w_x)_x + w_{xxx} = 0.$$

This equation arises in several physical applications: propagation of long waves in shallow water, one-dimensional nonlinear lattice-waves, vibrations in a nonlinear string, and ion sound waves in plasma.

It can be shown that any rapidly decaying (as $x \rightarrow +\infty$) function $F = F(x, y; t)$, which simultaneously satisfies the following two linear partial differential equations:

$$\begin{aligned} F_t + \sqrt{3}(F_{xx} - F_{yy}) &= 0, \\ F_{xxx} + F_{yyy} &= 0, \end{aligned}$$

generates a solution

$$w = 12 \frac{d}{dx} K(x, x; t)$$

of the Boussinesq equation, where $K(x, y; t)$ is a solution of the linear Gel'fand–Levitan–Marchenko integral equation

$$K(x, y; t) + F(x, y; t) + \int_x^\infty K(x, s; t)F(s, y; t) ds = 0.$$

Time t appears here as a parameter.

Example 4. Consider the *Kadomtsev–Petviashvili equation*

$$(w_t + w_{xxx} - 6ww_x)_x + 3aw_{yy} = 0.$$

It can be shown that any rapidly decaying (as $x \rightarrow +\infty$) function $F = F(x, z; y, t)$, which simultaneously satisfies the following two linear partial differential equations:

$$\begin{aligned}\sqrt{a} F_y + F_{xx} - F_{zz} &= 0, \\ F_t + 4F_{xxx} + 4F_{zzz} &= 0,\end{aligned}$$

generates a solution

$$w = -2 \frac{d}{dx} K(x, x; y, t)$$

of the Kadomtsev–Petviashvili equation, where $K = K(x, z; y, t)$ is a solution of the linear Gel'fand–Levitan–Marchenko integral equation

$$K(x, z; y, t) + F(x, z; y, t) + \int_x^\infty K(x, s; y, t) F(s, z; y, t) ds = 0.$$

Here the variables y and t are regarded as parameters.

⊙ References for Section 18.6: V. E. Zakharov and A. B. Shabat (1974), S. P. Novikov, S. V. Manakov, L. B. Pitaevskii, and V. E. Zakharov (1984), M. J. Ablowitz and P. A. Clarkson (1991), A. D. Polyaniin and V. F. Zaitsev (2004).

Supplements

Supplement 1

Elementary Functions and Their Properties

► Throughout Supplement 1 it is assumed that n is a positive integer, unless otherwise specified.

1.1. Power, Exponential, and Logarithmic Functions

1.1-1. Properties of the Power Function.

Basic properties of the power function:

$$x^\alpha x^\beta = x^{\alpha+\beta}, \quad (x_1 x_2)^\alpha = x_1^\alpha x_2^\alpha, \quad (x^\alpha)^\beta = x^{\alpha\beta},$$

for any α and β , where $x > 0$, $x_1 > 0$, $x_2 > 0$.

Differentiation and integration formulas:

$$(x^\alpha)' = \alpha x^{\alpha-1}, \quad \int x^\alpha dx = \begin{cases} \frac{x^{\alpha+1}}{\alpha+1} + C & \text{if } \alpha \neq -1, \\ \ln|x| + C & \text{if } \alpha = -1. \end{cases}$$

The Taylor series expansion in a neighborhood of an arbitrary point:

$$x^\alpha = \sum_{n=0}^{\infty} C_\alpha^n x_0^{\alpha-n} (x-x_0)^n \quad \text{for } |x-x_0| < |x_0|,$$

where $C_\alpha^n = \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}$ are binomial coefficients.

1.1-2. Properties of the Exponential Function.

Basic properties of the exponential function:

$$a^{x_1} a^{x_2} = a^{x_1+x_2}, \quad a^x b^x = (ab)^x, \quad (a^{x_1})^{x_2} = a^{x_1 x_2},$$

where $a > 0$ and $b > 0$.

Number e , base of natural (Napierian) logarithms, and the function e^x :

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = 2.718281\dots, \quad e^x = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n.$$

The formula for passing from an arbitrary base a to the base e of natural logarithms:

$$a^x = e^{x \ln a}.$$

The inequality

$$a^{x_1} > a^{x_2} \iff \begin{cases} x_1 > x_2 & \text{if } a > 1, \\ x_1 < x_2 & \text{if } 0 < a < 1. \end{cases}$$

The limit relations for any $a > 1$ and $b > 0$:

$$\lim_{x \rightarrow +\infty} \frac{a^x}{|x|^b} = \infty, \quad \lim_{x \rightarrow -\infty} a^x |x|^b = 0.$$

Differentiation and integration formulas:

$$(e^x)' = e^x, \quad \int e^x dx = e^x + C;$$

$$(a^x)' = a^x \ln a, \quad \int a^x dx = \frac{a^x}{\ln a} + C.$$

The expansion in power series:

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots = \sum_{k=0}^{\infty} \frac{x^k}{k!}.$$

1.1-3. Properties of the Logarithmic Function.

By definition, the logarithmic function is the inverse of the exponential function. The following equivalence relation holds:

$$y = \log_a x \iff x = a^y,$$

where $a > 0$, $a \neq 1$.

Basic properties of the logarithmic function:

$$a^{\log_a x} = x, \quad \log_a (x_1 x_2) = \log_a x_1 + \log_a x_2,$$

$$\log_a (x^k) = k \log_a x, \quad \log_a x = \frac{\log_b x}{\log_b a},$$

where $x > 0$, $x_1 > 0$, $x_2 > 0$, $a > 0$, $a \neq 1$, $b > 0$, $b \neq 1$.

The simplest inequality:

$$\log_a x_1 > \log_a x_2 \iff \begin{cases} x_1 > x_2 & \text{if } a > 1, \\ x_1 < x_2 & \text{if } 0 < a < 1. \end{cases}$$

For any $b > 0$, the following limit relations hold:

$$\lim_{x \rightarrow +\infty} \frac{\log_a x}{x^b} = 0, \quad \lim_{x \rightarrow +0} x^b \log_a x = 0.$$

The logarithmic function with the base e (*base of natural logarithms* or *Napierian base*) is denoted by

$$\log_e x = \ln x,$$

where $e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = 2.718281 \dots$

Formulas for passing from an arbitrary base a to the Napierian base e :

$$\log_a x = \frac{\ln x}{\ln a}.$$

Differentiation and integration formulas:

$$(\ln x)' = \frac{1}{x}, \quad \int \ln x \, dx = x \ln x - x + C.$$

Expansion in power series:

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots = \sum_{k=1}^{\infty} (-1)^{k-1} \frac{x^k}{k}, \quad |x| < 1;$$

$$\ln\left(\frac{x+1}{x-1}\right) = \frac{2}{x} + \frac{2}{3x^3} + \frac{2}{5x^5} + \dots = 2 \sum_{k=1}^{\infty} \frac{1}{(2k-1)x^{2k-1}}, \quad |x| > 1;$$

$$\ln x = 2\left(\frac{x-1}{x+1}\right) + \frac{2}{3}\left(\frac{x-1}{x+1}\right)^3 + \frac{2}{5}\left(\frac{x-1}{x+1}\right)^5 + \dots = 2 \sum_{k=1}^{\infty} \frac{1}{2k-1} \left(\frac{x-1}{x+1}\right)^{2k-1}, \quad x > 0.$$

1.2. Trigonometric Functions

1.2-1. Simplest Relations.

$$\begin{aligned} \sin^2 x + \cos^2 x &= 1, & \tan x \cot x &= 1, \\ \sin(-x) &= -\sin x, & \cos(-x) &= \cos x, \\ \tan x &= \frac{\sin x}{\cos x}, & \cot x &= \frac{\cos x}{\sin x}, \\ \tan(-x) &= -\tan x, & \cot(-x) &= -\cot x, \\ 1 + \tan^2 x &= \frac{1}{\cos^2 x}, & 1 + \cot^2 x &= \frac{1}{\sin^2 x}. \end{aligned}$$

1.2-2. Reduction Formulas.

$$\begin{aligned} \sin(x \pm 2n\pi) &= \sin x, & \cos(x \pm 2n\pi) &= \cos x, \\ \sin(x \pm n\pi) &= (-1)^n \sin x, & \cos(x \pm n\pi) &= (-1)^n \cos x, \\ \sin\left(x \pm \frac{2n+1}{2}\pi\right) &= \pm(-1)^n \cos x, & \cos\left(x \pm \frac{2n+1}{2}\pi\right) &= \mp(-1)^n \sin x, \\ \sin\left(x \pm \frac{\pi}{4}\right) &= \frac{\sqrt{2}}{2}(\sin x \pm \cos x), & \cos\left(x \pm \frac{\pi}{4}\right) &= \frac{\sqrt{2}}{2}(\cos x \mp \sin x), \\ \tan(x \pm n\pi) &= \tan x, & \cot(x \pm n\pi) &= \cot x, \\ \tan\left(x \pm \frac{2n+1}{2}\pi\right) &= -\cot x, & \cot\left(x \pm \frac{2n+1}{2}\pi\right) &= -\tan x, \\ \tan\left(x \pm \frac{\pi}{4}\right) &= \frac{\tan x \pm 1}{1 \mp \tan x}, & \cot\left(x \pm \frac{\pi}{4}\right) &= \frac{\cot x \mp 1}{1 \pm \cot x}, \end{aligned}$$

where $n = 1, 2, \dots$

1.2-3. Relations Between Trigonometric Functions of Single Argument.
--

$$\begin{aligned}\sin x &= \pm\sqrt{1-\cos^2 x} = \pm\frac{\tan x}{\sqrt{1+\tan^2 x}} = \pm\frac{1}{\sqrt{1+\cot^2 x}}, \\ \cos x &= \pm\sqrt{1-\sin^2 x} = \pm\frac{1}{\sqrt{1+\tan^2 x}} = \pm\frac{\cot x}{\sqrt{1+\cot^2 x}}, \\ \tan x &= \pm\frac{\sin x}{\sqrt{1-\sin^2 x}} = \pm\frac{\sqrt{1-\cos^2 x}}{\cos x} = \frac{1}{\cot x}, \\ \cot x &= \pm\frac{\sqrt{1-\sin^2 x}}{\sin x} = \pm\frac{\cos x}{\sqrt{1-\cos^2 x}} = \frac{1}{\tan x}.\end{aligned}$$

The sign before the radical is determined by the quarter in which the argument takes its values.

1.2-4. Addition and Subtraction of Trigonometric Functions.

$$\begin{aligned}\sin x + \sin y &= 2 \sin\left(\frac{x+y}{2}\right) \cos\left(\frac{x-y}{2}\right), \\ \sin x - \sin y &= 2 \sin\left(\frac{x-y}{2}\right) \cos\left(\frac{x+y}{2}\right), \\ \cos x + \cos y &= 2 \cos\left(\frac{x+y}{2}\right) \cos\left(\frac{x-y}{2}\right), \\ \cos x - \cos y &= -2 \sin\left(\frac{x+y}{2}\right) \sin\left(\frac{x-y}{2}\right), \\ \sin^2 x - \sin^2 y &= \cos^2 y - \cos^2 x = \sin(x+y) \sin(x-y), \\ \sin^2 x - \cos^2 y &= -\cos(x+y) \cos(x-y), \\ \tan x \pm \tan y &= \frac{\sin(x \pm y)}{\cos x \cos y}, \quad \cot x \pm \cot y = \frac{\sin(y \pm x)}{\sin x \sin y}, \\ a \cos x + b \sin x &= r \sin(x + \varphi) = r \cos(x - \psi).\end{aligned}$$

Here $r = \sqrt{a^2 + b^2}$, $\sin \varphi = a/r$, $\cos \varphi = b/r$, $\sin \psi = b/r$, and $\cos \psi = a/r$.

1.2-5. Products of Trigonometric Functions.

$$\begin{aligned}\sin x \sin y &= \frac{1}{2}[\cos(x-y) - \cos(x+y)], \\ \cos x \cos y &= \frac{1}{2}[\cos(x-y) + \cos(x+y)], \\ \sin x \cos y &= \frac{1}{2}[\sin(x-y) + \sin(x+y)].\end{aligned}$$

1.2-6. Powers of Trigonometric Functions.

$$\begin{aligned}\cos^2 x &= \frac{1}{2} \cos 2x + \frac{1}{2}, & \sin^2 x &= -\frac{1}{2} \cos 2x + \frac{1}{2}, \\ \cos^3 x &= \frac{1}{4} \cos 3x + \frac{3}{4} \cos x, & \sin^3 x &= -\frac{1}{4} \sin 3x + \frac{3}{4} \sin x, \\ \cos^4 x &= \frac{1}{8} \cos 4x + \frac{1}{2} \cos 2x + \frac{3}{8}, & \sin^4 x &= \frac{1}{8} \cos 4x - \frac{1}{2} \cos 2x + \frac{3}{8}, \\ \cos^5 x &= \frac{1}{16} \cos 5x + \frac{5}{16} \cos 3x + \frac{5}{8} \cos x, & \sin^5 x &= \frac{1}{16} \sin 5x - \frac{5}{16} \sin 3x + \frac{5}{8} \sin x,\end{aligned}$$

$$\begin{aligned}\cos^{2n} x &= \frac{1}{2^{2n-1}} \sum_{k=0}^{n-1} C_{2n}^k \cos[2(n-k)x] + \frac{1}{2^{2n}} C_{2n}^n, \\ \cos^{2n+1} x &= \frac{1}{2^{2n}} \sum_{k=0}^n C_{2n+1}^k \cos[(2n-2k+1)x], \\ \sin^{2n} x &= \frac{1}{2^{2n-1}} \sum_{k=0}^{n-1} (-1)^{n-k} C_{2n}^k \cos[2(n-k)x] + \frac{1}{2^{2n}} C_{2n}^n, \\ \sin^{2n+1} x &= \frac{1}{2^{2n}} \sum_{k=0}^n (-1)^{n-k} C_{2n+1}^k \sin[(2n-2k+1)x].\end{aligned}$$

Here $n = 1, 2, \dots$ and $C_m^k = \frac{m!}{k!(m-k)!}$ are binomial coefficients ($0! = 1$).

1.2-7. Addition Formulas.

$$\begin{aligned}\sin(x \pm y) &= \sin x \cos y \pm \cos x \sin y, & \cos(x \pm y) &= \cos x \cos y \mp \sin x \sin y, \\ \tan(x \pm y) &= \frac{\tan x \pm \tan y}{1 \mp \tan x \tan y}, & \cot(x \pm y) &= \frac{1 \mp \tan x \tan y}{\tan x \pm \tan y}.\end{aligned}$$

1.2-8. Trigonometric Functions of Multiple Arguments.

$$\begin{aligned}\cos 2x &= 2 \cos^2 x - 1 = 1 - 2 \sin^2 x, & \sin 2x &= 2 \sin x \cos x, \\ \cos 3x &= -3 \cos x + 4 \cos^3 x, & \sin 3x &= 3 \sin x - 4 \sin^3 x, \\ \cos 4x &= 1 - 8 \cos^2 x + 8 \cos^4 x, & \sin 4x &= 4 \cos x (\sin x - 2 \sin^3 x), \\ \cos 5x &= 5 \cos x - 20 \cos^3 x + 16 \cos^5 x, & \sin 5x &= 5 \sin x - 20 \sin^3 x + 16 \sin^5 x, \\ \cos(2nx) &= 1 + \sum_{k=1}^n (-1)^k \frac{n^2(n^2-1) \dots [n^2-(k-1)^2]}{(2k)!} 4^k \sin^{2k} x, \\ \cos[(2n+1)x] &= \cos x \left\{ 1 + \sum_{k=1}^n (-1)^k \frac{[(2n+1)^2-1][(2n+1)^2-3^2] \dots [(2n+1)^2-(2k-1)^2]}{(2k)!} \sin^{2k} x \right\}, \\ \sin(2nx) &= 2n \cos x \left[\sin x + \sum_{k=1}^n (-4)^k \frac{(n^2-1)(n^2-2^2) \dots (n^2-k^2)}{(2k-1)!} \sin^{2k-1} x \right], \\ \sin[(2n+1)x] &= (2n+1) \left\{ \sin x + \sum_{k=1}^n (-1)^k \frac{[(2n+1)^2-1][(2n+1)^2-3^2] \dots [(2n+1)^2-(2k-1)^2]}{(2k+1)!} \sin^{2k+1} x \right\}, \\ \tan 2x &= \frac{2 \tan x}{1 - \tan^2 x}, & \tan 3x &= \frac{3 \tan x - \tan^3 x}{1 - 3 \tan^2 x}, & \tan 4x &= \frac{4 \tan x - 4 \tan^3 x}{1 - 6 \tan^2 x + \tan^4 x},\end{aligned}$$

where $n = 1, 2, \dots$

1.2-9. Trigonometric Functions of Half Argument.

$$\sin^2 \frac{x}{2} = \frac{1 - \cos x}{2}, \quad \cos^2 \frac{x}{2} = \frac{1 + \cos x}{2},$$

$$\tan \frac{x}{2} = \frac{\sin x}{1 + \cos x} = \frac{1 - \cos x}{\sin x}, \quad \cot \frac{x}{2} = \frac{\sin x}{1 - \cos x} = \frac{1 + \cos x}{\sin x},$$

$$\sin x = \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}, \quad \cos x = \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}, \quad \tan x = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}}.$$

1.2-10. Differentiation Formulas.

$$\frac{d \sin x}{dx} = \cos x, \quad \frac{d \cos x}{dx} = -\sin x, \quad \frac{d \tan x}{dx} = \frac{1}{\cos^2 x}, \quad \frac{d \cot x}{dx} = -\frac{1}{\sin^2 x}.$$

1.2-11. Integration Formulas.

$$\int \sin x \, dx = -\cos x + C, \quad \int \cos x \, dx = \sin x + C,$$

$$\int \tan x \, dx = -\ln |\cos x| + C, \quad \int \cot x \, dx = \ln |\sin x| + C,$$

where C is an arbitrary constant.

1.2-12. Expansion in Power Series.

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots + (-1)^n \frac{x^{2n}}{(2n)!} + \cdots \quad (|x| < \infty),$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + \cdots \quad (|x| < \infty),$$

$$\tan x = x + \frac{x^3}{3} + \frac{2x^5}{15} + \frac{17x^7}{315} + \cdots + \frac{2^{2n}(2^{2n}-1)|B_{2n}|}{(2n)!} x^{2n-1} + \cdots \quad (|x| < \pi/2),$$

$$\cot x = \frac{1}{x} - \left(\frac{x}{3} + \frac{x^3}{45} + \frac{2x^5}{945} + \cdots + \frac{2^{2n}|B_{2n}|}{(2n)!} x^{2n-1} + \cdots \right) \quad (0 < |x| < \pi),$$

$$\frac{1}{\cos x} = 1 + \frac{x^2}{2} + \frac{5x^4}{24} + \frac{61x^6}{720} + \cdots + \frac{(-1)^n E_{2n}}{(2n)!} x^{2n} + \cdots \quad (|x| < \pi/2),$$

$$\frac{1}{\sin x} = \frac{1}{x} + \frac{x}{6} + \frac{7x^3}{360} + \cdots + \frac{(-1)^{n-1} 2(2^{2n-1}-1)B_{2n}}{(2n)!} x^{2n-1} + \cdots \quad (0 < |x| < \pi),$$

where B_n and E_n are Bernoulli and Euler numbers (see Supplements 11.1-3 and 11.1-4).

1.2-13. Representation in the Form of Infinite Products.

$$\sin x = x \left(1 - \frac{x^2}{\pi^2}\right) \left(1 - \frac{x^2}{4\pi^2}\right) \left(1 - \frac{x^2}{9\pi^2}\right) \cdots \left(1 - \frac{x^2}{n^2\pi^2}\right) \cdots$$

$$\cos x = \left(1 - \frac{4x^2}{\pi^2}\right) \left(1 - \frac{4x^2}{9\pi^2}\right) \left(1 - \frac{4x^2}{25\pi^2}\right) \cdots \left(1 - \frac{4x^2}{(2n+1)^2\pi^2}\right) \cdots$$

1.2-14. Euler and de Moivre Formulas. Relationship with Hyperbolic Functions.

$$e^{y+ix} = e^y(\cos x + i \sin x), \quad (\cos x + i \sin x)^n = \cos(nx) + i \sin(nx), \quad i^2 = -1,$$

$$\sin(ix) = i \sinh x, \quad \cos(ix) = \cosh x, \quad \tan(ix) = i \tanh x, \quad \cot(ix) = -i \coth x.$$

1.3. Inverse Trigonometric Functions

1.3-1. Definitions of Inverse Trigonometric Functions.

Inverse trigonometric functions (arc functions) are the functions that are inverse to the trigonometric functions. Since the trigonometric functions $\sin x$, $\cos x$, $\tan x$, $\cot x$ are periodic, the corresponding inverse functions, denoted by $\text{Arcsin } x$, $\text{Arccos } x$, $\text{Arctan } x$, $\text{Arccot } x$, are multi-valued. The following relations define the multi-valued inverse trigonometric functions:

$$\sin(\text{Arcsin } x) = x, \quad \cos(\text{Arccos } x) = x,$$

$$\tan(\text{Arctan } x) = x, \quad \cot(\text{Arccot } x) = x.$$

These functions admit the following verbal definitions: $\text{Arcsin } x$ is the angle whose sine is equal to x ; $\text{Arccos } x$ is the angle whose cosine is equal to x ; $\text{Arctan } x$ is the angle whose tangent is equal to x ; $\text{Arccot } x$ is the angle whose cotangent is equal to x .

The principal (single-valued) branches of the inverse trigonometric functions are denoted by

$$\arcsin x \equiv \sin^{-1} x \quad (\text{arcsine is the inverse of sine}),$$

$$\arccos x \equiv \cos^{-1} x \quad (\text{arccosine is the inverse of cosine}),$$

$$\arctan x \equiv \tan^{-1} x \quad (\text{arctangent is the inverse of tangent}),$$

$$\text{arccot } x \equiv \cot^{-1} x \quad (\text{arccotangent is the inverse of cotangent})$$

and are determined by the inequalities

$$-\frac{\pi}{2} \leq \arcsin x \leq \frac{\pi}{2}, \quad 0 \leq \arccos x \leq \pi \quad (-1 \leq x \leq 1);$$

$$-\frac{\pi}{2} < \arctan x < \frac{\pi}{2}, \quad 0 < \text{arccot } x < \pi \quad (-\infty < x < \infty).$$

The following equivalent relations can be taken as definitions of single-valued inverse trigonometric functions:

$$y = \arcsin x, \quad -1 \leq x \leq 1 \quad \iff \quad x = \sin y, \quad -\frac{\pi}{2} \leq y \leq \frac{\pi}{2};$$

$$y = \arccos x, \quad -1 \leq x \leq 1 \quad \iff \quad x = \cos y, \quad 0 \leq y \leq \pi;$$

$$y = \arctan x, \quad -\infty < x < +\infty \quad \iff \quad x = \tan y, \quad -\frac{\pi}{2} < y < \frac{\pi}{2};$$

$$y = \text{arccot } x, \quad -\infty < x < +\infty \quad \iff \quad x = \cot y, \quad 0 < y < \pi.$$

The multi-valued and the single-valued inverse trigonometric functions are related by the formulas

$$\text{Arcsin } x = (-1)^n \arcsin x + \pi n,$$

$$\text{Arccos } x = \pm \arccos x + 2\pi n,$$

$$\text{Arctan } x = \arctan x + \pi n,$$

$$\text{Arccot } x = \text{arccot } x + \pi n,$$

where $n = 0, \pm 1, \pm 2, \dots$

1.3-2. Simplest Formulas.

$$\begin{aligned}\sin(\arcsin x) &= x, & \cos(\arccos x) &= x, \\ \tan(\arctan x) &= x, & \cot(\operatorname{arccot} x) &= x.\end{aligned}$$

1.3-3. Some Properties.

$$\begin{aligned}\arcsin(-x) &= -\arcsin x, & \arccos(-x) &= \pi - \arccos x, \\ \arctan(-x) &= -\arctan x, & \operatorname{arccot}(-x) &= \pi - \operatorname{arccot} x,\end{aligned}$$

$$\arcsin(\sin x) = \begin{cases} x - 2n\pi & \text{if } 2n\pi - \frac{\pi}{2} \leq x \leq 2n\pi + \frac{\pi}{2}, \\ -x + 2(n+1)\pi & \text{if } (2n+1)\pi - \frac{\pi}{2} \leq x \leq 2(n+1)\pi + \frac{\pi}{2}, \end{cases}$$

$$\arccos(\cos x) = \begin{cases} x - 2n\pi & \text{if } 2n\pi \leq x \leq (2n+1)\pi, \\ -x + 2(n+1)\pi & \text{if } (2n+1)\pi \leq x \leq 2(n+1)\pi, \end{cases}$$

$$\begin{aligned}\arctan(\tan x) &= x - n\pi & \text{if } n\pi - \frac{\pi}{2} < x < n\pi + \frac{\pi}{2}, \\ \operatorname{arccot}(\cot x) &= x - n\pi & \text{if } n\pi < x < (n+1)\pi.\end{aligned}$$

1.3-4. Relations Between Inverse Trigonometric Functions.

$$\arcsin x + \arccos x = \frac{\pi}{2}, \quad \arctan x + \operatorname{arccot} x = \frac{\pi}{2};$$

$$\arcsin x = \begin{cases} \arccos \sqrt{1-x^2} & \text{if } 0 \leq x \leq 1, \\ -\arccos \sqrt{1-x^2} & \text{if } -1 \leq x \leq 0, \\ \arctan \frac{x}{\sqrt{1-x^2}} & \text{if } -1 < x < 1, \\ \operatorname{arccot} \frac{\sqrt{1-x^2}}{x} - \pi & \text{if } -1 \leq x < 0; \end{cases}$$

$$\arccos x = \begin{cases} \arcsin \sqrt{1-x^2} & \text{if } 0 \leq x \leq 1, \\ \pi - \arcsin \sqrt{1-x^2} & \text{if } -1 \leq x \leq 0, \\ \arctan \frac{\sqrt{1-x^2}}{x} & \text{if } 0 < x \leq 1, \\ \operatorname{arccot} \frac{x}{\sqrt{1-x^2}} & \text{if } -1 < x < 1; \end{cases}$$

$$\arctan x = \begin{cases} \arcsin \frac{x}{\sqrt{1+x^2}} & \text{for any } x, \\ \arccos \frac{1}{\sqrt{1+x^2}} & \text{if } x \geq 0, \\ -\arccos \frac{1}{\sqrt{1+x^2}} & \text{if } x \leq 0, \\ \operatorname{arccot} \frac{1}{x} & \text{if } x > 0; \end{cases}$$

$$\operatorname{arccot} x = \begin{cases} \arcsin \frac{1}{\sqrt{1+x^2}} & \text{if } x > 0, \\ \pi - \arcsin \frac{1}{\sqrt{1+x^2}} & \text{if } x < 0, \\ \arctan \frac{1}{x} & \text{if } x > 0, \\ \pi + \arctan \frac{1}{x} & \text{if } x < 0. \end{cases}$$

1.3-5. Addition and Subtraction of Inverse Trigonometric Functions.

$$\begin{aligned}\arcsin x + \arcsin y &= \arcsin(x\sqrt{1-y^2} + y\sqrt{1-x^2}) & \text{for } x^2 + y^2 \leq 1, \\ \arccos x \pm \arccos y &= \pm \arccos[xy \mp \sqrt{(1-x^2)(1-y^2)}] & \text{for } x \pm y \geq 0, \\ \arctan x + \arctan y &= \arctan \frac{x+y}{1-xy} & \text{for } xy < 1, \\ \arctan x - \arctan y &= \arctan \frac{x-y}{1+xy} & \text{for } xy > -1.\end{aligned}$$

1.3-6. Differentiation Formulas.

$$\begin{aligned} \frac{d}{dx} \arcsin x &= \frac{1}{\sqrt{1-x^2}}, & \frac{d}{dx} \arccos x &= -\frac{1}{\sqrt{1-x^2}}, \\ \frac{d}{dx} \arctan x &= \frac{1}{1+x^2}, & \frac{d}{dx} \operatorname{arccot} x &= -\frac{1}{1+x^2}. \end{aligned}$$

1.3-7. Integration Formulas.

$$\begin{aligned} \int \arcsin x \, dx &= x \arcsin x + \sqrt{1-x^2} + C, & \int \arccos x \, dx &= x \arccos x - \sqrt{1-x^2} + C, \\ \int \arctan x \, dx &= x \arctan x - \frac{1}{2} \ln(1+x^2) + C, & \int \operatorname{arccot} x \, dx &= x \operatorname{arccot} x + \frac{1}{2} \ln(1+x^2) + C, \end{aligned}$$

where C is an arbitrary constant.

1.3-8. Expansion in Power Series.

$$\begin{aligned} \arcsin x &= x + \frac{1}{2} \frac{x^3}{3} + \frac{1 \times 3}{2 \times 4} \frac{x^5}{5} + \frac{1 \times 3 \times 5}{2 \times 4 \times 6} \frac{x^7}{7} + \cdots + \frac{1 \times 3 \times \cdots \times (2n-1)}{2 \times 4 \times \cdots \times (2n)} \frac{x^{2n+1}}{2n+1} + \cdots \quad (|x| < 1), \\ \arctan x &= x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots + (-1)^{n-1} \frac{x^{2n-1}}{2n-1} + \cdots \quad (|x| \leq 1), \\ \operatorname{arctan} x &= \frac{\pi}{2} - \frac{1}{x} + \frac{1}{3x^3} - \frac{1}{5x^5} + \cdots + (-1)^n \frac{1}{(2n-1)x^{2n-1}} + \cdots \quad (|x| > 1). \end{aligned}$$

The expansions for $\arccos x$ and $\operatorname{arccot} x$ can be obtained from the relations $\arccos x = \frac{\pi}{2} - \arcsin x$ and $\operatorname{arccot} x = \frac{\pi}{2} - \arctan x$.

1.4. Hyperbolic Functions

1.4-1. Definitions of Hyperbolic Functions.

Hyperbolic functions are defined in terms of the exponential functions as follows:

$$\begin{aligned} \sinh x &= \frac{e^x - e^{-x}}{2}, & \cosh x &= \frac{e^x + e^{-x}}{2}, \\ \tanh x &= \frac{e^x - e^{-x}}{e^x + e^{-x}}, & \coth x &= \frac{e^x + e^{-x}}{e^x - e^{-x}}. \end{aligned}$$

1.4-2. Simplest Relations.

$$\begin{aligned} \cosh^2 x - \sinh^2 x &= 1, & \tanh x \coth x &= 1, \\ \sinh(-x) &= -\sinh x, & \cosh(-x) &= \cosh x, \\ \tanh x &= \frac{\sinh x}{\cosh x}, & \coth x &= \frac{\cosh x}{\sinh x}, \\ \tanh(-x) &= -\tanh x, & \coth(-x) &= -\coth x, \\ 1 - \tanh^2 x &= \frac{1}{\cosh^2 x}, & \coth^2 x - 1 &= \frac{1}{\sinh^2 x}. \end{aligned}$$

1.4-3. Relations Between Hyperbolic Functions of Single Argument ($x \geq 0$).

$$\begin{aligned}\sinh x &= \sqrt{\cosh^2 x - 1} = \frac{\tanh x}{\sqrt{1 - \tanh^2 x}} = \frac{1}{\sqrt{\coth^2 x - 1}}, \\ \cosh x &= \sqrt{\sinh^2 x + 1} = \frac{1}{\sqrt{1 - \tanh^2 x}} = \frac{\coth x}{\sqrt{\coth^2 x - 1}}, \\ \tanh x &= \frac{\sinh x}{\sqrt{\sinh^2 x + 1}} = \frac{\sqrt{\cosh^2 x - 1}}{\cosh x} = \frac{1}{\coth x}, \\ \coth x &= \frac{\sqrt{\sinh^2 x + 1}}{\sinh x} = \frac{\cosh x}{\sqrt{\cosh^2 x - 1}} = \frac{1}{\tanh x}.\end{aligned}$$

1.4-4. Addition and Subtraction of Hyperbolic Functions.

$$\begin{aligned}\sinh x + \sinh y &= 2 \sinh\left(\frac{x+y}{2}\right) \cosh\left(\frac{x-y}{2}\right), \\ \sinh x - \sinh y &= 2 \sinh\left(\frac{x-y}{2}\right) \cosh\left(\frac{x+y}{2}\right), \\ \cosh x + \cosh y &= 2 \cosh\left(\frac{x+y}{2}\right) \cosh\left(\frac{x-y}{2}\right), \\ \cosh x - \cosh y &= 2 \sinh\left(\frac{x+y}{2}\right) \sinh\left(\frac{x-y}{2}\right), \\ \sinh^2 x - \sinh^2 y &= \cosh^2 x - \cosh^2 y = \sinh(x+y) \sinh(x-y), \\ \sinh^2 x + \cosh^2 y &= \cosh(x+y) \cosh(x-y), \\ (\cosh x \pm \sinh x)^n &= \cosh(nx) \pm \sinh(nx), \\ \tanh x \pm \tanh y &= \frac{\sinh(x \pm y)}{\cosh x \cosh y}, \quad \coth x \pm \coth y = \pm \frac{\sinh(x \pm y)}{\sinh x \sinh y},\end{aligned}$$

where $n = 0, \pm 1, \pm 2, \dots$

1.4-5. Products of Hyperbolic Functions.

$$\begin{aligned}\sinh x \sinh y &= \frac{1}{2}[\cosh(x+y) - \cosh(x-y)], \\ \cosh x \cosh y &= \frac{1}{2}[\cosh(x+y) + \cosh(x-y)], \\ \sinh x \cosh y &= \frac{1}{2}[\sinh(x+y) + \sinh(x-y)].\end{aligned}$$

1.4-6. Powers of Hyperbolic Functions.

$$\begin{aligned}\cosh^2 x &= \frac{1}{2} \cosh 2x + \frac{1}{2}, & \sinh^2 x &= \frac{1}{2} \cosh 2x - \frac{1}{2}, \\ \cosh^3 x &= \frac{1}{4} \cosh 3x + \frac{3}{4} \cosh x, & \sinh^3 x &= \frac{1}{4} \sinh 3x - \frac{3}{4} \sinh x, \\ \cosh^4 x &= \frac{1}{8} \cosh 4x + \frac{1}{2} \cosh 2x + \frac{3}{8}, & \sinh^4 x &= \frac{1}{8} \cosh 4x - \frac{1}{2} \cosh 2x + \frac{3}{8}, \\ \cosh^5 x &= \frac{1}{16} \cosh 5x + \frac{5}{16} \cosh 3x + \frac{5}{8} \cosh x, & \sinh^5 x &= \frac{1}{16} \sinh 5x - \frac{5}{16} \sinh 3x + \frac{5}{8} \sinh x,\end{aligned}$$

$$\begin{aligned}\cosh^{2n} x &= \frac{1}{2^{2n-1}} \sum_{k=0}^{n-1} C_{2n}^k \cosh[2(n-k)x] + \frac{1}{2^{2n}} C_{2n}^n, \\ \cosh^{2n+1} x &= \frac{1}{2^{2n}} \sum_{k=0}^n C_{2n+1}^k \cosh[(2n-2k+1)x], \\ \sinh^{2n} x &= \frac{1}{2^{2n-1}} \sum_{k=0}^{n-1} (-1)^k C_{2n}^k \cosh[2(n-k)x] + \frac{(-1)^n}{2^{2n}} C_{2n}^n, \\ \sinh^{2n+1} x &= \frac{1}{2^{2n}} \sum_{k=0}^n (-1)^k C_{2n+1}^k \sinh[(2n-2k+1)x].\end{aligned}$$

Here $n = 1, 2, \dots$ and C_m^k are binomial coefficients.

1.4-7. Addition Formulas.

$$\begin{aligned}\sinh(x \pm y) &= \sinh x \cosh y \pm \sinh y \cosh x, & \cosh(x \pm y) &= \cosh x \cosh y \pm \sinh x \sinh y, \\ \tanh(x \pm y) &= \frac{\tanh x \pm \tanh y}{1 \pm \tanh x \tanh y}, & \coth(x \pm y) &= \frac{\coth x \coth y \pm 1}{\coth y \pm \coth x}.\end{aligned}$$

1.4-8. Hyperbolic Functions of Multiple Argument.

$$\begin{aligned}\cosh 2x &= 2 \cosh^2 x - 1, & \sinh 2x &= 2 \sinh x \cosh x, \\ \cosh 3x &= -3 \cosh x + 4 \cosh^3 x, & \sinh 3x &= 3 \sinh x + 4 \sinh^3 x, \\ \cosh 4x &= 1 - 8 \cosh^2 x + 8 \cosh^4 x, & \sinh 4x &= 4 \cosh x (\sinh x + 2 \sinh^3 x), \\ \cosh 5x &= 5 \cosh x - 20 \cosh^3 x + 16 \cosh^5 x, & \sinh 5x &= 5 \sinh x + 20 \sinh^3 x + 16 \sinh^5 x.\end{aligned}$$

$$\begin{aligned}\cosh(nx) &= 2^{n-1} \cosh^n x + \frac{n}{2} \sum_{k=0}^{[n/2]} \frac{(-1)^{k+1}}{k+1} C_{n-k-2}^{k-2} 2^{n-2k-2} (\cosh x)^{n-2k-2}, \\ \sinh(nx) &= \sinh x \sum_{k=0}^{[(n-1)/2]} 2^{n-k-1} C_{n-k-1}^k (\cosh x)^{n-2k-1}.\end{aligned}$$

Here C_m^k are binomial coefficients and $[A]$ stands for the integer part of the number A .

1.4-9. Hyperbolic Functions of Half Argument.

$$\begin{aligned}\sinh \frac{x}{2} &= \operatorname{sign} x \sqrt{\frac{\cosh x - 1}{2}}, & \cosh \frac{x}{2} &= \sqrt{\frac{\cosh x + 1}{2}}, \\ \tanh \frac{x}{2} &= \frac{\sinh x}{\cosh x + 1} = \frac{\cosh x - 1}{\sinh x}, & \coth \frac{x}{2} &= \frac{\sinh x}{\cosh x - 1} = \frac{\cosh x + 1}{\sinh x}.\end{aligned}$$

1.4-10. Differentiation Formulas.

$$\begin{aligned}\frac{d \sinh x}{dx} &= \cosh x, & \frac{d \cosh x}{dx} &= \sinh x, \\ \frac{d \tanh x}{dx} &= \frac{1}{\cosh^2 x}, & \frac{d \coth x}{dx} &= -\frac{1}{\sinh^2 x}.\end{aligned}$$

1.4-11. Integration Formulas.

$$\begin{aligned}\int \sinh x \, dx &= \cosh x + C, & \int \cosh x \, dx &= \sinh x + C, \\ \int \tanh x \, dx &= \ln \cosh x + C, & \int \coth x \, dx &= \ln |\sinh x| + C,\end{aligned}$$

where C is an arbitrary constant.

1.4-12. Expansion in Power Series.

$$\begin{aligned}\cosh x &= 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \cdots + \frac{x^{2n}}{(2n)!} + \cdots & (|x| < \infty), \\ \sinh x &= x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \cdots + \frac{x^{2n+1}}{(2n+1)!} + \cdots & (|x| < \infty), \\ \tanh x &= x - \frac{x^3}{3} + \frac{2x^5}{15} - \frac{17x^7}{315} + \cdots + (-1)^{n-1} \frac{2^{2n}(2^{2n}-1)|B_{2n}|x^{2n-1}}{(2n)!} + \cdots & (|x| < \pi/2), \\ \coth x &= \frac{1}{x} + \frac{x}{3} - \frac{x^3}{45} + \frac{2x^5}{945} - \cdots + (-1)^{n-1} \frac{2^{2n}|B_{2n}|x^{2n-1}}{(2n)!} + \cdots & (|x| < \pi), \\ \frac{1}{\cosh x} &= 1 - \frac{x^2}{2} + \frac{5x^4}{24} - \frac{61x^6}{720} + \cdots + \frac{E_{2n}}{(2n)!} x^{2n} + \cdots & (|x| < \pi/2), \\ \frac{1}{\sinh x} &= \frac{1}{x} - \frac{x}{6} + \frac{7x^3}{360} - \frac{31x^5}{15120} + \cdots + \frac{2(2^{2n-1}-1)B_{2n}}{(2n)!} x^{2n-1} + \cdots & (0 < |x| < \pi),\end{aligned}$$

where B_n and E_n are Bernoulli and Euler numbers (see Supplements 11.1-3 and 11.1-4).

1.4-13. Representation in the Form of Infinite Products.

$$\begin{aligned}\sinh x &= x \left(1 + \frac{x^2}{\pi^2}\right) \left(1 + \frac{x^2}{4\pi^2}\right) \left(1 + \frac{x^2}{9\pi^2}\right) \cdots \left(1 + \frac{x^2}{n^2\pi^2}\right) \cdots \\ \cosh x &= \left(1 + \frac{4x^2}{\pi^2}\right) \left(1 + \frac{4x^2}{9\pi^2}\right) \left(1 + \frac{4x^2}{25\pi^2}\right) \cdots \left(1 + \frac{4x^2}{(2n+1)^2\pi^2}\right) \cdots\end{aligned}$$

1.4-14. Relationship with Trigonometric Functions.

$$\sinh(ix) = i \sin x, \quad \cosh(ix) = \cos x, \quad \tanh(ix) = i \tan x, \quad \coth(ix) = -i \cot x, \quad i^2 = -1.$$

1.5. Inverse Hyperbolic Functions

1.5-1. Definitions of Inverse Hyperbolic Functions.

Inverse hyperbolic functions are the functions that are inverse to hyperbolic functions. The following notation is used for inverse hyperbolic functions:

$$\operatorname{arsinh} x \equiv \sinh^{-1} x \quad (\text{inverse of hyperbolic sine}),$$

$$\operatorname{arcosh} x \equiv \cosh^{-1} x \quad (\text{inverse of hyperbolic cosine}),$$

$$\operatorname{artanh} x \equiv \tanh^{-1} x \quad (\text{inverse of hyperbolic tangent}),$$

$$\operatorname{arcoth} x \equiv \coth^{-1} x \quad (\text{inverse of hyperbolic cotangent}).$$

Inverse hyperbolic functions can be expressed in terms of logarithmic functions:

$$\operatorname{arsinh} x = \ln(x + \sqrt{x^2 + 1}) \quad (x \text{ is any}); \quad \operatorname{arcosh} x = \ln(x + \sqrt{x^2 - 1}) \quad (x \geq 1);$$

$$\operatorname{artanh} x = \frac{1}{2} \ln \frac{1+x}{1-x} \quad (|x| < 1); \quad \operatorname{arcoth} x = \frac{1}{2} \ln \frac{x+1}{x-1} \quad (|x| > 1).$$

Here only one (principal) branch of the function $\operatorname{arcosh} x$ is listed, the function itself being double-valued. In order to write out both branches of $\operatorname{arcosh} x$, the symbol \pm should be placed before the logarithm on the right-hand side of the formula.

1.5-2. Simplest Relations.

$$\operatorname{arsinh}(-x) = -\operatorname{arsinh} x, \quad \operatorname{artanh}(-x) = -\operatorname{artanh} x, \quad \operatorname{arcoth}(-x) = -\operatorname{arcoth} x.$$

1.5-3. Relations Between Inverse Hyperbolic Functions.

$$\operatorname{arsinh} x = \operatorname{arcosh} \sqrt{x^2 + 1} = \operatorname{artanh} \frac{x}{\sqrt{x^2 + 1}},$$

$$\operatorname{arcosh} x = \operatorname{arsinh} \sqrt{x^2 - 1} = \operatorname{artanh} \frac{\sqrt{x^2 - 1}}{x},$$

$$\operatorname{artanh} x = \operatorname{arsinh} \frac{x}{\sqrt{1-x^2}} = \operatorname{arcosh} \frac{1}{\sqrt{1-x^2}} = \operatorname{arcoth} \frac{1}{x}.$$

1.5-4. Addition and Subtraction of Inverse Hyperbolic Functions.

$$\operatorname{arsinh} x \pm \operatorname{arsinh} y = \operatorname{arsinh} (x\sqrt{1+y^2} \pm y\sqrt{1+x^2}),$$

$$\operatorname{arcosh} x \pm \operatorname{arcosh} y = \operatorname{arcosh} [xy \pm \sqrt{(x^2-1)(y^2-1)}],$$

$$\operatorname{arsinh} x \pm \operatorname{arcosh} y = \operatorname{arsinh} [xy \pm \sqrt{(x^2+1)(y^2-1)}],$$

$$\operatorname{artanh} x \pm \operatorname{artanh} y = \operatorname{artanh} \frac{x \pm y}{1 \pm xy}, \quad \operatorname{artanh} x \pm \operatorname{arcoth} y = \operatorname{artanh} \frac{xy \pm 1}{y \pm x}.$$

1.5-5. Differentiation Formulas.

$$\frac{d}{dx} \operatorname{arsinh} x = \frac{1}{\sqrt{x^2 + 1}},$$

$$\frac{d}{dx} \operatorname{arcosh} x = \frac{1}{\sqrt{x^2 - 1}},$$

$$\frac{d}{dx} \operatorname{artanh} x = \frac{1}{1-x^2} \quad (x^2 < 1),$$

$$\frac{d}{dx} \operatorname{arcoth} x = \frac{1}{1-x^2} \quad (x^2 > 1).$$

1.5-6. Integration Formulas.

$$\int \operatorname{arcsinh} x \, dx = x \operatorname{arcsinh} x - \sqrt{1+x^2} + C,$$

$$\int \operatorname{arcosh} x \, dx = x \operatorname{arcosh} x - \sqrt{x^2-1} + C,$$

$$\int \operatorname{arctanh} x \, dx = x \operatorname{arctanh} x + \frac{1}{2} \ln(1-x^2) + C,$$

$$\int \operatorname{arcoth} x \, dx = x \operatorname{arcoth} x + \frac{1}{2} \ln(x^2-1) + C,$$

where C is an arbitrary constant.

1.5-7. Expansion in Power Series.

$$\operatorname{arcsinh} x = x - \frac{1}{2} \frac{x^3}{3} + \frac{1 \times 3}{2 \times 4} \frac{x^5}{5} - \dots + (-1)^n \frac{1 \times 3 \times \dots \times (2n-1)}{2 \times 4 \times \dots \times (2n)} \frac{x^{2n+1}}{2n+1} + \dots \quad (|x| < 1),$$

$$\operatorname{arcsinh} x = \ln(2x) + \frac{1}{2} \frac{1}{2x^2} + \frac{1 \times 3}{2 \times 4} \frac{1}{4x^4} + \dots + \frac{1 \times 3 \times \dots \times (2n-1)}{2 \times 4 \times \dots \times (2n)} \frac{1}{2nx^{2n}} + \dots \quad (|x| > 1),$$

$$\operatorname{arcosh} x = \ln(2x) - \frac{1}{2} \frac{1}{2x^2} - \frac{1 \times 3}{2 \times 4} \frac{1}{4x^4} - \dots - \frac{1 \times 3 \times \dots \times (2n-1)}{2 \times 4 \times \dots \times (2n)} \frac{1}{2nx^{2n}} - \dots \quad (|x| > 1),$$

$$\operatorname{arctanh} x = x + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^7}{7} + \dots + \frac{x^{2n+1}}{2n+1} + \dots \quad (|x| < 1),$$

$$\operatorname{arcoth} x = \frac{1}{x} + \frac{1}{3x^3} + \frac{1}{5x^5} + \frac{1}{7x^7} + \dots + \frac{1}{(2n+1)x^{2n+1}} + \dots \quad (|x| > 1).$$

© References for Supplement 1: M. Abramowitz and I. A. Stegun (1964), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1986), D. G. Zill and J. M. Dewar (1990), M. Kline (1998), R. Courant and F. John (1999), I. S. Gradshteyn and I. M. Ryzhik (2000), G. A. Korn and T. M. Korn (2000), C. H. Edwards and D. Penney (2002), D. Zwillinger (2002), E. W. Weisstein (2003), I. N. Bronshtein and K. A. Semendyayev (2004), M. Sullivan (2004), H. Anton, I. Bivens, and S. Davis (2005), R. Adams (2006).

Supplement 2

Finite Sums and Infinite Series

2.1. Finite Numerical Sums

2.1-1. Progressions.

Arithmetic progression:

$$1. \sum_{k=0}^{n-1} (a + bk) = an + \frac{bn(n-1)}{2}.$$

Geometric progression:

$$2. \sum_{k=1}^n aq^{k-1} = a \frac{q^n - 1}{q - 1}.$$

Arithmetic-geometric progression:

$$3. \sum_{k=0}^{n-1} (a + bk)q^k = \frac{a(1 - q^n) - b(n-1)q^n}{1 - q} + \frac{bq(1 - q^{n-1})}{(1 - q)^2}.$$

2.1-2. Sums of Powers of Natural Numbers Having the Form $\sum k^m$.

$$1. \sum_{k=1}^n k = \frac{n(n+1)}{2}.$$

$$2. \sum_{k=1}^n k^2 = \frac{1}{6}n(n+1)(2n+1).$$

$$3. \sum_{k=1}^n k^3 = \frac{1}{4}n^2(n+1)^2.$$

$$4. \sum_{k=1}^n k^4 = \frac{1}{30}n(n+1)(2n+1)(3n^2+3n-1).$$

$$5. \sum_{k=1}^n k^5 = \frac{1}{12}n^2(n+1)^2(2n^2+2n-1).$$

$$6. \sum_{k=1}^n k^m = \frac{n^{m+1}}{m+1} + \frac{n^m}{2} + \frac{1}{2}C_m^1 B_2 n^{m-1} + \frac{1}{4}C_m^3 B_4 n^{m-3} + \frac{1}{6}C_m^5 B_6 n^{m-5} + \dots.$$

Here the C_m^k are binomial coefficients and the B_{2k} are Bernoulli numbers (see Supplement 11.1-3); the last term in the sum contains n or n^2 .

2.1-3. Alternating Sums of Powers of Natural Numbers, $\sum (-1)^k k^m$.

1. $\sum_{k=1}^n (-1)^k k = (-1)^n \left[\frac{n-1}{2} \right]$; $[m]$ stands for the integer part of m .
2. $\sum_{k=1}^n (-1)^k k^2 = (-1)^n \frac{n(n+1)}{2}$.
3. $\sum_{k=1}^n (-1)^k k^3 = \frac{1}{8} [1 + (-1)^n (4n^3 + 6n^2 - 1)]$.
4. $\sum_{k=1}^n (-1)^k k^4 = (-1)^n \frac{1}{2} (n^4 + 2n^3 - n)$.
5. $\sum_{k=1}^n (-1)^k k^5 = \frac{1}{4} [-1 + (-1)^n (2n^5 + 5n^4 - 5n^2 + 1)]$.

2.1-4. Other Sums Containing Integers.

1. $\sum_{k=0}^n (2k+1) = (n+1)^2$.
2. $\sum_{k=0}^n (2k+1)^2 = \frac{1}{3} (n+1)(2n+1)(2n+3)$.
3. $\sum_{k=1}^n k(k+1) = \frac{1}{3} n(n+1)(n+2)$.
4. $\sum_{k=1}^n (k+a)(k+b) = \frac{1}{6} n(n+1)(2n+1+3a+3b) + nab$.
5. $\sum_{k=1}^n k k! = (n+1)! - 1$.
6. $\sum_{k=0}^n (-1)^k (2k+1) = (-1)^n (n+1)$.
7. $\sum_{k=0}^n (-1)^k (2k+1)^2 = 2(-1)^n (n+1)^2 - \frac{1}{2} [1 + (-1)^n]$.

2.1-5. Sums Containing Binomial Coefficients.

► Throughout Supplement 2.1-5, it is assumed that $m = 1, 2, 3, \dots$

1. $\sum_{k=0}^n C_n^k = 2^n$.
2. $\sum_{k=0}^n C_{m+k}^m = C_{n+m+1}^{m+1}$.
3. $\sum_{k=0}^n (-1)^k C_m^k = (-1)^n C_{m-1}^n$.

4.
$$\sum_{k=0}^n (k+1)C_n^k = 2^{n-1}(n+2).$$
5.
$$\sum_{k=1}^n (-1)^{k+1} k C_n^k = 0.$$
6.
$$\sum_{k=1}^n \frac{(-1)^{k+1}}{k} C_n^k = \sum_{m=1}^n \frac{1}{m}.$$
7.
$$\sum_{k=1}^n \frac{(-1)^{k+1}}{k+1} C_n^k = \frac{n}{n+1}.$$
8.
$$\sum_{k=0}^n \frac{1}{k+1} C_n^k = \frac{2^{n+1}-1}{n+1}.$$
9.
$$\sum_{k=0}^n \frac{a^{k+1}}{k+1} C_n^k = \frac{(a+1)^{n+1}-1}{n+1}.$$
10.
$$\sum_{k=0}^p C_n^k C_m^{p-k} = C_{n+m}^p; \quad m \text{ and } p \text{ are natural numbers.}$$
11.
$$\sum_{k=0}^{n-p} C_n^k C_n^{p+k} = \frac{(2n)!}{(n-p)!(n+p)!}.$$
12.
$$\sum_{k=0}^n (C_n^k)^2 = C_{2n}^n.$$
13.
$$\sum_{k=0}^{2n} (-1)^k (C_{2n}^k)^2 = (-1)^n C_{2n}^n.$$
14.
$$\sum_{k=0}^{2n+1} (-1)^k (C_{2n+1}^k)^2 = 0.$$
15.
$$\sum_{k=1}^n k (C_n^k)^2 = \frac{(2n-1)!}{[(n-1)!]^2}.$$

2.1-6. Other Numerical Sums.

1.
$$\sum_{k=1}^{n-1} \sin \frac{\pi k}{n} = \cot \frac{\pi}{2n}.$$
2.
$$\sum_{k=1}^n \sin^{2m} \frac{\pi k}{2n} = \frac{n}{2^{2m}} C_{2m}^m + \frac{1}{2}, \quad m < 2n.$$
3.
$$\sum_{k=0}^{n-1} (-1)^k \cos^m \frac{\pi k}{n} = \frac{1}{2} [1 - (-1)^{m+n}], \quad m = 0, 1, \dots, n-1.$$
4.
$$\sum_{k=0}^{n-1} (-1)^k \cos^n \frac{\pi k}{n} = \frac{n}{2^{n-1}}.$$

2.2. Finite Functional Sums

2.2-1. Sums Involving Hyperbolic Functions.

1.
$$\sum_{k=0}^{n-1} \sinh(kx + a) = \sinh\left(\frac{n-1}{2}x + a\right) \frac{\sinh(nx/2)}{\sinh(x/2)}.$$
2.
$$\sum_{k=0}^{n-1} \cosh(kx + a) = \cosh\left(\frac{n-1}{2}x + a\right) \frac{\sinh(nx/2)}{\sinh(x/2)}.$$
3.
$$\sum_{k=0}^{n-1} (-1)^k \sinh(kx + a) = \frac{1}{2 \cosh(x/2)} \left[\sinh\left(a - \frac{x}{2}\right) + (-1)^n \sinh\left(\frac{2n-1}{2}x + a\right) \right].$$
4.
$$\sum_{k=0}^{n-1} (-1)^k \cosh(kx + a) = \frac{1}{2 \cosh(x/2)} \left[\cosh\left(a - \frac{x}{2}\right) + (-1)^n \cosh\left(\frac{2n-1}{2}x + a\right) \right].$$
5.
$$\sum_{k=1}^{n-1} k \sinh(kx + a) = -\frac{1}{\sinh^2(x/2)} \left\{ n \sinh[(n-1)x + a] - (n-1) \sinh(nx + a) - \sinh a \right\}.$$
6.
$$\sum_{k=1}^{n-1} k \cosh(kx + a) = -\frac{1}{\sinh^2(x/2)} \left\{ n \cosh[(n-1)x + a] - (n-1) \cosh(nx + a) - \cosh a \right\}.$$
7.
$$\sum_{k=1}^{n-1} (-1)^k k \sinh(kx + a) = \frac{1}{\cosh^2(x/2)} \left\{ (-1)^{n-1} n \sinh[(n-1)x + a] \right. \\ \left. + (-1)^{n-1} (n-1) \sinh(nx + a) - \sinh a \right\}.$$
8.
$$\sum_{k=1}^{n-1} (-1)^k k \cosh(kx + a) = \frac{1}{\cosh^2(x/2)} \left\{ (-1)^{n-1} n \cosh[(n-1)x + a] \right. \\ \left. + (-1)^{n-1} (n-1) \cosh(nx + a) - \cosh a \right\}.$$
9.
$$\sum_{k=0}^n C_n^k \sinh(kx + a) = 2^n \cosh^n \frac{x}{2} \sinh\left(\frac{nx}{2} + a\right).$$
10.
$$\sum_{k=0}^n C_n^k \cosh(kx + a) = 2^n \cosh^n \frac{x}{2} \cosh\left(\frac{nx}{2} + a\right).$$
11.
$$\sum_{k=1}^{n-1} a^k \sinh(kx) = \frac{a \sinh x - a^n \sinh(nx) + a^{n+1} \sinh[(n-1)x]}{1 - 2a \cosh x + a^2}.$$
12.
$$\sum_{k=0}^{n-1} a^k \cosh(kx) = \frac{1 - a \cosh x - a^n \cosh(nx) + a^{n+1} \cosh[(n-1)x]}{1 - 2a \cosh x + a^2}.$$
13.
$$\sum_{k=1}^n \frac{1}{2^k} \tanh \frac{x}{2^k} = \coth x - \frac{1}{2^n} \coth \frac{x}{2^n}.$$
14.
$$\sum_{k=0}^{n-1} 2^k \tanh(2^k x) = 2^n \coth(2^n x) - \coth x.$$

2.2-2. Sums Involving Trigonometric Functions.

1.
$$\sum_{k=1}^n \sin(2kx) = \sin[(n+1)x] \sin(nx) \operatorname{cosec} x.$$

2.
$$\sum_{k=0}^n \cos(2kx) = \sin[(n+1)x] \cos(nx) \operatorname{cosec} x.$$
3.
$$\sum_{k=1}^n \sin[(2k-1)x] = \sin^2(nx) \operatorname{cosec} x.$$
4.
$$\sum_{k=1}^n \cos[(2k-1)x] = \sin(nx) \cos(nx) \operatorname{cosec} x.$$
5.
$$\sum_{k=0}^{n-1} \sin(kx+a) = \sin\left(\frac{n-1}{2}x+a\right) \sin \frac{nx}{2} \operatorname{cosec} \frac{x}{2}.$$
6.
$$\sum_{k=0}^{n-1} \cos(kx+a) = \cos\left(\frac{n-1}{2}x+a\right) \sin \frac{nx}{2} \operatorname{cosec} \frac{x}{2}.$$
7.
$$\sum_{k=0}^{2n-1} (-1)^k \cos(kx+a) = \sin\left(\frac{2n-1}{2}x+a\right) \sin(nx) \sec \frac{x}{2}.$$
8.
$$\sum_{k=1}^n (-1)^{k+1} \sin[(2k-1)x] = (-1)^{n+1} \frac{\sin(2nx)}{2 \cos x}.$$
9.
$$\sum_{k=1}^n (-1)^k \cos(2kx) = -\frac{1}{2} + (-1)^n \frac{\cos[(2n+1)x]}{2 \cos x}.$$
10.
$$\sum_{k=1}^n \sin^2(kx) = \frac{n}{2} - \frac{\cos[(n+1)x] \sin(nx)}{2 \sin x}.$$
11.
$$\sum_{k=1}^n \cos^2(kx) = \frac{n}{2} + \frac{\cos[(n+1)x] \sin(nx)}{2 \sin x}.$$
12.
$$\sum_{k=1}^{n-1} k \sin(2kx) = \frac{\sin(2nx)}{4 \sin^2 x} - \frac{n \cos[(2n-1)x]}{2 \sin x}.$$
13.
$$\sum_{k=1}^{n-1} k \cos(2kx) = \frac{n \sin[(2n-1)x]}{2 \sin x} - \frac{1 - \cos(2nx)}{4 \sin^2 x}.$$
14.
$$\sum_{k=1}^{n-1} a^k \sin(kx) = \frac{a \sin x - a^n \sin(nx) + a^{n+1} \sin[(n-1)x]}{1 - 2a \cos x + a^2}.$$
15.
$$\sum_{k=0}^{n-1} a^k \cos(kx) = \frac{1 - a \cos x - a^n \cos(nx) + a^{n+1} \cos[(n-1)x]}{1 - 2a \cos x + a^2}.$$
16.
$$\sum_{k=0}^n C_n^k \sin(kx+a) = 2^n \cos^n \frac{x}{2} \sin\left(\frac{nx}{2} + a\right).$$
17.
$$\sum_{k=0}^n C_n^k \cos(kx+a) = 2^n \cos^n \frac{x}{2} \cos\left(\frac{nx}{2} + a\right).$$
18.
$$\sum_{k=0}^n (-1)^k C_n^k \sin(kx+a) = (-2)^n \sin^n \frac{x}{2} \sin\left(\frac{nx}{2} + \frac{\pi n}{2} + a\right).$$
19.
$$\sum_{k=0}^n (-1)^k C_n^k \cos(kx+a) = (-2)^n \sin^n \frac{x}{2} \cos\left(\frac{nx}{2} + \frac{\pi n}{2} + a\right).$$

20.
$$\sum_{k=1}^n \left(2^k \sin^2 \frac{x}{2^k} \right)^2 = \left(2^n \sin^2 \frac{x}{2^n} \right)^2 - \sin^2 x.$$
21.
$$\sum_{k=0}^n \frac{1}{2^k} \tan \frac{x}{2^k} = \frac{1}{2^n} \cot \frac{x}{2^n} - 2 \cot(2x).$$

2.3. Infinite Numerical Series

2.3-1. Progressions.

1.
$$\sum_{k=0}^{\infty} aq^k = \frac{a}{1-q}, \quad |q| < 1.$$
2.
$$\sum_{k=0}^{\infty} (a + bk)q^k = \frac{a}{1-q} + \frac{bq}{(1-q)^2}, \quad |q| < 1.$$

2.3-2. Other Numerical Series.

1.
$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n+1} = \ln 2.$$
2.
$$\sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} = \frac{\pi}{4}.$$
3.
$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1.$$
4.
$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n(n+1)} = 1 - 2 \ln 2.$$
5.
$$\sum_{n=1}^{\infty} \frac{1}{n(n+2)} = \frac{3}{4}.$$
6.
$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n(n+2)} = -\frac{1}{4}.$$
7.
$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)(2n+1)} = \frac{1}{2}.$$
8.
$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}.$$
9.
$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2} = \frac{\pi^2}{12}.$$
10.
$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}.$$
11.
$$\sum_{n=1}^{\infty} \frac{1}{n^2 + a^2} = \frac{\pi}{2a} \coth(\pi a) - \frac{1}{2a^2}.$$

12.
$$\sum_{n=1}^{\infty} \frac{1}{n^2 - a^2} = -\frac{\pi}{2a} \cot(\pi a) + \frac{1}{2a^2}.$$
13.
$$\sum_{k=1}^{\infty} \frac{1}{k^{2n}} = \frac{2^{2n-1} \pi^{2n}}{(2n)!} |B_{2n}|; \quad \text{the } B_{2n} \text{ are Bernoulli numbers (see Supplement 11.1-3).}$$
14.
$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2n}} = \frac{(2^{2n-1} - 1) \pi^{2n}}{(2n)!} |B_{2n}|; \quad \text{the } B_{2n} \text{ are Bernoulli numbers.}$$
15.
$$\sum_{k=1}^{\infty} \frac{1}{(2k-1)^{2n}} = \frac{(2^{2n-1} - 1) \pi^{2n}}{2(2n)!} |B_{2n}|; \quad \text{the } B_{2n} \text{ are Bernoulli numbers.}$$
16.
$$\sum_{k=1}^{\infty} \frac{1}{k2^k} = \ln 2.$$
17.
$$\sum_{k=0}^{\infty} \frac{(-1)^k}{n^{2k}} = \frac{n^2}{n^2 + 1}.$$
18.
$$\sum_{k=0}^{\infty} \frac{1}{k!} = e = 2.71828 \dots$$
19.
$$\sum_{k=0}^{\infty} \frac{(-1)^k}{k!} = \frac{1}{e} = 0.36787 \dots$$
20.
$$\sum_{k=1}^{\infty} \frac{k}{(k+1)!} = 1.$$

2.4. Infinite Functional Series

2.4-1. Power Series.

1.
$$\sum_{k=0}^{\infty} x^k = \frac{1}{1-x}, \quad |x| < 1.$$
2.
$$\sum_{k=1}^{\infty} kx^k = \frac{x}{(1-x)^2}, \quad |x| < 1.$$
3.
$$\sum_{k=1}^{\infty} k^2 x^k = \frac{x(x+1)}{(1-x)^3}, \quad |x| < 1.$$
4.
$$\sum_{k=1}^{\infty} k^3 x^k = \frac{x(1+4x+x^2)}{(1-x)^4}, \quad |x| < 1.$$
5.
$$\sum_{k=0}^{\infty} (\pm 1)^k k^n x^k = \left(x \frac{d}{dx} \right)^n \frac{1}{1 \mp x}, \quad |x| < 1.$$
6.
$$\sum_{k=1}^{\infty} \frac{x^k}{k} = -\ln(1-x), \quad -1 \leq x < 1.$$
7.
$$\sum_{k=1}^{\infty} (-1)^{k-1} \frac{x^k}{k} = \ln(1+x), \quad |x| < 1.$$
8.
$$\sum_{k=1}^{\infty} \frac{x^{2k-1}}{2k-1} = \frac{1}{2} \ln \frac{1+x}{1-x}, \quad |x| < 1.$$

9.
$$\sum_{k=1}^{\infty} (-1)^{k-1} \frac{x^{2k-1}}{2k-1} = \arctan x, \quad |x| \leq 1.$$
10.
$$\sum_{k=1}^{\infty} \frac{x^k}{k^2} = - \int_0^x \frac{\ln(1-t)}{t} dt, \quad |x| \leq 1.$$
11.
$$\sum_{k=1}^{\infty} \frac{x^{k+1}}{k(k+1)} = x + (1-x) \ln(1-x), \quad |x| \leq 1.$$
12.
$$\sum_{k=1}^{\infty} \frac{x^{k+2}}{k(k+2)} = \frac{x}{2} + \frac{x^2}{4} + \frac{1}{2}(1-x^2) \ln(1-x), \quad |x| \leq 1.$$
13.
$$\sum_{k=0}^{\infty} \frac{x^k}{k!} = e^x, \quad x \text{ is any number.}$$
14.
$$\sum_{k=0}^{\infty} \frac{x^{2k}}{(2k)!} = \cosh x, \quad x \text{ is any number.}$$
15.
$$\sum_{k=0}^{\infty} (-1)^k \frac{x^{2k}}{(2k)!} = \cos x, \quad x \text{ is any number.}$$
16.
$$\sum_{k=0}^{\infty} \frac{x^{2k+1}}{(2k+1)!} = \sinh x, \quad x \text{ is any number.}$$
17.
$$\sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{(2k+1)!} = \sin x, \quad x \text{ is any number.}$$
18.
$$\sum_{k=0}^{\infty} \frac{x^{k+1}}{k!(k+1)} = e^x - 1, \quad x \text{ is any number.}$$
19.
$$\sum_{k=0}^{\infty} \frac{x^{k+2}}{k!(k+2)} = (x-1)e^x + 1, \quad x \text{ is any number.}$$
20.
$$\sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{k!(2k+1)} = \frac{\sqrt{\pi}}{2} \operatorname{erf} x, \quad x \text{ is any number.}$$
21.
$$\sum_{k=0}^{\infty} \frac{(k+a)^n}{k!} x^k = \left[\frac{d^n}{dt^n} \exp(at + xe^t) \right]_{t=0}, \quad x \text{ is any number.}$$
22.
$$\sum_{k=1}^{\infty} \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} x^{2k-1} = \tan x; \quad \text{the } B_{2k} \text{ are Bernoulli numbers, } |x| < \pi/2.$$
23.
$$\sum_{k=1}^{\infty} (-1)^{k-1} \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} x^{2k-1} = \tanh x; \quad \text{the } B_{2k} \text{ are Bernoulli numbers, } |x| < \pi/2.$$
24.
$$\sum_{k=1}^{\infty} \frac{2^{2k}|B_{2k}|}{(2k)!} x^{2k-1} = \frac{1}{x} - \cot x; \quad \text{the } B_{2k} \text{ are Bernoulli numbers, } 0 < |x| < \pi.$$
25.
$$\sum_{k=1}^{\infty} (-1)^{k-1} \frac{2^{2k}|B_{2k}|}{(2k)!} x^{2k-1} = \coth x - \frac{1}{x}; \quad \text{the } B_{2k} \text{ are Bernoulli numbers, } |x| < \pi.$$

2.4-2. Trigonometric Series in One Variable Involving Sine.

1.
$$\sum_{k=1}^{\infty} \frac{1}{k} \sin(kx) = \frac{1}{2}(\pi - x), \quad 0 < x < 2\pi.$$
2.
$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \sin(kx) = \frac{1}{2}x, \quad -\pi < x < \pi.$$
3.
$$\sum_{k=1}^{\infty} \frac{a^k}{k} \sin(kx) = \arctan \frac{a \sin x}{1 - a \cos x}, \quad 0 < x < 2\pi, |a| \leq 1.$$
4.
$$\sum_{k=0}^{\infty} \frac{1}{2k+1} \sin(kx) = \frac{\pi}{4} \cos \frac{x}{2} - \sin \frac{x}{2} \ln \left(\cot^2 \frac{x}{4} \right), \quad 0 < x < 2\pi.$$
5.
$$\sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} \sin(kx) = -\frac{1}{4} \cos \frac{x}{2} \ln \left(\cot^2 \frac{x+\pi}{4} \right) - \frac{\pi}{4} \sin \frac{x}{2}, \quad -\pi < x < \pi.$$
6.
$$\sum_{k=1}^{\infty} \frac{1}{k^2} \sin(kx) = -\int_0^x \ln \left(2 \sin \frac{t}{2} \right) dt, \quad 0 \leq x < \pi.$$
7.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k^2} \sin(kx) = -\int_0^x \ln \left(2 \cos \frac{t}{2} \right) dt, \quad -\pi < x < \pi.$$
8.
$$\sum_{k=1}^{\infty} \frac{1}{k(k+1)} \sin(kx) = (\pi - x) \sin^2 \frac{x}{2} + \sin x \ln \left(2 \sin \frac{x}{2} \right), \quad 0 \leq x \leq 2\pi.$$
9.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k(k+1)} \sin(kx) = -x \cos^2 \frac{x}{2} + \sin x \ln \left(2 \cos \frac{x}{2} \right), \quad -\pi \leq x \leq \pi.$$
10.
$$\sum_{k=1}^{\infty} \frac{k}{k^2 + a^2} \sin(kx) = \frac{\pi}{2 \sinh(\pi a)} \sinh[a(\pi - x)], \quad 0 < x < 2\pi.$$
11.
$$\sum_{k=1}^{\infty} (-1)^{k+1} \frac{k}{k^2 + a^2} \sin(kx) = \frac{\pi}{2 \sinh(\pi a)} \sinh(ax), \quad -\pi < x < \pi.$$
12.
$$\sum_{k=1}^{\infty} \frac{k}{k^2 - a^2} \sin(kx) = \frac{\pi}{2 \sin(\pi a)} \sin[a(\pi - x)], \quad 0 < x < 2\pi.$$
13.
$$\sum_{k=1}^{\infty} (-1)^{k+1} \frac{k}{k^2 - a^2} \sin(kx) = \frac{\pi}{2 \sin(\pi a)} \sin(ax), \quad -\pi < x < \pi.$$
14.
$$\sum_{k=2}^{\infty} (-1)^k \frac{k}{k^2 - 1} \sin(kx) = \frac{1}{4} \sin x + \frac{1}{2} x \cos x, \quad -\pi < x < \pi.$$
15.
$$\sum_{k=1}^{\infty} \frac{1}{k^{2n+1}} \sin(kx) = \frac{(-1)^{n-1} (2\pi)^{2n+1}}{2(2n+1)!} B_{2n+1} \left(\frac{x}{2\pi} \right),$$
 where $0 \leq x \leq 2\pi$ for $n = 1, 2, \dots$;
 $0 < x < 2\pi$ for $n = 0$; and the $B_n(x)$ are Bernoulli polynomials (see Supplement 11.18-1).
16.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k^{2n+1}} \sin(kx) = \frac{(-1)^{n-1} (2\pi)^{2n+1}}{2(2n+1)!} B_{2n+1} \left(\frac{x+\pi}{2\pi} \right),$$
 where $-\pi < x \leq \pi$ for $n = 0, 1, \dots$;
the $B_n(x)$ are Bernoulli polynomials.
17.
$$\sum_{k=1}^{\infty} \frac{1}{k!} \sin(kx) = \exp(\cos x) \sin(\sin x), \quad x \text{ is any number.}$$

18.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k!} \sin(kx) = -\exp(-\cos x) \sin(\sin x), \quad x \text{ is any number.}$$
19.
$$\sum_{k=0}^{\infty} \frac{1}{(2k)!} \sin(kx) = \sin\left(\sin \frac{x}{2}\right) \sinh\left(\cos \frac{x}{2}\right), \quad x \text{ is any number.}$$
20.
$$\sum_{k=0}^{\infty} \frac{(-1)^k}{(2k)!} \sin(kx) = -\sin\left(\cos \frac{x}{2}\right) \sinh\left(\sin \frac{x}{2}\right), \quad x \text{ is any number.}$$
21.
$$\sum_{k=0}^{\infty} \frac{a^k}{k!} \sin(kx) = \exp(k \cos x) \sin(k \sin x), \quad |a| \leq 1, \quad x \text{ is any number.}$$
22.
$$\sum_{k=0}^{\infty} a^k \sin(kx) = \frac{a \sin x}{1 - 2a \cos x + a^2}, \quad |a| < 1, \quad x \text{ is any number.}$$
23.
$$\sum_{k=1}^{\infty} k a^k \sin(kx) = \frac{a(1 - a^2) \sin x}{(1 - 2a \cos x + a^2)^2}, \quad |a| < 1, \quad x \text{ is any number.}$$
24.
$$\sum_{k=1}^{\infty} \frac{1}{k} \sin(kx + a) = \frac{1}{2}(\pi - x) \cos a - \ln\left(2 \sin \frac{x}{2}\right) \sin a, \quad 0 < x < 2\pi.$$
25.
$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \sin(kx + a) = \frac{1}{2}x \cos a + \ln\left(2 \cos \frac{x}{2}\right) \sin a, \quad -\pi < x < \pi.$$
26.
$$\sum_{k=1}^{\infty} \frac{\sin[(2k-1)x]}{2k-1} = \frac{\pi}{4}, \quad 0 < x < \pi.$$
27.
$$\sum_{k=1}^{\infty} (-1)^{k-1} \frac{\sin[(2k-1)x]}{2k-1} = \frac{1}{2} \ln \tan\left(\frac{x}{2} + \frac{\pi}{4}\right), \quad -\frac{\pi}{2} < x < \frac{\pi}{2}.$$
28.
$$\sum_{k=1}^{\infty} a^{2k-1} \frac{\sin[(2k-1)x]}{2k-1} = \frac{1}{2} \arctan \frac{2a \sin x}{1 - a^2}, \quad 0 < x < 2\pi, \quad |a| \leq 1.$$
29.
$$\sum_{k=1}^{\infty} (-1)^{k-1} a^{2k-1} \frac{\sin[(2k-1)x]}{2k-1} = \frac{1}{4} \ln \frac{1 + 2a \sin x + a^2}{1 - 2a \sin x + a^2}, \quad 0 < x < \pi, \quad |a| \leq 1.$$
30.
$$\sum_{k=1}^{\infty} (-1)^k \frac{\sin[(k+1)x]}{k(k+1)} = \sin x - \frac{1}{2}x(1 + \cos x) - \sin x \ln\left|2 \cos \frac{x}{2}\right|.$$
31.
$$\sum_{k=0}^{\infty} a^{2k+1} \sin[(2k+1)x] = \frac{a(1 + a^2) \sin x}{(1 + a^2)^2 - 4a^2 \cos^2 x}, \quad |a| < 1, \quad x \text{ is any number.}$$
32.
$$\sum_{k=0}^{\infty} (-1)^k a^{2k+1} \sin[(2k+1)x] = \frac{a(1 - a^2) \sin x}{(1 + a^2)^2 - 4a^2 \sin^2 x}, \quad |a| < 1, \quad x \text{ is any number.}$$
33.
$$\sum_{k=1}^{\infty} \frac{\sin[2(k+1)x]}{k(k+1)} = \sin(2x) - (\pi - 2x) \sin^2 x - \sin x \cos x \ln(4 \sin^2 x), \quad 0 \leq x \leq \pi.$$
34.
$$\sum_{k=1}^{\infty} (-1)^k \frac{\sin[(2k+1)x]}{(2k+1)^2} = \begin{cases} \frac{1}{4}\pi x & \text{if } -\frac{1}{2}\pi \leq x \leq \frac{1}{2}\pi, \\ \frac{1}{4}\pi(\pi - x) & \text{if } \frac{1}{2}\pi \leq x \leq \frac{3}{2}\pi. \end{cases}$$

2.4-3. Trigonometric Series in One Variable Involving Cosine.

1.
$$\sum_{k=1}^{\infty} \frac{1}{k} \cos(kx) = -\ln\left(2 \sin \frac{x}{2}\right), \quad 0 < x < 2\pi.$$

2.
$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \cos(kx) = \ln\left(2 \cos \frac{x}{2}\right), \quad -\pi < x < \pi.$$
3.
$$\sum_{k=1}^{\infty} \frac{a^k}{k} \cos(kx) = \ln \frac{1}{\sqrt{1 - 2a \cos x + a^2}}, \quad 0 < x < 2\pi, |a| \leq 1.$$
4.
$$\sum_{k=0}^{\infty} \frac{1}{2k+1} \cos(kx) = \frac{\pi}{4} \sin \frac{x}{2} + \cos \frac{x}{2} \ln\left(\cot^2 \frac{x}{4}\right), \quad 0 < x < 2\pi.$$
5.
$$\sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} \cos(kx) = -\frac{1}{4} \sin \frac{x}{2} \ln\left(\cot^2 \frac{x+\pi}{4}\right) + \frac{\pi}{4} \cos \frac{x}{2}, \quad -\pi < x < \pi.$$
6.
$$\sum_{k=1}^{\infty} \frac{1}{k^2} \cos(kx) = \frac{1}{12}(3x^2 - 6\pi x + 2\pi^2), \quad 0 \leq x \leq 2\pi.$$
7.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k^2} \cos(kx) = \frac{1}{12}(3x^2 - \pi^2), \quad -\pi \leq x \leq \pi.$$
8.
$$\sum_{k=1}^{\infty} \frac{1}{k(k+1)} \cos(kx) = \frac{1}{2}(x - \pi) \sin x - 2 \sin^2 \frac{x}{2} \ln\left(2 \sin \frac{x}{2}\right) + 1, \quad 0 \leq x \leq 2\pi.$$
9.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k(k+1)} \cos(kx) = -\frac{1}{2}x \sin x - 2 \cos^2 \frac{x}{2} \ln\left(2 \cos \frac{x}{2}\right) + 1, \quad -\pi \leq x \leq \pi.$$
10.
$$\sum_{k=1}^{\infty} \frac{1}{k^2 + a^2} \cos(kx) = \frac{\pi}{2a \sinh(\pi a)} \cosh[a(\pi - x)] - \frac{1}{2a^2}, \quad 0 \leq x \leq 2\pi.$$
11.
$$\sum_{k=1}^{\infty} \frac{1}{k^2 - a^2} \cos(kx) = -\frac{\pi}{2a \sin(\pi a)} \cos[a(\pi - x)] + \frac{1}{2a^2}, \quad 0 \leq x \leq 2\pi.$$
12.
$$\sum_{k=2}^{\infty} \frac{(-1)^k}{k^2 - 1} \cos(kx) = \frac{1}{2} - \frac{1}{4} \cos x - \frac{1}{2}x \sin x, \quad -\pi \leq x \leq \pi.$$
13.
$$\sum_{k=2}^{\infty} \frac{k}{k^2 - 1} \cos(kx) = -\frac{1}{2} - \frac{1}{4} \cos x - \cos x \ln\left(2 \sin \frac{x}{2}\right), \quad 0 < x < 2\pi.$$
14.
$$\sum_{k=1}^{\infty} \frac{1}{k^{2n}} \cos(kx) = \frac{(-1)^{n-1}(2\pi)^{2n}}{2(2n)!} B_{2n}\left(\frac{x}{2\pi}\right),$$
 where $0 \leq x \leq 2\pi$ for $n = 1, 2, \dots$;
the $B_n(x)$ are Bernoulli polynomials (see Supplement 11.18-1).
15.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k^{2n}} \cos(kx) = \frac{(-1)^{n-1}(2\pi)^{2n}}{2(2n)!} B_{2n}\left(\frac{x+\pi}{2\pi}\right),$$
 where $-\pi \leq x \leq \pi$ for $n = 1, 2, \dots$;
the $B_n(x)$ are Bernoulli polynomials.
16.
$$\sum_{k=0}^{\infty} \frac{1}{k!} \cos(kx) = \exp(\cos x) \cos(\sin x), \quad x \text{ is any number.}$$
17.
$$\sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \cos(kx) = \exp(-\cos x) \cos(\sin x), \quad x \text{ is any number.}$$
18.
$$\sum_{k=0}^{\infty} \frac{1}{(2k)!} \cos(kx) = \cos\left(\sin \frac{x}{2}\right) \cosh\left(\cos \frac{x}{2}\right), \quad x \text{ is any number.}$$
19.
$$\sum_{k=0}^{\infty} \frac{(-1)^k}{(2k)!} \cos(kx) = \cos\left(\cos \frac{x}{2}\right) \cosh\left(\sin \frac{x}{2}\right), \quad x \text{ is any number.}$$

20.
$$\sum_{k=0}^{\infty} \frac{a^k}{k!} \cos(kx) = \exp(a \cos x) \cos(a \sin x), \quad |a| \leq 1, \quad x \text{ is any number.}$$
21.
$$\sum_{k=0}^{\infty} a^k \cos(kx) = \frac{1 - a \cos x}{1 - 2a \cos x + a^2}, \quad |a| < 1, \quad x \text{ is any number.}$$
22.
$$\sum_{k=1}^{\infty} k a^k \cos(kx) = \frac{a(1 + a^2) \cos x - 2a^2}{(1 - 2a \cos x + a^2)^2}, \quad |a| < 1, \quad x \text{ is any number.}$$
23.
$$\sum_{k=1}^{\infty} \frac{1}{k} \cos(kx + a) = \frac{1}{2}(x - \pi) \sin a - \ln \left(2 \sin \frac{x}{2} \right) \cos a, \quad 0 < x < 2\pi.$$
24.
$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \cos(kx + a) = -\frac{1}{2}x \sin a + \ln \left(2 \cos \frac{x}{2} \right) \cos a, \quad -\pi < x < \pi.$$
25.
$$\sum_{k=1}^{\infty} \frac{\cos[(2k-1)x]}{2k-1} = \frac{1}{2} \ln \cot \frac{x}{2}, \quad 0 < x < \pi.$$
26.
$$\sum_{k=1}^{\infty} (-1)^{k-1} \frac{\cos[(2k-1)x]}{2k-1} = \frac{\pi}{4}, \quad 0 < x < \pi.$$
27.
$$\sum_{k=1}^{\infty} a^{2k-1} \frac{\cos[(2k-1)x]}{2k-1} = \frac{1}{4} \ln \frac{1 + 2a \cos x + a^2}{1 - 2a \cos x + a^2}, \quad 0 < x < 2\pi, \quad |a| \leq 1.$$
28.
$$\sum_{k=1}^{\infty} (-1)^{k-1} a^{2k-1} \frac{\cos[(2k-1)x]}{2k-1} = \frac{1}{2} \arctan \frac{2a \cos x}{1 - a^2}, \quad 0 < x < \pi, \quad |a| \leq 1.$$
29.
$$\sum_{k=1}^{\infty} \frac{\cos[(2k-1)x]}{(2k-1)^2} = \frac{\pi}{4} \left(\frac{\pi}{2} - |x| \right), \quad -\pi \leq x \leq \pi.$$
30.
$$\sum_{k=1}^{\infty} (-1)^k \frac{\cos[(k+1)x]}{k(k+1)} = \cos x - \frac{1}{2}x \sin x - (1 + \cos x) \ln \left| 2 \cos \frac{x}{2} \right|.$$
31.
$$\sum_{k=0}^{\infty} a^{2k+1} \cos[(2k+1)x] = \frac{a(1 - a^2) \cos x}{(1 + a^2)^2 - 4a^2 \cos^2 x}, \quad |a| < 1, \quad x \text{ is any number.}$$
32.
$$\sum_{k=0}^{\infty} (-1)^k a^{2k+1} \cos[(2k+1)x] = \frac{a(1 + a^2) \cos x}{(1 + a^2)^2 - 4a^2 \sin^2 x}, \quad |a| < 1, \quad x \text{ is any number.}$$
33.
$$\sum_{k=1}^{\infty} \frac{\cos[2(k+1)x]}{k(k+1)} = \cos(2x) - \left(\frac{\pi}{2} - x \right) \sin(2x) + \sin^2 x \ln(4 \sin^2 x), \quad 0 \leq x \leq \pi.$$

2.4-4. Trigonometric Series in Two Variables.

1.
$$\sum_{k=1}^{\infty} \frac{1}{k} \sin(kx) \sin(ky) = \frac{1}{2} \ln \left| \sin \frac{x+y}{2} \operatorname{cosec} \frac{x-y}{2} \right|, \quad x \pm y \neq 0, 2\pi, 4\pi, \dots$$
2.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k} \sin(kx) \sin(ky) = \frac{1}{2} \ln \left| \cos \frac{x+y}{2} \sec \frac{x-y}{2} \right|, \quad x \pm y \neq \pi, 3\pi, 5\pi, \dots$$
3.
$$\sum_{k=1}^{\infty} \frac{1}{k^2} \sin(kx) \sin(ky) = \begin{cases} \frac{1}{2}x(\pi - y) & \text{if } -y \leq x \leq y, \\ \frac{1}{2}y(\pi - x) & \text{if } y \leq x \leq 2\pi - y. \end{cases} \quad \text{Here } 0 < y < \pi.$$

4.
$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2} \sin(kx) \sin(ky) = \frac{1}{2}xy, \quad |x \pm y| \leq \pi.$$
5.
$$\sum_{k=1}^{\infty} \frac{a^k}{k} \sin(kx) \sin(ky) = \frac{1}{4} \ln \frac{4a \sin^2[(x+y)/2] + (a-1)^2}{4a \sin^2[(x-y)/2] + (a-1)^2}, \quad 0 < a < 1.$$
6.
$$\sum_{k=1}^{\infty} \frac{1}{k^2} \sin^2(kx) \sin^2(ky) = \frac{1}{2}\pi x, \quad 0 \leq x \leq y \leq \frac{\pi}{2}.$$
7.
$$\sum_{k=1}^{\infty} \frac{1}{k} \cos(kx) \cos(ky) = -\frac{1}{2} \ln |2(\cos x - \cos y)|, \quad x \pm y \neq 0, 2\pi, 4\pi, \dots$$
8.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k} \cos(kx) \cos(ky) = -\frac{1}{2} \ln |2(\cos x + \cos y)|, \quad x \pm y \neq \pi, 3\pi, 5\pi, \dots$$
9.
$$\sum_{k=1}^{\infty} \frac{1}{k} \sin(kx) \cos(ky) = \begin{cases} -\frac{1}{2} & \text{if } 0 < x < y, \\ \frac{1}{4}(\pi - 2y) & \text{if } x = y, \\ \frac{1}{2}(\pi - x) & \text{if } y < x < \pi. \end{cases} \quad \text{Here } 0 < y < \pi.$$
10.
$$\sum_{k=1}^{\infty} \frac{1}{k^2} \cos(kx) \cos(ky) = \begin{cases} \frac{1}{12} [3x^2 + 3(y-\pi)^2 - \pi^2] & \text{if } 0 \leq x \leq y, \\ \frac{1}{12} [3y^2 + 3(x-\pi)^2 - \pi^2] & \text{if } y \leq x \leq \pi. \end{cases}$$

Here $0 < y < \pi$.
11.
$$\sum_{k=1}^{\infty} \frac{(-1)^k}{k^2} \cos(kx) \cos(ky) = \begin{cases} \frac{1}{12} [3(x^2 + y^2) - \pi^2] & \text{if } -(\pi - y) \leq x \leq \pi - y, \\ \frac{1}{12} [3(x - \pi)^2 + 3(y - \pi)^2 - \pi^2] & \text{if } \pi - y \leq x \leq \pi + y. \end{cases}$$

Here $0 < y < \pi$.

⊙ References for Supplement 2: H. B. Dwight (1961), V. Mangulis (1965), E. R. Hansen (1975), I. S. Gradshteyn and I. M. Ryzhik (2000), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1986), D. Zwillinger (2002).

Supplement 3

Tables of Indefinite Integrals

► Throughout Supplement 3, the integration constant C is omitted for brevity.

3.1. Integrals Involving Rational Functions

3.1-1. Integrals Involving $a + bx$.

- $$\int \frac{dx}{a + bx} = \frac{1}{b} \ln |a + bx|.$$
- $$\int (a + bx)^n dx = \frac{(a + bx)^{n+1}}{b(n+1)}, \quad n \neq -1.$$
- $$\int \frac{x dx}{a + bx} = \frac{1}{b^2} (a + bx - a \ln |a + bx|).$$
- $$\int \frac{x^2 dx}{a + bx} = \frac{1}{b^3} \left[\frac{1}{2} (a + bx)^2 - 2a(a + bx) + a^2 \ln |a + bx| \right].$$
- $$\int \frac{dx}{x(a + bx)} = -\frac{1}{a} \ln \left| \frac{a + bx}{x} \right|.$$
- $$\int \frac{dx}{x^2(a + bx)} = -\frac{1}{ax} + \frac{b}{a^2} \ln \left| \frac{a + bx}{x} \right|.$$
- $$\int \frac{x dx}{(a + bx)^2} = \frac{1}{b^2} \left(\ln |a + bx| + \frac{a}{a + bx} \right).$$
- $$\int \frac{x^2 dx}{(a + bx)^2} = \frac{1}{b^3} \left(a + bx - 2a \ln |a + bx| - \frac{a^2}{a + bx} \right).$$
- $$\int \frac{dx}{x(a + bx)^2} = \frac{1}{a(a + bx)} - \frac{1}{a^2} \ln \left| \frac{a + bx}{x} \right|.$$
- $$\int \frac{x dx}{(a + bx)^3} = \frac{1}{b^2} \left[-\frac{1}{a + bx} + \frac{a}{2(a + bx)^2} \right].$$

3.1-2. Integrals Involving $a + x$ and $b + x$.

- $$\int \frac{a + x}{b + x} dx = x + (a - b) \ln |b + x|.$$
- $$\int \frac{dx}{(a + x)(b + x)} = \frac{1}{a - b} \ln \left| \frac{b + x}{a + x} \right|, \quad a \neq b. \quad \text{For } a = b, \text{ see Integral 2 with } n = -2 \text{ in Supplement 3.1-1.}$$
- $$\int \frac{x dx}{(a + x)(b + x)} = \frac{1}{a - b} (a \ln |a + x| - b \ln |b + x|).$$

4. $\int \frac{dx}{(a+x)(b+x)^2} = \frac{1}{(b-a)(b+x)} + \frac{1}{(a-b)^2} \ln \left| \frac{a+x}{b+x} \right|.$
5. $\int \frac{x dx}{(a+x)(b+x)^2} = \frac{b}{(a-b)(b+x)} - \frac{a}{(a-b)^2} \ln \left| \frac{a+x}{b+x} \right|.$
6. $\int \frac{x^2 dx}{(a+x)(b+x)^2} = \frac{b^2}{(b-a)(b+x)} + \frac{a^2}{(a-b)^2} \ln |a+x| + \frac{b^2 - 2ab}{(b-a)^2} \ln |b+x|.$
7. $\int \frac{dx}{(a+x)^2(b+x)^2} = -\frac{1}{(a-b)^2} \left(\frac{1}{a+x} + \frac{1}{b+x} \right) + \frac{2}{(a-b)^3} \ln \left| \frac{a+x}{b+x} \right|.$
8. $\int \frac{x dx}{(a+x)^2(b+x)^2} = \frac{1}{(a-b)^2} \left(\frac{a}{a+x} + \frac{b}{b+x} \right) + \frac{a+b}{(a-b)^3} \ln \left| \frac{a+x}{b+x} \right|.$
9. $\int \frac{x^2 dx}{(a+x)^2(b+x)^2} = -\frac{1}{(a-b)^2} \left(\frac{a^2}{a+x} + \frac{b^2}{b+x} \right) + \frac{2ab}{(a-b)^3} \ln \left| \frac{a+x}{b+x} \right|.$

3.1-3. Integrals Involving $a^2 + x^2$.
--

1. $\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \arctan \frac{x}{a}.$
2. $\int \frac{dx}{(a^2 + x^2)^2} = \frac{x}{2a^2(a^2 + x^2)} + \frac{1}{2a^3} \arctan \frac{x}{a}.$
3. $\int \frac{dx}{(a^2 + x^2)^3} = \frac{x}{4a^2(a^2 + x^2)^2} + \frac{3x}{8a^4(a^2 + x^2)} + \frac{3}{8a^5} \arctan \frac{x}{a}.$
4. $\int \frac{dx}{(a^2 + x^2)^{n+1}} = \frac{x}{2na^2(a^2 + x^2)^n} + \frac{2n-1}{2na^2} \int \frac{dx}{(a^2 + x^2)^n}; \quad n = 1, 2, \dots$
5. $\int \frac{x dx}{a^2 + x^2} = \frac{1}{2} \ln(a^2 + x^2).$
6. $\int \frac{x dx}{(a^2 + x^2)^2} = -\frac{1}{2(a^2 + x^2)}.$
7. $\int \frac{x dx}{(a^2 + x^2)^3} = -\frac{1}{4(a^2 + x^2)^2}.$
8. $\int \frac{x dx}{(a^2 + x^2)^{n+1}} = -\frac{1}{2n(a^2 + x^2)^n}; \quad n = 1, 2, \dots$
9. $\int \frac{x^2 dx}{a^2 + x^2} = x - a \arctan \frac{x}{a}.$
10. $\int \frac{x^2 dx}{(a^2 + x^2)^2} = -\frac{x}{2(a^2 + x^2)} + \frac{1}{2a} \arctan \frac{x}{a}.$
11. $\int \frac{x^2 dx}{(a^2 + x^2)^3} = -\frac{x}{4(a^2 + x^2)^2} + \frac{x}{8a^2(a^2 + x^2)} + \frac{1}{8a^3} \arctan \frac{x}{a}.$
12. $\int \frac{x^2 dx}{(a^2 + x^2)^{n+1}} = -\frac{x}{2n(a^2 + x^2)^n} + \frac{1}{2n} \int \frac{dx}{(a^2 + x^2)^n}; \quad n = 1, 2, \dots$
13. $\int \frac{x^3 dx}{a^2 + x^2} = \frac{x^2}{2} - \frac{a^2}{2} \ln(a^2 + x^2).$
14. $\int \frac{x^3 dx}{(a^2 + x^2)^2} = \frac{a^2}{2(a^2 + x^2)} + \frac{1}{2} \ln(a^2 + x^2).$
15. $\int \frac{x^3 dx}{(a^2 + x^2)^{n+1}} = -\frac{1}{2(n-1)(a^2 + x^2)^{n-1}} + \frac{a^2}{2n(a^2 + x^2)^n}; \quad n = 2, 3, \dots$
16. $\int \frac{dx}{x(a^2 + x^2)} = \frac{1}{2a^2} \ln \frac{x^2}{a^2 + x^2}.$

17. $\int \frac{dx}{x(a^2 + x^2)^2} = \frac{1}{2a^2(a^2 + x^2)} + \frac{1}{2a^4} \ln \frac{x^2}{a^2 + x^2}.$
18. $\int \frac{dx}{x(a^2 + x^2)^3} = \frac{1}{4a^2(a^2 + x^2)^2} + \frac{1}{2a^4(a^2 + x^2)} + \frac{1}{2a^6} \ln \frac{x^2}{a^2 + x^2}.$
19. $\int \frac{dx}{x^2(a^2 + x^2)} = -\frac{1}{a^2x} - \frac{1}{a^3} \arctan \frac{x}{a}.$
20. $\int \frac{dx}{x^2(a^2 + x^2)^2} = -\frac{1}{a^4x} - \frac{x}{2a^4(a^2 + x^2)} - \frac{3}{2a^5} \arctan \frac{x}{a}.$
21. $\int \frac{dx}{x^3(a^2 + x^2)^2} = -\frac{1}{2a^4x^2} - \frac{1}{2a^4(a^2 + x^2)} - \frac{1}{a^6} \ln \frac{x^2}{a^2 + x^2}.$
22. $\int \frac{dx}{x^2(a^2 + x^2)^3} = -\frac{1}{a^6x} - \frac{x}{4a^4(a^2 + x^2)^2} - \frac{7x}{8a^6(a^2 + x^2)} - \frac{15}{8a^7} \arctan \frac{x}{a}.$
23. $\int \frac{dx}{x^3(a^2 + x^2)^3} = -\frac{1}{2a^6x^2} - \frac{1}{a^6(a^2 + x^2)} - \frac{1}{4a^4(a^2 + x^2)^2} - \frac{3}{2a^8} \ln \frac{x^2}{a^2 + x^2}.$

3.1-4. Integrals Involving $a^2 - x^2$.

1. $\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right|.$
2. $\int \frac{dx}{(a^2 - x^2)^2} = \frac{x}{2a^2(a^2 - x^2)} + \frac{1}{4a^3} \ln \left| \frac{a+x}{a-x} \right|.$
3. $\int \frac{dx}{(a^2 - x^2)^3} = \frac{x}{4a^2(a^2 - x^2)^2} + \frac{3x}{8a^4(a^2 - x^2)} + \frac{3}{16a^5} \ln \left| \frac{a+x}{a-x} \right|.$
4. $\int \frac{dx}{(a^2 - x^2)^{n+1}} = \frac{x}{2na^2(a^2 - x^2)^n} + \frac{2n-1}{2na^2} \int \frac{dx}{(a^2 - x^2)^n}; \quad n = 1, 2, \dots$
5. $\int \frac{x dx}{a^2 - x^2} = -\frac{1}{2} \ln |a^2 - x^2|.$
6. $\int \frac{x dx}{(a^2 - x^2)^2} = \frac{1}{2(a^2 - x^2)}.$
7. $\int \frac{x dx}{(a^2 - x^2)^3} = \frac{1}{4(a^2 - x^2)^2}.$
8. $\int \frac{x dx}{(a^2 - x^2)^{n+1}} = \frac{1}{2n(a^2 - x^2)^n}; \quad n = 1, 2, \dots$
9. $\int \frac{x^2 dx}{a^2 - x^2} = -x + \frac{a}{2} \ln \left| \frac{a+x}{a-x} \right|.$
10. $\int \frac{x^2 dx}{(a^2 - x^2)^2} = \frac{x}{2(a^2 - x^2)} - \frac{1}{4a} \ln \left| \frac{a+x}{a-x} \right|.$
11. $\int \frac{x^2 dx}{(a^2 - x^2)^3} = \frac{x}{4(a^2 - x^2)^2} - \frac{x}{8a^2(a^2 - x^2)} - \frac{1}{16a^3} \ln \left| \frac{a+x}{a-x} \right|.$
12. $\int \frac{x^2 dx}{(a^2 - x^2)^{n+1}} = \frac{x}{2n(a^2 - x^2)^n} - \frac{1}{2n} \int \frac{dx}{(a^2 - x^2)^n}; \quad n = 1, 2, \dots$
13. $\int \frac{x^3 dx}{a^2 - x^2} = -\frac{x^2}{2} - \frac{a^2}{2} \ln |a^2 - x^2|.$
14. $\int \frac{x^3 dx}{(a^2 - x^2)^2} = \frac{a^2}{2(a^2 - x^2)} + \frac{1}{2} \ln |a^2 - x^2|.$

15. $\int \frac{x^3 dx}{(a^2 - x^2)^{n+1}} = -\frac{1}{2(n-1)(a^2 - x^2)^{n-1}} + \frac{a^2}{2n(a^2 - x^2)^n}; \quad n = 2, 3, \dots$
16. $\int \frac{dx}{x(a^2 - x^2)} = \frac{1}{2a^2} \ln \left| \frac{x^2}{a^2 - x^2} \right|.$
17. $\int \frac{dx}{x(a^2 - x^2)^2} = \frac{1}{2a^2(a^2 - x^2)} + \frac{1}{2a^4} \ln \left| \frac{x^2}{a^2 - x^2} \right|.$
18. $\int \frac{dx}{x(a^2 - x^2)^3} = \frac{1}{4a^2(a^2 - x^2)^2} + \frac{1}{2a^4(a^2 - x^2)} + \frac{1}{2a^6} \ln \left| \frac{x^2}{a^2 - x^2} \right|.$

3.1-5. Integrals Involving $a^3 + x^3$.

1. $\int \frac{dx}{a^3 + x^3} = \frac{1}{6a^2} \ln \frac{(a+x)^2}{a^2 - ax + x^2} + \frac{1}{a^2\sqrt{3}} \arctan \frac{2x-a}{a\sqrt{3}}.$
2. $\int \frac{dx}{(a^3 + x^3)^2} = \frac{x}{3a^3(a^3 + x^3)} + \frac{2}{3a^3} \int \frac{dx}{a^3 + x^3}.$
3. $\int \frac{x dx}{a^3 + x^3} = \frac{1}{6a} \ln \frac{a^2 - ax + x^2}{(a+x)^2} + \frac{1}{a\sqrt{3}} \arctan \frac{2x-a}{a\sqrt{3}}.$
4. $\int \frac{x dx}{(a^3 + x^3)^2} = \frac{x^2}{3a^3(a^3 + x^3)} + \frac{1}{3a^3} \int \frac{x dx}{a^3 + x^3}.$
5. $\int \frac{x^2 dx}{a^3 + x^3} = \frac{1}{3} \ln |a^3 + x^3|.$
6. $\int \frac{dx}{x(a^3 + x^3)} = \frac{1}{3a^3} \ln \left| \frac{x^3}{a^3 + x^3} \right|.$
7. $\int \frac{dx}{x(a^3 + x^3)^2} = \frac{1}{3a^3(a^3 + x^3)} + \frac{1}{3a^6} \ln \left| \frac{x^3}{a^3 + x^3} \right|.$
8. $\int \frac{dx}{x^2(a^3 + x^3)} = -\frac{1}{a^3x} - \frac{1}{a^3} \int \frac{x dx}{a^3 + x^3}.$
9. $\int \frac{dx}{x^2(a^3 + x^3)^2} = -\frac{1}{a^6x} - \frac{x^2}{3a^6(a^3 + x^3)} - \frac{4}{3a^6} \int \frac{x dx}{a^3 + x^3}.$

3.1-6. Integrals Involving $a^3 - x^3$.

1. $\int \frac{dx}{a^3 - x^3} = \frac{1}{6a^2} \ln \frac{a^2 + ax + x^2}{(a-x)^2} + \frac{1}{a^2\sqrt{3}} \arctan \frac{2x+a}{a\sqrt{3}}.$
2. $\int \frac{dx}{(a^3 - x^3)^2} = \frac{x}{3a^3(a^3 - x^3)} + \frac{2}{3a^3} \int \frac{dx}{a^3 - x^3}.$
3. $\int \frac{x dx}{a^3 - x^3} = \frac{1}{6a} \ln \frac{a^2 + ax + x^2}{(a-x)^2} - \frac{1}{a\sqrt{3}} \arctan \frac{2x+a}{a\sqrt{3}}.$
4. $\int \frac{x dx}{(a^3 - x^3)^2} = \frac{x^2}{3a^3(a^3 - x^3)} + \frac{1}{3a^3} \int \frac{x dx}{a^3 - x^3}.$
5. $\int \frac{x^2 dx}{a^3 - x^3} = -\frac{1}{3} \ln |a^3 - x^3|.$
6. $\int \frac{dx}{x(a^3 - x^3)} = \frac{1}{3a^3} \ln \left| \frac{x^3}{a^3 - x^3} \right|.$
7. $\int \frac{dx}{x(a^3 - x^3)^2} = \frac{1}{3a^3(a^3 - x^3)} + \frac{1}{3a^6} \ln \left| \frac{x^3}{a^3 - x^3} \right|.$

8. $\int \frac{dx}{x^2(a^3 - x^3)} = -\frac{1}{a^3x} + \frac{1}{a^3} \int \frac{x dx}{a^3 - x^3}.$
9. $\int \frac{dx}{x^2(a^3 - x^3)^2} = -\frac{1}{a^6x} - \frac{x^2}{3a^6(a^3 - x^3)} + \frac{4}{3a^6} \int \frac{x dx}{a^3 - x^3}.$

3.1-7. Integrals Involving $a^4 \pm x^4$.

1. $\int \frac{dx}{a^4 + x^4} = \frac{1}{4a^3\sqrt{2}} \ln \frac{a^2 + ax\sqrt{2} + x^2}{a^2 - ax\sqrt{2} + x^2} + \frac{1}{2a^3\sqrt{2}} \arctan \frac{ax\sqrt{2}}{a^2 - x^2}.$
2. $\int \frac{x dx}{a^4 + x^4} = \frac{1}{2a^2} \arctan \frac{x^2}{a^2}.$
3. $\int \frac{x^2 dx}{a^4 + x^4} = -\frac{1}{4a\sqrt{2}} \ln \frac{a^2 + ax\sqrt{2} + x^2}{a^2 - ax\sqrt{2} + x^2} + \frac{1}{2a\sqrt{2}} \arctan \frac{ax\sqrt{2}}{a^2 - x^2}.$
4. $\int \frac{dx}{a^4 - x^4} = \frac{1}{4a^3} \ln \left| \frac{a+x}{a-x} \right| + \frac{1}{2a^3} \arctan \frac{x}{a}.$
5. $\int \frac{x dx}{a^4 - x^4} = \frac{1}{4a^2} \ln \left| \frac{a^2 + x^2}{a^2 - x^2} \right|.$
6. $\int \frac{x^2 dx}{a^4 - x^4} = \frac{1}{4a} \ln \left| \frac{a+x}{a-x} \right| - \frac{1}{2a} \arctan \frac{x}{a}.$

3.2. Integrals Involving Irrational Functions

3.2-1. Integrals Involving $x^{1/2}$.

1. $\int \frac{x^{1/2} dx}{a^2 + b^2x} = \frac{2}{b^2} x^{1/2} - \frac{2a}{b^3} \arctan \frac{bx^{1/2}}{a}.$
2. $\int \frac{x^{3/2} dx}{a^2 + b^2x} = \frac{2x^{3/2}}{3b^2} - \frac{2a^2x^{1/2}}{b^4} + \frac{2a^3}{b^5} \arctan \frac{bx^{1/2}}{a}.$
3. $\int \frac{x^{1/2} dx}{(a^2 + b^2x)^2} = -\frac{x^{1/2}}{b^2(a^2 + b^2x)} + \frac{1}{ab^3} \arctan \frac{bx^{1/2}}{a}.$
4. $\int \frac{x^{3/2} dx}{(a^2 + b^2x)^2} = \frac{2x^{3/2}}{b^2(a^2 + b^2x)} + \frac{3a^2x^{1/2}}{b^4(a^2 + b^2x)} - \frac{3a}{b^5} \arctan \frac{bx^{1/2}}{a}.$
5. $\int \frac{dx}{(a^2 + b^2x)x^{1/2}} = \frac{2}{ab} \arctan \frac{bx^{1/2}}{a}.$
6. $\int \frac{dx}{(a^2 + b^2x)x^{3/2}} = -\frac{2}{a^2x^{1/2}} - \frac{2b}{a^3} \arctan \frac{bx^{1/2}}{a}.$
7. $\int \frac{dx}{(a^2 + b^2x)^2x^{1/2}} = \frac{x^{1/2}}{a^2(a^2 + b^2x)} + \frac{1}{a^3b} \arctan \frac{bx^{1/2}}{a}.$
8. $\int \frac{x^{1/2} dx}{a^2 - b^2x} = -\frac{2}{b^2} x^{1/2} + \frac{2a}{b^3} \ln \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$
9. $\int \frac{x^{3/2} dx}{a^2 - b^2x} = -\frac{2x^{3/2}}{3b^2} - \frac{2a^2x^{1/2}}{b^4} + \frac{a^3}{b^5} \ln \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$
10. $\int \frac{x^{1/2} dx}{(a^2 - b^2x)^2} = \frac{x^{1/2}}{b^2(a^2 - b^2x)} - \frac{1}{2ab^3} \ln \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$

11. $\int \frac{x^{3/2} dx}{(a^2 - b^2x)^2} = \frac{3a^2x^{1/2} - 2b^2x^{3/2}}{b^4(a^2 - b^2x)} - \frac{3a}{2b^5} \ln \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$
12. $\int \frac{dx}{(a^2 - b^2x)x^{1/2}} = \frac{1}{ab} \ln \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$
13. $\int \frac{dx}{(a^2 - b^2x)x^{3/2}} = -\frac{2}{a^2x^{1/2}} + \frac{b}{a^3} \ln \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$
14. $\int \frac{dx}{(a^2 - b^2x)^2x^{1/2}} = \frac{x^{1/2}}{a^2(a^2 - b^2x)} + \frac{1}{2a^3b} \ln \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$

3.2-2. Integrals Involving $(a + bx)^{p/2}$.

1. $\int (a + bx)^{p/2} dx = \frac{2}{b(p+2)}(a + bx)^{(p+2)/2}.$
2. $\int x(a + bx)^{p/2} dx = \frac{2}{b^2} \left[\frac{(a + bx)^{(p+4)/2}}{p+4} - \frac{a(a + bx)^{(p+2)/2}}{p+2} \right].$
3. $\int x^2(a + bx)^{p/2} dx = \frac{2}{b^3} \left[\frac{(a + bx)^{(p+6)/2}}{p+6} - \frac{2a(a + bx)^{(p+4)/2}}{p+4} + \frac{a^2(a + bx)^{(p+2)/2}}{p+2} \right].$

3.2-3. Integrals Involving $(x^2 + a^2)^{1/2}$.

1. $\int (x^2 + a^2)^{1/2} dx = \frac{1}{2}x(a^2 + x^2)^{1/2} + \frac{a^2}{2} \ln [x + (x^2 + a^2)^{1/2}].$
2. $\int x(x^2 + a^2)^{1/2} dx = \frac{1}{3}(a^2 + x^2)^{3/2}.$
3. $\int (x^2 + a^2)^{3/2} dx = \frac{1}{4}x(a^2 + x^2)^{3/2} + \frac{3}{8}a^2x(a^2 + x^2)^{1/2} + \frac{3}{8}a^4 \ln |x + (x^2 + a^2)^{1/2}|.$
4. $\int \frac{1}{x}(x^2 + a^2)^{1/2} dx = (a^2 + x^2)^{1/2} - a \ln \left| \frac{a + (x^2 + a^2)^{1/2}}{x} \right|.$
5. $\int \frac{dx}{\sqrt{x^2 + a^2}} = \ln [x + (x^2 + a^2)^{1/2}].$
6. $\int \frac{x dx}{\sqrt{x^2 + a^2}} = (x^2 + a^2)^{1/2}.$
7. $\int (x^2 + a^2)^{-3/2} dx = a^{-2}x(x^2 + a^2)^{-1/2}.$

3.2-4. Integrals Involving $(x^2 - a^2)^{1/2}$.

1. $\int (x^2 - a^2)^{1/2} dx = \frac{1}{2}x(x^2 - a^2)^{1/2} - \frac{a^2}{2} \ln |x + (x^2 - a^2)^{1/2}|.$
2. $\int x(x^2 - a^2)^{1/2} dx = \frac{1}{3}(x^2 - a^2)^{3/2}.$
3. $\int (x^2 - a^2)^{3/2} dx = \frac{1}{4}x(x^2 - a^2)^{3/2} - \frac{3}{8}a^2x(x^2 - a^2)^{1/2} + \frac{3}{8}a^4 \ln |x + (x^2 - a^2)^{1/2}|.$
4. $\int \frac{1}{x}(x^2 - a^2)^{1/2} dx = (x^2 - a^2)^{1/2} - a \arccos \left| \frac{a}{x} \right|.$

5. $\int \frac{dx}{\sqrt{x^2 - a^2}} = \ln|x + (x^2 - a^2)^{1/2}|.$
6. $\int \frac{x dx}{\sqrt{x^2 - a^2}} = (x^2 - a^2)^{1/2}.$
7. $\int (x^2 - a^2)^{-3/2} dx = -a^{-2}x(x^2 - a^2)^{-1/2}.$

3.2-5. Integrals Involving $(a^2 - x^2)^{1/2}.$

1. $\int (a^2 - x^2)^{1/2} dx = \frac{1}{2}x(a^2 - x^2)^{1/2} + \frac{a^2}{2} \arcsin \frac{x}{a}.$
2. $\int x(a^2 - x^2)^{1/2} dx = -\frac{1}{3}(a^2 - x^2)^{3/2}.$
3. $\int (a^2 - x^2)^{3/2} dx = \frac{1}{4}x(a^2 - x^2)^{3/2} + \frac{3}{8}a^2x(a^2 - x^2)^{1/2} + \frac{3}{8}a^4 \arcsin \frac{x}{a}.$
4. $\int \frac{1}{x}(a^2 - x^2)^{1/2} dx = (a^2 - x^2)^{1/2} - a \ln \left| \frac{a + (a^2 - x^2)^{1/2}}{x} \right|.$
5. $\int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a}.$
6. $\int \frac{x dx}{\sqrt{a^2 - x^2}} = -(a^2 - x^2)^{1/2}.$
7. $\int (a^2 - x^2)^{-3/2} dx = a^{-2}x(a^2 - x^2)^{-1/2}.$

3.2-6. Integrals Involving Arbitrary Powers. Reduction Formulas.

1. $\int \frac{dx}{x(ax^n + b)} = \frac{1}{bn} \ln \left| \frac{x^n}{ax^n + b} \right|.$
2. $\int \frac{dx}{x\sqrt{x^n + a^2}} = \frac{2}{an} \ln \left| \frac{x^{n/2}}{\sqrt{x^n + a^2} + a} \right|.$
3. $\int \frac{dx}{x\sqrt{x^n - a^2}} = \frac{2}{an} \arccos \left| \frac{a}{x^{n/2}} \right|.$
4. $\int \frac{dx}{x\sqrt{ax^{2n} + bx^n}} = -\frac{2\sqrt{ax^{2n} + bx^n}}{bnx^n}.$

► The parameters a , b , p , m , and n below in Integrals 5–8 can assume arbitrary values, except for those at which denominators vanish in successive applications of a formula. Notation: $w = ax^n + b$.

5. $\int x^m(ax^n + b)^p dx = \frac{1}{m + np + 1} \left(x^{m+1}w^p + npb \int x^m w^{p-1} dx \right).$
6. $\int x^m(ax^n + b)^p dx = \frac{1}{bn(p+1)} \left[-x^{m+1}w^{p+1} + (m+n+np+1) \int x^m w^{p+1} dx \right].$
7. $\int x^m(ax^n + b)^p dx = \frac{1}{b(m+1)} \left[x^{m+1}w^{p+1} - a(m+n+np+1) \int x^{m+n}w^p dx \right].$
8. $\int x^m(ax^n + b)^p dx = \frac{1}{a(m+np+1)} \left[x^{m-n+1}w^{p+1} - b(m-n+1) \int x^{m-n}w^p dx \right].$

3.3. Integrals Involving Exponential Functions

1. $\int e^{ax} dx = \frac{1}{a} e^{ax}.$
2. $\int a^x dx = \frac{a^x}{\ln a}.$
3. $\int x e^{ax} dx = e^{ax} \left(\frac{x}{a} - \frac{1}{a^2} \right).$
4. $\int x^2 e^{ax} dx = e^{ax} \left(\frac{x^2}{a} - \frac{2x}{a^2} + \frac{2}{a^3} \right).$
5. $\int x^n e^{ax} dx = e^{ax} \left[\frac{1}{a} x^n - \frac{n}{a^2} x^{n-1} + \frac{n(n-1)}{a^3} x^{n-2} - \dots + (-1)^{n-1} \frac{n!}{a^n} x + (-1)^n \frac{n!}{a^{n+1}} \right], \quad n=1, 2, \dots$
6. $\int P_n(x) e^{ax} dx = e^{ax} \sum_{k=0}^n \frac{(-1)^k}{a^{k+1}} \frac{d^k}{dx^k} P_n(x),$ where $P_n(x)$ is an arbitrary polynomial of degree n .
7. $\int \frac{dx}{a + b e^{px}} = \frac{x}{a} - \frac{1}{ap} \ln |a + b e^{px}|.$
8. $\int \frac{dx}{a e^{px} + b e^{-px}} = \begin{cases} \frac{1}{p\sqrt{ab}} \arctan \left(e^{px} \sqrt{\frac{a}{b}} \right) & \text{if } ab > 0, \\ \frac{1}{2p\sqrt{-ab}} \ln \left(\frac{b + e^{px}\sqrt{-ab}}{b - e^{px}\sqrt{-ab}} \right) & \text{if } ab < 0. \end{cases}$
9. $\int \frac{dx}{\sqrt{a + b e^{px}}} = \begin{cases} \frac{1}{p\sqrt{a}} \ln \frac{\sqrt{a + b e^{px}} - \sqrt{a}}{\sqrt{a + b e^{px}} + \sqrt{a}} & \text{if } a > 0, \\ \frac{2}{p\sqrt{-a}} \arctan \frac{\sqrt{a + b e^{px}}}{\sqrt{-a}} & \text{if } a < 0. \end{cases}$

3.4. Integrals Involving Hyperbolic Functions

3.4-1. Integrals Involving $\cosh x$.

1. $\int \cosh(a + bx) dx = \frac{1}{b} \sinh(a + bx).$
2. $\int x \cosh x dx = x \sinh x - \cosh x.$
3. $\int x^2 \cosh x dx = (x^2 + 2) \sinh x - 2x \cosh x.$
4. $\int x^{2n} \cosh x dx = (2n)! \sum_{k=1}^n \left[\frac{x^{2k}}{(2k)!} \sinh x - \frac{x^{2k-1}}{(2k-1)!} \cosh x \right].$
5. $\int x^{2n+1} \cosh x dx = (2n+1)! \sum_{k=0}^n \left[\frac{x^{2k+1}}{(2k+1)!} \sinh x - \frac{x^{2k}}{(2k)!} \cosh x \right].$
6. $\int x^p \cosh x dx = x^p \sinh x - p x^{p-1} \cosh x + p(p-1) \int x^{p-2} \cosh x dx.$
7. $\int \cosh^2 x dx = \frac{1}{2} x + \frac{1}{4} \sinh 2x.$
8. $\int \cosh^3 x dx = \sinh x + \frac{1}{3} \sinh^3 x.$

9. $\int \cosh^{2n} x \, dx = C_{2n}^n \frac{x}{2^{2n}} + \frac{1}{2^{2n-1}} \sum_{k=0}^{n-1} C_{2n}^k \frac{\sinh[2(n-k)x]}{2(n-k)}, \quad n = 1, 2, \dots$
10. $\int \cosh^{2n+1} x \, dx = \frac{1}{2^{2n}} \sum_{k=0}^n C_{2n+1}^k \frac{\sinh[(2n-2k+1)x]}{2n-2k+1} = \sum_{k=0}^n C_n^k \frac{\sinh^{2k+1} x}{2k+1}, \quad n = 1, 2, \dots$
11. $\int \cosh^p x \, dx = \frac{1}{p} \sinh x \cosh^{p-1} x + \frac{p-1}{p} \int \cosh^{p-2} x \, dx.$
12. $\int \cosh ax \cosh bx \, dx = \frac{1}{a^2 - b^2} (a \cosh bx \sinh ax - b \cosh ax \sinh bx).$
13. $\int \frac{dx}{\cosh ax} = \frac{2}{a} \arctan(e^{ax}).$
14. $\int \frac{dx}{\cosh^{2n} x} = \frac{\sinh x}{2n-1} \left[\frac{1}{\cosh^{2n-1} x} + \sum_{k=1}^{n-1} \frac{2^k(n-1)(n-2)\dots(n-k)}{(2n-3)(2n-5)\dots(2n-2k-1)} \frac{1}{\cosh^{2n-2k-1} x} \right], \quad n = 1, 2, \dots$
15. $\int \frac{dx}{\cosh^{2n+1} x} = \frac{\sinh x}{2n} \left[\frac{1}{\cosh^{2n} x} + \sum_{k=1}^{n-1} \frac{(2n-1)(2n-3)\dots(2n-2k+1)}{2^k(n-1)(n-2)\dots(n-k)} \frac{1}{\cosh^{2n-2k} x} \right] + \frac{(2n-1)!!}{(2n)!!} \arctan \sinh x, \quad n = 1, 2, \dots$
16. $\int \frac{dx}{a+b \cosh x} = \begin{cases} -\frac{\operatorname{sign} x}{\sqrt{b^2-a^2}} \arcsin \frac{b+a \cosh x}{a+b \cosh x} & \text{if } a^2 < b^2, \\ \frac{1}{\sqrt{a^2-b^2}} \ln \frac{a+b+\sqrt{a^2-b^2} \tanh(x/2)}{a+b-\sqrt{a^2-b^2} \tanh(x/2)} & \text{if } a^2 > b^2. \end{cases}$

3.4-2. Integrals Involving $\sinh x$.

1. $\int \sinh(a+bx) \, dx = \frac{1}{b} \cosh(a+bx).$
2. $\int x \sinh x \, dx = x \cosh x - \sinh x.$
3. $\int x^2 \sinh x \, dx = (x^2+2) \cosh x - 2x \sinh x.$
4. $\int x^{2n} \sinh x \, dx = (2n)! \left[\sum_{k=0}^n \frac{x^{2k}}{(2k)!} \cosh x - \sum_{k=1}^n \frac{x^{2k-1}}{(2k-1)!} \sinh x \right].$
5. $\int x^{2n+1} \sinh x \, dx = (2n+1)! \sum_{k=0}^n \left[\frac{x^{2k+1}}{(2k+1)!} \cosh x - \frac{x^{2k}}{(2k)!} \sinh x \right].$
6. $\int x^p \sinh x \, dx = x^p \cosh x - px^{p-1} \sinh x + p(p-1) \int x^{p-2} \sinh x \, dx.$
7. $\int \sinh^2 x \, dx = -\frac{1}{2}x + \frac{1}{4} \sinh 2x.$
8. $\int \sinh^3 x \, dx = -\cosh x + \frac{1}{3} \cosh^3 x.$
9. $\int \sinh^{2n} x \, dx = (-1)^n C_{2n}^n \frac{x}{2^{2n}} + \frac{1}{2^{2n-1}} \sum_{k=0}^{n-1} (-1)^k C_{2n}^k \frac{\sinh[2(n-k)x]}{2(n-k)}, \quad n = 1, 2, \dots$

- $$10. \int \sinh^{2n+1} x \, dx = \frac{1}{2^{2n}} \sum_{k=0}^n (-1)^k C_{2n+1}^k \frac{\cosh[(2n-2k+1)x]}{2n-2k+1} \\ = \sum_{k=0}^n (-1)^{n+k} C_n^k \frac{\cosh^{2k+1} x}{2k+1}, \quad n = 1, 2, \dots$$
- $$11. \int \sinh^p x \, dx = \frac{1}{p} \sinh^{p-1} x \cosh x - \frac{p-1}{p} \int \sinh^{p-2} x \, dx.$$
- $$12. \int \sinh ax \sinh bx \, dx = \frac{1}{a^2 - b^2} (a \cosh ax \sinh bx - b \cosh bx \sinh ax).$$
- $$13. \int \frac{dx}{\sinh ax} = \frac{1}{a} \ln \left| \tanh \frac{ax}{2} \right|.$$
- $$14. \int \frac{dx}{\sinh^{2n} x} = \frac{\cosh x}{2n-1} \left[-\frac{1}{\sinh^{2n-1} x} \right. \\ \left. + \sum_{k=1}^{n-1} (-1)^{k-1} \frac{2^k (n-1)(n-2) \dots (n-k)}{(2n-3)(2n-5) \dots (2n-2k-1)} \frac{1}{\sinh^{2n-2k-1} x} \right], \quad n = 1, 2, \dots$$
- $$15. \int \frac{dx}{\sinh^{2n+1} x} = \frac{\cosh x}{2n} \left[-\frac{1}{\sinh^{2n} x} \right. \\ \left. + \sum_{k=1}^{n-1} (-1)^{k-1} \frac{(2n-1)(2n-3) \dots (2n-2k+1)}{2^k (n-1)(n-2) \dots (n-k)} \frac{1}{\sinh^{2n-2k} x} \right] + (-1)^n \frac{(2n-1)!!}{(2n)!!} \ln \tanh \frac{x}{2}, \\ n = 1, 2, \dots$$
- $$16. \int \frac{dx}{a + b \sinh x} = \frac{1}{\sqrt{a^2 + b^2}} \ln \frac{a \tanh(x/2) - b + \sqrt{a^2 + b^2}}{a \tanh(x/2) - b - \sqrt{a^2 + b^2}}.$$
- $$17. \int \frac{Ax + B \sinh x}{a + b \sinh x} \, dx = \frac{B}{b} x + \frac{Ab - Ba}{b\sqrt{a^2 + b^2}} \ln \frac{a \tanh(x/2) - b + \sqrt{a^2 + b^2}}{a \tanh(x/2) - b - \sqrt{a^2 + b^2}}.$$

3.4-3. Integrals Involving $\tanh x$ or $\coth x$.

1. $\int \tanh x \, dx = \ln \cosh x.$
2. $\int \tanh^2 x \, dx = x - \tanh x.$
3. $\int \tanh^3 x \, dx = -\frac{1}{2} \tanh^2 x + \ln \cosh x.$
4. $\int \tanh^{2n} x \, dx = x - \sum_{k=1}^n \frac{\tanh^{2n-2k+1} x}{2n-2k+1}, \quad n = 1, 2, \dots$
5. $\int \tanh^{2n+1} x \, dx = \ln \cosh x - \sum_{k=1}^n \frac{(-1)^k C_n^k}{2k \cosh^{2k} x} = \ln \cosh x - \sum_{k=1}^n \frac{\tanh^{2n-2k+2} x}{2n-2k+2}, \quad n = 1, 2, \dots$
6. $\int \tanh^p x \, dx = -\frac{1}{p-1} \tanh^{p-1} x + \int \tanh^{p-2} x \, dx.$
7. $\int \coth x \, dx = \ln |\sinh x|.$
8. $\int \coth^2 x \, dx = x - \coth x.$
9. $\int \coth^3 x \, dx = -\frac{1}{2} \coth^2 x + \ln |\sinh x|.$

10. $\int \coth^{2n} x \, dx = x - \sum_{k=1}^n \frac{\coth^{2n-2k+1} x}{2n-2k+1}, \quad n = 1, 2, \dots$
11. $\int \coth^{2n+1} x \, dx = \ln |\sinh x| - \sum_{k=1}^n \frac{C_n^k}{2k \sinh^{2k} x} = \ln |\sinh x| - \sum_{k=1}^n \frac{\coth^{2n-2k+2} x}{2n-2k+2}, \quad n = 1, 2, \dots$
12. $\int \coth^p x \, dx = -\frac{1}{p-1} \coth^{p-1} x + \int \coth^{p-2} x \, dx.$

3.5. Integrals Involving Logarithmic Functions

1. $\int \ln ax \, dx = x \ln ax - x.$
2. $\int x \ln x \, dx = \frac{1}{2} x^2 \ln x - \frac{1}{4} x^2.$
3. $\int x^p \ln ax \, dx = \begin{cases} \frac{1}{p+1} x^{p+1} \ln ax - \frac{1}{(p+1)^2} x^{p+1} & \text{if } p \neq -1, \\ \frac{1}{2} \ln^2 ax & \text{if } p = -1. \end{cases}$
4. $\int (\ln x)^2 \, dx = x(\ln x)^2 - 2x \ln x + 2x.$
5. $\int x(\ln x)^2 \, dx = \frac{1}{2} x^2 (\ln x)^2 - \frac{1}{2} x^2 \ln x + \frac{1}{4} x^2.$
6. $\int x^p (\ln x)^2 \, dx = \begin{cases} \frac{x^{p+1}}{p+1} (\ln x)^2 - \frac{2x^{p+1}}{(p+1)^2} \ln x + \frac{2x^{p+1}}{(p+1)^3} & \text{if } p \neq -1, \\ \frac{1}{3} \ln^3 x & \text{if } p = -1. \end{cases}$
7. $\int (\ln x)^n \, dx = \frac{x}{n+1} \sum_{k=0}^n (-1)^k (n+1)n \dots (n-k+1) (\ln x)^{n-k}, \quad n = 1, 2, \dots$
8. $\int (\ln x)^q \, dx = x(\ln x)^q - q \int (\ln x)^{q-1} \, dx, \quad q \neq -1.$
9. $\int x^n (\ln x)^m \, dx = \frac{x^{n+1}}{m+1} \sum_{k=0}^m \frac{(-1)^k}{(n+1)^{k+1}} (m+1)m \dots (m-k+1) (\ln x)^{m-k}, \quad n, m = 1, 2, \dots$
10. $\int x^p (\ln x)^q \, dx = \frac{1}{p+1} x^{p+1} (\ln x)^q - \frac{q}{p+1} \int x^p (\ln x)^{q-1} \, dx, \quad p, q \neq -1.$
11. $\int \ln(a+bx) \, dx = \frac{1}{b} (ax+b) \ln(ax+b) - x.$
12. $\int x \ln(a+bx) \, dx = \frac{1}{2} \left(x^2 - \frac{a^2}{b^2} \right) \ln(a+bx) - \frac{1}{2} \left(\frac{x^2}{2} - \frac{a}{b} x \right).$
13. $\int x^2 \ln(a+bx) \, dx = \frac{1}{3} \left(x^3 - \frac{a^3}{b^3} \right) \ln(a+bx) - \frac{1}{3} \left(\frac{x^3}{3} - \frac{ax^2}{2b} + \frac{a^2x}{b^2} \right).$
14. $\int \frac{\ln x \, dx}{(a+bx)^2} = -\frac{\ln x}{b(a+bx)} + \frac{1}{ab} \ln \frac{x}{a+bx}.$
15. $\int \frac{\ln x \, dx}{(a+bx)^3} = -\frac{\ln x}{2b(a+bx)^2} + \frac{1}{2ab(a+bx)} + \frac{1}{2a^2b} \ln \frac{x}{a+bx}.$
16. $\int \frac{\ln x \, dx}{\sqrt{a+bx}} = \begin{cases} \frac{2}{b} \left[(\ln x - 2)\sqrt{a+bx} + \sqrt{a} \ln \frac{\sqrt{a+bx} + \sqrt{a}}{\sqrt{a+bx} - \sqrt{a}} \right] & \text{if } a > 0, \\ \frac{2}{b} \left[(\ln x - 2)\sqrt{a+bx} + 2\sqrt{-a} \arctan \frac{\sqrt{a+bx}}{\sqrt{-a}} \right] & \text{if } a < 0. \end{cases}$

$$17. \int \ln(x^2 + a^2) dx = x \ln(x^2 + a^2) - 2x + 2a \arctan(x/a).$$

$$18. \int x \ln(x^2 + a^2) dx = \frac{1}{2} [(x^2 + a^2) \ln(x^2 + a^2) - x^2].$$

$$19. \int x^2 \ln(x^2 + a^2) dx = \frac{1}{3} [x^3 \ln(x^2 + a^2) - \frac{2}{3} x^3 + 2a^2 x - 2a^3 \arctan(x/a)].$$

3.6. Integrals Involving Trigonometric Functions

3.6-1. Integrals Involving $\cos x$ ($n = 1, 2, \dots$).

$$1. \int \cos(ax + b) dx = \frac{1}{a} \sin(ax + b).$$

$$2. \int x \cos x dx = \cos x + x \sin x.$$

$$3. \int x^2 \cos x dx = 2x \cos x + (x^2 - 2) \sin x.$$

$$4. \int x^{2n} \cos x dx = (2n)! \left[\sum_{k=0}^n (-1)^k \frac{x^{2n-2k}}{(2n-2k)!} \sin x + \sum_{k=0}^{n-1} (-1)^k \frac{x^{2n-2k-1}}{(2n-2k-1)!} \cos x \right].$$

$$5. \int x^{2n+1} \cos x dx = (2n+1)! \sum_{k=0}^n \left[(-1)^k \frac{x^{2n-2k+1}}{(2n-2k+1)!} \sin x + \frac{x^{2n-2k}}{(2n-2k)!} \cos x \right].$$

$$6. \int x^p \cos x dx = x^p \sin x + px^{p-1} \cos x - p(p-1) \int x^{p-2} \cos x dx.$$

$$7. \int \cos^2 x dx = \frac{1}{2} x + \frac{1}{4} \sin 2x.$$

$$8. \int \cos^3 x dx = \sin x - \frac{1}{3} \sin^3 x.$$

$$9. \int \cos^{2n} x dx = \frac{1}{2^{2n}} C_{2n}^n x + \frac{1}{2^{2n-1}} \sum_{k=0}^{n-1} C_{2n}^k \frac{\sin[(2n-2k)x]}{2n-2k}.$$

$$10. \int \cos^{2n+1} x dx = \frac{1}{2^{2n}} \sum_{k=0}^n C_{2n+1}^k \frac{\sin[(2n-2k+1)x]}{2n-2k+1}.$$

$$11. \int \frac{dx}{\cos x} = \ln \left| \tan \left(\frac{x}{2} + \frac{\pi}{4} \right) \right|.$$

$$12. \int \frac{dx}{\cos^2 x} = \tan x.$$

$$13. \int \frac{dx}{\cos^3 x} = \frac{\sin x}{2 \cos^2 x} + \frac{1}{2} \ln \left| \tan \left(\frac{x}{2} + \frac{\pi}{4} \right) \right|.$$

$$14. \int \frac{dx}{\cos^n x} = \frac{\sin x}{(n-1) \cos^{n-1} x} + \frac{n-2}{n-1} \int \frac{dx}{\cos^{n-2} x}, \quad n > 1.$$

$$15. \int \frac{x dx}{\cos^{2n} x} = \sum_{k=0}^{n-1} \frac{(2n-2)(2n-4) \dots (2n-2k+2)}{(2n-1)(2n-3) \dots (2n-2k+3)} \frac{(2n-2k)x \sin x - \cos x}{(2n-2k+1)(2n-2k) \cos^{2n-2k+1} x} + \frac{2^{n-1}(n-1)!}{(2n-1)!} (x \tan x + \ln |\cos x|).$$

$$16. \int \cos ax \cos bx dx = \frac{\sin[(b-a)x]}{2(b-a)} + \frac{\sin[(b+a)x]}{2(b+a)}, \quad a \neq \pm b.$$

- $$17. \int \frac{dx}{a + b \cos x} = \begin{cases} \frac{2}{\sqrt{a^2 - b^2}} \arctan \frac{(a-b) \tan(x/2)}{\sqrt{a^2 - b^2}} & \text{if } a^2 > b^2, \\ \frac{1}{\sqrt{b^2 - a^2}} \ln \left| \frac{\sqrt{b^2 - a^2} + (b-a) \tan(x/2)}{\sqrt{b^2 - a^2} - (b-a) \tan(x/2)} \right| & \text{if } b^2 > a^2. \end{cases}$$
- $$18. \int \frac{dx}{(a + b \cos x)^2} = \frac{b \sin x}{(b^2 - a^2)(a + b \cos x)} - \frac{a}{b^2 - a^2} \int \frac{dx}{a + b \cos x}.$$
- $$19. \int \frac{dx}{a^2 + b^2 \cos^2 x} = \frac{1}{a\sqrt{a^2 + b^2}} \arctan \frac{a \tan x}{\sqrt{a^2 + b^2}}.$$
- $$20. \int \frac{dx}{a^2 - b^2 \cos^2 x} = \begin{cases} \frac{1}{a\sqrt{a^2 - b^2}} \arctan \frac{a \tan x}{\sqrt{a^2 - b^2}} & \text{if } a^2 > b^2, \\ \frac{1}{2a\sqrt{b^2 - a^2}} \ln \left| \frac{\sqrt{b^2 - a^2} - a \tan x}{\sqrt{b^2 - a^2} + a \tan x} \right| & \text{if } b^2 > a^2. \end{cases}$$
- $$21. \int e^{ax} \cos bx \, dx = e^{ax} \left(\frac{b}{a^2 + b^2} \sin bx + \frac{a}{a^2 + b^2} \cos bx \right).$$
- $$22. \int e^{ax} \cos^2 x \, dx = \frac{e^{ax}}{a^2 + 4} \left(a \cos^2 x + 2 \sin x \cos x + \frac{2}{a} \right).$$
- $$23. \int e^{ax} \cos^n x \, dx = \frac{e^{ax} \cos^{n-1} x}{a^2 + n^2} (a \cos x + n \sin x) + \frac{n(n-1)}{a^2 + n^2} \int e^{ax} \cos^{n-2} x \, dx.$$

3.6-2. Integrals Involving $\sin x$ ($n = 1, 2, \dots$).

1. $\int \sin(a + bx) \, dx = -\frac{1}{b} \cos(a + bx).$
2. $\int x \sin x \, dx = \sin x - x \cos x.$
3. $\int x^2 \sin x \, dx = 2x \sin x - (x^2 - 2) \cos x.$
4. $\int x^3 \sin x \, dx = (3x^2 - 6) \sin x - (x^3 - 6x) \cos x.$
5. $\int x^{2n} \sin x \, dx = (2n)! \left[\sum_{k=0}^n (-1)^{k+1} \frac{x^{2n-2k}}{(2n-2k)!} \cos x + \sum_{k=0}^{n-1} (-1)^k \frac{x^{2n-2k-1}}{(2n-2k-1)!} \sin x \right].$
6. $\int x^{2n+1} \sin x \, dx = (2n+1)! \sum_{k=0}^n \left[(-1)^{k+1} \frac{x^{2n-2k+1}}{(2n-2k+1)!} \cos x + (-1)^k \frac{x^{2n-2k}}{(2n-2k)!} \sin x \right].$
7. $\int x^p \sin x \, dx = -x^p \cos x + px^{p-1} \sin x - p(p-1) \int x^{p-2} \sin x \, dx.$
8. $\int \sin^2 x \, dx = \frac{1}{2}x - \frac{1}{4} \sin 2x.$
9. $\int x \sin^2 x \, dx = \frac{1}{4}x^2 - \frac{1}{4}x \sin 2x - \frac{1}{8} \cos 2x.$
10. $\int \sin^3 x \, dx = -\cos x + \frac{1}{3} \cos^3 x.$
11. $\int \sin^{2n} x \, dx = \frac{1}{2^{2n}} C_{2n}^n x + \frac{(-1)^n}{2^{2n-1}} \sum_{k=0}^{n-1} (-1)^k C_{2n}^k \frac{\sin[(2n-2k)x]}{2n-2k},$

where $C_m^k = \frac{m!}{k!(m-k)!}$ are binomial coefficients ($0! = 1$).

12. $\int \sin^{2n+1} x \, dx = \frac{1}{2^{2n}} \sum_{k=0}^n (-1)^{n+k+1} C_{2n+1}^k \frac{\cos[(2n-2k+1)x]}{2n-2k+1}.$
13. $\int \frac{dx}{\sin x} = \ln \left| \tan \frac{x}{2} \right|.$
14. $\int \frac{dx}{\sin^2 x} = -\cot x.$
15. $\int \frac{dx}{\sin^3 x} = -\frac{\cos x}{2 \sin^2 x} + \frac{1}{2} \ln \left| \tan \frac{x}{2} \right|.$
16. $\int \frac{dx}{\sin^n x} = -\frac{\cos x}{(n-1) \sin^{n-1} x} + \frac{n-2}{n-1} \int \frac{dx}{\sin^{n-2} x}, \quad n > 1.$
17. $\int \frac{x \, dx}{\sin^{2n} x} = -\sum_{k=0}^{n-1} \frac{(2n-2)(2n-4) \dots (2n-2k+2)}{(2n-1)(2n-3) \dots (2n-2k+3)} \frac{\sin x + (2n-2k)x \cos x}{(2n-2k+1)(2n-2k) \sin^{2n-2k+1} x} + \frac{2^{n-1}(n-1)!}{(2n-1)!!} (\ln |\sin x| - x \cot x).$
18. $\int \sin ax \sin bx \, dx = \frac{\sin[(b-a)x]}{2(b-a)} - \frac{\sin[(b+a)x]}{2(b+a)}, \quad a \neq \pm b.$
19. $\int \frac{dx}{a+b \sin x} = \begin{cases} \frac{2}{\sqrt{a^2-b^2}} \arctan \frac{b+a \tan x/2}{\sqrt{a^2-b^2}} & \text{if } a^2 > b^2, \\ \frac{1}{\sqrt{b^2-a^2}} \ln \left| \frac{b-\sqrt{b^2-a^2}+a \tan x/2}{b+\sqrt{b^2-a^2}+a \tan x/2} \right| & \text{if } b^2 > a^2. \end{cases}$
20. $\int \frac{dx}{(a+b \sin x)^2} = \frac{b \cos x}{(a^2-b^2)(a+b \sin x)} + \frac{a}{a^2-b^2} \int \frac{dx}{a+b \sin x}.$
21. $\int \frac{dx}{a^2+b^2 \sin^2 x} = \frac{1}{a\sqrt{a^2+b^2}} \arctan \frac{\sqrt{a^2+b^2} \tan x}{a}.$
22. $\int \frac{dx}{a^2-b^2 \sin^2 x} = \begin{cases} \frac{1}{a\sqrt{a^2-b^2}} \arctan \frac{\sqrt{a^2-b^2} \tan x}{a} & \text{if } a^2 > b^2, \\ \frac{1}{2a\sqrt{b^2-a^2}} \ln \left| \frac{\sqrt{b^2-a^2} \tan x + a}{\sqrt{b^2-a^2} \tan x - a} \right| & \text{if } b^2 > a^2. \end{cases}$
23. $\int \frac{\sin x \, dx}{\sqrt{1+k^2 \sin^2 x}} = -\frac{1}{k} \arcsin \frac{k \cos x}{\sqrt{1+k^2}}.$
24. $\int \frac{\sin x \, dx}{\sqrt{1-k^2 \sin^2 x}} = -\frac{1}{k} \ln |k \cos x + \sqrt{1-k^2 \sin^2 x}|.$
25. $\int \sin x \sqrt{1+k^2 \sin^2 x} \, dx = -\frac{\cos x}{2} \sqrt{1+k^2 \sin^2 x} - \frac{1+k^2}{2k} \arcsin \frac{k \cos x}{\sqrt{1+k^2}}.$
26. $\int \sin x \sqrt{1-k^2 \sin^2 x} \, dx = -\frac{\cos x}{2} \sqrt{1-k^2 \sin^2 x} - \frac{1-k^2}{2k} \ln |k \cos x + \sqrt{1-k^2 \sin^2 x}|.$
27. $\int e^{ax} \sin bx \, dx = e^{ax} \left(\frac{a}{a^2+b^2} \sin bx - \frac{b}{a^2+b^2} \cos bx \right).$
28. $\int e^{ax} \sin^2 x \, dx = \frac{e^{ax}}{a^2+4} \left(a \sin^2 x - 2 \sin x \cos x + \frac{2}{a} \right).$
29. $\int e^{ax} \sin^n x \, dx = \frac{e^{ax} \sin^{n-1} x}{a^2+n^2} (a \sin x - n \cos x) + \frac{n(n-1)}{a^2+n^2} \int e^{ax} \sin^{n-2} x \, dx.$

3.6-3. Integrals Involving $\sin x$ and $\cos x$.

- $\int \sin ax \cos bx \, dx = -\frac{\cos[(a+b)x]}{2(a+b)} - \frac{\cos[(a-b)x]}{2(a-b)}, \quad a \neq \pm b.$
- $\int \frac{dx}{b^2 \cos^2 ax + c^2 \sin^2 ax} = \frac{1}{abc} \arctan\left(\frac{c}{b} \tan ax\right).$
- $\int \frac{dx}{b^2 \cos^2 ax - c^2 \sin^2 ax} = \frac{1}{2abc} \ln \left| \frac{c \tan ax + b}{c \tan ax - b} \right|.$
- $\int \frac{dx}{\cos^{2n} x \sin^{2m} x} = \sum_{k=0}^{n+m-1} C_{n+m-1}^k \frac{\tan^{2k-2m+1} x}{2k-2m+1}, \quad n, m = 1, 2, \dots$
- $\int \frac{dx}{\cos^{2n+1} x \sin^{2m+1} x} = C_{n+m}^m \ln |\tan x| + \sum_{k=0}^{n+m} C_{n+m}^k \frac{\tan^{2k-2m} x}{2k-2m}, \quad n, m = 1, 2, \dots$

3.6-4. Reduction Formulas.

► The parameters p and q below can assume any values, except for those at which the denominators on the right-hand side vanish.

- $\int \sin^p x \cos^q x \, dx = -\frac{\sin^{p-1} x \cos^{q+1} x}{p+q} + \frac{p-1}{p+q} \int \sin^{p-2} x \cos^q x \, dx.$
- $\int \sin^p x \cos^q x \, dx = \frac{\sin^{p+1} x \cos^{q-1} x}{p+q} + \frac{q-1}{p+q} \int \sin^p x \cos^{q-2} x \, dx.$
- $\int \sin^p x \cos^q x \, dx = \frac{\sin^{p-1} x \cos^{q-1} x}{p+q} \left(\sin^2 x - \frac{q-1}{p+q-2} \right) + \frac{(p-1)(q-1)}{(p+q)(p+q-2)} \int \sin^{p-2} x \cos^{q-2} x \, dx.$
- $\int \sin^p x \cos^q x \, dx = \frac{\sin^{p+1} x \cos^{q+1} x}{p+1} + \frac{p+q+2}{p+1} \int \sin^{p+2} x \cos^q x \, dx.$
- $\int \sin^p x \cos^q x \, dx = -\frac{\sin^{p+1} x \cos^{q+1} x}{q+1} + \frac{p+q+2}{q+1} \int \sin^p x \cos^{q+2} x \, dx.$
- $\int \sin^p x \cos^q x \, dx = -\frac{\sin^{p-1} x \cos^{q+1} x}{q+1} + \frac{p-1}{q+1} \int \sin^{p-2} x \cos^{q+2} x \, dx.$
- $\int \sin^p x \cos^q x \, dx = \frac{\sin^{p+1} x \cos^{q-1} x}{p+1} + \frac{q-1}{p+1} \int \sin^{p+2} x \cos^{q-2} x \, dx.$

3.6-5. Integrals Involving $\tan x$ and $\cot x$.

- $\int \tan x \, dx = -\ln |\cos x|.$
- $\int \tan^2 x \, dx = \tan x - x.$
- $\int \tan^3 x \, dx = \frac{1}{2} \tan^2 x + \ln |\cos x|.$
- $\int \tan^{2n} x \, dx = (-1)^n x - \sum_{k=1}^n \frac{(-1)^k (\tan x)^{2n-2k+1}}{2n-2k+1}, \quad n = 1, 2, \dots$

5. $\int \tan^{2n+1} x \, dx = (-1)^{n+1} \ln |\cos x| - \sum_{k=1}^n \frac{(-1)^k (\tan x)^{2n-2k+2}}{2n-2k+2}, \quad n = 1, 2, \dots$
6. $\int \frac{dx}{a + b \tan x} = \frac{1}{a^2 + b^2} (ax + b \ln |a \cos x + b \sin x|).$
7. $\int \frac{\tan x \, dx}{\sqrt{a + b \tan^2 x}} = \frac{1}{\sqrt{b-a}} \arccos \left(\sqrt{1 - \frac{a}{b}} \cos x \right), \quad b > a, b > 0.$
8. $\int \cot x \, dx = \ln |\sin x|.$
9. $\int \cot^2 x \, dx = -\cot x - x.$
10. $\int \cot^3 x \, dx = -\frac{1}{2} \cot^2 x - \ln |\sin x|.$
11. $\int \cot^{2n} x \, dx = (-1)^n x + \sum_{k=1}^n \frac{(-1)^k (\cot x)^{2n-2k+1}}{2n-2k+1}, \quad n = 1, 2, \dots$
12. $\int \cot^{2n+1} x \, dx = (-1)^n \ln |\sin x| + \sum_{k=1}^n \frac{(-1)^k (\cot x)^{2n-2k+2}}{2n-2k+2}, \quad n = 1, 2, \dots$
13. $\int \frac{dx}{a + b \cot x} = \frac{1}{a^2 + b^2} (ax - b \ln |a \sin x + b \cos x|).$

3.7. Integrals Involving Inverse Trigonometric Functions

1. $\int \arcsin \frac{x}{a} \, dx = x \arcsin \frac{x}{a} + \sqrt{a^2 - x^2}.$
2. $\int \left(\arcsin \frac{x}{a} \right)^2 \, dx = x \left(\arcsin \frac{x}{a} \right)^2 - 2x + 2\sqrt{a^2 - x^2} \arcsin \frac{x}{a}.$
3. $\int x \arcsin \frac{x}{a} \, dx = \frac{1}{4}(2x^2 - a^2) \arcsin \frac{x}{a} + \frac{x}{4} \sqrt{a^2 - x^2}.$
4. $\int x^2 \arcsin \frac{x}{a} \, dx = \frac{x^3}{3} \arcsin \frac{x}{a} + \frac{1}{9}(x^2 + 2a^2) \sqrt{a^2 - x^2}.$
5. $\int \arccos \frac{x}{a} \, dx = x \arccos \frac{x}{a} - \sqrt{a^2 - x^2}.$
6. $\int \left(\arccos \frac{x}{a} \right)^2 \, dx = x \left(\arccos \frac{x}{a} \right)^2 - 2x - 2\sqrt{a^2 - x^2} \arccos \frac{x}{a}.$
7. $\int x \arccos \frac{x}{a} \, dx = \frac{1}{4}(2x^2 - a^2) \arccos \frac{x}{a} - \frac{x}{4} \sqrt{a^2 - x^2}.$
8. $\int x^2 \arccos \frac{x}{a} \, dx = \frac{x^3}{3} \arccos \frac{x}{a} - \frac{1}{9}(x^2 + 2a^2) \sqrt{a^2 - x^2}.$
9. $\int \arctan \frac{x}{a} \, dx = x \arctan \frac{x}{a} - \frac{a}{2} \ln(a^2 + x^2).$
10. $\int x \arctan \frac{x}{a} \, dx = \frac{1}{2}(x^2 + a^2) \arctan \frac{x}{a} - \frac{ax}{2}.$
11. $\int x^2 \arctan \frac{x}{a} \, dx = \frac{x^3}{3} \arctan \frac{x}{a} - \frac{ax^2}{6} + \frac{a^3}{6} \ln(a^2 + x^2).$

-
12. $\int \operatorname{arccot} \frac{x}{a} dx = x \operatorname{arccot} \frac{x}{a} + \frac{a}{2} \ln(a^2 + x^2).$
13. $\int x \operatorname{arccot} \frac{x}{a} dx = \frac{1}{2}(x^2 + a^2) \operatorname{arccot} \frac{x}{a} + \frac{ax}{2}.$
14. $\int x^2 \operatorname{arccot} \frac{x}{a} dx = \frac{x^3}{3} \operatorname{arccot} \frac{x}{a} + \frac{ax^2}{6} - \frac{a^3}{6} \ln(a^2 + x^2).$

© References for Supplement 3: H. B. Dwight (1961), I. S. Gradshteyn and I. M. Ryzhik (2000), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1986, 1988), D. Zwillinger (2002), I. N. Bronshtein and K. A. Semendyayev (2004).

Supplement 4

Tables of Definite Integrals

► Throughout Supplement 4 it is assumed that n is a positive integer, unless otherwise specified.

4.1. Integrals Involving Power-Law Functions

4.1-1. Integrals Over a Finite Interval.
--

$$1. \int_0^1 \frac{x^n dx}{x+1} = (-1)^n \left[\ln 2 + \sum_{k=1}^n \frac{(-1)^k}{k} \right].$$

$$2. \int_0^1 \frac{dx}{x^2 + 2x \cos \beta + 1} = \frac{\beta}{2 \sin \beta}.$$

$$3. \int_0^1 \frac{(x^a + x^{-a}) dx}{x^2 + 2x \cos \beta + 1} = \frac{\pi \sin(a\beta)}{\sin(\pi a) \sin \beta}, \quad |a| < 1, \beta \neq (2n+1)\pi.$$

$$4. \int_0^1 x^a (1-x)^{1-a} dx = \frac{\pi a(1-a)}{2 \sin(\pi a)}, \quad -1 < a < 1.$$

$$5. \int_0^1 \frac{dx}{x^a (1-x)^{1-a}} = \frac{\pi}{\sin(\pi a)}, \quad 0 < a < 1.$$

$$6. \int_0^1 \frac{x^a dx}{(1-x)^a} = \frac{\pi a}{\sin(\pi a)}, \quad -1 < a < 1.$$

$$7. \int_0^1 x^{p-1} (1-x)^{q-1} dx \equiv B(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}, \quad p, q > 0.$$

$$8. \int_0^1 x^{p-1} (1-x^q)^{-p/q} dx = \frac{\pi}{q \sin(\pi p/q)}, \quad q > p > 0.$$

$$9. \int_0^1 x^{p+q-1} (1-x^q)^{-p/q} dx = \frac{\pi p}{q^2 \sin(\pi p/q)}, \quad q > p.$$

$$10. \int_0^1 x^{q/p-1} (1-x^q)^{-1/p} dx = \frac{\pi}{q \sin(\pi/p)}, \quad p > 1, q > 0.$$

$$11. \int_0^1 \frac{x^{p-1} - x^{-p}}{1-x} dx = \pi \cot(\pi p), \quad |p| < 1.$$

$$12. \int_0^1 \frac{x^{p-1} - x^{-p}}{1+x} dx = \frac{\pi}{\sin(\pi p)}, \quad |p| < 1.$$

13. $\int_0^1 \frac{x^p - x^{-p}}{x-1} dx = \frac{1}{p} - \pi \cot(\pi p), \quad |p| < 1.$
14. $\int_0^1 \frac{x^p - x^{-p}}{1+x} dx = \frac{1}{p} - \frac{\pi}{\sin(\pi p)}, \quad |p| < 1.$
15. $\int_0^1 \frac{x^{1+p} - x^{1-p}}{1-x^2} dx = \frac{\pi}{2} \cot\left(\frac{\pi p}{2}\right) - \frac{1}{p}, \quad |p| < 1.$
16. $\int_0^1 \frac{x^{1+p} - x^{1-p}}{1+x^2} dx = \frac{1}{p} - \frac{\pi}{2 \sin(\pi p/2)}, \quad |p| < 1.$
17. $\int_0^1 \frac{dx}{\sqrt{(1+a^2x)(1-x)}} = \frac{2}{a} \arctan a.$
18. $\int_0^1 \frac{dx}{\sqrt{(1-a^2x)(1-x)}} = \frac{1}{a} \ln \frac{1+a}{1-a}.$
19. $\int_{-1}^1 \frac{dx}{(a-x)\sqrt{1-x^2}} = \frac{\pi}{\sqrt{a^2-1}}, \quad 1 < a.$
20. $\int_0^1 \frac{x^n dx}{\sqrt{1-x}} = \frac{2(2n)!!}{(2n+1)!!}, \quad n = 1, 2, \dots$
21. $\int_0^1 \frac{x^{n-1/2} dx}{\sqrt{1-x}} = \frac{\pi(2n-1)!!}{(2n)!!}, \quad n = 1, 2, \dots$
22. $\int_0^1 \frac{x^{2n} dx}{\sqrt{1-x^2}} = \frac{\pi}{2} \frac{1 \times 3 \times \dots \times (2n-1)}{2 \times 4 \times \dots \times (2n)}, \quad n = 1, 2, \dots$
23. $\int_0^1 \frac{x^{2n+1} dx}{\sqrt{1-x^2}} = \frac{2 \times 4 \times \dots \times (2n)}{1 \times 3 \times \dots \times (2n+1)}, \quad n = 1, 2, \dots$
24. $\int_0^1 \frac{x^{\lambda-1} dx}{(1+ax)(1-x)^\lambda} = \frac{\pi}{(1+a)^\lambda \sin(\pi\lambda)}, \quad 0 < \lambda < 1, \quad a > -1.$
25. $\int_0^1 \frac{x^{\lambda-1/2} dx}{(1+ax)^\lambda(1-x)^\lambda} = 2\pi^{-1/2} \Gamma(\lambda + \frac{1}{2}) \Gamma(1-\lambda) \cos^{2\lambda} k \frac{\sin[(2\lambda-1)k]}{(2\lambda-1) \sin k}, \quad k = \arctan \sqrt{a},$
 $-\frac{1}{2} < \lambda < 1, \quad a > 0.$

4.1-2. Integrals Over an Infinite Interval.

1. $\int_0^\infty \frac{dx}{ax^2 + b} = \frac{\pi}{2\sqrt{ab}}.$
2. $\int_0^\infty \frac{dx}{x^4 + 1} = \frac{\pi\sqrt{2}}{4}.$
3. $\int_0^\infty \frac{x^{a-1} dx}{x+1} = \frac{\pi}{\sin(\pi a)}, \quad 0 < a < 1.$
4. $\int_0^\infty \frac{x^{\lambda-1} dx}{(1+ax)^2} = \frac{\pi(1-\lambda)}{a^\lambda \sin(\pi\lambda)}, \quad 0 < \lambda < 2.$
5. $\int_0^\infty \frac{x^{\lambda-1} dx}{(x+a)(x+b)} = \frac{\pi(a^{\lambda-1} - b^{\lambda-1})}{(b-a) \sin(\pi\lambda)}, \quad 0 < \lambda < 2.$

6. $\int_0^{\infty} \frac{x^{\lambda-1}(x+c) dx}{(x+a)(x+b)} = \frac{\pi}{\sin(\pi\lambda)} \left(\frac{a-c}{a-b} a^{\lambda-1} + \frac{b-c}{b-a} b^{\lambda-1} \right), \quad 0 < \lambda < 1.$
7. $\int_0^{\infty} \frac{x^{\lambda} dx}{(x+1)^3} = \frac{\pi\lambda(1-\lambda)}{2\sin(\pi\lambda)}, \quad -1 < \lambda < 2.$
8. $\int_0^{\infty} \frac{x^{\lambda-1} dx}{(x^2+a^2)(x^2+b^2)} = \frac{\pi(b^{\lambda-2} - a^{\lambda-2})}{2(a^2 - b^2)\sin(\pi\lambda/2)}, \quad 0 < \lambda < 4.$
9. $\int_0^{\infty} \frac{x^{p-1} - x^{q-1}}{1-x} dx = \pi[\cot(\pi p) - \cot(\pi q)], \quad p, q > 0.$
10. $\int_0^{\infty} \frac{x^{\lambda-1} dx}{(1+ax)^{n+1}} = (-1)^n \frac{\pi C_{\lambda-1}^n}{a^{\lambda} \sin(\pi\lambda)}, \quad 0 < \lambda < n+1, \quad C_{\lambda-1}^n = \frac{(\lambda-1)(\lambda-2)\dots(\lambda-n)}{n!}.$
11. $\int_0^{\infty} \frac{x^m dx}{(a+bx)^{n+1/2}} = 2^{m+1} m! \frac{(2n-2m-3)!!}{(2n-1)!!} \frac{a^{m-n+1/2}}{b^{m+1}}, \quad a, b > 0, \quad n, m = 1, 2, \dots,$
 $m < b - \frac{1}{2}.$
12. $\int_0^{\infty} \frac{dx}{(x^2+a^2)^n} = \frac{\pi}{2} \frac{(2n-3)!!}{(2n-2)!!} \frac{1}{a^{2n-1}}, \quad n = 1, 2, \dots$
13. $\int_0^{\infty} \frac{(x+1)^{\lambda-1}}{(x+a)^{\lambda+1}} dx = \frac{1-a^{-\lambda}}{\lambda(a-1)}, \quad a > 0.$
14. $\int_0^{\infty} \frac{x^{a-1} dx}{x^b+1} = \frac{\pi}{b\sin(\pi a/b)}, \quad 0 < a \leq b.$
15. $\int_0^{\infty} \frac{x^{a-1} dx}{(x^b+1)^2} = \frac{\pi(a-b)}{b^2 \sin[\pi(a-b)/b]}, \quad a < 2b.$
16. $\int_0^{\infty} \frac{x^{\lambda-1/2} dx}{(x+a)^{\lambda}(x+b)^{\lambda}} = \sqrt{\pi}(\sqrt{a} + \sqrt{b})^{1-2\lambda} \frac{\Gamma(\lambda-1/2)}{\Gamma(\lambda)}, \quad \lambda > 0.$
17. $\int_0^{\infty} \frac{1-x^a}{1-x^b} x^{c-1} dx = \frac{\pi \sin A}{b \sin C \sin(A+C)}, \quad A = \frac{\pi a}{b}, \quad C = \frac{\pi c}{b}; \quad a+c < b, \quad c > 0.$
18. $\int_0^{\infty} \frac{x^{a-1} dx}{(1+x^2)^{1-b}} = \frac{1}{2} B\left(\frac{1}{2}a, 1-b-\frac{1}{2}a\right), \quad \frac{1}{2}a+b < 1, \quad a > 0.$
19. $\int_0^{\infty} \frac{x^{2m} dx}{(ax^2+b)^n} = \frac{\pi(2m-1)!!(2n-2m-3)!!}{2(2n-2)!! a^m b^{n-m-1} \sqrt{ab}}, \quad a, b > 0, \quad n > m+1.$
20. $\int_0^{\infty} \frac{x^{2m+1} dx}{(ax^2+b)^n} = \frac{m!(n-m-2)!}{2(n-1)! a^{m+1} b^{n-m-1}}, \quad ab > 0, \quad n > m+1 \geq 1.$
21. $\int_0^{\infty} \frac{x^{\mu-1} dx}{(1+ax^p)^{\nu}} = \frac{1}{pa^{\mu/p}} B\left(\frac{\mu}{p}, \nu - \frac{\mu}{p}\right), \quad p > 0, \quad 0 < \mu < p\nu.$
22. $\int_0^{\infty} (\sqrt{x^2+a^2} - x)^n dx = \frac{na^{n+1}}{n^2-1}, \quad n = 2, 3, \dots$
23. $\int_0^{\infty} \frac{dx}{(x+\sqrt{x^2+a^2})^n} = \frac{n}{a^{n-1}(n^2-1)}, \quad n = 2, 3, \dots$
24. $\int_0^{\infty} x^m (\sqrt{x^2+a^2} - x)^n dx = \frac{m! na^{n+m+1}}{(n-m-1)(n-m+1)\dots(n+m+1)}, \quad n, m = 1, 2, \dots,$
 $0 \leq m \leq n-2.$
25. $\int_0^{\infty} \frac{x^m dx}{(x+\sqrt{x^2+a^2})^n} = \frac{m! n}{(n-m-1)(n-m+1)\dots(n+m+1)a^{n-m-1}}, \quad n = 2, 3, \dots$

4.2. Integrals Involving Exponential Functions

1. $\int_0^{\infty} e^{-ax} dx = \frac{1}{a}, \quad a > 0.$
2. $\int_0^1 x^n e^{-ax} dx = \frac{n!}{a^{n+1}} - e^{-a} \sum_{k=0}^n \frac{n!}{k!} \frac{1}{a^{n-k+1}}, \quad a > 0, \quad n = 1, 2, \dots$
3. $\int_0^{\infty} x^n e^{-ax} dx = \frac{n!}{a^{n+1}}, \quad a > 0, \quad n = 1, 2, \dots$
4. $\int_0^{\infty} \frac{e^{-ax}}{\sqrt{x}} dx = \sqrt{\frac{\pi}{a}}, \quad a > 0.$
5. $\int_0^{\infty} x^{\nu-1} e^{-\mu x} dx = \frac{\Gamma(\nu)}{\mu^{\nu}}, \quad \mu, \nu > 0.$
6. $\int_0^{\infty} \frac{dx}{1 + e^{ax}} = \frac{\ln 2}{a}.$
7. $\int_0^{\infty} \frac{x^{2n-1} dx}{e^{px} - 1} = (-1)^{n-1} \left(\frac{2\pi}{p}\right)^{2n} \frac{B_{2n}}{4n}, \quad n = 1, 2, \dots; \quad \text{the } B_m \text{ are Bernoulli numbers (see Supplement 11.1-3).}$
8. $\int_0^{\infty} \frac{x^{2n-1} dx}{e^{px} + 1} = (1 - 2^{1-2n}) \left(\frac{2\pi}{p}\right)^{2n} \frac{|B_{2n}|}{4n}, \quad n = 1, 2, \dots; \quad \text{the } B_m \text{ are Bernoulli numbers.}$
9. $\int_{-\infty}^{\infty} \frac{e^{-px} dx}{1 + e^{-qx}} = \frac{\pi}{q \sin(\pi p/q)}, \quad q > p > 0 \text{ or } 0 > p > q.$
10. $\int_0^{\infty} \frac{e^{ax} + e^{-ax}}{e^{bx} + e^{-bx}} dx = \frac{\pi}{2b \cos\left(\frac{\pi a}{2b}\right)}, \quad b > a.$
11. $\int_0^{\infty} \frac{e^{-px} - e^{-qx}}{1 - e^{-(p+q)x}} dx = \frac{\pi}{p+q} \cot \frac{\pi p}{p+q}, \quad p, q > 0.$
12. $\int_0^{\infty} (1 - e^{-\beta x})^{\nu} e^{-\mu x} dx = \frac{1}{\beta} B\left(\frac{\mu}{\beta}, \nu + 1\right).$
13. $\int_0^{\infty} \exp(-ax^2) dx = \frac{1}{2} \sqrt{\frac{\pi}{a}}, \quad a > 0.$
14. $\int_0^{\infty} x^{2n+1} \exp(-ax^2) dx = \frac{n!}{2a^{n+1}}, \quad a > 0, \quad n = 1, 2, \dots$
15. $\int_0^{\infty} x^{2n} \exp(-ax^2) dx = \frac{1 \times 3 \times \dots \times (2n-1) \sqrt{\pi}}{2^{n+1} a^{n+1/2}}, \quad a > 0, \quad n = 1, 2, \dots$
16. $\int_{-\infty}^{\infty} \exp(-a^2 x^2 \pm bx) dx = \frac{\sqrt{\pi}}{|a|} \exp\left(\frac{b^2}{4a^2}\right).$
17. $\int_0^{\infty} \exp\left(-ax^2 - \frac{b}{x^2}\right) dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} \exp(-2\sqrt{ab}), \quad a, b > 0.$
18. $\int_0^{\infty} \exp(-x^a) dx = \frac{1}{a} \Gamma\left(\frac{1}{a}\right), \quad a > 0.$

4.3. Integrals Involving Hyperbolic Functions

1. $\int_0^{\infty} \frac{dx}{\cosh ax} = \frac{\pi}{2|a|}$.
2. $\int_0^{\infty} \frac{dx}{a + b \cosh x} = \begin{cases} \frac{2}{\sqrt{b^2 - a^2}} \arctan \frac{\sqrt{b^2 - a^2}}{a + b} & \text{if } |b| > |a|, \\ \frac{1}{\sqrt{a^2 - b^2}} \ln \frac{a + b + \sqrt{a^2 - b^2}}{a + b - \sqrt{a^2 - b^2}} & \text{if } |b| < |a|. \end{cases}$
3. $\int_0^{\infty} \frac{x^{2n} dx}{\cosh ax} = \left(\frac{\pi}{2a}\right)^{2n+1} |E_{2n}|$, $a > 0$; the E_m are Euler numbers (see Supplement 11.1-4).
4. $\int_0^{\infty} \frac{x^{2n} dx}{\cosh^2 ax} = \frac{\pi^{2n}(2^{2n} - 2)}{|a|(2a)^{2n}} |B_{2n}|$, the B_m are Bernoulli numbers (see Supplement 11.1-3).
5. $\int_0^{\infty} \frac{\cosh ax}{\cosh bx} dx = \frac{\pi}{2b \cos\left(\frac{\pi a}{2b}\right)}$, $b > |a|$.
6. $\int_0^{\infty} x^{2n} \frac{\cosh ax}{\cosh bx} dx = \frac{\pi}{2b} \frac{d^{2n}}{da^{2n}} \frac{1}{\cos\left(\frac{1}{2}\pi a/b\right)}$, $b > |a|$, $n = 1, 2, \dots$
7. $\int_0^{\infty} \frac{\cosh ax \cosh bx}{\cosh(cx)} dx = \frac{\pi}{c} \frac{\cos\left(\frac{\pi a}{2c}\right) \cos\left(\frac{\pi b}{2c}\right)}{\cos\left(\frac{\pi a}{c}\right) + \cos\left(\frac{\pi b}{c}\right)}$, $c > |a| + |b|$.
8. $\int_0^{\infty} \frac{x dx}{\sinh ax} = \frac{\pi^2}{2a^2}$, $a > 0$.
9. $\int_0^{\infty} \frac{dx}{a + b \sinh x} = \frac{1}{\sqrt{a^2 + b^2}} \ln \frac{a + b + \sqrt{a^2 + b^2}}{a + b - \sqrt{a^2 + b^2}}$, $ab \neq 0$.
10. $\int_0^{\infty} \frac{\sinh ax}{\sinh bx} dx = \frac{\pi}{2b} \tan\left(\frac{\pi a}{2b}\right)$, $b > |a|$.
11. $\int_0^{\infty} x^{2n} \frac{\sinh ax}{\sinh bx} dx = \frac{\pi}{2b} \frac{d^{2n}}{dx^{2n}} \tan\left(\frac{\pi a}{2b}\right)$, $b > |a|$, $n = 1, 2, \dots$
12. $\int_0^{\infty} \frac{x^{2n}}{\sinh^2 ax} dx = \frac{\pi^{2n}}{a^{2n+1}} |B_{2n}|$, $a > 0$; the B_m are Bernoulli numbers.

4.4. Integrals Involving Logarithmic Functions

1. $\int_0^1 x^{a-1} \ln^n x dx = (-1)^n n! a^{-n-1}$, $a > 0$, $n = 1, 2, \dots$
2. $\int_0^1 \frac{\ln x}{x+1} dx = -\frac{\pi^2}{12}$.
3. $\int_0^1 \frac{x^n \ln x}{x+1} dx = (-1)^{n+1} \left[\frac{\pi^2}{12} + \sum_{k=1}^n \frac{(-1)^k}{k^2} \right]$, $n = 1, 2, \dots$
4. $\int_0^1 \frac{x^{\mu-1} \ln x}{x+a} dx = \frac{\pi a^{\mu-1}}{\sin(\pi\mu)} [\ln a - \pi \cot(\pi\mu)]$, $0 < \mu < 1$.

5. $\int_0^1 |\ln x|^\mu dx = \Gamma(\mu + 1), \quad \mu > -1.$
6. $\int_0^\infty x^{\mu-1} \ln(1+ax) dx = \frac{\pi}{\mu a^\mu \sin(\pi\mu)}, \quad -1 < \mu < 0.$
7. $\int_0^1 x^{2n-1} \ln(1+x) dx = \frac{1}{2n} \sum_{k=1}^{2n} \frac{(-1)^{k-1}}{k}, \quad n = 1, 2, \dots$
8. $\int_0^1 x^{2n} \ln(1+x) dx = \frac{1}{2n+1} \left[\ln 4 + \sum_{k=1}^{2n+1} \frac{(-1)^k}{k} \right], \quad n = 0, 1, \dots$
9. $\int_0^1 x^{n-1/2} \ln(1+x) dx = \frac{2 \ln 2}{2n+1} + \frac{4(-1)^n}{2n+1} \left[\pi - \sum_{k=0}^n \frac{(-1)^k}{2k+1} \right], \quad n = 1, 2, \dots$
10. $\int_0^\infty \ln \frac{a^2+x^2}{b^2+x^2} dx = \pi(a-b), \quad a, b > 0.$
11. $\int_0^\infty \frac{x^{p-1} \ln x}{1+x^q} dx = -\frac{\pi^2 \cos(\pi p/q)}{q^2 \sin^2(\pi p/q)}, \quad 0 < p < q.$
12. $\int_0^\infty e^{-\mu x} \ln x dx = -\frac{1}{\mu} (\mathcal{C} + \ln \mu), \quad \mu > 0, \quad \mathcal{C} = 0.5772 \dots$

4.5. Integrals Involving Trigonometric Functions

4.5-1. Integrals Over a Finite Interval.

1. $\int_0^{\pi/2} \cos^{2n} x dx = \frac{\pi}{2} \frac{1 \times 3 \times \dots \times (2n-1)}{2 \times 4 \times \dots \times (2n)}, \quad n = 1, 2, \dots$
2. $\int_0^{\pi/2} \cos^{2n+1} x dx = \frac{2 \times 4 \times \dots \times (2n)}{1 \times 3 \times \dots \times (2n+1)}, \quad n = 1, 2, \dots$
3. $\int_0^{\pi/2} x \cos^n x dx = -\sum_{k=0}^{m-1} \frac{(n-2k+1)(n-2k+3)\dots(n-1)}{(n-2k)(n-2k+2)\dots n} \frac{1}{n-2k}$

$$+ \begin{cases} \frac{\pi}{2} \frac{(2m-2)!!}{(2m-1)!!} & \text{if } n = 2m-1, \\ \frac{\pi^2}{8} \frac{(2m-1)!!}{(2m)!!} & \text{if } n = 2m, \end{cases} \quad m = 1, 2, \dots$$
4. $\int_0^\pi \frac{dx}{(a+b \cos x)^{n+1}} = \frac{\pi}{2^n (a+b)^n \sqrt{a^2-b^2}} \sum_{k=0}^n \frac{(2n-2k-1)!! (2k-1)!!}{(n-k)! k!} \left(\frac{a+b}{a-b} \right)^k, \quad a > |b|.$
5. $\int_0^{\pi/2} \sin^{2n} x dx = \frac{\pi}{2} \frac{1 \times 3 \times \dots \times (2n-1)}{2 \times 4 \times \dots \times (2n)}, \quad n = 1, 2, \dots$
6. $\int_0^{\pi/2} \sin^{2n+1} x dx = \frac{2 \times 4 \times \dots \times (2n)}{1 \times 3 \times \dots \times (2n+1)}, \quad n = 1, 2, \dots$
7. $\int_0^\pi x \sin^\mu x dx = \frac{\pi^2}{2^{\mu+1}} \frac{\Gamma(\mu+1)}{[\Gamma(\mu+\frac{1}{2})]^2}, \quad \mu > -1.$

8. $\int_0^{\pi/2} \frac{\sin x \, dx}{\sqrt{1-k^2 \sin^2 x}} = \frac{1}{2k} \ln \frac{1+k}{1-k}.$
9. $\int_0^{\pi/2} \sin^{2n+1} x \cos^{2m+1} x \, dx = \frac{n! m!}{2(n+m+1)!}, \quad n, m = 1, 2, \dots$
10. $\int_0^{\pi/2} \sin^{p-1} x \cos^{q-1} x \, dx = \frac{1}{2} B\left(\frac{1}{2}p, \frac{1}{2}q\right).$
11. $\int_0^{2\pi} (a \sin x + b \cos x)^{2n} \, dx = 2\pi \frac{(2n-1)!!}{(2n)!!} (a^2 + b^2)^n, \quad n = 1, 2, \dots$
12. $\int_0^\pi \frac{\sin x \, dx}{\sqrt{a^2 + 1 - 2a \cos x}} = \begin{cases} 2 & \text{if } 0 \leq a \leq 1, \\ 2/a & \text{if } 1 < a. \end{cases}$
13. $\int_0^{\pi/2} (\tan x)^{\pm\lambda} \, dx = \frac{\pi}{2 \cos(\frac{1}{2}\pi\lambda)}, \quad |\lambda| < 1.$
14. $\int_0^a \frac{\cos(xt) \, dt}{\sqrt{a^2 - t^2}} = \frac{\pi}{2} J_0(ax), \quad J_0(z)$ is the Bessel function (see Supplement 11.6).
15. $\int_0^a \frac{t \sin(xt) \, dt}{\sqrt{a^2 - t^2}} = \frac{\pi}{2} a J_1(ax), \quad J_1(z)$ is the Bessel function.

4.5-2. Integrals Over an Infinite Interval.

1. $\int_0^\infty \frac{\cos ax}{\sqrt{x}} \, dx = \sqrt{\frac{\pi}{2a}}, \quad a > 0.$
2. $\int_0^\infty \frac{\cos ax - \cos bx}{x} \, dx = \ln \left| \frac{b}{a} \right|, \quad ab \neq 0.$
3. $\int_0^\infty \frac{\cos ax - \cos bx}{x^2} \, dx = \frac{1}{2}\pi(b-a), \quad a, b \geq 0.$
4. $\int_0^\infty x^{\mu-1} \cos ax \, dx = a^{-\mu} \Gamma(\mu) \cos\left(\frac{1}{2}\pi\mu\right), \quad a > 0, \quad 0 < \mu < 1.$
5. $\int_0^\infty \frac{\cos ax}{b^2 + x^2} \, dx = \frac{\pi}{2b} e^{-ab}, \quad a, b > 0.$
6. $\int_0^\infty \frac{\cos ax}{b^4 + x^4} \, dx = \frac{\pi\sqrt{2}}{4b^3} \exp\left(-\frac{ab}{\sqrt{2}}\right) \left[\cos\left(\frac{ab}{\sqrt{2}}\right) + \sin\left(\frac{ab}{\sqrt{2}}\right) \right], \quad a, b > 0.$
7. $\int_0^\infty \frac{\cos ax}{(b^2 + x^2)^2} \, dx = \frac{\pi}{4b^3} (1+ab)e^{-ab}, \quad a, b > 0.$
8. $\int_0^\infty \frac{\cos ax \, dx}{(b^2 + x^2)(c^2 + x^2)} = \frac{\pi (be^{-ac} - ce^{-ab})}{2bc(b^2 - c^2)}, \quad a, b, c > 0.$
9. $\int_0^\infty \cos(ax^2) \, dx = \frac{1}{2} \sqrt{\frac{\pi}{2a}}, \quad a > 0.$
10. $\int_0^\infty \cos(ax^p) \, dx = \frac{\Gamma(1/p)}{pa^{1/p}} \cos \frac{\pi}{2p}, \quad a > 0, \quad p > 1.$
11. $\int_0^\infty \frac{\sin ax}{x} \, dx = \frac{\pi}{2} \operatorname{sign} a.$

12. $\int_0^{\infty} \frac{\sin^2 ax}{x^2} dx = \frac{\pi}{2}|a|.$
13. $\int_0^{\infty} \frac{\sin ax}{\sqrt{x}} dx = \sqrt{\frac{\pi}{2a}}, \quad a > 0.$
14. $\int_0^{\infty} x^{\mu-1} \sin ax dx = a^{-\mu} \Gamma(\mu) \sin\left(\frac{1}{2}\pi\mu\right), \quad a > 0, \quad 0 < \mu < 1.$
15. $\int_0^{\infty} \sin(ax^2) dx = \frac{1}{2} \sqrt{\frac{\pi}{2a}}, \quad a > 0.$
16. $\int_0^{\infty} \sin(ax^p) dx = \frac{\Gamma(1/p)}{pa^{1/p}} \sin \frac{\pi}{2p}, \quad a > 0, \quad p > 1.$
17. $\int_0^{\infty} \frac{\sin x \cos ax}{x} dx = \begin{cases} \frac{\pi}{2} & \text{if } |a| < 1, \\ \frac{\pi}{4} & \text{if } |a| = 1, \\ 0 & \text{if } 1 < |a|. \end{cases}$
18. $\int_0^{\infty} \frac{\tan ax}{x} dx = \frac{\pi}{2} \operatorname{sign} a.$
19. $\int_0^{\infty} e^{-ax} \sin bx dx = \frac{b}{a^2 + b^2}, \quad a > 0.$
20. $\int_0^{\infty} e^{-ax} \cos bx dx = \frac{a}{a^2 + b^2}, \quad a > 0.$
21. $\int_0^{\infty} \exp(-ax^2) \cos bx dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} \exp\left(-\frac{b^2}{4a}\right).$
22. $\int_0^{\infty} \cos(ax^2) \cos bx dx = \sqrt{\frac{\pi}{8a}} \left[\cos\left(\frac{b^2}{4a}\right) + \sin\left(\frac{b^2}{4a}\right) \right], \quad a, b > 0.$
23. $\int_0^{\infty} (\cos ax + \sin ax) \cos(b^2x^2) dx = \frac{1}{b} \sqrt{\frac{\pi}{8}} \exp\left(-\frac{a^2}{2b}\right), \quad a, b > 0.$
24. $\int_0^{\infty} [\cos ax + \sin ax] \sin(b^2x^2) dx = \frac{1}{b} \sqrt{\frac{\pi}{8}} \exp\left(-\frac{a^2}{2b}\right), \quad a, b > 0.$

4.6. Integrals Involving Bessel Functions

4.6-1. Integrals Over an Infinite Interval.

1. $\int_0^{\infty} J_{\nu}(ax) dx = \frac{1}{a}, \quad a > 0, \operatorname{Re} \nu > -1.$
2. $\int_0^{\infty} \cos(xu) J_0(tu) du = \begin{cases} \frac{1}{\sqrt{t^2 - x^2}} & \text{if } x < t, \\ 0 & \text{if } x > t. \end{cases}$
3. $\int_0^{\infty} \sin(xu) J_0(tu) du = \begin{cases} 0 & \text{if } x < t, \\ \frac{1}{\sqrt{x^2 - t^2}} & \text{if } x > t. \end{cases}$
4. $\int_0^{\infty} \cos(xu) J_1(tu) du = \begin{cases} \frac{1}{t} & \text{if } x < t, \\ -\frac{t}{\sqrt{x^2 - t^2}(x + \sqrt{x^2 - t^2})} & \text{if } x > t. \end{cases}$
5. $\int_0^{\infty} \frac{\sin(tu) J_0(au)}{u^2 + b^2} du = \frac{\sinh(bt)}{b} K_0(ab), \quad b > 0, \quad 0 < t < a, \quad K_0(z) \text{ is the modified Bessel function (see Supplement 11.7).}$

6. $\int_0^\infty \frac{u \sin(tu) J_0(au)}{u^2 + b^2} du = \frac{\pi}{2} e^{-bt} I_0(ab)$, $b > 0$, $a < t < \infty$, $I_0(z)$ is the modified Bessel function.
7. $\int_0^\infty \frac{\sin(tu) J_1(au)}{u^2 + b^2} du = \frac{\pi}{2b} e^{-bt} I_1(ab)$, $b > 0$, $a < t < \infty$, $I_1(z)$ is the modified Bessel function.
8. $\int_0^\infty \frac{u \sin(tu) J_1(au)}{u^2 + b^2} du = \sinh(bt) K_1(ab)$, $b > 0$, $0 < t < a$, $K_1(z)$ is the modified Bessel function.
9. $\int_0^\infty \frac{J_1(au)}{\sqrt{u^2 + b^2}} du = \frac{1 - e^{-ab}}{ab}$, $a > 0$, $\operatorname{Re} b > 0$.

4.6-2. Other Integrals.

1. $\int_0^1 u J_0(xu) du = \frac{J_1(x)}{x}$.
2. $\int_0^a \frac{J_1(bx) dx}{\sqrt{a^2 - x^2}} = \frac{1 - \cos(ab)}{ab}$, $a > 0$.
3. $\int_0^t \frac{u J_0(xu) du}{\sqrt{t^2 - u^2}} = \frac{\sin(xt)}{x}$.
4. $\int_t^\infty \frac{J_1(xu) du}{\sqrt{u^2 - t^2}} = \frac{\sin(xt)}{x}$, $x > 0$, $t > 0$.

© References for Supplement 4: H. B. Dwight (1961), I. S. Gradshteyn and I. M. Ryzhik (2000), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1986, 1988), D. Zwillinger (2002), I. N. Bronshtein and K. A. Semendyayev (2004).

Supplement 5

Tables of Laplace Transforms

5.1. General Formulas

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
1	$af_1(x) + bf_2(x)$	$a\tilde{f}_1(p) + b\tilde{f}_2(p)$
2	$f(x/a), a > 0$	$a\tilde{f}(ap)$
3	$\begin{cases} 0 & \text{if } 0 < x < a, \\ f(x-a) & \text{if } a < x, \end{cases}$	$e^{-ap}\tilde{f}(p)$
4	$x^n f(x); n = 1, 2, \dots$	$(-1)^n \frac{d^n}{dp^n} \tilde{f}(p)$
5	$\frac{1}{x} f(x)$	$\int_p^{\infty} \tilde{f}(q) dq$
6	$e^{ax} f(x)$	$\tilde{f}(p-a)$
7	$\sinh(ax)f(x)$	$\frac{1}{2} [\tilde{f}(p-a) - \tilde{f}(p+a)]$
8	$\cosh(ax)f(x)$	$\frac{1}{2} [\tilde{f}(p-a) + \tilde{f}(p+a)]$
9	$\sin(\omega x)f(x)$	$-\frac{i}{2} [\tilde{f}(p-i\omega) - \tilde{f}(p+i\omega)], i^2 = -1$
10	$\cos(\omega x)f(x)$	$\frac{1}{2} [\tilde{f}(p-i\omega) + \tilde{f}(p+i\omega)], i^2 = -1$
11	$f(x^2)$	$\frac{1}{\sqrt{\pi}} \int_0^{\infty} \exp\left(-\frac{p^2}{4t^2}\right) \tilde{f}(t^2) dt$
12	$x^{a-1} f\left(\frac{1}{x}\right), a > -1$	$\int_0^{\infty} (t/p)^{a/2} J_a(2\sqrt{pt}) \tilde{f}(t) dt$
13	$f(a \sinh x), a > 0$	$\int_0^{\infty} J_p(at) \tilde{f}(t) dt$
14	$f(x+a) = f(x)$ (periodic function)	$\frac{1}{1-e^{ap}} \int_0^a f(x)e^{-px} dx$
15	$f(x+a) = -f(x)$ (antiperiodic function)	$\frac{1}{1+e^{-ap}} \int_0^a f(x)e^{-px} dx$
16	$f'_x(x)$	$p\tilde{f}(p) - f(+0)$
17	$f_x^{(n)}(x)$	$p^n \tilde{f}(p) - \sum_{k=1}^n p^{n-k} f_x^{(k-1)}(+0)$

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
18	$x^m f_x^{(n)}(x), m \geq n$	$\left(-\frac{d}{dp}\right)^m [p^n \tilde{f}(p)]$
19	$\frac{d^n}{dx^n} [x^m f(x)], m \geq n$	$(-1)^m p^n \frac{d^m}{dp^m} \tilde{f}(p)$
20	$\int_0^x f(t) dt$	$\frac{\tilde{f}(p)}{p}$
21	$\int_0^x (x-t)f(t) dt$	$\frac{1}{p^2} \tilde{f}(p)$
22	$\int_0^x (x-t)^\nu f(t) dt, \nu > -1$	$\Gamma(\nu+1) p^{-\nu-1} \tilde{f}(p)$
23	$\int_0^x e^{-a(x-t)} f(t) dt$	$\frac{1}{p+a} \tilde{f}(p)$
24	$\int_0^x \sinh[a(x-t)] f(t) dt$	$\frac{a \tilde{f}(p)}{p^2 - a^2}$
25	$\int_0^x \sin[a(x-t)] f(t) dt$	$\frac{a \tilde{f}(p)}{p^2 + a^2}$
26	$\int_0^x f_1(t) f_2(x-t) dt$	$\tilde{f}_1(p) \tilde{f}_2(p)$
27	$\int_0^x \frac{1}{t} f(t) dt$	$\frac{1}{p} \int_p^{\infty} \tilde{f}(q) dq$
28	$\int_x^{\infty} \frac{1}{t} f(t) dt$	$\frac{1}{p} \int_0^p \tilde{f}(q) dq$
29	$\int_0^{\infty} \frac{1}{\sqrt{t}} \sin(2\sqrt{xt}) f(t) dt$	$\frac{\sqrt{\pi}}{p\sqrt{p}} \tilde{f}\left(\frac{1}{p}\right)$
30	$\frac{1}{\sqrt{x}} \int_0^{\infty} \cos(2\sqrt{xt}) f(t) dt$	$\frac{\sqrt{\pi}}{\sqrt{p}} \tilde{f}\left(\frac{1}{p}\right)$
31	$\int_0^{\infty} \frac{1}{\sqrt{\pi x}} \exp\left(-\frac{t^2}{4x}\right) f(t) dt$	$\frac{1}{\sqrt{p}} \tilde{f}(\sqrt{p})$
32	$\int_0^{\infty} \frac{t}{2\sqrt{\pi x^3}} \exp\left(-\frac{t^2}{4x}\right) f(t) dt$	$\tilde{f}(\sqrt{p})$
33	$f(x) - a \int_0^x f(\sqrt{x^2 - t^2}) J_1(at) dt$	$\tilde{f}(\sqrt{p^2 + a^2})$
34	$f(x) + a \int_0^x f(\sqrt{x^2 - t^2}) I_1(at) dt$	$\tilde{f}(\sqrt{p^2 - a^2})$

5.2. Expressions with Power-Law Functions

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
1	1	$\frac{1}{p}$
2	$\begin{cases} 0 & \text{if } 0 < x < a, \\ 1 & \text{if } a < x < b, \\ 0 & \text{if } b < x. \end{cases}$	$\frac{1}{p}(e^{-ap} - e^{-bp})$
3	x	$\frac{1}{p^2}$
4	$\frac{1}{x+a}$	$-e^{ap} \text{Ei}(-ap)$
5	$x^n, \quad n = 1, 2, \dots$	$\frac{n!}{p^{n+1}}$
6	$x^{n-1/2}, \quad n = 1, 2, \dots$	$\frac{1 \cdot 3 \dots (2n-1)\sqrt{\pi}}{2^n p^{n+1/2}}$
7	$\frac{1}{\sqrt{x+a}}$	$\sqrt{\frac{\pi}{p}} e^{ap} \text{erfc}(\sqrt{ap})$
8	$\frac{\sqrt{x}}{x+a}$	$\sqrt{\frac{\pi}{p}} - \pi\sqrt{a} e^{ap} \text{erfc}(\sqrt{ap})$
9	$(x+a)^{-3/2}$	$2a^{-1/2} - 2(\pi p)^{1/2} e^{ap} \text{erfc}(\sqrt{ap})$
10	$x^{1/2}(x+a)^{-1}$	$(\pi/p)^{1/2} - \pi a^{1/2} e^{ap} \text{erfc}(\sqrt{ap})$
11	$x^{-1/2}(x+a)^{-1}$	$\pi a^{-1/2} e^{ap} \text{erfc}(\sqrt{ap})$
12	$x^\nu, \quad \nu > -1$	$\Gamma(\nu+1)p^{-\nu-1}$
13	$(x+a)^\nu, \quad \nu > -1$	$p^{-\nu-1} e^{-ap} \Gamma(\nu+1, ap)$
14	$x^\nu(x+a)^{-1}, \quad \nu > -1$	$k e^{ap} \Gamma(-\nu, ap), \quad k = a^\nu \Gamma(\nu+1)$
15	$(x^2 + 2ax)^{-1/2}(x+a)$	$a e^{ap} K_1(ap)$

5.3. Expressions with Exponential Functions

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
1	e^{-ax}	$(p+a)^{-1}$
2	$x e^{-ax}$	$(p+a)^{-2}$
3	$x^{\nu-1} e^{-ax}, \quad \nu > 0$	$\Gamma(\nu)(p+a)^{-\nu}$
4	$\frac{1}{x}(e^{-ax} - e^{-bx})$	$\ln(p+b) - \ln(p+a)$

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
5	$\frac{1}{x^2}(1 - e^{-ax})^2$	$(p+2a)\ln(p+2a) + p\ln p - 2(p+a)\ln(p+a)$
6	$\exp(-ax^2), \quad a > 0$	$(\pi b)^{1/2} \exp(bp^2) \operatorname{erfc}(p\sqrt{b}), \quad a = \frac{1}{4b}$
7	$x \exp(-ax^2)$	$2b - 2\pi^{1/2} b^{3/2} p \operatorname{erfc}(p\sqrt{b}), \quad a = \frac{1}{4b}$
8	$\exp(-a/x), \quad a \geq 0$	$2\sqrt{a/p} K_1(2\sqrt{ap})$
9	$\sqrt{x} \exp(-a/x), \quad a \geq 0$	$\frac{1}{2} \sqrt{\pi/p^3} (1 + 2\sqrt{ap}) \exp(-2\sqrt{ap})$
10	$\frac{1}{\sqrt{x}} \exp(-a/x), \quad a \geq 0$	$\sqrt{\pi/p} \exp(-2\sqrt{ap})$
11	$\frac{1}{x\sqrt{x}} \exp(-a/x), \quad a > 0$	$\sqrt{\pi/a} \exp(-2\sqrt{ap})$
12	$x^{\nu-1} \exp(-a/x), \quad a > 0$	$2(a/p)^{\nu/2} K_{\nu}(2\sqrt{ap})$
13	$\exp(-2\sqrt{ax})$	$p^{-1} - (\pi a)^{1/2} p^{-3/2} e^{a/p} \operatorname{erfc}(\sqrt{a/p})$
14	$\frac{1}{\sqrt{x}} \exp(-2\sqrt{ax})$	$(\pi/p)^{1/2} e^{a/p} \operatorname{erfc}(\sqrt{a/p})$

5.4. Expressions with Hyperbolic Functions

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
1	$\sinh(ax)$	$\frac{a}{p^2 - a^2}$
2	$\sinh^2(ax)$	$\frac{2a^2}{p^3 - 4a^2p}$
3	$\frac{1}{x} \sinh(ax)$	$\frac{1}{2} \ln \frac{p+a}{p-a}$
4	$x^{\nu-1} \sinh(ax), \quad \nu > -1$	$\frac{1}{2} \Gamma(\nu) [(p-a)^{-\nu} - (p+a)^{-\nu}]$
5	$\sinh(2\sqrt{ax})$	$\frac{\sqrt{\pi a}}{p\sqrt{p}} e^{a/p}$
6	$\sqrt{x} \sinh(2\sqrt{ax})$	$\pi^{1/2} p^{-5/2} (\frac{1}{2}p+a) e^{a/p} \operatorname{erf}(\sqrt{a/p}) - a^{1/2} p^{-2}$
7	$\frac{1}{\sqrt{x}} \sinh(2\sqrt{ax})$	$\pi^{1/2} p^{-1/2} e^{a/p} \operatorname{erf}(\sqrt{a/p})$
8	$\frac{1}{\sqrt{x}} \sinh^2(\sqrt{ax})$	$\frac{1}{2} \pi^{1/2} p^{-1/2} (e^{a/p} - 1)$
9	$\cosh(ax)$	$\frac{p}{p^2 - a^2}$

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
10	$\cosh^2(ax)$	$\frac{p^2 - 2a^2}{p^3 - 4a^2p}$
11	$x^{\nu-1} \cosh(ax), \quad \nu > 0$	$\frac{1}{2}\Gamma(\nu)[(p-a)^{-\nu} + (p+a)^{-\nu}]$
12	$\cosh(2\sqrt{ax})$	$\frac{1}{p} + \frac{\sqrt{\pi a}}{p\sqrt{p}} e^{a/p} \operatorname{erf}(\sqrt{a/p})$
13	$\sqrt{x} \cosh(2\sqrt{ax})$	$\pi^{1/2} p^{-5/2} (\frac{1}{2}p + a) e^{a/p}$
14	$\frac{1}{\sqrt{x}} \cosh(2\sqrt{ax})$	$\pi^{1/2} p^{-1/2} e^{a/p}$
15	$\frac{1}{\sqrt{x}} \cosh^2(\sqrt{ax})$	$\frac{1}{2} \pi^{1/2} p^{-1/2} (e^{a/p} + 1)$

5.5. Expressions with Logarithmic Functions

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
1	$\ln x$	$-\frac{1}{p}(\ln p + \mathcal{C}),$ $\mathcal{C} = 0.5772\dots$ is the Euler constant
2	$\ln(1 + ax)$	$-\frac{1}{p} e^{p/a} \operatorname{Ei}(-p/a)$
3	$\ln(x + a)$	$\frac{1}{p} [\ln a - e^{ap} \operatorname{Ei}(-ap)]$
4	$x^n \ln x, \quad n = 1, 2, \dots$	$\frac{n!}{p^{n+1}} (1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} - \ln p - \mathcal{C}),$ $\mathcal{C} = 0.5772\dots$ is the Euler constant
5	$\frac{1}{\sqrt{x}} \ln x$	$-\sqrt{\pi/p} [\ln(4p) + \mathcal{C}]$
6	$x^{n-1/2} \ln x, \quad n = 1, 2, \dots$	$\frac{k_n}{p^{n+1/2}} [2 + \frac{2}{3} + \frac{2}{5} + \dots + \frac{2}{2n-1} - \ln(4p) - \mathcal{C}],$ $k_n = 1 \cdot 3 \cdot 5 \dots (2n-1) \frac{\sqrt{\pi}}{2^n}, \quad \mathcal{C} = 0.5772\dots$
7	$x^{\nu-1} \ln x, \quad \nu > 0$	$\Gamma(\nu) p^{-\nu} [\psi(\nu) - \ln p], \quad \psi(\nu)$ is the logarithmic derivative of the gamma function
8	$(\ln x)^2$	$\frac{1}{p} [(\ln x + \mathcal{C})^2 + \frac{1}{6}\pi^2], \quad \mathcal{C} = 0.5772\dots$
9	$e^{-ax} \ln x$	$-\frac{\ln(p+a) + \mathcal{C}}{p+a}, \quad \mathcal{C} = 0.5772\dots$

5.6. Expressions with Trigonometric Functions

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^\infty e^{-px} f(x) dx$
1	$\sin(ax)$	$\frac{a}{p^2 + a^2}$
2	$ \sin(ax) , \quad a > 0$	$\frac{a}{p^2 + a^2} \coth\left(\frac{\pi p}{2a}\right)$
3	$\sin^{2n}(ax), \quad n = 1, 2, \dots$	$\frac{a^{2n}(2n)!}{p[p^2 + (2a)^2][p^2 + (4a)^2] \dots [p^2 + (2na)^2]}$
4	$\sin^{2n+1}(ax), \quad n = 1, 2, \dots$	$\frac{a^{2n+1}(2n+1)!}{[p^2 + a^2][p^2 + 3^2a^2] \dots [p^2 + (2n+1)^2a^2]}$
5	$x^n \sin(ax), \quad n = 1, 2, \dots$	$\frac{n! p^{n+1}}{(p^2 + a^2)^{n+1}} \sum_{0 \leq 2k \leq n} (-1)^k C_{n+1}^{2k+1} \left(\frac{a}{p}\right)^{2k+1}$
6	$\frac{1}{x} \sin(ax)$	$\arctan\left(\frac{a}{p}\right)$
7	$\frac{1}{x} \sin^2(ax)$	$\frac{1}{4} \ln(1 + 4a^2 p^{-2})$
8	$\frac{1}{x^2} \sin^2(ax)$	$a \arctan(2a/p) - \frac{1}{4} p \ln(1 + 4a^2 p^{-2})$
9	$\sin(2\sqrt{ax})$	$\frac{\sqrt{\pi a}}{p\sqrt{p}} e^{-a/p}$
10	$\frac{1}{x} \sin(2\sqrt{ax})$	$\pi \operatorname{erf}(\sqrt{a/p})$
11	$\cos(ax)$	$\frac{p}{p^2 + a^2}$
12	$\cos^2(ax)$	$\frac{p^2 + 2a^2}{p(p^2 + 4a^2)}$
13	$x^n \cos(ax), \quad n = 1, 2, \dots$	$\frac{n! p^{n+1}}{(p^2 + a^2)^{n+1}} \sum_{0 \leq 2k \leq n+1} (-1)^k C_{n+1}^{2k} \left(\frac{a}{p}\right)^{2k}$
14	$\frac{1}{x} [1 - \cos(ax)]$	$\frac{1}{2} \ln(1 + a^2 p^{-2})$
15	$\frac{1}{x} [\cos(ax) - \cos(bx)]$	$\frac{1}{2} \ln \frac{p^2 + b^2}{p^2 + a^2}$
16	$\sqrt{x} \cos(2\sqrt{ax})$	$\frac{1}{2} \pi^{1/2} p^{-5/2} (p - 2a) e^{-a/p}$
17	$\frac{1}{\sqrt{x}} \cos(2\sqrt{ax})$	$\sqrt{\pi/p} e^{-a/p}$
18	$\sin(ax) \sin(bx)$	$\frac{2abp}{[p^2 + (a+b)^2][p^2 + (a-b)^2]}$
19	$\cos(ax) \sin(bx)$	$\frac{b(p^2 - a^2 + b^2)}{[p^2 + (a+b)^2][p^2 + (a-b)^2]}$

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
20	$\cos(ax) \cos(bx)$	$\frac{p(p^2 + a^2 + b^2)}{[p^2 + (a + b)^2][p^2 + (a - b)^2]}$
21	$\frac{ax \cos(ax) - \sin(ax)}{x^2}$	$p \arctan \frac{a}{x} - a$
22	$e^{bx} \sin(ax)$	$\frac{a}{(p - b)^2 + a^2}$
23	$e^{bx} \cos(ax)$	$\frac{p - b}{(p - b)^2 + a^2}$
24	$\sin(ax) \sinh(ax)$	$\frac{2a^2 p}{p^4 + 4a^4}$
25	$\sin(ax) \cosh(ax)$	$\frac{a(p^2 + 2a^2)}{p^4 + 4a^4}$
26	$\cos(ax) \sinh(ax)$	$\frac{a(p^2 - 2a^2)}{p^4 + 4a^4}$
27	$\cos(ax) \cosh(ax)$	$\frac{p^3}{p^4 + 4a^4}$

5.7. Expressions with Special Functions

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^{\infty} e^{-px} f(x) dx$
1	$\operatorname{erf}(ax)$	$\frac{1}{p} \exp(b^2 p^2) \operatorname{erfc}(bp), \quad b = \frac{1}{2a}$
2	$\operatorname{erf}(\sqrt{ax})$	$\frac{\sqrt{a}}{p\sqrt{p+a}}$
3	$e^{ax} \operatorname{erf}(\sqrt{ax})$	$\frac{\sqrt{a}}{\sqrt{p}(p-a)}$
4	$\operatorname{erf}\left(\frac{1}{2}\sqrt{a/x}\right)$	$\frac{1}{p} [1 - \exp(-\sqrt{ap})]$
5	$\operatorname{erfc}(\sqrt{ax})$	$\frac{\sqrt{p+a} - \sqrt{a}}{p\sqrt{p+a}}$
6	$e^{ax} \operatorname{erfc}(\sqrt{ax})$	$\frac{1}{p + \sqrt{ap}}$
7	$\operatorname{erfc}\left(\frac{1}{2}\sqrt{a/x}\right)$	$\frac{1}{p} \exp(-\sqrt{ap})$
8	$\operatorname{Ci}(x)$	$\frac{1}{2p} \ln(p^2 + 1)$

No	Original function, $f(x)$	Laplace transform, $\tilde{f}(p) = \int_0^\infty e^{-px} f(x) dx$
9	$\text{Si}(x)$	$\frac{1}{p} \operatorname{arccot} p$
10	$\text{Ei}(-x)$	$-\frac{1}{p} \ln(p+1)$
11	$J_0(ax)$	$\frac{1}{\sqrt{p^2+a^2}}$
12	$J_\nu(ax), \quad \nu > -1$	$\frac{a^\nu}{\sqrt{p^2+a^2} (p+\sqrt{p^2+a^2})^\nu}$
13	$x^n J_n(ax), \quad n = 1, 2, \dots$	$1 \cdot 3 \cdot 5 \dots (2n-1) a^n (p^2+a^2)^{-n-1/2}$
14	$x^\nu J_\nu(ax), \quad \nu > -\frac{1}{2}$	$2^\nu \pi^{-1/2} \Gamma(\nu + \frac{1}{2}) a^\nu (p^2+a^2)^{-\nu-1/2}$
15	$x^{\nu+1} J_\nu(ax), \quad \nu > -1$	$2^{\nu+1} \pi^{-1/2} \Gamma(\nu + \frac{3}{2}) a^\nu p (p^2+a^2)^{-\nu-3/2}$
16	$J_0(2\sqrt{ax})$	$\frac{1}{p} e^{-a/p}$
17	$\sqrt{x} J_1(2\sqrt{ax})$	$\frac{\sqrt{a}}{p^2} e^{-a/p}$
18	$x^{\nu/2} J_\nu(2\sqrt{ax}), \quad \nu > -1$	$a^{\nu/2} p^{-\nu-1} e^{-a/p}$
19	$I_0(ax)$	$\frac{1}{\sqrt{p^2-a^2}}$
20	$I_\nu(ax), \quad \nu > -1$	$\frac{a^\nu}{\sqrt{p^2-a^2} (p+\sqrt{p^2-a^2})^\nu}$
21	$x^\nu I_\nu(ax), \quad \nu > -\frac{1}{2}$	$2^\nu \pi^{-1/2} \Gamma(\nu + \frac{1}{2}) a^\nu (p^2-a^2)^{-\nu-1/2}$
22	$x^{\nu+1} I_\nu(ax), \quad \nu > -1$	$2^{\nu+1} \pi^{-1/2} \Gamma(\nu + \frac{3}{2}) a^\nu p (p^2-a^2)^{-\nu-3/2}$
23	$I_0(2\sqrt{ax})$	$\frac{1}{p} e^{a/p}$
24	$\frac{1}{\sqrt{x}} I_1(2\sqrt{ax})$	$\frac{1}{\sqrt{a}} (e^{a/p} - 1)$
25	$x^{\nu/2} I_\nu(2\sqrt{ax}), \quad \nu > -1$	$a^{\nu/2} p^{-\nu-1} e^{a/p}$
26	$Y_0(ax)$	$-\frac{2}{\pi} \frac{\operatorname{Arsinh}(p/a)}{\sqrt{p^2+a^2}}$
27	$K_0(ax)$	$\frac{\ln(p+\sqrt{p^2-a^2}) - \ln a}{\sqrt{p^2-a^2}}$

⊙ References for Supplement 5: G. Doetsch (1950, 1956, 1958), H. Bateman and A. Erdélyi (1954), V. A. Ditkin and A. P. Prudnikov (1965), F. Oberhettinger and L. Badii (1973), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, Vol. 4).

Supplement 6

Tables of Inverse Laplace Transforms

6.1. General Formulas

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$\tilde{f}(p+a)$	$e^{-ax} f(x)$
2	$\tilde{f}(ap), \quad a > 0$	$\frac{1}{a} f\left(\frac{x}{a}\right)$
3	$\tilde{f}(ap+b), \quad a > 0$	$\frac{1}{a} \exp\left(-\frac{b}{a}x\right) f\left(\frac{x}{a}\right)$
4	$\tilde{f}(p-a) + \tilde{f}(p+a)$	$2f(x) \cosh(ax)$
5	$\tilde{f}(p-a) - \tilde{f}(p+a)$	$2f(x) \sinh(ax)$
6	$e^{-ap} \tilde{f}(p), \quad a \geq 0$	$\begin{cases} 0 & \text{if } 0 \leq x < a, \\ f(x-a) & \text{if } a < x. \end{cases}$
7	$p\tilde{f}(p)$	$\frac{df(x)}{dx}, \quad \text{if } f(+0) = 0$
8	$\frac{1}{p} \tilde{f}(p)$	$\int_0^x f(t) dt$
9	$\frac{1}{p+a} \tilde{f}(p)$	$e^{-ax} \int_0^x e^{at} f(t) dt$
10	$\frac{1}{p^2} \tilde{f}(p)$	$\int_0^x (x-t) f(t) dt$
11	$\frac{\tilde{f}(p)}{p(p+a)}$	$\frac{1}{a} \int_0^x [1 - e^{a(x-t)}] f(t) dt$
12	$\frac{\tilde{f}(p)}{(p+a)^2}$	$\int_0^x (x-t) e^{-a(x-t)} f(t) dt$
13	$\frac{\tilde{f}(p)}{(p+a)(p+b)}$	$\frac{1}{b-a} \int_0^x [e^{-a(x-t)} - e^{-b(x-t)}] f(t) dt$
14	$\frac{\tilde{f}(p)}{(p+a)^2 + b^2}$	$\frac{1}{b} \int_0^x e^{-a(x-t)} \sin[b(x-t)] f(t) dt$
15	$\frac{1}{p^n} \tilde{f}(p), \quad n = 1, 2, \dots$	$\frac{1}{(n-1)!} \int_0^x (x-t)^{n-1} f(t) dt$
16	$\tilde{f}_1(p)\tilde{f}_2(p)$	$\int_0^x f_1(t)f_2(x-t) dt$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
17	$\frac{1}{\sqrt{p}} \tilde{f}\left(\frac{1}{p}\right)$	$\int_0^\infty \frac{\cos(2\sqrt{xt})}{\sqrt{\pi x}} f(t) dt$
18	$\frac{1}{p\sqrt{p}} \tilde{f}\left(\frac{1}{p}\right)$	$\int_0^\infty \frac{\sin(2\sqrt{xt})}{\sqrt{\pi t}} f(t) dt$
19	$\frac{1}{p^{2\nu+1}} \tilde{f}\left(\frac{1}{p}\right)$	$\int_0^\infty (x/t)^\nu J_{2\nu}(2\sqrt{xt}) f(t) dt$
20	$\frac{1}{p} \tilde{f}\left(\frac{1}{p}\right)$	$\int_0^\infty J_0(2\sqrt{xt}) f(t) dt$
21	$\frac{1}{p} \tilde{f}\left(p + \frac{1}{p}\right)$	$\int_0^x J_0(2\sqrt{xt-t^2}) f(t) dt$
22	$\frac{1}{p^{2\nu+1}} \tilde{f}\left(p + \frac{a}{p}\right), \quad -\frac{1}{2} < \nu \leq 0$	$\int_0^x \left(\frac{x-t}{at}\right)^\nu J_{2\nu}(2\sqrt{axt-at^2}) f(t) dt$
23	$\tilde{f}(\sqrt{p})$	$\int_0^\infty \frac{t}{2\sqrt{\pi x^3}} \exp\left(-\frac{t^2}{4x}\right) f(t) dt$
24	$\frac{1}{\sqrt{p}} \tilde{f}(\sqrt{p})$	$\frac{1}{\sqrt{\pi x}} \int_0^\infty \exp\left(-\frac{t^2}{4x}\right) f(t) dt$
25	$\tilde{f}(p + \sqrt{p})$	$\frac{1}{2\sqrt{\pi}} \int_0^x \frac{t}{(x-t)^{3/2}} \exp\left[-\frac{t^2}{4(x-t)}\right] f(t) dt$
26	$\tilde{f}(\sqrt{p^2 + a^2})$	$f(x) - a \int_0^x f(\sqrt{x^2 - t^2}) J_1(at) dt$
27	$\tilde{f}(\sqrt{p^2 - a^2})$	$f(x) + a \int_0^x f(\sqrt{x^2 - t^2}) I_1(at) dt$
28	$\frac{\tilde{f}(\sqrt{p^2 + a^2})}{\sqrt{p^2 + a^2}}$	$\int_0^x J_0(a\sqrt{x^2 - t^2}) f(t) dt$
29	$\frac{\tilde{f}(\sqrt{p^2 - a^2})}{\sqrt{p^2 - a^2}}$	$\int_0^x I_0(a\sqrt{x^2 - t^2}) f(t) dt$
30	$\tilde{f}(\sqrt{(p+a)^2 - b^2})$	$e^{-ax} f(x) + be^{-ax} \int_0^x f(\sqrt{x^2 - t^2}) I_1(bt) dt$
31	$\tilde{f}(\ln p)$	$\int_0^\infty \frac{x^{t-1}}{\Gamma(t)} f(t) dt$
32	$\frac{1}{p} \tilde{f}(\ln p)$	$\int_0^\infty \frac{x^t}{\Gamma(t+1)} f(t) dt$
33	$\tilde{f}(p-ia) + \tilde{f}(p+ia), \quad i^2 = -1$	$2f(x) \cos(ax)$
34	$i[\tilde{f}(p-ia) - \tilde{f}(p+ia)], \quad i^2 = -1$	$2f(x) \sin(ax)$
35	$\frac{d\tilde{f}(p)}{dp}$	$-xf(x)$
36	$\frac{d^n \tilde{f}(p)}{dp^n}$	$(-x)^n f(x)$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
37	$p^n \frac{d^m \tilde{f}(p)}{dp^m}, \quad m \geq n$	$(-1)^m \frac{d^n}{dx^n} [x^m f(x)]$
38	$\int_p^\infty \tilde{f}(q) dq$	$\frac{1}{x} f(x)$
39	$\frac{1}{p} \int_0^p \tilde{f}(q) dq$	$\int_x^\infty \frac{f(t)}{t} dt$
40	$\frac{1}{p} \int_p^\infty \tilde{f}(q) dq$	$\int_0^x \frac{f(t)}{t} dt$

6.2. Expressions with Rational Functions

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$\frac{1}{p}$	1
2	$\frac{1}{p+a}$	e^{-ax}
3	$\frac{1}{p^2}$	x
4	$\frac{1}{p(p+a)}$	$\frac{1}{a} (1 - e^{-ax})$
5	$\frac{1}{(p+a)^2}$	$x e^{-ax}$
6	$\frac{p}{(p+a)^2}$	$(1 - ax) e^{-ax}$
7	$\frac{1}{p^2 - a^2}$	$\frac{1}{a} \sinh(ax)$
8	$\frac{p}{p^2 - a^2}$	$\cosh(ax)$
9	$\frac{1}{(p+a)(p+b)}$	$\frac{1}{a-b} (e^{-bx} - e^{-ax})$
10	$\frac{p}{(p+a)(p+b)}$	$\frac{1}{a-b} (a e^{-ax} - b e^{-bx})$
11	$\frac{1}{p^2 + a^2}$	$\frac{1}{a} \sin(ax)$
12	$\frac{p}{p^2 + a^2}$	$\cos(ax)$
13	$\frac{1}{(p+b)^2 + a^2}$	$\frac{1}{a} e^{-bx} \sin(ax)$
14	$\frac{p}{(p+b)^2 + a^2}$	$e^{-bx} \left[\cos(ax) - \frac{b}{a} \sin(ax) \right]$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
15	$\frac{1}{p^3}$	$\frac{1}{2}x^2$
16	$\frac{1}{p^2(p+a)}$	$\frac{1}{a^2}(e^{-ax} + ax - 1)$
17	$\frac{1}{p(p+a)(p+b)}$	$\frac{1}{ab(a-b)}(a-b + be^{-ax} - ae^{-bx})$
18	$\frac{1}{p(p+a)^2}$	$\frac{1}{a^2}(1 - e^{-ax} - axe^{-ax})$
19	$\frac{1}{(p+a)(p+b)(p+c)}$	$\frac{(c-b)e^{-ax} + (a-c)e^{-bx} + (b-a)e^{-cx}}{(a-b)(b-c)(c-a)}$
20	$\frac{p}{(p+a)(p+b)(p+c)}$	$\frac{a(b-c)e^{-ax} + b(c-a)e^{-bx} + c(a-b)e^{-cx}}{(a-b)(b-c)(c-a)}$
21	$\frac{p^2}{(p+a)(p+b)(p+c)}$	$\frac{a^2(c-b)e^{-ax} + b^2(a-c)e^{-bx} + c^2(b-a)e^{-cx}}{(a-b)(b-c)(c-a)}$
22	$\frac{1}{(p+a)(p+b)^2}$	$\frac{1}{(a-b)^2}[e^{-ax} - e^{-bx} + (a-b)xe^{-bx}]$
23	$\frac{p}{(p+a)(p+b)^2}$	$\frac{1}{(a-b)^2}\{-ae^{-ax} + [a + b(b-a)x]e^{-bx}\}$
24	$\frac{p^2}{(p+a)(p+b)^2}$	$\frac{1}{(a-b)^2}[a^2e^{-ax} + b(b-2a-b^2x+abx)e^{-bx}]$
25	$\frac{1}{(p+a)^3}$	$\frac{1}{2}x^2e^{-ax}$
26	$\frac{p}{(p+a)^3}$	$x(1 - \frac{1}{2}ax)e^{-ax}$
27	$\frac{p^2}{(p+a)^3}$	$(1 - 2ax + \frac{1}{2}a^2x^2)e^{-ax}$
28	$\frac{1}{p(p^2+a^2)}$	$\frac{1}{a^2}[1 - \cos(ax)]$
29	$\frac{1}{p[(p+b)^2+a^2]}$	$\frac{1}{a^2+b^2}\left\{1 - e^{-bx}\left[\cos(ax) + \frac{b}{a}\sin(ax)\right]\right\}$
30	$\frac{1}{(p+a)(p^2+b^2)}$	$\frac{1}{a^2+b^2}\left[e^{-ax} + \frac{a}{b}\sin(bx) - \cos(bx)\right]$
31	$\frac{p}{(p+a)(p^2+b^2)}$	$\frac{1}{a^2+b^2}[-ae^{-ax} + a\cos(bx) + b\sin(bx)]$
32	$\frac{p^2}{(p+a)(p^2+b^2)}$	$\frac{1}{a^2+b^2}[a^2e^{-ax} - ab\sin(bx) + b^2\cos(bx)]$
33	$\frac{1}{p^3+a^3}$	$\frac{1}{3a^2}e^{-ax} - \frac{1}{3a^2}e^{ax/2}[\cos(kx) - \sqrt{3}\sin(kx)],$ $k = \frac{1}{2}a\sqrt{3}$
34	$\frac{p}{p^3+a^3}$	$-\frac{1}{3a}e^{-ax} + \frac{1}{3a}e^{ax/2}[\cos(kx) + \sqrt{3}\sin(kx)],$ $k = \frac{1}{2}a\sqrt{3}$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
35	$\frac{p^2}{p^3 + a^3}$	$\frac{1}{3}e^{-ax} + \frac{2}{3}e^{ax/2} \cos(kx), \quad k = \frac{1}{2}a\sqrt{3}$
36	$\frac{1}{(p+a)[(p+b)^2 + c^2]}$	$\frac{e^{-ax} - e^{-bx} \cos(cx) + ke^{-bx} \sin(cx)}{(a-b)^2 + c^2}, \quad k = \frac{a-b}{c}$
37	$\frac{p}{(p+a)[(p+b)^2 + c^2]}$	$\frac{-ae^{-ax} + ae^{-bx} \cos(cx) + ke^{-bx} \sin(cx)}{(a-b)^2 + c^2},$ $k = \frac{b^2 + c^2 - ab}{c}$
38	$\frac{p^2}{(p+a)[(p+b)^2 + c^2]}$	$\frac{a^2e^{-ax} + (b^2 + c^2 - 2ab)e^{-bx} \cos(cx) + ke^{-bx} \sin(cx)}{(a-b)^2 + c^2},$ $k = -ac - bc + \frac{ab^2 - b^3}{c}$
39	$\frac{1}{p^4}$	$\frac{1}{6}x^3$
40	$\frac{1}{p^3(p+a)}$	$\frac{1}{a^3} - \frac{1}{a^2}x + \frac{1}{2a}x^2 - \frac{1}{a^3}e^{-ax}$
41	$\frac{1}{p^2(p+a)^2}$	$\frac{1}{a^2}x(1 + e^{-ax}) + \frac{2}{a^3}(e^{-ax} - 1)$
42	$\frac{1}{p^2(p+a)(p+b)}$	$-\frac{a+b}{a^2b^2} + \frac{1}{ab}x + \frac{1}{a^2(b-a)}e^{-ax} + \frac{1}{b^2(a-b)}e^{-bx}$
43	$\frac{1}{(p+a)^2(p+b)^2}$	$\frac{1}{(a-b)^2} \left[e^{-ax} \left(x + \frac{2}{a-b} \right) + e^{-bx} \left(x - \frac{2}{a-b} \right) \right]$
44	$\frac{1}{(p+a)^4}$	$\frac{1}{6}x^3 e^{-ax}$
45	$\frac{p}{(p+a)^4}$	$\frac{1}{2}x^2 e^{-ax} - \frac{1}{6}ax^3 e^{-ax}$
46	$\frac{1}{p^2(p^2 + a^2)}$	$\frac{1}{a^3} [ax - \sin(ax)]$
47	$\frac{1}{p^4 - a^4}$	$\frac{1}{2a^3} [\sinh(ax) - \sin(ax)]$
48	$\frac{p}{p^4 - a^4}$	$\frac{1}{2a^2} [\cosh(ax) - \cos(ax)]$
49	$\frac{p^2}{p^4 - a^4}$	$\frac{1}{2a} [\sinh(ax) + \sin(ax)]$
50	$\frac{p^3}{p^4 - a^4}$	$\frac{1}{2} [\cosh(ax) + \cos(ax)]$
51	$\frac{1}{p^4 + a^4}$	$\frac{1}{a^3\sqrt{2}} (\cosh \xi \sin \xi - \sinh \xi \cos \xi), \quad \xi = \frac{ax}{\sqrt{2}}$
52	$\frac{p}{p^4 + a^4}$	$\frac{1}{a^2} \sin\left(\frac{ax}{\sqrt{2}}\right) \sinh\left(\frac{ax}{\sqrt{2}}\right)$
53	$\frac{p^2}{p^4 + a^4}$	$\frac{1}{a\sqrt{2}} (\cos \xi \sinh \xi + \sin \xi \cosh \xi), \quad \xi = \frac{ax}{\sqrt{2}}$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
54	$\frac{1}{(p^2 + a^2)^2}$	$\frac{1}{2a^3} [\sin(ax) - ax \cos(ax)]$
55	$\frac{p}{(p^2 + a^2)^2}$	$\frac{1}{2a} x \sin(ax)$
56	$\frac{p^2}{(p^2 + a^2)^2}$	$\frac{1}{2a} [\sin(ax) + ax \cos(ax)]$
57	$\frac{p^3}{(p^2 + a^2)^2}$	$\cos(ax) - \frac{1}{2} ax \sin(ax)$
58	$\frac{1}{[(p+b)^2 + a^2]^2}$	$\frac{1}{2a^3} e^{-bx} [\sin(ax) - ax \cos(ax)]$
59	$\frac{1}{(p^2 - a^2)(p^2 - b^2)}$	$\frac{1}{a^2 - b^2} \left[\frac{1}{a} \sinh(ax) - \frac{1}{b} \sinh(bx) \right]$
60	$\frac{p}{(p^2 - a^2)(p^2 - b^2)}$	$\frac{\cosh(ax) - \cosh(bx)}{a^2 - b^2}$
61	$\frac{p^2}{(p^2 - a^2)(p^2 - b^2)}$	$\frac{a \sinh(ax) - b \sinh(bx)}{a^2 - b^2}$
62	$\frac{p^3}{(p^2 - a^2)(p^2 - b^2)}$	$\frac{a^2 \cosh(ax) - b^2 \cosh(bx)}{a^2 - b^2}$
63	$\frac{1}{(p^2 + a^2)(p^2 + b^2)}$	$\frac{1}{b^2 - a^2} \left[\frac{1}{a} \sin(ax) - \frac{1}{b} \sin(bx) \right]$
64	$\frac{p}{(p^2 + a^2)(p^2 + b^2)}$	$\frac{\cos(ax) - \cos(bx)}{b^2 - a^2}$
65	$\frac{p^2}{(p^2 + a^2)(p^2 + b^2)}$	$\frac{-a \sin(ax) + b \sin(bx)}{b^2 - a^2}$
66	$\frac{p^3}{(p^2 + a^2)(p^2 + b^2)}$	$\frac{-a^2 \cos(ax) + b^2 \cos(bx)}{b^2 - a^2}$
67	$\frac{1}{p^n}, \quad n = 1, 2, \dots$	$\frac{1}{(n-1)!} x^{n-1}$
68	$\frac{1}{(p+a)^n}, \quad n = 1, 2, \dots$	$\frac{1}{(n-1)!} x^{n-1} e^{-ax}$
69	$\frac{1}{p(p+a)^n}, \quad n = 1, 2, \dots$	$a^{-n} [1 - e^{-ax} e_n(ax)], \quad e_n(z) = 1 + \frac{z}{1!} + \dots + \frac{z^n}{n!}$
70	$\frac{1}{p^{2n} + a^{2n}}, \quad n = 1, 2, \dots$	$-\frac{1}{na^{2n}} \sum_{k=1}^n \exp(a_k x) [a_k \cos(b_k x) - b_k \sin(b_k x)],$ $a_k = a \cos \varphi_k, \quad b_k = a \sin \varphi_k, \quad \varphi_k = \frac{\pi(2k-1)}{2n}$
71	$\frac{1}{p^{2n} - a^{2n}}, \quad n = 1, 2, \dots$	$\frac{1}{na^{2n-1}} \sinh(ax) + \frac{1}{na^{2n}} \sum_{k=2}^n \exp(a_k x)$ $\times [a_k \cos(b_k x) - b_k \sin(b_k x)],$ $a_k = a \cos \varphi_k, \quad b_k = a \sin \varphi_k, \quad \varphi_k = \frac{\pi(k-1)}{n}$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
72	$\frac{1}{p^{2n+1} + a^{2n+1}}, \quad n = 0, 1, \dots$	$\frac{e^{-ax}}{(2n+1)a^{2n}} - \frac{2}{(2n+1)a^{2n+1}} \sum_{k=1}^n \exp(a_k x)$ $\times [a_k \cos(b_k x) - b_k \sin(b_k x)],$ $a_k = a \cos \varphi_k, \quad b_k = a \sin \varphi_k, \quad \varphi_k = \frac{\pi(2k-1)}{2n+1}$
73	$\frac{1}{p^{2n+1} - a^{2n+1}}, \quad n = 0, 1, \dots$	$\frac{e^{ax}}{(2n+1)a^{2n}} + \frac{2}{(2n+1)a^{2n+1}} \sum_{k=1}^n \exp(a_k x)$ $\times [a_k \cos(b_k x) - b_k \sin(b_k x)],$ $a_k = a \cos \varphi_k, \quad b_k = a \sin \varphi_k, \quad \varphi_k = \frac{2\pi k}{2n+1}$
74	$\frac{Q(p)}{P(p)},$ $P(p) = (p - a_1) \dots (p - a_n);$ $Q(p)$ is a polynomial of degree $\leq n - 1;$ $a_i \neq a_j$ if $i \neq j$	$\sum_{k=1}^n \frac{Q(a_k)}{P'(a_k)} \exp(a_k x),$ (the prime stand for the differentiation)
75	$\frac{Q(p)}{P(p)},$ $P(p) = (p - a_1)^{m_1} \dots (p - a_n)^{m_n};$ $Q(p)$ is a polynomial of degree $< m_1 + m_2 + \dots + m_n - 1;$ $a_i \neq a_j$ if $i \neq j$	$\sum_{k=1}^n \sum_{l=1}^{m_k} \frac{\Phi_{kl}(a_k)}{(m_k - l)! (l - 1)!} x^{m_k - l} \exp(a_k x),$ $\Phi_{kl}(p) = \frac{d^{l-1}}{dp^{l-1}} \left[\frac{Q(p)}{P_k(p)} \right], \quad P_k(p) = \frac{P(p)}{(p - a_k)^{m_k}}$
76	$\frac{Q(p) + pR(p)}{P(p)},$ $P(p) = (p^2 + a_1^2) \dots (p^2 + a_n^2);$ $Q(p)$ and $R(p)$ are polynomials of degree $\leq 2n - 2;$ $a_l \neq a_j, \quad l \neq j$	$\sum_{k=1}^n \frac{Q(ia_k) \sin(a_k x) + a_k R(ia_k) \cos(a_k x)}{a_k P_k(ia_k)},$ $P_m(p) = \frac{P(p)}{p^2 + a_m^2}, \quad i^2 = -1$

6.3. Expressions with Square Roots

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$\frac{1}{\sqrt{p}}$	$\frac{1}{\sqrt{\pi x}}$
2	$\sqrt{p-a} - \sqrt{p-b}$	$\frac{e^{bx} - e^{ax}}{2\sqrt{\pi x^3}}$
3	$\frac{1}{\sqrt{p+a}}$	$\frac{1}{\sqrt{\pi x}} e^{-ax}$
4	$\sqrt{\frac{p+a}{p}} - 1$	$\frac{1}{2} a e^{-ax/2} [I_1(\frac{1}{2} ax) + I_0(\frac{1}{2} ax)]$
5	$\frac{\sqrt{p+a}}{p+b}$	$\frac{e^{-ax}}{\sqrt{\pi x}} + (a-b)^{1/2} e^{-bx} \operatorname{erf}[(a-b)^{1/2} x^{1/2}]$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
6	$\frac{1}{p\sqrt{p}}$	$2\sqrt{\frac{x}{\pi}}$
7	$\frac{1}{(p+a)\sqrt{p+b}}$	$(b-a)^{-1/2} e^{-ax} \operatorname{erf}[(b-a)^{1/2} x^{1/2}]$
8	$\frac{1}{\sqrt{p}(p-a)}$	$\frac{1}{\sqrt{a}} e^{ax} \operatorname{erf}(\sqrt{ax})$
9	$\frac{1}{p^{3/2}(p-a)}$	$a^{-3/2} e^{ax} \operatorname{erf}(\sqrt{ax}) - 2a^{-1} \pi^{-1/2} x^{1/2}$
10	$\frac{1}{\sqrt{p}+a}$	$\pi^{-1/2} x^{-1/2} - ae^{a^2 x} \operatorname{erfc}(a\sqrt{x})$
11	$\frac{a}{p(\sqrt{p}+a)}$	$1 - e^{a^2 x} \operatorname{erfc}(a\sqrt{x})$
12	$\frac{1}{p+a\sqrt{p}}$	$e^{a^2 x} \operatorname{erfc}(a\sqrt{x})$
13	$\frac{1}{(\sqrt{p}+\sqrt{a})^2}$	$1 - \frac{2}{\sqrt{\pi}} (ax)^{1/2} + (1-2ax)e^{ax} [\operatorname{erf}(\sqrt{ax}) - 1]$
14	$\frac{1}{p(\sqrt{p}+\sqrt{a})^2}$	$\frac{1}{a} + \left(2x - \frac{1}{a}\right) e^{ax} \operatorname{erfc}(\sqrt{ax}) - \frac{2}{\sqrt{\pi a}} \sqrt{x}$
15	$\frac{1}{\sqrt{p}(\sqrt{p}+a)^2}$	$2\pi^{-1/2} x^{1/2} - 2axe^{a^2 x} \operatorname{erfc}(a\sqrt{x})$
16	$\frac{1}{(\sqrt{p}+a)^3}$	$\frac{2}{\sqrt{\pi}} (a^2 x + 1)\sqrt{x} - ax(2a^2 x + 3)e^{a^2 x} \operatorname{erfc}(a\sqrt{x})$
17	$p^{-n-1/2}, \quad n = 1, 2, \dots$	$\frac{2^n}{1 \cdot 3 \dots (2n-1)\sqrt{\pi}} x^{n-1/2}$
18	$(p+a)^{-n-1/2}$	$\frac{2^n}{1 \cdot 3 \dots (2n-1)\sqrt{\pi}} x^{n-1/2} e^{-ax}$
19	$\frac{1}{\sqrt{p^2+a^2}}$	$J_0(ax)$
20	$\frac{1}{\sqrt{p^2-a^2}}$	$I_0(ax)$
21	$\frac{1}{\sqrt{p^2+ap+b}}$	$\exp(-\frac{1}{2}ax) J_0[(b - \frac{1}{4}a^2)^{1/2} x]$
22	$(\sqrt{p^2+a^2}-p)^{1/2}$	$\frac{1}{\sqrt{2\pi x^3}} \sin(ax)$
23	$\frac{1}{\sqrt{p^2+a^2}} (\sqrt{p^2+a^2}+p)^{1/2}$	$\frac{\sqrt{2}}{\sqrt{\pi x}} \cos(ax)$
24	$\frac{1}{\sqrt{p^2-a^2}} (\sqrt{p^2-a^2}+p)^{1/2}$	$\frac{\sqrt{2}}{\sqrt{\pi x}} \cosh(ax)$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
25	$(\sqrt{p^2 + a^2} + p)^{-n}$	$na^{-n} x^{-1} J_n(ax)$
26	$(\sqrt{p^2 - a^2} + p)^{-n}$	$na^{-n} x^{-1} I_n(ax)$
27	$(p^2 + a^2)^{-n-1/2}$	$\frac{(x/a)^n J_n(ax)}{1 \cdot 3 \cdot 5 \dots (2n-1)}$
28	$(p^2 - a^2)^{-n-1/2}$	$\frac{(x/a)^n I_n(ax)}{1 \cdot 3 \cdot 5 \dots (2n-1)}$

6.4. Expressions with Arbitrary Powers

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$(p+a)^{-\nu}, \nu > 0$	$\frac{1}{\Gamma(\nu)} x^{\nu-1} e^{-ax}$
2	$[(p+a)^{1/2} + (p+b)^{1/2}]^{-2\nu}, \nu > 0$	$\frac{\nu}{(a-b)^\nu} x^{-1} \exp[-\frac{1}{2}(a+b)x] I_\nu[\frac{1}{2}(a-b)x]$
3	$[(p+a)(p+b)]^{-\nu}, \nu > 0$	$\frac{\sqrt{\pi}}{\Gamma(\nu)} \left(\frac{x}{a-b}\right)^{\nu-1/2} \exp\left(-\frac{a+b}{2}x\right) I_{\nu-1/2}\left(\frac{a-b}{2}x\right)$
4	$(p^2 + a^2)^{-\nu-1/2}, \nu > -\frac{1}{2}$	$\frac{\sqrt{\pi}}{(2a)^\nu \Gamma(\nu + \frac{1}{2})} x^\nu J_\nu(ax)$
5	$(p^2 - a^2)^{-\nu-1/2}, \nu > -\frac{1}{2}$	$\frac{\sqrt{\pi}}{(2a)^\nu \Gamma(\nu + \frac{1}{2})} x^\nu I_\nu(ax)$
6	$p(p^2 + a^2)^{-\nu-1/2}, \nu > 0$	$\frac{a\sqrt{\pi}}{(2a)^\nu \Gamma(\nu + \frac{1}{2})} x^\nu J_{\nu-1}(ax)$
7	$p(p^2 - a^2)^{-\nu-1/2}, \nu > 0$	$\frac{a\sqrt{\pi}}{(2a)^\nu \Gamma(\nu + \frac{1}{2})} x^\nu I_{\nu-1}(ax)$
8	$[(p^2 + a^2)^{1/2} + p]^{-\nu} = a^{-2\nu} [(p^2 + a^2)^{1/2} - p]^\nu, \nu > 0$	$\nu a^{-\nu} x^{-1} J_\nu(ax)$
9	$[(p^2 - a^2)^{1/2} + p]^{-\nu} = a^{-2\nu} [p - (p^2 - a^2)^{1/2}]^\nu, \nu > 0$	$\nu a^{-\nu} x^{-1} I_\nu(ax)$
10	$p[(p^2 + a^2)^{1/2} + p]^{-\nu}, \nu > 1$	$\nu a^{1-\nu} x^{-1} J_{\nu-1}(ax) - \nu(\nu+1)a^{-\nu} x^{-2} J_\nu(ax)$
11	$p[(p^2 - a^2)^{1/2} + p]^{-\nu}, \nu > 1$	$\nu a^{1-\nu} x^{-1} I_{\nu-1}(ax) - \nu(\nu+1)a^{-\nu} x^{-2} I_\nu(ax)$
12	$\frac{(\sqrt{p^2 + a^2} + p)^{-\nu}}{\sqrt{p^2 + a^2}}, \nu > -1$	$a^{-\nu} J_\nu(ax)$
13	$\frac{(\sqrt{p^2 - a^2} + p)^{-\nu}}{\sqrt{p^2 - a^2}}, \nu > -1$	$a^{-\nu} I_\nu(ax)$

6.5. Expressions with Exponential Functions

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$p^{-1}e^{-ap}, \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ 1 & \text{if } a < x. \end{cases}$
2	$p^{-1}(1 - e^{-ap}), \quad a > 0$	$\begin{cases} 1 & \text{if } 0 < x < a, \\ 0 & \text{if } a < x. \end{cases}$
3	$p^{-1}(e^{-ap} - e^{-bp}), \quad 0 \leq a < b$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ 1 & \text{if } a < x < b, \\ 0 & \text{if } b < x. \end{cases}$
4	$p^{-2}(e^{-ap} - e^{-bp}), \quad 0 \leq a < b$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ x - a & \text{if } a < x < b, \\ b - a & \text{if } b < x. \end{cases}$
5	$(p + b)^{-1}e^{-ap}, \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ e^{-b(x-a)} & \text{if } a < x. \end{cases}$
6	$p^{-\nu}e^{-ap}, \quad \nu > 0$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ \frac{(x-a)^{\nu-1}}{\Gamma(\nu)} & \text{if } a < x. \end{cases}$
7	$p^{-1}(e^{ap} - 1)^{-1}, \quad a > 0$	$f(x) = n \quad \text{if } na < x < (n+1)a; \quad n = 0, 1, 2, \dots$
8	$e^{a/p} - 1$	$\sqrt{\frac{a}{x}} I_1(2\sqrt{ax})$
9	$p^{-1/2}e^{a/p}$	$\frac{1}{\sqrt{\pi x}} \cosh(2\sqrt{ax})$
10	$p^{-3/2}e^{a/p}$	$\frac{1}{\sqrt{\pi a}} \sinh(2\sqrt{ax})$
11	$p^{-5/2}e^{a/p}$	$\sqrt{\frac{x}{\pi a}} \cosh(2\sqrt{ax}) - \frac{1}{2\sqrt{\pi a^3}} \sinh(2\sqrt{ax})$
12	$p^{-\nu-1}e^{a/p}, \quad \nu > -1$	$(x/a)^{\nu/2} I_\nu(2\sqrt{ax})$
13	$1 - e^{-a/p}$	$\sqrt{\frac{a}{x}} J_1(2\sqrt{ax})$
14	$p^{-1/2}e^{-a/p}$	$\frac{1}{\sqrt{\pi x}} \cos(2\sqrt{ax})$
15	$p^{-3/2}e^{-a/p}$	$\frac{1}{\sqrt{\pi a}} \sin(2\sqrt{ax})$
16	$p^{-5/2}e^{-a/p}$	$\frac{1}{2\sqrt{\pi a^3}} \sin(2\sqrt{ax}) - \sqrt{\frac{x}{\pi a}} \cos(2\sqrt{ax})$
17	$p^{-\nu-1}e^{-a/p}, \quad \nu > -1$	$(x/a)^{\nu/2} J_\nu(2\sqrt{ax})$
18	$\exp(-\sqrt{ap}), \quad a > 0$	$\frac{\sqrt{a}}{2\sqrt{\pi}} x^{-3/2} \exp\left(-\frac{a}{4x}\right)$
19	$p \exp(-\sqrt{ap}), \quad a > 0$	$\frac{\sqrt{a}}{8\sqrt{\pi}} (a - 6x)x^{-7/2} \exp\left(-\frac{a}{4x}\right)$
20	$\frac{1}{p} \exp(-\sqrt{ap}), \quad a \geq 0$	$\operatorname{erfc}\left(\frac{\sqrt{a}}{2\sqrt{x}}\right)$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
21	$\sqrt{p} \exp(-\sqrt{ap}), \quad a > 0$	$\frac{1}{4\sqrt{\pi}} (a-2x)x^{-5/2} \exp\left(-\frac{a}{4x}\right)$
22	$\frac{1}{\sqrt{p}} \exp(-\sqrt{ap}), \quad a \geq 0$	$\frac{1}{\sqrt{\pi x}} \exp\left(-\frac{a}{4x}\right)$
23	$\frac{1}{p\sqrt{p}} \exp(-\sqrt{ap}), \quad a \geq 0$	$\frac{2\sqrt{x}}{\sqrt{\pi}} \exp\left(-\frac{a}{4x}\right) - \sqrt{a} \operatorname{erfc}\left(\frac{\sqrt{a}}{2\sqrt{x}}\right)$
24	$\frac{\exp(-k\sqrt{p^2+a^2})}{\sqrt{p^2+a^2}}, \quad k > 0$	$\begin{cases} 0 & \text{if } 0 < x < k, \\ J_0(a\sqrt{x^2-k^2}) & \text{if } k < x. \end{cases}$
25	$\frac{\exp(-k\sqrt{p^2-a^2})}{\sqrt{p^2-a^2}}, \quad k > 0$	$\begin{cases} 0 & \text{if } 0 < x < k, \\ I_0(a\sqrt{x^2-k^2}) & \text{if } k < x. \end{cases}$

6.6. Expressions with Hyperbolic Functions

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$\frac{1}{p \sinh(ap)}, \quad a > 0$	$f(x) = 2n$ if $a(2n-1) < x < a(2n+1)$; $n = 0, 1, 2, \dots$ ($x > 0$)
2	$\frac{1}{p^2 \sinh(ap)}, \quad a > 0$	$f(x) = 2n(x-an)$ if $a(2n-1) < x < a(2n+1)$; $n = 0, 1, 2, \dots$ ($x > 0$)
3	$\frac{\sinh(a/p)}{\sqrt{p}}$	$\frac{1}{2\sqrt{\pi x}} [\cosh(2\sqrt{ax}) - \cos(2\sqrt{ax})]$
4	$\frac{\sinh(a/p)}{p\sqrt{p}}$	$\frac{1}{2\sqrt{\pi a}} [\sinh(2\sqrt{ax}) - \sin(2\sqrt{ax})]$
5	$p^{-\nu-1} \sinh(a/p), \quad \nu > -2$	$\frac{1}{2}(x/a)^{\nu/2} [I_\nu(2\sqrt{ax}) - J_\nu(2\sqrt{ax})]$
6	$\frac{1}{p \cosh(ap)}, \quad a > 0$	$f(x) = \begin{cases} 0 & \text{if } a(4n-1) < x < a(4n+1), \\ 2 & \text{if } a(4n+1) < x < a(4n+3), \end{cases}$ $n = 0, 1, 2, \dots$ ($x > 0$)
7	$\frac{1}{p^2 \cosh(ap)}, \quad a > 0$	$x - (-1)^n(x-2an)$ if $2n-1 < x/a < 2n+1$; $n = 0, 1, 2, \dots$ ($x > 0$)
8	$\frac{\cosh(a/p)}{\sqrt{p}}$	$\frac{1}{2\sqrt{\pi x}} [\cosh(2\sqrt{ax}) + \cos(2\sqrt{ax})]$
9	$\frac{\cosh(a/p)}{p\sqrt{p}}$	$\frac{1}{2\sqrt{\pi a}} [\sinh(2\sqrt{ax}) + \sin(2\sqrt{ax})]$
10	$p^{-\nu-1} \cosh(a/p), \quad \nu > -1$	$\frac{1}{2}(x/a)^{\nu/2} [I_\nu(2\sqrt{ax}) + J_\nu(2\sqrt{ax})]$
11	$\frac{1}{p} \tanh(ap), \quad a > 0$	$f(x) = (-1)^{n-1}$ if $2a(n-1) < x < 2an$; $n = 1, 2, \dots$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
12	$\frac{1}{p} \coth(ap), \quad a > 0$	$f(x) = (2n-1)$ if $2a(n-1) < x < 2an$; $n = 1, 2, \dots$
13	$\operatorname{Arcoth}(p/a)$	$\frac{1}{x} \sinh(ax)$

6.7. Expressions with Logarithmic Functions

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$\frac{1}{p} \ln p$	$-\ln x - C$, $C = 0.5772\dots$ is the Euler constant
2	$p^{-n-1} \ln p$	$(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} - \ln x - C) \frac{x^n}{n!}$, $C = 0.5772\dots$ is the Euler constant
3	$p^{-n-1/2} \ln p$	$k_n [2 + \frac{2}{3} + \frac{2}{5} + \dots + \frac{2}{2n-1} - \ln(4x) - C] x^{n-1/2}$, $k_n = \frac{2^n}{1 \cdot 3 \cdot 5 \dots (2n-1)\sqrt{\pi}}$, $C = 0.5772\dots$
4	$p^{-\nu} \ln p, \quad \nu > 0$	$\frac{1}{\Gamma(\nu)} x^{\nu-1} [\psi(\nu) - \ln x]$, $\psi(\nu)$ is the logarithmic derivative of the gamma function
5	$\frac{1}{p} (\ln p)^2$	$(\ln x + C)^2 - \frac{1}{6}\pi^2$, $C = 0.5772\dots$
6	$\frac{1}{p^2} (\ln p)^2$	$x [(\ln x + C - 1)^2 + 1 - \frac{1}{6}\pi^2]$
7	$\frac{\ln(p+b)}{p+a}$	$e^{-ax} \{ \ln(b-a) - \operatorname{Ei}[(a-b)x] \}$
8	$\frac{\ln p}{p^2 + a^2}$	$\frac{1}{a} \cos(ax) \operatorname{Si}(ax) + \frac{1}{a} \sin(ax) [\ln a - \operatorname{Ci}(ax)]$
9	$\frac{p \ln p}{p^2 + a^2}$	$\cos(ax) [\ln a - \operatorname{Ci}(ax)] - \sin(ax) \operatorname{Si}(ax)$
10	$\ln \frac{p+b}{p+a}$	$\frac{1}{x} (e^{-ax} - e^{-bx})$
11	$\ln \frac{p^2 + b^2}{p^2 + a^2}$	$\frac{2}{x} [\cos(ax) - \cos(bx)]$
12	$p \ln \frac{p^2 + b^2}{p^2 + a^2}$	$\frac{2}{x} [\cos(bx) + bx \sin(bx) - \cos(ax) - ax \sin(ax)]$
13	$\ln \frac{(p+a)^2 + k^2}{(p+b)^2 + k^2}$	$\frac{2}{x} \cos(kx) (e^{-bx} - e^{-ax})$
14	$p \ln \left(\frac{1}{p} \sqrt{p^2 + a^2} \right)$	$\frac{1}{x^2} [\cos(ax) - 1] + \frac{a}{x} \sin(ax)$
15	$p \ln \left(\frac{1}{p} \sqrt{p^2 - a^2} \right)$	$\frac{1}{x^2} [\cosh(ax) - 1] - \frac{a}{x} \sinh(ax)$

6.8. Expressions with Trigonometric Functions

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$\frac{\sin(a/p)}{\sqrt{p}}$	$\frac{1}{\sqrt{\pi x}} \sinh(\sqrt{2ax}) \sin(\sqrt{2ax})$
2	$\frac{\sin(a/p)}{p\sqrt{p}}$	$\frac{1}{\sqrt{\pi a}} \cosh(\sqrt{2ax}) \sin(\sqrt{2ax})$
3	$\frac{\cos(a/p)}{\sqrt{p}}$	$\frac{1}{\sqrt{\pi x}} \cosh(\sqrt{2ax}) \cos(\sqrt{2ax})$
4	$\frac{\cos(a/p)}{p\sqrt{p}}$	$\frac{1}{\sqrt{\pi a}} \sinh(\sqrt{2ax}) \cos(\sqrt{2ax})$
5	$\frac{1}{\sqrt{p}} \exp(-\sqrt{ap}) \sin(\sqrt{ap})$	$\frac{1}{\sqrt{\pi x}} \sin\left(\frac{a}{2x}\right)$
6	$\frac{1}{\sqrt{p}} \exp(-\sqrt{ap}) \cos(\sqrt{ap})$	$\frac{1}{\sqrt{\pi x}} \cos\left(\frac{a}{2x}\right)$
7	$\arctan \frac{a}{p}$	$\frac{1}{x} \sin(ax)$
8	$\frac{1}{p} \arctan \frac{a}{p}$	$\text{Si}(ax)$
9	$p \arctan \frac{a}{p} - a$	$\frac{1}{x^2} [ax \cos(ax) - \sin(ax)]$
10	$\arctan \frac{2ap}{p^2 + b^2}$	$\frac{2}{x} \sin(ax) \cos(x\sqrt{a^2 + b^2})$

6.9. Expressions with Special Functions

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
1	$\exp(ap^2) \text{erfc}(p\sqrt{a})$	$\frac{1}{\sqrt{\pi a}} \exp\left(-\frac{x^2}{4a}\right)$
2	$\frac{1}{p} \exp(ap^2) \text{erfc}(p\sqrt{a})$	$\text{erf}\left(\frac{x}{2\sqrt{a}}\right)$
3	$\text{erfc}(\sqrt{ap}), \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ \frac{\sqrt{a}}{\pi x \sqrt{x-a}} & \text{if } a < x. \end{cases}$
4	$e^{ap} \text{erfc}(\sqrt{ap})$	$\frac{\sqrt{a}}{\pi \sqrt{x} (x+a)}$
5	$\frac{1}{\sqrt{p}} e^{ap} \text{erfc}(\sqrt{ap})$	$\frac{1}{\sqrt{\pi(x+a)}}$
6	$\text{erf}(\sqrt{a/p})$	$\frac{1}{\pi x} \sin(2\sqrt{ax})$

No	Laplace transform, $\tilde{f}(p)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} \tilde{f}(p) dp$
7	$\frac{1}{\sqrt{p}} \exp(a/p) \operatorname{erf}(\sqrt{a/p})$	$\frac{1}{\sqrt{\pi x}} \sinh(2\sqrt{ax})$
8	$\frac{1}{\sqrt{p}} \exp(a/p) \operatorname{erfc}(\sqrt{a/p})$	$\frac{1}{\sqrt{\pi x}} \exp(-2\sqrt{ax})$
9	$p^{-a} \gamma(a, bp), \quad a, b > 0$	$\begin{cases} x^{a-1} & \text{if } 0 < x < b, \\ 0 & \text{if } b < x. \end{cases}$
10	$\gamma(a, b/p), \quad a > 0$	$b^{a/2} x^{a/2-1} J_a(2\sqrt{bx})$
11	$a^{-p} \gamma(p, a)$	$\exp(-ae^{-x})$
12	$K_0(ap), \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ (x^2 - a^2)^{-1/2} & \text{if } a < x. \end{cases}$
13	$K_\nu(ap), \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ \frac{\cosh[\nu \operatorname{Arcosh}(x/a)]}{\sqrt{x^2 - a^2}} & \text{if } a < x. \end{cases}$
14	$K_0(a\sqrt{p})$	$\frac{1}{2x} \exp\left(-\frac{a^2}{4x}\right)$
15	$\frac{1}{\sqrt{p}} K_1(a\sqrt{p})$	$\frac{1}{a} \exp\left(-\frac{a^2}{4x}\right)$

⊙ References for Supplement 6: G. Doetsch (1950, 1956, 1958), H. Bateman and A. Erdélyi (1954), I. I. Hirschman and D. V. Widder (1955), V. A. Ditkin and A. P. Prudnikov (1965), A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev (1992, Vol. 5).

Supplement 7

Tables of Fourier Cosine Transforms

7.1. General Formulas

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
1	$af_1(x) + bf_2(x)$	$a\check{f}_{1c}(u) + b\check{f}_{2c}(u)$
2	$f(ax), \quad a > 0$	$\frac{1}{a} \check{f}_c\left(\frac{u}{a}\right)$
3	$x^{2n} f(x), \quad n = 1, 2, \dots$	$(-1)^n \frac{d^{2n}}{du^{2n}} \check{f}_c(u)$
4	$x^{2n+1} f(ax), \quad n = 0, 1, \dots$	$(-1)^n \frac{d^{2n+1}}{du^{2n+1}} \check{f}_s(u), \quad \check{f}_s(u) = \int_0^\infty f(x) \sin(xu) dx$
5	$f(ax) \cos(bx), \quad a, b > 0$	$\frac{1}{2a} \left[\check{f}_c\left(\frac{u+b}{a}\right) + \check{f}_c\left(\frac{u-b}{a}\right) \right]$

7.2. Expressions with Power-Law Functions

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
1	$\begin{cases} 1 & \text{if } 0 < x < a, \\ 0 & \text{if } a < x \end{cases}$	$\frac{1}{u} \sin(au)$
2	$\begin{cases} x & \text{if } 0 < x < 1, \\ 2-x & \text{if } 1 < x < 2, \\ 0 & \text{if } 2 < x \end{cases}$	$\frac{4}{u^2} \cos u \sin^2 \frac{u}{2}$
3	$\frac{1}{a+x}, \quad a > 0$	$-\sin(au) \operatorname{si}(au) - \cos(au) \operatorname{Ci}(au)$
4	$\frac{1}{a^2+x^2}, \quad a > 0$	$\frac{\pi}{2a} e^{-au}$ (the integral is understood in the sense of Cauchy principal value)
5	$\frac{1}{a^2-x^2}, \quad a > 0$	$\frac{\pi \sin(au)}{2u}$
6	$\frac{a}{a^2+(b+x)^2} + \frac{a}{a^2+(b-x)^2}$	$\pi e^{-au} \cos(bu)$
7	$\frac{b+x}{a^2+(b+x)^2} + \frac{b-x}{a^2+(b-x)^2}$	$\pi e^{-au} \sin(bu)$
8	$\frac{1}{a^4+x^4}, \quad a > 0$	$\frac{1}{2} \pi a^{-3} \exp\left(-\frac{au}{\sqrt{2}}\right) \sin\left(\frac{\pi}{4} + \frac{au}{\sqrt{2}}\right)$

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
9	$\frac{1}{(a^2 + x^2)(b^2 + x^2)}, \quad a, b > 0$	$\frac{\pi}{2} \frac{ae^{-bu} - be^{-au}}{ab(a^2 - b^2)}$
10	$\frac{x^{2m}}{(x^2 + a)^{n+1}},$ $n, m = 1, 2, \dots; n + 1 > m \geq 0$	$(-1)^{n+m} \frac{\pi}{2n!} \frac{\partial^n}{\partial a^n} (a^{1/\sqrt{m}} e^{-u\sqrt{a}})$
11	$\frac{1}{\sqrt{x}}$	$\sqrt{\frac{\pi}{2u}}$
12	$\begin{cases} \frac{1}{\sqrt{x}} & \text{if } 0 < x < a, \\ 0 & \text{if } a < x \end{cases}$	$2\sqrt{\frac{\pi}{2u}} C(au),$ $C(u)$ is the Fresnel integral
13	$\begin{cases} 0 & \text{if } 0 < x < a, \\ \frac{1}{\sqrt{x}} & \text{if } a < x \end{cases}$	$\sqrt{\frac{\pi}{2u}} [1 - 2C(au)],$ $C(u)$ is the Fresnel integral
14	$\begin{cases} 0 & \text{if } 0 < x < a, \\ \frac{1}{\sqrt{x-a}} & \text{if } a < x \end{cases}$	$\sqrt{\frac{\pi}{2u}} [\cos(au) - \sin(au)]$
15	$\frac{1}{\sqrt{a^2 + x^2}}$	$K_0(au)$
16	$\begin{cases} \frac{1}{\sqrt{a^2 - x^2}} & \text{if } 0 < x < a, \\ 0 & \text{if } a < x \end{cases}$	$\frac{\pi}{2} J_0(au)$
17	$x^{-\nu}, \quad 0 < \nu < 1$	$\sin\left(\frac{1}{2}\pi\nu\right)\Gamma(1 - \nu)u^{\nu-1}$

7.3. Expressions with Exponential Functions

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
1	e^{-ax}	$\frac{a}{a^2 + u^2}$
2	$\frac{1}{x}(e^{-ax} - e^{-bx})$	$\frac{1}{2} \ln \frac{b^2 + u^2}{a^2 + u^2}$
3	$\sqrt{x}e^{-ax}$	$\frac{1}{2}\sqrt{\pi}(a^2 + u^2)^{-3/4} \cos\left(\frac{3}{2} \arctan \frac{u}{a}\right)$
4	$\frac{1}{\sqrt{x}}e^{-ax}$	$\sqrt{\frac{\pi}{2}} \left[\frac{a + (a^2 + u^2)^{1/2}}{a^2 + u^2} \right]^{1/2}$
5	$x^n e^{-ax}, \quad n = 1, 2, \dots$	$\frac{a^{n+1}n!}{(a^2 + u^2)^{n+1}} \sum_{0 \leq k \leq n+1} (-1)^k C_{n+1}^{2k} \left(\frac{u}{a}\right)^{2k}$
6	$x^{n-1/2} e^{-ax}, \quad n = 1, 2, \dots$	$k_n u \frac{\partial^n}{\partial a^n} \frac{1}{r\sqrt{r-a}},$ where $r = \sqrt{a^2 + u^2}, k_n = (-1)^n \sqrt{\pi/2}$
7	$x^{\nu-1} e^{-ax}$	$\Gamma(\nu)(a^2 + u^2)^{-\nu/2} \cos\left(\nu \arctan \frac{u}{a}\right)$

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
8	$\frac{x}{e^{ax} - 1}$	$\frac{1}{2u^2} - \frac{\pi^2}{2a^2 \sinh^2(\pi a^{-1}u)}$
9	$\frac{1}{x} \left(\frac{1}{2} - \frac{1}{x} + \frac{1}{e^x - 1} \right)$	$-\frac{1}{2} \ln(1 - e^{-2\pi u})$
10	$\exp(-ax^2)$	$\frac{1}{2} \sqrt{\frac{\pi}{a}} \exp\left(-\frac{u^2}{4a}\right)$
11	$\frac{1}{\sqrt{x}} \exp\left(-\frac{a}{x}\right)$	$\sqrt{\frac{\pi}{2u}} e^{-\sqrt{2au}} [\cos(\sqrt{2au}) - \sin(\sqrt{2au})]$
12	$\frac{1}{x\sqrt{x}} \exp\left(-\frac{a}{x}\right)$	$\sqrt{\frac{\pi}{a}} e^{-\sqrt{2au}} \cos(\sqrt{2au})$

7.4. Expressions with Hyperbolic Functions

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
1	$\frac{1}{\cosh(ax)}, \quad a > 0$	$\frac{\pi}{2a \cosh\left(\frac{1}{2}\pi a^{-1}u\right)}$
2	$\frac{1}{\cosh^2(ax)}, \quad a > 0$	$\frac{\pi u}{2a^2 \sinh\left(\frac{1}{2}\pi a^{-1}u\right)}$
3	$\frac{\cosh(ax)}{\cosh(bx)}, \quad a < b$	$\frac{\pi}{b} \left[\frac{\cos\left(\frac{1}{2}\pi ab^{-1}\right) \cosh\left(\frac{1}{2}\pi b^{-1}u\right)}{\cos(\pi ab^{-1}) + \cosh(\pi b^{-1}u)} \right]$
4	$\frac{1}{\cosh(ax) + \cos b}$	$\frac{\pi \sinh(a^{-1}bu)}{a \sin b \sinh(\pi a^{-1}u)}$
5	$\exp(-ax^2) \cosh(bx), \quad a > 0$	$\frac{1}{2} \sqrt{\frac{\pi}{a}} \exp\left(\frac{b^2 - u^2}{4a}\right) \cos\left(\frac{abu}{2}\right)$
6	$\frac{x}{\sinh(ax)}$	$\frac{\pi^2}{4a^2 \cosh^2\left(\frac{1}{2}\pi a^{-1}u\right)}$
7	$\frac{\sinh(ax)}{\sinh(bx)}, \quad a < b$	$\frac{\pi}{2b} \frac{\sin(\pi ab^{-1})}{\cos(\pi ab^{-1}) + \cosh(\pi b^{-1}u)}$
8	$\frac{1}{x} \tanh(ax), \quad a > 0$	$\ln \left[\coth\left(\frac{1}{4}\pi a^{-1}u\right) \right]$

7.5. Expressions with Logarithmic Functions

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
1	$\begin{cases} \ln x & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$	$-\frac{1}{u} \text{Si}(u)$

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
2	$\frac{\ln x}{\sqrt{x}}$	$-\sqrt{\frac{\pi}{2u}} \left[\ln(4u) + \mathcal{C} + \frac{\pi}{2} \right],$ $\mathcal{C} = 0.5772 \dots$ is the Euler constant
3	$x^{\nu-1} \ln x, \quad 0 < \nu < 1$	$\Gamma(\nu) \cos\left(\frac{\pi\nu}{2}\right) u^{-\nu} \left[\psi(\nu) - \frac{\pi}{2} \tan\left(\frac{\pi\nu}{2}\right) - \ln u \right]$
4	$\ln \left \frac{a+x}{a-x} \right , \quad a > 0$	$\frac{2}{u} [\cos(au) \text{Si}(au) - \sin(au) \text{Ci}(au)]$
5	$\ln(1 + a^2/x^2), \quad a > 0$	$\frac{\pi}{u} (1 - e^{-au})$
6	$\ln \frac{a^2 + x^2}{b^2 + x^2}, \quad a, b > 0$	$\frac{\pi}{u} (e^{-bu} - e^{-au})$
7	$e^{-ax} \ln x, \quad a > 0$	$-\frac{a\mathcal{C} + \frac{1}{2}a \ln(u^2 + a^2) + u \arctan(u/a)}{u^2 + a^2}$
8	$\ln(1 + e^{-ax}), \quad a > 0$	$\frac{a}{2u^2} - \frac{\pi}{2u \sinh(\pi a^{-1}u)}$
9	$\ln(1 - e^{-ax}), \quad a > 0$	$\frac{a}{2u^2} - \frac{\pi}{2u} \coth(\pi a^{-1}u)$

7.6. Expressions with Trigonometric Functions

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
1	$\frac{\sin(ax)}{x}, \quad a > 0$	$\begin{cases} \frac{1}{2}\pi & \text{if } u < a, \\ \frac{1}{4}\pi & \text{if } u = a, \\ 0 & \text{if } u > a \end{cases}$
2	$x^{\nu-1} \sin(ax), \quad a > 0, \nu < 1$	$\pi \frac{(u+a)^{-\nu} - u+a ^{-\nu} \text{sign}(u-a)}{4\Gamma(1-\nu) \cos(\frac{1}{2}\pi\nu)}$
3	$\frac{x \sin(ax)}{x^2 + b^2}, \quad a, b > 0$	$\begin{cases} \frac{1}{2}\pi e^{-ab} \cosh(bu) & \text{if } u < a, \\ -\frac{1}{2}\pi e^{-bu} \sinh(ab) & \text{if } u > a \end{cases}$
4	$\frac{\sin(ax)}{x(x^2 + b^2)}, \quad a, b > 0$	$\begin{cases} \frac{1}{2}\pi b^{-2} [1 - e^{-ab} \cosh(bu)] & \text{if } u < a, \\ \frac{1}{2}\pi b^{-2} e^{-bu} \sinh(ab) & \text{if } u > a \end{cases}$
5	$e^{-bx} \sin(ax), \quad a, b > 0$	$\frac{1}{2} \left[\frac{a+u}{(a+u)^2 + b^2} + \frac{a-u}{(a-u)^2 + b^2} \right]$
6	$\frac{1}{x} \sin^2(ax), \quad a > 0$	$\frac{1}{4} \ln \left 1 - 4 \frac{a^2}{u^2} \right $
7	$\frac{1}{x^2} \sin^2(ax), \quad a > 0$	$\begin{cases} \frac{1}{4}\pi(2a-u) & \text{if } u < 2a, \\ 0 & \text{if } u > 2a \end{cases}$
8	$\frac{1}{x} \sin\left(\frac{a}{x}\right), \quad a > 0$	$\frac{\pi}{2} J_0(2\sqrt{au})$

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
9	$\frac{1}{\sqrt{x}} \sin(a\sqrt{x}) \sin(b\sqrt{x}), a, b > 0$	$\sqrt{\frac{\pi}{u}} \sin\left(\frac{ab}{2u}\right) \sin\left(\frac{a^2 + b^2}{4u} - \frac{\pi}{4}\right)$
10	$\sin(ax^2), a > 0$	$\sqrt{\frac{\pi}{8a}} \left[\cos\left(\frac{u^2}{4a}\right) - \sin\left(\frac{u^2}{4a}\right) \right]$
11	$\exp(-ax^2) \sin(bx^2), a > 0$	$\frac{\sqrt{\pi}}{(A^2 + B^2)^{1/4}} \exp\left(-\frac{Au^2}{A^2 + B^2}\right) \sin\left(\varphi - \frac{Bu^2}{A^2 + B^2}\right),$ $A = 4a, B = 4b, \varphi = \frac{1}{2} \arctan(b/a)$
12	$\frac{1 - \cos(ax)}{x}, a > 0$	$\frac{1}{2} \ln \left 1 - \frac{a^2}{u^2} \right $
13	$\frac{1 - \cos(ax)}{x^2}, a > 0$	$\begin{cases} \frac{1}{2} \pi (a - u) & \text{if } u < a, \\ 0 & \text{if } u > a \end{cases}$
14	$x^{\nu-1} \cos(ax), a > 0, 0 < \nu < 1$	$\frac{1}{2} \Gamma(\nu) \cos\left(\frac{1}{2} \pi \nu\right) [u - a ^{-\nu} + (u + a)^{-\nu}]$
15	$\frac{\cos(ax)}{x^2 + b^2}, a, b > 0$	$\begin{cases} \frac{1}{2} \pi b^{-1} e^{-ab} \cosh(bu) & \text{if } u < a, \\ \frac{1}{2} \pi b^{-1} e^{-bu} \cosh(ab) & \text{if } u > a \end{cases}$
16	$e^{-bx} \cos(ax), a, b > 0$	$\frac{b}{2} \left[\frac{1}{(a + u)^2 + b^2} + \frac{1}{(a - u)^2 + b^2} \right]$
17	$\frac{1}{\sqrt{x}} \cos(a\sqrt{x})$	$\sqrt{\frac{\pi}{u}} \sin\left(\frac{a^2}{4u} + \frac{\pi}{4}\right)$
18	$\frac{1}{\sqrt{x}} \cos(a\sqrt{x}) \cos(b\sqrt{x})$	$\sqrt{\frac{\pi}{u}} \cos\left(\frac{ab}{2u}\right) \sin\left(\frac{a^2 + b^2}{4u} + \frac{\pi}{4}\right)$
19	$\exp(-bx^2) \cos(ax), b > 0$	$\frac{1}{2} \sqrt{\frac{\pi}{b}} \exp\left(-\frac{a^2 + u^2}{4b}\right) \cosh\left(\frac{au}{2b}\right)$
20	$\cos(ax^2), a > 0$	$\sqrt{\frac{\pi}{8a}} \left[\cos\left(\frac{1}{4} a^{-1} u^2\right) + \sin\left(\frac{1}{4} a^{-1} u^2\right) \right]$
21	$\exp(-ax^2) \cos(bx^2), a > 0$	$\frac{\sqrt{\pi}}{(A^2 + B^2)^{1/4}} \exp\left(-\frac{Au^2}{A^2 + B^2}\right) \cos\left(\varphi - \frac{Bu^2}{A^2 + B^2}\right),$ $A = 4a, B = 4b, \varphi = \frac{1}{2} \arctan(b/a)$

7.7. Expressions with Special Functions

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
1	$\text{Ei}(-ax)$	$-\frac{1}{u} \arctan\left(\frac{u}{a}\right)$
2	$\text{Ci}(ax)$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ -\frac{\pi}{2u} & \text{if } a < u \end{cases}$
3	$\text{si}(ax)$	$-\frac{1}{2u} \ln \left \frac{u + a}{u - a} \right , u \neq a$

No	Original function, $f(x)$	Cosine transform, $\check{f}_c(u) = \int_0^\infty f(x) \cos(ux) dx$
4	$J_0(ax), \quad a > 0$	$\begin{cases} \frac{1}{\sqrt{a^2 - u^2}} & \text{if } 0 < u < a, \\ 0 & \text{if } a < u \end{cases}$
5	$J_\nu(ax), \quad a > 0, \nu > -1$	$\begin{cases} \frac{\cos[\nu \arcsin(u/a)]}{\sqrt{a^2 - u^2}} & \text{if } 0 < u < a, \\ -\frac{a^\nu \sin(\pi\nu/2)}{\xi(u + \xi)^\nu} & \text{if } a < u, \end{cases}$ where $\xi = \sqrt{u^2 - a^2}$
6	$\frac{1}{x} J_\nu(ax), \quad a > 0, \nu > 0$	$\begin{cases} \nu^{-1} \cos[\nu \arcsin(u/a)] & \text{if } 0 < u < a, \\ \frac{a^\nu \cos(\pi\nu/2)}{\nu(u + \sqrt{u^2 - a^2})^\nu} & \text{if } a < u \end{cases}$
7	$x^{-\nu} J_\nu(ax), \quad a > 0, \nu > -\frac{1}{2}$	$\begin{cases} \frac{\sqrt{\pi} (a^2 - u^2)^{\nu-1/2}}{(2a)^\nu \Gamma(\nu + \frac{1}{2})} & \text{if } 0 < u < a, \\ 0 & \text{if } a < u \end{cases}$
8	$x^{\nu+1} J_\nu(ax),$ $a > 0, -1 < \nu < -\frac{1}{2}$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ \frac{2^{\nu+1} \sqrt{\pi} a^\nu u}{\Gamma(-\nu - \frac{1}{2})(u^2 - a^2)^{\nu+3/2}} & \text{if } a < u \end{cases}$
9	$J_0(a\sqrt{x}), \quad a > 0$	$\frac{1}{u} \sin\left(\frac{a^2}{4u}\right)$
10	$\frac{1}{\sqrt{x}} J_1(a\sqrt{x}), \quad a > 0$	$\frac{4}{a} \sin^2\left(\frac{a^2}{8u}\right)$
11	$x^{\nu/2} J_\nu(a\sqrt{x}), \quad a > 0, -1 < \nu < \frac{1}{2}$	$\left(\frac{a}{2}\right)^\nu u^{-\nu-1} \sin\left(\frac{a^2}{4u} - \frac{\pi\nu}{2}\right)$
12	$J_0(a\sqrt{x^2 + b^2})$	$\begin{cases} \frac{\cos(b\sqrt{a^2 - u^2})}{\sqrt{a^2 - u^2}} & \text{if } 0 < u < a, \\ 0 & \text{if } a < u \end{cases}$
13	$Y_0(ax), \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ -\frac{1}{\sqrt{u^2 - a^2}} & \text{if } a < u \end{cases}$
14	$x^\nu Y_\nu(ax), \quad a > 0, \nu < \frac{1}{2}$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ -\frac{(2a)^\nu \sqrt{\pi}}{\Gamma(\frac{1}{2} - \nu)(u^2 - a^2)^{\nu+1/2}} & \text{if } a < u \end{cases}$
15	$K_0(a\sqrt{x^2 + b^2}), \quad a, b > 0$	$\frac{\pi}{2\sqrt{u^2 + a^2}} \exp(-b\sqrt{u^2 + a^2})$

© References for Supplement 7: G. Doetsch (1950, 1956, 1958), H. Bateman and A. Erdélyi (1954), V. A. Ditkin and A. P. Prudnikov (1965), F. Oberhettinger (1980).

Supplement 8

Tables of Fourier Sine Transforms

8.1. General Formulas

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
1	$af_1(x) + bf_2(x)$	$a\check{f}_{1s}(u) + b\check{f}_{2s}(u)$
2	$f(ax), \quad a > 0$	$\frac{1}{a} \check{f}_s\left(\frac{u}{a}\right)$
3	$x^{2n} f(x), \quad n = 1, 2, \dots$	$(-1)^n \frac{d^{2n}}{du^{2n}} \check{f}_s(u)$
4	$x^{2n+1} f(ax), \quad n = 0, 1, \dots$	$(-1)^{n+1} \frac{d^{2n+1}}{du^{2n+1}} \check{f}_c(u), \quad \check{f}_c(u) = \int_0^\infty f(x) \cos(xu) dx$
5	$f(ax) \cos(bx), \quad a, b > 0$	$\frac{1}{2a} \left[\check{f}_s\left(\frac{u+b}{a}\right) + F_s\left(\frac{u-b}{a}\right) \right]$

8.2. Expressions with Power-Law Functions

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
1	$\begin{cases} 1 & \text{if } 0 < x < a, \\ 0 & \text{if } a < x \end{cases}$	$\frac{1}{u} [1 - \cos(au)]$
2	$\begin{cases} x & \text{if } 0 < x < 1, \\ 2-x & \text{if } 1 < x < 2, \\ 0 & \text{if } 2 < x \end{cases}$	$\frac{4}{u^2} \sin u \sin^2 \frac{u}{2}$
3	$\frac{1}{x}$	$\frac{\pi}{2}$
4	$\frac{1}{a+x}, \quad a > 0$	$\sin(au) \text{Ci}(au) - \cos(au) \text{si}(au)$
5	$\frac{x}{a^2+x^2}, \quad a > 0$	$\frac{\pi}{2} e^{-au}$
6	$\frac{1}{x(a^2+x^2)}, \quad a > 0$	$\frac{\pi}{2a^2} (1 - e^{-au})$
7	$\frac{a}{a^2+(x-b)^2} - \frac{a}{a^2+(x+b)^2}$	$\pi e^{-au} \sin(bu)$
8	$\frac{x+b}{a^2+(x+b)^2} - \frac{x-b}{a^2+(x-b)^2}$	$\pi e^{-au} \cos(bu)$

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
9	$\frac{x}{(x^2 + a^2)^n}, \quad a > 0, n = 1, 2, \dots$	$\frac{\pi u e^{-au}}{2^{2n-2}(n-1)! a^{2n-3}} \sum_{k=0}^{n-2} \frac{(2n-k-4)!}{k!(n-k-2)!} (2au)^k$
10	$\frac{x^{2m+1}}{(x^2 + a)^{n+1}},$ $n, m = 0, 1, \dots; 0 \leq m \leq n$	$(-1)^{n+m} \frac{\pi}{2n!} \frac{\partial^n}{\partial a^n} (a^m e^{-u\sqrt{a}})$
11	$\frac{1}{\sqrt{x}}$	$\sqrt{\frac{\pi}{2u}}$
12	$\frac{1}{x\sqrt{x}}$	$\sqrt{2\pi u}$
13	$x(a^2 + x^2)^{-3/2}$	$uK_0(au)$
14	$\frac{(\sqrt{a^2 + x^2} - a)^{1/2}}{\sqrt{a^2 + x^2}}$	$\sqrt{\frac{\pi}{2u}} e^{-au}$
15	$x^{-\nu}, \quad 0 < \nu < 2$	$\cos\left(\frac{1}{2}\pi\nu\right)\Gamma(1-\nu)u^{\nu-1}$

8.3. Expressions with Exponential Functions

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
1	$e^{-ax}, \quad a > 0$	$\frac{u}{a^2 + u^2}$
2	$x^n e^{-ax}, \quad a > 0, n = 1, 2, \dots$	$n! \left(\frac{a}{a^2 + u^2}\right)^{n+1} \sum_{k=0}^{[n/2]} (-1)^k C_{n+1}^{2k+1} \left(\frac{u}{a}\right)^{2k+1}$
3	$\frac{1}{x} e^{-ax}, \quad a > 0$	$\arctan \frac{u}{a}$
4	$\sqrt{x} e^{-ax}, \quad a > 0$	$\frac{\sqrt{\pi}}{2} (a^2 + u^2)^{-3/4} \sin\left(\frac{3}{2} \arctan \frac{u}{a}\right)$
5	$\frac{1}{\sqrt{x}} e^{-ax}, \quad a > 0$	$\sqrt{\frac{\pi}{2}} \frac{(\sqrt{a^2 + u^2} - a)^{1/2}}{\sqrt{a^2 + u^2}}$
6	$\frac{1}{x\sqrt{x}} e^{-ax}, \quad a > 0$	$\sqrt{2\pi} (\sqrt{a^2 + u^2} - a)^{1/2}$
7	$x^{n-1/2} e^{-ax}, \quad a > 0, n = 1, 2, \dots$	$(-1)^n \sqrt{\frac{\pi}{2}} \frac{\partial^n}{\partial a^n} \left[\frac{(\sqrt{a^2 + u^2} - a)^{1/2}}{\sqrt{a^2 + u^2}} \right]$
8	$x^{\nu-1} e^{-ax}, \quad a > 0, \nu > -1$	$\Gamma(\nu)(a^2 + u^2)^{-\nu/2} \sin\left(\nu \arctan \frac{u}{a}\right)$
9	$x^{-2} (e^{-ax} - e^{-bx}), \quad a, b > 0$	$\frac{u}{2} \ln\left(\frac{u^2 + b^2}{u^2 + a^2}\right) + b \arctan\left(\frac{u}{b}\right) - a \arctan\left(\frac{u}{a}\right)$

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
10	$\frac{1}{e^{ax} + 1}, \quad a > 0$	$\frac{1}{2u} - \frac{\pi}{2a \sinh(\pi u/a)}$
11	$\frac{1}{e^{ax} - 1}, \quad a > 0$	$\frac{\pi}{2a} \coth\left(\frac{\pi u}{a}\right) - \frac{1}{2u}$
12	$\frac{e^{x/2}}{e^x - 1}$	$-\frac{1}{2} \tanh(\pi u)$
13	$x \exp(-ax^2)$	$\frac{\sqrt{\pi}}{4a^{3/2}} u \exp\left(-\frac{u^2}{4a}\right)$
14	$\frac{1}{x} \exp(-ax^2)$	$\frac{\pi}{2} \operatorname{erf}\left(\frac{u}{2\sqrt{a}}\right)$
15	$\frac{1}{\sqrt{x}} \exp\left(-\frac{a}{x}\right)$	$\sqrt{\frac{\pi}{2u}} e^{-\sqrt{2au}} [\cos(\sqrt{2au}) + \sin(\sqrt{2au})]$
16	$\frac{1}{x\sqrt{x}} \exp\left(-\frac{a}{x}\right)$	$\sqrt{\frac{\pi}{a}} e^{-\sqrt{2au}} \sin(\sqrt{2au})$

8.4. Expressions with Hyperbolic Functions

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
1	$\frac{1}{\sinh(ax)}, \quad a > 0$	$\frac{\pi}{2a} \tanh\left(\frac{1}{2}\pi a^{-1}u\right)$
2	$\frac{x}{\sinh(ax)}, \quad a > 0$	$\frac{\pi^2 \sinh\left(\frac{1}{2}\pi a^{-1}u\right)}{4a^2 \cosh^2\left(\frac{1}{2}\pi a^{-1}u\right)}$
3	$\frac{1}{x} e^{-bx} \sinh(ax), \quad b > a $	$\frac{1}{2} \arctan\left(\frac{2au}{u^2 + b^2 - a^2}\right)$
4	$\frac{1}{x \cosh(ax)}, \quad a > 0$	$\arctan\left[\sinh\left(\frac{1}{2}\pi a^{-1}u\right)\right]$
5	$1 - \tanh\left(\frac{1}{2}ax\right), \quad a > 0$	$\frac{1}{u} - \frac{\pi}{a \sinh(\pi a^{-1}u)}$
6	$\coth\left(\frac{1}{2}ax\right) - 1, \quad a > 0$	$\frac{\pi}{a} \coth(\pi a^{-1}u) - \frac{1}{u}$
7	$\frac{\cosh(ax)}{\sinh(bx)}, \quad a < b$	$\frac{\pi}{2b} \frac{\sinh(\pi b^{-1}u)}{\cos(\pi ab^{-1}) + \cosh(\pi b^{-1}u)}$
8	$\frac{\sinh(ax)}{\cosh(bx)}, \quad a < b$	$\frac{\pi}{b} \frac{\sin\left(\frac{1}{2}\pi ab^{-1}\right) \sinh\left(\frac{1}{2}\pi b^{-1}u\right)}{\cos(\pi ab^{-1}) + \cosh(\pi b^{-1}u)}$

8.5. Expressions with Logarithmic Functions

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
1	$\begin{cases} \ln x & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$	$\frac{1}{u} [\text{Ci}(u) - \ln u - C],$ $C = 0.5772 \dots$ is the Euler constant
2	$\frac{\ln x}{x}$	$-\frac{1}{2}\pi(\ln u + C)$
3	$\frac{\ln x}{\sqrt{x}}$	$-\sqrt{\frac{\pi}{2u}} \left[\ln(4u) + C - \frac{\pi}{2} \right]$
4	$x^{\nu-1} \ln x, \quad \nu < 1$	$\frac{\pi u^{-\nu} \left[\psi(\nu) + \frac{\pi}{2} \cot\left(\frac{\pi\nu}{2}\right) - \ln u \right]}{2\Gamma(1-\nu) \cos\left(\frac{\pi\nu}{2}\right)}$
5	$\ln \left \frac{a+x}{a-x} \right , \quad a > 0$	$\frac{\pi}{u} \sin(au)$
6	$\ln \frac{(x+b)^2 + a^2}{(x-b)^2 + a^2}, \quad a, b > 0$	$\frac{2\pi}{u} e^{-au} \sin(bu)$
7	$e^{-ax} \ln x, \quad a > 0$	$\frac{a \arctan(u/a) - \frac{1}{2}u \ln(u^2 + a^2) - e^C u}{u^2 + a^2}$
8	$\frac{1}{x} \ln(1 + a^2 x^2), \quad a > 0$	$-\pi \text{Ei}\left(-\frac{u}{a}\right)$

8.6. Expressions with Trigonometric Functions

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
1	$\frac{\sin(ax)}{x}, \quad a > 0$	$\frac{1}{2} \ln \left \frac{u+a}{u-a} \right $
2	$\frac{\sin(ax)}{x^2}, \quad a > 0$	$\begin{cases} \frac{1}{2}\pi u & \text{if } 0 < u < a, \\ \frac{1}{2}\pi a & \text{if } u > a \end{cases}$
3	$x^{\nu-1} \sin(ax), \quad a > 0, -2 < \nu < 1$	$\pi \frac{ u-a ^{-\nu} - u+a ^{-\nu}}{4\Gamma(1-\nu) \sin\left(\frac{1}{2}\pi\nu\right)}, \quad \nu \neq 0$
4	$\frac{\sin(ax)}{x^2 + b^2}, \quad a, b > 0$	$\begin{cases} \frac{1}{2}\pi b^{-1} e^{-ab} \sinh(bu) & \text{if } 0 < u < a, \\ \frac{1}{2}\pi b^{-1} e^{-bu} \sinh(ab) & \text{if } u > a \end{cases}$
5	$\frac{\sin(\pi x)}{1-x^2}$	$\begin{cases} \sin u & \text{if } 0 < u < \pi, \\ 0 & \text{if } u > \pi \end{cases}$
6	$e^{-ax} \sin(bx), \quad a > 0$	$\frac{a}{2} \left[\frac{1}{a^2 + (b-u)^2} - \frac{1}{a^2 + (b+u)^2} \right]$
7	$x^{-1} e^{-ax} \sin(bx), \quad a > 0$	$\frac{1}{4} \ln \frac{(u+b)^2 + a^2}{(u-b)^2 + a^2}$
8	$\frac{1}{x} \sin^2(ax), \quad a > 0$	$\begin{cases} \frac{1}{4}\pi & \text{if } 0 < u < 2a, \\ \frac{1}{8}\pi & \text{if } u = 2a, \\ 0 & \text{if } u > 2a \end{cases}$

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
9	$\frac{1}{x^2} \sin^2(ax), \quad a > 0$	$\frac{1}{4}(u+2a) \ln u+2a + \frac{1}{4}(u-2a) \ln u-2a - \frac{1}{2}u \ln u$
10	$\exp(-ax^2) \sin(bx), \quad a > 0$	$\frac{1}{2} \sqrt{\frac{\pi}{a}} \exp\left(-\frac{u^2+b^2}{4a}\right) \sinh\left(\frac{bu}{2a}\right)$
11	$\frac{1}{x} \sin(ax) \sin(bx), \quad a \geq b > 0$	$\begin{cases} 0 & \text{if } 0 < u < a-b, \\ \frac{\pi}{4} & \text{if } a-b < u < a+b, \\ 0 & \text{if } a+b < u \end{cases}$
12	$\sin\left(\frac{a}{x}\right), \quad a > 0$	$\frac{\pi\sqrt{a}}{2\sqrt{u}} J_1(2\sqrt{au})$
13	$\frac{1}{\sqrt{x}} \sin\left(\frac{a}{x}\right), \quad a > 0$	$\sqrt{\frac{\pi}{8u}} [\sin(2\sqrt{au}) - \cos(2\sqrt{au}) + \exp(-2\sqrt{au})]$
14	$\exp(-a\sqrt{x}) \sin(a\sqrt{x}), \quad a > 0$	$a\sqrt{\frac{\pi}{8}} u^{-3/2} \exp\left(-\frac{a^2}{2u}\right)$
15	$\frac{\cos(ax)}{x}, \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ \frac{1}{4}\pi & \text{if } u = a, \\ \frac{1}{2}\pi & \text{if } a < u \end{cases}$
16	$x^{\nu-1} \cos(ax), \quad a > 0, \nu < 1$	$\frac{\pi(u+a)^{-\nu} - \text{sign}(u-a) u-a ^{-\nu}}{4\Gamma(1-\nu) \cos\left(\frac{1}{2}\pi\nu\right)}$
17	$\frac{x \cos(ax)}{x^2 + b^2}, \quad a, b > 0$	$\begin{cases} -\frac{1}{2}\pi e^{-ab} \sinh(bu) & \text{if } u < a, \\ \frac{1}{2}\pi e^{-bu} \cosh(ab) & \text{if } u > a \end{cases}$
18	$\frac{1 - \cos(ax)}{x^2}, \quad a > 0$	$\frac{u}{2} \ln\left \frac{u^2 - a^2}{u^2}\right + \frac{a}{2} \ln\left \frac{u+a}{u-a}\right $
19	$\frac{1}{\sqrt{x}} \cos(a\sqrt{x})$	$\sqrt{\frac{\pi}{u}} \cos\left(\frac{a^2}{4u} + \frac{\pi}{4}\right)$
20	$\frac{1}{\sqrt{x}} \cos(a\sqrt{x}) \cos(b\sqrt{x}), \quad a, b > 0$	$\sqrt{\frac{\pi}{u}} \cos\left(\frac{ab}{2u}\right) \cos\left(\frac{a^2+b^2}{4u} + \frac{\pi}{4}\right)$

8.7. Expressions with Special Functions

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
1	$\text{erfc}(ax), \quad a > 0$	$\frac{1}{u} \left[1 - \exp\left(-\frac{u^2}{4a^2}\right) \right]$
2	$\text{ci}(ax), \quad a > 0$	$-\frac{1}{2u} \ln\left 1 - \frac{u^2}{a^2} \right $
3	$\text{si}(ax), \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ -\frac{1}{2}\pi u^{-1} & \text{if } a < u \end{cases}$

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
4	$J_0(ax), \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ \frac{1}{\sqrt{u^2 - a^2}} & \text{if } a < u \end{cases}$
5	$J_\nu(ax), \quad a > 0, \nu > -2$	$\begin{cases} \frac{\sin[\nu \arcsin(u/a)]}{\sqrt{a^2 - u^2}} & \text{if } 0 < u < a, \\ \frac{a^\nu \cos(\pi\nu/2)}{\xi(u + \xi)^\nu} & \text{if } a < u, \end{cases}$ where $\xi = \sqrt{u^2 - a^2}$
6	$\frac{1}{x} J_0(ax), \quad a > 0, \nu > 0$	$\begin{cases} \arcsin(u/a) & \text{if } 0 < u < a, \\ \pi/2 & \text{if } a < u \end{cases}$
7	$\frac{1}{x} J_\nu(ax), \quad a > 0, \nu > -1$	$\begin{cases} \nu^{-1} \sin[\nu \arcsin(u/a)] & \text{if } 0 < u < a, \\ \frac{a^\nu \sin(\pi\nu/2)}{\nu(u + \sqrt{u^2 - a^2})^\nu} & \text{if } a < u \end{cases}$
8	$x^\nu J_\nu(ax), \quad a > 0, -1 < \nu < \frac{1}{2}$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ \frac{\sqrt{\pi}(2a)^\nu}{\Gamma(\frac{1}{2} - \nu)(u^2 - a^2)^{\nu+1/2}} & \text{if } a < u \end{cases}$
9	$x^{-1} e^{-ax} J_0(bx), \quad a > 0$	$\arcsin\left(\frac{2u}{\sqrt{(u+b)^2 + a^2} + \sqrt{(u-b)^2 + a^2}}\right)$
10	$\frac{J_0(ax)}{x^2 + b^2}, \quad a, b > 0$	$\begin{cases} b^{-1} \sinh(bu) K_0(ab) & \text{if } 0 < u < a, \\ 0 & \text{if } a < u \end{cases}$
11	$\frac{x J_0(ax)}{x^2 + b^2}, \quad a, b > 0$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ \frac{1}{2} \pi e^{-bu} I_0(ab) & \text{if } a < u \end{cases}$
12	$\frac{\sqrt{x} J_{2n+1/2}(ax)}{x^2 + b^2},$ $a, b > 0, \quad n = 0, 1, 2, \dots$	$\begin{cases} (-1)^n \sinh(bu) K_{2n+1/2}(ab) & \text{if } 0 < u < a, \\ 0 & \text{if } a < u \end{cases}$
13	$\frac{x^\nu J_\nu(ax)}{x^2 + b^2},$ $a, b > 0, \quad -1 < \nu < \frac{5}{2}$	$\begin{cases} b^{\nu-1} \sinh(bu) K_\nu(ab) & \text{if } 0 < u < a, \\ 0 & \text{if } a < u \end{cases}$
14	$\frac{x^{1-\nu} J_\nu(ax)}{x^2 + b^2},$ $a, b > 0, \quad \nu > -\frac{3}{2}$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ \frac{1}{2} \pi b^{-\nu} e^{-bu} I_\nu(ab) & \text{if } a < u \end{cases}$
15	$J_0(a\sqrt{x}), \quad a > 0$	$\frac{1}{u} \cos\left(\frac{a^2}{4u}\right)$
16	$\frac{1}{\sqrt{x}} J_1(a\sqrt{x}), \quad a > 0$	$\frac{2}{a} \sin\left(\frac{a^2}{4u}\right)$
17	$x^{\nu/2} J_\nu(a\sqrt{x}),$ $a > 0, \quad -2 < \nu < \frac{1}{2}$	$\frac{a^\nu}{2^\nu u^{\nu+1}} \cos\left(\frac{a^2}{4u} - \frac{\pi\nu}{2}\right)$

No	Original function, $f(x)$	Sine transform, $\check{f}_s(u) = \int_0^\infty f(x) \sin(ux) dx$
18	$Y_0(ax), \quad a > 0$	$\begin{cases} \frac{2 \arcsin(u/a)}{\pi \sqrt{a^2 - u^2}} & \text{if } 0 < u < a, \\ \frac{2 [\ln(u - \sqrt{u^2 - a^2}) - \ln a]}{\pi \sqrt{u^2 - a^2}} & \text{if } a < u \end{cases}$
19	$Y_1(ax), \quad a > 0$	$\begin{cases} 0 & \text{if } 0 < u < a, \\ -\frac{u}{a \sqrt{u^2 - a^2}} & \text{if } a < u \end{cases}$
20	$K_0(ax), \quad a > 0$	$\frac{\ln(u + \sqrt{u^2 + a^2}) - \ln a}{\sqrt{u^2 + a^2}}$
21	$xK_0(ax), \quad a > 0$	$\frac{\pi u}{2(u^2 + a^2)^{3/2}}$
22	$x^{\nu+1} K_\nu(ax), \quad a > 0, \nu > -\frac{3}{2}$	$\sqrt{\pi} (2a)^\nu \Gamma\left(\nu + \frac{3}{2}\right) u (u^2 + a^2)^{-\nu-3/2}$

⊙ References for Supplement 8: G. Doetsch (1950, 1956, 1958), H. Bateman and A. Erdélyi (1954), I. I. Hirschman and D. V. Widder (1955), V. A. Ditkin and A. P. Prudnikov (1965), F. Oberhettinger (1980).

Supplement 9

Tables of Mellin Transforms

9.1. General Formulas

No	Original function, $f(x)$	Mellin transform, $\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx$
1	$af_1(x) + bf_2(x)$	$a\hat{f}_1(s) + b\hat{f}_2(s)$
2	$f(ax), a > 0$	$a^{-s}\hat{f}(s)$
3	$x^a f(x)$	$\hat{f}(s+a)$
4	$f(1/x)$	$\hat{f}(-s)$
5	$f(x^\beta), \beta > 0$	$\frac{1}{\beta}\hat{f}\left(\frac{s}{\beta}\right)$
6	$f(x^{-\beta}), \beta > 0$	$\frac{1}{\beta}\hat{f}\left(-\frac{s}{\beta}\right)$
7	$x^\lambda f(ax^\beta), a, \beta > 0$	$\frac{1}{\beta}a^{-\frac{s+\lambda}{\beta}}\hat{f}\left(\frac{s+\lambda}{\beta}\right)$
8	$x^\lambda f(ax^{-\beta}), a, \beta > 0$	$\frac{1}{\beta}a^{\frac{s+\lambda}{\beta}}\hat{f}\left(-\frac{s+\lambda}{\beta}\right)$
9	$f'_x(x)$	$-(s-1)\hat{f}(s-1)$
10	$xf'_x(x)$	$-s\hat{f}(s)$
11	$f_x^{(n)}(x)$	$(-1)^n \frac{\Gamma(s)}{\Gamma(s-n)}\hat{f}(s-n)$
12	$\left(x\frac{d}{dx}\right)^n f(x)$	$(-1)^n s^n \hat{f}(s)$
13	$\left(\frac{d}{dx}x\right)^n f(x)$	$(-1)^n (s-1)^n \hat{f}(s)$
14	$x^\alpha \int_0^\infty t^\beta f_1(xt)f_2(t) dt$	$\hat{f}_1(s+\alpha)\hat{f}_2(1-s-\alpha+\beta)$
15	$x^\alpha \int_0^\infty t^\beta f_1\left(\frac{x}{t}\right)f_2(t) dt$	$\hat{f}_1(s+\alpha)\hat{f}_2(s+\alpha+\beta+1)$

9.2. Expressions with Power-Law Functions

No	Original function, $f(x)$	Mellin transform, $\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx$
1	$\begin{cases} x & \text{if } 0 < x < 1, \\ 2-x & \text{if } 1 < x < 2, \\ 0 & \text{if } 2 < x \end{cases}$	$\begin{cases} \frac{2(2^s-1)}{s(s+1)} & \text{if } s \neq 0, \\ 2 \ln 2 & \text{if } s = 0, \end{cases} \quad \text{Re } s > -1$
2	$\frac{1}{x+a}, \quad a > 0$	$\frac{\pi a^{s-1}}{\sin(\pi s)}, \quad 0 < \text{Re } s < 1$
3	$\frac{1}{(x+a)(x+b)}, \quad a, b > 0$	$\frac{\pi(a^{s-1}-b^{s-1})}{(b-a)\sin(\pi s)}, \quad 0 < \text{Re } s < 2$
4	$\frac{x+a}{(x+b)(x+c)}, \quad b, c > 0$	$\frac{\pi}{\sin(\pi s)} \left[\left(\frac{b-a}{b-c}\right)b^{s-1} + \left(\frac{c-a}{c-b}\right)c^{s-1} \right],$ $0 < \text{Re } s < 1$
5	$\frac{1}{x^2+a^2}, \quad a > 0$	$\frac{\pi a^{s-2}}{2 \sin(\frac{1}{2}\pi s)}, \quad 0 < \text{Re } s < 2$
6	$\frac{1}{x^2+2ax \cos \beta+a^2}, \quad a > 0, \beta < \pi$	$-\frac{\pi a^{s-2} \sin[\beta(s-1)]}{\sin \beta \sin(\pi s)}, \quad 0 < \text{Re } s < 2$
7	$\frac{1}{(x^2+a^2)(x^2+b^2)}, \quad a, b > 0$	$\frac{\pi(a^{s-2}-b^{s-2})}{2(b^2-a^2)\sin(\frac{1}{2}\pi s)}, \quad 0 < \text{Re } s < 4$
8	$\frac{1}{(1+ax)^{n+1}}, \quad a > 0, n = 1, 2, \dots$	$\frac{(-1)^n \pi}{a^s \sin(\pi s)} C_{s-1}^n, \quad 0 < \text{Re } s < n+1$
9	$\frac{1}{x^n+a^n}, \quad a > 0, n = 1, 2, \dots$	$\frac{\pi a^{s-n}}{n \sin(\pi s/n)}, \quad 0 < \text{Re } s < n$
10	$\frac{1-x}{1-x^n}, \quad n = 2, 3, \dots$	$\frac{\pi \sin(\pi/n)}{n \sin(\pi s/n) \sin[\pi(s+1)/n]}, \quad 0 < \text{Re } s < n-1$
11	$\begin{cases} x^\nu & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$	$\frac{1}{s+\nu}, \quad \text{Re } s > -\nu$
12	$\frac{1-x^\nu}{1-x^{n\nu}}, \quad n = 2, 3, \dots$	$\frac{\pi \sin(\pi/n)}{n\nu \sin(\frac{\pi s}{n\nu}) \sin[\frac{\pi(s+\nu)}{n\nu}]}, \quad 0 < \text{Re } s < (n-1)\nu$

9.3. Expressions with Exponential Functions

No	Original function, $f(x)$	Mellin transform, $\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx$
1	$e^{-ax}, \quad a > 0$	$a^{-s}\Gamma(s), \quad \text{Re } s > 0$
2	$\begin{cases} e^{-bx} & \text{if } 0 < x < a, \\ 0 & \text{if } a < x, \end{cases} \quad b > 0$	$b^{-s}\gamma(s, ab), \quad \text{Re } s > 0$
3	$\begin{cases} 0 & \text{if } 0 < x < a, \\ e^{-bx} & \text{if } a < x, \end{cases} \quad b > 0$	$b^{-s}\Gamma(s, ab)$
4	$\frac{e^{-ax}}{x+b}, \quad a, b > 0$	$e^{ab}b^{s-1}\Gamma(s)\Gamma(1-s, ab), \quad \text{Re } s > 0$
5	$\exp(-ax^\beta), \quad a, \beta > 0$	$\beta^{-1}a^{-s/\beta}\Gamma(s/\beta), \quad \text{Re } s > 0$

No	Original function, $f(x)$	Mellin transform, $\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx$
6	$\exp(-ax^{-\beta}), \quad a, \beta > 0$	$\beta^{-1} a^{s/\beta} \Gamma(-s/\beta), \quad \operatorname{Re} s < 0$
7	$1 - \exp(-ax^\beta), \quad a, \beta > 0$	$-\beta^{-1} a^{-s/\beta} \Gamma(s/\beta), \quad -\beta < \operatorname{Re} s < 0$
8	$1 - \exp(-ax^{-\beta}), \quad a, \beta > 0$	$-\beta^{-1} a^{s/\beta} \Gamma(-s/\beta), \quad 0 < \operatorname{Re} s < \beta$

9.4. Expressions with Logarithmic Functions

No	Original function, $f(x)$	Mellin transform, $\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx$
1	$\begin{cases} \ln x & \text{if } 0 < x < a, \\ 0 & \text{if } a < x \end{cases}$	$\frac{s \ln a - 1}{s^2 a^s}, \quad \operatorname{Re} s > 0$
2	$\ln(1 + ax), \quad a > 0$	$\frac{\pi}{sa^s \sin(\pi s)}, \quad -1 < \operatorname{Re} s < 0$
3	$\ln 1 - x $	$\frac{\pi}{s} \cot(\pi s), \quad -1 < \operatorname{Re} s < 0$
4	$\frac{\ln x}{x + a}, \quad a > 0$	$\frac{\pi a^{s-1} [\ln a - \pi \cot(\pi s)]}{\sin(\pi s)}, \quad 0 < \operatorname{Re} s < 1$
5	$\frac{\ln x}{(x + a)(x + b)}, \quad a, b > 0$	$\frac{\pi [a^{s-1} \ln a - b^{s-1} \ln b - \pi \cot(\pi s)(a^{s-1} - b^{s-1})]}{(b - a) \sin(\pi s)},$ $0 < \operatorname{Re} s < 1$
6	$\begin{cases} x^\nu \ln x & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$	$-\frac{1}{(s + \nu)^2}, \quad \operatorname{Re} s > -\nu$
7	$\frac{\ln^2 x}{x + 1}$	$\frac{\pi^3 [2 - \sin^2(\pi s)]}{\sin^3(\pi s)}, \quad 0 < \operatorname{Re} s < 1$
8	$\begin{cases} \ln^{\nu-1} x & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$	$\Gamma(\nu)(-s)^{-\nu}, \quad \operatorname{Re} s < 0, \nu > 0$
9	$\ln(x^2 + 2x \cos \beta + 1), \quad \beta < \pi$	$\frac{2\pi \cos(\beta s)}{s \sin(\pi s)}, \quad -1 < \operatorname{Re} s < 0$
10	$\ln \left \frac{1+x}{1-x} \right $	$\frac{\pi}{s} \tan\left(\frac{1}{2}\pi s\right), \quad -1 < \operatorname{Re} s < 1$
11	$e^{-x} \ln^n x, \quad n = 1, 2, \dots$	$\frac{d^n}{ds^n} \Gamma(s), \quad \operatorname{Re} s > 0$

9.5. Expressions with Trigonometric Functions

No	Original function, $f(x)$	Mellin transform, $\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx$
1	$\sin(ax), \quad a > 0$	$a^{-s} \Gamma(s) \sin\left(\frac{1}{2}\pi s\right), \quad -1 < \operatorname{Re} s < 1$
2	$\sin^2(ax), \quad a > 0$	$-2^{-s-1} a^{-s} \Gamma(s) \cos\left(\frac{1}{2}\pi s\right), \quad -2 < \operatorname{Re} s < 0$
3	$\sin(ax) \sin(bx), \quad a, b > 0, a \neq b$	$\frac{1}{2} \Gamma(s) \cos\left(\frac{1}{2}\pi s\right) [b^{-s} - (b+a)^{-s}],$ $-2 < \operatorname{Re} s < 1$

No	Original function, $f(x)$	Mellin transform, $\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx$
4	$\cos(ax), \quad a > 0$	$a^{-s}\Gamma(s) \cos(\frac{1}{2}\pi s), \quad 0 < \operatorname{Re} s < 1$
5	$\sin(ax) \cos(bx), \quad a, b > 0$	$\frac{\Gamma(s)}{2} \sin\left(\frac{\pi s}{2}\right) [(a+b)^{-s} + a-b ^{-s} \operatorname{sign}(a-b)],$ $-1 < \operatorname{Re} s < 1$
6	$e^{-ax} \sin(bx), \quad a > 0$	$\frac{\Gamma(s) \sin[s \arctan(b/a)]}{(a^2 + b^2)^{s/2}}, \quad -1 < \operatorname{Re} s$
7	$e^{-ax} \cos(bx), \quad a > 0$	$\frac{\Gamma(s) \cos[s \arctan(b/a)]}{(a^2 + b^2)^{s/2}}, \quad 0 < \operatorname{Re} s$
8	$\begin{cases} \sin(a \ln x) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$	$-\frac{a}{s^2 + a^2}, \quad \operatorname{Re} s > 0$
9	$\begin{cases} \cos(a \ln x) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$	$\frac{s}{s^2 + a^2}, \quad \operatorname{Re} s > 0$
10	$\arctan x$	$-\frac{\pi}{2s \cos(\frac{1}{2}\pi s)}, \quad -1 < \operatorname{Re} s < 0$
11	$\operatorname{arccot} x$	$\frac{\pi}{2s \cos(\frac{1}{2}\pi s)}, \quad 0 < \operatorname{Re} s < 1$

9.6. Expressions with Special Functions

No	Original function, $f(x)$	Mellin transform, $\hat{f}(s) = \int_0^\infty f(x)x^{s-1} dx$
1	$\operatorname{erfc} x$	$\frac{\Gamma(\frac{1}{2}s + \frac{1}{2})}{\sqrt{\pi} s}, \quad \operatorname{Re} s > 0$
2	$\operatorname{Ei}(-x)$	$-s^{-1}\Gamma(s), \quad \operatorname{Re} s > 0$
3	$\operatorname{Si}(x)$	$-s^{-1} \sin(\frac{1}{2}\pi s)\Gamma(s), \quad -1 < \operatorname{Re} s < 0$
4	$\operatorname{si}(x)$	$-4s^{-1} \sin(\frac{1}{2}\pi s)\Gamma(s), \quad -1 < \operatorname{Re} s < 0$
5	$\operatorname{Ci}(x)$	$-s^{-1} \cos(\frac{1}{2}\pi s)\Gamma(s), \quad 0 < \operatorname{Re} s < 1$
6	$J_\nu(ax), \quad a > 0$	$\frac{2^{s-1}\Gamma(\frac{1}{2}\nu + \frac{1}{2}s)}{a^s\Gamma(\frac{1}{2}\nu - \frac{1}{2}s + 1)}, \quad -\nu < \operatorname{Re} s < \frac{3}{2}$
7	$Y_\nu(ax), \quad a > 0$	$-\frac{2^{s-1}}{\pi a^s} \Gamma\left(\frac{s}{2} + \frac{\nu}{2}\right) \Gamma\left(\frac{s}{2} - \frac{\nu}{2}\right) \cos\left[\frac{\pi(s-\nu)}{2}\right],$ $ \nu < \operatorname{Re} s < \frac{3}{2}$
8	$e^{-ax} I_\nu(ax), \quad a > 0$	$\frac{\Gamma(1/2 - s)\Gamma(s + \nu)}{\sqrt{\pi} (2a)^s \Gamma(1 + \nu - s)}, \quad -\nu < \operatorname{Re} s < \frac{1}{2}$
9	$K_\nu(ax), \quad a > 0$	$\frac{2^{s-2}}{a^s} \Gamma\left(\frac{s}{2} + \frac{\nu}{2}\right) \Gamma\left(\frac{s}{2} - \frac{\nu}{2}\right), \quad \nu < \operatorname{Re} s$
10	$e^{-ax} K_\nu(ax), \quad a > 0$	$\frac{\sqrt{\pi} \Gamma(s - \nu)\Gamma(s + \nu)}{(2a)^s \Gamma(s + 1/2)}, \quad \nu < \operatorname{Re} s$

Supplement 10

Tables of Inverse Mellin Transforms

See Section 9.1 of Supplement 9 for general formulas.

10.1. Expressions with Power-Law Functions

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
1	$\frac{1}{s}, \operatorname{Re} s > 0$	$\begin{cases} 1 & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
2	$\frac{1}{s}, \operatorname{Re} s < 0$	$\begin{cases} 0 & \text{if } 0 < x < 1, \\ -1 & \text{if } 1 < x \end{cases}$
3	$\frac{1}{s+a}, \operatorname{Re} s > -a$	$\begin{cases} x^a & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
4	$\frac{1}{s+a}, \operatorname{Re} s < -a$	$\begin{cases} 0 & \text{if } 0 < x < 1, \\ -x^a & \text{if } 1 < x \end{cases}$
5	$\frac{1}{(s+a)^2}, \operatorname{Re} s > -a$	$\begin{cases} -x^a \ln x & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
6	$\frac{1}{(s+a)^2}, \operatorname{Re} s < -a$	$\begin{cases} 0 & \text{if } 0 < x < 1, \\ x^a \ln x & \text{if } 1 < x \end{cases}$
7	$\frac{1}{(s+a)(s+b)}, \operatorname{Re} s > -a, -b$	$\begin{cases} \frac{x^a - x^b}{b-a} & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
8	$\frac{1}{(s+a)(s+b)}, -a < \operatorname{Re} s < -b$	$\begin{cases} \frac{x^a}{b-a} & \text{if } 0 < x < 1, \\ \frac{x^b}{b-a} & \text{if } 1 < x \end{cases}$
9	$\frac{1}{(s+a)(s+b)}, \operatorname{Re} s < -a, -b$	$\begin{cases} 0 & \text{if } 0 < x < 1, \\ \frac{x^b - x^a}{b-a} & \text{if } 1 < x \end{cases}$
10	$\frac{1}{(s+a)^2 + b^2}, \operatorname{Re} s > -a$	$\begin{cases} \frac{1}{b} x^a \sin\left(b \ln \frac{1}{x}\right) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
11	$\frac{s+a}{(s+a)^2 + b^2}, \operatorname{Re} s > -a$	$\begin{cases} x^a \cos(b \ln x) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
12	$\sqrt{s^2 - a^2} - s, \quad \text{Re } s > a $	$\begin{cases} -\frac{a}{\ln x} I_1(-a \ln x) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
13	$\sqrt{\frac{s+a}{s-a}} - 1, \quad \text{Re } s > a $	$\begin{cases} aI_0(-a \ln x) + aI_1(-a \ln x) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
14	$(s+a)^{-\nu}, \quad \text{Re } s > -a, \nu > 0$	$\begin{cases} \frac{1}{\Gamma(\nu)} x^a (-\ln x)^{\nu-1} & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
15	$s^{-1}(s+a)^{-\nu},$ $\text{Re } s > 0, \text{Re } s > -a, \nu > 0$	$\begin{cases} a^{-\nu} [\Gamma(\nu)]^{-1} \gamma(\nu, -a \ln x) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
16	$s^{-1}(s+a)^{-\nu},$ $-a < \text{Re } s < 0, \nu > 0$	$\begin{cases} -a^{-\nu} [\Gamma(\nu)]^{-1} \Gamma(\nu, -a \ln x) & \text{if } 0 < x < 1, \\ -a^{-\nu} & \text{if } 1 < x \end{cases}$
17	$(s^2 - a^2)^{-\nu}, \quad \text{Re } s > a , \nu > 0$	$\begin{cases} \frac{\sqrt{\pi} (-\ln x)^{\nu-1/2} I_{\nu-1/2}(-a \ln x)}{\Gamma(\nu)(2a)^{\nu-1/2}} & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
18	$(a^2 - s^2)^{-\nu}, \quad \text{Re } s < a , \nu > 0$	$\begin{cases} \frac{(-\ln x)^{\nu-1/2} K_{\nu-1/2}(-a \ln x)}{\sqrt{\pi} \Gamma(\nu)(2a)^{\nu-1/2}} & \text{if } 0 < x < 1, \\ \frac{(\ln x)^{\nu-1/2} K_{\nu-1/2}(a \ln x)}{\sqrt{\pi} \Gamma(\nu)(2a)^{\nu-1/2}} & \text{if } 1 < x \end{cases}$

10.2. Expressions with Exponential and Logarithmic Functions

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
1	$\exp(as^2), \quad a > 0$	$\frac{1}{2\sqrt{\pi a}} \exp\left(-\frac{\ln^2 x}{4a}\right)$
2	$s^{-\nu} e^{-a/s}, \quad \text{Re } s > 0; a, \nu > 0$	$\begin{cases} \left \frac{a}{\ln x}\right ^{\frac{1-\nu}{2}} J_{\nu-1}(2\sqrt{a \ln x }) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
3	$\exp(-\sqrt{as}), \quad \text{Re } s > 0, a > 0$	$\begin{cases} \frac{(a/\pi)^{1/2}}{2 \ln x ^{3/2}} \exp\left(-\frac{a}{4 \ln x }\right) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
4	$\frac{1}{s} \exp(-a\sqrt{s}), \quad \text{Re } s > 0$	$\begin{cases} \text{erfc}\left(\frac{a}{2\sqrt{ \ln x }}\right) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
5	$\frac{1}{s} [\exp(-a\sqrt{s}) - 1], \quad \text{Re } s > 0$	$\begin{cases} -\text{erf}\left(\frac{a}{2\sqrt{ \ln x }}\right) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
6	$\sqrt{s} \exp(-\sqrt{as}), \operatorname{Re} s > 0$	$\begin{cases} \frac{a-2 \ln x }{4\sqrt{\pi} \ln x ^5} \exp\left(-\frac{a}{4 \ln x }\right) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
7	$\frac{1}{\sqrt{s}} \exp(-\sqrt{as}), \operatorname{Re} s > 0$	$\begin{cases} \frac{1}{\sqrt{\pi} \ln x } \exp\left(-\frac{a}{4 \ln x }\right) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
8	$\ln \frac{s+a}{s+b}, \operatorname{Re} s > -a, -b$	$\begin{cases} \frac{x^a - x^b}{\ln x} & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
9	$s^{-\nu} \ln s, \operatorname{Re} s > 0, \nu > 0$	$\begin{cases} \ln x ^{\nu-1} \frac{\psi(\nu) - \ln \ln x }{\Gamma(\nu)} & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$

10.3. Expressions with Trigonometric Functions

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
1	$\frac{\pi}{\sin(\pi s)}, 0 < \operatorname{Re} s < 1$	$\frac{1}{x+1}$
2	$\frac{\pi}{\sin(\pi s)}, -n < \operatorname{Re} s < 1-n,$ $n = \dots, -1, 0, 1, 2, \dots$	$(-1)^n \frac{x^n}{x+1}$
3	$\frac{\pi^2}{\sin^2(\pi s)}, 0 < \operatorname{Re} s < 1$	$\frac{\ln x}{x-1}$
4	$\frac{\pi^2}{\sin^2(\pi s)}, n < \operatorname{Re} s < n+1,$ $n = \dots, -1, 0, 1, 2, \dots$	$\frac{\ln x}{x^n(x-1)}$
5	$\frac{2\pi^3}{\sin^3(\pi s)}, 0 < \operatorname{Re} s < 1$	$\frac{\pi^2 + \ln^2 x}{x+1}$
6	$\frac{2\pi^3}{\sin^3(\pi s)}, n < \operatorname{Re} s < n+1,$ $n = \dots, -1, 0, 1, 2, \dots$	$\frac{\pi^2 + \ln^2 x}{(-x)^n(x+1)}$
7	$\sin(s^2/a), a > 0$	$\frac{1}{2} \sqrt{\frac{a}{\pi}} \sin\left(\frac{1}{4}a \ln x ^2 - \frac{1}{4}\pi\right)$
8	$\frac{\pi}{\cos(\pi s)}, -\frac{1}{2} < \operatorname{Re} s < \frac{1}{2}$	$\frac{\sqrt{x}}{x+1}$
9	$\frac{\pi}{\cos(\pi s)}, n - \frac{1}{2} < \operatorname{Re} s < n + \frac{1}{2}$ $n = \dots, -1, 0, 1, 2, \dots$	$(-1)^n \frac{x^{1/2-n}}{x+1}$
10	$\frac{\cos(\beta s)}{s \cos(\pi s)}, -1 < \operatorname{Re} s < 0, \beta < \pi$	$\frac{1}{2\pi} \ln(x^2 + 2x \cos \beta + 1)$

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
11	$\cos(s^2/a), \quad a > 0$	$\frac{1}{2} \sqrt{\frac{a}{\pi}} \cos(\frac{1}{4}a \ln x ^2 - \frac{1}{4}\pi)$
12	$\arctan\left(\frac{a}{s+b}\right), \quad \operatorname{Re} s > -b$	$\begin{cases} \frac{x^b}{ \ln x } \sin(a \ln x) & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$

10.4. Expressions with Special Functions

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
1	$\Gamma(s), \quad \operatorname{Re} s > 0$	e^{-x}
2	$\Gamma(s), \quad -1 < \operatorname{Re} s < 0$	$e^{-x} - 1$
3	$\sin(\frac{1}{2}\pi s)\Gamma(s), \quad -1 < \operatorname{Re} s < 1$	$\sin x$
4	$\frac{\sin(as)\Gamma(s)}{\operatorname{Re} s > -1, a < \frac{\pi}{2}}$	$\exp(-x \cos a) \sin(x \sin a)$
5	$\cos(\frac{1}{2}\pi s)\Gamma(s), \quad 0 < \operatorname{Re} s < 1$	$\cos x$
6	$\cos(\frac{1}{2}\pi s)\Gamma(s), \quad -2 < \operatorname{Re} s < 0$	$-2 \sin^2(x/2)$
7	$\cos(as)\Gamma(s), \quad \operatorname{Re} s > 0, a < \frac{\pi}{2}$	$\exp(-x \cos a) \cos(x \sin a)$
8	$\frac{\Gamma(s)}{\cos(\pi s)}, \quad 0 < \operatorname{Re} s < \frac{1}{2}$	$e^x \operatorname{erfc}(\sqrt{x})$
9	$\frac{\Gamma(a+s)\Gamma(b-s)}{-a < \operatorname{Re} s < b, a+b > 0}$	$\Gamma(a+b)x^a(x+1)^{-a-b}$
10	$\frac{\Gamma(a+s)\Gamma(b+s)}{\operatorname{Re} s > -a, -b}$	$2x^{(a+b)/2} K_{a-b}(2\sqrt{x})$
11	$\frac{\Gamma(s)}{\Gamma(s+\nu)}, \quad \operatorname{Re} s > 0, \nu > 0$	$\begin{cases} \frac{(1-x)^{\nu-1}}{\Gamma(\nu)} & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
12	$\frac{\Gamma(1-\nu-s)}{\Gamma(1-s)}, \quad \operatorname{Re} s < 1-\nu, \nu > 0$	$\begin{cases} 0 & \text{if } 0 < x < 1, \\ \frac{(x-1)^{\nu-1}}{\Gamma(\nu)} & \text{if } 1 < x \end{cases}$
13	$\frac{\Gamma(s)}{\Gamma(\nu-s+1)}, \quad 0 < \operatorname{Re} s < \frac{\nu}{2} + \frac{3}{4}$	$x^{-\nu/2} J_\nu(2\sqrt{x})$
14	$\frac{\Gamma(s+\nu)\Gamma(s-\nu)}{\Gamma(s+1/2)}, \quad \operatorname{Re} s > \nu $	$\pi^{-1/2} e^{-x/2} K_\nu(x/2)$

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
15	$\frac{\Gamma(s+\nu)\Gamma(1/2-s)}{\Gamma(1+\nu-s)},$ $-\nu < \operatorname{Re} s < \frac{1}{2}$	$\pi^{1/2} e^{-x/2} I_\nu(x/2)$
16	$\psi(s+a) - \psi(s+b),$ $\operatorname{Re} s > -a, -b$	$\begin{cases} \frac{x^b - x^a}{1-x} & \text{if } 0 < x < 1, \\ 0 & \text{if } 1 < x \end{cases}$
17	$\Gamma(s)\psi(s), \operatorname{Re} s > 0$	$e^{-x} \ln x$
18	$\Gamma(s, a), a > 0$	$\begin{cases} 0 & \text{if } 0 < x < a, \\ e^{-x} & \text{if } a < x \end{cases}$
19	$\Gamma(s)\Gamma(1-s, a), \operatorname{Re} s > 0, a > 0$	$(x+1)^{-1} e^{-a(x+1)}$
20	$\gamma(s, a), \operatorname{Re} s > 0, a > 0$	$\begin{cases} e^{-x} & \text{if } 0 < x < a, \\ 0 & \text{if } a < x \end{cases}$
21	$J_0(a\sqrt{b^2-s^2}), a > 0$	$\begin{cases} 0 & \text{if } 0 < x < e^{-a}, \\ \frac{\cos(b\sqrt{a^2-\ln^2 x})}{\pi\sqrt{a^2-\ln^2 x}} & \text{if } e^{-a} < x < e^a, \\ 0 & \text{if } e^a < x \end{cases}$
22	$s^{-1} I_0(s), \operatorname{Re} s > 0$	$\begin{cases} 1 & \text{if } 0 < x < e^{-1}, \\ \pi^{-1} \arccos(\ln x) & \text{if } e^{-1} < x < e, \\ 0 & \text{if } e < x \end{cases}$
23	$I_\nu(s), \operatorname{Re} s > 0$	$\begin{cases} -\frac{2^\nu \sin(\pi\nu)}{\pi F(x)\sqrt{\ln^2 x - 1}} & \text{if } 0 < x < e^{-1}, \\ \frac{\cos[\nu \arccos(\ln x)]}{\pi\sqrt{1-\ln^2 x}} & \text{if } e^{-1} < x < e, \\ 0 & \text{if } e < x, \end{cases}$ $F(x) = (\sqrt{-1-\ln x} + \sqrt{1-\ln x})^{2\nu}$
24	$s^{-1} I_\nu(s), \operatorname{Re} s > 0$	$\begin{cases} \frac{2^\nu \sin(\pi\nu)}{\pi\nu F(x)} & \text{if } 0 < x < e^{-1}, \\ \frac{\sin[\nu \arccos(\ln x)]}{\pi\nu} & \text{if } e^{-1} < x < e, \\ 0 & \text{if } e < x, \end{cases}$ $F(x) = (\sqrt{-1-\ln x} + \sqrt{1-\ln x})^{2\nu}$
25	$s^{-\nu} I_\nu(s), \operatorname{Re} s > -\frac{1}{2}$	$\begin{cases} 0 & \text{if } 0 < x < e^{-1}, \\ \frac{(1-\ln^2 x)^{\nu-1/2}}{\sqrt{\pi} 2^\nu \Gamma(\nu+1/2)} & \text{if } e^{-1} < x < e, \\ 0 & \text{if } e < x \end{cases}$
26	$s^{-1} K_0(s), \operatorname{Re} s > 0$	$\begin{cases} \operatorname{Arcosh}(-\ln x) & \text{if } 0 < x < e^{-1}, \\ 0 & \text{if } e^{-1} < x \end{cases}$
27	$s^{-1} K_1(s), \operatorname{Re} s > 0$	$\begin{cases} \sqrt{\ln^2 x - 1} & \text{if } 0 < x < e^{-1}, \\ 0 & \text{if } e^{-1} < x \end{cases}$

No	Direct transform, $\hat{f}(s)$	Inverse transform, $f(x) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \hat{f}(s)x^{-s} ds$
28	$K_\nu(s), \quad \operatorname{Re} s > 0$	$\begin{cases} \frac{\cosh[\nu \operatorname{Arcosh}(-\ln x)]}{\sqrt{\ln^2 x - 1}} & \text{if } 0 < x < e^{-1}, \\ 0 & \text{if } e^{-1} < x \end{cases}$
29	$s^{-1}K_\nu(s), \quad \operatorname{Re} s > 0$	$\begin{cases} \frac{1}{\nu} \sinh[\nu \operatorname{Arcosh}(-\ln x)] & \text{if } 0 < x < e^{-1}, \\ 0 & \text{if } e^{-1} < x \end{cases}$
30	$s^{-\nu}K_\nu(s), \quad \operatorname{Re} s > 0, \nu > -\frac{1}{2}$	$\begin{cases} \frac{\sqrt{\pi} (\ln^2 x - 1)^{\nu-1/2}}{2^\nu \Gamma(\nu + 1/2)} & \text{if } 0 < x < e^{-1}, \\ 0 & \text{if } e^{-1} < x \end{cases}$

⊙ References for Supplement 10: H. Bateman and A. Erdélyi (1954), V. A. Ditkin and A. P. Prudnikov (1965).

Supplement 11

Special Functions and Their Properties

► Throughout Supplement 11 it is assumed that n is a positive integer, unless otherwise specified.

11.1. Some Coefficients, Symbols, and Numbers

11.1-1. Binomial Coefficients.

Definitions (special cases):

$$C_n^k = \binom{n}{k} = \frac{n!}{k!(n-k)!}, \quad \text{where } k = 1, \dots, n;$$

$$C_a^0 = 1, \quad C_a^k = \binom{a}{k} = (-1)^k \frac{(-a)_k}{k!} = \frac{a(a-1)\dots(a-k+1)}{k!}, \quad \text{where } k = 1, 2, \dots$$

Here a is an arbitrary real number.

Definition (general case):

$$C_a^b = \frac{\Gamma(a+1)}{\Gamma(b+1)\Gamma(a-b+1)}, \quad \text{where } \Gamma(x) \text{ is the gamma function.}$$

Properties:

$$C_a^0 = 1, \quad C_n^k = 0 \quad \text{for } k = -1, -2, \dots \text{ or } k > n,$$

$$C_a^{b+1} = \frac{a}{b+1} C_{a-1}^b = \frac{a-b}{b+1} C_a^b, \quad C_a^b + C_a^{b+1} = C_{a+1}^{b+1},$$

$$C_{-1/2}^n = \frac{(-1)^n}{2^{2n}} C_{2n}^n = (-1)^n \frac{(2n-1)!!}{(2n)!!},$$

$$C_{1/2}^n = \frac{(-1)^{n-1}}{n2^{2n-1}} C_{2n-2}^{n-1} = \frac{(-1)^{n-1}}{n} \frac{(2n-3)!!}{(2n-2)!!},$$

$$C_{n+1/2}^{2n+1} = (-1)^n 2^{-4n-1} C_{2n}^n, \quad C_{2n+1/2}^{2n} = 2^{-2n} C_{4n+1}^{2n},$$

$$C_n^{1/2} = \frac{2^{2n+1}}{\pi C_{2n}^n}, \quad C_n^{n/2} = \frac{2^{2n}}{\pi} C_n^{(n-1)/2},$$

$$1 + C_n^1 + C_n^2 + \dots + C_n^n = 2^n,$$

$$1 - C_n^1 + C_n^2 - \dots + (-1)^n C_n^n = 0.$$

Here $(2n)!! = 2 \cdot 4 \cdot 6 \dots (2n)$, $(2n-1)!! = 1 \cdot 3 \cdot 5 \dots (2n-1)$, where $n = 1, 2, 3, \dots$ ($0!! = 1!! = 1$).

11.1-2. Pochhammer Symbol.

Definition:

$$(a)_n = a(a+1)\dots(a+n-1) = \frac{\Gamma(a+n)}{\Gamma(a)} = (-1)^n \frac{\Gamma(1-a)}{\Gamma(1-a-n)}.$$

Some properties ($k = 1, 2, \dots$):

$$\begin{aligned}(a)_0 &= 1, & (a)_{n+k} &= (a)_n(a+n)_k, & (n)_k &= \frac{(n+k-1)!}{(n-1)!}, \\ (a)_{-n} &= \frac{\Gamma(a-n)}{\Gamma(a)} = \frac{(-1)^n}{(1-a)_n}, & \text{where } a &\neq 1, \dots, n; \\ (1)_n &= n!, & (1/2)_n &= 2^{-2n} \frac{(2n)!}{n!}, & (3/2)_n &= 2^{-2n} \frac{(2n+1)!}{n!}, \\ (a+mk)_{nk} &= \frac{(a)_{mk+nk}}{(a)_{mk}}, & (a+n)_n &= \frac{(a)_{2n}}{(a)_n}, & (a+n)_k &= \frac{(a)_k(a+k)_n}{(a)_n}.\end{aligned}$$

11.1-3. Bernoulli Numbers.

The *Bernoulli numbers* are defined by the recurrence relation

$$B_0 = 1, \quad \sum_{k=0}^{n-1} C_n^k B_k = 0, \quad n = 2, 3, \dots$$

Numerical values:

$$B_0 = 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_4 = -\frac{1}{30}, \quad B_6 = \frac{1}{42}, \quad B_8 = -\frac{1}{30}, \quad B_{10} = \frac{5}{66}, \quad \dots, \\ B_{2m+1} = 0 \quad \text{for } m = 1, 2, \dots$$

All odd-numbered Bernoulli numbers but B_1 are zero; all even-numbered Bernoulli numbers have alternating signs.

The Bernoulli numbers are the values of Bernoulli polynomials at $x = 0$: $B_n = B_n(0)$.

Generating function:

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}, \quad |x| < 2\pi.$$

This relation may be regarded as a definition of the Bernoulli numbers.

The following expansions may be used to calculate the Bernoulli numbers:

$$\begin{aligned}\tan x &= \sum_{n=1}^{\infty} |B_{2n}| \frac{2^{2n}(2^{2n}-1)}{(2n)!} x^{2n}, & |x| &< \frac{\pi}{2}; \\ \cot x &= \sum_{n=0}^{\infty} (-1)^n B_{2n} \frac{2^{2n}}{(2n)!} x^{2n-1}, & |x| &< \pi.\end{aligned}$$

11.1-4. Euler Numbers.

The *Euler numbers* E_n are defined by the recurrence relation

$$\begin{aligned}\sum_{k=0}^n C_{2n}^{2k} E_{2k} &= 0 \quad (\text{even numbered}), \\ E_{2n+1} &= 0 \quad (\text{odd numbered}),\end{aligned}$$

where $n = 0, 1, \dots$

Numerical values:

$$E_0 = 1, \quad E_2 = -1, \quad E_4 = 5, \quad E_6 = -61, \quad E_8 = 1385, \quad E_{10} = -50251, \quad \dots, \\ E_{2n+1} = 0 \quad \text{for } n = 0, 1, \dots$$

All Euler numbers are integer, the odd-numbered Euler numbers are zero, and the even-numbered Euler numbers have alternating signs.

The Euler numbers are expressed via the values of Euler polynomials at $x = 1/2$: $E_n = 2^n E_n(1/2)$, where $n = 0, 1, \dots$

Generating function:

$$\frac{e^x}{e^{2x} + 1} = \sum_{n=0}^{\infty} E_n \frac{x^n}{n!}, \quad |x| < 2\pi.$$

This relation may be regarded as a definition of the Euler numbers.

Representation via a definite integral:

$$E_{2n} = (-1)^n 2^{2n+1} \int_0^{\infty} \frac{t^{2n} dt}{\cosh(\pi t)}.$$

11.2. Error Functions. Exponential and Logarithmic Integrals

11.2-1. Error Function and Complementary Error Function.

Definitions:

$$\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_0^x \exp(-t^2) dt \quad (\text{error function, also called probability integral}),$$

$$\operatorname{erfc} x = 1 - \operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_x^{\infty} \exp(-t^2) dt \quad (\text{complementary error function}).$$

Properties:

$$\operatorname{erf}(-x) = -\operatorname{erf} x; \quad \operatorname{erf}(0) = 0, \quad \operatorname{erf}(\infty) = 1; \quad \operatorname{erfc}(0) = 1, \quad \operatorname{erfc}(\infty) = 0.$$

Expansion of $\operatorname{erf} x$ into series in powers of x as $x \rightarrow 0$:

$$\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{k! (2k+1)} = \frac{2}{\sqrt{\pi}} \exp(-x^2) \sum_{k=0}^{\infty} \frac{2^k x^{2k+1}}{(2k+1)!!}.$$

Asymptotic expansion of $\operatorname{erfc} x$ as $x \rightarrow \infty$:

$$\operatorname{erfc} x = \frac{1}{\sqrt{\pi}} \exp(-x^2) \left[\sum_{m=0}^{M-1} (-1)^m \frac{\left(\frac{1}{2}\right)_m}{x^{2m+1}} + O(|x|^{-2M-1}) \right], \quad M = 1, 2, \dots$$

Integral:

$$\int_0^x \operatorname{erf} t dt = x \operatorname{erf} x - \frac{1}{2} + \frac{1}{2} \exp(-x^2).$$

11.2-2. Exponential Integral.

Definition:

$$\text{Ei}(x) = \int_{-\infty}^x \frac{e^t}{t} dt = - \int_{-x}^{\infty} \frac{e^{-t}}{t} dt \quad \text{for } x < 0,$$

$$\text{Ei}(x) = \lim_{\varepsilon \rightarrow +0} \left(\int_{-\infty}^{-\varepsilon} \frac{e^t}{t} dt + \int_{\varepsilon}^x \frac{e^t}{t} dt \right) \quad \text{for } x > 0.$$

Other integral representations:

$$\text{Ei}(-x) = -e^{-x} \int_0^{\infty} \frac{x \sin t + t \cos t}{x^2 + t^2} dt \quad \text{for } x > 0,$$

$$\text{Ei}(-x) = e^{-x} \int_0^{\infty} \frac{x \sin t - t \cos t}{x^2 + t^2} dt \quad \text{for } x < 0,$$

$$\text{Ei}(-x) = -x \int_1^{\infty} e^{-xt} \ln t dt \quad \text{for } x > 0,$$

$$\text{Ei}(x) = \mathcal{C} + \ln x + \int_0^x \frac{e^t - 1}{t} dt \quad \text{for } x > 0,$$

where $\mathcal{C} = 0.5772\dots$ is the Euler constant.Expansion into series in powers of x as $x \rightarrow 0$:

$$\text{Ei}(x) = \begin{cases} \mathcal{C} + \ln(-x) + \sum_{k=1}^{\infty} \frac{x^k}{k! k} & \text{if } x < 0, \\ \mathcal{C} + \ln x + \sum_{k=1}^{\infty} \frac{x^k}{k! k} & \text{if } x > 0. \end{cases}$$

Asymptotic expansion as $x \rightarrow \infty$:

$$\text{Ei}(-x) = e^{-x} \sum_{k=1}^n (-1)^k \frac{(k-1)!}{x^k} + R_n, \quad R_n < \frac{n!}{x^n}.$$

11.2-3. Logarithmic Integral.

Definition:

$$\text{li}(x) = \begin{cases} \int_0^x \frac{dt}{\ln t} & \text{if } 0 < x < 1, \\ \lim_{\varepsilon \rightarrow +0} \left(\int_0^{1-\varepsilon} \frac{dt}{\ln t} + \int_{1+\varepsilon}^x \frac{dt}{\ln t} \right) & \text{if } x > 1. \end{cases}$$

For small x ,

$$\text{li}(x) \approx \frac{x}{\ln(1/x)}.$$

For large x ,

$$\text{li}(x) \approx \frac{x}{\ln x}.$$

Asymptotic expansion as $x \rightarrow 1$:

$$\text{li}(x) = \mathcal{C} + \ln |\ln x| + \sum_{k=1}^{\infty} \frac{\ln^k x}{k! k}.$$

Relation to the exponential integral:

$$\text{li } x = \text{Ei}(\ln x), \quad x < 1;$$

$$\text{li}(e^x) = \text{Ei}(x), \quad x < 0.$$

11.3. Sine Integral and Cosine Integral. Fresnel Integrals

11.3-1. Sine Integral.

Definition:

$$\text{Si}(x) = \int_0^x \frac{\sin t}{t} dt, \quad \text{si}(x) = - \int_x^\infty \frac{\sin t}{t} dt = \text{Si}(x) - \frac{\pi}{2}.$$

Specific values:

$$\text{Si}(0) = 0, \quad \text{Si}(\infty) = \frac{\pi}{2}, \quad \text{si}(\infty) = 0.$$

Properties:

$$\text{Si}(-x) = -\text{Si}(x), \quad \text{si}(x) + \text{si}(-x) = -\pi, \quad \lim_{x \rightarrow -\infty} \text{si}(x) = -\pi.$$

Expansion into series in powers of x as $x \rightarrow 0$:

$$\text{Si}(x) = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} x^{2k-1}}{(2k-1)(2k-1)!}.$$

Asymptotic expansion as $x \rightarrow \infty$:

$$\text{si}(x) = -\cos x \left[\sum_{m=0}^{M-1} \frac{(-1)^m (2m)!}{x^{2m+1}} + O(|x|^{-2M-1}) \right] + \sin x \left[\sum_{m=1}^{N-1} \frac{(-1)^m (2m-1)!}{x^{2m}} + O(|x|^{-2N}) \right],$$

where $M, N = 1, 2, \dots$

11.3-2. Cosine Integral.

Definition:

$$\text{ci}(x) = - \int_x^\infty \frac{\cos t}{t} dt = \mathcal{C} + \ln x + \int_0^x \frac{\cos t - 1}{t} dt,$$

where $\mathcal{C} = 0.5772\dots$ is the Euler constant.

Expansion into series in powers of x as $x \rightarrow 0$:

$$\text{ci}(x) = \mathcal{C} + \ln x + \sum_{k=1}^{\infty} \frac{(-1)^k x^{2k}}{2k(2k)!}.$$

Asymptotic expansion as $x \rightarrow \infty$:

$$\text{ci}(x) = \cos x \left[\sum_{m=1}^{M-1} \frac{(-1)^m (2m-1)!}{x^{2m}} + O(|x|^{-2M}) \right] + \sin x \left[\sum_{m=0}^{N-1} \frac{(-1)^m (2m)!}{x^{2m+1}} + O(|x|^{-2N-1}) \right],$$

where $M, N = 1, 2, \dots$

11.3-3. Fresnel Integrals and Generalized Fresnel Integrals.
--

Fresnel sine and cosine integrals:

$$S(x) = \frac{1}{\sqrt{2\pi}} \int_0^x \frac{\sin t}{\sqrt{t}} dt = \sqrt{\frac{2}{\pi}} \int_0^{\sqrt{x}} \sin t^2 dt,$$

$$C(x) = \frac{1}{\sqrt{2\pi}} \int_0^x \frac{\cos t}{\sqrt{t}} dt = \sqrt{\frac{2}{\pi}} \int_0^{\sqrt{x}} \cos t^2 dt.$$

Expansion into series in powers of x as $x \rightarrow 0$:

$$S(x) = \sqrt{\frac{2}{\pi}} x \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{(4k+3)(2k+1)!},$$

$$C(x) = \sqrt{\frac{2}{\pi}} x \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(4k+1)(2k)!}.$$

Asymptotic expansion as $x \rightarrow \infty$:

$$S(x) = \frac{1}{2} - \frac{\cos x}{\sqrt{2\pi x}} P(x) - \frac{\sin x}{\sqrt{2\pi x}} Q(x),$$

$$C(x) = \frac{1}{2} + \frac{\sin x}{\sqrt{2\pi x}} P(x) - \frac{\cos x}{\sqrt{2\pi x}} Q(x),$$

$$P(x) = 1 - \frac{1 \times 3}{(2x)^2} + \frac{1 \times 3 \times 5 \times 7}{(2x)^4} - \dots, \quad Q(x) = \frac{1}{2x} - \frac{1 \times 3 \times 5}{(2x)^3} + \dots.$$

Generalized Fresnel sine and cosine integrals:

$$S(x, \nu) = \int_x^{\infty} t^{\nu-1} \sin t dt, \quad \operatorname{Re} \nu < 1;$$

$$C(x, \nu) = \int_x^{\infty} t^{\nu-1} \cos t dt, \quad \operatorname{Re} \nu < 1.$$

11.4. Gamma Function, Psi Function, and Beta Function

11.4-1. Gamma Function.

The *gamma function*, $\Gamma(z)$, is an analytic function of the complex argument z everywhere except for the points $z = 0, -1, -2, \dots$

For $\operatorname{Re} z > 0$,

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt.$$

For $-(n+1) < \operatorname{Re} z < -n$, where $n = 0, 1, 2, \dots$,

$$\Gamma(z) = \int_0^{\infty} \left[e^{-t} - \sum_{m=0}^n \frac{(-1)^m}{m!} t^m \right] t^{z-1} dt.$$

Simplest properties:

$$\Gamma(z+1) = z\Gamma(z), \quad \Gamma(n+1) = n!, \quad \Gamma(1) = \Gamma(2) = 1.$$

Fractional values of the argument:

$$\begin{aligned}\Gamma\left(\frac{1}{2}\right) &= \sqrt{\pi}, & \Gamma\left(n + \frac{1}{2}\right) &= \frac{\sqrt{\pi}}{2^n} (2n-1)!!, \\ \Gamma\left(-\frac{1}{2}\right) &= -2\sqrt{\pi}, & \Gamma\left(\frac{1}{2} - n\right) &= (-1)^n \frac{2^n \sqrt{\pi}}{(2n-1)!!}.\end{aligned}$$

Euler formula

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n! n^z}{z(z+1)\dots(z+n)} \quad (z \neq 0, -1, -2, \dots).$$

Symmetry formulas:

$$\begin{aligned}\Gamma(z)\Gamma(-z) &= -\frac{\pi}{z \sin(\pi z)}, & \Gamma(z)\Gamma(1-z) &= \frac{\pi}{\sin(\pi z)}, \\ \Gamma\left(\frac{1}{2} + z\right)\Gamma\left(\frac{1}{2} - z\right) &= \frac{\pi}{\cos(\pi z)}.\end{aligned}$$

Multiple argument formulas:

$$\begin{aligned}\Gamma(2z) &= \frac{2^{2z-1}}{\sqrt{\pi}} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right), \\ \Gamma(3z) &= \frac{3^{3z-1/2}}{2\pi} \Gamma(z) \Gamma\left(z + \frac{1}{3}\right) \Gamma\left(z + \frac{2}{3}\right), \\ \Gamma(nz) &= (2\pi)^{(1-n)/2} n^{nz-1/2} \prod_{k=0}^{n-1} \Gamma\left(z + \frac{k}{n}\right).\end{aligned}$$

Asymptotic expansion (*Stirling formula*):

$$\Gamma(z) = \sqrt{2\pi} e^{-z} z^{z-1/2} \left[1 + \frac{1}{12} z^{-1} + \frac{1}{288} z^{-2} + O(z^{-3})\right] \quad (|\arg z| < \pi).$$

11.4-2. Psi Function (Digamma Function).

Definition:

$$\psi(z) = \frac{d \ln \Gamma(z)}{dz} = \frac{\Gamma'_z(z)}{\Gamma(z)}.$$

The psi function is the logarithmic derivative of the gamma function and is also called the *digamma function*.

Integral representations ($\operatorname{Re} z > 0$):

$$\begin{aligned}\psi(z) &= \int_0^\infty [e^{-t} - (1+t)^{-z}] t^{-1} dt, \\ \psi(z) &= \ln z + \int_0^\infty [t^{-1} - (1-e^{-t})^{-1}] e^{-tz} dt, \\ \psi(z) &= -C + \int_0^1 \frac{1-t^{z-1}}{1-t} dt,\end{aligned}$$

where $C = -\psi(1) = 0.5772\dots$ is the Euler constant.

Values for integer argument:

$$\psi(1) = -\mathcal{C}, \quad \psi(n) = -\mathcal{C} + \sum_{k=1}^{n-1} k^{-1} \quad (n = 2, 3, \dots).$$

Functional relations:

$$\begin{aligned} \psi(z) - \psi(1+z) &= -\frac{1}{z}, \\ \psi(z) - \psi(1-z) &= -\pi \cot(\pi z), \\ \psi(z) - \psi(-z) &= -\pi \cot(\pi z) - \frac{1}{z}, \\ \psi\left(\frac{1}{2} + z\right) - \psi\left(\frac{1}{2} - z\right) &= \pi \tan(\pi z), \\ \psi(mz) &= \ln m + \frac{1}{m} \sum_{k=0}^{m-1} \psi\left(z + \frac{k}{m}\right). \end{aligned}$$

Asymptotic expansion as $z \rightarrow \infty$ ($|\arg z| < \pi$):

$$\psi(z) = \ln z - \frac{1}{2z} - \frac{1}{12z^2} + \frac{1}{120z^4} - \frac{1}{252z^6} + \dots = \ln z - \frac{1}{2z} - \sum_{n=1}^{\infty} \frac{B_{2n}}{2nz^{2n}},$$

where the B_{2n} are Bernoulli numbers.

11.4-3. Beta Function.

Definition:

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt,$$

where $\operatorname{Re} x > 0$ and $\operatorname{Re} y > 0$.

Relationship with the gamma function:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}.$$

Some properties:

$$\begin{aligned} B(x, y) &= B(y, x); \\ B(x, y+1) &= \frac{y}{x} B(x+1, y) = \frac{y}{x+y} B(x, y); \\ B(x, 1-x) &= \frac{\pi}{\sin(\pi x)}, \quad 0 < x < 1; \\ \frac{1}{B(n, m)} &= mC_{n+m-1}^{n-1} = nC_{n+m-1}^{m-1}, \end{aligned}$$

where n and m are positive integers.

11.5. Incomplete Gamma and Beta Functions

11.5-1. Incomplete Gamma Function.

Definitions:

$$\begin{aligned} \gamma(\alpha, x) &= \int_0^x e^{-t} t^{\alpha-1} dt, \quad \operatorname{Re} \alpha > 0, \\ \Gamma(\alpha, x) &= \int_x^{\infty} e^{-t} t^{\alpha-1} dt = \Gamma(\alpha) - \gamma(\alpha, x). \end{aligned}$$

Recurrence formulas:

$$\begin{aligned}\gamma(\alpha + 1, x) &= \alpha\gamma(\alpha, x) - x^\alpha e^{-x}, \\ \gamma(\alpha + 1, x) &= (x + \alpha)\gamma(\alpha, x) + (1 - \alpha)x\gamma(\alpha - 1, x), \\ \Gamma(\alpha + 1, x) &= \alpha\Gamma(\alpha, x) + x^\alpha e^{-x}.\end{aligned}$$

Special cases:

$$\begin{aligned}\gamma(n + 1, x) &= n! \left[1 - e^{-x} \left(\sum_{k=0}^n \frac{x^k}{k!} \right) \right], & n = 0, 1, \dots; \\ \Gamma(n + 1, x) &= n! e^{-x} \sum_{k=0}^n \frac{x^k}{k!}, & n = 0, 1, \dots; \\ \Gamma(-n, x) &= \frac{(-1)^n}{n!} \left[\Gamma(0, x) - e^{-x} \sum_{k=0}^{n-1} (-1)^k \frac{k!}{x^{k+1}} \right], & n = 1, 2, \dots\end{aligned}$$

Asymptotic expansions as $x \rightarrow 0$:

$$\begin{aligned}\gamma(\alpha, x) &= \sum_{n=0}^{\infty} \frac{(-1)^n x^{\alpha+n}}{n! (\alpha + n)}, \\ \Gamma(\alpha, x) &= \Gamma(\alpha) - \sum_{n=0}^{\infty} \frac{(-1)^n x^{\alpha+n}}{n! (\alpha + n)}.\end{aligned}$$

Asymptotic expansions as $x \rightarrow \infty$:

$$\begin{aligned}\gamma(\alpha, x) &= \Gamma(\alpha) - x^{\alpha-1} e^{-x} \left[\sum_{m=0}^{M-1} \frac{(1-\alpha)_m}{(-x)^m} + O(|x|^{-M}) \right], \\ \Gamma(\alpha, x) &= x^{\alpha-1} e^{-x} \left[\sum_{m=0}^{M-1} \frac{(1-\alpha)_m}{(-x)^m} + O(|x|^{-M}) \right] \quad \left(-\frac{3}{2}\pi < \arg x < \frac{3}{2}\pi \right).\end{aligned}$$

Asymptotic formulas as $\alpha \rightarrow \infty$:

$$\begin{aligned}\gamma(x, \alpha) &= \Gamma(\alpha) \left[\Phi(2\sqrt{x} - \sqrt{\alpha-1}) + O\left(\frac{1}{\sqrt{\alpha}}\right) \right], & \Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x \exp\left(-\frac{1}{2}t^2\right) dt; \\ \gamma(x, \alpha) &= \Gamma(\alpha) \left[\Phi(3\sqrt{\alpha}z) + O\left(\frac{1}{\alpha}\right) \right], & z = \left(\frac{x}{\alpha}\right)^{1/3} - 1 + \frac{1}{9\alpha}.\end{aligned}$$

Representation of the error function, complementary error function, and exponential integral in terms of the gamma functions:

$$\operatorname{erf} x = \frac{1}{\sqrt{\pi}} \gamma\left(\frac{1}{2}, x^2\right), \quad \operatorname{erfc} x = \frac{1}{\sqrt{\pi}} \Gamma\left(\frac{1}{2}, x^2\right), \quad \operatorname{Ei}(-x) = -\Gamma(0, x).$$

11.5-2. Incomplete Beta Function.

Definitions:

$$B_x(a, b) = \int_0^x t^{a-1} (1-t)^{b-1} dt, \quad I_x(a, b) = \frac{B_x(a, b)}{B(a, b)},$$

where $\operatorname{Re} a > 0$ and $\operatorname{Re} b > 0$, and $B(a, b) = B_1(a, b)$ is the beta function.

Symmetry property:

$$I_x(a, b) + I_{1-x}(b, a) = 1.$$

Recurrence formulas:

$$\begin{aligned} I_x(a, b) &= xI_x(a-1, b) + (1-x)I_x(a, b-1), \\ (a+b)I_x(a, b) &= aI_x(a+1, b) + bI_x(a, b+1), \\ (a+b-ax)I_x(a, b) &= a(1-x)I_x(a+1, b-1) + bI_x(a, b+1). \end{aligned}$$

11.6. Bessel Functions (Cylindrical Functions)

11.6-1. Definitions and Basic Formulas.

The *Bessel function of the first kind*, $J_\nu(x)$, and the *Bessel function of the second kind*, $Y_\nu(x)$ (also called the *Neumann function*), are solutions of the *Bessel equation*

$$x^2 y''_{xx} + xy'_x + (x^2 - \nu^2)y = 0$$

and are defined by the formulas

$$J_\nu(x) = \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{\nu+2k}}{k! \Gamma(\nu+k+1)}, \quad Y_\nu(x) = \frac{J_\nu(x) \cos \pi\nu - J_{-\nu}(x)}{\sin \pi\nu}. \quad (1)$$

The formula for $Y_\nu(x)$ is valid for $\nu \neq 0, \pm 1, \pm 2, \dots$ (the cases $\nu \neq 0, \pm 1, \pm 2, \dots$ are discussed in what follows).

The general solution of the Bessel equation has the form $Z_\nu(x) = C_1 J_\nu(x) + C_2 Y_\nu(x)$ and is called the *cylinder function*.

Some formulas:

$$\begin{aligned} 2\nu Z_\nu(x) &= x[Z_{\nu-1}(x) + Z_{\nu+1}(x)], \\ \frac{d}{dx} Z_\nu(x) &= \frac{1}{2}[Z_{\nu-1}(x) - Z_{\nu+1}(x)] = \pm \left[\frac{\nu}{x} Z_\nu(x) - Z_{\nu \pm 1}(x) \right], \\ \frac{d}{dx} [x^\nu Z_\nu(x)] &= x^\nu Z_{\nu-1}(x), \quad \frac{d}{dx} [x^{-\nu} Z_\nu(x)] = -x^{-\nu} Z_{\nu+1}(x), \\ \left(\frac{1}{x} \frac{d}{dx} \right)^n [x^\nu J_\nu(x)] &= x^{\nu-n} J_{\nu-n}(x), \quad \left(\frac{1}{x} \frac{d}{dx} \right)^n [x^{-\nu} J_\nu(x)] = (-1)^n x^{-\nu-n} J_{\nu+n}(x), \\ J_{-n}(x) &= (-1)^n J_n(x), \quad Y_{-n}(x) = (-1)^n Y_n(x), \quad n = 0, 1, 2, \dots \end{aligned}$$

Bessel functions for $\nu = \pm n \pm \frac{1}{2}$ ($n = 0, 1, 2, \dots$):

$$\begin{aligned} J_{1/2}(x) &= \sqrt{\frac{2}{\pi x}} \sin x, & J_{-1/2}(x) &= \sqrt{\frac{2}{\pi x}} \cos x, \\ J_{3/2}(x) &= \sqrt{\frac{2}{\pi x}} \left(\frac{1}{x} \sin x - \cos x \right), & J_{-3/2}(x) &= \sqrt{\frac{2}{\pi x}} \left(-\frac{1}{x} \cos x - \sin x \right), \end{aligned}$$

$$\begin{aligned}
 J_{n+1/2}(x) &= \sqrt{\frac{2}{\pi x}} \left[\sin\left(x - \frac{n\pi}{2}\right) \sum_{k=0}^{[n/2]} \frac{(-1)^k (n+2k)!}{(2k)! (n-2k)! (2x)^{2k}} \right. \\
 &\quad \left. + \cos\left(x - \frac{n\pi}{2}\right) \sum_{k=0}^{[(n-1)/2]} \frac{(-1)^k (n+2k+1)!}{(2k+1)! (n-2k-1)! (2x)^{2k+1}} \right], \\
 J_{-n-1/2}(x) &= \sqrt{\frac{2}{\pi x}} \left[\cos\left(x + \frac{n\pi}{2}\right) \sum_{k=0}^{[n/2]} \frac{(-1)^k (n+2k)!}{(2k)! (n-2k)! (2x)^{2k}} \right. \\
 &\quad \left. - \sin\left(x + \frac{n\pi}{2}\right) \sum_{k=0}^{[(n-1)/2]} \frac{(-1)^k (n+2k+1)!}{(2k+1)! (n-2k-1)! (2x)^{2k+1}} \right], \\
 Y_{1/2}(x) &= -\sqrt{\frac{2}{\pi x}} \cos x, & Y_{-1/2}(x) &= \sqrt{\frac{2}{\pi x}} \sin x, \\
 Y_{n+1/2}(x) &= (-1)^{n+1} J_{-n-1/2}(x), & Y_{-n-1/2}(x) &= (-1)^n J_{n+1/2}(x),
 \end{aligned}$$

where $[A]$ is the integer part of the number A .

Let $\nu = n$ be an arbitrary integer. The relations

$$J_{-n}(x) = (-1)^n J_n(x), \quad Y_{-n}(x) = (-1)^n Y_n(x)$$

are valid. The function $J_n(x)$ is given by the first formula in (1) with $\nu = n$, and $Y_n(x)$ can be obtained from the second formula in (1) by proceeding to the limit $\nu \rightarrow n$. For nonnegative n , $Y_n(x)$ can be represented in the form

$$Y_n(x) = \frac{2}{\pi} J_n(x) \ln \frac{x}{2} - \frac{1}{\pi} \sum_{k=0}^{n-1} \frac{(n-k-1)!}{k!} \left(\frac{2}{x}\right)^{n-2k} - \frac{1}{\pi} \sum_{k=0}^{\infty} (-1)^k \left(\frac{x}{2}\right)^{n+2k} \frac{\psi(k+1) + \psi(n+k+1)}{k!(n+k)!},$$

where $\psi(1) = -\mathcal{C}$, $\psi(n) = -\mathcal{C} + \sum_{k=1}^{n-1} k^{-1}$, $\mathcal{C} = 0.5772\dots$ is the Euler constant, and $\psi(x) = [\ln \Gamma(x)]'_x$ is the logarithmic derivative of the gamma function, also known as the digamma function.

Wronskians and similar formulas:

$$\begin{aligned}
 W(J_\nu, J_{-\nu}) &= -\frac{2}{\pi x} \sin(\pi\nu), & W(J_\nu, Y_\nu) &= \frac{2}{\pi x}, \\
 J_\nu(x)J_{-\nu+1}(x) + J_{-\nu}(x)J_{\nu-1}(x) &= \frac{2 \sin(\pi\nu)}{\pi x}, & J_\nu(x)Y_{\nu+1}(x) - J_{\nu+1}(x)Y_\nu(x) &= -\frac{2}{\pi x}.
 \end{aligned}$$

Here the notation $W(f, g) = fg'_x - f'_x g$ is used.

11.6-2. Integral Representations and Asymptotic Expansions.

The functions $J_\nu(x)$ and $Y_\nu(x)$ can be represented in the form of definite integrals (for $x > 0$):

$$\begin{aligned}
 \pi J_\nu(x) &= \int_0^\pi \cos(x \sin \theta - \nu\theta) d\theta - \sin \pi\nu \int_0^\infty \exp(-x \sinh t - \nu t) dt, \\
 \pi Y_\nu(x) &= \int_0^\pi \sin(x \sin \theta - \nu\theta) d\theta - \int_0^\infty (e^{\nu t} + e^{-\nu t} \cos \pi\nu) e^{-x \sinh t} dt.
 \end{aligned}$$

For $|\nu| < \frac{1}{2}$, $x > 0$,

$$J_\nu(x) = \frac{2^{1+\nu} x^{-\nu}}{\pi^{1/2} \Gamma(\frac{1}{2} - \nu)} \int_1^\infty \frac{\sin(xt) dt}{(t^2 - 1)^{\nu+1/2}},$$

$$Y_\nu(x) = -\frac{2^{1+\nu} x^{-\nu}}{\pi^{1/2} \Gamma(\frac{1}{2} - \nu)} \int_1^\infty \frac{\cos(xt) dt}{(t^2 - 1)^{\nu+1/2}}.$$

For $\nu > -\frac{1}{2}$,

$$J_\nu(x) = \frac{2(x/2)^\nu}{\pi^{1/2} \Gamma(\frac{1}{2} + \nu)} \int_0^{\pi/2} \cos(x \cos t) \sin^{2\nu} t dt \quad (\text{Poisson's formula}).$$

For $\nu = 0$, $x > 0$,

$$J_0(x) = \frac{2}{\pi} \int_0^\infty \sin(x \cosh t) dt, \quad Y_0(x) = -\frac{2}{\pi} \int_0^\infty \cos(x \cosh t) dt.$$

For integer $\nu = n = 0, 1, 2, \dots$,

$$J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(nt - x \sin t) dt \quad (\text{Bessel's formula}),$$

$$J_{2n}(x) = \frac{2}{\pi} \int_0^{\pi/2} \cos(x \sin t) \cos(2nt) dt,$$

$$J_{2n+1}(x) = \frac{2}{\pi} \int_0^{\pi/2} \sin(x \sin t) \sin[(2n+1)t] dt.$$

Asymptotic expansions as $|x| \rightarrow \infty$:

$$J_\nu(x) = \sqrt{\frac{2}{\pi x}} \left\{ \cos\left(\frac{4x - 2\nu\pi - \pi}{4}\right) \left[\sum_{m=0}^{M-1} (-1)^m (\nu, 2m) (2x)^{-2m} + O(|x|^{-2M}) \right] \right. \\ \left. - \sin\left(\frac{4x - 2\nu\pi - \pi}{4}\right) \left[\sum_{m=0}^{M-1} (-1)^m (\nu, 2m+1) (2x)^{-2m-1} + O(|x|^{-2M-1}) \right] \right\},$$

$$Y_\nu(x) = \sqrt{\frac{2}{\pi x}} \left\{ \sin\left(\frac{4x - 2\nu\pi - \pi}{4}\right) \left[\sum_{m=0}^{M-1} (-1)^m (\nu, 2m) (2x)^{-2m} + O(|x|^{-2M}) \right] \right. \\ \left. + \cos\left(\frac{4x - 2\nu\pi - \pi}{4}\right) \left[\sum_{m=0}^{M-1} (-1)^m (\nu, 2m+1) (2x)^{-2m-1} + O(|x|^{-2M-1}) \right] \right\},$$

where $(\nu, m) = \frac{1}{2^{2m} m!} (4\nu^2 - 1)(4\nu^2 - 3^2) \dots [4\nu^2 - (2m-1)^2] = \frac{\Gamma(\frac{1}{2} + \nu + m)}{m! \Gamma(\frac{1}{2} + \nu - m)}$.

For nonnegative integer n and large x ,

$$\sqrt{\pi x} J_{2n}(x) = (-1)^n (\cos x + \sin x) + O(x^{-2}),$$

$$\sqrt{\pi x} J_{2n+1}(x) = (-1)^{n+1} (\cos x - \sin x) + O(x^{-2}).$$

Asymptotic for large ν ($\nu \rightarrow \infty$):

$$J_\nu(x) \simeq \frac{1}{\sqrt{2\pi\nu}} \left(\frac{ex}{2\nu}\right)^\nu, \quad Y_\nu(x) \simeq -\sqrt{\frac{2}{\pi\nu}} \left(\frac{ex}{2\nu}\right)^{-\nu},$$

where x is fixed, and

$$J_\nu(\nu) \simeq \frac{2^{1/3}}{3^{2/3}\Gamma(2/3)} \frac{1}{\nu^{1/3}}, \quad Y_\nu(\nu) \simeq -\frac{2^{1/3}}{3^{1/6}\Gamma(2/3)} \frac{1}{\nu^{1/3}}.$$

Integrals with Bessel functions:

$$\int_0^x x^\lambda J_\nu(x) dx = \frac{x^{\lambda+\nu+1}}{2^\nu(\lambda+\nu+1)\Gamma(\nu+1)} F\left(\frac{\lambda+\nu+1}{2}, \frac{\lambda+\nu+3}{2}, \nu+1; -\frac{x^2}{4}\right), \quad \operatorname{Re}(\lambda+\nu) > -1,$$

where $F(a, b, c; x)$ is the hypergeometric series (see Supplement 11.10.1),

$$\begin{aligned} \int_0^x x^\lambda Y_\nu(x) dx &= -\frac{\cos(\nu\pi)\Gamma(-\nu)}{2^\nu\pi(\lambda+\nu+1)} x^{\lambda+\nu+1} F\left(\frac{\lambda+\nu+1}{2}, \nu+1, \frac{\lambda+\nu+3}{2}; -\frac{x^2}{4}\right) \\ &\quad - \frac{2^\nu\Gamma(\nu)}{\lambda-\nu+1} x^{\lambda-\nu+1} F\left(\frac{\lambda-\nu+1}{2}, 1-\nu, \frac{\lambda-\nu+3}{2}; -\frac{x^2}{4}\right), \quad \operatorname{Re} \lambda > |\operatorname{Re} \nu| - 1. \end{aligned}$$

11.6-3. Zeros of Bessel Functions.

Each of the functions $J_\nu(x)$ and $Y_\nu(x)$ has infinitely many real zeros (for real ν). All zeros are simple, except possibly for the point $x = 0$.

The zeros γ_m of $J_0(x)$, i.e., the roots of the equation $J_0(\gamma_m) = 0$, are approximately given by

$$\gamma_m = 2.4 + 3.13(m-1) \quad (m = 1, 2, \dots),$$

with a maximum error of 0.2%.

11.6-4. Orthogonality Properties of Bessel Functions.

1°. Let $\mu = \mu_m$ be positive roots of the Bessel function $J_\nu(\mu)$, where $\nu > -1$ and $m = 1, 2, 3, \dots$. Then the set of functions $J_\nu(\mu_m r/a)$ is orthogonal on the interval $0 \leq r \leq a$ with weight r :

$$\int_0^a J_\nu\left(\frac{\mu_m r}{a}\right) J_\nu\left(\frac{\mu_k r}{a}\right) r dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{1}{2}a^2 [J'_\nu(\mu_m)]^2 = \frac{1}{2}a^2 J_{\nu+1}^2(\mu_m) & \text{if } m = k. \end{cases}$$

2°. Let $\mu = \mu_m$ be positive zeros of the Bessel function derivative $J'_\nu(\mu)$, where $\nu > -1$ and $m = 1, 2, 3, \dots$. Then the set of functions $J_\nu(\mu_m r/a)$ is orthogonal on the interval $0 \leq r \leq a$ with weight r :

$$\int_0^a J_\nu\left(\frac{\mu_m r}{a}\right) J_\nu\left(\frac{\mu_k r}{a}\right) r dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{1}{2}a^2 \left(1 - \frac{\nu^2}{\mu_m^2}\right) J_\nu^2(\mu_m) & \text{if } m = k. \end{cases}$$

3°. Let $\mu = \mu_m$ be positive roots of the transcendental equation $\mu J'_\nu(\mu) + sJ_\nu(\mu) = 0$, where $\nu > -1$ and $m = 1, 2, 3, \dots$. Then the set of functions $J_\nu(\mu_m r/a)$ is orthogonal on the interval $0 \leq r \leq a$ with weight r :

$$\int_0^a J_\nu\left(\frac{\mu_m r}{a}\right) J_\nu\left(\frac{\mu_k r}{a}\right) r dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{1}{2}a^2 \left(1 + \frac{s^2 - \nu^2}{\mu_m^2}\right) J_\nu^2(\mu_m) & \text{if } m = k. \end{cases}$$

4°. Let $\mu = \mu_m$ be positive roots of the transcendental equation

$$J_\nu(\lambda_m b)Y_\nu(\lambda_m a) - J_\nu(\lambda_m a)Y_\nu(\lambda_m b) = 0 \quad (\nu > -1, m = 1, 2, 3, \dots).$$

Then the set of functions

$$Z_\nu(\lambda_m r) = J_\nu(\lambda_m r)Y_\nu(\lambda_m a) - J_\nu(\lambda_m a)Y_\nu(\lambda_m r), \quad m = 1, 2, 3, \dots,$$

satisfying the conditions $Z_\nu(\lambda_m a) = Z_\nu(\lambda_m b) = 0$ is orthogonal on the interval $a \leq r \leq b$ with weight r :

$$\int_a^b Z_\nu(\lambda_m r)Z_\nu(\lambda_k r)r \, dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{2}{\pi^2 \lambda_m^2} \frac{J_\nu^2(\lambda_m a) - J_\nu^2(\lambda_m b)}{J_\nu^2(\lambda_m b)} & \text{if } m = k. \end{cases}$$

5°. Let $\mu = \mu_m$ be positive roots of the transcendental equation

$$J'_\nu(\lambda_m b)Y'_\nu(\lambda_m a) - J'_\nu(\lambda_m a)Y'_\nu(\lambda_m b) = 0 \quad (\nu > -1, m = 1, 2, 3, \dots).$$

Then the set of functions

$$Z_\nu(\lambda_m r) = J_\nu(\lambda_m r)Y'_\nu(\lambda_m a) - J'_\nu(\lambda_m a)Y_\nu(\lambda_m r), \quad m = 1, 2, 3, \dots,$$

satisfying the conditions $Z'_\nu(\lambda_m a) = Z'_\nu(\lambda_m b) = 0$ is orthogonal on the interval $a \leq r \leq b$ with weight r :

$$\int_a^b Z_\nu(\lambda_m r)Z_\nu(\lambda_k r)r \, dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{2}{\pi^2 \lambda_m^2} \left[\left(1 - \frac{\nu^2}{b^2 \lambda_m^2}\right) \frac{[J'_\nu(\lambda_m a)]^2}{[J'_\nu(\lambda_m b)]^2} - \left(1 - \frac{\nu^2}{a^2 \lambda_m^2}\right) \right] & \text{if } m = k. \end{cases}$$

11.6-5. Hankel Functions (Bessel Functions of the Third Kind).

The *Hankel functions of the first kind and the second kind* are related to Bessel functions by

$$\begin{aligned} H_\nu^{(1)}(z) &= J_\nu(z) + iY_\nu(z), \\ H_\nu^{(2)}(z) &= J_\nu(z) - iY_\nu(z), \end{aligned}$$

where $i^2 = -1$.

Asymptotics for $z \rightarrow 0$:

$$\begin{aligned} H_0^{(1)}(z) &\simeq \frac{2i}{\pi} \ln z, & H_\nu^{(1)}(z) &\simeq -\frac{i}{\pi} \frac{\Gamma(\nu)}{(z/2)^\nu} & (\operatorname{Re} \nu > 0), \\ H_0^{(2)}(z) &\simeq -\frac{2i}{\pi} \ln z, & H_\nu^{(2)}(z) &\simeq \frac{i}{\pi} \frac{\Gamma(\nu)}{(z/2)^\nu} & (\operatorname{Re} \nu > 0). \end{aligned}$$

Asymptotics for $|z| \rightarrow \infty$:

$$\begin{aligned} H_\nu^{(1)}(z) &\simeq \sqrt{\frac{2}{\pi z}} \exp\left[i\left(z - \frac{1}{2}\pi\nu - \frac{1}{4}\pi\right)\right] & (-\pi < \arg z < 2\pi), \\ H_\nu^{(2)}(z) &\simeq \sqrt{\frac{2}{\pi z}} \exp\left[-i\left(z - \frac{1}{2}\pi\nu - \frac{1}{4}\pi\right)\right] & (-2\pi < \arg z < \pi). \end{aligned}$$

11.7. Modified Bessel Functions

11.7-1. Definitions. Basic Formulas.

The *modified Bessel functions of the first kind*, $I_\nu(x)$, and the *modified Bessel functions of the second kind*, $K_\nu(x)$ (also called the *MacDonald function*), of order ν are solutions of the *modified Bessel equation*

$$x^2 y''_{xx} + x y'_x - (x^2 + \nu^2)y = 0$$

and are defined by the formulas

$$I_\nu(x) = \sum_{k=0}^{\infty} \frac{(x/2)^{2k+\nu}}{k! \Gamma(\nu + k + 1)}, \quad K_\nu(x) = \frac{\pi}{2} \frac{I_{-\nu}(x) - I_\nu(x)}{\sin(\pi\nu)}$$

(see below for $K_\nu(x)$ with $\nu = 0, 1, 2, \dots$).

The modified Bessel functions possess the properties

$$\begin{aligned} K_{-\nu}(x) &= K_\nu(x); & I_{-n}(x) &= (-1)^n I_n(x), \quad n = 0, 1, 2, \dots \\ 2\nu I_\nu(x) &= x[I_{\nu-1}(x) - I_{\nu+1}(x)], & 2\nu K_\nu(x) &= -x[K_{\nu-1}(x) - K_{\nu+1}(x)], \\ \frac{d}{dx} I_\nu(x) &= \frac{1}{2}[I_{\nu-1}(x) + I_{\nu+1}(x)], & \frac{d}{dx} K_\nu(x) &= -\frac{1}{2}[K_{\nu-1}(x) + K_{\nu+1}(x)]. \end{aligned}$$

Modified Bessel functions for $\nu = \pm n \pm \frac{1}{2}$ ($n = 0, 1, 2, \dots$):

$$\begin{aligned} I_{1/2}(x) &= \sqrt{\frac{2}{\pi x}} \sinh x, & I_{-1/2}(x) &= \sqrt{\frac{2}{\pi x}} \cosh x, \\ I_{3/2}(x) &= \sqrt{\frac{2}{\pi x}} \left(-\frac{1}{x} \sinh x + \cosh x \right), & I_{-3/2}(x) &= \sqrt{\frac{2}{\pi x}} \left(-\frac{1}{x} \cosh x + \sinh x \right), \\ I_{n+1/2}(x) &= \frac{1}{\sqrt{2\pi x}} \left[e^x \sum_{k=0}^n \frac{(-1)^k (n+k)!}{k! (n-k)! (2x)^k} - (-1)^n e^{-x} \sum_{k=0}^n \frac{(n+k)!}{k! (n-k)! (2x)^k} \right], \\ I_{-n-1/2}(x) &= \frac{1}{\sqrt{2\pi x}} \left[e^x \sum_{k=0}^n \frac{(-1)^k (n+k)!}{k! (n-k)! (2x)^k} + (-1)^n e^{-x} \sum_{k=0}^n \frac{(n+k)!}{k! (n-k)! (2x)^k} \right], \\ K_{\pm 1/2}(x) &= \sqrt{\frac{\pi}{2x}} e^{-x}, & K_{\pm 3/2}(x) &= \sqrt{\frac{\pi}{2x}} \left(1 + \frac{1}{x} \right) e^{-x}, \\ K_{n+1/2}(x) &= K_{-n-1/2}(x) = \sqrt{\frac{\pi}{2x}} e^{-x} \sum_{k=0}^n \frac{(n+k)!}{k! (n-k)! (2x)^k}. \end{aligned}$$

If $\nu = n$ is a nonnegative integer, then

$$\begin{aligned} K_n(x) &= (-1)^{n+1} I_n(x) \ln \frac{x}{2} + \frac{1}{2} \sum_{m=0}^{n-1} (-1)^m \left(\frac{x}{2} \right)^{2m-n} \frac{(n-m-1)!}{m!} \\ &\quad + \frac{1}{2} (-1)^n \sum_{m=0}^{\infty} \left(\frac{x}{2} \right)^{n+2m} \frac{\psi(n+m+1) + \psi(m+1)}{m! (n+m)!}; \quad n = 0, 1, 2, \dots, \end{aligned}$$

where $\psi(z)$ is the logarithmic derivative of the gamma function; for $n = 0$, the first sum is dropped.

Wronskians and similar formulas:

$$\begin{aligned} W(I_\nu, I_{-\nu}) &= -\frac{2}{\pi x} \sin(\pi\nu), & W(I_\nu, K_\nu) &= -\frac{1}{x}, \\ I_\nu(x) I_{\nu+1}(x) - I_{-\nu}(x) I_{\nu-1}(x) &= -\frac{2 \sin(\pi\nu)}{\pi x}, & I_\nu(x) K_{\nu+1}(x) + I_{\nu+1}(x) K_\nu(x) &= \frac{1}{x}, \end{aligned}$$

where $W(f, g) = f g'_x - f'_x g$.

Modified Bessel functions can be expressed in terms of Bessel functions:

$$\begin{aligned} I_\nu(z) &= e^{-\pi\nu i/2} J_\nu(ze^{\pi i/2}) & (-\pi < \arg z \leq \pi/2); \\ I_\nu(z) &= e^{3\pi\nu i/2} J_\nu(ze^{-3\pi i/2}) & (\pi/2 < \arg z \leq \pi); \\ K_\nu(z) &= \frac{1}{2}\pi i e^{\pi\nu i/2} H_\nu^{(1)}(ze^{\pi i/2}) & (-\pi < \arg z \leq \pi/2); \\ K_\nu(z) &= -\frac{1}{2}\pi i e^{-\pi\nu i/2} H_\nu^{(2)}(ze^{-\pi i/2}) & (\pi/2 < \arg z \leq \pi). \end{aligned}$$

11.7-2. Integral Representations and Asymptotic Expansions.

The functions $I_\nu(x)$ and $K_\nu(x)$ can be represented in terms of definite integrals:

$$\begin{aligned} I_\nu(x) &= \frac{x^\nu}{\pi^{1/2} 2^\nu \Gamma(\nu + \frac{1}{2})} \int_{-1}^1 \exp(-xt)(1-t^2)^{\nu-1/2} dt & (x > 0, \nu > -\frac{1}{2}), \\ K_\nu(x) &= \int_0^\infty \exp(-x \cosh t) \cosh(\nu t) dt & (x > 0), \\ K_\nu(x) &= \frac{1}{\cos(\frac{1}{2}\pi\nu)} \int_0^\infty \cos(x \sinh t) \cosh(\nu t) dt & (x > 0, -1 < \nu < 1), \\ K_\nu(x) &= \frac{1}{\sin(\frac{1}{2}\pi\nu)} \int_0^\infty \sin(x \sinh t) \sinh(\nu t) dt & (x > 0, -1 < \nu < 1). \end{aligned}$$

For integer $\nu = n$,

$$\begin{aligned} I_n(x) &= \frac{1}{\pi} \int_0^\pi \exp(x \cos t) \cos(nt) dt & (n = 0, 1, 2, \dots), \\ K_0(x) &= \int_0^\infty \cos(x \sinh t) dt = \int_0^\infty \frac{\cos(xt)}{\sqrt{t^2 + 1}} dt & (x > 0). \end{aligned}$$

Asymptotic expansions as $x \rightarrow \infty$:

$$\begin{aligned} I_\nu(x) &= \frac{e^x}{\sqrt{2\pi x}} \left\{ 1 + \sum_{m=1}^M (-1)^m \frac{(4\nu^2 - 1)(4\nu^2 - 3^2) \dots [4\nu^2 - (2m-1)^2]}{m! (8x)^m} \right\}, \\ K_\nu(x) &= \sqrt{\frac{\pi}{2x}} e^{-x} \left\{ 1 + \sum_{m=1}^M \frac{(4\nu^2 - 1)(4\nu^2 - 3^2) \dots [4\nu^2 - (2m-1)^2]}{m! (8x)^m} \right\}. \end{aligned}$$

The terms of the order of $O(x^{-M-1})$ are omitted in the braces.

Integrals with modified Bessel functions:

$$\int_0^x x^\lambda I_\nu(x) dx = \frac{x^{\lambda+\nu+1}}{2^\nu(\lambda+\nu+1)\Gamma(\nu+1)} F\left(\frac{\lambda+\nu+1}{2}, \frac{\lambda+\nu+3}{2}, \nu+1; \frac{x^2}{4}\right), \quad \operatorname{Re}(\lambda+\nu) > -1,$$

where $F(a, b, c; x)$ is the hypergeometric series (see Supplement 11.10-1),

$$\begin{aligned} \int_0^x x^\lambda K_\nu(x) dx &= \frac{2^{\nu-1}\Gamma(\nu)}{\lambda-\nu+1} x^{\lambda-\nu+1} F\left(\frac{\lambda-\nu+1}{2}, 1-\nu, \frac{\lambda-\nu+3}{2}; \frac{x^2}{4}\right) \\ &+ \frac{2^{-\nu-1}\Gamma(-\nu)}{\lambda+\nu+1} x^{\lambda+\nu+1} F\left(\frac{\lambda+\nu+1}{2}, 1+\nu, \frac{\lambda+\nu+3}{2}; \frac{x^2}{4}\right), \quad \operatorname{Re} \lambda > |\operatorname{Re} \nu| - 1. \end{aligned}$$

11.8. Airy Functions

11.8-1. Definition and Basic Formulas.

The *Airy function of the first kind*, $\text{Ai}(x)$, and the *Airy function of the second kind*, $\text{Bi}(x)$, are solutions of the *Airy equation*

$$y''_{xx} - xy = 0$$

and are defined by the formulas

$$\begin{aligned}\text{Ai}(x) &= \frac{1}{\pi} \int_0^{\infty} \cos\left(\frac{1}{3}t^3 + xt\right) dt, \\ \text{Bi}(x) &= \frac{1}{\pi} \int_0^{\infty} \left[\exp\left(-\frac{1}{3}t^3 + xt\right) + \sin\left(\frac{1}{3}t^3 + xt\right)\right] dt.\end{aligned}$$

Wronskian: $W\{\text{Ai}(x), \text{Bi}(x)\} = 1/\pi$.

Relation to the Bessel functions and the modified Bessel functions ($x > 0$):

$$\begin{aligned}\text{Ai}(x) &= \frac{1}{3} \sqrt{x} [I_{-1/3}(z) - I_{1/3}(z)] = \pi^{-1} \sqrt{\frac{1}{3}x} K_{1/3}(z), \quad z = \frac{2}{3}x^{3/2}, \\ \text{Ai}(-x) &= \frac{1}{3} \sqrt{x} [J_{-1/3}(z) + J_{1/3}(z)], \\ \text{Bi}(x) &= \sqrt{\frac{1}{3}x} [I_{-1/3}(z) + I_{1/3}(z)], \\ \text{Bi}(-x) &= \sqrt{\frac{1}{3}x} [J_{-1/3}(z) - J_{1/3}(z)].\end{aligned}$$

11.8-2. Power Series and Asymptotic Expansions.

Power series expansions as $x \rightarrow 0$:

$$\begin{aligned}\text{Ai}(x) &= c_1 f(x) - c_2 g(x), \\ \text{Bi}(x) &= \sqrt{3} [c_1 f(x) + c_2 g(x)], \\ f(x) &= 1 + \frac{1}{3!}x^3 + \frac{1 \times 4}{6!}x^6 + \frac{1 \times 4 \times 7}{9!}x^9 + \dots = \sum_{k=0}^{\infty} 3^k \left(\frac{1}{3}\right)_k \frac{x^{3k}}{(3k)!}, \\ g(x) &= x + \frac{2}{4!}x^4 + \frac{2 \times 5}{7!}x^7 + \frac{2 \times 5 \times 8}{10!}x^{10} + \dots = \sum_{k=0}^{\infty} 3^k \left(\frac{2}{3}\right)_k \frac{x^{3k+1}}{(3k+1)!},\end{aligned}$$

where $c_1 = 3^{-2/3}/\Gamma(2/3) \approx 0.3550$ and $c_2 = 3^{-1/3}/\Gamma(1/3) \approx 0.2588$.

For large values of x , the leading terms of asymptotic expansions of the Airy functions are

$$\begin{aligned}\text{Ai}(x) &\simeq \frac{1}{2} \pi^{-1/2} x^{-1/4} \exp(-z), \quad z = \frac{2}{3}x^{3/2}, \\ \text{Ai}(-x) &\simeq \pi^{-1/2} x^{-1/4} \sin\left(z + \frac{\pi}{4}\right), \\ \text{Bi}(x) &\simeq \pi^{-1/2} x^{-1/4} \exp(z), \\ \text{Bi}(-x) &\simeq \pi^{-1/2} x^{-1/4} \cos\left(z + \frac{\pi}{4}\right),\end{aligned}$$

where $x > 0$.

TABLE 1
Special cases of the Kummer confluent hypergeometric function $\Phi(a, b; z)$

a	b	z	Φ	Conventional notation
a	a	x	e^x	
1	2	$2x$	$\frac{1}{x}e^x \sinh x$	
a	$a+1$	$-x$	$ax^{-a}\gamma(a, x)$	Incomplete gamma function $\gamma(a, x) = \int_0^x e^{-t}t^{a-1} dt$
$\frac{1}{2}$	$\frac{3}{2}$	$-x^2$	$\frac{\sqrt{\pi}}{2} \operatorname{erf} x$	Error function $\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_0^x \exp(-t^2) dt$
$-n$	$\frac{1}{2}$	$\frac{x^2}{2}$	$\frac{n!}{(2n)!} \left(-\frac{1}{2}\right)^{-n} H_{2n}(x)$	Hermite polynomial $H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} (e^{-x^2})$, $n = 0, 1, 2, \dots$
$-n$	$\frac{3}{2}$	$\frac{x^2}{2}$	$\frac{n!}{(2n+1)!} \left(-\frac{1}{2}\right)^{-n} H_{2n+1}(x)$	
$-n$	b	x	$\frac{n!}{(b)_n} L_n^{(b-1)}(x)$	Laguerre polynomial $L_n^{(\alpha)}(x) = \frac{e^x x^{-\alpha}}{n!} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha})$, $\alpha = b-1$, $(b)_n = b(b+1) \dots (b+n-1)$
$\nu + \frac{1}{2}$	$2\nu+1$	$2x$	$\Gamma(1+\nu)e^x \left(\frac{x}{2}\right)^{-\nu} I_\nu(x)$	Modified Bessel function $I_\nu(x)$
$n+1$	$2n+2$	$2x$	$\Gamma\left(n+\frac{3}{2}\right)e^x \left(\frac{x}{2}\right)^{-n-\frac{1}{2}} I_{n+\frac{1}{2}}(x)$	

11.9. Confluent Hypergeometric Functions

11.9-1. Kummer and Tricomi Confluent Hypergeometric Functions.

The *confluent hypergeometric functions* $\Phi(a, b; x)$ and $\Psi(a, b; x)$ are solutions of the *degenerate hypergeometric equation* (or *confluent hypergeometric equation*)

$$xy''_{xx} + (b-x)y'_x - ay = 0.$$

In the case $b \neq 0, -1, -2, -3, \dots$, the *Kummer confluent hypergeometric function* $\Phi(a, b; x)$ can be represented as Kummer's series:

$$\Phi(a, b; x) = 1 + \sum_{k=1}^{\infty} \frac{(a)_k}{(b)_k} \frac{x^k}{k!},$$

where $(a)_k = a(a+1) \dots (a+k-1)$, $(a)_0 = 1$.

Table 1 presents some special cases where Φ can be expressed in terms of simpler functions.

TABLE 2
Special cases of the Tricomi confluent hypergeometric function $\Psi(a, b; z)$

a	b	z	Ψ	Conventional notation
$1-a$	$1-a$	x	$e^x \Gamma(a, x)$	Incomplete gamma function $\Gamma(a, x) = \int_x^\infty e^{-t} t^{a-1} dt$
$\frac{1}{2}$	$\frac{1}{2}$	x^2	$\sqrt{\pi} \exp(x^2) \operatorname{erfc} x$	Complementary error function $\operatorname{erfc} x = \frac{2}{\sqrt{\pi}} \int_x^\infty \exp(-t^2) dt$
1	1	$-x$	$-e^{-x} \operatorname{Ei}(x)$	Exponential integral $\operatorname{Ei}(x) = \int_{-\infty}^x \frac{e^t}{t} dt$
1	1	$-\ln x$	$-x^{-1} \operatorname{li} x$	Logarithmic integral $\operatorname{li} x = \int_0^x \frac{dt}{t}$
$\frac{1-n}{2}$	$\frac{3}{2}$	x^2	$2^{-n} x^{-1} H_n(x)$	Hermite polynomial $H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} (e^{-x^2})$, $n = 0, 1, 2, \dots$
$\nu + \frac{1}{2}$	$2\nu + 1$	$2x$	$\pi^{-1/2} (2x)^{-\nu} e^x K_\nu(x)$	Modified Bessel function $K_\nu(x)$
$-\frac{\nu}{2}$	$\frac{1}{2}$	$\frac{1}{2} x^2$	$2^{-\nu/2} e^{x^2/4} D_\nu(x)$	Weber parabolic cylinder function $D_\nu(x)$
$\frac{1-\nu}{2}$	$\frac{3}{2}$	$\frac{1}{2} x^2$	$2^{(1-\nu)/2} x^{-1} e^{x^2/4} D_\nu(x)$	

The Tricomi confluent hypergeometric function $\Psi(a, b; x)$ is defined as follows:

$$\Psi(a, b; x) = \frac{\Gamma(1-b)}{\Gamma(a-b+1)} \Phi(a, b; x) + \frac{\Gamma(b-1)}{\Gamma(a)} x^{1-b} \Phi(a-b+1, 2-b; x).$$

Table 2 presents some special cases where Ψ can be expressed in terms of simpler functions.
Kummer transformation:

$$\Phi(a, b; x) = e^x \Phi(b-a, b; -x), \quad \Psi(a, b; x) = x^{1-b} \Psi(1+a-b, 2-b; x).$$

Linear relations for Φ :

$$\begin{aligned} (b-a)\Phi(a-1, b; x) + (2a-b+x)\Phi(a, b; x) - a\Phi(a+1, b; x) &= 0, \\ b(b-1)\Phi(a, b-1; x) - b(b-1+x)\Phi(a, b; x) + (b-a)x\Phi(a, b+1; x) &= 0, \\ (a-b+1)\Phi(a, b; x) - a\Phi(a+1, b; x) + (b-1)\Phi(a, b-1; x) &= 0, \\ b\Phi(a, b; x) - b\Phi(a-1, b; x) - x\Phi(a, b+1; x) &= 0, \\ b(a+x)\Phi(a, b; x) - (b-a)x\Phi(a, b+1; x) - ab\Phi(a+1, b; x) &= 0, \\ (a-1+x)\Phi(a, b; x) + (b-a)\Phi(a-1, b; x) - (b-1)\Phi(a, b-1; x) &= 0. \end{aligned}$$

Linear relations for Ψ :

$$\begin{aligned} \Psi(a-1, b; x) - (2a-b+x)\Psi(a, b; x) + a(a-b+1)\Psi(a+1, b; x) &= 0, \\ (b-a-1)\Psi(a, b-1; x) - (b-1+x)\Psi(a, b; x) + x\Psi(a, b+1; x) &= 0, \\ \Psi(a, b; x) - a\Psi(a+1, b; x) - \Psi(a, b-1; x) &= 0, \\ (b-a)\Psi(a, b; x) - x\Psi(a, b+1; x) + \Psi(a-1, b; x) &= 0, \\ (a+x)\Psi(a, b; x) + a(b-a-1)\Psi(a+1, b; x) - x\Psi(a, b+1; x) &= 0, \\ (a-1+x)\Psi(a, b; x) - \Psi(a-1, b; x) + (a-c+1)\Psi(a, b-1; x) &= 0. \end{aligned}$$

Differentiation formulas:

$$\begin{aligned} \frac{d}{dx}\Phi(a, b; x) &= \frac{a}{b}\Phi(a+1, b+1; x), & \frac{d^n}{dx^n}\Phi(a, b; x) &= \frac{(a)_n}{(b)_n}\Phi(a+n, b+n; x), \\ \frac{d}{dx}\Psi(a, b; x) &= -a\Psi(a+1, b+1; x), & \frac{d^n}{dx^n}\Psi(a, b; x) &= (-1)^n(a)_n\Psi(a+n, b+n; x). \end{aligned}$$

Wronskian:

$$W(\Phi, \Psi) = \Phi\Psi'_x - \Phi'_x\Psi = -\frac{\Gamma(b)}{\Gamma(a)}x^{-b}e^x.$$

The Tricomi confluent hypergeometric function for $b = n + 1$ ($n = 0, 1, 2, \dots$):

$$\begin{aligned} \Psi(a, n+1; x) &= \frac{(-1)^{n-1}}{n!\Gamma(a-n)} \left\{ \Phi(a, n+1; x) \ln x \right. \\ &\quad \left. + \sum_{r=0}^{\infty} \frac{(a)_r}{(n+1)_r} [\psi(a+r) - \psi(1+r) - \psi(1+n+r)] \frac{x^r}{r!} \right\} + \frac{(n-1)!}{\Gamma(a)} \sum_{r=0}^{n-1} \frac{(a-n)_r}{(1-n)_r} \frac{x^{r-n}}{r!}. \end{aligned}$$

Here the last sum is dropped for $n = 0$, $\psi(z) = [\ln \Gamma(z)]'_z$ is the logarithmic derivative of the gamma function,

$$\psi(1) = -\mathcal{C}, \quad \psi(n) = -\mathcal{C} + \sum_{k=1}^{n-1} k^{-1},$$

where $\mathcal{C} = 0.5772\dots$ is the Euler constant.

If $b < 0$, then the formula

$$\Psi(a, b; x) = x^{1-b}\Psi(a-b+1, 2-b; x)$$

is valid for any x .

For $b \neq 0, -1, -2, -3, \dots$, the general solution of the degenerate hypergeometric equation can be represented in the form

$$y = C_1\Phi(a, b; x) + C_2\Psi(a, b; x),$$

and for $b = 0, -1, -2, -3, \dots$, in the form

$$y = x^{1-b} [C_1\Phi(a-b+1, 2-b; x) + C_2\Psi(a-b+1, 2-b; x)].$$

11.9-2. Integral Representations and Asymptotic Expansions.

Integral representations:

$$\Phi(a, b; x) = \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 e^{xt} t^{a-1} (1-t)^{b-a-1} dt \quad (\text{for } b > a > 0),$$

$$\Psi(a, b; x) = \frac{1}{\Gamma(a)} \int_0^\infty e^{-xt} t^{a-1} (1+t)^{b-a-1} dt \quad (\text{for } a > 0, x > 0),$$

where $\Gamma(a)$ is the gamma function.

Asymptotic expansion as $|x| \rightarrow \infty$:

$$\Phi(a, b; x) = \frac{\Gamma(b)}{\Gamma(a)} e^x x^{a-b} \left[\sum_{n=0}^N \frac{(b-a)_n (1-a)_n}{n!} x^{-n} + \varepsilon \right], \quad x > 0,$$

$$\Phi(a, b; x) = \frac{\Gamma(b)}{\Gamma(b-a)} (-x)^{-a} \left[\sum_{n=0}^N \frac{(a)_n (a-b+1)_n}{n!} (-x)^{-n} + \varepsilon \right], \quad x < 0,$$

$$\Psi(a, b; x) = x^{-a} \left[\sum_{n=0}^N (-1)^n \frac{(a)_n (a-b+1)_n}{n!} x^{-n} + \varepsilon \right], \quad -\infty < x < \infty,$$

where $\varepsilon = O(x^{-N-1})$.

Integrals with confluent hypergeometric functions:

$$\int \Phi(a, b; x) dx = \frac{b-1}{a-1} \Psi(a-1, b-1; x) + C,$$

$$\int \Psi(a, b; x) dx = \frac{1}{1-a} \Psi(a-1, b-1; x) + C,$$

$$\int x^n \Phi(a, b; x) dx = n! \sum_{k=1}^{n+1} \frac{(-1)^{k+1} (1-b)_k x^{n-k+1}}{(1-a)_k (n-k+1)!} \Phi(a-k, b-k; x) + C,$$

$$\int x^n \Psi(a, b; x) dx = n! \sum_{k=1}^{n+1} \frac{(-1)^{k+1} x^{n-k+1}}{(1-a)_k (n-k+1)!} \Psi(a-k, b-k; x) + C.$$

11.9-3. Whittaker Confluent Hypergeometric Functions.

The *Whittaker confluent hypergeometric functions* (or *Whittaker functions*) $M_{k,\mu}(x)$ and $W_{k,\mu}(x)$ are linearly independent solutions of the *Whittaker equation*:

$$y''_{xx} + \left[-\frac{1}{4} + \frac{1}{2}k + \left(\frac{1}{4} - \mu^2 \right) x^{-2} \right] y = 0.$$

The Whittaker functions are expressed in terms of the Kummer and Tricomi confluent hypergeometric functions as

$$M_{k,\mu}(x) = x^{\mu+1/2} e^{-x/2} \Phi\left(\frac{1}{2} + \mu - k, 1 + 2\mu; x\right),$$

$$W_{k,\mu}(x) = x^{\mu+1/2} e^{-x/2} \Psi\left(\frac{1}{2} + \mu - k, 1 + 2\mu; x\right).$$

11.10. Gauss Hypergeometric Functions

11.10-1. Various Representations of the Gauss Hypergeometric Function.

The *Gauss hypergeometric function* (or *hypergeometric function*) $F(\alpha, \beta, \gamma; x)$ is a solution of the *Gaussian hypergeometric equation*

$$x(x-1)y''_{xx} + [(\alpha + \beta + 1)x - \gamma]y'_x + \alpha\beta y = 0.$$

For $\gamma \neq 0, -1, -2, -3, \dots$, the function $F(\alpha, \beta, \gamma; x)$ can be expressed in terms of the hypergeometric series:

$$F(\alpha, \beta, \gamma; x) = 1 + \sum_{k=1}^{\infty} \frac{(\alpha)_k (\beta)_k}{(\gamma)_k} \frac{x^k}{k!}, \quad (\alpha)_k = \alpha(\alpha+1)\dots(\alpha+k-1),$$

which certainly converges for $|x| < 1$.

If γ is not an integer, then the general solution of the hypergeometric equation can be written in the form

$$y = C_1 F(\alpha, \beta, \gamma; x) + C_2 x^{1-\gamma} F(\alpha - \gamma + 1, \beta - \gamma + 1, 2 - \gamma; x).$$

Table 3 shows some special cases where F can be expressed in term of elementary functions.

For $\gamma > \beta > 0$, the hypergeometric function can be expressed in terms of a definite integral:

$$F(\alpha, \beta, \gamma; x) = \frac{\Gamma(\gamma)}{\Gamma(\beta)\Gamma(\gamma-\beta)} \int_0^1 t^{\beta-1} (1-t)^{\gamma-\beta-1} (1-tx)^{-\alpha} dt,$$

where $\Gamma(\beta)$ is the gamma function.

11.10-2. Basic Properties.

Linear transformation formulas:

$$F(\alpha, \beta, \gamma; x) = F(\beta, \alpha, \gamma; x),$$

$$F(\alpha, \beta, \gamma; x) = (1-x)^{\gamma-\alpha-\beta} F(\gamma-\alpha, \gamma-\beta, \gamma; x),$$

$$F(\alpha, \beta, \gamma; x) = (1-x)^{-\alpha} F\left(\alpha, \gamma-\beta, \gamma; \frac{x}{x-1}\right),$$

$$F(\alpha, \beta, \gamma; x) = (1-x)^{-\beta} F\left(\beta, \gamma-\alpha, \gamma; \frac{x}{x-1}\right).$$

Gauss's linear relations for contiguous functions:

$$(\beta - \alpha)F(\alpha, \beta, \gamma; x) + \alpha F(\alpha + 1, \beta, \gamma; x) - \beta F(\alpha, \beta + 1, \gamma; x) = 0,$$

$$(\gamma - \alpha - 1)F(\alpha, \beta, \gamma; x) + \alpha F(\alpha + 1, \beta, \gamma; x) - (\gamma - 1)F(\alpha, \beta, \gamma - 1; x) = 0,$$

$$(\gamma - \beta - 1)F(\alpha, \beta, \gamma; x) + \beta F(\alpha, \beta + 1, \gamma; x) - (\gamma - 1)F(\alpha, \beta, \gamma - 1; x) = 0,$$

$$(\gamma - \alpha - \beta)F(\alpha, \beta, \gamma; x) + \alpha(1-x)F(\alpha + 1, \beta, \gamma; x) - (\gamma - \beta)F(\alpha, \beta - 1, \gamma; x) = 0,$$

$$(\gamma - \alpha - \beta)F(\alpha, \beta, \gamma; x) - (\gamma - \alpha)F(\alpha - 1, \beta, \gamma; x) + \beta(1-x)F(\alpha, \beta + 1, \gamma; x) = 0.$$

Differentiation formulas:

$$\frac{d}{dx} F(\alpha, \beta, \gamma; x) = \frac{\alpha\beta}{\gamma} F(\alpha + 1, \beta + 1, \gamma + 1; x),$$

$$\frac{d^n}{dx^n} F(\alpha, \beta, \gamma; x) = \frac{(\alpha)_n (\beta)_n}{(\gamma)_n} F(\alpha + n, \beta + n, \gamma + n; x),$$

$$\frac{d^n}{dx^n} [x^{\gamma-1} F(\alpha, \beta, \gamma; x)] = (\gamma - n)_n x^{\gamma-n-1} F(\alpha, \beta, \gamma - n; x),$$

$$\frac{d^n}{dx^n} [x^{\alpha+n-1} F(\alpha, \beta, \gamma; x)] = (\alpha)_n x^{\alpha-1} F(\alpha + n, \beta, \gamma; x),$$

where $(\alpha)_n = \alpha(\alpha+1)\dots(\alpha+n-1)$.

See Abramowitz and Stegun (1964) and Bateman and Erdélyi (1953, Vol. 1) for more detailed information about hypergeometric functions.

TABLE 3
Some special cases where the Gauss hypergeometric function $F(\alpha, \beta, \gamma; z)$
can be expressed in terms of elementary functions

α	β	γ	z	F
$-n$	β	γ	x	$\sum_{k=0}^n \frac{(-n)_k (\beta)_k}{(\gamma)_k} \frac{x^k}{k!}$, where $n = 1, 2, \dots$
$-n$	β	$-n-m$	x	$\sum_{k=0}^n \frac{(-n)_k (\beta)_k}{(-n-m)_k} \frac{x^k}{k!}$, where $n = 1, 2, \dots$
α	β	β	x	$(1-x)^{-\alpha}$
α	$\alpha + \frac{1}{2}$	$2\alpha + 1$	x	$\left(\frac{1 + \sqrt{1-x}}{2}\right)^{-2\alpha}$
α	$\alpha + \frac{1}{2}$	2α	x	$\frac{1}{\sqrt{1-x}} \left(\frac{1 + \sqrt{1-x}}{2}\right)^{1-2\alpha}$
α	$\alpha + \frac{1}{2}$	$\frac{3}{2}$	x^2	$\frac{(1+x)^{1-2\alpha} - (1-x)^{1-2\alpha}}{2x(1-2\alpha)}$
α	$\alpha + \frac{1}{2}$	$\frac{1}{2}$	x^2	$\frac{1}{2} [(1+x)^{-2\alpha} + (1-x)^{-2\alpha}]$
α	$\alpha + \frac{1}{2}$	$\frac{1}{2}$	$-\tan^2 x$	$\cos^{2\alpha} x \cos(2\alpha x)$
α	$\alpha - \frac{1}{2}$	2α	x	$2^{2\alpha-1} (1 + \sqrt{1-x})^{1-2\alpha}$
α	$\frac{1}{2}\alpha + 1$	$\frac{1}{2}\alpha$	x	$(1+x)(1-x)^{-\alpha-1}$
α	$2-\alpha$	$\frac{3}{2}$	$\sin^2 x$	$\frac{\sin[(2\alpha-2)x]}{(\alpha-1)\sin(2x)}$
α	$1-\alpha$	$\frac{3}{2}$	$\sin^2 x$	$\frac{\sin[(2\alpha-1)x]}{(\alpha-1)\sin(2x)}$
α	$1-\alpha$	$\frac{1}{2}$	$-x^2$	$\frac{(\sqrt{1+x^2}+x)^{2\alpha-1} + (\sqrt{1+x^2}-x)^{2\alpha-1}}{2\sqrt{1+x^2}}$
α	$1-\alpha$	$\frac{1}{2}$	$\sin^2 x$	$\frac{\cos[(2\alpha-1)x]}{\cos x}$
α	$-\alpha$	$\frac{1}{2}$	$-x^2$	$\frac{1}{2} [(\sqrt{1+x^2}+x)^{2\alpha} + (\sqrt{1+x^2}-x)^{2\alpha}]$
α	$-\alpha$	$\frac{1}{2}$	$\sin^2 x$	$\cos(2\alpha x)$
1	1	2	$-x$	$\frac{1}{x} \ln(x+1)$
$\frac{1}{2}$	1	$\frac{3}{2}$	x^2	$\frac{1}{2x} \ln \frac{1+x}{1-x}$
$\frac{1}{2}$	1	$\frac{3}{2}$	$-x^2$	$\frac{1}{x} \arctan x$
$\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{2}$	x^2	$\frac{1}{x} \arcsin x$
$\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{2}$	$-x^2$	$\frac{1}{x} \operatorname{arcsinh} x$
$n+1$	$n+m+1$	$n+m+l+2$	x	$\frac{(-1)^m (n+m+l+1)!}{n! l! (n+m)! (m+l)!} \frac{d^{n+m}}{dx^{n+m}} \left\{ (1-x)^{m+l} \frac{d^l F}{dx^l} \right\}$, $F = -\frac{\ln(1-x)}{x}$, $n, m, l = 0, 1, 2, \dots$

11.11. Legendre Polynomials, Legendre Functions, and Associated Legendre Functions

11.11-1. Legendre Polynomials and Legendre Functions.

The *Legendre polynomials* $P_n(x)$ and the *Legendre functions* $Q_n(x)$ are solutions of the second-order linear ordinary differential equation

$$(1 - x^2)y''_{xx} - 2xy'_x + n(n + 1)y = 0.$$

The *Legendre polynomials* $P_n(x)$ and the Legendre functions $Q_n(x)$ are defined by the formulas

$$P_n(x) = \frac{1}{n! 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n,$$

$$Q_n(x) = \frac{1}{2} P_n(x) \ln \frac{1+x}{1-x} - \sum_{m=1}^n \frac{1}{m} P_{m-1}(x) P_{n-m}(x).$$

The polynomials $P_n = P_n(x)$ can be calculated using the formulas

$$P_0(x) = 1, \quad P_1(x) = x, \quad P_2(x) = \frac{1}{2}(3x^2 - 1),$$

$$P_3(x) = \frac{1}{2}(5x^3 - 3x), \quad P_4(x) = \frac{1}{8}(35x^4 - 30x^2 + 3),$$

$$P_{n+1}(x) = \frac{2n+1}{n+1} x P_n(x) - \frac{n}{n+1} P_{n-1}(x).$$

The first five functions $Q_n = Q_n(x)$ have the form

$$Q_0(x) = \frac{1}{2} \ln \frac{1+x}{1-x}, \quad Q_1(x) = \frac{x}{2} \ln \frac{1+x}{1-x} - 1,$$

$$Q_2(x) = \frac{1}{4}(3x^2 - 1) \ln \frac{1+x}{1-x} - \frac{3}{2}x, \quad Q_3(x) = \frac{1}{4}(5x^3 - 3x) \ln \frac{1+x}{1-x} - \frac{5}{2}x^2 + \frac{2}{3},$$

$$Q_4(x) = \frac{1}{16}(35x^4 - 30x^2 + 3) \ln \frac{1+x}{1-x} - \frac{35}{8}x^3 + \frac{55}{24}x.$$

The polynomials $P_n(x)$ have the explicit representation

$$P_n(x) = 2^{-n} \sum_{m=0}^{[n/2]} (-1)^m C_n^m C_{2n-2m}^n x^{n-2m},$$

where $[A]$ stands for the integer part of a number A .

Integral representation of the Legendre polynomials (*Laplace integral*):

$$P_n(x) = \frac{1}{\pi} \int_0^\pi (x \pm \sqrt{x^2 - 1} \cos t)^n dt, \quad x > 1.$$

Integral representation of the Legendre polynomials (*Dirichlet–Mehler integral*):

$$P_n(\cos \theta) = \frac{\sqrt{2}}{\pi} \int_0^\theta \frac{\cos[(n + \frac{1}{2})\psi] d\psi}{\sqrt{\cos \psi - \cos \theta}}, \quad 0 < \theta < \pi, \quad n = 0, 1, \dots$$

Integral representation of the Legendre functions:

$$Q_n(x) = 2^n \int_x^\infty \frac{(t-x)^n}{(t^2-1)^{n+1}} dt, \quad x > 1.$$

Properties:

$$P_n(-x) = (-1)^n P_n(x), \quad Q_n(-x) = (-1)^{n+1} Q_n(x).$$

Recurrence relations:

$$(n+1)P_{n+1}(x) - (2n+1)xP_n(x) + nP_{n-1}(x) = 0,$$

$$(x^2-1)\frac{d}{dx}P_n(x) = n[xP_n(x) - P_{n-1}(x)] = \frac{n(n+1)}{2n+1}[P_{n+1}(x) - P_{n-1}(x)].$$

Values of the Legendre polynomials and their derivatives at $x = 0$:

$$P_{2m}(0) = (-1)^m \frac{(2m-1)!!}{2^m m!}, \quad P_{2m+1}(0) = 0, \quad P'_{2m}(0) = 0, \quad P'_{2m+1}(0) = (-1)^m \frac{(2m+1)!!}{2^m m!}.$$

Asymptotic formula as $n \rightarrow \infty$:

$$P_n(\cos \theta) \approx \left(\frac{2}{\pi n \sin \theta} \right)^{1/2} \sin \left[\left(n + \frac{1}{2} \right) \theta + \frac{\pi}{4} \right], \quad 0 < \theta < \pi.$$

The polynomials $P_n(x)$ (with natural n) have exactly n real distinct zeros; all zeros lie on the interval $-1 < x < 1$. The zeros of $P_n(x)$ and $P_{n+1}(x)$ alternate with each other. The function $Q_n(x)$ has exactly $n+1$ zeros, which lie on the interval $-1 < x < 1$.

The functions $P_n(x)$ form an orthogonal system on the interval $-1 \leq x \leq 1$, with

$$\int_{-1}^1 P_n(x)P_m(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{2}{2n+1} & \text{if } n = m. \end{cases}$$

The generating function for Legendre polynomials is

$$\frac{1}{\sqrt{1-2sx+s^2}} = \sum_{n=0}^{\infty} P_n(x)s^n \quad (|s| < 1).$$

The generating function for Legendre functions is

$$\frac{1}{\sqrt{1-2sx+s^2}} \ln \left[\frac{x-s+\sqrt{1-2sx+s^2}}{\sqrt{1-x^2}} \right] = \sum_{n=0}^{\infty} Q_n(x)s^n \quad (|s| < 1, x > 1).$$

11.11-2. Associated Legendre Functions with Integer Indices and Real Argument.

The associated Legendre functions $P_n^m(x)$ of order m are defined by the formulas

$$P_n^m(x) = (1-x^2)^{m/2} \frac{d^m}{dx^m} P_n(x), \quad n = 1, 2, 3, \dots, \quad m = 0, 1, 2, \dots$$

It is assumed by definition that $P_n^0(x) = P_n(x)$.

Properties:

$$P_n^m(x) = 0 \quad \text{if } m > n, \quad P_n^m(-x) = (-1)^{n-m} P_n^m(x).$$

The associated Legendre functions $P_n^m(x)$ have exactly $n-m$ real zeros, which lie on the interval $-1 < x < 1$.

The associated Legendre functions $P_n^m(x)$ with low indices:

$$P_1^1(x) = (1-x^2)^{1/2}, \quad P_2^1(x) = 3x(1-x^2)^{1/2}, \quad P_2^2(x) = 3(1-x^2),$$

$$P_3^1(x) = \frac{3}{2}(5x^2-1)(1-x^2)^{1/2}, \quad P_3^2(x) = 15x(1-x^2), \quad P_3^3(x) = 15(1-x^2)^{3/2}.$$

The associated Legendre functions $P_n^m(x)$ with $n > m$ are solutions of the linear ordinary differential equation

$$(1-x^2)y''_{xx} - 2xy'_x + \left[n(n+1) - \frac{m^2}{1-x^2} \right] y = 0.$$

The functions $P_n^m(x)$ form an orthogonal system on the interval $-1 \leq x \leq 1$, with

$$\int_{-1}^1 P_n^m(x) P_k^m(x) dx = \begin{cases} 0 & \text{if } n \neq k, \\ \frac{2}{2n+1} \frac{(n+m)!}{(n-m)!} & \text{if } n = k. \end{cases}$$

The functions $P_n^m(x)$ (with $m \neq 0$) are orthogonal on the interval $-1 \leq x \leq 1$ with weight $(1-x^2)^{-1}$, that is,

$$\int_{-1}^1 \frac{P_n^m(x) P_k^m(x)}{1-x^2} dx = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{(n+m)!}{m(n-m)!} & \text{if } m = k. \end{cases}$$

11.11-3. Associated Legendre Functions. General Case.

In the general case, the associated Legendre functions of the first and the second kind, $P_\nu^\mu(z)$ and $Q_\nu^\mu(z)$, are linearly independent solutions of the *Legendre equation*

$$(1-z^2)y''_{zz} - 2zy'_z + \left[\nu(\nu+1) - \frac{\mu^2}{1-z^2} \right] y = 0,$$

where the parameters ν and μ and the variable z can assume arbitrary real or complex values.

For $|1-z| < 2$, the formulas

$$P_\nu^\mu(z) = \frac{1}{\Gamma(1-\mu)} \left(\frac{z+1}{z-1} \right)^{\mu/2} F\left(-\nu, 1+\nu, 1-\mu; \frac{1-z}{2}\right),$$

$$Q_\nu^\mu(z) = A \left(\frac{z-1}{z+1} \right)^{\frac{\mu}{2}} F\left(-\nu, 1+\nu, 1+\mu; \frac{1-z}{2}\right) + B \left(\frac{z+1}{z-1} \right)^{\frac{\mu}{2}} F\left(-\nu, 1+\nu, 1-\mu; \frac{1-z}{2}\right),$$

$$A = e^{i\mu\pi} \frac{\Gamma(-\mu)\Gamma(1+\nu+\mu)}{2\Gamma(1+\nu-\mu)}, \quad B = e^{i\mu\pi} \frac{\Gamma(\mu)}{2}, \quad i^2 = -1,$$

are valid, where $F(a, b, c; z)$ is the hypergeometric series (see Supplement 11.10).

For $|z| > 1$,

$$P_\nu^\mu(z) = \frac{2^{-\nu-1}\Gamma(-\frac{1}{2}-\nu)}{\sqrt{\pi}\Gamma(-\nu-\mu)} z^{-\nu+\mu-1} (z^2-1)^{-\mu/2} F\left(\frac{1+\nu-\mu}{2}, \frac{2+\nu-\mu}{2}, \frac{2\nu+3}{2}; \frac{1}{z^2}\right)$$

$$+ \frac{2^\nu\Gamma(\frac{1}{2}+\nu)}{\Gamma(1+\nu-\mu)} z^{\nu+\mu} (z^2-1)^{-\mu/2} F\left(-\frac{\nu+\mu}{2}, \frac{1-\nu-\mu}{2}, \frac{1-2\nu}{2}; \frac{1}{z^2}\right),$$

$$Q_\nu^\mu(z) = e^{i\pi\mu} \frac{\sqrt{\pi}\Gamma(\nu+\mu+1)}{2^{\nu+1}\Gamma(\nu+\frac{3}{2})} z^{-\nu-\mu-1} (z^2-1)^{\mu/2} F\left(\frac{2+\nu+\mu}{2}, \frac{1+\nu+\mu}{2}, \frac{2\nu+3}{2}; \frac{1}{z^2}\right).$$

The functions $P_\nu(z) \equiv P_\nu^0(z)$ and $Q_\nu(z) \equiv Q_\nu^0(z)$ are called the *Legendre functions*.

For $n = 1, 2, \dots$,

$$P_\nu^n(z) = (z^2 - 1)^{n/2} \frac{d^n}{dz^n} P_\nu(z), \quad Q_\nu^n(z) = (z^2 - 1)^{n/2} \frac{d^n}{dz^n} Q_\nu(z).$$

Relations between associated Legendre functions:

$$P_\nu^\mu(z) = P_{-\nu-1}^\mu(z), \quad P_\nu^n(z) = \frac{\Gamma(\nu + n + 1)}{\Gamma(\nu - n + 1)} P_\nu^{-n}(z), \quad n = 0, 1, 2, \dots,$$

$$P_{\nu+1}^\mu(z) = \frac{2\nu + 1}{\nu - \mu + 1} z P_\nu^\mu(z) - \frac{\nu + \mu}{\nu - \mu + 1} P_{\nu-1}^\mu(z),$$

$$P_{\nu+1}^\mu(z) = P_{\nu-1}^\mu(z) + (2\nu + 1)(z^2 - 1)^{1/2} P_\nu^{\mu-1}(z),$$

$$(z^2 - 1) \frac{d}{dz} P_\nu^\mu(z) = \nu z P_\nu^\mu(z) - (\nu + \mu) P_{\nu-1}^\mu(z),$$

$$Q_\nu^\mu(z) = \frac{\pi}{2 \sin(\mu\pi)} e^{i\pi\mu} \left[P_\nu^\mu(z) - \frac{\Gamma(1 + \nu + \mu)}{\Gamma(1 + \nu - \mu)} P_\nu^{-\mu}(z) \right],$$

$$Q_\nu^\mu(z) = e^{i\pi\mu} \left(\frac{\pi}{2} \right)^{1/2} \Gamma(\nu + \mu + 1) (z^2 - 1)^{-1/4} P_{-\mu-1/2}^{-\nu-1/2} \left(\frac{z}{\sqrt{z^2 - 1}} \right), \quad \operatorname{Re} z > 0.$$

Integral representation for $\operatorname{Re}(-\mu) > \operatorname{Re} \nu > -1$:

$$P_\nu^\mu(z) = \frac{2^{-\nu} (z^2 - 1)^{-\mu/2}}{\Gamma(\nu + 1) \Gamma(-\mu - \nu)} \int_0^\infty (z + \cosh t)^{\mu - \nu - 1} (\sinh t)^{2\nu + 1} dt,$$

where z does not lie on the real axis between -1 and ∞ .

Integral representation for $\mu < 1/2$:

$$P_\nu^\mu(z) = \frac{2^\mu (z^2 - 1)^{-\mu/2}}{\sqrt{\pi} \Gamma(\frac{1}{2} - \mu)} \int_0^\pi (z + \sqrt{z^2 - 1} \cos t)^{\nu + \mu} (\sin t)^{-2\mu} dt,$$

where z does not lie on the real axis between -1 and 1 .

Integral representation for $\operatorname{Re} \nu > -1$ and $\operatorname{Re}(\nu + \mu + 1) > 0$:

$$Q_\nu^\mu(z) = e^{\pi\mu i} \frac{\Gamma(\nu + \mu + 1) (z^2 - 1)^{-\mu/2}}{2^{\nu+1} \Gamma(\nu + 1)} \int_0^\pi (z + \cos t)^{\mu - \nu - 1} (\sin t)^{2\nu + 1} dt,$$

where z does not lie on the real axis between -1 and 1 .

For $n = 0, 1, 2, \dots$,

$$P_\nu^n(z) = \frac{\Gamma(\nu + n + 1)}{\pi \Gamma(\nu + 1)} \int_0^\pi (z + \sqrt{z^2 - 1} \cos t)^\nu \cos(nt) dt, \quad \operatorname{Re} z > 0;$$

$$Q_\nu^n(z) = (-1)^n \frac{\Gamma(\nu + n + 1)}{2^{\nu+1} \Gamma(\nu + 1)} (z^2 - 1)^{-n/2} \int_0^\pi (z + \cos t)^{n - \nu - 1} (\sin t)^{2\nu + 1} dt, \quad \operatorname{Re} \nu > -1.$$

Note that $z \neq x$, $-1 < x < 1$, in the latter formula.

The *modified associated Legendre functions*, on the cut $z = x$, $-1 < x < 1$, of the real axis are defined by the formulas

$$\begin{aligned} \mathbf{P}_\nu^\mu(x) &= \frac{1}{2} [e^{\frac{1}{2}i\mu\pi} P_\nu^\mu(x + i0) + e^{-\frac{1}{2}i\mu\pi} P_\nu^\mu(x - i0)] \\ &= \frac{1}{\Gamma(1 - \mu)} \left(\frac{1 + x}{1 - x} \right)^{\mu/2} F\left(-\nu, 1 + \nu, 1 - \mu; \frac{1 - x}{2}\right), \\ \mathbf{Q}_\nu^\mu(x) &= \frac{1}{2} e^{-i\mu\pi} [e^{-\frac{1}{2}i\mu\pi} Q_\nu^\mu(x + i0) + e^{\frac{1}{2}i\mu\pi} Q_\nu^\mu(x - i0)] \\ &= \frac{\pi}{2 \sin(\pi\mu)} \left[\cos(\pi\mu) \mathbf{P}_\nu^\mu(x) - \frac{\Gamma(\nu + \mu + 1)}{\Gamma(\nu - \mu + 1)} \mathbf{P}_\nu^{-\mu}(x) \right]. \end{aligned}$$

Notation:

$$P_\nu(x) = P_\nu^0(x), \quad Q_\nu(x) = Q_\nu^0(x).$$

For $-1 < x < 1$, the modified associated Legendre functions can be represented in the form of the trigonometric series:

$$P_\nu^\mu(\cos \theta) = \frac{2^{\mu+1}}{\sqrt{\pi}} \frac{\Gamma(\nu + \mu + 1)}{\Gamma(\nu + \frac{3}{2})} (\sin \theta)^\mu \sum_{k=0}^{\infty} \frac{(\frac{1}{2} + \mu)_k (1 + \nu + \mu)_k}{k! (\nu + \frac{3}{2})_k} \sin[(2k + \nu + \mu + 1)\theta],$$

$$Q_\nu^\mu(\cos \theta) = \sqrt{\pi} 2^\mu \frac{\Gamma(\nu + \mu + 1)}{\Gamma(\nu + \frac{3}{2})} (\sin \theta)^\mu \sum_{k=0}^{\infty} \frac{(\frac{1}{2} + \mu)_k (1 + \nu + \mu)_k}{k! (\nu + \frac{3}{2})_k} \cos[(2k + \nu + \mu + 1)\theta],$$

where $0 < \theta < \pi$.

For $0 < x < 1$,

$$P_\nu^\mu(-x) = P_\nu^\mu(x) \cos[\pi(\nu + \mu)] - 2\pi^{-1} Q_\nu^\mu(x) \sin[\pi(\nu + \mu)],$$

$$Q_\nu^\mu(-x) = -Q_\nu^\mu(x) \cos[\pi(\nu + \mu)] - \frac{1}{2}\pi P_\nu^\mu(x) \sin[\pi(\nu + \mu)].$$

For $-1 < x < 1$,

$$P_{\nu+1}^\mu(x) = \frac{2\nu + 1}{\nu - \mu + 1} x P_\nu^\mu(x) - \frac{\nu + \mu}{\nu - \mu + 1} P_{\nu-1}^\mu(x),$$

$$P_{\nu+1}^\mu(x) = P_{\nu-1}^\mu(x) - (2\nu + 1)(1 - x^2)^{1/2} P_\nu^{\mu-1}(x),$$

$$P_{\nu+1}^\mu(x) = x P_\nu^\mu(x) - (\nu + \mu)(1 - x^2)^{1/2} P_\nu^{\mu-1}(x),$$

$$\frac{d}{dx} P_\nu^\mu(x) = \frac{\nu x}{x^2 - 1} P_\nu^\mu(x) - \frac{\nu + \mu}{x^2 - 1} P_{\nu-1}^\mu(x).$$

Wronskian:

$$P_\nu^\mu(x) \frac{d}{dx} Q_\nu^\mu(x) - Q_\nu^\mu(x) \frac{d}{dx} P_\nu^\mu(x) = \frac{k}{1 - x^2}, \quad k = 2^{2\mu} \frac{\Gamma(\frac{\nu+\mu+1}{2}) \Gamma(\frac{\nu+\mu+2}{2})}{\Gamma(\frac{\nu-\mu+1}{2}) \Gamma(\frac{\nu-\mu+2}{2})}.$$

For $n = 1, 2, \dots$,

$$P_\nu^n(x) = (-1)^n (1 - x^2)^{n/2} \frac{d^n}{dx^n} P_\nu(x), \quad Q_\nu^n(x) = (-1)^n (1 - x^2)^{n/2} \frac{d^n}{dx^n} Q_\nu(x).$$

11.12. Parabolic Cylinder Functions

11.12-1. Definitions. Basic Formulas.

The *Weber parabolic cylinder function* $D_\nu(z)$ is a solution of the linear ordinary differential equation:

$$y''_{zz} + \left(-\frac{1}{4}z^2 + \nu + \frac{1}{2}\right)y = 0,$$

where the parameter ν and the variable z can assume arbitrary real or complex values. Another linearly independent solution of this equation is the function $D_{-\nu-1}(iz)$; if ν is noninteger, then $D_\nu(-z)$ can also be taken as a linearly independent solution.

The parabolic cylinder functions can be expressed in terms of confluent hypergeometric functions as

$$D_\nu(z) = 2^{1/2} \exp\left(-\frac{1}{4}z^2\right) \left[\frac{\Gamma(\frac{1}{2})}{\Gamma(\frac{1}{2} - \frac{\nu}{2})} \Phi\left(-\frac{\nu}{2}, \frac{1}{2}; \frac{1}{2}z^2\right) + 2^{-1/2} \frac{\Gamma(-\frac{1}{2})}{\Gamma(-\frac{\nu}{2})} z \Phi\left(\frac{1}{2} - \frac{\nu}{2}, \frac{3}{2}; \frac{1}{2}z^2\right) \right].$$

For nonnegative integer $\nu = n$, we have

$$D_n(z) = \frac{1}{2^{n/2}} \exp\left(-\frac{z^2}{4}\right) H_n\left(\frac{z}{\sqrt{2}}\right), \quad n = 0, 1, 2, \dots;$$

$$H_n(z) = (-1)^n \exp(z^2) \frac{d^n}{dz^n} \exp(-z^2),$$

where $H_n(z)$ is the Hermitian polynomial of order n .

Connection with the error function:

$$D_{-1}(z) = \sqrt{\frac{\pi}{2}} \exp\left(\frac{z^2}{4}\right) \operatorname{erfc}\left(\frac{z}{\sqrt{2}}\right),$$

$$D_{-2}(z) = \sqrt{\frac{\pi}{2}} z \exp\left(\frac{z^2}{4}\right) \operatorname{erfc}\left(\frac{z}{\sqrt{2}}\right) - \exp\left(-\frac{z^2}{4}\right).$$

11.12-2. Integral Representations, Asymptotic Expansions, and Linear Relations.

Integral representations:

$$D_\nu(z) = \sqrt{2/\pi} \exp\left(\frac{1}{4}z^2\right) \int_0^\infty t^\nu \exp\left(-\frac{1}{2}t^2\right) \cos\left(z t - \frac{1}{2}\pi\nu\right) dt \quad \text{for } \operatorname{Re} \nu > -1,$$

$$D_\nu(z) = \frac{1}{\Gamma(-\nu)} \exp\left(-\frac{1}{4}z^2\right) \int_0^\infty t^{-\nu-1} \exp\left(-zt - \frac{1}{2}t^2\right) dt \quad \text{for } \operatorname{Re} \nu < 0.$$

Asymptotic expansion as $|z| \rightarrow \infty$:

$$D_\nu(z) = z^\nu \exp\left(-\frac{1}{4}z^2\right) \left[\sum_{n=0}^N \frac{(-2)^n \left(-\frac{\nu}{2}\right)_n \left(\frac{1}{2} - \frac{\nu}{2}\right)_n}{n!} \frac{1}{z^{2n}} + O(|z|^{-2N-2}) \right] \quad \text{for } |\arg z| < \frac{3\pi}{4},$$

where $(a)_0 = 1$, $(a)_n = a(a+1)\dots(a+n-1)$ for $n = 1, 2, 3, \dots$

Recurrence relations:

$$D_{\nu+1}(z) - zD_\nu(z) + \nu D_{\nu-1}(z) = 0,$$

$$\frac{d}{dz} D_\nu(z) + \frac{1}{2}zD_\nu(z) - \nu D_{\nu-1}(z) = 0,$$

$$\frac{d}{dz} D_\nu(z) - \frac{1}{2}zD_\nu(z) + D_{\nu+1}(z) = 0.$$

11.13. Elliptic Integrals

11.13-1. Complete Elliptic Integrals.

Complete elliptic integral of the first kind:

$$\mathbf{K}(k) = \int_0^{\pi/2} \frac{d\alpha}{\sqrt{1-k^2 \sin^2 \alpha}} = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-k^2 x^2)}}.$$

Complete elliptic integral of the second kind:

$$\mathbf{E}(k) = \int_0^{\pi/2} \sqrt{1-k^2 \sin^2 \alpha} d\alpha = \int_0^1 \frac{\sqrt{1-k^2 x^2}}{\sqrt{1-x^2}} dx.$$

The argument k is called the *elliptic modulus* ($k^2 < 1$).

Notation:

$$k' = \sqrt{1 - k^2}, \quad \mathbf{K}'(k) = \mathbf{K}(k'), \quad \mathbf{E}'(k) = \mathbf{E}(k'),$$

where k' is the *complementary modulus*.

Properties:

$$\begin{aligned} \mathbf{K}(-k) &= \mathbf{K}(k), & \mathbf{E}(-k) &= \mathbf{E}(k); \\ \mathbf{K}(k) &= \mathbf{K}'(k'), & \mathbf{E}(k) &= \mathbf{E}'(k'); \\ \mathbf{E}(k) \mathbf{K}'(k) + \mathbf{E}'(k) \mathbf{K}(k) - \mathbf{K}(k) \mathbf{K}'(k) &= \frac{\pi}{2}. \end{aligned}$$

Conversion formulas for complete elliptic integrals:

$$\begin{aligned} \mathbf{K}\left(\frac{1 - k'}{1 + k'}\right) &= \frac{1 + k'}{2} \mathbf{K}(k), \\ \mathbf{E}\left(\frac{1 - k'}{1 + k'}\right) &= \frac{1}{1 + k'} [\mathbf{E}(k) + k' \mathbf{K}(k)], \\ \mathbf{K}\left(\frac{2\sqrt{k}}{1 + k}\right) &= (1 + k) \mathbf{K}(k), \\ \mathbf{E}\left(\frac{2\sqrt{k}}{1 + k}\right) &= \frac{1}{1 + k} [2 \mathbf{E}(k) - (k')^2 \mathbf{K}(k)]. \end{aligned}$$

Representation of complete elliptic integrals in the form of series in powers of the modulus k :

$$\begin{aligned} \mathbf{K}(k) &= \frac{\pi}{2} \left\{ 1 + \left(\frac{1}{2}\right)^2 k^2 + \left(\frac{1 \times 3}{2 \times 4}\right)^2 k^4 + \dots + \left[\frac{(2n-1)!!}{(2n)!!}\right]^2 k^{2n} + \dots \right\}, \\ \mathbf{E}(k) &= \frac{\pi}{2} \left\{ 1 - \left(\frac{1}{2}\right)^2 \frac{k^2}{1} - \left(\frac{1 \times 3}{2 \times 4}\right)^2 \frac{k^4}{3} - \dots - \left[\frac{(2n-1)!!}{(2n)!!}\right]^2 \frac{k^{2n}}{2n-1} - \dots \right\}. \end{aligned}$$

Representation of complete elliptic integrals in the form of series in powers of the complementary modulus $k' = \sqrt{1 - k^2}$:

$$\begin{aligned} \mathbf{K}(k) &= \frac{\pi}{1 + k'} \left\{ 1 + \left(\frac{1}{2}\right)^2 \left(\frac{1 - k'}{1 + k'}\right)^2 + \left(\frac{1 \times 3}{2 \times 4}\right)^2 \left(\frac{1 - k'}{1 + k'}\right)^4 + \dots + \left[\frac{(2n-1)!!}{(2n)!!}\right]^2 \left(\frac{1 - k'}{1 + k'}\right)^{2n} + \dots \right\}, \\ \mathbf{K}(k) &= \ln \frac{4}{k'} + \left(\frac{1}{2}\right)^2 \left(\ln \frac{4}{k'} - \frac{2}{1 \times 2}\right) (k')^2 + \left(\frac{1 \times 3}{2 \times 4}\right)^2 \left(\ln \frac{4}{k'} - \frac{2}{1 \times 2} - \frac{2}{3 \times 4}\right) (k')^4 \\ &\quad + \left(\frac{1 \times 3 \times 5}{2 \times 4 \times 6}\right)^2 \left(\ln \frac{4}{k'} - \frac{2}{1 \times 2} - \frac{2}{3 \times 4} - \frac{2}{5 \times 6}\right) (k')^6 + \dots; \\ \mathbf{E}(k) &= \frac{\pi(1 + k')}{4} \left\{ 1 + \frac{1}{2^2} - \left(\frac{1 - k'}{1 + k'}\right)^2 + \frac{1^2}{(2 \times 4)^2} \left(\frac{1 - k'}{1 + k'}\right)^4 + \dots + \left[\frac{(2n-3)!!}{(2n)!!}\right]^2 \left(\frac{1 - k'}{1 + k'}\right)^{2n} + \dots \right\}, \\ \mathbf{E}(k) &= 1 + \frac{1}{2} \left(\ln \frac{4}{k'} - \frac{1}{1 \times 2}\right) (k')^2 + \frac{1^2 \times 3}{2^2 \times 4} \left(\ln \frac{4}{k'} - \frac{2}{1 \times 2} - \frac{1}{3 \times 4}\right) (k')^4 \\ &\quad + \frac{1^2 \times 3^2 \times 5}{2^2 \times 4^2 \times 6} \left(\ln \frac{4}{k'} - \frac{2}{1 \times 2} - \frac{2}{3 \times 4} - \frac{1}{5 \times 6}\right) (k')^6 + \dots. \end{aligned}$$

Differentiation formulas:

$$\frac{d\mathbf{K}(k)}{dk} = \frac{\mathbf{E}(k)}{k(k')^2} - \frac{\mathbf{K}(k)}{k}, \quad \frac{d\mathbf{E}(k)}{dk} = \frac{\mathbf{E}(k) - \mathbf{K}(k)}{k}.$$

The functions $\mathbf{K}(k)$ and $\mathbf{K}'(k)$ satisfy the second-order linear ordinary differential equation

$$\frac{d}{dk} \left[k(1-k^2) \frac{d\mathbf{K}}{dk} \right] - k\mathbf{K} = 0.$$

The functions $\mathbf{E}(k)$ and $\mathbf{E}'(k) - \mathbf{K}'(k)$ satisfy the second-order linear ordinary differential equation

$$(1-k^2) \frac{d}{dk} \left(k \frac{d\mathbf{E}}{dk} \right) + k\mathbf{E} = 0.$$

11.13-2. Incomplete Elliptic Integrals (Elliptic Integrals).

Elliptic integral of the first kind:

$$F(\varphi, k) = \int_0^\varphi \frac{d\alpha}{\sqrt{1-k^2 \sin^2 \alpha}} = \int_0^{\sin \varphi} \frac{dx}{\sqrt{(1-x^2)(1-k^2 x^2)}}.$$

Elliptic integral of the second kind:

$$E(\varphi, k) = \int_0^\varphi \sqrt{1-k^2 \sin^2 \alpha} d\alpha = \int_0^{\sin \varphi} \frac{\sqrt{1-k^2 x^2}}{\sqrt{1-x^2}} dx.$$

Elliptic integral of the third kind:

$$\Pi(\varphi, n, k) = \int_0^\varphi \frac{d\alpha}{(1-n \sin^2 \alpha) \sqrt{1-k^2 \sin^2 \alpha}} = \int_0^{\sin \varphi} \frac{dx}{(1-nx^2) \sqrt{(1-x^2)(1-k^2 x^2)}}.$$

The quantity k is called the *elliptic modulus* ($k^2 < 1$), $k' = \sqrt{1-k^2}$ is the *complementary modulus*, and n is the *characteristic parameter*.

Complete elliptic integrals:

$$\begin{aligned} \mathbf{K}(k) &= F\left(\frac{\pi}{2}, k\right), & \mathbf{E}(k) &= E\left(\frac{\pi}{2}, k\right), \\ \mathbf{K}'(k) &= F\left(\frac{\pi}{2}, k'\right), & \mathbf{E}'(k) &= E\left(\frac{\pi}{2}, k'\right). \end{aligned}$$

Properties of elliptic integrals:

$$\begin{aligned} F(-\varphi, k) &= -F(\varphi, k), & F(n\pi \pm \varphi, k) &= 2n\mathbf{K}(k) \pm F(\varphi, k); \\ E(-\varphi, k) &= -E(\varphi, k), & E(n\pi \pm \varphi, k) &= 2n\mathbf{E}(k) \pm E(\varphi, k). \end{aligned}$$

Conversion formulas for elliptic integrals (first set):

$$\begin{aligned} F\left(\psi, \frac{1}{k}\right) &= kF(\varphi, k), \\ E\left(\psi, \frac{1}{k}\right) &= \frac{1}{k} [E(\varphi, k) - (k')^2 F(\varphi, k)], \end{aligned}$$

where the angles φ and ψ are related by $\sin \psi = k \sin \varphi$, $\cos \psi = \sqrt{1-k^2 \sin^2 \varphi}$.

Conversion formulas for elliptic integrals (second set):

$$F\left(\psi, \frac{1-k'}{1+k'}\right) = (1+k')F(\varphi, k),$$

$$E\left(\psi, \frac{1-k'}{1+k'}\right) = \frac{2}{1+k'} [E(\varphi, k) + k'F(\varphi, k)] - \frac{1-k'}{1+k'} \sin \psi,$$

where the angles φ and ψ are related by $\tan(\psi - \varphi) = k' \tan \varphi$.

Transformation formulas for elliptic integrals (third set):

$$F\left(\psi, \frac{2\sqrt{k}}{1+k}\right) = (1+k)F(\varphi, k),$$

$$E\left(\psi, \frac{2\sqrt{k}}{1+k}\right) = \frac{1}{1+k} \left[2E(\varphi, k) - (k')^2 F(\varphi, k) + 2k \frac{\sin \varphi \cos \varphi}{1+k \sin^2 \varphi} \sqrt{1-k^2 \sin^2 \varphi} \right],$$

where the angles φ and ψ are related by $\sin \psi = \frac{(1+k) \sin \varphi}{1+k \sin^2 \varphi}$.

Trigonometric expansions for small k and φ :

$$F(\varphi, k) = \frac{2}{\pi} \mathbf{K}(k) \varphi - \sin \varphi \cos \varphi \left(a_0 + \frac{2}{3} a_1 \sin^2 \varphi + \frac{2 \times 4}{3 \times 5} a_2 \sin^4 \varphi + \dots \right),$$

$$a_0 = \frac{2}{\pi} \mathbf{K}(k) - 1, \quad a_n = a_{n-1} - \left[\frac{(2n-1)!!}{(2n)!!} \right]^2 k^{2n};$$

$$E(\varphi, k) = \frac{2}{\pi} \mathbf{E}(k) \varphi - \sin \varphi \cos \varphi \left(b_0 + \frac{2}{3} b_1 \sin^2 \varphi + \frac{2 \times 4}{3 \times 5} b_2 \sin^4 \varphi + \dots \right),$$

$$b_0 = 1 - \frac{2}{\pi} \mathbf{E}(k), \quad b_n = b_{n-1} - \left[\frac{(2n-1)!!}{(2n)!!} \right]^2 \frac{k^{2n}}{2n-1}.$$

Trigonometric expansions for $k \rightarrow 1$:

$$F(\varphi, k) = \frac{2}{\pi} \mathbf{K}'(k) \ln \tan \left(\frac{\varphi}{2} + \frac{\pi}{4} \right) - \frac{\tan \varphi}{\cos \varphi} \left(a'_0 - \frac{2}{3} a'_1 \tan^2 \varphi + \frac{2 \times 4}{3 \times 5} a'_2 \tan^4 \varphi - \dots \right),$$

$$a'_0 = \frac{2}{\pi} \mathbf{K}'(k) - 1, \quad a'_n = a'_{n-1} - \left[\frac{(2n-1)!!}{(2n)!!} \right]^2 (k')^{2n};$$

$$E(\varphi, k) = \frac{2}{\pi} \mathbf{E}'(k) \ln \tan \left(\frac{\varphi}{2} + \frac{\pi}{4} \right) + \frac{\tan \varphi}{\cos \varphi} \left(b'_0 - \frac{2}{3} b'_1 \tan^2 \varphi + \frac{2 \times 4}{3 \times 5} b'_2 \tan^4 \varphi - \dots \right),$$

$$b'_0 = \frac{2}{\pi} \mathbf{E}'(k) - 1, \quad b'_n = b'_{n-1} - \left[\frac{(2n-1)!!}{(2n)!!} \right]^2 \frac{(k')^{2n}}{2n-1}.$$

11.14. Elliptic Functions

An *elliptic function* is a function that is the inverse of an elliptic integral. An elliptic function is a doubly periodic meromorphic function of a complex variable. All its periods can be written in the form $2m\omega_1 + 2n\omega_2$ with integer m and n , where ω_1 and ω_2 are a pair of (primitive) half-periods. The ratio $\tau = \omega_2/\omega_1$ is a complex quantity that may be considered to have a positive imaginary part, $\text{Im } \tau > 0$.

Throughout the rest of this section, the following brief notation will be used: $\mathbf{K} = \mathbf{K}(k)$ and $\mathbf{K}' = \mathbf{K}(k')$ are complete elliptic integrals with $k' = \sqrt{1-k^2}$.

11.14-1. Jacobi Elliptic Functions.

When the upper limit φ of the incomplete elliptic integral of the first kind

$$u = \int_0^\varphi \frac{d\alpha}{\sqrt{1 - k^2 \sin^2 \alpha}} = F(\varphi, k)$$

is treated as a function of u , the following notation is used:

$$u = \operatorname{am} \varphi.$$

Naming: φ is the *amplitude* and u is the *argument*.

Jacobi elliptic functions:

$$\begin{aligned} \operatorname{sn} u &= \sin \varphi = \sin \operatorname{am} u && (\text{sine amplitude}), \\ \operatorname{cn} u &= \cos \varphi = \cos \operatorname{am} u && (\text{cosine amplitude}), \\ \operatorname{dn} u &= \sqrt{1 - k^2 \sin^2 \varphi} = \frac{d\varphi}{du} && (\text{delta amplitude}). \end{aligned}$$

Along with the brief notations $\operatorname{sn} u$, $\operatorname{cn} u$, $\operatorname{dn} u$, the respective full notations are also used: $\operatorname{sn}(u, k)$, $\operatorname{cn}(u, k)$, $\operatorname{dn}(u, k)$.

Simple properties:

$$\begin{aligned} \operatorname{sn}(-u) &= -\operatorname{sn} u, & \operatorname{cn}(-u) &= \operatorname{cn} u, & \operatorname{dn}(-u) &= \operatorname{dn} u; \\ \operatorname{sn}^2 u + \operatorname{cn}^2 u &= 1, & k^2 \operatorname{sn}^2 u + \operatorname{dn}^2 u &= 1, & \operatorname{dn}^2 u - k^2 \operatorname{cn}^2 u &= 1 - k^2, \end{aligned}$$

where $i^2 = -1$.

Jacobi functions for special values of the modulus ($k = 0$ and $k = 1$):

$$\begin{aligned} \operatorname{sn}(u, 0) &= \sin u, & \operatorname{cn}(u, 0) &= \cos u, & \operatorname{dn}(u, 0) &= 1; \\ \operatorname{sn}(u, 1) &= \tanh u, & \operatorname{cn}(u, 1) &= \frac{1}{\cosh u}, & \operatorname{dn}(u, 1) &= \frac{1}{\cosh u}. \end{aligned}$$

Jacobi functions for special values of the argument:

$$\begin{aligned} \operatorname{sn}\left(\frac{1}{2} \mathbf{K}, k\right) &= \frac{1}{\sqrt{1 + k'}}, & \operatorname{cn}\left(\frac{1}{2} \mathbf{K}, k\right) &= \sqrt{\frac{k'}{1 + k'}}, & \operatorname{dn}\left(\frac{1}{2} \mathbf{K}, k\right) &= \sqrt{k'}; \\ \operatorname{sn}(\mathbf{K}, k) &= 1, & \operatorname{cn}(\mathbf{K}, k) &= 0, & \operatorname{dn}(\mathbf{K}, k) &= k'. \end{aligned}$$

Reduction formulas:

$$\begin{aligned} \operatorname{sn}(u \pm \mathbf{K}) &= \pm \frac{\operatorname{cn} u}{\operatorname{dn} u}, & \operatorname{cn}(u \pm \mathbf{K}) &= \mp k' \frac{\operatorname{sn} u}{\operatorname{dn} u}, & \operatorname{dn}(u \pm \mathbf{K}) &= \frac{k'}{\operatorname{dn} u}; \\ \operatorname{sn}(u \pm 2\mathbf{K}) &= -\operatorname{sn} u, & \operatorname{cn}(u \pm 2\mathbf{K}) &= -\operatorname{cn} u, & \operatorname{dn}(u \pm 2\mathbf{K}) &= \operatorname{dn} u; \\ \operatorname{sn}(u + i\mathbf{K}') &= \frac{1}{k \operatorname{sn} u}, & \operatorname{cn}(u + i\mathbf{K}') &= -\frac{i}{k} \frac{\operatorname{dn} u}{\operatorname{sn} u}, & \operatorname{dn}(u + i\mathbf{K}') &= -i \frac{\operatorname{cn} u}{\operatorname{sn} u}; \\ \operatorname{sn}(u + 2i\mathbf{K}') &= \operatorname{sn} u, & \operatorname{cn}(u + 2i\mathbf{K}') &= -\operatorname{cn} u, & \operatorname{dn}(u + 2i\mathbf{K}') &= -\operatorname{dn} u; \\ \operatorname{sn}(u + \mathbf{K} + i\mathbf{K}') &= \frac{\operatorname{dn} u}{k \operatorname{cn} u}, & \operatorname{cn}(u + \mathbf{K} + i\mathbf{K}') &= -i \frac{k'}{k \operatorname{cn} u}, & \operatorname{dn}(u + \mathbf{K} + i\mathbf{K}') &= ik' \frac{\operatorname{sn} u}{\operatorname{cn} u}; \\ \operatorname{sn}(u + 2\mathbf{K} + 2i\mathbf{K}') &= -\operatorname{sn} u, & \operatorname{cn}(u + 2\mathbf{K} + 2i\mathbf{K}') &= \operatorname{cn} u, & \operatorname{dn}(u + 2\mathbf{K} + 2i\mathbf{K}') &= -\operatorname{dn} u. \end{aligned}$$

Periods, zeros, poles, and residues (see Table 4).

TABLE 4
Periods, zeros, poles, and residues of the Jacobian elliptic functions
($m, n = 0, \pm 1, \pm 2, \dots; i^2 = -1$)

Functions	Periods	Zeros	Poles	Residues
$\operatorname{sn} u$	$4m\mathbf{K} + 2n\mathbf{K}'i$	$2m\mathbf{K} + 2n\mathbf{K}'i$	$2m\mathbf{K} + (2n+1)\mathbf{K}'i$	$(-1)^m \frac{1}{k}$
$\operatorname{cn} u$	$(4m+2n)\mathbf{K} + 2n\mathbf{K}'i$	$(2m+1)\mathbf{K} + 2n\mathbf{K}'i$	$2m\mathbf{K} + (2n+1)\mathbf{K}'i$	$(-1)^{m-1} \frac{i}{k}$
$\operatorname{dn} u$	$2m\mathbf{K} + 4n\mathbf{K}'i$	$(2m+1)\mathbf{K} + (2n+1)\mathbf{K}'i$	$2m\mathbf{K} + (2n+1)\mathbf{K}'i$	$(-1)^{n-1}i$

Double-argument formulas:

$$\begin{aligned}\operatorname{sn}(2u) &= \frac{2 \operatorname{sn} u \operatorname{cn} u \operatorname{dn} u}{1 - k^2 \operatorname{sn}^4 u} = \frac{2 \operatorname{sn} u \operatorname{cn} u \operatorname{dn} u}{\operatorname{cn}^2 u + \operatorname{sn}^2 u \operatorname{dn}^2 u}, \\ \operatorname{cn}(2u) &= \frac{\operatorname{cn}^2 u - \operatorname{sn}^2 u \operatorname{dn}^2 u}{1 - k^2 \operatorname{sn}^4 u} = \frac{\operatorname{cn}^2 u - \operatorname{sn}^2 u \operatorname{dn}^2 u}{\operatorname{cn}^2 u + \operatorname{sn}^2 u \operatorname{dn}^2 u}, \\ \operatorname{dn}(2u) &= \frac{\operatorname{dn}^2 u - k^2 \operatorname{sn}^2 u \operatorname{cn}^2 u}{1 - k^2 \operatorname{sn}^4 u} = \frac{\operatorname{dn}^2 u + \operatorname{cn}^2 u (\operatorname{dn}^2 u - 1)}{\operatorname{dn}^2 u - \operatorname{cn}^2 u (\operatorname{dn}^2 u - 1)}.\end{aligned}$$

Half-argument formulas:

$$\begin{aligned}\operatorname{sn}^2 \frac{u}{2} &= \frac{1}{k^2} \frac{1 - \operatorname{dn} u}{1 + \operatorname{cn} u} = \frac{1 - \operatorname{cn} u}{1 + \operatorname{dn} u}, \\ \operatorname{cn}^2 \frac{u}{2} &= \frac{\operatorname{cn} u + \operatorname{dn} u}{1 + \operatorname{dn} u} = \frac{1 - k^2}{k^2} \frac{1 - \operatorname{dn} u}{\operatorname{dn} u - \operatorname{cn} u}, \\ \operatorname{dn}^2 \frac{u}{2} &= \frac{\operatorname{cn} u + \operatorname{dn} u}{1 + \operatorname{cn} u} = (1 - k^2) \frac{1 - \operatorname{cn} u}{\operatorname{dn} u - \operatorname{cn} u}.\end{aligned}$$

Argument addition formulas:

$$\begin{aligned}\operatorname{sn}(u \pm v) &= \frac{\operatorname{sn} u \operatorname{cn} v \operatorname{dn} v \pm \operatorname{sn} v \operatorname{cn} u \operatorname{dn} u}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}, \\ \operatorname{cn}(u \pm v) &= \frac{\operatorname{cn} u \operatorname{cn} v \mp \operatorname{sn} u \operatorname{sn} v \operatorname{dn} u \operatorname{dn} v}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}, \\ \operatorname{dn}(u \pm v) &= \frac{\operatorname{dn} u \operatorname{dn} v \mp k^2 \operatorname{sn} u \operatorname{sn} v \operatorname{cn} u \operatorname{cn} v}{1 - k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}.\end{aligned}$$

Table 5 presents conversion formulas for Jacobi elliptic functions. If $k > 1$, then $k_1 = 1/k < 1$. Elliptic functions with real modulus can be reduced, using the first set of conversion formulas, to elliptic functions with a modulus lying between 0 and 1.

Descending Landen transformations (Gauss's transformations):

$$\operatorname{sn}(u, k) = \frac{(1 + \mu) \operatorname{sn}(v, \mu^2)}{1 + \mu \operatorname{sn}^2(v, \mu^2)}, \quad \operatorname{cn}(u, k) = \frac{\operatorname{cn}(v, \mu^2) \operatorname{dn}(v, \mu^2)}{1 + \mu \operatorname{sn}^2(v, \mu^2)}, \quad \operatorname{dn}(u, k) = \frac{\operatorname{dn}^2(v, \mu^2) + \mu - 1}{1 + \mu - \operatorname{dn}^2(v, \mu^2)},$$

where

$$\mu = \left| \frac{1 - k'}{1 + k'} \right|, \quad v = \frac{u}{1 + \mu}.$$

TABLE 5

Conversion formulas for Jacobi elliptic functions. Full notation is used: $\operatorname{sn}(u, k)$, $\operatorname{cn}(u, k)$, $\operatorname{dn}(u, k)$

u_1	k_1	$\operatorname{sn}(u_1, k_1)$	$\operatorname{cn}(u_1, k_1)$	$\operatorname{dn}(u_1, k_1)$
ku	$\frac{1}{k}$	$k \operatorname{sn}(u, k)$	$\operatorname{dn}(u, k)$	$\operatorname{cn}(u, k)$
iu	k'	$i \frac{\operatorname{sn}(u, k)}{\operatorname{cn}(u, k)}$	$\frac{1}{\operatorname{cn}(u, k)}$	$\frac{\operatorname{dn}(u, k)}{\operatorname{cn}(u, k)}$
$k'u$	$i \frac{k}{k'}$	$k' \frac{\operatorname{sn}(u, k)}{\operatorname{dn}(u, k)}$	$\frac{\operatorname{cn}(u, k)}{\operatorname{dn}(u, k)}$	$\frac{1}{\operatorname{dn}(u, k)}$
iku	$i \frac{k'}{k}$	$ik \frac{\operatorname{sn}(u, k)}{\operatorname{dn}(u, k)}$	$\frac{1}{\operatorname{dn}(u, k)}$	$\frac{\operatorname{cn}(u, k)}{\operatorname{dn}(u, k)}$
$ik'u$	$\frac{1}{k'}$	$ik' \frac{\operatorname{sn}(u, k)}{\operatorname{cn}(u, k)}$	$\frac{\operatorname{dn}(u, k)}{\operatorname{cn}(u, k)}$	$\frac{1}{\operatorname{cn}(u, k)}$
$(1+k)u$	$\frac{2\sqrt{k}}{1+k}$	$\frac{(1+k) \operatorname{sn}(u, k)}{1+k \operatorname{sn}^2(u, k)}$	$\frac{\operatorname{cn}(u, k) \operatorname{dn}(u, k)}{1+k \operatorname{sn}^2(u, k)}$	$\frac{1-k \operatorname{sn}^2(u, k)}{1+k \operatorname{sn}^2(u, k)}$
$(1+k')u$	$\frac{1-k'}{1+k'}$	$\frac{(1+k') \operatorname{sn}(u, k) \operatorname{cn}(u, k)}{\operatorname{dn}(u, k)}$	$\frac{1-(1+k') \operatorname{sn}^2(u, k)}{\operatorname{dn}(u, k)}$	$\frac{1-(1-k') \operatorname{sn}^2(u, k)}{\operatorname{dn}(u, k)}$

Ascending Landen transformations:

$$\operatorname{sn}(u, k) = (1+\sigma) \frac{\operatorname{sn}(v, \mu) \operatorname{cn}(v, \mu)}{\operatorname{dn}(v, \mu)}, \quad \operatorname{cn}(u, k) = \frac{1+\sigma}{\mu} \frac{\operatorname{dn}^2(v, \mu) - \sigma}{\operatorname{dn}(v, \mu)}, \quad \operatorname{dn}(u, k) = \frac{1-\sigma}{\mu} \frac{\operatorname{dn}^2(v, \mu) + \sigma}{\operatorname{dn}(v, \mu)},$$

where

$$\mu = \frac{4k}{(1+k)^2}, \quad \sigma = \left| \frac{1-k}{1+k} \right|, \quad v = \frac{u}{1+\sigma}.$$

Representation Jacobi functions in the form of power series in u :

$$\operatorname{sn} u = u - \frac{1}{3!}(1+k^2)u^3 + \frac{1}{5!}(1+14k^2+k^4)u^5 - \frac{1}{7!}(1+135k^2+135k^4+k^6)u^7 + \dots,$$

$$\operatorname{cn} u = 1 - \frac{1}{2!}u^2 + \frac{1}{4!}(1+4k^2)u^4 - \frac{1}{6!}(1+44k^2+16k^4)u^6 + \dots,$$

$$\operatorname{dn} u = 1 - \frac{1}{2!}k^2u^2 + \frac{1}{4!}k^2(4+k^2)u^4 - \frac{1}{6!}k^2(16+44k^2+k^4)u^6 + \dots,$$

$$\operatorname{am} u = u - \frac{1}{3!}k^2u^3 + \frac{1}{5!}k^2(4+k^2)u^5 - \frac{1}{7!}k^2(16+44k^2+k^4)u^7 + \dots.$$

These functions converge for $|u| < |\mathbf{K}(k')|$.

Representation Jacobi functions in the form of trigonometric series:

$$\operatorname{sn} u = \frac{2\pi}{k\mathbf{K}\sqrt{q}} \sum_{n=1}^{\infty} \frac{q^n}{1-q^{2n-1}} \sin \left[(2n-1) \frac{\pi u}{2\mathbf{K}} \right],$$

$$\operatorname{cn} u = \frac{2\pi}{k\mathbf{K}\sqrt{q}} \sum_{n=1}^{\infty} \frac{q^n}{1+q^{2n-1}} \cos \left[(2n-1) \frac{\pi u}{2\mathbf{K}} \right],$$

$$\operatorname{dn} u = \frac{\pi}{2\mathbf{K}} + \frac{2\pi}{\mathbf{K}} \sum_{n=1}^{\infty} \frac{q^n}{1+q^{2n}} \cos \left(\frac{n\pi u}{\mathbf{K}} \right),$$

$$\operatorname{am} u = \frac{\pi u}{2K} + 2 \sum_{n=1}^{\infty} \frac{1}{n} \frac{q^n}{1+q^{2n}} \sin\left(\frac{n\pi u}{K}\right),$$

where $q = \exp(-\pi K'/K)$, $K = K(k)$, $K' = K(k')$, and $k' = \sqrt{1-k^2}$.

Derivatives:

$$\frac{d}{du} \operatorname{sn} u = \operatorname{cn} u \operatorname{dn} u, \quad \frac{d}{du} \operatorname{cn} u = -\operatorname{sn} u \operatorname{dn} u, \quad \frac{d}{du} \operatorname{dn} u = -k^2 \operatorname{sn} u \operatorname{cn} u.$$

Integrals:

$$\begin{aligned} \int \operatorname{sn} u \, du &= \frac{1}{k} \ln(\operatorname{dn} u - k \operatorname{cn} u) = -\frac{1}{k} \ln(\operatorname{dn} u + k \operatorname{cn} u), \\ \int \operatorname{cn} u \, du &= \frac{1}{k} \arccos(\operatorname{dn} u) = \frac{1}{k} \arcsin(k \operatorname{sn} u), \\ \int \operatorname{dn} u \, du &= \arcsin(\operatorname{sn} u) = \operatorname{am} u. \end{aligned}$$

The arbitrary additive constant C in the integrals is omitted.

11.14-2. Weierstrass Elliptic Function.

The *Weierstrass elliptic function* (or *Weierstrass \wp -function*) is defined as

$$\wp(z) = \wp(z|\omega_1, \omega_2) = \frac{1}{z^2} + \sum_{m,n} \left[\frac{1}{(z-2m\omega_1-2n\omega_2)^2} - \frac{1}{(2m\omega_1+2n\omega_2)^2} \right],$$

where the summation is assumed over all integer m and n , except for $m = n = 0$. This function is a complex, double periodic function of a complex variable z with periods $2\omega_1$ and $2\omega_2$:

$$\begin{aligned} \wp(-z) &= \wp(z), \\ \wp(z + 2m\omega_1 + 2n\omega_2) &= \wp(z), \end{aligned}$$

where $m, n = 0, \pm 1, \pm 2, \dots$ and $\operatorname{Im}(\omega_2/\omega_1) \neq 0$. The series defining the Weierstrass \wp -function converges everywhere except for second-order poles located at $z_{mn} = 2m\omega_1 + 2n\omega_2$.

Argument addition formula:

$$\wp(z_1 + z_2) = -\wp(z_1) - \wp(z_2) + \frac{1}{4} \left[\frac{\wp'(z_1) - \wp'(z_2)}{\wp(z_1) - \wp(z_2)} \right]^2.$$

The Weierstrass function $\wp = \wp(z, g_2, g_3) = \wp(z|\omega_1, \omega_2)$ is defined implicitly by the elliptic integral:

$$z = \int_{\wp}^{\infty} \frac{dt}{\sqrt{4t^3 - g_2t - g_3}} = \int_{\wp}^{\infty} \frac{dt}{2\sqrt{(t-e_1)(t-e_2)(t-e_3)}}.$$

The parameters g_2 and g_3 are known as the *invariants*.

The parameters e_1, e_2, e_3 , which are the roots of the cubic equation $4z^3 - g_2z - g_3 = 0$, are related to the half-periods ω_1, ω_2 and invariants g_2, g_3 by

$$\begin{aligned} e_1 &= \wp(\omega_1), & e_2 &= \wp(\omega_1 + \omega_2), & e_3 &= \wp(\omega_2), \\ e_1 + e_2 + e_3 &= 0, & e_1e_2 + e_1e_3 + e_2e_3 &= -\frac{1}{4}g_2, & e_1e_2e_3 &= \frac{1}{4}g_3. \end{aligned}$$

Homogeneity property:

$$\wp(z, g_2, g_3) = \lambda^2 \wp(\lambda z, \lambda^{-4} g_2, \lambda^{-6} g_3).$$

The Weierstrass \wp -function can be expanded into a Laurent series:

$$\wp(z) = \frac{1}{z^2} + \frac{g_2}{20} z^2 + \frac{g_3}{28} z^4 + \frac{g_2^2}{1200} z^6 + \frac{3g_2g_3}{6160} z^8 + \dots = \frac{1}{z^2} + \sum_{k=2}^{\infty} a_k z^{2k-2},$$

$$a_k = \frac{3}{(k-3)(2k+1)} \sum_{m=2}^{k-2} a_m a_{k-m} \quad \text{for } k \geq 4, \quad 0 < |z| < \min(|\omega_1|, |\omega_2|).$$

The Weierstrass \wp -function satisfies the first-order and second-order nonlinear differential equations:

$$(\wp'_z)^2 = 4\wp^3 - g_2\wp - g_3,$$

$$\wp''_{zz} = 6\wp^2 - \frac{1}{2}g_2.$$

Direct and inverse representations of the Weierstrass elliptic function via Jacobi elliptic functions:

$$\wp(z) = e_1 + (e_1 - e_3) \frac{\text{cn}^2 w}{\text{sn}^2 w} = e_2 + (e_1 - e_3) \frac{\text{dn}^2 w}{\text{sn}^2 w} = e_3 + \frac{e_1 - e_3}{\text{sn}^2 w};$$

$$\text{sn } w = \sqrt{\frac{e_1 - e_3}{\wp(z) - e_3}}, \quad \text{cn } w = \sqrt{\frac{\wp(z) - e_1}{\wp(z) - e_3}}, \quad \text{dn } w = \sqrt{\frac{\wp(z) - e_2}{\wp(z) - e_3}};$$

$$w = z\sqrt{e_1 - e_3} = \mathbf{K} z / \omega_1.$$

The parameters are related by

$$k = \sqrt{\frac{e_2 - e_3}{e_1 - e_3}}, \quad k' = \sqrt{\frac{e_1 - e_2}{e_1 - e_3}}, \quad \mathbf{K} = \omega_1 \sqrt{e_1 - e_3}, \quad i \mathbf{K}' = \omega_2 \sqrt{e_1 - e_3}.$$

11.15. Jacobi Theta Functions

11.15-1. Series Representation of the Jacobi Theta Functions. Simplest Properties.

The *Jacobi theta functions* (or *elliptic theta functions*) are defined by the following series:

$$\vartheta_1(v) = \vartheta_1(v, q) = \vartheta_1(v|\tau) = 2 \sum_{n=0}^{\infty} (-1)^n q^{(n+1/2)^2} \sin[(2n+1)\pi v] = i \sum_{n=-\infty}^{\infty} (-1)^n q^{(n-1/2)^2} e^{i\pi(2n-1)v},$$

$$\vartheta_2(v) = \vartheta_2(v, q) = \vartheta_2(v|\tau) = 2 \sum_{n=0}^{\infty} q^{(n+1/2)^2} \cos[(2n+1)\pi v] = \sum_{n=-\infty}^{\infty} q^{(n-1/2)^2} e^{i\pi(2n-1)v},$$

$$\vartheta_3(v) = \vartheta_3(v, q) = \vartheta_3(v|\tau) = 1 + 2 \sum_{n=0}^{\infty} q^{n^2} \cos(2n\pi v) = \sum_{n=-\infty}^{\infty} q^{n^2} e^{2i\pi n v},$$

$$\vartheta_4(v) = \vartheta_4(v, q) = \vartheta_4(v|\tau) = 1 + 2 \sum_{n=0}^{\infty} (-1)^n q^{n^2} \cos(2n\pi v) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2} e^{2i\pi n v},$$

where v is a complex variable and $q = e^{i\pi\tau}$ is a complex parameter (τ has a positive imaginary part).

The Jacobi theta functions are periodic entire functions that possess the following properties:

$$\begin{aligned} \vartheta_1(v) & \text{ odd, has period } 2, \text{ vanishes at } v = m + n\tau; \\ \vartheta_2(v) & \text{ even, has period } 2, \text{ vanishes at } v = m + n\tau + \frac{1}{2}; \\ \vartheta_3(v) & \text{ even, has period } 1, \text{ vanishes at } v = m + (n + \frac{1}{2})\tau + \frac{1}{2}; \\ \vartheta_4(v) & \text{ even, has period } 1, \text{ vanishes at } v = m + (n + \frac{1}{2})\tau. \end{aligned}$$

Here $m, n = 0, \pm 1, \pm 2, \dots$

Remark. The theta functions are not elliptic functions. The very good convergence of their series allows the computation of various elliptic integrals and elliptic functions using the relations given above in Supplement 11.15-1.

11.15-2. Various Relations and Formulas. Connection with Jacobi Elliptic Functions.

Linear relations (first set):

$$\begin{aligned} \vartheta_1\left(v + \frac{1}{2}\right) &= \vartheta_2(v), & \vartheta_2\left(v + \frac{1}{2}\right) &= -\vartheta_1(v), \\ \vartheta_3\left(v + \frac{1}{2}\right) &= \vartheta_4(v), & \vartheta_4\left(v + \frac{1}{2}\right) &= \vartheta_3(v), \\ \vartheta_1\left(v + \frac{\tau}{2}\right) &= ie^{-i\pi\left(v + \frac{\tau}{4}\right)}\vartheta_4(v), & \vartheta_2\left(v + \frac{\tau}{2}\right) &= e^{-i\pi\left(v + \frac{\tau}{4}\right)}\vartheta_3(v), \\ \vartheta_3\left(v + \frac{\tau}{2}\right) &= e^{-i\pi\left(v + \frac{\tau}{4}\right)}\vartheta_2(v), & \vartheta_4\left(v + \frac{\tau}{2}\right) &= ie^{-i\pi\left(v + \frac{\tau}{4}\right)}\vartheta_1(v). \end{aligned}$$

Linear relations (second set):

$$\begin{aligned} \vartheta_1(v|\tau + 1) &= e^{i\pi/4}\vartheta_1(v|\tau), & \vartheta_2(v|\tau + 1) &= e^{i\pi/4}\vartheta_2(v|\tau), \\ \vartheta_3(v|\tau + 1) &= \vartheta_4(v|\tau), & \vartheta_4(v|\tau + 1) &= \vartheta_3(v|\tau), \\ \vartheta_1\left(\frac{v}{\tau}\middle|\frac{1}{\tau}\right) &= \frac{1}{i}\sqrt{\frac{\tau}{i}}e^{i\pi v^2/\tau}\vartheta_1(v|\tau), & \vartheta_2\left(\frac{v}{\tau}\middle|\frac{1}{\tau}\right) &= \sqrt{\frac{\tau}{i}}e^{i\pi v^2/\tau}\vartheta_4(v|\tau), \\ \vartheta_3\left(\frac{v}{\tau}\middle|\frac{1}{\tau}\right) &= \sqrt{\frac{\tau}{i}}e^{i\pi v^2/\tau}\vartheta_3(v|\tau), & \vartheta_4\left(\frac{v}{\tau}\middle|\frac{1}{\tau}\right) &= \sqrt{\frac{\tau}{i}}e^{i\pi v^2/\tau}\vartheta_2(v|\tau). \end{aligned}$$

Quadratic relations:

$$\begin{aligned} \vartheta_1^2(v)\vartheta_2^2(0) &= \vartheta_4^2(v)\vartheta_3^2(0) - \vartheta_3^2(v)\vartheta_4^2(0), \\ \vartheta_1^2(v)\vartheta_3^2(0) &= \vartheta_4^2(v)\vartheta_2^2(0) - \vartheta_2^2(v)\vartheta_4^2(0), \\ \vartheta_1^2(v)\vartheta_4^2(0) &= \vartheta_3^2(v)\vartheta_2^2(0) - \vartheta_2^2(v)\vartheta_3^2(0), \\ \vartheta_4^2(v)\vartheta_4^2(0) &= \vartheta_3^2(v)\vartheta_3^2(0) - \vartheta_2^2(v)\vartheta_2^2(0). \end{aligned}$$

Representation of the theta functions in the form of infinite products:

$$\begin{aligned} \vartheta_1(v) &= 2q_0q^{1/4}\sin(\pi v)\prod_{n=1}^{\infty}[1 - 2q^{2n}\cos(2\pi v) + q^{4n}], \\ \vartheta_2(v) &= 2q_0q^{1/4}\cos(\pi v)\prod_{n=1}^{\infty}[1 + 2q^{2n}\cos(2\pi v) + q^{4n}], \\ \vartheta_3(v) &= q_0\prod_{n=1}^{\infty}[1 + 2q^{2n-1}\cos(2\pi v) + q^{4n-2}], \\ \vartheta_4(v) &= q_0\prod_{n=1}^{\infty}[1 - 2q^{2n-1}\cos(2\pi v) + q^{4n-2}], \end{aligned}$$

where $q_0 = \prod_{n=1}^{\infty}(1 - q^{2n})$.

Representations of Jacobi elliptic functions in terms of the theta functions:

$$\operatorname{sn} w = \frac{\vartheta_3(0)\vartheta_1(v)}{\vartheta_2(0)\vartheta_4(v)}, \quad \operatorname{cn} w = \frac{\vartheta_4(0)\vartheta_2(v)}{\vartheta_2(0)\vartheta_4(v)}, \quad \operatorname{dn} w = \frac{\vartheta_4(0)\vartheta_3(v)}{\vartheta_3(0)\vartheta_4(v)}, \quad w = 2\mathbf{K}v.$$

The parameters are related by

$$k = \frac{\vartheta_2^2(0)}{\vartheta_3^2(0)}, \quad k' = \frac{\vartheta_4^2(0)}{\vartheta_3^2(0)}, \quad \mathbf{K} = \frac{\pi}{2}\vartheta_3^2(0), \quad \mathbf{K}' = -i\tau\mathbf{K}.$$

TABLE 6

The Mathieu functions $ce_n = ce_n(x, q)$ and $se_n = se_n(x, q)$ (for odd n , functions ce_n and se_n are 2π -periodic, and for even n , they are π -periodic); definite eigenvalues $a = a_n(q)$ and $a = b_n(q)$ correspond to each value of parameter q

Mathieu functions	Recurrence relations for coefficients	Normalization conditions
$ce_{2n} = \sum_{m=0}^{\infty} A_{2m}^{2n} \cos 2mx$	$qA_2^{2n} = a_{2n}A_0^{2n};$ $qA_4^{2n} = (a_{2n}-4)A_2^{2n} - 2qA_0^{2n};$ $qA_{2m+2}^{2n} = (a_{2n}-4m^2)A_{2m}^{2n}$ $- qA_{2m-2}^{2n}, \quad m \geq 2$	$(A_0^{2n})^2 + \sum_{m=0}^{\infty} (A_{2m}^{2n})^2$ $= \begin{cases} 2 & \text{if } n=0 \\ 1 & \text{if } n \geq 1 \end{cases}$
$ce_{2n+1} = \sum_{m=0}^{\infty} A_{2m+1}^{2n+1} \cos(2m+1)x$	$qA_3^{2n+1} = (a_{2n+1}-1-q)A_1^{2n+1};$ $qA_{2m+3}^{2n+1} = [a_{2n+1}-(2m+1)^2]A_{2m+1}^{2n+1}$ $- qA_{2m-1}^{2n+1}, \quad m \geq 1$	$\sum_{m=0}^{\infty} (A_{2m+1}^{2n+1})^2 = 1$
$se_{2n} = \sum_{m=0}^{\infty} B_{2m}^{2n} \sin 2mx,$ $se_0 = 0$	$qB_4^{2n} = (b_{2n}-4)B_2^{2n};$ $qB_{2m+2}^{2n} = (b_{2n}-4m^2)B_{2m}^{2n}$ $- qB_{2m-2}^{2n}, \quad m \geq 2$	$\sum_{m=0}^{\infty} (B_{2m}^{2n})^2 = 1$
$se_{2n+1} = \sum_{m=0}^{\infty} B_{2m+1}^{2n+1} \sin(2m+1)x$	$qB_3^{2n+1} = (b_{2n+1}-1-q)B_1^{2n+1};$ $qB_{2m+3}^{2n+1} = [b_{2n+1}-(2m+1)^2]B_{2m+1}^{2n+1}$ $- qB_{2m-1}^{2n+1}, \quad m \geq 1$	$\sum_{m=0}^{\infty} (B_{2m+1}^{2n+1})^2 = 1$

11.16. Mathieu Functions and Modified Mathieu Functions

11.16-1. Mathieu Functions.

The Mathieu functions $ce_n(x, q)$ and $se_n(x, q)$ are periodical solutions of the *Mathieu equation*

$$y''_{xx} + (a - 2q \cos 2x)y = 0.$$

Such solutions exist for definite values of parameters a and q (those values of a are referred to as eigenvalues). The Mathieu functions are listed in Table 6.

The Mathieu functions possess the following properties:

$$ce_{2n}(x, -q) = (-1)^n ce_{2n}\left(\frac{\pi}{2} - x, q\right), \quad ce_{2n+1}(x, -q) = (-1)^n se_{2n+1}\left(\frac{\pi}{2} - x, q\right),$$

$$se_{2n}(x, -q) = (-1)^{n-1} se_{2n}\left(\frac{\pi}{2} - x, q\right), \quad se_{2n+1}(x, -q) = (-1)^n ce_{2n+1}\left(\frac{\pi}{2} - x, q\right).$$

Selecting sufficiently large number m and omitting the term with the maximum number in the recurrence relations (indicated in Table 6), we can obtain approximate relations for eigenvalues a_n (or b_n) with respect to parameter q . Then, equating the determinant of the corresponding homogeneous linear system of equations for coefficients A_m^n (or B_m^n) to zero, we obtain an algebraic equation for finding $a_n(q)$ (or $b_n(q)$).

For fixed real $q \neq 0$, eigenvalues a_n and b_n are all real and different, while

$$\begin{aligned} \text{if } q > 0 \text{ then } & a_0 < b_1 < a_1 < b_2 < a_2 < \dots; \\ \text{if } q < 0 \text{ then } & a_0 < a_1 < b_1 < b_2 < a_2 < a_3 < b_3 < b_4 < \dots. \end{aligned}$$

The eigenvalues possess the properties

$$a_{2n}(-q) = a_{2n}(q), \quad b_{2n}(-q) = b_{2n}(q), \quad a_{2n+1}(-q) = b_{2n+1}(q).$$

Tables of the eigenvalues $a_n = a_n(q)$ and $b_n = b_n(q)$ can be found in Abramowitz and Stegun (1964, chap. 20).

The solution of the Mathieu equation corresponding to eigenvalue a_n (or b_n) has n zeros on the interval $0 \leq x < \pi$ (q is a real number).

Listed below are two leading terms of asymptotic expansions of the Mathieu functions $\text{ce}_n(x, q)$ and $\text{se}_n(x, q)$, as well as of the corresponding eigenvalues $a_n(q)$ and $b_n(q)$, as $q \rightarrow 0$:

$$\text{ce}_0(x, q) = \frac{1}{\sqrt{2}} \left(1 - \frac{q}{2} \cos 2x \right), \quad a_0(q) = -\frac{q^2}{2} + \frac{7q^4}{128};$$

$$\text{ce}_1(x, q) = \cos x - \frac{q}{8} \cos 3x, \quad a_1(q) = 1 + q;$$

$$\text{ce}_2(x, q) = \cos 2x + \frac{q}{4} \left(1 - \frac{\cos 4x}{3} \right), \quad a_2(q) = 4 + \frac{5q^2}{12};$$

$$\text{ce}_n(x, q) = \cos nx + \frac{q}{4} \left[\frac{\cos(n+2)x}{n+1} - \frac{\cos(n-2)x}{n-1} \right], \quad a_n(q) = n^2 + \frac{q^2}{2(n^2-1)} \quad (n \geq 3);$$

$$\text{se}_1(x, q) = \sin x - \frac{q}{8} \sin 3x, \quad b_1(q) = 1 - q;$$

$$\text{se}_2(x, q) = \sin 2x - q \frac{\sin 4x}{12}, \quad b_2(q) = 4 - \frac{q^2}{12};$$

$$\text{se}_n(x, q) = \sin nx - \frac{q}{4} \left[\frac{\sin(n+2)x}{n+1} - \frac{\sin(n-2)x}{n-1} \right], \quad b_n(q) = n^2 + \frac{q^2}{2(n^2-1)} \quad (n \geq 3).$$

Asymptotic results as $q \rightarrow \infty$ ($-\pi/2 < x < \pi/2$):

$$a_n(q) \approx -2q + 2(2n+1)\sqrt{q} + \frac{1}{4}(2n^2 + 2n + 1),$$

$$b_{n+1}(q) \approx -2q + 2(2n+1)\sqrt{q} + \frac{1}{4}(2n^2 + 2n + 1),$$

$$\text{ce}_n(x, q) \approx \lambda_n q^{-1/4} \cos^{-n-1} x \left[\cos^{2n+1} \xi \exp(2\sqrt{q} \sin x) + \sin^{2n+1} \xi \exp(-2\sqrt{q} \sin x) \right],$$

$$\text{se}_{n+1}(x, q) \approx \mu_{n+1} q^{-1/4} \cos^{-n-1} x \left[\cos^{2n+1} \xi \exp(2\sqrt{q} \sin x) - \sin^{2n+1} \xi \exp(-2\sqrt{q} \sin x) \right],$$

where λ_n and μ_n are some constants independent of the parameter q , and $\xi = \frac{1}{2}x + \frac{\pi}{4}$.

11.16-2. Modified Mathieu Functions.

The modified Mathieu functions $\text{Ce}_n(x, q)$ and $\text{Se}_n(x, q)$ are solutions of the *modified Mathieu equation*

$$y''_{xx} - (a - 2q \cosh 2x)y = 0,$$

with $a = a_n(q)$ and $a = b_n(q)$ being the eigenvalues of the Mathieu equation (see Supplement 11.16-1).

The modified Mathieu functions are defined as

$$\text{Ce}_{2n+p}(x, q) = \text{ce}_{2n+p}(ix, q) = \sum_{k=0}^{\infty} A_{2k+p}^{2n+p} \cosh[(2k+p)x],$$

$$\text{Se}_{2n+p}(x, q) = -i \text{se}_{2n+p}(ix, q) = \sum_{k=0}^{\infty} B_{2k+p}^{2n+p} \sinh[(2k+p)x],$$

where p may be equal to 0 and 1, and coefficients A_{2k+p}^{2n+p} and B_{2k+p}^{2n+p} are indicated in Supplement 11.16-1.

11.17. Orthogonal Polynomials

All zeros of each of the orthogonal polynomials $\mathcal{P}_n(x)$ considered in this section are real and simple. The zeros of the polynomials $\mathcal{P}_n(x)$ and $\mathcal{P}_{n+1}(x)$ are alternating.

For Legendre polynomials see Supplement 11.11-1.

11.17-1. Laguerre Polynomials and Generalized Laguerre Polynomials.

The Laguerre polynomials $L_n = L_n(x)$ satisfy the second-order linear ordinary differential equation

$$xy''_{xx} + (1-x)y'_x + ny = 0$$

and are defined by the formulas

$$L_n(x) = \frac{1}{n!} e^x \frac{d^n}{dx^n} (x^n e^{-x}) = \frac{(-1)^n}{n!} \left[x^n - n^2 x^{n-1} + \frac{n^2(n-1)^2}{2!} x^{n-2} + \dots \right].$$

The first four polynomials have the form

$$L_0(x) = 1, \quad L_1(x) = -x + 1, \quad L_2(x) = \frac{1}{2}(x^2 - 4x + 2), \quad L_3(x) = \frac{1}{6}(-x^3 + 9x^2 - 18x + 6).$$

To calculate $L_n(x)$ for $n \geq 2$, one can use the recurrence formulas

$$L_{n+1}(x) = \frac{1}{n+1} [(2n+1-x)L_n(x) - nL_{n-1}(x)].$$

The functions $L_n(x)$ form an orthonormal system on the interval $0 < x < \infty$ with weight e^{-x} :

$$\int_0^\infty e^{-x} L_n(x) L_m(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ 1 & \text{if } n = m. \end{cases}$$

The generating function is

$$\frac{1}{1-s} \exp\left(-\frac{sx}{1-s}\right) = \sum_{n=0}^{\infty} L_n(x) s^n, \quad |s| < 1.$$

The generalized Laguerre polynomials $L_n^\alpha = L_n^\alpha(x)$ ($\alpha > -1$) satisfy the equation

$$xy''_{xx} + (\alpha+1-x)y'_x + ny = 0$$

and are defined by the formulas

$$L_n^\alpha(x) = \frac{1}{n!} x^{-\alpha} e^x \frac{d^n}{dx^n} (x^{n+\alpha} e^{-x}) = \sum_{m=0}^n C_{n+\alpha}^{n-m} \frac{(-x)^m}{m!} = \sum_{m=0}^n \frac{\Gamma(n+\alpha+1)}{\Gamma(m+\alpha+1)} \frac{(-x)^m}{m!(n-m)!}.$$

Notation: $L_n^0(x) = L_n(x)$.

Special cases:

$$L_0^\alpha(x) = 1, \quad L_1^\alpha(x) = \alpha + 1 - x, \quad L_n^{-n}(x) = (-1)^n \frac{x^n}{n!}.$$

To calculate $L_n^\alpha(x)$ for $n \geq 2$, one can use the recurrence formulas

$$L_{n+1}^\alpha(x) = \frac{1}{n+1} [(2n+\alpha+1-x)L_n^\alpha(x) - (n+\alpha)L_{n-1}^\alpha(x)].$$

Other recurrence formulas:

$$L_n^\alpha(x) = L_{n-1}^\alpha(x) + L_n^{\alpha-1}(x), \quad \frac{d}{dx} L_n^\alpha(x) = -L_{n-1}^{\alpha+1}(x), \quad x \frac{d}{dx} L_n^\alpha(x) = nL_n^\alpha(x) - (n+\alpha)L_{n-1}^\alpha(x).$$

The functions $L_n^\alpha(x)$ form an orthogonal system on the interval $0 < x < \infty$ with weight $x^\alpha e^{-x}$:

$$\int_0^\infty x^\alpha e^{-x} L_n^\alpha(x) L_m^\alpha(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{\Gamma(\alpha+n+1)}{n!} & \text{if } n = m. \end{cases}$$

The generating function is

$$(1-s)^{-\alpha-1} \exp\left(-\frac{sx}{1-s}\right) = \sum_{n=0}^{\infty} L_n^\alpha(x) s^n, \quad |s| < 1.$$

11.17-2. Chebyshev Polynomials and Functions.

The *Chebyshev polynomials of the first kind* $T_n = T_n(x)$ satisfy the second-order linear ordinary differential equation

$$(1-x^2)y''_{xx} - xy'_x + n^2y = 0 \quad (1)$$

and are defined by the formulas

$$\begin{aligned} T_n(x) &= \cos(n \arccos x) = \frac{(-2)^n n!}{(2n)!} \sqrt{1-x^2} \frac{d^n}{dx^n} [(1-x^2)^{n-\frac{1}{2}}] \\ &= \frac{n}{2} \sum_{m=0}^{[n/2]} (-1)^m \frac{(n-m-1)!}{m!(n-2m)!} (2x)^{n-2m} \quad (n = 0, 1, 2, \dots), \end{aligned}$$

where $[A]$ stands for the integer part of a number A .

An alternative representation of the Chebyshev polynomials:

$$T_n(x) = \frac{(-1)^n}{(2n-1)!!} (1-x^2)^{1/2} \frac{d^n}{dx^n} (1-x^2)^{n-1/2}.$$

The first five Chebyshev polynomials of the first kind are

$$T_0(x) = 1, \quad T_1(x) = x, \quad T_2(x) = 2x^2 - 1, \quad T_3(x) = 4x^3 - 3x, \quad T_4(x) = 8x^4 - 8x^2 + 1.$$

The recurrence formulas:

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x), \quad n \geq 2.$$

The functions $T_n(x)$ form an orthogonal system on the interval $-1 < x < 1$ with weight $(1-x^2)^{-1/2}$:

$$\int_{-1}^1 \frac{T_n(x)T_m(x)}{\sqrt{1-x^2}} dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{1}{2}\pi & \text{if } n = m \neq 0, \\ \pi & \text{if } n = m = 0. \end{cases}$$

The generating function is

$$\frac{1-sx}{1-2sx+s^2} = \sum_{n=0}^{\infty} T_n(x) s^n \quad (|s| < 1).$$

The functions $T_n(x)$ have only real simple zeros, all lying on the interval $-1 < x < 1$.

The normalized Chebyshev polynomials of the first kind, $2^{1-n}T_n(x)$, deviate from zero least of all. This means that among all polynomials of degree n with the leading coefficient 1, it is the maximum of the modulus $\max_{-1 \leq x \leq 1} |2^{1-n}T_n(x)|$ that has the least value, the maximum being equal to 2^{1-n} .

The *Chebyshev polynomials of the second kind* $U_n = U_n(x)$ satisfy the second-order linear ordinary differential equation

$$(1-x^2)y''_{xx} - 3xy'_x + n(n+2)y = 0$$

and are defined by the formulas

$$\begin{aligned} U_n(x) &= \frac{\sin[(n+1)\arccos x]}{\sqrt{1-x^2}} = \frac{2^n(n+1)!}{(2n+1)!} \frac{1}{\sqrt{1-x^2}} \frac{d^n}{dx^n} (1-x^2)^{n+1/2} \\ &= \sum_{m=0}^{[n/2]} (-1)^m \frac{(n-m)!}{m!(n-2m)!} (2x)^{n-2m} \quad (n = 0, 1, 2, \dots). \end{aligned}$$

The first five Chebyshev polynomials of the second kind are

$$U_0(x) = 1, \quad U_1(x) = 2x, \quad U_2(x) = 4x^2 - 1, \quad U_3(x) = 8x^3 - 4x, \quad U_4(x) = 16x^4 - 12x^2 + 1.$$

The recurrence formulas:

$$U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x), \quad n \geq 2.$$

The generating function is

$$\frac{1}{1-2sx+s^2} = \sum_{n=0}^{\infty} U_n(x)s^n \quad (|s| < 1).$$

The Chebyshev polynomials of the first and second kind are related by

$$U_n(x) = \frac{1}{n+1} \frac{d}{dx} T_{n+1}(x).$$

The *Chebyshev functions of the second kind*,

$$\begin{aligned} U_0(x) &= \arcsin x, \\ U_n(x) &= \sin(n \arccos x) = \frac{\sqrt{1-x^2}}{n} \frac{dT_n(x)}{dx} \quad (n = 1, 2, \dots), \end{aligned}$$

just as the Chebyshev polynomials, also satisfy the differential equation (1).

The first five Chebyshev functions are

$$\begin{aligned} U_0(x) &= 0, \quad U_1(x) = \sqrt{1-x^2}, \quad U_2(x) = 2x\sqrt{1-x^2}, \\ U_3(x) &= (4x^2-1)\sqrt{1-x^2}, \quad U_5(x) = (8x^3-4x)\sqrt{1-x^2}. \end{aligned}$$

The recurrence formulas:

$$U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x), \quad n \geq 2.$$

The functions $U_n(x)$ form an orthogonal system on the interval $-1 < x < 1$ with weight $(1-x^2)^{-1/2}$:

$$\int_{-1}^1 \frac{U_n(x)U_m(x)}{\sqrt{1-x^2}} dx = \begin{cases} 0 & \text{if } n \neq m \text{ or } n = m = 0, \\ \frac{1}{2}\pi & \text{if } n = m \neq 0. \end{cases}$$

The generating function is

$$\frac{\sqrt{1-x^2}}{1-2sx+s^2} = \sum_{n=0}^{\infty} U_{n+1}(x)s^n \quad (|s| < 1).$$

11.17-3. Hermite Polynomials and Functions.

The *Hermite polynomials* $H_n = H_n(x)$ satisfy the second-order linear ordinary differential equation

$$y''_{xx} - 2xy'_x + 2ny = 0$$

and are defined by the formulas

$$H_n(x) = (-1)^n \exp(x^2) \frac{d^n}{dx^n} \exp(-x^2) = \sum_{m=0}^{[n/2]} (-1)^m \frac{n!}{m!(n-2m)!} (2x)^{n-2m},$$

where $[A]$ stands for the integer part of a number A .

The first five polynomials are

$$H_0(x) = 1, \quad H_1(x) = 2x, \quad H_2(x) = 4x^2 - 2, \quad H_3(x) = 8x^3 - 12x, \quad H_4(x) = 16x^4 - 48x^2 + 12.$$

Recurrence formulas:

$$H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x), \quad n \geq 2;$$

$$\frac{d}{dx} H_n(x) = 2nH_{n-1}(x).$$

Integral representation:

$$H_{2n}(x) = \frac{(-1)^n 2^{2n+1}}{\sqrt{\pi}} \exp(x^2) \int_0^\infty \exp(-t^2) t^{2n} \cos(2xt) dt,$$

$$H_{2n+1}(x) = \frac{(-1)^n 2^{2n+2}}{\sqrt{\pi}} \exp(x^2) \int_0^\infty \exp(-t^2) t^{2n+1} \sin(2xt) dt,$$

where $n = 0, 1, 2, \dots$

The functions $H_n(x)$ form an orthogonal system on the interval $-\infty < x < \infty$ with weight e^{-x^2} :

$$\int_{-\infty}^{\infty} \exp(-x^2) H_n(x) H_m(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \sqrt{\pi} 2^n n! & \text{if } n = m. \end{cases}$$

Generating function:

$$\exp(-s^2 + 2sx) = \sum_{n=0}^{\infty} H_n(x) \frac{s^n}{n!}.$$

Asymptotic formula as $n \rightarrow \infty$:

$$H_n(x) \approx 2^{\frac{n+1}{2}} n^{\frac{n}{2}} e^{-\frac{n}{2}} \exp(x^2) \cos\left(\sqrt{2n+1} x - \frac{1}{2}\pi n\right).$$

The *Hermite functions* $h_n(x)$ are introduced by the formula

$$h_n(x) = \exp\left(-\frac{1}{2}x^2\right) H_n(x) = (-1)^n \exp\left(\frac{1}{2}x^2\right) \frac{d^n}{dx^n} \exp(-x^2), \quad n = 0, 1, 2, \dots$$

The Hermite functions satisfy the second-order linear ordinary differential equation

$$h''_{xx} + (2n + 1 - x^2)h = 0.$$

The functions $h_n(x)$ form an orthogonal system on the interval $-\infty < x < \infty$, with

$$\int_{-\infty}^{\infty} h_n(x) h_m(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \sqrt{\pi} 2^n n! & \text{if } n = m. \end{cases}$$

11.17-4. Jacobi Polynomials.

The *Jacobi polynomials*, $P_n^{(\alpha,\beta)}(x)$, are solutions of the second-order linear ordinary differential equation

$$(1-x^2)y''_{xx} + [\beta - \alpha - (\alpha + \beta + 2)x]y'_x + n(n + \alpha + \beta + 1)y = 0$$

and are defined by the formulas

$$\begin{aligned} P_n^{(\alpha,\beta)}(x) &= \frac{(-1)^n}{2^n n!} (1-x)^{-\alpha} (1+x)^{-\beta} \frac{d^n}{dx^n} \left[(1-x)^{\alpha+n} (1+x)^{\beta+n} \right] \\ &= 2^{-n} \sum_{m=0}^n C_{n+\alpha}^m C_{n+\beta}^{n-m} (x-1)^{n-m} (x+1)^m, \end{aligned}$$

where the C_b^a are binomial coefficients.

The generating function:

$$2^{\alpha+\beta} R^{-1} (1-s+R)^{-\alpha} (1+s+R)^{-\beta} = \sum_{n=0}^{\infty} P_n^{(\alpha,\beta)}(x) s^n, \quad R = \sqrt{1-2xs+s^2}, \quad |s| < 1.$$

The Jacobi polynomials are orthogonal on the interval $-1 \leq x \leq 1$ with weight $(1-x)^\alpha (1+x)^\beta$:

$$\int_{-1}^1 (1-x)^\alpha (1+x)^\beta P_n^{(\alpha,\beta)}(x) P_m^{(\alpha,\beta)}(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{2^{\alpha+\beta+1}}{\alpha + \beta + 2n + 1} \frac{\Gamma(\alpha + n + 1) \Gamma(\beta + n + 1)}{n! \Gamma(\alpha + \beta + n + 1)} & \text{if } n = m. \end{cases}$$

For $\alpha > -1$ and $\beta > -1$, all zeros of the polynomial $P_n^{(\alpha,\beta)}(x)$ are simple and lie on the interval $-1 < x < 1$.

11.17-5. Gegenbauer Polynomials.

The *Gegenbauer polynomials* (also called *ultraspherical polynomials*), $C_n^{(\lambda)}(x)$, are solutions of the second-order linear ordinary differential equation

$$(1-x^2)y''_{xx} - (2\lambda + 1)xy'_x + n(n + 2\lambda)y = 0$$

and are defined by the formulas

$$\begin{aligned} C_n^{(\lambda)}(x) &= \frac{(-2)^n}{n!} \frac{\Gamma(n + \lambda) \Gamma(n + 2\lambda)}{\Gamma(\lambda) \Gamma(2n + 2\lambda)} (1-x^2)^{-\lambda+1/2} \frac{d^n}{dx^n} (1-x^2)^{n+\lambda-1/2} \\ &= \sum_{m=0}^{[n/2]} (-1)^m \frac{\Gamma(n-m+\lambda)}{\Gamma(\lambda) m! (n-2m)!} (2x)^{n-2m}. \end{aligned}$$

Recurrence formulas:

$$\begin{aligned} C_{n+1}^{(\lambda)}(x) &= \frac{2(n+\lambda)}{n+1} x C_n^{(\lambda)}(x) - \frac{n+2\lambda-1}{n+1} C_{n-1}^{(\lambda)}(x); \\ C_n^{(\lambda)}(-x) &= (-1)^n C_n^{(\lambda)}(x), \quad \frac{d}{dx} C_n^{(\lambda)}(x) = 2\lambda C_{n-1}^{(\lambda+1)}(x). \end{aligned}$$

The generating function:

$$\frac{1}{(1-2xs+s^2)^\lambda} = \sum_{n=0}^{\infty} C_n^{(\lambda)}(x) s^n.$$

The Gegenbauer polynomials are orthogonal on the interval $-1 \leq x \leq 1$ with weight $(1-x^2)^{\lambda-1/2}$:

$$\int_{-1}^1 (1-x^2)^{\lambda-1/2} C_n^{(\lambda)}(x) C_m^{(\lambda)}(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{\pi \Gamma(2\lambda + n)}{2^{2\lambda-1} (\lambda + n) n! \Gamma^2(\lambda)} & \text{if } n = m. \end{cases}$$

11.18. Nonorthogonal Polynomials

11.18-1. Bernoulli Polynomials.

The *Bernoulli polynomials* $B_n(x)$ are introduced by the formula

$$B_n(x) = \sum_{k=0}^n C_n^k B_k x^{n-k} \quad (n = 0, 1, 2, \dots),$$

where C_n^k are the binomial coefficients and B_n are Bernoulli numbers (see Supplement 11.1-3).

The Bernoulli polynomials can be defined using the recurrence relation

$$B_0(x) = 1, \quad \sum_{k=0}^{n-1} C_n^k B_k(x) = nx^{n-1}, \quad n = 2, 3, \dots$$

The first six Bernoulli polynomials are given by

$$\begin{aligned} B_0(x) &= 1, & B_1(x) &= x - \frac{1}{2}, & B_2(x) &= x^2 - x + \frac{1}{6}, & B_3(x) &= x^3 - \frac{3}{2}x^2 + \frac{1}{2}x, \\ B_4(x) &= x^4 - 2x^3 + x^2 - \frac{1}{30}, & B_5(x) &= x^5 - \frac{5}{2}x^4 + \frac{5}{3}x^3 - \frac{1}{6}x. \end{aligned}$$

Basic properties:

$$\begin{aligned} B_n(x+1) - B_n(x) &= nx^{n-1}, & B'_{n+1}(x) &= (n+1)B_n(x), \\ B_n(1-x) &= (-1)^n B_n(x), & (-1)^n E_n(-x) &= E_n(x) + nx^{n-1}, \end{aligned}$$

where the prime denotes a derivative with respect to x , and $n = 0, 1, \dots$

Multiplication and addition formulas:

$$\begin{aligned} B_n(mx) &= m^{n-1} \sum_{k=0}^{m-1} B_n\left(x + \frac{k}{m}\right), \\ B_n(x+y) &= \sum_{k=0}^n C_n^k B_k(x) y^{n-k}, \end{aligned}$$

where $n = 0, 1, \dots$ and $m = 1, 2, \dots$

The generating function is expressed as

$$\frac{te^{xt}}{e^t - 1} \equiv \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} \quad (|t| < 2\pi).$$

This relation may be used as a definition of the Bernoulli polynomials.

Fourier series expansions:

$$\begin{aligned} B_n(x) &= -2 \frac{n!}{(2\pi)^n} \sum_{k=1}^{\infty} \frac{\cos(2\pi kx - \frac{1}{2}\pi n)}{k^n} & (n = 1, 0 < x < 1; \quad n > 1, 0 \leq x \leq 1); \\ B_{2n-1}(x) &= 2(-1)^n \frac{(2n-1)!}{(2\pi)^{2n-1}} \sum_{k=1}^{\infty} \frac{\sin(2k\pi x)}{k^{2n-1}} & (n = 1, 0 < x < 1; \quad n > 1, 0 \leq x \leq 1); \\ B_{2n}(x) &= 2(-1)^n \frac{(2n)!}{(2\pi)^{2n}} \sum_{k=1}^{\infty} \frac{\cos(2k\pi x)}{k^{2n}} & (n = 1, 2, \dots, \quad 0 \leq x \leq 1). \end{aligned}$$

Integrals:

$$\begin{aligned} \int_a^x B_n(t) dt &= \frac{B_{n+1}(x) - B_{n+1}(a)}{n+1}, \\ \int_0^1 B_m(t) B_n(t) dt &= (-1)^{n-1} \frac{m! n!}{(m+n)!} B_{m+n}, \end{aligned}$$

where m and n are positive integers and B_n are Bernoulli numbers.

11.18-2. Euler Polynomials.

Definition:

$$E_n(x) = \sum_{k=0}^n C_n^k \frac{E_k}{2^n} \left(x - \frac{1}{2}\right)^{n-k} \quad (n = 0, 1, 2, \dots),$$

where C_n^k are the binomial coefficients and E_n are Euler numbers (see Supplement 11.1-4).

The first six Euler polynomials are given by

$$\begin{aligned} E_0(x) &= 1, & E_1(x) &= x - \frac{1}{2}, & E_2(x) &= x^2 - x, & E_3(x) &= x^3 - \frac{3}{2}x^2 + \frac{1}{4}, \\ E_4(x) &= x^4 - 2x^3 + x, & E_5(x) &= x^5 - \frac{5}{2}x^4 + \frac{5}{2}x^2 - \frac{1}{2}. \end{aligned}$$

Basic properties:

$$\begin{aligned} E_n(x+1) + E_n(x) &= 2x^n, & E'_{n+1} &= (n+1)E_n(x), \\ E_n(1-x) &= (-1)^n E_n(x), & (-1)^{n+1} E_n(-x) &= E_n(x) - 2x^n, \end{aligned}$$

where the prime denotes a derivative with respect to x , and $n = 0, 1, \dots$

Multiplication and addition formulas:

$$\begin{aligned} E_n(mx) &= m^n \sum_{k=0}^{m-1} (-1)^k E_n\left(x + \frac{k}{m}\right), & n &= 0, 1, \dots, \quad m = 1, 3, \dots; \\ E_n(mx) &= -\frac{2}{n+1} m^n \sum_{k=0}^{m-1} (-1)^k E_{n+1}\left(x + \frac{k}{m}\right), & n &= 0, 1, \dots, \quad m = 2, 4, \dots; \\ E_n(x+y) &= \sum_{k=0}^n C_n^k E_k(x) y^{n-k}, & n &= 0, 1, \dots \end{aligned}$$

The generating function is expressed as

$$\frac{2e^{xt}}{e^t + 1} \equiv \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \quad (|t| < \pi).$$

This relation may be used as a definition of the Euler polynomials.

Fourier series expansions:

$$\begin{aligned} E_n(x) &= 4 \frac{n!}{\pi^{n+1}} \sum_{k=0}^{\infty} \frac{\sin\left((2k+1)\pi x - \frac{1}{2}\pi n\right)}{(2k+1)^{n+1}} \quad (n = 0, 0 < x < 1; \quad n > 0, 0 \leq x \leq 1); \\ E_{2n}(x) &= 4(-1)^n \frac{(2n)!}{\pi^{2n+1}} \sum_{k=0}^{\infty} \frac{\sin\left((2k+1)\pi x\right)}{(2k+1)^{2n+1}} \quad (n = 0, 0 < x < 1; \quad n > 0, 0 \leq x \leq 1); \\ E_{2n-1}(x) &= 4(-1)^n \frac{(2n-1)!}{\pi^{2n}} \sum_{k=0}^{\infty} \frac{\cos\left((2k+1)\pi x\right)}{(2k+1)^{2n}} \quad (n = 1, 2, \dots, 0 \leq x \leq 1). \end{aligned}$$

Integrals:

$$\begin{aligned} \int_a^x E_n(t) dt &= \frac{E_{n+1}(x) - E_{n+1}(a)}{n+1}, \\ \int_0^1 E_m(t) E_n(t) dt &= 4(-1)^n (2^{m+n+2} - 1) \frac{m! n!}{(m+n+2)!} B_{m+n+2}, \end{aligned}$$

where $m, n = 0, 1, \dots$ and B_n are Bernoulli numbers. The Euler polynomials are orthogonal for even $n + m$.

Connection with the Bernoulli polynomials:

$$E_{n-1}(x) = \frac{2^n}{n} \left[B_n \left(\frac{x+1}{2} \right) - B_n \left(\frac{x}{2} \right) \right] = \frac{2}{n} \left[B_n(x) - 2^n B_n \left(\frac{x}{2} \right) \right],$$

where $n = 1, 2, \dots$

⊙ References for Supplement 11: H. Bateman and A. Erdélyi (1953, 1955), N. W. McLachlan (1955), M. Abramowitz and I. A. Stegun (1964), W. Magnus, F. Oberhettinger, and R. P. Soni (1966), H. Buchholz (1969), S. Yu. Slavyanov and W. Lay (2000), D. Zwillinger (2002), A. D. Polyavin and V. F. Zaitsev (2003), E. W. Weisstein (2003).

Supplement 12

Some Notions of Functional Analysis

12.1. Functions of Bounded Variation

12.1-1. Definition of a Function of Bounded Variation.

1°. Let $f(x)$ be a function defined on a finite segment $[a, b]$. Consider an arbitrary partition of the segment by the points

$$a = x_0 < x_1 < x_2 < \cdots < x_{n-1} < x_n = b$$

and construct the sum

$$v = \sum_{k=0}^{n-1} |f(x_{k+1}) - f(x_k)| \quad (1)$$

whose terms are absolute values of the increments of $f(x)$ on each segment of the partition.

If, for all partitions, the sums (1) are bounded by a constant independent of the partition, one says that the function $f(x)$ has *bounded variation* on the segment $[a, b]$. The supremum of all such sums over all partitions is called the *total variation* of the function $f(x)$ on the segment $[a, b]$. The total variation is denoted by

$$\bigvee_a^b f(x) = \sup\{v\}.$$

A function $f(x)$ is said to have bounded variation on the infinite interval $[a, \infty)$ if it is a function of bounded variation on any finite segment $[a, b]$ and its total variation on $[a, b]$ is bounded by a constant independent of b . By definition,

$$\bigvee_a^\infty f(x) = \sup_{b>a} \left\{ \bigvee_a^b f(x) \right\}.$$

2°. In the above definitions, the continuity of the function $f(x)$ is not mentioned. A continuous function (without additional conditions) may have bounded or unbounded variation.

Example. Consider the continuous function

$$f(x) = \begin{cases} x \cos \frac{\pi}{2x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0 \end{cases}$$

and the partition of the segment $[0, 1]$ by the points

$$0 < \frac{1}{2n} < \frac{1}{2n-1} < \cdots < \frac{1}{3} < \frac{1}{2} < 1.$$

Then the sums (1) corresponding to this partition have the form

$$v_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Therefore, $\bigvee_0^1 f(x) = \infty$.

12.1-2. Classes of Functions of Bounded Variation.

Next, we list some common classes of functions of bounded variation.

1. Any bounded monotone function has bounded variation. Its total variation on the segment $[a, b]$ is defined by $\bigvee_a^b f(x) = |f(b) - f(a)|$.

Remark. The last statement is true for infinite intervals $(-\infty, a]$ and $[a, \infty)$; in the latter case, the total variation is equal to $\bigvee_a^\infty f(x) = |f(\infty) - f(a)|$.

2. Suppose that $f(x)$ is a bounded function on $[a, b]$ and this segment can be divided into finitely many parts

$$[a_k, a_{k+1}] \quad (k = 0, 1, \dots, m-1; \quad a_0 = a, \quad a_m = b),$$

so that the function $f(x)$ is monotone on each part. Then $f(x)$ has bounded variation on $[a, b]$.

Remark. This statement is also true for infinite segments.

3. Let $f(x)$ be a function on a finite segment $[a, b]$ satisfying the *Lipschitz condition*

$$|f(x_1) - f(x_2)| \leq L|x_1 - x_2|,$$

for any x_1 and x_2 in $[a, b]$, where L is a constant. Then $f(x)$ has bounded variation and $\bigvee_a^b f(x) \leq L(b-a)$.

4. Let $f(x)$ be a function on a finite segment $[a, b]$ with a bounded derivative $|f'(x)| \leq L$, where $L = \text{const}$. Then, $f(x)$ is of bounded variation and $\bigvee_a^b f(x) \leq L(b-a)$.

5. Let $f(x)$ be a function on $[a, b]$ or $[a, \infty)$ and suppose that $f(x)$ can be represented as an integral with variable upper limit,

$$f(x) = c + \int_a^x \varphi(t) dt,$$

where $\varphi(t)$ is an absolutely continuous function on the interval under consideration. Then $f(x)$ has bounded variation and

$$\bigvee_a^b f(x) = \int_a^b |\varphi(x)| dx.$$

Corollary. Suppose that $\varphi(t)$ on a finite segment $[a, b]$ or $[a, \infty)$ is integrable, but not absolutely integrable. Then the total variation of $f(x)$ is infinite.

12.1-3. Properties of Functions of Bounded Variation.

Here, all functions are considered on a finite segment $[a, b]$.

1. Any function of bounded variation is bounded.

2. The sum, difference, or product of finitely many functions of bounded variation is a function of bounded variation.

3. Let $f(x)$ and $g(x)$ be two functions of bounded variation and $|g(x)| \geq K > 0$. Then the ratio $f(x)/g(x)$ is a function of bounded variation.

4. Let $a < c < b$. If $f(x)$ has bounded variation on the segment $[a, b]$, then it has bounded variation on each segment $[a, c]$ and $[c, b]$; and the converse statement is true. In this case, the following additivity condition holds:

$$\bigvee_a^b f(x) = \bigvee_a^c f(x) + \bigvee_c^b f(x).$$

5. Let $f(x)$ be a function of bounded variation of the segment $[a, b]$. Then, for $a \leq x \leq b$, the variation of $f(x)$ with variable upper limit

$$F(x) = \bigvee_a^x f(x)$$

is a monotonically increasing bounded function of x .

6. Any function $f(x)$ of bounded variation on the segment $[a, b]$ has a left-hand limit $\lim_{x \rightarrow x_0-0} f(x)$ and a right-hand limit $\lim_{x \rightarrow x_0+0} f(x)$ at any point $x_0 \in [a, b]$.

12.1-4. Criteria for Functions to Have Bounded Variation.

1. A function $f(x)$ has bounded variation on a finite segment $[a, b]$ if and only if there is a monotonically increasing bounded function $\Phi(x)$ such that for all $x_1, x_2 \in [a, b]$ ($x_1 < x_2$), the following inequality holds:

$$|f(x_2) - f(x_1)| \leq \Phi(x_2) - \Phi(x_1).$$

2. A function $f(x)$ has bounded variation on a finite segment $[a, b]$ if and only if $f(x)$ can be represented as the difference of two monotonically increasing bounded functions on that segment: $f(x) = g_2(x) - g_1(x)$.

Remark. The above criteria are valid also for infinite intervals $(-\infty, a]$, $[a, \infty)$, and $(-\infty, \infty)$.

12.1-5. Properties of Continuous Functions of Bounded Variation.

1. Let $f(x)$ be a function of bounded variation on the segment $[a, b]$. If $f(x)$ is continuous at a point x_0 ($a < x_0 < b$), then the function $F(x) = \bigvee_a^x f(x)$ is also continuous at that point.

2. A continuous function of bounded variation can be represented as the difference of two continuous increasing functions.

3. Let $f(x)$ be a continuous function on the segment $[a, b]$. Consider a partition of the segment

$$a = x_0 < x_1 < x_2 < \cdots < x_{n-1} < x_n = b$$

and the sum $v = \sum_{k=0}^{n-1} |f(x_{k+1}) - f(x_k)|$. Letting $\lambda = \max |x_{k+1} - x_k|$ and passing to the limit as $\lambda \rightarrow 0$, we get

$$\lim_{\lambda \rightarrow 0} v = \bigvee_a^b f(x).$$

12.2. Stieltjes Integral

12.2-1. Basic Definitions.

Let $f(x)$ and $\varphi(x)$ be functions defined on an interval $[a, b]$. Let us partition this interval into n elementary subintervals defined by a set of points $\{x_0, x_1, \dots, x_n\}$ such that $a = x_0 < x_1 < \cdots < x_n = b$. Each subinterval $[x_{k-1}, x_k]$ will be characterized by its length $\Delta x_k = x_k - x_{k-1}$ and an arbitrarily chosen point $\xi_k \in [x_{k-1}, x_k]$. Let us make up a *Stieltjes integral sum*

$$s_n = \sum_{k=1}^n f(\xi_k) \Delta_k \varphi(x),$$

where $\Delta_k \varphi(x) = \varphi(x_k) - \varphi(x_{k-1})$ is the increment of the function $\varphi(x)$ on the k th elementary subinterval.

If there exists a limit of the integral sums s_n , as the number of subintervals n increases indefinitely so that the length of every subinterval Δx_k vanishes, and this limit depends on neither the way the interval $[a, b]$ was partitioned nor the way the points ξ_k were selected, then this limit is called the *Stieltjes integral* of the function $f(x)$ with respect to the function $\varphi(x)$ over the interval $[a, b]$:

$$\int_a^b f(x) d\varphi(x) = \lim_{\lambda \rightarrow 0} s_n \quad \left(\max_{1 \leq k \leq n} \Delta x_k \rightarrow 0 \text{ as } n \rightarrow \infty \right).$$

Then $f(x)$ is called an *integrable function* with respect to $\varphi(x)$, and $\varphi(x)$ is called an *integrating function*.

The Stieltjes integral is a generalization of the Riemann integral; the latter corresponds to the special case $\varphi(x) = x + \text{const.}$

12.2-2. Properties of the Stieltjes Integral.

The Stieltjes integral has properties analogous to those of the definite Riemann integral:

- 1) $\int_a^b d\varphi(x) = \varphi(b) - \varphi(a)$;
- 2) $\int_a^b [Af(x) \pm Bg(x)] d\varphi(x) = A \int_a^b f(x) d\varphi(x) \pm B \int_a^b g(x) d\varphi(x)$;
- 3) $\int_a^b f(x) d[A\varphi(x) \pm B\psi(x)] = A \int_a^b f(x) d\varphi(x) \pm B \int_a^b f(x) d\psi(x)$;
- 4) $\int_a^b f(x) d\varphi(x) = \int_a^c f(x) d\varphi(x) + \int_c^b f(x) d\varphi(x) \quad (a < c < b)$.

It is assumed that all integrals on the left- and right-hand sides exist.

THEOREM (MEAN VALUE). *If a function $f(x)$ satisfies inequalities $m \leq f(x) \leq M$ on an interval $[a, b]$ and is integrable with respect to an increasing function $\varphi(x)$, then*

$$\int_a^b f(x) d\varphi(x) = \mu[\varphi(b) - \varphi(a)],$$

where $m < \mu < M$.

12.2-3. Existence Theorems for the Stieltjes Integral.

The existence of the Stieltjes integral and its reduction to the Riemann integral is established by the following theorem.

THEOREM 1. *If $f(x)$ is continuous on $[a, b]$ and $\varphi(x)$ has a bounded variation* on $[a, b]$, then the integral $\int_a^b f(x) d\varphi(x)$ exists.*

* A function $\varphi(x)$ is said to have a *bounded variation* on an interval $[a, b]$ if there exists a number $M > 0$ such that for any set of points $a = x_0 < x_1 < \dots < x_n = b$ the inequality $\sum_{k=1}^n |\varphi(x_{k+1}) - \varphi(x_k)| < M$ holds (see also Supplement 12.1).

THEOREM 2. Let $f(x)$ be integrable on $[a, b]$ in the sense of Riemann and let $\varphi(x)$ satisfy the Lipschitz condition

$$|\varphi(x_2) - \varphi(x_1)| < K|x_2 - x_1|,$$

where x_1 and x_2 are arbitrary points of the interval $[a, b]$ and K is a fixed positive constant. Then the function $f(x)$ is integrable with respect to the function $\varphi(x)$.

THEOREM 3. Let $f(x)$ be integrable on $[a, b]$ in the sense of Riemann and let $\varphi(x)$ be differentiable and have an integrable derivative on $[a, b]$. Then the function $f(x)$ is integrable with respect to the function $\varphi(x)$ and, moreover,

$$\int_a^b f(x) d\varphi(x) = \int_a^b f(x)\varphi'(x) dx,$$

where the integral on the right-hand side is understood in the sense of Riemann.

Remark. If a function $f(x)$ is integrable on an interval $[a, b]$ with respect to a function $\varphi(x)$, then, vice versa, the function $\varphi(x)$ is also integrable with respect to the function $f(x)$ on $[a, b]$. Owing to this property, the functions $f(x)$ and $\varphi(x)$ are interchangeable in Theorems 1 and 2.

THEOREM 4. Let $f(x)$ be continuous on $[a, b]$ and let $\varphi(x)$ have an absolutely integrable derivative $\varphi'(x)$ everywhere on $[a, b]$, except, perhaps, finitely many points. Let, in addition, the function $\varphi(x)$ undergo a jump discontinuity at finitely many points

$$a = c_0 < c_1 < \cdots < c_m = b.$$

Then the Stieltjes integral exists and is calculated as

$$\int_a^b f(x) d\varphi(x) = \int_a^b f(x)\varphi'(x) dx + f(a)[\varphi(a+0) - \varphi(a)] \\ + \sum_{k=1}^{m-1} f(c_k)[\varphi(c_k+0) - \varphi(c_k-0)] + f(b)[\varphi(b) - \varphi(b-0)],$$

where the right-hand side contains a Riemann integral. Note the presence of terms outside the integral on the right-hand side, where, apart from the ordinary jumps of the function $\varphi(x)$ at the internal points of discontinuity, there are terms with one-sided jumps at the endpoints (if there is no jump at either endpoint, the corresponding term vanishes).

The Stieltjes integral is useful for finding static moments, moments of inertia, and some other distributed quantities on an interval $[a, b]$, where, apart from continuous distributions, there are concentrated quantities like point masses that correspond to a discontinuous function $\varphi(x)$ with finite jumps.

12.3. Lebesgue Integral*

12.3-1. Riemann Integral and the Lebesgue Integral.

The space $C[a, b]$ of continuous functions on a finite interval $[a, b]$ is a metric space with the metric

$$\rho(f, g) = \int_a^b |f(x) - g(x)| dx,$$

where the integral is understood in the sense of Riemann. It is well known that this metric space is incomplete, in the sense that there is a Cauchy sequence (with respect to this metric) that does not converge to any element of $C[a, b]$. One can consider a formal completion $L[a, b]$ of the space $C[a, b]$ in this metric. The space $L[a, b]$ is wider than $C[a, b]$ and the problem is to describe the structure of its elements. It turns out that $L[a, b]$ consists of the so-called *summable* or *Lebesgue integrable* functions. Below, we briefly describe a version of the Lebesgue integration theory.

* Supplement 12.3 was written by G. A. Yosifian.

12.3-2. Sets of Zero Measure. Notion of "Almost Everywhere".

Let $[a, b]$ be a finite interval on the real axis x .

A set $A \subset [a, b]$ is called a *set of zero measure* if for any ε it can be covered by finitely many or countably many intervals whose joint length is less than ε .

In particular, any finite or countable set of points on $[a, b]$ is a set of zero measure on $[a, b]$.

The union of finitely many (or countably many) sets of zero measure is a set of zero measure.

A set $B \subset [a, b]$ is called a *set of full measure* on $[a, b]$ if its complement $[a, b] \setminus B$ is a set of zero measure on $[a, b]$.

If some property holds for all points of a segment $[a, b]$ except points of some set of zero measure, one says that this property holds *almost everywhere* on $[a, b]$, or holds *for almost all* $x \in [a, b]$, or holds on a set of full measure.

A function is said to be *defined almost everywhere* on $[a, b]$ if it is defined at all points of $[a, b]$ except points forming a set of zero measure on $[a, b]$.

Let $f_n(x)$ be a sequence of functions defined almost everywhere on $[a, b]$. One says that *the sequence $f_n(x)$ converges to a function $f(x)$ almost everywhere on $[a, b]$* as $n \rightarrow \infty$ if there is pointwise convergence $f_n(x) \rightarrow f(x)$ for almost all $x \in [a, b]$; in other words, if there is pointwise convergence on a set of full measure.

12.3-3. Step Functions and Measurable Functions.

A *partition* of a segment $[a, b]$ is a system of intervals (x_i, x_{i+1}) , $i = 0, 1, \dots, k$, such that $a = x_0 < x_1 < \dots < x_k = b$.

A *step function* on $[a, b]$ is a function that takes a constant value on every interval (x_i, x_{i+1}) of some partition of $[a, b]$.

A *measurable function* $f(x)$ on $[a, b]$ is a function that is defined and finite almost everywhere on $[a, b]$ and can be represented as the pointwise limit (almost everywhere) of a sequence of step functions; in other words, there is a sequence of step functions $f_n(x)$ such that $f_n(x)$ converges to $f(x)$ almost everywhere on $[a, b]$ as $n \rightarrow \infty$.

Since measurable functions are defined almost everywhere, two such functions are identified if they coincide on a set of full measure.

Obviously, any step function is measurable. Many properties of step functions can be transferred to measurable functions. In particular:

(i) All step functions on $[a, b]$ form a linear space, i.e., if f, g are step functions, then their linear combination $\alpha f + \beta g$ is a step function. It follows that all measurable functions on $[a, b]$ form a linear space.

(ii) The product of two step functions is a step function, and accordingly, the product of two measurable functions is a measurable function.

(iii) The ratio of two step functions is a step function, provided that the denominator is different from zero. The ratio of two measurable functions is a measurable function, provided that the denominator differs from zero almost everywhere on $[a, b]$.

(iv) The absolute value $|h(x)|$ of a step function $h(x)$ is a step function. The absolute value of any measurable function is also a measurable function.

(v) Let $f(x), g(x)$ be measurable functions, then the functions

$$h_1(x) = \max\{f(x), g(x)\}, \quad h_2(x) = \min\{f(x), g(x)\}$$

are measurable. In particular, for any measurable function $f(x)$, the functions

$$f^+(x) = \max\{f(x), 0\}, \quad f^-(x) = \max\{0, -f(x)\}$$

are measurable. The functions f^+ and f^- are called the *positive part* and the *negative part* of f , respectively.

Any continuous function on $[a, b]$ (or even a piecewise continuous function) is measurable.

12.3-4. Definition and Properties of the Lebesgue Integral.

Let $h(x)$ be a step function on the interval $[a, b]$ taking constant values h_1, \dots, h_k on mutually disjoint segments $\Delta_1, \dots, \Delta_k$ into which $[a, b]$ is divided by points $a = x_0 < x_1 < \dots < x_k = b$. The *integral* of such a step function $h(x)$ is defined by

$$Ih = \int_{[a,b]} h(x) dx = \sum_{j=1}^k h_j |\Delta_j|,$$

where $|\Delta_j|$ is the length of the interval Δ_j .

For a sequence of function $g_n(x)$ on $[a, b]$, we write $g_n \nearrow g$ if g_n converge to a function g almost everywhere on $[a, b]$ and the numerical sequence $g_n(x)$ is monotonically increasing for almost all $x \in [a, b]$.

DEFINITION 1. A function $f(x)$ on $[a, b]$ is said to belong to the class L^+ if it can be represented as the limit (in the sense of convergence almost everywhere) of a monotonically increasing sequence of step functions $h_n \nearrow f$ and the integrals of these step functions are bounded by the same constant: $Ih_n \leq C$.

Any function of class L^+ is measurable. Continuous functions belong to L^+ .

The *integral* of $f \in L^+$ is defined by the formula

$$If = \lim_{n \rightarrow \infty} Ih_n,$$

where $h_n \nearrow f$ is the sequence from Definition 1 of the class L^+ .

The value If for $f \in L^+$ does not depend on the sequence of step functions $h_n \nearrow f$.

DEFINITION 2. A function $\phi(x)$ on $[a, b]$ is called *summable* or *Lebesgue integrable* on $[a, b]$, (or simply, *integrable*) if it can be represented in the form

$$\phi = f - g, \quad \text{for some } f, g \in L^+.$$

The set of all summable functions is denoted by L .

Properties of summable functions:

(i) if $f, g \in L$, then any linear combination $\alpha f + \beta g$ belongs to L ; in other words, L is a linear space;

(ii) if $f \in L$, then $|f| \in L$;

(iii) if $f, g \in L$ and $h_1(x) = \max\{f(x), g(x)\}$, $h_2(x) = \min\{f(x), g(x)\}$, then $h_1, h_2 \in L$.

DEFINITION 3. The *integral* of a summable function $\phi \in L$ is defined by

$$I\phi = If - Ig, \quad \text{where } \phi = f - g, \quad f, g \in L^+.$$

The value $I\phi$ does not depend on the representation $\phi = f - g$.

Properties of the integral of summable functions:

(i) $I(\phi_1 + \phi_2) = I\phi_1 + I\phi_2$ for any $\phi_1, \phi_2 \in L$;

(ii) $I(\alpha\phi) = \alpha I\phi$ for any $\phi \in L$ and any scalar α ;

(iii) if $f, g \in L$ and $f(x) \geq g(x)$ almost everywhere, then $If \geq Ig$.

THEOREM 1. Any Riemann integrable function on $[a, b]$ (in particular, any continuous function on $[a, b]$) is Lebesgue integrable, and its Riemann integral coincides with its Lebesgue integral.

For a sequence $\phi_n \in L$ such that $\phi_n \rightarrow \phi$ almost everywhere, it cannot be claimed, in general, that $I\phi_n \rightarrow I\phi$. For example, consider the sequence

$$\phi_n(x) = \begin{cases} n \sin nx & \text{for } 0 \leq x \leq \frac{\pi}{n}, \\ 0 & \text{for } \frac{\pi}{n} \leq x \leq \pi. \end{cases}$$

It is easy to verify that $\phi_n(x) \rightarrow 0$ for any $x \in [0, \pi]$, but $I\phi_n = 2$.

An important result with regard to integrating pointwise convergent sequences is the following theorem.

THEOREM 2 (LEBESGUE THEOREM ON DOMINATED CONVERGENCE). Let ϕ_n be a sequence of summable functions that converges to a function ϕ almost everywhere and satisfies the condition

$$|\phi_n(x)| \leq \phi_0(x) \in L.$$

Then ϕ is a summable function and $I\phi = \lim_{n \rightarrow \infty} I\phi_n$. In particular, $I\phi = \lim_{n \rightarrow \infty} I\phi_n$ if the functions ϕ_n are uniformly bounded.

Some important properties of measurable and summable functions:

(i) If ϕ is a measurable function that satisfies (almost everywhere) the inequality

$$-\phi_0 \leq \phi \leq \phi_0 \in L.$$

Then $\phi \in L$.

(ii) The limit of a sequence of measurable functions that converges almost everywhere to a finite limit is a measurable function.

(iii) (Fatou lemma.) If $\phi_n \geq 0$ is a sequence of summable functions, $\phi_n \rightarrow \phi$ almost everywhere, and $I\phi_n \leq C$, then ϕ is a summable function and $0 \leq I\phi \leq C$.

(iv) If $\phi_0(x) \geq 0$ is a summable function such that $I\phi_0 = 0$, then $\phi_0 = 0$ almost everywhere.

THEOREM 3 (FISCHER–RIESZ). The space L endowed with the norm

$$\|\phi\| = I(|\phi|)$$

is a Banach space.

THEOREM 4. The space L is the completion of the space $C[a, b]$ with respect to the norm

$$\|f\| = \int_a^b |f(x)| dx.$$

In other words, continuous functions form a dense set in L .

12.3-5. Measurable Sets.

A set $A \subset [a, b]$ is called *measurable* if its characteristic function

$$\chi_A(x) = \begin{cases} 1 & \text{for } x \in A, \\ 0 & \text{for } x \in [a, b] \setminus A \end{cases}$$

is measurable.

The integral of the characteristic function of a measurable set $A \subset [a, b]$ is called the *measure* of A and is denoted by $\mu(A)$, i.e.,

$$\mu(A) = \int_{[a,b]} \chi_A(x) dx.$$

In particular, for a set B of zero measure, we have $\mu(B) = 0$.

Measurable sets have the following properties:

(i) the union $A = \bigcup A_j$ of finitely many or countably many measurable sets A_1, \dots, A_n, \dots is a measurable set; moreover, if the sets A_j are mutually disjoint, i.e., $A_j \cap A_i = \emptyset$ for all $i \neq j$, then

$$\mu(A) = \mu(A_1) + \dots + \mu(A_n) + \dots;$$

(ii) the intersection $A = \bigcap A_j$ of finitely many or countably many measurable sets A_1, \dots, A_n, \dots is a measurable set;

(iii) the difference $A = B \setminus C$ of measurable sets B, C is a measurable set, in particular, the complement of B , i.e., $[a, b] \setminus B$, is a measurable set;

(iv) any interval $[\alpha, \beta]$, $(\alpha, \beta]$, (α, β) , $[\alpha, \beta)$ is a measurable set and its measure is equal to its length $\beta - \alpha$;

(v) any open and any closed set on $[a, b]$ is measurable.

12.3-6. Integration Over Measurable Sets.

So far, the domain of integration has been the interval $[a, b]$. It is easy to extend the notion of integral to any measurable set $E \subset [a, b]$.

A function ϕ is called *summable* (or *integrable*) on a measurable set E if the function $\chi_E(x)\phi(x)$ is summable on $[a, b]$, where χ_E is the characteristic function of E . The *integral* of ϕ over E is defined by

$$\int_E \phi dx = \int_{[a,b]} \chi_E(x)\phi(x) dx = I(\chi_E\phi).$$

This integral has the following *additive property*: if ϕ is summable on a set $E = E_1 \cup E_2 \cup \dots$, where E_1, E_2, \dots are mutually disjoint measurable sets, then ϕ is summable on each E_j and

$$\int_E \phi dx = \int_{E_1} \phi dx + \int_{E_2} \phi dx + \dots.$$

12.3-7. Case of an Infinite Interval.

The above considerations pertain to functions defined on a finite interval $[a, b]$. It is not very difficult to extend the above theory to the cases of intervals $[a, \infty)$, $(-\infty, b]$, or $(-\infty, \infty)$.

In all these cases, a *step function* is defined as a function taking constant values on finitely many finite intervals $\Delta_j = (x_j, x_{j+1})$ ($x_j < x_{j+1}$) and on the rest of the infinite interval, it is supposed to be equal to zero. A *measurable function* is a function $\phi(x)$ that is the limit (almost everywhere on every finite segment) of a sequence of step functions. The *integral* of a step function $h(x)$ taking values h_j on an interval Δ_j of length $|\Delta_j|$ ($j = 1, \dots, k$) is naturally defined by the formula

$$Ih = \sum_{j=1}^k h_j |\Delta_j|.$$

The class L^+ consists of all functions $f(x)$ that can be represented as the limit of an increasing sequence of step functions $f_n(x)$ with bounded integrals. The class L is defined as the set of differences $\phi = f - g$, $f, g \in L^+$. The results formulated above for a finite interval can be easily extended to the case infinite intervals.

12.3-8. Case of Several Variables.

We limit ourselves to functions of two variables $\phi(x, y)$ defined on a rectangle $D = \{a_1 \leq x \leq b_1, a_2 \leq y \leq b_2\}$.

A set $A \subset D$ is called a *set of zero measure* in D if for any ε the set D can be covered by a finite or countable system of rectangles $D_j = \{a_1^{(j)} \leq x \leq b_1^{(j)}, a_2^{(j)} \leq y \leq b_2^{(j)}\}$ whose joint area does not exceed ε .

A partition of D is a system of mutually disjoint open rectangles $D_1, \dots, D_k \subset D$ such that $D = \overline{D}_1 \cup \dots \cup \overline{D}_k$, where \overline{D}_j is the closure of D_j .

A *step function* on D is a function that takes constant values on each rectangle D_j of some partition of D , $D = \overline{D}_1 \cup \dots \cup \overline{D}_k$.

The *integral* of a step function $h(x)$ with values h_j on the rectangles D_j of some partition is defined as

$$Ih = \sum_{j=1}^k h_j |D_j|,$$

where $|D_j|$ is the area the rectangle of D_j .

As in the one-dimensional case, the *class* L^+ is the set of functions f such that f is a limit (almost everywhere on D) of a sequence of step functions f_n with uniformly bounded integrals. The *class* L of *summable functions* is again defined as the set of differences $\phi = f - g$, $f, g \in L^+$.

The properties formulated above for the one-dimensional case are obviously modified in the case of two dimensions. However, in the two-dimensional case, there is the question of the reduction of an integral over a two-dimensional domain D to a double integral over linear segments, and also the question of changing the order of double integration. The answers to these questions are given by the following theorem.

THEOREM 5 (FUBINI THEOREM). *Let $\phi(x, y)$ be a summable function on a rectangle $D = \{a_1 \leq x \leq b_1, a_2 \leq y \leq b_2\}$. Then:*

- (i) *regarded as a function of the argument x for a fixed y , this function is integrable in x for almost all y ;*
- (ii) *its integral over the interval $a_1 \leq x \leq b_1$, denoted by $I_x \phi(x, y)$, is a summable function of y on the interval $a_2 \leq y \leq b_2$;*
- (iii) *the integral over D can be reduced to a double integral in which the order of integration can be changed:*

$$I\phi = I_y \{I_x \phi(x, y)\} = I_x \{I_y \phi(x, y)\}.$$

As in the one-dimensional case, a set $G \subset D$ is called *measurable* if its characteristic function $\chi_G(x, y)$ is measurable and the integral of ϕ over G is defined by the formula

$$\int_G \phi \, dx \, dy = \int_D \chi_G(x, y) \phi(x, y) \, dx \, dy.$$

12.3-9. Spaces L_p .

For a measurable set G and $p > 0$, the *class* $L_p(G)$ consists of all measurable functions $f(x)$ on G for which $|f|^p$ is summable on G , i.e.,

$$\int_G |f|^p \, dx < \infty.$$

For any $p > 0$, this class of functions is a linear space.

For $p \geq 1$, the class $L_p(G)$ is a Banach space (complete normed space) with the norm

$$\|f\|_p = \left(\int_G |f|^p \right)^{1/p}.$$

The set of continuous functions is dense in the Banach space $L_p(G)$, i.e., for any $f \in L_p$, there is a sequence of continuous functions f_n such that $\|f - f_n\|_p \rightarrow 0$ as $n \rightarrow \infty$.

Let $p > 1$, $q > 1$ be real numbers such that $p^{-1} + q^{-1} = 1$. For $f \in L_p$, $g \in L_q$, the product fg is summable on G and the *Hölder inequality* holds:

$$\int_G fg \, dx \leq \|f\|_p \|g\|_q.$$

12.4. Linear Normed Spaces

12.4-1. Linear Spaces.

A *linear space* or a *vector space* L over the field of real or complex numbers (called the *field of scalars*) is a nonempty set of elements (also called *vectors*) for which two operations are defined: addition of elements and their multiplication by scalars. To be more precise: for any two elements $x, y \in L$, there is a unique element $z \in L$, called their *sum* and denoted by $z = x + y \in L$, and for any scalar α (real or complex) and any element $x \in L$ there is a unique element y , called the product of α and x and denoted by $y = \alpha x$, so that for these two operations the following axioms hold:

I. Axioms for addition of vectors:

- 1) $x + y = y + x$ (commutative property);
- 2) $x + (y + z) = (x + y) + z$ (associative property);
- 3) there is an element $0 \in L$ such that $x + 0 = x$ for all $x \in L$ (existence of zero);
- 4) for any $x \in L$, equation $x + y = 0$ is solvable; the element y is called the *opposite* of x and is denoted by $-x$, so that $x + (-x) = 0$ (existence of an opposite element).

II. Axioms relating addition of vectors with their multiplication by scalars:

- 5) $\alpha(\beta x) = (\alpha\beta)x$ for any vector $x \in L$ and any scalars α, β ;
- 6) $1 \cdot x = x$ for any $x \in L$;
- 7) $(\alpha + \beta)x = \alpha x + \beta x$ for any $x \in L$ and any scalars α, β ;
- 8) $\alpha(x + y) = \alpha x + \alpha y$ for any scalar α and any vectors $x, y \in L$.

If the field of scalars is the set of real numbers, then L is called a *real linear space*. If the field of scalars is the set of all complex numbers, then L is called a *complex linear space*.

Elements (vectors) y_1, y_2, \dots, y_n of a linear space L are called *linearly dependent* if there exist scalar coefficients $\alpha_1, \alpha_2, \dots, \alpha_n$ such that at least one of them is different from zero and $\alpha_1 y_1 + \alpha_2 y_2 + \dots + \alpha_n y_n = 0$. Otherwise, vectors y_1, y_2, \dots, y_n are called *linearly independent*.

A nonempty subset \bar{L} of a linear space L is called its *subspace* if for any $x, y \in \bar{L}$ and any scalars α, β , we have $\alpha x + \beta y \in \bar{L}$.

12.4-2. Linear Normed Spaces.

A linear space L is called a *normed space* if any element $y \in L$ is associated with a real number $\|y\| \geq 0$, called the *norm* of y , so that the following properties (axioms of a linear normed space) hold:

- 1) $\|y\| = 0$ if and only if $y = 0$;
- 2) $\|\lambda y\| = |\lambda| \|y\|$ for any scalar λ (homogeneity of the norm);
- 3) $\|y_1 + y_2\| \leq \|y_1\| + \|y_2\|$ (triangle inequality).

A sequence $\{y_n\}$ of elements of a normed space L is called *convergent* to an element y_0 if $\|y_0 - y_n\| \rightarrow 0$ as $n \rightarrow \infty$.

12.4-3. Space of Continuous Functions $C(a, b)$.

The linear normed space $C(a, b)$ consists of all continuous functions $y(x)$ on the interval $[a, b]$, with the norm defined by

$$\|y\| = \max_{a \leq x \leq b} |y(x)|.$$

The distance between two functions in this space has the form

$$\rho(y_1, y_2) = \max_{a \leq x \leq b} |y_1(x) - y_2(x)|.$$

The convergence of a sequence of functions $\{y_n\}$ in the space $C(a, b)$ to an element $y_0(x)$ means uniform convergence of the functions $y_n(x)$ to $y_0(x)$.

12.4-4. Lebesgue Space $L_p(a, b)$.

The linear normed space $L_p(a, b)$ ($p \geq 1$) consists of all measurable functions $y(x)$ on (a, b) such that $|y(x)|^p$ is integrable (has finite integral) on $[a, b]$, and the norm in $L^p(a, b)$ is defined by

$$\|y\| = \left(\int_a^b |y(x)|^p dx \right)^{1/p}.$$

Convergence $y_n \rightarrow y_0$ in $L_p(a, b)$ means that

$$\int_a^b |y_n(x) - y_0(x)|^p dx \rightarrow 0.$$

Remark 1. Functions $y_1(x)$ and $y_2(x)$ in $L_p(a, b)$ that coincide almost everywhere (i.e., may differ only on a set of zero measure) are identified.

Remark 2. With regard to the space $L_2(a, b)$, see also Subsection 9.1-1.

12.4-5. Hölder Space $C_\alpha(0, 1)$.

The normed linear space $C_\alpha(0, 1)$ is the set of all functions $y(x)$ defined on the interval $[0, 1]$ and satisfying the *Hölder condition* with exponent α ($0 < \alpha \leq 1$):

$$|y(x_1) - y(x_2)| \leq A|x_1 - x_2|^\alpha \quad (0 \leq x_1, x_2 \leq 1).$$

The norm of a function $y(x)$ in $C_\alpha(0, 1)$ is introduced by the formula

$$\|y\| = |y(0)| + \sup_{0 \leq x_1, x_2 \leq 1} \frac{|y(x_1) - y(x_2)|}{|x_1 - x_2|^\alpha}.$$

12.4-6. Space of Functions of Bounded Variation $V(0, 1)$.

The normed linear space $V(0, 1)$ is the set of all functions of bounded variation (see Supplement 12.1) on the interval $[0, 1]$.

The norm of $y(x)$ in $V(0, 1)$ is introduced by

$$\|y\| = |y(0)| + \bigvee_0^1 y(x).$$

12.5. Euclidean and Hilbert Spaces. Linear Operators in Hilbert Spaces

12.5-1. Preliminary Remarks.

The mathematical concept of a Hilbert space generalizes the notion of Euclidean space in a way that extends methods of vector algebra from the two-dimensional plane and three-dimensional space to infinite-dimensional spaces. In more formal terms, a Hilbert space is an inner product space—an abstract vector space in which distances and angles can be measured—which is “complete,” meaning that if a sequence of vectors approaches a limit, then that limit is guaranteed to be in the space as well.

Geometric intuition plays an important role in many aspects of Hilbert space theory. An element of a Hilbert space can be uniquely specified by its coordinates with respect to an orthonormal basis, in analogy with cartesian coordinates in the plane. This means that Hilbert space can also usefully be thought of in terms of infinite sequences that are square-summable. Linear operators on a Hilbert space are likewise fairly concrete objects: in good cases, they are simply transformations that stretch the space by different factors in mutually perpendicular directions.

12.5-2. Euclidean and Hilbert Spaces.

A *Euclidean space* E is a (real or complex) linear space endowed with a scalar product $\{x, y\} \mapsto (x, y)$, i.e., a mapping of $E \times E$ into the field of real or complex numbers satisfying the following conditions:

$$\begin{aligned} (x, y) &= \overline{(y, x)} \text{ for all } x, y \in E, \\ (x + y, z) &= (x, z) + (y, z) \text{ for all } x, y, z \in E, \\ (\lambda x, y) &= \lambda(x, y) \text{ for all } x, y \in E \text{ and all (real or complex) } \lambda, \\ (x, x) &\geq 0 \text{ for all } x \in E, \text{ and } x = 0 \iff (x, x) = 0. \end{aligned}$$

Here the bar over a complex number denotes its complex conjugate.

For a Euclidean space E , the formula

$$\|x\| = \sqrt{(x, x)}, \quad x \in E,$$

defines a norm on E . Therefore, any Euclidean space can be regarded as a normed space. Vectors $x, y \in E$ are called *orthogonal* if $(x, y) = 0$. A set of nonzero vectors $\{e_i, i \in I\} \subset E$ (here I is a set of indices) is called an *orthogonal system in E* if e_i and e_j are orthogonal for all $i \neq j, i, j \in I$. An orthogonal system $\{e_i, i \in I\}$ is called an *orthonormal system* if $\|e_i\| = 1$ for any $i \in I$. A system of vectors $\{e_i, i \in I\}, e_i \in E$, is called *complete* if any $x \in E$ can be approximated in the norm of E (with any given accuracy) by finite linear combinations of the vectors e_i , i.e., for any $\varepsilon > 0$ there is a finite linear combination $\sum_i c_i e_i$ such that $\left\|x - \sum_i c_i e_i\right\| < \varepsilon$.

A normed linear space is called a *complete space* or a *Banach space* if the Cauchy criterion holds for that space, namely, for any sequence $\{x_n, n \in \mathbb{N}\}, x_n \in G$ (here \mathbb{N} is the set of all positive integers) the following conditions are equivalent:

- there exists an $x_0 \in G$ such that $\lim_{n \rightarrow \infty} \|x_n - x_0\| = 0$;
- for any $\varepsilon > 0$, there exists an $N \in \mathbb{N}$ such that $\|x_n - x_m\| < \varepsilon$ for all $m, n > N$.

A complete Euclidean space is called a *Hilbert space*. An orthogonal system $\{e_i, i \in I\}$ in a Hilbert space E is complete if and only if the only vector in E orthogonal to every vector of the system $\{e_i, i \in I\}$ is the zero-vector. For a closed linear subspace L in a Hilbert space E , the symbol L^\perp denotes the set of all vectors $y \in E$ such that $(x, y) = 0$ for all $x \in L$. The set L^\perp is a closed linear subspace of E called the *orthogonal complement* of L . Any vector $x \in E$ can be uniquely represented as a sum $x = y + z$, where $y \in L$ and $z \in L^\perp$. In particular, the orthogonal complement of L^\perp coincides with L .

THEOREM 1. Any closed subspace of a Hilbert space is either finite-dimensional or is itself a Hilbert space.

A Hilbert space H is said to be represented as a *direct sum* of its orthogonal subspaces M_1, M_2, \dots, M_n ,

$$H = M_1 \oplus M_2 \oplus \dots \oplus M_n,$$

if for any $f \in H$ there exist $h_1 \in M_1, \dots, h_n \in M_n$ such that $f = h_1 + \dots + h_n$, and any element of M_i is orthogonal to any element of M_k for $i \neq k$.

THEOREM 2. Any element $f \in H$ can be uniquely represented in the form $f = h_1 + h_2 + \dots + h_n$, where $h_j \in M_j$.

COROLLARY. If $\{\varphi_n^i\}$ are complete orthonormal systems in the subspaces M_i , then the union of all $\{\varphi_n^i\}$ is a complete orthonormal system in H .

12.5-3. Linear Operators in Hilbert Spaces.

Given two linear spaces L and L_1 any mapping

$$y = \mathbf{A}x \quad (x \in L, y \in L_1)$$

of subset of L (possibly L itself) into L_1 is called *operator* (from L to L_1). The operator \mathbf{A} is said to be *linear* if

$$\mathbf{A}(\alpha x + \beta y) = \alpha \mathbf{A}x + \beta \mathbf{A}y.$$

Let $D_{\mathbf{A}}$ be the set of all $x \in L$ for which \mathbf{A} is defined. Then $D_{\mathbf{A}}$ is called the *domain (of definition)* of operator \mathbf{A} . Although in general $D_{\mathbf{A}}$ need not equal L , we will always assume that $D_{\mathbf{A}}$ is a linear subspace of L , i.e., that $x, y \in D_{\mathbf{A}}$ implies $\alpha x + \beta y \in D_{\mathbf{A}}$ for all α and β .

The operator \mathbf{A} is said to be *continuous at the point* $x_0 \in D$ if, given any neighborhood V of the point $y_0 = \mathbf{A}x_0$, there is a neighborhood U of the point x_0 such that $\mathbf{A}x \in V$ for all $x \in U \cap D_{\mathbf{A}}$. We say that the operator \mathbf{A} is *continuous* if it is continuous at every point $x_0 \in D_{\mathbf{A}}$.

Suppose L and L_1 are normed linear spaces. Then it is easy to see that \mathbf{A} is continuous if and only if, given any $\varepsilon > 0$, there is a $\delta > 0$ such that

$$\|x - y\| < \delta \quad (x, y \in D_{\mathbf{A}})$$

implies

$$\|\mathbf{A}x - \mathbf{A}y\| < \varepsilon.$$

Given a bounded linear operator mapping a normed linear space L into another linear space L_1 , the number $\|\mathbf{A}\| = \sup_{\|x\| < 1} \|\mathbf{A}x\|$, equal to the least upper bound of $\|\mathbf{A}x\|$ on the closed unit sphere

$\|x\| < 1$, is called the *norm* of \mathbf{A} .

The norm $\|\mathbf{A}\|$ has the following properties:

$$\begin{aligned} \|\mathbf{A}\| &= \sup_{x \neq 0} \frac{\|\mathbf{A}x\|}{\|x\|}, \\ \|\mathbf{A}x\| &\leq \|\mathbf{A}\| \|x\| \quad \text{for all } x \in L. \end{aligned}$$

An *eigenvalue* of a linear operator \mathbf{A} in H is defined as a scalar μ for which there is $x \neq 0$ such that $\mathbf{A}x = \mu x$. The element x in this relation is called an *eigenvector* or an *eigenfunction* of \mathbf{A} corresponding to the eigenvalue μ . The set of all eigenvalues of \mathbf{A} is called *spectrum* of \mathbf{A} , and all other values of μ are said to be *regular (points)*.

If H is Hilbert space, then by the adjoint of an operator \mathbf{A} mapping H into H , we mean the operator \mathbf{A}^* defined by $(\mathbf{A}x, y) = (x, \mathbf{A}^*y)$ for all $x, y \in H$.

A bounded linear operator \mathbf{A} mapping a Hilbert space H into itself is said to be *self-adjoint* if $\mathbf{A} = \mathbf{A}^*$, i.e., if $(\mathbf{A}x, y) = (x, \mathbf{A}y)$ for all $x, y \in H$.

An operator \mathbf{A} in H is *positive definite* if for all nonzero x , $(\mathbf{A}x, x) > 0$.

Let \mathbf{A} be a linear operator mapping a Hilbert space H into itself. Then \mathbf{A} is completely continuous if and only if:

1) \mathbf{A} maps every relatively compact set in the weak topology into a relatively compact set in the strong topology;

2) \mathbf{A} maps every weakly convergent sequence into a strongly convergent sequence.

THEOREM 3. *All eigenvalues of a self-adjoint operator in H are real, and eigenvectors corresponding to different eigenvalues are orthogonal.*

THEOREM 4. *The set of all eigenvalues of a compact operator in H is no more than countable. Zero is the only possible limit point of this set.*

THEOREM 5. *All eigenvalues of a compact self-adjoint positive definite operator in H are positive.*

THEOREM 6 (HILBERT–SCHMIDT). *Let \mathbf{A} be a compact self-adjoint linear operator in a Hilbert space H . Then there is an orthonormal system of eigenvectors $\{\phi_n\}$ corresponding to eigenvalues $\{\mu_n\}$ ($\mu_n \neq 0$) such that each element $\xi \in H$ can be uniquely represented in the form*

$$\xi = \sum_k c_k \phi_k + \xi',$$

where $\xi' \in \text{Ker } \mathbf{A}$, i.e., $\mathbf{A}\xi' = 0$. Moreover,

$$\mathbf{A}\xi = \sum_k \mu_k c_k \phi_k,$$

and if the system $\{\phi_n\}$ is infinite, then $\lim_{n \rightarrow \infty} \mu_n = 0$.

COROLLARY. *If zero is not an eigenvalue of the operator \mathbf{A} , then the system $\{\phi_n\}$ is complete in H . In particular, for a compact self-adjoint positive definite operator, this system forms a basis in H .*

Suppose that a Hilbert space H is represented as a direct sum of its two orthogonal closed subspaces: $H = H_1 \oplus H_2$. Thus each element $h \in H$ can be uniquely represented in the form $h = h_1 + h_2$ ($h_i \in H_i$, $i = 1, 2$).

An operator $\mathbf{P}_i : H \rightarrow H_i$ defined by the relation $\mathbf{P}_i h = h_i$ is called the *orthogonal projector* of H onto H_i ($i = 1, 2$). Obviously $\mathbf{P}_2 = \mathbf{I} - \mathbf{P}_1$, where \mathbf{I} is the identity operator.

Any orthogonal projector is a linear continuous self-adjoint operator in H .

Orthogonal projectors have the following properties:

$$\mathbf{P}_i h_i = h_i, \quad \mathbf{P}_1 h_2 = \mathbf{P}_2 h_1 = 0, \quad \mathbf{P}_1 \mathbf{P}_2 h = 0, \quad \|\mathbf{P}_i\| = 1.$$

THEOREM 7. *In a Hilbert space H an operator of orthogonal projection onto a subspace is compact if and only if this subspace has a finite dimension.*

THEOREM 8. *A linear operator P on H is an orthogonal projector if and only if \mathbf{P} is self-adjoint and satisfies the condition $\mathbf{P}(\mathbf{P}x) = \mathbf{P}x$ for any $x \in H$ (i.e., $\mathbf{P}^2 = \mathbf{P}$).*

THEOREM 9. *A linear combination of compact operators is a compact operator.*

THEOREM 10. *If \mathbf{A} is a compact operator and \mathbf{B} is a bounded linear operator, then the operators $\mathbf{A}\mathbf{B}$ and $\mathbf{B}\mathbf{A}$ are compact.*

© References for Supplement 12: L. V. Kantorovich and G. P. Akilov (1964), R. Edwards (1965), M. G. Krein (1972), M. Reed and B. Simon (1972), W. Rudin (1973), K. Yosida (1980), A. N. Kolmogorov and S. V. Fomin (1999), B. M. Levitan (2001), A. D. Polyaniin and A. V. Manzhirov (2007), http://en.wikipedia.org/wiki/Hilbert_space.

References

- Ablowitz, M. J. and Clarkson, P. A.**, *Solitons, Non-linear Evolution Equations and Inverse Scattering*, Cambridge Univ. Press, Cambridge, 1991.
- Abramowitz, M. and Stegun, I. A. (Editors)**, *Handbook of Mathematical Functions With Formulas, Graphs and Mathematical Tables*, National Bureau of Standards, Washington, 1964.
- Acrivós, A. and Shambre, P. L.**, Laminar boundary layer flows with surface reactions, *Ind. Eng. Chem. Fundam.*, Vol. 49, pp. 1025–1029, 1957.
- Adams, R.**, *Calculus: A Complete Course, 6th Edition*, Pearson Education, Toronto, 2006.
- Agarwal, R. P., O'Regan, D., and Wong, P. J. Y.**, *Positive Solutions of Differential, Difference and Integral Equations*, Springer-Verlag, New York, 1998.
- Aggarwala, B. D. and Nasim, C.**, Steady-state temperature in a quarter plane, *Int. J. Math. & Math. Sci.*, Vol. 19, pp. 371–380, 1996.
- Akhiezer, N. I. and Glazman, I. M.**, *Theory of Linear Operators in Hilbert Space* [in Russian], Nauka, Moscow, 1966.
- Alexandrov, V. M.**, Asymptotic methods in the mechanics of continuous media: problems with mixed boundary conditions, *Appl. Math. and Mech. (PMM)*, Vol. 57, No. 2, pp. 321–327, 1993.
- Alexandrov, V. M. and Kovalenko, E. V.**, *Problems With Mixed Boundary Conditions in Continuum Mechanics* [in Russian], Nauka, Moscow, 1986.
- Alexandrov, V. M., and Manzhairov, A. V.**, Two-dimensional integral equations in applied solid mechanics, *Applied Mechanics and Technical Physics*, No. 5, pp. 146–152, 1987.
- Alexandrov, V. M. and Mkhitaryan, S. M.**, *Contact Problems for Bodies with Thin Coatings and Interlayers* [in Russian], Nauka, Moscow, 1983.
- Andreev, A. V.**, A method for numerical solution of complete singular integral equations with complex power-law singularities, *Mech. Solids*, Vol. 41, No. 1, pp. 76–87, 2006.
- Andreev, A. V.**, A method of determination of exponential type complex singularities in the solutions of singular integral equations with generalized kernels and conjugate variables, *Mechanics of Solids*, 2007 (to be published).
- Andreev, A. V.**, Development of direct numerical integration methods of one-dimensional integro-differential equations in mechanics, *Mech. Solids*, Vol. 42, No. 2, pp. 209–222, 2007.
- Andreev, A. V.**, Direct numerical method for solving singular integral equations of the first kind with generalized kernels, *Mech. Solids*, Vol. 40, No. 1, pp. 104–119, 2005.
- Antimirov, M. Ya.**, *Applied Integral Transforms*, American Mathematical Society, Providence, Rhode Island, 1993.
- Anton, H., Bivens, I., and Davis, S.**, *Calculus: Early Transcendental Single Variable, 8th Edition*, John Wiley & Sons, New York, 2005.
- Arutyunyan, N. Kh.**, A plane contact problem of creep theory, *Appl. Math. and Mech. (PMM)*, Vol. 23, No. 5, pp. 901–924, 1959.
- Arutyunyan, N. Kh.**, *Some Problems in the Theory of Creep*, Pergamon Press, Oxford, 1966.
- Arutyunyan, N. Kh., Manzhairov, A. V.**, *Contact Problems in the Theory of Creep* [in Russian], 1990, Izd-vo NAN RA, Erevan, 1999.
- Arutyunyan, N. Kh., Manzhairov, A. V., and Naumov V. E.**, *Contact Problems in Mechanics of Growing Solids* [in Russian], Nauka, Moscow, 1991.

- Atkinson, K. E.**, *Numerical Solution of Integral Equations of the Second Kind*, Cambridge Univ. Press, Cambridge, 1997.
- Babenko, Yu. I.**, *Heat and Mass Transfer: A Method for Computing Heat and Diffusion Flows* [in Russian], Khimiya, Moscow, 1986.
- Bakhvalov, N. S.**, *Numerical Methods* [in Russian], Nauka, Moscow, 1973.
- Bateman, H.**, On the numerical solution of linear integral equations, *Proc. Roy. Soc. (A)*, Vol. 100, No. 705, pp. 441–449, 1922.
- Bateman, H. and Erdélyi, A.**, *Higher Transcendental Functions. Vol. 1*, McGraw-Hill Book Co., New York, 1953.
- Bateman, H. and Erdélyi, A.**, *Higher Transcendental Functions. Vol. 2*, McGraw-Hill Book Co., New York, 1953.
- Bateman, H. and Erdélyi, A.**, *Higher Transcendental Functions. Vol. 3*, McGraw-Hill Book Co., New York, 1955.
- Bateman, H. and Erdélyi, A.**, *Tables of Integral Transforms. Vol. 1*, McGraw-Hill Book Co., New York, 1954.
- Bateman, H. and Erdélyi, A.**, *Tables of Integral Transforms. Vol. 2*, McGraw-Hill Book Co., New York, 1954.
- Beerends, R. J., ter Morschem, H. G., and van den Berg, J. C.**, *Fourier and Laplace Transforms*, Cambridge University Press, Cambridge, 2003.
- Bellman, R. and Cooke, K. L.**, *Differential–Difference Equations*, Academic Press, New York, 1963.
- Bellman, R. and Roth, R.**, *The Laplace Transform*, World Scientific Publishing Co., Singapore, 1984.
- Belotserkovskii, S. M. and Lifanov I. K.**, *Method of Discrete Vortices*, CRC Press, Boca Raton–New York, 1993.
- Beyer, W. H.**, *CRC Standard Mathematical Tables and Formulae*, CRC Press, Boca Raton, 1991.
- Bitsadze, A. V.**, *Integral Equation of the First Kind*, World Scientific, Singapore, 1995.
- Bochner, S.**, *Lectures on Fourier Integrals*, Princeton Univ. Press, Princeton, 1959.
- Bochner, S. and Chandrasekharan, K. C.**, *Fourier Transforms*, Princeton Univ. Press, Princeton, 1949.
- Bracewell, R.**, *The Fourier Transform and Its Applications, 3rd Edition*, McGraw-Hill, New York, 1999.
- Brakhage, H., Nickel, K., and Rieder, P.**, Auflösung der Abelschen Integralgleichung 2. Art, *ZAMP*, Vol. 16, Fasc. 2, S. 295–298, 1965.
- Bronshstein, I. N. and Semendyayev, K. A.**, *Handbook of Mathematics, 4th Edition*, Springer-Verlag, Berlin, 2004.
- Brunner, H.**, *Collocation Methods for Volterra Integral and Related Functional Differential Equations*, Cambridge University Press, Cambridge, 2004.
- Brychkov, Yu. A. and Prudnikov, A. P.**, Integral Transforms, In: *Mathematical Encyclopedia*, Vol. 2, pp. 589–590 [in Russian], Sovetskaya Entsiklopediya, Moscow, 1979.
- Brychkov, Yu. A. and Prudnikov, A. P.**, *Integral Transforms of Generalized Functions*, Gordon & Breach Sci. Publ., New York, 1989.
- Buchholz, H.**, *The Confluent Hypergeometric Function*, Springer-Verlag, New York, 1969.
- Bueckner, H. F.**, On a class of singular integral equations, *J. Math. Anal. Appl.*, Vol. 14, pp. 392–426, 1966.
- Busbridge, I. W.**, Dual integral equations, *Proc. Lond. Math. Soc.*, Vol. 44, pp. 115–129, 1938.
- Butkovskii, A. G.**, *Characteristics of Systems With Distributed Parameters* [in Russian], Nauka, Moscow, 1979.
- Cochran, J. A.**, *The Analysis of Linear Integral Equations*, McGraw-Hill Book Co., New York, 1972.
- Collatz, L.**, *The Numerical Treatment of Differential Equations*, Springer-Verlag, Berlin, 1960.

- Corduneanu, C.**, *Integral Equations and Applications*, Cambridge Univ. Press, Cambridge–New York, 1991.
- Corduneanu, C.**, *Integral Equations and Stability of Feedback Systems*, Academic Press, New York, 1973.
- Courant, R. and Hilbert, D.**, *Methods of Mathematical Physics. Vol. 1.*, Interscience Publ., New York, 1953.
- Courant, R. and Hilbert, D.**, *Methods of Mathematical Physics, Vol. 2*, Wiley-Interscience, New York, 1989.
- Courant, R. and John, F.**, *Introduction to Calculus and Analysis, Vol. 1*, Springer-Verlag, New York, 1999.
- Davenport, W. B. and Root, W. L.**, *An Introduction to the Theory of Random Signals and Noise*, McGraw-Hill Book Co., New York, 1958.
- David G. and Journé J.-L.** A boundedness criterion for generalized Calderón–Zygmund operators, *Ann. of Math.*, Vol. 120, pp. 371–397, 1984.
- Davis, B.**, *Integral Transforms and Their Applications*, Springer-Verlag, New York, 1978.
- Debnath, L. and Bhatta, B.**, *Integral Transforms and Their Applications, 2nd Edition*, Chapman & Hall/CRC Press, Boca Raton, 2007.
- Delves, L. M. and Mohamed J. L.**, *Computational Methods for Integral Equations*, Cambridge Univ. Press, Cambridge–New York, 1985.
- Demidovich, B. P., Maron, I. A., and Shuvalova E. Z.**, *Numerical Methods. Approximation of Functions and Differential and Integral Equations* [in Russian], Fizmatgiz, Moscow, 1963.
- Ditkin, V. A. and Prudnikov, A. P.**, *Integral Transforms and Operational Calculus*, Pergamon Press, New York, 1965.
- Doetsch, G.**, *Einführung in Theorie und Anwendung der Laplace-Transformation*, Birkhäuser Verlag, Basel–Stuttgart, 1958.
- Doetsch, G.**, *Handbuch der Laplace-Transformation. Anwendungen der Laplace-Transformation*, Birkhäuser Verlag, Basel–Stuttgart, 1956.
- Doetsch, G.**, *Handbuch der Laplace-Transformation. Theorie der Laplace-Transformation*, Birkhäuser Verlag, Basel–Stuttgart, 1950.
- Doetsch, G.**, *Introduction to the Theory and Application of the Laplace Transformation*, Springer-Verlag, Berlin, 1974.
- Duduchava, R.**, *Singular Integral Equations with Fixed Singularities*, Teubner, Leipzig, 1979.
- Dunford, N. and Schwartz J.**, *Linear Operators. Part III. Spectral Operators*, Interscience Publ., New York, 1971.
- Dwight, H. B.**, *Tables of Integrals and Other Mathematical Data*, Macmillan, New York, 1961.
- Dzhuraev, A.**, *Methods of Singular Integral Equations*, J. Wiley, New York, 1992.
- Edwards, C. H. and Penney, D.**, *Calculus, 6th Edition*, Pearson Education, Toronto, 2002.
- Erdogan, F. E.**, *Complex Function Technique*, In: *Continuum Physics* (Ed. A. C. Eringen), Vol. 2, pp. 523–603, Academic Press, New York, 1975.
- Erdogan, F. E., Gupta, G. D., and Cook, T. S.**, The Numerical Solutions of Singular Integral Equations, *Mechanics of Fracture. Vol. 1. Methods of Analysis and Solutions of Crack Problems*, pp. 368–425, Noordhoff Intern. Publ., Leyden, 1973.
- Estrada, R. and Kanwal, R. P.**, *Singular Integral Equations*, Birkhäuser, Boston, 1999.
- Fenyő, S. and Stolle H. W.**, *Theorie und Praxis der Linearen Integralgleichungen, Bd. 3*, Birkhäuser Verlag, Basel, 1984.
- Fock, V. A.**, Some integral equations of mathematical physics, *Doklady AN SSSR*, Vol. 26, No. 4–5, pp. 147–151, 1942.
- Frank-Kamenetskii, D. A.**, *Diffusion and Heat Transfer in Chemical Kinetics* [in Russian], Nauka, Moscow, 1987.
- Gakhov, F. D.**, *Boundary Value Problems* [in Russian], Nauka, Moscow, 1977.
- Gakhov, F. D.**, *Boundary Value Problems*, Dover Publ., New York, 1990.

- Gakhov, F. D. and Cherskii, Yu. I.**, *Equations of Convolution Type* [in Russian], Nauka, Moscow, 1978.
- Gantmakher, F. R.**, Non-symmetric Kellogg Kernels [in Russian], Doklady RAN, Vol. 1, No. 3, 1936.
- Gantmakher, F. R. and Krein, M. G.**, *Oscillating Matrices and Small Vibrations of Mechanical Systems* [in Russian], Gostekhizdat, Moscow, 1950.
- Gohberg, I. C. and Krein, M. G.**, *The Theory of Volterra Operators in a Hilbert Space and Its Applications* [in Russian], Nauka, Moscow, 1967.
- Golberg, A. (Editor)**, *Numerical Solution of Integral Equations*, Plenum Press, New York, 1990.
- Gorenflo, R. and Vessella, S.**, *Abel Integral Equations: Analysis and Applications*, Springer-Verlag, Berlin–New York, 1991.
- Goursat, E.**, *Cours d'Analyse Mathématique, III*, 3^{me} éd., Gauthier–Villars, Paris, 1923.
- Gradshteyn, I. S. and Ryzhik, I. M.**, *Tables of Integrals, Series, and Products, 6th Edition*, Academic Press, New York, 2000.
- Griffith, J. L.**, On the Hankel J -, Y and H transforms, *Proc. Amer. Math. Soc.*, Vol. 9, No. 5, pp. 738–741, 1958.
- Gripenberg, G., Londen, S.-O., and Staffans, O.**, *Volterra Integral and Functional Equations*, Cambridge Univ. Press, Cambridge–New York, 1990.
- Guenther, R. B. and Lee, J. W.**, *Partial Differential Equations of Mathematical Physics and Integral Equations*, Dover Publications, New York, 1996.
- Gupalo, Yu. P., Polyanin, A. D., and Ryzantsev, Yu. S.**, *Mass and Heat Transfer of Reacting Particles with the Flow* [in Russian], Nauka, Moscow, 1985.
- Hackbusch, W.**, *Integral Equations: Theory and Numerical Treatment*, Birkhäuser Verlag, Boston, 1995.
- Hansen, E. R.**, *A Table of Series and Products*, Prentice Hall, Englewood Cliffs, London, 1975.
- Hirschman, I. I. and Widder, D. V.**, *The Convolution Transform*, Princeton Univ. Press, Princeton, New Jersey, 1955.
- Iovane, G., Lifanov, I.K., and Sumbatyan, M.A.**, On direct numerical treatment of hypersingular integral equations arising in mechanics and acoustics, *arXiv:math-ph/0301034 v1 26 Jan 2003*, pp. 1–19, 2003.
- Jentzsch, R.**, Über Integralgleichungen mit positivem Kern, *J. Math.*, Vol. 141, p. 235, 1912.
- Jerry, A. J.**, *Introduction to Integral Equations With Applications*, Marcel Dekker, New York–Basel, 1985.
- Joseph, D. D.**, *Stability of Fluid Motions*, Springer-Verlag, Berlin, 1976.
- Kalandiya, A. I.**, *Mathematical Methods of Two-Dimensional Elasticity*, Mir Publ., Moscow, 1973.
- Kamke, E.**, *Differentialgleichungen: Lösungsmethoden und Lösungen, Bd. 1*, B. G. Teubner, Leipzig, 1977.
- Kantorovich, L. V. and Akilov, G. P.**, *Functional Analysis in Normed Spaces*, Macmillan, New York, 1964.
- Kantorovich, L. V. and Krylov, V. I.**, *Approximate Methods of Higher Analysis*, Interscience Publ., New York, 1958.
- Kanwal, R. P.**, *Linear Integral Equations*, Birkhäuser Verlag, Boston, 1996.
- Karlin, S.**, *Total Positivity, Vol. 1*, Stanford Univ. Press, Stanford, 1968.
- Karon, J. M.**, The sign-regularity properties of a class of Green's functions for ordinary differential equations, *J. Diff. Equations*, Vol. 6, p. 484, 1969.
- Kellogg, O. D.**, Orthogonal functions sets arising from integral equations, *Amer. J. Math.*, Vol. 40, p. 145, 1918.
- Kellogg, O. D.**, The oscillation on functions of an orthogonal set, *Amer. J. Math.*, Vol. 38, p. 1, 1916.
- Kline, M.**, *Calculus: An Intuitive and Physical Approach, 2nd Edition*, Dover Publications, New York, 1998.

- Kolmogorov, A. N. and Fomin, S. V.**, *Elements of the Theory of Functions and Functional Analysis*, Dover Publications, New York, 1999.
- Kolmogorov, A. N. and Fomin, S. V.**, *Introductory Real Analysis*, Prentice Hall, Englewood Cliffs, 1970.
- Kondo, J.**, *Integral Equations*, Clarendon Press, Oxford, 1991.
- Korn, G. A. and Korn, T. M.**, *Mathematical Handbook for Scientists and Engineers, 2nd Edition*, Dover Publications, New York, 2000.
- Krantz, S. G.**, *Handbook of Complex Variables*, Birkhäuser, Boston, 1999.
- Krasnosel'skii, M. A.**, *Topological Methods in the Theory of Nonlinear Integral Equations*, Macmillan, New York, 1964.
- Krasnov, M. L., Kiselev, A. I., and Makarenko, G. I.**, *Problems and Exercises in Integral Equations*, Mir Publ., Moscow, 1971.
- Krasnov, M. L.**, *Integral Equations: Introduction to the Theory* [in Russian], Nauka, Moscow, 1975.
- Krein, M. G. (Editor)**, *Functional Analysis, 2nd Edition* [in Russian], Nauka, Moscow, 1972.
- Krein, M. G.**, Integral equations on a half-line with kernels depending upon the difference of the arguments [in Russian], *Uspekhi Mat. Nauk*, Vol. 13, No. 5 (83), pp. 3–120, 1958.
- Krein, M. G.**, Non-symmetric oscillating Green's functions of ordinary differential operators [in Russian], *Doklady AN SSSR*, Vol. 25, p. 643, 1939.
- Kress, R.**, *Linear Integral Equations, 2nd Edition*, Springer-Verlag, New York, 1999.
- Kress, R.**, *Numerical Analysis*, Springer-Verlag, New York, 1998.
- Krylov, V. I., Bobkov, V. V., and Monastyrnyi, P. I.**, *Introduction to the Theory of Numerical Methods. Integral Equations, Ill-Posed Problems, and Improvement of Convergence* [in Russian], Nauka i Tekhnika, Minsk, 1984.
- Kythe, P. K. and Puri, P.**, *Computational Methods for Linear Integral Equations*, Birkhäuser Verlag, Basel, 2002.
- Ladopoulos, E. G.**, *Singular Integral Equations: Linear and Non-Linear Theory and Its Applications in Science and Engineering*, Springer-Verlag, New York, 2000.
- Lavrentiev, M. M.**, *Some Improperly Posed Problems of Mathematical Physics*, Springer-Verlag, New York, 1967.
- Lavrentiev, M. M., Romanov, V. G., and Shishatskii, S. P.**, *Ill-Posed Problems of Mathematical Physics and Analysis* [in Russian], Nauka, Moscow, 1980.
- Lebedev, N. N.**, *Special Functions and Their Applications*, Prentice Hall, Englewood Cliffs, 1965.
- LePage, W. R.**, *Complex Variables and the Laplace Transform for Engineers*, Dover Publications, New York, 1980.
- Levitan, B. M.**, Hilbert space, In: *SpringerLink Encyclopaedia of Mathematics*, 2001.
- Lifanov, I. K.**, *Singular Integral Equations and Discrete Vortices*, VSP, Amsterdam, 1996.
- Lifanov, I. K., Poltavskii, L. N., and Vainikko, G. M.**, *Hypersingular and Singular Integral Equations and Their Applications*, Chapman & Hall/CRC Press, Boca Raton, 2004.
- Linkov, A. M.**, *Boundary Integral Equations in Elasticity Theory*, Kluwer Academic Publ., Dordrecht, 2002.
- Linz, P.**, *Analytical and Numerical Methods for Volterra Equations*, SIAM, Philadelphia, 1987.
- Lovitt, W. V.**, *Linear Integral Equations*, Dover Publ., New York, 1950.
- Magnus, W., Oberhettinger, F., and Soni, R. P.**, *Formulas and Theorems for the Special Functions of Mathematical Physics, 3rd Edition*, Springer-Verlag, Berlin, 1966.
- Mandal, B. N. and Mandal, N.**, *Advances in Dual Integral Equations*, Chapman & Hall/CRC Press, Boca Raton, 1999.
- Mangulis, V.**, *Handbook of Series for Scientists and Engineers*, Academic Press, New York, 1965.
- Manzhairov, A. V.**, A method of solution of two-dimensional integral equations of axisymmetric contact problems for solids with complex rheology, *Applied Mathematics and Mechanics*, Vol. 49, No. 6, pp. 1019–1025, 1985.

- Manzhurov, A. V.**, Contact problems for inhomogeneous aging viscoelastic solids, in I. I. Vorovich and V. M. Alexandrov (Editors), *Mechanics of Contact Interactions* [in Russian], pp. 549–565, Fizmatlit, Moscow, 2001.
- Manzhurov, A. V.**, Mixed integral equations of contact mechanics and tribology, in N. F. Morozov (Editor), *Mixed Problems of Solid Mechanics. Materials of the Vth Russian Conference with International Participation* [in Russian], pp. 222–226, Izd-vo Saratov. Un-ta, Saratov, 2005.
- Manzhurov, A. V. and Kazakov, K. E.**, Plane and axisymmetric contact problems for viscoelastic aging solids with surface-inhomogeneous coatings, in *Problems in Solid and Rock Mechanics. Collection of Papers Devoted to the 75th Birthday of E. I. Shemyakin* [in Russian], pp. 411–422, Fizmatlit, Moscow, 2006.
- McLachlan, N. W.**, *Bessel Functions for Engineers*, Clarendon Press, Oxford, 1955.
- McLean, W.**, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge Univ. Press, Cambridge, 2000.
- Mikhailov, L. G.**, *Integral Equations With Homogeneous Kernel of Degree -1* [in Russian], Donish, Dushanbe, 1966.
- Mikhlin, S. G. (Editor)**, *Linear Equations of Mathematical Physics*, Holt, Rinehart and Winston, New York, 1967.
- Mikhlin, S. G.**, *Linear Integral Equations*, Hindustan Publ. Corp., Delhi, 1960.
- Mikhlin, S. G. and Prössdorf, S.**, *Singular Integral Operators*, Springer-Verlag, Berlin–New York, 1986.
- Mikhlin, S. G. and Smolitskiy K. L.**, *Approximate Methods for Solution of Differential and Integral Equations*, American Elsevier Publ. Co., New York, 1967.
- Miles, J. W.**, *Integral Transforms in Applied Mathematics*, Cambridge Univ. Press, Cambridge, 1971.
- Moiseyev, N. G., and Popov G. Ya.**, The antiplane problem of a crack with edges touching planes where the constants of elasticity change, *J. Appl. Math. and Mech. (PMM)*, Vol. 58, No. 4, pp. 713–725, 1994.
- Morse, P. M. and Feshbach, H.**, *Methods of Theoretical Physics, Vol. 1*, McGraw-Hill, New York, 1953.
- Muhly, P. S. and Xia, J.**, Calderón–Zygmund operators, local mean oscillation and certain automorphisms of the Toeplitz algebra, *Amer. J. Math.*, Vol. 117, pp. 1157–1201, 1995.
- Müntz, H. M.**, *Integral Equations* [in Russian], GTTI, Leningrad, 1934.
- Murphy, G. M.**, *Ordinary Differential Equations and Their Solutions*, D. Van Nostrand, New York, 1960.
- Muskhelishvili N. I.**, *Singular Integral Equations: Boundary Problems of Function Theory and Their Applications to Mathematical Physics*, Dover Publ., New York, 1992.
- Naidenov, V. I. and Polyaniin, A. D.**, Certain nonisothermal flows of fluid, *J. Appl. Mech. & Tech. Physics*, Vol. 31, No. 3, pp. 419–428, 1990.
- Nasim, C. and Aggarwala, B. D.**, On some dual integral equations, *Indian J. Pure Appl. Math.*, Vol. 15, pp. 323–340, 1984.
- Naylor, D.**, On an integral transform, *Int. J. Math. & Math. Sci.*, Vol. 9, No. 2, pp. 283–292, 1986.
- Nikiforov, A. F. and Uvarov, V. B.**, *Special Functions of Mathematical Physics*, Birkhäuser Verlag, Basel, 1988.
- Nikol'skii S. M.**, *Quadrature Formulas* [in Russian], Nauka, Moscow, 1979.
- Noble, B.**, *Methods Based on Wiener–Hopf Technique for the Solution of Partial Differential Equations*, Pergamon Press, London, 1958.
- Noble, B.**, The solution of Bessel function dual integral equations by a multiplying factor method, *Proc. Camb. Phil. Soc.*, Vol. 59, pp. 351–362, 1963.
- Novikov, S. P., Manakov, S. V., Pitaevskii, L. B., and Zakharov, V. E.**, *Theory of Solitons. The Inverse Scattering Method*, Plenum Press, New York, 1984.
- Oberhettinger, F.**, *Tables of Bessel Transforms*, Springer-Verlag, New York, 1972.

- Oberhettinger, F.**, *Tables of Fourier Transforms and Fourier Transforms of Distributions*, Springer-Verlag, Berlin, 1980.
- Oberhettinger, F.**, *Tables of Mellin Transforms*, Springer-Verlag, New York, 1974.
- Oberhettinger, F. and Badii, L.**, *Tables of Laplace Transforms*, Springer-Verlag, New York, 1973.
- Oldham, K. B. and Spanier, J.**, *The Fractional Calculus*, Academic Press, London, 1974.
- Paley, E. A. C. and Wiener, N.**, *Fourier Transforms in the Complex Domain*, Amer. Math. Soc., New York, 1934.
- Petrov, A. G.**, Asymptotic expansions for thin axisymmetric cavities, *J. Appl. Mech. Tech. Phys.*, Vol. 27, No. 5, pp. 667–672, 1986.
- Petrovskii, I. G.**, *Lectures on the Theory of Integral Equations*, Graylock Press, Rochester, 1957.
- Petrovsky, I. G.**, *Lectures on Partial Differential Equations*, Dover Publications, New York, 1991.
- Pinkus, A. and Zafrany, S.**, *Fourier Series and Integral Transforms*, Cambridge University Press, Cambridge, 1997.
- Pipkin, A. C.**, *A Course on Integral Equations*, Springer-Verlag, New York, 1991.
- Polyanin, A. D. and Manzhirov, A. V.**, *Handbook of Integral Equations, 1st Edition*, CRC Press, Boca Raton, 1998.
- Polyanin, A. D. and Manzhirov, A. V.**, *Handbook of Mathematics for Engineers and Scientists*, Chapman & Hall/CRC Press, Boca Raton, 2007.
- Polyanin, A. D. and Manzhirov, A. V.**, Method of model solutions in the theory of linear integral equations [in Russian], *Doklady AN*, Vol. 354, No. 1, pp. 30–34, 1997.
- Polyanin, A. D. and Sergeev, Yu. A.**, Convective diffusion to a reacting particle in a fluid. Nonlinear surface reaction kinetics, *Int. J. Heat Mass Transfer*, Vol. 23, No. 9, pp. 1171–1182, 1980.
- Polyanin, A. D. and Zaitsev, V. F.**, *Handbook of Exact Solutions for Ordinary Differential Equations, 2nd Edition*, Chapman & Hall/CRC Press, Boca Raton, 2003.
- Polyanin, A. D. and Zaitsev, V. F.**, *Handbook of Nonlinear Partial Differential Equations*, Chapman & Hall/CRC Press, Boca Raton, 2004.
- Polyanin, A. D. and Zhurov, A. I.**, A solution method for some classes of nonlinear integral, integro-functional, and integro-differential equations, From Website *EqWorld—The World of Mathematical Equations*, 2007, <http://eqworld.ipmnet.ru/en/methods/ie/ie-meth3.htm>.
- Porter, D. and Stirling, D. S. G.**, *Integral Equations: A Practical Treatment, from Spectral Theory to Applications*, Cambridge Univ. Press, Cambridge–New York, 1990.
- Poularikas, A. D. (Editor)**, *The Transforms and Applications Handbook, 2nd Edition*, CRC Press, Boca Raton, 2000.
- Precup, R.**, *Methods in Nonlinear Integral Equations*, Springer-Verlag, New York, 2006.
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., and Flannery, B. P.**, *Numerical Recipes in C. The Art of Scientific Computing, 2nd Edition*, Cambridge Univ. Press, Cambridge, 1992.
- Privalov, I. I.**, *Integral Equations* [in Russian], ONTI, Moscow–Leningrad, 1935.
- Prössdorf, S. and Silbermann, B.**, *Numerical Analysis for Integral and Related Operator Equations*, Birkhäuser Verlag, Basel–Boston, 1991.
- Prudnikov, A. P., Brychkov, Yu. A., and Marichev, O. I.**, *Integrals and Series, Vol. 1, Elementary Functions*, Gordon & Breach Sci. Publ., New York, 1986.
- Prudnikov, A. P., Brychkov, Yu. A., and Marichev, O. I.**, *Integrals and Series, Vol. 2, Special Functions*, Gordon & Breach Sci. Publ., New York, 1986.
- Prudnikov, A. P., Brychkov, Yu. A., and Marichev, O. I.**, *Integrals and Series, Vol. 3, More Special Functions*, Gordon & Breach Sci. Publ., New York, 1988.
- Prudnikov, A. P., Brychkov, Yu. A., and Marichev, O. I.**, *Integrals and Series, Vol. 4, Direct Laplace Transforms*, Gordon & Breach, New York, 1992.
- Prudnikov, A. P., Brychkov, Yu. A., and Marichev, O. I.**, *Integrals and Series, Vol. 5, Inverse Laplace Transforms*, Gordon & Breach, New York, 1992.
- Reed, M. and Simon, B.**, *Methods of Modern Mathematical Physics, Vol. 1: Functional Analysis*, Academic Press, New York, 1972.

- Richtmyer, R.**, *Principles of Advanced Mathematical Physics, Vol. 1*, Springer-Verlag, Berlin, 1978.
- Riesz, F. and Sz.-Nagy, B.**, *Functional Analysis*, Ungar, New York, 1955.
- Rooney, P. G.**, A generalisation of an inversion formula for the Gauss transformation, *Can. Math. Bull.*, Vol. 6, No. 1, 1963.
- Rudin, W.**, *Functional Analysis*, McGraw-Hill, New York, 1973.
- Sakhnovich, L. A.**, *Integral Equations With Difference Kernels on Finite Intervals*, Birkhäuser Verlag, Basel–Boston, 1996.
- Samko, S. G.**, *Hypersingular Integrals and Their Applications*, Gordon & Breach Sci. Publ., New York, 2000.
- Samko, S. G., Kilbas, A. A., and Marichev, O. I. (Editors)**, *Fractional Integrals and Derivatives. Theory and Applications*, Gordon & Breach Sci. Publ., New York, 1993.
- Sarason, D.**, Functions of vanishing mean oscillation, *Trans. Amer. Math. Soc.*, Vol. 207, pp. 391–405, 1975.
- Savruk, M. P., Madenci, E., and Shkarayev S.**, Singular integral equations of the second kind with generalized Cauchy-type kernels and variable coefficients, *Int. J. Numer. Meth. Eng.*, Vol. 45, pp. 1457–1470, 1999.
- Savruk, M. P., Osiv, P. N., and Prokopchuk I. V.**, *Numerical Analysis in Plane Problems of Crack Theory* [in Russian], Naukova Dumka, Kiev, 1989.
- Schmeidler, W.**, *Integralgleichungen mit Anwendungen in Physik und Technik*, Leipzig, 1950.
- Sherman, D. I.**, On a singular equation and its application to some problems of elasticity theory, *Izv. AN ArmSSR*, Vol. 22, No. 3, pp. 3–8, 1969.
- Slavyanov, S. Yu. and Lay, W.**, *Special Functions: A Unified Theory Based on Singularities*, Oxford University Press, Oxford, 2000.
- Smirnov, N. S.**, *Introduction to the Theory of Nonlinear Integral Equations* [in Russian], Gostekhizdat, Moscow, 1951.
- Smirnov, V. I.**, *A Course in Higher Mathematics. Vol. 4. Part 1* [in Russian], Nauka, Moscow, 1974.
- Smirnov, V. I. and Lebedev, N. A.**, *Functions of a Complex Variable: Constructive Theory* [in Russian], Nauka, Moscow–Leningrad, 1964.
- Smithies, F.**, *Integral Equations*, Cambridge Univ. Press, Cambridge, 1958.
- Sneddon, I.**, *Fourier Transforms*, Dover Publications, New York, 1995.
- Sneddon, I.**, *The Use of Integral Transforms*, McGraw-Hill, New York, 1972.
- Soni, K.**, Fractional integrals and Hankel transforms, *Duke Math. J.*, Vol. 35, No. 2, pp. 313–319, 1968.
- Srivastava, H. M. and Buschman, R. G.**, *Convolution Integral Equations with Special Function Kernels*, Wiley Eastern, New Delhi, 1977.
- Sullivan, M.**, *Trigonometry, 7th Edition*, Prentice Hall, Englewood Cliffs, 2004.
- Sutton, W. G. L.**, On the equation of diffusion in a turbulent medium, *Proc. Roy. Soc., Ser. A*, Vol. 138, No. 988, pp. 48–75, 1943.
- Sveshnikov, A. G. and Tikhonov, A. N.**, *Theory of Functions of a Complex Variable* [in Russian], Nauka, Moscow, 1970.
- Szegő, G.**, *Orthogonal Polynomials*, AMS, Providence, 1975.
- Theocaris, P. S., and Ioakimidis, N. I.**, A method of numerical solution of Cauchy-type singular integral equations with generalized kernels and arbitrary complex singularities, *J. Comp. Phys.*, Vol. 30, No. 3, pp. 309–323, 1979.
- Theocaris, P. S. and Ioakimidis, N. I.**, Stress intensity factors at the tips of an antiplane shear crack terminating at a bimaterial interface, *Int. J. of Fracture*, Vol. 13, pp. 549–552, 1977.
- Tikhonov, A. N. and Arsenin, V. Ya.**, *Methods for the Solution of Ill-Posed Problems* [in Russian], Nauka, Moscow, 1979.
- Tikhonov, A. N. and Samarskii, A. A.**, *Equations of Mathematical Physics*, Dover Publications, New York, 1990.
- Titchmarsh, E. C.**, Hankel transforms, *Proc. Cambridge Phil. Soc.*, Vol. 21, pp. 463–473, 1923.

- Titchmarsh, E. C.**, *Introduction to the Theory of Fourier Integrals*, 3rd Edition, Chelsea Publishing, New York, 1986.
- Titchmarsh, E. C.**, *Theory of Fourier Integrals*, Oxford Univ. Press, Oxford, 1937.
- Tricomi, F. G.**, *Integral Equations*, Dover Publ., New York, 1985.
- Tslaf, L. Ya.**, *Variational Calculus and Integral Equations* [in Russian], Nauka, Moscow, 1970.
- Tuan, Vu Kim**, Some integral transforms of Fourier convolution type, *Soviet Math. Dokl.*, Vol. 37, No. 3, pp. 669–673, 1988.
- Uflyand, Ya. S.**, *Dual Integral Equations Method in Problems of Mathematical Physics* [in Russian], Nauka, Leningrad, 1977.
- Van der Pol, B. and Bremmer, H.**, *Operational Calculus Based on the Two-Sides Laplace transform*, Cambridge Univ. Press, Cambridge, 1955.
- Vasilieva, A. D. and Tikhonov, A. N.**, *Integral Equations* [in Russian], Izdat. Moskovskogo Universiteta, Moscow, 1989.
- Verlan', A. F. and Sizikov V. S.**, *Integral Equations: Methods, Algorithms, and Programs* [in Russian], Naukova Dumka, Kiev, 1986.
- Vinogradov, I. M. (Editor)**, *Encyclopedia of Mathematics. Vol. 2* [in Russian], Sovetskaya Entsiklopediya, Moscow, 1979.
- Vladimirov, V. S.**, *Equations of Mathematical Physics* [in Russian], Nauka, Moscow, 1981.
- Volterra, V.**, *Theory of Functionals and of Integral and Integro-Differential Equations*, Dover Publ., New York, 1959.
- Vorovich, I. I., Aleksandrov, V. M., and Babeshko, V. A.**, *Nonclassical Mixed Problems in the Theory of Elasticity* [in Russian], Nauka, Moscow, 1974.
- Watson, G. N.**, *A Treatise on the Theory of Bessel Functions*, 2nd Edition, Cambridge Univ. Press, Cambridge, 1952.
- Weisstein, E. W.**, *CRC Concise Encyclopedia of Mathematics*, 2nd Edition, CRC Press, Boca Raton, 2003.
- Whittaker, E. T. and Watson, G. N.**, *A Course of Modern Analysis*, Cambridge Univ. Press, Cambridge, 1958.
- Wiarda, G.**, *Integralgleichungen unter besonderer Berücksichtigung der Anwendungen*, Verlag und Druck von B. G. Teubner, Leipzig und Berlin, 1930.
- Widder, D. V.**, *An Introduction to Transform Theory*, Acad. Press, New York, 1971.
- Widder, D. V.**, The Stieltjes transform, *Trans. Amer. Math. Soc.*, Vol. 43, No. 1, pp. 7–60, 1939.
- Wimp, J.**, Integral transforms involving Whittaker function, *Glasnik Matematički*, Vol. 6, No. 1, pp. 67–70, 1971.
- Yosida, K.**, *Functional Analysis*, 6th Edition, Springer-Verlag, Berlin, 1980.
- Zabreyko, P. P., Koshelev, A. I., et al.**, *Integral Equations: A Reference Text*, Noordhoff Int. Publ., Leyden, 1975.
- Zakharov, V. E. and Shabat, A. B.**, A scheme for the integration of nonlinear evolutionary equations of mathematical physics by the inverse scattering method [in Russian], *Funkts. Analiz i ego Prilozh.*, Vol. 8, No. 3, pp. 43–53, 1974.
- Zill, D. G. and Dewar, J. M.**, *Trigonometry*, 2nd Edition, McGraw-Hill, New York, 1990.
- Zwillinger, D.**, *CRC Standard Mathematical Tables and Formulae*, 31st Edition, CRC Press, Boca Raton, 2002.
- Zwillinger, D.**, *Handbook of Differential Equations*, Academic Press, San Diego, 1989.

Index

A

- Abel equation
 - first kind, 10
 - generalized, 519, 527
 - generalized, first kind, 531
 - generalized, second kind, 141, 548
 - second kind, 138
- Abel problem, 520
- Abel type two-dimensional equation, 15
- absolutely continuous function, 529
- abstract Hilbert space, 873
- Airy equation, 1023
- Airy function, 1023
 - asymptotic expansions, 1023
 - definition, 1023
 - first kind, 1023
 - power series, 1023
 - second kind, 1023
- algebraic equations
 - linear, infinite system, 858, 861, 864, 868, 971
 - linear, infinite system, symmetric matrix, 850, 853
- alternating sums of powers of natural numbers, 920
- alternative, Fredholm, 637, 638, 643
 - symmetric equations, 643
- alternative Fourier transform, 512
- amplitude, 1039
- analysis, functional, 1055
- analytic continuation theorem, 595, 714
- application of integral equations to differential equations, 875
- approach, Carleman–Vekua, 778
- approximate methods
 - nonlinear equations, constant integration limits, 826
 - nonlinear equations, variable integration limit, 811
- approximate solution, 688, 693
- approximate values of eigenvalues, Hilbert–Schmidt kernel, 845
- approximating a kernel, 687
- approximation
 - characteristic values, 646
 - eigenfunctions, Hilbert–Schmidt operator, 868, 872
 - eigenvalues, Hilbert–Schmidt operator, 868, 872
 - kernel, 687
 - Lanczos, 798
 - approximation (*continued*)
 - method, successive, 566, 579, 632, 633, 811, 826, 876
 - solution, 854
- arbitrary functions, 111, 191, 278, 357, 406, 410, 413, 437, 444, 456
- arbitrary parameters, 408, 411, 433, 453
- arbitrary powers, 12, 139, 223, 317, 939, 977
- arccosine, 176, 344
- arccotangent, 178, 347
- arcsine, 177, 345
- arctangent, 178, 346
- argument, complicated, 227, 346, 254
- Arutyunyan equation, 198
- associated Legendre functions, 107, 271, 1031, 1032, 1033
 - first kind, 1032
 - general case, 1032
 - integer indices, real argument, 1031
 - modified, 1033
 - second kind, 1032
- asymmetric form
 - Fourier cosine transform, 514
 - Fourier sine transform, 515
 - Fourier transform, 512
- asymptotic expansions, 509, 1017, 1022–1024, 1035
 - Airy functions, 1023
 - Bessel functions, 1017
 - modified Bessel functions, 1022
 - parabolic cylinder functions, 1035
 - Tricomi confluent hypergeometric functions, 1024
- asymptotic methods, 618
 - equations with logarithmic singularity, 618
- auxiliary conditions, 843, 845, 851, 856, 862, 869, 870
- auxiliary equation, 546, 550, 551, 527
 - application, 527
 - first kind, 550
 - second kind, 551
- auxiliary integral conditions, 841–843
- auxiliary results, 784
- axioms for addition, vectors, 1065
- axioms for addition and multiplication by scalars, vectors, 1065
- axis, real, 575, 713
 - Hölder condition, 575
 - Sokhotski–Plemelj formulas, 713

B

Banach space, 1062, 1065, 1067
 base, Napierian, 906
 base of Napierian logarithm, 905
 base of natural logarithm, 905, 906
 basis
 abstract space, 844, 863
 Euclidean space, 857, 869
 Hilbert space, 857, 867, 869
 Hilbert space, special, 869
 orthonormal, 855, 856
 Bateman method, 689
 general scheme, 689
 special cases, 690
 Bernoulli numbers, 1008
 Bernoulli polynomials, 1052
 Bessel's formula, 1018
 Bessel equation, 1016
 modified, 1021
 Bessel function, 88, 187, 264, 269, 353, 958, 1016
 asymptotic expansions, 1017
 definitions, 1016
 first kind, 261, 297, 1016
 integral representations, 1017
 modified, 97, 189, 269, 355, 1021
 modified, first kind, 266, 1021
 modified, second kind, 266, 1021
 orthogonality properties, 1019
 second kind, 264, 299, 1016
 third kind, 1020
 zeros, 1019
 beta function, 1012, 1014
 incomplete, 1014, 1015
 bifurcation point, nonlinear integral equations, 834, 835
 bilinear series, 640
 iterated kernels, 642
 binomial coefficients, 909, 920, 1007
 Boas transform, 250
 boundary conditions, 887
 boundary value problem
 first, 895, 896
 Hilbert, 742
 linear, representation, 892
 n th-order differential equations, 882
 ODEs, 881
 ODEs, reduction to Fredholm equations, 881
 ODEs, reduction to Volterra equations, 877
 Riemann, 595, 714
 second, 895, 897
 second-order differential equations, 883
 bounded closed domain, 839
 bounded set, 866
 closed, 842
 bounded variation function, 1055, 1058
 classes, 1056
 criteria, 1057
 definition, 1055
 properties, 1056, 1057

Boussinesq equation, 900
 Bubnov–Galerkin method, 697
 Buchholz transform, 274
 Bueckner equation, 801

C

$C(a, b)$, space of continuous functions, 1066
 $C_\alpha(0, 1)$, Hölder space, 1066
 calculation of eigenvalues, 877
 canonical factorization, 680
 canonical form, 805–807
 Hammerstein equation, 807
 canonical function, nonhomogeneous Riemann problem, 605
 Carleman equation, 243, 590
 Carleman method
 characteristic equations, 761
 equation, convolution type, first kind, 606
 equation, convolution type, second kind, 660
 equation, difference kernels, 610
 Carleman–Vekua regularization, 778
 Cauchy criterion, 1067
 Cauchy integral, 708
 Cauchy kernel, 707, 757
 characteristic equation, 761
 complete singular equation, 757
 equation on real axis, 743
 general singular equation, first kind, 745
 generalized, 783
 integral equation, 743, 757
 Cauchy principal value, 709
 Cauchy problem
 first-order ODEs, 875, 876
 ODEs, reduction to integral equations, 875
 second-order ODEs, 876
 special n th-order linear ODE, 876
 Cauchy residue theorem, 504
 Cauchy–Schwarz–Bunyakovsky inequality, 501
 Cauchy type and Fourier integrals, 592
 Cauchy type integral, 708
 Cauchy-type kernel, 751, 753
 characteristic equation, 758, 761
 Cauchy kernel, 761
 exceptional case, 767
 Hilbert kernel, 769
 real axis, 765
 transposed, 758, 764
 characteristic operator, 758
 transposed, 758
 characteristic value, 301, 625, 637, 639, 645, 697
 approximation, 646
 extremal properties, 644
 system, 640
 Chebyshev formula, 535
 Chebyshev functions, 1049
 Chebyshev nodes, 748
 Chebyshev polynomial
 first kind, 109, 1048
 second kind, 750, 1049

- closed-form solution
 case of constant coefficients, 770
 general case, 771
- closed bounded set, 842
- closed domain, bounded, 839
- closed kernel, 578
- coefficient
 binomial, 909, 920, 1007
 discontinuous, 739
 rational, 601, 723
 Riemann problem, 596, 718
 undetermined, 692
- collocation method, 692, 693, 815
 hypersingular integral equation, 755
- collocation points, 693
- combination
 elementary functions, 73, 255
 hyperbolic functions, 39
 trigonometric functions, 63, 252
- compact operator, 842, 843, 1069
 self-adjoint, 843
 self-adjoint positive, 873
 self-adjoint positive definite, 1069
- compact self-adjoint operator, 843
- compact self-adjoint positive definite operator, 1069
 eigenvalues, 1069
- compact self-adjoint positive operator, 873
- compactness of integral operator, sufficient condition, 842
- compatibility condition, 896, 897
- complementary error function, 1009, 1025
- complementary modulus, 1036, 1037
- complete elliptic integral
 first kind, 1035
 second kind, 1035
- complete equation
 generalized Cauchy kernel, 783
 Hilbert kernel, 780
- complete kernel, 578
- complete orthonormal system of functions, 844, 855
- complete singular integral equation, 757, 770, 772
 Cauchy kernel, 757
 Hilbert kernel, 759, 780
 regularization method, 772
 solution methods, 757
- complete space, 1067
- complete system, 1067
- complete system of eigenfunctions, 640
- complex linear space, 1065
- complicated argument, 227, 246, 254
- concentration, 890
 integral equation, 890
 integral equation, numerical method, 891
- condition
 auxiliary integral, 841–843, 845, 851, 856, 862, 869, 870
 boundary, 887
 compatibility, 896, 897
 condition (*continued*)
 Hölder, 709, 1066
 Hölder, real axis, 575
 Lipschitz, 709, 1056, 1059
 normality, 596
 sufficient for compactness of integral operator, 842
- confluent hypergeometric equation, 1024
- confluent hypergeometric function, 107, 1024
 Kummer, 1024
 Tricomi, 1024, 1025
 Tricomi, asymptotic expansions, 1024
 Tricomi, integral representations, 1024
 Whittaker, 1027
 Wronskian, 1026
- conjugate kernels, 582
- connected domain, 731
- constant
 Euler, 533, 1013, 1017, 1026
 Hölder, 709
 eigenfunctions, 696, 699
 resolvent, 633
- continuation
 analytic, 714
- continuity, principle, 714
- continuous function of real argument
 values in Banach space, 840
 values in Hilbert space, 840
 values in space of functions square integrable over a closed bounded set, 842
 values in space of functions square integrable over a ring-shaped domain, 841
 values in space of square integrable functions, 840
- continuous operator, 1068
- contour, smooth, 708
- convergence
 almost everywhere, 1060
 mean-square, 501
- convergent series, 509
- convolution theorem, 507, 513
- convolution type, 574, 606, 660, 669
- coordinate functions, 693, 697
- cosine, 46, 166, 246, 335, 558, 928
 hyperbolic, 22, 154, 238, 327
- cosine integral, 87, 258, 1011
- cosine transform, 514
- cotangent, 62, 175, 252, 343
 hyperbolic, 38, 162, 242, 333
- criterion, Cauchy, 1067
- Crum transform, 268
- curves, open, Riemann problem, 734
- cuspidal point, 708
- cylinder function, 1016
- cylindrical function, 1016
 definitions, 1016

D

- De Moivre formulas, 911
- definite integrals, tables, 951

- definition
 Cauchy type integral, 708
 hyperbolic functions, 913
 degenerate hypergeometric equation, 1024
 degenerate kernel, 111, 191, 278, 357, 519, 522,
 539, 540–543, 569, 573, 589, 625, 627, 631,
 810, 817
 general, 523, 628
 simplest, 627
 density, potential, 893
 derivative
 fractional, definition, 529
 fractional, left-sided, 529
 fractional, properties, 530
 fractional, right-sided, 529
 integrable, fractional, 531
 logarithmic of gamma function, 1017, 1021
 Riemann–Liouville, 529
 determinant, Fredholm, 636
 method, 635
 difference kernel, 114, 203, 283, 372, 519, 524,
 539, 544, 573, 574, 586, 625, 610, 626, 655,
 683
 entire axis, 655
 finite interval, 683
 weak singularity, 588
 differential equation
 n th-order, boundary value problems, 882
 ordinary, 527, 547, 686, 875, 877
 ordinary, linear, 881
 second-order, boundary value problems, 883
 differential equation and Volterra integral
 equations, 877
 differentiating, method for integral equations, 820
 differentiation
 fractional, method, 529
 method, 564, 583, 810
 differentiation formulas, 910, 913, 916, 917
 diffusion flux, integral equations, 890
 digamma function, 1013, 1017
 direct sum of orthogonal subspaces, 845, 863,
 869
 Dirichlet–Mehler integral, 1030
 Dirichlet problem
 exterior, 896
 interior, 895
 reduction to integral equations, 895, 896
 discontinuous coefficient, 739
 divisor transform, 269
 Dixon equation, 136
 domain
 bounded closed, 839
 circular, 841
 multidimensional, 839
 one-dimensional, 839
 ring-shaped, 841, 855, 862
 double layer potential, 893
 Gauss formula, 894
 dual integral equation
 first kind, 295, 575, 610
 first kind, exact solutions, 613
 dual integral equation (*continued*)
 reduction to Fredholm equation, 615
 second kind, 627
 second kind, convolution type, 669
- ## E
- eigenfunctions, 301, 625, 639, 834, 867
 construction, 696, 699
 extremal properties, 644
 Fredholm equation, second kind, 694
 Hilbert–Schmidt kernel, 854, 856, 858, 861,
 864, 865
 Hilbert–Schmidt operator, 871
 kernel, 844
 linear operator, 1068
 nonlinear equation, 834
 nonlinear operator, 834
 system, 640
 system, complete, 640
 system, incomplete, 640
 eigenvalues, 301, 625, 834
 calculation, 877
 compact self-adjoint positive definite operator,
 1069
 Hilbert–Schmidt kernel, 854, 856, 858, 861,
 864, 865
 Hilbert–Schmidt operator, 871
 kernel, 844
 linear operator, 1068
 matrix, 845, 848, 856, 859, 861, 868, 872
 operator, 867
 positive, 648
 self-adjoint operator, 1069
 eigenvectors of matrix, orthonormal, 845, 848,
 856, 859, 868, 872
 eigenvectors of self-adjoint operator, 1069
 electrostatic problem, Roben, 897
 elementary functions, 73, 255, 257, 348, 905
 combinations, 179
 properties, 905
 elements
 linearly dependent, 1065
 linearly independent, 843, 1065
 elliptic function, 1038
 Jacobi, 1039
 Weierstrass, 1042
 elliptic integral, 1035, 1036
 complete, 1035
 complete, first kind, 1035
 complete, second kind, 1035
 first kind, 1037
 incomplete, 1037
 second kind, 1037
 third kind, 1037
 elliptic modulus, 1037
 elliptic theta functions, 1043
 entire axis, equation, 574, 586, 587, 626, 655
 equation
 Abel, first kind, 10
 Abel, generalized, 519, 527
 Abel, generalized, first kind, 531

equation (*continued*)

- Abel, generalized, second kind, 141, 548
- Abel, second kind, 138
- Abel type, first kind, 15
- Abel type, two-dimensional, 15
- Airy, 1023
- Arutyunyan, 198
- auxiliary, 546
- auxiliary, application, 527
- auxiliary, first kind, 550
- auxiliary, second kind, 551
- Bessel, 1016
- Bessel, modified, 1021
- Boussinesq, 900
- Bueckner, 801
- Carleman, 243, 590
- Cauchy kernel, complete, 757
- Cauchy kernel, first kind, 707
- Cauchy kernel, first kind, real axis, 743
- Cauchy kernel, general of first kind, 745
- Cauchy kernel, simplest of first kind, 707, 743
- Cauchy kernel, simplest of first kind, real axis, 743
- characteristic, 758, 761
- characteristic, Cauchy kernel, 761
- characteristic, exceptional case, 767
- characteristic, Hilbert kernel, 769
- characteristic, real axis, 765
- characteristic, transposed, 758, 764
- compact self-adjoint and positive definite operator, 843
- complete, generalized Cauchy kernels, 783
- complete, Hilbert kernel, 780
- complete singular, 757, 770, 772
- complete singular, Cauchy kernel, 757
- complete singular, regularization method, 772
- confluent hypergeometric, 1024
- contain arbitrary functions, 410, 413
- contain arbitrary parameters, 408, 411
- contain modulus, 278, 583
- contain unknown function of complicated argument, 227, 254
- convolution type, first kind, 574
- convolution type, first kind, Carleman method, 606
- convolution type, second kind, 626, 655, 657
- convolution type, second kind, Carleman method, 660
- degenerate kernel, 111, 191, 278, 357, 522, 540–543
- degenerate kernel, nonlinear, method of differentiation, 810
- difference kernel, 114, 203, 283, 372, 524, 544, 574, 586, 626, 685
- difference kernel, Carleman method, 610
- difference kernel, entire axis, 655
- difference kernel, finite interval, 683, 685
- difference kernel, weak singularity, 588
- differential, 875, 877
- differential, n th-order, boundary value problems, 882

equation (*continued*)

- differential, ordinary, 527, 547, 686
- differential, ordinary, linear, 881
- differential, second-order, boundary value problems, 883
- diffusion flux, 890
- Dixon, 136
- dual, first kind, 295, 575, 610
- dual, first kind, exact solutions, 613
- dual, reduction to Fredholm equation, 615
- dual, second kind, 627
- dual, second kind, convolution type, 669
- eigenfunctions, Fredholm equation, second kind, 694
- elasticity, 621
- entire axis, 574, 586, 587, 626, 655
- exact methods, 588–592
- exact solutions, 3–500
- exponential nonlinearity, 411, 467
- finite interval, 683, 685
- finite interval, first kind, 744
- first kind, 3, 519, 591, 624
- first kind, reduction to equations of second kind, 591
- first kind, weak singularity, 574
- Fredholm, degenerate kernel, second kind, 627
- Fredholm, first kind, 573, 623
- Fredholm, second kind, 625, 685, 698, 701
- Fredholm, second kind, system, 701
- Fredholm, second kind on contour, 759
- Fredholm, spectrum, 760
- Fredholm, symmetric kernel, second kind, 639
- Fredholm and dual equations, 615
- Fredholm and Green's function, 881
- function of complicated argument, 246
- Gaussian hypergeometric, 1028
- Gelfand–Levitan–Marchenko, 900
- Gelfand–Levitan–Marchenko type, 898
- general degenerate kernel, 523
- generalized Abel, 519, 527
- generalized Abel, first kind, 531
- generalized Abel, second kind, 141, 548
- generalized Cauchy kernel, complete, 783
- generalized Schlomilch, equation, generalized Schlämilch 254
- Hammerstein, canonical form, 807
- Hammerstein, first kind, 807
- Hammerstein, second kind, 807
- Hammerstein, second kind, degenerate kernel, 817
- Hammerstein type, 807
- Hilbert kernel, complete, 759, 780
- Hilbert kernel, first kind, 707, 746
- Hilbert kernel, general of first kind, 708, 747
- Hilbert kernel, simplest of first kind, 707, 746
- Hilbert kernel, simplest of first kind, complete, 759
- Hilbert–Plessner, 255
- homogeneous, 301, 502, 539, 625, 627, 637, 708, 751
- hyperbolic nonlinearity, 414, 468

equation (*continued*)

hypergeometric, 1028
 hypergeometric, confluent, 1024
 hypergeometric, degenerate, 1024
 hypersingular, Cauchy-type kernel, first kind, 751
 hypersingular, Cauchy-type kernel, general of first kind, 751
 hypersingular, Cauchy-type kernel, simplest of first kind, 231, 751, 753
 hypersingular, collocation method, 755
 hypersingular, Hilbert-type kernel, first kind, 751
 hypersingular, Hilbert-type kernel, general of first kind, 751
 hypersingular, Hilbert-type kernel, simplest of first kind, 255, 754
 hypersingular, numerical methods, 754
 infinite integration limit, first kind, 537
 infinite limits of integration, second kind, 702
 Kadomtsev–Petviashvili, 901
 kernel contains arbitrary functions, 111, 191, 278, 357
 kernel contains arbitrary powers, 12
 kernel contains combinations of elementary functions, 73, 179, 255, 348
 kernel contains combinations of various functions, 565
 kernel contains exponential functions, 15, 144, 231, 320
 kernel contains higher-order polynomials in arguments, 6
 kernel contains hyperbolic functions, 22, 154, 238, 327
 kernel contains inverse trigonometric functions, 66, 176, 344
 kernel contains logarithmic functions, 42, 45, 164, 242, 334
 kernel contains power-law functions, 4, 45, 127, 217, 301
 kernel contains rational functions, 7
 kernel contains special functions, 86, 187, 258, 353
 kernel contains square roots, 9
 kernel contains sum of exponential functions, 564
 kernel contains sum of hyperbolic functions, 564
 kernel contains sum of trigonometric functions, 564
 kernel contains trigonometric functions, 46, 166, 246, 335
 kernel cubic in arguments, 5
 kernel linear in arguments, 4
 kernel quadratic in arguments, 4
 Korteweg–de Vries, 899
 Korteweg–de Vries, modified, 900
 Krein's method, 588
 Lalesco–Picard, 323
 Laplace, 893
 Laplace, potentials, properties, 892

equation (*continued*)

Laplace, potentials, types, 892
 Legendre, 1032
 linear, constant integration limits, 502
 linear, constant integration limits, first kind, 217, 502, 573
 linear, constant integration limits, second kind, 301, 502, 625
 linear, first kind, 502
 linear, operator methods, 549
 linear, second kind, 502
 linear, solution methods, 519, 539, 573, 625
 linear, structure of solutions, 502
 linear, variable integration limit, first kind, 3, 502
 linear, variable integration limit, second kind, 127, 502
 linear and nonlinear PDEs, 898
 logarithmic nonlinearity, 419, 472
 logarithmic singularity, 618
 logarithmic singularity, asymptotic methods, 618
 Mathieu, 1045
 Mathieu, modified, 1046
 method of differentiating, 564, 583, 820
 mixed multidimensional, bounded set, projection method, 866
 mixed multidimensional, closed bounded set, 842
 mixed multidimensional, Fredholm operator, 842
 mixed multidimensional, Hilbert–Schmidt operator, 869
 mixed multidimensional, integral operators of Volterra and Hilbert–Schmidt types, 866
 mixed multidimensional, integral operators of Volterra and Schmidt types, 866
 mixed multidimensional, methods of solving, 839–874
 mixed multidimensional, Schmidt operator, 843
 mixed multidimensional, Schmidt operator, equivalent form, 843
 mixed multidimensional, symmetric Fredholm kernel, 842
 mixed operator, 866, 869
 mixed operator, auxiliary conditions, 869
 mixed two-dimensional, circular domain, 841
 mixed two-dimensional, finite interval, 840
 mixed two-dimensional, finite interval, methods of solving, 843–854
 mixed two-dimensional, Hilbert–Schmidt kernel and auxiliary conditions, finite interval, 845
 mixed two-dimensional, Hilbert–Schmidt kernel and auxiliary conditions, ring-shaped domain, 856
 mixed two-dimensional, Hilbert–Schmidt kernel and given right-hand side, finite interval, 843
 mixed two-dimensional, Hilbert–Schmidt kernel and given right-hand side, ring-shaped domain, 855

equation (*continued*)

mixed two-dimensional, ring-shaped domain, 841
 mixed two-dimensional, ring-shaped domain, methods of solving, 855–866
 mixed two-dimensional, Schmidt kernel, 841
 mixed two-dimensional, Schmidt kernel, equivalent form, 842
 mixed two-dimensional, Schmidt kernel and auxiliary conditions, ring-shaped domain, 862
 mixed two-dimensional, Schmidt kernel and given right-hand side, finite interval, 848
 modified Bessel, 1021
 modified Korteweg–de Vries, 900
 modified Mathieu, 1046
 Nekrasov, 836
 nonhomogeneous, 502, 539, 627, 708, 751
 nonhomogeneous, positive solutions, 649
 nonhomogeneous, solution, 642
 nonlinear, 805, 807, 834, 899
 nonlinear, bifurcation points, 834, 835
 nonlinear, constant integration limits, 806, 829
 nonlinear, constant integration limits, approximate methods, 826
 nonlinear, constant integration limits, exact methods, 817
 nonlinear, constant integration limits, first kind, 433
 nonlinear, constant integration limits, numerical methods, 826
 nonlinear, constant integration limits, second kind, 453
 nonlinear, degenerate kernels, 817
 nonlinear, eigenfunctions, 834
 nonlinear, existence theorems, 830
 nonlinear, uniqueness theorems, 830
 nonlinear, variable integration limit, 805
 nonlinear, variable integration limit, approximate methods, 811
 nonlinear, variable integration limit, exact methods, 809
 nonlinear, variable integration limit, first kind, 393
 nonlinear, variable integration limit, numerical methods, 811
 nonlinear, variable integration limit, second kind, 403
 nonlinear, Volterra, 805
 nonlinear, with parameter, local solutions, 835
 nonlinearity, general form, 399, 425, 447, 477
 nonnegative kernel, 648
 nonsymmetric kernel, first kind, 580
 one-sided, first kind, 574
 one-sided, second kind, 626
 operator, general projection problem, 873
 operator, mixed, 866, 869
 operator, mixed with auxiliary conditions, 869
 operator, “quadratic”, 552
 operator, solution, 553
 ordinary differential, 527, 547, 686

equation (*continued*)

parameter, 625
 Picard–Goursat, 134
 Poisson, 894
 power-law nonlinearity, 408, 464
 power-law nonlinearity that contains arbitrary functions, 444
 quadratic nonlinearity, 819
 quadratic nonlinearity that contains arbitrary functions, 397, 406, 437, 456
 quadratic nonlinearity that contains arbitrary parameters, 393, 403, 433, 453
 “quadratic” operator, 552
 reducible to symmetric equation, 647
 renewal, 203
 right-hand side, 519, 539, 573, 625
 right-hand side, special, 555
 Schlömilch, 254, 452, 825
 Schlömilch, generalized, 254
 Schmidt integral operator, 843
 Schmidt kernel, 843, 859, 863
 Schmidt kernel and auxiliary conditions, finite interval, 851
 Schmidt kernel and auxiliary conditions, ring-shaped domain, 862
 Schmidt kernel and given right-hand side, finite interval, 848
 Schmidt kernel and given right-hand side, ring-shaped domain, 859
 Schmidt operator, 869
 second kind, 591
 second kind, operator method, 654
 semiaxis, 574, 587, 626, 657
 simplest hypersingular, Cauchy-type kernel, first kind, 231, 753
 simplest hypersingular, Hilbert-type kernel, first kind, 255, 754
 single kernel, first kind, 574, 626
 singular, 228, 255, 319, 344, 707
 singular, Bueckner type, 801
 singular, complete, 757, 770, 772
 singular, first kind, 707, 743
 singular, generalized kernel, 792
 singular, numerical solution, 799
 singular, transposed, 758
 singular, two-dimensional, 231
 skew-symmetric, 647
 solution methods, 501–901
 special right-hand side, 555
 surface concentration, 890
 surface concentration, numerical method, 891
 symmetric, 639, 647
 symmetric, Fredholm alternative, 643
 symmetric kernel, 639
 symmetric kernel, first kind, 577
 system, 701
 transposed, 573, 575, 625, 627, 637
 transposed of characteristic equation, 764
 Tricomi, 319, 769, 769
 Tricomi–Gellerstedt, 320
 trigonometric nonlinearity, 420, 473

- equation (*continued*)
- “truncated” first kind, 549
 - two kernels, first kind, 574, 607
 - two kernels, second kind, 626, 664
 - Urysohn, 806, 832
 - Urysohn, first kind, 806, 829
 - Urysohn, first kind, special, method, 821
 - Urysohn, second kind, 806
 - Urysohn, second kind, degenerate kernel, 818
 - Urysohn, second kind, special, method, 822
 - Urysohn type, 806
 - variable integration limit, 3
 - variable lower integration limit, first kind, 537
 - variable lower integration limit, second kind, 570
 - Volterra, 549, 805, 877
 - Volterra, first kind, 519, 524, 565
 - Volterra, first kind, connection with Volterra equations of second kind, 524
 - Volterra, first kind, existence of solution, 519
 - Volterra, first kind, Hammerstein form, 806
 - Volterra, first kind, problems, 520
 - Volterra, first kind, uniqueness of solution, 519
 - Volterra, first kind, Urysohn form, 805, 815
 - Volterra, Hammerstein form, 806
 - Volterra, nonlinear, 805
 - Volterra, quadratic nonlinearity, 809
 - Volterra, reduction to Wiener–Hopf equation, 528
 - Volterra, second kind, 524, 539, 565
 - Volterra, second kind, connection with Volterra equations of first kind, 524
 - Volterra, second kind, Hammerstein form, 816
 - Volterra, second kind, sequence, 855
 - Volterra, second kind, sequence of independent, 853, 865, 872
 - Volterra, second kind, Urysohn form, 805
 - Volterra, sequence, 844, 850, 862
 - Volterra, sequence of independent, 847, 858
 - Volterra, Urysohn form, 805, 811, 814, 816
 - weak singularity, 519
 - weak singularity, first kind, 532, 574
 - weak singularity, second kind, 625
 - weakly singular kernel, 532
 - Whittaker, 1027
 - Wiener–Hopf, 574, 626, 679
 - Wiener–Hopf, first kind, 285, 574, 538, 606
 - Wiener–Hopf, Krein’s method, 679
 - Wiener–Hopf, second kind, 373, 547, 571, 626, 660, 679
 - Wiener–Hopf, second kind, exceptional case, 678
 - Wiener–Hopf, second kind, homogeneous, 672
 - Wiener–Hopf, second kind, index, 661
 - Wiener–Hopf, second kind, nonhomogeneous, 677
 - Wiener–Hopf, second kind, solution, 681
 - Wiener–Hopf, Volterra equation, 528
 - equidistant surface, method, 891
 - equilibrium potential, 897
 - equivalent regularization, problem, 776
 - Erdélyi–Kober operators, 532
 - error function, 86, 258, 549, 1009, 1024
 - complementary, 1009, 1025
 - estimates for spectral radius, 649
 - Euclidean space, 845, 857, 863, 869, 1067
 - basis, 857, 869
 - Euler constant, 533, 1013, 1017, 1026
 - Euler formula, 911, 1013
 - Euler numbers, 1008
 - Euler polynomials, 1053
 - exceptional case
 - characteristic equation, 767
 - regularization, 779
 - Riemann problem, 605, 727
 - Wiener–Hopf equation, second kind, 678
 - existence theorems, 875
 - nonlinear equations, 830
 - Stieltjes integral, 1058
 - expansion, asymptotic, 509
 - Airy functions, 1023
 - Bessel functions, 1017
 - modified Bessel functions, 1022
 - parabolic cylinder functions, 1034
 - Tricomi confluent hypergeometric functions, 1024
 - expansion in power series, 910, 913, 916, 918
 - exponent, growth, 505
 - exponential form, 555
 - exponential function, 15, 73, 77, 78, 144, 151, 179–181, 231, 234, 236, 257, 320, 326, 348, 349, 419, 564, 905, 940, 954, 963, 978, 984, 990, 998, 1002
 - properties, 905
 - exponential integral, 86, 258, 1009, 1010, 1025
 - exponential nonlinearity, 411, 467
 - exponents, singularity, 787, 789
 - expressions with
 - arbitrary powers, 977
 - exponential functions, 963, 978, 984, 990, 998, 1002
 - hyperbolic functions, 964, 979, 985, 991
 - logarithmic functions, 965, 980, 985, 992, 999, 1002
 - power-law functions, 963, 983, 989, 998, 1001
 - rational functions, 971
 - special functions, 967, 981, 987, 993, 1000, 1004
 - square roots, 975
 - trigonometric functions, 966, 981, 986, 992, 999, 1003
 - exterior Dirichlet problem, 896
 - reduction to integral equations, 896
 - exterior Neumann problem, 897
 - reduction to integral equations, 896
- F**
- factorization, 597, 674, 676, 677, 679, 720, 723
 - canonical, 680
 - factorization problem, 676, 679
 - Feller potential, 226

- Feller transform, 226
 field of scalars, 1065
 finite functional sums, 922
 finite interval, 683, 840, 843
 equation, 683, 685
 integrals, 951, 956
 mixed equations, 840
 finite numerical sums, 919
 finite sums, 919
 finitely many singular points, 507
 first-order ODEs, 875, 876
 first boundary value problem, 895, 896
 Fischer–Riesz, theorem, 1062
 flow
 fluid, 888
 nonisothermal in plane channel, 884
 fluid flow, 888
 flux, diffusion integral equations, 890
 form
 canonical, 805–807
 canonical of Hammerstein equation, 807
 equivalent of mixed multidimensional equation
 with Schmidt operator, 843
 equivalent of mixed two-dimensional equation
 with Schmidt kernel, 842
 exponential, 555
 Hammerstein, for Volterra equation, 806
 Hammerstein, for Volterra equation of first
 kind, 806
 Hammerstein, for Volterra equation of second
 kind, 816
 polynomial, 553
 quadratic, 644
 Urysohn, for Volterra equation, 805, 811, 814,
 816
 Urysohn, for Volterra equation of first
 kind, 805, 815
 Urysohn, for Volterra equation of second
 kind, 805
 form of infinite products, representation, 910, 916
 formula
 Bessel's, 1018
 Chebyshev, 535
 Euler, 1013
 Fourier inversion, 512
 Gauss, 535
 Gauss, for double layer potential, 894
 Gauss, for volume potential, 894
 Green's, 895
 Hilbert inversion, 746
 Hopf–Fock, 683
 Kontorovich–Lebedev inversion, 516
 Meijer inversion, 516
 Poincaré–Bertrand, 714
 Poisson's, 1018
 Post–Widder, 510
 quadrature, 534, 815
 Sokhotski–Plemelj, 713, 785
 Stirling, 1013
 formulas
 addition, 909, 915
 formulas (*continued*)
 calculation, 504
 De Moivre, 911
 differentiation, 910, 913, 916, 917
 Euler, 911
 integration, 910, 913, 916, 918
 quadrature, 534, 793
 reduction, 907, 939, 947
 Sokhotski–Plemelj, for real axis, 713
 Fourier cosine transform, 514, 518
 asymmetric form, 514
 Parseval's relation, 514
 tables, 983
 Fourier integral
 left, 594
 one-sided, 593, 594
 relationships with Cauchy type integral, 592
 right, 594
 Fourier inversion formula, 512
 Fourier sine transform, 514, 518
 asymmetric form, 515
 Parseval's relation, 515
 tables, 989
 Fourier transform, 235, 511, 512, 518, 658
 alternative, 512
 asymmetric form, 512
 definition, 512
 inverse, 512
 inversion formula, 512
 properties, 513
 rational, 685
 fractional derivative, 529
 definition, 529
 integrable, 531
 left-sided, 529
 properties, 530
 right-sided, 529
 fractional differentiation, method, 529
 fractional integral
 definition, 529
 left-sided, 529
 properties, 530
 Riemann–Liouville, 529
 right-sided, 529
 fractional integration, 548
 by parts, 529
 operator, 529
 semigroup property, 529
 fractional order, integral, 529
 fractional powers, 138
 fracture mechanics, 791
 Fredholm alternative, 637, 638
 symmetric equations, 643
 Fredholm determinant, 636
 method, 635
 Fredholm equation, 615, 881
 degenerate kernel, second kind, 627
 first kind, 573, 623
 second kind, 625, 685, 698, 701
 second kind, on contour, 759
 second kind, system, 701

- Fredholm equation (*continued*)
 spectrum, 760
 symmetric kernel, second kind, 639
- Fredholm kernel, 573, 625, 839–841
 positive definite, 840
 positive definite, symmetric, 866
 symmetric definite, 840
 symmetric positive, 841
 symmetric positive definite, 866
- Fredholm minor, 636
- Fredholm operator, 758, 842
 symmetric kernel, generalization, 843
- Fredholm theorems, 637, 702, 777
- Fresnel cosine integral, 1012
 generalized, 1012
- Fresnel integrals, 87, 258, 1011, 1012
 generalized, 1012
- Fresnel sine integral, 1012
 generalized, 1012
- Fubini theorem, 1064
- full measure, set, 1060
- function
 absolutely continuous, 529
 Airy, 1023
 arccosine, 66
 arccotangent, 71
 arcsine, 68
 arctangent, 70
 associated Legendre, 107, 271, 1030–1033
 associated Legendre, first kind, 1032
 associated Legendre, general case, 1032
 associated Legendre, integer indices and real argument, 1031
 associated Legendre, second kind, 1032
 Bessel, 88, 187, 264, 269, 353, 958, 1016
 Bessel, asymptotic expansions, 1017
 Bessel, definitions, 1016
 Bessel, first kind, 261, 297, 1016
 Bessel, integral representations, 1017
 Bessel, modified, 97, 189, 269, 355, 1021
 Bessel, modified, first kind, 266, 1021
 Bessel, modified, second kind, 266, 1021
 Bessel, orthogonality properties, 1019
 Bessel, second kind, 264, 299, 1016
 Bessel, third kind, 1020
 Bessel, zeros, 1019
 beta, 1012, 1014
 beta, incomplete, 1014, 1015
 canonical of nonhomogeneous Riemann problem, 605
 Chebyshev, 1049
 complementary error, 1009, 1025
 confluent hypergeometric, 107, 1024
 confluent hypergeometric, Kummer, 1024
 confluent hypergeometric, Tricomi, 1024
 confluent hypergeometric, Whittaker, 1027
 confluent hypergeometric, Wronskian, 1026
 cosine, 46
 cotangent, 62
 cylinder, 1016
 cylindrical, 1016
 function (*continued*)
 digamma, 1013, 1017
 elementary, 73, 179, 255, 257, 348
 elementary, properties, 905
 elliptic, 1038
 elliptic, Jacobi, 1039
 elliptic, Weierstrass, 1042
 elliptic theta, 1043
 error, 86, 258, 549, 1009, 1024
 error, complementary, 1009, 1025
 exponential, 15, 73, 77, 78, 144, 151, 179–181, 213, 234, 236, 257, 320, 326, 348, 349, 419, 564, 905, 940, 954, 963, 978, 984, 990, 998, 1002
 exponential, properties, 905
 gamma, 260, 1012
 gamma, incomplete, 88, 260, 1014, 1024, 1025
 gamma, logarithmic derivative, 1017, 1021
 Gauss hypergeometric, 275, 1028
 generalized Riemann zeta, 277
 generating, 555, 580
 generating, power-law, 557
 generating contain cosines, 558
 generating contain sines, 558
 generating of exponential form, 555
 Green's, 881–883
 Hankel, 1020
 Hankel, first kind, 265
 Hankel, second kind, 265
 harmonic, 893
 Hermite, 1050
 hyperbolic, 22, 73, 83, 84, 154, 164, 179, 185, 186, 238, 255, 327, 334, 348, 351, 352, 564, 911, 913, 922, 940, 955, 964, 979, 985, 991
 hyperbolic, inverse, 917
 hyperbolic, of half argument, 915
 hyperbolic, of multiple argument, 915
 hypergeometric, 1028
 hypergeometric, confluent, 107, 1024
 hypergeometric, confluent, Wronskian, 1026
 hypergeometric, Gauss, 275, 1028
 hypergeometric, Kummer confluent, 272
 hypergeometric, Tricomi confluent, 273, 1025
 hypergeometric, Whittaker confluent, 274, 1027
 incomplete beta, 1014, 1015
 incomplete gamma, 88, 260, 1014, 1024, 1025
 index, 595
 influence, 577, 882
 integrable, 501, 502, 1058
 integrable, Lebesgue, 1059, 1061
 inverse hyperbolic, 917
 inverse trigonometric, 66, 176, 344, 911, 948
 irrational, 937
 Jacobi elliptic, 1039
 Jacobi elliptic, connection with Jacobi theta functions, 1044
 Jacobi theta, 110, 1043
 Jacobi theta, connection with Jacobi elliptic functions, 1044
 Jacobi weight, 793
 Kummer confluent hypergeometric, 272, 1024

function (*continued*)

Lebesgue integrable, 1059, 1061
 left, 594
 Legendre, 270, 1030
 Legendre, associated, 107, 271, 1030–1033
 Legendre, associated, first kind, 1032
 Legendre, associated, second kind, 1032
 Legendre, modified associated, 1033
 Legendre, spherical of first kind, 299
 Legendre, Wronskians, 1034
 logarithmic, 42, 45, 77, 83, 85, 164, 165, 180, 185, 187, 242, 244, 255, 256, 334, 335, 349, 351, 353, 905, 943, 955, 965, 980, 985, 992, 999, 1002
 logarithmic, properties, 906
 MacDonald, 266, 1021
 Mathieu, 1045, 1046
 Mathieu, modified, 1046
 measurable, 1060
 modified associated Legendre, 1033
 modified Bessel, 97, 189, 269, 355, 1021
 modified Bessel, asymptotic expansions, 1022
 modified Bessel, definitions, 1021
 modified Bessel, first kind, 266, 1021
 modified Bessel, integral representations, 1022
 modified Bessel, second kind, 266, 1021
 modified Mathieu, 1046
 multivalued, 711
 Neumann, 1016
 of complicated argument, 227, 346, 254
 one-sided, 594
 parabolic cylinder, 276, 1034
 parabolic cylinder, asymptotic expansions, 1035
 parabolic cylinder, basic formulas, 1034
 parabolic cylinder, definitions, 1034
 parabolic cylinder, integral representations, 1035
 parabolic cylinder, linear relations, 1035
 parabolic cylinder, Weber, 1034
 power, properties, 905
 power-law, 4, 45, 127, 151, 165, 217, 236, 244, 301, 326, 335, 419, 951, 963, 983, 989, 998, 1001
 power-law generating, 557
 psi, 1012, 1013
 rational, 7, 136, 220, 314, 933, 971
 rational, inverse transforms, 506
 Riemann zeta, generalized, 277
 special, 86, 111, 187, 258, 277, 353, 967, 981, 987, 993, 1000, 1004
 special, properties, 1007
 spherical, Legendre of first kind, 299
 square integrable, 501, 502
 Struve, 264, 299, 516, 518
 summable, 1059, 1061
 summable, integral, 1061
 tangent, 60
 theta, Jacobi, 1043
 total variation, 1055
 Tricomi confluent hypergeometric, 273, 1024, 1025

Tricomi confluent hypergeometric, asymptotic expansions, 1024
 Tricomi confluent hypergeometric, integral representations, 1024
 trigonometric, 78, 84, 85, 166, 176, 181, 186, 187, 246, 252, 256, 295, 335, 344, 349, 352, 353, 564, 907, 922, 944, 956, 966, 981, 986, 992, 999, 1003
 trigonometric, inverse, 176, 344, 911, 948
 trigonometric, of half argument, 909
 trigonometric, of multiple arguments, 909
 trigonometric, of single argument, relations, 908
 trigonometric, powers, 908
 Weber, 88
 Weber parabolic cylinder, 1034
 Weierstrass elliptic, 1042
 weight, Jacobi, 793
 Whittaker, 1027
 Whittaker confluent hypergeometric, 274, 1027
 function of bounded variation, 1055
 function of real argument
 values in Banach space, continuous, 840
 values in Hilbert space, continuous, 840
 values in space of functions square integrable functions, continuous, 841
 values in space of functions square integrable over closed bounded set, continuous, 842
 values in space of functions square integrable over ring-shaped domain, continuous, 841
 function of several variables, 839
 functional analysis, some notions, 1055
 functional series, infinite, 925
 functional sums, finite, 922
 functions
 coordinate, 693, 697
 measurable, 1060
 of bounded variation, 1055, 1058, 1066
 orthogonal, 582
 power, 905
 real-valued, multidimensional, 839
 with finitely many singular points, 507
 fundamental solution, 881

G

Galerkin method, 582
 gamma function, 260, 1012
 incomplete, 88, 260, 1014, 1024, 1025
 logarithmic derivative, 1017, 1021
 Gauss formula, 535
 for double layer potential, 894
 for volume potential, 894
 Gauss hypergeometric functions, 275, 1028
 Gauss transform, 237
 Gaussian hypergeometric equation, 1028
 Gegenbauer polynomials, 1051
 Gelfand–Levitan–Marchenko equation, 900
 general degenerate kernel, 523
 general equation of first kind with Cauchy kernel, 745
 general hypersingular equation of first kind with Cauchy-type kernel, finite interval, 751

- general hypersingular equation of first kind with
 Hilbert-type kernel, 751
 general projection problem, 873
 special case, 846, 852, 857, 870
 general scheme
 Bateman method, 689
 method of quadratures, 568
 quadrature method for Fredholm equations of
 second kind, 698
 solving of dual integral equations, 611
 successive approximation method, 566
 general singular equation of first kind with Hilbert
 kernel, 708, 747
 generalization of Fredholm integral operator with
 symmetric kernel, 843
 generalized Abel equation, 519, 527
 first kind, 531
 second kind, 141, 548
 generalized Cauchy kernel, 783
 generalized Fresnel cosine integral, 1012
 generalized Fresnel integral, 1012
 generalized Fresnel sine integral, 1012
 generalized Jentzsch theorem, 648
 generalized kernel of integral equation, 783
 generalized Laguerre polynomials, 1047
 generalized Liouville theorem, 595, 714
 generalized Mehler–Fock transform, 271
 generalized Riemann zeta function, 277
 generalized Schlömilch equation, 254
 generating function, 555, 580
 containing cosines, 558
 containing sines, 558
 exponential form, 555
 power-law, 557
 Green’s formula, 895
 Green’s function, 881–883
 growth exponent, 505
- ## H
- Hammerstein equation, 807, 817, 830
 canonical form, 807
 degenerate kernel, second kind, 817
 first kind, 807
 second kind, 807
 Hammerstein form, Volterra equation, 806
 first kind, 806
 second kind, 816
 Hankel function, 1020
 first kind, 265
 second kind, 265
 Hankel transform, 261, 515, 518
 Parseval’s relation, 515, 516
 Hardy transform, 264
 harmonic function, 893
 Hartley transform, 252, 518
 Hermite functions, 1050
 Hermite interpolation polynomial, 716
 Hermite polynomial, 108, 1024, 1025, 1050
 Hilbert boundary value problem, 742
 Hilbert inversion formula, 746
 Hilbert kernel, 707, 780
 characteristic equation, 769
 complete singular equation, 759, 780
 equation, 759
 equations of first kind, 746
 Hilbert–Plessner equation, 255
 Hilbert problem, 742
 Hilbert–Schmidt kernel, 841, 843, 845, 853, 855,
 856, 860
 approximate values of eigenvalues, 845
 eigenfunctions, 854, 856, 858, 861, 864, 865
 eigenvalues, 854, 856, 858, 861, 864, 865
 Hilbert–Schmidt operator, 842, 843, 866, 871
 approximation for eigenfunctions, 868, 872
 approximation for eigenvalues, 868, 872
 eigenfunctions, 871
 eigenvalues, 871
 Hilbert–Schmidt theorem, 641, 1069
 Hilbert–Schmidt theory, 843
 Hilbert space, 839, 845, 857, 863, 867, 869,
 1067
 abstract, 873
 basis, 857, 867, 869
 linear operators, 1067, 1068
 special basis, 869
 Hilbert transform, 228, 255, 518, 743
 Hilbert transform on semiaxis, 229
 Hilbert-type kernel, 751, 754
 Hölder condition, 709, 1066
 Hölder condition on real axis, 575
 Hölder constant, 709
 Hölder inequality, 1065
 Hölder space $C_\alpha(0, 1)$, 1066
 homogeneous integral equation, 301, 502, 539,
 625, 627, 637, 708, 751
 homogeneous problem, 596, 602, 742
 homogeneous problem solution, 720
 homogeneous Wiener–Hopf equation, second
 kind, 672
 Hopf–Fock formula, 683
 hyperbolic cosine, 22, 154, 238, 327
 hyperbolic cotangent, 38, 162, 242, 333
 hyperbolic function, 22, 73, 83, 84, 154, 179, 185,
 186, 238, 255, 327, 334, 348, 351, 352, 564,
 911, 913, 922, 940, 955, 964, 979, 985, 991
 combinations, 164
 half argument, 915
 multiple argument, 915
 hyperbolic nonlinearity, 414, 468
 hyperbolic sine, 28, 156, 238, 329
 hyperbolic tangent, 36, 161, 241, 332
 hypergeometric equation, 1028
 confluent, 1024
 degenerate, 1024
 hypergeometric function, 1028
 confluent, 107, 1024
 confluent, Kummer, 272, 1024
 confluent, Tricomi, 1024, 1025
 confluent, Whittaker, 274, 1027
 confluent, Wronskian, 1026
 Gauss, 275, 1028

- hypergeometric function (*continued*)
 Gauss, basic properties, 1028
 Kummer confluent, 272, 1024
 Tricomi confluent, 273
 Whittaker confluent, 274, 1027
- hypergeometric series, 1028
- hypersingular equation, 751
 Cauchy-type kernel, 751, 753
 collocation method, 755
 first kind, Cauchy-type kernel on finite interval, 751
 first kind, Hilbert-type kernel, 751
 Hilbert-type kernel, 751, 754
 numerical methods, 754
 simplest of first kind, Cauchy-type kernel, 231, 753
 simplest of first kind, Hilbert-type kernel, 255, 754
- hypersingular integral
 definition, 751
 in sense of Hadamard principal value, 752
- |
- identities, integral, 895
 identity operator, 842, 873
 ill-posed problem, 623
 general notions, 623
 incomplete beta function, 1014, 1015
 incomplete elliptic integrals, 1036
 incomplete gamma function, 88, 260, 1014, 1024, 1025
 incomplete kernel, 578
 incomplete system of eigenfunctions, 640
 indefinite integrals, tables, 933
 independent elements, linearly, 843, 1063
 index, 603, 661, 664
 notion, 716
 index of function, 595
 index of Riemann problem, 596, 731
 index of Wiener–Hopf equation, 661
 inequality
 Cauchy–Schwarz–Bunyakovsky, 501
 Hölder, 1063
 triangle, 501
 infinite functional series, 925
 infinite numerical series, 924
 infinite products, 910, 916
 infinite system of linear algebraic equations, 858, 861, 864, 868, 971
 infinite system of linear algebraic equations with symmetric matrix, 850, 853
 influence function, 577, 882
 inner product, 501, 644
 integrable fractional derivative, 531
 integrable function, 501, 502, 1056
 Lebesgue, 1059
- integral
 Cauchy, 708
 Cauchy type, 708
- integral (*continued*)
 Cauchy type, relationships with Fourier integral, 592
 complete elliptic, 1035
 complete elliptic, first kind, 1035
 complete elliptic, second kind, 1035
 cosine, 87, 258, 1011
 definite, tables, 951
 Dirichlet–Mehler, 1030
 elliptic, 1035, 1036
 elliptic, complete, 1035
 elliptic, first kind, 1036
 elliptic, incomplete, 1036
 elliptic, second kind, 1036
 elliptic, third kind, 1036
 exponential, 86, 258, 1009, 1010, 1025
 Fourier, left, 594
 Fourier, one-sided, 593, 594
 Fourier, relationships with Cauchy type integral, 592
 Fourier, right, 594
 fractional, definition, 529
 fractional, left-sided, 529
 fractional, properties, 530
 fractional, Riemann–Liouville, 529
 fractional, right-sided, 529
 fractional order, 529
 Fresnel, 87, 258, 1011, 1012
 Fresnel, generalized, 1012
 Fresnel cosine, 1012
 Fresnel cosine, generalized, 1012
 Fresnel sine, 1012
 Fresnel sine, generalized, 1012
 hypersingular, definition, 751
 hypersingular, in sense of Hadamard principal value, 752
 incomplete elliptic, 1036
 indefinite, tables, 933
 involving arbitrary powers, 939
 involving Bessel functions, 958
 involving exponential functions, 940, 954
 involving hyperbolic functions, 940, 955
 involving inverse trigonometric functions, 948
 involving irrational functions, 937
 involving logarithmic functions, 943, 955
 involving power-law functions, 951
 involving rational functions, 933
 involving trigonometric functions, 944, 956
 Jacobi weight function, 793
 Laplace, 1030
 Lebesgue, 1057
 Lebesgue, definition, 1059
 Lebesgue, properties, 1059
 left Fourier, 594
 logarithmic, 258, 1009, 1010, 1025
 Mehler, 299, 615
 one-sided Fourier, 593, 594
 probability, 1009
 Riemann, 1057
 Riemann–Liouville fractional, 529
 right-sided fractional, 529

- integral (*continued*)
 right Fourier, 594
 sine, 87, 258, 1011
 singular, 709
 singular, principal value, 709
 step-function, 1059
 Stieltjes, 1055, 1056
 Stieltjes, basic definitions, 1055
 Stieltjes, existence theorems, 1056
 Stieltjes, properties, 1056
 summable function, 1059
 integral conditions, auxiliary, 841–843
 integral equation, *see* equation
 integral identities, 895
 integral operator
 compactness, sufficient condition, 842
 Fredholm, 842
 Fredholm, symmetric kernel, 843
 Hilbert–Schmidt, 842, 843, 866
 positive definite, 842
 positive definite kernel, 843
 Schmidt, 843, 866
 self-adjoint, 842, 843
 spectral radius, 649
 symmetric kernel, 843
 Volterra, 842
 integral representations
 Bessel functions, 1017
 modified Bessel functions, 1022
 parabolic cylinder functions, 1034
 Tricomi confluent hypergeometric functions, 1024
 integral sum, Stieltjes, 1055
 integral transform, *see* transform
 integrand
 contain exponential functions, 419
 contain power-law functions, 419
 nonlinearity, 414–416, 418, 420, 422–424, 467–470, 472–475
 integration
 fractional, 548
 fractional, by parts, 529
 fractional, operator, 529
 fractional, semigroup property, 529
 interior Dirichlet problem, 895
 reduction to integral equations, 895
 interior Neumann problem, 895
 reduction to integral equations, 895
 interpolation nodes, 534
 interpolation polynomial
 Hermite, 716
 Lagrange, 748
 inverse Fourier transform, 512
 inverse hyperbolic functions, 917
 inverse Laplace transforms, tables, 969
 inverse Mellin transform, 510, 1001
 inverse transform
 rational functions, 506
 representation as asymptotic expansions, 509
 representation as convergent series, 509
 inverse trigonometric function, 66, 176, 344, 911, 948
 inversion formula
 Hilbert, 746
 Kontorovich–Lebedev, 516
 Meijer, 516
 inversion of functions with finitely many singular points, 507
 investigation of differential equations, 875
 irrational functions, 937
 iterated kernel, 566, 632
 bilinear series, 642
 iteration process, 811, 814
- ## J
- Jacobi elliptic function, 1038
 connection with Jacobi theta functions, 1042
 Jacobi polynomials, 1049
 Jacobi theta function, 110, 1042
 connection with Jacobi elliptic functions, 1042
 properties, 1042
 relations and formulas, 1042
 series representation, 1042
 Jacobi weight function, 793
 Jentzch theorem, generalized, 648
 Jordan lemma, 505
 jump problem, 596
- ## K
- K -transform, 518
 Kadomtsev–Petviashvili equation, 901
 Kellogg's method for finding characteristic values
 in case of symmetric kernel, 645
 kernel
 approximation, 687
 Cauchy, 707, 757
 Cauchy, characteristic equation, 761
 Cauchy, complete singular integral equation, 757
 Cauchy, generalized, 783
 Cauchy, integral equations, 757
 Cauchy-type, 751, 753
 closed, 578
 complete, 578
 conjugate, 582
 containing arbitrary functions, 111, 191, 278, 357
 containing arbitrary powers, 12, 139, 223, 317
 containing arccosine, 66, 176, 344
 containing arccotangent, 71, 178, 347
 containing arcsine, 68, 177, 345
 containing arctangent, 70, 178, 346
 containing associated Legendre functions, 107, 271
 containing Bessel functions, 88, 187, 353
 containing Bessel functions of first kind, 261, 297
 containing Bessel functions of second kind, 264, 299
 containing Chebyshev polynomials, 109

kernel (*continued*)

containing combination of Bessel and modified Bessel functions, 269

containing combination of Bessel functions, 264

containing combination of elementary functions, 179, 255, 348

containing combination of hyperbolic functions, 39, 164, 334

containing combination of trigonometric functions, 63, 176, 252, 344

containing combination of various functions, 565

containing confluent hypergeometric functions, 107

containing cosine, 46, 166, 246, 335

containing cosine integral, 87

containing cosine integrals, 258

containing cotangent, 62, 175, 252, 343

containing elementary functions, 257

containing error function, 86, 258

containing exponential function, 15, 19, 73, 77, 78, 144, 151, 179–181, 231, 234, 236, 257, 320, 326, 348, 349

containing exponential integral, 86, 258

containing fractional powers, 138

containing Fresnel integral, 87, 258

containing gamma function, 260

containing Gauss hypergeometric function, 275

containing Hermite polynomial, 108

containing higher-order polynomial in arguments, 6, 133, 311

containing hyperbolic cosine, 22, 154, 238, 237

containing hyperbolic cotangent, 38, 162, 242, 333

containing hyperbolic function, 22, 73, 83, 84, 154, 179, 185, 186, 238, 255, 327, 348, 351, 352

containing hyperbolic sine, 28, 156, 238, 329

containing hyperbolic tangent, 36, 161, 241, 332

containing incomplete gamma function, 88, 260

containing integer powers of arguments, 220

containing inverse trigonometric function, 66, 176, 344

containing Jacobi theta functions, 110

containing Kummer confluent hypergeometric function, 272

containing Laguerre polynomial, 110

containing Legendre function, 270

containing Legendre polynomial, 105

containing Legendre spherical function of first kind, 299

containing logarithmic function, 42, 45, 77, 83, 85, 164, 165, 180, 185, 187, 242, 244, 255, 256, 334, 335, 349, 351, 353

containing logarithmic integral, 258

containing modified Bessel function, 97, 189, 355

containing modified Bessel function of first kind, 266

kernel (*continued*)

containing modified Bessel function of second kind, 266

containing other special function, 111, 277

containing parabolic cylinder function, 276

containing power-law function, 4, 19, 45, 127, 151, 165, 217, 236, 244, 301, 326, 335

containing rational function, 7, 136, 220, 314

containing sine, 52, 169, 247, 337

containing sine integral, 87, 258

containing special function, 86, 187, 258, 353

containing square roots, 9, 222

containing square roots powers, 138

containing sum of exponential functions, 564

containing sum of hyperbolic functions, 564

containing sum of trigonometric functions, 564

containing tangent, 60, 174, 251, 342

containing Tricomi confluent hypergeometric function, 273

containing trigonometric function, 46, 78, 84, 85, 166, 181, 186, 187, 246, 256, 295, 335, 349, 352, 353

containing Whittaker confluent hypergeometric function, 274

cubic in arguments, 5, 132, 307

degenerate, 111, 191, 278, 357, 519, 522, 539, 540–543, 569, 573, 589, 625, 627, 631, 810, 817

degenerate, general, 523

degenerate, general case, 628

degenerate, simplest, 627

difference, 114, 203, 283, 372, 519, 524, 539, 544, 573, 574, 586, 610, 625, 626, 655, 683

difference, on entire axis, 655

difference, with weak singularity, 588

eigenfunction, 844

eigenvalue, 844

Fredholm, 573, 625, 839–841

Fredholm, positive definite, 840

Fredholm, positive definite, symmetric, 866

Fredholm, symmetric definite, 840, 841

general degenerate, 523

generalized, 783

Hilbert, 707, 780

Hilbert, characteristic equation, 769

Hilbert, complete singular integral equation, 759, 780

Hilbert, integral equations, 759

Hilbert–Schmidt, 841, 843, 845, 853, 855, 856, 860

Hilbert–Schmidt, approximate values of eigenvalues, 845

Hilbert–Schmidt, eigenfunctions, 854, 856, 858, 861, 864, 865

Hilbert–Schmidt, eigenvalues, 854, 856, 858, 864, 865

Hilbert-type, 751, 754

incomplete, 578

iterated, 566, 632

iterated, bilinear series, 642

linear in arguments, 4, 127, 217, 301

- kernel (*continued*)
 logarithmic, 519, 588
 nondegenerate, 589, 631
 nonnegative, 648
 nonsymmetric, 580, 647
 of integral equation, 519, 573, 625
 of integral transform, 503
 orthogonal, 634
 oscillation, 651
 oscillation, definition, 651
 oscillation, theorems, 651
 polar, 519, 532, 574, 588
 positive definite, 641
 quadratic in arguments, 4, 129, 219, 304
 resolvent, 844
 Schmidt, 582, 841, 848, 851, 859, 860, 862
 simplest degenerate, 627
 singular, weakly, 532
 spectral radius, 649
 stochastic, 654
 symmetric, 573, 577, 625, 639, 645
 symmetric, resolvent, 644
 trace, 646
 transformation, method, 532
 Volterra, 839
 weakly singular, 532
 with logarithmic singularity, 533
 with rational Fourier transforms, 685
 with weak singularity, 519, 532, 574, 588, 625
- Kontorovich–Lebedev inversion formula, 516
 Kontorovich–Lebedev transform, 267, 516, 518
 Korteweg–de Vries equation, 899
 modified, 900
 Krein’s method, 588, 683
 for integral equations, 588
 for Wiener–Hopf equations, 679
 Kummer confluent hypergeometric function, 272, 1024
 Kummer series, 1024
 Kummer transformation, 1025
- L**
- L_2 -norm, 501
 Lagrange interpolation polynomial, 748
 Laguerre polynomial, 110, 1024, 1045
 generalized, 1045
 Lalesco–Picard equation, 323
 Lanczos approximation, 798
 Laplace equation, 893
 potentials, properties, 892
 Laplace integral, 1030
 Laplace transform, 235, 505, 511, 518, 524, 544, 658, 809
 definition, 505
 inverse, tables, 969
 inversion formula, 505
 properties, 507
 solution method, 524
 tables, 961
 two-side, 234, 518
- large λ , solution, 619
 Lavrentiev regularization method, 621
 layer potential, single, 893
 least squares method, 695
 description, 695
 normal system, 695
 Lebedev transform, 269
 Lebesgue integrable function, 1059
 Lebesgue integral, 1057
 definition, 1059
 properties, 1059
 Lebesgue space $L_p(a, b)$, 1064
 Lebesgue theorem on dominated convergence, 1060
 left-sided fractional derivative, 529
 left-sided fractional integral, 529
 left Fourier integral, 594
 left function, 594
 left regularization, 775
 method, 775
 left regularizer, 703
 Legendre equation, 1032
 Legendre functions, 270, 1030
 associated, 107, 271, 1030
 associated, first kind, 1032
 associated, modified, 1033
 associated, second kind, 1032
 modified associated, 1033
 Wronskians, 1034
 Legendre polynomials, 105, 856, 1030
 orthonormal, 844
 Legendre spherical functions, first kind, 299
 lemma, Jordan, 505
 limit theorems, 507
 linear algebraic equations
 infinite system, 858, 861, 864, 868, 971
 infinite system with symmetric matrix, 850, 853
 linear boundary value problems, representation, 892
 linear equation, 898
 constant integration limits, 502
 first kind, 502
 first kind, constant integration limits, 217, 573
 first kind, variable integration limit, 3
 operator methods, 549
 second kind, 502
 second kind, constant integration limits, 301, 625
 second kind, variable integration limit, 127
 solution methods, 519, 539, 573, 625
 structure of solutions, 502
 variable integration limit, 502
 linear normed spaces, 1063
 linear operator, 502, 1066
 eigenfunction, 1066
 eigenvalue, 1066
 linear operators in Hilbert spaces, 1065, 1066
 linear ordinary differential equations, 881
 linear relations of parabolic cylinder functions, 1034

linear space, 1063
 complex, 1063
 real, 1063
 linear superposition principle, 502
 linearly dependent elements, 1063
 linearly independent elements, 843, 1063
 Liouville theorem, generalized, 714
 Lipschitz condition, 709, 1054, 1057
 local solutions of nonlinear integral equation with
 parameter, 835
 logarithm
 Napierian, base, 905
 natural, base, 905
 logarithmic derivative of gamma function, 1017,
 1021
 logarithmic function, 42, 45, 77, 83, 85, 164, 165,
 180, 185, 187, 242, 244, 255, 256, 334, 335,
 349, 351, 353, 905, 943, 955, 965, 980, 985,
 992, 999, 1002
 properties, 906
 logarithmic integral, 258, 1009, 1010, 1025
 logarithmic kernel, 519, 588
 logarithmic nonlinearity, 419, 472
 logarithmic singularity, 533, 618
 kernel, 533
 L_p , spaces, 1062
 $L_p(a, b)$, Lebesgue space, 1064

M

MacDonald function, 266, 1021
 mass transfer to particle in fluid flow complicated
 by surface reaction, 888
 Mathieu equation, 1043
 modified, 1045
 Mathieu function, 1043, 1044
 modified, 1043, 1045
 matrix
 eigenvalues, 845, 848, 856, 859, 861, 868, 872
 eigenvectors, orthonormal, 845, 848, 856, 859,
 868, 872
 orthonormal eigenvectors, 845, 848, 856, 859,
 868, 872
 mean-square convergence, 501
 measurable function, 1058
 measurable set, 1060
 integration, 1061
 measure
 full, set, 1058
 zero, set, 1058
 measure of set, 1061
 mechanics, fracture, 791
 Mehler–Fock transform, 270, 518
 generalized, 271
 Mehler integral, 299, 615
 Meijer inversion formula, 516
 Meijer transform, 266, 516, 517
 Mellin transform, 510, 511, 518, 587, 657, 658
 definition, 510
 inverse, 510
 inverse, tables, 1001

Mellin transform (*continued*)
 inversion formula, 510
 properties, 511
 tables, 997
 method
 approximation, successive, 566
 Bateman, 689
 Bateman, general scheme, 689
 Bateman, special cases, 690
 Bubnov–Galerkin, 697
 Bubnov–Galerkin, description, 697
 Carleman, for characteristic equations, 761
 Carleman, for equations of convolution type of
 first kind, 606
 Carleman, for equations with difference
 kernels, 610
 Carleman, for integral equations of convolution
 type of second kind, 660
 collocation, 692, 693, 815
 collocation, for solving hypersingular integral
 equation, 755
 exact, 588
 Galerkin, 582
 Kellogg's, for finding characteristic values in
 case of symmetric kernel, 645
 Krein's, 588, 683
 Krein's, for integral equations, 588
 Krein's, for Wiener–Hopf equations, 679
 Multhopp–Kalandiya, 747
 Newton–Kantorovich, 813, 814, 827
 Newton–Kantorovich, modified, 814, 827
 nonlinear equations with constant integration
 limits, exact, 817
 nonlinear equations with variable integration
 limit, exact, 809
 operator, 549, 654
 operator, for solving integral equations of
 second kind, 654
 Picard, 876
 projection, for solving mixed equations on
 bounded set, 866
 quadrature, 698, 816, 829
 quadrature, general scheme, 698
 regularization, 704
 regularization, for complete singular integral
 equations, 772
 regularization, for equations with infinite limits
 of integration, 702
 regularization, Lavrentiev, 621
 regularization, Tikhonov, 622, 829
 solution, Laplace transform, 524
 successive approximation, 566, 811, 826
 successive approximation, general scheme, 566
 successive approximation, resolvent, 566
 Tikhonov regularization, 829
 trace, for approximation of characteristic
 values, 646
 Wiener–Hopf, 671
 Wiener–Hopf, scheme, 676
 Zakharov–Shabat, 898

- method based on solution of auxiliary equation, 546
- method for
- solving “quadratic” operator equations, 552
 - special Urysohn equations of first kind, 821
 - special Urysohn equations of second kind, 822
- method of
- approximating kernel by degenerate one, 687
 - differentiating, for integral equations, 820, 564, 583
 - differentiation, 564, 583, 810
 - differentiation, for nonlinear equations with degenerate kernel, 810
 - equidistant surface, 891
 - fractional differentiation, 529
 - fractional integration, for generalized Abel equation, 548
 - Fredholm determinants, 635
 - Fredholm determinants, 635
 - integral transforms, 586, 655, 809, 819
 - least squares, 695
 - least squares, description, 695
 - least squares, normal system, 695
 - left regularization, 775
 - model solutions, 559, 655, 659
 - model solutions, description, 560
 - numerical integration of equation for surface concentration, 891
 - quadratures, 534, 568, 698
 - quadratures, algorithm based on trapezoidal rule, 536
 - quadratures, general scheme, 535, 568
 - quadratures, trapezoidal rule, 568
 - replacing kernel by degenerate kernel, 687
 - right regularization, 775
 - successive approximations, 579, 632, 633, 811, 876
 - successive approximations, for ODEs, 876
 - transformation of kernel, 532
- methods
- approximate, for nonlinear equations with constant integration limits, 826
 - approximate, for nonlinear equations with variable integration limit, 811
 - asymptotic, 618
 - asymptotic, for solving equations with logarithmic singularity, 618
 - exact, for integral equations, 588
 - exact, for nonlinear equations with constant integration limits, 817
 - exact, for nonlinear equations with variable integration limit, 809
 - for solving complete singular integral equations, 757
 - for solving equations with difference kernels on finite interval, 683
 - for solving integral equations, 499
 - for solving linear equations, 519, 539, 573, 625
 - for solving multidimensional mixed integral equations, 839
 - for solving nonlinear integral equations, 805
- methods (*continued*)
- for solving singular integral equations of first kind, 707
 - integral equations of first kind, 707
 - numerical, for hypersingular equations, 754
 - numerical, for nonlinear equations with constant integration limits, 826
 - numerical, for nonlinear equations with variable integration limit, 811
 - of solving mixed integral equations on finite interval, 843
 - of solving mixed integral equations on ring-shaped domain, 855
 - operator, for solving linear integral equations, 549
 - regularization, 621
- minor, Fredholm, 636
- mixed equation, 839
- bounded set, projection method, 866
 - circular domain, 841
 - closed bounded set, 842
 - finite interval, 840
 - Hilbert–Schmidt kernel, finite interval, 843
 - Hilbert–Schmidt kernel, ring-shaped domain and given right-hand side, 855
 - multidimensional, 839
 - multidimensional, solution methods, 839
 - on finite interval, methods of solving, 843
 - on ring-shaped domain, methods of solving, 855
 - ring-shaped domain, 841
 - Schmidt kernel and auxiliary conditions on ring-shaped domain, 862
 - Schmidt Kernel and given right-hand side on interval, 848
- mixed multidimensional equation
- Fredholm operator, 842
 - Schmidt operator, 843
 - Schmidt operator, equivalent form, 843
 - symmetric Fredholm kernel, 842
 - Volterra and Hilbert–Schmidt types operators, 866
 - Volterra and Schmidt types operators, 866
- mixed operator equation, 866, 869
- with given right-hand side, 866
- mixed operator equations with auxiliary conditions, 869
- mixed two-dimensional equation, Schmidt kernel, 841
- Schmidt kernel, equivalent form, 842
- model solution
- cosine-shaped right-hand side, 563
 - exponential right-hand side, 561
 - power-law right-hand side, 562
 - sine-shaped right-hand side, 562
 - method, 559, 655, 659
- modified associated Legendre functions, 1033
- modified Bessel equation, 1021
- modified Bessel function, 97, 189, 269, 355, 1021
- asymptotic expansions, 1022

modified Bessel function (*continued*)
 definitions, 1021
 first kind, 266, 1021
 integral representations, 1022
 second kind, 266, 1021
 modified Korteweg–de Vries equation, 900
 modified Mathieu function, 1043, 1045
 modified Newton–Kantorovich method, 814, 827
 modulus, 278, 583
 complementary, 1036
 elliptic, 1036
 Multhopp–Kalandiya method, 747
 multidimensional domain, 839
 integration, 839
 multidimensional equation, mixed, 839
 Fredholm operator, 842
 integral operators of Volterra and Hilbert–Schmidt types, 866
 integral operators of Volterra and Schmidt types, 866
 Schmidt operator, 843
 solution methods, 839
 symmetric Fredholm kernel, 842
 multidimensional real-valued functions, 839
 multiply connected domain, 731
 multivalued functions, 711

N

Napierian base, 906
 Napierian logarithms, base, 905
 natural logarithms, base, 905
 natural numbers, powers, sums, 919
 Nekrasov equation, 836
 Neumann function, 1016
 Neumann problem
 exterior, reduction to integral equations, 896
 interior, 895
 interior, reduction to integral equations, 895
 Neumann series, 567, 633
 Newton–Kantorovich method, 813, 814, 827
 modified, 814, 827
 nodes
 Chebyshev, 748
 interpolation, 534
 quadrature, 534
 nondegenerate kernel, 589, 631
 nonhomogeneous equation, 502, 539, 627, 708, 751
 positive solutions, 649
 solution, 642
 nonhomogeneous problem, 604, 742
 solution, 721
 nonhomogeneous Riemann problem, canonical function, 605
 nonhomogeneous Wiener–Hopf equation of second kind, 677
 nonisothermal flow in plane channel, 884
 nonlinear equation, 807, 834, 899
 bifurcation points, 834, 835
 constant integration limits, 806, 829

nonlinear equation (*continued*)
 constant integration limits, approximate methods, 826
 constant integration limits, exact methods, 817
 constant integration limits, numerical methods, 826
 degenerate kernel, 817
 degenerate kernel, method of differentiation, 810
 eigenfunctions, 834
 existence theorems, 830
 first kind with constant limits of integration, 433
 parameter, local solutions, 835
 second kind with variable limit of integration, 403
 second kind with constant limits of integration, 453
 solution methods, 805
 uniqueness theorems, 830
 variable limit of integration, 805
 variable limit of integration, approximate methods, 811
 variable limit of integration, exact methods, 809
 variable limit of integration, numerical methods, 811
 nonlinear operator, eigenfunctions, 834
 nonlinear PDEs, 898
 nonlinear problem of nonisothermal flow in plane channel, 884
 nonlinear Volterra integral equation, 805
 nonlinearity, 414–416, 418, 467–470, 472–475
 exponential, 411, 467
 general form, 399, 425, 447, 477
 hyperbolic, 414, 468
 logarithmic, 419, 472
 power-law, 408, 444, 464
 quadratic, 393, 397, 403, 406, 437, 453, 456
 trigonometric, 420, 473
 nonnegative kernels, 648
 nonorthogonal polynomials, 1050
 nonsymmetric kernel, 580, 647
 norm, 501, 644, 839
 L_2 , 501
 operator, 1066
 normal system of method of least squares, 695
 normality condition, 596
 normed space, 1063
 linear, 1063
 notion of almost everywhere, 1058
 notion of index, 716
 n th-order differential equations, boundary value problems, 882
 n th-order linear ODE, 876
 number e , 905, 906
 numbers, 1007
 Bernoulli, 1008
 Euler, 1008
 natural, powers, sums, 919
 numerical integration, method, 891

numerical methods for hypersingular equations, 754
 numerical methods for nonlinear equations with constant integration limits, 826
 numerical methods for nonlinear equations with variable limit of integration, 811
 numerical series, 924
 infinite, 924
 numerical solution, singular equations, 799
 generalized kernels, 792
 numerical sums, 921
 finite, 919

O

ODE

first-order, 875, 876
 method of successive approximations, 876
 n th-order, linear, 876
 second-order, 876

Olevskii transform, 276

one-dimensional domain, 839

 integration, 839

one-sided equation, 574, 626

one-sided Fourier integrals, 593, 594

one-sided function, 594

open curves, 734

 Riemann problem, 734

operator

compact, 842, 843, 1067
 compact, self-adjoint, 843
 compact, self-adjoint positive, 873
 compact, self-adjoint positive definite, 1067
 compact, self-adjoint positive definite, eigenvalues, 1067
 Erdélyi–Kober, 532
 Fredholm, 758, 842
 Fredholm, symmetric kernel, generalization, 843
 Hilbert–Schmidt, 842, 843, 866, 871
 Hilbert–Schmidt, approximation for eigenfunctions, 868, 872
 Hilbert–Schmidt, approximation for eigenvalues, 868, 872
 Hilbert–Schmidt, eigenfunction, 871
 Hilbert–Schmidt, eigenvalues, 871
 identity, 842, 873
 integral, characteristic, 758
 integral, characteristic, transposed, 758
 integral, compactness, sufficient condition, 842
 integral, continuous, 1066
 integral, domain, 1066
 integral, domain of definition, 1066
 integral, eigenvalues, 867
 integral, positive definite, 842
 integral, self-adjoint, 842, 843, 1067
 integral, self-adjoint, eigenvalues, 1067
 integral, self-adjoint, eigenvectors, 1067
 integral, spectral radius, 649
 integral, spectrum, 1066
 integral, transposed, 758
 integral, transposed characteristic, 758

operator (*continued*)

 integral with positive definite kernel, 843
 integral with symmetric kernel, 843
 linear, 502, 1066
 linear, eigenfunction, 1066
 linear, eigenvalue, 1066
 linear in Hilbert spaces, 1065, 1066
 nonlinear, eigenfunctions, 834
 norm, 1066
 orthogonal projection, 1067
 point, continuous, 1066
 positive definite, 842, 1067
 regular, 758
 regularizing, 703
 Schmidt, 843, 866
 singular, 758
 singular, certain properties, 772
 Volterra, 842, 873

operator equation

 general projection problem, 873
 general projection problem, 873
 mixed, 866, 869
 mixed with auxiliary conditions, 869
 “quadratic”, 552
 solution, 553

operator method, 549, 654

operator method for solving integral equations of second kind, 654

operator of fractional integration, 529

operator of orthogonal projection, 846, 852, 857, 563, 870

order, fractional, integral, 529

ordinary differential equations, 527, 547, 686
 linear, 881

orthogonal function, 582

orthogonal kernels, 634

orthogonal polynomials, 1045

 system, 795

orthogonal projection, operator, 846, 852, 857, 563, 870, 1067

orthogonal projector, 1067

orthogonal subspaces, 873

 direct sum, 845, 863, 869

orthogonal system, 1065

orthogonal vectors, 1065

orthogonality properties of Bessel functions, 1019

orthonormal basis, 855, 856

orthonormal eigenvectors of matrix, 845, 848, 856, 859, 868, 872

orthonormal Legendre polynomials, 844

orthonormal system, 1065

 complete, 844, 855

oscillation kernel, 651

 definition, 651

 theorems, 651

P

\wp -function, Weierstrass, 1041

Paley–Wiener transform, 260

- parabolic cylinder function, 276, 1034
 asymptotic expansions, 1034
 basic formulas, 1034
 definitions, 1034
 integral representations, 1034
 linear relations, 1034
 Weber, 1034
- parameter of integral equation, 625
- parameters, arbitrary, 408, 411, 433, 453
- Parseval's relation
 Fourier cosine transform, 514
 Fourier sine transform, 515
 Hankel transform, 515, 516
- particular solutions of PDEs, 887
- PDEs, nonlinear, 898
- PDEs with boundary conditions
 third kind, 887
 third kind, reduction to integral equations, 887
- permutator, 654
- Picard–Goursat equation, 134
- Picard method, 876
- Pochhammer symbol, 1007
- Poincaré–Bertrand formula, 714
- point
 bifurcation, 835
 bifurcation of nonlinear integral equations, 834, 835
 collocation, 693
 cuspidal, 708
 regular, 1066
 singular, 507
- point operator, continuous, 1066
- Poisson's formula, 1018
- Poisson equation, 894
- polar kernel, 519, 532, 574, 588
- polynomial
 Bernoulli, 1050
 Chebyshev, 109, 1047
 Chebyshev, second kind, 750
 Euler, 1051
 Gegenbauer, 1050
 generalized Laguerre, 1045
 Hermite, 108, 1024, 1025, 1048
 higher-order in arguments, 6, 133, 311
 interpolation, Hermite, 716
 interpolation, Lagrange, 748
 Jacobi, 1049
 Lagrange interpolation, 748
 Laguerre, 110, 1024, 1045
 Laguerre, generalized, 1045
 Legendre, 105, 856, 1030
 Legendre, orthonormal, 844
 nonorthogonal, 1050
 orthogonal, 1045
 orthogonal, system, 795
 orthonormal Legendre, 844
 ultraspherical, 1050
- polynomial form, 553
- positive definite Fredholm kernel, 840
 symmetric, 866
- positive definite integral operator, 842
- positive definite kernel, 641
- positive definite operator, 1067
- positive eigenvalue, 648
- positive Fredholm kernel, symmetric, 841
- positive solutions of nonhomogeneous integral equation, 649
- Post–Widder formula, 510
- potential
 density, 893
 double layer, 893
 double layer, Gauss formula, 894
 equilibrium, 897
 Feller, 226
 Laplace equation, 892
 Laplace equation, properties, 892
 layer, single, 893
 Riesz, 226
 Roben, 897
 single layer, 893
 volume, 893
 volume, Gauss formula, 894
- power-law functions, 4, 45, 127, 151, 165, 217, 236, 244, 301, 326, 335, 419, 951, 963, 983, 989, 998, 1001
- power-law generating function, 557
- power-law nonlinearity, 408, 464
- power-law nonlinearity that contain arbitrary functions, 444
- power function, 905
 properties, 905
- power series, 925
 expansion, 910, 913, 916, 918
- power series in parameter, 632
- power series of Airy functions, 1023
- powers, arbitrary, 139, 223, 317, 939, 977
- powers, fractional, 138
- powers of natural numbers, sums, 919
- principal value
 curvilinear integral, 712
 singular curvilinear integral, 712
 singular integral, 709
- principle
 linear superposition, 502
 superposition, linear, 502
- principle of argument, 714
- principle of continuity, 714
- probability integral, 1009
- problem
 Abel, 520
 boundary value, first, 895, 896
 boundary value, for n th-order differential equations, 882
 boundary value, for ODEs, 877, 881
 boundary value, for second-order differential equations, 883
 boundary value, linear, representation, 892
 boundary value, Riemann, 595
 boundary value, second, 895, 897
 Cauchy, for ODEs, reduction to integral equations, 875
 Cauchy, for second-order ODEs, 876

- problem (*continued*)
 Cauchy, for special n th-order linear ODE, 876
 Dirichlet, exterior, reduction to integral equations, 896
 Dirichlet, interior, 895
 Dirichlet, interior, reduction to integral equations, 895
 electrostatic, Roben, 897
 factorization, 676, 679
 general projection, 873
 general projection, for operator equation, 873
 general projection, special case, 846, 852, 857, 870
 Hilbert, 742
 Hilbert, boundary value, 742
 homogeneous, 596, 602, 742
 homogeneous, solution, 720
 ill-posed, 623, 624
 ill-posed, general notions, 623
 interior Dirichlet, 895
 interior Dirichlet, reduction to integral equations, 895
 interior Neumann, 895
 interior Neumann, reduction to integral equations, 895
 jump, 596
 linear boundary value, representation, 892
 Neumann, exterior, reduction to integral equations, 896
 Neumann, interior, 895
 Neumann, interior, reduction to integral equations, 895
 nonhomogeneous, 604, 742
 nonhomogeneous, solution, 721
 nonhomogeneous Riemann, canonical function, 605
 nonlinear of nonisothermal flow in plane channel, 884
 projection, general, for operator equation, 873
 projection, general, special case, 846, 852, 857, 870
 Riemann, 596, 685, 714
 Riemann, boundary value, 595
 Riemann, coefficient, 596, 718
 Riemann, discontinuous coefficient, 739
 Riemann, exceptional cases, 727
 Riemann, for half-plane, 725
 Riemann, for open curves, 734
 Riemann, for real axis, 592
 Riemann, general case, 741
 Riemann, index, 596, 731
 Riemann, multiply connected domain, 731
 Riemann, nonhomogeneous, canonical function, 605
 Riemann, open curves, 734
 Riemann, right-hand side, 596, 718
 Riemann, statement, 718
 Riemann, with discontinuous coefficient, 739
 Riemann, with rational coefficients, 723
 Roben electrostatic, 897
 second boundary value, 895, 897
- problem (*continued*)
 tautochrone, 520
 well-posed, 623
 well-posed, general notions, 623
 problem of equivalent regularization, 776
 problem with rational coefficients, 601
 process, iteration, 811, 814
 product
 infinite, 910, 916
 inner, 501, 644
 scalar, 839
 progressions, 919, 924
 projection, orthogonal, operator, 846, 852, 857, 563, 870
 projection method for solving mixed equations on bounded set, 866
 projection problem
 general, for operator equation, 873
 general, special case, 846, 852, 857, 870
 projector, orthogonal, 1067
 properties
 basic of Gauss hypergeometric functions, 1028
 certain of singular operators, 772
 orthogonality of Bessel functions, 1019
 property, semigroup of fractional integration, 529
 psi function, 1012, 1013
- ## Q
- quadratic form, 644
 quadratic nonlinearity, 393, 397, 403, 406
 containing arbitrary functions, 437, 456
 containing arbitrary parameters, 433, 453
 quadrature formula, 534, 793, 815
 quadrature method, 698, 816, 829
 general scheme, 698
 quadrature nodes, 534
 quadratures, method, 534, 568, 698
 method, algorithm based on trapezoidal rule, 536
 method, general scheme, 535
- ## R
- radius
 spectral, estimates, 649
 spectral, of integral operator, 649
 spectral, of kernel, 649
 rational coefficients, 601, 723
 rational Fourier transforms, 685
 rational functions, 7, 136, 220, 314, 933, 971
 inverse transforms, 506
 reaction, surface, 888
 real-valued functions, multidimensional, classes, 839
 real axis
 Hölder condition, 575
 Sokhotski–Plemelj formulas, 713
 real linear space, 1063
 rectangle rule, 534
 recurrent relations, 636

reduction formulas, 907, 939, 947
 regular operator, 758
 regular points, 1066
 regular value, 301, 625, 637
 regularization, 774
 Carleman–Vekua, 778
 equivalent, problem, 776
 left, 775
 left, method, 775
 right, 776
 right, method, 775
 regularization in exceptional cases, 779
 regularization method, 621, 704
 complete singular integral equations, 772
 equations with infinite limits of integration, 702
 Lavrentiev, 621
 Tikhonov, 622, 829
 regularizer, 774
 left, 703
 right, 704
 regularizing operators, 703
 relation
 linear of parabolic cylinder functions, 1034
 Parseval's, Fourier cosine transform, 514
 Parseval's, Fourier sine transform, 515
 Parseval's, Hankel transform, 515, 516
 recurrent, 636
 relations between Mellin, Laplace, and Fourier transforms, 511
 remainder, 534
 renewal equation, 203
 representation
 Bessel functions, 1017
 form of infinite products, 910, 916
 Gauss hypergeometric functions, 1028
 inverse transforms as asymptotic expansions, 509
 inverse transforms as convergent series, 509
 modified Bessel functions, 1022
 parabolic cylinder functions, 1034
 series of Jacobi theta functions, 1042
 Tricomi confluent hypergeometric functions, 1024
 residual, 692
 residue theorem, Cauchy, 504
 residues, 504
 resolvent, 539, 567, 626, 633, 635
 construction, 633
 kernel, 844
 symmetric kernel, 644
 results, auxiliary, 784
 Riemann boundary value problem, 595, 714
 Riemann integral, 1057
 Riemann–Liouville derivatives, 529
 Riemann–Liouville fractional integrals, 529
 Riemann problem, 596, 685, 714
 coefficient, 596, 718
 exceptional cases, 727
 for half-plane, 725
 for multiply connected domain, 731
 for open curves, 734

Riemann problem (*continued*)
 for real axis, 592
 general case, 741
 index, 596, 731
 nonhomogeneous, canonical function, 605
 right-hand side, 596, 718
 statement, 718
 with discontinuous coefficient, 739
 with rational coefficients, 723
 Riemann zeta function, generalized, 277
 Riesz potential, 226
 Riesz–Schauder theory, 843
 Riesz transform, 226
 right-hand side, 757
 equation, 519, 573, 625
 integral equation, 539
 Riemann problem, 596, 718
 special, 555
 right-sided fractional derivative, 529
 right-sided fractional integral, 529
 right Fourier integral, 594
 right function, 594
 right regularization, 776
 method, 775
 right regularizer, 704
 ring-shaped domain, 841, 855, 862
 Roben electrostatic problem, 897
 Roben potential, 897
 roots, square, 138, 222, 975
 rule
 rectangle, 534
 Simpson's, 534
 trapezoidal, 534, 568

S

scalar, 1063
 scalar product, 839
 scalars, field, 1063
 scheme
 general, Bateman method, 689
 general, method of quadratures, 568
 general, successive approximation method, 566
 Schlömilch equation, 254, 452, 825
 generalized, 254
 Schmidt integral operator, 843, 866
 Schmidt kernel, 582, 841, 848, 851, 859, 860, 862
 Schmidt operator, 866
 second-order differential equations, boundary value problems, 883
 second-order ODEs, 876
 second boundary value problem, 895, 897
 segment, finite, equation, 683, 685
 self-adjoint operator, 842, 843, 1067
 eigenvalues, 1067
 eigenvectors, 1067
 semiaxis
 equation, 574, 587, 626, 657
 Hilbert transform, 229
 semigroup property of fractional integration, 529

- sequence of independent Volterra equations, 847, 858
- sequence of independent Volterra equations of second kind, 853, 865, 872
- sequence of Volterra equations, 844, 850, 862
- sequence of Volterra equations of second kind, 855
- series
- bilinear, 640
 - bilinear, iterated kernels, 642
 - convergent, 509
 - functional, infinite, 925
 - hypergeometric, 1028
 - infinite, 919
 - infinite functional, 925
 - infinite numerical, 924
 - Kummer, 1024
 - Neumann, 567, 633
 - numerical, 924
 - numerical, infinite, 924
 - power, 913, 925
 - power, expansion, 910, 916, 918
 - power in parameter, 632
 - power of Airy functions, 1023
 - trigonometric, in one variable, involving cosine, 928
 - trigonometric, in one variable, involving sine, 927
 - trigonometric, in two variables, 930
- series representation of Jacobi theta functions, 1042
- set, 866
- bounded, closed, 842
 - closed bounded, 842
 - measurable, 1060
 - measure, 1061
- set of full measure, 1058
- set of zero measure, 1058
- sets, measurable, 1060
- measurable, integration, 1061
 - zero measure, 1058
- several variables, function, 839
- side
- right-hand, 757
 - right-hand, of equation, 519, 573, 625
 - right-hand, of integral equation, 539
 - right-hand, of Riemann problem, 596
 - right-hand, of Riemann problem, 718
 - right-hand, special, 555
- simple hypersingular equation of first kind with Cauchy-type kernel, 231
- simple hypersingular equation of first kind with Hilbert-type kernel, 255
- simplest degenerate kernel, 627
- simplest equation with Cauchy kernel, 743
- simplest hypersingular equation for first kind with Hilbert-type kernel, 754
- simplest singular equation of first kind with Hilbert kernel, 707, 746
- Simpson's rule, 534
- sine, 52, 169, 247, 337, 558, 927
- hyperbolic, 28, 156, 238, 329
- sine integral, 87, 258, 1011
- sine transform, Fourier, Parseval's relation, 515
- single layer potential, 893
- singular curvilinear integral, principal value, 712
- 228, 255, 319, 344
 - Bueckner type, 801
 - Cauchy kernel, complete, 757
 - Cauchy kernel, first kind, 707
 - complete, 757, 770, 772
 - first kind, 743
 - generalized kernels, 792
 - generalized kernels, direct numerical solution, 792
 - Hilbert kernel, 759
 - Hilbert kernel, complete, 759, 780
 - numerical solution, 799
 - simplest of first kind with Hilbert kernel, 707, 746
 - transposed, 758
 - two-dimensional, 231
- singular equations of first kind, 707
- singular integral, 709
- principal value, 709, 712
- singular kernel, weakly, 532
- singular operator, 758
- singular operators, certain properties, 772
- singular points, 507
- singularities, solutions, 783
- singularity
- logarithmic, 533, 618
 - logarithmic, kernel, 533
 - weak, 574, 588, 625
 - weak, kernel, 519, 532, 574, 588, 625
- singularity exponents, 787, 789
- skew-symmetric integral equation, 647
- small λ solution, 620
- smooth contour, 708
- Sokhotski–Plemelj formula, 713, 785
- Sokhotski–Plemelj formulas for real axis, 713
- solution
- approximate, 688, 693
 - approximation, 854
 - convolution representation, 526
 - direct numerical of singular integral equations with generalized kernels, 792
 - exact of simple hypersingular equation with Cauchy-type kernel, 753
 - exact of simple hypersingular equation with Hilbert-type kernel, 754
 - fundamental, 881
 - homogeneous problem, 720
 - integral equations, exact, 1–500
 - model, cosine-shaped right-hand side, 563
 - model, exponential right-hand side, 561
 - model, power-law right-hand side, 562
 - model, sine-shaped right-hand side, 562
 - nonhomogeneous problem, 721
 - numerical, of singular integral equations, 799

- solution (*continued*)
- simple hypersingular equation with Cauchy-type kernel, exact, 753
 - simple hypersingular equation with Hilbert-type kernel, exact, 754
 - stable, 623
 - trivial, 502
- solution method, Laplace transform, 524
- solution method based on Laplace transform, 544
- solution of auxiliary equation, method, 546
- solution of generalized Abel equation, 531
- solution of operator equations of polynomial form, 553
- solutions
- closed-form, case of constant coefficients, 770
 - closed-form, general case, 771
 - fundamental, 881
 - local of nonlinear integral equation with parameter, 835
 - model, method, 559, 655, 659
 - particular of PDEs, 887
 - positive of nonhomogeneous integral equation, 649
- solutions of dual integral equations, general scheme, 611
- solutions of nonlinear PDEs, representation in terms of solutions of linear integral equations, 898
- solutions singularities, 783
- solving linear equations, methods, 519, 539
- solving “quadratic” operator equations, 552
- Sonine transform, 114
- space
- Banach, 1065
 - basis, 844, 863
 - complete, 1065
 - complex linear, 1063
 - Euclidean, 845, 857, 863, 869, 1065
 - Euclidean, basis, 857, 869
 - Hilbert, 839, 845, 857, 863, 867, 869, 1065
 - Hilbert, abstract, 873
 - Hilbert, basis, 857, 867, 869
 - Hilbert, linear operators, 1065, 1066
 - Hilbert, special basis, 869
 - Hölder $C_\alpha(0, 1)$, 1064
 - Lebesgue $L_p(a, b)$, 1064
 - linear, 1063
 - linear, complex, 1063
 - linear, normed, 1063
 - linear, real, 1063
 - normed, 1063
 - normed linear, 1063
 - real linear, 1063
 - vector, 1063
- space L_p , 1062
- space of continuous functions $C(a, b)$, 1064
- space of functions of bounded variation $V(0, 1)$, 1064
- special basis of Hilbert space, 869
- special case of general projection problem, 846, 852, 857, 870
- special functions, 86, 111, 187, 258, 277, 353, 967, 981, 987, 993, 1000, 1004
- calculations, 797
 - properties, 1007
- special right-hand side, 555
- special Urysohn equations of first kind, method, 821
- special Urysohn equations of second kind, method, 822
- spectral radius, estimates, 649
- spectral radius of integral operator, 649
- spectral radius of kernel, 649
- spectrum of Fredholm integral equation, 760
- spectrum of operator, 1066
- spherical functions, Legendre of first kind, 299
- square integrable function, 501, 502
- square root, 9, 138, 222, 975
- stable solution, 623
- statement of Riemann problem, 718
- step-function, 1058
- integral, 1059
- Stieltjes integral, 1055, 1056
- basic definitions, 1055
 - existence theorems, 1056
 - properties, 1056
- Stieltjes integral sum, 1055
- Stieltjes transform, 221
- Stirling formula, 1013
- stochastic kernel, 654
- structure of solutions to linear integral equations, 502
- Struve function, 264, 299, 516, 518
- subspace, 1063
- orthogonal, 873
 - orthogonal, direct sum, 845, 863, 869
- successive approximation method, 566, 579, 632, 633, 811, 826, 876
- for ODEs, 876
 - general scheme, 566
 - resolvent, 566
- sufficient condition for compactness of integral operator, 842
- sum
- contain binomial coefficients, 920
 - contain integers, 920
 - finite, 919
 - finite functional, 922
 - finite numerical, 919
 - functional, finite, 922
 - integral, Stieltjes, 1055
 - involving hyperbolic functions, 922
 - involving trigonometric functions, 922
 - numerical, 921
 - numerical, finite, 919
 - of exponential functions, 564
 - of hyperbolic functions, 564
 - of orthogonal subspaces, direct, 845, 863, 869
 - of powers of natural numbers, 919, 920
 - of powers of natural numbers, alternating, 920
 - of trigonometric functions, 564
 - Stieltjes integral, 1055

- summable function, 1059
 integral, 1059
 superposition principle, linear, 502
 surface, equidistant, method, 891
 surface concentration
 equation, method of numerical integration, 891
 integral equations, 890
 surface reaction, 888
 symbol, Pochhammer, 1007
 symbols, 1007
 symmetric definite Fredholm kernel, 840
 symmetric equation, 639, 647
 Fredholm alternative, 643
 symmetric kernel, 573, 577, 625, 639, 645
 resolvent, 644
 symmetric positive definite Fredholm kernel, 866
 symmetric positive Fredholm kernel, 841
 system
 complete, 1065
 complete orthonormal, 855
 Fredholm integral equations of second kind, 701
 infinite of linear algebraic equations, 858, 861, 864, 868, 971
 infinite of linear algebraic equations with symmetric matrix, 850, 853
 normal of method of least squares, 695
 orthogonal, 1065
 orthonormal, 1065
 orthonormal, complete, 855
 Volterra integral equations, 549
 system of characteristic values, 640
 system of eigenfunctions, 640
 complete, 640
 incomplete, 640
 system of equations, 701
 reduction to single equation, 701
 system of Fredholm equations of second kind, 701
 system of functions
 complete orthonormal, 844
 orthonormal, complete, 844
 system of orthogonal polynomials, 795
- T**
- tables of definite integrals, 951
 tables of Fourier cosine transforms, 983
 tables of Fourier sine transforms, 989
 tables of indefinite integrals, 933
 tables of inverse Laplace transforms, 969
 tables of inverse Mellin transforms, 1001
 tables of Laplace transforms, 961
 tables of Mellin transforms, 997
 tangent, 60, 174, 251, 342
 hyperbolic, 36, 161, 241, 332
 tautochrone problem, 520
 terms of potentials, 892
 theorem
 analytic continuation, 595, 714
 Cauchy residue, 504
 theorem (*continued*)
 convolution, 507, 513
 existence, 875
 existence, for nonlinear equations, 830
 existence, for Stieltjes integral, 1056
 Fischer–Riesz, 1060
 Fredholm, 637, 702, 777
 Fubini, 1062
 generalized Jentzch, 648
 generalized Liouville, 595, 714
 Hilbert–Schmidt, 641, 1067
 Jentzch, generalized, 648
 Lebesgue on dominated convergence, 1060
 limit, 507
 residue, Cauchy, 504
 uniqueness, 875
 uniqueness, for nonlinear equations, 830
 theory
 Hilbert–Schmidt, 843
 Riesz–Schauder, 843
 theta functions, Jacobi, 110, 1042
 Tikhonov regularization method, 622, 829
 total variation of function, 1053
 trace method for approximation of characteristic values, 646
 trace of kernel, 646
 transform
 alternative Fourier, 512
 Boas, 250
 Bochner, 263, 518
 Buchholz, 274
 cosine, Fourier, Parseval’s relation, 514
 Crum, 268
 divisor, 269
 Feller, 226
 Fourier, 235, 511, 512, 518, 658
 Fourier, alternative, 512
 Fourier, asymmetric form, 512
 Fourier, definition, 512
 Fourier, inverse, 512
 Fourier, inversion formula, 512
 Fourier, properties, 513
 Fourier, rational, 685
 Fourier cosine, 514, 518
 Fourier cosine, asymmetric form, 514
 Fourier cosine, Parseval’s relation, 514
 Fourier cosine, tables, 983
 Fourier sine, 514, 518
 Fourier sine, asymmetric form, 515
 Fourier sine, Parseval’s relation, 515
 Fourier sine, tables, 989
 Gauss, 237
 generalized Mehler–Fock, 271
 Hankel, 261, 515, 518
 Hankel, Parseval’s relation, 515, 516
 Hardy, 264
 Hartley, 252, 518
 Hilbert, 228, 255, 518, 743
 Hilbert, on semiaxis, 229
 integral, 503, 515
 integral, kernel, 503

- transform (*continued*)
- integral, method, 586, 655, 809, 819
 - integral, table, 517
 - inverse, 503
 - inverse, representation as asymptotic expansions, 509
 - inverse, representation as convergent series, 509
 - inverse Fourier, 512
 - inverse Laplace, tables, 969
 - inverse Mellin, 510
 - inverse Mellin, tables, 1001
 - inverse of rational functions, 506
 - kernel, 503, 586, 655, 809, 819
 - Kontorovich–Lebedev, 267, 516, 518
 - Laplace, 235, 505, 511, 518, 524, 544, 658, 809
 - Laplace, definition, 505
 - Laplace, inverse, tables, 969
 - Laplace, inversion formula, 505
 - Laplace, properties, 507
 - Laplace, solution method, 524
 - Laplace, tables, 961
 - Laplace, two-side, 234, 518
 - Lebedev, 269
 - Mehler–Fock, 270, 518
 - Mehler–Fock, generalized, 271
 - Meijer, 516, 517
 - Mellin, 510, 511, 518, 587, 657, 658
 - Mellin, definition, 510
 - Mellin, inverse, 510
 - Mellin, inverse, tables, 1001
 - Mellin, inversion formula, 510
 - Mellin, properties, 511
 - Mellin, tables, 997
 - Olevskii, 276
 - Paley–Wiener, 260
 - rational Fourier, 685
 - Riesz, 226
 - sine, Fourier, Parseval’s relation, 515
 - Sonine, 114
 - Stieltjes, 221
 - table, 517
 - two-side Laplace, 234, 518
 - Weber, 265, 518
 - Weierstrass, 237, 518
- transformation, Kummer, 1025
- transformation of kernel, method, 532
- transposed characteristic equation, 758
- transposed characteristic operator, 758
- transposed equation, 573, 575, 625, 627, 637
- transposed equation of characteristic equation, 764
- transposed operator, 758
- transposed singular equation, 758
- trapezoidal rule, 534, 568
- triangle inequality, 501
- Tricomi confluent hypergeometric function, 273, 1024, 1025
- asymptotic expansions, 1024
 - integral representations, 1024
- Tricomi equation, 319, 769
- Tricomi–Gellerstedt equation, 320
- trigonometric functions, 46, 78, 84, 85, 166, 181, 186, 187, 246, 252, 256, 295, 335, 344, 349, 352, 353, 564, 907, 922, 944, 956, 966, 981, 986, 992, 999, 1003
- addition, 908
 - combinations, 176
 - inverse, 176, 344, 911, 948
 - inverse, addition, 912
 - inverse, relations, 912
 - inverse, subtraction, 912
 - of half argument, 909
 - of multiple arguments, 909
 - of single argument, relations, 908
 - powers, 908
 - products, 908
 - relationship, 916
 - subtraction, 908
 - sum, 564
- trigonometric nonlinearity, 420, 473
- trigonometric series
- in one variable, involving cosine, 928
 - in one variable, involving sine, 927
 - in two variables, 930
- trivial solution, 502
- two-dimensional equation of Abel type, 15
- two-dimensional integral equation, mixed with Schmidt kernel, 841
- two-dimensional singular equation, 231
- two-side Laplace transform, 234, 518
- type, convolution, 574, 606, 660, 669

U

- ultraspherical polynomials, 1050
- undetermined coefficients, 692
- uniqueness theorems, 875
- uniqueness theorems for nonlinear equations, 830
- unknown function of complicated argument, 227, 246, 254
- Urysohn equation, 806, 832
- first kind, 806, 829
 - second kind, 806
 - second kind with degenerate kernel, 818
 - special of first kind, method, 821
 - special of second kind, method, 822
- Urysohn form
- Volterra equation, 805, 811, 814, 816
 - Volterra equation, first kind, 805, 815
 - Volterra equation, second kind, 805

V

- value
- approximate of eigenvalues of Hilbert–Schmidt kernel, 845
 - Cauchy principal, 709
 - characteristic, 301, 625, 637, 639, 645, 697
 - characteristic, approximation, 646
 - characteristic, extremal properties, 644
 - characteristic, system, 640

- value (*continued*)
- in Banach space, continuous function of real argument, 840
 - in Hilbert space, continuous function of real argument, 840
 - in space of functions square integrable over closed bounded set, continuous function of real argument, 842
 - in space of functions square integrable over ring-shaped domain, continuous function of real argument, 841
 - in space of square integrable functions, continuous function of real argument, 840
 - regular, 301, 625, 637
- variable integration limit, 3, 805, 809, 811
- variable limit of integration, 3, 805, 809, 811
- variable lower integration limit, 537, 570
- variable lower limit of integration, 537, 570
- variables, several, function, 839
- variation, total, of function, 1053
- variation function, bounded, 1056
- vector, 1063
- axioms for addition, 1063
 - axioms relating addition of vectors with their multiplication by scalars, 1063
 - orthogonal, 1065
- vector space, 1063
- Volterra equation, 549, 805, 877
- first kind, 519, 524, 565
 - first kind, connection with Volterra equations of second kind, 524
 - first kind, existence of solution, 519
 - first kind, in Hammerstein form, 806
 - first kind, in Urysohn form, 805, 815
 - first kind, problems, 520
 - first kind, uniqueness of solution, 519
 - Hammerstein form, 806
 - nonlinear, 805
 - quadratic nonlinearity, 809
 - reduction to Wiener–Hopf equation, 528
 - second kind, 524, 539, 565
 - second kind, connection with Volterra equations of first kind, 524
 - second kind, in Urysohn form, 805
 - second kind, of Hammerstein form, 816
 - second kind, reduction to Volterra equations of first kind, 565
 - second kind, sequence, 855
 - second kind, sequence of independent, 853, 865, 872
 - sequence, 844, 850, 862
 - sequence of independent, 847, 858
- Volterra equation (*continued*)
- systems, 549
 - Urysohn form, 805, 811, 814, 816
- Volterra integral operator, 842
- Volterra kernel, 839
- Volterra operator, 873
- volume potential, 893
- Gauss formula, 894
- ## W
- weak singularity, 574, 588, 625
- kernel, 519, 532, 574, 588, 625
- weakly singular kernel, 532
- Weber function, 88
- Weber parabolic cylinder function, 1034
- Weber transform, 265, 518
- Weierstrass elliptic function, 1041
- Weierstrass \wp -function, 1041
- Weierstrass transform, 237, 518
- weight function, Jacobi, 793
- well-posed problem, 623
- general notions, 623
- Whittaker confluent hypergeometric function, 274, 1027
- Whittaker equation, 1027
- Wiener–Hopf equation, 574, 626, 679
- first kind, 285, 538, 574, 606
 - Krein’s method, 679
 - second kind, 373, 547, 571, 626, 660, 679
 - second kind, exceptional case, 678
 - second kind, homogeneous, 672
 - second kind, index, 661
 - second kind, nonhomogeneous, 677
 - second kind, solution, 681
 - Volterra equation, 528
- Wiener–Hopf method, 671
- scheme, 676
- Wronskian, confluent hypergeometric function, 1026
- Wronskian, Legendre function, 1034

Y

Y-transform, 516, 518

Y_ν -transform, 264

Z

Zakharov–Shabat method, 898

zero measure, set, 1058

zeros of Bessel functions, 1019