



<http://www.biodiversitylibrary.org>

**Botanical Museum leaflets, Harvard University.**

Cambridge, Mass. :Botanical Museum, Harvard University, 1932-  
<http://www.biodiversitylibrary.org/bibliography/7542>


**v.24 (1974-1976):** <http://www.biodiversitylibrary.org/item/31873>

Page(s): Page 113, Page 114, Page 115, Page 116, Page 117, Page 118, Page 119

Contributed by: Missouri Botanical Garden  
Sponsored by: Missouri Botanical Garden

Generated 2 February 2011 1:02 PM  
<http://www.biodiversitylibrary.org/pdf3/005713600031873>

This page intentionally left blank.



*Orupl.*

# BOTANICAL MUSEUM LEAFLETS

## HARVARD UNIVERSITY

CAMBRIDGE, MASSACHUSETTS, OCTOBER 31, 1975

VOL. 24, No. 6

### NUTRITIONAL VALUE OF COCA

BY

JAMES A. DUKE<sup>1</sup>, DAVID AULIK<sup>2</sup> AND  
TIMOTHY PLOWMAN<sup>3</sup>

Leaves of wond'rous nourishment  
Whose Juice Succ'd in, and to the Stomach tak'n  
Long Hunger and long Labour can sustain;  
From which our faint and weary Bodies find  
More Succor, more they cheer the drooping Mind,  
Than can your Bacchus and your Ceres join'd.

—ABRAHAM COWLEY in Mortimer's  
*History of Coca*

*Abstract.* Coca leaves (*Erythroxylum Coca* Lam.) from Chapare, Bolivia, compared to an average of 50 other Latin American vegetable products, are higher in calories (305 per 100 g compared to 279), protein (18.9 g: 11.4 g), carbohydrate (46.2 g: 37.1 g), fiber (14.4 g: 3.2 g), ash (9.0 g: 2.0 g), calcium (1540 mg: 99 mg), phosphorus (911 mg: 270 mg), iron (45.8 mg: 3.6 mg), vitamin A (11,000 IU: 135 IU), and riboflavin (1.91 mg: 0.18 mg). Coca was lower than the average for the 50 plant foods in oil content (5.0 g per 100 g compared to 9.9 g), moisture (6.5 g: 40.0 g), thiamin (0.35 mg: 0.38 mg), niacin (1.3 mg: 2.2 mg), and ascorbic acid

<sup>1</sup>Chief, Plant Taxonomy Laboratory, Plant Genetics and Germplasm, Institute, Agricultural Research Service, Beltsville, Maryland.

<sup>2</sup>WARF Institute, Inc., Box 2599, Madison, Wisconsin.

<sup>3</sup>Research Associate in Economic Botany, Botanical Museum, Harvard University, Cambridge, Massachusetts.

(1.4 mg: 13.0 mg). Ingestion of 100 g of the Bolivian coca leaves tested would more than satisfy the Recommended Dietary Allowance for reference man and woman of calcium, iron, phosphorus, vitamin A, vitamin B<sub>2</sub> and vitamin E. However, the leaves also contain alkaloids and may harbor pesticide residues.

Unable to establish the nutritional value of coca leaves (*Erythroxylum Coca* Lam.) after consulting many sources, we obtained a one kilogram sun-dried sample from San Francisco, Province of Chapare, Bolivia, in June, 1974. Using methods listed in References and Notes, we obtained the following nutritional analysis (1): calories, 305 per 100 g; moisture, 6.5 g; protein, 18.9 g; carbohydrate, 46.2 g; fat, 5.0 g; vitamin A, 11,000 IU (as beta-carotene); vitamin C, 1.4 mg; vitamin B<sub>1</sub> (thiamine), 0.35 mg; vitamin B<sub>2</sub> (riboflavin), 1.9 mg; niacin, 1.29 mg; calcium, 1,540 mg; iron, 45.8 mg; vitamin E, 43.5 IU (as d-alpha tocopherol); vitamin B<sub>6</sub>, 0.508 mg; folic acid, 0.130 mg; vitamin B<sub>12</sub>, 1.05 mcg; iodine, 5.0 mcg; phosphorus, 911 mg; magnesium, 213 mg; zinc, 2.70 mg; copper, 1.21 mg; biotin, 0.0863 mg; pantothenic acid, 0.684 mg; and sodium, 40.6 mg. The analysis of other elements by emission spectroscopy yielded the following amounts: potassium, 2.02 g per 100 g; aluminum, 39.5 mg; barium, 4.67 mg; strontium, 9.71 mg; boron, 5.35 mg; zinc, 2.70 mg; manganese, 6.65 mg; and chromium, 0.359 mg.

Surprised by the high values, especially in calcium and iron, we tabulated nutritional averages for other plant products ingested by Latin Americans (Table 1). Compared with an average from ten nuts and oilseeds (2) (*Sesamum indicum*, *Terminalia Catappa*, *Prunus Amygdalus*, *Corylus* spp., *Arachis hypogaea*, *Castanea* spp., *Bertholletia excelsa*, *Helianthus annuus*, *Anacardium occidentale* and *Inga* spp.), the San Francisco coca leaves

were higher in protein, carbohydrate, ash, calcium, phosphorus, iron, vitamin A and riboflavin. Coca was lower in calories, moisture, fat, thiamin, niacin, and vitamin C. Compared with an average of ten pulses (*Vigna unguiculata*, *Cicer arietinum*, *Cajanus Cajan*, *Pisum sativum*, *Vicia Faba*, *Phaseolus vulgaris*, *Dolichos Lablab*, *Lens* spp., *Glycine Max* and *Lupinus mutabilis*), coca was equal in fat; higher in fiber, ash, calcium, phosphorus, iron, vitamin A, and riboflavin; and lower in calories, moisture, protein, carbohydrate, thiamin, niacin, and vitamin C. Compared with an average of ten cereals (*Amaranthus caudatus*, *Oryza sativa*, *Avena sativa*, *Chenopodium pallidicaule*, *Chenopodium Quinoa*, *Hordeum vulgare*, *Secale cereale*, *Coix Lachryma-jobi*, *Zea Mays* and *Triticum aestivum*), coca was higher in protein, fat, fiber, ash, calcium, phosphorus, iron, vitamin A, riboflavin, and vitamin C; lower in calories, moisture, carbohydrate, thiamin, and niacin. Compared with an average of ten vegetables (*Canna edulis*, *Capsicum* spp., *Allium sativum*, *Arracacha xanthorrhiza*, *Ipomoea Batatas*, *Cyclanthera pedata*, *Cucurbita maxima*, *Allium Cepa*, *Brassica oleracea*, and *Tropaeolum tuberosum*) and an average of ten fruits (*Persea americana*, *Ananas comosus*, *Musa sapientum*, *Cocos nucifera*, *Passiflora mollissima*, *Annona Cherimolia*, *Prunus persica*, *Fragaria* spp., *Annona muricata*, and *Ficus Carica*), coca was high on all counts except moisture and vitamin C.

The present coca analyses are comparable to an average of three earlier coca analyses from Bolivia (3) and three recently reported from Peru (4). Frequent reports that coca has no nutritional value should be re-evaluated in view of these findings. The comparatively high nutritional values for coca are due partly to the fact that the leaves are dry (less than 10% moisture) when purchased, whereas most other foods are higher in moisture.

In most areas where it is used, coca should be considered a masticatory since it is not wholly consumed by the chewer. Typically, the leaves are first moistened in the mouth with saliva, then formed into a quid with the tongue and pushed into the upper cheek cavity. They are then sucked to extract the rich, green juice which is subsequently swallowed. Usually some form of alkali is added to facilitate this extraction. When the chew is exhausted, it is usually spat out. Thus, the full complement of nutrients present in the coca leaf is not consumed entirely, and the nutritional amounts reported here may be somewhat higher than the amounts actually ingested by the coca chewer. To our knowledge, no studies have been made on the nutritional value of the swallowed extract.

In the Colombian Amazon, a variation of coca use is practiced by several tribes. Coca leaves are pulverized to a fine powder along with the ashes of *Pourouma* or *Cecropia* leaves. The mixture is placed in the mouth on the gums and inner cheeks and is eventually swallowed (8). In this case, and in instances where a coca chewer swallows his quid, the full complement of the leaf nutrients would be ingested. There is essentially no difference here between the use of coca and the direct consumption of food, in terms of nutrition.

The amounts of coca consumed may contribute significantly to the diets of Andean coca chewers (5). If the average chewer ingests 60 g of Peruvian coca per day (5,6), he more than satisfies his requirements for calcium, even without the supplemental ash or lime usually added to the coca quid (6). No other food in the INCAP Food Composition Tables (2) approaches coca for calcium content (1,789 mg). Other food items are high in calcium: sesame seeds, 1212 mg per 100 g; spinach flour, 488 mg; leaves of *Laurus nobilis*, 803 mg; leaves of *Justicia pecto-*

*ralis*, 663 mg; West Indian Almond (*Terminalia Catappa*), 497 mg; powdered skim milk, 1,301 mg; whole milk, 921 mg; and alligator meat, 1,231 mg; but none equals coca. Few food plants can satisfy the calcium and iron in the Recommended Dietary Allowance (RDA) of reference man ingesting 100 g. The Bolivian coca leaves reported here do satisfy the RDA.

Coca leaves may, however, contain 0.25 to 2.25% toxic alkaloids, including benzoylecgonine, benzoyltropine, cinnamylcocaine, cocaine, cuscohygrine, dihydroxytropine, hygrine, hygroline, methylcocaine, methylecgonidine, nicotine, tropacocaine, and  $\alpha$ - and  $\beta$ -truxilline (4,7). These alone could make the nutritious coca leaf undesirable as a source of nutrients. The average coca chewer could also ingest 442 mg of copper in a year if the San Francisco leaves are typical; but this amount is not excessive.

Many coca growers in both high (Chapare, Bolivia) and low (Yungas, Bolivia) rainfall areas may use insecticides. The leaves are not intentionally washed, and harvest and curing are timed to avoid rainfall. In considering coca for human consumption, the leaves should be analyzed for insecticide residues.

Although coca leaves contain relatively high levels of certain nutrients, the presence of alkaloids and the possible presence of insecticide residues suggest caution in coca chewing.

## REFERENCES AND NOTES

1. Methods: Protein ( $\% \text{ N} \times 6.25$ ), Association of Official Analytical Chemists (A.O.A.C.), 11th Ed., 16. 1970; moisture, vacuum oven, A.O.A.C., 122. 1970; ash, A.O.A.C., 123. 1970; fat, A.O.A.C., 129. 1970; carbohydrates, by difference; calories, by calculation; vitamin A, Moore & Ely, Ind. Eng. Chem. Anal. Ed., 13: 600. 1941; vitamin C, J. Biol. Chem., 147: 399. 1943; vitamin B<sub>1</sub>, A.O.A.C., 771. 1970; vitamin B<sub>2</sub>, A.O.A.C., 789. 1970; niacin, A.O.A.C., 787. 1970; elemental, J.A.O.A.C., 51: 1003. 1968; vitamin E, Acta Chemica Scandinavica 11: 34-43. 1957; vitamin B<sub>6</sub> (*Streptococcus carlsbergensis*), Atkins, Schultz, Williams & Frey, Ind. & Eng. Chem., Anal. Ed., 15: 141. 1943; folic acid, A.O.A.C., 786. 1970; vitamin B<sub>12</sub>, U.S.P. 17: 864. 1965; iodine, ashing, A.O.A.C., 674. 1970; colorimetry, W.T. Binnerts, Anal. Chemica Acta 10: 78. 1954; biotin (*Lactobacillus arabinosus*), Wright & Skeggs, Proc. Soc. Exp. Biol. & Med., 56: 95. 1944; pantothenic acid, Nielands & Strong, Arch. Biol., 19: 2. 1948.
2. Wu Leung, W. & M. Flores. 1961. Tabla de Composición de Alimentos para Uso en América Latina. Instituto de Nutrición de Centro América y Panamá (INCAP) and Interdepartmental Committee on Nutrition for National Defense (ICNND). U.S. Government Printing Office. Washington, D.C.
3. del Granado, J.T. 1931. Plantas Bolivianas. Arno Hermanos. La Paz.
4. Machado, E. 1972. El género *Erythroxylon* en el Perú. Raymondiana 5: 5-101.
5. Hanna, J.M. 1974. Coca Leaf Use in Southern Peru: Some Bio-social Aspects. Am. Anthropologist 76 (2): 281-296.
6. Baker, P.T. & R.B. Mazess. 1963. Calcium: Unusual Sources in the Highland Peruvian Diet. Science 142: 1466-7.
7. Willaman, J.J. & B.G. Schubert. 1961. Alkaloid-bearing plants and their Contained Alkaloids. U.S. Department of Agriculture. Washington, D.C.
8. Schultes, R.E. 1957. A new method of coca preparation in the Colombian Amazon. Bot. Mus. Leaflet Harvard Univ. 17 (9): 241-246.



TABLE 1

Nutritional Comparison per 100 g of Coca Leaves with other Latin American Plant Foods.

FOOD ITEM	# in sample	Cal	H <sub>2</sub> O g	Prot. g	Fat g	Carb. g	Fiber g	Ash g	Ca mg	P mg	Fe mg	Vit A <sup>1</sup> IU	Thia mg	Rib mg	Nia mg	Vit C mg
San Francisco coca	(1)	305	6.5	18.9	5.0	46.2	14.4	9.0	1540	911	45.8	11,000	0.35	1.91	1.29	1.4
Bolivia coca	(3)	— <sup>2</sup>	8.8	—	1.6	42.4	8.0	5.3	—	—	—	—	—	—	—	—
Peru coca	(3)	—	10.3	18.7	—	—	17.5	4.6	2038	363	7.9	9,000	0.81	1.55	6.17	—
COCA AVERAGE	(7)	—	8.5	18.8	3.3	44.3	13.3	6.3	1789	637	26.8	10,000	0.58	1.73	3.7	1.4
PLANT FOOD AVERAGE	(50)	279	40.0	11.4	9.9	37.1	3.2	2.0	99	270	3.6	135	0.38	0.18	2.2	13.0
Nuts & Seeds	(10)	521	9.9	16.8	36.0	28.2	3.6	3.1	273	522	4.3	17	0.78	0.28	5.2	2.1
Pulses	(10)	354	11.3	25.4	5.0	55.1	5.5	3.3	102	398	7.1	20	0.58	0.24	2.25	1.9
Cereals	(10)	352	11.5	11.7	3.7	71.0	4.0	2.1	74	346	4.8	13	0.41	0.25	2.7	0.8
Vegetables	(10)	74	87.3	1.8	0.4	16.9	1.5	0.9	26	52	1.2	595	0.09	0.05	1.0	31.0
Fruits	(10)	93	79.6	1.2	4.5	14.1	1.4	0.7	20	33	0.8	35	0.05	0.06	0.08	29.0

<sup>1</sup> as beta-carotene<sup>2</sup> no data

The following text is generated from uncorrected OCR.

[Begin Page: Page 113]

- ANICAL MUSEUM LEAFLETS

HARVARD UNIVERSITY

CAM IHDIE, MASSA (IISFTTS, )(:IO i .I 31, 1975 Vol. 24, No. (

N TITUTIONAL VALUI OF COCA  
BY

.liAM:ES A. 1)UKi-•, DAVID AUIl'K" ANI  
TIM(OTHY PI)IW)MAN;

Leaves of wond'rous nourishment  
Whose Juice Succ'd in, and to the Stomach tak'n  
Long Hunger and long Labour can sustain:  
From which our faint and weary Bodies find  
More Succor, more they cheer the drooping Mind,  
Than can your Bacchus and your Ceres join'd.

-AHAIllAM (COWI.IY in Mortimer's  
lis.tory/ ql' Cora

,blstract. Coca leaves (*Erythroxi.v/lui* (Cc){• ILamn.)  
from Chapare. Bolivia, compared to an average of 50  
other Latin American vegetable products, are higher in  
calories (305 per 100 g compared to 279), protein (18.9  
g: 11.4 g). carbohydrate (46.2 g: 37.1 g), fiber (14.4 g:  
3.2 g). ash (9.0 g: 2.0 g), calcium (1540 mg: 99 mg).  
phosphorus (911 nng: 270 mg), iron (45.8 mg: 3.6 mg).  
vitamin A (11.000 II': 135 I1'), and riboflavin (1.91  
img: 0.18 mg). Coca was lower than the average for the  
50 plant foods in oil content (5.0 g per 100 g comlpared  
to 9.9 g), moisture (6.5 g: 40.0 g), thianiin (0.35 mrg:  
0.38 rmg), niacin (1.3 mug : 2. 2 mg), and ascorbic acid  
Chief, Plant Taxonony Laboratory, Plant Genetics and Germ-  
plasm, Institute, Agricultural Research Service, Beltsville, Maryland.  
'WA RF Institute, Inc., Box ,2599, Madison, Wisconsin.  
" Research Associate in Economic Botany, Botanical Museum,  
llarvard University, Cambridge, Massachusetts.

[ 113 ]

[Begin Page: Page 114]

(1. - mg : 133.0 rug). Ingestion of 100 g of the Bolivian coca leaves tested would more than satisfy the recommended Allowance for reference man and women of calcium, iron, phosphorus, vitamin B<sub>1</sub>, vitamin B<sub>2</sub>, vitamin B<sub>6</sub>, vitamin B<sub>12</sub> and folic acid. However, the leaves also contain alkaloids and may harbor pesticide residues.

In order to establish the nutritional value of coca leaves (from the Yungas of the Cochabamba Province, Bolivia, in 1974), we obtained a one kilogram sun-dried sample from San Francisco, Province of Tarija, Bolivia, in 1974. The analytical methods listed in References and Notes, we obtained the following nutritional analysis (1): calories, 305 per 100 g; moisture, 11.5 g; protein, 18.1 g; carbohydrate, 42.2 g; fat, 3.0 g; vitamin A, 11,000 IU (as beta-carotene); vitamin C, 1.4 mg; vitamin B<sub>1</sub>, 0.3 mg; vitamin B<sub>2</sub>, 0.3 mg; vitamin B<sub>6</sub>, 1.9 mg; niacin, 1.29 mg; calcium, 134 mg; iron, 14.3 mg; vitamin K, 13.3 IU (as d-alpha-tocopherol); vitamin E, 0.308 mg; folic acid, 0.13 mg; vitamin B<sub>12</sub>, 1.0 mcg; iodine, 5.0 mcg; phosphorus, 111 mg; magnesium, 213 mg; zinc, 0.70 mg; cobalt, 1.21 mcg; biotin, 0.08 mcg; pantoic acid, 0.81 mg; and sodium, 10.1 mg. The analysis of other elements by emission spectroscopy yielded the following amounts: potassium, 0.02 g per 100 g; aluminum, 39.5 mg; barium, 4.7 mg; strontium, 0.71 mg; boron, 5.35 mg; zinc, 2.70 mg; manganese, 0.39 mg.

Surprised by the high values, especially in calcium and iron, we tabulated nutritional averages for other plant products ingested by Latin Americans (Table 1). Compared with an average from ten nuts and oilseeds (2) (Saxena and Pandey, 1974), the coca leaves (1) are rich in calcium, iron, phosphorus, and potassium. The coca leaves are also rich in vitamins B<sub>1</sub>, B<sub>2</sub>, B<sub>6</sub>, B<sub>12</sub>, and folic acid. The coca leaves are also rich in potassium, calcium, iron, phosphorus, and magnesium. The coca leaves are also rich in vitamins B<sub>1</sub>, B<sub>2</sub>, B<sub>6</sub>, B<sub>12</sub>, and folic acid. The coca leaves are also rich in potassium, calcium, iron, phosphorus, and magnesium.

were higher in protein, carbohydrate, ash, calcium, phosphorus, iron, vitamin A and riboflavin. Coca was lower in calories, moisture, fat, thiamin, niacin, and vitamin C. Compared with an average of ten pulses (Guinea, Chickpea, Bean, Lentil, Pea, Broad Bean, Soybean, Mung Bean, Pigeon Pea, and Lupin), coca was equal in fat; higher in fiber, ash, calcium, phosphorus, iron, vitamin A, and riboflavin; and lower in calories, moisture, protein, carbohydrate, thiamin, niacin, and vitamin C. Compared with an average of ten cereals (Wheat, Rice, Corn, Sorghum, Millet, Barley, Oats, Rye, Buckwheat, and Quinoa), coca was higher in protein, fat, fiber, ash, calcium, phosphorus, iron, vitamin A, riboflavin, and vitamin C; lower in calories, moisture, carbohydrate, thiamin, and niacin. Compared with an average of ten vegetables (Cauliflower, Broccoli, Spinach, Carrot, Potato, Onion, Garlic, Tomato, Eggplant, and Zucchini), coca was higher in protein, fat, fiber, ash, calcium, phosphorus, iron, vitamin A, riboflavin, and vitamin C; lower in calories, moisture, carbohydrate, thiamin, and niacin. Compared with an average of ten fruits (Apple, Banana, Orange, Grape, Peach, Pear, Plum, Raisin, Strawberry, and Watermelon), coca was high on all counts except moisture and vitamin C.

The present coca analyses are comparable to an average of three earlier coca analyses from Bolivia (3) and three recently reported from Peru (4). Frequent reports that coca has no nutritional value should be re-evaluated in view of these findings. The comparatively high nutritional values for coca are due partly to the fact that the leaves are dry (less than 10% moisture) when purchased, whereas most other foods are higher in moisture.

[ 11.

[Begin Page: Page 116]

In most areas where it is used, coca should be considered a masticatory since it is not wholly consumed by the chewer. Typically, the leaves are first moistened in the mouth with saliva, then formed into a quid with the tongue and pushed into the upper cheek cavity. They are then sucked to extract the rich, green juice which is subsequently swallowed. Usually some form of alkali is added to facilitate this extraction. When the chew is exhausted, it is usually spat out. Thus, the full complement of nutrients present in the coca leaf is not consumed entirely, and the nutritional amounts reported here may be somewhat higher than the amounts actually ingested by the coca chewer. To our knowledge, no studies have been made (on the nutritional value of the swallowed extract).

In the Colombian Amazon, a variation of coca use is practiced by several tribes. Coca leaves are pulverized to a fine powder along with the ashes of *Ipourouina* or *Crocipia* leaves. The mixture is placed in the mouth on the gums and inner cheeks and is eventually swallowed (8). In this case, and in instances where a coca chewer swallows his quid, the full complement of the leaf nutrients would be ingested. There is essentially no difference here between the use of coca and the direct consumption of food, in terms of nutrition.

The amounts of coca consumed may contribute significantly to the diets of Andean coca chewers (5). If the average chewer ingests 10 g of Peruvian coca per day (5,6), he not only satisfies his requirements for calcium, even without the supplementary calcium usually added to the coca quid (1). No other food in the I NCAP Food Composition Tables (1) approaches coca for calcium content (1.78 mg). Other food items are high in calcium: sesame seeds, 1212 mg per 100 g; spinach flour, 488 mg; leaves (of *Laurus nobilis*, 80; fig : leaves of *Juiceira* / *co-*

[ 1 ; 11

[Begin Page: Page 117]

ralis, (1.3 mg; West Indian Almond (1 Ter/Inalia Catap-

pa), 497 mg powdered skim milk, 1,301 mg; whole milk, 921 mg; and alligator meat, 1,231 mg: but none equals coca. Few food plants can satisfy the calcium and iron in the Recommended Dietary Allowance (RDA) of reference man ingesting 100 g. The Bolivian coca leaves reported here do satisfy the RDA.

Coca leaves may, however, contain 0.25 to 2.25% toxic alkaloids, including benzoylecgonine, benzoylecgonine, cinnamylecgonine, cocaine, cuscohygrine, dihydroxytropine, hygrine, hygroline, methylcocaine, n-methylecgonidine, nicotine, tropacocaine, and  $\alpha$ - and  $\beta$ -truxilline (4,7). These alone could make the nutritious coca leaf undesirable as a source of nutrients. The average coca chewer could also ingest 442 mg of copper in a year if the San Francisco leaves are typical: but this amount is not excessive.

Many coca growers in both high (Chlapare, Bolivia) and low (Yungas, Bolivia) rainfall areas may use insecticides. The leaves are not intentionally washed, and harvest and curing are timed to avoid rainfall. In considering coca for human consumption, the leaves should be analyzed for insecticide residues.

Although coca leaves contain relatively high levels of certain nutrients, the presence of alkaloids and the possible presence of insecticide residues suggest caution in coca chewing.

[117

[Begin Page: Page 118]

#### INTERNATIONAL NOTES

1. Methods: Protein (N x 6.25), Association of Official Analytical Chemists (A.O.A.C.), 11th Ed., 116. 1970: moisture, vacuum oven, A.O.A.C., 122. 1970; ash, A.O.A.C., 123. 1970: fat, A.O.A.C., 129. 1970: carbohydrates, by difference: calories, by calculation: vitamin A, Moore & Ely, Ind, Eng. Chem. Anal. ed., 1: 6(1900). 191- : vitamin C, J. Biol. Chem., 11-47 : 899. 191- : vitamin B<sub>1</sub>, A.O.A.C., 771. 1970: vitamin I, A.O.A.C., 5,9. 1970: niacin, A.O.A.C., 7S7. 1970: elemental, A.O.A.C., 1 : 100. 1961S: vitamin 1:., Acta Chemica Scandinavica 11 : ; - I:

19.K : vitamin 13, ( Streplacoccus. cnr/lshrgei.vi s), Atkins, Schultz, Williams & Frey, Intl. & Irng. Chern., Anal. l.d., 1 : 41. 1918: folic acid, A.O.A.C , 7S . 1970: vitarmin B , ".S. l. 17: S 1.. 19ti,: iodine, ashing, A.O.A.C., i 174. 1970: colorimetry, W .T. Binnerts, Anal. C(hemica Acta 10: 7 S. 1.)! : biotin ( l.actofbah illus arlbhi, o.isus), \rigit & Skeggs. l'roc. Soc. 1xp. Biol. & Bled., (ti: 9.3. 191-1: pantothenic acid, Nielsands & Strong, Arch. Biol., 19: 2. 19 8.

,. Wu leung, Wl. & M. Flores. 1961. 'abla de (omposicion de Alimentos para l so en America .Latina. Instituto de Nutriciin de Centro America vy Panan;i (IN('AP) and Interdepartmental Committee on Nutrition for National l)efense (('N N l). .S. ( iovernmient Printing (t)ice. \ashington, l).C.

8. del (iranado, J.T'. 19381. Plantas Holivianas. Arno llerrnanos. La Paz.

l. Machado, E. 1972. ,l genero l)'rythroxylyon en el i'eri. Raymondiana 5 : ,- 101.

. lanna, .J.M. 197 l.. Coca Leaf l se in Southern Peru : Some Bio-social Aspects. Am. Anthropologist 7; ('): ' 1- 9 l- ti.

ti. Baker, P.T. & lt. B. Maezss. 19ti;. Calcium : Inusual Sources in the hhighland Peruvian l)iet. Science 1 4 : 14titi-7

t. Willaman, J.,1. & B.G( . Schubert. l 9i 1. Alkaloid-bearing plants and their Contained Alkaloids. .S. l)epartment of Agriculture. Washington, D).C.

S. Scihultes, R. 1. 19.7. A new nrethod of coca preparation in the Colombian Amazon. Bot. Mus. Leaf, l larvard lUniv. 17 (91 : 21-'2 hl.

t ri c-

0 0  
:2 9 '9

-N -4 O 0

CM ifl 0. 0-< cx  
N 0

co tf iN LO 'sC  
\*n N 0 0  
0066

C"MCN CM o  
oC o 0

t-- LO4 1

r- 0 M  
N

1& m i N c  
G -! r1 -! 6  
UNM f r~ t

IMc

--r w N

-t~ 10 N. CC C7



0 CC

l 1 t-

d:c M m

a o

tl n

-r

1 -  
t C.

0  
oa

LC  
0-

cc 00 cc N M  
n rn to M

Nt-

r--4 CM N.i U

N - -o c -

vU ttj , j r\_  
N M l'  
N\L) N . r--

t6 6 J - -  
tri-- Nco  
cl

h cn

N )O  
nC 1")

0 0 0 0 0

[ 119 ]

rU

0  
CT  
0

ro  
J  
Sc;

I?  
^)~

C

cr

C  
. r