

K6+8.574

BIBLIOTECA UCM



5302357871

564.192

TRE

TREATISE ON INVERTEBRATE PALEONTOLOGY

Prepared under Sponsorship of

The Geological Society of America, Inc.


The Paleontological Society The Society of Economic Paleontologists and Mineralogists

The Palaeontographical Society The Palaeontological Association

Directed and Edited by
RAYMOND C. MOORE

Revisions and Supplements Directed and Edited by
CURT TEICHERT

Editorial Assistants, LAVON McCORMICK, ROGER B. WILLIAMS



BIBLIOTECA DE PALEONTOLOGIA		
Armario	Estante	Número
U	3	15

Part N
VOLUME 2 (of 3)
MOLLUSCA 6
BIVALVIA

UNIVERSIDAD DE MADRID
FACULTAD DE CIENCIAS
Sección de GEOLOGICAS
Registro de Libros
Bibliotecas

N.º Registro 2371 vol 2

By †L. R. COX, N. D. NEWELL, D. W. BOYD, C. C. BRANSON, RAYMOND CASEY,
ANDRÉ CHAVAN, A. H. COOGAN, COLETTE DECHASEAUX, C. A. FLEMING, FRITZ
HAAS, L. G. HERTLEIN, E. G. KAUFFMAN, A. MYRA KEEN, AURÈLE LAROCQUE,
A. L. McALESTER, R. C. MOORE, C. P. NUTTALL, B. F. PERKINS, H. S. PURI, L. A.
SMITH, T. SOOT-RYEN, H. B. STENZEL, E. R. TRUEMAN, RUTH D. TURNER, and
JOHN WEIR

THE GEOLOGICAL SOCIETY OF AMERICA, INC.
and
THE UNIVERSITY OF KANSAS

1969





PART N

BIVALVIA

By †L. R. COX, N. D. NEWELL, D. W. BOYD, C. C. BRANSON, RAYMOND CASEY, ANDRÉ CHAVAN, A. H. COOGAN, COLETTE DECHASEAUX, C. A. FLEMING, FRITZ HAAS, L. G. HERTLEIN, E. G. KAUFFMAN, A. MYRA KEEN, AURÈLE LAROCQUE, A. L. McALESTER, R. C. MOORE, C. P. NUTTALL, B. F. PERKINS, H. S. PURI, L. A. SMITH, T. SOOT-RYEN, H. B. STENZEL, E. R. TRUEMAN, RUTH D. TURNER, and JOHN WEIR

VOLUME 2

Subclass HETERODONTA Neumayr, 1884

[*nom. transl.* NEWELL, 1965 (ex Heterodonta NEUMAYR, 1884; unspecified higher category)] [Diagnosis by N. D. NEWELL]

Mantle lobes more or less joined and usually possessing developed siphons; cardinal area commonly present, prosodetic or divided into lunule and escutcheon; ligament opisthodetic, with or without separate resilium; without lithodesma and rarely with external accessory shelly pieces; nepionic stage usually without taxodont provinculum or adult byssus; generally possessing hinge plate and teeth differentiated into distinct cardinals and laterals; shell structure crossed lamellar, complex, or prismatic, never nacreous; gills eulamellibranchiate. *M.Ord.-Rec.*

Order VENEROIDA

H. Adams & A. Adams, 1856

[*nom. correct.* NEWELL, 1965 (pro order Veneracea H. & A. ADAMS, 1856)] [= [Heterodonta AUCT. (*partim*)] [Diagnosis by N. D. NEWELL]]

Commonly equivalve, isomyarian; dentition differentiated into standardized cardinals and laterals, or few cardinals only; posterior laterals, where present, lie entirely behind ligament; anterior laterals, if present, formed early in ontogeny by separation from cardinals of anterior part of hinge lamellae; habit active or nestling, rarely sedentary burrowers, or sessile. *M.Ord.-Rec.*

Superfamily BABINKACEA Horný, 1960

[*nom. transl.* McALESTER, 1965, ex Babinkidae HORNÝ, 1960]
Characters of family. *M.Ord.*

Family BABINKIDAE Horný, 1960

[Materials for this family prepared by A. L. McALESTER, Yale University]

Shell anteriorly expanded, compressed; two small cardinal teeth in LV, one large cardinal tooth in RV; elongate, subequal adductor muscle scars, eight rounded pedal muscle scars in line between top of adductors; pallial line nonsinuate. *M.Ord.*

Babinka BARRANDE, 1881 (p. 31) [**B. prima*; OD]. Characters of family. *M.Ord.*, Eu.(Czech.).— FIG. E1.1. **B. prima*; Czech.(Osek); 1a, LV int. mold, $\times 2.7$ (Newell, n); 1b,c, LV and RV hinges, latex casts, $\times 2.7$ (McAlester, n).

Superfamily LUCINACEA Fleming, 1828

[*nom. transl.* ANTON, 1839 (ex Lucinidae FLEMING, 1828)] [Materials for this superfamily prepared by ANDRÉ CHAVAN, Seyssel (Ain), France]

Shell generally equivalve, subcircular to oval or subtrigonal; commonly smooth externally or concentrically sculptured on internal layer of straight radial ribs, many genera with two dorsal folds which delimit anterior and posterior areas; beaks normally small, prosogyrous to orthogyrous; lunule generally more extended on RV; escutcheon ill-defined. Hinge with typical tooth formula 3a, 3b/2, 4b, but 3a may be obliterated, lateral laminae (if present) duplicate in LV only; integripalliate. Animal with more or less elongate to vermiform foot, rudimentary labial palps, and one or two siphonal apertures. [Marine or estuarine.] *Sil.-Rec.*

The alphabetically arranged generic descriptions in each family-group division of the Lucinacea are




FIG. E1. Babinkidae (p. N491).

accompanied by numbers inclosed by square brackets. Such numbers indicate position in the sequence of generic taxa given with the respective families or subfamilies for the purpose of recording CHAVAN's arrangement, designed to reflect "natural relationships" of these taxa as inferred by him.

Family LUCINIDAE Fleming, 1828

[*nom. correct.* D'ORBIGNY, 1837 (*pro* Lucinidae FLEMING, 1828)]

Subcircular to ovate or trapezoidal in outline, shell commonly thickened, concentric sculpture tending to be irregular, strong ribs alternating with weaker ones, radial ribs (if present) stronger laterally; lunule more or less asymmetrical. Hinge with two cardinals, anterior right covered in some, lateral laminae commonly duplicate in LV; ligament marginal to inframarginal, without separated resilium; shell internally more or less punctate or grooved; anterior adductor scar with ventral extension separate from pallial line, forming moderately to strongly divergent or elongate digitation. Margin crenate or smooth. *Sil.-Rec.*

Subfamily LUCININAE Fleming, 1828

[*nom. transl.* CHAVAN, herein (*ex* Lucinidae FLEMING, 1828)]

Shell generally solid, more or less lenticular, convex, with concentric and commonly

radial sculpture. Anterior scars short. *L. Jur.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Lucina*.—2. *Lucinisca*.—3. *Recurvella*.—4. *Volupia*.—5. *Here*.—6. *Herella*.—7. *Linga*.—8. *Illesca*.—9. *PleuroLucina*.—10. *Bellilucina*.—11. *Parvilucina*.—12. *Cavilinga*.—13. *MicroLoripes*.—14. *Callucinella*.—15. *Callucina*.—16. *Pseudolucinisca*.—17. *Striolucina*.—18. *Callucinopsis*.—19. *Barbierella*.—20. *Pillucina*.—21. *Wallucina*.—22. *Loripes*.—23. *Nevenulora*.—24. *Jagonoma*.—25. *Megaxinus*.—26. *Pteromyrtea*.—27. *Ctena*.—28. *Talocodakia*.—29. *Epicodakia*.—30. *Luciniola*.—31. *Codakia*.—32. *Epilucina*.—33. *Cavilucina*.—34. *Mesolinga*.

Lucina BRUGUIÈRE, 1797 [**Venus jamaicensis* SPENGLER, 1784; SD GRAY, 1847] [*non Lucina* LAMARCK, 1799] [= *Egraca* LEACH in GRAY, 1852 (obj.); *Dentilucina* P. FISCHER, 1887 (obj.); *Phacoides* AGASSIZ, 1845]. Medium-sized to large, subtrapezoidal, more or less flattened, with well-marked dorsal areas; sculpture of somewhat evenly spaced concentric lamellae with some areas smoother than remainder of disc. Cardinals two, straight, 3a obsolete, 3b and 4b oblique and compressed, laterals with posterior ones distant; anterior scars elongate; inner shell margin with fine crenulations which tend to vanish. *U.Cret.-Rec.*, Eu.-Afr.-Asia-N.Am.

L. (Lucina) [1] [= *L. (Lepilucina)* OLSSON, 1965 (type, *L. (L.) gratis*; OD)]. Concentric lamellae well spaced, stronger posteriorly; dorsal areas clearly marked; lunule asymmetrical, elongate, narrow. Cardinals oblique, somewhat weakened; anterior scars distant from pallial line; inner margin feebly crenulate. *U.Cret.-Rec.*, Eu.-Afr.-Asia.—FIG. E2,6. **L. (L.) jamaicensis* (SPENGLER), *Rec.*, USA (Fla.); 6a-c, LV and RV hinges, LV ext., $\times 0.8$ (513, Perry).

L. (Lucinisca) DALL, 1901 [2] [**L. nassula* CONRAD, 1846; OD]. Sculpture reticulate, posterior area less distinct than anterior; lunule somewhat depressed, slightly asymmetrical. Cardinals well marked; anterior scars elliptical, in part close to pallial line; inner margin strongly dentate. *Mio.-Rec.*, N.Am.-Asia (Japan-Philip.).—FIG. E2,5. **L. (L.) nassula* CONRAD, *Rec.*, USA (Fla.); LV ext., $\times 1.5$ (Perry).—FIG. E2,8. *L. (L.) cribriaria* SAY, *Mio.*, USA (S.Car.); 8a,b, LV int., RV int., $\times 3.6$ (98).

Barbierella CHAVAN, 1938 [19] [**Lucina barbieri* DESHAYES, 1858;¹ OD]. Small, subtrigonal, sculpture strong, cancellate, with well-marked posterior

¹ New taxa introduced by DESHAYES in 1858 as indicated by first publication of descriptions are not correctly cited, as by some authors, as 1857, when plates alone were published. The names without accompanying descriptions used in 1857 were *nomina nuda*. Likewise, names of taxa used by DESHAYES on plates published in 1858 are correctly cited as 1860, when descriptions first were given.




FIG. E2. Lucinidae (Lucininae) (p. N492, N494).

- area; lunule deeply concave. Shell margin with broad internal undulations. *Eoc.-Rec.*, Eu.-Ind.O. (Mauritius).—FIG. E2,3. **B. barbieri* (DESHAYES), *M.Eoc.*(Lutet.), France; LV int., $\times 3.2$ (Chavan, n).
- Callucina** DALL, 1901 [**Lucina radians* CONRAD, 1841; OD]. Medium-sized to large, generally suborbicular, dorsal areas obsolete; lunule slightly asymmetrical. Cardinals broad but weakened in some; anterior scars tending to be narrowed; shell margin minutely crenulate internally. *L.Cret.* (*Apt.*)-*Rec.*, Eu.-Afr.-Asia-N.Am.-Australia.
- C. (Callucina)** [15]. Rounded to trigonal, moderately convex; sculpture concentric, locally lamellose, more or less marked, with weak intercalated radial ribs; lunule asymmetrical, not excavated. Cardinals commonly weakened, *3a* practically obsolete; laterals distant, minute; anterior scars moderately short; shell margin finely crenulate internally. *L.Cret.* (*Apt.*)-*Rec.*, Eu.-Afr.-N.Am.-W. Asia.—FIG. E2,9. *C. (C.) hoernesii* (DESHAYES), *Eoc.*, France; *9a,b*, LV int., RV int., $\times 2.8$ (98).
- C. (Callucinopsis)** CHAVAN, 1959 [18] [**Lucina foucardi* DESHAYES, 1858; OD]. Moderately convex; dense concentric ribbing; lunule excavated. Hinge teeth present, *3a* and *A1* approximate; *3b* bifid; anterior scars rather long, subparallel to pallial line but distant from it; shell margin very finely denticulate internally. *U.Cret.* (*Senon*)-*Rec.*, Eu.-N.Afr.-E.Afr.—FIG. E2,1. **C. (C.) foucardi* (DESHAYES), *Paleoc.*, France; *1a,b*, LV hinge, RV int., $\times 2$ (Chavan, n).
- C. (Pseudolucinisca)** CHAVAN, 1959 [16] [**Lucina lacteola* TATE, 1897 (*pro. L. lactea* A. ADAMS, 1855 (*non Tellina lactea* POLI, 1795=*L. lactea* LAMARCK, 1818) (= *L. concentrica* ADAMS & ANGAS, 1863, *non* LAMARCK, 1806); OD]. Globose, with reticulate sculpture; lunule asymmetrical excavated, partly obliterating tooth 2; *3b* trigonal, laterals lacking; anterior scars short, distant from pallial line; inner margin of shell with long crenulations. *Oligo.-Rec.*, Eu.-Australia.—FIG. E2,4. **C. (P.) lacteola* (TATE), *Rec.*, Australia; *4a,b*, LV and RV hinges, $\times 0.6$ (Chavan, n).
- C. (Striolucina)** SACCO, 1901 [17] [**Dentilucina (S.) persolida*; OD]. Large, convex, with irregular lamellose concentric ribs; lunule elongate, ligament broad. Hinge with cardinals and feeble laterals; anterior scars long, moderately narrow; shell margin finely crenulated internally. *Eoc.-Plio.*, S.Eu.
- Cavilucina** P. FISCHER, 1887 [33] [**Lucina sulcata* LAMARCK, 1806 (1808); OD]. Rather small, rounded-oblong, thick, somewhat inflated; surface with concentric strong ribs; lunule deeply sunken, projecting below along hinge margin and backward, obliterating hinge; escutcheon wanting. Only anterior laterals developed; anterior scars short, very divergent, narrow; shell margin smooth internally. *Eoc.*, Eu.—FIG. E2,10. **C. sulcata* (LAMARCK), *M.Eoc.*, France; *10a,b*, LV and RV hinges, $\times 2.4$ (Chavan, n).
- Codakia** SCOPOLI, 1777 [**Chama codak* ADANSON, 1757 (*errore pro C. codok*, pre-Linnaean, invalid = **Venus orbicularis* LINNÉ, 1758; M) [= *Lenticularia* SCHUMACHER, 1817 (type, *L. punctata* = *Venus punctata* LINNÉ, 1758; M); *Lenticularia* GRAY, 1847 (missp.); *Chama* MARTINI in MÖRCH, 1853 (*non* LINNÉ, 1758); *Antilla* DE GREGORIO, 1885; *Codakia* P. FISCHER, 1887; *Lintellaria* BUCQUOY, DAUTZENBERG, & DOLLFUS, 1898 (missp.); *Pexocodakia* IREDALE, 1930 (type, *Lucina rugifera* REEVE, 1850)]. Medium-sized to large, lenticular, compressed, slightly inequilateral, no differentiated areas; concentric or radial sculpture predominant; small deep lunule in RV; ligament broadly sunken. Anterior laterals and cardinals well marked, posterior laterals obsolete or small; anterior scars moderately short and broad, divergent; shell margin smooth. *U.Jur.-Rec.*, Eu.-Am.-Asia-Australia-Pac.
- C. (Codakia)** [31]. Generally large, sculpture reticulate; lunule small, very asymmetrical, depressed; ligamentary groove broad. Posterior laterals obsolete. *Paleoc.-Rec.*, Eu.-Am.-Asia-Australia-Pac.—FIG. E2,11. **C. (C.) orbicularis* (LINNÉ), *Rec.*, W.Indies; *11a*, LV ext., $\times 0.8$ (REEVE); *11b,c*, LV and RV hinges, $\times 0.5$ (513).
- C. (Epilucina)** DALL, 1901 [32] [**Lucina californica* CONRAD, 1837; OD]. Medium-sized, with concentric sculpture only; lunule rather large, asymmetrical; ligamentary groove of moderate extent. Posterior laterals small. *U.Jur.-Rec.*, Eu.-N.Am.—FIG. E3,13. **C. (E.) californica* (CONRAD), *Rec.*, USA (Calif.); *13a,b*, LV int., RV int., $\times 1.7$ (98).
- Ctena** MÖRCH, 1861 [**Lucina pectinata* CARPENTER, 1857 (*non* C. B. ADAMS, 1852); SD DALL, BARTSCH & REHDER, 1938] [= *Jagonia* RECLUZ, 1869 (type, *Venus eburnea* GMELIN, 1790; OD)]. Transversely ovate to elliptical, anteriorly enlarged, compressed; beaks pointed; sculpture regularly reticulated; no differentiated areas; lunule concave. Cardinals short or thin, laterals not duplicate, originating under hinge plate; anterior scars narrowed at ends. *M.Eoc.-Rec.*, Eu.-N.Am.-Pac.-Ind. O.-W.Afr.-Australia.
- C. (Ctena)** [27]. Moderately convex, somewhat inequilateral, with well-marked radial ribs and lunule. Cardinals, short but normally developed, laterals strong; shell margin finely denticulated internally. *M.Eoc.-Rec.*, Eu.-N.Am.-Pac.-Ind. O.-W.Afr.—FIG. E3,3. *C. (C.) eburnea* (GMELIN), *Rec.*, Atlantic; *3a,b*, LV and RV hinges, enl. (513).
- C. (Talocodakia)** IREDALE, 1936 [28] [**Epicodakia kennethi*; OD]. Compressed, very inequilateral, radial riblets at ends only; lunule obsolete; ligamentary groove shallow. Cardinals small,




FIG. E3. Lucinidae (Lucininae) (p. N494, N496, N498).

laterals small, distant; anterior scars short, reniform, moderately divergent; shell margin smooth internally. *Oligo-Rec.*, Australia.

Epicodakia IREDALE, 1930 [29] [**E. consettiana* (= *Lucina minima* TENISON-WOODS, 1876, non ROEMER, 1836); OD]. Compressed, transversely elliptical to ovate, anteriorly enlarged; with dichotomous radial ribs predominating over concentric ribs; ligament in deep groove. Cardinals and laterals well marked, latter duplicate on LV; anterior scars elliptical, divergent, rounded at ends; marginal internal crenulations of shell almost obsolete. *Pleist.-Rec.*, Australia-N.Am.-Asia.—FIG. E3,9. *E. divergens* (PHILIPPI), *Rec.*, Japan; 9a,b, LV int., RV int., $\times 2.3$ (119, mod.).—FIG. E2,2. **E. consettiana*, *Rec.*, Australia; LV ext., $\times 1.35$ (Iredale).

Here GABB, 1866 [**Lucina* (H.) *richtofeni* (= *Lucina excavata* CARPENTER, 1857); SD STOLICZKA, 1871]. Medium-sized to small, rounded, with weak anterior and posterior areas; concentric ribs tending to vanish; lunule deeply excavated, covering part of hinge; anterior scars somewhat short, attenuated; internal shell margin with minute denticles. *Paleoc.-Rec.*, W.Eu.-W.N.Am.-N.Afr.

H. (Here) [5]. Concentric ribs fine, but well-marked; lunule rather broadly rounded, obliterating only anterior cardinal, posterior ones well marked; anterior laterals minute, posterior laterals distant; anterior scars divergent at their ends; internal shell margin finely crenulated. *Eoc.-Rec.*, W.Eu.-W.N.Am.—FIG. E3,1. **H. (H.) excavata* (CARPENTER), Mio., USA (Calif.); 1a,b, LV int., RV int., $\times 2.25$; 1c, LV ext., $\times 1.5$ (98; Gabb, 1869).

H. (Herella) CHAVAN, 1942 [6] [**Lucina levesquei* D'ORBIGNY in DESHAYES, 1858; OD] [= *?Tuberculina* DE GREGORIO, 1882 (*nom. dub.*)]. Sculpture weakly cancellate; lunule narrow, projecting under beaks, covering obsolete RV cardinals, those of LV tuberculiform; laterals feeble; anterior scars short, wholly divergent; internal shell margin almost smooth. *Paleoc.-Oligo.*, W.Eu.-N.Afr.—FIG. E2,7. **H. (H.) levesquei* (D'ORBIGNY), L.Eoc. (Cuis.), France (Paris basin); 7a-c, LV int., RV int., LV ext., $\times 3.6$ (Chavan, n; 257).

Linga DE GREGORIO, 1884 [**Lucina columbella* LAMARCK, 1819; SD SACCO, 1889] [= *Quasilucina* STEWART, 1930 (type, *Lucina carinifera* CONRAD, 1833; OD)]. Medium-sized, rounded, tumid; concentric ribbing more or less lamellose; lunule sunken, short, cordiform. Hinge teeth strong, short, LV laterals duplicate; anterior scars moderately short; shell margin internally denticulated. *?Paleoc., Eoc.-Rec.*, Eu.-Am.-Afr.-Asia-Australia.

L. (Linga) [7]. Rounded; concentric sculpture nearly uniform consisting of numerous equidistant lamellae; dorsal angulation rounded.

Hinge teeth 3a and 2 present; shell margin crenulated internally. *?Paleoc., Eoc.-Rec.*, W.Eu.-W.Afr.-N.Am.-C.Am.—FIG. E3,7,8. **L. (L.) columbella* (LAMARCK) *basteroti* (AGASSIZ), L. Mio. (Burdigal.), France (Gironde); 7a,b, LV int., RV int., $\times 1.8$; 8, LV-ext., $\times 0.8$ (98).

L. (Bellucina) DALL, 1901 [10] [**Parvilucina eucosmia* DALL, 1901 (= *Lucina pisum* REEVE, Aug. 1850; non SOWERBY, 1836; nec D'ORBIGNY, 1841; nec PHILIPPI, Apr. 1850) (= **Lucina semperiana* ISSEL, 1869); OD] [= *Cardiolucina* SACCO, 1901 (type, *Cardium agassizi* MICHELOTTI, 1847; = *Cardita agassizi* MICHELOTTI, 1839; OD)]. Small, rounded, very tumid; sculpture of heavy concentric ribs, crossed, but not interrupted, by radiating riblets; lunule small, not immersed. Complete lucinoid hinge with AIV and PIV feebly marked; anterior scars broad, short; shell margin deeply but finely crenulated internally. *M.Eoc.-Rec.*, Eu.-Afr.-Asia (Japan)-Australia.—FIG. E3,2,4. **L. (B.) semperiana* (ISSEL), *Rec.*, Ind.O.; 2, LV ext., $\times 1.4$ (Reeve); 4a,b, LV and RV hinges, enlarged (513).

L. (Illesca) OLSSON, 1932 [8] [**Phacoides* (Here) *andersoni* OLSSON, 1930; OD]. Smaller than *L. (Linga)*, with concentric sculpture of strong well-spaced folds. Hinge teeth 3a and 2 obliterated by totally immersed lunule; anterior scars ovate, short; internal shell margin finely crenulated. *U.Eoc.*, S.Am. (Peru).—FIG. E4,3. *L. (I.) andersoni* (OLSSON); 3a, LV ext., $\times 2.4$; 3b,c, LV int., RV int., $\times 3.6$, $\times 4.8$ (Olsson, 1932).

L. (PleuroLucina) DALL, 1901 [9] [**Lucina leucocyma* DALL, 1886 (1889); OD] [= *Dallucina* OLSSON & HARBISON, 1953 (type, *Lucina* (Here) *amabilis* DALL, 1898; OD); *L. (P.) undata* OLSSON, 1965 (type, *L. (P.) follis*; OD)]. Smaller than *Linga*, oblong, more or less trigonal to quadrate, with sharp dorsal angulation; radial sculpture laterally strong; lunule not immersed. Cardinal 2 larger than 4b; broadly crenulated to foliated shell margin internally. *Oligo.-Rec.*, C. Am.-N.Am.—FIG. E4,7a. *L. (P.) quadricostata* DALL, Mio., Jamaica; LV ext., $\times 6$ (1005).—FIG. E4,7b,c. *L. (P.) undata* (CARPENTER), *Rec.*, Mex.; 7b,c, LV int., RV int., $\times 3.6$ (98).

Loripes POLI, 1791 [22] [**Tellina lactea* POLI, 1791 (non LINNÉ, 1758) (= *Amphidesma lucinalis* LAMARCK, 1818); SD GRAY, 1847] [*non Loripes* SCHWEIGER, 1820] [= *Ligula* MENKE, 1830 (non MONTAGU, 1803); *Lucinida* D'ORBIGNY, 1846 (type, *L. cryptella*, = *L. brasiliensis*, = *Lucina brasiliensis* PHILIPPI, 1848; OD)]. Lenticular, compressed, somewhat gibbous; sculpture feeble, concentric; areas almost obsolete; ligament internal. Hinge teeth comprising 1 RV cardinal and 2 weak LV laterals; anterior scars subelliptical; internal shell margin smooth. *Oligo.-Rec.*, Eu.-Afr.-S.Am.-Japan.—FIG. E4,1. **L. lucinalis* (LA-




FIG. E4. Lucinidae (Lucininae) (p. N496, N498-N499).

MARCK), Rec., Eng.(Sussex); LV int., $\times 2.4$ (905a).

Luciniola SKEAT & MADSEN, 1898 [30] [**Venus pumila* GOLDFUSS, 1840; OD]. Small, subtrigonal, posteriorly enlarged, moderately convex, thick, with pointed beaks, with coarse concentric sculpture; lunule small. Hinge as in *Epicodakia* but with external ligament; anterior scars subparallel to pallial line. *L.Jur.(U.Lias.)*, Eu.(Ger.-Denm.). —FIG. E3,12. **L. pumila* (GOLDFUSS), Denm.; 12a,b, RV hinge, LV int., $\times 3.2$ (Skeat & Madsen, 1898).

Megaxinus BRUGNONE, 1880 [**Lucina rostrata* PECCHIOLI, 1864; OD] [= *Stewartia* OLSSON & HARBISON, 1953 (type, *Lucina anodonta* SAX, 1824; OD)]. Irregularly lenticular to oblong compressed; hinge broad, almost edentulous, with remnants of 3b and 4; anterior scar broad, juxtaposed to pallial line; shell margin internally smooth. *Palcos.-Rec.*, Eu.-Am.-N.Z.

M. (Megaxinus) [25]. Very irregular, thickened and gibbous, with ill-defined areas; lunule profoundly penetrating into broad trigonal hinge plate. Anterior scars almost continuous to pallial line. *Oligo.-Rec.*, Eu.-Am.-Red Sea.—FIG. E3, 5. **M. (M.) rostratus* (PECCHIOLI), Plio., Italy; 5a,b, LV int., RV int., $\times 1.6$ (Pecchioli, 1864).

M. (Pteromyrta) FINLAY, 1926 [26] [**Cyclina dispar* HUTTON, 1873 (= *Lucinida laevifoliata* MARSHALL & MURDOCH, 1919); OD]. Thin, rounded, with alate areas; lunule moderately depressed, obliterating anterior cardinals; posterior ones more or less marked; anterior scars wholly contiguous to pallial line. *U.Cret.-Pleist.*, N.Z.—FIG. E4,10. **M. (P.) dispar* (HUTTON), Plio., N.Z.; 10a,b, LV int., RV int., $\times 2.2$ (Chavan, n).

Mesolinga CHAVAN, 1951 [34] [**Lucina plebeia* CONTEJEAN, 1859 (non GIEBEL, 1856) (= **M. typica* CHAVAN, 1952); OD]. Globose, anteriorly produced and rounded, inequilateral; with concentric approximate ribs; ligament marginal, narrow. One right cardinal, 2 left ones, and duplicate right, simple left, tuberculiform laterals; reniform anterior scars narrowed at their end; shell margin internally smooth. *U.Jur.-Cret.*, Eu.-N.Am.-?Asia.—FIG. E3,11. **M. typica* CHAVAN, U.Jur. (Astart.), France (Calvados); 11a,b, LV int., RV int., $\times 2.4$ (Chavan, n).

Nevenulora IREDALE, 1930 [**Lucinida hilaira* HEDLEY, 1916; OD]. Moderately convex; concentric sculpture of more or less vanishing alternate distant lamellae and finer intercalated striae; lunule well marked. Hinge with cardinals and laterals, both evanescent in some forms; anterior scars moderately short; shell margin smooth internally. *U.Jur.(Raurac.)-Rec.*, Eu.-Australia-N.Z.

N. (Nevenulora) [23]. Irregularly striated, almost smooth, anteriorly angular, posteriorly rounded; lunule moderately concave. Teeth small;




FIG. E5. Lucinidae (Lucininae) (p. N498-N499).

anterior scars somewhat divergent. *L.Eoc.-Rec.*, Eu.-Australia.—FIG. E4,8. **N. hilaira* (HEDLEY), Rec., Australia; 8a,b, LV int., RV hinge, $\times 2$ (Chavan, n).


N. (Jagonoma) CHAVAN, 1946 [24] [**Lucina circumcisa* ZITTEL & GOUBERT, 1861; OD] [= *Dilora* MARWICK, 1948 (type, *D. lorea*; OD)]. Rounded at both ends; sculpture of alternate stronger and smaller concentric fine ribs. Hinge teeth well developed, 3b narrow, not bifid; anterior scars not widely diverging. *U.Jur.(Raurac.)-Plio.*, Eu.-N.Z.—FIG. E3,6. **N. (J.) circumcisa* (ZITTEL & GOUBERT), Astart., France; 6a,b, LV int., RV hinge, $\times 1.6$ (Chavan, n).

Parvilucina DALL, 1901 [**Lucina tenuisculpta* CARPENTER, 1864; OD]. Rather small, rounded, inflated, with finely concentric sculpture, radial striae weakening medially, anterior and posterior areas feebly marked. Thin, but almost complete hinge, with 3a but AIV and PIV obsolete; anterior scars more or less broad and short. *U.Cret.(Senon.)-Rec.*, Eu.-Afr.-N.Am.-Australia.

P. (Parvilucina) [11]. Sculpture finely reticulate; ligament marginal. Hinge teeth comprising short 3b and distant laterals; internal shell margin very finely crenulated. *U.Cret.(Senon.)-Rec.*, W.Eu.-N.Am.—FIG. E3,10. **P. (P.) tenuisculpta* (CARPENTER), Rec., USA (Calif.); 10a,b, LV int., RV int., $\times 4.4$ (98).

P. (Callucinella) CHAVAN, 1961 [14] [**Lucina albella* LAMARCK, 1806]. Lenticular, depressed; with fine irregular superficial concentric sculpture; areas obsolete; lunule long; ligament marginal. Hinge teeth comprising broad and bifid 3b and distant laterals; anterior scars rather narrow, short; internal shell margin feebly crenulated to smooth. *M.Eoc.-Plio.*, W.Eu.-Australia.

Veneroida—Lucinacea



—FIG. E5,1. **P. (C.) albella* (LAMARCK), M. Eoc.(Lutet.), France; 1a,b, LV and RV hinges, $\times 2$ (Chavan, n).

P. (Cavilinga) CHAVAN, 1937 [12] [**Lucina trisulcata* CONRAD, 1841; OD]. Rounded, trigonal, high, inequilateral, with approximate concentric close ribs and some annular rings; ligament inframarginal; lunule deep. Cardinal 3b curved, laterals approximate; anterior scars divergent, somewhat elongated. *U.Cret.-Rec.*, W.Eu.-S.Afr.-N.Am.—FIG. E5,2. **P. (C.) trisulcata* (CONRAD), Mio., USA(Md.); 2a,b, LV int., RV int., $\times 6.5$ (98).

P. (Microroripes) COSSMANN, 1910 (1912) [13] [**Lucina dentata* DEFRANCE, 1823; OD]. Rounded, globose, small; sculpture irregularly concentric; ligament internal, oblique. Anterior scars short; shell margin strongly dentated internally. ?*U. Cret.*(Senon.), *Paleoc.-Rec.*, Eu.-W.Afr.—FIG. E4,4. **P. (M.) dentata* (DEFRANCE), L.Mio. (Burdigal.), S.France; 4a,b, LV int., RV int., $\times 4$ (Chavan, n).

Pillucina PILSBRY, 1921 [20] [**P. spaldingi*; OD]. Rounded, moderately convex; sculpture of concentric and radial lines laterally strengthened, medially evanescent. Hinge short, with 2 and 3b trigonal and one almost obsolete right anterior lateral; anterior scars moderately short; internal margin of shell undulated. *Plio.-Rec.* Australasia-Pac.-Asia-Red Sea.

P. (Pillucina). Lateral tooth *AIII* small; anterior scar rather elongate. *Plio.-Rec.*, Pac.-Asia-Red Sea.—FIG. E4,9. *P. fischeriana* (ISSEL), Rec., Red Sea; 9a,b, LV int., RV int., $\times 5.6$ (98).

P. (Sydlorina) IREDALE, 1930 [**S. symbolica* (= **Codakia pisidium* HEDLEY, 1914, non DUNKER, 1860); OD]. Sculpture stronger than in *P. (Pillucina)*, anterior scar shorter and more rounded, *AIII* rather stout. Rec., Australia-Japan.

Recurvella CHAVAN, 1937 [3] [**Lucina dolabra* CONRAD, 1833; OD]. Small, subtrigonal, compressed, with high, recurved beaks; areas well marked; lunule symmetrical. Hinge high, without laterals; anterior scars short, divergent. *M.Eoc.-Oligo.* or ?*Plio.*, W.Eu.-N.Am.-?Asia.—FIG. E4,5. **R. dolabra* (CONRAD), M.Eoc.(Claiborn.), USA (Ala.); 5a, RV ext., $\times 2.4$; 5b,c, RV int., LV int., $\times 4.8$ (98; Harris, 1919).

Volupia DEFRANCE, 1829 [4] [**V. rugosa*; OD] [= *Gradilucina* COSSMANN, 1902 (type, *Lucina tabulata* DESHAYES, 1858; OD)]. Small, oblong, tumid, with prominent curved beaks; sculpture of strong annular rings; areas deep. Cardinals strong, tuberculiform; anterior scars short, ovate-oblong; internal margin of shell crenate. *M.Eoc.-U.Eoc.*, Eu.(France).—FIG. E4,6. *V. tabulata* (DESHAYES), Eoc.(Barton.), France(Paris basin); 6a, LV ext., $\times 6$; 6b,c, LV int., RV int., $\times 6$ (257).

Wallucina IREDALE, 1930 [21] [**Lucina jacksoniensis* SMITH, 1885; OD]. Rounded, inflated; sculpture of almost uniform concentric and radial lines. Hinge with cardinals broader than in *Pillucina*, 1 right anterior and 2 marked, posterior laterals; anterior scars broad, short; finely denticulated internal shell margin. *Mio.-Rec.*, Australia-Pac.-Asia-W.Afr.—FIG. E4,2a,b. *W. erythraea* (ISSEL), Red Sea; 2a,b, LV int., RV int., $\times 3.2$ (Chavan, n).—FIG. E4,2c. **W. jacksoniensis* (SMITH), Australia; RV ext., $\times 2.4$ (852).

Subfamily MYRTEINAE Chavan, new subfamily

Thin, more or less quadrangular or transverse, moderately compressed shell with concentric sculpture; anterior scars medium-sized. *L.Jur.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Mesomiltha*.—2. *Monitilora*.—3. *Prophetilora*.—4. *Gonimyrtea*.—5. *Myrtea*.—6. *Myrteopsis*.—7. *Perampliata*.—8. *Gardnerella*.—9. *Milthona*.—10. *Lucinoma*.—11. *Discomiltha*.

Myrtea TURTON, 1822 [**Venus spinifera* MONTAGU, 1803; M] [= *Cyrachaea* LEACH, 1819 (obj.); *Eulopia* DALL, 1901 (type, *Lucina sagrinata* DALL, 1886; OD); ?*Notomyrtea* IREDALE, 1924 (type, *Myrtea botanica* HEDLEY, 1918, = *Tellina brazieri* SOWERBY, 1883, non 1869; OD)]. Transversely elliptical to quadrangular, flattened; areas obsolete; sculpture of concentric, posteriorly elevated ribs with intercalated vermiculate radials in some; beaks pointed; lunule and escutcheon narrow and straight. Cardinals narrow, with 3b, 2, and 4b; RV laterals very long (those of LV on margin), originating from under plate; reniform short scars; internal shell margin smooth. ?*U.Cret.*, *Oligo.-Rec.*, Eu.-Australia-N.Z.-Asia-N.Am.

M. (Myrtea) [5] [= *Myrtaea* DALL, 1901]. Concentrically ribbed; ligament external. Teeth well developed; laterals of same length anteriorly and posteriorly. ?*U.Cret.*, N.Z.; *Oligo.-Rec.*, Eu.-N.Am.-Asia-Australia.—FIG. E6,1; E7,1. **M. spinifera* (MONTAGU), Rec., Medit.; E6,1, RV ext., $\times 0.7$ (513, Reeve); E7,1a, RV int., $\times 1$ (905a); E7,1b,c, LV and RV hinges, $\times 1$ (513).

M. (Myrteopsis) SACCO, 1901 [6] [**M. taurolaevis* (= *M. magnotaurina*); OD]. Finely striated. Ligament semi-internal. Teeth minute, *AI* shorter than *PI*. *Mio.*, S.Eu.

Discomiltha CHAVAN, 1952 [11] [**D. oehlerti* BIGOT in CHAVAN, 1952; OD]. Moderately large, suborbicular, flattened, thin; sculpture of well-spaced feeble, concentric ribs and finer striae; posterior area well marked; lunule very asymmetrical, narrow, long, projecting ligament inframarginal. Hinge teeth comprising only 2 weak cardinals




FIG. E6. Lucinidae (Myrteinae) (p. N499-N501).

partly covered by lunular expansion; shell internally punctate and grooved; anterior scars elongate, narrow, distant from pallial line; shell margin smooth internally. *U. Jur.*, Eu.—FIG. E6.6. **D. oehlerti* BIGOT, Astart., France (Calvados); 6a,b, LV int., RV int., $\times 1$ (Chavan, n).

Gardnerella CHAVAN, 1951 [8] [**Myrtaea waltonensis* GARDNER, 1926; OD]. Large, inequilateral, more or less produced anteriorly, truncated or rounded posteriorly; moderately compressed; sculpture of irregular weak concentric ribs; lunule long. Cardinals thin, 1 right, 2 left, and right laterals; anterior scars rather broad, only slightly divergent; shell internally punctate, with smooth margin. *L. Eoc.-Mio.*, Eu.-N.Afr.-N.Am.—FIG. E6.5. *G. cosmanni* (CHAVAN), *L. Eoc.*, France; 5a,b, LV int., ext., $\times 0.9$ (Chavan, 1941).

Gonimyrttea MARWICK, 1929 [4] [**Loripes concinnus* HUTTON, 1885 (= *L. icterica* LAMY, 1920; non *Lucina concinna* ADAMS, 1870; nec *L. icterica* REEVE, 1850); OD] [= *Alucinoma* HABE, 1958 (type, *A. soyogae*; OD)]. Small, subquadrangular, compressed, sculpture fine, concentric; lunule depressed, asymmetrical. Hinge tooth 2 more or less adherent to its edge, 3a lacking, 3b present, *Al* small; anterior scars reniform, short; shell margin internally smooth. *Paleoc.-Rec.*, Eu.-N.Afr.-N.Am.-C. Am.-N. Z.-Australia-Japan.—FIG. E7.3. *G. galeottiana* (NYST), *Eoc.* (Barton.), Belg.; 3a,b, LV ext., int., $\times 4$ (Glibert, 1936).

Lucinoma DALL, 1901 [10] [**Lucina filosa* STIMPSON, 1851; OD] [= *Triodonta* GRAY, 1851 (non BORY, 1827; nec MULSANT, 1842; nec AGASSIZ, 1846) (obj.); *Thiatira* LEACH, 1819]. Medium-sized to large, lenticular, moderately convex, posterior margin rectilinear; with regularly spaced annular ribs and intercalated finer ones; lunule lanceolate, long, not sunken or bent. Two compressed teeth in each valve, 2 and 3b bifid, and feeble laminar *Al*; anterior scars narrow, long, diverging; shell margin internally smooth. *Oligo.-Rec.*, Eu.-Red Sea-N.Am.-Pac.-Australia-Japan.—FIG. E6.8. *L. borealis* (LINNÉ), *Rec.*, Eu.; 8a,b, LV and RV hinges, $\times 1$ (513).

Mesomiltha CHAVAN, 1938 [1] [**Lucina pulchra* ZITTEL & GOUBERT, 1861; OD]. Medium-sized, transversely ovate-rounded, posteriorly depressed; with fine, regular equidistant concentric ribs and very fine intercalated radial lines; lunule lanceolate projecting below in dentiform elongation; long ligament. Hinge teeth comprising 1 RV, 2 LV cardinals, and thin distant laterals; anterior scars curved, detached at their ends; shell margin internally smooth. *L. Jur.* (*Lias.*)-*U. Jur.*, ?*Cret.*, Eu.—FIG. E6.2. **M. pulchra* (ZITTEL & GOUBERT), *U. Jur.* (Astart.), France (Calvados); 2a,b, LV int., RV int., $\times 2$ (Chavan, n).

Milthona MARWICK, 1931 [9] [**M. glomerosa*; OD]. Moderately large, inequilateral, anteriorly rounded, posteriorly enlarged, with concentric

coarse, wide-spaced ribs; not differentiated posterior area; lunule somewhat depressed, cordiform. Hinge teeth 2 and 3b bifid, 4b strong, laterals lacking; anterior scars narrowly linguiform, divergent; shell margin smooth internally. *Mio.*, N.Z.—FIG. E6, 7. **M. glomerosa*; RV int., $\times 0.9$ (595).

Monitilora IREDALE, 1930 [**Lucina ramsayi* SMITH, 1885 (= *Loripes icterica* ANGAS, 1867, non *Lucina icterica* REEVE, 1850)]. Rounded, with concentric regularly spaced ribs and radial intercalated lines; lunule lanceolate, its edge forming ridge along hinge plate; ligament elongated, inframarginal. Single weak cardinal in each valve, RV lateral obsolete; anterior scars narrow, distant from pallial line; internal rugosities present, margin smooth. *Paleoc.-Rec.*, Eu.-Australia-Asia.

M. (Monitilora) [2]. Medium-sized, circular to elliptical in outline, relatively tumid, dorsal area not distinguished; lunule straight-edged; scars moderately long; shell internally punctate. *Paleoc.-Rec.*, Eu.-Australia-Asia.—FIG. E6,4. *M. elegans* (DEFRANCE), *Eoc.*, France; 4a,b, LV int., RV int., $\times 2.7$ (98).

M. (Propetilora) IREDALE, 1930 [3] [**P. arizela*; OD]. Large, ovate, enlarged backward, com-




FIG. E7. Lucinidae (Myrteinae) (p. N499-N501).




FIG. E8. Lucinidae (Milthinae) (p. N502, N504).

pressed, with faint dorsal area; lunule bent inward. Hinge almost obsolete; scars linguiform, relatively long; shell internally pustulose. *Paleoc.-Rec.*, Eu.-Australia.—FIG. E6,3. **M. (P.) arizela*, *Rec.*, Queensl.; LV int., $\times 1$ (Iredale, 1930).

?**Perampliata** ARKELL, 1936 [7] [*pro Ampliata* ARKELL, 1934, p. 282 (non WAGNER, 1907)] [**Tellina ampliata* PHILLIPS, 1829, p. 324; OD]. Large, compressed, with close-spaced concentric ribs; beaks small; lunule lacking, escutcheon broad. Stout short hinge plate with 1 RV and 2 LV cardinals; pallial line and anterior scars unknown. *Jur.*, Eu.(G.Brit.-France).—FIG. E7,2. **P. ampliata* (PHILLIPS), *U.Jur.*(Corall.), Eng.; LV int., $\times 0.7$ (Arkell, 1936).

Subfamily MILTHINAE Chavan,
new subfamily

Shell relatively solid, generally compressed. Sculpture concentric, faint, irregular to vanishing; anterior scars long. *Sil.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Phenacocyclas*.—2. *Pterolucina*.—3. *Myrtucina*.

5302357871

4. *Jagolucina*.—5. *Saxolucina*.—6. *Plastomiltha*.—7. *Claibornites*.—8. *Codalucina*.—9. *Miltha*.—10. *Recticardo*.—11. *Trinitasia*.—12. *Ilionia*.—13. *Eomiltha*.—14. *Gibbolucina*.—15. *Pseudomiltha*.—16. *Zorrira*.—17. *Eamesiella*.—18. *Pegophysema*.—19. *Eophysema*.—20. *Anodontia*.—21. *Cavatidens*.—22. *Loripinus*.

Miltha H. & A. ADAMS, 1857 [**Lucina childreni* GRAY, 1825; OD] [= *Miltheoidea* MARWICK, 1930 (type, *Miltha neozelandica* MARSHALL & MURDOCK, 1921; OD)]. Discoidal, flattened; sculpture of unequal concentric striae; ligament on enlarged nymph. Hinge teeth *3b* and 2 somewhat bifid; shell margin smooth internally. *U.Cret.-Rec.*, Eu.-N.Am.-Australia-N.Z.

M. (Miltha) [9]. Large, subcircular to ovate-oblong, slightly inequivalve; surface smooth or lamellose, with faint areas; lunule asymmetrical, striated. Anterior scars long, club-shaped. *U.Cret.-Rec.*, Eu.-N.Am.-Australia-N.Z.—FIG. E8,2. **M. (M.) childreni* (GRAY), *Rec.*, Brazil; LV hinge, $\times 1$ (513, Sowerby).

M. (Recticardo) COSSMANN, 1908 [10] [**Phacoides (R.) rutoti*; OD]. Medium-sized, subelliptical; surface with concentric dense striation without differentiated areas; lunule depressed, almost symmetrical, its projecting edge prolonged by ridge-like anterior lateral tooth, posterior lateral weak; anterior scars relatively short, penetrating upward between lunular edge and anterior lateral. *U.Cret.-Paleoc.*, Eu.-N.Am.—FIG. E9,3. **M. (R.) rutoti* (COSSMANN), *Paleoc.*, Belg.; *3a,b*, LV and RV hinges, $\times 2$ (98).

Anodontia LINK, 1807 [**A. alba* (= **Venus edentula* LINNÉ, 1758); OD]. Rounded, tumid, slightly inequilateral; sculpture of irregular growths and very fine radial striae; anterior area more or less marked, posterior area obsolete; beaks prosogyrous; lunule ill-defined; ligament oblique, sunken. Hinge edentulous except for faint tuberculiform cardinal; anterior scars broad, short, very divergent; shell margin internally smooth. *Eoc.-Rec.*, Eu.-Asia-N.Am.-Pac.-Australia-Afr.

A. (Anodontia) [20]. Medium-sized to large, globose, rounded in front, slightly truncate posteriorly; surface with concentric and radial lines, anterior area ill-defined; lunule depressed; ligament sunken but inserted upon cardinal elongation. *Eoc.-Rec.*, Eu.-Asia-N.Am.-Pac.O.-Australia.—FIG. E9,6. **A. edentula* (LINNÉ), *Rec.*, Ind. O.; RV int., $\times 1.3$ (98).

A. (Cavatidens) IREDALE, 1930 [21] [**C. omissa*; OD]. Small, globose, laterally truncate, with concentric sculpture only, dorsal area distinct; beaks high; ligament sunken on cardinal process. *Rec.*, Australia.

A. (Loripinus) DE MONTEROSATO, 1883 [22] [**Lucina fragilis* PHILIPPI, 1836; SD VON MARTENS, 1884]. Small, very globose, thin, anteriorly

rounded, not depressed; ligament strongly depressed, without cardinal elongation or process. Hinge tooth *3b* directed backward. *Eoc.-Rec.*, S. Eu.-Afr.-Asia-Japan-?N.Am.

Claibornites STEWART, 1930 [**Lucina rotunda* LEA, 1833; OD]. Medium-sized, lenticular, flattened; sculpture of concentric striae, dorsal areas obsolete; lunule lanceolate. Hinge with narrow cardinals and strong anterior laterals; anterior scars elongate, narrow; shell margin internally smooth. *Paleoc.-Oligo.*, Eu.-N.Am.

C. (Claibornites) [7]. Thick; sculpture coarsely concentric; areas weak; ligament on large depressed nymph, gibbous internally. Hinge as in *Saxolucina*, but with *3a* less obsolete and *3b* only obscurely bifid, *A1* stronger; anterior scars long. *Eoc.*, N.Am.—FIG. E9,1. **C. (C.) rotunda* (LEA), *M.Eoc.*, USA (Ala.); *1a,b*, LV int., RV hinge, $\times 0.9$ (Harris, 1919).

C. (Codalucina) STEWART, 1930 [8] [**Lucina uncinata* DEFRANCE, 1823; OD]. Thin, sculpture finely concentric; no areas; ligament deeply sunken, in broad groove. Hinge with well-marked cardinals and anterior laterals, weak posterior laterals; scars relatively long, linguiform. *Paleoc.-Oligo.*, Eu.-N.Am.—FIG. E9,2. **C. (C.) uncinata* (DEFRANCE), *Paleoc.*, France; LV int., $\times 1.3$ (98).

Eamesiella CHAVAN, 1951 [17] [*pro Pseudolucina* CHAVAN, 1947 (*non* WILCKENS, 1909)] [**Lucina corrugata* DESHAYES, 1843; OD]. Subquadrangular, rather inflated; surface with distant concentric lamellose waves. Edentulous; stout nymph; shell margin smooth internally. ?*Eoc.* or *Oligo.*, *Rec.*, Afr.-Australia-Eu.—FIG. E8,3. **E. corrugata* (DESHAYES), *Rec.*, Pac.; *3a,b*, LV ext., int., $\times 2$ (Alcock & Anderson, 1897).

Gibbolucina COSSMANN, 1904 [**Venus callosa* LAMARCK, 1806; OD] [= ?*Elathia* ISSEL, 1869 (type, *E. arconatii*; OD) (*gen. dub.*)]. Irregularly compressed; sculpture of coarse lamellae or ribs, with narrow deep areas; lunule concave, short, broad. Trigonal hinge plate with *3a* thin, *3b*, strongly bilobate, *2*, *4b* present, laterals lacking; short stout nymph; fine radial internal threads; anterior scars elongate, narrow, subparallel to pallial line; shell margin internally smooth. *Cret.-Rec.*, Eu.-Am.-Australia-Afr.

G. (Gibbolucina) [14]. Small, moderately convex, subtrigonal, gibbose; lunule depressed, partly covering ill-defined cardinals, *3a* obsolete; short scars. *L.Eoc.-Rec.*, Eu.-Afr.-Australia-?Red Sea.—FIG. E8,1; E10,2. **G. (G.) callosa* (LAMARCK), *M. Eoc.*, France; E8, *1a,b*, RV ext., int., $\times 1$ (257); E10, *2a,b*, RV ext., int., $\times 1$ (257).

G. (Eomiltha) COSSMANN, 1910 (1912) [13]. [**Lucina contorta* DEFRANCE, 1823; OD]. Medium-sized, almost flat, transversely subrhomboidal; with more or less elevated concentric lamellae; lunule not sunken. Cardinals well defined;




FIG. E9. Lucinidae (Milthinae) (p. N502, N504-N506).




FIG. E10. Lucinidae (Milthinae) (2); Divaricellinae (1,4-5); (Genus Uncertain) (3) (p. N502, N504, N506, N508).

scars relatively long. *Cret.-Rec.*, Eu.-Am.-E.Afr.—FIG. E9,5. **G. (E.) contorta* (DEFRANCE), *Paleoc.*, France; 5a, RV ext., $\times 0.7$; 5b,c, RV int., LV hinge, $\times 1$ (Chavan, n).

Illionia BILLINGS, 1875 [12] [**I. canadensis*; SD WHITEAVES, 1884] [= *ProLucina* DALL, 1896 (type, *Lucina prisca* HISINGER, 1837; OD); *Platymermis* NOETLING, 1883]. Irregularly rhomboidal, transverse, compressed, with postumbonal sulcus; lunule concave, long. Anterior scars long, narrow, linguiform. *Sil.*, N.Eu.-N.Am.—FIG. E9,9. *I. prisca* (HISINGER), Sweden; LV int. mold, $\times 0.7$ (1026).

Jagolucina CHAVAN, 1939 [4] [**Lucina concava* DEFRANCE, 1823; OD]. Rather inflated, rounded, inequilateral, without areas but with irregular growths; beaks prosogyrous; large lunule moderately asymmetrical. Two cardinals in each valve, 2 and 3b clearly bifid, with distant well-marked laterals; anterior scars very divergent, attenuated, short; shell margin internally smooth. *Paleoc.-Eoc.*, Eu.—FIG. E11,1. **J. concava* (DEFRANCE), L. Eoc., France; 1a,b, RV int., LV int., $\times 2$ (98).

Myrtucina VOKES, 1939 [3] [**Lucina roseburgensis* HENDON in TURNER, 1938; OD]. Medium-sized, compressed, transversely rounded, with irregular lamellose growths and well-marked areas. Cardinals broadly trigonal, 3b almost bifid, 2 and 4b somewhat divergent, laterals on hinge plate comprising strong *AI* and faint *PI*; nymph elongate; shell margin internally smooth. *Eoc.*, Eu.-N.Am.—FIG. E8,4. **M. roseburgensis* (HENDON), USA (Ore.); 4a,b, LV and RV hinges, $\times 3$ (Vokes, 1939).

Pegophysema STEWART, 1930 [18] [**Lucina schrammi* CROSSE, 1876; OD]. Rounded, tumid, slightly inequilateral; sculpture of irregular growths, with well-marked anterior area, posterior area obsolete; lunule narrow, depressed. Cardinal plate triangular without protuberances; anterior scars scarcely divergent, long and narrow; shell margin internally smooth. *Eoc.-Rec.*, N.Am.-N.Afr.-Eu. [= *Lissosphaira* OLSSON, 1961 (type, *Lucina spherica* DALL & OCHSNER, 1928; OD) (subj.).]

P. (Pegophysema). [18]. Subcircular; surface with coarse concentric ribs and well-marked, subalate, anterior area. Trigonal hinge plate, without teeth; anterior scars narrow. *Oligo.-Rec.*, N.Am.-Eu.—FIG. E9,12. *P. (P.) chrysostoma* (MEUSCHEN), *Rec.*, W.Indies; RV int., $\times 0.9$ (98).

P. (Eophysema) STEWART, 1930 [19] [**Lucina subvexa* CONRAD, 1832 (1846); OD]. Subovate, anteriorly produced, posteriorly enlarged; surface with concentric and radial striae, anterior area vanishing at adult stage. Elongate hinge plate with ill-defined transverse small tooth and left posterior lateral; relatively broad anterior scars. *Eoc.*, ?*Oligo.*, N.Am.-N.Afr.-Eu.

Phenacocyclus LA ROCQUE, 1950 [1] [**P. pohli*; OD]. Irregularly rhomboidal, oblong, compressed; surface only concentrically striated, anteriorly plicate and with deep posterior area corresponding to notch in ventral margin; lunule small; escutcheon narrow. Probably one small anterior lateral and two presumably small cardinals; sigmoid internal groove; falciform, enlarged anterior scars; shell margin internally smooth. *Dev.*, N.Am.-Eu.—FIG. E9,7. **P. pohli*, M.Dev., USA (Mich.); 7a,b, RV int., ext., $\times 0.7$ (530).

Pseudomiltha P. FISCHER, 1887 [**Lucina gigantea* DESHAYES, 1825; OD]. Large, flattened, transversely subquadrangular to rounded, with small erect beaks; surface concentrically ribbed or smooth. Teeth obsolete or absent; short stout nymph; interior punctate, with also fine radial threads and low flange bordering pallial line; long, narrow anterior scars; shell margin internally smooth. ?*Jur.*, *Eoc.-Mio.*, Eu.-Asia-S.Am.-W.Indies.

P. (Pseudomiltha) [15]. Broadly elliptical to ovate; dorsal areas lacking. Hinge totally edentulous. ?*Jur.*, *Eoc.-Oligo.*, Eu.-Asia.—FIG. E11,3. **P. (P.) gigantea* (DESHAYES), M.Eoc., France; 3a,b, RV int., ext., $\times 0.7$ (Deshayes, 1825).

P. (Zorrita) OLSSON, 1932 [16] [**P. (Z.) petersoni*; OD]. Transversely subquadrangular, with deep posterior area. Unequal faint internal ribs. Remnants of 3b and 4b present. *Eoc.-Mio.*, Peru-Jamaica-Eu.—FIG. E9,10. **P. (Z.) petersoni*; LV int., $\times 0.9$ (Chavan, n).

Pterolucina CHAVAN, 1942 [2] [**Lucina coeloprocta* COSSMANN, 1887; OD]. Medium-sized to large, rounded trigonal, relatively compressed, with




FIG. E11. Lucinidae (Milthinae) (p. N504-N505).

locally lamellose growths and well-marked areas; lunule depressed; ligament semi-internal on relatively short nymph. Cardinal *3b* weak, oblique, thin, *2* and *4b* scarcely divergent, *A1* faint, *P1* lacking; anterior scars distant from pallial line, subparallel to it; shell margin internally smooth. *U.Cret.-Mio.*, *?Plio.*, Eu.-Asia-N.Afr.—FIG. E9,4. **P. coeloprocta* (COSSMANN), U. Eoc. (Barton.), France; *Aa-c*, RV ext., LV int., RV int., $\times 1$ (160). *Saxolucina* STEWART, 1930 [**Lucina saxorum* LAMARCK, 1806 (1808); OD]. Medium-sized to large, discoidal to subtrigonal, flattened, anteriorly somewhat acuminate, posteriorly truncate; sculpture consisting of lamellose concentric ribs or outgrowths; areas narrow, anterior marked by lines of punctuations, posterior obsolete; lunule projecting backward. Anterior scars very narrow, long; shell margin internally smooth. *?U.Cret.*, *Paleoc.-Plio.*, *?Rec.*, Eu.-Afr.-Am.-Eu.

S. (Saxolucina) [5]. Medium-sized; lunule covering part of teeth, but not really excavated. Cardinals *3b* and *2* partly bifid, *4b* present, *3a* and laterals more or less obsolete; nymph curved; shell commonly pustulose internally. *Eoc.*, Eu.; *?Rec.*, Afr.—FIG. E9,8. **S. (S.) saxorum* (LAMARCK), *Eoc.*, France; *8a,b*, LV int., RV int., $\times 1.7$ (98).

S. (Plastomiltha) STEWART, 1930 [6] [**Cyclas clai-bornensis* CONRAD, 1865; OD] [= *Armi-miltha* OLSSON & HARBISON, 1953 (type, *Lucina disciformis* HEILPRIN, 1886; OD)]. Large, lamellose, with depressed lunule. Straight nymph. Cardinals *3b* and *2* strongly bifid, *3a* well marked, shell internally punctate. *?U.Cret.*, *Paleoc.-Plio.*, C.Am.-N.Am.-Eu.—FIG. E11,2. **S. (P.) clai-bornensis* (CONRAD), *Eoc.*, USA (Ala.); *2a,b*, RV int., LV int., $\times 1$ (98).

?Trinitasia MAURY, 1928 [11] [**Thyasira sancti-*

andreae MAURY, 1925; OD]. Trigonal, subequilateral, convex, ventral margin rounded; surface smooth, posterior area marked by angulation. Cardinals 3 (incompletely known), strong, 1 on RV, 2 on LV; scars apparently not detached, linguiform, and curved. *Mio.*, Trinidad.—FIG. E9,11. **T. sanctiandreae* (MAURY); LV int., $\times 0.75$ (Chavan, n).

Subfamily DIVARICELLINAE Glibert, 1967

Shell convex, rounded, with divaricate or undulating external sculpture. *L.Eoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Divalinga*.—2. *Viaderella*.—3. *Stchepinskya*.—4. *Paralucinella*.—5. *Lucinella*.—6. *Divalucina*.—7. *Boeuvia*.—8. *Divaricella*.—9. *Egracina*.—10. *Pompholigina*.—11. *Eodivaricella*.

Divaricella VON MARTENS, 1880 [**D. angulifera*; OD]. Rounded, with small lunule. Hinge with cardinals and incomplete or obsolete laterals; anterior scars flexuous, narrowly elongate; shell margin internally smooth but incised or exceeded by terminations of ribs. *Plio.-Rec.*, Afr.-Ind.O.-Asia-Australia-C.Am.

D. (Divaricella) [8]. Solid, divaricated by angular ribs. Lateral teeth, *AII*, *PII*, and *PIII* obsolete; terminations of ribs exceeding margin. *Plio.-Rec.*, Asia-Australia-E.Afr.—FIG. E12,1; E10,1. **D. (D.) angulifera*, *Rec.*, Ind.O.(Mauritius); E12, 1a,b, RV hinge, LV int., $\times 2$ (107); E10,1, RV ext., $\times 1$ (von Martens).

D. (Egracina) CHAVAN, 1951 [9] [**Tellina dentata* Woods, 1815; OD]. Relatively flattened, divaricated by large flattened ribs, with narrow interspaces. Hinge with *3a*, *AI*, *AII*, *PII*, *PIII* obsolete; terminations of ribs incising margin. *Pleist.-Rec.*, C.Am.-Afr.—FIG. E12,5. *D. (E.) dentata collignoni* CHAVAN, Afr.; 5a-c, LV ext., RV hinge, LV int., $\times 2.5$ (107).

Boeuvia CHAVAN, 1948 [7] [**Lucina pulchella* AGASSIZ, 1845; OD] [= ?*Bourdotia* DALL, 1901 (type, *Lucina bourdoti* COSSMANN, 1882; OD) (*nom. dub.*)]. Not much inflated; divaricated by numerous flattened ribs with narrow interspaces. Incomplete hinge without *AI* and *PI*; *3a*, *4b*, *AIV*, *PII* and *PIV* all obsolete or lacking; ligament partly internal; anterior scars strongly divergent from pallial line; shell margin internally smooth. *L.Eoc.-Oligo.*, N.Afr.-Eu.—FIG. E12,3; E10,5. **B. pulchella* (AGASSIZ), M.Eoc.(Lutet.), France; E12, 3a,b, RV hinge, LV int., $\times 2$ (107); E10,5, RV ext., $\times 1$ (Deshayes).

Divalinga CHAVAN, 1951 [**Lucina quadrisulcata* D'ORBIGNY, 1846; OD]. Orbicular, inflated; divaricated by broad flattened ribs with narrow interspaces; lunule slightly depressed, dissymmetric; ligament external. Hinge with well-developed cardinals and laterals but no *AI* and *PI*; anterior

scars short; shell margin internally denticulate. *L.Eoc.-Rec.*, Eu.-N.Am.-C.Am.-Afr. [= *Cyclas* MÖRCH, 1853 (*non* LAMARCK, 1799) (type, *Lucina quadrisulcata* D'ORBIGNY, 1846; SD CHAVAN, herein).]

D. (Divalinga) [1]. Hinge teeth *3a* and *AIV* feeble; anterior scars slightly divergent from pallial line. *Oligo.-Rec.*, W.Eu.-N.Am.-C.Am.—FIG. E12,9. **D. (D.) quadrisulcata* (D'ORBIGNY), *Rec.*, USA(Fla.); 9a,b, RV hinge, LV int., $\times 2$ (107).

D. (Stchepinskya) CHAVAN, 1951 [3] [**Lucina rigaultiana* DESHAYES, 1858; OD]. Inflated; lunule elongate. Hinge teeth *3a* and *PIV* feeble, anterior and posterior laterals both distant; anterior scars short, acutely diverging from pallial line; shell margin weakly crenulate internally. *L.Eoc.-U.Eoc.*, W.Eu.-N.Afr.—FIG. E12,11. **D. (S.) rigaultiana* (DESHAYES), M.Eoc. (Auvers.), France; 11a,b, RV hinge, LV int., $\times 3$ (107).

D. (Viaderella) CHAVAN, 1951 [2] [**Divaricella perparvula* DALL, 1901 (= *Lucina pisum* PHILIPPI, 1857; *non* SOWERBY, 1836; *nec* D'ORBIGNY, 1841; *nec* REEVE, 1850)]. Inflated; sculpture obsolete on angle of divergence, posterior lamellae distant. Hinge teeth *PIV* feeble, *PI* lacking (also *AI*); anterior scars short, acutely pointed and diverging. *Mio.*, W.Eu.; *Rec.*, N.Am.-S.Afr.-W. Eu.—FIG. E12,10. **D. (V.) perparvula* (DALL), USA(Calif.); 10a,b, RV hinge, LV int., $\times 2$ (107).

Divalucina IREDALE, 1936 [6] [**Lucina (Cyclas) cumingi* ADAMS & ANGAS, 1863; OD]. Large, solid, moderately inflated; surface divaricated by broad flattened ribs with narrow interspaces; lunule narrowly elongate. Hinge with large cardinals (2 bifid) and *AII*, *AIII* (*AI* lacking, *AIV* feeble), posterior laterals obsolete; anterior scars narrowly elongate, not divergent from pallial line; shell margin smooth internally. *Oligo.-Rec.*, Asia (Ceylon-Japan)-Australia-N.Z.—FIG. E12,4. **D. cumingi* (ADAMS & ANGAS), *Rec.*, N.Z.; 4a,b, RV hinge, LV int., $\times 1.5$ (107).

Lucinella DI MONTEROSATO, 1883 [5] [**Tellina divaricata* LINNÉ, 1758 (= *Lucina commutata* PHILIPPI, 1836); OD]. Relatively small, solid, with posterior area, sculpture undulated. Hinge with cardinals and laterals, *3a* and *PII* feeble, *AI* and *PI* lacking; ligament deeply internal; anterior scars not much divergent from pallial line; shell margin crenulate internally. *Mio.-Rec.*, W.Eu.-S. Eu.-Ind.O.(Mauritius).—FIG. E12,7. **L. divaricata* (LINNÉ), *Rec.*, France; 7a,b, RV hinge, LV int., $\times 4$ (107).

Paralucinella CHAVAN, 1951 [4] [**Lucina undulata* LAMARCK, 1806; OD]. Relatively small; sculpture undulated. Hinge narrow, dentition as in *Stchepinskya*, but *3a* lacking; ligament elongated; anterior




FIG. E12. Lucinidae (Divaricellinae) (p. N506, N508).




FIG. E13. Thyasiridae (p. N508).

scars short, rounded; shell margin crenulated internally. *Oligo.*, W.Eu.—FIG. E12,8. *P. undulata* (LAMARCK), Stamp., France; 8a,b, RV hinge, LV int., $\times 4.5$ (107).

Pompholigina DALL, 1901 [**Lucina gibba* GRAY, 1825; OD]. Tumid, with subspiral beaks; surface divaricated by elevated rounded ribs with wide interspaces. Hinge apparently cyclodont, with cardinals and several laterals. *M.Eoc.-Rec.*, W.Afr.

P. (Pompholigina) [10]. Hinge with 3a and 4b well developed; anterior scars short, pointed, divergent from pallial line; shell margin smooth internally. *Rec.*, W.Afr.—FIG. E12,2; E10,4. **P. (P.) gibba* (GRAY); E12,2a,b, RV hinge, LV int., $\times 2$ (107); E10,4, LV ext., $\times 1$ (Reeve, 1850).

P. (Eodivaricella) CHAVAN, 1951 [11] [**Divaricella oppenheimi* NEWTON, 1923; OD]. Lunule excavated; nepionic sculpture concentric, then divaricate. Anterior scars narrow; middle part of internal shell margin undulated. *M.Eoc.*, W.Afr. (Nigeria).—FIG. E12,6. **P. (E.) oppenheimi* (NEWTON); LV int., $\times 3.3$ (107).

LUCINIDAE GENERA DUBIA

Austriella TENISON-WOODS, 1881 [**A. sordida*; OD]. Like *Eamesiella* but with an inconspicuous arcuate tooth and a falciform scar. [May be an eroded form, possibly an edentulous unguinid.] *Rec.*, Australia.

Elathia ISSEL, 1869 [**E. arconatii*; OD]. *Rec.*, Red Sea [See *Gibbolucina*.]—FIG. E10,3. **E. arconatii*; 3a,b, LV ext., RV int., $\times 1$ (Issel).

?**Freila** DE GREGORIO, 1930.

Jagonella SELLI, 1944 [**J. gortanii*; OD]. (*Nom. nud.*)

Levimyrtaea OLSSON, 1965 [**Myrtaea (L.) inconspicua*; OD]. Possibly equivalent of *Gonimyrtaea* or *Myrteopsis*. *Mio.*, Ecuador.

Lucinigenus RENIER, 1894 [Rejected, ICZN].

Family THYASIRIDAE Dall, 1901

[=Cryptodontidae DALL, 1895]

Trigonal to subquadrangular or obliquely rounded, thin, with dorsal anterior depression and more or less well-marked posterior area; beaks small, acute, prosogyrate; ligament and resilium juxtaposed, united, former inframarginal, latter deeply sunken in ill-defined depression or socket; no nymph. Surface concentrically ribbed or smooth. Hinge with protruding right lunular edge forming one or two small ill-defined tuberosities, with intermediate left and corresponding sockets; muscle scars elongate, superficial. *M.Trias.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Thyasira*.—2. *Conchocele*.—3. *Philis*.—4. *Maoriithyas*.—5. *Tauraxinus*.—6. *Mendicula*.—7. *Axinulus*.—8. *Parathyasira*.—9. *Axinopsida*.—10. *Adontorhina*.—11. *Leptaxinus*.—12. *Genaxinus*.—13. *Storthodon*.

Thyasira LEACH in LAMARCK, 1818 [**Amphidesma flexuosa* LAMARCK, 1818 (= **Tellina flexuosa* MONTAGU, 1803); OD] [= *Axinus* SOWERBY, 1821 (type, *A. angulatus*; OD); *Cryptodon* TURTON, 1822; *Bequania* LEACH in BROWN, 1827; *Ptychina* PHILIPPI, 1836 (obj.); *Prothyasira* IREDALE, 1930 (type, *P. peroniana*; OD) (obj.)]. Obliquely trigonal, with sharp dorsal folds and well-marked posterior area. Concentrically sculptured. *Cret.-Rec.*, Eu.-N.Am.-Pac.-Australia.

T. (Thyasira) [1]. Small, oblique, with lunular margin only slightly curved, protruding in right valve, followed by resilium in moderately short, broadly trigonal depression. A minute pseudo-cardinal on left valve. *Cret.-Rec.*, Eu.-Pac.-Australia.—FIG. E13,1; E14,8. **T. (T.) flexuosa* (MONTAGU), *Rec.*, *Medit.*; E13,1, RV int., $\times 1$ (305); E14,8, RV hinge, $\times 1$ (513).

T. (Conchocele) GABB, 1866 [2] [**C. disjuncta*; OD]. Very oblique, enlarged backward; lunular margin nearly straight, followed by resilium in long narrow depression; anterior muscle scar broad. *Oligo.-Rec.*, N.Am.-N.Pac.—FIG. E14,4.

**T. (C.) disjuncta* (GABB), Rec., USA(Calif.); 4a,b, RV ext., int., $\times 0.7$ (Keen, 1939).
T. (Philis) P. FISCHER, 1861 [3] [**P. Cumingi*; OD]. Small, somewhat oblique; deeply excavated lunule in spoon-shaped socket under beak, its edge U-shaped in front of ill-defined depression. Rec., Australasia-Japan.—FIG. E14,6. **T. (P.) cumingi*, Moluccas; LV int., enl. (Chavan, n).

Adontorhina BERRY, 1947 [10] [**A. cyclia*; OD]. Small, resembling *Axinopsida* but hinge teeth replaced by indefinite ridges and denticles on somewhat reflected lunular and escutcheonal margins. Anterior pseudocardinals tuberculiform. *Pleist.-Rec.*, W.N.Am.—FIG. E14,11. **A. cyclia*, Pleist., USA(Calif.); 11a,b, LV int., ext., $\times 10$ (Berry, 1917).




FIG. E14. Thyasiridae (p. N508-N511).

- laterals. *Pleist.-Rec.*, Australia-N.Z.-Japan.—FIG. E13,3; E14,13. *P. resupina neozelandica* IREDALE, N.Z.; E13,3, LV ext., $\times 5$ (Iredale, 1930); E14, 13a-c, RV ext., int., LV int., $\times 1$ (Fleming, 1950).
- Storhodon** GIEBEL, 1856 [13] [**S. liscaviensis*; OD]. Oblong, gibbous, with well-marked posterior area, beaks pointed, prosogyrous; hinge with pyramidal quadrangular tooth under beaks and nearly similar one beneath it; marginal ligament in flattened area. *M.Trias.*, Eu.—FIG. E14, 1. **S. liscaviensis*, Ger.; RV int. (Chavan, n).
- Tauraxinus** SACCO, 1901 [**T. miorugosus*; OD]. Flabelliform, with quite obsolete areas, lunule depressed or sunken, ligamentary groove somewhat narrow and elongate; surface with concentric wavy sculpture. *Mio.-Rec.*, Eu.-S.Pac.
- T. (Tauraxinus)** [5]. Anterior end attenuated, produced; beaks small; lunule long, depressed. *Mio.*, Eu.—FIG. E14,9. **T. (T.) miorugosus*, Italy; 9a,b, RV ext., int., enl. (Chavan, n).
- T. (Mendicula)** IREDALE, 1924 [6] [**M. memorata* (= *Lucina induta* HEDLEY, 1907, non STOLICZKA, 1887); OD]. Anterior end rounded, beaks prominent, prosogyrate; lunule not defined but corresponding anterior margin projecting backward behind beaks. *Rec.*, S.Pac.-Australia.—FIG. E14,5. **T. (M.) memorata* (IREDALE), Australia; 5a,b, RV ext., hinge, enl. (Chavan, n).
- Family MACTROMYIDAE Cox, 1929**
[=Mactromyidae Cox, 1935]
- Globose, outline trigonal to elliptical, commonly elongate transversely; beaks more or less prominent, prosogyrous; lunule ill-defined, ligament in broadened, distinctly marginal groove with somewhat sunken resilium; surface with predominant concentric sculpture. Hinge with primitive laminae in cyclodont pattern; anterior adductor scars ovate, not digitate; some shells with shallow pallial sinus. *Dev.-Rec.*
- Arrangement of generic taxa by CHAVAN.—1. *Mactromya*.—2. *Clisocolus*.—3. *Macromyopsis*.—4. *Mactromyella*.—5. *Linetia*.—6. *Thetis*.—7. *Sexta*.—8. *Bathycorbis*.—9. *Plesiocyprinella*.—10. *Palaeolucina*.—11. *Montanaria*.—12. *Paracyclas*. [Insert above, 1a. *Unicardium*; 6a. *Cordiula*.]
- Mactromya** AGASSIZ, 1843 [1] [**M. rugosa* (= **Lutraria concentrica* MÜNSTER in GOLDFUSS, 1840; SD HERRMANNSEN, 1847)]. Transversely elliptical, nearly equilateral, with strong subequidistant concentric ribs; RV hinge with anterior thickening and shallow socket. *Jur.(Bathon.-Portland.)*, Eu.-E.Afr.—FIG. E15,11. **M. concentrica* (MÜNSTER), U.Jur., Ger., LV ext., $\times 1.25$ (Chavan, n, after Goldfuss).
- Bathycorbis** IREDALE, 1930 [8] [**Chione despecta* HEDLEY, 1904; OD]. Transversely trigonal, small, nearly equilateral; surface with strong concentric ribs, lacinate, intercalated striae; LV hinge with 2 superposed tuberculiform teeth and posterior thickening in front of ligament, RV hinge with single rounded tooth and laminar processes in inverted V above it. *Rec.*, Australia.—FIG. E15,8. **B. despecta* (HEDLEY); LV int., $\times 12$ (119).
- Clisocolus** GABB, 1869 [2] [**Loripes dubia* GABB, 1864; OD]. Trigonal to rounded, somewhat inequilateral, beaks high, with lamellar striae; resilium sunken; hinge with ill-defined virguliform cardinal and anterior lateral in each valve. *Cret.*, W.N.Am.-Eu.—FIG. E15,1; E16,1. **C. dubius* (GABB), Chico Gr., USA(Calif.); E15,1a,b, RV and LV hinges (119); E16,1a,b, RV and LV hinges, $\times 1.5$; E16,1c, LV ext., $\times 1.5$ (333).
- Cordiula** MEYER, 1887 [6a] [**Unicardium? eoecense*; OD]. Similar to *Thetis* in outline and sculpture, but without oblique internal rib; three faint medioposterior pallial sinuosities. One left tubercular cardinal and obsolete laterals. Anterior scar elliptical, posterior elliptical-cordate. *Oligo.*, USA (Miss.).
- Linetia** CHAVAN, 1959 [5] [**Mactromya Caumonti* CHAVAN, 1959 (ex AGASSIZ, MS, 1845) (= *Mactromya calliope* ROLLIER, 1913, non THÉVENIN, 1909, ex D'ORBIGNY, 1850); OD]. Large, rounded ovate, very inequilateral, with irregular concentric ribbing and radial striation laterally; hinge with *AIII-3* in inverted V with triangular 2 and distinct 4b below. *Jur.(Aalen.-Callov.)*, Eu.—FIG. E15,9. **L. caumonti* (AGASSIZ), Bajoc., W.France; 9a-d, LV ext., int., LV and RV hinges, $\times 0.8$ (119).
- Mactromyopsis** CHAVAN, 1959 [**Unicardium hemirhytideum* COSSMANN, 1905; OD]. Subquadrate to subcirculate, rather thick, with strong concentric ribbing; hinge with single cardinal in each valve attached to its anterior lamina, elevated posterior muscle scar. *Jur.(Charmouth.-Callov.)*, Eu.
- M. (Mactromyopsis)** [3]. Subquadrate; RV hinge with well-developed cardinal 3. *Jur.(Bajoc.-Callov.)*, Eu.—FIG. E15,7. **M. (M.) hemirhytidea* (COSSMANN), Bajoc., W.France; LV hinge, $\times 1.25$ (119).
- M. (Mactromyella)** CHAVAN, 1959 [4] [**Unicardium inflatum* THÉVENIN, 1909 (ex D'ORBIGNY, 1850); OD]. Subcircular, with rather regular ribbing; RV hinge with evanescent cardinal 3, overhanging well-developed *A1*. *Jur.(Charmouth.-Bajoc.)*, Eu.—FIG. E15,12. **M. (M.) inflata* (THÉVENIN), Bajoc., W.France; RV int., $\times 1.25$ (119).
- Montanaria** SPRIESTERSBACH, 1909 [11] [**M. ovata* (= **Pleurophorus devonicus* BEUSHAUSEN, 1884); SD HAFFER, 1959]. Rounded oval to elongate, very inequilateral, beaks small, anterior, surface




FIG. E16. Mactromyidae (p. N511-N512).

with fine concentric growth lines only; hinge with *AIII*, *3a-3b/4a*, *2*, (*4b*); *2* and *3* grooved to bifid, some shells with *AIII* and *4a* weak or lacking; anterior scars ovate. *Dev.*, Eu.—FIG. E15,10. **M. devonica* (BEUSHAUSEN), Remscheid beds, Ger.; *10a*, RV hinge (Beushausen); *10b,c*, RV and LV hinges (Chavan, n).

?*Palaeolucina* CHAO, 1927 [10] [**P. carbonaria*; OD]. Mostly elliptical, with somewhat low beaks, distinct ovate lunule, and long narrow escutcheon; surface with lamellose concentric ribs; hinge with 3 teeth in LV, median one bifid; *2* oblique teeth and marginal anterior lamella in RV; muscle scars unknown. *Carb.*, China.—FIG. E15,4. **P. carbonaria*; *4a,b*, LV and RV hinges, enl. (Chavan, n, after Chao).

Paracyclas HALL, 1843 [12] [**P. elliptica* (= *Lucina proavia* GOLDFUSS, 1840); OD]. Ovate, beaks low, surface with concentric ribs; RV hinge apparently with *2* oblique superposed rounded tubercles separated by sockets, superior one smaller than inferior, flat area in front of them and possibly linear anterior lateral; broad and long ligament depression; anterior muscle scars elliptical. *Dev.*, N. Am. - Eu. — FIG. E15,5. *P. marginata*

(MAURER), Ger.; *5a,b*, LV hinges, enl. (Chavan after Maurer).

Plesiocyprinella HOLDHAUS, 1918 [9] [**P. carinata*; OD]. Cordate, rounded, beaks prominent; RV hinge with *AIII*, *3a*, *3b* hooked, LV hinge with large, low, subtrigonal *2* and thinner, oblique *4b*. *Perm.*, S.Am.—FIG. E15,3; E16,2. **P. carinata*, Brazil; E15,3*a,b*, LV and RV hinges, $\times 0.8$ (119); E16,2*a-e*, LV ext., ant., RV ant., hinge, LV hinge, all $\times 1.5$ (Holdhaus).

Sexta STEPHENSON, 1954 [7] [**S. navicula*; OD]. Rounded, convex, relatively low beaks, no lunule, ligament external; surface nearly smooth; hinge with anterior laterals, median depression, faint posterior laterals, RV with minute cardinal; adductor scars small, subequal. *U.Cret.(Cenoman.)*, N.Am.—FIG. E15,6. **S. navicula*, USA(Tex.); *6a,b*, RV and LV hinges, $\times 2$ (119).

Thetis J. DE C. SOWERBY, 1826 [6] [**T. major*; M (= *Corbula laevigata* SOWERBY, 1818, non *Thetis* OKEN, 1815, inval., nec C. B. ADAMS, 1845, nec H. ADAMS & A. ADAMS, 1856)] [= *Thetironia* STOLICZKA, 1870 (obj.); *Fimbriella* STOLICZKA, 1871 (obj.)]. Subcircular, tumid, with prominent nearly orthogyrus beaks; surface with concentric

and laterally radial sculpture; ligament thin; hinge with *AIV* continued above and behind tubercular 2 as hooked tooth, RV with 2 superposed tuberculiform cardinals. Oblique internal rib resembling pallial sinus. *Cret. (Alb.-Cenoman.)*, Eu.-N.Am. —FIG. E15,2. **T. laevigata* (SOWERBY), L.Cret., Eng.; 2a,b, RV and LV hinges, $\times 1.25$ (119).

[The American species, known as *Thetiopsis* MEEK, shows a venerid hinge.]

Unicardium D'ORBIGNY, 1850 (1849) [1a] [**Corbula cardioides* PHILLIPS, 1829; SD STOLICZKA, 1871]. Like *Mactromya* but more tumid, rounded; broadly inflated prosogyrous beaks; irregular lamellose concentric ribbing. Hinge with right faint cardinal tubercle; long straight nymph. *Jur.*, W.Eu.

Family FIMBRIIDAE Nicol, 1950

[=Corbidae DALL, 1895]

Transversely elliptical or ovate, beaks rounded; lunule and escutcheon generally well marked; surface bearing primary radial ribs superposed by smooth or concentrically sculptured external layer; marginal ligament on nymph. Hinge with massive, trigonal or tubercular, generally entire medial teeth in more or less cycloodont pattern, anterior laterals commonly and posterior laterals generally developed; muscle scars ovate to reniform, short, without digitation. *Carb.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Fimbria*.—2. *Scaldia*.—3. *Schafhaeutlia*.—4. *Sphaeriola*.—5. *Haastina*.—6. *Cyclopellatia*.—7. *Sphaera*.—8. *Mutiella*.—9. *Parvicorbis*.

Fimbria MEËRLE VON MÜHLFELD, 1811 [1] [**F. magna* (= *Venus fimbriata* LINNÉ, 1758); OD] [non *Fimbria* BOHADSCH, 1761 (invalid)] [= *Corbis* CUVIER, 1817 (obj.); *Idothea* SCHUMACHER, 1817 (non FABRICIUS, 1796) (obj.)]. Transversely elliptical, subequilateral, thick, beaks prosogyrous; lunule small, lanceolate, ligament external but partly sunken; surface with strong reticulate sculpture, concentric ribs medially dominant; hinge with 2 cardinals in each valve, 3b grooved, anterior laterals near, posterior laterals remote; interior with very small pallial sinus and crenulate margin. *M.Jur. (Bathon.)-Rec.*, Eu.-N.Am.-Pac.-India-Australia.—FIG. E17,1. **F. fimbriata* (LINNÉ), *Rec.*, Eu.; 1a-c, RV ext., int., LV int., $\times 1$ (674). *Cyclopellatia* COSSMANN, 1907 [6] [**C. acrodontia*; OD]. Rounded, nearly equilateral, low prosogyrous beaks; surface with close-spaced radial ribs crossed by concentric striations which predominate posteriorly; hinge broad, thick, LV with marginal anterior approximate to upper lateral (*AIV*), strong trigonal 2, oblique 4b, strong *PII* not very remote; ligament marginal. *L.Jur. (Barrem.)*, Eu.—FIG. E18,4. **C. acrodontia*, S.




FIG. E17. Fimbriidae (p. N513).

France; 4a,b, LV ext., $\times 1.25$, hinge, $\times 2.5$ (Chavan, n, after Cossmann).

Haastina MARWICK, 1953 [5] [**Heminajas? haastiana* WILCKENS, 1918; OD]. Transversely elliptical, slightly inequilateral, beaks high, prosogyrous; surface with concentric striation; hinge with 3 diverging cardinals, median bifid, posterior weak on RV, posterior laterals strong; muscle scars ovate. *U.Jur. (Oxford.)*, N.Z.—FIG. E18,3. **H. haastina* (WILCKENS); RV int., $\times 1.5$ (Chavan, n, after Marwick).

Mutiella STOLICZKA, 1871 [8] [**Corbis rotundata* D'ORBIGNY, 1843; OD]. Globose, high, ovate, beaks broad, nearly orthogyrous, with long marginal ligament; surface finely reticulate; LV hinge with *AIV* marginal, elongate and small remote tubercular *All*, 2 deeply bifid, 4b long and stout. *U.Cret.*, Eu.-S.Asia (India).—FIG. E18,5. **M. rotundata* (D'ORBIGNY), Senon., France; LV hinge, $\times 1.5$ (Chavan, n, after d'Orbigny).

Parvicorbis COSSMANN, 1892 [9] [*pro Bernayia* COSSMANN, 1887 (non *Bernaya* JOUSSEAUME, 1884)] [**Bernayia subarata* COSSMANN, 1887; SD CHAVAN,




FIG. E18. Fimbriidae (p. N513-N514).

herein]. Small, transversely elliptical, slightly gaping at both ends; lunule narrow, escutcheon well impressed, nymph short; surface with concentric to reticulate sculpture; hinge with small pointed cardinals, bilobate 2, remote anterior and posterior laterals; inner shell margin smooth. *Eoc.*, Eu.—FIG. E18,6a. **P. subarata* (COSSMANN), France (Paris Basin); 6a, LV int., $\times 4.5$ (Chavan, n, after Cossmann).—FIG. E18,6b. *P. goodallina* (COSSMANN), France (Paris Basin); 6b, RV hinge, $\times 8$ (Chavan, n, after Cossmann).

Scaldia DE RYCKHOLT, 1847 [2] [**S. lambotteana*; SD STOLICZKA, 1871]. Rounded, short, slightly inequilateral, somewhat prosogyrous prominent beaks; escutcheon narrow, nymph elongate; surface concentrically striate; hinge with 2 cardinals in each valve, 2 and 3b strong, 3a laminar, 4b weak; inner shell margin smooth. *Carb.*, Eu.—FIG. E18,8. **S. lambotteana*, Belg.; 8a,b, LV and RV hinges, $\times 1.25$ (Chavan, n).

Schafhaeutlia COSSMANN, 1897 [3] [*pro Gonodon* SCHAFFHAEUTL, 1863 (non HELD, 1837)] [**Gonodon ovatum* SCHAFFHAEUTL, 1863; OD]. Broadly elliptical, short, very thick, slightly inequilateral; surface with concentric sculpture; hinge with cyclodont conical cardinals and irregular anterior lateral; anterior muscle scar small. *U.Trias.*, Eu.-S.Am.—FIG. E18,7; E19,1. *S. mellingi* (HAUER); E18,7a,b, RV and LV hinges, $\times 1$ (Chavan, n, after Hauer); E19,1a-c, LV int., RV int., both valves ant., $\times 1$ (Schafhaeutl, 1863). [= *Gonodus* MUSKETOV, 1913 (missp.).]

Sphaera SOWERBY, 1822 [7] [**S. corrugata*; SD STOLICZKA, 1871] [= *Palaeocorbis* CONRAD, 1869 (obj.)]. Rounded, inflated, solid, subequilateral, with inflated nearly orthogyrous beaks; surface with corrugated concentric ribs; hinge strong, cyclodont, RV with strong anterior lateral, subhorizontal anterior cardinal, and arcuate posterior lateral, LV with anterior lateral fused with anterior cardinal and ill-defined posterior lateral, each valve with 2 remote posterior laterals. *L. Cret.*, Eu.—FIG. E18,1. **S. corrugata*, Neocom., France; 1a,b, RV and LV hinges, $\times 2$ (Chavan, n, after d'Orbigny & Woods).

Sphaeriola STOLICZKA, 1871 [4] [**S. madridi* (= *Cardium madridi* D'ARCHIAC, 1843); OD]. Globose, rounded, subequilateral, with prominent nearly orthogyrous rounded beaks, ligament short, broad; hinge with 2 teeth in each valve, anterior one (AIV, AIII -3a) subhorizontal and elongate, posterior one (2, 3b) trigonal, arcuate, stout, inner shell margin with denticles not corresponding to apparent ribs; sculpture of concentric sulci on radial internal ribs. *Jur. (Domer.-Bathon.)*, Eu.-N.Z.-Japan.—FIG. E18,2. **S. madridi* (D'ARCHIAC), Bathon., Eng.; 2a,b, RV and LV hinges, $\times 1.25$ (Chavan, n, after Morris & Lycett).

Family UNGULINIDAE

Adams & Adams, 1857

[= *Diplodontidae* DALL, 1895]

Outline subtrigonal to oblong, rounded or obliquely ovate, beaks generally low, dorsal angulations weak or absent, ligament and resilium marginal; surface smooth, punctate, or with fine concentric sculpture; hinge with two cardinals, medial one bifid, and incomplete or evanescent laterals; muscle scars irregular, anterior scar elongate and not detached from pallial line, posterior scar large. *U.Cret.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Ungulina*.—2. *Numella*.—3. *Felania*.—4. *Brachymeris*.—5. *Microstagon*.—6. *Diplodonta*.—7. *Zemysina*.—8. *Felaniella*.—9. *Zemysia*.




FIG. E19. Fimbriidae (p. N514).

—10. *Bruetia*.—11. *Phlyctiderma*.—12. *Timothyus*.—13. *Cycladicama*.—14. *Toralimysia*.

Ungulina ROISSY, 1805 [1] [**U. rubra* (= *Tellina cuneata* SPENGLER, 1782); SD GRAY, 1847] [= *Unguline* DAUDIN in BOSCH, 1801 (vernac.); *Clotho* BASTEROT, 1825 (non WALCKENAER, 1805, nec FAUJAS, 1808) (type, *C. unguiformis*; OD)]. Oblong, thick, irregular; lunule lacking, external ligament extended backward, with groove adjacent to shorter socket occupied by broad resilium which partly covers nymph and obliterates 4b; large oblong subequal muscle scars. *Oligo.*(*Aquitain.*)-*Rec.*, Eu.-W.Afr.—FIG. E20,8. **U. cuneata* (SPENGLER), *Rec.*, Senegal; 8a,b, LV int., RV hinge, enl. (121).

Brachymeris CONRAD, 1875 [4] [**B. alta*; OD] [= *Arene* CONRAD, 1875 (non ADAMS & ADAMS, 1854)]. Obliquely oblong, nearly orthogyrous beaks; lunule lacking, narrow marginal ligament and resilium in short groove; hinge with generally bifid 2 and 3b, and oblique 3a and 4b, anterior laterals moderately developed; posterior ones weak; anterior muscle scar very broad. *U.Cret.*, N.Am.—FIG. E20,7. **B. alta*, USA(N.Car.); 7a,b, RV int., LV hinge, $\times 3$ (Chavan, 1960).

Bruetia CHAVAN, 1962 [10] [**Lucina radians* MELLEVILLE, 1843 (non CONRAD, 1841) (= *L. subadians* D'ORBIGNY, 1850); OD]. Subtrigonal to subquadrate, somewhat thick; narrow resilium and flat nymph in front and resting on short ligament; hinge with 2 bifid cardinals in each valve, anterior laterals and AIV-4a developed marginally; internal radial lines generally well marked; anterior muscle scar falciform, posterior scar shorter and ovate. *Paleoc.-Mio.*, Eu.-C.Am.—FIG. E20, 5. *B. subadians* (D'ORBIGNY), *L.Eoc.*, France (Paris Basin); 5a,b, LV int., RV hinge, $\times 3$ (Chavan, 1960).

Cycladicama VALENCIENNES in ROUSSEAU, 1854 [**C. luciniformis* (= *Cyrenoida oblonga* HANLEY, 1844

(*nom. nud.*), 1856); OD] [= *Joannisiella* DALL, 1895 (pro *Joannisia* DALL, 1895, non MONTEROSATO, 1884, nec KIEFER, 1894) (obj.)]. Transversely sub-ovate, attenuate forward, somewhat angular and truncate rearward, with external flattened area defined by medioposterior weak angulation; ligament in groove adjacent to shorter resilium in broader socket; surface concentrically striate; hinge with 2 cardinals in each valve and single more or less distinct anterior lateral; anterior muscle scar irregularly extended and doubles, elongate. *U.Eoc.-Rec.*, Pac.-W.Afr.-Japan-?Eu.

C. (Cycladicama) [13]. Broad, normal in convexity; hinge with laminar anterior laterals. *U. Eoc.-Rec.*, Pac.-W.Afr.-Japan-?Eu.—FIG. E20, 11. **C. (C.) oblonga* (HANLEY), *Rec.*, Philip. Is.; 11a,b, RV int., LV hinge, $\times 2$ (Chavan, 1960).

C. (Toralimysia) IREDALE, 1936 [14] [**T. excentrica* (= *Joannisiella sphaericula* HEDLEY, 1906, non *Cyrenella sphaericula* DESHAYES, 1854); OD]. Strongly convex; hinge with broad cardinals and obsolete laterals. *Rec.*, Australia-?Eu.—FIG. E20,3. **C. (T.) excentrica* IREDALE; 3a,b, RV and LV hinges, enl. (Chavan, 1960).

Diplodonta BRONN, 1831 [**Venus lupinus* BROCCHI, 1814 (non LINNÉ, 1858) (= **Tellina rotundata* MONTAGU, 1803) (= *D. rotundata* var. *aequilateralis* CERULLI, 1909); SD HERRMANNSEN, 1846] [= ?*Taras* RISSO, 1826 (type, *T. antiquatus*; OD) (*nom. dub.*); *Mysia* LEACH in BRONN, 1827 (non LAMARCK, 1818) (obj.); *Glocomene* LEACH in GRAY, 1852 (type, *Tellina rotundata* MONTAGU, 1803; SD CHAVAN, 1962); ?*Mittrea* GRAY, 1864 (type, *Diplodonta brasiliensis* MITTRE, 1850)]. Suborbicular, convex, inequilateral, beaks prosogyrous; ligament on medium-sized flattened nymph with narrow resilium on its anterior extremity; hinge with 2 well-developed oblique cardinals in each valve and weak 5b; anterior muscle scars sinuate, narrower than posterior scars. *Paleoc.-Rec.*, Eu.-N.Am.-Pac.-W.Afr.

D. (Diplodonta) [6]. Moderately convex; hinge with somewhat narrow short teeth; anterior muscle scars not elongate. *Paleoc.-Rec.*, Eu.-N.Am.-Pac.—FIG. E20,10. **D. (D.) rotundata* (MONTAGU), *Rec.*, Medit.; 10a,b, LV int., RV hinge, $\times 3$ (Chavan, 1960).

D. (Zemysina) FINLAY, 1926 [1927] [7] [**Zemysia (Z.) globus* FINLAY, 1927; OD]. Very convex; hinge with longer and more bifid teeth and more elongate muscle scar than in *D. (Diplodonta)*. *Eoc.-Rec.*, Eu.-N.Am.-Pac.—FIG. E21, 3. *D. (Z.) walli* (WOODRING), *Mio.*, Jamaica; 3a,b, RV ext., int., $\times 4$ (1005).

Felania RECLUZ, 1851 [3] [**Venus diaphana* GMELIN, 1790; OD]. Sublenticular, thin, medium-sized; lunule small, external ligament adjacent to long resilium lying in groove; hinge with 2 cardi-

nals and single anterior lateral in each valve; anterior muscle scars less elongate than posterior scars. *Rec.*, W.Afr.—FIG. E20,2. **F. diaphana* (GMELIN), Senegal; 2a,b, LV int., RV hinge, enl. (Chavan, 1960).

Felaniella DALL, 1899 [**Mysia (Felania) usta* GOULD, 1861; OD]. Subquadrangular to rounded,

inequilateral, posteromedially enlarged and flattened, rather thin; ligament in marginal groove, small resilium in elliptical socket at top of nymph; hinge with 2 subvertical or oblique cardinals in each valve; anterior muscle scars somewhat irregular, slightly narrower than posterior scars, pallial line doubled. *U.Cret.-Rec.*, N.Am.-W.Pac.-Eu.-Afr.




FIG. E20. Ungulinidae (p. N515-N517).

F. (Felaniella) [8]. Ligament almost external, resilium restricted; hinge with somewhat oblique teeth; muscle scars rather narrow. *Paleoc.-Rec.*, N.Am.-Pac.-Eu.—FIG. E20,1. **F. (F.) usta* (GOULD), *Rec.*, Japan; 1a,b, LV int., RV hinge, $\times 2$ (Chavan, 1960).

F. (Zemysia) FINLAY, 1926 [1927] [**Lucina zelandica* GRAY, 1835; OD]. Broader and more rounded than *F. (Felaniella)*; ligament somewhat sunken, resilium extended, partly covering long nymph; hinge with nearly vertical teeth; muscle scars relatively broad. *U.Cret.-Rec.*, S.Pac.-S.Afr.-N.Am.-Eu.—FIG. E21,1. *F. (Z.) acclinis* CONRAD, *Mio.*, USA(Md.); 1a,b, RV int., LV hinge, enl. (121).

Microstagon COSSMANN, 1896 (1899) [5] [*pro Goodallia* DESHAYES, 1860 (*non* TURTON, 1822)] [**Goodallia herouvalensis* DESHAYES, 1860; SD DALL, 1903]. Small, very obliquely oblong, antero-medially elongated; ligament narrow and long, on nymph adjacent to shorter resilium lying in groove; hinge with thick 2 and oblique 3b, oblique 3a, AIII, 4b, and obsolete PII; anterior muscle scars well marked, irregularly elongate, enlarged. *L. Eoc. - Rec.*, Eu.-C. Am.-N. Am.-Japan. — FIG. E21,2. *M. miliare* (LAMARCK), *Eoc.*, France (Paris Basin); 2a,b, LV, RV int., $\times 2$ (121). [= *Minipisum* YABE, 1961 (type, *M. japonicum*; M) (subj.).]

Numella IREDALE, 1924 [2] [**Mysia (Felania) adamsi* ANGAS, 1867; OD]. Ovately oblong, concentrically striate; ligament next to deep resilium, extended narrowly forward and broadly rearward; hinge with elongate 4b; anterior muscle scars longer and narrower than others. *Mio.-Rec.*, Australia.—FIG. E20,4. **N. adamsi* (ANGAS), *Rec.*; 4a,b, LV int., RV hinge, $\times 3$ (Chavan, 1960).

Phlyctiderma DALL, 1899 [11] [**Diplodonta semiaspera* PHILIPPI, 1836; OD]. Globose, rounded, inequilateral, posteriorly enlarged, externally punctate, pustulose, or reticulate; ligament long, in groove, resilium narrow; hinge with somewhat projecting bifid 2 and 3b, and oblique 3a, 4b, AIII; posterior muscle scars very broad. *Mio.-Rec.*, N. Am.-W.Pac.-Japan.—FIG. E20,9. **P. semiaspera* (PHILIPPI), *Rec.*, USA(Fla.); 9a,b, LV int., RV hinge, $\times 4$ (Chavan, 1960).

Timothyus HARRIS & PALMER, 1946 [12] [**Sphaerella bulla* CONRAD, 1865; OD] [= *Sphaerella* CONRAD, 1838 (*non* SOMMERFELT, 1834) (type, *S. subvexa*; OD)]. Very globose, rounded, inequilateral, posteriorly enlarged; ligament and resilium adjacent, with long rounded nymph; surface smooth; hinge with 2 and 3b deeply bifid and projecting strongly (2 subvertically, posterior half of 3b almost horizontally), 3a and 4b oblique; anterior muscle scars narrow, much elongated. *L. Eoc.-Rec.*, N.Am.-?Eu.—FIG. E20,6a. *T. subvexus* (CONRAD), *Mio.*, USA(Md.); LV int., enl.




FIG. E21. Ungulinidae (p. N515, N517).

(Chavan, 1960).—FIG. 20,6b,c. **T. bulla* (CONRAD), *Mio.*, USA(Md.); 6b,c, LV and RV hinges, enl. (Chavan, 1960).

GENUS DUBIUM

Pegmapex BERRY, 1960 [**P. phoebe*; M]. Thin, smooth, orbicular; true hinge plate continued posteriorly with flat expansive shelf formed by narrow, sharply angular, deeply sunken escutcheon. Two cardinals in each valve, posterior right and anterior left larger and bifid. *Rec.*, N.Am.(Mex.).

Family CYRENOIDIDAE Adams & Adams, 1857

[=Cyrenellidae FISCHER, 1882]

Suborbicular, posteriorly enlarged, inequilateral, thin, inflated, with rounded prosogyrous beaks; lunule lacking, escutcheon present or absent, ligament and resilium developed marginally; surface concentrically striate; hinge with anterior LV and RV cardinals forming posterior bent part of their original anterior lateral, elongate, forming pattern AI-1, AIII-3, AII-2, with 4b simple and minute, posterior laterals lacking; muscle scars large, elongate, without digitation. *Rec.*




FIG. E22. Cyrenoididae (p. N518).

Cyrenoida JOANNIS, 1835 [**C. dupontia*; OD] [= *Cyrenella* DESHAYES, 1835; *Cyrenoides* SOWERBY, 1839; *Cyrenodonta* ADAMS & ADAMS, 1857]. Characters of family. *Rec.*, Afr.-N.Am.-C.Am.—FIG. E22, 1a-d. **C. dupontiae* [= *C. dupontia* (*lapsus*)], W.Afr.; 1a,b, LV ext., RV hinge, $\times 1.25$ (Sowerby, 1852); 1c,d, LV and RV hinges, enl. (Lamy).—FIG. E22, 1e,f. *C. rhodopyga* VON MARTENS, Congo; 1e,f, LV and RV int., $\times 0.8$ (114).

Superfamily CHAMACEA Lamarck, 1809

[*nom. correct.* ICZN, 1957, Opinion 484 (*pro* Camacea DE BLAINVILLE, 1825) (=camacées LAMARCK, 1809)] [Chamacea MENKE, 1830, not recognized by ICZN] [Materials for this superfamily prepared by MYRA KEEN]

Sculpture normally well developed, concentric or radial or both; shell cemented to substrate by one valve, at least temporarily; beaks prosogyrate, ligament parivincular; hinge degenerate in appearance, with at least one large cardinal tooth in either valve, two cardinals and weak laterals in some; muscle scars two, large, subequal; pallial line entire. ?*U.Cret.*, *Paleoc.-Rec.*

Family CHAMIDAE Lamarck, 1809

[*nom. transl.*, BRODERIP, 1839 (*ex* Camacea DE BLAINVILLE, 1825) (=camacées LAMARCK, 1809)] [Chamidae GRAY, 1823, not recognized by ICZN]

Characters of superfamily. ?*U.Cret.*, *Paleoc.-Rec.*

Chama LINNÉ, 1758 [**C. lazarus*; SD CHILDREN, 1823 (ICZN, Opinion 484, 1957)] [= *Jataronus* BRUGUIÈRE, 1792, AUCTT. (*nom. nud.*); *Maceris*

MODEER, 1793 (*obj.*); *Cameola* RAFINESQUE, 1815 (*nom. null.*); *Lacinea* SOWERBY, 1842 (*obj.*); ?*Hellia* SCHAFHÄUTL, 1863 (type, *H. gryphus*; *Cret.*, Ger.; M)]. Attached by LV throughout life; ornamentation foliaceous. ?*U.Cret.*, *Paleoc.-Rec.*, tropics-Eu.-N.Am.

C. (Chama). Concentric ornamentation of distinctive flattened spines in irregular radial rows. *L.Eoc.-Rec.*, Eu.-Am.-SW.Pac.—FIG. E23, 1. **C. (C.) lazarus*, *Rec.*, E.Indies; 1a-c, RV ext., LV int., RV hinge, $\times 0.5$ (124b, 684).

C. (Cipliacella) VINCENT, 1928 [**Chama pulchra* RAVN, 1902; M] [= *Ciplyella* VINCENT, 1930 (*obj.*); *Ciplelyella* (*nom. null.*)]. Attachment scar obscure; otherwise as in *C. (Chama)*. *Paleoc. (Dan.)*, Eu.

C. (Psilopus) POLI, 1795 [**Chama gryphoides* LINNÉ, 1758; M] [= *Psilopoderma* POLI, 1795 (*obj.*); *Psilotus*, *Psilopoderma* (*nom. null.*)]. Concentric sculpture of small, even, flat scales. *Rec.*, IndoPac.

Arcinella SCHUMACHER, 1817 [**Chama arcinella* LINNÉ, 1767; T] [not preoccupied by *Arcinella* OKEN, 1815, rejected ICZN, Opinion 417] [= *Echinochama* FISCHER, 1887 (*obj.*)]. Shell attached in early growth stages only; ornamentation spinose. *Mio.-Rec.*, SE.USA-C.Am.-S.Am.—FIG. E23, 2. **A. arcinella* (LINNÉ), *Rec.*, Carib., 2a-c, RV ext., LV, RV int., $\times 0.7$ (675, 684).

Pseudochama ODHNER, 1917 [**Chama cristella* LAMARCK, 1819; SD GARDNER, 1923]. Shell attached by RV. *Oligo.-Rec.*, tropics.

P. (Pseudochama). No *Pll* in adult; nepionic shell with concentric sculpture only. *Oligo.-Rec.*, *Medit.-C.Am.-SW.Pac.*—FIG. E23, 3. **P. (P.) cristella* (LAMARCK), *Rec.*, E.Indies; LV ext., $\times 0.5$ (Reeve, 1847).

P. (Eopseuma) ODHNER, 1919 [**P. (E.) pusilla*; M]. Small; nepionic shell with concentric and radial sculpture; *Pll* persisting in adult. *Rec.*, E. Indies.

Superfamily LEPTONACEA Gray, 1847

[*nom. transl.* DALL, 1900 (*ex* Leptonidae GRAY, 1847)] [= *Erycinacea* FISCHER, 1887] [Materials for this superfamily prepared by ANDRÉ CHAVAN]

Marine, byssiferous, with three apertures in mantle—anterior (buccal), median (pedal), and posterior (anal); shell equi-valve, commonly thin and more or less covered by mantle, with small resilium, indenting hinge plate or not, rarely with marginal ligament. Cardinal teeth commonly tubercular, contrasting with elongated laterals and partly atrophied by incomplete individualization of tooth 1; hinge, therefore, para-




FIG. E23. Chamidae (p. N518).

cyrenoid (114). [Burrowing.] ?Cret., Paleoc.-Rec.

The alphabetically arranged generic descriptions in each family-group division of the Leptonacea are accompanied by numbers inclosed by square brackets. Such numbers indicate position in the sequence of generic taxa given with the respective families or subfamilies for the purpose of recording CHAVAN'S arrangement, designed to reflect "natural relationships" of these taxa as inferred by him.

Family ERYCINIDAE Deshayes, 1850

[=Laseadae GRAY, 1847]

Shell somewhat flattened; hinge plate indented under beaks, small, lucinoid; cardinals and laterals present on both valves, duplicate on RV in some; internal ligament in triangular, rather ill-bounded socket, its adjacent marginal part on small nymph in some forms. Paleoc.-Rec.

Arrangement of generic taxa by CHAVAN.—1. *Erycina*.—2. *Hemilepton*.—3. *Scacchia*.—4. *Austroscintilla*.—5. *Lucinaxinus*.—6. *Goodallioipsis*.—7. *Amerycina*.—8. *Parvikellia*.—9. *Litigiella*.—10. *Lasaea*.—11. *Pythina*.—12. *Melliteryx*.—13. *Myllita*.—14. *Zemyllita*.—15. *Myllitella*.—16. *Arthritica*.—17. *Semierycina*.—18. *Erycinopsis*. [Insert above, 12a. ?*Clathroconcha*.]

Erycina LAMARCK, 1805 [**E. pellucida*; SD STOLICZKA, 1871 [=*Migonitis* RAFINESQUE, 1815 (obj.); *Eryx* SWAINSON, 1840 (obj.)]. Transversely elliptical or ovate, externally smooth or with faint radial striae. Anterior RV cardinal faintly developed, other ones trigonal; elongated laterals. Ligament extended backward. Paleoc.-Plio., ?Rec., Eu.-?S.Afr.-N.Am.-C.Am.

E. (*Erycina*) [1]. Rather small and thin, transversely ovate to elliptical, commonly subequilateral, smooth or faintly striated radially. Hinge narrow; posterior LV cardinal small, but not minute, lamelliform narrow laterals parallel to margin. Paleoc.-Mio., Eu.—FIG. E24,1. **E. (E.) pellucida*, Eoc.(Lutet.), France; 1a,b, LV and RV hinges, $\times 4.8$ (Chavan, n).

E. (*Hemilepton*) COSSMANN, 1911 (1912) [2] [**H. longifossula*; OD]. Subelliptical, smooth. Hinge





FIG. E24. Erycinidae (p. N519-N520).



- stout, anterior RV cardinal in direct prolongation of elongated *AIII*; posterior RV and anterior LV obliquely trigonal; posterior LV very minute; posterior lateral stout and distant from ligament, much extended backward. *Eoc.-Plio.*, ?*Rec.*, Eu.-N.Am.-C.Am.-?S.Afr.—FIG. E25,8. *E. (H.) longifossula*, Mio. (Aquitain.), France; *8a,b*, LV and RV hinges, much enl. (Cossmann, 1913).
- Amerycina** CHAVAN, 1959 [7] [**Erycina colpoica* DALL, 1913; OD]. Transversely elliptical, very inequilateral, with opisthogyrous beaks and anterior side by far longest, concentrically furrowed. Long anterior laterals; short posterior; in each valve, one trigonal somewhat bilobate cardinal; anterior right quite indistinct and posterior left narrow against well-defined resilium. ?*Eoc.*, *Mio.-Rec.*, N. Am. (Gulf Calif.)-W.C.Am.-S.Am.—FIG. E25,7. **A. colpoica* (DALL), Gulf Mexico; *7a,b*, RV and LV hinges, $\times 6$ (Chavan, n).
- Arthritica** FINLAY, 1927 [16] [**Kellia bifurca* WEBSTER, 1908; OD]. Ovate, moderately flattened, smooth. Anterior RV and posterior LV cardinals indistinct; *2, 3b*, angular, small, in front of moderately short ligamentary hollow; laterals simple, remote, *2* diverging internal furrows. *Mio.-Rec.*, N.Z.—FIG. E25,13. **A. bifurca* (WEBSTER); *Rec.*, *13a,b*, LV int., RV hinge, $\times 7.2$ (Chavan, n).
- ?**Clathroconcha** COEN, 1934 (12a) [**C. istriensis*; OD]. Small, slightly convex, inequilateral, anteriorly elongated; minute prosogyrous beaks. Sculpture of strong radial ribs, without any dichotomy, but crossed by concentric lamellae. RV hinge with elongate straight laterals and 2 median sockets, anterior one virguliform and deeper, posterior one trigonal, in front of both is one acute thin tooth (or socket's margin). Pallial line broadened; inner margin crenate. *Rec.*, Adriatic.
- Goodalliopsis** DE RAINCOURT & MUNIER-CHALMAS, 1863 [6] [**G. orbignyi* = *Erycina terminalis* DESHAYES, 1860; M]. Obliquely oblong, small; finely striated, very inequilateral, posterior side broadest. Anterior cardinals in prolongation of *AII* and *AIII*; *AI* rather long, posterior cardinals narrow, in front of oblique socket; posterior laterals strong, more elongate than anterior ones. *M.Eoc.*, Eu.—FIG. E24,2. **G. terminalis* (DESHAYES), Lutet., France; *2a,b*, LV int., RV hinge, $\times 16$ (Cossmann, 1913; 259).
- Lasaea** BROWN, 1827 [10] [**Cardium rubrum* MONTAGU, 1803; M] [= *Cycladina* CANTRAINE, 1835 (type, *C. adansoni*; SD GRAY, 1847) (*non* BERTHOLD in LATREILLE, 1827); *Lasea* GRAY, 1842 (*nom. van.*) (obj.); *Anapa* GRAY (type, *Poronia petitiana* RÉCLUZ, 1847); *Poronia* RÉCLUZ, 1843 (type, *P. adansoni*; OD)]. Transversely rounded, minute, inflated, inequilateral, smooth, anterior side longest, beaks straight. Cardinals small, all (except *4b*) partly fused to anterior laterals, which duplicate on both valves; ligament interno-mar-
- ginal, with minute narrow socket. *Eoc.-Rec.*, cosmop.—FIG. E25,12. **L. rubra* (MONTAGU), *Rec.*, Eng.; *12a,b*, LV int., RV hinge, $\times 10$ (Chavan, n).
- Litigiella** DI MONTEROSATO, 1909 [9] [**Erycina cuenoti* LAMY, 1908 (= *Lepton glabrum* P. FISCHER, 1873); OD]. Ovate, inequilateral, somewhat produced and angular anteroventrally, beaks low, hinge plate flat, broad, with rather small, ill-separated laterals and narrow long cardinal in each valve in front of wide hollow for receipt of internal part of broad ligament. *Mio.-Rec.*, Eu.-S. Am.-C.Am.-Australia.—FIG. E25,11. **L. glabra* (FISCHER), *Rec.*, Medit.; *11a,b*, LV hinge and RV int., enl. (Lamy, 1908). [*?Eoc.*, S.Afr.]
- Lucinaxinus** CERULLI-IRELLI, 1909 [5] [**Thyasira (L.) reticulata*; OD]. Subequilateral rounded, with high but not prominent beaks; externally, finely reticulated, laterally with dichotomy of radials. Hinge narrow, RV posterior cardinal oblique, LV anterior angularly fused to *AI*, LV posterior obscure; long marginal laterals; ligament in small oblique socket. *Mio.-Plio.*, Eu.—FIG. E25,4. **L. reticulatus* (CERULLI-IRELLI), Italy (Calabr.); *4a,b*, LV hinge and RV int.; *4c*, LV ext., enl. (Cerulli-Irelli, 1909).
- Melliteryx** IREDALE, 1924 [12] [**Erycina acupuncta* HEDLEY, 1909; OD]. Transversely trigonal; externally punctate. Small cardinals and strong laterals, posterior ones larger and duplicate; narrow ligamentary socket and marginal flattening. *Plio.-Rec.*, Japan-Australia.—FIG. E24,3. **M. acupuncta* (HEDLEY), *Rec.*, Australia; *3a,b*, LV hinge and RV int.; *3c*, RV ext., $\times 3.2$ (365; Hedley, 1909).
- Myllita** D'ORBIGNY & RÉCLUZ, 1850 [13] [**Erycina deshayesii* RÉCLUZ; OD]. Transversely subelliptical, rather broad, thick; divaricate strong curved ribs, small beaks. Cardinals well defined, RV anterior in prolongation of *AIII*; well-marked subparallel laterals, duplicate on RV, anterior ones longest; ligament internomarginal, with both socket and nymph. *Plio.-Rec.*, SW.Pac.-Australia.
- M. (Myllita)**. Moderately transverse, thick, beaks not inflated. Both valves with well-developed cardinals and strong elongate laterals. *Plio.-Rec.*, SW.Pac. - Australia.—FIG. E25,9. **M. (M.) deshayesii* D'ORBIGNY & RÉCLUZ, *Rec.*, Australia; *9a,b*, LV int., RV hinge; *9c*, all $\times 4$ (Chavan, n).
- M. (Zemyllita)** FINLAY, 1927 [14] [**Pythina stowei* HUTTON, 1877; OD]. Very transverse, with rounded beaks. Curved cardinals, only 1 well developed on each valve and short bent laterals. *Plio.-Rec.*, N.Z.-Australia.—FIG. E25,6. **M. (Z.) stowei* HUTTON, N.Z.; *6a,b*, LV and RV hinges, $\times 5.2$; *6c*, LV juv. ext., $\times 2.4$ (Chavan, n).
- Myllitella** FINLAY, 1927 [15] [**M. vivens*; OD]. Circular, with undulated divaricate ribs and con-




FIG. E25. Erycinidae (p. N519-N520, N522).




FIG. E26. Erycinidae (p. N522).

centric growths. Small cardinals and somewhat remote well-marked elongate laterals; hinge plate scarcely indented by internal part of ligament. *Oligo.-Rec.*, N.Z.—FIG. E25,3. **M. vivens*, *Rec.*, N.Z.; 3a,b, LV int., RV hinge, $\times 11$; 3c, LV ext., $\times 6.4$ (Chavan, n).

Parvikellia LASERON, 1956 [8] [**P. isolata*; OD]. Oval, thin, polished, inflated; with faint growths. RV hinge with subumbonal resilifer, simple cardinal in front with slight hook, faint marginal anterior laterals, prominent, posterior; LV hinge (*P. depressa*) with 2 narrow cardinals and 2 laterals, anterior one ascending toward beak as 3rd anterior cardino-lateral. *Rec.*, Australia.—FIG. E26,1a,b. **P. isolata*; 1a,b, RV ext., hinge, enl. (Laseron, 1956).—FIG. E26,1c,d. *P. depressa* LASERON; 1c,d, LV ext., hinge, enl. (Laseron, 1956).

Pythina HINDS, 1844 [11] [**P. deshayesiana*; OD]. Transversely but irregularly trigonal, with divaricate ribs. Very small cardinals and somewhat strong short laterals, posterior ones duplicate on RV; narrow ligamentary hollow and marginal flattening. *U.Eoc.-Rec.*, Eu.-Asia.—FIG. E25,5. **P. deshayesiana*, *Rec.*, Philip. Is.; 5a,b, LV int., RV hinge; 5c, LV ext., $\times 3.2$ (Chavan, n).

Scacchia PHILIPPI, 1844 [**Tellina elliptica* SCACCHI, 1833 (*non* BROCCHI, 1814) (= **Loripes ellipticus* SCACCHI, 1836); SD GRAY, 1847]. Somewhat trigonal, smooth, inequilateral; anterior side long-

est. Cardinals rather small (3b, 2, 4b), laterals simple, one on each side, rising apparently from under side of plate; ligament in oblong socket adjacent to well-marked nymph. *Eoc.-Rec.*, Eu.

S. (Scacchia) [3] Lateral teeth rather narrow and remote, cardinals small. *Eoc.-Rec.*, Eu.—FIG. E25,10. **S. (S.) elliptica* (SCACCHI), *Rec.*, *Medit.*; 10a,b, LV and RV int., $\times 6$ (Cerulli-Irelli, 1909; Cossmann, 1913).

S. (Austroscintilla) KAUTSKY, 1939 [4] [**A. meieri*; OD]. Broad flat laterals, strong right posterior cardinal. *Mio.*, Eu.

Semierycina DE MONTEROSATO in COSSMANN, 1911 (1912) [**Lepton prismaticum* DI MONTEROSATO, 1878; OD]. Small, translucent, transversely sub-elliptical. Hinge with 1 cardinal developed on each valve, its lower part pointed; anterior approximate and posterior remote laterals. *Paleoc.-Rec.*, W.Eu.

S. (Semierycina) [17]. Shell more ovate and beaks farther forward than in *Erycinopsis*. *Mio.-Rec.*, Eu.—FIG. E25,1. **S. (S.) prismaticum* (DI MONTEROSATO), *Rec.*, *Medit.*; 1a,b, LV and RV int., $\times 14.5$ (Chavan, n).

S. (Erycinopsis) CHAVAN, 1959 [18] [**Erycina semipeecten* COSSMANN, 1887; OD]. Trigonal-rounded, concentrically ribbed except on anterior radiated area. RV with strong laterals and 1 vertically projecting cardinal, in front of deep oblique cut. *Paleoc.-Eoc.*, Eu. (France).—FIG. E25,2. **S. (E.) semipeecten* (COSSMANN), *Eoc.*, France; RV int., $\times 5.5$ (160).

Family KELLIIDAE Forbes & Hanley, 1848

[*nom. correct.* SOWERBY, 1862 (pro Kelliidae FORBES & HANLEY, 1848)]

Shell variable in convexity. Hinge plate indented under more or less prominent beaks, hinge paracyrenoid with obliquely trigonal cardinals, RV with 1 strongest, which is almost entirely differentiated from its origin, obliterating 3b, LV with generally two unequal pointed cardinals, the anterior one (2) upward, posterior one downward; no distinctly elongate anterior laterals, but strong remote posteriors present. *Paleoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Kellia*.—2. *Pseudolepton*.—3. *Pseudokellya*.—4. *Aligena*.—5. *Radobornia*.—6. *Paraborniola*.—7. *Diplodontina*.—8. *Lasaeokellya*.—9. *Kaneoha*.—10. *Micropolia*.—11. *Cicatellia*.—12. *Pseudopythina*.—13. *Bornia*.—14. *Lionellia*.—15. *Borniola*.—16. *Byssobornia*.—17. *Scinillula*.—18. *Nesobornia*.—19. *Marikellia*.—20. *Mioerycina*. [Insert above, 4a. *Odontogena*.]

- Kellia** TURTON, 1822 [1] [**Mya suborbicularis* MONTAGU, 1803; SD RÉCLUZ, 1844] [= *Chironia* DESHAYES, 1839 (type, *C. laperousii*; M); *Oronthea* GRAY (ex LEACH, MS) 1852 (obj.)]. Ovate, slightly angular and strongly convex, with concentric striation. Cardinal *3a* in prolongation of *AIII*, *1* stout and oblique, *2a* angular, curved, *2b* conical, pointed; strong posterior laterals; *PI*, *PII* and faint *PIII*; wide cut for ligament. *Eoc.-Rec.*, cosmop.—FIG. E27,5. **K. suborbicularis* (MONTAGU), *Rec.*, Spain; *5a,b*, LV int., RV hinge, $\times 4$ (Chavan, n).
- Aligena** LEA, 1846 (1843, *nom. nud.*) [**Amphidesma aequata* CONRAD, 1843 (= *A. striata* LEA, 1846); SD DALL, 1900]. Irregularly trigonal to quadrangular rounded, high, smooth. Hinge with small *3a*, stout *1*, curved *2a* in prolongation of anterior lateral, with small *2b* behind it; wide oblique ligament and single strong remote posterior lateral on each valve. *Mio.-Rec.*, Eu.-Afr.-N.Am.
- A. (**Aligena**) [4]. Subvertically pointed, more or less virguliform right cardinal; posterior laterals of moderate strength. *Mio.-Rec.*, Eu.-?Afr.-N.Am.—FIG. E27,7. **A. (A.) aequata* (CONRAD), Mio., USA (Va.); *7a,b*, LV and RV int., $\times 4$ (Chavan, n).
- A. (**Odontogena**) COWAN, 1964 [4a] [**A. (O.) borealis*; OD]. Oblique right cardinal, stronger posterior laterals. *Rec.*, N.Am. (W.Can.).
- Bornia** PHILIPPI, 1836 [**B. corbuloides* (= *Erycina corbuloides* BIVONA MS. in PHILIPPI); SD STOLICZKA, 1871 (= *Cyclas sebetia* COSTA, 1836)] [= ? *Ceratobornia* DALL, 1899 (type, *B. (C.) longipes* = *Lepton longipes* STIMPSON, 1855; OD)]. Transversely trigonal to trapezoidal, slightly inequilateral, flattened, shining, smooth or with faint infra-externally placed radials, crenulating inner margin. *Paleoc.-Rec.*, cosmop.
- B. (**Bornia**) [13]. Hinge with narrow oblique well-marked *1*, *2a*, *2b*, and traces of *AIII*, *AII* in front of them; short well-defined resilium and moderately strong *PI*, strong *PII*, traces of *PIII*, all closely behind ligament. *Paleoc.-Rec.*, Eu., N.Am.-S.Ind.O.—FIG. E27,6. **B. (B.) sebetia* (COSTA), *Rec.*, Italy; *6a,b*, LV int., RV hinge, $\times 6.8$ (Chavan, n).
- B. (**Borniola**) IREDALE, 1924 [15] [**B. lepida* HEDLEY, 1906; OD]. Transversely elliptical, striated, inequilateral, with pointed beaks. *Rec.*, Australia.
- B. (**Byssobornia**) IREDALE, 1936 [16] [**B. filosa* HEDLEY, 1902; OD]. Radial striae conspicuous on entire surface. Oblique cardinals and very short posterior laterals. *Rec.*, Australia.
- B. (**Lionelita**) JOUSSEAUME, 1888 [14] [**L. lionelita* (= *Erycina denticulata* DESHAYES, 1855, var. *lionelita* JOUSSEAUME, 1888, in CHAVAN, 1953); OD]. Tooth *1* almost straight, terminating at right angle to its origin in *A1*; *3a* minute, but discernible; *2b* trigonal pointed; resilium oblique, with strong laterals behind it. *Rec.*, Ind.O.—FIG. E27,10. **B. (L.) denticulata lionelita* (DESHAYES, 1855), Obock; *10a,b*, LV and RV int.; *10c*, RV ext., $\times 2.4$ (114).
- Cicatellia** LASERON, 1956 [11] [**C. indenta*; OD]. Transversely oval, small, somewhat inequilateral; ventral margin flexuous; broad, prosogyrous, inflated beaks; surface microscopically wrinkled. LV with conical median cardinal and 2 rather faint laterals, rounded protuberant prodissoconch. *Rec.*, Australia, ?*Mio.-Rec.*, USA (Fla.).—FIG. E28,3. **C. indenta*; *3a,b*, LV ext., hinge, enl. (531).
- Diplodontina** STEPELL, 1899 [7] [**D. tumbesiana*; M]. Transversely ovate, moderately broad, with concentric striae. LV showing trigonal bilobate cardinal, its posterior part smaller than on *Kellia*; short anterior lamina and broad oblique resilium with parallel thickening above it; posterior margin enlarged. *Rec.*, W.S.Am.—FIG. E27,3. **D. tumbesiana*, Peru; LV int., $\times 4$ (Stempel, 1899).
- Kaneoha** DALL, BARTSCH & REHDER, 1938 [9] [**K. rosea*; OD]. Very small, rounded, smooth, somewhat inflated. *A1-1* and *3b* coalescent under marginal *AIII*; *2* hooked with long *AII*, not detached from it; stout resilium and rather long posterior laterals (duplicate on RV). *Rec.*, Hawaii.—FIG. E27,1. **K. rosea*; *1a,b*, LV and RV hinge, $\times 32$ (Dall, Bartsch & Rehder, 1938).
- Lasaeokellya** COSSMANN & PEYROT, 1911 (1912) [8] [**Kellia (L.) cestasensis*; OD]. Orbicular, subequilateral, slightly convex, with lamellar growths. RV with oblique *1*, fused to anterior side of *3b* into bifid inequilateral unit; resilium oblique, scarcely indenting inferior margin of plate; posterior laterals toothlike. *Mio.*, Eu. (France).—FIG. E29,2. *L. cestasensis*, Burdigal, France; RV int., $\times 6.4$ (165).
- Marikellia** IREDALE, 1936 [19] [**Kellia solida* ANGAS, 1877; OD]. Transversely subquadrangular, solid, rather inflated, with irregular concentric lines, small beaks. Hinge broad, with stout trigonal *1* and *2b*, oblique ill-defined *2a* and *3a*; also somewhat oblique long resilium indenting hinge plate, posterior laterals strong. *Rec.*, E.Afr.-Australia.—FIG. E27,4. **M. solida* (ANGAS), Australia; *4a,b*, LV and RV int., $\times 4$ (Chavan, n).
- Micropolia** LASERON, 1953 [10] [**M. typica*; OD]. Minute, translucent, subtrigonal and subequilateral, smooth. LV hinge with tooth pointed backward and 2 bow-shaped laterals; RV with 2 elongated laterals. *Rec.*, Australia.—FIG. E28,2. **M. typica*; *2a-c*, LV ext., LV and RV hinges, enl. (531).
- Mioerycina** KAUTSKY, 1939 [20] [**Erycina letochai* HOERNES, 1859; OD]. Transversely elliptical, sinuous in middle, almost equilateral. Hinge with short *1* and *2b*; *AIII-3a* and *2a-AII* being undifferentiated from margin while very small *3b* occurs behind top of *1*; oblique spoon-shaped resilium and long posterior laterals. *Mio.*, Eu. (Austria).




FIG. E27. Kelliidae (p. N523, N525-N526).

—FIG. E27,8. **M. letochai* (HOERNES), M.Mio., Austria; 8a,b, LV int. and RV hinge, enl. (456).
Nesobornia DALL, BARTSCH & REHDER, 1938 [18] [**N. bartschi* CHAVAN, herein (pro *Erycina* (*Poronia*) *ovata* GOULD, 1850, non GRAY, 1825, nec PHILIPPI, 1836)]. Subtrapezoidal rounded, somewhat inequilateral, obliquely elongated backward, shining and smooth; beaks pointed with 2 small cardinals on each valve: 3a, 1, 2a, 2b, with faint short anterior prolongations of the former and somewhat backward produced resilium, posterior laterals well-marked. ?*M.Eoc.-Rec.*, Hawaii-E.Afr.
 —FIG. E27,13. **N. bartschi* CHAVAN, Rec., Hawaii; 13a,b, LV int. and RV hinge, $\times 3.2$ (Chavan, n).

Pseudokellyya PELSENEER, 1903 [3] [**Kellia cardiiformis* E. SMITH, 1885; OD]. Globular, high, thin, with radial riblets; beaks inflated. Hinge very narrow, LV with 2 small cardinals in prolongation of short anterior lateral thickening and RV with single cardinal, then ligamentary cut and posterior thickening. *Rec.*, Antarctic.—FIG. E29,1. **P. cardiiformis* (SMITH); LV int., $\times 4.8$ (852).

Pseudolepton COSSMANN, 1895 [2] [**Lepton insigne* MAYER in HÖRNES, 1864; OD]. Subquad-rangular to ovate, with diverging striae or granular laminae. Hinge with 3a, 3b, oblique 2a, 2b poorly




FIG. E29. Kelliidae (p. N523, N525).

developed; remote posterior laterals *PI*, *PII*. *Mio.*, ?*Rec.*, Eu.-?S.Afr.—FIG. E27,2. **P. insigne* (MAYER), *Mio.*, Aus.; 2a,b, LV and RV hinges, enl. (Cossmann, 1913; 456).

Pseudopythina P. FISCHER in DI MONTEROSATO, 1884 [12] [**Kellia macandrewi* FISCHER, 1867 = *Corallophaga setosa* DUNKER, 1864; M] [= ?*Bornioopsis* HABE, 1959 (type, *B. tsurumaru*; OD)]. Transversely trigonal, very inequilateral, anteriorly attenuated and elongate; enlarged and rounded backward, smooth. No anterior prolongation of 1, 2 oblique and well marked; 3a small, 2b minute; with oblique short resilium and nymph; posterior laterals just behind them, short, placed marginally. *Plio.-Rec.*, W.Eu.-NW.Pac.—FIG. E27,12. **P. setosa* (DUNKER), Rec., Portugal; 12a,b, LV int. and RV hinge, $\times 6.8$ (Chavan, n).

Radobornia DALL, BARTSCH & REHDER, 1938 [**R. arafia*; OD]. Transversely rounded, inequilateral, beaks rather prominent; RV with small laminar 3a and strong hooked 1, LV with curved 2a-*All* and curved approximate 2b, each valve with one remote lamellar posterior lateral; long internal resilium and marginal nymph. *Rec.*, Hawaii.

R. (Radobornia) [5]. Transversely ovate, thin, externally punctate; ventral margin rounded. *Rec.*, Hawaii.—FIG. E27,9. **R. (R.) arafia* DALL, BARTSCH & REHDER; 9a,b, LV and RV hinges, $\times 2.8$; 9c, LV ext., $\times 1.6$ (Dall, Bartsch, and Rehder, 1938).

R. (Paraborniola) HABE, 1958 [6] [**P. matsumotoi*; OD]. Transversely elongate, ventral margin straight. Hinge as in *R. (Radobornia)* with 2a thinner and 2b strongly trigonal. *Rec.*, Japan.—FIG. E28,1. **R. (P.) matsumotoi* (HABE); 1a,b, RV ext., hinges, $\times 1$ (388).

Scintillula JOUSSEAUME, 1888 [17] [**S. scintillula*; OD]. Transversely subelliptical almost equilateral, inflated, slightly enlarged backward, finely striated. Hinge with marginal 3a, slightly continued by 3b above and behind hooked trigonal 1, itself in prolongation of *All*; similarly hooked 2a-*All* followed by tubercular small 2b on opposite valve; and, perhaps, also minute 4b; oblique short re-




FIG. E28. Kelliidae (p. N523, N525).

silium and much elongated faint posterior laterals. *Rec.*, Ind.O.—FIG. E27,11. **S. scintillula*, Obock; 11a,b, LV and RV int.; 11c, RV ext., ×4 (114).

Family LEPTONIDAE Gray, 1847

Shell commonly compressed; hinge plate not or scarcely indented by resilium, under very small beaks. Paracyrenoid hinge of

neponic appearance, oblique strongest cardinals (right 1 and left 2a) being not separated from their anterior original lamellae (AI, AII), elongating in front of them. Posterior laterals only slightly longer than anteriors. Pallial line irregular and distant from ventral margin. ?*Cret.*, *Paleoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Lepton*.—2. *Planikellia*.—3. *Divarikellia*.—




FIG. E30. Leptonidae (p. N527).

4. *Vermitexta*.—5. *Anomalokellia*.—6. *Platomysia*.—7. *Semeloidea*.—8. *Temblornia*.—9. *Solecardia*.—10. *Properycina*.—11. *Potidoma*.

Lepton TURTON, 1822 [**Solen squamosus* MONTAGU, 1803; SD HERRMANNSEN, 1846] [= *Eupoleme* GRAY (ex LEACH, MS), 1852 (obj.)]. Ovate-subquadrangular, flattened. Hinge with 2 narrow cardinals on each valve, of which anterior ones (1, 2a) are gradually strengthened straight from anterior small crest; 3a very oblique; 4b narrow, both small and thin; resilium broad and trigonal; posterior laterals (duplicate on right) just behind and above it, lamellar. ?*Cret.*, *Eoc.-Rec.*, W.Eu.-W.N.Am.

L. (Lepton). [1]. Ovate, with slight oblique truncation; punctate. Hinge with elongate laterals and 4b almost vertical. ?*Cret.*, *Eoc.-Rec.*, Eu.—FIG. E30,9. **L. (L.) squamosum* (MONTAGU), *Rec.*, Ireland; 4a,b, LV int. and RV hinge, $\times 4$ (Chavan, n).

L. (Divarikellia) COSSMANN, 1887 [3] [**Erycina nitida* CAILLAT, 1834; OD]. Ovate-rounded, with divaricate riblets. Anterior laterals inflexed, but moderately long, 4b obscure, elongate posterior laterals. *Eoc.*, W.Eu.—FIG. E30,5. **L. (D.) nitidum* (CAILLAT), Lutet., France; 5a,b, LV int. and RV hinge, $\times 6.4$ (Chavan, n).

L. (Planikellia) COSSMANN, 1887 [2] [**Erycina radiolata* LAMARCK, 1805; OD]. Ovate-rounded, with radial striae. Short anterior laterals, inflexed at their strengthening into cardinals, 4b oblique backward; stouter posterior laterals. *Eoc.*, W.Eu.—FIG. E30,3. **L. (P.) radiolatum* (LAMARCK), Lutet., France; 3a,b, LV int. and RV hinge, $\times 4.8$ (Chavan, n).

Anomalokellia COSSMANN, 1887 [5] [**Erycina catalaunensis* COSSMANN, 1883; M]. Transversely elliptic-trigonal, small, with concentric striae. Hinge with marginal faint *AIII-3a*, stout angular *AI-1*, thinner angular *AII-2a* and subsymmetric oblique 4b. Stout angular resilium; remote laterals. *Paleoc.*, Eu.(France).—FIG. E30,4. **A. catalaunensis* (COSSMANN), Thanet., France; 4a,b, LV int. and RV hinge, $\times 6.4$ (Cossmann, 1913).

Platomysia HABE, 1951 [6] [**P. rugata*; OD]. Transversely subelliptical, with concentric regular undulations; 2 small cardinals, anterior one prolonged by marginal small crest. *Rec.*, Japan-N.Atl.-?N.Sea.—FIG. E30,1. **P. rugata*; LV int., $\times 0.8$ (365).

Properycina CERULLI-IRELLI, 1908 [**P. mariana*; OD]. Subtrigonal to ovate, with small beaks. Long subsymmetrical anterior laterals and only 1 oblique cardinal (4b) distinct from anterior and posterior laterals (both duplicate on right valve). Tooth 1 almost undifferentiated from *AI*. *M.Eoc.-Rec.*, W. Eu.-Atl.-Medit.

P. (Properycina) [10]. Anterior and posterior margins oblique; 4b quite distinct, laterals al-

most straight. *U.Eoc.-U.Plio.*, W.Eu.—FIG. E30,7. **P. (P.) mariana*, Calabr., Italy; 1a,b, LV int. and RV hinge, enl. (Cerulli-Irelli, 1909).

P. (Potidoma) DEROUX, 1961 [11] [**Lepton subtrigonum* FOLIN & PERRIER (ex JEFFREYS MS), 1873; OD]. Anterior and posterior margins nearly horizontal, 4b appressed against anterior cardino-lateral 2-*AII*; laterals sinuate. *M.Eoc.-Rec.*, W. Eu.-Atl.-Medit.

Semeloidea BARTRUM & POWELL, 1928 [**S. donaciformis*; M]. Trigonal to subquadrangular, thin, with laterally, several radial riblets; LV with 2 well-marked cardinals, posterior one passing above top of anterior one, obliquely attenuated onward; 1 on RV prolonged angularly into small crest under elevated margin; curved, posterior laterals. *Eoc.-Rec.*, *Medit.-N.Z.-C.Eu.*

S. (Semeloidea) [7]. Hinge plate arched under resilium, 4b posteriorly curved, 2 rather strong. Posterior laterals long. *Oligo.-Rec.*, Eu.-N.Z.—FIG. E30,2a. **S. (S.) donaciformis* BARTRUM & POWELL, *Rec.*, N.Z.; LV int., enl. (Bartrum & Powell, 1928).—FIG. E30,2b. **S. (S.) geoffroyi* (PAYRAUDEAU), *Rec.*, *Medit.*; RV hinge, $\times 4$ (Chavan, n).

S. (Temblornia) KEEN, 1943 [8]. [**Donax triangulatus* ANDERSON & MARTIN, 1914; OD]. Short cardinals and shallow small resilium, so that posterior laterals, just behind it, look like cardinals; hinge plate straight under resilium. *Eoc.-Plio.*, N.Am.-S.Am.—FIG. E30,6. **S. (T.) triangulata* (ANDERSON & MARTIN), Mio., USA (Calif.); 6a,b, LV int. and RV hinge, $\times 7.8$ (Chavan, n).

Solecardia CONRAD, 1849 [9] [**S. eburnea*; M]. Medium-sized, transversely elliptical, moderately convex, punctate, with crenulated margin, hinge with oblique cardinals, anteriorly differentiated from short anterior crest; trigonal resilium and oblique short posterior laterals just above and behind it. *Plio.-Rec.*, N.Am.-E.Afr.—FIG. E30,8. **S. eburnea*, *Rec.*, Mex.; LV int., $\times 2.4$ (Chavan, n).

Vermitexta LASERON, 1956 [4] [**V. garrardi*; OD]. Nearly equilateral, small, slightly elongated anteriorly; translucent, thin, moderately inflated; sculpture of faint growths and of microscopic short radially arranged broken corrugations. Ligament small; small median cardinal and anterior one, prolonged into long marginal lateral; moderately long laminar posterior laterals. ?*Eoc.*, Eu.; *Rec.*, Australia.—FIG. E30,10. **V. garrardi*, *Rec.*, Australia; 10a,b, LV int., RV hinge, enl. (531).

Family MONTACUTIDAE Clark, 1855

Shell more or less convex, rounded, inequilateral; resilium in large defined hollow




FIG. E31. Montacutidae (p. N529-N533).

or socket under beaks. Hinge with no true separated cardinals, but only thickened, generally subsymmetrical laterals, commonly bent up at their umbonal end so as to form distant toothlike hook, therefore no more than one pseudocardinal on each valve; posterior laterals commonly duplicate on one valve. *Eoc.-Rec.*

Generic arrangement by CHAVAN.—1. *Montacuta*.—2. *Devonia*.—3. *Entovalva*.—4. *Sciobereitia*.—5. *Pythinella*.—6. *Tellimya*.—7. *Nipponomysella*.—8. *Laubriereia*.—9. *Namnesia*.—10. *Decipula*.—11. *Kelliopsis*.—12. *Fronsella*.—13. *Neaeromya*.—14. *Isorobitella*.—15. *Orobitella*.—16. *Axinodon*.—17. *Merignacia*.—18. *Montacutona*.—19. *Issina*.—20. *Kona*.—21. *Lasaeoneaera*.—22. *Virmysella*.—23. *Rochefortula*.—24. *Mysella*.—25. *Rochefortia*.—26. *Malvinasia*.—27. *Curvemysella*.—28. *Sphaerumbonella*.—29. *Mancikellia*.—30. *Pileatona*.—31. *Nipponomontacuta*.—32. *Fastimysia*.—33. *Thecodonta*.—34. *Pristes*.—35. *Barrimysia*.—36. *Callomysia*. [Insert above, 3a. *Conchentopyx*; 31a. *Tahunanuia*.]

Montacuta TURTON, 1822 [1] [**Ligula substriata* MONTAGU, 1808; SD HERRMANNSEN, 1846] [= *Sphenalia* S. WOOD, 1874 (obj.); *Coriareus* HEDLEY, 1907 (type, *C. vitreus*; OD); *Montaguia* BRONN, 1848 (obj.)]. Small, subquadrangular, rounded in front, truncated backward, inflated; with radial striae or distant riblets and concentric lamellar growths. Rounded prosogyrous beaks. Elongate laterals, more or less duplicate on right valve (upper margin scarcely differentiating into *AIII* and *PIII*); anterior ones stronger. Small resilium. *Eoc.-Rec.*, Eu.—FIG. E31,1. **M. substriata* (MONTAGU), *Rec.*, Britain; 1a,b, LV int. and RV hinge, $\times 8$; 1c, RV ext., $\times 4$ (Chavan, n).

Axinodon VERRILL and BUSH, 1898 [16] [**A. ellipticus* (= *Kellia symmetros* JEFFREYS, 1876); OD] [= *Kelliola* DALL, 1899 (obj.)¹]. Ovately trans-

¹ Identity of *K. symmetros* with *A. ellipticus* established by Dr. K. OCKELMANN.

verse, rounded, orthogyrous; LV hinge with rounded 2, RV with tuberculiform ill-defined 1; chondrophore oblique. *Rec.*, N.Am.-N.Atl.—FIG. E32,7. **A. symmetros* JEFFREYS, USA (Mass.); 7a,b, LV and RV hinges (Chavan, n).

Barrimysia IREDALE, 1929 [**Rochefortia excellens* HEDLEY, 1912; OD]. Very transverse, with striae crenulating inner margin. Two well-marked laterals, subsymmetrical on both sides of deep somewhat oblique resilium. *Rec.*, Australia-Japan.

B. (*Barrimysia*) [35]. Outline trigono-elliptical, nearly equilateral. *Rec.*, Australia.

B. (*Callomysia*) HABE, 1951 [36] [**C. matsuii*; OD]. Subquadrangular transverse, inequilateral, with concentric lines; also oblique riblets at lateral ends. *Rec.*, Japan.—FIG. E31,5. *B. (*C. matsuii*) (HABE); 5a,b, LV int., ext., $\times 1.6$ (365).




FIG. E32. Montacutidae (p. N529-N530, N532-N533).

Conchentopyx BARNARD, 1964 (3a) [**C. granulosa*; M]. Shell internal, semioval, thin; umbo at anterior quarter; prodissoconch prominent, circular. Coarse growth lines and faint radial striae; inner margin smooth; hinge edentulous; no anal siphon. *Rec.*, S.Afr.

Curvemysella HABE, 1959 [27] [**Mysella paula* (= *Pythina paula* A. ADAMS, 1856); OD]. Transversely elongated, obliquely sloping at each end, with interior and posterior dorsal angulation; medially depressed. Hinge narrow, with lateral lamina on each side of resilial depression. *Rec.*, N. Australia-New Guinea-Japan.—FIG. E33,1. **C. paula* (A. ADAMS), Japan; 1a,b, RV ext., int., $\times 1$ (Habe, 1959).

- Decipula** FRIELE, 1875 [10] [**D. ovata* (= *Montacuta tenella* LOVÉN, 1846); OD]. Transversely oval, thin, inequilateral; beaks low, orthogyrous; anterior side elongated. Long narrow anterior laterals, obliquely modified into right toothlike and left enlarged termination; triangular oblique resilium. *Plio.-Rec.*, Atl.-Medit.—FIG. E32,1. **D. tenella* (LOVÉN), *Rec.*, N.Sea; 1a,b, RV and LV hinges; 1c, LV ext., all enl. (829).
- Devonia** WINCKWORTH, 1930 [2] [*pro Synapticola* MALARD, 1904 (non VOIGHT, 1892)] [**Synapticola perrieri* MALARD, 1904; OD]. Transversely subquadrangular, very inequilateral, elongated onward and truncated at both sides. Hinge with a marginal ill-defined *All* and, behind a broad resilial hollow, a strong *Pll*. *Eoc.-Rec.*, W.Eu.—FIG. E31,6. **D. perrieri* (MALARD), *Rec.*, Britain (Devon); LV int., $\times 8.8$ (905a).
- ‡**Entovalva** VÖLTZKOW, 1890 [3] [**E. mirabilis*; M] [‡=*Cycladoconcha* SPAERCK, 1932 (type, *C. amboinensis*; M)]. Like *Devonia* but shell completely internal, with more distinct teeth. *Rec.*, Zanzibar.
- Fastimysia** IREDALE, 1929 [32] [**Rochefortia viastellata* HEDLEY, 1909; OD]. Trigono-elliptical, very transverse, inequilateral, shortly sloping under beaks, then obliquely truncated; strongly acuminate at opposite side. Deep resilium in semielliptical socket just under beaks. Almost subsimilar elongated flexuous laterals on both sides. *Rec.*, Australia.—FIG. E31,14. **F. viastellata* (HEDLEY); 14a,b, LV ext., hinge, $\times 11.5$ (433).
- Fronsella** LASERON, 1956 [12] [**F. adipata*; OD]. Transversely subquadrangular, to ovately transversal, moderately inflated, anterior side longer. Hinge narrow, bearing in each valve small anterior conical teeth, anterior margin somewhat laminar, and, after broad ligamentary indentation of plate, lamellar rather strong posterior lateral. *Rec.*, Australia-Japan.—FIG. E33,4b,c. **F. adipata*, Australia; 4b,c, LV ext. and hinge, enl. (Laseron, 1956).—FIG. E33,4a. *F. oshimai* HABE, Japan; LV hinge, enl. (Habe, 1958).
- Issina** JOUSSEAUME, 1898 [19] [**I. issina*; OD]. Subquadrangular, small, very inequilateral, posterior side much elongated. On each valve, 2 anterior laterals, marginal one curved, internal one shorter and pointed; and one long curved posterior lateral. Very narrow, but well-marked resilium. *Rec.*, Red Sea.—FIG. E31,18. **I. issina*; 18a,b, LV int. and RV hinge, $\times 8$ (Melvill & Standen, 1914).
- Kelliopsis** VERRILL & BUSH, 1898 [11] [**Montacuta elevata* STIMPSON, 1851; OD] Ovately rounded, inequilateral, anterior side longest. On each valve, anterior and posterior toothlike short stout lateral, prolonged by thickening of margin with small sub-trigonal resilium between them. *Rec.*, Am.—FIG. E31,4. **K. elevata* (STIMPSON), USA; 4a,b, LV and RV int., enl. (937).
- Kona** DALL, BARTSCH & REHDER, 1938 [20] [**K. bucki*; OD]. Subelliptical, small inequilateral. Resilium stout and short, on well-developed shelf. On each valve single short subperpendicular cardinal-like process and 2 (anterior and posterior) unequal true laterals, both well marked. *Rec.*, Hawaii.—FIG. E31,15. **K. bucki*; 15a,b, LV and RV int., $\times 8$ (Dall, Bartsch, & Rehder, 1938).
- Lasaoneaera** COSSMANN, 1913 [21] [**Corbula radiata* DESHAYES, 1824; M]. Moderately small, anteriorly rounded, posteriorly acuminate. Regular radial ribs, of which, on each valve, is an irregularly thick anterior lateral and, backward, an obliquely divided posterior one. *Eoc.*, Eu.(France).—FIG. E32,3. **L. radiata* (DESHAYES), Lutet., France; 3a,b, LV and RV int., $\times 3.3$; 3c, LV ext., $\times 2.7$ (Deshayes, 1837).
- Laubriereia** COSSMANN, 1887 [8] [**Erycina emarginata* DESHAYES, 1860; OD]. Small, trigonal-rounded to subquadrangular, very inequilateral, concentrically striated, anterior side much elongated. Narrow hinge with, on each valve, marginal long anterior lateral enlarged and bent into short, toothlike, additive one under the beak; also marginal posterior lateral. Between them, semi-circular socket for resilium. *Eoc.*, ?*Plio.*, W.Eu.—FIG. E31,7. **L. emarginata* (DESHAYES), Lutet., France; LV int., enl. (259).
- Malvinasia** COOPER & PRESTON, 1910 [26] [**M. arthuri*; M]. Subtrigonal, small inequilateral, posterior side produced. On each valve, anterior long marginal lateral, accompanied by shorter oblique one, toothlike on right valve, hooklike and joined to marginal on left. Subperpendicular trigonal resilium and one sinuate posterior lateral. *Rec.*, Falkland Is.—FIG. E31,17. **M. arthuri*; 17a,b, RV hinge and LV int., enl. (Chavan, n).
- Mancikellia** DALL, 1899 [29] [*pro Zoe* MONTEROSATO, 1878 (non PHILIPPI, 1840)] [**Zoe pumila* DI MONTEROSATO, 1878 (= *Montacuta pumila* S. WOOD, 1840); OD]. Small convex inequilateral rounded shell, anteroventrally enlarged. Hinge with well-marked laterals, anterior ones separated by broad socket from thickened margin and completed, under beak (in right valve only) by a stout toothlike termination; posterior laterals slightly shorter than anterior and separated from them by large oblique hollow resilifer. *Plio.-Rec.*, W.Eu.-Medit.—FIG. E31,11. **M. pumila* (WOOD), *Rec.*, S.Sea; 11a,b, LV hinge and RV int., $\times 8$ (Friele, 1886).
- Merignacia** COSSMANN, 1914 [17] [**Pseudolepton (M.) pleurodesmatoides* COSSMANN & PEYROT, 1914; OD]. Small, ovate-rounded, flattened, inequilateral, little longer anteriorly. Marginal sinuous anterior lateral on each valve and a toothlike hook appressed to it (stouter on the right). Superficial resilial socket. Moderately flat posterior lateral. Very dissymmetric scars. *Mio.*, Eu.(France).—FIG. E31,8. **M. pleurodesmatoides* (Coss-




FIG. E33. Montacutidae (p. N529-N532).

MANN & PEYROT), Burdigal., France; 8a,b, LV int. and RV hinge, enl. (Cossmann & Peyrot, 1914).
Montacutona YAMAMOTO & HABE, 1959 [18] [**M. mutsuwanensis*; OD]. Suborbicular, compressed, with polished surface, marked only by weak

growths. Two solid diverging teeth and a resilial gap between them. Lateral hinge margins more or less thickened and lamellar. *Rec.*, Japan.—FIG. E33,2. **M. mutsuwanensis*; RV ext., int., $\times 4$ (Habe & Yamamoto, 1959).

Mysella ANGAS, 1877 (Aug.) [**M. anomala*; OD] [= *Petricola* GRAY, 1825 (*non* LAMARCK, 1801) (type, *Mya bidentata* MONTAGU, 1803; SD GRAY, 1847, subj.)]. Transversely trigono-elliptical, inequilateral, shortened in front, obliquely elongate backward. Approximate concentric ribs. Well-marked laterals, posterior right strongest, anterior right marginal, 1 minute bordering it, 2 oblique and long. Prominent long chondrophore³ and trigonal resilium. *Mio.-Rec.*, cosmop.

M. (Mysella) [24]. Resilium without crests; chondrophore obliquely elongate, 2 long under *fAl*, 1 long, very thin and oblique more or less truncate at both ends. *Mio.-Rec.*, cosmop.—FIG. E31,9. **M. anomala*, *Rec.*, Vict.; 9a,b, LV int. and RV hinge, $\times 8$ (Chavan, n).

M. (Rochefortia) [25] VÉLAIN, 1877 (post-Nov.) [?1878] [**R. australis*; M]. Resilium with crests as bifid tooth; chondrophore stouter and shorter, with minute crest at its top on RV; 1 subvertical and *Al* attenuated, thin; 2 almost marginal. More or less rounded at both ends. *Plio.-Rec.*, Ind.O. (St. Paul Is.)-Australia-N.Am.—FIG. E31, 3. **R. australis* (VÉLAIN), *Rec.*, St. Paul Is.; 3a,b, LV int., RV hinge, much enl. (Vélain, mod.).

Namnetia COSSMANN, 1905 [9] [**N. discoides*; M]. Oblong-ovate, smooth, with flattened beaks and very small hinge plate. Marginal left curved and 2 right marginal and minute, infraumbonal anterior laterals. Restricted short resilium and, on each valve, long marginal posterior lateral. No nymph. Anterior adductor clublike. *Eoc.*, Eu. (France).—FIG. E31,13. **N. discoides*, Lutet., France; 13a,b, LV int. and RV hinge, $\times 2.7$ (Cossmann, 1905).

Neaeromya GABB, 1873 [**N. quadrata*; M]. Transversely subquadrangular to trigonal, finely striated. Right strong anterior laminar tooth and oblique posterior enlarged margin; oblique resilium between them; left thinner oblique anterior. *Mio.-Rec.*, C.Am.-N.Am.-Eu.

N. (Neaeromya) [13]. Subquadrangular, anterior side elongate, inequilateral. Right anterior tooth long, left anterior prolonged into margin. *Mio.-Rec.*, USA (Calif.)-C. Am.-Eu.—FIG. E31,12. **N. (N.) quadrata* GABB; 12a,b, LV and RV hinges, $\times 2$ (Pilsbry, 1921).

N. (Isorbitella) KEEN, 1962 [14] [**N. (I.) singularis*; OD]. Ovate-trigonal, subequilateral. Resilium less oblique than in *Oorbitella*, being

³In such shells, the chondrophore (usually cited as *Pl*, *Pll*) functions also as posterior laterals, with its edge laminar and posterior part depressed (*Pl*, *Pll*) in RV, the contrary in LV (*fPl*, *fll*) so that the other, remote, marginal true laterals may be given as *Plll* and *Pllv*.




FIG. E34. Montacutidae (p. N531-N532).

almost parallel to the hinge margin, and showing a median ridge. No gap between the cardinal and the resilifer. *Rec.*, W.Mex.—FIG. E34, 2. **N. (I.) singularis*; 2a,b, LV, RV hinges, $\times 10$ (Keen, 1962).

N. (Orobitella) DALL, 1900 [15] [**Montacuta floridana* DALL, 1899; OD]. Anterior margin broadly rounded, elongate, thus very inequilateral. Right laminar tooth rather small. *Mio.-Rec.*, N. Am.—FIG. E31,16. **N. (O.) floridana* (DALL), *Rec.*, USA(Fla.); RV int., enl. (Dall, 1900).

Nipponomontacuta YAMAMOTO & HABE, 1961 [31] [**N. actinariophila*; M]. Transversely ovate, subequilateral, anterior end shorter and narrower; moderately convex. Sculpture of growth lines and of several anteroventral radial cords. LV hinge with 2 diverging teeth, posterior one stouter, resilifer between them; lateral margins enlarged, posterior groove for *PIII*. *Rec.*, Japan.—FIG. E33,5. **N. actinariophila*; 5a,b, RV ext., LV hinge, enl. (Habe & Yamamoto, 1961).

?**Nipponomysella** HABE, 1959 [7] [**Montacuta oblongata* YOKOYAMA, 1922; OD]. Quite transversely inequilateral, donaciform, with growth lines. Looks like *Tellimya*, but hinge said to be edentulous in right valve and with two unequal cardinals in left. *Rec.*, Japan.—FIG. E33,3a. **N. oblongata* (YOKOYAMA); RV ext., $\times 3$ (Habe, 1959). —FIG. E33,3b. *N. obesa* HABE; LV int., $\times 7.5$ (Habe, 1960).

Pileatona LASERON, 1956 [30] [**P. compressa*; OD]. Small, thin, translucent, subtrigonal, with nearly erect beaks and fine growths. Hinge with narrow subumbonal resilifer and 2 obscure subsymmetric laterals. *Rec.*, Australia.—FIG. E33,6. **P. com-*

pressa; 6a,b, LV ext., hinge, enl. (Laseron, 1956). **Pythinella** DALL, 1899 [5] [**Montacuta cuneata* VERRILL & BUSH, 1898; OD]. Transversely subtrigonal, very inequilateral, elongated posteriorly and attenuated at both sides. Orthogyrous beaks. Hinge with rather long and narrow right, shorter and stouter left, laterals. Resilium restricted to the upper part of a wide arch under the beaks. *Rec.*, N.Am.—FIG. E34,1. **P. cuneata* (VERRILL & BUSH), Cape Hatteras; 1a-c, LV ext., hinges, $\times 11.3$ (937).

Rochefortula FINLAY, 1927 [23] [**Rochefortia reniformis*; SUTER, 1909; OD]. Transversely subtrigonal, rounded and sinuate ventrally, ornamented by concentric ribs, regularly crossed by concentric striae. Prosogyrous moderately high beaks. Trigonal well-defined resilium; left anterior and posterior laterals subsymmetric; right anterior oblique, the posterior somewhat remote. *Mio.-Rec.*, Australia-N.Z.-S.Atl.—FIG. E31,2. **R. reniformis* (SUTER), N.Z.; 2a,b, LV int. and RV hinge, $\times 11.5$; 2c, LV ext., enl. (Chavan, n).

?**Scioberetia** BERNARD, 1895 [4] [**S. australis*; M]. Subcircular, inequilateral, very minute; several crenulations on margin before and behind resilial socket. *Rec.*, S.Am.-Antarctica.

Sphaerumbonella COEN, 1933 [28] [**S. brunellii*; OD]. Ovate, globose, thick, slightly truncated laterally. Rounded beaks. Two short strong subsymmetric right laterals, anterior and posterior, on enlarged margin and on both sides of oblique deep socket. On left valve more internally placed laterals under sockets of right ones. *Rec.*, Red Sea. —FIG. E32,4. **S. brunellii*; 4a,b, LV and RV int., both enl. (Coen, 1933).

Tahunanuia POWELL, 1952 [31a] [**T. alata*; OD]. Thin, trapezoidal, very inequilateral, anteriorly short and rounded, posteriorly broadened and truncated. Irregular distant concentric waves. Hinge with single tooth in each valve, right one subvertically projecting, left one subhorizontally curved. Oblique broad resilifer. Slight radial ridge bordering inner margin of both scars. *Rec.*, N.Z.

Tellimya BROWN, 1827 [6] [**Mya ferruginosa* MONTAGU, 1808; SD GRAY, 1847] [= *Tellinomya* AGASSIZ, 1846 (*emend.*)]. Shell subelliptical, very inequilateral, elongated and attenuated onward, shortly truncated backward. Externally striated. Slightly opisthogyrous beaks, under which lies small deep resilial socket. Oblique anterior laterals, right one long, attenuated, subhorizontal, left one shorter and more oblique. Short, strong, almost vertical resilium. *Mio.-Rec.*, W.Eu.-Am.—FIG. E32,2. **T. ferruginosa* (MONTAGU), *Rec.*, USA(Mass.); 2a,b, LV int. and RV hinge, $\times 4$ (937 mod., and Chavan, n).

Thecodonta A. ADAMS, 1864 [**T. sieboldi*; M]. Transversely rounded, convex, very inequilateral, posterior part much elongated. Beaks prosogyrous. Sculpture concentric. Anterior left lateral curved,

gradually increasing subparallel to margin, then straightened along oblique resilium. Posterior laterals remote, thickened. *Rec.*, W.Pac.-E.Pac.

T. (Thecodonta) [33]. Obliquely elongate, with projecting beaks. Anterior LV lateral strengthened at right angle along small resilium, posterior laterals unequally remote. *Rec.*, W.Pac.—FIG. E32,5. **T. (T.) sieboldi* ADAMS, Japan; LV int., $\times 4.5$ (Chavan, n).

T. (Pristes) CARPENTER, 1864 (*non Pristis* LATHAM, 1794) [34] [**Pristes oblonga* CARPENTER, 1864; OD] [= *Pristiphora* CARPENTER, 1866 (*non* BLANCHARD, 1835); *Serridens* DALL, 1899]. More transverse than oblique, with scarcely projecting beaks and arcuate anterior, transversely serrated, laterals; duplicate subparallel left posteriors. *Rec.*, E.Pac.—FIG. E32,6. **T. (P.) oblonga* (CARPENTER), USA (Calif.); 6*a,b*, LV hinge and RV int., $\times 9$ (Chavan, n).

Virmysella IREDALE, 1930 [22] [**V. spernax*; M]. Trigono-elliptical, rather flattened, solid, inequilateral, small. Externally sculptured by alternating concentric lines and growths. A deep trigonal socket bordered anteriorly by laminar bent left and hooklike right lateral; posteriorly, by single right and 2 left ones. *Rec.*, Australia-Alaska-N.Eu.—FIG. E31,10. **V. spernax*; 10*a,b*, LV and RV hinges, $\times 6$ (Iredale, 1930).

Family GALEOMMATIDAE Gray, 1840

[*nom. correct.* DALL, 1899 (*pro* Galeommidae GRAY, 1847)]
[= Galeommidae GRAY, 1840]

Shell slightly convex or flattened, commonly transverse, irregular or gaping. Resilium ill-defined, accompanied by marginal ligament. Hinge irregular, with small to quite minute, tuberculiform cardinals, hooking or projecting at termination of flexuous somewhat obscure laterals, and crossing themselves from one valve to another around resilium, in more or less cycloid disposition. *U.Eoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Galeomma*.—2. *Paralepida*.—3. *Amphilepida*.—4. *Galeommella*.—5. *Libratula*.—6. *Ephippodonta*.—7. *Ephippodontoana*.—8. *Ephippodontina*.—9. *Thyreopsis*.—10. *Levanderia*.—11. *Lactemiles*.—12. *Scintilla*.—13. *Cymatioa*.—14. *Ambuscintilla*.—15. ?*Scintillorbis*.—16. *Spaniorinus*.—17. *Scintillona*.—18. *Barclaya*.—19. *Leiochasmea*.—20. *Achasmea*.—21. *Passya*.—22. *Phlyctaenachlamys*.—23. *Turquetia*.—24. *Cyamionema*.—25. *Vasconiella*.—26. *Divariscintilla*. [Insert above, 3*a*. ?*Uncidens*; 9*a*. *Aclistothyra*; 9*b*. *Coleoconcha*; 27. *Tryphomya*.]
Galeomma SOWERBY in TURTON, 1825 [1] [**G. turtoni* SOWERBY, 1825; SD GRAY, 1847] [= *Parth-*

enope SCACCHI, 1833 (type, *P. formosa*; OD); *Hirudinaria* SCACCHI, 1833 (type, *H. alba*; M)]. Irregularly quadrangular-transverse, thin, moderately inequilateral, gaping, externally finely radiated or smooth. Hinge margin straight or nearly so; beaks low with lateral thickenings on both sides, raised up at their infraumbonal junction, there crossed with tubercular very minute cardinal. Resilium inframarginal, oblique, small. *Rec.*, W.Eu.-Medit.

G. (Galeomma). Distorted, ventrally gaping; fine external ribs crenulating inner margin. Long narrow hinge plate with cardinals not distinctly separated from lateral thickenings. *Rec.*, W.Eu.-Medit.-W.Afr.—FIG. E35,4. **G. (G.) turtoni* SOWERBY, Spain; 4*a,b*, LV and RV int., $\times 4.8$ (Chavan, n).

G. (Amphilepida) DALL, 1899 [3] [**G. politum* DESHAYES, 1855; OD]. Transversely subelliptical, smooth, gaping. Dental process forming oblique T on right valve (3*a*, 1, 3*b*) and oblique M on left (2*b*, 2*a*, All), set apart from enlarged laterals. Oblique resilium under sinuous nymph. *Rec.*, Ind.O.—FIG. E35,11. *G. (A.) obockense* JOUSSEAUME, Obock; 11*a,b*, LV and RV int.; 11*c*, RV ext.; all $\times 4$ (114).

G. (Paralepida) DALL, 1899 [2] [**G. formosum* DESHAYES, 1855; OD] [= *Lepirodes* P. FISCHER, 1887 (obj.) (*non Lepyrodes* GUÉNÉE, 1854)]. Largely gaping, with faint radial riblets. Dental process on both sides of ligament, distinctly separated from lateral thickenings. *Rec.*, Australia-Japan.

Aclistothyra MCGINTY, 1955 [9*a*] [**A. atlantica*; OD]. Thin, nearly flat, broadly gaping; transversely semiovate, rounded at both ends. Minute granulations, no radial ornament. Hinge plate straight, slightly thicker medially, without developed teeth. External ligament. *Rec.*, USA (Fla.).
Ambuscintilla IREDALE, 1936 [14] [**A. praemium*; M]. Short, broadly and ovately rounded, somewhat transverse, smooth. Small cardinal developed on each valve and short oblique resilium and nymph. Lateral processes thickened but not separated from margin. *Rec.*, Australia-?Ind.O.—FIG. E35,5. **A. praemium*; LV int., enl. (436).

Barclaya A. ADAMS, 1875 [18] [**Scintilla incerta* DESHAYES, 1863; M] [= *Barclaya* ?AUCT., 1874, Zool. Rec., p. 184; *Barclaya* P. FISCHER, 1887 (*err.*)]. Subtrapezoidal, almost subrectangular, somewhat inequilateral, ventrally moderately rounded. Sculptured by concentric lines and numerous radial riblets. On the left valve, 2 long laterals with sockets above them, subparallel to margin, separated by resilium and 2 minute tuberculiform cardinal teeth. *Rec.*, E.Afr.—FIG. E35, 8. **B. incerta* (DESHAYES), Réunion; 8*a*, LV int., $\times 3.6$; 8*b*, RV ext., $\times 2.1$ (Deshayes, 1863).

Coleoconcha BARNARD, 1964 (9*a*) [**C. opalina*; M].




FIG. E35. Galeommatidae (p. N533, N535-N537).

Shell internal, transversely, rather narrowly semi-ovate, thin, translucent; umbo slightly in front of middle; prodissoconch prominent, subcircular. Concentric grooves only; tiny granules at both ends corrugating lateral surfaces and crenulating margins. Hinge line straight, very slightly thickened, edentulous; no adductors. *Rec.*, S.Afr.

Cymatioa BERRY, 1964 [13] [*pro Crenimargo* BERRY, 1963 (*non* COSSMANN, 1902)] [**Crenimargo electilis* BERRY, 1963; OD]. Shell resembling *Solecardia* in outline but surface microscopically punctate, with several low, ripple-like radial ribs making ventral margin gently crenate; adductor scars high, pallial line interrupted or dotlike, showing through shell; hinge in RV with small pointed cardinal tooth and long posterior lateral, ligament socket large; hinge of LV with thin anterior laterals, 2 small cardinals and pustular *Pl.* *Rec.*, W. C. Am.-Red Sea.—FIG. E36,6. **C. electilis* (BERRY); 6a,b, RV and LV hinges, $\times 8$ (Berry, 1964); 6c, *C. sp.*, Red Sea; LV int., $\times 3.3$ (Chavan, n).

?**Cyamionema** MELVILL & STANDEN, 1914 [24] [**Cyamiun (C.) decoratum*; M]. Papyraceous, quadrangular, upper and ventral margins subparallel; anterior side short and rounded, posterior end obliquely truncated. Lirate on ventral region. Two minute conical right teeth, one more prominent left, and no separated laterals. Ligament external, elongate. *Rec.*, S.Atl. (Falkland Is.).—FIG. E35,1. **C. decoratum*; 1a, RV ext.; 1b,c, LV and RV int., all $\times 2.8$ (Melvill & Standen, 1914). [?Sportellid.]

Ehippodonta TATE, 1889 [**Scintilla? lunata* TATE, 1887; SD MITCHELL, 1890]. Semielliptical, laterally angular, most extended not far from straight hinge margin, ventrally rounded. Externally striated, papillose or reticulated. Ligament bounded on each side by obtuse tooth with bifid crown and long laterals. *Rec.*, Australia.

E. (Ehippodonta) [6]. Almost semielliptical in outline, with symmetric sides. Hinge margin only slightly shorter than oblique diameter. Radial papillae only developed toward external margin. Inner margin plain. *Rec.*, Australia.—FIG. E35, 2. **E. (E.) lunata* (TATE); RV ext., $\times 2.4$ (Tate, 1889).

E. (Ehippodontina) KURODA, 1945 [8] [**E. (E.) murakamii*; OD]. Irregularly semielliptical hinge margin straight, noticeably shorter than angular diameter. Surface finely reticulated, hinge almost edentulous, with well-marked thickened laterals only. *Rec.*, Japan.—FIG. E36,4. **E. (E.) murakamii*; LV int., $\times 3.1$ (Kuroda, 1945).

E. (Ehippodontoana) HABE, 1951 [7] [**E. macdougalli* TATE, 1889; OD]. Almost semicircular, hinge margin diametral. Radial papillae on whole surface, radial riblets indenting ventral margin deeply. *Rec.*, Australia.—FIG. E35,9. **E. (E.)*




FIG. E36. Galeommatidae (p. N535-N536).

macdougalli TATE; 9a,b, LV and RV int.; 9c, RV ext., all $\times 4$ (Chavan, n).

Galeommella HABE, 1958 [4] [**G. utinomii*; OD]. Ellipsoidal, moderately convex, not gaping; beaks small, elevated; radial ribs distinct, laterally weakened and less widely spaced than medially. Inner margin crenulated. *Rec.*, Japan.—FIG. E36,1. **G. utinomii*; RV ext., $\times 5$ (Habe, 1958).

Lactemiles IREDALE, 1931 [11] [**Scintilla strangei* DESHAYES, 1856; OD]. Medium-sized, elliptical, thin, with shining surface ornamented only by faint striae. Tubercular short crossed cardinals: 3a, 1, 3b, forming T; 2a, 2b, making inverted V,

- prolonged by somewhat short lateral, parallel to margin. Oblique nymph and resilial socket. *Rec.*, Australia-Pac.—FIG. E35,13. **L. strangei* (DESHAYES), Pac.; 13a,b, LV and RV int., $\times 2.8$ (Chavan, n). [Beaks prosogyrous.]
- Leiochasmea** DALL, BARTSCH & REHDER, 1938 [**Solecardia* (*Scintilla*) *chascax* PILSBRY, 1921; OD]. Small, elliptical, thin, smooth or with coarse growths. Beaks pointed. Hinge line straight. Two marginal thickenings separated by resilium in trigonal depression and by external support of ligament. *Rec.*, Hawaii.
- L. (Leiochasmea)** [19]. Ventrally gaping. *Rec.*, Hawaii.—FIG. E35,15. **L. (L.) chascax* (PILSBRY); 15a,b, RV int. and LV hinge, $\times 5.1$ (Dall, Bartsch, & Rehder, 1938).
- L. (Achasmea)** [19] DALL, BARTSCH & REHDER, 1938 [**Solecardia* (*Scintilla*) *thaanumi* PILSBRY, 1921; OD]. Without ventral gape. *Rec.*, Hawaii—?Japan.
- Levandieria** STURANY, 1905 [10] [**L. erythraensis*; M]. Narrowly transverse, hinge margin straight, almost diametral; angular at lateral ends, almost straight and somewhat concave ventrally. Externally sculptured with zigzag striae. No teeth developed. *Rec.*, Red Sea.—FIG. E35,6. **L. erythraensis*; 6a,b, both valves int., ext., $\times 1.6$ (Sturany, 1925).
- Libratula** PEASE, 1865 [5] [**L. plana*; M]. Smooth, semilunar, flat, with straight serrate cardinal margin and median ligament. Not gaping. *Rec.*, Pac.
- ?**Passya** DESHAYES, 1858 [21] [**P. eugenii*; OD]. Subquadrangular, flat, very irregular in outline and inequilateral. Beaks small, pointed, anterior side angular, very short; posterior side sloping down, its end strongly acuminate and somewhat rostrate. Ventral margin very convex in middle. Hinge margin thickened in long sinuate curved lamellae. Ligament external. *U.Eoc.*, Eu.(France).—FIG. E35,14. **P. eugenii*; 14a,b, LV int. and RV hinge, $\times 4.8$ (258).
- Phlyctaenachlamys** POPHAM, 1939 [22] [**P. lysiosquillina*; OD]. Very inequilateral, rounded in front, medioposteriorly much acuminate, angular. Regularly sculptured. Hinge with tuberculiform cardinals and elongate lateral. *Rec.*, Australia.—FIG. E36,5. **P. lysiosquillina*; 5a, RV ext., $\times 9$; 5b,c, LV and RV hinges, $\times 16.5$ (Popham, 1928-29).
- Scintilla** DESHAYES, 1856 [12] [**S. philippinensis*; SD STOLICZKA, 1871]. Largely subelliptical, somewhat straightened on hinge margin. Surface granular, nacreous. Sinuous faint anterior and posterior laterals. Two ill-defined unequal cardinals on right valve and one, more or less obscure, on left. ?*Eoc.*, W.Eu.; *Rec.*, Pac.-Ind.O.—FIG. E36, 2. **S. philippinensis*, Penang; 2a,b, LV int. and RV hinge, enl. (Chavan, n).
- Scintillona** FINLAY, 1927 [17] [**Spaniorinus zelandicus* ODHNER, 1924] [=? *Varotoga* IREDALE, 1931 (subj.) (type, *Solecardia cryptozoica* HEDLEY, 1917)]. Transversely ovate, rounded at both ends. Right tubercular prominent small cardinal, left, slightly prominent, oblique one, prolonged into obscure lamella (fitting in socket of opposite valve). Resilium followed by marked nymph. *L.Mio.-Rec.*, Australia-N.Z.—FIG. E36,3. **S. zelandica* (ODHNER), N.Z.; 3a,b, LV hinge and RV int., $\times 4.6$ (Chavan, n).
- ?**Scintillorbis** DALL, 1899 [15] [**Scintilla crispata* P. FISCHER, 1873; OD]. Orbicular, inequilateral, compressed, very thin. Radial and concentric sculpture. External obsolete ligament and stouter resilium. Small tooth on each valve. *Rec.*, Atl.
- Spaniorinus** DALL, 1899 (1900) [16] [**Solecardia* (*S.*) *cossmanni* DALL, 1900; OD]. Transversely subovate, compressed; beaks median; anterior side gradually narrowing, posterior side remaining broad and somewhat truncated. Subvertical tooth in RV, in front of oblique resilium, with traces of shorter teeth on each side; oblique tooth on LV, no distinct laterals on enlarged margins. *Eoc.-Plio.*, W.Eu.-N.Am.—FIG. E35,12. **S. cossmanni* (DALL), USA (Va.); 12a,b, LV hinge and RV int., $\times 3.6$ (Dall, 1900).
- Thyreopsis** A. ADAMS, 1868 [9] [**T. coralliophila*; M]. Trigonal, angular dorsally and at extremities, slightly inequilateral. Valves widely gaping. Hinge apparently edentulous. *Rec.*, Ind.O.(Mauritius).—FIG. E35,3. **T. coralliophila*; 3a,b, LV ext., dorsal, $\times 1.2$ (H. Adams, 1868).
- Tryphomyax** OLSSON, 1961 (27) [**T. lepidiformis*; OD]. Small, thin, flat, subovate, somewhat enlarged posteriorly and with more or less pronounced ventral notch. Sculpture of low radial riblets, cancelled by minute concentrics. RV cardinal knob-like; LV with 2 cardinals, anterior one larger. Internal ligament, in small resilifer. *Rec.*, Panama.
- ?**Turquetia** VÉLAIN, 1877 [23] [**T. fragilis* VÉLAIN, 1876 (1877); M] [= *Turquetia* VÉLAIN, 1876 (*nom. nud.*)]. Convex, very inequilateral, transversely ovate, rounded and enlarged anteriorly, shortened, obliquely truncated and folded posteriorly. Beaks orthogyrous. Hinge margin thickened on anterior elongation, then showing small tubercular tooth separated by 2 sockets. Ligament external. *Rec.*, S.Atl. (St.Paul Is.).—FIG. E35,10. **T. fragilis*; 10a,b, RV ext., int., enl. (Chavan, n). [?Sportellid.]
- ?**Uncidens** COEN, 1934 (3a) [**U. arupinensis*; OD]. Small, thin, slightly convex, equivalve, subequilateral, subelliptical, with slight posterior truncation; no lunule or escutcheon. Sculpture of coarse concentric ribs and faint radial lines. RV hinge with anterior lateral, strong median cardinal with rounded median deep concavity and oblique posterior cardinal; semiinternal ligament. LV with 2 rounded, anterior and median cardinal processes,

located in concavities in front of and in RV tooth; and posterior oblique cardinal. Inner margin smooth. *Rec.*, Adriatic.

Vasconiella DALL, 1899 [**Vasconia jeffreysiana* P. FISCHER, 1873; OD] [*pro Vasconia* FISCHER, 1875, *partim*]. Irregularly rounded, ventrally lobated by deep concave sinuosity. Sculpture of unequal lamellar growths. Small conical tooth under submedian beak, followed by resilium and ligamentary attachment. *Rec.*, Atl.-Medit.-N.Z.

V. (*Vasconiella*) [25]. Slightly oblique, with rather strong rounded tooth. *Rec.*, Atl.-Medit.—FIG. E35,7. **V. (V.) jeffreysiana* (FISCHER), Cape Breton Is.; 7a,b, RV ext., int., $\times 3.2$ (Fischer, 1875).

V. (*Divariscintilla*) POWELL, 1932 [26] [**D. maoria*; OD]. Strongly oblique; with smaller, angular tooth. *Rec.*, N.Z.

GENERA DUBIA

Autonoe LEACH, 1852 (*non* RAFINESQUE, 1815) (*nom. nud.*).

Bilobaria PELSENER, 1911. Shell undescribed. Type lost.

Hyalokellia HABE, 1960 [**H. polita*; OD]. Described as bean-shaped, posteriorly produced; hinge narrow, with RV small cardinal and long posterior lateral, LV very weak anterior cardinal and posterior lateral. Unfigured. ?Erycinid. *Rec.*, Japan.

Soyokellia HABE, 1958 [**S. compressa*; OD]. Unfigured. Said to be oblong and inequilateral, rounded at both ends, hinge with single anterior and 2 posterior teeth on right valve, resilium narrow. ?Erycinid. *Rec.*, Japan.

Superfamily CHLAMYDOCONCHACEA Dall, 1884

[*nom. transl.* KEEN, herein (*ex* Chlamydoconchidae DALL, 1884)] [Materials for this superfamily prepared by MYRA KEEN]

Shells internal, without pallial or adductor scars or hinge teeth; enclosed in two sacs within the mantle, umbones loosely joined by an abortive ligament; prodissoconchs prominent, persistent, at dorsal ends of valves; animal sluglike, ovoid, covered by tough, papillose mantle, foot protruding anteriorly through orifice, a second orifice posteriorly for anal opening. *Rec.*

Family CHLAMYDOCONCHIDAE Dall, 1899

Characters of superfamily. *Rec.*

Chlamydoconcha DALL, 1884 [**C. orcutti*; OD]. Characters of superfamily. *Rec.*, N.Am.—FIG.




FIG. E37. Chlamydoconchidae (p. N537).

E37,1. **C. orcutti*, USA (Calif.); 1a, animal, valve visible dorsally through transparent mantle, $\times 1.5$; 1b-e, shell, showing RV ext., LV int., LV ext., RV int., all $\times 10$ (1a, Williams, 1949; 1b-e, Los Angeles County Museum specimens, Keen, n).

Superfamily CYAMIACEA Philippi, 1845

[*nom. transl.* THIELE, 1934 (*ex* Cyamiidae PHILIPPI, 1845)] [Materials for this superfamily prepared by ANDRÉ CHAVAN]

Shell equivalve, commonly somewhat thickened, mostly with resilium adjacent to nymph, slightly hollowing hinge plate; upper hinge elements distinct, elongate and hooked at their termination above inferior ones; laterals distinct, *AI* (or *I*) and *AII* (or *2*) present. Marine; foot with a byssal gland; two posterior apertures in mantle. *Jur.-Rec.*

The alphabetically arranged generic descriptions in each family-group division of the Cyamiacea are accompanied by numbers inclosed by square brackets. Such numbers indicate position in the sequence of generic taxa given with the respective families or subfamilies for the purpose of recording CHAVAN's arrangement, designed to reflect "natural relationships" of these taxa as inferred by him.

Family CYAMIIDAE Philippi, 1845

[=Perricrinidae MARWICK, 1927]

Shell small, more or less angular, with broad hinge plate scarcely or not at all hollowed by oblique well-defined resilium;




FIG. E38. Cyamiidae (p. N538-N539).

cardinals narrowly elongate, diverging, some (commonly 2b) bilobate; laterals remote, some with taxodont tendency. *Mio.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Cyamium*.—2. *Cyamiomacra*.—3. *Reloncavia*.—4. *Cyamiocardium*.—5. *Perrierina*.—6. *Legrandina*.—7. *Ptychocardia*.—8. *Dicranodesma*.—9. *Lutetina*.

Cyamium PHILIPPI, 1845 [1] [**C. antarcticum*; M]. Transversely elongate, inequilateral, rounded in front, somewhat obliquely extended and faintly truncated backward. Left hinge with remnants of *All*, 2a, bifid 2b, and obsolete 4b; right with remnants of 3a, 1, 3b; posterior laterals remote, more or less indistinct. *Rec.*, S.Am.-Antarctica.—FIG. E38,5. **C. antarcticum*, S.Am.(Patag.); 5a,b, LV hinge, RV int., $\times 5$ (Chavan, n).

Cyamiocardium SOOT-RYEN, 1951 [4] [**Cyamium denticulatum* SMITH, 1907; M]. Rounded, with inflated beaks, radiating sculpture. Hinge as in *Cyamiomacra* but without elongate well-developed laterals and with broader 2a, shorter 2a, and 3b obscurely bifid. Denticulated inner margin. *Rec.*, Antarctica-Chile-?N.Z.-?Australia.—FIG. E39,1. **C. denticulatum* (SMITH); 1a, LV int., $\times 20$; 1b,c, RV and LV hinges, $\times 10$ (858).


Cyamiomacra BERNARD, 1897 [2] [**C. problematica*; M] [= *Heteromacra* LAMY, 1906 (type, *Macra* (H.) *laminifera*)]. Subtrigonal, short, slightly inequilateral; angular in front, enlarged and somewhat obliquely sloping backward; surface smooth. Hinge with 2 well-developed long parallel anterior laterals in each valve, coalescent 3a, 1, 3b, curved *All*, largely bifid 2, 4b, and faint remote posterior lateral; very oblique narrow resilium. ?*Pleist.*, *Rec.*, ?Australia-?N.Z.-Antarctica.—FIG. E38,7. **C. problematica*, Stewart Is.; 7a,b, RV int., LV hinge, $\times 10$ (Chavan, n).

Dicranodesma DALL, 1899 (1900) [8] [**Mysella calvertensis* GLENN in DALL, 1900]. Transversely and very obliquely trigonal, inequilateral, shortly rounded in front, acuminate backward; beaks prosogyrous. Hinge moderately stout, RV with anterior lateral and trigonal 1 having remnants of 3 around its top, LV with oblique inverted V (2-*All*), all in front of oblique resilium, moderately broad, long posterior lateral. *Mio.*, N.Am. (Md.).—FIG. E38,3. **D. calvertensis* (GLENN); 3a,b, RV hinge, LV int., $\times 4$ (Chavan, n).

Legrandina TATE & MAY, 1901 [6] [**L. bernardi*; OD]. Largely ovate, somewhat inequilateral, anteriorly attenuated, posteriorly enlarged; radially ribbed. Hinge with 1, 3a-b, *All* in prolongation of 2, scarcely bifid, 4b and several oblique, subparallel, taxodont-like laterals, RV with only 2 right anterior but 4 posterior ones, LV with 1 to 3 laterals; crenulated margin; resilium in spoonlike pit. *Rec.*, S.Pac.(Tasmania).—FIG. E38,2. **L. bernardi*; 2a,b, LV hinge, RV int., $\times 20$ (Chavan, n).

Lutetina VÉLAIN, 1876 (1877) [9] [**L. antarctica*; OD]. Largely subovate, relatively thick, anterior side shorter and narrower than broadened posterior; ventral margin convex, shining surface with faint concentric ribs; beaks small. Hinge with narrow oblique cardinals and lamellar anterior lateral; *All* 3a obsolete, 1 on inferior margin of plate, 3b, *All*, 2a, 2b well separated, oblique resilium and each valve with strong posterior lateral. *Rec.*, Atl.(St.Paul Is.).—FIG. E38,6; E39,2. **L. antarctica*; E38,6a,b, RV hinge, LV int., enl. (Chavan, after Vélain, 1876); E39,2a,b, LV and RV hinges, $\times 32$ (Bernard).

Perrierina BERNARD, 1897 [5] [**P. taxodonta*; M]. Transversely ovate, very inequilateral, anterior side much shorter than posterior. Hinge with 1, 3a-b, curved *All*, 2a coalescent with 2b, 4b present, and



several oblique, subparallel, taxodont-like laterals; oblique narrow resilium. *Plio.-Rec.*, N.Z.-Antarctica.—FIG. E38,4. **P. taxodonta*, Rec., Stewart Is.; 4a,b, LV and RV hinges, $\times 10$ (Chavan, n). *Ptychocardia* THIELE, 1912 [7] [**P. vanhoeffeni*; OD]. Oblong-angular, convex, with 3 radial folds which undulate margin, also concentric ribbing; prominent prosogyrous beaks. Hinge with lamellar curved *AIII*, and tubercular *AIV*, *AI-1* and *AII-2*, each one in inverted V; rounded deep resilium. Rec., Antarctica.—FIG. E38,1. **P. vanhoeffeni*; 1a-c, RV and LV hinges, LV ext., much enl. (Chavan, n).

Reloncavia SOOT-RYEN, *nom. subst.* herein [3] [*pro Kingiella* SOOT-RYEN, 1957 (*non* SEGUY, 1937)] [**Kingiella chilénica* SOOT-RYEN, 1957; OD]. Ovate, radial sculpture distinct, ventral margin crenate. Outer ligament conspicuous and resilium in oblique groove. Two RV cardinals, 3b grooved; central triangular, not bilobate, 2 and 4a, 4b. Elongate laterals. Rec., S.Am.(Chile). —FIG. E39,3. **K. chilénica*; 3a, LV ext., $\times 0.7$; 3b,c, RV and LV hinges, $\times 24$ (Soot-Ryen, 1959).

Family TURTONIIDAE Clark, 1855

Small, prosogyrous, very inequilateral; with rather narrow hinge plate bearing tubercular cardinals and both anterior and posterior laterals; ligament external. Four mantle folds and no outer demibranchs. *Mio.-Rec.*

Turtonia ALDER, 1848 [**Venus minutus* FABRICIUS, 1780; OD]. Anteriorly short, convex, posteriorly acuminate; LV hinge with subhorizontal *AII* adjacent to feebly bifid tubercular 2, laminar *AIV* and minute *4b* above and behind it; RV with *AI* and *AIII* prolonged by *1* and *AIII-3a* by tubercular *3b*, hinge plate hollowed backward, posterior laterals very distant and minute. *Mio.-Rec.*, N.Eu.-Greenl.-Alaska-Japan.—FIG. E40,3. **T. minuta* (FABRICIUS), Rec., North Sea; 3a,b, RV hinge, LV int., $\times 10$ (Chavan, n).

[This genus has been placed in the Veneracea by K. OCKELMANN (*Ophelia* I, no. 1, p. 121-145, 1964) but, apart of several anatomical differences, its hinge is not cyrenoid (with hollowed plate, *4b* minute, *AIV* developed, and tubercular cardinals).]

Family SPORTELLIDAE Dall, 1899

[=Basterotidae WOODRING, 1925]

Small to medium-sized, shell commonly thickened, more or less rounded in outline; hinge plate broad, scarcely or not at all hollowed by internal part of ligament, which is principally marginal, lying on nymph; cardinals differently developed, *1* trigonal and strong, obliterating *3*, tooth *2* oblique, *4b*

FIG. E39. Cyamiidae (p. N538-N539).




FIG. E40. Turtonidae (3); Sportellidae (1-2,4-9) (p. N539-N541).

obsolete in some, laterals incomplete. *Jur.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Sportella*.—2. *Fabella*.—3. *Vokesella*.—4.

Cerullia.—5. *Hindsia*.—6. *Angusticardo*.—7. *Isoconcha*.—8. *Benthoquetia*.—9. *Lasaeina*.—10. *Grundensia*.—11. *Anisodonta*.—12. *Fulcrella*.—13. *Basterotia*.—14. *Basterotella*.—15. *Ensitellops*.—16. *Hitia*.

Sportella DESHAYES, 1858 [1] [*Psammotea dubia* DESHAYES, 1824; OD]. Largely elliptical, almost equilateral, more or less flattened and thickened, smooth. Hinge with laminar narrow curved *AIII-3a-3b*, latter posteriorly adjacent to strong trihedral *I*; laminar *AI* at inferior angle of trigonal oblique *2a, 2b* at its top and *4b* at right angle, both minute; long callous nymph. *Paleoc.-Plio.*, Eu.-N.Am.

S. (Sportella). Only slightly inequilateral; hinge plate almost unhollowed, with short anterior laterals. *Paleoc.-Mio.*—FIG. E40,2. **S. (S.) dubia* (DESHAYES), M.Eoc.(Lutet.), France; *2a,b*, LV hinge, RV int., $\times 2$ (Chavan, n).

S. (Fabella) CONRAD, 1863 [2] [*Amphidesma constricta* CONRAD, 1841; M]. Inequilateral; hinge plate hollowed, broad, stout, with long subparallel anterior laterals (*AIII, AI, AII*). *Mio.-Plio.*, ?*Rec.*, N.Am.-SW.Eu.-Medit.

Angusticardo COSSMANN, 1887 [6] [*Poromya rotundata* DESHAYES, 1857; SD COSSMANN, 1913]. Short ovate, with distinct *3a* and *2b*, trigonal *I*, moderately strong posterior *4b*, long anterior laterals, short nymph; no ventral sinuosity. *Eoc.*, W. Eu.—FIG. E40,5. **A. rotundata* (DESHAYES), M.Eoc.(Lutet.), France; *5a,b*, LV hinge, RV int., $\times 6$ (Chavan, n).

Anisodonta DESHAYES, 1858 [11] [*A. complanatum*; M]. Subquadrangular, narrowly transverse, anterior side short, attenuated, posterior area long, broad, and vertically truncated; with medioposterior external carina. Hinge with irregular anterior laterals, partly fused to anterior cardinals, trigonal broad *I*, oblique *2*, other ones obsolete, posterior laterals long; with very narrow resilium and short flat broad nymph. *Paleoc.-Rec.*, cosmop.—FIG. E41,3. **A. complanatum*, *Paleoc.*, France; *3a,b*, LV int., RV hinge, $\times 3.7$ (Chavan, n).

Basterotia C. MAYER in HÖRNES, 1859 [*B. corbuloides* HÖRNES; M] [=*Eucharis* RÉCLUZ, 1850 (non LATREILLE, 1804) (type, *Corbula quadrata* HINDS, 1843; OD)]. Strongly convex, somewhat angular, short, subtrapezoidal, very inequilateral, with very prominent beaks; ventral margin convex, slightly concave in middle. Hinge with one projecting cardinal on each valve, 2 hooklike, *I* trigonal; *3a* oblique, faint; with short nymph and enlarged posterior margin. *Mio.-Rec.*, cosmop.

B. (Basterotia) [13]. Dorsally angular. *Mio.-Rec.*, cosmop.—FIG. E40,1. **B. (B.) corbuloides* HÖRNES, Helvet., Aus.; *1a,b*, RV hinge, LV int., enl. (Chavan, n).

B. (Basterotella) OLSSON & HARBISON, 1953 [14] [*Pleurodesma floridana* DALL, 1903; OD]. Scarcely angular, irregularly striated; RV coarsely




FIG. E41. Sportellidae (p. N540-N541).

granular. Hinge with distinct resilium beneath nymph. *Mio.-Rec.*, Eu.-Am.—FIG. E41,4. **B. (B.) floridana* (DALL), *Plio.*, USA (Fla.); RV hinge, $\times 3.3$ (Chavan, n).

Cerullia CHAVAN, 1953 [4] [*pro Scintillula* CERULLI-IRELLI, 1909 (*non* JOUSSEAUME, 1888)] [**Solecardia* (*Scintillula*) *intermedia* CERULLI-IRELLI, 1909; SD CHAVAN, 1953]. Subovate to subelliptical, acuminate forward, slightly truncated backward, somewhat inequilateral. Hinge with long marginal *AIII*, strong oblique curved 1 and 2, others more or less obsolete, moderately short resilium and nymph; faint left posterior lateral. *Paleoc.-Plio.*, ?*Rec.*, Eu.-?N.Am.—FIG. E41,6. **C. intermedia* (CERULLI-IRELLI), *U.Plio.*, Italy; RV int., enl. from $\times 7$ orig. (Chavan, n).

Ensitellops OLSSON & HARBISON, 1953 [15] [**Amphidesma protexta* CONRAD, 1841; OD]. Solenoid, much elongated posteriorly, externally somewhat pustular. Strong curved *All*, 2 and 1 trigonal, elongated posterior lateral. *Mio.-Rec.*, Am.—FIG. E41,2. **E. protexta* (CONRAD), *Mio.*, USA (Va.); 2a,b, LV and RV hinges, $\times 4$ (Chavan, n).

Fulcrella COSSMANN, 1886 [12] [**Poromya paradoxa* DESHAYES, 1857; OD]. Broadly ovate-transverse, slightly inequilateral. Very long regular anterior laterals, fused to anterior cardinals; well-marked narrow posterior right cardinal and resilial hollow under maginal ligament; remote faint posteriors. *Eoc.-Rec.*, Eu.-N.Am.-Pac.—FIG. E40,6. **F. paradoxa* (DESHAYES), *L.Eoc.*, France; 6a,b, RV hinge, LV int., $\times 5.3$ (Chavan, n).

Grundensia KAUTSKY, 1939 [10] [**G. adametzi*; OD]. Subquadrangular, inequilateral, with broad flat hinge, narrowed in its middle, no detached anteriors, *All-2a* bifid, 2b short, 3a, 1, small, obsolete 3b, all small and narrow; long posterior laterals; broad resilium and marginal ligament. *Mio.* (*Helvet.*), Austria.—FIG. E41,1. **G. adametzi*; 1a,b, LV int., RV hinge, enl. (Chavan, n).

Hindsella STOLICZKA, 1871 [5] [*pro Hindsia* DESHAYES, 1858 (*non* H. & A. ADAMS, 1853)] [**Modiola arcuata* LAMARCK, 1807; OD]. Moderately small, inequilateral, transversely elongated, arcuate in middle and rounded at lateral ends. Hinge with long *AIII* and *All* grooves, rounded tuberculiform 1, curved 2; nymph relatively elongate. *Paleoc.-U.Eoc.*—FIG. E40,8. **H. arcuata* (LAMARCK), *M.Eoc.* (Lutet.), France; 8a,b, RV hinge, LV int., $\times 4$ (Chavan, n).

Hitia DALL, BARTSCH & REHDER, 1938 [16] [**H. ovalis*; OD]. Transversely elliptical, inflated, smooth. Knoblike cardinals; with 1, 2a and 2b and 3a and 3b in an inverted V, latter in prolongation of *AIII*, above top of 1; marginal *All* and broad ligament, resilium and nymph; remote faint marginal left lateral. *Rec.*, Hawaii.—FIG. E40,4. **H. ovalis*; 4a,b, RV hinge, LV int., $\times 6$ (Chavan, n).


Isoconcha DAUTZENBERG & FISCHER in PELSENEER, 1911 [**I. sibogai*; SD PRASHAD, 1932]. Transversely elliptical, almost subequilateral, concentrically striated, inflated, small. Hinge with one developed cardinal on each valve and corresponding socket, under beak; ligament external, no distinct resilium. *Rec.*, Australasia.

I. (Isoconcha) [7]. Regularly elliptical, sinuated in middle, subequilateral (755). *Rec.*, Australasia.

I. (Benthonetia) IREDALE, 1930 [8] [**Turquetia integra* HEDLEY, 1907; OD] [= *Austroturquetia* COTTON, 1930]. Irregularly elliptical, somewhat inequilateral. *Rec.*, N.Z.-Australia.—FIG. E41,5. **I. (B.) integra* (HEDLEY); RV int., enl. (Chavan, n). [May be an erycinid with faintly marked laterals.]

Lasaeina COSSMANN, 1910 [1912] [9] [**Lasaea saucatsensis* COSSMANN, 1896; OD]. Ovately rounded, rather broad, smooth. Hinge with 2 cardinals on each valve; 3a faint, 3b strong, *All-2*, 4b. Faint anterior and posterior lateral; narrow resilium, moderately long nymph. *Mio.*, Eu.—FIG. E40,9. **L. saucatsensis* (COSSMANN), Burdigal., France; 9a,b, LV hinge, RV int., $\times 7.4$ (Chavan, n).

Vokesella CHAVAN, 1952 [3] [**V. inopinata*; OD]. Subelliptical, thin, slightly inequilateral, with distinct right anterior *All-1-3b* trigonal, oblique, right-angled *All-2*, a faint 4b; small oblique resilium and prominent long nymph. *Jur.*, Eu. (W. France).—FIG. E40,7. **V. inopinata*; 7a,b, RV int., LV int., $\times 4.7$ (Chavan, n).






FIG. E42. Neoleptonidae (p. N542-N543).

Family NEOLEPTONIDAE Thiele, 1934

Shell minute, rounded, with more or less broad hinge plate, arched under internal ligament. Hinge without completely developed cardinals or with one, but with long laminar anterior and posterior arched, commonly hooked laterals on both sides. *Plio.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Neolepton*.—2. *Neodavisia*.—3. *Notolepton*.—4. *Pachykellya*.—5. *Puyseguria*.—6. *Calvitium*.—7. *Jousseaumiella*.—8. *Epilepton*.—9. *Arculus*.

Neolepton DI MONTEROSATO, 1875 [1] [**Lepton sulcatulum* JEFFREYS, 1859; SD CROSSE, 1884]. Suborbicular, somewhat truncated backward, almost equilateral. RV hinge with well-developed duplicate right anterior and successive posterior laterals, LV hinge with 2-*All* angular, *AIV* marginal, also *PII*. *Plio.-Rec.*, Eu.—FIG. E42,5. **N. sulcatulum* (JEFFREYS), *Rec.*, *Medit.*; 2*a,b*, RV hinge, LV int., enl. (Chavan, n).

Arculus DI MONTEROSATO, 1909 [9] [**Lepton sykesi* CHASTER, 1895; M]. Oval, subrhomboidal, rather convex, inequilateral, concentrically striated. Each

valve with extremely minute erect cardinal and anterior and posterior laterals. *Rec.*, G.Brit. (Guernsey).—FIG. E42,4. **A. sykesi* (CHASTER); 4*a,b*, RV int., LV ext., $\times 18$ (905*a*).

Calvitium LASERON, 1953 [6] [**C. glabra*; OD]. Minute, thin, inequilateral, smooth, with large dome-shaped prodissoconch. Internal ligament; LV with small rounded anteriorly placed cardinal with large fold of hinge, bearing hardly a tooth behind it and prominent lateral, RV with 2 prominent laterals. *Rec.*, Australia.—FIG. E43,3. **C. glabrum*, New S.Wales; 3*a*, RV ext., enl.; 3*b,c*, LV and RV hinges, $\times 12$ (531).

Epilepton DALL, 1899 [8] [**Lepton clarkiae* CLARK, 1852; M]. Broadly ovate, pellucid, inequilateral; anterior side twice length of posterior. Hinge with well-marked laterals, duplicate on RV and single distinct oblique tooth in front of resilium. *Rec.*, Eu. (N. Sea-Medit.).—FIG. E43,2. **E. clarkiae* (CLARK), Eng.; RV int., much enl. (Chavan, n).

Jousseaumiella BOURNE, 1907 [7] [*pro Jousseaumiella* BOURNE, 1906 (non SACCO, 1894)] [**Jousseaumiella heterocyathi* BOURNE, 1906; SD CHAVAN, herein]. Trigonal, inequilateral, concentrically ribbed. Hinge with duplicate laterals and single vertical slightly inequilateral cardinal on RV, 2 narrow cardinals and marginal successive laterals on LV. *Rec.*, India (Andaman I.).—FIG. E43,1. **J. heterocyathi* (BOURNE); 1*a,b*, LV and RV hinges, much enl. (76).

Neodavisia CHAVAN, *nom. subst.*, herein [2] [*pro Davisia* COOPER & PRESTON, 1910 (non DEL GUERCIO, 1909; nec BARNES & MACDUNNOUGH, 1913)] [**Davisia cobbi* COOPER & PRESTON; M]. Rounded, somewhat angular in front, slightly inequilateral. Hinge with strong laterals, duplicate on RV, subparallel, *A1* and *All* continued by short *3a* and 2; resilium small, scarcely oblique, submedian. *Rec.*, SW. Atl. (Falkland Is.).—FIG. E42,6. **N. cobbi* (COOPER & PRESTON); 6*a,b*, LV hinge, RV int., much enl. (Chavan, n).

Notolepton FINLAY, 1927 [3] [**Kellia antipoda* FILHOL, 1880; OD]. Trigonal-rounded, scarcely inequilateral, concentrically striated, with duplicate successive anterior and posterior laterals, *A1* short. ?*Mio.*, *Rec.*, Australasia-S. Atl.—FIG. E42,3. **N. antipodum* (FILHOL); 3*a,b*, LV and RV hinges, much enl. (Chavan, n).

Pachykellya BERNARD, 1897 [4] [**P. edwardsi*; M]. Somewhat obliquely oblong, thick, with prominent beaks and broad hinge, central resilium on both sides of which 2 laterals on each valve are hooked, curved or bifid at their termination. *Pleist.-Rec.*, N.Z. (Stewart Is.).—FIG. E42,2. **P. edwardsi* BERNARD; 4*a,b*, RV hinge, LV int., $\times 33$ (Chavan, n).

Puyseguria POWELL, 1927 [5] [**P. cuneata*; OD]. Very oblique and inequilateral, ovately rounded, anteriorly produced, posteriorly shortened; beaks




FIG. E43. Neoleptonidae (p. N542).

opisthogyrous. Hinge with long anterior laterals and 2 cardinals (2 bifid) of which anterior one is in prolongation of lateral, short posterior laterals. *Plio.-Rec.*, N.Z.—FIG. E42, 1. *P. wanganuica* (POWELL), U.Plio.; 1a, b, LV hinge, RV int., $\times 27$ (Chavan, n).

Superfamily CARDITACEA Fleming, 1820

[*nom. transl.* MENKE, 1830 (*ex* Carditidae FLEMING, 1820)]
[Materials for this superfamily prepared by ANDRÉ CHAVAN]

Shell trigonal to cordiform, trapezoidal, or mytiliform, with radial external sculpture predominating at least locally over concentric and with internal layer of straight radial riblets, which invariably crenulate margin if relatively strong; with more or less marked medioposterior rib or angulation; lunule small, commonly depressed; escutcheon somewhat ill defined; beaks proogyrate, tending to be rounded. Hinge of curved lucinoid type with two unequal teeth (2, 4b) in LV, 3b oblique; 5b very thin, if present; *PI* and *AIV* lacking, other laterals tuberculiform or remote, producing in some shells hinge of cyrenoid aspect; ligament internal or external, latter type inserted on well-marked nymph; shell integripalliate; pedal scars distinct. Animal bysiferous, with open mantle lacking communication between branchial aperture and pedal slitlike outlet; gills large, unequal, united posteriorly. [Marine.] ?*Ord.*, *Dev.-Rec.*

The alphabetically arranged generic descriptions in each family-group division of the Carditacea are accompanied by numbers inclosed by square brackets.

Such numbers indicate position in the sequence of generic taxa given with the respective families or subfamilies for the purpose of recording CHAVAN's arrangement, designed to reflect "natural relationships" of these taxa as inferred by him.

Family PERMOPHORIDAE van de Poel, 1959 (1895)

[*pro* Pleurophoridae DALL, 1895 (Code, 1961, Art. 40)]
[=Kalenteridae MARWICK, 1953; Redoniidae BABIN, 1966]

Medium-sized to large, trapezoidal to modioliform, very inequilateral; radial ribs tending to be obsolete on anterior part of surface, internal margin smooth; with marginal ligament and long nymph. Cardinals partly obsolete, tuberculiform, or much elongated; 5b and anterior laterals lacking in most, posterior laterals remote; that of LV strongest; anterior and pedal scars on thickened buttress. ?*Ord.*, ?*Dev.*, *L.Carb.-U.Cret.*

Subfamily PERMOPHORINAE van de Poel, 1959 (1895)

[*nom. transl.* CHAVAN, herein (*ex* Permophoridae VAN DE POEL, 1959) (=Pleurophoridae DALL, 1895)]

Anterior margin more or less rounded, posterior truncate, with medioposterior angulation and generally several posterior ribs. Hinge with moderately elongate 3b, trigonal 2, narrow 4b appressed in some shells to inferior edge of nymph. ?*Ord.*, ?*Dev.*, *L.Carb.-L.Jur.*

Arrangement of generic taxa by CHAVAN.—1. *Permophorus*.—2. *Curionia*.—3. *Pseudopermophorus*.—4. *Celtoides*.—5. *Triaphorus*.—6. *Kalentera*.—7. *Redonia*.—8. *Neischajewia*.—9. *Rimmyjimina*.—10. *Pleurophorella*.

Permophorus CHAVAN, 1954 [1] [*pro* *Pleurophorus* KING, 1844 (*non* MULSANT, 1842)] [**Arca costata* BROWN, 1841; M]. Subrectangular, solid, with low beaks near anterior end; sculpture rugose concentric, with several posterior radial riblets; lunule and escutcheon well developed. LV hinge with broadly tuberculiform 2, weak, but distinct, 4b and 2 broad, distant, oblique, posterior laterals; RV hinge with 3a as rounded obsolescent tubercle, 3b oblique, strong, sharply defined and posterior lateral submarginal; deep reniform anterior scar in front of myophoric buttress and shallow pedal scar just below 2. *L.Carb.-Perm.*, cosmop.—FIG. E44, 1. *P. albequus* (BEEDE), *L.Perm.*, USA (Tex.); 1a, b, RV int., LV int., $\times 2$, $\times 3$ (665).—FIG. E45, 3. **P. costatus* (BROWN), *Perm.* (Upper Magnesian Ls.), Souter Point, Durham, Eng.; 3a, latex cast int. RV, $\times 3.3$ (Logan, 1964, fig. 8); showing two cardinal teeth; 3b, latex cast LV,

×3.3 (Logan, 1964, fig. 9), showing two cardinal teeth (Newell, n).¹

¹The type specimens of *P. costatus* are lost and there has been uncertainty about the characteristics and limits of

the genus *Permophorus* (Logan, 1964, 548). Unpublished studies by NEWELL suggest that more than one species of *Permophorus* may occur in the British faunas and that the genus is highly variable in American faunas. Fig. E45,3a, above (613.62, Hancock Mus., Newcastle upon Tyne) is here designated as neotype. [NEWELL]




FIG. E44. Permophoridae (Permophorinae) (p. N543-N547).




FIG. E45. Permophoridae (Permophorinae) (p. N543-N545).

Celtoides NEWELL, 1957 [4] [**C. unioniformis*; OD]. Transversely elongate, large, thick-shelled, inequilateral, surface unornamented; lunule small, escutcheon broad. Hinge with broad *3b*, stronger than *2* and bearing radial furrow, both trigonal, *4b* obsolete, *PII* long; with myophoric buttress of moderate size. *Perm.*, N.Am.—FIG. E44,6. **C. unioniformis*, USA (Wyo.); *6a,b*, RV int., LV int., $\times 1$ (669).

Curionia ROSSI RONCHETTI, 1965 [2] [**Myoconcha curionii* HAUER, 1857; OD]. Subovate, very inequilateral, with rounded beaks; lunule broadly developed, covering anterior side of cardinal *2*; ornament of concentric striae. Hinge with trigonal oblique *2* and *3b*; very oblique narrow *4b*, distant posterior laterals, that of RV quite marginal. Ovate anterior scar on buttress, broad posterior scar superficial. *Trias.*(*Carn.-Rhaet.*), Eu.(Italy-Switz.-Ger.).—FIG. E45,2. **C. curionii* (HAUER), *Carn.*, Italy; *2a,b*, RV int., LV int., $\times 2$ (Rossi Ronchetti & Allesinaz, 1965).

Kalentera MARWICK, 1953 [6] [**K. mackayi*; OD]. Transversely suboval, large, beaks near front end; sculpture concentric, posterior radial lines vanishing. Hinge with short conical *2*, deeply oblique *3b*,

curved *4b*, and remote, very strong, broad posterior laterals; anterior adductor on buttress extending backward to hinge, beneath which is very deep pedal scar bounded above by toothlike ridge resembling branch of *3b* on RV. *L.Jur.*, N.Z.—FIG. E46,4. **K. mackayi*; *4a,b*, RV int., LV int., $\times 1$ (599).

Netschajewia YAKOVLEV, 1925 [8] [**Mytilus pallasi* DE VERNEUIL, 1845 (= *Pleurophorus modioliformis* KING, 1844); OD]. Medium-sized, trapezoidal, relatively narrow, convex, front end angular, produced, rear end acuminate; sculpture of irregular growth lines; lunule and escutcheon lacking. Hinge with obscure *3b* and quite remote posterior laterals, stronger on LV. *Perm.*, Eu.-Greenl.—FIG. E44,3. **N. modioliformis* (KING), USSR; *3a,b*, LV int., RV int., $\times 2$ (669).

?*Pleurodapis* CLARKE, 1913 [**P. multincta*; OD]. Elongate ovate, beaks subanterior, umbones low; hinge line straight, ligament external; dentition unknown; surface with strong rounded ridge from beaks to anterior margin, ending in marginal notch; 4 or 5 divergent radial ridges of varying strength from beaks to posterior margin, delimiting shallow indentations of shell margin. *Dev.*, S.Am.




FIG. E46. Permophoridae (Permophorinae) (4); (Myoconchinae) (1-3,5) (p. N545, N547-N548).

(Ponta Grossa & Santa Cruz, Brazil).—FIG. E44, 7. **P. multincta*, Ponta Grossa, Brazil; $\times 1$ (138). [L'AROCQUE]

?*Pleurophorella* GIRTY, 1904 [10] [**P. papillosa*; OD]. Externally like *Permophorus* but somewhat less inequilateral, thin, ornamented only by minute papillae and weak concentric lines; lunule and escutcheon sharply defined. Interior unknown. *U. Penn.*, N.Am. (Tex.).

Pseudopermophorus CIRIACKS, 1963 [3] [**P. annettae*; OD] [?=*Protrete* GIRTY, 1908 (type, *P. texana*; OD)]. Shell heavy, slightly inequivalve, RV more convex than LV, subquadrate, tapering slightly posteriorly; ventral margin gently indented medially at the termination of a broad medial sulcus; posterior and anterior margins broadly rounded; escutcheon deep, elongate, more prominent on the RV; lunule deeply invaginated as an ovoid cavity (brood pouch?) immediately above the anterior adductor; shell surface ornamented with coarse, rounded concentric fila, devoid of radial ornamentation; dentition 3*b*, (5*b*), (PIII)/2, 4*b*, PII with 5*b* barely distinguishable from the ligament nymph. *Perm.* (Park City), USA (Mont.)-Japan.—FIG. E45, 1. **P. annettae*; 1*a,b*, LV int., RV int., $\times 2$; 1*c*, LV int., $\times 1$ (132). [NEWELL]

?*Redonia* ROUAULT, 1851 [7] [**R. deshayesiana*; SD P. FISCHER, 1886]. Subelliptical, with prominent beaks relatively far forward. Hinge with arcuate subhorizontal 2, subelliptical 3*b*, very long curved and transversely striated, 4*b* and 5*b* parallel to margin; posterior laterals fused together, anterior adductor on broad buttress. [Doubtfully belongs in Modiomorphidae.] *Ord.*, Eu. (France).—FIG. E44, 4. **R. deshayesiana*, Brittany; 4*a,b*, LV int., RV int., $\times 1$; 4*c,d*, LV and RV hinges, enl. (Chavan, n).

Rimmyjimina CHRONIC, 1952 [7] [**R. arcula*; OD]. Small, trapezoidal, relatively broad, convex, front more or less rounded, rear truncate; sculpture of irregular growth lines; lunule small, oval, well defined; narrow escutcheon bearing fine ligament groove. Hinge with narrow 3*b*, posterior LV lateral long and broad. *Perm.*, N.Am.-Eu. (Aus.).—FIG. E44, 2. **R. arcula*, USA (Ariz.); 2*a,b*, RV int., LV int., $\times 5$; 2*c,d*, RV and LV hinges, $\times 3.5$ (669).

Triaphorus MARWICK, 1953 [3] [**Pleurophorus zelandicus* TRECHMANN, 1918; OD]. Transversely oval, prosogyrous beaks almost terminal; irregular concentric sculpture, radial ridges only on posterior surface; lunule deep, escutcheon well de-

fined. Hinge with very weak oblique *3a*, strong oblique 2 and *3b*, long, weak *4b* welded to long nymph; strong posterior laterals remote; anterior adductor on buttress below anterior cardinals, shallow pedal scar on their lower side. *U.Trias.*, N.Z.—FIG. E44,5. **T. zelandicus* (TRECHMANN), Otamitan; *5a-c*, RV ext., LV ext., RV hinge, $\times 1$ (599).

Subfamily MYOCONCHINAE Newell, 1957

[*nom. transl.* CHAVAN, herein (*ex* Myoconchidae NEWELL, 1957)]

Acuminate anteriorly, regularly enlarged posteriorly, mytili- to modioliform, angulated posterodorsally, surface with many fine radial threads, at least toward rear. Hinge with elongate *3b*, extremely narrow, linear 2 and *4b*, rather faint, *4b* fused with lower edge of nymph. ?*M.Dev.*, *Perm.-U.Cret.*

Arrangement of generic taxa by CHAVAN.—1. *Myoconcha*.—2. *Modiolina*.—3. *Daharina*.—4. *Pseudomyoconcha*.—5. *Stuichburia*.—6. *Pleurophorina*. [Insert above, 4a. ?*Pseudosanguinolites*.]

Myoconcha J. DE C. SOWERBY, 1824 [**M. crassa*; M]. Modioliform, with long, narrow submarginal angulation bordered by furrow, surface with very fine numerous riblets. Cardinals 2 and *4b* thin, *3b* stronger, *Pll* well developed, behind long nymph; myophoric buttress broad. ?*Perm.*, *L.Jur.* (*Lias.*)-*U.Cret.* (*Senon.*), cosmop.

M. (Myoconcha) [1] [= ?*Labayophorus* LIKHAREV, 1939 (type, *L. magnus*; M)]. Large, with terminal beaks, ribs vanishing toward anterior side. Cardinals nearly straight, those of LV very thin, low, *3b* generally trigonal; myophoric buttress low, moderate in extent. ?*Perm.*, *L.Jur.* (*Lias.*)-*U.Cret.* (*Senon.*), cosmop.—FIG. E46,2. **M. (M.) crassa*, *M.Jur.* (*Bajoc.*), France; *2a-c*, RV ext., hinge, LV hinge, $\times 1$ (Chavan, n).

M. (Modiolina) J. MÜLLER, 1851 [2] [**M. bosqueti* (= *Lithodomus discrepans* MÜLLER, 1847); OD]. Small, beaks not quite terminal, whole surface with fine radial and concentric riblets. Cardinals thin, 2 sinuate, *3b* somewhat curved; nymph relatively narrow; myophoric buttress broad. *U.Cret.* (*Senon.*), Eu.—FIG. E46,1. **M. (M.) discrepans* (MÜLLER), Campan., Neth.; *1a,b*, RV and LV hinges, $\times 1$ (Chavan, n).

Daharina DUBAR, 1948, p. 170 [3] [**Myoconcha (Daharina) gentili*; M]. Large, gibbose, thick-shelled, oval, obliquely elongated, smooth, with anterior, terminal beaks; posterodorsal region rather flattened, winglike; anteroventral margins with byssal gape; umbonal angle occupied in each valve by broad, thick hinge plate, posterior part of which bears in RV broad, elongate cardinal tooth, received in socket adjacent to nymph in





FIG. E47. Permophoridae (Myoconchinae) (p. N547).

LV; no lateral teeth; anterior adductor scar deep, on shell wall below hinge plate. *L.Jur.* (*U.Pliensb.*), Afr. (Morocco).—FIG. E47,1. **D. gentili*; *1a,b*, LV ext., LV int. (broken post. region restored), $\times 0.4$ (Dubar, 1948). [Cox]

?*Modiella* HALL, 1883, p. 4 [**Pterinea pygmaea* CONRAD, 1842; M]. Similar externally to *Myoconcha* SOWERBY (1824), *Modiolina* MÜLLER (1851), and *Netschajewia* YAKOVLEV (1925); hinge details unknown. *M.Dev.*, USA (N.Y.). [NEWELL]

?*Pleurophorina* LIKHAREV, 1925 [6] [**Pleurophorus simplex* LIKHAREV, 1925 (= *Modiola simplex* KEYSERLING, 1846); M]. Externally like *Permophorus* but reported to lack cardinal 2. *Perm.*, USSR.

Pseudomyoconcha ROSSI RONCHETTI, 1966 (1967) [4] [**Myoconcha lombardica* HAUER, 1857; OD].




FIG. E48. Carditidae (Carditinae) (p. N548).

Very inequilateral, modioliform, more or less sinuate ventrally, strongly convex dorsally. Terminal curved beaks. External concentric irregular growths and radial lines. Broad anterior buttress. LV with 2, *P*II, no developed 4*b*; RV with 3*b*, well marked *P*III. *M.Trias.-U.Trias.*, Eu.(Italy-Ger.-Hung.)-Japan.—FIG. E46,5. **P. lombardica* (HAUER), *U.Trias.*, Italy; 5*a,b*, LV, RV int., $\times 1$ (Rossi Ronchetti & Allasinaz, 1966).

?*Pseudosanguinolites* PATTE, 1929 (4*a*) [**P. douvillei*; M]. Transversely much elongated and curved, anterior side angular, posterior side more or less rounded; shell thick, with almost terminal, anterior beaks. Trapezoidal striated area above hinge plate, which in RV bears long posterior cardinal parallel to upper margin; external long narrow ligament; anterior part of hinge unknown, but elliptical anterior scar and ogival posterior one can be seen. *Dev.*, IndoChina.

?*Stahlia* E. FISCHER, 1915, p. 219 [**S. persica*; M]. Rectangular, elongate, not oblique; umbones terminal; evenly and rather strongly inflated; ornament of radial ribs, most prominent on dorsal part of shell, and concentric folds; anterior adductor scar moderately large, below beak; dentition unknown; said to gape posteriorly, but appearance of gape possibly due to imperfect preservation of type. [Possibly synonym of *Myoconcha*.] *L.Jur.(Toarc.)*, SW.Asia(Iran). [Cox]

Stutchburia ETHERIDGE, JR., 1900 [5] [**Orthonota? costata* MORRIS, 1845; OD]. Large, transversely subquadrate, ornamented posteriorly by coarse ribs; lunule and escutcheon developed. Cardinals 2 and 3*b* obsolete, likewise laterals *PI* and *P*III but *P*II well marked. *Perm.*, cosmop.—FIG. E46, 3. **S. costata* (MORRIS), Artinsk., Australia; RV ext., $\times 1$ (669).

Family CARDITIDAE Fleming, 1828

[*nom. correct.* COSSMANN, 1914 (*pro* Carditidae FLEMING, 1828)]

Small to large, trapezoidal or rounded, with strong radial ribs and shell margin invariably crenulated internally; ligament external. Hinge with faint 3*a* and thin 5*b*, anterior laterals tuberculiform; anterior scars set on platform. *Dev.-Rec.*

Subfamily CARDITINAE Fleming, 1828

[*nom. transl.* CHAVAN, herein (*ex* Carditidae FLEMING, 1828)]

Mytiliform, with long 3*b* and obsolete laterals. *Paleoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1.

Cardita.—2. *Jesonia*.—3. *Beguina*.

Cardita BRUGUIÈRE, 1792 [**Chama calyculata* LINNÉ, 1758; SD GRAY, 1847].¹ Transversely inequilateral, trapezoidal or modioliform, with nodulose radial ribs. Hinge with obliquely trigonal divergent cardinals in LV and faint anterior laterals. *Paleoc.-Rec.*, cosmop.

C. (Cardita) [1] [= *Arcinella* OKEN, 1815 (rejected by ICZN, 1956, Opinion 417, as non-binominal) (*non* SCHUMACHER, 1817) (obj.); *Mytilocardita* ANTON, 1839 (obj.)]. Relatively small, short; lunule not depressed. Cardinal 3*b* elongated backward-inward. *Paleoc.-Rec.*, Eu.-Afr.-Asia-Australia.—FIG. E48,1. **C. (C.) calyculata* (LINNÉ), *Rec.*, *Medit.*; 1*a-c*, RV ext., RV int., LV hinge, $\times 1$ (7; Chavan, n).

C. (Jesonia) GRAY, 1847 [2] [**Perna jeson* ADANSON, 1757, p. 217 (invalid, pre-Linnaean) (= *Cardita senegalensis* REEVE, 1843); OD] [= *Mytilicardes* DE BLAINVILLE, 1824 (vernacular); *Jesonia* GRAY, 1840 (*nom. nud.*); *Mytilocardia* AGASSIZ, 1847 (obj.); *Mytilocardia* TRYON, 1872 (obj.)]. Medium-sized, notably elongated, with commonly squamose ribs. Cardinal 3*b* scaleniform, prolonged only backward, *AI* lacking but posterior laterals moderately distinct; lunule sunken. *Oligo.-Rec.*, cosmop.

Beguina RÖDING, 1798 [3] [**B. nephritica* (= *Chama phrenetica* BORN, 1780, = **C. semi-orbiculata* LINNÉ, 1758); M] [= *Azarella* GRAY, 1854]. Broad modioliform, compressed, with fine intersecting growth lines and disparate radial riblets; beaks stretched out to cover anterior part of hinge. Posterior cardinals much elongated, laterals wanting. *Rec.*, Afr.-Asia-Polynesia.—FIG. E48,2. **B. semi-orbiculata* (LINNÉ), Ind.O.; 2*a,b*, RV ext., int., $\times 0.8$ (124*b*).

Subfamily CARDITAMERINAE Chavan, new subfamily

More or less trigonal or transversely trapezoidal, with strong radial ribs. Cardinal 3*a*

¹ According to International Code (Art. 69.a.iii), type species is *Cardita variegata* BRUGUIÈRE, 1792; SD FLEMING, 1818. In opinion of CHAVAN, FLEMING's designation was not explicit and therefore not to be recognized.—Ed.

present, *3b* V-shaped, laterals fairly well developed. *U.Trias.*(*Carn.*)-*Rec.*




FIG. E49. Carditidae (Carditamerinae) (p. N549-N551).

Arrangement of generic taxa by CHAVAN.—1. *Carditamera*.—2. *Lazariella*.—3. *Glans*.—4. *Centrocardita*.—5. *Goossensia*.—6. *Carditella*.—7. *Carditellona*.—8. *Cyclocardia*.—9. *Scalaricardia*.—10. *Plionema*.—11. *Vimentum*.—12. *Vetericardiella*.—13. *Miodontiscus*.—14. *Tutcheria*.—15. *Choniocardia*.—16. *Carditellopsis*.—17. *Arcturellina*.—18. *Izumicardia*.—19. *Pleuromeris*.—20. *Cossmannella*.—21. *Cardiocardita*.—22. *Bathycardita*. [Insert above, 11a. *Fenestricardita*; 20a. *Cretocardia*.]

Carditamera CONRAD, 1838 [**Cypricardia arata* CONRAD, 1832; OD]. Transversely subrectangular to trapezoidal, solid, somewhat compressed, with subparallel dorsal and ventral margins; sculpture of flabellate ribs, posterior ones unequal; lunule oblique. Cardinal teeth strong, laterals (except *AIII*, *AIV*, and in some shells *PIV*) well defined. *U.Eoc.-Rec.*, *W.Eu.-Afr.-N.Am.*

C. (*Carditamera*) [1] [= *Lazaria* GRAY, 1854 (type, *Cardita radiata* SOWERBY, 1833; SD DALL, 1903)]. Elongate, posterior margin rounded or with slight straight truncation; ribs rugose. Hinge with faint *PIV*. *U.Eoc.-Rec.*, *N.Am.-W.Eu.-?E. Afr.*—FIG. E49,6a,b. **C.* (*C.*) *arata arata* (CONRAD), *Mio.*, *USA(N.Car.)*; 6a,b, *RV ext.*, *LV int.*, $\times 1$ (Conrad, 1839).—FIG. E49,6c. *C.* (*C.*) *arata verdevilla* GARDNER, *Mio.*, *USA(N.Car.)*; *RV int.*, $\times 0.8$ (Gardner, 1943). [= *Byssomera* OLSSON, 1961 (type, *Cardita affinis* SOWERBY, 1832; OD).]

C. (*Lazariella*) SACCO, 1899 [2] [**Cardita subalpina* MICHELOTTI, 1839; OD]. Shorter than *C.* (*Carditamera*), with concave anal truncation, more widely spaced ribs which are subquadrate and strong. Cardinals *3a* and *3b* scaleniform, laterals *A1* and *P111* obsolete. *L.Mio.(Aquitain.)-Rec.*, *SW.Eu.-W.Afr.*—FIG. E49,1. *C.* (*L.*) *hippopaea* (BASTEROT), *Aquitain.*, *S.France*; 1a-c, *RV ext.*, *int.*, *LV int.*, $\times 1$ (165).

Arcturellina CHAVAN, 1951 [17] [*nom. subst. pro Arcturella* CHAVAN, 1941 (*non* SARS, 1897)] [**Venericardia asperula* DESHAYES, 1825; OD]. Small to medium-sized, inequilateral, trapezoidal to rounded, moderately convex; with regularly striate or squamose radial ribs; beaks small, rounded; lunule relatively convex and long, bounded forward by oblique furrow. Hinge with very thin *3a*, almost obsolete, and trigonal to subrectangular *3b* with subvertical anterior face and backward extension; *A11* linear, *P11* and *P111* faint; shell margin deeply indented internally by ends of ribs. *Paleoc.-Rec.*, *Eu.-Afr.-S.Am.-Australia*.—FIG. E49,3. **A. asperula* (DESHAYES), *M.Eoc.* (Lutet.), *France(Paris basin)*; 3a, *LV ext.*, $\times 1$ (Deshayes, 1837); 3b,c, *LV and RV hinges*, ca. $\times 1.8$ (101).

Carditella E. A. SMITH, 1881 [**C. pallida*; SD DALL, 1903]. Small, with flabellate ribs crossed

by concentric growth lines; lunule elongate; ligament small, external, adjacent to restricted resilium just behind beak and slightly narrowing tops of 3*b* and 4*b*. Each valve with 2 unequal cardinals




FIG. E50. Carditidae (Carditamerinae) (p. N550-N551).

and 2 remote, well-defined laterals. *Plio.-Rec.*, S. Am. - Australia - E. Asia (Japan - Formosa)-S. Atl. (Tristan da Cunha).

C. (Carditella) [6]. Slightly inequilateral, mostly trigonal, with broad, close-spaced ribs. Cardinal 3*a* fused to lunular margin. *Rec.*, S.Am.-Japan. —FIG. E49,5. **C. (C.) pallida*, S.Am. (Port Rosario); 5*a-c*, LV ext., int., RV int., ca. $\times 4$ (Smith, 1881).

C. (Carditellona) IREDALE, 1936 [7] [**C. angasi* E. A. SMITH, 1885; OD]. Very inequilateral, obliquely rounded, with well-spaced squamose ribs. Hinge with 3*a* more distinct and *Al* shorter than in *C. (Carditella)*. *Plio.-Rec.*, Australia-W. Pac.O.—FIG. E49,2. **C. (C.) angasi* SMITH, Australia (Port Jackson); RV ext., much enl. (852).

Cardiocardita ANTON, 1839 [**Chama ajar* ADANSON, 1757 (invalid, pre-Linnaean) (= *Cardita ajar* BRUGUIÈRE, 1792); SD HERRMANNSEN, 1846]. Inequilateral, subtrapezoidal, rounded in front, obliquely truncate at rear; with strong nodular or echinate ribs; beaks low. Hinge with trihedral cardinals, 3*a* very small, 3*b* straight, sharp-sided, at mid-hinge; anterior laterals minute, almost obsolete. *M.Eoc.-Rec.*, SW.Eu.-Afr.-Pac.O.-Australia.

C. (Cardiocardita) [21] [= *Cardiocardites* DE BLAINVILLE, 1825 (vernacular); *Agaria* GRAY, 1840 (obj.); *Azaria* TRYON, 1872 (*nom. null.*); *Divergidens* EAMES, 1957 (type, *Cardita tripartitocostata* NEWTON, 1922; OD)]. Anteriorly angular, ribs commonly well spaced and sharply defined; lunule slightly depressed. *M.Eoc.-Rec.*, SW. Eu.-Afr.-Pac.O.-N.Z.—FIG. E50,4. **C. (C.) ajar* (BRUGUIÈRE), *Rec.*, W.Afr.(Senegal); 4*a,b*, RV ext., int., $\times 1$ (7).

C. (Bathycardita) IREDALE, 1925 [22] [**Cardita raouli* ANGAS, 1872; OD]. More rounded than *C. (Cardiocardita)* and with less sharply delimited, spinose ribs; lunule deep. *Rec.*, Australia. —FIG. E50,5. **C. (B.) raouli* (ANGAS), Tasmania; 5*a,b*, RV ext., int., $\times 1$ (431).

Choniocardia COSSMANN, 1904 [**Venericardia (C.) oppenheimi*; OD]. Relatively small, depressed, with reticulate or striate ribs; lunule broad, well defined, flat and somewhat depressed. Hinge with nearly symmetrical LV cardinals, 3*b* straight trigonal, 3*a* oblique; anterior and posterior laterals present. *L.Eoc.-Rec.*, Eu.-Australia-E.Asia (Japan).

C. (Choniocardia) [15]. Rounded trigonal to subtrapezoidal, truncate posteriorly; with finely reticulate ribs; lunule oblique, sunken. *L.Eoc.-U.Eoc.*, W.Eu.—FIG. E50,6. **C. (C.) oppenheimi* (COSSMANN), *M.Eoc.*(Lutet.), W.France; RV int., $\times 3$ (Chavan, n).

C. (Carditellopsis) IREDALE, 1936 [16] [**Carditella elegantula* TATE & MAY, 1901; OD]. Rounded trigonal, solid, slightly inequilateral, tending

- to be slightly oblong with front side produced and no definite anterior lamellae; with broad, flattened ribs crossed by concentric striae; lunule not depressed. *M.Eoc.-Rec.*, W.Eu.-W.Pac.O.-Australia.—FIG. E49,4. **C. (C.) elegantula* (TATE & MAY), *Rec.*, Australia; 4*a,b*, RV ext., LV int., much enl. (Tate & May, 1901).
- Cossmannella** MAYER-EYMAR, 1896 [20] [**Cardita (C.) aegyptiaca* FRAAS" of MAYER-EYMAR, 1896 (= *Cardium aegyptiacum* MAYER, 1896, non FRAAS, 1867; = *Cardita fajumensis* OPPENHEIM, 1903); OD] [= *Ameqiglanis* EAMES, 1957 (type, *Cardita costaenodulosis* NEWTON, 1922; OD)]. Transversely trigonal, subrounded in front, acuminate in rear; with flat ribs separated by narrow interspaces. Hinge with 3*b* nearly isosceles triangular, 2 and 4*b* subsymmetrical, anterior RV laterals obsolete. *All* and remote posterior lamellae well marked. *Paleoc.-L.Mio.*, Afr.-Asia (India).—FIG. E50,3. *C. costaenodulosis* NEWTON, *M.Eoc.*, W.Afr. (Nigeria); 3*a-c*, RV ext., int., LV int., ×2 (99).
- Cretocardia** CONRAD, 1877 (20a) [**Cardita jaquinoti* STOLICZKA, 1871 (= ?*Cardium jaquinoti* D'ORBIGNY, 1847) (= *Cardita orbicularis* FORBES, 1846, non SOWERBY, 1825); OD]. Subquadrangular, inflated, very inequilateral, with tripartite somewhat tubercular ribs separated by deep trough. Diverging thick 3*a* and 3*b*, 2 small, tubercular and median, 4*b* very long, rather thin, curved; minute tuberculiform laterals. *U.Cret.*, India.
- Cyclocardia** CONRAD, 1867 [**Cardita borealis* CONRAD, 1831; SD STOLICZKA, 1871]. Subtrigonal or short trapezoidal to cordiform, thickened or somewhat compressed, ventral margin well rounded; regular radial ribs may be closely spaced, crossed by numerous equidistant growth lines; beaks very small, tending to be erect. Hinge with very faint *All* and *PIII*, other laterals virtually obsolete; cardinals strong, 3*b* straight in front, 2 straight, equilateral. *U.Cret. (Cenoman.)-Rec.*, cosmop.
- C. (Cyclocardia)** [8] [= *Arcturus* HUMPHREY in GRAY, 1839 (non BERTHOLD in LATREILLE, 1827; nec CURTIS, 1830) (obj.); *Bendeglans* EAMES, 1957 (type, *Cardita costaeirregularis* NEWTON, 1922; OD)]. Juvenile-stage ribs narrow, rounded, equally perlate, later becoming flattened and broadened, crossed by close-spaced growth lines; lunule large, smooth, slightly convex. *M.Eoc.-Rec.*, cosmop.—FIG. E50,1*a*. **C. (C.) borealis* (CONRAD), *Rec.*, USA (Mass.); LV ext., ×0.8 (Gould, 1841).—FIG. E50,1*b,c*. *C. (C.) granulata* (SAY), *Mio.*, USA (Md.); 1*b,c*, LV and RV hinges, enl. (101). [= *Strophocardia* OLSSON, 1961 (type, *Venericardia megastrophia* GRAY, 1825; OD).]
- C. (Plionema)** CONRAD, 1872 [10] [**Astarte guerangeri* D'ORBIGNY, 1843; M]. Small, fine radial ribs very numerous; lacking distinct lunule. Hinge with 2 and 4*b* almost equally strong and oblique, 2*b* short; laterals wanting. *U.Cret. (Cenoman.)*, Eu.—FIG. E50,2. **C. (P.) guerangeri* (D'ORBIGNY), W.France; 2*a,b*, LV ext., int., enl. (695).
- C. (Scalaricardita)** SACCO, 1899 [9] [**Miodon (S.) scalaris* (= *Venericardia scalaris* J. DE C. SOWERBY, 1825); OD]. Ribs everywhere closely spaced, in young stage flat and regularly incised by furrows which may be slightly oblique; lunule elongate, flat, ill-defined. Hinge with remote, faint anterior laterals and longer, linear posterior laterals; 3*b* trigonal-rectangular, 2 vertical, 4*b* oblique. *Oligo.-Neog.*, Eu. (Aus.).
- C. (Vimentum)** IREDALE, 1925 [11] [**Cardita dilecta* E. A. SMITH, 1885; OD]. Transversely ovate, numerous rounded regular ribs crossed by very close-spaced concentric riblets; lunule cordiform, relatively elongate. Cardinal 3*b* conical. *Plio.-Rec.*, Eu. (W. France) - Australia.—FIG. E49,7. **C. (V.) dilecta* (SMITH), *Rec.*, Australia (Bass Str.); 7*a,b*, LV lat., dorsal, enl. (852).
- Fenestricardita** CASEY, 1961 [11a] [**Venus? fenestrata* FORBES, 1845; OD]. Transversely trapezoidal, posteriorly elongated. Regularly reticulated by narrow concentric and squamose radial ribs. Hinge with marginal 3*a*, high trigonal 3*b*, oblique 5*b*, narrow subvertical 2 and strong sharp, oblique 4*b*; marginal *All*, long well-marked *PIII*; depressed ligament. *L.Cret.-M.Cret.*, W.Eu.
- Glans** MEGERLE, 1811 [**Chama trapezia* LINNÉ, 1767; SD HERRMANNSEN, 1846]. Quadrangular trapeziform, convex, inequilateral, with nodulose or squamose ribs; lunule with convex margin. Hinge as in *Carditamera* but *AIII* and *PII* faint and *PIII* less developed. Shell with well-marked medioposterior dorsal angulation. *Paleoc.-Rec.*, Eu.-Asia-N.Afr.-E.Afr.-Australia-C.Am.
- G. (Glans)** [3]. Truncate posteriorly, with nodulose or granular rounded ribs, posterior ones unequal in some species. Lateral teeth well developed. *Paleoc.-Rec.*, Eu.-Asia-N.Afr.-E.Afr.-C.Am.—FIG. E51,5. **G. (G.) trapezia* (LINNÉ); *Rec.*, *Medit.*; 5*a-c*, RV ext., RV and LV hinges, ×3 (89*a*, 101).
- G. (Centrocardia)** SACCO, 1899 [4] [**G.? aculeata* (= *Chama aculeata* POLI, 1795); OD]. Posterior margin rounded, with equal squamose or echinate ribs. Hinge with almost obsolete small laterals. *U.Eoc.-Rec.*, Eu.-E.Afr.-Australia.—FIG. E51,1. *G. (C.) aequicostata* (COSSMANN), *U.Eoc.*, France (Paris basin); 1*a,b*, RV ext., int., ×1 (160).
- Goossensia** COSSMANN, 1885 [5] [**G. plicatuloides* (= *Cardita irregularis* DESHAYES, 1860); M]. Relatively small, subtrapeziform, sinuate, very inequilateral, with concentric growth lines intersecting radial ribs which commonly are squamose; lunule depressed. Cardinals and laterals present, 3*b* trigonal; shell margin internally irregularly denticulate. *Paleoc.-Eoc.*, W.Eu.—FIG. E51,7. **G. irregularis* (DESHAYES), *M.Eoc. (Lutet.)*, France




FIG. E51. Carditidae (Carditamerinae) (p. N551-N553).

(Paris basin); 7a,b, RV ext., int., $\times 3$ (259); 7c, LV int., $\times 2.5$ (160).

Venericardia ICHIKAWA in ICHIKAWA & MAEDA, 1963, p. 119 [18] [**I. parva* ICHIKAWA & MAEDA, 1963; OD]. Moderately small, roundly subquadrate, with prominent prosogyrous beaks; surface with strong, regular, approximate radial ribs. Ventral hinge margin straight, hinge with short, trigonal 3b, lamellar 3a and 5b, AII horizontal, long and straight, reaching lower anterior angle of trigonal 2b, AI horizontal, then ascending obliquely beneath lunular margin, 4b present; posterior laterals tuberculiform. *U.Cret.(U.Senon.)*, Japan.—FIG. E52,1. **I. parva* ICHIKAWA & MAEDA, Campan.; 1a,b, LV int., RV int., $\times 1.5$ (426).

Miodontiscus DALL, 1903 [13] [*nom. subst. pro Miodon* CARPENTER, 1863 (non DUMÉRIL, 1859; nec SANDBERGER, 1870)] [**Miodon prolongatus* CARPENTER, 1863; OD]. Small, obliquely oblong, with strongly prosogyrate beaks and long, ill-defined lunule; sculpture of broad radial ribs crossed by concentric furrows. Cardinals oblique, scalariform, 2 short and subparallel to 4b, which is much elongated, like 3b; anterior laterals obsolete, posteriors very faint. *Plio.-Rec.*, W.N.Am.-Japan.—FIG. E51,2. **M. prolongatus* (CARPENTER), *Rec.*, USA(Calif.); 2a-c, LV ext., int., RV hinge, $\times 4$ (Keen, 1939).

Pleuromeris CONRAD, 1867 [19] [**P. decemcostata* (= **Venericardia tridentata* SAY, 1826, *decemcostata* CONRAD, 1867; non *Cardita tridentata* REEVE, 1843); OD] [= *?Cycloglans* GORODISKI & FRENEIX, 1959 (1960) (type, *Glans (C.) schencki* FRENEIX ex GORODISKI & FRENEIX, MS; OD)]. Rounded trigonal, small, solid, relatively inflated, with erect beaks; radial ribs granular to squamose, large, tending to square transverse profile; lunule elongate, depressed or slightly convex. Hinge with long, partly undetached 3a, trigonal 3b, and short, stout 2 and 4b; anterior laterals narrow but well defined AIV lacking, PII, PIII, PIV present. *?Eoc.*, *Mio.-Rec.*, N.Am.-Australia-?C.Afr.—FIG. E51,3. **P. tridentata decemcostata* CONRAD, *Mio.*, USA (N.Car.); 3a-c, LV ext., RV int., LV int., $\times 2.2$ (Gardner, 1943).

Tutcheria COX, 1946 [14] [**Cardium submulticostatum* D'ORBIGNY, 1850 (pro *C. multicostatum* PHILLIPS, 1829, non BROCCHI, 1814; OD)]. Quadrate to rounded, small, with large, depressed lunule and false escutcheon. Hinge strong, with trigonal, short, entire 3b, 2, and 4b; strong tuberculose laterals. *U.Trias.(Carn.)-L.Jur.(Aalen.)*, Eu.-S.Am.-N.Z.—FIG. E51,4. *T. cingulata* (GOLDFUSS), *L. Jur.(Lias)*, Eng.; 4a-c, LV ext., LV int., RV int., $\times 8$ (187).

Vetericardiella CHAVAN, new genus, herein [12] [*nom. subst. pro Vetericardia* AUCTT. (non CONRAD, 1872)] [**Astarte crenalirata* CONRAD, 1860; OD]. Trigonal, slightly inequilateral, with radial ribs




FIG. E52. Carditidae (Carditamerinae) (p. N552).

regularly cut off by strong, deeply separated concentric furrows; lunule large, impressed. Elongate laterals on each side of hinge, *3b* short, isosceles triangular, *2* and *4b* equally oblique. *U.Cret.* (Senon.), N.Am.—FIG. E51,6a,b. **V. crenalirata* (CONRAD), USA (Tenn.); 6a,b, LV ext., int., $\times 4$ (951).—FIG. E51,6c. *V. webervillensis* (STEPHENSON), USA (Tex.); RV int., $\times 3$ (889).

[*Vetericardia* is a generic name invalidly proposed by CONRAD (1872, p. 52) as replacement of *Vetocardia* CONRAD, 1868 (Feb., 1869, p. 246), which is an objective junior synonym of *Pseudocardia* CONRAD, 1866 (p. 103) because published as a substitute name for the latter. A type species of *Pseudocardia* was not designated by CONRAD and STOLICZKA's (1871, p. 283) somewhat casual mention of *Venericardia dupiniana* (= *Cardita dupiniana* D'ORBIGNY, 1843) as a form that "can fairly be taken" as the type of *Vetocardia* (and hence of *Pseudocardia*) fails to qualify with provisions of the Code (1961, art. 67,c) as an explicit subsequent designation of a type species. A valid fixation of the type species of *Pseudocardia* was first made by CHAVAN (1952, p. 117) in naming "*C. Smidti* HORN." (presumed to be an erroneous citation of *Cardium schmidti* HÖRNES), eligible as one of the species originally assigned to the genus by CONRAD. STEPHENSON (1941, p. 175) invalidly designated *Astarte crenalirata* CONRAD (1860) as type species of *Vetericardia* CONRAD. A new name consequently is required for the genus based on *A. crenalirata*; *Vetericardiella* is chosen.]

Subfamily MIODOMERIDINAE Chavan,
new subfamily

Small, oblong, astartiform shells with immature ligament inframarginal. Large hinge plate with oblique *3b* and lacking *3a*. *Paleoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Miodomeris*.—2. *Chavanella*.—3. *Pteromeris*.—4. *Coripia*.

Miodomeris CHAVAN, 1936 [*Eomiodon* (M.) *cossmanni* CHAVAN, 1936; SD CHAVAN, 1938]. Compressed, with close-spaced concentric ribs and radial threads developed mainly at both ends;

lunule with convex margin. Hinge with cardinals and laterals. *Paleoc.-U.Eoc.*, Eu. (France-Belg.).

M. (Miodomeris) [1]. Moderately thickened, oblique, with radial lines distinct; ligament marginal. Laterals relatively faint, posterior ones not very remote. *M.Eoc.-U.Eoc.*, Eu. (France).—FIG. E53,1. **M. (M.) cossmanni* (CHAVAN), Lutet., W.France; 1a, LV ext., $\times 10$ (96); 1b,c, RV and LV hinges, enl. (96).

M. (Chavanella) JAWORSKI, 1938 [2] [*pro Eomiodon* CHAVAN, 1936 (non Cox, 1935)] [**Miodon semen* COSSMANN, 1908; OD]. Thickened, only slightly oblique, radial ribs faint; ligament inframarginal. Hinge with subsymmetrical *2* and *4b*, laterals well marked, posterior ones remote. *Paleoc.-U.Eoc.*, Eu. (Belg.-France).—FIG. E53, 3. **M. (C.) semen* (COSSMANN), Paleoc. (Mont.), Belg.; 3a,b, LV and RV hinges, enl. (97).

Pteromeris CONRAD, 1862 [**Cardita perplana* CONRAD, 1841 (= *Astarte radians* CONRAD, 1845); OD] [*non Pteromeris* CONRAD, 1865]. Somewhat compressed, oblong, obliquely rounded; with concentric and radial ribbing; lunule ill-defined. Hinge with obsolete posterior laterals. *L.Mio. (Aquit.)-Rec.*, W.Eu.-N.Am.-N.Z.

P. (Pteromeris) [3]. Radial ribs predominating over concentric ones laterally and around beaks; ligament marginal. *L.Mio. (Aquit.)-Rec.*, W.Eu.-




FIG. E53. Carditidae (Miodomeridinae) (p. N553-N554).




FIG. E54. Carditidae (Palaeocarditinae) (p. N554).

N.Am.-N.Z.—FIG. E53,4. *P. (P.) perplana abbreviata* (CONRAD), Mio., USA(N.Car.); 4a,b, RV ext., int., $\times 3.8$; 4c, LV int., $\times 4$ (Gardner, 1943).

P. (Coripia) DE GREGORIO, 1885 [4] [**Cardita (C.) corbis* (= *Cardita corbis* PHILIPPI, 1836, = *C. minuta* SCACCHI, 1836); M]. Sculpture uniformly reticulate; ligament partly internal. Mio. (Helvet.)-Rec., W.Eu.—FIG. E53,2. **P. (C.) corbis* (PHILIPPI), Rec., Medit.; 2a,b, LV and RV hinges, enl. (Chavan, n).

Subfamily PALAEOCARDITINAE Chavan
new subfamily

Posteriorly enlarged to trigoniform, beaks orthogyrous or opisthogyrous, tending to be

recurved. Hinge with high, pointed 3b. *L.Dev.(Downton.)-L.Jur.*

Arrangement of generic taxa by CHAVAN.—1. *Palaeocardita*.—2. *Carditomantea*.—3. *Schizocardita*.—4. *Pseudopsis*.

Palaeocardita CONRAD, 1867 [1] [**Cardita austriaca* (= **Cardium austriacum* HAUER, 1853); OD]. Inequilateral, enlarged posteriorly, with granulose or tripartite ribs; beaks almost orthogyrous. Hinge with well-developed cardinals and distant LV posterior lateral. *Trias.*, N.Z.-Eu.—FIG. E54,1a-d. *P. crenata* (MÜNSTER), M.Trias.(Ladin.), Aus. (Tyrol); 1a,b, LV and dorsal views of both valves, $\times 1$ (Münster, 1841); 1c,d, LV int., RV int., enl. (Laube, 1865).—FIG. E54,1e. **P. austriaca* (HAUER), U.Trias.(Rhaet.), Aus.(Tyrol); RV int., $\times 1$ (Stoppani, 1860-65).

Carditomantea QUENSTEDT, 1929 [2] [**C. spinata*; OD]. Trapezoidal, inequilateral, enlarged and rounded posteriorly, with medioposterior angulation; ribs spinose. Hinge with stout trigonal 3b and well-marked 2 and 4b; hinge plate enlarged downward. *L.Dev.(Downton.)*, Arctic O.(Spitz.). —FIG. E54,3. **C. spinata*; 3a, LV ext., $\times 1$; 3b, RV hinge, $\times 4$ (Quenstedt, 1929).

Pseudopsis COX, 1946 [4] [**P. astonensis*; OD]. Relatively small, subquadrate, posterior margin vertically truncate, exterior sharply bicarinate, covered by fine radial threads, growth stages defined; beaks strongly prosogyrous; lunule small, deep; escutcheon lacking. Each valve with single anterior lateral, in LV prolonged by 2, cardinal 4b oblique, 3b short; PI weak, PII present. *L.Jur.* (Lias.), Eu.(Eng.-France).—FIG. E54,2. *P. deslongchampsii* (TATE), M.Lias., Eng.; 2a-c, LV ext., int., RV int., enl. (187).

Schizocardita KÖRNER, 1937 [3] [**S. cristata*; OD]. Trigoniform, rounded in front, acuminate and biangulated posteriorly; beaks angular, prominent, slightly opisthogyrous. LV with 2 subequal diverging cardinals and single long, remote posterior lateral; RV with small anterior cardinal oblique, and trigonal, high, bifid median one, with base acuminate onward; long, curved, marginal lateral. *Trias.*, S.Am.(Peru).—FIG. E54,4. **S. cristata*; LV int., $\times 2$ (482).

Subfamily VENERICARDIINAE Chavan,
new subfamily

Outline subtrapezoidal to rounded trigonal, beaks strongly prosogyrous, with penetrating lunule. Hinge with laminar 3a, other cardinals curved, laterals approximate, almost obsolete. ?*U.Cret.(Senon.)*, *Paleoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Venericardia*.—2. *Venericor*.—3. *Leuroactis*.




FIG. E55. Carditidae (Venericardiinae) (p. N555-N556).

—4. *Pacificor*.—5. *Megacardita*.—6. *Trapezicardita*.

Venericardia LAMARCK, 1801 [**V. imbricata* (= *Venus imbricata* GMELIN, 1791); SD SCHMIDT, 1818]. Rounded trigonal, inequilateral, thick-shelled, with numerous radial ribs evenly elevated at young stage but becoming flattened and enlarged in adults. Hinge with compressed *3a* and *3b* higher than long. ?*U.Cret.* (*Senon.*), *Paleoc.-Eoc.*, Eu.-Afr.-N.Am.

V. (Venericardia) [1]. Subquadrangular, truncated almost vertically; ribs at juvenile stage rounded, in adults squarish and squamose; lunule very small. Cardinals narrow, high, with 2 and *4b* subparallel, elongate, *All* indistinct. ?*U.Cret.*

(*Senon.*), *Paleoc.-Eoc.*, Eu.-Afr.-N.Am.-N.Z.—FIG. E55,4. **V. (V.) imbricata* (GMELIN); M.Eoc. (Lutet.), France (Paris basin); *4a,b*, LV ext., int., $\times 0.8$ (Deshayes, 1837).

V. (Leuroactis) STEWART, 1930 [3] [**Venericardia pilsbryi*; OD]. Obliquely subtrigonal; immature ribs trifold, later becoming rounded and vanishing rapidly, replaced by numerous growth lines; beaks erect; lunule subvertical. Hinge with high, sharp cardinals, anterior lateral wanting. *L.Eoc.*, N.Am.—FIG. E55,3. **V. (L.) pilsbryi*, USA (Ala.); *3a,b*, RV ext., int., $\times 0.5$ (892).

V. (Pacificor) VERASTEGUI, 1953 [4] [**V. (P.) mulleri*; OD]. Subtrigonal to rounded; immature ribs trifold, later becoming simple and rounded,




FIG. E56. Carditidae (Glyptoactininae) (p. N556-N557).

nodulose to irregularly striated. *Paleoc.-Eoc.*, N. Am.-?Eu.

V. (Venericor) STEWART, 1930 [2] [**Venericardia planicosta* LAMARCK, 1799 (1806); OD]. Obliquely subtrigonal; immature ribs angular and spaced evenly, in adults flat and low, approximate, and finally vanishing; lunule depressed. Cardinals stout, 2 subtrigonal, 4b curved and thinner, 3b oblique; *All* tuberculiform, quite distinct, *PII* superficial. *Paleoc.-Eoc.*, Eu.-N.Am. —FIG. E55,2. **V. (V.) planicosta* LAMARCK, M.Eoc.(Lutet.), France(Paris basin); 2a,b, LV ext., int., $\times 0.5$ (Deshayes, 1837).

Megacardita SACCO, 1899 [5] [**Venericardia jouanneti* BASTEROT, 1825; OD]. Transversely elliptical, very inequilateral, with rounded ribs becoming approximated in adult stage; beaks rounded, prominent. Hinge with weak 3a, very oblique 3b larger than high, 2 and 4b present; *All* weak, other laterals obsolete. ?*Eoc.*, *Oligo.-Rec.*, Eu.-Afr.-Australia-N.Z. —FIG. E55,1. **M. jouanneti* (BASTEROT), Mio.(Helvet.), S.France; 1a-c, LV ext., int., RV int., $\times 0.5$ (165).

Trapezicardita CASEY, 1961 (6) [**Cypricardia squamosa* KEEPING, 1883; OD]. Quadrate, gibbous, very inequilateral, with obtuse medioposterior angulation; sculpture of squamous distant concentric laminae and radiating striae; prominent incurved anterior beaks; deeply impressed cordiform lunule. Hinge with 2 obtuse LV cardinals and 1 in RV; no developed laterals. Inner margin crenate. *L.Cret.*, W.Eu.

Subfamily CARDITESINAE¹ Chavan,
new subfamily

Short trapezoidal to rounded, tumid, with strong, more or less scaly ribs separated by V-shaped furrows; lunule enveloping, very convex, with 3a and 3b arched. *L.Cret.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Glyptoactis*.—2. *Claibornicardia*.—3. *Baluchicardia*.—4. *Ludbrookia*.—5. *Xenocardita*.—6. *Paraglans*.—7. *Cardites*.

Cardites LINK, 1807 [*non* LAMARCK, 1801 (spelling error)] [7] [**Chama antiquata* LINNÉ, 1758; M] [= *Cardita* MEGERLE, 1811 (*non* BRUGUIÈRE, 1792); *Actinobolus* MÖRCH, 1853 (*non* WESTWOOD, 1842) (obj.)]. Trapezoidal to rounded, with broad, striate or squamose ribs and short anal truncature; beaks tumid; lunule very convex, depressed, cordiform. Hinge with very faint 3a, other cardinals well marked; *All* small, *PII* minute. *L. Eoc.-Rec.*, Eu.-Asia-Australia-Pac. O. —FIG. E56,1. **C. antiquatus* (LINNÉ), *Rec.*, *Medit.*; 1a, RV ext., $\times 1$; 1b,c, RV and LV hinges, $\times 1.5$ (89a, 101).

Glyptoactis STEWART, 1930 [**Venericardia hadra*

¹ For avoidance of homonymy with Carditinae, based on *Cardita*, the spelling Carditesinae is introduced, approval being sought from ICZN as directed by the zoological Code (Art. 55a).




FIG. E57. Carditidae (Glyptoactininae) (p. N557).

DALL, 1903; OD]. Short trapezoidal, with high nodulose or echinate ribs; small, irregularly convex lunule extending under and slightly behind beaks. Hinge with very low 2, thin, long 4*b*, rather indistinct 3*a*, and strong, oblique 3*b*; *All* minute tuberculiform, close to 2, *Pll* faint. *U.Cret.* (*Senon.*)-*Mio.*, N.Am.-Eu.-Afr.

G. (Glyptoactis) [1]. Short, very convex, with prominent beaks, nodulose ribs, and marked posterior depression. Cardinal 2 very low. *Oligo-Mio.*, N.Am.—FIG. E56,3. **G. (G.) hadra* (DALL), *Mio.*, USA (Fla.); 3*a*, LV ext., $\times 1$; 3*b,c*, LV and RV hinges, $\times 1$ (229, Chavan, n).

G. (Baluchicardia) CHAVAN, herein (*ex* RUTSCH, 1944) [3] [**Cardita beaumonti* D'ARCHIAC & HAIME, 1854; OD]. Very short in front, elongate behind, with broadly tripartite ribs, median part nodulose. *U.Cret.* (*Senon.*)-*Paleoc.*, Afr.-N. Am.

G. (Claibornicardia) STENZEL & KRAUSE, 1957 [2] [**Venericardia alticosta* CONRAD, 1833; OD]. More developed on both sides and with lower beaks than *G. (Glyptoactis)*; ribs tripartite, posteriorly echinate; lunule less depressed and 2 more developed. *Eoc.*, N.Am.-Eu.

Ludbrookia CHAVAN, 1951 [4] [**Venericardia dupiniana* (= **Cardita dupiniana* D'ORBIGNY, 1843); OD] [= **Pseudocardia* STOLICZKA, 1871 (*non* CONRAD, 1866)] [See note under *Venericardiella*]. Short trapezoidal, with fine echinate ribs, posterior depression, and straight anal truncature; large sinuous lunule enveloping 2. Cardinals strong, 2 and 3*b* trigonal, 4 oblique, prominent; *Pll* broad, relatively weak, other laterals obscure. Projecting RV posterior margin. *L.Cret.* (*Alb.-Cenoman.*), Eu. (France)-N.Am.—FIG. E57,3. **L. dupiniana* (D'ORBIGNY), *Alb.-Cenoman.*; 3*a*, LV ext., $\times 1$ (695); 3*b,c*, LV and RV hinges, $\times 1.5$ (Chavan, n).

Paraglans CHAVAN, 1941 [6] [**Cardium calcitrapoides* LAMARCK, 1806; OD]. Subtrapezoidal to subquadrate, small, with broad posterior truncature; ribs rounded or partly spinose, simple; long convex lunule with broadened inner anterior margin. Hinge with oblique 3*b*, very thin 5*b*, short, rounded trigonal 2, and long, oblique 4*b*; laterals marked but weak, *All* nearly obsolete. *Paleoc.-Oligo.*, W.Eu.—FIG. E56,2; E57,2. **P. calcitrapoides* (LAMARCK), *M.Eoc.* (Lutet.), France (Paris basin); E56,2*a,b*, RV and LV hinges, enl.; E57,2, RV ext., $\times 2$ (Deshayes, 1837).

Xenocardita VOKES, 1946 [5] [**X. lacunaris* (= **Cardita lacunar* HAMLIN, 1884); OD]. Subtrapezoidal, with broad anal truncature; ribs spinose, posterior ones lamellose; lunule small, with sinuous anterior margin. Hinge with long, subhorizontal 3*a*, low, trigonal 3*b*, oblique 5*b*, curved oblique 4*b* above small trigonal 2; anterior ridge in both valves independent of cardinals but bearing obsolete laterals, LV with 2 posterior ones




FIG. E58. Carditidae (Thecaliinae) (p. N557-N558).

remote, RV with elevated margin. *L.Cret.* (*Apt.*), Lebanon.—FIG. E57,1. **X. lacunar* (HAMLIN); 1*a-c*, LV ext., int., RV int., $\times 3$ (945).

Subfamily THECALIINAE Dall, 1903

Transversely trapezoidal, ventral margin invaginated by incubatory chamber in female. *Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Thecalia*.—2. *Milneria*.

Thecalia ADAMS & ADAMS, 1858 [1] [**Cardita concamerata* BRUGUIÈRE, 1792 (= **C. concamerata* CHEMNITZ, 1784, not binominal); M]. Trapeziform, with rounded nodulose ribs, posterior ones flabelliform; incubatory chamber internal, ovate. Each valve with 2 cardinals, including long trigonal, sinuate 3*b*, and faint but distinct anterior laterals, posterior laterals obsolete. *Rec.*, S.Afr.-Austria.—FIG. E58,1. **T. concamerata* (BRUGUIÈRE), S.Afr.; 1*a,b*, RV ext., LV int., $\times 1.5$; 1*c*, LV hinge, enl. (85).

Milneria DALL, 1881 [2] [*pro Ceropsis* DALL, 1871 (*non* SOLIER, 1839)] [**Ceropsis minima* DALL, 1871; OD]. Narrowly trapeziform, very inequilateral, with 2 median posterior angulations, finely echinate ribs, concentric lines; beaks orthogyrous. Hinge with 3*b* isosceles triangular, 2 and 4*b* subsymmetrical, 3*a* and 5*b* obsolete; posterior laterals remote, anterior ones lacking. Incubatory chamber external. *Rec.*, W.N.Am. (Calif.).—FIG. E58,2. **M. minima* (DALL); 2*a*, RV ext., $\times 4$; 2*b,c*, both valves, ventral, dorsal, enl. (305); 2*d,e*, hinges, enl. (305).




FIG. E59. Condylardiidae (Condylardiinae) (p. N558-N559).

Subfamily UNCERTAIN

Pseudocardia CONRAD, 1866 [**Cardium schmidti* HÖRNES, 1861; SD CHAVAN, 1952] [= *Vetocardia* CONRAD, 1868 (1869) (*nom. van.*) (obj.); *Vetericardia* CONRAD, 1872 (*nom. van.*) (obj.)] [see note under *Vetericardiella*]. *Genus dubium.* Mio., Eu.

Family CONDYLOCARDIIDAE
F. Bernard, 1897

Minute, trigonal, trapeziform or ovate, commonly higher than long; radial ribs more or less marked but may be hidden by concentric sculpture. Spondyliiform hinge margin, internal ligament resilial edge partly obliterating cardinal 4b, behind which posterior tooth (5b-6b) is fused to long posterior laterals. *Eoc.-Rec.*

Subfamily CONDYLOCARDIINAE F. Bernard, 1897

[*nom. transl.* CHAVAN, herein (*ex* Condylardiidae F. BERNARD, 1897)]

Hinge margin more or less spondyliiform, resilium wholly internal, both laterals long and hooked. Prodissoconch saucer-shaped. *Eoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Condylocardia*.—2. *Benthocardiella*.—3. *Condyllocuna*.—4. *Radiocondyla*.—5. *Glibertia*.—6. *Micromeris*.—7. *Mesocuna*.—8. *Erycinella*.—9. *Carditopsis*.—10. *Cunanax*.—11. *Particondyla*. [Insert above, 6a. *Americuna*.]

Condylocardia F. BERNARD, 1896 [1] [**C. sanctipauli* MUNIER-CHALMAS in BERNARD, 1896 (= *C. pauliana* BERNARD, 1897); SD COSSMANN, 1902] [= *Hippella* MÖRCH, 1861 (type, *H. hippopus*; M) (*nom. oblit.*, ICZN pend.)]. Somewhat obliquely trigonal; radial ribs rounded, low, flabellate, crossed by concentric striae. Hinges with 3 cardinals in RV, 2 in LV: *AIII-3a, 3b, 5b, AII, 2a, 4b, 6b, PII, AI* much elongated. *M.Eoc.-Rec.*, S.Pac.-C.Am.-St.Atl.-Eu.—FIG. E59,2. **C. sanctipauli* MUNIER-CHALMAS, Rec., St.Paul Is.; 2a-c, LV ext., LV and RV hinges, much enl. (41).

Americuna KLAPPENBACH, 1962 (6a) [**A. besnardi*; OD]. Obliquely inequilateral, pyriform, with smooth well-marked prodissoconch. Dense concentric ribbing. RV (described as LV) with single cuneiform median tooth and 2 long laterals, separated from margin by narrow furrow; LV (described as RV) with 2 cardinals, 2 prominent, 4b thinner and 2 marginal laterals. Ligament in broad socket. Margin denticulate. *Rec.*, Brazil.

Benthocardiella POWELL, 1930 [2] [**B. pusilla*; OD]. Rounded trigonal, smooth to concentrically ribbed, large prodissoconch. Hinge plate thick, 3 RV cardinals, 2 in LV; hooked anterior and posterior duplicate laterals; resilium spoon-shaped. ?*M.Eoc.*, *Plio.-Rec.*, Australasia-?Eu. (France). —FIG. E59,1. **B. pusilla*, N.Z.; 1a-c, LV ext., LV int., RV int., much enl. (Powell, 1930).

Carditopsis E. A. SMITH, 1881 [9] [**Cardita flabellum* REEVE, 1843; OD]. Transversely trigonal, subequilateral, anteriorly only slightly produced, flabellate, granulose radial ribs well marked; beaks rounded. Hinge plate relatively narrow, with small teeth, 2, 3b, rounded median resilium, 5b, 6b (or *PI, PII*) behind socket nearly symmetrical with anterior cardinals; laterals very elongate, *AI* strongest. *Rec.*, S.Am. (Chile). —FIG. E60,1. **C.*




FIG. E60. Condylardiidae (Condylardiinae) (p. N558-N559).

Veneroida—Carditacea




FIG. E61. Condylardiidae (Condylardiinae) (p. N559-N560).

flabellum (REEVE), Valparaiso; 1a-c, RV ext., LV and RV hinges, $\times 4$ (Chavan, n).

Condyllocuna IREDALE, 1930 [3] [**Condylocardia projecta* HEDLEY, 1902; OD]. Transversely trigonal, very inequilateral, anteriorly elongated; sculpture of concentric waves. Cardinals 3a weak, 3b strong, 5b V-shaped, 2 oblique, 4b V-shaped, 6b stout; well-marked RV anterior and LV pos-

terior duplicate laterals; resilium well compressed. *Mio.-Rec.*, Australia-N.Z.—FIG. E59. [**G. projecta* (HEDLEY), *Rec.*, Australia; 3a-c, LV ext., int., RV int., enl. (397).

Cunanax IREDALE, 1936 [10] [**Cuna pisum* HEDLEY, 1908; OD]. Comparatively large, thick, proogyrous, strongly inequilateral, outline trigonal; approximated broad, low radial ribs transversely striated. LV teeth diverging subequally, also RV 3a and PI (or 5b), 3b bifurcate; anterior lateral remote. Lunule and escutcheon distinct; prodissoconch indistinct. *Plio.-Rec.*, Australasia.—FIG. E61.7. **C. pisum* (HEDLEY), *Rec.*, Australia; LV ext., much enl. (Hedley, 1908).

Erycinella CONRAD, 1845 [8] [**E. ovalis*; M] [= *Triodonta* VON KOENEN, 1893 (non BORY, 1827; nec Mulsant, 1842; nec Agassiz, 1846; nec Willesmere, 1885) (type, *T. clara*; SD GLIBERT, 1945)]. Ovate, somewhat enlarged anteromedially; radial ribs low, tending to become obsolete; both marginal ligament and rounded trigonal resilium. Hinge with oblique, elongate 2, scalariform stout 3b and 4b fused to edge of resilium socket, with elongate PI and PII just behind it, AII, PIII remote. *Oligo.-Rec.*, N.Am.-Eu.-S.Afr.—FIG. E61, 6. **E. ovalis*, *Mio.*, USA (Va.); 6a, b, RV and LV hinges, enl. (Gardner, 1943).

Glibertia VAN DEN MEULEN, 1951 [5] [**G. prosperi*; OD]. Very small, tumid, ovate, elongated




FIG. E62. Condylardiidae (Cuninae) (p. N560-N561).

anteriorly; concentric sculpture almost obsolete; flat prodissoconch. Hinge consisting of several projecting callosities, long *AIII-3a*, stout short *3b*, rounded *2*, *5b* behind broad resilium, *6b*, *PIII*, *PI* in successive crests. *Plio.*, Neth.—FIG. E61,5. **G. prosperi*; *5a,b*, RV and LV hinges, much enl. (Chavan, n).

Micromeris CONRAD, 1866 [**M. minutissima* (= *Astarte minutissima* LEA, 1833); OD] [= *Pteromeris* CONRAD, 1865 (non CONRAD, 1862) (obj.)]. Oblong, with pointed beaks and cordate broad lunule; sculpture vertical rather than truly radial; resilium somewhat restricted. Relatively narrow cardinals and very long laterals. *Eoc.-Rec.*, N.Am.-Australia.

M. (Micromeris) [6]. Trigonal, inequilateral, elongated anteromedially; with vertical ribs and concentric striae. Hinge with obliquely trigonal *2* and *3b*, very thin *3a*, *4b*, *5b*; shell margin strongly crenulate internally. *Eoc.*, N.Am.; *Rec.*, Australia.—FIG. E61,2. **M. (M.) minutissima* (LEA), *M.Eoc.*, USA (Ala.); *2a*, LV ext.; *2b,c*, RV and LV hinges; much enl. (Chavan, n).

M. (Mesocuna) LASERON, 1953 [7] [**M. saza*; OD]. Ovately trigonal, subequilateral; with rugose axial ribs and concentric lines. Hinge with massive cardinals and long, strong laterals; shell margin denticulate internally. *Rec.*, Australia.—FIG. E61,1. **M. (M.) saza*; *1a*, RV ext.; *1b,c*, RV int., LV hinge; much enl. (531).

Particondyla LASERON, 1953 [11] [**P. cuneata*; OD]. Obliquely trigonal, oblong; sculpture of radial ribs; prodissoconch saucer-shaped. Deep hinge plate with 1 cardinal on RV, 2 on LV, laterals not prominent. *Rec.*, Australia.—FIG. E61,3. **P. cuneata*; *3a*, LV ext.; *3b,c*, RV and LV hinges; much enl. (531).

Radiocondyla IREDALE, 1936 [4] [**R. arizela* (= *Condylocardia porrecta* HEDLEY & MAY, 1908) (non HEDLEY, 1906); OD]. Ovately oblong, relatively narrow; with broad radial ribs intersected by distant concentric furrows; prodissoconch small. Hinge with tuberculiform approximated short teeth of spondyloid appearance. *Mio.-Rec.*, Australia.—FIG. E61,4a. **R. arizela*, *Rec.*; RV ext., enl. (Chavan, n).—FIG. E61,4b,c. *R. jacksonensis* LASERON, *Rec.*; *4b,c*, LV ext., hinge, much enl. (531).

Subfamily CUNINAE Chavan, new subfamily

Hinge margin more or less rounded; prodissoconch also rounded, commonly ill-defined; resilium inframarginal; laterals incomplete. *M.Eoc.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Cuna*.—2. *Saltocuna*.—3. *Ovacuna*.—4. *Propecuna*.—5. *Volupicuna*.—6. *Hamacuna*. [Insert above, 4a. *Goniocuna*.]

Cuna HEDLEY, 1902 [1] [**C. concentrica*; OD]. Narrowly trigonal, commonly obliquely angular; with concentric sculpture overlapping radial ribs which strongly crenulate internal margin. Hinge with median *3b* subtriangular, posteriorly acuminate at base; *2* and *4b* nearly symmetrical but *4b* posteriorly restricted by resilium which does not reach inferior margin of plate; *AI*, *PII* thin, long, other laterals fused to margin; pallial line impressed. *M.Eoc.-Rec.*, N.Am.-Australia.—FIG. E62,2. **C. concentrica*, *Rec.*, Australia; *2a,b*, LV and RV hinges, much enl. (Chavan, n).

Goniocuna KLAPPENBACH, 1962 (4a) [**Cuna dalli* VANATTA, 1903; OD]. Subtriangular, inequilateral, very small, with concentric ribbing; hinge strong and broad, with low, long anterior and trigonal median cardinal in each valve, ligament in pit; bordered by posterior cardinal and by short rather faint posterior lateral. Adductor scars rather large; inner margin apparently smooth. *Rec.*, USA (Miss.-Fla.).

Hamacuna COTTON, 1931 [6] [**Cuna hamata* HEDLEY & MAY, 1908; OD]. Very oblique, anteromedially much elongated, with concave front margin and curved prosogyrous beaks, compressed, thick-shelled; surface with concentric sculpture and radial lines toward ventral margin; faint lunule and escutcheon. Hinge with curved bilobate *3b*, arched *2* and *4b* converging upward under beak, long curved *AIII-3a*, relieving remote *AI*; resilium narrow, elongate, and long nymph present. ?*Pleist.*, *Rec.*, Australasia.—FIG. E62,4. **H. hamata* (HEDLEY & MAY), Australia; *4a-c*, LV ext., int., RV int., much enl. (Chavan, n).


Saltocuna IREDALE, 1936 [**Cuna particula* HEDLEY, 1902; OD]. Broadly rounded obliquely, truncate posteriorly; sculpture coarse concentric; prodissoconch relatively large. Narrow hinge with short posterior laterals and relatively low cardinals. *Plio.-Rec.*, Australasia.

S. (Saltocuna) [2]. Very inequilateral, prodissoconch distinct. Hinge with strong *3b*, broader than high, and well-developed laterals. *Plio.-Rec.*, Australasia.—FIG. E62,6. **S. (S.) particula* (HEDLEY), *Rec.*, Australia; *6a,b*, LV ext., RV hinge, much enl. (Chavan, n).


S. (Ovacuna) LASERON, 1953 [3] [**Kellia atkinsoni* TENISON-WOODS, 1876; OD]. Less inequilateral than *S. (Saltocuna)*, concentric sculpture faint, prodissoconch not prominent. Hinge with distinctly bifurcate *3b* and weak laterals; pallial line impressed. *Rec.*, Australia.—FIG. E62,1. *S. (O.) solida* (COTTON); *1a-c*, RV ext., RV and LV hinges, much enl. (531).

S. (Propecuna) COTTON, 1931 [4] [**Cardita obliquissima* TATE, 1887; OD]. Very inequilateral, oblique grooves cutting across growth lines, also with radial posterior grooves and dorsoventral angulation. *Rec.*, Australia.—FIG. E62,3. **S.*

(*P.*) *obliquissima* (TATE); LV ext., much enl. (Chavan, n).




1b



1a

Yavorskiella



2b

2a

Aenigmoconcha

Volupicuna IREDALE, 1936 [5] [**Carditella delta* TATE & MAY, 1900; OD]. Highly triangular, slightly inequilateral, rather deep, with strong, well-spaced, pectinate radial ribs. Hinge with bifurcate cardinals *3b*, *2*, *4b* (subspherical), elongate, thin *3a-III*- and remote *AI*; pallial line obsolete; resilium quite inframarginal, slightly depressed. *Mio.-Rec.*, Australasia.—FIG. E62,5. **V. delta* (TATE & MAY), *Rec.*, Australia; *5a-c*, LV ext., LV and RV hinges, much enl. (Chavan, n).

Family UNCERTAIN

?*Aenigmoconcha* BENEDICTOVA, 1950 [**A. obliqua*; OD]. Shell subrhomboidal, of considerable size, umbones anteriorly directed, situated at varying distance from end. Hinge margin straight, or only slightly curved. Valves equally but not strongly inflated by diagonal elevation running from umbo to posteroventral angle; clearly delimited from steep posterior slope; valve surface almost flat near upper posterior angle, simulating obtuse auricle. Anterior slope of oblique elevation merging gradually with rest of valve; surface sculptured by narrow, rather closely set radial costae, strongest anteriorly, becoming weaker over diagonal ridge, tending to disappear on upper posterior field; costae of internal mold thin, linear, widely separated toward margins, strongest anteriorly and absent from posterior region; diagonal ridge bearing single narrow rib, simple or bifurcating downward. Flat surfaces of wide costal interspaces in mold marked by conspicuous concrescent lines. Internal characters unknown. [Nonmarine.] *L. Perm.*, USSR (W.Sib.).—FIG. E63,2. **A. obliqua*; *2a,b*, RV ext. molds, $\times 1$ (Ragozin, 1955). [WEIR]

?*Blairiella* MILLER & GURLEY, 1893 [**B. sedaliensis*; OD]. At present unrecognizable. *L. Carb.*, USA (Mo.). [NEWELL]

?*Yavorskiella* KHALFIN, 1950 [**Yavorskia skoki* FEDOTOV, 1938; OD]. Shell short, subrhomboidal, subelliptical or quadrilateral, narrowing toward lower posterior angle. Umbones submedian to anterior, broadly based, swollen, directed more or less anteriorly. Valves moderately convex, ill-defined diagonal elevation running from umbones to posterior extremity; shell sculpture of numerous strong radial costae, varying in number and width, widest interspaces occurring on oblique elevation; costae thinner and more closely spaced in front, rather indefinite on posterodorsal surface. Radial costation intersected by well-spaced growth folds. Internal characters unknown. [Nonmarine.] *L. Perm.*, USSR (W.Sib.).—FIG. E63,1a. **Y. skoki* (FEDOTOV); LV ext., $\times 2$ (Ragozin, 1955).—FIG. E63,1b. *Y. analoga* (BENEDICTOVA), RV mold, $\times 2$ (Ragozin, 1955). [WEIR]

FIG. E63. Family Uncertain (p. N561).

Superfamily CRASSATELLACEA Férussac, 1822

[*nom. transl.* NEWELL, 1965 (*ex* Crassatellidae FÉRUSSAC, 1822)]

[Materials for this superfamily prepared by ANDRÉ CHAVAN except as stated otherwise]

Animal with fully open mantle, or communication between branchial and pedal openings; unequal branchiae; papillate mantle edge. Shell trigonal, trapezoidal, or rounded, with concentric external sculpture, costate to striate, vanishing in some, except in Cardiniidae with internal layer of straight or arcuate radial riblets, which may be reflected as crenulations of inner shell margin. Lunule and escutcheon generally distinct; beaks prosogyrate and pointed. Hinge lucinoid, with *3b* median and *5b* commonly present on right valve, thus more or less cyrenoid in appearance. Lateral teeth laminar in many forms; no *PI* developed except in Cardiniidae and Myophoricardiidae. Ligament external or internal. Integripalliate or with very faint sinuosity. Pedal scars well marked (*110*, *111*). [Marine.] *Ord.-Rec.*

The alphabetically arranged generic descriptions in each family-group division of the Crassatellacea are accompanied by numbers inclosed by square brackets. Such numbers indicate position in the sequence of generic taxa given with the respective families or subfamilies for the purpose of recording CHAVAN's arrangement, designed to reflect "natural relationships" of these taxa as inferred by him.

Family ASTARTIDAE d'Orbigny, 1844

[=Crassinidae GRAY, 1840]

Trigonal rounded to subquadrangular in outline. Concentric ribbed to smooth, always ribbed around beaks. Successive internal layers of radial riblets periodically set up, under concentric outer layer. Ligament external or inframarginal, with nymph and without defined internal resilial pit. ?*M. Ord.*, *Dev.-Rec.*

Subfamily ASTARTINAE d'Orbigny, 1844

[*nom. transl.* CHAVAN, herein (*ex* Astartidae D'ORBIGNY, 1844)]

Shell trigonal or transversely trapezoidal, with more or less curved beaks; hinge without distinct *AIV*, other laterals (when present) in prolongation of cardinals. ?*M. Ord.*, *Dev.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Astarte*.—2. *Leckhamptonia*.—3. *Isocrassina*.—4. *Grotriana*.—5. *Ashtarotha*.—6. *Bythiamena*.—7. *Digitariopsis*.—8. *Tridonia*.—9. *Nicania*.—10. *Rictocyma*.—11. *Nicaniella*.—12. *Trautscholdia*.—13. *Astartella*.—14. *Eodon*.—15. *Kaibabella*.—16. *Eoastarte*.—17. *Astartopsis*.—18. *Astartopsis*.—19. *Matheria*.—20. *Praeconia*.—21. *Megapraeconia*.—22. *Yabea*.—23. *Neocrassina*.—24. *Coelastarte*.—25. *Prorokia*.—26. *Parisiella*.—27. *Sita*.—28. *Goodallia*.—29. *Ancliffia*.—30. *Gonilia*.—31. *Digitaria*. [Insert above, 1a. *Carinastarte*; 10a. *Filatovallea*; 16a. *Astartellopsis*; 25a. *Seendia*; 30a. *Ensio*.]


Astarte J. SOWERBY, 1816 [**Venus scotica* MATON & RACKETT, 1807 (= **Pectunculus sulcatus* DA COSTA, 1778, var. *scotica* MATON & RACKETT, 1807); OD]. Trigono-elliptical to subtrapezoidal, with regularly spaced concentric ribs or undulations, at least around prominent, but small, beaks. Inner margin denticulated or not. Hinge with well-developed cardinals, except for *3a*, faint, and *6b*, lacking; and with well-marked *AI*. Other laminae (*AII*, *PII*, *PIII*) partly obsolete. Narrow nymph behind *5b*. *Jur.-Rec.*, cosmop.

A. (Astarte) [1] [= *Crassina* LAMARCK, 1818 (obj.)]. Subquadrangular, inequilateral, posterior side enlarged and truncated; rounded strong persistent ribs. Remote lateral teeth and laminar oblique cardinals. Impressed lunule and escutcheon. *M. Jur.-Rec.*, cosmop.—FIG. E64, 1. **A. (A.) sulcata* (DA COSTA), *scotica* (MATON & RACKETT, Rec.), Scot.; 1a, RV ext., $\times 1.2$ (511); 1b,c, LV and RV hinges, ca. $\times 2.5$ (Maton & Rackett, 1807).

A. (Ashtarotha) DALL, 1903 [5] [**A. undulata* SAY, 1824; OD]. Ovatotrigonal, flattened, thick; depressed around the pointed beaks; posterior side commonly cuneiform. Sculpture of strong concentric waved ribs, in many forms vanishing on disc. Large, depressed, well-defined lunule and flat escutcheon. Teeth strong; elongated marginal laterals. *Neog.*, N.Am.-W.Eu.—FIG. E64, 2a. **A. (A.) undulata* SAY, Mio., USA (Va.); LV ext. $\times 1.2$ (Gardner, 1943).—FIG. E64, 2b,c. *A. (A.) undulata vaginulata* DALL, Mio. USA (Va.); 2b,c, RV int., ext., $\times 1.2$ (Gardner, 1943).

A. (Bythiamena) GARDNER, 1926 [6] [**A. (B.) isosceles*; OD]. Narrowly trigonal, high. Thick equidistant concentric ribs. Much excavated, but restricted lunule, obliterating *3a*; other cardinals strong; posterior laterals long. *Mio.*, USA (Fla.).—FIG. E64, 7. **A. (B.) isosceles*; 7a-c, LV ext., int., RV int., $\times 3.6$ (Gardner, 1926).

A. (Carinastarte) HINSCH, 1952 [1a] [**A. reimersii* SEMPER in RAVN, 1907; OD]. Like *A. (Astarte)* in outline, but with irregular ribbing and a



medioposterior angulation. Less remote, straighter and stouter posterior laterals, more vertical 4b. *Oligo.-Plio.*, Eu.

A. (*Digitariopsis*) CHAVAN, 1952 [7] [**A. grateloupi* DESHAYES, 1843; OD]. Trapezoidal, rounded, rather small, with irregular concentric fine ribs, commonly medially flattened and laterally oblique. Lunule and escutcheon narrow, long, defined only by their smooth surface. Hinge with well-marked long laterals, 2 very oblique forward. *Neog.*, W.Eu.-Japan.—FIG. E64,5. **A. (D.) grateloupi* DESHAYES, L.Mio. (Burdigal.), France; 5a, RV ext., $\times 2.8$; 5b,c, LV and RV hinges, $\times 3.6$ (Chavan, n).

A. (*Grotriania*) SPEYER, 1860 [4] [**Grotriania semicostata*; M]. Like *Isocrassina*, but with faint, medioposterior radial lines, more curved beaks, very large, excavated, concave lunule and flattened escutcheon. Hinge with 3a fused to margin; other cardinals very narrow; long obsolete laterals. *Oligo.*, Eu. (Ger.).—FIG. E64,3. **A. (G.) semicostata* (SPEYER); 3a-c, RV ext., LV int., RV int., $\times 2.4$ (Speyer, 1860).

A. (*Isocrassina*) CHAVAN, 1950 [3] [**A. castanea* SAY, 1830 (= **Venus castanea* SAY, 1822; OD)]. Subtrigonal-rounded, almost equilateral; thick; strong close-set umbonal ribs passing to low, rounded, spaced concentric undulations. Large shallow lunule. Hinge thick, with high cardinals and short laterals. *U.Eoc.-Rec.*, N.Am.-W.Eu.—FIG. E64,6. **A. (I.) castanea* (SAY), Rec., USA (N.J.); 6a-c, RV ext., int., LV int., $\times 1.2$ (Say, 1830). [= *Laevastarte* HINSCH, 1952 (type, *Tellina fusca* POLI, 1791; OD).]

A. (*Leckhamptonia*) COX & ARKELL, 1948 [2] [**Hiatella interlineata* LYCETT, 1850; OD]. Transversely trapezoidal, very inequilateral, rather small. Narrow sharp spaced ridges. Shallow lunule and escutcheon. Hinge with obscure laterals; 4b, small. *Jur.*, W.Eu.—FIG. E64,4. **A. (L.) interlineata* (LYCETT), M.Jur. (Bathon.), Eng.; 4a,b, RV ext., hinge, $\times 2.7$ (645; Chavan, n).

Astartella HALL, 1858 [13] [**A. vera* HALL, 1858; SD MILLER, 1889]. Transversely rhomboidal to ovate, thick, rather small; laterally truncated. Surface with concentric furrows and distant lamellae. Cordiform impressed lunule. Valves margin crenulate. RV hinge with narrow, trigonal 3b, oblique 5b, and marginal laterals; LV hinge with 2 partly fused as hook with raised (*All*) lunular margin, 4b, strong, broad and oblique, and *Pll* elongate. Scars small. *U.Carb. (Penn.)-Perm.*, Eu.-N.Am.-?Australia.—FIG. E65,2a,b. **A. vera*, Penn., USA; 2a,b, RV ext., both valves dorsal, $\times 1$ (Hall, 1858).—FIG. E65,2c,d. *A. concentrica* (CONRAD), Penn., USA (Mo.); 2c,d, LV and RV hinges, $\times 1.3$ (Girty, 1927). [See also Fig. E75,2.]

FIG. E64. Astartidae (Astartinae) (p. N562-N563).

Astartellopsis BEURLIN, 1954 (16a) [**A. prosoclina* (= *Astarte* cf. *triasina* "ROEMER" in COWPER REED, 1929); OD]. Trigonal, with high, acutely pointed, prominent prosogyrous beaks, anterior side de-

pressed, angular where it joins broadly convex ventral margin, posterior margin convex; long lunule. No dorsal carina, but broadly spaced, rather strong concentric waves. Hinge with curved




FIG. E65. Astartidae (Astartinae) (p. N563, N565-N568).

RV cardinals, 3a marginal, prolonged into straight anterior lateral fused to margin; 2 trihedral, strong, 4b long and curved, short nymph. *Permotrias.*, Brazil-Ger.-Japan.

Astartopsis VON WÖHRMANN, 1889 [17] [**Myophoria richthofeni* STUR, 1868; M]. Small, irregularly subquadrate, deep, with distant concentric ribs and strong posterior fold. Prominent beaks. Lunule and escutcheon broad and flat. Well-marked cardinals, obscure laterals. [Considered by Cox as junior subjective synonym of *Myophoriopsis* VON WÖHRMANN (see Myophoriopidae in this superfamily).] *U.Trias.*, Eu.(Aus.).—FIG. E65,1. **A. richthofeni* (STUR), Aus.; 1a, LV ext., $\times 4$; 1b,c, LV int., hinge, $\times 2$; 1d,e, RV hinge, int., $\times 2$ (1001).

Astartopsis DE LORIOL, 1891 [18] [**A. elongata*; SD CHAVAN, 1952]. Rounded, convex, thin; with concentric growths. Hinge with remote well-marked anterior laterals and widely divergent cardinals. Ovate superficial scars. Margin finely denticulated. *Jur.(L.Lusitan.)*, Eu.(Switz.).—FIG. E65,4. **A. elongata*; 4a, LV ext., $\times 0.7$; 4b,c, LV and RV hinges, $\times 1$ (550).

Digitaria S. WOOD, 1853 [31] [**Astarte digitaria* WOOD, 1853 (= *Digitaria vulgaris* WOOD, 1853) (= *Tellina digitaria* LINNÉ, 1758; T); SD LAMY, 1920] [= *Woodia* DESHAYES, 1860 (obj.); *Parvati* SEMPER, 1862 (invalid name)]. Ovate, inequilateral, rather small; more or less excentric sculpture. No lunule or escutcheon. Rounded beaks. Two strongly divergent cardinals on each valve and flexuous thin, elongate laterals. *Oligo.-Rec.*, W.Eu.-S. Afr.—FIG. E65,3. **D. digitaria* (LINNÉ), Rec., Port.; 3a, RV ext., $\times 2.5$; 3b,c, LV and RV hinges, $\times 5$ (Chavan, n).

Eoastarte CIRIACKS, 1963 [16] [**E. subcircularis*; OD]. Shell ovate or subcircular, inequilateral, prosocline; beaks prominent, anteriorly placed, prosogyre; shell surface smooth, dentition: 3b, 5b/2, 4b without laterals; 3b and 4b heavy, subtriangular, steeply inclined to hinge; without laterals; 2 short, slender, and 5b elongate, slender, both inclined and nearly parallel to adjacent shell margins; ligament opisthodontic; ligament nymphs and grooves short and shallow; adductor scars deep, subovate, dorsally extended, located high under hinge; anterior adductor pit bounded above by strong buttress which is fused to cardinal plate; pallial line simple. *L.Perm.*, USA(Wyo.).—FIG. E66,1. **E. subcircularis*, Grandeur Member, Park City F.; 1a,b, RV ext., LV int., $\times 4$ (132). [NEWELL]

Eodon HALL in MILLER, 1877 [14] [*pro Microdon* CONRAD, 1842, non AGASSIZ, 1833] [**Microdon bellastratus* CONRAD, 1842; M] [= *Microdonella* OEHLERT, 1881 (obj.)]. Trapezoidal, flattened, inequilateral, with medioposterior angulation and marginal sinuosity; concentric striated. Hinge

less developed than on *Astartella*, with *Alli-2*, *4b*, and *Pli* in LV. *Dev.-Carb.*, USA-W.Eu.—FIG. E66A,1a. **E. bellastratus* (CONRAD), M.Dev., USA; LV ext., $\times 1$ (146).—FIG. E66A,1b. *E. tremulus* (DE RYCKHOLT), Carb., Belg.; LV int., $\times 2$




FIG. E66. Astartidae (Astartinae) (1), (Opinae) (2) (p. N565, N572).




FIG. E66A. Astartidae (Astartinae) (p. N565-N566).

(Chavan, n).—FIG. E66A, 2. *E. sp.*, Dev., Ger.; LV int., diag. [classified as *Cypricardella* by HAFFER] (Haffer, 1959).

Filatovaellá MERKLIN, 1959 (10a) [*pro Astartella* FILATOVA, 1958, non HALL & WHITNEY, 1958] [**Astarte (Astartella) ioani* FILATOVA, 1958; OD]. Subtrapezoidal, short, thick, slightly inequilateral and angular backward; sculpture of numerous regular rounded ribs. Hinge plate broad, with 3 well-marked cardinals in RV, 2 in LV and thin *6b*; *4b* being subvertical; marginal obtuse RV anterior and LV posterior lateral; inner margin finely crenulated. *Rec.*, Kamchatka.

Gonilia STOLICZKA, 1871 [**Lucina? bipartita* PHILIPPI, 1836 (non DEFRANCE, 1823) (= **Astarte bipartita* PHILIPPI, 1844, non SOWERBY, 1829); (= *A. calliglypta* DALL, 1903); OD]. Rather small, rounded; divaricate sculpture. Ill-defined lunule. Two cardinals on each valve; lateral teeth long and narrow, anteriorly in RV but posteriorly in LV. Crenate margin. *Jur.-Rec.*, Medit.

G. (Gonilia) [30]. Suborbicular, with rather incurved low beaks; small lunule, no posterior area. Long narrow *P11*. *Rec.*, Medit.—FIG. E65, 6. *G. calliglypta* (DALL), Sicily; *6a*, LV ext., $\times 5$; *6b,c*, LV hinge, RV int., $\times 8$ (Chavan, n; Philippi, 1847).

G. (Ensio) COX, 1962 [30a] [**Prychomya agassizi* LYCETT, 1850; OD]. With a trigonal tendency, erected beaks; narrow, obtusely bordered lunule and escutcheon; *4b* somewhat grooved. *Jur.*, Eng.-Afr. (Borneo).

Goodallia TURTON, 1822 [**Mactra triangularis* MONTAGU, 1803; SD HERMANNSEN, 1847]. More or less trigonal, high, small. Anterior right lateral extending upward to beak, *3b*, *5b*; 2 and *4b*, *P11*. Broad scars. Margin crenate or not. *M.Jur.* (*Bathon.*)-*Rec.*, Eu.-Atl.

G. (Goodallia) [28] [= *Mactrina* BROWN, 1827 (obj.); *Mactroidea* BROWN, 1827]. Trigonal, smooth; almost orthogyrous; *3b* strongly bifid; *A11* indistinct, *P11* elongated. Margin crenate in adult stages. *Mio. (Burdigal.)-Rec.*, W.Eu.—FIG. E65, 5; E66A, 3. ***G. (G.) triangularis** (MONTAGU); *Rec.*, Atl., E65, 5a, LV ext., $\times 6$; E65, 5b, RV hinge, enl. (511; Turton, 1822); *Rec.*, Eng. (Sussex), E66A, 3a, b, RV ext., LV int., $\times 22$ (905a).

G. (Ancliffia) COX & ARKELL, 1948 [29] [**Astarte pumila* J. DE C. SOWERBY, 1824; OD]. Subtrigonal-rounded, with prominent curved beaks; finely striated. Cordiform shallow lunule and posterior area. Trigonal cardinals, obscure *A11*, short *P11*. Broad scars. Margin usually crenulate. *M.Jur.*, Eu.—FIG. E65, 8. ***G. (A.) pumila** (J. DE C. SOWERBY), Bathon., Eng.; LV int., $\times 4$ (Chavan, n).

Kaibabella H. CHRONIC, 1952 [15] [**K. curvilineata*; OD]. Ovatotrigonal, compressed, with rounded concentric ribs. Lunule and escutcheon poorly defined. Anterior cardinals in prolongation of obsolete laterals: *3a*, *4b*, strong; *5b*, *6b* present. No posterior laminae. Inner margin apparently smooth. *Perm.*, USA (Ariz.).—FIG. E65, 9. ***K. curvilineata**; *9a*, LV ext., $\times 2$; *9b,c*, LV and RV hinges, $\times 2$ (128).

?Matheria BILLINGS, 1858 [19] [**M. tener*; SD MILLER, 1889]. Subrectangular, very inequilateral; numerous concentric ribs and several growth-furrows; prosogyrous rounded beaks; concave lunule. Hinge with 2 LV and 1 RV distinct cardinals; short right lunular margin. [May belong in Cyrtodontidae; see Cyrtodontacea.] *M.Ord.*, N.Am. (Can.).—FIG. E65, 10. ***M. tenera**; *10a,b*, LV ext., int., $\times 1$ (618).

Megapraeconia CHAVAN, 1952 [21] [**Hippopodium bajocense* THÉVENIN, 1909 (ex D'ORBIGNY, 1850); OD]. Trapezoidal to ovate, large, very inequilateral and thick, posteriorly attenuated. Concentric irregular ribbing. Cordiform deep lunule under the curved beaks. Strong anterior scar. Much extended hinge plate with 2 high, massive, cardinals on each valve and obsolete short posterior laterals. Smooth margin. *Jur. (Bajoc.-Lusitan.)*, ?*L.Cret. (Neocom.)*, W.Eu.—FIG. E67, 2. ***M. bajocensis** (THÉVENIN), Bajoc., W.France; *2a,b*, RV int., LV int., $\times 0.5$ (Thévenin, 1909).




FIG. E67. Astartidae (Astartinae) (p. N565-N567).

Neocrassina FISCHER, 1886 [1887] [*nom. subst. pro Crassinella* BAYLE, 1878 (non GUPPY, 1874)] [**Astarte obliqua* DESHAYES, 1830 (= *Cypricardia obliqua* LAMARCK, 1819); SD DALL, 1903] [= *Puschia* ROUILLIER & VOSSYNSKI, 1847 (subj.) (type, *Astarte planata* SOWERBY, 1816; M); *Oreada* ROUILLIER & VOSSYNSKI, 1847 (*nom. nud.*); *Pruvostia* AGRAWAL, 1955 (subj.) (type, *Astarte* (*P. freneixae*))]. Ovate or trapezoidal, inequilateral, rather thick; concentric ribs commonly vanishing on adults. Two trigonal cardinals and strong posterior lateral on each valve. Broad flattened nymph; no 5b. Margin denticulated or not. *L.Jur.* (*U.Lias.*)-*U.Cret.* (*Turon.*), Eu.-Madag.-S.Afr.

N. (*Neocrassina*) [23]. Slightly depressed lunule; external escutcheon; *Pll* considerably behind

nymph. *L.Jur.* (*U.Lias.*)-*L.Cret.*, Eu.-Madag.—FIG. E67, 1. **N.* (*N.*) *obliqua* (LAMARCK), Bajoc., W.France; 1a-c, RV ext., int., LV int., $\times 0.5$ (Bayle, 1879).

N. (*Coelastarte*) BÖHM, 1893 [24] [**Astarte excavata* SOWERBY, 1819; OD]. Excavated lunule; obliquely penetrating, flat escutcheon; compressed cardinals; *Pll* just behind nymph. *L.Jur.* (*Aalen.*)-*U.Cret.* (*Turon.*), Eu.-Madag.-S.Afr.—FIG. E67, 3. **N.* (*C.*) *excavata* (SOWERBY), Bajoc., W. France; 3a,b, RV int., LV int., $\times 0.6$ (Böhm, 1893).

Nicaniella CHAVAN, 1945 [**Astarte communis* ZITTEL & GOUBERT, 1861; OD]. Trigonal or trapezoidal, small. Concentric ribs regular anteriorly and medially, faint or irregular posteriorly and ventrally. Inner margin usually crenate. Broad well-defined lunule and escutcheon. Hinge with small and strong cardinals, no 3a; laminar laterals; short narrow nymph. *M.Jur.*-*Paleoc.*, ?*Plio.*, N.Eu.-W.Eu.-Japan.

N. (*Nicaniella*) [11]. Trigonal to subquadrangular, with moderately prominent beaks; convex or straight posterior margin. Well-marked laterals. *U.Jur.*-*Paleoc.*, ?*Plio.*, Eu.—FIG. E65, 12. **N.* (*N.*) *communis* (ZITTEL & GOUBERT), *U.Jur.* (Sequan.), W.France; 12a,b, LV ext., int., $\times 4$; 12c, RV hinge, $\times 5.1$ (Chavan, 1945; Zittel & Goubert, 1861).

N. (*Trautscholdia*) COX & ARKELL, 1948 [12] [**Astarte cordata* TRAUTSCHOLD, 1861; OD]. Cordate, with high inflated beaks; more or less flexuous anterior, concave posterior margin. Strong *Al*; narrow cardinals. Impressed cordiform lunule and lanceolated escutcheon. *M.Jur.*-*U.Jur.*, NW.Eu.—FIG. E65, 13. **N.* (*T.*) *cordata* (TRAUTSCHOLD), *Jur.*, USSR; 13a,b, LV ext., int., $\times 0.7$; 13c, both valves, ant., $\times 0.7$ (Rouillier, 1847).

Parisiella COSSMANN, 1887 [26] [**P. ambigua*; M]. Very oblique, inequilateral, small and flattened. Fine concentric ribbing. Each valve with 2 well-marked divergent cardinals, 3b lobate; lateral remote, rather faint. Posterior scar somewhat raised. Smooth margin. *M.Eoc.* (*Lutet.*), Eu. (France).—FIG. E65, 7. **P. ambigua*, Paris basin; 7a-c, RV ext., int., LV int., $\times 10$ (160).

Praeonia STOLICZKA, 1871 [20] [**Astarte terminalis* ROEMER, 1842 (= *A. terminalis* DESHAYES, 1839-42, = *Hippopodium gibbosum* THÉVENIN, 1909) (*ex d'ORBIGNY*, 1850); OD] [= *Theveninia* ROMAN, 1921 (obj.)]. Subtrapezoidal, compressed, thick, very inequilateral, with median depression and posterior enlargement. Numerous concentric rounded ribs and more or less equidistant growth furrows. Small deep concave lunule. Trigonal 3b, with faint *AIII-3a* in front; 2, 4b and remote, elongated *Pll*. Margin irregularly crenate. *Jur.* (*L.Lias.*-*Lusitan.*), Eu. (France)-Japan.—FIG. E65, 18a,b. **P. terminalis* (DESHAYES), *M.Jur.* (Bajoc.),

W. France; *18a*, RV view, ext., $\times 0.7$; *18b*, RV int., $\times 1$ (257; Thévenin, 1909).—FIG. E65, 18c. *P. sarthacensis* COSSMANN, M. Jur. (Bathon.), W. France; LV int., $\times 1$ (Cossmann, 1914).

Prorokia BÖHM, 1893 [25] [**Cardita ovalis* QUENSTEDT, 1852; OD] [= *Pachytypus* MUNIER-CHALMAS (ex FISCHER, MS), 1887 (type, *Cardita problematica* BUVIGNIER, 1852; OD)]. Subelliptical, thick, rather small, concentrically ribbed. No lunule or escutcheon. Widely divergent cardinals. *3b* largely trigonal, *3a*, 2 and *4b* long and thin; remote posterior laterals, no anterior ones. Raised posterior scar. Crenate margin. M. Jur. (Bathon.)-U. Jur. (Kimmeridg.), Eu.—FIG. E65, 11. *P. problematica* (BUVIGNIER), U. Jur. (Raurac.), E. France; *11a-c*, LV ext., int., RV int., $\times 2$ (Buvignier, 1852).

? **Seendia** CASEY, 1961 (25a) [**Crassatella saxoneti* PICTET & ROUX, 1847; OD]. Transversely inequilateral, subrectangular, thick, compressed. Narrow well-defined lunule. Concentric ridges and faint radial lines; margin crenate. Narrow hinge, with 2 cardinals in each valve, *3b* much larger; ligament said to be sunken; posterior adductor surelevated. [Possibly belongs to Eryphyllinae.] *L. Cret.*, W. Eu.

Sita SEMPER, 1862 [27] [**Woodia crenulata* DESHAYES, 1860; SD CHAVAN, herein] [= *Crenimargo* COSSMANN, 1902 (obj.)]. Ovate, smooth, rather small. Each valve with 2 cardinals, LV anterior and RV posterior ones strongly lobate; no laterals. Ovately elongated scars. Margin obliquely crenulate. *Eoc.*, W. Eu.—FIG. E65, 14. **S. crenulata* (DESHAYES), M. Eoc. (Lutet.), France (Paris basin); *14a*, LV ext., $\times 4$; *14b, c*, U. Eoc. (Barton.), Belg., LV int., RV int., enl. (Glibert, 1936).

[Although originally indicated to be Sanskrit name, *Sita* is here accepted as a properly latinized designation with priority over *Crenimargo*, which was based on misinterpretation of the same species.]

Tridonta SCHUMACHER, 1817 [**T. borealis* (= *Venus borealis* CHEMNITZ, 1784, invalid non-binom. ICZN); M] [= *Tridonta* AGASSIZ, 1846 (*nom. null.*)]. Transversely quadrangular to rounded; closely spaced concentric ribs at least in early growth stages; inner margin usually smooth. Hinge lacking *AIV* and *PI*; strong short *AI* and *PII*. Broad nymph behind small *6b*. [Localized in cold and temperate waters.] *L. Cret.-Rec.*, Arctic-N.N. Am.-Japan-Eu.

T. (**Tridonta**) [8]. Medium-sized, transversely inequilateral; concentric ribs more or less vanishing on the disc. Flattened long lunule. *L. Cret.* (*Neocom.*)-*Rec.*, Atl.—FIG. E65, 17. **T.* (*T.*) *borealis*, *Rec.*, North Sea; *17a*, LV ext., $\times 1$; *17b, c*, LV and RV hinges, $\times 0.5$ (511; Sowerby, 1854).

T. (**Nicania**) LEACH, 1819 [9] [**N. banksii* (= *Venus montagui* DILLWYN, 1817 *var. banksii*); SD GRAY, 1847]. Smaller, less inequilateral, with persistent concentric rounded ribs. Very long more depressed lunule, obliterating *3a*; large, somewhat

bifid *3b*; posterior cardinals laminar; *AI* strong and *PII* remote, springing from under hinge plate. *Pleist.-Rec.*, circumpolar-N. Eu.-N.N. Am.-Japan.—FIG. E65, 16. **T.* (*N.*) *montagui* (DILLWYN), *Rec.*, USA (Mass.); *16a, b*, LV ext., RV int., $\times 1$ (Gould, 1841).

T. (**Rictocyma**) DALL, 1872 [10] [**R. mirabilis*; M] [= *Rhectocyma* VON MARTENS, 1873, emend.]. With concentric striae and larger ribs, becoming broadly irregular, oblique, and interrupted on disc. *Plio.-Rec.*, Arctic.—FIG. E65, 15. **T.* (*R.*) *mirabilis* (DALL), *Rec.*, Unga Is.; LV ext., $\times 3$ (Dall, 1872).

Yabea HAYAMI, 1965 [22] [**Astarte shinanoensis* YABE & NAGAO in YABE, NAGAO, & SHIMIZU, 1926; OD]. Very inequilateral, strongly convex, trapezoidal, with prominent prosogyrous beaks; deeply depressed lunule; margin crenulated; *3a* short, curved, *3b*, *5b* narrow, not clearly separated from the nymph; 2, *4b*; well developed both anterior and long posterior laterals. *L. Cret.*, Japan.

Subfamily ERIPHYLLINAE Chavan, 1952

Shell lenticular or crassatelliform, commonly with pointed beaks. Hinge with *AIV* reaching almost to beak, its socket forming narrow groove in front of *3a*; other laterals (when present) also elongate and laminar. *Dev.-Eoc.*

Arrangement of generic taxa by CHAVAN.—1. *Eriphyla*.—2. *Dozyia*.—3. *Herzogina*.—4. *Amphiaraus*.—5. *Bruniastarte*.—6. *Eriphylopsis*.—7. *Disparilia*.—8. *Crassatellopsis*.—9. *Astartemya*.—10. *Freiastarte*.—11. *Crassatellina*.—12. *Cardiniopsis*.—13. *Lirodiscus*.—14. *Crustuloides*.

Eriphyla GABB, 1867 [**E. umbonata* GABB, 1864; OD]. Almost equilateral, rounded; smooth or finely lamellar. Obliquely depressed lunule; narrow escutcheon. Hinge with 2 cardinals on each valve, *4b* not reaching inferior margin of plate but posteriorly fused to nymphal thickening; *AIII*, *AIV*, *PII*, *PIII* present. Shallow pallial inflexion. Margin smooth or crenulate. *Cret.*, W.N. Am.-W. Eu.-Japan. [= *Mikayoella* HAYAMI, 1965 (type, *Astarte mikayoensis* NAGAO in YABE, 1927; OD).]

E. (**Eriphyla**) [1]. Thickened, high; *3a* fused to *AIII*, *3b* not very oblique, *4b* reduced to its upper part; posterior laterals not far behind cardinals. *Cret.*, W.N. Am.-W. Eu.—FIG. E68, 1. *E.* (*E.*) *lapidis* (PACKARD), U. Cret., USA (Calif.); *1a, b*, LV and RV hinges, $\times 0.75$ (748).

E. (**Dozyia**) BOSQUET in DEWALQUE, 1868 [2] [**D. lenticularis* (= *Lucina lenticularis* GOLDFUSS, 1840); M]. Broader and thinner than *E.* (*Eriphyla*); *3a* separated from *AIII*, *3b* very oblique, *4b* widened posteriorly; posterior laterals




FIG. E68. Astartidae (Eriphylinae) (p. N568-N571).

remote. *U.Cret.*(*Senon.*), W.N.Am.-W.Eu.—
FIG. E68,6. **E. (D.) lenticularis* (GOLDFUSS),
Campan., Neth.; 6a-c, LV ext., int., RV int.,
×0.7 (415).

Amphiararus VOKES, 1946 [**A. seleniscus*; OD].
Flattened and rounded, inequilateral, smooth. Al-
most flat lunule and sharply limited escutcheon.
Hinge with *AIII-3a* under *AIV*; *3b* trigonal, *5b*,
and broad nymphal plate; remote posterior laterals.
M. Jur.(*Bajoc.*)-*L. Cret.*(*Apt.*), M. East (Lebanon)-
W.Eu.

[Supposition by Cox that *Amphiararus* is a junior synonym
of *Eomiodon* Cox, 1935, classed in the family Eomiodonti-
dae of the Glossacea, is firmly rejected by CHAVAN, who
points out that the shell of *Amphiararus* is lenticular, con-
trasting strongly with the peculiar trigonal-transverse or-
nament. Further, *Eomiodon* has long sinuate lateral teeth,
no *AIV* groove above *3a*, and bears *PI*, a tooth which is
lacking in *Amphiararus* and basically in other astartids.—

Ed.] [CHAVAN states (11 January, 1967): "*Amphiararus* has
been treated by several authors as a small neomiodontid
(arcticacean), but its small, remote *fAIV* above a stout,
noncorbiculoid *AIII*, as well as its *PIII*, instead of *PI*,
astartoid cardinals, and lenticular form, are characteristic
of the Eriphylinae."]

A. (Amphiararus) [4]. Transversely ovate. Broad
AIII-3a; *3b* very oblique; strong right posterior
ridge or lateral. *L.Cret.*(*Apt.*), Lebanon.—FIG.
E68,2. **A. (A.) seleniscus* VOKES; 2a,b, RV ext.,
hinge, ×0.75, ×1 (945).

A. (Bruniastarte) CHAVAN, 1952 [5] [**Astarte*
thoas THÉVENIN, 1909; OD]. Subquadrangular.
AIV straight and raised; *AIII-3a* thinner; *3b* not
very oblique, *PIII* faint. *M. Jur.*(*Bajoc.*), Eu.(W.
France).—FIG. E68,5. **A. (B.) thoas* (THÉVE-
NIN); LV int., ×1 (110, 111).

Astartemya STEPHENSON, 1941 [**A. fentressensis*;
OD]. Subtrigonal to ovate, flattened; prominent




FIG. E69. Astartidae (Eriphylinae) (p. N570).

curved beaks. Concentric waves or ribs and fine growth lines. Depressed lunule, flat escutcheon. Hinge with long laterals prolonged upward, commonly to beaks; 2 oblique cardinals on each valve, 5b obsolete. Flat nymphal plate in front of narrow crest and furrow. Slight pallial impression. *U.Cret.*, ?*Paleoc.*, cosmop.

A. (Astartemya) [9]. Higher than long, subovate; shallow undulations; grooved cardinals; laterals bounded by flangelike plate, anterior ones reaching beaks. Broad nymph. Fine marginal crenulations. *U.Cret.* (*Senon.*), cosmop.—FIG. E68,8. **A. (A.) fentressensis* STEPHENSON, USA (Tex.); 8a-c, RV ext., int., LV int., $\times 0.7$ (889).

A. (Freiastarte) CHAVAN, 1952 [10] [*nom. subst. pro Freia* BÖHM, 1884 (*non* CLAPARÉDE & LACHMANN, 1858)] [**Astarte coelata* MÜLLER, 1847

(= **Astarte similis* MÜNSTER in GOLDFUSS, 1837 [1840]); OD]. Small, subtrigonal; concentric distant ribs. Anterior laterals not completely reaching beaks, posterior ones thickened behind, 5b conspicuous. Ligamentary plate on a superficial reed-shaped area. Margin strongly crenate, sometimes smooth. *Cret.*, and possibly *Paleoc.*, W. Eu.-S. Afr.-India. — FIG. E68,9. **A. (F.) similis* (MÜNSTER in GOLDFUSS), Campan., Neth.; 9a-c, RV ext., RV and LV hinges, $\times 0.7$ (110, 111, 415).

Cardiniopsis STANTON, 1895 [12] [**C. unioides*; OD]. Largely transverse, anteriorly rounded and posteriorly narrowed. Hinge with one prominent median RV cardinal; 3a, 5b, faint, and 2 sub-symmetric LV teeth. Broad long nymph, no laterals except for weak AIV in front of 2. Well-impressed pedal scars. *L.Cret.*, USA (Calif.). — FIG. E69,1. **C. unioides*; 1a, LV int., $\times 0.3$; 1b, both valves, dorsal, $\times 0.3$; 1c, RV ext., $\times 0.3$; 1d, RV hinge, $\times 0.5$ (877).

Crassatellina MEEK, 1871 [11] [**C. oblonga*; OD]. Transversely subtrapezoidal, rather small. Well-marked anterior lateral prolonged up to beak, in front of cardinals; 2 and 3b bifid. Elongated nymph. No posterior laterals. *U.Cret.*, USA (N. Dak.-Kans.). — FIG. E68,12. **C. oblonga*, USA (Kan.); 12a, LV ext., $\times 1$; 12b,c, LV int., RV int., $\times 2$ (Meek & Hayden, 1856).

Crassatellopsis BEUSHAUSEN, 1895 [8] [**C. hauchecornei*; M]. Sublenticular, almost smooth. Pointed beaks. Hinge with stout trigonal anterior LV cardinal and obliquely directed RV one; posterior left and anterior right, both thinner. Very narrow, almost linear resilium apparently present. Anterior scar somewhat elliptical. *Dev.*, Eu. (Ger.). — FIG. E70,1. **C. hauchecornei*, Rheinland; 1a-c, LV int., RV int., LV int., $\times 1$ (47).

Disparilia CHAVAN, 1953 [7] [**Astarte disparilis* D'ORBIGNY, 1843; OD]. Trigonal-transverse, posteriorly narrowed; concentric somewhat lamellar ribs. Hinge with almost obsolete laterals, narrow 3a and 5b, 2, stout 3b and 4b. External ligament sunken into recess of escutcheon, simulating resilial socket. Crenate margin. *Cret.*, W. Eu.-N. Afr.-Lebanon. — FIG. E68,11. **D. disparilis* (D'ORBIGNY), *L.Cret.* (Neocom.), E. France; 11a,b, LV ext., int., $\times 1.5$ (115, 695).

Eriphylopsis MEEK, 1876 [6] [**Eriphyla gregaria* MEEK & HAYDEN, 1856; OD]. Subtrapezoidal to rounded, rather small; concentrically undulate, lunule relatively broad. Hinge with elongate laterals, RV anterior and LV posterior ones duplicate, and anterior cardinals more or less in prolongation of their laterals; stout posterior cardinals. *U.Cret.*, W. Eu.-S. Eu.-W. N. Am. — FIG. E68,3. **E. gregaria* (MEEK & HAYDEN), USA (Mont.); 3a,b, LV hinge, RV hinge, $\times 2$ (Chavan, n); 3c, RV ext., $\times 1.5$ (Meek & Hayden, 1856).




FIG. E70. Astartidae (Eriphyllinae) (p. N570).

Herzogina CHAVAN, 1952 [3] [**Astarte herzogii* KRAUSS, 1850 (= **Cytherea herzogii* HAUSSMANN, 1837); OD]. Rounded, moderately convex, with strong concentric ribs. Depressed lunule. Hinge with marginal *AIV*, *AIII* almost perpendicular to *3a*, *3b* curved onwards on its lower part; a broad depression for *4b* and the nymph; remote posterior laterals. Crenate margin. *L.Cret.* (*Neocom.*), S.Afr.—FIG. E68,10. **H. herzogii* (HAUSSMANN); *10a-c*, LV ext., int., RV int., $\times 0.7$ (Krauss, 1850).

Lirodiscus CONRAD, 1869 [**Astarte tellinoides* CONRAD, 1833; SD DALL, 1903]. More or less transversely rounded and compressed. Concentrically sculptured. Flattened beaks and escutcheon. Two cardinals on each valve; anterior laterals obliquely directed; posterior ones prolonged under ligamentar extension. Well-impressed scars. Margin commonly denticulate. *Eoc.*, USA.

L. (Lirodiscus) [13]. Thickened, with strong concentric ribs and growth striae; with posterior fold. Broad lunule and escutcheon. Flat nymph with undulating growth striae, extending up from posterior margin to and above *3b* and *4b*. Anterior left laterals bifurcate under lunule. *Eoc.*, USA.—FIG. E68,7. **L. (L.) tellinoides* (CONRAD), *M.Eoc.*, Ala.; *7a-c*, LV ext., int., RV int., $\times 1$ (Harris, 1919).

L. (Crustuloides) G. D. HARRIS, 1919 [14] [**Crassatellites (Scambula) psychopterus* DALL, 1903; M]. Thin, much compressed; irregularly striated, undulating peripherally; lunule and escutcheon linear. Hinge plate enlarged in front, with one straight anterior lateral; posterior ones somewhat broadened. *L.Eoc.*, USA (Miss.).—FIG. E68,4. **L. (C.) psychopterus* (DALL); *4a-c*, LV ext., int., RV int., $\times 1$ (Harris, 1919).

Subfamily OPINAE Chavan, 1952

[*nom. correct.* herein (*pro* Opininae CHAVAN, 1952)]

Very high beaks; anterior laterals obsolete; high, trigonal cardinals. Oblong or transverse carinated shell, with a defined posterior area (50). *Dev.-U.Cret.*

Arrangement of generic taxa by CHAVAN.—1. *Opis*.—2. *Trigonopsis*.—3. *Pachyopsis*.—4. *Coelopsis*.—5. *Cryptocoelopsis*.—6. *Prosocoelus*.—7. *Tripleura*.—8. *Prosocoelogeton*.—9. *Heteropsis*.—10. *Trigonastarte*.—11. *Seebachia*.—12. *Opisoma*.

Opis DEFRANCE, 1825 (1824, *nom. nud.*) [**Trigonia cardissoides* LAMARCK, 1819; M]. Obliquely oblong, with prominent beaks. Ventral margin curved in front of medioposterior undulation; posterior margin then truncated. Concentric ribbing. More or less shallow lunule. Large hinge, without laterals; one strong RV trigonal and 2 LV cardinals, commonly with their lateral faces striated; *4b* stronger than 2. Ligament narrow. Margin more or less crenate. *L.Jur.* (*Lias.*)-*U.Cret.*, Eu.-Madag.-N.Am.-Japan.

O. (Opis) [1]. Subtrigonal, rounded in front and ventrally, inflated, obliquely elongated. Erect beaks. Lunule large, somewhat depressed. Ribs closely spaced. Hinge with *3b* triangular and high. *L.Jur.* (*Lias.*)-*U.Cret.*, Eu.-N.Am.-Madag.—FIG. E71,2. **O. (O.) cardissoides* (LAMARCK), *U.Cret.* (Cenoman.), Belg.; *2a,b*, RV ext., ant.; *2c,d*, LV hinge view, ant.; *2e*, LV int. mold, $\times 1$ (50; d'Archiac, 1847).

O. (Pachyopsis) BIGOT, 1895 [3] [**O. ponderosa* DESLONGCHAMPS, 1883; OD]. Much inflated, thick; ventrally (instead of posteroventrally) elongated. Ribs crossed by median and postmedian angulation. Beaks rounded and conspicuously prosogyrous. Very shallow lunule. Rudimentary anterior LV cardinal. Margin crenate. *Jur.* (*Bajoc.-Lusitan.*), W.Eu.—FIG. E72,3. **O. (P.) ponderosa* DESLONGCHAMPS, *Bajoc.*, W. France; *3a,b*, RV ext., int., $\times 0.7$ (50).

O. (Trigonopsis) FISCHER (*ex* MUNIER-CHALMAS, MS), 1887 [2] [**Opis similis* D'ORBIGNY, 1844 (= *Cardita similis* SOWERBY, 1819); M]. Subtrapezoidal, angular in front; inflated, but slightly curved forward; narrow regularly spaced ribs; strong medioposterior angulation. Rounded beaks. Shallow lunule; *3b* largely and more or less obtusely triangular. *L.Jur.-L.Cret.*, W.Eu.—FIG. E71,3. *O. (T.) praesimilis* COSSMANN, *M.Jur.* (*Bajoc.*), France; *3a-c*, RV ext., int., LV int., $\times 3$ (161).

Coelopsis FISCHER (*ex* MUNIER-CHALMAS, MS), 1887 [**Opis lunulata* MILLER" (*errore pro* **Cardita lunulata* J. SOWERBY, 1819); M]. Subquadrangular to subtrapezoidal, with almost enrolled prosogyrous beaks and large depressed well-defined lunule. Concentric ribbing; anteromedian and medioposterior carinas. Relatively restricted hinge with one strong RV and 2 LV narrow cardinals, without lateral striations; 2 very rudimentary. Denticulated margin. *M.Trias.-L.Cret.*, Eu.

C. (Coelopsis) [4]. Without chamber individualized in front of teeth. *M.Trias.-L.Cret.*, Eu.—FIG. E71,1a. **C. (C.) lunulata* (SOWERBY), *M.Jur.*

(Bathon.), France; RV ext., $\times 3$ (Cossmann, 1918).—FIG. E71, 1b-d. *C. (C.) affinis* LAUBE, M.Trias.(Ladin.), Aus.(Tyrol); 1b,c, LV ant. hinge; 1d, RV hinge; all $\times 3$ (58).

C. (Cryptocoelopsis) BITTNER, 1895 [5] [**Opis locularis* BITTNER, 1895 (= ?*O. affinis* LAUBE, 1865, "female form"); SD CHAVAN, herein].




FIG. E71. Astartidae (Opinae) (p. N571-N573).

With a wide chamber individualized between the lunular margin and the anterior margin of the hinge plate, thus duplicated. *M.Trias. (Ladin.), Eu.(Aus.)*.—FIG. E72, 2. **C. (C.) locularis* (BITTNER), Tyrol; LV hinge, $\times 3$ (58).

Heteropsis BÖHM, 1893 [9] [**Opis carinata* QUENSTEDT, 1858; OD]. Higher than long, inequilateral, with medioposterior angulation; finely ribbed. Large flattened or shallow lunule. Prosogyrous oblique prominent, but small beaks. Hinge with anterior laterals partly developed and fused to anterior cardinals, 3b narrow; posterior ones rudimentary. Smooth inner margin. *Jur.(Bathon.-Lusitan.), Eu.*—FIG. E72, 1a,b. **H. carinata* (QUENSTEDT), Lusitan., Ger.; 1a,b, LV ext., int., $\times 2$ (Buvignier, 1852).—FIG. E72, 1c. *H. raulinea* (BUVIGNIER), U.Jur.(U.Oxford.), E.France; RV int., $\times 1$ (Böhm, 1893).

Opisoma STOLICZKA, 1871 [12] [**Opis paradoxa* BUVIGNIER, 1852 (= **Cardium paradoxum* BUVIGNIER, 1843); OD]. Trigonal, extremely oblong, with an anteromedian strong carina. Opisthogyrous erect beaks. Concentric striation. Three RV and 2 LV much elongated narrow cardinals; 6b fused to narrow nymph. Anterior scar on a long platform. *Jur.(Infralias-Raurac.), Eu.(France)*.—FIG. E72, 4. **O. paradoxum* (BUVIGNIER), U.Jur. (Lusitan.), E.France; 4a-c, LV ext., int., RV int., $\times 0.7$ (Buvignier, 1852).

Prosocoelus KEFERSTEIN, 1857, p. 155 [**Venus prisca* ROEMER, 1843; SD HAFFER, 1959]. Medium-sized, ovate, gibbous, umbones high, prosogyrous, more or less anterior; lunule very deep, just below beaks; without lateral teeth; RV with strong cardinal tooth received between 2 teeth of LV, anterior of which is ill-defined conical and posterior rather thin; with several diagonal folds and furrows on mid-posterior area or lacking ornament. *Dev., Eu.-Spitz.*

P. (Prosocoelus) [6]. Short subquadrate, nearly smooth. *L.Dev., Ger.*—FIG. E66, 2. **P. (P.) priscus* (ROEMER); 2a,b, RV and LV hinges, $\times 1$; 2c, RV int., $\times 0.3$ (Haffer, 1959).


P. (Tripleura) SANDBERGER, 1889 [7] [**T. pesanseris* (= **Grammysia pesanseris* ZEIL & WIRTGEN, 1851); OD]. Posteriorly elongate, with several radial folds. *L.Dev., Ger.*

P. (Prosocoelogeton) QUENSTEDT, 1926 [8] [**P. (P.) lenticularis*; OD]. With distinct 3a along lunular margin, 4b shorter and stouter than 3a. *U.Dev., Spitz.*

Seebachia HOLUB & NEUMAYR, 1881 [11] [**Astarte bronni* KRAUSS, 1850; OD]. Transversely trigonal, large and very inequilateral; anterior end rounded, posterior much elongated with dorsal angulation. Concentric ribbing. Lunule and escutcheon developed. Cardinals unequal, striated on their lateral faces, 3b strongest; anterior laterals small, posterior ones absent. Scars impressed, posterior one

on platform. Margin crenate. *L.Cret.(Neocom.)*, S.Afr.-India.—FIG. E71,4. **S. bronni* (KRAUSS), S.Afr.; 4a-c, LV ext., int., RV int., $\times 0.5$ (Krauss, 1850).

Trigonastarte BIGOT, 1895 [10] [**Astarte trigonalis* J. DE C. SOWERBY, 1824; OD] [= *Opisastarte* FRECH & MEYER, 1922 (type, *Astarte trigonalis* SOWERBY, 1824; SD CHAVAN, herein)]. Transversely and inequilaterally trigonal, shortened forward. Prosogyrous beaks; somewhat depressed lunule. Concentric ribbing with medioposterior angulation. One strong RV trigonal and 2 LV cardinals, striated on their lateral faces; poorly marked anterior right and posterior left laterals. Crenulated margin. *M.Jur.*, W.Eu.-Indon.—FIG. E72,5. **T. trigonalis* (SOWERBY), Bajoc., W.France; 5a-c, RV ext., int., LV int., $\times 0.7$ (50, 870).



Family CRASSATELLIDAE Férussac, 1822

[=Crassatellitidae DALL, 1895]

Subquadrangular to trigonal in outline, rounded in front, more or less truncated posteriorly. Concentrically ribbed to smooth. Internal layer of radial riblets straight and continuous (so far as observable). Ligament internal, in pit, which commonly obliterates upper part of tooth 4b; narrowly marginal nymphal ridge behind pit. *Dev.-Rec.*

Subfamily CRASSATELLINAE Férussac, 1822

[*nom. transl.* CHAVAN, 1952 (*ex* Crassatellidae FÉRUSSAC, 1822)]

Prosogyrous or orthogyrous beaks; anterior laterals not passing up in front of anterior cardinals. Cardinals divergent. Resilium well developed. *Dev.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Crassatella*.—2. *Pachythaerus*.—3. *Landinia*.—4. *Indocrassatella*.—5. *Salaputium*.—6. *Oriocrassatella*.—7. *Cypricardella*.—8. *Uddenia*.—9. *Crassatina*.—10. *Chattonia*.—11. *Talabrica*.—12. *Eucrassatella*.—13. *Hybolophus*.—14. *Spissatella*.—15. *Bathytormus*.—16. *Anthonya*.

Crassatella LAMARCK, 1799 [**Maetra cygnaea* LAMARCK, 1799 (*non* CHEMNITZ, 1782) (= *C. gibba* LAMARCK, 1801 = *Venus ponderosa* GMELIN, 1791); SD SCHMIDT, 1818] [= ?*Crassatellites* KRUEGER, 1823 (type, *C. sinuatus* (*non* *Crassatella sinuata* LAMARCK, 1818); M) (*nom. dub.*); *Roissya* "LESSON" SCHAUFUSS, 1869]. Subtrapezoidal, thick; prosogyrous beaks. Concentric ribbing and posterior angulation. Lunule and escutcheon deeply sunken. Resilial pit large, but not reaching lower margin of plate. Scars broad, anterior one reniform, posterior one ovate and truncate. Valve margins finely crenulate. *M.Cret.(Turon.)-Mio.*, Eu.-N.Am.

C. (Crassatella) [1]. High, irregular, with more or less vanishing ribs. Short anterior laterals. Post-

FIG. E72. Astartidae (Opinae) (p. N571-N573).




FIG. E73. Crassatellidae (Crassatellinae) (p. N573-N574).

resilial ridge prolonged by posterior laterals. *U. Cret. (Turon.)-Mio. (Helvet.)*, Eu.-N.Am.—FIG. E73,5. *C. (C.) ponderosa* (GMELIN), M.Eoc., France (Paris basin); 5a-c, LV ext., int., RV int., $\times 0.5$ (Chavan, n; Deshayes, 1837).

C. (Landinia) CHAVAN, 1952 [3] [**C. landinensis* NYSTR, 1843; OD]. Transversely elongate, obliquely truncate posteriorly; compressed. Rounded ribs. Anterior teeth long, 2 slightly oblique forward. Strong postresilial ridge, prolonged by remote posterior laterals. *U. Cret. (Senon.)-U. Eoc.*, Eu.-N. Am.-?Palestine.—FIG. E73,1. **C. (L.) landinensis* NYSTR, L.Eoc., France (Paris basin); 1a,b, LV ext., int., $\times 0.7$ (Deshayes, 1837).

C. (Pachythaerus) CONRAD, 1869 [2] [**C. vindinensis* D'ORBIGNY, 1843; OD]. With prominent dorsal slope and well-marked posterior angulation. Ribs commonly lamellar. 3b trigonal, pointed; 4b curved, continuing far below pit. Postresilial ridge almost fused with margin of escutcheon above laterals. *M. Cret.-M. Eoc.*, Eu.-N.Am.-Japan.—FIG. E73,4. **C. (P.) vindinensis* D'ORBIGNY, L.Turon., W.France; 4a, LV ext., $\times 0.75$; 4b,c, LV int., RV int., $\times 1.4$ (695; Chavan, n).

Anthonya GABB, 1864 [16] [**A. cultriformis*; M]. Narrowly and transversely inequilateral, flattened; anteriorly rounded, posteriorly much elongated and tapering. Suborthogyrous beaks. Oblique and narrow lunule, flattened escutcheon. Broad obsolete upper anterior lateral, 2 well-developed cardinals, largely oblique resilium and long posterior lateral. *Cret., ?Eoc.*, N.Am.-W.Eu.-Afr.-Asia-Australia-Japan.—FIG. E73,2. **A. cultriformis*, *U. Cret.*, USA (Calif.); 2a-c, RV ext., hinge, LV hinge, $\times 1$ (333).

Bathytormus STEWART, 1930 [15] [**Crassatella protexta* CONRAD, 1832; OD] [= *Crenocrassatella* HABE, 1951 (type, *Crassatella foveolata* SOWERBY, 1870; OD) (subj.)]. Transversely subtriangular, anterior side rounded, posterior attenuated, in many shells ventrally narrowed. Very small, feebly prosogyrous beaks. Anterior RV lateral laminar, cardinals oblique, very large pit pushing them forward and extending downward to margin of plate and with thin ridge behind it; remote strong laminar laterals. Large scars. Valve margins usually crenulate. *U. Cret.-Rec.*, W.Eu.-N.Am.-NE.Mex.-Asia (India-Japan)-Ghana.—FIG. E73,3. **B. protextus* (CONRAD), M.Eoc., USA (Ala.); 3a-c, LV ext., int., RV int., $\times 0.7$ (Harris, 1919).

Crassatina KOBELT, 1881 [**Crassatella triquetra* "SOWERBY" REEVE (1842) 1843; OD] [= *Crassatina* WEINKAUFF, 1881 (*nom. nud.*)]. Subtriangular to subquadrate, rather small; prosogyrous beaks. Concentric ribs divided or vanishing backward. Narrow lunule and escutcheon. Anterior RV cardinal more or less prolonging laterals; posterior narrow cardinals and elongate laterals. Trigonal resilium. *Paleoc.-Rec.*, Eu.-Asia (Japan)-Afr.




FIG. E74. Crassatellidae (Crassatellinae) (p. N574-N577).




FIG. E75. Crassatellidae (Crassatellinae) (p. N575-N576).

C. (Crassatina) [9]. Valve margins crenulate; shell rather thick. Anterior RV cardinal in prolongation of lateral; *3b* slightly curved forward; well-marked ligamentary ridge in prolongation of laterals. *U.Eoc.-Rec.*, Eu.-Afr.-Asia.—FIG. E74,1a. **C. (C.) triquetra* (REEVE), Rec., W.Afr.; RV ext., $\times 1$ (Reeve, 1843).—FIG. E74,1b. *C. (C.)* sp. cf. *C. bronni* (WOOD), U.Eoc., Eng.; RV hinge, $\times 0.7$ (100).

C. (Chattonia) MARWICK, 1928 [10] [**C. animula*; OD]. Valve margins smooth. Anterior RV cardinal more distinct than lateral; *3b* curved forward; ligamentary ridge and posterior laterals separated. *Paleoc.-Rec.*, W.Eu.-Australia-N.Z.—FIG. E74,2a-c. **C. (C.) animula* MARWICK, Oligo., N.Z.; 2a-c, LV ext., int., RV int., $\times 4$ (593).—FIG. E74,2d. *C. (C.) bartonensis* (WOOD), U.Eoc. (Barton.), Eng.; RV hinge, enl. (100).

Cypricardella HALL, 1858 [7] [**C. subelliptica*; SD MILLER, 1889]. Subtrapezoidal, flattened, rather small, with somewhat lamellar concentric ribs. Hinge plate narrow; laterals present, long and straight; anterior RV and posterior LV cardinals thin; *3b* stout, virguliform, 2 more or less fused to termination of *A11*. Resilium in front of flat nymph. Elliptical small scars. *Dev.-Carb.*, W. Eu.-N.Am.-India.—FIG. E75,1. **C. subelliptica*, Miss., Iowa; 1a,b, topotypes, LV, RV hinges, $\times 20$, $\times 12$ (Newell, n).—FIG. E75,3. *C.* sp., L.Dev.,






FIG. E76. Crassatellidae (Crassatellinae) (p. N576).

Ger.; LV int., diagr. (Haffer, 1959).—FIG. E75, 2. *Astartella vera* HALL, U.Penn., Tex.; LV int., $\times 2$ [for comparison with *Cypricardella*] (Newell, n).—FIG. E74, 7. *C. baudeti* CHAVAN, Carb., Belg.; 7a, RV ext., $\times 1.3$ (Hall, 1858); 7b,c, RV and LV hinges, $\times 1.3$ (109).

Eucrassatella IREDALE, 1924 [**Crassatella kingicola* LAMARCK, 1805; OD]. Transversely subtrigonal and inequilateral, large, thickened; with rounded ventral margin. Concentric ribs vanishing ventrally, stronger and closer toward more or less orthogyrus beaks. Cardinals strong, posterior ones straight in front of large pit which extends downward to lower margin of plate; laterals large. Inner margin smooth. *Paleoc.-Rec.*, Antarctic-N.Z.-W.Eu.-Australia-N.Am.-S.Am.

E. (Eucrassatella) [12]. Rounded, with almost prosogyrous beaks; posterior side truncate. Anterior laterals elongated, cardinals oblique. *Paleoc.*, W.Eu.; *Oligo.-Rec.*, N.Z.-Australia.—FIG. E74, 6. **E. (E.) kingicola* (LAMARCK), Australia; 6a-c, RV ext., RV and LV hinges, $\times 0.7$ (509).

E. (Hybolophus) STEWART, 1930 [13] [**Crassatella gibbosa* SOWERBY, 1832; OD]. Flattened, with somewhat opisthogyrous beaks; shell tapering backward. Very short anterior laterals; 3a well separated from lunular margin. *Mio.-Rec.*, N.Am.-S.Am.—FIG. E74, 5. **E. (H.) gibbosa* (SOWERBY), Rec., S.Am.(W.Coast); 5a-c, LV ext., int., RV int., $\times 0.7$ (Reeve, 1841, 1843).

E. (Spissatella) FINLAY, 1926 [14] [**Crassatella trilli* HUTTON, 1873; OD]. Flattened, with almost orthogyrus flat beaks; shell acuminate, scarcely truncate posteriorly. Narrow lunule. Marginal 3a. *M.Eoc.-Rec.*, N.Am.-Australia-N.Z.-Antarctica.—FIG. E76, 2. **E. (S.) trilli* (HUTTON), L.Mio., N.Z.; 2a,b, RV ext., int., $\times 0.75$; 2c, RV ext., $\times 0.75$ (Suter, 1914).

Indocrassatella CHAVAN, 1952 [4] [**Crassatella indica* E. SMITH, 1895; OD]. Subtrapezoidal, inflated, moderately thin. Concentric regular rounded ribs; no dorsal angulations. 3a almost hidden under lunular margin; resilial pit small, behind upper part of 4b. Long, narrow posterior ridge and LV lateral lamina. Very small, rounded, partly truncated scars. Valve margins finely crenulate. *Rec.*, Arabian Sea.—FIG. E76, 3. **I. indica* (SMITH); 3a-c, LV ext., int., both valves (dorsal), $\times 1$ (Alcock & Anderson, 1897).

Oriocrassatella ETHERIDGE, JR., 1907 [**O. stokesi*; OD] [= *Procrassatella* YAKOVLEV, 1928 (subj.) (type, *Schizodus planus* GOLOVINSKY, 1869)]. Transversely inequilateral, elongated backward. Hinge with 2 strong bifid, 3a long, 3b narrow, 4b curved in front of rather large pit, with post-resilial ridge isolating upper part of ligament. Long posterior laterals. *U.Carb.-Perm.*, Aus.-Asia-Eu.-New S. Wales - Australia-Indon.-USSR-Greenl.—FIG. E76, 4a,b. *P. plana* (GOLOVINSKI), Perm., USSR; 4a,b, RV int., LV int., $\times 0.7$ (1010).—FIG. E76, 4c. **O. stokesi*, W.Australia; RV ext., $\times 0.7$ (Dickins, 1956).

Salaputium IREDALE, 1924 [5] [**Crassatella fulvida* ANGAS, 1871; OD]. Subtrigonal, ventrally rounded, flattened; rather small. Concentric ribbing. Resilial pit adjacent to apophysis of posterior cardinal, not reaching lower margin of plate. Laterals very long, especially anterior ones. Anterior scar large, subreniform. Valve margins finely crenulate. *Neog.-Rec.*, Australia-Indon.—FIG. E76, 1. **S. fulvidum* (ANGAS), Rec., Australia; 1a-c, LV ext., int., RV int., $\times 2$ (Angas, 1863; Chavan, n).

Talabrica IREDALE, 1924 [11] [**Crassatella aurora* ADAMS & ANGAS, 1863; OD]. Transversely subelliptical, compressed, with small orthogyrus beaks. Concentric ribs not divided backward but




FIG. E77. Crassatellidae (Scambulinae) (p. N577-N578).

irregular ventrally. Escutcheon narrow, ill-defined. Cardinals oblique, 4b not narrowed by pit; anterior laterals elongate, clearly distinguishable from cardinals, posterior laterals remote. Margin feebly crenulate. *U.Plio.-Rec.*, N.Z.-Australia.—FIG. E74,3. **T. aurora* (ADAMS & ANGAS), *Rec.*, Tasmania; 3a, LV ext., $\times 1$ (Angas, 1863); 3b,c, LV int., RV int., $\times 1.5$ (Chavan, n).

Uddenia STEPHENSON, 1941 [8] [**Gouldia conradi* WHITFIELD, 1885; OD]. Subquadrate to subtrigonal, small; anteriorly rounded, posteriorly attenuated and truncated. Concentric, spaced, narrow and irregular ribs. Orthogyrous beaks. Marginal narrow anterior, large and grooved posterior cardinals; long and remote posterior laterals. Small scars. Valve margins grooved and smooth. *U.Cret.*, N.Am.—FIG. E74,4. **U. conradi* (WHITFIELD), Emscher., USA (N.Car.); 4a, LV ext., $\times 2.5$; 4b,c, LV and RV hinges, $\times 8$, $\times 5$ (887).

Subfamily SCAMBULINAE Chavan, 1952

Strongly opisthogyrous, or orthogyrous beaks; anterior laterals reaching beaks in

front of anterior cardinals. Subparallel, narrow cardinals; reduced resilium in narrow pit (109). *L.Cret.-Rec.*

Arrangement of generic taxa by CHAVAN.—1. *Scambula*.—2. *Remondia*.—3. *Mediraon*.—4. *Stearnsia*.—5. *Crassinella*.

Scambula CONRAD, 1869 [1] [**S. perplana*; M]. Subtrigonal, inequilateral, much compressed, with convex ventral and concave posterior margins. Pointed beaks. Cardinals approximate and striated; short resilium and ligamentary support above and behind top of LV posterior cardinal; broad and very long laterals on both valves. Small anterior scar. Valve margins crenulate. *U.Cret.*, N.Am.—FIG. E77,3. **S. perplana*, Senon., USA (Tenn.); 3a,b, LV ext., int., $\times 2$ (951).

Crassinella GUPPY, 1874 [5] [**Crassatella martinicensis* d'ORBIGNY in SAGRA, 1853; M] [= *Pseudoriphyla* FISCHER, 1887 (obj.); *Gouldia* AUCT. (non C. B. ADAMS, 1847)]. Trigonal-rounded, short, compressed, small; anterior side rounded, posterior somewhat longer, angular at its end. Concentric lamellar undulations. Narrow lunule; posterior margin strongly concave, bounded by broad escutcheon. Cardinals long and slender, in front of deep, large resilium protruding backward; very long anterior RV and posterior LV laterals. Smooth inner margin. *M.Eoc.-Rec.*, Am.-W.Indies.—FIG. E77,4a. **C. martinicensis* (d'ORBIGNY), *Rec.*, Cuba; RV ext., $\times 10$ (d'Orbigny in Sagra, 1853).—FIG. E77,4b,c. *C. branneri* (ARNOLD), Pleist., USA (Calif.); 4b,c, RV and LV hinges, enl. (100).

Remondia GABB, 1869 [**R. furcata*; M]. Transversely subtrapezoidal, anteriorly rounded, posteriorly elongated, with sinuous truncation. Concentric external undulations and medioposterior angle. Cardinals and laterals curved. Posterior part of 4b on minor elevation. Inner shell margin crenulate. *L.Cret.*, W.Eu.-N.Am.

R. (Remondia) [2]. Much elongated posteriorly. Teeth arcuate; with nymphal ridge and another one in front of it, which divides resilium. Lunule and escutcheon excavated. *L.Cret.*, W.Eu.-N.Am.—FIG. E77,5. **R. (R.) furcata* GABB, *L.Cret.*, Mexico; 5a,b, LV ext., hinge, $\times 0.75$, $\times 0.7$ (333, 877).

R. (Mediraon) VOKES, 1946 [3] [**M. divaricatum*; OD]. Shorter backward, much flattened. Resilium not divided, barely separated from nymphal ridge. *L.Cret.*, W.Eu.-Lebanon-N.Am.—FIG. E77,2. **R. (M.) divaricata* (VOKES), Apt., Lebanon; 2a,b, RV ext., hinge, $\times 1.5$ (945).

Stearnsia WHITE, 1887 [4] [**S. robbinsi*; OD]. Subtrigonal, compressed, almost equilateral; laterally angulated; ventrally rounded, posteroventrally narrowed. Concentric undulations on disc, with sharp posterior carina. Long narrow lunule and




FIG. E78. Cardiniidae (p. N578-N580).

longer escutcheon. Beaks angular, appressed, almost orthogyrous. Approximate strong cardinals and long laterals; resilium narrowly separated from ligament by septum. Valve margins smooth. *L.Cret.*, W.Eu.-N.Am.—FIG. E77, 1. **S. robbinsi*, USA(Tex.); 1a,b, LV ext., int., $\times 0.7$ (White, 1887).

Family CARDINIIDAE Zittel, 1881

[emend. Cox, 1961] [=Cypricardiniidae ULRICH, 1897]
[Materials for this family prepared by L. R. Cox and
ANDRÉ CHAVAN except as recorded otherwise]

Ovate or subtrigonal, more or less inequilateral, thick-shelled, equivalve, convexity weak to moderate. Lunule and escutcheon well defined in more typical genera. Ligament opisthodontic, external although deeply sunk in some forms owing to steep slope of escutcheon. Cardinal teeth low, not more than two in each valve, 3b oblique, 2 more or less fused to margin, more or less obsolete in many forms. Laterals strong, obliquely increasing in prominence, posterior ones of RV never duplicated on LV (no PIV) and entirely posterior to ligamental nymphs. Anterior laterals pres-

ent or absent. Adductor scars deep, subequal; accessory scars rarely present. Pallial line simple. Ornament coarsely concentric or lacking. [Marine.] *Ord.-Rec.*

Besides *Cardinia*, a heterodont genus, ZITTEL originally included in the Cardiniidae several nonheterodont genera which now are referred to the Anthracosiidae and Pachycardiidae (201).

Cardinia AGASSIZ, 1841 [**Unio listeri* J. SOWERBY, 1817; SD ICZN, Op. 292] [=*Thalassides* BERGER, 1833 (suppressed by ICZN); *Ginorga* GRAY, 1840 (*nom. nud.*); *Sinemuria* DE CHRISTOL, 1841 (suppressed by ICZN); *Cardinea* STUTCHBURY, 1842 (*nom. null.*); *Pachyodon* STUTCHBURY, 1842 (*non* VON MEYER, 1838); *Dihora* "Gray" ANON., 1842 (obj.); *Thalassites* QUENSTEDT, 1843 (*nom. van.*) (*non* SWAINSON, 1837); *Storthodon* "Brown" ZITTEL, 1881 (*non* GIEBEL, 1856) (obj.)]. Medium-sized to large, ovate to cuneiform. Lunule and escutcheon bordered by more or less distinct ridges, steeply inclined, in some forms lying almost in plane of commissure of valves, ligamental nymph thus deeply sunk, although not internal; lunule commonly extending to posterior side of beak. Single weak, radially elongated cardinal

tooth (3b) present in RV in some species (but indistinguishable in others), and received in shallow recess in LV. Lateral teeth heavy, LV posterior and RV anterior ones each with tubercle-like termination received in recess between corresponding




FIG. E79. Cardiniidae (p. N579-N580).




FIG. E80. Cardiniidae (p. N579).

tooth and dorsal margin of opposite valve. Ornament of concentric ribs or imbrications, or lacking. *U. Trias. (Carn.) - U. Lias. (Toarc.)*, cosmop. [Records from Bajocian based on poor material and doubtfully acceptable].—FIG. E78,2. *C. hybrida* (SOWERBY), L.Lias., Eng.; 2a-c, RV ext., LV int., RV int., $\times 1$ (Cox, n).

Balantioselena SPEDEN, 1962, p. 96 [**B. gairi*; OD]. Small, subrectangular, ovate or subtriangular; umbo placed well forward, prosogyrous; lunule deep, no distinct escutcheon; nymph prominent; cardinal teeth well developed, 3b in RV received between 2 and 4b in LV; one anterior lateral in each valve, that of RV tuberculiform; 2 slightly oblique posterior laterals in RV, 1 in LV. *M.Trias. (Ladin.)*, N.Z.—FIG. E79,4. **B. gairi*; 4a-c, LV ext. (holotype), RV int., LV int., all $\times 5.5$ (Speden, 1962).

Cypricardinia HALL, 1859 [*pro C. lamellosa* HALL, 1859; *non C. lamellosa (Sanguinolaria lamellosa)* GOLDFUSS, 1840] [**C. halli* BEUSHAUSEN, 1897; OD] [=?*Synopleura* MEEK, 1871 (type, *Cypricardinia? carbonaria* MEEK, 1871, L. Carb.)]. RV more convex than LV, very inequilateral; modioliform in shape, beaks anterior; outline variable, wider posteriorly; umbonal slope prominent, often obtusely angular; surface with strong imbricating concentric equidistant lamellose undulations and commonly with radiating costellae; valves crossed obliquely by shallow byssal sulcus; dentition (AIII-3a), 3b, 5b, PI, PIII / (AIV), AII-2, 4b, PII. *Ord. (Llandeilo)*, Eu.; *Sil. (Medinan)-L. Perm.*, cosmop.—FIG. E80,1a,b. *C. lamellosa* (GOLDFUSS), M.Dev., Ger.; 1a,b, LV and RV hinges, enl. (Chavan, n, from Beushausen).—FIG. E80,1c,d. **C. halli* BEUSHAUSEN, L.Dev. (L. Helderberg.), USA (N.Y.); 1c,d, LV ext., both valves dorsal, $\times 3$ (373).

Isopristes NICOL & ALLEN, 1953 [**I. crassus*; OD]. Thick, subquadrate, beaks prosogyrate, located nearly at anterior end of dorsal border; no lunule. Sculpture of strong equidistant spaced concentric




FIG. E80A. Cardiniidae (p. N580).

folds and small, closely spaced radial riblets. Anterior scar rather small and deep; posterior larger. LV with 2 oblique teeth, RV with 1 small round anterior lateral; posterior hinge not preserved; apparently a nymph. Inner margin crenate. *U.Trias.* (Nor.), S.Am.(Peru).—FIG. E79,3. **I. crassus*; 3a-c, RV and LV int., LV ext., $\times 0.7$ (Nicol & Allen, 1953).

?*Minepharus* TOKUYAMA, 1958 [**Palaeopharus* (*Minepharus*) *triadicus*; OD]. Shell elliptical, tapering slightly anteriorly; umbo prosogyre; ligament area wide, transversely striate; lunule small but distinct; anterior adductor scar strong; posterior scar larger and less distinct; dentition consists of 1 RV and 2 LV cardinals and anterior and posterior lateral; 2b not clearly separated from margin of lunule; 4b elongated, slender and welded to depressed triangular nymph; socket 3b deep, fairly short; socket PI profound; PIII less distinct than PI; narrow furrow divides anterior lateral AIII; ornament consisting of radial costae and lines of growth. *U.Trias.*, Japan.—FIG. E79,1. **M. triadicus* (TOKUYAMA); Yamaguchi Prefecture; LV int., $\times 1$ (A. Tokuyama, 1958). [NEWELL].

Nidarica COX, 1961 [**Cardinia slatteri* WILSON & CRICK, 1889]. Small-medium in size, *Opis*-like, with strongly prosogyrous, terminal umbones from which prominent, angular carinae pass to extremities of concave ventral margin. Lunule deep, with

angular posterior end; nymphs not quite so deeply sunk as in *Cardinia*. No distinct cardinal teeth, but lunular marginal region of LV projects and is much thickened, this projection being received in corresponding recess in RV. Lateral teeth as in *Cardinia*. Ornament of wide-spaced growth imbrications. *L.Jur.*(*M.Lias.*,*Domer.-U.Lias.*, basal *Toarc.*), Eng.—FIG. E78,3. *N. slatteri* (WILSON & CRICK), basal *Toarc.*; LV ext., $\times 1$ (Cox, n).

Pseudastarte COSSMANN, 1921, p. 17 [**Astarte* (*Pseudastarte*) *emarginata*; M]. Very small, trigonally ovate, beaks at anterior quarter of length; convexity feeble; lunule shallower than in *Cardinia*, escutcheon narrow; cardinal tooth (3b) of RV well developed, bifid; laterals as in *Cardinia*. *L.Jur.*(*Lias.*,*Hettang.*), Eu.(France).—FIG. E79, 2. **P. emarginata* (COSSMANN); RV int., $\times 5.3$ (Chavan, n).

Tellidorella BERRY, 1963 [**T. cristulata*; M] [= *Liortarte* OLSSON, 1964 (type, *L. paphia*; OD)]. Small, solid, flattened, acutely subequilateral, ventrally rounded. Sculpture of laminate concentric ridges, each one cristate on carinations which bound long narrow lunule and escutcheon; intervals radially striate or ridged. Anterior scar reniform, posterior one rounded. Median massive LV cardinal, anterior one almost directly prolonged into long anterior upper lateral; upper and lower laterals separated by socket; distant posterior laterals forming edges of socket; LV hinge with 2 cardinals and long, marginal, anterior and posterior lateral; short external ligament. *Mio.-Rec.*, S. Am.(Ecuador)-N. Am.(Mexico).—FIG. E80A, 2. **T. cristulata*, *Rec.*, Mexico; RV int., $\times 10$ (Keen, n).—FIG. E80A,1. *T. paphia* (OLSSON), *Mio.*, Ecuador; 1a-c, RV int., LV int., RV ext., $\times 6$ (Olsson, 1964).

[This small shell is a "living fossil," having all morphological characters of the Cardiniidae, among which are the right duplicate, V-shaped posterior laterals, lack of distinct 5b, a long AI, no marginal AIV and PIV.]

Torastarte MARWICK, 1953 [**T. bensoni*; OD]. Trigonally ovate, astartiform; deep lunule in front of beak, ligamental nymphs relatively heavy, deeply sunk, with corresponding downward arching of posterodorsal margin. Strong RV tubercular laterals, those of LV marginal, 2, 3b, 4b narrowly oblique and partly covered. Ornament of unequal closely spaced concentric ribs. *U.Trias.*(*Otapir.*=*Rhaet.*)-*L.Jur.*(*Lias.*), N.Z.—FIG. E78,1. **T. bensoni*, *Rhaet.*; 1a-c, LV ext., LV int., RV int., $\times 1$ (593).

Family MYOPHORICARDIIDAE Chavan in Vokes, 1967

[Materials for this family prepared by L. R. Cox and ANDRÉ CHAVAN]

Small to small-medium, equivalve, sub-trigonal to trapezoidal, subequilateral to strongly inequilateral with umbones anterior




FIG. E81. Myophoricardiidae (p. N581-N582).

to mid-length; commonly carinate posteriorly, with posterior end truncate; ligament external, opisthodontic, short; hinge fundamentally with one anterior and one posterior lateral in each valve, and two (perhaps three) cardinals in LV, two cardinals in RV, anterior one small, some cardinals commonly suppressed, so that only one may be present in single valve (either LV or RV) and two in other, or only one in each valve; lateral teeth lamelliform, those of LV formed by projections of dorsal margin and received in sockets between corresponding duplicate laterals of RV and dorsal margin; pallial line entire; surface smooth or with concentric riblets. *Trias*.

Much disagreement has existed as to the systematic position of genera placed in this new family, and consequently, in notations applicable to their cardinal teeth. Provisional association of the family with the Crassatellacea implies that the two cardinal teeth of the LV are interpreted as 2 and 4b and those of the RV as 3a and 3b. ODHNER (683, p. 582), however, considered the hinge to be of "cyrenoid" type, interpreting the teeth of the LV as 2a and 2b, those of the RV as 1 and 3b; he cited a statement by FRECH that a third (very weak) cardi-

nal (4b) can be detected in some specimens of the LV of *Myophoriopsis*. ODHNER therefore considered that the genus should be regarded as an early member of the Corbiculacea. The fact that some specimens of *Myophoriopsis* have only one cardinal in the LV, received between two in the RV (Fig. E81, 1b,c), with this arrangement reversed in other specimens (Fig. E81, 1g,h), is not attributable to hinge inversion (since arrangement of the lateral teeth remains the same) but to great variability in the development of individual teeth and to incomplete preservation. It is owing to this variability that the interpretation of the elements of the hinge presents difficulty.

Myophoricardium VON WÖHRMANN, 1889, p. 226 [**M. lineatum*; M] [= *Myophorocardium* NEUMAYR, 1891 (*nom. null.*)]. Trapeziform to subtrigonal, slightly inequilateral, length only slightly exceeding height; posterior ridge obtuse, no lunule or escutcheon; LV with strong anterior and weak posterior tooth, almost equally divergent and bordering recess for single strong tooth of RV; anterior laterals weak; surface almost smooth. *U. Trias*. (*Carn.*), Eu. (N. Alps-S. Alps-Sicily-Hung.)-SW. Asia (Jordan)-China (Yunnan).—FIG. E81. 2. **M. lineatum*; Carinthia (2a,b), N. Alps (2c,d); 2a, LV ext., 2b, dorsal view, both $\times 1.3$; 2c,d, LV and RV hinges, both $\times 3$ (58).

Myophoriopsis VON WÖHRMANN, 1889, p. 221 [**Myophoria lineata* VON MÜNSTER, 1841, p. 88 (= *Lyrodon lineatus* VON MÜNSTER in GOLDFUSS, 1837-40); M] [= *Myophoriopsis* NEAVE, 1940 (*nom. null.*)]. Subtrigonal, moderately to strongly inequilateral, with prominent posterior carina; lunule and escutcheon well developed; either valve with single strong cardinal tooth in median position, received between 2 divergent cardinals in other valve; cardinal teeth transversely ridged; laterals well developed; flank ornamented with concentric riblets. *M. Trias.* (Ladin.)-*U. Trias.* (Nor.), Eu. (N. Alps-S. Alps-Hung.-Italy-Balkans)-Afr. (Libya)-SW. Asia (Jordan)-Indon. (Sumatra).—FIG. E81, 1a-d. **M. lineata* (VON MÜNSTER), M. Trias. (Ladin.), S. Tyrol; 1a, LV ext., $\times 1.3$; 1b, LV int., $\times 2$; 1c, RV int., $\times 2$; 1d, teeth of RV viewed from above, $\times 2$ (58).—FIG. E81, 1e-h. *M. rosthorni* (BOUÉ), U. Trias. (Carn.), Carinthia; 1e, LV ext., $\times 2$; 1f, dorsal view, $\times 2$; 1g, LV int., $\times 2$; 1h, RV int., $\times 2$ (58).

Pseudocorbula E. PHILIPPI, 1898, p. 168 [**Nucula gregaria* MÜNSTER in GOLDFUSS, 1837, p. 152; SD DIENER, 1923, p. 186] [= ?*Raetolucina* OSSWALD, 1930 (type, *Corbula alpina* WINKLER, 1859; OD); *Myophoriaemorphis* MAZAROVICH, 1939 (no diagnosis) (type, *Cucullaea nuculiformis* ZENKER)]. Transversely subtrigonal, slightly to strongly inequilateral; with or without weak posterior carina; lunule and escutcheon present; LV with strong anterior cardinal and in some specimens weak posterior cardinal; RV with single strong posteriorly directed cardinal received in recess between LV cardinals, accompanied by well-developed long laterals duplicated in RV, teeth without transverse ridges; surface smooth. *L. Trias.*-*U. Trias.*, Eu. (Ger.-Spain-S. Alps-Sardinia)-S. Am. (Brazil-Uruguay).—FIG. E81, 3. **P. gregaria* (VON MÜNSTER), M. Trias., M. Muschelkalk, W. Ger.; 3a-c, LV ext., int. and dorsal view; 3d, RV int., all $\times 3$ (Hohenstein, 1913).

Family HIPPOPODIIDAE Cox, new family

[Materials for this family prepared by L. R. Cox with addition by AURÉLE LA ROCQUE]

Medium-sized to large, thick-shelled, irregularly reniform or pyriform, gibbose, with anteriorly placed, terminal or subterminal, prosogyrous beaks, well separated in full-grown specimens; valve margins closed; anisomyarian, posterior adductor scar moderately large, anterior scar smaller, deep, situated in anteroventral lobe; pallial line virtually entire; internal marginal region much broadened anteroventrally by growth laminae, edges of which form rugose




FIG. E82. Hippopodiidae (p. N583).

area continued by broad hinge plate bearing cardinal teeth and recesses as follows: adjacent to nymph in RV, broad tooth which is triangular and ridgelike in some specimens but rectangular and flat-topped in others, and anterior to which is broad, shallow recess for similar tooth of LV; RV with weak posterior lateral tooth, LV with or without posterior lateral; ligament external, opisthodontic; nymph strong, arched; exterior of shell with very irregular and pronounced growth rugae. ?*Dev.*, *L.Jur.-U.Jur.*

The single cardinal tooth of the RV is presumably to be interpreted as *3b* and the main cardinal of the LV as *2*. An angular edge projecting from the face of the nymph in some LV's is possibly to be regarded as a second cardinal (*4b*), but the dentition is very variable and the interpretation of minor ridges doubtful.

This family is known only by its type genus,¹ which is remarkable for its discontinuous distribution. Abundant in northwestern Europe (particularly in England) from the Hettangian to the Pliensbachian stages of the Lower Jurassic, it has otherwise been met with only in the uppermost Jurassic of one area in East Africa. Although the East African species was originally made the type of a genus named *Epihippopodium*, differences between it and Lower Jurassic specimens are of no more than specific importance. }

Hippopodium J. SOWERBY, 1819, p. 91 [**H. ponderosum*; M] [= *Epihippopodium* DIETRICH, 1933, p. 71 (type, *E. quenstedti*)]. Characters of family. *L.Jur.* (Hettang.-Pliensbach.), Eu.; *U.Jur.* (Tithon.), E.Afr.—FIG. E82, 1. **H. ponderosum*, *L.Jur.* (Pliensbach.), SW.Eng.; *1a, b*, LV ext. and dorsal view, $\times 0.8$ (202); *1c-e*, int. of 2 RV's showing variation in cardinal dentition and 1 LV, all $\times 0.82$ (Cox, n).

?**Tusayana** STOYANOW, 1948 [**T. cibola*; OD]. Nearly circular, beaks almost in middle of hinge line; surface smooth, umbones moderate, beaks prosogyre, lunule small; ligament external; hinge plate thick; LV with 2 cardinal teeth, one of which is bifid, other long and simple, lying behind anterior adductor muscle; RV with 1 amorphous cardinal tooth; 1 posterolamellar tooth in each valve. *Dev.* (Island Mesa beds, Jerome F.), USA (Ariz.).—FIG. E83, 1. **T. cibola*; *1a, b*, LV int., RV int., $\times 1$ (Stoyanow, 1948). [LAROCQUE]

¹ Inclusion of the Paleozoic *Tusayana* in this family is very dubious.—Ed.




FIG. E83. Hippopodiidae (p. N583).

Superfamily CARDIACEA Lamarck, 1809

[*nom. transl.* GILL, 1871 (ex family Cardiacea GOLDFUSS, 1820) (=cardiacées LAMARCK, 1809)] [Materials for this superfamily prepared by MYRA KEEN]

Sculpture normally radial, with change of ribbing pattern in most forms on posterior slope; hinge with two conical cardinal teeth, those in LV of unequal size, anterior larger, in RV fused to some extent; lateral teeth distant from cardinals, anterior laterals wanting in some groups. Pallial line entire in marine forms; some brackish-water forms with pallial sinus and long siphons. *U.Trias.-Rec.*

Family CARDIIDAE Lamarck, 1809

[*nom. correct.*, BRODERIP, 1839 (ex Cardiacea GOLDFUSS, 1820) (=cardiacées LAMARCK, 1809)]

Ligament parivincular, external, typically short. Adductor scars subequal. Hinge with two nonbifid cardinal teeth in either valve, cruciform in arrangement; lateral teeth present, one anterior, one posterior in LV, two anterior, one posterior in RV. *U.Trias.-Rec.*

Subfamily CARDIINAE Lamarck, 1809

[*nom. transl.* STOLICZKA, 1870 (ex family Cardiacea GOLDFUSS, 1820) (=cardiacées LAMARCK, 1809)]

Semicircular to quadrangular or, rarely, elliptical; rib ornamentation along rib crests, as beading or furrowing, or in intercostal spaces, never arising from sides of ribs; posterior margin digitate or crenulate;

hinge, with few exceptions, nearly straight (deviating less than 25 degrees from straight line), relatively long, posterior cardinal in LV elevated. *U.Trias.-Rec.*

Cardium LINNÉ, 1758 [**C. costatum*; SD CHILDREN, 1823] [= *Bucardites* VON SCHLOTHEIM, 1820 (*nom. van.*); *Cordium* GISTL, 1848 (*nom. null.*); *Tropidocardium* RÖMER, 1865 (*obj.*)]. Ribs either




FIG. E84. Cardiidae (Cardiinae) (p. N585-N586).

- keeled or spinose; cardinal *4b* elevated (458). *Mio.-Rec.*, S.Eu.-W.Afr.
- C. (Cardium)**. Shell gaping at posterior margin; cardinal *4b* vertically above 2 or nearly so. *Mio.-Rec.*, France-W.Afr.—FIG. E84,2. **C. (C.) costatum* LINNÉ, *Rec.*, W.Afr.; *2a-c*, LV ext., int., RV int., $\times 0.25$ (Tryon).
- C. (Bucardium)** GRAY, 1853 [**Cardium ringens* BRUGUIÈRE, 1789; SD VON VEST, 1875] [=Ringicardium FISCHER, 1887 (obj.)]. Posterior ribs strongly digitate; no posterior gape. *L.Mio.-Rec.*, S.France-W.Afr.—FIG. E84,3. **C. (B.) ringens* BRUGUIÈRE, *Rec.*, W.Afr.; *3a,b*, LV ext., int., $\times 1$ (Tryon).
- Acanthocardia** GRAY, 1851 [**Cardium aculeatum* LINNÉ, 1758; SD STOLICZKA, 1870] [=Orbis DE BLAINVILLE, 1825 (non MÜLLER, 1767, obj.); M]; *Archicardium* SANDBERGER, 1863 (obj.); *Eucardium* FISCHER, 1887 (obj.); *Sphaerocardium* COEN, 1933 (type, *Cardium paucicostatum* SOWERBY, 1841, non DESHAYES, 1838; SD KEEN, 1937); *Sphaerocardium* COEN, 1933 (nom. null.). Oblique-quadrate; cardinal teeth in LV partially fused at base; ribs nodose to spinose. *U.Cret.-Rec.*, Eu.-Asia-N.Am.-S.Am.
- A. (Acanthocardia)**. Ribs spinose (458). *U.Oligo.-Rec.*, S.Eu.-Medit.—FIG. E84,5. **A. (A.) aculeata* (LINNÉ), *Rec.*, Medit.; *5a,b*, LV ext., int., $\times 0.5$ (Tryon).
- A. (Agnocardia)** STEWART, 1930 [**Cardium clai-bornense* ALDRICH, 1911; OD]. Ribs numerous, flat-topped, with hollow, A-shaped spines (892). *Eoc.-Mio.*, N.Am.-S.Am.
- A. (Incacardium)** OLSSON, 1944 [**Cardium mellisum*; OD]. Posterior area set off by sinuation; ribs with A-shaped spines on central and anterior slopes; hinge relatively short. *U.Cret.*, Peru.—FIG. 84,10. **A. (I.) mellisa* (OLSSON); *10a,b*, RV ext., int., $\times 1$ (Olsson).
- A. (Rudicardium)** COEN, 1915 [**Cardium tuberculatum* LINNÉ, 1758; SD KEEN, 1937]. Ribs noded. *Mio.-Rec.*, Medit.
- A. (Schedocardia)** STEWART, 1930 [**Cardium hatcherigbeense* ALDRICH, 1886; OD] [=Africofragum EAMES, 1957 (type, *Fragum (A.) newtoni*; OD)]. Less oblique than *A. (Acanthocardia)*; spines on ribs weak to obsolete, especially anteriorly (892). *Paleoc.-Eoc.*, SE.Asia-Afr.-Eu.-N.Am.-S.Am.
- Granocardium** GABB, 1869 [**Cardium carolinum* D'ORBIGNY, 1844; SD STEWART, 1930]. Smooth-ribbed to spinose, with 1 to 3 intercalary ribs between primaries or with internal pits in intercostal spaces (892). *L.Cret.-U.Cret.*, N.Am.-S.Am.-Afr.-Eu.-Asia-N.Z.
- G. (Granocardium)**. Elliptical, hinge nearly straight, wide; intercalary ribs 2 to 3. *L.Cret.-U.Cret.*, cosmop.—FIG. E84,11. **G. (G.) carolinum* (D'ORBIGNY), *U.Cret.*, France; LV ext., $\times 1$ (Orbigny).
- G. (Criocardium)** CONRAD, 1870 [**Cardium dumosum* CONRAD, 1870; SD STOLICZKA, 1871] [=Criocarpium BÖHM, 1884 (nom. null.); Cardea WHITFIELD, 1885 (obj.)]. Quadrate, hinge straight, relatively long; intercalary ribs tending to be single rows of small spines between ribs. *U.Cret.*, N.Am.-Eu.—FIG. E84,1. **G. (C.) dumosum* (CONRAD), USA(N.J.); *1a-c*, RV ext., LV int., RV int., $\times 1$ (Wade).
- G. (Ethmocardium)** WHITE, 1880 [**Cardium speciosum* MEEK & HAYDEN, 1857 (non ADAMS & REEVE, 1850) (=C. whitei DALL, 1900); OD]. Small, thin, intercostal spaces pitted within. *U.Cret.*, N.Am.-S.Pac.—FIG. E84,6. **G. (E.) whitei* (DALL), USA(Mont.); LV int., $\times 2$ (Keen).
- Loxocardium** COSSMANN, 1886 [**Cardium formosum* DESHAYES, 1858; SD CROSSE, 1886]. Nearly equilateral, posterior margin slightly truncate; posterior ribs notched; rib sculpture of fine looped cross-threads or A-shaped nodes; hinge relatively short (458). *Eoc.-Mio.*, Eu.—FIG. E84,9. **L. formosum* (DESHAYES), *Eoc.*, France; *9a,b*, LV ext., int., $\times 2$ (Deshayes).
- Parvicardium** MONTEROSATO, 1884 [**Cardium parvum* PHILIPPI, 1844 (non DA COSTA, 1778) (=C. exiguum commutatum BUCQUOY, DAUTZENBERG & DOLLFUS, 1892); SD CROSSE, 1885]. Small, hinge weak, cardinal teeth minute, fused at base in LV; ribs sculptured with heavy cross-threads or spines; resembling *Plagiocardium (Papillicardium)* but with weaker hinge and sharper sculpture (458). *Eoc.-Rec.*, Eu.-N.Am.
- Plagiocardium** COSSMANN, 1886 [**Cardium granulosum* LAMARCK, 1805; SD CROSSE, 1887]. Elliptical-oblique, hinge arched; rib sculpture of bead-like nodes. *Paleoc.-Rec.*, Eu.-Afr.-E.Indies-N.Z.
- P. (Plagiocardium)**. Shell medium in size; cardinal teeth nearly equidistant between lateral teeth (458). *Paleoc.-Mio.*, Eu.-E.Indies.
- P. (Maoricardium)** MARWICK, 1944 [**Cardium spatiosum* HUTTON, 1873; OD]. Shell large, heavy; anterior section of hinge shorter than posterior (598). *Oligo.-Rec.*, Afr.-E.Indies-N.Z.—FIG. E84,8. **P. (M.) spatiosum* (HUTTON), Plio., N.Z.; LV hinge, $\times 0.5$ (598).
- P. (Papillicardium)** SACCO, 1899 [**Cardium papillosum* POLI, 1795; OD]. Small, resembling *Parvicardium* but with heavier hinge and with beaded rather than spinose sculpture on ribs (458). *Eoc.-Rec.*, Eu.-W.Asia.
- Septocardia** HALL & WHITFIELD, 1877 [**S. typica*; OD] [=Pascoella COX, 1949 (type, *P. peruviana*; OD)]. Thick-shelled, ribs beaded, incremental lines conspicuous in interspaces as cross-striae; ligament inserted in broad and slightly oblique furrow bordered by nymph with pointed summit; hinge plate wide, anteriorly forming deep cavity for reception of anterior adductor muscle; anterior lateral teeth small, placed at junction of muscle-




FIG. E85. Cardiidae (Cardiinae) (p. N585-N586).

cavity flange and hinge plate proper, cardinal teeth with 2 salient, *4b* lamellar, sloping obliquely backward from beak, *3a-3b* joined above; posterior lateral teeth well developed; pallial line entire, posterior muscle scar ovate, not impinging on hinge plate; inner margins strongly crenulate. [Hitherto allocated to Carditidae, this group seems rather to be the ancestral stock in Cardiacea, but with close affinities to *Palaeocardita*, which is contemporaneous at some localities in N.Am.] *U. Trias. (Nor.)*, N.Am.-S.Am.—FIG. E85, 1. **S. typica*; *1a-c*, RV ext., LV int., RV int., $\times 1$ (Cox); *1d,e* (Alaska), RV and LV hinges, $\times 1.3$ (Keen, n), from USGS specimens); *1f-h* (USA, Nev.), RV ext., LV and RV hinges, all enl. (Hall & Whitfield).

Vepricardium IREDALE, 1929 [**V. pulchricostatum*; OD]. Like *Bucardium* in outline but larger, posterior ribs not digitate; dorsal margins with smooth areas (458). *U. Cret.-Rec.*, Australia-Afr.-Eu.-S.Am. **V. (Vepricardium)**. Rib sculpture of close-set spines. *Paleoc.-Rec.*, Eu.-Afr.-E.Indies-Australia.—FIG. E84, 12. **V. (V.) pulchricostatum* IREDALE, Rec., Australia; *12a-c*, RV ext., LV int., RV int., $\times 0.6$ (433).

V. (Hedecardium) MARWICK, 1944 [**Cardium waitakiense* SUTER, 1907; OD]. Ribs smooth, finer on posterior area, interspaces linear (598). *Eoc.-L.Mio.*, N.Z.-Australia-Burma.—FIG. E84, 4. **V. (H.) waitakiense* (SUTER), *Oligo.*, N.Z.;

4a-c, LV ext., LV and RV hinges, $\times 0.5$ (598). **V. (Orthocardium)** TREMLETT, 1950 [**Cardium porulosum* SOLANDER, 1766; OD]. Rib-sculpture tending to form vertical frill at rib crest; outline more quadrate than in *V. (Vepricardium)*, hinge line narrower and straighter. *M.Eoc.-U.Eoc.*, Eu.

V. (Perucardia) OLSSON, 1944 [**Cardium bruegeni*; OD]. Ribs with granular nodes. *U. Cret.*, Peru.—FIG. E84, 7. **V. (P.) bruegeni* (OLSSON); *7a,b*, LV ext., hinge, $\times 0.5$ (Olsson).

Subfamily TRACHYCARDIINAE Stewart, 1930

Ovate but not oblique; rib ornamentation normally of spines or imbricating scales along posterior sides of ribs; hinge relatively short, either nearly straight or sharply bent; cardinal teeth unequal in size, *All* with socket above; posterior margin notched to digitate. ?*Eoc.*, *Oligo.-Rec.*

Trachycardium MÖRCH, 1853 [**Cardium isocardia* LINNÉ, 1758; SD VON MARTENS, 1870] [= *Kathocardia* TUCKER & WILSON, 1932 (type, *Cardium (K.) acinense*; OD)]. Outline nearly equilateral, height greater than length; hinge heavy, short, nearly straight. ?*Eoc.*, *Oligo.-Rec.*, tropic Am.

T. (Trachycardium). Sculpture of imbricating scales over entire shell, reduced to beads anteriorly (458). *Oligo.-Rec.*, W.Indies-E.C.Am.-W.C.Am.-S.Am.—FIG. E86, 5. ***T. (T.) isocardia** (LINNÉ), Rec., W.Indies; *5a-c*, LV ext., int., RV int., $\times 0.5$ (*5a,c*, Perry; *5b*, Keen).

T. (Dallocardia) STEWART, 1930 [**Cardium quadragenarium* CONRAD, 1837; OD]. Large shells with ornamentation of nonimbricating spines along posterior sides of ribs (892). *U. Oligo.-Rec.*, N.Am.-S.Am.

T. (Mexicardia) STEWART, 1930 [**Cardium procerum* SOWERBY, 1833; OD]. Ribs scaly in young, nearly smooth in adults; hinge heavy, short; posterior margin somewhat digitate (892). ?*Eoc.*, *Mio.-Rec.*, N.Am.-S.Am.

T. (Phlogocardia) STEWART, 1930 [**Cardium belcheri* BRODERIP & SOWERBY, 1829; OD]. Central ribs with frills along posterior sides, anterior ribs with knobby spines (892). *Mio.-Rec.*, W.C. Am.

Acrosterigma DALL, 1900 [**Cardium dalli* HEILPRIN, 1887; OD]. Elliptical, hinge strongly bent; cardinal teeth unequal in size (458). *U. Oligo.-Rec.*, N.Am.-S.Am.-Eu.-E.Indies-S.Pac.

A. (Acrosterigma). Hinge bent to angle of 60 degrees or more; ribs smooth except for slight pitting along edges of incised intercostal spaces. *Mio.-Plio.*, USA (Fla.)-W.Indies.—FIG. E86, 6. ***A. (A.) dalli** (HEILPRIN), *Plio.*, USA (Fla.); *6a,b*, LV ext., int., $\times 0.25$ (Cooke).

A. (Ovicardium) MARWICK, 1944 [**Trachy-*

cardium (O.) *rossi*; OD]. Ovate, intercostal spaces cross-striate, with vertical sides; hinge with oblique groove behind 4b (598). *Plio.*, N.Z.-Australia.

A. (*Regozara*) IREDALE, 1936 [**R. olivifer*; OD]. Hinge less angulate than in A. (*Acrosterigma*); sculpture of small scales or cross-threads on posterior sides of ribs. *Neog.-Rec.*, E.Indies-Australia.




FIG. E86. Cardiidae (Trachycardiinae) (5-6, 10); (Fraginae) (2-3, 7, 11), (Protocardiinae) (1, 4, 8-9, 12) (p. N586, N588-N589).

- A. (Vasticardium)** IREDALE, 1927 [**V. nebulosum* (ex MARTYN, nonbinom.); OD]. Resembling *A. (Acrosterigma)* but eaves of ribs more strongly sculptured; posterior margin notched (458). *U. Oligo.-Rec.*, IndoPac.-N.Am.-C.Am.-Eu.
- Papyridea** SWAINSON, 1840 [**Cardium soleniforme* BRUGUIÈRE, 1789; SD GRAY, 1847]. Shell longer than high, gaping at both ends, ribs spinose; hinge short; posterior margin notched (458). *Mio.-Rec.*, tropic Am.—FIG. E86,10. **P. soleniformis* (BRUGUIÈRE), Rec., W.Indies; 10a-c, LV ext., hinge, RV int., $\times 1$ (10a,c, Perry; 10b, Keen).
- Subfamily FRAGINAE Stewart, 1930**
- Posterior slope set off by low ridge to sharp keel, postero-ventral margin angulate; rib ornamentation of threads or imbricating spines, intercostal spaces narrow, cross-striate; hinge short, bent more than 25 degrees from horizontal; cardinal teeth mostly unequal in size. *Oligo.-Rec.*
- Fragum** RÖDING, 1798 [**Cardium fragum* LINNÉ, 1758; T] [= *Hemicardia* SPENGLER, 1799, AUCT. (invalidly proposed)]. Triangular; hinge with cardinals subequal, peg-shaped; rib ornamentation threadlike (458). *Mio.-Rec.*, E.Indies.
- F. (Fragum)**. Lunular area not depressed. *Mio.-Rec.*, E.Indies.—FIG. E86,11. **F. (F.) fragum* (LINNÉ), Rec., E.Indies; 11a,b, RV ext., LV int., $\times 1$ (Keen).
- F. (Lunulicardia)** GRAY, 1853 [**Cardium retusum* LINNÉ, 1758; M] [= *Opisocardium* BAYLE, 1879 (obj.)]. Lunule very deeply impressed. *Pleist.-Rec.*, Japan-IndoPac.
- Corculum** RÖDING, 1798 [**Cardium cardissa* LINNÉ, 1758; $\frac{1}{2}$ SD VON MARTENS, 1870] [= *Cardissa* MEGERLE VON MÜHLFELD, 1811 (obj.); *Hemicardia* FLEMING, 1818 (obj.); *Hemicardium* SCHWEIGGER, 1820 (obj.)]. Foreshortened, posterior slope nearly plane, bounded by carina, of glassy texture; sculpture on ribs faint (458). Rec., E.Indies.—FIG. E86,7. **C. cardissa* (LINNÉ); 7a,b, both valves ant., LV int., $\times 0.5$ (7a, Tryon, 7b, Keen).
- Ctenocardia** H. ADAMS & A. ADAMS, 1857 [**Cardium hystrix* REEVE, 1844 (non LIGHTFOOT, 1786) (= *Fragum symbolicum* IREDALE, 1929); SD DALL, 1900]. With imbricating or tubular spines on ribs; hinge with cardinals subequal, midway between laterals (458). *Mio.-Rec.*, E.Indies-S.Afr.
- C. (Ctenocardia)**. Outline quadrate, not oblique. *Mio.-Rec.*, E.Afr.-E.Indies.—FIG. E86,3. *C. (C.) symbolica* (IREDALE), Rec., E.Indies; 3a,b, RV ext., LV int., $\times 1$ (3a, Tryon; 3b, Keen).
- C. (Afrocardium)** TOMLIN, 1931 [**Fragum (A.) shepstonense*; OD]. Oblique, umbonal ridge indistinct (458). *Pleist.-Rec.*, S.Afr.-E.Indies.
- C. (Microfragum)** HABE, 1951 [**Cardium festivum* DESHAYES, 1854; OD]. Quadrate, with little or no sculpture on ribs. Rec., E.Indies-Japan.
- Trigoniocardia** DALL, 1900 [**Cardium graniferum* BRODERIP & SOWERBY, 1829; OD]. Ovate to quadrate; hinge asymmetric, anterior section shorter than posterior. *Oligo.-Rec.*, tropic Am.
- T. (Trigoniocardia)**. Ovate, mostly small, sculpture beaded, with intercostal striations. *Oligo.-Rec.*, C. Am.—FIG. E86,2. **T. (T.) granifera* (BRODERIP & SOWERBY), Rec., W.Mex.; 2a,b, RV ext., LV int., $\times 2$ (Keen & Frizzell).
- T. (Americardia)** STEWART, 1930 [**Cardium medium* LINNÉ, 1758; OD]. Quadrate, medium in size, ribs and intercostal spaces smooth (892). *L. Mio.-Rec.*, C.Am.
- T. (Apiocardia)** OLSSON, 1961 [**Cardium obovale* SOWERBY, 1833; OD]. Rounded-ovate, ribs nearly smooth. *Plio.-Rec.*, W.C.Am.-S.Am.
- Subfamily PROTOCARDIINAE Keen, 1951**
- Rounded quadrate, beaks nearly central; posterior slope well defined by umbonal ridge, its radial ribs tending to be spinose; hinge long, slightly arched, cardinals 2 and 3b larger, peglike, curved upward, sinuating lower border of hinge plate; pallial line mostly entire, some with small sinus near posterior adductor scar. *U.Trias.(Rhaet.)-Rec.*
- Protocardia** VON BEYRICH, 1845 [**Cardium hillanum* SOWERBY, 1813; SD HERRMANNSEN, 1847] [= *Hassbergia* KRUMBECK, 1939 (invalidly proposed, without type designation)]. Posterior slope with radial ribs in most, remainder of shell with more or less well developed concentric ribs (893, 1006). *U.Trias.-U.Cret.*, Eu.-N.Am.-S.Am.-Afr.
- P. (Protocardia)**. Anterior concentric and posterior radial ribs well developed. *U.Trias.-U.Cret.*, Eu.-Asia-Afr.-N.Am.-S.Am.—FIG. 86,4. **P. (P.) hillana* (SOWERBY), L.Cret., Eng.; 4a-c, LV ext., int., RV hinge, $\times 0.5$ (1006).
- P. (Brevicardium)** STEPHENSON, 1941 [**B. fragile*; OD]. With faint radial ribs crossing concentric sculpture of central and anterior slopes (889). *U.Cret.*, N.Am.-Eu.-Asia.
- P. (Globocardium)** HAYAMI, 1965 [**Cardium sphaeroideum* FORBES, 1845; OD]. Globose, with weak radial ribs. *L.Cret.*, Eu.-Japan.
- P. (Leptocardia)** MEEK, 1876 [**Cardium subquadratum* EVANS & SHUMARD, 1857; SD DALL, 1901]. Small, polished, posterior radial ribs faint; marginal crenulations strong near junction of posterior and central slopes (458). *L.Cret.-U.Cret.*, N.Am.
- P. (Pachycardium)** CONRAD, 1869 [**Cardium spillmani* CONRAD, 1858; SD DALL, 1900]. Relatively short, with high narrow beaks; concentric sculpture fine, radial sculpture covering up to half central slope; hinge thickened (458). *L.Cret.-U.Cret.*, Afr.-Eu.-Asia-N.Am.-S.Am.

- P. (Tendagurium)** DIETRICH, 1933 [**Cardium* (*T.*) *propebanneianum*; SD KEEN, 1937]. Radial sculpture wanting on posterior slope (458). *Jur.-Cret.*, Afr.-Eu.-E.Asia.
- Integricardium** ROLLIER, 1912 [**Cardium dupinianum* D'ORBIGNY, 1844; OD]. Elliptical to elongate-quadrate; sculpture obsolete; hinge as in *Protocardia* (795). *M.Jur.-U.Cret.*, Eu.-N.Afr.-W.Asia-N.Am.
- I. (Integricardium)**. Elliptical, without umbonal ridge. *M.Jur.-U.Cret.*, Eu.-N.Afr.-W.Asia.—FIG. 86,12. *I. (*I.*) *dupiniana* (D'ORBIGNY), U.Cret., France; *12a,b*, both valves dorsal, RV hinge, $\times 0.6$ (d'Orbigny).
- I. (Onestia)** McLEARN, 1933 [**Laevicardium onestae* McLEARN, 1931; OD]. Elongate-quadrate; lateral teeth small; external ligament in deep groove (458). *L.Cret.*, Can.
- Jurassicardium** COSSMANN, 1906 [**Cardium* (*J.*) *axonense*; SD KEEN, 1937]. Posterior-dorsal margin rectangular; posterior ribs not spinose, other radial riblets indistinct (458). *Jur.*, Eu.(France).—FIG. 86,1. **J. axonense* (COSSMANN); *1a-c*, RV ext., int., LV hinge, $\times 2$ (458).
- Nemocardium** MEEK, 1876 [**Cardium semiasperum* DESHAYES, 1858; SD SACCO, 1899] [= *Awadia* ABBASS, 1962 (type, *N. (A.) magharensis*; OD)]. Sculpture radial throughout, strong on posterior slope (459). *L.Cret.-Rec.*, Eu.-Asia-S.Pac.-N.Am.-S.Am.
- N. (Nemocardium)**. Central and anterior slopes nearly smooth, marginal crenulations changing abruptly in size at posterior-ventral junction. *L.Cret.-Rec.*, cosmop.—FIG. 86,9. **N. (N.) semiasperum* (DESHAYES), Eoc., France; *9a-c*, LV ext., int., RV hinge, $\times 1$ (Deshayes).
- N. (Arctopratalum)** KEEN, 1954 [**N. (A.) griphus*; OD]. Ovate-trigonal, hinge short. *Oligo-Mio.*, Japan-W.N.Am.
- N. (Discors)** DESHAYES, 1858 [**Cardium discors* LAMARCK, 1805 (*non* MONTAGU, 1803) (= **C. parisiense* D'ORBIGNY, 1850); T] [= *Hemidiscors* ROVERETO, 1898 (type, *Cardium (H.) rugiferum*; OD)]. Ovate, smaller than *N. (Nemocardium)*; oblique secondary striae crossing ribs of central and anterior slopes, posterior ribs faint. *Eoc.-Mio.*, Eu.-Asia.
- N. (Divaricardium)** DOLLFUS & DAUTZENBERG, 1886 [**Cardium discrepans* BASTEROT, 1825; SD COSSMANN, 1886]. More quadrate than *N. (Discors)*, oblique secondary sculpture on posterior slope; central slope nearly smooth. *Oligo-Plio.*, Eu.-Asia.
- N. (Frigidocardium)** HABE, 1951, p. 147 [**Cardium eos* KURODA, 1929; OD] [= *Erigidocardium* HABE, 1951, p. 152 (*nom.null.*)]. Radial sculpture beaded throughout, alternate ribs higher. *Rec.*, Japan-IndoPac.
- N. (Keenaea)** HABE, 1951 [**N. samarangae* MAKIYAMA, 1934; OD]. Resembling *N. (Pratalum)* but with secondary concentric sculpture on posterior slope. *Oligo.-Rec.*, Japan-W.N.Am.
- N. (Lophocardium)** FISCHER, 1887 [**Cardium cumingii* BRODERIP, 1833; OD]. Fragile, sculpture of faint sinuous radial and concentric threads; periostracum forming laminar keel at junction of posterior and central slopes; cardinal teeth well developed, lateral teeth obsolescent (226). ?*Eoc.*, *Mio.-Rec.*, C.Am.-S.Am.
- N. (Lyrocardium)** MEEK, 1876 [**Cardium lyratum* SOWERBY, 1841; SD DALL, 1900] [= *Amphicardium* VON MARTENS, 1880 (obj.)]. Like *N. (Discors)* but more quadrate, oblique secondary sculpture more widely and evenly spaced (458). *Pleist.-Rec.*, E.Indies.—FIG. 86,8. **N. (L.) lyratum* (SOWERBY), *Rec.*, E.Indies; *8a,b*, LV ext., RV int., $\times 0.5$ (Tryon).
- N. (Microcardium)** THIELE, 1934 [**Cardium peramabile* DALL, 1881; SD KEEN, 1937]. Small, with secondary concentric sculpture on central and anterior slopes both in intercostal spaces and as beads along ribs. *Mio.-Rec.*, C.Am.
- N. (Pratalum)** IREDALE, 1924 [**Cardium thetidis* HEDLEY, 1902; OD]. More equilateral than *N. (Nemocardium)*; radial ribs evident throughout but coarser on posterior slope (459). *L.Cret.-Rec.*, Eu.-N.Z.-Australia.
- N. (Trifaricardium)** HABE, 1951 [**Cardium nomurai* KURODA & HABE, 1951; OD]. Near *N. (Microcardium)* but secondary concentric sculpture on anterior slope only. *Rec.*, Japan.
- N. (Varicardium)** MARWICK, 1944 [**Cardium patulum* HUTTON, 1873; OD]. Heavy, with strong but irregular secondary concentric ridges anteriorly and ventrally overriding radial ribs; ribs of posterior slope without tubercles or spines (598). *Mio.*, N.Z.

Subfamily LAEVICARDIINAE Keen, 1936

Elliptic-oblique; rib ornamentation of looped threads or small nodes, not spines; ribs of posterior slope weaker than those of central and anterior slopes or obsolescent; posterior margin wavy rather than notched; hinge long and arched (line joining laterals and cardinals bends more than 25 degrees); cardinal teeth somewhat unequal in size, anterior left lateral blade-like. *Eoc.-Rec.*

Laevicardium SWAINSON, 1840 [**Cardium oblongum* GMELIN, 1791; SD STOLICZKA, 1871] [= *Liocardium* AGASSIZ, 1846 (*nom.van.*); *Exocardium* OLSSON, 1964 (type, *Cardium ecuadoriale* OLSSON, 1932; OD)]. Ribs faintly to moderately developed, weaker on posterior slope; smooth to threaded or noded. *Eoc.-Rec.*, Eu.-Atl.-Pac.

L. (Laevicardium). Smooth, nongaping; cardinals unequal. *Eoc.-Rec.*, Eu.-W.Atl.-E.Pac.-W.Pac.—FIG. E87,2. **L. (L.) oblongum* (GMELIN), *Rec.*,




FIG. E87. Cardiidae (Laevicardiinae) (p. N589-N590).

Eu.; 2a-c, RV ext., LV int., RV int., $\times 0.5$ (2a-c, 89a; 2b, Keen).

L. (Dinocardium) DALL, 1900 [**Cardium robustum* LIGHTFOOT, 1786; OD]. Ribbing on anterior slope and beaks stronger than in *L. (Laevicardium)*; ornamentation of threadlike nodes crossing ribs (892). *Eoc.-Rec.*, N.Am.-S.Am.

L. (Fulvia) GRAY, 1853 [**Cardium apertum* BRUGUIÈRE, 1789; M]. Thin-shelled, ventricose, slightly gaping posteriorly; cardinal teeth subequal. *Mio.-Rec.*, Japan-E.Indies-Alaska.

Cerastoderma POLI, 1795 [**Cardium edule* LINNÉ, 1758; SD VON MARTENS, 1870] [= *Cerastes* POLI,

1795 (non LAURENTI, 1768); *Edulicardium* MONTEROSATO, 1923 (obj.)]. Oblique-quadrangular, hinge only moderately arched; beaks orthogyrate; ribs moderately strong on posterior slope, sculptured with cross-threads (242). *U.Oligo.-Rec.*, Eu.—FIG. E87,3. **C. edule* (LINNÉ), *Rec.*, Eu.; 3a,b, LV ext., int., $\times 1$ (3a, Tryon; 3b, Keen).

Clinocardium KEEN, 1936 [**Cardium nuttallii* CONRAD, 1837; OD]. Oblique-elliptical, hinge well arched; beaks prosogyrate, ribs weak on posterior slope, ornamented with cross-threads; cardinal 4b weak, left posterior lateral double (458). *U.Mio.-Rec.*, N.Pac.-NW.Atl.—FIG. E87,4. **C. nuttallii* (CONRAD), *Rec.*, USA (Calif.); 4a-c, RV ext., LV int., RV hinge, $\times 0.5$ (Keen).

Serripes GOULD, 1841 (ex BECK, MS) [**Cardium groenlandicum* BRUGUIÈRE, 1789; M] [= *Aphrodite* LEA, 1834 (non HÜBNER, 1816)]. Sculpture almost obsolete; shell medium-sized to large, inequilateral; cardinal teeth weak in adults or absent, laterals also evanescent (458). *Mio.-Rec.*, N.Atl.-N.Pac.-Japan.—FIG. E87,1. **S. groenlandicus* (BRUGUIÈRE), *Rec.*, Arctic O.; 1a-c, LV ext., LV and RV hinges, $\times 0.5$ (Tryon).

Family LAHILLIIDAE Finlay & Marwick, 1937

Orbicular to oblong in outline; surface smooth, without radial sculpture; internal margins smooth, not crenulate; ligament external, on large nymph. Cardinal teeth two in each valve, arranged as in Cardiidae; anterior lateral teeth absent, posterior laterals one in each valve. *U.Cret.-Mio.*

Lahillia COSSMANN, July, 1899 [*pro Iheringia* COSSMANN, April 1899, *nom. correct. pro Theringia* COSSMANN, Jan. 1899 (non KEYSERLING, 1891) et *pro Amathusia* PHILIPPI, 1887 (non FABRICIUS, 1807)] [**Amathusia angulata* PHILIPPI, 1887; SD FINLAY & MARWICK, 1937]. Medium-sized to large, ovate, thin; hinge plate broad, thick. *U.Cret.-Mio.*, S.Am.-N.Z.-Australia-Antarctic.

L. (Lahillia). Pallial line entire (598). *U.Cret.-Mio.*, S.Am.-N.Z.-Australia-Antarctic.—FIG. E88,5. **L. (L.) angulata* (PHILIPPI), *Oligo.*, ?*Mio.*, Chile; 5a,b, RV ext., int., $\times 0.5$ (5a, Philippi; 5b, Wilckens).

L. (Lahilleona) FINLAY & MARWICK, 1937 [**L. neozelanica* MARSHALL & MURDOCH, 1923; OD]. With well-defined rounded pallial sinus (598). *U.Cret.*, N.Z.—FIG. E88,6. **L. (L.) neozelanica* MARSHALL & MURDOCH; 6a-c, LV ext., int., RV hinge, $\times 0.5$ (304).

Family LYMNOCARDIIDAE Stoliczka, 1870

[*nom. transl.* (as Adacnidae), VON VEST, 1875 (ex *Lymnocardiinae* STOLICZKA, 1870)]

Shells ribbed to smooth, small to medium-sized, many with alations on dorsal margin, some with shell gaping. Normal hinge with two cardinals in either valve, obsolescent in some; lateral teeth distant, varying from large to obsolete; pallial line entire or sinuate to a varying extent. [Brackish-water forms, perhaps of polyphyletic origin, characterized by great variability.] *Mio.-Rec.*

The classification here is adapted from an outline by EBERSIN (1965), the most complete modern summary of named taxa; however, differential diagnoses of many nominal taxa are yet unavailable.

Subfamily LYMNOCARDIINAE Stoliczka, 1870

Shells oblique, umbones in front of midline; hinge with well-developed anterior lateral teeth. Pallial line normally entire, valves mostly not gaping. *Mio.-M.Plio.*

Lymnocardium STOLICZKA, 1870 [**Cardium haueri* HÖRNES, 1861; OD] [= *Limnocardium* (nom. null.)]. Long-quadrate; ribs few, smooth or with spines; cardinals 2 and 3b present. *M.Mio.-M.Plio.*, Eu.

L. (*Lymnocardium*). Hinge teeth and sculpture well developed. *M.Mio.* (*Sarmat.*)-*L.Plio.* (*Ruman.*), E.Eu.—FIG. E88, 4. **L.* (*L.*) *haueri* (HÖRNES), *L.Plio.*, Aus.; 4a-c, LV ext., LV hinge, RV int., $\times 0.5$ (Andrussov, 1903).

L. (*Arpadiocardium*) EBERSIN, 1947 [**L.* (*A.*) *peregrinum*; OD]. Hinge nearly edentulous; pallial line slightly sinuate. *L.Plio.* (*Pont.*), S.USSR.

L. (*Bosphoricardium*) EBERSIN, 1947 [**Cardium emarginatum* DESHAYES, 1838 (non DESHAYES, 1820) (= *C. aperturatum* DESHAYES, 1857); OD]. Ovate, ribs numerous, low; cardinal teeth weak. *L.Plio.* (*Pont.*), S.USSR.

L. (*Ecericardium*) EBERSIN, 1947 [**L.* (*E.*) *ecericum*; OD]. Nearly smooth, ribs low; lateral teeth somewhat reduced. *M.Plio.* (*Kimmer.-Kuiálnik.*), S.USSR.

L. (*Euxinocardium*) EBERSIN, 1947 [**L.* (*N.*) *subsymiense* ANDRUSSOV, 1903; OD]. Ovate, ribs triangular in section, cardinals weak. *L.Plio.* (*Pont.*), S.USSR.

L. (*Moquicardium*) EBERSIN, 1947 [**L.* (*M.*) *moquicum* SENINSKY, 1905; OD]. Subrounded to ovate-triangular; ribbed within; pallial line raised anteriorly. *M.Plio.* (*Kimmer.*), S.USSR.

L. (*Nargicardium*) EBERSIN, 1947 [**L.* (*N.*) *nargicardium*; OD]. Ribs widely spaced, triangular in section; hinge with weak cardinals, strong anterior laterals, obsolete posterior laterals. *L.Plio.* (*Pont.*), S.USSR.

L. (*Pannonicardium*) STEVANOVIĆ, 1951 [**L.* (*P.*) *du-*

micici GORIANOVIC-KRAMBERGER, 1899 (fide EBERSIN, 1965)]. *L.Plio.* (*Pont.*), S.Eu.




FIG. E88. Lahillidae (5-6); Lymnocardiidae (Lymnocardiinae) (3-4), (Didacninae) (1,7), (Adacninae) (2,8) (p. N590-N593).

- L. (Tauricardium) EBERSIN, 1947 [**L. subsquamulosum* ANDRUSSOV, 1903; OD]. Ribs few, well developed anteriorly. *Plio.(Pont.-Kimmer.)*, S. USSR.
- Budmania BRUSINA, 1897 [**Adacna meisi* BRUSINA, 1884; SD EBERSIN, 1965]. Ribs few, narrow, crested; hinge weak. *L.Plio.(Pont.)*, S.Eu.
- Eoprosodacna DAVIDASCHVILI, 1934 [**Cardium (E.) karilicum*; M] [= *Limnopappia* SCHLICKUM, 1962 (type, *L. schuetti*; OD)]. *L.Mio.-M.Mio.*, USSR (S.Caucasus).
- E. (Eoprosodacna). *L.Mio.-M.Mio.*, USSR (S.Caucasus).
- E. (Succuridacna) KOROBKOV, 1954 [**Cardium goriense* DAVIDASCHVILI, 1934; OD] [= *Limnopagettia* SCHLICKUM, 1963 (type, *Cardium friabile* KRAUSS, 1852; OD)]. Ribs widely spaced. *L.Mio.-M.Mio.*, USSR (S.Caucasus).
- Horiodacna SABBA STEFANESCU, 1896 [**H. rumana*; M]. Smooth, beaks prosogyrate; shell ribbed internally; hinge with rudimentary cardinal in 1 valve, fossette in other. *L.Plio.(Pont.)*, Eu. (Rumania).
- Limnodacna EBERSIN, 1936 [**L. cristulata*; OD]. Posterior slope set off by several carinate ribs; hinge of LV reduced to 1 weak cardinal. *M.Plio.(Kimmer.)*, S.USSR.
- Pachydacna EBERSIN, 1955 [*pro Natella* EBERSIN, 1949 (non WATSON, 1934)] [**Natella natella* EBERSIN, 1949; M]. Ribs few, hinge heavy. *M.Plio.(Kimmer.-Kuialnik.)*, Euxine Basin, S.USSR.
- Prionopleura EBERSIN, 1949 [**Prosodacna prionopleura* WASSOIEVICH & EBERSIN, 1931 (ex ANDRUSSOV MS); T]. Ribs keeled, crossed by raised regular growth lamellae. *M.Plio.(Kimmer.)*, S.USSR.
- Prosodacna TOURNOUËR, 1882 [**Cardium macrodon* DESHAYES, 1838; OD] [= *Psilodon* COBALESCU, 1883 (non PERTY, 1830); *Pseudoprosodacna* GILLET, 1943 (fide EBERSIN, 1959)]. Ovate, oblique, beaks prosogyrate; exterior ribbing weak, interior ribs strong; hinge with cardinals wanting, laterals *Al*, *All* thick, *Alli* thin; pallial line entire. *L.Plio.-M.Plio.*, S.USSR.
- P. (Prosodacna). *L.Plio.-M.Plio.(Pont.-Kuialnik.)*, S.USSR.—FIG. E88,3. *P. (*P.*) *macrodon* (DESHAYES), *Plio.*, Euxine Basin, S.Eu.-S.USSR, *3a,b*, RV ext., int., $\times 0.4$ (Davidaschvili).
- P. (Metadacna) [**P. metoica* DAVIDASCHVILI, 1930; OD]. *M.Plio.*, Azov area, S.USSR.
- P. (Prosochiasta) EBERSIN, 1959 [**P. prosochiasta* ANDRUSSOV, 1917; OD]. *L.Plio.(Pont.)*, S.USSR.
- P. (Prosodacnomya) EBERSIN, 1959 [**Cardium rostratum* SINZOV, 1900 (non DE KONINCK, 1841) (= *Prosodacna sinzovi* ANDRUSSOV, 1917); OD]. *L.Plio.(Pont.)*, S.USSR.
- Stylodacna SABBA STEFANESCU, 1896 [**Psilodon heberti* COBALESCU, 1883; OD] [= *Stylodacna* DOLLFUS, 1905 (nom. null.)]. Internal ribs heavy, lamellar; hinge without cardinals, laterals small, lamellar. *M.Plio.*, Eu. (Rumania).

Subfamily DIDACNINAE Ebersin, 1962

Umbones subcentral; lateral teeth wanting. *L.Plio.(Pont.)-Rec.*

Didacna EICHWALD, 1838 [**Cardium trigonioides* PALLAS, 1771; SD STOLICZKA, 1870]. Rounded-trigonal, slightly truncate; pallial line entire. *L.Plio.(Pont.)-Rec.*, S.USSR.

D. (Didacna). *Rec.*, Caspian Sea.—FIG. E88,7. *D. (*D.*) *trigonioides* (PALLAS); *7a,b*, LV ext., RV int., $\times 0.5$ (Reeve, 1843).

D. (Crassadacna) EBERSIN, 1962 [**Cardium crassatellatum* DESHAYES, 1838; OD]. *Plio.(Pont.-Kimmer.)*, S.USSR.

D. (Pontalmyra) SABBA STEFANESCU, 1896 [**P. placida*; SD SACCO, 1899] [= *Tschaudia* DAVIDASCHVILI, 1956 (type, *Didacna tschaudae* ANDRUSSOV, 1910; OD)]. *Plio.(Pont.-Kuialnik.)*, S. USSR-S.Eu.

Brachiodacna EBERSIN, 1964 [**Cardium negativum* ANDRUSSOV, 1909; OD]. *L.Plio.(Pont.)*, S.USSR (fide EBERSIN, 1965).

Caladacna ANDRUSSOV, 1917 [**Cardium steindachneri* BRUSINA, 1884; M] [= *Kaladacna* ANDRUSSOV, 1923 (type, *Cardium fittoni* D'ORBIGNY, 1845; OD)]. Ribs triangular in section, conspicuous, irregular. *Plio.(Pont.-Kimmer.)*, S.USSR.

Didacnomya ANDRUSSOV, 1923 [**Cardium vulgaris* SINZOV, 1876 (non DA COSTA, 1778) (?= *Monodacna vulgaris planior* KRESTOVNIKOV, 1928); OD]. *Plio.(Pont.-Kuialnik.)*, S.USSR.

Oraphocardium EBERSIN, 1949 [**Phyllocardium oraphense* DAVIDASCHVILI, 1930; OD]. Like *Phyllocardium* but posterior margin rectangular instead of oblique. *Plio.(Pont.-Kimmer.)*, S.USSR.

Oxydacna DAVIDASCHVILI, 1930 [**O. tenericardo*, ex ANDRUSSOV MS; M]. Ribs narrow at crests. *M.Plio.(Kimmer.)*, S.USSR.

Phyllocardium FISCHER, 1887 [**Cardium planum* DESHAYES, 1838; M] [= *Phyllicardium* AUCTION. (nom. null.)]. Ovate, tending toward alation of dorsal margin. *Plio.(Pont.-Kimmer.)*, S.USSR.—FIG. E88,1. *P. *planum* (DESHAYES), *L.Plio.*, S. USSR; *1a-c*, LV ext., RV int., LV int., $\times 1$ (Andrussov).

Plagiodacna ANDRUSSOV, 1903 [**Cardium carinatum* DESHAYES, 1838 (non BRONN, 1831) (= *C. angulosum* DESHAYES, 1857); SD CROSSE, 1905]. Rhomboidal, with umbonal ridge. *Plio.(Pont.-Kuialnik.)*, S.USSR.

Pseudocatillus ANDRUSSOV, 1903 [**Cardium pseudocatillus* BARBOT-DE-MARNY, 1869; T]. Shell somewhat compressed, ribs weak. *Plio.(Pont.-Kuialnik.)*, S.USSR.

Pteradacna ANDRUSSOV, 1907 [**Cardium edentulum* DESHAYES, 1838 (non MONTAGU, 1808) (= *C. subedentulum* D'ORBIGNY, 1852; M)]. Ribs numerous, with threadlike nodes; dorsal margin alate; hinge edentulous. *M.Plio.(Kimmer.)*, S.USSR.

Stenodacna ANDRUSSOV, 1923 [**Cardium angusti-*

costatum ROUSSEAU, 1842; M]. *M.Plio.*(Kimmer.), S.USSR.

Subfamily PARADACNINAE Ebersin, 1964

Shells very thin and fragile, mostly edentulous; pallial line entire or nearly so. *Plio.* (Pont.-Kimmer.).

Paradacna ANDRUSSOV, 1909 [pro *Abichia* ANDRUSSOV, 1907 (non GEMMELLARO, 1888)] [**Cardium abichi* HÖRNES, 1874; M]. Elongate, trapezoidal, posterior slope smooth, posterior ribs carinate, wide-spaced. *Plio.*(Pont.-Kimmer.), S.USSR.

Arcicardium FISCHER, 1887 [**Cardium acardo* DESHAYES, 1838; M]. Shell flattened, trigonal, very inequilateral, edentulous. *L.Plio.*, USSR (Crimea).

Chartoconcha ANDRUSSOV, 1907 [**Cardium bayerni* HÖRNES, 1874; M]. Smooth, hinge edentulous. *Plio.*(Pont.-Kiuialnik.), S.USSR.

Panticapaea ANDRUSSOV, 1923 [**Cardium duboisi* MAYER, 1856; M]. *U.Plio.*(Kimmer.), S.USSR.

Papyrocardium GABUNIA, 1953 [**P. fragilicostatum* (fide EBERSIN, 1965)]. *U.Plio.*(Kimmer.), S.USSR.

Parvidacna STEVANOVIĆ, 1951 [**P. planicostata* (fide EBERSIN, 1965)]. *L.Plio.*(Pont.), S.USSR.

Subfamily ADACNINAE von Vest, 1875

[nom. transl. EBERSIN, 1965 (ex Adacnidae von VEST, 1875)]

Hinge weak to edentulous; pallial sinus present in most; some forms gaping at one or both ends. *U. Plio.* (*Aktschagyl.-Apscheron.*)-*Rec.*

Adacna EICHWALD, 1838 [**Glycimeris laeviuscula* EICHWALD, 1829 (?=*A. laeviuscula* EICHWALD, 1838); SD von VEST, 1875]. Edentulous; valves thin, gaping at both ends; pallial sinus large. *U. Plio.*(*Apscheron.*)-*Rec.*, Caspian Sea-S.USSR.—FIG. E88,2. **A. laeviuscula* (EICHWALD), *Rec.*, Caspian Sea; 2a,b, RV ext., LV int., ×1 (Chenu, 1857).

Apscheronia ANDRUSSOV, 1903 [**Cardium propinquum* EICHWALD, 1841 (non MÜNSTER, 1837) (= *C. proximum* DESHAYES, 1857); M]. Shell surface smooth, without ribs. *U.Plio.*(*Apscheron.*), S.USSR.

Avicardium KOLESNIKOV, 1950 [**Cardium nikitini* ANDRUSSOV, 1902; OD]. *U.Plio.*, Caspian Basin.

Caspicardium ASTAFIEVA, 1955 [**Cardium trapezinum* ANDRUSSOV, 1923 (fide EBERSIN, 1965)]. *U.Plio.*(*Apscheron.*), S.USSR.

Catilloides ANDRUSSOV, 1923 [**Monodacna catilloides*; OD]. *U.Plio.*(*Apscheron.*), S.USSR.

Hypanis EICHWALD, 1838 (ex PANDER MS) [**Cardium plicatum*; M]. Resembling *Adacna* but with wide-spaced sharp ribs. *U.Plio.*(*Apscheron.*), S.USSR.

Hyrcania KOLESNIKOV, 1950 [**Didacna hyrcana*




FIG. E89. Lahillidae (Pseudocarditinae) (p. N593).

ANDRUSSOV, 1923; OD] [= *Turkmena* POPOV, 1956 (type, *Didacna turkmena* ANDRUSSOV, 1923; OD); *Irinia* POPOV, 1956 (type, *Didacna turkmena major* ANDRUSSOV, 1923; OD)]. *U.Plio.*(*Apscheron.*), S.USSR.

H. (Hyrcania). *U.Plio.*(*Apscheron.*), S.USSR.

H. (Didacnoides) ASTAFIEVA, 1955 [**Monodacna didacnoides* ANDRUSSOV, 1923; T (fide EBERSIN, 1965)]. *U.Plio.*(*Apscheron.*), S.USSR.

H. (Hyrcanomya) ASTAFIEVA, 1955 [**Monodacna bakuana* ANDRUSSOV, 1923 (fide EBERSIN, 1965)]. *U.Plio.*(*Apscheron.*), S.USSR.

Monodacna EICHWALD, 1938 [**Corbula caspia* EICHWALD, 1829; SD von VEST, 1875]. Rounded-quadrate, ribs broad; hinge with cardinals 2 and 3b, no laterals; pallial sinus small. *U.Plio.*(*Apscheron.*)-*Rec.*, Caspian-S.USSR.—FIG. E88, 8. **M. caspia* (EICHWALD), *Rec.*, Caspian Sea; RV ext., ×1 (124b).

Parapscheronia EBERSIN, 1955 [**Apscheronia volarovi* ANDRUSSOV, 1923; M]. *U.Plio.*(*Apscheron.*), S.USSR.

Plagiadacnopsis ANDRUSSOV, 1923 [**Monodacna (P.) isseli*; OD]. *U.Plio.*(*Apscheron.*), S.USSR.

Subfamily PSEUDOCARDITINAE Ebersin, 1964

Outline oblique; hinge lacking laterals and right anterior cardinal; pallial line entire. *Plio.*

Pseudocardita OPPENHEIM, 1918 [**Cardium (P.) bukowskii*; SD KEEN, 1937]. Oblique, inequilateral, ribs broad, rounded; hinge plate wide, teeth stout, nymphs elongate. *Neog.*(?Pont.), Asia Minor.—FIG. E89,1. **P. bukowskii* (OPPENHEIM), *Plio.*, S.Turkey; 1a-c, RV ext., LV int., RV int., ×1 (Oppenheim, 1918).

Subfamily UNCERTAIN

Diversicostata EBERSIN & WASSOIEVIĆ, 1930 [**Monodacna maxima*, ex ANDRUSSOV MS; SD SALISBURY,

1932 (proposed as an undefined "section" and later rejected as invalid by EBERSIN)].

Myocardia VON VEST, 1861 [**M. truncata*; SD VON VEST, 1875]. Hinge resembling that of *Cardium*

but shells with the pallial sinus of *Adacna*. *Mio.*, Eu.(Hung.).

Prophylicardium JEKELIUS, 1944 [**P. soceni* (fide Zool. Record, 1946)]. *Mio.*, Eu.(Rumania).

Replidacna JEKELIUS, 1944 [**R. carasi* (fide Zool. Record, 1946)]. *Mio.*, Eu.(Rumania).

Uniocardium CAPELLINI, 1880 [**U. meneghini*; M]. *Plio.*, Eu.(Italy).

Superfamily TRIDACNACEA Lamarck, 1819

[*nom. transl.* DALL, 1895 (ex *Tridacnidae* GOLDFUSS, 1820) (=tridacnées LAMARCK, 1819)] [Materials for this superfamily prepared by MYRA KEEN]

Medium-sized to very large shells, with radial ribbing; byssal gape in most; hinge with two oblique lamellar cardinal teeth and one or more lateral teeth; soft parts "rotated" within shell as special adaptation to mode of life among corals; posterior adductor and large posterior pedal retractor muscle scar centrally located, anterior adductor scar wanting; pallial line entire. ?*U. Cret., Tert.-Rec.*

Family TRIDACNIDAE Lamarck, 1819

[*nom. correct.* MENKE, 1830 (pro *Tridacnadae nom. transl. et correct.* FLEMING, 1828, ex Tribe *Tridacnae* GOLDFUSS, 1820) (=tridacnées LAMARCK, 1819)]

Radial ribs smooth or with spinose ornamentation. ?*U. Cret., Tert.-Rec.*

Tridacna BRUGUIÈRE, 1797 [**Chama gigas* LINNÉ, 1758; SM LAMARCK, 1799] [= *Tridachnes* RÖDING, 1798 (obj.); *Tridacnodites* KRÜGER, 1823 (obj.); *Dinodacna* IREDALE, 1937 (obj.)]. Large to massive, ribs few, strong; anterior lateral teeth wanting. ?*U. Cret., Tert.-Rec.*, Eu.-E.Indies-Afr.-S.Pac.

T. (Tridacna). Ribs deeply folded; umbones subcentral. Shells unattached but among reef corals. *L.Mio.-Rec.*, Eu.-E.Indies.—FIG. E90,6. **T. (T.) gigas* (LINNÉ), *Rec.*, E.Indies; LV ext., ×0.05 (562; 797).

T. (Chametracea) MÖRCH, 1853 [**Tridacna crocea* LAMARCK, 1819; SD IREDALE, 1937] [= *Floodacna* IREDALE, 1937 (type, *T. squamosa* LAMARCK, 1819; OD); *Sepidacna* IREDALE, 1937 (type, *T. trouhthoni* = *Tridachnes maxima* RÖDING, 1798; OD); *Vulgodacna* IREDALE, 1937 (type, *T. maxima* var. *fossor* HEDLEY, 1921 = *Tridachnes maxima* RÖDING, 1798; OD)]. Valves equilateral to strongly inequilateral, umbones anterior to mid-line; sculpture tending to be scaly. Boring into coral. ?*U. Cret., Tert.-Rec.*, Afr.-Eu.-E.Indies.

T. (Persikima) IREDALE, 1937 [**P. whiteleyi* (= *Tridachnes derasa* RÖDING, 1798); OD]. Um-




FIG. E90. Tridacnidae (p. N594-N595).

bones posterior to mid-line, shell strongly inequilateral. *Rec.*, S.Pac.

Avicularium GRAY, 1853 [*"Cardium aviculare* LAM." (= *Cardita avicularia* LAMARCK, 1805); SD DALL, 1900] [= *Lithocardium* WOODWARD, 1854 (obj.)]. Triangular, higher than long, ribs evident but nearly smooth. *M.Eoc.-L.Oligo.*, Eu.-W.Indies.—FIG. E90,1. **A. avicularium* (LAMARCK), Eoc., France; 1*a,b*, LV int., ext., $\times 0.5$ (Cossmann & Pissarro, 1904).

Byssocardium MUNIER-CHALMAS, 1882 [**Cardium emarginatum* DESHAYES, 1829; OD]. With an anterior gape bordered by plaits, as in *Tridacna*, but shell more triangular. *M.Eoc.-L.Mio.*, Eu.—FIG. E90,4. **B. emarginatum* (DESHAYES), Eoc., France; 4*a,b*, RV ext., int., $\times 0.7$ (Deshayes).

Gonicardium VASSEUR, 1880 [**Cardium rachitis* DESHAYES, 1829; SD KEEN, 1937]. Oblique-ovate, spinose, anterior lateral teeth wanting. *M.Eoc.-U.Eoc.*, Eu.—FIG. E90,2. **G. rachitis* (DESHAYES), Eoc., France; 2*a*, RV int., $\times 0.7$; 2*b*, LV ext., $\times 1$ (Cossmann & Pissarro, 1904).

Hippopus LAMARCK, 1799 [**Chama hippopus* LINNÉ, 1758; M] [= *Pelvis* MEGERLE, 1811 (obj.); *Cerceis* GISTL, 1848 (obj.) (*nom. var.*)]. Like *Tridacna* in general form but byssal gape closed in adult. *Mio.-Rec.*, E.Indies-W.Indies.—FIG. E90,5. **H. hippopus* (LINNÉ), *Rec.*, E.Indies; RV ext., $\times 0.3$ (Stasek, 1962).

Sawkinsia COX, 1941 [**S. matleyi*; OD]. Resembling *Tridacna* but not gaping; anterior lateral teeth present. *U.Eoc.-U.Oligo.*, Carib.—FIG. E90,3. **S. matleyi*, Eoc., Jamaica; RV int., $\times 1$ (Cox, 1941).

Superfamily MACTRACEA Lamarck, 1809

[*nom. transl.* DALL, 1895 (ex family Mactracea GRAY, 1823) (=mactracées LAMARCK, 1809)] [Materials for this superfamily prepared by MYRA KEEN]

Thin-shelled, porcelaneous; hinge with inverted V-shaped cardinal tooth in LV, two cardinals in RV; lateral teeth and accessory cardinal laminae present in most groups; external ligament small or wanting, internal ligament or resilium seated in socket-like resilifer; pallial sinus normally well developed (507, 510). *U.Cret.-Rec.*

Family MACTRIDAE Lamarck, 1809

[*nom. correct.* SWAINSON, 1835 (pro family Mactracea GRAY, 1823) (=mactracées LAMARCK, 1809)]

Periostracum present, glossy; shell smooth or concentrically sculptured; valves slightly gaping; pallial line with sinus; siphons of animal united to their tips (510). *U.Cret.-Rec.*

Subfamily MACTRINAE Lamarck, 1809

[*nom. transl.* ADAMS & ADAMS, 1856 (ex family Mactracea GRAY, 1823) (=mactracées LAMARCK, 1809)]

Subequilateral, nearly closed; hinge well developed, two cardinals of RV somewhat joined; siphons wholly retractile within shell. *U.Cret.-Rec.*

Mactra LINNÉ, 1767 [**Cardium stultorum* LINNÉ, 1758; SD FLEMING, 1818 (+ICZN, Dir. 72, as of GRAY, 1847)] [= *Trigonella* DA COSTA, 1778 (obj.; SD WINCKWORTH, 1926); *Deikea* MAYER, 1872 (type, *M. gallensis* MAYER, 1867; SD KEEN, herein); *Colorimactra* IREDALE, 1929 (type, *M. queenslandica* E. A. SMITH, 1914; OD); *Tele-mactra* IREDALE, 1929 (type, *M. obesa* REEVE, 1854, ex DESHAYES MS; OD)]. Trigonal to oval, somewhat inflated, subequilateral; lunule and escutcheon delimited; ligament separated from resilium by shelly lamina; lateral teeth smooth; pallial sinus oval. *Eoc.-Rec.*, cosmop.

M. (Mactra). Lunule and escutcheon not set off by groove; hinge of LV with *AII* and *4b*; RV with *3a-3b* not fused above, laterals doubled; pallial sinus rounded, not deep. *Rec.*, circumtrop.—FIG. E91,2. **M. (M.) stultorum* (LINNÉ), *Medit.*; 2*a*, LV ext., $\times 0.5$ (Reeve, 1854); 2*b,c*, LV and RV hinges, $\times 1$ (510).

M. (Allomactra) TOMLIN, 1931 [*pro Heteromactra* COSSMANN & PEYROT, 1909 (*non* LAMY, 1906)] [**M. (H.) grateloupi* COSSMANN & PEYROT, 1909, ex DESHAYES MS; OD]. Trigonal, flattened, smooth, resembling *M. (Eomactra)* but with wider cardinal plate, resilifer pyriform, *4b* wanting, also nymph; pallial sinus long and broad. *Mio. (Burdigal.)*, Eu.—FIG. E91,3. **M. (A.) grateloupi* COSSMANN & PEYROT, France; 3*a,b*, RV ext., int., $\times 1$; 3*c*, LV hinge, $\times 2$ (164).

M. (Andrussella) KOROBKOV, 1954 [**M. acute-carinata* ANDRUSSOV, 1902; OD]. Posterior slope meeting central slope in sharp keel that may be prolonged into spout. *Plio.*, E.Eu.—FIG. E91,1. **M. (A.) acute-carinata* ANDRUSSOV, S.Caucasus; 1*a,b*, RV, LV ext., $\times 1.5$; 1*c*, RV hinge, $\times 5$ (Andrussov, 1902).

M. (Austromactra) IREDALE, 1930 [**A. caloundra*; OD]. Trigonal, equivalve, sculpture of laminar concentric ridges wavy at ends; hinge broad; pallial sinus small, rounded. *Rec.*, S.Pac.—FIG. E91,6. **M. (A.) caloundra* (IREDALE), Australia; 6*a,b*, LV ext., int., $\times 1$ (Iredale, 1930).

M. (Avimactra) ANDRUSSOV, 1905 [**M. (A.) aviculoides*; M]. Trigonal, dorsal margin long and straight, ventral attenuated by radial fold; hinge with large resilifer. *Plio.*, E.Eu.—FIG. E91,5. **M. (A.) aviculoides*, S.Caucasus; 5*a*, RV ext., $\times 1$; 5*b*, RV hinge, $\times 2$ (Andrussov, 1905).

M. (Barymactra) COSSMANN & PEYROT, 1909 [**Mactra burdigalensis* MAYER, 1864; OD]. Large, thick, short, hinge teeth close-set; pallial sinus

short, slightly descending, line doubled, bordered above by row of punctate scars. *Eoc.-Mio.*, Eu.-SW.Afr.—FIG. E91,4. **M. (B.) burdigalensis* MAYER, Mio., France; 4*a,b*, RV int., LV int., $\times 0.5$ (164).

M. (Coelomactra) DALL, 1895 [**M. violacea* GMELIN, 1791; OD]. Valves thin, inflated, dorsal

areas grooved, resilifer not roofed at apex; hinge plate oblique, teeth not concentrated, laterals long, thin, cardinals buttressed; pallial sinus short, high. *Rec.*, Pac.—FIG. E92,6. **M. (C.) violacea* GMELIN, E.Indies; 6*a*, LV ext., $\times 0.5$ (Reeve, 1854), 6*b,c*, LV and RV hinges, $\times 2$ (510).

M. (Cryptomactra) ANDRUSSOV, 1902 [**Lucina*




FIG. E91. Mactridae (Mactrinae) (p. N595-N596).

pesanseris MAYER-EYMAR, 1857; M]. Small, trapezoidal, with 1 or 2 strong radial folds, making ventral margin sinuous. *Mio.* (*Sarmat.*), E.Eu.—
FIG. E92, 1. **M. (C.) pesanseris* (MAYER-EYMAR), U.Mio., S.Caucasus; 1a, b, LV ext., RV ext., $\times 1$;

1c, d, RV and LV hinges, $\times 1.5$ (Andrussov, 1902).

M. (Cyclomactra) DALL, 1895 [**M. tristis* "GRAY" (*err. pro* **M. tristis* REEVE, 1854); OD]. Like *M. (Mactroderma)* but subcircular, compressed, liga-




FIG. E92. Mactridae (Mactrinae) (p. N596-N598).

- ment almost submerged though separate from resilium. *Plio.-Rec.*, S.Pac.—FIG. E92,5. **M. (C.) tristis* REEVE, *Rec.*, N.Z.; LV ext., $\times 0.7$ (Reeve, 1854).
- M. (Electomacra)** IREDALE, 1930 [**M. parkesiana* HEDLEY, 1902; OD] [=Electomacra, spelling error]. Small, short, hinge wide and heavy for size of shell. *Rec.*, S.Pac.—FIG. E92,7. **M. (E.) parkesiana* HEDLEY, Australia; 7a,b, RV ext., int., $\times 1.5$; 7c, LV hinge, $\times 5$ (Hedley, 1902).
- M. (Eomacra)** COSSMANN in COSSMANN & PEYROT, 1909 [**M. basteroti* MAYER, 1853; OD]. Small, triangular, smooth except for concentric ribs on lunule and escutcheon; resilifer small, not projecting beyond margin of hinge plate. *Eoc.-Plio.*, Eu.—FIG. E92,2. **M. (E.) basteroti* MAYER, Mio., France; 2a,b, RV int., ext., $\times 1$ (164).
- M. (Longimacra)** FINLAY, 1928 [**M. elongata* QUOY & GAIMARD, 1835; OD]. Long, concentric ribs stronger at ends; sinus deep, muscle scars large. *Rec.*, S.Pac.—FIG. E92,9. **M. (L.) elongata* QUOY & GAIMARD, N.Z.; LV ext., $\times 0.5$ (Reeve, 1854).
- M. (Macrinula)** GRAY, 1853 [**M. plicaria* (=Macra plicataria LINNÉ, 1767); M] [=Macretrella GRAY, 1853 (type, *M. striatula* LINNÉ, 1767; M); *Papyrina* MÖRCH, 1853 (obj.; SD KEEN, herein)]. Valves concentrically plicate, inequilateral, posterior slope set off by ridge; resilifer large and narrow, hinge teeth not concentrated, cardinal in LV reinforced by apophysis, with accessory lamella in front. *L.Mio.-Rec.*, Pac.—FIG. E92,4. **M. (M.) plicataria* LINNÉ, *Rec.*, E.Indies; 4a, LV ext., $\times 0.5$ (Reeve, 1854); 4b,c, LV and RV hinges, $\times 2$ (510).
- M. (Mactroderma)** DALL, 1894 [**M. velata* PHILIPPI, 1848; OD]. Anterior end shorter than posterior; pallial sinus large and round. *Rec.*, N. Am.-S. Am.—FIG. E92,8. **M. (M.) velata* PHILIPPI, W.C. Am.; 8a, LV ext., $\times 0.3$ (Reeve, 1854); 8b,c, LV and RV hinges, $\times 1$ (510).
- M. (Mactrona)** MARWICK, 1952 [*pro Mactrula* MARWICK, 1948 (non RISSO, 1826)] [**M. (Mactrula) mula* MARWICK, 1948; OD]. Exteriorly like *M. (Macra)* but with sunken ligament and large cardinal of *M. (Cyclomacra)*; pallial sinus small, rounded. *Plio.*, S.Pac.—FIG. E92,3. **M. (M.) mula* MARWICK, N.Z.; 3a,b, LV ext., int., $\times 0.4$ (Marwick, 1948).
- M. (Mactrotoma)** DALL, 1894 [**M. fragilis* GME-LIN, 1791; OD]. With thin, silky periostracum, posterior area set off by angular ridge over which periostracum is wrinkled; ligament long, resilifer large, shallow. *Rec.*, circumtrop.—FIG. E93,11. **M. (M.) fragilis* GME-LIN, Carib.; 11a, LV ext., $\times 0.5$ (Reeve, 1854); 11b,c, LV and RV hinges, $\times 2$ (510).
- M. (Maorimacra)** FINLAY, 1928 [**M. ordinaria* E. A. SMITH, 1898; OD]. Small, outline *Corbula*-like, posterior end longer. *Rec.*, S.Pac.—FIG. E93,4. **M. (M.) ordinaria* SMITH, N.Z.; LV ext., $\times 2$ (Smith, 1898).
- M. (Micromacra)** DALL, 1894 [**M. californica* CONRAD, 1837; M]. Shell small, solid, beaks and umbones with undulating concentric ribs; hinge as in *M. (Mactrotoma)*. *Rec.*, W.N. Am.-C. Am.—FIG. E93,3. **M. (M.) californica* CONRAD, Panama; LV ext., $\times 1$ (Reeve, 1854).
- M. (Nannomacra)** IREDALE, 1930 [**M. jacksonensis* E. A. SMITH, 1885; OD]. Thin, glassy, concentrically striate, trigonal; hinge narrow, delicate; pallial sinus shallow. *Rec.*, S.Pac.—FIG. E93,6. **M. (N.) jacksonensis* SMITH, Australia, 6a,b, RV int., dorsal, $\times 1.5$ (852).
- M. (Sarmatimacra)** KOROBKOV, 1954 [**M. vitaliana* D'ORBIGNY, 1845; OD]. Subtrigonal, umbones wide; hinge fairly massive; lamina between ligament and resilifer toothlike; pallial sinus short, rounded, pallial line deeply incised, vertically striate. *Mio.*, E. Eu.
- M. (Simomacra)** DALL, 1894 [**M. dolabriformis* CONRAD, 1837; M]. Outline flattened cuneiform, inequilateral; pallial sinus smaller than in *M. (Mactrotoma)*, siphonal gape small. *Rec.*, W.N. Am.-C. Am.—FIG. E93,7. **M. (S.) dolabriformis* CONRAD, USA (Calif.); 7a, LV ext., $\times 0.3$ (Dall, 1894); 7b,c, RV and LV hinges, $\times 1$ (510).
- M. (Stiphromacra)** BÖHM, 1929 [**M. (S.) welwitschi*; OD]. Hinge wide but teeth very small. *Eoc.*, W. Afr.—FIG. E93,10. **M. (S.) welwitschi*, Angola; 10a-c, RV ext., LV and RV hinges, $\times 0.5$ (Böhm, 1929).
- M. (Tumbeziconcha)** PILSBRY & OLSSON, 1935 [**M. thracioides* ADAMS & REEVE, 1848; OD]. Resembling *M. (Macrinula)* as to sculpture but lacking posterior ridge; concentric ribs intersected by oblique striae on posterior slope. *Rec.*, W.C. Am.—FIG. E93,2. **M. (T.) thracioides* ADAMS & REEVE, Peru; LV ext., $\times 1$ (Pilsbry & Olsson, 1935).
- Aliomacra** STEPHENSON, 1952 [1953] [**A. compressa*; OD]. More compressed than *Macra*; ligament short, external, with internal part on surface sloping down and forward from nymph; pallial sinus rounded, moderate, not confluent. *U. Cret.*, N. Am.—FIG. E93,9. **A. compressa*, USA (Tex.); 9a, RV ext., $\times 1$; 9b,c, RV and LV hinges, $\times 1.5$ (Stephenson, 1953).
- Cymbophora** GABB, 1869 [**Macra ashburnerii* GABB, 1864; OD] [=Veleda CONRAD, 1870 (non BLACKWALL, 1859) (type, *V. lintea* = *Cardium lintea* CONRAD, 1860, non CONRAD, 1855 = *Cymbophora intoxicata* HANNA, 1924)]. Trigonal, resilifer narrow and shallow, bordered by accessory lamella; lateral teeth strong, close to beaks. *U. Cret.*, E. N. Am.-W. N. Am.-Eu.—FIG. E93,8. **C. ash-*

burnerii (GABB), USA (Calif.); LV hinge, $\times 1$ (748).

Diaphoromactra IREDALE, 1930 [*Hemimactra versicolor* TATE, 1887; OD]. Small, solid, oblique, an-

terior end short, posterior with umbonal ridge; lateral teeth partially or entirely wanting in LV; pallial sinus broad and shallow. *Rec.*, S.Pac.—FIG. E93, I. **D. versicolor* (TATE), *Rec.*, Australia;




FIG. E93. Mactridae (Mactrinae) (p. N598-N601).

1*a,b*, RV ext., both valves dorsal, $\times 1.5$ (Tate, 1887).
 Geltena STEPHENSON in VOKES, 1946 [*G. subequilatera*; OD]. Subcircular to broadly subovate; resilifer very small; nymph plate heavy; hinge plate

with deep grooves above laterals. *U.Cret.*, Eu.-N.Am.-W.Asia.—FIG. E93,5. **G. subequilatera*, USA(Tex.); 5*a*, LV ext., $\times 1$; 5*b,c*, $\times 1$, LV and RV hinges, $\times 1.3$ (Stephenson in Vokes, 1946).
 Harvella GRAY, 1853 [*Maetra elegans* SOWERBY,




FIG. E94. Mactridae (Mactrinae) (p. N601).

- 1825; M]. Resembling *Mactrellona* but with spaced concentric undulations in front of keel; anterior lateral teeth short. *Rec.*, W.C.Am.—FIG. E93,12. **H. elegans* (SOWERBY); 12a-c, RV ext., int., LV hinge, $\times 1$ (H. Adams & A. Adams, 1856).
- Mactrellona** MARKS, 1951 [**Mactra alata* SPENGLER, 1802; OD] [= *Mactrella* AUCTION (non GRAY, 1853)]. Trigonal, inequilateral, thin, inflated, posterior slope set off by keel, umbones prominent; hinge concentrated, anterior laterals short. *Mio.-Rec.*, E.C.Am.-W.C.Am.—FIG. E94,8. **M. alata* (SPENGLER), *Rec.*, Carib.; 8a, RV ext., $\times 0.5$; 8b,c, RV and LV hinges, $\times 1$ (H. Adams & A. Adams, 1856).
- Mulinia** GRAY, 1837 [**M. typica* (= *Mactra edulis* KING & BRODERIP, 1832); T]. Outline as in *Mactra* but ligament entirely internal, not separated from resilium by shelly ridge; shell not gaping; pallial sinus short, small. Estuarine. *Rec.*, W.C.Am.-E.C.Am.-S.Am.-E.Afr.—FIG. E94,9. **M. edulis* (KING & BRODERIP), S.Am.; 9a, LV ext., $\times 1$ (Carcelles, 1950); 9b,c, RV and LV hinges, $\times 2$ (510).
- Mulinoides** OLSSON, 1944 [**M. chilca*; OD]. Resembling *Cymbophora* but shell more solid, rounded, lateral teeth stronger, less lamellar, resilifer more mactroid in form; posterior dorsal area well marked, set off by ridge; lunule present. *U.Cret.*, W.S.Am.—FIG. E94,5. **M. chilca*; 5a, RV ext., Peru; $\times 0.5$; 5b,c, LV and RV hinges, $\times 1$ (Olsson, 1944).
- Priscomactra** STEPHENSON, 1952 [1953] [**P. cymba*; OD]. Ligament and hinge as in *Geltena* but outline subtrigonal and mactroid. *U. Cret.*, N.Am.—FIG. E94,3. **P. cymba*, USA (Tex.); 3a, LV ext., $\times 1$; 3b,c, LV and RV hinges, $\times 1.3$ (Stephenson, 1953).
- Pseudocardium** GABB, 1866 [**Cardium gabbii* RÉMOND, 1863 (= *Mulinia densata* CONRAD, 1857); M]. Shell solid, rounded-trigonal; hinge with long substrate or pitted laterals; pallial sinus short, rounded. *Oligo.-Rec.*, W.N.Am.-E.Asia.—FIG. E94,10. **P. densatum* (CONRAD), Plio., USA (Calif.); 10a-c, LV ext., int., RV int., $\times 0.5$ (Packard, 1916).
- Rangia** DES MOULINS, 1832 [**R. cyrenoides* (= *Gnathodon cuneatus* SOWERBY, 1831); M] [= *Gnathodon* SOWERBY, 1831 (non GOLDFUSS, 1820); *Colombia* RANG, 1834 (nom.nud.); *Perissodon* CONRAD, 1863 (type, *Mactra clathrodonta* CONRAD, 1833; SD DALL, 1894)]. Shell somewhat resembling *Mulinia* but with lateral teeth elongate, curved, cross-striate, posterior pair longer; pallial sinus small. [Estuarine.] ?*Paleoc.-Mio.-Rec.*, N.Am.-S.Am.-Eu.
- R.** (**Rangia**). Anterior lateral teeth hooked. *Mio.-Rec.*, E.N.Am.-W.N.Am.-S.Am.—FIG. E94,2. **R. (R.) cuneata* (SOWERBY), *Rec.*, USA (La.); 2a,b, LV ext., int., $\times 0.5$ (Dall, 1894).
- R.** (**Miorangia**) DALL, 1894 [**Gnathodon johnsoni* DALL, 1892; OD]. Small, extremely inequilateral, with cardinals reversed, superior pair in LV; pallial sinus obsolete. *Mio.*, E.N.Am.—FIG. E94,7. **R. (M.) johnsoni* (DALL), USA (Miss.); LV ext., $\times 1.5$ (Dall, 1894).
- R.** (**Rangianella**) CONRAD, 1868 [**Gnathodon trigonum* PETIT, 1853 (= *Mactra mendica* GOULD, 1851); M]. Small, nearly equilateral, lateral teeth short, straight, subequal, faintly rugose; pallial sinus inconspicuous or obsolete. Estuarine to marine. *Rec.*, W.C.Am.-N.Am.—FIG. E94,1. **R. (R.) mendica* (GOULD), W.C.Am.; LV int., $\times 1$ (Dall, 1894).
- R.** (?**Praerangia**) COSSMANN, 1908 [**P. minuscula*; OD]. Resembling *Rangia* in form but much smaller, with shallower sinus. *Paleoc.(Mont.)*, Eu.—FIG. E94,4. **R. (?P.) minuscula* (COSSMANN), Belg.; 4a-c, RV ext., LV int., RV int., $\times 7$ (Cossmann, 1908).
- Scalpomactra** FINLAY in MARWICK, 1928 [**Mactra scalpellum* REEVE, 1854; OD]. Resembling *Spisula* but lateral teeth not striate; resilium stout, oblique, triangular below, produced above as long spike curved to front, with minute ligament attached at top and united nearly its full length. *Plio.-Rec.*, S.Pac.—FIG. E94,6. **S. scalpellum* (REEVE), *Rec.*, N.Z.; LV ext., $\times 1$ (Powell, 1957).
- Scissodesma** GRAY, 1837 [**Mactra spengleri* LINNÉ, 1767; SD GRAY, 1847] [= *Schizodesma* GRAY, 1838 (nom.van.)]. Ligament only partially external, in deep slit, no lamina separating it from resilium. *Rec.*, Pac.—FIG. E95,8. **S. spengleri* (LINNÉ), E.Indies; 8a-c, RV ext., int., LV hinge, $\times 1$ (H. Adams & A. Adams, 1856); 8d-e, LV, RV hinges (510).
- Spisula** GRAY, 1837 [*“*Mactra solida* MONTAGU” (= *Cardium solidum* LINNÉ, 1758); SD GRAY, 1847] [= *Spisulina* FISCHER, 1887 (type, *Mactra truncata* MONTAGU, 1808; M)]. Trigonal to ovate, not gaping, concentrically striate; lunule and escutcheon delimited; ligament and resilium not separated by shelly lamella; pallial sinus oval. *Tert.-Rec.*, cosmop.
- S.** (**Spisula**). Lunule and escutcheon concentrically grooved; lateral teeth striate. *Rec.*, Atl.—FIG. E95,4. **S. (S.) solida* (LINNÉ), Eng.; 4a-c, RV int., ext., LV hinge, $\times 1$ (H. Adams & A. Adams, 1856).
- S.** (**Crassula**) MARWICK, 1948 [**Mactra aequilateralis* DESHAYES, July 1854 (= *M. aequilatera* REEVE, May 1854); OD]. Posterior area bounded by sharp ridge, sculpture only on anterior end, not wavy. *Rec.*, S.Pac.—FIG. E95,5. **S. (C.) aequilatera* (REEVE), N.Z.; LV ext., $\times 0.5$ (Reeve, 1854).
- S.** (**Crepispisula**) EAMES, 1957 [**S. (C.) amekien-sis*; OD]. Near *Scissodesma* in outline but with concentric sculpture, posterior carina less marked, ligament slit smaller, lateral teeth not crenulate; lunule and escutcheon sulcate. *Eoc.*, W.Afr.—

FIG. E95,7. **S. (C.) amekiensis*, Nigeria; 7a,b, RV int., LV int., $\times 2$ (Eames, 1957).

S. (Hemimactra) SWAINSON, 1840 [**Mactra gigantea* LAMARCK, 1818 (= *M. solidissima* DILLWYN, 1817); M]. Large, ovate-trigonal, with

dorsal areas not sulcate; anterior arm of cardinal in RV confluent with ventral lamina, cardinals markedly compressed; laterals striate. *Tert.-Rec.*, E.N.Am.-Atl.—FIG. E95,2. **S. (H.) solidissima* (DILLWYN), Rec., USA (Mass.); 2a, LV ext.,




FIG. E95. Mactridae (Mactrinae) (p. N601-N603).




FIG. E96. Mactridae (Mactrinae) (p. N603-N604).

×0.3 (Gould-Binney, 1870); 2*b,c*, RV and LV hinges (510).

S. (Mactromeris) CONRAD, 1868 [**Mactra ovalis* GOULD, 1840 (*non* SOWERBY, 1817) (= **M. polynyma* STIMPSON, 1860); SD STOLICZKA, 1871] [= *Mactrodesma* CONRAD, 1869 (type, *M. ponderosa* CONRAD, 1830 [*non* EICHWALD, 1830] = *M. subponderosa* D'ORBIGNY, 1852; M)]. With fibrous periostracum; lateral teeth smooth, cardinals not compressed, anterior arm of 3*a* not confluent with ventral lamina. *Mio.-Rec.*, W.N.Am.-Japan-N.Atl.—FIG. E95,3. **S. (M.) polynyma* (STIMPSON), *Rec.*, N.Atl.; 3*a*, LV ext., ×0.5 (217); 3*b,c*, LV and RV hinges, ×1 (510).

S. (Notospisula) IREDALE, 1930 [**Gnathodon par-*

vum PETIT, 1853; OD]. Small, thick, trigonal, tumid, rounded anteriorly, pointed posteriorly; lateral teeth ridged; pallial sinus small to obsolete. *Rec.*, S.Pac.—FIG. E95,1. **S. (N.) parva* (PETIT), Australia; 1*a*, LV ext., ×1 (Petit, 1853); 1*b,c*, RV and LV hinges, ×3 (397).

S. (Oxyperas) MÖRCH, 1853 [**Mactra triangularis* LAMARCK, 1818" (*non* MONTAGU, 1803) (= **M. lentiginosa* GOULD, 1852); M]. Triangular, with well-marked concentric folds; pallial sinus deep. *Rec.*, Pac.—FIG. E95,6. *S. (O.) lentiginosa* (GOULD), ?E.Indies; 6*a*, RV ext., ×0.7 (510); 6*b,c*, LV and RV hinges, ×2 (510).

S. (Pseudoxyperas) SACCO, 1901 [**P. proaspera*; OD]. Differing from *S. (Oxyperas)* by being

less trigonal; lateral lamellae striate; 4a present; pallial sinus long and oval. *Mio.*(*Burdigal.*)-*Rec.*, Eu.—FIG. E96,6. **S. (P.) proaspersa* (SACCO), *Mio.*, Italy; 6a-c, LV ext., LV and RV hinges, $\times 1$ (Sacco, 1901).

- S. (Ruellia)** COSSMANN, 1914 [**Mactra bernayi* COSSMANN, 1886; OD]. Resembling *S. (Pseudoxyperas)* but with lower beaks and umbones; hinge with 4a, as in *S. (Oxyperas)*, 2a-2b not fused, laterals not striate. *Eoc.*, Eu.—FIG. E96,2. **S. (R.) bernayi* (COSSMANN), France; 2a-c, LV ext., RV, LV int., $\times 1.3$ (Cossmann, 1886).
- S. (Spisulona)** MARWICK, 1948 [**Mactra crassitesta* FINLAY, 1927; OD]. Larger than *S. (Spisula)*; hinge with strong spur; pallial sinus shallow. *Plio.*, S.Pac.—FIG. E96,4. **S. (S.) crassitesta* (FINLAY), N.Z.; 4a,b, LV ext., int., $\times 1$ (Hutton, 1893).
- S. (Standella)** GRAY, 1853 [**Mactra striatella* LAMARCK, 1818; SD STOLICZKA, 1871] [= *Leptospisula* DALL, 1895 (obj.)]. Thin, inflated, beaks undulated, dorsal areas smooth; gaping; hinge concentrated, laterals smooth, resilifer not roofed; pallial sinus large, deep. *Rec.*, E.S.Am.-W.Afr.—FIG. E96,3. **S. (S.) striatella* (LAMARCK), W.Afr.; 3a-c, RV ext., int., LV hinge, $\times 1$ (H. Adams & A. Adams, 1856).
- S. (Stereomacra)** STEWART, 1930 [**Schizodesma abscissa* GABB, 1866; OD]. Like *Scissodesma* but shell larger, heavier, laterals longer; resilium shallow. *U.Mio.-Plio.*, W.N.Am.—FIG. E96,5. **S. (S.) abscissa* (GABB), *Mio.*, USA (Calif.); 5a, LV hinge, $\times 0.5$ (Packard, 1916); 5b, RV ext., $\times 0.3$ (892).
- S. (Symmorphomacra)** DALL, 1894 [**Mactra falcata* GOULD, 1851; M]. With cardinal teeth prominent, thin, posterior arm overhanging resilifer, accessory teeth present, hinge plate flat. *Mio.-Rec.*, W.N.Am.—FIG. E96,1. **S. (S.) falcata* (GOULD), *Rec.*, USA (Calif.); 1a-c, RV ext., LV int., RV int., $\times 0.7$ (Packard, 1916).

Subfamily LUTRARIINAE Adams & Adams,
1856

Inequilateral, shell widely gaping; hinge somewhat irregular, concentrated, laterals tending to be obsolete; resilifer broadly open. Siphons of animal not retractile within shell, covered to tips with rough epidermis. *Mio.-Rec.*

Lutraria LAMARCK, 1799 [**Mya lutraria* LINNÉ, 1758; T] [= *Cacophonia* GISTEL, 1848 (*nom. van., pro Lutraria*); *Eustylon* GISTEL, 1848 (*nom. van., pro Cacophonia*); *Lutaria* MAYER, 1875 (*nom. null.*)]. Elongate, thin, gaping at each end; ligament short; no lunule or escutcheon. *Mio.-Rec.*, cosmop.

L. (Lutraria). Outline elliptical; hinge with 2a-2b

heavy, prominent. *Mio.-Rec.*, Eu.-IndoPac.—FIG. E97,5. **L. (L.) lutraria* (LINNÉ), *Rec.*, *Medit.*, 5a-c, RV ext., int., LV hinge, $\times 1$ (H. Adams & A. Adams, 1856); 5d-e, LV and RV hinges, $\times 2$ (510).

L. (Goniomacra) MAYER, 1867 [**L. impar* REEVE, 1854; OD]. Elongate, subquadrangular, with concentric folds, especially on anterior and posterior slopes. *Rec.*, Pac.—FIG. E97,6. **L. (G.) impar* REEVE, Australia; LV ext., $\times 0.5$ (Reeve, 1854).

L. (Lutromacra) IREDALE, 1929 [**L. impedita* (*pro L. elongata* GRAY, 1837) (*non* MÜNSTER, 1835); OD]. Posterior end somewhat pointed. *Rec.*, Pac.—FIG. E97,8. **L. (L.) impedita* IREDALE, Australia; LV ext., $\times 0.5$ (Reeve, 1854).

L. (Lutrophora) DALL, 1895 [**Mactra complanata* GMELIN, 1791; OD]. Elongate, evenly rounded at both ends, with texture and surface of *Raeta*. *Rec.*, Pac.—FIG. E97,2. **L. (L.) complanata* (GMELIN), E.Indies; 2a,b, LV and RV hinges, $\times 2$ (510).

L. (Psammophila) BROWN, 1827 (*ex* LEACH MS, *in synon.*) [**L. solenoides* LAMARCK, 1801 (= *Mya oblonga* GMELIN, 1791); M]. Posterior dorsal margin concave; hinge with 2a-2b weak; ligament partially separated from resilium. *Rec.*, Atl.—FIG. E97,9. **L. (P.) oblonga* (GMELIN), Eu.; 9a,b, RV ext., LV int., $\times 1$ (124).

Eastonia GRAY, 1853 [**Mactra rugosa* HELBLING, 1779; M]. Radiately ribbed or striate except at ends, ovate; pallial sinus deep. *Rec.*, Eu.-*Medit.*—FIG. E97,7. **E. rugosa* (HELBLING), Eu.; 7a, RV ext., $\times 0.7$ (H. Adams & A. Adams, 1856); 7b,c, LV and RV hinges, $\times 1$ (510).

Heterocardia DESHAYES, 1855 [**H. gibbosula*; SD STOLICZKA, 1871]. Short, dorsal slope arched, surface concentrically striate, with vermiculate wrinkling; ligament as in *Mactra*; posterior laterals in each valve. *Rec.*, Pac.—FIG. E97,1. **H. gibbosula*, Philip. Is.; 1a-c, LV int., RV hinge, LV ext., $\times 1$ (H. Adams & A. Adams, 1856).

Meropesta IREDALE, 1929 [**M. meridiana*; OD] [= *Merope* H. & A. ADAMS, 1856 (*non* NEWMAN, 1838) (type, *Mactra nicobarica* GMELIN, 1791; SD STOLICZKA, 1871); *Standella* AUCTT. (*non* GRAY, 1853)]. Resembling *Eastonia* but thinner, ovate, with radiating sculpture; hinge deep; pallial sinus broadly rounded, long, reaching mid-line of shell. *Rec.*, IndoPac.—FIG. E97,4. **M. nicobarica meridiana*, Australia; LV ext., $\times 0.5$ (Allan, 1950).

Tresus GRAY, 1853 (Jan.) [**Lutraria maxima* MIDENDORFF, 1849 (*non* JONAS, 1844) (= *L. nuttalli* CONRAD, 1837); M] [not preoccupied by "*Trésus*" WALCKENAER, 1833, vernacular and *nom. nud.*; = *Cryptodon* CONRAD, 1837 (*non* TURTON, 1822; obj.); *Schizothaerus* CONRAD, 1853 (Feb.; obj.; M)]. Large, ovate, ventricose, gaping posteriorly;

hinge teeth small, resilifer large; ligament separated from resilium by shelly plate. *U.Mio.-Rec.*, W.N.Am.-E.Asia.—FIG. E97,3. **T. nuttalli* (CONRAD), *Rec.*, USA (Calif.); 3a, RV ext., $\times 0.2$ (Stanford Univ. specimen); 3b,c, LV and RV hinges, $\times 1$ (510).

Subfamily PTEROPSELLINAE Keen, new name (1894)

[*nom. subst.* KEEN, herein (*pro* Pteropsinae DALL, 1894, based on junior homonym)]

Shell thin, valves subequilateral, nearly closed; hinge feeble, concentrated, laterals




FIG. E97. Mactridae (Lutrariinae) (p. N604-N605).




FIG. E98. Mactridae (Pteropsellinae) (p. N606).

much reduced or partially obsolete. Siphons of animal wholly retractile, naked. *Eoc.-Rec.*

Pteropsella VOKES, 1956 [*pro Pteropsis* CONRAD, 1860 (*non* RAFINESQUE, 1814)] [**Lutraria papyria* CONRAD, 1833; OD] [= *Kymatox* STENZEL & KRAUSE, 1957 (type, *K. praelapidus*; OD)]. Ovate, posteriorly attenuate; with undulating concentric sculpture; resiliifer ovate; hinge plate deeply grooved. *Eoc.*, SE.N.Am.—FIG. E98,4. **P. papyria* (CONRAD), USA (Ala.); 4a,b, LV ext., int., $\times 0.5$ (Stenzel & Krause, 1957).

Anatina SCHUMACHER, 1817 [**A. pellucida* (= **Mactra anatina* SPENGLER, 1802); M] [= *Labiosa* MÖLLER, 1832 (*ex* SCHMIDT MS) (*nom. van. pro Anatina*); *Cypricia* GRAY, 1847 (*obj.*); *Leucoparia* MAYER, 1867 (*nom. van. pro Cypricia*)]. Thin, ovate, anterior end rounded, posterior end gaping, set off by ridge; lunule and escutcheon evident; external ligament separated from resilium by lamella. *Rec.*, W.C.Am.-E.N.Am.-E.S.Am.—FIG. E98,2. **A. anatina* (SPENGLER), E.S.Am. (?Brazil); 2a,b, RV ext., LV int., $\times 0.7$ (836).

Raeta GRAY, 1853 [**Mactra campechensis* GRAY, 1825 (= **L. plicatella* LAMARCK, 1818); M] [= *Lovellia* MAYER, 1867 (*obj.*)]. Resembling *Anatina*, but more convex, compressed posteriorly, sculpture of concentric plications; dorsal margin not reflected. *Eoc.-Rec.*, Eu.-E.N.Am.-W.C.Am.-S.Am.

R. (Raeta). Larger than *Anatina*, sculpture coarse, posterior end subrounded; hinge with posterior laterals, as in *Anatina*. *Eoc.-Rec.*, Eu.-E.N.Am.-W.C.Am.-S.Am.—FIG. E98,3. **R. (R.) plicatella* (LAMARCK), *Rec.*, W.Indies; 3a-c, LV ext., int., RV hinge, $\times 1$ (H. Adams & A. Adams, 1856).

R. (Raetella) DALL, 1898 [**Labiosa (R.) tenuis* (*ex* ADAMS MS); OD]. Small, thin, dorsal areas well defined, shell inflated; hinge with no lateral lamellae; pallial sinus short, rounded. *Rec.*, E.Asia.

R. (Raetellops) HABE, 1952 [**Poromya pulchella* ADAMS & REEVE, 1850; OD]. Shell small, sculpture coarse; posterior end rostrate; posterior lateral teeth well developed. *Rec.*, E.Asia.—FIG. E98,1. **R. (R.) pulchella* (ADAMS & REEVE), Japan; 1a-d, RV ext., int., RV and LV hinges, $\times 1.5$ (Habe, 1952).

R. (Raetina) DALL, 1898 [**Labiosa (R.) indica*; OD]. Small, posterior end attenuate; posterior laterals wanting. *Rec.*, Ind.O.—FIG. E98,5. **R. (R.) indica* (DALL), India; LV int., $\times 1$ (Dall, 1925).

Subfamily ZENATIINAE Dall, 1895

Inequilateral, compressed; hinge concentrated, irregular, laterals tending to become obsolete; resiliifer bent out of plane of hinge plate, somewhat adherent to valve. *Eoc.-Rec.*

Zenatia GRAY, 1853 [**Lutraria zelandica* GRAY, 1837 (= **L. acinaces* QUOY & GAIMARD, 1835); M] [= *Metabola* MAYER, 1867 (*nom. van., pro Zenatia*)]. Quadratic, periostracum conspicuous; ends gaping, beaks nearer anterior end; valves reinforced by radial buttress below hinge; pallial sinus very deep. *Mio.-Rec.*, S.Pac.

Z. (Zenatia). Moderately elongate, not conspicuously narrow. *Rec.*, S.Pac.—FIG. E99,5. **Z. (Z.) acinaces* (QUOY & GAIMARD), N.Z.; 5a, RV

ext., $\times 1$ (124b); 5b,c, LV and RV hinges, $\times 2$ (510).

Z. (Zenatiopsis) TATE, 1879 [**Zenatiopsis angustata*; M]. Outline narrower than *Z. (Zenatia)*. *Mio.-Rec.*, S.Pac.—FIG. E99,3. **Z. (Z.) angus-*

tata (TATE), Mio., Australia; 3a,b, RV ext., LV int., $\times 1$ (Tate, 1879).

Z. (Zenatraria) BEU, 1966 [**Z. (Z.) vellai*; OD]. *Pleist.*, N.Z.

Darcinia CLARK & DURHAM, 1946 [**D. colombiana*; OD]. With posterior umbonal ridge as in *Darina*; sculpture and posterior gape as in *Eastonia*; hinge plate with deeply excavated resilifer, heavy cardinals, well-developed anterior laterals and nearly obsolete posterior laterals. *Eoc.*, W.S.Am.—FIG. E99,2. **D. colombiana*, Colombia; 2a, RV ext., $\times 1$; 2b, RV hinge, $\times 1.5$ (Clark & Durham, 1946).

Darina GRAY, 1853 [**Erycina solenoides* KING, 1832; M]. Elongate, thin, with periostracum; beaks back of mid-line; both ends gaping; ligament short, deep, resilifer large, resting on ray-like buttress; hinge teeth weak; pallial sinus deep. *Rec.*, W.S.Am.—FIG. E99,1. **D. solenoides* (KING), 1a, LV ext., $\times 1$ (H. Adams & A. Adams, 1856); 1b,c, LV and RV hinges, $\times 2$ (510).

Resania GRAY, 1853 (Jan.) [**R. lanceolata*; M] [= *Vanganella* GRAY, 1853 (Apr.) (type, *V. taylorii*, = *R. lanceolata*; M); *Laminaria*, *Myomactra* MAYER, 1867 (nom. van. pro *Vanganella*, *Resania*)]. Elongate, anterior end longer, smooth, gaping; ligament not set off by shelly ridge, resilifer large, oblique, resting on thickened radial rib; second rib behind anterior adductor scar; pallial sinus short, broad. *Rec.*, S. Pac.—FIG. E99,4. **R. lanceolata*, N.Z.; 4a,b, RV ext., LV int., $\times 1$ (124b).

Family ANATINELLIDAE Gray, 1853

[nom. correct. THIELE, 1934 (pro *Anatinellidae* GRAY, 1853)]

Shell inflated, gaping; hinge with prominent narrow resilifer, narrow cardinal and accessory lamella in each valve; laterals wanting; pallial line without sinus (510). *Rec.*

Anatinella SOWERBY, 1833 [**A. sibbaldii* (= *Mya nicobarica* GMELIN, 1791); M]. Thin, sculpture of concentric striae and fine radial lines; ligament short, separated from resilium by shelly plate. *Rec.*, IndoPac.—FIG. E100,2. **A. nicobarica* (GMELIN), Japan; 2a,b, RV ext., int., $\times 1$ (Kuroda, 1951).

Family CARDILIIDAE Fischer, 1887

Equivalve, higher than long, beaks inverted, subspiral; external ligament on strong nymph, resilium in large, oblique resilifer; hinge with strong inverted V-shaped cardinal in LV, triangular and weaker lamellar cardinal in RV; pallial line entire (510). ?*Eoc.*, *Oligo.-Rec.*




FIG. E99. Mactridae (Zenatiinae) (p. N606-N607).




FIG. E100. Anatinellidae (2); Cardiliidae (1, 3-4)
(p. N607-N608).

Cardilia DESHAYES, 1835 [*Isocardia semisulcata* LAMARCK, 1819; SD HERRMANNSEN, 1846] [= *Leptina* PICTET, 1855 (non MEIGEN, 1830; obj.)]. Sculpture of radial ribs over at least part of shell; cardinal teeth on buttress below resilifer; one lateral present in RV; posterior muscle scar on myophore or flange projecting from dorsal margin of shell. *L.Mio.-Rec.*, IndoPac.—FIG. E100,4. **C. semisulcata* (LAMARCK), *Rec.*, Japan; 4a-d, LV ext., LV and RV hinges, both valves ant., $\times 1.3$ (Kuroda, 1951).

Cardilona MARWICK, 1943 [*C. bensoni*; OD]. Like *Cardilia* in form and sculpture, with strong resilifer, but lacking hinge plate, cardinal teeth, and posterior myophore. *Oligo.*, S.Pac.—FIG. E100,3. **C. bensoni*, N.Z.; 3a,b, RV ext., int., $\times 0.8$ (Marwick, 1943).

?**Hemicyclonosta** MICHELIN, 1828 [*H. michelini* "DESHAYES" (= *Hemicyclodonta* DESHAYES, 1850, ex MS); M] [= *Hemicyclostera* BRONN, 1838, *Hemicyclostera* PAETEL, 1875 (nom. null.), spelling errors] [The evidence for valid publication by MICHELIN is inconclusive, although accepted by

his contemporaries; should the work be rejected, the generic name may revert to synonymy with *Cardilia*]. Resembling *Cardilia* but without radial sculpture. *Eoc.-Plio.*, Eu.—FIG. E100,1. **H. michelini*, Eoc., France, 1a-c, LV ext., RV int., LV int., $\times 1.3$ (259).

Family MESODESMATIDAE Gray, 1839

[*nom. correct.* DALL, 1895 (pro *Mesodesmidae* GRAY, 1839)
[= *Paphiinae* H. ADAMS & A. ADAMS, 1856]]

Shells cuneiform, more or less compressed, disproportionately heavy. Siphons retractile, naked, nearly or completely separated (507, 510). *Eoc.-Rec.*

Subfamily MESODESMATINAE Gray, 1840

[*nom. transl. et correct.* DALL, 1895 (ex *Mesodesmidae* GRAY, 1840)]

Texture porcelaneous, periostracum conspicuous; ligament small or obsolete, resilium narrow, oblique; hinge in Eocene forms with cardinal lamella in RV, weak or obsolete in Recent forms but long narrow cardinal in LV becoming stronger, crossing apex of resilifer. *Eoc.-Rec.*

Mesodesma DESHAYES, 1832 [*Mactra donacia* LAMARCK, 1818; SD ANTON, 1839] [= "*Amphidesma* LAM." AUCT. (non LAMARCK); *Ceronia* GRAY, 1853 (type, *Erycina denticulata* GRAY, 1825, = *Mactra deaurata* TURTON, 1822; M)]. Donaciform, posterior end short; hinge strong, ligament short, internal, resilifer with raised margins; pallial sinus well marked. *Rec.*, N.Am.-S.Am.-Pac.

M. (Mesodesma). Posterior end obliquely truncate; resilifer broad; lateral teeth striate. *Rec.*, Eu.-E.N.Am.-W.S.Am.-Pac.—FIG. E101,13. ***M. (M.) donacium** (LAMARCK), Chile; 13a,b, LV ext., RV int.; $\times 1$ (124b).

M. (Amesodesma) IREDALE, 1930 [*A. perfuga*; OD]. Posterior end somewhat quadrate; resilifer narrow. *Rec.*, S.Pac.—FIG. E101,2. ***M. (A.) perfuga** (IREDALE), Australia; RV int., $\times 1$ (Iredale, 1930).

Atactodea DALL, 1895 [pro *Paphia* LAMARCK, 1799 (non RÖDING, 1798)] [*Mactra glabrata* GMELIN, 1791; OD]. Subtrigonal, strong, smooth or concentrically sculptured; hinge strong, ligament submarginal; resilium narrow; pallial sinus short. *Rec.*, IndoPac.—FIG. E101,4. ***A. glabrata** (GMELIN), E.Indies; 4a-c, RV int., LV hinge, RV ext., $\times 1$ (H. Adams & A. Adams, 1856).

?**Ceroniola** WILCKENS, 1904 [*Cultellus australis* GABB, 1860; M]. Resembling *Mesodesma* in form; shell gaping at each end; ligamental pit not well developed; hinge with 2 laterals but cardinal area not clear. *U.Tert.*, S.Am.—FIG. E101,8. ***C. australis** (GABB), Chile; 8a-c, LV ext., int., RV int., $\times 1$ (Wilckens, 1904).

Donacilla PHILIPPI, 1836 (ex LAMARCK, vernacular)
 [**D. lamarckii* (= *Amphidesma donacilla* LAMARCK, 1818) (= *Mactra cornea* POLI, 1795); M]

[= "*Donacilla* DE BLAINVILLE, 1819," AUCTT., proposed in synonymy]. Ligament marginal, obsolete; laterals not sulcate, anterior lateral long,




FIG. E101. Mesodesmatidae (Mesodesmatinae) (2,4,7-8,11-14), (Davilinae) (1,3), (Erviliinae) (5-6,9-10) (p. N608-N610).

- posterior short; margin of resilial pit toothlike. *Rec.*, Eu.-S.Pac.—FIG. E101,15. **D. cornea* (POLI), Eu.; 15a, RV ext., $\times 1.3$; 15b,c, RV int., LV hinge, $\times 0.7$ (H. Adams & A. Adams, 1856).
- Mactropsis* CONRAD, 1854 [*pro Triquetra* CONRAD, 1846 (*non* DE BLAINVILLE, 1828)] [**Erycina aequorea* CONRAD, 1833; SD DALL, 1895]. Thick, cardinal plate heavy, cardinals distinct, with subequal arms, laterals striate; ligament and resilium combined, near dorsal border; pallial sinus small but distinct. *Eoc.*, E.N.Am.—FIG. E101,7. **M. aequorea* (CONRAD), USA(Ala.); 7a-c, RV ext., int., LV int., $\times 1.5$ (Harris, 1919).
- ?*Myadesma* CLARK, 1922 [**M. dalli*; OD]. Posterior end short, beaks opisthogyrate; resilifers directed anteriorly, large in LV, sunken in RV; pallial sinus small, broad. *Eoc.-Mio.*, W.N.Am.—FIG. E101,14. **M. dalli*, Oligo., Vancouver Is.; 14a,b, LV int., ext., $\times 0.5$; 14c, RV hinge, $\times 1$ (Clark, 1922).
- Paphis* LESSON, 1830 [**P. roissyana* (= **Mya australis* GMELIN, 1791); OD] [= **Machaena* GRAY, 1843, *ex* LEACH MS (obj.); M]. Ovate-elongate, subequilateral; ligament internal, small, resilifer long, narrow, projecting; pallial sinus angular, small, or wanting. *Rec.*, S.Pac.—FIG. E101,12. **P. australis* (GMELIN), N.Z.; 12a-c, RV int., LV hinge, RV ext., $\times 1$ (H. Adams & A. Adams, 1856).
- Taria* GRAY, 1853 [**T. stokesii* (= **Mesodesma lata* DESHAYES, 1843) (?= **M. quoyi* DESHAYES, 1832); M] [= **Taria* H. ADAMS & A. ADAMS, 1856, AUCTT. (obj., KOBELT, 1881)]. Resembling *M. (Mesodesma)* but more equilateral, lateral teeth smooth. *Rec.*, S.Pac.—FIG. E101,11. **T. quoyi* (DESHAYES), N.Z.; 11a,b, LV ext., RV int., $\times 1$ (124b).

Subfamily DAVILINAE Dall, 1895

Outline somewhat rounded; pallial line without sinus. *Rec.*

- Davila* GRAY, 1853 [**D. polita* (= **Mesodesma planum* HANLEY, 1843); M]. Smooth, compressed, periostracum thin; ligament small, nearly marginal, resilium narrow, elongate; cardinal tooth large and prominent in LV, obsolete in RV. *Rec.*, IndoPac.—FIG. E101,3. **D. plana* (HANLEY), Philip. Is.; RV ext., $\times 1$ (Reeve, 1854).
- Anapella* DALL, 1895 [**Mesodesma triquetrum* HANLEY, 1843; OD]. Resembling *Atactodea* in form; hinge with cardinal tooth bifid in RV, laterals smooth. *Rec.*, W.Pac.-S.Pac.—FIG. E101,1. **A. triquetra* (HANLEY), S.Australia; 1a-c, RV int., ext., LV hinge, $\times 1$ (H. Adams & A. Adams, 1856).

Subfamily ERVILIINAE Dall, 1895

Small, thin, equilateral; ligament marginal, subobsolete, resilium small; hinge

concentrated, laterals small, one cardinal in either valve, RV larger, LV bifid; pallial sinus distinct. *Pleist.-Rec.*

- Ervilia* TURTON, 1822 [**Mya nitens* MONTAGU, 1806; M]. Concentrically striate, periostracum inconspicuous; ligament obsolete. *Pleist.-Rec.*, Atl.-Medit.—FIG. E101,10. **E. nitens* (MONTAGU), *Rec.*, USA(Fla.); 10a,b, RV ext., int., $\times 2.5$ (Smith, 1937).
- Argyrodonax* DALL, 1911 [**A. haycocki*; M]. Concentrically sculptured; external ligament feeble, resilium narrow but strong; muscle scars pronounced; pallial sinus large and deep. *Rec.*, W. Indies.
- Caccella* GRAY, 1853 [**C. horsfieldii*; OD]. Relatively large, elongate, with fine concentric striae; periostracum thick, brown; ligament obsolete; resilium small; pallial sinus not deep. [Estuarine.] *Rec.*, Pac.—FIG. E101,9. **C. horsfieldii*; 9a-c, LV ext., int., int., $\times 3$; 9d, RV hinge, $\times 1$ (specimens, British Museum, N.H.).
- Rochefortina* DALL, 1924 [**Rochefortia* (R.) *semele* (= **Ervilia sandwichensis* E. A. SMITH, 1885); OD]. Sculpture radial and concentric; with minute escutcheon; pallial sinus moderately deep. *Rec.*, Pac.—FIG. E101,6. **R. sandwichensis* (SMITH), Hawaiian Is.; 6a,b, LV ext., hinge, $\times 8$ (852).
- Spondervilia* IREDALE, 1930 [**Ervilia australis* ANGAS, 1877 (= **E. biscalpta* GOULD, 1861); OD]. Sculpture radial at ends, with fine wavy concentric threads on central slope. *Rec.*, Pac.—FIG. E101,5. **S. biscalpta* (GOULD), Japan; 5a,b, LV int., RV hinge, $\times 3$ (Habe, 1952).

Superfamily SOLENACEA Lamarck, 1809

[*nom. transl.* TRYON, 1884 (*ex* family Solenacea GRAY, 1823) (=solenacées LAMARCK, 1809)] [Materials for this superfamily prepared by MYRA KEEN]

Valves cylindrical to flattened, elongate, gaping at both ends, hinge weak, with one to three small cardinal teeth; lateral teeth, if present, weak and laminar; pallial sinus short in most forms. *L.Cret.-Rec.*

Family SOLENIDAE Lamarck, 1809

[*nom. correct.* LEACH, 1823 (*pro* family Solenacea GRAY, 1823) (=solenacées LAMARCK, 1809)]

Beaks terminal or nearly so; hinge with only one tooth in either valve; siphons of animal fused, foot modified for rapid digging in sand. *L.Eoc.-Rec.*

- Solen* LINNÉ, 1758 [**S. vagina*; SD SCHUMACHER, 1817] [= **Vagina* MEGERLE VON MÜHLFELD, 1811 (obj., T); *Solenia* OKEN, 1823 (*nom. null.*); *Listera* GRAY, 1852, *ex* LEACH MS (*non* TURTON,




FIG. E102. Solenidae (p. N611).

1822) (obj.); *Fistula* MÖRCH, 1853, ex MARTINI MS (type, *S. brevis* HANLEY, 1842, ex GRAY MS; SD KEEN, herein)]. Long, nearly straight; anterior margin truncate, smooth within; anterior adductor muscle scar elongate. *Eoc.-Rec.*, N.Am.-Eu.-Pac.—FIG. E102,2. **S. vagina*, Rec., Eu.; 2a-c, LV ext., RV hinge, LV hinge, $\times 0.4$ (124b).

Solena MÖRCH, 1853 [**Solen obliquus* SPENGLER, 1794; SD STOLICZKA, 1871] [= *Hypogella* GRAY, 1854 (type, *S. ambiguus* LAMARCK, 1818, = *S. obliquus*; SD DALL, 1900)]. Anterior end somewhat produced beyond beaks, obliquely truncate; anterior adductor scar short. *Eoc.-Rec.*, Asia-Eu.-Am.

S. (Solena). Anterior margin thickened within, smooth outside. *Oligo.-Rec.*, E.Asia-Eu.-C.Am.—FIG. 102,3. **S. (S.) obliquus* (SPENGLER), Rec., Carib.; LV ext., $\times 0.5$ (124b).

S. (Eosolen) STEWART, 1930 [**Solen plagiaulax* COSSMANN, 1906 (nom. subst. pro *S. obliquus* SOWERBY, 1844, non SPENGLER, 1794); OD]. With external oblique groove near anterior end. *Eoc.*, Eu.-N.Am.—FIG. E102,4. **S. (E.) plagi-aulax* (COSSMANN), M.Eoc., France; 4a-c, RV ext., RV hinge, LV hinge, $\times 0.5$ (Deshayes, 1860).

S. (Plectosolen) CONRAD, 1866 [**Solen gracilis* SOWERBY, 1844; SD STOLICZKA, 1871]. Dorsal margin slightly curved, anterior end rounded, set off by furrow. *L.Eoc.-M.Eoc.*, Eu.-W.N.Am.—FIG. E102,1. **S. (P.) gracilis* (SOWERBY), Eng.; 1a,b, LV ext., both valves, dorsal, $\times 1$; 1c,d, RV, LV hinge, $\times 2.5$ (Sowerby, 1844).

Family CULTELLIDAE Davies, 1935

Valves wider and more compressed than in Solenidae, beaks mostly not terminal; hinge with one to three cardinal teeth. *L. Cret.-Rec.*

Cultellus SCHUMACHER, 1817 [**C. magnus* (= *Solen lacteus* SPENGLER, 1794); M]. Oblong, ends rounded, gaping; hinge with 1 cardinal in RV, 2 in LV, posterior bifid; pallial sinus small. *L.Eoc.-Rec.*, Eu.-Asia-IndoPac.

C. (Cultellus). Large, with rib above rounded anterior adductor scar. *L.Eoc.-Rec.*, Eu.-C.Asia-IndoPac.—FIG. E103,2. **C. (C.) lacteus* (SPENGLER), Rec., E.Indies; RV int., $\times 0.7$ (Woodward).

C. (Cultrensis) COEN, 1933 [**C. (C.) adriaticus*; M]. Small and thin, dorsal margin straight, with 2 furrows from beak to ventral margin. *Rec.*, Medit.

Ensis SCHUMACHER, 1817 [**E. magnus* (= *Solen ensis* LINNÉ, 1758); T] [= *Ensatella* SWAINSON, 1840 (obj.)]. Beaks terminal or nearly so, dorsal margin slightly curved; 2 cardinal teeth or cardinal chevrons present; anterior adductor scar elongate. *L.Eoc.-Rec.*, N.Am.-Eu.—FIG. E103,6. **E. ensis* (LINNÉ), Rec., Medit; 6a,b, LV ext., RV int., $\times 0.3$ (89a).

Leptosolen CONRAD, 1865 [**Siliquaria biplicata* CONRAD, 1858; M] [= *Solenaria* STOLICZKA, 1870 (?non RAFINESQUE, 1815) (type, *Leguminaria affinis* EICHWALD, 1867; M)]. Like small *Siliqua* with radial sulcus and strong internal rib; ligament long and narrow. *L.Cret.-U.Cret.*, N.Am.-E.Eu.—FIG. E103,9. **L. biplicata* (CONRAD), U.Cret., USA(Ala.); 9a,b, LV ext., int., $\times 1$ (Wade).

Neosolen GHOSH, 1920 [**N. aquae-dulcioris*; M]. Small, thin, translucent, truncate in front, pos-




FIG. E103. Cultellidae (p. N611-N613).

teriorly rounded; RV with 1/2 small, long tooth; anterior adductor scar long, triangular, posterior small, rounded. *Rec.*, S.Asia. [Estuarine.]—FIG. E103,10. **N. aquaedulcoris*, India; 10a,b, RV ext., int., $\times 1$ (Ghosh).

Ospriasolen CONRAD, 1868 [**Cultellus cretaceus* GABB, 1860; M] [= *Ospirasolen*, spelling error]. Elongate, gaping, pallial line with deep sinus. *U. Cret.*, N.Am.—FIG. E103,11. **O. cretaceus* (GABB), USA(N.J.); 11a,b, RV int. mold, both valves dorsal, $\times 1$ (Weller, 1907).

Pharella GRAY, 1854 [**Solen javanicus* LAMARCK, 1818; SD KOBELT, 1881]. Subcylindrical, rounded at ends, ventral margin contracted, umbones in front of mid-line; anterior adductor scar elongate, subtrigonal; hinge with 2 cardinals in RV, 3 in LV. *Plio.-Rec.*, E.Indies.—FIG. E103,5. **P. javanica* (LAMARCK), *Rec.*, Java; LV ext., $\times 1$ (Fischer).

Phaxas GRAY, 1852 (ex LEACH MS) [**S. pellucidus* PENNANT, 1777; M] [= *Subcultellus* GHOSH, 1920 (obj.)]. Dorsal margin curved, ends of shell

rounded; beaks near anterior end; anterior adductor scar long and triangular, posterior scar small, sinus short. *Eoc.-Rec.*, Asia-Eu.-E.Indies.

P. (Phaxas). Shell small and thin. *Eoc.-Rec.*, E. Asia-Eu.—FIG. E103,1. **P. (P.) pellucidus* (PENNANT), *Rec.*, Eng.; 1a, LV ext., $\times 1$ (Locard); 1b,c, RV int., LV hinge, $\times 1.3$ (Stanford Univ. specimen).

P. (Ensiculus) H. ADAMS, 1860 [**Solen cultellus* LINNÉ, 1758; OD] [= *Cultellus* AUCTION (non SCHUMACHER)]. Larger and more solid than in *P. (Phaxas)*. *Rec.*, E.Indies.—FIG. E103,3. **P. (E.) cultellus* (LINNÉ); LV ext., $\times 0.5$ (124b).

?**Senis** STEPHENSON, 1952 [1953] [**S. elongatus*; OD]. Long-quadrangular, beaks slightly in front of mid-line; ligament external, long, narrow, nymph thin and prominent; hinge smooth, without teeth. *U. Cret.*, N.Am.—FIG. E103,7. **S. elongatus*, USA(Tex.); LV ext., $\times 1.5$ (890).

Siliqua MEGERLE VON MÜHLFELD, 1811 [**Solen radiatus* LINNÉ, 1758; M] [= *Aulus* OKEN, 1815,

AUCT. (in work rejected by ICZN, 1956, Op. 417) (obj.; SD HERRMANNSEN, 1846); *Leguminaria* SCHUMACHER, 1817 (obj.; M); *Solecurtoides* DESMOULINS, 1832 (obj.; SD DALL, 1900); *Machaera* GOULD, 1841 (non CUVIER, 1832) (type, *Solen costatus* SAY, 1822; SD DALL, 1900)]. Thin-shelled, moderately large, with internal rib; pallial sinus widely rounded, fairly deep. *Eoc.-Rec.*, N. Am.-Eu.-Pac.-Asia.

S. (Siliqua). Internal rib nearly vertical, from beaks to ventral margin. *Eoc.-Rec.*, N. Am.-Eu.-E. Indies. —FIG. E103,8. *S. (S.) *radiata* (LINNÉ), *Rec.*, E. Indies; LV ext., $\times 0.3$ (124b).

S. (Neosiliqua) HABE, 1965 [**Aulus winterianus* DUNKER, 1853; OD]. Internal rib narrow, running diagonally from beaks to anteroventral margin. *Rec.*, W. Pac.

?**Tanysiphon** BENSON, 1858 [**T. rivalis*; M]. Resembling *Glaucanome* in Veneracea, small, thin, long-ovate, with greenish periostracum; beaks in front of mid-line, ligament short; hinge with 3 teeth in either valve, anterior 2 smaller, posterior tooth flattened, resembling a resilifer; pallial sinus large, rounded, extending beyond mid-line. *Rec.*, S. Asia. [Estuarine.] —FIG. E103,4. **T. rivalis*, India; LV ext., $\times 1$ (Benson, 1858).

Superfamily TELLINACEA de Blainville, 1814

[*nom. transl.* DALL, 1895 (ex family Tellinacées DE BLAINVILLE, 1814)] [Materials for this superfamily prepared by MYRA KEEN except as otherwise indicated]

Shells mostly inequilateral with external ligament or, if sunken, in pit on hinge plate, not in chondrophore; cardinal hinge teeth two in either valve, tending to be bifid, lateral teeth well developed in most families; adductor muscle scars connected by pallial line with distinct sinus. Animal with two elongate siphons that are not fused. *U. Trias.-Rec.*

Family TELLINIDAE de Blainville, 1814

[*nom. correct.* SWAINSON, 1840 (pro family Tellinacées DE BLAINVILLE, 1814)]

Somewhat elongate; ligament external; valves more or less unequal, most forms with posterior flexure, especially in RV (822). *L. Cret.-Rec.*

Subfamily TELLININAE de Blainville, 1814

[*nom. transl.* H. ADAMS & A. ADAMS, 1856 (ex family Tellinacées DE BLAINVILLE, 1814)]

Lateral teeth present in at least one valve; sculpture various. *L. Cret.-Rec.*

Tellina LINNÉ, 1758 [**T. radiata*; SD CHILDREN, 1823] [= *Musculus* MÖRCH, 1853 (ex MARTINI, nonbinom.) (non RÖDING, 1798); *Liotellina* FISCHER, 1887 (obj.); *Tellinarius*, *emend.*]. Characteristics of subfamily. [Most modern attempts to divide *Tellina*, s.l., into a number of genera, each with several subgenera, abound in inconsistencies. Evolutionary history of the family in different ocean basins seems to have resulted in many parallel forms or homeomorphs, so that subdivision on one set of characters (e.g., shell outline) runs counter to that on another (e.g., hinge details or musculature). A proper review being beyond the scope of the present summary, the expedient of a conservative classification is adopted herein, treating most generic taxa as subgenera of *Tellina*.] ?*Cret.*, *Tert.-Rec.*, cosmop.

T. (Tellina). Smooth, polished, hinge with 2 cardinals and 2 laterals in either valve. *Rec.*, E. N. Am. W. Indies. —FIG. E104,11. **T. (T.) radiata* LINNÉ, W. Indies; 11a,b, LV ext., RV hinge, $\times 0.7$ (Chenu).

T. (Abranda) IREDALE, 1924 [**A. rex* (pro *T. elliptica* SOWERBY, 1868) (non BROCCHI, 1814); OD]. Small, compressed, elliptical; concentrically ridged behind the angle setting off posterior slope. *Rec.*, Australia. —FIG. E104,1. **T. (A.) rex* (IREDALE); LV ext., $\times 1$ (Sowerby in Reeve).

T. (Acorylus) OLSSON & HARBISON, 1953 [**T. suberis* DALL, 1900; OD]. Small, solid, sub-ovate, flexed; hinge stout, RV with 2 cardinals, 2 laterals, LV with 1 cardinal, no laterals; pallial sinus deep, widely confluent with pallial line. *Plio.*, E. N. Am. —FIG. E104,7. **T. (A.) suberis* DALL, USA (Fla.); 7a, RV ext., $\times 3$ (Olsson & Harbison); 7b, RV int., $\times 4$ (Dall, 1900).

T. (Angulus) MEGERLE VON MÜHLFELD, 1811 [**T. lanceolata* GMELIN, 1791; SD GRAY, 1847]. Of moderate size, elongate, posterior end pointed but not twisted; hinge with *Al* strong, near cardinals, other laterals wanting, cardinals small. *Rec.*, Pac. —FIG. E104,8. **T. (A.) lanceolata* GMELIN, E. Indies; 8a, RV ext., $\times 1$ (Chenu), 8b, RV hinge, $\times 2$ (Olsson & Harbison).

T. (Arcopagia) BROWN, 1827 (ex LEACH MS) [**T. crassa* PENNANT, 1777; SD HERRMANNSEN, 1846] [= *Cydippe* GRAY, 1852 (ex LEACH MS) (non ESCHSCHOLTZ, 1829) (obj.)]. Rounded, somewhat inflated, concentrically ridged, with 2 laterals in RV; pallial sinus deep, rounded, not confluent. [See also *T. (Sinuosipagia)*.] ?*Cret.*, *Eoc.-Rec.*, Eu.-N. Afr. —FIG. E104,6. **T. (A.) crassa* PENNANT, *Rec.*, *Medit.*; 6a,b, LV lat., $\times 0.5$ (Reeve); 6c, RV int., $\times 0.5$ (Davies).

T. (Arcopaginula) LAMY, 1918 [**T. inflata* GMELIN, 1791; M]. Medium-sized, white, smooth, posterior area set off by rib; inequivalve, LV larger; *Al* and *All* short, *Pl* long, *Pll* weak; pallial sinus confluent. *Rec.*, W. Pac. —FIG.

E104,12. **T. (A.) inflata* GMELIN, Japan; 12a,b, RV ext., int., $\times 1$ (Habe).
T. (Arcopella) THIELE, 1934 [**T. balaustina* LINNÉ, 1758; M]. Near *T. (Arcopagia)* but

smaller, posterior end shorter; lateral teeth weak in LV. *Rec.*, Eu.—FIG. E104,5. **T. (A.) balaustina* LINNÉ, France; 5a-c, RV ext., int., LV int., $\times 1$ (89a).




FIG. E104. Tellinidae (Tellininae) (p. N613-N615).

- T. (Cadella)** DALL, BARTSCH, & REHDER, 1939 [**T. lechriogramma* MELVILLE, 1893; OD]. Small, resembling *T. (Semelangulus)* in outline and sculpture but with normal ligament; also resembling *T. (Moerella)* but with smaller pallial sinus. *Rec.*, Indo-Pac.—FIG. E104,2, **T. (C.) lechriogramma* MELVILLE, India; LV ext., $\times 1.5$ (Melville).
- T. (Clathrotellina)** THIELE, 1934 [**T. pretiosa* DESHAYES, 1855 (*non* EICHWALD, 1830)=*T. pretium* SALISBURY, 1934; M]. Small, with cancellate sculpture, otherwise resembling *T. (Merisca)*. *Rec.*, IndoPac.—FIG. E104,9. *T. (C.) pretium* SALISBURY, E.Indies; RV ext., $\times 2$ (Reeve).
- T. (Cyclotellina)** COSSMANN, 1887 [**Donax lunulata* LAMARCK, 1805; OD]. Shape as in *T. (Arcopagia)* but with posterior lateral in LV; pallial sinus smaller, partly confluent, angular, joined to adductor by linear scar. *Eoc.-Rec.*, Eu.-Carib.-IndoPac.—FIG. E104,4. **T. (C.) lunulata* (LAMARCK), Eoc., France; 4*a,b*, RV ext., int., $\times 1$ (Deshayes).
- T. (Elliptotellina)** COSSMANN, 1887 [**Donax tellinella* LAMARCK, 1805; OD]. Near *T. (Tellinella)* but smaller, with lateral teeth weak or wanting in LV. *L.Eoc.-M.Eoc.*, Eu.—FIG. E104,3. **T. (E.) tellinella* (LAMARCK), France; 3*a-e*, RV ext., int., LV ext., int., RV int., $\times 2$ (Cossmann).
- T. (Elpidollina)** OLSSON, 1961 [**T. decumbens* CARPENTER, 1865; OD]. Thin, subtrigonal, somewhat inflated, cardinal teeth small, laterals large in RV, pallial sinus large and deep. *Rec.*, W.C. Am.—FIG. E105,11. **T. (E.) decumbens* CARPENTER, Panama; 11*a,b*, LV ext., RV int., $\times 0.7$ (Reeve; Olsson).
- T. (Eurytellina)** FISCHER, 1887 [**T. punicea* BORN, 1790; M]. Medium-sized, elongate, posterior end not rostrate; sculpture concentric; hinge with 4*b* small, laterals in LV not as strong as in RV, anterior laterals close to cardinals, posterior distant; pallial sinus confluent with pallial line, approaching or touching anterior adductor. *Mio.-Rec.*, E.N.Am.-W.N.Am.-S.Am.—FIG. E105,1. **T. (E.) punicea* BORN, *Rec.*, W.Indies; 1*a-c*, LV ext., RV and LV hinges, $\times 1$ (Chenu).
- T. (Fabulina)** GRAY, 1851 [**T. fabula* GMELIN, 1791; SD WINGKORTH, 1932] [= *Tellinula* MÖRCH, 1853 (*ex* CHEMNITZ) (obj.)]. Ovate, with oblique sculpture on RV except at ends; hinge with only 1 lateral, *AI*, near cardinals. [See also *T. (Scissula)*.] *Mio.-Rec.*, Eu.—FIG. 105,7. **T. (F.) fabula* GMELIN, *Rec.*, Medit.; 7*a,b*, LV ext., RV ext., $\times 1$ (Reeve).
- T. (Finlayella)** LAWS, 1933 [**F. sinuaris*; OD]. Small, pallial sinus asymmetrically developed, deeper in LV than in RV. *Mio.-Plio.*, S.Pac.—FIG. E105,8. **T. (F.) sinuaris* (LAWS), Mio., N.Z.; 8*a-c*, RV ext., LV int., RV int., $\times 2$ (Laws).
- T. (Gastranopsis)** COSSMANN, 1906 [**G. bureauii*; OD]. Resembling *Gastrana* but with lateral teeth in RV. *Eoc.*, Eu.—FIG. E105,2. **T. (G.) bureauii* (COSSMANN), France; 2*a-d*, RV ext., LV int., LV ext., RV int., $\times 3$ (Cossmann).
- T. (Hemimetis)** THIELE, 1934 [**T. plicata* VALENCIENNES, 1827; M]. Thin, somewhat lenticular, resembling *Apolymetis* but with lateral teeth and a lunule in RV; sculpture of concentric lamellae; ligament narrow; pallial sinus not confluent. *Rec.*, IndoPac.—FIG. E105,5. **T. (H.) plicata* VALENCIENNES, Japan; 5*a,b*, LV ext., RV int., $\times 0.5$ (Habe).
- T. (Hertellina)** OLSSON, 1961 [**T. nicoyana* HERTLEIN & STRONG, 1949; OD]. Outline as in *Sanguinolaria* but hinge and oblique sculpture as in *T. (Scissula)*. *Rec.*, W.C.Am.—FIG. E105,3. **T. (H.) nicoyana* HERTLEIN & STRONG, W. Costa Rica; 3*a-c*, RV ext., LV and RV hinges, $\times 1$ (Hertlein & Strong).
- T. (Homalina)** STOLICZKA, 1870 [**T. triangularis* "Chemnitz" DILLWYN, 1817 (*non* GMELIN, 1791) (= *T. trilatera* GMELIN, 1791); OD]. Thin, trigonal, posterior end slightly longer; hinge weak, anterior lateral teeth short, close to cardinals, posterior laterals at end of ligament; pallial sinus deep, confluent. [See also *T. (Macomona)*.] *Rec.*, E.Indies.—FIG. E105,9. **T. (H.) trilatera* GMELIN, E.Indies; 9*a,b*, LV ext., RV ext., $\times 1$ (Reeve, 1867).
- T. (Iraqitellina)** DANCE & EAMES, 1966 [**I. iraqensis*; OD]. *Rec.*, Persian Gulf.
- ?**T. (Jactellina)**. [See *Exotica (Jactellina)*, subfamily Macominac.]
- T. (Laciolina)** IREDALE, 1937 [**T. quoyi* SOWERBY in REEVE, 1868; OD] [= *Bosempra* BROWN, 1844, *ex* LEACH MS (invalidly proposed in synonymy of *Tellina depressa* GMELIN, 1791)]. Large, oblique, smooth, posterior end shorter; ligament long, somewhat sunken. *Rec.*, Pac.O.-W.Atl.-Eu.-N.Am.—FIG. E104,10. *T. (L.) incarnata* LINNÉ, France; 10*a,b*, RV ext., $\times 1$; 10*c*, RV int., $\times 1$ (89*a*).—FIG. E105,4. **T. (L.) quoyi* SOWERBY, N.Australia; RV ext., $\times 0.7$ (Reeve, 1868).—FIG. E106,15. *T. (L.) ochracea* (CARPENTER), W.Mexico; 15*a*, RV ext., $\times 0.5$; 15*b*, LV hinge, enl. (Hertlein & Strong, 1940-51).
- ?**T. (Loxoglypta)** DALL, BARTSCH, & REHDER, 1939 [See *Exotica (Loxoglypta)*, subfamily Macominac.]
- T. (Lyratellina)** OLSSON, 1961 [**T. lyra* HANLEY, 1844; OD]. Elliptical, posterior end unflexed; escutcheon deep and narrow, below valve margin. *Mio.-Rec.*, NW.S.Am.-W.C.Am.—FIG. E105,10. **T. (L.) lyra* HANLEY, *Rec.*, Panama; 10*a*, LV ext., $\times 0.7$; 10*b,c*, LV hinge, RV hinge, $\times 2$ (Olsson, 1961).
- T. (Macaliopsis)** COSSMANN, 1887 [**T. barrandei* DESHAYES, 1857; SD DALL, 1900]. Rounded-trapezoidal, posterior end truncate, sculpture of spaced concentric ridges; cardinals and laterals

strong, remote; pallial sinus narrow, only slightly confluent. *Paleoc.-Mio.*, Eu.-E.C.Am.—
 FIG. E107,6. **T. (M.) barrandei* DESHAYES, Eoc.,
 France; 6*a,b*, LV ext., RV int., $\times 1$ (Deshayes).
T. (Macomona) FINLAY, 1927 [**T. liliana* IRE-

DALE, 1915; OD]. Resembling *T. (Homalina)*
 but larger, compressed, posterior end twisted and
 somewhat pointed; RV with 2 cardinals and 2
 laterals, posterior weak; pallial sinus deep, con-
 fluent, reaching to anterior muscle scar. *Tert.-*




FIG. E105. Tellinidae (Tellininae) (p. N615-N618).

Rec., Australia.—FIG. E105,6. **T. (M.) liliana* IREDALE, Rec., N.Z.; LV ext., $\times 0.3$ (Powell).
T. (Merisca) DALL, 1900 [**T. crystallina* WOOD, 1815—*T. crystallina* SPENGLER, 1798; OD].

Small to moderate-sized, posterior end rostrate, with basal margin emarginate in front of rostrum; sculpture of distant thin concentric lamellae; hinge without laterals in LV; pallial sinus




FIG. 106. Tellinidae (Tellininae) (p. N615-N620, N625).




FIG. E107. Tellinidae (Tellininae) (p. N615-N616, N618-N619).

deep, confluent. *Mio.-Rec.*, E.C.Am.-W.C.Am.—FIG. E105, 12. **T. (M.) cristallina* SPENGLER, *Rec.*, W.Indies; RV ext., $\times 1$ (Dall).

T. (Moerella) FISCHER, 1887 [**T. donacina* LINNÉ, 1758; M] [= *Donacilla* GRAY, 1851 (*non* PHILIPPI, 1836) (obj.); SD SALISBURY, 1934]; *Moera* H. & A. ADAMS, 1856 (*non* HUEBNER, 1819) (obj.; SD STOLICZKA, 1871); *Bathytellina* KURODA & HABE in HABE, 1958 (type, *B. citrocarnea*). Rather small, elongate-ovate, posterior end obscurely rostrate; sculpture of concentric threads; hinge with 3a bifid, 2 very small, laterals strong, anterior pair closer to cardinals; pallial sinus confluent, almost touching anterior adductor scar. *L.Eoc.-Rec.*, Eu.-N.Am.-Pac.—FIG. E107, 7. **T. (M.) donacina* LINNÉ, *Rec.*, Eu.; 7a-d, LV

ext., RV ext., LV and RV hinges, $\times 1$ (Chenu). [= *Asbjornsenia* FRIELE, 1886 (type, *A. striata*; M). Dr. OCKELMANN of Helsingør, Denmark, has studied the type specimens and has reported that they represent young *Tellina* (*Moerella*) *pygmaea* LOVÉN.]

T. (Nitidotellina) SCARLATO, 1965 [**Tellina nitidula* DUNKER, 1860; OD]. *Rec.*, E.Asia.

T. (Obtellina) IREDALE, 1929 [**T. bougei* SOWERBY, 1909; OD]. Beaks near posterior end; surface with fine oblique lines and irregular spines on posterior margin; pallial sinus largely confluent. *Rec.*, Pac.—FIG. E107, 2. **T. (O.) bougei* SOWERBY, New Caledonia; LV ext., $\times 1.5$ (Sowerby).

T. (Omala) SCHUMACHER, 1817 [**O. inaequalvis*

- Tellina hyalina* GMELIN, 1791; M]. Ovate, thin, pellucid, ligament long, sunken; hinge with *AI* close to cardinals. *Rec.*, Pac.—FIG. E107,1. **T. (O.) hyalina* GMELIN, Japan; 1a-c, RV ext., int., hinge, $\times 0.5$ (Habe).
- T. (Oudardia)** MONTEROSATO, 1884 [**T. oudardii* PAYRAUDEAU, 1826 =*T. compressa* BROCCHI, 1814; OD]. Resembling *T. (Moerella)* in form but with a thick internal rib, radiating from beak to ventral margin in front of mid-line; surface with or without fine oblique lines of sculpture. ?*Eoc.*, *Oligo.-Rec.*, Eu.-W.N.Am.—FIG. E107,4. **T. (O.) compressa* BROCCHI, Plio., Eu.; 4a-c, RV ext., LV int., RV int., $\times 1$ (Hörnes).
- T. (Peronaca)** POLI, 1791 [**T. planata* LINNÉ, 1758; SD STOLICZKA, 1870] [= *Peronacoderma* POLI, 1795 (obj.; SD KEEN, herein)]. Ovate, posteriorly rounded, inequivalvae, LV flatter; sculpture of incremental lines and a few fine radials; lateral teeth weak; ligament long, depressed; pallial sinus confluent. *U.Oligo.-Rec.*, Eu.—FIG. E107,5. **T. (P.) planata* LINNÉ, Mio., Aus., 5a-c, LV ext., int., RV int., $\times 0.7$ (Zittel).
- T. (Peronidia)** DALL, 1900 [**T. albicans* GMELIN, 1791; OD]. Elongate, solid, compressed, subequivalve, without lunule, escutcheon long, narrow; shell with fine concentric grooves, stronger anteriorly; lateral teeth weak. *Eoc.-Rec.*, Eu.-W.N.Am.-E.Asia.—FIG. E107,3. **T. (P.) albicans* GMELIN, *Rec.*, Algeria; 3a-c, RV ext., LV int., RV int., $\times 0.7$ (89a).
- T. (Pharaonella)** LAMY, 1918 [**T. pharaonis* HANLEY, 1844; SD SALISBURY, 1934]. Elongate, posterior end strongly rostrate, with a double radial rib. *Rec.*, Pac.—FIG. E106,1. **T. (P.) pharaonis* HANLEY, Red Sea; RV ext., $\times 0.5$ (Reeve).
- T. (Phylloda)** SCHUMACHER, 1817 [**P. aurea* =*T. foliacea* LINNÉ, 1758; M]. Large, thin, ovate, posterior dorsal margin serrate, posterior slope with spinose radials; with 2 laterals in RV, laterals weak or absent in LV. *U.Oligo.-Rec.*, Eu.-Pac.—FIG. E106,12. **T. (P.) foliacea* LINNÉ, *Rec.*, E.Indies; LV ext., $\times 0.5$ (Chenu).
- T. (Phyllodella)** HERTLEIN & STRONG, 1949 [**T. insculpta* HANLEY, 1844; OD]. Like *T. (Eurytelina)* but with the foliations of *T. (Phyllodina)* on posterior slope. *Rec.*, W.C.Am.—FIG. E106,3. **T. (P.) insculpta* HANLEY; RV ext., $\times 1$ (Hertlein & Strong).
- T. (Phyllodina)** DALL, 1900 [**T. squamifera* DESHAYES, 1855; OD]. With concentric sculpture forming spines along posterior dorsal margin; pallial line not confluent. *Oligo.-Rec.*, E.C.Am.-W.C.Am.—FIG. E106,4. **T. (P.) squamifera* DESHAYES, *Rec.*, W.Indies; RV ext., $\times 1$ (Reeve).
- T. (Pinguitellina)** IREDALE, 1927 [**T. robusta* HANLEY, 1844; OD]. Small, inflated, smooth, posterior slope set off by an angle; pallial sinus long. *Rec.*, Pac.—FIG. E106,5. **T. (P.) robusta* HANLEY, NE.Australia; 5a,b, LV int., RV ext., $\times 2$ (Iredale).
- T. (Pistris)** THIELE, 1934 [*pro Pristis* LAMY, 1918 (non LINCK, 1790)] [**T. pristis* LAMARCK, 1818; M] [= *Pristipagia* IREDALE, 1936 (type, *P. gemonia*; OD)]. Small, relatively high, ligament deep; sculpture of wavy ribs and fine impressed radial lines; pallial sinus free; resembling *T. (Serratina)* but higher, with stronger sculpture. *Rec.*, Pac.—FIG. E106,16. **T. (P.) pristis* LAMARCK, E.Indies; LV ext., $\times 1$ (Reeve).
- T. (Pseudarcopagia)** BERTIN, 1878 [**T. decussata* LAMARCK, 1818 (non WOOD, 1815) =*T. victoriae* GATLIFF & GABRIEL, 1914; SD DALL, 1900]. Lenticular, sculpture decussate; hinge with 2 laterals in either valve. [See also *T. (Arcopagia)*, *T. (Sinuosipagia)*.] *Rec.*, Pac.—FIG. E106,11. **T. (P.) victoriae* GATLIFF & GABRIEL, Australia; LV ext., $\times 0.5$ (Reeve).
- T. (Punipagia)** IREDALE, 1930 [**T. subelliptica* SOWERBY, 1868, in REEVE (non MEEK & HAYDEN, 1857) =*T. hypelliptica* SALISBURY, 1934; OD] [= *Punigapia* THIELE, 1934, spelling error]. Small, rounded, ligament short, somewhat sunken; pallial sinus deep, triangular, about half confluent. *Rec.*, Pac.—FIG. E106,2. *T. (P.) hypelliptica* SALISBURY, Australia; 2a,b, LV ext., $\times 1$ (Reeve).
- T. (Quadrans)** BERTIN, 1878 [**T. gargadia* LINNÉ, 1758; SD DALL, 1900]. Shell thin, posterior slope wrinkled, the margin with spines; ligament sunken; laterals feeble, 2 in RV, wanting in LV, pallial sinus confluent below. *Rec.*, Pac.—FIG. E106,7. **T. (Q.) gargadia* LINNÉ, E.Indies; LV ext., $\times 1$ (Reeve).
- T. (Quidnipagus)** IREDALE, 1929 [**Cochlea palatam* Martyn" IREDALE, 1929 (= *T. rugosa* BORN, 1778, non PENNANT, 1777); OD]. Somewhat attenuate posteriorly, lunule and escutcheon small; surface with irregular wrinkles and radial furrows; pallial sinus half free. *Rec.*, Pac.—FIG. E106,13. **T. (Q.) palatam* (IREDALE), E.Indies; 13a,b, LV ext., RV ext., $\times 0.5$ (Reeve).
- T. (Scissula)** DALL, 1900 [**T. decora* SAY, 1826 =*T. similis* SOWERBY, 1806; OD]. Ovate-quadrate, surface of both valves with fine oblique grooving not in harmony with incremental lines [see also *T. (Fabulina)*]. *Plio.-Rec.*, E.C.Am.-W.C.Am.—FIG. E106,9. **T. (S.) similis* SOWERBY, Plio., USA (Fla.); RV ext., $\times 2$ (Olsson & Harbison).
- T. (Scutarcopagia)** PILSBRY, 1918 [**T. scobinata* LINNÉ, 1758; OD]. Lenticular, inflated, with scaly or granose sculpture. *Pleist.-Rec.*, IndoPac.—FIG. E106,8. **T. (S.) scobinata* LINNÉ, *Rec.*; 8a, NE.Australia, LV ext., $\times 0.5$ (Iredale); 8b, E.Indies, LV int., $\times 0.5$ (Chenu).
- T. (Semelanguilus)** IREDALE, 1924 [**T. tenuilirata* SOWERBY, 1867, in REEVE; OD]. Small, ovate,

somewhat cuneiform, anterior end longer; sculpture of concentric striae; ligament sunken, almost internal; hinge of RV with 2 laterals, wanting in LV. *Rec.*, Pac.—FIG. E106,10. **T. (S.) tenuilirata* SOWERBY, SE.Australia; LV ext., $\times 1$ (Reeve).

T. (Serratina) PALLARY, 1922 [**T. serrata* BROCCHI, 1814; OD] [= *Striotellina* THIELE, 1934 (obj.)]. Relatively high for length, ovate, sculpture concentric, radial ribs obsolete. *Mio.-Rec.*, Eu.-E.Atl.—FIG. E106,6. **T. (S.) serrata*

BROCCHI, Mio., France; 6a,b, RV int., ext., $\times 1$ (Cossmann & Peyrot).

T. (Sinuosipagia) COSSMANN, 1921 [**T. colpodes* BAYAN, 1873; OD]. Resembling *T. (Arcopagia)* but muscle impressions more unequal and lateral teeth more sharply truncate. *Eoc.*, Eu.—FIG. E108,6. **T. (S.) colpodes* BAYAN, France; 6a,b, RV ext., int., $\times 0.8$ (Cossmann).

T. (Tellinangulus) THIELE, 1934 [**T. aethiopica* THIELE & JAECKEL, 1931; M]. Small, asymmetrically rounded, somewhat rostrate; sculpture




FIG. E108. Tellinidae (Tellininae) (p. N620-N621).

- of few concentric lamellae, interspaces faintly striate radially; hinge with *All* and *PI* wanting, *3a* short. *Rec.*, E.Afr.—FIG. E108,1. **T. (T.) aethiopica* THIELE & JAECKEL, E.Afr.; LV ext., $\times 4$ (Thiele & Jaekel).
- T. (Tellinella)** MÖRCH, 1853 [**T. virgata* LINNÉ, 1758; SD STOLICZKA, 1870] [= *Capsa* LAMARCK, 1799 (non BRUGUIÈRE, 1797) (type, *T. angulata* LINNÉ, 1767; M); *Eutellina* FISCHER, 1887 (obj.); *Maoritellina* FINLAY, 1927 (type, *T. charlottae* SMITH, 1885; OD); *Tellinota* IREDALE, 1936 (type, *T. roseola*; OD)]. Elongate, pointed behind, posterior slope with 1 or 2 radial ribs; sculpture of fine concentric lamellae. *Oligo.-Rec.*, Eu.-W.Indies-Pac.—FIG. E108,10. **T. (T.) virgata* LINNÉ, *Rec.*, E.Indies; *10a,b*, RV int., ext., $\times 0.5$ (Habe).
- T. (Tellinidella)** HERTLEIN & STRONG, 1949 [**Tellinides purpureus* BRODERIP & SOWERBY, 1829; OD]. Resembling *T. (Eurytellina)* in form but with a thin shell; hinge with *PI* weak. *Rec.*, W.C.Am.—FIG. E108,12. **T. (T.) purpurea* (BRODERIP & SOWERBY), W.C.Am.; *12a,b*, RV int., ext., $\times 0.6$ (Gray).
- T. (Tellinides)** LAMARCK, 1818 [**Tellinides timorensis*; M]. Rectangular, smooth, posterior slope obliquely truncate, with a slight flexure. *Rec.*, E. Indies.—FIG. E108,9. **T. (T.) timorensis* (LAMARCK); *9a,b*, LV ext., RV int., $\times 1$ (Blainville).
- T. (Zearcopagia)** FINLAY, 1927 [**T. disculus* DESHAYES, 1855; OD]. Lenticular, concentrically grooved; hinge with *All* remote from cardinals; resembling *T. (Pseudarcopagia)* but without radial sculpture. *Rec.*, Pac.—FIG. E108,7. **T. (Z.) disculus* DESHAYES, Philip. Is.; RV ext., $\times 1$ (Reeve).
- Aenona** CONRAD, 1870 [**Tellina eufulensis* CONRAD, 1860; SD STOLICZKA, 1871]. Thin, fragile, compressed, nearly equilateral and smooth; dorsal margins of shell serving as lateral teeth; pallial sinus broad, not confluent. *U.Cret.*, N.Am.—FIG. E108,11. **A. eufulensis* (CONRAD), USA (Tenn.); *11a,b*, RV ext., RV int., $\times 2$ (Wade).
- Agnomyax** STEWART, 1930 [**Tellina monilifera* GABB, 1864; OD]. Resembling *Linearia* but with radial ribs on posterior slope only; posterior end diagonally truncate; hinge with cardinals directed ventrally, *3b* large and bifid. *U.Cret.*, W.N.Am.-?S.Asia-?Eu.—FIG. E108,5. **A. monilifer* (GABB), USA (Calif.); *5a*, RV hinge, $\times 5$ (Stewart); *5b*, RV ext., $\times 2$ (Gabb).
- Arcopagella** MEEK, 1871 [**A. mactroides*; M]. Resembling *Tellina (Arcopagia)* but with anterior and posterior teeth in both valves; surface smooth. *U.Cret.*, N.Am.—FIG. E108,2. **A. mactroides*, USA (Kans.); *2a,b*, RV int., LV int., $\times 1$ (Meek).
- Arcopagiopsis** COSSMANN, 1886 [**Tellina pustula* DESHAYES, 1825; SD SALISBURY, 1934]. Resembling *Tellina (Arcopagia)* but pallial sinus horizontal.
- Eoc.*, Eu.—FIG. E108,4. **A. pustula* (DESHAYES), M.Eoc., France; *4a,b*, RV ext., int., $\times 2$ (Cossmann).
- Barytellina** MARWICK, 1924 [**B. crassidens*; OD]. Shell thick; hinge with posterior laterals and *3b* strong. *Plio.*, N.Z.—FIG. E108,8. **B. crassidens*, N.Z.; *8a-c*, RV int., LV int., RV ext., $\times 1$ (Marwick).
- Hercodon** CONRAD in KERR, 1875 (issued separately, 1873) [**H. ellipticus*; M]. Elliptical, surface with irregular growth lines and numerous fine radials; hinge with *3a* angular, long, oblique, *2a* bifid, *4b* linear; pit for resilium deeply impressed; pallial sinus broad, rounded. *U.Cret.*, N.Am.—FIG. E108,3. **H. ellipticus*, USA (N.Car.); *3a-c*, LV int., RV ext., RV int., $\times 0.7$ (Stephenson, 1926).
- Linearia** CONRAD, 1860 [**L. metastrata*; M] [= *Oene* CONRAD, 1873 (type, *O. plana*; M); *Aene*, spelling error]. Ovate, beaks nearly central; sculpture of beaded radials, strongest at ends; hinge with 2 cardinals directed forward, dorsal margins of RV beveled to serve as laterals, with double grooves in LV. *L.Cret.-U.Cret.*, Eu.-N.Am.-Afr.
- L. (Linearia)**. Umbones rather high, inner ventral margin finely crenulate. *L.Cret.-U.Cret.*, N.Am.-Eu.-N.Afr.—FIG. E109,11. **L. (L.) metastrata* CONRAD, *U.Cret.*, USA (Tenn.); *11a,b*, RV ext., int., $\times 2.5$ (Wade).
- L. (Liothyris)** CONRAD in KERR, 1875 [?1873] [**L. (L.) carolinensis*; M]. Umbones lower and shell longer and smoother than in *L. (Linearia)*; ventral margin smooth; pallial sinus deep. *U.Cret.*, N.Am.—FIG. E109,7. **L. (L.) carolinensis*, USA (Tenn.); *7a-c*, RV int., LV ext., LV int., $\times 0.8$ (Wade).
- Nelltia** STEPHENSON, 1953 ("1952") [**N. stenzeli*; OD]. Resembling *Tellina* but not flexed, posterior end shorter; cardinal teeth weaker, anterior laterals distant, short, weak; pallial sinus relatively short. *U.Cret.*, N.Am.—FIG. E109,8. **N. stenzeli*, USA (Tex.); *8a-c*, RV ext., int., LV hinges, int., $\times 1$ (Stephenson).
- Palaeomoera** STOLICZKA, 1870 [**Tellina strigata* GOLDFUSS, 1840; M]. Like *T. (Phylloida)* in form; sculpture of fine, reticulate radial ribs; hinge with one lamellar anterior cardinal in each valve, bifid in RV, posterior cardinals not traceable in either valve; laterals less distinct. *U.Cret.*, Eu.-N.Afr.—FIG. E109,4. **P. strigata* (GOLDFUSS), Ger.; *4a-c*, RV ext., LV int., RV int., $\times 1$ (Holzapfel, 1889).
- Solyma** CONRAD, 1870 [**S. lineolatum*; M]. Ovate-quadrate, smooth; hinge in LV with one bifid cardinal, margins of shell serving as laterals., *U.Cret.*, N.Am.—FIG. E109,1. **S. lineolatum*; *1a,b*, USA (N.J.), RV ext., int., $\times 1$; *1c*, USA (Tex.), hinge, $\times 3$ (Whitfield).
- Strigilla** TURTON, 1822 [**Tellina carnaria* LINNÉ, 1758; SD GRAY, 1847] [= *Limicola* GRAY, 1852, ex LEACH MS (non KOCH, 1816) (obj.); *Strigillina*, spelling error, STOLICZKA, 1870]. Lenticular, sculp-

tured with oblique riblets; shells mostly tinged with pink. *Oligo.-Rec.*, Eu.-W.Indies-Afr.-C.Am.-S.Am.

S. (*Strigilla*). Oblique ribbing over most of shell, with one or more lines of flexure; pallial sinus somewhat discrepant. *Mio.-Rec.*, Eu.-W.Indies.—FIG. E109,5. **S.* (*S.*) *carnaria* (LINNÉ), *Rec.*, Carib.; LV ext., $\times 1$ (Woodward).

S. (*Aeretica*) DALL, 1900 [**Tellina senegalensis* HANLEY, 1844 (*non* GMELIN, 1791) = *S. polyaulax* TOMLIN & SHACKLEFORD, 1915; OD]. Oblique sculpture on anterior part of shell only, posterior slope smooth; pallial line connecting adductors in

one valve, falling short in the other. *Oligo. (Aquitana.)-Rec.*, Eu.-W.Afr.—FIG. E109,10.

**S.* (*A.*) *polyaulax* TOMLIN & SHACKLEFORD, *Rec.*, W.Afr.; RV ext., $\times 1.5$ (Nicklès).

S. (*Pisostrigilla*) OLSSON, 1961 [**Tellina pisiformis* LINNÉ, 1758; OD]. Small, oblique sculpture with a line of sharp flexure or zigzag bends along center of posterodorsal area. *Mio.-Rec.*, E.C.Am.-W.C.Am.—FIG. E109,6. **S.* (*P.*) *pisiformis* (LINNÉ), *Rec.*, USA (Fla.); LV ext., $\times 2$ (Stanford Univ. specimen).

S. (*Rombergia*) DALL, 1900 [**S. rombergii* MÖRCH, 1853; OD]. Sculpture as in *S.* (*Strigilla*)




FIG. E109. Tellinidae (Tellininae) (p. N621-N623).

- but pallial sinus alike in both valves, falling short of uniting adductors. *Rec.*, W.Indies.
- S. (Simplistrigilla)** OLSSON, 1961 [**S. (S.) strata* (= **Strigilla serrata* MÖRCH, 1860); OD]. Oblique striations without any lines of flexure. *Rec.*, W.C. Am.-S. Am.—FIG. E109,3. **S. (S.) serrata*, Ecuador; LV ext., $\times 3$ (Olsson).
- Tellidora** H. ADAMS & A. ADAMS, 1856 [**Tellina burneti* BRODERIP & SOWERBY, 1829; SD STOLICZKA, 1870 [= *Tellinodora* PAETEL, 1875 (spelling error)]. Compressed, inequilateral, ovate-trigonal, surface with spaced concentric lamellae ending in marginal serrations. *Plio.-Rec.*, E.N. Am.-W.C. Am.—FIG. E109,12. **T. burneti* (BRODERIP & SOWERBY), *Rec.*, W.C. Am.; 12a,b, RV ext., LV int., $\times 1$ (Chenu).
- Tellinimera** CONRAD, 1860 [**T. eborea*; SD GARDNER, 1916] [= *Tellimera* CONRAD, 1870, *emend.* (*nom. van.*)]. More compressed and inequilateral than *Aenona*, surface with sharp concentric striations; cardinal plate channeled posteriorly, with a small pit anteriorly; hinge with 2a V-shaped, 4b bifid. *U. Cret.*, E.N. Am.—FIG. E109,9. **T. eborea*, USA (Md.); 9a,b, RV ext., both valves dorsal, $\times 2$ (Gardner).
- Telliura** OLSSON, 1944 [**Tellidora (T.) peruana*; OD]. Resembling *Tellidora* in serrate dorsal margin and hinge but outline more equilateral, with close-spaced concentric riblets or undulations. *U. Cret.*, W.S. Am.—FIG. E109,2. **T. peruana* (OLSSON), Peru; RV ext., $\times 1$ (Olsson).
- Subfamily MACOMINAE Olsson, 1961**
- Sculpture generally more subdued than in Tellininae; hinge without lateral teeth. *Eoc.-Rec.*
- Macoma** LEACH, 1819 [**M. tenera* (= *Tellina calcarea* GMELIN, 1791); M] [= *Macroma*, *Macrotoma*, spelling errors; *Limecola* BROWN, 1844 (*ex* LEACH MS) (in synonymy of *Tellina solidula* PULTENEY, 1799 = *T. balthica rubra* DA COSTA, 1778); *Macomopsis* SACCO, 1901 (type, *Tellina elliptica* BROCCHI, 1814; OD); ?*Pulvinus* SCARLATO, 1965 (type, *Tellina micans* HANLEY, 1844; OD)]. Inequilateral, mostly thin, with deciduous periostracum; posterior end slightly twisted; pallial sinus tending to be discrepant, larger in one valve. *Eoc.-Rec.*, cosmop.
- M. (Macoma)**. Ovate-trigonal, shell texture somewhat chalky. *Mio.-Rec.*, N. Eu.-N. Am.-NE. Asia.—FIG. E110,4. **M. (M.) calcarea* (GMELIN), *Rec.*, Arctic; 4a, RV ext., $\times 0.8$ (Sars); 4b, RV int., $\times 0.5$; 4c, both valves, $\times 0.8$ (Davies).
- M. (Austromacoma)** OLSSON, 1961 [**Solen constrictus* BRUGUIÈRE, 1792; OD]. Texture not chalky; pallial sinus large, high, pointed under beak, connected to anterior adductor scar at its lower end. *Plio.-Rec.*, E.N. Am.—FIG. E110,7. **M. (A.) constricta* (BRUGUIÈRE), *Rec.*, Carib.; 7a,b, LV ext., int., $\times 1$ (Philippi).
- M. (Bendemacoma)** EAMES, 1957 [**Peronaea nigeriensis* NEWTON, 1922; OD]. Large, high, solid, oblique; hinge with 3a grooved; pallial sinus tongue-shaped, narrow, confluent for half its length. *Eoc.*, Afr.—FIG. E110,11. **M. (B.) nigeriensis* (NEWTON), Nigeria; 11a-c, LV ext., int., RV int., $\times 0.5$ (Newton).
- M. (Cymatoica)** DALL, 1890 [**M. (C.) occidentalis* = *Tellina undulata* HANLEY, 1844; SD DALL, 1900]. Small, elongate, with undulating oblique sculpture. *Mio.-Rec.*, E.C. Am.-W.C. Am.—FIG. E110,5. **M. (C.) undulata* (HANLEY), *Rec.*, W. Mex.; LV ext., $\times 2$ (Dall).
- M. (Macoploma)** PILSBRY & OLSSON, 1941 [**M. (M.) ecuadoriana*; OD]. Area of posterior slope near dorsal margin granulated. *Plio.-Rec.*, W.C. Am.-S. Am.—FIG. E110,6. **M. (M.) ecuadoriana*, *Plio.*, Ecuador; LV ext., $\times 0.6$ (Pilsbry & Olsson).
- M. (Panacoma)** OLSSON, 1942 [**M. (P.) chiriquiensis*; OD]. Sculpture of a few small granules and strong raised concentric threads separated by flat interspaces; pallial sinus wide, confluent below. *Mio.-Plio.*, W.C. Am.—FIG. E110,8. **M. (P.) chiriquiensis*, W. Panama; 8a,b, LV ext., RV ext., $\times 1$ (Olsson).
- M. (Pinguimacoma)** IREDALE, 1936 [**Pinguimacoma hemicilla*; OD]. Small, thin, resembling *Tellina (Pinguitellina)* in form but without laterals. *Rec.*, Australia.—FIG. E110,9. **M. (P.) hemicilla* (IREDALE), E. Australia; 9a,b, RV ext., hinge, $\times 3$ (Iredale).
- M. (Psammacoma)** DALL, 1900 [**Psammotaea candida* LAMARCK, 1818; OD]. Elongate, smooth, posterior flexure obsolete, periostracum delicate; ligament and resilium wholly external; pallial sinus only half confluent. *Mio.-Rec.*, E.N. Am.-Pac.—FIG. E110,12. **M. (P.) candida* (LAMARCK), *Rec.*, E. Indies; 12a,b, LV ext., int., $\times 0.5$ (Philippi).
- M. (Rexithaerus)** TRYON, 1869 *ex* CONRAD MS [**Tellina secta* CONRAD, 1837; SD DALL, 1900]. Large, compressed, nonchalky; ligament strong, deep-set, dorsal margin produced upward behind it. *Mio.-Rec.*, W.N. Am.—FIG. E110,10. **M. (R.) secta* (CONRAD), *Rec.*, USA (Calif.); RV ext., $\times 0.5$ (Reeve).
- M. (Rostrimacoma)** SALISBURY, 1934 [**Panopea cancellata* SOWERBY *in* REEVE, 1873; OD]. Large, anterior end produced; surface cancellate. *Rec.*, E. Atl.—FIG. E110,3. **M. (R.) cancellata* (SOWERBY), W. Afr.; RV ext., $\times 0.3$ (Salisbury).
- M. (Salmacoma)** IREDALE, 1929 [**Salmacoma vappa*; OD]. Ovate, inequivalve, posterior flexure pronounced. *Rec.*, S. Pac.—FIG. E110,2. **M. (S.) vappa* (IREDALE), *Rec.*, NE. Australia; RV ext., $\times 1$ (Iredale).

M. (Scissulina) DALL, 1924 [**Tellina dispar* CONRAD, 1837; OD]. With fine oblique sculpture on one valve. *Rec.*, Pac.—FIG. E110,1. **M. (S.) dispar* (CONRAD), USA (Hawaii); 1a, RV ext., ×1 (Sowerby); 1b,c, detail of posterior end (1c, LV; 1b, RV), ×6 (Dall, Bartsch, Rehder).

M. (Tenuconcha) DALL, 1921 [**Psammacoma (T.) brasiliensis*; OD] [= *Psammothalia* OLSSON, 1961 (type, *Tellina cognata* C. B. ADAMS, 1852; OD)]. Resembling *M. (Scissulina)* but more quadrate, with oblique sculpture on both valves. *Rec.*, W.C.Am.-E.S.Am.—FIG. E111,8. **M.*




FIG. E110. Tellinidae (Macominae) (p. N623-N624).




FIG. E111. Tellinidae (Macominae) (p. N624-N626).

(*T.*) *brasiliana* (DALL), Brazil; 8a,b, RV ext., int., $\times 1$ (Stanford spec.).—FIG. E106,14. *M.* (*T.*) *cognata* (C. B. ADAMS), Peru; 14a,d, LV ext., int., $\times 1$; 14b,c, LV and RV hinges, enl. (688).

Apolymetis SALISBURY, 1929 [pro *Polymetis* SALISBURY, 1929 (non WALSINGHAM, 1908), pro *Metis* H. ADAMS & A. ADAMS, 1856 (non PHILIPPI, 1843)] [**Tellina meyeri* PHILIPPI, 1846, ex DUNKER MS; M]. Subcircular, compressed, flexure and median furrow pronounced, ligament long, embedded; sculpture of spaced concentric ribs. *Mio.-Rec.*, E.Indies.—FIG. E111,3. **A. meyeri* (PHILIPPI), *Rec.*, E.Indies; 3a-c, RV ext., both valves dorsal, LV int., $\times 0.5$ (Philippi, 1846).
Bartrumia MARWICK, 1934 [**Raeta tenuiplicata*

BARTRUM, 1919; OD]. Shorter than *Macoma* (*Rostrimacoma*), more inflated; ligament sunken as in *Psammotreta*. *Mio.*, S.Pac.—FIG. E111,5. **B. tenuiplicata* (BARTRUM), L.Mio., N.Z.; 5a, LV ext., $\times 1$; 5b, hinge, enl. (Marwick, 1934).
Exotica LAMY, 1918 (ex JOUSSEAUME MS) [**E. exotica* (?=*Tellina triradiata* A. ADAMS, 1870); T]. Like *T.* (*Moerella*) in form but lacking lateral teeth; sculpture tending to be oblique, fine. *Rec.*, Pac.-Ind.O.-E.Indies.

E. (*Exotica*). Surface smooth or with concentric striae only. *Rec.*, Ind.O.—FIG. E112,1. **E.* (*E.*) *triradiata* (A. ADAMS), Red Sea; RV ext., $\times 2$ (Adams, 1870).

?*E.* (*Jactellina*) IREDALE, 1929 [**Tellina obliquaria* DESHAYES, 1855; OD]. Sculpture of fine oblique




FIG. E112. Tellinidae (Macominae) (p. N625-N628).

ribs in both valves. *Rec.*, E.Indies.—FIG. E111, 1. **E. (I.) obliquaria* (DESHAYES), E.Indies; 1a,b, RV ext., LV ext., $\times 1.5$ (Reeve, 1868).

?*E. (Loxoglypta)* DALL, BARTSCH, & REHDER, 1939 [**Tellina obliquilineata* CONRAD, 1837; OD]. Oblique sculpture replaced at anterior end of shell by concentric grooves. *Rec.*, Pac.—FIG. E111, 4. **E. (L.) obliquilineata* (CONRAD), USA (Hawaii); detail of sculpture, $\times 12$ (Dall, Bartsch, & Rehder, 1939).

Florimetis OLSSON & HARBISON, 1953 [**Tellina in-tastriata* SAY, 1826; OD] [= *Apolymetis* AUCT. (*partim*, see *Psammotreta*)]. Quadrate, posteriorly with strong flexure, central slope with broad median furrow; sculpture of incremental lines; ligament deep; pallial sinus large. *U.Mio.-Rec.*, W.Atl.-E.Pac.-E.Indies.—FIG. E111, 7. **F. in-*

tastriata (SAY), *Rec.*, USA (Fla.); 7a,b, RV ext., LV int., $\times 0.5$; 7c, hinge, $\times 1.7$ (689).

Gastrana SCHUMACHER, 1817 [**G. donacina* = *Tellina abildgaardiana* SPENGLER, 1798 (= *T. matadoa* GMELIN, 1791); SD BUCQUOY, DAUTZENBERG, & DOLLFUS, 1898] [= *Diodonta* DESHAYES, 1846 (*non* HARTMAN, 1843); *Fragilia* DESHAYES, 1848 (type, *Tellina fragilis* LINNÉ, 1758; M)]. Oblique, thin, surface with irregular concentric lamellae; hinge with 3b bifid; pallial sinus rounded, deep, not confluent. *Eoc.-Rec.*, Eu.-W.Afr.—FIG. E111, 6. **G. matadoa* (GMELIN), *Rec.*, W.Afr.; 6a, LV ext., $\times 1$ (Fischer, 1942); 6b,c, LV ext., RV int., $\times 1$ (Nicklès, 1950).

Heteromacoma HABE, 1952 [**Tellina irus* HANLEY, 1845; OD] [= *Gastrana* AUCT. (*partim*)]. Ovate-trigonal, RV with lunular area, LV with corresponding emargination; ligament long, in sunken groove; adductor scars unequal; pallial sinus large, angular, slightly discrepant, about three-fourths confluent. [Type species has been confused by authors with *Macoma inquinata* (DESHAYES, 1854) from W.N.Am. and "*Gastrana*" *yantaiensis* (CROSSE & DEBEAUX, 1863) from China; see also *Sinomacoma*.] *Rec.*, W.Pac.—FIG. E111, 9. **H. irus* (HANLEY), Japan; 9a,b, LV ext., RV int., $\times 0.5$ (Habe, 1952).

Leporimetis IREDALE, 1930 [**Tellina spectabilis* HANLEY, 1844; OD]. Oblique-trigonal, posterior strongly flexed, incremental sculpture roughened; pallial sinus nearly meeting anterior adductor scar. *Mio.-Rec.*, W.Pac.—FIG. E111, 10. **L. spectabilis* (HANLEY), *Rec.*, Japan; 10a,b, RV ext., RV int., $\times 0.5$ (Habe, 1952).

Macalia H. ADAMS, 1860 [**Tellina bruguieri* HANLEY, 1844; M] [= *Tellinungula* RÖMER, 1873 (obj.; M); *Macalina* HABE, 1952 (*nom. null.*)]. Trigonal, smooth, hinge with markedly large cardinal teeth; adductor scars unequal. *Rec.*, W.Pac.—FIG. E112, 5. **M. bruguieri* (HANLEY), Japan; 5a-c, RV hinge, RV ext., LV int., $\times 1$ (Habe, 1952).

Psammotreta DALL, 1900 [**Tellina aurora* HANLEY, 1844; OD] [= *Cydippina* DALL, 1900 (type, *T. brevifrons* SAY, 1834; OD); *Scrobiculina* DALL, 1900 (type, *Scrobicularia viridotincta* CARPENTER, 1856; OD); *Schumacheria* COSSMANN, 1902 (*nom. van. pro Scrobiculina*); *Apolymetis* AUCT. (*partim*, see *Florimetis*)]. More elongate than *Macoma*, ligament and resilium sunken, almost internal, adductor scars unequal; pallial sinus partly confluent. *Mio.-Rec.*, C.Am.-S.Am.-W.Pac.

P. (Psammotreta). Shorter than *Macoma* (*Psammacoma*); anterior end not inflated. *Rec.*, E.C.Am.-W.C.Am.-S.Am.—FIG. E111, 2. **P. (P.) aurora* (HANLEY), W.C.Am.; 2a,b, RV ext., LV ext., $\times 1$ (822).

P. (Ardeamya) OLSSON, 1961 [**Tellina columbien-sis* HANLEY, 1844; OD]. Elongate-elliptical, beaks small, narrow, rising above dorsal margin. *Rec.*,

W.C.Am.-S.Am.—FIG. E112,4. **P. (A.) columbiensis* (HANLEY), Peru; RV ext., $\times 0.5$ (688).

P. (Pseudometis) LAMY, 1918 [**Tellina truncata* PHILIPPI, 1843 (*non* LINNÉ, 1767 = *T. praerupta* SALISBURY, 1934); SD SALISBURY, 1934]. Resembling *P. (Psammotreta)* but higher, anterior end more inflated; surface smooth; posterior flexure weak; ligament less immersed than in *Florimetis*; pallial sinus about one-third confluent.

Mio.-Rec., E. C. Am.-W. C. Am.-W. Pac.—FIG. E112,3. **P. (P.) praerupta* (SALISBURY), Rec., E. Indies; 3a-c, LV ext., int., RV hinge, $\times 0.5$ (Philippi, 1843).

P. (Tellinimactra) LAMY, 1918 [**Tellina edentula* SPENGLER, 1798; OD]. Thin, more ovate and less strongly flexed than *Florimetis*, lacking central depressed area; ligament and resilium large; pallial sinus larger than in *P. (Psammotreta)*. Rec., W. Pac.—FIG. E112,2. **P. (T.) edentula*




FIG. E113. Donacidae (p. N628).

(SPENGLER), Japan; 2a-c, LV ext., int., hinge, $\times 0.7$ (Habe, 1952).

Sinomacoma YAMAMOTO & HABE, 1950 [**Fragilia yantaiensis* CROSSE & DEBEAUX, 1863; OD]. Trigonal, heavy shelled, ligament in long groove; area in front of beaks beveled, wide; posterior end not twisted. *Rec.*, E.Asia.—FIG. E112,6. **S. yantaiensis* (CROSSE & DEBEAUX), NE.China; 6a,b, LV ext., RV int., $\times 0.5$ (Crosse & Debeaux, 1863).

Family DONACIDAE Fleming, 1828

Trigonal shells, medium-sized to small, solid, inequilateral, opisthogyrate; hinge with two cardinals and well-developed laterals; pallial sinus normally present. *U. Cret.-Rec.*

Donax LINNÉ, 1758 [**D. rugosa*; SD SCHUMACHER, 1871] [= *Donaciaris* DUMÉRIL, 1806 (obj.); *Donacina* FÉRUSAC, 1822 (*nom. nud.*, in *synon.*)]. Radial sculpture present in most, at least as marginal crenulations; periostracum wanting. *L.Eoc.-Rec.*, cosmop.

D. (Donax). Sturdy, with radial sculpture, posterior area concentrically wrinkled; internal margin strongly denticulate. *Rec.*, circumtrop.—FIG. E113,1. **D. (D.) rugosus* LINNÉ, W.Afr.; 1a,b, LV ext., RV int., $\times 1$ (124b).

D. (Capsella) GRAY, 1851 [**Tellina polita* POLI, 1795 (= **T. variegata* GMELIN, 1791); M] [= *Peronaeoderma* MÖRCH, 1853 (*non* POLI, 1795) (obj.; SD SALISBURY, 1934)]. Elongate-ovate, polished, without radial sculpture, internal margin smooth. *Rec.*, Eu.-W. N. Am.—FIG. E113,8. **D. (C.) variegatus* (GMELIN), *Medit.*; 8a-c, LV ext., LV int., RV int., $\times 1$ (89a).

D. (Chion) SCOPOLI, 1777 [**D. denticulata* LINNÉ, 1758; M]. Solid, with punctate radial sculpture, posterior slope truncate, roughened by scaly sculpture. *L.Eoc.-Rec.*, Eu.-Carib.-E. Asia-S. Pac.—FIG. E113,3. **D. (C.) denticulatus* LINNÉ, *Rec.*, W.Indies; 3a,b, LV ext., RV int., $\times 1$ (Woodward; Chenu).

D. (Cuneus) DA COSTA, 1778 [**Cuneus vittatus*; SD WINCKWORTH, 1926] [= *Cunerus*, spelling error]. Radiately ribbed, with some oblique striae crossing central slope; posterior truncation weak. *Mio.-Rec.*, Eu.—FIG. E113,9. **D. (C.) vittatus* (DA COSTA), *Rec.*, Eu.; LV ext., $\times 1$ (124b).

D. (Deltachion) IREDALE, 1930 [**D. virilis*; OD]. Small, high, posterior side truncate; pallial sinus 0.75 length of shell. *Rec.*, S.Pac.—FIG. E113,12. **D. (D.) virilis* IREDALE, E.Australia; 12a,b, RV ext., int., $\times 2$ (Iredale, 1930).

D. (Grammatodonax) DALL, 1900 [**D. madagascariensis* WOOD, 1828 ("Lamarck" by error); OD]. Short, compressed, surface furrowed with oblique ribs; RV with 1 cardinal, LV with 2.

Rec., Afr.—FIG. E113,5. **D. (G.) madagascariensis* WOOD, S.Afr.; LV ext., $\times 1$ (124b).

D. (Hecuba) SCHUMACHER, 1817 [**Venus scortum* LINNÉ, 1758; SD HERRMANNSEN, 1847]. Large, with conspicuous posterior carina; sharp groove in dorsal margin of RV in front of *Al. Rec.*, E. Indies.—FIG. E113,11. **D. (H.) scortum* (LINNÉ), E.Indies; 11a-c, LV ext., $\times 0.5$; LV and RV hinges, enl. (124b).

D. (Latona) SCHUMACHER, 1817 [**D. cuneatus* LINNÉ, 1758; M]. Radials reduced to fine threadlike striae except on posterior slope, which may be rugose; interior margin smooth or with minute radial wrinkles. *Rec.*, E.Indies.—FIG. E113,10. **D. (L.) cuneatus* LINNÉ, E.Indies; 10a,b, RV ext., LV int., $\times 1$ (124b).

D. (Machacrodonax) ROEMER, 1870 [**D. scalpellum* GRAY, 1823; SD DALL, 1900] [= *?Platydona* DALL, 1900 ("*D. finchii* SOWERBY," *nom. inquirend.*)]. Elongate, posterior end short, set off by ridge; shell slightly gaping at both ends. *Mio.-Rec.*, E. C. Am.-W. C. Am.-IndoPac.—FIG. E113,4. **D. (M.) scalpellum* GRAY, *Rec.*, Red Sea; LV ext., $\times 1$ (Reeve, 1854).

D. (Paradonax) COSSMANN & PEYROT, 1910 [**D. transversus* DESHAYES, 1830; OD]. Nearly smooth, radials on central slope only; smaller than *D. (Donax)*, with lower beaks; posterior laterals in RV strong. *Oligo.-Rec.*, Eu.-Carib.—FIG. E113,2. **D. (P.) transversus* DESHAYES, *Mio.*, France; 2a,b, LV and RV hinges, enl.; 2c-e, RV ext., LV int., RV int., $\times 1$ (Cossmann & Peyrot, 1911).


D. (Plebidonax) IREDALE, 1930 [**D. deltoides* LAMARCK, 1818; OD]. Relatively large, nearly smooth, posterior end set off by a weak carina; 1 massive bifid cardinal in RV, 2 in LV, anterior laterals wanting; pallial sinus half the length of shell. *Rec.*, S.Pac.—FIG. E113,7. **D. (P.) deltoides* LAMARCK, Australia; 7a,b, LV int., RV int., $\times 0.5$ (Sowerby).

D. (Serrula) MÖRCH, 1853 [**D. trunculus* LINNÉ, 1758; SD STOLICZKA, 1870]. Elongate, smooth, without posterior carination; posterior margin smooth within; RV without lateral teeth. *Plio.-Rec.*, Eu.—FIG. E113,6. **D. (S.) trunculus* LINNÉ, *Rec.*, *Medit.*; LV ext., $\times 1$ (124b).

D. (Tentidonax) IREDALE, 1930 [**D. veruinus* HEDLEY, 1913; OD]. Small, elongate, sides slanting, smooth except for wrinkle-ridged posterior area. *Rec.*, Australia.

Egerella STOLICZKA, 1870 [*pro Egeria* LEA, 1833 (*non* ROISSY, 1805)] [**Egeria subtrigonia* LEA, 1833; SD STOLICZKA, 1871]. Thin, small, variable, radial sculpture faint, interior margins finely serrate; lateral teeth wanting. *Paleoc.-Eoc.*, Eu.-E.N.Am.—FIG. E114,1. **E. subtrigonia* (LEA), *Eoc.*, USA (Ala.); 1a-c, LV ext., int., RV int., $\times 3$ (Harris, 1919).


Galatea BRUGUIÈRE, 1797 (Genus without species) [**Galathea radiata* LAMARCK, 1805 (= **Venus*



Veneroida—Tellinacea

paradoxa BORN, 1778); SM LAMARCK, 1805] [=*Egeria* ROISSY, 1805 (*pro Galatea*) (*nom. van.*); *Potamophila* SOWERBY, 1821 (*obj.*); *Megadesma* BOWDICH, 1822 (*obj.*); *Galateola* FLEMING, 1828 (*nom. nud.*); *Galataea*, *Galathea*, *Galateia*, spelling

errors]. Large, thick, trigonal, with a smooth, heavy periostracum; hinge plate massive; inner ventral margin smooth; pallial sinus small. *Rec.*, W.Afr.—FIG. E114,3. **G. paradoxa* (BORN); RV int., $\times 1$ (Nicklès, 1950).



Hemidonax MÖRCH, 1870 [**Donax pictus* TRYON, 1870 (= **Cardium donaciforme* BRUGUIÈRE, 1792); M] [= *Donacocardium* VON VEST, 1875 (*obj.*); OD]. Sculpture of smooth but elevated radial ribs, inner margin coarsely crenate; hinge somewhat as in *Donax* but teeth larger. *Rec.*, E. Indies.—FIG. E114,4. **H. donaciformis* (BRUGUIÈRE); LV ext., $\times 1$ (Reeve, 1844).

Iphigenia SCHUMACHER, 1817 [**Donax laevigata* GMELIN, 1791; M] [= *Fischeria* BERNARDI, 1860 (*non* ROBINEAU-DESVOIDY, 1830; *Finlaria*, spelling error); *Profischeria* DALL, 1903 (*pro Fischeria*; type, *F. delessertii* BERNARDI, 1860; OD); *Iphigenia*, *Ephigenia*, spelling errors]. Thin, trigonal to ovate, with a periostracum, smooth; LV with 2 subequal cardinals, no laterals, RV with 1 weak and 3b bifid, laterals 2, elongate and weak; pallial sinus large. *Mio.-Rec.*, W.Afr.-E.C.Am.-W.C.Am.-S.Am.—FIG. E114,2. **I. laevigata* (GMELIN), *Rec.*, W.Afr.; 2a,b, LV ext., RV int., $\times 1$ (Nicklès, 1950).

?*Macrodonax* OLSSON, 1944 [**M. peruviana*; OD]. Large, solid, sculpture discrepant, posterior area with nodose radial ribs, remainder of shell concentrically ribbed; hinge heavy, with anterior and posterior laterals; pallial area unknown. *U.Cret.* (*Senon.*), S.Am.—FIG. E114,5. **M. peruviana*, Peru; 5a,b, RV ext., int., $\times 0.25$ (Olsson, 1944).

Notodonax FERUGLIO, 1936 [**Donax (N.) annae-eugeniae*; OD]. Donaciform, solid, with a posterior-dorsal angulation; margin smooth within; hinge with 2 cardinals and 2 laterals; pallial sinus small. *U.Cret.*, S.Am.-N.Am.-SW.Asia.

N. (*Notodonax*). Surface with a few concentric furrows, lateral teeth relatively short; pallial sinus wavy to angular. *U.Cret.*, S.Am.—FIG. E114,7. **N. (N.) annae-eugeniae* (FERUGLIO), Patag.; 7a,b, LV ext., RV int., $\times 1$ (Feruglio, 1935).

N. (*Protodonax*) VOKES, 1945 [**Protodonax elongatus*; OD]. Surface smooth, lateral teeth somewhat longer than in *N. (Notodonax)*, pallial sinus angular. *U.Cret.*, N.Am.-SW.Asia.—FIG. E114,6. **N. (P.) elongatus* (VOKES), USA (Wyo.); 6a, LV hinge, $\times 1$; 6b, RV hinge, $\times 2$ (1944).

Family PSAMMOBIIDAE Fleming, 1828

[*nom. correct.* FISCHER, 1887 (*pro Psammobiidae* FLEMING, 1828)] [= *Garidae* STOLICZKA, 1871 (*nom. transl.* STEWART, 1930, *ex* *Garinae* STOLICZKA, 1871); *Asaphidae* WINCKWORTH, 1932 (*non* BURMEISTER, 1843)]

Shells inequilateral, slightly gaping in most forms, especially at posterior end; hinge with one to three cardinal teeth, lat-

FIG. E114. Donacidae (p. N628-N629).

erals weak to wanting; ligament on nymph; pallial sinus present. *U.Cret.-Rec.*

Ovate-trapezoidal, gape small or absent. *U.Cret.-Rec.*

Subfamily PSAMMOBIINAE Fleming, 1828

[*nom. transl.* E. A. SMITH, 1885 (*ex* Psammobiidae FLEMING, 1828)]

Gari SCHUMACHER, 1817 [**G. vulgaris* (= *Solen amethystus* WOOD, 1815) (ICZN pend.)] [= *Psammotaea* LAMARCK, 1818 (type, *P. donacina*; SD




FIG. E115. Psammobiidae (Psammobiinae) (p. N630-N631).

- CHILDREN, 1823); *Capsella* DESHAYES, 1855 (non GRAY, 1851); *Milligaretta* IREDALE, 1936 (type, *M. venta*; OD)]. Elongate-ovate to quadrate, smooth to strongly sculptured, posterior end wider than anterior. *Eoc.-Rec.*, cosmop.
- G. (Gari).** Posterior slope weakly set off; pedal muscle scars 1, oval, in front of cardinal teeth; sculpture of concentric growth lamellae, overlain by oblique striae in some. *Mio.-Rec.*, Eu.-Asia. —FIG. E115,7. *G. (*G.*) *amethystus* (WOOD), *Rec.*, E.Indies; LV ext., $\times 1$ (Wood, 1815).
- G. (Amphipsammus)** COSSMANN, 1914 [*Psammobia lamarckii* DESHAYES, 1857; M]. Posterior slope with radial riblets, shell otherwise smooth; lateral lamellae wanting; sinus free, not confluent. *Eoc.-Oligo.*, Eu.—FIG. E115,11. *G. (*A.*) *lamarckii* (DESHAYES), *Eoc.*, France; 11a,b, LV ext., int., $\times 0.5$ (Deshayes, 1825).
- G. (Azor)** J. SOWERBY, ex LEACH MS, 1824 [*Sanguinolaria compressa*; M]. Small, smooth, rectangular, pallial sinus rounded, pallial line not confluent. *Eoc.*, Eu.—FIG. E115,10. *G. (*A.*) *compressa* (SOWERBY), Eng.; 10a-c, RV ext., int., LV int., $\times 1$ (Brit. Mus. Guide, 1959).
- G. (Dysmea)** DALL, BARTSCH, & REHDER, 1939 [*Solen occidentis* GMELIN, 1791; OD]. Smooth except for concentric riblets on flaring posterior end; pallial sinus confluent for half its length. *Rec.*, IndoPac.—FIG. E115,6. *G. (*D.*) *occidens* (GMELIN), E.Indies; 6a,b, RV ext., LV int., $\times 0.3$ (Chemnitz, 1782).
- G. (Garum)** DALL, 1898 [*Psammobia filosa* CONRAD, 1853; M]. Sculpture of fine concentric grooves; pallial sinus short. *Eoc.*, N.Am.-Eu.—FIG. E115,13. *G. (*G.*) *filosa* (CONRAD), M.Eoc., USA (Ala.); 13a,b, RV ext., int., $\times 1$ (Stanford Univ. specimen).
- G. (Gobraeus)** BROWN, 1844 (in synonym.), ex LEACH, MS [*Solen vespertinus* GMELIN, 1791 (= *Tellina depressa* PENNANT, 1777); M] [= *Psammocola* DE BLAINVILLE, 1825, AUCTT. (non DE BLAINVILLE, 1824) (obj.)]. More gaping than *G. (Gari)*, nearly smooth, concentric sculpture of growth lines only, posterior slope with superficial radial striae; 2 subequal cardinals in RV, anterior cardinal bifid in LV, posterior weak; pallial sinus large, partly confluent. *Eoc.-Rec.*, Eu.-N.Am.-S.Am.—FIG. E115,8. *G. (*G.*) *depressa* (PENNANT), *Rec.*, Medit.; LV ext., $\times 0.6$ (89a).
- G. (Grammatomya)** DALL, 1898 [*Psammobia squamosa* LAMARCK, 1818; M]. Entire surface obliquely grooved, coarser on dorsal posterior area. *Rec.*, IndoPac.—FIG. E115,5. *G. (*G.*) *squamosa* (LAMARCK), Borneo; LV ext., $\times 1$ (Woodward, 1854).
- G. (Kermadysmea)** POWELL, 1958 [*K. galathea*; OD]. Resembling *G. (Dysmea)* but with concentric ribs, granular. *Rec.*, S.Pac.—FIG. E115, 12. *G. (*K.*) *galathea* (POWELL), Kermadec Is.; 12a,b, LV ext., RV int., $\times 0.7$; 12c, sculpture, enl.; 12d,e, RV and LV hinges, enl. (Powell, 1958).
- G. (Psammobella)** GRAY, 1851 [*Psammobia costulata* TURTON, 1822; SD KOBELT, 1881]. Small, smooth except for radial ribs on posterior slope; hinge feeble, pallial sinus deep, partly confluent. *Rec.*, Eu.—FIG. E115,3. *G. (*P.*) *costulata* (TURTON), Eng.; RV ext., $\times 1$ (Forbes & Hanley, 1848).
- G. (Psammobia)** LAMARCK, 1818 [*Tellina fervensis* GMELIN, 1791; SD CHILDREN, 1822, as *T. ferroensis*] [= *Haplomochlia* GISTEL, 1848 (nom. van.) (obj.)]. Posterior slope with strong ridge; pedal muscle scars 2 to 3, below cardinal teeth. *Rec.*, Eu.—FIG. E115,9. *G. (*P.*) *fervensis* (GMELIN), Eng.; 9a,b, RV ext., LV int., $\times 1$ (Wood, 1815).
- G. (Psammodonax)** COSSMANN, 1887 [*Psammobia caillati* DESHAYES, 1857; OD]. Inequilateral, posterior end short, posterior slope radially striate; hinge without laterals; pallial sinus large, passing beaks, oval, only partly confluent. *L.Eoc.-U.Eoc.*, Eu.—FIG. E115,4. *G. (*P.*) *caillati* (DESHAYES), M.Eoc., France; LV int., $\times 1$ (Deshayes, 1864).
- G. (Psammoica)** DALL, 1900 [*Psammobia appendiculata* DESHAYES, err. pro *Solen appendiculata* LAMARCK, 1806; OD] [= *Macropsammus* COSSMANN, 1902 (nom. van.)]. Small, smooth, compressed, truncate behind; ligamental nymph prominent; pallial sinus long, not confluent. *Eoc.*, Eu.—FIG. E115,2. *G. (*P.*) *appendiculata* (LAMARCK), M.Eoc., France; 2a,b, RV ext., int., $\times 1$ (Deshayes, 1825).
- G. (Psammotaena)** DALL, 1900 [*Psammobia effusa* LAMARCK (= *Solen effusa* LAMARCK, 1806); OD]. Resembling *G. (Amphipsammus)* but with posterior slope smooth. *Eoc.*, Eu.—FIG. E115,1. *G. (*P.*) *effusa* (LAMARCK), M.Eoc., France; 1a,b, LV ext., int., $\times 0.7$ (Deshayes, 1864).
- Amphichaena** PHILIPPI, 1847 [*A. kindermanni*; M] [= *Amphidona* MÖRCH, 1858 (nom. null.)]. Subcylindrical, inner margin arcuate at anterior end, smooth or nearly so posteriorly; pallial sinus short, not reaching mid-line. *Pleist.-Rec.*, W.C.Am.—FIG. E116,11. **A. kindermanni*, *Rec.*, W. Mexico; 11a-c, LV int., RV int., LV ext., $\times 1$ (R. Palmer & L. Hertlein, 1926).
- Asaphinella** COSSMANN, 1886 [*Capsa minima* DESHAYES, 1857; OD]. Outline as in *Gari* but less inequilateral, adult shells small. *Eoc.*, Eu.
- A. (Asaphinella).** Oblong, smooth. *Eoc.*, Eu.—FIG. E116,1. **A. (A.) minima* (DESHAYES), France; 1a-c, LV ext., LV int., RV hinge, $\times 5$ (Deshayes, 1857).
- A. (Herouvalia)** COSSMANN, 1891 [*A. semitexta* COSSMANN, 1886; OD]. Minute, sculptured strongly reticulate, especially on posterior slope. *Eoc.*, Eu.—FIG. E116,2. **A. (H.) semitexta*




FIG. E116. Psammobiidae (Psammobiinae) (p. N630-N631, N633).

- COSSMANN, France; 2a,b, LV int., ext., $\times 4$; 2c, LV hinge, enl. (160).
- Asaphis** MODEER, 1793 [*Venus deflorata* LINNÉ, 1758; M] [= *Capsa* BRUGUIÈRE, 1797 (obj.; SD SCHMIDT, 1818); *Corbula* RÖDING, 1798 (non BRUGUIÈRE, 1797) (obj.; SD WINCKWORTH, 1930); *Capsula* SCHUMACHER, 1817 (obj.; M); *Psammocola* DE BLAINVILLE, 1824 (obj.; SD BUCQUOY, DAUTZENBERG, & DOLLFUS, 1895); *Procos* GISTEL, 1848 (nom. van. pro *Capsa*); *Acapis* PAETEL, 1875 (nom. null.)]. Shells of medium size, with well-developed radial ribs, elliptical, inequilateral, not gaping; hinge plate well developed. *Mio.-Rec.*, S.Am.-IndoPac.-Indies.
- A. (Asaphis)**. Hinge plate narrow to moderate in width; ribs numerous, rounded in section. *Rec.*, E.Indies-W.Indies.—FIG. E116,3. **A. (A.) deflorata* (LINNÉ), W.Indies; 3a,b, LV ext., RV int., $\times 0.5$ (124b).
- A. (Asaphinoides)** HODSON, 1931 [**A. (A.) cantaurana* F. HODSON in HODSON & HODSON; OD]. Hinge plate wide, ribs few, sharply triangular in section. *L.Mio.*, S.Am.—FIG. E116,12. **A. (A.) cantaurana*, Venez.; 12a-c, LV int., ext., dorsal, $\times 1$ (Hodson & Hodson, 1931).
- A. (Heteroglypta)** VON MARTENS, 1880 [**Psammobia contraria* DESHAYES, 1863; M]. Anterior and posterior slopes with oblique ribs that intersect central ribs at angle, posterior ribs coarser. *Rec.*, IndoPac.—FIG. E116,10. **A. (H.) contraria* (DESHAYES), Ind.O.; 10a,b, RV ext., int., $\times 1$ (Deshayes, 1863).
- Ascitellina** MARWICK, 1928 [**A. donaciformis*; OD]. Small, posterior end short, sculpture of concentric ridges; hinge with posterior cardinals bifid, laterals weak or absent; ligament external, nymphs narrow, somewhat sunken posteriorly; pallial sinus large, nearly reaching anterior adductor, partly confluent. *Oligo.-Rec.*, S.Pac.—FIG. E116,7. **A. donaciformis*, Oligo., N.Z.; 7a,b, LV ext., int., $\times 3$ (593).
- Heterodonax** MÖRCH, 1853 [**Tellina bimaculata* LINNÉ, 1758; SD KOBELT, 1881]. Rounded-quadrate to ovate, with concentric growth striae only, variously colored; hinge teeth large, laterals wanting. *Plio.-Rec.*, C.Am.-N.Am.—FIG. E116,5. **H. bimaculatus* (LINNÉ), *Rec.*, W.Indies; 5a,b, LV ext., RV int., $\times 1$ (124b).
- Liodonax** FISCHER, 1887 [**Donax auversiensis* DESHAYES, 1858; SD COSSMANN & PEYROT, 1911]. Trigonal, posterior end short; sculpture of fine radial ribs except on posterior slope. *Eoc.*, Eu.—FIG. E116,4. **L. auversiensis* (DESHAYES), France; 4a,b, LV ext., int., $\times 1$; 4c, surface sculpture, enl. (Deshayes, 1860).
- Macrosolen** ZITTEL, 1883, ex MAYER-EYMAR MS. [**Sanguinolaria hollowaysii* SOWERBY, 1817; M] [= *Latosiliva* DE GREGORIO, 1894 (type, *Solen plicatus* VON SCHAUROTH, 1865; OD)]. Elongate, oval, flaring behind, beaks subanterior; not gaping; posterior slope with a radial furrow; hinge with 2 diverging cardinals in either valve; ligament on a nymph; pallial sinus large. *L.Eoc.-Mio.*, Eu.-Asia-N.Afr.—FIG. E116,9. **M. hollowaysii* (SOWERBY), *Eoc.*, Eng.; 9a-c, LV int., RV int., LV ext., $\times 0.4$ (Brit.Mus.Guide, 1959).
- Orbicularia** DESHAYES, 1850 [**Solen orbiculatus* WOOD, 1828; OD] [= *Elizia* GRAY, 1854 (obj.)]. Orbicular, nearly flat; pallial sinus not confluent. *Rec.*, E.Indies.—FIG. E116,8. **O. orbiculata* (WOOD), 8a, LV ext., $\times 1$; 8b,c, RV and LV hinges, enl. (124b).
- Rhectomyax** STEWART, 1930 [**Asaphis undulata* GABB, 1864; OD]. Small, ovate-quadrate, beaks nearly central; not gaping; sculpture of concentric striae except for undulating radial ribs at ends of shell; hinge with 1 cardinal in each valve and a toothlike nymph. *U.Cret.*, W.N.Am.—FIG. E116,6. **R. undulatus* (GABB), USA (Calif.); 6a,b, RV ext., RV hinge, $\times 4$ (892); 6c,d, RV and LV hinges, $\times 5$ (specimen, Univ. Calif., Los Angeles).

Subfamily SANGUINOLARIINAE

Grant & Gale, 1931

[nom. transl. KEEN, herein (ex Sanguinolariidae GRANT & GALE, 1931)]

Smooth or nearly so, inequilateral and more or less inequivalved; pallial sinus well developed. *Mio.-Rec.*

Sanguinolaria LAMARCK, 1799 [**Solen sanguinolentus* GMELIN, 1791; M] [= *Lobaria* SCHUMACHER, 1817 (non MÜLLER, 1776; obj.); *Isarcha* GISTEL, 1848 (nom. van.) (obj.)]. Ovate; periostracum of varying development. *Mio.-Rec.*, N.Am.-S.Am.-IndoPac.

S. (Sanguinolaria). Somewhat inflated, slightly gaping, posterior end pointed; valves nearly equal in size; shell tending to be tinged with red. *Mio.-Rec.*, N.Am.-S.Am.—FIG. E117,3. **S. (S.) sanguinolenta* (GMELIN), *Rec.*, Carib.; LV ext., $\times 1$ (124b).

S. (Hainania) SCARLATO, 1965 [**S. (H.) tchangsii*; OD]. *Rec.*, E.Asia.

S. (Nuttallia) DALL, 1900 [**S. nuttallii* CONRAD, 1837; OD]. Large, rounded; RV flattened; shell gaping posteriorly; hinge with posterior cardinal obsolete in LV; pallial sinus somewhat detached. *Mio.-Rec.*, E.Asia-W.N.Am.—FIG. E117,5. **S. (N.) nuttallii* (CONRAD), *Rec.*, USA (Calif.); RV ext., $\times 0.3$ (307).

S. (Psammospaerica) JOUSSEAUME, 1894 [**P. psammospaerita*; M] [= *Psammospaerita*, spelling error]. Like *S. (Sanguinolaria)* but without any gape. *Rec.*, Ind.O.

S. (Psammotella) HERRMANNSEN, 1852 (ex DE BLAINVILLE, 1826, vernac.) [**Tellina rufescens* "CHEMNITZ" (nonbinom.) (= **T. cruenta* LIGHTFOOT, 1786; M)]. Elongate-ovate, LV flattened, somewhat rostrate posteriorly; without periostra-




FIG. E117. Psammobiidae (Sanguinolariinae) (p. N633-N634).

cum; valves pinkish-buff in color. *Rec.*, W.C.Am.—FIG. E117,1. **S. (P.) cruenta* (LIGHTFOOT), Brazil; LV ext., RV int., $\times 0.5$ (Chemnitz, 1782).

S. (Psammotellina) FISCHER, 1887 [**Psammotella ambigua* REEVE, 1857 (ex DESHAYES MS); M] [= *Psammotella* H. ADAMS & A. ADAMS, 1856 (nom. nud.), REEVE, 1857 (non HERRMANNSEN, 1852); *Flavomala* IREDALE, 1936 (type, *Solen biradiatus* WOOD, 1815; OD)]. Thin, with a deciduous periostracum, asymmetrical but not rostrate posteriorly, tending to be tinged bluish-purple. *Rec.*, IndoPac.—FIG. E117,2. **S. (P.) ambigua* (REEVE), E.Indies; RV ext., $\times 0.5$ (783).

S. (Soletellina) DE BLAINVILLE, 1824 [**S. radiata* (= **Solen diphos* LINNÉ, 1771); M] [= "*Psammotaea* LAMARCK" AUCT. (non LAMARCK); *Florisarka* IREDALE, 1936 (type, *F. onuphria*, = *Soletellina donacioides* REEVE, 1857; OD)]. Elongate-ovate, posterior end somewhat rostrate,

ventral margin sinuate; periostracum dark-colored, shell yellowish to violet, variously rayed with darker stripes. *Mio.-Rec.*, IndoPac.—FIG. E117, 4. **S. (S.) diphos* (LINNÉ), *Rec.*, E.Indies; 4a, LV ext., $\times 0.7$; 4b, RV int., $\times 0.5$ (4a, 124b; 4b, 1007).

Family UNICARDIOPSIDAE Vokes, 1967

[=Unicardiidae FISCHER, 1887 (*partim*)] [Materials for this family prepared by ANDRÉ CHAVAN]

Shell medium-sized, transversely ovate, inflated, anteriorly produced and rounded, posteriorly truncated, very inequilateral; with prominent opisthogyrous beaks; escutcheon well marked; surface with coarse concentric ribs. Hinge showing general tellinacean tendency to develop anteriorly, with tooth 2 small oblique, 3b tubercular and superficial, 4b thin; semicircular resilium adjacent to prominent flat nymph; integripalliate but with superior diverticulation of posterior scar. *M.Jur.-U.Jur.*

Unicardiopsis CHAVAN, 1962 [**Unicardium aceste* D'ORBIGNY, 1850; OD]. Characters of family. *M. Jur.-U.Jur.*, Eu.—FIG. E118,1. **U. aceste* (D'ORBIGNY), *U.Jur.*(Sequan.), France; 1a, LV int. (112); 1b, LV ext., $\times 2$ (Chavan, 1950); 1c, RV int., ca. $\times 2$ (Chavan, n).

Family QUENSTEDTIIDAE Cox, 1929

Equivalve, subequilateral, rectangular, compressed, ligament external, in long pit; hinge plate flat, narrow. *L.Jur.-M.Jur.*




FIG. E118. Unicardiopsidae (p. N634).

Quenstedtia MORRIS & LYCETT, 1855 [**Pullastra oblita* PHILLIPS, 1829; SD STOLICZKA, 1871] [= *Corbicella* MORRIS & LYCETT, 1855 (type, *Corbis* (*C.*) *bathonica*; SD STOLICZKA, 1871)]. Sculpture of irregular concentric ribs; hinge with 1 cardinal in LV, socket and faint *All* in front of it; RV with 1 socket and incipient cardinal; lateral teeth otherwise wanting; posterior adductor scars rounded, anterior elongate and sinuate; pallial sinus small, rounded. *L.Jur.-M.Jur.*, Eu.—FIG. E119,3a. **Q. oblita* (PHILLIPS), *M.Jur.*, Eng.; LV ext., $\times 1$ (Phillips, 1829).—FIG. E119,3b,c. *Q. rodborensis* (LYCETT, 1851), *M.Jur.*, Eng.; 3b,c, RV int., LV int., $\times 0.5$ (Arkell, 1934).

Family ICANOTIIDAE Casey, 1961

Equivalve, closed, compressed, somewhat oblong, anterior and narrower; smooth to radially ribbed, especially on posterior slope; ligament on nymphs; hinge with two cardinals in either valve, no laterals; pallial sinus large, deep, rounded. *Cret.*(*Hauteriv.-Maastricht.*).

Icanotia STOLICZKA, 1870 [**Psammobia impar* ZITTEL, 1865; OD]. Ovate-elongate, anterior end short; radial sculpture well-developed except on lunular and escutcheonal areas. *Cret.*(*Apt.-Maastricht.*), Eu.-N.Am.-Asia.—FIG. E119,1. **I. impar* (ZITTEL), *U.Cret.*, Ger.; LV ext., $\times 0.7$ (Zittel-Eastman).

Scittila CASEY, 1961 [**S. nasuta*; OD]. Ovate-quadrate, without lunule and with incipient escutcheon; beaks subcentral, posterior margin obliquely truncate, posterior slope set off by low ridge; central slope with low radial furrow sinuating ventral margin; sculpture weak to wanting. *L.Cret.*(*Hauteriv.-Apt.*), Eu.—FIG. E119,2. **S. nasuta*, Eng., $\times 1$; 2a-c, LV ext., LV and RV hinges, $\times 2$ (Casey).

Family SCROBICULARIIDAE Adams & Adams, 1856

Resembling Semelidae, with sunken resilium, but without lateral teeth on hinge; posterior end without evident flexure (508). *Eoc.-Rec.*

Scrobicularia SCHUMACHER, 1815 [**S. calcarea* (= **Trigonella plana* DA COSTA, 1778); SD BUCQUOY, DAUTZENBERG, DOLLFUS, 1898] [= *Ligula* MONTAGU, 1808 (*non* BLOCK, 1782) (obj.); *Arenaria* MEGERLE VON MÜHLFELDT, 1811 (*non* BRISSON, 1760) (obj.); *Lavignonus* FÉRUSAC, 1821 (*ex* CUVIER, 1817, *vernac.*) (obj.), *Lavigno*, *Lavignon*, *Lavignona*, spelling errors; *Listera* TURTON, 1822 (obj.); *Lutricola* DE BLAINVILLE, 1824 (type, *L. compressa*, = *Trigonella plana* DA COSTA, 1778;




FIG. E119. Quenstedtiidae (3); Icanotiidae (1-2); Scrobiculariidae (4-5) (p. N635-N636).

SD KEEN, herein) (obj.); *Calcinella* DESHAYES, 1830 (*pro Ligula, partim*); *Carinella* SOWERBY, 1839 (*non* JOHNSTON, 1833) (obj.); *Martinea* BUCQUOY, DAUTZENBERG, & DOLLFUS, 1898 (*ex* DA COSTA MS.) (obj.)]. Smooth, compressed, lenticular, with well-developed chondrophores. *Eoc.-Rec.*, Eu.-E.Indies.—FIG. E119,4. **S. plana* (DA COSTA), *Rec.*, Eu.; LV ext. showing siphons, $\times 1$ (H. Adams & A. Adams).




FIG. E120. Semelidae (p. N636-N637).

Scrobiculabra WOOD, 1877 [**S. dulwichiensis*; SD COSSMANN, 1887]. Posterior end rostrate, set off by ridge; hinge of RV with 2 large cardinal teeth. *Eoc.*, Eu.—FIG. E119, 5. **S. dulwichiensis*, Eng.; 5a, RV ext., $\times 2$; 5b, RV hinge, enl. (Wood).

Family SEMELIDAE Stoliczka, 1870

Shells mostly well sculptured, with slight posterior flexure; ligament both external and internal, resilium sunken or lodged in small chondrophore; hinge with two cardinal teeth, laterals present in most forms; pallial sinus large, rounded (508). *Eoc.-Rec.*

Semele SCHUMACHER, 1817 [**S. reticulata* (= **Tel-*

lina proficua PULTENEY, 1799); M] [= *Amphidesma* LAMARCK, 1818 (type, *A. variegata*; SD CHILDREN, 1822); *Syndesmyella* SACCO, 1901 (type, *S. plicovoides*; OD); *Elegantula* DE GREGORIO, 1884 (type, *S. fazisa*, = *Amphidesma striata* REEVE, 1853; M)]. Mostly large shells, with anterior end longer than posterior; sculpture radial, concentric, or oblique; resilium in ovoid depression of hinge plate; lateral teeth stronger in RV than in LV, cardinals subequal; pallial sinus ascending obliquely, pallial line not confluent. *Eoc.-Rec.*, N.Am.-S.Am.-Eu.-Pac.—FIG. E120, 5. **S. proficua* (PULTENEY), *Rec.*, Carib.; 5a, b, LV int., RV hinge, $\times 1$ (7c).

Abra LAMARCK, 1818 (ex LEACH MS) (proposed in synonymy) [**Mactra tenuis* MONTAGU, 1818; SD

- HERRMANNSEN, 1846] [= *Habra*, *nom. null.*; *Orixa*, *Dorvillea* GRAY, 1852 (*ex* LEACH, MS) (obj.; M); *Lutricularia* MONTEROSATO, 1884 (type, *Erycina ovata* PHILIPPI, 1836, *non* GRAY, 1825, = *Syndosmya segmentum* RÉCLUZ, 1843; SD CROSSE, 1885); *Abrina* HABE, 1952 (type, *Abra kanamarui* KURODA, 1951; OD)]. Thin, small resilium in a chondrophore that projects slightly into the shell cavity; pallial line partially confluent. *Eoc.-Rec.*, cosmop.
- A. (Abra).** Trigonal, beaks high, pointed; lateral teeth weak. *Mio.-Rec.*, Eu.-N.Am.-S.Am.-Pac.—FIG. E120,1. **A. (A.) tenuis* (MONTAGU), *Rec.*, Eng.; LV ext. showing siphons and foot, $\times 1$ (7c).
- A. (Iacra)** H.ADAMS & A.ADAMS, 1856 [**Scrobicularia (I.) seychellarum* A.ADAMS, 1856; M] [= *Strigillina* DUNKER, 1861 (type, *S. lactea*)]. Surface divaricately sculptured; posterior flexure strong. *Rec.*, IndoPac.
- A. (Syndosmya)** RÉCLUZ, 1843 [**Mactra alba* WOOD, 1802; SD WOODWARD, 1854] [= *Sinodesmia*, *Sinodesmya*, *Syndesmya*, *Syndomya*, spelling errors]. Hinge as in *A. (Abra)* but shell more ovate; lateral teeth stronger. *Eoc.-Rec.*, E.N.Am.-Eu.—FIG. E120,8. **A. (S.) alba* (WOOD), *Rec.*, Eng.; *8a,b*, LV ext., LV ext., $\times 1$ (Forbes & Hanley, 1854).
- Cumिंगia** SOWERBY, 1833 [**C. lamellosa*; GRAY, 1847] [= *Harpax* GISTEL, 1848 (*non* PARKINSON, 1811) (obj.); *Mikrola* MEYER, 1887 (type, *M. mississippiensis*; M); *Cumिंगia* (spelling error)]. Outline irregular, due to nestling habit, generally rounded in front, angular posteriorly; resilifer large, cardinal teeth small, laterals large in RV, obsolescent in LV, pallial sinus partially confluent. *Eoc.-Rec.*, N.Am.-S.Am.-IndoPac.—FIG. E120,3. *C. mutica*, *Rec.*, Ecuador; *3a,b*, RV int., LV hinge, $\times 1$ (H.Adams & A.Adams, 508).
- Leptomya** A. ADAMS, 1864 [**Neaera cochlearis* HINDS, 1844; SD STOLICZKA, 1871]. Smooth, posterior end more or less pointed; pallial sinus large, partially confluent. *Rec.*, S.Pac.-Japan-C.Am.
- L. (Leptomya).** Posterior end rostrate; lateral teeth of moderate length. *Rec.*, W.Pac-S.Pac.-W.C.Am.—FIG. E120,7. **L. (L.) cochlearis* (HINDS), E.Indies; LV ext., $\times 1$ (Hanley, 1882).
- L. (Leptomylaria)** HABE, 1960 [**Leptomylaria trigona*; OD]. Posterior end rounded, anterior widely produced; lateral teeth long. *Rec.*, Japan.—FIG. E120,4. **L. (L.) trigona* (HABE); *4a,b*, RV ext., int., $\times 4$ (Habe, 1960).
- Montrouzieria** SOUVERBIE, 1863 [*pro* *Montrouzieria* SOUVERBIE, 1863 (*non* BIGOT, 1860)] [**Montrouzieria clathrata*; OD] [= *Eumontrouzieria* HEDLEY, 1915 (obj.)]. Small, trapezoidal, with radial striae; chondrophores triangular, external ligament short. *Rec.*, Pac.-E.Indies.
- M. (Montrouzieria).** Hinge with 2 cardinals in RV, 1 in LV, 1 lateral in each. *Rec.*, Pac.—FIG. 120,11. **M. (M.) clathrata* SOUVERBIE, *Rec.*, N.Caledon., *11a-c*, RV ext., LV int., RV int., $\times 2$ (Souverbie).
- M. (Lonoa)** DALL, BARTSCH & REHDER, 1939 [**L. hawaiiensis*; OD]. With strong concentric sculpture in addition to fine radial ribs; resilifer shelflike. *Rec.*, Pac.—FIG. E120,10. **M. (L.) hawaiiensis* (DALL, BARTSCH & REHDER), Hawaii; *10a-c*, LV ext., int., RV int., $\times 4$ (Dall, Bartsch & Rehder).
- M. (Thyellisca)** VOKES, 1956 [*pro* *Thyella* H. ADAMS, 1866 (*non* WALLENGREN, 1858)] [**Thyella pulchra* H.ADAMS, 1866; OD]. Lateral teeth wanting. *Rec.*, E.Indies.
- Semelina** DALL, 1900 [**Amphidesma nuculoides* CONRAD, 1841; OD]. Small, sculpture concentric only; hinge of LV without laterals and posterior cardinal, chondrophore short. *L.Mio.-Rec.*, E.USA-Carib.—FIG. E120,9. **S. nuculoides* (CONRAD), *Rec.*, USA(Fla.); *9a-c*, LV ext., LV int., RV int., $\times 4$ (Gardner).
- Septeuilia** COSSMANN, 1914 [**Scrobicularia bezanconi* COSSMANN, 1887; SD KEEN, herein]. Small, elongate, posterior end shorter, truncate; hinge with 1 large nonbifid cardinal and a lamellar lateral in LV. *Eoc.*, Eu.—FIG. E120,2. **S. bezanconi* (COSSMANN), France; *2a-c*, LV ext., int., RV hinge, $\times 3$ (160).
- Theora** H.ADAMS & A.ADAMS, 1856 [**Neaera lata* HINDS, 1843; SD STOLICZKA, 1871]. Compressed, smooth, hyaline, with a posterior gape; hinge weak. *Rec.*, E.Indies.
- T. (Theora).** Adductor scars elongate; cardinal teeth present, LV with no laterals. *Rec.*, E.Indies.—FIG. E120,6. **T. (T.) lata* HINDS; *6a-c*, RV int., LV ext., hinge, $\times 1$ (H.Adams & A.Adams, 508).
- T. (Endopleura)** A.ADAMS, 1864 [**T. lubrica* GOULD, 1861; M]. With one cardinal bifid; inside of shell with a radial rib. *Rec.*, E.Indies.
- T. (Souleyetia)** RÉCLUZ, 1869 [**S. moulinisii*; M]. Resembling *T. (Theora)* but without hinge teeth; chondrophore oblique. *Rec.*, E.Indies.

Family SOLECURTIDAE d'Orbigny, 1846

[=Pharinae ADAMS & ADAMS, 1856]

Elongate-quadrate, widely gaping at both ends; hinge plate weak, narrow; pallial sinus shallow to deep (223). *L.Eoc.-Rec.*

Subfamily SOLECURTINAE d'Orbigny, 1846

[*nom. transl.* GHOSH, 1920 (*ex* Solecurtidae d'ORBIGNY, 1846)]

Beaks subcentral. *L.Eoc.-Rec.*

Solecurtus DE BLAINVILLE, 1824 [**Solen strigilatus* LINNÉ, 1758; SD DESHAYES, 1829] [= *Psammosolen* RISSO, 1826 (obj.); *Macha* OKEN, 1835

(obj.); *Silex* QUOY & GAIMARD, 1835 (type, *S. albus* = *Solen rhombus* SPENGLER, 1794; M); *Cyrtosolen* HERRMANNSEN, 1848 (*nom.van. pro*

Solecurtus); *Adasius* GRAY, 1852 (*ex LEACH MS*) (obj.); *Solecurellus* GHOSH, 1920 (type, *Solen dombeyi* LAMARCK, 1818; MJ]. Sculptured with




FIG. E121. Solecurtidae (Solecurtinae) (1,3-6,8); (Novacutininae) (2,7) (p. N637-N639).



Veneroida—Tellinacea

spaced oblique striae; hinge without lateral teeth, cardinals 2 in RV, 1 in LV, pallial sinus large; ligamental nymph long. *Eoc.-Rec.*, Atl.-Pac.—FIG. E121,8. **S. strigilatus* (LINNÉ), *Rec.*, *Medit.*; 8a-e, RV ext., LV ext., RV int., LV int., both valves dorsal, $\times 0.7$ (89a).

Azorinus RÉCLUZ, 1869 [**Solen coarctatus* GMELIN, 1791 (= **S. chamasolen* DA COSTA, 1778); M] [= *Azor* BROWN, 1844 (ex LEACH MS) (non SOWERBY, 1824) (obj.); *Zozia* WINKWORTH, 1930 (obj.; OD)]. Hinge with 2 cardinal teeth in either valve. *L.Eoc.-Rec.*, Eu.-Asia.

A. (Azorinus). Surface smooth. *Plio.-Rec.*, Eu.—FIG. E121,1. **A. (A.) chamasolen* (DA COSTA), *Rec.*, *Medit.*; 1a-c, RV ext., LV int., RV int., $\times 0.5$ (89a).

A. (Zoziella) EAMES, 1951 [**A. (Z.) punjabensis*; OD]. Small, with undulating concentric folds and 2 central radial furrows. *L.Eoc.*, Asia.—FIG. E121,3. **A. (Z.) punjabensis*, Pak., RV ext., $\times 1.5$ (288).

Pharus BROWN, 1844 (in synonymy) [**Solen legumen* LINNÉ, 1758; M] [= *Polia* D'ORBIGNY, 1845 (non OCHSENHEIMER, 1816) (obj.); *Ceratisolen* FORBES & HANLEY, 1848 (obj.); *Erratisolen*, *Seratisolen* AUCT. (nom.null.); *Artusius* GRAY, 1852 (ex LEACH MS) (obj.)]. Cylindrical, ligament sub-central, no ligamental nymph; hinge with 2 cardinals in LV, 1 in RV, long low anterior lateral and short projecting posterior lateral; pallial sinus short. *L.Mio.-Rec.*, Eu.—FIG. E121,5. **P. legumen* (LINNÉ), *Rec.*, *Medit.*; 5a-d, LV ext., RV int., LV int., RV int., $\times 0.7$ (89a).

Tagelus GRAY, 1847 [*“*Sol. guinensis*” (non *Solen guinensis* HANLEY, 1842) (pro “*Solen tagel* ADANSON, 1757” of GRAY) = **Solen adansonii* BOSCH, 1801 (better known as *Solecurtus angulatus* SOWERBY, 1874); OD] [= *Siliquaria* SCHUMACHER, 1817 (non BRUGUIÈRE, 1789) (type, *S. notata*, = *Solen plebeius* LIGHTFOOT, 1786; M); *Cultellus* CONRAD, 1837 (non SCHUMACHER, 1817)]. Narrower than *Solecurtus*, surface smooth; periostracum present; teeth 2 in either valve, simple; pallial sinus deep, mostly reaching to or beyond vertical mid-line, partly confluent with pallial line. [Estuarine to marine.] *Oligo.-Rec.*, N.Am.-S.Am.-Eu.-W.Afr.

T. (Tagelus). Medium-sized to large, without any median rib. *Oligo.-Rec.*, N.Am.-S.Am.-Eu.-W.Afr.—FIG. E121,6. **T. (T.) adansonii* (Bosc), *Rec.*, W.Afr.; RV ext., $\times 0.5$ (Fischer-Piette, 1942).

T. (?Clunaculum) DALL, 1899 [**Solecurtus mollis* SOWERBY, 1874 (ex GOULD MS); OD]. Valves obliquely constricted; pallial sinus confluent with pallial line, not reaching beaks; posterior adductor scar triangular. *Rec.*, ?S.Am.

T. (Mesopleura) CONRAD, 1868 [**Solen bidentatus* SPENGLER, 1794 (= **S. divisus* SPENGLER, 1794); SD STOLICZKA, 1871] [= *Subtagelus* GHOSH, 1920

(obj.)]. Smaller and thinner than *T. (Tagelus)* with a central rib, especially in young (obsolete in some adults), showing exteriorly as dark line; pallial sinus relatively short, nymphs long. *Plio.-Rec.*, E.N.Am.-W.N.Am.-C.Am.—FIG. E121,4. **T. (M.) divisus* (SPENGLER), *Rec.*, W.Indies; LV ext. and RV int., $\times 1$ (Chemnitz, 1795).

Subfamily NOVACULININAE Ghosh, 1920

[nom. correct. YONGE, 1949 (ex Novaculinae GHOSH, 1920)]

Long-cylindrical, resembling Solenidae but with soft parts as in Solecurtinae; beaks nearly at anterior end, hinge with two or three cardinal teeth; pallial sinus small to large. *Rec.*

Novaculina BENSON, 1830 [**N. gangetica*; M] [= *Loncosilla* RAFINESQUE, 1831 (type, *L. solenoides*; M); *Navaculina* (nom.null.)]. Shell width uniform throughout; periostracum present; hinge with 3 cardinals in LV, 2 in RV in most; pallial sinus of moderate depth. [Freshwater.] *Rec.*, S. Asia.—FIG. E121,2. **N. gangetica*, India; 2a,b, RV int., LV int., $\times 1$ (1007; Prashad, 1924).

Sinonovacula PRASHAD, 1924 [**Solen constrictus* LAMARCK, 1818; OD]. Larger than *Novaculina*, more equivalve, beaks lower, pallial sinus shorter and broader; valves obliquely compressed near middle. [Estuarine.] *Rec.*, E.Asia.—FIG. E121,7. **S. constrictus* (LAMARCK), IndoChina; LV int., $\times 0.5$ (Prashad, 1924).

Family SOWERBYIDAE Cox, 1929

[= Isodontidae ARKELL, 1934]

Inequilateral to oblique in outline; hinge with large lateral teeth and normally with two cardinals. ?*Trias.*, *U.Jur.*

Sowerbya D'ORBIGNY, 1850 [**S. crassa*; M] [= *Isodonta* BUVIGNIER, 1851 (type, *I. deshayesia*; M); *Isodon* (emend. DOUVILLÉ, 1912) (non SAY, 1822)]. Lateral teeth long and large; shell resembling *Mactra* as to form but without chondrophore; cardinal teeth 2 in either valve; pallial sinus evident, rounded. *U.Jur.* (Oxford.), Eu.—FIG. E122,2. **S. crassa*, France; 2a-c, RV ext., int., LV int., $\times 1$ (Cottreau).

?**Rhaetidia** BITTNER, 1895 [**R. zittelii*; OD]. Resembling *Sowerbya* in outline but hinge with large lateral teeth only, no cardinals; pallial line entire. *Trias.*, Eu.—FIG. E122,1. **R. zittelii*, U.Trias., Alps; 1a, RV ext., $\times 2$; 1b, both valves dorsal, $\times 2$; 1c, RV ext., $\times 1$ (Bittner).

Family TANCREDIIDAE Meek, 1864

[Materials for this family prepared by L. R. Cox]

Shell small to medium-sized, smooth or with weak radial ornament, equivalve,




FIG. E122. Sowerbyidae (p. N639).

feebly to moderately inflated, ovate or sub-trigonal, more or less elongate, subequilateral to strongly inequilateral, in latter case with anterior end longer; ligament external, opisthodontic, usually short; cardinal teeth fundamentally two in each valve (2, 4b in LV, 3a, 3b in RV), but decreased to single tooth in one or both valves in some forms by extreme reduction of LV posterior (4b), of RV anterior (3a), or of both, and increased to three in some LV's of *Eodonax* by development of thin ridge (4a?) adjoining lunular margin; single posterior lateral (PII) present in LV and received, below dorsal margin of RV, in socket which may lie between two distinct lateral teeth (PIII, PI); anterior laterals, if distinguishable, consisting of laminar projections from margins; adductor scars small; pallial line simple or with shallow sinus. *U.Trias.-U.Cret.*

Tancredia LYCETT, 1850, p. 407 [**T. donaciformis*; SD MORRIS & LYCETT, 1855, p. 91]. Subtrigonal, with anterior end of shell tapering to more or less acute extremity; subequilateral to strongly inequilateral; posterior margins commonly gaping; adductor scars circular or oval, subequal, placed in rather dorsal position; pallial line remote from ventral margin, bending abruptly upward posteriorly. *U.Trias.-L.Cret.*, cosmop.

T. (*Tancredia*) [= *Hettangia* TERQUEM in BUVIGNIER, 1852, p. 14] (type, *H. deshayesea*; SD

TERQUEM, 1855, p. 290)]. Mostly rather large, subequilateral, high and obliquely truncated posteriorly, subangular anteriorly; posterior slope ridged; lunule narrow and elongate, bordered by ridge; cardinal teeth moderately strong, 2 usually distinguishable in both valves; posterior laterals relatively strong; LV anterior lateral a laminar projection of dorsal margin, RV anterior laterals rarely distinguishable. *L.Jur.(L.Lias.)-U.Jur.(Oxford.)*, cosmop.—FIG. E123,4a-e. **T.* (*T. donaciformis*) LYCETT, M.Jur.(Aalen.), France; 4a-e, LV ext., both valves dorsal, RV int., LV int. and int. mold showing pallial line, all $\times 1$ (Benecke, 1905).—FIG. E123,4f,g. *T.* (*T. coxi*) CHAVAN, U.Jur.(U.Oxford.), France; 4f,g, RV int., LV int., $\times 10$ (Chavan, 1950).

T. (*Corburella*) LYCETT, 1850, p. 422 [**Corbula curtansata* PHILLIPS, 1829, p. 128; OD]. Medium-sized subequilateral, high and rounded posteriorly, rostrate anteriorly, ventral margin sinuate; posterior slope not ridged; no lunule; only 1 cardinal tooth in each valve; LV posterior lateral small but elongate; anterior laterals not distinguishable. *M.Jur.-U.Jur.*, cosmop.—FIG. E123,6. **T.* (*C. curtansata*) (PHILLIPS), U.Jur.(Oxford.), Eng.; LV ext., $\times 1$ (19f).

T. (*Isotancredia*) CHAVAN, 1950, p. 12 [**Tancredia extensa* LYCETT, 1850, pl. 11, fig. 9; OD]. Small, elongate, subtrigonal, subequilateral, with dorsal margins straight and sloping, anterior more steeply than posterior; posterior end of shell rather low, obliquely truncated, anterior end cuneiform; posterior slope ridged; cardinal teeth 1 in each valve; posterior laterals moderately prominent; RV anterior lateral present but weak. *M.Jur.-U.Jur.*, cosmop.; *L.Cret.*, S.Afr.—FIG. E123,2. **T.* (*I. extensa*) LYCETT, M.Jur.(Bathon.), Eng.; LV ext., $\times 1$ (Morris & Lycett, 1855).

T. (*Paratancredia*) CHAVAN, 1950, p. 12 [**T. (P.) brasili*; OD]. Small to medium-sized, compressed, inequilateral, less elongated than in other subgenera, high and obliquely truncated posteriorly, tapering to some extent anteriorly, although less than in other subgenera; cardinal teeth 2 in each valve, anterior ones elongated; posterior laterals small, RV anterior lateral distinct, elongate. *M.Jur.-U.Jur.*, cosmop.—FIG. E123,1. **T. (P.) brasili*, U.Jur.(Oxford.), France; 1a,b, LV int., RV int., $\times 3$, $\times 7$ (Chavan, 1950).

T. (*Palaeomya*) ZITTEL & GOUBERT, 1861, p. 194 [**Palaeomya deshayesi* ZITTEL & GOUBERT, 1861 (non BUVIGNIER) (= *Tancredia corallina* ZITTEL, 1881, p. 97); M] [= *Rosenbuschia* RÖDER, 1882, p. 97 (type, *R. typica*; M)]. Elongate, very inequilateral, anterior end much longer than posterior end, tapering; LV anterior cardinal an elongate lamina; posterior laterals prominent; RV anterior lateral distinct, elongate. *L.Jur.-U.Jur.*, cosmop.—FIG. E123,3. **T. (P.) corallina* (ZITTEL & GOUBERT), U.Jur.(Oxford.), France; 3a-c, RV




FIG. E123. Tancrediidae (p. N640-N642).

ext., LV int., RV int., all $\times 2$ (Zittel & Goubert, 1861).

Corbicellopsis COX, 1929, p. 577 [**Corbis laevis* J. DE C. SOWERBY, 1827, p. 156; OD] [= *Corbicella* AUCT. (non MORRIS & LYCETT, 1855) (cf. synonymy of *Quenstedtia*)]. Of medium to large-size, ovate, anterior taper less pronounced than in *Tancredia*; equilateral or with beaks anterior to mid-length; no posterior gape; nymphs prominent; cardinal teeth 1 or 2 in each valve; posterior laterals well developed except in smaller species; anterior laterals weak, indistinguishable in some forms; adductor scars small, subequal, placed in rather dorsal position; pallial line remote from ventral margin, bending up abruptly to meet posterior adductor. *M. Jur. (Bajoc.)-L. Cret., Eu.*—FIG. E123,7. **C. laevis*, U. Jur. (Oxford), Eng.; 7a,b, RV ext. and hinge, $\times 1$ (19f).

Eodonax COX, 1929, p. 584 [**Sowerbya dukei* MORRIS & LYCETT in DAMON, 1860, p. 172; OD]. Medium-sized, subovate or subtriangular, more or less pointed anteriorly, rounded or obliquely truncated posteriorly; subequilateral or with beaks just posterior to mid-length; no gape; nymphs prominent; 2 moderately strong, grooved cardinal teeth in LV, also thin ridge in some specimens (possibly 3rd cardinal) adjoining margin of lunule; 2 cardinals in RV, anterior tooth thin, posterior one strong and grooved, median ridge in recess between it and nymph corresponding to groove in LV posterior cardinal; lateral teeth weak; adductor scars small, dorsally placed; pallial line remote from margin, with small sinus. *U. Jur.-L. Cret., Eu.*—FIG. E123,8. **E. dukei* (MORRIS & LYCETT), U. Jur. (Portland), Eng.; 8a, LV ext., $\times 1$ (171); 8b,c, LV int., RV int., $\times 1$ (Cox, n).

Meekia GABB, 1864, p. 191 [**M. sella*; SD STOLICZKA, 1871, p. 312]. Subovate except for sharp anterior rostrum, slightly to moderately inequilateral, anterior end shorter; valve margins closed posteriorly in earlier, gaping in later species; anterior margins with narrow gape below rostrum; lunule lanceolate; surface unornamented or with weak radial lines; cardinal teeth 2 in each valve, on short hinge plate, anterior cardinal of LV broad, prominent, received in deep socket between smaller, subequal cardinals of RV; posterior cardinal of LV parallel to nymph; posterior laterals weak, 1 in LV received between 2 in RV; no anterior laterals; shelly internal buttress runs in anteroventral direction from beneath hinge plate, passing below anterior adductor, pallial line entire. *L. Cret. (Alb.)-U. Cret. (Maastricht.)*, W.N. Am.-Japan.

M. (Meekia). Relatively thick and heavy; pronouncedly rostrate, with strongly concave anterodorsal margin; inflation moderately strong; surface with more or less distinct radial ornament and with microscopic punctations; anterior cardinal of LV peglike. *Cret. (Alb.-Maastricht.)*,




FIG. E124. Superfamily and family Uncertain (p. N642-N643).

W.N. Am.-Japan.—FIG. E123,9. **M. (M.) sella* GABB, U. Cret. (Maastricht.), USA (Calif., Martinez), 9a-c, LV ext., LV int., RV int., $\times 1$ (Saul & Popenoe, 1962).

M. (Mygallia) SAUL & POPENOE, p. 962, p. 302 [**Meekia mygale*; OD]. Differs from *M. (Meekia)* in thinner, more elongated and less inflated shell, less marked anterior rostrum, absence of surface punctations, and more elongate anterior cardinal tooth of LV. *Cret. (Alb.-Maastricht.)*, W.N. Am.-Japan.

Sakawanella ICHIKAWA, 1950, p. 245 [**S. triadica*; OD]. Small elongate-oblong, beaks just anterior to mid-length; LV with 1 cardinal tooth, RV with 2, anterior one ill-defined; posterior laterals elongate; adductor scars and pallial line as in *Tancredia*; interior of shell with 2 grooves diverging from below beak. *U. Trias., Japan.*—FIG. E123,5. **S. triadica*; LV int. mold, $\times 2$ (Ichikawa, 1950).


Doubtful TELLINACEA

Cowperesia MENDES, 1952, p. 86 [**Pseudocorbula anceps* REED, 1935, p. 34; OD]. Suboval or subtriangular, tapering posteriorly; equivale or slightly inequivale; umbones low, orthogyre, submedian; surface ornamented with concentric ridges or relatively smooth; posterior umbonal ridge rounded; lunule and escutcheon obscure; dentition consisting of 1 cardinal tooth below beak of RV and corresponding socket; shallow pallial sinus present. [In view of CHAVAN, this genus is refer-


able to the Tellinidae. KEEN agrees that it may be an ancestral tellinacean but considers it inappropriately classifiable in the Tellinidae.] *Permo-Trias.*, S.Am.(S.Brazil).—FIG. E124,1. **C. anceps* (REED), Corumbatai, Passa Dois F., S.Brazil; 1a,b, LV ext. and int., ×2 (Mendes, 1952). [NEWELL]

Terraia COX, 1934, p. 269 [**Solenomorpha altissima* HOLDHAUS, 1919, p. 12; OD] [= *Jacquesia* MENDES, 1944, p. 62 (type, *Myophoriopsis brasiliensis* REED, 1929; OD); *Holdhausiella* MENDES, 1952, p. 94 (type, *Sanguinolites elongatus* HOLDHAUS, 1919; OD); *Maackia* MENDES, 1954, p. 100 (type, *M. contorta*; OD); *Oliveiraia* MENDES, 1954, p. 103 (type, *Thracia pristina* REED, 1929; SD NEWELL, herein)]. Trigonally ovate, thick-shelled, equivalve, inequilateral, not greatly inflated; posterior end tapering to low, vertically truncated extremity, without gape; umbones obtusely angular, contiguous, placed at about anterior 0.3 of length;


lunule and escutcheon long, narrow, limited by carinae which in most specimens are sharp, although in some shells escutcheon carina may die out posteriorly; posterior slope with prominent elevated carina, separated from escutcheon by narrow, slightly concave area; surface with growth rugae which may produce denticulations on posterior carina; RV with single blunt tooth placed just behind umbo, narrow shallow groove separating this tooth from short nymph, which supported external, opisthodontic ligament; between tooth and anterior margin is shallow triangular recess; posterior margin somewhat thickened; LV with deep median recess for reception of RV tooth, bordered by narrow weak tooth separated by shallow groove from nymph; well-defined lateral teeth lacking; anterior adductor with low reinforcing buttress (clavicle); pallial line simple. *Permo-Trias.*, S.Am.(Brazil-N.Uruguay).—FIG. E124,2. **T. altissima* (HOLDHAUS), Estrada Nova F., Urug. (Rivera Prov.); 2a-c, LV int., RV ext., RV int., ×2 (Cox, 1934). [NEWELL]



1 Dreissenomya 2 Sinucongeria



4a 4b Congeria



5a 5b Mytilopsis

FIG. E125. Dreissenidae (p. N644).

Superfamily DREISSENACEA Gray in Turton, 1840

[*nom. transl.* GILL, 1871 (ex Dreissenidae GRAY in Turton, 1840)] [Materials for this superfamily prepared by MYRA KEEN]

Mytiliform to quadrate, beaks anterior or terminal; interior of shell not nacreous; ligament sunken, hinge edentulous; beak cavity bridged by septum or myophore; posterior adductor muscle scar long; periostracum well developed. Animal byssiferous; siphons two; gills reticulate. *Eoc.-Rec.*

Family DREISSENIDAE Gray in Turton, 1840

[*nom. correct.* GRAY, 1847 (ex Dreissenadac GRAY in Turton, 1840)]

Characters of superfamily. *Eoc.-Rec.*

Dreissena BENEDEN, 1835 [*nom. correct.* ICZN Op. 872 (pro *Driessena* BENEDEN, 1835)] [**Mytilus polymorphus* PALLAS, 1771 (errore pro *Mytilus*); M] [= *Tichogonia* ROSSMAESSLER, 1835 (obj.); *Dithalmia* JAY, 1835 (*nom. nud.*); *Mytilina* CANTRAINE, 1837 (non BORY DE ST. VINCENT, 1824) (obj.); *Coelogonia* BRONN, 1837 (type, *Mytilus brardii* BRONGNIART, 1823; M); *Mytilomya* BRONN, 1838 (obj.); *Mytilimax* SCHAUFUSS, 1869 (obj.); *Dreissena*, *Dreisseina*, *Dreissencia*, *Dreissenia*, *Dreissensa*, *Dreissensia*, *Dreissina*, *Dreistena*, *Dreysseina*, *Dreysseina*, *Dreysseusia*, *Dreysseusia*, *Dreysseusia*, *Driessensia* AUCTT. (*nom.null.*)]. Mytiliform, anteriorly compressed, with periostracum; septum with single adductor muscle scar. *Eoc.-Rec.*, Eu.-Afr.

D. (*Dreissena*). Smooth, pallial line entire. *Eoc.-Rec.*, Eu.-Afr.—FIG. E125,3. **D.* (*D.*) *polymorpha* (PALLAS), *Rec.*, Eu.; oblique anteroventral view of entire animal, $\times 1$ (Fischer).

D. (*Dreissenomya*) FUCHS, 1870 [**Dreissenomya schroeckingeri*; OD] [= *Congeriomya* ANDRUSOV, 1897 (*nom.van.*); *Dreissensiomya* FISCHER, 1886 (*nom.null.*); *Dreissenomya* NEUMAYR, 1891 (*nom.null.*);]. Septum obsolete; pallial line deeply sinuate: *Plio.*, E.Eu.—FIG. E125,1. **D.* (*D.*) *schroeckingeri* (FUCHS), Hungary, RV int., $\times 0.5$ (Papp).

D. (*Prodreissensia*) ROVERETO, 1898 [**D.* (*P.*) *perandoi*; OD]. Shell sculptured with longitudinal ribs. *Oligo.* (*Tongr.*), Eu.

D. (*Sinucongeria*) LÖRENTHEY, 1894 [**Congeria arcuata* FUCHS, 1870; OD]. Like *D.* (*Dreissenomya*) but more trigonal, septum larger, pallial sinus smaller. *Plio.*, E.Eu.—FIG. E125,2. **D.* (*S.*) *arcuata* (FUCHS), Hungary; RV int., $\times 1$ (Papp).

Congeria PARTSCH, 1835 [**C. subglobosa*; SD PILSBRY, 1911] [= *Enocephalus* MÜNSTER, 1831 (*nom.nud.*)]. Quadrangle, smooth, thick-shelled; septum with 2 scars, for anterior adductor and pedal retractor muscles; pallial line entire. *L.Oligo.-Plio.*, Eu.-W.Asia.—FIG. E125,4. **C. subglobosa*, Mio., Aus.; *4a,b*, ant. view of both valves, LV int., $\times 0.3$ (Fischer).

Mytilopsis CONRAD, 1858 [**Mytilus leucophaeatus* CONRAD, 1831; SD DALL, 1898] [= *Praxis* H. ADAMS & A. ADAMS, 1857 (*non* GUENÉE, 1852); *Mytiloides*, spelling error]. Mytiliform, as in *Dreissena*, but septum backed by a myophore with 2 scars, as in *Congeria*. *U.Oligo.-Rec.*, W.S.Am.-Eu.-Afr.-E.Indies.—FIG. E125,5. **M. leucophaeatus* (CONRAD), *Rec.* USA (Va.); *5a,b*, RV ext., LV int., $\times 1.5$ (Keen, n).

Superfamily GAIMARDIACEA Hedley, 1916

[*nom. transl.* FLEMING, herein (*ex* Gaimardiidae HEDLEY, 1916)] [Materials for this superfamily prepared by C. A. FLEMING]

Small, thin, equivalve, ventricose, with anterior beaks, surface smooth or radially ribbed; integripalliate; anterior retractor separate from adductor. Ligament external or sunken, opisthodontic; hinge weak, basically with single LV cardinal and bifid RV cardinal (tooth 1 invariably beneath bifid 3), LV and RV anterior laterals and LV posterior lateral varyingly reduced. Foot byssiferous, with creeping disc; mantle triforic, fry-developed in gills. [Commonly attached to floating seaweed.] (685, 908.) *Mio.-Rec.*

Family GAIMARDIIDAE Hedley, 1916

[*nom. correct.* ODHNER, 1924, p. 63 (*pro* Gaimardiidae HEDLEY, 1916, p. 26, *nom. imperf.*, *nom. subst. pro* Modiolarciidae, *nom. correct. pro* Modiolarciidae J. E. GRAY in M. E. GRAY, 1857, p. 25)] [Materials for this family prepared by C. A. FLEMING, New Zealand Geological Survey]

Characters of superfamily. *Mio.-Rec.*

Gaimardia GOULD, 1852, p. 459 [**Modiola trapesina* LAMARCK, 1819, p. 114; M] [= *Modiolarca* GRAY, 1847, p. 199 (*non* GRAY, 1843, p. 259) (*obj.*); *Phaseolicama* ROUSSEAU, 1854, p. 116 (*type*, *P. magellanica*; M); *Gaimarda* GRAY, 1855, p. 108; *Phascolicama* GRAY, 1855, p. 108]. Trapezoid, smooth, anteriorly rostrate, up to 32 mm. long, generally gaping anteroventrally. Teeth commonly reduced; anterior and posterior accessory marginal teeth in some species. *L.Pleist.-Rec.*, N.Z.-Patagonia-Subantarct. Is.—FIG. E126,4. **G. trapesina* (LAMARCK), *Rec.*, Magellan Strait; *4a-c*, RV ext., LV int., RV int., $\times 3$ (C. A. Fleming, n).

Costokidderia FINLAY, 1927, p. 457 [**Kidderia costata* ODHNER, 1924, p. 68; OD]. Resembling *Kidderia* but with strong radial ribs on median and posterior parts of shell, imbricated by growth striae and crenulating margins. *U.Pleist.-Rec.*, N.Z.—FIG. E126,2. **C. costata* (ODHNER), *Rec.*, Auckland Is.; *2a,b*, LV ext., RV int., $\times 9$ (685).

?*Eugaimardia* COTTON, 1931, p. 63 [*pro* *Neogaimardia* COTTON, 1931, p. 341 (*non* ODHNER, 1924, p. 64)] [**Neogaimardia perplexa* COTTON, 1931, p. 341; M]. Similar to *Neogaimardia* but lacking ventral sinuosity and anterior rostrum; hinge more robust, veneriform; ligament external. [Systematic position doubtful.] *Rec.*, S.Australia.—FIG. E126,3. **E. perplexa* (COTTON); *3a,b*, RV ext., hinge, $\times 9$ (B. C. Cotton).

Kidderia DALL, 1876, p. 46 [**K. minuta*; M]. Small, solid, elongate-oval, lacking ventral sinuosity and gape of *Gaimardia*, with sunken subinternal ligament; some species not rostrate. *Mio.-Rec.*, N.Z.-Fuegia-Subantarctic Is.—FIG. E126,1. **K. minuta*, *Rec.*, Kerguelen; *1a-c*, RV ext., RV int., LV int., $\times 9$ (C. A. Fleming, n).

Neogaimardia ODHNER, 1924, p. 64 [**Kellia rostellata* TATE, 1889, p. 63; OD]. Small, rostrate, with anteroventral sinus and gape; ligament internal; right cardinal embraced by hooklike left cardinal; accessory marginal teeth present. *L.Pleist.-Rec.*, N.Z.-S.Australia.—FIG. E126,5. **N. rostellata* (TATE), *Rec.*, Victoria; *5a,b*, LV and RV hinges showing accessory marginal teeth (*a-c*, *a₁*, *b₁*) and teeth (Bernard's notation), $\times 9$ (685).

Superfamily ARCTICACEA Newton, 1891

[*nom. transl.* HABE, 1951 (*ex* Arcticidae NEWTON, 1891)] [= *Trapezacea* HABE, 1951] [Materials for this superfamily prepared by MYRA KEEN with additions as recorded]




FIG. E126. Gaimardiidae (p. N644).

Shells inequilateral, mostly equivalve, beaks well forward, spirally twisted only in a few; surface smooth or with some concentric (rarely any radial) ribbing; ligament external; hinge of the form termed cyprinoid by authors, having two or three cardinal teeth in each valve and well-developed laterals in most; teeth tending to radiate from beaks; pallial line normally entire (sinuate in few). *M.Dev.-Rec.*

Family ARCTICIDAE Newton, 1891

[=Cyprinidae D'ORBIGNY, 1844 (non BONAPARTE, 1841); Venilidae DALL, 1889; Veniellinae DALL, 1895] [Family-group

names based on *Cyprina* LAMARCK, 1818 and *Venilia* MORTON, 1833 are invalid (Code, Art. 11e)] [Materials for this family prepared by MYRA KEEN and RAYMOND CASEY]

Equivalve, inequilateral, closed; shell evenly inflated or with lateral carinae; sculpture, if present, mostly of fine concentric riblets (radial threads in a few); ligament on nymphs, insertion grooves incised; hinge formula=*AI AIII 3a 1 3b 5b PI PIII/AII 2a 2b 4b PII PIV*, not all teeth fully developed, especially *AII 1 5b, PI PIII 2a, PII* and *PIV*; valve margins smooth within or feebly crenulate; adductor muscle scars subequal; pallial line entire or slightly sinuate. *U.Trias.-Rec.*

- Arctica** SCHUMACHER, 1817 [**A. vulgaris* (= **Venus islandica* LINNÉ, 1767); M] [Not preoccupied by *Arctica* MÖHRING, 1758 (ICZN, rejected work)] [= *Cyprina* LAMARCK, 1818 (obj.; SD CHILDREN, 1823); *Armida* GISTEL, 1848 (non RISSO, 1826) (obj.); *Asmidia* (nom. null.); *Nympha* MÖRCH, 1853 (p. 36, in synonym.) (obj.) (non MÖRCH, 1853, p. 25); *Cypriniadea* ROVERETO, 1900 (nom. van., pro *Cyprina*) (obj.)]. Solid, smooth, ovate, periostracum well developed; no lunule or escutcheon; hinge formula, *AI AIII 3a 1 3b PI/AII 2a 2b 4b PII*, with *PI* elongate, its edge striate, *PII* merged into margin, *2a* scarcely differentiated from *AII*, well separated from *2b*, *1* small, tubercular, *3b* broad, bifid, *AI* and *AII* short, crenulate. *L.Cret. (Alb.)-Rec.*, Eu.-N.Atl.-N.Am.—FIG. E127,1. **A. islandica* (LINNÉ), Plio., Eng.; *1a,b*, RV ext., int., $\times 0.3$ (Wood, 1853).
- Agapella** VOKES, 1946 [**A. rotunda*; OD]. Smooth, subcircular to subovate, with prominent umbones; hinge with narrow *3b* and large hook-shaped *2b*; evidence as to posterior lateral teeth and pallial sinus incomplete. *L.Cret. (Apt.)*, SW.Asia.—FIG. E127,5. **A. rotunda*, Syria; *5a,b*, RV and LV hinges, $\times 1.5$ (after 945).
- Anisocardia** MUNIER-CHALMAS, 1863 [**A. elegans*; M] [= *Apocardia* DOLLFUS, 1863 (obj.); *Cardiodonta* LAUBE, 1867 (ex STOLICZKA MS) (type, *C. balinensis* (?M); SD COX, 1947)]. Ovate to subtrigonal or trapezoidal; superficial lunule present in some but no escutcheon; hinge formula *AI 1 3a 3b PI/AII 2a 2b PII*, with *3b* bifid, *1* strongly projecting, *2b* chevron-shaped, *2a* not well differentiated from *AI*, *PI* strong, elongate, *PII* merged into margin; pallial line posteriorly truncate. *M. Jur. (Bajoc.)-L.Cret. (Apt.)*, Eu.-Afr.
- A. (Anisocardia)**. Subtrigonal, posterior end more or less truncate or rostrate, umbones prominent, beaks strongly prosogyrate; surface with radial threads; inner ventral margin crenulate; anterior lateral teeth short. *M. Jur. (Bajoc.)-L.Cret. (Apt.)*, Eu.-E.Afr.—FIG. E127,3. **A. (A.) elegans* MUNIER-CHALMAS, U.Jur. (Kimmeridg.), France; *3a*, LV ext., $\times 0.7$ (1026); *3b,c*, LV and RV hinges, $\times 1$ (Cox, 1947).
- A. (Antiquicyprina)** CASEY, 1952 [**Cyprina loweana* MORRIS & LYCETT, 1854; OD]. Ovate, with trapezoidal tendency and obtuse posterior carina; hinge with anterior lateral teeth longer than in *A. (Anisocardia)*, tooth *1* less prominent. *M. Jur. (Bathon.)*, Eu.—FIG. E127,9. **C. sarthacensis* (MORRIS & LYCETT), Eng.; *2a*, LV ext., $\times 0.7$; *2b,c*, RV and LV hinges, $\times 0.7$ (92).
- A. (Collignonicardia)** MAHMOUD, 1955 [**A. (C.) simplex*; OD]. Sculpture of fine, serrate concentric riblets; shell trigonal, with blunt posterior carina; hinge unknown. *Cret.*, Egypt.
- Ceolocyprina** DOUVILLÉ, 1921 [**C. sarthacensis*; OD]. Trigonal-ovate, globose, with faint radial striae, inner ventral margin crenulate; lunule deeply sunken, inhibiting development of anterior hinge teeth; posterior lateral teeth not observed. *M. Jur. (Bathon.)*, Eu.—FIG. E127,9. **C. sarthacensis*, France; LV int., $\times 0.7$ (after Douvillé, 1921).
- Dietrichia** RECK, 1921 [**D. parvula*; OD]. Ovate, smooth, evenly inflated; lunule deep; dentition imperfectly known, posterior lateral teeth apparently wanting. *Jur.*, E.Afr.—FIG. E127,6. **D. parvula*; *6a,b*, RV int., both valves ant., $\times 2$ (Reck).
- Epicypriina** CASEY, 1952 [**Venus angulata* J. SOWERBY, 1814; OD]. Ovate, with trapezoidal tendency and obtuse posterior ridge; escutcheon deep, narrow, limited by sharp carinae; hinge formula, *AI (III) 1 3a 3b PI/AII 2a 2b 4b PII*, with *1* strong, conical, anterior to *3a*, anterior laterals rugose, nymphs with interlocking rugosities; pallial line simple. *L. Cret. (Apt.-Alb.)*, Eu.-Afr.-W. Indies-S. Am.—FIG. E127,8. **E. angulata* (SOWERBY), Alb., Eng.; *8a*, LV ext., $\times 0.3$; *8b,c*, RV and LV hinges, $\times 1$ (92).
- Etea** CONRAD, 1875 [**E. carolinensis*; M]. Elongate-subovate, tapering and truncate posteriorly, umbones well forward, posterior ridge strong, posterior area flat or concave; no lunule or escutcheon; hinge longer, narrower, less massive and less arched than in *Veniella*, with smooth lateral teeth. *U. Cret.*, N.Am.—FIG. E127,4. **E. carolinensis*, USA (N.Car.); *4a,b*, LV ext., RV int., $\times 1$ (Stephenson).
- ?**Fissilunula** ETHERIDGE, 1902 [**Cytherea clarkei* MOORE, 1870; M]. Large, massive, beaks prosogyrate; lunule with median sulcus; hinge with *1* cardinal in RV; pallial sinus present, small. *U. Cret.*, Australia.—FIG. E127,11. **F. clarkei* (MOORE), New S. Wales; *11a,b*, RV hinge, RV ext., $\times 0.3$, $\times 0.2$ (Etheridge, 1902).
- Hartwellia** KITCHIN, 1926 [**Astarte hartwellensis* J. DE C. SOWERBY, 1845; M] [= *Adalantia* SEELEY, 1864 (non MEIGEN, 1800) (obj.)]. Ovate or trapezoidal, posterior slope flattened, limited by ridge; beaks small and pointed; lunular area excavated, bounded by overhanging ridges; escutcheon deep, with carinate edges; sculpture *Astarte*-like in young; hinge formula, *AI AIII 1 3a 3b PIII/AII 2a 2b 4b PII*, with laterals well developed, cross-striate, *3a* and *3b* united by tuberosity, nymphs with rugose callosities; pallial line feebly sinuate. *U. Jur.-L. Cret.*, Eu.-N.Asia-N.Atl.
- H. (Hartwellia)**. Ovate-trapezoidal, posterior ridge distinct; hinge with teeth *1* and *2b* pyramidal. *U. Jur. (Kimmeridg.)-L. Cret. (Neocom.)*, NW.Eu.-N.Asia-N.Atl.—FIG. E127,10. **H. (H.) hartwellensis* (SOWERBY), Kimmeridg., Eng.; *10a-c*, LV ext., LV and RV hinges, $\times 0.5$ (Casey, 1952).
- H. (Tealbya)** CASEY, 1952 [**Cyprina tealbiensis* WOODS, 1907; OD]. Externally like *H. (Hartwellia)* in young but becoming trigonal-ovate with ill-defined posterior ridge. *L. Cret. (Neocom.)*,

Eu.—FIG. E127,7. **H. (T.) tealbiensis* (Woods), Eng.; 7a,b, LV int., RV hinge, $\times 0.7$ (Casey).

BUVIGNIER, 1852; SD COSSMANN, 1921]. Suborbicular or ovate, evenly inflated or with feeble posterior carina; lunule superficial, mostly bounded by impressed line; escutcheon wanting; surface

Isocyprina RÖDER, 1882 [**Cardium cyreniforme*




FIG. E127. Arcticidae (p. N646-N647).

smooth or with fine concentric lines; hinge formula, *AI(III) (1) (3a) 3b PI/AII (2a) 2b 4b (PII)* (teeth indicated in parentheses reduced or obsolete), lamina *AII (+2a)* joined to *2b* by vinculum, *PII* merged into margin, *PI* strong, elongate. *U. Trias.-U. Jur.*, Eu.-S.Am.

I. (Isocyprina). Hinge with tooth *2a* differentiated from *All. Jur.* (*Lias.-Oxford*), Eu.—FIG. E128, 2. **I. (I.) cyreniformis* (BUVIGNIER), *U. Jur.* (*Oxford*), Eng.; *2a-c*, LV ext., LV and RV hinges, $\times 0.7$ (Casey).

I. (Eotrapezium) DOUVILLÉ, 1913 [*Mesodesma germari* DUNKER, 1844; OD] [= *Raetolucina* OSSWALD, 1929 (type, *Corbula alpina* WINKLER, 1859; M)]. Suborbicular or elongate-ovate; hinge with cardinal tooth *1* scarcely differentiated from *AI*, no *2a*, lamina *AIII-3a* suppressed in most. *U. Trias.-L. Jur.* (*Lias.*), Eu.-S.Am.—FIG. E128, 1. **I. (E.) germari* (DUNKER), *L. Jur.*, Ger.; *1a,b*, LV and RV hinges, $\times 0.7$ (Cox).

I. (Venericyprina) CASEY, 1952 [**I. (V.) argillacea*; OD]. Externally like *I. (Isocyprina)*, posterior end elongate or rostrate in some; hinge with lateral teeth cross-striate, lamina *AIII-3a* well developed, *2a* distinct at end of *AII*. *U. Jur.* (*Kimmeridg.*)-*L. Cret.* (*Apt.*), Eu.—FIG. E128, 3. **I. (V.) argillacea*, *U. Jur.*, Eng.; *3a,b*, RV int., LV hinge, $\times 2$ (Casey).

?Izumia ICHIKAWA & MAEDA, 1963 [**I. trapezoidalis*; OD]. Outline subtrapezoidal, surface nearly smooth but with fine radial lirae beneath outermost layer, causing crenulation of inner margins; hinge plate well developed, cardinal teeth veneroid in appearance, lateral teeth distinct; pallial line not exposed in type material. *U. Cret.*, E.Asia.—FIG. E128, 14. **I. trapezoidalis*, Japan; *4a,b*, LV int., RV int., $\times 2$ (Ichikawa & Maeda).

?Loparia OPPENHEIM, 1901 [**L. katzeri*; M]. Small, obliquely trapezoidal, with angulation setting off posterior slope; surface smooth except for some thick, elevated, distant concentric ridges; hinge unknown. *Eoc.*, Eu.—FIG. E128, 6. **L. katzeri*, Bosnia; RV ext., $\times 3$ (Oppenheim).

Microcyprina COSSMANN, 1921 [**Cyprina (M.) newvillei*; OD]. Outline cordiform, surface smooth; hinge as in *Arctica* but *AI* not crenulate, *1* bifid, *2a*, *2b* thin. *Eoc.*, Eu.—FIG. E128, 4. **M. newvillei* (COSSMANN), France; *4a,b*, RV ext., int., $\times 1$ (Cossmann, 1921).

?Mokattamia MAYER, 1890 [**M. agassizi*; M]. Rounded-ovate, resembling Mesozoic genus *Ptychomya* but shorter, with larger posterior muscle scars. *Eoc.*, Egypt. [Unfigured.]

Petalocardia VINCENT, 1925 [**Venus pectinifera* SOWERBY, 1823; SD GLIBERT, 1936]. Small, with distant spaced concentric lamellae and fine radial riblets, inner margin crenulate; hinge as in *Arctica* but with only 2 cardinal teeth in either valve. *Eoc.-Oligo.*, Eu.—FIG. E128, 5. **P. pectini-*

fera (SOWERBY), *Eoc.*, France; *5a,b*, RV int., LV ext., $\times 4$ (Glibert, 1936).

Plesiocyprina FISCHER, 1887 (ex MUNIER-CHALMAS MS) [**P. gaudryi*; OD]. Trapezoidal, sharply carinate; surface smooth; hinge as in *Isocyprina* (*Eotrapezium*). *U. Trias.* (*Rhaet.*)-*U. Jur.* (*Callov.*), Eu.-E.S.Am.—FIG. E128, 7. **P. gaudryi*, *U. Jur.*, France; *7a,b*, LV int., RV int., $\times 1.7$ (Fischer).

Procyprina CASEY, 1952 [**P. venusta*; OD]. Trigonal-ovate, posterior end more or less truncate; hinge as in *Arctica* but *3b* narrow, anterior laterals longer. *U. Jur.* (*Oxford.*)-*L. Cret.* (*Valangin.*), Eu.—FIG. E128, 8. **P. venusta*, *L. Cret.* (Neocom.), Eng.; *8a*, LV ext., $\times 0.7$; *8b,c*, LV and RV hinges, $\times 2$ (Casey).

Pronoella FISCHER, 1887 [*pro Pronoe* AGASSIZ, 1843 (non GUÉRIN, 1838)] [**Venulites trigonellaris* VON SCHLOTHEIM, 1820, AUCTT.; ?M]. Ovate, trigonal, or trapezoidal, evenly inflated or with posterior carina; lunule present, escutcheon absent or poorly defined; hinge formula, *AI AIII 1 3a 3b PI/AII 2a 2b 4b (PII)*, with *PII* represented only by a thickened or projected shell margin; pallial line simple or with shallow sinus. *Jur.*, Eu.

P. (Pronoella). Anterior lateral teeth strong; *1* and *2a* stoutly triangular; *2a* laminar or wedge-shaped. *Jur.* (*Lias.-Portland.*), Eu.—FIG. E128, 9. **P. (P.) trigonellaris* (VON SCHLOTHEIM), Ger.; *9a*, RV ext., $\times 0.5$ (Casey); *9b,c*, LV and RV hinges, $\times 0.7$ (Cox, 1947).

P. (Gythemon) CASEY, 1952 [**P. elongata* Cox, 1944; OD]. Elongate, beaks subterminal; lunule deeply impressed; anterior lateral teeth short; hinge plate deep, with narrow, wedge-shaped cardinals. *M. Jur.* (*Bajoc.*), Eu.—FIG. E128, 10. **P. (G.) elongata* Cox, Eng.; *10a*, RV ext., $\times 0.5$; *10b,c*, RV and LV hinges, $\times 0.7$ (Casey).

Proveniella CASEY, 1952 [**Cyprina meyeri* Woods, 1913; OD]. Ovate-trapezoidal, umbones moderately to strongly prominent; hinge formula as in *Veniella*, lateral teeth cross-striate, *1* co-laminar with *AI*; pallial line entire. *Cret.*, Eu.—FIG. E128, 12. **P. meyeri* (Woods), *L. Cret.* (*Apt.*), Eng.; *12a*, LV ext., $\times 0.5$; *12b,c*, RV and LV hinges, $\times 1.3$ (Casey).

Pseudotrapezium FISCHER, 1887 [**Cypricardia bathonica* D'ORBIGNY, 1850 (= **C. cordiformis* DESHAYES, 1830); M]. Subtrigonal to cuneiform, umbones prominent, beaks strongly prosogyrate, posterior carination strong; hinge with anterior laterals short, *1* strong, restricting development of *2a* in LV; *3a* and *3b* united into a single angulate structure. *Jur.* (*Lias.-Portland.*), Eu.—FIG. E128, 13. **P. cordiforme* (DESHAYES), *M. Jur.* (*Bathon.*), Eng.; *13a*, RV ext., $\times 0.7$; *13b,c*, hinges, $\times 0.5$ (Casey).

Pygocardia FISCHER, 1887 [**Cyprina tumida* NYST, 1835 (= **Venus rustica* SOWERBY, 1818); M]. Globose to quadrate; hinge with anterior lateral teeth stronger than in *Arctica*, anterior cardinal

weak, posterior laterals long, curved, *Oligo-Plio.*, Eu.—FIG. E128, 11. **P. rustica* (SOWERBY), Plio., Eng.; 11a-c, RV ext., LV int., RV int., $\times 0.5$ (Wood).

Rollierella COSSMANN, 1924 [*pro Rollierella* COSSMANN, 1923 and COSSMANN, 1924 (*non* COSSMANN, 1920)] [**Isocardia laubei* ROLLIER, 1913 (*pro I. cordata* LAUBE, 1867, *non* BUCKMAN, 1845); OD].




FIG. E128. Arcticidae (p. N648-N650).

Orbicular, gibbous, evenly inflated; hinge as in *Pseudotrapezium*. *M. Jur. (Bathon.)*, Eu.—FIG. E128,15. **R. laubei* (ROLLIER), Ger.; 15a,b, RV ext., int., $\times 1$ (Laube).

Schedotrapezium STEWART, 1930 [**Trapezium carinatum* GABB, 1864; OD]. Trapezoidal, umbones well forward, posterior ridge strong; lunule and escutcheon wanting; hinge in RV with slender 3b and laminar 3a diverging at obtuse angle from below beak, 1 laminar, more or less parallel with 3a, dentition of LV unknown. *U.Cret.*, N.Am.-?W.Afr.-?Eu.—FIG. E129,1. **S. carinatum* (GABB), USA (Calif.); 1a, RV ext., $\times 3$; 1b, RV hinge, $\times 8$ (Stewart).

Somarctica TAMURA, 1960 [**Arctica* (*S.*) *abukumensis*; OD]. Exterior as in *Arctica*; hinge with 2a present, 3a small. *U. Jur.*, E.Asia.—FIG. E129, 3. **S. abukumensis* (TAMURA), Japan; LV hinge, $\times 0.7$ (Tamura).

Staffinella CASEY, 1952 [**Protomiodon staffinensis* ANDERSON & COX, 1948; OD]. Trigonal-ovate, umbones not prominent, lunule and escutcheon wanting; hinge like that of *Hartwellia* (*s.s.*), but with teeth 1 and 2a poorly differentiated from laterals; pallial line truncate posteriorly. *M. Jur. (Bathon.)*, Eu.—FIG. E129,4. **S. staffinensis* (ANDERSON & COX), Scot.; 4a, LV ext., $\times 0.7$; 4b,c, RV int., LV hinge, $\times 1.7$ (Casey).

Tenea CONRAD, 1870 [**Mysia parilis* CONRAD, 1860; M]. Subcircular to subovate; umbones produced; hinge with 3b broad, deeply bifid, 2b chevron-shaped, 1 and 2a attached to vestigial *Al* and *All*, no posterior laterals; pallial sinus narrow, deep. *U.Cret.*, N.Am.—FIG. E129,7. **T. parilis* (CONRAD), USA (Tex.); 7a, LV ext., $\times 1$; 7b,c, RV and LV hinges, $\times 2$ (Stephenson).

Tortarticia CASEY, 1961 [**Isocardia similis* J. DE C. SOWERBY, 1826; OD]. Trigonal-ovate, well inflated, beaks prominent, spirally enrolled, lunular area depressed, escutcheon limited by blunt carinae; hinge with *All* pustular, 1 spoon-shaped, 3b bifid, united with 3a into single curved strongly opisthoclinal tooth lying almost horizontally, 2a and 2b nodular, separated by constriction, posterior laterals close behind nymph. *L.Cret. (Alb.)*, Eu.—FIG. E129,5. *T. similis* (SOWERBY), France; RV hinge, $\times 0.7$ (Casey).

Vectianella CASEY, 1952 [**Tellina vectiana* FORBES, 1845; OD]. Small, trigonal-ovate, tapering posteriorly, compressed, umbones subcentral, no lunule, escutcheon long and narrow, bounded by obtuse ridges; sculpture concrescent, finely lineate; hinge with 1 and 3a large, laminar, 3b minute, 2b stoutly triangular, bifid, anterior laterals obsolescent. *L.Cret. (Apt.)*, NW.Eu.—FIG. E129,2. **V. vectiana* (FORBES), Eng.; 2a, LV ext., $\times 0.7$; 2b,c, LV and RV hinges, $\times 1$ (Casey).

Veniella STOLICZKA, 1870 [*pro Venilia* MORTON, 1833 (*non* DUPONCHEL, 1829)] [**Venilia conradi* MORTON, 1833; M] [= *Goniosoma* CONRAD, 1869 (*non* PERTY, 1833) (type, *G. inflata*; M); *Cicatreia*

STOLICZKA, 1870 (type, *Cyprina cordialis*; OD); *Roudairia* MUNIER-CHALMAS, 1881 (type, *R. drui*; M) (*Roudaireia*, *nom. null.*); *Trigonocardia* ZITTEL, 1881 (in synonymy of *Roudairia*); *Platopsis* WHITFIELD, 1891 (type, *Opis undata* CONRAD, 1852; SD WOODWARD, 1892) (*Platopsis*, *nom. null.*); *Venetia*, *Venniella*, *nom. null.*]. Subtrapezoidal and moderately ventricose in young, obliquely subtrigonal and strongly ventricose in adult, umbones prominent, beaks prosogyrate, posterior carina strong; surface with distant thickened concentric lamellae; hinge massive, laterals cross-striate, 3a and 3b united in hook-shaped structure inserted posterior to tubercular tooth 1; pallial line truncate posteriorly. *U.Cret.*, Asia-Afr.-W. Indies-N.Am.-S.Am.—FIG. E129,6. **V. conradi* (MORTON), USA (Miss.); 6a,b, LV int., RV hinge, $\times 1$ (Casey, n).

Venilicardia STOLICZKA, 1870 [**Cyprina bifida* ZITTEL, 1865; OD]. Like *Arctica* externally; hinge as in *Epicypriina* but with 3a and 3b united to form hook-shaped structure, 1 strong, tubercular, placed below 3a, nymphs and lateral teeth smooth. *Cret.*, Eu.-Afr.-S. Asia-S. Am.—FIG. E129,8. **V. bifida* (ZITTEL), Ger.; 8a,b, RV ext., int., $\times 0.5$ (Zittel).

Family BERNARDINIDAE Keen, 1963

[Materials for this family prepared by MYRA KEEN]

Small to minute shells with ligament partially to entirely internal; hinge with two or three cardinal teeth and two or more laterals; pallial line entire. *Rec.*

Bernardina DALL, 1910 [**B. bakeri*; OD]. Centrally sculptured, thin, prodissoconch set off by ridge in some; posterior dorsal margin of RV fitting into shallow groove in LV; hinge with 2 cardinal teeth in RV, 3 in LV, anterior lateral strong in LV, 2 flexuous laterals in RV; resilium in pit behind cardinals. *Rec.*, W.N.Am.-C.Am.—FIG. E130,10. **B. bakeri*, USA (Calif.); 10a-c, RV ext., int., LV int., $\times 12$ (specimen, Stanford coll.).

Halodakra OLSSON, 1961 [**Circe subtrigona* CARPENTER, 1857; OD]. Ovate, inequilateral, smooth or with concentric striae; posterior area with chevron-shaped color markings in some; ligament in shallow resilifer, somewhat sunken; hinge with 3 or more teeth in cardinal area, posterior lateral behind resilifer more or less well developed; posterior margins of shell grooved to receive opposite valve; pallial line indistinct, apparently entire. *Rec.*, W.N.Am.-S.Am.—FIG. E130,7. **H. subtrigona* (CARPENTER), Ecuador; 7a,b, LV and RV hinges, $\times 13$; 7c, LV ext., $\times 7$ (688).

Family EULOXIDAE Gardner, 1943

[Materials for this family prepared by MYRA KEEN]

Ovate, inequilateral, not elongate; hinge




FIG. E129. Arcticidae (p. N650).

with two or three cardinal teeth, no posterior laterals; pallial line entire. *Mio.*

Euloxa CONRAD, 1863 [**Venus latisulcata* CONRAD, 1840; M]. Sculpture of concentric undulations; posterior slope set off by radial ridge; hinge with 3 cardinals in LV, 2 in RV. *Mio.*, E.N.Am.—FIG. E130,6. **E. latisulcata* (CONRAD), USA (Va.); 6a-c, LV ext., RV int., LV int., $\times 1$ (Nicol, 1953). *Cabralista* KEEN, *nom. subst.* herein [*pro Cabralia* BÖHM, 1899 (*non* MOORE, 1886)] [**Cabralia schmitzi* BÖHM, 1899; M]. Ovate to trapezoidal, with both radial and concentric sculpture; hinge

with 3 cardinals in either valve, no lateral teeth; pallial line entire. *Mio.*, E.Atl.—FIG. E130,9. **Cabralista schmitzi* (BÖHM), Salvages Is.; 9a-c, LV ext., LV and RV hinges, $\times 1$ (Böhm, 1899).

Family KELLIPELLIDAE Fischer, 1887

[*nom. correct.* DALL, 1900 (*pro Kellyellidae* FISCHER, 1887)]
[Materials for this family prepared by MYRA KEEN]

Small to minute, equivalve, mostly sub-orbicular; not gaping; ligament mostly external, some with resilium or fossette under beak; hinge incompletely developed, anterior

lamellae adjacent to cardinals, posterior lamellae not constant; inner margins smooth; pallial line entire (854). *Tert.-Rec.*

Kelliella M. Sars, 1870 [**K. abyssicola* (= **Venus miliaris* PHILIPPI, 1844); M] [= *Kelliella* M. Sars, 1865, 1868, *AUCT.* (nom. nud.); *Kellyella*




FIG. E130. Bernardinidae (7,10); Euloxidae (6,9); Kelliellidae (1-5,8,11-12) (p. N650-N653).

- FISCHER, 1887 (*nom. null.*). Minute, rounded-ovate; lunule circumscribed; hinge with 2 teeth in LV, 1 cardinal and anterior lateral in RV. *Tert.-Rec.*, Atl.-Medit. [Deep-water.]. — FIG. E130,4. **K. miliaris* (PHILIPPI), *Rec.*, Norway; 4a, LV ext., $\times 6$; 4b,c, RV and LV hinges, enl. (829).
- Allopagus** STOLICZKA, 1871 [**Hippagus leanus* DESHAYES, 1860; OD]. Minute, thin, ovate, very inequilateral, tumid, beaks small; hinge of RV with 1 cardinal in front of beak, LV with cardinal below. *M.Eoc.*, Eu.—FIG. E130,12. **A. leanus* (DESHAYES), France; 12a,b, RV ext., int., $\times 10$ (259).
- Alveinus** CONRAD, 1865 [**A. minutus*; M]. Smooth; ligamental nymph feeble; resilium in cavity under beaks; hinge of RV with 2 parallel teeth; LV with single tooth bent or A-shaped; posterior margin of RV grooved. *U.Eoc.-Mio.*, Eu.-N.Am.—FIG. E130,11. **A. minutus*, *Eoc.*, USA (La.-Miss.); 11a-c, LV ext., int., RV int., $\times 14$ (388).
- ?**Eocrassina** COSSMANN, 1914 [**Parisiella veneriformis* COSSMANN, 1907; OD]. Resembling *Lutetia* but nymph larger, hinge teeth more regular, 1 and 3b in RV, 2a, 2b, and PII in LV; margin even, not grooved; muscle scars large, high in shell. *U.Eoc.*, Eu.—FIG. E130,5. **E. veneriformis* (COSSMANN), France; 5a,b, LV int., RV int., $\times 10$ (Cossmann, 1907).
- Lutetia** DESHAYES, 1860 [**L. parisiensis*; SD STOLICZKA, 1871]. Globose, ligament external, on nymph, resilium wanting; adductor muscle scars small, equal, oval. *L.Eoc.-Mio.*, Eu.-N.Am.-Asia. **L. (Lutetia)**. Hinge with 3 teeth in either valve, RV with posterior tooth nearly marginal, central tooth L-shaped, anterior tooth deflected downward, teeth in LV laminar; margin of RV grooved. *L.Eoc.-Oligo.*, Eu.-N.Am.—FIG. E130,1. **L. (L.) parisiensis* DESHAYES, *Eoc.*, France; 1a-c, LV ext., int., RV hinge, $\times 10$ (Deshayes, 1858).
- L. (Spaniodontella)** ANDRUSSOV, in GOLUBIATNIKOV, 1902 [*pro Spaniodon* REUSS, 1867 (*non* PICTET, 1851)] [**Spaniodon nitidus* REUSS, 1867; M] [= *Davidaschwilia* MERKLIN, 1950 (type, *Spaniodontella intermedia* "ANDRUSSOV" GOLUBIATNIKOV, 1902; OD)]. Slightly larger than *L. (Lutetia)*, hinge teeth relatively larger, especially central teeth; hinge plate wider. *Mio.*, Eu.-SW.Asia.—FIG. E130,2. **L. (S.) nitida* (REUSS), Ger.; 2a,b, RV int., LV int., $\times 7$ (Reuss, 1867).
- Pauliella** MUNIER-CHALMAS, 1895 [**P. bernardi*; M]. Ovate, sculpture concentric; hinge with 3 cardinal teeth in either valve, 2 posterior laterals, 3 anterior lateral laminae. *Rec.*, Ind.O.—FIG. E130,8. **P. bernardi*, St. Paul I.; 8a, RV ext., $\times 5$; 8b, RV hinge, $\times 10$ (44).
- Warrana** LASERON, 1953 [**W. dielasma*; OD]. High-ovate, with concentric sculpture; prodissoconch distinct; ligament internal; hinge of LV with thin arched cardinal and linear obscure laterals. *Rec.*, Australia.—FIG. E130,3. **W. dielasma*, SE. Australia; 3a,b, RV ext., LV int., $\times 18$ (531).

Family NEOMIODONTIDAE Casey, 1955

[Materials for this family prepared by RAYMOND CASEY]

Shells subtrigonal to ovate, smooth, inner margins smooth; hinge lucinoid, formula: *AI AIII 3a 3b 5b PI PIII/AII 2b 4b PII*, with cardinal teeth entire, laterals typically long and corbiculoid, teeth 5b, PIII not always developed. [Habitat, fresh-water or seas of reduced salinity.] *L.Jur.-U.Cret.*

Neomiodon FISCHER, 1887 [*pro Miodon* SANDBERGER, 1871 (*non* DUMÉRIL, 1859)] [**Cyclas medius* J. DE C. SOWERBY, 1826; SD DALL, 1903] [= *Bidentina* OPPENHEIM, 1895 (obj.; *pro Miodon*); *Miodontopsis* DALL, 1903; *Protomiodon* ANDERSON & COX, 1948 (type, *Cyrena brycei* TATE, 1873; OD)]. Subtrigonal, ovate to suborbicular, with no definite lunule or escutcheon; smooth or with weak concentric sculpture, posterior slope more or less ridged; hinge formula, *AI AIII 3a 3b PI PIII/AII 2b 4b PII*, laterals minutely cross-striate; pallial line entire. *M.Jur.(Bathon.)-L.Cret.(Weald.)*, Eurasia.—FIG. E131,8. **N. medius* (SOWERBY), U.Jur.(Purbeck.), Eng.; 8a-d, LV ext., RV int., LV hinge, LV dorsal, $\times 1.5$ (Casey).

Crenotrapezium HAYAMI, 1958 [**C. kurumense*; OD]. Externally resembling a cuneiform *Neomiodon* but with hinge of *Eomiodon*. *L.Jur.(Lias.)*, ?*L.Cret.(Weald.)*, E.Asia.—FIG. E131,1. **C. kurumense*, Lias, Japan; 1a-c, LV ext., dorsal hinge, $\times 1$; 1d, RV hinge, $\times 2$ (Hayami).

Cyrenopsis ETHERIDGE, 1902 [**Macra meeki* = *Unicardium meeki* ETHERIDGE, 1892; OD]. Subtrigonal to suborbicular, evenly inflated; no lunule, escutcheon long and deep; sculpture concentric; hinge-formula, *AI AIII 3a 3b PI/AII 2b 4b PII*, with smooth laterals; pallial line entire. *L.Cret.(Apt.)*, Australia.—FIG. E131,3. **C. meeki* (ETHERIDGE); 3a,b, RV ext., LV int., $\times 0.7$ (Jack and Etheridge).

Eomiodon COX, 1935 [**E. indicus*; OD] [= *Protocyprina* VOKES, 1946 (type, *Astarte libanotica* FRAAS, 1878; OD)]. Gibbous-trigonal, cuneiform, or suborbicular, lunule and escutcheon impressed, smooth, lunule limited by furrow, escutcheon by ridge; sculpture concrescent *Astarte*-like in young; hinge as in *Neomiodon* except for presence of small 5b and absence of PIII; pallial line truncate posteriorly. *L.Jur.(Lias.)-U.Cret.(?Cenoman.)*, Eu.-N.Afr.-Asia-N.Am.—FIG. E131,6. **E. indicus*, M.Jur., India; 6a,b, LV and RV hinges, $\times 1$ (Cox).

Musculiopsis MACNEIL, 1939 [**M. russelli*; OD].

Subrectangular, posteriorly truncate; lunular area excavated, escutcheon wanting; young with posterior carina; surface nearly smooth; hinge of LV with strong *2b*, slender *4b*, short *All* and long *Pll*

addressed to margin, dentition of RV imperfectly known; pallial line apparently entire. *L.Cret.*, N. Am.—FIG. E131,7. **M. russelli*, USA (Nev.); 7a, RV ext., $\times 2$; 7b, LV int., $\times 3$ (MacNeil).




FIG. E131. Neomiodontidae (1,3-4,6-8); Pollicidae (2,9); Ptychomyidae (5) (p. N653-N655).

Myrene CASEY, 1955 [**M. fittoni*; OD]. Cuneiform, trigonal-ovate, or trapezoidal; lunule and escutcheon not clearly defined; posterior slope ridged or carinate, posterior area flattened, with or without second ridge or carina; surface smooth or with concentric riblets; hinge formula, *AI AIII 3a 3b 5b PI/AO AII 2b 4b PO PII*, laterals cross-striate in most; pallial line entire. *U. Jur. (Oxford.)-L. Cret. (Ryaz.)*, Eu.-NE. Asia.—FIG. E131,4. **M. fittoni*, L. Cret., Eng.; 4a,b, RV and LV hinges, $\times 2$ (Casey).

Family POLLICIDAE Stephenson, 1953

[Materials for this family prepared by MYRA KEEN]

Elongate, plump, subelliptical, ligament external, opisthodontic; inner margins of shell smooth, shape of muscle scars and pallial line unknown. *U. Cret.*

Pollex STEPHENSON, 1953 [**P. obesus*; OD]. Sculpture of faint radial lines, hinge with 3 cardinal teeth in either valve; anterior cardinal with 2 cusps; posterior lateral tooth long, lamellar. *U. Cret.*, N. Am.—FIG. E131,9. **P. obesus*, Cenoman., USA (Tex.); 9a, LV ext., $\times 1$; 9b, RV int., $\times 1.3$; 9c, LV int., $\times 3$ (Stephenson).

Neritra STEPHENSON, 1954 [**N. polliciformis*; OD]. Resembling *Permophorus* (Permophoridae) in form but with hinge somewhat more like that of *Pollex*, except that anterior cardinal in RV is narrower and central cardinal broader. *U. Cret.*, N. Am.—FIG. E131,2. **N. polliciformis*, USA (N.J.); 2a, RV ext., $\times 1$; 2b,c, int., LV and RV hinges, $\times 3$ (Stephenson).

Family PTYCHOMYIDAE Keen, new family

Elongate-ovate shells of solid texture, with well-developed radial sculpture; hinge of somewhat veneroid aspect but also with astartoid resemblances, the formula, *AIII 3a 1 3b PIII/AII 2a 2b 4b PII*. ?*U. Jur.*, L. Cret.-*U. Cret.*

Ptychomya AGASSIZ, 1842 [**P. plana*; M] [= *Radioconcha* CONRAD, 1869 (type, *Crassatella robinaldina* D'ORBIGNY, 1844 [= *P. plana*]; SD KEEN, herein)]. Ovate-trapezoidal; ligamentary depression broad and long; hinge with 2a-2b joined above; pallial line truncate. ?*U. Jur.*, L. Cret.-*U. Cret.*, Eu.-N. Am.-S. Am.-Afr.

P. (Ptychomya). Radial sculpture becoming divaricate on anterior and posterior slopes, rib ends showing as crenulations of inner margins. ?*U. Jur.*, L. Cret.-*U. Cret.*, Eu.-N. Am.-S. Am.-Afr.—FIG. E131,5. **P. (P.) plana* AGASSIZ, L. Cret., Eng.; 5a, LV ext., $\times 0.5$; 5b, RV hinge, $\times 1$ (Woods, 1907).

P. (Pleuroconcha) CONRAD, 1872 [**Crassatella "gallieni"* (i.e., **Crassatella galliennei* D'ORBIGNY, 1844); M]. Radial sculpture not divaricate, *U. Cret.*, Eu.

Family TRAPEZIIDAE Lamy, 1920

[= *Lithophagellidae* COSSMANN, 1910; *Libitinae* THIELE, 1924] [Materials for this family prepared by MYRA KEEN]

Elongate, beaks near anterior end; hinge plate narrowed, normally with two cardinals in either valve, one posterior and one small anterior lateral; pallial line mostly entire. ?*L. Cret.*, *U. Cret.-Rec.*

Trapezium MEGERLE VON MÜHLFELD, 1811 [**T. perfectum* (= *Chama oblonga* LINNÉ, 1758); SD STEWART, 1930] [= *Libitina* SCHUMACHER, 1817 (type, *L. bicarinata*; M); *Cypricardia* LAMARCK, 1819 (type, *C. guinaica*, = *Chama oblonga* LINNÉ; SD CHILDREN, 1823)]. Oblong, solid, ligament external. Nestling in crevices. *Eoc.-Rec.*, N. Am.-Eu.-IndoPac.-Ind. O.

T. (Trapezium). Surface with radiating striae; inner margin smooth. *Eoc.-Rec.*, E. N. Am.-Eu.-E. Indies.—FIG. E132,9. **T. (T.) oblongum* (LINNÉ), Rec., E. Indies; 9a,b, LV ext., RV int., $\times 1$ (de Blainville, 1825).

T. (Glossocardia) STOLICZKA, 1870 [**Cypricardia obesa* REEVE, 1843; OD]. More quadrate than *T. (Trapezium)*, radial striae not present; hinge teeth more strongly developed. *Rec.*, IndoPac.—FIG. E132,1. **T. (G.) obesum* (REEVE); 1a,b, RV int., LV int., $\times 0.5$ (305).

T. (Neotrapezium) HABE, 1951 [**Cardita sublaevigata* LAMARCK, 1819; OD]. Radial sculpture wanting; shell quadrangular, compressed, distorted by nestling habit; posterior lateral teeth weak and small. *Rec.*, IndoPac.—FIG. E132,2. **T. (N.) sublaevigatum* (LAMARCK), Ind. O.; 2a,b, LV ext., RV int., $\times 1.5$ (512).

Coralliophaga DE BLAINVILLE, 1824 [**C. carditoidea* (= *Chama coralliophaga* GMELIN, 1791); M] [= *Lithophagella* GRAY, 1854 (type, *Cardita dactyla* BRUGUIÈRE, 1792; OD)]. Thin, translucent, sculpture of radiating striae or granules; hinge reduced. *Eoc.-Rec.*, Eu.-W. Indies-E. Indies-N. Am.

C. (Coralliophaga). Posterior end with elevated concentric ridges; pallial line sinuate. *Eoc.-Rec.*, Eu.-W. Indies-E. Indies.—FIG. E132,6. **C. (C.) coralliophaga* (GMELIN), Rec., W. Indies; 6a-c, RV int., ext., LV hinge, $\times 1$ (7).

C. (Oryctomya) DALL, 1898 [**C. (O.) claibornensis*; OD]. Surface granular; hinge with 1 cardinal in either valve; pallial sinus distinct, angular. *Eoc.-Mio.*, E. N. Am.—FIG. E132,4. **C. (O.) claibornensis*, Eoc., USA (Ala.); 4a, LV ext., $\times 1$; 4b, enl. (223).

?**Corbiculopsis** WHITFIELD, 1891 [**C. birdi*; M]. Trapezoidal, wider posteriorly, somewhat carinate;




FIG. E132. Trapeziidae (p. N655-N657).

hinge imperfectly known, 2a and 3b bifid. *L.Cret.* (*Apt.*), SW.Asia.—FIG. E132,10. **C. birdi*, Syria; 10a-c, RV ext., RV and LV hinges, $\times 0.7$ (Whitfield).

?*Isorropodon* STURANY, 1896 [**I. perplexum*; M]. Ovate, inequivalve, LV smaller than RV; posterior area set off by radial ridge. [Deep-water.] *Rec.*, *Medit.*—FIG. E132,5. **I. perplexum*, off Egypt; 5a-c, RV ext., RV and LV hinges, $\times 2.5$ (Sturany, 1896).

Kitsonia EAMES, 1957 [**Coralliophaga eocenica* NEWTON, 1922; OD]. Thin, elongate, smooth, ligament external, long; hinge with larger teeth than in *Coralliophaga*, 2 cardinals in RV, posterior bifid, 1 lamellar cardinal in LV; adductor impres-

sions large; pallial sinus 0.4 of shell length. *Eoc.*, W.Afr.—FIG. E132,8. **K. eocenica* (NEWTON), Nigeria; 8a,b, LV ext., RV int., $\times 2.5$; 8c, LV hinge, $\times 5$ (Eames, 1957).

Pseudopleurophorus CHAVAN, 1954 [**P. rochi*; OD]. Ovate-trapezoidal, surface with growth rugae only, posterior area set off by ridge; hinge with strong posterior lateral in RV received between duplicate laterals in LV, 3a wanting. *U.Cret.*, C.Afr.—FIG. E132,7. **P. rochi*, Tchad; 7a-c, LV ext., int., RV hinge, $\times 1.5$ (Chavan).

Tortucardia OLSSON, 1944 [**Glossocardia* (*T.*) *stephensoni*; OD]. Oblong, with strong posterior carina, beaks small; escutcheon narrow; hinge with 2a, 2b, and 4b stout, subequal, *All* close to

margin, dentition of RV imperfectly known. *U. Cret.(Maastricht.)*, S.Am.—FIG. E132,3. **T. stephensoni* (OLSSON), Peru; LV hinge, $\times 1$ (Olsson).

Family MECYNODONTIDAE Haffer, 1959

[Materials for this family prepared by AURÈLE LAROCQUE]

Characters of *Mecynodon*. M.Dev.

Mecynodon KEFERSTEIN, 1857 [**Megalodus carinatus* GOLDFUSS, 1837; SD HAFFER, 1859] [= *Mecynodon* FRENCH, 1889; *Mecynodus* BEUSHAUSEN, 1895 (obj.)]. Rhomboidal, with strong postumbonal ridge; surface concentrically striate or smooth; dental formula: 1 3 1 III/2a 2p II. M.Dev., Eu.(Ger.-Eng.); M.Dev.(Onondag.), N.Am.—FIG. E133,1. **M. carinatus* (GOLDFUSS), Dev. (*Stringocephalus* Ls.), Ger.; 1a,b, RV ext., int., approx. $\times 1$ (47); 1c,d, LV and RV hinges (diagram.) (Haffer, 1959); 1e, RV int., $\times 1$ (Haffer, 1959).




FIG. E133. Mecynodontidae (p. N657).

Superfamily GLOSSACEA Gray, 1847

[*nom. transl.* HABE, 1951 (ex Glossidae GRAY, 1847)] [= Isocardiacea GRAY, 1842 (*nom. transl. et correct.* DALL, 1895, ex Isocardiadae GRAY, 1842); Arcticacea AUCT. (*partim*)] [Family-group names based on *Isocardia* LAMARCK, 1799, are invalid (Code, Art. 11e)] [Materials for this superfamily prepared by MYRA KEEN and RAYMOND CASEY with additions as recorded]

Shells inequilateral, mostly equivalve, beaks well forward, spirally twisted in most; surface smooth or with some concentric (rarely any radial) ribbing; ligament external; hinge of the form termed cyprinoid by authors, having 2 or 3 cardinal teeth in each valve and well-developed laterals in most; teeth tend to be parallel to hinge margins; pallial line normally entire (sinuate in a few). *U.Trias.-Rec.*

Family GLOSSIDAE Gray, 1847

[= Isocardiadae GRAY, 1842] [Materials for this family prepared by MYRA KEEN and RAYMOND CASEY]

Rotund to cordiform, not gaping, with prosogyrate to gyrate beaks; no lunular groove; ligament and resilium in deep groove; hinge with two lamellar cardinals in either valve, variable in form, laterals inconstant; adductor scars equal; internal margin smooth; pallial line entire. *Paleoc.-Rec.*

Glossus POLI, 1795 [**G. rubicundus* (= *Cardium humanum* LINNÉ, 1758); M] [= *Cuculla* SEBA, 1781 (*nonbinom.*); *Glossoderma* POLI, 1795 (obj.);

Isocardia LAMARCK, 1799 (obj.; SD CHILDREN, 1823); *Buccardium* MEGERLE VON MÖHLFELD, 1811 (obj.); *Bucardia* SCHUMACHER, 1817 (obj.); *Tychocardia* ROEMER, 1868 (obj.)]. Beaks gyrate, lunular area depressed. *Paleoc.-Rec.*, Eu.-N.Atl.-W.Asia-IndoPac.

G. (Glossus). Cordiform, smooth, ligament long. *L.Oligo.-Rec.*, Eu.-N.Atl.—FIG. E134,4. **G. (G.) humanus* (LINNÉ), Rec., N.Atl.; 4a, LV ext., $\times 0.5$; 4b,c, RV and LV hinges, $\times 1$ (124b).

G. (Aralocardia) VYALOV, 1937 [**Isocardia eichwaldiana* ROMANOVSKY, 1890; OD]. Beaks strongly coiled; concentric sculpture strong especially on umbones. *U.Eoc.-L.Oligo.*, Eu.-W.Asia.—FIG. E134,1. *G. (A.) multicostatus* (NYST, 1845), Oligo., Ger.; 1a,b, LV ext., hinge, $\times 0.5$ (Von Koenen, 1893).

G. (Cytherocardia) SACCO, 1900 [**Isocardia cytheroides* MAYER, 1868; OD]. Smaller than *G. (Glossus)*, more ovate, beaks less twisted, hinge thinner. *Eoc.-Mio.*, Eu.—FIG. E134,9. **G. (C.) cytheroides* (MAYER), Mio., France; LV ext., $\times 1$ (Mayer, 1868).

G. (Meiocardia) H.ADAMS & A.ADAMS, 1857 [**M. moltkiana* "SPENGLER" (= *Chama moltkiana* GMELIN, 1791); SD STOLIČKA, 1870] [= *Miocardia* (*nom.null.*)]. Surface concentrically grooved. *Paleoc.-Rec.*, Eu.-IndoPac.—FIG. E134,7. **G. (M.) moltkianus* (GMELIN), Rec., E.Indies; LV ext., $\times 1$ (124b).

G. (Miocardiella) SACCO, 1904 [**M. taurinensis*; OD]. Small, subtrigonal, beaks prosogyrate, not coiled. *Tert.*, Eu.—FIG. E134,3. **G. (M.) taurinensis* (SACCO), Neog., Italy; 3a,b, LV ext., RV ext., $\times 3$ (Sacco, 1904).




FIG. E134. Glossidae (p. N657-N658).

G. (Miocardiopsis) GLIBERT, 1936 [*Anisocardia eocaenica* BAYAN, 1873; OD]. Ovate, beaks not twisted; hinge with 1 cardinal absent in RV and only 1 anterior lateral. *Eoc.*, Eu.—FIG. E134,5. **G. (M.) eocaenicus* (BAYAN), Belg.; 5a, RV ext., $\times 2$; 5b,c, RV and LV hinges, $\times 3$ (Glibert, 1936).

G. (Sulcocardia) ROVERETO, 1898 [*Isocardia justinensis* MAYER, 1893; OD]. Small, oblique, not carinate, with incised concentric grooves, no lunule; hinge of LV with 2 cardinals, 2a medium-sized, 2b stronger, obliquely arcuate. *Oligo.*, Eu.—FIG. E134,6. **G. (S.) justinensis* (MAYER), Italy; LV ext., $\times 2$ (Mayer, 1893).

?**Blagroveia** COX, 1933 [*B. corrugata*; OD]. Trigonal-ly ovate, inequilateral, shell thin; sculpture of concentric undulations; lunule deeply set, forming

lamina within shell cavity; hinge with 2 cardinal teeth in either valve, no laterals; pallial line and muscle scar outlines not evident in type material. *Eoc.*, Asia.—FIG. E134,2. **B. corrugata*, M.Eoc., India; 2a,b, RV ext., both valves dorsal, $\times 1$ (Cox, 1933).

?**Rakhia** EAMES, 1951 [*R. trapezoidalis*; OD]. Quadrate, thin-shelled, umbones with concentric undulations; hinge and interior unknown. *L.Eoc.*, Asia.—FIG. E134,8. **R. trapezoidalis*, Pak.; RV ext., $\times 1$ (288).

Family DICEROCARDIIDAE Kutassy, 1934

[Materials for this family prepared by L. R. Cox]

Shell medium-sized to large, equivalve,




FIG. E135. Dicerocardiidae (p. N660).

with prominent umbones and prosogyrous, more or less anteriorly placed beaks; ligament external, opisthodontic; cardinal teeth usually two in both valves, but in some genera only one is present in one or other valve; teeth mostly oriented more or less longitudinally (i.e., parallel to hinge line) and well elongated; no posterior lateral teeth; pallial line entire; adductor scars inconspicuous, not observed so far in most genera; surface smooth or with growth rugae only. *U.Trias.-U.Cret.*

With exception of *Dicerocardium*, the genera included in this family strongly resemble the Cenozoic genus *Glossus* (= *Isocardia*) both in external characters and in dentition, except that they lack posterior lateral teeth. Separation of the family from the Glossidae is based partly on the view that the two groups originated independently during different geological periods and that the resemblances between them are due to convergence. The Dicerocardiidae are thought to be derived from the Megalodontidae.

Dicerocardium STOPPANI, 1865, p. 248 [**D. jani*; SD STOLICZKA, 1871, p. 229] [= *Diceratocardium* FISCHER, 1887 (*nom.van.*)]. Medium-sized to large, trigonal, with angular and usually sharply pointed ventral margin; truncated anteriorly, attenuated and in some specimens sharply rostrate posteriorly; umbones narrow, widely separated and even out-turned in some specimens; each valve with sharp carina running from beak to angle of ventral margin; hinge teeth elongate, 1 in LV, 2 in RV, extending posteriorly from below beak in direction parallel with hinge margin. *U.Trias.(Nor.-Rhaet.)*, N. Alps-S. Alps-Hung.-Sicily-Himalayas. — FIG. E135,1a. **D. jani*, Nor., N. Italy; viewed from ant. end, $\times 0.3$ (Stoppani, 1865), — FIG. E135,1b. *D. curionii* STOPPANI, Nor., Sicily; LV ext., $\times 0.5$ (Di Stefano, 1912). — FIG. E135,1c,d. *D. dolomiticum* (LORETZ), Nor., N. Italy, 1c,d, LV int. mold and cast from it showing dentition, $\times 1$ (Frech, 1904).

Agelasina RIEDEL, 1932, p. 57 [**A. plenodonta*; M]. Medium-sized, ovate, higher than long, with strongly prosogyrous and incoiled, terminal beaks; moderately and evenly inflated, without umbonal ridges; both valves with 2 longitudinally elongated cardinal teeth extending posteriorly from below beak, more posterior (upper) one slightly narrower; those of LV received respectively in recesses behind those of RV; muscle scars and pallial line not observed. *U.Cret.(Coniac.-Maastricht.)*, W. Afr. — FIG. E135,2. **A. plenodonta*, Senon.,

Cameroons; 2a,b, RV ext., RV int., $\times 0.7$ (Riedel, 1932).

Ambocardia BERINGER, 1949, p. 212 [**Isocardia planidorsata* ZITTEL, 1865, p. 140; OD]. Medium-sized, rectangularly ovate, with strongly prosogyrous and incoiled, almost terminal beaks; with weak posterior and anterior umbonal ridges persisting to ventral margin but not forming distinct carinae; LV with stout, arcuate, longitudinally elongated main cardinal tooth adjoining lower margin of hinge plate in median position and lamellar posterior cardinal close to nymph; RV with strong, cuneiform anterior cardinal and slightly oblique, lamellar posterior cardinal; lateral teeth not observed. *U.Cret.(Senon.)*, Eu. — FIG. E135,3. **A. planidorsata* (ZITTEL), Senon., Aus.; 3a,b, LV ext. and dentition, $\times 1$ (Zittel, 1865).

Cornucardia KOKEN, 1913, p. 34 [**Craspedodon hornigii* BITTNER, 1901, p. 8; M] [= *Craspedodon* BITTNER, 1901 (*non* DOLLO, 1883); *Conucardia* DIENER, 1923 (*nom.null.*)]. Medium-sized, ovate, higher than long, with strongly prosogyrous and incoiled beaks; valves not carinate but with posterior radial sulcus; hinge plate massive; RV with median arcuate, longitudinally elongated tooth received by LV in recess between 2 similarly elongated teeth, upper weak, lower strong and situated along margin of hinge plate; posterior adductor scar not observed. *U.Trias.*, N. Alps-S. Alps-Hung.-Timor. — FIG. E136,4. **C. hornigii* (BITTNER), Hung.; 4a-c, LV int., RV int., RV ext., $\times 0.7$ (60).

Megalocardia BERINGER, 1949, p. 212 [**Isocardia merrilli* HAMLIN, 1884, p. 43; OD]. Large medium-sized, trigonally ovate, with strongly prosogyrous and incoiled, terminal beaks; valves with weak posterior carina; known only as internal mold, so that details of dentition are uncertain. *L.Cret.(Alb.)*, Syria.

Physocardia WÖHRMANN, 1894, p. 671 [**P. ogilviae*; M]. Medium-sized, ovate, inequilateral; beaks strongly prosogyrous and slightly incoiled; no posterior carina; 2 more or less elongate, parallel, longitudinal teeth in each valve, extending on both sides of beak, upper tooth in RV relatively weak. *U.Trias.(Carn.)*, N. Alps-S. Alps. — FIG. E136,3. **P. ogilviae*, S. Tyrol; 3a, broken shell viewed from right side; 3b,c, LV int., RV int., showing dentition, $\times 0.7$ (Wöhrmann, 1894).

Platycardia BERINGER, 1949, p. 212 [**Isocardia zitteli* BÖHM, 1883, p. 502; OD]. Medium-sized, obliquely ovate, with very strongly prosogyrous and incoiled beaks; not carinate posteriorly; LV with 2 arcuate, longitudinally elongated teeth extending back from below beak and separated by recess for reception of single tooth of RV. *U.Jur.(Tithon.)*, Czech. — FIG. E136,2. **P. zitteli* (BÖHM); 2a,b, RV ext., LV dentition, $\times 1$ (Böhm, 1883).

Pseudisocardia DOUVILLÉ, 1913, p. 459 [*Isocardia cordata* BUCKMAN in MURCHISON, 1844, p. 98; OD] [= *Ankistrocardia* BERINGER, 1949, p. 212 (obj.)]. Medium-sized, ovate, evenly inflated, with strongly prosogyrous and incoiled beaks which are placed well anteriorly but are not terminal; RV with elongate, arcuate, longitudinal posterior cardinal

tooth joined at its anterior end to apex of short anterior cardinal, base of which lies along margin of hinge plate; these teeth are received in LV in recess between 2 arcuate teeth, lower one bordering margin of hinge plate. [Thickening of the lunular margin has been regarded by DOUVILLÉ as an anterior lateral tooth.] *M.Jur.*, Eu.—FIG.




FIG. E136. Dicerocardiidae (p. N660-N662).

E136,1. **P. cordata* (BUCKMAN); 1a,b, Bajoc., Eng., RV ext., RV int., $\times 1$ (Benecke, 1905); 1c-e, Aalen., France; LV, RV, and RV hinges, $\times 1$ (277).

Family CERATOMYOPSIDAE Cox, 1964

[Materials for this family prepared by L. R. Cox]

Shell small, medium- to medium-sized, ovate or subtrigonal, tall, gibbose, with prominent umbones and strongly prosogyrous, incoiled and outturned beaks; valves equally inflated but slightly dissimilar in some specimens, RV beak situated in more anterior position than that of LV; mostly thin-shelled, but with shell wall thickened locally in some specimens; valve margins not gaping; dorsal margins of valves in exact juxtaposition, not overlapping as in Ceratomyidae; ligament external, opisthodontic, its functional part inserted in each valve in groove adjoining posterodorsal margin, which groove is so deeply impressed that its track forms internal ridge coiling around to margin from below beak and giving rise in many specimens to corresponding groove on internal mold of each valve; hinge teeth absent; adductor scars and pallial line superficial, their impressions not yet observed on any of numerous internal molds studied; surface of shell ornamented with weak concentric or oblique ribs. *M.Jur.-U.Jur.*

Representatives of this family have been included in the Ceratomyidae by most modern authors, although before the erection of the type genus its species were mostly referred to *Glossus* [*Isocardia*] and not to *Ceratomya* [*Ceromya*]. The hinge structure, with the juxtaposed posterodorsal margins joined by a simple external but deeply inserted, opisthodontic ligament, is quite different from that characteristic of the Ceratomyidae, in which the margin of the right valve overlaps that of the left and a sub-internal ligament lies between them (Fig. E137,1; F18,1f). The lack of hinge teeth and the usually thin shell distinguish the members of this family from the Glossidae [*Isocardiidae*] and the Dicerocardiidae, some genera of which are rather similar to them in external morphology. While most species of the Ceratomyopsidae are known only as internal molds, specimens of *Cerato-*




FIG. E137. Ceratomyopsidae (p. N662, N664).

myopsis undulata (MORRIS & LYCETT), from the English Bathonian, retain their shell, so that it has been possible to investigate its hinge structure and to confirm the conclusions of various previous workers that it is edentulous. The systematic position of the Ceratomyopsidae is uncertain, but (by analogy with certain Lucinacea which lack hinge teeth) it is suggested that they are edentulous Glossacea with a purely superficial resemblance to the Ceratomyidae.

Ceratomyopsis COSSMANN, 1915, p. 7 [*nom. subst. pro Ceratomyopsis* DE LORIO, 1897 (*non* MEEK, 1872)] [**Ceratomyopsis helveticus (sic)* DE LORIO, 1897, p. 80; SD ROLLIER, 1913, p. 269]. Shell oval, with small posterior lobe separated from main part of flank by sulcus; ornament of concentric or oblique undulations of small amplitude; other characters as defined for family. *M.Jur.-U.Jur.* (*Bajoc.-Kimmeridg.*), Eu.-Asia-Afr.—FIG. E137, 1. **C. helvetica* (DE LORIO), U.Jur. (U.Oxford.), Switz.; 1a,b, lat. and dorsal views of int. molds showing impressions of ligamental grooves, $\times 1$ (de Loriol, 1897).

Tellurimya Cox, 1964, p. 41 [**Cardium telluris* LAMARCK, 1819, p. 19; OD]. Shell small medium-sized, subtrigonal, higher than long, with prominent, almost median ridge dividing its surface into 2 parts, each bearing series of flattened, oblique ribs separated by narrow furrows; 2 series of ribs meeting along ridge at acute, downward-pointing angle; regularly spaced growth furrows commonly divide ribs into rectangular or rhomboid segments. *M.Jur.* (*Bathon.-Callov.*), France-




FIG. E138. Vesicomidae (p. N664).

NE.Afr.-E.Afr.—FIG. E137,2. **T. telluris* (LAMARCK), Callov., E.Afr.; 2*a,b*, lat. and ant. views, $\times 1$ (Müller, 1900).

Family VESICOMYIDAE Dall, 1908

[*nom. correct.* KEEN, herein (*pro Vesicomycidae*)]

[Materials for this family prepared by MYRA KEEN]

Shell ovate to elongate; lunule incised in most; hinge with up to three teeth, not clearly differentiated into cardinals and laterals; with or without pallial sinus. *Oligo-Rec.*

Vesicomya DALL, 1886 [**Callocardia atlantica* SMITH, 1885; OD]. Ovate, inequilateral, smooth. *Mio.-Rec.*, Atl.-Eu.-E.Indies-W.Indies.

V. (*Vesicomya*). Periostracum polished; lunule bounded by groove; hinge resembling *Glossus* (*Meiocardia*) without laterals. *Mio.-Rec.*, Atl.-Eu.-E.Indies. [Deep-water.]—FIG. E138,1. *V. (*V.*) *atlantica* (SMITH), *Rec.*, Atl.; 1*a,b*, RV int., dorsal, $\times 5$ (852).

V. (*Callogonia*) DALL, 1889 [**Callocardia (Callogonia) leeana*; M]. Lunule without border; pallial sinus shallow but acute. *Rec.*, Atl.-W.Indies. [Deep-water.]—FIG. E138,9. *V. (*C.*) *leeana* (DALL), W.Indies; 9*a*, LV int., 9*b,c*, LV and RV hinges, $\times 2$ (217).

V. (*Veneriglossa*) DALL, 1886 [**Cytherea (V.) vesica*; M]. Beaks as in *Glossus* but hinge veneroid, young resembling V. (*Vesicomya*); pallial sinus of moderate size. *Rec.*, Carib.—FIG. E138,12. *V. (*V.*) *vesica* (DALL); RV ext., $\times 1.5$ (Dall, 1890).

V. (*Waisiuconcha*) BEETS, 1942 [**W. alberdinae*; OD]. Resembling *Lioconcha* (*Veneridae*) in outline; LV with 3 cardinal teeth; pallial line entire. *Mio.-Rec.*, W.N.Am.-E.Indies—FIG. E138,2. *V. (*W.*) *alberdinae* (BEETS), Mioplio., Celebes; 2*a,b*, LV int., LV ext., $\times 1.5$; 2*c*, LV hinge, $\times 2.5$ (Beets, 1942).

Archivesica DALL, 1908 [**Callocardia gigas* DALL, 1890; OD]. Modioliform, inflated, lunule not set off; hinge plate short; pallial line with small sinus, descending almost vertically from posterior adductor scar. *Neog.-Rec.*, W.C.Am.-Japan.

A. (*Archivesica*). Hinge teeth tending to radiate. *Rec.*, W.C.Am. [Deep-water.]—FIG. E138,6. *A. (*A.*) *gigas* (DALL), Gulf Calif.; LV int., $\times 0.3$ (Dall, 1908).

A. (*Akebiconcha*) KURODA, 1943 [**Akebiconcha kawamurai*; M]. Smaller than A. (*Archivesica*), less inflated; hinge with teeth more vertical. *Neog.-Rec.*, Japan.—FIG. E138,7. *A. (*A.*) *kawamurai* (KURODA); 7*a-c*, LV ext., RV int., LV int., $\times 0.5$ (Kuroda, 1943).

Calyptogena DALL, 1891 [**C. pacifica*; M]. Elongate, smooth, of earthy texture, with periostracum; escutcheon present, no lunule; ligament external,

deep-seated; pallial line entire. *Mio.-Rec.*, N.Am.-Japan.

C. (*Calyptogena*). Hinge with 1 anterior and 1 posterior lateral tooth in either valve, RV cardinal triangular, LV cardinal hook-shaped. *Mio.-Rec.*, W.N.Am.-Japan.—FIG. E138,11. *C. (*C.*) *pacifica*, DALL, *Rec.*, USA (Alaska); 11*a,b*, RV ext., int., $\times 0.7$ (Dall, 1895).

C. (*Ectenagena*) WOODRING, 1938 [**C. elongata* DALL, 1916; M]. Shell thinner and longer than C. (*Calyptogena*), anterior tooth wanting in RV. *Rec.*, W.N.Am.—FIG. E138,8. *C. (*E.*) *elongata* DALL, USA (Calif.); 8*a,b*, LV ext., RV hinge, $\times 1$ (Dall, 1921; Woodring, 1938).

C. (*Phreagena*) WOODRING, 1938 [**P. lasia*; OD]. Hinge teeth more vertical than in C. (*Calyptogena*). *Plio.*, W.N.Am.—FIG. E138,10. *C. (*P.*) *lasia* (WOODRING), USA (Calif.); 10*a,b*, RV ext., LV hinge, $\times 1$ (Woodring, 1938).

Hubertschenckia TAKEDA, 1953 [**Tapes exoensis* YOKOYAMA, 1890; OD]. Thick, elongate; prosogyrate beaks well forward; escutcheon present; hinge of LV with two cardinals and posterior lateral; RV with 2 teeth parallel to margin, one cardinal bifid, posterior lateral slender, fitting grooved dorsal margin of LV; pallial line slightly sinuate. *U.Oligo.*, Japan.—FIG. E138,4. *H. *exoensis* (YOKOYAMA), Hokkaido; 4*a,b*, LV ext., part int. mold, $\times 0.5$; 4*c*, part RV int. mold and LV int., $\times 0.7$ (Yokoyama, 1890; specimen, Univ. Hokkaido).

Pleurophopsis PALMER, 1919 [**P. unioides*; M] [= *Pleurophoropsis* COSSMANN, 1920 (*nom.van.*)]. Elongate, sculpture of concentric growth lines, LV hinge with 2 cardinals, posterior large, anterior slender; RV hinge with 2 subequal cardinals; anterior adductor scar embedded, bordered by flange above; pallial line entire. *Oligo.*, W.Indies-C.Am.-N.W.S.Am.—FIG. E138,5. *P. *unioides*, Trinidad; RV ext., $\times 0.5$ (Palmer, 1919).

Pliocardia WOODRING, 1925 [**Anomalocardia bowdeniana* DALL, 1903; OD]. Resembling *Vesicomya*, with lunule circumscribed, but posterior end of shell rostrate; hinge with double anterior laterals, no posterior laterals, 1 cardinal in either valve, broad and ill-defined; pallial line with small sinus. *Mio.*, Carib.—FIG. E138,3. *P. *bowdeniana* (DALL), Jamaica; 3*a-c*, RV ext., int., LV int., $\times 4$ (1005).

Superfamily CORBICULACEA Gray, 1847

[*nom. transl.* TRYON, 1882 (*ex Corbiculidae* GRAY, 1847)]
[= *Cyrenacea* GRAY, 1840] [Materials for this superfamily prepared by MYRA KEEN and RAYMOND CASEY]

Rounded-trigonal to ovate, porcelaneous; sculpture concentric, striate; ligament external; hinge with up to three cardinal teeth

in either valve, pivotal cardinal in RV; pallial line entire or with small sinus. ?*L. Jur., M. Jur.-Rec.*

Small to moderately large shells; shell material dense, with well-developed periostracum; hinge commonly with strong posterior and anterior lateral teeth, somewhat serrate. [Recent forms in brackish to fresh water; fossil forms also in marine environments.] ?*L. Jur., M. Jur.-Rec.*

Family CORBICULIDAE Gray, 1847

[*nom. correct.* DALL, 1889 (*pro* Corbiculidae GRAY, 1847, ICZN pend.)] [=Cyrenidae GRAY, 1840] [Materials for this family prepared by MYRA KEEN and RAYMOND CASEY]




FIG. E139. Corbiculidae (p. N666).

- Corbicula** MERGELE VON MÜHLFELD, 1811 [**Tellina fluminalis* MÜLLER, 1774 (designated ICZN, 1955)] [= *Cyrena* LAMARCK, 1818 (type, *C. cor*; SD CHILDREN, 1823); *Eucorbicula* CROSSE & FISCHER, 1894 (genus without cited species); *Serrilaminula* LINDHOLM, 1933 (type, *Corbicula minima* VON MARTENS, 1874; OD)]. Rounded-trigonal; concentric sculpture present; lateral teeth mostly serrate. *L.Cret.-Rec.*, cosmop.
- C. (Corbicula)**. Lateral teeth long, pallial line entire. *U.Cret.-Rec.*, Asia-Afr.-Eu.-N.Am.-W.Pac.—FIG. E139,8. **C. (C.) fluminalis* (MÜLLER), *Rec.*, Asia Minor; *8a-c*, LV ext., RV int., LV int., $\times 1$ (Kennard & Woodward).
- C. (Corbiculella)** VON IHERING, 1907 [**Corbicula (C.) tenuis*; M]. Ovate, hinge weak, laterals short, not serrate. *?U.Mio., Plio.-Pleist.*, S.Am.
- C. (Corbiculina)** DALL, 1903 [**Corbicula angasi* PRIME, 1864; OD]. Small; viviparous; living in fresh water. *Rec.*, Orient.
- C. (Cyanocyclus)** DE BLAINVILLE, 1818 [**Tellina limosa* MATON, 1811; SD DALL, 1903] [= *Neocorbicula* FISCHER, 1887 (type, *Cyrena variegata* D'ORBIGNY, 1835; OD)]. With small pallial sinus; animal viviparous. *?Eoc., Plio.-Rec.*, S.Am.—FIG. E139,9. **C. (C.) limosa* (MATON), *Rec.*, Brazil; *9a,b*, LV int., RV int., $\times 2$ (Parodiz & Hennings, 1965).
- C. (Cyrenodonax)** DALL, 1903 [**C. formosana*; OD]. Small, thin, trigonal, beak at posterior 3rd; smooth, inflated. *Pleist.-Rec.*, E.Asia.
- C. (Donacopsis)** SANDBERGER, 1872 [**Cyrena acutangularis* DESHAYES, 1858; SD DALL, 1903]. Donaciform, posterior end shorter; pallial sinus short, wide. *Paleoc.-Eoc.*, Eu.—FIG. E139,5. **C. (D.) acutangularis* (DESHAYES), *Eoc.*, France; *5a-c*, LV ext., RV int., LV int., $\times 1$ (Cossmann & Pissarro).
- C. (Leptesthes)** MEEK, 1871 [**Cyrena fracta* MEEK, 1870; OD]. Subtrigonal to long-ovate; lunule defined by incised line; pallial sinus short, wide, triangular. *U.Cret.*, C.USA.—FIG. E139,7. **C. (L.) fracta* (MEEK), USA(Wyo.); RV ext., $\times 0.5$ (Meek).
- C. (Loxoptychodon)** SANDBERGER, 1872 [**Cyrena intermedia* MELLEVILLE, 1843; SD DALL, 1903]. Subtrigonal, anterior end shorter; cardinals 2 in LV; anterior laterals shorter than posterior; pallial line broadly sinuous. *Paleoc.-L.Eoc.*, Eu.—FIG. E139,1. **C. (L.) intermedia* (MELLEVILLE), *L.Eoc.*, Eng; *1a,b*, LV int., ext., $\times 1$ (Morris).
- C. (Paracorbicula)** KOBAYASHI & SUZUKI, 1939 [**Corbicula sanchuensis* YABE & NAGAO, 1926; OD]. Obliquely ovate to subcircular; posterolateral teeth long and crenulate; pallial line distinctly sinuate. *L.Cret.* ("Wealden"), Japan.—FIG. E-139,2. **C. (P.) sanchuensis* YABE & NAGAO; *2a,b*, RV ext., int. mold, $\times 0.5$ (Yabe & Nagao).
- C. (Tellinocyclus)** DALL, 1903 [**Cyrena tellinella* DESHAYES, 1825; OD]. Like *C. (Donacopsis)* but with short, distant lateral teeth. *Paleoc.-Eoc.*, Eu.—FIG. E139,4. **C. (T.) tellinella* (DESHAYES), Sparnac., France; *4a-c*, LV ext., RV int., LV int., $\times 1$ (Cossmann & Pissarro).
- Acyrena** LEBEDEV, 1958 [**A. jenissijensis*; OD?]. *M.Jur.*, USSR(W.Sib.).
- Batissa** GRAY, 1853 [**Cyrena tenebrosa* HINDS, 1842; SD STOLICZKA, 1871]]. Large, heavy, ovate; lateral teeth serrate to crenulate. *M.Jur.-Rec.*, Orient.
- B. (Batissa)**. Pallial line entire or nearly so. *U.Jur.-Rec.*, Asia-E.Indies.—FIG. E139,10. **B. (B.) tenebrosa* (HINDS), *Rec.*, Philip. Is.; LV ext., $\times 0.5$ (Hinds).
- B. (Cyrenobatissa)** SUZUKI & OYAMA, 1943 [**Corbicula subsulcata* CLESSIN, 1878; OD]. Laterals shorter than in *B. (Batissa)*, nymph and ligament not so large. *Eoc.-Rec.*, Orient.
- B. (Tetoria)** KOBAYASHI & SUZUKI, 1937 [**B. (T.) yokoyamai*; OD]. Of medium size, subelliptical to subcircular; pallial line with deep, narrow ascending sinus. *M.Jur.*, Japan-China.—FIG. E139,6. **B. (T.) yokoyamai*, Japan; RV ext., $\times 0.5$ (Kobayashi & Suzuki).
- Costocyrena** HAYAMI, 1965, ex MATSUMOTO & KANMERA, 1952, MS [**C. matsumotoi*; OD]. Shell radially ribbed; hinge resembling that in *Eomiodon*. *L.Cret.*, Japan.
- Dentonia** STEPHENSON, 1953 [**Cytherea leveretti* CRAGIN, 1893; OD]. Broadly subovate to subtrigonal; lunule long, bounded by feeble line; umbonal ridge weak; escutcheon wanting; lateral teeth smooth, anterior teeth short; pallial line with posterior truncation. *U.Cret.(Cenom.)-Eoc.*, N.Am.-France.—FIG. E139,11. **D. (C.) leveretti* (CRAGIN), *U.Cret.*, USA(Tex.); *11a*, RV ext., $\times 1$ (Stephenson); *11b,c*, RV int., LV int., $\times 1$ (Casey, after Stephenson).
- Ditypodon** SANDBERGER, 1875 [**Cyrena suesii*; M]. Oval, anterior end shorter; 1 stout cardinal in either valve; lateral teeth smooth, anterior short and stout, posterior longer; pallial line obtusely sinuous. *U.Mio.-L.Plio.*, Eu.—FIG. E139,3. **D. suesii* (SANDBERGER), *Plio.*, Italy; *3a-c*, LV hinge, LV int., LV ext., $\times 2$ (Sandberger).
- Eocallista** DOUVILLÉ, 1921 [**Venus brongniarti* RÖMER, 1836 (non PAYRAUDEAU, 1826) (= *V. caudata* GOLDFUSS, 1840); SD FRIZZELL, 1936]. Trigonal-ovate; no lunule or escutcheon; lateral teeth smooth, *PII* merged into margin, *PIII* absent; *2a* and *I* attached to laterals; pallial line with posterior truncation. *M.Jur.-U.Jur.*, Eu.
- E. (Eocallista)**. Cardinal teeth, except *3b*, entire; anterior laterals short, *AIII* wanting. *Jur.(Bathon-Portland.)*, Eu.—FIG. E140,10. **E. (E.) caudata* (GOLDFUSS), *U.Jur.*, France; *10a,b*, LV ext., both valves dorsal, $\times 0.5$ (Romer); *10c,d*, RV and LV hinges, $\times 1$ (Casey, from specimens).
- E. (Hemicorbicula)** CASEY, 1955 [**Cyclas parva* J. DE C. SOWERBY, 1836; OD]. Cardinal teeth 1,




FIG. E140. Corbiculidae (p. N666, N668-N669).

- 3*b*, 2*b*, and 4*b* more or less bifid; 2*a* entire or grooved; anterior laterals moderately long, *AIII* incipient. *U.Jur.*-(*Purbeck.*), Eu.—FIG. E140, 11. **E. (H.) parva* (SOWERBY); *U.Jur.*, Eng.; 11*a,b*, RV and LV hinges, $\times 2$ (Casey, from specimens).
- Filosina** CASEY, 1955 [**F. gregaria*; OD]. Trigonal-ovate or subrectangular; no lunule or escutcheon; lateral teeth finely striate, *PII* merged into margin, *PIII* wanting; pallial line truncate posteriorly but not sinuate. *L.Cret.*(*Wealden-Apt.*), Eu.-Syria.—FIG. E140, 14. **F. gregaria*, *L.Cret.*(*Wealden*), Eng.; 14*a-c*, slab, $\times 1$, RV and LV hinges, enl. (Casey).
- Fulpia** STEPHENSON, 1946 [**F. pinguis*; OD]. Trigonal-ovate; umbo subangular; posterior ridge weak; lunule bounded by impressed line; lateral teeth finely striate; pallial sinus narrow, deep, ascending. *U.Cret.*(*Cenoman.*), N.Am.—FIG. E140, 1. **F. pinguis*, USA(Tex.); 1*a*, LV ext., $\times 1$ (Stephenson); 1*b,c*, LV int., RV int., $\times 1$ (after Stephenson).
- Isodomella** KOBAYASHI & SUZUKI, 1939 [**Cyrena shiroensis* YABE & NAGAO, 1926 (= **C. naumanni* NEUMAYR, 1890); OD]. Subtrapezoidal, rounded in front, truncate behind; posterior-lateral teeth longer than anterolaterals; 2*b* and 1 long, wedge-shaped; 3*b* very thin; pallial line simple. *L.Cret.* ("Wealden"), Japan.—FIG. E140, 2. **I. naumanni* (NEUMAYR); RV ext., $\times 1$ (Kobayashi & Suzuki).
- Nemetia** CASEY, 1955 [**Platopsis triangularis* WHITFIELD, 1891; OD]. Subtrigonal, with strong posterior angulation; no lunule or escutcheon; cardinal teeth entire, 1 and 2*a* attached to laterals; *PIII* wanting. *L.Cret.*(*Apt.*), Syria.—FIG. E140, 5. **N. triangularis* (WHITFIELD), Syria; 5*a,b*, LV and RV hinges, $\times 2$ (after Vokes).
- ?**Pharodina** STEPHENSON, 1953 [**P. ferrana*; OD]. Obliquely trigonal; posterior slope subangular; umbo prominent, rounded; beaks strongly prosogyrate; hinge as in *Nemetia* but with anterior laterals shorter and distinctly separated from 2*a* and 1. *U.Cret.*(*Cenoman.*), N.Am.—FIG. E140, 13. **P. ferrana*, USA(Tex.); 13*a-c*, LV ext., LV int., RV int., $\times 1$ (890).
- Polymesoda** RAFINESQUE, 1828 [**Cyclas caroliniana* Bosc, 1801; OD] [= *Egetaria* MÖRCH, 1860 (type, *P. (E.) pullastra*; M); *Cyprinella* GABB, 1864 (non GIRARD, 1856); *Diodus* GABB, 1868 (pro *Cyprinella*) (type, *Cyprinella tenuis* GABB, 1864; M); *Leptosiphon* FISCHER, 1872 (obj.); *Americana* CLESSIN, 1879 (obj.)]. Medium-sized to large; beaks prominent, inturned; inequilateral; cardinal teeth 3, LV with 1 anterior and 1 long posterior lateral against 2 in RV. *Eoc.-Rec.*, C.Am.-Japan-S.Am.-Carib.
- P. (Polymesoda)**. Pallial line with deep, narrow sinus. *Rec.*, E.N.Am.-E.C.-Am.-W.C.Am.—FIG. E140, 4. **P. (P.) caroliniana* (Bosc), USA(Fla.); 4*a,b*, RV ext., LV int., $\times 1$ (Bosc, 1801).
- P. (Egeta)** H. ADAMS & A. ADAMS, 1858 [**Cyrena anomala* DESHAYES, 1858; SD BAKER, 1930] [*pro Anomala* DESHAYES, 1855 (non VON BLOCK, 1799)]. Ovate-elongate, periostracum velvety; pallial sinus small. *Rec.*, C.Am.-S.Am.
- P. (Geloia)** GRAY, 1842 [genus without nominal species] [**Cyrena zeylanica* LAMARCK, 1818 (= **Venus coaxans* GMELIN, 1791); SD GRAY, 1847] [= *Isodoma* DESHAYES, 1858 (type, *I. cyprinoides*; M); *Indica* CLESSIN, 1879 (obj.)]. Large, lateral teeth smooth and short; pallial line entire. *Eoc.-Rec.*, Orient.—FIG. E140, 7. **P. (G.) coaxans* (GMELIN), *Rec.*, E.Indies; 7*a,b*, LV ext., RV int., $\times 0.25$ (124).
- P. (Neocyrena)** CROSSE & FISCHER, 1894 [**Cyrena nicaraguana* PRIME, 1869; SD BAKER, 1930]. Surface concentrically corrugated, periostracum smooth, shining. *Rec.*, E.C.Am.-W.C.Am.
- P. (Pseudocyrena)** BOURGUIGNAT, 1854 [**Cyclas maritima* D'ORBIGNY in SAGRA, 1842; OD] [= *Cyrenocapsa* FISCHER, 1872 (type, *Cyrena floridana* CONRAD, 1816; M)]. Thin-shelled, pallial sinus minute; living in brackish to salt water. *Rec.*, Carib.
- ?**Pseudasaphis** MATSUMOTO, 1938 [**P. japonicus*; OD]. Small, elongate, subtrigonal; lunular region excavated; surface with radial costae; laterals long, smooth. *Cret.*, Japan.—FIG. E140, 9. **P. japonicus*; LV ext., $\times 1$ (Matsumoto).
- Sogamosa** PILSBRY & OLSSON, 1935 [**S. cyrenoides*; OD]. Like *Polymesoda* but hinge with no lateral teeth on flat plate. *U.Eoc.*, S.Am.—FIG. E140, 3. **S. cyrenoides*, Colomb.; 3*a,b*, RV ext., LV hinge, $\times 1$ (Pilsbry & Olsson).
- ?**Soleilletia** BOURGUIGNAT, 1885 [**S. abbadiana*; SD PILSBRY & BEQUAERT, 1927]. Shell thin, fragile; lateral teeth wanting. *Rec.*, Afr.
- Veloritina** MEEK, 1872 [**Cyrena durkeei* MEEK, 1870; OD] [= *Mesocorbicula* SUZUKI & OYAMA, 1943 (type, *Corbicula tetoriensis* KOBAYASHI & SUZUKI, 1939; OD)]. Gibbous trigonal; lunular and ligamentary areas deeply depressed; pallial line posteriorly truncate. Scarcely sinuate; hinge like that of *Filosina*. *M.Jur.-U.Cret.*, Japan-N.Am.—FIG. 140, 6. **V. durkeei* (MEEK), *U.Cret.*, USA (Mont.); 6*a,b*, LV ext., both valves dorsal, $\times 0.5$ (White).
- Villorita** GRIFFITH & PIDGEON, 1834 [**Cyrena cyprinoides* GRAY, 1825; OD] [= *Velorita* GRAY, AUCTT.]. Thick, solid, triangular; lateral teeth large, especially anterior, triangular, finely striate. *Rec.*, Orient.—FIG. E140, 8. **V. cyprinoides* (GRAY); 8*a,b*, RV ext., LV int., $\times 1$ (Chenu).
- ?**Yokoyamaina** HAYAMI, 1958 [**Cyrena elliptica* YOKOYAMA, 1904 (non DUNKER, 1843, = *Y. hayamii* KEEN & CASEY, herein, new name); OD]. Valves inequilateral, elliptical, inflated; surface

nearly smooth, few radial threads ventrally; hinge formula, *Al 3a 3b Pl/AlI 2 4b PII*, with 2 and 3*b* conical, 3*a* and 4*b* close to nymph, anterior laterals tubercular, posterior short and weak; adductor scars small, subequal, impressed, pallial line distinctly sinuate. *Jur. (Hettang.)*, E.Asia.—FIG. E140, 12. **Y. hayamii* KEEN & CASEY, Japan; 12*a-c*, RV and LV int. molds, RV ext., $\times 0.7$ (Hayami, 1958).

Family PISIDIIDAE Gray, 1857

[*nom. correct.*, ICZN, 1955, ex Pisidiidae; = Cycladidae FORBES & HANLEY, 1853; Sphaeriidae JEFFREYS, 1862] [Materials for this family prepared by MYRA KEEN, with advice on technical details of classification from Mr. PETER DANCE, British Museum (Natural History)]

Small to minute shells, oval or quadrate to subtriangular; shell texture thin to opaque, some forms appearing porous. Ligament partially to completely immersed (rarely external); hinge curved, narrow, with anterior and posterior lateral teeth in both valves; cardinal teeth small, not more than two, those of RV straight or united into inverted V, of LV wholly separate. [Fresh-water.] ?*U. Jur.*, *Cret.-Rec.*

Pisidium PFEIFFER, 1821 [**Tellina amnica* MÜLLER, 1774; SD GRAY, 1847 (+ICZN, Op. 335, 1955)] [= *Euglesa* GRAY, 1840, AUCT. (*nom. invalid.*, in *synon.*); *Galileja* DA COSTA, 1840, AUCT. (type, *G. tenebrosa*, *spec. dub.*; M); *Pera* SOWERBY, 1842, ex ALDER, 1831 [in *synon.*] (obj.; M); *Pisum* "MEGERLE" GRAY, 1847, et AUCT. (*non* MEGERLE VON MÜHLFELD, 1811, *nom. dub.*); *Cordula* GRAY, 1852 (ex LEACH MS) (obj.); *Cycladina* CLESSIN, 1871 (*non* CANTRAINE, 1835); *Fossarina* CLESSIN in WESTERLUND, 1873 (*non* ADAMS & ANGAS, 1864); *Fluminina* WESTERLUND, 1873 (obj.); *Flumininea*, *Fluminea* (*nom. null.*); *Rivulina* CLESSIN in WESTERLUND, 1873 (type, *Pisidium supinum* SCHMIDT, 1851; SD CLESSIN, 1879); *Amnicana* FAGOT, 1892 (obj.; SD BOETTGER, 1961); ?*Pusillana* FAGOT, 1892 (type, *Tellina pusilla* GMELIN, 1791, SD BOETTGER, 1961, *spec. dub.*); *Roseana* FAGOT, 1892 (type, *Pisidium roseum* SCHOLTZ, 1843; SD BOETTGER, 1961); *Casertiana* FAGOT, 1892 (type, *Cardium casertanum* POLI, 1795; SD BOETTGER, 1961); *Henslowiana* FAGOT, 1892 (type, *Tellina henslowana* SHEPPARD, 1825; SD BOETTGER, 1961); *Cyclocalyx* DALL, 1903 (type, *P. scholtzii* CLESSIN, 1873; OD); *Cymatocyclas* DALL, 1903 (type, *P. compressum* PRIME, 1851; OD); *Tropidocyclas* DALL, 1903 (type, *Tellina henslowana* SHEPPARD, 1825; OD); *Clessinia* PIAGET, 1913 (*non* DÖRING, 1877); *Pseudeupera* GERMAIN, 1913 (type, *Pisidium landeroini* GERMAIN, 1909; OD); *Lacustrina* STERKI, 1916 (type, *Pisidium idahoense* ROPER, 1890; M);




FIG. E141. Pisidiidae (p. N669-N670).

Fontinalina STERKI, 1916 (type, *Cyclas fontinalis* DRAPARNAUD, 1801 (= *Cardium casertanum* POLI, 1795); M); *Eupisidium* ODHNER, 1921 (obj.; SD BOETTGER, 1961); *Cletella* STRAND, 1928 (*pro Clessinia* PIAGET) (type, *Pisidium profundum* CLESSIN, 1877; OD); *Australpera* IREDALE, 1943 (type, *P. etheridgii* SMITH, 1882; OD); *Glacipisum* IREDALE, 1943 (type, *G. kosciusko*; OD *nom. nud.*); *Speleopisidium* ZHADIN, 1952 (type, *P. subterraneum* ZHADIN, 1932; SD BOETTGER, 1961)]. Small, globose or subtriangular, inequilateral, beaks nearer posterior end; lateral teeth double in RV, single in LV; cardinals 2 in LV, 1 in RV; animal with single siphon. *U. Cret.-Rec.*, cosmop.

P. (Pisidium). Animal with 2 pairs of gills. *U. Cret.-Rec.*, N.Am.-Eurasia.—FIG. E141, 1. **P. (P.) amnicum* (MÜLLER), *Rec.*, Eu.; 1*a-c*, RV ext., int., LV int., $\times 3$ (294).

P. (Afropisidium) KUIPER, 1962 [**P. lepus* KUI-

- PER, 1957 (= **P. pirothi* JICKELI, 1881); OD]. With one pair of gills; ligament external, its fossa turned outward. *Rec.*, Afr.-Asia-N.Z.
- P. (Neopisidium)** ODHNER, 1921 [**P. torquatum* STELFOX, 1918 (= **P. moitessierianum* PALADILHE, 1866); SD HABE, 1951]. Gills reduced to one pair, ligament internal, enclosed in fossette. *Rec.*, E.Afr.-N.Asia-N.Eu.-?N.Am.
- P. (Odhneripisidium)** KUIPER, 1962 [**P. stewarti* PRESTON, 1909; OD]. Shell and animal as in *P. (Neopisidium)* but ligament deeply immersed. *Rec.*, Asia-W.Eu.
- Byssanodonta** D'ORBIGNY, 1846 [**B. paranensis*; M]. Rhomboidal, very inequilateral, cardinal teeth evanescent, anterior lateral tooth present, low. *Rec.*, S.Am.—FIG. E141,4. **B. paranensis*, Argentina; 4a-c, RV ext., LV int., RV hinge, ×3 (4a, U.S. Natl. Museum specimen; 4b, Klappenbach, 1960; 4c, unpublished drawing).
- Eupera** BOURGUIGNAT, 1854 [**Pisidium* (E.) *moquinianum* (= ?*Cyclas modioliformis* ANTON, 1837); M] [= *Limosina* CLESSIN, 1872 (non MACQUART, 1835); *Clessinella* WAAGEN, 1905 (type, *Sphaerium sturanyi*; M)]. Resembling *Byssanodonta* but with cardinal teeth evident in at least LV. *Eoc.-Rec.*, Eurasia-N.Am.-S.Am.-Afr.—FIG. E141,3a,b. **E. moquiniana* (BOURGUIGNAT), *Rec.*, Brazil, 3a,b, RV hinge, LV ext., ×4 (Bourguignat, 1854).—FIG. E141,3c. *E. platenensis* DOELLO-JURADO, *Rec.*, Argentina; LV int., ×4 (Klappenbach, 1960).
- Sphaerium** SCOPOLI, 1777 [**Tellina cornea* LINNÉ, 1758 (fixed, ICZN Opinion 94)] [= *Cyclas* BRUGUIÈRE, 1798 (genus without named species; type, *C. rivicola* LAMARCK, 1818; SD CHILDREN, 1823); *Cornea* MEGERLE VON MÜHLFELD, 1811 (obj.); *Corneocyclas* DE BLAINVILLE, 1818 (type, *Cyclas cornea* "DRAPARNAUD" (= *C. rivicola* LAMARCK, 1818); SD PILSBRY & BEQUAERT, 1927); *Amesoda* RAFINESQUE, 1820 (type, *Cyclas similis* SAY, 1819; SD HANNIBAL, 1912); *Cycladites* KRUEGER, 1823 (obj.); ?*Euglesa* GRAY, 1852 ex LEACH MS (type, *E. henslowiana*; *spec. dub.*); *Cyrenastrum* BOURGUIGNAT, 1854 (type, *Cyclas solida* NORMAND, 1844; M); *Cycladella* CARPENTER, 1865 (type, *C. papyracea*; M); *Corneola* WESTERLUND, 1873 (non HELD, 1837); *Serratisphaerium* GERMAIN, 1909 (type, *Sphaerium courteti*; M); *Trigonosphaerium* KOBELT, 1913 (type, *S. alticola*; M); *Sphaerinova* IREDALE, 1943 (type, *Sphaerium macgillivrayi* SMITH, 1882; OD)]. Oval, quadrate, bluntly triangular in some, shell moderately solid, concentrically striate; beaks nearly median; animal with 2 siphons. ?*U.Jur.*; *Cret.-Rec.*, Eu.-N.Am.-Holarctic-Afr. [= *Sphaeriastrum* BOURGUIGNAT, 1854 (type, *Cyclas rivicola* LAMARCK, 1818; SD KOBELT, 1881).]
- S. (Sphaerium)**. Beaks rounded, not set off by projecting caps. ?*U.Jur.*, *Cret.-Rec.*, Holarctic-Afr.—FIG. E141,2a-c. **S. (S.) corneum* (LINNÉ), *Rec.*, Eng.; 2a, LV ext., ×2 (Jeffreys); 2b,c, LV and RV hinges, ×2 (294).
- S. (Musculium)** LINK, 1807 [**Tellina lacustris* MÜLLER, 1774; M] [= *Phymesoda* RAFINESQUE, 1820 (obj.); *Calyculina* CLESSIN, 1872 (obj.); *Carneola* WESTERLUND, 1873 (obj.); *Primella* COOPER, 1890 (obj.) (type, *Cyclas calyculata* DRAPARNAUD, 1805 = *Tellina lacustris* MÜLLER, 1774; SD KEEN, herein)]. Hinge plate weak, embryonal shell prominent, commonly set off by a groove. *Mio.-Rec.*, Eu.-N.Am.—FIG. E141,5. **S. (M.) lacustre* (MÜLLER), *Rec.*, Eng.; 5a,b, LV and RV hinges, ×2 (294).
- S. (Pseudocorbicula)** DAUTZENBERG, 1908 [**P. alluaudi*; M]. Like *Corbicula* in form but with hinge of *Sphaerium*. *Rec.*, Afr.
- S. (Sulcastrum)** STERKI, 1930 [**Sphaerium sulcatum* LAMARCK, 1818; OD]. Densely microscopically rugulose. *Rec.*, N.Am.-Eu.

Superfamily VENERACEA Rafinesque, 1815

[*nom. correct.* MENKE, 1830 (pro *Veneridia* RAFINESQUE, 1815)] [Materials for this superfamily prepared by MYRA KEEN]

Ovate shells, ornamentation predominantly concentric but also radial in some, with spines or lamellae, especially near posterior slope; beaks anterior, prosogyrate. Ligament external, opisthodontic. Cardinal hinge teeth generally three in either valve; pallial sinus usually present. *L.Cret.-Rec.*

Family VENERIDAE Rafinesque, 1815

[*nom. transl. et correct.* LEACH, 1819 (ex *Veneridia* RAFINESQUE, 1815)]

Lunule and escutcheon usually well developed. Cardinal teeth three in either valve, with 1 and 2b commonly thicker than 3a and 2a; posterior lateral teeth feeble or wanting, anterior laterals present in some groups, absent in others; pallial sinus varying in size and shape. *L.Cret.-Rec.*

This family probably is polyphyletic in origin. Subfamily divisions here used do not necessarily reflect genetic relationships but are adopted for convenience in arrangement.

Subfamily VENERINAE Rafinesque, 1815

[*nom. transl. et correct.* SWAINSON, 1840 (ex *Veneridia* RAFINESQUE, 1815)]

Sculpture usually both radial and concentric. Anterior lateral tooth (AII) present in left valve. *M.Eoc.-Rec.*

Venus LINNÉ, 1758 [**V. verrucosa*; SD GRAY, 1847 (ICZN, 1950)] [= *Venusarius* DUMÉRIL, 1805 (obj.); *Clausina* BROWN, 1827 (obj.); *Ventricola* RÖMER, 1867 (obj.)]. Escutcheon smooth, large, beveled in LV; hinge with *All* small to pustular;

pallial sinus short. *Oligo-Rec.*, Eu.-Afr.-E.Indies-N.Am.

V. (Venus). Sculpture of concentric ribs intersected over part of surface by divaricate radial ribs; *All* minute (711). *Oligo.(Aquitain.)-Rec.*,




FIG. E142. Veneridae (Venerinae) (1-4,7-9) (Circinae) (5-6,10) (p. N671-N673).

- S.Eu.-N.Afr.—FIG. E142,1. **V. (V.) verrucosa* LINNÉ, Rec., Medit.; 1a-c, LV ext., RV and LV hinges, $\times 0.5$ (124b).
- V. (Antigona)** SCHUMACHER, 1817 [**A. lamellaris*; OD] [= *Omphalocathrum* MÖRCH, 1853 (obj.)]. With radiating sculpture underlying raised concentric lamellae; All small but evident; lunule sunken (329a). Rec., E.Indies.
- V. (Ventricoloidea)** SACCO, 1900 [**Cytherea multilamella* LAMARCK, 1818; OD]. Radial sculpture wanting; All elongate (329a). *Oligo.-Rec.*, Eu.-E.Indies-E.N.Am.
- ?**Ameghinomya** VON IHERING, 1907 [**Chione argentina* VON IHERING, 1897; OD]. Sculpture concentric to cancellate; escutcheon wanting; pallial sinus small. *Tert.*, ?*Rec.*, S.S.Am.
- Circomphalus** MÖRCH, 1853 [**Venus plicata* GME-LIN, 1791 (non BARBUT, 1788) (= **V. foliaceo-lamellosa* DILLWYN, 1817); SD SACCO, 1900] [= *Mioclauisnella* KAUTSKY, 1936 (invalidly proposed)]. Trigonal, compressed, cordate, lunule impressed, escutcheon well defined, larger in LV; sculpture of raised lamellae (329a). *Mio.-Rec.*, Eu.-W.Afr.—FIG. E142,2. **C. foliaceolamellosa* (DILLWYN), Rec., Medit.; RV ext., $\times 0.5$ (124).
- Dosina** GRAY, 1835 [**D. zelandica*; SD FRIZZELL, 1936] [= *Dorsina* GRAY, 1840 (nom. null.); *Dosinula* FINLAY, 1926 (obj.)]. Concentric sculpture predominating; All small to moderate; lunule small, sculptured; pallial sinus short, triangular. *M.Eoc.-Rec.*, S.Pac.
- D. (Dosina)**. Medium in size; hinge plate broad; marginal crenulations strong. *U.Oligo.-Rec.*, N.Z.—FIG. E142,3. **D. (D.) zelandica* GRAY, N.Z.; 3a-c, LV ext., int., RV hinge, $\times 0.5$ (592).
- D. (Hina)** MARWICK, 1927 [**Marama (H.) pinguis*; OD]. Cardinals 2a-2b not joined (592). *M.Eoc.-L.Mio.*, N.Z.
- D. (Kuia)** MARWICK, 1927 [**Chione vellicata* HUTTON, 1873; OD]. Like *D. (Dosina)* but All markedly larger (592). *L.Oligo.-Mio.*, N.Z.
- D. (Marama)** MARWICK, 1927 [**M. murdochi*; OD]. Hinge plate narrow, 2a-2b joined; marginal crenulations fine (592). *U.Eoc.-L.Plio.*, N.Z.
- D. (Plurigen)** FINLAY, 1930 [**P. phenax*; OD]. Early sculpture thin lamellae later becoming heavy sulcations at ends of shell (392a). *U.Mio.-Rec.*, N.Z.
- Globivenus** COEN, 1934 [**Venus effossa* PHILIPPI, 1836; M]. Like *Ventricolaria* but with lunule also channeled, escutcheon not beveled in LV (329a). Rec., Medit.—FIG. E142,4. **G. effossa* (PHILIPPI), Italy; 4a-c, LV ext., int., RV int., $\times 1$ (Specimen, Station Oceanographique, Monaco).
- Melosia** DALL, 1915 [**Cytherea glyptoconcha* DALL, 1903; M] [= *Artena* CONRAD, 1870 (non WALKER, 1858); *Netara* FRIZZELL, 1936 (pro *Artena*) (type, *Cytherea staminea* CONRAD, 1839; OD)]. Trigonal, escutcheonal area large, nearly smooth, equal in both valves; sculpture of concentric lamellae. *Mio.*, E.N.Am.—FIG. E142,7. **M. glyptoconcha* (DALL), USA(Fla.); LV ext., $\times 1$ (Dall, 1900).
- Periglypta** JUKES-BROWNE, 1914 [pro *Cytherea* RÖDING, 1798 (non FABRICIUS, 1794)] [**Venus puerpera* LINNÉ, 1758; OD] [= *Proxichione* IREDALE, 1929 (type, *P. materna*, OD); *Tigammona* IREDALE, 1930 (type, *T. persimilis*; OD)]. Quadrate, heavy, pallial sinus rounded; sculpture cancellate (455a). *Oligo.-Rec.*, Eu.-E.N.Am.-W.N.Am.-Pac.—FIG. E142,8. **P. puerpera* (LINNÉ), Rec., Pac.; LV ext., $\times 0.5$ (124b).
- Ventricolaria** KEEN, 1954 [**Venus rigida* DILLWYN, 1817; OD] [= *Ventricola* AUCT. (non RÖMER, 1867)]. Globose, lunule depressed, escutcheon smooth, beveled in LV; sculpture predominantly concentric. *Oligo.-Rec.*, E.N.Am.-W.N.Am.-Medit.-Pac.—FIG. E142,9. **V. rigida* (DILLWYN), Rec., Carib.; 9a,b, LV ext., RV int., $\times 0.5$ (711).

Subfamily CIRCINAE Dall, 1896

Equivalve, subequilateral; pallial line nearly entire. Cardinals smooth or faintly grooved; anterior laterals present. Surface sculptured, usually with dichotomous radial ribbing on some part of shell. *Paleoc.-Rec.*

Circe SCHUMACHER, 1817 [**C. violacea* (= **Venus scripta* LINNÉ, 1758); M]. Trigonal, umbones low; lunule and escutcheon narrow. *Oligo.-Rec.*, Eu.-E.Indies-Pac.-Australia-Japan.

C. (Circe). Compressed, ligament deeply sunken, radial sculpture weak, usually on umbonal area only (329a). *L.Oligo.-Rec.*, Eu.-E.Indies.—FIG. E142,10. **C. (C.) scripta* (LINNÉ), Rec., Pac.; 10a,b, RV ext., LV int., $\times 1$ (Sowerby, 1851-53).

C. (Circentia) JOUSSEAUME, 1888 [**Venus arabica* DILLWYN, 1817 (ex CHEMNITZ, nonbinom.); OD]. Convex, sculpture feebly concentric, inner margins smooth (329a). Rec., Pac.

C. (Fluctiger) IREDALE, 1924 [**F. royanus*; OD]. Rec., Australia.

C. (Laevicirce) HABE, 1951 [**L. soyoae*; OD] (365). Rec., Japan.

C. (Microcirce) HABE, 1951 [**Meretrix gordonis* YOKOYAMA, 1927; OD]. Small, All long, with groove posterior to it (365). *Pleist.-Rec.*, Japan.

C. (Parmulophora) DALL, 1915 [pro *Parmulina* DALL, 1902 (non PÉNARD, 1902)] [**Venus corrugata* DILLWYN, 1817; OD]. Inner margins finely crenulate (329a). Rec., Pac.

C. (Privigna) DALL, BARTSCH, & REHDER, 1938 [**C. (P.) pilsbryi*; OD] (659). Rec., Hawaii.

C. (Redicirce) IREDALE, 1936 [**R. mistura*; OD] (659). Rec., Australia.

Gafrarium RÖDING, 1798 [**Venus pectinata* LINNÉ; SD DALL, 1902] [= *Crista* RÖMER, 1857 (obj.)]. Sculpture of nodose radial ribs dichotomous medial-

ly; pallial sinus small. *L.Mio.-Rec.*, Pac.—FIG. E142,6. **G. pectinatum* (LINNÉ), *Rec.*, E.Indies; 6a-c, LV ext., int., RV int., $\times 1$ (711).

Gouldia C. B. ADAMS, 1847 [**Thetis cerina* ADAMS, 1845; SD VON MARTENS, 1882]. Small, lunule long, escutcheon wanting; pallial sinus small. *Paleoc.-Rec.*, Eu.-N. Am.-Carib.-Japan-Pac.-Australia.

G. (Gouldia). Sculpture fine, concentric or reticulate. *Paleoc.-Rec.*, Eu.-E.N.Am.—FIG. E142,5.

**G. (G.) cerina* (ADAMS), *Rec.*, Carib.; 5a,b, RV ext., int., $\times 2$ (217).

G. (Crenocirce) HABE, 1960 [**Dorisca (C.) picta*; OD]. With crenulate inner ventral margin; otherwise like *Dorisca*. *Rec.*, Japan.

G. (Dorisca) DALL, BARTSCH, & REHDER, 1938 [**D. cooki*; OD]. Radial sculpture stronger, more divergent (659). *Rec.*, Pac.

G. (Gouldiopa) IREDALE, 1924 [**G. australis* ANGAS, 1865; OD]. Smooth (329a). *Rec.*, Australia.

Subfamily SUNETTINAE Stoliczka, 1870

Ligament in a deeply excavated escutcheon; shell surface smooth or with concentric ribbing; hinge with elongate *All. Eoc.-Rec.*

Sunetta LINK, 1807 [**Donax scripta* LINNÉ, 1758; SD DALL, 1902] [= *Cuneus* MEGERLE VON MÜHLFELD, 1811 (non DA COSTA, 1778); *Meroë* SCHUMACHER, 1817 (type, *Venus meroë* LINNÉ, 1758; OD); *Sunemeroë* IREDALE, 1930 (type, *Sunetta adelinae* ANGAS, 1868, ?=*S. scripta* (LINNÉ); OD)]. Elongate-oval, beaks commonly posterior to mid-line; hinge with 4b short, 3b smooth and entire; pallial sinus ample, rounded; inner margin crenulate. *U.Oligo.-Rec.*, W.Pac.-Eu.

S. (Sunetta). Elongate, sculpture concentric. *U. Oligo.-Rec.*, Eu.-W.Pac.-S.Pac.—FIG. E143,10. **S. (S.) scripta* (LINNÉ), *Rec.*, Pac.; 10a,b, RV ext., LV int., $\times 1$ (Sowerby, 1851-53).

S. (Cyclosunetta) FISCHER-PIETTE, 1939 [*pro Sunettina* JOUSSEAUME, 1891 (non PFEIFFER, 1869)] [**Sunettina sunettina* JOUSSEAUME, Sept. 1891 (= *Sunetta contempta* SMITH, June, 1891); OD]. Short, escutcheon shallow (659). *Rec.*, W. Pac.

S. (Sunettina) PFEIFFER, 1869 [**Cytherea solanderii* GRAY, 1825; M] [= *Solanderina* DALL, 1902 (obj.)]. Inflated, smooth (659). *Rec.*, Pac.

Meroena JUKES-BROWNE, 1908 [**Cytherea trigonula* DESHAYES, 1825; OD]. Nearly equilateral, ligament only moderately deep; hinge with 3b deeply grooved; inner margin smooth (455). *Eoc.*, Eu.—FIG. E143,9. **M. trigonula* (DESHAYES), France; 9a, RV ext., $\times 1$; 9b,c, RV and LV hinges, $\times 1.5$ (Cossmann & Pissarro, 1904).

Subfamily MERETRICINAE Gray, 1847

[*nom. correct.* FISCHER, 1887 (*pro Meretricina* GRAY, 1847)]

Elongate-ovate; sculpture subdued or wanting; hinge with cardinal teeth tending to radiate, anterior lateral teeth in RV commonly flanked with denticles above and below. *U.Cret.-Rec.*

Meretrix LAMARCK, 1799 [**Venus meretrix* LINNÉ, 1758; T] [= *Cytherea* LAMARCK, 1805 (non FABRICIUS, 1794); *Nympha* MÖRCH, 1853 (non FITZINGER, 1826)]. Large, trigonal, smooth, lunule and escutcheon faint; teeth large; pallial sinus broad. *U.Mio.-Rec.*, E.Indies.—FIG. E143,1. **M. meretrix* (LINNÉ), *Rec.*, E.Indies; 1a-c, RV int., LV int., RV ext., $\times 0.5$ (711).

Aeora CONRAD, 1870 [**A. cretacea*; M]. Smooth, hinge with nymphs smooth or weakly rugose, *All* distant from 2a; pallial sinus deep, rounded, somewhat ascending. *U.Cret.*, E.N.Am.—FIG. E143,3. **A. cretacea*, *Cret.*, USA(N.J.); 3a-c, LV ext., RV int., LV int., $\times 1$ (711).

Bassinaria MARWICK, 1928 [**B. macclurigi*; OD]. Resembling *Bassina* but RV with deep pit for reception of *All. M.Plio.*, N.Z.—FIG. E143,11. **B. macclurigi*; RV int., $\times 0.7$ (593).

Eomeretrix TURNER, 1938 [**Pitaria martini* DICKERSON, 1914; OD]. Like *Meretrix* but with V-shaped pallial sinus and a well-defined lunule (695). *Eoc.*, W.N.Am.

Grateloupia DESMOULINS, 1828 (*emend.*) [**G. donaciformis* (= *Donax irregularis* BASTEROT, 1825)] [= *Gratelupia* DESMOULINS, 1828 (*nom. imperf.*)]. With parallel ridges on nymphs. *Eoc.-Mio.*, Eu.-N.Am.-S.Am.

G. (Grateloupia). Pallial sinus long and acute. *Oligo. (Aquit.)-Mio. (Helvet.)*, Eu.—FIG. E143,7. **G. (G.) irregularis* (BASTEROT), *Mio.*, France; RV int., $\times 1$ (1007).

G. (Cytheriopsis) CONRAD, 1865 [**Cytherea hydana* CONRAD, 1833; OD] [= *Grateloupina* DALL, 1902 (obj.)]. Pallial sinus small (711). *Eoc.-Mio.*, E.N.Am.-S.Am.

G. (Xenoloupia) CLARK & DURHAM, 1946 [**X. carmenensis*; OD]. With a flangelike escutcheon (695). *Eoc.*, S.Am.

Meretrissa JUKES-BROWNE, 1908 [**Cytherea depressa* DESHAYES, 1858; OD]. Small, nearly smooth; hinge weak; pallial sinus short, rounded (455). *Oligo.*, Eu.—FIG. E143,6. **M. depressa* (DESHAYES), France; 6a,b, LV int., RV int., $\times 4$ (Keen, n, Stanford Univ. Coll.).

Tivela LINK, 1807 [**Venus tripla* LINNÉ, 1771; SD KOBELT, 1881] [= *Trigona* MEGERLE VON MÜHLFELD, 1811 (non JURINE, 1807)]. Medium-sized to large, trigonal, smooth; nymph bifid or trifid, resembling cardinal tooth. *Eoc.-Rec.*, N.Am.-C. Am.-S.Am.-Afr.-Ind.O.

T. (Tivela). Medium in size, inner margin smooth. *Mio.-Rec.*, E.N.Am.-W.C.Am.-W.Afr.-Ind.O.—FIG. E143,4. **T. (T.) tripla* (LINNÉ), *Rec.*, W. Afr.; 4a,b, LV ext., RV int., $\times 1$ (124b; 1007).

T. (Comus) COX, 1930 [*Venus platyaulax* TOMLIN, 1924; OD]. Sculptured with heavy concentric undulations (329a). *Pleist.-Rec.*, SE.Afr.

STEARNS, MS) (= *Venus isabelleana* D'ORBIGNY, 1846); OD]. Inner margins crenulate (329a). *Rec.*, E.S.Am.

T. (Eutivela) DALL, 1891 [*T. perplexa* (ex

T. (Pachydesma) CONRAD, 1854 [*pro Trigonella*




FIG. E143. Veneridae (Sunettinae) (9-10), (Meretricinae) (1-8, 11) (p. N673-N675).

- CONRAD, 1837 (*non DA COSTA*, 1778)] [**Cytherea crassatelloides* CONRAD, 1837 (= *Donax stultorum* MAWE, 1823); M]. Large and heavy, inner margins smooth; nymph not bifid but set off from *4b*. *L.Eoc.-Rec.*, E. N. Am.-W. N. Am.—FIG. E143,8. **T. (P.) stultorum* (MAWE), *Rec.*, USA (Calif.); LV int., $\times 0.25$ (711; Keen, n, Stanford Univ. Coll.).
- T. (Planitivela)** OLSSON, 1961 [**Cytherea planulata* BRODERIP & SOWERBY, 1829; OD]. Compressed, lunule elliptical, nymphal area forming high plate set off by deep groove. *Rec.*, W.C.Am.-S.Am.
- Tivelina** COSSMANN, 1886 [**Cytherea rustica* DESHAYES, 1825; SD CROSSE, 1886]. Small, compressed cardinal teeth short; pallial sinus small, rounded, ascending (329a). *L.Eoc.-L.Mio.*, Eu.—FIG. E143,5. **T. rustica* (DESHAYES), *Eoc.*, France; *5a-c*, RV ext., RV and LV hinges, $\times 2$ (Cossmann & Pissarro, 1904).
- Transennella** DALL, 1883 [**Cytherea (T.) conradina* DALL, 1883; OD]. Small, pallial sinus rounded; valve margins tangentially grooved (711). *Eoc.-Rec.*, E.N.Am.-W.N.Am.-S.Am.—FIG. E143,2. **T. conradina* (DALL), *Rec.*, Carib.; *2a,b*, RV ext., int., $\times 2$ (222; Dall, 1900, 1904).
- Subfamily PITARINAE Stewart, 1930**
- Inequilateral, beaks anterior. Cardinal teeth not tending to radiate, anterior laterals well developed. *L.Cret.-Rec.*
- Pitar** RÖMER, 1857 [**Venus tumens* GMELIN, 1791; M] [= *Caryatis* RÖMER, 1862 (*non HÜBNER*, 1816); *Pitaria* DALL, 1902 (*nom. van.*)]. Oval or subtriangular, smooth or finely concentrically lamellate; lunule superficial, escutcheon not defined; hinge with *2b* triangular, joined to a thin *2a*, *3a* and *1* separate. *Eoc.-Rec.*, cosmop.
- P. (Pitar)**. Smooth or finely striate, pallial sinus deep and pointed; *4b* confluent with nymph. *Eoc.-Rec.*, N.Am.-W.Afr.-Pac.—FIG. E144,10. **P. (P.) tumens* (GMELIN), *Rec.*, W.Afr.; *10a,b*, RV ext., int., $\times 1$ (711).
- P. (Calpitarina)** JUKES-BROWNE, 1908 [**Cytherea sulcataria* DESHAYES, 1825; OD]. With fine concentric lamellae; nymphs striate; pallial sinus short, rounded (455). *Eoc.*, Eu.-W.N.Am.-SE. Asia.
- P. (Costellipitar)** HABE, 1951 [**Caryatis chordata* RÖMER, 1876; OD]. Small, concentric ribs evident (365). *Rec.*, Pac.-Japan.
- P. (Hyphantosoma)** DALL, 1902 [**Cytherea carbasea* GUPPY, 1866; OD]. With zigzag concentric grooves (228). *Oligo.-Plio.*, W.Indies-N.Z.
- P. (Hysteroconcha)** DALL, 1902 [*pro Dione* GRAY, 1847 (*non HÜBNER*, 1819)] [**Venus dione* LINNÉ, 1758; OD]. Like *P. (Lamelliconcha)* but with posterior area bordered by spines (228). *U.Oligo.-Rec.*, C.Am.—FIG. E144,9. **P. (H.) dione* (LINNÉ), *Rec.*, Carib.; *9a,b*, LV ext., RV int., $\times 1$ (1007).
- P. (Katherinella)** TEGLAND, 1929 [**Calloccallista arnoldi* WEAVER, 1916; OD]. Lunule pouting; *All* bladelike, buttressed under upper margin of hinge plate (329a). *Oligo.-Rec.*, W.N.Am.
- P. (Lamelliconcha)** DALL, 1902 [**Cytherea concinna* SOWERBY, 1835; OD]. With thin concentric lamellae; hinge plate excavated and attenuated behind, nymphs longitudinally striate; pallial sinus obtuse (711). *Eoc.-Rec.*, N.Am.-W.N.Am.-S.Am.
- P. (Meisenia)** MAKIYAMA, 1936 [**M. tateiwai*; OD]. Lunule wanting (329a). *Mio.*, Korea.
- P. (Nanopitar)** REHDER, 1943 [**P. (N.) pilula*; OD]. Small, smooth, rounded, inner margins smooth (695). *Rec.*, Carib.
- P. (Omnivenus)** PALMER, 1927 [**Cytherea discoidalis* CONRAD, 1833; OD]. Resembling *P. (Tinctora)*; nymphs rugose (711). *Eoc.*, E.N.Am.
- P. (Pitarella)** PALMER, 1927 [**Calloccardia gatunensis* DALL, 1903; OD]. *All* much reduced (711). *Eoc.-Mio.*, Carib.
- P. (Pitarenus)** REHDER & ABBOTT, 1951 [**Pitaria cordata* SCHWENDEL, 1951; OD]. Ventral margin crenulate; hinge as in *P. (Pitarella)* (695). *Rec.*, Carib.
- P. (Pitarina)** JUKES-BROWNE, 1913 [**Cytherea citrina* LAMARCK, 1818; OD]. Nymphs smooth; *4b* wholly free and oblique; pallial sinus short (329a). *Rec.*, Pac.
- P. (Rhabdopitaria)** PALMER, 1927 [**Calloccardia astartoides* GARDNER, 1923; OD]. Shell smooth but middle layer radially ribbed; inner ventral margin finely crenulate; nymphs smooth (711). *M.Eoc.*, E.N.Am.
- P. (Tinctora)** JUKES-BROWNE, 1914 [*pro Callizonata* JUKES-BROWNE, 1913 (*non DOUBLEDAY*, 1846-50)] [**Cytherea vulnerata* BRODERIP, 1835; OD] [= *Jukes-Brownia* COSSMANN, 1920 (obj.); *Callizonata* STRAND, 1926 (obj.)]. Thick-shelled, round, glossy; valve margins crenulate (455a). *Rec.*, W.C.Am.
- Amiantis** CARPENTER, 1864 [**Cytherea callosa* CONRAD, 1837; M]. Large, thick-shelled; nymphs rugose; pallial sinus generally pointed. *Eoc.-Rec.*, N.Am.-S.Am.
- A. (Amiantis)**. Concentric ribs commonly anastomosing (711). *U.Oligo.-Rec.*, E.N.Am.-W.N.Am.—FIG. 39,1. **A. (A.) callosa* (CONRAD), *Rec.*, USA (Calif.); *1a,b*, RV int., LV ext., $\times 0.5$ (Keen, n; Stanford Univ. Coll.).
- A. (Eucallista)** DALL, 1902 [**Cytherea purpurata* LAMARCK, 1818; OD]. Thinner than *A. (Amiantis)*, ribs weak on posterior (711). *Rec.*, E.N. Am.-S.Am.
- A. (Venidia)** CLARK & DURHAM, 1946 [**V. steinekei*; OD]. Smooth, escutcheon stronger than in *A. (Amiantis)*, lunule weaker (695). *Eoc.*, W.S. Am.
- Anofia** REYMENT, 1955 [**A. aro*; OD]. Lenticular,




FIG. E144. Veneridae (Pitarinae) (p. N675, N677-N679).

resembling *Naulia* but with cardinals more separated; an excavated ligamental pit in front of nymph. *U.Cret.*, W.Afr.

Aphrodina CONRAD, 1869 [**Meretrix tippiana* CONRAD, 1858; OD] [= *Callistina* JUKES-BROWNE, 1908 (type, *Venus plana* SOWERBY, 1813; OD)]. Cardinal teeth divergent, *All* rugose; pallial sinus deep, ascending. *L.Cret.-U.Eoc.*, S.Am.-N.Z.-N.Am.-Eu.-Afr.

A. (Aphrodina). Escutcheon wanting. *L.Cret.-U.Eoc.*, Eu.-Afr.-N.Am.-S.Am.—FIG. E145, *A. *A. (A.) tippiana* (CONRAD), *U.Cret.*, USA (N.Car.); *4a-c*, RV ext., int., LV hinge, $\times 0.5$ (887).

A. (Sechurina) OLSSON, 1944 [**A. (S.) australis*; OD]. With well-marked escutcheon (695). *Cret.*, S.Am.(Peru).

A. (Tikia) MARWICK, 1927 [**Callista thomsoni* WOODS, 1917; OD]. With strong concentric





FIG. E145. Veneridae (Pitarinae) (p. N678-N679).

- ridges; *All* extremely long, with a knobbed posterior end (592). *Cret.*, N.Z.
- Callista** POLI, 1791 [**Venus chione* LINNÉ, 1758; SD MEEK, 1876] [= *Callistoderma* POLI, 1795 (obj.)]. Glossy, with or without sculpture; pallial sinus wide, horizontal, pointed; *3b* narrow. *Paleoc.-Rec.*, Eu.-N.Am.-C.Am.-Carib.-Asia-N.Z.-E.Indies.
- C. (Callista)**. Smooth, ovate; *3b* with a slight groove (711). *Eoc.-Rec.*, E.N.Am.-Eu.—FIG. E144,7. **C. (C.) chione* (LINNÉ), *Rec.*, *Medit.*; *7a-c*, LV ext., int., RV int., $\times 0.5$ (592).
- C. (Chionella)** COSSMANN, 1886 [*pro Chione* GRAY, 1838 (non MEGERLE VON MÜHLFELD, 1811)]. [**Cytherea ovalina* DESHAYES, 1858; SD CROSSE, 1886] [= *Paradione* DALL, 1909 (obj.)]. Small, nearly smooth, with obsolete concentric folds (392a). *Eoc.-Oligo.*, Eu.-C.Am.
- C. (Costacallista)** PALMER, 1927 [**Venus erycina* LINNÉ, 1758; OD]. Sculpture strong, of flat concentric ridges; hinge plate excavated, *2a* grooved (711). *Paleoc.-Rec.*, E.N.Am.-W.N.Am.-Asia-Eu.-N.Z.
- C. (Macrocallista)** MEEK, 1876 [**Venus gigantea* GMELIN, 1791 (= **V. nimbose* LIGHTFOOT, 1786)]. Elongate-ovate, smooth or with concentric grooves; *3b* not grooved (711). *Eoc.-Rec.*, E.N.Am.-W.N.Am.-E.Indies.—FIG. E144,6. **C. (M.) nimbose* (LIGHTFOOT), *Rec.*, *Carib.*; LV ext., $\times 0.25$ (124b).
- C. (Microcallista)** STEWART, 1930 [**Cytherea proxima* DESHAYES, 1858; OD]. No groove on *2a* (892). *Eoc.*, Eu.-E.N.Am.-W.N.Am.
- Callocardia** A. ADAMS, 1864 [**C. guttata*; M] [= "*Callocallista* Adams," WEAVER, 1916 (*nom. null.*)]. Thin; hinge plate narrow, excavated, 2 cardinals in each valve united as curved arches. *Eoc.-Rec.*, N.Am.-Asia-E.Indies-Eu.
- C. (Callocardia)**. Very thin-shelled, pallial sinus

- nearly obsolete. *Rec.*, E.N.Am.-W.N.Am.-E.Asia.
—FIG. E144,3. **C. (C.) guttata* ADAMS, *Rec.*,
Korea; LV hinges, $\times 2$ (711).
- C. (Agriopoma)** DALL, 1902 [**Cytherea texasiana*
DALL, 1892; OD] [= *Agriodesma* DALL, 1916
(*nom. null.*; *non* DALL, 1909)]. Dull white,
thicker than *C. (Callocardia)*; pallial sinus
sharply angular (711). *M.Eoc.-Rec.*, E.N.Am.
- C. (Aphrodora)** JUKES-BROWNE, 1914 [*pro Leuco-*
thea JUKES-BROWNE, 1913 (*non* MERTENS, 1833)]
[**Callista birsi* PRESTON, 1905; OD]. Hinge
teeth weak (455a). *Rec.*, Ceylon.
- C. (Atopodonta)** COSSMANN, 1886 [**Venus con-*
formis DESHAYES, 1858; SD CROSSE, 1886]. Small
but not extremely thin; pallial line entire (329a).
Eoc.-Mio., Eu.-E.Indies.
- C. (Nitidavenus)** VOKES, 1939 [**Cytherea nitida*
DESHAYES, 1858; OD]. Lunule large, impressed,
escutcheon wanting; pallial sinus well developed.
L.Eoc., Eu.-W.N.Am.—FIG. E144,4. **C. (N.)*
nitida (DESHAYES), *Eoc.*, France; *4a-c*, RV ext.,
RV and LV hinges, $\times 1$ (Vokes, 1939).
- Calva** POPENOE, 1937 [**C. regina*; OD]. Resembling
Trigonocallista but more elongate, with bifid *3b*,
nymph smooth, laterals close to cardinals. *L.Cret.-*
U.Cret., Eu.-N.Am.
- C. (Calva)**. With escutcheon or depressed dorsal
area. *L.Cret.-U.Cret.*, Eu.-N.Am. (92).—FIG.
E145,1. **C. (C.) regina* POPENOE, *U.Cret.*, USA
Calif.; *1a-c*; LV ext., LV and RV hinges, $\times 1$
(748).
- C. (Chimela)** CASEY, 1952 [**Venus caperata* J. DE
C. SOWERBY, 1826; OD]. Dorsal area not de-
pressed (92). *L. Cret. (Alb.)*, Eu.
- Dollfusia** COSSMANN, 1886 [**D. crassa*; M]. Re-
sembling *Pelecycora* in outline but small, hinge with
3b bifid; pallial sinus short, rounded, wide (329a).
Eoc., Eu.
- Dosiniopsis** CONRAD, 1864 [**D. meeki*; SD TATE,
1868]. Large, smooth, lenticular; anterior lateral
teeth and sockets rugose, near cardinals, *3b* deeply
bifid; pallial sinus deep and angular, not ascending.
Paleoc.-Eoc., E.N.Am.-W.N.Am.-Eu. — FIG.
E144,2. **D. meeki*, *Eoc.*, USA (Md.); *2a-c*, LV int.,
RV int., LV ext., $\times 0.5$ (711).
- Gilbertharrisella** F. HODSON & H. HODSON, 1927
[**Pitaria lyncei*; OD]. Like *Pitar* but thick-shelled,
sculpture of concentric folds; lunule large; *All*
peglike; heavy ridge behind anterior muscle scar
(695). *U.Cret.*, S.Am. (Venez.).
- Lepidocardia** DALL, 1902 [**Chione floridella* GRAY,
1838; OD]. Small, compressed, smooth, poster-
iorly attenuated; hinge short, teeth crowded (228).
Rec., W.Afr.
- Lioconcha** MÖRCH, 1853 [**Venus castrensis* LINNÉ,
1758; SD STOLICZKA, 1870]. Hinge strong, re-
sembling *Pitar*; lunule present; pallial sinus shal-
low. *Plio.-Rec.*, Pac.-E.Indies.
- L. (Lioconcha)**. Surface smooth, with zigzag
streaks or spots of color (329a). *Plio.-Rec.*, Pac.
- L. (Sulcilioconcha)** HABE, 1951 [**Cytherea philip-*
pinarum HANLEY, 1844; OD]. With concentric
ribs (365). *Rec.*, Japan.-E.Indies.
- Loxo** DAILEY & POPENOE, 1966 [**L. decore*; OD].
Smaller, longer than *Calva*. *U.Cret.*, N.Am.
- Marwickia** FINLAY, 1930 [*pro Finlaya* MARWICK,
1927 (*non* THEOBALD, 1903)] [**Finlaya parthiana*
MARWICK, 1927; OD]. Oval, lunule concave,
sculpture of weak concentric grooves; *2b* large,
All distant from *2a* (329a). *U.Cret.*, N.Z.
- Megapitaria** GRANT & GALE, 1931 [**Cytherea aur-*
antica SOWERBY, 1831; OD]. Like *Pitar* but much
larger and heavier (329a). *Plio.-Rec.*, W.C.Am.
- Mesocallista** COX, 1952 [**Meretrix andersoni* NEW-
TON, 1909; OD]. Ovate, lunule shallow, narrow,
incised; hinge as in *Aphrodina* but *All* not corruga-
ted. *U.Cret.*, Afr.-Asia-Eu.-S.Am.-N.Am.
- M. (Mesocallista)**. Small, sculpture weak, concen-
tric (695). *U.Cret.*, W.Afr.-India-Eu.-W.S.Am.
—FIG. E145,6. **M. (M.) andersoni* (NEW-
TON), W.Afr.; LV int., $\times 1$ (Cox, 1952).
- M. (Larma)** STEPHENSON, 1953 [**Callistina (L.)*
munda; OD]. Larger, with irregular concentric
ribs bearing secondary threads and some fine
radial lines (695). *U.Cret. (Cenoman.)*, E.N.Am.
- Nagaocella** HAYAMI, 1965 [**Dosiniopsis corrugata*
NAGAO, 1934; OD]. Near *Dosiniopsis* but smooth
marginated within. *L.Cret.*, Japan.
- Naulia** COX, 1952 [**N. orbicularis*; OD]. More
orbicular than *Aphrodina*; hinge with *3b* not so
conspicuously bifid, *4b* down-curved distally, hinge
plate flattened behind *4b*; pallial sinus shallower
(695). *U.Cret.*, W.Afr.
- Notocallista** IREDALE, 1924 [**Cytherea kingi* GRAY,
1827; OD]. Sculptured with irregular concentric
grooves and ridges; otherwise like *Callista*. *Oligo-*
Rec., Australia-N.Z.
- N. (Notocallista)**. Large, periostracum thick. *Rec.*,
Australia.—FIG. E144,5. **N. (N.) kingi*
(GRAY); *5a-c*, RV hinge, LV int., ext., $\times 0.5$
(597).
- N. (Fossacallista)** MARWICK, 1938 [**Paradione*
parki MARWICK, 1926; OD]. Ligament deeply
sunken (597). *Oligo.-Mio.*, N.Z.
- N. (Striacallista)** MARWICK, 1938 [**Cytherea mul-*
istriata SOWERBY, 1852; OD]. Small, periostra-
cum thin, extremities of shell concentrically
grooved (597). *Oligo.-Rec.*, N.Z.-Australia.
- Pelecycora** DALL, 1902 [**Cytherea hatchetigbeensis*
ALDRICH, 1886; OD] [= *Sinodia* JUKES-BROWNE,
1908 (type, *Artemis trigona* REEVE, 1850; OD)].
Ovately trigonal, inflated, lunule faint. *Eoc.-Rec.*,
Eu.-Asia-N.Am.-Afr.
- P. (Pelecycora)**. Pallial sinus long, angular, as-
cending; cardinal teeth divergent, *All* well re-
moved, near lower margin of hinge plate. *Eoc.-*
Rec., Eu.-Asia-N.Am.—FIG. E144,1. **P. (P.)*
hatchetigbeensis (ALDRICH), *Eoc.*, USA (Ala.);
1a,b, RV int., LV int., $\times 1$ (711).
- P. (Cordiopsis)** COSSMANN, 1910 [**Venus increas-*

Veneroida—Veneracea



- sata* SOWERBY, 1817 (non BROCCHI, 1814) = *V. suborbicularis* GOLDFUSS, 1841; OD] [=? *Cyprinopsis* CONRAD, 1869 (non FITZINGER, 1832) (type, *Artemis elliptica* SOWERBY in SMITH, 1847 = *Cyprina isocardoides* LAMARCK, 1818; OD; Mio., Eu.)]. Pallial sinus moderate to short, rounded, cardinal teeth not markedly divergent, *All* near *2a*, not marginal (912). *Eoc.-Mio.*, Eu.-Afr.-Asia.
- P. (Sinodiopsis)** EAMES, 1957 [**S. coxi*; OD]. Hinge plate thinner than in *P. (Pelecycora)*, *2a* not grooved. *Eoc.*, W.Afr.
- Pseudamiantis** KURODA, 1933 [**Meretrix tauyensis* YOKOYAMA, 1927; OD]. Resembling *Amiantis* but concentric sculpture more regular, intersected by fine radial lines; pallial sinus rounded (365). *Plio.*, Japan.
- Pseudaphrodina** CASEY, 1952 [**Venus ricordeana* D'ORBIGNY, 1845; OD]. Compressed, with carinate escutcheon; cardinals widely diverging. *L.Cret.*, Eu.—FIG. E145,3. **P. ricordeana* (D'ORBIGNY), Eng.; *3a,b*, LV int., RV ext., $\times 1$ (92).
- Resatrix** CASEY, 1952 [**R. dolabra*; OD]. Trigonal ovate, smooth or concentrically striate, lunule faint; hinge with *All* scarcely detached from or joined to *2a*. *L.Cret.*, Eu.
- R. (Resatrix)**. *All* and *Alll* long, laminar, *Pll* also present. *L.Cret.*, Eu.—FIG. E145,2. **R. (R.) dolabra* (CASEY), Eng.; *2a-c*, LV hinge, RV int., LV ext., $\times 1$ (92).
- R. (Dosiniopsella)** CASEY, 1952 [**R. (D.) cantiana*; OD]. With *2b* bifid, *All* striate (92). *L.Cret.*, Eu.
- R. (Vectorbis)** CASEY, 1952 [**Venus vectensis* FORBES, 1845; OD]. Laterals present in young but weak to absent in adults (92). *L.Cret.*, Eu.
- Rohea** MARWICK, 1938 [**Paradione trigonalis* MARWICK, 1927; OD]. Sculpture of concentric grooves; hinge plate flat, *3a* long, *All* distant from cardinals (597). *Oligo.*, N.Z.
- Saxidomus** CONRAD, 1837 [**S. nuttalli*; M] [= *Exocallista* KAMADA, 1962 (ex KIRA, 1955, *nom. invalid.*) (type, *S. brevisiphonatus* CARPENTER, 1865; OD)]. Large, oval, concentrically corrugated, without lunule or escutcheon; hinge curved, with irregular teeth (228). *U.Oligo.-Rec.*, W.N.Am.-Japan.—FIG. E144,8. **S. nuttalli*, Rec., USA (Calif.); *8a,b*, LV ext., RV int., $\times 0.5$ (124b).
- Trigonocallista** RENNIE, 1930 [**Meretrix umzambiensis* WOODS, 1906; OD]. Trigonal; lunule and large escutcheon well marked; *3b* entire, *All* very long, nymph rugose (92). *U.Cret.*, S.Afr.—FIG. E145,5. **T. umzambiensis* (WOODS); *5a,b*, LV ext., int., $\times 0.5$ (WOODS, 1906).

Subfamily SAMARANGIINAE Keen,
new subfamily

Shells inequilateral, quadrate, resembling some Arctiidae, surface unsculptured;

hinge veneroid, with *All* pustular, *2b* tending to be deeply bifid; pallial line entire. *Rec.*

Samarangia DALL, 1902 [**Venus quadrangularis* ADAMS & REEVE, 1850; OD] [= *Petroderma* KURODA, 1945 (type, *P. thaanumi*; OD)]. Periostracum present, encrusted with agglutinated sandy material. *Rec.*, E.Indies.—FIG. E146,7. **S. quadrangularis* (ADAMS & REEVE), E.China Sea; *7a-c*, LV int., RV int., RV ext., $\times 0.5$ (Kuroda, 1945).

Subfamily DOSINIINAE Deshayes, 1853

[*nom. correct.* ADAMS & ADAMS, 1858 (ex Dosiniana DESHAYES, 1853)]

Equivalve, lenticular, concentrically striate; hinge strong, with *All* present. *U.Cret.-Rec.*

Dosinia SCOPOLI, 1777 [**Chama dosin* ADANSON, 1757 (= *Venus concentrica* BORN, 1778, fide FISCHER-PIETTE, 1942); M] [= *Dosinidia* DALL, 1902 (obj.); *Bonartemis* IREDALE, 1929 (type, *B. stabilis*; OD); *Pardosinia* IREDALE, 1929 (type, *P. colorata*; OD); *Fallartemis* IREDALE, 1930 (type, *F. armina*; OD); *Meridosinia* IREDALE, 1930 (type, *M. nedigna*; OD); *Semelartemis* IREDALE, 1930 (type, *S. aetha*; OD)]. Compressed, nearly orbicular; lunule well defined. *L.Eoc.-Rec.*, cosmop.

D. (Dosinia). With periostracum; escutcheon wanting; sculpture of concentric grooves, none lamellose. *L.Oligo.-Rec.*, tropics-Atl.-Pac.—FIG. E146,3. **D. (D.) concentrica* (BORN), Rec., USA (Fla.); *3a,b*, LV ext., int., $\times 0.5$ (711).

D. (Asa) BASTEROT, 1825 [**Venus lincta* PULTENEY, 1799 (= *V. lupinus* LINNÉ, 1758, fide FISCHER-PIETTE, 1942); M?] [= *Arctoë* RISSO, 1826 (obj.); *Dosinia, s.s.*, AUCT. (non SCOPOLI)]. Lunule deep, escutcheon narrow, bordered by lamellose sculpture (329a). *Mio.-Rec.*, Eu.-W.Afr.

D. (Austrodosinia) DALL, 1902 [**Cytherea anus* PHILIPPI, 1848; OD]. Escutcheon with strong crest; pallial sinus horizontal (228). *U.Oligo.-Rec.*, N.Z.

D. (Dosinella) DALL, 1902 [**Cytherea angulosa* PHILIPPI, 1847; OD]. Lunule shallow; pallial sinus deep (228). *Rec.*, SW.Pac.

D. (Dosinisca) DALL, 1902 [**Artemis alata* REEVE, 1859; OD]. Lunule and escutcheon pouting; radial sculpture sometimes present (288). *Rec.*, N.Pac.-S.Pac.

D. (Dosinorbis) DALL, 1902 [**Artemis bilunulata* GRAY, 1838; OD]. Lunule bounded by raised cords (228). *Rec.*, Japan.

D. (Kakahuia) MARWICK, 1927 [**D. (K.) suteri*; OD]. Roundly trigonal; lunule shallow (592). *M.Eoc.*, N.Z.

D. (Kanecharaia) MAKIYAMA, 1936 [**D. kanecharai* YOKOYAMA, 1926; OD]. Cardinal teeth not bifid (392a). *Mio.*, Korea.




FIG. E146. Veneridae (Samarangiinae) (7), (Dosiniinae) (3-4), (Cycliniinae) (1-2,5-6) (p. N679-N681).

D. (*Kereia*) MARWICK, 1927 [**D. greyi* ZITTEL, 1864; OD]. Sculpture of spaced sharp lamellae (592). *L.Eoc.-Rec.*, N.Z.

D. (*Pectunculus*) DA COSTA, 1778 [**P. capillaceus* (= **Venus exoleta* LINNÉ, 1758; SD JUKES-BROWNE, 1911)] [= *Arthemis*, *Arthemiderma* POLI, 1795 (obj.; M) (*Artemis* AUCTT., nom. van.); *Orbiculus* MEGERLE VON MÜHLFELD, 1811 (obj.; SD DALL, 1902); *Exoleta* BROWN, 1827 (obj.; T); *Cerana* GISTEL, 1848 (obj.; pro *Artemis*); *Ampithaea* GRAY (ex LEACH MS), 1852 (obj.; M)]. Escutcheon wanting; pallial sinus narrow. *Oligo.-Rec.*, S.Eu.

D. (*Phacosoma*) JUKES-BROWNE, 1912 [**Artemis*

japonica REEVE, 1850; OD]. Escutcheon pouting, defined by raised ridges; margins of RV grooved posteriorly (329a). *Rec.*, N.Pac.-S.Pac.

D. (*Raina*) MARWICK, 1927 [**D. (R.) bensoni*; OD]. Lunule shallow; 3b much broader than 1 (592). *M.Oligo.-M.Plio.*, N.Z.

Dosinobia FINLAY & MARWICK, 1937 [**Dosinia ongleyi* MARWICK, 1927; OD]. Resembling *Dosinia* but with less sunken lunule; hinge teeth are more spread out, *All* scarcely separate from 2a (695). *U.Cret.*, N.Z.—FIG. E146,4. **D ongleyi* (MARWICK), *Cret.*, N.Z.; 4a-c, LV and RV hinges, RV ext., $\times 1$ (592).

Subfamily CYCLININAE Frizzell, 1936

Like *Dosiniinae* in form but without anterior lateral teeth or incised lunule; sculpture concentric with few faint radial traces. *L.Cret.-Rec.*

Cyclina DESHAYES, 1850 [**Venus sinensis* GMELIN, 1791; SD DALL, 1902] [= *Eocyclina* DALL, 1908 (obj.)]. Sculpture weak; lunule faint or absent. *Oligo.-Rec.*, Asia.

C. (*Cyclina*). No lunule or escutcheon; inner ventral margin crenate. *Oligo.-Rec.*, Asia (228).—FIG. E146,2. **C. (C.) sinensis* (GMELIN), *Rec.*, China; 2*a,b*, LV ext., RV int., $\times 0.5$ (124*b*).

C. (*Cyclinorbis*) MAKIYAMA, 1926 [**C. (C.) lunulata*; OD]. With faint lunular area, beaks more central than in *C. (Cyclina)*, sculpture finer, entirely concentric; inner ventral margin smooth (329*a*). *L.Mio.*, Korea.

Cyclinella DALL, 1902 [**Dosinia tenuis* RÉCLUZ, 1852; OD]. Margins smooth, faint lunular area present; 3*b* bifid. *Eoc.-Rec.*, E.N.Am.-W.N.Am.—FIG. E146,6. **C. tenuis* (RÉCLUZ), *Rec.*, Carib.; 6*a,b*, RV int., LV int., $\times 1$ (711).

Cyprimeria CONRAD, 1864 [**Cytherea excavata* MORTON, 1833; M]. Subcircular, valves flattened, bent to left posteriorly; lunule wanting, escutcheon deep; sculpture weakly concentric; cardinal teeth diverging, nymphs rugose; pallial line hardly sinuated; inner margins smooth. *L.Cret.-Eoc.*, N.Am.-Eu.—FIG. E146,1. **C. excavata* (MORTON), *Cret.*, USA(N.J.); 1*a-c*, LV ext., RV hinge, LV int., $\times 0.5$ (Weller, 1907).

Frigichione FLETCHER, 1938 [**Chione permagna* TATE, 1900; OD]. Concentric sculpture strong but not lamellar; middle layer of shell with radial ribs; pallial sinus shallow. *U.Tert.*, Antarctic.

F. (*Frigichione*). Hinge relatively heavy and massive, 2*a* stronger than 2*b* (695). *U.Tert.*, Antarctic.—FIG. E146,5. **F. (F.) permagna* (TATE); 5*a,b*, LV ext., int., $\times 0.8$ (Fletcher, 1938).

F. (*Paleomarcia*) FLETCHER, 1938 [**P. tatei*; OD]. Hinge thinner than in *F. (Frigichione)* (Fletcher, 1938). *U.Tert.*, Antarctic.

Luciplota OLSSON, 1942 [**L. panamensis*; OD]. Margins smooth; no lunule; hinge as in *Cyclina* but without ligamental nymphs (695). *L.Plio.*, C.Am.

Subfamily GEMMINAE Dall, 1902

Small, polished, with marginal grooves and denticles simulating posterior and anterior lateral teeth; inner ventral margin crenulate. *Eoc.-Rec.*

Gemma DESHAYES, 1853 [**Venus gemma* TOTTEN, 1834; T] [= *Totteniana* PERKINS, 1869 (obj.)]. Ovate, pallial sinus triangular (695). *Eoc.-Rec.*, E.N.Am.—FIG. E147,6. **G. gemma* (TOTTEN),

Rec., W.Atl.; 6*a,b*, RV ext., int., $\times 10$ (Dall, 1900).

Parastarte CONRAD, 1862 [**Astarte triquetra* CONRAD, 1846; OD] [= *Callicistronia* DALL, 1883 (obj.)]. Trigonal, ligament high, beneath beaks; pallial sinus small (711). *Mio.-Rec.*, E.N.Am.—FIG. E147,1. **P. triquetra* (CONRAD), *Rec.*, W.Atl.; 1*a-c*, LV int., ext., RV int., $\times 16$ (Dall, 1883).

Plesiastarte FISCHER, 1887 [*pro Anomala* COSSMANN, 1886, non VON BLOCK, 1799] [**Cyrena crenulata* DESHAYES, 1858; OD]. Subtrigonal, concentrically striate; inner margins minutely crenulate; pallial line hardly flexuous. *Eoc.*, Eu.—FIG. E147,5. **P. crenulata* (DESHAYES), France; 5*a-c*, RV ext., RV int., LV hinge, $\times 7$ (Deshayes, 1858).

?*Rohini* SEMPER, 1862 (*nom. dub.*) [**"Woodia lamellosa* SANDBERGER" (?*nom. nud.*; = *Venus woodiaeformis* SANDBERGER, 1863); M]. Resembling *Digitaria* in Astartidae but with a veneroid hinge and small pallial sinus; shell surface striate to cancellate. *Oligo.*, Eu.

Subfamily CLEMENTIINAE Frizzell, 1936

[*nom. transl.* KEEN, 1951 (*ex* Clementiidae FRIZZELL, 1936)]

Shell thin, inequilateral; without escutcheon; sculpture subdued or wanting; inner ventral margin smooth; hinge without lateral teeth. *U.Cret.-Rec.*

Clementia GRAY, 1842 [**Venus papyracea* GRAY, 1825; SD GRAY, 1847] [= *Blainvillia* HUPÉ, 1854 (non ROBINEAU-DESVOIDY, 1830)]. Ovate, concentrically waved; no lunule; pallial sinus deep. *Eoc.-Rec.*, N.Am.-Eu.-Asia-Afr.-Pac.

C. (*Clementia*). Hinge thin with deep niche in front of anterior cardinals; pallial sinus wide (711). *Eoc.-Rec.*, N.Am.-Eu.-Asia-Afr.—FIG. E147,2. **C. (C.) papyracea* (GRAY), *Rec.*, Pac.; 2*a,b*, LV ext., RV int., $\times 0.5$ (124*b*).

C. (*Egesta*) CONRAD, 1845 [**Venus inoceriformis* WAGNER, 1840; M]. Like *C. (Clementia)* but larger, heavier, pallial sinus narrower; posterior slope truncate (711). *Mio.-Rec.*, E.N.Am.-W.N.Am.-Japan.

C. (*Terentia*) JUKES-BROWNE, 1914 [**C. granulifera* SOWERBY, 1852; OD] [= *Euterentia* COSSMANN, 1920 (obj.)]. With irregular divaricate or reticulate striae (455*a*). *Rec.*, Pac.

Compsomyax STEWART, 1930 [**Clementia subdiphana* CARPENTER, 1864; OD]. Shell thicker than in *Clementia*, undulating sculpture wanting; hinge with 3*b* strongly bifid. *Eoc.-Rec.*, W.N.Am.-S.Am.-Japan.—FIG. E147,7. **C. subdiphana* (CARPENTER), *Rec.*, USA(Wash.); 7*a,b*, RV int., ext., $\times 0.5$ (219; Jukes-Browne, 1913).

Psathura DESHAYES, 1858 [**Erycina fragilis* LAMARCK, 1805; M]. Hinge teeth small, 3*b* bifid and 1 grooved; pallial sinus wanting. *Eoc.*, Eu.—FIG. E147,4. **P. fragilis* (LAMARCK), France; 4*a-c*, RV and LV hinges, LV ext., $\times 1$ (124*b*).




FIG. E147. Veneridae (Gemminae) (1,5-6), (Clementiinae) (2-4,7) (p. N681-N682).

Thetiopsis MEEK, 1875 [*Venus circularis* MEEK & HAYDEN, 1856; M]. Small, sculpture faint, concentric only; pallial sinus sharply ascending; hinge with *2b* stout. *U.Cret.*, N.Am.—FIG. E147,3. **T. circularis* (MEEK & HAYDEN), USA(Mont.); *3a-c*, LV int., RV and LV hinges, $\times 1$ (Keen, Stanford Univ. Coll.; Meek, 1876).

Subfamily TAPETINAE Adams & Adams, 1857

Ovate to elongate, shell surface somewhat polished, inner margins smooth on at least posterior third; hinge plate narrow, with cardinals *3a* entire, *3b* normally entire, others frequently bifid; lateral teeth wanting. *L.Cret.-Rec.*

Tapes MEGERLE VON MÜHLFELD, 1811 [*Venus literata* LINNÉ, 1758; M] [= *Parembola* RÖMER, 1857 (obj.; M)]. Oblong, rhomboidal, somewhat

compressed, higher posteriorly; sculpture weak; lunule incised, escutcheon bordered by low carina. *Mio.-Rec.*, E.Indies-Afr.-Eu.-Pac.-Asia.

T. (Tapes). Surface concentrically grooved. *Pleist.-Rec.*, W.Pac.—FIG. E148,4. **T. (T.) literatus* (LINNÉ), Rec., E.Indies; *4a-c*, LV ext., RV and LV hinges, $\times 0.5$ (124b).

T. (Myrsopsis) SACCO, 1900 [*Venerupis pernarum* BONELLI in MICHELOTTI, 1839; OD]. Ovoid; sculpture irregular; ligament in deep pit excavated in nymph. *Mio.-Plio.*, Eu.

T. (Ruditapes) CHIAMENTI, 1900 [*Venus decussata* LINNÉ; SD DALL, 1902] [= *Amygdala* RÖMER, 1857 (*non* GRAY, 1825); M]. Sculpture weakly decussate posteriorly, obscure anteriorly. *Mio.-Rec.*, Eu.-W.Pac.

T. (Siratoria) OTUKA, 1937 [*Paphia siratoriensis* OTUKA, 1934; OD]. Radial and concentric ribs coarse, of equal strength. *Mio.*, Japan.

- T. (Taurotapes)** SACCO, 1900 [**Venus craverii* MICHELOTTI, 1847; OD]. Elongate, concentrically with irregular striae; left cardinal teeth not joined above, *1* bifid, *3b* widely divergent. *Mio.*, Italy.
- Cyclorisma** DALL, 1902 [*pro Cyclothyris* CONRAD, 1875 (*non* M'COY, 1844)] [**Cyclothyris carolinensis* CONRAD, 1875; M]. Ovate to subcircular, concentrically striate; hinge with *2a* and *2b* of equal size, *3b* deeply bifid; pallial sinus narrow, deep. *U.Cret.*, E.N.Am.—FIG. E148,10. **C. carolinensis* (CONRAD), *U.Cret.*, USA(N.Car.); *10a,b*, LV int., ext., $\times 0.5$ (Dall).
- Cyclorisma** MARWICK, 1927 [**C. woodsii*; OD]. Like *Dosinia* but without anterior lateral teeth; anterior cardinals divergent. *U.Cret.*, N.Z.—FIG. E148,9. **C. woodsii*; *9a-c*, RV, LV, and RV hinges, $\times 1$ (592).
- Eumarcia** IREDALE, 1924 [**Venus fumigata* SOWERBY, 1853; OD]. Moderate in size to large; lunule weakly incised, escutcheon not defined; inner margins smooth, pallial sinus ample. *L.Oligo.-Rec.*, N.Z.-Australia.
- E. (Eumarcia)**. Surface smooth. *M.Oligo.-Rec.*, N.Z.-Australia.
- E. (Atamarcia)** MARWICK, 1927 [**E. (A.) sulcifera*; OD]. Concentric grooves well spaced, polished. *L.Oligo.-M.Plio.*, N.Z.—FIG. E148, *6a,b*, **E. (A.) sulcifera*, *U.Oligo.*; *6a,b*, LV ext., RV hinge, $\times 0.5$ (592).—FIG. E148, *6c*. *E. (A.) curia* (HUTTON), *U.Oligo.*; LV hinge, $\times 0.5$ (592).
- E. (Opimarcia)** MARWICK, 1948 [**E. (O.) healyi*; OD]. Markedly inflated; sculpture fine and wavy. *Plio.*, N.Z.
- Eurhomalea** COSSMANN, 1920 [*pro Rhomalea* JUKES-BROWNE, 1914 (*non* KOCH, 1837)] [**Venus rufa* LAMARCK, 1818; OD]. Quadrate, nearly smooth, with faint radial striae; inner margins smooth. *Rec.*, W.S.Am.
- Flaventia** JUKES-BROWNE, 1908 [**Venus ovalis* SOWERBY, 1827; OD]. Ovate, lunule well defined; *3b* long and deeply bifid, its laminae of unequal length (228). *U.Cret.*, Eu.-W.N.Am.—FIG. E148, 2. **F. ovalis* (SOWERBY), Eng.; *2a-d*, RV int., LV ext., both valves dorsal, LV hinge, $\times 1$ (1008).
- Gomphina** MÖRCH, 1853 [**Venus undulosa* LAMARCK, 1818; M]. Trigonal, nearly equilateral, smooth; dorsal margins grooved and beveled beyond hinge plate; pallial sinus short, rounded. *L.Plio.-Rec.*, W.Pac.-Australia-N.Z.-Japan-S.Atl.
- G. (Gomphina)**. Valves relatively solid, tumid; hinge with *4b* and right nymph rugose (228). *Rec.*, W.Pac.-Australia.
- G. (Gomphinella)** MARWICK, 1927 [**G. maorum* SMITH, 1902; OD]. Small, beaks subcentral. *L.Plio.-Rec.*, N.Z.—FIG. E148,3. **G. (G.) maorum* SMITH, Rec.; *3a-c*, RV ext., RV int., LV int., $\times 4$ (592).
- G. (Jukesena)** IREDALE, 1915 [*pro Acolus* JUKES-BROWNE, 1913 (*non* FÖRSTER, 1856)] [**Psephis foveolata* COOPER & PRESTON, 1910; OD]. Small, pallial line only slightly flexed. *Rec.*, S.Atl.
- G. (Macridiscus)** DALL, 1902 [**Donax aequilatera* SOWERBY, 1825; OD]. Thinner than *G. (Gomphina)*; nymphs smooth. *Pleist.-Rec.*, Japan.
- Irus** SCHMIDT, 1818 [*ex* OKEN, 1815; nonbinom., ICZN Op. 417] [**Donax irus* LINNÉ, 1758; T]. Subquadrate, with raised concentric lamellae intersected by weaker radial ribs; hinge weak, cardinals *1*, *2a*, *2b*, *3b* bifid; shell commonly distorted by growth in crevices. *Oligo.-Rec.*, Eu.-C.Am.-N.Am.-N.Z.
- I. (Irus)**. Lamellae well spaced. *Oligo.-Rec.*, Eu.-W.N.Am.—FIG. E148,8. **I. (I.) irus* (LINNÉ), Rec., France; *8a*, LV ext., $\times 1$; *8b,c*, RV and LV hinges, $\times 2$ (124*b*; Keen, n, Stanford Univ. Coll.).
- I. (Notirus)** FINLAY, 1928 [*pro Irona* FINLAY, 1927 (*non* SCHIÖDTE & MEINERT, 1883)] [**Venerupis reflexa* GRAY, 1843; OD]. Concentric lamellae closely spaced. *Pleist.-Rec.*, N.Z.
- I. (Notopaphia)** OLIVER, 1923 [**Venerupis elegans* DESHAYES, 1854; OD]. Radial sculpture fine, concentric lamellae closely spaced; lunule incised; inner ventral margin crenulate anteriorly; pallial sinus nearly horizontal, angular. *Pleist.-Rec.*, N.Z.
- I. (Paphonotia)** HERTLEIN & STRONG, 1948 [**Petricolaria* SOWERBY, 1834; OD]. Like *I. (Notopaphia)* but radial sculpture coarser, concentric lamellae wider spaced; hinge with *2a* large; pallial sinus ascending. *Rec.*, W.C.Am.
- Katelsysia** RÖMER, 1857 [**Venus scalarina* LAMARCK, 1818; SD KOBELT, 1881] [= *Catelsysia* AUCTION, *nom. null.*]. Concentric ribbing irregular anteriorly, radial sculpture weak; pallial sinus medium-sized to short; cardinal teeth radiating. *?Paleoc.*; *Eoc.-Rec.*, Eu.-N.Am.-Pac.-Japan-Australia.
- K. (Katelsysia)**. Compressed, markedly inequilateral; pallial sinus short. *?Paleoc.*; *Eoc.-Rec.*, Eu.-E.N.Am.-Pac.—FIG. E148,5. **K. (K.) scalarina* (LAMARCK), Rec., Australia; *5a-c*, LV ext., RV int., LV int., $\times 1$ (711).
- K. (Nipponomarcia)** IKEBE, 1941 [**K. (N.) nakamurai*; OD]. Hinge as in *K. (Katelsysia)* but otherwise resembling *Eumarcia*. *Mio.*, Japan.
- K. (Textivenus)** COSSMANN, 1886 [**Venus texta* LAMARCK, 1905; SD DALL, 1902]. Sculpture reticulate to zigzag. *Eoc.*, Eu.-E.N.Am.
- Legumen** CONRAD, 1858 [**L. ellipticus*; SD STOLICZKA, 1871] [= *Baroda* STOLICZKA, 1870 (type, *Venus fragilis* D'ORBIGNY, 1845; *non* FABRICIUS, 1780; OD)]. Elongate, slender, subelliptical, beaks at anterior fourth; sculpture concentric only; no lunule or escutcheon; hinge plate narrow; pallial sinus moderate. *L.Cret.-U.Cret.*, E.N.Am.-W.N.Am.-India-Eu.—FIG. E148,1. **L. ellipticus*, *U.Cret.*, USA(Miss.); *1a,b*, LV and RV hinges, $\times 1$; *1c*, RV ext., $\times 0.5$ (889).




FIG. E148. Veneridae (Tapetinae) (p. N682-N685).

Liocyma DALL, 1870 [*Venus fluctuosa* GOULD, 1841; OD]. Oblique-oval, surface concentrically striate; hinge with 1 and 2b feebly grooved, nymphs smooth; pallial sinus short, rounded. *Pleist.-Rec.*, N.Atl.-N.Pac.—FIG. E148,7. **L. fluctuosa* (GOULD), *Rec.*, NW.Atl.; RV int., $\times 1$ (Jukes-Browne, 1913).

Marcia H.ADAMS & A.ADAMS, 1857 [*Venus pinguis* CHEMNITZ" (*nonbinom.*) (= **V. opima* GMELIN, 1791); SD KOBELT, 1881] [= *Hemi-*

tapes RÖMER, 1864 (obj.; SD STOLICZKA, 1870); *Levimarcia* COX, 1927 (obj.; OD)]. Ovate-cordate, inflated, smooth; lunule convex, escutcheon depressed but not sharply delimited; cardinal teeth radiating, 1, 2b, 3b bifid; pallial sinus oval, horizontal. *Eoc.-Rec.*, N.Am.-Afr.-E.Indies-Australia-Eu.

M. (Marcia). Of medium size; pallial sinus well developed. *L.Mio.-Rec.*, Afr.-E.Indies.—FIG. E148,11. **M. (M.) opima* (GMELIN), *Rec.*,

- E.Indies; 11a,b, RV int., LV ext., $\times 1$ (Jukes-Browne, 1913).
- M. (Gomphomarcia)** KAUTSKY, 1929 [**G. abeli*; OD]. Subtrigonal, lunule and escutcheon well defined. *Mio.*, Eu.
- M. (Granicorium)** HEDLEY, 1906 [**G. indutum*; OD]. Lunule not well defined. *Rec.*, Australia.
- M. (Mercimonia)** DALL, 1902 [**Venus bernayi* COSSMANN, 1886; OD]. Hinge with 2b weak; pallial sinus small. *Eoc.-Mio.*, Eu.-E.N.Am.-W.N.Am.
- M. (Similivenerus)** COSSMANN, 1910 [**Venus solida* DESHAYES, 1824 (non SCHRÖTER, 1802) (= *S. insolida* KEEN, 1954); OD]. Small, lunule lanceolate, escutcheon beveled, teeth crowded. *Eoc.-Oligo.*, Eu.
- Paphia** RÖDING, 1798 [**P. alapapilionis* (= *Venus rotundata* LINNÉ, 1758); SD DALL, 1902] [= *Textrix* RÖMER, 1857 (non SUNDEVALL, 1833; obj., SD STOLICZKA, 1870); *Paratapes* STOLICZKA, 1870 (pro *Textrix*); *Eutapes* CHIAMENTI, 1900 (obj.; OD); *Acritopaphia* IREDALE, 1936 (type, *A. transfusa*; OD)]. Elongate, compressed, surface glossy. *Oligo.-Rec.*, Pac.-N.Z.-Asia-Eu.
- P. (Paphia)**. Nearly smooth, with faint concentric ribbing. *Plio.-Rec.*, W.Pac.—FIG. E149,2. **P. (P.) rotundata* (LINNÉ), *Rec.*; 2a-c, LV ext., RV and LV hinges, $\times 0.5$ (Marwick, Chenu).
- P. (Callistotapes)** SACCO, 1900 [**Venus vetula* BASTEROT, 1825; OD] [= *Hemitapes* AUCTT., non RÖMER]. Concentric ridges close, well defined. *Oligo.(Aquit.)-Rec.*, Eu.-Asia-N.Z.
- P. (Protapes)** DALL, 1902 [**Venus gallus* GMELIN, 1791; OD]. Concentric ribs evident; shell shorter and more tumid than *P. (Paphia)*. *Rec.*, W.Pac.
- Paraesa** CASEY, 1952 [**Venus faba* J. DE C. SOWERBY, 1827; OD]. Ovate, concentricly striate, lunule faint; resembling *Flaventia* but with 3b less deeply bifid, 2a not wedge-shaped, 2b reaching base of hinge plate. *L.Cret.(U.Alb.)-U.Cret.(Senon.)*, Eu.—FIG. E149,1. **P. faba* (SOWERBY), *L.Cret.*, Eng.; 1a-c, LV and RV hinges, LV ext., $\times 1$ (92).
- Psephidia** DALL, 1902 [pro *Psephis* CARPENTER, 1865 (non GUENÉE, 1854)] [**Chione lordi* BAIRD, 1863; OD] [non *Psephidia* POMEL, 1872 (nom. nud.)]. Small, ovate-trigonal, polished, sculpture faint, concentric; lunule narrow, feeble; escutcheon wanting; pallial sinus distinct, angular. *Plio.-Rec.*, W.N.Am.—FIG. E149,4. **P. lordi* (BAIRD), *Rec.*, USA(Wash.); 4a-c, LV ext., RV int., LV int., $\times 3$ (228).
- Sinonia** STEPHENSON, 1953 [**S. levis*; OD]. Elongate-ovate, nearly smooth, with irregular concentric ridges; no lunule or escutcheon; pallial sinus small, directed upward. *U.Cret.(Cenoman.)*, USA(Tex.).—FIG. E149,5. **S. levis*; 5a-c, RV int., LV hinge, ext., $\times 1$ (Stephenson, 1953).
- Venerella** COSSMANN, 1886 [**Venerupis hermonvillensis* DESHAYES, 1858; SD CROSSE, 1886]. Like




FIG. E149. Veneridae (Tapetinae) (p. N685-N686).

- Marcia* but hinge teeth divergent, extending beyond edge of narrow hinge plate (329a). *Eoc.*, Eu.-Asia.
- Veneritapes** COSSMANN, 1886 [**Psammobia bervillei* DESHAYES, 1858; M]. Smooth, anteriorly attenuated; teeth small, 3a weak or obsolete (329a). *Eoc.*, Eu.
- Venerupis** LAMARCK, 1818 [**Venus perforans* MONTAGU, 1803 (= *Venus saxatilis* FLEURIAU DE BELLEVUE, 1802); SD CHILDREN, 1823] [= *Pullastra* SOWERBY, 1826 (type, *Venus pullastra* MONTAGU, 1803; T); *Myrsus* H. ADAMS & A. ADAMS, 1853 (type, "*M. corrugata* DESHAYES," = *Venus corrugata* GMELIN, 1791; SD STOLICZKA, 1870); *Politiitapes* CHIAMENTI, 1900 (type, *Venus aurea* GMELIN, 1791; SD DALL, 1900); *Aureitapes*, *Cyaneitapes* CHIAMENTI, 1900 (type, *V. aurea* GMELIN, 1791; SD KEEN, 1954)]. Ovate-elliptic; sculpture of irregular radial and concentric or

slightly zigzag riblets, with stronger concentric lamellae at ends. *L.Eoc.-Rec.*, Eu.-Afr.-N.Z.

V. (*Venerupis*). Sculpture present throughout; distortion, due to growth in crevices, common. *Plio.-Rec.*, Eu.-W.Afr.—FIG. E149,3. *V. (*V.*) *saxatilis* (FLEURIAU DE BELLEVUE), *Rec.*, Eu.; 3*a*,*b*, RV ext., LV int., ×1 (Brown).

V. (*Paphirus*) FINLAY, 1927 [**Venus largillierti* PHILIPPI, 1847; OD]. Radial sculpture absent on posterior slope. *L.Eoc.-Rec.*, N.Z.

Subfamily CHIONINAE Frizzell, 1936

[*nom. transl.* KEEN, 1951 (*ex* Chionidae FRIZZELL, 1936)]

Ovate-trigonal, inequilateral, sculpture cancellate; lunule normally present; escutcheon, if present, beveled; inner margins usually crenulate; hinge plate and teeth well developed, cardinals 2*a* and 1 large; pallial sinus mostly short, ascending. *M.Eoc.-Rec.*

Chione MEGERLE VON MÜHLFELD, 1811 [**Venus dysera* LINNÉ (of GMELIN, 1791) (*non* LINNÉ, 1758) (= **V. cancellata* LINNÉ, 1767); SD GRAY, 1847]. Cordate; lunule and escutcheon mostly well defined; concentric sculpture strong, frilled in some forms. *Oligo.-Rec.*, N.Am.-S.Am.-C.Am.-N.Z.

C. (*Chione*). Pallial sinus small; cardinal teeth smoothly or only faintly grooved. *U.Oligo.-Rec.*, E.N.Am.-W.N.Am.-S.Am.—FIG. E150,1. *C. (*C.*) *cancellata* (LINNÉ), *Rec.*, Carib.; 1*a-c*, RV ext., int., LV int., ×1 (711).

C. (*Austrovenus*) FINLAY, 1927 [**Venus stutchburii* GRAY" (= **V. stutchburii* WOOD, 1828); OD]. Sculpture of low radial ribs crossed by irregular concentric lamellae (695). *M.Plio.-Rec.*, N.Z.

C. (*Chionista*) KEEN, 1958 [**Venus fluctifraga* SOWERBY, 1853; OD]. Lunule and escutcheon wanting; concentric sculpture irregular, beaded. *Pleist.-Rec.*, W.N.Am.

C. (*Chionopsis*) OLSSON, 1932 [**Venus amathusia* PHILIPPI, 1844; OD] [= *Gnidiella* PARKER, 1949 (type, *Venus gnidia* BRODERIP & SOWERBY, 1829; OD)]. Pallial sinus well developed; one or more cardinal teeth in each valve bifid or strongly grooved (329*a*). *Mio.-Rec.*, E.N.Am.-W.N.Am.

C. (*Iliochione*) OLSSON, 1961 [**Venus subrugosa* WOOD, 1828; OD] [= *Anomalocardia* AUCTT. (*non* SCHUMACHER)]. Sculpture of undulating concentric folds, mostly obsolete toward ventral margin. *Rec.*, W.C.Am.-S.Am.

C. (*Lirophora*) CONRAD, 1863 [**Circumphalus athleta* (= **Venus latilirata* CONRAD, 1841); SD DALL, 1902]. Concentric lamellae of broad thickened rolls that may coalesce medially. *Oligo.-Rec.*, E.N.Am.-W.N.Am.—FIG. E150,2. *C. (*L.*) *latilirata* (CONRAD), *Mio.*, USA(Md.); RV ext., ×1 (Glenn, 1904).

C. (*Panchione*) OLSSON, 1964 [**Grateloupia mactropsis* CONRAD, 1865; OD]. Intermediate between *C. (Lirophora)* and *C. (Iliochione)*, with foliaceous posterior area of former and outline of latter. *Mio.*, C.Am.—FIG. E150,3. *C. (*P.*) *mactropsis* (CONRAD), Panama; LV ext., ×1 (Olsson, 1964).

C. (*Securella*) PARKER, 1949 [**Venus securis* SHUMARD, 1858; OD]. Ligament deeply sunken; concentric sculpture underlain by radial, appearing cancellate on eroded surfaces; 2*b* deeply grooved with posterior part larger and higher than anterior (695). *Oligo.-Plio.*, NW.N.Am.

Anomalocardia SCHUMACHER, 1817 [**Venus flexuosa* LINNÉ, 1767; OD] [= *Triquetra* DE BLAINVILLE, 1828 (type, *Venus flexuosa* LINNÉ, 1767; SD ANTON, 1839); *Cryptogramma* MÖRCH, 1853 (obj.); *Murcia* RÖMER, 1857 (*non* KOCH, 1835)]. Shell thick, with undulating concentric folds crossed by radial riblets; lunule large, impressed. *Mio.-Rec.*, C.Am.-S.Am.-Afr.-E.Indies-W.Pac.

A. (*Anomalocardia*). Radial sculpture subordinate; 4*b* rugose. *Mio.-Rec.*, E.C.Am.-S.Am.-W.Afr.—FIG. E150,5. *A. (*A.*) *flexuosa* (LINNÉ), *Rec.*, W.Afr.; 5*a,b*, LV int., LV ext., ×1 (711).

A. (*Anomalodiscus*) DALL, 1902 [**Cytherea squamosa* LAMARCK" (= **Venus squamosa* LINNÉ, 1758); OD]. Sculpture reticulate, ventral margin crenulate, hinge teeth without rugosities (228). *U.Mio.-Rec.*, E.Indies.

A. (*Cryptonemella*) KURODA & HABE, 1951 [*pro Cryptonema* JUKES-BROWNE, 1914 (*non* BIGSBY, 1868)] [**A. (C.) producta (pro Venus impressa* ANTON, 1839; *non* DE SERRES, 1829); OD]. Sculpture weak, radial, internal margin nearly smooth (695). *Rec.*, W.Pac.

Bassina JUKES-BROWNE, 1914 [**Venus paucilamellata* "SOWERBY" (i.e., *Mercenaria paucilamellata* DUNKER, 1858) (= **V. pachyphylla* JONAS, 1839); OD]. Resembling *Chione* but without escutcheon; inner margins crenate; 2*b* large, 3*b* grooved. *Oligo.-Rec.*, E.Indies-Australia-N.Z.

B. (*Bassina*). Thick-shelled; sculpture of thin, spaced concentric lamellae (592). *Mio.-Rec.*, E.Indies-S.Australia.

B. (*Callanaitis*) IREDALE, 1917 [**Venus yatei* GRAY, 1835; OD] [= *Salacia* JUKES-BROWNE, 1914 (*non* LAMOUREUX, 1816) (type, *Venus lamellata* LAMARCK, 1818; OD); *Eusalacia* COSSMANN, 1920 (*pro Salacia*)]. Thin-shelled, somewhat quadrate, compressed, concentric lamellae strong. *Oligo.-Rec.*, N.Z.—FIG. E150,9. *B. (*C.*) *yatei* (GRAY), *Rec.*, N.Z.; 9*a-c*, LV int., ext., RV hinge, ×0.5 (592).

Chamelea MÖRCH, 1853 [**Venus gallina* LINNÉ, 1758; SD BUCQUOY, DAUTZENBERG, & DOLLFUS, 1893] [= *Ortygia* BROWN, 1827 (*non* BOIE, 1826) (obj.); SD HERRMANNSEN, 1847); *Hermione* GRAY, 1852, *ex* LEACH MS (*non* MEIGEN, 1800)]. With narrow close concentric lamellae, no radial sculp-

ture; cardinal teeth not bifid. *Oligo.-Rec.*, E.N.Am.-Eu.—FIG. E150,7. **C. gallina* (LINNÉ), *Rec.*, *Medit.*; 7a-c, LV ext., LV and RV hinges, $\times 1$ (711).

Clausinella GRAY, 1851 [**Chione fasciata*" (i.e., **Pectunculus fasciatus* DA COSTA, 1778); M] [=*Zucleica* GRAY, 1852, ex LEACH MS (obj.);

Anaitis RÖMER, 1857 (non DUPONCHEL, 1778)]. Sculpture of recurved, spaced concentric ridges; inner margins crenate. *Oligo.-Rec.*, Eu.-E.N.Am.-N.Z.—FIG. E150,4. **C. fasciata* (DA COSTA), *Rec.*, *Medit.*; 4a-c, LV ext., RV and LV hinges, $\times 1$ (711).

Hinemoana MARWICK, 1927 [**Chione accuminata*




FIG. 150. Veneridae (Chioninae) (p. N686-N688).

- HUTTON, 1873; OD]. Sculpture concentric, with fine subsurface radial threads; RV with grooved margin simulating posterior lateral tooth (592). ?*Oligo.*, N.Z.
- Humilaria** GRANT & GALE, 1931 [**Venus kenneleyi* REEVE, 1863; OD]. With heavy concentric ribs, no radial sculpture; inner margins crenate; 2a large, bifid in some; pallial sinus short, angular. *Plio.-Rec.*, W.N.Am.—FIG. E150,6. **H. kenneleyi* (REEVE), *Rec.*, USA (Wash.); 6a-c, RV ext., int., LV hinge, $\times 0.5$ (288; Keen, n, Stanford Univ. Coll.).
- Mercenaria** SCHUMACHER, 1817 [**M. violacea* (= *Venus mercenaria* LINNÉ, 1758); T] [= *Cras-sivenus* PERKINS, 1869 (obj.)]. Like *Chione* in form but larger, radial sculpture obsolete; ligamental nymphs rugose. *Oligo.-Rec.*, N.Am.-Japan.—FIG. E150,8. **M. mercenaria* (LINNÉ), *Rec.*, W.Atl.; 8a-c, LV ext., RV and LV hinges, $\times 0.5$ (711).
- Placamen** IREDALE, 1925 [**Venus placida* PHILIPPI, 1844; OD]. Concentric sculpture as in *Lirophora*; lunule deep, heart-shaped (329a). *Mio.-Rec.*, Australia.
- Protothaca** DALL, 1902 [**Chama thaca* MOLINA, 1782; OD]. Ovate, convex, sculpture reticulate; lunule normally present; escutcheon absent or in LV only; 2b and 1 bifid; pallial sinus moderate in size, pointed. *Mio.-Rec.*, N.Am.-S.Am.-Asia-Carib.-C.Am.
- P. (Protothaca)**. Radial sculpture predominating on anterior and middle slopes, concentric on posterior; inner margins crenate (228). *Mio.-Rec.*, N.Am.-S.Am.-Japan.—FIG. E150,10. **P. (P.) thaca* (MOLINA), *Rec.*, W.S.Am.; 10a,b, RV ext., int., $\times 0.7$ (Philippi, 1847).
- P. (Antinioche)** OLSSON, 1961 [**Nioche (A.) beili*; OD]. Sculpture as in *Periglypta* but hinge and escutcheon as in *P. (Leukoma)*. *Rec.*, W.C.Am.-S.Am.
- P. (Callithaca)** DALL, 1902 [**Tapes tenerrima* CARPENTER, 1857; OD]. Sculpture uniformly reticulate; lunule feeble, escutcheon wanting; inner margins smooth (228). *Mio.-Rec.*, W.N.Am.-Japan.
- P. (Colonche)** OLSSON, 1961 [**C. ecuadoriana*; OD]. Resembling *P. (Antinioche)* but lunule and escutcheon are wanting. Valves inflated. *Rec.*, W.S.Am.
- P. (Leukoma)** RÖMER, 1857 [**Venus granulata* GMELIN, 1791; SD KOBELT, 1881] [= *Nioche* HERTLEIN & STRONG, 1948 (type, *Venus asper- rima* SOWERBY, 1835; OD); not preoccupied by *Leucoma* HÜBNER, 1822]. Radial ribs fine, concentric sculpture reduced; lunule radially ribbed, incised; left valve with beveled escutcheon (329a). *Mio.-Rec.*, Carib.-E.C.Am.-W.C.Am.
- P. (Notochione)** HERTLEIN & STRONG, 1948 [**Venus columbiensis* SOWERBY, 1835; OD]. Ribs heavy, predominantly radial; shell rounded-trigonal (695). *Neog.-Rec.*, Japan-W.C.Am.-S. Am.
- P. (Novathaca)** HABE, 1951 [**Chione euglypta* SOWERBY, 1914; OD]. Radial sculpture weak, absent at ends of shell (365). *Rec.*, Japan.
- P. (Protocallithaca)** NOMURA, 1937 [**Venus adamsi* REEVE, 1863; OD]. Sinus shorter and more pointed than in *P. (Callithaca)* (365). *Plio.-Rec.*, Japan.
- P. (Tropithaca)** OLSSON, 1961 [**Venus grata* SAX, 1830; OD]. Resembling *P. (Protothaca)* but smaller, sculpture weaker, escutcheon reduced; color variability extreme. *Rec.*, W.C.Am.-S.Am.
- P. (Tuangia)** MARWICK, 1927 [**Venus crassicosta* DESHAYES, 1835; OD]. Radial ribs heavy, shell elongate to rounded-quadrate (592). *Plio.-Rec.*, Japan-N.Z.
- Tawera** MARWICK, 1927 [**Venus spissa* DESHAYES, 1835; OD]. Resembling *Chamelea* but lunule less deeply sunken; cardinal teeth more divergent. *L.Eoc.-Rec.*, N.Z.
- T. (Tawera)**. Sculpture of levelled smooth ridges. *L.Eoc.-Rec.*—FIG. E151,7. **T. (T.) spissa* (DESHAYES), *Rec.*, N.Z.; 7a-c, LV int., RV int., LV ext., $\times 1$ (592).
- T. (Turia)** MARWICK, 1927 [**Turia chattonensis*; OD]. Small, sculpture of thin, spaced concentric lamellae with irregular crests (592). *L.Eoc.-L.Mio.*, N.Z.
- Timoclea** BROWN, 1827 [**Venus ovata* PENNANT, 1777; M] [= *Pasiphæ* GRAY (ex LEACH, MS), 1852 (obj.); *Parvivenus* SACCO, 1900 (type, *Venus marginata* HÖRNES, 1861; OD); *Veremolpa* IREDALE, 1930 (type, *V. ethica*; OD)]. Small, rather compressed, sculpture predominantly radial; escutcheon smooth. *U.Oligo.-Rec.*, Eu.-C.Am.-W. Pac.-Australia-E.Indies.
- T. (Timoclea)**. Concentric sculpture present as nodes on ribs. *U.Oligo.(Aquitana)-Rec.*, Eu.-E.C.Am.-W.C.Am.-W.Pac.—FIG. E151,5. **T. (T.) ovata* (PENNANT), *Rec.*, *Medit.*; 5a-b, RV int., LV int., $\times 2$; 5c-e, RV ext., RV and LV hinges, $\times 1.3$ (711; Keen, n, Stanford Univ. Coll.).
- T. (Chioneryx)** IREDALE, 1924 [**Venus striatissima* SOWERBY, 1853 (= *Erycina cardioides* LAMARCK, 1818); M]. Substrate posteriorly; lunule striate, colored (329a). *Rec.*, Australia.
- T. (Glycydonta)** COTTON, 1936 [**Venus marica* LINNÉ, 1758; OD]. Concentric sculpture of raised lamellae on posterior slope (695). *Pleist.-Rec.*, E.Indies-Australia.

Family PETRICOLIDAE Deshayes, 1839

Oval, with no lunule or escutcheon; sculpture radial or obsolete. Hinge without lateral teeth; three cardinals in left valve, 2b bifid, right valve with 1 and 3b only; pallial sinus well developed. *Eoc.-Rec.*

Petricola LAMARK, 1801 [**P. costata* (= **Venus lapicida* GMELIN, 1791); SD SCHMIDT, 1818] [= *Naranio* GRAY, 1853 (obj.; SD LAMY, 1922)]. With radial sculpture, at least in young; commonly

distorted by nestling. *Eoc.-Rec.*, N.Am.-W.Pac.-Australia-Asia-Eu.-Carib.

P. (Petricola). Sculpture of fine radial, divaricate, or zigzag riblets (515). *Rec.*, E.N.Am.-W.N.Am.-W.Pac.—FIG. E151,3. **P. (P.) lapicida* (GMELIN), Carib.; 3a,b, LV ext., RV int., $\times 1$ (Brown, 1844).

P. (Claudiconcha) FISCHER, 1887 [**Venus monstrosa* GMELIN, 1791; M]. Sculpture cancellate, lamellose posteriorly; posterior margin thick; RV may overlap LV ventrally. *Rec.*, W.Pac.-Australia.

P. (Lajonkairia) DESHAYES, 1854 [**Venerupis decussata* (= **V. lajonkairii* PAYRAUDEAU, 1826); T] [= *Lajonkairia* AUCTT. (nom. null.)]. Sculpture cancellate; boring habit not developed. *Oligo.-Rec.*, Eu.-Japan.—FIG. E151,2. **P. (L.) lajonkairii* (PAYRAUDEAU), *Rec.*, Medit.; 2a,b, LV and RV hinges, $\times 2$ (515).

P. (Petricolaria) STOLICZKA, 1870 [**P. pholadiformis* LAMARCK, 1818; OD] [= *Gastranella* VERRILL, 1872 (type, *G. tumida*; M)]. Elongate, with nodose radial sculpture; hinge plate narrow, teeth small, 2a may be obsolete (515). *Oligo. (Aquitain.)-Rec.*, Eu.-E.N.Am.-W.N.Am.—FIG. E151,6. **P. (P.) pholadiformis* (LAMARCK), *Rec.*, E.N.Am.; RV int., $\times 1$ (1007).

P. (Petricolirus) HABE, 1951 [**P. aequistriata* SOWERBY, 1874; OD]. Radial striae subequal, numerous, over entire shell; posterior end truncate (365). *Rec.*, Japan.

P. (Pseudoirus) HABE, 1951 [**P. mirabilis* DESHAYES, 1853; OD]. Shell thick, compressed, not burrowing (365). *Rec.*, Japan.

P. (Rupellaria) FLEURIAU, 1802 [**Venus lithophaga* RETZIUS, 1786; M] [= *Choristodon* JONAS, 1844 (type, *C. typicum*; M)]. Sculpture of coarse radial riblets. *Eoc.-Rec.*, Eu.-E.N.Am.-W.N.Am.-W.Pac.-Japan.—FIG. E151,4. **P. (R.) lithophaga* (RETZIUS), *Rec.*, Medit.; 4a, RV int., $\times 1$; 4b,c, RV and LV hinges, $\times 2$ (515, 1007).

P. (Velargilla) IREDALE, 1931 [**Naranio rubiginosa* ADAMS & ANGAS, 1863; OD]. Thinner than *P. (Petricola)*; burrowing in ooze, not boring (695). *Rec.*, Australia.

Mysia LAMARCK, 1818 (ex LEACH MS) [**Venus undata* PENNANT, 1777; M] [= *Lucinopsis* FORBES & HANLEY, 1848 (obj.)]. Smooth, resembling *Cyclinella* but with only 2 cardinal teeth in RV (515). *Rec.*, Eu.-Australia.—FIG. E151,8. **M. undata* (PENNANT), *Rec.*, Eu.; RV int., $\times 1$ (695).

Pleiorytis CONRAD, 1862 [**P. ovata* (= *Petricola centenaria* CONRAD, 1833); M]. Shell thin, hinge with 3b weak to obsolete. *Mio.*, E.N.Am.—FIG. E151,1. **P. centenaria* (CONRAD), USA (Md.); 1a,b, LV int., RV ext., $\times 0.5$ (Glenn, 1904).

Family COOPERELLIDAE Dall, 1900

Shell thin, ovate-quadrate, moderately inflated; hinge without lateral teeth, right




FIG. E151. Veneridae (Chioninae) (5,7); (Petricolidae) (1-4,6,8) (p. N688-N689).




FIG. E152. Cooperellidae (1); Glauconomidae (2); Rzehakiidae (3) (p. N690).

valve with two thin cardinals, 1 and bifid 3b; left valve with three cardinals, 2b bifid; ligament depressed, seated on laminar nymph; pallial sinus wide. *Mio.-Rec.*

Cooperella CARPENTER, 1864, p. 639 [*Oedalia* (C.) *scintillaeformis*, p. 639 (= *O. subdiaphana*, p. 639); M (synonymy of generic and specific names established by first reviser, DALL, 1900) [= *Oedalia* CARPENTER, 1864 (non *Oedalea* MEIGEN, 1820); *Oedalina* CARPENTER, 1865 (pro *Oedalia*)]. *Mio.-Rec.*

C. (Cooperella). Hinge with 2b deeply divided; pallial sinus deep, its apex rounded. *Mio.-Rec.*, E.N.Am.-W.N.Am.-S.Am.—FIG. E152,1. ***C. (C.) subdiaphana** (CARPENTER), *Rec.*, USA (Calif.); 1a, RV int., $\times 1.3$; 1b,c, LV and RV hinges, $\times 2$ (515; Arnold, 1903).

C. (Cooperellopsis) WOODRING, 1925 [*C. (C.) thaumasta*; OD]. More quadrate and inflated than *C. (Cooperella)*; 2b not deeply bifid; pallial sinus obscure (695). *Mio.*, Carib.

Family GLAUCONOMIDAE Gray, 1853

[= *Glaucomyidae* CHENU, 1862; *Glaucomyidae*, *Glauconomidae* (spelling errors)]

Elongate shells, smooth, with conspicuous periostracum; hinge with three cardinal teeth in either valve, lateral teeth wanting. *Rec.*

Glaucnome GRAY, 1828 [**G. chinensis*; M] [= *Glaucomya* BRONN, 1838 (obj.); *Glaucomya* WOODWARD, 1854 (nom. van.); *Glaucnoma* CHENU, 1862 (nom. null.); *Glaucnometta* IREDALE, 1936 (type, *G. plankta*; OD); *Glaucnomella* ALLAN, 1950 (nom. null.)]. Periostracum light to dark green, smooth or wrinkled. [Habitat, marine to brackish water.] *Rec.*, IndoPac.-Australia. —FIG. E152,2. ***G. chinensis**, E. Indies; 2a,b, LV ext., int., $\times 1$ (Habe, 1954); 2c, RV int., $\times 1$ (305).

Family RZEHAKIIDAE Korobkov, 1954

[= *Onchophoridae* DAVIDASCHVILI, 1934]

Ovate, moderately small, strongly inequilateral, nearly smooth, beaks low; hinge lacking lateral teeth, with two cardinals in right valve (1 and 3b), and three in left (2a, 2b, and 4b); anterior adductor muscle scar deeply sunken, bordered by ridge posteriorly; pallial sinus shallow (716). [Apparently aberrant brackish-water descendants of marine Tapetinae.] *M.Mio.-U.Mio.*

Rzehakia KOROBKOV, 1954 [pro *Oncophora* RZEHAK, 1882 (non DIESING, 1851)] [**Oncophora socialis* RZEHAK, 1882; OD]. Cardinal teeth slightly bifid. *M.Mio.-U.Mio.*, E.Eu.—FIG. E152,3. ***R. socialis** (RZEHAK), *Mio.*, Czech.; 3a-c, RV int., ext., LV int., $\times 1$ (Rzehak).

Order MYOIDA Stoliczka, 1870

[nom. correct. NEWELL, 1965 (ex order Myacea STOLICZKA, 1870)] [Adapedonta AUCTION, in part] [Diagnosis by N. D. NEWELL]

Thin-shelled, burrowing forms with well-developed siphons; strongly inequilateral, equivalve or inequivalve; isomyarian or anisomyarian; one cardinal tooth in each valve, or edentulous; lunule and escutcheon absent, or poorly developed; shell not nacreous. *Carb.-Rec.*

Suborder MYINA Stoliczka, 1870

[nom. transl. et correct. NEWELL, 1965 (ex order Myacea STOLICZKA, 1870)] [Diagnosis by N. D. NEWELL]

Hinge edentulous or with one cardinal tooth on each valve; ligament external, borne on well-marked nymphs, in some forms with internal resilium; sinupalliate. *Perm.-Rec.*

Superfamily MYACEA Lamarck, 1809

[nom. transl. GILL, 1871 (ex family Myacea GOLDFUSS, 1820) (= *myaires* LAMARCK, 1809)] [Materials for this superfamily prepared by MYRA KEEN]




FIG. E153. Myidae (p. N691-N692).

Valves elongate or ovate, subequal, porcelaneous to chalky, with thin periostracum; hinge margin mostly without teeth; ligament mainly internal; valve margins smooth; pallial sinus, with some exceptions, well developed. *U. Jur.-Rec.*

Family MYIDAE Lamarck, 1809

[*nom. correct.* BRODERIP, 1839 (*pro* family Myacea GOLDFUSS, 1820) (=myaires LAMARCK, 1809)] [=Myacidae D'ORBIGNY, 1845]

Elongate, chalky, mostly burrowing forms with wide posterior gape; chondrophore present in LV. *Paleoc.-Rec.*

Mya LINNÉ, 1758 [**M. truncata*; SD CHILDREN, 1822 (ICZN Op. 94, Dir. 72)] [= *Hiatula* MODEER, 1793; SD WINCKWORTH, 1935 (obj.);

Myarius DUMÉNIL, 1806 (*nom. van.*); *Myes*, AUCT. (*nom. van.*)]. Of medium size, chondrophore in LV large; sculpture of concentric growth lines. *Oligo.-Rec.*, Eu.-N.Am.-Japan.

M. (Mya). Chondrophore with triangular anterior portion in front of anterior ridge. *Oligo.-Rec.*, Eu.-N.Am.-Japan.—FIG. E153, 10. **M. (M.) truncata* LINNÉ, Rec., Eng.; 10a-c, LV ext., LV and RV hinges, $\times 1$ (Jeffreys).

M. (Arenomya) WINCKWORTH, 1930 [**M. arenaria* LINNÉ, 1758; OD]. Chondrophore bounded anteriorly by ridge; posterior end of shell elongate-ovate. *Mio.-Rec.*, Eu.-N.Am.-Japan.—FIG. E153, 9. **M. (A.) arenaria* LINNÉ, Rec., N.Atl.; 9a-c, LV ext., int., RV hinge, $\times 0.5$ (H. Adams & A. Adams).

Cryptomya CONRAD, 1848 [**Sphaenia californica* CONRAD, 1837; M]. Hinge as in *Mya*, shell narrowly gaping, somewhat compressed; chondro-

- phore in RV appressed under beaks; shell thin and nearly smooth. *Mio.-Rec.*, N.Am.-S.Am.-Asia-Pac.
- C. (Cryptomya)**. Chondrophore smooth in front; pallial line entire. *Mio.-Rec.*, W.N.Am.-S.Am.-E. Asia.—FIG. E153,4. *C. (*C.*) *californica* (CONRAD), *Rec.*, USA (Calif.); 4a,b, RV ext., LV int., $\times 0.7$ (after Olsson, 1961).
- C. (Venatomya)** IREDALE, 1930 [*Sphaenia elliptica* A. ADAMS, 1851; OD]. Hinge with small 2a in front of chondrophore; pallial sinus of varying size. *Rec.*, Pac.—FIG. E153,2. *C. (*V.*) *elliptica* (A. ADAMS), Australia; 2a-c, RV ext., RV int., LV int., $\times 1$ (Hedley, 1913).
- Paramya** CONRAD, 1861 [*pro Myalina* CONRAD, 1845 (*non* DE KONINCK, 1842)] [*Myalina subovata* CONRAD, 1845; M]. Small, smooth, quadrate, with chondrophore in either valve. *Mio.-Rec.*, SE.N.Am.—FIG. E153,3. *P. *subovata* (CONRAD), *Mio.*, USA (Va.); 3a,b, RV ext., LV int., $\times 5$ (Glenn, 1904).
- Platyodon** CONRAD, 1837 [*Mya* (*P.*) *cancellata*; M] [= *Cryptodonta* CARPENTER, 1864 (obj.) (in synonymy)]. Resembling *Mya* but with chondrophore smaller and with concentric sculpture formed by raised growth striae; pallial sinus large and deep. *Mio.-Rec.*, Afr.-N.Am.-S.Am.
- P. (Platyodon)**. Shell 30 mm. or more in length. *Mio.-Rec.*, W.Afr.-W.N.Am.—FIG. E153,6. *P. (*P.*) *cancellatus* (CONRAD), *Rec.*, USA (Calif.); 6a,b, LV int. (oblique), ext., $\times 0.5$ (Grant & Gale).
- P. (Austroplatyodon)** OLSSON, 1961 [*P.* (*A.*) *australis*; OD]. Shell less than 6 mm. in length. *Rec.*, W.S.Am.—FIG. E153,7. *P. (*A.*) *australis*, Ecuador; LV int., $\times 5$ (688).
- Sphenia** TURTON, 1822 [*S. binghami*; SD GRAY, 1847] [= *Sphaena*, *Sphaenia*, spelling errors; *Tyleria* H. ADAMS & A. ADAMS, 1854 (type, *T. fragilis*; M)]. Small to minute, shell quadrate but apt to be distorted by nesting habit; pallial sinus small to large. *Paleoc.-Rec.*, Eu.-N.Am.-E.Asia.—FIG. E153,1. *S. *binghami*, *Rec.*; Eng.; 1a, LV and RV hinge; $\times 1$ (Woodward); 1b,c, RV and LV hinges, $\times 4$, $\times 6$ (Lamy).
- Tugonia** RÉCLUZ, 1846 [*Mya anatina* GMELIN, 1791 (*ex* "Le Tugon" ADANSON, 1757); SD GRAY, 1847] [= *Tagonia*, spelling error]. Globose, gaping posteriorly; posterior area set off by constriction; chondrophores subequal; pallial sinus shallow. *Eoc.-Rec.*, Eu.-N.Am.-IndoPac.-Asia-Australia.
- T. (Tugonia)**. Sculpture reticulate; shell of moderate size. *Oligo.-Rec.*, Eu.-IndoPac.—FIG. E153,8. *T. (*T.*) *anatina* (GMELIN), *Rec.*, W. Afr.; 8a-c, LV int., ext., RV hinge, $\times 1$ (H. Adams & A. Adams).
- T. (Antiguamya)** EFFINGER, 1938 [*Saxicava arnoldi* DICKERSON, 1917; OD]. Umbonal grooves marked, posterior area sharply set off, larger than in *T. (Tugonia)*. *Eoc.-L.Oligo.*, W.N.Am.—FIG. E153,5. *T. (*A.*) *arnoldi* (DICKERSON), *Oligo.*, USA (Ore.); LV int., $\times 1.5$ (Effinger).
- T. (Distugonia)** IREDALE, 1936 [*D. inopinata*; OD]. Sculpture weak, of concentric growth lines; posterior area small, attenuated. *Rec.*, Australia-E.Asia.
- T. (Tugonella)** JOUSSEAUME, 1891 [*Tugonia divaricata* REEVE, 1863; M]. Inflated, small, sculpture faint; chondrophore in RV smaller than in *T. (Tugonia)*; pallial sinus small. *Rec.*, Indo-Pac.
- T. (Tugoniopsis)** DALL, 1898 [*T. (T.) compacta*; OD]. Small to minute, short, chondrophore in front of dentiform projection; posterior adductor scar large and deep; pallial sinus moderate. *Mio.*, SE.N.Am.—FIG. E153,11. *T. (*T.*) *compacta*, USA (Fla.); LV int., $\times 3$ (Dall, 1898).

Family CORBULIDAE Lamarck, 1818

[*nom. correct.* BRODERIP, 1839 (*ex* Corbulidées LAMARCK, 1818, vernac.) [= *Corbularia* GRAY, 1823; *Corbulidae* FLEMING, 1828]]

Small to moderate-sized sturdy shells, normally with resilifer in one valve; valves inequilateral, LV tending to be smaller than RV; pallial sinus small to wanting. *U.Jur.-Rec.*

Subfamily CORBULINAE Gray, 1823

[*nom. transl.* STOLICZKA, 1870 (*ex* family *Corbularia* GRAY, 1823)]

LV only slightly smaller than RV, mostly with projecting chondrophore; posterior end somewhat rostrate. *U.Jur.-Rec.*

Corbula BRUGUIÈRE, 1797 [*C. sulcata* LAMARCK, 1801; SD SCHMIDT, 1818] [= *Aloidis* MEGERLE VON MÜHLFELD, 1811 (obj.)]. Moderately inflated, smooth to concentrically ribbed. *Cret.-Rec.*, Afr.-Eu.-Asia-IndoPac.-C.Am.-N.Am.-S.Am.

C. (Corbula). Solid, strongly ribbed, rostrate, LV without projecting chondrophore but with posterior cardinal tooth and ligamental pit, RV with posterior lateral. *Rec.*, W.Afr.—FIG. E154,5. *C. (*C.*) *sulcata* LAMARCK, Senegal; 5a, RV ext., $\times 1$; 5b, LV ext., $\times 1$; 5c,d, RV int., LV int., $\times 1$ (944).

C. (Anisocorbula) IREDALE, 1930 [*Corbula macgillivrayi* SMITH, 1885; OD]. Elongate-quadrate, with sharp umbonal keel, ventral margin sinuous; pallial sinus obsolete. *Rec.*, IndoPac.—FIG. E154,8. *C. (*A.*) *macgillivrayi* SMITH, New Guinea; 8a,b, RV int., ext., $\times 1$ (944).

C. (Bicorbula) FISCHER, 1887 [*C. gallica* LAMARCK, 1801; M]. Large, inequivalved, keel obsolescent; sculpture weak; pallial sinus broad and shallow. *Paleoc.-Rec.*, Eu.-Afr.-Asia.—FIG. E154,10. *C. (*B.*) *gallica* LAMARCK, *Eoc.*, France; 10a-e, RV ext., int., dorsal, LV ext., int., $\times 1$ (944).

C. (Bothrocorbula) GABB, 1873 [*C. viminea* GUPPY, 1866; M]. Moderately large, thick, sculpture of concentric waves and sparse radial threads;

ligamental pit broad in RV, one small dentiform projection and deep lunular pit in LV. *Mio.*, E.C. Am.-W.Indies.—FIG. E154,7. **C. (B.) viminea* GUPPY, Jamaica; 7a-c, LV ext., int., RV int., X1.5 (944).

C. (*Caryocorbula*) GARDNER, 1926 [**Corbula alabamiensis* LEA, 1833; OD]. Subquadrate, both valves acutely keeled posteriorly, concentrically rugose; pallial sinus short. *Eoc.-Rec.*, N.Am.-S. Am.-E.Asia.—FIG. E154,2. **C. (C.) alabami-*




FIG. E154. Corbulidae (Corbulinae) (p. N692-N694).




FIG. E155. Corbulidae (Corbulinae) (p. N694).

ensis LEA, Eoc., USA (Ala.); 2a,b, LV dorsal, RV ext., $\times 1.5$ (944); 2c,d, RV int., LV int., $\times 1$ (944; Stenzel *et al.*, 1957).

C. (*Cuneocorbula*) COSSMANN, 1886 [**Corbula biangulata* DESHAYES, 1857; SD DALL, 1898]. Small, thin, birostrate, concentric sculpture weak, valves subequal; pallial sinus small. Eoc., Eu.-N.Am.—FIG. E154,3. **C. (C.) biangulata* DESHAYES, France; 3a,b, RV ext., int., $\times 2.5$ (944).

C. (*Cuspicorbula*) OLSSON, 1928 [**Corbula busera*; OD]. Solid, inflated, coarsely sculptured, posterior end contracted and set off by deep sinus. Eoc., S.Am.—FIG. E154,6. **C. (C.) busera* OLSSON, Peru; RV ext., $\times 1.5$ (944).

?C. (*Eoursivivas*) OTA, 1964 [**C. matsumotoi* HABE, 1960; OD]. Hinge lacking dorsolateral

ridge on chondrophore. Cret. (*L. Neocom.*), Japan.—FIG. E155,3. **C. (E.) matsumotoi* HABE, Kumamoto Pref.; 3a-d, LV ext., RV int., LV int., hinge, $\times 1.5$ (Ota, 1964).

C. (*Flexicorbula*) CHAVAN, 1947 [**Varicorbula (F.) vokesi*; OD]. Resembling C. (*Varicorbula*) but beaks prosogyrate. Cret. (*Campan.*), W. Asia.—FIG. E154,1. **C. (F.) vokesi* (CHAVAN), Palestine; 1a-d, RV int., LV int., both valves (LV view), RV ext., $\times 1.5$ (105).

C. (*Hexacorbula*) OLSSON, 1932 [**C. hexacyma* BROWN & PILSBRY, 1913; OD]. Resembling C. (*Bothrocorbula*) but lacking lunular pit. Mio.-Rec., E.C.Am.-W.C.Am.—FIG. E154,4. **C. (H.) hexacyma* BROWN & PILSBRY, Mio., E. Panama; LV ext., $\times 3$ (944).

C. (*Juliacorbula*) OLSSON & HARBISON, 1953 [**C. cubaniana* D'ORBIGNY, 1853 (= *C. knoxiana* C. B. ADAMS, 1852); OD]. Valves solid, similar in size and shape, rectangular, rostrum bordered by keel. Mio.-Rec., E. C. Am.-W. C. Am.-S. Am.—FIG. E154,9. **C. (J.) knoxiana* ADAMS, Rec., W. Indies; 9a-c, LV ext., post., ant., $\times 2.5$ (Dall, 1889).

?C. (*Nipponicorbula*) OTA, 1964 [**N. mifunensis*; OD]. Sculpture discrepant, cancellate in RV, concentric in LV. U.Cret., Japan.—FIG. E155,1. **C. (N.) mifunensis* (OTA), Kumamoto Pref.; 1a-d, RV ext., LV ext., RV int., LV int., $\times 2.5$ (Ota, 1964).

C. (*Notocorbula*) IREDALE, 1930 [**N. vicaria*; OD]. Large, chondrophore bipartite; adductor scars large. Rec., Pac.—FIG. E156,1. **C. (N.) vicaria* (IREDALE), Australia; 1a-c, RV ext., LV ext., RV int., $\times 1$ (944).

?C. (*Pulsidis*) OTA, 1964 [**P. nagatoensis*; OD]. Resembling *Caryocorbula* but chondrophore without median ridge. L.Cret., Japan.—FIG. E155, 2. **C. (P.) nagatoensis* (OTA), Yamaguchi Pref.; 2a-d, RV ext., LV ext., RV int., LV int., $\times 2.5$ (Ota, 1964).

C. (*Serracorbula*) OLSSON, 1961 [**S. tumaca*; OD]. Solid, heavy, subequivalve, convex, ventral margin evenly serrate; adductor scars large. Rec., W.C.Am.-S.Am.—FIG. E156,4. **C. (S.) tumaca* (OLSSON), W. Colombia; LV int., $\times 2.2$ (688).

C. (*Solidicorbula*) HABE, 1949 [**C. erythrodon* LAMARCK, 1818; OD]. Heavy, convex, solid; LV with knob at posterior end; chondrophore in LV and resilifer in RV divided by ridge; muscle scars deep, pallial sinus wanting. Rec., W. Pac.—FIG. E157,1. **C. (S.) erythrodon* LAMARCK, Japan; 1a-c, LV ext., RV int., LV int., $\times 1$ (Habe, 1949).

C. (*Tenuicorbula*) OLSSON, 1932 [**C. tenuis* SOWERBY, 1833; OD]. Thin, subequilateral, posterior end set off by keel; sculpture strong, of concentric threads. Mio.-Rec., E.C.Am.-W.C.Am.-W.S.Am.—FIG. E156,3. **C. (T.) tenuis* SOWERBY, Rec., W. Colom.; 3a,b, LV ext., RV dorsal, $\times 1$ (688).

C. (Varicorbula) GRANT & GALE, 1931 [**Tellina gibba* OLIVI, 1792; OD] [= *Corbula* AUCTT. (non BRUGUIÈRE)]. Small, sculpture discrepant, RV with concentric ribbing, LV with faint radials and concentric growth striae; pallial sinus small

or wanting. *L.Eoc.-Rec.*, E.N.Am.-Eu.-E.Pac.-W. Pac.—FIG. E156,8. **C. (V.) gibba* (OLIVI), *Rec., Eng.*; 8a-d, RV ext., int., LV int., both valves (LV view), $\times 2$ (944).

C. (Vokesula) STENZEL & TWINING in STENZEL,




FIG. E156. Corbulidae (Corbulinae) (p. N694-N696).

KRAUSE & TWINING, 1957 [**C. aldrichi* MEYER, 1885, var. *smithvillensis* HARRIS, 1895; OD]. Inflated, valves discrepant, RV larger, concentrically ribbed, LV nearly smooth; pallial sinus small, nearly vertical. *L.Eoc.-L.Oligo.*, E.N.Am.—FIG. E156,7. **C. (V.) smithvillensis* HARRIS, M.Eoc., USA(Tex.); 7a-c, both valves (LV view), LV int., RV ext., $\times 2.5$ (Stenzel & Twining, 1957).

Anapteris VAN WINKLE [PALMER], 1919 [**A. regalis*; M]. LV with winglike flare anteriorly; both valves with posterior area set off by keel. *Eoc.*, E.N.Am.—FIG. E156,9. **A. regalis*, USA (Va.); 9a,b, LV ext., LV int., $\times 3$; 9c, RV ext., $\times 2.8$ (944).

Corbulomima VOKES, 1945 [**C. nuciformis*; OD]. Without special structures in either valve for reception of resilifer, otherwise resembling *C. (Corbula)*. *U.Jur.-L.Cret.(Apt.)*, Eu.-W.Asia.—FIG. E156,2. **C. nuciformis*, Lebanon; 2a-c, LV int., RV int., both valves (LV view), $\times 2$ (944).

Panamicorbula PILSBRY, 1932 [**Potamomya inflata* C. B. ADAMS, 1852; OD]. Thin, inflated, not rostrate posteriorly; hinge with toothlike lateral buttresses to dorsal grooves of RV. *Rec.*, C.Am.—FIG. E156,6. **P. inflata* (C. B. ADAMS), W.Panama; 6a, ext. RV, $\times 1$; 6b,c, LV and RV hinges, $\times 2$ (688).

Physoida PALLARY, 1900 [**Corbula physoides* DESHAYES, 1845-48; OD]. Small, smooth, fragile, with low umbonal ridge. *Rec.*, N.Afr.—FIG. E156,11. **P. physoides* (DESHAYES), Algeria; 11a-c, RV ext., LV hinge, RV int., $\times 3$ (944).

Potamocorbula HABE, 1955 [**Corbula amurensis* SCHRENCK, 1867; OD]. Hinge of RV with strongly projecting tooth in front of deep pit; LV with longitudinal keel on chondrophore; periostracum present, thick. *Rec.*, NE.Asia.—FIG. E156,5. **P. amurensis* (SCHRENCK), Korea; 5a,b, RV ext., LV int., $\times 1$ (Habe, 1955).

Ursirivus VOKES, 1945 [*nom. subst. pro Anisorhynchus* CONRAD in MEEK & HAYDEN, 1871 (*non* SCHOENHERR, 1842)] [**Corbula pyriformis* MEEK, 1871; OD]. Large, pyriform, with deep lunule and small but definite pallial sinus. *U.Cret.*, N.Am.—FIG. E156,10. **U. pyriformis* (MEEK), USA (Wyo.); 10a-c, RV ext., int., LV int., $\times 1$ (944).

Subfamily CAESTOCORBULINAE Vokes, 1945

Valves discrepant in shape, LV more equilateral and less rostrate than RV; accessory siphonal plate posterior to LV fitting into rostrum of RV; projecting chondrophore of hinge of LV. *Cret.-Eoc.*

Caestocorbula VINCENT, 1910 [**Corbula henckeliusiana* NYST, 1836 (as *C. henckeliusi*, spelling error); OD] [= *Ficusocorbula* KOROBKOV, 1954 (type, *Corbula ficus* SOLANDER, 1766); OD]. *L.Cret.-Eoc.*, Eu.-N.Am.-Asia-Afr.




FIG. E157. Corbulidae (Corbulinae) (p. N694, N696).

C. (Caestocorbula). Siphonal plate obliquely trapezoidal, with distinct median ridge; pallial sinus well developed, broad and rounded. *L.Cret.(Apt.)-Eoc.*, E.N.Am.-Eu.-Asia.—FIG. E158,6. **C. (C.) henckeliusiana* (NYST), Eoc., Belg.; both valves (LV view), $\times 1$ (Vincent, 1910; 944).

C. (Parmicorbula) VOKES, 1944 [**Corbula neaeroides* BLANCKENHORN, 1890; OD]. Siphonal plate small, rectangular, with faint median groove; otherwise resembling *C. (Caestocorbula)*. *U.Cret.-Eoc.*, Eu.-Asia-E.N.Am.-Afr.—FIG. E158,3. **C. (P.) neaeroides* (BLANCKENHORN), U.Cret., Lebanon; 3a-d, RV int., ext., both valves (LV views), $\times 1.5$ (944).

Subfamily CORBULAMELLINAE Vokes, 1945

Small, valves subtrigonal, lacking both posterior rostrum and umbonal ridge. Posterior adductor on raised, spoon-shaped lamella. *Cret.*

Corbulamella MEEK & HAYDEN, 1857 [**Corbula? gregaria*; OD]. Umbones inflated, slightly prosogyrate; sculpture of fine concentric lines; pallial sinus shallow. *Cret.*, N.Am.—FIG. E158,1. **C. gregaria* (MEEK & HAYDEN), U.Cret., USA (S. Dak.); 1a-d, RV int., ext., both valves (LV view), ant. view, $\times 5$ (944).

Subfamily LENTIDIINAE Vokes, 1945

Small, essentially tellinid in external shape, RV lacking hinge plate; cardinal tooth on subumbonal thickening that projects from interior of valve. *Paleoc.-Rec.*

Lentidium CRISTOFORI & JAN, 1832 [genus without named species] [**L. maculatum* (= **Tellina mediterranea* COSTA, 1829); SM] [= *Corbulomima* NYST, 1844 (type, *Corbula complanata* SOWERBY, 1822; SD HERRMANNSEN, 1847)]. Shell thin, chondrophore projecting; muscle scars small and impressed, pallial sinus broad and shallow. *Paleoc.-Rec.*, Eu.-Asia-Atl.-Medit.

L. (Lentidium). Beaks central or posterior to midline. *Paleoc.-Rec.*, Eu.-E.Atl.-Medit.—FIG. E158,




FIG. E158. Corbulidae (Caestocorbulinae) (3,6), (Corbulamellinae) (1), (Lentidiinae) (4-5), Pachydoniinae (2,7), Subfamily Uncertain (8) (p. N696-N698).

5. **L. (L.) mediterraneum* (COSTA), Rec., Italy; 5a,b, RV ext., int., $\times 4$ (944).

L. (Janschinella) MERKLIN, 1961 [**L. (J.) garetzkii*; OD]. Anterior end short, beaks well forward of mid-line. *M.Oligo-L.Mio.*, C.Asia-W.Eu. —FIG. E158,4. **L. (J.) garetzkii*, Oligo., USSR; 4a-c, RV ext., int., LV hinge, $\times 2$ (Merklin).

Subfamily PACHYDONTINAE Vokes, 1945

Valves tending to be twisted and distorted, ligamental area attaching resilium to lateral rather than dorsal face of resilifer. *Oligo.-Plio.*

Pachydon GABB, 1868 [**P. obliqua*; SD VOKES, 1944] [= *Anisothyris* CONRAD, 1871 (obj.); *Pachyodon*, nom. van., MEEK, 1878 (non VON MEYER, 1838)]. Sculpture of weak concentric ridges; outline cordate, umbonal ridge narrow; pallial sinus broad, shallow. *Plio.*, S.Am. —FIG. E158,7. **P. obliquus*, Peru, 7a-e, RV int., LV int., RV ext., both valves ant., LV view, $\times 2$ (944).

Tiza DE GREGORIO, 1890 [**Corbula? amara* (= **C. aliformis* CONRAD, 1885); M]. More quadrate than *Pachydon*, somewhat donaciform; pallial line angular at junction with posterior adductor scar. *Oligo.*, E.N.Am. —FIG. E158,2. **T. aliformis* (CONRAD), USA (Miss.); 2a-d, RV ext., int., RV ext., LV int., $\times 1$ (944).

Subfamily UNCERTAIN

?*Coquandia* SEGUENZA, 1882 [**C. italica*; SD KEEN,




FIG. E159. Erodonidae (2); Pleurodesmatidae (4), Raetomyidae (5-6); Spheniopsidae (1,3) (p. N698-N699).

herein]. Relatively large, hinge with a *Mya*-like chondrophore, known only from internal molds; pallial line with a small sinus; shell not gaping. *U.Cret.(Cenoman.)*, Eu.—FIG. E158,8. **C. italica*, Italy; 8a,b, RV int. mold, both valves int. molds dorsal, $\times 0.7$ (Seguenza, 1882).

Harlea, Raleta, Tomala GRAY, 1842 (*nomina nuda*). Described as Corbulidae but without citation of species names. Unrecognizable from descriptions. *Semicorbula* COSSMANN & PEYROT, 1909 [**S. nadali*; OD]. *Oligo.(Aquitain.)*, Eu. Based on single RV, indeterminate from figure.

Family ERODONIDAE Winckworth, 1932

Resembling Corbulidae but with chondrophore of LV broad, projecting, as in *Mya*. *U.Eoc.-Rec.*

Erodonia BOSCH (*ex* DAUDIN, MS), 1801 [**E. macroides*; SD DALL, 1898] [= *Potamomya* J. SOWERBY, 1835 (type, *Mya plana* SOWERBY, 1814; SD KEEN, herein); *Azara* D'ORBIGNY, 1842 (obj.); *Phaenomya* WEAVER & PALMER, 1922 (type, *P. vaderensis*; OD)]. Ovate-trapezoidal in outline; umbo of LV higher than RV; inequivalve, RV larger; hinge of RV with 2 thin cardinals bordering triangular resilial pit; pallial sinus small. [Habitat mainly brackish-water.] *U.Eoc.-Rec.*, N.Am.-C. Am.-N.Eu.-SE.S.Am.—FIG. E159,2. **E. macroides*, Rec., Arg.; 2a,b, LV int., RV int., $\times 0.6$ (Woodward).

Family PLEURODESMATIDAE Cossmann & Peyrot, 1909

Small, porcelaneous, quadrate to trapezoidal, mostly smooth or with laminar growth striae; hinge with one cardinal tooth in either valve and long excavated cavity on posterior dorsal hinge margin for resilium; pallial line entire. *U.Oligo.-Mio.*

Pleurodesma HÖRNES, 1859 [**P. mayeri*; M]. Beaks prosogyrate; resilifer bordered by low lamina below; muscle scars and pallial line inconspicuous. *U.Oligo.(Aquitain.)-Mio.(Burdigal.)*, Eu.—FIG. E159,4. **P. mayeri*, Mio., Austria; 4a-c, LV int., RV int., LV ext., $\times 1$; 4d, RV hinge, $\times 5$ (after Hörnes, 1859).

Family RAETOMYIDAE Newton, 1919

[= *Amotapidae* OLSSON, 1928]

Ovate, thin, sculptured with undulating concentric ribs, posteriorly compressed and gaping; hinge with large projecting chondrophore in LV, smaller sunken chondrophore in RV entering umbonal cavity vertically. Resembling mactrid group *Raeta* exteriorly but with myacean hinge. *Eoc.*

Raetomya NEWTON, 1919 [**Lovellia schweinfurthi* MAYER-EYMAR, 1887; OD]. Ovate-quadrate, chondrophore in LV 3-lobed. *Eoc.*, NE.Afr.-W.C.Afr. —FIG. E159,5. **R. schweinfurthi* (MAYER-EYMAR), U.Eoc., Nigeria; 5a, RV ext., $\times 0.5$; 5b,c, LV hinge, $\times 1$; 5d, RV hinge, $\times 1$ (Newton, 1919).

Amotapus OLSSON, 1928 [**Perna arbolensis* WOODS, 1922; OD]. Elongate-ovate, posterior end short; chondrophore in LV rounded, not conspicuously lobed. *Eoc.*, W.S.Am. —FIG. E159,6. **A. arbolensis* (WOODS), M.Eoc., Peru; 6a,b, LV hinge, RV ext., 0.5 (6a, Stanford specimen; 6b, Olsson).

Family SPHENIOPSISIDAE Gardner, 1928

Small, ovate-triangular, RV with 2 hinge teeth, LV edentulous, with resilial socket; pallial sinus rounded. *M.Eoc.-Rec.*

Spheniopsis SANDBERGER, 1863 [**Corbula scalaris* BRAUN in WALCHNER, 1851; OD]. Trigonal, posteriorly rostrate; smooth to concentrically undulate; anterior hinge tooth in RV triangular, posterior lamellar. *M.Eoc.-Mio.*, Eu.-E.N.Am. —FIG. E159, 1. **S. scalaris* (BRAUN), Oligo., Aus.; 1a-c, LV int., ext., RV int., $\times 5$ (Sandberger).

Grippina DALL, 1912 [**G. californica*; OD]. Ovate-trigonal, finely concentrically sculptured, with radial ridges bounding lunule and escutcheon. *Rec.*, W.N.Am.-W.C.Am. —FIG. E159,3. **G. californica*, USA (Calif.); 3a-c, RV ext., int., LV int., $\times 8$ (Oldroyd, 1925, and specimen, Stanford Univ.).

Superfamily GASTROCHAENACEA Gray, 1840

[*nom. trans.* THIELE, 1934 (*ex Gastrochaenidae* GRAY, 1840)]
[Materials for this superfamily=prepared by MYRA KEEN]

Burrowing shells, lying free within linear cavity; hinge edentate, valves broadly gaping, porcelaneous (517). *U.Jur.-Rec.*

Family GASTROCHAENIDAE Gray, 1840

[*nom. correct.* ADAMS & ADAMS, 1856 (*pro Gastrochaenadae* GRAY, 1840)]

Characters of superfamily. *U.Jur.-Rec.*

Gastrochaena SPENGLER, 1783 [**G. cuneiformis*; SD CHILDREN, 1822] [= *Chaena* PHILIPSSON, 1788 (*ex* RETZIUS, MS) (*nom. van.*); *Rocellaria* DE BLAINVILLE, 1829 (type, *Gastrochaena modiolina* LAMARCK, 1818, = *Mya dubia* PENNANT, 1777, M); *Gastrochaenolites* LEYMERIE, 1842 (*nom. van.*); *Gastrochaenoecium*, *Gastrochaenites* BRONN, 1848 (*nom. van.*); *Cucurbitula* GOULD, 1861 (type, *Fistulana lagenula* LAMARCK, 1801, = *Gastrochaena*




FIG. E160. Gastrochaenidae (p. N699-N700).

cymbium SPENGLER, 1783; M); *Dufoichaena* LAMY, 1925 (*ex* JOUSSEAUME, MS) (type, *Gastrochaena dentifera* DUFO, 1840; M)]. Cavity flask-shaped, in rock or shell; shell triangular to quadrate in outline. ?*U.Jur.*, *U.Cret.-Rec.*, N.Am.-W. Indies-Eu.-E.Indies.

G. (*Gastrochaena*). Shell smooth or nearly so, oval, tube short, formed of successive increments. ?*U.Jur.*, *U.Cret.-Rec.*, N.Am.-Eu.-circumtrop. —FIG. E160,5. **G.* (*G.*) *cuneiformis* SPENGLER, *Rec.*, IndoPac.; 5a,b, LV ext., ventral, $\times 1$ (Reeve).

G. (*Spengleria*) TRYON, 1862 [**G. mytiloides* LAMARCK, 1818; SD STOLICZKA, 1871]. Valves divided by oblique furrow, posterior area with vertical striae. *Rec.*, E.Indies-W.Indies. —FIG. E160,4. **G.* (*S.*) *mytiloides* LAMARCK, E.Indies; 4a,b, LV ext., ventral, $\times 1$ (Reeve).

Eufistulana EAMES, 1951 [**Gastrochaena mumia* SPENGLER, 1783; OD] [= *Fistulana* BRUGUIÈRE,

1789 (*non* MÜLLER, 1776); SD FLEMING, 1818]. Elongate valves enclosed in adventitious tubular structure; widely gaping. *Eoc.-Rec.*, Eu.-N.Afr.-IndoPac.—FIG. E160,1. **E. mumia* (SPENGLER), *Rec.*, E.Indies; both valves, LV view, $\times 0.5$ (Fischer).

Gastrochaenopsis CHAVAN, 1952 [**Gastrochaena uncostata* DESLONGCHAMPS, 1838; OD]. Posterior cardinal margin patulous, accentuated by 2 dorsal carinae. *U.Jur.*, Eu.—FIG. E160,2. **G. uncostata* (DESLONGCHAMPS), France; RV ext., $\times 1$ (Chavan).

Kummelia STEPHENSON, 1937 [**Gastrochaena americana* GABB, 1860; OD]. Shell resembling *Gastrochaena* but ventral margin broadly rounded, not truncate; tube cylindrical, not tapering, annular, rough, somewhat sinuous. *Eoc.*, E.N.Am.—FIG. E160,3. **K. americana* (GABB), USA (N.J.); 3a,b, tube, RV ext., $\times 0.5$ (Stephenson).

Superfamily HIATELLACEA Gray, 1824

[*nom. transl.* NEWELL, 1965 (*ex* Hiatellidae GRAY, 1824) [Materials for this superfamily prepared by MYRA KEEN except as otherwise recorded]]

Quadrangle to trapezoidal, valves slightly to widely gaping; habit nestling or burrowing; hinge with one or two weak teeth, ligament on nymph; pallial sinus mostly well developed (516). *Perm.-Rec.*

Family HIATELLIDAE Gray, 1824

[*nom. correct.* WINGKORTH, 1932 (*pro* Hyatellidae GRAY, 1824)] [=Saxicavidae SWAINSON, 1835; Glycimeridae DESHAYES, 1839 (*nom. correct.* PETIT DE LA SAUSSAYE, 1869, *pro* Glycimeridés)]

Characters of superfamily. *Perm.-Rec.*

Hiatella BOSCH (*ex* DAUDIN, MS), 1801 [**H. monoperta* (=Mya *arctica* LINNÉ, 1767); SD WINCKWORTH, 1932] [=Saxicava FLEURIAU DE BELLEVUE, 1802 (*obj.*), *Laxicava* CONRAD, 1855 (*nom. null.*); *Haicana* SACCO, 1901 (*nom. null.*); *Clotho* FAUJAS DE SAINT FOND, 1808 (*nom. dub.*); *Didonta* SCHUMACHER, 1817 (*obj.*) (type, *Solen minutus* LINNÉ, 1767, =Mya *arctica*; M); *Byssonina* DE BLAINVILLE, 1817 (*ex* CUVIER, *vernac.*) (type, *Mytilus pholadis* LINNÉ, 1771), *Bissomya*, *Bissomyia*, *Bys-somia*, *Byssomya*, spelling errors; *Biapholus* LAMARCK (*ex* LEACH, MS), 1818 (in synonymy of *Mytilus rugosus* LINNÉ, 1767); *Biapholus*, *Biopholeus* LEACH, 1847 (*nom. null.*); *Pholeobia* LEACH, 1819 (type, *Mytilus rugosus* LINNÉ, 1767; M); *Agina* TURTON, 1822 (*obj.*; M); *Rhomboides* DE BLAINVILLE, 1824 (*non* GOLDFUSS, 1820); *Coromya* BROWN, 1844 (*obj.*, in synonymy); *Spongyophylla* BRUSINA (*ex* NARDO, MS), 1866 (*obj.*; M). Small, trapezoidal, with thin perio-

stracum; hinge teeth best developed in juvenile stages. *U.Jur.-Rec.*, Asia-Eu.-N.Am.

H. (Hiatella). Outline rendered irregular by nestling habit, in general with posterior end wider than anterior, gape small. *Oligo.-Rec.*, E.Asia-N.Am.-Eu.—FIG. E161,1. **H. (H.) arctica* (LINNÉ), *Rec.*, France, 1a-d, LV ext., int., RV int., ext., $\times 1$ (89a).

H. (Pseudosaxicava) CHAVAN, 1952 [**P. bernardi*; OD]. Hinge teeth weak, muscle scars elliptical. *U.Jur.*, Eu.—FIG. E161,2. **H. (P.) bernardi* (CHAVAN), France; 2a-c, RV ext., int., LV int., $\times 2$ (Chavan).

Capistrocardia TATE, 1887 [**C. fragilis*; OD]. Rounded quadrangle to elliptical, with internal radial ridge; not gaping; hinge with 2 teeth in LV, none in RV; pallial sinus small or wanting. *Tert.*, Australia.—FIG. E161,3. **C. fragilis*; 3a,b, LV ext., int., $\times 2$ (Tate).

Cyrtodaria REUSS, 1801 (*ex* DAUDIN, *vernac.*) [**Mya siliqua* SPENGLER, 1793; SD VOKES & COX, 1961] [=Cyrtodera DE BLAINVILLE, 1825 (*nom. null.*); *Glycimeris* LAMARCK, 1801 (*non* LAMARCK, 1799)]. Elongate, with thick, dark periostracum; hinge with 2 teeth, nearly central; pallial sinus small, irregular. *Paleoc.-Rec.*, Eu.-Arctic.—FIG. E161,10. **C. siliqua* (SPENGLER), *Rec.*, N.Atl.; 10a,b, RV int. with LV hinge, RV ext., $\times 1$ (H. Adams & A. Adams).

Panopea MENARD, 1807 (ICZN, *pend.*) [**P. faujasi* (=Mya *glycimeris* BORN, 1778); SD FLEMING, 1818] [=Glycimeris LAMARCK, 1799 (*non* DA COSTA, 1778); *Panope* MENARD, 1807 (*nom. null.*); *Panodea* OKEN, 1817 (*nom. null.*); *Panopaea* LAMARCK, 1818 (*nom. null.*); *Myopsis* AGASSIZ, 1840 (type, *Mya mandibula* J. SOWERBY, 1813; SD COX, 1964); *Heteromya* MAYER, 1884 (type, *P. (H.) lessepsi*; M)]. Medium-sized to large, elongate, gaping, beaks subcentral; ligamental nymph large, high; pallial sinus wide. ?*Trias.*, *L.Cret.-Rec.*, Eu.-N.Am.-S.Pac.

P. (Panopea). Hinge with 1 small cardinal tooth in either valve. ?*Trias.*, *L.Cret.-Rec.*, Eu.-S.Pac.-N.Am.—FIG. E161,7. **P. (P.) glycimeris* (BORN), *Rec.*, *Medit.*; 7a,b, LV ext., RV hinge, $\times 0.5$ (124b).

P. (Degrangia) COSSMANN & PEYROT, 1909 [**Panopaea fischerina* MAYER, 1861; OD]. Resembling *P. (Panopea)* in outline and shape of ligamental nymph and pallial sinus but hinge with 2 strong cardinal teeth in either valve. ?*Eoc.*, *Mio.*, Eu.—FIG. E161,8. **P. (D.) fischerina* (MAYER), *Mio.*, France; 8a,b, LV and RV hinges, $\times 3$; 8c, RV ext., LV int., $\times 1$ (Mayer; Cossmann & Peyrot).

P. (Panomya) GRAY, 1857 [**Mya norvegica* SPENGLER, 1793 (not preoccupied by *M. norvegica* GMELIN, 1791); M] [=Chaenopaea MAYER, 1885 (*obj.*; OD)]. Smaller than *P.*

(*Panopea*), with median depressed area bordered by broadly rounded ridges; pallial line interrupted, forming series of oval scars. ?*L.Cret.* (*Neocom.*), *Tert.-Rec.*, Eu.-N.Am.-Arctic.—FIG. E161,9. **P. (P.) norvegica* (SPENGLER), *Rec.*, N.Atl.; 9a,b, LV int. with RV hinge, LV ext., ×1 (H. Adams & A. Adams).

Roxoa MENDES, 1952, p. 103 [**Anoplophora intrigans* MENDES, 1944; OD]. Elongate, subbelliptical, equivalve, compressed, with or without weak posterior umbonal fold; beaks prosogyre, small, situated well forward; lunule and escutcheon lacking; surface smooth with well-defined lateral sulcus and ventral sinus; edentulous; pallial line with deep sinus. [In view of CHAVAN, this genus is allied to *Cowperesia* MENDES and at least doubtfully assignable to the Tellinidae.]. *Perm.*(*Corumbatae* F.), S.Am.(S.Brazil).—FIG. E161,5. **R. intrigans* (MENDES); 5a,b, RV ext., RV int. mold, ×1 (MENDES, 1952), [NEWELL]

Saxicavella FISCHER, 1878 [**Mytilus plicatus* GMELIN" of MONTAGU, 1808 (*non* GMELIN, 1791)




FIG. E161. Hiattellidae (p. N700-N702).

(=**S. jeffreysi* WINCKWORTH, 1930); M] [= *Arcinella* PHILIPPI, 1844 (*non* SCHUMACHER, 1817); *Arcinellina*, spelling error]. Thin, small, inequilateral, gaping, subtrapezoidal; hinge edentulous or with small conical cardinal in RV and socket in LV; ligament short, external; pallial line slightly sinuous, pallial sinus large but not deep. *U.Oligo.-Rec.*, Eu.-W.N.Am.—FIG. E161, A. **S. jeffreysi* WINCKWORTH, *Rec.*, N.Atl.; 4a-c, LV int., RV int., LV ext., $\times 3$ (Jeffreys, 1865).

Suborder PHOLADINA

H. Adams & A. Adams, 1858

[*nom. transl. et correct.* NEWELL, 1965 (*ex order* Pholadacea H. ADAMS & A. ADAMS, 1858)] [= *Adesmacea* AUCT.] [Diagnosis by N. D. NEWELL]

Characters of superfamily Pholadacea.
? *Carb., Jur.-Rec.*

Superfamily PHOLADACEA

Lamarck, 1809

[*nom. transl. et correct.* VOKES, 1967, p. 326 (*ex* Pholadaires LAMARCK, 1809)] [= family Pholadaria RAFINESQUE, 1815; *Adesmacea* DE BLAINVILLE, 1825, p. 577; *Cladopoda* GRAY, 1847, p. 187] [Materials for this superfamily prepared by RUTH D. TURNER, Museum of Comparative Zoology, Harvard University]

Bivalves characterized by closed mantle and truncate, more or less circular foot developed as suction disc. Shells inequilateral, dorsal margin reflected anterior to beak, forming attachment area for anterior adductor muscle which works counter to posterior adductor when animal is boring. Hinge teeth lacking, but small chondrophore and internal ligament usually present. Ventral adductor muscle or at least thickening of muscles of ventral margin of mantle located at union of anterior slope and disc, forming pivotal point opposite umbones on which valves rock when boring. Gills usually elongate, with two demibranchs except in the *Xylophaginae* and *Teredinidae*. Apophyses found in all Pholadacea except in some *Pholadidae* (*Jouannetiinae*, *Xylophaginae*). [The *Pholadidae* are characterized by the presence of one or more accessory plates and the *Teredinidae* by the presence of pallets. In the *Pholadidae* the intestine traverses the heart, whereas in all *Teredinidae*, except *Kuphinae*, it does not.]
? *Carb., Jur.-Rec.*

The Pholadacea is a highly specialized assemblage of bivalves adapted for boring into relatively hard substrata such as peat,

wood, nuts, woody plant stems, stiff clays, soft or friable rocks, corals, and shells. Though most are marine, some species occur in brackish and even fresh water.

Semidiagrammatic sketches of the anatomy of the *Pholadidae* and *Teredinidae* given in Figure E162 show the relationship of the two families and will be referred to when defining some genera. As a result of close approximation of the anterior and posterior adductor muscles in the *Teredinidae*, it can be seen that the visceral mass has been extended posteriorly in a long loop, with the mouth and anus remaining in normal position, that the ventricle is anterior to the auricles, and that the kidney is dorsal to the heart. Recent work by TURNER on anatomy of the *Teredinidae* has shown some striking differences within the group which have necessitated the redefining of some genera and the reassignment of several species. It has also resulted in firm establishment of the subfamily *Kuphinae*.

Among living forms, the *Teredinidae*, *Xylophaginae*, and the genus *Martesia* of the *Martesiinae* are found normally only in wood or other plant materials.

The classification of living species in this superfamily is not difficult if complete specimens are available, but when dealing with fossil forms it is often impossible to assign them to a subfamily. The reason for this is that the shells may be deformed, either as a result of the material in which they were boring or as a result of processes of fossilization. Many fossil species and genera are based only on internal casts; in others the accessory plates are lacking, the siphonoplax is broken off, and the apophyses are usually imbedded so as not to be seen at all. Using only the outer surface of the valve, one finds it impossible definitely to distinguish some young *Jouannetia* from young *Xylophaga* or even some *Teredinidae*. The variation in size and shape of individuals of any one species of pholad or teredinid is extremely great, depending upon conditions under which the animal was living and on its age. Therefore, it is usually necessary to have large series in order to understand a species adequately. Such series are seldom available in fossil forms, and it is indeed seldom that a growth series, together with

complete adults showing characters of both the inner and outer surface of the valves, is found. It is little wonder, then, that the assignment of fossil genera to the well-defined subfamilies based on living species has been done either provisionally or not at all.

In assigning fossil genera to subfamilies, the substrata in which they were found and the depth of the burrows are of prime importance. In the fossilized state, valves of the young of some species of *Martesia*, *Xylophaga*, and the Teredinidae—all wood-borers—could be indistinguishable, but a




FIG. E162. Morphological features of Pholadidae (1) illustrated by semidiagrammatic drawing of soft parts of *Zirfaea crispata* LINNÉ with mantle partially cut away, ca. $\times 2$, and of Teredinidae (2) by similar drawing of *Psiloteredo healdi* BARTSCH, ca. $\times 2.5$ (Turner, 1965).




FIG. E163. Side views of LV (1, exterior, 2, interior) of diagrammatic composite shell showing morphological features of Pholadidae (921a).

Martesia burrow is only slightly longer than the shell, whereas *Xylophaga* may make burrows five to ten times the length of the shells, and the burrows of *Teredinidae* are long and wormlike (see footnote, p. N721).

The presence of an umbonal-ventral sulcus on the outer surface of the valve or the corresponding ridge on the inner surface immediately eliminates all Pholadinae except *Zirfaea*, which is a transition genus between this subfamily and the Martesiinae.

The presence of a callum immediately places a genus in the Martesiinae or Jouannetiinae. This is not always easy to determine, for after the animal dies the valves remain in place, though the accessory plates may move or disintegrate and the

burrow is completely filled with silt. This filling protrudes beyond the valves and fills the anterior end of the burrow. When the entire mass is removed from the substratum, and particularly if it is an internal cast, the resulting specimen appears to have a callum. If a true callum is present, a central division should be evident, for the two halves of the callum do not fuse but are joined by a band of periostracum (in the case of the Martesiinae) or the line of overlap of the callum should be evident (e.g., *Jouannetia*). On the basis of such considerations the fossil genera have been placed in what seems to be the appropriate subfamily and the Recent genus to which they appear to be related is indicated.




FIG. E164. Morphological features of Teredinidae.—1, *Bankia*, entire animal, ca. $\times 2$.—2, Enlarged views of valves; 2a, outer view of right valve; 2b, inner view of right valve; 2c, anterior view of opposed valves showing the condyles on which the valves rock and the pedal gape.—3. Composite diagrammatic sketch of a *Bankia* pallet showing all characters (923; Clench & Turner, 1946).

In view of the large number of specialized terms used in describing the Pholadacea, a glossary is included, along with diagrammatic sketches indicating the specialized characteristics (Fig. E163, E164).

Special Morphological Terms Used For Pholadacea

apophyses. Calcareous, styloid projections, one in each valve, extending from beneath umbo and serving for attachment of pedal muscles.

callum. Closure of pedal gape in the Martesiinae and Jouannetiinae. It may be partial or complete, entirely calcareous or composed partly of periostracum. Generally halves of callum do not quite meet and are joined by a periostracal fold with only a minute pore remaining open.

cephalic hood. Thickened fleshy fold of mantle covering umbonal area.

chimney. Tube of agglutinated particles produced as result of boring activities. It fits over posterior end of shell and in some species extends anteriorly nearly to umbos. [Occurs in *Parapholas* and *Xylophaga*.]

chondrophore. Projection of hinge area for support of internal ligament. Chondrophore of RV a small swelling with central depression, that of LV a small shelflike projection.

hypoplax. Long, narrow, ventral plate covering space between valves and joined to ventral margin of valves by periostracal fold.

mesoplax. Transverse plate, usually wider than long, which straddles valves at the umbos and protects posterior portion of anterior adductor muscle. It usually originates ventral to the muscle, is calcareous, and may be in one or two pieces.

metaplax. Long, narrow plate covering gap between valves on dorsal margin behind mesoplax and joined to valves by periostracal fold.




FIG. E165. Accessory plates of genera in Pholadinae (921a).

pallets. Pair of paddle-shaped, calcareous and periostracal organs located at base of siphons in Terebinthidae and functioning to close entrance of burrow. They may be in one piece or composed of numerous elements.

protoplax. Simple, nearly flat, elongate dorsal plate which rests on top of anterior adductor muscle but does not enclose it. It may be calcareous or periostracal, in one piece, or divided longitudinally.

siphonal tube. Tube fused to siphonoplax and composed of secretions of mantle and agglutinated particles produced as result of boring. [Occurs in *Pholadidea*.]

siphonoplax. Accessory structure secreted by mantle and added to posterior end of valves in adult stage. It may be calcareous, periostracal, or combination of both, and may be variously formed.

Family PHOLADIDAE Lamarck, 1809

[*nom. correct.* VOKES, 1967, p. 326 (*pro* family *Pholadaires* LAMARCK, 1809, p. 319)] [= *Pholadaria* RAFINESQUE, 1815, p. 146; *Pholadariae* LATREILLE, 1825, p. 223; *Pholadae* FLEMING, 1828, p. 410, 456; *Pholadidae* GRAY, 1847, p. 20; *Pholadoidea* AGASSIZ, 1847, p. 286; *Pholidae* SWAINSON, 1835, p. 30; *Pholadae* MENKE, 1828, p. 73]

Shells elongate to globular, with slitlike to nearly circular pedal gape which may or

may not be closed by callum in adult; anterior portion of valves with imbricate or denticulate concentric ridges and commonly ribbed radially; umbonal-ventral sulcus may separate anterior slope from disc; posterior slope not generally demarked, but clearly defined in *Parapholas*; accessory plates present in all forms but varying in number. Dorsal and ventral condyles and internal umbonal-ventral rib (reflection of umbonal-ventral sulcus of outer surface of valve) present or absent; pallial sinus usually deep. Siphons variable, usually united, capable of considerable extension, smooth or papillose, and commonly enclosed in periostracal sheath, may or may not be completely retracted within valves when animal contracts; foot well developed, usually truncate and adapted for suction, atrophying in adults of callum-producing forms. [Members of this family bore into stiff clays or muds, shales, friable or soft rock, shells, poor grade cement, wood, nuts, or other plant products.] ?*Carb., Jur.-Rec.*




FIG. E166. Pholadidae (Pholadinae) (p. N707-N708).

Subfamily PHOLADINAE Lamarck, 1809

[*nom. transl. et correct.* VOKES, 1967, p. 326 (*ex family Pholadaires* LAMARCK, 1809, p. 319)] [=Pholadina GRAY, 1847, p. 187]

Shell elongate, rounded or beaked anteriorly and not closed by callum in adult; number of accessory plates variable, but hypoplax always lacking and siphonoplax lacking except in *Talona*, where a periostracal siphonoplax occurs, making this genus a transition form with the Martesiinae; apophyses present; condyles and umbonal-ventral rib and sulcus absent except in *Zirfaea*, which has weak rib, ventral condyles, and sulcus barely visible in adult,

thus also seeming to be transitional with the Martesiinae. Foot well developed, not atrophying in adult. Animal not capable of retraction within shell. Gill with two demibranchs. Accessory plates in Pholadinae shown in Figure E165. *Cret.-Rec.*

Pholas LINNÉ, 1758, p. 669 [**P. dactylus*; SD CHILDREN, 1822, p. 82] [=Pholadites DAVILA, 1767, p. 169 (refers to D'ARGENVILLE, 1757, pl. 26, fig. H, =**P. dactylus* LINNÉ) (obj.); *Polas* GRONOVIVUS, 1781, p. 259 (*nom. null.*); *Hypogaea* POLI, 1791, p. 29 (obj.); *Hypogaeoderma* POLI, 1795, p. 251, 257 (obj.); *Pholas* LAMARCK, 1799, p. 90 (*non* LAMARCK, 1801, p. 127) (obj.); *Phloas* TURTON, 1802, p. 172 (*nom. null.*); *Pholadites* VON SCHLOTHEIM, 1813, p. 105 (*nom. van.*); *Pholalites* SCHLAEPFER, 1821, p. 278 (*nom. null.*); *Xylotrya* MENKE, 1830, p. 121 (obj.); *Dactylina* GRAY, 1847, p. 187 (*non* ZBORZEWSKI, 1843) (obj.); *Pholadarius* HERRMANNSEN, 1852, Suppl., p. 105 (*nom. null.*); *Pholalithes* PAETEL, 1875, p. 160 (*nom. null.*); *Pragmopholas* FISCHER, 1887, p. 1133 (obj.)]. Rock borers occurring in temperate and tropical seas. Shell elliptical, rounded or beaked anteriorly with septate umbonal reflections and 3 dorsal accessory plates; protoplax thin, calcareous, and divided longitudinally; mesoplax transverse, calcareous, and solid; metaplax calcareous, long and narrow. *Cret.-Rec.*, E.Atl.-W. Atl.-E.Pac.-IndoPac.

P. (*Pholas*). Shell beaked anteriorly, nuclei of divided protoplax near posterior outer margin. *Mio.-Rec.*, Eu.-N.Afr.—FIG. E165,5; E166,1.




FIG. E167. Pholadidae (Pholadinae) (p. N708).




FIG. E168. Pholadidae (Pholadinae) (p. N708).

**P. (P.) dactylus* LINNÉ, Rec., Malta; E165,5a, entire specimen (dorsal) showing protoplax, mesoplax, and metaplax in place; E165,5b, long, divided protoplax (dorsal); E165,5c,d, mesoplax (dorsal) showing variations; E165,5e, metaplax (dorsal); E166,1a,b, RV int. and LV ext. showing septate umbonal reflection, $\times 0.5$ (921a).

P. (Monothyra) TRYON, 1862, p. 205 [**Pholas orientalis* GMELIN, 1790, p. 3216; OD]. Shell beaked anteriorly; protoplax in one piece, with central nucleus. [The protoplax of this species has twice been described as a *Scutum*.] *Cret.-Rec.*, IndoPac.—FIG. E167,1. **P. (M.) orientalis* GMELIN, Rec., Singapore; 1a,b, RV ext., int., $\times 1.2$; 1c,d, inner and outer sides of protoplax, $\times 1.3$ (Turner, n).

P. (Thovana) GRAY, 1847, p. 187 [**Pholas oblongatus* SAY, 1822, p. 320 (= **P. campechiensis* GMELIN, 1790, p. 3216); OD] [= *Gitocentrum* TRYON, 1862, p. 203 (obj.); *Gitoventrum* PAETEL, 1875, p. 86 (*nom. null.*)]. Shell rounded anteriorly, nuclei of divided protoplax anterior and more or less centrally located. *Mio.-Rec.*, W.Atl.-E.Pac.—FIG. E165,6; E168,1. **P. (T.) campechiensis* GMELIN, Rec., Trinidad; E165,6a,b, long, divided protoplax (dorsal, lat.); E165,6c,d, mesoplax (dorsal) in normal position and tipped forward to show basal flange; E165,6e, metaplax (dorsal); E168,1a,b, LV ext., int., $\times 0.75$ (921a).

Barnea LEACH in Risso, 1826, p. 376 [**B. spinosa* Risso, 1826, p. 376 (= **Pholas candida* LINNÉ, 1758, p. 669); OD] [= *Barnia* GRAY, 1840, p. 150; 1842, p. 76 (*nom. null.*); *Barnia* LEACH, 1852, p. 254 (obj.); *Holopholas* FISCHER, 1887, p. 1133 (obj.)]. Shell broadly oval to elliptical, beaked or rounded anteriorly; accessory plate a simple, calcareous, lanceolate protoplax; umbonal reflection simple; pedal gape ranging from narrow slit to broad oval. Sculpture consisting of concentric ridges and radial ribs. [Rock and peat borers.] *Mio.-Rec.*, cosmop.

B. (Barnea). Shells elliptical, rounded at both ends with slitlike pedal gape and slight posterior gape.

Mio.-Rec., E.Atl.-IndoPac.—FIG. E165,1; E169,1. **B. (B.) candida* (LINNÉ), Rec., France; E165,1a-c, protoplax (dorsal, ventral, lat.); E169,1a-c, LV ext., RV int., opposed valves (dorsal) with protoplax in place, $\times 1$ (921a).

B. (Anchomasa) LEACH, 1852, p. 253 [**Anchomasa pennatiana* LEACH, p. 253 (= **Pholas parva* PENNANT, 1777, p. 77); OD] [= *Anchomosa* LAMY, 1925, p. 79 (*nom. null.*); *Barnea (Umitateka)* HABE, 1952, p. 241 (type, *Pholadomya japonica* YOKOYAMA, 1920, p. 106, = *Barnea dilatata* SOULEYET; OD)]. Shell broadly to narrowly elliptical, beaked anteriorly with large oval pedal gape, rounded to truncate posteriorly with moderate to large posterior gape. *Plio.-Rec.*, cosmop.—FIG. E165,2; E170,2. *B. (A.) truncata* (SAY), Rec., USA (Mass.); E165,2a-c, protoplax (dorsal, ventral, lat.); E170,2a,b, LV ext., int., $\times 1$; E170,2c,d, entire animal, dorsal view showing siphons and protoplax in place and ventral view showing foot, $\times 1$ (921a).—FIG. E170,1. **B. (A.) parva* (PENNANT), Rec., Eng.; 1a,b, LV ext., RV int., $\times 1$ (921a).

Clavipholas CONRAD, 1868, p. 728 [**Pholas pectorosa* CONRAD, 1852, p. 200; OD]. Shell elongate, subtrigonal in outline, gaping slightly anteriorly, gape apparently closed by callum in adult; umbonal-ventral ridge and sulcus present, narrow and extending in sinuous curve from umbo to ventral margin. Sculpture on anterior slope consisting of strong radial ribs and rather weak concentric ridges. Accessory plates, apophyses, etc., unknown. *U.Cret.*, N.Am.—FIG. E171,1. **C. pectorosa* (CONRAD), USA (N.J.); 1a,b, dorsal and lat. views of holotype, $\times 1.5$ (889).




FIG. E169. Pholadidae (Pholadinae) (p. N708).




FIG. E170. Pholadidae (Pholadinae) (p. N708).

Cyrtopleura TRYON, 1862, p. 201 [**Pholas crucifera* SOWERBY, 1849, p. 489 (= **Pholas cruciger* SOWERBY, 1834, p. 69); SD STOLICZKA, 1871, p. 22]. Shell rounded or beaked anteriorly; accessory plates consisting of thin, largely chitinous, triangular to T-shaped protoplax and heavy transverse, calcareous mesoplax which may be in one or two pieces; umbonal reflection well separated from umbos and supported at posterior end where sockets are formed for reception of anterior projections of mesoplax; apophyses broad and flattened, or more or less spoon-shaped. [Clay, peat, and soft rock borers.] *Mio.-Rec.*, E.Pac.-E.N.Am.-C.Am.-E.S. Am.

C. (*Cyrtopleura*). Shell beaked anteriorly, with broad oval pedal gape. *Rec.*, E.Pac.—FIG. E165, 3; E171, 2. **C.* (*C.*) *cruciger* (SOWERBY), W. Panama; E165, 3a, protoplax (dorsal); E165, 3b, mesoplax (dorsal); E171, 2a, b, LV ext., RV int., $\times 1$; E171, 2c, d, dorsal and ventral views of opposed valves showing large chondrophore and small apophysis, $\times 1$ (921a).

C. (*Scobinopholas*) GRANT & GALE, 1931, p. 431 [*nom. subst. pro Scobina* BAYLE, 1880 (*non* LEPELETIER, 1825; *nec* WADE, 1917) (*nom. van. pro Pholas* LAMARCK, 1801, p. 127)] [**Pholas costata* LINNÉ, 1758, p. 669; OD] [= *Pholas* LAMARCK, 1801, p. 127 (*non* LINNÉ, 1758; *nec* LAMARCK, 1799) (type, *Pholas costata* LINNÉ, 1758) (obj.); *Pholadigenus* RENIER, 1807, pl. vii (type, *P. costata* LINNÉ; OD, work rejected, ICZN Opinion 427); *Pholas* ADAMS & ADAMS, 1856, p. 325 (*non* LINNÉ, 1758) (type, *Pholas costata* LINNÉ, 1758; SD BAYLE, 1880) (obj.); *Leuconyx* ADAMS & ADAMS, 1863, p. 18; 1865, p. 755 (type, *L. tayleriana* ADAMS & ADAMS, = *P. costata* LINNÉ) (based on apophysis only) (*nom. oblit.*); *Leuconyx* CROSSE, 1868, p. 298 (*nom. null.*)]. Shell rounded anteriorly with slitlike pedal gape. [The isolated apophysis of this species has been described as a *Capulus* and as noted above, the genus *Leuconyx* was created for them.] *Mio.-Rec.*, E.N.Am.-E.S.Am.—FIG. E165, 4; E172, 1. **C.* (*S.*) *costata* (LINNÉ), *Rec.*, USA

(Fla.); 165, *4a,b*, protoplax and mesoplax (dorsal); E172, *1a,b*, LV ext., RV int., $\times 0.7$; E172, *1c,d*, mesoplax (dorsal, ventral), $\times 1.7$; E172, *1e-g*, RV apophysis showing cavity and curvature, $\times 1.7$ (921a).




FIG. E171. Pholadidae (Pholadinae) (p. N708-N709).

Talona GRAY, 1842, p. 76, 91 [**Pholas clausa* HANLEY, 1842, p. 6, 336 (= *Pholas explanata* SPENGLER, 1792, p. 91); OD] [= *Talona* GRAY, 1840, p. 154 (*nom. nud.*); = *Talena* SOWERBY, 1849, p. 498 (*nom. null.*)]. Shells elongate, rounded and closed anteriorly, rounded and gaping posteriorly; reflection of dorsal margin anterior to umbos narrow and nearly erect; umbonal-ventral sulcus lacking. Sculpture consisting of concentric ridges over entire surface and radial ribs on anterior slope and disc, ribs imbricate where they cross ridges; apophyses small, short, and thin. Mesoplax small, transverse, and composed of 2 triangular pieces. Posterior end of valves with reflected lobes which are composed largely of periostracum (seldom present in dried specimens) and form incipient siphonoplax. [Probably a transition genus between *Cyrtopleura*, which has a similar mesoplax and very thin protoplax composed almost entirely of periostracum and the Martesiinae, which have a siphonoplax.] *Rec.*, W.Afr.—FIG. E173, 1. **T. explanata* (SPENGLER), Senegal; *1a,b*, RV ext., LV int.; *1c,d*, ventral and dorsal views of opposed valves showing incurving of valves, remains of siphonoplax, and left half of divided mesoplax remaining in position; all $\times 1$ (Turner, n).

Zirfaea LEACH in GRAY, 1842, p. 76 [**Pholas crispata* LINNÉ, 1776, p. 111; SD GRAY, 1847, p. 188] [= *Zirfaea* GRAY, 1840, p. 154 (*nom. nud.*); *Thurlosia* CATLOW & REEVE, 1845, p. 3 (obj.); *Zirphaea* LEACH, 1852, p. 252 (obj.); *Zyrphaea* CLESSIN, 1891, p. 7 (*nom. null.*)]. Shells elongate-oval in outline, beaked anteriorly, rounded to truncate posteriorly, widely gaping at both ends and with weak umbonal-ventral sulcus and internal ridge which may not be evident in older specimens; apophyses solid, commonly broad and spoon-shaped; single accessory plate (mesoplax), small and more or less triangular in outline. [Soft clay, mud and peat borers.] *Mio.-Rec.*, N.Atl.-N. Pac.—FIG. E165, 7; E174, 1. **Z. crispata* (LINNÉ), *Rec.*; E165, *7a-c*, mesoplax (dorsal, ventral, lat.) (921a); E174, 1, USA (Mass.) (*1a-c*), USA (N.H.) (*1d-f*); E174, *1a*, RV int. showing umbonal-ventral ridge and apophysis, $\times 1.5$; E174, *1b*, LV ext. showing umbonal-ventral sulcus, $\times 1.5$; E174, *1c*, LV int. showing muscle scars, $\times 1$; E174, *1d*, dorsal view of entire specimen showing anterior adductor muscle covered only by periostracum with small mesoplax at posterior end, $\times 1$; E174, *1e*, side view of entire specimen showing foot and extent of periostracal covering on siphons, $\times 1$; E174, *1f*, ventral view of entire specimen showing foot, anterior muscular collar of mantle, and extent to which siphons can be contracted, $\times 1$ (921a).

Zirlona FINLAY, 1930, p. 257 [**Pholadidea increnata* MARWICK, 1929, p. 914; OD]. Shell rounded anteriorly and only slightly gaping. Umbonal-ventral ridge and sulcus present. Anterior slope




FIG. E172. Pholadidae (Pholadinae) (p. N709).

sculptured with strong regularly spaced concentric lamellae which are slightly crenulate; disc and posterior slope sculpture with concentric ridges; radial sculpture lacking. Accessory plates unknown. *Tert.* (Chatton F.), N.Z.—FIG. E175, 1. **Z. increnata* (MARWICK); LV ext. (holotype), $\times 3$ (594).

Subfamily MARTESIINAE Grant & Gale, 1931

Shell elongate to globular, beaked and gaping anteriorly in young stage, completely closed by calcareous callum in adult in all genera except *Chaceia*, which produces only a partial callum; number of accessory plates variable but always lacking protoplax, apophyses, condyles, umbonal-ventral rib and sulcus present; siphonal tube, chimney, and periostracal laminae on posterior slope present in some forms. Foot well developed in young working stage but atrophied in adult. Siphons capable of complete retraction within shell except in *Chaceia*, which is transitional with Pholadinae. [Species of Martesiinae have been reported from Car-

boniferous strata, but, since the accessory plates are unknown, these are unassignable to subgenera. Plates typical of the subfamily are illustrated in Figure E176.] ?*Carb.*, *Jur.-Rec.*

Martesia SOWERBY, 1824, p. 2 [**Pholas clavata* LAMARCK, 1818, p. 446 (= **P. striata* LINNÉ, 1758, p. 669); OD] [= *Mactresia* GRAY, 1840, p. 154; *Mactesia* PAETEL, 1875, p. 119 (*nom. null.*); *Martesiella* VERRILL & BUSH, 1898, p. 777 (type, *M. (M.) fragilis*; OD); *Hiata* ZETEK & McLEAN, 1936, p. 110 (type, *H. infelix*, = young *M. striata* (LINNÉ); OD) (obj.); *Mesopholas* TAKI & HABE, 1943, p. 109 (type, *M. intusgranosa*, = *M. striata* (LINNÉ); OD) (obj.); *Diploplax* BARTSCH & REHDER, 1945, p. 10 (type, *M. (D.) americana*, = young *M. striata* (LINNÉ); OD) (obj.)]. Shells beaked and widely gaping anteriorly in young; adult with callum; beaks sinuously to sharply truncated, giving young *Teredo*-like appearance. Protoplax lacking, mesoplax variable, circular to cuneiform in outline; metaplax and hypoplax long and narrow, pointed anteriorly, pointed truncate or divided posteriorly. [Small wood-borers.] ?*Carb.* (subgenus unknown), *Jur.-Rec.*, world-wide, temp-




FIG. E173. Pholadidae (Pholadinae) (p. N710).

trop. [= *Martesia* DE BLAINVILLE, 1825, p. 578, 632 (obj.).]

M. (Martesia). Mesoplax of adult broadly oval to circular, cushion-like in adult; flat, semicircular, and located posteriorly beneath anterior adductor muscle in young; metaplex and hypoplex pointed or truncate posteriorly. ?*Carb.*, *Jur.-Rec.*, temp. and trop. seas of world.—FIG. E176,2; E177,1. **M. (M.) striata* (LINNÉ), *Rec.*; E176,2, diagram; E177,1, Trinidad (1a-c), USA (Fla.) (1d-g); E176,2a-c, mesoplax of adult (dorsal, ventral, lat.); E176,2d,e, mesoplax of young (dorsal, ventral) (921b); E177,1a, dorsal view of adult with mesoplax and metaplex in place, ant. to right, $\times 1.2$; E177,1b, ventral view of adult showing callum and hypoplex, ant. to left, $\times 1.2$; E177,1c, side view of adult, ant. to left, $\times 1.2$; E177,1d,e, LV ext., RV int. of young specimens; E177,1f, dorsal view of opposed valves with small, flat mesoplax of young in place; E177,1g,

ventral view of opposed valves showing large pedal gape; all $\times 1.5$ (921b).

M. (Particoma) BARTSCH & REHDER, 1945, p. 5 [**Pholas cuneiformis* SAY, 1822, p. 322; OD]. Mesoplax cuneiform, similar in shape in young and adult; metaplex and hypoplex long, narrow, forked posteriorly and extending over posterior end of valve where they meet and fuse. ?*Carb.*, *Jur.-Rec.*, W.Atl.-E.Pac.—FIG. E176,3; E178,1. **M. (P.) cuneiformis* (SAY), *Rec.*, USA (N.Y.); E176,3a-c, mesoplax (dorsal, ventral, lat.) (921b); E178,1a, dorsal view of adult with mesoplax and forked metaplex in place; E178,1b, dorsal view of young specimen showing similarity of mesoplax to that of adult; E178,1c, ventral view of adult showing forked hypoplex; E178,1d, side view of adult showing fused metaplex and hypoplex; E178,1e, ventral view of opposed valves of young specimen showing pedal gape, apophyses, and chondrophore; E178,1f, RV ext. of young specimen; E178,1g, LV int. of young specimen; all $\times 2.5$ (921b).

Aspidopholas FISCHER, 1887, p. 1137 [**Pholas scutata* DESHAYES, 1824, p. 252; OD] [= *Scutigera* COSSMANN, 1886, p. 25 (non LATREILLE, 1803); *Calyptopholas* LAMY, 1927, p. 180 (type, *Pholadidea (Calyptopholas) cheveyi* LAMY, 1927, p. 180, = *Pholas obtecta* SOWERBY, 1849; OD)]. Shells similar to *Pholadidea*, beaked and gaping anteriorly in young stage; partially closed by callum in adult; umbonal-ventral sulcus and ridge narrow and rather weak; anterior slope sculptured by closely set, finely imbricate concentric ridges and weak radial ribs. Apophyses short, solid, and broad at free end; mesoplax greatly produced, forming cap which may cover anterior end of adult; siphonoplax small, calcareous extension of valves which may be covered with periostracum. Siphons may be retracted completely. [Coral-borers.] *Eoc.*, Eu.; *Rec.*, IndoPac.—FIG. E179,1a. **A. scutata* (DESHAYES), *Eoc.*, France (Paris Basin); specimen showing large mesoplax and lining of tube covering posterior end of shell, $\times 3$ (259, pl. 6, fig. 5).—FIG. E179,1b,c. *A. cheveyi* (LAMY), *Rec.*, Annam; 1b, adult with dorsal plate removed; 1c, specimen with greatly enlarged mesoplax in place; both $\times 1.5$ (520).

Chacea TURNER, 1955, p. 66 [**Pholas ovoidea* GOULD, 1851, p. 87; OD]. Shell beaked anteriorly, gaping widely in young stage, with pronounced umbonal ventral sulcus and partial callum in adult. Mesoplax small, transverse in one piece and *Zirfaea*-like. No other accessory plates present. [Rock borers.] *Rec.*, E.Pac.—FIG. E176,1; E179,2; E180,1. **C. ovoidea* (GOULD), USA (Calif.); E176,1a-c, mesoplax (dorsal, ventral, lat. with ant. at left) (921b); E179,2a,b, LV ext., RV int. of young specimens, $\times 1$; E179,2c, ventral view of opposed valves of young specimen, $\times 1$ (921b); E180,1a,




FIG. E174. Pholadidae (Pholadinae) (p. N710).

side view of entire adult showing partial callum and siphons, $\times 0.6$; E180, 1b, ventral view of young specimen showing foot, large anterior gape, and meeting of valves at ventral condyle, $\times 0.6$; E180, 1c, d, LV ext., RV int. of adult specimens, $\times 0.6$ (921b).

Diplothyra TRYON, 1862, p. 449 [**D. smithii*; OD] [= *Schroteria* TRYON, 1862, p. 221 (nom. oblit.)] (type, *Pholas cordata* TRYON, 1862, p. 221, = *Pholas cordata* GMELIN, 1790) (refers to SCHROTER, 1786, v. 3, p. 554, pl. 9, fig. 22-24)]. Shell *Martesia*-like but with beaks obliquely truncated; mesoplax subquadrate to subtriangular in adult, extending anteriorly between dorsal extension of callum; metaplex and hypoplex pointed anteriorly and forked posteriorly, forked portions extending over posterior margins of valves and fusing with one another. [Small shell- and rock-borers, usually found in oysters and *Haliotis*]. *Rec.*, W.Atl.-E.Pac.

—FIG. E176, 5; E181, 1-4. **D. smithii*, USA (N.Y.); E176, 5a-c, mesoplax of adult (dorsal, ventral, lat.); E176, 5d, e, mesoplax of young (lat., dorsal) (921b); E181, 1a, b, split shell of *Crasso-*




FIG. E175. Pholadidae (Pholadinae) (p. N710-N711).




FIG. E176. Accessory plates of genera in Martesiinae (921b).

straea virginica (LINNÉ) showing attack of *D. smithii*, $\times 1$; E181, 2a-c, ventral and 2 dorsal views showing extension of callum around mesoplax and differences in shape of mesoplax; $\times 3.5$; E181, 3a,b, RV ext., RV int. of adult, $\times 3.5$; 4a,b, RV ext., LV int. of young specimen, $\times 4.5$ (921b).

Eutylus VINCENT, 1891, p. 164, pl. 4, fig. 1-4 [**Pholadomya cuneata* SOWERBY, 1844, p. 40; OD] [= *Phenacomya* DALL, 1898, p. 823 (obj.)]. Probably close to *Pholadidea*. Anterior slope sculptured with rather weak concentric ridges and only a few rather widely spaced radial ribs; umbonal-ventral sulcus weak; anterior margin sinuous; pedal gape small; callum ribbed, showing extension of radial

ribs of anterior slope. Dorsal plates unknown. Apophyses probably present. [Mud-dweller.] *Eoc.*, Eu.—FIG. E182, 1. **E. cuneata* (SOWERBY), Belg.; 1a, RV ext.; 1b, ant. view of opposed valves; 1c, int. cast showing pallial sinus; 1d, dorsal view of opposed valves; all $\times 1$ (941).

Formosulus ZHIZHENKO, 1934, p. 44 [**F. lucidus*; OD] [= *Formesulus* ZHIZHENKO, 1939 (errore NEAVE, p. 736)]. Shell small, more or less rectangular in outline, beaked anteriorly and divided into 3 distinct areas by 2 furrows extending from umbos in ventral margin. [Description and figure poor. Described as a hiatellid bivalve (Hiatellacea) but now regarded by Russian paleon-

tologists as probably nonmolluscan, a cirriped plate, fide VOKES, 1967, p. 346.] *Mio.*, USSR (Ciscaucasia).—FIG. E161,6. **F. lucidus*; LV ext., $\times 10$ (Zhizhchenko).

Goniochasma MEEK, 1864, p. 34 [**Xylophaga simpsoni* MEEK & HAYDEN, 1857, p. 141; OD]. Shell *Martesia*-like, ovate-oblong with beak truncated at sharp angle; umbonal-ventral sulcus and ridge oblique, internal ridge faintly crenulate; posterior ridges very oblique, broad, rounded and not extending to margin. Mesoplax present (889, p. 249). [Is probably a young *Martesia*. From fossilized wood.] *U.Cret.*, N.Am.-India.—FIG. E182,2. **G. simpsoni* (MEEK & HAYDEN), USA (Idaho); 2a, LV ext., $\times 2$; 2b, cast of holotype, $\times 1$ (609, pl. 30, fig. 9a-b).

Heteropholas FISCHER, 1887, p. 1136 [**Pholas xylophagina* DESHAYES, 1860, p. 142; OD]. Shell *Martesia*-like but with beaks obliquely to sinuously truncated; umbonal-ventral sulcus very oblique and with 2 additional grooves on outer surface, one anterior, other median. Accessory plates and apophyses unknown. *Eoc.*, Eu.—FIG. E183,1. **H. xylophagina* (DESHAYES), France (Paris Basin); 1a,b, LV ext., int., $\times 2$ (259).

Lignopholas TURNER, 1955, p. 98 [**L. clappi*; SD TURNER, 1956, p. 188]. Shell *Martesia*-like, posterior slope elongated and having series of fringed, overlapping, thin, periostracal lamellae; mesoplax divided longitudinally; other accessory plates lacking; apophyses long and thin, chondrophore prominent. [Small wood borers, in brackish and fresh water.] *Rec.*, W.Atl.-IndoPac.—FIG. E184,1-3. **L. clappi*, Nicaragua; 1a, adult, lat. view, ant. to left; 1b, dorsal view of adult, ant. to right, showing divided mesoplax (holotype), $\times 4$; 2a, young LV ext., ant. to left; 2b, young RV int., ant. to left (paratype), $\times 6$; 3a-c, dorsal, ventral, lat. views of mesoplax of adult, $\times 6$; 3d, dorsal view of mesoplax of young, anterior for all to left, $\times 6$ (921b).

Opertochasma STEPHENSON, 1952, p. 139 [**O. venustum*; OD]. Shell *Martesia*-like but with 2 radial grooves extending from beak to ventral margin in area of umbonal-ventral ridge; posterior slope sculptured. Mesoplax long and narrow, divided longitudinally and extending anteriorly in adult in 2 lobes over umbos; hypoplax lacking. [Probably close to *Parapholas*.] *U.Cret.*, N.Am.—FIG. E183,2. **O. venustum*, Woodbine F., USA (Tex.); 2a, LV ext., 2b-d, dorsal, ant., and ventral views of opposed valves; 2a-c, $\times 3$; 2d, $\times 2$ (890).

Parapholas CONRAD, 1848, p. 121 [**Pholas californica* CONRAD, 1837, p. 226; OD] [= *Paraphola* PAETEL, 1875, p. 152 (*nom. null.*)]. Shell divided into 3 distinct areas, anterior slope, disc, and posterior slope being clearly marked; posterior slope having series of overlapping periostracal plates; accessory plates consisting of oval mesoplax, elongate metaplax, and hypoplax, which may




FIG. E177. Pholadidae (Martesiinae) (p. N712).

or may not be forked posteriorly; siphonoplax lacking; chimney not attached to valves but fitting over end of valves and composed of fine particles which have been ejected from siphons and cemented together. [Moderate-sized to rather large rock-borers.] *Cret.-Rec.*, E. Pac.-Indo Pac.-W. Afr.—FIG. E176,6; E185,1. **P. californica* (CONRAD), *Rec.*, USA (Calif.); E176,6a-c, metaplax (dorsal, ventral, lat.); E176,6d,e, hypoplax (dorsal, ventral); E176,6f-h, mesoplax of adult (dorsal, ventral, lat.); E176,6i,j, mesoplax of young (dorsal, ventral); E185,1a, side view of adult, $\times 1$; E185,1b, ventral view of young specimen showing foot, $\times 1$; E185,1c, adult RV int., $\times 1$; E185,1d, ventral view of adult showing hypoplax in place, $\times 0.7$; E185,1e, dorsal view of adult showing mesoplax and metaplax in place, $\times 0.7$; E185,1f, chimney of specimen in fig. 1d, $\times 0.7$. [Figures show variation in size of adults] (921b).




FIG. E178. Pholadidae (Martesiinae) (p. N712).

Penitella VALENCIENNES in ABEL DU PETIT-THOUARS, 1846, pl. 24 [**P. conradi*; SD HABE, 1952, p. 243] [= *Penicilla* CONRAD, 1854, p. 335 (*nom. null.*); *Navea* GRAY, 1851, p. 385 (type, *N. subglobosa* GRAY, 1851, p. 385; SD TURNER, herein)]. Shell oval in outline, beaked anteriorly in young stage with pronounced umbonal-ventral sulcus, callum partial or complete. Mesoplax *Zirfaea*-like in young but producing dorsal covering in adult; siphonoplax variable, present or absent; no other accessory plates present. [*Navea* GRAY was based upon a young form. Shale- and rock-borers.] *Mio.-Rec.*, E.Pac.-N.Pac.—FIG. E176,4; E186, 1-4. **P. conradi* VALENCIENNES; *Rec.*, USA (Calif.); E176,4a-c, mesoplax of adult (dorsal, ventral, lat.); E176,4d,e, mesoplax of young (dorsal, ventral); E186,1, adult specimens in substrata (type from VALENCIENNES), $\times 1$; E186,2a,b, adult LV ext. taken from shale and RV int. of same specimen showing muscle scars and siphonoplax, $\times 3$; E186,3a, dorsal view of adult showing mesoplax in position, $\times 3$; E186,3b,c, dorsal and LV int. views of adult taken from *Haliotis*, $\times 2$; E186,4a,b, LV ext. and RV int. of young specimen taken from shale, $\times 4.5$ (921b).

Pholadidea TURTON, 1819, p. 147 [**P. loscombiana*; OD] [= *Pholadidoidea* DE BLAINVILLE, 1826, p. 535 (obj.); *Pholidea* SWAINSON, 1835, p. 30 (obj.); *Pholadidae* SOWERBY, 1839, p. 81 (*nom. null.*); *Pholidae* SWAINSON, 1840, p. 364 (*nom. null.*); *Talonella* GRAY, 1851, p. 385 (type, *T. tridens* GRAY, 1843; OD); *Cadmusia* GRAY, 1852, p. 254 (obj.); *Pholameria* CONRAD, 1865, p. 2 (type, *Pholas triquetra* CONRAD, 1848, p. 127, pl.

13, fig. 3; OD); *Pholididae* PAETEL, 1875, p. 160 (*nom. null.*)]. Shell beaked anteriorly in young stage, closed by callum which extends dorsally over beaks in adult; rounded to truncate posteriorly, divided into 2 areas by prominent umbonal-ventral sulcus; posterior slope not clearly defined. Mesoplax variable, divided longitudinally in young but may be greatly produced in adult; incipient metaplast and hypoplast present or absent, not existing as separate plates but resulting from deposition of calcium in periostracum uniting valves posterior to umbos; siphonoplax commonly tubular, composed largely of periostracum and fused; siphonal tube present in most species. Apophyses short, fragile and not broadening at free end. [Shale, soft-rock and coral-borers.] *Eoc.-Rec.*, E.Pac.-IndoPac.-E.Atl.

P. (*Pholadidea*). Umbonal reflection raised well above umbos; mesoplax small, lacking basal portion; siphonoplax cuplike; siphonal tube lacking. *Eoc.-Rec.*, E.Atl.-IndoPac.—FIG. E187,1-3. **P.* (*P.*) *loscombiana* TURTON, *Rec.*, Eng.; 1a,b, adult RV ext., and LV int., showing small apophysis and large chondrophore, $\times 1.5$; 2, dorsal view of opposed valves with mesoplax and siphonoplax in place, $\times 1.5$; 3, lateral view of adult with siphonoplax, $\times 1.5$ (921b).


P. (*Hatasia*) GRAY, 1851, p. 385 [**Pholas melanura* SOWERBY, 1834, p. 70; SD STOLICZKA, 1871, p. 21] [= *Hatoisia* SCHAUFUSS, 1869, p. 17 (*nom. null.*); *Hatasia* PAETEL, 1875, p. 91 (*nom. null.*)]. Umbonal reflection closely appressed, mesoplax in young stage flat, semi-circular and in one piece, upper part, added in

adult stage, divided longitudinally, young basal part extending beyond it; siphonoplax composed largely of periostracum, variable in shape and size. Siphonal tube attached to siphonoplax. *Rec. E.Pac.*—FIG. E188,1. **P. (H.) melanura* (SOWERBY), Panama (1a,b), Mexico (1c); 1a, adult LV ext. with siphonoplax; 1b, adult RV int.; 1c, dorsal view of adult showing extension of callum over beaks and mesoplax, all $\times 1$ (921b).—FIG. E188,2. *P. (H.) tubifera* (SOWERBY), Panama; lateral view of adult showing siphonal tube attached to the siphonoplax, $\times 2$ (921b).


Ramsetia STEPHENSON, 1941, p. 250 [**R. whitfieldi*; OD]. Shell *Martesia*-like, beaked and gaping anteriorly in young but closed by callum in adult; beaks diagonally truncated; umbonal-ventral ridge and sulcus present. Sculpture consisting of coarse concentric ridges which are crossed on anterior slope by weakened radial ribs. Accessory plates unknown. *U.Cret.*, N.Am.—FIG. E183,3. **R. whitfieldi*, Navarro Gr., USA (Tex.); 3a, LV ext. (holotype); 3b, RV ext. of another specimen; 3c, dorsal view of opposed valves; all $\times 1$ (889).

Teredina LAMARCK, 1818, p. 438 [**Fistulana personata* LAMARCK, 1806, p. 429, 1808, pl. 43, fig. 6-7; SD CHILDREN, 1823, p. 81] [= *Fistulana* LAMARCK, 1806 (non LAMARCK, 1789) (obj.); *Teridina* SOWERBY, 1839, p. 117 (nom. null.)]. Shell *Teredo*-like, particularly in young stage, but anterior gape closed by callum in adult; umbos covered by large, 4-lobed mesoplax which extends anteriorly; apophyses present; long, thick calcareous tube fused to valves anteriorly and in some shells divided by longitudinal partitions posteriorly; calcareous tube probably homologous with that found in some *Pholadidea*, to which seemingly it is most closely related. *U.Cret.-M.Mio.*, Eu.—FIG. E189,1-6. **T. personata* (LAMARCK), France (Paris Basin); 1, entire specimen; 2a,b, dorsal and ventral views of opposed valves, mesoplax removed; 3a,b, dorsal and ant. views of opposed valves showing lobed mesoplax; 4, internal view of shells showing apophyses; 5, abnormal post. end of tube with 2 openings; 6a,b, posterior ends of tubes showing internal ribbing; all $\times 1$ (259, pl. 3, fig. 10-15, 17-20).


Xylophomya WHITFIELD, 1902, p. 75, pl. 27-29 [**X. laramiensis*; OD]. Shell somewhat *Xylophaga*-like in general outline, anterior gape being closed by broad shieldlike plate (callum) in adult form; widely gaping posteriorly, with posterior margin of valves prolonged by thin, smooth tongue-shaped extensions, which probably fuse with lining of tube. Narrow umbonal-ventral sulcus present. Dorsal margin of RV with 4 minute toothlike denticles anterior to umbo. [Wood-boring.] *Cret.*, USA (Wyo.).—FIG. E190,1. **X. laramiensis*, Laramie Gr.; 1a, outline showing form of RV; 1b, composite sketch from several




1a



1b




1c




2a

Chaceia



2b



2c

FIG. E179. Pholadidae (Martesiinae) (p. N712).

specimens to show sculpture and callum; 1c, ant. end (reconstr.); 1d, ant. teeth as seen on fragment chipped from between beaks; 1e, outline of dorsal view of ant. end; 1a-c, 1e, $\times 1$; 1d, $\times 3$ (982).

Subfamily JOUANNETIINAE Tryon, 1862

[nom. correct. TURNER, 1955 (pro Jouannetinae TRYON, 1862)]

Shells in young stage more or less globu-




FIG. 180. Pholadidae (Martesiinae) (p. N712-N713).

lar, equivalved, beaked and widely gaping anteriorly; apophyses lacking; umbonal-ventral rib and sulcus present but weak; adult valves may or may not be equal; if inequivalved, callum closing pedal gape usually greatly produced anteriorly and entirely calcareous; if equivalved, callum flat and composed largely of periostracum. Rudimentary mesoplax present in some forms. Siphonoplax calcareous and variable in shape, unequal in *Jouannetia*. Foot well developed in young stage, atrophied in adult. Siphons capable of complete retraction within shell. Gills with two demibranchs. *U. Cret.-Rec.*

Jouannetia DESMOULINS, 1828, p. 244 [*]. *semicaudata*; OD] [= *Jouannetia* GRAY, 1840, p. 154 (*nom. van.*); *Juannetia* D'ORBIGNY, 1846, p. 737 (*nom. null.*); *Juannetia* DESMAREST, in CHENU, 1859, p. 231 (*nom. null.*)]. Shells in young stage equivalved, beaked and widely gaping anteriorly, closed posteriorly; mesoplax, if present, in young shell more or less semicircular in outline and located in normal position transversing both valves, but displaced in adult by unequal growth of callum on LV; callum very large, that of LV being largest and overlapping callum of RV. Siphonoplax of LV, if present, always much smaller than that of RV, L siphonoplax always smooth, while R may be pectinate or smooth. Shell with or without an inwardly projecting lamina for the attachment of posterior adductor and pedal mus-

cles. [Shale- and soft-rock-borers.] *U.Cret.-Rec.*, cosmop., temp. and trop. seas.

J. (*Jouannetia*). Shell with special lamina for attachment of muscles; margin of siphonoplax smooth. *U.Cret.-Rec.*, cosmop. in temp. and trop. seas.—FIG. E191, 1. *J. (*J.*) *semicaudata* DESMOULINS, ?Eoc., France (Mérignac, Bordeaux); *1a, b*, right and left sides of complete adult show-

ing overlap of callum from LV; *1c*, adult RV int. showing siphonoplax; *1d*, adult LV int. showing lack of siphonoplax; *1e-f*, RV int. and LV int. without callum, showing laminae for attachment of posterior adductor muscle; *1g*, ant. view of opposed valves without callum; *1h*, complete specimens as found in calcareous rock; all $\times 1$ (648).




FIG. E181. Pholadidae (Martesiinae) (p. N713-N714).




FIG. E182. Pholadidae (Martesiinae) (p. N714-N715).

J. (Pholadopsis) CONRAD, 1849, p. 156 [*Pholadopsis pectinata*; OD] [=Triumphalia SOWERBY, 1849, p. 500 (type, *Pholas globosa* QUOY) (=Pholas globulosa QUOY & GAIMARD; SD SOWERBY, 1850); Triumphalia PAETEL, 1890, p. 5 (nom. null.); Triumphala CLESSIN, 1892, p. 33 (nom. null.)]. Shells without special lamina for attachment of muscles; margin of siphonoplax pectinate; mesoplax small and displaced in adult. *Paleoc.-Rec.*, Eu.-W.Atl.-E.Pac.-IndoPac.—FIG. E192,1. ***J. (P.) pectinata** (CONRAD), *Rec.*, USA (Calif.); 1a, adult RV ext. showing callum not extending to mid-line and pectinate siphonoplax; 1b, LV int. showing greatly enlarged callum which overlaps RV, smaller smooth siphonoplax, and posterior adductor muscle scar just posterior to umbo; 1c, dorsal view of opposed valves showing greatly enlarged LV callum which has displaced mesoplax; all $\times 1.5$ (1a,b, holotype; 1c, paratype) (921b, 922).

Nettastomella CARPENTER, 1865, p. 202 [*Nettastoma darwinii* SOWERBY in CARPENTER, 1865 (=Pholas rostrata VALENCIENNES, 1846, non darwinii SOWERBY); OD] [=Nettastoma CARPENTER, 1864, p. 637 (non Nettastoma RAFINESQUE, 1810¹); Nettastomella PAETEL, 1875, p. 138 (nom.

null.); Nettastoma LAMY, 1926 (nom. null.)]. Large pedal gape of young only partially closed by calcareous callum which forms band along anterior margin of shell in adult, equal on both valves or much wider on LV; periostracal callum with minute central pore filling most of pedal gape in adult. Siphonoplax, present on both valves or on RV only, may be short or long, straight, diverging or tapering and tubelike. Mesoplax, if present, small and fused with dorsal extension of callum on LV. [Shale- and soft-rock-borers.] *Plio.-Rec.*, N.Pac.-S.Atl.-S.Pac.—FIG. E193,1; E194,1. ***N. rostrata** (VALENCIENNES), *Rec.*, USA (Calif.); E193,1a, large LV ext. with wide callum and short siphonoplax; E193,1b, small LV ext. with long siphonoplax; E193,1c, dorsal view of opposed valves; E193,1d, RV int. with siphonoplax broken off; E193,1e, ventral view of specimen beginning to produce siphonoplax; all $\times 2.5$; E194,1, specimen with LV removed to show muscles, siphons and periostracal callum with open central pore, $\times 4.5$ (921b, 922).

Scyphomya DALL, 1898, p. 822 [*Pholas semicostata* H. C. LEA, 1844, pl. 24, fig. 1; OD]. Shell similar to young *Jouannetia* (*Pholadopsis*). [The location of the type of *P. semicostata* LEA is unknown and specimens labeled with this name by DALL proved to be young *J. (Pholadopsis)*, a genus not known to occur in the Atlantic at the time DALL was working. *Scyphomya* probably is a synonym of *Pholadopsis*.] *Tert.-Rec.*, W.Atl.

¹ Although according to the present rules of Zoological Nomenclature a name differing in one letter is not considered a homonym, this was not the case at the time CARPENTER himself instituted a new name for his *Nettastoma*. Since *Nettastomella* has been in general use since CARPENTER introduced it in 1865 it is being retained in accordance with Article 23b.

Subfamily XYLOPHAGAINAE Purchon, 1941


[*nom. correct.* TURNER, herein (*pro* Xylophaginae TURNER, 1955, *non* LATREILLE, 1807) *nom. transl. ex* Xylophagidae LATREILLE, 1807 (*nom. correct.* SCHINER, 1868, *pro* family Xylophagi LATREILLE, 1807, Insecta)] [=Xylophaginidae PURCHON, 1941]

Shells small, globular, beaked and gaping anteriorly throughout life; beaks truncated at nearly right angles, giving shell a *Teredo*-like appearance. Apophyses absent. Anterior slope sculptured with numerous rows of finely denticulated ridges, as in *Teredinidae*; disc and posterior slope sculptured only by growth lines and low concentric ridges; umbonal-ventral ridge and sulcus well developed; ventral condyle usually present but weak; dorsal condyle lacking; chondrophore well developed. Accessory plates consisting only of small divided mesoplax, which varies greatly in shape. Posterior adductor muscle scar large, located high on posterior slope. Foot well developed, truncate and not atrophying in adult. Siphons commonly capable of retraction within shell, and quite variable. Excurrent siphon may be shorter than incurrent siphon and in some species exists only as series of lappets on dorsal side of incurrent siphon. Gills with single demibranch. Length of the tunnel ranging from slightly longer than the shell to about ten times the length of the valves, the posterior end often lined with agglutinated particles ejected from the excurrent siphon forming a chimney similar to that in *Parapholas*.¹ *U.Cret.-Rec.*


Attempts have been made to erect several genera on the basis of dorsal plate characters and type of siphons, but KNUDSEN (1961) has shown that these do not group naturally and that on the basis of our present knowledge there is no way to divide *Xylophaga*.

Xylophaga TURTON, 1822, p. 258 [**Teredo dorsalis* TURTON, 1819, p. 185; OD] [= *Xilophaga* GEINITZ, 1845, p. 397 (*nom. null.*); *Xylotomea* DALL, 1898, p. 821 (new name *pro* *Xylophaga* TURTON [*non* *Xylophagus* MEUSCHEN, 1778; *nec* MEIGEN, 1803]—name not needed as *Xylophaga* and *Xylophagus* are not considered homonyms); *Xilotoma*


GIGOUX, 1934, p. 285 (*nom. null.*); *Protoxylophaga* TAKI & HABA, 1945, p. 112 (type, *X. tomlini* PRAESHAD; OD); *Neoxylophaga* TAKI & HABA, 1950, p.



Heteropholas



Opertochasma



Ramsetia

¹ A remarkable new genus of bathyal Xylophaginae was discovered while this work was in press. An account of it is being prepared now for publication. The two species (Western Atlantic and Eastern Pacific) are very *Teredo*-like in habit, making tunnels over 15 cm. long, the posterior two-thirds of which have a thin calcareous lining. If wood containing species of this genus were fossilized it probably would be impossible to distinguish them from teredinids.




FIG. E184. Pholadidae (Martesiinae) (p. N715).

46 (type, *X. rikuzenica* TAKI & HABE; OD); *Meta-xylophaga* TAKI & HABE, 1950, p. 47 (type, *M. supplicata* TAKI & HABE; OD). Shell globose, *Teredo*-like, apophyses lacking, mesoplax divided and variable in shape and size. Chondrophore and internal ligament present. Siphons variable. [Small borers usually found in water-logged plant material in deep water.] *U.Cret.-Rec.*, cosmop.—FIG. E195,1,2. **X. dorsalis* (TURTON), Rec., Scot.; E195,1a, LV ext. showing umbonal-ventral sulcus; E195,1b, LV int. showing muscle scars and umbonal-ventral ridge; E195,1c,d, ant. and dorsal views of opposed valves showing pedal gape, condyles, chondrophores, and mesoplax in place; E195,2a,b, dorsal and ventral views of mesoplax (921b).

Xylophagella MEEK, 1864, p. 34 [**Xylophaga elegantula* MEEK & HAYDEN, 1857, p. 141; OD]. Shell *Xylophaga*-like, globose, beaks sharply truncated; umbonal-ventral sulcus and ridge present and extending from umbos in nearly straight line to ventral margin; internal ridge strongly crenate; second posterior internal ridge extending from umbos obliquely to posterior ventral margin. Accessory plates and apophyses not known. Burrows without shelly lining. *U.Cret.*, N.Am.—FIG. E190,2. **X. elegantula* (MEEK & HAYDEN), USA (Idaho); 2a, LV ext.; 2b, LV int. cast; 2c-e, post., ant., post. views of opposed valves; 2f, side view of holotype, $\times 1$ (2a-e, $\times 3.5$) (609).

Family TEREDINIDAE Rafinesque, 1815

[*nom. correct.* DEKAY, 1843, p. 249 (*pro* *Teredaria* RAFINESQUE, 1815, p. 148)] [= *Teredinites* LATREILLE, 1825, p. 224; *Teredinadae* FLEMING, 1828, p. 409; *Teredina* GRAY, 1847, p. 188; *Teredidae* CARPENTER, 1861, p. 248; *Teredinadae* TRYON, 1862, p. 459 (*nom. null.*)]

Shells greatly reduced, covering only anterior tip of long wormlike animal (Fig. E164,1a). Burrow usually with calcareous lining which varies greatly in thickness and is in some species divided or has septa at distal end. Valves divided into three parts as shown in Figure E164,1b,c. Condyles (on which valves rock during boring process), umbonal-ventral ridge and apophyses prominent. Chondrophore and small internal ligament present in all species examined by author. Siphons separate or united, relatively short and with pair of pallets inserted at base which serve to close end of burrow when siphons are withdrawn. Gill with only single demibranch. Intestine not transverse heart, except in *Kuphinae*. Visceral mass extending in loop posterior to shell, as shown in Figure E162,2. Caecum present, except in the *Kuphinae*. ?*Cret.*, *Paleoc.-Rec.*

The classification of genera is based on the morphology of the soft parts and on the

pallets. The species are usually distinguished on the basis of pallets and siphons. The shells are almost useless in classification as they are extremely variable, depending upon the age of the specimen and the ecologic conditions affecting it. Similar shell forms may appear in widely separated genera.

Though many fossil species have been named, some as early as the Lower Cre-

taceous, most have been instituted on the basis of shells and tubes only, and consequently they cannot be placed definitely in any Recent genus. Pallets of *Bankia* have been recorded from the Paleocene of Iraq (ELLIOTT, 1963, p. 315) and DURHAM & ZULLO (1961, p. 1) described a *Bankia* from the Middle Oligocene of Washington. *Nototeredo* has recently been discovered by A. M.




FIG. E185. Pholadidae (Martesiinae) (p. N715).




FIG. E186. Pholadidae (Martesiinae) (p. N716).

CVANCARA in the Cannonball Formation (Paleocene) of Morton Co., North Dakota (TURNER, 1966, p. 16, fig. 3; CVANCARA 1966, p. 350, pl. 9, fig. 12, 28). Pallets of *Bankia*, *Teredo*, *Teredora*, and *Nototeredo* are known from the Eocene of the Paris Basin, the Upper Eocene of Barton, Hampshire, England, and from Burgenland Formation, Austria. The fossil record is poor, but from what is known it may be assumed that all Recent genera existed in the Eocene and that the family was worldwide in distribution, for wherever sufficient fossilized wood is found in a marine horizon, evidence

of borers is also eventually found. Unfortunately, the characters which distinguish the species, particularly in the genus *Bankia*, are invested mainly in the periostracum which covers the calcareous portion of the pallets, and this is not preserved in fossils. Pallets, devoid of periostracum, are similar in appearance, and so it is usually impossible to relate fossil forms definitely with any one living species.


Inferred relationships of Recent genera of the Teredinidae are shown diagrammatically in Figure E196.

Subfamily TEREDININAE Rafinesque, 1815

[*nom. transl.* STOLICZKA, 1871, p. 11 (*ex Teredaria* RAFINESQUE, 1815)]


Wood-borers with posterior adductor muscle much larger than anterior one and without muscular collar surrounding body posterior to shell. Calcareous tube lining burrow, varying greatly in thickness but never as heavy as in Kuphinae. Digestive tract elongate, esophagus short, anterior portion of stomach and crystalline style in front of posterior adductor muscle; caecum and intestine extending posteriorly; intestine not transversing heart but passing beneath it, around posterior adductor muscle, and opening into anal canal. Length of gills, as well as size and type of stomach and posterior caecum, variable. These are important in generic classification and probably indicate type of feeding. Pallets nonsegmental in structure, composed of calcareous base covered by periostracum; blade increased in size by addition of material over entire surface. *Eoc.-Rec.*

Teredo LINNÉ, 1758, p. 651 [**T. navalis*; SD ICZN, 1926, opin. 94] [= *Teredigenus* RENIER, 1807, pl. vii (obj.) (ICZN, work rejected); *Toredo* MAY, 1929, p. 652 (*nom. null.*)]. Pallets variable in shape, but with blade always in one piece, periostracum usually thin and closely adhering to calcareous base (in some species may extend beyond calcareous portion distally as narrow border, but never forms cap, as in *Lyrodus*); blade usually sheathing stalk. Siphons usually separate. Young, in all species examined to date, carried in brood



Pholadidea

FIG. E187. Pholadidae (Martesiinae) (p. N716).




Hatasia

FIG. E188. Pholadidae (Martesiinae) (p. N716-N717).

pouch dorsal to gills. Anatomy generally similar to *Lyrodus*. [The genus has been divided into numerous subgenera which are probably of little value, as intergrades are constantly being found.

These, however, are listed and their type species figured.] *Eoc.*, Eu.(Eng.-Belg.-France)-*Rec.*, cosmop.



T. (Teredo) [= *Zopoteredo* BARTSCH, 1923, p. 96 (type, *Teredo* (*Zopoteredo*) *clappi*; OD); *T. (Austroteredo)* HABE, 1952, p. 249 (type, *Teredo* (*Teredo*) *parksi* BARTSCH, 1921, p. 28; OD)]. Pallets white with thin pale yellow to dark brown periostracum covering distal portion of blade; distal margin of outer face U- or V-shaped, but varying from U-shaped to nearly straight on inner margin. *Eoc.-Rec.*, cosmop.—FIG. E197,1. **T. (T.) navalis* LINNÉ, USA(N.Y.); 1a,b, LV ext. and int.; 1c,d, pallet, outer and inner faces (923).

T. (Coeloteredo) BARTSCH, 1923, p. 99 [**T. (C.) mindanensis*; OD]. Pallet blade wedge-shaped, convex on outer surface, concave to flat on inner surface and with deep cup extending nearly to stalk, which is short and swollen at base. *Rec.*, IndoPac.—FIG. E197,2. **T. (C.) mindanensis*, Philip.; 2a,b, RV ext. and int.; 2c-e, pallet, outer and inner faces and side (holotype) (923).


T. (Pingoteredo) IREDALE, 1932, p. 30 [**Teredo shawi* IREDALE, 1932, p. 30; OD] [= *Pingoteredo* HABE, 1952, p. 250 (*nom. null.*)]. Pallets similar in shape to *Teredo* (*Teredo*) but with thick periostracum extending beyond calcareous base, both laterally and distally (best seen with transmitted light). *Rec.*, temp. and trop. seas.—FIG. E197,3. **T. (P.) shawi* IREDALE, Australia; 3a,b, LV ext. and int.; 3c-e, pallet, outer and inner faces and side; 3f,g, pallet of young, outer and inner faces (paratypes) (923).

Bactronophorus TAPPARONE-CANEFRI, 1877, p. 290 [*nom. subst. pro Calobates* GOULD, 1862, p. 283 (*non* KAUP, 1829)] [**Teredo thoracites* GOULD, 1862, p. 283; OD]. Pallets asymmetrical, basal portion of blade more or less triangular, with shallow cup from which issues pustulose, calcareous, dagger-like blade. Siphons united except at tip, surrounded by wide mantle collar. Heart anterior, extending from posterior adductor muscle about 0.3 length of animal. Digestive system with globular stomach and large, thin-walled caecum. Gills broad and flattened ventrally with well-developed food groove. *Rec.*, IndoPac.—FIG. E198,1. **B. thoracites* (GOULD), Australia; 1a,b, RV ext. and int.; 1c,d, pallet, outer and inner faces (923).


Dicyathifer IREDALE, 1932, p. 28 [**Teredo manni* WRIGHT in CALMAN, 1920, p. 395 (= *D. caroli* IREDALE, 1936, p. 38, = *T. manni* WRIGHT, 1866, p. 565); OD] [= *Pseudodicyathifer* TCHANG, TSI & LI, 1958 (type, *Teredo manni* WRIGHT, 1866; OD)]. Pallets simple, solid, almost entirely calcareous, blade more or less triangular in outline, stalk long and heavy. Inner face of blade flat, outer face convex with shallow cup which is partially or almost completely divided by central, longitudinal ridge.

FIG. E189. Pholadidae (Martesiinae) (p. N717).

Shell with very broad anterior slope, disc narrow and posterior slope reduced in mature specimens, typical in young. Tube thick, solid and divided posteriorly. Siphons separate. Mantle collar around



Xylophomya



Xylophagella

FIG. E190. Pholadidae (Martesiinae) (1), (Xylophagainae) (2) (p. N717, N722).




FIG. E191. Pholadidae (Jouanettiinae) (p. N718-N719).

siphons narrow, rather muscular. Gills broad, truncated and extending from posterior tip of gonads to base of siphons. Branchial groove lateral and well developed. Labial palps attached. Stomach globular, caecum short and bulbous, intestine making simple loop around it; digestive gland large, extending posteriorly beyond caecum. Gonads mainly posterior to digestive glands. Heart anterior, about half length of animal; ventricle short and broad; auricles long, tapering and red-brown in color. Kidney large and dorsal to heart. *Rec.*, IndoPac.—FIG. E199, 1. **D. manni* (WRIGHT), Singapore; 1a, b, LV ext., int. (apophysis broken); 1c, RV ant.; 1d, LV post.; 1e, f, pallet, outer and inner faces; 1g, entire animal, showing anatomical characters of genus with section cut out, $\times 2$ (923). *Lyrodus* GOULD in GOULD & BINNEY, 1870, p. 34 [*Teredo chlorotica* GOULD, 1870, p. 33 (= *T. pedicellata* DE QUATREFAGES, 1849, p. 26, non DÖRING, 1885); OD] [= *Teredops* BARTSCH, 1921, p. 26 (type, *T. diegensis* BARTSCH, 1916, p. 48; OD); *Cornuteredo* DALL, BARTSCH & REHDER, 1938, p. 209 (type, *T. (Cornuteredo) milleri* DALL, BARTSCH, & REHDER, 1938, p. 210, = *T. affinis* DESHAYES, 1863, p. 6; OD)]. Pallets with calcareous base and pronounced brown to blackish periostracal cap which can readily be separated from base; calcareous portion narrowly to broadly rounded at distal end and commonly marked with growth lines. Distal margin of periostracal




FIG. E192. Pholadidae (Jouannetiinae) (p. N720).

cap may be straight, broadly curved, or have lateral horns; cap may be solid or have bubble-like cavity which usually breaks open in old specimens, allowing cavity to become filled with debris which solidifies and produces a knoblike process. Siphons short and separate, gills long, bladelike, stomach elongate, caecum large, intestine making simple loop around caecum. Young carried in brood pouch dorsal to gills until veliger stage. *Rec.*, cosmop. in temp. and trop. seas.—FIG. E200, 1-3. **L. pedicellatus* (DE QUATREFAGES), Br. Is. (1), USA (Calif.) (2), USA (Hawaii) (3); 1a,b, LV ext., int.; 1c-e, pallet, outer and inner faces, lat.; 2a-e, corresponding views of holotype of *L. diegensis* (type species of *Teredops*); 3a-d, RV ext., int., outer and inner face of pallet, holotype of *L. milleri* (=lectotype of *L. affinis*, type species of *Cornuteredo*) (923).

Neoterredo BARTSCH, 1920, p. 69 [**Teredo* (*Neoterredo*) *reynei* BARTSCH; OD]. Pallets simple, broadly oval, solid, slightly cupped at distal end and

usually greatly eroded. Posterior end of animal with 2 long fleshy lappets on dorsal surface. *Rec.*, N.S. Am.-W.Afr.—FIG. E201,1; E202,1. **N. reynei* (BARTSCH), Surinam; E201,1, entire animal (61 cm. long); E202,1a,b, RV ext., int. (apophysis lost); E202,1c-e, pallet, outer and inner faces, side view (holotype) (923).

Psiloteredo BARTSCH, 1922, p. 36 [**Teredo dilatata* STIMPSON, 1851, p. 113 (= *T. megotara* HANLEY, 1848, p. 77); OD] [= *Dactyloteredo* MOLL, 1941, p. 193 (type, *Teredo megotara* HANLEY, 1848, p. 77; SD MOLL, 1952, p. 83)]. Pallets broad to elongate oval in outline, solid, almost entirely calcareous, and with short stalk. Blade thick at base, becoming thin at distal margin, slightly concave on inner face, convex on outer; outer face variable, usually with moderate to deep thumb-nail-like depression, paddle-like, or with slight depression and 2 finger-like projections. Valves with prominent condyles; the posterior slope moderate to large and usually flaring. Tubes probably concamerated at posterior end. Siphons united for most of their length. Gills extending from base of siphons to visceral mass; intestine making simple loop around the caecum. *Rec.*, cosmop. in temp. and trop. seas.—FIG. E162,2. *P. healdi* (BARTSCH), Surinam; entire animal showing major features of anatomy (923).—FIG. E203,2a-d. **P. megotara* (HANLEY), Brit. Is. (lectotype); 2a,b, LV ext., RV int., 2c,d, outer and inner face of pallet (923).—FIG. E203,2e-h. *P. megotara excisa* (JEFFREYS), Brit. Is. (lectotype) [=deformed *P. megotara*]; 2e,f, LV ext., LV int. showing typical large dorsal condyle but reduced deformed posterior slope; 2g,h, outer and inner face of pallet (923).




FIG. E193. Pholadidae (Jouannetiinae) (p. N720).




FIG. E194. Pholadidae (Jouannetiinae) (p. N720).

Teredora BARTSCH, 1921, p. 26 [**Teredo malleolus* TURTON, 1822, p. 255; OD] [= *Malleolus* GRAY, 1847, p. 188 (obj.) (*non* RAFINESQUE, 1815; *nec* EHRENBERG, 1838)]. Pallets narrowly to broadly oval in outline with short stalk; blade thin, convex on outer, concave on inner face, outer face with small to large, usually deep, thumbnail-like depression which is marked with broadly curved, concentric growth lines. Thickened area at base of pallet smooth. Stalk extending only to base of

depression. Distal end of tube concaerated. Shell usually with small, high posterior slope. Siphons united. Gill lamellae bladlike, extending length of animal from base of siphons to mouth. Heart anterior, with short broad ventricle and small auricles. Stomach transitional, caecum doubling on itself. Intestine short, making only simple loop around caecum. *Rec.*, cosmop.—FIG. E204, 1, 2. **T. malleolus* (TURTON), Eng.; 1a, b, LV ext., RV int.; 1c-e, pallet, outer and inner face, side view; 1f, g, pallet, outer and inner face of lectotype (malformed); 2, semidiagrammatic view of entire animal showing anatomical characters of genus (from SÉNÉGAL) (923).

Teredothyra BARTSCH, 1921, p. 26 [**Teredo (Teredothyra) dominicensis* BARTSCH, 1921, p. 26, 30; OD] [= *Ungoteredo* BARTSCH, 1927, p. 544 (type, *Teredo (Ungoteredo) matacotana* BARTSCH, 1927, p. 544; OD); *T. (Idioteredo) TAKI & HABE*, 1945, p. 115 (type, *Teredo (Teredothyra) smithii* BARTSCH, 1927, p. 540)]. Blade of pallets composed of 2 elements, basal portion forming broad to elongate cup with secondary divided conelike element within it; stalk of pallet, which is sheathed by basal cone, extending into blade only as far as base of inner cone. Siphons long and separated. Gill long, lamellae thin and bladlike; heart anterior, extending beneath posterior adductor muscle. Digestive system with globular stomach, small caecum which folds on itself, and short intestine. *Rec.*, warm temp. and trop seas.—FIG. E205, 1, 2. **T. dominicensis* (BARTSCH), W. Indies; 1a, b, LV ext., int.; 1c, distal end of tube showing division and thickening; 1d, f, outer faces of pallets; 1e, g, inner faces of pallets; 1h, side view of pallet (1a-h, holotype of *T. atwoodi* BARTSCH, = *T. dominicensis*, adult); 2a, b, RV ext., int.; 2c, distal end of tube; 2d-f, pallet, outer and inner face, side




FIG. E195. Pholadidae (Xylophaginae) (p. N721-N722).




FIG. E196. Diagrammatic representation of relationships of living genera of Teredinidae (923).

view (2a-f, *T. dominicensis*, holotype, young specimens) (923).

Uperotus GUETTARD, 1770, p. 126 [**Teredo clava* GMELIN, 1791, p. 3748; SD LAMY, 1927, p. 275] [= *Fistulana* LAMARCK, 1799, p. 90 (obj.) (non MÜLLER, 1776, nec BRUGUIÈRE, 1789; *Fistulanigenus* RENIER, 1807, pl. 7 (obj.) (ICZN, work rejected); *Guetera* GRAY, 1840, p. 42 (nom. nud.), 1847, p. 188 (type, *Fistulana corniformis* LAMARCK, 1799, = *T. clava* GMELIN, 1791; OD); *Hyperotus* HERRMANNSEN, 1847, p. 671 (nom. van.); *Hyperotis* PAETEL, 1875, p. 99 (nom. null.); *Guetera* H. ADAMS & A. ADAMS, 1856, p. 333 (nom. null.); *Glumebra* IREDALE, 1936, p. 42 (obj.) (type, *G. elegans*, = *T. clava* GMELIN; OD)]. Pallets oval to rectangular in outline with short heavy stalk, basal portion of blade nearly smooth, distal portion with pronounced radiating ribs. Shells with anterior and posterior slopes greatly reduced or typical in shape, with small, high posterior slope. [Species in this genus are usually obtained from dredged nuts or wood or from drift cast ashore. They are probably not an intertidal group. The anatomy of this species in this genus is very close to that of *Teredora*, gill extending from base of siphons to mouth (Fig. E204,2).] *Rec.*, IndoPac-

E.Pac.-E.Atl.—FIG. E206,1. **U. clavus* (GME-LIN), Australia (Queensl.) (from a nut); 1a,b, RV (holotype of *Glumebra elegans* IREDALE), ext., int., $\times 5$; 1c, cluster of tubes, $\times 0.9$; 1d-f, pallet, outer and inner face, side view, $\times 3$ (435).—FIG. E206, 2a-g. *U. lieberkindi* ROCH (from dredged wood) off Rio de Oro, Afr.; 2a,b, RV (holotype) ext., int.; 2c-g, pallet, outer and inner face, side view (923).

Zachia BULATOV & RZHAVSHCHIKOV. [RJAFTSCHIKOFF], 1933, p. 166 [**Z. zenkewitschi*; SD HABE, 1952, p. 255]. Shells greatly reduced; pallets similar to *Spathoteredo*. [Species living in roots of *Phylospadix*, apparently stenomorphic and malformed.] *Rec.*, N.Pac.—FIG. E207,1. **Z. zenkewitschi*, USSR (Vladivostok); 1a, ant. view of opposed valves, $\times 20$; 1b, RV ext., $\times 20$; 1c,d, outer and inner face of pallet, $\times 40$ (1a-d, holotype) (90).

Subfamily BANKIINAE Turner, 1966

Wood-borers with pallets composed of numerous segments which may be very closely packed and indistinct (e.g., *Nototeredo*, *Spathoteredo*), distinct but fused




FIG. E197. Teredinidae (Teredininae) (p. N725-N726).




FIG. E198. Teredinidae (Teredininae) (p. N726).

(e.g., *Nausitora*), or widely spaced and conelike (e.g., *Bankia*). Blade of pallet increased in size by addition of new segments at distal end. Valves similar to those of Teredininae. *Paleoc.-Rec.*

Bankia GRAY, 1842, p. 76 [**Teredo bipalmulata* LAMARCK, 1801, p. 129; SD GRAY, 1847, p. 188] [= *Xylotrya* LEACH in GRAY, 1847, p. 188 (non MENKE, 1830; nec GRAY, 1842) (type, *Teredo bipalmulata* LAMARCK, 1801; OD); *Xylotrya* DE QUATREFAGES, 1849, p. 28 (nom. null.); *Xylotria* DESHAYES, 1860, p. 114 (nom. null.); *Xylotya* VON MARTENS, 1880, p. 331 (nom. null.); *Hylotrya* CLESSIN, 1893, p. 82 (nom. null.)]. Pallets elongate, composed of numerous conelike elements built upon central stalk which extends length of blade, cones separate and easily removed from stalk, base of cones calcareous, with periostracal covering extended as border which varies greatly in width, may be smooth or fringed, and may be produced laterally as awns. [The genus has been divided into numerous subgenera on the basis of ornamentation of the periostracal border and the presence or absence of awns, but these taxa prob-




FIG. E199. Teredinidae (Teredininae) (p. N726-N727).

ably have small value owing to the presence of intergrades. However the subgenera are listed and a figure of the type species given.] The periostracum is not preserved in fossils, so usually the best one can do is refer to a fossil as *Bankia* sp. *Paleoc.-Rec.*, cosmop.

B. (*Bankia*) GRAY, 1842 [= *Bankiura* MOLL, 1952, p. 8, 42, 85 (obj.)]. Margin serrated on inner face, smooth on outer face and produced in unequal lateral awns, those on one side being 2 to 3 times length of other. ?*Paleoc.*, *Rec.*, IndoPac.

—FIG. E208,1. **B. (B.) bipalmulata* (LAMARCK), New Hebrides; 1a,b, pallet, outer and inner faces (920).

B. (*Bankiella*) BARTSCH, 1921, p. 25-26 [**Bankia (Bankiella) mexicana* (= *B. gouldi* BARTSCH, 1908); OD]. Margins of cones smooth, lateral awns long and thin. Margin on inner face wide, forming "web" between awns. *Rec.*, cosmop.—FIG. E208,2. **B. (B.) gouldi* BARTSCH, USA (Va.); 2a,b, pallet, outer and inner faces (920).

B. (*Clupibankia*) MOLL, 1952, p. 42, 85 [**Bankia*




FIG. E200. Teredinidae (Teredininae) (p. N727-N728).




FIG. E201. Teredinidae (Teredininae) (p. N728).

barthelowi BARTSCH, 1927, p. 537; OD]. Calcareous portion of cones wide on inner face so that cones overlap one another; margin serrate on outer face, smooth on inner face. *Rec.*, IndoPac.—FIG. E208,4. **B. (C.) barthelowi* BARTSCH, Philip. Is.; outer face of pallet (923).

B. (Liliobankia) CLENCH & TURNER, 1946, p. 17 [**B. (L.) katherinae* (= *B. campanellata* MOLL & ROCH, 1931); OD]. Margin of cones wide, smooth and produced laterally to form broad triangular awns. Embryonic cones separated. *Rec.*, subtrop. and trop. seas.—FIG. E208,5. **B. (L.) campanellata* MOLL & ROCH; 5a,b, pallet, outer and inner faces (920).

B. (Lyrodobankia) MOLL, 1941, p. 200 [**Nausitora kamiyai* ROCH, 1929, p. 17 (= young *B. carinata* GRAY, 1827); SD TURNER, 1966] [= *Bankiopsis* CLENCH & TURNER, 1946 (type, *B. (B.) caribbea* (= *B. carinata* GRAY) (obj.);

OD)]. Margin smooth, equal width on both inner and outer faces, and produced laterally into short, blunt points. Embryonic cones closely packed and covered by periostracal cap. *Rec.*, temp. and trop. seas.—FIG. E208,3. **B. (L.) carinata* (GRAY), USA (Fla.) (3a,b), Japan (3c,d); 3a,b, pallet, outer and inner face (holotype of *B. (Bankiopsis) caribbea* CLENCH & TURNER (920); 3c,d, pallet, outer and inner face (holotype of *Nausitora kamiyai* ROCH, = young *B. carinata* GRAY) (923).

B. (Neobankia) BARTSCH, 1921, p. 26 [**B. (N.) zeteki*; OD] [= *Deviobankia* IREDALE, 1932, p. 33 (type, *Bankia debenhami* IREDALE, 1932; OD)]. Margin of cones wide, even, serrated on both inner and outer faces and not produced laterally to form awns. *Rec.*, warm temp. and trop. seas.—FIG. E208,6. **B. (N.) zeteki*, Panama; 6a,b, pallet, outer and inner faces (920).

B. (Plumulella) CLENCH & TURNER, 1946, p. 22 [**B. fimbriatula* MOLL & ROCH, 1931; OD]. Margin of cones serrated on both inner and outer faces and produced laterally into long serrated awns. *Rec.*, temp. and trop. seas.—FIG. E208,7. **B. (P.) fimbriatula* MOLL & ROCH, Haiti; 7a,b, pallet, outer and inner faces (920).

Nausitora WRIGHT, 1864, p. 452 [**N. dunlopei*; OD] [= *Nausitoria* TRYON, 1868, p. 20 (nom. null.); *Nausitora* SOWERBY, 1887, pl. 469, fig. 3 (nom. null.); *Nausitora* ROCH & MOLL, 1929,




FIG. E202. Teredinidae (Teredininae) (p. N728).




FIG. E203. Tereidinidae (Teredininae) (p. N728, N737).

p. 17 (*nom. null.*); *Inequarista* IREDALE, 1932, p. 37 (type, *Nausitora messeli* IREDALE, = *N. dunlopei* WRIGHT; OD); *Nausitorea* MOLL, 1952, p. 84 (type, *T. fusticula* JEFFREYS, 1860, p. 125;




FIG. E204. Teredinidae (Teredininae) (p. N729).




FIG. E205. Teredinidae (Teredininae) (p. N729-N730).

OD)]. Pallets elongate, composed of closely packed, fused, conelike elements built on central stalk; periostracal covering commonly extended as awns, particularly on proximal portion of blade, distal portion of blade with asymmetrical, papillose, calcareous covering which is usually worn off in old specimens. Siphons united for about three-fourths of their length. Gills short, broad and flat; heart small and posterior, located at anterior end of

gills posterior to visceral mass. Stomach elongate; caecum very large, intestine making simple loop around it. *Rec.*, trop. and subtrop. seas, usually in brackish water.—FIG. E209, 1. **N. dunlopei*, Australia; 1a,b, LV ext., int.; 1c,d, pallet, outer and inner faces (923).

Nototeredo BARTSCH, 1923, p. 100 [**Teredo* (*Nototeredo*) *edax* HEDLEY, 1895, p. 501; OD] [= *Phyloterredo* ROCH, 1937, p. 169 (type, *Teredo norvegica* SPENGLER, 1792, p. 102; OD)]. Pallets elongate oval to broadly oval in outline with short stalk; blade thin, convex on outer and concave on inner face; composed of soft, friable calcareous material laid down in closely packed segments separated by thin layers of periostracum which may extend laterally as small awns, entire surface of blade with thin layer of periostracum. Stalk extending length of blade. Siphons united for about





FIG. E207. Teredinidae (Teredininae) (p. N730).



Uperotus




FIG. E206. Teredinidae (Teredininae) (p. N730).

half their length. Gills about 0.4 length of animal, lamellae broadly U-shaped. Labial palps free at tips. Stomach globular, caecum large, and intestine extremely long and convoluted. Heart short, located at anterior end of gills, posterior to visceral mass; ventricle short and broad, auricles short; posterior aorta long, extending from ventricle to posterior adductor muscle. *Paleoc.-Rec.*, worldwide.—FIG. E203, 1. **N. edax* (HEDLEY), *Rec.* S.Australia; 1a,b, LV ext., int. (paratype); 1c,d, pallet, outer and inner face; 1e, post. end of young specimen (923).

Spathoterredo MOLL, 1928, p. 282 [**Teredo bataviana* MOLL & ROCH, 1931, p. 207; (= *T. obtusa* SIVICKIS, 1928, p. 290); SD TURNER, 1966, p. 122] [= *Spathoterredo* ROCH, 1937, p. 173 (obj.)]. Blade of pallets more or less rectangular in outline, truncated at anterior end and sheathing stalk for about 0.3 of its length. Stalk extending well up into blade which is thick at anterior end and tapers posteriorly. Periostracum dark brown, usually appearing as band across middle of blade on outer face, distal portion of outer face covered with pustulose, calcareous incrustation. Siphons with muscular collar at their base, united for most of their length, incurrent siphon with rather elaborate branched papillae. Gills long, narrowly U-shaped. Heart median, ventricle broad, auricles long and tapering. Stomach elongate, caecum large, gonads dorsal and posterior to it, intestine making loop around it. *Rec.*, IndoPac.-S.Atl.—FIG. E210, 1. **S. obtusa* (SIVICKIS), Java; 1a,b, LV ext., int.; 1c-e, pallet, outer and inner face, side view (923).




FIG. E208. Teredinidae (Bankiinae) (p. N732-N734).

Subfamily KUPHINAE Tryon, 1862

[Kuphinae TRYON, 1862, p. 455]

Large mud- and possibly wood-borers having greatly reduced shells and strong muscular collar surrounding body at posterior end of shell; anterior and posterior adductor muscles nearly equal in size; condyles, chondrophore and ligament reduced, and posterior end of shell covered by muscular collar. Digestive tract greatly reduced, esophagus extending posterior to muscular collar, crystalline style and stomach located at posterior end of visceral mass;




FIG. E209. Teredinidae (Bankiinae) (p. N734, N736).




FIG. E210. Teredinidae (Bankiinae) (p. N737).

caecum lacking. [Unlike all other Teredinidae, the heart has a large ventricular bulb. The intestine passes through this bulb and around the posterior adductor muscle; it becomes imbedded in the thickened mantle for the length of the pericardial cavity and then opens into the epibranchial chamber behind the heart. The Kuphinae apparently are all filter-feeders.] ?*Eoc., Rec.*

Kuphus GUETTARD, 1770, p. 139 [**Serpula polythalamia* LINNÉ, 1758; SD GRAY, 1847, p. 188] [= *Furcella* LAMARCK, 1801, p. 104 (obj.); *Septaria* LAMARCK, 1818, p. 436 (non FÉRUSSAC, 1807) (obj.); *Cloissonaria* FÉRUSSAC, 1822, p. xiv (referring to cloissonnaire and *Septaria* LAMARCK); *Clausaria* MENKE, 1828, p. 73; *Cloissonaria* ADAMS, 1858, p. 648 (nom. van.); *Cloissonaria* PAETEL, 1875, p. 46 (nom. null.); *Cloissonaria* PAETEL, 1890, p. 6 (nom. van.); *Cuphus* AGASSIZ, 1846, p. 108 (nom. van.); *Cyphus* FISCHER, 1887,




FIG. E211. Teredinidae (Kuphinae); 1, *Kuphus*, entire animal, $\times 0.5$; 2, same, morphology of soft parts (923).

p. 1158 (*nom. van.*); *Kyphus* HERRMANNSEN, 1847, p. 569 (*nom. van.*). Pallets simple, solid, almost entirely calcareous; blade triangular in outline with long heavy stalk, flat on inner face and with shallow cup. Shells relatively small, denticulated ridges of anterior slope greatly reduced; posterior slope reduced; tube very thick, solid, divided posteriorly. [The morphology of the soft parts is shown in Figure E211,2. Though the genus has been reported from the middle Eocene, this is on the basis of the tubes only. The pallets and shells of *Kuphus* have never been found as fossils.] ?*Eoc.*, *Rec.*, IndoPac.—FIG. E211,1,2. **K. polythalamia* (LINNÉ), *Rec.*, Solomon Is.; 1a, sketch of entire animal 50.5 cm. long, $\times 0.5$; 2a, semidiagrammatic long. sec. showing relationship of organs; 2b, ant. portion enl. (923).

Subfamily UNCERTAIN

Genera based on shells and tubes only cannot be classified as to subfamily within the Teredinidae.

Eoteredo BARTSCH, 1923, p. 98 [**E. philippinensis*; OD]. Name based on shells only, distinguished by projection of apophyses from beneath shelf


rather than umbos. [This is an age factor which may be found in the shells of several genera.] *Rec.*, Philip. Is.—FIG. E212,1. **E. philippinensis*, Mindanao; 1a,b, LV ext., RV int. (apophysis broken) (holotype), $\times 6$ (923).

Terebrimya STEPHENSON, 1952, p. 141 [**T. lamarana*; OD]. Shells typical of family; pallets unknown. *U.Cret.*(*Cenoman.*), USA (Tex.).—FIG. E213,1. **T. lamarana*, Woodbine F.(Templeton Mbr.), Lamar Co.; 1a,b, LV ext., RV ext., in matrix (holotype), $\times 4$; 1c, tube, $\times 1$ (890).


Teredinopsis FUCH, 1878, p. 39 [**T. problematica*; OD]. Based on shells and cluster of tubes similar to *Uperotus clavus*; pallets unknown. *Quat.*, Suez.—FIG. E214,1. **T. problematica*; 1a,b, LV ext., int., $\times 3$; 1c, anterior view of opposed valves; 1d, cluster of tubes; all $\times 3$ (1a-c, holotype) (332).

Teredolites LEYMERIE, 1842, p. 2 [**T. clavatus*; OD] [= *Teredolites* LEYMERIE, 1841, p. 341 (*nom. nud.*); *Teredolithes* HERRMANNSEN, 1852, p. 131 (*nom. null.*)]. Name based on tubes only. [Close to *Uperotus clavus* GMELIN.] *Cret.*, Eu.—FIG. E214,2. **T. clavatus*, France (Spatangues, Dept. l'Aube); 2a,b, basal and lat. views of cluster of tubes (holotype), $\times 1$ (540).

Teredolithus BARTSCH, 1930, p. 460. [Collective



1a



Eoteredo


1b

[=*Polarthus* STOLICZKA, 1871, p. 14 (*nom. null.*); *Kummelia* STEPHENSON, 1936, p. 60 (*obj.*)].
Belongs to Gastrochaenidae.


Order HIPPURITOIDA Newell,
1965

[Diagnosis by N. D. NEWELL]

Rudists and pachydonts; thick-shelled, aberrant heterodonts, primitively equivalved, with few thick and amorphous hinge teeth; mainly attached and strongly inequi-



1a



1b

1c

Terebrimya

FIG. E212. Teredinidae (Subfamily Uncertain)
(p. N740).

group name for species of Teredinidae that cannot be placed generically.]

Turnus GABB, 1864, p. 145 [**T. plenus*; OD]. Shell thin, pholadiform; internally bearing rib which rises in apex, and passes downward and backward to basal margin in same manner as umbonal groove and posterior to it. Accessory plates unknown. Tube simple, thin. [This genus was placed in the Teredininae by GABB and considered to be a link with the Pholadidae. Until more material is available the genus cannot be positioned.] *Cret.*, N.Am.—FIG. E213,2. **T. plenus*, USA (Calif.); (holotype), $\times 2$ (333).

INVALID GENERA

The following names for hypothetical genera are invalid (ICZN Code, 1961, Art. 1).


Bicornia MAY, 1929, p. 642, 665.

Microvexillum MAY, 1929, p. 642, 665.

Proteredo MAY, 1929, p. 664, 665.

GENERA ERRONEOUSLY ASSIGNED
TO TEREDINIDAE

Polorthus GABB, 1861, p. 366 [**P. americana*; OD]



2

Turnus

FIG. E213. Teredinidae (Subfamily Uncertain)
(p. N740-N741).




FIG. E214. Teredinidae (Subfamily Uncertain) (p. N740).

valved, tending to gyrate or conical, operculate and edentulous forms resembling solitary corals. *M.Sil.-U.Cret.(Maastricht.)*.

The Hippuritoida are cosmopolitan bivalves especially characteristic of low latitudes where they dominate extensive reefs.

Superfamily MEGALODONTACEA Morris & Lycett, 1853

[Diagnosis by N. D. NEWELL]

Characters of family Megalodontidae. *M.Sil.-L.Cret.*

Family MEGALODONTIDAE Morris & Lycett, 1853

[*nom. correct.* ZITTEL, 1881 (*pro* Megalodontidae MORRIS & LYCETT)] [Materials for this family prepared by L. R. COX (Mesozoic) and AURELE LAROCQUE (Paleozoic)]

Shell mostly medium-sized to large, thick, equivalve except in some species of *Megalodon* and in *Pomarangina*, gibbose, subtrigonal or ovate, usually with prosogyrous

beaks; hinge plate massive, bearing in each valve one to several cardinal teeth which are commonly rather amorphous and not all of which diverge from beak; lateral teeth present in some genera; ligament external, opisthodontic, nymphs heavy; pallial line entire; anterior adductor scar deep, situated in relatively dorsal and marginal position, just below anterior end of hinge plate; surface usually smooth or at most with weak concentric folds, but with radial ribs in a few forms. *M.Sil.-L.Cret.*

Megalodon J. DE C. SOWERBY, 1827, p. 131 [**M. cucullatus*; M] [= *Megalodon* LINK, 1830 (*nom. null.*); *Megalodonta* GOLDFUSS, 1832 (*nom. van.*); *Megalodus* GOLDFUSS, 1837 (*nom. van.*) (*non* RAFINESQUE, 1815); *Eumegalodon* GUEMBEL, 1862 (*obj.*)]. Shell medium-sized to large, subtrigonal or ovate, gibbose; beaks anterior, strongly prosogyrous; hinge plate very massive; cardinal teeth varying in number, from 1 or 2 in LV and 1 to 3 in RV, owing to suppression or duplication of elements of dentition; posterior adductor scar on slightly raised platform; surface smooth or con-

centrically rugose (Dechaseaux in Piveteau, 1952, p. 293, 333). *Dev.-U.Trias.(Rhaet.)*.

M. (Megalodon). Equivalve, valves more or less carinate posteriorly; LV with strong, elongate, acutely triangular, commonly grooved posterior cardinal tooth close to posterodorsal margin and tuberculiform anterior cardinal tooth (represented by 2 unequal teeth in some specimens) near anterior end of hinge plate; RV with strong, subtriangular, commonly pustulose or transversely grooved, mesially placed cardinal tooth received in deep, broad recess between 2 main teeth of LV, and in some specimens with very thin, elongate posterior cardinal; traces of elongate lateral tooth adjacent to posterior margin beyond hinge plate seen in some RV's. *Dev.-Trias., cosmop.*—FIG. E215,4. **M. (M.) cucullatus* SOWERBY, *Dev.*, Paffrath, Ger.; LV int., showing dentition, $\times 1$ (379).

M. (Neomegalodon) GUEMBEL, 1862, p. 362 [**Cardium triquetrum* WULFEN, 1793, p. 48; SD STOLICZKA, 1871, p. 275] [= *Tauroceras* SCHAFHAEUTL, 1854 (*non* HOPE, 1840); *Neomegalodus* FRECH, 1904 (*nom. van.*)]. Equivalve or slightly inequivalve, presence of posterior carina variable; hinge plate narrowing posteriorly; LV without distinct posterior cardinal tooth but with 1 or 2 anterior teeth representing tuberculiform tooth of *M. (Megalodon)*; when 2 such teeth are present these are commonly subequal and radially elongated; RV with variable, acutely triangular posterior cardinal tooth received in recess behind tooth or teeth of LV and with 2 anterior teeth which are commonly subequal and radially elongated. *U.Trias., cosmop.*—FIG. E216,1. **M. (Neomegalodon) triqueter* (WULFEN), *U.Trias. (Rhaet.)*, Aus.; 1a,b, RV int., LV int., showing dentition, $\times 1$ (Guembel, 1862).

Conchodon STOPPANI, 1865, p. 246 [**C. infraliasicus* (= *Lycodus cor* SCHAFHAEUTL, 1863, p. 375); M] [= *Lycodus* SCHAFHAEUTL, 1863 (*non* QUENSTEDT, 1856); *Conchodus* TAUSCH, 1891 (*nom. van.*) (*non* M'COY, 1848); *Lycodes* DALL, 1900 (*nom. null.*)]. Very large, thick-shelled, about as high as long, with strongly prosogyrous and incoiled beaks and projecting anterior extremity; valves sharply carinate posteriorly; hinge plate massive; RV with massive arcuate or irregularly tripartite main tooth which is situated near anterodorsal margin and is received in broad socket in LV between arms of horseshoe-shaped tooth; in some specimens second and weaker tooth lies between this main tooth and margin in RV; posterior part of hinge plate without distinct teeth; no lateral teeth; anterior adductor scar almost marginal, just below main tooth; posterior scar not observed; surface of shell with weak concentric folds. *U.Trias.(Rhaet.)*, Eu.(N.Alps-S. Alps-Hung.-Carpathians).—FIG. E216,3. **C. cor* (SCHAFHAEUTL), S.Alps; 3a, int. mold, $\times 0.3$




FIG. E215. Megalodontidae (p. N742-N743, N745).

(Stoppani, 1865); 3b, ant. part of RV hinge showing tripartite main tooth, $\times 0.3$ (Frech, 1904).

Cumularia SPRIESTERSBACH, 1919, p. 467 [**C. circularis*; M]. Small, round-oval, not carinate; umbones depressed, well anterior; 1 stoutly triangular cardinal tooth in each valve, that in RV stronger and fitting behind tooth of LV; no laterals; anterior adductor scar small and deep, posterior scar larger, rounded. *M.Dev.*, Eu.(Ger.).—FIG. E215,5. **C. circularis*, ?Honseler beds, Schaf-




FIG. E216. Megalodontidae (p. N743, N745).

feld; 5a, LV int. mold, $\times 1$; 5b, RV dentition, enl. (872).

Eomegalodus SPRIESTERSBACH, 1915, p. 53 [**E. fuchsi*; M] [= *Eomegalodon* COSSMANN, 1922 (*nom. van.*)]. Medium-sized, thick, elliptical, not carinate; umbones anterior, not prominent; lunule small; hinge very thick; LV with 2 cardinal teeth, posterior one heavy, and broad, cuneiform posterior lateral; RV with 1 strong and 1 weak cardinal and lamellar posterior lateral; lower margin of hinge plate of RV thickened into large toothlike pad which projects into LV; anterior adductor scar small, elliptical, deep, striated; posterior scar large, deep; sculpture of fine concentric lines. *M.Dev.*, Eu.(Ger.).—FIG. E215,3. **E. fuchsi*, U.Honseler beds, Lenneschiefer, Verneis, near Haspe; RV int., showing dentition and toothlike pad below hinge, $\times 1$ (871).

Ferrazia REED, 1932, p. 480 [**F. cardinalis*; M]. Equivalve; beaks subanterior, strongly prosogyre, incurved; lunule and escutcheon well defined; surface angulated and ornamented with strong radial carinae; each valve with large rounded, longitudinally oval cardinal tooth on narrow thickened hinge plate; ligament opisthodontic, parivincular. *Perm.*, S.Am.(S.Brazil).—FIG. E216,1. **F. cardinalis*, Corumbatai F., Rio Claro; 1a,b, RV ext. and hinge views, $\times 1$ (Reed, 1932). [NEWELL]

Krumbeckia DIENER, 1915, p. 131 [**K. tambangensis* (= *Pomarangina* aff. *haydeni* DIENER, KRUMBECK, 1914, p. 256); M]. Medium-sized, oval, gibbose, not carinate; beaks prosogyrous, at anterior 5th of length; hinge plate heavy; RV with single heavy, broadly arched tooth, almost median in position; teeth of LV and muscle scars unknown; ornament of concentric ribs. *U.Trias.*, Indon.(Sumatra).—FIG. E216,2. **K. tambangensis*; 2a,b, ant. view of shell and RV int. showing dentition, $\times 1$ (Krumbeck, 1914).

Megalomoidea COX, 1964, p. 43 [**Megalomus canadensis* HALL, 1852, p. 343; OD] [= *Megalomus* HALL, 1852 (*non* RAMBUR, 1842); *Megalomys* FISCHER, 1886 (*nom. van.*; *non* TROUËSSART, 1881); *Megalomus* KIRK, 1927 (*nom. null.*)]. Large, reniform-ovate, thick-shelled, especially in hinge region and along V-shaped area extending posteriorly from it across middle of each valve; valves not carinate; beaks anterior, prosogyrous; hinge plate massive, not undercut, broad and projecting in middle, where in each valve it bears group of transversely elongated teeth variable in number (2-6) and strength; anterior adductor scar as in typical Megalodontidae, 2 small accessory scars on its posterodorsal side; posterior adductor scar obscure; surface smooth except for growth lines (Kirk, 1927, p. 4). *M.Sil.(Guelph)*, E.N.Am.-Alaska.—FIG. E215,2. **M. canadensis* (HALL), Ont.(Elora); RV int., $\times 0.7$ (Kirk, 1927).

Pachyrisma MORRIS & LYCETT, 1850, p. 399 [**P. grande*; M] [= *Cardilla* LYCETT, 1848 (*nom. nud.*, or error for *Cardilla* DESHAYES, 1835); *Pachyerisma* BAYAN, 1874 (*nom. van.*)]. Medium-sized to large, equivalve, thick-shelled, ovate to subtrigonal, with strongly prosogyrous, more or less anteriorly placed beaks; valves carinate posteriorly; hinge plate, ligamental nymph and teeth heavy; single strong, mesially placed cardinal in LV received in recess between strong posterior cardinal and weak (in some specimens ill-defined) anterior cardinal in RV; well-developed, tuberculiform lateral tooth present near each end of hinge plate in LV and received in sockets above corresponding laterals in RV; anterior adductor scar deep, situated just below anterior lateral tooth. *L.Jur.-U.Jur.*, Eu.

P. (Pachyrisma). Large, higher than long, beaks terminal in some but not all species; posterior adductor scar on lamina emerging from beneath hinge plate; surface smooth or with traces of radial ribbing. *M.Jur.-U.Jur.*, Eu.—FIG. E217, 2a-c. **P. (P.) grande* MORRIS & LYCETT, M.Jur. (Bathon.), Eng.; 2a, RV ext.; 2b, LV hinge, post. lat. tooth broken away; 2c, RV hinge, post. lat. tooth broken away; all $\times 0.7$ (645).—FIG. E217,2d. *P. sp. cf. P. (P.) beaumonti* ZEUSCHNER, U.Jur.(Tithon.), Czech.; RV hinge showing lat. teeth, $\times 0.5$ (Böhm, 1870).

P. (Durga) G. BÖHM, 1884, p. 191 [**D. nicolisi*; OD]. Medium-sized, trigonally ovate, longer than high, tapering posteriorly; posterior carina sharp; posterior adductor scar not observed; surface smooth, except for growth rugae. *L.Jur.(L.Lias.)*, Eu.(N.Italy-France).—FIG. E217,3. **P. (D.) nicolisi*, N.Italy; 3a, RV ext., $\times 0.7$; 3b, RV hinge, defective posteriorly, $\times 1$ (Böhm, 1884).

P. (Pachymegalodon) GUEMBEL, 1862, p. 375 [**Bucardites chamaeformis* VON SCHLOTHEIM, 1820, p. 208; M] [= *Pachymegalodus* TAUSCH, 1890 (*nom. van.*)]. Medium-sized, higher than long, acutely subtrigonal; posterior carina sharp, with broad radial sulcus in front of it; posterior adductor scar on lamina, as in *P. (Pachyrisma)*; surface smooth, except for growth rugae. *L.Jur.(L.Lias.)*, Eu.(S.Alps).—FIG. E217,1. **P. (P.) chamaeforme* VON SCHLOTHEIM, Yugosl.; 1a,b, RV ext., RV int., $\times 1$ (Tausch, 1890).

Pachyrismella COX, 1964, p. 43 [**Cardium septiferum* BUVIGNIER, 1843, p. 230; OD]. Medium-sized, subequilateral, slightly longer than high, *Cardium*-like; beaks not strongly prosogyrous; hinge plate and teeth less heavy and irregular than in *Pachyrisma*; strong acutely angular posterior cardinal and weak or scarcely developed anterior cardinal present in both valves, main tooth of RV fitting in front of that of LV; lateral teeth and adductor scars as in *Pachyrisma*, posterior adductor on lamina emerging from be-

low hinge plate; radial ornament absent or obscure. *U. Jur. (U. Oxford.)*, Eu.-SW. Asia (Lebanon).
 —FIG. E218, 2. **P. septifera* (BUVIGNIER), U.

Oxford., France; *2a, b*, LV hinge, RV ext.; *2c*, RV int., all $\times 0.7$ (Buvignier, 1852).
Paramegalodus COX, new genus [*ex* KUTASSY, 1934,




FIG. E217. Megalodontidae (p. N745).

p. 52 (*nom. nud.*, type species cited but no generic diagnosis) [**Dicerocardium eupalliatum* FRECH, 1904, p. 51; OD]. Large, tall, trigonal, hornlike, with very prominent, terminal, pointed,

prosogyrous umbones; internal mold with rounded rib running from near anterior margin almost to umbo, then curving around and descending to posterior end of ventral margin; dentition un-




FIG. E218. Megalodontidae (p. N745-N748).

known. *U.Trias.* (*Rhaet.*), Eu. (Hung.)-Asia (Himalayas).—FIG. E218, 3a. **P. eupalliatu*s (FRECH), Hung.; ant. view of int. mold, $\times 0.3$.—FIG. E218, 3b, c. *P. incisus* (FRECH), Hung.; 3b, post. view, 3c, RV int. mold, $\times 0.7$ (all Vigh, 1914).

Pinzonella REED, 1932, p. 482 [**P. illusa*; SD Cox, herein] [= *Pinzonellopsis* MENDES, 1944, p. 58 (type, *Pachycardia occidentalis* REED, 1929; M)]. Small for family, orbicular or ovate, strongly inequilateral, beaks fairly anterior, not strongly incurved; valves weakly carinate; hinge plate relatively broad and massive, with large, rounded-triangular cardinal tooth in each valve, that of RV fitting in front of LV tooth; weak lamellar lateral tooth close to posterodorsal margin distinguishable in LV of some specimens; surface smooth. *Perm.* (originally considered *Trias.*), *Corumbatai beds*, S. Am. (Brazil).—FIG. E218, 1. **P. illusa*, São Paulo State; 1a-d, LV ext. and int., RV ext. and int., all $\times 1.3$ (Mendes, 1952).

?*Pomarangina* DIENER, 1908, p. 64 [**P. haydeni*; M]. Ovate, inequilateral, moderately gibbose, thick-shelled, some specimens slightly inequivalve, with RV larger; beaks prosogyrous and incurved; valves carinate posteriorly; dentition and musculature unknown. *U.Trias.*, Asia (Himalayas). *Protodicer*s G. BÖHM, 1891, p. 19 [**Megalodon pumilus* GUEMBEL, 1862, p. 367 (as *M. triqueter* var. *pumilus*); OD] [= *Protodicer*s BERINGER, 1949 (*nom. null.*)]. Of small-medium size, sub-trigonal, about as long as high, with strongly prosogyrous, terminal beaks; valves sharply carinate posteriorly; hinge plate and teeth heavy; LV with weak, elongate, marginal posterior cardinal tooth with its apex close to beak and horseshoe-shaped anterior cardinal which is well separated from beak; between them is large, deep, triangular socket for reception of strong RV posterior cardinal, and within 2 arms of anterior tooth is socket for simple RV anterior cardinal; RV an-




FIG. E219. Megalodontidae (p. N748-N749).

Hippuritoida—Hippuritacea—Introduction

terior lateral is feebly developed but posterior laterals are absent; anterior adductor scar deep, located as in *Pachyrisma*; posterior adductor scar not observed; surface smooth. *L.Jur.*(*L.Lias.*), Eu. (N.Italy-France).—FIG. E219,1. **P. pumilum* (GUEMBEL), N.Italy; 1a, LV ext., 1b, LV hinge, both $\times 1$ (Tausch, 1890); 1c, RV hinge, $\times 1$ (Böhm, 1891).

Pterocardia BAYAN, 1874, p. 338 (ex FAVRE, *nom. nud.*) [**Cardium buwignieri* DESHAYES, 1857, p. 49 (syn., *C. corallinum* LEYMERIE, 1847, non LINNÉ, 1758); SD COSSMANN, 1906, p. 292] [= *Pterocardium* ROLLIER, 1912 (*nom. van.*)]. Large, thick-shelled, subrectangular, higher than long, inequilateral, posteriorly sublobate; umbones prominent, beaks anterior to median, but almost orthogyrous; ornament of radial ribbing; hinge plate massive and moderately broad; LV with stoutly triangular, anteriorly placed cardinal, short anterior lateral close to it, and more elongate and distant posterior lateral; RV with 2 obliquely aligned cardinals, posterior and lower one stoutly triangular, anterior and marginal one relatively small, together with anterior and posterior laterals; posterior adductor scar on raised lamina emerging from below hinge plate. *M.Jur.*(*Bathon.*)-*L.Cret.* (*Apt.*), Eu.-SW.Asia(Syria)-Mex.—FIG. E219,2. **P. buwignieri* (DESHAYES), *U.Jur.*(*U.Oxford.*), Swiss Jura; 2a,b, RV ext. and int.; 2c, LV int., $\times 0.7$ (de Loriol, 1891).

Superfamily HIPPURITACEA Gray, 1848

[*nom. transl.* NEWELL, 1965 (ex Hippuritidae GRAY, 1848)] [Materials for this superfamily prepared initially by COLETTE DECHASEAUX, typescript in French translated by R. C. MOORE reviewed and improved by L. R. COX with advice received from L. J. CHUBB (mainly on occurrence of Caribbean rudists); subsequently (1965-67) extensively revised, with additions of new typescript and illustrations produced by A. H. COOGAN and B. F. PERKINS, as noted in different families. The final version has been referred to the original author to allow expression of any disagreements. The Editor is responsible for integrating the text and illustrations and for accepting the taxonomic framework adopted in the *Treatise*, but with advice received from N. D. NEWELL. Illustrations with reference number 252 are taken by permission from *Traité de paléontologie*, MASSON et CIE (DECHASEAUX, 1952).] [Diagnosis of superfamily by COLETTE DECHASEAUX]

Inequivalve bivalves, most commonly attached to substrate by either right or left valve, but in rare instances free; solitary or gregarious. Hinge with two teeth and one socket in free valve and two sockets and one tooth in attached valve, except in *Diceras*, in which right valve invariably has two teeth and left valve only one. Two adductor muscles present, inserted directly on shell wall or on projecting myophores. *U.Jur.*(*U.Oxford.*)-*U.Cret.*(*Maastricht.*).

INTRODUCTION

By COLETTE DECHASEAUX

[Centre National de la Recherche Scientifique, Laboratoire de Paléontologie à la Sorbonne, Paris, France]

Members of the order Hippuritoida and superfamily Hippuritacea collectively are designated commonly as rudists. They were solitary or gregarious bivalve mollusks which generally were attached to the substrate by the extremity of one of their valves. Their highly varied external and internal morphological features separate them widely from all other representatives of the class, whether fossil or Recent.

From the time (1775) when PICOT DE LAPEIROUSE collected and figured fossil shells "called horns by the country folk because of their close resemblance to actual horns" (*Hippurites*) and "petrified vase-shaped oysters" (*Sphaerulites*) until 1825, naturalists were uncertain as to how they should be classified (Fig. E220). The diagnosis given in 1819 by LAMARCK (505, v. 6, p. 230), who introduced the term "les Rudistes," well illustrates the ignorance of contemporary scholars concerning the organization of these "shelly objects appearing to have some affinities with the oysters, but clearly distinguished from this group in that there is no trace of hinge, ligament, or muscle scars, and no indication as to where these were situated."

LAMARCK included certain brachiopods in his "rudists," but he omitted the hippurites because he considered them to be Cephalopoda. PICOT DE LAPEIROUSE was the first to include these "petrified oysters" with bivalves. CUVIER (1817) also suggested a relationship with this group, without giving convincing reasons. It was not until 1825 that DESHAYES and DE BLAINVILLE established that *Sphaerulites* and *Hippurites* should be classified as Bivalvia. This conclusion was opposed, however, by various naturalists, who continued to class the rudists as brachiopods, corals, or cirripeds. The study of rudists has been continued by S. P. WOODWARD, SAEMANN, BAYLE, PARONA, TOUCAS, DOUVILLÉ, and many others, who have made great advances in understanding these fossils; even so, many problems presented by these curious forms are not yet solved.






FIG. E220. Morphological features of rudists (hippuritids and radiolitids).

1. *Hippurites radiosus* DES MOULINS, U.Cret. (Maastricht.), France; 1a, view of operculiform FV showing posterior myophore (left), posterior tooth (middle), anterior tooth (right), and anterior myophore, $\times 0.45$; 1b, interior of AV from above, $\times 0.45$ (252).
2. *Hippurites bioculatus* LAMARCK, U.Cret. (L. Campan.), France (Pyrenees) (Montagne des Cornes); transv. sec. at level of AV with projecting teeth and myophore of FV, $\times 0.9$ (910).
3. *Praeradiolites cylindraceus* DES MOULINS, U. Cret. (Maastricht.), France; lat view of FV showing myophores and elongate teeth, $\times 0.5$ (911).
4. *Radiolites angeoides* PICOT DE LAPEIROUSE, U. Cret. (low.Campan.), France, Pyrenees (Montagne des Cornes); side view of both valves showing on exterior of AV positions of *Eb* and *Sb* siphonal bands, $\times 0.7$ (911).
5. *Sphaerulites foliaceus* LAMARCK, U.Cret. (Cenoman.), France (Bouche-du-Rhone); transv. sec. at level of AV with projecting teeth and myophores of FV, $\times 0.3$ (911).
6. *Hippurites striatus* DEFRANCE, U.Cret., France (Rennes-les-Bains), side view of AV with cuts for transv. secs. (not figured) near top, $\times 1.5$ (269).

[Explanation: *am*, *pm*, anterior and posterior myophores; *Eb* and *Sb*, siphonal bands; *Ep*, *Sp*, internal pillars farthest and nearest *L*; *L*, ligamental ridge; 1, 3, anterior and posterior teeth of FV; 2, tooth of AV; 1', 3', sockets in AV for reception of correspondingly numbered FV teeth.]

The earliest known rudists are found in Upper Jurassic (Rauracian) strata in France and elsewhere, and the latest ones lived during Maastrichtian times,¹ near the close of the Cretaceous Period. In intervening ages they spread widely, becoming highly diversified, and, according to present knowledge, were distributed along a belt extending between the latitudes of southern Sweden, which contains the most northerly rudist-bearing deposits, and Madagascar, which is the most southerly region where such fossils have been collected. Among the most prolific occurrences of rudists are those in Texas, Mexico, the Antilles, Portugal, France, the Alps, Italy, Yugoslavia, Turkey, Syria and Iran.

RUDIST MORPHOLOGY

By B. F. PERKINS²

[Louisiana State University, Baton Rouge, Louisiana]

SHELL SIZE AND SHAPE

Although the first rudists were only slightly inequivalve, their descendants very early became strongly so, with the two valves of individuals usually differing greatly from each other in size, shape, and shell wall structure. The inequivalve form in rudists was developed according to three basic plans (Fig. E221): 1) the ^{upper} attached valve coiled and larger of the two, the smaller, free valve flat or slightly convex and lidlike; 2) the attached valve conical and larger of the two, the smaller, free valve flat, lidlike, conical or coiled; and 3) the free valve coiled and larger of the two, the smaller, attached valve conical or only slightly coiled.

¹ It is now known that beds in Catalonia (Spain), once thought to be Danian in age and the latest known deposits containing rudists, belong in fact to the Maastrichtian, and are no younger than rudist-bearing beds of that stage found in other areas (L. P. MANGIN, 1963, 21 Sess. Internat. Geol. Congr., Norden, 960, pt. 27, p. 59).

² The contributions by PERKINS are adapted in part from Shell Development Company reports and are published with permission of the Company. He acknowledges with gratitude the opportunity provided by Shell Development Company to carry out the work on which his contribution is based. He also acknowledges the invaluable assistance of T. D. ALTMAN, J. F. LAVERDE, and J. W. MOORE in preparation of illustrations and of APHRODITE MAMOULIDES and BERNICE MELDE for library work.

Conical or gently curved rudist shells range in length from 2 cm. (*Monopleura*) to more than 2 m. (*Titanosarcolites*). They range in diameter from about 8 mm. (*Monopleura*) to more than 0.6 m. (*Bournonia*). Coiled rudist valves range in coil diameter from about 2 cm. (*Toucasia*) to at least 30 cm. (*Caprinuloidea*). Shell wall thickness of rudists varies from less than 2 mm. (*Monopleura*, *Toucasia*) to 10 cm. or more (*Durania*).

Terminology of the principal sections which show the internal features and shell-wall structure through different shapes of rudist shells is summarized in Figure E222.


ORIENTATION

In descriptive morphology of rudists the terms right valve (RV), left valve (LV), dorsal, ventral, anterior and posterior have been used commonly, but the application of these to many rudists is not so obvious as with normal bivalves. Use of the terms in rudist descriptions depends upon a comparison between the shells of rudists and living Chamidae.

Among living cemented Chamidae certain species are almost invariably attached by the RV or oppositely by the LV, although some species are fixed to the substrate indiscriminately by either valve (1024a). Comparison of attached LV's of some specimens with attached RV's of other specimens shows that they are mirror images with similar dentition. Comparison of free left with free RV's reveals that they, also, are mirror images. The similarity of dentition in attached valves of Chamidae was first noted by MUNIER-CHALMAS (652a), who designated the commoner condition of attachment by the LV as "normal" and the contrasting condition of attachment as "inverse." He recognized that the designation of a valve as left or right does not identify its dentition. Consequently, he proposed that in attached Chamidae the free valves be designated α and the attached valves β .

MUNIER-CHALMAS considered the inversion in *Chama* and conditions in the rudists as comparable. Accepting this, DOUVILLÉ (267a) recognized two series of rudists: a "normal" series in which the LV is at-

Bivalvia—Heterodonta



1
attached valve (AV) larger and coiled;
free valve (FV) smaller and flat,
slightly convex, or operculate

2
attached valve (AV) larger and
conical; free valve (FV) smaller and
flat, conical, coiled, or operculate

3
attached valve (AV) smaller and conical,
or slightly coiled; free valve (FV)
larger and coiled

FIG. E221. Principal rudist shell forms (FV, free valve; AV, attached valve) (Perkins, n).

- A. *Toucasia patagiata* WHITE, L.Cret.(Alb.), USA (Texas), $\times 1$.
- B. *Radiolites angeiodes* (PICOT DE LAPEIROUSE), U. Cret.(Campan.), France, $\times 1$.
- C. *Caprinuloidea* sp., L.Cret.(Alb.), USA (Texas), $\times 0.4$.
- D. *Monopleura marcida* WHITE, L.Cret.(Alb.), USA (Texas), $\times 0.4$.
- E. *Coralliochama orcutti* WHITE, U.Cret.(Senon.), Mexico, $\times 0.3$.
- F. *Sellaea* sp., L.Cret.(Alb.), USA (Texas), $\times 0.6$.




FIG. E222. Principal sections (polished or thin) through different shapes of rudist shells (*pc*, commissural plane) (Perkins, n).




FIG. E223. *Dicerias arietinum* LAMARCK (Dicerati-
dae), U.Jur.(Raurac.), France (Coulanges-sur-
Yonne); 1a, LV (FV) int., 1b, RV (AV) int.,
×0.5 (modified after 246). [Trace of ligamental
groove shown in solid, broken and dotted line.
Solid segment=ligamental groove parallel to com-
missure and probably containing functional liga-
ment at stage shown; broken line=segment of
groove visible in this view but below the level of
the commissure; dotted line=part of groove con-
cealed in this view below umbones and probably
not containing a functional ligament at stage shown.
Explanation as for Fig. E220; 2', socket for recep-
tion of tooth 2.]

tached and an "inverse" series in which the RV is attached. Paleontologists generally have followed the conclusions of MUNIER-CHALMAS and DOUVILLÉ. DECHASEAUX (252), on the other hand, has interpreted the teeth in these two series as not homologous, basing her opinion on a comparison of *Valletia*, earliest rudist, attached by the RV, with *Dicerias*. She has pointed out that teeth of the free LV of *Valletia* (see Fig. E250,4a) are not comparable to those in the RV of *Dicerias* (Fig. E223). The posterior tooth in *Valletia* is very weak, whereas in *Dicerias* it is quite strong. The posterior tooth of rudists attached by the RV becomes larger in successively younger genera and at least by Albian time is as prominent as in *Dicerias*. DECHASEAUX argued that inversion is only apparent in rudists and, consequently, she abandoned the former classification of the group into normal and inverse series.

In many rudists orientation in terms of right, left, dorsal, ventral, anterior and posterior can be made in the same manner (i.e., with the hinge line up and the beaks directed away from the observer) as for a normal bivalve. In conical forms such as the Radiolitidae and Hippuritidae, how-

ever, which show no evidence of coiling, orientation by this method is impossible. Without proof in some groups paleontologists have assumed that attachment was by the RV in Monopleuridae, Caprotinidae, Caprinidae, Hippuritidae and Radiolitidae, and by the LV in Heterodiceratinae, Plesiodiceratinae, Epidiceratinae, and Requieriidae. Based on this assumption, parts distinguished as anterior, posterior, ventral and dorsal can be determined. However, because of the uncertain identification of rightness or leftness in many rudist shells, it seems desirable to designate the two valves as attached valve (AV) and free valve (FV) in descriptions of rudist genera and species.

Anterior, posterior, dorsal, and ventral can be retained as convenient orientation terms, but it is important to recognize that these terms are applicable to conical shells only if assumption is made regarding rightness or leftness of the valves. Investigation of young shells of rudists which show no evidence of coiling in the adult may provide the ontogenetic information necessary to settle the question.

TEETH AND SOCKETS

Hinge structure is similar in all rudists. Except for *Dicerias*, rudists have two teeth in the free valve separated by a socket which receives the single tooth of the attached valve. In *Dicerias* two teeth are present in the attached RV and a single tooth occurs in the free LV (Fig. E223). Rudist hinge teeth vary from feeble tubercles weakly aligning the two valves to long, heavy straight or curved (Fig. E220,1a,3; E224,3) projections firmly guiding the opening and closing of the valves. The teeth and their corresponding sockets may be smooth or grooved.

Three principal notation systems for rudist dentition have been proposed, and several minor variations of these have been used (Table E1). The first system introduced by DOUVILLÉ (267a) has been used most commonly by rudist workers and has not been replaced completely by any later proposed notation system. This terminology

diceras *foretas*
char.
become
opinion




FIG. E224. Morphological features of rudists (coral-like shells with operculiform upper valve).

1. *Hippuritella maestri* VIDAL, U.Cret.(Santon.), France, Pyrenees (Montsech); side view of AV, $\times 1.7$ (269).
2. *Distefanella lombricalis* D'ORBIGNY, U.Cret. (Turon.), France (Charente); 2a-d, side views of 4 AV's showing shape variations and position of Eb and Sb siphonal bands, $\times 0.7$ (911).
3. *Caprinuloidea perfecta* PALMER, U.Cret.(Cenoman.), Mexico; side view of both valves (cap-like FV above), $\times 0.35$ (714).

[Explanation: Eb, Sb, siphonal bands farthest and nearest ligament; Ep, Sp, internal pillars.]

apparently was not derived from nomenclature used previously for other bivalves and was not intended by DOUVILLÉ to express homology. The second system used by DOUVILLÉ (278a) was derived from terminology of MUNIER-CHALMAS (652b) and BERNARD (37, 40). It expressed DOUVILLÉ's interpretation of the homologous relationships of rudist dentition. His last system (282) is only a slightly modified version of the second. Most other notation systems, such as that used by DI STEFANO (882a, 882b) and one used by KLINGHARDT (475), are minor variations of DOUVILLÉ's first nomenclature.

DECHASEAUX (245, 246, 252) noted the impossibility of distinguishing cardinal and lateral teeth in rudists, and consequently, the difficulty of applying with certainty the dental notation of MUNIER-CHALMAS and BERNARD. She proposed using the numerical system of BERNARD, without the implications of homology, however. In this system the teeth are numbered from front to back (i.e., 1, 2, 3) and their corresponding sockets are indicated by the same numbers with the addition of a prime accent (i.e., 1', 2', 3'). By this system tooth 2 of the Diceratinae is in the free LV and teeth 1 and 3 are in the attached RV. In all other rudists teeth 1 and 3 are in the free valve (FV) and tooth 2 is in the attached valve (AV), whether the LV or RV is attached.

This simplified notation without homologous implications first used by DECHASEAUX is followed in the *Treatise*.

LIGAMENTAL STRUCTURES

The ligament is not preserved in fossil rudists, but its original location is marked by a ligamental groove or furrow on the exterior of the shell, a ligamental cavity within the shell wall, or a ligamental ridge or pit on the interior of the shell. The ligament was external in the most archaic rudists, but it was progressively submerged into the shell wall of other groups, and in many advanced groups it was entirely internal or lost completely.

The ligament in the Diceratidae, Requiiniidae, and some Monopleuridae (e.g., *Valletia*, *Gyropleura*) lay in a groove parallel to the commissure. As the valves grew,

TABLE E1. Notation Systems for Rudist Teeth and Sockets Employed by Various Authors.

		DICERAS					
		A	RV	P	A	LV	P
		+	O	+	O	+	O
DOUVILLÉ, 1886 (267a)	Teeth	B'		B		N	
	Sockets		n		b'		b
DOUVILLÉ, 1918 (278a)	Teeth	AI		3b		AII	
	Sockets		AII'		AI'		3b'
DOUVILLÉ, 1935 (282)	Teeth	AI		3		AII-2	
	Sockets		AII'-2'		AI'		3'
DI STEFANO, 1889, 1899 (882a, 882b)	Teeth						
	Sockets						
KLINGHARDT, 1928 (475)	Teeth						
	Sockets						
DECHASEAUX, 1952 (252)	Teeth	1		3		2	
	Sockets		2'		1'		3'

EXPLANATION: + = tooth, O = socket; A, anterior; M, median; P, posterior.

the strong tangential growth component caused separation and coiling away from the commissure of the umbones and consequently the ligament was split so that only the most recently formed portion was en-

tire and paralleled the commissure, the location of the previously formed sections remaining as a coiled groove in each valve extended from the commissure to the beak (Fig. E223). Quite probably only the un-




FIG. E225. Diagram showing effect of tangential growth component on ligament, arrows denoting increased effect (1024a).

[Explanation: L-L, longitudinal axis; T-T, transverse axis; broken lines indicate divergence of umbones induced by interumbonal growth consequent on increasing effect of tangential growth component.—a. Primitive amphidetic ligament, longitudinally disposed and anteroposteriorly symmetrical.—b. Anteriorly separated and coiled ligament of *Glossus*, bilaterally but not anteroposteriorly symmetrical.

—b₁. Ligament of Chamidae, bilaterally asymmetrical.—c. Coiled ligament of Diceratidae, almost bilaterally symmetrical.—c₁. Ligament of *Requienia* bilaterally symmetrical.—d. Secondarily straightened ligament of *Caprina*, functionless and now disposed transversely.—d₁. Bilaterally asymmetrical ligament of Hippuritidae and Radiolitidae.]

TABLE E1. (continued)

RUDISTS WITH LEFT VALVE ATTACHED						RUDISTS WITH RIGHT VALVE ATTACHED					
	RV			LV			RV			LV	
A	M	P	A	M	P	A	M	P	A	M	P
+	O	+	O	+	O	O	+	O	+	O	+
B'		B		N		N		B'		B	
	n		b'		b	b'		b		n	
AI		3b		AII		3B		AII		PII	
	AII'		AI'		3b'	AII'		PII'		3b'	
AI		3		AII		3		AII		PIV	
	AII'		AI'		3'	AII'		PVI'		3'	
						N		D'		D	
						d'		d		n	
						N		B ₁		B	
						al ₁		al		N	
1		3		2		2		1		3	
	2'		1'		3'	1'		3'		2'	

split part of the ligament approximately parallel to the commissure was functional, and it is quite likely that the older functionless part disintegrated while the animal was still alive, as does the older functionless part of the ligament in some living oysters.

In all other rudists with conical, openly coiled or slightly twisted attached valves and coiled, conical, or operculiform free valves the long axis of the ligament, when present, was approximately normal to the plane of commissure. The reorientation of the ligament from a parallel-to-commissure to a normal-to-commissure structure is a result chiefly of the influence of the tangential component of growth. The effects of the tangential growth component on the ligament have been summarized diagrammatically by YONGE (Fig. E225).

In rudists with the ligament oriented normal to the commissure the ligament may have been functionless. This type of ligament, if functional, was certainly subject more to tensional stresses than to bending stresses of a parallel-to-commissure ligament. This normal-to-commissure ligament may have served only as a stringlike guide holding the dorsal margins of the valves together but not strongly counteracting contraction of the adductor muscles.

In the Monopleuridae (excluding *Valletia* and *Gyropleura*) the ligament of the AV lay in a groove normal to the commissure. The ligamental groove is shallow and does not penetrate the shell wall (*Petalodontia*, Fig. E252,4a,b) or it is deep and penetrates the shell wall so that an internal groove or cavity is developed (*Himeraelites*, E252,5b).

In monopleurids with operculiform FV's (*Monopleura*, E250,3b; *Araeopleura*, E250, 2b; *Petalodontia*, E250,1a,b) the FV ligamental position may be marked externally by a slight infolding or notching of the shell wall, but in some no external feature can be related to the ligament. Internally a shallow rounded or elongate ligamental pit is found between the dorsal margin and the tooth socket (2'). The FV ligament in species of *Monopleura* with the FV coiled in a low spiral (E251,3a; E252,1a) is represented externally by a groove which extends from the beak to the commissure and leads internally to a shallow ligamental cavity. In the elevated FV of *Himeraelites* (E252,5c) an external ligamental groove extends from the beak to the commissure and opens internally into a deep ligamental pit.

In the AV of caprotinids the ligament location is marked externally by a deep groove which may open internally into a




FIG. E226. *Caprinuloidea* sp. (Caprinidae), L.Cret. (Alb.), USA (Texas); FV int. showing ligamental cavity, $\times 1.3$ (Perkins, n).

ligamental cavity (*Sellaea*, Fig. E255,2b; *Polyconites*, E254,1b). A similar external groove and internal ligamental cavity are found in conical or coiled caprotinid FV's (*Sellaea*, Fig. 255,2a). The ligament location in caprotinids with operculate FV's may be marked externally by a short, shallow groove which corresponds internally to a ligamental pit (*Polyconites*, E254,1a; *Chaperia*, E253,1a).

The ligament position is defined in most caprinids by a shallow to deep external groove in both valves (*Caprinula*, E257,4d; *Caprinuloidea*, E258,3; *Planocaprina*, E259,3a; *Plagioptychus*, E260,2a; *Coralliochama*, E258,2a). The ligamental groove may open internally into a narrow or wide ligamental cavity (*Sabinia*, E262,3a,b; *Caprinula*, E257,4c; *Caprina*, E257,2a; *Caprinuloidea* sp., E226). In some caprinids, however, there is no external groove or internal structure which can be related to the ligament (*Dictyoptychus*, E256,2a-c).

In most hippuritids the ligamental position in the AV is marked by an external furrow and an internal ligamental ridge, but in some species (e.g., *Hippurites bioculata*, Fig. E220,2 and Fig. E263,1b) no internal ridge occurs and an external furrow is not developed. In other species a well-developed ligamental ridge is not matched by a distinct external furrow. The external furrow, when present, extends from the apex to the commissure and may be broadly

(E220,1b) or narrowly rounded (E220,6). In transverse section the ligamental ridge may be a low, rounded swelling (*Batolites*, E237,11), a sharp triangle (*Hippurites*, E220,1b) or rounded triangle (*Tetracionites*, E264,5), a long, parallel-sided projection with a truncated, rounded or swollen tip (*Torreites*, E238,5; E265), or a beaded (moniliform) ray (*Barrettia*, E227,2). An internal ligamental ridge and external furrow are present in the FV of some hippuritids (*Hippurites*, E220,1a) which correspond to the ridge and furrow of the AV. Hippuritids have been studied most commonly from transverse sections because separate, free valves are rare. As a result, little is known about the ligamental structures of the FV and only those features of the AV which are exposed in transverse sections are well known. It seems likely that the ligament in hippuritids was not functional.

An internal ligamental ridge is present in the AV of all Radiolitinae and in *Sauvagesia* (Sauvagesiinae), *Praelapeirousia* (Lapeirousiinae) and *Joufia* (subfamily uncertain). An external groove (*Paronella*, E269,4) corresponding to the ligamental ridge is not found commonly in radiolitids. In transverse section the ridge may be long, slender and parallel-sided (*Pseudopolyconites*, E266,3), bilobed (*Kuehnia*, E267,5), or truncate (*Neoradiolites*, E269,6). An internal ligamental ridge is known in the FV of *Praeradiolites* (E220,3; E269,2e) *Eoradiolites*, *Sauvagesia* and *Joufia*. The FV ligamental ridge is straight and narrow and may terminate in a ligamental pit. A ligamental ridge may be developed in the FV of other genera which have ridges in the AV. The ligament of most radiolitids probably was functionless, but in some (*Praeradiolites*, *Eoradiolites*, and others) it may have served as a stringlike guide.

SIPHONAL STRUCTURES

Structures presumably related to siphons are present in many rudists. Siphonal structures include bands, fossettes, pillars, pseudopillars and oscules. These occur in pairs when present and have been designated by DOUVILLÉ as "E" and "S" according to their positions. The posterodorsal structures

Hippuritoida—Hippuritacea—Morphology




FIG. E227. Morphological features of rudists (shell structure) (hippuritids).

1. *Pironaea polystylus* (PIRONA), U.Cret. (Maas-tricht.), Italy; transv. sec. of type (AV) showing radial inward projecting partitions formed of bundled calcite prisms, $\times 0.5$ (269).
2. *Barrettia monilifera* WOODWARD, U.Cret. (Maas-tricht.), Jamaica; transv. sec. of type (AV) showing radial beaded structure in thick shell wall, $\times 0.5$ (269).

[Explanation as for Fig. E220.]

nearer the ligamental position were presumed by DOUVILLÉ to mark the location of the excurrent siphon and hence were designated "S" for *sortie* (French, excurrent). The ventral structures farther from the ligamental position were presumed to represent the incurrent siphon location and

were named "E" for *entré* (French, in-current). DOUVILLÉ's interpretation of these structures has been questioned by YONGE (1027a) who presented a biologically convincing alternative explanation (discussed elsewhere in this introduction by COOGAN). However, DOUVILLÉ's symbols "E" and "S" have been used generally and are retained in the *Treatise* but modified by a lower case letter to denote the particular structure. For example, *Eb* and *Sb* mark bands, *Ep* and *Sp* denote pillars, *Es* and *Ss* indicate pseudopillars, and *Ef* and *Sf* indicate fossettes. Oscules are designated *Eo* and *So*.

Siphonal bands (Fig. E224,1,2; E254,2b) are external features of the AV (and, in some genera, the FV) of Radiolitidae, a few Requieniidae (*Requienia*, *Toucasia*) and one Caprotinidae (*Horiopleura*). Siphonal bands are broad to narrow, salient or superficial to deeply impressed areas with the long dimension normal to the commissure and extending from the apex of the valve to the commissure. In Requieniidae the bands are broad areas marked by slight deflections of the growth lines (*Requienia*) and slight depression of the area (*Toucasia*). The siphonal bands of *Horiopleura* (Caprotinidae) (Fig. E254,2b) are smooth depressed areas on an otherwise radially ribbed shell.

The siphonal bands of the AV of most Radiolitidae are smooth, shallow, depressed areas as in *Radiolites* (Fig. E267,2). However, in some radiolitids the bands may be finely and longitudinally costulate (*Sauvagesia*) (Fig. E266,4), broadly costulate (*Medeella* (*Medeella*)), marked by strongly convex-upward folds (*Praeradiolites*) (Fig. E269,2) or occur in deeply sunken grooves (*Thyrastylon*) (Fig. E243,2). The siphonal bands of *Medeella* (*Fossulites*) (Fig. E243,1) are salient areas which mark cylindrical siphonal fossettes within the shell wall. In *Tampsia* (Fig. E272,2) sharp inflections define the bands and *Eb* is marked by a deep slit cutting the shell wall almost to the inner layer of the shell. In many radiolitids no external feature of the FV corresponds to bands of the AV, although in some genera (*Hardaghia*) bands are present in the FV. In *Medeella* (Fig. E243,1a; E267,1c) inflected crescentic slits or arches in the FV correspond to bands and fossettes




FIG. E228. *Sellaea* sp. (Caprotinidae), L.Cret. (Alb.), USA (Texas); long. sec. both valves showing mantle-shell cavity open without tabulae (*p-c*; commissural plane), $\times 0.7$ (Perkins, n).

of the AV and in *Thyrastylon* openings or oscules through the FV correspond to the deeply sunken bands of the AV.

In hippuritids infoldings of the AV shell wall produced ridges normal to the commissure (Fig. E220, 1*b*, 2). These ridges or pillars are usually three in number, one marking the ligamental position and the other two presumably siphonal in location. The pillars are matched in the FV by openings, oscules, through the valve (Fig. E263, 1*a*, 2*b*).

In *Parasauvagesia* (Radiolitinae) and in most Lapeifousiinae development of cellular structure (Fig. E273, 6) in the outer shell wall produces pillar-like ridges (termed pseudopillars) on the inner shell wall. Oscules in the FV correspond to pseudopillars in the AV. The function of pillars, pseudopillars, and oscules is discussed by COOGAN in "Evolutionary Trends in Rudist Hard Parts," and in the systematic sections on Hippuritidae and Radiolitidae.




FIG. E229. *Eoradiolites* sp. (Radiolitidae), L.Cret. (Alb.), USA (Texas); natural long. sec. of two AV's showing closely spaced tabulae in mantle-shell cavity, $\times 0.7$ (Perkins, n).

MANTLE-SHELL CAVITY

The mantle-shell cavity of rudists is a single or subdivided cavity within each of the two valves. The cavities of the valves are nearly equal in size in the Diceratidae and in many Caprotinidae and Caprinidae. In the Requienniidae, Monopleuridae, Radiolitidae, and Hippuritidae the shell cavity of the AV is much more elongate normal to the commissure and is larger than that of the FV. In the shells of species which have operculiform FV's the cavity of that valve is reduced to a shallow depression. In the Caprotinidae, Caprinidae, some Requienniidae and Monopleuridae the shell cavity consists of a principal cavity which contained the visceropodal mass and gills along with accessory cavities which probably were lined only by the mantle. In rudists having relatively thin or compact shell walls (e.g., Diceratidae, Requienniidae, Monopleuridae, and Caprotinidae) the shell cavity is large in proportion to the cross-sectional area of the valve. In rudists with thick shell walls (Hippuritidae) or spongy shell wall structure (Radiolitidae, Caprinidae) the principal shell cavity may be very small in proportion to the valve's cross-sectional area.

The shell cavity may be open throughout its length from the commissure to the um-




FIG. E230. *Hippurites bioculatus* LAMARCK, U.Cret. (L.Campan.), France (Corbières); long. sec. both valves showing tabulae in mantle-shell cavity of AV, $\times 1.3$ (Perkins, n).




FIG. E231. *Eoradiolites davidsoni* HILL (Radiolitidae), L.Cret.(Alb.), USA(Texas); long. sec. both valves showing mantle-shell cavity size in relation to shell size, $\times 0.7$ (Perkins, n).

bonal area (Fig. E228), but in Caprinidae, Radiolitidae, and Hippuritidae the shell cavity commonly is divided by transverse tabulae concave toward the commissure (Fig. E229-E231). In caprinids in which the shell cavity is divided into a principal cavity and accessory cavities all cavities may be tabulate but the tabulae in these various spaces do not occur at common levels. The tabulae close off earlier-formed portions of the shell cavity and spaces remain as vacant chambers. In some Hippuritidae the earlier-formed part of the shell cavity is filled from the umbo toward the commissure by dense shell material. The tabulae and dense shell material reduce the available space in the cavity for the animal which was small in relation to size of the shell in many caprinids, radiolitids, and hippuritids (Fig. E231).

ACCESSORY CAVITIES

Accessory cavities were first defined by DOUVILLÉ (267a) from studies of *Caprotina* (Caprotinidae). These cavities separate anterior and posterior myophore plates and

teeth from the shell wall in caprotinids and caprinids. They are developed chiefly in the FV but also occur in the AV of some genera.

Accessory cavities are large in most caprotinids but may be reduced greatly (*Chaperia*, *Retha*). They are usually oval or lunate in section parallel to the commissure and in some shells are divided by thin, radial plates normal to the commissure. Accessory cavities occur in all caprotinid genera but are not limited to this family, for accessory cavities are found in *Bayleia* and *Bayleioidea* (Requeniidae), *Himeraelites* (Monopleuridae), and in most Caprinidae.

The posterior accessory cavity is connected with the posterior socket in the AV of Western Hemisphere caprinid genera (e.g., *Amphitriscoelus*, *Coalcomana*, *Caprinuloidea*) and has a characteristic shape in transverse section. MACGILLAVRY (561) first noted this distinction of Western Hemisphere caprinids and described the cavity as "more or less shaped like an ant's larva, or better, like the ornament one often encounters on oriental tapestry" (Fig. E241; E257,1a; E232). The AV posterior accessory cavity in some caprotinids (e.g., *Sellaea*) is connected also with the posterior socket, which, in turn, is connected to the ligamental cavity (Fig. E239,3a; E255,2b).

A myophorous accessory cavity is present in the FV of some caprotinids (e.g., *Pachytraga*, Fig. E253,4) and caprinids (e.g., *Caprina*, Fig. E256,1; *Caprinula*, Fig. E257,4b; *Caprinuloidea*, *Plagioptychus*, Fig. E259,5b). These accessory cavities are usually connected to the tooth socket (2') and may be only partially separated from the principal shell cavity. The cavity is formed by a vertical plate connecting the anteroventral margin of the anterior tooth (1) with the anteroventral shell wall. The plate does not reach the commissure in most species, so that at this level the principal cavity is not separated from the myophorous accessory cavity. The anterior muscle insertion of the FV is on the anterior or anteroventral cavity and the cavity received the anterior myophore plate of the AV when the valves were closed.

Accessory cavities occur in the AV of some hippuritids, radiolitids, and caprinids




FIG. E232. *Caprinuloidea* sp. (Caprinidae), L.Cret. (Alb.), USA (Texas); AV int. showing characteristic shape of posterior accessory cavity and connecting posterior socket (3') in western hemisphere caprinids, $\times 0.7$ (Perkins, n).

between the cardinal structures and the dorsal shell wall.

DOUVILLÉ postulated that accessory cavities developed as the valve increased in height and the teeth and myophore plates became erect to the point that these structures no longer increased in size at the same rate as the shell diameter. As a result, a cavity developed between the muscle insertion surfaces and teeth on one side and the shell wall on the other. Consequently, a much larger but much less dense shell was formed.

Accessory cavities converge toward the apex of the shell. Steinkerns of the cavities appear as clusters of cones attached to each other by their bases and are the "birostres" of 18th and early 19th century French paleontologists.

PALLIAL CANALS

Longitudinal, thin-walled canals, called **pallial canals**, in the shell wall characterize the Caprinidae. These canals may be present in either or both valves of a shell, and although they occur only in the shell wall of most caprinids they may be in the dental apparatus of some genera (e.g., *Caprinuloidea* and *Titanosarcolites*, Fig. E232). The canals are bounded by radial, transverse and tangential plates lying normal to the commissure. The radial plates may be simple, bifurcate or many-branched, and the trans-

verse and tangential plates may be regularly or irregularly arranged. In transverse section pallial canals may be approximately quadrangular (e.g., *Amphitriscoelus*), regularly oval to pyriform (e.g., *Coalcomana*, *Caprinuloidea*), or irregularly polygonal (e.g., *Coralliochama*). The pallial canals of some caprinids are traversed by thin tabulae concave toward the commissure.

In some shells a distinction cannot be made between pallial canals and divided accessory cavities. This observation caused DOUVILLÉ to speculate that pallial canals originated by the invasion of the pallial region by canals formed in advanced stages of the subdivision of accessory cavities.

Development of abundant pallial canals in caprinids produced a thick, spongy, porous shell wall which probably was rapidly deposited and which contained a minimum amount of calcium carbonate for its size and strength.

Pallial canals are discussed further in the section on shell-wall structures (Caprinidae).

MUSCLE INSERTIONS

The two adductor muscles in rudists were inserted on the shell wall, joined to projecting plates or myophores or within myophorous sockets. Within an individual rudist the type of muscle insertion may be different for the two valves or for the two muscles and are not necessarily symmetrical. For example, in Diceratidae the anterior muscle insertions in both valves may be on the shell wall, but the posterior insertions are on myophore plates. In Radiolitidae the FV muscle insertions are on projecting myophores but the AV insertions are on the shell wall (Table E2).

Shell-wall insertions may be superficial, on weakly to strongly depressed areas or on thickened areas.

Platelike prolongations of the cardinal platform are muscle insertion sites in some Diceratidae (e.g., *Heterodicerias*, Fig. E244, 1; *Mesodicerias*, Fig. E244, 2a,b), Requiiniidae (e.g., *Kugleria*, Fig. E246, 6a) and Monopleuridae (e.g., *Monopleura*, Fig. E252, 1). The insertion surfaces on plates of this type are parallel or nearly parallel

TABLE E2. Summary of Muscle Insertions in Rudists.

FAMILY	ATTACHED VALVE		FREE VALVE	
	ANTERIOR INSERTION	POSTERIOR INSERTION	ANTERIOR INSERTION	POSTERIOR INSERTION
DICERATIDAE	1) Shell wall or 2) prolongation of cardinal platform	1) Plate projecting above shell wall or 2) prolongation of cardinal platform	1) Shell wall or 2) on prolongation of cardinal platform	1) Plate projecting above shell wall, 2) prolongation of cardinal platform, or 3) plate extending beneath cardinal platform
REQUIENIIDAE	Shell wall	1) Shell wall or 2) on plate extending beneath cardinal platform	1) Shell wall or 2) on prolongation of cardinal platform	1) Plate extending beneath or above cardinal platform, 2) prolongation of cardinal platform, 3) plate separated from shell wall by accessory cavity, or 4) plate parallel to shell wall and attached to it by plate normal to shell wall
MONOPLEURIDAE	Extension of cardinal platform as thickened area or buttress	Extension of cardinal platform as thickened area or buttress	Extension of cardinal platform as thickened area, buttress or myophore plate	1) Extension of cardinal platform as thickened area, buttress or myophore plate or 2) on erect separate myophore plate or lamina
CAPROTINIDAE	1) Shell wall or 2) on plate separated from shell wall by accessory cavity	1) Shell wall or 2) on plate separated from shell wall by accessory cavity	1) Shell wall or 2) on plate separated from shell wall by accessory cavity	Shell wall or on plate separated from shell wall by accessory cavity
CAPRINIDAE	1) Shell wall or 2) on plate separated from shell wall by accessory cavity	1) Shell wall or 2) on plate separated from shell wall by accessory cavity	1) Shell wall or 2) on plate separated from shell wall by accessory cavity	1) Shell wall, 2) on plate separated from shell wall by accessory cavity, or 3) on projecting myophore which fits into socket of AV
HIPPURITIDAE	Shell wall	In socket which receives FV myophore	Projecting myophore or buttress	Myophore which fits into AV socket
RADIOLITIDAE	Shell wall	Shell wall	Projecting myophore or buttress	Projecting myophore or buttress

to the commissure. These plates are attached to the shell wall along their front (for the anterior plate) and back (for the posterior plate) edges and curve around the shell cavity. Similar plates which are not connected directly to the cardinal platform are

developed in the Requieniidae. These plates either pass above (e.g., RV posterior of *Requenia*, Fig. E246,1b) or below (e.g., LV posterior of *Bayleia*, Fig. E246,2a; LV posterior of *Apricardia*, Fig. E246,3) the cardinal platform.

Erect myophore plates are present in many Caprotinidae (e.g., *Sellaea*, Fig. E255, 2b) and most Caprinidae (e.g., *Amphitriscoelus*, Fig. E257, 1, *Schiosia*, Fig. E262, 5). Plates of this type are normal to the commissure and extend from the hinge to the posteroventral and anteroventral shell wall. The plates are separated from the anterior and posterior shell wall by simple or subdivided accessory cavities (described above).

Strongly projecting myophorous apophyses or myophores are characteristic of the FV of most Radiolitidae and Hippuritidae. Myophores of this sort may be elongate and toothlike (e.g., *Hippurites*, Fig. 220, 1a) or thick-based with the tip swollen into a knob or boss (e.g., *Praeradiolites*, Fig. 220, 3). The two myophores in a shell may be similar in form and subequal in size (e.g., *Praeradiolites*, Fig. 220, 3), or they may be quite unsymmetrical in size and shape as in *Hippurites* (Fig. 220, 1a).

A muscle buttress is intermediate between an insertion on a thickened shell wall area and a well-developed projecting myophore. A buttress is an elevated or pendent structure in which the insertion surface usually is only slightly inclined to the shell wall as in the FV of *Agriopleura* (Fig. E237, 1). Buttresses are usually developed as extensions of the cardinal platform to which they remain attached. They are found in the FV of some Radiolitidae and in the AV of some species of Monopleuridae (Fig. E233).

In hippuritids the posterior muscle insertion area in the AV is on the inner wall of a socket dorsal to *Sp* (e.g., *Hippurites*, Fig. 220, 1b). When the valves are closed the toothlike posterior myophore of the FV fits into the myophorous socket of the AV.

MECHANISM FOR OPENING AND CLOSING VALVES

By COLETTE DECHASEAUX

The almost constant occurrence of attachment scars for anterior and posterior adductor muscles indicates that the rudists could open and close their valves. In all except the Hippuritidae and Radiolitidae the apparatus for this purpose, consisting of

a ligament, well-developed teeth (not, however, forming long projections), and muscle scars, situated either directly on the shell wall or on more or less projecting apophyses (myophores), shows that the two valves could be opened by a hinging movement as in an ordinary lamellibranch. The form of the teeth in some shells (e.g., *Plagioptychus*) may have required a slight rotational movement of the free valve with respect to the fixed valve in order to disengage the expanded extremity of a tooth from its socket.

The problem of opening the shell is more involved when we consider the Hippuritidae and Radiolitidae, in which the teeth and myophores form elongate projections (Fig. E220, 1, 3). The free valve in this group forms a lid to the attached valve, its long teeth and myophores being received in elongate sockets, so that only vertical movements are possible. The free valve, when raised above the fixed one, is guided by grooves along the teeth and also (in some forms) along the myophores and by corresponding ridges along the sides of their sockets (Fig. E220, 1, 3). It may be suggested that the free valve could be raised by increased pressure inside the shell, brought about by expansion of the soft parts such as the foot. When the valves were separated the muscles were stretched, but when internal pressure was relaxed they would resume their normal length, thus closing the shell.

MODE OF LIFE

By COLETTE DECHASEAUX

Some rudists lived as solitary individuals (Fig. E224), whereas others were gregarious, their accumulated shells now forming dense masses called "rudist reefs." Except for certain radiolitids the form of which suggests that they lay resting on a flattened side on the sea bottom, and also some others with an equivalve shell lacking an attachment scar, and thus possibly capable of some free movement, all rudists grew attached, either intermingled with other organisms, particularly corals, or, when in an erect position, in clusters of varying magnitude. The earliest known species occur in coral reefs, forming an ele-

diante (tooth)
|
vicarities




FIG. E233. *Monopleura* sp. (Monopleuridae), L. Cret. (Alb.), USA (Texas); dorsal oblique view of AV showing muscle buttresses and external ligamental groove, $\times 1.3$ (Perkins, n).

ment of about the same importance as less specialized bivalves, gastropods, echinoderms, etc.

Hippuritidae and Radiolitidae are also associated locally with corals (as in eastern Serbia), but then other elements of the usual coral reef fauna are absent. Generally they occur in colonies of a single species or in assemblages containing only a small number of species. Near Angoulême, in France, certain Turonian beds which can be traced for several dozen meters contain only *Distefanella lombricalis* (D'ORBIGNY) (Fig. E224,2), apart from a very impoverished microfauna.

ORIGIN AND EXTINCTION

By COLETTE DECHASEAUX

According to present knowledge, the appearance of *Diceras* in Late Jurassic (Rauracian) time marks the beginning of the history of the rudists, a history extending almost to the close of the Cretaceous and including the development of a host of highly varied forms. Two hypotheses have been advanced as to the origin of *Diceras*. According to DOUVILLÉ, its ancestor was a bivalve living where its earliest representatives are found and having some of its distinctive characters. These requisites, it seemed to him, were possessed by *Pterocardium*, a genus in which the ornament consists of radial costae and in which the posterior muscle was attached to a projecting lamella. He supposed that the newly hatched young of *Pterocardium* became

cemented to the substrate with the aid of the foot to avoid being carried away by currents during early life, and that as the shell developed it became twisted about the point of fixation so as to acquire a spiral form, the number of hinge teeth being reduced to three. Such modifications led ultimately to the genus *Diceras*.

The other hypothesis (by DECHASEAUX) is based on the tendency of the posterior elements of the hinge of *Megalodon* and *Protodiceras* to become reduced. In *Megalodon* (Devonian-Upper Triassic) one (or in some shells two) of these teeth are absent. In *Protodiceras* (Lower Jurassic) which normally has fewer hinge elements than *Megalodon*, a tendency toward reduction of the teeth on the posterior side occurs also, the final stage corresponding to the hinge of *Diceras*. Thus, beginning in the Devonian, a tendency toward development of the *Diceras* type of hinge is shown by *Megalodon* and *Protodiceras*. In support of this hypothesis we may note that in early growth stages of *Diceras* the posterior muscle was attached in the same way as in *Megalodon*.

Among the last surviving rudists were three radiolitids and a hippuritid which lived in Catalonia during the Maastrichtian Stage. They were not solitary bivalves, but on the contrary formed large "rudist reefs." Their abrupt extinction is the more difficult to explain since in this region deposition of calcareous sediment had continued uninterruptedly from the Coniacian onward, some species nevertheless disappearing while others persisted and new ones appeared; then, at the same moment, apparently all became extinct.

In Somalia, where the succession has been studied by TAVANI, Upper Cretaceous limestones pass without a break into lower Eocene limestone, yet no rudists occur in beds higher than Upper Cretaceous. The nature of the sediments gives no indication of a possible change in ecological conditions, so that we are entirely ignorant of factors responsible for extinction of the rudists and other groups that disappeared; more particularly, we are unable to ascertain what changes accounted for the fact that new species of these groups ceased to appear to replace those that died out. All we can say is that extinction of the group took place at this time.

CLASSIFICATION

By COLETTE DECHASEAUX

The hypothesis is here accepted that the rudists were derived from the Megalodontidae, and they are considered to form a major taxon of the Bivalvia to which STEINMANN assigned the name Pachyodonta, and for which the name Hippuritoida is adopted in the *Treatise*. The rudists constitute the superfamily Hippuritacea. It is thought probable that the Chamacea are not descended from them but are sessile heterodonts. In early ontogeny the shell of chamids is unattached and regular both in outline and in the nature of its concentric ornament. Much has yet to be learned of the early ontogeny of the rudists, but we have no knowledge of the existence of a comparable stage in this group.

To a large extent, the classification of the rudists in families has been based on the various characters (fixation, hinge structure, etc.) which have been discussed in preceding paragraphs. Certain special internal features (accessory cavities, canals, pillars, pseudopillars) allow other families to be defined, as stated in the systematic descriptions which follow.

A tabular summary of the classification adopted in the *Treatise*, showing the number of genera and subgenera (for example, "2; 4" denoting 2 genera and 4 subgenera) in each division, and recording known stratigraphic distribution, follows.

Divisions of Hippuritacea

- Hippuritacea (superfamily) (115;3). *U. Jur.* (*U. Oxford.*)-*U. Cret.* (*U. Maastricht.*)
 Diceratidae (8). *U. Jur.* (*U. Oxford.*)-*L. Cret.* (*Valangin.*)
 Diceratinae (1). *U. Jur.* (*U. Oxford.*)-*L. Kimmeridg.*)
 Heterodiceratinae (3). *U. Jur.* (*Tithon.*)-*L. Cret.* (*Valangin.*)
 Plesiodiceratinae (2). *U. Jur.* (*Oxford.*-*Kimmeridg.*)
 Epidiceratinae (2). *U. Jur.* (*Oxford.*)-*L. Cret.* (*Valangin.*)
 Requieniidae (8;2). *U. Jur.* (*Tithon.*)-*U. Cret.* (*Maastricht.*)
 Monopleuridae (8). *L. Cret.* (*Valang.*)-*U. Cret.* (*Maastricht.*)
 Caprotinidae (8). *L. Cret.* (*Neocom.*)-*U. Cret.* (*Turon.*)
 Caprinidae (23). *L. Cret.* (*Barrem.*)-*U. Cret.* (*Maastricht.*)

- Hippuritidae (12). *U. Cret.* (*Turon.*-*Maastricht.*)
 Radiolitidae (39; 1)^a. *L. Cret.* (*Barrem.*)-*U. Cret.* (*Maastricht.*)
 Radiolitinae (14; 1). *L. Cret.* (*Barrem.*)-*U. Cret.* (*Maastricht.*)
 Biradiolitinae (7). *U. Cret.* (*Turon.*-*Maastricht.*)
 Sauvagesiinae (7). *L. Cret.* (*Alb.*)-*U. Cret.* (*Maastricht.*)
 Lapeirousiinae (8). *U. Cret.* (*Santon.*-*Maastricht.*)
 Subfamily Uncertain (3).
 Family Uncertain (9).

EVOLUTIONARY TRENDS IN RUDIST HARD PARTS

By A. H. COOGAN

[COOGAN'S present affiliation is with Kent State University, Kent, Ohio. His *Treatise* contributions are based mainly on studies made while he was employed by the Humble Oil & Refining Company and Esso Production Research Company, Houston, Texas, to the staffs and management of which he acknowledges assistance and encouragement. Special appreciation is expressed to R. M. JEFFORDS, R. D. WOODS, and the late H. N. FISK.]

The morphological diversity of rudists is a stony monument to the plasticity of the bivalve mantle as it operated in adapting to a variety of reef, bank, and shallow-shelf environmental niches available on extensive sea-covered continental platforms of the Late Jurassic and Cretaceous. Broadly speaking, two groups of rudists can be recognized with respect to their rapidity of morphological change. The conservative, little-changing, and for the most part older families are the Diceratidae, Requieniidae, Monopleuridae, and Caprotinidae. The more plastic, rapidly changing, adaptive families are the Caprinidae, Radiolitidae, and Hippuritidae (Fig. E234). In the conservative group distinctions between families involve slight differences in dentition and musculature, different modes of attachment, variations in shape and size, and the development of internal structures. In the more plastic families these same differences occur at the subfamily or generic level and in addition dramatic changes are observed in wall structure, mechanism of opening the valves, and development of structures for lightening and strengthening the shell and for support of the siphons.

Evolutionary trends with morphological significance can be discussed selectively in terms of dentition, ligamental and muscular supports, wall structure, siphonal structures and pillars. Several instances of parallel evolutionary trends in one or more of

^a Includes 3 radiolitids of unknown affinity.




FIG. E234. Late Jurassic and Cretaceous stratigraphic distribution of rudist families, more plastic and rapidly evolving ones distinguished by oblique-ruled pattern (Coogan, n).








FIG. E235. Diagram indicating effects of tangential components in growth of various bivalve shells—Glossidae (1), Chamidae (2), Hippuritacea (3-6) (1024a).

1. *Glossus*; 1a, lying on LV; 1b, lying on RV.
2. Chamids; 2a, normal *Chama* viewed from dorsal side, lying on LV; 2b, inverse *Pseudochama*, lying on RV.
3. *Heterodicerias* (Diceratidae), with "normal" dentition, viewed from dorsal side, attached by LV.
4. *Requienia* (Requieniidae), with "normal" dentition, viewed from dorsal side, attached by LV.
5. *Caprina* (Caprinidae), attached by RV; 5a, *C. adversa*, with almost planospirally coiled LV, viewed from dorsal side; 5b, same, viewed from left side.
6. *Hippurites* sp. (Hippuritidae), attached by RV; 6a, long. sec.; 6b, posterodorsal side view showing position of Ep and Sp pillars inside shell.

[Ligament shown plain where exposed, indicated by broken line where enclosed by shell. Arrows mark locations of inhalant and exhalant apertures. Explanation: am, pm, anterior and posterior myophores; Ep, Sp, internal pillars; lig, ligament.]

these features are known from different families and subfamilies. YONGE (1024a) has recently emphasized a highly developed tangential component of shell growth in rudists and an associated lateral extension and coiling of the umbonal regions of the shell with attachment as the cause of great bilateral asymmetry in rudists (Fig. E235).

DENTITION

Across the whole spectrum of rudist families runs a trend toward modification of dentition from the "primitive" diceratid type (Fig. E236,2), which itself is a modification of heterodont dentition by reduction in tooth number, to extremes of the caprinid, radiolitid, and hippuritid dentitions.

From the simple, slightly unequally sized teeth of *Diceras* a trend in the radiolitids is directed toward enlargement and elongation of the FV teeth accompanied by reduction in size or loss of the AV tooth, or both. For example, in the oldest radiolitid genus, *Agriopleura* (Fig. E237,1), the FV teeth are already elongate. *Eoradiolites*, a younger genus, has similar FV teeth, as does *Praeradiolites* (Fig. E220,3), but has a narrow finger-like AV tooth (Fig. E237,2). A reduction in the AV tooth occurs in the bioradiolitine line (e.g., *Distefanella*, Fig. E237,8) and complete loss of the tooth in other bioradiolitines (e.g., *Biradiolites*, Fig. E237,4). A parallel evolutionary trend of reduction and loss of the AV tooth occurs in the Lapeirousiinae (e.g., *Dubertretia*, Fig. E237,7). In the hippuritids the FV teeth are elongate (e.g., *Hippurites*, Fig. E220,1a), as in the radiolitids, and the AV tooth is reduced in size but not lost (e.g., *Hippurites*, E220,2; *Barrettia*, Fig. E237,10).

(On facing page).

FIG. E236. Morphological features of rudists (diceratids).

1. *Diceras rotundatum* BAYLE, U.Jur.(Raurac.), France (Coulanges-sur-Yonne); 1a, exterior, front view (AV at left) of type specimen showing strongly coiled beaks, $\times 0.7$; 1b, interior of FV showing single tooth (2), $\times 0.7$ (246).
2. Diagrammatic sketches of interiors of FV (2a) and AV (2b) showing teeth and sockets and muscle insertions (246).

[Explanation as for Fig. E220; 2', socket corresponding to tooth 2.]

In the Caprinidae distinct trends are less clearly discernible among the considerable diversity of tooth types. In some genera the teeth are large and porous (e.g., *Caprinuloides*, Fig. E237,3), and located nearly in the center of the shell. In others they are reduced, apparently solid, and placed marginally (e.g., *Caprinula*, Fig. E237,5).

LIGAMENT SUPPORTS

Ligament supports of calcite and in some fossils calcite molds or casts of the flexible ligament are preserved in many rudists. In the primitive diceratid condition the ligament was coiled. In fact, it was already greatly modified from the primitive amphidetic ligament (Fig. E225). A general trend toward reduction in size of ligamental supports partly results from the strong tangential growth component and consequent interumbonal growth of the ligament which may become functionless. The ligament moves to the outside of the shell as rudists evolved from slightly unequally coiled diceratid-requieniid shape to extremely unequal vase-like forms of the radiolitids and hippuritids. Among radiolitids an internal ligament support occurs (e.g., *Eoradiolites*, Fig. E237,2), which becomes reduced in size and eventually is lost in branches of three separate radiolitid subfamilies. The bioradiolitines, which probably evolved from one of the radiolitine genera, have no internal ligament ridge (e.g., *Biradiolites*, Fig. E237,4; *Chiapasella*, Fig. E238,1). In the Sauvagesiinae the ligamental ridge is present in the older genus *Sauvagesia* (Fig. E238,4) but is lost in other genera (e.g., *Durania*, Fig. E237,9; *Tampsia*, Fig. E238,2). In the Lapeirousiinae a similar loss occurs. *Praelapeirousia* has a weak ligament ridge and other genera lack it entirely (e.g., *Lapeirousia*, *Dubertretia*, Fig. E237,7). In the caprinids the ligament has a strong external expression in the form of a groove and commonly is curved internally. The hippuritids, starting from a Turonian shell with a relatively small ligament ridge formed by infolding of the shell wall (e.g., *Hippurites*, Fig. E220,1b), developed enormously elongate and thick ligament pillars (e.g., *Vaccinites*, Fig. E237,6; *Torreites*, Fig. E238,5). In another hippuritid line the




FIG. E237. Morphological features of rudists (dentition).

1. *Agriopleura blumenbachi* (STUDER) (Hippuritiidae), L.Cret.(Barrem.), France (Brouzet); side view of FV showing projecting teeth, $\times 0.6$ (911).
2. *Eoradiolites davidsoni* (HILL) (Radiolitidae), L.Cret.(mid.Alb.), USA(Texas); transv. sec. AV, $\times 1.9$ (Coogan, n).
3. *Caprinuloidea* sp. (Caprinidae), U.Cret.(Cenoman.), Mexico; schematic transv. sec. FV (714).
4. *Biradiolites canaliculatus* D'ORBIGNY (Radiolitidae), U.Cret.(Coniac.), France; transv. sec. AV, $\times 0.6$ (277).

(Continued on facing page).

ligament ridge was modified to a thin bulbous pillar (compare *Batolites*, Fig. E237,11, and *Pironaea*, Fig. E238,6). In a parallel line of evolution (*Praebarrettia*, Fig. E238,3 to *Barrettia*, Fig. E237,10) the ligament support was drawn out gradually until it became narrowed into a moniliform ray.

MUSCLE SUPPORTS

Adductor muscle supports of calcite, the **myophores**, are relatively stable morphological characters in the rudists. In the conservative diceratids the muscles are attached to part of the cardinal area or connect with it (e.g., *Diceras*, Fig. E236,2). Modifications of this scheme occur in the requieniids. As the shape of the AV valve changed markedly to vasselike, the myophores shifted laterally (e.g., *Eoradiolites*, Fig. E237,2; *Hippurites*, Fig. E220,2), while the apophyses became massive, grooved, elongate downward projections of the FV (Fig. E220, 1a,3).

In the radiolitids the elongation of the FV myophores occurs early in the Cretaceous. *Agriopleura* (Fig. E237,1), a Barremian and younger genus slightly evolved from the monopleurids, has short massive apophyses. In younger radiolitids (e.g., *Praeradiolites*, Fig. E220,3, Albian) the apophyses are elongate smooth and displaced 90 degrees from the ligament crest (Fig. 238,4). Still younger genera evolved highly serrated lateral apophyses margins as can be seen in *Dubertretia* (Fig. E237,7) and *Dechaseauxia* (Fig. E239,1b). The earlier evolutionary imprint of twisting still may be manifest internally in highly specialized genera and where it is marked by

a greater extension of the anterior apophysis in the radiolitids (e.g., *Chiapasella*, Fig. E238,1) and hippuritids (e.g., *Praebarrettia*, Fig. E238,3; *Torreites*, E238,5).

Adductor muscle supports in the caprotinids and caprinids took an entirely different turn. In these families the development of marginal accessory cavities and canals between the body cavity and anterior wall provided a thin, long interior plate or wall for muscle attachment which can be seen in *Sellaea* (Fig. E239,3) and *Schiosia* (Fig. E239,2). In the caprotinids this wall became a projecting plate in *Polyconites* (Fig. E239,4), whereas in *Titanosarcolithes* (Fig. E262,4b) the posterior myophore was a raised apophysis in the FV that fits into an alveole in the AV.

WALL STRUCTURE

Wall structure is a useful basis for classification in rudist families and accordingly several trends are discussed here briefly.

The radiolitids, which characteristically have a celluloprismatic wall with a thick outer layer of hollow prisms, display a major trend toward less dense, more open network outer shell walls in younger taxa. Early genera (e.g., *Agriopleura*, Fig. E240, 5; *Sphaerulites*, Fig. E240,2) have a compact reticulate outer wall. Younger genera (e.g., *Eoradiolites*, Fig. E240,1) have a compact to loosely open rectangular network. *Sauvagesia* (Fig. E238,4), *Chiapasella* (Fig. E238,1) and others have an open polygonal network. More complicated mixed wall structure of compact and open occurs in some later Cretaceous genera (e.g., *Joufia*, Fig. E240,6, and *Colveraia*, Fig. E240,4). Wall structure appears to be a relatively

(Fig. E237).

5. *Caprinula* sp. (Caprinidae), U.Cret.(Cenoman.), Eu.; schematic transv. sec. FV (252).
6. *Vaccinites marticensis* DOUVILLÉ (Hippuritidae), U.Cret.(Coniac.), France; transv. sec. at level of AV with projecting teeth and posterior myophore of FV, $\times 1$ (910).
7. *Dubertretia kelleri* COX (Radiolitidae), U.Cret. (Maastricht.), Syria; transv. sec. at level of AV viewed toward commissure, $\times 0.9$ (468).
8. *Distefanella salmoiraghi* PARONA (Radiolitidae), U.Cret., Italy; transv. sec. AV, $\times 0.6$ (719).
9. *Durania cornupastoris* DES MOULINS (Radiolitidae), U.Cret.(Turon.), France; transv. sec. at level of AV with projecting teeth and myophores of FV, $\times 0.6$ (268).
10. *Barrettia monilifera* WOODWARD (Hippuritidae), U.Cret.(Campan.), W.Indies; transv. sec. AV, $\times 0.3$ (269).
11. *Batolites organisans* DE MONTFORT (Hippuritidae), U.Cret.(Santon.), S.France (Pyrénées, Rennes-les-Bains); transv. sec. at level of AV with projecting teeth and posterior myophore of FV, $\times 2$ (269).

[Explanation as for Fig. E220; also *Es*, *Ss*, pseudopillars.]




FIG. E238. Morphological features of rudists (ligament supports).

1. *Chiapasella cubensis* RUTTEN (Radiolitidae), U.Cret.(Maastricht.), Cuba; transv. sec. at level of AV with projecting teeth and myophores of FV, $\times 1.5$ (810).
2. *Tampsia bishopi* STEPHENSON (Radiolitidae), U. Cret.(?Maastricht.), Mexico, 2a, colony viewed from above, $\times 0.17$; 2b, transv. sec. ant. part of AV, $\times 1$ (886).
3. *Praebarrettia sparcilirata* (WHITFIELD) (Hippuritidae), U.Cret.(Maastricht.), Cuba; transv. sec. AV, $\times 0.3$ (67).
4. *Sauvagesia sharpei* (BAYLE) (Radiolitidae), U. Cret.(Turon.), Port.; transv. sec. at level of AV with projecting teeth and myophores of FV, $\times 0.6$ (268).
5. *Torreites sanchezi* (DOUVILLÉ) (Hippuritidae), U.Cret., Cuba; transv. sec. AV, $\times 0.6$ (281).
6. *Pironaea polystylus* (PIRONA) (Hippuritidae), U.Cret.(Maastricht.), Italy; transv. sec. at level of AV with projecting teeth and posterior myophore of FV, $\times 0.3$ (269).

[Explanation as for Fig. E220.]

conservative character which nevertheless has a certain adaptive potential.

Among the caprinids the pattern of marginal canal development in the shell wall shows a remarkable tendency to repeat a sequence of increasing complexity in several lines of caprinid evolution (MACGILLAVRY, 561). Early caprinid genera in different lines commonly have simple undivided or bifurcating radial plates that make a pattern of marginal pyriform canals. As the plates became increasingly polyfurcated,

rows of marginal polygonal canals appeared on the periphery of the shell. In the Early Cretaceous of the Western Hemisphere, marginal canal development can be traced in the sequence of genera *Amphitriscoelus*, *Planocaprina*, *Coalcomana*, and *Caprinuloidea* (Fig. E241), spanning the Aptian to late Albian. In the Eastern Hemisphere an analogous sequence of canal development occurs in mainly the Cenomanian genera *Caprina*, *Schiosia*, *Orthoptychus*, and *Sphaerucaprina* (Fig. E241). Similar de-




FIG. E239. Morphological features of rudists (muscle insertions).

1. *Dechaseauxia costata* TAVANI (Radiolitidae), U.Cret.(Maastricht.), Somalia; 1a, side view of AV, $\times 0.2$; 1b,c, transv. sec. AV, $\times 3.3$, $\times 4.6$ (904).
2. *Schiosia schiosensis* BÖHM (Caprinidae), U.Cret., Alps; schematic transv. sec. FV (Dechaseaux, after Parona).
3. *Sellaea orbignyi* DI STEFANO (Caprotinidae), U. Cret.(Cenoman.), Italy; 3a,b, schematic transv. secs. AV and FV (Dechaseaux, after di Stefano).

velopment of canal patterns in the line *Offneria* (Fig. E260,4)-*Caprinula* (Fig. E257,4b) and the line *Plagiopychus* (E260,2b)-*Mitrocaprina* (E260,3) confirm the highly adaptive nature of the development of multiple marginal canals by polyfurcation of radial plates. The superficial homeomorphy between genera such as *Sphaerucaprina* and *Caprinuloidea* or *Caprina* and *Planocaprina* was a source of taxonomic confusion among workers who first tried to identify Western Hemisphere rudists.

SIPHONAL STRUCTURES

External opening to the shallow sea of predators, contaminants, and fluctuating salinities and oxygen content is served by the siphons. Expected adaptations induced by environmental pressure are well displayed in siphonal structures of the radiolitids and hippuritids. In the radiolitids the siphonal positions are marked by bands or grooves in both valves; this is the usual morphological expression of the siphons. In a few genera of different subfamilies additional intramural siphonal structures are developed. YONGE's (1027a) stimulating comparison of chamid and rudist soft parts calls for a different interpretation of the position of siphonal openings than commonly accepted (KLINGHARDT, 475). YONGE recognized water currents and cleaning currents (Fig. E242) but did not place them necessarily as coincident with the position of the siphonal bands or the pillars *E* and *S*.

In the radiolitine genus *Medeella* (Fig. 243,7) the siphonal supports take the form of cylindrical structures within the AV outer wall. As continuation of this feature, cylindrical fossettes are developed in the AV and operculiform arches in the FV of *Medeella* (*Fossulites*) (Fig. E243,1). Similar protective structures for the FV siphons develop in the bioradiolitine genus *Thyrastylon* (Fig. E243,2). The AV bands in

(Continued.)

4. *Polyconites operculatus* ROULLAND (Caprotinidae), L.Cret., Eu.; interior FV (Dechaseaux, after 267b).

[Explanation as for Fig. E220.]




FIG. E240. Morphological features of rudists (Radiolitidae) (wall structure).

1. *Eoradiolites davidsoni* (HILL), L.Cret.(mid. Alb.), USA(Texas); part of transv. sec. showing AV inner and outer wall structure, $\times 2$ (Coogan, n).
2. *Sphaerulites patera* ARNAUD, U.Cret.(low Turon.), France; 2a, AV from above, $\times 0.3$; 2b, side view of AV siphonal bands, $\times 0.3$ (23).
3. *Hardaghia quadrata* TAVANI, U.Cret.(Maastricht.), Somalia; AV from above, $\times 0.7$ (904).
4. *Colveria variabilis* KLINGHARDT, U.Cret.(Maastricht.), Italy; transv. sec. FV near commissure viewed from above, $\times 0.5$ (475).
5. *Agriopleura blumenbachi* (STUDER), L.Cret.(Barrem.), Switz.; transv. sec. at level of AV with projecting teeth and myophores of FV, $\times 0.7$ (Pictet & Campiche, in 911).
6. *Joufia reticulata* BÖHM, U.Cret.(Maastricht.), Italy; transv. sec. at level of AV with projecting teeth and myophores of FV, $\times 0.2$ (853). [Explanation as for Fig. E220.]

this genus are deeply sunken and in the FV the brim of the valve forms oval oscules above the AV siphonal bands. The sub-family Lapeirousiinae is characterized by intramural siphonal structures. In the AV these structures generally consist of tubular bands of polygonal cells (e.g., *Vautrinia*, Fig. E243,5-6) called pseudopillars. In the FV there are oscules or openings for the siphons.

Finally, an analogous development of oscules occurs in the hippuritid genus *Yvaniella* (Fig. E243,3-4) where a tubercle on the FV surface is traversed by two large

canals. It is noteworthy that intramural siphonal structures are a Late Cretaceous phenomenon.


HIPPURITID PILLARS

The basic hippuritid stock, exemplified by *Hippurites*, has three extensions of the AV wall internally which are called pillars (Fig. E220,1b). The arrangement of the three pillars (usually lettered L, S (sortie) and E (entrée) following DOUVILLÉ carries through from Turonian to Maastrichtian in widespread and morphologically conserva-


EUROPE

NORTH AMERICA

VENTRAL
RADIAL PLATES




Sphaerucaprina (Cenoman.)




Caprinuloidea (mid. Alb.)

Tangential plates
cross radial plates




Orthoptychus (Cenoman.)




Coalcomana
(low. Alb.)

Trifurcating
radial plates




Schiosia (U. Cenoman.-Senon.)



Planocaprina (Apt.)

Bifurcating
radial plates



Caprina (Barrem.-Cenoman.)



Amphitriscoelus (Apt.)

Simple unbranched
radial plates

FIG. E241. Morphological features of rudists (wall structure). Marginal canals and ventral radial plates in four European and four North American caprinid genera; AV transverse sections, $\times 0.9$ (Coogan, n).




FIG. E242. Morphological features of rudists, AV interiors of *Hippurites* (1) and *Radiolites* (2) with indications of inhalant and exhalant water currents (full-line arrows) and cleaning currents (broken-line arrows); possible disposition of visceral mass and foot marked by light broken lines, teeth sockets solid black (1024a). [Explanation: *am*, *pm*, anterior and posterior myophores; *lig*, ligament.]

tive genera such as *Hippurites*, *Hippuritella* (Turon.-Maastricht., Fig. E263,4), and *Vaccinites* (Turon.-Maastricht., Fig. E237, 6). A trend toward increasing the number and length of pillars in both European and American lines of hippuritids is recognized. In the European line the increase in pillar number begins in *Batolites* (Fig. E237,11) which has numerous folds that do not project as far inward as the three pillars *L*, *S*, and *E*. In *Pironaea* (Campan.-Maastricht., Fig. E227,1), there are more than 12 pillars of lengths equal to those of *L*, *E*, and *S*. Many of these accessory pillars are elongate and reach 0.7 the distance to the center point of the AV. In the American line the marked increase in pillar num-

ber (15 to 20) occurs in *Praebarrettia* (Santon.-Maastricht., Fig. E238,3). The pillars extend toward the center and become narrow and beadlike (moniliform) in late growth stages. The final stage of this development occurs in *Barrettia* (Campan.-Maastricht.) where the AV is filled with 60 or more elongate beaded pillars or rays (Fig. E237,10). Lengthening of the pillars may occur independently of increase in pillar number as is evident in *Vaccinites* and *Torreites* (Fig. E238,5).

SYSTEMATIC DESCRIPTIONS

Family DICERATIDAE Dall, 1895

[Materials for this family prepared by COLETTE DECHASEAUX and A. H. COOGAN (subfamily divisions not recognized by DECHASEAUX, introduced by COOGAN)]

One valve or both hornlike, coiled, with umbo directed anteriorly and outward. Valves generally subequal, attached by RV or unequal, attached by LV with free valve more or less operculiform. Surface ornamented by striae or finely radiating ribs. AV with two unequal teeth, small one anterior and large one posterior, separated by horseshoe-shaped socket; FV with strong conical tooth bearing small socket on its anterior face, bordered behind by large socket. Anterior muscle inserted in each valve to shell wall, posterior muscle insertion on plate that projects above shell (except in Heterodiceratinae in which AV muscles were attached to elongations of cardinal platforms). Long ligamentary groove between beak and most posterior element of hinge. *U. Jur. (Oxford.)-L. Cret. (Valangin.)*.

This most ancient family of the rudists comprises genera or subgenera which are defined partly by mode of attachment of the shell. Fixation by the right valve is observable only in *Diceras*, whereas fixation is by the left valve in *Epidiceras* and *Heterodiceratas*.

Subfamily DICERATINAE Dall, 1895

[*nom. transl.* COOGAN, herein (*ex* Diceratidae DALL, 1895)]

Shell attached by RV (AV), which has large conical posterior tooth; LV (FV) tooth hook-shaped. *U. Jur. (U. Oxford.-L. Kimmeridg.)*.




FIG. E243. Morphological features of rudists (siphonal structures).

1. *Medeella (Fossilites) undaesaltus* ASTRE (Radiolitidae), U.Cret.(Coniac.), France (Aude); 1a, oblique side view of both valves, $\times 0.7$; 1b, side view of AV, $\times 0.7$; 1c, transv. sec. AV, $\times 0.7$ (24).
 2. *Thyrastylon adhaerens* (WHITFIELD) (Radiolitidae), U.Cret.(Maastricht.), Jamaica; 2a, adjoined shells, one at left showing oscules (So and Eo) in FV, $\times 0.7$; 2b, both valves, FV with Eo and So oscules, $\times 0.7$; 2c, transv. sec. AV, $\times 4$ (131).
 3. *Yvaniella maestrichtiensis* (MILOVANOVIC) (Hippuritidae), U.Cret.(Maastricht.), Yugosl.; vertical sec. FV and upper part of AV, $\times 0.7$ [c, canal of central excrescence; p, pore] (624).
 4. Same, E.Serbia; oblique view of both valves, $\times 0.6$ (628).
 - 5-6. *Vautrinia syriaca* (VAUTRIN) (Radiolitidae), U. Cret.(Maastricht.), Syria; 5, transv. sec. part of AV at level of pseudopillar, weathered, showing small columns in relief above prismatic tissue, $\times 0.7$ (933); 6a, lat. view AV, $\times 0.3$ (933); 6b, transv. sec. AV, $\times 0.5$ (933).
 7. *Medeella* sp. (Radiolitidae), U.Cret., France; transv. sec. AV showing part of cylindrical siphonal structure, $\times 0.6$ (24).
- [Explanation: c, canal; Eb, Sb, siphonal bands; Eo, So, oscules; Ep, Sp, internal pillars farthest and nearest L; L, ligament ridge; p, pore.]




FIG. E244. Diceratidae (Diceratinae) (4); (Heterodiceratinae) (1-3), (Epidiceratinae) (5) (p. N778-N779). [Explanation: *am*, *pm*, anterior and posterior myophores; 1, 3, anterior and posterior teeth of FV; tooth of AV; 1', 3', sockets in AV for reception of correspondingly numbered FV teeth; 2', socket for reception of tooth 2.]

Diceratidae LAMARCK, 1805, p. 299 [*D. arietinum*; M] [= *Diceratia* OKEN, 1815 (*nom. van.*) (ICZN rejected publ.); *Dicerata* RAFINESQUE, 1815 (*nom. van.*)]. Shell robust, valves subequal, both with partially coiled beaks. *U.Jur.*(*U.Oxford.-L.Kimmeridg.*), Eu.-N.Afr.—FIG. E244, 4. **D. arietinum*, Raurac., France (Saint-Mihiel); 4a, b (LV=FV), $\times 1$ (246). [Also Fig. E236, 1-2; E223, 1a, b.]

Subfamily HETERODICERATINAE Pchelintsev, 1959

[*nom. transl.* COOGAN, herein (*ex* Heterodiceratidae PCHELINTSEV, 1959)]

Valves large, inequivalve, attached by LV (AV); RV (FV) smaller, may be operculiform, posterior tooth in RV large, trapezoidal to triangular, projecting upward; LV (AV) tooth arcuate, muscles attached to elongation of cardinal platform. *U.Jur.* (*Tithon.*)-*L.Cret.* (*Valangin.*).

Heterodiceratidae MUNIER-CHALMAS in HEBERT, 1870, p. 116 [*Diceratid luci* DEFRANCE, 1819, p. 177;

M] [= *Pseudodiceratid* GEMMELLARO, 1876, p. 50 (type, *Diceratid carinatum* GEMMELLARO; SD COX herein)]. FV small, may be operculiform, both muscle insertions of FV on elongations of cardinal platform and curved outer edge of body cavity, not on prominent myophore. *U.Jur.* (*Tithon.*)-*L.Cret.* (*Valangin.*), Eu.-USSR.—FIG. E244, 1a. **H. luci* (DEFRANCE) *communis* (BÖHM), *Jur.* (Portland.), France; LV, $\times 0.5$ (246).—FIG. E244, 1b, c. *H. sp.*; diagram showing features of hinge; 1b, c, AV, FV (252).

Mesodiceratidae PCHELINTSEV, 1959, p. 82 [*M. enissalense*; OD]. Differs from *Heterodiceratid* in having operculiform FV and posterior muscle insertions of FV attached to cardinal platform by narrow subdental bridge. *U.Jur.* (*Oxford.-L.Kimmeridg.*), USSR (Crimea).—FIG. E244, 2. **M. enissalense*; int. view, 2a, b, AV, FV, $\times 0.7$ (722).

Paradiceratidae PCHELINTSEV, 1959, p. 96 [*Chama speciosa* MÜNSTER in GOLDFUSS, 1837, p. 205; OD]. FV not operculiform, posterior cardinal tooth trapezoidal to triangular, posterior muscle insertions of AV on elongation of cardinal platform, anterior separate. *U.Jur.* (*Tithon.*), Eu.-USSR

(Crimea).—FIG. E244,3. *P. alsuense* Pchelintsev; 3a, AV, $\times 0.7$; 3b, both valves diagram (722).

Subfamily PLESIODICERATINAE Pchelintsev, 1959

[nom. transl. COOGAN, herein (ex Plesiodiceratidae Pchelintsev, 1959)]

Shell irregular, attached by larger LV (AV). RV (FV) usually operculiform, beaks moderately curved, posterior muscle sertion of FV extending toward and behind posterior tooth, posterior cardinal tooth strongly curved, reaching laterally nearly across valve. *U. Jur. (Oxford.-Kimmeridg.)*.

Plesiodiceras MUNIER-CHALMAS, 1882, p. 478 [**Diceras valfinense* BÖHM, 1881, p. 160; OD]. FV with large cardinal and smaller but prominent anterior tooth; posterior muscle insertion on cardinal platform, extending behind posterior tooth. *U. Jur. (Up. Oxford.-Kimmeridg.)*, Eu.—FIG. E245,2a. *P. sp.*, diagram showing features of FV hinge (246).—FIG. E245,2b,c. *P. muensteri* (GOLDFUSS), Kimmeridg., France; 2b,c, AV, FV, $\times 1$ (246).

Eodiceras Pchelintsev, 1959, p. 37 [**Diceras ursicinum* THURMANN, 1852, p. 278; OD]. Shell small, moderately inequivalve; FV hinge primitive, posterior tooth large, transversely elongate subtriangular, anterior muscle insertion extending partly onto hinge area. *U. Jur. (Oxford.)*, Switz.-USSR.—FIG. E245,1b,c. **E. ursicinum* (THURMANN), Lusit., Crimea; 1b, ant., 1c, post., $\times 1$ (722).—FIG. E245,1a. *E. eximium* (BAYLE), Jur. (Kimmeridg.), Switz.; FV, $\times 1$ (de Loriol in 722).

Subfamily EPIDICERATINAE Rengarten, 1950

[nom. transl. COOGAN, herein (ex Epidiceratidae RENGARTEN, 1950)]

Shell with triangular to oval outline, weakly inequivalve, attached by LV (AV), dentition massive, close to megalodontid, cardinal area flattened, teeth and sockets set obliquely. *U. Jur. (Oxford.) - L. Cret. (Valangin.)*.

Epidiceras DOUVILLÉ, 1936, p. 332 [**Diceras sinistrum* DESHAYES, 1824, p. 466; SD DECHASEAUX, 1952, p. 326]. Shell large, posterior part of valves flattened, furrowed; FV operculiform or elevated, smaller than AV. *U. Jur. (Oxford.-Kimmeridg.)*, Eu.-USSR.—FIG. E244,5. **E. sinistrum* (DESHAYES), Raurac., France; FV, $\times 1$ (246).

Megadiceras Pchelintsev, 1959, p. 72 [**Diceras beyrichi porrecta* BÖHM, 1883; OD]. Much larger than *Epidiceras*, beaks drawn out, posterior muscle insertion in FV extended under cardinal area. *U. Jur. (Tithon.)-L. Cret. (Valangin.)*, Eu.-USSR.




FIG. E245. Diceratidae (Plesiodiceratinae) (p. N779). [Explanation: am, pm, anterior and posterior myophores; 1, 3, anterior and posterior teeth of FV; 2', socket in FV for reception of tooth 2 of AV.]

Family REQUIENIIDAE Douvillé, 1914

[=Bayleidae MUNIER-CHALMAS, 1873 (rejected under Code Art. 23,b)] [Materials for this family prepared by COLETTE DECHASEAUX and B. F. PERKINS]

Inequivalve, attached by coiled LV (AV) which invariably is larger than operculiform RV (FV); hinge of *Diceras* type, with single tooth in LV (AV) and two teeth in RV (FV), although toothlike thickening behind posterior socket of LV (AV) may be present in some shells (*Requienia*, *Matheronia*),




FIG. E246. Requieniidae (p. N781, N783). [Explanation: 1, 3, anterior and posterior teeth of FV; 2, tooth of AV.]

posterior tooth of RV (FV) generally more developed than anterior; siphonal bands more or less clearly present on posterior side of LV (AV); anterior muscle insertion on shell wall in LV (AV) and either on shell wall or prolongation of cardinal platform in RV (FV); posterior muscle inserted to shell wall in LV (AV) or on myophore plate beneath cardinal platform and on variously developed plate in RV (FV). *U. Jur. (Tithon.)-U. Cret. (Maastricht.)*.

Genera comprised by the family Requeniidae are distinguished from one another by relative size of the teeth and by form and position of the posterior muscle insertions. They retain some features of the Diceratidae, but new characters make their appearance in them, for example, so-called siphonal bands (*Requienia*, *Toucasia*) and accessory cavities (*Bayleia*). These occur also in other rudist families, siphonal bands being a constant feature of the Radiolitidae and accessory cavities constituting an essential attribute of the Caprotinidae.

Requienia MATHERON, 1843, p. 102 [**Chama ammonia* GOLDFUSS, 1837, p. 205; OD] [= *Requeniites* MATHERON, 1843 (obj.)]. Highly inequivalve; AV coiled in spiral of several whorls; FV forming flat spiral; ornament of growth wrinkles and fine radial ribs; sinuosities of growth lines on posterior side of AV define 2 siphonal bands; slight thickening behind posterior socket in AV of some species suggests tooth; AV muscle attachments on shell wall; FV anterior muscle insertion on shell wall, posterior muscle insertion on plate projecting above cardinal platform; pallial line entire. *L. Cret. (Valangin.)-U. Cret. (Senon.)*, Eu.-N. Afr.-S. Am.-N. Am.-E. Afr.—FIG. E247, 3. **R. ammonia* (GOLDFUSS), *L. Cret. (U. Barrem.)*, France (Brouzet); both valves, $\times 0.5$ (278a).—FIG. E246, 1. *R. sp.*, diagrams showing features of hinge, 1a, b, AV and FV int., $\times 0.7$ (252).

Apricardia GUÉRANGER, 1853, p. 36 [**A. carinata*; M]. External shell form as in *Toucasia*; AV hinge unknown; posterior tooth of FV large; posterior muscle insertion in both valves on plate passing below cardinal platform, plate of FV small and oblique to valve surface. *U. Cret. (Cenoman.-Senon.)*, Eu.-N. Afr.-E. Afr.-Syria-Mexico.—FIG. E246, 3. *A. sp.*, diagram showing features of FV hinge, $\times 0.7$ (252).

Bayleia MUNIER-CHALMAS, 1873, p. 74 [**B. pouechi*; OD]. External shell form, AV hinge and AV arched ligament ridge as in *Toucasia*; FV large coil; anterior muscle insertion on shell wall in both valves; AV posterior muscle insertion on plate

passing beneath cardinal platform; FV posterior muscle insertion on projecting plate thickened distally and attenuated proximally; FV with 3 small posterior accessory cavities, 1 below ligament and 2 others separated by muscle attachment plate. *U.Cret.(Campan.-Maastricht.)*, S. France-Alg.—FIG. E246,2. *Bayleia* sp., diagram showing features of hinge; 2a, AV; 2b, FV, $\times 0.7$ (252).

Bayleioidea PALMER, 1928, p. 35 [**B. clivi*; OD]. External shell form similar to that of *Toucasia*; AV loosely coiled, keeled; FV nonspiral semioperculiform; hinge greatly reduced; AV posterior muscle insertion on projecting plate, FV posterior muscle insertion on thin(?) plate separated from shell wall by accessory cavity; anterior muscle insertions unknown. *U.Cret.(Turon.)*, Mexico.—FIG. E246,4. **B. clivi*; both valves, $\times 0.7$ (Perkins, n; courtesy L. G. Hertlein).

Kugleria BOUWMAN, 1938, p. 41 [**Toucasia steinmanni* SCHNARRENBERGER, 1901, p. 20; SD PERKINS herein]. External shell form similar to that of *Toucasia*; AV usually carinate, spirally coiled; FV flattened with several spiral whorls; posterior tooth of FV large and strongly projecting, anterior tooth reduced; FV posterior muscle insertion on plate forming extension of cardinal platform, anterior muscle insertion larger than posterior. *U.Cret.(Cenoman.)*, Eu.(Italy)-W.Indies(Trinidad).—FIG. E246,6. *K. macgillavryi* BOUWMAN, Trinidad; 6a, FV int.; 6b, FV post. profile; 6c, FV ext., $\times 1$ (after 78).—FIG. E247,4. **K. steinmanni* (SCHNARRENBERGER), Italy(Appennines); ant. view of both valves, $\times 0.5$ (834a).

Matheronia MUNIER-CHALMAS, 1873, p. 74 [**Caprotina virginiae* GRAS, 1852, p. 32; OD]. Inequivalve, AV low spiral coil, FV operculiform or convex; AV with small posterior tooth; FV anterior tooth weak or absent, FV posterior tooth large and projecting; FV muscle insertions on prolongation of cardinal platform; shell wall thick and lamellose. *U.Jur.(Tithon.)-U.Cret.(Cenoman.)*, Eu.

M. (Matheronia). AV muscle insertions on shell wall except in earliest known species, *M. (M.) salevensis* JOUKOWSKY & FAVRE, Tithon., Alps, in which posterior AV muscle insertion on myophore plate of *Diceras* type. *L.Cret.(Valangin.)-U.Cret.(Cenoman.)* (but unknown *Hauteriv.-Alb.*), Eu.—FIG. E247,5a. *M. (M.) munieri* PAQUIER, *L.Cret.(Barrem.)*, France; ext. of both valves, $\times 0.5$ (278a).—FIG. E247,5b,c. **M. (M.) virginiae* (GRAS), *L.Cret.(Urgon.)*, France; 5b-c, AV and FV int., $\times 0.5$ (after 716b).

M. (Monnieria) PAQUIER, 1898, p. 343 [**Monnieria romani* PAQUIER; M]. AV commonly keeled, ornamented with concentric wrinkles in some species; AV muscle insertions on shell wall. *U.Jur.(Tithon.)*, S.France.—FIG. E247,2. **M. (M.) romani* (PAQUIER), France(Gard); 2a,b,

AV int., ext., $\times 0.75$; 2c,d, AV and FV int. (diagram.), $\times 0.4$ (after 716a).

M. (Hypelasma) PAQUIER, 1898, p. 846 [**Hypelasma collioti* PAQUIER, 1898; M]. Very inequivalve; AV anterior muscle insertion on shell wall, posterior muscle insertion on weakly developed plate passing under cardinal platform (*Diceras* type). *U.Jur.(Tithon.)*, Eu.(S.France).—FIG. E247,1. **M. (H.) collioti* (PAQUIER), France (Gard); 1a,b, both valves and AV int., $\times 0.75$ (after 716a).

Pseudotoucasia DOUVILLÉ, 1911, p. 82 [**Toucasia santanderensis* DOUVILLÉ, 1889, p. 632; M]. Posterior muscle insertion of FV on plate lying parallel to shell wall surface and attached to shell wall by plate normal to the surface; teeth as in *Toucasia*. *L.Cret.(Apt.)*, Eu.(Spain).—FIG. E249,1. **P. santanderensis* (DOUVILLÉ); oblique section RV showing elevated myophore plate, $\times 0.5$ (after 267c).

Toucasia MUNIER-CHALMAS, 1873, p. 74 [**Requienia carinata* MATHERON, 1843, p. 104; OD] [= *Caprotina* D'ORBIGNY, 1842 (suppression proposed) ICZN pend.]. Valves keeled, carinate, or frilled with shallow siphonal bands on posterior side of AV; *Diceras*-type hinge but AV tooth less projecting and AV anterior socket shallower than in *Diceras*; anterior muscle insertions on shell wall in both valves, slightly depressed in AV; FV posterior muscle insertion on plate projecting obliquely from shell wall, with upper edge free or attached to lower surface of cardinal platform; posterior part of AV posterior muscle insertion on shell wall, anterior part raised and forming plate which passes under cardinal platform, as in *Diceras*. *L.Cret.(Barrem.)-U.Cret.(Cenoman.)*, Eu.-N.Afr.-N.Am.—FIG. E246,5; E249,2. **T. carinata* (MATHERON); E246,5, both valves, $\times 0.7$ (Dechaseaux, n); E249,2a,b, AV and FV int., $\times 1.5$; E249,2c, FV transv. sec. showing post. myophore, $\times 0.5$ (E249,2a,b, after 716b; E249,2c, after 267c).—FIG. E248,1. *Toucasia* sp.; 1a,b, AV and FV int. (diagram) showing hinge features (252).

Family MONOPLEURIDAE Munier-Chalmas, 1873

[= Gyropleuridae MACGILLAVRY, 1935] [Materials for this family prepared by COLETTE DECHASEAUX and B. F. PERKINS]

Inequivalve (except *Valletia*); RV (AV) attached, larger; LV (FV) with two equal or unequal teeth; RV (AV) with single tooth; muscle insertions generally on extensions of cardinal platform. *L.Cret.(Valangin.)-U.Cret.(Maastricht.)*, *U.Eoc.*

Monopleura MATHERON, 1843, p. 105 [**M. varians*; SD KUTASSY, 1934, p. 113] [= *Dipilidia* MATHERON, 1843, p. 111 (type, *D. unisulcata*)]. AV




FIG. E247. Requieniidae (p. N781, N783).




FIG. E248. Requieniidae (p. N783). [Explanation: 1, 3, anterior and posterior teeth of FV; 2, tooth of AV.]

conical, spirally twisted; FV operculiform or coiled in low spiral; exterior ornamented with unequal ribs extending from apex to commissure; 2 teeth of FV conical, subequal; single tooth of AV oval in section, usually transverse to cardinal platform; ligamental groove arched according to degree of spiral twisting or coiling; muscle insertions in both valves on prolongations of cardinal platform. *L. Cret. (low. Valangin.) - U. Cret. (Maastricht.)*, Eu.-N.Am.-Jamaica.—FIG. E250, 3a; E252, 1. **M. varians*, Barrem., France (Brouzet); E250, 3a, both valves, $\times 1$; 252, 1a, b, AV and FV int., $\times 1$ (both after 278a).—FIG. 250, 3b. *M. marcida* WHITE, Alb., USA (Texas); dorsal view of both valves, $\times 1$ (Perkins, n).—FIG. E251, 3. *Monopleura* sp.; 3a, b, FV and AV int. (diagram.) showing hinge features (252).

Araeopleura Cox, 1965, p. 731 [*pro Stenopleura* Počta, 1889, p. 35 (*non* STEBBING, 1888)] [**Plagioptychus angustissimus* Počta, 1887, p. 204; SD KUTASSY, 1934, p. 127]. AV compressed coil, flattened along anterior-posterior axis; FV claviform; FV teeth weak, separated by deep socket, muscle insertions on thickened areas of shell wall extending from cardinal platform along margins of body cavity; AV teeth and muscle insertions unknown. *U. Cret. (Cenoman.)*, Eu. (Czech.).—FIG. E250, 2. **A. angustissimus* (Počta); 2a, b, AV ant., FV int., $\times 1$ (741c).

Gyropleura DOUVILLÉ, 1887, p. 768 [**Requienia cenomanensis* D'ORBIGNY, 1850, p. 261; OD]. AV exogyrid; FV operculiform or coiled in low spiral; teeth anterior, FV teeth subequal, small; AV posterior muscle insertion on plate posterior to posterior socket and forming extension of cardinal platform. *L. Cret. (Valangin.) - U. Cret. (Maastricht.)*, Eu.-Jamaica-N.Am. (Texas).—FIG. E250, 5. *Gyropleura* sp.; 5a, b, FV and AV int. (diagram.) showing hinge features (252).


Himeraelites DI STEFANO, 1889, p. 1 [**Monopleura (Himeraelites) vultur*; SD KUTASSY, 1934, p. 123]. AV conical, straight; FV curved, elevated,

beak overhanging AV; FV, teeth subequal, AV tooth round to oval in section; cardinal platform extended toward valve interior in both valves; FV posterior muscle insertion on exterior surface of elevated plate or lamina. *U. Cret. (Cenoman.)*, Eu. (Italy)-Mexico.—FIG. 252, 5a. **H. vultur* (DI STEFANO), Sicily; both valves, $\times 0.5$ (882a).—FIG. 252, 5b, c. *H. douvillei* (DI STEFANO), Sicily; 5b, c, AV and FV int., $\times 0.75$ (after 882a). ?*Paramonopleura* KOROBKOV in KOROBKOV & MAKARENKO, 1967, p. 135 [**P. ukrainica* KOROBKOV & MAKARENKO; OD]. Shell small; AV exogyrid, with cuneiform tooth, 1' circular, 3' divided by low ridge; FV operculiform, tooth 1 high, curved, tooth 3 marginal, biapical, socket deep. *U. Paleoc.*, Cherkasskaya Oblast, USSR.—FIG. E251, 1. **P. ukrainica*; 1a, b, FV and AV ext., $\times 2.7$; 1c, d, FV and AV int., $\times 5.3$ (Korobkov & Makarenko, 1967). [B. F. PERKINS. See additions, p. N868]

Petalodontia Počta, 1889, p. 61 [**Hippurites gemari* GEINITZ, 1840, p. 60; SD KÜHN, 1932, p. 121]. Differs from *Monopleura* in having FV anterior tooth much larger than posterior; FV muscle insertions on myophore plates which are joined to teeth and project around sides of body cavity. *U. Cret. (Cenoman.)*, Eu. (Czech.)-Mexico.—FIG. E250, 1. *P. planoperculata* (Počta),



FIG. E249. Requieniidae (p. N783).



Czech.; 1a,b, LV int. and dorsal view, $\times 1$ (741c).—FIG. E252,4. *P. felixi* DOUVILLÉ, Mexico (Coalcoman); 4a,b, transv. secs. parallel to commissure at level of AV with projecting teeth of FV myophore well below and close to it, $\times 0.5$ (after 270a).

Simacia POČTA, 1889, p. 39 [**S. minima*; M]. AV conical, curved or coiled, small; AV tooth broad, swollen; anterior socket rounded and deep, posterior socket lacking; FV unknown. *U.Cret.* (Cenoman.), Eu.(Czech.).—FIG. E252,2. **S. minima*; 2a,b, AV lat. and int. views, $\times 1.5$ (after 741c).

Valletia MUNIER-CHALMAS, 1873, p. 74 [**V. tombecki*; OD]. Externally similar to *Diceras*; FV anterior tooth much larger than posterior; muscle insertions in both valves on extensions of cardinal platform. *L.Cret.*(Valangin.), Eu.—FIG. E250,4; E252,3. **V. tombecki*, Alps; E250,4a,b, FV and AV int., $\times 1$ (Dechaseaux, n); E252,3a,b, AV and FV int., $\times 0.75$ (after 652a).—FIG. E251,2. *Valletia* sp.; FV and AV int. (diagram.) showing hinge features (252).

Family CAPROTINIDAE Gray, 1848

[*nom. correct.* GILL, 1871 (*pro* Caprotinidae GRAY, 1848)] [*=Heterocaprinidae* MUNIER-CHALMAS, 1873] [Materials for this family prepared by COLETTE DECHASEAUX and B. F. PERKINS]

Inequivalve; RV (AV) attached; LV (FV) with two subequal or unequal teeth; RV (AV) with single tooth; LV (FV) with cavities (accessory cavities) between shell wall and posterior muscle insertion and in some species between anterior muscle insertion and shell wall; similar but smaller cavities may be present in the AV. *L.Cret.* (Neocom.)-*U.Cret.*(Turon.).

Caprotina D'ORBIGNY, 1850, p. 236 [**Caprina striata* D'ORBIGNY, p. 170; ICZN pend.¹] AV conical, spirally twisted, anterior muscle insertion on extension of cardinal platform along shell wall, posterior muscle insertion on posterior surface of plate extending from cardinal platform to posterior shell wall, posterior accessory cavity shallow and undivided; FV conical, strongly coiled, teeth not well developed, anterior muscle insertion on extension of cardinal platform and posterior on plate, separated from shell wall by two accessory cavities. *L.Cret.*(Neocom.)-*U.Cret.*(Turon.), Eu.-

¹ The genus *Caprotina*, as erected by D'ORBIGNY in 1842, contained 10 nominal species none of which is included in the genus by modern authors, and it would be a senior synonym of *Toucasia* MUNIER-CHALMAS, 1873, by strict application of the International Code. It would only be retained in the sense in which it has come to be used by attributing it to D'ORBIGNY, 1850, and by declaring *C. striata* to be its type species, setting aside any prior designations. An application to this effect is being submitted to the International Commission. [Note by L. R. Cox.]

FIG. E250. Monopleuridae (p. N783-N785). [Explanation: 1, 3, anterior and posterior teeth of FV; 2, tooth of AV.]

N.Afr.—FIG. E253,2. *Caprotina* sp.; 2a,b, FV and AV int. (diagram.) showing hinge features (252).

Chaperia MUNIER-CHALMAS (1873, *nom. nud.*), 1882, p. 493 [**Caprina costata* D'ORBIGNY, 1839, p. 107; OD]. Differs from *Caprotina* in having slightly convex, operculiform FV; posterior accessory cavities of FV small and inconspicuous. *U. Cret. (Cenoman.)*, Eu. (S. France)-Mexico.—FIG. E253,1. **C. costata* (D'ORBIGNY), France; 1a,b, FV and AV int., $\times 2$; 1c, cluster of individual shells, $\times 1$ (695).

Horiopleura MUNIER-CHALMAS in DOUVILLÉ, 1889, p. 639 [**Monopleura lamberti* MUNIER-CHALMAS in DOUVILLÉ; SD PAQUIER, 1905, p. 56] [= *Oriopleura* DE LACVIVIER, 1884 (*nom. nud.*)]. AV exogyriiform, ornamented with radial ribs except on siphonal bands which are smooth; AV posterior muscle insertion on a thickened platform somewhat projecting over body cavity, normal to straight line formed by tooth, sockets, and anterior muscle insertion platform; FV operculiform with a narrow, shallow posterior accessory cavity. ?*L. Cret. (Barrem.)*, *L. Cret. (Apt.)*-*U. Cret. (Cenoman.)*, Eu.-N.Afr.-India-Mexico.—FIG. E254,2a,c. **H. lamberti* (MUNIER-CHALMAS), Alb., Spain (Pyrenees); 2a,b, AV int. and ventral ext., $\times 0.75$; 2c, AV vert. sec., $\times 0.4$ (2a,b, after 267d; 2c, after 561).—FIG. E254,2d. *H. baylei* (COQUAND), Apt., Spain (Pyrenees); FV int., $\times 0.75$ (after 267d).

Pachytraga PAQUIER, 1900, p. 337 [**Sphaerulites paradoxa* PICTET & CAMPICHE, 1869, p. 48; SD PAQUIER, 1905, p. 61]. AV conical, spirally twisted, tooth strongly projecting, posterior muscle insertion on posterior face of vertical plate connecting posteroventral side of tooth with posteroventral shell wall, posterior accessory cavity present, anterior muscle insertion on thickened area of shell wall, anterior accessory cavity may be present and divided into large canals by radial plates; FV conical, openly coiled, anterior tooth much larger than posterior, anterior muscle insertion on anterior face of plate connecting large anterior tooth with anteroventral shell wall, anterior accessory cavity simple and shallow or subdivided into large shallow canals by radial plates, posterior muscle insertion on shell wall, vertical plate connects ventral edge of anterior tooth with shell wall at ventral margin of posterior muscle insertion forming a large cavity connected to tooth socket but separated from shell cavity. *L. Cret. (Urgonian facies)*, Eu. (S. France-Switz.-Bulg.)-Jamaica.—FIG. E253,4; E254,3. **P. paradoxa* (PICTET & CAMPICHE); E253,4, FV int. (diagram.), $\times 0.7$ (Dechaseaux, n); E254,3a,b, AV sec. and ant. accessory cavities (canals), $\times 0.5$ (716c).




FIG. E251. Monopleuridae (p. N783-N785). [Explanation: 1, 3, anterior and posterior teeth of FV; 2, tooth of AV.]

Polyconites ROULLAND, 1830, p. 166 (genus without nominal species) [**P. operculatus* DOUVILLÉ, 1887, p. 778 (= *Radiolites polyconilites* D'ORBIGNY, 1842, p. 181); SM] [= *Polyconilites* D'ORBIGNY, 1842 (*nom. van.*); *Heterocaprina* MUNIER-CHALMAS, 1873 (obj.)]. AV conical, resembling *Monopleura*, tooth transverse, muscle insertions on thickened areas of shell wall, no accessory cavities; FV operculiform, anterior muscle insertion extends from cardinal platform along the shell wall, posterior muscle insertion on a recumbent plate projecting posteriorly from cardinal platform, several accessory cavities as in *Caprotina*. *L. Cret. (Apt.)*-*U. Cret. (Cenoman.)*, Eu.-Syria.—FIG. E254,1. **P. operculatus* (DOUVILLÉ), Cenoman., France; 1a,b, FV and AV int., $\times 0.75$ (after 267b). [Also, Fig. E229,4.]

Praecaprotina YABE & NAGAO, 1926, p. 126 [**Horiopleura yaegashii* YEHARA, 1920, p. 39; OD]. AV




FIG. E252. Monopleuridae (p. N783-N785). [Explanation: *am*, *pm*, anterior and posterior myophores; 1, 3, anterior and posterior teeth of FV; 2, AV tooth.]

irregularly conical, tooth strongly projecting, anterior muscle insertion on extension of cardinal platform on thickened area of shell wall, posterior muscle insertion extends as erect plate from tooth to posteroventral shell wall, large posterior accessory cavity may be divided in early stages by radial plates; FV convex, teeth subequal, elongate anterior muscle insertion extends from cardinal platform along shell wall and is separated from it by shallow groove, posterior muscle insertion on thin nearly vertical plate, large posterior accessory cavity extends from dorsal to ventral border, vertical plate connects ventral edge of anterior tooth with shell wall at ventral margin of posterior muscle insertion forming large cavity connected to socket but separated from shell cavity as in *Pachytraga*. *L.Cret.*, Japan.—FIG. E255,3. **P. yaegashii* (YEHARA); 3a,b, AV and FV int., $\times 0.75$ (after 1009a).

Retha COX, 1965, p. 731 [*pro Ethra* MATHERON, 1878 (*non* LAPORTE, 1833)] [**Ethra munieri* MATHERON; SD PAQUIER, 1905, p. 59]. Valves in elongate loose spiral, FV larger, AV with deep ligamental furrow; AV tooth slightly projecting, anterior muscle insertion on thickened area of shell wall, posterior muscle insertion on posterior face of plate extending from cardinal platform to posteroventral shell wall, posterior accessory cavity wide and shallow; FV muscle insertions on thickened areas of shell wall extending from cardinal platform, no accessory cavities, vertical plate connects posteroventral edge of anterior tooth with shell wall at ventral margin of posterior muscle attachment forming large cavity connected to socket but separated from shell cavity as in *Pachytraga*. *L.Cret.*(*Urgonian facies*), Eu.(S.France). —FIG. E253,3; E255,1. **R. munieri* (MATHERON); E253,3a,b, FV and AV int., $\times 1$ (E253,3a, Dechaseaux after Paquier; E253,3b, 716c); E255, 1, both valves, $\times 0.5$ (600b).

Sellaea DI STEFANO, 1889, p. 20 [**Caprotina zitteli* DI STEFANO, 1889, p. 28; SD KUTASSY, 1934, p. 142]. Both valves conical, AV straight, FV coiled; AV anterior muscle insertion on extension of cardinal platform and separated from shell wall by large accessory cavity divided by vertical radial plates into small cavities which in turn may be subdivided into small canals by 2 sets of vertical plates lying parallel to the shell wall, posterior myophore plate extending from tooth to posterior shell wall; FV with 1 elongate deep anterior accessory cavity and 2 smaller but deep posterior accessory cavities. *L.Cret.*(*Albian*)-*U.Cret.*(*Cenoman.*), Sicily-Apennines-USA(Texas)-Mexico.—FIG. E255,2a,b. **S. zitteli* (DI STEFANO),




FIG. E253. Caprotinidae (p. N785, N787). [Explanation: 1, 3, anterior and posterior teeth of FV; tooth 2 of AV; 2', socket of FV corresponding to tooth 2 of AV; am, pm, anterior and posterior myophores.]

Cenoman., Sicily; 2a,b, FV int., AV sec., $\times 0.75$ (after 882a).—FIG. E255,2c. *Sellaea* sp., Alb., USA(Texas), both valves, post. ext., $\times 0.5$ (Perkins, n).

Family CAPRINIDAE d'Orbigny, 1850

[=Caprinellidae GILL, 1871; Trechmannellidae Cox, 1933; incl. Anomoptychidae Vokes, 1967] [Materials for this family prepared by COLETTE DECHASEAUX and B. F. PERKINS with acknowledgment of assistance by L. R. COX]

Inequivalve, RV (AV) attached, some species attaining large size; teeth and myophores well developed, massive in some species; accessory cavities present in most




FIG. E254. Caprotinidae (p. N785, N787).




FIG. E255. Caprotinidae (p. N787, N789).

genera; pallial canals in pallial region of one or both valves. *L.Cret.(Urgonian facies)*-*U.Cret.(Maastricht.)*.

In the Caprinidae generic distinctions have been based mainly upon the form and disposition of the pallial canals and accessory cavities. Both the canals and accessory cavities may vary in shape and arrangement, depending upon the distance from the apex that a section is made, and such variations rarely have been taken into account in the definition of caprinid genera. Many of the presently rather arbitrarily defined caprinid genera must be restudied before a phylogenetically sound classification of the assemblage can be derived.

[In original manuscript (modified inappreciably by L. R. COX) DECHASEAUX divided caprinid genera into four informal groups, as follows: 1) forms considered to be most typical representatives of the family (*Caprina*, *Caprinula*, *Neocaprina*, *Orthoptychus*, *Paracaprinula*, *Plagioptrychus*, *Praecaprina*, *Sphaerucaprina*), 2) variants possibly representing expression of regional differentiation (*Amphitriscoelus*, *Antillocaprina*, *Caprinuloidea*, *Coalcomana*, *Kipia*, *Mitrocaprina*, *Offneria*, *Planocaprina*, *Sabina*, *Schiosia*), 3) forms considered to exhibit approach to the Radiolitidae (*Coralliochama*, *Dictyoptrychus*, *Ichthyosarcolites*, *Titanosarcolites*), and 4) imperfectly known

forms questionably assignable to the Caprinidae only because of the presence of accessory cavities or canals (*Bicornucopina*, *Cryptaulia*, *Immanitas*, *Lithocalamus*, *Palus*). In my opinion and that of PERKINS, it is sufficient simply to record the grouping of genera as listed and preferably to arrange systematic descriptions alphabetically after the type genus. Group 4 of DECHASEAUX is given under the heading "Superfamily and Family Uncertain."—EDITOR.]

Caprina C. D'ORBIGNY, 1822, p. 105 [**C. adversa*; SD PAQUIER, 1905, p. 69] [= *Gemellaria* MUNIER-CHALMAS, 1873, p. 75 (type, *Caprina communis* GEMMELLARO, 1865); *Cornucaprina* FUTTERER, 1892, p. 87 (type, *Schiosia carinata* BÖHM, 1892)]. AV conical, straight, smaller than FV, which is conical and strongly coiled; AV with projecting tooth anterior to ligamental groove, anterior muscle insertion on thickened area of shell wall, posterior insertion on subrect plate separated from shell wall by accessory cavity divided into smaller cavities by simple radial vertical plates; FV anterior muscle insertion on plate connecting anterior tooth with shell wall, large anterior accessory cavity divided by thin, vertical radial and transverse plates, posterior muscle attachment on shell wall, vertical plate connects anteroventral edge of anterior tooth with ventral shell wall forming a large cavity separated from body cavity; FV with one or two series of pyriform pallial canals in anterior, ventral and posterior regions. *L.Cret. (Urgonian facies)-U.Cret. (Cenoman.)*, Eu.-N.Afr.-N.Am.—FIG. E256,1; E257,2. **C. adversa*, Cenoman., France; E256,1, FV sec., $\times 0.8$; E257,2a, AV sec., $\times 0.5$; E257,2b, both valves, $\times 0.11$ (E256,1; E257,2a, after 267c; E257,2b, 695).

Amphitriscoelus HARRIS & HODSON, 1922, p. 130 [**A. waringi*; M]. Both valves elongate, incurved but not coiled, small; AV tooth narrow, elongate in section, anterior socket large, posterior socket small but joined to large accessory cavity which is undivided, pallial canals more or less rectangular or rounded-triangular limited to anterior, dorsal and posterior regions; FV with small anterior accessory cavity, vertical wall joins anterior tooth with anteroventral border of posterior muscle insertion forming cavity separate from shell cavity, pallial canals same shape and arrangement as in AV. *L.Cret. (Apt.)*, Trinidad-USA (Texas).—FIG. E257,1. **A. waringi*, Trinidad; 1a,b, AV sec., viewed from apex so that posterior is on left, and FV sec., $\times 0.7$ (Perkins, n).

Antilocaprina TRECHMANN, 1924, p. 407 [**Caprinella occidentalis* WHITFIELD, 1897, p. 193; M¹]. AV more or less conical, FV with coiled beak; FV with conical anterior tooth and crescentic or sigmoidal posterior tooth; internal ligamental groove; no accessory cavities; both valves with numerous small, mostly tabulate, polygonal or oval canals in inner shell wall layers, in teeth and in myophorous plates. *U.Cret. (U.Senon.-Maastricht.)*, W. Indies (Antilles).—FIG. E257,3. **A. occidentalis* (WHITFIELD), Maastricht., Jamaica; 3a, both valves, $\times 0.2$; 3b,c, AV oblique view and FV int., $\times 0.3$ (911a).

Caprinula D'ORBIGNY, 1847, p. 269 (genus without nominal species); 1850, p. 187 [**Caprina boissyi* D'ORBIGNY, 1839, p. 169; SM]. AV conical, straight; FV conical, loosely coiled; muscle insertions as in *Caprina*; anterior and posterior accessory cavities in both valves and divided by thin radial vertical plates; vertical plate divides cavity joined to socket from shell cavity as in *Caprina*; pallial canals include inner polygonal series and outer pyriform series, polygonal series decreases in diameter from interior of valves outward. *L.Cret. (Alb.)-U.Cret. (Cenoman.-Turon.)*, Eu.-Syria-N.Am.—FIG. E257,4. **C. boissyi* (D'ORBIGNY), Turon., France; 4a,b, FV secs. near commissure and nearer apex, $\times 1$; 4c, AV sec. near commissure, $\times 1$ (after 267c); 4d, both valves, $\times 0.5$ (695).

Caprinuloidea PALMER, 1928, p. 59 [**C. perfecta*; OD]. AV greatly elongated, straight to curved; FV much smaller, open to tightly coiled; teeth in both valves large and projecting, those of FV commonly strongly curved; AV tooth, sockets, and posterior accessory cavity similar to those of *Amphitriscoelus*; FV with vertical plate joining anterior tooth with ventral shell wall forming cavity separated from body cavity, no accessory cavities; pallial canals in both valves throughout shell wall, generally comprise inner polygonal series and outer pyriform series, rounded or polygonal canals may be present in myophorous plates and in teeth; shell cavity, tooth sockets, accessory cavity, and pallial canals may be tabulate. *L.Cret. (Alb.)*, Mexico-USA (Texas)-W. Indies (Cuba-Jamaica).—FIG. E258,3. **C. perfecta*, Mexico; FV int. (plate joining ant. tooth with shell wall broken), $\times 0.7$ (Perkins, n). [Also Fig. E224,3.]

Coalcomana HARRIS & HODSON, 1922, p. 132 [**Caprina ramosa* BÖHM, 1898, p. 327; M]. Resembles *Caprinuloidea* in external form and internal structures; differs in having pyriform pallial canals only. *L.Cret. (Alb.)-U.Cret. (Cenoman.)*,

¹ The genus was founded on specimens misidentified by TRECHMANN as *Caprinella occidentalis* and which Dr. L. J. CHUBB is describing as a new species, but the conception of the genus is not affected. [Note by L. R. Cox.]




FIG. E256. Caprinidae (p. N790, N795).




FIG. E257. Caprinidae (p. N790, N792-N793).

USA (Texas) - Mexico-W. Indies (Cuba).—FIG. E258,1. **C. ramosa* (BÖHM), Cenoman., Mexico (Coalcoman); 1a,b, FV and AV secs., $\times 0.8$ (after 270a).

Coralliochama WHITE, 1885, p. 9 [**C. orcutti*; M]. Inequivalve; AV conical, tall; FV convex, beak broad and incurved; ligament marked by inflection of outer shell wall layer into inner layer; AV muscle insertions on shell wall, no accessory cavities; AV shell wall with small poly-

gonal, closely tabulate canals throughout; FV anterior tooth joined to ventral shell wall by vertical plate as in *Plagiptychus*, no accessory cavities; FV shell wall with wide inner area of polygonal canals and marginal series of radially elongated pyriform canals, closely tabulate as in AV. *U.Cret. (Senon.)*, USA (Calif.)-Mexico.—FIG. E258,2. **C. orcutti*, Mexico (Baja Calif.); E258,2a, both valves, $\times 0.3$; E258,2b,c, AV and FV transv. secs., $\times 0.7$ (Perkins, n).




FIG. E258. Caprinidae (p. N792-N795).




FIG. E259. Caprinidae (p. N795-N797). [Explanation: *am*, anterior myophore; *L*, ligamental ridge.]

- Dictyoptychus** DOUVILLÉ, 1905, p. 178 [*pro Polyptychus* DOUVILLÉ, 1904, p. 248 (*non* HUEBNER, 1816)] [**Polyptychus morgani* DOUVILLÉ, 1904; M] [= *Anomoptychus* DOUVILLÉ, 1906, p. 56 (obj.); *Trechmanella* COX, 1933, p. 388 (obj.)]. Very large, inequivalve; AV conical, slightly incurved, with 2 longitudinal swellings possibly representing siphonal bands; FV much depressed, with eccentric summit; ligamental structures absent, hinge teeth produced, myophorous plates elongate; FV with large accessory cavity behind anterior myophore; AV with accessory cavities outside cardinal structures; AV with thick compact external layer and internal layer with narrow elongate canals separated by bifurcate radial plates as in *Plagioptychus*. *U.Cret.(U.Senon.)*, SW.Asia (Iran).—FIG. E256,2. *D. persica* (COX); 2a, both valves, $\times 0.16$; 2b, FV transv. sec., $\times 0.5$; 2c, transv. sec. at level of AV with projecting teeth and myophores (inner black areas) of FV, $\times 0.5$ (2a, 176; 2b,c, after 176).
- Ichthyosarcolithus** DESMAREST, 1817, p. 51 [**I. triangularis*; M] [= *Ichthyosarcolithus* HERMANNSEN, 1846 (*nom. van.*); *Caprinella* D'ORBIGNY, 1847, p. 269 (obj.)]. Very large, inequivalve, strongly carinate; AV curved or loosely coiled with whorls not in contact; FV curved in same direction as AV, smaller; teeth of FV fused with myophores to form single plate which fits into notches in AV shell cavity wall, radiolitic fashion; no ligament; shell cavity obliquely tabulate with tabulae of AV sloping away from commissure toward convex side and those of FV sloping away from commissure toward concave side; both valves with small, rounded-polygonal canals irregularly distributed and few large round canals in some areas, canals not tabulate. *L.Cret.(Urgonian facies)-U.Cret.(Turon.)*, Eu.-N.Afr.-?Iran-Canada-Cuba.—FIG. E258,4. **I. triangularis*, Cenoman.; 4a, both valves, $\times 0.2$; 4b, transv. sec., $\times 0.3$ (695).
- Kipia** HARRIS & HODSON, 1922, p. 133 [**K. trinitaria*; M]. AV conical, short; FV capuloid; both valves with rectangular pallial canals formed by simple, thin radial plates and restricted to anterior and ventral shell wall, which is thin where pallial canals are lacking, myophores imperfectly known. *L.Cret.(Apt.)*, W.Indies (Trinidad).—FIG. E259, 1. **K. trinitaria*; 1a,b, FV from above (shell wall eroded) and transv. sec., $\times 1.5$ (388a).
- Mitrocaprina** BÖHM, 1895, p. 102 [**Coralliochama bayani* DOUVILLÉ, 1888, p. 725; OD]. Very inequivalve; AV smaller, more or less exogyiform, attachment area very large; FV slightly to highly elevated, coiled; AV posterior muscle insertion on obliquely erect plate, posterior accessory cavity present and not always separated from posterior tooth socket, no pallial canals; FV teeth projecting slightly, no accessory cavities, pallial canals comprise 1 or 2 inner rows of polygonal canals and 1 or more outer rows of pyriform or radially elongate canals. *U.Cret.(Turon.-Maastricht.)*, Eu.-W. Indies (Jamaica-Cuba).—FIG. E259,2a. **M. bayani* (DOUVILLÉ), Turon., France; FV int., $\times 0.7$ (267c).—FIG. E259,2b,c. *M. vidaldi* DOUVILLÉ, Turon., Spain; 2b,c, FV and AV int., $\times 1$ (272).—FIG. E260,3. *M. tschoppi* (Palmer), Maastricht., Cuba; transv. sec., at level of FV with projecting tooth of AV, $\times 0.5$ (561).
- Neocaprina** PLENIČAR, 1961, p. 41 [**N. nanosi*; OD]. Known only from transverse sections of FV; anterior muscle insertion larger than posterior; anterior and posterior accessory cavities divided into smaller rectangular or oval cavities by thin radial vertical plates; pallial canals rectangular in section and formed by simple radial plates which bifurcate near outer shell wall anteriorly and posteriorly. *U.Cret.(Cenoman.-Turon.)*, Eu. (Yugosl.).—FIG. E259,6. **N. nanosi*, Turon.; 6a,b, FV transv. sec. and hinge area sec., $\times 1$ (741b).
- Offneria** PAQUIER, 1905, p. 82 [**O. rhodanica*; OD]. External form resembling *Praeacaprina*; AV conical, FV curved; cardinal apparatus and myophore arrangement similar to those of *Praeacaprina*; AV with elongate, curved posterior accessory cavity divided by thin radial plates and in some species by discontinuous plate parallel to shell wall, anterior accessory cavity divided similarly, marginal canals when present in ventral areas indistinguishable from subdivided accessory cavities; FV with marginal canals usually around entire valve but may be missing in ventral area, polygonal and pyriform canals present but arrangement irregular. *L.Cret.(Urgonian facies)*, Eu. (S.France).—FIG. E260,4a,b. **O. rhodanica*; 4a,b, FV and AV transv. sec., $\times 0.75$ (after 716c).—FIG. E260, 4c. *O. interrupta* PAQUIER; FV transv. sec., $\times 0.75$ (after 716c).
- Orthoptychus** FUTTERER, 1892, p. 91 [**O. striatus*; M]. AV conical, small; FV strongly inflated with beak incurved; distinguished from *Sphaerucaprina* by presence in FV posterior shell wall of some septa transverse to radial plates. *U.Cret.(Cenoman.)*, Eu. (Alps-Apennines).—FIG. E260, 1. **O. striatus*; 1a,b, FV and AV int., $\times 0.5$ (332a).
- Paracaprlnula** PIVETEAU, 1939, p. 34 [**P. syriaca*; M]. Resembles *Plagioptychus* externally but less inequivalve; anterior pallial canals polygonal as in *Caprinula*, posterior canals of very unequal size limited by more or less oblique transverse plates. *U.Cret.(Maastricht.)*, SW.Asia (Syria).—FIG. E259,4. **P. syriaca*; 4a, both valves, $\times 0.2$; 4b, shell wall transv. sec., $\times 2$ (741a).
- Plagioptychus** MATHERON, 1843, p. 114 [**P. paradoxus*; SD KUTASSY, 1934, p. 172]. AV conical and straight or low and twisted; external ligamental groove; FV strongly convex with recurved beak; AV tooth erect and strongly projecting, muscle insertions on shell wall, no pallial canals;

FV muscle insertions on thickened plates projecting above cardinal platform, anterior tooth smaller than posterior, separated by deep socket, vertical plate extending from anterior tooth to ventral shell wall forming cavity separate from

shell cavity as in *Caprina*, pallial canals comprise single series of pyriform canals in oldest species but in other species radial plates branch, forming several series of more or less pyriform canals, no accessory cavities. *U.Cret. (Cenoman.-Maastricht.)*,




FIG. E260. Caprinidae (p. N795-N797). [Explanation: 2, tooth of AV.]

Eu.-N. Afr.-Mexico-W.Indies (Jamaica-Cuba). — FIG. E259,5. *P. arnaudi* DOUVILLÉ, Turon., France; 5a,b, AV and FV int., $\times 0.7$ (267c).—FIG. E260,2a. **P. paradoxus*, Turon., France; both valves, $\times 0.4$ (600a).—FIG. E260,2b. *P. toucasianus* MATHERON, Turon., France; FV transv. sec., $\times 0.5$ (after 267c).

Planocaprina PALMER, 1928, p. 64 [**P. trapezoides*; OD]. AV conical, straight to slightly curved; FV larger, coiled in open spiral; AV tooth marginal, posterior accessory cavity large and joined to posterior socket as in *Amphitriscoelus*; FV anterior tooth large, curved and marginal, posterior tooth rudimentary; pyriform pallial canals occurring in single series in both valves; polygonal canals absent. ?*L.Cret.(Alb.)-U.Cret.(Cenoman.)*, Mexico-?USA (Texas).—FIG. E259, 3. **P. trapezoides*, Mexico; 3a, both valves, $\times 0.3$; 3b, AV transv. sec., $\times 1.3$; 3c, FV int., $\times 0.7$ (3a, Dechaseaux after 714; 3b,c, Perkins, n).

Praecaprina PAQUIER, 1905, p. 72 [**P. varians*; OD]. External form resembling *Caprina* but much smaller, FV coiling rarely more than 1 whorl, ventral region of both valves marked by longitudinal depression bordered by 2 swellings; AV tooth strongly projecting, both muscle insertions on plates projecting from cardinal platform and separated from shell wall by accessory cavities usually divided into smaller cavities by vertical radial plates; FV anterior tooth larger, erect and median in position, muscle insertions on shell wall and extend from cardinal platform, vertical plate connects anterior tooth with ventral shell wall as in *Caprina*, rounded or oval pallial canals formed by simple radial plates in anterior, posterior and posteroventral areas. *L.Cret.(Urgonian facies)*, Eu.-W.Indies (Trinidad).—FIG. E261,2; E262,2. **P. varians*, France; E261,2a, both valves, $\times 0.5$; E261,2b,c, FV and AV int., $\times 0.7$; E262,2, FV transv. sec., $\times 0.75$ (716c).

Rousselia DOUVILLÉ, 1898, p. 151 [**R. guilhoti*; M]. AV straight, conical; FV strongly convex; general form and hinge resemble *Monopleura*; FV anterior tooth large, crescentic, near shell cavity, posterior tooth small, marginal; FV muscle insertions on raised thickened areas of shell wall; no ligamental groove or cavity; AV with rounded to polygonal, tabulate canals in some marginal areas. *U.Cret.(Maastricht.)*, Eu.(Pyrenees).—FIG. E261,1. **R. guilhoti*, Spain (Lerida); AV int., $\times 1$ (272).

Sabinia PARONA, 1909, p. 303 [**S. aniensis*; SD KUTASSY, 1934, p. 169]. Externally resembling *Plagioptychus*; both valves without accessory cavities; AV pallial canals polygonal to rounded throughout shell wall, few pyriform canals in outer part but not forming continuous series; FV pallial canals comprise an inner wide area of polygonal canals of nearly uniform size and outer marginal row of pyriform canals. *U.Cret.(Senon.)*,




FIG. E261. Caprinidae (p. N797).

Eu.(Apennines-W.Serbia-Turkey).—FIG. E262, 3a. **S. aniensis*, Maastricht., Apennines; FV sec., $\times 0.75$ (after 719b).—FIG. E262,3b. *S. sublacensis* PARONA, Maastricht., Apennines; AV transv. sec., $\times 0.75$ (after 719b).

Schiosia BÖHM, 1892, p. 144 [**S. schiosensis*; OD]. Very inequivalve; AV low, strongly curved; FV larger, elongate, coiled; accessory cavities present in FV and possibly also in AV; FV pallial canals include single pyriform series in anterior, ventral and posterior areas, few polygonal and rounded canals in hinge area; AV interior not known. *U. Cret. (Cenoman.-Senon.)*, Eu.(S. Alps)-NW.Asia Minor (Bithynia).—FIG. E262,5. **S. schiosensis*, Turon., S.Alps; FV transv. sec., $\times 0.75$ (after 719a).


Sphaerucaprina GEMMELLARO, 1865, p. 212 [**S. woodwardi*; M] [= *Sphaerocaprina* DOUVILLÉ,

1910 (*nom. van.*)]. AV resembling that of *Caprina*; FV strongly convex, caplike, beak incurved but not coiled; AV anterior muscle inser-


tion on shell wall, posterior on curved vertical plate extending from tooth to ventral margin, large posterior accessory cavity, no pallial canals;



Sphaerucaprina





Praecaprina



Sabinia



Schiosia



Titanosarcolites

FIG. E262. Caprinidae (p. N797-N799).

Hippuritoida—Hippuritacea—Systematic Descriptions



FV with dorsal and anterior accessory cavities, vertical wall connecting anterior tooth to ventral shell wall as in *Caprina*, anterior, ventral and posterior areas with 1 or 2 rows of polygonal pallial canals and outer row of pyriform pallial canals. *L.Cret.(Alb.)-U.Cret.(Cenoman.)*, Eu.-N. Am.-W. Indies (Jamaica).—FIG. E262,1. * *S. woodwardi*, Cenoman., Italy; 1a,b, AV and FV transv. secs., $\times 0.75$ (after 274a).

Titanosarcolites TRECHMANN, 1924, p. 397 [**Caprinella gigantea* WHITFIELD, 1897, p. 194; M] [= *Diatretus* DOUVILLÉ, 1926, p. 132 (obj.)]. Very large, subequivalve, with both valves curved or coiled; dorsal and anterior surfaces with rounded longitudinal ridges separated by angular furrows, posterior and ventral surfaces with angular ridges or flanges separated by rounded grooves; ligamental groove internal, comma-shaped in transverse section; cardinal apparatus as in *Antillocaprina*, tooth alveoles tabulate; posterior muscle insertion in FV on raised apophysis which fits into alveole of AV, anterior muscle insertion superficial; shell cavity small, tabulate; small polygonal canals throughout shell wall, teeth and myophores; large tubular canals, rounded, oval or reniform, present on anteroventral side only. *U. Cret.(Maastricht.)*, USA (Texas)-Mexico-W. Indies (Jamaica-Cuba-Puerto Rico-Virgin Is.).—FIG. E262,4a. * *T. giganteus* (WHITFIELD), Jamaica; AV ext., $\times 0.35$ (980a).—FIG. E262,4b. *Titanosarcolites* sp.; AV transv. sec., $\times 0.75$ (Perkins, n).

Family HIPPURITIDAE Gray, 1848

[Materials for this family prepared by COLETTE DECHASEAUX and A. H. COOGAN, with acknowledgment of assistance by L. R. COX]

Shell conical to elongate-cylindrical and inequivalve. RV (AV) larger than LV (FV) which is operculiform, slightly convex or flat. Internally RV (AV) shell wall has compact smooth inner layer and thicker outer layer which extends toward interior in three or more folds (ligamental ridge and two or more pillars or rays). Pillar nearer ligamental ridge is designated as "S," using the notation devised by DOUVILLÉ and accepted by most authors, and the other (generally more elongate) as "E" (Fig. E237, 11). Teeth and myophore well developed. H-shaped (in transv. sec.) tooth stands between two sockets, anterior one receiving anterior tooth of LV (FV), and between it and inner wall of valve is an accessory cavity. On posterior side of RV (AV), between tooth and pillar *Sp*, are two sockets, that nearer tooth 2 for reception of posterior tooth of LV (AV), and that nearer

Sp is for posterior myophore of LV (FV) (Fig. E220,2). Posterior muscle insertion in RV (AV) on side of socket dorsal to *Sp*, and anterior muscle insertion marked by superficial scar between anterior socket and pillar *Ep*. Cavities or canals very rare in RV (AV). LV (FV) may be porous and commonly has two oval openings (oscles) (Fig. E263,2b). In longitudinal section RV (AV) in genera of this family shows curved partitions with concave side facing upward toward living chamber. *U. Cret.(Turon.-Maastricht.)*.

All elements observable in morphological study of the Hippuritidae (pores, ligamental ridge, teeth, pillars) occur in combinations that permit different genera to be defined. Important for determination of genera and species is the sector (*L-Sp-Ep*) occupied by the three principal folds—the ligamental ridge (*L*) and the pillars (*Sp, Ep*).

The pillars, pores and oscles are the most curious features of the hippuritids, and paleontologists have long tried to interpret them. DESHAYES and WOODWARD supposed that the pillars served for muscle attachment, but suggested different ways in which this may have been accomplished. BAYLE observed that the three folds have the same structure, and he considered that the muscles were not attached to them but to the wall of the shell. DOUVILLÉ, extending the conclusions reached from his work on *Sphaerulites* and genera of the Requieniidae and Monopleuridae to *Hippurites*, suggested that the pillars correspond to siphonal zones, and accordingly designated them as "S" (*sortie*, excurrent) and "E" (*entrée*, incurrent). This opinion was accepted widely, with its implication that a hippuritid possessed two siphons serving respectively for outflow and inflow of water currents. YONGE (1024a) has advanced other interpretations based on analogies with *Chama*. The presence of oscles in the FV exactly above extremities of the pillars was held to support this hypothesis, for a hippuritid with closed valves could be provided continuously with water circulation through siphons opening outward at the oscles. Oacles are not a constant feature of all Hippuritidae, however, and, according to some accounts, where they exist in earlier

growth stages they may later become obliterated.

Study of the structure of the pillars by means of thin sections has led to alternative hypotheses. KLINGHARDT (475aa) and WIONTZEK (1000) observed cavities within the pillars and suggested that each pillar

was a "siphonal tube" which enclosed the siphon of the living animal. No orifice, however, has been observed at the base of the pillars (Fig. E263,1c) through which a siphon could have protruded or a current of water have passed, and it seems probable that the cavities are of secondary origin.




FIG. E263. Hippuritidae (p. N801-N802). [Explanation: *Eo*, *So*, oacles; *Ep*, *Sp*, internal pillars; *L*, ligamental ridge.]

Actually, the pillars are folds of the external layer of the shell (MILOVANOVIĆ, 622, 625; KÜHN, 491), similar to other folds known in various hippuritids. A whole series of observations throws doubt on the existence of siphons in the hippuritids, and the hypothesis that long ones were enclosed in the pillars should be rejected.

CHUBB (131) has suggested that the sole function of the pillars was to block the oscules of the upper valve when this was pulled down tightly by the adductor muscles, thus interrupting the passage of the respiratory currents through them when necessity arose (Fig. E263,1e). The oscules themselves, he suggested, evolved in consequence of the gregarious habit of the hippurites (and other rudists in which they are found), their presence enabling interchange of water between the mantle cavity and the sea above the animal when interchange from the side may have been difficult or impossible.

Another theory recently has been advanced by VOGEL (942), who suggested that the pillars acted as supports of sensory organs or of accessory gills. As there are no living bivalves at all like the hippuritids, it is difficult to decide what role the observed morphological characters played in the biology of the animal.

If the siphons were short or lacking, circulation of water was perhaps provided by the pores. In addition, some authors have ascribed to the pores the function of a filter that screened off impurities harmful to the animal. The arrangement of canals in the FV is complex, but their organization, especially in *Yvaniella* (Fig. E243,4) seems to indicate that water circulation was made possible to some extent by this system of perforations when the valves were closed.

As the pillars blocked the oscules completely when the shell was closed, the hippuritids must have had to raise the FV slightly to allow circulation of water for respiration and feeding. The FV is veritably a lid placed over the AV in such a manner that the long teeth and posterior myophore are engaged by the long vertical sockets of the AV. Because of this, any rocking movement about the ligament is impossible and only elevation of the FV above the AV permits separation of one from the other.

This lifting could have been effected by pressure in the shell interior, produced possibly by inflation of a soft organ such as the foot.

Hippurites LAMARCK, 1801, p. 104 [**H. bioculata*; M] [= *Orthoceratites* LAMARCK, 1799, p. 81 (suppressed ICZN Op. 613); *Cornucopina* THOMSON, 1802, p. 245 (generic description without nominal species); *Coralliolites* VON SCHLOTHEIM, 1813, p. 36 (type, *C. orthoceratoides*; SD COOGAN, herein); *Pachynus* RAFINESQUE, 1815, p. 140 (obj.); *Dorbignya* WOODWARD, 1862, p. 375 (obj.); *Orbignia* STOLICZKA, 1871 (nom. van.); *Orbignya* FISCHER, 1887, p. 1064 (nom. van.); non *Hyppurites* VON SCHLOTHEIM, 1820, p. 351]. AV nearly cylindrical, smooth, bearing 2 furrows in some but not all species; ligamental ridge short or absent; pillars *Ep* and *Sp* distinct, *L* absent or slightly curved to pointed inflexion of wall; sector *L-Sp-Ep* greater than one-quarter circumference; FV nearly flat, with 2 oscules and simple linear, well-separated pores. *U.Cret. (Turon.-Maastricht)*, Eu.-NE. Afr. (Somalia)-Asia-N. Am.-Antilles.—FIG. E263,1a-c. **H. bioculata*, *U.Cret. (Senon.)*, France; 1a, FV ext., $\times 1.3$ (269); 1b, transv. sec. viewed toward commissure, AV center pseudo-pillars *Sp* and *Ep* dark, $\times 1.7$; 1c, detail ant. part polarized light, $\times 2.7$; 1d, pillar *Sp*, polarized light, $\times 13$; 1e, long. sec. AV, pillars white, oscular indentations above pillars, $\times 2$ (942).—FIG. E263,1f. *H. canaliculatus* ROLLAND DU ROUAN, *U.Cret.*, France; linear pores in FV, $\times 7$ (269). [Also Fig. E220,1-2,6.]

Barrettia WOODWARD, 1862, p. 372 [**B. monilifera*; M] [= *Barretia* MUNIER-CHALMAS, 1873 (nom. null.)]. Adult shell very large; AV elongate, recumbent, distinguished by up to 60 or more beaded (moniliform) rays in transverse section; *Sp* and *Ep* distinguishable by their swollen extremities, that of *Sp* being circular and that of *Ep* elongate elliptical; *L* indentifiable by reference to teeth; portion of valve between *L* and siphonal pillars greater than 0.3 of shell circumference; internal layer strongly developed, forming cellular tissue between folds and tabular below shell cavity. FV convex, operculiform, consisting of central boss from which radiate numerous rods, oscule fitting over pillar *Sp*, no *Ep* oscule. *U.Cret. (Campan.-Maastricht.)*, W. Indies-Mexico.—FIG. E263,3. **B. monilifera*, Jamaica; AV from above, $\times 0.2$ (981). [Also Fig. E227,2; E237,10.]

Batolites DE MONTFORT, 1808, p. 334 [**B. organisans*; OD] [= *Batholites* GABB, 1862 (nom. van.); *Bihippurites* FUTTERER, 1896, p. 263 (type, *B. plicatus* FUTTERER)]. AV long, cylindrical, slender, shaped like organ pipe; internally like *Hippurites*, but differs in having outer shell layer with numerous folds which, however, do not give rise to supplementary pillars projecting toward




FIG. E264. Hippuritidae 1, 3-6; unrecognizable (2) (p. N802-N803). [Explanation: *Ep*, *Sp*, internal pillars; *L*, ligamental ridge; *pm*, posterior myophore; *R*, anterior (extra) pillar; 1, 3, anterior and posterior teeth of FV.]

shell interior; 3 distinct pillars; *L* short, area of pillars about 0.3 of shell circumference. FV with 2 oscules and linear pores. *U.Cret.(Santon.)*, France-Austria-USSR.—FIG. E263, 2. **B. organisans*, Pyrenees; 2*a*, *b*, AV ext. and FV ext. (same individual), $\times 0.7$, $\times 2$ (269). [Also Fig. E237, 11.]

Hippuritella DOUVILLÉ, 1908, p. 268 [**Hippurites maestrei* VIDAL, 1878, p. 99; OD]. Differs from *Hippurites* in having simple or denticulate pores in FV. *U.Cret.(Turon.-Maastricht.)*, Eu.-N.Afr. NE. Afr.(Somalia)-Asia-N. Am.—FIG. E263, 4*a*. **H. maestrei* (VIDAL), Santon., France (Pyrenees); AV int. view, $\times 2$ (269).—FIG. E263, 4*b*. *H. sulcatoides* DOUVILLÉ, Campan., France; polygonal pores FV, $\times 3.5$ (269). [Also Fig. E224, 1.]

Parastroma DOUVILLÉ, 1926, p. 133 [**P. sanchezi*; M]. AV wall mainly of small vesicular domelike folded tabulae; rays almost effaced, stems of rays *Ep* and *Sp* rudimentary near outer shell layer, occupying more than 0.3 of shell circumference. FV unknown. *U.Cret.(Campan.-Maastricht.)*, Cuba.—FIG. E263, 5. **P. sanchezi*; details of AV wall, 5*a*, transv. sec., 5*b*, long. sec., $\times 3$ (280).—FIG. E264, 6. *P. guitari* (PALMER), Maastricht, Cuba; transv. sec. at level of AV with projecting teeth of FV, $\times 0.4$ (561).

Pironea MENEHINI in PIRONA, 1868, p. 511 [**Hippurites polystylus* PIRONA, 1868, p. 508;

M] [= *Pironea* MENEHINI, 1868 (*nom. nud.*); *Pyronea* MUNIER-CHALMAS, 1873 (*nom. null.*); *Pironea* BLANCKENHORN, 1934 (*nom. null.*); *Peronea* WHITFIELD, 1897 (*nom. null.*)]. Attaining large size; surface of AV with numerous longitudinal furrows; outer shell layer with many infoldings which give rise to secondary pillars or rays, some as prominent as ligamental ridge (*L*) and pillars (*Sp*, *Ep*), other smaller infoldings located between the more prominent ones and between *Sp* and *Ep*; sector *L-Ep-Sp* less than 0.25 of circumference value. FV flattened, with denticulate pores. *U.Cret.(Campan.-Maastricht.)*, Eu.-Afr.-Syria-Asia-N. Am.-Cuba.—FIG. E238, 6. **P. polystylus* (PIRONA), Maastricht, Italy; transv. sec. AV, $\times 0.3$ (269).

Praebarrettia TRECHMANN, 1924, p. 395 [**Barrettia sparcilirata* WHITFIELD, 1897, p. 245; M]. Shell moderate in size. Differs from *Barrettia* in having fewer (15 to 40) rays which become moniliform only in late growth stages; rays *L*, *Sp*, *Ep* clearly distinguished, occupying more than 0.3 of shell circumference. *U.Cret.(?Santon.-Maastricht.)*, W. Indies-S. Am.—FIG. E264, 4. **P. sparcilirata* (WHITFIELD), Jamaica; AV from above, $\times 0.3$ (981). [Also Fig. E238, 3.]

Rhodensia SÉNESSE, 1939, p. 227 [**R. mutans*; OD]. AV with 3 distinct pillars, *Ep*, *Sp*, and *L*; pillar *L* elongate, about equal to *Ep*, set at small angle

to teeth; pillars occupying more than 0.25 circumference of shell. FV pores linear, marginally subpolygonal. *U.Cret.(Up.Turon.)*, Eu. (France).—Fig. E264,3. **R. mutans*; AV transv. sec., $\times 0.7$ (839).

Tetracionites ASTRE, 1931, p. 269 [**T. mozambicus*; M]. AV small, with 4 external furrows corresponding to 4 internal folds, *Ep*, *Sp*, *L*, and anteriorly placed extra pillar (*R*); FV unknown. *U.Cret.(Coniac.)*, France-Madag.—Fig. E264,5. **T. mozambicus*; diagram. sec. AV, $\times 1.3$ (22).

Torreites PALMER, 1933, p. 99 [**Hippurites (Vaccinites) sanchezii* DOUVILLÉ, 1927, p. 27; OD]. AV cylindroconical without well-marked external furrows; internally pillars *Ep*, *Sp*, *L* unequal, *L* very long, longer than shell radius; *Sp* twice length of *Ep*; sector *L-Sp-Ep* about 0.25 of circumference; outer shell layer thin, forming less well-developed folds than in *Batolites*. FV depressed-conical, with 3 radial grooves corresponding to *L*, *Sp* and *Ep* without pores, imperforate outer layer and canaliferous inner layer. *U.Cret.(?Santon-Campan.)*, Cuba.—Fig. E265,1. **T. sanchezii* (DOUVILLÉ); transv. sec. AV, $\times 0.6$ (281). [Also Fig. E238,5.]

Vaccinites FISCHER, 1887, p. 1064 [**Hippurites cornuvaccinum* BRONN, 1831, p. 374; M] [= *Pseudovaccinites* SÉNESSE, 1947, p. 40 (type, *P. pseudolatus major* SÉNESSE)]. AV cylindroconical, large; 3 distinct pillars, *Ep*, *Sp*, and *L* occupying less than 0.25 of shell circumference; pillar *L* generally long, angle between *L* and teeth less than 45 degrees; ligamental ridge generally truncate, *Sp* and especially *Ep* tending to be contracted proximally; inner margin of outer shell layer undulating. FV flat to slightly convex; pores reticulate or denticulate, 2 oscules. *U.Cret.(Turon-Maastricht.)*, Eu.-Asia-N. Afr.-W. Indies.—Fig. E264,1a. *V. corbaricus* (DOUVILLÉ), Coniac, S. France; reticulate pores, $\times 7$ (269).—Fig. E264, 1b. *V. marticensis* (DOUVILLÉ), Coniac, S. France; FV ext., $\times 1$ (910). [Also Fig. E237,6.] [SÉNESSE, who classified the hippuritids solely on the basis of shape of the FV pores, has maintained that, since their pores are polygonal, species of the group that includes the type of *Vaccinites* are more closely related to *Hippurites* than to the remaining groups included by TOUCAS and others in *Vaccinites*. Accordingly, he has proposed to unite all hippuritids with reticulate, subreticulate or denticulate pores in a genus named *Pseudovaccinites*. The classification of TOUCAS is accepted here, however.]

Yvaniella MILOVANOVIĆ, 1938, p. 129 [pro *Ivania* MILOVANOVIĆ, 1936, p. 33 (non FISCHER ex BAYLE, 1885)] [**Yvania mastrichtiensis* MILOVANOVIĆ, 1936; M]. AV like *Hippurites*. FV specialized, concave, traversed by simple canals which open on outer surface as small pores and on inner surface as enlarged funnel-shaped pores separated by minutely rugose ridges; at center of valve is tubercle traversed by 2 large canals, walls of which contain fine canals, some opening as minute pores on surface of tubercle. *U.Cret.(Maastricht.)*, Eu.(Yugosl.).—Fig. E243,3-4.




FIG. E265. Hippuritidae (p. N803). [Explanation: *am*, *pm*, anterior and posterior myophores; *Ep*, *Sp*, internal pillars; *L*, ligamental ridge; *1*, *3*, sockets in AV corresponding to teeth in FV.]

**Y. mastrichtiensis* (MILOVANOVIĆ); 3, long. sec. FV (624); 4, drawing showing both valves from above, $\times 0.7$ (628).

UNRECOGNIZABLE HIPPURITID GENERA

Chiapasia MÜLLERRIED, 1933, p. 268 (*nom. nud.*).
Pseudobarrettia MÜLLERRIED, 1931, p. 255 [**P. chiapasensis* (*nom. nud.*); OD]. Based on pathological anomaly in growth of AV pillar in individuals of a *Barrettia*-like hippuritid. *U.Cret.(Campan.)*, Mexico (Chiapas).—Fig. E264,2. **P. chiapasensis*; transv. sec. of AV showing pathological duplication of pillar *L* (scale not given, ca. $\times 1.3$ (651)).

Family RADIOLITIDAE Gray, 1848

[Materials for this family prepared by COLETTE DECHASEAUX and A. H. COOGAN, with acknowledgment of assistance by L. R. Cox]

Inequivalve, RV (AV) conical, LV (FV) operculiform; surface of RV (AV) without furrows but generally with two concave, flat, or convex siphonal bands (*Eb*, *Sb*) separated by interband; ornament of siphonal bands and interband unlike that of remainder of surface (Fig. E266,4); in LV (FV), oscules present in a few genera; structure of shell characteristically celluloprismatic; outer layer thick, cellular in texture, walls of cellulites formed by radial and transverse funnel-shaped plates which intersect to form hollow prisms with rec-




FIG. E266. Morphological features of Radiolitidae.

- Bournonia*, U.Cret.(Maastricht.), France; *1a*, *B. fourtaui* DOUVILLÉ, transv. sec. of part of wall showing quadrangular network of celluloprismatic structure, $\times 0.7$ (251); *1b*, *B. bournoni* DES MOULINS, AV long. sec. showing reticulate shell wall and on inside of valve fluted groove for reception of sliding tooth of FV, $\times 0.7$ (251).
 - Lapeirousia jouanneti* DES MOULINS, U.Cret. (Maastricht.), France (Charente); *2a*, transv. sec. of part of shell wall showing polygonal cell structure, $\times 3.3$; *2b*, tang. sec. perpendicular to plane of *2a* sec., $\times 3.3$ (Dechaseaux, n).
 - Pseudopolyconites parvus* MILOVANOVIĆ, U.Cret. (Maastricht.), Serbia; transv. sec. at level of AV with projecting teeth and myophores of FV, $\times 0.7$ (623).
 - Sauvagesia nicaisei* COQUAND, var. *villei* ARNAUD, U.Cret.(Cenoman.), France (Charente); side view ext. AV showing longitudinally fluted surface and position of external finely ribbed *Eb* and *Sb* siphonal bands, $\times 0.7$ (911).
 - Joufia reticulata* BÖHM, U.Cret.(Senon.), Italy; *5a,b*, long. and transv. secs. of part of FV shell wall showing polygonal cell structure, $\times 6.7$ (905).
- [Explanation: *L*, ligament ridge; *am*, *pm*, ant. and post. myophores; *Eb*, *Sb*, siphonal bands; *1*, *3*, FV teeth, *2*, AV tooth.]

tangular or polygonal bases; at lip of valve these prisms form network with polygonal or quadrangular mesh (Fig. E266, *1a-2a,b*). Interior of RV (AV) without pillars, but in some forms with low, broad, longitudinal projections of shell wall termed "pseudopillars"; hinge, with two teeth in LV (FV) and one tooth in RV (AV); ligamental ridge present or absent; teeth and myo-

phores of LV (FV) forming thick laminae, fluted in some species (Fig. E220,3); hinge of RV (AV) commonly reduced to sockets into which teeth of LV (FV) slide (Fig. E266, *1b*); muscles attached to wall of shell in RV (AV); many genera with accessory cavities outside cardinal structures (Fig. E266,3). *L. Cret. (Barrem.) - U. Cret. (Maastricht.)*.

The major features of the Radiolitidae, siphonal bands and their ornament, presence or absence of a ligamental ridge, and myophore apophyses and wall structure, are characters by which different genera in the family are distinguished. The siphonal

bands and pseudopillars have structure similar to that of other parts of the shell, differing chiefly in the greater development of certain layers of cellules. The function of the pseudopillars is still uncertain; the supposition that these structures were re-




FIG. E267. Radiolitidae (Radiolitinae) (p. N806-N808). [Explanation: *Eb*, *Sb*, siphonal bands.]




FIG. E268. Radiolitidae (Radiolitinae) (p. N807-N808).

lated to respiratory or digestive organs is unproved. So far as known, however, all forms with true pseudopillars have oscules in the FV.

Subfamily RADIOLITINAE Gray, 1848

[*nom. transl.* DOUVILLÉ, 1903 (*ex* Radiolitidae GRAY, 1848)]

Wall of RV (AV) composed of quadrangular prismatic cells; posterior ligament present. *L. Cret. (Barrem.) - U. Cret. (Maastricht.)*.

Radiolites LAMARCK, 1801, p. 130 [**Ostracites angeiodes* PICOT DE LAPEIROUSE, 1781, p. 40; M] [= *Euradiolites* WIONTZEK, 1934, p. 11 (presumed obj.)], AV conical, ornamented with strong longitudinal folds over whole valve, siphonal bands smooth accentuations of regular folds, outer wall structure coarsely reticulate. FV small, conical, strongly convex or rarely flat, myophore apophyses present. *U. Cret. (Cenoman.-Maastricht.)*, Eu.-N.

Afr.-NE.Asia-N.Am.—FIG. E267,2. **R. angeiodes* (DE LAPEIROUSE), U. Santon., France (Pyrenees), 2a, AV lat. post., $\times 0.9$; 2b,c, AV from above and below, $\times 2.2$ (23). [Also Fig. E220,4.]

Agriopleura KÜHN, 1932, p. 78 [*pro Agria* MATHÉRON, 1878, pl. C-8 (*non* ROBINEAU-DESVOIDY, 1830)] [**Hippurites blumenbachi* STUDER, 1834, p. 107; OD] [= ?*Aptyxites* BÖHM, 1919, p. 74 (type, *Radiolites muelleri* WEGNER, 1905, p. 193)]. AV elongate, straight or slightly arched, commonly longitudinally ribbed; surface with 2 unequal concave furrows bordered by longitudinal swellings; siphonal bands in furrows; ligamental ridge well defined. FV wall structure poorly known, probably finely lamellar; operculiform, flat or strongly concave; hinge with 2 slender teeth, thickened areas at their base constituting muscle scars; no myophore apophyses. [Lack of myophore apophyses separates *Agriopleura* from other Radiolitinae.] *L. Cret. (Barrem.) - U. Cret. (Maastricht.)*, Eu.-N.Am.—FIG. E267,3. **A. blumen-*

bachi (STUDER), Barrem., France; 3a, FV ext., $\times 0.9$; 3b, both valves lat. view, $\times 0.9$ (911). [Also Fig. E237,1; E240,5.]

Eoradiolites DOUVILLÉ, 1909, p. 77 [**Radiolites davidsoni* HILL, 1893, p. 106; OD]. AV conical elongate, prominent smooth siphonal bands, large ribbed sockets in dorsal inner wall separated from

shell cavity by lateral extension ("transverse septum") of cardinal platform from which lower valve tooth arises, outer wall of coarsely reticulate cells. FV small, operculiform, concave to flat, of fine concentric laminations, myophore apophyses present. *L.Cret.(Alb.)-U.Cret.(Turon.)*, Eu.-Afr.-Asia-N.Am.—FIG. E267,4. **E. davidsoni*




FIG. E269. Radiolitidae (Radiolitinae) (p. N808-N810). [Explanation: *am*, *pm*, anterior and posterior myophores; *Eb*, *Sb*, siphonal bands; *L*, ligamental ridge; 1, 3, teeth of FV; 2, tooth of AV.]

- (HILL), Alb., USA (Texas); 4a, both valves; 4b, AV lat. view of siphonal band Eb, $\times 0.9$ (Coogan, n). [Also Fig. E237,2; E240,1.]
- Gorjanovicia** POLŠAK, 1968, p. 201 [**G. costata*; OD]. AV elongate, cylindrical-conical, slender, longitudinal ribs salient, smooth or with fine zigzag growth lines, commonly pointed and separated by finely costulate furrows; siphonal zones large wide bands, pseudopillars clearly imprinted on internal face have special structure of folds with concave inflexions or are concentric or radially disposed. Regular shell structure lamellar, ligament triangular. FV slightly convex and costulate. Differs from *Medeella* in lacking external folds and in having differently shaped siphonal zones. *U.Cret. (Santon.-L. Campan.)*, Yugosl. (Istria).—FIG. E268,1. **G. costata*; 1a, lat. view, 1b, transv. sec., $\times 0.7$; 1c, tang. sec., AV, $\times 4.3$ (746b).
- Kühnia** MILOVANOVIĆ, 1956, p. 132 [**K. serbica*; OD]. Known only by AV; ligamental ridge well developed, bilobate at extremity; no pseudopillars; circle of tubular cavities in inner part of thick outer shell layer, prismatic cellular structure of which is well defined. *U.Cret. (Santon.-Campan. boundary)*, Serbia.—FIG. E267,5. **K. serbica*; transv. sec. AV, $\times 0.45$ (627).
- Medeella** PARONA, 1924, p. 64 [**Radiolites zignana* PIRONA, 1869, p. 419; OD]. AV elongate, cylindrical, longitudinal folds regularly costulate, shell surface covered by fine growth lines, outer wall thick; cylindrical siphonal structure in outer wall of polygonal cells, touching body cavity; myophore apophyses present. *U.Cret. (Turon.-Coniac.-?Santon.)*, Eu.
- M. (Medeella)**. Siphonal bands broadly costulate; FV convex; siphons marked by inflected crescentic slits. *U.Cret. (Turon.-Coniac.-?Santon.)* Eu.—FIG. E267,1. **M. (M.) zignana* (PIRONA), Italy; 1a-c, views ventral, lat., and from above, $\times 0.9$ (740); 1d, AV transv. sec., $\times 1.8$ (720). [Also Fig. E243,7.]
- M. (Fossulites)** ASTRE, 1957, p. 42 [**Medeella undaesaltus* ASTRE, 1954; M]. Longitudinal folds mark surface, siphonal bands salient, smooth or costate, cylindrical siphonal fossettes or cavities in AV outer wall. FV operculiform, flattened, siphons marked by longitudinal arches. Fossettes distinctive. *U.Cret. (L. Coniac.)*, France (Aude).—FIG. E243,1. **M. (F.) undaesaltus*; 1a, both valves; 1b, lat. view AV, 1c, transv. sec. AV, $\times 0.9$ (24).
- Neoradiolites** MILOVANOVIĆ, 1935, p. 97 [**N. serbicus*; M]. AV conical, ornamented by fine costae, siphonal zones marked by slight inflexions, valve outline irregularly circular, outer wall thick, of coarsely prismatic cells. FV capuloid, superficially like *Robertella*, but with ligament. *U.Cret. (Maastricht.)*, Yugosl.—FIG. E269,6. **N. serbicus*; AV transv. sec., at level of AV with projecting teeth of FV and part of dorsal overlap of FV, $\times 0.75$ (623).
- Parasauvagesia** Cox, 1960, p. 428 [**P. cappadociensis*; OD]. AV depressed conical, longitudinally costate, without well-differentiated siphonal bands; short ligamental ridge and 2 weakly developed pseudopillars present. FV with central domelike elevation and broad, flat brim; no oscules. *U.Cret. (Campan. or Maastricht.)*, Turkey.—FIG. E269,1; E270,7. **P. cappadociensis*; E269,1a,b, lat. and top view, $\times 0.2$; E269,1c, AV transv. sec. shell, $\times 1$; E270,7, AV transv. sec., $\times 0.55$ (199).
- Paronella** WIONTZEK, 1934, p. 26 [**P. volzanensis*; M]. AV cylindrical, transverse septum separating shell cavity from tooth sockets; teeth, myophores, outer wall structure and FV unknown; differs from *Distefanella* in having ligament. *U.Cret. (L. Santon.)*, SE.Eu. (Italy).—FIG. E269,4. **P. volzanensis*; AV transv. sec., $\times 0.75$ (998).
- Praeradiolites** DOUVILLÉ, 1902 [1903], p. 467 [**Radiolites fleuriau* D'ORBIGNY, 1842, p. 181; OD]. AV conical, externally with smooth folds in form of stack of inverted cones, in younger species developed into undulating chevron pattern; siphonal folds convex, plates bent upward, separated by projecting fold; outer wall structure coarsely reticulate. FV operculiform, convex to flat; myophore apophyses present. Stacked, undulating siphonal band folds distinctive. *L.Cret. (Alb.)-U.Cret. (Maastricht.)*, Eu.-N.Afr.-NE.Asia-N.Am.—FIG. E269,2a-c. **P. fleuriau* (D'ORBIGNY), Cenoman., France (Charente); 2a,b, post. lat. and ant. views, $\times 0.75$; 2c, view from above, $\times 0.75$ (911).—FIG. E269,2d. *P. toucasi* (D'ORBIGNY), Santon., France; AV lat. view, $\times 0.75$ (911).—FIG. E269,2e. *P. cylindraceus* (DES MOULINS), Maastricht., France; FV int., $\times 0.45$ (911). [Also Fig. E220,3.]
- Pseudopolyconites** MILOVANOVIĆ, 1935, p. 132 [**P. parvus*; SD MILOVANOVIĆ, 1937, p. 4]. AV conical, shell surface nearly smooth, strong longitudinal folds absent, siphons marked by broad folds, internal inflexion present at incurrent position; ligamental ridge long, displaced dorsally; outer wall of 2 layers, internal prismatic and normal prismatic; siphonal zones with hornlike transverse partitions. FV weakly convex, teeth and myophores present, shell wall 2 layers. *U.Cret. (Maastricht.)*, Yugosl. (Serbia).—FIG. E266,3. **P. parvus*; transv. sec. AV, $\times 1$ (623).
- Radiolitella** DOUVILLÉ, 1904, p. 533 [**Chama forojuliensis* PIRONA, 1869, p. 431; OD]. AV conical, with ligamental ridge and siphonal bands; outer wall in juvenile stages of coarse polygonal cells that become rectangular in adult. FV cap-shaped with strong siphonal bands. *U.Cret. (Santon.-Maastricht.)*, Eu. (Italy-Spain).—FIG. E269,3. **R. forojuliensis* (PIRONA), Maastricht., Italy; 3a, lat. view, $\times 0.75$; 3b,c, FV ext., AV transv. sec., $\times 2$ (272).
- [PARONA (1923, p. 61) stated that DOUVILLÉ in designating *Chama forojuliensis* PIRONA as the type of *Radiolitella* figured a specimen from Colle di Medea in the Sorbonne (Paris) collection which corresponds with PIRONA's figures

of *Sphaerulites guiscardiana* PIRONA, 1869, not with *Chama forojuliensis* PIRONA. KÜHN, 1932, accepted PARONA's evaluation listing *Sphaerulites guiscardiana* PIRONA (= *Radiolitella forojuliensis* DOUVILLÉ, non PIRONA) as the type of *Radiolitella*.]

Robertella COSSMANN, 1904, p. 254 [pro *Mouretia*

DOUVILLÉ, 1903, p. 480 (non G. B. SOWERBY, 1835)] [**Mouretia arnaudi* DOUVILLÉ, 1903; M] [= *Sarlatia* DOUVILLÉ, 1910, p. 29 (obj.)]. Nearly equivalve (AV slightly larger than FV), each




FIG. E270. Radiolitidae (Radiolitinae) (7); (Biradiolitinae) (1-3,5); (Sauvagesiinae) (4,6) (p. N810-N812). [Explanation: *am*, *pm*, anterior and posterior myophores; *Eb*, *Sb*, siphonal bands; *Es*, *Ss*, pseudo-pillars; *L*, ligamental ridge; *1*, *3*, anterior and posterior teeth of FV.]

valve capuloid in form; shell wall very thin, without well-marked siphonal bands (only 2 very weak undulations or depressions); no attachment scar at extremity of AV, shell seemingly free, resting on sea bottom; cardinal structures typically radiolite, with ligamental ridge and myophore apophyses. *U.Cret. (Santon.)*, France (Sarlat).—FIG. E269,5. **R. arnaudi* (DOUVILLÉ); 5a, FV side view, $\times 1.5$; 5b,c, AV int. and side view, $\times 0.7$ (271).

Sphaerulites LAMARCK, 1819, p. 231 (ex DELAMÉTHÉRIE, 1805, *vernac.*) [**S. foliaceus*; M] [= *Sphaerulites* DE BLAINVILLE, 1824 (*nom. null.*)]. AV broad flattened sphere, external folds undulating slack; siphonal zones marked by sinuses, bordered and separated by folds; internally 2 sinus folds indenting shell cavity; shell structure probably coarsely reticulate. FV flattened, myophore apophyses present, broad, compressed. Foliaceous shape of AV distinctive. *L.Cret. (Apt.)-U.Cret. (Turon.)*, Eu.-N.Afr.—FIG. E220,5. **S. foliaceus*, Cenoman., France; AV transv. sec., $\times 0.3$ (911). [Also Fig. E240,2.]

Subfamily BIRADIOLITINAE Douvillé, 1902

RV (AV) wall of quadrangular prismatic cells, posterior ligament absent. *U.Cret. (Turon.-Maastricht.)*.

Biradiolites D'ORBIGNY, 1850, p. 230 [**B. canaliculatus* D'ORBIGNY, 1850; SD ICZN pend.] [= *Eubiradiolites* COOGAN, 1966, p. 763 (*obj.*)]. AV conical, more or less produced, straight or arched; siphonal bands smooth, slightly depressed; 1 rib in interband; outer wall of fine cells; AV tooth absent. FV operculiform convex to concave with ornament like that of AV. *U.Cret. (Turon.-Maastricht.)*, Eu.-N. Afr.-Asia-N. Am.-Antilles.—FIG. E270,1. **B. canaliculatus*, Coniac., France; side view, $\times 0.75$ (911). [Also Fig. E237,4.]

Bournonia FISCHER (ex BAYLE, MS), 1887, p. 1067 [**Sphaerulites bournoni* DES MOULINS, 1826 (1827), p. 271; M]. AV conical, small to very large ornamented by folds and sporadically by collared horns as in *Praeradiolites*; siphonal bands salient, undulating smooth or ribbed; anterior dorsal fold may project markedly. FV operculiform, convex. *U.Cret. (Turon.-Maastricht.)*, Eu.-N.Afr.-Asia-W.Indies.—FIG. E270,5. **B. bournoni* (DES MOULINS), Maastricht., France (Dordogne); lat. view, $\times 0.4$ (251). [Also Fig. E266, 1.]

Distefanella PARONA, 1901, p. 205 [**D. salmoiraghii*; SD PARONA, 1912, p. 283] [= *Stefanella* DOUVILLÉ, 1901, p. 101 (*nom. van.*)]. AV subcylindrical, elongate, uniformly costate, siphonal bands smooth to finely striate, outer wall thin; internally septum connects dental sockets, dorsal cavity large; AV tooth rudimentary. FV operculiform, cap-shaped. *U.Cret. (Turon.-Senon.)*, Italy-Yugosl.-?Jamaica.

—FIG. E237,8. **D. salmoiraghii*, Italy; transv. sec. AV, $\times 0.7$ (719). [Also Fig. E224,2.]

Milovanovicia POLŠAK, 1968 [**M. heraki*; OD]. AV cylindro-conical, very elongate, straight or slightly curved. External ornamentation of few large massive, elongate folds separated by deep furrows; both with fine longitudinal striae. Siphonal bands costulate as in *Sauvagesia* and *Durania*. Pseudopillars not developed, ligamental crest absent. Shell structure lamellar. FV unknown. Differs from *Petkovicia* in its lamellar shell structure, costulate siphonal bands, external ornamentation, and absence of pseudopillars. *U.Cret. (Turon.)*, Yugosl. (Dalmatia).—FIG. E271,1. **M. heraki*; 1a, lat. view, 1b, transv. sec. AV, $\times 0.53$; 1c, transv. sec. showing detailed lamellar structure, $\times 4.8$ (746a).

Parabournonia DOUVILLÉ, 1927, p. 55 [**P. hispida*; M]. Shell small; differs from *Bournonia* in having AV wall of denticulated structure of parallel plates connected by spines, according to DOUVILLÉ. Considered as indistinguishable from *Bournonia* or *Biradiolites* (810). *U.Cret. (Maastricht.)*, W. Indies (Cuba).—FIG. E270,3. **P. hispida*; 3a, transv. sec. AV, $\times 0.75$; 3b, detail of AV outer wall, $\times 3$ (281).

Synodontites PIRONA, 1867, p. 840 [**S. stoppaniana*; M] [= *Synodontites* STOLICZKA, 1871 (*nom. null.*)]. AV pyramidal, elongate, base quadrangular; siphonal bands narrow, smooth; teeth displaced toward center of shell cavity, fused together except at apex. FV operculiform, flat to concave, ornamented like AV. Held to differ from *Biradiolites* by having fused teeth. *U.Cret. (Santon.)*, Eu. (Italy-France).—FIG. E270,2. **S. stoppaniana*, Italy; 2a, AV lat. view, $\times 0.75$; 2b,c, FV ext. and side view, $\times 0.75$ (739).

Thyrastylon CHUBB, 1956, p. 36 [**Radiolites adhaerens* WHITFIELD, 1897, p. 188; OD]. AV conical or distorted by anterior attachment; siphonal bands deeply sunken, interband rounded. FV conical, convex, brim of valve forms oval oscules above AV siphonal areas; oscules distinctive. *U.Cret. (Maastricht.)*, Jamaica-Cuba-Guatemala-Iran.—FIG. E243,2. **T. adhaerens* (WHITFIELD), Jamaica; 2a, 2 shells, upper showing oscule *So* in FV; lower only AV, $\times 0.7$; 2b, both valves, FV with oscules *Eo* and *So* enclosed, $\times 0.7$; 2c, transv. sec. AV, $\times 4$ (131).

Subfamily SAUVAGESIINAE Douvillé, 1908

RV (AV) wall composed of cells which are prismatic in longitudinal section but polygonal in transverse section. *L.Cret. (Alb.)-U.Cret. (Maastricht.)*.

Sauvagesia CHOFFAT (ex BAYLE, MS), 1886, p. 31 [**Sphaerulites sharpei* BAYLE, 1857, p. 638; SD DOUVILLÉ, 1903, p. 474]. AV conical to cylindro-conical, ornamented with longitudinal ribs;




FIG. E271. Radiolitidae (Bioradiolitinae) (p. N810). [Explanation: *Eb*, *Sb*, siphonal bands.]

siphonal bands finely costulate folds, separated by interband that generally bears 2 costae. FV operculiform, with radial folds; ligamental ridge present, most marked in earlier forms. *U.Cret.(Alb.)-U.Cret.(Maastricht.)*, Eu.-N.Afr.-N.Am.-Antilles. —FIG. E238,4. **S. sharpei* (BAYLE), Turon., Port.; AV, transv. sec., $\times 0.6$ (268). [Also Fig. E266,4.]

[The generic name *Sauvagesia*, attributed to BAYLE, was also published in 1886 by DOUVILLÉ (Bull. Soc. Géol. France, Ser. 3, v. 14, p. 393), who cited only one species, *Sphaerulites lusitanicus* BAYLE, 1857, as an example. DOUVILLÉ was misled by a misidentified specimen, for, as pointed out by TOUCAS (919, p. 81), this species does not belong to the group to which it was intended to apply the generic name. DOUVILLÉ himself (268, p. 669; also *ibid.*, ser. 4, v. 2, p. 464) had previously admitted his mistake and the fact that CHOFFAT had interpreted *Sauvagesia* correctly. The best course seems to be to attribute the name to CHOFFAT and to apply for the suppression of its publication by DOUVILLÉ, should this prove to have priority.]

Apulites TAVANI, 1958, p. 173 [**A. giganteus*; M]. AV very elongate, with longitudinal costae; outer wall thin, of subquadrangular cells; siphonal zones marked by finely costate folds; ligamental ridge lacking. FV unknown. May be a biradiolite. *U.Cret.(Senon.)*, SE.Eu.(Italy). —FIG. E270,4. **A. giganteus*; 4a, AV lat. view, $\times 0.75$; 4b, AV

transv. sec., $\times 1.1$; 4c, detail of AV wall, $\times 4$ (905).

Chiapasella MÜLLERRIED, 1931, p. 243 [**Coralliochama radiolitiiformis* TRECHMANN, 1924, p. 406; M]. AV conical, short, expanding rapidly, ornamented with longitudinal folds among which striated siphonal zone may be distinguished. Outer wall of irregular polygonal cells; cortex folded into wall locally along periphery. FV high and coiled as in a caprinid, convex or flattened, shell material brown, outer layer of club-shaped canals, inner layer cellular. *U.Cret.(Maastricht.)*, Mexico-Cuba-Jamaica. —FIG. E270,6. **C. radiolitiiformis* (TRECHMANN), Cuba; 6a, AV transv. sec. at level of AV with projecting teeth and myophores of FV, $\times 4$; 6b, AV transv. sec. of infolding, $\times 3$ (67). [Also Fig. E238,1.]

Dechaseauxia TAVANI, 1949, p. 21 [**D. costata*; M]. AV conical to cylindrical, ornamented by salient longitudinal costae, siphonal bands narrow, marked externally by costate sulci, internally consisting of polygonal cells in regular parallel rows; siphonal zones connected next to shell cavity by 2 rows of polygonal cells; ligamental ridge absent.

FV unknown; genus distinguished by continuous 2-row polygonal wall, siphonal structure and lack of ligament. *U.Cret.* (*Maastricht.*), N.Afr. (*Somalia*).
 —FIG. E239,1. **D. costata*; 1a, lat. view, $\times 0.2$;

1b, transv. sec. viewed toward commissure, $\times 3$;
 1c, AV detail of wall structure, $\times 4.5$ (904).
Durania DOUVILLÉ, 1908, p. 309 [**Hippurites cornupastoris* DES MOULINS, 1827, p. 288; OD]




FIG. E272. Radiolitidae (Sauvagesiinae) (1-2,4), (Lapeirousiinae) (3) (p. N811-N813). [Explanation: *Eb*, *Sb*, siphonal bands; *Eo*, *So*, oscules; *i*, incisure.]

- [=*Duranius* DOUVILLÉ, 1910 (*nom. null.*). AV cylindrical, short or elongate, siphonal bands concave, smooth or ribbed, ligamental ridge absent; bifurcating radial furrows on upper surface of outer wall in many species. FV operculiform. *L. Cret. (Alb.)-U. Cret. (Maastricht.)*, Eu.-N.Afr.-Asia-S.Am.-N.Am.—FIG. E272,4. **D. cornupastoris* (DES MOULINS), *U. Cret. (Turon)*, France; 4*a,b*, FV ext., side view both valves, $\times 0.7$ (911). [Also Fig. E237,9.]
- Hardaghia** TAVANI, 1949, p. 19 [**H. quadrata*; M]. AV pyramidal, depressed toward quadrangular base, ornamented by thick radial ribs; siphonal bands weakly marked externally, internally *Sb* marked by deep sulcus; ligamental ridge lacking. FV flat, siphonal bands deep, narrow grooves; teeth and myophores absent. Distinguished by pyramidal shape and lack of cardinal apparatus. *U. Cret. (Maastricht.)*, E. Afr. (Somalia)—FIG. E272,1. **H. quadrata*; 1*a,b*, FV ext. and lat. view both valves, $\times 0.7$ (904). [Also Fig. E240,3.]
- Tampsia** STEPHENSON, 1922, p. 4 [**T. bishopi*; OD]. AV conical, elongate outer wall thick, cells subrectangular to polygonal, siphons marked by inflexions, anterior *Eb* shallow and extended by narrow depressed slit that cuts outer wall to body cavity; dentition, muscle attachments unknown. FV small bosslike cap, weakly folded over AV shell cavity with thin brim over peripheral surface AV. *U. Cret. (Campan.-Maastricht.)*, Mexico-W. Indies (Cuba)—FIG. E272,2*a*. *Tampsia* sp., Mexico (Cardenas); FV ext., $\times 0.7$ (Coogan, n; courtesy R. Meyers, Univ. Texas)—FIG. E272, 2*b*. **T. bishopi*, Mexico; AV transv. sec. showing *Sb* and *Eb* inflexions and incisure (*i*), $\times 0.7$ (Dechaseaux after Stephenson). [Also Fig. E238, 2.]
- Subfamily LAPEIROUSIINAE Kühn, 1932**
- RV (AV) wall of irregular polygonal cells, ligamental ridge reduced or absent, siphonal zones marked by development of tubular structure (pseudopillars) in outer wall. LV (FV) with oscules marking siphonal openings. *U. Cret. (Santon.-Maastricht.)*.
- Lapeirousia** BAYLE, 1878, p. 110 [**Sphaerulites jouanneti* DES MOULINS, 1827, p. 246; M] [= *Lapeirouseia* KÜHN, 1932 (*nom. van.*)]. AV cylindrical to depressed convex, wall thick, pseudopillars well developed, bulge into shell cavity; externally siphonal bands indented, shell ribbed; ligamental ridge absent. FV operculiform, flat or convex, internally with 2 oscules. *U. Cret. (Santon.-Maastricht.)*, Eu.-Asia-N.Afr.—FIG. E272,3. **L. jouanneti* (DES MOULINS), *U. Maastricht.*, France (Charente); 3*a,b*, side view and FV ext., $\times 0.33$ (911). [Also Fig. E266,2.]
- Dubertretia** COX, 1965, p. 731 [*pro Kellera* MILOVANOVIĆ, 1938, p. 137 (*non* GURNEY, 1928)] [**Dubertretia kelleri* COX (= *Lapeirousia jouanneti* KELLER, 1933, *non* DES MOULINS sp.); OD] [= *Kellera* MILOVANOVIĆ, 1957 (*nom. nud.*)]. AV arched cone, surface with strong folds like *Praeradiolites* which indent half of outer wall; siphonal bands concave, pseudopillars subequal, well developed, ligamental ridge lacking, no lower valve tooth. LV unknown. *U. Cret. (Maastricht.)*, Syria.—FIG. E273,1. **D. kelleri*; 1*a,b*, AV transv. sec. and lat. view, $\times 0.5$ (468). [Also Fig. E237, 7.]
- Katzeria** SLIŠKOVIĆ, 1966, p. 176 [**K. hercegovinaensis*; OD]. AV conical-cylindrical; outer wall nearly without lamellae, those present of radiate structure; inner edge of outer wall has one row of massive prismatic cells next to inner wall; pseudopillars triangular in cross section, with radial and concentric lamellae. FV unknown. May be a bioradiolite. Differs from *Petkovicia* in having prismatic cells on inner edge of outer wall. *U. Cret. (low. Campan.-Maastricht.)*, Eu. (Yugosl.).—FIG. E273,3. **K. hercegovinaensis*; AV transv. sec., $\times 3$ (847*a*).
- Lapeirousella** MILOVANOVIĆ, 1938, p. 89 [**L. orientalis*; OD]. AV conical with more or less regular longitudinal ribs; no ligamental ridge, pseudopillars weakly developed and projecting only slightly, with simpler structure than in *Praelapeirousia*; pseudopillar *Ss* with large prisms at center and small ones at sides. *U. Cret. (Campan.)*, Yugosl. (Serbia)—FIG. E273,6. **L. orientalis*; AV transv. sec. pseudopillar *Ss*; $\times 11$ (626).
- Osculigera** KÜHN, 1932, p. 165 [**O. cleggi*; OD]. AV shaped like *Lapeirousia*, with 2 primary pseudopillars and multiple secondary warty bumps irregularly spaced in outer wall forming small swelling of inner edge where they touch shell cavity, ligamental ridge absent; FV unknown. Distinguished by multiple small pseudopillars. *U. Cret. (Senon.)*, SW. Asia (Iran)—FIG. E273,2. **O. cleggi*; AV transv. sec., $\times 0.5$ (488).
- Petkovicia** KÜHN & PEJOVIĆ, 1959, p. 979 [**P. prima*; OD]. AV high-conical, strongly ribbed, not foliaceous; siphonal bands broad, flattened, separated by furrow on which is 1 rib; outer wall of shell wall very finely cellular, slight internal bulges corresponding to siphonal zones, which are differentiated within wall of shell, narrowing from interior to their outer, convex border. *U. Cret. (Santon.)*, Yugosl. (Serbia)—FIG. E273,5. **P. prima*; 5*a,b*, AV lat. view, and transv. sec. 2 cm. below commissure, $\times 0.75$, $\times 1.3$; 5*c*, drawing of 5*b*, $\times 1.3$ (495).
- Praelapeirousia** WIONTZEK, 1934, p. 28 [**P. rossmati*; M]. AV conical?, siphonal bands externally round projecting ribs, internally of few large cells, not swelling into shell cavity, interband broad; reduced ligamental ridge present; lower valve tooth




FIG. E273. Radiolitidae (Lapeirousiinae) (p. N813, N815). [Explanation: *am*, *pm*, anterior and posterior myophores; *Eb*, *Sb*, siphonal bands; *Es*, *Ss*, pseudopillars.]

absent. FV flat, concentrically layered; ribs few; 2 oscules mark siphonal opening. *U.Cret.(Santon)*, Eu.-N.Afr.—FIG. E273,4. **P. kossmati*; AV transv. sec., $\times 2$ (998).

Vautrinia MILOVANOVIĆ, 1938, p. 86 [**Lapeirousia syriaca* VAUTRIN, 1933, p. 31; OD]. AV conical, ornamented by ribs, no siphonal bands externally, but internally 2 prominent pseudopillars swell inner edge of wall; AV tooth unknown; outer wall of regular radial folds of complex irregular structure. FV operculiform, convex, of compact lamellae, with 2 oscules. *U.Cret.(Maastricht.)*, SW.Asia (Syria-Turkey-Iran).—FIG. E274,2. **V. syriaca* (VAUTRIN), Syria; AV transv. sec. of pseudopillar, $\times 0.5$ (933). [Also Fig. E243,5-6.]

Subfamily UNCERTAIN

Arnaudia FISCHER (ex BAYLE, MS), 1887, p. 1064 [**Hippurites arnaudi* COQUAND, 1859, p. 985; M]. AV conical, ornamented by longitudinal costae; siphonal bands weak broad inflexions internally and externally; ligamental ridge lacking; outer wall thin, compact, detail structure unknown. FV operculiform, concave, ornamented with radial ribs. Long considered a hippuritid, it lacks pillars in lower valve and pores in upper valve. *U.Cret.(Senon.)*, France (Charente).—FIG. E274,3. **A. arnaudi* (COQUAND); 3a,b, side view of shell cluster and transv. sec. at level of AV with pendent teeth and post. myophore of FV, $\times 0.7$, $\times 1.1$ (269, 910).

[*Arnaudia* has been universally classified with the Hippuritidae since BAYLE described it and was so considered by DOUVILLÉ (269) and TOUCAS (910). KÜHN (493) reexamined the type in the École des Mines in Paris and concluded that *Arnaudia* is a radiolitid, probably evolved from *Agriopleura*.]

Colveria KLINGHARDT, 1921, p. 23 [**C. variabilis*; OD]. Shell depressed, both valves with numerous longitudinal ribs, siphonal bands convex; hinge teeth weak, muscle scars on wall of shell; outer layer of AV composed of irregular hollow prisms, that of FV not prismatic but laminated; next layer (pseudocanal layer) in FV penetrating more or less deeply into valve interior and riddled with elliptical cavities which may invade teeth and myophores. Structure of this layer constitutes an essential generic character. *U.Cret.(Maastricht.)*, Eu.(NE.Italy).—FIG. E274,1. **C. variabilis*; lat. view both valves, $\times 0.7$ (475). [Also Fig. E240, 4.]

Joufia BÖHM, 1897, p. 180 [**J. reticulata*; M]. AV conical, nearly smooth, ligamental ridge present, siphonal bands weak inflexions or lacking; outer wall of 2 layers; inner finely reticulate lamellar layer and outer layer with canals. FV oval, cap-like, of 3 shell layers, inner layer of lamellar reticulate structure, middle layer of polygonal canals arranged 4 together with one in center, and outermost layer of concentric laminae; myophores present. Wall structure distinctive. *U.Cret.(Senon.-Maastricht.)*, Eu.(Italy).—FIG. E266,5.




FIG. E274. Radiolitidae (Lapeirousiinae) (2), (Subfamily uncertain) (1,3) (p. N815). [Explanation: am, pm, anterior and posterior myophores; Eb, Sb, siphonal bands; 1, 3, teeth of FV; 2, tooth of AV.]

**J. reticulata*; 5a,b, FV polygonal cells long. and transv. secs., $\times 6.7$ (905). [Also Fig. E240,6.]

UNRECOGNIZABLE RADIOLITID GENERA

Acardo BRUGUIÈRE, 1797, pl. 172 (generic name without nominal species, with illustration). Used by CUVIER in 1830 to include the genera *Radiolites*, *Sphaerulites*, *Hippurites*, *Batolites* and *Calceola*.

Birostrites LAMARCK, 1819, p. 230 [**B. inaequiloba*, an internal mold]. Held by ROQUAN (1841) to

be internal mold of *Sphaerulites* and by TOUCAS (1907-09) of *Praeradiolites hoeninghausi*.




FIG. E275. Family Uncertain (p. N817).

- Bisphaerulites** BAYLE, 1857, p. 680 [*nom. nud.*].
Diplidia MATHERON, 1842, p. 111 [**Diplidia unisulcata*; M]. Poorly known, type species probably represented by mold of *Praeradiolites coquandi* BAYLE.
Jodamia DEFRANCE, 1822, p. 230 [**J. bilinguis*; SD KÜHN, 1932]. Internal mold, possibly of *Praeradiolites* (Toucas, 1907-09).
Laskarevia MILOVANOVIĆ, 1960, p. 371 [*nom. nud.*, generic name without nominal species].
Rajkaia MILOVANOVIĆ, 1960, p. 367 [*nom. nud.*, generic name without nominal species].
Rhytoides ZEKELI, 1854, p. 206 [*nom. nud.*, generic name without nominal species].
Trommia KLINGHARDT, 1935, p. 38 [*nom. nud.*, generic name without nominal species, with figure].

Family UNCERTAIN

[Materials for this section prepared by B. F. PERKINS and A. H. COOGAN]

- Anodontopleura** FELIX, 1891, p. 167 [**A. speciosa*; M]. Externally resembles *Monopleura*; hinge and muscle insertions unknown. *L.Cret.(Neocom.)*, Mexico.—FIG. E275,6. **A. speciosa*; both valves, dorsal view, $\times 0.5$ (301a).
Baryconites PALMER, 1928, p. 51 [**B. multilineatus*; OD]. AV robust, conical, shell wall thick, externally with 2 wide flat siphonal bands separated by narrow ridge, cardinal platform large, anterior muscle insertion large and triangular, large posterior muscle insertion elongate and narrow, anterior tooth socket much larger than posterior, tooth erect and radially elongate, shell cavity small and septate, ligamental groove mostly internal; FV unknown. [This genus has been considered to be related to *Polyconites* and *Horiopleura* but irregularly spaced radial plates in dorsal shell wall suggest a possible relationship to radiolitids.] *U.Cret.(Cenoman.)*, Mexico.—FIG. E275,5. **B. multilineatus*; 5a,b, AV int., post. view, $\times 0.35$ (Perkins, n, courtesy L. G. Hertlein).
Bicornucopina HOFMANN, 1912, p. 219 [**B. petersi*; M] [= *Bicornucaprina* NEAVE, 1939 (*nom. null.*)]. Inequivalve; AV elongate, conical, slightly curved; FV strongly curved; wall of both valves with system of radial canals toward interior and smaller canals toward exterior, as in *Caprinula*; hinge insufficiently known to determine affinities. *L.Cret.(mid-Neocom.)*, Eu.(Hung.).—FIG. E275,2. **B. petersi*; AV ext., $\times 0.5$ (412a).
Cryptaulia POČTA, 1889, p. 50 [**C. triangulum*; SD KUTASSY, 1934, p. 181]. AV straight or slightly curved; longitudinal canals in shell wall; hinge unknown. *U.Cret.(Cenoman.)*, Eu.(Czech.).
Immanitas PALMER, 1928, p. 28 [**I. anahuacensis*; OD]. Equivalve, very large, both valves curved to loosely coiled; surface with 3 ridges separated by furrows, ridges correspond to 3 cavities extending length of both valves near surface; hinge teeth rudimentary, on concave side of shell; shell




FIG. E276. Family Uncertain (p. N817). [Explanation: *am*, *pm*, anterior and posterior myophores; *Eb*, *Sb*, siphonal bands; *L*, ligament; 1, 3, anterior and posterior teeth of FV.]

- cavity small, tabulate; shell wall composed of small, rounded or polygonal prisms often septate near interior. *U.Cret.(Cenoman.)*, Mexico.—FIG. E275,1. **I. anahuacensis*; 1a, both valves, $\times 0.2$ (714); 1b, valve sec., $\times 0.35$ (Perkins, n, courtesy L. G. Hertlein).
Lithocalamus LUPHER & PACKARD, 1930, p. 207 [**L. colonicus*; M]. Known only by AV; gregarious; hinge tooth weak; no accessory cavities; shell wall with 2 layers corresponding to inner and middle layers of *Coralliochama* shell except that cross sections of canals are more rounded and outer row of larger, elliptical canals is present. *U.Cret.(Senon.)*, USA(Ore.).—FIG. E275,3. **L. colonicus*; AV secs., $\times 0.5$ (552).
Palus PALMER, 1928, p. 28 [**P. corrugata*; OD]. Only AV known, straight, with small tooth; muscle insertions superficial; tooth sockets reduced to grooves; shell wall composed of 3 layers, middle one with polygonal canals and possibly accessory cavities. *U.Cret.(Cenoman.)*, Mexico.—FIG. E275,4. **P. corrugata*; 4a,b, AV transv. sec. and lat. view, $\times 1.5$, $\times 1$ (Perkins, n, courtesy L. G. Hertlein).
Pileochama PARONA, 1901, p. 211 [**P. cremai*; M]. AV cylindrical, FV capuliform; resembling *Biradiolites* except for presence of accessory cavity behind each muscle insertion of FV. ?*U.Cret.(Cenoman.-Senon.)*, range poorly defined; Italy (Apennines).—FIG. E276,1. **P. cremai*, transv. sec. AV, $\times 0.7$ (719).
Tepeyacia PALMER, 1928, p. 46 [**T. corrugata*; OD]. AV small, conical, flattened, ornamented externally with angular ribs; siphonal bands *Eb* and *Sb* deep grooves, ligamental ridge present, no myophores observed, hinge area weakly developed; wall of two layers; thin outer finely lamellar layer and thicker massive opaque inner layer. FV unknown. May be a monopleurid. *L.Cret.(?Alb.)*, Mexico-W.Indies(?Cuba-?Jamaica).—FIG. E276,2. **T. corrugata*, Mexico; transv. sec., $\times 0.7$ (714).

Subclass ANOMALODESMATA

Dall, 1889

[nom. transl. et correct. KEEN, 1963 (ex Anomalodesmacea DALL, 1889)] [Diagnosis by N. D. NEWELL]

Relatively short to elongate, nestling to fossorial forms with ventrally fused mantle lobes; equivalve to subequivalve; generally approximately isomyarian; hinge margin characteristically thickened or enrolled, edentulous, or with one amorphous tooth and corresponding socket in one or each valve, lateral teeth absent; ligament opisthodontic or absent, where present associated in many genera with separate resilium and lithodesma; many genera said to have nacreous endostracum; living forms eulamelli-branchiate; a few, septibranchiate. ?*L.Ord.*, *M.Ord.-Rec.*

Recent and many fossil representatives, extending far back into the early Paleozoic (e.g., *Rytimya*, *Cuneomya*, *Sphenolium*) have radial rows of very small tubercles. It may be that this delicate ornament, easily worn away, was present in most members of the subclass. Clearly this is a group of great antiquity and possibly it should include genera placed provisionally in the Cryptodonta and elsewhere. The Anomalodesmata include many of the Desmodonten of NEUMAYR and all of PELSENEER's Septibranchiata.

Order PHOLADOMYOIDA

Newell, 1965

[Diagnosis by N. D. NEWELL]

Ordinal characters same as those of subclass Anomalodesmata. ?*L.Ord.*, *M.Ord.-Rec.*

Superfamily EDMONDIACEA

King, 1850

[nom. transl. NEWELL, 1965 (ex Edmondiidae KING, 1850)] [Materials for this superfamily prepared by N. D. NEWELL]

Equivalve, isomyarian, generally more or less inequilateral; ovoid; ligament opisthodontic, external, short, attached in well-defined groove and supported by strong nymphs and thin hinge plate; essentially edentulous. *U.Dev.-U.Perm.*

Family EDMONDIIDAE King, 1850

Equivalve, evenly convex shells with incurving prosogyre or orthogyre beaks; sur-

face generally without radial ornamentation, nearly smooth or rugose, lunule and escutcheon lacking or obscure; pallial line entire or with very shallow sinus. *U.Dev.-U.Perm.*

Edmondia DE KONINCK, 1841, p. 66 [**Isocardia unioniformis* PHILLIPS, 1936; OD] [= *Allorisma* KING, 1844, p. 315 (type, *Sanguinolaria sulcata* PHILLIPS, 1936; OD); *Aediculus* GISTL, 1848, p. 9 (obj.); *Broeckia* DE KONINCK, 1885, p. 19 (type, *B. latissima*; SD NEWELL, herein); *Pseudedmondia* FISCHER, 1887, p. 1100 (type, *P. puzosi* DE KONINCK, 1842; M); *Edmondiella* CHERNYSHEV, 1950, p. 74 (type, *Sanguinolaria sulcata* PHILLIPS, 1836; OD); *Allorismiella* ASTAFIEVA-URBAJTI, 1962, p. 36 (type, "*Allorisma sulcata* of HIND, 1899" = *Hiatella sulcata* FLEMING, 1828 and *Sanguinolaria sulcata* PHILLIPS, 1836; OD). Ovoid to elongate elliptical, evenly gibbous, without lateral carinae or sulci; margins closed; beaks small, prosogyre, situated one-fourth to one-third behind anterior margin; surface generally with somewhat irregular concentric ridges or growth lines; hinge plate internally reinforced by nearly parallel internal ridge. *U.Dev.-U.Perm.*, cosmop.—FIG. F1,4a. **E. unioniformis* (PHILLIPS), L.Carb., Ire.; LV ext., $\times 1$ (Hind, 1899).—FIG. F1,4b. *E. sp.*, L.Perm. (Getaway), Guadalupe Mts., USA (Tex.); RV int. $\times 1$ (Newell, n).

Cardiomorpha DE KONINCK, 1841, p. 101 [**Isocardia oblonga* SOWERBY, 1825; SD WOODWARD, 1854] [= *Isoculia* M'COY, 1862, explan. pl. 8, fig. 15 (type, *I. corrugata*; M)]. Similar to *Edmondia* but with very prominent, strongly prosogyre umbones and beaks; surface smooth; internal characters unknown. L.Carb., Eu.-N.Am.—FIG. F1,5. **C. oblonga* (SOWERBY), Ire.; 5a-c, LV ext., ant., hinge views, $\times 0.5$ (Hind, 1898).

Superfamily PHOLADOMYACEA

Gray, 1847

[nom. transl. N. D. NEWELL, 1965] [Materials for this superfamily prepared by N. D. NEWELL, with additions as recorded]

Equivalve, ovoid to elongate shells, generally elongate, with simple, external ligament. ?*L.Ord.*, *M.Ord.-Rec.*

Family ORTHONOTIDAE S. A. Miller,

1877

[Materials for this family prepared by N. D. NEWELL and AURÈLE LA ROCQUE]

Soleniform Paleozoic bivalves characterized by concentric folds and posterior gape; lunule and escutcheon generally absent; hinge and internal characters unknown. *M.Ord.-M.Dev.*




FIG. 11. Edmondiidae (4-5); Grammysiidae (1-3,6) (p. N818, N822-N823).

Orthonota CONRAD, 1841, p. 50 [**O. undulata*; SD CONRAD, 1866]. Very elongate, more than 4 times longer than height, gaping slightly posteriorly; beaks about 0.2 length behind front margin; anterior surface with concentric growth lines and, on postumbonal slope, strong concentric undulations and radial folds; ligament enclosed in linear fold on cardinal margin; edentulous. *M.Ord.* (Llandeil.-Caradoc.)-*M.Dev.* (Hamilton.), N.Am.-Eu.—FIG. F2,2. **O. undulata*, *M.Dev.* (Hamilton.), USA (N.Y.); bivalve specimen, composite mold, $\times 1$ (379).

Palaeosolen HALL, 1885, p. 483 [**Orthonota siliquoidea* HALL, 1870; M]. Similar to *Orthonota* but with beaks nearly terminal; postumbonal slope marked by shallow diagonal sulcus and umbonal ridge; surface marked by fine growth lines which may become regular and coarse on postumbonal area; posterior end quadrately truncate, widely gaping; anterior end with narrow gape. *L.Dev.-M.Dev.*, Eu. (Ger.)-USA.—FIG. F2,1. **P. siliquoidea* (HALL), *M.Dev.* (Hamilton.), USA (N.Y.); RV, composite ext.-int. mold, $\times 1$ (379).

Family GRAMMYSIIDAE S. A. Miller, 1877

[=Sanguinolitidae S. A. MILLER, 1877] [Materials for this family prepared by N. D. NEWELL and AURÈLE LA ROCQUE]

Oval to elongate forms characterized by smooth surface or concentric ornamentation;

radial ribs commonly subordinate; ligament elongate; nymphs obscure; lunule and escutcheon commonly present; posterior umbonal ridge well defined in majority of genera; pallial line entire; hinge edentulous. [A poorly known, probably heterogeneous grouping of convenience.] ?*L.Ord.*, *M.Ord.-U.Perm.*

Grammysia DE VERNEUIL, 1847, p. 696 [**G. hamiltonensis* (= **Pterinea bisulcata* CONRAD, 1838); M]. Ovoid, gibbous; lunule and escutcheon well defined; characteristically with concentric folds and 2 oblique radial sulci with intervening fold extending backward from beaks to near middle of ventral margin; edentulous; pallial line simple, radially striate. *L.Dev.-U.Dev.*, cosmop.—FIG. F3, 10. **G. bisulcata* (CONRAD), *M.Dev.*, USA (N.Y.); 10a,b, lat. and front views of RV int. mold, $\times 0.7$ (379).

Alula GIRTY, 1912, p. 3 [**A. squamulifera*; OD]. Externally similar to *Solenomorpha* but with fine radial costellae over surface in front of posterior umbonal carina; dorsal margin gently concave, ventral margin convex; RV with single platelike anterior and posterior tooth with possibly small rounded cardinal tooth below beaks; pallial line unknown. *L.Perm.*, N.Am.—FIG. F3,12. **A. squamulifera*, Lykins F., USA (Colo.); RV ext., $\times 2$ (Girty, 1912).




FIG. F2. Orthonotidae (p. N819).

?*Cimitaria* HALL & WHITFIELD, 1869, p. 66 [**Cypri-cardites recurvus* CONRAD, 1842; SD HALL, 1885]. Shell elongate, subrectangular, falcate or trapezoid; anterior end short and rounded; posterior end elongate, truncate; umbonal ridge angular and commonly strongly defined; surface marked by strong concentric lines of growth and by fine radial costellae; lateral sulcus extends slightly backward from umbones to basal margin; hinge marked by narrow, elongate ligament groove; lunule and escutcheon well defined; internal features and hinge unknown. [COOPER, 1931.] *M.Dev.-U.Dev.*, E.U.S.A.—FIG. F3,7. **C. recurva* (CONRAD), *M.Dev.*(Hamilton.), N.Y.; 7a,b, cardinal view and RV ext., $\times 0.5$ (379).

Cuncemya HALL & WHITFIELD, 1875, p. 90 [**C. miamiensis*; OD] [= *Ceromyopsis* MEEK, 1872, p. 327 (type, *Cardiomorpha? obliquata* MEEK, 1872); *Sphenomya* HALL, 1883, pl. 62 (type, *Grammysia cuneata* HALL, 1883; OD)]. Subrhomboidal, tapering backward for maximum height at prominent, nearly terminal umbones and strongly incurved beaks to obliquely truncate posterior margin; lunule and escutcheon well developed; posterior umbonal ridge rounded and broad; lateral sulcus broad, faint, oblique; anterior surface with uniform concentric ornamentation; posterior slope smooth; without marginal gape. *M.Ord.*(Trenton.)-*M.Dev.*(Hamilton.), N.Am.—FIG. F3,3. **C. miamiensis*, *U.Ord.*, USA (Ohio); 3a,b, hinge view and RV ext., $\times 1$ (381).

?*Davidia* HICKS, 1873, p. 49 [**D. ornata*; SD NEWELL, herein]. Subtriangular, with submedian beaks and rounded, nasute, anterior and posterior extremities; posterior and anterior umbonal carinae near cardinal border; surface smooth except for faint radial costellae over ?posterior end of shell. *L.Ord.*(Tremadoc.), Wales.—FIG. F3,4. **D. ornata*, St. Davids; 4a, ?LV ext., $\times 1$; 4b, surface, enl. (406).

?*Glossites* HALL, 1885, p. 49 [**G. lingualis*; OD] [= ?*Elymella* HALL, 1885, p. 50 (type, *E. nuculoides*; OD)]. Shell elliptical, beaks anterior, small, appressed; hinge line long, gently arcuate; posterior umbonal ridge rounded; surface smooth or sublamellose, with or without faint radial costellae; lunule and escutcheon distinct; muscle scars shallow. *L.Dev.*(Chapman Ss.), USA (Maine)-*U.Dev.* USA(N.Y.)-Eu.-Afr.; *L.Miss.*, USA (Iowa-Ohio). —FIG. F3,6. **G. lingualis*, *U.Dev.*(Chemung), USA(N.Y.); RV ext., $\times 1$ (379).

Grammysioidea WILLIAMS & BREGER, 1916, p. 133 [**G. princiana*; OD]. Similar to *Grammysia*, but lacks regular concentric ornamentation and either lacks radial sulcus or shows only single shallow lateral sulcus. *Sil.*(Clinton.)-*U.Dev.*, USA (Nev.)-E.N.Am.-S.Am.-Eu.—FIG. F3,5. **G. princiana*, *L.Dev.*(Moose R. Ss.), USA(Maine); RV ext., $\times 0.5$ (986).

?*Leinzia* MENDES, 1949, p. 11 [**Solenomorpha similis* HOLDHAUS, 1918; OD]. Elongate pteroid, with anterior auricle; strong umbonal carina in front of narrow, flattened, smooth or carinate siphonal area; body of shell in front of umbonal carina with coarse, concentric ornamentation; LV with single cardinal socket; laterals lacking. *Perm.*, S.Am. (S.Brazil).—FIG. F4,5. **L. similis* (HOLDHAUS), Passa Dois F.; LV ext., $\times 1$ (Mendes, 1949).

?*Palaeocorbula* REED, 1932, p. 63 [**P. difficilis*; M]. Moderately inflated, strongly inequivalve, RV smaller than LV, suboval, longer than high; anterior end projecting, sharply rounded; umbones submedian; surface ornamented with numerous, regular, sharp, equidistant, concentric ridges, some of which abruptly thin out or coalesce at weak anterior sulcus in front of which concentric ridges are only about half as numerous; lunule lacking; escutcheon well defined, narrow; interior unknown. *L.Perm.*, India(Kashmir).—FIG. F3,9. **P. difficilis*, agglomeratic slates, 9a,b, lat. and post. views of RV, $\times 1$ (Reed, 1932).

?*Pholadella* HALL, 1869, p. 63 [**P. newberryi*; SD HALL, 1885] [= *Ceromyopsis* MEEK, 1871, p. 71 (type, *Allorisma pleuropistha* MEEK, 1871; OD); *Grammysiopsis* CHERNYSEV, 1950 (type, *G. irregularis*; OD)]. Shell elongate, gibbous; beaks about 0.25 distance behind anterior margin; umbones and umbonal ridge prominent; anterior end rounded or obliquely truncate; laterally compressed posterior truncate; broad, lateral sulcus extending directly downward from beaks to ventral indentation; surface marked by strong concentric undulations which may be confined to anterior part of shell; distinct costae originating at umbo extend across central or greater portion of shell body, leaving anterior and cardinal area free of radial ornamentation; cardinal margin marked by broad ligament groove, lunule and escutcheon well defined; hinge, pallial and adductor impressions unknown. *L.Dev.*(Coblentz.), Ger.; *L.Miss.*, E.U.S.A.; *M.Penn.*-

U.Penn., C.USA; *U.Carb.*, USSR(Urals).—FIG. F3,11. **P. newberryi*, L.Miss.(Waverly.), USA (Ohio); LV ext., X1 (379). ?*Promacrus* MEEK, 1871 p. 4 [**P. nasutus*; OD]. Anterior end attenuated and produced; posterior

end higher and obliquely truncated; beaks ranging in position from median to slightly anterior or posterior; umbonal ridge well defined; without lunule and escutcheon; surface smooth or in some shells with radial costellae over umbonal region;




FIG. F3. Grammysiidae (p. N819-N823).




FIG. F4. Grammysiidae (5); Megadesmidae (1-4) (p. N820, N825-N827).

hinge and pallial line unknown. *L.Miss.*, N.Am.
 —FIG. F1,6. **P. nasutus*, *L.Miss.*(Chouteau), USA(Mo.); LV ext., $\times 0.6$ (Meek, 1872).

?*Prothyris* MEEK, 1871, p. 4 [**P. elegans*; OD] [= *Prothyris* MEEK, 1869, p. 172 (*nom. nud.*)]. Elongate-oblong, smooth or weakly costellate; gaping slightly behind, or closed and widely gaping in front, where hiatus is increased by notch; beaks depressed and very near anterior end, with small ridge extending from anterior side of each beak to corner of anterior notch; dorsal margin without escutcheon or lunule; pedentulous. *L.Dev.-U.Perm.*, W.Eu.-N.Am.-S.Am.(Brazil).

P. (Prothyris). Without posteroventral notch or sulcus. *L.Dev.-U.Perm.*, W.Eu.-EUSA-C.USA.

—FIG. F1,2. **P. (P.) elegans* MEEK, U.Penn. (Willard Sh.), USA(Neb.); RV ext., lectotype; $\times 2$ (Newell, n).

P. (Paraprothyris) CLARKE, 1913 [**P. (P.) knodi*; M]. Similar to *P. (Prothyris)* but with broad posterior sulcus ending in marginal notch, surface with 2 sets of concentric ornamentation forming herringbone pattern below posterior sulcus. *L.Dev. (Ponta Grossa Sh.)*, E.Am.(Brazil). —FIG. F1,1. **P. (P.) knodi*; bivalved individual, LV below, $\times 1$ (138).

?*Protomya* HALL, 1885, p. 52 [*pro Palaeomya* HALL, 1885, p. 52 (*non* ZITTEL & GOUBERT, 1861)] [**P. oblonga*; OD]. Elongate, oval; beaks submedian, incurved, prominent; umbonal slope




FIG. F5. Morphological features of *Megadesmus*.—A, B. *M. nobilissimus*, semidiagrammatic long. sec. of LV showing shell layers, $\times 1.5$, and long. sec. of inner layer showing lamellae secreted around sand grain, $\times 7.5$.—C, D. *M. globosus*, superposed serial secs. of hinge viewed from post. end (C) and ant. view of umbonal secs. of LV showing socket (D), $\times 1$.—E. *M. nobilissimus*, transv. sec. of ligament. [Explanation: a, b, outer and inner shell layers; c, periostracum; d, beak; e, tooth; f, socket; g, nymph; h, ligament groove; i, j, k, inner, outer, and periostracal layers of ligament.] (Runnegar, 1965).

gibbous above; surface with fine growth lines and strong concentric undulations; cardinal line long, nearly straight; ligament external; hinge and pallial line unknown. *M.Dev.*, USA (N.Y.).—FIG. F3,8. **P. oblonga*, Hamilton, N.Y.; LV int. mold, $\times 0.7$ (379).

Sanguinolites M'COY, 1844, p. 47 [**S. discors*; SD STOLICZKA, 1871] [= *Cercomyopsis* SANDBERGER, 1881, p. 247 (type, *C. acutirostris*; OD); *Sphenotus* HALL, 1885, p. 33 (type, *Sanguinolites arcaiformis* HALL & WHITFIELD, 1869; SD S.A. MILLER, 1889); ?*Eopleurophorus* ELIAS, 1957, p. 780 (type, *Cypricardia? tricostata* PORTLOCK, 1843; OD)]. Shell elongate; umbones small, gibbous, incurved and contiguous, placed in anterior 0.2 of shell; anterior convex part of shell with coarse concentric ornamentation, sharply set off from flattened post-umbonal area by 1 or more radial carinae; lunule and escutcheon well developed; pallial line entire. *U.Dev.-Perm.*, cosmop.—FIG. F3,1. **S. discors*, L.Carb., Ire.; RV ext., $\times 1$ (M'COY, 1844).

Sedgwickia M'COY, 1844, p. 61 [**S. attenuata*; SD STOLICZKA, 1871] [= ?*Fuchsella* DAHMER, 1942, p. 145 (type, *Sphenotus maillicuxi* ASSELBERGHS, 1930; M)]. Elongate, more or less gibbous; anterior end rounded, posterior edge subtruncate; umbones prominent, placed well forward, slightly proso-

gyre; surface smooth or with low concentric ridges; defined ligament nymphs behind beaks; lunule, escutcheon and radial carinae lacking; hinge unknown, ?edentulous. [The relationship between this genus and *Myonia* DANA, 1847, is uncertain.] *Miss.-Perm.*, cosmop.—FIG. F1,3. **S. attenuata*, L.Carb., Marsden, Eng.; RV ext., $\times 1$ (M'COY, 1844).

Solenomorpha COCKERELL, 1903, p. 559 [*pro Solenopsis* M'COY, 1844, p. 47 (non WESTWOOD, 1840)] [**Solenopsis minor* M'COY; M]. Subcuneiform, tapering posteriorly from maximum height at beaks near anterior end; anterior end closed; posterior end gaping slightly; umbonal carina conspicuous; narrow lunule and escutcheon present; surface smooth, without radial ornamentation; hinge unknown; pallial line entire. *L.Dev.-U.Perm.*, cosmop.—FIG. F3,2. **S. minor* (M'COY), L.Carb., Ire.; LV ext., $\times 1$ (M'COY, 1844).

Family MEGADESMIDAE Vokes, 1967

[*nom. correct.* NEWELL, herein (*pro Megadesmatidae* VOKES, 1967, *nom. subst. pro Pachydomidae* FISCHER, 1886, based on invalid generic name, ICZN Code, Art. 11e)]

[Materials for this family prepared by N. D. NEWELL]

Similar to Edmondiidae but many genera more clearly adapted for burrowing; liga-




FIG. F6. Musculature and pallial lines shown on internal molds of early to late species of *Megadesmus* and *Astartila* (stratigraphic ranges indicated by heavy vertical lines).—A. *M. globosus*.—B. *M. nobilissimus*.—C. *M. gryphoides*.—D. *A. intrepida*.—E. *A. elegans*.—F. *M. grandis*. [Explanation: 1, ant. adductor; 2, pedal protractor; 3, ant. pedal retractor; 4, umbonal pedal retractor; 5, post. pedal retractor; 6, post. adductor.] (Runnegar, 1965).

ment short, nymphs very prominent with a single blunt tooth in RV and rarely in LV. *U.Carb.-Perm.*

Some of the more important features of shell structure, hingement, and musculature are illustrated in Figures F5-7, based on studies by RUNNEGAR (1965, 805a).

Megadesmus SOWERBY, 1839, p. 14 [**M. globosus*; SD WOODWARD, 1854, p. 262] [= *Pachydomus* MORRIS, 1845, p. 271 (*nom. van. pro Megadesmus* SOWERBY, 1839, *non Megadesma* BOWDICH, 1822); *Cleobis* DANA, 1847, p. 154 (type, *Cleobis grandis* DANA, 1847; SD NEWELL, 1956, p. 10)]. Large

equivalve, asymmetrical shells ornamented with coarse to medium imbricate growth lamellae and fine to medium concentric ribs; with sporadic obscure radial costae; micro-ornament consisting of radial rows of minute papillae, hinge with dental process in RV produced by fold in dorsal anterior valve margin, and socket in LV resulting from depression of valve edge; ligament external, opisthodontic, parivincular, supported by strong nymphs set in well-defined escutcheon; adductors subequal, the anterior smaller; 2 pedal muscle scars (protractor and retractor) situated above anterior adductor, and single retractor adjoining posterior adductor; third retractor scar in and on anterior side

of umbonal cavity; pallial line simple, radially striated; shell inside pallial line commonly pitted by insertions of mantle; shell and ligament of 3 layers: periostracum, outer lamellar layer and inner (?nacreous) layer (Runnegar, 1965). *Perm.*, Australia-Tasmania-India-Argentina.—FIG. F8, 4a-c. **M. globosus* (SOWERBY), L.Perm. (Allandale F.), Harper's Hill, Hunter Valley, New S. Wales; 4a, LV view of shells, $\times 0.5$ (Sowerby, 1838); 4b, dorsal view of lectotype, $\times 0.5$ (Runnegar); 4c, RV ext., $\times 0.5$ (805a).—FIG. F8, 4d. *M. gryphoides* DE KONINCK, E.Australia; latex cast of interior showing that front of dorsal margin slightly overlaps that of LV, ant. part of which fits into shallow groove in front of LV tooth, $\times 0.5$ (Runnegar, n).

Astartila DANA, 1847, p. 155 [**A. intrepida*; SD STOLICZKA, 1871, p. xix]. Small, ovate shells with gently convex posterior umbonal slopes and low, forwardly directed beaks; ornament varying from regular imbricate growth lamellae to irregularly spaced concentric threads; radial ornament of low, wide costae may be present; shell surface covered with radial rows of minute papillae; ligament external and opisthodontic, attached behind well-developed nymphs set in narrow escutcheon; lunule usually absent; RV with single upright tooth derived from fold in valve margin; LV with deep to shallow socket of type found in *Megadesmus* specimens with well-developed radial ornament having weakly developed megadesmid dentition; adductor scars subequal, the anterior smaller and deeper, commonly having several lobes on the posterior-dorsal edge; a single, deep, oval pedal scar occurring above anterior adductor and a large pedal retractor scar attached to upper edge of posterior adductor; a third pedal scar on rear side of umbonal cavity; pallial line simple and radially striate. L.Perm., Australia.

[*Astartila* differs from *Megadesmus* in the following respects. 1) Because of the narrow, deep escutcheon, the upper margin of the valve behind the umbo is gently convex. In *Megadesmus* the escutcheon is broader and less steeply inclined so that it is mechanically possible for the umbones to project above the posterior margin of the valves. 2) Only one pedal scar occurs above the anterior adductor. The scar that is present is ovoid and deeply impressed so that it is almost certainly the scar that has been termed the anterior pedal retractor in *Megadesmus*. In one specimen of *Astartila intrepida* there is a differentiation of the edge of the adductor which may represent the protractor scar. However, as the edge of the adductor is commonly lobed, it is probable that this is merely an exaggerated development of a single lobe. The absence of a scar does not necessarily imply the absence of a pedal protractor muscle since BLOOMER (1901, p. 40) has described a species of *Solen* in which the protractor muscle is attached to the anterior adductor and not to the shell. If this were true in species of *Megadesmus* or *Astartila* it may account for the more or less complete merging of the protractor and adductor scars (RUNNEGAR, 1965, 805a).]

A. (*Astartila*). Species lacking well-developed radial ornamentation. L.Perm., Australia.—FIG. F8, 3. **A. (A.) intrepida* DANA, U.Marine, Wollongong, New S. Wales; 3a-c, RV ext., latex cast of hinge int., shell front, $\times 1$; 3d, hinge ext., $\times 0.7$ (668).

A. (*Pleurikodonta*) RUNNEGAR, p. 247, 1965 [**P. elegans*, OD]. Internally like *A. (Astartila)* but ornamented with obscure radial costae which are reflected by inflections in growth lamellae and




FIG. F7. Muscle scars and pallial lines on internal molds of representative Megadesmidae and Pholadomyidae (Runnegar, 1965).—A. *Astartila (Astartila) intrepida* DANA.—B. *A. (Pleurikodonta) elegans* RUNNEGAR.—C. *Chaenomya leavenworthensis* (MEEK).—D. *Megadesmus gryphoides* (DE KONINCK).—E. *Chaenomya?* sp.—F. *Pyramus laevis* (SOWERBY).—G. *Myonia valida* DANA.—H. *Vacunella* sp. cf. *V. waterhousei* (DUN).—I. *Pholadomya gigantea* (SOWERBY).—J. *Vacunella curvata* (MORRIS). [Explanation: a, muscles of uncertain significance; ar, anterior retractor; ava, accessory ventral adductor; g, siphonal gape; p, protractor; pd, pedal gape; pr, posterior protractor; ur, umbonal retractor.]




FIG. F8. Megadesmidae (p. N824-N826).

crenulations in valve margins; megadesmid dentition of *A. (Astartila)* wholly or partially replaced by costae which interlock in front of beaks. *L.Perm.*(*M.Bowen Beds*), Queensl.—FIG. F4,3. **A. (P.) elegans* (RUNNEGAR); 3a-c, ventral, ant. and oblique views of latex cast, $\times 1$ (Runnegar, 1965).

Casterella MENDES, 1952, p. 100 [**C. gratiosa*; M]. Similar to *Megadesmus* but with pallial sinus and thin hinge plate. *Perm.*(*Corumbatai F.*), S.Am. (S.Brazil).—FIG. F8,2. **C. gratiosa*; 2a,b, RV ext. and hinge views, $\times 1.5$ (Mendes, 1952).

?*Crassiconcha* NECHAEV, 1894, p. 316 [**C. stuckenbergi*; M]. Equivalve; smooth; beaks submedian; lunule and escutcheon well marked; front end projecting, narrowly rounded; lateral surfaces divided approximately into thirds by broad shallow sulci; deep pallial sinus; hinge unknown. *U.Perm.*, Eu.(E.USSR)-E.Greenl.—FIG. F8,1. **C. stuckenbergi*, Kazan., USSR; 1a, hinge view and 1b, ext., $\times 0.7$ (Nechaev, 1894).

Myonia DANA, 1847, p. 158 [**M. elongata*; SD FLETCHER, 1932] [= *Maeonia* DANA, 1849 (*nom. van. pro Myonia* DANA, 1847)]. Shell elongate,

lenticular, with small umbones; tapering backward as viewed from above and side; posterior end obliquely truncate; obscure sulcus extending from beaks to ventral margin, usually bounded behind by posterior umbonal ridge; cardinal teeth lacking or obscure; margins not gaping; pallial line continuous, but radially striate; mantle within pallial line attached to shell by scattered deep pits. *U.Carb.-Perm.*, New S. Wales-Kashmir-S. Am. (Brazil)-USSR (NE.Sib.).

[*Myonia* is placed in the Megadesmidae rather than in the Edmondiidae because of its very strong outward rolled nymphs; at least some examples (Fig. F4,2b) show the usual cardinal boss characteristics of the subfamily.]

M. (Myonia). Umbonal ridge rounded or with weak carina. *U.Carb.-Perm.*, New S. Wales-Brazil-Sib.—FIG. F4,2a. **M. (M.) elongata* DANA, Gerringong, New S. Wales; LV ext., $\times 0.34$ (668). —FIG. F4,2b. *M. (M.) carinata*, U. Marine Ser., south coast, New S. Wales; RV int., $\times 0.4$ (Newell, n).

M. (Pachymyonia) DUN, 1932, p. 411 [**P. morrisii* ETHERIDGE, 1919; OD]. Umbonal ridge very strong, angular. *Perm.*, New S. Wales.—FIG. F4,1. **M. (P.) morrisii* (ETHERIDGE), L. Marine Ser., New S. Wales; 1a,b, RV ext., hinge view, $\times 0.5$ (Newell, n).

Pyramus DANA, July 1847, p. 156 [**P. myiformis*; SD NEWELL, 1856] [= *Notomya* M'COY, Nov. 1847, p. 303 (type, *N. securiformis*; SD STOLICZKA, 1871); *Pyramia* DANA, 1849 (*nom. van. pro Pyramus* DANA, 1847); *Clarkia* DEKONINCK, 1877 (obj.)]. Similar to *Myonia* but with slight posterior gape and shallow pallial sinus; pallial line not striate and interior of shell not pitted within pallial line; 1 large tooth in each valve below beaks. *L.Perm. (U. Marine)*, New S. Wales; *Perm.?* (*Corumbatai* F.), S. Am. (S. Brazil). —FIG. F4,4. **P. myiformis*, L. Perm. (U. Marine), Blackhead, New S. Wales; 4a,b, latex cast viewed from right; int. same, RV; both $\times 1$ (668).

Family PHOLADOMYIDAE Gray, 1847

[*nom. correct.* KING, 1850 (*pro Pholadomyidae* GRAY, 1847)] [= Arcomyidae FISCHER, 1887] [Materials for this family prepared by L. R. COX, with additions by N. D. NEWELL]

Shell equivale (except in *Girardotia*), with wide range of size, ovate, oblong or subtrigonal, subequilateral to (more commonly) strongly inequilateral, moderately to strongly inflated; most forms gaping at posterior end and some with narrower anterior gape; ostracum thin (except in some *Pachymya* and in *Machomya*), nacreous, although commonly altered in fossilization; surface bearing small pustules, aligned radially or concentrically in many forms. Hinge edentulous; cardinal margin thickened below and behind beak and bearing

strong nymph in each valve for support of external, opisthodontic ligament; small internal ligament (resilium) also present below beak in Recent type species of *Pholadomya*, but not observed in any fossil forms, internal characters of most being inadequately known; pallial sinus present in most forms but absent in a few. *Miss.-Rec.*

FISCHER recognized two distinct families, Pholadomyidae and Arcomyidae, supposing the fine surface pustules present in the latter group to be absent in the former. It is now known that the surface is finely pustulose in some Lower Tertiary *Pholadomya* (Fig. F9,1f), however, and it seems probable that this delicate ornament, which is easily worn away, was present both in the holotype of the Recent type species and in numerous fossil representatives of the genus.

Pholadomya G.B. SOWERBY, 1823 [**P. candida*; SD GRAY, 1847, p. 194] [= *Pholadomya* FLEMING, 1828 (*nom. null.*); *Pholadomia* SWAINSON, 1835 (*nom. null.*); *Phaladomya* MEEK, 1858 (*nom. null.*); *Pholadomye* DESHAYES, 1860 (*nom. null.*); *Pholdomya* CONRAD, 1865 (*nom. null.*); *Pholodomya* HECTOR, 1886 (*nom. null.*); *Pholadomya* DALL, 1905 (*nom. null.*)]. Medium-sized to large, ovate to subtrigonal, strongly inequilateral, ventricose, most so anteriorly; umbones broadly rounded to subangular, of variable prominence, more or less anteriorly placed; valves gaping posteriorly to variable extent, anterior gape narrow or absent; pustulation of surface very delicate; ornament usually strongly developed, but weak in some forms and consisting of radial ribs or ridges, commonly bearing tubercles, and of concentric undulations or rugae; pallial sinus broad, of moderate depth. *U.Trias.-Rec.*, cosmop.

[Although several Recent species from various parts of the world have been described under *Pholadomya*, the type species is the only one resembling the numerous fossil forms in size and shape. The remainder are relatively small shells, some of which belong to the genus *Panacca* DALL (see below), while some others, particularly certain very small forms from the Antarctic, should probably be included in one or more new genera. All known Recent Pholadomyidae are moderately deep- to deep-water forms, whereas the Mesozoic species of the family are found commonly in shallow-water sediments. No satisfactory subgeneric classification of the numerous species of *Pholadomya* has yet been achieved and a thorough revision of the genus would result in the recognition of many more subgeneric groups than those here distinguished.]

P. (Pholadomya) [= *Flabellomya* ROLLIER, 1911, p. 231 (type, *Lutraria ambigua* J. SOWERBY, 1819, p. 48; SD COX, herein)]. Elongate-ovate, with broad and not prominently protruding umbones; dorsal umbonal ridge, bordering an escutcheon, present or absent. *U.Trias.-Rec.*, cosmop.—FIG. F9,1a-d. **P. (P.) candida* G.B. SOWERBY, Rec., Carib.; 1a-c, holotype, RV ext., RV int. (showing pallial line), and dorsal view of both valves (all




FIG. F9. Pholadomyidae (p. N827-N829).

×0.7); *1d*, hinge region of RV showing ligamental nymph and shallow rhombic subumbonal pit for internal ligament, small, obtusely angular projection of margin below it (×2) (Cox, n). —FIG. F9, *1e*. *P. (P.) ambigua* (J. SOWERBY), L. Jur. (Lias.), Eng.; LV ext., ×0.7 (Cox, n). —FIG. F9, *1f*. *P. (P.) leonensis* STENZEL, KRAUSE & TWINING, Eoc., USA (Tex.); pustulose ornament, ×4 (Stenzel, Krause & Twining, 1957).

P. (Bucardiomya) ROLLIER in COSSMANN, 1912, p. 215 [*pro Bucardia* ROLLIER, 1911, p. 231 (non SCHUMACHER, 1817)] [**Pholadomya bucardium* AGASSIZ, 1842, p. 77; SD Cox, herein] [= *Bureiomya* VORONETZ, 1937, p. 58 (type, *Pholadomya polymorpha* D'ORBIGNY, 1850, p. 360; SD Cox, herein)]. Subtrigonal or obliquely oval, not greatly elongated; umbones prominent and subangular in most species; no escutcheon; strength

of radial ribbing very variable. *L. Jur. (Toarc.)-L. Tert.*, cosmop.—FIG. F10,1a. **P. (B.) bucardium* AGASSIZ, M. Jur. (Bathon.), Switz.; LV ext., $\times 0.7$ (Moesch, 1874).—FIG. F10,1b,c. *P. (B.) lirata* (J. SOWERBY), M. Jur. (Bathon.), Eng.;

1b,c, LV ext., both valves dorsal, $\times 1$ (Cox, n). —FIG. F10,1d. *P. (B.) margaritacea* (J. SOWERBY), U. Eoc., Italy; RV ext., $\times 1$ (Moesch, 1874).

P. (Procardia) MEEK, 1871, p. 184 [**Isocardia?*]




FIG. F10. Pholadomyidac (p. N828-N830).




FIG. F11. Pholadomyidae (p. N830-N832).

hodgei MEEK, 1871, p. 183; M]. Relatively short, subtrigonal or oblong, with prominent umbones and strongly incurved, prosogyrous beaks; anterior end short, truncated, flattened or impressed, with still more impressed lunule under beaks; escutcheon distinctly bordered; flank ornamented with narrow radial costae and concentric ridges. *U. Jur. (Callov.)-U. Cret. (Turon.)*, cosmop.—FIG. F10,2. *P. (P.) decussata* (MANTELL), *U. Cret.*

(Cenoman.), Eng.; 2a-c, RV ext., both valves dorsal and ant. views, $\times 1$ (Woods, 1909). *Anomalopleuroides* COX, 1962 [*pro Anomalopleura* LEONARDI, 1948, p. 62 (non KLEINE, 1916)] [**Anomalopleura elisae* LEONARDI, 1948; M]. Small-medium sized, elongate-ovate, fairly strongly inflated, with wide posterior gape; umbones obtusely rounded, scarcely protruding, placed at about anterior third of length; posterior two-thirds of shell




FIG. F12. Pholadomyidae (p. N831).

ornamented with narrow, closely spaced radial ribs, anterior third with more widely spaced, oblique, steep ribs which meet last rib of radial series in acute angle; internal characters unknown. *U.Trias.*, Austria(S.Tirol).—FIG. F11,6. **A. elisae* (LEONARDI); LV ext., $\times 1.3$ (Leonardi, 1948).

Argyromya FISCHER, 1887, p. 1166 [**Glycimeris margaritacea* LAMARCK, 1818, p. 458; OD (cited as "*Panopaea margaritacea* DESHAYES")] [= *Aigyromya* DONCIEUX, 1911 (*nom. null.*)]. Medium-sized, oblong, subequilateral, moderately inflated, most so posteriorly, where very wide gape occurs; anterior gape narrow; umbones narrow, scarcely protruding; surface with irregular growth rugae; pustules minute, radially aligned, seen only in earlier growth stages; pallial sinus narrow and deep. *M.Eoc.*, Eu.(France).—FIG. F11,1. **A. margaritacea* (LAMARCK); 1a-c, RV ext., RV int., and dorsal view of both valves, all $\times 1$ (Deshayes, 1856).

Chaenomya MEEK, 1864, p. 42 [**Allorisma? leavenworthensis* MEEK & HAYDEN, 1858; M]. Shell smooth or ornamented with concentric folds; umbones prominent, situated about 0.25 of shell length from anterior margin; subcylindrical at beaks; without lateral sulcus or ventral sinus; siphonal gape extreme, equal to cross section of shell, or flaring slightly; pallial sinus deep. *Penn.-Perm.*, USA(Mid-Cont.); *Perm.*, Australia.—FIG. F12,2. **C. leavenworthensis* (MEEK & HAYDEN), U.Penn.; 2a,b, LV lat. and hinge view of both valves, $\times 1$; 2c, detail of shell surface, enl., (Meek, in Meek & Hayden, 1864). [NEWELL]

Cortinia LEONARDI, 1948, p. 63 [**C. catharinae*; M]. Small, obliquely oval, inequilateral, *Pholadomya*-like but with obtuse ridge running from umbo toward posteroventral angle; ornamented with pustulose radial threads and with undulations which follow growth lines or are slightly oblique to them; internal characters unknown. *U.Trias.*, Austria(S.Tirol).—FIG. F11,2. **C. catharinae*; RV ext., $\times 2$ (Leonardi, 1948).

Cosmomya HOLDHAUS, 1913, p. 446 [**C. egregia*; M]. Medium-sized, moderately inflated, with broadly rounded, well protruding, rather anteriorly placed umbo; ornament consisting of posterior series of steep, oblique ribs, some zigzagging posteriorly and bending up in V anteriorly, and of anterior series of ribs which are only slightly oblique to growth lines and meet the posterior series along shallow groove descending from umbo to ventral margin; internal characters unknown. "Below Werfen beds," i.e. pre-Lower Trias.¹ Asia(Niti Pass, Himalayas), (Dickins & Shah, 1965).

Exochorhynchus MEEK & HAYDEN, 1864, p. 42 [**Allorisma? altirostrata* MEEK & HAYDEN, 1858; M]. Similar to *Wilkingia* but with terminal beaks; hinge and interior unknown. *Miss.-Penn.*, W.Eu.-N.Am.—FIG. F12,1. **E. altirostratus*, U.Penn., USA(Kans.), LV ext., $\times 0.7$ (Meek & Hayden, 1864). [NEWELL]

¹ The type species of *Cosmomya*, thought by HOLDHAUS to be derived from Jurassic rocks, is probably Permian and seems to be congeneric with *Palaeocosmomya*, fide J. M. DICKINS (ed.).

- Girardotia** DE LORIOI, 1903, p. 133 [**G. elegans*; M]. Small to medium-sized, oval, slightly to moderately inequilateral, gibbose, with wide posterior and narrow anterior gape; slightly inequivalve, RV smaller; umbones broadly rounded, scarcely protruding; RV with radial groove, LV with ridge in some species and groove in others, running from umbo to posterior part ventral margin; ornament consisting of narrow concentric ribs present on entire surface or on anterior part only, and of concentric ridges; internal characters unknown. *M.Jur.-U.Jur.(Bajoc.-Kimmeridg.)*, Eu.—FIG. F11,3. **G. elegans*, U.Jur.(Oxford.), France; 3a,b, RV ext., and dorsal view of both valves showing gape, both $\times 1$ (de Loriol, 1903).
- Goniomya** AGASSIZ, 1841, p. 275 [**Mya angulifera* J. SOWERBY, 1819, p. 46 (= **Mya intersectans* SMITH, 1817, p. 92; SD HERRMANNSEN, 1847, p. 486)] [= *Gonomya* AGASSIZ, 1838 (nom. nud.); *Gonyomya* HAUER, 1853 (nom. van.); *Gonimya* NECHAEV, 1894 (nom. null.); *Goniomya* PCHELINTSEV, 1924 (nom. null.)]. Small to medium-sized, ovate, moderately to strongly elongate and inequilateral, with broadly rounded to subangular umbones protruding slightly above dorsal margin; beaks almost orthogyrous; inflation moderate; posterior gape relatively wide, anterior gape narrow; diagonal ridge obtuse, rounded off; escutcheon shallow, its bordering ridges ill-defined except near umbones; anterior and posterior parts of surface with discordant ribbing; hinge structure unknown; pallial sinus present. *L.Jur.(L.Lias.)-Eoc.*
- G. (Goniomya)** [= *Lysianassa* MÜNSTER, 1838 (non EDWARDS, 1830); *Rhombomya* SACCO, 1901, p. 133 (type, *Lysianassa rhombifera* GOLDFUSS, 1840, p. 264)]. Ornament consisting of anterior and posterior series of steep oblique ribs inclined toward one another; in some species these meet along line from umbo to ventral margin forming series of V's, and in others they are separated by intervening rib parallel with growth lines or by almost smooth area. *L.Jur.(L.Lias.)-U.Cret.(Turon.)*, cosmop.; *Eoc.*, Eu.(Italy).—FIG. F11,4. **G. (G.) intersectans* (SMITH), M.Jur.(Bathon.), Eng.; 4a,b, LV ext., dorsal of both valves, $\times 1$ (Cox, n).
- G. (Deltamya)** BURMEISTER, 1914, p. 21 [**D. jboehmi*; M]. Ribs on posterior and median parts of surface parallel with growth lines, those on anterior part slightly oblique to them but weak. *U.Cret.(Senon.)*, Eu.(Ger.).—FIG. F11,5. **G. (D.) jboehmi* (BURMEISTER); LV ext., $\times 1$ (Burmeister, 1914).
- Homomya** AGASSIZ, 1843, p. 154 [**Maetra gibbosa* J. SOWERBY, 1813, p. 91; SD HERRMANNSEN, 1847, p. 541]. Resembling *Pholadomya* (*Pholadomya*) in shape except that in some species umbones are less prominent; without radial ribbing, unless in earliest stages of growth; hinge margin smooth, uninterrupted; pallial sinus deep. *M.Trias.(Muschelkalk)-U.Jur.(Poriland.)*, cosmop. [The commoner Muschelkalk species which have been referred to this genus seem better included in *Pachymya* (*Arcomya*).]—FIG. F13,3. **H. gibbosa* (SOWERBY), M.Jur.(Bathon.), Eng.; 3a,b, LV ext., both valves dorsal (holotype), $\times 0.7$ (Cox, n).
- Machomya** DE LORIOI, 1868, p. 517 [**Panopaea dunkeri* D'ORBIGNY, 1850, p. 47 (pro *Solen jurensis* DUNKER, 1847, p. 131, incorrectly treated by D'ORBIGNY as a secondary homonym of "*Panopaea jurensis* BRONGNIART, 1821," presumably a mistake for *Mya? jurassi* BRONGNIART, 1821); M]. Resembling more compressed and elongate species of *Pachymya* (*Arcomya*) in external characters; interior of each valve with low oblique rib near anterior margin; ostracum moderately thick; details of hinge and pallial line unknown. *M.Jur.-U.Jur.(Bathon.-Kimmeridg.)*, Eu.—FIG. F13,2. **M. jurensis* (DUNKER), U.Jur.(Kimmeridg.), France; 2a-c, LV ext., both valves dorsal; LV ext. showing impression of internal rib; all $\times 1$ (de Loriol, 1868).
- ?**Molukkana** KRUMBECK, 1923, p. 219 [**M. seranensis*; M]. Oblong, strongly inequilateral, with broad, moderately protruding, almost orthogyrous, well anteriorly placed umbones and rounded-off diagonal umbonal ridge; ostracum thick, ligamental nymph solid; hinge edentulous; pallial line and adductor scars unknown; ornament of irregular concentric undulations. [Possibly inseparable from *Pachymya* (*Pachymya*) but imperfectly known.] *U.Trias.(Nor.)*, Indon.
- Neoburmesia** YABE & SATO, 1942, p. 251 [**N. iwakiensis*; OD]. Medium-sized, elongate-oblong, with prominent, anteriorly placed, broadly rounded umbones well incurved to orthogyrous beaks; strongly inflated, most so at mid-length; posterior gape moderately wide; strong carina in each valve from umbo to posteroventral angle, delimiting broad, elongate, partly concave posterior area, while on other side of umbo weaker radial ridge delimits small anterior area; escutcheon long, narrow, bordered by distinct ridges; median part of surface bearing radial and concentric ribs with tubercles at their intersections; posterior and anterior areas with weak concentric ornament only; internal characters unknown. *U.Jur.*, Japan.—FIG. F13,1. **N. iwakiensis*; 1a,b, LV ext. and dorsal view of both valves, $\times 1$ (Tamura, 1960).
- Osteomya** MOESCH, 1874, p. 19 [**Mya dilata* PHILLIPS, 1829, p. 155 (cited as "*dilatata*"); M] [= *Goniomeris* CHOFFAT, 1893, p. 37 (type, *G. gaudryi*; M); *Uromya* ROLLIER, 1913, p. 262 (type, *Mya dilata* PHILLIPS; SD COSSMANN, 1921, p. 27); *Eurychasma* COSSMANN, 1915, p. 9 (type, *E. combesi*; M)]. Medium-sized, elongate-oblong, strongly

inequilateral, with broad, only slightly protruding, opisthogyrous umbones placed from anterior third to quarter of length; some specimens upcurved posteriorly; inflation moderately strong; shell trun-

cated vertically posteriorly and with wide gape corresponding to complete cross section of shell; anterior gape absent or very narrow; ornament consisting of weak concentric undulations, irregu-




FIG. F13. Pholadomyidae (p. N832).

larly arranged except on anterodorsal part of surface; pustules relatively conspicuous, in closely spaced radial rows; internal characters unknown. *L.Jur.(Toarc.)-U.Jur.(Callov.)*, Eu.-Madag.-E.Afr. —FIG. F14,1. **O. dilata* (PHILLIPS), M.Jur. (Bajoc.), Eng.; 1*a,b*, LV ext., dorsal view of both valves, $\times 1$ (Cox, n).

Pachymya J. DE C. SOWERBY, 1826, p. 1 [**P. gigas*; M]. Oblong, moderately to strongly inequilateral, with broad, only slightly protruding, almost orthogyrus umbones; posterior gape slight to moderate; most species with umbonal ridge, crossing most inflated part of shell, running diagonally to posteroventral corner; pallial line distant from shell margins; pallial sinus (where known) small or absent; surface unornamented in most species except for fine, radially aligned pustules, but with concentric undulations in some Triassic forms. *M.Trias.-U.Cret.*, cosmop.

P. (Pachymya) [= *Cratomya* ROLLIER, 1913, p. 287 (type, *Arcomya liesbergensis* ROLLIER, 1913, p. 286; M)]. Medium-sized to large, with umbones placed well anteriorly; moderately to strongly inflated, with diagonal ridge prominent; escutcheon broad, scarcely impressed, bordered by obtuse umbonal ridges; ostracum relatively thick in many species; hinge margin much thickened near beak and commonly with irregular longitudinal rugosities; ligamental nymphs stout; pallial sinus small or absent. *M.Trias.-U.Cret. (Cenoman.)*, cosmop.—FIG. F14,2*a,b*. **P. (P.) gigas* SOWERBY, *U.Cret.(Cenoman.)*, France; 2*a,b*, LV ext., and dorsal view of both valves, $\times 0.5$ (d'Orbigny, 1843-47).—FIG. F14,2*c*. *P. (P.) austiniensis* SHUMARD, *L.Cret.(Alb.)*, USA (Tex.); RV int. showing hinge and pallial line, $\times 0.45$ (Stanton, 1947).—FIG. F14,2*d*. *P. (P.) latissima* (AGASSIZ), *U.Jur.(Oxford)*, Switz.; LV int., showing pallial line, $\times 0.7$ (Rollier, 1913).

P. (Arcomya) ROEMER (ex AGASSIZ, MS), 1839, p. 43 [**Solen helveticus* ROEMER (ex THURMANN, MS), 1839; M]. Medium-sized, relatively weakly inflated; position of umbones variable; hinge structure not observed; pallial line (where known, but not seen in type species) without sinus. *M. Trias.-U.Cret.(Turon.)*, cosmop. [The Triassic species have been included mostly in *Homomya*.] —FIG. F15,2*a,b*. **P. (A.) helvetica* (ROEMER), *U.Jur.(Kimmeridg.)*, Switz.; 2*a,b*, RV ext. and dorsal views of both valves, $\times 1$ (Agassiz, 1843). —FIG. F15,2*c*. *P. (A.)* sp., *U.Jur.*, Scot.; RV int. mold showing pallial line, $\times 1$ (Cox, n).

P. (Trichomyella) COX, *nom. subst.* herein [pro *Trichomya* CRICKMAY, 1936, p. 558 (non IHERING, 1900, p. 87)] [**Trichomya amphitrite* CRICKMAY, 1936; OD]. Resembling *P. (Arcomya)* but without diagonal ridge; radial lines of pustules conspicuous. *U.Jur.(Callov.)*, USA (Wyo.).—FIG. F15,1. **P. (T.) amphitrite* (CRICKMAY); RV ext., $\times 1.3$ (Crickmay, 1936).

Palaeocosmomya FLETCHER, 1946, p. 401 [**P. teichertii*; OD]. Medium-sized, ovate or oblong, variably elongated; with broadly rounded, unprotruding or slightly protruding umbones situated at or in front of the anterior third of length, and with feebly prosogyrous beaks; inflation moderate, posterior gape wide; escutcheon distinctly bordered; posterior 0.7 of surface bearing 2 convergent series of steep, oblique ribs which meet to form V's, anterior third with broader ribs which are only slightly oblique to growth lines; radial groove crossing latter series of ribs (that is, not marking boundary between them and V-shaped ribs) runs almost perpendicularly from umbo to ventral margin; internal characters unknown. [Regarded by DICKINS & SHAH (1965) as subgenus of *Cosmomya*.] *Perm.*, Australia-Greenland.—FIG. F15,7. **P. teichertii*; LV ext., $\times 1$ (Cox, n).

Panacca DALL, 1905, p. 143 [pro *Aporema* DALL, 1903, p. 1532 (non SCUDDER, 1890)] [**Pholadomya arata* VERRILL, 1881, p. 301; OD] [= *Notomya* COTTON, 1931, p. 342 (non M'COY, 1847) (type, *Pholadomya tasmanica* HEDLEY & MAY, 1914, p. 132; OD); *Panacea* SALISBURY, 1932 (*nom. null.*)]. Shell small for family, gibbous, subtriangular, inequilateral, anterior end more or less truncated and flattened transversely; anterior margin commonly sinuate and meeting anterodorsal margin in well-defined angle; beaks slightly prosogyrous; posterior gape narrow or absent; most species ornamented with unevenly spaced, narrow radial ridges absent from anterior end of shell (one species with radial threads only); surface pustules very fine; pallial line with shallow sinus; hinge margin simple, very little thickened; ligament entirely external. *Rec.*, cosmop. [Deep water. About seven known species are typical; *Pholadomya loveni* JEFFREYS and *P. pacifica* DALL, which some authors have included in *Panacca*, disagree with the above diagnosis in being less trigonal and inequilateral.] —FIG. F15,3*a,b*. **P. arata* (VERRILL), *W.Atl.*; side and ant. views RV, $\times 1$ (Verrill, 1882, 1884).—FIG. F15,3*c,d*. *P. locardi* DALL, *E.Atl.*; LV ext. and RV int., $\times 1.3$ (Locard, 1898).

Parilimya MELVILL & STANDEN, 1899, p. 202 [**Pholadomya (Parilimya) haddoni*; M]. Small, oval, subequilateral, evenly and only moderately inflated; margins with narrow anterior but no posterior gape; ornament of weak radial threads; pallial line not observed; ligament entirely external; dorsal margin with toothlike thickening in front of beak in both valves. *Rec.*, Torres Straits.

Pentagrammysia CHERNYSHEV, 1950, p. 22 [**P. altaica*, OD]. Superficially similar to *Undulomya* FLETCHER, 1946, in form and ornamentation but without posterodorsal furrows; posterior gape obscure, if present. *L.Carb.*, Kuznetsk Basin, USSR (Sib.).—FIG. F16,4. **P. altaica*; RV ext., $\times 1$ (Ebersin, 1960). [NEWELL]

Praeundulomya DICKINS, 1957, p. 10 [**P. concentrica*; OD]. Medium-sized, oblong, elongate, with broad, scarcely protruding, well anteriorly placed umbones and prosogyrous beaks; inflation

moderate, posterior gape narrow; escutcheon narrow, elongate, distinctly bordered; flank with 2 dorsal grooves like those of *Undulomya* but shallower; ornament consisting of irregular undula-




FIG. F14. Pholadomyidae (p. N833-N834).




FIG. F15. Pholadomyidae (p. N834-N835, N837-N838).




FIG. F16. Pholadomyidae (p. N834, N837-N838).

tions, which, although almost concentric, are slightly oblique to growth lines on most parts of surface; V-shaped ribs absent; internal characters unknown. *L.Perm.*, Australia-?Kashmir-S. Am.(Peru).—FIG. F15,6. **P. concentrica*, W. Australia; 6*a,b*, LV ext. (imperfect anteriorly) and dorsal view of both valves, $\times 1$ (Dickins, 1957).

Tetorimya HAYAMI, 1959, p. 159 [**T. carinata*; OD]. Medium-sized, trapezoidal, highest near posterior end, well inflated, most so along diagonal (which may form distinct ridge) from umbo to

posteroventral angle of shell; posterior gape narrow or absent; umbo narrow, protruding, placed well anteriorly and incurved to prosogyrous beak; anterior part of ventral margin with broad sinus to which radial sulcus of flank corresponds; pallial line with deep sinus; surface with coarse growth rugae; internal features of hinge unknown. *U.Jur.*, Japan-Alaska-USSR.—FIG. F15,5*a*. **T. carinata*, Japan; RV ext., $\times 1$ (392).—FIG. F15,5*b*. *T. panderi* (EICHWALD), Alaska; LV ext., $\times 1$ (Eichwald, 1871).

?*Triplicosta* COOPER, 1897, p. 333 [**Pholadomya*

(*Triplacosta*) *progressiva*; M]. Medium-sized, ovate, not much elongated, gibbose, strongly inequilateral, with broadly rounded, well anteriorly placed umbones; with moderate posterior gape; ornament of rounded radial ribs which are relatively broad on the posterior part of the surface and margins and clearly arranged on the anterior part; internal characters unknown; ostracum relatively thick, not nacreous in known fossil specimens. [While included in *Pholadomya* it is also suggested that this species may be related to *Cardita* or *Petricola*.] ?*Eoc.*, USA (Calif.).—FIG. F16,2. **T. progressiva* (COOPER); 2*a,b*, RV ext., both valves ant., ×1 (Cooper, 1897).

Undulomya FLETCHER, 1946, p. 391 [**U. pleio-pleura*; OD]. Medium-sized, elongate-ovate, with broad, unprotruding, well anteriorly placed umbones and prosogyrous beaks; inflation moderate, posterior gape moderately wide; escutcheon shallow; dorsal part of flank with 2 grooves running from umbo to posterior margin; ornament consisting of 2 convergent series of straight or curved ribs which meet along line from umbo to ventral margin; ribs of posterior series strongly oblique to growth lines, while obliquity of anterior series is variable and they are almost concentric in some specimens; internal characters unknown. *Perm.*, Australia-Afr. (Madag.).—FIG. F15,4. **U. pleio-pleura*, W.Australia; LV ext., ×0.7 (Dickins, 1956).

Vacunella WATERHOUSE, 1956, p. 277 [**Allorisma curvatum* MORRIS, 1845; OD]. Similar to *Chaenomya* but with more rounded ventral margin, maximum shell convexity near mid-length and relatively small posterodorsal gape; shell interior marked by scattered mantle insertions; edentulous; ligament nymphs heavy and rolled outward. *Perm.*, E.Australia.—FIG. F16,3. **V. curvatum* (MORRIS), L.Perm., Wollongong, New S. Wales; 3*a*, top view of LV int. mold; 3*b*, same, int. hinge, latex cast showing interior of amphidetic ligament; 3*c*, same showing pallial sinus and mantle insertions, all ×0.4 (Newell, n). [NEWELL]

Wilkingia WILSON, 1959, p. 401 [**Venus elliptica* PHILLIPS, 1836; OD] [= *Allorisma* AUCT. (non KING, 1850); ?*Tellinomorpha* DE KONINCK, 1885, p. 90 (type, *T. cuneiformis*; OD)]. Elongate oval, subcylindrical at beaks; beaks placed 0.2 to 0.25 behind front margin; anterior and posterior extremities rounded, dorsal and ventral margins subparallel; broad, shallow sulcus extending ventrally from umbones across anterior half of valves; escutcheon and elongate lunule present; umbonal ridge broad and poorly developed; valves marked by regular broad and rounded concentric folds; radial rows of small papilli present, especially on posterodorsal area; adductor scars shallow, without buttresses; shallow pallial sinus present; with or without narrow posterior gape; ligament nymphs short and weak; hinge edentulous. *Miss.-Perm.*,

cosmop.—FIG. F16,1. **W. elliptica* (PHILLIPS), L.Carb., Eng. (Redesdale); 1*a,b*, RV and cardinal views, ×1 (Wilson, 1959). [NEWELL]

Family BURMESIIDAE Healey, 1908

[Materials for this family prepared by L. R. Cox]

Medium-sized, equivalve, thin-shelled, of moderate convexity, well elongated, not strongly inequilateral; hinge line elongate, umbones broad and protruding only slightly above it; each valve with narrow, spoonlike chondrophore extending into shell cavity from below beak; no hinge teeth; pallial line and adductor scars not observed; surface elaborately ornamented with radial riblets together with concentric folds and threads, radial riblets predominating on middle of each valve; whole surface also covered with rows of minute granules. *U. Trias.-L.Jur. (Lias.)*.

Burmesia HEALEY, 1908, p. 58 [**B. latouchii*; SD DIENER, 1923, p. 247]. Elongate-ovate, beaks very slightly anterior to mid-length; valves gaping slightly in front, more widely behind; posterior end of shell ornamented with concentric folds and subordinate radial threads, anterior end commonly bearing depressed folds which are almost perpendicular to hinge line and cross growth lines obliquely. *U. Trias.-L. Jur. (Lias.)*, Asia (Jordan-Burma-Indonesia-Japan).—FIG. F17,1. *B. lirata* HEALEY, U.Trias. (Rhaet.), Burma; pair of valves, RV below, ×1 (Healey, 1908).

Prolaria HEALEY, 1908, p. 60 [**P. sollasi*; M]. Presumed posterior end of shell forming rostrum-like projection truncated at its extremity and situated in relatively dorsal position with deep sinus below it; sharp sigmoidally curved carina runs in each valve from umbo to lower corner of this projection; predominant radial ribbing confined to posteromedian part of surface; strong, irregular concentric folds with subordinate radial threads cover anterior half of surface, while posterior projection is unornamented except for growth lines. *U. Trias. (Rhaet.)*, Asia (Burma-Armenia-Indon.).—FIG. F17,2*a,b*. **P. sollasi*, Burma; 2*a*, part of RV ext., showing posterior projection, 2*b*, more complete but crushed RV, both ×1 (Healey, 1908).—FIG. F17,2*c,d*. *P. armenica* ROBINSON, Armenia; 2*c,d*, RV ext. and dorsal view of both valves, post. extremity broken away, ×1 (Kiparivsova, 1947).

Family CERATOMYIDAE Arkel, 1934

[= *Ceromyidae* FISCHER, 1887 (invalid because based on junior homonym)] [Materials for this family prepared by L. R. Cox]

Shell ovate, longer than high, inequilateral, moderately to strongly inflated, some

Pholadomyoidea—Pholadomyacea




FIG. F17. Burmesiidae (p. N838).

specimens inequivalve; valve margins closed or with narrow posterior gape; shell wall rather thin; beaks prosogyrous; no demarcated lunule or escutcheon; ligament opisthodontic, subinternal, located between reflected and thickened posterodorsal margin of LV and overlapping margin of RV, which has subinternal thickening protruding into cavity of valve, giving rise in *Cera-*

tomya and *Gresslya* to slit extending back from beak on internal mold of many specimens; true hinge teeth absent, replaced by thickenings or protuberances of dorsal margins; pallial line variable; surface with variously oriented plications or unornamented except for minute pustules present in some forms. *U.Trias.-U.Jur., ?Mio.*

The hinge apparatus is so closely similar in *Ceratomya* and *Gresslya* that their union in the same family seems justified, especially as the earlier species of *Ceratomya* (e.g., *C. petricosa* of the Lias) lack the external ribbing which renders some of the later species so much unlike *Gresslya* in appearance. Nevertheless, the two genera differ greatly in the form of the pallial line.

It seems most improbable that the Miocene genus *Ceromyella* was (as thought by its author, SACCO) a survivor of the Ceratomyidae, but, for lack of definite evidence as to its affinities, it is here listed under this family, but with a query.

Ceratomya SANDBERGER, 1864, p. 16 [*nom. subst. pro Ceromya* AGASSIZ, 1842] [*non Ceromya* ROBINEAU-DESVOIDY, 1830] [**Isocardia excentrica* ROEMER, 1836, p. 106; ICZN pend.] [= *Ceromia* STOLICZKA, 1871 (*nom. null.*)]. Medium-sized to large, ovate, gibbose, especially anteriorly, with strongly prosogyrous and in some species incoiled, more or less anteriorly placed beaks; some specimens slightly inequivalve, with umbo of either valve extending more anteriorly than that of other; no gape of valve margins; posterodorsal margin of RV with thin outer flange separated from subinternal thickening by groove in which ligament was inserted, corresponding groove on upper (outer) side of reflected margin serving same purpose in LV; anterodorsal margin of RV much thickened below beak in some species; adductor scars small; pallial line bending up sharply posteriorly, shallowly sinuate in some species; surface smooth or (more commonly) with concentric or oblique undulations. *L.Jur.-U.Jur.*, cosmop.—FIG. F18,1a. **C. excentrica* (ROEMER), *U.Jur.* (*L. Kimmeridg.*), France; LV ext., $\times 1$ (549).—FIG. F18,1b-d. *C. aalensis* (QUENSTEDT), *M.Jur.* (*Bajoc.*), France; 1b-d, LV int., RV int., and int. mold showing pallial line and adductor scars, all $\times 0.7$ (Benecke, 1905).—FIG. F18,1e. *C. petricosa* (SIMPSON), *L.Jur.* (*M.Lias.*), Eng.; LV ext. of a smooth species, $\times 1.3$ (Cox, n).—FIG. F18,1f. *C. bajociana* (D'ORBIGNY), *M.Jur.* (*Bajoc.*), Eng.; transv. sec. through hinge region short distance behind beaks, LV and RV as oriented, former position of ligament dotted, $\times 2$ (Cox, n).

?*Ceromyella* SACCO, 1901, p. 133 [**C. miotaurina*; M]. Like *Ceratomya* in shape but only 6 mm. long; internal characters unknown. *Mio.*(*Helvet.*), N.Italy.

Gresslya AGASSIZ, 1843, p. 202 [**Lutraria gregaria* ZIETEN, 1833, p. 85; SD HERRMANNSEN, 1847, p. 490, as "*Lutraria gregaria* GOLDFUSS"] [=Gres-

slyia BRONN, 1848 (*nom. van.*); *Gresslya* D'ORBIGNY, 1850 (*nom. null.*); *Gressluya* PAETEL, 1875 (*nom. null.*)]. Ovate, usually with cuneiform tendency, moderately elongate, umbones not prominent, that of RV slightly higher than LV; valves with narrow posterior gape; hinge structure as defined for family, slit extending back from





FIG. F18. Ceratomyidae (p. N839-N841).

beak on internal molds of RV even better defined and more constant than in *Ceratomya*; both valves in some specimens with a weak tooth-like callosity of the antero-dorsal margin near the beak; pallial line with a deep sinus; surface without con-

spicuous ornament but bearing very fine pustules or puncta radially aligned. *L. Jur.-U. Jur.*, cosmop. —FIG. F18,2a,b. *G. peregrina* (PHILLIPS), U. Jur. (Oxford), Eng.; 2a,b, LV int. mold showing pallial sinus, dorsal view of both valves showing slit extending back from beak in RV, $\times 1.3$ (Cox, n). —FIG. F18,2c. *G. gregaria* (ZIETEN), M. Jur. (Bajoc), Ger.; LV ext., $\times 1$ (Schmidtill, 1926). —FIG. F18,2d. *G. abducta* (PHILLIPS), M. Jur. (Bajoc.), Eng.; transv. sec. through hinge region short distance behind beaks, LV and RV as oriented, edge marked "X" giving rise to the slit on internal molds; $\times 3.3$ (Cox, n).



?*Pholadomyocardia* SZAJNOCHA, 1889, p. 88 [**P. jelskii*; M]. Large, inequivalve, gibbose, slightly longer than high; valves with a broadly trigonal body and a flattened winglike postero-dorsal extension; umbones placed well anteriorly, protruding moderately, prosogyrous, the right one higher than the left; surface with a few broad, irregular, concentric folds; internal characters unknown. *Jur.*, S. Am. (Peru). —FIG. F19,2. **P. jelskii*, 2a,b, LV ext. and ant. view of both valves, $\times 0.5$ (Szajnocha, 1889).

Pteromya MOORE, 1861, p. 505 [**P. crowcombeia*; SD STOLICZKA, 1871, p. xv]. Subovate, subequilateral or inequilateral to a varying extent, not strongly inflated, slightly to moderately inequivalve, right valve the more gibbose and with its umbo the more elevated; valve margins not gaping; umbones broad, protruding very little; posterodorsal and posterior margins forming continuous, strongly convex curve or else meeting in obtuse angle, in which case very obtuse posterior ridge may be present in one or both valves; hinge structure as in *Ceratomya* and *Gresslya*; adductor scars and pallial line not yet observed, the latter probably without a sinus; surface with concentric undulations or ridges, or merely with coarse growth threads. *U. Trias. (L. Rhaet.)-L. Jur. (L. Hettang.)*, Eu. —FIG. F19,1a,b. **P. crowcombeia*, L. Rhaet., Eng.; 1a,b, LV ext., and RV ext., of different specimens, $\times 1$ (Cox, n). —FIG. F19,1c,d. *P. tatei* (RICHARDSON & TUTCHER), L. Hettang., Eng.; 1c,d, RV ext. with and without posterior ridge, all $\times 1$ (Cox, n).

Family MYOPHOLADIDAE Cox, 1964

[Materials for this family prepared by L. R. Cox]

Shell medium-sized, elongate-ovate, some specimens with slight tendency to be rostrate posteriorly; umbones broadly rounded, scarcely protruding, situated anterior to mid-length and strongly incurved to slightly prosogyrous beaks; inflation of shell strong below umbones, diminishing regularly in posterior direction; valve margins with

FIG. F19. Ceratomyidae (p. N841).




FIG. F20. Myopholadidae (p. N842).

rather wide posterior and narrow anterior gape; ornament of narrow radial ribs which are widely separated on anterior part of surface and relatively closely arranged on median part; in some species strong and fairly closely arranged ribs continue almost to posterior extremity, but in others posterior end of shell is sparsely ribbed or almost smooth; dorsal margin of RV overlapping that of LV, with subinternal ligament lying between them, as in *Ceratomya* and *Gresslya*; sharp edge of thickening of dorsal margin of right valve which projects into cavity of valve gives rise to groove ex-

tending back from beak on internal mold, as in those two genera; adductor scars and pallial line unknown; ostracum thin, probably originally nacreous internally. *M. Jur.* (*Bathon.*)-*L. Cret.* (*U. Alb.*).

Myopholas DOUVILLÉ, 1907, p. 107 [*Pholadomya multicostata* AGASSIZ, 1842, p. 52; OD]. Characters and distribution of family. *M. Jur.*-*L. Cret.*, Eu.—FIG. F20, *1a, b*. **M. multicostata* (AGASSIZ), *U. Jur.* (*Kimmeridg.*), Switz.; *1a, b*, RV ext. and dorsal view of both valves, $\times 1$ (Cox, n).—FIG. F20, *1c*. *M. ledouxi* DOUVILLÉ, *L. Cret.* (*U. Alb.*), France; dorsal view of both valves showing groove on RV of internal mold, $\times 1$ (Douvillé, 1907).

Family PLEUROMYIDAE Dall, 1900

[*nom. correct.* ZITTEL, 1903 (*pro* Pleuromyidae DALL, 1900, *ex* ZITTEL MS)] [Materials for this family prepared by L. R. Cox]

Shell of medium size, equivalve, oval, oblong or trapeziform, moderately to strongly inflated, with narrow to moderately wide posterior gape; umbones situated within anterior half and usually well toward anterior end of shell, broadly rounded, not strongly protruding; no demarcated dorsal areas; no true hinge teeth; main part of ligament external, opisthodic, with short supporting nymphs; below and slightly anterior to the beak of each valve is short, rounded protuberance of margin, slightly hollowed out above, and behind protuberance is small niche; protuberance of RV fits above that of the left, a small internal ligament possibly occupying the space between them; postero-dorsal margin of RV overlapping that of LV; pallial sinus deep; ostracum thin, probably originally nacreous internally. Surface with or without concentric ribbing and with a delicate ornament of radial rows of minute pustules. *Trias.*-*L. Cret.*

Pleuromya AGASSIZ, 1842, p. 439 [**Mya gibbosa* J. DE C. SOWERBY, 1823, p. 19 (= **Donacites alduini* BRONGNIART, 1821, p. 555); M] [= *Myacites* VON SCHLOTHEIM, AUCTT.¹; *Anoplomya* KRAUS, 1843, p. 130 (type, *A. lutraria*; M); *Anoplomya* KRAUS, 1850 (*nom. null.*); *Hapalomya* ROEDER, 1882, p. 102 (type, *H. fragilis*; M); *Hapalomya* ROEDER, 1882 (*nom. null.*); ?*Fogiella* KRUMBECK,

¹ This generic name, applied particularly to shells of this genus by some early authors, merely denoted a fossil *Mya* and is not available for purposes of nomenclature (Zool. Code, 1961, Art. 20).




FIG. F21. Pleuromyidae (p. N842-N843).

1913, p. 57 (type, *F. deningeri*; M)]. Characters of family. *Trias.-L.Cret.*, cosmop.—FIG. F21, 1a. **P. alduini* (BRONGNIART), U.Jur. (U. Oxford Clay), Eng.; RV ext., $\times 1$ (19f).—FIG. F21, 1b. *P. uniformis* (J. SOWERBY), U.Jur. (Oxford), Eng.; RV composite ext.-int., showing pallial line, etc. (not retouched), $\times 1$ (Cox, n).—FIG. F21, 1c. *P. uniformis*, U.Jur. (Corallian Beds), Eng.; LV hinge tilted slightly toward observer, showing protuberance just anterior to beak, $\times 2$ (Cox, n).—FIG. F21, 1d. *P. marginata* AGASSIZ, M.Jur. (Bathon.), France; diagram showing relation between protuberances of hinge margins of valves, LV on left, enl. (Douvillé, 1907).—FIG. F21, 1e, f. *P. uniformis*, M.Jur. (Inf. Oolite), Eng.; 1e, f, transverse sections through dorsal region of slightly separated valves, on left, 1e just behind, 1f just in front of beaks, $\times 2$ (Cox, n).

Superfamily PANDORACEA Rafinesque, 1815

[*nom. transl.* STEWART, 1930 (ex Pandoridae RAFINESQUE, 1815)] [Materials for this superfamily prepared by MYRA KEEN with additions as recorded]

Sedentary to burrowing forms, shell material nacreous, at least inner layer; mostly thin, somewhat elongate or gaping, inequivalve; hinge without regular heterodont

dentition but hinge margin variously reinforced by buttresses or denticles; ligament and resilium sheathed with a calcareous layer or lithodesma in many forms. *U. Trias.-Rec.*

Family PANDORIDAE Rafinesque, 1815

[*nom. correct.* GRAY, 1840 (pro family Pandoracia RAFINESQUE, 1815)] [Materials for this family prepared by MYRA KEEN]

Compressed, inequivalve, dorsal border of one valve overlapping other; ligament wanting, resilium internal, reinforced in some by elongate lithodesma; dorsal margin edentulous but with laminar buttresses beside the resilium; pallial line entire. *Oligo.-Rec.*

Pandora BRUGIÈRE, 1797 (genus without species) [**Solen inequivalvis* LINNÉ, 1758; SM LAMARCK, 1799] [= *Calopodium* RÖDING, 1798 (obj.); *Truina* BROWN, 1827 (type, *T. solenoidea*, = *Solen pinna* MONTAGU, 1803; M)]. Of medium size, flat, thin, RV flatter than LV, dorsal border convex; anterior end rounded, posterior rostrate. *Oligo.-Rec.*, N. Am.-S. Am.-C. Am.-Eu.-Pac.-Ind. O. P. (*Pandora*). Sculpture of RV feebly concentric; without lithodesma. *Oligo.-Rec.*, W. N. Am.-Eu.-




FIG. F22. Pandoridae (1,3-7); Cleidothaeridae (2)
(p. N843-N844).

W.Pac.—FIG. F22,3. **P. (P.) inaequivalvis* (LINNÉ), Rec., Eu.; 3a-c, LV int., ext., RV hinge, $\times 1$ (7c).

P. (Clidiophora) CARPENTER, 1864 [**P. claviculata* CARPENTER, 1856; OD]. With 3 laminae in either valve; lithodesma present. *Mio.-Rec.*, E.N. Am.-W.N.Am.-C.Am.—FIG. F22,5. **P. (C.) claviculata* CARPENTER, Rec., W.Mex.; 5a,b, RV ext., LV int., $\times 0.7$ (Stanford Univ. specimen).

P. (Foveadens) DALL, 1915 [**F. panamensis*; OD]. Hinge with 2 diverging laminae in RV, posterior longer and higher; LV with 2 short laminae close together; sheet of shelly material bridging bases of laminae, forming cavities between them. *Rec.*, W.C.Am.—FIG. F22,4. **P. (F.) panamensis* (DALL); 4a,b, holotype, RV ext., LV int., $\times 2$ (U.S. National Museum).

P. (Frenamya) IREDALE, 1930 [**Coelodon patulus* TATE, 1889; OD] [= *Coelodon* CARPENTER, 1865 (non AUDINET-SERVILLE, 1882) (type, *P. ceylanica* SOWERBY, 1835; SD STOLICZKA, 1871)]. Sculpture of concentric wrinkles; LV with laminae united by transverse plate; no lithodesma. *Rec.*, W.C.Am.-S.Pac.-W.Pac.-Ind.O.—FIG. F22,6. **P. (F.) patula* (TATE), S.Australia, 6a,b, RV int, LV hinge, $\times 2$ (Cotton, 1938).

P. (Heteroclidus) DALL, 1903 [**P. punctata* CONRAD, 1837; OD]. With 1 laminar buttress in LV, 3 in RV; lithodesma present. *Plio.-Rec.*, W.N.Am.-C.Am.—FIG. F22,7. **P. (H.) punctata* CONRAD, Rec., USA (Calif.); 7a,b, LV ext., int., $\times 1$ (Stanford Univ. specimen).

P. (Pandorella) CONRAD, 1863 [**P. arenosa* CONRAD, 1834; M] [= *Kennerlia* CARPENTER, 1864 (type, *K. bicarinata*; SD STOLICZKA, 1870) (*Kennerleya*, *Kennerleyia*, *Kennerlyia*, spelling errors)]. With radial ribbing in RV; lithodesma wanting. *Mio.-Rec.*, E.N.Am.-W.N.Am.-S.Am.—FIG. F22,1. **P. (P.) arenosa* CONRAD, Mio., USA (Va.); 1a,b, LV ext., int., $\times 1$ (Conrad, 1838).

Family CLEIDOTHAERIDAE Hedley, 1918

[=Chamostridae FISCHER, 1887] [Materials for this family prepared by MYRA KEEN]

Shell attached by right valve; one cardinal tooth in LV; ligament wanting but resilium present, submerged, with lithodesma. *Mio.-Rec.*

Cleidothaerus STUTCHBURY, 1830 [**C. chamoides* (= *Chama albida* LAMARCK, 1819; M] [= *Chamostraea* HERRMANNSEN, 1846 (latinization of "Chamostrée" DE BLAINVILLE, 1825, ex ROISSY MS); OD (obj.); "*Chamostraea* ROISSY, 1805, AUCTT." (fictitious reference)]. Inequivalve, fixed on anterior slope; hinge of RV with large pit; lithodesma long, curved. *Mio.-Rec.*, Australasia.—FIG. F22,2. **C. albidus* (LAMARCK), Rec., S. Australia; 2a,b, LV int., RV int., $\times 0.5$ (307).

Family LATERNULIDAE Hedley, 1918

[=Anatinidae GRAY, 1840 (invalid, based on junior homonym); Cercomyacidae CRICKMAY, 1936] [Materials for this family prepared by MYRA KEEN and L.R. COX]

Subequivalve, elongate, gaping posteriorly; umbones not at all or only slightly pro-

truding, with transverse external slit, or with internal plate represented by slit on internal molds; hinge edentulous; ligament (where known) on two projecting spoon-shaped chondrophores, each supported by thin oblique buttress (or clavicle); pallial sinus broad; shell thin, subnacreous internally (526). *U.Trias.-Rec.*

Laternula RÖDING, 1798, p. 155 [**L. anatina* (= *Solen anatinus* LINNÉ, 1758, p. 673); SD GRAY, 1847, p. 190] [= *Auriscalpium* MEGERLE VON MÜHLFELD, 1811 (obj.; M); *Anatina* LAMARCK, 1818 (non SCHUMACHER, 1817) (obj.; T); *Butor* GISTEL, 1848 (non FORSTER, 1827) (pro *Anatina* LAMARCK); *Butorella* STRAND, 1928 (pro *Butor* GISTEL)]. Thin subcylindrical, posterior end roundly gaping. *U.Cret.-Rec.*, N.Am.-Eu.-Pac.-Ind.O.

L. (Laternula). Surface nearly smooth; posterior gape wide. *U.Cret.-Rec.*, N.Am.-Eu.-S.Pac.-Ind.O.—FIG. F23,1. **L. (L.) anatina* RÖDING, *Rec.*, E.Indies; *1a,b*, LV int., with RV hinge, LV ext., $\times 0.7$ (307).

L. (Laternulina) HABE, 1952, p. 266 [**Anatina japonica* LISCHKE, 1872, p. 107 (= *A. flexuosa* REEVE, 1863, sp. 5); OD]. Surface with wrinkled undulations; gape narrower than in *L. (Laternula)*. *Rec.*, W.Pac.—FIG. F23,2. **L. (L.) flexuosa* (REEVE), Japan; LV view of shell, $\times 1$ (Habe, 1952).

Anatimya CONRAD, 1860, p. 276 [**Pholadomya (A.) anteradiata*; SD SHIMER & SHROCK, 1944, p. 414]. Oblong, subequilateral, with concentric furrows anteriorly and radiating lines posteriorly; hinge unknown. *U.Cret.*, E.N.Am.—FIG. F23,3. **A. anteradiata* (CONRAD), USA(N.J.); LV view, $\times 0.7$ (Whitfield, 1907).

Cercomya AGASSIZ, 1843, p. 143 [**C. pinguis*; OD] [= ?*Kercomya* GRESSLY, 1838 (*nom. nud.*)]. Elongate, subequilateral to distinctly inequilateral, compressed, with tapering posterior end, upcurved in some species; umbones level with posterodorsal margin or almost so; posterior umbonal ridges well defined; umbonal slit or impression of internal umbonal plate observed in some species; hinge and pallial line unknown; surface bearing minute, radially aligned pustules. *U.Trias.-Cret.*, cosmop.

C. (Cercomya). Flanks with concentric folds, posterodorsal area smooth. *U.Trias.-Cret.*, cosmop.—FIG. F23,7a,b. **C. (C.) pinguis* AGASSIZ, M.Jur.(Bajoc.), Switz.; *7a,b*, LV ext., and dorsal views, $\times 1$ (Agassiz, 1843).—FIG. F23,7c,d. *C. (C.)* sp. aff. *C. (C.) gurgilis* (PICTET & CAMPICHE), L. Cret.(Alb.), Eng.; *7c,d*, RV ext., and dorsal view, $\times 0.7$ (Woods, 1909).

C. (Capillimya) CRICKMAY, 1936, p. 558 [**Capillimya capillifera*; OD]. With punctate radial striae on posteroventral sector and concentric

folds on rest of surface. *Jur.*, Eu.-N.Am.—FIG. F23,6. *C. (C.) striata* AGASSIZ, *U.Jur.*(L. Kimmeridg.), France; *6a,b*, RV ext. and dorsal views, $\times 1$ (de Loriol, 1872).

Clistoconcha SMITH, 1910, p. 217 [**C. insignis*; M]. Small, irregularly oblique-ovate; chondrophores small, buttress large; valves united in adult except for siphonal gape. *Rec.*, S.Afr.—FIG. F23,5. **C. insignis*; *5a,b*, LV ext., RV int., $\times 4$ (Kuehnelt, 1958).

Periplomya CONRAD, 1870 (July 7), p. 76 [*nom. subst. pro Leptomya* CONRAD, 1867, p. 15 (non ADAMS, 1864)] [**Periploma applicata* CONRAD, 1858, p. 324; SD GARDNER, 1916, p. 633] [= *Pliomya* STOLICZKA, 1870 (Sept. 1) (pro *Leptomya* CONRAD)]. Like *Laternula*, but with chondrophores tapering and attached to buttress; slit anterior, oblique, long; surface with concentric ribbing. *U.Cret.*, E.N.Am.—FIG. F23,4. **P. applicata* (CONRAD), USA(Tenn.); RV ext., $\times 1$ (951).

Platymyoidea COX, 1964, p. 42 [*nom. subst. pro Platymya* AGASSIZ, 1843, p. 180 (non ROBINEAU-DESVOIDY, 1830)] [**Platymya dilatata* AGASSIZ; OD]. Ovate-oblong, not tapering posteriorly, compressed, with broad, scarcely protruding umbones between middle and anterior third of length; internal umbonal plate well developed; hinge unknown; ornament of irregular concentric folds strongest at anterior end of shell. *L.Jur.-L.Cret.*, Eu.(Switz.-France-Eng.).—FIG. F23,8. *P. rostrata* (AGASSIZ), L.Cret.(Neocom.), France; *8a,b*, LV ext. and dorsal views, $\times 0.7$ (696).

Plectomya DE LORIO, 1868, p. 525 [**Tellina rugosa* RÖMER, 1836, p. 120 (non PENNANT, 1777) (= *Anatina subrugosa* D'ORBIGNY, 1850, p. 49)] [= ?*Rhynchomya* AGASSIZ, 1843, p. 152 (type, *Cercomya gibbosa* AGASSIZ, possibly founded on imperfect specimens of *A. subrugosa*)]. Oblong, compressed, some specimens with trigonal tendency; umbones near mid-length; oblique internal umbonal plate present; hinge and pallial line unknown; ornament consisting of folds which follow or are slightly oblique to growth lines and are best developed at ends of shell; minute, radially aligned granules are also present. *U.Jur.*(*U.Kimmeridg.*) - *L.Cret.*(*Apt.*), Eu.(Ger.-Switz.-France-Eng.)-Afr.(Congo).—FIG. F23,9. **P. subrugosa* (D'ORBIGNY), *U.Jur.*(Portland.), France; *9a,b*, LV ext. and both valves showing variability, $\times 1$ (de Loriol, 1868, 1872).

Family LYONSIDAE Fischer, 1887

[Materials for this family prepared by MYRA KEEN]

Thin, subnacreous, inequivalve, beaks not slit; hinge edentulous; ligament sunken or submarginal, lithodesma present, median; pallial sinus distinct (508, 522). *Eoc.-Rec.*

Lyonsia TURTON, 1822 [**Mya striata* MONTAGU, 1815 (= **M. norvegica* GMELIN, 1791); M] [= *Hiatella* BROWN, 1827 (non BOSC, 1801) (obj.); *Magdala* BROWN, 1827 (obj.; M); *Osteodesma* DE-SHAYES, 1830, AUCTT. (non DE BLAINVILLE, 1825);

Myatella BROWN, 1833 (obj.; M); *Pandorina* SCACCHI, 1833 (non BORY ST. VINCENT, 1827) (obj.; M)]. Sculpture of radial lines. *Eoc.-Rec.*, W.N.Am.-Eu.
L. (*Lyonsia*). Elongate, posterior end attenuate,




FIG. F23. Laternulidae (p. N845).

- slightly gaping; sculpture of radial striations and raised threads. *Eoc.-Rec.*, N.Am.-Eu.—FIG. F24,1. **L. (L.) norwegica* (GMELIN), *Rec.*, Eng.; 1a,b, LV ext., int., 1c, RV hinge, $\times 1$ (7c).
- L. (Bentholyonsia) HABE, 1952** [**Allogramma (B.) teramachii* HABE, 1952; OD]. Sculpture of radial granules but not ribs; lithodesma large; posterior end short, gaping. *Rec.*, W.Pac.—FIG. F24,3. **L. (B.) teramachii* (HABE), Japan; 3a-d, RV ext., int., LV and RV hinges, $\times 1$ (Habe, 1952).
- L. (Endomargarus) COSSMANN, 1887** [**L. heberti* DESHAYES, 1857; OD]. Shape irregular because of nestling habit; sculpture of radial punctuations. *Eoc.*, Eu.—FIG. F24,2. **L. (E.) heberti* DESHAYES, France; 2a,b, LV, RV int., $\times 3$ (160).
- Allogramma DALL, 1903** [**Lyonsia formosa* JEFFREYS, 1881; OD]. Valves with radial and vertical undulations; posterior end gaping. *Rec.*, Eu.-N.Am.-NW.Afr.-Pac.—FIG. F24,6. **A. formosum* (JEFFREYS), NW.Afr.; 6a-c, LV ext., int., dorsal view of both valves, $\times 2$ (852).
- Entodesma PHILIPPI, 1845** [**E. chilensis*; M] [= *Tetragonostea* HERRMANNSEN, 1849 (*ex* DESHAYES, vernac.) (type, *Mya solemyalis* LAMARCK, 1818; M); *Philippina* DALL & SIMPSON, 1901 (type, *Lyonsia beana* D'ORBIGNY, 1853; OD)]. Periostracum thickened; shell irregularly rhomboidal, deformed by nestling; resilifer in each valve relatively small. *Rec.*, N.Am.-S.Am.-S.Pac.
- E. (Entodesma).** Small to medium-sized, not gaping; periostracum adherent; commensal with sponges and tunicates. *Rec.*, W.Indies-W.S.Am.-N.Am.-S.Pac.—FIG. F24,4. **E. (E.) chilense* PHILIPPI, W.S.Am.; LV ext., $\times 1$ (124b).
- E. (Agriodesma) DALL, 1909** [**Lyonia saxicola* BAIRD, 1863; M]. Large, gaping ventrally; periostracum conspicuously thick, leathery, friable; ligament large. *Rec.*, W.N.Am.—FIG. F24,7. **E. (A.) saxicola* (BAIRD), USA (NW. coast); 7a,b, LV ext., RV int., $\times 0.3$ (Reeve, 1875).
- E. (Phlycticoncha) BARTSCH & REHDER, 1940** [*nom. subst. pro Phlyctiderma* BARTSCH & REHDER, 1939 (*non* DALL, 1899)] [**Lyonsia lucasana* BARTSCH & REHDER, 1939; OD]. Narrowed anteriorly, periostracum thin; surface with radial rows of pustules obsolete near margins. *Rec.*, W.C.Am.-W.S.Am.—FIG. F24,10. **E. (P.) lucasanum* (BARTSCH & REHDER), W.Mexico; 10a,b, LV ext., RV int., $\times 1$ (Stanford Univ. specimen).
- Mytilimeria CONRAD, 1837** [**M. nuttallii*; M]. Thin, fragile, ventricose, with thin, adherent periostracum; beaks subspiral; hinge edentulous but with one or more ligamental pits; muscle scars small, pallial sinus inconspicuous. *Rec.*, W.N.Am.—FIG. F24,8. **M. nuttallii*, USA (Calif.); 8a,b, RV int., ext., $\times 1$ (7c).
- Ostomya CONRAD, 1874** [**O. papyria*; M] [= *Himella* H. ADAMS, 1860 (*non* DALLAS, 1852) (type, *H. fluviatilis*; M); *Anticorbula* DALL, 1898 (*pro Himella*); *Guianadesma* MORRISON, 1943 (type, *G. sinuosum*; OD)]. Thin, concentrically plicate; hinge with spoon-shaped process and small tooth near apex in LV; RV with oblique chondrophore. [Estuarine.] *U.Oligo.-Rec.*, S.Am.—FIG. F24,5. **O. papyria*, Plio., Peru; RV ext., $\times 2$ (Conrad).

Family MARGARITARIIDAE Vokes, 1964

[Materials for this family prepared by MYRA KEEN]

Shell moderately large, subcylindrical, gaping, nearly smooth except for some widely spaced radial ribs and surface irregularities; inner layer nacreous; hinge area wide but without teeth, beaks low; adductor muscle scars unequal in size, posterior elongate, anterior rounded; pallial line entire or only slightly sinuate. *Mio.*

Margaritaria CONRAD, 1849 [**Pholadomya abrupta* CONRAD, 1832; M] [= *Actinomya* MAYER, 1870 (obj.; OD)]. Differing from *Panopea*, which it resembles, by edentulous hinge, radial ribbing, and nacreous shell material. *Mio.*, E.N.Am.—FIG. F24,9. **M. abrupta* (CONRAD), USA (Md.); 9a,b, LV ext., RV int., $\times 0.5$ (Glenn, 1904).

Family MYOCHAMIDAE Bronn, 1862

[Materials for this family prepared by MYRA KEEN]

Inequivalve, free or sessile, subnacreous; hinge edentulous, dorsal margins overlapping; ligament external or wanting, resilium internal; pallial sinus small. *Mio.-Rec.*

Myochama STUTCHBURY, 1830 [**M. anomioides*; M]. Sessile, attached by RV and modified in form by surface of attachment. *Mio.-Rec.*, Australasia.—FIG. F25,1. **M. anomioides*; 1a,b, LV ext., RV int., $\times 1$ (305).

Myadora GRAY, 1840 [**Pandora brevis* SOWERBY, 1829; M] [= *Myodora*, spelling error]. Free, resembling *Pandora*, with flattened cardinal area on either side of beaks. *Mio.-Rec.*, Australasia-W.Pac.

M. (Myadora). Surface irregular but not squamose; RV convex, LV flat; hinge with triangular resilifer; pallial sinus rather shallow. *Mio.-Rec.*, Australasia-W.Pac.—FIG. F25,3. **M. (M.) brevis* (SOWERBY), *Rec.*, Australia; 3a-c, RV int., LV ext., RV int., $\times 1$ (124b, 305).

M. (Hunkydora) FLEMING, 1948 [**Thracia transenna* SUTER, 1913 (= **T. australica novozelandica* REEVE, 1859); OD]. Surface squamose or granular, with low concentric folds; hinge with bladeli-like resilifer obliquely directed backward;

pallial sinus broader, deeper, and more open than in *M. (Myadora)*. *U.Plio.-Rec.*, N.Z.—FIG. F25,4. **M. (H.) novozelandica* (REEVE), *Rec.*; 4a, LV hinge, $\times 4$; 4b,c, LV ext., int., $\times 1.3$ (Fleming, 1948).

M. (Myadoropsis) HABE, 1960 [**Thracia trans-*

montana YOKOYAMA, 1922; OD]. Somewhat inflated; hinge with 2 distinct toothlike knobs below umbo in RV; pallial sinus deep. *Pleist.-Rec.*, W.Pac.-Australasia.—FIG. F25,2. **M. (M.) transmontana* (YOKOYAMA), *Rec.*, Japan; 2a,b, RV int., ext., $\times 2.5$ (Habe, 1960).




FIG. F24. Lyonsiidae (1-8,10); Margaritariidae (9) (p. N846-N847).




FIG. F25. Myochamidae (p. N847-N848).

Family PERIPLOMATIDAE Dall, 1895

[nom. correct. DALL, 1900 (pro Periplomidae DALL, 1895)]
 [Materials for this family prepared by MYRA KEEN]

Subnacreous, strongly inequivalve; beaks with a slit; hinge edentulous, resilium in two spoon-shaped chondrophores directed anteriorly or downward; pallial sinus mostly wide but shallow (524). *U.Cret.-Rec.*

Periploma SCHUMACHER, 1817 [**P. inaequivalvis* (= **Corbula margaritacea* LAMARCK, 1801); M]. Thin, RV more convex than LV and overlapping it, surface granular, beaks opisthogyrate; chondrophores supported by clavicle in most forms. *U.Cret.-Rec.*, cosmop.

P. (Periploma). Ovate-quadrangle, smooth; anterior muscle scar long and narrow, posterior small and crescentic; pallial sinus short, rounded. *U.Cret.-Rec.*, N.Am.-S.Am.-E.Asia-Eu.—FIG. F26, 1. **P. (P.) margaritaceum* (LAMARCK), Rec., W.Indies; 1a-c, LV int., LV view of both valves, RV hinge, $\times 1$ (H. Adams & A. Adams).

P. (Aelga) SLODKEVICH, 1935 [**Tellina besshoensis* YOKOYAMA, 1924; OD]. Large, ovate-quadrangle, with undulating concentric sculpture; chondrophores massive and shallow, clavicle long, curved. *Oligo.-Mio.*, NE.Asia.—FIG. F26, 7. **P. (A.) besshoense* (YOKOYAMA), Mio., Japan; 7a, b, RV ext., dorsal view of both valves, $\times 0.5$ (Makiyama, 1957).




FIG. F26. Periplomatidae (p. N849-N850).

- at margins; inequivalve, LV smaller. *Rec.*, W.C.Am.—FIG. F26,3. **P. (A.) pentadactylus*, W. Panama; *3a,b*, RV ext., int., $\times 1$ (Pilsbry & Olsson, 1935).
- P. (Halistrepta)** DALL, 1904 [**P. sulcata*; M]. Sculpture of irregular concentric ribs. *Rec.*, W.N. Am.-W.Pac.—FIG. F26,6. **P. (H.) sulcatum*, USA (Calif.); RV ext., $\times 0.8$ (Dall).
- P. (Offadesma)** IREDALE, 1930 [**P. angasi* CROSSE & FISCHER, 1864; OD]. Large, thin, oblique, chondrophore directed downward; pallial sinus triangular, deep, narrow. *Rec.*, S.Pac.-W.Pac.—FIG. F26,4. **P. (O.) angasi* CROSSE & FISCHER, S.Australia; *4a,b*, LV ext., post. view of both valves, $\times 0.5$ (Cotton, 1938).
- P. (Pentaloma)** IREDALE, 1930 [**P. micans* HEDLEY, 1901; OD]. Posterior area rostrate, set off by rib; periostracum thin, obliquely ridged; shell with concentric undulations and 1 or 2 radial furrows; chondrophore small, directed downward. *Rec.*, S.Pac.—FIG. F26,5. **P. (P.) micans* HEDLEY, E.Australia; *5a-c*, RV ext., dorsal, int., $\times 1.5$ (396); *5d*, RV hinge, $\times 7$ (Hedley).
- Cochlodesma** COUTHOUY, 1839 [**Anatina leana* CONRAD, 1831; SD HERRMANNSEN, 1847 (Mar.)] [= *Aperiploma* HABE, 1952 (OD, obj.)]. Lenticular, subequilateral; lithodesma present; chondrophore buttressed. *Mio.-Rec.*, Eu.-Atl.-Pac.
- C. (Cochlodesma)**. Slightly inflated, surface not granulose; lithodesma cartilaginous. *Mio.-Rec.*, Eu.-W.Atl.-S.Pac.-NW.Pac.—FIG. F26,2. **C. (C.) leanum* (CONRAD), *Rec.*, USA (Mass.); RV ext., $\times 1$ (Gould & Binney, 1870).
- C. (Bontaea)** BROWN, 1844 (*ex* LEACH MS) (in synonymy of *Ligula praetenuis* [**Chama praetenuis* PULTENEY, 1799; M] [= *Galaxura* GRAY, 1852 (*ex* LEACH MS) (M, obj.); *Calcaraea* RÉCLUZ, 1868 (M, obj.)]. Less rounded than *C. (Cochlodesma)*, flatter, posterior somewhat attenuated; surface granulose; lithodesma small. *Plio.-Rec.*, N.Eu.-S.Eu.—FIG. F26,8. **C. (B.) praetenuis* (PULTENEY); *Rec.*, Eng., *8a*, LV ext., $\times 1$ (Jeffreys, 1869); *Rec.*, Norway, *8b,c*, LV and RV hinges, $\times 2$ (857).
- Family THRACIIDAE Stoliczka, 1870**
- [*nom. transl.* DALL, 1903 (*ex* Thraciinae STOLICZKA, 1870)] [= *Osteodesmacea* DESHAYES, 1839 (invalid family-group name, Code Art. 11, e); *Osteodesmidae* HÖRNES, 1859 (invalid family-group name, Code Art. 11, e)] [Materials for this family prepared by MYRA KEEN]
- Smooth, nonnacreous, inequivalve (RV larger), surface granular in most; hinge edentulous; chondrophore directed obliquely toward posterior end; pallial line with sinus (510). *Jur.-Rec.*
- Thracia** SOWERBY, 1823 (*ex* LEACH MS) [**T. pubescens* LAM.] (= **Mya pubescens* PULTENEY, 1799); SD ANTON, 1839] [= *Throna* CARPENTER, 1859 (*nom. null.*); *Osteodesma* DE BLAINVILLE, 1825 (OD, obj.); *Odoncineta* COSTA, 1829 (type, *O. papyracea*; M); *Odontocineta*, *Cinetodonta* (*nom. van. pro* *Odoncineta*); *Corymya* AGASSIZ, 1843 (type, *Mya depressa* SOWERBY, 1823, *non* DONOVAN, 1802, = *Corimya pinguis* AGASSIZ, 1845; M; *Jur.*, Eu.); *Corimya*, *Coromya* (*nom. null.*); *Homoeodesma* FISCHER, 1887 (type, *T. conradi* COUTHOUY, 1839; M); *Eximiothracia* IREDALE, 1924 (type, *T. speciosa* ANGAS, 1869; OD); *Cetothrax* IREDALE, 1949 (type, *T. alciopae* ANGAS, 1872, = *Anatina imperfecta* LAMARCK, 1818; OD)]. Oblong, somewhat inflated, nearly equilateral, posterior end broadly truncate and set off by low ridge; periostracum present in some. *Jur.-Rec.*, Eu.-N.Am.-Pac.-Atl.-Afr.-Arctic.
- T. (Thracia)**. Centrally striate; surface granulate; beaks touching and perforate; ligament external; cardinal plate with fissure occupied by lithodesma; chondrophore moderately long. *Jur.-Rec.*, Eu.-N.Am.-Pac.—FIG. F27,11. **T. (T.) pubescens* (PULTENEY), *Rec.*, Eng.; *11a-c*, LV int., RV hinge, LV view both valves, $\times 1$ (H. Adams & A. Adams, 1858).
- T. (Crassithracia)** SOOT-RYEN, 1941 [**T. crassa* BECHER (= **T. septentrionalis* JEFFREYS, 1872); OD]. Solid, smooth, periostracum present, thin, shining, except on posterior truncation; chondrophore very narrow, horizontal. *Rec.*, N.Atl.—FIG. F27,2. **T. (C.) septentrionalis* JEFFREYS, Norway; *2a*, RV ext., $\times 0.5$; *2b,c*, RV and LV hinges, enl. (857).
- T. (Ixartia)** GRAY, 1852 (*ex* LEACH MS) [**Mya distorta* MONTAGU, 1803; M] [= *Rupicola* FLEURIAU, 1802 (*non* BRISSON, 1760) (obj.); *Pelopia* H. ADAMS, 1868 (*non* MEIGEN, 1800) (type, *P. brevifrons*; M); *Rupicilla* SCHAUFUSS, 1869 (*pro* *Rupicola*)]. Distorted by nestling habit, inequilateral; chondrophore free from hinge plate posteriorly; pallial sinus large. *Rec.*, Eu.-W.N.Am.-S.Afr.—FIG. F27,3. **T. (I.) distorta* (MONTAGU); Eng.; *3a*, LV ext., $\times 1$ (Jeffreys, 1869); Norway, *3b,c*, hinge, lat. and ventral views, $\times 2$ (857).
- T. (Lampeia)** MACGINITIE, 1959 [**T. (L.) adamsi*; OD]. Chondrophore buttressed, forming depression under beaks. *Rec.*, Arctic.—FIG. F27,8. **T. (L.) adamsi*, USA (Alaska); *8a,b*, LV ext., RV ext., $\times 1$; *8c,d*, RV and LV hinges, $\times 3$ (MacGinitie, 1959).
- T. (Trigonothracia)** YAMAMOTO & HABE, 1959 [**T. (T.) nomurai*; OD]. Small, thin, chondrophore small, trigonal, directed downward. *Rec.*, W.Pac.—FIG. F27,5. **T. (T.) nomurai*, Japan; *5a,b*, LV ext., int., $\times 2$ (Yamamoto & Habe, 1959).
- Asthenothaerus** CARPENTER, 1864 [**A. villosior*; M]. Edentulous, surface somewhat granular; cartilage below umbones, not in chondrophore; pallial sinus large. *Rec.*, E.C.Am.-W.N.Am.—FIG.

F27,10. **A. villosior*, W.Mex.; 10a-c, LV ext., int., RV int., $\times 2.5$ (specimen, U.S.Natl.Mus.).
Bushia DALL, 1886 [**B. elegans*; M]. Resembling *Asthenothaerus* but surface without granulations;

ligament external; apices solid within, supporting resilium. *Rec.*, W.Indies-C.Am.-S.Am.—FIG. F27,12a. **B. elegans*, W.Indies; LV ext., $\times 3$ (Dall, 1889).—FIG. F27,12b,c. *B. panamensis*




FIG. F27. Thraciidae (p. N850-N852).

- DALL, Panama; 12*b,c*, RV int., LV oblique view inside apex, $\times 2$ (specimen, U.S.Nat.Mus.).
- Cyathodonta** CONRAD, 1849 [**C. undulata*; M]. With strong undulating and somewhat oblique folds; beaks nonperforate; resilium small, chondrophore short, rounded, prominent, lithodesma thin, semicircular, vertical. *Pleist.-Rec.*, N.Am.-C.Am.-S.Am.—FIG. F27,7. **C. undulata*, Rec., W.Mex.; 7*a,b*, RV ext., LV int., $\times 0.5$ (Stanford Univ. specimen).
- Parvithracia** FINLAY, 1927 [**P. suteri* (*pro Montacuta triquetra* SUTER, 1913) (*non* VERRILL & BUSH, 1898); OD]. Small, like miniature *Thracia* with deep pallial sinus. *Mio.-Rec.*, S.Pac.—FIG. F27,1. **P. suteri*, Rec., N.Z.; 1*a,b*, LV ext., RV int., $\times 10$ (modified from Suter, 1913).
- Phragmorisma** TATE, 1894 [**Thracia watsoni* SMITH, 1885; SD LAMY, 1931]. Inequivalve, RV larger, resembling *Thracia* but with large subumbonal plate in each valve hanging from umbo; pallial sinus broad. *Oligo.-Rec.*, Australasia.—FIG. F27,9. **P. watsoni* (SMITH), Rec., N.Australia; 9*a,b*, RV ext., int., $\times 0.5$ (852).
- Thracidora** IREDALE, 1924 [**Thraciopsis arenosa* HEDLEY, 1904; OD]. Oblong, equilateral, ventricose, gaping behind; ligament external; shell surface radially granular. *Rec.*, S.Pac.—FIG. F27,6. **T. arenosa* (HEDLEY), Australia; 6*a,b*, RV ext., hinge, $\times 6$ (Hedley, 1904).
- Thraciopsis** TATE & MAY, 1900 [*pro Alicia* ANGAS, 1867 (*non* JOHNSON, 1861)] [**Alicia angustata* ANGAS, 1867; SD STOLICZKA, 1871] [=?*Thracidentula* GARRARD, 1961 (type, *T. perulae*; OD)]. Inequivalve, quadrate, very inequilateral, posterior end short and truncate; beaks not fissured; interior subnacreous; hinge with posterior callosity in RV and a socket in LV, with marginal plate anteriorly; resilium below beaks under large triangular lithodesma; pallial sinus large. *Rec.*, S.Pac.—FIG. F27,4. **T. angustata* (ANGAS), Australia; 4*a-c*, LV ext., RV int., LV hinge, $\times 2$ (Angas, 1867).

Superfamily POROMYACEA Dall, 1886

[*nom. transl.* DALL, 1895 (*ex* Poromyidae DALL, 1886)]
[Materials for this superfamily prepared by MYRA KEEN]

Rounded to elongate, generally not gaping; hinge with somewhat developed cardinal and lateral teeth; resilium reinforced by lithodesma; pallial sinus small. Gills scantily reticulate or even absent; mantle lobes united. *Cret.-Rec.*

Family POROMYIDAE Dall, 1886

Rounded-quadrate; shell surface granulate in most, tending toward radial ribbing;

interior subnacreous to nacreous; ligament external; hinge with cardinal tooth in one valve. *Cret.-Rec.*

- Poromya** FORBES, 1844 [**P. anatinoides* (= **Corbula granulata* NYST & WESTENDORP, 1839); M] [= *Embla* LOVÉN, 1847 (type, *E. korenii*; M, = *Corbula granulata* NYST & WESTENDORP, 1839); *Ectorisma* TATE, 1892 (type, *E. granulata*; OD; = *P. laevis* SMITH, 1885); *Questimya* IREDALE, 1930 (type, *P. undosa* HEDLEY & PETTERD, 1906; OD)]. Nacreous, thin, posterior slope set off by angle; surface granular. *Cret.-Rec.*, cosmop.
- P. (Poromya)**. Hinge with strong cardinal in RV, socket in LV, and ridgelike *Pil*. *Cret.-Rec.*, Eu.-E.N.Am.-W.N.Am.-worldwide (deep water).—FIG. F28,1. **P. (P.) granulata* (NYST & WESTENDORP), Rec., Atl.; 1*a,b*, RV ext., LV hinge, $\times 3$ (124*b*).
- P. (Cetomya)** DALL, 1889 [**P. elongata* DALL, 1886; SD GLIBERT, 1936]. Hinge teeth obsolete in adult; pallial sinus wanting. *Rec.*, Atl.—FIG. F28,4. **P. (C.) elongata* DALL; LV ext., $\times 1$ (Dall, 1889).
- P. (Dermatomya)** DALL, 1889 [**P. (D.) mactroides*; M]. Surface not granular; hinge strong; pallial sinus developed. *Rec.*, W.N.Am.-S.Am.-S.Afr. (deep water).—FIG. F28,5. **P. (D.) mactroides*, Chile; LV ext., $\times 1.5$ (Dall, 1890).
- P. (Mioporomya)** SACCO, 1901 [**M. taurinensis*; OD]. Larger than *P. (Poromya)*, with heavier hinge and strong radial sulci. *Mio.*, Eu.—FIG. F28,10. **P. (M.) taurinensis* (Sacco), Italy; 10*a-d*, RV ext., int., LV int., ext., $\times 1$ (Sacco, 1901).
- Cetoconcha** DALL, 1886 [**Lyonsia bulla* DALL, 1878; OD] [= *Silenia* SMITH, 1885 (*non* RYE, 1873) (type, *S. sarsii*; M)]. Differing from *Poromya* in having fossettes smaller and upturned, resilium almost external, ligament obsolescent; hinge teeth weak or absent except cardinal in RV, present in young, weaker in adult. *Rec.*, Atl.-S.Pac. (deep water).—FIG. F28,8. **C. bulla* (DALL), Atl.; 8*a,b*, LV int., ext., $\times 2$ (Dall, 1889).
- Cymella** MEEK, 1864 [**Pholadomya undata* MEEK & HAYDEN, 1856; OD]. Small, ovate, with many regular concentric undulations, crossed midway by some radial lines impressed on crests of waves. *U.Cret.*, N.Am.—FIG. F28,7*a*. **C. undata* (MEEK & HAYDEN), USA(Mont.); LV ext., $\times 1$ (Meek, 1876).—FIG. F28,7*b,c*. *C. bella* CONRAD, USA(N.J.); 7*b,c*, RV int., LV int., $\times 1$ (Meek, 1876).
- Liopistha** MEEK, 1864 [**Cardium elegantulum* ROEMER, 1852 (*non* BECK, 1842) (= **L. elegantulata* VOKES, 1956); OD]. Thin, ovate, posterior end somewhat attenuate; hinge with 2 cardinal teeth, no laterals; ligament external; pallial area indistinct. *U.Cret.*, N.Am.-Afr.-Asia.
- L. (Liopistha)**. Sculpture of fine radial ribs except on dorsal part of posterior slope. *U.Cret.*,

N.Am.-Afr.—FIG. F28,2. **L. elegantulata* VOKES, USA(Tex.); 2a, RV ext., $\times 1$; 2b, sculpture, ext. (Roemer, 1852).

L. (*Psilomya*) WHITE, 1874 (ex MEEK MS) [**L. (P.) meekii*; M]. Radial ribs present as rows of granules or spines. U.Cret., N.Am.-Asia.—FIG. F28,3. **L. (P.) meekii*, USA(Utah); RV ext., $\times 1$ (White, 1874).

Neeroporomya COSSMANN, 1887 [**Corbula argentea* LAMARCK, 1807; OD]. Shell subnacreous, without surface punctations; posterior slope with concentric folds; single tooth in each valve, no in-

ternal fossette or posterior lamella; pallial line sinuous. Eoc., Eu.—FIG. F28,6. **N. argentea* (LAMARCK), France; 6a, LV ext., $\times 3$ (Vincent, 1898); 6b,c, RV, LV hinges, $\times 3$ (Cossmann, 1914).

Pseudocrepidaria EAMES, 1951 [**Crepidaria lakienensis* Cox, 1938; OD]. Resembling *Crepidaria* in outline but larger; subequivalve; sculpture of strong concentric folds; shell very thin, nacreous within. Eoc., Asia.—FIG. F28,9. **P. lakienensis* (Cox), NW.India; 9a,b, both valves dorsal, RV ext., $\times 1$ (Cox, 1938).




FIG. F28. Poromyidae (p. N852-N853).

Family CUSPIDARIDAE Dall, 1886

Small to medium-sized, thin, ovate, mostly strongly rostrate; hinge with external ligament; resilium on small resilifer; hinge teeth present or absent. Carnivorous pelecypods, gill lamellae forming muscular partition across mantle cavity. *U.Cret.-Rec.*

Cuspidaria NARDO, 1840 [**C. typus* (= *Tellina cuspidata* OLIVI, 1792); M] [= *Neaera* GRIFFITH, 1834 (non ROBINEAU-DESVOIDY, 1830) (type, *N. chinensis*; M); *Aulacophora* JEFFREYS, 1882 (non CHEVROLAT, 1842) (type, *N. lamellosa* SARS, 1858; SD KEEN, herein); *Vulcanomya* DALL, 1886 (type, *V. smithii*, nom. dub.; M)]. Inequivalve, LV more convex; hinge with resilifer and one or more teeth; posterior end of shell strongly rostrate. *U.Cret.-Rec.*, Eu.-N.Am.-Atl.-Pac.-Medit.-IndoPac.

C. (Cuspidaria). Hinge with single posterior lateral tooth in RV; exterior of shell feebly sculptured; resilium in chondrophore or fossette that is inclined and attached by posterior edge. *U.Cret.-Rec.*, Eu.-N.Am.—FIG. F29,5. **C. (C.) cuspidata* (OLIVI), *Rec.*, *Medit.*; 5a-c, LV int., ext., RV hinge, $\times 1$ (H. Adams & A. Adams, 1858).

C. (Halonympha) DALL, 1886 [**Neaera claviculata* DALL, 1881; OD]. Smooth or concentrically striate; hinge with single tooth, acute cardinal in RV; fossette small, central; posterior clavicular rib in each valve. *Rec.*, *Atl.*—FIG. F29,6. **C. (H.) claviculata* (DALL), *Carib.*; 6a,b, RV ext., int., $\times 2$ (216).

C. (Leiomya) A. ADAMS, 1864 [**Neaera adunca* GOULD, 1861; M]. Smooth, hinge with anterior cardinal in each valve; RV with anterior and posterior laterals, LV without laterals. *Rec.*, *Pac.*

C. (Pseudoneaera) STURANY, 1902 [**P. thaumasia*; M] [= *Bendoneaera* GOSSMANN, 1904 (nom. van.)]. Rostrum short, pointed; hinge with shallow fossette, anterior and posterior laminae in RV and denticle in LV in front of fossette. *Rec.*, *IndoPac.*—FIG. F29,12. **C. (P.) thaumasia* (STURANY), *Red Sea*; 12a-c, LV ext., LV and RV hinges, $\times 5$ (Sturany, 1902).

C. (Rhinoclama) DALL, 1886 [*pro Rhinomya* A. ADAMS, 1864 (non ROBINEAU-DESVOIDY, 1830)] [**Neaera philippinensis* HINDS, 1843; SD STOLICZKA, 1871] [= *Luzonia* DALL, 1890 (obj.; OD)]. Surface concentrically striate, not granulate; hinge without cardinal teeth. *Rec.*, *Pac.*—FIG. F29,13. **C. (R.) philippinensis* (HINDS), *Philip.Is.*; 13a-c, RV ext., int., LV int., $\times 7.5$ (type specimens, British Museum).

C. (Tergulina) NOSZKY, 1939 [**Neaera (T.) sulcosa*; M]. Surface with fine concentric ribs. *Oligo.*, *Eu.*—FIG. F29,4. **C. (T.) sulcosa*

(NOSZKY), *Hung.*; RV ext., $\times 1.3$ (Noszky, 1939).

C. (Tropidomya) DALL, 1886 [*pro Tropidophora* JEFFREYS, 1882 (non TROSCHEL, 1847)] [**Neaera abbreviata* FORBES, 1843; SD DALL, 1886] [= *Goniophora* JEFFREYS, 1883 (non PHILLIPS, 1848) (*pro Tropidophora* JEFFREYS, 1882, non TROSCHEL, 1847)]. With anterior cardinal in each valve, no lateral teeth. *Rec.*, *Atl.-Medit.*—FIG. F29,3. **C. (T.) abbreviata* (FORBES), *Medit.*; RV ext., $\times 3$ (Forbes & Hanley, 1848).

Austroneaera POWELL, 1937 [**A. brevisstris*; OD]. Thin, smooth, rostrum poorly developed; LV edentulous, RV with laterals; sinus broad, rounded. *Rec.*, *S.Pac.*—FIG. F29,7. **A. brevisstris*, *N.Z.*; 7a-c, RV ext., int., LV int., $\times 5$ (Powell, 1937).

Boriesia DONCIEUX, 1911 [**B. cossmanni*; OD]. Shape and sculpture somewhat as in *Cardiomya*; hinge unknown. *Paleoc.*, *Eu.*—FIG. F29,10. **B. cossmanni*, *Ypres.*, *France*; 10a,b, RV ext., both valves dorsal, $\times 2$ (Doncieux, 1911).

Cardiomya A. ADAMS, 1864 [**Neaera gouldiana* HINDS, 1843; M] [= *Spathophora* JEFFREYS, 1882 (non AMYOT & SERVILLE, 1843) (type, *Neaera curta* JEFFREYS, 1876; SD KEEN herein)]. Surface radially ribbed. *U.Cret.-Rec.*, *N.Am.-C.Am.-Pac.*

C. (Cardiomya). RV with prominent posterior lateral tooth. *U. Cret.-Rec.*, *N.Am.-Pac.*—FIG. F29,2. **C. (C.) gouldiana* (HINDS), *Rec.*, *E.Indies*; RV ext., $\times 1.5$ (124b).

C. (Bowdenia) DALL, 1903 [**Cuspidaria distira*; OD]. Without laterals; margin of RV grooved. *Mio.*, *E.C.Am.*—FIG. F29,1. **C. (B.) distira* (DALL), *Jamaica*; 1a, LV ext., $\times 6$ (Dall, 1903); 1b,c, LV int., RV int., $\times 6$ (1005).

?**Fabagella** GOSSMANN, 1887 [**Corbula faba* DESHAYES, 1824; OD]. Carinate, umbones acute, lunule deep; hinge edentulous, fossette triangular. *Eoc.*, *Eu.*—FIG. F29,9. **F. faba* (DESHAYES), *France*; 9a,b, LV int., RV hinge, $\times 3$ (160).

Myonera DALL, 1886 [**Neaera paucistriata* DALL, 1885; OD]. Sculpture radiating or concentric; hinge without teeth; fossette vertical or posteriorly directed. *Rec.*, *Atl.-Pac.*—FIG. F29,8. **M. paucistriata* (DALL), *Atl.*; LV ext., $\times 2$ (Dall, 1890).

Plectodon CARPENTER, 1864 [**P. scaber*; OD]. Surface granular; hinge with pseudocardinal under small external ligament; resilium below beaks, not in fossette. *Plio.-Rec.*, *E.N.Am.-W.N.Am.*—FIG. F29,11. **P. scaber*; 11a-c, LV ext., int., RV int., $\times 2$ (Stanford Univ. specimen).

Family VERTICORDIIDAE Stoliczka, 1871

[=Euciroidae DALL, 1894]

Small (few more than 25 mm. in diameter), cordate, nacreous within, more or less

strongly ribbed radially; hinge mostly weak or with one or two low teeth. *Paleoc.-Rec.*

Verticordia SOWERBY, 1844 (ex WOOD MS) [**Hippagus? cardiiformis* SOWERBY, 1844; M] [= *Trigonulina* D'ORBIGNY, 1846 (type, *T. ornata*; M) *Iphigenia* COSTA, 1850 (non SCHUMACHER, 1817) (type, *Hippagus acuticostatus* PHILIPPI, 1844; OD); *Hippella*, AUCTT. (non MÖRCH, 1861)]. Small to minute, ligament external; ribs well developed. *Paleoc.-Rec.*, Eu.-Carib.-N.Am.-C.Am.-Pac.

V. (*Verticordia*). Ribs raised, well spaced; lunule deep in LV, shallower in RV; hinge with cardinal in RV, lateral behind beaks in LV. *Paleoc.-Rec.*, Eu.-Carib.-W.N.Am.-C.Am.-W.Pac.—FIG. F30, 2. **V. (V.) cardiiformis* (SOWERBY), Plio., Eng.; 2a,b, LV ext., int., ×2 (1004).

V. (*Spinospella*) IREDALE, 1930 [**V. ericia* HEDLEY, 1911; OD]. Borders of ribs prickly; lunule wanting. *Rec.*, Pac.—FIG. F30,5. **V. (S.) ericia* HEDLEY, S.Australia; 5a,b, RV ext., hinge, ×4; 5c, surface detail, ext. (Cotton, 1938).

V. (Vertambitus) IREDALE, 1930 [**V. vadosa*




FIG. F29. Cuspidariidae (p. N854).

HEDLEY, 1907; OD]. Minute, solid, granose, hinge heavy. *Rec.*, S.Pac.—FIG. F30,7. **V. (V.) vadosa* HEDLEY, Australia; 7a-d, LV ext., dorsal, LV and RV hinges, $\times 10$ (Hedley, 1907). *V. (Vertisphaera)* IREDALE, 1930 [**V. cambrica*; OD]. Thin, nearly smooth, ribs as fine radial threads; hinge narrow. *Rec.*, S.Pac.—FIG.

F30,6. **V. (V.) cambrica* IREDALE, Australia; 6a,b, LV dorsal, int., $\times 2.5$ (Hedley, 1907). *Euciroa* DALL, 1881 (in synonym.) [**Verticordia elegantissima* DALL, 1881; M]. Relatively large, with numerous granular ribs; lunule and escutcheon present; resiliary ossicle triangular, wider behind. *Rec.*, Atl.-Pac.




FIG. F30. Verticordiidae (p. N854-N857).

E. (*Euciroa*). Valves subequal, hinge well developed. *Rec.*, Atl.-Pac.—FIG. F30,3. *E. (*E.*) *elegantissima* (DALL), Carib.; 3a,b, LV ext., int., $\times 2$ (216).

E. (*Acreuciroa*) THIELE & JAECKEL, 1931 [*E. (*A.*) *rostrata*; M]. Hinge weak; LV broadened. *Rec.*, Pac.—FIG. F30,4. *E. (*A.*) *rostrata*, Japan; 4a-c, RV ext., RV and LV hinges, $\times 0.6$ (Thiele & Jaekel, 1931).

Halicardia DALL, 1895 [**Mytilimeria flexuosa* VERRILL, 1881; OD] [= *Halocconcha*, AUCTT., spelling error]. Dorsal margin flaring at ends; surface of shell granular; hinge obsolete, lunule in RV larger; lithodesma asymmetrical, right limb longer; shell slightly inequivalve. *Rec.*, Atl., Pac.—FIG. F30,11. *H. *flexuosa* (VERRILL), N.Atl.; RV ext., $\times 1$ (Verrill, 1881).

Halicardissa DALL, 1913 [**Verticordia perplicata* DALL, 1890; OD]. Resembling *Halicardia* in form, few-ribbed; soft parts as in *Verticordia*. *Rec.*, Pac.—FIG. F30,10. *H. *perplicata* (DALL), Galapagos Is.; RV ext., $\times 1$ (Dall, 1890).

Haliris DALL, 1886 [**Verticordia fischeriana* DALL, 1881; OD]. Globose, lunule shallow; hinge of RV as in *Verticordia*, LV with single small cardinal and short stout lateral behind beak; resiliary ossicle short, quadrate. *Rec.*, Atl.-Pac.

H. (*Haliris*). Ventral margin even, shell rounded. *Rec.*, Atl.-Pac.—FIG. F30,9. *H. (*H.*) *fischeriana* (DALL), W.Atl.; 9a,b, RV hinge, ext., $\times 3$ (216).

H. (*Setaliris*) IREDALE, 1930 [**Verticordia setosa* HEDLEY, 1907; OD]. Smaller than H. (*Haliris*), more quadrate, ventral margin sinuous. *Rec.*, S.Pac.—FIG. F30,8. *H. (*S.*) *setosa* (HEDLEY), N.Z.; 8a,b, RV ext., dorsal, $\times 5$ (Hedley, 1904).

Kuriniua MARWICK, 1942 [**Trigonia areolata* MARSHALL, 1919; OD]. Ribless area on posterior slope, crossed only by low concentric ridges. *Eoc.*, N.Z.—FIG. F30,12. *K. *areolata* (MARSHALL); 12a,b, RV int., ext., $\times 1.3$ (Marwick, 1942).

Lyonsiella G. SARS, 1872 (ex M. SARS MS) (in synonym.) [**Pecchiolia abyssicola* SARS, 1872; M]. Quadrate, thin, ligament internal, with lithodesma; hinge weak to edentulous; pallial line entire. *Rec.*, Atl.-Pac.

L. (*Lyonsiella*). Surface ribbed, ribs granular in some. *Rec.*, Atl.-Pac.—FIG. F30,1. *L. (*L.*) *abyssicola* (SARS), N.Atl.; 1a-c, LV ext., both valves dorsal, RV int., $\times 4$ (829).

L. (*Proagorina*) IREDALE, 1930 [*L. *quadrata* HEDLEY, 1907; OD]. Surface granular, with some radial furrows, granules prickly; hinge edentulous. *Rec.*, S.Pac.—FIG. F30,13. *L. (*P.*) *quadrata* HEDLEY, Australia; 13a,b, RV ext., int., $\times 6$ (Hedley, 1907).

Laevicordia SEGUENZA, 1876 [**Verticordia* (*L.*) *orbiculata* SEGUENZA, 1876; SD SOOT-RYEN, 1966]. Resembling *Policordia* but more rounded, with fine surface granulation. *Plio.-Rec.*, Eu.-Atl.-Pac.




FIG. F31. Verticordiidae (p. N857).

Pecchiolia SAVI & MENEHINI in MURCHISON, 1850 [*P. *argentea* (= *Chama arietina* BROCCHI, 1814); M]. Resembling *Verticordia* but larger, ribs lower, beaks higher, spirally twisted; hinge tooth larger, more solid. *Eoc.-Plio.*, Eu.—FIG. F31,2. *P. *arietina* (BROCCHI), Plio., Italy; 2a-c, LV ext., int., RV int., $\times 1$ (Hörnes, 1870).

Policordia DALL, BARTSCH, & REHDER, 1939 [*P. *diomedea*; OD] [= *Laevicordia*, AUCTT. (non SEGUENZA, 1876)]. Small, ovate, nearly smooth, with fine radial ribs; ligament in groove, with lithodesma below it; hinge edentulous; pallial sinus small, not deep. *Rec.*, Atl.-Pac.—FIG. F31,1. *P. *diomedea*, Hawaii; 1a-c, RV int., LV ext., int., $\times 2.5$ (Dall, Bartsch, & Rehder, 1939).

Superfamily CLAVAGELLACEA d'Orbigny, 1844

[*nom. transl.* THIELE, 1934 (ex Clavagellidae D'ORBIGNY, 1843) [Materials for this superfamily prepared by MYRA KEEN and L. A. SMITH]]

Shell nacreous, free when young, degenerate in adult; with one or both valves at least partially embedded in elongate calcareous tube, an adaptation for burrowing; hinge-

plate wanting; ligament external. *U.Cret.* (*Turon.*)-*Rec.*

Family CLAVAGELLIDAE d'Orbigny, 1843

[=Aspergillidae GRAY, 1858]

Tube anteriorly rounded or discoid and smooth or fringed with simple or branching tubules, or partially closed by calcareous disc having several pedal foramina. *U.Cret.* (*Turon.*)-*Rec.*

Clavagella LAMARCK, 1818 [**C. echinata*; SD CHILDREN, 1823] [=Bacilia GRAY, 1858 (ex VALENCIENNES MS) (obj.); *Clavigella*, spelling error]. One valve never merging with tube and both adductors persistent in adult. *U.Cret.* (*Turon.*)-*Rec.*, cosmop.

C. (Clavagella). Siphonal end of tube simple; tube free, elongate, clavate, compressed and symmetrical in shape; with irregular spinelike tubules on anterior portion of tube. *U.Cret.* (*Turon.*)-*Rec.*, Eu. - India - Australasia.—FIG. F32,1. **C. (C.) echinata* LAMARCK, Eoc.(Lutet.-Barton.), Paris Basin; $\times 0.4$ (823).

C. (Bryopa) GRAY, 1847 [**C. aperta* SOWERBY, 1823; OD] [=Tiria DE GREGORIO, 1886 (obj.; SD SMITH, 1962)]. Siphonal end of tube periodically expanded, anterior end smooth; with small very short tubules through tube around valves. *U.Oligo.*-(*Aquitanian*)-*Rec.*, Medit. - Ind.O. - Pac.—FIG. F32,8. **C. (B.) aperta* SOWERBY, Rec., Malta; $\times 0.3$ (124b).

C. (Dacosta) GRAY, 1858 [**C. australis* SOWERBY, 1829; M]. Siphonal end of tube not expanded, anterior end smooth, rounded; with small very short tubules through tube around valves. *Rec.*, W.Pac.—FIG. F32,9. **C. (D.) australis* SOWERBY, Australia; $\times 0.6$ (Sowerby in Reeve, 1872, mod.).

C. (Stirpulina) STOLICZKA, 1870 [**C. coronata* DESHAYES, 1824; OD] [=Styrpulina, spelling error; may prove to be synonym of *Tubolana* BIVONA-BERNARDI, 1832 (type, *T. digitata*, =*Aspergillum bacillaris* DESHAYES, 1830, M); *Tubulana* (nom. null.)]. Siphonal end of tube periodically expanded, anterior end with tubules formed only in terminal corona; tube long and with more or less distinct anterior slit. *U.Cret.* (*Turon.*)-*Rec.*, N.Am.-Eu.-Asia.—FIG. F32,2. **C. (S.) coronata* DESHAYES, *U.Eoc.* (*Barton.*), Paris Basin; $\times 0.67$ (124b).

Humphreyia GRAY, 1858 [**Aspergillum strangei* A. ADAMS, 1854; M] [=Humphreysia, spelling error]. Tube twisted and irregularly square in cross section; both valves united into single plate forming most of anterior baglike cavity. *Rec.*, W.Pac.—FIG. F32,7. **H. strangei* (A. ADAMS), Australia; $\times 0.8$ (124b).

Penicillus BRUGUIÈRE, 1789 [**P. javanus* (=Serpula penis LINNÉ, 1758); SD HABE, 1952] [=Brechites GUETTARD, 1770 (nonbinom.); *Penicilli* DACOSTA, 1776 (vernacular) (*Penecillus*, *Peni-*




FIG. F32. Clavagellidae (p. N858-N859).

cellus, nom. null.); *Verpa* RÖDING, 1798 (obj.); *Aquaria* PERRY, 1811 (type, *A. radiata*; SD SMITH, 1962, obj.); *Arytene* OKEN, 1815 (obj.; rejected ICZN, 1956); *Bunodus* DE BLAINVILLE, 1817 (nom. nud.) (ex GUETTARD, nonbinom.); *Arytaena* OKEN, 1817 (obj.) (*Arythaena*, *Arytene*, spelling errors); *Aspergillum* LAMARCK, 1818 (obj.) (*Adaspergillum*, *Aspergillum*, *Aspergillus*, spelling errors); *Clepsydra* SCHUMACHER, 1817 (obj.) (*Clepydra*, spelling error)]. Both valves merging with tube; tube circular in cross section; anterior adductor degenerate, posterior adductor absent in adult. *U.Oligo.*(*Aquitan.*)-*Rec.*, IndoPac.-Australia-Pac.-Japan-Eu.

P. (Penicillus). Siphonal end of tube simple; anterior end fringed with single row of distinct simple tubules; anterior disc with slit. *U.Oligo.*(*Aquitan.*)-*Rec.*, IndoPac.-Australia.—FIG. F32,3. **P. (P.) penis* (LINNÉ), *Rec.*, Singapore; $\times 0.3$ (Haas, 1935).

P. (Foegia) GRAY, 1847 [*Aspergillum novaezealandiae*] (= **P. novaezealandiae* BRUGUIÈRE, 1789; OD]. Umbo almost covered with swollen prominence; fringe indistinct, of short thick tubules, formed like hole in disc. *U.Oligo.*(*Aquitan.*)-*Rec.*, IndoPac.-W.Pac.—FIG. F32,6. **P. (F.) novaezealandiae* BRUGUIÈRE, *Rec.*, Australasia; $\times 0.8$ (Reeve, 1860).

P. (Pseudobrechites) MAGNE, 1941 [*Aspergillum leognanum* HOENINGHAUS, 1827; OD]. Similar to *P. (Penicillus)*, but fringe tubules shorter, less distinct, and anterior disc without central slit. *U.Oligo.*(*Aquitan.*), Eu.—FIG. F32,5. **P. (P.) leognanus* (HOENINGHAUS); $\times 0.67$ (Peyrot, 1920).

P. (Warnea) GRAY, 1858 [*Aspergillum australe* CHENU, 1843; SD STOLICZKA, 1871]. Tube cylindrical, siphonal end with series of plaited ruffles; fringe distinct, of single series of thick simple tubules. *Phio.*-*Rec.*, Red Sea-Australasia-Japan.—FIG. F32,4. **P. (W.) australis* (CHENU), *Rec.*, Australasia; $\times 0.55$ (305).

Subclass UNCERTAIN

Order CONOCARDIOIDA Neumayr, 1891

[nom. correct NEWELL, 1965 (ex order Conocardiiden NEUMAYR, 1891)] [Materials for this order prepared by C. C. BRANSON, AURÈLE LAROCQUE, and N. D. NEWELL]

Characters of genus *Conocardium*. *M. Ord.-U.Perm.*, ?*U.Trias.*

Superfamily CONOCARDIACEA Miller, 1889

[nom. transl. NEWELL, 1965 (ex Conocardiidae MILLER, 1889)]

Characters of genus *Conocardium*. *M. Ord.-U.Perm.*, ?*U.Trias.*

Family CONOCARDIIDAE Miller, 1889

Characters of genus *Conocardium*. *M. Ord.-U.Perm.*, ?*U.Trias.*

Only *Conocardium* is recognized currently within this poorly understood group. Hence, the family, superfamily, and ordinal traits are identical with those of the genus.

Conocardium BRONN, 1835, p. 92 [**Cardium elongatum* SOWERBY, 1812 (= *Conchiolithus rostratus* MARTIN, 1809); M] [= *Pleurorhynchus* PHILLIPS, 1836, p. 210 (type, *Cardium hibernicum* SOWERBY, 1812; SD BRANSON, LAROCQUE & NEWELL, herein); *Lichas* STEININGER, 1837, p. 231 (type, *L. antiquus*; M); *Hippocardia* BROWN, 1843, p. 97 (type, *Cardium hibernicum* SOWERBY, 1812; M); *Rhipidocardium* FISCHER, 1887, p. 1036 (type, *Conocardium amygdala* BARRANDE, 1881; M)]. Equivalve, alate anteriorly, produced posteriorly in elongate, tubular rostrum; shell surface minutely ornamented; shell material thick, inlaid by vertically set partitions which form more or less submerged radial ribs appearing as external ribs in some species, or weathering in relief by exfoliation of smooth outer prismatic layer and terminating as alternating denticulations at ventral margin; 1 or more ridges or septa in each valve curve upward and inward along hinge to form conical, forward directed receptacles of unknown function; hinge ankylosed in mature individuals, edentulous; some species distinctly carinate behind beaks and at least in some species carinae serving as attachment for collar of prismatic texture (*Schleppe*, *éventail*, fringe, flange, *Kragen*, hood); ventral anterior margin gaping, denticulate. *M.Ord.*(*Chazyan.*)-*U.Perm.*(*Capitan.*), ?*U.Trias.*, cosmop.—FIG. G1,1. **C. rostratum* (MARTIN), L.Carb., Eng.; *1a-c*, hinge, ventral and LV views of holotype, $\times 1$ (Hind, 1900); *1d*, RV ext., $\times 1$ (Hind, 1900).—FIG. G1,2. *C. herculeum* (DEKONINCK), L.Carb., Belg.(Tournai); *2a,b*, LV ext., LV int., $\times 1$ (Hind, 1900); *2c*, hinge view of another individual, $\times 1$ (*a*, ligament furrow?; *b*, muscle scar? and septum; *x*, ligament furrow) (Hind, 1900).—FIG. G1,3. *C. ventricosum* (HALL), M.Dev., USA(Iowa); *3a,b*, restoration seen from ventral side; both approx. $\times 1$ (BRANSON, n).

BIVALVE GENERA UNCERTAINLY ASSIGNED TO FAMILIES

Families, and likewise genera tentatively assigned to them, are arranged alphabetically. All are considered to be nomenclaturally sound but some may be *nomina oblita*.




FIG. G1. Conocardiidae (p. N859).

ANTHRACOSIIDAE (p. N406)

Amnigeniella BETEKHTINA, 1966¹ (?1967) [*Anthraconauta kumsassiana* RAGOZIN, 1960; OD]. *U.Carb.-L.Perm.*, USSR (W.Sib.). [Not an anthracosiid, but may be near *Procopievskia* (nonmarine Myalinidae).] [WEIR]

ARCIDAE (p. N250)

Arcrites SCHLOEPPER, 1821.
Calliarca CONRAD, 1866.
Cucullea MICHELIN, 1838.
Cyphoxis RAFINESQUE, 1818.
Hataiarca NODA, 1966, p. 114. (as subgenus of *Anadara*) [*H. kakchataensis* HATAI & NISIYAMA, 1949; OD]. *Mio.-Rec.*, Japan.
Kikaiarca NODA, 1966, p. 127 [*Anadara (Kikaiarca) Kikaiizimana* NONURA & ZINBO, 1934; OD]. *Pleist. (Ryukyu Is.)*, Japan.
Pectinarca SACCO, 1898.
Polyodonta MEGERLE, 1811.
Scaphura GRAY, 1842.

Thyas GRAY, 1857.

Tosarca NODA, 1965, p. 120 [*Anadara (Tosarca) tosaensis* NODA, 1965; OD]. *Plio.*, Japan.

ARCTICIDAE (p. N645)

Ambonicardia WHITFIELD, 1885.
Bradicardia LORIOU, 1891.
Bruntrutia COSSMANN, 1902.
Kobya LORIOU, 1901.

AVICULOPECTINIDAE (p. N335)

Schizopecten KOBAYASHI, 1936.

CAPRINIDAE (p. N789)

Caprinus MONTFORT, 1810.

CARDIIDAE (p. N583)

Avardaria ANDRUSSOV, 1923.
Callicardia CONRAD, 1873.
Cardiarlus DUMERIL, 1806.
Cartissa HERMANNSEN, 1848.

CHAMIDAE (p. N518)

Camostraca DESHAYES, 1830.

CORBICULIDAE (p. N665)

Limnocyrena MARTINSON, 1961. *Mesoz.*

ETHERIIDAE (p. N466)

Caillaudiana BOURGUIGNAT, 1880.
Chambardiana BOURGUIGNAT, 1880.
Niloticiana BOURGUIGNAT, 1880.

GRAMMYSIIDAE (p. N819)

Goldfussia CASTELNAU, 1843.
Rhombocardia MEEK & WORTHEN, 1866.

HIPPURITIDAE (p. N799)

Campiloceratites FORTIS, 1778.

LIMOPSIDAE (p. N264)

Feliciella LAMY, 1934.

LUCINIDAE (p. N492)

Loripoderma POLI, 1795.

LYMNOCARDIIDAE (p. N590)

Suchumca SENINSKI, 1905.

MACTRIDAE (p. N595)

Cycladina BERTHOLD, 1827.
Villarita DUNKER, 1846.

MACTROMYIDAE (p. N511)

Ferrata ROEDER, 1882 (subj. syn. of *Unicardium*, according to Cox).

MARGARITIFERIDAE (p. N414)

Damaris SWAINSON, 1823.

MODIOMORPHACEA FAMILY UNCERTAIN
(p. N393)

Coxiconcha BABIN, 1966, p. 281 [**Lyonsia britannica* ROUAULT, 1851; OD]. *Tert.(Dandeil.)*, Eu. (France).

MODIMORPHIDAE (p. N393)

Taimyria LUTKEVICH, 1951 [**T. taimyrensis*; OD]. *L.Perm.*, USSR (W.Sib.). [WEIR]

MUTELIDAE (p. N463)

Berpolis LEACH, 1825.
Dentina MEGERLE, 1841.

MYALINIDAE (p. N289)

Hoplomytilus SANDBERGER, 1856.
Myalinoptera FRECH, 1891.

MYTILIDAE (p. N271)

Callitriche POLI, 1791.
Callitricoderma POLI, 1798.
Ciboticola IREDALE, 1939.
Dactylus MÖRCH, 1861.
Pisima MÖRCH, 1834.
Pseudomodiolus BETEKHTINA, 1966 (?1967) [**?* (First included species, *Anthraconauta vulgaris* KHALFIN; first fully described species, *P. ella* BETEKHTINA, 1967)]. [WEIR]
Septiola BITTNER, 1895.
Sinomytilus THIELE, 1934.
Stenolema DALL, BARTSCH & REHDER, 1939.

PARALLELODONTIDAE (p. N256)

Areocuculla COSSMANN, 1923.

PECTINACEA FAMILY UNCERTAIN (p. N332)

Protopalliolium SADYKOV, 1962, p. 66 [**P. kasanli*; OD]. *Dev.*, USSR (Kazakhstan).

PTERIIDAE (p. N302)

Unionum DALL, 1890 (new name for *Unionium* LINK, 1807).

PTERINEIDAE (p. N298)

Stainieria MAILLIEUX, 1930.

RADIOLITIDAE (p. N803)

Enargetes FISCHER DE WALDHEIM, 1830.

REQUIENIIDAE (p. N779)

Arietina CONRAD, 1853.

TEREDINIDAE (p. N722)

Teredarius DUMERIL, 1806.

TRIDACNIDAE (p. N594)

Tridacnoides KRUEGER, 1823.

UNIONIDAE (p. N415)

Abakaniella BETEKHTINA, 1966 (?1967) [**Anthracomya magna* CHERNYSHEV; OD]. *M.Carb.*, USSR (W.Sib.). [WEIR]
Brussiella BETEKHTINA, 1966 (?1967) [**B. curta*; OD]. *U.Perm.*, USSR (Sib.). [WEIR]
Sinomya POGOREVICH, 1951 [**S. kriegeri*; OD]. USSR (W.Sib.). [WEIR]

UNIONACEA FAMILY UNCERTAIN (p. N411)

- Abbreviatiana BOURGUIGNAT, 1881.
Acalliana BOURGUIGNAT, 1881.
Adamiana BOURGUIGNAT, 1881.
Aegericiana LOCARD, 1889.
Alpecanusiana LOCARD, 1889.
Amnicusiana LOCARD, 1889.
Anatiniana BOURGUIGNAT, 1881.
Arealiana BOURGUIGNAT, 1881.
Arnouldiana LOCARD, 1890.
Arrosiana LOCARD, 1890.
Asticusiana LOCARD, 1889.
Ateriana LOCARD, 1889.
Avoniana BOURGUIGNAT, 1881.
Aximedia RAFINESQUE, 1820.
Baryana LOCARD, 1889.
Batavusiana LOCARD, 1889.
Beccariana LOCARD, 1890.
Berenguieriana LOCARD, 1889.
Brebissoniana LOCARD, 1889.
Briandiana BOURGUIGNAT, 1881.
Brotiana BOURGUIGNAT, 1881.
Camuriana LOCARD, 1890.
Carvalhoiana LOCARD, 1890.
Coelatura CONRAD, 1853.
Collobiana LOCARD, 1890.
Complanatiana LOCARD, 1890.
Courquiniana LOCARD, 1889.
Crassiana LOCARD, 1889.
Cygnaeana BOURGUIGNAT, 1881.
Cyrtusiana LOCARD, 1889.
Depressiana BOURGUIGNAT, 1881.
Despontainiana LOCARD, 1889.
Diplasma RAFINESQUE, 1831.
Ellipsaria RAFINESQUE, 1820.
Ellipsopsiana BOURGUIGNAT, 1881.
Ellipsopsiana LOCARD, 1889.
Elongatiana LOCARD, 1890.
Embiana LOCARD, 1890.
Eydyusiana LOCARD, 1889.
Falsusiana LOCARD, 1889.
Fusculusiana LOCARD, 1889.
Gallandriana BOURGUIGNAT, 1881.
Gallicusiana LOCARD, 1889.
Gastrodiana BOURGUIGNAT, 1881.
Gestrouana LOCARD, 1890.
Glyciana BOURGUIGNAT, 1881.
Granigeriana LOCARD, 1889.
Gravidiana LOCARD, 1890.
Heckingiana LOCARD, 1889.
Hemisolasma RAFINESQUE, 1831.
Hispaniana LOCARD, 1889.
Holandriana LOCARD, 1889.
Humbertiana LOCARD, 1890.
Idriniana LOCARD, 1890.
Illuviosiana FAGOT, 1885.
Incrassatiana LOCARD, 1890.
Jacqueminiana LOCARD, 1889.
Joannisiana LOCARD, 1889.
Jourdheuiliana BOURGUIGNAT, 1881.
Jousseauwana LOCARD, 1889.
Lacanniana LOCARD, 1880.
Lapidus SIMPSON, 1900.
Lemotheuxiana LOCARD, 1889.
Letourneuxiana LOCARD, 1890.
Leucosilla RAFINESQUE, 1831.
Limnoica GRAY, 1857.
Locardiana LOCARD, 1889.
Lusitaniana BOURGUIGNAT, 1881.
Machadoiana LOCARD, 1890.
Macilentiana BOURGUIGNAT, 1881.
Macrosteniana LOCARD, 1890.
Mancusiana LOCARD, 1889.
Mariana LOCARD, 1889.
Mariolana LOCARD, 1890.
Melasiana LOCARD, 1889.
Meretricisiana LOCARD, 1889.
Meretrixiana LOCARD, 1890.
Metaptera RAFINESQUE, 1820.
Milletiana BOURGUIGNAT, 1881.
Moquiniana LOCARD, 1889.
Moreletiana LOCARD, 1889.
Mucidusiana LOCARD, 1889.
Nanusiana LOCARD, 1889.
Nemrodia PALLARY, 1939.
Notopala COTTON, 1935.
Notopalena IREDALE, 1943.
Nubilusiana LOCARD, 1889.
Ogerieniana LOCARD, 1890.
Ovuliana LOCARD, 1890.
Pammegaliana BOURGUIGNAT, 1881.
Parreyssiana BOURGUIGNAT, 1884.
Penchinatiana LOCARD, 1889.
Perniana LOCARD, 1889.
Picardiana BOURGUIGNAT, 1881.
Pisaniana LOCARD, 1889.
Piscimaliana BOURGUIGNAT, 1881.
Platteniciana BOURGUIGNAT, 1881.
Platyrrhynchoidiana LOCARD, 1889.
Ponderosiana BOURGUIGNAT, 1881.
Pseudocaelatura GERMAIN, 1921.
Pseudoglyciana LOCARD, 1890.
Rayana LOCARD, 1890.
Requieniana LOCARD, 1889.
Rhombeidiana LOCARD, 1889.
Rockanaia DE MORRETES, 1941.
Rossmassleriana BOURGUIGNAT, 1881.
Rossmassleriana LOCARD, 1890.
Rostratiana BOURGUIGNAT, 1881.
Rostratiana LOCARD, 1889.
Rumanicana LOCARD, 1890.
Sandriana LOCARD, 1889.
Schizostoma SCHLUETER, 1838.
Scrupeana LOCARD, 1890.
Securilla DROUET, 1855.
Simonisiana LOCARD, 1889.
Socardiana LOCARD, 1889.
Spengleriana BOURGUIGNAT, 1881.
Sperchinusiana LOCARD, 1889.

Spondaeana LOCARD, 1890.
 Sturmiiana LOCARD, 1890.
 Tricassiniana LOCARD, 1890.
 Tumidusiana LOCARD, 1889.
 Turtoniana LOCARD, 1889.
 Unioniformiana LOCARD, 1890.
 Ventricosiana BOURGUIGNAT, 1881.
 Vietuliana LOCARD, 1890.
 Viliacana LOCARD, 1889.
 Villanana FUGOT, 1892.
 Westerlundiana BOURGUIGNAT, 1881.

VENERIDAE (p. N670)

Radiocrista DALL, 1901 [**Venus pulcherrima* DESHAYES, 1860; OD]. DALL assigned this genus to Veneracea, but the age and locality of the type was unknown at the time of description. It could be a *Protocardia* or a *Trigonia*. [KEEN]

CLASS, ORDER, AND FAMILY UNCERTAIN

- Allocardium HALL, 1883.
 Anomianella DE RYCKHOLT, 1847, p. 45 [**A. proteus*; M]. *L.Carb.*(*Tournais.*), Eu.(Belg.). [NEWELL]
 Arnoldia MAYER, 1887 [**A. consecta*; M].
 Aviculomya HOLZAPFEL, 1889 [**A. peralata*; OD]. Shell thin, much swollen; beaks pointed, spirally enrolled, small; anterior wing rather large, continuing as ridge on dorsal margin behind beaks; posterior wing very large, not distinctly separated from main body of shell. Dentition and pallial line unknown. Ligament probably external, lying between shell surface and ridgelike prolongation of anterior wing. *L.Carb.*, Eu.(Ger.).—FIG. H1, 10. **A. peralata*, Liebstein (near Herborn); RV ext., $\times 1$ (Holzapfel, 1889). [LAROQUE]
 Barcoona FINLAY, 1926, p. 526 [*pro Pachydomella* ETHERIDGE, JR., 1907, p. 325 (*non* ULRICH, 1891)] [**Pachydomella chutus* ETHERIDGE; M]. Small (length 8 mm. or less), subtriangular, subequilateral, longer than high, equivalve, compressed, relatively thick-shelled; umbones obtusely angular, scarcely incurved; hinge supposedly edentulous; muscle scars and pallial line unknown; surface unornamented except for well-marked growth rugae. *L.Cret.*, Australia(Queensl.).—FIG. H1, 2. **B. chutus* (ETHERIDGE); LV, $\times 3$ (Etheridge, 1907). [COX]
 Binghami BROWN, 1827 [**B. paradoxus*; M].
 Bitubulites BLUMENBACH, 1803.
 Blainvilla CHIAJE, 1829.
 Bleta DE GREGORIO, 1930, p. 17 [**B. elegans*; M]. Genus founded on strongly convex, acutely triangular, subequilateral valve about 25 mm. high with strong growth rings; internal characters unknown. *U.Trias.*, Sicily. [COX]
 Carnidia BITTNER, 1901, p. 6 [**C. panonica*; SD DIENER, 1923, p. 223]. Small smooth, very gibbose, trigonally subquadrate, only slightly inequilateral, with well-marked posterodorsal angle and diagonal ridge delimiting narrow posterior area; posterior margins with broad gape; internal characters unknown. *U.Trias.*, Eu.(Hung.-Austria).—FIG. H1, 3. **C. panonica*, Veszprém, Hung.; 3a, RV lat.; 3b, dorsal view; 3c, posterior end showing gape, all $\times 3$ (Bittner, 1901). [COX]
 Ceratoconcha KRAMBERGER-GORJAHOVIC, 1889.
 Conchites SCHLAEPFER, 1821.
 Costulopteria PAUL, 1941.
 Cruciella KOKEN, 1913, p. 35 [**Nucula inflata* WISSMANN in MÜNSTER, 1841, p. 20 (*non* J. DE C. SOWERBY, 1827) (= *Megalodon klipsteini* BITTNER, 1895, p. 22; OD)]. Small to medium-sized, gibbose, ovately cuneiform, inequilateral, slightly inequivalve; umbones at anterior 0.25 or 0.3 of length, that of LV higher than RV; lunule wide, cordiform; escutcheon deep, with well-defined bordering ridges; weak diagonal ridge commonly present; ligament apparently external, opisthodontic; hinge margin thickened below beak in RV, forming toothlike process received in recess of LV, hinge otherwise edentulous; pallial line entire; shell wall rather thin; exterior smooth or with weak concentric folds. *U.Trias.*(*Carn.*), Austria(S.Tyrol).—FIG. H1, 1. **C. klipsteini* (BITTNER), Heiligenkreuz; 1a, LV lat.; 1b, ant. view showing greater prominence of LV umbo, both $\times 2$ (Bittner, 1895). [COX]
 Cyclostrea GENTIL, 1902.
 Cyrenolimopsis HABE, 1953.
 Delia DE LORIO, 1891, p. 246 [**D. amaena*; M]. Medium-sized, equivalve, moderately and evenly inflated, suboval, slightly longer than high; umbones moderately prominent, just posterior to mid-length; lunule present but no bordered escutcheon; ligament external, opisthodontic; internal characters unknown; surface smooth except for growth lines. *U.Jur.*(*Oxford.*), Eu.(Switz.).—FIG. H1, 12. **D. amaena*; 12a, RV lat.; 12b, dorsal view, both $\times 1$ (de Loriol, 1891). [COX]
 Diptyxis OPPENHEIM, 1889.
 Donacites GMELIN, 1793.
 Dorsomya DE RYCKHOLT, 1847, pl. 10, fig. 20 [**D. dorsata*; M]. [NEWELL]
 Ectenoptera ULRICH, 1894, p. 485 (*nom. nud.*). [NEWELL]
 Elasmatium CLARKE, 1904, p. 293 [**E. gowan-dense*; OD]. *U.Dev.*(*Naples*), USA(N.Y.). [NEWELL]
 Elliptoidea TASCH, 1961 [**E. vulgaris*; OD]. Similar to stunted *Palaeonodonia*. *Perm.*(*Leonard.*), USA(Kans.). [WEIR]
 Ensia DE GREGORIO, 1930, p. 21 [**Pleuromya?* (*Ensia*) *pumila*; M]. Genus founded on small, sublunate, smooth shell with moderately prominent, almost terminal umbo; internal characters unknown. *U.Trias.*, Eu.(Sicily). [COX]
 Gibboconcha DE GREGORIO, 1930, p. 30 [**G. sicula*;

Bivalvia





FIG. H1. Class, Order and Family Uncertain (p. N864).

- M]. Genus founded on fragment of small, concentrically ribbed, subtrigonal, strongly inequilateral shell; internal characters unknown. Possibly young *Cardinia* or *Astarte*. *L.Jur.(Lias.)*, Sicily. [Cox]
- Gingillum** DE GREGORIO, 1930, p. 27 [**G. lindum*; M]. Medium-sized, smooth, obliquely oval, higher than long, with narrow submedian umbo and short, arcuate dorsal margin; internal characters unknown. *L.Jur.(Lias.)*, Italy (Sicily). [Cox]
- Guebhardia** COSSMANN, 1905, p. 841 [**G. veneriformis*; M]. Small, oblong-oval, subequilateral, moderately inflated, with broad, slightly protruding umbo and posterior umbonal ridge bordering narrow, radially costulate posterior area; surface otherwise smooth; anterior end with shallow radial sulcus forming lower border of slightly swollen anterodorsal region; ligament external, opisthodontic; 1 strong cardinal tooth in each valve, in LV posterior to that of RV, which occupies median position below beak; 1 short, thick posterior lateral in each valve; adductor scars & pallial line unknown. *M.Jur.(Bathon.)*, Eu.(S. France).—FIG. H1,4. **G. veneriformis*, Courmes, Alpes-Maritimes; *4a,b*, LV lat., ext., int., both $\times 1.3$ (Cossmann, 1905). [Cox]
- Hadrodon** YEN, 1952.
- Ischyrinia** BILLINGS, 1866, p. 52 [**I. winchelli*; SD S. A. MILLER, 1889] [= ?*Technophorus* MILLER, 1889, p. 541 (type, *T. faberi*; M)]. Equivalve, inequilateral; 2 strong ridges radiating from beak to extended posteroventral extremity; hinge and musculature unknown. *M.Ord.-U.Ord.*, E.N.Am.—FIG. H1,5. **I. winchelli*, U.Ord., Can. (Anticosti I.); RV ext., $\times 1$ (Billings, 1866). [NEWELL]
- Laevipinna** PAUL, 1941.
- Laubeia** BITTNER, 1895, p. 116 [**Cardita strigilata* KLIPSTEIN, 1845, p. 255; M]. Small, slightly inequivalve, LV larger than RV; oval, strongly inequilateral; umbones prosogyrate, presence of diagonal ridge variable, deep escutcheon bordered by sharp ridge present in each valve; LV with 2 teeth, more dorsal of which is bifid, consisting of short posterior limb directed radially from beak and larger, longitudinally directed anterior limb, 2nd tooth below and parallel to this anterior limb; RV with single bifid tooth consisting of short radial posterior limb and longer longitudinal anterior limb; lateral teeth absent; pallial line and adductor scars unknown; surface smooth. *U.Trias.(Carn.)*, Austria (S. Tyrol).—FIG. H1,7. **L. strigilata* (KLIPSTEIN), St. Cassian; *7a*, LV ext.; *7b*, dorsal view; *7c*, hinge teeth of LV; *7d*, hinge teeth of RV; all $\times 1.3$ (Bittner, 1895). [Cox]
- Libyaconchus** HASSAN, 1957, p. 135 [**L. parvus*; OD]. Small (< 5 mm. long), suborbicular, gibbose, slightly inequivalve; umbones broadly rounded, that of RV more prominent; beaks prosogyrous; RV with strong cardinal tooth, recurved in socket of LV; pallial line obscure; pedal retractor scars conspicuous, paired, on posterodorsal side of anterior adductor; ornament of strong radial ribs, some specimens also with radial sulcus posterior to mid-length in each valve, sulcus of RV with median rib. *U.Cret. or Paleoc.(Dan.)*, Esna Shales, Egypt.—FIG. H1,6. **L. parvus*; *6a*, RV, int. mold; *6b*, dorsal view of bivalve int. mold, showing pedal retractor scars; both $\times 5$ (Hassan, 1957). [Cox]
- Liocardia** AGASSIZ, 1842, p. 26 [Virtual *nom. nud.* without included nominal species.] Said to be close to "*Ceromya*" [*Ceratomya*] in external characters, but closer to *Isocardia* in other respects [Cox]
- Macrodonia** COSTA, 1845.
- Megagyrypha** SPREISTERSBACH, 1922, p. 419 [**M. dahmeri*; OD]. *L.Dev.*, Eu.(Ger.). [LA ROCQUE]
- Mellaca** MABILE, 1899.
- Modioptera** SCHINDEWOLF, 1924, p. 275 (*nom. nud.*). [NEWELL]
- Nothamusium** HIND, 1904, p. 159 [**N. radiatum*; SD NEWELL, herein]. Equivalve, very inequilateral, obliquely ovate, prosogyre with beaks about 0.3 distance behind anterior extremity of hinge; posteroanterior margin subquadrate; free border ornamented by bifurcating flattened costae; hinge unknown. *L.Cam.*, Eu.(Ire.-Eng.-Scot.).—FIG. H1,8. **N. radiatum*, Eng.(Staffords.); RV ext., $\times 1$ (Hind, 1904). [NEWELL]
- Omalia** DE RYCKHOLT, 1847 (*nom. oblit.*), *L.Carb.*, Eu.(Belg.). [NEWELL]
- Onkogrypha** SPRIESTERSBACH, 1925, p. 418 [**Mego-grypha (Onkogrypha) cullmanni*; M]. *L.Dev.*, Eu.(Ger.). [NEWELL]
- Opisenia** ETALLON, 1862.
- Osteophorus** PHILIPPI, 1893.
- Ostreavicula** BLANCKENHORN, 1934, p. 179 [**O. dayi*; M]. Large, quadrate, oysterlike, LV feebly convex, RV slightly concave; no clearly demarcated posterior wing; RV with anterior auricle bordered posteriorly by narrow incision running from beak toward anterior margin; hinge structure and musculature unknown; surface bearing narrow dichotomous ribs. *U.Cret.(Turon.)*, Syria. [Cox]
- Paradoxia** KRUMBECK, 1923, p. 101 [**P. timorensis*; M]. Known only by single valve, possibly LV; with attachment area at tip; large, very thick-shelled, oval, higher than long, with large, down curved, hooklike ?posterior wing with median external crest; broad, dorsoventrally elongated cardinal area, without median depression, occupying posterior half of inner side and curved in posterior direction of dorsal end; visceral cavity shallow, single adductor scar occupying position dorsal to median; surface with uncertain rugae and a few wartlike protuberances. *L.Jur.(mid. Lias., Pliensbach.)*, E.Indies (Timor).—FIG. H1, 11. **P. timorensis*; *11a,b*, ?LV ext., int., $\times 0.7$ (Krumbeck, 1923). [Cox]
- Patogocardia** DOELLO-JURADO, 1946.
- Planktomya** SIMROTH, 1896 [**P. henseni*; OD]. *Rec.*, Atl.






FIG. H2. Class, Order and Family Uncertain (p. N866).

Plicatostylus LUPHER & PACKARD, 1930, p. 204 [**P. gregarius*; OD]. Known mainly or solely from attached valve (AV) (= ?LV, ?RV) which is large, greatly elongated, acutely conical or columnar, straight or slightly curved, attached at apex; cross section almost semicircular, one side with broad, flat, or slightly concave area (interpreted as ligamental area) extending whole length of AV between 2 riblike projections and bearing numerous low, flat-topped longitudinal ridges varying in width and separated by narrow intervals; convex (probably ventral) side, bearing coarse, upward-arched undulations at distant intervals and growth lines curved in conformity with them, indicating that commissure of valves was very oblique, sloping upward from hinge margin, which presumably was located at upper end of ligamental area; body chamber greatly elongated, subelliptical in cross section, with one side close to ligamental area, other side separated from outer surface of AV by thick wall, which has

compact outer layer formed of rather coarse calcite prisms normal to surface, and thick, probably originally vesicular, inner layer of uncertain structure; hinge edentulous, its margin thin; 2 adductor scars present, one located on thick projecting buttress in upper part of body chamber on one side, other limited by low ridge; free valve (FV) imperfectly known or unknown, probably a flat, relatively thin structure. [This colonial reef-building bivalve, with largest illustrated specimen 48 cm. (19 in.) in height, was assigned by LUPHER & PACKARD (1930) to their monotypic Plicatostylidae. By other authors it has been referred to the Caprinidae and with small doubt seems to be a rudist.] *L. jur.* (low. Pleiensch.), N.Am. (Ore.)-S.Am. (Peru).—FIG. H2, 4. **P. gregarius*, Robertson Formation, USA (Ore.); 4a, presumed dorsal side of AV showing ?ligamental area, $\times 0.4$; 4b, opposite (?ventral) side of same specimen showing uparched undulations. $\times 0.4$; 4c, assemblage of AVs indicating colonial mode of life, $\times 0.3$ (Cox, n). [Cox]

- [In the original description of the genus it was stated that the upper valve was curved and cap-shaped, with a prominent umbo and broad ligamental area like that of the lower valve. Dr. R. L. LUPHER has informed me that the specimens originally thought to be upper valves probably are small deformed lower valves and that the upper valve (not yet found complete) seems to have been as here stated.]
- Pracardiomya** AWAD, 1952, p. 7 [**P. sinaitica*; M]. Rather small, elongate-ovate, beaks at anterior third of length, escutcheon lanceolate; posterior area defined by faint diagonal groove with corresponding internal ridge; integripalliate; LV with triangular chondrophore, RV with corresponding pit; 5 irregular transverse teeth in each valve, forming protuberances from hinge margin on posterior side of beaks. *U.Cret.(Cenoman.)*, Sinai. [COX]
- Preroderma** KURODA, 1945.
- Productea** TASCH, *nom. subst.* herein [*pro Productae* TASCH, 1961 (*nom. vet.*)] [**P. dunbari* TASCH, herein (*nom. subst. pro Productae dunbaris* TASCH, 1961, *nom. vet.*); OD]. *L.Perm.(Leonard.)*, USA (Kans.). [TASCH]
- Psammoconcha** TOMMASI, 1896, p. 61 [**P. servini*; M]. Medium-sized, elongate-oval, anterior margin more abruptly curved than posterior; subequilateral, with broad, scarcely projecting umbones; obscure diagonal ridge present in some specimens; internal characters unknown. *L.Trias.*, Eu.(S.Alps). [COX]
- Pseudomya** ROEDER, 1882, p. 103 [**P. rarissima*; OD]. Founded on single imperfect elongate, oval, inequilateral LV; muscle scars and pallial line unknown; prominent spoon-shaped chondrophore, extending upwards, and with groove behind it, situated just anterior to beak. *U.Jur.(Oxford.)*, Eu.(Alsace). [COX]
- Ptychodes** FISCHER DE WALDHEIM, 1848.
- Rebusum** DE GREGORIO, 1930, p. 27 [**R. drepanense*; M]. Small to medium-sized, smooth, obliquely oval, higher than long, subalate, compressed, recalling some Limidae in outline but apparently lacking cardinal area; internal characters unknown. *L.Jur.(Lias.)*, Eu.(Sicily). [COX].
- Rhenanomya** LA ROCQUE, *nom. subst.* herein [*pro Rhenania* FUCHS, 1915, p. 53 (*non* WAAGEN, 1907)] [**Rhenania tumida* FUCHS, 1915; OD]. Shell ventricose, elongate oval; umbones near anterior end, prosogyrous, strongly incurved; lunule small; hinge curved, prolonged posteriorly; LV with 3, RV with 2 to 3 cardinal teeth; lateral teeth lacking; ligament opisthodontic and external. *L.Dev.*, Eu.(Ger.).—FIG. H1,9. **R. tumida* (FUCHS), Coblenz.; 9a, LV ext. showing outline and position of beaks, $\times 0.7$; 9b, RV dentition, $\times 0.7$ (Fuchs, 1915). [LA ROCQUE]
- Rhinchotropis** MEEK, 1872.
- Rostrortotus** TASCH, 1961 [**R. dissimilis*; OD]. ?Esteriid or possibly pathologic bivalve. *Perm.(Leonard.)*, USA (Kans.). [NEWELL]
- Sanoarca** STEPHENSON, 1952.
- Sollea** DAHMER, 1949.
- Spinarcullaea** CHAVAN, 1952.
- Spondiolites** REICHENBACH, 1828.
- Stagnestesta** TASCH, 1961 [**S. solitaria*; OD]. ?Possibly estheriid. *Perm.(Leonard.)*, USA (Kans.). Stunted bivalves from deposits of temporary, variable saline pools. [NEWELL]
- Stalagmina** DENCKMANN, 1887, p. 89 [**S. koeneni*; M]. Small (8 mm.), obliquely suborbicular, inequilateral; umbones prosogyrate but scarcely protruding; small lunule present; dentition consisting in each valve of elongate posterior lateral and about 3 small teeth of taxodont appearance close to beak; surface smooth. *L.Jur.(up.Lias.-Toarc.)*, Eu.(Goslar, Ger.).—FIG. H2,3. **S. koeneni*; 3a, LV ext., $\times 1$; 3b, RV showing impression of teeth, $\times 1$ (Denckmann, 1887). [COX]
- Taeniodon** DUNKER, 1848, p. 179 [**T. ellipticus*; M]. Small (20 mm.) equivalve, oval, smooth, moderately inequilateral; LV with curved, upwardly concave, lamellar tooth just posterior to beak, with recess between it and margin for reception of thickened margin of RV; ligament external; muscle scars and pallial line unknown. *L.Jur.(Hettang.)*, Eu.(Ger.).—FIG. H2,2. **T. ellipticus*; 2a,b, LV ext., int., $\times 1$ (Dunker, 1848).
- Tatella** ETHERIDGE, 1901, p. 27 [**Corbicella? maranoana* ETHERIDGE, 1892, p. 471; OD]. Medium-sized, elongate, subrectangular or ovate, subequilateral, compressed, gaping slightly at both ends; RV with 2 short subumbonal teeth, LV with 1 blunt tooth recurved between them; no lateral teeth; anterior adductor scars elongate pyriform, oblique, posterior ones rounded; pallial line with small sinus; no ornament except concentric rugae. *L.Cret.*, Australia.—FIG. H2,1. **T. maranoana*, Maranoa River, Queensl.; 1a,b, LV, ext., hinge, $\times 1$ (Jack & Etheridge, 1892). [COX]
- Taxocardia** OLSSON & HARBISON, 1953.
- Verticpronus** HEDLEY, 1904 [**V. mytilus*; OD]. Although assigned to Philobryidae by some authors, it was provisionally assigned to Carditidae and is so listed by POWELL and other Australasian workers. [KEEN]
- Zoreia** BRUSINA, 1907 [**Z. sanguinea*; M]. ?Pleist., Yugosl. Unfigured, minute (2 mm. long); description follows that of a species of *Tapes* in Veneridae, but it is so vague that one cannot be sure the author intended it in that family. [KEEN]

GENERA OF DOUBTFULLY MOLLUSCAN AFFINITIES

(With cross references to *Treatise* Part R)

- Anatifopsis** BARRANDE, 1872 (*see p. R329*).
- Aptychopsis** WOODWARD, 1872 (*see p. R329*).
- Discinocaris** WOODWARD, 1866 (*see p. R329*).
- Eoischyrina** KOBAYASHI, 1933.
- Euchasmella** KOBAYASHI, 1933.
- Euclusaea** GRAY, 1852. Brachiopod?

Fordilla BARRANDE, 1881.
 Lebescontia JONES & WOODWARD, 1899 (see p. R326).
 Lingulocaris SALTER, 1866 (see p. R330).
 Modiolooides WALCOTT, 1889.
 Myocaris SALTER, 1864 (see p. R331).
 Ozomia WALCOTT, 1924.
 Paleodora FLEMING, 1957.
 Pinnocaris ETHERIDGE, 1878 (see p. R331).
 Pseudoeuchasma KOBAYASHI, 1933.
 Pseudotechnophorus KOBAYASHI, 1933.
 Ribeirella SCHUBERT & WAAGEN, 1903.
 Ribeiria SHARPE, 1853.
 Spirodomus BEECHER, 1886, p. 162 [**S. insignis*;
 OD, M]. *Miss. (Waverly)*, USA (Pa., Warren Co.).
 Probably not a mollusk. [NEWELL]
 Wanwania KOBAYASHI, 1933.
 Wanwanella KOBAYASHI, 1933.
 Wanwanoidea KOBAYASHI, 1933.

UNRECOGNIZABLE GENERA

Aegilops HALL, 1850, p. 179 [**A. subcarinata*;
 M]. *M. Ord. (Trenton)*, USA (N.Y.). [Possibly a
 cyrtodontid (p. N248).]
 Burriera LASERON, 1910 [**B. dunii*; OD]. *L. Perm.*
 (*U. Marine, Wandrawan.*), Australia (N.S. Wales).
 [NEWELL]
 Dolabra M'COY, 1844. *Carb.* [May belong to Myo-
 phoridae.]
 Hemisterias RAFINESQUE, 1832, p. 142 [**H. quad-*
riloba; M]. ?*Penn.*, Sherman Cr., Allegheny Mts.,
 USA (Pa.). [NEWELL]
 Mactrula RISSO, 1826.
 Pleureterites RAFINESQUE, 1832, p. 142 [**P. lateri-*
stria; SD NEWELL, herein]. ?*Penn.*, Sherman Cr.,
 Allegheny Mts., USA (Pa.). [NEWELL]

Preavicula WILLIAMS & BREGER, 1916, p. 205
 [**Megambonia oblonga* HALL, 1895; OD]. *L. Dev.*,
 USA (N.Y.). [NEWELL]
 Protopecten HIND, 1910 [**P. vimineus*, **P. crenu-*
latus]. Probably a brachiopod. *Sil. (Llandov.)*,
 Eu. (Eng.). [NEWELL]
 Sarka BARRANDE, 1881, p. 150 [**S. infelix*; M].
 Possibly a brachiopod. *M. Dev.*, Eu. (Boh.). [NEW-
 ELL]
 Telistrophis RAFINESQUE, 1832, p. 142 [**T. torsala*;
 OD, M]. ?*Penn.*, Sherman Cr., Allegheny Mts.,
 USA (Pa.). [NEWELL]

Family UNCERTAIN

Paramonopleura KOROBKOV in KOROBKOV & MAKAR-
 ENKO, 1967 [addition by B. F. PERKINS (see p.
 N783)]. Anterior muscle insertion of AV on shell
 wall, posterior insertion on prolongation of cardinal
 platform and separated from 3' by low ridge;
 muscle insertions of FV on thickened areas of shell
 wall. [Note.—If the conclusions of KOROBKOV &
 MAKARENKO are correct, this genus would be the
 only known Tertiary rudist. It was assigned to
 the Monopleuridae by KOROBKOV, but without data
 on larger, more mature shells, such placement is
 doubtful. The size of the type specimens and their
 occurrence in "glauconitic sandy-argillaceous de-
 posits with intercalations of shelly detritus" suggest
 that they may be juveniles reworked from older
 deposits.] [Ref.—KOROBKOV, I. A., & MAKARENKO,
 D. YE., 1967, *Paramonopleura gen. nov. (Rudis-*
tae) iz verkhnego Paleotsena Ukrainy: Paleont.
Zhurnal, no. 4, p. 135-138, 1 fig. [*Paramonopleura*
gen. nov. (Rudistae) from the upper Paleocene of
the Ukraine: Paleont. Jour., 1968 (July), no. 4,
 p. 109-112, 1 fig. (transl. Am. Geol. Inst.).]

REFERENCES

Abbott, R. T.

- (1) 1955, *American sea shells*: 541 p., 40 pl., D. Van Nostrand Co., Inc. (Princeton, N.J.).
- (2) 1958, *The marine mollusks of Grand Cayman Island, British West Indies*: Acad. Nat. Sci. Philadelphia, Mon., no. 11, 138 p.
- (3) 1962, *Sea shells of the world*: 160 p., 790 text fig., Golden Press (New York).

Adams, Arthur

- (4) 1860, *On some new genera and species of Mollusca from Japan*: Annals & Mag. Nat. History, ser. 3, v. 5, p. 299-303.
- (5) 1868, *Note sur quelques nouveaux genres de mollusques du Japon*: Jour. Conchyliologie, ser. 3, v. 8, v. 16, p. 40-56, pl. 4.

Adams, Henry

- (6) 1866, *List of land and freshwater shells collected by Mr. E. Bartlett on the upper Amazon and on the River Ucayali, eastern Peru, with descriptions of new species*: Zool. Soc. London, Proc., p. 440-445, pl. 38.

—, & Adams, Arthur

- (7) 1853-58, *The genera of recent Mollusca*: (a) v. 1(1853-54), 484 p.; (b) v. 2(1854-58), 661 p.; (c) v. 3(1858), 136 pl. (London).

Agassiz, Alexander

- (8) 1840, *Mémoire sur les moules des mollusques vivants et fossiles*: Soc. Sci. Nat. Neuchâtel, Mém., v. 2, no. 4, 48 p., 9 pl.

Agassiz, L. J. R.

- (9) 1841, *Etudes critiques sur les mollusques fossiles. Mémoire sur les Trigonies*: 58 p., 11 pl. (Neuchâtel).
- (10) 1852, *Über die Gattungen unter den Nord-Amerikanischen Najaden*: Arch. Naturgesch., v. 18, p. 41-52.

Allen, J. A.

- (11) 1963, *Ecology and functional morphology of molluscs*: Oceanogr. Mar. Biol., Ann. Rev., v. 1, p. 253-288.

Amalitsky, W. P.

- (12) 1891, *On the age of the Unionidae*: Soc. Nat. de Varsovie, Comptes Rendus (Sec. Biol.), no. 7, p. 1-5.
- (13) 1892, *Ueber die Anthrocosien der Permianformation Russlands*: Palaeontographica; v. 39, p. 125-214, pl. 19-23.
- (14) 1895, *A comparison of the Permian freshwater Lamellibranchiata from Russia with those from the Karroo System of S. Africa*: Geol. Soc. London, Quart. Jour., v. 51, p. 337-351, pl. 12-13.

Ancey, C. F.

- (15) 1887, *On the generic name of a remarkable bivalve shell found in the Congo*: Conch. Exch., v. 2, p. 22.

Anderson, F. W., & Cox, L. R.

- (16) 1940, *Notes in the molluscan fauna of the Great Estuarine Series of Skye*: Royal Phys. Soc. Edinburgh, Proc. v. 23, p. 105-122, 3 pl.

Andrussov, N. I.

- (17) 1910, *Studien über Brackwassercardiden: Didacna (Erste Hälfte)*: Acad. Impér. Sci. St. Pétersbourg, Mém., sér. 8, v. 25, no. 8, 84 p., 5 text fig., 10 pl.

Anthony, R. L. F.

- (18) 1907, *Étude monographique des Aetheriidae (anatomie, morphogénie, systématique)*: Soc. Royale Zool. Belgique, Ann. v. 41, p. 322-356, 20 text fig., pl. 10-11.

Arkell, W. J.

- (19) 1927(1929)-35(1936), *A monograph of British Corallian Lamellibranchia*: Palaeontograph. Soc. London, (a) pt. 1(1929), p. 1-72, pl. 1-4; (b) pt. 2(1930), p. 73-104, pl. 5-8; (c) pt. 3(1931), p. 105-132, pl. 9-12; (d) pt. 4(1932), p. 133-180, pl. 13-20; (e) pt. 5(1933), p. 181-228, pl. 21-28; (f) pt. 6(1934), p. 229-276, pl. 29-36; (g) pt. 7(1934), p. 277-324, pl. 37-44; (h) pt. 8(1935), p. 325-350, pl. 45-49; (i) pt. 9(1936), p. 315-376, pl. 50-56.

- (20) 1930, *The generic position and phylogeny of some Jurassic Arcidae*: Geol. Mag., v. 67, p. 297-319, 337-352, pl. 14-16.

- (21) 1934, *The oysters of the Fuller's Earth; and on the evolution and nomenclature of the Upper Jurassic Catinulas and Gryphaeas*: Cotteswold Nat. Field Club, Proc., v. 25, p. 21-68, pl. 1-6.

Astre, Gaston

- (22) 1931, *Existence d'Hippurites dans les terrains sénoniens de Madagascar*: Soc. Histoire Nat. Toulouse, Bull., v. 61, p. 269-275, pl. 15.
- (23) 1954, *Radiolitiidés nord-pyrénéens*: Soc. Géol. France, Mém., new ser., v. 33, no. 3-4, Mém. 71, p. 1-140, text fig. 1-34; pl. 1-8.
- (24) 1957, *Recherches sur l'organisation et la vie des radiolitiidés*: Same, Mém, new sér., v. 36, Mém. no. 80, 88 p., 19 text fig.

Atkins, Daphne

- (25) 1936-38, *On the ciliary mechanisms and interrelationships of lamellibranchs*: Micro. Sci., Quart. Jour., v. 79, p. 181-308(1936), 339-445(1937), pl. 10-11; v. 80(1938), p. 321-436, pl. 29.

Babin, Claude

- (25a) 1966, *Mollusques bivalves et céphalopodes du Paléozoïque armoricain*: 470 p., 18 pl., Imprimerie Commerciale et Administrative (Brest).

Baker, F. C.

- (26) 1928, *The fresh water Mollusca of Wisconsin. Pt. II, Pelecypoda*: Wisconsin Geol. Nat. History Survey, Bull. 70, pt. 2, p. 1-495, text fig. 203-299, pl. 29-105.

Barrande, Joachim

- (27a) 1881a, *Système silurien du centre de la Bohême, première partie, recherches paléontologique, volume 6, classe des mollusques, ordre des acéphalés*: xxiv+342 p., 48 pl. (quarto edit.) published by the author and editor (Prague and Paris).
- (27b) 1881b, *Acéphalés. Études locales et comparatives. Extraits du Système silurien du centre de la Bohême*, v. 6, xxxii+536 p., 10 pl. (irregular numbers) (octavo edit. reprinted from quarto edit. with different pagination), published by the author and editor (Prague and Paris).

Barrois, C. E.

- (28) 1891, *Mémoire sur la faune du grès armoricain*: Soc. Géol. Nord, Ann., v. 19, p. 134-237, pl. 1-5.

Bartsch, Paul

- (29) 1922, *A monograph of the American shipworms*: U.S. Natl. Museum, Bull. 122, 51 p., 37 pl.
- (30) 1927, *The shipworms of the Philippine Islands*: Same, Bull. 100, v. 2, pt. 5, p. 533-554, pl. 53-60.

Basse, Éliane

- (31) 1933, *Paléontologie de Madagascar. XVIII. Faune malacologique du Crétacé supérieur du Sud-ouest de Madagascar*: Ann. Paléontologie, v. 21, pt. 3-4, p. 1(89)-80(168), pl. 1-9.

Bassler, R. S.

- (32) 1915, *Bibliographic index of American Ordovician and Silurian fossils*: Smithsonian Inst., U.S. Natl. Museum, Bull. 92, v. 1-2, 1521 p., 4 pl.
- (33) 1919, *The Cambrian and Ordovician deposits of Maryland*: Maryland Geol. Survey, Cambrian & Ordovician vol., 424 p., 58 pl.

Bayle, E.

- (33a) 1855, *Notice sur la structure de coquilles du genre Hippurites*: Soc. Helvétique Sci. Nat., Actes (1855), p. 177-190.

Bédé, Paul

- (34) 1932, *Le genre Anodonta au Maroc*: Soc. Sci. Nat. Maroc, Bull., v. 12, p. 223-227, 3 text fig.

Bellardi, Luigi

- (35) 1875, *Monografia delle nuculidi trovate finora nei terreni terziari del Piemonte e della Liguria*: Il R. Liceo Gioberti nell' Anno scolastico 1874-75, 32 p., 1 pl. (Torino).

Bennison, G. M.

- (36) 1962, *Palaeontological and physical evi-*

dence of the palaeoecology of some early species of non-marine lamellibranchs: Liverpool & Manchester Geol. Jour., v. 3, pt. 1, p. 41-50.

Bernard, Félix

- (37) 1895, *Première note sur le développement et la morphologie de la coquille chez les lamellibranchs*: Soc. Géol. France, Bull., v. 23, pt. 3, p. 104-154.
- (38) 1896, *Deuxième note sur le développement et la morphologie de la coquille chez les lamellibranchs (taxodontes)*: Same, v. 24, pt. 3, p. 54-82.
- (39) 1896, *Troisième note sur le développement et la morphologie de la coquille chez les lamellibranchs (anisomyaires)*: Same, Bull., v. 24, pt. 3, p. 412-449.
- (40) 1897, *Quatrième et dernière note sur le développement et la morphologie de la coquille chez les lamellibranchs*: Same, Bull., v. 25, pt. 3, p. 559-566.
- (41) 1897, *Études comparatives sur la coquille des lamellibranchs. Condylocardia*: Jour. Conchyliologie, v. 44, no. 3, p. 149-207, pl. 6.
- (42) 1897, *Études comparatives sur la coquille des lamellibranchs. II. Les genres Philobrya et Hochstetteria*: Same, v. 45, p. 6-47, pl. 1.
- (43) 1897, *Anatomie de Chlamydoconcha orcutti Dall, lamellibranche à coquille interne*: Ann. Sci. Nat. sér. 8, v. 4, p. 221-252, pl. 12.
- (44) 1898, *Recherches ontogéniques et morphologiques sur la coquille des lamellibranchs*: Same (Zool.), v. 8, p. 1-208, pl. 1-12.

Betekhtina, O. A.

- (44a) 1966 (?1967), *Verkhne-paleozoiskie Nemorskije Peletsipody Sibiri i vostochnovo Kazakhstana*: Nauka, 220 p., 45 text fig., 20 pl., 2 tables. [Upper Palaeozoic non-marine pelecypods of Siberia and Eastern Kazakhstan.]

Beurlen, Karl

- (45) 1944, *Beiträge zur Stammesgeschichte der Muscheln*: Bayer. Akad. Wiss., Sitzungsber., no. 1-2, p. 133-145.

Beushausen, H. E. L.

- (46) 1884, *Beiträge zur Kenntniss des Oberharzer Spiriferensandsteins und seiner Fauna*: Geol. Spezialk. Preuss. Thüring. Staat., Abhandl., v. 6, no. 1, 133 p., atlas 6 pl.
- (47) 1895, *Die Lamellibranchiaten des rheinischen Devon mit Ausschluss der Aviculiden*: K. Preuss. Geol. Landesanst., Abhandl., new ser., v. 17, p. 1-514, pl. 1-38.

Beyrich, H. E.

- (48) 1861, *Ueber das Vorkommen von Posidonien in baltischen Juragesteinen*: Deutsch. Geol. Gesell., Zeitschr., v. 13, p. 143-145.

Bigot, A. P. D.

- (49) 1889, *Note sur quelques pélecypodes des*

- grès siluriens de l'Ouest de la France: Soc. Géol. France, Bull., sér. 3, v. 17, p. 791-801, pl. 22-23.
- (50) 1895, *Contributions à l'étude de la faune jurassique de Normandie. 2^e Mémoire. Sur les Opis*: Soc. Linnéenne Normandie, Mém., v. 18, p. 153-191, 2 pl.
- Billings, Elkanah**
- (51) 1858, *New genera and species of fossils from the Silurian and Devonian formations of Canada*: Can. Nat. & Geol., v. 3, p. 419-444, 24 text fig.
- (52) 1865, *Palaeozoic fossils*: Geol. Survey Canada, v. 1, 426 p., 399 text fig.
- (53) 1866, *Catalogue of the Silurian fossils of Anticosti*: Same, Publ. no. 427, 93 p., 28 text fig.
- (54) 1874, *Palaeozoic fossils*: Same, v. 2, pt. 1, 144 p., 85 text fig., 9 pl.
- (55) 1874, *On some new genera and species of Palaeozoic Mollusca*: Can. Naturalist, new ser., v. 7, p. 301-302.
- Bittner, Alexander**
- (56) 1891, *Triaspetrefakten aus Balia in Kleinasien*: K. K. Geol. Reichsanst. Wien, Jahrb., v. 41, p. 97-116, pl. 1-3.
- (57) 1894, *Einige Bemerkungen zu A. Rothpletz's "Ein geologischer Querschnitt durch die Ostalpen"*: Same, Verhandl., v. 28, no. 3, p. 87-102.
- (58) 1895, *Lamellibranchiaten der Alpenen Trias. I Theil: Revision der Lamellibranchiaten von St. Cassian*: Same, Abhandl., v. 18, no. 1, 235 p., 24 pl.
- (59) 1900, *Ueber nachtriadische Verwandte der Gattung Mysidioptra*: Same, Verhandl. 1900, p. 207-208.
- (60) 1901, *Lamellibranchiaten aus der Trias des Bakonyer Waldes*: Wiss. Erforsch. Balatonsees, Result., v. 1, pt. 1, Anhang, Palaeont., v. 2, pt. 3, 107 p., 9 pl.
- (61) 1901, *Ueber Pseudomonotis telleri und verwandte Arten der unteren Trias*: Same, Jahrb., v. 1, p. 559-592, pl. 22-23.
- Blainville, H. M. D. de**
- (62) 1814, *Mémoire sur la classification méthodique des animaux mollusques, et établissement d'une nouvelle considération pour y parvenir*: Soc. Philomath. Paris, Bull., p. 175-180.
- Böhm, Georg**
- (63) 1903, *Eurydesma und Leiomyalina*: Centralblatt Mineralogie, Geologie u. Paläontologie, p. 296-300.
- Böhm, Johannes**
- (64) 1904, *Ueber die obertriadische Fauna der Baereninsel*: K. Svenska Vetenskapsakad., Handl., v. 37, no. 3, 76 p., 71 pl.
- (65) 1914, *Zur Gattung Pleurophorus King und Myoconcha Sowerby*: K. Preuss. Geol. Landesanst., Jahrb., v. 35, p. 549-561.
- Boettger, C. R.**
- (66) 1961, *Zur Systematik der in die Gattung Pisidium C. Pfeiffer gerechneten Muscheln*: Archiv Molluskenkunde, v. 90, no. 4/6, p. 227-248.
- Boissevain, H., & MacGillivray, H. J.**
- (67) 1932, *Some remarks on Barrettia sparcilirata Whitfield and Chiapasella radiolitifformis (Trechmann)*: K. Nederland. Akad. Wetensch., Proc., v. 35, p. 1303-1312, 6 text fig.
- Bonetto, A. A.**
- (68) 1951, *Acerca de las formas larvales de Mutelidae Ortman*: Jornadas Icticas, Santa Fé, Argentine, 8 p., 5 text fig.
- Borissiak, A. A.**
- (69) 1904-09, *Die Pelecypoden der Jura-Ablagerungen im europäischen Russland*: Comité Geol. St. Petersburg, Mém., new ser., (a) I. Nuculidae, pt. 11(1904), 49 p., 3 pl.; (b) II. Arcidae, pt. 19(1905), 63 p., 4 pl.; (c) III. Mytilidae, pt. 29(1906), 35 p., 2 pl.; (d) IV. Aviculidae, pt. 44(1909), 26 p., 2 pl.
- , & **Ivanov, E.**
- (70) 1917, *Les pélecypodes des couches jurassiques de la Russie d'Europe. V. Pectinidae*: Comité Géol. Pétrograd, Mém., 143, 59 p., 4 pl.
- Boss, K. J.**
- (71) 1965, *A new mollusk (Bivalvia, Erycinidae) commensal on the stomatopod crustacean Lysiosquilla*: Am. Museum Nat. History, Novitates, no. 2215, 11 p., 3 text fig.
- Bourguignat, J. R.**
- (72) 1877, *Descriptions de deux nouveaux genres algériens suivies d'une classification des familles et des genres de mollusques terrestres et fluviatiles du système européen*: Soc. Phys. Natur. Toulouse, Bull., v. 3, 1875-76, p. 49-101.
- (73) 1879, *Description de diverses espèces terrestres et fluviatiles et de différents genres de mollusques de l'Egypte, de l'Abyssinie, de Zanzibar, du Sénégal et du centre de l'Afrique*: 54 p. (Paris).
- (74) 1885, *Monographie d'un nouveau genre d'Acéphales du lac Tanganika*: Soc. Mal. France, Bull., v. 2, p. 1-12, pl. 1.
- (75) 1885, *Espèces nouvelles et genres nouveaux découverts par les Rév. Pères missionnaires dans les grands lacs africains Oukéréwé et Tanganika*: 39 p. (Paris).
- Bourne, G. C.**
- (76) 1906, *On Jousseaumia*: in W. A. Herdman, Report to the government of Ceylon on the pearl oyster fisheries of the gulf of Manaar, pt. 5, p. 243-266, 3 pl., Royal Society (London).

Bouwman, L. A. H.

- (77) 1937, *Sur une espèce nouvelle du genre Sabinia (caprinidés)*: Akad. Ned. Wetensch., Proc., v. 40, no. 5, p. 449-453, text fig. 1-10.
 (78) 1938, *Sur un genre nouveau (Kugleria) de la famille des requienidés pachyodontes*: Same, v. 41, no. 4, p. 418-429, 1 text fig.

Branson, C. C.

- (79) 1935, *Fresh-water invertebrates from the Morrison (Jurassic?) of Wyoming*: Jour. Paleontology, v. 9, p. 514-522, pl. 56-57.
 (80) 1942, *Parallelodon, Grammatodon, and Beushausenia*: Same, v. 16, p. 247-249.
 (81) 1942, *Correction of homonyms in the lamelli-branch genus Conocardium*: Same, v. 16, no. 3, p. 387-392, pl. 59.

Branson, E. B.

- (82) 1909, *The fauna of the residuary Auburn chert of Lincoln County, Missouri*: Acad. Sci. St. Louis, Trans., v. 18, no. 4, p. 39-52, pl. 7.

Broadhurst, F. M.

- (83) 1959, *Anihraconia pulchella sp. nov. and a study of palaeoecology in the Coal Measures of the Oldham Area of Lancashire*: Geol. Soc., London, Quart. Jour., v. 114, p. 523-545, text fig. 1-6.

Brown, Thomas

- (84) 1837-49, *Illustrations of the fossil conchology of Great Britain and Ireland*: viii+273 p., 98 pl.

Bruguère, J. B.

- (85) 1792, *Sur une nouvelle coquille du genre de l'anodontite (Anodontites crispata)*: Soc. Histoire Nat. Paris, Jour., v. 1, p. 131-136, pl. 8.

Brusina, Spiridion

- (86) 1897, *Matériaux pour la faune malacologique néogène de la Dalmatie, de la Croatie et de la Slavonie avec des espèces de la Bosnie, de l'Herzégovine et de la Serbie*: Djela Jugosl. Akad., Znanosti Umjetnosti, v. 18, ser. 6, 43 p., 21 pl.
 (87) 1902, *Iconographia molluscorum fossilium in tellure tertiaria Hungariae, Croatiae, Slavoniae, Dalmatiae, Bosniae, Herzegovinae et Bulgariae inventorum*: X, atlas, 30 pl. (Agram).
 (88) 1905, *Revision des Dreissensidae vivants du système européen*: Jour. Conchyliologie, v. 53, no. 3, p. 272-297.

Buchsbaum, Ralph

- (89) 1948, *Animals without backbones*: revised edit., 405 p., illus., Univ. Press (Chicago).

Bucquoy, E., Dautzenberg, Philippe, & Dollfus, G. F.

- (89a) 1887-98, *Les mollusques marins du Rousillon*: v. 2, Pélécy-podes, 884 p., 99 pl., J.-B. Baillièrre et fils (Paris).

Bulatoff, G. A., & Rjabtschikoff, P. I.

- (90) 1933, *Eine neue Gattung aus der Familie der Tereidinidae aus dem Japanischen Meer*: Zool. Anzeiger, v. 104, p. 165-175, text fig. 1-12.

Butts, Charles

- (91) 1926, *The Paleozoic rocks*: in Geology of Alabama, Geol. Survey Alabama, Spec. Rept. no. 14, p. 41-230, pl. 3-76.

Casey, Raymond

- (92) 1952, *Some genera and subgenera, mainly new, of Mesozoic heterodont lamellibranchs*: Malacol. Soc. London, Proc., v. 29, pt. 4, p. 121-176, 100 text fig., pl. 7-9.
 (93) 1961, *The stratigraphical palaeontology of the Lower Greensand*: Palaeontology, v. 3, pt. 4, p. 487-621, pl. 77-84.

Caster, K. E.

- (94) 1930, *Higher fossil faunas of the upper Allegheny*: Bull. Am. Paleontology, v. 15, no. 58, p. 143-256 (1-114), pl. 22-49 (43-70).

Chao, Y. T.

- (95) 1927, *Fauna of the Taiyuan Formation of North China—Pelecypoda*: Palaeontologia Sinica, ser. B, v. 9, pt. 3, 57 p., 4 pl.

Chavan, André

- (96) 1936, *Sur la signification paléontologique du genre Pteromeris Conrad. Discussion des espèces égarées rapportées à ce genre*: Soc. Géol. France, Bull., sér. 5, v. 6, p. 251-59.
 (97) 1936, *Les carditidés du Calcaire de Mons*: Musée Royal Histoire Nat. Belgique, Bull., v. 12, no. 39, 13 p., 1 pl.
 (98) 1937-38, *Essai critique de classification des lucines*: Jour. Conchyliologie, v. 81, p. 133-153, 198-216, 237-282; v. 82, p. 59-99, 105-130, 215-243.
 (99) 1938, *Un genre africain de carditidés: Cossmannella Mayer-Eymar*: Soc. Géol. France, Bull., sér. 5, v. 8, p. 3-10, text fig.
 (100) 1939, *Sur quelques Crassatellidae tertiaires: Chattonia, Crassatina, Crassinella*: Musée Royal Histoire Nat. Belgique, Bull., v. 15, no. 34, 36 p., 2 text fig.
 (101) 1941, *Sur deux nouveaux groupes de Carditidae*: Jour. Conchyliologie, v. 84, p. 97-102, fig.
 (102) 1942, *Sur deux nouveaux groupes de Lucinidae*: Soc. Géol. France, Comptes Rendus, no. 7, p. 61-63.
 (103) 1946, *Notes sur les Jagonia (lamellibranches)*: Muséum Natl. Histoire Nat., Bull., v. 18, no. 1, p. 87-90.
 (104) 1946, *Nouvelles notes sur les Jagonia (lamellibranches)*: Same, Bull., v. 18, no. 4, p. 345-47.
 (105) 1947, *L'évolution des faunes marines de mollusques dans le Nord-Ouest de l'Europe*,

- de la fin du Crétacé à celle de l'Éocène: Soc. Géol. France, Bull., sér. 5, v. 16, p. 193-212.
- (106) 1949, *Remarques sur la signification climatique des mollusques marins fossiles*: Same, Bull., sér. 5, v. 19, p. 507-512.
- (107) 1951, *Essai critique de classification des Divaricella*: Inst. Royal Sci. Nat. Belgique, Bull., v. 27, no. 18, p. 1-27, 27 text fig.
- (108) 1951, *Les affinités du genre Microstagon Cossmann*: Cahiers Géologiques, Seyssel, no. 8, p. 70-71.
- (109) 1952, *Mélanges paléontologiques 2. Distinction et classement des crassatellidés*: Cahiers Géologiques Thoiry, v. 14, p. 117-120, 1 fig.
- (110) 1952, *Mélanges paléontologiques 3. Distinction et classement des astaridés*: Same, v. 15, p. 123-127, 2 text fig.
- (111) 1952, *Mélanges paléontologiques 4. Sur Astarte thoas d'Orbigny*: Same, v. 15, p. 127-128, 1 fig.
- (112) 1952, *Les pélicypodes des sables astartiens de Cordebugle (Calvados)*: Soc. Paléont. Suisse, Mém., v. 69, p. 1-132, 82 text fig., pl. 1-4.
- (113) 1952, *Nomenclatural notes on carditids and lucinids*: Washington Acad. Sci., Jour., v. 42, no. 4, p. 116-122.
- (114) 1953, *Notes de malacologie africaine*: Revue Zool. Bot. Africaines, v. 47, no. 1-2, p. 132-141, 12 text fig.
- (115) 1953, *Sur Astarte disparilis d'Orbigny*: Cahiers géologiques Thoiry, v. 19, p. 164-165, 1 text fig.
- (116) 1954, *Les Pleurophorus et genres voisins*: Cahiers Géologiques Seyssel, no. 22, p. 200.
- (117) 1956, *Remarques sur l'origine des Lucinacea (Mollusques Pélicypodes)*: Soc. Géol. France, Comptes Rendus Somm., no. 4, p. 163-164.
- (118) 1957, *Observations sur les Myoconcha*: Same, no. 41, p. 415.
- (119) 1959, *Essai critique de classification des Mactromyidae*: Same, no. 52, p. 505-508; no. 53, p. 509-512.
- (120) 1960(1959), *Remarques sur la charnière des Erycinacea et des Cyamiacea*: Soc. Géol. France, Bull., 7e sér., v. 1, p. 712-718.
- (121) 1962, *Essai critique de classification des Ungulinidae*: Inst. Royal Sci. Nat. Belgique, Bull., v. 38, no. 23, p. 1-23, 14 text fig.
- Chemnitz, J. H.**
 (122) 1784-85, *Neues systematisches Conchylien-Cabinet*: v. 7, 356 p., pl. 37-69; v. 8, 372 p., pl. 70-102, Bauer & Raspe (Nürnberg).
- Chen, Chu-chen**
 (123) 1964, *On the occurrence of Halobia fauna from the Ganzhi region, western Sichuan (Szechuan) and its significance*: Acta Palaeont. Sinica, v. 12, no. 1, p. 66-78.
- Chenu, H. G.**
 (124) 1859-62, *Manuel de conchyliologie et de paléontologie conchyliologique*: (a) v. 1, vii+508 p. (1859), (b) v. 2, 327 p. (1862), Masson (Paris).
- Chernyshev [Tchernyschew], B. I.**
 (125) 1950, *Semeistvo Grammysiidae iz verkhne-paleozoiskikh otlozhenii SSSR*: Akad. Nauk Ukrain. SSR, Kiev, Inst. Geol. Nauk, Trudy, Ser. Strat. Paleont., no. 1, p. 3-91, 14 pl. [The family Grammysiidae from the upper Paleozoic deposits of the USSR.]
 (126) 1951, *Semeistvo Ledidae iz Kamennougolnykh otlozhenii SSSR*: Same, no. 2, p. 3-40, 2 pl. [The family Ledidae from the Carboniferous deposits of the USSR.]
- Choffat, Paul**
 (127) 1885-88, *Déscription de la faune jurassique de Portugal. Mollusques lamelibranches. Ordre Asiphonida*: Travaux Géol. Portugal, Mém., 76 p., 19 pl.
- Chronic, Halka**
 (128) 1952, *Molluscan fauna from the Permian Kaibab Formation, Walnut Canyon, Arizona*: Geol. Soc. America, Bull., v. 63, p. 95-166, pl. 1-11.
- Chubb, L. J.**
 (129) 1955, *A revision of Whitfield's type specimens of the rudist mollusks from the Cretaceous of Jamaica, British West Indies*: Am. Museum Nat. History, Novitates, no. 1713, p. 1-15.
 (130) 1956, *Some rarer rudists from Jamaica, British West Indies*: Palaeont. Americana, v. 4, no. 26, p. 1-24, pl. 1-5, text fig. 1-4.
 (131) 1956, *Thyrastylon, a new rudist genus from the Upper Cretaceous of Guatemala, the Antilles, and Persia, with a discussion of the functions of rudist oscules and pillars*: Same, v. 4, no. 27, p. 33-48, pl. 6, 7.
- Ciriacks, K. W.**
 (132) 1963, *Permian and Eotriassic bivalves of the middle Rockies*: Am. Museum Nat. History, Bull., v. 125, art. 1, 100 p., 16 pl.
- Clapp, W. F., & Kenk, Roman**
 (133) 1963, *Marine borers. An annotated bibliography*: xii+1136 p., Office of Naval Research, Dept. of Navy (Washington, D.C.).
- Clark, B. L.**
 (134) 1934, *A new genus and two new species of Lamelibranchiata from the middle Eocene of California*: Jour. Paleontology, v. 8, no. 3, p. 270-272, pl. 37.
- Clarke, J. M.**
 (135) 1899, *Molluscos devonianos do Estado do*

- Pará, Brazil: Archiv. Museu Nac. Rio de Janeiro, v. 10, p. 49-174, pl. 3-8.
- (136) 1904, *Naples fauna in western New York*: N.Y. State Museum, 57th Ann. Rept., v. 3, Mem. 6, p. 199-454, 20 pl.
- (137) 1909, *Early Devonian history of New York and eastern North America*: Same, Mem. 9, 250 p.
- (138) 1913, *Fosseis Devonianos do Parana*: Serv. Geol. & Mineral. Brasil, Mon., v. 1, 353 p., 2 text fig., 27 pl.
- , & Ruedemann, Rudolf
- (139) 1903, *Guelph fauna in the state of New York*: N.Y. State Museum, Mem. 5, 195 p., 21 pl.
- Clench, W. J., & Turner, R. D.**
- (139a) 1946, *The genus Bankia in the western Atlantic*: Johnsonia, v. 2, no. 19, p. 1-28, pl. 1-16.
- Clessin, Stefan**
- (140) 1873-76, *Die Gattung Anodonta nebst den übrigen Najaden mit unvollkommenem Schloss*: in *Abbildungen nach der Natur mit Beschreibungen*. Begonnen von Dr. H. C. Kuester, fortgesetzt und vollendet von S. Clessin, Martini-Chemnitz, Syst. Conch. Cab. (2), v. 9, no. 1, p. 65-287, pl. 17-87.
- Cockerell, T. D. A.**
- (141) 1903, *Some homonymous generic names*: Nautilus, v. 16, p. 118.
- Connolly, M. W. K.**
- (142) 1939, *A monographic survey of South-African non-marine Mollusca*: South African Museum, Ann., v. 33, 660 p., 58 text fig., 19 pl.
- Conrad, T. A.**
- (143) 1834, *Description of a new genus of fresh-water shells*: Acad. Nat. Sci. Philadelphia, Jour., v. 7, p. 178-180.
- (144) 1835-38, *Monograph of the family Unionidae or Najades of Lamarck (Fresh water bivalve shells)*: 110 p., 60 pl. (unfinished) (Philadelphia).
- (145) 1841, *On the palaeontology of the state of New York*: N.Y. State Geol. Survey, 5th Ann. Rept., p. 25-27.
- (146) 1842, *Descriptions of new species of organic remains belonging to the Silurian, Devonian and Carboniferous systems of the United States*: Acad. Nat. Sci. Philadelphia, Jour., v. 8, p. 183-190.
- (147) 1853, *Synopsis of the family of Naiades of North America; with notes and a table of some of the genera and subgenera*: Same, Proc., v. 6, p. 243-269, 449.
- (148) 1855, *Observations on the Eocene deposit of Jackson, Mississippi, with descriptions of thirty-four new species of shells and corals*: Same, Proc., v. 7, p. 257-263. (Figured in B. L. C. Wailes, Report on the geology and agriculture of Mississippi, 1854.)
- (149) 1857, *Descriptions of two new genera of shells*: Same, Proc., v. 11, p. 165-166.
- (150) 1862, *Descriptions of new genera, subgenera and species of Tertiary and Recent shells*: Same, Proc., 1862, p. 284-291.
- (151) 1865, *Descriptions of new Eocene shells from Enterprise, Mississippi*: Am. Jour. Conchology, v. 1, p. 137-141, 1 pl.
- (152) 1865, *Remarks on the genera Monocondylaea d'Orbigny, and Pseudodon Gould, with a synopsis of the latter*: Same, v. 1, p. 232-233.
- (153) 1865, *Description of a new genus of Unionidae*: Same, v. 1, p. 224.
- (154) 1867, *Descriptions of new genera and species of fossil shells*: Am. Jour. Conchology, v. 3, p. 8-16.
- (155) 1869, *Notes on Recent and fossil shells, with descriptions of new genera*: Same, v. 4, p. 246-249.
- (156) 1869, *Descriptions of Miocene, Eocene, and Cretaceous shells*: Same, v. 5, p. 39-45.
- (157) 1869, *Descriptions of new fossil Mollusca, principally Cretaceous*: Same, v. 5, p. 96-103.
- Conrath, P.**
- (158) 1888, *Über einige silurischen Pelecypoden*: Akad. Wiss. Wien, Sitzungsber., v. 96, p. 40-51, pl. 1-28.
- Cooper, G. A.**
- (159) 1931, *Concerning the authorship of the "Preliminary notice of the lamellibranch shells of the upper Helderberg, Hamilton and Chemung groups, etc., Part 2"*: Washington Acad. Sci., Jour., v. 21, p. 459-467.
- Cossmann, Maurice**
- (160) 1887, *Catalogue illustré des coquilles fossiles de des environs de Paris*: Soc. Royale Belgique, Ann., v. 22, no. 2ème, 214 p., 11 pl. (same, 1913, v. 49 (5ème append.), p. 19-238, pl. 1-8).
- (161) 1912, *Sur l'évolution des Trigones*: Ann. Paléontologie, v. 7, p. 59-84.
- (162) 1920, *Rectifications de nomenclature*: Revue Critique de Paléozoologie, v. 24, no. 2, p. 81-83.
- (163) 1925, *Essais de paléonchologie comparée*: v. 13, 345 p., 11 pl., Cossmann (Paris).
- , & Peyrot, Adrien
- (164) 1909, *Conchologie néogénique de l'Aquitaine (Pelecypoda, outline of classification)*: Soc. Linnéenne Bordeaux, Actes, v. 63, p. 89-100.
- (165) 1912, *Conchologie néogénique de l'Aquitaine*: Same, Actes, v. 66, p. 121-324.
- Cotton, B. C.**
- (166) 1930, *Pelecypoda of the "Flindersian" re-*

- gion, southern Australia, No. 1: S. Austral. Museum Records, v. 4, p. 223-240, 15 text fig.
- (167) 1961, *South Australian Mollusca. Pelecypoda*: 363 p., 351 text fig., Government Printer (Adelaide, South Australia).
- , & **Gabriel, C. J.**
- (168) 1931, *Australian Unionidae*: Royal Soc. Victoria, Proc., v. 44, p. 155-160, pl. 16.
- , & **Godfrey, F. K.**
- (169) 1938, *The Mollusca of South Australia. Part I. The Pelecypoda*: 314 p., 340 text fig. (Adelaide).
- Cox, L. R.**
- (170) 1925, *The fauna of the basal shell-bed of the Portland stone, Isle of Portland*: Dorset Nat. History, Proc., v. 46, p. 113-172, 5 pl.
- (171) 1929, *A synopsis of the Lamellibranchia and Gastropoda of the Portland beds of England. Part 1. Lamellibranchia*: Dorset Nat. History & Arch. Soc., Proc., v. 50, p. 131-202, pl. 1-6.
- (172) 1930, *On British fossils named by William Smith*: Ann. & Mag. Nat. History, ser. 10, v. 6, p. 287-304, pl. 12.
- (173) 1931, *New lamellibranch genera from the Tethyan Eocene*: Malacol. Soc. London, Proc., v. 19, pt. 4, p. 177-187, text fig. 1-2, pl. 20-21.
- (174) 1932, *Lamellibranchia from the Karroo Beds of the Ruhuhu Coalfields, Tanganyika Territory*: Geol. Soc. London, Quart. Jour., v. 88, p. 623-633, pl. 39-40.
- (175) 1933, *The evolutionary history of the rudists*: Geologists' Assoc. London, Proc., v. 44, p. 379-388.
- (176) 1934, *On the structure of the Persian rudist genus Trechmannella (formerly Polyptychus), with a description of a new species*: Malacol. Soc. London, Proc., v. 21, pt. 1, p. 42-66, text fig. 1-27, pl. 8.
- (177) 1935, *Jurassic Gastropoda and Lamellibranchia*: in *The Mesozoic palaeontology of British Somaliland*, p. 148-197, pl. 14-21.
- (178) 1936, *Karoo Lamellibranchia from Tanganyika Territory and Madagascar*: Geol. Soc. London, Quart. Jour., v. 92, p. 32-57, pl. 4-5.
- (179) 1937, *Notes on Jurassic Lamellibranchia. 1. On the occurrence of the genus Palaeoneilo in the Jurassic of Great Britain*: Malacol. Soc. London, Proc., v. 22, p. 190-193, pl. 15.
- (180) 1937, *Notes on Jurassic Lamellibranchia. 2. On Indogrammatodon, a new subgenus from the Jurassic of the Indo-African province*: Same, v. 22, p. 194-198, pl. 15-16.
- (181) 1937, *Notes on Jurassic Lamellibranchia. 3. On a new Trigonina and other species from Tanganyika Territory*: Same, v. 22, p. 198-203, pl. 16.
- (182) 1937, *Notes on Jurassic Lamellibranchia. 5. On a new subgenus of Mytilus and a new Mytilus-like genus*: Same, v. 22, p. 339-348, pl. 17.
- (183) 1940, *The Jurassic lamellibranch fauna of Kuchh (Cutch)*: Geol. Survey India, Mem., Palaeont. Indica, ser. 9, v. 3, pt. 3, 157 p., 10 pl.
- (184) 1942, *Notes on Jurassic Lamellibranchia. 8. On the genus Velata Quenstedt*: Malacol. Soc. London, Proc., v. 25, p. 119-124, pl. 4.
- (185) 1943, *The English upper Lias and Inferior Oolite species of Lima*: Same, Proc., v. 25, pt. 5-6, p. 151-187, pl. 6-29.
- (186) 1944, *On Pseudolimea Arkell*: Same, v. 26, pt. 2-3, p. 74-88, pl. 2-3.
- (187) 1946, *Tutcheria and Pseudopsis, new lamellibranch genera from the Lias*: Same, Proc., v. 27, p. 34-48, pl. 3-4.
- (188) 1946, *The fossils and age of the Sekondi series (Upper Palaeozoic) of the Gold Coast*: Annals & Mag. Nat. History, ser. 11, v. 13, p. 602-624, pl. 12-13.
- (189) 1947, *The lamellibranch family Cyprinidae in the lower oolites of England*: Malacol. Soc. London, Proc., v. 27, pt. 4, p. 141-184, pl. 8-10.
- (190) 1949, *On Linsdallia, a genus of Lamellibranchia from the Inferior Oolite*: Same, v. 27, pt. 6, p. 251-252 (Jan. 14, 1949).
- (191) 1951, *The lamellibranch genus Schizodus and other Palaeozoic Myophoriidae*: Geol. Mag., v. 88 (Sept.-Oct.), p. 362-371.
- (192) 1952, *Notes on the Trigonidae, with outlines of a classification of the family*: Malacol. Soc. London, Proc., v. 29, p. 45-70, pl. 3-4.
- (193) 1952, *The Jurassic lamellibranch fauna of Cutch (Kuchh), no. 3, Families Pectinidae, Amusiidae, Plicatulidae, Limidae, Ostreidae and Trigonidae (Suppl.)*: Geol. Survey India, Mem., Palaeont. Indica, ser. 9, v. 3, pt. 4, p. 1-128, pl. 1-12.
- (194) 1954, *Taxonomic notes on Isognomonidae and Bakevellidae*: Malacol. Soc. London, Proc., v. 31, pt. 2, p. 46-49, 1 fig. (Dec. 31, 1954).
- (195) 1955, *On the affinities of the Cretaceous lamellibranch genera Trigonoides and Hoffetrigonia*: Geol. Mag., v. 92, no. 4 (July-Aug.), p. 345-349.
- (196) 1959, *The geological history of the Protobranchia and the dual origin of taxodont Lamellibranchia*: Malacol. Soc. London, Proc., v. 33, pt. 5, p. 200-209.
- (197) 1960, *Thoughts on the classification of the*

- Bivalvia*: Same, v. 34, pt. 2, p. 60-88, 2 text fig.
- (198) 1960, *The preservation of moulds of the intestine in fossil Nuculana (Lamellibranchia) from the Lias of England*: Palaeontology, v. 2, pt. 2, p. 262-269, pl. 40.
- (199) 1960, *Two new radiolitids (rudistid Lamellibranchia) from the Upper Cretaceous of Turkey*: British Museum (Nat. History), Bull., v. 4, no. 9, p. 425-433, 1 text fig., pl. 61-62.
- (200) 1961, *New genera and subgenera of Mesozoic Bivalvia*: Palaeontology, v. 4, pt. 4, 1961, p. 592-598.
- (201) 1961, *Observations on the family Cardiniidae (Class Bivalvia)*: Malacol. Soc. London, Proc., v. 34, pt. 6, p. 325-339 (Dec.).
- (202) 1962, *Tertiary Bivalvia from Libya*: Palaeontology, v. 5, pt. 1, p. 1-8, pl. 1-5.
- (203) 1963, *The Rhaetic-Hettangian bivalve genus Pteromya Moore*: Same, v. 6, pt. 3, p. 582-595, pl. 79-80.
- , & Arkell, W. J.
- (204) 1948-49, *A survey of the Mollusca of the British Great Oolite series, primarily a nomenclatorial revision of the monographs by Morris and Lycett (1851-55), Lycett (1863) and Blake (1905-1907)*: Palaeontograph Soc., pt. 1 (1948), xiii+48 p.; pt. 2 (1949), xiv-xxiv+49-105 p.
- , & Eames, F. E.
- (205) 1956, *Some Tertiary Pectinacea from East Africa, Persia, and the Mediterranean region*: Malacol. Soc. London, Proc., v. 32, pt. 1 & 2, p. 1-68.
- Crickmay, C. H.**
- (206) 1932, *Contributions toward a monograph of the Trigoniidae*: Am. Jour. Sci., v. 24, p. 443-464.
- Cumings, E. R.**
- (207) 1908, *The stratigraphy and paleontology of the Cincinnati series of Indiana*: Dept. of Geol. & Nat. Res. Indiana, 32nd Ann. Rept., p. 607-1188, 55 pl.
- (208) 1922, *Nomenclature and description of the geological formations of Indiana*: in Handbook of Indiana Geology, publ. 21, by W. N. Logan et al., p. 403-570, text fig. 1-31 (Indianapolis).
- Cuvier, G. L. C. F. D.**
- (209) 1797(1798), *Tableau élémentaire de l'histoire naturelle des animaux*: xvi+710 p., 14 pl. (Paris).
- (210) 1800, *Leçons d'anatomie comparée*: v. 1, xxxi+521 p., 9 pl. (Paris).
- Cvancara, A. M.**
- (210a) 1966, *Revision of the fauna of the Cannonball Formation (Paleocene) of North and South Dakota*: Univ. Michigan, Museum Paleontology, Contrib., 20, p. 277-370, pl. 5-11.
- Dahmer, Georg**
- (211) 1917-21, *Studien über die Fauna des Oberharzer Kahlebergsandsteins. I. and II.*: Preuss. Geol. Landesanst., Jahrb., (a) v. 37 (1917), pt. 1, p. 443-526, 4 text fig., pl. 36-39; (b) v. 40 (1921), pt. 2, p. 161-306, 7 text fig., pl. 6-17.
- (212) 1922, *Die Fauna der obersten Koblenzschichten am Nordwestrand der Dillmulde*: Same, Jahrb., v. 42, p. 655-693, 4 text fig., pl. 9-10.
- (213) 1936, *Die Fauna der Siegener Schichten von Unkel (Bl. Königswinter)*: Same, v. 56, p. 633-671, 1 text fig., pl. 44-48.
- (214) 1946, *Revidiertes Verzeichnis der Versteinerungen des Oberharzer Kaleberg-Sandsteins (Unter-Devon)*: Senckenbergiana, v. 27, p. 166-187, 10 text fig.
- (215) 1948, *Die Fauna des Koblenzquarzits (Unterdevon, Oberkoblenz-Stufe) vom Kühkopf bei Koblenz*: Same, v. 29, p. 115-136, 25 text fig.
- Dall, W. H.**
- (216) 1886, *Reports on the results of dredging, under the supervision of Alexander Agassiz, in the Gulf of Mexico (1877-78) and in the Caribbean Sea (1879-80), by the U.S. Coast-Survey Steamer "Blake," Lieut.-Commander C. D. Sigsbee, U.S.N., and Commander J. R. Barlett, U.S.N., Commanding. XXIX, Report on the Mollusca,—Part I. Brachiopoda and Pelecypoda*: Harvard Univ., Museum Comp. Zoology, Bull., v. 12, no. 6, p. 171-318, 9 pl.
- (217) 1889, *Scientific results of explorations U.S. Fish Comm. Albatross. No. VII. Preliminary report on the collection of Mollusca and Brachiopoda obtained in 1887-88*: U.S. Natl. Museum, Proc., v. 12, p. 217-362.
- (218) 1889, *On the hinge of pelecypods and its development, with an attempt toward a better subdivision of the group*: Am. Jour. Sci., v. 38, no. 3, p. 445-462.
- (219) 1895-1903, *Contributions to the Tertiary fauna of Florida, with especial reference to the Miocene Silex beds of Tampa and the Pliocene beds of Caloosahatchie river*: Wagner Free Inst. Sci., Philadelphia, Trans., v. 3, pt. 3(1895), p. 483-565; pt. 4(1898), p. 571-947, pl. 23-35; pt. 5(1900), p. 949-1218, pl. 36-47; pt. 6(1903), p. 1219-1654, pl. 48-60.
- (220) 1895, *Maclridae and Mesodesmatidae*: Malacol. Soc. London, Proc., v. 1, pt. 5, p. 203-213.

- (221) See 219.
- (222) 1896-1900, *Pelecypoda*: in K. A. von Zittel, Text-book of palaeontology (transl. & ed. C. R. Eastman), v. 1, p. 346-429, text fig. 588-787, Macmillan & Co. (New York, London).
- (223) 1898, *Synopsis of the Recent and Tertiary Psammobiidae of North America*: Acad. Nat. Sci. Philadelphia, Proc., v. 50, p. 57-62.
- (224) See 219.
- (225) 1899, *Synopsis of the Recent and Tertiary Leptonacea of North America and the West Indies*: U.S. Natl. Museum, Proc., v. 21, no. 1177, p. 873-897, pl. 87-88.
- (226) 1901, *Synopsis of the family Cardidae and of the North American species*: Same, Proc., v. 23, no. 1214, p. 381-392.
- (227) 1901, *Synopsis of the Lucinacea and of the American species*: Same, Proc., v. 23, no. 1237, p. 779-834, pl. 39-42.
- (228) 1902, *Synopsis of the family Veneridae and of the North American species*: Same, Proc., v. 26, p. 335-412, pl. 12-16.
- (229) 1902, *Synopsis of the Carditacea and of the American species*: Acad. Nat. Sci. Philadelphia, Proc., v. 54, p. 696-716.
- (230) 1903, *Synopsis of the family Astartidae, with a review of the American species*: U.S. Natl. Museum, Proc., v. 26, p. 933-951, pl. 61-63.
- (231) 1903, *Review of the classification of the Cyrenacea*: Biol. Soc. Washington, Proc., v. 16, p. 5-8.
- (232) 1908, *The Mollusca and the Brachiopoda*: Harvard Univ., Museum Comp. Zoology, Bull., v. 43, no. 6, p. 205-487.
- (233) 1908, *Descriptions of new species of mollusks from the Pacific coast of the United States, with notes on other mollusks from the same region*: U.S. Natl. Museum, Proc., v. 34, no. 1610, p. 245-257.
- (234) 1913, *Pelecypoda*: in K. A. von Zittel, Text-book of palaeontology, transl. & ed. C. R. Eastman, rev. edit., v. 1, p. 421-507, fig. 637-835, Macmillan & Co. (London, New York).
- Darragh, T. A.**
- (235) 1964, *A preliminary revision of the living species of Eucrassatella (Pelecypoda: Crassatellidae)*: Malacol. Soc. Australia, Jour., no. 8, p. 3-9, 3 pl.
- Dautzenberg, Philippe**
- (236) 1927, *Mollusques provenant des campagnes scientifiques du Prince Albert 1^{er} de Monaco dans l'Océan Atlantique et dans le Golfe de Gascogne*: Résult. Campagnes Scientifiques accomplies sur son Yacht par Albert 1^{er} Prince de Monaco: pt. 72, 398 p., 9 pl. (Monaco).
- Davidaschwili, L. C.**
- (237) 1930, *Kimmeriyskiy yarus. Rukovodiashchiye iskopaemiye neftenosnykh rayonov Krymsko-Kavkazskoy oblasti. VIII: Gosudarst. Issledov. Nefti. Inst., Trudy (Transactions of the State Petroleum Institute), no. 6, 44 p., 5 pl. (Nauchno-Tekhnicheskoe upravleniye VSNKh, no. 322, SSSR: Moscow). [Kimmerian beds. Characteristic fossils of the oil districts of the Crimea and Caucasus. no. 8.]*
- (238) 1931, *Ponticheskii yarus. Rukovodiashchiye iskopaemiye neftenosnykh rayonov Krymsko-Kavkazskoy oblasti. VII: Same, no. — [not given], 56 p., 7 pl. [Pontian beds. Characteristic fossils of the oil districts of the Crimea and Caucasus, no. 7.]*
- (238) 1932, *Sarmatskiy yarus. Rukovodiashchiye iskopaemiye neftenosnykh rayonov Krymsko-Kavkazskoy oblasti. V: Same, no. — [not given], 77 p., 10 pl. [Sarmatian beds. Characteristic fossils of the oil districts of the Crimea and Caucasus, no. 5.]*
- , & **Krestovnikov, V. N.**
- (240) 1931, *Duabskiye plasty. Rukovodiashchiye iskopaemiye neftenosnykh rayonov Krymsko-Kavkazskoy oblasti. IX: Gosudarst. Issledov. Nefti. Inst. [Trans. State Petroleum Institute], Trudy, no. 10, p. 27, pl. 3. [Duab beds. Characteristic fossils of the oil districts of the Crimea and Caucasus, no. 9.]*
- Davies, A. M.**
- (241) 1933, *The bases of classification of the Lamellibranchia*: Malacol. Soc. London, Proc., v. 20, p. 322-326.
- (242) 1935, *Tertiary faunas: a text-book for oil-field palaeontologists and students of geology. Vol. I. The composition of Tertiary faunas*: xi + 406 p., 568 text fig., Thomas Murby & Co. (London).
- Dechaseaux, Colette**
- (243) 1936, *Limidés jurassiques de l'est du Bassin de Paris*: Musée Royal Histoire Nat. Belgique, Mém., ser. 2, pt. 8, 58 p., 3 pl.
- (244) 1936, *Pectinidés jurassiques de l'est du Bassin de Paris*: Ann. Paléontologie, v. 25, 148 p., 10 pl.
- (245) 1939, *Megalodon, Pachyerisma, Prodiceras, Dicerias, Pterocardium et l'origine des Dicerias*: Soc. Géol. France, Bull., v. 7, no. 4, p. 96-114, pl. 2 (also sér. 5, v. 9, p. 207-218, pl. 15).
- (246) 1941, *Rudistes I, Dicerias et Heterodicerias*: Same, Mém., new ser., v. 20, Mém. 44, 55 p., 10 text fig.
- (247) 1941, *Les Lamellibranches. I-Rudistes (fasc. 1)*: Catalogue des échantillons types et figurées de la collection de paléontologie,

- École National Supérieure des Mines, 11 p. (Paris).
- (248) 1943, *La notion d'inversion ne peut s'appliquer aux Rudistes*: Soc. Géol. France, Comptes Rendus Somm. Séanc., p. 67-69.
- (249) 1947, *Bandes siphonales, piliers et siphons des rudistes*: Same, Bull., sér. 5, v. 17, p. 425-435.
- (250) 1948, *Le problème de l'extinction des groupes étudié chez les rudistes*: Revue Scientifique, 86° année, pt. 2, no. 3289, p. 83-86.
- (251) 1949, *Essai sur la paléobiologie des rudistes: Le genre Bournonia*: Ann. Paléontologie, v. 35, p. 119-130, pl. 1-4, text fig. 1-3.
- (252) 1952, *Classe des Lamellibranches*: in Jean Piveteau (ed.), *Traité de paléontologie*, v. 2, 790 p., Masson et Cie (Paris) (Rudistae, p. 323-364, text fig. 163-215, 2 pl.).
- (253) 1954, *Forme et milieu chez quelques rudistes*: Ann. Hébert Haug, v. 8, p. 191-195, 1 text fig.
- (254) 1954, *Étude sur un rudiste de Syrie: Le genre Vautrinia Milovanovic*: Moyen-Orient, Notes & Mém., v. 5, p. 173-190, text fig. 1-11, pl. 1-4.
- Delgado, J. F. N.**
 (255) 1904, *Faune cambrienne du Haut-Alemtejo (Portugal)*: Comm. Géol. Portugal, Commun., v. 5, p. 307-374, pl. 1-6.
- Depéret, Charles, & Roman, Frederic**
 (256) 1902-28, *Monographie des pectinides néogènes de l'Europe et des régions voisines*: Soc. Géol. France, Paléont., Mém. 26, (a) v. 10, pt. 1 (1902), p. 1-74, text fig. 1-33, pl. 1-8; (b) v. 13, pt. 2 (1905), suppl., p. 75-104, text fig. 34-43, pl. 9(6)-11(8); (c) v. 18, pt. 2 (1910), p. 105-139, text fig. 44-60, pl. 12(8)-17(13); (d) v. 19, pt. 1, p. 139-168 (1912), text fig. 61-71, pl. 18(1)-23(6); (e) new ser., Mém. 10, v. 4, pt. 4 (1928), p. 169-194, text fig. 1-10, pl. 24(20)-28(24). [For completion of Mém. see Roger, J.]
- Deshayes, G. P.**
 (256a) 1825, *Quelques observations sur les genres Hippurite et Radiolite*: Ann. Sci. Nat., ser. 1, v. 5, p. 205-211.
 (257) 1839-53, *Traité élémentaire de conchyliologie*: v. 1, pt. 1, p. 1-272 (1839); xii + 273-368 p. (1853); v. 2, 384 p. (not completed), 1853; plates and explanation 1838-1853 (dates uncertain).
 (258) 1855 (1856), *Descriptions de nouvelles espèces du genre Erycina*: Zool. Soc. London, v. 23, p. 181-183.
 (259) 1856 (1860-66), *Description des animaux sans vertèbres découverts dans le Bassin de Paris*: v. 1, 912 p., 89 pl., v. 2, 968 p., 107 pl.; J.-B. Baillière et fils (Paris).
- Dickins, J. M.**
 (260) 1960, *Characters and relationships of the Mesozoic pelecypod Pseudavicula*: Palaeontology, v. 3, pt. 3, p. 392-396.
 (261) 1960, *The Permian leiopteriid Merismopteria and the origin of the Pteriidae*: Same, v. 3, pt. 3, p. 387-391, 1 text fig.
 (262) 1961, *Eurydesma and Peruwispira from the Dwyka beds of South Africa*: Same, v. 4, pt. 1, p. 138-148, pl. 18.
 (263) 1963, *Permian pelecypods and gastropods from Western Australia*: Austral. Bur. Mineral Resources, Bull. 63, 150 p., 26 pl.
- _____, & **Shah, S. C.**
 (264) 1965, *The pelecypod Undulomya, Cosmomya, and Palaeocosmomya in the Permian of India and Western Australia*: Geol. Soc. Australia, Jour., v. 12, p. 253-260, pl. 16-17.
- Diener, Carl**
 (265) 1923, *Lamellibranchiata triadica*: Fossilium Catalogus, 1, Animalia, v. 19, 259 p.
- Dix, Emily, & Trueman, A. E.**
 (266) 1931, *Some non-marine lamellibranchs from the upper part of the Coal Measures*: Geol. Soc. London, Quart. Jour., v. 87, p. 180-211, text fig. 1-8, pl. 17.
- Dodge, Henry***
 (267) 1952, *A historical review of the mollusks of Linnaeus. Part 1. The classes Loricata and Pelecypoda*: Am. Museum Nat. History, Bull., v. 100, art 1, p. 1-263.
- Douvillé, Henri**
 (267a) 1886, *Essai sur la morphologie des rudistes*: Soc. Géol. France, Bull., sér. 3, v. 14, p. 389-404, 19 text fig.
 (267b) 1887, *Sur quelques formes peu connues de la famille des chamides*: Soc. Géol. France, Bull., sér. 3, v. 15, p. 756-802, 18 text fig., pl. 28-31.
 (267c) 1888, *Études sur les caprines*: Same, Bull., sér. 3, v. 16, p. 699-730, 9 text fig., pl. 22-25.
 (267d) 1889, *Sur quelques rudistes du terrain crétacé inférieur des Pyrénées*: Same, Bull., sér. 3, v. 17, p. 627-653, 18 text fig., pl. 15, 16.
 (268) 1891, *Sur les caractères des Sauvagesia*: Same, Bull., sér. 3, v. 19, p. 669-672.
 (269) 1891-97, *Études sur les rudistes*: Same, Mém., Paleont. Mém., no. 6, 236 p., 71 text fig., 34 pl.
 (270) 1896, *Observations sur la charnière des lamellibranches hétérodontes*: Same, Bull., v. 24, no. 3, p. 26-28.
 (270a) 1900, *Sur quelques rudistes Américains*:

- Same, Bull., sér. 3, v. 28, p. 205-221, 17 text fig.
- (271) 1902, *Sur un nouveau genre de radiolites*: Same, Bull., sér. 4, v. 2, p. 478-482, pl. 15.
- (272) 1904, *Sur quelques rudistes à canaux*: Same, Bull., sér. 4, v. 4, p. 519-538, 4 text fig., pl. 13-14.
- (273) 1907, *Les lamellibranches cavicoles ou desmodontes*: Same, Bull., v. 7, no. 4, p. 96-114, pl. 2.
- (274) 1907, *Études sur les lamellibranches. Vulsellidés*: Ann. Paléontologie, v. 2, pt. 3, p. 97(1)-119(23), text fig. 1-11, pl. 15(1)-16(2).
- (274a) 1910, *Rudistes de Sicilie, d'Algérie, d'Égypte, du Liban et de la Perse*: Soc. Géol. France, Paléont., Mém., no. 41, p. 1-83, 77 text fig., pl. 1-7.
- (275) 1911, *Observations sur les ostréidés. Origine et classification*: Same, Bull., sér. 4, v. 10, p. 635-646, pl. 10-11.
- (276) 1912, *Un essai de classification phylogénique des lamellibranches*: Acad. Sci. Paris, Comptes Rendus, v. 154, p. 1677-1682.
- (277) 1913, *Classification des lamellibranches*: Soc. Géol. France, Bull., v. 12, no. 4, p. 419-467.
- (278) 1913, *Sur quelques rudistes du Liban et sur l'évolution des biradiolitins*: Same, Bull., sér. 4, v. 13, p. 409-421, 1 pl.
- (278a) 1918, *Le Barrémien supérieur de Brouzet*, part 3, *Les rudistes*: Same, Paléont. Mém., no. 52, 19 p., 10 text fig., 4 pl.
- (279) 1921, *La charnière dans les lamellibranches hétérodontes et son évolution*: Same, sér. 4, v. 21, p. 116-124.
- (280) 1926, *Quelques fossiles du Crétacé supérieur de Cuba*: Same, Bull., sér. 4, v. 26, p. 127-138, 4 text fig., pl. 7-8.
- (281) 1927, *Nouveaux rudistes du Crétacé de Cuba*: Same, Bull., sér. 4, v. 27, p. 49-56, text fig., 1-2, pl. 4.
- (282) 1935, *Les rudistes et leur évolution*: Same, Bull., sér. 5, v. 5, p. 319-358, text fig. 1-26, pl. 15. (Contains bibliography listing large part of author's works on rudistids.)
- Driscoll, E. G.**
- (282a) 1966, *Morphology and evolution of certain Paleozoic Nuculanidae from the midcontinental United States*: Narod. Musea Praze, Sborník, v. 22B, p. 1-26.
- (282b) 1966, *Hinge structure and shell perforation in an upper Paleozoic nuculoid genus*: Malacol. Soc. London, Proc., v. 37, p. 93-96.
- Duméril, A. M. C.**
- (283) 1806, *Zoologie analytique, ou méthode naturelle de classification des animaux*: xxxii + 344 p. (Paris).
- Eagar, R. M. C.**
- (284) 1946, *The hinge of certain non-marine lamellibranchs from the Lenisulcata Zone of the Coal Measures*: Geol. Mag., v. 83, p. 1-19, text fig. 1-7, pl. 1-2.
- (285) 1948, *Variation in shape of shell with respect to ecological station. A review dealing with the recent Unionidae and certain species of the Anthracosiidæ in Upper Carboniferous times*: Royal Soc. Edinburgh, Proc. sec. B, v. 63, p. 130-148, text fig. 1-9.
- (286) 1952, *Growth and variation in the non-marine lamellibranch fauna above the Sand Rock Mine of the Lancashire Millstone Grit*: Geol. Soc. London, Quart. Jour., v. 107, p. 339-373, text fig. 1-16.
- (287) 1956, *Naming non-marine lamellibranchs*: Syst. Assoc., Publ. no. 2, p. 111-116, 1 text fig.
- Eames, F. E.**
- (288) 1951, *A contribution to the study of the Eocene in Western Pakistan and Western India. B. The description of the Lamellibranchia from standard sections in the Rakhi Nala and Zinda Pir areas of the Western Punjab and in the Kohat district*: Royal Soc. London, Philos. Trans., ser. B, no. 627, v. 235, p. 311-482.
- Ebersin, A. G.**
- (289) 1965, *Sistema i flogeniya solonovatovodnykh kardiid*: Akad. Nauk SSSR, Zool. Inst., Mollyuski—voprosi; Teoreticheskoy i prikladnoy malakologii. Tezisy dokladov., Sbornik vtoroy [Second Conference on Study of Mollusca: Problems of theoretical and applied malacology: Abstr., v. 2], p. 11-25, 5 text fig. [System and phylogeny of brackish-water Cardiidae.]
- Eichwald, E. I. von**
- (290) 1860, *Lethaea Rossica ou paléontologie de la Russie*: v. 1, pt. 2, xix + p. 269-1657 (Atlas, 1859), Schweizerbart (Stuttgart).
- (291) 1866-69, *Lethaea Rossica, ou paléontologie de la Russie, v. 2, Période moderne*: 1304 p., atlas of 40 pl. (Stuttgart).
- Elias, M. K.**
- (292) 1956, *Upper Mississippian and Lower Pennsylvanian formations of south central Oklahoma*: in Petroleum geology of Southern Oklahoma, v. 1, p. 56-134, 6 pl., Am. Assoc. Petrol. Geol. (Tulsa).
- (293) 1957, *Late Mississippian fauna from the Redoak Hollow Formation of southern Oklahoma, Part 3*: Jour. Paleontology, v. 3, p. 737-784, 1 text fig., pl. 89-97.
- Ellis, A. E.**
- (294) 1940, *The identification of the British species of Pisidium*: Malacol. Soc. London, Proc., v. 24, p. 44-88, pl. 1-6.

- (295) 1947, *Freshwater bivalves* (Mollusca [Unionacea]): Linnean Soc. London, Synopses Brit. Fauna, no. 5, p. 1-39, 17 text fig., pl. 1-12.
- (296) 1956, *Proposed use of the plenary powers to validate the currently accepted usage of the generic name "Unio" Philippon, 1788 (Class Pelecypoda)*: Bull. Zool. Nomencl., v. 11, p. 337-343.
- Emmons, Ebenezer**
(297) 1842, *Geology of New York*: Nat. History New York, pt. 2, 437 p. (Albany).
- Etheridge, Robert**
(298) 1872, *Description of the Palaeozoic and Mesozoic fossils of Queensland*: Geol. Soc., London, Quart. Jour., v. 28, p. 317-350, pl. 13-25.
(299) 1888, *The invertebrate fauna of the Hawkesbury - Wianamatta Series (Beds above the productive coal measures) of New South Wales*: Geol. Survey New South Wales, Mem., Paleontology, no. 1, p. 1-21, pl. 1-2.
- , & **Dun, W. S.**
(300) 1906, *A monograph of the Carboniferous and Permo-Carboniferous Invertebrata of New South Wales, vol. 2. Pelecypoda, part 1. The Paleopectens*: Geol. Survey New Wales, Mem., Palaeont. no. 5, 39 p., 16 pl.
- Fabre, S. J.**
(301) 1942, *Catalogue des types de la collection Philippe Matheron*: Musée Histoire Nat. Marseille, Bull., v. 2, no. 2, p. 121-140.
- Felix, J.**
(301a) 1891, *Versteinerungen aus der mexicanischen Jura- und Kreideformation*: Palaeontographica, v. 37, p. 140-194, pl. 22-30.
- Férussac, A. E. de**
(302) 1822, *Tableaux systématiques des animaux mollusques*: 111 p. (Paris and London).
- Finlay, H. J.**
(303) 1926, *A further commentary on New Zealand Mollusca systematics*: New Zealand Inst., Trans., v. 57, p. 320-485.
- , & **Marwick, John**
(304) 1937, *The Wangaloan and associated molluscan faunas of Kaitangata-Green Island subdivision*: New Zealand Geol. Survey, Paleont. Bull., no. 15, 1140 p., 17 pl.
- Fischer, P. H.**
(305) 1880-87, *Manuel de conchyliologie et de paléontologie conchyliologique*: xxv + 1369 p., 23 pl., F. Savy (Paris).
- , & **Crosse, Hippolyte**
(306) 1880-1902, *Études sur les mollusques terrestres et fluviatiles du Mexique et du Guatemala*: Zool. Histoire Faune Amérique Centrale et Mexique, Rech., v. 7, pt. 2, 731 p., pl. 32-72.
- Fitch, J. E.**
(307) 1953, *Common marine bivalves of California*: California Dept. Fish & Game, Fish Bull. no. 90, 102 p., 63 text fig.
- Fleming, C. A.**
(308) 1957, *Lower Devonian Pelecypoda from Reefton, New Zealand*: Royal Soc. New Zealand, Trans., v. 85, pt. 1, p. 135-140, text fig. 1-4, pl. 14-15.
(309) 1959, *Notes on New Zealand Recent and Tertiary mussels (Mytilidae)*: Same, v. 37, pt. 1 & 2, p. 165-178, text fig. 1-3, pl. 13-14.
- Fleming, John**
(310) 1822, *The philosophy of zoology*: v. 2, 618 p., 5 pl. (Edinburgh).
(311) 1828, *A history of British animals, exhibiting the descriptive characters and systematical arrangement of the genera and species of quadrupeds, birds, reptiles, fishes, Mollusca and Radiata of the United Kingdom; including the indigenous, extirpated, and extinct kinds; together with periodical and occasional visitants*: xxiii + 554 p. (Edinburgh).
- Fletcher, H. O.**
(312) 1945, *A new genus Glyptoleda and a revision of the genus Nuculana from the Permian of Australia*: Australian Museum, Records, v. 21, p. 293-312, pl. 19-22.
- , & **Dun, W. S.**
(313) 1966, *Contributions on the Permo-Carboniferous Aviculopectinidae of New South Wales*: Australian Museum, Records, v. 42, p. 1-34, pl. 1-16.
- Foerste, A. F.**
(314) 1895 (1893), *Fossils of the Clinton group in Ohio and Indiana*: Geol. Survey Ohio, Rept., v. 7, pt. 2, p. 516-601, 14 text fig., pl. 25-37a.
(315) 1909, *Fossils from the Silurian formations of Tennessee, Indiana and Illinois*: Denison Univ., Sci. Lab., Bull., v. 14, p. 61-107, pl. 1-4.
(316) 1914, *Lorraine faunas of New York and Quebec*: Same, Bull., v. 17, p. 247-339.
(317) 1916, *Notes on Cincinnati fossil types*: Same, Bull., v. 18, p. 285-355.
(318) 1924, *Upper Ordovician faunas of Ontario and Quebec*: Geol. Survey Canada, Mem. 138, 255 p., 46 pl.
(319) 1931, *The Silurian fauna of Kentucky*: in *The Paleontology of Kentucky*, W. R. Jillson (ed.), Kentucky Geol. Survey, ser. 6, v. 36, p. 167-212, pl. 17-26.
- Franc, André**
(320) 1949, *Unionidae d'Afrique française recueillies par Th. Monod*: Jour. Conchyliologie, v. 89, p. 157-187, 10 text fig.

- (321) 1960, *Classe de bivalves*: in *Traité de zoologie*, P.-P. Grassé (ed.), p. 1845-2219 (Paris).
- Frech, Fritz**
- (322) 1899, *Über Mecynodon und Myophoria*: *Deutsch. Geol. Gesell., Zeitschr.*, v. 41, p. 127-138, pl. 11.
- (323) 1891, *Die devonischen Aviculiden Deutschlands, ein Beitrag zur Systematik und Stammesgeschichte der Zweischaler*: *Geol. Spezialk. Preuss., Abhandl.*, v. 9, 253 p., 23 text fig., 5 pl., atlas of 18 pl.
- Frierson, L. S.**
- (324) 1911, *Notes on oriental Unionidae*: *Nautilus*, v. 24, p. 97-98.
- (325) 1914, *Observations on the genus Symphynota, Lea*: Same, v. 28, p. 40.
- (326) 1914, *Remarks on classification of the Unionidae*: Same, v. 28, p. 6-8.
- (327) 1922, *Observations on the genus Margaritana with a new sub-genus*: Same, v. 36, p. 42-44.
- (328) 1927, *A classified and annotated check list of the North American najades*: 111 p. (Shreveport, Louisiana).
- Fritz, M. A.**
- (329) 1951, *The pelecypod genus Vlasta in the Ordovician of North America*: *Jour. Paleontology*, v. 25, p. 696-698, 4 text fig.
- Frizzell, D. L.**
- (329a) 1936, *Preliminary reclassification of veneracean pelecypods*: *Musée Royal Histoire Nat. Belgique, Bull.*, v. 12, no. 34, 84 p.
- Fuchs, Alexander**
- (330) 1915, *Der Hunsrückschiefer und die Unterkoblenzschichten am Mittelrhein (Loreleigegend). I. Teil. Beitrag zur Kenntnis der Hunsrückschiefer und Unterkoblenzfauna der Loreleigegend*: *K. Preuss. Geol. Landesanst., Abhandl., new ser.*, no 79, 79 p., 18 pl.
- (331) 1920, *Beitrag zur Kenntnis der Devonfauna der Verse und der Hobracker Schichten des sauerländischen Faciesgebietes*: Same, *Jahrb.*, v. 39, pt. 1 (for 1918), p. 58-95, pl. 5-9.
- Fuchs, Theodor**
- (332) 1878, *Die geologische Beschaffenheit der Landenge von Suez*: *K. Akad. Wiss. Wien, Denkschr., Math.-Natur. Classe*, v. 38, part 2, p. 25-42, pl. 1-3.
- Futterer, Karl**
- (332a) 1892, *Die Oberen Kreidebildungen der Umgebung des Lago di Santa Croce in den Venetianer Alpen*: *Paläont. Abhandl., Jena*, new ser. 2, pt. 1, p. 1-123, 25 text fig., pl. 1-12.
- Gabb, W. M.**
- (333) 1864, *Description of the Cretaceous fossils*: *Geol. Survey California, Palaeont.*, v. 1, sec. 4, p. 55-243, pl. 9-32.
- Gardner, J. A.**
- (334) 1916, *Mollusca, Brachiopoda, and Vermes*: in W. B. Clark, *Systematic paleontology of the Upper Cretaceous deposits of Maryland*, *Maryland Geol. Survey, Upper Cretaceous vol.*, p. 518.
- Gill, E. D., & Darragh, T. A.**
- (335) 1963, *Evolution of the Zenatiinae (Macrtridae: Lamellibranchiata)*: *Royal Soc. Victoria, Proc.*, v. 77, pt. 1, p. 177-190, 1 text fig., pl. 28-31.
- Gillet, Suzette**
- (336) 1922, *Étude de la faune lamellibranches du calcaire à spatangues (Hauterivien supérieur)*: *Soc. Sci. Historiques Nat. l'Yonne, Bull.*, v. 75, pt. 2, p. 45-108, pl. 1-5.
- (337) 1924, *Études sur les lamellibranches néocomiens*: *Soc. Géol. France, Mém., new sér.*, v. 1, pt. 3-4, *Mém.* 3, p. 1-224, pl. 1-2.
- Girty, G. H.**
- (338) 1909, *The Manzano Group of the Rio Grande Valley*: *U.S. Geol. Survey, Bull.* 389, p. 1-120, pl. 6-12.
- (339) 1911, *On some new genera and species of Pennsylvanian fossils from the Wewoka formation of Oklahoma*: *New York Acad. Sci., Ann.*, v. 21, p. 99-156.
- Goldfuss, G. A.**
- (340) 1820, *Handbuch der Zoologie*: v. 3, no. 1, xlvi + 696 p., 2 pl. (Nürnberg).
- (341) 1833-40, *Petrefacta Germaniae*: pt. 2, (a) p. 1-68, pl. 72-96 (1833); (b) lief. 5, p. 69-140, pl. 97-121 (1836); (c) lief. 6, p. 141-224, pl. 122-146 (1837); (d) lief. 7, p. 225-312, pl. 147-165 (1840), Arnz (Düsseldorf).
- Gosselet, J., Barrois, C., Leriche, M., Crépin, A., Pruvost, P., & Dubois, G.**
- (342) 1920, *Description de la faune siluro-dévonnienne de Liévin*: *Soc. Géol. du Nord, Mém.*, v. 6, pt. 2, p. 71-225, pl. 10-17.
- Gould, A. A.**
- (343) 1844, *Descriptions of two species of Anodon, from the river Salween, in British Burmah, sent by Rev. F. Mason*: *Boston Soc. Nat. History, Proc.*, v. 1, p. 160-161.
- Graham, A.**
- (344) 1949, *The molluscan stomach*: *Royal Soc. Edinburgh, Trans.*, v. 61, p. 737-778.
- Grant, U. S. IV, & Gale, H. R.**
- (345) 1931, *Catalogue of the marine Pliocene and Pleistocene Mollusca of California*: *San Diego Soc. Nat. History, Mem.*, v. 1, 1036 p., 32 pl.

Gray, J. E.

- (346) 1821, *A natural arrangement of Mollusca, according to internal structure*: London Med. Repos., v. 15, p. 229-239.
- (347) 1840, *Synopsis of the contents of the British Museum*: 42nd. edit., 370 p., Brit. Museum (London).
- (348) 1847, *A list of the genera of Recent Mollusca, their synonyms and types*: Zool. Soc. London, Proc., v. 15, p. 129-219.

Gregorio, Antonio de

- (349) 1914, *Su taluni moluschi di acqua dolce di America*: Naturalista Siciliano, v. 22, p. 32-72, pl. 3-12.

Grobben, Carl

- (350) 1892, *Beiträge zur Kenntniss der Baues von Cuspidaria (Neaera) cuspidata Olivi, nebst Betrachtungen über das System der Lamelibranchiaten*: Zool. Inst. Univ. Wien, Arbeiten, v. 4, p. 101-146, 4 pl.
- (351) 1894, *Zur Kenntniss der Morphologie, der Verwandtschaftsverhältnisse und des Systems der Mollusken*: Akad. Wiss. Wien, Sitzungsber., v. 103, p. 61-86.

Guillaume, Louis

- (352) 1928, *Revision des posidonomyes jurassiques*: Soc. Géol. France, Bull., sér. 4, v. 27, p. 217-234, pl. 10.

Haas, Fritz

- (353) 1910, *Pseudunio, neues Genus für Unio sinuatus Lam.*: Deutsch. Malakozool. Gesell., Nachr., v. 42, p. 181-183.
- (354) 1910-20, *Die Unioniden*: Martini-Chemnitz, Ill. Conch. Cab. (2), v. 9, Abt. 2, p. 1-344, pl. 1-73 (unfinished).
- (355) 1912, *Neue Najaden*: Deutsch. Malakozool. Gesell., Nachr., v. 42, p. 97-103.
- (356) 1913, *Neue Najadengattungen*: Same, v. 45, p. 33-38.
- (357) 1914, *Prohyriopsis, neue Gattung für Unio stolatus Mart.*: Same, v. 46, p. 76-78.
- (358) 1920, *Die Neumayrschen Najaden aus der Ausbeute des Grafen Széchenyi*: Senckenbergiana, v. 2, p. 146-151.
- (359) 1929, *Bemerkungen über mittelamerikanische Najaden*: Same, v. 11, p. 310-344, 21 text fig.
- (360) 1929-56, *Bivalvia*: in H. G. Bronn, Klassen und Ordnungen des Tierreichs, v. 3, Abt. 3, pt. 1 (1829-35), p. 1-984, text fig. 1-563; pt. 2 (1937-56), p. 1-466, text fig. 1-189 (Leipzig).
- (361) 1931, *Versuch einer kritischen Sichtung der südamerikanischen Najaden an Hand der Sammlung des Senckenberg-Museums. III.*: Senckenbergiana, v. 13, p. 87-110, text fig. 33-37.
- (362) 1936, *Binnen-Mollusken aus Inner-Afrika, hauptsächlich gesammelt von Dr. F. Haas*

während der Schomburg-Expedition in den Jahren 1931-32: Senckenberg. Naturf. Gesell., Abhandl., v. 431, p. 1-156, 17 text fig., 8 pl.

- (363) 1940, *A tentative classification of the Palearctic unionids*: Field Museum Nat. History, Zool. ser., v. 24, p. 115-141.
- (364) 1955, *Unionella Haas 1913, vorweggenommen durch Unionella Etheridge, 1888*: Arch. Molluskenkunde, v. 84, p. 212.

Habe, Tadashige

- (365) 1951-53, *Genera of Japanese shells; Pelecypoda and Scaphopoda* (in Japanese): 326 p., 770 text fig. (Kyoto).
- (366) 1961, *Coloured illustrations of shells of Japan (II)*: 182 p., 66 pl., Hoikusha Publ. (Osaka, Japan).
- (367) 1965, *Family Solenidae in Japan and its adjacent areas*: Venus, Japan. Jour. Malacol., v. 23, no. 4, p. 188-197, pl. 12-13.

Hajkr, Oldřich, Prantl, Ferdinand, & Růžička, Bohuslav

- (368) 1961, *Biometrical study of the outline of some mytiloid pelecypods*: Acta Museum Nat. Prague, p. 81-95, pl. 7.

Hall, James

- (367) 1847, *Palaeontology*: Geol. Survey of New York, v. 1, xxiii + 338 p., 87 pl. (Albany).
- (370) 1850, *Description of new species of fossils, and observations upon some other species previously not well known, from the Trenton Limestone*: N. Y., 3rd Ann. Rept. Regents Univ., p. 173-183, pl. 1-5.
- (371) 1852, *Palaeontology, v. 2, containing descriptions of the organic remains of the lower middle division of the New-York system (equivalent in part to the middle Silurian rocks of Europe)*: Nat. History of New York, 362 p., text fig., 85 pl. (Albany).
- (372) 1859, *Palaeontology, v. 3, pt. 1, containing descriptions and figures of the organic remains of the lower Helderberg group and the Oriskany sandstone*: Geol. Survey New York, 1855-59, 532 p., text fig. (Albany).
- (373) 1861, *Containing descriptions and figures of the organic remains of the lower Helderberg group and the Oriskany sandstone*: Same, v. 3, 1855-1859, pt. 2, 120 pl.
- (374) 1867, *Account of some new or little known species of fossils from rocks of the age of the Niagara group, 2. Descriptions of some new fossils from the Niagara group*: N.Y. State Cab. Nat. History, 20th Ann. Rept., p. 308-401, text fig., pl. 10(1)-23(14).
- (375) 1869, *Preliminary notice of the lamelibranchiate shells of the upper Helderberg, Hamilton and Chemung groups, with oth-*

- ers from the Waverly sandstones: pt. 2, 97 p., N.Y. State Museum (Albany).
- (376) 1883, *Lamellibranchiata, plates and explanations*: Nat. History New York, Paleont., v. 5, pt. 1, 20 p., 79 pl.
- (377) 1884, *Preliminary notice of the lamellibranchiate shells of the upper Helderberg, Hamilton, and Chemung groups, Pt. 1*: N.Y. State Museum Nat. History, Ann. Rept. 35, p. 215-406, 11 pl.
- (378) 1884, *Lamellibranchiata I, descriptions and figures of the Monomyaria of the upper Helderberg, Hamilton, and Chemung Groups*: N.Y. Geol. Survey, Paleont., v. 5, pt. 1, p. 1-268, pl. 1-33, 81-92.
- (379) 1885, *Lamellibranchiata II, descriptions and figures of the Dimyaria of the upper Helderberg, Hamilton, Portage and Chemung Groups*: Same, v. 5, pt. 1, p. 269-562, pl. 34-80, 93-96.
- , & Meek, F. B.
- (380) 1856, *Descriptions of new species of fossils from the Cretaceous formations of Nebraska*: Am. Acad. Arts & Sci., Mem., new ser., v. 5, p. 379-411.
- , & Whitfield, R. P.
- (381) 1875, *Fossils of the Hudson River Group (Cincinnati formations)*: Geol. Survey Ohio, Rept., v. 2, pt. 2, p. 67-161, pl. 1-13.
- Hanley, S. C. T.**
- (382) 1842-56, *An illustrated and descriptive catalogue of Recent bivalve shells*: 392 p., 24 pl., William & Norgate (London).
- (383) 1860, *On some species of Nuculaceae in the collection of Hugh Cuming, Esq.*: Zool. Soc. London, Proc., pt. 28, p. 370-372.
- , & Theobald, W.
- (384) 1870-76, *Conchologia Indica: illustrations of the land and freshwater shells of British India*: 65 p., 160 pl. (London).
- Hanna, M. A.**
- (385) 1927, *An Eocene invertebrate fauna from the La Jolla Quadrangle, California*: Univ. California, Geol. Sci. Bull., v. 16, p. 247-398, pl. 24-57.
- Hannibal, Harold**
- (386) 1912, *A synopsis of the Recent and Tertiary freshwater Mollusca of the California province, based upon an ontogenetic classification*: Malacol. Soc. London, Proc., v. 10, p. 112-165, pl. 5-8.
- Harris, G. D.**
- (387) 1896, *The Midway Stage*: Bull. Am. Paleontology, v. 1, no. 4, p. 115-272.
- (388) 1920, *The genera Lutetia and Alveinus, especially as developed in America*: Palaeont. Americana, v. 1, no. 2, p. 105-118, 17 pl.
- , & Hodson, Floyd
- (388a) 1922, *The rudistids of Trinidad*: Palaeont. Americana, v. 1, no. 3, p. 119-162, pl. 18-28.
- , & Palmer, K. V. W.
- (389) 1946, *The Mollusca of the Jackson Eocene of the Mississippi embayment*: Bull. Am. Paleontology, v. 30, no. 117, p. 1-206.
- Hatai, Kotoru, & Nisiyama, Syozo**
- (390) 1952, *Check list of Japanese Tertiary marine Mollusca*: Tohoku Univ., Sci. Rept., ser. 2, spec. vol., no. 3, 464 p.
- Hayami, Itaru**
- (391) 1958, *Taxonomic notes on Cardinia with description of a new species from the Lias of western Japan*: Tokyo Univ., Jour. Fac. Sci., v. 2, no. 11, p. 115-130, pl. 11.
- (392) 1959, *Late Jurassic isodont and myacid pelecypods from Makito, central Japan*: Japanese Jour. Geology & Geography, v. 30, p. 151-167, pl. 13.
- (393) 1960, *Jurassic inoceramids in Japan*: Univ. Tokyo, Jour. Fac. Sci., sec. 2, v. 12, pt. 2, p. 277-328.
- (394) 1961, *On the Jurassic pelecypod faunas in Japan*: Same, sec. 2, v. 12, pt. 2, p. 243-343.
- (395) 1965, *Lower Cretaceous marine pelecypods of Japan, Part 1: Kyushu Univ.*, Mem. Fac. Sci., ser. D, Geol., v. 15, no. 2, p. 221-349, pl. 27-52.
- Hedley, Charles**
- (396) 1901, *Some new or unfigured shells*: Australian Museum, Records, v. 4, p. 22-27.
- (397) 1902-03, *Scientific results of the trawling expedition of H. M. C. S. "Thetis", Mollusca*: Same, Mem. 4, pt. 5 (1902), p. 285-324; pt. 6 (1903), p. 325-402.
- (398) 1906, *Results of dredging on the continental shelf of New Zealand*: New Zealand Inst., Trans., v. 38, p. 68-76.
- (399) 1914(1915), *Studies on Australian Mollusca, Part 12*: Linnean Soc. New South Wales, Proc., v. 39, p. 695-755, pl. 77-85.
- Heinz, R.**
- (400) 1932, *Aus der neuen Systematik der Inoceramen*: Mitteil. Mineral.- Geol. Staatsinst. Hamburg, v. 13, 26 p.
- Henderson, Junius**
- (401) 1935, *Fossil non-marine Mollusca of North America*: Geol. Soc. America, Spec. Paper 3, viii+313 p.
- Herrmannsen, A. N.**
- (402) 1846-52, *Indicis generum malacozoorum primordia*: v. 1, xxvii+637 p. (1846-47); v. 2, xxiv-xlii+717 p. (1847-49); suppl., v+140 p. (1852) (Cassel).

Hertlein, L. G., & Strong, A. M.

- (403) 1940, *Mollusks from the West Coast of Mexico and Central America*: Zoologica, N.Y. Zool. Soc., v. 25, no. 4, p. 418-420.

Hertwig, C. W. T. R.

- (404) 1895, *Lehrbuch der Zoologie*: 3te Auflage, xii+599 p. (Jena).

Heude, P. M.

- (405) 1878-86, *Conchyliologie fluviatile de la province de Nanjing et de la Chine centrale*: 72 pl. with text (Paris).

Hicks, Henry

- (406) 1873, *On the Tremadoc rocks in the neighbourhood of St. David's, South Wales and their fossil contents*: Geol. Soc. London, Quart. Jour., v. 29, p. 39-52, pl. 3-5.

Hind, Wheelton

- (407) 1896-1905, *A monograph of the British Carboniferous Lamellibranchiata*: Palaeontograph. Soc., (a) v. 1 (1896-1900), 486 p., 54 pl.; (b) v. 2 (1901-05), 222 p., 25 pl.

- (408) 1910, *The lamellibranchs of the Silurian rocks of Girvan*: Royal Soc. Edinburgh, Trans., v. 47, p. 479-548, pl. 1-5.

Hirase, Shintaro, & Taki, Iwao

- (409) 1951, *A handbook of illustrated shells— from Japanese Islands and their adjacent territory*: 134 pl., Maruzen Co., Ltd. (Tokyo).

Hitchcock, Edward

- (410) 1865, *Supplement to the ichnology of New England. A report . . . in 1863*: 96 p., 20 pl. (Boston). (Appendix B.—Descriptive catalogue of the specimens in the Hitchcock Collection of Amherst College.)

Hörnes, Moritz

- (411) 1851-70, *Die fossilen Mollusken des Tertiärbeckens von Wien*: K. K. Geol. Reichsanst. Wien, Abhandl., v. 3, pt. 1, 736 p., 52 pl., 1 map; pt. 2, 479 p., 85 pl.

Hörnes, Rudolf

- (412) 1882, *Ueber die Analogien des Schlossapparates von Megalodus, Dicerias und Caprina*: Geol. Reichsanst. Wien, Verhandl., p. 179-181.

Hofmann, Károly

- (412a) 1913, *Die Lamellibranchiaten der mittel-neokömen Schichten des Mecsekgebirges*: K. Ungar. Geol. Reichsanst., Jahrb., v. 20, pt. 5, p. 211-252, 5 text fig., pl. 5-7.

Holzappel, Eduard

- (413) 1882, *Die Goniatiten-Kalke von Adorf in Waldeck*: Palaeontographica, v. 28, no. 6, p. 225-262, pl. 44-49.

- (414) 1889, *Die Cephalopoden-führenden Kalke des unteren Carbon von Erdbach-Breitscheid bei Herborn*: Paläont. Abhandl., v. 5, new ser., v. 1, no. 1, p. 1-74, 8 pl.

- (415) 1889, *Die Mollusken der Aachener Kreide*. 3. Classe: Lamellibranchiata: Palaeontographica, v. 35, p. 139-263, pl. 8-29.

Horný, Radvan

- (416) 1960, *On the phylogeny of the earliest pelecypods (Mollusca)*: Ústřed. Ústavu Geol., Českosl. Věstník, v. 35, p. 479-482.

Howse, Richard

- (417) 1878, *Preliminary notice of the occurrence of Archanodon (Anodonta) južesi (Forbes) in the Lower Carboniferous rocks of north Northumberland*: Nat. History Trans. Northumberland, Durham and Newcastle-on-Tyne, v. 7, p. 173-175, pl. 14.

Hüffner, Ernst

- (418) 1916, *Beiträge zur Kenntnis des Devons von Bithynien. Auf Grund der Aufsammlungen von Herrn Dr. Endriss bearbeitet*: K. Preuss. Geol. Landesanst., Jahrb., v. 37, pt. I, p. 260-357, 1 text fig., pl. 28-30.

Hussey, R. C.

- (419) 1926, *The Richmond Formation of Michigan*: Univ. Michigan, Museum Geology, Contrib., v. 2, art. 8, p. 113-187, 12 text fig., pl. 1-11.

Hutton, F. W.

- (420) 1893, *The Pliocene Mollusca of New Zealand*: Linnean Soc. New South Wales, Macleay Mem. vol., p. 35-92, pl. 6-9.

Hyde, J. E.

- (421) 1953, *The Mississippian formations of central and southern Ohio. Edited by Mildred Fisher Marple*: Ohio, Div. Geol. Survey, Bull. 51, 355 p., 54 pl.

Ichikawa, Koichiro

- (422) 1949, *Trigonucula (nov.) and other taxodont pelecypods from the Upper Triassic of the Sakawa Basin in Shikoku, Japan*: Japanese Jour. Geology & Geography, v. 21, p. 267-272, pl. 10.

- (423) 1958, *Zur Taxionomia und Phylogenie der triadschen "Pteriidae" (Lamellibranch.)*: Palaeontographica, v. 3, Abt. A, p. 131-212, pl. 21-24.

———, & Maeda, Yasuo

- (424) 1958, *Late Cretaceous pelecypods from the Izumi group, part 1, Cucullaeidae (Pleurogrammatodon, nov., Nanonavis, and Indogrammatodon)*: Osaka City Univ., Jour. Inst. Polytech., ser. G, v. 3, p. 61-74.

- (425) 1958, *Late Cretaceous pelecypods from the Izumi group, part 2, Orders Taxodontida, Prionodontida, Dysodontida, Desmodontida, Adapodontida*: Same, ser. G, v. 4, p. 71-122.

- (426) 1963, *Late Cretaceous pelecypods from the Izumi group, part 3, Order Heterodontida (1)*: Same, Jour. Geosci., v. 7, art. 5, p. 113-145.

Ihering, Hermann von

- (427) 1891-92, *Anodonta und Glabaris*: Zool. Anzeiger, v. 14 (1891), p. 474-484; v. 15 (1892), p. 1-5, 1 text fig.
- (428) 1893, *Najaden von S. Paulo und die geographische Verbreitung der Süßwasserfaunen von Südamerika*: Arch. Naturgesch., v. 59, v. 1, p. 45-140, pl. 3-4.
- (429) 1900, *On the South American species of Mytilidae*: Malacol. Soc. London, Proc., v. 4, pt. 2, p. 84-98.

Imlay, Ralph

- (430) 1959, *Succession and speciation of the pelecypod Aucella*: U.S. Geol. Survey, Prof. Paper 314G, p. 155-169.

Iredale, Tom

- (431) 1924, *Results from Roy Bell's molluscan collections*: Linnean Soc. New South Wales, v. 49, p. 179-278, pl. 33-36.
- (432) 1925, *Mollusca from the continental shelf of eastern Australia*: Australian Museum Records, v. 14, p. 243-270, pl. 41-43.
- (433) 1929, *Mollusca from the continental shelf of eastern Australia, Part 2*: Same, v. 17, no. 4, p. 157-189, pl. 38-41.
- (434) 1934, *The fresh-water mussels of Australia*: Austral. Zool., v. 8, p. 57-78, pl. 3-6.
- (435) 1936, *Queensland cobra or shipworms*: in *Destruction of timber by marine organisms in the port of Brisbane*, Queensland Forest Service, Bull. no. 12, p. 31-44, pl. 1-2.
- (436) 1936, *Australian molluscan notes nr. 2*: Australian Museum Records, v. 19, no. 5, p. 267-340.
- (437) 1939, *Great Barrier Reef expedition 1928-1929*: British Museum (Nat. History), Sci. Repts., v. 5, no. 6 (Mollusca, pt. 1), p. 209-425, 7 pl.
- (438) 1951, *Recent palaeontology*: Austral. Zool., v. 11, pt. 4, p. 347-350.

Isberg, Orvar

- (439) 1934, *Studien über Lamellibranchiaten des Leptaenakalkes in Dalarna, Beitrag zu einer Orientierung über die Muschelfauna im Ordovicium und Silur*: 492 p., 9 text fig., 32 pl. (Lund).

Jack, R. L., & Etheridge, Robert

- (440) 1892, *The geology and palaeontology of Queensland and New Guinea*: xxx+768 p., atlas of 68 pl.

Jackson, R. T.

- (441) 1890, *Phylogeny of the Pelecypoda; the Aviculidae and their allies*: Boston Soc. Nat. History, Mem. IV, p. 277-400, pl. 23-30.

Jacob, Charles

- (442) 1939, *La vie et l'oeuvre de H. Douvillé, 1846-1937*: Soc. Géol. France, Bull., sér. 5,

v. 9, p. 321-591 (with complete bibliography of Douvillé's papers).

Jameson, H. L.

- (443) 1901, *On the identity and distribution of the mother-of-pearl oysters; with a revision of the subgenus Margaritifera*: Zool. Soc. London, Proc., v. 1, p. 372-394, text fig. 92-95.

Jaworski, Erich

- (444) 1913, *Beiträge zur Stammesgeschichte der Austern*: Zeitschr. Induk. Abstamm. u. Vererbungs., v. 9 (1913), p. 192-215, pl. 6-7.
- (445) 1914, *Beiträge zur Kenntnis der Lias-Volen Südamerikas und der Stammesgeschichte der Gattung Vola*: Paläont. Zeitschr., v. 1, p. 273-320.
- (446) 1922, *Die marine Trias in Südamerika*: Neues Jahrb., Beil.-Bd. 47, p. 93-200, pl. 4-6.
- (447) 1951, *Review of H. B. Stenzel, Nomenclatural synopsis of supraspecific groups of the family Ostreidae*: Zentralbl. Geologie & Paläontologie, pt. 2, Hist. Geol. & Paläont. 1951, no. 3, p. 648-650.

Jeffreys, J. G.

- (448) 1876, *New and peculiar Mollusca of the Pecten, Mytilus and Arca families procured in the "Valorous" Expedition*: Annals & Mag. Nat. History, ser. 4, v. 18, p. 424-436.
- (449) 1879, *On the Mollusca procured during the "Lightning" and "Porcupine" Expeditions, 1868-70 (Part II)*: Zool. Soc. London, Proc., v. 1879, p. 553-588, pl. 45, 46.

Jensen, A. S.

- (450) 1912, *Lamellibranchiata*: Danish Ingolf-Expedition, Vol. II: 5 (pt. 1), 119 p., 4 pl. (Copenhagen).

Johnson, C. W.

- (451) 1934, *List of marine Mollusca of the Atlantic and Gulf Coasts*: rev. edit., 236 p., 45 pl., Houghton Mifflin Co. (Boston).

Jourdy, Emile

- (452) 1924, *Histoire naturelle des exogyres*: Ann. Paléontologie, p. 1-104, pl. 1-11.

Jousseume, F.-P.

- (453) 1897, *Description d'une coquille nouvelle*: Naturaliste, Ann. 19, no. 257, p. 265.

Jukes-Browne, A. J.

- (454) 1905, *A review of the genera of the family Mytilidae*: Malacol. Soc. London, Proc., v. 6, pt. 4, p. 211-224.
- (455) 1908, *On the genera of Veneridae represented in the Cretaceous and older Tertiary deposits*: Same, Proc., v. 8, p. 148-177, pl. 6.
- (455a) 1914, *A synopsis of the Veneridae. Pts. 1 and 2*: Same, v. 8, p. 58-94.

Kautsky, Fritz

- (456) 1939, *Die Erycinen des niederösterreichischen Miocäns*: Naturhist. Museums Wien, Ann., v. 50, p. 584-671, pl. 19-22.

Kawaguti, Siro, & Baba, Kikutarô

- (457) 1959, *A preliminary note on a two-valved sacoglossan gastropod, Tamonovalva limax, n.gen., n.sp., from Tamano, Japan*: Okayama Univ., Biol. Jour., v. 5, p. 177-184.

Keen, A. M.

- (458) 1937, *Nomenclatural units of the pelecypod family Cardiidæ*: Musée Royal Histoire Nat. Belgique, v. 13, no. 7, p. 1-22.
- (459) 1950, *Notes on the history of Nemocardium (Family Cardiidæ)*: Jour. Conchylologie, v. 90, no. 1, p. 23-29.
- (460) 1951, *Outline of a proposed classification of the pelecypod family Cardiidæ*: Conch. Club South. California, Min., p. 6-8.
- (461) 1951, *Outline of a proposed classification of the pelecypod family Veneridæ*: Same, p. 1-10.
- (462) 1958, *Marine shells of tropical West America*: 624 p., Stanford University Press (Stanford, Calif.).
- (463) 1959, *Some side notes on "Sea shells of tropical North America"*: The Veliger, v. 2, no. 1, p. 1-3.
- (464) 1963, *Marine molluscan genera of western North America. An illustrated key*: 126 p., Stanford University Press (Stanford, Calif.).
- , & **Frizzell, D. L.**
- (465) 1939, *Illustrated key to west North American pelecypod genera*: p. 3-28, Stanford Univ. Press (Stanford, Calif.).

Keferstein, Wilhelm

- (466) 1857, *Ueber einige deutsche devonische Conchiferen aus der Verwandtschaft der Trigoniacen und Carditaceen*: Deutsch. Geol. Gesell., Zeitschr., v. 9, p. 148-162, pl. 4.

Kegel, Wilhelm

- (467) 1925, *Bemerkungen über die obersilurische Zweischaler-Gattung Rhombopteria und ihre Verwandten*: Centralbl. Mineralogie, Geologie u. Paläontologie, Jahrg. 1925, Abt. B, p. 295-303, text fig.

Keller, Alexandre

- (468) 1933, *Sur quelques rudistes du Djebel Ansaryeh et de l'Amanus (Syrie septentrionale)*: in Contrib. Étude Géol. de la Syrie Septentrionale; Notes et Mém. Haut Com. Rep. France en Syrie et Liban, Serv. Trav. Pub. Sect. Étude Géol., p. 45-52, text fig. 12-13, pl. 3-7.

Kimura, Toshio

- (469) 1951, *Some pectinids and limids from the Jurassic Torinosu Group in Japan*: Univ. Tokyo, Jour. Fac. Sci., sec. 2, v. 8, pt. 7, p. 337-350.

Kindle, E. M.

- (470) 1904, *The stratigraphy and paleontology of the Niagara of northern Indiana*: Indiana, Dept. Geology & Nat. Res., 28th Ann. Rept., p. 397-486, pl. 1-25.

King, William

- (471) 1844, *On a new genus of Palaeozoic shells*: Ann. & Mag. Nat. History, ser. 1, v. 14, p. 313-317.

Kira, Tetsuaki

- (472) 1962, *Shells of the western Pacific in color*: 224 p., 72 pl., Hoikusha Publ. Co. (Osaka, Japan).

Kirk, Edwin

- (473) 1927, *Tanaodon, a new molluscan genus from the Middle Devonian of China*: U.S. Natl. Museum, Proc., v. 70, art. 12, p. 1-4, pl. 1.
- (474) 1927, *Pycnodesma, a new molluscan genus from the Silurian of Alaska*: Same, Proc., v. 71, art. 20, p. 1-9, pl. 1-2.

Klinghardt, Franz

- (475) 1928, *Die Rudisten, Teil IV, Atlas und eingehende Figurenbeschreibung*: 24 pl., Selbstverlag (private publ. by author) (Berlin, Spandau).
- (475aa) 1931, *Die Rudisten, Teil III, Biologie und Beobachtungen an anderen Muscheln*: 60 p., 6 pl., privately publ. (Berlin).

Knudsen, Jørgen

- (475a) 1961, *The bathyal and abyssal Xylophaga (Pholadidae, Bivalvia)*: Galathea Report, no. 5, 163-209, fig. 1-41 (Copenhagen).

Kobayashi, Teichii

- (476) 1933, *Faunal study of the Wanwanian (basal Ordovician) Series, with special notes on the Ribeiridae and the ellesmereoceroids*: Tokyo Univ., Jour. Fac. Sci., sec. 2, v. 3, p. 249-328, pl. 1-10.

———, & **Amano, Masahisa**

- (477) 1955, *On the Pseudoquadratae trigonians, Steinmannella, in the Indo-Pacific Province*: Japanese Jour. Geology & Geography, v. 37, no. 3-4, p. 193-208.

———, & **Ichikawa, Koichiro**

- (478) 1951, *On Palaeopharus, a Late Triassic pelecypod genus*: Palaeont. Soc. Japan, Trans. & Proc., new ser., no. 1, p. 7-12.

———, & **Suzuki, Koiti**

- (479) 1936, *Non-marine shells of the Nakdong-Wakino Series*: Japan. Geol. Geogr., Jour., v. 13, p. 243-259, pl. 27-29.

———, & **Tokuyama, Akira**

- (480) 1959, *Daonella in Japan*: Univ. Tokyo, Jour. Fac. Sci., sec. 2, v. 12, pt. 1, p. 1-30.

Kobelt, Wilhelm

- (481) 1913, in E. A. Rossmäessler, *Iconographie der Land- und Süßwasser-Mollusken, mit vorzüglicher Berücksichtigung der europäi-*

- schen noch nicht abgebildeten Arten, fortges.: new ser., v. 19, 59 p., pl. 511-540, Arnold (Dresden, Leipzig), Constenable (Leipzig), Kreidel (Wiesbaden).
- Koerner, Karl**
 (482) 1937, *Marine (Cassianer-Raibler) Trias am Nevado de Acrotambo (Nord.-Peru)*: Palaeontographica, v. 86, pt. A, p. 145-237, pl. 10-14.
- Koninck, L. G. de**
 (483) 1841-44, *Description des animaux fossiles qui se trouvent dans le terrain carbonifère de Belgique*: iv+650 p., Atlas pl. A-H, 1-55 (Liège).
- Korobkov, I. A.**
 (484) 1954, *Spravochnik i metodicheskoe Rukovodstvo po tretichnym molluskam Platinchatozhabernye*: Gosud. Nauchno-tech. Issledov. Nefti. Gorno-toplivnoi lit-ry, Leningradskoi Otdelenie, 444 p., 96 pl. (in Russian). [*Handbook on and systematic guide to the Tertiary Mollusca, Lamellibranchia.*]
- Krestovnikov, V. N.**
 (485) 1931, *Kuyal'nitskiy yarus. Rukovodiashchiye iskopaemiye neftenosnykh rayonov Krymsko-Kavkazskoy oblasti. X*: Gosudarst. Issledov. Nefti. Inst., Trudy [Transactions of the State Petroleum Institute], no. 11, 24 p., 2 pl. [*Kuialnik beds, Characteristic fossils of the oil districts of the Crimea and Caucasus, no. 10.*]
- Krumbeck, Lothar**
 (486) 1923, *Brachiopoden, Lamellibranchiaten und Gastropoden aus der oberen Trias der Insel Seran (Mittel-Seran)*: Palaeontographica, Suppl. 4, pt. 3, p. 185-246, pl. 12-15.
 (487) 1924, *Die Brachiopoden, Lamellibranchiaten und Gastropoden der Trias von Timor. 2. Palaeontologischer Teil*: Paläontologie von Timor, no. 13, 275 p., 20 pl.
- Kühn, Othmar**
 (488) 1932, *Rudistae from eastern Persia*: Geol. Survey India, Records, v. 66, pt. 1, p. 151-179, text fig. 1-5, pl. 1-2.
 (489) 1933-49, *Stratigraphie und Paläogeographie der Rudisten*: Neues Jahrb. Mineralogie, Geologie, Paläontologie, Beilage, v. 70 (1933), p. 227-250, text fig. 7-8, pl. 9-10; v. 78 (1937), p. 268-284, text fig. 1-8; v. 86 (1942), p. 450-480, text fig. 1-4, pl. 28-30; v. 89 (1945), p. 167-194, text fig. 1-13, pl. 26; v. 90 (1949), p. 267-316, text fig. 1-2, pl. 14-15.
 (490) 1935, *Die Gattung Pironea Menegh.*: Zentralbl. Mineralogie, Geologie, u. Paläontologie, Abt. B, no. 9, p. 353-368, text fig. 1-6.
 (491) 1937-41, *Morphologisch-anatomische Untersuchungen an Rudisten*: Same, Abt. B, no. 5 (1937), p. 229-240; no. 12 (1941), p. 362-371, text fig. 1-5.
- (492) 1943, *Über Rudisten aus Sizilien, Lykien und Syrien*: Neues Jahrb. Geologie u. Paläontologie, Monatshefte, Abt. B, no. 1, p. 23-29, text fig. 1-2.
 (493) 1944, *Die Gattung Arnaudia Bayle*: Same, Abt. B, no. 2-3, p. 61-67.
 (494) 1952, *Rudiste nouveau du Vraconien des Pyrénées*: Univ. Grenoble, Lab. Géol., Travaux, v. 30, p. 79-80, text fig. 1-3.
- _____, & Pejovic, Desanka
 (495) 1959, *Zwei neue Rudisten aus Westserbien*: Österr. Akad. Wiss., Sitzungsber., Mat.-naturwiss. Kl., Abt. 1, v. 168, p. 979-989, 4 text fig., 4 pl.
- Kuester, H. C.**
 (496) 1838, *Die Gattung Anodonta nebst den übrigen Najaden mit unvollkommenem Schloss*: Martini-Chemnitz, Syst. Conch. Cab. (2), v. 9, pt. I, p. 1-64, pl. 1-21.
 (497) 1858-59, *Die Flussperlmuscheln (Unio et Hyria) in Abbildungen nach der Natur, mit Beschreibungen*: Same, v. 9, pt. 2, 318 p., 100 pl.
- Kuiper, J.-G.-J.**
 (498) 1962, *Note sur la systématique des pisidies*: Jour. Conchyliologie, v. 102, pt. 2, p. 53-57.
- Kumpera, Otakar, Prantl, Ferdinand, & Růžicka, Bokulav**
 (499) 1960, *Revision of the Nuculanidae from the Ostrava-Karviná District*: Národ. Musea Praze, Sborník, v. 16, B, p. 17-122.
- Kuroda, Tokubei, & Habe, Tadashige**
 (500) 1952, *Check list and bibliography of the Mollusca of Japan*: 210 p., Hosokawa Printing Co. (Tokyo).
- Lamarck, J. B. de**
 (501) 1799, *Prodrome d'une nouvelle classification des coquilles, comprenant une rédaction appropriée des caractères génériques, et l'établissement d'un grand nombre de genres nouveaux*: Soc. Histoire Nat. Paris, Mém. v. 1, p. 63-91.
 (502) 1801, *Système des animaux sans vertèbres*: viii + 432 p. (Paris).
 (503) 1809, *Philosophie zoologique, ou exposition des considérations relatives à l'histoire naturelle des animaux, la diversité de leur organisation et des facultés qu'ils en obtiennent, aux causes physiques qui main tiennent en eux la vie, et donnent lieu aux mouvements qu'ils exécutent; enfin, à celles qui produisent les unes les sentiments, et les autres l'intelligence de ceux qui en sont doués*: 2 vols., p. 1-422, 1-473 (Paris).
 (504) 1812, *Extrait du cours de zoologie du Muséum d'Histoire Naturelle . . .*: 127 p. (Paris).

- (505) 1818-19, *Histoire naturelle des animaux sans vertèbres*: v. 5, 612 p; v. 6, no. 1, 343 p. (Paris).
- Lameere, August**
- (506) 1903, *Sur l'évolution des mollusques*: Soc. Royal Zool. Malacol. Belgique, Ann., v. 38, p. xxx-xlvi.
- Lamy, Édouard**
- (507) 1914, *Révision des Mesodesmatidae vivants du Muséum d'Histoire Naturelle de Paris*: Jour. Conchyliologie, v. 62, no. 1, p. 1-74, 16 text fig., 1 pl.
- (508) 1914, *Révision des Scrobiculariidae vivants du Muséum d'Histoire Naturelle de Paris*: Same, v. 61, no. 3, p. 243-368, 7 text fig., 1 pl.
- (509) 1917, *Révision des Crassatellidae vivants du Muséum d'Histoire Naturelle de Paris*: Same, v. 62, no. 4, p. 197-270, 10 text fig., pl. 6.
- (510) 1917-18, *Révision des Macridae vivants du Muséum d'Histoire Naturelle de Paris*: Same, v. 63, no. 3, p. 173-275, pl. 6; no. 4, p. 291-411, 30 text fig., pl. 7.
- (511) 1919, *Révision des Astartidae vivants du Muséum d'Histoire Naturelle de Paris*: Same, v. 64, no. 2, p. 70-119, 3 text fig.
- (512) 1920, *Révision des Cypricardiacea et des Isocardiacea vivants du Muséum d'Histoire Naturelle de Paris*: Same, v. 64, pt. 4, p. 259-307, text fig.
- (513) 1920-21, *Révision des Lucinacea vivants du Muséum d'Histoire Naturelle de Paris*: Same, v. 65, p. 71-122, 169-222 (1920), p. 233-318, 335-388 (1921).
- (514) 1921, *Révision des Carditacea vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 66, p. 218-353, pl. 7-8.
- (515) 1922, *Révision des Petricola vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 67, no. 4, p. 309-359, pl. 3.
- (516) 1925, *Révision des Saxicavidae vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 68, p. 261-283.
- (517) 1925, *Révision des Gastrochaenidae vivants du Muséum d'Histoire Naturelle de Paris*: Same, v. 68, p. 284-319.
- (518) 1925-26, *Révision des Pholadidae vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 69, p. 19-51, 79-103 (1925); p. 136-168, 193-222 (1926).
- (519) 1927, *Révision des Myidae vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 70, p. 151-185, 4 text fig.
- (520) 1927, *Description d'une Pholade nouvelle de la côte d'Annam*: Muséum Natl. Histoire Nat., Bull., v. 33, p. 180-183, text fig. 1-3.
- (521) 1927, *Révision des Teredimidae vivants du Muséum National d'Histoire Naturelle de Paris*: Jour Conchyliologie, v. 70, p. 201-284.
- (522) 1928, *Révision des Lyonsiidae vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 72, p. 237-264, 2 text fig.
- (523) 1931, *Révision des Thraciidae vivants du Muséum d'Histoire Naturelle de Paris*: Same, v. 75, p. 213-241, 285-302.
- (524) 1931, *Révision des Periplomatidae vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 75, p. 303-321.
- (525) 1934, *Révision des Pandoridae vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 78, pt. 2, p. 95-125.
- (526) 1934, *Révision des Anatina vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 78, no. 3, p. 145-168, 1 pl.
- (527) 1936-37, *Révision des Mytilidae vivants du Muséum National d'Histoire Naturelle de Paris*: Same, v. 80, p. 66-102, 107-198, 229-295, 307-363.
- (528) 1938, *Mission Robert Ph. Dollfus en Egypte—Mollusca*: Inst. Egypte, Mém., v. 37, Testacea, p. 1-89.
- Lankester, E. R.**
- (529) 1883, *Mollusca*: in Encyclopaedia Britannica, v. 16, edit. 9, p. 632-695 (London).
- LaRocque, Aurèle**
- (530) 1950, *Pre-Devonian Devonian pelecypods of Michigan*: Univ. Michigan, Museum Paleontology, Contrib., v. 7, no. 10, p. 271-366, 19 pl.
- Laseron, C. F.**
- (531) 1953, *Minute bivalves from New South Wales*: Australian Museum, Records, v. 23, no. 2, p. 1-53, text fig. 1-44.
- (532) 1956, *New South Wales mussels. A taxonomic review of the family Mytilidae from the Peronian zoogeographical province*: Austral. Zool., v. 12, pt. 3, p. 263-283, 53 text fig.
- Latreille, P. A.**
- (533) 1825, *Familles naturelles du règne animal, exposées succinctement et dans un ordre analytique avec l'indication de leurs genres*. . . : 570 p. (Paris).
- Lea, Isaac**
- (534) 1832-74, *Observations on the genus Unio*: 13 vols. and index vol., 293 pl., printed by J. Kay, Jr. & Co. (Philadelphia).
- (535) 1870, *A synopsis of the family Unionidae*: 4th edit., 184 p. (see p. 72, footnote), H. C. Lea (Philadelphia).
- Lebedev, I. B.**
- (536) 1944, *O nekotorykh plastinchatozhabernykh iz kol'chuginskoy svity Kuzbassa*: Akad. Nauk. Sci. SSSR, Izvestiya, Cl. Biol., no. 2, p. 78-88, 1 pl. [On some lamellibranchs from the Kolchugino Series of the Kuznetsk Basin.]

- Lebküchner, R.**
(537) 1933, *Die Trigonien des süddeutschen Jura*: Palaeontographica, v. 77, p. 1-119, pl. 1-16.
- LeLoup, Eugène**
(538) 1950, *Lamellibranches, Res. Scient. Explor. Hydrobiol. Lac Tanganika (1946-47)*: Inst. Royal Sci. Nat. Belgique, Bull., v. 3, pt. 1, p. 1-153, 61 text fig., 8 p.
- Leroy, Pierre**
(539) 1940, *The late-Cenozoic unionids of China*: Geol. Soc. China, Bull., v. 19, p. 393-453, 21 text fig.
- Leymerie, M. A.**
(540) 1842, *Suite du mémoire sur le terrain Crétacé du Département de l'Aube*: Soc. Géol. France, Mém., v. 5, pt. 1, p. 1-34, pl. 1-13.
- Lichkov, B.**
(541) 1912, *Trigonies mésozoïques de Manghycklak*: Obshch. Estest., Zapiski, Kiev, v. 22, p. 89-145, pl. 11, 12.
- Lindahl, Josua**
(542) 1906, *Orthography of the names of the naiades*: Cincinnati Soc. Nat. History, Jour., v. 20, p. 235-243.
- Lindholm, W. A.**
(543) 1925, *Anodonta arcaeformis Heude im Süd-Ussuri-Gebiet*: Arch. Molluskenkunde, v. 57, p. 137-139.
(544) 1932, *Mollyuski iz sredne-Pljotsenovykh presnovodnykh otlozheniy Yugo-zapadnoy Sibiri*: Vsesoy. Geol. -Razved. Obed. SSSR, Trudy, pt. 238, p. 1-50, pl. 1-6. [Mollusken aus mittelplozänen Süßwasserablagerungen Südwest-Sibiriens.]
(545) 1932, *Süßwassermollusken aus pliozaenen Ablagerungen am Fluss Irtysh*: Union. Geol. Prosp. Service USSR, Trans., pt. 239, p. 1-27, pl. 1-7.
- Link, H. F.**
(546) 1806-08, *Beschreibung der Naturalien-Sammlung der Universität zu Rostock*: p. 1-160 (1806), pl. 1-30, 1-38 (1807), pl. 1-38 (1808) (Rostock).
- Lintz, Joseph, Jr.**
(547) 1958, *The fauna of the Ames and Brush Creek Shale of the Conemaugh Formation of western Maryland*: Jour. Paleontology, v. 32, p. 97-112, 2 text fig., pl. 16.
- Logan, J. A.**
(548) 1964, *The dentition of the Durham Permian pelecypod Permophorus costatus (Brown)*: Palaeontology, v. 7, p. 281-285.
- Loosanoff, V. L., Davis, H. C., & Chanley, P. E.**
(548a) 1966, *Dimensions and shapes of larvae of some marine bivalve mollusks*: Malacologia, v. 4, no. 2, p. 351-435.
- Loriol, C. L. P. de**
(549) 1875, *Description des fossiles. Mollusques acéphales*: in Monographie paléontologique et géologique des étages supérieurs de la formation jurassique des environs de Boulogne-sur-Mer, Soc. Phys. Histoire Nat. Genève, Mém., v. 24, p. 1-326, pl. 11-26.
(550) 1889-93, *Études sur les mollusques des couches coralligènes inférieures du Jura Bernois*: Soc. Paléont. Suisse, Mém., v. 16-19, 419 p., 37 pl.
- Lucas, Gabriel**
(551) 1952, *Étude microscopique et pétrographique de la coquille des Lamellibranches*: in Jean Piveteau, *Traité de paléontologie*, v. 2, p. 246-261, text fig. 17-34, Masson (Paris).
- Lupher, R. L., & Packard, E. L.**
(552) 1930, *The Jurassic and Cretaceous rudistids of Oregon*: Univ. Oregon Publ., Geol. Ser., v. 1, no. 3, p. 201-212, 1 text fig., pl. 1-6.
- Lycett, John**
(553) 1950, (1), *On Trichites, a fossil genus of bivalve mollusks*: Ann. & Mag. Nat. History, ser. 2, v. 5, p. 343-347, pl. 10.
(554) 1872-83, *A monograph of the British fossil Trigoninae*: Palaeontograph. Soc. London, 245 + 19 p., 41 + 4 pl.
- McAlester, A. L.**
(555) 1962, *Upper Devonian pelecypods of the New York Chemung Stage*: Peabody Museum Nat. History, Bull. 16, 88 p., 32 pl.
(556) 1963, *Revision of the type species of the Ordovician nuculoid pelecypod genus Tancrediopsis*: Same, Postilla, no. 74, p. 1-19, text fig. 1-80 (on 4 pl.).
(557) 1965, *Systematics, affinities, and life habits of Babinka, a transitional Ordovician lucinoid bivalve*: Palaeontology, v. 8, pt. 2, p. 231-246, pl. 26-28.
(558) 1966, *Evolutionary and systematic implications of a transitional Ordovician lucinoid bivalve*: Malacologia, v. 3, p. 433-439.
(558a) 1968, *Type species of Paleozoic nuculoid bivalve genera*: Geol. Soc. America, Mem. 105, 143 p., 36 pl., 2 tab.
- M'Coy, Frederick**
(559) 1852, *Detailed systematic description of the British Palaeozoic fossils in the Geological Museum of the University of Cambridge*: in A. Sedgwick & F. M'Coy, p. 185-406, 6 pl. (London & Cambridge).
- McFarlan, A. C.**
(560) 1931, *The Ordovician fauna of Kentucky*: in *The paleontology of Kentucky*, W. R. Jillson (ed.), p. 47-165, pl. 1-16.
- MacGillavry, H. J.**
(561) 1937, *Geology of the province Camaguey, Cuba, with revisional studies of rudist paleontology*: Utrecht Rijks-Univ., Geogr. Geol. Meded., Phys. Geol. Reeks Thèse no. 14, p. 1-168, pl. 1-10, 1 map.

McLean, R. A.

- (562) 1947, *A revision of the pelecypod family Tridacnidae*: Notulae Naturae, Acad. Nat. Sci. Philadelphia, no. 195, p. 1-7, text fig. 1-12.

McLearn, F. H.

- (563) 1918, *The Silurian Arisaig Series of Arisaig, Nova Scotia*: Am. Jour. Sci., v. 45, p. 126-140.
- (564) 1924, *Palaeontology of the Silurian rocks of Arisaig, Nova Scotia*: Geol. Survey Canada, Mem. 137, 180 p., 30 pl.

MacLennan, R. M.

- (565) 1944, *Hinge structure in Carbonicola pseudobusta Trueman and related species*: Geol. Mag., v. 81, p. 1-14, text fig. 1-4, 1 pl.

McMichael, D. F.

- (566) 1926, *Notes on the freshwater mussels of New Guinea*: Nautilus, v. 70, p. 38-48, pl. 2.

—, & Hiscock, I. D.

- (567) 1958, *A monograph of the freshwater mussels (Mollusca: Pelecypoda) of the Australian region*: Austral. Jour. Mar. & Freshwater Res., v. 9, no. 3, p. 372-508, 18 text fig., 19 pl.

MacNeil, F. S.

- (568) 1935, *Fresh-water mollusks from the Catahoula Sandstone (Miocene) of Texas*: Jour. Paleontology, v. 9, p. 10-17, pl. 3-5.
- (569) 1936, (generic description) in F. K. Morris, *Central Asia in Cretaceous time*: Geol. Soc. America, Bull., 47, p. 1477-1534, 5 text fig., pl. 1-2.
- (570) 1937, *The systematic position of the pelecypod genus Trinacria*: Washington Acad. Sci., Jour., v. 27, p. 452-458.
- (571) 1965, *Evolution and distribution of the genus Mya and Tertiary migrations of Mollusca*: U.S. Geol. Survey, Prof. Paper 483-G, 51 p., 3 text fig., 11 pl.

MacPherson, J. H., & Gabriel, C. J.

- (572) 1962, *Marine mollusks of Victoria*: xv + 475 p., 486 text fig., Handbook No. 2, National Museum of Victoria, Melbourne Univ. Press & Natl. Museum Victoria (Melbourne).

Mailleux, Eugene

- (573) 1920, *Note sur quelques groupes de mollusques acéphales des terrains paléozoïques*: Soc. Belge de Géologie, Paléontologie et d'Hydrologie, Bull., v. 29, p. 140-150, 3 text fig.
- (574) 1931, *La faune des grès et schistes de Solières (siegenien moyen)*: Musée Royal Histoire Nat. Belgique, Mém., no. 51, 90 p., 2 pl.
- (575) 1932, *Le genre Limoptera J. Hall dans le Dévonien de l'Ardenne*: Same, Bull., v. 8, no. 10, p. 1-18, pl. 1-2.

- (576) 1936, *La faune des schistes de Matagne (Frasnien supérieur)*: Same, Mém., v. 77, p. 1-74, pl. 1.

- (577) 1937, *Les lamellibranches du Dévonien inférieur de l'Ardenne*: Same, Mém., v. 71, p. 1-273, pl. 1-14.

Maire, Victor, & Dechaseaux, Collette

- (578) 1937, *Sur quelques limidés de l'oolithe ferrugineuse de Talant (Côte-d'Or)*: Soc. Géol. France, Bull., v. 5, no. 6, p. 439-446.

March, M. C.

- (579) 1912, *A discussion of the general classification of the Pelecypoda*: Ann. & Mag. Nat. History, v. 10, no. 8, p. 91-166, pl. 3.

Marshall, W. B.

- (580) 1916, *A new genus of species of naiad from the James River at Huron, South Dakota*: Nautilus, v. 29, p. 133-135, pl. 4.
- (581) 1922, *New pearly fresh-water mussels from South America*: U.S. Natl. Museum, Proc., v. 61, art. 16, 9 p., 3 pl.
- (582) 1927, *A new genus and two new species of South American fresh-water mussels*: Same, v. 71, art. 6, 4 p., 2 pl.
- (583) 1928, *New fossil pearly fresh-water mussels from deposits on the Upper Amazon of Peru*: Same, v. 74, art. 3, p. 1-7, pl. 1.
- (584) 1929, *New fossil land and fresh-water mollusks from the Reynosa Formation of Texas*: Same, v. 76, art. 1, 6 p., 1 pl.
- (585) 1931, *Ruganodontites, a new subgenus of South American pearly fresh-water mussels*: Nautilus, v. 45, p. 16-20.
- (586) 1934, *Two new species of pearly fresh-water mussels*: Washington Acad. Sci., Jour., v. 24, p. 78-81, 6 text fig.

—, & Bowles, E. O.

- (587) 1932, *New fossil fresh-water mollusks from Ecuador*: U.S. Natl. Museum, Proc., v. 82, art. 5, 7 p., 1 pl.

Martens, Edouard von

- (588) 1890-1901, *Land and freshwater Mollusca*: Biologia Centrali-Americana, 706 p., 44 pl.
- (589) 1897, *Beschalte Weichtiere Deutsch-Ost-Afrikas*: Ostafrika, v. 4, 308 p., text fig., 7 pl.

Martinson, G. G.

- (590) 1953, *Novye Unionidy iz Verkhnemelovykh Olozhenii Mongolii*: Akad. Nauk. USSR, Doklady, v. 89, p. 167-170, 3 text fig. [New unionids from Upper Cretaceous deposits in Mongolia.]

Marwick, John

- (591) 1924, *Zealeda and Barytellina, new fossil molluscan genera from New Zealand*: Malacol. Soc. London, Proc., v. 16, p. 25-26, text fig. 1-7.
- (592) 1927, *The Veneridae of New Zealand*: New Zealand Institute, Trans., v. 57, p. 567-635, pl. 34-54, text fig.

- (593) 1928, *The Tertiary Mollusca of the Chatham Islands including a generic revision of the New Zealand Pectinidae*: Same, Trans. & Proc., v. 58, pt. 4, p. 432-506, text fig. 1-148.
- (594) 1929, *Tertiary molluscan fauna of Chatton, Southland*: Same, v. 59, p. 903-926, text fig. 1-75.
- (595) 1931, *The Tertiary Mollusca of the Gisborne District*: New Zealand Geol. Survey, Palaeont. Bull., no. 13, 177 p., 18 pl.
- (596) 1935, *Some new genera of the Myalinidae and Pteriidae of New Zealand*: Royal Soc. New Zealand, Trans., v. 65, p. 295-303, pl. 34-36.
- (597) 1938, *Notocallista and its allies*: Same, v. 68, p. 60-81, pl. 10-14.
- (598) 1944, *New Zealand fossil and Recent Cardiidæ (Mollusca)*: Same, Trans., v. 74, pt. 3, p. 255-272, pl. 25-37.
- (599) 1953, *Division and faunas of the Hokonui System (Triassic and Jurassic)*: New Zealand Geol. Survey, Paleont. Bull. 21, 141 p., 3 text fig., 17 pl.
- Masuda, Koichiro**
(600) 1962, *Tertiary Pectinidae of Japan*: Tohoku Univ., Sci. Rept., ser. 2 (Geol.), v. 33, no. 2, 238 p., 27 pl.
- Matheron, Philippe**
(600a) 1843, *Catalogue méthodique et descriptif des corps organisés fossiles du Département des Bouches-du-Rhône et lieux circonvoisins*: 269 p., 41 pl. (Marseilles).
- (600b) 1878, *Recherches paléontologiques dans le midi de la France*: pt. 3, pl. C-14 (Marseilles).
- Meek, F. B.**
(601) 1864, *Check list of the invertebrate fossils of North America. Cretaceous and Jurassic*: Smithsonian Misc. Coll., no. 177 (in v. 7), 40 p.
- (602) 1864, *Remarks on the family Pteriidae (=Aviculidae), with descriptions of some new fossil genera*: Am. Jour. Sci., ser. 2, v. 37, p. 212-220.
- (603) 1871, *On some new Silurian crinoids and shells*: Same, ser. 3, v. 2, p. 295-302.
- (604) 1872, *Preliminary paleontological report, consisting of lists of fossils, with descriptions of some new types. . . .*: U. S. Geol. Survey Wyoming, Prelim. Rept., 1870, p. 285-318.
- (605) 1872, *Descriptions of new western Palaeozoic fossils, mainly from the Cincinnati Group of the Lower Silurian series of Ohio*: Acad. Nat. Sci. Philadelphia, Proc. for 1871, p. 308-336.
- (606) 1873, *Descriptions of invertebrate fossils of the Silurian and Devonian systems*: Geol. Survey Ohio, Rept., v. 1, pt. 2, Palaeontology, p. 1-243, pl. 1-23.
- (607) 1875, *A report on some of the invertebrate fossils of the Waverly Group and Coal Measures of Ohio*: Same, v. 2, Geology and Palaeontology, part 2, Geology and Palaeontology, part II, Palaeontology, p. 269-347, pl. 10-20.
- , & **Hayden, F. V.**
(608) 1865, *Palaeontology of the upper Missouri: Invertebrates*: Smithsonian Contrib. Knowledge, v. 14, no. 172, vii + 135 p., 5 pl.
- (609) 1876, *A report on the invertebrate Cretaceous and Tertiary fossils of the Upper Missouri country*: U.S. Geol. Survey of the Territories, Rept., v. 9, 629 p., 45 pl.
- , & **Worthen, A. H.**
(610) 1868, *Geology and palaeontology*: Geol. Survey Illinois, v. 3, 574 p., 20 pl.
- (611) 1869, *Descriptions of new Carboniferous fossils from the western states*: Acad. Nat. Sci. Philadelphia, Proc., p. 137-172.
- (612) 1870, *Descriptions of new species and genera of fossils from the Palaeozoic rocks of western states*: Same, Proc., p. 22-56.
- Ménégaux, Auguste**
(613) 1889, *Sur la branche des lamellibranches et sur sa comparaison avec celle des scutibranches*: Soc. Philomath. Paris, Bull., v. 1, no. 8, p. 137-144.
- Mennessier, Guylot**
(614) 1949, *Sur la présence de rudistes dans un sédiment hautement ligniteux à Piolenc (Vaucluse)*: Soc. Géol. France, Comptes Rendus Somm. Séanc., no. 14, p. 215-217.
- Miller, S. A.**
(615) 1874, *Genus Anomalodonta*: Cincinnati, Quart. Jour. Sci., v. 1, p. 16-18, text fig. 7-9.
- (616) 1881, *Description of new species of fossils from the Hudson River group, and remarks upon others*: Cincinnati Soc. Nat. History, Jour., v. 4, p. 316-319, pl. 8.
- (617) 1882, *Description of three new species and remarks upon others*: Same, v. 5, p. 116-117, pl. 5.
- (618) 1889, *North American geology and palaeontology*: 664 p., 1194 text fig.
- (619) 1897, *Second appendix to North American geology and palaeontology*: p. 719-793, text fig. 1266-1458.
- , & **Faber, C. L.**
(620) 1894, *New species of fossils from the Hudson River group, and remarks upon others*: Same, v. 17, p. 22-33, pl. 1.
- Milovanović, Branislav**
(621) 1933, *Les problèmes paléobiologiques et bio-stratigraphiques des rudistes*: Serv. Géol. Yougoslavie., Mém., p. 1-194.

- (622) 1934, *Les rudistes de la Yougoslavie*: Geol. An. Balk. Poluost., v. 12, pt. 2, p. 275-308, pl. 1-5 (in Serbian; French summ., p. 302-308).
- (623) 1935, *Novi rudisti Srbije*: Srpske Kral'ev. Akad., Glasnik, v. 166, p. 47-125, 31 text fig. [*New Serbian rudists*.]
- (624) 1936, *Yvania maestrichtiensis n. sp. et son importance pour la formation brusque des formes nouvelles*: Geol. An. Balk. Poluost., v. 13, p. 28-54, 3 text fig., 2 pl.
- (625) 1938, *La structure et la fonction des piliers des hippurites*: Same, v. 15, p. 256-267, pl. 9.
- (626) 1938, *Anatomie comparée et relations phylogénétiques des Lapeirouseinae*: Acad. Sci. math.-nat. (Belgrade), Bull., sec. B, Sci. Nat., no. 4, p. 85-152, text fig. 1-29, pl. 1-6.
- (627) 1956, *Kuechnia nov. gen. i aberantni rodovi*: Zavod Geol. Geof. Srbije, Vestnik, v. 12, p. 131-144, 2 text-fig., 1 pl. [*Kuechnia, nov. gen. and aberrant genera*.]
- (628) 1957, *Sifonalna zona rudista*: Same, Vestnik, v. 13, p. 115-148, 2 text fig., 1 pl. [*The siphonal zone of the rudists*.]
- Modell, Hans**
- (629) 1931, *Die Najaden der oberbayrischen Cyrenenschichten*: Palaeontographica, v. 75, p. 63-80, pl. 13-14.
- (630) *Das natürliche System der Najaden*: Arch. Molluskenkunde, v. 74, p. 161-191, pl. 5-7.
- (631) 1943, *Tertiäre Najaden, III*: Same, v. 75, p. 107-117, pl. 1.
- (632) 1950, *Tertiäre Najaden, IV*: Same, v. 79, p. 25-36, 4 text fig., pl. 6-7.
- (633) 1950, *Südamerikanische Najaden der Gattungen Castalia, Schlesiella und Ecuadorea*: Same, v. 79, p. 135-146, pl. 11.
- Möller, H. P. C.**
- (634) 1842, *Index molluscorum Groenlandiae*: Naturhist. Tidsskr., v. 4, no. 1, p. 76-97.
- Mörch, O. A. L.**
- (635) 1853, *Catalogus conchyliorum quae reliquit D. Alphonso d'Aguirra et Gadea, Comes de Yoldi, regis Daniae cubiculariorum princeps, ordinis Danebrogici in prima classe et ordinis tertii eques. Fasc. secundus, Acephala*: 74 p. (Copenhagen).
- (636) 1857, *Prodromus faunae molluscorum (sic) Grönlandiae*: 28 p. (Copenhagen).
- Moll, Friedrich**
- (637) 1941, *Uebersicht über die Terediniden des Museums für Naturkunde zu Berlin*: Gesell. Naturforsch. Freunde Berlin, Sitzungsber., v. 10, p. 152-225.
- (638) 1941, *Teredinidae*: in Fossilium Catalogus I: Animalia, Pars 95, 89 p., W. Junk (s'Gravenhage).
- (639) 1942, *Die fossilen Terediniden und ihre Beziehung zu den rezenten Arten*: Palaeontographica, v. 94, Abt. A, no. 3-6, p. 134-153, pl. 24-26.
- Monod, Théodore, Nicklès, Maurice, & Moll, Friedrich**
- (640) 1952, *Xylophages et pétricoles ouest africains*: Inst. Français d'Afrique Noire, Cat., v. 8, 145 p., 12 pl.
- Montagne, D. G.**
- (641) 1941, *Geologie und Paläontologie der Umgebung von Sestanovac Dalmatien*: Utrecht Rijks Univ., Geogr. Geol. Meded., Phys. Geol. Reeks. ser. 2, no. 1, 93 p., 8 pl., 3 maps.
- Monterosato, T. Allery de**
- (642) 1875, *Nuova rivista delle conchiglie Mediterranee*: Accad. Sci. Lett. Arti, Palermo, Atti (new ser.), v. 5, 50 p.
- Moret, Léon**
- (643) 1936, *Durania delphinensis, nouvelle espèce de rudiste du Vraconien de la Fauge, près Grenoble, et révision du genre Durania Douvillé*: Soc. Sci. Daughiné, Bull., v. 55 (for 1935), p. 1-24, pl. 1.
- Morretes, F. L. de**
- (644) 1945, *Rochanaia gutmansii nov. gen. n. sp., novo lamellibranquio de Brasil*: Arquivos Inst. Biol. São Paulo, v. 12 (1941), p. 75-79, pl. 31-32.
- Morris, John, & Lycett, John**
- (645) 1853-54, *A monograph of the Mollusca from the Great Oolite, chiefly from Minchinhampton and the coast of Yorkshire, part II, Bivalves*: Palaeontograph. Soc. London, 147 p., 15 pl.
- Morton, John**
- (646) 1963, *The molluscan pattern: evolutionary trends in a modern classification*: Linnean Soc. London, Proc., Sess. 174, p. 1, p. 53-72.
- Mossevitich, N. A. K.**
- (647) 1928, *Academie des Sciences de l'Union des Républiques Sovietique Socialistes, Matériaux de la Commission pour l'Étude de la République Autonome Sovietique Socialiste laKonte, Liv. 19, Beiträge zur Systematic Ökologie, und Verbreitung der Recenter und Fossilen Yoldia artica Gray*: p. 1-44, 1 pl. (Leningrad).
- Moulins, Charles des**
- (648) 1827, *Description de trois genres nouveaux de coquilles fossiles du terrain tertiaire de Bordeaux*: Soc. Linnéenne Bordeaux, Bull. Hist. Nat., v. 2, p. 226-255, pl. 5-7.
- (649) 1832, *Description d'un nouveau genre de coquille vivante, bivalve, des mers du Chili*: Soc. Linnéenne de Bordeaux, Actes, v. 5 (Feb.), p. 83-92.
- Müller, J. M.**
- (650) 1847, *Monographie der Petrefacten der*

- Aachener Kreideformation*: pt. 1 (1847), 48 p., 2 pl. (Bonn, Germany).
- Müllerried, F. K. G.**
 (651) 1931, *Sobre un anomalia en las invaginaciones de las valvas de algunas Hippuritidae*: An. Inst. Biol. Mexico, v. 2, p. 255-261, 2 text fig.
 (652) 1933, *Die Beziehungen der Pachyodonten Amerikas zu denen der alten Welt*: Geol. Rundschau, Sonderband, v. 23a, p. 267-271.
- Munier-Chalmas, E. C. P. A.**
 (652a) 1882, *Études critiques sur les rudistes*: Soc. Géol. France, Bull., sér. 3, v. 10, p. 472-492, 1 text fig., pl. 10-11.
 (652b) 1895, *Note préliminaire sur le développement de la charnière chez les mollusques acéphales*: Same, Bull., sér. 3, v. 23, p. 73-80.
- Murchison, R. I.**
 (653) 1839, *The Silurian system*: Part 1, 768 p., John Murray (London).
- Murray, H. D., & Leonard, A. B.**
 (654) 1962, *Handbook of unionid mussels in Kansas*: Museum Nat. History, Univ. Kansas, Misc. Publ. 28, p. 1-184, 42 text fig., 45 pl.
- Nagao, Takumi**
 (655) 1938, *Some molluscan fossils from the Cretaceous deposits of Hokkaido and Japanese Saghalien, Pt. 1, Lamellibranchiata and Scaphopoda*: Hokkaido Univ., Geol. and Min., Jour. Fac. Sci., ser. 4, pt. 1-2, p. 116-142, pl. 14-16.
- Nakazawa, Keiji**
 (656) 1952-55, *A study on the pelecypod fauna of the upper Triassic Nabae group in the northern part of Kyoto Prefecture, Japan*: Kyoto Imp. Univ., Mem. Coll. Sci., ser. B, pt. 1 (1952), pectinids and limids, v. 20, no. 2, p. 95-106; pt. 2 (1954), bavevelliids, v. 21, no. 2, p. 213-222; pt. 3 (1955), halobiids and others, v. 22, no. 2, p. 243-260.
 (657) 1960, *Permian and Eo-Triassic Myophoriidae from the Maizuru zone, southwest Japan*: Japanese Jour. Geology & Geography, v. 31, no. 1, p. 49-62.
 (658) 1961, *Early and Middle Triassic pelecypod fossils from the Maizuru zone, southwest Japan*: Kyoto Imp. Univ. Mem. Coll. Sci., ser. B, v. 27, no. 3, Geol. & Min., p. 249-291, pl. 12-14.
- Neave, S. A.**
 (659) 1939-50, *Nomenclator zoologicus, a list of the names of genera and subgenera in zoölogy from the tenth edition of Linnaeus 1758 to the end of 1935*: v. 1, A-C (1939), 957 p.; v. 2, D-L (1939), 1025 p.; v. 3, M-P (1940), 1065 p.; v. 4 (1940), Q-Z and supplement, 758 p.; v. 5 (1950, 1936-45), 308 p.
- Neumayr, Melchior**
 (660) 1883, *Zur Morphologie des Bivalvenschlosses*: K. K. Akad. Wiss. Wien, Nat. Math. Cl., Sitzungsber., Abth. 1, v. 88, no. 1, p. 385-418.
 (661) 1889, *Über die Herkunft der Unioniden*: Same, v. 98, p. 5-27, pl. 1-3.
 (662) 1891, *Beiträge zu einer morphologischen Eintheilung der Bivalven*: Same, Denkschr., v. 58, p. 701-801.
- Newell, N. D.**
 (663) 1937(1938), *Late Paleozoic pelecypods: Pectinacea*: Kansas State Geol. Survey, Publ., v. 10, no. 1, 123 p., 20 pl.
 (664) 1940, *Paleozoic pelecypods Myalina and Naiadites*: Am. Jour. Sci., v. 238, p. 286-295, 2 text fig., 2 pl.
 (665) 1940, *Invertebrate fauna of the late Permian Whitehorse Sandstone*: Geol. Soc. America, Bull. 51, p. 261-336, 2 text fig., 10 pl.
 (666) 1942, *Late Paleozoic pelecypods: Mytilacea*: Kansas State Geol. Survey, Publ., v. 10, no. 2, 115 p., 20 pl.
 (667) 1954, *Status of invertebrate paleontology, 1953, V. Mollusca*: Harvard Univ., Museum Comp. Zoology, Bull., v. 112, p. 161-172.
 (668) 1956, *Primitive desmodont pelecypods of the Australian Permian*: Am. Museum Nat. History, Novitates, no. 1799, p. 1-13.
 (669) 1957, *Notes on certain primitive heterodont pelecypods*: Same, v. 1857, p. 1-14, 3 text fig.
 (670) 1960, *The origin of the oysters*: Internat. Geol. Congress, Rept. 21st Sess., Norden, pt. 22, p. 81-86.
- Newton, R. B.**
 (671) 1909, *Cretaceous Gastropoda and Pelecypoda from Zululand*: Royal Soc. South Africa, Trans., v. 1, p. 1-106, pl. 1-9.
- Nicol, David**
 (672) 1944, *Observations on Pseudomonitis, a late Paleozoic pelecypod*: Nautilus, v. 57, p. 90-93.
 (673) 1947, *The genotype of Isoarca (Class Pelecypoda)*: Jour. Paleontology, v. 21, p. 344-345, 4 text fig.
 (674) 1950, *Origin of the pelecypod family Glycymeridae*: Same, v. 24, no. 1, p. 89-98.
 (675) 1932, *Nomenclatorial review of genera and subgenera of Chamidae*: Washington Acad. Sci., Jour., v. 42, no. 5, p. 154-156.
 (676) 1953, *Review of the living species of Echinochama*: Same, v. 43, no. 11, p. 386-388, 4 text fig.

- (677) 1955, *Morphology of Astartella, a primitive heterodont pelecypod*: Jour. Paleontology, v. 29, p. 155-158, 4 text fig.
- , **Desborough, G. A., & Solliday, J. R.**
- (678) 1959, *Paleontologic record of the primary differentiation in some major invertebrate groups*: Washington Acad. Sci., Jour., v. 49, no. 10, p. 351-366.
- Noda, H.**
- (678a) 1966, *The Cenozoic Arcidae of Japan*: Tohoku Univ., Sci. Repts., ser. 2, v. 38, no. 1, p. 1-161, pl. 1-14, 35 tables.
- Noetling, Fritz**
- (679) 1899, *Notes on the morphology of the Pelecypoda*: India Geol. Survey, Mem., Palaeont. Indica, new ser., v. 1-2, 57 p., pl. 2-5.
- Oberling, J. J.**
- (680) 1955, *Shell structure of west American Pelecypoda*: Washington Acad. Sci., Jour., v. 45, p. 128-130, 2 text fig.
- (681) 1964, *Observations on some structural features of the pelecypod shell*: Naturforsch. Gesell. Bern., Mittell., new ser. v. 20, 63 p.
- Odhner, N. H.**
- (682) 1912, *Morphologische und phylogenetische Untersuchungen über die Nephridien der Lamellibranchien*: Zeitschr. Wiss. Zoologie, v. 100, p. 287-391, 40 text fig.
- (683) 1918, *Zur Kenntnis der Homologien des Bivalvenschlosses*: Geol. Fören. Stockholm, Förhandl., v. 40, p. 562-590.
- (684) 1919, *Studies on the morphology, the taxonomy, and the relation of Recent Chamidae*: K. Svenska Vetenskapsakad., Handl., v. 59, no. 3, 102 p., 8 pl.
- (685) 1924, *Papers from Dr. Th. Mortensen's Pacific Expedition 1914-16. XIX New Zealand Mollusca*: Vidensk. Medd. Dansk Naturh. Foren, v. 77, p. 1-90, 24 text fig., pl. 1-2.
- (686) 1925, *Shells from the San Men Series*: Geol. Survey China, Palaeont. Sinica, ser. B, v. 6, p. 1-20, pl. 1-5.
- Öpik, A. A.**
- (687) 1930, *Beiträge zur Kenntnis der Kukuruse-(C₂-C₃) Stufe in Eesti*: Tartu Ülikooli Geoloogia-Instituudi Toimetused (Publ. Geol. Inst. Tartu), no. 24, 34 p., 11 text fig., 4 pl.
- Olsson, A. A.**
- (688) 1961, *Mollusks of the tropical eastern Pacific. Panamic - Pacific Pelecypoda*: Paleont. Res. Inst., 574 p., 86 pl. (Ithaca, New York).
- , **& Harbison, A.**
- (689) 1953, *Pliocene Mollusca of southern Florida with special reference to those from north Saint Petersburg*: Acad. Nat. Sci. Philadelphia, Mon. 8, p. 1-304, 63 pl.
- , **& Wurtz, C. B.**
- (690) 1951, *New Colombian naiades, with observations on other species*: Acad. Nat. Sci. Philadelphia, Notulae Naturae, no. 239, 9 p., 12 text fig.
- Oppenheim, Paul**
- (691) 1892, *Ueber einige Brackwasser und Binnen-Mollusken aus der Kreide und dem Eocän Ungarns*: Deutch. Geol. Gesell., Zeitschr., v. 44, p. 697-818, pl. 31-36.
- (692) 1903, *Zur Kenntnis alttertiärer Faunen in Aegypten. I. Der Bivalven erster Teil*: Palaeontographica, v. 30, no. 3, p. 1-164, pl. 1-17.
- Orbigny, Alcide d'**
- (693) 1835, *Synopsis terrestrium et fluviatilium molluscorum, in suo per Americam Meridionalem itinere collectorum*: Mag. Zool., v. 5, cl. 5, no. 61-62, 44 p.
- (694) 1839-47, *Voyage dans l'Amerique meridionale pendant 1826-1833*: Zoologie, v. 4-6, 758 p., 85 pl.
- (695) 1844 (1843), *Paléontologie française terrains crétacés, v. 3, Mollusques*: 807 p., pl. 237-489, G. Masson (Paris).
- (696) 1845, *Système jurassique (Étage oxfordien). Mollusques*: in Géologie de la Russie d'Europe et des montagnes de l'Oural, by R. I. Murchison, E. de Verneuil, & A. de Keyserling, v. 2, p. 419-488, pl. 28-42 (London and Paris).
- (697) 1851, *Note sur un nouveau genre de coquille lamellibranche d'eau douce découvert dans les rivières de la Nouvelle Grenade par M. Acosta*: Rev. Mag. Zool., v. 3, p. 183-188.
- Oria, M.**
- (698) 1933, *Observations sur des Ostreidae de l'Oxfordien de Normandie*: Soc. Linnéenne Normandie, Bull., v. 8, no. 5, p. 19-76, pl. 1-3.
- Ortmann, A. E.**
- (699) 1912, *Notes upon the families and genera of the najades*: Carnegie Museum, Ann., v. 8, p. 222-365, 28 text fig., pl. 18-20.
- (700) 1912, *Cumberlandia, a new genus of naiades*: Nautilus, v. 26, p. 13-14.
- (701) 1914, *Study in najades*: Same, v. 28, p. 28-34, 41-47.
- (702) 1921, *South American naiades. A contribution to the knowledge of the fresh-water mussels of South America*: Carnegie Museum, Mem. 8, p. 451-670, 4 text fig., pl. 34-48.
- (703) 1922, *A new genus and species of American naiades*: Nautilus, v. 36, p. 1-6, pl. 1.

- , & Walker, B.
(704) 1912, *A new North American naiad*: *Nautilus*, v. 25, p. 97-100, pl. 8.
- Owen, Gareth**
(705) 1955, *Observations on the stomach and digestive diverticula of the Lamellibranchia. I. The Anisomyaria and Eulamellibranchia*: *Micro. Sci., Quart. Jour.*, v. 96, p. 517-537.
(706) 1959, *The ligament and digestive system in the taxodont bivalves*: *Malacol. Soc. London, Proc.*, v. 33, pt. 5, p. 215-223.
(707) 1959, *Observations on the Solenacea with reasons for excluding the family Glaucomyidae*: *Royal Soc. London, Philos. Trans.*, B, v. 242, p. 59-97.
- , Trueman, E. R., & Yonge, C. M.
(708) 1953, *The ligament in the Lamellibranchia*: *Nature*, v. 171, p. 73-75.
- Oyama, Katsura, Mizuno, Atsuyuki, & Sakamoto, Toru**
(709) 1960, *Illustrated handbook of Japanese Paleogene molluscs*: 244 p., 71 pl., Geol. Survey Japan, printed by Dai-Nippon Printing Co. & Hanshuchi Printing Co., Ltd. (place of publication not stated).
- Pallary, P.**
(710) 1924, *Supplément à la faune malacologique terrestre et fluviatile de l'Égypte*: *Inst. Égypt., Mém.*, v. 7, p. 1-61, 7 text fig., pl. 1-4.
- Palmer, K. V. W.**
(711) 1927, *The Veneridae of eastern America, Cenozoic and Recent*: *Palaeont. Americana*, v. 1, p. 209-522, 45 pl.
(712) 1938, *Neocene Spondyli from the southern United States and tropical America*: Same, v. 2, no. 8, p. 147-162, pl. 16-18.
- , & Brann, D. C.
(713) 1965, *Catalogue of the Paleocene and Eocene Mollusca of the southern and eastern United States, part 1: Pelecypoda, Amphineura, Pteropoda, Scaphopoda and Cephalopoda*: *Bull. Am. Paleontology*, v. 48, no. 218, 443 p., 3 pl.
- Palmer, R. H.**
(714) 1928, *The rudistids of southern Mexico*: *Calif. Acad. Sci., Occas. Papers*, v. 14, 137 p., 8 text fig., 18 pl.
(715) 1933, *Nuevo rudistes de Cuba*: *Revista Agric. Com. Trabajo (Habana)*, Año 14, no. 15-16, p. 95-125, 1 text fig., pl. 1-10.
- Papp, Adolph**
(716) 1955, *Bemerkungen über Vorkommen und Variabilität der Bivalvengattung Onco-phora*: *Geol. Bundesanst., Verhandl.*, 1955, no. 2, p. 120-133, 1 pl.
- Paquier, M. V.**
(716a) 1898, *Sur quelques dicératinés nouveaux du Tithonique*: *Soc. Géol. France, Bull.*, sér. 3, v. 25, p. 843-851, 2 text fig., pl. 23.
(716b) 1903, *Les rudistes urgoniens, part I*: Same, *Paléont. Mém.*, no. 29, p. 1-46, 4 text fig., pl. 1-4.
(716c) 1905, *Les rudistes urgoniens, Part II. Série inverse*: Same, *Paléont. Mém.*, no. 29, p. 49-95, 7 text fig., pl. 7-12.
- Paris, E. T.**
(717) 1911, *Notes on some species of Gervillia from the Lower and Middle Jurassic rocks of Gloucestershire*: *Cotteswold Nat. Field Club, Proc.*, v. 17, p. 237-256, pl. 28-29.
- Parona, C. F.**
(718) 1889, *Studio monografico della fauna Rabbiana della Lombardia*: 156 p., 13 pl., Premiata Tipographia Fratelli Fusi (Pavia).
(719) 1901, *Le rudiste e le camacee di S. Polo Matese*: [R.] *Accad. Sci. Torino, Mem., Cl. Sci. Fis. Mat. e Nat.*, ser. 2, v. 50, p. 197-214, pl. 1-3.
(719a) 1908, *Saggio per uno studio sulle caprinidi dei calcari scogliera (orizzonte del Col dei Schiosi) nelle Prealpi Venete orientali*: *R. Accad. Lincei, Mem.*, ser. 5, v. 7, p. 319-346, 30 text fig.
(719b) 1909, *Notizie sulla fauna a rudiste della pietra di Subiaco nella valle dell'Aniene*: *Soc. Geol. Italiana*, v. 27, p. 299-310, 3 text fig., 1 pl.
(720) 1923, *Osservazioni sopra alcune specie della fauna a rudiste del Colle di Medea (Friuli)*: [R.] *Accad. Sci. Torino, Atti, Cl. Sci. Fis. Mat. e Nat.*, v. 59, p. 143-152, 2 text fig.
- Patte, Etienne**
(721) 1926, *Études paléontologiques relatives à la géologie de l'est du Tonkin (Paléozoïque et Trias)*: *Serv. Géol. Indo-Chine, Bull.*, v. 15, pt. 1, 241 p., 12 pl.
- Pchelintsev, V. F.**
(722) 1959, *Rudisty Mezozoya gornogo Kryma*: *Geol. Muzei A. P. Karpinskii, Akad. Nauk SSR*, 178 p., 67 text fig., 43 pl. [*Mesozoic rudistids of the Crimean Range*.]
- Pelseneer, Paul**
(723) 1891, *Contribution à l'étude des lamellibranches*: *Arch. Biologie*, v. 11, p. 147-312, pl. 6-23.
(724) 1906, *Mollusca*: in *A treatise on zoology*, v. 5, E. Ray Lankester (ed.), 355 p., A. & C. Black (London).
(725) 1911, *Les lamellibranches de l'expédition du Siboga, partie anatomique*: *Siboga Expéd. Mon.*, pt. 61, v. 53a, 125 p., 26 pl. (Leiden).
- Peron, Alphonse**
(726) 1906, *Études paléontologiques sur les terrains du département de l'Yonne. Les pélé-*

- cypodes rauraciens et séquaniens*: Soc. Sci. Historiques Nat. l'Yonne, Bull. 59, p. 33-266, pl. 1-11.
- Perrier, J. O. E.**
(727) 1897[1896], *Traité de zoologie*: v. 4, p. 1345-2140, Masson et Cie (Paris).
- Perry, L. M.**
(728) 1940, *Marine shells of the southwest coast of Florida*: Bull. Am. Paleontology, v. 26, no. 95, 260 p., 39 pl.
- , & **Schwengel, J. S.**
(729) 1955, *Marine shells of the western coast of Florida*: Paleont. Research Inst., 318 p., 56 pl., (Ithaca, N.Y.).
- Pfab, Leo**
(730) 1934, *Revision der Taxodonta des böhmischen Silurs*: Palaeontographica, v. 80, Abt. A, p. 195-253, 14 text fig., pl. 2-4.
- Pfannenstiel, Max**
(731) 1928, *Organisation und Entwicklung der Gryphäen*: Palaeobiologica, v. 1, p. 381-418.
- Philippi, Emil**
(732) 1899-1900, *Beiträge zur Morphologie und Phylogenie der Lamellibranchier*: Deutsch. Geol. Gesell., Zeitschr., (a) I. Ueber *Hinnites* und *Velopecten*, v. 50, p. 597-622, pl. 29; (b) II. Zur Stammgeschichte der Pectiniden, v. 52, p. 64-117, fig. 1-24; (c) III. *Lima* und ihre Untergattungen, v. 52, p. 619-639, pl. 24.
- Philippi, R. A.**
(733) 1853, *Handbuch der Conchyliologie und Malacozoologie*: xx + 547 p., Eduard Anton (Halle).
(734) 1887, *Die Tertiären und Quartären Versteinerungen Chiles*: p. 198-199 (*Monopleura Philippi*, novum genus?), pl. xxxvii, fig. 12, F. A. Brockhaus (Leipzig).
- Phillips, John**
(735) 1848, *The Malvern hills, compared with the Palaeozoic districts of Abberley, Woolhope, May Hill, Tortworth, and Usk*: Geol. Survey Great Britain, Mem., v. 2, pt. 1, p. 1-330.
- Pilsbry, H. A.**
(736) 1893, *Notes on the genera of Unionidae and Mutelidae*: Nautilus, v. 7, p. 30.
(737) 1922, *On the nomenclature of certain North American naiades*: in A. E. Ortman & B. Walker, Univ. Michigan, Museum Zoology, Occas. Paper, no. 112, 75 p.
- , & **Bequaert, J.**
(737a) 1927, *The aquatic mollusks of the Belgian Congo*: Am. Museum Nat. History, v. 53, art. 2, p. 69-602, pl. 10-77.
- Ping, Chi**
(738) 1931, *On some new fossil mollusks of Mongolia*: Geol. Soc. China, Bull., v. 10, p. 229-246, 31 text fig., pl. 1-2.
- Pirona, G. A.**
(739) 1867, *Synodontites, nuovo genere di rudiste*: Istit. Veneto Sci., Lett. Arti, Atti, ser. 3, v. 12, p. 833-846, pl. 12.
(740) 1869, *Le ippuritidi del Colle di Medea nel Friuli*: Same, Mem., v. 14, p. 397-435, pl. 17-25.
- Piveteau, Jean**
(741) 1938, *Recherches sur la morphologie des rudistes*: Jour. Conchyliologie, v. 83, p. 1-10, text fig. 1-7.
(741a) 1939, *Recherches sur la morphologie des rudistes*: Same, v. 83, p. 27-34, 5 text fig.
- Pleničar, Mario**
(741b) 1961, *Stratigrafski razvoj Krednih plasti na Južnem Primorskem in Notranjskem*: Geol. Razprave Poročila, v. 6, p. 22-145, 33 text fig., pl. 1-16.
- Počta, Filip**
(741c) 1889, *O rudistech, Vymřelé čeledi mlžů z Českého Křídového Útvaru*: K. České společnosti Nauk, Rozpravy, v. 7, no. 3, p. 1-92, 5 text fig., 1-6 pl.
- Poel, Luc van de**
(742) 1959, *Faune malacologique du Hervien*: Inst. Roy. Sci. Nat. Belgique, Bull., v. 35, no. 15 & 16, p. 1-26, 1-28, 1 pl.
- Pohl, E. R.**
(743) 1929, *The Devonian of Wisconsin, Pt. 1. Lamellibranchiata*: Publ. Museum Milwaukee, Bull., v. 11, p. 1-100, pl. 1-14.
- Pohlig, Hans**
(744) 1880, *Maritime Unionen*: Palaeontographica, v. 27, p. 105-127, pl. 13-14.
- Pojeta, John, Jr.**
(745) 1926, *The pelecypod genus Byssonychia as it occurs in the Cincinnati at Cincinnati, Ohio*: Palaeont. Americana, v. 4, p. 171-216.
(746) 1966, *North American Ambonychiidae (Pelecypoda)*: Same, v. 5, no. 36, 241 p., 47 pl.
- Polšák, Ante**
(746a) 1967[?1968], *Milovanovicia, un nouveau genre de rudistes de la famille des Radiolitidés*: Soc. Géol. France, Bull., ser. 7, v. 9, p. 758-761.
(746b) 1968, *Macrofauna Cretacée de l'Istrie meridionale (Yougoslavie)*: Palaeont. Jugoslavica, v. 8, 210 p., 85 pl.
- Pompeckj, J. F.**
(747) 1901, *Ueber Aucellen und Aucellen-ähnliche Formen*: Neues Jahrb., Beil.-Bd., v. 14, p. 319-366, pl. 15-17.
- Popenoe, W. P.**
(748) 1937, *Upper Cretaceous Mollusca from southern California*: Jour. Paleontology, v. 11, p. 379-402, pl. 45-49.

Powell, A. W. B.

(749) 1935, *New Recent and Tertiary Nuculanidae from New Zealand*: Malacol. Soc. London, Proc., v. 21, p. 252-255, 7 text fig.

Prantl, Ferdinand, Kumpera, O., & Růžicka, Bohuslav

(750) 1960, *Revision of the Nuculanidae from the Ostrava-Karvina district (Pelecypoda)*: Acta Musée Nat. Pragae, p. 17-124.

———, & Růžicka, Bohuslav

(751) 1954, *Straba nov. gen., nový mlž z českého Devonu*: Národ. Musea v. Praze (Acta Musei Nat.), Sborník, v. 10B, p. 1-29, 4 pl. [*Straba nov. gen., a new Devonian pelecypod from Bohemia*.]

(752) 1955, *Mnataia nov. gen., nový mlž z českého Devonu*: Same, v. 11B, no. 5, Geol. & Paleont., no. 1, 17 p., 1 pl. [*Mnataia nov. gen., a new Devonian pelecypod from Bohemia*.]

Prashad, Bains

(753) 1919, *On the generic position of some Asiatic Unionidae*: Indian Museum, Rec., v. 16, p. 403-411, 5 text fig.

(754) 1922, *A revision of the Burmese Unionidae*: Same, v. 24, p. 91-111, pl. 2.

(755) 1932, *The Lamellibranchia of the Siboga expedition, Systematic Part II Pelecypoda Siboga Exped. No. 53 c*: 353 p., 9 pl., E. J. Brill, Ltd. (Leiden).

Preston, H. B.

(756) 1915, *Mollusca (freshwater Gastropoda and Pelecypoda): The fauna of British India, including Ceylon and Burma*: 244 p., 29 text fig. (London).

Přibyl, Alois, & Růžicka, Bohuslav

(757) 1953, *Neklania nov. nom. pro Dalila Barrande, 1881 (Lamellibranchiata)*: Zvlátní Otisk Přírodov. sborn. Ostrav. Kraje, Rozník 14 (1953), c. 21-Tisk, Ostravske tiskarny, n. p. provozovna 05 Opava, p. 57-59.

Pruvost, Pierre

(758) 1930, *La faune continentale du terrain houiller de la Belgique*: Musée Royal Histoire Nat. Belgique, Mém., no. 44, 282 p., 14 pl.

Purchon, R. D.

(759) 1955, *The structure and function of the British Pholadidae (rock-boring Lamellibranchia)*: Zool. Soc. London, Proc., v. 124, p. 859-911.

(760) 1956, *A note on the biology of Brechites penis (L.) (Lamellibranchia)*: Linnean Soc. London (Zool.), Jour., v. 53, p. 43-54.

(761) 1956, *The stomach in the Protobranchia and Septibranchia (Lamellibranchia)*: Zool. Soc. London, Proc., v. 127, p. 511-525.

(762) 1957, *The stomach in the Filibranchia and Pseudolamellibranchia*: Same, v. 129, p. 27-60.

(763) 1958, *The stomach in the Eulamellibranchia; stomach types IV and V*: Same, v. 131, p. 487-525.

(764) 1958, *Phylogeny in the Lamellibranchia*: Cent. & Bicent. Cong. Biol., Proc., Singapore, p. 69-82, 2 tables, 8 text fig.

(765) 1959, *Phylogenetic classification of the Lamellibranchia, with special reference to the Protobranchia*: Malacol. Soc. London, Proc., v. 33, p. 224-230, 3 text fig.

(766) 1963, *Phylogenetic classification of the Bivalvia, with special reference to the Septibranchia*: Same, v. 35, p. 71-80.

Quenstedt, Werner

(767) 1926, *Mollusken aus den Redbay- und Greyhook-Schichten*: Result. Norske Statsunderstøttede Spitsbergeneksped. (Skrifter om Svalbard og Ishavet), no. 11, p. 1-107, 6 text fig., 4 pl., 1 map.

Quoy, J. R. C., & Gaimard, J. P.

(768) 1835, *Voyage de découvertes de d'Astrolabe*: Zoologie, 954 p., 9 pl.

Rafinesque, C. S.

(769) 1819, *Prodrome de 70 nouveaux genres d'animaux découverts dans l'intérieur des États-Unis d'Amérique, durant l'année 1819*: Jour. Physique Chimie Histoire Nat. Élément. (Paris), v. 88, p. 423-428.

(770) 1820, *Monographie des coquilles bivalves fluviales de la rivière Ohio, contenant douze genres et soixante-huit espèces*: Ann. Gén. Sci. Phys. Bruxelles, v. 5, p. 287-322, pl. 80-82.

(771) 1831, *Continuation of a monograph of the bivalve shells of the River Ohio and other rivers of the western states*: 8 p. (Brussels).

Raincourt, J. B. P. de

(772) 1877, *Descriptions d'espèces nouvelles du bassin de Paris*: Soc. Géol. France, Bull., ser. 3, v. 5, p. 329-333, pl. 4.

Ranson, Gilbert

(773) 1942, *Note sur la classification des Ostreidés*: Soc. Géol. France, Bull., v. 5, no. 12, p. 161-164.

Rawitz, Bernhard

(774) 1887, *Das zentrale Nervensystem der Acephalen*: Jena Zeitschr. Naturwiss., v. 20, p. 384-460, pl. 25-29.

Raymond, P. E.

(775) 1916, *The Pelecypoda of the Chazy formation*: Carnegie Museum, Annals, v. 10, p. 325-343, pl. 29-30.

Redlich, K. A.

(776) 1899, *The Cambrian fauna of the eastern Salt-Range*: Geol. Survey India, Mem.,

- Palaeont. Indica (new ser.), Mem. 1, art. 1, p. 1-12, pl. 1.
- Reed, F. R. C.**
 (777) 1902, *Woodwardian Museum notes: Salter's undescribed species*: VII, Geol. Mag., ser. 2, dec. 4, v. 9, p. 145-148, pl. 7.
 (778) 1915, *Palaeontologia Indica*: Geol. Survey India, Mem. (new ser.), v. 6, Mem., no. 1, p. 1-98, pl. 1-12.
 (779) 1924, *Some new Ordovician and Silurian fossils*: Geol. Mag., v. 61, p. 25-31, pl. 3.
 (780) 1927, *Palaeontological notes on the Silurian inlier of Woolhope*: Geol. Soc. London, Quart. Jour., v. 83, p. 531-550, 7 text fig., pl. 41, 42.
 (781) 1934, *Downtonian fossils from the Anglo-Welsh area*: Same, v. 90, p. 571-584, pl. 20, 21.
 (782) 1946, *Notes on some lamellibranchs from Quarrel Hill, Girvan*: Geol. Mag., v. 83, p. 201-205, pl. 14.
- Reeve, L. A.**
 (783) 1856-58, *Conchologia Iconica*: v. 10, *Mytilus*, 9 pl., *Modiola*, 9 pl., Reeve Brothers (London).
- Reinhart, P. W.**
 (784) 1935, *Classification of the pelecypod family Arcidae*: Musée Royal Histoire Nat. Belgique, Bull., v. 11, no. 13, p. 1-68.
- Rennie, J. V. L.**
 (785) 1836, *Lower Cretaceous Lamellibranchia from northern Zululand*: South African Museum, Ann., v. 31, p. 277-391, pl. 37-55.
- Retzius, A. J.**
 (786) 1788, *Dissertatio historico-naturalis sistens nova testaceorum genera*: 23 p. (Lund).
- Rice, E. L.**
 (787) 1897, *Die systematische Verwertbarkeit der Kiemen bei den Lamellibranchiaten*: Jena Zeitschr. Naturwiss., v. 31, p. 29-89, pl. 3-4.
- Ridewood, W. G.**
 (788) 1903, *On the structure of the gills of lamellibranchs*: Royal Soc. London, Philos. Trans., B, v. 194, p. 147-284.
- Risso, J. A.**
 (789) 1826, *Histoire naturelle des principales productions de l'Europe méridionale*: v. 4, viii + 439 p., 12 pl., F. G. Levrault (Paris).
- Rochebrune, A. T. de**
 (790) 1886, *Sur quelques lamellibranches nouveaux provenant du Congo et de ses tributaires*: Soc. Malacol. France, Bull., v. 3, p. 1-14, pl. 1.
 (791) 1904, *Sur deux genres nouveaux d'Unionidae propres à la Sénégambie*: Muséum Histoire Nat. Paris, Bull., v. 10, p. 460-463, text fig. 1-2.
- Röding, P. F.**
 (792) 1798, *Museum Boltenianum sive Catalogus cimeliorum e tribus regnis naturae quae olim collegerat Joa. Fried. Bolten, M.D.p.d., Pars Secunda*: viii + 119 p., Typis Johan Christi Trapii (Hamburg).
- Roger, Jean**
 (793) 1939, *Le genre Chlamys dans les formations néogènes de l'Europe. Conclusions générales sur la répartition géographique et stratigraphique des pectinidés du tertiaire récent*: Soc. Géol. France, Mém. 40, new ser., v. 17, no. 2-4, p. 1-294, text fig. 1-113, pl. 1(6)-28(33).
- Rogers, M. J.**
 (794) 1965, *A revision of the species of nonmarine Bivalvia from the upper Carboniferous of eastern North America*: Jour. Paleontology, v. 39, no. 4, 663-686, pl. 83-85.
- Rollier, Louis**
 (795) 1912-17, *Fossiles nouveaux ou peu connus des terrains secondaires (Mésozoïque) du Jura*: Schweiz. Palaeont. Gesell., Abhandl., (a) pt. 2 (1912), v. 38, p. 35-146, pl. 5-12; (b) pt. 4 (1914), v. 40, p. 321-343; (c) pt. 5 (1917), v. 42, p. 503-696, pl. 33-40.
 (796) 1923, *Rectification de nomenclature*: Rev. Critique Paléozoologie, v. 27, p. 67.
- Rosewater, Joseph**
 (797) 1965, *The family Tridacnidae in the Indo-Pacific*: Indo-Pacific Mollusca, v. 1, no. 6, p. 347-396, pl. 263-293 (Philadelphia).
- Rossmassler, E. A., & Kobelt, Wilhelm**
 (798) 1835-1915, *Inconographie der Land- und Süßwasser-Mollusken, mit vorzüglicher Berücksichtigung der europäischen noch nicht abgebildeten Arten*: v. 1-3, by E. A. Rossmassler, Leipzig (1835-54); v. 4-7 & 2 (1-20), by W. Kobelt, Wiesbaden (1876-1915), Arnold (Dresden, Leipzig), Kreidel (Wiesbaden).
- Roth, R., Newell, N. D., & Burma, B. H.**
 (799) 1941, *Permian pelecypods in the lower Quartermaster Formation, Texas*: Jour. Paleontology, v. 15, p. 312-317, pl. 45.
- Roth von Telegd, K.**
 (800) 1914(1915), *Ein oberoligozäne Fauna aus Ungarn*: Geologica Hungarica, v. 1, pt. 1, p. 5-77.
- Rouault, Marie**
 (801) 1851, *Mémoire sur le terrain paléozoïque des environs de Rennes*: Soc. Géol. France, Bull., sér. 2e, v. 8, p. 358-399, 8 text fig.
- Rowland, H. I. T.**
 (802) 1938, *A preliminary survey of the nomenclatural units of the Tertiary Pectinidae*: Musée Royal Histoire Nat. Belgique, Bull., v. 14, no. 49, p. 1-9.

Roy, S. K.

- (803) 1941, *The upper Ordovician fauna of Fro-bisher Bay, Baffin Land*: Field Museum Nat. History, Geol., Mem., v. 2, p. 1-212, 146 text fig.

Ruedemann, Rudolf

- (804) 1912, *The lower Siluric shales of the Mohawk Valley*: New York State Museum, Bull. 162, 151 p., 30 text fig., 10 pl.
 (805) 1926, *The Utica and Lorraine formations of New York, Part 2, Systematic paleontology, No. 2, Mollusks, crustaceans and eurypterids*: Same, no. 272, p. 5-227, 26 text fig., pl. 1-28.

Runnegar, Bruce

- (805a) 1965, *The bivalves Megadesmus Sowerby and Astartila Dana from the Permian of Eastern Australia*: Geol. Soc. Australia, v. 12, pt. 2, p. 227-252, fig. 1-2, pl. 12-15.
 (806) 1966, *Systematics and biology of some desmodont bivalves from the Australian Permian*: Geol. Soc. Australia, Jour., v. 13, no. 2, p. 373-386.

Russell, L. S.

- (807) 1932, *Mollusca from the McMurray Formation of northern Alberta*: Royal Soc. Canada, Trans., ser. 3, v. 26, sec. 4, p. 37-44, pl. 1.
 (808) 1934, *Reclassification of the fossil Unionidae (fresh-water mussels) of western Canada*: Canad. Field Nat., v. 48, p. 1-4.
 (809) 1935, *Fauna of the upper Milk River beds, southern Alberta*: Royal Soc. Canada, Trans., ser. 3, v. 29, sec. 4, p. 115-128, pl. 1-5.

Rutten, M. G.

- (810) 1936, *Rudistids from the Cretaceous of northern Santa Clara Province, Cuba*: Jour. Paleontology, v. 10, p. 134-142, 4 text fig.

Růžicka, Bohuslav

- (811) 1949, *Pteriidae Thiele Českého siluru a devonu (Lamellibranchiata)*: Sborník Vysoké školy banské v Ostravě, příloha Přírod. sborník Ostrav. Kraje 1949, c. 3, p. 97-118, 5 text fig.
 (812) 1949, *Pterineidae Dall du Silurien et du Dévonien tchèques (Lamellibranchiata)*: Acad. Tchèque Sci., Bull. Internatl., 1949; extr. texte tchèque publié dans la revue Rozpravy II, tridy Ceske akad., v. 1e, no. 13, p. 1-27, 1 text fig., 1 pl.
 (813) 1950, *Leiopteriidae Maillieux Českého siluru a devonu (Lamellibranchiata)*: Czech., Stat. Geol. Ustav., Sborník, v. 17, Paleont. ser., 55 p., 1 pl. [*Leiopteriidae Maillieux of the Siluro-Devonian of Bohemia (Lamellibranchiata)*.]
 (814) 1951, *Cypricardinia Hall Českého siluru (Lamellibranchiata)*: Přírod. Sborník Ostrav. Kraje (Acta Rerum Naturalium Dis-

trictus Ostraviensis), v. 12, no. 3, p. 304-312. [*Cypricardinia Hall of the Silurian of Bohemia (Lamellibranchiata)*.]

- (815) 1951, *Pectinidae Lamarck Českého siluru a devonu*: Same, v. 12, no. 2, p. 170-177. [*Pectinidae Lamarck of the Siluro-Devonian of Bohemia*.]

———, & Prantl, Ferdinand

- (816) 1960, *Types of some of Barrande's pelecypods (Barrandian)*: Casopis Národ. Musea (Praze), v. 1, p. 48-55.
 (817) 1960, *The Silurian representatives of the genus Paracyclas Hall, 1843, from Bohemia (Pelecypoda)*: Acta Musei Nat. Pragae, v. 16B, no. 5, p. 201-223.
 (818) 1961, *Boiomytilus nov. gen., new Devonian pelecypod from Bohemia*: Same, Sborník, v. 17, p. 74-80.

———, & Hajkr, O.

- (819) 1961, *Contribution to the study of the ontogeny of nuculanid pelecypods*: Acta Musei Nat. Pragae, v. 17, p. 57-73.

———, Přibyl, Alois, & Prantl, Ferdinand

- (820) 1959, *Some pectinoid pelecypods from the Silurian and Devonian of central Bohemia*: Acta Musei Nat. Pragae, v. 15, B, no. 1, p. 1-47, pl. 1-11.

Sacco, Federico

- (821) 1897-98, *I molluschi dei terreni terziari del Piemonte e della Liguria* (a) pt. 24 (1897), Pectinidae; p. 1-116, pl. 1-21; (b) pt. 30 (1897), p. 1-203, pl. 1-31; (c) pt. 26 (1898), Fam. Arcidae, Pectunculidae, Limopsidae, Nuculidae, Ledidae, Malletidae, 70 p., 12 pl. (Torino).

Salisbury, A. E.

- (822) 1934, *On the nomenclature of Tellinidae*: Malacol. Soc. London, Proc., v. 21, no. 2, p. 74-91, pl. 9-14.

Salter, J. W.

- (823) 1864, *Note on the fossils from the Budleigh Salterton pebble-bed*: Geol. Soc. London, Quart. Jour., v. 20, p. 286-302, pl. 15-16.

Sandberger, C. L. F. von

- (824) 1870-75, *Die Land- und Süßwasser-Conchylien der Vorwelt*: viii + 1000 p., 36 pl. (Wiesbaden).

Sandberger, Guido, & Sandberger, F. C.

- (825) 1850-56, *Die Versteinerungen des rheinischen Schichten-Systems in Nassau*: xiv + 564 p., atlas of 39 pl. (Wiesbaden).

Sardeson, F. W.

- (826) 1896, *The Saint Peter sandstone*: Minnesota Acad. Nat. Sci., Bull., v. 4, p. 64-88, pl. 2-4.
 (827) 1939, *Early pelecypod Vanuxemia in Minnesota*: Pan Am. Geologist, v. 71, p. 283-293, pl. 19.
 (828) 1939, *Early pelecypod Cyrtodonta in Minnesota*: Same, v. 71, p. 337-346, pl. 21.

Sars, G. O.

- (829) 1878, *Bidrag til kundskaben om Norges arktiske fauna: I. Mollusca regionis arcticae norvegiae. Oversigt over de i Norges arktiske region forekommende bløddyr*: 466 p., 52 pl., A. W. Brøgger (Christiania).

Savage, T. E.

- (830) 1917, *Stratigraphy and paleontology of the Alexandrian series in Illinois and Missouri*: Illinois State Geol. Survey, Bull., no. 23, p. 67-160, pl. 3-9.

Say, Thomas

- (831) 1818, *Description of a new genus of freshwater bivalve shells*: Acad. Nat. Sci. Philadelphia, Jour., v. 1, p. 459-460.

Schenck, H. G.

- (832) 1934, *Classification of nuculid pelecypods*: Musée Royal Histoire Nat. Belgique, Bull., v. 10, no. 20, p. 1-78, pl. 1-5.
- (833) 1936, *Nuculid bivalves of the genus Acila*: Geol. Soc. America, Spec. Paper, no. 4, 149 p., 18 pl.

Schlosser, Max

- (834) 1902, *Die Fauna des Lias und Dogger in Franken und der Oberpfalz*: Deutsch. Geol. Gesell., Zeitschr., v. 53, p. 513-569, pl. 16.

Schnarrenberger, C.

- (834a) 1901, *Ueber die Kreideformation der Monte d'Ocre-Kette in den aquilaner Abruzzen*: Naturforsch. Gesell., Freiburg i. Br., Berichte, v. 11, p. 176-214, pl. 1-4.

Schumacher, C. F.

- (835) 1816, *Over conchyliologiske systemer*: K. Dansk. Vidensk. Selsk., Förhandl., p. 7.
- (836) 1817, *Essai d'un nouveau système des habitations des vers testacés*: 287 p., 22 pl. (Copenhagen).

Scopoli, J. A.

- (837) 1777, *Introductio ad historiam naturalem sistentes genera lapidum, plantarum et animalium, hactenus detecta, caracteribus essentialibus donata, in tribus divisa, subinde ad leges naturae*: 506 p. (Prague).

Seguenza, Giuseppe

- (838) 1877, *Nuculidi terziarie rinvenute nelle provincie meridionali d'Italia*: Accad. Lincei, Atti, ser. 3, v. 1, p. 1163-1200.

Sénesse, Pierre

- (839) 1939, *Rhedensia, serie d'hippurites intermédiaire entre Orbignya et Vaccinites*: Soc. Histoire Nat. Toulouse, Bull., v. 73, p. 226-249, 32 text fig., pl. 4.

Servain, G.

- (840) 1890, *Des acéphales lamellibranches fluviales du système européen*: Soc. Malacol. France, Bull., v. 7, p. 281-323, pl. 5-7.

Sharpe, Daniel

- (841) 1853, *Description of Zoophyta and Mol-*

lusca: in Carlos Ribeiro, On the Carboniferous and Silurian formations of the neighbourhood of Bussaco in Portugal, Geol. Soc. London, Quart. Jour., v. 9, p. 146-158, pl. 7-9.

- (842) 1856, *Description of fossils from the secondary rocks of Sunday River and Zwartkop River*: Geol. Soc. London, Trans., v. 2, no. 7, p. 193-203, pl. 22, 23, 28.

Sieverts [Sieverts-Doreck], Hertha

- (843) 1933, *Beiträge zur Paläontologie des Ostindischen Archipels. IX. Jouannetia cumingi (Sowerby) aus dem Pliocän von Timor*: Neues Jahrb. Mineralogie, Geologie, Paläontologie, Abt. B, Beilage-Band 71, p. 267-302, text fig. 1-2.

Simpson, C. T.

- (844) 1898, [no title] in F. C. Baker, The Mollusca of the Chicago area. The Pelecypoda, Chicago Acad. Sci. Bull., v. 3, pt. 1, p. 1-130, pl. 1-27.
- (845) 1900, *Synopsis of the najades or pearly freshwater mussels*: U. S. Natl. Museum, Proc., v. 22, p. 501-1044, pl. 18.
- (846) 1912, *A descriptive catalogue of the naiades, or pearly fresh-water mussels*: 1540 p. (Detroit).

Simpson, G. G.

- (847) 1945, *The principles of classification and a classification of mammals*: Am. Museum Nat. History, Bull., v. 85, 350 p.

Slišković, T.

- (847a) 1966, *Katzeria hercegovinaensis n. gen. n. sp. aus den obersenenonischen Ablagerungen der Südherzegowina*: Sci. Conseil. Acad. RSF Yougoslav., Bull., sec. A., v. 2, p. 176-177, 2 text fig.

Slodkevich [Slodkewitsch], W. S.

- (848) 1938, *Tretichnye peletsipody Dal'nego Vostočka*: Acad. Sci. USSR, Paleont. Inst., Paleont. USSR, v. 10, pt. 3, no. 18, 508 p. [Tertiary Pelecypoda from the Far East, Part I, Paleontology of USSR.]

Smith, E. A.

- (849) 1875, *Descriptions of some new shells from Kerguelen's Island*: Ann. & Mag. Nat. History, ser. 4, v. 16, p. 67-73.
- (850) 1879, *Mollusca*: Royal Soc. London, Philos. Trans., v. 168, p. 167-192, pl. 9.
- (851) 1884, *Report on the zoological collections made in the Indo-Pacific during the voyage of H. M. S. 'Albert' 1881-82, Mollusca*: British Museum (Nat. History), Rept., p. 34-116, pl. 4-7.
- (852) 1885, *Report on the Lamellibranchiata collected by H. M. S. Challenger during the years 1873-1876*: Rept. Scientific Results Explor. Voyage H. M. S. Challenger, Zoology, v. 13, p. 240.

Snethlage, Emilie

- (853) 1905, *Ueber die Gattung Joufia G. Boehm*: Naturforsch. Gesell. Freiburg i. Br., Berichte, v. 16, p. 1-9, 2 text fig., 2 pl.

Solem, Alan

- (854) 1954, *Living species of the pelecypod family Trapeziidae*: Malacol. Soc. London, Proc., v. 31, pt. 2, p. 64-86, 2 pl.

Soot-Ryen, Helen

- (855) 1966, *The Middle Ordovician of Norway. 21. A new pelecypod of the genus Palaeosolen Hall, 1885*: Norsk Geol. Tidsskrift, v. 46, p. 21-26.

Soot-Ryen, Tron

- (856) 1832, *Pelecypoda, with a discussion of possible migrations of Arctic pelecypods in Tertiary times*: Norweg. North Polar Exped. with the "Maud," 1918-25, Sci. Results, v. 5, no. 12, p. 1-35, pl. 1-2.

- (857) 1941, *Northern pelecypods in the collection of Tromsø Museum. 1. Order Anomalodesmacea, families Pholadomyidae, Thraciidae and Periplomatidae*: Tromsø Museum Aarshefter, Naturhist. Avd. no. 17, v. 61 (1938), no. 1, 41 p., 10 pl.

- (858) 1951, *Scientific results of the Norwegian Antarctic expeditions 1927-28 no. 32, Antarctic pelecypods*: Norske Videnskapsakad., no. 32, 46 p., 13 text fig., 1 pl.

- (859) 1953, *Choromytilus, a new genus in the Mytilidae*: Revista Soc. Malacol. Carlos de la Torre, v. 8, no. 3, p. 121-122.

- (860) 1955, *A report on the family Mytilidae (Pelecypoda)*: Allan Hancock Pacific Exped., v. 20, no. 1, 1,175 p., 78 text fig., 10 pl.

Sowerby, G. B.

- (861) 1833, *Characters of new species of Mollusca and Conchifera collected by Mr. Cuming*: Zool. Soc. London, Proc., 1832, p. 194-202.

- (862) 1864-68, *Monograph of the genus Unio*: in L. A. Reeve, *Conchologia Iconica*, v. 16, 96 pl.

- (863) 1867-70, *Monograph of the genus Anodon*: in L. A. Reeve, *Conchologia Iconica*, v. 17, 37 pl. with text.

- (864) 1868, *Monograph of the genus Mycetopus*: in L. A. Reeve, *Conchologia Iconica*, v. 16, 4 pl. with text.

- (865) 1868, *Monograph of the genus Iridina*: in L. A. Reeve, *Conchologia Iconica*, v. 16, 2 pl. with text.

- (866) 1869, *Monograph of the genus Hyria*: in L. A. Reeve, *Conchologia Iconica*, v. 17, 5 pl. with text.

- (867) 1869, *Monograph of the genus Castalia*: in L. A. Reeve, *Conchologia Iconica*, v. 17, 3 pl. with text.

- (868) 1872, *Monograph of the genus Etheria*: in

L. A. Reeve, *Conchologia Iconica*, v. 18, 2 pl. with text.

- (869) 1872, *Monograph of the genus Mulleria and Barilettia*: in L. A. Reeve, *Conchologia Iconica*, v. 18, 1 pl. with text.

Sowerby, J. de C.

- (870) 1823-25, *The mineral conchology of Great Britain*: v. 5, 168 p., pl. 407-503, the author (London).

Spriestersbach, Julius

- (871) 1915, *Neue oder wenig bekannte Versteinerungen aus dem rheinischen Devon, besonders aus dem Lenneschiefer*: Preuss. Geol. Landesanst., Abhandl., new ser., no. 80, 80 p., 28 pl.

- (872) 1919, *Neue Versteinerungen aus dem Lenneschiefer*: Same, Jahrb., v. 38, pt. 1, p. 434-512, 3 text fig., pl. 24-39.

- (873) 1920, *Die Stellung von Montanaria Spriestersbach und Crassatellopsis Beushausen*: Same, Jahrb., v. 39, pt. 1, p. 32-40, pl. 3.

- (874) 1925, *Die Oberkoblenszschichten des Bergischen Landes und Sauerlandes*: Same, Jahrb., v. 45, p. 367-450, 1 text fig., pl. 10-17.

———, & Fuchs, Alexander

- (875) 1909, *Die Fauna der Remscheider Schichten*: Preuss. Geol. Landesanst., Abhandl., new ser., no. 58, p. 1-81, 11 pl.

Staesche, K.

- (876) 1926, *Pectiniden des schwäbischen Jura*: Geol. Paläont. Abhandl., new ser., v. 15, p. 1-136, pl. 1-6.

Stanton, T. W.

- (877) 1897, *On the genus Remondia Gabb, a group of Cretaceous bivalve mollusks*: U.S. Natl. Museum, Proc., v. 19, p. 299-301, pl. 26.

Stasek, C. R.

- (878) 1961, *The ciliation and function of the labial palps of Acila castrensis (Protobranchia, Nuculidae)*: Zool. Soc. London, Proc., v. 137, no. 4, p. 511-538.

- (879) 1963, *Orientation and form in the bivalved Mollusca*: Jour. Morphology, v. 112, no. 3, p. 195-214.

- (880) 1963, *Synopsis and discussion of the association of ctenidia and labial palps in the bivalved Mollusca*: The Veliger, North Calif. Malacozool. Club, Quart., v. 6, no. 2, p. 91-97.

Stefanescu, Sabba

- (881) 1889, *Études sur les terrains tertiaires de la Roumanie. Contribution à l'étude des faunes sarmatiques, pontiques et levantins*: Soc. Géol. France, Paléont. Mém., v. 4, no. 15, 147 p., 11 pl.

Stefanini, Giuseppe

- (882) 1939, *Molluschi del Guiralias della Somalia*.

- Gasteropodi e Lamellibranchi*: Palaeont. Italica, v. 32, suppl. 4, p. 103-270, pl. 13-27.
- Stefano, Giovanni di**
 (882a) 1889, *Studi stratigrafici i palaeontologici sul sistema cretaceo della Sicilia I. Gli strati con Caprotina di Termini-Imerese*: R. Accad. Sci. Palermo, Atti, v. 10, p. 1-44, pl. 1-11.
 (882b) 1899, *Studi stratigrafici e paleontologici sul sistema Cretaceo della Sicilia*: Palaeont. Italica, Mem. Paleont., v. 4, p. 1-46, pl. 1-5.
- Steinmann, Gustav**
 (883) 1903, *Einführung in die Paläontologie*: ix + 446 p. (Leipzig).
- Stempell, Walter**
 (884) 1898, *Beiträge zur Kenntnis der Nuculiden*: Zool. Jahrb., Jena, suppl. v. 4 (Fauna Chilensis), p. 339-430, pl. 22-25.
- Stenzel, H. B.**
 (885) 1947, *Nomenclatural synopsis of supraspecific groups of the family Ostreidae (Pelecypoda, Mollusca)*: Jour. Paleontology, v. 21, no. 2, p. 165-185.
- Stephenson, L. W.**
 (886) 1922, *Some Upper Cretaceous shells of the rudistid group from Tamaulipas, Mexico*: U.S. Natl. Museum, Proc., v. 61, art. 1, p. 1-28, 15 pl.
 (887) 1923, *The Cretaceous formations of North Carolina. Pt. I*: North Carolina Geol. & Econ. Survey Publ., v. 5, 604 p., 102 pl.
 (888) 1934, *The genus Diploschiza from the Upper Cretaceous of Alabama and Texas*: Jour. Paleontology, v. 8, no. 3, p. 273-280, pl. 38.
 (889) 1941, *The larger invertebrate fossils of the Navarro Group of Texas*: Univ. Texas Publ. 4101, 641 p., 95 pl.
 (890) 1952, *Larger invertebrate fossils of the Woodbine Formation (Cenomanian) of Texas*: U.S. Geol. Survey, Prof. Paper 242, 226 p., 59 pl.
- Stewart, B. H.**
 (891) 1920, *The stratigraphy and palaeontology of Toronto and vicinity, Part 1, The Pelecypoda*: Ontario Dept. Mines, 29th Ann. Rept., v. 29, pt. 6, p. 5-58, pl. 1-5.
- Stewart, R. B.**
 (892) 1930, *Gabb's California Cretaceous and Tertiary type lamellibranchs*: Acad. Nat. Sci. Philadelphia, Spec. Publ., no. 3, p. 1-314, text fig. 1-5, pl. 1-17.
- Stoliczka, Ferdinand**
 (893) 1870-71, *Cretaceous fauna of southern India, v. 3, The Pelecypoda, with a review of all known genera of this class, fossil and Recent*: Geol. Survey India, Palaeont. Indica, ser. 6, v. 3, 537 p., 50 pl.
- Strand, Embrik**
 (894) 1932, *Miscellanea nomenclatoria zoologica et palaeontologica*: Folia Zool. Hydrobiol., v. 4, p. 133-147.
- Strong, A. M., & Hertlein, L. G.**
 (895) 1937, *New species of Recent mollusks from the coast of western North America*: California Acad. Sci., Proc., ser. 4, no. 22, p. 159-178.
- Suzuki, Koiti**
 (896) 1941, *A new naiad Unio (Nipponaias) ryosekiana n. subgen., and n. sp., from the Lower Cretaceous of Japan*: Geol. Soc. Japan, Jour., v. 48, p. 410-413, text fig. 1-3.
- , & Oyama, K.
 (897) 1943, *Ueberblick über die Corbiculiden Ostasiens*: Venus, Japan Jour. Malacology, v. 12, no. 3-4, p. 138-149.
- Swainson, William**
 (898) 1840, *A treatise on malacology or shells and shell-fish*: The Cabinet Cyclopaedia, Nat. History, 420 p., 130 text fig., Longman (London).
- Swartz, C. K., & Prouty, W. F.**
 (899) 1923, *Pelecypoda*: in Systematic paleontology of Silurian deposits, Maryland Geol. Survey, Silurian vol., p. 464-481, pl. 27, 28.
- Taki, Iwao, & Habe, Tadashige**
 (900) 1955, *Pholadidae in Japan*: Illustrated catalogue of Japanese shells, Tokubei Kuroda (ed.), v. 2, no. 2, p. 7-15, pl. 2 (Kyoto, Japan).
- Tamura, Minoru**
 (901) 1959, *Trigoniidae, Ostreidae, Bakevelliidae, Pteriidae, Cardiidae and Astartidae from the Upper Jurassic Sakamoto Formation in central Kyushu, Japan*: Palaeont. Soc. Japan, Trans. Proc., new ser., no. 33, p. 23-32.
- Tasch, Paul**
 (902) 1953, *Cause and paleoecological significance of dwarfed fossil marine invertebrates*: Jour. Paleontology, v. 27, p. 356-444, 6 text fig., pl. 49.
- Tate, Ralph**
 (903) 1867, *On the oldest known species of Exogyra, with a description of the species*: Geol. Nat. Hist. Repertory, v. 1, p. 378-380.
- Tavani, Guido**
 (904) 1949, *Rudiste ed altri molluschi cretacei della Migiurtinia (Africa orientale)*: Palaeont. Italica, v. 46, new ser., v. 16, p. 1-40, text fig. 1-2, pl. 1-4.
 (905) 1958, *Rudiste del Cretaceo delle Puglie (Italia Meridionale)*: Palaeont. Soc. India, Jour., v. 3, p. 170-177, pl. 26-29.

Tebble, Norman

- (905a) 1966, *British bivalve seashells*: 212 p., 110 text fig., trustees of the British Museum (Natural History) (London).

Teppner, W. von

- (906) 1922, *Fossilium catalogus I: Animalia, C. Diener (ed.). Pars. 15: Lamellibranchiata tertiaria "Anisomyaria."* II: p. 67-296, W. Junk (Berlin).

Thiadens, A. A.

- (907) 1936, *On some caprinids and a mono-pleurid from southern Santa Clara, Cuba*: [K.] Ned. Akad. Wetensch. Amsterdam, v. 39, Proc., no. 9, p. 1-12, text fig. 1-5.

Thiele, Johannes

- (908) 1935, *Handbuch der systematischen Weichtierkunde*: v. 2, p. 779-1154, text fig. 784-897, Gustav Fischer (Jena).

Tomlin, J. R. le B.

- (909) 1928, *Pseudavicula Simpson preoccupied*: Nautilus, v. 42, p. 66.

Toucas, Aristide

- (910) 1903-04, *Études sur la classification et l'évolution des hippurites*: Soc. Géol. France, Mém., Paléont. Mém., no. 30, 128 p., 17 pl.
(911) 1907-09, *Études sur la classification et l'évolution de Radiolitidae*: Same, Mém., no. 36, 132 p., 24 pl.

Trechmann, C. T.

- (911a) 1924, *The Cretaceous limestones of Jamaica and their Mollusca*: Geol. Mag., v. 61, no. 9, p. 385-409, 1 text fig., pl. 22-26.

Tremlett, W. E.

- (912) 1953, *English Eocene and Oligocene Veneridae, Parts 1, II*: Malacol. Soc. London, Proc., v. 30, pt. 1, p. 1-21, pl. 1-4; pt. 3, p. 55-71, pl. 9-14.

Troedsson, G. T.

- (913) 1929, *On the Middle and Upper Ordovician faunas of northern Greenland*: Pt. II, Meddel. Grønland, v. 72, p. 1-197, 56 pl.

Trueman, A. E.

- (914) 1938, *On certain Anthracomyas from the Similis-Pulchra Zone of the Coal Measures*: Geol. Mag., v. 75, p. 459-469, text fig. 1-3.

, & Weir, John

- (915) 1945-60, *A monograph of the British Carboniferous non-marine Lamellibranchia*: Palaeontograph. Soc. London, (a) Part 1 (1945), p. i-xxxii, 1-18, pl. 1-4; (b) Part 2 (1946), p. 19-44, pl. 5-7; (c) Part 3 (1947), p. 45-74, pl. 8-12; (d) Part 4 (1948), p. 75-100, pl. 13, 14; (e) Part 5 (1951), p. 101-152, pl. 15-19; (f) Part 6 (1952), p. 153-170, pl. 20-22; (g) Part 7 (1954), p. 171-206, pl. 23-26; (h), Part 8 (1955), p. 207-242, pl. 27-29; (i) Part 9 (1956), p. xxxiii-xxxiv, 243-271, pl. 30-31.

Trueman, E. R.

- (916) 1952, *Observations on the ligament of Nucula*: Malacol. Soc. London, Proc., v. 29, p. 201-205.

Tryon, G. W.

- (917) 1862, *On the classification and synonymy of the Recent species of Pholadidae*: Acad. Nat. Sci. Philadelphia, Proc., p. 191-221.
(918) 1862, *Monograph of the family Teredidae*: Same, Proc., p. 453-482.

Tullberg, S. A.

- (919) 1881, *Ueber Versteinerungen aus den Aucellen-Schichten Novaja-Semljas*: Bihang Svenska Vetenskapsakad. Handl., v. 6, no. 3, p. 1-25, pl. 1-2.

Turner, R. D.

- (920) See 139a.
(921) 1954-55, *The family Pholadidae in the western Atlantic and the eastern Pacific*: Same, (a) pt. I (1954), Pholadinae, v. 3, no. 33, p. 1-64, pl. 1-34; (b) pt. II (1955) Martesiinae, Jouannetiinae and Xylophaginae, v. 3, no. 34, p. 65-160, pl. 35-93.
(922) 1962, *Nettastomella japonica Yokoyama in North America and notes on the Pholadidae*: Harvard Univ., Museum Comp. Zoology, Occas. Papers on Mollusks, v. 2, p. 289-308, pl. 47-53.
(923) 1966, *A survey and illustrated catalogue of the Teredidae*: Same, spec. publ. 265 p., 25 text fig., 64 pl.

Twenhofel, W. H.


- (924) 1928, *Geology of Anticosti Island*: Geol. Survey of Canada, Mem. 154, 481 p., 1 text fig., 60 pl.
(925) 1938, *Geology and paleontology of the Mingan Islands, Quebec*: Geol. Soc. America, Spec. Papers, no. 11, 132 p., 1 text fig., 24 pl.

Ulrich, E. O.

- (926) 1890, *New Lamellibranchiata, No. 2, on two new genera and six new species*: Am. Geologist, v. 6, p. 173-181, 15 text fig.
(927) 1892, *New Lamellibranchiata, No. 4, descriptions of one new genus and eight new species*: Same, v. 10, p. 96-104, pl. 7.
(928) 1892, *New Lower Silurian Lamellibranchiata, chiefly from Minnesota rocks*: Geol. Nat. History Survey Minnesota, 19th Ann. Rept., p. 211-248, 32 fig.
(929) 1895 (1893), *New and little known Lamellibranchiata from the lower Silurian rocks of Ohio and adjacent states*: Geol. Survey Ohio, Rept., v. 7, pt. 2, p. 627-693, 2 text fig., pl. 45-56.
(930) 1897, *The Lower Silurian Lamellibranchiata of Minnesota*: Geol. Nat. History Survey, Minnesota, Final Rept., v. 3, pt. 2, p. 475-628, text fig. 35-45, pl. 35-46 (abstr.,

- Minnesota Univ. Quart. Bull. 2, p. 89-91, 1894).
- (931) 1924, *Modiodesma*, n. gen. Ulrich, and the genotype of *Modiolopsis* Hall: Geol. Survey Canada, Mem. 138. p. 183-190.
- Utterback, W. I.**
(932) 1915-16, *The naiades of Missouri*: Am. Midland Naturalist, v. 4, p. 41-53, 97-152, 182-204, 244-273. 311-327, 339-354, 387-400, 432-464, 23 pl.
- Vautrin, Henri**
(933) 1933, *Sur quelques formes nouvelles de rudistes recueillis en Syrie Septentrionale*: Haut-Comm. Syrie et Liban, Notes et Mém., Républ. Française, p. 29-43, text fig. 5-10, pl. 1-5.
- Verneuil, P. E. P. de**
(934) 1845, *Mollusques*: in R. I. Murchison, E. de Verneuil, & A. de Keyserling; Géologie de la Russie d'Europe; v. 2, no. 3, Paléontologie, p. 37-376.
(935) 1847, *Note sur le parallélisme des roches dépôts paléozoïques de l'Amérique septentrionale avec ceux de l'Europe, suivie d'un tableau des espèces fossiles communes aux deux continents, avec l'indication des étages où elles se rencontrent, et terminée par un examen critique de chacune de ces espèces*: Soc. Géol. France, Bull., 2e sér., v. 4, p. 646-710, 3 text fig.
- Verrill, A. E.**
(936) 1897, *A study of the family Pectinidae, with a revision of the genera and subgenera*: Connecticut Acad. Arts & Sci., Trans., v. 10, pt. 1, p. 41-96, pl. 16-21, 1 text fig.
- _____, & **Bush, K. J.**
(937) 1897 (1898), *Revision of the genera of Ledidae and Nuculidae of the Atlantic Coast of the United States*: Am. Jour. Sci., ser. 4, v. 3, no. 153, no. 13, p. 51-63, text fig. 1-22.
(938) 1898, *Revision of the deep-water Mollusca of the Atlantic coast of North America with descriptions of new genera and species*: U.S. Natl. Museum, Proc., v. 20, p. 775-901.
- Vest, Wilhelm von**
(939) 1866, *Über Margaritana bonelli Fer.*: Siebenbürg. Ver. Naturwiss., Verhandl. u. Mitteil., v. 17, p. 193-201, 2 pl.
(940) 1899, *Ueber die Bildung und Entwicklung des Bivalven-Schlusses*: Same, Verhandl. & Mitteil., v. 48, p. 25-153, pl. 1-3.
- Vincent, E. G.**
(941) 1891 (1892), *Contribution à la paléontologie de l'Éocène Belge Pholadidae*: Soc. Royale Malacol. Belgique, Mém., v. 26, p. 164-168, pl. 4.
- Vogel, Klaus**
(942) 1960, *Zu Struktur und Funktion der Siphonalpfeiler der Hippuriten (Lamellibranchiata)*: Paläont. Zeitschr., v. 34, p. 275-294, 6 text fig., pl. 21-25.
(943) 1962, *Muscheln mit Schlosszähnen aus dem spanischen Kambrium und ihre Bedeutung für die Evolution der Lamellibranchiaten*: Akad. Wiss. & Lit. Mainz., Math.-Naturwiss. Kl., Abhandl., Jahrg. 1962, no. 4, p. 193-244, 5 pl.
- Vokes, H. E.**
(944) 1945, *Supraspecific groups of the pelecypod family Corbulidae*: Am. Museum Nat. History, Bull., v. 86, art. 1, p. 1-32, pl. 1-4.
(945) 1946, *Contributions to the paleontology of the Lebanon mountains. Republic of Lebanon; part 3: The pelecypod fauna of the "Olive locality" (Aptian) at Abeih*: Same, Bull., v. 87, art. 3, p. 139-216, text fig. 1, pl. 1-10.
(946) 1949, *The hinge and marginal pectinations of Nuculoidea opima (Hall), type of Nuculoidea Williams and Breger*: Washington Acad. Sci., Jour., v. 39, no. 11, p. 361-363.
(947) 1954, *Some primitive fossil pelecypods and their possible significance*: Same, v. 44, p. 233-236.
(948) 1963, *Studies on Tertiary and Recent giant Limidae*: Tulane Studies in Geology, v. 1, no. 2, p. 73-123.
(949) 1964, *Margaritariidae, new family (Pelecypoda) and description of two new species*: Same, v. 2, no. 4, p. 135-141, 1 pl.
(949a) 1967, *Genera of the Bivalvia: a systematic and bibliographic catalogue*: Bull. Am. Paleontology, v. 51, no. 232, p. 105-394.
- Waagen, Lukas**
(950) 1907, *Die Lamellibranchiaten der Pachycardientuffe der Seiser Alm*: Geol. Reichsanst. Wien, Abhandl., v. 18, no. 2, 180 p., 19 text fig., pl. 25-34.
- Wade, Bruce**
(951) 1926, *The paleontology of the Coon Creek beds of the Ripley Formation*: U.S. Geol. Survey, Prof. Paper 137, 272 p., 72 pl.
- Walcott, C. D.**
(952) 1889, *Stratigraphic position of the Olenellus fauna in North America and Europe*: Am. Jour. Sci., ser. 3, v. 38, p. 29-42.
- Walker, Bryant**
(953) 1917, *The method of evolution in the Unionidae*: Univ. Michigan, Museum Zoology Occas. Paper, v. 45, p. 1-10.
(954) 1918, *A synopsis of the classification of the fresh-water Mollusca of North America, North of Mexico, and a catalogue of the more recently described species, with notes*:

References



- Univ. Michigan, Museum Zoology, Misc. Publ., no. 6, p. 1-213, 233 text fig.
- (955) 1918, *The Mollusca*: in H. B. Ward & G. C. Whipple, Fresh-water biology, p. 957-1020, text fig. 1403-1534, John Wiley & Sons (New York).
- Warmke, G. L., & Abbott, R. T.**
 (956) 1961, *Caribbean shells*: 346 p., 44 pl., Livingstone Publ. Co. (Narberth, Pa.).
- Waterhouse, J. B.**
 (957) 1960, *Some Carnian pelecypods from New Zealand*: Royal Soc. New Zealand, Trans., v. 88, pt. 3, p. 425-442, pl. 19-25.
 (958) 1963, *Etheripecten, a new aviculopectinid genus from the Permian*: New Zealand Jour. Geol. Geophys., v. 6, no. 2, p. 193-196, 5 text fig.
 (959) 1963, *New Zealand species of the Permian bivalve Atomodesma Beyrich*: Palaeontology, v. 6, pt. 4, p. 699-717, pl. 100-105.
 (959a) 1966, *On the validity of the Permian bivalve family Pachydomidae Fischer 1887*: Geol. Soc. Australia, v. 13, no. 2, p. 543-559, pl. 15-17.
- Watson, Hugh**
 (960) 1930, *On the anatomy and affinities of Plicatula*: Malacol. Soc. London, Proc., v. 19, p. 25-30, pl. 5.
- Weigelt, Johannes**
 (961) 1922, *Die Bedeutung der Jugendformen karbonischer Posidonomyen für ihre Systematik*: Palaeontographica, v. 64, p. 43-130.
- Weir, John**
 (962) 1938, *Palaeanodonta from Kenya and Burma*: Hunterian Museum Univ. Glasgow, Mon. 5, p. 12-15, text fig. 1-9, pl. 1.
 (963) 1945, *A review of recent work on the Permian non-marine lamellibranchs and its bearing on the affinities of certain non-marine genera of the Upper Palaeozoic*: Geol. Soc. Glasgow, Trans., v. 20, p. 291-340, text fig. 1-5.
 (964) 1960-66, *A monograph of British Carboniferous non-marine Lamellibranchia*: Palaeontograph. Soc., (a) pt. 10, p. 273-320, pl. 32-33; (b) pt. 11 (1966), p. 321-372, pl. 34-41.
- Weller, Stuart**
 (965) 1903, *A report on the Paleozoic paleontology of New Jersey*: Geol. Survey New Jersey, Rept. Paleontology, v. 3, 462 p., 53 pl.
- Wenz, Wilhelm**
 (966) 1927, *Paläontologischer Teil*: in K. Krejci & W. Wenz, Beobachtungen im Schacht I auf Grube Bucea, Câmpina, Centralbl. Mineralogie, 1927B, p. 418-423, 5 text fig.
 (967) 1928, *Rhombunio Germain, 1911 = Psilunio S. Stefanescu, 1896*: Arch. Molluskenkunde, v. 60, p. 269-270.
- (968) 1942, *Die Mollusken des Pliozäns der rumänischen Erdöl-Gebiete als Leitversteinerungen für die Aufschluss-Arbeiten*: Senckenbergiana, v. 24, 293 p., 71 pl.
- Westerlund, C. A.**
 (969) 1890, *Malacozoa acephala: Fauna der in der palaearktischen Region (Europa Kaukasien, . . .) lebenden Binnenconchylien*: 319 p. (Berlin).
- Whidborne, G. F.**
 (970) 1883, *Notes on some fossils, chiefly Mollusca, from the Inferior Oolite*: Geol. Soc. London, Quart. Jour., v. 39, 1883, no. 156, p. 487-540, pl. 15-19.
- White, K. M.**
 (971) 1942, *The pericardial cavity and the pericardial gland of the Lamellibranchia*: Malacol. Soc. London, Proc., v. 25, p. 37-88, 41 text fig.
- Whiteaves, J. F.**
 (972) 1893 (1894), *Note on the recent discovery of large Unio-like shells in the Coal Measures at the South Joggins, N.S.*: Royal Soc. Canada, Trans., v. 11, sec. 4, p. 21-24, pl. 1.
 (973) 1903, *Description of a new species of Matheria, from the Trenton limestone at Ottawa*: Ottawa Naturalist, v. 17, p. 32-34.
 (974) 1904, *Preliminary list of fossils from the Silurian (Upper Silurian) rocks of the Ekwan River and Sutton Mill Lakes, Keewatin, collected by D. B. Dowling in 1901, with descriptions of such species as appear to be new*: Geol. Survey Canada, Ann. Rept., 24 p.
 (975) 1906, *The fossils of the Silurian (Upper Silurian) rocks of Keewatin, Manitoba, the North Eastern shore of Lake Winnipegosis, and the Lower Saskatchewan River*: Same, Palaeozoic fossils, v. 3, pt. 4, p. 243-298.
 (976) 1908, *Notes on the Pelecypoda or bivalve Mollusca of the Chazy formation in Canada, with descriptions of one new genus and four new species from the Chazy sandstone at the Hog's Back, near Ottawa*: Ottawa Naturalist, v. 22, p. 105-115, pl. 3.
- Whitfield, R. P.**
 (977) 1882, *On the fauna of the Lower Carboniferous limestones of Spergen Hill, Indiana, with a revision of the descriptions of its fossils hitherto published, and illustrations of the species from the original type series*: Am. Museum Nat. History, Bull., v. 1, no. 3, art. 5, p. 39-97, pl. 6-9.
 (978) 1882, *Palaeontology*: Wisconsin Geol. Survey (1873-79), v. 4, pt. 3, p. 161-363.
 (979) 1885, *Brachiopoda and Lamellibranchiata of the Raritan clays and greensand marls of New Jersey*: U.S. Geol. Survey, Mon., no. 9, xx + 338 p.

- (980) 1893, *Contributions to the paleontology of Ohio*: Geol. Survey Ohio, Rept., v. 7, pt. 2 (1893), p. 407-494, 2 text fig., pl. 1-13.
- (980a) 1897, *Descriptions of species of Rudistae from the Cretaceous rocks of Jamaica, W. I.*: Am. Museum Nat. History, Bull., v. 9, art. 12, p. 185-196, pl. 6-22.
- (981) 1897, *Observations on the genus Barretia Woodward, with descriptions of two new species*: Same, Bull., v. 9, art. 20, p. 233-246, 2 text fig., pl. 27-38.
- (982) 1902, *Description of a new Teredo-like shell from the Laramie group*: Same, Bull., v. 16, p. 73-76, pl. 28-29.
- Williams, H. S.**
- (983) 1908, *On the revision of the mollusk genus Pterinea Goldfuss*: U.S. Natl. Museum, Proc., v. 34, p. 83-90.
- (984) 1912, *Some new Mollusca from the Silurian formations of Washington County, Maine*: Same, Bull., v. 42, p. 381-398, pl. 49-50.
- (985) 1913, *New species of Silurian fossils from the Edmunds and Pembroke formations of Washington County, Maine*: Same, Proc., v. 45, p. 319-352, pl. 29-31.
- , & **Breger, C. L.**
- (986) 1916, *The fauna of the Chapman Sandstone of Maine, including descriptions of some related species from the Moose River sandstone*: U.S. Geol. Survey, Prof. Paper no. 89, 347 p., 27 pl.
- Williams, M. Y.**
- (987) 1919, *The Silurian geology and faunas of Ontario Peninsula, and Manitoulin and adjacent islands*: Geol. Survey Canada, Mem. 111, 195 p., 34 pl.
- Wilson, A. E.**
- (988) 1921, *The range of certain lower Ordovician faunas of the Ottawa valley, with descriptions of some new species*: Geol. Survey Canada, Bull. 33, p. 19-57, 7 text fig., 3 pl.
- Wilson, R. B.**
- (989) 1958, *A revision of the Carboniferous lamellibranchs Edmondia punctatella (Jones) and 'Estheria' youngii Jones*: Geol. Survey Great Britain, Bull., no. 15, p. 21-28, 1 pl.
- (990) 1959, *Wilkingia gen. nov. to replace Alorisma for a genus of upper Palaeozoic lamellibranchs*: Palaeontology, v. 1, pt. 4, p. 401-404, pl. 71.
- (991) 1960, *A revision of the types of the Carboniferous lamellibranch species erected by J. Fleming*: Geol. Survey Great Britain, Bull., no. 16, p. 110-126.
- Wilton, C. P. N.**
- (992) 1830, *Memoir on the geology of the shore of the Severn, in the parish of Awre, Gloucestershire*: Quart. Jour. Sci., Lit. Art (Jan.-June), p. 64-73, 1 text fig. (London).
- Winchell, Alexander**
- (993) 1863, *Descriptions of fossils from the yellow sandstone lying beneath the "Burlington Limestone," at Burlington, Iowa*: Acad. Nat. Sci. Philadelphia, Proc., p. 2-25.
- , & **Marcy, Oliver**
- (994) 1865, *Enumeration of fossils collected in the Niagara limestone at Chicago, Illinois; with descriptions of several new species*: Boston Soc. Nat. History, Mem., v. 1, p. 81-114.
- Winckworth, R. A.**
- (995) 1929, *Marine Mollusca from South India and Ceylon. III. Pinna. With an index to the Recent species of Pinna*: Malacol. Soc. London, Proc., v. 18, pt. 6, p. 276-297, text fig. 1-5.
- (996) 1930, *Notes in nomenclature. 6. Lima and allied genera*: Same, Proc., v. 19, pt. 3, p. 115-116.
- (997) 1932, *The British marine Mollusca*: Jour. Conchology, v. 19, p. 211-252.
- Wiontzek, Herbert**
- (998) 1934, *Rudisten aus der oberen Kreide des mittleren Isonzogebietes*: Palaeontographica, v. 80, Abt. A, p. 1-84, text fig. 1-21, pl. 1-4.
- (999) 1935, *Zur Organisation der Radiolitengattungen Bournonia und Medeella*: Zentralbl. Mineralogie, Geologie, Paläontologie, Abt. B, no. 3, p. 90-96, text fig. 1-3.
- (1000) 1937, *Ueber altbekannte Hippuriten aus Südfrankreich und von Gosau*: Paläont. Zeitschr., v. 19, p. 327-341, text fig. 1-4, pl. 24-28.
- Wöhrmann, S. F. von**
- (1001) 1889, *Die Fauna der sogenannten Cardita- und Raibler-Schichten in den nord-tiroler und bayrischen Alpen*: K. K. Geol. Reichsanst. Wien., Jahrb., v. 39, p. 181-260, pl. 5-10.
- (1002) 1893, *Ueber die systematische Stellung der Trigoniden und die Abstammung der Nayaden*: Same, Jahrb., v. 43, p. 1-28.
- Wood, S. V.**
- (1003) 1840, *On the fossil shells of the Crag*: Charlesworth's Mag. Nat. History, v. 4, p. 230 (London).
- (1004) 1850, *A monograph of the Crag Mollusca*: v. 2 (Bivalves), Palaeontograph. Soc. London, 341 p., 31 pl. (London).
- Woodring, W. P.**
- (1005) 1925, *Miocene Mollusca from Bowden Jamaica, pelecypods and scaphopods*: Carnegie Inst. Washington, Publ. 366, p. 1-564, pl. 1-40.

- Woods, Henry**
 (1006) 1908, *A monograph of the Cretaceous Lamellibranchia of England* (v. 2, pt. 5): Palaeontograph. Soc. London, v. 62, p. 194-207, pl. 30-33.
- Woodward, S. P.**
 (1007) 1851-56, *A manual of the Mollusca*: xvi + 486 p., 25 pl., J. Weale (London).
 (1007a) 1855, *On the structure and affinities of the Hippuritidae*: Geol. Soc. London, Quart. Jour., v. 11, p. 40-61, pl. 3-5.
- Worthen, A. H., & Meek, F. B.**
 (1008) 1875, *Palaeontology of Illinois, Section II, Descriptions of invertebrates*: Geol. Survey Illinois, v. 6, p. 481-532, pl. 23-33.
- Wrigley, A. G.**
 (1009) 1936, *An "Aenigma" from the London Clay*: Malacol. Soc. London, Proc., v. 22, no. 1, p. 23-26, pl. 5.
- Yabe, Hisakatsu, & Nagao, Takumi**
 (1009a) 1926, *Praecaprotina, nov. gen., from the Lower Cretaceous of Japan*: Tohoku Imper. Univ., Sci. Repts., ser. 2 (Geol.), v. 9, p. 21-24, pl. 7.
- Yakovlev, N. N.**
 (1010) 1928, *Procrassatella, un genre nouveau du permien*: Soc. Paléont. Russie, Ann., v. 7, p. 121-125.
- Yen, Teng-Chien**
 (1011) 1946, *On Lower Cretaceous fresh-water mollusks of Sage Creek, Wyoming*: Acad. Nat. Sci. Philadelphia, Notula Naturae, no. 166, p. 1-13, 13 text fig.
- Yonge, C. M.**
 (1012) 1926, *The digestive diverticula in the lamellibranchs*: Royal Soc. Edinburgh, Trans., v. 54, p. 703-718.
 (1013) 1928, *Structure and function of the organs of feeding and digestion in the septibranchs, Cuspidaria and Poromya*: Royal Soc. London, Philos. Trans., ser. B, v. 216, p. 221-263.
 (1014) 1936, *The evolution of the swimming habit in the Lamellibranchia*: Musée Histoire Nat. Belgique, Mém., v. 2, no. 3, p. 77-100.
 (1015) 1939, *The protobranchiate Mollusca: a functional interpretation of their structure and evolution*: Royal Soc. London, Philos. Trans., ser. B, Biol. Sci. no. 566, v. 230, p. 79-147, pl. 15.
- (1016) 1946, *On the habits and adaptations of Aloidis (Corbula) gibba*: Marine Biol. Assoc. United Kingdom, Jour., v. 26, p. 358-376.
 (1017) 1947, *The pallial organs in the aspidobranch gastropods and their evolution throughout the Mollusca*: Royal Soc. London, Philos. Trans., ser. B, v. 232, p. 443-518.
 (1018) 1949, *On the structure and adaptations of the Tellinacea, deposit feeding Eulamellibranchia*: Same, ser. B, v. 234, p. 29-76.
 (1019) 1953, *The monomyarian condition in the Lamellibranchia*: Royal Soc. Edinburgh, Trans., v. 62, p. 443-478.
 (1020) 1953, *Form and habit in Pinna carnea*: Royal Soc. London, Philos. Trans., ser. B, v. 237, p. 335-374.
 (1021) 1957, *Mantle fusion in the Lamellibranchia*: Stazione Zool. Napoli, Pubbl., v. 29, p. 151.
 (1022) 1959, *The status of the Protobranchia in the bivalve Mollusca*: Malacol. Soc. London, Proc., v. 33, p. 210-214.
 (1023) 1960, *General characters of mollusca: in Treatise on invertebrate paleontology, Raymond C. Moore (ed.), Part I, Mollusca 1, p. 113-136* (New York).
 (1024) 1962, *On Etheria elliptica Lam. and the course of evolution, including assumption of monomyarianism, in the family Etheriidae (Bivalvia: Unionacea)*: Royal Soc. London, Philos. Trans., ser. B, v. 244, p. 423-458.
 (1024a) 1967, *Form, habit and evolution in the Chamidae (Bivalvia) with reference to conditions in the rudists (Hippuritacea)*: Royal Soc. London, Philos. Trans., ser. B, v. 252, p. 49-105, text fig. 1-31.
- Zittel, K. A. von**
 (1025) 1881-85, *Handbuch der Palaeontologie. I Abt. Palaeozoologie*: v. 2, 893 p., Oldenbourg (München, Leipzig).
 (1026) 1913, *Text-book of Palaeontology*: transl. C. R. Eastman, 2nd edit., v. 1, 839 p., 1594 text fig., Macmillan & Co., Ltd. (New York, London).

ADDENDUM

LIMITATIONS OF BERNARD AND MUNIER-CHALMAS SYSTEM
FOR BIVALVE HINGE NOTATION

DONALD W. BOYD and NORMAN D. NEWELL

[Modified from D. W. Boyd and N. D. Newell, 1968; prepared under NSF Grant GB-6905X]

A method of designating homologous heterodont hinge teeth was devised independently and synthesized late in the 19th century by BERNARD (1895-98) and MUNIER-CHALMAS. It is described by Cox in a previous section (p. N53-55) of this *Treatise* Part. Their notation has been widely recommended, but it has not been accepted universally in practice. In many cases their method of naming teeth is difficult to apply because homologies among dissimilar bivalves commonly are not demonstrable readily; the homologous teeth must be recognized and correlated before the notation can be applied properly.

According to these workers, cardinal tooth 1 is an ontogenetically modified part of anterior lateral *AI*, 2 of *AII*, and so on. Both 2 and 3 commonly divide to form *2a*, *2b*, and *3a*, *3b*, respectively. But, if growth series are lacking, how can these teeth be identified? Furthermore, tooth *4b*, although correlated with *AIV*, is not, in our experience, accompanied by *4a*, and *5b* apparently does not have a counterpart in *5a*. Details of these ontogenetic modifications presumably were determined empirically, but they have never been well documented and need to be confirmed.

The recognition of homologies and the tracing of phylogenetic trends in bivalve teeth are complicated by the fact that non-occluding, or incompletely occluding, ridges and furrows along the hinge plate commonly are excluded from consideration in the dental notation by most investigators. An obsolescent or poorly developed tooth ridge, or the shell margin itself, commonly is not specified in the dental formula. Indeed, many students of heterodont bivalves designate as teeth only those that are completely embraced by a socket, or furrow, of the complementary valve. In this narrow definition, a weakly or incompletely oc-

cluding ridge may not be accepted as a hinge tooth. This is not only illogical but fails to provide for the recognition of newly appearing or obsolescent teeth.

Semantic difficulties aside, it is well established that teeth of the heterodonts are "variable" and that new elements may be added and old ones lost during evolution. It has not been demonstrated, however, that these historical events are recapitulatory in ontogenies of living bivalves, as supposed by BERNARD. In fact, it may be that they are rarely recapitulatory.


We believe the system of BERNARD and MUNIER-CHALMAS, with its stress on homologies, is misapplied commonly when used for bivalve taxa in which ontogeny and phylogeny are known poorly.

OBJECTIVE PLAN FOR HINGE ILLUSTRATION AND NOTATION


We propose a flexible, objective method of recording morphological details of the hinge teeth that can be converted readily to a modified version of the BERNARD system whenever homologies are secure. The new system is equally applicable for both heterodont and nonheterodont hinges.

The conventional orientation in illustrating bivalve hinges and the arrangement of dental formulas are extremely confusing. Furthermore, the idea that dental homologies in related bivalves can be ascertained by matching hinge teeth in numerical sequence certainly is not warranted unless supported by independent evidence such as graded growth series or stratigraphic sequences.


Consequently, we find merit in illustrating matched pairs of valves (Fig. BN1) in a manner that facilitates direct comparison with a simplified and objective nota-




posterior anterior
 RV 1 0 (1)—e r 1 0 1 0 1——1 0 1
 LV 1 0 1——e r 1 0 1 0——(1) 0 1



RV 1 0 (1)—n 1 0 1 0 (1)——1 0 1
 LV 1 0 1——n 0 1 0 1 0——(1) 0 1



RV 1 0 (1)—e s r 1 0 1——1
 LV 1 0 1——e s r (1) 0 1 0——(1) 0 1



RV 1 0 (1)—n (1) 0 1 0——1 0 1
 LV 1 0 1——n 0 1 0 (1)——(1) 0 1

FIG. BNI. Crassatellacean hinges with notation of morphological elements and proposed formulas for them.

1. *Hybolophus speciosus* (ADAMS), Recent crassatellid from Florida; 1a,b, RV and LV int., $\times 2$, with formulas for hinges.

2. *Oriocrassatella elongata* BOYD & NEWELL, Permian crassatellid from Park City Formation, Wyoming; 2a,b, RV and LV int., $\times 2$, with formulas for hinges.

3. *Astarte castanea* SAY, Recent astartid from shallow water off Atlantic Highlands, New Jersey; 3a,b, RV and LV int., $\times 2$, with formulas for hinges.

4. *Astartella aueri* BOYD & NEWELL, Permian astartid from Park City Formation, Wyoming; 4a,b, RV and LV int., $\times 3$, with formulas for hinges.

[EXPLANATION: C, cardinal tooth; e, elastic-ligament area; L, lateral tooth; n, nymph; r, resilifer; s, septum; 0, space between adjacent teeth or ridges; 1, tooth or potential articular ridge; parentheses (), inconspicuous or doubtful tooth; horizontal line, boundary between lateral and cardinal teeth.]

tion. For use with poorly understood groups, our proposed method has the merit of being free from implications of uncertain homologies by avoiding specification of homologues during preliminary studies. The two valves of a shell are illustrated beak to beak, with the right valve (RV) above the left valve (LV) (Fig. BN1). Conventionally, the BERNARD formula places the LV below, and our suggested orientation is a concession to this convention. Thus, the posterior parts of both valves lie on the left. The notation system, a modification of that of STEINMANN as interpreted by DALL (1913), has been devised to be compared directly with such illustrations.

The RV hinge is expressed by the upper of two lines of symbols, and, in both lines, the symbols are arranged from left to right to reflect a traverse along the hinge from the posterior extremity to the anterior extremity. An arabic numeral one "1" represents each projection and ridge of an alternating series along the dorsal part of the shell. These alternate from one valve to the other but they do not necessarily occlude. Inconspicuous or dubious teeth are indicated between parentheses. A zero "0" represents a space between two teeth or ridges. Generally, but not invariably, these spaces function as sockets or grooves. In any case, they indicate the position of an articulating ridge or prominence on the opposite valve. Horizontal lines are used to delimit the cardinal from the lateral series. For the crassatellaceans, the letters "r," "s," "n," and "e" are added to represent the position of the resilium, septum, nymph, and elastic ligament, respectively.

CASE HISTORY: CRASSATELLACEAN HINGE

A review of past and present interpretations of hinge features in the heterodont superfamily Crassatellacea illustrates the limitations of the BERNARD system of hinge notation, as well as the uncertain phyletic significance of grades of hinge complexity.

BERNARD's research on the ontogeny and comparative morphology of the hinges of bivalve mollusks was cut short by his early death, and, to our knowledge, his work has not been subsequently brought under

critical review. Essential documentation of his views about dental homologies among living crassatellaceans is lacking, so we are unable to confirm or deny his conclusion. However, we note that BERNARD and many subsequent investigators have recognized certain peculiarities in the dentition of crassatellaceans. Some authors have placed the superfamily with the lucinoids, others with the cyrenoids. DAVIES (1935), following BERNARD (1895) and LAMY (1917), considered the Crassatellacea FÉRUSSAC, 1822 (=Astartacea D'ORBIGNY, 1844), to be of lucinoid derivation, superficially assuming a cyrenoid aspect as a result of evolutionary acquisition of an additional cardinal tooth in the RV.

DAVIES (op. cit.) segregated the Crassatellacea and Carditacea known to him from other lucinoids on the basis of their hinge characteristics and KOROBKOV (1954) applied the ordinal name "Astartedonta" to these bivalves. Our studies indicate, however, that this segregation undesirably would dismember the lines usually classed as Astartidae and Crassatellidae, and distribute them between two of KOROBKOV's orders, the Lucinodonta and the Astartedonta. Consequently, although we acknowledge the utility of "lucinoids" and "cyrenoids" as grades of hinge complexity, there is little to commend them as bases for separate taxa. It appears to us more probable that they represent iterative rather than phyletic grades of complexity. As is shown below, the characteristics of ancestral crassatellaceans do ally them with primitive lucinoids. But post-Paleozoic forms achieve a more advanced evolutionary status superficially similar to, but not quite like, the cyrenoid grade. Furthermore, the hinges of later members of the Crassatellidae and the Astartidae resemble one another more than they do those of their Paleozoic ancestors—an example, apparently, of evolutionary convergence. Thus, some doubt is cast on the integrity of the superfamily Crassatellacea as usually constituted.

CRASSATELLID DENTAL FORMULAS

In the notation suggested above, the hinge of *Hybolophus speciosus*, a living

crassatellid, can be characterized as given in the two-line formulas accompanying Figure BN1,1a,b.

BERNARD and later students have supposed that the RV cardinals correspond to 5b, 3b, and 3a, and the two LV cardinals, to 4b and 2. We cannot find convincing documentation in the literature that this notation is firmly based on studies of ontogenetic development, or that these teeth are homologous with those of other heterodonts conventionally given the same numerals. Similarly, little agreement is found with respect to the lateral teeth. For example, DARRAGH (1965, p. 98) cited only two laterals, AII and PI, in *Eucrassatella*, although he mentioned AI and PII (*ibid.*, p. 100, 102), in comments on certain species of that genus. Authors using the BERNARD system have not specified as lateral teeth all the eight ridges indicated as such in our interpretation of *Hybolophus* (Fig. BN1,1). LAMY (1917, p. 200) utilized AI, AII, AIII, PI, PII, and PIV. In several diagrams he designated a strong posterior lateral between PII and PIV as PI. It is not clear whether he so intended, because, in the BERNARD system, the position between PII and PIV should be occupied by PIII. There is precedent, however, in one of BERNARD's diagrams (1895, p. 121) for the misplacement of a lateral tooth where he places AII above AI. We assume that this position was unintentional.

Recent crassatellids (indeed, all post-Paleozoic ones known to us), possess more cardinals and laterals than does the Paleozoic *Oriocrassatella* (Fig. BN1,2). Furthermore, the strong septum that divides the resilium from the ligament in this genus is obscure or lacking in post-Paleozoic crassatellids in which the ligament and the resilium are in contact.

In living crassatellids, as in all the astartids known to us, the anterodorsal margin of the RV overlaps that of the LV. The reverse situation exists in *Oriocrassatella*. The homologies of *Hybolophus* and *Oriocrassatella* are not at all evident and will remain uncertain until intermediate forms, if they exist, are discovered. We favor the hinge analysis of the Permian genus given with Figure BN1,2a,b.

ASTARTID DENTAL FORMULAS

Although astartids lack the internal ligament of crassatellids, the dentition of modern representatives of the two families is similar. We interpret the hinge of a Recent shell (*Astarte castanea*) as shown with Figure BN1,3a,b.

DAVIES (1935, p. 156), following BERNARD's interpretation of the hinge in the Astartidae, designated the cardinal teeth of the RV as 5b, 3b, and 3a, and those of the LV as 4b and 2.

By contrast, NICOL (1955, p. 157) interpreted the hinge in *Astarte* as cyrenoid. He designated the large central tooth on the RV as the pivotal tooth 1, and termed the bracketing teeth of the LV 2b and 2a. He showed (*ibid.*, fig. 4, p. 157) on the RV a posterior cardinal which he designated 3b. The lack of an anterior cardinal 3a in his diagram is perplexing, because the LV (his fig. 1) shows an appropriate anterior socket.

In the present study, a survey of Cenozoic astartids in the collections of the American Museum of Natural History indicates that the shell shown in Figure BN1,3, is characteristic in possessing three cardinal teeth in the RV. The anterior cardinal, omitted from NICOL's diagram, is recognizable in nearly all RVs examined. The posterior cardinal, although commonly well developed, is virtually absent from three Eocene species investigated. Both of these teeth are generally small. In cases in which one is especially poorly developed, the corresponding socket is more conspicuous than the tooth.

The hinge of the Paleozoic genus *Astartella* is represented by the formulas given with Figure BN1,4a,b.

It appears that *Astarte* characteristically has the anterior cardinal in the RV, whereas *Astartella* has the anterior cardinal in the LV, as do members of the Myophoriidae and of the Scaphellinidae. The outer anterior lateral of the RV extends to the beak, an unusual and primitive feature in heterodonts.

INTRODUCTION OF NEW TEETH IN CRASSATELLACEAN HINGE

The lucinoid hinge grade appears at least as early as the Middle Ordovician (McALESTER, 1965), whereas the cyrenoid hinge was rare or lacking before the Mesozoic. These differences in stratigraphic distribution lend support to the view of BERNARD, DAVIES, and others that the cyrenoid hinge was derived by elaboration of the lucinoid hinge. BERNARD concluded that the splitting of the pivotal (central and principal) cardinal tooth in LVs of lucinoids gave rise to two teeth separated by a new socket. The latter receives a new pivotal cardinal tooth of the RV. If this change has been a phyletic as well as an ontogenetic innovation, it may have been heralded by a broadening and bilobation of tooth 2 before actual bifurcation. Bilobation of the pivotal tooth of the LV is a characteristic feature of many late Paleozoic pelecypods (e.g., *Schizodus*) and suggests a possible pattern in a trend toward multiplication of hinge teeth.

BERNARD, DAVIES, and others have supposed that the three cardinal teeth of living crassatellacean RVs correspond to *5b*, *3b*, and *3a*, whereas the two cardinal teeth of LVs were designated as *4b* and 2. We have given thought to NICOL's (1955) view that the central cardinal on the RV may be cyrenoid 1, rather than the lucinoid *3b*, and we are unable at present to resolve this problem. The Permian crassatellaceans clearly are similar to lucinoids in possessing two cardinal teeth in each valve, whereas the post-Paleozoic forms generally have a third tooth, commonly identified as *5b*, at the posterior end of the RV series. This tooth is represented in the Paleozoic in *Astartella* but not in *Oriocrassatella*. Although *Astartella* has the cardinal just cited, it has only two, not three, cardinals in the RV. The pivotal tooth of the Paleozoic forms is perceptibly bifid in *Oriocrassatella*. The evolutionary bifurcation of this tooth and the appearance of a new pivotal tooth, 1, in the opposite valve conceivably could result in hinge development very similar to that which characterizes living crassatellaceans.

According to this interpretation, the crassatellaceans would consist of Paleozoic lucinoids, and post-Paleozoic forms of cyrenoid aspect lacking one tooth typical of the cyrenoid dentition. This suggestion is perhaps more attractive for the crassatellids than for the astartids, because the pivotal tooth in *Astartella* is not bifid.

In any case, the two families require a manipulation of different teeth, in order to derive for both a cyrenoid post-Paleozoic hinge from a Paleozoic lucinoid ancestor. The cyrenoid pivotal tooth would originate within the posterior cardinal socket of the RV in astartids but in the anterior cardinal socket in crassatellids. The two cardinal teeth of the astartid RV would then change numbers in transition from Paleozoic to post-Paleozoic (*5b* to *3b*, and *3b* to *3a*), whereas the enumeration of crassatellid cardinal teeth of the RV (*3a* and *3b*) would remain constant. Finally, the two teeth of the LV would originate from a splitting of the posterior cardinal (*4b*) of *Astartella*, but from the anterior cardinal (2) in *Oriocrassatella*.

Even for crassatellids, the preceding idea is suspect, because both Paleozoic and post-Paleozoic LVs have only two cardinals. The hypothesis recognizes an evolutionary trend toward an increase in the number of cardinal teeth and suggests that a shift in hinge symmetry is accomplished by a splitting of the Paleozoic pivotal tooth of the LV to accommodate a new pivotal tooth in the RV. However, this interpretation requires the disappearance of a pre-existing tooth, the LV posterior cardinal. In the interest of simplicity, it appears more likely that steps in the development of the cyrenoid hinge involved only the appearance of new teeth, rather than contemporaneous appearance and disappearance. In our present state of knowledge, it is more reasonable to suppose that the two cardinals of the LV in *Oriocrassatella* are homologous with those in modern crassatellids, than that the posterior one has disappeared while the anterior one evolved into two. Likewise, although it is possible that the two teeth of Paleozoic RVs are homologous with the anterior and posterior elements of the three-toothed post-Paleozoic RVs, it seems

more likely that they are homologous with the anterior and central ones, leaving the posterior rather than the middle one as a post-Paleozoic innovation.

This alternative invites speculation that the posterior tooth in question, commonly designated as *5b*, developed as a wall related to the posterior margin of the resilifer. The border of the resilium in *Oriocrassatella* is reflected in a few valves by an obscure shelf along the anterior margin of the resilifer. This shelf occupies a position analogous to that of the posterior cardinal in post-Paleozoic specimens. Cardinal *5b* in living species is more variable in size and orientation than the other cardinals, and the LV lacks a corresponding socket in some species.

REFERENCES

Bernard, Félix

1895, *Première note sur le développement et la morphologie de la coquille chez les lamellibranches*: Soc. Géol. France, Bull., v. 23, pt. 3, p. 104-154.

1896, *Deuxième note sur le développement et la morphologie de la coquille chez les lamellibranches (taxodontes)*: Same, v. 24, pt. 3, p. 54-82.

1896, *Troisième note sur le développement et la morphologie de la coquille chez les lamellibranches (anisomyaires)*: Same, v. 24, pt. 3, p. 412-449.

1897, *Quatrième et dernière note sur le développement et la morphologie de la coquille*

chez les lamellibranches: Same, v. 24, pt. 3, p. 559-566.

1898, *Recherches ontogéniques et morphologiques sur la coquille des lamellibranches*: Ann. Sci. Nat. (Zool.), Paris, v. 8, p. 1-208, pl. 1-12.

Dall, W. H.

1913, *Pelecypoda*: in Zittel, K. A. von, Textbook of paleontology, translated and edited by C. R. Eastman (revised edit.). London and New York, Macmillan Co., v. 1, p. 422-507.

Darragh, T. A.

1965, *Revision of the species of Eucrassatella and Spissatella in the Tertiary of Victoria and Tasmania*: Roy. Soc. Victoria, Proc., v. 78, pt. 1, p. 95-114, pl. 12-15.

Davies, A. M.

1935, *Tertiary faunas*: London, Thomas Murby and Co., v. 1, x + 406 p., 565 fig.

Korobkov, I. A.

1954, *Spravochnik i metodicheskoe Rukovodstvo po tretichnym mollyuskam Platinchatozhabernye*: Gosud. Nauchno-tech. Issledov. Nefti Gornotoplivnoi litry, Leningradskoi Otdelenie, 444 p., 96 pl. (in Russian) (*Handbook on and systematic guide to Tertiary Mollusca. Lamellibranchia.*)

Lamy, Édouard

1917, *Révision des Crassatellidae vivants du Muséum d'Histoire Naturelle de Paris*: Jour. Conchyliologie, v. 62, no. 4, p. 197-270, pl. 6.

McAlester, A. L.

1965, *Systematics, affinities, and life habits of Babinka, a transitional Ordovician lucinoid bivalve*: Palaeontology, v. 8, p. 231-246, 3 pl.

Nicol, David

1955, *Morphology of Astartella, a primitive heterodont pelecypod*: Jour. Paleontology, v. 29, no. 1, p. 155-158.

INDEX

Italicized names in the following index are considered to be invalid; those printed in roman type, including morphological terms, are accepted as valid. Suprafamilial names are distinguished by the use of full capitals and author's names are set in small capitals with an initial large capital. Page references having chief importance are in boldface type (as N327). Some divergences in classification reflect differences of authors concerning validity of nomenclature.

- Abakaniella, N861
Abarbatia, N252
Abathoceramus, N315
 Abbreviatiana, N862
Abichia, N593
 Abiella, N408, N409
Abisa, N332
 Abra, N45, N63, N199, N636
 Abranda, N613
Abrina, N636
 Acalliana, N862
 Acanthocardia, N51, N585
 Acanthopecten, N336
Acanthotrigonia, N487
Acapis, N633
 Acar, N252
Acardo, N815
 accessory cavities (rudists), N761
 accessory muscle, N102
 accessory muscle scars, N38
 accessory plates, N64
 accessory ventral adductor, N36
 ACEPHALA, N3, N227
 ACÉPHALÉS DIMYAIRES, N119
 ACÉPHALÉS MONOMYAIRES,
 N120
 ACEPHALOPHORA, N121
 Acesta, N386
 Acharax, N243
 Achasma, N536
 Acila, N45, N69, N231
 acline, N102
 Aclistothyra, N533
Acolus, N683
 Acorylus, N613
 Acostaea, N466
Acostea, N466
 Acreuciroa, N857
Acritopaphia, N685
Acroceramus, N317
 Acrostrigma, N586
 Acteocinidae, N195
Actinobolus, N556
Actinoceramus, N315
Actinochlamys, N355
 Actinodesma, N299
 actinodont, N51, N102
 ACTINODONTA, N118, N393
 Actinodonta, N111-N112, N399
Actinodontacea, N210
 ACTINODONTOIDA, N393
 Actinodontophora, N470
 Actinodontophoridae, N219, N468
Actinomya, N399, N847
 Actinonaias, N451
Actinopecten, N350
Actinophorus, N311
Actinoptera, N302
Actinopterella, N302
 Actinopteria, N302
 Actinopterina, N299
 Actinotrigonia, N478
Actionaias, N455
 Acuticosta, N416
 Acyrena, N666
 Adacna, N593
 Adacnarca, N270
Adacnidae, N590, N593
 Adacninae, N220, N593
 Adamiana, N862
 ADAMS & ADAMS, N208
 Adamussium, N349
 ADAPEDONTA, N118, N214,
 N690
 adaptation, N138
Adasius, N638
 adductor muscle, N31, N102,
 N177
 adductor scar, N102
Adesmacea, N118, N210, N702
 Adipicola, N278
 ADKINS, N210
 Adontorhina, N509
 Adrana, N237
Adranaria, N234
 Adranella, N239
Adspergillum, N859
 Adula, N7, N278
 Adulomya, N243
Aediculus, N818
 Aegericiana, N862
 Aegilops, N868
Aeglia, N450
 Aelga, N849
Aene, N621
Aenigma, N383
 Aenigmoconcha, N561
 Aenona, N621
 Aeora, N673
 Aequipecten, N142, N161-N162,
 N355
Aequipectus, N355
Aequivalvis, N355
 Aeqvioldia, N240
 Aeretica, N622
Aetheria, N466
Africofragum, N585
 Afrocardium, N588
 Afroparresia, N416
Afropisidium, N669
 Agapella, N646
Agaria, N550
 Agelasina, N660
Agina, N700
 AGLOSSA, N227
 Agnocardia, N585
 Agnomyax, N621
 Agonisca, N489
Agria, N806
Agriodesma, N678, N847
 Agriopleura, N764, N769-N771,
 N774, N806, N815
 Agriopoma, N678
 Aguilerella, N306
 Aguileria, N308
 Ahtioconcha, N299
Aigyromya, N831
 Akebiconcha, N664
Alasmedonta, N447
Alasmesodonta, N447
 Alasmidonta, N447
 Alasmidontinae, N219, N447
Alasminota, N448
Alasmisodonta, N447
Alasmodon, N447
Alasmodonta, N447
 alate, N102
Alathyria, N461
 Albimanus, N849
 Alectryonia, N380
Alicia, N852
 Aligena, N523
 Aliomactra, N598
 alivincular, N60, N102
 ALLEN, N188
 Allocardium, N863
Alloceramus, N315, N317
 Alloidesma, N400
 Alloidesmatidae, N113, N120,
 N219, N400
 Allogramma, N847
 Allomactra, N595
 allomorphism, N102
 Allonychia, N285
 Allopagus, N653
Allorisma, N115, N818, N838
Allorismacea, N119
Allorismiella, N818
 Alnifia, N299
Aloidis, N692
 Alpecanusiana, N862
 alternative sexuality, N29
Alucinoma, N500
 Alula, N819
 Alveinus, N653

- AMALITSKY, N404, N409
Amathusia, N590
Amblasmodon, N447
 Amblema, N437
 Ambocardia, N660
 Ambonicardia, N860
 AMBONODONTA, N118-N119
 Ambonychia, N147, N211, N219, N285
 Ambonychiacea, N63, N285
 Ambonychiidae, N112-N113, N147-N148, N177, N219, N285
Ambonychinia, N287
 Ambonychiopsis, N285
 Ambuscintilla, N533
 Ameghinomya, N672
Ameigilans, N551
 AMEODONTIDA, N118
Americana, N668
 Americardia, N588
 Americuna, N558
 Amerycina, N520
Amesoda, N670
 Amesodesma, N608
 Amiantis, N39, N675
Amita, N297
Amnicana, N669
 Amnicusiana, N862
Amnigenia, N402, N404
 Amnigeniella, N860
Amoebea, N355
 Amonotis, N343
Amotapidae, N698
 Amotapus, N699
 Amphiarus, N569
Amphicardium, N589
 Amphichaena, N631
 Amphicoelia, N285
Amphidesma, N608, N636
 amphidetic, N60, N102
Amphidona, N631
 Amphijanira, N355
 Amphilepida, N533
 Amphinaias, N437
 Amphipsammus, N631
 Amphitriscoelus, N761-N762, N764, N772, N789-N790
Amphithaea, N680
Amusiidae, N348
Amusiopecten, N367
 Amusium, N142, N160-N162, N199, N348-N349
 Amusium group, N349
 Amusiopecten, N117, N348, N367
Amusium, N350
Amusyopecten, N367
Amygdala, N682
Amygdalonia, N457
 Amygdalum, N132, N278
 anachomata, N103
 Anadara, N85, N150, N254
 Anadarinae, N218, N254
Anadontina, N447
Anaitis, N687
Anapa, N520
 Anapachydiscus, N197
 Anapella, N610
Anaplomya, N842
Anaplophora, N468
 Anapteris, N696
 Anatifopsis, N867
 Anatimya, N845
 Anatina, N606
Anatina, N845
Anatinacea, N118-N119, N210
 Anatinella, N607
Anatinelladae, N607
 Anatinellidae, N220, N607
 Anatiniana, N862
Anatinidae, N119, N844
 Anatipecten, N355
 anatomy, N134
Anaucella, N374
Ancella, N374
 Anchomasa, N708
Anchomosa, N708
 Ancliffia, N566
 Anderkenia, N299
 Andrussella, N595
Anelasmodon, N447
Anelasmodontia, N447
 Angarodon, N410
Angellum, N248
 angle of incidence, N85
 Anguipecten, N365
 Angulus, N613
Angustella, N308
 Angusticardo, N540
 Anisocardia, N117, N646
 Anisocorbula, N692
 Anisodonta, N540
 ANISOMYARIA, N34, N118-N119, N121, N214
 anisomyarian, N33, N103
 ANISOMYARIER, N118
Anisoperna, N322
Anisorhynchus, N696
Anisothyris, N697
Ankistrocardia, N661
 Annachlamys, N355
 Anningella, N376
Anningia, N376
 Annuliconcha, N337
Anodon, N442
 anodont, N103
 Anodonta, N22, N30, N51, N63-N64, N95, N173-N174, N181-N182, N402-N403, N442
Anodonte, N442
Anodontes, N442
Anodontophora, N468
 Anodontia, N51, N164, N167, N502
Anodontigenus, N442
Anodontina, N442
 Anodontinae, N219, N442
 Anodontites, N463
Anodontites, N463
 ANODONTODA, N227
 Anodontoides, N444
Anodontophora, N468
 Anodontopleura, N817
 Anodontopsis, N400
Anodota, N442
 Anofia, N675
Anomala, N668, N681
 Anomalocardia, N686
Anomalocardia, N254, N686
 Anomalocoelia, N289
 ANOMALODESMACEA, N2, N118, N210, N214, N818
 ANOMALODESMATA, N63, N113-N114, N118, N211, N213-N215, N218, N221, N881
 Anomalodiscus, N686
 Anomalodonta, N287
 Anomalokellia, N527
Anomalolina, N391
Anomalopleura, N830
 Anomalopleuroides, N830
 Anomia, N4, N8, N21, N29, N33, N37-N38, N47, N132, N147-N148, N155, N383
 Anomiacea, N15, N18, N35, N98, N119-N120, N211-N212, N219, N383
 Anomianella, N863
 Anomiidae, N12, N58, N76, N119-N120, N148, N212, N219, N383
Anominia, N383
Anomolima, N39
Anomoptychidae, N787
Anomoptychus, N795
Anonica, N302
 Anopaea, N317
 Anopisthodon, N489
Anoplomya, N842
Anoplophora, N468
Anoplophoria, N468
Anoptera, N287
 Antediplodon, N458
 anterior, N42, N103
 anterior adductors, N31
 anterior laterals, N49
 anterior lateral tooth, N103
 anterior pallial arteries, N24
 anterior pedal retractor, N37
 anterior protractors, N38
 anterior retractor muscle of the visceral mass, N38
 anterodorsal margin, N42, N103
 anteroposterior axis, N79
 Antetrichoma, N271
 ANTHONY, N79
 Anthonya, N574
Anthracomya, N293, N407
 Anthraconaia, N114, N293, N406
 Anthraconauta, N293
Anthraconeilopsis, N233
Anthraconneilo, N233
Anthracoptera, N295
 Anthracosia, N63, N114, N404, N406-N407
 Anthracosiacea, N63, N219,

- N401-N402, N404
Anthracosidae, N409
 Anthracosiidae, N219, N404,
 N406, N860
 Anthracosphaerium, N406-N407
Anticorbula, N847
 Antigonina, N168, N672
 Antiguamya, N692
 Antijanira, N355
 Antijanira group, N355
Antijanirula, N355
Antilla, N494
 Antilocaprina, N789-N790
 Antinioche, N688
 Antipecten, N356
 Antipleura, N247
 Antipleuridae, N218, N247
Antipleuriden, N247
 Antipleurinae, N218, N247
 Antiquicyprina, N646
 Antiquilima, N386
 Anulipecten, N333
 anus, N5, N22
Anusium, N349
Aparcthyria, N420
Aperiploma, N850
 Aphanais, N317
 Aphrodina, N676
Aphrodite, N590
 Aphrodora, N678
 Apocardia, N588
 Apiotrigonia, N478
Aplodon, N465
Apocardia, N646
 Apolymetis, N625
Apolymetis, N626
 apophyses, N705
 apophysis, N37, N103
Aporema, N834
 Aporrhaidae, N195
 Aporrhais, N195, N199-N200
Appius, N445
 Apricardia, N763, N780
 Apscheronia, N593
Aptolinter, N258
 Aptychopsis, N867
Aptyxites, N806
 Apulites, N811
Aquaria, N859
 Araeopleura, N757, N783
 Aralocardia, N657
 Arca, N8, N16, N30, N38, N42,
 N47, N61-N63, N77, N79,
 N86, N90, N116, N132, N137,
 N141, N144, N150-N151,
 N195, N199, N211, N250
 ARCACEA, N248
 Arcacea, N16, N18, N63, N73,
 N81, N120, N210-N211, N218,
 N250
Arcaceae, N250
 Arcavicula, N304
Archaeodon, N260
 ARCHAEOGASTROPODA, N17
 Archanodon, N114, N402, N404
 Archanodontacea, N211, N219,
 N401-N402, N404
 Archanodontidae, N219, N404
Archicardium, N585
 Archivesica, N664
 Arcicardium, N593
 Arcidae, N12, N16, N37, N70,
 N98, N118-N120, N150, N173,
 N195, N212, N214, N218
 N250, N860
 Arcidens, N448
 Arcidopsis, N412, N416
 Arcinae, N218, N250
Arcinella, N518
Arcinella, N548, N701
Arcinellina, N701
 Arclites, N860
 ARCOIDA, N63, N99, N112,
 N212, N214, N217-N218,
 N248
 Arcomya, N116, N834
Arcomyidae, N827
 Arcomytilus, N116, N271
 Arconaia, N412, N418
 Arcopagella, N621
 Arcopagia, N23, N613
 Arcopaginula, N613
 Arcopagiopsis, N621
 Arcopella, N614
 Arcoperna, N275
Arcoptera, N252, N265
 Arcopsis, N263
 Arctica, N62, N195, N645
 Arcticacea, N16, N211, N221,
 N644
Arcticacea, N657
 Arcticeramus, N317
 Arctiidae, N116-N117, N195,
 N221, N644-N645, N860
 arcticoid ("cyprinoid") type, N56
 arcticoid type, N103
Arctinula, N354
Arctoë, N679
 Arctopratalum, N589
 Arctostrea, N154, N171
 Arctotis, N345
Arcturella, N549
 Arcturellina, N549
Arcturus, N551
 Arcuatula, N132, N141, N163,
 N275
Arcuatula, N304
 Arcullaea, N268
 Arcullaeinae, N218, N268
 Arculus, N542
 Ardeamya, N626
 area, N46
 Arealiana, N862
Arenaria, N635
Arene, N515
 Arenomya, N691
 Areocuculla, N861
Aretinula, N354
Argina, N254
Arginarca, N254
Arginella, N261
 Arginopsis, N261
Argoderma, N366
Argodesma, N366
 Argopecten, N356
Argus, N366
 Argyrodonax, N610
 Argyromya, N831
 Arietina, N861
 Arisaigia, N233
 Aristerella, N394
Aristoceramus, N317
 Arkansia, N448
Armida, N646
Armimiltha, N505
 Arnaudia, N815
 Arnoldia, N863
 Arnoldina, N444
 Arnouldiana, N862
Arotonaias, N455
 Arpadicardium, N591
 Arroiana, N862
Artemis, N680
Artena, N672
Arthemiderma, N680
Arthemis, N680
 Arthritica, N520
 Arthropteron, N463
Artusius, N639
 Arvella, N275
Arytaena, N859
Arytene, N859
Arythaena, N859
 Asa, N679
Asaphidae, N629
 Asaphinella, N631
 Asaphinoides, N633
 Asaphis, N633
 Ascitellina, N633
Ashcroftia, N260
 Ashtarotha, N562
 Asiatotrigonia, N480
 ASIPHONIDA, N118-N119
Asmidia, N646
 Asoella, N337
Aspalima, N265
 Aspatharia, N463
Asperarca, N252
Aspergillidae, N858
Aspergillum, N859
Aspergillum, N859
Aspergillus, N859
 Aspidopholas, N712
Assellbergia, N299
Astartacea, N118-N119
 Astarte, N44, N55, N57, N63,
 N73, N116, N166, N171,
 N195, N199, N562, N909
 ASTARTEDONTA, N118, N910
 Astartella, N114, N563, N909,
 N911-N912
 Astartellopsis, N564
 Astartemya, N569
 Astartidae, N12, N57, N70, N76,
 N114, N116, N173, N195,
 N199, N220, N562, N910-
 N911
 Astartila, N825

- Astartinac, N220, N562
 Astartopsis, N565
 Astatoceramus, N315
ASTHENODONTA, N118
Asthenodonta, N402-N404
ASTHENODONTIDA, N214
 Asthenothaerus, N850
 Asticusiana, N862
 Atactodea, N608
Atalanta, N646
 Atamarcia, N683
 Ateriana, N862
Athletoceramus, N315
 Athlopecten, N357
 Atomodesma, N115, N289
 Atopodonta, N678
 Atreta, N378
 atria, N24
 Atrina, N132, N162-N163, N283
 attachment scars (rudist), N764
Aucella, N374
Aucellidae, N374
 Aucellina, N376
Aucellinae, N374
 Augea, N293
Augustaia, N302
Aulacoceramus, N315
Aulacomya, N320
Aulacomya, N271
Aulacomyella, N343
Aulacophora, N854
Aulacopleurum, N489
Aulus, N612
Aupouria, N270
Aureitapes, N685
 auricle, N24, N46, N103
 auricular crura, N53, N103
 auricular sulcus, N103
 auriculate, N103
Auriscalpium, N845
Aurora, N420
Australima, N386
Australpera, N669
 Austriella, N508
 Austrodozinia, N679
Austrolima, N386
 Austromacoma, N623
 Austromactra, N595
 Austromytilus, N273
 Austroneaera, N854
 Austronucula, N231
 Austroplatyodon, N692
Austropteria, N303
Austrosarepta, N266
 Austrosintilla, N522
Austroteredo, N726
Austrotindaria, N235
 Austrotrigonia, N480
Austroturquetia, N541
 Austrovenus, N686
AUTOLAMELLIBRANCHIATA,
 N118
 Autonoe, N537
 Avardaria, N860
 Avicardium, N593
Avicula, N302
Aviculae, N302
 Avicularca, N346
 Avicularium, N594
Aviculidae, N119, N212, N297
Aviculipinna, N283
 Aviculolima, N386
Aviculomonotis, N342
 Aviculomya, N863
 Aviculomyalina, N290
 Aviculopecten, N113, N115,
 N137, N143-N144, N336
 Aviculopectinidae, N113-N115,
 N143-N144, N219, N335,
 N860
 Aviculopectininae, N219, N336
 Aviculoperna, N308
Aviculoperna, N322
Aviculopinna, N283
 Aviculovulsa, N329
 Avimactra, N595
 Avoniana, N862
Awadia, N589
 axes, N78
 Aximedia, N862
 Axinactis, N267
Axinaea, N267
Axinaeinae, N267
Axinaeoderma, N267
 Axinodon, N529
 Axinopsida, N195, N199, N510
Axinopsis, N475, N510
 Axinulus, N9, N510
Axinus, N508
 axis, N16, N78
Azara, N698
Azarella, N548
Azaria, N550
 Azor, N631
Azor, N639
 Azorinus, N639
 Babinka, N38, N109-N113, N182,
 N491
 Babinkacea, N206, N211, N220,
 N491
 Babinkidae, N220, N491
Bacilia, N858
 Bactronophorus, N726
 Baculites, N196, N197, N200
 Badiotella, N386
 Bakevellia, N115, N306
 Bakevelliidae, N114-N116, N219,
 N306
 Bakevelloides, N306
Bakewellia, N306
 Balantioselena, N579
 Baluchicardia, N557
 Balwantia, N418
 Bankia, N8, N65, N143, N723,
 N732
 Bankiella, N733
 Bankiinae, N221, N730
Bankiopsis, N734
Bankiura, N733
Baphia, N332
Barbala, N445
Barbarca, N252
Barbata, N445
 Barbatia, N116, N132, N141,
 N150-N151, N252
Barbatiella, N262
Barbatirus, N252
 Barbierella, N492
 Barbosaia, N299
Barcinia, N244
 Barclaya, N533
Barclaya, N533
 Barcoona, N863
Bariosta, N437
 Barnea, N65, N708
Barnia, N708
Baroda, N683
 BARRANDE, N109, N113, N118
Barretia, N801
 Barrettia, N157, N758-N759,
 N769-N771, N776, N801
 Barrimysia, N529
 BARROIS, N112
 bars, N69
Bartelletia, N467
 Bartlettia, N467
 Bartrumia, N625
 Baryana, N862
 Baryconites, N817
 Barymactra, N595
Barynaias, N426
Baryosta, N437
 Barytellina, N621
 basal margin, N103
 Bassina, N686
 Bassinaria, N673
 Basterotella, N540
 Basterotia, N540
Basterotidae, N539
 Batavusiana, N862
Bathmoceramus, N317
Batholites, N801
Bathyamussium, N351
 Bathyarcia, N254
 Bathycardita, N550
 Bathycorbis, N511
Bathyspinula, N235
Bathytellina, N618
 Bathytormus, N574
 Batissa, N666
 Batolites, N758, N770-N771,
 N776, N801
 BAYLE, N485, N749, N799,
 N811, N815
 Bayleia, N761, N763, N780
Bayleidae, N779
 Bayleioidea, N761, N781
 bead, N69, N103
 beak, N42, N103
 beak cavities, N412
Beaumontia, N329
 Becoariana, N862
 Beguina, N548
Belchlamys, N355
 Bellucina, N496
Beloceramus, N220
Bendeglans, N551

- Bendemacoma, N623
Bendoneacera, N854
 BENETT, N21
Bentharca, N254
Benthocardiella, N558
Bentholyonsia, N847
Benthoquetia, N541
Bequania, N508
Berenguieriana, N862
 BERNARD, N53, N55, N57, N94,
 N98-N100, N118, N120, N209,
 N322, N471, N755, N908,
 N910-N912
 BERNARD & MUNIER-CHALMAS
 system, N908
Bernardina, N650
Bernardinidae, N221, N650
Bernayia, N513
Berpolis, N861
Besairieia, N317
Beushausenella, N302
Beushausenia, N256, N258, N299
bialate, N103
Biapholus, N700
Bicorbula, N692
Bicornia, N741
Bicornuacprina, N817
Bicornucopina, N790, N817
Bicrenula, N233
Bidentina, N653
Bihippurites, N801
Bilobaria, N537
Bineurus, N431
Binghami, N863
 biogeography (Thyasira), N195
Biopholeus, N700
Biradiolites, N158, N769-N770,
 N810
Biradiolitinae, N221, N766, N810
birostres, N762
Birostrina, N315
Birostrites, N816
Bisphaerulites, N817
Bissomya, N700
Bisulcus, N470
Bittnerella, N265, N338
Bittneria, N376
Bitubulites, N863
 BIVALVIA, N3, N4, N109, N130,
 N195, N197, N205, N213,
 N217-N218, N227, N766
 bladder, N25
Blagroveia, N658
Blainvilla, N863
 BLAINVILLE, DE, N118, N749
Blainvillia, N681
Blairella, N561
Blanckenhornia, N370
Blanfordina, N346
Bleta, N863
 blood sinuses, N24
Bodmania, N249
 body, N11, N46
 body of shell, N103
Boehmiceramus, N315
Boeuvia, N506
 BøGGILD, N74, N76, N209
Boiomytilus, N290
Bonartemis, N679
Bontaea, N850
Boriesia, N854
 boring bivalves, N168
Borissia, N235
Bornia, N523
Borniola, N523
Bosempra, N615
Bositra, N9, N343
Bosphoricardium, N591
Bothrocorbula, N692
Botula, N34, N169, N278
Botulina, N275
Botulopa, N279
Botulopsis, N275
Bouleigmus, N329
Bourdotia, N506
Bournonia, N751, N804, N810
bourrelet, N103
Boussacia, N266
Bowdenia, N854
 BOWDICH, N326
Brachidontes, N132, N147-N148,
 N170, N273
Brachiodacna, N592
 BRACHIOPODA, N4
Brachtechlamys, N366
Brachyanodon, N444
Brachymeris, N515
Brachyodontes, N273
Bradicardia, N860
branchitellum, N103
Brazzaea, N418
Brebissoniana, N862
Brechites, N858
Breviarca, N263
Brevicardium, N588
Brevimalleus, N329
Brevinucula, N231
Briandiana, N862
 BRILL, N71
Broeckia, N818
 BRONN, N3, N118
 brood chamber, N29
 brood protection, N91
Brotiana, N862
Bruetia, N515
 BRUGUIÈRE, N385
Bruniastarte, N569
Bruntrutia, N860
Brussiella, N861
Bruzzaea, N418
Bryopa, N858
Bryophila, N269
Bucardia, N657, N828
Bucardiomya, N828
Bucardites, N584
Bucardium, N585
Buccardium, N657
Buchia, N116-N117, N374
Buchiidae, N76, N219, N374
Buchiola, N244
Buchotrigonia, N480
Budmania, N592
Bullata, N437
Bullella, N447
Bulloideus, N460
Bunodus, N859
 BUONANNI, N4
Burckhardtia, N312
Bureiomya, N828
Burmesia, N838
Burmesiidae, N221, N838
Burricera, N868
Burtonia, N466
Bushia, N851
Busycon, N173
Butor, N845
Butorella, N845
Butovicella, N248
Butovicellidae, N218, N248
buttress, N103, N183
byssal foramen, N103
byssal gape, N47, N103
byssal gland, N15
byssal groove, N15
byssal notch, N47, N103
byssal pit, N15
byssal sinus, N103
Byssanodonta, N670
byssate closely attached bivalves,
 N146
byssate epifaunal nestlers, N150
byssate fissure-dwellers, N151
byssate free-swinging bivalves,
 N144
byssiferous, N103
Byssosarca, N250
Byssobornia, N523
Byssocardium, N595
Byssodesma, N394
Byssomia, N700
Byssomya, N700
Byssonia, N700
Byssonychia, N285
Byssopteria, N287
byssus, N8, N14, N103
Bythiamena, N562
Cabralia, N651
Cabralista, N651
Cacophonia, N604
Cadella, N615
Cadmusia, N716
Cadomia, N233
Caecella, N610
Caclatura, N418
Caenonaias, N425
Caestocorbula, N65, N696
Caestocorbulinae, N221, N696
Cafferria, N420
Caillaudiana, N861
Caladacna, N592
Calcaraca, N850
Calceola, N448
Calcinella, N635
Callanaitis, N686
Calliarca, N860
Callicardia, N860
Callicistronia, N681

- Calliscapha*, N463
 Callista, N86, N168, **N677**
Callistina, N676
Callistoceramus, N315
Callistoderma, N677
 Callistotapes, N685
 Callithaca, N688
 Callitriche, N861
 Callitricoderma, N861
 Callitrigonia, N485
Callizona, N675
Callizonata, N675
 Calloarca, N252
Callocallista, N677
 Callocardia, N101, **N677**
 Callodonta, N394
 Callogonia, N664
Callolima, N386
 Callomysia, N529
 Callonaia, N458
 Callucina, N494
 Callucinella, N498
 Callucinopsis, N494
 callum, N65, N102, N705
Calobates, N726
Caloosarca, N256
Calopodium, N843
 Calorhadia, N240
 Calpitaria, N675
 Calva, N678
 Calvitium, N542
Calyculina, N670
 Calyptogena, N664
Calyptopholas, N712
Camacea, N518
Cameola, N518
Cameronia, N464
 Camostraea, N861
 Campiloceratites, N861
 Campitonectes, N352
Camponectes, N352
Campstonectes, N352
Camptoceramus, N315
 Camptochlamys, N352
 Camptonectes, N351
 Camptonectes group, N351
 Camuriana, N862
 cancellate, N103
 Caneyella, N299
 Canthyria, N420
 Capillimya, N845
 Capistrocardia, N700
 Caprina, N756, N758, N761,
 N768, N772-N773, N789-
 N790
 Caprinella, N790, N795
Caprinellidae, N787
 Caprinidae, N78, N221, N754,
 N758, N760-N761, N763-
 N764, N766, N769, **N787**,
 N789-N790, N860
 Caprinula, N758, N761, N769-
 N770, N773, N789-N790
 Caprinuloidea, N158, N751-
 N752, N755, N758, N761-
 N762, N769-N770, N772-
 N773, N789-N790
 Caprinus, N860
 Caprotina, N761, N784
Caprotina, N781
Caprotinadae, N784
 Caprotinidae, N221, N754, N759-
 N761, N763-N764, N766, **N784**
Capsa, N621, N633
 Capsaeformis, N452
 Capsella, N628, N631
Capsula, N633
 Cara, N254
 Carbonarca, N257
 Carbonicola, N114, N404, N406-
 N407
Carbonicola, N407
Carbonicolidae, N406, N409
Cardea, N585
 Cardiacea, N16, N63, N117-
 N118, N121, N211, N220,
 N583
 Cardiarius, N860
 Cardiidae, N13, N16, N34, N38,
 N70, N113, N116-N118, N159,
 N166, N168, N195, N220,
 N583, N860
 Cardiinae, N220, N583
 Cardilia, N608, N745
 Cardiliidae, N220, N607
Cardilla, N745
 Cardilona, N608
 cardinal area, N46, N103
 cardinal axis, N40, N103
 cardinal costa, N103
 cardinal crura, N53, N103
 cardinal platform, N103
 cardinal teeth, N49, N103
Cardinea, N578
 Cardinia, N116, N468, N578
 Cardiniidae, N116, N220, N468,
 N562, N578
 Cardinioides, N468
 Cardiniopsis, N570
Cardinopecten, N364
 Cardiocardita, N550
Cardiocardites, N550
Cardiodonta, N646
 Cardiola, N245
Cardiolaria, N229
Cardiolidae, N245
 Cardiolinae, N218, N245
 Cardiolopsis, N245
Cardiolucina, N496
 Cardiomorpha, N112, N115,
 N818
 Cardiomya, N854
Cardiopsis, N245
Cardissa, N588
 Cardita, N44, N57, N67, N548
Cardita, N556
 Carditacea, N16, N119, N121,
 N211, N220, N543, N910
Carditadade, N548
 Carditamera, N549
 Carditamerinae, N220, N548
 Carditella, N549
 Carditellona, N550
 Carditellopsis, N550
 Cardites, N556
 Carditesinae, N220, N556
 Carditidae, N12, N70, N73,
 N116-N117, N220, N543,
 N548
 Carditinae, N220, N548
 Carditomantea, N554
 Carditopsis, N558
 Cardium, N4, N6, N9, N29, N30,
 N38, N60-N63, N77, N86, N91,
 N93, N166, N181-N182, N195,
 N584
 carina, N103
 Carinastarte, N562
 carinate, N103
Carinella, N635
Caritodens, N299
Carneola, N670
 Carnidia, N862
 Carolia, N15, N383
 CARPENTER, N720
Carpenteria, N380
 CARTER, N136, N138, N168
cartilage, N103
 Cartissa, N860
 Carunculina, N451
 Carvalhoiana, N862
Caryatis, N675
 Carydiidae, N219, N400
 Carydium, N114, N400
 Caryocorbula, N693
Casertiana, N669
 CASEY, N55, N209
 Caspicardium, N593
 Cassianella, N311
 Cassianellidae, N115, N219, N310
 Castalia, N458
 Castaliella, N458
 Castalina, N458
 Castalioides, N459
 Casterella, N826
Casteria, N322
 Cataceramus, N315
 catachomata, N103
 Catella, N258
 Catella group, N258
Catelysia, N683
 Catilloides, N593
Catillopecten, N354
Catillum, N315
Catillus, N315
 caudate, N103
 Caudiculatus, N420
 Cavatidens, N502
 Cavilinga, N499
 Cavilucina, N494
 Celtoides, N545
 cementation, N8
 cemented epifaunal bivalves, N153
Centralhyria, N461
 Centrocordita, N551
 cephalic hood, N705
Cephaloceramus, N317
 CEPHALOPODA, N197

- Cerana*, N680
Cerastes, N590
 Cerastoderma, N11, N590
Ceratisolen, N639
Ceratobornia, N523
Ceratoceramus, N315
Ceratoconcha, N862
Ceratomya, N69, N90, N662, N839
 Ceratomyidae, N221, N838
 Ceratomyopsidae, N221, N662
 Ceratomyopsis, N662
Cerceis, N595
Cercomyacidae, N844
Cercomya, N45, N845
Cercomyopsis, N820, N823
 cerebropleurial ganglia, N26
Ceromia, N839
Ceromya, N839
Ceromyella, N839-N840
Ceromyidae, N838
Ceromyopsis, N662
Ceronia, N608
Ceroniola, N608
Ceropsis, N557
Cerullia, N541
Cervilleia, N308
Cetoconcha, N852
Cetomya, N852
Cetothrax, N850
Chacea, N711-N712
Chaena, N699
Chaenocardia, N340
Chaenocardiinae, N219, N339
Chaenocardiola, N295
Chaenodomus, N394
Chaenomya, N114, N831
Chaenopaea, N700
Chalmasia, N329
Chalmys, N355
Chama, N57, N88, N101, N141, N154-N155, N157, N518, N751, N768, N799, N808
Chama, N494
Chamacea, N16, N119, N121, N211, N220, N518
Chambardia, N463
Chambardiana, N861
Chamberlainia, N420
Chamelea, N686
Chametrachea, N594
Chamidae, N8, N13, N36-N37, N57, N72-N73, N101, N116, N154, N220, N518, N751, N756, N768, N861
Chaminae, N518
Chamostraea, N844
Chamostrea, N13
Chamostrea, N844
Chamostreidae, N844
Chaperia, N758, N761, N785
Chartoconcha, N593
Chattonia, N575
Chavanella, N553
Chelidoneura, N463
Chelidonopsis, N463
Chelidonura, N463
 CHEMNITZ, N389
Chevronaias, N460
 chevron groove, N103
Chiapasella, N769, N771-N772, N811
Chiapasia, N803
Chimaera, N283
Chimacoderma, N283
Chimela, N678
 chimney, N705
Chion, N628
Chione, N77, N86, N168, N686
Chione, N677
Chionella, N677
Chioneryx, N688
Chioninae, N221, N686
Chionista, N686
Chionopsis, N686
Chironia, N523
Chlamydella, N353
Chlamydina, N355
Chlamydoconcha, N14, N537
 Chlamydoconchacea, N211, N220, N537
 Chlamydoconchidae, N220, N537
Chlamys, N23, N33, N38, N47-N48, N69, N101, N115-N116, N132, N141-N142, N150-N153, N160-N162, N170, N175, N355
 Chlamys group, N355
Chloromya, N273
 CHOFFAT, N811
 chomata, N103
 Chondrodontidae, N73, N219
 chondrophore, N49, N103, N705
Chondrostea, N415
Choniocardia, N550
Choristodon, N689
Choromytilus, N34, N37, N274
Christadens, N420
 CHUBB, N158, N801
Cibota, N250
Ciboticola, N861
Cicatella, N523
Cicatrea, N650
 cicatrix, N103
 Ciclopekten, N352
 ciliated discs, N17
Cimitaria, N820
Cinclidoceramus, N320
Cinetodonia, N850
Cipliacella, N518
Ciplyella, N518
 Circe, N672
 Circenita, N672
 Circinae, N221, N672
Circlimopa, N265
Circumphalus, N40, N672
 circulatory system, N24
Cista, N443
Citharoceramus, N317
Cladoceramus, N320
Cladopoda, N702
 Claibornicardia, N557
 Claibornites, N502
Clamys, N355
Claraia, N115, N337
 CLARKE, N9
Clarkia, N827
 classification, N205, N218; rudist, N766; Unionacea, N414
Clathroconcha, N520
Clathrolima, N346
Clathrotellina, N615
Claudiconcha, N689
Clausaria, N739
Clausina, N510, N671
Clausinella, N687
 claustra, N412
Clavagella, N67, N858
 Clavagellacea, N16, N76, N118-N119, N211, N222, N857
 Clavagellidae, N13, N67, N222, N858
Clavellatae, N477
 clavicle, N103
 Clavicosta, N337
Clavigella, N858
 Clavipholas, N708
Clavitrigonia, N485
Clavotrigonia, N485
Cleidophorus, N233
 Cleidothaeridae, N222, N844
Cleidothaerus, N844
Cleionychia, N287
 Clementia, N681
Clementiinae, N221, N681
Clencharia, N235
 Cleobis, N115
Cleobis, N824
Cleone, N445
Clepsydra, N859
Clepydra, N859
Clessinella, N670
Clessinia, N669
Cletella, N669
 Clidiophora, N844
Climacoceramus, N315
Climacotrigonia, N480
Clinocardium, N590
Clinopistha, N228
Clione, N446
 Cliopteria, N299
 Clisocolus, N511
Clistoconcha, N845
Cloissonaria, N739
Cloissonaria, N739
 closed, N103
Clossonaria, N739
Clossonaria, N739
Clotho, N515, N700
 Clunaculum, N639
Clupibankia, N733
Cnesterium, N240
Cnidiella, N686
Cnisma, N265
 Coalcomana, N761-N762, N772, N789-N790
Cobracephalus, N299
 Cochentopyx, N529

- Cochlodesma, N63, N850
 Codakia, N86, N164, N167, N494
 Codalucina, N502
Codokia, N494
 Coe, N29
 Coelastarte, N567
 Coelatura, N862
 Coelocyprina, N646
Coelodon, N844
Coelogonia, N643
 Coelomactra, N596
 Coelopsis, N571
 Coeloteredo, N726
 coiled umbones, N45
 Cokeria, N437
 Coleoconcha, N533
 collar, N65
Colleopterum, N442
 Collignonicardia, N646
 Collobiana, N862
 COLLOCONCHIDA, N118, N214, N217
Collopterum, N442
Colombia, N601
 Colonche, N688
 coloration, N70
Colorimactra, N595
Colpoceramus, N321
 Colpomya, N394
Columba, N465
 Colveraia, N771, N774, N815
 COMFORT, N11, N72
Comitileda, N235
 commarginal, N103
 commissure, N42, N103, N170
Complanaria, N448
Complanatiana, N862
 complex prismatic structure, N74
 complex structure, N76
Complicachlamys, N366
 composite prismatic structure, N74
 compressed, N103
 Compsomyax, N681
Comptovallium, N366
 Comus, N674
 concentric, N69, N103
Conchacea, N119
 Conchentopyx, N529
 CONCHIFERA, N3, N227
 conchiolin, N58, N74, N103
 Conchites, N863
 Conchocele, N508
 Conchodon, N743
 Conchodromus, N452
 Conchodus, N116
Conchodus, N743
 CONCHOPHORA, N227
 Condylocardia, N558
 Condylocardiidae, N220, N222, N558
 Condylocardiinae, N220, N558
Condyloceramus, N321
 Condylucuna, N559
 Congeria, N72, N117, N644
 Congeriomorpha, N287
Congeriomya, N644
 Conocardicea, N121, N211, N222, N859
 Conocardiidae, N113, N222, N859
 CONOCARDIIDEN, N859
 CONOCARDIOIDA, N218, N222, N859
Conocardiopsis, N297
 Conocardium, N112-N113, N115, N168, N859
 CONRAD, N553
 Conradilla, N452
 consecutive sexuality, N29
 Contradens, N420
Conucardia, N660
 convexity, N103
Convexopecten, N369
 COOKE, N119
 Cooperella, N690
 Cooperellidae, N221, N689
 Cooperellopsis, N690
 Coquandia, N698
Coralichlamys, N355
 Corallidomus, N113, N168
Corallidomus, N397
Coralliochama, N752, N758, N762, N789, N793
Coralliolithes, N801
Coralliophaga, N655
Corallospondylus, N380
Coramya, N700
Corbicella, N635, N642
 Corbicellopsis, N642
 Corbicula, N21, N666
 Corbiculacea, N16, N211, N221, N664
Corbiculadae, N665
 Corbiculella, N666
 Corbiculidae, N13, N56, N96, N221, N664-N665, N861
 Corbiculina, N666
 corbiculoid, N56
 corbiculoid type, N104
 Corbiculopsis, N655
Corbidae, N513
 Corbis, N86
Corbis, N513
 Corbula, N55, N72-N73, N100, N116, N170, N195, N633, N692
Corbula, N695
 Corbulamella, N696
 Corbulamellinae, N221, N696
Corbularia, N692
 Corbulidae, N65, N73, N117-N118, N195, N221, N692
 Corbulinae, N221, N692
 Corbulomima, N696
Corbulomya, N696
 Corburella, N640
 Corculum, N90, N588
 cordate, N104
Cordiceramus, N317
 Cordiopsis, N678
 Cordiula, N511
Cordium, N584
Cordula, N669
Coriarius, N529
Corimya, N850
 Coripia, N554
 CORMOPODA, N227
 Cornea, N670
 Cornellites, N138, N144
Cornellites, N302
Corneocyclas, N670
Corneola, N670
Cornucaprina, N790
 Cornucardia, N660
Cornucopina, N801
Cornuteredo, N727
Coromya, N850
 corselet, N46, N104
 Cortinia, N831
Corymbichlamys, N357
Corymya, N850
 Cosa, N270
Cosatova, N270
 Cosmetodon, N258
Cosmetopsis, N265
Cosmioceramus, N315
 Cosmogoniophora, N396
 Cosmogoniophorina, N111, N397
 Cosmomya, N831
 Cosmopseudodon, N431
Cossmannella, N551
 costa, N69, N104
 Costacallista, N677
 Costanaia, N438
 Costanuculana, N235
Costatae, N477
 Costatoleda, N237
 Costatoria, N473
 costella, N69, N104
 Costellacea, N386
 Costellipitar, N675
 Costelloleda, N235
 Costigervillia, N308
 Costocyrena, N666
 Costokidderia, N644
 costule, N104
 Costulopteria, N863
 Courquiniana, N862
 Cowperesia, N115, N642
 Cox, N1, N146, N174-N175, N208, N212-N215, N273, N404, N409, N488
 Coxiconcha, N861
Cranatula, N325
Craspedodon, N660
Craspedodonta, N445
 Crassadacna, N592
 Crassatella, N12, N46, N48-N49, N166, N171, N173, N573
 Crassatellacea, N16, N63, N118, N211, N220, N562, N910
 crassatellacean hinge, N910
 Crassatellidae, N12, N57, N70, N114, N173, N220, N573, N910
 Crassatellina, N570
 Crassatellinae, N220, N573

- Crassatellites*, N573
Crassatellitidae, N573
Crassatellopsis, N114, N570
Crassatina, N574
Crassialaria, N302
Crassiana, N862
Crassiconcha, N826
Crassina, N562
Crassinella, N577
Crassinella, N567
Crassinidae, N562
Crassitesta, N445
Crassithracia, N850
Crassivenus, N688
Crassostrea, N5, N77, N131, N154-N157, N174
Crassula, N601
Cratis, N270
Cratomya, N834
Cremnoceramus, N315
crenate, N104
Crenatula, N325
Crenatuligenus, N325
Crenella, N274
Crenellinae, N218, N274
Crenellodon, N274
Crenimargo, N535, N568
Crenipecten, N341
Crenocirce, N673
Crenocrassatella, N574
Crenodonta, N437
Crenomytilus, N271
Crenotrapezium, N653
crenulations, N171
Crepispisula, N601
Cretocardia, N551
 CRICKMAY, N483
Cricoceramus, N315
Criocardium, N585
Criocarpium, N585
Crista, N672
Cristadens, N420
Cristaria, N444
crossed-lamellar, N104
crossed-lamellar structure, N74
Cruciella, N863
cruciform muscle, N36, N104
crus, N53, N103
 CRUSTACEA, N4, N109
Crustuloides, N571
Crypogenia, N452
Cryptaulia, N790, N817
Cryptina, N472
Cryptochasma, N258
Cryptocoelopsis, N572
Cryptodon, N508, N604
cryptodont, N104
 CRYPTODONTA, N62, N63, N112-N113, N118, N211, N213-N214, N217-N218, N241
Cryptodonta, N692
 CRYPTODONTEN, N118, N241
 CRYPTODONTIDA, N214
Cryptodontidae, N508
Cryptogramma, N686
Cryptomactra, N597
Cryptomya, N691
Cryptonema, N686
Cryptonemella, N686
Cryptopecten, N357
Cryopseudodon, N432
crystalline style, N21
Ctena, N494
Ctenamusium, N350
ctenidia, N16
Cteniopleurium, N347
Ctenocardia, N588
Ctenoconcha, N233
Ctenodesma, N420
ctenodont, N51, N104
 CTENODONTA, N118
Ctenodonta, N110, N111-N112, N118, N165, N228
Ctenodontacea, N211, N218, N227
Ctenodontella, N233
Ctenodontidae, N218, N228, N409
Ctenoides, N386
ctenolium, N104
Ctenostreon, N116, N385-N386
Cuculla, N657
Cucullaea, N32, N50, N69, N73, N116, N183, N260
Cucullaearca, N252
Cucullaeidae, N218, N259
Cucullaeinae, N259
Cucullana, N260
Cucullaria, N258
Cucullaria group, N258
Cucullastis, N260
Cucullea, N860
Cucullella, N233
Cucullifera, N315
Cucullona, N260
Cucullopsis, N258
Cucumaria, N420
Cucumeria, N420
Cucumerunio, N420
Cucurbitula, N699
Cultellidae, N220, N611
Cultellus, N92, N611
Cultellus, N612, N639
Cultrensis, N611
Cultrioopsis, N308
Culunana, N238
Cumberlandia, N414
Cumingia, N637
Cumularia, N743
Cuna, N560
Cunanax, N559
Cuneamya, N113, N818, N820
Cunearca, N256
cuneiform, N104
Cuneigervillia, N308
Cuneocorbula, N694
Cuneolus, N273
Cuneopsida, N441
Cuneopsis, N420
Cunerus, N628
Cuneus, N628
Cuneus, N673
Cunicula, N420
Cuninae, N220, N560
Cuphus, N739
Curionia, N545
Curricula, N420
Curtonotus, N473
Curvemysella, N529
Curvirimula, N295
Curvirostrum, N257
Curvula, N283
Curvulites, N283
Cuspicorbula, N694
Cuspidaria, N9, N195, N199, N854
Cuspidariidae, N13, N73, N195, N222, N854
 CUVIER, N208, N383, N749
 CVANGARA, N724
Cyamiacea, N11, N16, N211, N220, N537
Cyamiidae, N220, N537
Cyamiocardium, N538
Cyamiomactra, N538
Cyamionema, N535
Cyamium, N538
Cyaneitapes, N685
Cyanocyclus, N666
Cyathodonta, N852
Cycladella, N670
Cycladicama, N515
Cycladidae, N669
Cycladina, N861
Cycladina, N520, N669
Cycladites, N670
Cycladoconcha, N530
Cyclas, N670
Cyclina, N681
Cyclinella, N681
Cyclininae, N221, N681
Cyclinorbis, N681
Cyclocalyx, N669
Cyclocardia, N551
Cycloceramus, N315
Cycloclamys, N353
Cycloconcha, N133, N138, N165, N399
Cycloconchacea, N211, N219, N399
Cycloconchidae, N112, N216, N219, N399
cyclodont, N104
 CYCLODONTA, N118
Cycloglans, N552
Cyclomactra, N597
Cyclomya, N460
Cyclonaias, N439
Cyclopecten, N353
Cyclopellatia, N513
Cyclorisma, N683
Cyclorismina, N683
Cyclostrea, N863
Cyclostreon, N378
Cyclosunetta, N673
Cyclotellina, N615
Cyclothyris, N683
Cyclopecten, N353

- Cydippe*, N613
Cydrinina, N626
Cygnacana, N862
Cylichna, N195
Cylindrica, N424
Cymatioa, N535
Cymatoceramus, N321
Cymatocyclas, N669
Cymatoica, N623
Cymatonota, N133, N137-N138, N165, N394
Cymbophora, N197, N200, N598
Cymella, N852
Cyphoxis, N860
Cyphus, N739
Cypricardella, N114, N575
Cypricardia, N655
Cypricardiacea, N119
Cypricardinia, N111, N113-N114, N579
Cypricardites, N112, N248
Cypricia, N606
Cyprimeria, N681
Cyprina, N646
Cyprinella, N668
Cypriniadea, N646
Cyprinidae, N645
Cyprinopsis, N679
Cyprogenia, N452
Cyrachaea, N499
Cyrena, N666
Cyrenacea, N119, N664
Cyrenastrum, N670
Cyrenella, N518
Cyrenellidae, N517
Cyrenidae, N665
CYRENINA, N118
Cyrenobatissa, N666
Cyrenocapsa, N668
Cyrenodonax, N666
CYRENODONTA, N118
Cyrenodonta, N518
cyrenoid type, N56, N104, N118
Cyrenoida, N518
Cyrenoides, N518
Cyrenoididae, N220, N517
Cyrenolimopsis, N863
Cyrenopsis, N653
Cyrilla, N269
Cyrollista, N269
Cyrollona, N269
Cyrtoceramus, N317
Cyrtodaria, N700
Cyrtodera, N700
Cyrtodonta, N111-N112, N133, N138, N248
Cyrtodontacea, N120, N211, N218, N248
Cyrtodontarca, N299
Cyrtodontidae, N112-N113, N216, N218, N248, N868
Cyrtodontopsis, N287
Cyrtodontula, N249
Cyrtonaias, N450
Cyrtonais, N450
Cyrtopinna, N283
Cyrtopleura, N708, N709-N710
Cyrtorostra, N346
Cyrtosolen, N638
Cyrtusiana, N862
Cystia, N389
Cystigenus, N389
Cytherea, N672-N673
Cytheriopsis, N673
Cytherocardia, N657
Cytherodon, N473

Dacosta, N858
Dacrydium, N195, N279
Dacryomya, N237
Dactylina, N707
Dactyloceramus, N320
Dactyloteredo, N728
Dactylus, N861
Daharina, N547
Dalacia, N325
Dalila, N245
DALL, N49, N53, N57-N58, N112, N209, N212-N214, N910
Dalliconcha, N308
Dalliella, N446
Dallocardia, N586
Dallucina, N496
Damalis, N414
Damaris, N861
Daonella, N115, N344
Daphne, N250
Daphnoderma, N250
Darcinia, N607
Darina, N607
DARRAGH, N911
Darteplicatula, N378
Datta, N314
Dattidae, N219, N314
Davidaschvilia, N653
Davidia, N112, N820
DAVIES, N120, N209, N348, N910-N912
Davila, N610
Davilinae, N220, N610
Davisia, N542
Dceruska, N394
Decadopecten, N365
Decarambis, N448
Decatopecten, N365
Decatopecten group, N365
Deceptrix, N229
DECHASEAUX, N213, N754-N756, N790
Dechaseauxia, N771, N773, N811
Dechenia, N393
Decipula, N530
Decurambis, N448
Degrangia, N700
Deikeya, N595
Deletopecten, N354
DELGADO, N110
Delia, N863
Delphinonaias, N450
Deltachion, N628
Deltamya, N832

Deltopecten, N341, N354
Deltopectinidae, N219, N341
Demanetia, N289
demarcation line, N87, N104
demibranch, N16
Deminucula, N235
Dendopecten, N366
dental formulas, astartid, N911; *crassatellid*, N910
Dentaliidae, N195
Dentalium, N195, N197, N200
denticle, N70, N104, N171
Denticosa, N270
denticulation, N70, N104
Dentilucina, N492
Dentimodiolus, N274
Dentina, N861
Dentipecten, N365
dentition, N104, N172; *rudist*, N769
Dentonia, N666
Dentoperna, N306
Deperetia, N367
Depressiana, N862
Dermatomya, N852
Desertella, N120, N467
Desertellidae, N219, N467
DESHAYES, N749, N799
DESMODONTA, N119
DESMODONTEN, N119
Despontaniana, N862
Destacar, N252
Deuteromya, N382
Deviobankia, N734
Devonia, N530
Dexiobia, N246
Dexiobiinae, N218, N246
Diabolica, N269
Diachora, N378
diagenodont, N104
diagonal ridge, N46, N104
Dianchora, N378
Dianisotis, N445
Diaphoromactra, N599
Diaphragmella, N302
Diatretus, N799
Diaurora, N420
Diberus, N276
Diceras, N46, N89, N90, N116, N120, N749, N754, N756, N765, N768-N769, N771, N776, N778
Dicerata, N778
Diceratia, N778
Diceratidae, N52, N73, N221, N755-N756, N760, N762-N763, N766, N776
Diceratinae, N221, N766, N776
Diceratocardium, N660
Dicerocardiidae, N221, N660
Dicerocardium, N660
Dicranodesma, N538
Dicranodonta, N260
Dictyoceramus, N320
Dictyoptychus, N758, N789, N795
Dictys, N295

- Dicyathifer, **N726**
 Didacna, **N592**
 Didacninae, **N220, N592**
 Didacnoides, **N593**
 Didacnomya, **N592**
Didimacar, **N262**
Didonta, **N700**
 Didymoceras, **N197, N200**
 Didymotis, **N344**
 Dietrichia, **N646**
 digestive gland, **N21**
 digestive system, **N21**
 Digitaria, **N565**
 Digitariopsis, **N563**
Digoniomya, **N394**
Dihora, **N578**
Dilecta, **N296**
Dilora, **N498**
Diluvarca, **N254**
Dimeroceramus, **N315**
Dimya, **N21, N34, N382**
Dimyaciadae, **N382**
 DIMYARIA, **N34, N119**
 dimyarian, **N32, N104**
Dimyarina, **N382**
 Dimyidae, **N12, N16, N119, N219, N382**
 Dimyodon, **N382**
Dimyodus, **N382**
Dimyopsis, **N378**
 Dinocardium, **N166, N590**
Dinodacna, **N594**
 Diodonta, **N626**
Diodontopteria, **N300**
Diodus, **N668**
 DIOGENODONTA, **N119**
Dione, **N675**
Dionisotis, **N445**
 Diotis, **N344**
Dipilidia, **N781**
 Diplasma, **N862**
 Dipleurites, **N344**
 Dipleurodonta, **N394**
Diplidia, **N817**
 Diplodon, **N450**
 Diplodonta, **N63, N515**
 Diplodontidae, **N206**
Diplodontidae, **N514**
 Diplodontina, **N523**
 Diplodontites, **N464**
Diploplax, **N711**
 Diplopseudodon, **N434**
Diploschiza, **N378**
 Diplothyra, **N174, N712**
Dipsada, **N445**
Dipsas, **N444**
Dipsax, **N445**
Dipsus, **N445**
Dipterophora, **N302**
 Diptyxis, **N863**
 direct development, **N91**
 directive plane, **N85**
 directive rib, **N85, N88, N104**
 directive spiral, **N104**
 disc, **N46, N104**
 Discinocaris, **N867**
Discoceramus, **N317**
 Discomiltha, **N499**
 Discomya, **N439**
 Disconaias, **N450**
 discordant margins, **N104**
 Discors, **N589**
Disnomia, **N452**
 Disparilia, **N570**
 dissoconch, **N99, N104**
 Distefanella, **N755, N765, N769-N770, N810**
 Distugonia, **N692**
Dithalmia, **N643**
 DITHYRA, **N227**
 Ditichia, **N228**
 Ditypodon, **N666**
 Divalinga, **N506**
 Divalucina, **N506**
 Divaricardium, **N589**
 divaricate, **N104**
 Divaricella, **N69, N506**
 Divaricellinae, **N220, N506**
Divaricolima, **N386**
 Divarikellia, **N527**
 Divarilima, **N389**
 Divariscintilla, **N537**
Divergidens, **N550**
 Diversicostata, **N593**
 DODD, **N77**
 DODGE, **N46, N378**
 Dolabra, **N868**
 Dolichopteron, **N299**
Doliolabis, **N277**
 Dollfusia, **N678**
Donacarius, **N628**
Donacocardium, **N629**
 Donacidae, **N13, N76, N119, N138, N220, N628**
 Donacilla, **N609**
Donacilla, **N618**
Donacina, **N628**
 Donacites, **N863**
 Donacopsis, **N666**
 Donax, **N21, N32, N38, N42, N63, N67, N166, N168, N628**
Dorbignya, **N801**
 Dorisca, **N673**
 dorsal, **N42, N104**
Dorsina, **N672**
 dorsomedian scars, **N38**
 Dorsomya, **N863**
Dorvillea, **N636**
 Dosina, **N672**
 Dosinella, **N679**
 Dosinia, **N42, N44, N86, N167, N679**
Dosinia, **N679**
Dosiniana, **N679**
Dosinidia, **N679**
 Dosiniinae, **N221, N679**
 Dosiniopsella, **N679**
 Dosiniopsis, **N678**
 Dosinisca, **N679**
 Dosinobia, **N680**
 Dosinorbis, **N679**
Dosinula, **N672**
 DOUVILLÉ, **N55, N112, N209, N212-N214, N471, N745-N746, N749, N751, N758, N761, N799, N808, N811, N815**
Dozierella, **N300**
 Dozyia, **N568**
Dreissena, **N643**
Dreisseina, **N643**
Dreissena, **N9, N13, N32, N63, N96, N117, N147-N148, N643**
 Dreissenacea, **N16, N63, N211, N221, N643**
Dreissenadae, **N643**
Dreissencia, **N643**
Dreissenia, **N643**
 Dreissenidae, **N95, N117, N147, N221, N643**
Dreissenomya, **N644**
Dreissensa, **N643**
Dreissensia, **N643**
Dreissensiomya, **N644**
Dreissina, **N643**
Dreistena, **N643**
Drepanoceramus, **N315**
Dreysseina, **N643**
Dreysseina, **N643**
Dreysseomya, **N644**
Dreysseusia, **N643**
Dreysseusia, **N643**
Driessena, **N643**
Driessensia, **N643**
Dromus, **N452**
 Dualina, **N247**
 Dubertretia, **N769-N771, N813**
Dufoichaena, **N699**
 DUMÉRIL, **N208**
 Dunbarella, **N334**
 Duplipecten, **N357**
 duplivincular, **N104**
 duplivincular ligament, **N60**
 Durania, **N154, N157-N158, N751, N769-N770, N812**
Duranius, **N812**
 Durania, **N158**
 Durga, **N745**
 DURHAM & ZULLO, **N723**
Dux, **N247**
Dvorecia, **N297**
Dymia, **N382**
Dymya, **N382**
Dypsas, **N445**
 Dysmea, **N631**
 Dysnomia, **N452**
 dysodont, **N53, N104**
 DYSODONTA, **N119-N120**
 Dysodonta, **N233**
 DYSODONTEN, **N119**
Dysonomia, **N452**
Dystactella, **N228**
 Eamesiella, **N502**
 ear, **N104**
 Eastonia, **N604**
 EBERSIN, **N213**
 Eburneopecten, **N352**

- Eburneopecten group, N352
Eburnopecten, N352
Ecericardium, N591
Echinochama, N101, N154
Echinochama, N518
Echinopecten, N372
Echinotis, N346
Echinotus, N346
Echion, N383
Echionoderma, N383
 ecology (Thyasira), N190
Ectenagena, N664
Ectenocardiomorpha, N394
Ectenodesma, N299
Ectenoptera, N863
Ectorisma, N852
Ecuadorea, N460
Edentula, N326, N442
 edentulous, N104
Edmondia, N112, N115, N408, N818
 Edmondiacea, N211, N221, N818
Edmondiella, N818
 Edmondiidae, N113, N221, N818
Edulicardium, N590
Egerella, N628
Egeria, N628-N629
Egesta, N681
Egeta, N668
Egetaria, N668
 EGGLETON & DAVIS, N57
Egraca, N492
Egracina, N506
Ekstadia, N233
Elasmatium, N863
Elasmodophora, N287
Elasmogena, N448
Elasmogona, N448
Elathia, N508
Elathia, N502
 ELATIOBRANCHIATA, N227
 ELATIOCEPHALA, N227
Electomactra, N598
Electrina, N304
Electroma, N304
Electromactra, N598
Elegantarca, N265
Elegantinia, N473
Elegantula, N636
 ELEUTHERORHABDA, N119
Eleutherospondylos, N378
Eligma, N329
Eligmidae, N326
Eligmus, N116, N329
Elimata, N389
Ellipsaria, N862
Ellipsopsiana, N862
Elliptio, N420
Elliptioidea, N863
Elliptioideus, N425
Elliptiotellina, N615
Elongaria, N422
Elongatiana, N862
Elpidollina, N615
Eltopera, N380
Elymella, N820
 emarginate, N104
Embiana, N862
Embla, N852
Empleconia, N265
Enanticeramus, N317
Enantiostreon, N115, N380
Enargetos, N861
Endocostea, N183, N317
Endodesma, N168
Endodesma, N394
Endomargarus, N847
Endopleura, N637
Enigmonia, N383
Enkelbergia, N287
Ennucula, N231
Enocephalus, N644
Ensatella, N611
Ensia, N863
Ensiculus, N612
Ensidents, N423
 ensiform, N104
Ensigerilleia, N308
Ensio, N566
Ensis, N30, N32, N37, N63, N90, N164, N166-N167, N170, N173, N611
Ensitellops, N541
Enteropleura, N344
 entire, N104
Entodesma, N28, N58, N847
Entoliidae, N219, N346
Entoliopsis, N349
Entolium, N71, N346
Entomonotis, N374
Entovalva, N11, N14, N530
entré, N759
Eoastarte, N565
Eocallista, N666
Eocamptonectes, N339
Eocrassina, N653
Eocyclina, N681
Eodicerus, N779
Eodiplodon, N460
Eodivaricella, N508
Eodon, N565
Eodonax, N642
Eoinoceramus, N320
Eoischyrina, N867
Eolimea, N389
Eolymnium, N416
Eomactra, N598
Eomegalodon, N114
Eomegalodon, N745
Eomegalodus, N745
Eomeretrix, N673
Eomiltha, N502
Eomiodon, N569, N653
Eomiodon, N553
Eonaias, N439
Eonavicula, N116, N252
Eontia, N261
Eopecten, N72, N716, N373
Eophysema, N504
Eopinctada, N304
Eopleurophorus, N823
Eoprosodacna, N592
Eopseuma, N518
Eopteria, N244
Eoradiolites, N758, N760-N761, N769-N771, N774, N806
Eoschizodus, N473
Eosolen, N611
Eoteredo, N740
Eotrapezium, N116, N648
Eotrigonia, N480
Eoursivivas, N694
Ephigenia, N629
Ephippium, N383
Ephippodonta, N14, N535
Ephippodontina, N535
Ephippodontoana, N535
 epibranchial chamber, N20
Epiceramus, N315
Epicodakia, N496
Epicyprina, N646
 epidermis, N104
Epidiceras, N776, N779
Epidiceratidae, N779
Epidiceratinae, N221, N754, N766, N779
 epifaunal Bivalvia, N141
Epilhippopodium, N583
Epilepton, N542
Epilobasma, N452
Epilucina, N494
Epioblasma, N452
Epirisis, N254
 EPSTEIN & LOWENSTAM, N134
Eptolea, N235
Equichlamys, N357
 equilateral, N45, N104
Equipecten, N355
 equivalve, N104
 erect, N69
Ereila, N508
Eridonychia, N285
Erigidocardium, N589
Eriphyla, N568
Eriphyliinae, N220, N568
Eriphylopsis, N570
Erodona, N698
Erodonidae, N221, N698
Erratisolen, N639
Ervilia, N610
Erviliinae, N220, N610
Erycina, N519
Erycinacea, N11, N84, N98
Erycinacea, N518
Erycinella, N559
Erycinidae, N13, N220, N519
Erycinopsis, N522
Eryx, N519
Escalima, N390
 escutcheon, N46, N104
 escutcheon ridge, N46, N104
Eselaevitrigonia, N481
 N414, N466, N861
Esmerarca, N254
 esophagus, N21
Estellacar, N263
Etea, N646
Ethaeria, N466

- Etherea*, N466
 Etheria, N60, N62-N63, N72, N466
 Etheriidae, E9, E73, E118, E219, N466, N861
Etheriigenus, N466
 Etheripecten, N337
 Ethmocardium, N585
Ethra, N787
Etolium, N346
Euanodonta, N442
Eubiradiolites, N810
Euburneopecten, N352
 Eucallista, N675
Eucardium, N585
Eucharia, N540
 Euchasma, N295
 Euchasmella, N867
 Euchondria, N340
Euchondriidae, N340
 Euchondriinae, N219, N340
 Euciroa, N49, N856
Euciroidae, N854
 Euclusaea, N867
Eucorbicula, N666
 Eucrassatella, N576, N911
 EUEDESMODONTIDA, N119, N214
Eufira, N464
 Eufistulana, N699
 Eugaimardia, N644
Eugenoceramus, N320
Euglesa, N669-N670
 EULAMELLIBRANCHIA, N18, N118-N120, N210, N212, N214
 EULAMELLIBRANCHIÉS, N119
Eulopia, N499
 Euloxa, N651
 Euloxidae, N221, N650
 Eumarcia, N683
Eumegalodon, N742
Eumicrotis, N341
Eumodiolus, N278
Eumontrouzieria, N637
Eumorphotis, N115, N337
Eumulleria, N466
Eumytilus, N271
 Eupera, N670
 Euphenax, N329
Euphira, N464
Euphrata, N442
Euphyoceramus, N315
Eupisidium, N669
Eupoleme, N527
 Euptera, N308
Euradiolites, N806
 Eurhomalea, N683
Eurinia, N420
Euryanodon, N463
Euryceramus, N320
Eurychasma, N832
 Eurydesma, N115, N291
 Eurydesmatidae, N115
 Eurymya, N299
 Eurymyella, N394
Eurynaia, N420
Eurynia, N420
 Eurytellina, N615
Eusalacia, N686
Eusebia, N261
Eustylon, N604
Eutapes, N685
 EUTAXODONTA, N118-N119
 EUTAXODONTIDA, N214
Eutellina, N621
Euterentia, N681
 Euthydesma, N245
 Eutivela, N674
 Eutrephoceras, N197
 Eutylus, N714
 Euvola, N367
 Euxinocardium, N591
 evolution, N109, N129, N183; rudist, N766; Thyasira, N200
 Excellichlamys, N366
 excurrent, N104, N759
 exhalant, N104
 exhalant current, N11
Eximiothracia, N850
 Exitopinna, N283
Exocardium, N589
Exocholeda, N235
 Exochorhynchus, N831
Exodiberus, N276
 Exogyra, N45, N70, N72, N116, N154, N156-N157
 exogyrate, N104
 Exogyrinae, N219
Exoleta, N680
 Exosiperna, N276
 Exotica, N625
 Exputens, N331
 external features, N40
 extinction (rudists), N765
 Eydyusiana, N862
 eyes, N29
Ezocallista, N679
 Ezonuculana, N237

 Fabagella, N854
 Fabella, N540
 Fabulina, N615
 falciform, N104
 Falcimytilus, N279
Fallartemis, N679
 Falsusiana, N862
 fascicle, N104
 fasciculated, N104
 Fasciculiconcha, N337
 Fascinewellipecten, N333
 Fastimysia, N530
 FAVRE, N485
 fecal pellets, N22
 Felania, N515
 Felaniella, N516
 Felicia, N267
 Feliciella, N861
Felipes, N359
Fenestella, N383
 Fenestricardita, N551
 Ferganoconcha, N410
 Ferganoconchidae, N117, N219, N410
 Ferrata, N861
 Ferrazia, N115, N745
 FÉRUSSAC, N208
 fibrous layers, N58
 fibrous ligament, N104
Ficusocorbula, N696
 filaments, N16
 Filatovaella, N566
 FILIBRANCHIA, N18, N119-N120, N210, N212, N214
 FILIBRANCHIÉS, N119
Filius, N233
 Filosina, N668
 Fimbria, N513
Fimbriella, N512
 Fimbridae, N117, N220, N513
Finlaria, N629
Finlaya, N678
 Finlayella, N615
 FISCHER, N79, N118, N210, N827
Fischeria, N629
 Fissilunula, N646
Fistula, N610
 Fistulana, N67
Fistulana, N699, N717, N730
Fistulanigenus, N730
 flabelliform, N104
 Flabellipecten, N369
Flabellomya, N827
 flank, N46, N104
Flavamussium, N350
 Flaventia, N683
Flavomala, N634
 FLEMING, N208
 Flexicorbula, N694
Flexipecten, N357
Flexiptis, N424
 Flexopecten, N357
Flodacna, N594
 Florimetis, N626
Florisarka, N634
 Fluctiger, N672
Fluminea, N669
Fluminina, N669
Flumininea, N669
 fluted, N104
 Fluviolanatus, N279
 Foegia, N859
Fogiella, N842
 folds, N69, N104
 foliated structure, N76
 FOLIOBRANCHES, N119
 FOLIOBRANCHIA, N119
 Follmannia, N287
Folmannella, N302
Fontinalina, N669
 food groove, N20
 foot, N5, N11, N14, N104
 Foramelina, N326
 foramen, N104
 Fordilla, N109, N868
Formesulus, N714
 Formosulus, N714
 Fortipecten, N371

- Fossacallista, N678
 Fossarina, N669
 fosset, N104
 fossette, N105
 Fossicula, N464
 Fossula, N464
 Fossilarca, N263
 Fossilites, N759, N773, N777,
 N808
 Foveadens, N844
 Fractoceramus, N320
 Fragilia, N626
 Fraginae, N220, N588
 Fragum, N588
 FRANC, N212-N214
 FRECH, N581
 free-living epifaunal bivalves,
 N159
 Freia, N570
 Freiastarte, N570
 Frenamya, N844
 Frenouelliella, N478
 fresh-water bivalves, N95
 Friersonia, N453
 Frigichione, N681
 Frigidocardium, N589
 Fronsella, N530
 Fuchsella, N823
 Fugleria, N252
 Fulcrella, N541
 fulcrum, N105, N412
 Fulgida, N278
 Fulpia, N668
 Fulvia, N590
 Fundella, N329
 funnel plates, N105
 Furcella, N739
 Fusconia, N439
 Fusconia, N439
 Fusculusiana, N862
 fusion layer, N58, N105

 Gabillotia, N444
 Gabinarca, N263
 Gafrarium, N67, N672
 Gaiderope, N378
 Gaimarda, N644
 Gaimardia, N644
 Gaimardiacea, N211, N221, N644
 Gaimardiadae, N644
 Gaimardiidae, N221, N644
 Galactella, N263
 Galataea, N629
 Galatea, N628
 Galateia, N629
 Galateola, N629
 Galathea, N629
 Galaxura, N850
 Galeomma, N533
 Galeommatidae, N220, N533
 Galeommella, N535
 Galeommidae, N533
 Galileja, N669
 Gallandriana, N862
 Gallicusiana, N862
 ganglia, N26

 gape, N47, N105
 Gardnerella, N500
 Gari, N45, N630
 Garidae, N629
 Garinae, N629
 Garum, N631
 Gastrana, N626
 Gastrana, N626
 Gastranella, N689
 Gastranopsis, N615
 gastric shield, N22
 gastrochaena, N67, N699
 Gastrochaenacea, N211, N221,
 N699
 Gastrochaenadae, N699
 Gastrochaenidae, N118, N221,
 N699, N741
 Gastrochaenites, N699
 Gastrochaenoecium, N699
 Gastrochaenolites, N699
 Gastrochaenopsis, N700
 GASTRODEUTEIA, N119
 Gastrodiana, N862
 gastro-intestinal arteries, N24
 GASTROPEMPTA, N119
 GASTROPODA, N4, N79, N195,
 N197
 GASTROPROTEIA, N119
 GASTROTETARTIKA, N119
 GASTROTRITEIA, N119
 GEINITZ, N120
 Geloia, N668
 Geltena, N600
 Gemellima, N391
 Gemma, N159, N164, N166-
 N167, N681
 Gemmellaria, N790
 Gemminae, N221, N681
 Genaxinus, N510
 geometry of shell, N84
 Geratrigonia, N480
 Gerlus, N380
 Germanoceramus, N317
 Gervilla, N308
 Gervilla, N308
 Gervillaria, N308
 Gervillea, N308
 Gervilleia, N308
 Gervilleioperna, N325
 Gervillella, N308
 Gervillella, N45, N47, N115-
 N116, N308
 Gervillia, N42, N115, N117,
 N144, N308
 Gervilliopsis, N308
 Gestrouana, N862
 Geukensia, N279
 Gibboconcha, N863
 Gibbolucina, N502
 Gibbomodiola, N278
 Gibbonucula, N231
 Gibbopleura, N244
 Gibbosula, N439
 Gigantopecten, N358
 Gilbertharrisella, N678
 Gilbertwhitea, N258

 gill, N5
 gill retractor, N38
 gill retractor muscle, N105
 Gingillum, N865
 Ginorga, N578
 Girardotia, N832
 Girtyana, N238
 Girtypecten, N143, N338
 Gitocentrum, N708
 Gitoventrum, N708
 Glabaris, N463
 Glabrae, N477
 Glabris, N463
 Glabropecten, N357
 Glacipisum, N669
 Glans, N551
 Glaucion, N385
 Glaucoderma, N303
 Glaucomya, N690
 Glaucomyidae, N690
 Glaucanoma, N690
 Glaucanome, N63, N690
 Glaucanomeidae, N690
 Glaucanomella, N690
 Glaucanometta, N690
 Glauconomidae, N221, N690
 Glauconomya, N690
 Glauconomyidae, N690
 Glaucus, N303
 Glebula, N453
 GLIBERT & VAN DE POEL, N331
 Glibertia, N559
 Globivenus, N672
 Globocardium, N588
 globose, N105
 glochidium, N96
 Glochidium, N442
 Glomene, N515
 Glomus, N239
 Gloria, N245
 Gloripallium, N357
 Glossacea, N16, N63, N211,
 N221, N657
 Glossidae, N221, N657, N768
 Glossites, N820
 Glossocardia, N655
 Glossoderma, N657
 Glossus, N45, N51, N58, N60,
 N63, N88, N90, N657, N660,
 N662, N756, N768
 Glumebra, N730
 Glyciana, N862
 Glycilima, N265
 Glycimeridae, N119, N700
 Glycimeris, N700
 Glycydonta, N688
 Glycymeracea, N120
 Glycymerella, N267
 Glycymeridae, N267
 Glycymerididae, N12, N159,
 N218, N267
 Glycymeridinae, N218, N267
 Glycymeris, N38, N44, N60,
 N62-N63, N77, N86-N87,
 N99, N141, N159, N183,
 N267

- Glycymerita, N267
Glycymerula, N267
Glyptamusium, N350
 Glyptoactis, N556
Glyptocardia, N244
Glyptodesma, N299
 Glyptoleda, N239
 GMELIN, N389
Gnathodon, N601
Gnathonoceramus, N317
Gnesioceramus, N315
 gnomon, N84
 Gobraeus, N631
 Goldfussia, N861
 Gomphina, N683
 Gomphinella, N683
 Gomphomarcia, N685
 gonads, N29
Gonamblyus, N437
 Gonidea, N423
 Gonilia, N566
Gonimya, N832
 Gonimyrtea, N500
 Goniocardium, N595
 Gonioceramus, N315
 Gonioclasma, N715
 Goniocuna, N560
Goniodon, N233
Goniodus, N306
 Goniomactra, N604
Goniomeris, N832
Goniomya, N68-N69, N117, N832
Goniomyaa, N832
 Goniophora, N394
Goniophora, N854
Goniophorella, N297
 Goniophorina, N396
Goniosoma, N650
Gonodon, N514
Gonomya, N832
Gonyomya, N832
 Goodallia, N57, N566
 Goodalliopsis, N520
 Goossensia, N551
 Gorjanovicia, N808
 Gosseletia, N111, N287
Gotodonta, N228
 Gouldia, N54, N673
Gouldia, N577
 Gouldiopa, N673
Gradilucina, N499
 Grammatodon, N116, N258
 Grammatodon group, N258
 Grammatodonax, N628
Grammatodontidae, N257
 Grammatodontinae, N218, N257
 Grammatomya, N66, N631
 Grammysia, N819
 Grammysiidae, N221, N819, N861
 Grammysioidea, N820
Grammysiopsis, N820
Grandaxinaea, N267
Grandiarca, N254
 Grandidieria, N418
Grandipecten, N358
 Granicorium, N685
 Granigeriana, N862
Granoarca, N252
 Granocardium, N585
Graptamusium, N350
Graptomaias, N425
 Grateloupia, N673
Grateloupina, N673
Gratelupia, N673
 Gravidiana, N862
 GRAY, N118, N208
 Gregariella, N275
Greslya, N840
Gressluya, N840
 Gresslya, N117, N839, N840
Gresslyia, N840
 Grippina, N699
 GROBBEN, N120
 grooves, N69
 Grottriana, N563
 growth lines, N67, N105
 growth ruga, N67, N105
 growth thread, N105
 Gruenewaldia, N473
 Grundensia, N541
 — Gryphaea, N45, N76, N89-N90, N154, N156-N157
 gryphaeate, N105
 Gryphaeidae, N219
 Gryphaeinae, N219
 Gryphellina, N389
Gryphorhynchus, N311
 Guebhardia, N865
 Guerangeria, N397
Guetera, N730
Guettera, N730
Guianadesma, N847
 Guizhoupecten, N338
 Gyropleura, N755, N783
Gyropleuridae, N781
 Gythemon, N648
 HAAS, N3, N208, N213
 Haasica, N464
 Haasiella, N444
 Haasodonta, N423
 Haastina, N513
 habitat (Thyasira), N190
 habitats and food (Bivalvia), N5
Habra, N636
 Hadrodon, N865
 Haenleinia, N317
Haicana, N700
Haidaia, N485
 Hainania, N633
 Halicardia, N857
 Halicardissa, N857
 Haliotis, N88
 Haliris, N857
 Halistrepta, N850
 Halobia, N115, N344
 Halobiidae, N115
Halobiidae, N342
Halocochia, N857
 Halodakra, N650
Halonanus, N264
 Halonympha, N854
 Hamacuna, N560
Hapalomia, N842
Hapalomya, N842
Haploceramus, N315
Haplomochlia, N631
Haploscaptha, N315
Haplothaerus, N463
 hard parts, N40
 Hardaghia, N759, N774, N813
 Harlea, N698
Harmandia, N462
Harpax, N378, N637
 HARRINGTON, N111
 Hartwellia, N646
 Harvella, N600
Hassbergia, N588
Hastasia, N716
 Hataiarca, N860
 Hataiyoldia, N239
 Hatasia, N716
Hatoisia, N716
Haumea, N357
 haustorial larva, N96
Hawaiarca, N252
 HEATH, N38
 Heckingiana, N862
 Hecuba, N138, N168, N628
 Hedecardium, N586
 Hefteria, N473
 height, N42, N105
 Heikea, N249
 HEINZ, N315
 helicospirals, N86
 Heligmina, N329
 Heligmopsis, N329
Heligmus, N329
Helioceramus, N320
Hellia, N518
 HEMIBRANCHIA, N119
 Hemicardia, N588
Hemicardia, N588
Hemicardium, N588
 Hemicorbicula, N666
Hemicyclodonta, N608
 Hemicyclonosta, N608
Hemicycloster, N608
Hemicyclostera, N608
 HEMIDAPEDONTA, N119
Hemidiscors, N589
Hemidon, N441
 Hemidonax, N629
Hemilastena, N424
 Hemilepton, N519
 Hemimactra, N602
 Hemimetis, N72, N615
 Heminajas, N473
Hemiodon, N447
 Hemipecten, N38, N354
Hemiplacuna, N383
 Hemiplicatula, N383
 Hemisolasma, N862
Hemistena, N424
 Hemisterias, N868
Hemitapes, N684-N685
Henslowiana, N669
 hepatic arteries, N24

- Hercodon, N621
 Hercynella, N247
 Here, N496
 Herella, N496
Heritschia, N367
 hermaphroditism, functional, N29
 hermaphroditism, protandrous, N29
Hermione, N686
Heroceramus, N315
 Herouvalia, N631
 HERRMANNSEN, N119
 Hertellina, N615
 HERTWIG, N118
 Herzogina, N571
 Heteranomia, N385
Heterocaprina, N785
Heterocaprinidae, N784
 Heterocardia, N604
Heteroceramus, N315
 Heteroclidus, N844
HETEROCOCHIA, N119, N214
 Heterodicerias, N762, N768, N776, N778
Heterodiceratidae, N778
 Heterodiceratinae, N221, N754, N778
 Heterodonax, N633
 heterodont, N52, N105
HETERODONTA, N62-N64, N118-N120, N210-N211, N213-N215, N218, N220, N491
HETERODONTEN, N119
HETERODONTIDA, N118-N119, N214
 Heteroglypta, N633
 Heteromacoma, N626
Heteromactra, N538, N595
Heteromya, N700
HETEROMYARIA, N118-N119
 heteromyarian, N34, N105
Heteropecten, N336
 Heteropholas, N715
 Heteropis, N572
 Heterotrigonia, N478
 Heterunio, N416
Hettangia, N640
 Heudeana, N423
 Hexacorbula, N694
Hiata, N711
 Hiatella, N7, N32, N169-N170, N195, N700
Hiatella, N846
 Hiatellacea, N211, N221, N700
 Hiatellidae, N52, N76, N195, N221, N700
Hiatula, N691
 HICKS, N112
Hijitrigonia, N488
 Hilberia, N357
 Hilgardia, N237
Himanthopoda, N327
Himantopoda, N327
Himella, N847
 Himeraelites, N757, N761, N783
Himotopoda, N327
 Hina, N672
 HIND, N407
Hindsia, N541
 Hindsella, N541
 Hinemoana, N687
 hinge, N42, N105, N412;
 crassatellacean, N910-N912
 hinge axis, N40, N105
 hinge line, N105
 hinge margin, N40, N105
 hinge notation, N908
 hinge plate, N49, N105, N412
 hinge structures, N96
 hinge teeth, N47, N105
Hinnia, N364
 Hinnites, N16, N60, N101, N142, N154, N156, N170, N208, N314
 Hinnites group, N364
Hinnus, N364
Hippagus, N274
Hippella, N558, N855
Hippocardia, N859
 Hippochaeta, N322
 Hippomya, N397
 Hippopodiidae, N220, N582
 Hippopodium, N116, N583
 Hippopus, N595
 Hippuritacea, N211, N221, N749, N766, N768
 Hippuritella, N755, N776, N802
 Hippurites, N76, N749-N750, N758, N760, N764, N768-N769, N771, N774, N776, N799, N801, N803
 hippuritid pillars, N774
 Hippuritidae, N53, N221, N749, N754, N756, N760-N761, N763-N764, N766, N799, N815, N861
HIPPURITOIDA, N214, N218, N221, N741, N766
Hirudinaria, N533
Hirundigenus, N303
 Hispaniana, N862
 Hitia, N541
 Hochstetteria, N270
Hochstetteria, N270
 Hochstetterina, N270
 Hoernesia, N115, N310
 Hoernesiella, N312
 Hoferia, N265
Hoffetrigonia, N489
 Hokonua, N376
 Holandriana, N862
 HOLDHAUS, N831
Holdhausiella, N643
Holocraspedum, N380
Holopholas, N708
 Homalina, N615
Homaloceramus, N315
Homoeodesma, N850
 homogeneous structure, N76
 Homomya, N116-N117, N832
HOMOMYARIA, N119
 homomyarian, N32, N105
 Honeoyea, N295
Honeymania, N229
 Hoplomylus, N861
 Hoploscapites, N197, N200
 Horiiodacna, N592
 Horiopleura, N759, N785
 Hormomya, N273
Horusia, N418
 Hubertschenckia, N664
 Humbertiana, N862
 Humilaria, N688
 Humphreyia, N858
Humphreysia, N858
 Hunkydora, N847
 Huxleyia, N269
 Hyalokellia, N537
 Hyalopecten, N354
Hyatelladae, N700
 HYATT, N83, N100
 Hybolophus, N576, N909, N911
Hydrodaetes, N308
Hylopecten, N354
Hylotrya, N732
 Hynnites, N364
 hyote spines, N105
 Hypanis, N593
 Hypelasma, N781
Hyperotis, N730
Hyperotus, N730
 Hyphantosoma, N675
Hypochaeta, N322
Hypogaea, N707
Hypogaeoderma, N707
Hypogella, N611
 hypoplax, N64, N105, N705
 hypostracum, N105
 Hypotrema, N326
 Hypoxytoma, N345
Hyppurites, N801
 Hyrcania, N593
 Hyrcanomya, N593
Hyria, N457
Hyriana, N457
 Hyridella, N460
Hyridella, N461
Hyridunio, N461
 Hyriinae, N219, N457
 Hyriopsis, N423
Hyris, N457
 Hysteroconcha, N68, N138, N168, N675
 Iacra, N637
 Ibotrigonia, N480
 Icanotia, N635
 Icanotiidae, N220, N635
 Ichthyosarcolithes, N789, N795
Ichthyosarcolithus, N795
Idas, N279
 Idasola, N279
Idioceramus, N315
Idioteredo, N729
 Idonearca, N260
Idothea, N513
 Idriniana, N862

- Iheringella*, N464
Iheringia, N590
 Iliochione, N686
 Illesca, N496
 Illionia, N504
 Illuviosiana, N862
Ilyiria, N457
 imbricate, N69, N105
 Immanitas, N790, N817
Imparilarca, N256
Inaequiceramus, N315
Inaequiptecten, N358
 Incacardium, N585
 Incanopsis, N261
 Incrassatiana, N862
 incremental line, N105
 incurrent, N105, N759
Indacar, N254
Indica, N668
 Indocrassatella, N576
 Indogrammatodon, N69, N116, N258
Indonaia, N428
 Indopecten, N372
 Indoplacuna, N385
 Indopseudodon, N434
 Indotrigonia, N481
Inequarista, N736
 inequilateral, N45, N105
 inequivalve, N105
 inequivalve bivalves, N72
 infaunal bivalves, N141, N163
 inflated, N105
 inflation, N42, N105
 inhalant, N105
 inhalant current, N11
 inner demibranch, N116
 inner lamellae, N18
 inner ligament, N105
 Inoceramidae, N76, N117, N138, N142-N144, N147, N175, N199, N219, N314
Inoceraminae, N314
Inoceramus, N72, N76, N78, N115, N143-N144, N197, N200, N315
Inoceromya, N317
 Inoperna, N278
 Integricardium, N589
 INTEGRIPALLIATA, N119
 integripalliate, N36, N105
 INTÉGROPALÉALES, N119
 INTEGROPALLIALIA, N119
 interband, N105
 interdendum, N105
 interior shell structures, N175
 internal calcareous structures, N47
 interspace, N69, N105
 interval, N69, N105
 INTESTINA, N4
 intestine, N22
Intomodesma, N290
 inverse, N105
 Inversidens, N423
 Iotrigonia, N481
Iouannetia, N718
Iphegenia, N629, N855
 Iphigenia, N629
 Iraqitellina, N615
 IREDALE, N208
Iridea, N441
Iridella, N460, N464
 Iridina, N464
Irinia, N593
Irona, N683
 Irus, N683
Isarcha, N633
 Ischadium, N273
 Ischyrinia, N865
 Ischyrodonta, N400
 Isoarca, N241
 Isoarcidae, N218, N241
 Isocardia, N21
Isocardia, N657
Isocardiacea, N118, N121, N657
Isocardiadae, N657
 Isoconcha, N541
 Isocrassina, N563
Isoculia, N818
 Isocyprina, N117, N647
Isodoma, N668
 Isodomella, N668
Isodon, N639
 isodont, N53, N105
 ISODONTA, N119
Isodonta, N639
Isodontidae, N639
 ISOFILIBRANCHIA, N119
 ISOFILIBRANCHIDA, N214
Isognoma, N322
 Isognomon, N33, N37-N38, N45, N60-N61, N63, N78, N115-N116, N132, N143-N147, N151-N153, N174-N175, N179, N182, N322
 Isognomonidae, N76, N115, N138, N143-N144, N151, N219, N321
Isognomontidae, N322
 Isogonum, N322
Isolea, N237
 Isolimea, N391
 Isomonina, N383
 isomyarian, N32, N105
Isonuculana, N237
 Isopristes, N579
 Isorobitella, N531
 Isorropodon, N656
 Isotancredia, N640
 Issina, N530
Ivania, N803
 Ixartia, N850
 Izumia, N648
 Izumicardia, N552
 JACKSON, N79, N110
Jacksonarca, N252
 Jacqueminiana, N862
Jacquesia, N643
 Jactellina, N615, N625
 Jagolucina, N504
 Jagonella, N508
Jagonia, N494
 Jagonoma, N498
Jahnia, N247
 Janeia, N243
Janeira, N243
Janera, N366
Janira, N366
 Janschinella, N697
Janupecten, N363
Jaronia, N418
Jataronus, N518
 JAWORSKI, N36
Jaworskia, N367
 Jaworskiella, N481
 JEFFERIES & MINTON, N9
 Jesonia, N548
Joachymcardium, N244
Joachymia, N297
Joannina, N280
Joannisia, N515
 Joannisiana, N862
Joannisiella, N515
Jodamia, N81
 Jolya, N278
 JONES, N408
 Jouannetia, N65, N702, N704, N718
 Jouannetiinae, N221, N702, N704, N717
Jouannetinae, N717
 Joufia, N758, N774, N804, N815
 Jourdheuiliana, N862
 Jousseameana, N862
Jousseaumia, N542
 Jousseaumiella, N542
Juanetia, N718
Juannetia, N718
 Jugosus, N446
Jukes-Brownia, N675
 Jukesena, N683
 Juliacorbula, N694
Junomia, N235
 Jupiteria, N235
 Jurassicardium, N589
 Juxtamusium, N357
 Kaibabella, N566
 Kakahuia, N679
Kaladacna, N592
 Kalayoldia, N240
 Kalentera, N545
Kalenteridae, N543
 Kalliphenga, N418
Kamaleda, N235
 Kanecharaia, N679
 Kaneoha, N523
 Katadesmia, N241
 Katelysia, N683
 Katherinella, N675
Kathocardia, N586
 Katzeria, N813
 KAUFFMAN, N135, N156, N184
 keel, N105
 Keenaea, N589
Kefersteinia, N473
Keleria, N813
Kelleria, N813

- Kellia, N13, N523
Kelliadae, N522
 Kelliella, N652
Kelliella, N652
Kelliellidae, N221, N651
Kelliidae, N220, N522
Kelliola, N529
Kelliopsis, N530
Kellyella, N652
Kellyellidae, N651
Kennerleya, N844
Kennerleyia, N844
Kennerlia, N844
Kennerlyia, N844
Kercomya, N845
 Kereia, N680
Kermadysmea, N631
Kidderia, N644
 kidneys, N24
Kidodia, N468
 Kija, N411
Kikaiarca, N860
Kinerkaella, N293
Kinerkaellina, N295
Kipia, N789, N795
Kistinaia, N418
Kitsonia, N656
 KLINGHARDT, N755-N756
 KLINGHARDT & WIONTZEK, N800
Klinoptera, N301
 KOBAYASHI, N488
Kobayashiites, N310
Kobyia, N860
Kochia, N302
Kochiidae, N219, N302
Kochiinae, N302
Kodynia, N302
Koenenia, N233
Kolymia, N290
Kona, N530
 KONINGCK, DE, N114
 KOROBKOV, N910
Korobkovitrigonia, N481
Kotorapecten, N370
Kralovna, N244
Krumbeckia, N745
Krumbeckiella, N314
 KÜHN, N815
Kuehnia, N758, N808
Kugleria, N762, N781
Kuia, N672
Kumatrigonia, N478
Kummelia, N700
Kummelia, N741
Kuphinae, N221, N702, N739
Kuphus, N67, N739
Kurinuia, N857
Kymatox, N601
Kyphus, N740

Labayophorus, N547
 labial palps, N5, N16, N18
Labiosa, N606
Labis, N277
Lacanniciana, N862
Lacinea, N518

Laciolina, N615
Lactemiles, N535
Lacustrina, N669
 LADD, N136
Laevicardiinae, N220, N589
Laevicardium, N14, N16, N589
Laevicirce, N672
Laevicordia, N857
Laevipinna, N865
Laevirostris, N418
Laevitrigonia, N21, N481
Laevitrigonia, N481
Lahilleona, N590
Lahillia, N590
Lahilliidae, N220, N590
Lajonkaira, N689
Lajonkairia, N689
 LAMARCK, N208, N389, N476, N749
 LAMEERE, N118
lamella, N17-N18, N105
 lamellar layer, N58
 lamellar ligament, N105
LAMELLIBRANCHES, N213
LAMELLIBRANCHIA, N3-N4, N213
LAMELLIBRANCHIATA, N3, N213, N227
Lamelliconcha, N675
Lamellidens, N423
 lamelliform, N105
Lamellileda, N237
Lamellinucla, N231
Lamellodonta, N51, N110-N111, N400
Lamellodontidae, N216, N219, N400
 lamina, N105
Laminaria, N607
Lampeia, N850
Lamproscapha, N463
Lamprotula, N439
Lampsilinae, N219, N450
Lampsilis, N450
 LAMY, N910-N911
Lamya, N275
Lanceolaria, N424
 lanceolate, N105
Landinia, N574
Langsonella, N310
Lanilda, N378
Lanistes, N276
Lanistina, N276
 LANKESTER, N208
Lapeirouseia, N813
Lapeirousella, N813
Lapeirousia, N769, N804, N813
Lapeirousinae, N221, N758, N760, N766, N769, N813
Lapidosus, N862
Larkinia, N254
Larma, N678
 larval development, N91
 larval stage (Unionacea), N413
Lasaea, N13, N21, N520
Lasacina, N541

Lasaeokellya, N523
Lasaeoneaera, N530
Lasea, N520
Laseadae, N519
lasidium larva, N96
Laskarevia, N817
Lasmigona, N448
Lasmonos, N454
Lastena, N424
Lastenes, N424
Latemaria, N391
 lateral hinge system, N98, N105
 lateral tooth, N49, N106
Laternula, N49, N845
Laternulacea, N119
Laternulidae, N13, N37, N76, N222, N844
Laternulina, N845
Latiarca, N260
Latitrigonia, N478
Latona, N628
Latosiliqua, N633
 LATREILLE, N118, N208
Laubeia, N865
Laubriereia, N530
Lavigno, N635
Lavignon, N635
Lavignona, N635
Lavignonus, N635
Laxicava, N700
Lazaria, N549
Lazariella, N549
Lebescontia, N868
 lecithotrophic larval development, N91
Leckhamptonia, N563
Lecompteus, N279
Leda, N235
Ledaspina, N235
Ledella, N235
Ledidae, N235
Ledina, N237
Ledina, N237
Ledopsis, N229
 left valve, N42, N106
Legrandina, N538
Legumen, N683
Legumenaia, N424
Legumia, N455
Leguminaia, N412, N424
Leguminaria, N613
Leila, N465
Leinzia, N820
Leiochasmea, N536
Leiomya, N854
Leiomyalina, N291
Leionucula, N231
Leiopecten, N334
Leiopectinella, N334
Leiopectinidae, N219, N334
Leiopteria, N71, N138, N144, N301
Leiosolenus, N278
Leiovirgus, N436
Lembulus, N20, N164-N165, N167, N236

- Lembulus*, N231
Lemiox, N452
 Lemotheuxiana, N862
Lenella, N317
 length, N42, N106
 Lens, N428
 lenticular, N106
Lenticularia, N494
 Lentidiinae, N221, N696
Lentidium, N696
Lentillaria, N494
Lentipecten, N348-N349
 Lentipecten group, N349
Lepidocardia, N678
Lepidodesma, N446
Lepilucina, N492
Lepirodes, N533
Leporimetis, N626
Leproconcha, N325
Leptaena, N113
Leptanodonta, N446
Leptaxinus, N510
Leptesthes, N666
Leptina, N608
Leptocardia, N588
Leptochondria, N115, N338
Leptodea, N453
Leptodesma, N112-N113, N299
Leptodomus, N475
Leptomya, N637
Leptomya, N845
Leptomylaria, N637
Lepton, N527
Leptonacea, N16, N119, N121, N206, N211, N220, N518
Leptonaias, N425
 Leptonidae, N220, N526
Leptopecten, N132, N138, N142-N144, N146, N357
Leptosiphon, N668
Leptosolen, N183, N611
Leptospatha, N463
Leptospisula, N604
Leptotrigonia, N488
Leptynoconcha, N297
Lepyrodes, N533
 Letourneuxiana, N862
Leuconix, N709
Leuconyx, N709
Leucoparia, N606
Leucosilla, N862
Leucothea, N678
 Leukoma, N688
Leuroactis, N555
Levanderia, N536
Leviconcha, N475
Levimarcia, N684
Levimyrtaea, N508
Lexingtonia, N425
Libitina, N655
Libitiniidae, N655
Libratula, N536
Libyaconchus, N865
Lichas, N859
Liebea, N291
 ligament, N4, N58, N106;
 rudist, N764
 ligament fulcrum, N106
 ligament groove, N106
 ligament pit, N106
 ligament ridge, N106
 ligament supports (rudist), N769
 ligament suture, N46, N106
 ligamental structures (rudist), N755
 ligamenture, N174
Lignopholas, N715
Ligula, N496, N635
Ligumea, N454
Ligumia, N454
Ligumina, N455
Lilangina, N312
Liliobankia, N734
 Lima, N37-N38, N42, N58, N132, N141-N142, N150-N153, N170, N175, N197, N385
Limacea, N15, N18, N36, N47, N121, N211-N212, N219, N385
Limacula, N389
Limaea, N389
Limanomia, N334
Limanomva, N334
Limarca, N270
Limaria, N389
Limaria, N385
Limaridia, N385
Limatula, N389
Limatulella, N389
Limatulina, N336
Limea, N389
Limecola, N623
Limicola, N621
 Limidae, N9, N12, N15, N29, N36-N37, N76, N114-N116, N151, N153, N159-N160, N162, N195, N210, N212, N219, N385
Limigenus, N385
Limipecten, N339
Limnadea, N455
Limnaea, N415
Limnaeoderma, N415
Limnoarca, N391
Limnobasilissa, N423
Limnocardium, N591
Limnocyrena, N861
Limnodacna, N592
Limnoica, N862
Limnopageta, N592
Limnopappia, N592
Limnoperna, N273
Limnoscapha, N423
Limoarca, N391
Limopsacea, N63, N211, N218, N264
Limopsidae, N62, N218, N264, N861
Limopsilla, N265
Limopsis, N38, N47, N58, N60-N63, N79, N265
Limopsista, N265
Limoptera, N113, N138, N301
Limosina, N670
Limula, N389
Linatulina, N389
Linearia, N621
 lines, N69
Linettia, N511
Linga, N496
Lingulocaris, N868
 LINK, N389
 LINNÉ, N4, N45, N208
Linotrigonia, N483
Linsdallia, N325
Lintellaria, N494
Linter, N264
Lintoxia, N439
Linucula, N231
Lioberus, N276
Liocardia, N865
Liocardium, N589
Lioconcha, N678
Liocyma, N684
Liodonax, N633
Lionelita, N523
Liodistha, N852
Liostrea, N116, N331
Liotellina, N613
Liothyris, N621
Liotrigonia, N473
Liouvillea, N444
 LIPOCEPHALA, N227
 LIPODONTA, N119-N120
Lipodonta, N442
 LIPODONTIDA, N214
 lips, N21
Lirodiscus, N571
Liropecten, N357
Lirophora, N686
Lirotarte, N499, N580
Lison, N84, N86, N88
Lissarca, N266
Lissarcula, N265
Lissochlamis, N354
Lissochlamys, N354
Lissopecten, N357
Listera, N610, N635
Litharca, N254
 Lithotidae, N219
Lithobia, N397
Lithocalamus, N790, N817
Lithocardium, N595
 lithodesma, N49, N106
Lithodroma, N278
Lithodomus, N276
Lithoglyphus, N276
Lithophaga, N7, N114, N141, N165, N169, N276
Lithophagella, N655
Lithophagellidae, N655
 Lithophaginae, N219, N276
Lithophagus, N276
Lithorhadia, N237
Litigiella, N520
Litschkovitrigonia, N487
Lituyapecten, N371
 liver, N21

- Lobaria*, N633
 lobes, N11
Locardiana, N862
 logarithmic spiral, N85
Loncosilla, N639
Longimactra, N598
 longitudinal, N106
 longitudinal vein, N25
Lonoa, N637
Loparia, N648
Lopatinia, N261
Lopha, N70, N115-N116, N155-N156, N171, N178, N380
Lophinae, N219
Lophocardium, N589
Lophonychia, N287
Loringella, N265
Loripes, N4, N54, N496
Loripinus, N502
Loripoderma, N861
Lorttiella, N460
Lostena, N424
Lovellia, N606
 LOWENSTAM, N77, N134
Loxocardium, N585
Loxoglypta, N615, N626,
Loxopleurus, N439
Loxopteria, N302
Loxoptychodon, N666
Lucasiella, N380
Lucina, N21, N30, N32, N44,
 N77, N79, N111, N164, N167-
 N168, N195, N197, N199-
 N200, N492
Lucinacea, N16, N63, N98, N113,
 N119, N121, N167-N168,
 N181, N183, N188, N199,
 N206, N211, N220, N491
Lucinadae, N492
Lucinaxinus, N520
Lucinella, N506
Lucinidae, N6, N12-N14, N16,
 N35-N36, N51, N76, N113,
 N117, N195, N200, N206,
 N220, N492, N861
Lucinigenus, N508
 LUCININA, N119
Lucininae, N220, N492
Luciniola, N498
Lucinisca, N492
 LUCINODONTA, N120, N910
 lucinoid type, N56, N106, N120
Lucinoma, N164, N167, N500
Lucinopsis, N689
Luciploma, N681
Ludbrookia, N557
Lunarca, N254
Lunatia, N197-N200
Lunopsis, N265
Lunulacardiidae, N219, N295
Lunulacardium, N295
lunule, N46, N106
Lunulicardia, N588
Lusitaniana, N862
Lutaria, N604
Luteacarnea, N437
Luteamusium, N350
Lutetia, N653
Lutetina, N538
Lutraria, N6, N13, N48, N62-
 N63, N604
Lutrariinae, N220, N604
Lutricola, N635
Lutricularia, N636
Lutromactra, N604
Lutrophora, N604
Luzonia, N854
Lycettia, N273
Lycodes, N743
Lycodus, N743
Lymnadea, N455
Lymnadia, N455
Lymnium, N415
Lymnocardiidae, N117, N220,
 N590, N861
Lymnocardiinae, N220, N591
Lymnocardium, N591
Lyntoxia, N439
Lyonsia, N21, N63, N846
Lyonsiella, N857
Lyonsiidae, N13, N16, N222,
 N845
Lyratellina, N615
Lyridon, N478
Lyriodon, N478
Lyriomyophoria, N475
Lyriopecten, N113, N334
Lyrocardium, N589
Lyrodesma, N50, N112-N113,
 N165, N471
Lyrodesmatidae, N112-N113,
 N219, N471
Lyrodobankia, N734
Lyrodon, N478
Lyrodus, N727
Lyronucula, N471
Lyropecten, N357
Lyropectin, N357
Lysianassa, N832

Maackia, N643
Mabellarca, N254
Macalia, N626
Macalina, N626
Macaliopsis, N615
Maccoyella, N346
Maceris, N518
 MACGILLAVRY, N761
Macha, N638
Machadoiana, N862
Machaena, N610
Machaera, N613
Machaerodonax, N628
Machomya, N832
Machrochlamys, N358
Macilentiana, N862
Macoma, N9, N22, N42, N44-
 N45, N77, N166, N195, N199,
 N623
Macominae, N220, N623
Macomona, N616
Macomopsis, N623
Macoploma, N623
Macridiscus, N683
Macrocallista, N71, N164, N166,
 N677
Macrochlamis, N358
Macrochlamys, N358
 MACROCILIOBRANCHIA, N120
Macrodesma, N250
Macrodon, N256
Macrodonax, N629
Macrodonata, N865
Macrodonella, N257
Macrodontidae, N256
Macrodonus, N256
Macroma, N623
Macropsammus, N631
Macrosolen, N633
Macrosteniana, N862
Macrotoma, N623
Mactesia, N711
Mactra, N595
Mactracea, N16, N63, N84,
 N121, N220, N595
Mactrella, N598, N601
Mactrellona, N68, N601
Mactresia, N711
Mactridae, N16, N76, N119,
 N220, N595, N861
Mactrina, N566
Mactrinae, N220, N595
Mactrinula, N598
Mactroderma, N598
Mactrodesma, N603
Mactroidea, N566
Mactromeris, N603
Mactromya, N511
Mactromyella, N511
Mactromyidae, N220, N511,
 N861
Mactromyopsis, N511
Mactrona, N598
Mactropsis, N610
Mactrotoma, N598
Mactrula, N868
Maetrula, N598
Madagiceramus, N317
Madrela, N332
Madrila, N332
Maconia, N826
Magaleda, N235
Magdala, N846
Magnavicula, N303
Magnonaias, N439
Mailleuxia, N299
Mailleuxia, N256
Maitaia, N289
Maizuria, N306
Malagasitrigonia, N481
Malayomaorica, N376
Mallacea, N326
Malleacea, N326
Malleaceae, N326
malleacées, les, N326
Malleidae, N76, N116, N120,
 N219, N326
Malleolus, N327, N929

- Malleoperma*, N322
Malletia, N13, N20, N62-N63, N99, N165, N167-N168, N233
Malletiidae, N231
Malletiella, N233
Mallettiidae, N214, N218, N231
Mallettinae, N231
Malleus, N38, N42, N71, N327
Malligenus, N327
Malvinasia, N530
Malvufundus, N329
Maminka, N295
Manaia, N267
Mancikellia, N530
Mancusiana, N862
Manoceramus, N317
Mantellina, N389
Mantellum, N385, N389
Mantacula, N314
 mantle, N11, N106
 mantle cavity, N5, N11
 mantle isthmus, N14
 mantle-shell cavity (rudist), N760
Manupecten, N359
Manzanella, N269
Manzanellidae, N218, N269
Maoria, N314
Maoricardium, N585
Maorimactra, N598
Maoritellina, N621
Maorithyas, N510
Maoritrigonia, N483
Marama, N672
Marcia, N684
Margariona, N382
Margarita, N304, N415
Margaritana, N414
Margaritaninae, N414
Margaritanopsis, N415
Margaritaria, N847
Margaritariidae, N222, N847
Margaritifera, N11, N78, N304, N414
Margaritifera, N219, N412, N414
Margaritiphora, N304
Margaron, N415
Margartifera, N304, N414
Margatifera, N414
 marginal carina, N46, N106
 marginal folds, N171
Mariana, N862
Marikellia, N523
 marine bivalves, N91
Mariolana, N862
Marshalliella, N464
Martensnais, N425
Martesia, N143, N702, N704, N711
Martesiella, N711
Martesiinae, N221, N702, N704, N711
Martinea, N635
 MARTINSON, N488
Marwickia, N678
Masudapecten, N370
Matercula, N295
Matheria, N249, N566
Matheronia, N781
Matsumotoa, N264
Mauricia, N273
 McALESTER, N111
 M'COY, N118
Mclearnia, N373
Mecinodon, N657
Mecynodon, N114, N657
Mecynodontidae, N221, N657
Mecynodus, N657
Medeella, N759, N773, N777, N808
 median muscle scar, N38
Medionidus, N455
Mediraon, N577
Meekia, N642
Megacardita, N556
Megacucullaea, N261
Megadesma, N629, N824
Megadesmatidae, N823
Megadesmidae, N115, N221, N823
Megadesmus, N115, N824
Megadiceras, N779
Megadomus, N448
Megagrypha, N865
Megaladon, N742
Megalocardia, N660
Megaloceramus, N315
Megalodon, N50, N78, N114, N116, N120, N742, N765
Megalodontia, N742
Megalodontacea, N211, N221, N742
Megalodontidae, N116, N221, N742, N766
Megalodus, N742
Megalomoidea, N113, N745
Megalomus, N745
Megalomys, N745
Megalonaias, N439
Megaloniidae, N742
Megalonioidea, N439
Megambonia, N397
Megapitaria, N678
Megapraeonia, N566
Megaptera, N287, N455
Megatrigonia, N483
Megaxinus, N498
Megayoldia, N241
Meiocardia, N657
Meisenia, N675
Melasia, N862
Melaxinaea, N267
Meleagrina, N304
Meleagrinnella, N116, N346
Melina, N322
Melinidae, N119, N322
Mellaca, N865
Melliteryx, N520
Melosia, N672
Mendicula, N511
 MENKE, N118
Menuites, N197
Meotolima, N386
Mercenaria, N11, N164, N166, N168, N171, N179, N181, N688
Mercimonia, N685
Meretricina, N673
Meretricinae, N221, N673
Meretricisiana, N862
Meretrissa, N673
Meretrix, N86, N673
Meretrixiana, N862
Meridosinia, N679
Merignacia, N530
Merisca, N617
Merismopteria, N301
Merismopteria, N299
Meroë, N673
Meroena, N673
Merope, N604
Meropesta, N604
Mesafra, N425
Mesanodon, N444
Mesocallista, N678
Mesocibota, N252
 mesoconch, N106
Mesocorbicula, N668
Mesocuna, N560
Mesodesma, N608
Mesodesmatidae, N13, N119, N220, N608
Mesodesmatinae, N220, N608
Mesodesmidae, N608
Mesodiceras, N762, N778
Mesolinga, N498
Mesomiltha, N500
Mesonaia, N450
Mesopeplum, N366
Mesopholas, N711
 mesoplax, N64, N106, N705
Mesopleura, N639
Mesosaccella, N237
Mesotrigonia, N483
Metabola, N606
Metadacna, N592
 metamorphosis, N98
Metapalaeoneilo, N233
 metaplast, N64, N106, N705
Metaptera, N862
Metaxylophaga, N722
Metis, N625
Mewalonus, N745
Mexicardia, N586
Micatula, N377
Micramussium, N351
Microcallista, N677
Microcardium, N589
 MICROCILIOBRANCHIA, N120
Microcirce, N672
Microcondylaea, N424
Microcondylus, N424
Microcucullaea, N254
Microcyprina, N648
Microdon, N565
Microdonella, N565
Microdonta, N408

- Microdontella, N409
 Microdontia, N461
 Microdontidae, N219, N407
 Microfragum, N588
 Microripipes, N499
 Micromactra, N598
 Micromeris, N560
 Micromya, N455
 Micromytilus, N270
 Micronaias, N422
 Micropolia, N523
 Micropteria, N299
 Microstagon, N517
 Microtrigonia, N483
 Microvexillum, N741
 Microyoldia, N239
 Middalaya, N475
 Migonitis, N519
 Migranaia, N441
 Mikrola, N637
 Mila, N296
 Milletiana, N862
 Milligareta, N631
 Milneria, N73, N557
 Milovanovicia, N810
 Miltha, N502
 Miltheoidea, N502
 Miltthinae, N220, N501
 Miltthona, N500
 Mimachlamys, N355
 Mimarcaria, N252
 Mimetostreon, N346
 Mimoceramus, N315
 Minepharus, N580
 Minetrigonia, N483
 Minnivola, N371
 Minormalletia, N233
 Miocardia, N657
 Miocardiella, N657
 Miocardiopsis, N658
 Mioclausinella, N672
 Miodomeridinae, N220, N553
 Miodomeris, N553
 Miodon, N552, N653
 Miodontiscus, N552
 Miodontopsis, N653
 Mioerycina, N523
 Miophoria, N473
 Mioporomya, N852
 Miorangia, N601
 Mirapecten, N366
 Miratacar, N252
 Miridas, N278
 Misidioptera, N391
 Mitriodon, N463
 Mitrocaprina, N773, N789, N795
 Mittraea, N515
 Mixtipecten, N359
 Mixtopecten, N359
 Miyagipecten, N360
 Mizuhopecten, N370
 Mnataia, N256
 mode of life (rudist), N764
 MODELL, N414
 Modiella, N547
 Modiella, N278
 Modioconcha, N407
 Modiodesma, N397
 Modiola, N114, N278
 Modiolarca, N13
 Modiolarca, N276, N644
 Modiolarcadae, N644
 Modiolarcidae, N644
 Modiolaria, N276
 Modiolatus, N278
 modioliform, N106
 Modioliformia, N252
 Modiolina, N547
 Modiolinae, N219, N278
 Modiolini, N278
 Modiolodon, N111, N397
 Modioloides, N109, N868
 Modiolopsidae, N113, N393
 Modiolopsis, N110, N397
 Modiolula, N279
 Modiolus, N23, N33-N34, N79,
 N114, N116, N132, N141,
 N144, N147-N148, N163,
 N170, N182, N278
 Modiolusia, N278
 Modiomorpha, N393
 Modiomorphacea, N116, N120,
 N211, N219, N393, N861
 Modiomorphidae, N111-N113,
 N137, N214, N216, N219,
 N393, N861
 MODIOMORPHOIDA, N214,
 N217, N219, N393
 Modioptera, N865
 Moera, N618
 Moerella, N23, N618
 Mokattamia, N648
 MOLLUSCA, N109, N130
 Molukkana, N832
 Moncetia, N463
 Monocondylaea, N465
 Monelagmus, N454
 Monelasmus, N454
 Monia, N58, N385
 Monica, N385
 moniliform, N106
 Monitilora, N501
 Monnieria, N781
 Monocondyla, N465
 Monocondylaea, N465
 Monocondylea, N465
 Monocondyloia, N465
 Monocondylus, N465
 Monodacna, N593
 Monodonta, N447
 Monodontina, N434
 Monoetia, N463
 MONOMYARIA, N118, N120
 monomyarian, N34, N106
 MONOPLACOPHORA, N109,
 N111
 Monopleura, N751-N752, N757,
 N762, N765, N781
 Monopleura, N235, N787
 Monopleuridae, N221, N754-
 N755, N757, N760-N764,
 N766, N781, N799
 Monoptera, N297
 Monopteriidae, N219, N297
 Monothyra, N708
 Monotidae, N115, N154, N219,
 N373
 Monotinae, N373
 Monotis, N374
 Montacuta, N21, N38, N529
 Montacutidae, N220, N527
 Montacutona, N531
 Montaguia, N529
 Montanaria, N114, N511
 Montrouzieria, N637
 Montrouzieria, N637
 Moquicardium, N591
 Moquiniana, N862
 Moreletiana, N862
 morphology, N134, N175; rudist,
 N751; Thyasira, N185
 MORTON, N208, N213
 MORTON & YONGE, N212-N213
 Mouretia, N809
 mouth, N5, N21
 Mrassiella, N294
 Mucidusiana, N862
 Muelleria, N466
 Mulinarca, N263
 Mulinia, N601
 Mulinoides, N601
 Mulleria, N208
 Mulleria, N466
 Mulletia, N326
 Multidentata, N241
 multivincular, N106
 multivincular ligament, N60
 MUNIER-CHALMAS, N53, N99,
 N751, N755, N908
 Murcia, N686
 Murraia, N470
 MURRAY, N45
 muscles (attachment), N37
 muscle buttress, N764
 muscle insertions (rudists),
 N762-N763
 muscle platforms, N183
 muscle scar, N106; rudist, N764
 muscle supports (rudist), N771
 musculature, N30, N176
 Musculiopsis, N653
 Musculium, N670
 Musculus, N9, N15, N23, N116,
 N276
 Musculus, N613
 Mutela, N96, N463
 Mutelidae, N13, N219, N412,
 N414, N463, N861
 Mutelina, N463
 Mutiella, N513
 Mweruella, N418
 Mya, N4, N6, N11, N14-N15,
 N35, N49, N60, N62-N64,
 N142, N164, N166, N170-
 N171, N173-N174, N195, N691
 MYACEA, N690
 Myacea, N16, N63, N115, N118-

- N119, N210-N211, N221,
N690
Myacidae, N691
Myacites, N842
Myadesma, N610
Myadora, N847
Myadoropsis, N848
Myalina, N63, N111, N114-
 N115, N143, N289
Myalina, N692
Myalinella, N291
Myalinidae, N113-N114, N143,
 N216, N219, N289; marine,
 N289; nonmarine, N291, N404,
 N406, N861
Myalinodonta, N301
Myalinoptera, N861
Myalinopterella, N287
Myapalmula, N278
Myarius, N691
Myatella, N846
Mycetopoda, N465
Mycetopodella, N465
Mycetopus, N465
Myes, N691
Mygallia, N642
Myidae, N73, N195, N221, N691
 MYINA, N221, N690
Myisca, N415
Mylioides, N317
Myllita, N520
Myllitella, N520
Myocardia, N594
Myocaris, N868
Myochama, N13, N72, N847
Myochamidae, N72, N222, N847
Myochlamys, N355
Myoconcha, N547
Myoconchidae, N114
Myoconchidae, N547
Myoconchinae, N220, N547
Myodora, N847
Myoforceps, N278
 MYOIDA, N214, N218, N221,
 N690
Myomactra, N607
Myonera, N854
Myonia, N115, N826
Myoparo, N274
Myopholadidae, N221, N841
Myopholas, N842
myophore, N32, N53, N106,
 N764, N771
myophore plates, N764
Myophorella, N68, N483
Myophoria, N472
Myophoriaemorphis, N582
Myophoricardiidae, N220, N562,
 N580
Myophoricardium, N581
Myophorieae, N472
Myophorigonia, N485
Myophoriidae, N114-N115, N219,
 N472
Myophoriogonia, N485
Myophoriopsis, N582
Myophoriopsis, N582
Myophorocardium, N581
Myopsis, N700
myostracum, N76, N106
Myrene, N655
Myrina, N278
Myrsopsis, N682
Myrsus, N685
Myrtaea, N499
Myrtea, N195, N199, N499
Myrteinae, N220, N499
Myrteopsis, N499
Myrtucina, N504
Mysca, N415
Mysella, N531
Mysia, N689
Mysia, N515
Mysidia, N280
Mysidiella, N280
Mysidiellidae, N219, N280
Mysidioptera, N115, N391
Mytella, N273
Mytilacea, N15, N18, N63, N76,
 N112, N120-N121, N210-
 N212, N214, N218, N271
Mytilarca, N147, N287
Mytilarius, N271
Mytilaster, N273
Mytilicardes, N548
Mytilicardia, N548
Mytilicardita, N548
Mytiliconcha, N280
Mytilidae, N9, N12, N23, N27,
 N34-N35, N37, N114, N116,
 N119, N163, N173, N175,
 N195, N218, N271, N861
Mytilidia, N271
Mytilidonta, N273
mytiliform, N106
Mytiligenus, N271
Mytilimeria, N58, N847
Mytilina, N643
Mytilinae, N218, N271
Mytilisepta, N274
Mytillicés, N270
Mytilocardia, N548
Mytiloceramus, N317
Mytiloconcha, N271, N273, N280
 MYTILOIDA, N76, N214, N217,
 N218, N270
Mytiloida, N317
Mytiloides, N142-N143, N147,
 N317
Mytiloides, N644
Mytilomorpha, N394
Mytilomya, N643
Mytiloperna, N325
Mytiloperna, N325
Mytilops, N280
Mytilopsis, N644
Mytilus, N5, N8-N9, N11, N14-
 N15, N17-N18, N23, N25-
 N26, N28-N29, N34, N37,
 N45, N47, N50, N58, N59-
 N61, N63-N64, N74, N77,
 N91, N99, N100, N114,
 N117, N132, N141, N147-
 N148, N155, N175, N271
Mytolimax, N643
Mytulus, N271
nacreous, N74, N106
nacreous layer, N412
Naiia, N462
 NAIADACEA, N120
Naiadacea, N120, N210, N411
 NAIADIDA, N214
Naiadina, N330
Naiadites, N114, N295, N406
Naiadopsis, N394
Naidea, N455
Najadites, N295, N404, N407
Nakamuranaia, N470
Namnetia, N531
Nanaochlamys, N360
Nannomactra, N598
Nannonaia, N423
Nanonavis, N258
Nanopitar, N675
Nanusiana, N862
Narano, N689
Nargicardium, N591
Nasus, N434
Natella, N592
nates, N45, N106
Nathorstella, N287
Naucitora, N734
Naulia, N678
Nausitora, N734
Nausitorea, N736
Nausitoria, N734
Navaculina, N639
Navea, N716
Navicula, N250
Nayadina, N330
Nayadina, N442
Neacra, N854
Neaeromya, N531
Neaeroporomya, N853
Neamnigenia, N402, N404
Necklania, N245
Neilo, N233
Neilonella, N235
Neithea, N73, N117, N371
Neithea group, N371
Neitheops, N371
Nelltia, N621
Nemetia, N668
Nemoarca, N254
Nemocardium, N44, N589
Nemodon, N150, N258
Nemrodia, N862
Neoactinodonta, N399
Neobakevellia, N306
Neobankia, N734
Neoburmesia, N832
Neocaprina, N789, N795
Neocardia, N270
Neocatillus, N315
Neocorbicula, N666
Neocrassina, N567

- Neocyrena, N668
 Neodavisia, N542
Neodipecten, N360
 Neogaimardia, N644
Neogaimardia, N644
 Neoinoceramus, N320
 Neolepton, N542
 Neoleptonidae, N220, N542
 Neomegalodon, N743
Neomegalodus, N743
 Neomiodon, N117, N653
 Neomiodontidae, N221, N653
Neopisidium, N670
Neopleurodon, N269
 Neoradiolites, N758, N808
 Neoschizodus, N475
 Neosiliqua, N613
 Neosolen, N611
 NEOTAXODONTA, N120
 Neoteredo, N728
 Neotrapezium, N655
 Neotrigonia, N30, N52, N471, N485
Neoxylophaga, N721
 nephridia, N24
 Nephritica, N425
 Nephronaias, N425
 nepioconch, N106
 nepionic, N106
Nerithea, N371
 Neritra, N655
 nervous system, N26
 Nesobornia, N525
Nesonaia, N436
Netara, N672
Netastoma, N720
Netastomella, N720
 Netschajewia, N545
Nettastoma, N720
Nettastomella, N720
 NEUMAYR, N209, N213
 Nevenulora, N498
Nevesta, N244
Newaagia, N380
 NEWELL, N38, N57-N58, N137, N141, N173-N175, N182, N214, N544
 Newellipecten, N333
 Newsomella, N301
 NEWTON, N71
Niäa, N462
Niäca, N462
 Nicania, N568
Nicaniella, N567
 Nicol, N911-N912
 Nidarica, N580
 Niloticiana, N861
Niobe, N475
Nioche, N688
 Nippononaia, N426
Nipponarca, N252
Nipponicorbula, N694
Nipponitrigonia, N485
Nipponolimopsis, N266
Nipponomarcia, N683
Nipponomontacuta, N532
Nipponomysella, N532
Nipponopecten, N360
 Nitia, N426
 Nitidavenus, N678
 Nitidotellina, N618
 Nodipecten, N41, N67, N360
Nodopecten, N360
 nodose, N106
Nodularia, N415
Nodularidia, N415
 Noetia, N150, N261
Noetidae, N261
Noetiella, N261
Noetiidae, N100, N218, N261
 Noetiinae, N218, N261
 Noetiopsis, N261
Noidea, N455
Noramyia, N261
 Nordenskjöldia, N258
 normal, N106
 notation systems (rudist), N756
Nothamusium, N865
 Notirus, N683
 Notobotula, N280
 Notocallista, N678
 Notochione, N688
Notochlamys, N366
Notocorbula, N694
Notodonax, N629
Notolepton, N542
Notolimea, N391
Notomyia, N827, N834
Notomyrtea, N499
Notomytilus, N270
 Notopala, N862
 Notopapena, N862
 Notopaphia, N683
Notoscabrotrigonia, N487
Notospisula, N603
Nototeredo, N723, N737
Nototrigonia, N485
Notovola, N307
Novaculina, N639
Novaculinae, N639
Novaculininae, N220, N639
Novathaca, N688
Nubilusiana, N862
Nucinella, N269
Nucinellidae, N269
Nucula, N6, N19, N21-N23, N27, N45, N50, N63, N84, N99, N141, N164-N167, N171, N195, N197, N199-N200, N230
 NUCULACEA, N227
Nuculacea, N9-N10, N16-N18, N20, N38, N63, N76, N98, N111-N113, N118-N120, N173, N210-N211, N218, N229
Nuculana, N13, N24, N63, N84, N99, N167, N195, N199, N235
Nuculanacea, N63, N118, N211, N218, N231
Nuculanella, N231
Nuculanidae, N120, N195, N199, N214, N218, N231, N235
Nucularia, N233
Nuculavus, N231
Nuculidae, N12, N70, N76, N100, N120, N173, N195, N199, N214, N218, N230
Nuculina, N269
Nuculinidae, N269
Nuculites, N112, N233
Nuculocardia, N274
Nuculochlamys, N239
 NUCULOIDA, N214, N217-N218, N227
Nuculoidea, N231
Nuculoma, N231
Nuculopsis, N231
Nuculopsis, N237
Nucundata, N239
Nucunella, N266
Nudiola, N278
Nuggetia, N376
Numella, N517
Nuttallia, N633
Nyassa, N407
Nyassunio, N427
 nymph, N46, N49, N106
Nympha, N646, N673
nymphae, N46
 OBERLING, N76, N209
Oblicaria, N455
Oblimopa, N265
Obliquarca, N252
Obliquaria, N455
Obliquata, N437
oblique, N106
Obliquipecten, N340
obliquity, N106
Obovalis, N434
Obovaria, N455
Obtellina, N618
Occultamussium, N350
 OCKELMANN, N199
Odatelia, N424
 ODHNER, N57, N471, N581
Odhneripisidium, N670
Odoncineta, N850
Odontocineta, N850
Odontoperna, N306
Oedalia, N690
Oedalina, N690
Oene, N621
Offadesma, N49, N850
Offneria, N773, N789, N795
Overieniana, N862
Oistotrigonia, N483
Okribella, N411
Okunominetania, N475
Oleanella, N302
Olegija, N233
Oligodon, N409
 OLIGODONTA, N120
 OLIGOSTIGMATA, N604
Oliveiraia, N643
Omala, N618

- Omalia, N865
 Omnivenus, N675
Omphaloclatrum, N672
Onchophoridae, N690
Oncoceramus, N317
Oncophora, N690
 Onestia, N589
 Onkogrypha, N865
Onodon, N442
Ontaria, N245
 ontogeny, N91
Onychia, N302
Onychocardium, N295
Oopecten, N358
Operculella, N383
Opertochasma, N715
Opimarcia, N683
Opinae, N220, N571
Opis, N44, N116-N117, N571
Opisastarte, N573
Opisenia, N865
Opisocardium, N588
Opisoma, N572
 opisthoclinal, N51, N106
Opisthocoele, N245
 opisthodontic, N60, N106
 opisthogyrate, N45, N106
Opistholobata, N287
Opisthoptera, N287
Opisthotrigonia, N485
Oppenheimia, N369
Oppenheimiopecten, N369
Opsiceramus, N321
Opularca, N252
Oracardia, N247
 oral groove, N21
Oraphocardium, N592
 orbicular, N106
Orbicularia, N633
Orbiculus, N680
Orbignia, N801
 ORBIGNY, D', N208, N785
Orbignya, N801
Orbipecten, N334
Orbis, N585
Oreada, N567
Oretia, N382
 orientation, N78; rudist, N751
 origin (rudists), N765
Oriocrassatella, N576, N909, N911-N912
Oriopleura, N785
Orixa, N636
 ornament, N142
Ornithopecten, N339
 oro-anal axis, N79
Orobitella, N532
Oronthea, N523
Orophoceramus, N315
Orpheoceramus, N317
Orthocardium, N586
Orthoceramus, N315
Orthoceratites, N801
 orthoclinal, N51, N106
 ORTHOCONCHAE, N120
 ORTHOCONQUES, N120-N121
- Orthodesma*, N397
 orthodont, N106
Orthodontiscus, N400
 orthogyrate, N45, N106
Orthomyalina, N289
Orthonaiadites, N295
Orthonota, N819
Orthonotella, N397
Orthonotidae, N221, N818
Orthonymus, N437
Orthoptychus, N772, N789, N795
Orthotrigonia, N488
Orthoyoldia, N241
Ortmanniana, N450
Ortonella, N249
Ortygia, N686
Oryctomya, N655
 oscule, N106, N760
Osculigera, N813
 osphradia, N27
Ospirasolen, N613
Ospriasolen, N612
 ossiculum, N49, N106
Osteodesma, N846, N850
Osteodesmacea, N850
Osteodesmidae, N850
Osteomya, N832
Osteophorus, N865
 ostia, N18
Ostomya, N847
Ostracites, N380
Ostracum, N74, N106
Ostrea, N11, N29, N58, N60, N63-N64, N78, N91, N93-N94, N98, N141, N154, N171, N174-N175, N177, N197, N200, N208, N386, N389
Ostreacea, N16, N18, N36, N63, N98, N118, N120, N210-N212, N215, N219
Ostreavicula, N865
Ostreidae, N8, N12, N16, N27, N35-N38, N72-N73, N76, N78, N98, N115-N117, N142, N154, N156, N171, N173-N174, N183, N212, N219
Ostreina, N219
Ostreinae, N219
Ostreinella, N331
Ostrenomia, N378
Ostreopecten, N386
Otapiria, N339
 otocysts, N27
Otysma, N283
Oudardia, N619
 outer calcareous layer, N11
 outer demibranch, N16
 outer lamellae, N18
 outer ligament, N107
Ovacuna, N560
Ovalarca, N264
Ovaleda, N239
 ovate, N107
 override, N107
Ovicardium, N586
Ovuliana, N862
- OWEN, N87-N88, N174, N177, N179, N212
Oxisma, N283
Oxybeloceras, N197, N200
Oxyceramus, N315
Oxydacna, N592
Oxynaia, N426
Oxyperas, N603
Oxypteria, N339
Oxysma, N283
Oxytoma, N68-N69, N73, N116-N117, N144, N344
Oxytomidae, N76, N219, N344
Oxytominae, N344
Ozomia, N868
- Pachecoa*, N264
Pachites, N378
Pachoya, N280
Pachyanodon, N463
Pachycardia, N468
Pachycardiidae, N219, N401, N467
Pachycardium, N588
Pachydacna, N592
Pachydesma, N674
Pachydomella, N863
Pachydomidae, N823
Pachydomus, N824
Pachydon, N697
Pachyodontinae, N221, N697
Pachyerisma, N745
Pachykellya, N542
Pachymegalodon, N745
Pachymegalodus, N745
Pachymya, N116, N834
Pachymyonia, N827
Pachymytilus, N291
Pachynaias, N455
Pachynus, N801
Pachyodon, N461, N578
 pachyodont, N52, N107
 PACHYDONTA, N120, N766
 PACHYDONTIDA, N214
Pachyopsis, N571
Pachyperna, N326
Pachypteria, N114, N342
Pachyrisma, N50, N116, N745
Pachyrismella, N745
Pachyta, N378
Pachytes, N378
Pachythaerus, N574
Pachytos, N378
Pachytraga, N761, N785
Pachytus, N378
Pachytypus, N568
Pacificor, N555
Pacipecten, N360
Pacitrigonia, N485
Packardella, N487
Pacycyardia, N468
Palaeonodonta, N115, N404, N406-N407
Palaearca, N248
 PALAEOBRANCHIA, N119-N120

- Palaeocardia, N287
 Palaeocardita, N116, N120, N554
 Palaeocarditinae, N220, N554
 PALAEOCONCHA, N112, N118, N214
 Palaeoconcha, N210, N229
 PALAEOCONCHAE, N120
 Palaeocorbis, N514
 Palaeocorbula, N820
 Palaeocosmomya, N114, N831, N834
 Palaeocucullaea, N256
 PALAEOHETERODONTA, N62-N63, N120, N211, N213-N215, N217, N219, N393
 PALAEOOLAMELLIBRANCHIA, N120
 Palaeolima, N114, N391
 Palaeolucina, N512
 Palaeomoera, N621
 Palaeomutela, N51, N115, N404, N409
 Palaeomutelidae, N219, N402, N404, N409
 Palaeomya, N640
 Palaeomya, N822
 Palaeoneilo, N111, N233, N409
 Palaeonucula, N231
 Palaeopecten, N113, N334
 Palaeopharus, N470
 Palaeopinna, N301
 Palaeopleiodon, N409
 Palaeopteria, N301
 Palaeosolen, N819
 PALAEOOTAXODONTA, N63, N99, N120, N211, N213-N214, N217-N218, N227
 PALAEOOTAXODONTIDA, N214
 Palanatina, N393
 Paleodora, N868
 Paleomarcia, N681
 Palestinarca, N261
 Paleyoldia, N237
 Palindonaia, N427
 pallets, N8, N65, N107, N706
 pallial, N107
 pallial canals (rudists), N762
 pallial line, N11, N35, N107, N176, N179
 pallial muscles, N11, N35
 pallial nerve, N26
 pallial region, N107
 pallial retractor muscles, N36, N107
 pallial sinus, N13, N36, N107
 Palliolium, N354
 Palliorum, N354
 Pallium, N365-N366
 Palmoxytoma, N345
 palp pouch, N19
 palp proboscides, N18
 Palus, N790, N817
 Pammegaliana, N862
 Panacca, N827, N834
 Panacea, N834
 Panacoma, N623
 Panamicorbula, N696
 Panchione, N686
 Pandora, N21, N91, N366, N843
 Pandoracea, N63, N211, N222, N843
 Pandoracia, N843
 Pandorella, N844
 Pandoridae, N222, N843
 Pandorina, N846
 Panenka, N113
 Panenka, N244
 Panis, N310
 Pannonocardium, N591
 Panodea, N700
 Panomya, N700
 Panopaea, N700
 Panopaeacea, N118
 Panope, N166, N170
 Panope, N700
 Panopea, N13, N267, N700
 Pantaita, N244
 Panticapaea, N593
 PANTODONTA, N120
 PANTODONTIDA, N214
 Paphia, N66, N71, N79, N685
 Paphia, N608
 Paphidae, N119
 Paphies, N610
 Paphiinae, N608
 Paphirus, N686
 Paphonotia, N683
 Papiilicardium, N585
 Papyridea, N588
 Papyrina, N598
 Papyrocardium, N593
 Paraborniola, N525
 Parabournonia, N810
 Paracaprinula, N789, N795
 Paracardium, N113
 Paracardium, N244
 Paraceramus, N315
 Paracorbicula, N666
 Paracyclas, N113-N114, N512
 Paradacna, N593
 Paradacninae, N220, N593
 Paradiceras, N778
 Paradiione, N677
 Paradonax, N628
 Paradoxia, N865
 Paraesa, N685
 PARAFILIBRANCHIA, N120
 Paraglans, N557
 Paragrammatodon, N259
 Parainoceramus, N320
 Paralepida, N533
 Parallelodon, N111-N112, N114, N116, N137, N150, N256
 Parallelodontacea, N120
 Parallelodontidae, N120, N216, N218, N256, N861
 Parallelodontinae, N218, N256
 Parallelodus, N398
 Parallelopipedum, N254
 Paralucinella, N506
 Paramegalodus, N746
 Paramodiola, N398
 Paramonopleura, N783, N868
 Paramusium, N350
 Paramussium, N350
 Paramya, N692
 Paramytilarca, N287
 Paranodonta, N427
 Paranoetia, N262
 Paranomia, N148, N383
 Parapecten, N371
 Paraphola, N715
 Parapholas, N706, N715
 Paraphtonia, N398
 Paraplacuna, N383
 Paraprothyris, N822
 Parapscheronia, N593
 Paraptyx, N244
 Pararca, N244
 Parasauvagesia, N760, N808
 Parastarte, N681
 Parastroma, N802
 Paratancredia, N640
 Paratapes, N685
 Parathyasira, N510
 Paraucellina, N376
 Paraviculidae, N310
 Pardosinia, N679
 Parembola, N682
 Pareptera, N454
 Parilimya, N834
 Parimalleus, N329
 Parinomya, N383
 Parisiella, N567
 parivincular, N60, N107
 PARKINSON, N326
 Parmicorbula, N696
 Parmulina, N672
 Parmulophora, N672
 PARONA, N749, N808
 Paronella, N758, N808
 Paronomia, N383
 Paropsis, N301
 Parreysia, N427
 Parreysiana, N862
 Parthenope, N533
 Particoma, N712
 Particondyla, N560
 Parunio, N439
 Parvamussium, N350
 Parvati, N565
 Parviamussium, N350
 Parvicardium, N585
 Parvicorbis, N513
 Parvidacna, N593
 Parvikellia, N522
 Parvilucina, N498
 Parvimalleus, N329
 Parviperna, N322
 Parvithracia, N852
 Parvivenus, N688
 Parvochlamys, N354
 Pascoella, N585
 Pasiphae, N688
 Paslucina, N496
 Passya, N536
 Patagocardia, N865
 Pater, N244

- Patinopecten, N370
 Patinopecten subgroup, N370
 Patro, N383
Patrocardia, N297
Patros, N383
Patularia, N463
 Pauliella, N653
Paxydonta, N461
 Paxyodon, N461
 pearls, N78
Pecchiolia, N857
 Pecten, N10-N11, N22-N23, N29, N42, N63-N64, N69, N73, N77, N85-N87, N90, N117, N135, N141, N161-N162, N173-N175, N177, N197, N199, N366
 Pecten group, N366
 Pecten subgroup, N366
Pectenacea, N332
Pectenina, N332, N348
Pectenidae, N348
Pecten, N366
Pectinacea, N9, N15-N16, N18, N36, N46-N47, N63, N69, N72, N76, N81, N98, N113-N114, N116, N119-N120, N211-N212, N217, N219, N332, N861
Pectinarca, N860
Pectinatarca, N254
Pectinella, N354
Pectinia, N348
Pectinida, N348
Pectinidae, N9, N12, N16, N22, N29, N46, N63, N69-N70, N72-N73, N76, N101, N115, N117, N119, N131, N138, N142-N143, N151, N153-N154, N159-N160, N162, N170-N171, N173, N197, N199, N212, N219, N348
Pectinides, N348
Pectinigenus, N366
Pectinites, N366
Pectinium, N366
Pectinucula, N231
Pectinula, N355
Pectunculidae, N267
Pectunculina, N265
Pectunculopsis, N267
Pectunculus, N680
Pectunculus, N267
Pectunpennaceus, N267
 pedal artery, N24
 pedal elevator muscle, N38, N107
 pedal ganglia, N26
 pedal gape, N47, N107
 pedal levator muscle, N107
 pedal musculature, N181
 pedal protractor muscle, N38, N107
 pedal retractor muscle, N107
 pedalbyssal musculature, N181
Pedalion, N322
Pedinus, N364
 pediveliger stage, N95
 Pedum, N47, N364
Pegassella, N299
Pegias, N448
Pegias, N448
Pegophysema, N504
Pelecypora, N678
 PELECYPODA, N3-N4, N213, N227
Pelopia, N850
 PELSENEER, N17-N18, N143, N210, N212
Pelvis, N595
Penchinatiana, N862
Pendaloma, N850
Penecilli, N858
Penecillus, N858
Penicellus, N858
Penicilla, N716
Penicillus, N67, N102, N858
Penita, N453
Penitella, N716
Pennaria, N283
Pentadina, N300
Pentagrammysia, N834
Peplum, N355
Pera, N366, N669
Perampliata, N501
Peraptera, N299
Pergamidia, N314
Pergamidiidae, N219, N313
 pericardial artery, N24
 pericardial cavity, N22, N24
Pericylindrica, N424
Periglypta, N672
 periostracum, N11, N73, N107, N412
Periploma, N49, N849
Periplomatidae, N222, N849
Periplomidae, N849
Periplomya, N845
Perissodon, N601
Perlamer, N304
Permophoridae, N114, N216, N220, N543
Permophorinae, N220, N543
Permophorus, N57, N543
Perna, N77, N273
Perna, N322
Pernadae, N322
Pernaria, N322
Pernaridia, N322
Perniana, N862
Pernidae, N322
Pernigenus, N322
Pernomytilus, N271
Pernomytilus, N322
Pernopecten, N70, N114, N347
Peronaea, N90, N619
Peronaecoderma, N619, N628
Peronea, N802
Peronidia, N619
 PERRIER, N119
Perrierina, N538
Perrierinidae, N537
Perrisonota, N235
Persikima, N594
Peruarca, N268
Perucardia, N586
Petalocardia, N648
Petalodontia, N757, N783
Pethopecten, N360
Petkovicia, N813
Petrasma, N243
Petricola, N13, N63, N689
Petricola, N531
Petricolaria, N689
Petricolidae, N221, N688
Petricolirus, N689
Petroderma, N679
Pettersia, N268
Pexocodakia, N494
Phacosides, N164, N492
Phacosoma, N680
Phaenodesmia, N233
Phaenomya, N698
Phaladomya, N827
Pharaonella, N619
Pharaonia, N418
Pharella, N612
Pharinae, N637
Pharodina, N668
Pharomytilus, N278
Pharus, N32, N639
Phascolicama, N644
Phaseolicama, N644
Phaseolus, N238
Phaxas, N612
Phelopteria, N310
Phenacocyclas, N504
Phenacomya, N714
Phestia, N238
Phialopecten, N360
Philine, N195
Philinidae, N195
Philipiella, N380
 PHILIPPI, N208, N348
Philippia, N367
Philippiella, N269, N380
Philippina, N847
Philis, N509
Philobrya, N269
Philobryidae, N218, N269
Philobryinae, N269
Phloas, N707
Phlogocardia, N586
Phlyctaenachlamys, N536
Phlycticoncha, N847
Phlyctiderma, N517
Phlyctiderma, N847
 PHOLADACEA, N702
Pholadacea, N16, N32, N58, N63, N118-N119, N211, N221, N702
Pholadae, N706
Pholadaria, N702, N706
Pholadariae, N706
Pholadarius, N707
Pholadeae, N706
Pholadella, N820
Pholadiadae, N706
Pholadidae, N13, N32, N36-N38,

- N62, N65, N165, N221, N702, N706
 Pholadidea, N716
 Pholadidae, N716
 Pholadidoidea, N716
 Pholadigenus, N709
 PHOLADINA, N221, N702
 Pholadina, N707
 Pholadinae, N221, N707, N711
 Pholadites, N707
 Pholadoidae, N706
 Pholadomia, N827
 Pholadomorpha, N398
 Pholadomya, N43, N116-N117, N827
 Pholadomyacea, N211, N221, N818
 Pholadomyadae, N827
 Pholadomya, N827
 Pholadomye, N827
 Pholadomyidae, N13, N76, N114, N116, N118, N221, N827
 Pholadomyocardia, N841
 PHOLADOMYOIDA, N214, N218, N221, N818
 Pholadomya, N827
 Pholadopsis, N720
 Pholalites, N707
 Pholalithes, N707
 Pholameria, N716
 Pholas, N7-N8, N65, N169, N173, N707
 Pholas, N707, N709
 Pholdomya, N827
 Pholeobia, N700
 Pholidae, N706
 Pholidaea, N716
 Pholidea, N716
 Pholididae, N716
 Pholodomya, N827
 Phragmorisma, N852
 Phreagena, N664
 Phrynelima, N265
 Phthonia, N280
 Phyllicardium, N592
 Phyllocardium, N592
 Phylloda, N619
 Phyllodella, N619
 Phyllodina, N619
 Phyllonaias, N450
 Phylloteredo, N737
 phylogeny (Thyasira), N200
 Phymesoda, N670
 Physetomya, N394
 Physocardia, N660
 Physoida, N696
 Physunio, N428
 Picardiana, N862
 Pichleria, N266
 PICOT DE LAPEIROUSE, N749
 Pilea, N453
 Pileatona, N532
 Pileochama, N817
 pillar, N107, N760
 Pillucina, N499
 Pilsbryconcha, N446
 Pina, N283
 Pinctada, N8, N14, N33, N38, N42, N71, N73-N74, N78, N132, N138, N143-N149, N151-N153, N178, N304
 Pinctadina, N304
 Pingoteredo, N726
 Pinguimacoma, N623
 Pinguitellina, N619
 Pinguiteredo, N726
 Pinna, N8, N10, N33, N42, N58, N60, N63, N71, N77, N78, N114, N132, N141, N179, N283
 Pinnacea, N15, N18, N35, N63, N76, N121, N211-N212, N219, N281
 Pinnarius, N283
 Pinnidae, N12, N37, N76, N114, N162-N163, N210, N212, N219, N281
 Pinnigena, N285
 Pinnigenus, N283
 Pinnites, N283
 Pinnocaris, N868
 Pinnogena, N285
 Pinnopsis, N295
 Pinnula, N283
 Pintadina, N304
 Pinua, N283
 Pinuigena, N285
 Pinzonella, N115, N748
 Pinzonellopsis, N748
 PIRONA, N808
 Pironaea, N759, N771-N772, N776, N802
 Pironaia, N802
 Pironea, N802
 Pisaniana, N862
 Piscimaliana, N862
 Pisiadiadae, N669
 Pisiidiidae, N96, N221, N669
 Pisidium, N12, N669
 Pisima, N861
 PISOSTRIGILLA, N622
 Pisotrigonia, N487
 Pistris, N619
 Pisum, N669
 Pitar, N168, N675
 Pitarella, N675
 Pitarenus, N675
 Pitaria, N675
 Pitarina, N675
 Pitarinae, N221, N675
 PITTENDRIGH, N138
 Pixis, N354
 Placamen, N688
 Placenta, N383
 Placenticeras, N197
 Placopecten, N160-N161, N360
 Placuanomia, N385
 Placumonya, N385
 Placuna, N15, N383
 Placunanomia, N385
 Placunanomya, N385
 Placunema, N384
 Placunidae, N120, N383
 Placunopsis, N380
 Placunopsis, N380
 Plagia, N306
 Plagiarca, N252
 Plagiastoma, N391
 Plagiocardium, N585
 Plagioctenium, N356
 Plagiodacna, N592
 Plagiodacnopsis, N593
 Plagiodon, N464
 Plagiola, N455
 Plagiolopsis, N455
 Plagiopecten, N357
 Plagiopectenium, N357
 Plagioplecten, N758, N761, N764, N773, N789, N795
 Plagiostoma, N47, N71, N391
 Plagiostomata, N391
 Plagiostomatites, N391
 Planikellia, N527
 Planimodiola, N276
 Planitivela, N675
 Planktomya, N865
 planktotropic development, N91
 Planocaprina, N758, N772-N773, N789, N797
 Plastomiltha, N505
 Platinopecten, N370
 Platipecten, N355
 Platiris, N464
 Platomyia, N527
 Platopsis, N650
 Platopsis, N650
 Platteniciana, N862
 Platycardia, N660
 Platyceramus, N315
 Platydonax, N628
 Platymermis, N504
 Platymya, N845
 Platymyoidea, N845
 Platynaias, N448
 Platyodon, N7, N169, N692
 Platyrrhynchoidiana, N862
 Platyris, N464
 Plebidonax, N628
 Plectodon, N854
 Plectomerus, N437
 Plectomya, N845
 Plectosolen, N611
 Pleiaceramus, N315
 Pleiodon, N38, N51, N465
 Pleiorytis, N689
 Pleodon, N465
 Plesiastarte, N681
 Plesielliptio, N431
 Plesiocyprina, N648
 Plesiocyprinella, N115, N512
 Plesiodiceras, N779
 Plesiodiceratidae, N779
 Plesiodiceratinae, N221, N754, N766, N779
 Plesiopecten, N364
 Plesiopinna, N283
 Plethobasus, N428
 Pletholophus, N446

- Plethomytilus*, N287
 Pleureterites, N868
 Pleurikodonta, N825
 Pleurobema, N428
 Pleuroconcha, N655
PLEUROCONCHAE, N120
 Pleurodapis, N545
 Pleurodesma, N698
 Pleurodesmatidae, N221, N698
Pleurodon, N269
PLEURODONTA, N120
Pleurodonta, N244
 Pleurogrammatodon, N258
Pleurolocina, N496
Pleuromeris, N552
Pleuromya, N116-N117, N842
Pleuromyacidae, N842
Pleuromyidae, N221, N842
Pleuromysidia, N374
Pleuronaia, N437
Pleuronectes, N349
Pleuronectia, N349-N350
Pleuronectites, N71, N339
Pleuronovecten, N350
Pleurophopsis, N664
Pleurophorella, N546
Pleurophoridae, N543
Pleurophorina, N547
Pleurophoropsis, N664
Pleurophorus, N543
Pleurorhynchus, N859
Pleurotrigonia, N478
plica, N107
Plicacesta, N386
plicate canals, N26
Plicatostylidae, N866
Plicatostylus, N866
Plicatounio, N429
Plicatula, N35, N53, N70, N71, N101, N154, N348, N377
Plicatulidae, N12, N16, N34-N35, N37, N115, N154, N219, N377
Plicatuligenus, N377
Plicatulinae, N377
Plicatulopecten, N378
Plicomya, N845
Pliconaias, N439
Pliocardia, N664
Pliodon, N465
PLIODONTA, N120
Plionema, N551
Plumulella, N734
Plurigenis, N672
Plycatula, N377
Pododesmus, N148, N385
Podopsis, N378
 ПОЖЕТА, N181
Polarthus, N741
Polas, N707
Polia, N639
Policordia, N857
Folidevcia, N239
Polittapes, N685
Politoleda, N236
Pollex, N655
Pollicidae, N221, N655
- Polorthus, N741
Polyconilites, N785
Polyconites, N758, N771, N773, N785
Polymesoda, N668
Polymetis, N625
Polynema, N252
Polynemamussium, N351
Polydonta, N860
POLYPLACOPHORA, N79
Polyptychus, N795
Pomarangina, N748
Pompeckjina, N289
Pompholigina, N508
Ponderosiana, N862
Pontalmyra, N592
Popenaias, N426
 POPENOE & FINDLAY, N57
porcelaneous, N74, N107
Poroleda, N237
Poromya, N9, N852
Poromyacea, N10, N16, N18, N210-N211, N222, N852
Poromyidae, N13, N222, N852
Poronia, N520, N525
Porterius, N259
Portlandella, N239
Portlandia, N195, N199, N239
Posidonia, N9, N113, N342, N408
Posidoniella, N291
Posidoniella, N343
Posidoniidae, N219, N342
Posidoniinae, N342
Posidonomya, N408
Posidonotis, N344
Posodonomya, N342
posterior, N42, N107
posterior adductor, N31
posterior area, N107
posterior lateral, N49
posterior lateral tooth, N107
posterior pallial arteries, N24
posterior pedal retractor, N37
posterior retractor of the visceral mass, N38
posterior ridge, N46, N107
posterior slope, N46, N107
posterodorsal margin, N42, N107
Postligata, N269
postneanic changes, N100
Potamida, N441
Potamila, N454
Potamilus, N454
Potamocorbula, N696
Potamomya, N697
Potamophila, N629
Potiarca, N256
Potidoma, N527
Potodoma, N415
Potomida, N441
Praearca, N235
Praeastartacea, N210
Praebarretia, N771-N772, N776, N802
Praecaprina, N789, N797
- Praecaprotina*, N785
Praecardiacea, N211, N218, N243
Praecardiidae, N113, N218, N243
Praecardiinae, N218, N244
PRAECARDIOIDA, N214, N217-N218, N243
Praecardiomya, N867
Praecardium, N113, N244
Praeconia, N567
Praectenodonta, N228
Praegonia, N485
PRAEHETERODONTA, N120
Praelapeirousia, N758, N769, N813
Praeleda, N229
Praelima, N244
Praelucina, N113
Praelucina, N244
Praemytilus, N280
Praenomalodonta, N289
Praenucula, N229
Praenuculidae, N218, N229
Praeostrea, N247
Praeradiolites, N750, N758-N759, N764, N769, N771, N808
Praerangia, N601
Praesacella, N237
Praeundulomya, N114, N835
Pragmopholas, N707
Prantliella, N221
Pratulium, N589
Praxis, N644
préastartidés, N120
Praevicula, N868
PRÉHÉTERODONTES, N120
Preroderma, N867
Pressidens, N430
Pressodonta, N448
primary ligament, N58, N107
primary riblets, N69, N107
Primella, N670
Prionia, N241
PRIONODESMACEA, N119-N120, N210, N214
prionodont, N107
PRIONODONTA, N120
Prionopleura, N592
Priscomactra, N601
Prisconaiia, N475
prismatic, N107
prismatic calcareous layer, N412
prismatic structure, N74
Prismatro, N383
Prisodon, N457
Prisodontopsis, N430
Pristes, N533
Pristigлома, N239
Pristipagia, N619
Pristiphora, N533
Pristis, N533, N619
Privigna, N672
Proagorina, N857
Procardia, N829
Procopievskia, N294, N860
Procos, N633
Procrassatella, N575

- Procyprina*, N648
Prodiceras, N748
Prodiplodon, N461
prodissoconch, N42, N96, N107
prodissoconch I, N94, N107
prodissoconch II, N94, N107
Prodreissensia, N644
Productae, N867
Productea, N867
Profischeria, N629
Proheligmus, N329
Prohinnites, N72, N102, N364
Prohyriopsis, N430
Prolaria, N838
Prolasmidonta, N448
Prolobella, N398
Prolucina, N113-N114
ProLucina, N504
Promacrus, N821
Promantellum, N389
Promyalina, N291
Promyophorella, N485
Promvtilus, N280
Pronoe, N648
Pronoella, N117, N648
Pronocula, N231
Prooxytoma, N346
Proparresya, N441
Propeamusium, N350
Propeamusium, N9, N160-N161, N195, N199, N348, N350
Propeannessium, N350
Propecuna, N560
Propehydella, N460
Propeleda, N237
Properycina, N527
Propetilus, N276
Prophetilora, N501
Prophyllicardium, N594
Propreamusium, N350
Proptera, N455
Prorhynchus, N256
Prorokia, N568
Protrigonia, N485
Prosochasma, N295
Prosochiasta, N592
prosocline, N51, N107
Prosocoelogeton, N572
Prosocoelus, N114, N120, N572
prosocoelous, N107
Prosodacna, N592
Prosodacnomya, N592
prosodetic, N107
prosogyrate, N45, N107
Prosogyrotrigonia, N485
Prosoleptus, N233
prosopon, N107
Prospodylidae, N341
Prospodylus, N342
Protamusium, N346
Protamusium, N346
Protapes, N685
Protarca, N269
Protelliptio, N431
Proteoceramus, N315
Proteopecten, N357
Proteredo, N741
Prothyasira, N508
Prothyris, N822
Prothyris, N822
PROTOBRANCHIA, N17, N18, N118-N120, N210, N212, N214
PROTOBRANCHIËS, N119-N120
Protocallithaca, N688
Protocardia, N46, N83, N116, N588
Protocardiinae, N220, N588
Protocyprina, N653
Protodicerax, N748, N765
Protodonax, N629
Protohydridella, N461
Protomiodon, N653
Protomusium, N346
Protomya, N822
Protonoetia, N262
Protonucula, N234
Protopalliolum, N861
Protopecten, N113, N868
Protopis, N280
protoplax, N64, N107, N706
Protopsinae, N605
Protoschizodon, N475
Protoschizodus, N475
protostracum, N94, N107
protostyle, N22
Prothaca, N688
Protunio, N470, N488
Protoxylophaga, N721
Protrete, N546
Protunio, N431
Proveniella, N648
provinculum, N98, N107
Proxichione, N672
Pruvostiella, N567
Prysodon, N457
Psammacoma, N623
Psammobella, N26, N631
Psammobia, N631
Psammobiadae, N629
Psammobiidae, N13, N38, N57, N119-N120, N629
Psammobiinae, N220, N630
Psammocola, N631, N633
Psammococoncha, N867
Psammodonax, N631
Psammoica, N631
Psammophila, N604
Psammosolen, N638
Psammosphaerica, N633
Psammotaea, N630, N634
Psammotaena, N631
Psammotella, N633
Psammotella, N634
Psammotellina, N634
Psammothalia, N624
Psammotreta, N626
Psathura, N681
Psephidia, N685
Psephis, N685
Pseuanodonta, N446
Pseudacesta, N391
Pseudamiantis, N679
Pseudamusium, N354
Pseudamussium, N9, N27, N354
Pseudamussium, N354
Pseudanodonta, N446
Pseudaphrodina, N679
Pseudarca, N234
Pseudarcopagia, N619
Pseudasaphis, N668
Pseudastarte, N580
Pseudaucella, N377
Pseudavicula, N346
Pseudavicula, N430
Pseudaviculopecten, N144, N334
Pseudaxinea, N267
Pseudaxinus, N471
Pseudedmondia, N818
Pseudentolium, N348-N349
Pseuderiphyla, N577
Pseudeuopera, N669
Pseudisocardia, N661
Pseudoammussium, N354
Pseudoanodonta, N446
Pseudobakewellia, N299
Pseudobaphia, N442
Pseudobarrettia, N803
Pseudobrechites, N859
Pseudocaelatura, N862
Pseudocardia, N558
Pseudocardia, N553, N557
pseudocardinal, N51, N107
Pseudocardinia, N410
Pseudocardiniidae, N117, N219, N410
Pseudocardita, N593
Pseudocarditinae, N220, N593
Pseudocardium, N601
Pseudocatillus, N592
Pseudochama, N57, N518, N768
Pseudocorbicula, N670
Pseudocorbula, N582
pseudoctenodont, N51, N107
PSEUDOCTENODONTA, N120
Pseudoctenostreon, N386
Pseudocucullaea, N261
Pseudocuspidaria, N853
Pseudocyrena, N668
Pseudocyrtodonta, N233
Pseudodicerax, N778
Pseudodidymotis, N344
Pseudodon, N412, N431
Pseudodontideus, N446
Pseudodontoideus, N446
Pseudodontopsis, N425
Pseudodus, N431
Pseudoeuchasma, N868
pseudofeces, N19
Pseudogervilleia, N306
Pseudoglomus, N234
Pseudoglyciana, N862
Pseudogrammatodon, N259
Pseudoheligmus, N332
Pseudohyria, N461
Pseudoirus, N689
Pseudokellya, N525
PSEUDOLAMELLIBRANCHIA, N120, N212

- PSEUDOLAMELLIBRANCHIÉS*,
 N120
pseudolateral, N51, N107
Pseudoleodopsis, N229
Pseudoleguminaia, N425
Pseudolepton, N525
Pseudolimea, N391
Pseudolucina, N391
Pseudolucinisca, N494
Pseudomacrodon, N258
Pseudomalletia, N233
Pseudometis, N627
Pseudomiltha, N504
Pseudomodiolus, N861
Pseudomonotidae, N144, N219,
 N341
Pseudomonotinae, N341
Pseudomonotis, N72, N114, N116,
 N154, N341
Pseudomülleria, N467
Pseudomulleria, N467
Pseudomuseum, N354
Pseudomutela, N463
Pseudomya, N867
Pseudomyalina, N291
Pseudomyoconcha, N547
Pseudomyophorella, N485
Pseudomytiloides, N45, N320
Pseudoneaera, N854
Pseudoon, N455
Pseudopachymytilus, N291
Pseudopalliorum, N350
Pseudopecten, N372
Pseudopecten group, N371
Pseudopermophorus, N546
pseudopillar, N107, N760
Pseudopis, N554
Pseudoplacuna, N385
Pseudoplacunopsis, N378
Pseudopleurophorus, N656
Pseudopolyconites, N758, N804,
 N808
Pseudoportlandia, N239
Pseudoprosodacna, N592
Pseudoptera, N310
Pseudopteria, N310
Pseudopythina, N525
Pseudosanguinolites, N301, N548
Pseudosaxicava, N700
Pseudospatha, N466
pseudotaxodont, N51, N107
Pseudotechnophorus, N868
Pseudotoucasia, N781
Pseudotrapezium, N117, N648
Pseudovaccinites, N803
Pseudovola, N371
Pseudoxyperas, N603
Pseudunio, N415
Psiloconcha, N394
Psilodon, N592
Psilomya, N853
Psilonychia, N289
Psilopoderma, N518
Psilopus, N518
Psiloteredo, N728
Psilotrigrionia, N486
Psilotus, N518
Psilunio, N441
Psoronaia, N442
Psorula, N437
Psuedodicyathifer, N726
Ptenoides, N386
Pteradacna, N592
Pteranodon, N442
Pterelectroma, N304
Pteria, N14, N37-N38, N42,
 N60-N63, N71, N100, N132,
 N137-N138, N141, N143-
 N147, N149, N151, N155,
 N175, N182, N197, N200,
 N302
Pteriacea, N15, N18, N34-N35,
 N46-N47, N63, N72, N76,
 N81, N84, N98, N113, N115,
 N121, N138, N200, N210-
 N211, N219, N297
Pteriadae, N302
Pteriidae, N12, N27, N35, N37,
 N70, N76, N143-N144, N151,
 N212, N219, N297, N302,
 N861
PTERIINA, N219, N285
Pterinea, N112-N113, N137-
 N138, N144, N149, N299
Pterineidae, N113, N144, N216,
 N219, N298, N861
Pterinella, N304
Pterinites, N283
Pterinopecten, N63, N113, N332
Pterinopectinella, N143, N334
Pterinopectinidae, N143, N219,
 N332
Pterinoperna, N304
Pterinopsis, N299
PTERIOIDA, N76, N115, N214,
 N217, N219, N285
PTERIOMORPHIA, N15, N63,
 N99, N118, N120-N121, N211,
 N213-N215, N217-N218, N248
Pterocardia, N116, N749
Pterocardium, N749, N765
Pterochaenia, N297
PTEROCONCHIDA, N121, N214
Pterolucina, N504
Pteromeris, N553
Pteromeris, N553, N560
Pteromya, N841
Pteromyrtea, N498
Pteronitella, N301
Pteronites, N283
Pteronitina, N301
Pteroperna, N45, N70, N304
Pteropsella, N606
Pteropsellinae, N220, N605
Pteropsis, N606
Pterosygna, N448
Pterosyna, N448
Pterotrigrionia, N117, N487
Ptilotrigrionia, N487
Ptychina, N508
Ptychobranchus, N456
Ptychocardia, N539
Ptychoderma, N455
Ptychodes, N867
Ptychodesma, N250
Ptychomya, N655
Ptychomyidae, N221, N655
Ptychopteria, N302
Ptychorhynchoideus, N434
Ptychorhynchus, N434
Ptychostolis, N231
Puella, N244
Pugilarca, N252
Pullastra, N685
Pulsella, N332
Pulsidis, N694
Pulvinites, N326
Pulvinitidae, N219, N326
Pulvinus, N623
puncta, N69
punctate, N107
punctiform scars, N38
punctum, N107
Punigapia, N619
Punipagia, N619
PURCHON, N208, N210, N212-
 N213
Purpurina, N463
Puschia, N567
Pusillana, N669
pustules, N69
Pustulosa, N437
Puyseguria, N542
Pyanomya, N398
Pycnodonte, N154, N178
Pycnodonteinae, N219
Pyganodon, N444
Pygocardia, N648
Pyramia, N827
Pyramus, N827
Pyrenomoeus, N233
Pyrona, N802
Pythina, N522
Pythinella, N532
Pyxio, N354
Pyxis, N354
Quadrans, N619
quadrante, N107
Quadratonucula, N235
Quadratotrigrionia, N488
Quadrula, N436
Quadrulinae, N219, N436
Quacsithyria, N420
Quantulopinna, N283
Quasilucina, N496
Quendreda, N279
Quenstedt muscle, N38, N108
Quenstedtia, N635
Quenstedtiidae, N220, N634
Questimya, N852
Quidnipagus, N619
Quincuncina, N442
Quoieccchia, N487
radial, N69, N108
Radiatodonta, N398
Radiatula, N428

- Radioconcha*, N655
 Radiocondyla, N560
 Radiocrista, N863
 Radiolitella, N808
 Radiolites, N750, N752, N759,
 N776, N806
 Radiolitidae, N53, N78, N221,
 N754, N756, N760-N764,
 N766, N803, N806, N861
 Radiolitinae, N221, N758, N760.
 N766, N806
 Radiopecten, N361
 Radobornia, N525
 Radula, N385
 Radulidae, N385
 Radulonectes, N352
 Radulopecten, N362
 Raeta, N606
 Raetella, N606
 Raetellops, N606
 Raetina, N606
 Raetolucina, N582, N648
 Raetomya, N699
 Raetomyidae, N221, N698
 RAFINESQUE, N208
 RAGOZIN, N409
 Raikaia, N817
 Raina, N680
 Rakhia, N658
 Raleta, N698
 Ramsetia, N717
 Rangia, N601
 Rangianella, N601
 RANSON, N98
 Rasia, N254
 RAUP, N84, N88
 Rayana, N862
 Rebusum, N867
 Recticardo, N502
 Rectidens, N434
 Rectodontia, N409
 rectum, N22
 Recurvella, N499
 Redicirce, N672
 Redonia, N111-N112, N399,
 N546
 Redoniidae, N543
 REES, N97
 Regalilima, N391
 Regina, N244
 Regozara, N587
 Remondia, N577
 Renatus, N418
 Reneus, N418
 Reniella, N332
 renogenital pore, N29
 Replidacna, N594
 reproductive organs, N29
 Requienia, N756, N759, N763,
 N768, N780
 Requieniana, N862
 Requeniidae, N221, N754-N755,
 N759-N763, N766, N779,
 N799, N861
 Requienites, N780
 Resania, N607
 Resatrix, N679
 resilia, N61
 resiliifer, N108
 resiliifer, N49, N108
 resiliium, N49, N108
 Retha, N761, N787
 reticulate, N108
 Reticulatus, N426
 Retroceramus, N320
 Reubenia, N312
 Rexithaerus, N623
 Rhabdopitaria, N675
 Rhabdoporphorus, N434
 Rhachidoceramus, N317
 Rhadinoceramus, N321
 Rhaetavicula, N304
 Rhaetidia, N639
 Rhectocyma, N568
 Rhectomyax, N633
 Rhedensia, N802
 Rhenania, N475
 Rhenania, N867
 Rhenanomya, N867
 Rhinchotropis, N867
 Rhinoclama, N854
 Rhinomya, N854
 Rhipidocardium, N859
 Rhipidodonta, N460
 Rhomalea, N683
 Rhomboidiana, N862
 Rhombocardia, N861
 rhomboidal, N108
 Rhomboidella, N276
 Rhomboides, N700
 Rhombomya, N832
 Rhombopteria, N334
 Rhombunio, N441
 Rhombuniopsis, N442
 Rhynchomya, N845
 Rhynchomytilus, N276
 Rhynchopterus, N305
 Rhynchotropis, N249
 rhythmic consecutive sexuality,
 N29
 Rhytidonaia, N418
 Rhytimya, N113
 Rhytoides, N817
 ribs, N69, N108
 Ribeirella, N868
 Ribeiria, N868
 Ribeiriidae, N110
 riblets, N69, N108
 Rictocyma, N568
 RIDWOOD, N210, N348
 right valve, N42, N108
 Rimmyjimina, N546
 Rinetrigonina, N487
 Ringicardium, N585
 Rivulina, N669
 Robaia, N237
 Robertella, N809
 Rocellaria, N699
 Rochanaia, N467
 Rochefortia, N531
 Rochefortina, N610
 Rochefortula, N532
 Rockanaia, N862
 Roemeria, N302
 Rogeria, N371
 Rohea, N679
 Rohini, N681
 Roissya, N573
 Rollierella, N649
 Rollieria, N237
 Rollieria, N649
 Rombergia, N622
 Roncania, N391
 ROQUAN, N816
 Roseana, N669
 Rosenbuschia, N640
 ROSEWATER, N162, N169
 Rossmassleriana, N862
 rostrate, N108
 Rostratiana, N862
 Rostrimacoma, N623
 ROSTROCONCHIDA, N121
 Rostroperna, N325
 Rostrotortus, N867
 Rotundaria, N439
 Roudaireia, N650
 Roudairia, N650
 Rousseauia, N299
 Rousselia, N797
 Roxxoa, N701
 Rudicardium, N585
 RUDISTA, N121
 Rudistae, N119
 RUDISTES, N214
 Rudistes, les, N749
 Ruditapes, N682
 RUDWICK, N136-N137, N140,
 N146, N156, N171-N172
 Ruellania, N418
 Ruellia, N604
 Ruganodontites, N463
 Rugifera, N448
 Rugoshyria, N461
 Rumanicana, N862
 RUNNEGAR, N217, N824
 Rupellaria, N689
 Rupicilla, N850
 Rupicola, N850
 Rurynea, N420
 Rutitrigonia, N487
 Rutotia, N339
 Ryderia, N239
 Ryenella, N276
 Rytia, N441
 Rytimya, N818
 Rzehakia, N690
 Rzehakiidae, N221, N690
 Sabbaia, N441
 Sabinia, N758, N789, N797
 Saccella, N237
 SAEMANN, N749
 Saffordia, N399
 sagittal plane, N42, N108
 Sainschandia, N489
 Saintia, N385
 Saintiopsis, N378
 Sakawanella, N642
 Salacia, N686
 Salaputium, N576
 Salebrolabis, N276

- Salmacoma, N623
Saltermya, N293
 Saltocuna, N560
Salweyia, N245
 Samacar, N256
Samanoetia, N261
 Samarangia, N679
 Samarangiinae, N221, N679
 SANDBERGER, N414
 Sandriana, N862
 Sanguinolaria, N633
 Sanguinolariinae, N220, N633
 Sanguinolites, N112-N115, N823
Sanguinolitiidae, N819
 Sanoarca, N867
 SARDESON, N112
 Sarepta, N239
 Sarka, N868
Sarlatia, N809
 Sarmatimactra, N598
 Saturnia, N235
 Sauvagesia, N758-N759, N769,
 N772, N804, N810
 Sauvagesiinae, N221, N758,
 N766, N769, N810
Savignyarca, N252
 Sawkinsia, N595
Saxicava, N700
 Saxicavella, N701
Saxicavidae, N700
 Saxidomus, N679
 Saxolucina, N505
Sayunio, N424
 Scabies, N434
Scabrae, N477
 Scabrotrigonia, N487
 Scacchia, N522
Scaechlamys, N355
 Scaeoleda, N237
 Scalaricardita, N551
 Scaldia, N514
 scales, N69, N108
Scalenaria, N437
 Scalenilla, N453
 scalloped, N70, N108
 Scalpomactra, N601
 Scambula, N577
 Scambulinae, N220, N577
 Scapharca, N256
 Scaphellina, N476
 Scaphellinidae, N219, N476
 Scaphites, N197
Scaphitrigon, N485
Scaphogonia, N485
Scaphoideae, N477
 SCAPHOPODA, N195, N197
 Scaphotrigonia, N487
 Scaphula, N256
 Scaphura, N860
Scapularca, N263
 Schafhaeutlia, N514
 Schedocardia, N585
 Schedotrapezium, N650
 SCHENCK, N38
 Schepmania, N442
 Schiosia, N764, N771-N773,
- N789, N797
Schistodesmus, N434
 Schizocardita, N554
Schizoceramus, N315
Schizocleithrum, N420
Schizodesma, N601
Schizodon, N475
 schizodont, N52, N108
 SCHIZODONTA, N118, N120-
 N121, N214
 SCHIZODONTIDA, N214
 Schizodus, N115, N475, N912
 Schizopecten, N860
 Schizostoma, N862
Schizothaerus, N604
 Schlesiella, N460
 SCHMIDT, N172
Schroteria, N712
Schumacheria, N626
 Scintilla, N536
 Scintillona, N536
 Scintillorbis, N536
 Scintillula, N525
Scintillula, N541
 Scioberetia, N14, N532
 Scissodesma, N601
 Scissula, N619
Scissula, N240
 Scissulina, N624
 Scittila, N635
Scobina, N709
 Scobinopholas, N709
Scolianodon, N463
Scolioceramus, N315
 Scrobiculabra, N636
 Scrobicularia, N21, N45, N635
 Scrobiculariidae, N13, N62, N220,
 N635
Scrobiculariinae, N635
Scrobiculina, N626
 Scrupeana, N862
 Sculptunio, N416
 sculpture, N69, N108
 Scutarcopagia, N619
Scutigera, N712
 Scutum, N64
 Scyphomya, N720
 Sechurina, N676
 secondary riblet, N69, N108
 Sectiarca, N256
 Sectipecten, N363
 Securella, N686
 Securilla, N862
 Sedgwickia, N823
 Seebachia, N572
 Seendia, N568
Seguminaia, N424
 SEILACHER, N382
 Selenimyalina, N291
Selenoceramus, N315
 Sellaea, N752, N757-N758,
 N760-N761, N764, N771,
 N773, N787
Sellaria, N384
 Semelangulus, N619
Semelartemis, N679
- Semele, N49, N636
 Semelidae, N62, N195, N199,
 N220, N636
 Semelina, N637
 Semeloidea, N527
 Semicorallidomus, N113, N399
 Semicorbula, N698
 Semierycina, N522
 semi-infaunal bivalves, N141,
 N162
 Semimodiola, N271
 Semimytilus, N274
 Semipallium, N67, N365
Semipeecten, N354
Semiplicatula, N383
 Semuridia, N314
Senectidens, N265
 SÉNESSE, N803
 Senilia, N256
 Senis, N612
 sense organs, N26
 sensory tentacles, N29
Sepidacna, N594
 Septaria, N739
Septaria, N739
 Septeuilia, N637
 SEPTIBRANCHIA, N18, N119-
 N121, N210, N214
 Septifer, N32, N147, N174, N274
 Septihoernesia, N312
 Septimyalina, N291
 Septiula, N861
 Septocardia, N585
 Serania, N391
Seratisolen, N639
 Sergipia, N320
 Serracorbula, N694
 Serratina, N620
Serratisphoerium, N670
Serratovola, N367
Serridens, N533
Serrilaminula, N666
 Serripecten, N363
 Serripes, N590
 Serrula, N628
Servitor, N235
Servatrina, N283
Sestra, N247
 Setaliris, N857
 Setotrigonia, N487
 Sexta, N512
 sexual dimorphism, N73
 Shanina, N247
 Shaninopsis, N247
 Sheldonella, N262
 shell, N40, N141; rudist, N751;
 Thyasira, N185; unionacean,
 N412
 shell characters, N83
 shell sculpture, N67
 shell wall, N73
Shistodesma, N434
Shistodesmus, N434
 SHROCK & TWENHOFEL, N143
 Sibireconcha, N411
Silenia, N852

- Silex*, N638
Silicula, N238
Siliqua, N612
Siliquarca, N234
Siliquaria, N639
Silurina, N247
Silurocardium, N244
Simacia, N784
Similipecten, N354
Similivenu, N685
Similodonta, N230
Simomactra, N598
Simonaias, N425
Simonisiana, N862
Simplistrigilla, N623
 SIMPSON, N215
Simpsonaias, N448
Simpsonella, N446
Simpsonia, N420
Simpsoniconcha, N448
Sincyclonema, N347
Sinectenolium, N365
Sinemuria, N578
Sinodesmia, N637
Sinodesmya, N637
Sinodia, N678
Sinodiopsis, N679
Sinomacoma, N628
Sinomya, N861
Sinomylus, N861
Sinonia, N685
Sinovacula, N639
Sinsyclonema, N347
Sintoxia, N454
Sinucongeria, N644
Sinuospagia, N620
 SINUPALEALES, N121
 SINUPALLIALIA, N121
 SINUPALLIATA, N121
sinupalliate, N36, N108
sinus, N108
siphon, N12, N108
siphonal area, N83, N108
siphonal band, N108
siphonal retractor muscles, N13, N36, N108
siphonal structures (rudist), N758, N773
siphonal tube, N706
 SIPHONIDA, N121
siphonoplax, N65, N108, N706
Siratoria, N682
Sita, N568
Slava, N102, N245
Sluha, N235
Sluzka, N112
Sluzka, N247
Smodingoceramus, N315
Socardiana, N862
socket, N108
Sogamosa, N668
Solamen, N276
Solanderina, N673
Soldania, N252
Solecardia, N527
Solecurtellus, N638
Solecurtidae, N220, N637
Solecurtinae, N220, N637
Solecurtoides, N613
Solecurtus, N32, N66, N637
Soleilletia, N668
Solemya, N7, N12, N30, N38, N63, N112, N114, N141, N164-N168, N212, N214, N217, N242
Solemyacea, N17, N63, N118-N119, N211, N218, N241
Solemyacidae, N242
Solemyaria, N243
Solemyarina, N243
Solemyidae, N114, N119-N120, N212, N214, N217-N218, N242
 SOLEMYOIDA, N214, N217-N218, N241
Solen, N13, N137, N166, N168, N610
Solena, N611
Solenacea, N16, N63, N118, N121, N211, N220, N610
Solenaia, N435
Solenaria, N611
Solenella, N233
Solenia, N610
Solenidae, N36, N38, N52, N60, N119, N137, N220, N610
Solenimya, N242
Solenomorpha, N114, N823
Solenomya, N20-N21
Solenomya, N242
 SOLENYMYACEA, N241
Solenomyacidae, N242
Solenomyadae, N242
Solenomyidae, N242
Solenopsis, N823
Solenymia, N242
Solenymya, N242
Soletellina, N634
Solidicorbula, N694
Sollea, N867
Solyma, N621
Somapecten, N347
Somapteria, N305
Somarctica, N650
Soror, N247
sortie, N759
Souleyetia, N637
Sowerbya, N639
Sowerbyidae, N220, N639
Sowteria, N250
Soyokellia, N537
Spanila, N297
Spaniodon, N653
Spaniodontella, N653
Spaniorinus, N536
Spatha, N38
Spatha, N463
Spathella, N399
Spathella, N463
Spathophora, N854
Spathopsis, N463
Spathoteredo, N737
Spathoteredo, N737
Speleopisidium, N669
Spengleria, N699
Spengleriana, N862
Spencoceramus, N320
Sperchinosiana, N862
Sphaena, N692
Sphaenia, N692
Sphaera, N514
Sphaerella, N517
Sphaeriidae, N5
Sphaeriidae, N669
Sphaerinova, N670
Sphaeriola, N514
Sphaerium, N11, N57, N670
Sphaerocaprina, N797
Sphaerocardium, N585
Sphaeroceramus, N317
Sphaerucaprina, N772-N773, N789, N797
Sphaerulites, N749-N750, N771, N774, N799, N808, N810-N811
Sphaerumbonella, N532
Sphenalia, N529
Sphenia, N692
Spheniopsidae, N221, N699
Spheniopsis, N699
Sphenoceramidae, N314
Sphenoceramus, N320
Sphenolium, N113, N818
Sphenolium, N397
Sphenomya, N820
Sphenonaias, N426
Sphenotomomorpha, N399
Sphenotrigonia, N487
Sphenotus, N823
Spherulites, N810
Sphoerocardium, N585
Spincullaea, N867
spine, N69, N108
Spinearca, N263
Spineilo, N235
Spinospella, N855
Spinula, N235
Spinularca, N258
Spirodomus, N868
spirogyrate, N46, N108
Spissatella, N576
Spisula, N601
Spisulina, N601
Spisulona, N604
Spixiconcha, N465
Spodylus, N378
Spondaeana, N863
Spondervilia, N610
Spondilus, N378
Sponditolites, N867
Spondylidae, N8, N12, N35, N37, N72, N76, N119, N154, N173, N212, N219, N378
Spondyligenus, N378
Spondylopecten, N364
Spondylus, N16, N29, N47, N49-N50, N53, N60, N63, N137, N154-N157, N168, N170,

- teeth and sockets (rudists), N754, N756
- Tegillarca*, N254
- TEIXEIRA, N110
- Telederma*, N436
- Telemactra*, N595
- TELEODESMACEA, N119-N121, N210, N214
- teleodont, N108
- TELEODONTA, N121
- Telistrophis, N868
- Tellidora, N623
- Tellidorella, N580
- Tellimera*, N623
- Tellimya, N532
- Tellina, N4, N11, N19, N21, N30, N36, N38, N58-N60, N62-N64, N77, N86, N141, N166, N181, N613
- Tellinacea, N16, N36, N63, N84, N98, N119, N121, N211, N220, N613
- Tellinacées*, N613
- Tellinangulus, N620
- Tellinarius*, N613
- Tellinella, N51, N68, N71, N621
- Tellinidae, N13, N16, N23, N36, N38, N73, N195, N220, N613, N643
- Tellinidella, N621
- Tellinides, N621
- Tellinimactra, N627
- Tellinimera, N623
- Tellininae, N220, N613
- Tellinocyclas, N666
- Tellinodora*, N623
- Tellinomorpha*, N838
- Tellinomya*, N228, N532
- Tellinopsis, N229
- Tellinota*, N621
- Tellinula*, N615
- Tellinungula*, N626
- Tellipiura, N623
- Tellurimya, N662
- Temblornia, N527
- Temnoconcha*, N624
- Tendagurium, N589
- Tenea, N650
- Tenka, N297
- Tentidonax, N628
- Tenuicorbula, N694
- Tenuipteria, N310
- Tenuis*, N297
- Tepeyacia, N817
- Tepidoleda*, N239
- Terebrimya, N740
- Teredaria*, N722, N725
- Teredarius, N861
- Teredidae*, N722
- Teredigenus*, N725
- Teredina, N717
- Teredina*, N722
- Teredinadae*, N722
- Teredinidae, N8, N11, N13, N65, N173-N174, N221, N702, N722, N861
- Teredininae, N221, N725
- Teredinites*, N722
- Teredinopsis, N740
- Teredo, N4, N8, N13, N65, N67, N142-N143, N165, N169-N170, N704, N724-N725
- Teredolites, N740
- Teredolites*, N740
- Teredolithes*, N740
- Teredolithus, N470
- Teredops*, N727
- Teredora, N724, N729-N730
- Teredothyra, N729
- Terentia, N681
- Teretileda*, N235
- Tergulina, N854
- Teridina*, N717
- terminal, N45, N108
- Terquemia, N380
- Terquemiidae, N72, N115, N219, N380
- Terraia, N115, N643
- Terua, N280
- TESTACEA, N4
- Tethyoceramus*, N315
- Tetinka, N297
- Tetoria, N666
- Tetorimya, N837
- Tetracionites, N758, N803
- Tetragonostea*, N847
- Tetraplodon*, N458
- Textivenus, N683
- Textrix*, N685
- Thalassides*, N578
- Thalassites*, N578
- Thecalia, N73, N557
- Thecaliinae, N220, N557
- Thecodonta, N532
- Theliderma*, N436
- Theora, N195, N199, N637
- Theringia*, N590
- Thestyleda, N237
- Thetiopsis, N682
- Thetironia*, N512
- Thetis, N512
- Thevenimia*, N567
- Thiatira*, N500
- thickness, N108
- THIELE, N3, N213-N214
- THOMPSON, N84
- Thoracoceramus*, N320
- THORAL, N110, N112
- Thorslundia, N250
- Thovana, N708
- Thracia, N13, N117, N850
- Thracidentula*, N852
- Thracidora, N852
- Thraciidae, N73, N222, N850
- Thraciinae*, N850
- Thraciopsis, N852
- thread, N69, N108
- Throna*, N850
- Thronacar*, N252
- Thurlosia*, N710
- Thyas, N860
- Thyasira, N135, N164, N167, N180, N184-N187, N189-N194, N195-N202, N508
- Thyasiridae, N9, N52, N195, N202, N206, N220, N508
- Thyella*, N637
- Thyellisca, N637
- Thyrastylon, N158, N759, N773, N777, N810
- Thyreopsis, N536
- Tiaraconcha*, N245
- Tibialectus*, N275
- Tigammona*, N672
- Tikia, N676
- Timoclea, N688
- Timoria*, N314
- Timothyunus, N517
- Tinctora, N675
- Tindaria, N235
- Tindariopsis, N235
- Tiria*, N858
- Tirolidia, N391
- Titanosarcolithes, N158, N751, N762, N771, N789, N799
- Tivela, N673
- Tivellina, N675
- Tiza, N697
- Toechomya, N475
- Tolmaia*, N302
- Tomala, N698
- Tommasina, N280
- Toralimysia, N515
- Torastarte, N580
- Toredo*, N725
- Torinosulcatella, N259
- Torreites, N758, N769, N771-N772, N776, N803
- Tortarctica, N650
- Tortocardia, N656
- Torulosa, N453
- Tosapecten, N371
- Tosarca, N860
- Totteniana*, N681
- TOUCAS, N749, N803, N811, N816
- Toucasia, N154, N751-N752, N759, N780, N781, N784-N785
- Toxelasma*, N451
- Toxolasma*, N451
- trace elements, N77
- Trachycardiinae, N220, N586
- Trachycardium, N586
- Trachyceramus*, N317
- Transennella, N675
- Transitrigonia*, N487
- transposed hinge, N56, N108
- transverse, N108
- Trapezacea*, N644
- Trapezicardita, N556
- trapeziform, N108
- Trapeziidae, N221, N655
- Trapezium, N86, N655
- trapezoidal, N108
- Trapezoides, N435
- Trautscholdia, N567
- Trechmanella*, N795

- Trechmannellidae*, N787
 Tresus, N604
 Triaphorus, N546
 Tricassiniana, N863
 Trichites, N76, N284
 Trichomusculus, N276
 Trichomya, N274
Trichomya, N834
 Trichomyella, N834
Tridachnes, N594
 Tridacna, N7, N11, N34, N47, N69-N70, N79, N118, N135, N159, N169, N594
 Tridacnacea, N16, N118, N121, N211, N220, N594
Tridacnadae, N594
Tridacnae, N594
 Tridacnidae, N13, N220, N594, N861
Tridacnodites, N594
 Tridacnoides, N861
 Tridonta, N568
 Trifurcardium, N589
Trigomyophoria, N473
Trigona, N673
 trigonal, N108
 Trigonarca, N269
 Trigonastarte, N573
trigonées, les, N471, N476
Trigonella, N473, N595, N674
Trigonellites, N472
 Trigonina, N38, N63, N477-N478
 Trigoniacca, N16, N18, N46, N52, N56, N63, N76, N114, N120, N210-N211, N219, N471
Trigoniana, N476
 Trigonidae, N12, N52, N83, N115-N117, N119-N121, N212, N220, N476-N477
 Trigonocardia, N588
 TRIGONIOIDA, N214, N218-N219, N471
 Trigonoides, N489
 Trigonoididae, N220, N488
 Trigonocallista, N679
Trigonocardia, N650
Trigonocoelia, N265
 Trigonodesma, N264
 Trigonodon, N434
 Trigonodus, N468
 Trigonopsis, N571
Trigonosphaerium, N670
 Trigonothracia, N850
 Trigonucula, N231
Trigonulina, N855
 Trinacria, N264
Trinacriella, N264
 Trinacriinae, N218, N264
 Trinitasia, N505
Triodonta, N500, N559, N568
Triumphala, N720
Triumphalia, N720
 Tripleura, N572
 Triplicosta, N837
 Tripilodon, N461
Triquetra, N457, N610, N686
 Triquetrana, N462
 Trisidos, N72, N254
Trisis, N254
 Trisulcus, N471
 Tritogonia, N442
Triumphalia, N720
Trochoceramus, N320
 trochophore, N91
Trommia, N817
Tropidocardium, N584
Tropidocyclus, N669
 Tropidomya, N854
Tropidophora, N854
 Tropinuculites, N235
Tropiphora, N475
 TROPIPODA, N227
 Tropithaca, N688
 TRUEMAN, N174
 TRUEMAN, BRAND, & DAVIS, N166
 Truncacila, N231
 truncate, N108
 Truncilla, N457
 Truncillopsis, N453
Trutina, N843
 Tryphomyax, N536
Tschaudia, N592
 Tuangia, N688
 tubercles, N69
Tuberculina, N496
 Tuberunio, N416
Tubolana, N858
Tubulana, N858
 tubular protective structures, N65
Tuceta, N267
Tucetilla, N267
Tucetona, N252, N267
Tucetonella, N252
Tucetopsis, N267
Tudes, N327
 Tugonella, N692
 Tugonia, N692
 Tugoniopsis, N692
 Tumbeziconcha, N67, N598
 tumid, N108
Tumidusiana, N863
 TUREKIAN & ARMSTRONG, N77
 Turia, N688
Turkmena, N593
 Turkmenia, N321
 Turnus, N741
 Turquetia, N536
 Turritella, N195, N199 ✓
 Turritellidae, N195
 Turtonia, N539
 Turtoniana, N863
 Turtoniidae, N220, N539
 Tusayana, N583
 Tutcheria, N552
 Tutuella, N411
Tychoecardia, N657
Tyleria, N692
 Tylophora, N399
 typhlosoles, N22
 Uddenia, N577
 ULRICH & BASSLER, N109
 umbo, N45, N108
 umbonal angle, N108
 umbonal cavity, N45, N109
 umbonal depression, N109
 umbonal fold, N109
 umbonal pole, N45, N109
 umbone, N45
Umitakea, N708
 Uncidens, N536
 Undatimusculus, N276
Undulatae, N477
 undulations, N69
 Undulomya, N114, N838
Ungoteredo, N729
 Ungulina, N515
Unguline, N515
 Ungulinidae, N220, N514
 Uniandra, N423
Unicardiidae, N634
 Unicardiopsidae, N220, N634
 Unicardiopsis, N634
 Unicardium, N513
Uniigenus, N415
 Unio, N96, N402-N403, N406, N415
 Uniocardium, N594
 Uniomerus, N435
Uniona, N468 ✓
 Unionacea, N9, N16, N38, N52, N63, N76, N96, N118, N120, N209, N211, N219, N401, N411, N862
 Unionea, N435
Unionea, N415
 Unionella, N471
Unionella, N435
 Unionerius, N435
 Unionidae, N5, N12, N38, N51, N57, N70, N116, N219, N411-N412, N414-N415, N861
 Unioniformiana, N863
Unionina, N468
 Unioninae, N219, N415
 Unionites, N468
Unionium, N304, N861
 UNIONOIDA, N214, N217, N219, N401
Unionopsis, N447
 Unionum, N861
Uniopsis, N447
 Uperotus, N730, N740
 UREY, N134
 Uromya, N832
 Ursirivus, N696
Ustularca, N252
Utela, N463
 Utschamiella, N411
Utterbachia, N444
Utterbackia, N444
 Utterbackiana, N444
 Vaccinites, N769-N770, N776, N803
 Vacunella, N838
Vagina, N610

- Valletia, N754-N755, N781, N784
 Valsella, N332
 valve, N40, N109
 valve gape, N170
 Vanderschliea, N449
 Vanganella, N607
 Vanuxemia, N112, N250
 Variamussium, N350
 Varicardium, N589
 Varicorbula, N695
 Variopecten, N350
 Varotoga, N536
 Vasconella, N266
 Vasconia, N537
 Vasconiella, N537
 Vasticardium, N588
 Vaticinaria, N510
 Vaugonia, N488
 Vautrinia, N774, N777, N815
 Vectianella, N650
 Vectorbis, N679
 Velargilla, N689
 Velata, N373
 Velates, N373
 Velatopecten, N373
 Veldidenella, N344
 Veleda, N598
 Velesunio, N5, N461
 Velutuceta, N267
 veliconch, N94, N109
 veliconcha, N94
 veliger, N93
 Velopecten, N373
 Velorita, N668
 Veloritina, N668
 velum, N93
 Velunio, N428
 Venatomya, N692
 VENERACEA, N491
 Veneracea, N13, N16, N63, N121, N211, N221, N670
 Venerella, N685
 Venericardia, N57, N555
 Venericardiinae, N220, N554
 Venericor, N556
 Venericyprina, N648
 Veneridae, N13, N16, N52, N56, N73, N76, N117, N138, N141, N159, N166, N168, N173, N175, N221, N670, N863
 Veneridia, N670
 Veneriglossa, N664
 Venerinae, N221, N670
 Veneritapes, N685
 VENEROIDA, N214, N218, N220, N491
 Venerupis, N685
 Venetia, N650
 Venidia, N675
 Veniella, N650
 Veniellinae, N645
 Venilia, N645, N650
 Venilicardia, N70, N72, N650
 Venilidae, N645
 Venniella, N650
 venous system, N24
 Ventalium, N371
 ventral, N42, N109
 ventricle, N24
 Ventricola, N671-N672
 Ventricularis, N672
 Ventricoloidea, N672
 ventricose, N109
 Ventricosiana, N863
 ventromedian scars, N38
 Venus, N13, N21, N63, N671
 Venusarius, N671
 Venusta, N297, N450
 Venustaconcha, N450
 Vepricardium, N586
 Veprichlamys, N355
 Veremolpa, N688
 Verilarca, N263
 VERMES, N4
 VERMES TESTACEA
 BIVALVIA, N227
 Vermitexta, N527
 Verpa, N859
 Versipella, N265
 Vertambitus, N855
 Verticipronus, N867
 Verticordia, N855
 Verticordiidae, N13, N222, N854
 Vertipecten, N364
 Vertisphaera, N856
 Vertumnia, N333
 Vesicomya, N664
 VESICOMYACIDAE, N664
 Vesicomysidae, N221, N664
 VEST, VON, N118
 Veteranella, N239
 Vetericardia, N552
 Vetericardiella, N552
 Vetoarca, N264
 Vetocardia, N553
 Vetulonaca, N436
 Vetulonaia, N436
 Vevoda, N247
 Viaderella, N506
 Vietuliana, N863
 Vigorniella, N250
 Vilasina, N276
 Viliaeana, N863
 Vilianana, N863
 Villarita, N861
 Villorita, N668
 Villosa, N450
 Vimentum, N551
 vinculum, N109
 Virgula, N463
 Virgus, N436
 Virmysella, N533
 visceral ganglia, N26
 visceral mass, N11
 Vitracar, N252
 Vlasta, N247
 Vlastiden, N247
 Vlastinae, N218, N247
 Vogel, N110, N801
 VOKES, N213, N386
 Vokesella, N541
 Vokesula, N695
 Vola, N366
 Volachlamys, N357
 Volsella, N278, N332
 Volsellina, N280
 Volupia, N499
 Volupicuna, N561
 Volviceramus, N142, N321
 Vulcanomya, N854
 Vulcella, N332
 Vulgodacna, N594
 Vulsella, N11, N332
 Vulsellidae, N38, N212
 Vulsellidae, N326
 Vulselligenus, N332
 Vulsellina, N332
 Vulsellopsis, N329
 Vulsinella, N332
 Waagenoperna, N326
 Waiuiuconcha, N664
 Wakullina, N383
 WALCOTT, N109
 wall structure (rudist), N771
 WALLER, N170
 Wallucina, N499
 Wanwania, N868
 Wanwanella, N868
 Wanwanoidea, N868
 Warburgia, N250
 Warnea, N859
 Warrana, N653
 water pores, N18
 WERNER, N93
 Westerlundiana, N863
 Westralunio, N461
 Weyla, N73, N116, N371
 WHITEAVES, N402
 Whiteavesia, N399
 Whitella, N249
 Wilkingia, N114-N115, N838
 WILSON, N21, N408
 wing, N46, N109
 WÖHRMANN, N409
 Woodia, N565
 WOODWARD, N21, N119, N208, N749, N799
 WRIGLEY, N100
 x-axis, N89
 Xenamussium, N351
 Xenocardita, N557
 Xenoceramus, N315
 Xenoloupia, N673
 xenomorphic, N109
 xenomorphism, N109
 Xilophaga, N721
 Xilotoma, N721
 Xiphoceramus, N321
 Xylophaga, N702, N704, N721
 Xylophagella, N722
 Xylophaginae, N221, N702, N721
 Xylophagi, N721
 Xylophagidae, N721
 Xylophaginae, N721

