

Useful
PRESCRIPTIONS

Compiled by
CLOYCE WILSON, M. D.

Published By
LLOYD BROTHERS, PHARMACISTS, INC.
CINCINNATI, OHIO
MARCH 1935

Foreword

The prescriptions comprising the major portion of this little volume have been carefully selected from a large collection contributed by many practicing physicians. Through years of study and clinical experience, they have been found applicable and serviceable in the treatment of the sick.

On the concluding pages the indications for each Specific Medicine mentioned in the prescriptions are briefly stated and the average dose given. It is hoped these pages will prove of value by enabling the physician to vary the prescription when needed, or to choose a single remedy adapted to the case under treatment.

We believe the recipients of this volume will join in our expression of appreciation to Dr. Cloyce Wilson for his painstaking work in assembling and classifying these prescriptions; to Dr. Byron H. Nellans for constructive suggestions, and to the many physicians who have contributed prescriptions that have proved of value in their daily practice.

Copyright 1935
By
LLOYD BROTHERS, PHARMACISTS, INC.
Cincinnati, Ohio

LLOYD BROTHERS PHARMACISTS, INC.

Contents

Preface	7
Skin	19
Ear and Upper Respiratory System	26
Lower Respiratory System	31
Alimentary Tract	39
Cardio Vascular System	53
Nervous System	58
Genito-Urinary System	64
Reproductive Systemó Female	72
Neuro-Muscular System and Joints	79
Alteratives and Tonics	84
Specific Medicines	89
Precipitates in Specific Medicines	91
Compatibility	91
Colloidums	92
Comparative Strength of Specific Medicines	93
Vehicle for Dilution	95
Indications and Doses of Specific Medicines	97

Preface

CLOYCE WILSON, M.D.

The physician is ever alert to find improved methods of treating the sick. He welcomes suggestions. Yet experience in his own practice teaches him the dependability of certain remedies in certain conditions. As physicians pass on their own experiences, the knowledge of practical medicine is enriched.

In the following pages are presented prescriptions which have been found efficacious in the treatment of disease and disease-expressions. This presentation represents the best of those gleaned from the correspondence of hundreds of physicians successfully engaged in the actual practice of medicine. They are the fruits of actual clinical experience.

These prescriptions are presented, not as hard and fast formulae, which must be strictly followed, but as *suggestive forms* which will aid the prescriber in formulating his own methods of prescribing. The careful prescriber uses as few remedies as possible in his prescriptions. The ideal method, possibly is to exhibit but one drug at a time. But certain remedies have been found to be synergistic, and experience has shown the wisdom of their intelligent combination.

PREFACE

Many drugs are known to have affinities for certain tissues, groups of tissues (organs), or bodily systems. Knowledge of this selective action aids in their proper application. This knowledge is imperative to proper understanding of drug therapy. Disease names are given only as aids in finding the indicated treatment. We do not believe in the treatment of the disease by name only, but rather the treatment of the underlying wrongs of the disease, as expressed by tangible signs and symptoms. The physician must necessarily study carefully the case in hand, to arrive at a correct working diagnosis, to understand the underlying pathology, the changes to be expected, the duration, the susceptibility to treatment and possible ultimate cure, as well as the possibility of complications and involvement of other structures.

It will be noted that many of the prescriptions are applicable to chronic lesions. Comparatively few inquiries have come regarding the treatment of the acute or the sub-acute diseases; it is in the field of chronic disease that the practitioner finds his deepest problems. These answers have been taken from *Gleaners*, Nos. 1 to 42, from correspondence connected with the publishing of the *Gleaner*, from textbooks and active journals of medicine, and from prescriptions furnished by physicians engaged in the day-by-day practice of their profession, under actual

PREFACE

conditions, in the office, at the bedside, in the hospital and in the clinic.

It would not be practicable to attempt to arrange a complete list of prescriptions, covering the wide range of disease expressions. It is hoped that those given will form the basis for an intelligent method of treatment, though many times requiring change to meet the requirements of the particular case in hand.

There are, of course, limits to the therapeutic action of medicines. Structure is but little affected directly, by any form of medication. We may hope, however, to correct disordered function, thus restoring the patient's condition to a more nearly normal state. By so correcting abnormal function, indirectly, many wrongs with a pathological basis may be righted.

Medicines are not to be expected to supplant conservative and intelligent surgery, dietetics, hygiene and other forms of therapeutics recognized and used by all physicians; but presuppose a close alliance with all these forms of treatment. Medicines are a means to an end, the instruments by which we work our art. And so considered are a source of confidence and reliance to the physician, profit and well-being to the sufferer. So far as drugs are intelligently and understandingly used, they are capable

of immense benefit. The physician must recognize definitely, where one form of treatment begins and another ends.

The dosage of drugs varies according to the effects desired from their use. The amounts given may be modified to suit the individual case. When drugs are administered for their physiological effect, a safe rule to follow when prescribing for children, is Young's Rule. To find the dose for a child: Divide the age of the child by the age plus 12. The resulting fraction will approximate the fraction of the adult dose required.

All individuals are not susceptible alike to drug influence. Some react in an atypical manner. This individual reaction is known as idiosyncrasy, and can only be known by experience with the particular patient. Any drug which is known to be capable of this effect must be used in small amounts until the patient's reaction is known. It is often difficult to impress upon the average student of therapeutics, that the small dose frequently repeated does have value in treating the sick.

Many other factors determine the dosage. The general condition of the patient, age, sex, habits, interval between doses, presence or absence of food in stomach, are but a few of the factors, which must be considered in making up an intelligent prescription.

The amounts given in the following prescriptions have been found correct in practice with the average patient, but may require change in certain instances. The doses given upon the Specific Medicine labels have been found safe and effective in years of actual use, and should not be exceeded without care and a full understanding of possible untoward and even poisonous effects. The Specific Medicines are active energetic preparations, representing the desirable medicinal qualities of the drugs from which they are prepared. "The dosage should be the smallest possible quantity that will cure disease." (Dr. J. M. Scudder)

GENERAL CONSIDERATIONS

So large a proportion of the medical profession follows the natural method of drug selection, many times unconsciously, that it has been deemed a necessary result of this tendency to formulate and present reasons for the selection of drugs in certain disease manifestations. This knowledge comes from long experience and careful observation. It is not sufficient to merely diagnose the name of the disease, praiseworthy as that may be. The study must go further and determine how far and in what ways the condition of the patient has departed from the normal. The most successful prescriber is he who treats the conditions which actually are present.

That which we call disease may always be divided into its component parts. In our system of medicine agents are used to correct, influence or return to normal those functions which have become abnormal. We are not always able to influence all disease expressions at one time, but by proper observation we may determine the underlying conditions, as shown by clinical signs and symptoms, and in time, by proper medication, in a kindly manner, and by working with natural laws in mind overcome the entire disease complex.

"Disease is not an entity, something to be expelled from the body, but is a method of life. It cannot be purged from the body, vomited from the stomach, strained from the skin and kidneys, or exorcised by counter-irritation. Such means may do good when they remove offending substances, or establish a function that is deficient, but they should have a rational use.

"The life of disease is not the life of health, neither in the whole body, nor in the individual part. It is always exhaustive, impairing the life now and later, by faulty renewal of tissue. We study disease, therefore, as an impairment of living, and we treat the diseased body as a living body which has been placed in such position that its life has become enfeebled or deranged.

"Disease has certain expressions which we call signs or symptoms, as health has certain expressions. We find the manifestations of health uniform and consistent and we can hardly mistake their meaning. So in disease, the expressions of morbid life are uniform and constant, and do not vary in different individuals. If we determine in any given case, the expression of diseased life, we shall find it the same in all cases. It has been claimed and tacitly admitted, that symptoms of disease are so changeable and inconstant that they could not be depended upon with any certainty. This is certainly true to those who make their diagnosis according to the usual nosology, and then prescribe at the name. For, as many diverse pathological conditions would be grouped under each name, the symptoms would of course vary, and the treatment would show the element of uncertainty in so marked a manner that idiosyncrasy would be called upon to explain the trouble.

"We propose to study the expressions or symptoms of disease with reference to the administration of remedies. It is a matter of scientific interest to know the exact character of a lesion but it is also important to know the exact relationship of drug action to disease expression, and how the one will oppose the other and restore health. If we have determined

such relationship, we have determined it in all disease alike, in all persons, and for all time to come.

"The first lesson in pathology we learn, is, that disease is a wrong life. The first in diagnosis is that wrong life finds a distinct and uniform expression in the outward manifestations cognizable to one's senses. The first lesson in therapeutics, that remedies are uniform in their action; the conditions being the same, the action is always the same.

"All disease has a distinct expression in the outer man. We have a standard of color for the general surface or for special parts. It is in the skin at large, and in those places where the circulation is less free, where it is very superficial and free, as in the nails and veins. We have a standard color for lips, gums, teeth, tongue, mucous membrane. We have a standard in touch, smoothness, resistance, elasticity, size and association. We have a standard respiration, in frequency fullness and depth and freedom, which is distinct and clear, which we know of ourselves, by observation. We have a standard circulation which we know at the radial artery, under the finger, as well as by the expression of the parts associated with the heart in action. We have a standard condition of the digestive organs, which we know by the general expression of the muscles of the mouth, whether the man has a good appetite, good digestion, and good

bloodmaking. Constipation will be shadowed forth in the outer man.

"Bloodmaking and nutrition must have a distinct expression and a standard for measurement will be readily formed, for all functional activity is dependent upon nutrition. The standard of healthy excretion by skin, kidneys and bowels, is readily found by observation, not upon the sick but upon the well, not under the influence of drugs.

"We desire to determine the nature and locality of disease, its pathology, but we also wish to determine those agencies which will oppose disease and restore health to therapeutics.

"The Brain: Determination of blood is marked by bright eyes, contracted pupils, moderate contraction of facial muscles. The direct remedy is Gelsemium. Congestion is marked by dull eyes, expressionless face, dilated pupils, inclination to sleep, eventually coma. The direct remedy Belladonna. Inflammation is marked by greater contraction of the muscles, especially of the orbits and frontal region, deeper flushing of face and constriction of tissues. Here the direct remedies are Aconite and Gelsemium or Veratrum. Gastro-intestinal, a white line around mouth, with fullness of tissue, Podophyllum, Hydrastis or Santonin: With constriction and thinning of tissue, Aconite and Ipecac.

"Bright red flush, left cheek, in acute disease, indicates a lesion of sympathetic and spinal innervation. Rhus tox. is the remedy. Deeper flush right cheek, lesions of serous membranes and fibrous tissues, Bryonia is the remedy. Undue blueness of superficial veins, Iron is the remedy. Deeper redness of tongue and mucous membranes calls for tincture Chloride of Iron.

"Skin: Brownish discoloration, associated with wrongs of digestion, Podophyllum and Hydrastis. Brownish, dull and opaque, associated with leaden tinge of adjacent skin, Chelidonium. Podophyllin if tissue show fullness. Sallowiness, with slight tinge of green calls for copper. Deepened tint of skin, dull and opaque, arsenic or phosphorus. Distinct want of pigment, not anemia, Sulphur as a restorative.

"Tongue. Form: Elongated pointed express irritation and determination of blood to gastro-intestinal tract. It is associated with excitation of the nerve centers. Remedies are, Aconite, Ipecac, Acid Hydrocyanic Dil., or small doses of Rhubarb and Bismuth. Full tongue, broad and thick, shows atony, Nux Vom. Coating: Full, heavy, more at base, emetics. Yellowness, wrongs of hepatic function, calling for Nux Vom., Podophyllum, Hydrastis. Dull leaden coating, Chelidonium. Bright redness, tip and edges, sedatives: Aconite, Ipecac, Rhus tox. Broad pallid

tongue, want of alkali, usually Sodium Bicarb. If associated with wrongs of the muscular system, Potassium Bicarbonate. If of the cellular tissues, some form of lime. Deep red tongue, contracted and dry; calls for acid, usually Dilute Hydrochloric Acid. Deep red or deep violet with fullness, showing sepsis, Baptisia. Full color with violet shade, Acid Nitric Dil. Dull leaden color, Phosphorus or the hypophosphites. Large thick in center, incurved edges, dull blue or leaden color, Arsenic.

"Dirty gray or dirty white, Sodium Sulphite.

"Moisture of tongue assures us that Quinine or opium may be used, if indicated.

"Contraction shows excited nervous system.

"Fullness shows atony, wrongs of the sympathetic nervous system.

"The Pulse:

"The full pulse, with strength, Veratrum.

"The full doughy pulse, Lobelia. Lacks vibration.

"The full open pulse, Podophyllum.

"The full vibratile pulse, Gelsemium.

"The full oppressed pulse, Belladonna.

"The small pulse, Aconite.

"The small vibratile pulse, Aconite with Gelsemium.

"Pulse with want of power, Digitalis.

"Sharp stroke of pulse, with tremulous waves, Rhus tox.

“Sharp stroke with even small vibratile current between strokes, Bryonia.

“Small open pulse, wave drops suddenly, Cactus, Pulsatilla.

“Irregularity, soft and open, Cactus.

“Steady vibratile pulse without marked waves, Macrotys.

“Temperature is but an expression of disease. If disease is favorably influenced by medicines, it tends to fall or rise to the normal standard. As is pulse, so is temperature. Certain remedies influence temperature through the nervous system: Rhus tox., Gelsemium, Bryonia, Belladonna, Nux Vomica, Nitric Acid.

“Periodicity: Whatever may be the cause, Quinine is the usual remedy.”

(Excerpt, *Specific Diagnosis*, Dr. J. M. Scudder)

SKIN

Acne:

℞	Sp. Med. Nux Vom.	gtt. x.
	Sp. Med. Berberis	ʒij.
	Sp. Med. Pulsatilla	ʒj.
	Glycyrrhiza	q.s. ʒiv. M.

Sig: A teaspoonful before meals and at bedtime.

Gl. 40, p. 1266

Acne:

Alterative:

℞	Sp. Med. Berberis	ʒiij.
	Echafolta	ʒiij.
	Sp. Med. Iris	ʒj.
	Elix. Simplex	q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 3, p. 105

Skin Lesions Accompanying Anemia:

℞	Lloyd's Iron.	
	Fowler's Solution Arsenic	āā. ʒj.
	Aq. Dest.	q.s. ʒvi. M.

Sig: A teaspoonful in water after meals.

Gl. 33, p. 1066

Boils:

(a)

- ℞ Syr. Stillingia Comp. ℥iv.
Sig: A teaspoonful every three hours.

(b)

- ℞ Calcium Sulphide gr. vj.
 Sacch. Lact. gr. lx. M.
 Div. in Chart. No. xxiv.
Sig: A powder four times a day.

Gl. 2, p. 21

Recurrent Boils:

Furunculosis:

- ℞ Sp. Med. Echinacea ℥iv.
 Sp. Med. Phytolacca gtt. xx.
 Fl. Ext. Cascara Arom. ℥j.
 Elix. Simplex q.s. ℥iv. M.

Sig: A teaspoonful before meals and at bedtime.

A. W. Hobby, M.D.

Dermatitis Venenata—Rhus Poisoning, etc.:

- ℞ Sp. Med. Grindelia ℥ss.
 Glycerin ℥i.
 Calamine Lotion q.s. ℥iv. M.

Sig: Apply locally four times daily.

G. W. DeMuth, M.D.

Rhus Poisoning:

(a) Locally:

- ℞ Fl. Ext. Grindelia robusta ℥iv.
 Sp. Med. Lobelia ℥ij.
 Sodium Bicarbonate ℥ij.
 Aq. Dest. q.s. ℥viii. M.
Sig: Cleanse areas with soap and water; apply mixture upon gauze dressing.

(b) Locally:

- ℞ Sp. Med. Belladonna gtt. x.
 Alcohol.
 Aq. Dest. āā. q.s. ℥iv. M.
Sig: Apply locally on gauze dressings, keep moist.
 Gl. 35, pp. 1118-1119

Acute Eczema:

- ℞ Sp. Med. Berberis ℥j.
 Sp. Med. Rhus tox. gtt. xx.
 Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every three hours.

Wm. P. Best, M.D.

Itching Eczema:

- ℞ Sp. Med. Apis.
 Sp. Med. Rhus tox āā. gtt. v.
 Sp. Med. Echinacea ℥ij.
 Aq. Dest. q.s. ℥iij. M.
Sig: A teaspoonful every four hours.

Locally Echafolta Cream.

Gl. 36, p. 1156

Infantile Eczema:

(a) Locally:

℞ Ungt. Hydrar. Ammon. ʒiij.
 Ungt. Aq. Rosa q.s. ʒj. M. F. Ungt.

Sig: Apply twice daily.

(b) Internally:

℞ Sp. Med. Apis. gtt. x.
 Sp. Med. Rhus tox. gtt. iij.
 Aq. Dest. q.s. ʒiv. M

Sig: A teaspoonful three times a day.

Gl. 41, p. 1293

Chronic Eczema:

Alterative:

℞ Fowler's Solution Arsenic ʒss.
 Sp. Med. Nux Vom. gtt. x.
 Sp. Med. Berberis ʒj.
 Glyconda.
 Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful after each meal.

Gl. 41, p. 1305

Chronic Eczema:

Alterative:

℞ Sp. Med. Berberis ʒiij.
 Sp. Med. Iris ʒj.
 Sp. Med. Rumex ʒiij.
 Elix. Simplex q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 7, p. 261

Ringworm:

℞ Tr. Iodine ʒij.
 Sp. Med. Ac. Sulphurous ʒiv.
 Aq. Dest. q.s. ʒiv. M.

*Sig: Apply to ringworm.**May be used as throat swab in aphthous conditions.*

Gl. 17, p. 587

Furuncle:

Also useful in Ulcer Dressing.

℞ Sp. Med. Echinacea
 Bismuth Subnitrate āā. ʒj.
 Lanolin
 White Petrolatum āā. q.s. ʒj. M. F. Ungt.

Sig: Apply locally twice daily.

Gl. 39, p. 1248

Hair Tonic, Scudder Formula:

℞ Sp. Med. Jaborandi ʒj.
 Sp. Med. Polymnia ʒj.
 Bay Rum q.s. ʒiv. M.

Sig: Rub into scalp once a day.

Gl. 6, p. 237

Herpes Progenitalis:

℞ Sp. Med. Echinacea ʒij.
 Dist. Hamamelis ʒj.
 Aq. Dest. q.s. ʒiij. M.

Sig: Apply frequently. Allow to dry. Dust with powdered talcum.

Gl. 17, p. 590

Hyperhidrosis:

℞ Sp. Med. Belladonna gtt. x.
 Sp. Med. Ergot ʒij.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.
 Gl. 23, p. 783

Pruritis Ani:

Local Treatment:

℞ Sp. Med. Thuja ʒj.
Sig: Apply on cotton for ten minutes on alternate days.
 G. W. DeMuth, M.D.

Pruritis Ani:

Internal:

℞ Sp. Med. Apis
 Sp. Med. Rhus tox. āā. gtt. v.
 Sp. Med. Collinsonia gtt. x.
 Aq. Dest. q.s. ʒiij. M.
Sig: A teaspoonful after meals and at bedtime.
 Gl. 37, p. 1181

Pruritis, Neurosis:

℞ Sp. Med. Apis gtt. xxx.
 Sp. Med. Pulsatilla ʒj.
 Glyconda
 Aq. Dest. q.s. āā. ʒiij. M.
Sig: A teaspoonful every four hours.
 Gl. 41, p. 1296

Prickly Heat:

℞ Echafolta (Iodized) ʒss.
 Ac. Boric ʒij.
 Dist. Hamamelis ʒiv.
 Aq. Dest. q.s. ʒviiij. M. F. Sol.
Sig: Apply to skin, allow to dry.
 Gl. 17, p. 610

Urticaria:

℞ Sp. Med. Belladonna
 Sp. Med. Apis āā. gtt. v.
 Magnesium Sulph. ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every hour until relieved.
 Gl. 4, p. 158

Urticaria:

Long Standing:

℞ Sp. Med. Belladonna gtt. v.
 Sp. Med. Pulsatilla gtt. x.
 Magnesium Sulph. ʒss.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every two hours.
 Gl. 7, p. 266

Venereal Warts:

℞ Ac. Salicylic gr. xxx.
 Sp. Med. Thuja ʒj. M. Ft. Sol.
Sig: Apply to warts with camel's hair pencil.
 Gl. 17, p. 590

EAR AND UPPER RESPIRATORY SYSTEM

Acute Coryza—Abortive:

℞ Sp. Med. Gelsemium
Sig: Fifteen drops at 8 P. M. and at 10 P. M.
 E. G. Sharp, M.D.

Acute Rhinitis:

℞ Sp. Med. Belladonna gtt. x.
 Sp. Med. Euphrasia ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every two hours to effect.
 Wm. P. Best, M.D.

Acute Laryngitis—Hoarseness:

℞ Sp. Med. Collinsonia ʒj.
 Sp. Med. Stillingia ʒiij.
 Glycerin ʒiv.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every two hours.
 T. D. Adlerman, M.D.

Acute Laryngitis—Internally:

℞ Sp. Med. Belladonna gtt. v.
 Sp. Med. Sanguinaria gtt. x.
 Elix. Aromat. q.s. ʒiv. M.
Sig: A teaspoonful every hour.

Acute Laryngitis—Inhalation:

℞ Tr. Benzoin Co. gtt. xx.
*Sig: Add to pint boiling water. Inhale through
 mouth until steaming of water stops.*
 A. W. Hobby, M.D.

Croupous Laryngitis:

℞ Sp. Med. Aconite gtt. v.
 Comp. Stillingia Lin. ʒss.
 Glycerin ʒj.
 Aq. Dest. q.s. ʒiij. M.
Sig: A teaspoonful every hour.
 Geo. C. Porter, M.D.

Croup:

℞ Comp. Stillingia Liniment
 Alcohol
 Glycerin āā. ʒj. M.
*Sig: Five to ten drops on sugar, repeated as re-
 quired.*
 Gl. 12, p. 424

Catarrhal Congestion, Nose and Throat:

℞ Sodium Borate gr. xxij.
 Ac. Salicylic gr. jss.
 Colorless Hydrastis,
 Lloyd's gtt. xxij.
 Aq. Dest. q.s. ʒj. M.
Sig: Use in atomizer three times a day.
*Patient instructed not to expose himself to dust
 or changes of temperature for one hour after using.*
This may be used also as nose drops.
 Gl. 32, p. 1043

Profuse Rhinorrhoea with Sneezing:

℞ Sp. Med. Belladonna gtt. iij.
 Sp. Med. Euphrasia ʒiij.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

Cloyce Wilson, M.D.

Acute Tonsillitis:

℞ Sp. Med. Aconite gtt. v.
 Sp. Med. Belladonna gtt. v.
 Sp. Med. Phytolacca gtt. lx.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours—children according to age.

Cloyce Wilson, M.D.

Acute Tonsillitis:

℞ Sp. Med. Aconite gtt. x.
 Sp. Med. Phytolacca gtt. xxx.
 Sp. Med. Macrotys ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Rudolph Wagner, M.D.

Tonsillitis—Pharyngitis, Uvulitis:

℞ Sp. Med. Apis gtt. v.
 Sp. Med. Phytolacca ʒss.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour:

Paint throat with Pinus Canadensis, red.

Cloyce Wilson, M.D.

Tonsillitis with exudate:

℞ Sp. Med. Echinacea ʒiij.
 Sp. Med. Baptisia ʒj.
 Sp. Med. Phytolacca gtt. xx.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

T. D. Hollingsworth, M.D.

Ulcerative Pharyngitis—Internally:

℞ Sp. Med. Aconite gtt. v.
 Sp. Med. Phytolacca ʒj.
 Sp. Med. Echinacea ʒij.
 Asepsin gr. x.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

J. M. Billman, M.D.

Ulcerative Pharyngitis. Locally:

Swab throat with:

℞ Sp. Med. Ac. Carbolic ʒj.
 Tr. Iodine ʒiv.
 Glycerin
 Aq. Dest. āā. q.s. ʒij. M. Ft. Sol.

J. M. Billman, M.D.

Hypertrophied Tonsils:

(A) Swab tonsils with Sp. Med. Thuja:

(B) Spray with:

℞	Sp. Med. Aconite	ʒi.
	Sp. Med. Phytolacca	ʒss.
	Sp. Med. Thuja	ʒij.
	Glycerin	ʒiij.
	Aq. Dest.	q.s. ʒiv. M.

*Sig: As throat spray three times a day.***Post Nasal Drip:**

℞	Sp. Med. Aconite	gtt. v.
	Sp. Med. Apis	gtt. x.
	Sp. Med. Ambrosia	ʒss.
	Aq. Dest.	q.s. ʒiij. M.

Sig: A teaspoonful every two hours.

W. W. Klement, M.D.

Acute Otitis Media:

℞	Sp. Med. Aconite	gtt. v.
	Sp. Med. Bryonia	gtt. v.
	Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every hour.

A. W. Hobby, M.D.

LOWER RESPIRATORY SYSTEM**Asthma—Sensitization:**

℞	Sp. Med. Belladonna	gtt. v.
	Sp. Med. Lobelia	ʒj.
	Glyconda	q.s. ʒiij. M.

Sig: A teaspoonful every hour until relieved.

Gl. 39, p. 1249

Catarrhal Asthma:

℞	Sp. Med. Sanguinaria	
	Sp. Med. Ipecac	āā. gtt. x.
	Sp. Med. Echinacea	ʒij.
	Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 42, p. 1329

Cardiac Asthma:

℞	Sp. Med. Cactus	
	Sp. Med. Lobelia	āā. ʒj.
	Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 42, p. 1330

Bronchial Asthma:

℞	Sp. Med. Lobelia	
	Sp. Med. Asthma Weed	āā. ʒj.
	Sp. Med. Gelsemium	ʒss.
	Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

Cloyce Wilson, M.D.

Bronchial Asthma:

Hypodermatic injections of Subculoyd Lobelia, gtt. x. to gtt. xxx. to cut short the paroxysms. Repeat every half hour to every three hours. Signs governing the dose:

The severity of the bronchial spasm.

The amount of dyspnea.

Short labored breathing with suffocation.

When paroxysms are lessened:

℞ Sp. Med. Lobelia gtt. xxx.
Sp. Med. Asthma Weed gtt. lx.
Aq. Dest. q.s. ʒiij. M.

Sig: A teaspoonful every one, two or three hours to effect.

Gl. 31, p. 1009

Bronchial Asthma—Congestive:

℞ Sp. Med. Belladonna gtt. v.
Sp. Med. Lobelia gtt. xxx.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour to effect.

Gl. 36, p. 157

Bronchitis—Sub-acute:

℞ Sp. Med. Bryonia
Sp. Med. Ipecac āā. gtt. x.
Sp. Med. Aesculus
Sp. Med. Lobelia āā. ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Rudolph Wagner, M.D.

Bronchitis—Asthmatic Type:

℞ Sp. Med. Lobelia
Sp. Med. Grindelia āā. ʒj.
Sp. Med. Sanguinaria ʒss.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Gl. 38, p. 1201

Bronchitis—1. Retarded Secretion:

℞ Sp. Med. Asclepias ʒi. to ij.
Sp. Med. Bryonia gtt. xij.
Ammonium Chloride ʒij.
Syr. Ac. Citric ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

T. D. Adlerman, M.D.

Bronchitis—2. Constriction, Dyspnea:

℞ Sp. Med. Lobelia ʒj.
Sp. Med. Sanguinaria ʒss.
Sp. Med. Asclepias ʒj.
Syr. Pruni Virginiana q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

T. D. Adlerman, M.D.

Bronchitis—3. Dry, Spasmodic Cough:

℞ Sp. Med. Drosera ʒiss.
Sp. Med. Sticta ʒiss.
Sp. Med. Bryonia gtt. xij.
Syr. Tolu q.s. ʒiij. M.

Sig: A teaspoonful every three hours.

T. D. Adlerman, M.D.

Cough—Dry, Hacking:

℞ Sp. Med. Bryonia
 Sp. Med. Sanguinaria āā. gtt. xv.
 Glycerin ℥ss.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

E. P. Zeumer, M.D.

Influenza—Myalgia:

℞ Sp. Med. Bryonia
 Sp. Med. Gelsemium āā. gtt. xx.
 Sp. Med. Eupatorium
 Sp. Med. Macrotys āā. ℥j.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Rudolph Wagner, M.D.

Influenza—Cough:

℞ Sp. Med. Aconite gtt. v.
 Sp. Med. Lobelia
 Sp. Med. Macrotys
 Sp. Med. Echinacea āā. ℥j. M.

Sig: Ten drops in a teaspoonful water every hour.

W. W. Klement, M.D.

Tonic—Post-Influenza:

℞ Sp. Med. Nux Vom. gtt. vij.
 Sp. Med. Hydrastis ℥j.
 Lloyd's Iron ℥jss.
 Aq. Dest. q.s. ℥iij. M.

Sig: A teaspoonful before meals and at bedtime.

B. H. Nellans, M.D.

Hay Fever:

℞ Sp. Med. Pulsatilla ℥j.
 Sp. Med. Eupatorium ℥ij.
 Sp. Med. Euphrasia ℥iv.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

W. L. LeBoy, M.D.

Hay Fever:

(a)

℞ Sp. Med. Belladonna gtt. x.
 Sp. Med. Euphrasia ℥iij.
 Aq. Dest. q.s. ℥iij. M.

Sig: A teaspoonful every hour until dryness of mouth occurs, then every three hours.

(b) Felter Treatment:

1.

℞ Subculoyd Lobelia ℥ij.
 Spt. Ammonia Aromat. ℥ij.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

2.

℞ Quinine Sulph. ℥j.
 Aq. Dest. q.s. ℥iv. M.
 Ac. Hydrochloric q.s. to make a solution.

Sig: A teaspoonful every three hours, alternating with ℞ No. 1.

Gl. 31, p. 1008

La Grippe:

1. To increase elimination:

℞ Sp. Med. Rhus tox. gtt. xv.
 Sp. Med. Macrotys ʒij.
 Sp. Med. Apocynum ʒss.
 Elix. Aromat. ʒiv. M.

Sig: A teaspoonful every four hours.

2. Tonic:

℞ Sp. Med. Nux Vom. gtt. xx.
 Sp. Med. Berberis ʒij.
 Elix. Calisaya and Iron q.s. ʒiv. M.

Sig: A teaspoonful after meals and on retiring.

Gl. 10, p. 371, 372

Pertussis:

℞ Sp. Med. Belladonna gtt. v.
 Sp. Med. Lobelia
 Sp. Med. Drosera āā. gtt. xx.
 Sp. Med. Passiflora ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours for children four to eight years old.

W. L. LeBoy, M.D.

Pertussis:

℞ Gold and Sodium Chloride gr. j.
 Aq. Dest. q.s. ʒj. M. Ft. Sol.

Sig: Five to ten drops, according to age, every four hours.

W. N. Mundy, M.D.

Chronic Pleurisy:

℞ Sp. Med. Aesculus
 Sp. Med. Lobelia āā. ʒj.
 Syr. Simplex
 Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 5, p. 218

Pleurisy:

Hot dry skin:

℞ Sp. Med. Asclepias ʒi.
 Sp. Med. Bryonia gtt. x.
 Sp. Med. Gelsemium gtt. x.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Cloyce Wilson, M.D.

Acute Pleurisy:

Early stages:

℞ Sp. Med. Bryonia gtt. x.
 Sp. Med. Aconite gtt. v.
 Sp. Med. Lobelia gtt. xx.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

Cloyce Wilson, M.D.

Pleurodynia

Intercostal neuralgia:

℞ Sp. Med. Pulsatilla ʒi.
 Sp. Med. Asclepias ʒss.
 Sp. Med. Belladonna gtt. v.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Cloyce Wilson, M.D.

Pleurisy:

Irritative cough. Sore muscular structures:

℞ Sp. Med. Ipecac gtt. x.
 Sp. Med. Sanguinaria gtt. x.
 Sp. Med. Macrotys ʒi.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Cloyce Wilson, M.D.

ALIMENTARY TRACT**Acid Bowel Disorders:**

℞ Sp. Med. Apis gtt. xv.
 Sp. Med. Eryngium
 Sp. Med. Frag. Sumach āā. ʒj.
 Glyconda
 Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful every two hours with a glass of water.

Gl. 41, p. 1296

Anorexia:

℞ Sp. Med. Nux Vom. gtt. x.
 Sp. Med. Hydrastis ʒj.
 Sp. Med. Piper Meth. ʒij.
 Glyconda q.s. ʒiv. M.

Sig: A teaspoonful before each meal.

Rudolph Wagner, M.D.

Atonic Gastritis—Gastralgia:

℞ Sp. Med. Nux Vom. gtt. xv.
 Sp. Med. Belladonna gtt. x.
 Sp. Med. Capsicum gtt. v.
 Aq. Dest. q.s. ʒiii. M.

Sig: A teaspoonful every hour.

Geo. C. Porter, M.D.

Atony of Digestive Tract:

℞ Sp. Med. Nux Vom. gtt. x.
 Sp. Med. Hydrastis gtt. lx.
 Glyconda q.s. ℥iv. M.
Sig: A teaspoonful before meals and at bedtime.
 Gl. 36, p. 1162

Hepatic Insufficiency and Stasis of Colon:

(Gaseous Distension):

℞ Sp. Med. Chionanthus ℥ij.
 Sp. Med. Podophyllum gtt. xx.
 Sp. Med. Cascara ℥iv.
 Syr. Simplex q.s. ℥iv. M.
Sig: A teaspoonful three times a day.
 Gl. 7, p. 277

Chronic Colitis and Sigmoiditis:

℞ Sp. Med. Geranium ℥iv.
 Sp. Med. Krameria gtt. xxx.
 Glyconda q.s. ℥iv. M.
Sig: A teaspoonful every two to four hours.
 Gl. 32, p. 1037

Chronic Colitis:

℞ Sp. Med. Nux Vom. gtt. xv.
 Sp. Med. Dioscorea ℥j.
 Sp. Med. Geranium ℥iv.
 Glyconda q.s. ℥iv. M.
*Sig: A teaspoonful every three hours to effect, then
 before meals and at bedtime.*
*Locally to rectum, 20% Solution Krameria by
 proctoscope, on alternate days.*
 Gl. 33, p. 1070

Colitis—Acute:

℞ Sp. Med. Aconite gtt. v.
 Sp. Med. Ipecac gtt. x.
 Sp. Med. Colocynth gtt. v.
 Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every two hours.
 Gl. 33, p. 1082

Colitis Subacute or Chronic:

℞ Sp. Med. Geranium ℥iv.
 Glyconda q.s. ℥iv. M.
Sig: A teaspoonful every four hours.
 Gl. 33, p. 1082

Mucous Colitis:

℞ Sp. Med. Geranium ℥ij.
 Dest. Hamamelis ℥ij.
 Elix. Lact. Pepsin q.s. ℥iv. M.
Sig: A teaspoonful every three hours.
 Gl. 17, p. 585

Acute Catarrhal Cholecystitis:

℞ Sp. Med. Chionanthus ℥j.
 Podophyllum Laxative,
 DeWees' ℥ss.
 Glyconda q.s. ℥iv. M.
*Sig: A teaspoonful an hour after meals and at bed-
 time.*
 Gl. 39, p. 1241

Cholera Morbus:

℞ Sp. Med. Dioscorea ʒij.
 Sp. Med. Colocynth gtt. v.
 Glyconda ʒii.
 Glycerin ʒj.
 Aq. Dest. q.s. ʒiv. M.

*Sig: A teaspoonful every half hour for four doses.
 then a teaspoonful every four hours.*

Gl. 16, p. 559

Cholecystitis:

Chronic, non-operative:

℞ Sp. Med. Chelidonium ʒj.

Sig: Ten drops in water every three hours.

P. A. DeOgny, M.D.

Cholecystitis:

Spasmodic Hepatic Pain:

℞ Sp. Med. Chionanthus ʒij.
 Sp. Med. Dioscorea ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every half-hour to two hours.

W. P. Best, M.D.

Colic:

℞ Sp. Med. Dioscorea

*Sig: A teaspoonful in one-quarter glass hot water
 every fifteen minutes for three doses.*

P. A. DeOgny, M.D.

Colic:

Infantile:

℞ Sp. Med. Colocynth, gtt. v.
 1/10 strength
 Liq. Taka-diastrase ʒj.
 Glyconda ʒss.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every half hour to one hour.

C. W. Beaman, M.D.

Colitis:

℞ Sp. Med. Chionanthus ʒj.
 Podophyllum Laxative,
 Dewees' ʒj.
 Glyconda q.s. ʒiv. M.

Sig: A teaspoonful after meals.

Gl. 41, p. 1307

Congestive Headache:

Intestinal Origin:

℞ Sp. Med. Nux Vom. gtt. v.
 Aq. Dest. q.s. ʒij. M.

Sig: A teaspoonful every two hours.

B. H. Nellans, M.D.

Constipation from Atony:

℞ Sp. Med. Chionanthus ʒj.
 Sp. Med. Nux Vom. gtt. x.
 Podophyllum Laxative,
 Dewees'
 Glyconda āā. q.s. ʒiv. M.

Sig: A teaspoonful an hour after each meal.

Gl. 41, p. 1291

Constipation from Atony:

℞ Sp. Med. Nux Vom. gtt. x.
 Podophyllum Laxative,
 Dewees' ʒj.
 Glyconda q.s. ʒiv. M.

Sig: A teaspoonful every four hours to effect, then twice daily.

Gl. 39, p. 1232

Constipation (Glandular Deficiency):

℞ Sp. Med. Chionanthus ʒiij.
 Sp. Med. Nux Vom. gtt. xxx.
 Sp. Med. Leptandra ʒij.
 Glyconda ʒiij.
 Aq. Dest. q.s. ʒvi. M.

Sig: A teaspoonful every four hours.

Gl. 20, p. 723

Diarrhoea:**Atonic:**

℞ Sp. Med. Nux Vom. gtt. v.
 Sp. Med. Geranium ʒij.
 Aq. Dest. q.s. ʒij. M.

Sig: A teaspoonful every hour until relieved, then every three hours.

B. H. Nellans, M.D.

Diarrhoea:**(Irritation):**

℞ Sp. Med. Aconite gtt. ij.
 Sp. Med. Ipecac gtt. v.
 Aq. Dest. q.s. ʒij. M.

Sig: A teaspoonful every hour to effect, then every two hours.

B. H. Nellans, M.D.

Diarrhoea—Tenesmus:

℞ Tr. Opii Camph. ʒij.
 Sp. Med. Colocynth gtt. v.
 Sp. Med. Dioscorea gtt. x.
 Glyconda q.s. ʒiii. M.

Sig: A teaspoonful every hour to effect.

B. Billman, M.D.

Sialogogue:

℞ Sp. Med. Phytolacca ʒj.
 Sp. Med. Jaborandi gtt. xv.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful four times a day.

Cloyce Wilson, M.D.

Dry Tongue and Mouth with Anorexia:

℞ Sp. Med. Nux Vom. gtt. v.
 Sp. Med. Lobelia ʒj.
 Syr. Simplex q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Cloyce Wilson, M.D.

Dysentery:

- ℞ Sp. Med. Aconite
 Sp. Med. Colocynth āā. gtt. v.
 Aq. Dest. q.s. ℥iij. M.

Sig: A teaspoonful every fifteen minutes for first two hours, every half hour for second two hours, and every hour thereafter. Restrict fluids.

W. W. Klement, M.D.

Gallstone Colic:

- ℞ Sp. Med. Chionanthus ℥ij.
 Sp. Med. Leptandra ℥j.
 Elix. Lact. Pepsin q.s. ℥iv. M.

Sig: A teaspoonful every two to four hours.

Gl. 5, p. 202

Gall Stones Diathesis:

To improve function of liver and its appendages:

- ℞ Sp. Med. Chionanthus ℥ij.
 Sp. Med. Leptandra ℥j.
 Sp. Med. Podophyllum gtt. xx.
 Elix. Simplex q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Gl. 2, p. 26

Flatulent Dyspepsia:

- ℞ Sp. Med. Nux Vom. gtt. x.
 Sp. Med. Hydrastis ℥ss.
 Elix. Lact. Pepsin q.s. ℥iij. M.

Sig: A teaspoonful one half hour after each meal.

Gl. 41, p. 1299

Intestinal Indigestion:**Gaseous Fermentation:**

- ℞ Lloyd's Hydrastis ℥ss.
 Sp. Med. Nux Vom. gtt. xv.
 Sp. Med. Xanthoxylum ℥j.
 Glyconda ℥j.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two to four hours.

Cloyce Wilson, M.D.

Indigestion, Flatulence:

- ℞ Sp. Med. Nux Vom. gtt. x.
 Sp. Med. Hydrastis ℥j.
 Glyconda q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Gl. 39, p. 1251

Gastric Disorders in the Aged:

- ℞ Sp. Med. Amygdalus
 Sp. Med. Apocynum āā. gtt. xv.—xx.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Gl. 28, p. 934

Gastric Hyperacidity:

- ℞ Sp. Med. Belladonna gtt. v.
 Sp. Med. Nux Vom. gtt. x.
 Glyconda q.s. ℥iv. M.

Sig: A teaspoonful after each meal.

Gl. 41, p. 1293

Hemorrhoids (Internal Bleeding):

℞ Sp. Med. Collinsonia gtt. x.
 Sp. Med. Aesculus ʒij.
 Aq. Dest. q.s. ʒiij. M.
Sig: A teaspoonful every hour, as needed.
 B. Billman, M.D.

Hemorrhoids (Internal Medication):

℞ Sp. Med. Collinsonia ʒiij.
 Sp. Med. Cascara ʒiij.
 Dist. Hamamelis ʒij.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.
 Gl. 8, p. 319

Intestinal Intoxication:

℞ Sp. Med. Dioscorea ʒij.
 Sp. Med. Collinsonia ʒij.
 Sp. Med. Bryonia gtt. x.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.
 Gl. 30, p. 995

Stomatitis:

℞ Sp. Med. Aconite gtt. v.
 Sp. Med. Phytolacca gtt. x.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.
 Gl. 16, p. 547

Indigestion and Heartburn:

℞ Sp. Med. Nux Vom. gtt. xv.
 Sp. Med. Pulsatilla ʒj.
 Lloyd's Iron ʒij.
 Aq. Dest. q.s. ʒiv. M.
*Sig: A teaspoonful one half hour before meals
 and at bedtime.*

Cloyce Wilson, M.D.

Jaundice:

Congestion of Liver:
 ℞ Sp. Med. Chionanthus
 Sp. Med. Chelidonium āā. ʒij.
 Fl. Ext. Cascara Arom. ʒss.
 Glyconda q.s. ʒiv. M.
Sig: A teaspoonful every two hours.

J. M. Billman, M.D.

Jaundice:

Catarrhal:
 ℞ Sp. Med. Nux Vom. gtt. v.
 Sp. Med. Chionanthus ʒj.
 Aq. Dest. q.s. ʒij. M.
Sig: A teaspoonful every three hours.
*Sodium Phosphate as a laxative in conjunction
 with above.*

B. H. Nellans, M.D.

Liver:

Fullness, Torpor:

℞ Sp. Med. Hydrastis	ʒjss.
Sp. Med. Hyoscyamus	ʒijss.
Sp. Med. Chionanthus	q.s. ʒj. M.

Sig: Five drops in water after meals.

E. P. Zeumer, M.D.

Liver—Enlargement:

Congestion:

℞ Sp. Med. Dioscorea	ʒiij.
Sp. Med. Chionanthus	ʒiv.
Glyconda	q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 7, p. 267

Liver:

Podophyllum Leptandra Compound:

℞ Res. Podophyllum (podophyllin)	gr. xxx.
Res. Leptandra (leptandrin)	gr. lx.
Powd. Capsicum	gr. x.
Sodium bicarbonate	gr. c. M. Trit.
Div. in Caps. No. 40	

*Sig: One at bedtime.**Indications: Full broad tongue, coated at the base, yellowish. Fullness of tissue. Soreness under right scapula. Dizziness with constipation. Full veins.*

Gl. 4, p. 142

O. S. Coffin, M.D.

Hypoacidity (Low Gastric Acidity):

℞ Sp. Med. Nux Vom.	gtt. x.
Sp. Med. Hydrastis	ʒss.
Sp. Med. Chionanthus	ʒj.
Elix. Simplex	q.s. ʒiij. M.

Sig: A teaspoonful one half hour after meals and at bedtime.

Gl. 41, p. 1308

Alcoholic Gastritis:

℞ Sp. Med. Nux Vom.	gtt. viij.
Sp. Med. Hydrastis	ʒj.
Tr. Capsicum	gtt. xv.—xxx.
Aq. Dest.	q.s. ʒiij. M.

Sig: A teaspoonful every two hours to effect, then every four hours.

B. H. Nellans, M.D.

Persistent Vomiting:

℞ Sp. Med. Amygdalus	ʒj.
Aq. Dest.	q.s. ʒiv. M.

Sig: Teaspoonful every half hour until relieved.

Gl. 37, p. 1196

Sigmoid:

Chronic Atony:

℞ Sp. Med. Gelsemium	ʒj.
Sp. Med. Polytrichum	ʒiij.
Elix. Lactated Pepsin	q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

P. A. DeOgny, M.D.

Splenic Enlargement:

℞ Sp. Med. Polymnia ʒi.
 Sp. Med. Ceanothus ʒij.
 Elix. Aromatic q.s. ʒiv. M.
Sig: A teaspoonful every three hours.
 Gl. 5, p. 196

Stomachic Tonic:

Children:
 ℞ Sp. Med. Nux Vom. gtt. ij.
 Lloyd's Iron gtt. xxx.
 Glycerin ʒij.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful before meals and at bedtime.
 C. W. Beaman, M.D.

Wrongs of Digestion and Assimilation in Aged:

℞ Sp. Med. Xanthoxylum ʒij.
 Sp. Med. Ipecac gtt. x.
 Glyconda ʒss.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.
 Gl. 7, p. 259

CARDIO VASCULAR SYSTEM**Angina Pectoris:**

℞ Sp. Med. Cactus ʒij.
 Sp. Med. Crataegus ʒss.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours, or oftener.
 Gl. 29, p. 954

Cardiac Dilatation:

℞ Sp. Med. Cactus ʒj.
 Sp. Med. Crataegus ʒij.
 Lloyd's Iron ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.
 Gl. 30, p. 982

Palpitation of Heart (Menopause):

℞ Sp. Med. Cactus ʒj.
 Sp. Med. Pulsatilla āā. ʒj.
 Sp. Med. Ignatia gtt. x.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful as needed.
 Gl. 32, p. 1033

Phlebitis with Venous Stasis:

℞ Sp. Med. Aconite ʒj.
 Sp. Med. Belladonna āā. gtt. v.
 Sp. Med. Arnica ʒss.
 Glyconda q.s. ʒiv. M.
Sig: A teaspoonful every four hours.
 Gl. 39, p. 1252

Tachycardia (Neurosis):

℞ Sp. Med. Cactus ʒjss.
 Sp. Med. Lobelia ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 39, p. 1243

Angioneurotic Edema:

(Bold Hives):

℞ Sp. Med. Apis gtt. xxxij.
 Sp. Med. Berberis ʒiv.
 Sp. Med. Rhus tox. gtt. x.
 Aq. Dest. q.s. ʒiij. M.

Sig: A teaspoonful every two hours.

T. D. Adlerman, M.D.

Cardiac Distress with Flatulence:

℞ Sp. Med. Cactus ʒj.
 Spt. Vini Rect. ʒiij.
 Glyconda ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every ten minutes until relieved.

O. C. Welbourn, M.D.

Cardiac Neurosis:

Heart Pains:

℞ Sp. Med. Bryonia gtt. x.
 Sp. Med. Cactus ʒij.
 Glyconda
 Aq. Dest. āā. q.s. ʒij. M.

Sig: A teaspoonful after meals and at bedtime.

Gl. 41, p. 1285

Congestive Chills:

℞ Tr. Capsicum and Myrrh ʒj.

Sig: Ten drops in one ounce hot water. Repeat at half-hour intervals if required.

Rudolph Wagner, M.D.

Hemorrhage:

Passive:

℞ Oil Cinnamon
 Oil Erigeron āā. ʒj.
 Alcohol q.s. ʒj. M.

Sig: Twenty drops, repeated as needed.

J. M. Billman, M.D.

Heart—Tonic:

℞ Sp. Med. Cactus
 Sp. Med. Crataegus
 Sp. Med. Hyoscyamus āā. q.s. ʒj. M.

Sig: Fifteen drops in water three times a day.

P. K. Morse, M.D.

Heart Symptoms in Hysteria:

℞ Sp. Med. Cactus
 Sp. Med. Pulsatilla āā. ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 34, p. 1106

Cardiac Dropsy:

℞ Sp. Med. Apocynum ʒj. to ʒij.
 Potassium Acetate ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

In addition, fifteen drops Sp. Med. Crataegus four times a day.

Gl. 34, p. 1095

Hypertension:

Pulse, full and bounding:

℞ Sp. Med. Gelsemium ʒj.
 Sp. Med. Veratrum gtt. xx.
 Aq. Dest. q.s. ʒiv. M.
 Wm. P. Best, M.D.

Hypertension—Nervous Symptoms:

℞ Sp. Med. Gelsemium gtt. xxx.
 Glyconda q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 40, p. 1276

Hypotension (Low Blood Pressure):

℞ Sp. Med. Nux Vom. gtt. xv.
 Sp. Med. Belladonna gtt. x.
 Lloyd's Iron ʒiij.
 Aq. Dest. q.s. ʒiij. M.

Sig: A teaspoonful before meals and at bedtime.

Gl. 40, p. 1267

Mitral Regurgitation:

℞ Sp. Med. Apocynum ʒj.
 Sp. Med. Echinacea āā. ʒiv.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 41, p. 1262

Mitral Insufficiency with Tachycardia:

℞ Sp. Med. Cactus ʒj.
 Sp. Med. Aconite gtt. iij.
 Glyconda ʒij.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two to four hours.

In high arterial tension, use Sp. Med. Veratrum gtt. x, instead of Aconite: if highly nervous use Sp. Med. Gelsemium, gtt. x, instead of either Aconite or Veratrum.

Gl. 10, p. 366

Cardio-Renal Insufficiency with Edema:

℞ Sp. Med. Apocynum ʒj. to ʒiij.
 Glyconda
 Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 39, p. 1242

Valvular Insufficiency:

(Dyspnea)

℞ Sp. Med. Apocynum ʒiij.
 Sp. Med. Cactus ʒij.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Gl. 6, p. 245

NERVOUS SYSTEM

Alcohol Addiction:

℞ Sp. Med. Black Haw ʒijss.
 Sp. Med. Capsicum gtt. v.
 Glyconda q.s. ʒij. M.

Sig: Teaspoonful every four hours.

Gl. 41, p. 1287

Convulsions in Children:

℞ Sp. Med. Gelsemium
 Sp. Med. Lobelia
 Potassium Bromide āā. ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour or half hour.

H. W. Felter, M.D.

℞ Sp. Med. Lobelia
 Sp. Med. Gelsemium āā. ʒj.
 Sp. Med. Solanum ʒss.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every five minutes for six doses, then every two hours for the day.

Gl. 35, p. 1120

Delirium:

(Febrile):

℞ Sp. Med. Jaborandi ʒij.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour until quieted.

John J. Sutter, M.D.

Epilepsy:

℞ Sp. Med. Gelsemium
 Sp. Med. Oenanthe Croc. āā. ʒj.
 Sp. Med. Conium Mac. ʒij.
 Ess. Pepsin q.s. ʒiv. M.
Sig: A teaspoonful every three hours.

Gl. 3, p. 7

Hiccough:

1. ℞ Sp. Med. Lobelia ʒjss.
 Emuls. Turpentine q.s. ʒiv. M.
Sig: A teaspoonful in water every four hours.

2. ℞ Sp. Med. Lobelia ʒiv.
 Aq. Dest. ʒjss. M.
Sig: A teaspoonful every three hours.

Cloyce Wilson, M.D.

Insomnia:

℞ Sp. Med. Gelsemium gtt. xx.
 Sp. Med. Passiflora ʒij.
 Glyconda
 Aq. Dest. āā. q.s. ʒiv. M.
*Sig: A teaspoonful an hour after meals and a
 tablespoonful at bedtime.*

Gl. 41, p. 1299

Insomnia:

(Alcoholism) :

℞	Sp. Med. Passiflora	ʒij.
	Sp. Med. Capsicum	gtt. x.
	Sp. Med. Zingiber	
	Sp. Med. Lupulin	āā. ʒij.
	Elixir Peptenzyme	q.s. ʒiij. M.

Sig: A teaspoonful every hour to effect.

T. D. Adlerman, M.D.

Insomnia:

(Worry) :

℞	Sp. Med. Passiflora	ʒjss.
	Sp. Med. Avena	ʒiij.
	Sp. Med. Hyoscyamus	ʒj.
	Glyconda	q.s. ʒiij. M.

Sig: A teaspoonful every two hours, from 6 P.M. until bedtime.

T. D. Adlerman, M.D.

Migraine:

℞	Sp. Med. Belladonna	gtt. v.
	Sp. Med. Bryonia	gtt. x.
	Sp. Med. Macrotys	ʒj.
	Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every three hours or oftener.

Gl. 15, p. 533

Morphine Substitute:

℞	Sp. Med. Gelsemium	ʒij.
	Sp. Med. Passiflora	ʒss.
	Potassium Bromide	ʒij.
	Elix. Aromat.	q.s. ʒiv. M.

Sig: A teaspoonful. Repeat in one to four hours, if needed.

Gl. 9, p. 344

Mountain Sickness:

℞	Sp. Med. Lobelia	gtt. x.
	Sp. Med. Gelsemium	gtt. xxx.
	Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful as required.

Gl. 28, p. 934

Neuralgia:

Facial :

℞	Sp. Med. Gelsemium	gtt. xxxv.
	Sp. Med. Rhus tox	gtt. x.
	Sp. Med. Plantago	ʒjss.
	Aq. Dest.	q.s. ʒiij. M.

Sig: A teaspoonful as needed according to case.

T. D. Adlerman, M.D.

Neuralgia:

Periodic Trifacial :

℞	Sp. Med. Cannabis	gtt. xx.
	Sp. Med. Piscidia	
	Sp. Med. Cinchona	āā. ʒj.
	Glyconda	q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Rudolph Wagner, M.D.

Neuritis:

℞ Sp. Med. Rhus tox. gtt. x.
 Sp. Med. Bryonia gtt. v.
 Sp. Med. Aconite gtt. iii.
 Elix. Lact. Pepsin q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

Gl. 4, p. 159

Neuritis:

℞ Sp. Med. Apis
 Sp. Med. Bryonia
 Sp. Med. Rhus tox. āā. gtt. x.
 Sp. Med. Gelsemium gtt. xxx.
 Glyconda ℥j.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

F. L. Thomas, M.D.

Nerve Sedative:

℞ Sp. Med. Hyoscyamus ℥j.
 Strontium Bromide ℥ss.
 Glycerin ℥ss.
 Glyconda ℥ij.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

G. W. DeMuth, M.D.

Neurasthenia:

℞ Sp. Med. Cannabis gtt. xx.
 Sp. Med. Pulsatilla ℥ss.
 Sp. Med. Matricaria ℥j.
 Sp. Med. Passiflora ℥ij.
 Sp. Med. Avena ℥iv.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Rudolph Wagner, M.D.

GENITO-URINARY SYSTEM

Cystitis:

(Acute, with frequent and painful urination):

℞ Sp. Med. Belladonna gtt. x.
Sp. Med. Eryngium
Sp. Med. Erigeron āā. ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful three times a day.

Rudolph Wagner, M.D.

Cystitis:

(Excessive burning):

℞ Sp. Med. Elaterium gtt. x.
Sp. Med. Gelsemium gtt. xv.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour. Patient to lie in bed and drink copious amounts of water.

T. D. Hollingsworth, M.D.

Cystitis:

(Sub acute):

℞ Sp. Med. Belladonna gtt. x.
Sp. Med. Fragrant Sumach ʒj.
Glyconda
Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful after meals and at bedtime.

Gl. 42, p. 1319

Cystitis:

(Chronic):

℞ Sp. Med. Cannabis ʒss.
Sp. Med. Fragrant Sumach ʒiij.
Sp. Med. Saw Palmetto ʒij.
Elix. Simplex q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Gl. 2, p. 15

Cystitis:

(Atony)

℞ Sp. Med. Gelsemium gtt. xxv.
Sp. Med. Piper Meth. ʒjss.
Sp. Med. Saw Palmetto ʒij.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

P. R. Tindall, M.D.

Epididymitis:

(Orchitis):

℞ Sp. Med. Aconite gtt. x.
Sp. Med. Belladonna gtt. x.
Sp. Med. Phytolacca ʒj.
Sp. Med. Pulsatilla ʒss.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Libradol applied externally.

Rudolph Wagner, M.D.

Incontinence Urine (Aged):

1. ℞ Sp. Med. Fragrant Sumach ʒj. to ʒiij.
Glyconda
Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 41, p. 1292

2. ℞ Sp. Med. Fragrant Sumach
Sp. Med. Thuja āā. ʒij.
Sp. Med. Cannabis gtt. xx.
Elix. Simplex q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Children for nocturnal incontinence, according to age.

Gl. 1, p. 13

Incontinence of Urine:

- ℞ Sp. Med. Belladonna gtt. x.
Sp. Med. Fragrant Sumach ʒj.
Glyconda
Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 42, p. 1333

Nocturnal Enuresis:

(Children):

- ℞ Sp. Med. Fragrant Sumach ʒj.
Aq. Dest. q.s. ʒiij. M.

Sig: A teaspoonful four times a day, last dose at bedtime.

Restrict fluids after 4 P. M.

B. H. Nellans, M.D.

Nocturnal Enuresis:

(Prostatic origin):

- ℞ Sp. Med. Belladonna gtt. v.
Sp. Med. Thuja ʒj.
Sp. Med. Saw Palmetto ʒiij.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

B. Billman, M.D.

Scalding Urine:

- ℞ Sp. Med. Cantharis
Sp. Med. Apis āā. gtt. v.
Sp. Med. Eryngium ʒij.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour.

Cloyce Wilson, M.D.

Tenesmus, Strangury, Urgency:

Frequency of Urination:

- ℞ Sp. Med. Apis gtt. x.
Sp. Med. Gelsemium gtt. xxx.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every one to two hours.

A. Harry Crum, M.D.

Injection for Gonorrhoea (Sub Acute) :

1. ℞ Zinc Sulphate gr. v.
Lloyd's Hydrastis ℥ij.
Glycerin ℥ij.
Aq. Dest. q.s. ℥iv. M.

Sig: Use as injection three times a day.

More astringent, add Tannic acid in place of Zinc Sulphate.

W. B. Church, M.D.

Internal Medication for Gonorrhoea :

2. ℞ Sp. Med. Staphisagria ℥ij.
Sp. Med. Pulsatilla ℥j.
Elix. Aromatic q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Gl. 6, p. 228

Painful Urination (Gonorrhoea) :

- ℞ Sp. Med. Apis gtt. v.
Sp. Med. Eryngium ℥j.
Aq. Dest. q.s. ℥iij. M.

Sig: A teaspoonful every three hours.

Gl. 33, p. 1078

Acute Prostatitis :

- ℞ Sp. Med. Staphisagria ℥j.
Sp. Med. Gelsemium ℥ss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Gl. 34, p. 1091

Enlarged Prostate :

- ℞ Sp. Med. Gaultheria ℥jss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Gl. 36, p. 1145

Enlarged Prostate with Hemorrhoids :

- ℞ Sp. Med. Collinsonia
Sp. Med. Staphisagria āā. ℥j.
Glyconda
Aq. Dest. āā. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Gl. 41, p. 1286

Enlargement Prostate with Irritable Bladder :

- ℞ Sp. Med. Fragrant Sumach ℥ij.
Sp. Med. Staphisagria ℥j.
Sp. Med. Saw Palmetto ℥iij.
Elix. Simplex q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Gl. 7, p. 269

Chronic Prostatitis :

- ℞ Sp. Med. Fragrant Sumach
Sp. Med. Staphisagria āā. ℥j.
Glyconda q.s. ℥iij. M.

Sig: A teaspoonful every four hours.

Gl. 40, p. 1279

Chronic Prostatitis with Cystitis:

℞ Sp. Med. Belladonna gtt. x.
 Sp. Med. Thuja ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.
 Gl. 37, p. 1193

Prostatic Hypertrophy:

℞ Sp. Med. Saw Palmetto ʒvj.
 Sp. Med. Staphisagria
 Sp. Med. Ergot āā. ʒij.
 Sp. Med. Piper Meth. ʒv.
 Elix. Lact. Pepsin q.s. ʒiv. M.
Sig: A teaspoonful after each meal.
 W. L. LeBoy, M.D.

Prostate:

(Irritation, sexual hyperesthesia):

℞ Sp. Med. Staphisagria ʒj.
 Sp. Med. Salix Nigra Aments ʒiv.
 Sp. Med. Saw Palmetto ʒj.
 Glyconda q.s. ʒiv. M.
Sig: A teaspoonful four times a day.
 E. P. Zeumer, M.D.

Prostate:

(Enlargement with irritation):

℞ Sp. Med. Gaultheria ʒj.
 Sp. Med. Phytolacca ʒij.
 Sp. Med. Macrotys ʒj.
 Glyconda
 Aq. Dest. q.s. ʒvj. M.
Sig: A teaspoonful four times a day.
 P. K. Morse, M.D.

Retention of Urine:

℞ Sp. Med. Gelsemium ʒss.
 Sp. Med. Conium gtt. xx.
 Sp. Med. Fragrant Sumach ʒij.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.
 Gl. 3, p. 68

Retention of Urine:

(Vegetable Catheter, Old Men):

℞ Sp. Med. Staphisagria gtt. xxx.
 Sp. Med. Gelsemium ʒss.
 Sp. Med. Gaultheria q.s. ʒj. M.
Sig: Five drops every three hours.
 P. A. DeOgny, M.D.

Spermatorrhoea:

Emissions:

℞ Sp. Med. Oenanthe gtt. v.
 Sp. Med. Cannabis gtt. xx.
 Sp. Med. Pulsatilla gtt. xxx.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful before meals and at bedtime.
 Rudolph Wagner, M.D.

REPRODUCTIVE SYSTEM—FEMALE

Abortion:

℞ Codein Sulph. gr. ij.
 Sp. Med. Black Haw ʒss.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours until pain is relieved, then as required.

W. N. Mundy, M.D.

Amenorrhoea:

(Suppression):

℞ Sp. Med. Macrotys
 Sp. Med. Black Haw āā. ʒij.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

Gl. 10, p. 368

Delayed Menstruation:

Functional:

℞ Sp. Med. Macrotys gtt. xxv.
 Sp. Med. Caulophyllum gtt. xlv.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

P. R. Tindall, M.D.

Irregular Menstruation:

℞ Sp. Med. Pulsatilla ʒj.
 Sp. Med. Viburnum ʒss.
 Sp. Med. Macrotys gtt. xl.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful four times a day.

Gl. 38, p. 1220

Excessive Uterine Hemorrhage:

Flooding:

℞ Sp. Med. Capsella
 Sp. Med. Geranium āā. ʒiv.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour until checked, then every four hours.

Gl. 20, p. 708

Uterine Hemorrhage:

Menopause:

℞ Sp. Med. Geranium ʒiv.
 Sp. Med. Gossypium gtt. xx.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Study also Capsella, Ceanothus, Geranium.

Cloyce Wilson, M.D.

Amenorrhoea and Dysmenorrhoea:

℞ Sp. Med. Macrotys ʒj.
 Sp. Med. Pulsatilla ʒss.
 Sp. Med. Helonias ʒj.
 Aq. Dest. q.s. ʒiij. M.

Sig: A teaspoonful every two hours.

Geo. C. Porter, M.D.

Dysmenorrhoea :

℞ Sp. Med. Dioscorea
 Sp. Med. Viburnum āā. ʒj.
 Sp. Med. Tiger Lily ʒij.
 Glyconda q.s. ʒiv. M.

Sig: A teaspoonful every three hours, beginning three days before expected period. Repeat monthly.

Rudolph Wagner, M.D.

Dysmenorrhoea—Anemia :

℞ Sp. Med. Ignatia gtt. v.
 Sp. Med. Pulsatilla ʒj.
 Lloyd's Iron ʒij.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful before meals and at bedtime.

Cloyce Wilson, M.D.

Menorrhagia :

(a) Painful; two days before usual period :

℞ Sp. Med. Black Haw
 Sp. Med. Macrotys āā. ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

(b) When menstruation is established, if excessive :

℞ Sp. Med. Pulsatilla ʒj.
 Sp. Med. Hydrastis ʒss.
 Aq. Dest. q.s. ʒiij. M.

Sig: A teaspoonful every four hours.

Gl. 42, p. 1323

Menorrhagia :

℞ Sp. Med. Ergot
 Sp. Med. Erigeron āā. ʒj.
 Sp. Med. Capsella ʒiij.
 Aq. Cinnamomi q.s. ʒiv. M.

Sig: A teaspoonful every three hours, starting second day of menstruation.

Rudolph Wagner, M.D.

Menorrhagia

℞ Sp. Med. Black Haw ʒss.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

W. N. Mundy, M.D.

Metrorrhagia :

℞ Sp. Med. Belladonna gtt. x.
 Sp. Med. Geranium ʒij.
 Dist. Hamamelis
 Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful every half hour to three hours.

Gl. 8, p. 315

Dysmenorrhoea :

Functional :

℞ Sp. Med. Viburnum gtt. xxx.
 Sp. Med. Dioscorea ʒiss.
 Sp. Med. Ignatia. gtt. iv.
 Aq. Dest. q.s. ʒiij. M.

Sig: A teaspoonful every three hours.

B. H. Nellans, M.D.

Dysmenorrhoea:

(a) Congestive:

℞	Sp. Med. Belladonna	gtt. v.
	Sp. Med. Gossypium	gtt. x.
	Sp. Med. Macrotys	gtt. x.
	Aq. Dest.	q.s. ℥iv. M.

Sig: A teaspoonful every half hour until relieved.

(b) One week before expected period:

℞	Sp. Med. Black Haw	℥ij.
	Sp. Med. Macrotys	
	Sp. Med. Pulsatilla	āā. ℥ss.
	Aq. Dest.	q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Gl. 7, p. 260

Menopause:

Nervous Symptoms:

℞	Sp. Med. Gelsemium	gtt. xx.
	Sp. Med. Gossypium	gtt. xxx.
	Sp. Med. Pulsatilla	gtt. x.
	Aq. Dest.	q.s. ℥iv. M.

Sig: A teaspoonful every two to four hours.

Gl. 9, p. 345

Menopause:

Circulatory Disturbances:

℞	Sp. Med. Belladonna	gtt. v.
	Sp. Med. Ipecac.	gtt. x.
	Sp. Med. Lycopus	gtt. xx.
	Dist. Hamamelis	℥ij.
	Aq. Dest.	q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Gl. 7, p. 263

Vicarious Menstruation:

℞	Sp. Med. Senecio	℥j.
	Aq. Dest.	q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Gl. 33, p. 1068

Leucorrhoea:

(a) Internal Medication:

℞	Sp. Med. Pulsatilla	℥j.
	Sp. Med. Hydrastis	℥ss.
	Aq. Dest.	q.s. ℥iij. M.

Sig: A teaspoonful every four hours.

(b) Tampon:

℞	Pulv. Alum	gr. xx.
	Zinc Sulph.	gr. x.
	Lloyd's Hydrastis	℥ijss.
	Glycerin	q.s. ℥iv. M. Ft. Sol.

Sig: Use as vaginal tampon on alternate days.

(c) Suppositories:

℞	Hydrastine Hydrochloride	gr. j.
	Ichthyol	gr. v.
	Ol. Theobrom.	q.s.
	Ft. Suppos. No. j, D. T. D. No. xii.	

Sig: Insert at bedtime.

Gl. 39, p. 1240

Salpingitis:

Pain, Fallopian Tubes:

℞ Sp. Med. Colocynth gtt. x.
 Sp. Med. Dioscorea
 Sp. Med. Tiger Lily
 Sp. Med. Viburnum āā. ʒj.
 Glyconda q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Rudolph Wagner, M.D.

Eclampsia:

Subculoyd Veratrum hypodermatically, ten to twenty minims, every hour, increasing the dose as the case requires.

Gl. 2, p. 15

NEURO-MUSCULAR SYSTEM AND JOINTS

Articular Rheumatism:

(Chronic) :

℞ Sp. Med. Colchicum gtt. xx.
 Sp. Med. Bryonia gtt. x.
 Sp. Med. Piscidia
 Sp. Med. Eupatorium āā. ʒj.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful after each meal.

Rudolph Wagner, M.D.

Chronic Arthritis:

℞ Sp. Med. Colchicum
 Sp. Med. Bryonia āā. gtt. xxx.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 42, p. 1321

Chronic Rheumatoid Arthritis:

℞ Sp. Med. Phytolacca ʒj.
 Glyconda
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful three times a day.

Locally, Libradol applied at bedtime to remain all night.

Gl. 40, p. 1259

Rheumatoid Arthritis:

Acute:

℞ Sp. Med. Rhus tox.	gtt. x.
Sp. Med. Bryonia	gtt. xx.
Sp. Med. Eupatorium	ʒj.
Sp. Med. Echinacea	ʒv.
Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every hour.

W. L. LeBoy, M.D.

Rheumatism:

℞ Sp. Med. Phytolacca	ʒij.
Sp. Med. Macrotys	ʒss.
Sp. Med. Cactus	ʒj.
Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every two to four hours.

Gl. 10, p. 360

Ten grains Sodium Salicylate every two hours to saturation: follow with twenty grains sodium bicarbonate every four hours, both to be taken with large quantities of water.

Gl. 10, p. 361

Muscular Rheumatism:

Myalgia:

℞ Sp. Med. Bryonia	gtt. x.
Sp. Med. Macrotys	ʒj.
Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 7, p. 269

Locally to Inflamed Joints:

℞ Ol. Gaultheria	ʒiij.
Tr. Arnica	
Spt. Turpentine	āā. ʒj.
Lin. Saponis	q.s. ʒiv. M. Ft. Lin.

Sig: Apply without friction, cover with cotton.

Cloyce Wilson, M.D.

Pleurodynia:

℞ Sp. Med. Bryonia	gtt. x.
Sp. Med. Gelsemium	ʒj.
Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

Wm. P. Best, M.D.

Chorea:

℞ Sp. Med. Gelsemium	gtt. vj.
Sp. Med. Lobelia	
Sp. Med. Macrotys	āā. gtt. xv.
Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every two hours for children 4 to 8 years.

W. L. LeBoy, M.D.

Chorea:

℞ Sp. Med. Gelsemium	ʒss.
Sp. Med. Cypripedium	ʒij.
Sp. Med. Pulsatilla	ʒj.
Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every two to four hours.

Gl. 7, p. 274

Chorea:

℞ Sp. Med. Belladonna gtt. v.
 Sp. Med. Macrotys ʒij.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.
 Gl. 42, p. 1318

Chorea:

℞ Sp. Med. Solanum gtt. xxx.
 Sp. Med. Macrotys ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.
 Gl. 41, p. 1298

Lumbago:

℞ Sp. Med. Hydrangea ʒiv.
 Sp. Med. Macrotys ʒij.
 Potassium Acetate ʒij.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every two to four hours.
 W. N. Mundy, M.D.

Lumbago:

℞ Sp. Med. Bryonia gtt. x.
 Sp. Med. Macrotys ʒij.
 Sodium Salicylate ʒij.
 Elix. Simplex q.s. ʒiv. M.
Sig: A teaspoonful every four hours.
 Gl. 12, p. 431

Sciatica:

1. ℞ Sp. Med. Bryonia gtt. v.
 Sp. Med. Macrotys ʒss.
 Sp. Med. Gelsemium gtt. xv.
 Aq. Dest. q.s. ʒij. M.
Sig: A teaspoonful every two hours.
 Gl. 40, p. 1271
2. ℞ Sp. Med. Bryonia gtt. x.
 Sp. Med. Piscidia ʒij.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every hour until relieved.
3. ℞ Sp. Med. Macrotys
 Sp. Med. Apocynum āā. ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every two hours.
 Gl. 39, p. 1250
4. ℞ Fowler's Solution Arsenic
 Sp. Med. Gelsemium āā. ʒij.
 Sp. Med. Bryonia ʒss.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful in one-third glass water after each meal.
 Cloyce Wilson, M.D.

Torticollis:

℞ Sp. Med. Gelsemium gtt. xxx.
 Sp. Med. Macrotys ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every hour or three hours.
 Gl. 34, p. 1107

ALTERNATIVES AND TONICS

Alterative:

Blood Dyscrasia:

℞	Echafolta	ʒss.
	Sp. Med. Rumex	ʒss.
	Sp. Med. Iris	ʒj.
	Elix. Simplex	q.s. ʒvi. M.

Sig: A teaspoonful every three hours.

Gl. 4, p. 143

Alterative:

Chronic Eczema:

℞	Sp. Med. Phytolacca	
	Sp. Med. Corydalis	
	Sp. Med. Berberis	āā. ʒss.
	Syr. Simplex	q.s. ʒvj. M.

Sig: A teaspoonful three times a day.

Gl. 36, p. 1155

Alterative:

℞	Potassium Iodide	gr. xxx.
	Sp. Med. Stillingia	ʒiv.
	Sp. Med. Phytolacca	ʒij.
	Sp. Med. Rumex	ʒiv.
	Glyconda	q.s. ʒiv. M.

Sig: A teaspoonful three times a day.

Gl. 25, p. 841

Alterative:

Accessory in treatment of Syphilis:

℞	Sp. Med. Berberis	ʒiij.
	Potassium Iodide	ʒij.
	Glyconda	q.s. ʒiv. M.

Sig: A teaspoonful an hour after meals followed by a glass of water.

Gl. 41, p. 1297

Alterative:

Glandular Enlargement:

Acute, Subacute:

℞	Sp. Med. Iris	
	Sp. Med. Phytolacca	āā. ʒj.
	Podophyllum, Laxative	ʒss.
	Dewees'	
	Aq. Dest.	q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Gl. 42, p. 1331

Tonic and Alterative—Malaria:

(a)

- ℞ Sp. Med. Eupatorium
 Sp. Med. Cornus
 Lloyd's Hydrastis āā. ʒj.
 Glyconda
 Aq. Dest. āā. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

Gl. 25, p. 853

(b)

- ℞ Sp. Med. Chionanthus ʒiv.
 Quinine Bisulphate ʒij.
 Elix. Lactated Pepsin q.s. ʒiv. M.

*Sig: A teaspoonful every three hours from 6 A.M.
 to 6 P.M.*

Gl. 25, p. 854

(c)

- ℞ Quinine Sulph. gr. xxx.
 Hydrastin Phos. gr. x.
 Pulv. Capsicum gr. v. M.
 Div. in Caps. no. vj.

Sig: A capsule every two hours for three doses.

Gl. 20, p. 733

Alterative:

Pus:

- ℞ Sp. Med. Echinacea ʒiv.
 Sp. Med. Baptisia ʒij.
 Glyconda q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Gl. 39, p. 1238

Anemia:

(Simple Tonic):

- ℞ Sp. Med. Echinacea
 Sp. Med. Berberis āā. ʒj.
 Lloyd's Iron ʒij.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful before meals and at bedtime.

Gl. 39, p. 1231

Asthenia:

- ℞ Sp. Med. Nux Vom. gtt. x.
 Sp. Med. Belladonna gtt. v.
 Lloyd's Iron ʒjss.
 Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful before meals and at bedtime.

Gl. 39, p. 1246

Diabetes:

(a) Excessive flow of Urine:

- ℞ Sp. Med. Fragrant Sumach ʒiij.
 Glycerin ʒiv. M.

Sig: A teaspoonful every three hours.

(b) Imbalance Digestive Glands:

- ℞ Podophyllum Laxative
 Dewees'

Sig: A teaspoonful twice daily.

Gl. 3, p. 103

Goiter:

Simple—to prevent, to reduce:

- Sp. Med. Spongia, gtt. v. daily for two weeks,
 once a year.

John J. Sutter, M.D.

Thyroid Enlargement:

(a) Simple Enlargement :

℞ Sp. Med. Iris gtt. xv.-xxv.
 Aq. Dest. q.s. ℥iv. M.
 (Shake Label)

Sig: A teaspoonful four times a day.

Gl. 32, p. 1042

(b) Exophthalmic :

℞ Sp. Med. Ergot 3j.
 Sp. Med. Passiflora 3ij. to 3iv.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful four times a day.

Gl. 32, p. 1050

(c) Enlargement with Anemia and Cardiac Lesions:

℞ Sp. Med. Cactus 3j.
 Lloyd's Iron 3jss.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful four times a day.

Gl. 32, p. 1052

Enlarged Thyroid Women—Simple Enlargement :

Wrongs of reproductive system :

℞ Sp. Med. Phytolacca 3ij.
 Sp. Med. Iris
 Sp. Med. Macrotys āā. 3j.
 Elix. Aromatic q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Gl. 4, p.

SPECIFIC MEDICINES

BY JOHN THOMAS LLOYD

Specific Medicines with few exceptions, are liquid preparations of vegetable drugs. They are designed to carry the desirable medicinal qualities of the drugs in therapeutic balance and to eliminate incompatible and antagonistic constituents. They are permanent preparations which, with reasonable care do not deteriorate in any climate.

Specific Medicines are supplied in 4, 8 and 16 ounce square bottles. Each bottle is plainly labeled with the indications for the remedy and with the dose usually dispensed.

Drugs Used in Making Specific Medicines.—Each Specific Medicine is made from a certain part of the plant supplying its name. The bark, the root, the leaf, the catkin, the gum, the wood, the seed, the flower, the fruit or the stem is employed. In some cases the green drug is used, in others the dried. The process of curing is as painstaking as the preparation of tobacco for the market. It is a fallacy to believe that the finer qualities of all drugs must be lost or lessened by drying. On the contrary, in some instances they are developed only by the process of curing. The same principle applies to certain well known foods such as tea, coffee and chocolate, and to tobacco. These, we all know, are useless until proper curing and aging have re-arranged natural structures and developed flavor and aroma that did not exist in the green plant.

The careful and intelligent selection and treatment of the crude drug is the first step in the preparation of a fine medicinal product.

Drug Structures.-Every drug contains many related interstructural constituents. Each part of a plant is an intercellular complexity. No one constituent represents a whole drug. In many cases a drug's dominating substance is less valuable for the uses of the medicinal preparation than the less energetic constituents which it overshadows when in natural proportion. The art of selecting the desirable structures, or limiting those undesirable, is generally a neglected feature of plant pharmacy. *The most conspicuous alkaloid or resin of a drug, though usually viewed as all-important, if present in too great amount, may be an enemy to a preparation's finer qualities.* The study of each drug and each part of a drug with intent of balancing its important interstructural complexity is a part of the pharmacy of Specific Medicines.

Colors.-The coloring matter of a drug is not often a therapeutic factor, but it is usually so intimately associated with the structures that impart its medicinal qualities that separation without injury to the product is impossible. The bark of a root is brown even though the fresh root pulp is white. This bark imparts a brown or red color if the drug be thoroughly represented, for to remove the bark is often to remove the drug's most energetic part.

Inert colors in different lots of drugs may vary according to the soil, sunlight, heat or moisture in which the plant grew. Accordingly, the shade of different batches of Specific Medicines may differ slightly. It is the aim to keep Specific Medicines of constant therapeutic value, regardless of the amount of coloring matter imparted from the crude drug to the preparation.

Precipitates in Specific Medicines.-In marked contrast to vegetable liquid preparations of former times, Specific Medicines, unless allowed to evaporate, seldom form precipitates. It must be remembered that the proportions of the ingredients of the menstrua and the dissolved drug constituents are exactly balanced. *If a bottle is left uncorked or with loose stopper, one constituent of the menstruum evaporates faster than another.* Thus the proportions are changed and precipitates may occur. Evaporation is the most common cause of precipitation in Specific Medicines.

If precipitates do occur in Specific Medicines, the bottle should be shaken before using. In this way the full value of the medicine will enter each prescription.

Compatibility.-Any combination or mixture of Specific Medicines may be made in prescriptions without change of therapeutic power. The physician can with confidence combine all Specific Medicines that are not physiologically antagonistic.

Some Specific Medicines have opposite action, and, accordingly, are classed as physiological incompatibles. For example, remedies that increase secretion and remedies that decrease secretion are physiologically antagonistic. "Hence," in the words of Dr. A. F. Stephens, "to administer gelsemium and belladonna at one and the same time is to administer two drugs that must necessarily act to neutralize each other, with the possible result of one gaining a slight ascendancy through greater activity or larger dosage, which effect, however, must fall short of the normal force of the remedy."

Precipitates in Prescriptions.—Every Specific Medicine designated "Colloidum" will mix clear with water in the dose recommended on the label.

Most Specific Medicines not designated "Colloidum" precipitate when mixed with water. In these preparations there are oils, resins or other constituents of medicinal value that are not water soluble, or else the valuable qualities are so intimately associated with insoluble substances that separation has as yet proved impossible. When precipitates occur in prescriptions they are a valuable part of the medicine and the prescription should be stirred or shaken before each dose is taken.

Colloidums.—Vegetable structures in their natural settings are in non-crystalline form known as "colloids," a term devised by Graham in 1861. For many

years we have studied the colloidal nature of natural vegetable structures, with the result that we are able to offer many Specific Medicines with the non-crystalline (colloidal) structure of the drug transferred unchanged to the liquid solvent of the medicine. These Specific Medicines are designated "Colloidums."

Colloidum Specific Medicines will mix with water, glycerin or syrup without milkiness or precipitation. This remarkable pharmaceutical quality is achieved without lessening or in any way altering the therapeutic values of the preparations. In accomplishing this object each drug has been studied as a thing unto itself, and distinct manufacturing processes have been worked out for each Specific Medicine. Years have elapsed since we perfected the first "Colloidum," but our studies in this field have not lessened.

Strength of Specific Medicines.—Physicians sometimes ask the strength of Specific Medicines compared to tinctures, fluid extracts or the specialties of other manufacturers. Simple though this question may seem on first thought, the comparison in most cases can not be easily made.

As stated, Specific Medicines are made with apparatus designed by us and used in no other pharmaceutical laboratories. Also the menstrua as well as the manipulative processes have resulted from our long and painstaking study.

To understand how two preparations of the same drug can differ in their qualities, let us offer a very simple hypothetical example. Suppose that the crude drug contains equal amounts of two familiar constituents, shellac and sugar. In this simple case, if an extract be made with strong alcohol the product will contain shellac but little or no sugar. If, on the other hand, an aqueous menstruum is used, the sugar will be easily extracted but the shellac will remain in the drug. A half and half mixture of alcohol and water will dissolve some sugar and a small amount of shellac, but will not dissolve them in the same proportion. With even a slight change in the alcoholic per cent of the menstruum, there will be a decided change in the proportion of the two ingredients.

No plant is as simple as the imaginary example cited, yet this principle is the same in all. In making medicines from vegetable drugs we not only have to do with the action of solvents on simple constituents of different soluble qualities like shellac or sugar, but must also deal with the complex reactions of the extracted constituents themselves.

From the example cited, which illustrates but one of many factors determining the quality of medicines, it should be readily understood that preparations made by different methods and with menstrooms of different strengths must differ not alone in the amount of the dominant energetic constituent

which determines strength, but also in the proportion of less energetic constituents, which is an important factor in determining *quality*. In manufacturing Specific Medicines it has ever been our aim to produce balanced preparations in which the energy of no constituent overshadows the milder action of less energetic principles. The quality of Specific Medicines can not be gauged by the quantity or action of any one constituent or any one separate, but must be determined by the therapeutically balanced relation of all of the drug constituents desirable for the purpose for which the preparation is designed.

Vehicle for Dilution.-Water is the most frequently used vehicle for Specific Medicines in prescriptions. When the taste of the medicine is not disagreeable to the patient and when the prescription is not to be left for a considerable length of time, no vehicle is superior to water.

Like most all other organic substances and most inorganics, when greatly diluted with water Specific Medicines may sour. To prevent souring when the prescription is not to be completely consumed within a few days, as well as to overcome or mask bitter and disagreeable flavors, nothing else known to us will so perfectly fill the requirements for a vehicle as Glyconda. Glyconda is described more at length in the following paragraph.

Glyconda As a Vehicle.-When an alkaline agent is not contra-indicated, Glyconda is unsurpassed as a

vehicle for prescribed medicines. Its deep, brilliant, wine-red color adds to the appearance of the prescription, while its characteristic flavor, free from the sweetness of sugar or syrup, is almost always pleasant to the sick. Its power of masking bitter and unpleasant flavors is remarkable. Even such intensely bitter flavors as *Nux Vomica* and *Colocynth* in medicinal doses are overcome or greatly lessened when prescribed in *Glyconda*.

Almost all medicines not dependent upon an acid menstruum may be prescribed in *Glyconda*. In our experience there is no menstruum in which all medicines will mix without precipitation. While precipitates do occur when some medicines are mixed with *Glyconda*, the precipitate is almost always light and flocculent. Even such heavily resinous preparations as *Jalap*, *Xanthoxylum* and *Podophyllum* do not form the heavy, sticky precipitates that characterize them when mixed with water.

When *Glyconda* is the vehicle, even in mid-summer, prescriptions need not be replaced on account of fermentation. *Glyconda* will neither freeze nor ferment.

INDICATIONS AND DOSES OF IMPORTANT SPECIFIC MEDICINES

SPECIFIC MEDICINE ACID CARBOLIC (Phenol) —
Specific Indications: A broad, moist tongue, cadaverous odor of breath, gastric fermentation.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Locally, as an antiseptic, diluted with three parts water.

Usual Dosage :

℞	Sp. Med. Ac. Carbolic	gtt. x.
	Syrup or Water	q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE ACID HYDROCHLORIC — *Specific Indications:* Deep red, dry and contracted tongue, with brownish coating, brown sordes on teeth and tongue. Tongue contracted, fissured and brown. Pungent heat of skin. Slow digestion with deficient gastric juice acidity.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞	Sp. Med. Ac. Hydrochloric	gtt. x.
	Aq. Dest.	q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE ACID SULPHUROUS (A Compound) — A supersaturated alkaline aqueous solution of Sulphur Dioxide.

Specific Indications: Full relaxed tissues, deep redness, sticky unhealthy discharges. Sweetish mawkish odor of breath and excretions, increased, viscid saliva. Full, broad tongue, atonic, normally red, with glutinous coat, transparent or dirty-brownish, effaced papillae. Sepsis and debility.

Locally in parasitic skin diseases.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞	Sp. Med. Ac. Sulphurous	℥iij.
	Aq. Dest.	q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Locally, 3j, diluted with four to twenty parts water,

SPECIFIC MEDICINE ACONITE—*Specific Indications*: Small frequent pulse, with impaired circulation. Dryness of skin, lack of tone of capillary circulation. Acute inflammations of mucous membranes of nose, throat and larynx.—H. W. Felter, M.D. Mat. Med. Phar. Ther.

Usual Dosage :

℞ Sp. Med. Aconite gtt. v.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every hour.

SPECIFIC MEDICINE AESCULUS (Buckeye) — *Specific Indications*: Constriction of chest, with spasmodic cough; of intestines, near umbilicus, with colic; of rectum, with tightness, with or without haemorrhoids.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Aesculus ℥j.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE AMBROSIA — *Specific Indications*: Sneezing, with excessive irritation of mucous membranes, nose, throat, mouth, urethra, bowels, with free mucous discharge.—J. S. Niederkorn, M.D. Reference Book.

Usual Dosage :

℞ Sp. Med. Ambrosia ℥ij.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE AMYGDALUS—*Specific Indications*: Gastric and abdominal tenderness. Elongated, pointed tongue, reddened tip and edges, prominent papillae. Nausea and vomiting. Irritative cough, colds and bronchitis.—John King, M.D. Amer. Disp.

Usual Dosage :

℞ Sp. Med. Amygdalus gtt. xx.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every half hour.

SPECIFIC MEDICINE APIS—*Specific Indications*: Frequent urination, with scanty burning urine. Burning, itching skin, urticarial in character. Edema of mucous membranes, sore throat, coughs and colds.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Apis gtt. x.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE APOCYNUM—*Specific Indications*: Local edema, especially of feet and legs, pitting on pressure; feeble heart action, and defective renal elimination.—Finley Ellingwood, M.D. Am. Mat. Med. and Ther.

Usual Dosage :

℞ Sp. Med. Apocynum gtt. xxx.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE ASCLEPIAS (Pleurisy Root)—*Specific Indications*: Skin hot, but inclined to moisture, face flushed, vascular excitement of bronchial region, scanty urine; serous or synovial inflammation.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Asclepias gtt. xxx.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

ASEPSIN—A definite sodium compound, a white crystalline powder, prepared from oil of wintergreen. Soluble in water, hot or cold, decomposed by acids.

Specific Indications: Fermentation and putrefaction; pale tongue and dusky discoloration, of throat and tongue, feeble capillary circulation, with tendency to breaking down of tissue.

Usual Dosage :

℞ Asepsin gr. v.
Milk Sugar ℥j. M.

Div. in Chart. No. xij.

Sig: A powder every four hours with one-quarter glass water.

Locally, a five per cent. solution may be used as a wash or irrigation.

H. W. Felter, M.D. Mat. Med. Pharm. Ther.

SPECIFIC MEDICINE ASTHMA WEED (Euphorbia Puli-fer)—*Specific Indications*: Dyspnea; spasmodic action of respiratory muscles, with bronchial irritation.—John King, M.D. Amer. Disp.

Usual Dosage :

℞ Sp. Med. Asthma Weed ℥j.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every hour.

SPECIFIC MEDICINE AVENA (Common Oat)—*Specific Indications:* Nervous exhaustion of convalescence, cardiac weakness with nervous depression.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Avena ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE BAPTISIA—*Specific Indications:* Fullness of tissue, with dusky, leaden, purplish discoloration; tendency to ulceration and sloughing. Face swollen and bluish, enfeebled circulation, fetid discharges.—H. W. Felter M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Baptisia gtt. xx.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE BAROSMA (Buchu)—*Specific Indications:* Acid urine, with constant desire to urinate; vesico-renal irritation, with copious mucous or muco-purulent discharges.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Barosma ʒiv.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE BELLADONNA—*Specific Indications:* Dull expressionless face, dilated pupils, impaired capillary circulation of skin and mucous membrane, soft oppressed easily compressed pulse. Urinary incontinence. Spasm of involuntary muscles.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Belladonna gtt. v.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE BERBERIS—*Specific Indications:* Chronic blood dyscrasia, with scaly skin eruptions and impaired nutrition and waste.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :
 ℞ Sp. Med. Berberis ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE BLACK HAW (Viburnum Prunifolium) — *Specific Indications:* Uterine irritability and hyperesthesia, uterine colic, with severe lumbar and Pelvic cramps.—H. W. Felter, M.D. Mat. Med. Phar. Ther.

Usual Dosage :
 ℞ Sp. Med. Black Haw ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE BRYONIA—*Specific Indications:* Sharp cutting pain in serous or synovial membranes, increased by pressure or motion; moderately full vibratile pulse; hacking, racking explosive cough.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Bryonia gtt. x.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE CACTUS—*Specific Indications:* Impaired heart action, feeble, irregular tumultuous, with mental depression, apprehension and praecordial oppression.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Cactus ʒj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE CANNABIS—*Specific Indications:* Great nervous depression, mental illusions, spasmodic neuralgia. Genito-urinary irritation with tenesmus and scalding frequent urination.—John King, M.D. Amer. Disp.

Usual Dosage :
 ℞ Sp. Med. Cannabis gtt. xxx.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE CAPSELLA—*Specific Indications:* Passive haemorrhage and discharge of mucous membrane, especially of female generative organs. Prolonged and recurring menorrhagia.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Capsella ʒijj.
 Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE CAPSICUM—*Specific Indications:* Marked depression with feeble pulse and scanty secretions. Tongue dry and harsh, salivary secretions suppressed.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Capsicum gtt. ij.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

Local use: Stimulant, rubefacient, counter-irritant.

Preparations: Emplastrum Capsici.

Capsicum occurs also in Compound Lobelia Powder, and in Libradol, and in Compound Tincture of Myrrh and Capsicum.

SPECIFIC MEDICINE CASCARA (Rhamnus Purshiana)—*Specific Indications:* Constipation, due to neglect, or to nervous or muscular atony of the bowels.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Cascara ℥j.
Glyconda
Aq. Dest. aa. q.s. ℥iv. M.

Sig: A teaspoonful every four hours to effect, then reduce the dosage.

SPECIFIC MEDICINE CAULOPHYLLUM—*Specific Indications:* Uterine heaviness and sense of soreness in legs, with pelvic congestion. Sluggish labor pains.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Caulophyllum 3ss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Related Preparations: Leontin, a 1% solution of Leontin, a glucosid, the emmenagogue principle of caulophyllum, in water, Dose 5 to 15 drops in water or syrup.

Syrup Mitchella Compound, Uterine tonic, Dose, one ounce.

SPECIFIC MEDICINE CEANOTHUS—*Specific Indications:* Gastric and hepatic disorders with splenic hypertrophy, expressionless countenance, sallow doughy skin. Catarrhal conditions with profuse mucous flow. Antihemorrhagic.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Ceanothus ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE CHELIDONIUM—*Specific Indications:* Full, pale, sallow tongue and membranes; skin sallow, sometimes greenish. Hepatic congestion with light pasty stools; fullness in right hypochondrium, with tense throbbing to right shoulder.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Chelidonium ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE CHIONANTHUS—*Specific Indications:* Jaundice of skin and conjunctiva. Hepatic tenderness upon deep pressure, light clay-colored stools, high colored urine.—H. W. Felter M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Chionanthus ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE CINCHONA—*Specific Indications:* Periodicity; soft open pulse, tongue moist and cleaning, skin soft and moist, nervous system free from irritation.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Cinchona ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE CINNAMOMUM—*Specific Indications:* Passive, haemorrhage. Gastric irritation with flatulence. Antiseptic, preservative, a flavor.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Cinnamomum ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE COLCHICUM—*Specific Indications:* Gouty, rheumatoid or lithemic diathesis, characterized by muscular involvement of a sharp, tearing, cutting quality, aggravated by heat and pressure.—J. W. Fyfe, M.D. Spec. Medication.

Usual Dosage :

℞ Sp. Med. Colchicum gtt. xxx.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE COLLINSONIA—*Specific Indications:* Atony of venous circulation, irritation and constriction of mucous membrane of larynx, hoarseness. **Gastro-intestinal** irritation with sluggish portal circulation.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Collinsonia ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE COLOCYNTH — *Specific Indications:* Abdominal distress, cutting, boring in character. Tormina and tenesmus, gaseous distension, dry scybalous stools. — H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Colocynth gtt. v.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour.

SPECIFIC MEDICINE CONIUM—*Specific Indications:* Nervous excitation and restlessness with or without pain or distress.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Conium ʒss.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE CORNUS—*Specific Indications:* Periodicity, with feeble relaxed tissues, weak pulse, subnormal temperature.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Cornus ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every hour. In intermittent fevers, ten to fifteen drops of the *Specific Medicine Cornus* may be given in water every hour as required.

SPECIFIC MEDICINE CORYDALIS—*Specific Indications:* Blood dyscrasias, sluggish digestion, deficient glandular secretions. Disturbed menstruation, following exhausting diseases.—Finley Ellingwood, M.D. Amer. Mat. Med. Ther.

Usual Dosage :

℞ Sp. Med. Corydalis ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE CRATAEGUS — *Specific Indications:* Cardiac weakness and palpitation, irregular intermittent pulse, with increased rate, dyspnoea and nervous depression.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Crataegus ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE CYPRIPEDIUM—*Specific Indications:* Insomnia, irritability, neuralgia, restlessness, muscular twitching and tremor, due to atony.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Cypripedium ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE DIGITALIS—*Specific Indications:* Weak, rapid, irregular heart action, jugular fullness; labored, accelerated breathing; edema, anasarca, ascites, scanty high-colored urine.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Digitalis gtt. xxx.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE DIOSCOREA—*Specific Indications:* Spasmodic abdominal colic, nausea, with skin and conjunctiva yellow. Twisting boring distress centered at umbilicus.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Dioscorea ʒss.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE DROSERA—*Specific Indications:* Dryness of air passages, with spasmodic frequent cough, explosive in character.—John King, M.D. Amer. Disp.

Usual Dosage :

℞ Sp. Med. Drosera ʒss.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE ELATERIUM—*Specific Indications:* Chronic cystitis, with soreness in neck of bladder; micturition followed by violent cramp-like aching extending from bladder to thighs and pelvis. Ascites of hepatic or abdominal origin, in the plethoric.—John King, M.D. Amer. Disp.

Usual Dosage :
 ℞ Sp. Med. Elaterium gtt. x.
 Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE ECHINACEA—*Specific Indications:* Systemic sepsis, tendency to boils and to formation of semi-internal multiple cellular abscesses, with adynamia and asthenia. Foul discharges with emaciation. Dirty brownish or bluish tongue, with sordes. Skin and mucous membranes, dull bluish or purplish in color.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Echinacea ʒj.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

Echinacea is applied locally in aphthous and herpetic eruptions; it is deodorant, antiseptic, and slightly anaesthetic.

Related Preparations: Echafolta, for external use. Echafolta Cream, mild soothing ointment.

SPECIFIC MEDICINE ERGOT—*Specific Indications:* Active haemorrhage. Post-partum haemorrhage with uterine inertia. Venous fullness, mental apathy, cold surface, low blood pressure.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Ergot ʒj.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

In active haemorrhage, one drachm Specific Medicine Ergot repeated as conditions require.

SPECIFIC MEDICINE ERIGERON—*Specific Indications:* Free discharge from mucous membranes. Passive capillary haemorrhage. Choleraic discharges, sudden, gushing and watery, attended by cramping and distress.—John King, M.D. Amer. Disp.

Usual Dosage :
 ℞ Sp. Med. Erigeron ʒj.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

Oil of Erigeron, capillary or passive haemorrhage: haematuria, haemoptysis, epistaxis, haematemesis, metrorrhagia. Dose 1 to 60 drops.

SPECIFIC MEDICINE ERYNGIUM—*Specific Indications:* Irritability of bladder and urethra, with burning and frequent desire to urinate, aching extending to loins. Scanty, scalding urine.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :
 ℞ Sp. Med. Eryngium ʒss.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE EUPATORIUM—*Specific Indications:* Large full pulse, current showing small waves, skin, hot and full, with tendency to moisture. Deep-seated aching in bones with general bodily aching. Hoarseness, cough, soreness of chest. Urine turbid and high-colored.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Eupatorium ʒj.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE EUPHRASIA—*Specific Indications:* Acute irritating inflammation of mucous membranes of eyes and upper respiratory passages, with acrid watery discharges.—Finley Ellingwood, M.D. Amer. Mat. Med. and Ther.

Usual Dosage :
 ℞ Sp. Med. Euphrasia ʒj.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE FRAGRANT SUMACH (Rhus Aromatica)—*Specific Indications:* Pulse small and feeble, stools profuse, abdomen flabby, tongue pale, trembling and moist, languor and lassitude. Nocturnal enuresis. Frequent micturition with enlarged prostate.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Fragrant Sumach ʒj.
 Glyconda
 Aq. Dest. āā. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE GAULTHERIA—*Specific Indications:* Irritation of bladder and prostate, undue sexual excitement. Early stages of nephritis.—H. W. Felter M.D. Mat. Med. Pharm. Ther.

Usual Dosage :
 ℞ Sp. Med. Gaultheria ʒj.
 Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE GELSEMIUM—*Specific Indications:* Hyperemia. Bright eyes, contracted pupils, nervous unrest. Tremors with nervous excitement and marked temperature. Dysuria with scanty urine. Dryness of parturient canal with thin rigid os uteri.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Gelsemium gtt. xv.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE GERANIUM—*Specific Indications:* Relaxed mucous tissues, with profuse debilitating discharges. Diarrhoea with constant desire to defecate. Passive haemorrhage.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Geranium ʒij.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE GOSSYPIUM—*Specific Indications:* Delayed menstruation? backache, and sense of fullness and dragging in pelvis. Difficult micturition, with weight and fullness in bladder.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Gossypium ʒss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE GRAVEL ROOT (Eupatorium purpureum)—*Specific Indications:* Functional derangements of urinary organs, scanty, burning urination.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Gravel Root ʒij.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE GRINDELIA—*Specific Indications:* Dyspnea with cyanosis in the plethoric individual, attended by mucous rales and ineffectual expectoration.—Lyman Watkins, M.D. Compend Pract. Med.

Usual Dosage :

℞ Sp. Med. Grindelia ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Locally applied in dilution as treatment for rhus poisoning, and to slow-healing ulcers.

SPECIFIC MEDICINE GUAIAIACUM (Guaiac)—*Specific Indications:* Dryness and stiffness of throat, in incipient inflammation of tonsils and pharynx, painful deglutition and dribbling of saliva.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Guaiacum ʒj.
Glycerine ʒss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

As a gargle, add ʒij to glass warm water.

SPECIFIC MEDICINE HAMAMELIS—*Specific Indications:* Full tissues, venous stasis, excessive mucous flow.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Hamamelis ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Local use, sprains, contusions, minor wounds, haemorrhoids. Inflamed or contused skin or mucosa, venous stasis.

**SPECIFIC MEDICINE HELLEBORUS NIGER (Black Hel-
lebores)**—*Specific indications:* Flashes of heat, burning of sur-
faces, sensitiveness of perineal structures. Weak, rapid irregu-
lar heart action, low arterial tension. Jelly-like, mucoid bowel
evacuations, dullness and stupor.—J. W. Fyfe, M.D. Specific
Medication.

Usual Dosage :

℞ Sp. Med. Helleborus Niger gtt. v.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE HELONIAS—*Specific Indications:* Re-
laxation of uterine tissue, with pelvic fullness, irritability, de-
spendency and mental torpor.—H. W. Felter, M.D. Mat. Med.
Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Helonias ʒss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE HUMULUS (Hops)—*Specific Indica-
tions:* Nervousness, irritability, insomnia. Acid eructations.
Vesical irritation.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Humulus ʒij.
Glycerin ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Related Preparation: Specific Medicine Lupulin.

SPECIFIC MEDICINE HYDRANGEA—*Specific Indications:* Vesical and urethral irritation with dull aching in back, urine tinged with blood.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Hydrangea 3j.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE HYDRASTIS (Golden Seal)—*Specific Indications:* Relaxed mucous membranes, with feeble circulation, and profuse mucous flow of thick, tenacious, yellowish or greenish-yellow character. Gastric irritability and anorexia.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Hydrastis 3j.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every four hours.

Related Preparations: Colorless Hydrastis (Lloyd's Hydrastis), Dose, 5 to 15 drops. Also used locally.

Hydrastine Muriate (Berberine Hydrochloride). Average dose gr. ij.

SPECIFIC MEDICINE HYOSCYAMUS—*Specific Indications:* Nervous irritability with insomnia and restlessness. Dilated pupils, flushed face. Low muttering delirium. Urinary incontinence in the feeble and the aged.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Hyoscyamus gtt. x.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE IGNATIA—*Specific Indications:* Atony of female generative organs with nervous debility. Deep-seated dull soreness in epigastrium extending to right shoulder.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Ignatia gtt. v.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE IPECACUANHA (Ipecac)—*Specific Indications:* Irritation, long pointed tongue, reddened tip and edges, nausea and vomiting. Increased bronchial secretion and hoarseness.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Ipecacuanha gtt. x.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE IRIS—*Specific Indications:* Enlarged soft, yielding lymphatic tissue. Gastro-intestinal irritation, burning in the epigastrium, acid eructations.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Iris 3ss.
Glycerin ℥ss.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE JABORANDI (Pilocarpus)—*Specific Indications:* Skin and mucous membranes hot and dry, with deficient secretion. Pulse, full, hard, sharp and strong; dry parched tongue.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Jaborandi 3ss.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE JALAP—*Specific Indications:* Sluggish bowel action from deficient secretion of intestinal glands. Edema and anasarca, with colonic stasis, tormina and tenesmus.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Jalap gtt. xx.
Glyconda
Aq. Dest. āā. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Related Preparation: Compound Jalap Powder. (Antibilious Physic).

Dose, 60 grains in hot water.

SPECIFIC MEDICINE JUGLANS—*Specific Indications:* Gastro-intestinal irritation with acid eructations and flatulence. Tenesmus with burning fetid alvine discharges. Chronic vesicular skin disease with free discharge.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Juglans 3ss.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE KRAMERIA—*Specific Indications:* Relaxed mucous membranes with mucous discharges and full relaxed skin.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Krameria 3j.
Aq. Dest. q.s. ℥iv. M.
Sig: A teaspoonful every four hours.

Locally it is applied to relaxed tissue.

SPECIFIC MEDICINE LEPTANDRA—*Specific Indications:* Drowsiness, dizziness, mental depression, with tenderness and heaviness in hepatic region. Tongue coated markedly white, nausea, skin yellow, extremities cold, dull frontal headache, clay-colored stools.—John King, M.D. American Disp.

Usual Dosage:

℞ Sp. Med. Leptandra ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE LOBELIA—*Specific Indications:* Fullness of tissue, with turgid veins and tense arterial flow; labored doughy pulse, labored breathing, praecordial oppression; mucous bronchial accumulations.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

In Labor, thick, rigid unyielding os uteri and vaginal tissues.

Usual Dosage:

℞ Sp. Med. Lobelia ʒss.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

Related Preparations: Subculoyd Lobelia, dose 1 to 20 minims subcutaneously.

Compound Emetic Powder, a local application in chest conditions. Libradol, a cataplasma, for painful inflamed conditions.

SPECIFIC MEDICINE LUPULIN—*Specific Indications:* Nervous irritability, cerebral hyperemia with insomnia. Mental irritability associated with wrongs of reproductive system. Fermentative indigestion with acid eructations.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Lupulin ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Related Preparation: Specific Medicine Humulus.

SPECIFIC MEDICINE LYCOPUS—*Specific Indications:* Vascular excitation with rapid, tumultuous heart action lacking power, impaired breathing and tendency to small passive haemorrhage. Morbid wakefulness with active but weak circulation.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Lycopus ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE MACROTYS—*Specific Indications:* Heavy, tense muscular soreness, tense drawing muscular contraction. Deep-seated boring distress of stomach, bowels or uterus. Muscular involvement of the rheumatoid diathesis.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Macrotys ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE MATRICARIA—*Specific Indications:* Nervous irritability, fretfulness, muscular twitching: Fetid, feculent, greenish alvine discharges, associated with flatulence, colic and anal excoriation.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Matricaria ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE MITCHELLA—*Specific Indications:* Atony of female reproductive organs, with tardy menstruation, and pelvic dragging, tenderness and pressure.—J. W. Fyfe, M.D. Spec. Med.

Usual Dosage:

℞ Sp. Med. Mitchella ʒss.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Related Preparations: Syrup Mitchella comp. (Compound syrup of Partridgeberry). Uterine tonic. Dose one drachm to one ounce.

SPECIFIC MEDICINE MYRICA (Myrrh)—*Specific Indications:* Full, relaxed, spongy mucous membrane with increased secretions. Tenacious secretions, offensive and irritating.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage:

℞ Sp. Med. Myrica ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every three hours.

Related Preparation: Compound Tincture Myrrh and Capsicum, stimulant.

SPECIFIC MEDICINE NUX VOMICA—*Specific Indications:* Atony. Tongue pallid and uncoated or heavy pasty yellowish coating. Yellowish tinge to skin and conjunctiva, sallow line around mouth. Fullness and dull pain in right shoulder pointing toward umbilicus.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Nux Vomica gtt. v.
Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE OENANTHE—*Specific Indications:* Spasmodic seizures, epileptiform in character, with anemia of brain and cord.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Oenanthe gtt. x.
Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.
Note—Colloidal Oenanthe is twice the strength of the Specific Medicine.

SPECIFIC MEDICINE PASSIFLORA INCARNATA—*Specific Indications:* Irritation and atony of brain and nervous system with insomnia.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Passiflora Incarnata ʒj.
Aq. Dest. qs. ʒiv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE PHOSPHORUS—*Specific Indications:* Nervous exhaustion and atony. Vesical and urethral irritation with mucoid discharges, fullness and dragging in perineum. Bowel discharges slimy and frothy, with tympanites. Delayed bone resolution.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Phosphorus gtt. xx.
Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE PHYSOSTIGMA—*Specific Indications:* Feeble, tremulous pulse, cool surface, cold extremities? contracted pupils. Small rapid pulse, with constriction and difficult breathing. Relaxed muscular walls of stomach and bowels with impaired secretion.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Physostigma gtt. v.
Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE PHYTOLACCA—*Specific Indications:* Enlargement and inflammation of glandular structures, mucous membranes pallid. Impaired glandular secretion and function.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Phytolacca gtt. xxx.
Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE PIPER METHYSTICUM (Kava Kava)—*Specific Indications:* Irritation, inflammation, atony of urinary passages., painful micturition, scanty and irregular. Pale edematous tissues.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Piper Methysticum ʒj.
Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE PISCIDIA—*Specific Indications:* Insomnia and nervous unrest, spasm, pain, nervous irritability; neuralgias of trifacial and cervical plexuses.—C. S. Amidon, M.D.

Usual Dosage :

℞ Sp. Med. Piscidia ʒj.
Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE PLANTAGO MAJOR—*Specific Indications:* Nocturnal enuresis in children, with pale abundant urine, irritation and relaxation of sphincter vesicae.—John King, M.D. Amer. Disp.

Usual Dosage :

℞ Sp. Med. Plantago Major ʒj.
Aq. Dest. q.s. ʒiv. M.
Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE PODOPHYLLUM—*Specific Indications:* Fullness of tissue, fullness of superficial veins, dirty yellowish coating of tongue.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Podophyllum gtt. xx.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Related Preparations: Sp. Med. Podophyllin Triturate, dose 1 to 10 grains. Podophyllum Laxative (Dewees'), Dose, one-half to 1 drachm.

SPECIFIC MEDICINE POLYMNIA—*Specific Indications:* Full, sodden, inelastic tissues. Splenic and hepatic enlargement, with fullness and weight in epigastric region. Impairment of function of parts supplied by coeliac axis.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Polymnia 3ss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE POLYTRICHUM—*Specific Indications:* Vesical irritation, scanty painful urination with anasarca.—J. W. King, M.D. Amer. Disp.

Usual Dosage:

℞ Sp. Med. Polytrichum ʒij.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE PRUNUS (Wild Cherry Bark)—*Specific Indications:* Irregular or intermittent heart action; convulsive action due to overstrain. Irritation of stomach with cough, bronchial irritation. Impaired appetite and atonic indigestion. Lack of muscular tone.—J. W. Fyfe, M.D. Spec. Med.

Usual Dosage:

℞ Sp. Med. Prunus gtt. xxx.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

Related Preparation: Syrupus Pruni Virginianae. A syrup used as a tonic and sedative in cases where prunus is indicated, also as a vehicle for cough mixtures.

SPECIFIC MEDICINE PULSATILLA—*Specific Indications:* Nervousness, fear of impending danger. Nervous symptoms depending upon wrongs of reproductive system. Pasty, creamy-white coating upon tongue.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Pulsatilla ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE RHAMNUS CALIFORNICA—*Specific Indications:* Rheumatoid conditions, depending upon wrongs of intestinal tract, atonic in character. Rheumatoid diathesis.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Rhamnus Californica ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE RHEUM (Rhubarb)—*Specific Indications:* Gastric irritation, elongated, reddened tongue, nausea and vomiting. Irritative diarrhoea, tenderness on pressure. Light-colored acrid fecal discharges.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Rheum 3ss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

Related Preparation: Glyconda, Neutralizing Cordial. Dosage 1 to 4 drachms.

SPECIFIC MEDICINE RHUS TOX—*Specific Indications:* Long, pointed tongue, prominent papillae, heat redness, unrest. Moderately sharp, quick pulse, sometimes vibratile. Urethral burning with urinary dribbling. Inflammations, red swollen surface, deep burning pain.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Rhus tox gtt. v.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE RUMEX (Yellow Dock)—*Specific Indications:* Vitiated blood, with skin disorders; low glandular and cellular deposits, with tendency to ulceration. Dyspnea, with epigastric fullness and pectoral distress. Anorexia, with disturbed nutrition.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Rumex ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE SALIX NIGRA AMENTS—*Specific Indications:* Sexual erethism. Vesical and urethral irritation from venereal excess.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Salix Nigra Aments ʒj.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE SANGUINARIA—*Specific Indications:* Burning, itching of throat, air passages, hot dry and swollen. Sense of constriction in throat, difficult deglutition. Bronchial irritation with increased secretion. Uneasiness and burning in stomach. - J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Sanguinaria gtt. x.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

Related Preparations: Sanguinarine Nitrate, mixed alkaloids of Sanguinaria, dose, 1/30 to 1/4 grain.

Trituration Sanguinarine Nitrate (1: 100) dose 1 to 10 grains.

SPECIFIC MEDICINE SAW PALMETTO—*Specific Indications:* Relaxation of tissues, copious mucous secretion. Genito-urinary distress and dribbling. Impaired appetite, digestion and assimilation. - H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Saw Palmetto ℥j.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours,

SPECIFIC MEDICINE SCUTELLARIA (Skullcap)—*Specific Indications:* Nervous twitchings from mental or physical exhaustion, tremors, restlessness. - H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Scutellaria ℥j.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE SENECIO—*Specific Indications:* Atony and relaxation of pelvic contents with discharge, weight and vascular engorgement. - H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Senecio ℥j.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE SODIUM PHOSPHATE—*Specific Indications:* Deficient biliary excretion, pale membranes, tongue dirty and moist, or with yellow coat or dry fur upon a pallid tongue; sallow skin intestinal stasis. - H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

One to sixty grains well diluted with water.

SPECIFIC MEDICINE SODIUM SULPHITE—*Specific Indications:* Tongue broad and pallid, pasty whitish or yellowish coating. Pallid mucous tissue with light and pasty exudate. — J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Sodium Sulphite gr. xx.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every two hours.

SPECIFIC MEDICINE SOLANUM CAROLINENSE—*Specific Indications:* Epileptiform and choreiform seizures. Cough spasmodic in character. - J. W. Fyfe, M.D. Specific Medication.

Usual Dosage :

℞ Sp. Med. Solanum car. ʒss.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE SPIKENARD—*Specific Indications:* Irritation of mucous surfaces of respiratory tract, wheezing respiration, expectoration of tough stringy mucus. — Lyman Watkins, M.D. Comp. Prac. Med.

Usual Dosage :

℞ Sp. Med. Spikenard ℥j.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every three hours.

SPECIFIC MEDICINE SPONGIA—*Specific Indications:* Simple enlargement of thyroid and cervical glands. Laryngeal irritation with dry catarrhal cough. - H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Spongia gtt. xx.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE SQUILL—*Specific Indications:* Cough, with scanty expectoration. Cardiac edema, general asthenia, scanty high-colored urine. Hypotension and cardiac atony. — H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage :

℞ Sp. Med. Squill gtt. xx.
Aq. Dest. q.s. ℥iv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE STAPHISAGRIA—*Specific Indications:* Chronic irritation and inflammation of genito-urinary tract, scalding urination, prostatorrhoea and urinary incontinence.—J. W. Fyfe, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Staphisagria ʒss.
Aq. Dest. q.s. ad. ʒiv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE STICTA PULMONARIA—*Specific Indications:* Short hacking cough, pain in shoulder extending to neck and back of head. Soreness of pectoral muscles increased by deep breathing.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage:

℞ Sp. Med. Sticta Pulmonaria ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE STILLINGIA—*Specific Indications:* Feeble tissues with impaired waste and repair. Mucosa red, glistening and tumid, scanty secretion. Laryngeal irritation, hoarseness and cough. Skin lesions with ichorous discharge.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Stillingia ʒss.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

Related Preparation: Compound Stillingia Liniment, dose one drop on sugar or in simple syrup.

SPECIFIC MEDICINE STRAMONIUM—*Specific Indications:* Cerebral irritation, with or without delirium, face congested, red and bloated, restlessness and fearfulness.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Stramonium gtt. v.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE STROPHANTHUS—*Specific Indications:* Irregular heart action, praecordial pain, dyspnea. Edema. Heart lesions showing atony and weakness of cardiac muscle.—J. W. Fyfe, Specific Medication.

Usual Dosage:

℞ Sp. Med. Strophanthus gtt. x.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE THUJA—*Specific Indications:* Vesical irritation and atony, urinary incontinence. Atonic enuresis. Catarrhal flow from bladder or genitalia.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Thuja ʒj.
Glycerin ʒss.
Aq. Dest. q.s. Z-iv. M.

Sig: A teaspoonful every four hours.

External Use: Stimulant, sub-astringent, deodorant, antiseptic. Used in the treatment of verrucous growths, haemorrhoids, fistula, hydrocele, hernia.

Related Preparation: Aqueous Thuja, chiefly used locally: dose, internally 1 to 30 drops.

Long's Thuja, an ointment.

SPECIFIC MEDICINE TIGER LILY—*Specific Indications:* Uterine congestion, weight and downward pressure in lower abdomen. Reflex nausea and headache from uterine causes.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage:

℞ Sp. Med. Tiger Lily ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE TRITICUM REPENS (Couch Grass)—*Specific Indications:* Irritation of urinary system, with catarrhal discharge, aching in back, dysuria and tenesmus.—J. W. Fyfe, M.D. Specific Medication.

Usual Dosage:

℞ Sp. Med. Triticum Repens ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every four hours.

SPECIFIC MEDICINE VALERIAN—*Specific Indications:* Cerebral anemia, mental depression, despondency, mild spasmodic movements.—H. W. Felter, M.D. Mat. Med. Pharm. Ther.

Usual Dosage:

℞ Sp. Med. Valerian ʒj.
Aq. Dest. q.s. ʒiv. M.

Sig: A teaspoonful every two hours.

