

61/3486/A




Henry Brabazon Urmston.

★ GEMSTONES-OF-NORTH-AMERICA-PROSPECTING-FOR-GEM ★

★ OF-GEMS-& GEM-CUTTING ★

★ STONES-AND MINERALS ★

★ MINERALOGY-EMERALD-AND-OTHER-BERYLS-CATALOG ★



EX LIBRIS
JOHN SIN KAN KAS



3460 @ 22.-

X # 5.50 1966


Friedländer


9⁰⁰

Elizth Armstrong

1831







3L
RL015206

NEW
DESCRIPTIVE CATALOGUE
OF
Minerals;

WITH
DIAGRAMS OF THEIR SIMPLE FORMS:

Intended for the
USE OF STUDENTS,
IN
THE CLASSIFICATION OF MINERALS,
AND THE
Arrangement of Collections.

BY JOHN MAWE.

Honorary Member of the Mineralogical Society of Jena; Member of the Royal Geological Society; First Administrator and Mineralogist to His Majesty the King of Portugal, &c. &c. Author of the Mineralogy of Derbyshire; Travels in South America and through the Gold and Diamond Districts of Brazil; Treatise on Diamonds and Precious Stones; Familiar Lessons on Mineralogy and Geology, &c. &c.

THE SEVENTH EDITION.



LONDON:

PRINTED FOR AND SOLD BY THE AUTHOR, 149, STRAND;
AND LONGMAN, HURST, REES, ORME, BROWN, AND GREEN,
PATERNOSTER-ROW.

1826.

*W. M^cDowall, Printer, Pemberton Row,
Gough Square.*

SIMPLE FORMS.




The Author is indebted to Professor Clarke's excellent Syllabus for the List of Names.



TETRAHEDRON.


Having three sides and a base.



Blende, Zinc Sulphuret.
Grey Copper, Sulphuret.

Pyritous Copper, Sulphuret.


CUBE.



Amphigene? (Leucite). Copper, Native.
Analcime. Copper, Red Oxide.
Boracite. Cobalt, Arsenical.

Fluor Spar.	Lead, Sulphuret
Gold, Native.	Lead, Molybdate.
Iron Pyrites, Super-sulphuret.	Mellite.
Iron, Arseniate.	Silver, Muriate.


RHOMBOID.



Chabasite.	Lime, Carbonate.
Corundum.	Lime, Magnesian Carbonate.
Copper, Phosphate.	Lime, Sulphate.
Iron, Oxide.	Silver, Antimonial Sulphuret.
Lead, Chromate.	Quartz.


RECTANGULAR PRISM.

Having four sides and a rectangular base.




Andalusite.	Mesotype.
Apophyllite.	Meionite.
Antimony, Sulphuret.	Mercury, Muriate.
Chrysolite.	Micarelle. (Pinite).
Cymophane.	Scapolite.
Epidote.	Stilbite.
Harmotome.	Tin, Oxide.
Idocrase.	Uranium, Oxide.
Lead, Muriate.	Wolfram.
Macle.	Zinc, Sulphate.
Manganese, White Oxide.	Zircon.

RIGHT RHOMBOIDAL PRISM.




Barytes, Sulphate.	Mica.
Gadolinite.	Strontian, Sulphate.
Yenite.	Topaz.
Iron, Phosphate.	Tremolite.
Iron, Arsenical.	Prehnite.
Lime, Sulphate.	

OBLIQUE RHOMBOIDAL PRISM.




Axinite.	Hornblende.
Copper, Sulphate.	Iron, Oxide.
Copper, Blue Carbonate.	Lime, Carbonate.
Corundum.	Scapolite.
Euclase.	Sphene, Siliceo-calcareous Ti-
Feldspar.	tanium.

OCTAHEDRON.



Amalgam, Native.	Lead, Sulphate.
Anatase, Titanium Oxide.	Lead, Molybdate.
Bismuth, Native.	Lead, Carbonate.
Ceylonite.	Lead, Sulphuret.
Copper, Native.	Lime, Fluuate.
Copper, Arseniate.	Lime, Tungstate.
Copper, Muriate.	Mellite.
Copper, Red Oxide.	Silver, Native.
Diamond.	Silver, Sulphuret.
Gold, Native.	Spinelle.
Iron, Chromate.	Sulphur.
Iron, Oxide.	Tin, Oxide.
Iron Pyrites, Super-sulphuret.	Zinc, Sulphuret.


HEXAGON AND HEXAHEDRAL PRISM.



Actynolite	Lime, Carbonate.
Arragonite.	Lime, Sulphate.
Barytes, Carbonate.	Lime, Phosphate.
Barytes, Sulphate.	Mercury, Red Sulphuret.
Corundum.	Mica.
Copper, Arseniate.	Molybdenum, Sulphuret.
Dichroite.	Nepheline.
Dioptase.	Pyroxene.
Emerald.	Quartz.
Epidote.	Schorlite.
Grenatite.	Silver, Antimonial Sulphuret.
Hornblende.	Strontian, Carbonate.
Kyanite.	Strontian, Sulphate.
Lead, Arseniate.	Tourmaline*.
Lead, Phosphate.	

* Three sides are so diminished as to give to the hexagonal the appearance of a triangular prism with bevelled or rounded edges. In some specimens the lateral planes are so numerous that the crystal assumes a rounded form with a striated surface, resembling some varieties of aqua-marine.


DODECAHEDRON,
With Isosceles triangular faces.



Corundum.

Quartz.

DODECAHEDRON,
With Rhomboidal faces.



Amalgam, Native.
 Amphigene? (Leucite).
 Ceylonite.
 Copper, Red Oxide.
 Copper, Grey.

Garnet.
 Gold, Native.
 Lime, Fluates, *rare*.
 Mellite.
 Sodalite.

VOCABULARY,

CONTAINING

AN EXPLANATION OF THE TERMS COMMONLY USED
IN MINERALOGY.

- Acicular*, long and slender crystals.
- Aggregate*, several substances adhering together.
- Alloy*, a composition of two or more metals.
- Amalgam*, a combination of two metals, one of which is Mercury.
- Amorphous*, without any regular form.
- Anhydrous*, without water.
- Arborescent*, or *Dendritic*, resembling a tree or shrub.
- Bevelled*—A crystal is said to be bevelled when its edges are replaced by two planes, which are separated only by an edge.
- Brittle*, when the particles fly off in cutting or breaking.
- Botryoidal*, resembling a bunch of grapes.
- Canaliculated*, when the surface is indented by deep channels.
- Capillary*, hair-like.
- Cellular*, exhibiting cells, or pores.
- Chatoyant*, a changeable light, as seen in the eye of a cat.
- Cleavage*, is the division of a crystal in the direction of its natural joints.
- Columnar*, a term used to express a substance of a column-like form, sometimes applied to aggregated crystals.

- Compact*, a mineral is so termed when no particular parts are discernible.
- Concretion* signifies a small distinct mass.
- Conchoidal*, when the fractured surface has smooth, shallow hollows resembling the inside of a shell.
- Coralloidal*, resembling a coral.
- Cuneiform*, having the form of a wedge.
- Decrepitate*, is a term applied to minerals which fly into particles with a crackling noise, when exposed to heat.
- Decomposed*, the consequence of chemical action, which naturally affects some minerals, generally those substances which contain an unusual portion of sulphur.
- Decrement*, relating to structure.
- Dentiform*, tooth-shaped.
- Dendritic*, see Arborescent.
- Disseminated*, when one mineral is found embedded in another.
- Disintegrated*, a term used to express the falling to pieces of a substance.
- Double Refraction* is the property possessed by some minerals, of presenting two images of any object seen through them.
- Drusy*, when the surface is covered with minute aggregated crystals.
- Earthy*, a term relating to fracture and texture.
- Efflorescence*, very small fibres, or powder, produced by decomposition.
- Elastic*, any substance which after being bent returns to its original form, is said to be elastic. Mica is elastic.
- Embedded*, a mineral found in another substance.
- Fascicular*, when small groups of crystals slightly diverge in two opposite directions, like a bundle of pliant rods tied tight in the middle.
- Filiform*, wire or thread-like.

- Foliated*, in parallel layers, like the leaves of a book.
- Fracture*, is the surface a mineral presents when broken, as, compact, foliated, conchoidal, earthy, &c.
- Frangible*, relates to the degree of force necessary to break or separate one piece from another. Thus calcareous spar, from the ease with which it may be broken, is said to be fragile; emery or basalt, tough, &c.
- Friable*, when the particles of a mineral loosely cohere.
- Gangue* or *Matrix*, is the substance in or upon which a mineral is found, or embedded.
- Geode*, a hollow ball, the interior often lined with crystals.
- Glance*, shining.
- Hæmatitic*, having the *structure* of the Red Hæmatites. (Hæmatites signifies blood-red).
- Hepatic*, resembling the liver in color or form.
- Indeterminate*, minerals which have no definable form.
- Iridescent*, tarnish exhibiting prismatic colors.
- Lamellar*—A mineral is said to have a lamellar structure when it can be fractured into thin plates.
- Lenticular*, having a double convex surface, resembling a lens.
- Malleable*, capable of being beaten out with a hammer.
- Mammillated*, in spherical excrescences or elevations.
- Matrix*—See Gangue.
- Nodular*, in irregular globular masses.
- Opaque*, when *no* light is transmitted.
- Phosphorescence*, is a feeble light (unattended by heat), which is emitted by some minerals when thrown on a hot substance, (as in fluor), or when rubbed together, (as in quartz).
- Porous*, see Cellular.
- Plumose*, down-like.
- Pseudomorphous*, is applied to those minerals which exhibit a form peculiar to substances of a different genus

or species; and which they have obtained by coating these substances, or filling up cavities which were formerly occupied by them.

Pulverulent, in a state of powder or dust; feebly cohering.

Radiated, when crystals are so formed as to diverge from a centre.

Ramose, branch-like.

Reniform, kidney-shaped.

Reticulated, when the fibres intersect each other; net-like.

Scopiform, when a number of small crystals are aggregated in tufts or bundles.

Semi-transparent, when objects are seen *indistinctly* through a mineral.

Specular, having a smooth shining surface, like a mirror.

Spicular, in thorn-like fibres.

Stellated, when the fibres diverge from a common centre, and form a circle; star-like.

Striated, when the surface is marked by parallel lines.

Tabular, is applied to crystals which are nearly flat.

Translucent, when the substance transmits light; this is sometimes only observable on the edges of a mineral.

Transparent, when objects may be seen *distinctly* through the mineral.

Trapezoidal, having twenty-four faces, each a trapezium.

Truncated, when the edge or solid angle of a crystal appears to have been cut off.

Tuberous, exhibiting round bumps, bulb-like.

Vesicular, porous, sponge-like.

Vitreous, glassy.

M A W E ' S

NEW

DESCRIPTIVE CATALOGUE.



Metals.



PLATINA, PALLADIUM, IRIDIUM.

THESSE substances generally occur in small irregularly formed grains, flat, angular, or blunted, having apparently acquired this form by attrition; rarely crystallized. They are found with granular gold, in Peru, Mexico, and Brazil. When associated, it requires a well accustomed eye to discriminate one from the other.

PLATINA occurs in considerable quantities, and sometimes in rough lumps, larger than a pea. It resembles silver in color; it may be melted with arsenic, and is soluble in nitro-muriatic acid: when pure, it is the heaviest substance known.—*Sp. Gr.* 17; *purified*, 23.

PALLADIUM is very rare, and is found with Platina, with which it is alloyed. It is delicately striated, and has the color of lead: it melts with sulphur.

IRIDIUM is also extremely rare, and occurs with the preceding in small, flat, foliated grains, of a shining steel color: it is alloyed with Osmium, and melts with nitre. From either of the preceding it may be distinguished by its texture and extreme hardness; a grain placed on a polished iron surface, and struck with a hammer, will make an indention in both.

NATIVE PLATINA. In rough irregular lumps.
In flat, angular, or blunted grains.

PALLADIUM. In delicate folia, surface striated.

IRIDIUM. In shining grains, edges striated.
Imperfectly crystallized.

Oxide of Iridium. In black ferruginous grains.



GOLD.

TWO species are contained in this Genus, viz. Native Gold, and Electrum, or Gold and Silver combined.

Gold occurs in rounded lumps of various sizes; also disseminated in quartz and ferruginous substances, but generally in grains of irregular forms, and dust of a yellow or reddish yellow color. It is also found crystallized* in cubes, octahedrons, and many other forms, as well as foliated, ramified, capillary, reticulated, &c. It is very generally dispersed throughout the globe, and particu-

* The finest crystals known are in the Author's collection, which he brought from the gold mines of Brazil.

larly in South America; also in Ireland, Scotland, Cornwall, &c. but is never found pure in its natural state.

NATIVE GOLD is of a pale or deep yellow color, and sometimes tarnished, according as it is less or more alloyed with silver or copper. It is always ductile, easily melted, and does not change color in heated nitric acid. Its proper solvent is nitro-muriatic acid. *Sp. Gr.* 17 to 19.

ELECTRUM occurs in quartz, &c. in patches of a pale yellow color, and sometimes contains one third silver.

NATIVE GOLD. Massive, generally rounded.

Disseminated, embedded in quartz, &c.

Ramified, reticulated.

Crystallized in cubes, or modified.

Crystallized in octahedrons, or modified.

In rhombic dodecahedrons.

In aggregated crystals, confused.

In tetrahedrons, or serrated prisms.

Foliated.

Crystallized in three, four, six, or eight sided tables.

In delicate prisms, or capillary.

In coarse pieces, as broken from the matrix.

In granular, angular, and blunted grains.

Gold dust, yellow, or reddish yellow.

ELECTRUM. In superficial patches.

SILVER

IS rarely or never found pure; it is combined with many of the metals (particularly with lead) and two of the acids, which become volatilized by a continuance of the heat of the blow-pipe, and a bead of pure silver will remain. It is soluble in nitric acid. The mines of this metal are very numerous, but the most productive are those of Mexico and Peru; it is met with in Devonshire and Cornwall, and has been found in Scotland. It is also extracted from lead.

NATIVE SILVER occurs massive, ramose, capillary, reticulated, serrated, foliated, &c. and crystallized in various forms. Its color is nearly white, but is subject to tarnish resembling copper, and becomes blackish. It is ductile, a little harder than lead, and may easily be known by being soft and tough to the knife. *Sp. Gr.* 10.

Auriferous Silver contains a portion of gold. It generally occurs in superficial patches of a yellowish white color, and is sometimes tarnished.

ANTIMONIAL SILVER is found massive, and crystallized in striated cubes, but more generally disseminated. It has a bright white color, and contains from sixty to seventy per cent. of silver.

ARSENICAL SILVER is sometimes massive or disseminated, also crystallized. It has a dark metallic lustre, and is extremely heavy and brittle. It contains iron, and a large portion of arsenic.

BISMUTHIC SILVER occurs disseminated, and embedded, in fascicular crystals. It has a slight purple tint, and is alloyed with lead.

NATIVE SILVER. Massive.

Disseminated.

Crystallized in cubes, or variously modified.

In octahedrons, or modified.

In dodecahedrons.

Trapezoidal—with twenty-four faces.

In hexagonal prisms.

In four-sided prisms.

In three-sided prisms.

Tabular, in three or six sided tables.

Capillary, in delicate curls.

Ramose.

Foliated.

In delicate folia, superficial.

Variety.

Auriferous Silver. Superficial.

ANTIMONIAL SILVER. Massive.

Disseminated.

Crystallized in cubes.

ARSENICAL SILVER. Massive.

Crystallized or disseminated.

BISMUTHIC SILVER. Crystallized.

Disseminated, or fascicular.

Sulphurets of Silver.

THE following ores contain a larger portion of sulphur, with antimony, iron, copper, lead, &c. and less of silver

than the preceding; they are also brittle and harder, and more difficult to melt. They occur massive, disseminated, and crystallized. Their general color is that of polished iron after it has become dull, approaching to lead grey. They are the most common ores of silver, particularly in Europe, and present a great variety.

BLACK SULPHURET OF SILVER is found coating quartz, massive, disseminated, reticulated, and crystallized; it sometimes resembles soot, and is often associated with native silver in Calcareous Spar. It is of a dark color, approaching black, and has, when cut or broken, a metallic lustre. It is malleable, soft to the knife, and easily melts before the blow-pipe. It contains from sixty to seventy per cent. of silver, the remainder sulphur.

GREY SILVER ORE occurs massive, disseminated, also in regular and confused crystals. Its color is steel or lead grey. Its alloys are with difficulty volatilized before the blow-pipe.

DARK RED SILVER ORE. This beautiful mineral occurs massive, disseminated, dendritic, cellular, in superficial patches, and crystallized: it has a strong black-red metallic lustre. It contains about sixty per cent. of silver, with antimony and sulphur.

Light Red Silver Ore occurs massive, disseminated, dendritic, reniform, and crystallized. It is of a bright red color, beautifully translucent. It easily melts before the blow-pipe, emitting fumes of antimony and sulphur, leaving a bead of silver.

WHITE SILVER ORE is said to be lighter colored than lead ore; it contains a large portion of lead, antimony, and sulphur, with less than ten per cent. of silver, and is probably an argentiferous ore of lead.

SULPHURET OF SILVER. Massive and compact.

- Disseminated with quartz, &c.
- Crystallized in cubes or modified.
- In octahedrons, or variously modified.
- In garnet-formed dodecahedrons.
- Variety in aggregated crystals.
- Tabular, in three or six sided tables.
- Indeterminately crystallized.
- Variety.
- Pulverulent, sooty.

GREY SILVER ORE. Massive.

- Disseminated.
- Crystallized in six-sided prisms.
- Variety, with or without pyramid.
- In double six-sided pyramids.
- Tabular.
- Confusedly crystallized, aggregated
- Indeterminately crystallized.

DARK RED SILVER ORE. Massive.

- Crystallized in six-sided prisms.
- In six-sided prisms with pyramids.
- In six-sided prisms and double pyramids.
- Variety, pyramidal.
- In four or eight-sided prisms.
- Dendritic.
- Acicular.
- Capillary.
- Disseminated with pyrites.
- Variety, foliated.

Light Red Silver Ore. Massive.

Disseminated.

In equi-angular six-sided prisms.

Variety, with double pyramids.

In six-sided prisms, acuminated.

In double six-sided pyramids.

In acicular crystals.

WHITE SILVER ORE.

MURIATE of SILVER occurs massive, disseminated, in delicate veins and superficial layers; also crystallized. It is frequently dark brown, sometimes black, green of various shades, and yellow; the fresh fracture generally resembles gum arabic, but it soon acquires a metallic hue. It is easily indented by the nail, and melts even in the flame of a candle. It is mineralized by muriatic acid; whence its name: from being soft and translucent, it is also called Horn-silver.

CARBONATE OF SILVER is very scarce, and occurs in one mine only. It is of an iron grey color, heavy, and yields above seventy per cent. of silver, twelve of carbonic acid, with sulphur and antimony.

MURIATE OF SILVER. Massive.

Crystallized in cubes.

Variouly modified.

In octahedrons.

Modified.

Disseminated, yellow, green, or brown.

Superficial.

Earthy

Variety, radiated.

Variety.

CARBONATE OF SILVER.


MERCURY.

OF this mineral there are five different ores, viz. Native Mercury, Native Amalgam, Muriate or Horn Mercury, Hepatic or Brown Cinnabar, and Red Cinnabar.

Mercury generally occurs in a fluid state, and has a strong silver-white lustre. Its presence in any of its ores may be easily detected, by heating a small portion under a gold coin, or plate of polished copper; the mercury will be volatilized by the heat, and condensed by the cold metal, to which it will adhere in minute globules. The ores of mercury are confined to a few localities, but wherever they have been found, they occur in great abundance. The principal mines are situated in Almaden, Deux-ponts, Idria, and Peru. This metal is of great use in the arts, medicine, &c. It is employed

most extensively in South America in extracting the silver and gold from their various combinations.

NATIVE MERCURY occurs in minute or large globules, also disseminated in cinnabar, clay-slate, bituminous schist, sand-stone, &c. *Sp. Gr. above 13.*

NATIVE AMALGAM—Mercury and silver in various proportions, is sometimes crystallized, very heavy, and soft, and has a silver white color. The proportion of silver is generally above twenty per cent. When rubbed on a plate of polished metal, it leaves a silvery trace. Amalgam may be formed by uniting a globule of mercury with gold or silver leaf. *Sp. Gr. above 10.*

MURIATE OF MERCURY, or *Horn Mercury*, quicksilver combined with muriatic acid, occurs coating ores of mercury, and lining cavities, and sometimes crystallized. It has generally the appearance and color of gum-arabic. It is so soft that a pin may be stuck into it.

HEPATIC CINNABAR is found massive or disseminated; it is very heavy; and its fracture is fine grained, and sometimes slaty. Its color varies from red to brown.

RED CINNABAR occurs in beautiful diaphanous crystals, sometimes having a metallic lustre; and another variety, called *Native Vermilion*, is in the state of powder. Its color is dark or light red. It contains portions of sulphur and iron.

* * * The mercury of commerce is principally obtained from the two latter ores.

NATIVE MERCURY. Disseminated on sandstone.

In globules on bituminous schist, &c.

In globules with cinnabar, &c.

In globules with pyrites.

Variety.

NATIVE AMALGAM. Crystallized in octahedrons.

Crystallized in dodecahedrons, or modified.

Crystallized and variously modified.

Superficial, disseminated.

In capillary veins.

Variety.

MURIATE OF MERCURY. Spongeous, soft, disseminated.

Crystallized in cubes on ores of Mercury.

Variety, in aggregated crystals, confused.

Crystallized in octahedrons, or modified.

In capillary veins.

Superficially disseminated.

HEPATIC CINNABAR. Massive.

Slaty texture, curved, foliated.

Disseminated.

RED CINNABAR. Massive.

Crystallized in six-sided prisms.

In rhombs, or modified.

Variety.

Disseminated.

In tabular crystals.

In flat lenticular rhombs

Bright Red Cinnabar. Massive.

Disseminated.

Foliated.

Vermilion. Pulverulent.

COPPER—*Sp. Gr. 7.*

THIS metal is in such general use, that it is unnecessary to describe its appearance or color. It is easily dissolved in nitric acid, to which it communicates a green color, and may be precipitated in a metallic state by a rod of iron. The ores of copper are very numerous, and are found in almost all parts of the world, particularly in Cornwall, Chili, and Japan. When alloyed with zinc, it forms brass. Silver and gold coin are debased by it, and of jeweller's gold it often forms the greatest proportion.

NATIVE OR VIRGIN COPPER occurs massive, disseminated, and crystallized, and in a variety of other forms. It has a bright or dark red color, as it is less or more tarnished. It is extremely ductile, always soft to the knife, and, if scratched, exhibits a bright metallic lustre. It is found in great abundance in the mines of Cornwall, as well as in Chili, where it is accompanied by a small portion of gold.

NATIVE COPPER. Massive.

Foliated.

Ramose.

Crystallized in cubes, modified.

In four-sided prisms.

In hexagonal prisms.

In double six-sided pyramids, elongated.

In three-sided pyramids.

In aggregated crystals, indeterminate.

Dendritic, serrated.

Moss-like, interwoven.

Reticulated.

Disseminated.

Earthy.

Variety.

Sulphurets of Copper.

GREY COPPER—*Copper Glance*, occurs massive and crystallized, of a dark color, approaching bluish black; it is often tarnished, and has sometimes a metallic lustre. It is soft, has occurred feebly malleable, and is easily melted. It frequently contains above eighty per cent. of copper, with a small portion of sulphur, either with or without iron. To this species belongs the *Nail-head copper* of Cornwall, viz. two hexagonal pyramids, united at the base, truncated by a plane on the apex, supported on a long hexagonal prism, and sometimes variously modified.

Tennantite occurs massive, and crystallized, of a lead grey color, inclining to black, with shining surface; it is rather harder than the preceding, of which it is merely a variety.

Fahlertz Grey Copper occurs disseminated, compact, and crystallized, and is often covered with pyrites. It has a dull iron color, and contains antimony, iron, silver, and a portion of sulphur. The black variety I have placed with the silvers, as it is generally melted for that metal.

PURPLE OR VARIEGATED COPPER ORE is found massive, botryoidal, and crystallized. It has a beautiful iridescent appearance, exhibiting various shades of purple,

blue, green, &c. from which it is commonly called *Peacock-ore*. It is soft to the knife, and melts easily, emitting sulphurous vapors. It contains from fifty to sixty per cent. of copper, with iron and sulphur.

COPPER PYRITES occurs in great abundance, massive, disseminated, stalactitic, and crystallized, of a pale yellow color. It is harder and more difficult to melt than the preceding, but is easily dissolved in nitric acid. It contains from nine to twenty-five or thirty per cent. of copper, with iron, sulphur, and sometimes arsenic.

GREY COPPER ORE. Compact.

Variety, massive.

Crystallized in hexagonal prisms.

Variety, confusedly crystallized.

In double six-sided pyramids.

Variety tabular.

In dodecahedrons.

Variety, aggregated.

Tennantite. Massive.

In rhomboidal dodecahedrons.

In cubes.

In octahedrons, or modified.

Fahlertz Grey Copper. Compact.

Massive, with bright metallic lustre.

Disseminated.

Crystallized in tetrahedrons.

In tetrahedrons, modified on the angles.

Variety, modified on the edges.

Variety, aggregated.

VARIEGATED COPPER ORE. Massive.

Variety, color deep steel blue.

Botryoidal, mammillated, stalactitic.

Yellow Copper Ore.

Crystallized in cubes, inclining to curvilinear.

In cubes intersecting, or variously modified.

In aggregated crystals, iridescent.

Disseminated.

COPPER PYRITES. Massive.

Disseminated.

Botryoidal or stalactitic.

Crystallized in tetrahedrons.

In tetrahedrons, modified on the angles.

In tetrahedrons, modified on the edges.

In octahedrons, or modified.

In dodecahedrons, or modified.

Tabular, lamellar.

Indeterminately crystallized, aggregated.

Disseminated with lead ore.

BLACK COPPER ORE is sometimes loosely cohesive, soot-like; it is also massive and compact. Its color approaches black. It is very heavy, and easily fused, tinging borax green, which, when cold, becomes hollow, and appears like a scoria. The massive variety is soft, shows a metallic white streak, and is almost ductile. This ore has been called an oxide of copper.

BLACK COPPER ORE. Massive.

Variety, with metallic lustre.

Loosely cohesive.

Variety.

WHITE COPPER ORE—*Arsenical Copper*, is a rare mineral; it occurs massive and disseminated, and has a granular texture; it is heavy, brittle, and soft to the knife. Its color is yellowish white, with bright metallic lustre, or tarnished. Before the blow-pipe it exhales arsenical vapors.

ARSENICAL COPPER. Massive.

Disseminated, granular.

RED OXIDE OF COPPER occurs massive, foliated, crystallized, capillary, and earthy. Its color varies from bright red to dull brown, and has sometimes a metallic lustre. It easily fuses, and may be detected by nitric acid. It is generally associated with native copper.

Ferruginous Red Copper is found massive and earthy, It is heavy, soft to the knife, and has a dull red color. It contains a large portion of iron, and is more difficult to melt than the preceding.

RUBY COPPER ORE. Massive.

Disseminated.

Foliated, indeterminately crystallized.

In minute crystals, aggregated.

Crystallized in octahedrons.

Variety, modified on the angles.

Variety, modified on the edges.

Variety, grouped, reticulated.

In cubes, variously modified.

In bright, capillary red prisms.

Variety, crossing at right angles.

Variety, plumose.

Earthy, disseminated.

Variety, dull brown.

Brick Red Copper. Massive.

Earthy, often coated by green copper ore.

Variety, brown, ferruginous.

HYDRO-CARBONATE OF COPPER. This species occurs in foliated, globular masses, lamellar, disseminated, stalactitic, crystallized, and earthy, upon ferruginous substances. Its colors are various shades of blue, from light to dark. It is soft, effervesces with nitric acid; and, before the flame of the blow-pipe, tinges borax green, leaving a small bead of copper.

GREEN CARBONATE OF COPPER—*Malachite.* This beautiful species occurs in a variety of forms, massive, disseminated, botryoidal, stalactitic, crystallized, and coating red copper; also earthy: the fracture is zoned, fibrous or radiated. Its colors are of various shades, from pale to dark green, and sometimes even blue. Before the blow-pipe it burns with a green flame, and then melts into a bead of copper.

HYDRO CARBONATE OF COPPER. Massive.

Disseminated, foliated.

Crystallized in rhombic prisms.

Variouly modified.

In hexagonal tabular crystals.

Variety.

In four or eight-sided prisms.

Variety.

In diverging spicula.

Earthy Blue Copper Ore. Massive.

Variety, disseminated.

Fibrous.

Pitch-like.

Scoriaceous; sometimes black.

Botryoidal.

MALACHITE. Massive, and compact.

Cellular.

Mammillated, zoned.

Crystallized in hexagonal prisms.

In prisms with pyramids.

In octahedrons, coating red copper.

Variety, variously modified.

In rhombic dodecahedrons.

Variety, in aggregated crystals.

Variety, variously modified.

Tabular, in delicate four-sided crystals.

Blue variety.

Fibrous.

Variety, diverging.

Earthy.

Green and blue, disseminated.

SILICIOUS COPPER—*Crysocolla*, occurs massive, botryoidal, cellular, stalactitic, and incrusting native copper and ores of copper; it has often a slag-like appearance, and is very brittle. Its color varies from dark olive to white and blue green. It contains oxide of copper, silex, and water.

CRYSOCOLLA. Massive.

Botryoidal.

Stalactitic.

Variety.

Slag-like.

DIOPTASE is extremely rare, and is found crystallized upon limestone. It is of a beautiful emerald green color; contains thirty per cent. copper, forty lime, and twenty-eight silic.

DIOPTASE. Upon the matrix.

Crystallized in hexagonal prisms.

Variety, in rhombic dodecahedrons.

MURIATE OF COPPER occurs in compact radiated masses, lamellar prisms, and arenaceous. Its color is shining green and blackish green; before the blow-pipe it produces a beautiful and continued blue flame. It is found at Atacama, in Chili, and contains seventy per cent. copper, twelve muriatic acid, and fifteen water.

PHOSPHATE OF COPPER is found compact in mammillar concretions, and disseminated (on quartz or chalcidony), also crystallized. Its color is green, or dark green, with blackish spots. It burns with a green flame, melts into a fluid, which extends on the charcoal, and when cool becomes crystallized, and of a brownish red color. It contains sixty to seventy per cent. of copper, the residue phosphoric acid.

ARSENIATE OF COPPER. This interesting mineral occurs massive, disseminated, crystallized, fibrous, in mammillary concretions, asbestos-like, and earthy; and

has a foliated fracture. Its color presents various shades of green. Before the blow-pipe it swells, and easily melts, with slight explosions, emitting strong fumes of arsenic, and leaves a globule of copper.

Tabular Arseniate of Copper occurs in flat hexahedral crystals, the sides alternately inclining; color, beautiful green.

Octahedral Arseniate of Copper occurs in oblate crystals, of a sky blue or emerald green color.

Prismatic Arseniate of Copper. In prisms, of various shades of color, but commonly brown, or blackish-green.

MURIATE-OF COPPER: Massive.

Disseminated in ferruginous earth.

Crystallized in rhombic prisms.

Variety, variously modified.

Fibrous, stellated, diverging.

Arenaceous. (*Atacamite*).

PHOSPHATE OF COPPER. Compact.

Disseminated in delicate veins, &c.

Mammillar.

Fibrous, diverging.

Crystallized in rhombs.

In octahedrons.

Variety.

In rhombic prisms, with pyramids.

Variety.

In aggregated crystals, or curvilinear.

ARSENATE OF COPPER. Massive,

Foliated.

Crystallized in rhombic prisms

In rhombic prisms with dihedral pyramids.

Fibrous.

Disseminated.

Capillary.

In mammillary concretions.

Variety.

Hematitic, fracture silky.

Asbestos-like.

Earthy.

Tabular Arseniate of Copper.

— In flat six-sided crystals.

Octahedral Arseniate of Copper.

Crystallized in blue oblate octahedrons.

Variety, green.

Foliated, or disseminated.

Variety, aggregated.

In delicate veins.

Prismatic Arseniate of Copper. In three-sided prisms.

Variety, curvilinear, flat, foliated.

In four-sided prisms.

In rhombs, elongated.

Variety, diverging.

In delicate prisms, with pyramids.

Variety, in capillary prisms.

IRON.

THIS useful mineral is so generally distributed, that there are but few substances into which it does not enter, or associate with. It is found in all countries, in a variety of forms, and of different colors. Its presence may in general be detected by the magnet, especially after having been exposed to the flame of the blow-pipe or a red heat.

NATIVE IRON *is said* to have been met with massive, and in plates of a grey color, with a granular fracture, soft and malleable. Of this ore there are two varieties—

Terrestrial Native Iron, which is feebly malleable, and *Meteoric Native Iron*, which is found in large lumps, and coating earthy substances, of a rusty brown exterior. Its streak has a shining metallic lustre, and the fracture exhibits delicate, bright folia. These varieties are strongly magnetic, and contain nickel.

NATIVE IRON. Massive.

In laminæ.

Terrestrial Native Iron.

Meteoric Native Iron. Massive.

Cellular, containing olivine.

Coating earthy nodules.

In white folia, embedded.

Sulphurets of Iron.

IRON PYRITES occurs in great abundance, massive, crystallized, &c. It is brittle, and hard to the knife. The color is of various shades of yellow, tarnished, and sometimes beautifully iridescent.

Hepatic Pyrites, is found reniform, globular, &c. the fracture generally diverging. It is of a liver-brown color; the fresh fracture is yellowish.

Magnetic Pyrites, as its name implies, is attracted by the magnet; it occurs massive, embedded, and crystallized, and has generally the color of tarnished copper.

IRON PYRITES. Massive.

Granular.

Stalactitic, mammillated.

Radiated, diverging.

Disseminated.

Crystallized in cubes, smooth or striated.

Variety, variously modified.

In octahedrons.

In octahedrons modified, aggregated.

In dodecahedrons.

Variety, crystals striated.

In isocahedrons.

Capillary, embedded in calcareous spar.

Hepatic Pyrites. Crystallized in prisms.

Variety, aggregated crystals.

Nodular.

Variety.

Earthy.

Magnetic Pyrites. Massive.

Disseminated.

Crystallized.

In decomposition.

Oxides of Iron.

NATURAL LOADSTONE is massive, compact, earthy, and strongly magnetic. Its color is black or brown.

IRON GLANCE occurs massive, disseminated, and crystallized. Its color is that of shining steel, sometimes beautifully iridescent, especially in the specimens from Elba.

Micaceous Iron Ore is found in masses composed of aggregated and disseminated delicate shining laminæ. It is unctuous to the touch, and has a metallic-grey color, with a red tinge.

RED IRON ORE occurs sometimes compact, but generally in crusts or masses, which are more or less friable; the color is brownish red. Of this ore there are several varieties, viz.

Scaly Red Iron Ore occurs in loose scaly particles, and sometimes pulverulent, of a red color: it is extremely light, and has been called *Iron Froth*.

Compact Red Iron Glance is found massive, and sometimes in supposititious crystals, which have a feeble lustre; its color approaches black or reddish brown.

Red Hematites occurs in large masses, reniform, globular, botryoidal, stalactitic, &c. of a brownish-red color, and often fibrous, with metallic lustre.

Red Ochre occurs in friable masses, also coating ores of iron; its color is brownish red.

Reddle is found massive; it is adhesive, soft, unctuous to the touch, and soils the fingers: color dull red.

BROWN IRON ORE is more or less compact, and friable.

Scaly Brown Iron Ore is composed of delicate particles, and occurs sometimes with Brown Hematites.

Compact Brown Iron Ore occurs massive and crystallized; its color presents various shades of brown.

Brown Hematites is found stalactitic, reniform, or coralloidal, with a structure radiated, and silky lustre; color brownish black.

Brown Ochre occurs in dull earthy masses, soft, soiling the fingers: color, yellowish brown.

BLACK IRON ORE occurs massive, reniform, stalactitic, and in distinct concretions.

Black Hematites is also found massive and reniform, the fracture is delicately fibrous, with a strong metallic lustre.

CLAY IRON STONE occurs massive and reniform, and containing vegetable impressions; its color is dull brown of various shades. This is the common ore of England, and occurs in beds above coal.

Columnar Clay Iron Stone. In distinct long prisms, straight or curved; it is generally friable, and of a red-brown color; appears as having been exposed to heat.

Pea Iron Ore occurs aggregated, in pea-like concretions, with ochre; its color is light or dark brown.

Lenticular Iron Stone is in aggregated masses, composed of lenticular particles; color, yellowish or reddish brown.

Nodular Iron Stone occurs in spherical or reniform masses, composed of concentric layers enveloping loose earth, and sometimes, in the specimens from Brazil, gold and diamonds.

BOG IRON ORE occurs in amorphous or rounded masses, more or less compact; its color is brown of different shades, and approaching black. *Meadow* and *Morass Iron Ore* are varieties of this species, and are formed by deposits from stagnant water.

Pitchy Iron Ore occurs in soft, shining crusts, which resemble pitch.

Umber is a fine deposit, of a brown color; it is used as a pigment.

TITANIOUS IRON ORE is found massive, of a shining greyish black color; it contains oxide of Titanium.

NATURAL LOADSTONE. Massive.

Coarsely granular.

Variety.

Crystallized in octahedrons.

Arenaceous.

Minutely crystallized, granular.

Earthy, decomposed.

IRON GLANCE. Compact.

In flat double three-sided pyramids.

In three-sided pyramids, angles truncated.

Variety.

Variety, approaching octahedrons.

Variety.

In foliated hexagonal tables.

Variety.

In double six-sided pyramids.

Lenticular

Iron Mica. Massive.

In delicate six-sided tables.

Aggregated.

Disseminated.

RED IRON ORE. Compact.

In crusts, coating other substances.

Scaly Red Iron Ore. Encrusting red hematites.

Variety, earthy, pulverulent.

Compact Red Iron Glance. Massive.

In supposititious crystals.

Variety.

Red Hematites. Massive.

Reniform, mammillar.

Zoned, fracture radiated.

Variety, fibrous.

Red Ochre. Friable, with hematites.

Reddle. Massive.

BROWN IRON ORE. Massive and compact.

Friable.

Scaly Brown Iron Ore.

Compact Brown Iron Ore. In crusts, on iron ochre.

Brown Hematites. Massive.

Reniform, mammillated.

Fibrous or radiated.

Brown Ochre. Massive.

BLACK IRON ORE. Massive.

In distinct concretions.

Black Hematites. Massive.
Variety, stalactitic.

CLAY IRON STONE. Massive.
Variety, with vegetable impressions.
Containing veins of calcareous spar. (*Septaria*).
Variety.

Columnar Clay Iron Stone. In long prisms.

Pea Iron Ore. In pea-like concretions.

Lenticular Clay Iron Stone. Massive.
Granular, aggregated.
Earthy.

Nodular Iron Stone. Reniform.
In hollow nodules.
Variety, enveloping Gold or Diamonds.
Variety, showing several stages of decomposition.

BOG IRON ORE. Compact.
Earthy.

Pitchy Iron Ore. Massive.
Incrusting.

Umber. Massive.
Earthy.

TITANIOUS IRON ORE. Massive.
Variety, fracture granular.

CARBONATE OF IRON, or *Sparry Iron Ore*, occurs amorphous, crystallized, and lamellar; structure foliated; its color is yellowish white, brown, or reddish brown.

PHOSPHATE OF IRON occurs crystallized on pyrites, (Cornwall); also in delicate folia, stellated on earthy carbonate of lime, (Derbyshire); it has a Prussian-blue color, is soft, and easily melts.

Earthy Blue Iron Ore occurs in friable masses, sometimes compact, and coating ores of iron; wood is often impregnated with it: color, light blue.

CHROMATE OF IRON is found in aggregated concretions; massive, crystallized, and in shining grains; its color approaches black. It is slightly magnetic; before the blowpipe it gives a green flame, and tinges borax of the same color.

ARSENATE OF IRON occurs in small aggregated crystals. Its color is generally green. It easily melts, emitting fumes of arsenic.

Cupreous Arseniate of Iron occurs in aggregated crystals, of a bluish-green color.

SPARRY IRON ORE. Amorphous.

Crystallized in hexagonal prisms.

In rhombs, or double pyramids.

Lenticular.

Variety.

Red Sparry Iron Ore. Rhombic.

In aggregated crystals.

Granular.

Yellow Sparry Iron Ore. Massive.

Indeterminately crystallized.

PHOSPHATE OF IRON. Crystallized on pyrites.

In rhombic prisms.

In hexagonal prisms.

Variouly modified.

Foliated.

Variety.

Tabular.

Disseminated.

Earthy Phosphate of Iron. Massive.

Pulverulent.

Variety.

CHROMATE OF IRON. Massive.

Disseminated.

Crystallized in octahedrons.

Variety.

Granular.

ARSENIATE OF IRON. Crystallized in cubes.

Variety, color brown.

Cupreous Arseniate of Iron. In rhombic crystals.

Variety, aggregated, globular.

MANGANESE.

THE ores of Manganese present great diversity in external characters; but its presence in any substance may easily be detected by melting it with borax and a little nitre, which will form a violet glass. It is much used in the arts, particularly in bleaching, in making glass, and for producing oxygen gas. It is generally combined with oxygen, and more rarely with sulphur, and carbonic or phosphoric acids. It occurs in great abundance in Devonshire, Cornwall, Derbyshire, and Scotland, often associated with ores of iron.

GREY MANGANESE occurs massive, crystallized, foliated, compact, and earthy. The color is steel grey or iron black; it has a metallic lustre; which characters generally pervade all the varieties.

Radiated Grey Manganese occurs stalactitic and crystallized; it is often tarnished.

Foliated Grey Manganese is found massive, disseminated, and crystallized, with a foliated fracture.

Compact Grey Manganese occurs massive and botryoidal; it is soft, and stains the fingers.

Earthy Grey Manganese occurs massive and in crusts, sometimes friable. It is used for producing oxygen gas, and effervesces with the acids.

Wad, a variety of the preceding, is found stalactitic, botryoidal, reticulated, and pulverulent: it is frequently associated with other minerals, and is used as a pigment.

BLACK MANGANESE is found massive, disseminated, or crystallized, also in nests with quartz, &c. lustre, that of iron tarnished.

GREY MANGANESE. Massive.

In delicate acicular crystals.

Variety.

Radiated Grey Manganese. Stalactitic.

In small crystals, structure radiated or fibrous.

Foliated Grey Manganese. Massive.

Crystallized in acicular prisms.

Compact Grey Manganese. Amorphous.

Stalactitic.

Earthy Grey Manganese. Massive.

In crusts, disseminated.

Wad. Massive.

Fibrous, color brown.

Pulverulent.

BLACK MANGANESE. Massive.

Stalactitic.

Botryoidal.

Disseminated.

Crystallized in elongated octahedrons.

Dendritic, on indurated marl, &c.

Variety.

SULPHURET OF MANGANESE OCCURS massive, of a dark color, approaching black: its fracture has a shining metallic lustre, which soon becomes tarnished; in melting, it emits sulphurous vapors.

PHOSPHATE OF MANGANESE OCCURS massive, fracture uneven or foliated. It has a shining black or brownish

color, is hard, and melts before the blow-pipe into a black enamel.

CUPREOUS MANGANESE occurs massive, botryoidal or disseminated, of a bluish black color; with a shining resinous lustre; and conchoidal fracture.

CARBONATE OF MANGANESE is composed of carbonate of lime, with oxide of manganese; and occurs massive, crystallized, and fibrous; fracture, foliated; color, pale rose-red.

Compact Red Manganese occurs combined with feldspar, is heavy, and has generally a rose-red color.

Fibrous Red Manganese occurs reticulated, in delicate acicular crystals; color, brownish red.

SULPHURET OF MANGANESE. Massive.

PHOSPHATE OF MANGANESE. Massive.

CUPREOUS MANGANESE. Massive,

Botryoidal.

Disseminated.

CARBONATE OF MANGANESE. Massive,

Crystallized.

Fibrous,

Compact Red Manganese. Massive.

Variety, with black veins.

Fibrous Carbonate of Manganese. Reticulated.

Embedded.

Stellated.

TITANIUM

GENERALLY occurs combined with the oxides of other metals. It is found in various parts, in alluvial deposits, in the form of black grains; also imbedded and crystallized. It is almost infusible, but when reduced, it has a dark copper-red color.

MENACHANITE has a black sand-like appearance, resembling gunpowder. It is feebly attracted by the magnet, and is infusible by the blowpipe.

ISERINE occurs in larger grains, and more spherical than the preceding; the color is brownish black.

NIGRINE is also found in rounded or angular grains; its color approaches black, its lustre is semi-metallic; it is not attracted by the magnet.

SPHENE occurs in amorphous masses, in grains, and crystals, also imbedded; it presents a great variety of colors, but they may in general be referred to brown or yellow; one variety has a green or greyish-green tinge, with a shining lustre, and often coated by chlorite.

RUTILE is very heavy, and occurs massive, and in beautiful striated and capillary prisms, intersecting each other; of a brown and red color, also gold-yellow. A variety of this species, from its hair-like appearance, has been called *Cheveux de Venus*.

OCTAHEDRITE occurs in elongated octahedrons, which are often variously modified, of an indigo-blue color: it is generally associated with adularia and quartz.

CRICHTONITE is found crystallized in very acute rhomboids, not unlike specular iron ore; it is harder than Octahedrite, and is generally associated with it.

MENACHANITE. In grains.

ISERINE. In spherical grains.

NIGRINE. In grains.

In rolled pieces.

BROWN SPHENE. Imbedded.

In very oblique four-sided prisms.

Variety, with dihedral summits.

Pale-green Sphene. Imbedded.

In oblique four-sided prisms.

Variety.

In oblique flat rhombs.

In double crystals, canaliculated.

In blade-like crystals.

RUTILE. Massive.

Disseminated.

In striated prisms.

Variety, geniculated.

In aggregated crystals.

In capillary prisms, embedded in crystal.

Variety, curved. (*Cheveux de Venus*).

Acicular. (*Fleche d'Amour*).

Variety, reticulated.

OCTAHEDRITE. In elongated octahedrons.

Variety, solid angles truncated.

Variety, modified.

CRICHTONITE. In acute rhomboids.

LEAD.

THIS is one of the most abundant of metals, and occurs in large and small veins, in almost every rock formation, combined with sulphur, oxygen, and many of the acids, and generally contains silver. The ores of lead are easily reduced by the blow-pipe, and dissolved in nitric acid. Its uses in the arts are numerous and well known. There are many mines of this metal in England; the principal are situated in Northumberland, about Alston Moor; Durham; the West Riding of Yorkshire; Matlock, and throughout the Peak in Derbyshire; Devonshire, and Cornwall.

The ores of Devonshire and Cornwall are particularly rich in Silver.

SULPHURET OF LEAD—*Galena*, occurs massive, in veins, and crystallized; its structure is foliated, fibrous, granular, or compact. It has a bright lead-grey color, sometimes beautifully iridescent. It melts easily before the blow-pipe, emitting fumes of sulphur.

Blue Lead Ore occurs massive, and in crystals, with a rough surface, sometimes pulverulent. It melts before the blow-pipe, burning with a bluish flame. It is rare.

Antimonial Lead Ore occurs reticulated and crystallized. Its color approaches tin-white; it contains lead, a large portion of antimony, and sulphur.

Triple Sulphuret of Lead—*Bournonite*, occurs massive and crystallized. Its color is shining steel-grey. It is composed of sulphur, lead, antimony, and copper; and derives its name from the discoverer, Count Bournon,

Cobaltic Lead Ore is found disseminated with ores of Cobalt, in minute aggregated crystals, and of a grey color.

GALENA. Massive.

Disseminated.

Fibrous, or striated.

Granular, fine or coarse.

Foliated.

Crystallized in cubes.

Variety, modified.

In octahedrons.

Variety, modified.

Variety, argentiferous.

In brilliant aggregated prisms.

Variety, reticulated and indeterminately formed.

Specular. (*Slickenside*).

Blue Lead Ore. Massive.

Crystallized in hexagonal prisms.

Pulverulent.

Variety, coating Galena.—Derbyshire.

Antimoniated Lead Ore. Reticulated.

In aggregated crystals.

Bournonite. Massive.

Crystallized in four-sided prisms.

Tabular, variously modified.

Variety, disseminated.

Cobaltic Lead Ore. In minute crystals.

Variety, disseminated.

CARBONATE OF LEAD—*Sparry Lead Ore*, occurs in compact masses, crystallized, fibrous, laminar, and earthy. It is more or less translucent, and in color presents various shades of white; it is easily reduced to a globule of lead.

Compact Carbonate of Lead occurs massive and crystallized. It is semi-translucent; color, white or cream-yellow, sometimes approaching a metallic lustre; and is commonly associated with galena.

Earthy Carbonate of Lead occurs massive, scaly, or coarsely granular, and friable, sometimes presenting a glistening fracture; its color is brown, green, or blue, of various shades.

Black Lead Ore occurs massive, disseminated, corroded, cellular, and crystallized. Its color is bluish or greyish black.

MURIATE OF LEAD is extremely rare, and occurs crystallized; color, wine-yellow, of various shades.

PHOSPHATE OF LEAD occurs massive, stalactitic, reniform, and crystallized. Its color varies from green to brown and yellow. It is very heavy, and has a glistening fracture.

ARSENATE OF LEAD occurs reniform, crystallized, and capillary. It has a yellowish brown color, and easily fuses, emitting arsenical fumes.

SULPHATE OF LEAD occurs disseminated and crystallized, of a yellowish-grey color, and is very easily fused. It also occurs reniform in Siberia.

MOLYBDATE OF LEAD occurs crystallized, and very rarely massive. Its color is some shade of yellow.

CHROMATE OF LEAD—*Red Lead*, occurs massive, disseminated, and crystallized. It has a beautiful red color, and a splendid exterior.

Green Chromate of Lead occurs earthy and crystallized, usually associated with red lead and manganese.

CARBONATE OF LEAD.

Compact Carbonate of Lead. Massive.

Laminated, micaceous.

Foliated.

Crystallized in three or four-sided prisms.

Variety, canaliculated.

In six-sided prisms.

Variety, modified.

Variety.

In double hexagonal pyramids.

Variety, modified.

In aggregated acicular crystals, fibrous.

Variety, coated by green or blue carbonate of copper.

Variety, coated with brown oxide of iron.

Earthy Carbonate of Lead. Massive.

Scaly, or coarsely granular.

Friable, earthy.

Black Lead Ore. Massive.

Crystallized in six-sided prisms.

Cellular.

Coating carbonate of lead.

MURIATE OF LEAD. In rectangular four-sided prisms.

Variety, angles truncated.

Variety, edges truncated.

Variety, with pyramids.

PHOSPHATE OF LEAD. Massive.

Disseminated.

Crystallized in six-sided prisms.

Variety, modified.

In acicular crystals.

Variety, dendritic.

Botryoidal.

Brown Phosphate of Lead. Massive.

Crystallized in hexagonal prisms.

In double hexagonal pyramids.

Acicular.

ARSENIATE OF LEAD. In reniform masses.

Crystallized in hexagonal prisms.

Variety, in double pyramids.

In capillary fibres.

Earthy, in crusts.

Variety.

SULPHATE OF LEAD. Massive.

Crystallized in octahedrons.

Variety, minutely crystallized.

Disseminated.

Variety, cuneiform.

Tabular.

Reniform Sulphate of Lead. Massive.*Earthy Sulphate of Lead.*

MOLYBDATE OF LEAD. Massive.

Crystallized in perfect octahedrons.

Variety, modified.

In four or eight-sided tables.

Variety, modified.

CHROMATE OF LEAD. Massive.

Crystallized in four-sided prisms.

Variety, with pyramids.

Variety, modified.

In obtuse rhombs.

Pulverulent.

Green Chromate of Lead. Earthy.

In acicular crystals.

ZINC.

THE ores of this metal are generally found associated with lead. They are combined with sulphur and oxygen, and with carbonic and sulphuric acids. From the external appearance of some of the ores, the presence of a metal would not be suspected; but it may readily be discovered by first roasting the ore, and then fusing it with copper filings, with which it will form brass:—this is one of the uses to which this metal is applied.

RED OXIDE OF ZINC occurs massive and disseminated, also indeterminately crystallized. Its color is red, tinged with yellow, or brownish.

ELECTRIC CALAMINE has a stony appearance, and occurs massive, stalactitic, and botryoidal; it is very hard and heavy, and becomes electric by heat: the color is dull grey.

CALAMINE occurs massive, stalactitic, and crystallized, also compact and earthy: its colors are yellowish-grey and brown, sometimes green. A variety with a striated diverging fracture, and resinous lustre, contains cadmium, and occurs in a mine near Matlock.

Cupriferous Calamine is composed of delicate diverging spicula, with a silky lustre; its green color is owing to the presence of carbonate of copper.

SULPHURET OF ZINC—*Blende*, occurs massive, disseminated, and seldom regularly crystallized: its colors are yellow, brown, and black, of various shades. It accompanies almost every metal; some varieties become phosphorescent by friction.

Yellow Blende occurs massive and crystallized, of a resin-yellow color; its fracture is foliated.

Brown Blende differs from the preceding only in color, which is reddish-brown.

Black Blende has also the same forms, and is of a shining or dull black color.

Green Blende occurs disseminated with galena, and has a shining metallic lustre.

RED OXIDE OF ZINC. Massive.

Disseminated.

Indeterminately crystallized.

ELECTRIC CALAMINE. Massive.

Stalactitic or botryoidal.

Crystallized in flat six-sided prisms.

Variety.

In acute octahedrons.

Variety.

CALAMINE. Massive.

Stalactitic, or reniform.

Crystallized in obtuse rhomboids.

Variety, in aggregated crystals.

In four-sided tables.

Striated, composed of delicate prisms.

Compact.

Coating calcareous spar or fluor.

Variety, in supposititious crystals.

Earthy.

Cupriferous Calamine. In capillary tufts.

SULPHURET OF ZINC. Massive and foliated.

Disseminated.

In minute shining crystals.

Yellow Blende. Massive and foliated.

Crystallized in dodecahedrons.

In twin crystals.

Brown Blende. Massive.

Crystallized in tetrahedrons.

In twin crystals.

In octahedrons.

In dodecahedrons.

In aggregated crystals.

Disseminated.

Black Blende. Massive.

Crystallized in tetrahedrons.

In octahedrons.

In dodecahedrons.

In aggregated crystals.

In double crystals.

Green Blende. Disseminated with galena.

TIN.

THIS genus contains only three species; its localities are not numerous, but wherever it has been found, it occurs in abundance. It is fused with difficulty.

SULPHURET OF TIN occurs massive and disseminated; its color approaches steel-grey, with shades of yellow, and has a metallic glistening lustre. It is peculiar to Cornwall, and is often called *Bell-metal Ore*.

OXIDE OF TIN is found massive, disseminated, and crystallized, also in rounded lumps, when it is called *Stream Tin*: its color is dark brownish black, and is extremely heavy.

Wood Tin occurs in small mammillated masses, has a diverging, fibrous structure: its color varies from red to brown. It is very heavy, and occurs in alluvial soil*.

SULPHURET OF TIN. Massive.

Disseminated.

OXIDE OF TIN. Massive.

Disseminated in clay-slate.

Variety, in quartz or granite.

Crystallized in octahedrons.

Crystallized in four-sided prisms.

Variety, with pyramids.

* A remarkably fine piece, beautifully zoned, and weighing 10½ ounces, from Mexico, was once in the Author's possession.

In double six-sided pyramids.

Variety, modified.

In twin crystals.

In aggregated crystals.

In delicate capillary crystals.

Detached crystals, broken from the matrix.

Wood Tin. In mammillary masses.

In fragments, zoned, diverging.

Variety, wood-like.

Variety, in globular concretions, embedded.

Variety.

BISMUTH.

IS not found in great abundance; it occurs in veins with silver, cobalt, arsenic, &c. It is extremely easy of fusion, and is used in making pewter, solder, &c.

NATIVE BISMUTH occurs massive and crystallized. It has a whitish metallic lustre, and its fresh fracture is beautifully iridescent, resembling the hues of a pigeon's neck.

SULPHURET OF BISMUTH occurs massive, disseminated, and in delicate crystals, often acicular. The color approaches tin-white, or tarnished with a yellow tinge; it melts in the flame of a candle.

NEEDLE ORE is found amorphous and in grey acicular crystals, embedded in quartz, and often surrounding wire-like gold.

CUPREOUS BISMUTH ORE is found massive and disseminated; its color is lead-grey, approaching tin-white. It contains Bismuth, copper, and sulphur.

BISMUTH OCHRE is found in an earthy state, with Native Bismuth and ores of Cobalt. Its color is yellowish and greenish grey.

NATIVE BISMUTH. Massive.

Disseminated.

Crystallized in octahedrons.

In double three-sided pyramids.

In tabular crystals, striated.

Granular.

SULPHURET OF BISMUTH. Massive.

Disseminated.

In delicate acicular crystals.

NEEDLE ORE. Amorphous.

In striated prisms.

Variety, covering filiform gold.

Variety, disseminated in quartz.

CUPREOUS BISMUTH ORE. Massive.

Disseminated.

BISMUTH OCHRE. Earthy.

Variety.

TELLURIUM

HAS only been found in Transylvania, where it occurs in delicate veins, and is alloyed with gold, silver, &c. Its lustre varies from bright to that of tarnished iron. In nitric acid it forms a limpid solution, and easily melts before the blow-pipe, emitting a peculiar pungent odor.

NATIVE TELLURIUM occurs massive and disseminated; it is fine grained, generally of a white-color, with a metallic lustre.

Graphic Tellurium occurs foliated, crystallized, and reticulated, but is generally superficially distributed on quartz, with a moss-like appearance.

Yellow Tellurium is often accompanied by gold, manganese, &c. It is white, with a yellow tinge, and sometimes slightly iridescent.

Black Tellurium occurs generally foliated; and crystallized, of an iron-black color; it is soft and flexible. This variety contains about 10 per cent. gold.

NATIVE TELLURIUM. Massive, granular.

Disseminated.

Graphic Tellurium. Superficially disseminated.

Crystallized in four-sided prisms.

In six-sided prisms, acuminated.

Foliated.

Yellow Tellurium. Disseminated.

Variety.

Black Tellurium. Massive.

In delicate folia, disseminated.

In tabular crystals.

ANTIMONY

IS found in veins, and generally accompanied by Blende. It often resembles Galena, but is not so heavy: it is mineralized by arsenic, sulphur, and oxygen; it easily melts, discharging thick white vapors. It is used in type-metal, and in various medicinal preparations.

NATIVE ANTIMONY is found massive, rarely distinctly crystallized: it has a shining tin-white color, with a granular and foliated fracture. It is found in Dauphiny.

GREY ANTIMONY occurs massive, disseminated, and crystallized, with a compact, radiated, and foliated structure; it has a shining lead color, sometimes iridescent.

Nickeliferous Antimonial Ore occurs massive and disseminated; it has the color of steel or lead, and is often tarnished. It is harder than the preceding varieties, and contains antimony, nickel, arsenic, sulphur, iron, and lead.

OXIDE OF ANTIMONY—*White Antimony*, occurs crystallized, of a white or yellowish white color.

Antimony Ochre is found earthy, of a yellow color of different shades, upon ores of antimony.

RED ANTIMONY occurs in amorphous masses, and in delicate crystals, of a deep red or purple color.

Tinder Ore is an earthy variety of the preceding; the color is reddish brown; structure tinder-like.

NATIVE ANTIMONY. Massive, granular, or foliated.
Indeterminately crystallized.

GREY ANTIMONY. Massive.

- Disseminated.
- Compact.
- Fibrous.
- Foliated.
- Crystallized in oblique prisms.
- In acicular crystals.
- Variety, iridescent.
- Capillary, approaching black.
- Plumose, coating quartz, &c.
- Variety.

Nickeliferous Antimonial Ore. Massive.

- Disseminated.

WHITE ANTIMONY. In flat four-sided tables.

- In acicular crystals.
- Variety, stellated.

Antimony Ochre. Massive.

- Variety, earthy.

RED ANTIMONY. Amorphous.

- In delicate crystals.
- Variety, stellated.

Tinder Ore. Earthy.

- Friable.

MOLYBDENA.

THIS genus contains but one species, which is combined with sulphur. It is with great difficulty reduced to a metallic state.

MOLYBDENA occurs massive, disseminated, and crystallized, and in flexible laminæ. It has the color of black lead, and leaves a greenish streak on porcelain.

Molybdena Ochre is rare; it is found disseminated, and coating molybdena, of a yellow color.

MOLYBDENA. Massive.

Disseminated.

Crystallized in hexahedral tables.

Variety, acuminated.

Variety, in hexagonal plates.

Molybdena Ochre. Disseminated.

Friable.

Encrusting molybdena.

COBALT.

THE ores of this metal are found in the primitive and alluvial formations. Before the blow-pipe it emits arsenical vapours, and tinges borax intense blue. It is used in enamel painting; the blue on earthenware and porcelain is produced by this substance; it is found in Cornwall, but the best is imported from Saxony.

ARSENICAL COBALT occurs massive, disseminated, reticulated, and crystallized; the color of the fracture is tin-white, which tarnishes by exposure.

GREY COBALT occurs massive and disseminated. It has a grey color; this like the preceding species becomes tarnished by exposure.

COBALT GLANCE occurs massive, disseminated, and crystallized, of a shining white color. The finest crystals are found in Sweden, and are of great variety.

SULPHURET OF COBALT occurs massive and disseminated, sometimes botryoidal, of a pale steel-grey color, which becomes reddish when tarnished.

OXIDE OF COBALT is generally found friable, of various colors, from brownish black to yellowish grey.

Black Cobalt Ochre is composed of dull black particles, earthy, and often accompanied by brown and yellow varieties, which occur in alluvial soil.

RED COBALT, or *Arseniate of Cobalt*, is found in small botryoidal masses, and in velvet-like balls, frequently efflorescent, and of a reddish white or peach-red color.

Radiated Red Cobalt occurs disseminated and crystallized, in beautiful folia, of a peach-red color.

Slaggy Red Cobalt occurs in crusts of a brownish red color, and has a scoriaceous appearance.

ARSENICAL COBALT. Massive.

Disseminated.
Reticulated.
Crystallized in cubes.
In octahedrons.
Variety, aggregated.

GREY COBALT. Massive.

Disseminated.

COBALT GLANCE. Massive.

Disseminated.
Crystallized in cubes.
Variety, modified.
In octahedrons.
Variety, modified.
In dodecahedrons.
Variety, pentagonal.
In icosahedrons.
In detached crystals.
Variously modified.

SULPHURET OF COBALT. Massive.

Disseminated.
Botryoidal.

OXIDE OF COBALT. Earthy.

Black Cobalt Ochre. Earthy.

Brown Cobalt Ochre. Amorphous.

Yellow Cobalt Ochre. Amorphous.
Variety, corroded.

RED COBALT. In masses.
Disseminated.
Botryoidal.

Radiated Red Cobalt. Massive.
Disseminated.
Crystallized in flat four-sided prisms.
In double six-sided pyramids.
Acicular, radiated.

Slaggy Red Cobalt Ochre. In crusts.
Coating ores of cobalt.

NICKEL.

IS not very abundant. It is found in combination with meteoric iron, and associated with copper. It is difficult to melt, and emits arsenical fumes; it gives a green color to nitric acid. With copper it forms *Petit Or*.

NATIVE NICKEL is found in long capillary crystals, of a brass-yellow color, and tarnished.

ARSENICAL NICKEL—*Copper Nickel*, occurs in compact amorphous masses, which have a reddish copper color.

BLACK ORE OF NICKEL is said to be an earthy substance, of a blackish color, which gives a green solution in nitric acid.

OXIDE OF NICKEL—*Nickel Ochre*, occurs earthy, and in efflorescences, of a light-green color.

NATIVE NICKEL. In capillary crystals, intersecting.

In diverging groups, or reticulated.

COPPER NICKEL. Massive.

Foliated.

Variety, with nickel ochre.

Disseminated.

BLACK ORE OF NICKEL. Earthy.

NICKEL OCHRE. Earthy.

Coating ores of nickel.

ARSENIC.

THIS mineral is very generally diffused. It is found combined with sulphur and oxygen, and often enters into ores of other metals, which have been already described. The odor of garlic, which it emits when struck with a hammer, or heated before the blow-pipe, together with its *rapid* volatilization, will distinguish it from other metallic ores.

NATIVE ARSENIC occurs in amorphous or tuberosc masses, also in plates. Its fresh fracture has a tin-white color, which soon becomes blackish; it has a bright metallic lustre, with a foliated, fibrous, or granular structure, and is very heavy.

ARSENICAL PYRITES occurs massive, disseminated, and crystallized, generally of a pale yellow color; it is sometimes argentiferous, and of a silver-white color. It melts easily, emitting fumes of arsenic and sulphur.

SULPHURET OF ARSENIC—*Realgar*, occurs massive, disseminated, and crystallized, and sometimes earthy, of a scarlet or orange-red color. It melts in the flame of a candle, and is extremely friable.

Orpiment occurs massive, structure foliated or granular, and in small crystals; it has a shining yellow color and a foliated or slaty fracture; it is soft and flexible.

NATIVE ARSENIC. Massive.

Reniform.

Laminated.

ARSENICAL PYRITES. Massive.

Disseminated.

Crystallized in oblique four-sided prisms.

Variety, modified.

In flat octahedrons.

Variety, in aggregated crystals

Argentiferous. Disseminated.

Embedded in lithomarge.

Variety.

SULPURET OF ARSENIC—*Realgar.* Massive.

Disseminated.

Crystallized in rhombic four-sided prisms.

Variety, variously modified.

In friable concretions.

Orpiment. Massive, foliated.

In minute crystals.

In granular concretions.

OXIDE OF ARSENIC occurs stalactitic, crystallized, capillary, and earthy; color, various shades of white.

PHARMACOLITE occurs in minute cotton-like balls; its color is generally reddish-white, and its fracture stellular. Arsenic acid 50, lime 25, and water.

OXIDE OF ARSENIC. Stalactitic.

Botryoidal.

Crystallized in beautiful octahedrons.

In delicate tabular crystals.

Capillary, in snow-white silky filaments.

Earthy, encrusting other substances.

PHARMACOLITE. In balls.

In aggregated capillary crystals.

TUNGSTEN

IS found combined with oxygen, lime, and iron. It is associated with Tin in primitive rocks, is infusible before the blow-pipe, and nearly insoluble in the acids. It is extremely heavy.

TUNGSTEN occurs massive and crystallized, of a dull white color, and shining lustre. It contains acid of tungsten 60 to 70, with lime.

WOLFRAM occurs massive, disseminated, and crystallized; it has a black shining metallic hue, structure foliated, and yields a *red* streak to the knife. It contains tungstic acid 60 to 70, with iron and manganese.

TUNGSTEN. Massive.

Crystallized in octahedrons.

Variety, modified, detached, or aggregated.

In four-sided tables.

Aggregated, coating other substances.

WOLFRAM. Massive.

Disseminated.

Crystallized in four-sided prisms.

Variety, tabular.

Variety, modified.

Variety, embedded.

URANIUM.

THIS beautiful mineral has only been found combined with oxygen. Both of its species are infusible, but easily dissolved in diluted nitric acid, to which they communicate a yellow color.

BLACK OXIDE OF URANIUM—*Pitch Ore*, occurs massive, of a black color; it is extremely heavy, and yields to the knife. It contains oxide of uranium 60 to 70, with lead and iron.

GREEN OXIDE OF URANIUM—*Uranite*, occurs in delicate and beautiful crystals, generally aggregated, of an emerald-green color.

Uranium Ochre occurs in small nodular masses, and in crusts frequently upon pitch ore; it has a yellow color of various shades, is soft, and breaks with a glimmering lustre.

PITCH ORE. Massive.

Variety, on uranium ochre.

Variety.

URANITE. Crystallized in four-sided tables.

Variety, detached.

Variety, elegantly grouped.

Variety, aggregated.

Variety, pyramidal.

Variety.

Uranium Ochre. In small nodular masses.

Pulverulent, or disseminated with pitch ore.

TANTALUM—*Columbium*.

SINCE the time Mr. Hatchett discovered this mineral, in a specimen from America, it has not been met with, till recently. It is found combined with oxygen, and an earth called Yttria, and is very hard.

TANTALITE occurs imbedded, in striated fragments, and crystallized. It is of a black color, and resembles wolfram, but its streak is *brown* or *grey*. It contains tantalum 60 to 90, with iron and manganese.

Ytthro-tantalite occurs in flat or angular masses, sometimes imbedded; color, black. It contains tantalum, yttria, and iron.

GADOLINITE occurs massive and disseminated, generally imbedded in quartz; color, black, often red-brown on the edges*.

TANTALITE. Imbedded in striated masses in granite.
Crystallized in irregular prisms.

Ytthro-tantalite. Imbedded in oblique prisms.
In flat angular pieces.

GADOLINITE. Massive.
Disseminated in particles, embedded.
Variety, encrusted.

*This mineral would perhaps be better removed to the class of earths, its most characteristic ingredient being Yttria.

CERIUM.

OF this mineral there are only two species, both oxides. They are infusible before the blow-pipe.

CERITE occurs massive and disseminated, of a pale lilac, or deep reddish brown color.

ALLANITE occurs massive and crystallized, of a brownish black color; it is often accompanied by tremolite. Oxide of cerium 34, iron 25, silica 35, with lime and alumine.

CERITE. Massive.

Disseminated.

Variety.

ALLANITE. Massive.

Crystallized in four-sided prisms.

In six-sided prisms.

Variety, aggregated.

CHROME

HAS only lately been found in the state of an oxide. It occurs massive or earthy, of a dull green color. It is with difficulty reduced to a metallic state; to glass it communicates a bright and permanent green.

SILICIOUS OXIDE OF CHROME. Massive.

Earthy.

SELENUM

IS a new metal, which has lately been discovered in an ore of copper from the Hartz.

SELENIURET OF COPPER has a black appearance, and occurs disseminated in patches in foliated carbonate of lime, not unlike schorl in white feldspar.

CADMIUM

IS also a newly discovered metal, and much resembles tin. It is extracted from a striated black blende, found in the Hartz, also from two varieties of calamine found in Derbyshire.

CADMIFEROUS BLENDE. In striated masses.

Cadmiferous Calamine. Stellated.

Variety, crystallized.

Metallic Salts.

THE substances of this class have metallic bases, and easily dissolve in water.

SULPHATE OF IRON—*Copperas*, occurs massive, crystallized, and stalactitic; color, generally green or brown. It is formed by the absorption of oxygen, during the decomposition of iron pyrites.

SULPHATE OF COPPER—*Blue Vitriol*, occurs massive, stalactitic, and crystallized, of a blue or bluish green color.

SULPHATE OF ZINC—*White Vitriol*, occurs stalactitic and crystallized; color, greyish or greenish white.

SULPHATE OF COBALT occurs in crusts, and pulverulent, of a pale green color. With borax it gives a fine blue.

SULPHATE OF IRON. Crystallized in rhombs.

In octahedrons.

Stalactitic.

SULPHATE OF COPPER. Massive and stalactitic.

Crystallized in rhombs.

Variety, modified.

SULPHATE OF ZINC. Stalactitic.

Crystallized in four-sided prisms.

Acicular.

SULPHATE OF COBALT. Pulverulent.

In crusts.

In white balls.

DIAMOND.

THE diamond is often found with a polished surface, although more commonly rough. It generally occurs distinctly crystallized, also indeterminate and round. The primitive form is an octahedron. It has never been discovered in its matrix, though sometimes embedded in indurated alluvial soil. Its color is generally pale grey, but sometimes brown, green, yellow, rose-red, and blue. It has a metallic-like lustre; it may be split in four directions, is incomparably hard, and burns with a brilliant light in oxygen gas. The diamond forms the most beautiful and perfect series of crystallization. Those diamonds which are unfit to be cut, are employed by glaziers, or pounded for the use of engravers and lapidaries, and are called *Bort*.

DIAMOND. Rounded, spheroidal, *veiny*.

In octahedrons, (*primitive form*).

Variety, modified.

Crystallized in cubes. (*Extremely rare*).

Variety, modified.

In dodecahedrons, rhomboidal.

Variety, modified.

In tetrahedrons.

Variety, modified.

Variety, curvilinear.

In twin crystals.

Triangular, hemitrope, *veiny*.

Variety.

Earthy Minerals.



ZIRCON.

OF this genus there are two varieties; they contain a peculiar earth, called *Zirconia*, with silex; and when polished, somewhat resemble diamonds of inferior quality. They occur in the alluvial soil of Ceylon.

ZIRCON is found crystallized, also in rounded fragments, generally of a grey or brown color, also blue, yellow, and pink.

Hyacinth differs from the preceding only in color, which is a shining red-brown.

ZIRCON. In rectangular four-sided prisms, with pyramids.

Variety, modified.

In rounded fragments.

Embedded.

Hyacinth. Crystallized in four-sided prisms.

Variety, with pyramids.

Variety, modified.

In dodecahedrons.

Embedded.

In rounded and angular grains.

RUBY.

AUTOMALITE occurs embedded in chlorite-talc, in perfect octahedrons, of a dark green color, approaching black.

CEYLONITE—*Pleonaste*, occurs crystallized, and in rounded grains, of a dull bluish or red color, and frequently both red and blue in the same specimen. It is found with ruby, in the beds of rivers in Ceylon.

SPINELLE occurs crystallized; color, most beautiful bright red. It is sometimes embedded in granular and foliated carbonate of lime.

SAPPHIRE. The varieties of this species are the hardest of the earthly substances, and, next to the diamond, the most valuable. The finer stones are called *Oriental*, because they have always been found in India. The colors are blue, red, violet, green, and yellow; and sometimes the same specimen exhibits two or three colors, which are frequently distinct. *Sp. Gr.* 4. Alumine 98.

Blue Sapphire is fine dark blue or party-coloured, also pale blue or clouded; it occurs crystallized.

Oriental Ruby—Red Sapphire, is of a deep red color, also of different shades, generally with a bluish hue; some rubies exceed the diamond in value. It occurs crystallized and amorphous.

EMERY is nearly allied in chemical composition to the preceding and following varieties: it has a brown color, is very compact, and difficult to break; it occurs with mica, in the isle of Naxos. When reduced to powder, it is used by lapidaries for polishing and cutting facets in precious stones.

CORUNDUM is of various colors, with a high metallic

lustre, but generally grey or greenish white, and sometimes pink and blue; it occurs massive, detached, embedded, and crystallized. It is nearly allied, both in form and texture, to the oriental stones.

CHRYSOBERYL has a dull gold-yellow color of different shades, inclining to green, and sometimes reddish brown. It occurs amorphous and crystallized.

Cymophane is distinguished from the preceding by a chatoyant light on the surface. These two varieties are found with diamonds, in Brazil, and contain above 80 per cent. of alumine.

* * For a more particular account of the gems, see the author's Treatise on Diamonds and Precious Stones.

AUTOMALITE. Crystallized in octahedrons.

Variety, modified.

Variety, hemitrope.

CEYLONITE. Crystallized in octahedrons.

Variety, modified, partly coloured.

In rounded particles.

SPINELLE. Crystallized in octahedrons.

Variety, modified on the angles.

Variety, modified on the edges.

In dodecahedrons.

Variety modified.

Hemitrope, triangular.

In twin-crystals.

In tabular crystals.

Variety.

Indeterminately crystallized.

Blue Spinelle. In octahedrons, embedded.

Variety.

SAPPHIRE. In hexagonal prisms.

In double six-sided pyramids.

Variety, with double pyramids.

Variety, rounded.

Girasol, opalescent.

Asteria reflects a star of six rays*, *chatoyant*.

Variety, white or pale violet.

Variety, reddish white.

Oriental Ruby. In hexagonal prisms.

In double hexagonal pyramids.

Variety, indistinctly formed.

Variety, rounded.

Variety, exhibiting a chatoyant star of six rays.

Oriental Topaz. Color, pale yellow.

Oriental Amethyst. Color, violet-blue. (*Very rare*).

Oriental Emerald. Color, green. (*Extremely rare*).

EMERY. Massive and compact.

CORUNDUM. Massive, amorphous.

Embedded, sometimes in fibrolite.

Crystallized in six-sided prisms, detached.

Variety, modified.

In hexahedral pyramids.

In double hexahedral pyramids.

In rhombic dodecahedrons.

Variety.

Brown Corundum. Amorphous.

Crystallized in six-sided prisms.

In hexahedral pyramids.

*A [matchless specimen of this variety the Author sold to Count Bournon, for the private collection of his late Majesty Louis XVIII.

Blue Corundum. In six-sided prisms.

Variety.

Red Corundum.

Yellow Corundum.

CHRYSOBERYL. Amorphous, in rolled pieces.

Crystallized in four-sided prisms.

Variety, modified.

Cymophane. Amorphous.

Variety, crystallized.



SCHORL FAMILY.

THE substances which compose this family have generally a striated surface, and appear as if composed of an aggregation of long fibres. The specimens generally exhibit cross rents.

TOPAZ occurs in rolled masses, and in striated rhombic prisms, having sometimes one or both extremities terminated by pyramids; the cross fracture is always foliated. It is composed of alumine, silica, and fluoric acid. It is found in Brazil, Ceylon, New Holland, Saxony, Siberia, Cornwall, &c.

Brazil Topaz occurs crystallized, of a deep wine-yellow color, sometimes with a pink tinge, and pink; when heated, it becomes rose-red.

Saxon Topaz occurs in crystals, embedded or detached, of a pale yellow color.

PYCNITE—*Schorlite*, is found embedded in granite, of a greyish or reddish white color. It occurs crystallized.

PYROPHYSOLITE has a dull exterior, and a greenish white color; it occurs in irregular prisms, embedded in quartz. Its powder phosphoresces on hot coals.

EUCLASE is pale green, bluish green, and blue; it occurs crystallized, and is very rare.

EMERALD has a green color, which migrates into various shades of white; it occurs in rounded fragments, and crystallized, sometimes exhibiting transverse striæ.

Beryl is yellowish green, and sometimes greenish blue; it occurs in crystals, longitudinally striated.

A variety of a sea-green color is called *Aqua-marine*. Some beautiful specimens of this variety have been recently found in the Moran Mountains, in Ireland, accompanied with crystals of smoky quartz, and decomposed granite.

IOLITE is of a dull blue and yellowish brown color, as viewed in different directions. It occurs amorphous and crystallized. From the property of exhibiting two colors, it has been called *Dichroite*.

SCHORL generally occurs in black acicular crystals, which are sometimes aggregated, forming irregular three and six-sided prisms, longitudinally striated; it is also found fibrous, compact, and disseminated.

Precious Tourmaline is of various colors, green, blue, red, yellow, and blackish blue (*Indicolite*). It occurs crystallized.

Common Tourmaline differs from the preceding in color, which is a fine shining black.

Rubellite has a red color of various shades; it occurs crystallized and embedded, sometimes in the centre of the blue and green tourmalines.

EPIDOTE.—*Pistazite*. Its color varies from blackish green to pale green. It occurs crystallized or granular.

Zoisite has a smoky-grey color, with a pearly lustre; it occurs in crystals, though rarely disseminated. An earthy variety is said to have occurred, of a pale reddish white color, with a shining lustre, and friable.

AXINITE has generally a brown color, with a violet tinge. It occurs embedded, and crystallized in striated rhombic tables, which have a high lustre, and resemble the edge of an axe; whence its name.

BRAZIL TOPAZ. Crystallized in rhombic prisms.

Variety, with four-sided summits.

Variety, with pyramid, modified.

Variety.

Pink Topaz, generally with a brownish tint.

Blue Topaz, rounded, rarely crystallized.

White Topaz. Crystallized in rhombic prisms.

Variety, modified.

In rolled pieces.

Variety, exhibiting the fracture.

Saxon Topaz. Crystallized in rhombic prisms.

Variety with pyramid.

Variety, modified on the lateral edges.

Variety, modified on the terminal edges.

Variety, modified on the pyramids.

Variety, embedded.

Siberian Topaz. Embedded.

Variety.

PYCNITE. In short prismatic concretions.
Crystallized in long hexagonal prisms.

PYROPHYSOLITE.

EUCLASE. In rhombic prisms, with pyramids.
Variety, modified.
In fragments.

EMERALD. Crystallized in six-sided prisms.
Variety, truncated on the terminal edges.
Variety, modified.
Variety, pale green.
Variety, the green and white distinct.
Variety, rounded and polished.

Beryl. Crystallized in long six-sided prisms, striated.
Variety, with pyramids, or modified.
Aqua-marine. In six-sided prisms.
Variety, with pyramids.
Variety, modified.
Variety, in the matrix.

IOLITE. Amorphous.
Crystallized.
Embedded in feldspar.

SCHORL. Massive, fibrous.
Crystallized in three-sided prisms.
In six-sided prisms.
In acicular crystals.
Variety, aggregated.

Precious Tourmaline. In three-sided prisms.

Variety, the edges truncated or bevelled.

Variety, with pyramids.

In six-sided prisms.

Variety.

Common Tourmaline. In three-sided prisms.

Variety with pyramids.

In six-sided prisms.

Variety.

Rubellite. Crystallized in three-sided prisms.

In cylindrical prisms, embedded in quartz.

Variety, embedded in green tourmaline.

Variety, wine yellow (*very rare*).

Indicolite. Crystallized in three-sided prisms.

Variety, modified.

Variety, cylindrical.

EPIDOTE. Crystallized in oblique four-sided prisms.

In six-sided prisms.

Variety, with pyramids.

Variety, modified.

In aggregated acicular prisms.

Granular.

Zoisite. Massive.

Crystallized in oblique prisms.

Variety, indeterminate.

Variety, fibrous.

Variety, friable.

AXINITE. Crystallized in rhombic tables.

Variety, modified.

Disseminated.

GARNET FAMILY.

THE substances which compose the following division are generally crystallized. The crystals are commonly modifications of the four-sided prism, the rhomboidal dodecahedron, or the trapezohedron, and their planes are usually smooth. They contain silica, alumine, and lime, with a small portion of iron.

LEUCITE is of a greyish-white and red color; it occurs embedded, granular, and crystallized.

VESUVIAN—*Idocrase*, has a resin-brown color, with a shining lustre; it occurs crystallized, generally associated with mica, schorl, and garnets.

GEHLENITE. It has generally a grey or yellow color; it only occurs in crystals, which have a rough surface.

EGERAN has a deep brown color; it occurs crystallized.

GROSSULAR has a yellowish-green color; it occurs crystallized and massive.

GARNET, color, red, of various shades, occurs crystallized, and in fragments or grains. It easily melts before the blow-pipe.

Precious Garnet is of a blackish or bluish red color.

Topazolite is of a bright yellow color; it usually accompanies muscite. It occurs crystallized, and is nearly allied to the precious garnet.

Pyrope has a dark cherry-red color, and occurs in rounded and angular concretions.

Common Garnet is brown, of various shades; it occurs massive, and in large dodecahedrons. It becomes magnetic after being heated. The brown variety is melted in Bohemia as an ore of iron.

Colophonite resembles resin in color and lustre: it occurs crystallized and in aggregated concretions. It is not so heavy as the garnet.

Melanite is a black variety of the garnet; it is always crystallized in dodecahedrons.

Aplome is generally of a dull green color; it occurs in rhomboidal dodecahedrons, striated in the direction of their shorter diagonals.

GRENATITE—*Staurotide*. Color, dark brown; it occurs in prismatic crystals, which often intersect each other in the form of a cross, sometimes at right angles, and sometimes obliquely. It is frequently associated with *Kyanite*.

CINNAMON STONE has a brownish or yellowish red color, and resinous lustre; it is found in grains or fragments.

ALLOCHROITE. Color, yellowish grey and greenish. It occurs massive, has a resinous lustre, gives fire with steel, and melts before the blow-pipe.

LEUCITE. In twenty-four-sided crystals.

Variety, modified.

Embedded in lava.

Variety.

In granular concretions.

VESUVIAN. Crystallized in four-sided prisms.

Variety, lateral edges truncated.

Variety, terminated by four-sided pyramids.

Variety, pyramids truncated.

Variety, aggregated.

GEHLENITE. In rectangular flat prisms.

EGERAN. In rectangular prisms.

Variety, modified on the edges.

GROSSULAR. In smooth dodecahedrons.

Variety, trapezoidal.

Variety, modified.

PRECIOUS GARNET. In rhomboidal dodecahedrons.

Variety, truncated on all its edges.

Variety, in trapezoids.

Variety, modified.

Variety, embedded.

Topazolite. Crystallized in dodecahedrons.

Variety, modified.

Pyrope. In rounded or angular pieces.

Common Garnet. Massive.

In dodecahedrons.

Variety, modified.

Variety, embedded.

Colophonite. In aggregated concretions.

Crystallized in dodecahedrons.

Melanite. Crystallized in dodecahedrons.

Variety, edges truncated.

Aplome. In dodecahedrons.

Variety.

GRENATITE. In short oblique four-sided prisms.

Variety, the acute lateral edges truncated.

In long hexagonal prisms, intersecting.

CINNAMON STONE. Massive.

Variety, fragments.

ALLOCHROITE. Massive.

QUARTZ FAMILY.

AMETHYST, color, violet-blue, occurs massive and crystallized. The *common* variety has a dull violet hue, intermixed with red. The colors are seldom uniformly diffused, but appear in spots, or only in parts of the specimen. It occurs crystallized and disseminated.

Rock Crystal occurs massive and crystallized, and is perfectly transparent.

Common Quartz occurs massive and crystallized, sometimes exhibiting the primitive form, which is a rhomb. It is translucent: the colors are various, generally some shade of white or red, which will appear from the numerous sub-varieties.

Avanturine. Color, red-brown. It is interspersed by minute spots of yellow mica, which shine with a gold-like lustre.

Pseudomorphous Quartz is formed in cavities formerly occupied by crystals of other minerals, and presents the cube, octahedron, rhomb, &c. which have generally a dull surface, and are often hollow.

Float Quartz is cellular and spongy, and floats on water.

Flexible Sandstone is supposed to owe its flexibility to flattened grains of quartz, resembling mica. It is not elastic, but bends by its own weight.

Rhombic Quartz—*Fontainebleau Sandstone*, has a yellowish white color, and is found massive and crystallized in rhombs.

Prase, color, dark and dull green, occurs massive and crystallized on quartz.

Ferruginous Quartz is opaque, or translucent at the edges only; it owes its colors to the oxide of iron; it occurs in small aggregated crystals, which become magnetic by heat.

Cat's Eye (quartz with amianthus) is generally pale grey, and exhibits a chatoyant play of light.

Hornstone is of various colors, commonly grey or greenish; it occurs massive and in supposititious crystals. Its fracture is splintery or conchoidal; it has a shining lustre. A variety, called *Chert*, contains petrifications; it is used in the potteries.

Leelite, a variety of the preceding, is of a reddish color.

Woodstone has various colors, but most commonly some shade of grey or brown. It occurs massive, and has the appearance of wood. It is evidently formed by the infiltration of silicious particles, as the vegetable fibres decay.

Flinty Slate has a dull, smoke-grey color. It occurs massive, and is frequently traversed by veins of quartz. It is difficult to break.

Basanite, a finer variety, is used as a touch-stone to try the purity of gold.

Flint is of various colors, and is too generally known to require description. The interior of the spherical masses is often coated by crystals of quartz.

PRECIOUS AMETHYST. Massive.

Crystallized in six-sided pyramids.

Variety, in six-sided prisms.

Variety.

Common Amethyst. Disseminated.

Crystallized in six-sided pyramids.

Variety, in six-sided prisms.

Variety.

Rock Crystal. Massive.

Variety, iridescent.

Variety, containing chlorite.

Variety, with actynolite.

Variety, with rutile, &c.

Crystallized in six sided prisms, with pyram

Variety, in double pyramids.

Variety, modified.

Variety.

Common Quartz. Massive, and compact.

Variety, fibrous.

Variety, granular.

Imbedded, in rhombic crystals.

Crystallized in six-sided pyramids, aggregated.

In double six-sided pyramids.

In six-sided prisms, with six-sided summits.

Cap Quartz. A six-sided pyramidal nucleus, coated, on which a silicious deposit is subsequently formed.

Babel-tower Quartz. Prism upon prism, diminishing; (lately discovered at the Land's End).

Smoky Quartz. Massive.

Crystallized.

Black Quartz.

Yellow Quartz—Topazine Crystal.—*Cairn-gorm.*

Blue Quartz—Sappharine.

Red Quartz—Hyacinth of Compostella.

Rose Quartz, red or white-red.

Milky Quartz—Hyaline. Color, pale bluish pink.

Crystallized.

Paper Quartz. In foliated leaves.

Avanturine.

Pseudomorphous Quartz. In cubes.

In octahedrons.

In rhombs.

Variety.

Float Quartz. In cellular masses.

Variety.

Variety, foliated.

Flexible Sandstone. In laminæ. (Brazil).

Variety, from China.

Fontainebleau Sandstone. Massive.

Crystallized in rhombs.

Variety.

Prase. Massive.

Crystallized in six-sided prisms, with pyramids.

Variety, radiated, or fibrous.

Ferruginous Quartz. Massive.

In six-sided prisms, with pyramids.

Variety, reddish.

Variety, black or brown.

Cat's Eye. Compact.

Spheroidal, polished.

Hornstone. Massive.

In pseudomorphous cubes, rhombs.

Variety, foliated.

Variety, fine-grained, fracture conchoidal.

Leelite. Compact.

Chert. Marine remains petrified.

Variety, compact.

Variety.

Woodstone. Massive.

Variety, striped, wood-like.

Variety.

Flinty Slate. Massive.

Stratified.

Basanite—Lydian Stone. Massive.

Variety, with veins of quartz.

Flint.

CHALCEDONY is a silicious substance, of great diversity of color, but generally grey or bluish grey, which sometimes appears in spots or stripes. It rarely occurs in large masses, but is commonly found in veins, or filling cavities in other minerals; more generally stalactitic and mammillated. It also forms pseudomorphous crystals. It is translucent, and receives a fine polish; which characters pervade the following varieties.

Mocha Stone is a beautiful variety of the preceding, and contains dendritic appearances.

Carnelian. Its usual colors are blood-red, flesh-red, and white, which sometimes appear in spots or stripes; it occurs in masses, and stalactitic or mammillated.

Onyx has a deep clove-brown color, with concentric white or grey veins of chalcedony.

Sardonyx has the same color as the preceding, and is sometimes yellow, but is generally clouded, and without the white zones.

Plasma is of a green color, rather dull and dark, frequently with spots of white or yellow.

Chrysoprase has a pale green color, of various shades, and lighter than the preceding.

Heliotrope—*Blood Stone*, has a pleasing dark-green color, with red or yellow spots. It loses its colors before the blow-pipe.

Hyalite resembles chalcedony, except in its surface, which appears like gum-arabic. It occurs in crusts on decomposed basalt or porous wacke.

Silicious Tufa has a white color, and contains leaves and stems of plants mineralized. It is extremely friable and light: from the hot springs in Iceland.

Fiorite Pearl Sinter. Color, generally greyish white. It occurs stalactitic and botryoidal.

CHALCEDONY. Mammillated.

Stalactitic.

Botryoidal.

Veiny, striated.

Amorphous.

Crystallized in supposititious cubes.

Mocha Stone. Arborescent.

Variety, spotted.

Variety, red.

Variety.

Carnelian. Massive.

Variety, composed of red and white layers.

Mammillated.

Variety.

Onyx. Oriental.

Variety, from Germany.

Variety.

Sardonyx.

Plasma.

Chrysoprase.

Heliotrope. With red spots.

With yellow spots.

Hyalite. Encrusting basalt.

Variety, on porous wacke.

Silicious Tufa. With vegetable impressions.

Variety.

Fiorite Pearl Sinter. Massive.

Botryoidal.

Stalactitic.

OPAL. This beautiful species, when fine, is highly valued, and ranks with the first class of precious stones. Its varieties have a vitreous appearance.

Precious Opal is of a milk-white or pale blue color, diaphanous, and exhibits beautiful chatoyant and iridescent colors. It occurs in delicate veins and patches, also disseminated.

Girasol or *Fire Opal*, has a red, yellow, and greenish color, with a flame-like iridescence. It is found in Mexico.

Hydrophane is considered to be a variety of the opal, which, after the absorption of water, becomes diaphanous, and sometimes opalescent, but less splendid than the preceding.

Common Opal. Color, white, yellow, red, and bluish. It is brittle, very light, and has a vitreous lustre.

Cacholong. Color, milk or greyish white. It is opaque, and often associated with varieties of opal and chalcledony.

Semi Opal is of various colors, generally yellow, white, or brown. It is distinguished from common opal by being heavier and less brilliant.

Jasper Opal is of a red-brown or yellow color, sometimes spotted.

Wood Opal occurs of various colors, generally light yellow, and has a wood-like appearance, and conchoidal fracture. The finest specimen known was brought by Dr. Clarke from Hungary.

PRECIOUS OPAL. Disseminated in delicate veins.

In the matrix.

Detached.

Variety. *Harlequin Opal.*

Variety, in distinct patches. *Golden Opal.*

Girasol.

Hydrophane.

Variety, opalescent.

Common Opal. Variety, pale blue.

Variety, reddish brown.

Cacholong. Massive.

Variety, stratified with chalcledony.

With other varieties of opal.

Semi Opal. Amorphous.

Tuberose.

Variety.

Jasper Opal. In masses.

Variety, spotted.

Wood Opal. Massive.

Variety, wood penetrated by opal.

MENILITE is of various colors, generally brown and grey, and sometimes blue on the surface; it occurs in tuberosse masses, embedded in adhesive slate, from Mount Menil, near Paris; whence its name.

MENILITE. Tuberosse.

Embedded in adhesive slate.

TURQUOIS. This precious substance has a sky-blue color. It generally occurs botryoidal and in very delicate veins in limestone; it is opaque, and yields with difficulty to the file. It is very much used in jewellery, and has hitherto only been found in Persia.

TURQUOIS. Botryoidal.

Variety, in delicate veins, disseminated.

Variety.

JASPER is very universally distributed, and of infinite variety. Its colors are numerous, but most generally red, brown, or yellow. It never occurs crystallized. It is opaque; and contains portions of iron, to which its colors may be attributed.

Egyptian Jasper occurs of various colors, generally red or brown, with curvilinear delineations, concentric stripes, or black spots, often exhibiting curious *lusus naturæ*, dendritic appearances, &c.

Striped Jasper. This substance is of a brownish red color, with green bands passing through it.

Porcelain Jasper has a grey or dull blue color, sometimes black, with a vitreous lustre, and appears as if it had been acted upon by heat.

Common Jasper occurs of various colors, as red, yellow, brown, green, and black; it is compact, heavy, and breaks with much difficulty; fracture, conchoidal.

Pudding-stone is composed of rounded pebbles, gravel, &c. cemented together by oxide of iron and silice.

RED EGYPTIAN JASPER. In spheroidal masses.

Variety.

Brown Egyptian Jasper. With concentric zones.

Variety, exhibiting *lusus naturæ*, &c.

Striped Jasper. Massive.

Variety.

Porcelain Jasper. Massive.

Variety, vitrified.

Common Jasper. Massive and compact.

Variety.

Sinopal. Color, red.

Pudding-stone. Aggregated pebbles.

Variety.

AGATE. This beautiful substance is generally composed of chalcedony, quartz, carnelian, &c. Its colors are various, and finely contrasted, sometimes arranged in concentric zones and angular lines: from its great hardness it is capable of receiving a fine polish.

Striped Agate is composed of alternate layers of chalcedony, quartz, and amethyst, which are straight or curved.

Agate Breccia is composed of different fragments apparently cemented together by oxide of iron.

Agate Jasper is an assemblage of agate and jasper, of various colors, generally white, red, and yellow.

Fortification Agate is marked with angular lines, so disposed as to represent a fortification.

Landscape Agate. The colors are arranged so as to have a resemblance to a landscape.

Moss Agate is of various colors, generally yellow or red, with moss-like fibres.

Oriental Agate. Color, generally grey and clouded; it often contains dendritic appearances.

Blue Agate has a clouded bluish appearance.

Petrifaction Agate is composed of marine substances silicified.

PITCH STONE.

SOME of the substances which compose this family are supposed to be of volcanic origin.

OBSIDIAN. Its color approaches black; it occurs massive and compact, with a conchoidal fracture. It is translucent, and from its vitreous appearance has been called *Volcanic Glass*.

Marchanite, a variety of the preceding, occurs in rounded masses, of a smoke-grey color.

PITCH STONE generally resembles pitch in color and fracture, but is sometimes red, green, brown, and approaching black; the exterior is often decomposed: it is lighter, and melts more easily than the preceding.

Pitch Stone Porphyry is composed of feldspar and other substances, embedded in pitch stone, generally of a green color.

PEARL STONE is commonly of a smoke-grey color, with a shining lustre; it occurs massive, and has a spheroidal structure; it is brittle.

SPHÆROLITE occurs in botryoidal masses, of a brown or grey color, with faintly glimmering lustre; it appears to be nearly allied to the preceding.

PUMICE. Its color is light or dark grey; it occurs in cellular masses, sometimes intermixed with obsidian; it is so light as to float in water, and is used for polishing. It is found in great abundance in the volcanic islands.

Porphyritic Pumice is generally of a grey color, and contains feldspar and mica.

OBSIDIAN. Massive.

Variety, stratified.

Variety, iridescent.

Variety, enveloping glassy feldspar.

Variety.

Marekanite. In rounded pieces.

Variety.

PITCH STONE. Massive.

Variety, brown.

Variety, green.

Variety, red.

Variety.

Pitch Stone Porphyry. Massive.

Variety.

PEARL STONE. Massive.

Variety.

SPHERULITE. Botryoidal.

Variety.

PUMICE. In cellular masses.

Variety, fibrous.

Variety, glassy, containing obsidian.

Porphyritic Pumice.

ZEOLITE.

THIS family derives its name from the property of its species intumescing under the action of the blow-pipe.

PREHNITE. Its color is green, which migrates into white. It occurs massive and crystallized, with a foliated or fibrous structure; it easily melts before the blow-pipe.

ZEOLITE. Color, generally white, red, or green; it occurs massive and crystallized, and forms a beautiful series; its fracture is foliated, radiated, or fibrous.

Earthy Zeolite occurs massive, and in friable crusts, with other varieties of zeolite, and often filling cells in amygdaloid.

Fibrous Zeolite occurs massive, reniform, and in capillary crystals.

Needle Zeolite occurs massive and in delicate crystals, aggregated.

Radiated Zeolite is found massive, botryoidal, and crystallized; the fracture is beautifully radiated.

Foliated Zeolite—Stilbite, occurs massive, disseminated, and crystallized. It has a shining foliated fracture.

APOPHYLLITE. Its color is generally white or reddish white: it resembles calcareous spar; it occurs massive and crystallized. It exfoliates in acids, and in the flame of a candle; it easily melts before the blow-pipe into a white enamel.

CUBICITE. Color, greyish or reddish white; it occurs crystallized, seldom massive; it is sometimes translucent.

CHABASITE. Color, greyish white; it occurs crystallized; it melts into a porous white mass. A green variety of this substance is extremely rare.

HARMATOME. Its color is generally white; it occurs crystallized, the crystals sometimes maced, which character distinguishes the following variety.

Cross Stone. Its color is generally white, sometimes yellowish red, translucent; it occurs in tabular crystals, which intersect each other, forming a cross; whence its name.

LAUMONITE. The color of this rare mineral is snow or greyish white; it occurs massive and crystallized; it is liable to decompose, unless kept constantly in water.

DIPYRE. Its color is pearl-grey, with a shining lustre; it occurs massive, and in minute crystals, embedded, sometimes disintegrated; it phosphoresces and melts before the blow-pipe.

NATROLITE is of an ochre-yellow color; it occurs massive, beautifully zoned, and in capillary crystals; it contains a large portion of natron.

ALLOPHANE. Is found massive, disseminated, and botryoidal; the color is usually blue or white; it has a glistening lustre, and imperfectly conchoidal fracture.

PREHNITE. Massive.

Crystallized in short four-sided prisms.

Variety, truncated on the lateral edges.

In four-sided prisms, acicular.

Variety, aggregated.

Fibrous.

Embedded in wacke.

Variety.

ZEOLITE.

Earthy Zeolite. Massive.

In crusts.

In amygdaloid.

Fibrous Zeolite. Massive.

In reniform balls.

In delicate fibres.

Needle Zeolite. Massive.

In long four-sided prisms.

Variety, modified.

Radiated Zeolite. In four-sided prisms, with dihedral summits.

Foliated Zeolite. Massive.

Crystallized in four-sided tables.

In six-sided tables.

Variety, modified.

APOPHYLLITE. Massive.

Crystallized in four-sided prisms.

Variety, modified.

In four-sided tables.

Indeterminately crystallized.

CUBICITE. Crystallized in cubes.

Variety, modified on the edges.

Variety, modified on the angles.

Variety, trapezoidal.

CHABASITE. Crystallized in rhombs.

Variety, modified.

Green Chabasite.

HARMATOME. Crystallized in four-sided prisms.

Variety, with dihedral pyramids.

Cross Stone. In four-sided prisms with pyramids, macle.

Variety, red.

Variety, yellow.

Variety.

LAUMONITE. Crystallized in oblique prisms.

Variety, modified.

Variety, spicular.

DIPYRE. Embedded.

In minute crystals, disseminated.

Disintegrated (*very rare*).

NATROLITE. In veins.

Variety, embedded.

In delicate capillary crystals.

Diverging.

ALLOPHANE. Massive.

Variety, disseminated.

Variety, botryoidal.

WAVELLITE.

THIS family derives its name from Dr. Wavel, its discoverer. It consists of two species, which are composed of alumine, phosphoric acid, and water.

WAVELLITE. Its colors are yellowish and brownish grey. It occurs in spherical balls, which have a stellular fracture, sometimes iridescent.

BRAZILIANITE. Its color is darker than the preceding; it occurs massive, botryoidal, and crystallized; its fracture is feebly radiated; it is extremely rare, and was discovered by the Author, at Villa Rica, in Brazil.

WAVELLITE. In spherical balls.

Variety, stellated.

Crystallized in oblique four-sided prisms.

BRAZILIANITE. Massive, stellular.

Botryoidal.

Crystallized in flat rhombic prisms.



AZURE STONE FAMILY.

THE substances which compose this family are of a blue color; they generally gelatinize in acid.

LAPIS LAZULI is of a light or dark blue color; it occurs massive and disseminated, very rarely crystallized.

It is generally associated with pyrites; it melts before the blow-pipe.

AZURITE—*Lazulite*, is of a lighter blue color than the preceding; it occurs massive, and in crystals, embedded.

HAUYNE is of a dark or pale blue color; it occurs in granular concretions, crystallized, and disseminated in basalt and feldspar.

BLUE SPAR is of a pale blue color; it occurs massive and disseminated; it is hard, and has a splintery fracture. It is found associated with quartz, mica, and garnets. It has hitherto been found only in Asia.

LAPIS LAZULI. Massive.

With iron pyrites.

Disseminated in spots.

Crystallized in rhombic dodecahedrons. (*Rare*).

AZURITE. Embedded.

Crystallized in oblique prisms.

In four-sided pyramids.

HAUYNE. Disseminated in feldspar.

Crystallized in rhombic dodecahedrons.

In granular concretions.

BLUE SPAR. Massive.

Disseminated.

FELDSPAR FAMILY.

THIS is a very numerous and interesting family, and inclined to crystallize in forms derived from the four-sided prism. The colors are various; most of its species have a foliated and glistening fracture.

ANDALUSITE is of a reddish brown or pale grey color. It occurs massive and in crystals, which are sometimes embedded in mica slate; it is hard, and scratches glass.

SAUSSURITE is of a white, grey, or green color; it occurs massive and disseminated.

CHIASTOLITE—*Macle*, is of a yellowish white color; it occurs in crystals intersecting each other, forming a cross, sometimes hollow, but generally filled with clay-slate, in which they are embedded.

INDIANITE is of a grey color; it occurs massive, and in granular concretions, with corundum embedded.

ADULARIA has generally a white or dull white color; it occurs massive and crystallized. A rare variety of this substance from Ceylon is called *Moonstone*, which, when cut in a convex form, exhibits a chatoyant light.

Glassy Feldspar is of a greyish white color; it occurs in crystals, embedded, and appears cracked in various directions.

Labrador Feldspar is of a dull grey color, and exhibits most beautiful opalescent colors; it is massive and compact, and translucent on the edges.

Common Feldspar is of various colors, flesh-red, grey, or white, more rarely blue or green. It occurs massive and crystallized; it is one of the constituents of granite.

Disintegrated Feldspar is of a light grey color, and is a variety of the preceding, sometimes passing into clay.

Compact Feldspar is of a white or grey color; it occurs massive and crystallized.

Radiated Feldspar—*Albite*, has a white or greenish white color; it occurs massive and crystallized, with a fibrous or radiated structure.

SPODUMENE—*Triphane*, is of a pale green color; it occurs massive and disseminated; it is translucent. Before the blow-pipe it exfoliates in gold-like scales, then melts into a green enamel.

SCAPOLITE—*Paranthine* or *Wernerite*. Its general color is grey, or greenish grey, sometimes red; it occurs massive and crystallized; often associated with mica, chlorite, and magnetic iron; its structure is compact, foliated, and radiated.

BERGMANITE is of a greenish grey or flesh-red color; it occurs massive; its fracture is fibrous, curved, and stellular; it scratches feldspar.

ELAOLITE—*Fettstein*. Color, dull bluish green, passing into grey; it occurs massive. It melts into a white glass before the blow-pipe.

PETALITE has a greyish color; it occurs massive; it has a glistening lustre, and is translucent; the fracture is foliated. It contains the new alkali Lithia.

SODALITE is of a dark muddy green color; it occurs massive and crystallized; it contains 25 per cent. soda, and a small portion of muriatic acid.

MEIONITE is of a greyish white color, and translucent; it occurs massive and crystallized; it is very easily fused.

NEPHELINE is of a white color, sometimes tinged with

yellow or green; it occurs crystallized, and generally accompanied by mica: it becomes clouded in nitric acid.

ICE SPAR has a greyish white color resembling ice; it occurs massive, cellular, and crystallized; it is associated with mica, hornblende, and the preceding varieties. It is found at Monte Somma.

ANDALUSITE. Massive.

Disseminated.

Crystallized in four-sided prisms.

Embedded in mica-slate.

SAUSSURITE. Massive.

Disseminated.

In rolled pieces.

CHIASTOLITE. Crystallized in four-sided prisms.

Disseminated in clay slate.

Variety, hollow.

Variety, filled with clay-slate.

INDIANITE. Massive.

In granular concretions.

Variety, containing corundum.

ADULARIA. Massive.

In oblique four-sided prisms, with pyramids.

Variety, modified.

Moonstone.

Variety.

Glassy Feldspar. In four-sided prisms, embedded.

Labrador Feldspar. Massive.

Variety, blue or blue and green.

Variety, flame-coloured, margined.

Variety, color disseminated in small patches.

Variety.

Common Feldspar. Massive.

Crystallized in rhombic prisms.

Variety, embedded in granite.

In twin crystals.

Blue Feldspar. Massive.

Crystallized.

Green Feldspar. Massive.

Crystallized.

Avanturine Feldspar. Massive.

Variety, slaty.

Disintegrated Feldspar. Massive.

Variety, disseminated.

Variety, decomposing.

Compact Feldspar. Massive.

Crystallized in oblique four-sided prisms.

Albite. Massive.

Variety, radiated.

Crystallized in four-sided prisms with pyramids.

SPODUMENE. Massive.

Crystallized.

Disseminated.

SCAPOLITE. Massive.

Crystallized in oblique four sided prisms.

Variety, modified.

Foliated Scapolite. In oblique four-sided prisms.

Radiated Scapolite.

BERGMANITE. Massive.

Crystallized.

ELAOLITE. Massive.

PETALITE. Massive.

SODALITE. Massive.

Crystallized in rhomboidal dodecahedrons.

MEIONITE. Massive.

Crystallized in rectangular four-sided prisms.

Variety, aggregated.

Variety, disseminated.

NEPHELINE. Massive.

Crystallized in six-sided prisms.

Variety, aggregated.

Variety, disseminated.

ICE SPAR. Massive.

Crystallized in six-sided tables.

Cellular.

CLAY FAMILY.

THE varieties which compose this class of minerals have an earthy fracture, and emit an argillaceous smell when breathed upon; they are never found in any regular form: the colors are dull.

ALUMINITE has a white or yellowish white color: it occurs in reniform masses, and adheres feebly to the tongue.

Alum Stone is of various colors, generally greyish or reddish white; it occurs massive and porous; it is brittle, and is found in volcanic craters.

Porcelain Earth is generally of a white color; it occurs massive and compact; it is probably a deposit of decomposed feldspar, silica, &c.

COMMON CLAY—Loam, is of a yellowish grey color; it occurs massive, and sometimes indurated; it is an alluvial deposit; it adheres strongly to the tongue.

Potters' Clay. Its colors are greyish or yellowish white; it occurs massive, sometimes slaty, and semi-indurated; it is the common clay of which earthenware, pipes, &c. are made.

Variiegated Clay. Its color is some shade of red or brown; it occurs massive.

Slate Clay. Color, approaching black; it occurs massive, and has a slaty or earthy structure; it is soon decomposed. It generally contains vegetable impressions.

Adhesive Slate has a grey color; it occurs massive; it exfoliates by exposure to the atmosphere, but becomes

compact on immersion in water: it adheres strongly to the tongue; whence its name.

Polishing Slate is of a snow or yellowish white color; it occurs massive. It appears to be a fine deposit of silica with alumine.

Tripoli—Rotten Stone, has a dull brown color; it occurs earthy and friable; it is a decomposed limestone, or alluvial deposit.

FLOAT STONE is of a yellowish grey color; it occurs massive, and appears to be a transition from flint.

ALUM SLATE. Its color approaches black; it occurs massive, often covered by a white efflorescence of alum.

BITUMINOUS SLATE. Its color is black, or brownish black; it occurs massive; it is hard, and soon decomposes by exposure to the atmosphere.

DRAWING SLATE is of a black color; it occurs massive and compact; it is used for crayons.

WHET SLATE. Its color is generally grey, yellowish, or greenish; it occurs massive, and is of a fine texture.

CLAY SLATE. Its colors are various, from grey to black or red. It occurs massive, and is frequently traversed by delicate veins of tin. The common variety of this substance is used for slating houses.

MICA FAMILY.

THE principal characteristic of this family is its foliated and glistening appearance, sometimes approaching splendid; it has a tendency in all its crystallizations to the hexagonal form; it is difficultly fusible before the blow-pipe.

LEPIDOLITE. Its color is peach-red, which migrates into greenish yellow or white; it occurs massive and crystallized; it is composed of delicate shining scaly particles.

MICA is of various colors, but generally white, grey, brown, or black; it occurs massive, disseminated, and crystallized; it is easily divisible into *the thinnest* laminæ, which are perfectly flexible.

PINITE is of a blackish green color; it occurs massive and crystallized, and is generally found embedded in granite.

CHLORITE. Its color is some shade of green; it occurs massive, disseminated, and crystallized; it is unctuous to the touch; its texture is earthy, slaty, or foliated.

Common Chlorite is of a dark dull green color; it occurs massive.

Chlorite Slate has a blackish green color; it occurs massive and compact, with a slaty texture. It probably passes into the earthy variety.

Foliated Chlorite is of a dark or light green color; it occurs massive and crystallized.

LEPIDOLITE. Massive.

Crystallized in six-sided prisms.

Variety, of a greenish hue.

Variety, pale pink.

MICA. Massive.

Foliated.

Disseminated.

Crystallized in rhomboidal four-sided prisms.

In rectangular six-sided prisms.

In four or six-sided tables.

Variety,

PINITE. Massive.

Crystallized in six-sided prisms.

Variety, truncated on the edges or angles.

CHLORITE. Massive.*Chlorite Slate.* Massive.*Foliated Chlorite.* Crystallized in six-sided tables.

Aggregated.

Variety, massive.

Earthy.

Variety.

LITHOMARGE FAMILY.

THE substances which compose this family are rarely found crystallized; they are generally unctuous to the touch, which is a property common to all substances containing a considerable portion of magnesia; they are in general tough, though not hard; the colors are seldom bright.

GREEN EARTH is of a dull green color; it occurs massive, and in nodules, filling cells in amygdaloid; it contains a large portion of potass.

PIMELITE is of a dull green color; it occurs earthy, more or less indurated, and contains 15 per cent. of oxide of nickel.

LITHOMARGE is of a snow-white color; it occurs massive and disseminated; it is soft, also indurated; it adheres to the tongue, and falls to powder in water.

MOUNTAIN SOAP is of a dark brown color; it occurs massive, generally in cells, in trap rocks.

YELLOW EARTH has an ochre-yellow color; it occurs massive; is very soft, and adheres to the tongue.

CIMOLITE is of a greyish white color; it occurs massive, and is found in the island of Cimolia. It was highly prized in medicine by the ancients.

KOLLYRITE has a reddish or greyish white color; it occurs massive, is soft, and strongly adheres to the tongue.

GREEN EARTH. Massive.

Disseminated in amygdaloid.

Globular.

PIMELITE. Massive.

Disseminated.

LITHOMARGE. Massive.

Disseminated.

Indurated Lithomarge. Massive

MOUNTAIN SOAP. Massive.

YELLOW EARTH. Massive.

CIMOLITE. Massive. (Impressed with a seal).

COLLYRITE. Massive.

SOAP-STONE FAMILY.

NATIVE MAGNESIA is of a white or greyish white color; it occurs massive; fracture, foliated or radiated; it is soft, and adheres slightly to the tongue.

Valentianite is a variety of the preceding; it occurs compact, in large crystals, and in rounded pieces; it has a splintery or conchoidal fracture. It was brought by Lord Valentia from the Red Sea.

MAGNESITE—*Carbonate of Magnesia*. Its color approaches cream-yellow, often spotted; it occurs vesicular.

MEERSCHAUM has a greyish white color; it occurs massive; it is soft, very light, and adheres strongly to the tongue.

BOLE is of various colors, generally red; it occurs massive: when put in water, it falls to pieces with a hissing noise.

LEMNIAN EARTH has a yellow-grey or white color; it has a fine earthy fracture. The specimens hitherto received are from the Isle of Lemnos, and are impressed with a seal.

FULLERS' EARTH has a greenish grey color; it occurs massive, and has a dull earthy appearance; it falls into powder in water; it is friable.

STEATITE—*Soap-stone*, has a mottled soap-like appearance; it occurs massive, and in pseudomorphous crystals; it is generally soft, sometimes indurated, and has a greasy feel.

FIGURE STONE—*Agalmatolite*, has generally a grey color, sometimes mottled; it comes from China, carved

in grotesque figures: it feels greasy, and differs from steatite in not containing magnesia; it is also much harder.

NATIVE MAGNESIA. Massive.

Variety

Valentianite. Crystallized.

Variety.

MAGNESITE. Massive.

Tuberose.

Vesicular.

MEERSCHAUM. Massive.

Tuberose.

BOLE. Massive.

Disseminated.

LEMNIAN EARTH. Massive.

With impressions.

FULLERS' EARTH. Massive.

Stratified.

STEATITE. Massive.

Crystallized in four or six-sided prisms.

In rhomboidal dodecahedrons.

Variety, indurated.

FIGURE STONE. Massive.

Variety, coloured.

TALC FAMILY.

THIS family presents great diversity in color, texture, and general appearance; they all contain a large portion of magnesia.

NEPHRITE has a dull light-green color; it occurs in rounded masses, of a fine compact texture; it is translucent on the edges, is moderately hard, and receives a high polish.

Jade—Axe Stone, has a dark green color; it occurs massive, and is more translucent than the preceding. The specimens received in this country, come from the South Seas in the form of hatchets, idols, and various ornaments.

SERPENTINE has a dull color, generally green, brown, white, or red, often intermixed in the same specimen; it occurs massive; it frequently contains veins of asbestos, and some varieties have so large a portion of iron as to be magnetic.

Precious Serpentine. Its color is dark green, frequently spotted; it occurs massive; it is translucent, and easily yields to the knife.

POT-STONE is generally of a greenish-grey color, often spotted; it occurs massive; it is translucent on the edges, and soft; it is worked into culinary utensils, in which shape it always comes to this country.

TALC—*Venetian Talc*, has a greenish-white color, with a shining pearly lustre; it occurs massive, and in delicate foliated crystals, also radiated; it is peculiarly soft and agreeable to the touch; it is distinguished from mica by its inflexibility.

Compact Talc is of a dull white color; it occurs massive; it forms the base of rouge, and communicates a softness to the skin without any pernicious effect.

Columnar Talc has generally a greenish-grey color; it is composed of thin prismatic folia aggregated, forming six-sided prisms.

Nacrite—Earthy Talc, has generally a greenish color, with pearly lustre, it is composed of delicate scales, and is friable.

ASBESTOS. Its color is greenish-white or grey; it occurs massive, in delicate veins, and frequently disseminated in calcareous spar, quartz, &c.

Amianthus is a fine variety of the preceding, and has a silky lustre; it is composed of delicate flax-like fibres.

Amianthoide has an olive-green color; it occurs in filaments, accompanied by carbonate of lime, feldspar, quartz, &c. It is flexible and elastic.

Byssolite appears to be a variety of the preceding, and occurs in short and stiff filaments.

Rock Cork has a greyish or cream like color; it occurs in laminar masses, with a porous structure; some varieties very much resemble leather.

Rock Wood has generally a wood-brown color; it occurs massive and compact, and has a ligneous appearance.

NEPHRITE. Massive.

In rolled pieces.

Jade. Massive.

SERPENTINE. Massive, spotted.

Variety, containing veins of asbestos.

Variety.

Precious Serpentine. Massive.

POT-STONE. Massive.

VENETIAN TALC. Massive.

Disseminated and radiated.

Crystallized in six-sided tables.

Compact Talc. Massive.

Variety.

Columnar Talc. In common talc, folia aggregated.

Variety, disseminated.

Variety.

Nacrite. Compact.

Friable.

ASBESTOS. Massive.

Disseminated.

Variety.

Amianthus. In delicate flax-like fibres.

Variety, disseminated.

Variety, detached.

Amianthoide. In filaments.

Disseminated.

Byssolite. In short perpendicular filaments.

Disseminated.

Rock Cork. Massive.

Lamellar.

Rock Wood. Massive.

Ligneous.

Variety.

HORNBLLENDE FAMILY

CONTAINS many minerals which resemble some of the schorl family; a little practice will, however, easily distinguish them; some varieties are characterized by a strong pseudo-metallic lustre. They are soft to the knife, and when abraded have a dull greenish hue.

HORNBLLENDE. Its color is green or greenish-black; it occurs massive and crystallized.

Hornblende Slate is of a blackish-green color; it occurs massive, has a glistening lustre, and slaty fracture.

Basaltic Hornblende has a velvet-black appearance; it occurs crystallized, embedded in basalt.

ACTYNOLITE is of various colors, generally white or green; it occurs asbestos-like, massive, and crystallized.

Asbestos Actynolite has a snow-white color; it occurs massive, or aggregated in delicate spicula; it is always rough and coarse to the touch.

Common Actynolite has a green color of various shades; it occurs massive, disseminated, and crystallized.

Pargasite. Color, bottle-green; it occurs in concretions differing in size, also imperfectly crystallized in calcareous spar.

Glassy Actynolite is green of various shades, with considerable lustre; it occurs in crystals embedded, often separated by rents.

TREMOLITE is generally of a light color, never green, which distinguishes it from the two preceding species; it occurs massive and crystallized, with a pearly lustre.

Asbestos Tremolite is of a white color, variously tinged; it occurs massive, with a fibrous and stellular structure; it phosphoresces on hot coals.

Common Tremolite has a greyish-white color, it occurs massive and in crystals, embedded, longitudinally streaked.

Glassy Tremolite. Color, greyish or yellowish white; it occurs massive and crystallized, and has a shining lustre; it also frequently occurs granular.

Baikalite is of the same color as the preceding, of which it is a variety. It is found at the lake Baikal, whence its name.

SAPPARE—Kyanite. Its colors are yellow, white, and sky-blue of various shades; it occurs massive, disseminated, and crystallized, often embedded in mica-slate, and associated with granatite.

Rhatizite is of a white, green, or yellow color; it occurs in aggregated crystalline masses.

SCHILLER SPAR has a black-green color; it occurs in patches, with a splendent lustre, embedded in serpentine.

DIALLAG has a green color of various shades; it occurs massive, disseminated, and granular; it is translucent on the edges, and is very hard.

BRONZITE has a bronze-brown color, with shades of yellow; it occurs massive and disseminated, and has a strong metallic lustre.

ANTHOPHYLLITE is of an intermixed brown and grey color, with semi-metallic lustre; it occurs massive and in aggregated crystals, sometimes fibrous.

HYPERSTENE. Its color is brownish, with a copper-like lustre; it occurs massive; it breaks into rhombic fragments.

HORNBLLENDE. Massive.
 Crystallized in four-sided prisms.
 Variety, truncated.
 Embedded.

Hornblende Slate. Massive.

Basaltic Hornblende. Crystallized in six-sided prisms.
 Variety, modified.

ACTYNOLITE.

Asbestos Actynolite. Massive.
 In delicate spicula, fibrous.

Common Actynolite. Massive.
 Disseminated.
 Fibrous or granular.

Pargasite.

Glassy Actynolite. Massive.
 Disseminated.
 Crystallized in rhomboidal four-sided prisms.

TREMOLITE.

Asbestos Tremolite. Massive.
 Variety, fibrous or radiated.

Compact Tremolite. Massive.
 Crystallized in oblique four-sided prisms.
 Variety, modified.

Glassy Tremolite. Massive.

In acicular crystals.

Baikalite.

SAPPARE. Massive.

Disseminated.

Crystallized in oblique four-sided prisms.

Variety, modified, twin crystals.

Variety, embedded in mica slate.

Rhatizite. Massive.

Foliated, diverging.

SCHILLER SPAR. Massive.

Disseminated in patches.

DIALLAGI. Massive.

Disseminated.

Granular. (*Smaragdite*).

BRONZITE. Massive.

Disseminated.

ANTHOPHYLLITE. Massive.

Disseminated, fibrous.

In reed-like crystals.

HYPERSTENE. Massive.

In granular or lamellar concretions.

In curved laminae.

CHRYSOLITE FAMILY

IS composed of substances whose general color is green or black; the crystallizations are commonly derived from the four-sided prism; they have a foliated structure, and generally a vitreous lustre.

SAHLITE is green of various shades; it occurs massive and crystallized, sometimes disseminated in Tisee marble.

AUGITE has a blackish green or black color; it occurs massive, in rounded grains, and crystallized; it has an uneven or conchoidal fracture, and has different degrees of lustre.

Coccolite has a green color of various shades; it occurs granular, aggregated, crystallized, and disseminated; it has a glistening lustre, and is translucent on the edges.

DIOPSIDE is green of various shades; it occurs finely crystallized, generally accompanied by the following sub-species.

Mussite has a light green color; it occurs in aggregated crystals, and has a silky abestos-like appearance, with radiated fracture.

CHRYSOLITE has an oil-green color; it occurs generally embedded, filling cavities, rarely crystallized.

OLIVINE. Its color is olive-green, sometimes brown or black; it occurs massive, in granular concretions, also crystallized and embedded.

YENITE—Leivrite. Its color approaches black; it occurs massive, fascicular, and crystallized; it is very heavy; it has hitherto been found only in Elba.

SAHLITE. Massive.

Disseminated in Ticee marble.

Crystallized in rectangular four-sided prisms.

AUGITE. Massive.

In roundish grains.

Crystallized in four or eight-sided prisms.

Variety, modified.

Variety, embedded.

COCOLITE. Massive.

Disseminated, granular.

Crystallized in four or six-sided prisms.

DIOPSIDE. Massive.

Disseminated.

Crystallized in oblique four or eight-sided prisms.

Variety, modified.

Mussite. Massive.

In aggregated acicular fibres.

CHRYSLITE. In angular pieces.

Crystallized in four-sided prisms.

Variety, modified.

OLIVINE. Massive, granular.

In rectangular four-sided prisms.

Variety, embedded.

YENITE. Massive.

Radiated, fascicular.

Crystallized in four-sided prisms.

Variety, modified.

BASALT FAMILY.

BASALT has a dull blackish color; it occurs massive, in large columnar prisms.

WACKE is of a grey or brown color; it occurs massive and vesicular.

AMYGDALOID is a variety of wacke, having the cells filled with zeolite, green earth, calcareous spar, &c.

CLINK STONE is green of various shades; it occurs massive and compact; it has a slaty structure, and rings when struck with a hammer, whence its name.

BASALT. Massive.

Variety, compact.

Variety, granular.

Variety.

WACKE. Massive.

Vesicular.

AMYGDALOID. Massive.

Containing green earth, zeolite, or calcareous spar.

CLINK STONE. Massive.

DOLOMITE FAMILY.

THE substances which compose this family contain large portions of carbonate of lime and magnesia; they effervesce feebly in acids.

DOLOMITE has a snow-white color; it occurs granular and massive, often with realgar and pyrites disseminated.

Bitter Spar—*Rhomb Spar*, has a yellowish color; it occurs in rhombic crystals, embedded in chlorite-slate.

Magnesian Limestone has a light yellow or brown color; it occurs massive and botryoidal, and has a glistening fracture.

Flexible Limestone occurs massive; it is a variety of the preceding, and in thin slices is very flexible.

MIEMITE is of a green color; it occurs massive and crystallized, sometimes embedded in alabaster.

PEARL SPAR—*Brown Spar*, is of various colors, generally grey, white, pink, or brown, with a pearly lustre; it occurs massive, crystallized, stalactitic, and mammillated; it has a foliated or fibrous structure.

GURHOFITE has a snow-white or pink color; it occurs massive; it is dull, hard, and brittle.

DOLOMITE. Massive.

Crystallized in rhombs.

Variety, with pyrites disseminated.

Variety, with realgar.

Variety.

Bitter Spar. Massive.

Crystallized in rhombs, embedded.

Magnesian Limestone. Massive.

Variety, with vegetable impressions.

Botryoidal.

Flexible Limestone. Massive.

MIEMITE. Massive.

Crystallized in flat double three-sided pyramids.

PEARL SPAR. Massive.

Variety, rose-colored.

Fibrous or foliated.

Stalactitic or mammillated.

Crystallized in rhombs.

Variety, modified.

Variety.

GURHOFITE. Massive, snow-white.

Pink.

Blackish, in patches.

Variety.

LIMESTONE FAMILY.

THE various members of this family are universally diffused. Independently of their use in the arts, they contribute largely to the fertility of the soil; and their chemical agencies are essentially beneficial to animal and vegetable life.

TABULAR SPAR has a greyish white color; it occurs massive and crystallized, often disseminated in Cinnamon-stone; it is rather hard and brittle; in nitric acid it effervesces for a moment, and then granulates.

SLATE SPAR—*Schiefer Spar*. Its color is white of various shades; it occurs massive and disseminated, also in distinct concretions, and lamellar; it has a pearly lustre, and slaty fracture.

APHRITE. Its color is white; it occurs massive and disseminated, of a slaty or sparry structure; it has an earthy texture, and effervesces violently in acid.

AGARIC MINERAL—*Rock-Milk*, is of a yellowish or white color; it occurs in crusts and tuberos pieces; it is a pure carbonate of lime, and dissolves in acid.

CHALK. Its color is snow-white; it occurs massive and disseminated.

COMMON LIMESTONE. Its color is grey; it occurs massive and compact; it burns to lime, and effervesces in acid.

Oolite—*Roe-stone*, is of a yellowish brown color; it occurs massive, composed of minute globular concretions, resembling the roe of a fish.

Granular Limestone—Marble, is of a white color, and occurs massive; it is much used in the arts, for statuary, &c.

Tiree Marble is of a flesh-red color; sahlite and titanium are often embedded in it.

Mona Marble has a white and green color, and much resembles verde antique.

Shell Limestone is of various colors, generally dark; it is composed of fossil shells.

Lumachella—Fire Marble, is a variety of the preceding; it is composed of shells, which have a brilliant opalescent lustre.

Coralloid Limestone is generally of a dark color; it is found massive, and composed of corals, zoophites, &c. some varieties much resemble madreporæ.

Calcareous Spar. The crystallizations of this substance far exceed those of any other in number, beauty, and complexity. Count Bournon has described nearly 1000 varieties: they are arranged in three divisions, arising 1st. from the acute six-sided pyramid (primitive rhomb); 2d. the six-sided prism; and 3d. the three-sided pyramid; and so beautiful is the connexion, that the termination of the third series gradually approaches the first, forming, as it were, a complete circle. The color is generally yellowish white; the transparent variety exhibits double refraction.

Fibrous Limestone—Satin Spar, is snow-white; it occurs massive and compact, composed of short aggregated fibres; it is often associated with pyrites.

Stalactite has a white, yellow, green, or brown color; it occurs massive, botryoidal, mammillary, reniform, &c. it is formed by precipitation, and hangs like icicles from the roofs of caverns.

Calcareous Tuffa is grey of various shades; it occurs massive, cellular, and ramose, and is generally found precipitated upon substances immersed in water strongly impregnated with carbonate of lime.

Pea Stone is generally brownish-white; it is composed of rounded pea-like concretions.

LUCULLITE—*Black Marble*. Its color is intense black; it is capable of receiving a very high polish, and is in much estimation for vases, chimney-pieces, &c.

Swine Stone is of a bluish-grey color, and clouded; it occurs in granular masses, and when rubbed, emits a very disagreeable odor.

MARL is of various colors, generally greyish-white; it occurs massive, and frequently contains impressions of fish and dendritic appearances.

ARRAGONITE. Its colors are white, yellow, or grey; it occurs massive, crystallized, arborescent, and stalactitic, with a fibrous or radiated structure, and pearly lustre; it is sometimes embedded in granular gypsum.

TABULAR SPAR. Massive.

Crystallized in rectangular four-sided tables.

SLATE SPAR. Massive.

Disseminated.

Crystallized.

APHRITE. Massive.

Disseminated.

Tuberosc, friable.

AGARIC MINERAL. Pulverulent.

CHALK. Massive.

Variety, with organic remains.

COMMON LIMESTONE. Massive.

Oolite. Massive, in granular concretions.

Granular Limestone. Massive.

Tiree Marble. Massive.

Containing sahlite or titanium.

Mona Marble. Massive.

Shell Limestone. Massive.

Variety.

Lumachella. Massive, iridescent.

Coralloid Limestone. Massive.

Variety.

Calcareous Spar.

SIX-SIDED PRISM.

- 1 Terminated by planes perpendicular to the axis.
- 2 No. 1. The alternate terminal edges truncated.
- 3 No. 2. Truncations meeting and forming three-sided acuminations.
- 4 No. 3. Prism short, forming pentagonal dodecahedron.
- 5 No. 1. All the terminal edges truncated.
- 6 No. 5. Truncations meeting and forming six-sided acuminations.
- 7 No. 1. With terminating planes set on the lateral edges.
- 8 No. 7. Terminal planes meeting and forming pyramid.

- 9 Prism short and flattened, acumination on alternate lateral edges, forming rhombic dodecahedron (*rare*).
- 10 No. 1. The alternate lateral planes broader.
- 11 No. 10. The alternate lateral planes meeting and forming three-sided prism.
- 12 No. 1. Prism shortened, forming six-sided table.
- 13 No. 3. Prism wanting, the acuminations meet and form a very obtuse rhomb.
- 14 No. 1. Alternate lateral planes inclined towards the axis, forming a prism six-sided at one end, and three-sided at the other.

RHOMB.

- 15 Primitive rhomb (angles 105. 5, and 74. 55.)
- 16 Rhomb, with modifications on the edges, very nearly approaching the cube (angles 92. 18, and 87. 42.)
- 17 Rhomb having on each face a flat four-sided pyramid set on the edges of the rhomb.
- 18 Rhomb having on each face a flat four-sided pyramid set on the angles of the rhomb.

SIX-SIDED PYRAMID.

- 19 Perfect.
- 20 With alternate faces broader.
- 21 Double, the alternate edges longer, junction of the bases forming a zigzag line.

The above are the principal *simple modifications*, which are liable to almost innumerable varieties, arising from supernumerary planes, and in the rhombic and pyramidal sections from the various angles under which the planes meet. From so extensive a field it is difficult to select,

but the following few may be pointed out as interesting, and likely to fall under the notice of the young collector.

- 22 No. 3. With two additional planes set obliquely on the ends of the alternate lateral planes.
- 23 No. 3. With additional planes on the solid angles.
- 24 No. 3. With the lateral and terminal edges of the prism truncated.
- 25 No. 8. The summit of the pyramid again acuminate by six planes.
- 26 No. 25. The alternate edges of the original pyramid truncated.
- 27 No. 10. The broader planes truncated on the terminal edges.
- 28 No. 12. The tables truncated on the lateral edges.
- 29 No. 12. The tables truncated on the terminal edges.
- 30 Rhomb varying from very obtuse to very acute.
- 31 No. 21. The summits of the pyramids acuminate by three or six planes.
- 32 No. 21. With truncations on the angles, formed by the junction of the bases, (when these truncating planes meet, No. 8 is formed).

When the rhombic forms have their edges rounded, the crystal becomes lenticular, and should the student wish to obtain the primitive rhomb for examination or comparison, it will arise from the mechanical fracture of any of the crystallized varieties, the fragments always having angles whose measurement coincides with that of the primitive.

Fibrous Limestone. Massive.

Variety, with veins of pyrites.

Stalactite. Massive, zoned, agate-like.

Botryoidal.

Reniform or mammillated.

Variety, tubiform.

Stalactitic.

Variety, green.

Calcareous Tuffa. Massive.

Coating, cellular.

Variety.

Variety.

Pea Stone. Massive.

In pea-like concretions.

Variety.

LUCULLITE—Black Marble. Massive.

Variety, with sparry veins.

Swine Stone. Massive, granular.

Variety.

MARL. Compact.

Bituminous, containing petrifications.

Variety, with impressions.

Variety.

ARRAGONITE. Massive.

Fibrous or radiated.

Variety, arborescent or stalactitic.

Crystallized in six-sided prisms.

Variety, modified.

In very acute double six-sided pyramids.

Variety.

APATITE FAMILY.

OF this family there are only two species; they are composed of lime and phosphoric acid; and commonly phosphoresce when placed on hot coals.

APATITE is of a grey, green, or bluish color; it occurs crystallized, sometimes massive, and disseminated, generally embedded.

Moroxite—*Asparagus Stone*, has a green or blue color; it occurs disseminated and crystallized; it is translucent, some of its varieties do not phosphoresce.

PHOSPHORITE is white, or yellow, and reddish white; it occurs massive, earthy, stalactitic, and crystallized; it has sometimes a radiated or fibrous fracture.

APATITE. Massive.

Disseminated.

In crystals, aggregated.

Variety.

Crystallized in six-sided prisms.

Variety, truncated.

Variouly colored.

Asparagus Stone. Massive,

Disseminated.

Crystallized in six-sided prisms.

Variety, with pyramids.

Variety.

PHOSPHORITE. Massive.

Stalactitic.

Reniform.

Earthy.

FLUOR FAMILY.

THE substances of this family are composed of fluoric acid and lime; when powdered and digested with sulphuric acid, the fluoric acid escapes in the form of gas. By means of this acid the beautiful operation of etching upon glass is performed. It decrepitates on the application of heat, and becomes phosphorescent when thrown on hot coals.

COMPACT FLUOR has a bluish-grey, or greenish-white color; it is found massive, and gives a white streak with the knife; it is of rare occurrence.

CHLOROPHANE. This beautiful and rare species is of a brown or pale violet color; it occurs massive; fracture, foliated. It is beautifully phosphorescent.

FOLIATED FLUOR. Its colors are various, as white, blue, yellow, green, and pink, frequently two or more of them appear in the same specimen, often zoned; it occurs massive and crystallized in a variety of beautiful forms, the primitive of which is the octahedron; it admits of perfect cleavage. By exposure to heat, the color of the blue variety becomes a beautiful purple, and vanishes if the heat is continued. A variety of this species is manufactured into beautiful vases, &c.

EARTHY FLUOR. Its color is light purple or deep blue; it occurs massive, and coating the other varieties; it is very friable, and sometimes striated.

Argillaceous Fluor is of a brown color; it occurs in small detached cubes, and is generally found in decomposed amygdaloid.

COMPACT FLUOR. Massive.

CHLOROPHANE. Massive.

Foliated.

FOLIATED FLUOR. Massive.

Variety.

- 1 Cube perfect.
- 2 Cube truncated on the edges.
- 3 No. 2. Truncations effacing the original planes of the cube, and forming a rhombic dodecahedron.
- 4 Rhombic dodecahedron, having each face bent in the direction of its shorter diagonal, crystal assuming a more globular form, and having twenty-four triangular faces.
- 5 The edges of the last form indistinct; the crystal becomes spheroidal.
- 6 Cube truncated on the solid angles.
- 7 No. 6. Truncations effacing the original planes of the cube, and forming an octahedron. (This is the primitive form).
- 8 Octahedron truncated on the edges.
- 9 Cube having each solid angle replaced by three planes.
Variety, angles replaced by six planes.
- 10 Cube bevelled on the edges.
- 11 No. 10, the bevelments effacing the original planes of the cube, and forming a cube having a four-sided pyramid on each face.

* * * It will easily be perceived that a strong connection exists between Nos. 4 and 11, each consisting of twenty-four triangular faces, which, taken in pairs, produce rhombs; but from the different angles under which the faces meet, their forms are dissimilar; and the general

aspect of the crystals would appear to a casual observer as not possessing the slightest resemblance.

Idem. In cubes with indented faces.

Variety, with pyrites in the interior.

In cubes disseminated.

Variety, margined with barytes.

Variety.

In minute cubes aggregated.

Topazine. Yellow.

Sappharine. Blue.

False Emerald. Green.

Crystallized in cubes.

Variety, with low four-sided pyramid on each face.

Variety, with planes on the edges.

Variety.

Variety, containing water.

False Ruby. Pink.

Variety.

EARTHY FLUOR. Massive.

Variety, granular.

Variety, disseminated.

Variety, striated.

Argillaceous Fluor. In detached cubes.



GYPSUM FAMILY.

GYPSUM, when calcined, forms Plaster of Paris; the translucent varieties become opaque in the flame of a candle.

EARTHY GYPSUM has a dull yellowish brown appearance, sometimes with red and white veins; it is massive, and has a granular structure; it is loosely coherent.

Compact Gypsum is of a snow-white color; it occurs massive, and is used in sculpture.

Fibrous Gypsum has a snow-white color; it occurs massive and in silky fibres; it is often translucent.

Foliated Gypsum. Its color is generally white, grey, or reddish; it occurs massive and crystallized.

Plumose Gypsum is of a snow-white color; it occurs in capillary curls, or mammillated.

Selenite is generally transparent; its colors are dull white; it occurs massive, disseminated, and crystallized.

ANHYDRITE is of a bluish white color, sometimes deep red; it occurs massive, and has a fibrous or radiated structure.

Cube Spar—Sparry Anhydrite, is pearl white, of different shades; it occurs massive and crystallized; it is frequently embedded in a scaly or granular variety.

VULPINITE is of a bluish-white color; it occurs massive, and has a granular and foliated fracture; it is harder than the preceding, and is composed of sulphate of lime and silica.

GLAUBERITE has a greyish-white color; it occurs in minute crystals embedded in rock-salt; it is soluble in water.

EARTHY GYPSUM. Massive.

Variety, granular.

Compact Gypsum. Massive.

Variety.

Fibrous Gypsum. In fibrous masses.
Variety.

Foliated Gypsum. Massive.
Crystallized in six-sided prisms.

Plumose Gypsum. In delicate spicula, or curls.
Variety, mammillated.
Variety.

Selenite. Massive.
Crystallized in six-sided prisms.
In twin crystals.
Variety, acicular.
Variety.

ANHYDRITE. Massive.
Fibrous or radiated.

Cube Spar. Massive.
Crystallized in rectangular four-sided prisms.
In six-sided prisms.
In eight-sided prisms.

VULPINITE. Massive.
Disseminated.

GLAUBERITE. Disseminated.
Crystallized in four-sided prisms.

BORACITE FAMILY.

THERE are only two species in this family, both of which are of rare occurrence; they contain silica from 80 to 90, with boracic acid, lime, and water.

DATHOLITE. Its color is pale greenish white; it occurs massive, disseminated, and crystallized; also botryoidal and in delicate concretions, when it is called *Botryolite*.

BORACITE has an opaque grey color, and is always crystallized, the crystals occur embedded in gypsum.

DATHOLITE. Massive.

Disseminated.

Crystallized in oblique four-sided prisms.

Botryolite. Massive.

Botryoidal.

Radiated or fibrous.

BORACITE. Crystallized in cubes.

Variety, truncated on the edges.

Variety, truncated on the angles.

Variety, truncated on the edges and angles.

Variety.

BARYTE FAMILY.

THE two species of this family—Carbonate and Sulphate of Barytes, may easily be distinguished from other earthy substances by their great weight, whence their name.

CARBONATE OF BARYTES. Color, yellowish white or grey; it occurs massive, disseminated, and crystallized; its longitudinal fracture is striated or radiated, and has a glistening lustre. It is soluble in diluted muriatic acid, and melts before the blow-pipe.

SULPHATE OF BARYTES—*Heavy Spar.* Its colors are yellow, white, brown, grey, and blue, of various shades; it occurs massive, disseminated, and crystallized; also reniform, hepatic, dendritic, and columnar; its structure is compact, foliated, or earthy; it is transparent, translucent, or opaque; it often contains delicate veins and patches of galena.

CARBONATE OF BARYTES. Massive.

Crystallized in six-sided prisms.

In double six-sided pyramids.

Variety.

SULPHATE OF BARYTES. Massive, compact.

Variety, with galena.

Variety, earthy or granular.

Variety, lamellar.

Variety, arborescent.

Idem. Radiated.

Variety, reniform.

Variety, hepatic.

Crystallized in rectangular four-sided tables.

Variety, with bevelled edges.

Variety, edges and angles truncated.

In oblique four-sided tables.

In four or six-sided prisms.

Variety, bevelled.

Variety, transparent.

Variety, acicular.

Variety, columnar.

STRONTIAN FAMILY.

CARBONATE OF STRONTIAN—*Strontianite*. Its color is green of various shades, or brown; it occurs massive, disseminated, and crystallized, and has a radiated or fibrous fracture. It is generally associated with sulphate of barytes. Its powder gives a beautiful red color to the flame of spirits of wine; it phosphoresces at a high temperature before the blow-pipe. It is extremely rare, and has hitherto been only met with at Strontian.

SULPHATE OF STRONTIAN—*Celestine*, is of a white, sky-blue, or bluish grey color, sometimes reddish; it also occurs massive and crystallized, with a foliated and fibrous structure.

STRONTIANITE. Massive, green.

Variety, disseminated with sulphate of barytes.

Crystallized in acicular six-sided prisms.

Variety.

Variety, brown.

SULPHATE OF STRONTIAN. Massive.

Variety, compact.

In irregular four, six, or eight-sided tables.

In rectangular four-sided prisms.

Variety with pyramids.

Radiated.

Fibrous.

Variety.

Granular.



CRYOLITE.

THE color of this rare substance is generally snow-white, but it is sometimes brown from the presence of iron; it occurs massive and disseminated; its fracture is foliated, with a glistening lustre; it is translucent. It contains a large quantity of soda, which causes its easy fusibility. It derives its name from melting like ice. It has hitherto been found only in Greenland.

CRYOLITE. Massive.

Disseminated.

Ferruginous.

Inflammables.

NATIVE SULPHUR has a bright yellow color; it occurs massive, disseminated, and crystallized in octahedrons; the finest varieties are from Coneil, in Spain.

Volcanic Sulphur. Color, yellow of various shades; it occurs stalactitic, spongy, and granular; also in aggregated crystals.

BITUMINOUS FAMILY.

NAPHTHA. It is composed of carbon, hydrogen, and oxygen; it takes fire at the approach of flame.

PETROLIUM is reddish or blackish brown; it is thick, floats on the water, and may be seen oozing from various strata. It is sometimes precipitated on limestone.

ELASTIC BITUMEN. Color, blackish, greenish, and yellowish brown; it occurs massive, filling holes in limestone; it is peculiar to Castleton, Derbyshire.

Indurated Bitumen. Color, brown; it occurs massive, and botryoidal; fracture, perfectly conchoidal, with a shining lustre; it is brittle. It is found in Derbyshire.

COAL FAMILY.

BITUMINOUS WOOD—*Bovey Coal*—*Brown Coal*, has a dark brown color, and a ligneous appearance; it burns with a disagreeable smell.

Earthy Brown Coal. Its colors are yellowish or blackish brown; it occurs with the above.

Alum Earth—*Bituminous Shale*, is blackish brown; it occurs massive, and is scarcely inflammable.

Cannel Coal occurs massive and compact, of a black color; fracture, conchoidal. It is sometimes formed into vases, &c.

BLACK COAL—*Jet*, occurs massive, sometimes with impressions.

Newcastle Coal—*Slate Coal*. It occurs massive and foliated, often with pyrites disseminated.

Shining Coal is of a black color, often beautifully iridescent, and is sometimes called *Peacock Coal*; it occurs massive; it is fragile, and has a foliated fracture.

Slaty Coal. Color, black, and sometimes shining grey; it often contains layers of charcoal, in regular strata. It is found in Derbyshire and other coal counties.

Foliated Coal occurs massive, and is very soft and light; it contains fine folia of pyrites, which are sometimes silver white.

GLANCE COAL has a peculiar iron-black and tempered steel-like appearance; it occurs massive and vesicular; it has a conchoidal fracture. It is also found slaty and columnar.

GRAPHITE—PLUMBAGO.

SCALY PLUMBAGO. Its color is dark steel-grey, with metallic lustre; it occurs massive and disseminated, rarely crystallized; it is very soft.

Compact Plumbago differs from the preceding only in structure. It is used for pencils, crayons, &c. It is better known by the name of Black Lead.

MINERAL CHARCOAL. Color, black; it occurs in thin layers, disseminated in peculiar varieties of coal.

RESIN FAMILY.

AMBER. Color, yellow, or yellowish white, and reddish; it occurs in rounded pieces, with a rough exterior, sometimes decomposed. It is found on the Norfolk coast. The variety from Mozambique often envelopes insects.

HONEY STONE has a yellow color; it occurs embedded in grains, or crystallized in flat octahedrons, in brown or wood coal, and is very rare.

RETIN ASPHALT. Colors, yellowish and reddish-brown; it occurs massive; it burns with a fragrant odor. It contains resin 55, asphalt 42.

FOSSIL COPAL. Its color is yellow or brown; it occurs in rounded pieces, and appears to be a variety of retin asphalt, but approaches more nearly to gum; it has sometimes a resinous lustre.

Earthy Salts.

SALTS, with an earthy base, dissolve in water, and frequently diliquesce by the humidity of the atmosphere.

NATIVE ALUM has a white color, and occurs stalactitic, efflorescent, and in capillary crystals. It contains alumine 18, oxide of iron from 5 to 10, with sulphuric acid and water.

EPSOM SALT is white of various shades; it occurs massive and tuberosc; it contains sulphuric acid 33, magnesia 19, and water.

Alkaline Salts.

SALTS OF SODA.

THESE salts are composed of soda, variously combined.

CARBONATE OF SODA—*Natron*. Its color is yellowish white; it occurs in efflorescences, and acicular, or radiated.

SULPHATE OF SODA—*Glauber Salt*, is white of various shades; it occurs stalactitic, reniform, botryoidal, and crystallized.

MURIATE OF SODA—*Rock Salt*. Its colors are white and grey; it occurs massive, disseminated, reniform, and stalactitic; structure, foliated or fibrous.

BORAX. Its colors are yellowish or greenish white; it occurs crystallized, and has the property of double refraction. It contains boracic acid 40, soda 10, and water.

SASSOLIN. Its colors are greyish or yellowish white; it occurs in grains, crusts, or small corroded pieces in extinct volcanoes; it is very light.

SALT OF POTASH.

NITRE is white of various shades; it occurs in flakes or capillary crystals. It contains nitrate of potash 45, and lime.

SALTS OF AMMONIA.

THESE salts contain muriate of ammonia 98.

VOLCANIC SAL AMMONIAC. Its color is white, grey, green, or yellow; it occurs stalactitic, efflorescent, and sometimes crystallized; it may be known by its pungent taste. It is found in volcanic countries.

SULPHATE OF AMMONIA. Color, yellowish grey, or yellow; it occurs in mealy crusts or stalactitic.

APPENDIX.

Many of the substances described in this Appendix are very imperfectly known; the greater part of them have not been analyzed, others are separated from species in which they have been formerly included, and probably many will ultimately prove to be varieties of substances already described.

ABRAZITE occurs in globular masses and crystallized in octahedrons; its color is greyish white, sometimes blue or red. It has been called *Zeagonite*, and *Gismondine*—from its having been first noticed by Gismondi. It is composed of silex, lime, alumine, magnesia, and oxide of iron. Its locality is Cape di Bove.

ALLAGITE. Said to occur massive, with a conchoidal fracture, of a brown and green color. The green hue soon changes to dark grey or black, and the brown into pink. It is supposed to be an oxide of manganese.

AMBLYGONITE occurs massive and crystallized in oblique four-sided prisms; color, various shades of light green, spotted externally. It has a vitreous lustre, an uneven fracture, and appears to be a variety of spodumene.

ARFVEDSONITE. A variety of hornblende. Color, black; it is irregularly crystallized. From Greenland.

BLOEDIT. A soft substance of fibrous texture, of a red color; it is found transparent, but becomes opaque by exposure. It is named after Bloede, a mineralogist at Dresden.

BREWSTERITE. A substance formerly described as apophyllite, but differing from it in the measurement of its angles; it is found in Scotland, and named after Dr. Brewster.

- BUCHOLZITE.** Said to occur amorphous, spotted black and white, with glistening lustre, and of lamellar structure, resembling feldspar. It is from the Tyrol; and named after Bucholz, the chemist. It is composed of silix, alumine, potash, and oxide of iron.
- CALAMITE.** A substance resembling tremolite, which occurs in translucent, rhombic prisms, striated; and of a light green color. It is from Sweden.
- CARINTHIN** occurs in coarse granular concretions of a greenish black color; it is translucent, and has a conchoidal fracture. It is the Amphibole laminaire of Haüy, and may be arranged at the head of the hornblende family.
- CEREOLITE.**—Of this substance little is known. *De Dree*, (*Musée Mineralogique*, page 17), says, it derives its name from its resemblance to wax, of which it often has the softness, and that the variety found in lava has been erroneously considered as steatite.
- CHLOROPAL.** Said to occur earthy, with conchoidal fracture, of a green color.
- CHLOROPHÆITE.** A substance found embedded in amygdaloid; the fresh fracture has a green color of various shades, but turns darker after being exposed a short time, and ultimately becomes black; it is very soft.
- COMPTONITE** occurs in small four or eight-sided prisms, colorless, semi-transparent, harder than apatite. It is very nearly allied to the mesotypes, and was found about five years since in the vicinity of Vesuvius.
- CONITE** occurs massive, stalactitic, and encrusting; its color is various shades of grey. It becomes brown by exposure to the atmosphere; is brittle, and has an uneven fracture. It appears to be a calcareous carbonate of Magnesia.
- CONDRODITE** occurs in small grains of an orange color,

with resinous lustre, disseminated in an imperfectly crystallized carbonate of lime. It has been recently imported from Pargas, in Finland; and, excepting in color, has a considerable resemblance to Pargasite, from the same place. This substance has also been called Brucite and Maclureite.

COUZERANITE. Said to occur in rectangular prisms of a black or blue color.

CRONSTEDITE is said to be found massive and crystallized, and composed of dark colored fibres, with shining lustre; in thin laminæ it is slightly elastic. It is found in Bohemia.

DESMINE is but little known; it occurs in small silky tufts in lava, from the vicinity of the Rhine, and is sometimes accompanied by Haüyne.

DIASPORE has been found in masses composed of curvilinear laminæ, and in cellular masses, with shining pearly lustre, of a greenish grey color; it contains 80 per cent. alumine, 3 iron, and 17 water.

DOMITE. The substance to which this name has been given, occurs massive, of a white or greyish white color; sometimes tinged yellow; mica is frequently embedded in it.

DYSODILE occurs massive, compact, or laminated, of a greenish grey or yellow color; it is fragile; when under the blow-pipe it emits a most powerful fetid odor; it is from Melili in Sicily.

EUCHYSIDERITE. A newly described substance, found in prisms, the surface of which is brilliant, similar to augite, but without regular terminations. It is from Norway.

EUDEALITE is of a red color, and accompanied by sodalite and hornblende. It is composed of silica, zirconia, and soda: from Greenland.

FAHLUNITE is found massive, and imperfectly crystallized, in four or six-sided prisms: it is of a brownish black color, with an uneven or flat conchoidal fracture and resinous lustre. It seems nearly allied to precious serpentine, and has been called Tricklarite.

FIBROLITE occurs in fibrous masses, and sometimes prismatic; its colors are white or grey. It has a glistening lustre, and is harder than quartz. It is found in the East Indies, forming the matrix for corundum.

FUSCITE is found massive and crystallized in prisms, sometimes aggregated, of a greyish or greenish black color; it is found near Arendahl in Norway, in rolled masses of granular quartz.

GIBBSITE is described as occurring stalactitic and in aggregated, tuberous masses. Its color is white, grey, or greenish; it is composed of alumine and water.

GIESECKITE occurs crystallized in six-sided prisms, of a dark green, or brownish green color, with an uneven dull fracture, and is moderately hard, affording a nearly white streak with the knife.

HATCHETINE occurs of a yellowish or greenish color, in small quartz crystals; it is found flaky, with an appearance like spermaceti or bee's wax; it has a glistening or pearly lustre, is sometimes translucent, and sometimes dull and opaque; it is soft like tallow. It was discovered by the Rev. J. J. Conybeare, and is named after the celebrated chemist, Mr. Hatchett.

HEDENBERGITE. A substance of a blackish green color, very much resembling hornblende. It is massive; and according to Hedenberg, by whom it was analyzed, consists of silex, alumine, water, iron, manganese, and carbonic acid.

HELVIN is a crystallized substance of a red brown, or pale yellow color: it occurs in a dull green matrix of chlorite. Its form is said to be a tetrahedron modified.

HISINGERITE is said to be found massive, of a black color, according to Berzelius (by whom it was named); it contains oxide of iron, silex, alumine, oxide of manganese, and a small portion of magnesia.

HOLMITE. This substance, as described by the late Dr. Clarke, occurred in the form of an oblique rhombic prism; it consisted of lime 27, carbonic acid 21, alumine $6\frac{1}{2}$, silex $6\frac{1}{2}$, oxide of iron 29, and water 10. It was found among the paving stones at Cambridge.

HORN MANGAN. This substance is described as being compact, with conchoidal fracture, translucent on the edges, glistening, and becoming brilliant by exposure; its colors are white, grey, brown, and greenish blue; by analysis it consists of protoxide of manganese, silex, carbonic acid, water, and oxide of iron.

HUMBOLDTINE (Oxalate of Iron) is described as occurring in small flattish masses, and crystallized, of a bright yellow color; it is so soft that it may be easily scratched by the nail. It consists of protoxide of iron and oxalic acid, and is supposed to result from the decomposition of succulent plants.

HUMITE. Occurs in very small crystals of a deep reddish brown color; transparent, with a shining lustre. It is named in honor of Sir A. Hume. It is found on Mount Somma.

JEFFERSONITE. This substance occurs massive and crystallized, of a dark brown or green color, disseminated with red oxide of zinc; it is translucent on the edges.

KARPHOLITE. Described as being of a yellow color; it occurs in minute fibrous crystals, radiated, also amorphous; it is generally translucent, with glistening lustre. It is found in Bohemia. It consists of silex, alumine, oxide of manganese, protoxide of iron, and water.

KILLINITE is found in layers dispersed through a very

light-colored granite: the color is a brownish yellow, and the structure compact or slightly lamellar. It is moderately hard, and yields a light-colored streak.

KNEBELITE occurs massive, of various colors; the surface, cellular and uneven, with glistening lustre; it consists of siliceous, protoxide of iron and manganese.

KONILITE is found in the form of a white powder, gritty; it consists chiefly of siliceous, with a small portion of lime; it is easily fusible with the blow-pipe.

LATROBITE is described of a pale pink color; found massive and imperfectly crystallized. It is supposed to be a variety of feldspar. It was received by the Rev. C. I. Latrobe from Amitok Island, on the coast of Labrador.

LENZINITE is described as occurring massive, opaline, and argillaceous; of a milk white color, tinged yellow by oxide of iron. The fracture is dull and earthy.

LEUTTRITE resembles the Domite already described; it is of loose texture, and harsh to the touch; its color is greyish white; small fragments of mica are sometimes embedded in it.

LIGURITE. This substance is said to occur in oblique rhombic prisms, of an apple green color, sometimes externally specked; it falls to pieces in water; it is composed of siliceous, alumine, and lime.

LIMBILITE. Is described as being compact, of a yellow color. It was found in the Volcanic Hill at Limbourg.

LYTHRODES. This substance is found massive and disseminated, of various shades of red or brown, sometimes with green or yellow spots. It is from Norway.

MARGARITE. Described as occurring in masses of a greyish white color, and in delicate laminae, intersecting in every direction; it much resembles mica, of which it is probably a variety.

- MONTMARTRITE** occurs in masses of a yellowish or greenish grey color. It consists of sulphate of lime 83, with carbonate of lime 17, and may be placed after the other varieties of gypsum.
- NECRONITE** occurs in small masses of a clear white or bluish white color, with silky lustre on the exterior. It has much the appearance of feldspar, of which it will probably prove to be only a variety.
- OMPHACITE** is a lightish green variety of augite, which was separated by Werner, who considered it a distinct species.
- ORTHITE** has a general resemblance to gadolinite; it is composed of about 19.5 oxide of Cerium, 32 siliceous earth, 15 alumina, 12.5 oxide of iron, with manganese, yttria, lime, and water.
- PHOTIZITE** is described as being a compact substance, and varies in color from white to yellowish brown, green, and red of different shades. It appears to be a variety of manganese.
- PICOLITE** is a name given to a mineral found in the Pyrenees, which has not yet been analyzed, but whose external characters are very similar to those of Gadolinite.
- PICROLITE** is described as occurring massive, of a green or yellow color, with a fibrous structure. It is said to be composed principally of carbonate of magnesia.
- POLYHALLITE** is massive, with a fibrous structure, of a yellowish brick-red color. It is composed of common and anhydrous sulphates of lime, sulphate of potash, and anhydrous sulphate of magnesia: it is the fibrous anhydrite of some authors.
- PYRALLOLITE.** This substance is found massive and crystallized in flat rhombic prisms, resembling tremolite; it has a greenish tinge, but becomes yellow by ex-

posure; it consists of silex, alumine, lime, magnesia, and oxides of iron and manganese; it is found with augite, feldspar, and scapolite, in calcareous spar. Finland.

PYRENITE. The substance described under this name is said to be a variety of garnet; it occurs in small rhombic dodecahedrons, which have a black color, and glisten externally.

PYROSMAILITE occurs massive, or in six-sided prisms; it is of a liver brown color, and has a shining lustre, with a glimmering fracture. The composition appears to be oxides of iron and manganese, with silex, and about 14 per cent. of sub-muriate of iron.

RATHOFFITE. The substance under this name appears to be a variety of garnet; it occurs in rhombic dodecahedrons having its angles replaced; it has a brownish black color; it was brought from Sweden.

RHODONITE. Is said to occur massive and compact, of a pink, red, or yellowish white color, with fibrous structure and splintery fracture, and shining lustre; it consists of protoxide of manganese, silex, carbonic acid, alumine, water, and oxide of iron.

SIDERITE. This substance is described as being compact, of a greyish or greenish blue color; it is nearly as hard as quartz, and has a resinous lustre. It is found in granular gypsum, near Salzburg.

SIDEROCLEPTE. This substance is described as being massive, of a yellowish green color, translucent; it yields to the nail, and has a greasy lustre. It was found in the cavities of lava at Brigaw.

SKOLEZITE. Is described as being massive and crystallized; it is nearly transparent, and colorless; under the flame of the blow-pipe it twists itself up like a worm, (whence its name from the Greek in allusion to that property). It is found at Parges in Finland.

SKORODITE is usually crystallized in broad four-sided prisms, terminated on each extremity by four-sided pyramids; its color is green, passing into brown; the lustre is somewhat vitreous, the fracture uneven, and in composition it seems to be an impure arseniate of iron, free from copper.

SORDAWALITE is described as being compact; its color is nearly black, sometimes grey or green; by exposure it becomes red externally. It is opaque, with conchoidal fracture. It is composed of silex, alumine, magnesia, protoxide of iron, phosphoric acid, and water. It is found at Sordawala, whence its name.

SPAK. The mineral frequently described under this name is merely fibrous rock salt.

STILPNOSIDERITE occurs massive, small reniform, and lamellar; it is of a brownish black color, splendent, and opaque, with a nearly perfect conchoidal fracture. It is probably a variety of meadow iron ore.

STROMNITE—BARYSTRONTIANITE has been found in masses of a greyish or yellowish white color; and appears disintegrated externally; it is translucent on the edges, brittle, and soft. It is composed of carbonate of strontian, sulphate of barytes, carbonate of lime, and oxide of iron; it derives its latter name from containing barytes and strontian; and stromnite from being found at Stromness.

SUCCINITE occurs in globular masses, the size of a pea, of an amber yellow color, translucent; it is considered an amorphous variety of topazolite. From Piedmont.

THOMSONITE. The substance so named occurs in masses; it is colorless and translucent; it has a radiated structure, and pearly lustre. It was formerly called Mesotype, but has been newly named after Dr. Thomson. It is from Scotland.

THULITE. The mineral under this name occurs in crystalline masses, of a rose red color; it is sometimes found in masses with quartz, fluor, and idocrase. The fracture is uneven or slightly lamellar, with a somewhat resinous lustre.

TURNERITE. The substance to which this name has been given, occurs in small crystals of a yellowish brown color, translucent, brilliant on the exterior; it was considered a variety of sphene, but contains no titanium; it consists of alumine, lime, magnesia, and small portions of iron and silex. It has sometimes been called pictite; it is from Dauphine.

VAUQUELINITE—*Chromate of Lead and Copper.* This substance occurs in minute aggregated crystals, of a black color, sometimes tinged green; it consists of oxide of lead, oxide of copper, and chromic acid. It is found in Siberia with chromate of lead.

WODAN PYRITES is found in vesicular masses of a tin white color, passing into brown; its lustre is shining and metallic; the fracture uneven: it is opaque, brittle, and rather harder than fluor, and is said to contain 20 per cent. of a new metal, called Wodanium, united with sulphur, arsenic, iron, and nickel.

WOLLASTONITE. The substance so named occurs in small masses, externally of a brown color, internally transparent and colorless, or of a flesh color. This mineral is only a variety of tabular spar.

ZURLITE is said to occur in rectangular prisms, and in roundish masses; its color is green, and fracture conchoidal; it is compact. It is found with calcareous spar on Mount Vesuvius.

SYNONYMES.



Acanticonite, another name for *Epidote*, from *Arendahl*.

Agaphite, *Iohnite*, and *Calaité*, are other names for *Turquoise*.

Albin, *Cleavelandite*, for *Albite*.

Almandine, another name for *Precious Garnet*.

Aphrizite, for *Black Tourmaline*.

Cordierite, for *Iolite*.

Endellione, for *Bournonite*.

Euhairite, for *Seleniuret of Copper*.

Fassaite, *Pygrom*, for *Diopside*.

Gabronite, for *Compact Scapolite*.

Gahnite, for *Automalite*.

Keffehil, for *Meerschaum*.

Koupholite, for *Prehnite*.

Latialite, for *Haiÿne*.

Lhergolite, for *Coccolite*.

Liehuites, for *Marble*.

Kaolin, for *Porcelain Clay of China*.

Loboite, is a variety of *Idocrase*.

Malacolite, another name for *Sahlite*.

Maltha, for *Bitumen*.

Mesolite, for *Needle Zcolite*.

Miarcite, for *Bitterspar*, from *Miarka* in *Siberia*.

Novaculite, for *Whetslate*.

Pelium, *Peliome*, for *Iolite*.

Physalite, for *Pyrophyssalite*.

Picropharmacolite is a variety of *Pharmacolite*.

Romanzovite is a variety of *Cinnamon Stone* from *Finland*.

Sarcolite is a variety of *Analcime*, so called from its being of a flesh-red color.

Scorza is another name for *Granular Epidote*.

Siberite, for *Rubellite*, from *Siberia*.

Steinheilite, for *Dichroite*.

Surturbrand, for *Brown Coal*, so called in *Iceland*.

Trona, for *Natron*, *Carbonate of Soda*.

Geology.



ROCKS are either simple or compound; the simple are those which consist entirely, or at least essentially, of one mineralogical species, and are therefore arranged amongst the simple minerals. Such are limestone, gypsum, serpentine, rock salt, and coal.

The compound rocks are formed of two or more mineralogical species, variously aggregated, in different proportions, and differing in magnitude.

According to Werner, there are three distinct classes of rocks, formed at different and very distant periods: viz. primitive, transition, and secondary, or floetz.

PRIMITIVE ROCKS,

AS the term implies, are such as were formed first; they do not contain fossil remains, and are supposed to have had their origin before the creation of animal or vegetable substances, and to be the result of chemical precipitation.

GRANITE is a crystalline aggregate, consisting of quartz, feldspar, and mica, promiscuously arranged.

GNEISS—*Slaty Granite*, consists of quartz, feldspar, hornblende, or mica, stratified, waved, or in patches; the structure is commonly slaty.

MICA SLATE consists of quartz and mica, generally laminated. This rock is considered to pass into clay slate.

CLAY SLATE. This is a simple rock, often lamellar.

PRIMITIVE LIMESTONE is that which contains no fossil remains, and is generally crystalline in its structure.

PRIMITIVE TRAP is principally composed of hornblende, and sometimes feldspar, generally dark coloured, forming a crystalline aggregate.

SERPENTINE is a simple rock; it contains hornblende, and sometimes veins of asbestos and steatite; it is frequently spotted.

PORPHYRY is a compact indurated substance, not unlike jasper; it contains crystals of feldspar, embedded; color, generally red, green, or brown.

SIENITE. This rock consists of feldspar and hornblende; it is of various colors, as reddish, dull green, &c. as the feldspar or hornblende may predominate.

TOPAZ ROCK is an aggregate of feldspar and quartz, containing topazes, and frequently schorl.

QUARTZ ROCK is compact; its varieties frequently contain schorl and other substances.

FLINTY SLATE is a black compact substance of close texture; it often contains veins of quartz.

PRIMITIVE GYPSUM is massive, and occurs with the preceding rocks.

WHITE STONE is apparently a variety of fine-grained granite, chiefly composed of granular feldspar.

GRANITE.

Common Granite. Large-grained.

Variety, containing schorl.

Variety.

Variety.

Variety, small grained.

Variety.

Variety.

Graphic Granite. Exhibiting the appearance of Hebrew letters.

Variety.

Porphyritic Granite. Containing large crystals of feldspar embedded in fine-grained granite.

Orbicular Granite.

Granite, the feldspar of which is decomposing and forming clay.

Variety.

GNEISS. Stratified.

Variety.

Variety, with garnets.

Variety, containing hornblende.

Variety.

MICA SLATE. Waved.

Variety, slaty.

Variety.

Variety, with crystals of garnet embedded.

Variety, with schorl embedded.

Variety.

CLAY SLATE.

Common Slate.

Variety, with chialstolite embedded.

Variety, with veins of tin ore.

Variety, with pyrites.

Variety.

PRIMITIVE LIMESTONE.

Tiree Marble. Massive, containing sahlite.

Dolomite. Granular limestone.

Variety.

Carrara Marble.

Variety.

PRIMITIVE TRAP. Fine grained.

Variety.

Variety, with feldspar.

Variety.

SERPENTINE.

Variety, dark coloured.

Variety, spotted.

Variety.

Variety, with asbestos.

Variety.

PORPHYRY.

Red Porphyry.

Brown Porphyry.

Variety.

Green Porphyry. (Oophites of Pliny).
Variety.

SIENITE. With red feldspar and green hornblende.
Variety.
Variety.

TOPAZ ROCK. Massive.
Variety, with topazes embedded.

QUARTZ ROCK. Massive.
Variety.

Schorl Rock. Massive.
Variety, schorl embedded in quartz.

FLINTY SLATE.
Variety.

PRIMITIVE GYPSUM.

WHITE STONE.

Thus ends the series of what are termed primitive rocks, according to Werner's theory; but they must be supposed to form infinite variety in the actual proportions of their constituents, and (notwithstanding the regular arrangement of authors) to have undergone great alterations from various causes. The formation of what are considered to be primitive and transition rocks is by no means a determined point.

TRANSITION ROCKS.

THE Transition Formation, as it is termed, according to Werner, contains only four distinct species, which are supposed to have been formed during the transition of the earth from its uninhabitable to its habitable state. Hence they sometimes contain the first traces of organic remains, partaking of both chemical and mechanical deposit; and connecting the primitive with the Floetz Formation.

It is not easy to form any thing like a correct opinion of the alteration substances undergo, on being exposed to the action of water or atmospherical changes, for a series of ages. The rocks of this class may be supposed to be originally placed at the base of the primitive, filling ravines, or skirting mountains.

TRANSITION LIMESTONE is fine grained, and is most common in Devonshire, where it fills ravines between clay-slate; it sometimes contains traces of fossil remains.

TRANSITION TRAP—*Green Stone*. This substance forms great variety, and is considered to be composed of feldspar and hornblende, in different proportions, and of different colors. It is less crystalline than the primitive order.

GREY WACKE is a mechanical deposit, in which are embedded fragments of primitive rock, in coarse and fine particles; it is sometimes slaty, and probably migrates into an earthy sandstone. It is extremely variable in its appearance and texture, and is considered to be of great extent.

TRANSITION FLINTY SLATE is stratified chert or flint.

TRANSITION LIMESTONE.

Variety, green.

Variety, red.

Variety, with organic remains.

TRAP—*Greenstone.*

Variety.

Variety, with veins.

GREY WACKE.

Variety, composed of clay-slate and fragments of primitive rocks.

Variety.

Variety.

Grey Wacke Slate.

Variety, fine grained.

Variety, with organic remains.

Variety.

Transition Flinty Slate.

Variety, stratified.

* * The two last Rocks are not well understood; geologists by no means agree in describing their characters and components. The same may be said of old red sandstone, the first in the following series.

FLOETZ OR FLAT FORMATION.

CONSISTS of a series of rocks hereafter enumerated, that are supposed to be the result of decomposition, containing animal and vegetable remains, and formed by aqueous deposit. This formation fills up vast tracts betwixt elevated rocks of the primitive or transition class.

OLD RED SANDSTONE is an aggregate extremely variable in appearance, consisting chiefly of silicious particles produced from the debris of the preceding formations, on which it is considered to rest. Its structure varies from very fine to very coarse.

FLOETZ LIMESTONE is generally of a dull color and compact texture; it contains more or less of fossil remains, such as echinites, madrepores, zoophytes, &c. and often entirely composed of marine remains. It is evidently formed under water. It is a formation of great extent, and is highly metalliferous, and regularly stratified.

TOADSTONE—TRAP alternates with the Derbyshire floetz limestone. It migrates into a variety of amygdaloid, and decomposes into clay.

FIRST GYPSUM is not of great extent, though in considerable abundance. In some cases it rises into small hills, and fills cavities; it is always accompanied with clay and marl.

VARIEGATED SANDSTONE is colored by the oxide of iron, red or yellow of various shades. It is a deposit, more or less pure; it is marked with lines, and often stratified.

SHELL LIMESTONE is evidently of more modern formation than the limestones before mentioned; it is almost wholly composed of shells and marine substances, such as bivalves, ammonites, &c.

THIRD SANDSTONE is of modern formation, and may be considered the uppermost, and what is daily forming by accumulation on the banks of rivers, &c.

SECOND GYPSUM occurs filling cavities, and surrounded by clay, sandstone, &c. It is fibrous, and very soft.

ROCK SALT. Its situation is peculiar; it is very widely diffused, being under some varieties of sandstone, and above others. It is accompanied by gypsum and clay. In this country the beds of rock salt are about fifty to sixty fathoms deep, and are peculiar to Cheshire.

CHALK FORMATION is of considerable extent; it contains a great variety of animal, vegetable, and marine remains; also flint in great abundance, and pyrites. It is of different qualities, regularly stratified, and contains beds of sand, marl, and clay.

BITUMINOUS SHALE, which forms a large range in Derbyshire, is not even noticed in the Wernerian series. Its place is directly above the floetz limestone.

COARSE GRIT lies immediately upon the preceding, and under the coal formation. It is a granular aggregate, containing feldspar, sometimes schorl, and oxide of iron. It is considered to be both a mechanical and chemical deposit. It often resembles old red sandstone.

COAL FORMATION occurs alternating with shale and sandstone, accompanied by beds of clay iron-stone.

NEWEST TRAP is probably an earthy homogeneous substance; it is not met with in our coal formation, except in a disintegrated state, or decomposed into clay. There are substances, called Whinstones, intersecting

the coal formation, which occur in large veins or dykes. They vary in color and texture.

Varieties of basalt, trap-tuff, green-stones, amygdaloid, pitch-stones, and jaspers, are said to belong to this formation.

OLD RED SANDSTONE. Fine-grained.

Variety, coarse.

Variety.

FLOETZ LIMESTONE. Massive.

Variety, with fossil remains.

Coralloidal.

Variety.

Botryoidal.

Variety, black.

TOADSTONE—TRAP. Compact.

Variety.

Variety.

Variety, containing pyrites.

Variety.

Variety, in decomposition.

Amygdaloid.

Variety.

Variety.

Variety, in decomposition.

Wacke, Cellular.

Variety.

Variety, partially cellular.

FIRST GYPSUM.

Variety.

VARIEGATED SANDSTONE. Red.

Variety, yellow.

Variety, green.

Variety.

SHELL LIMESTONE.

Variety.

Variety.

THIRD SANDSTONE.

Variety.

SECOND GYPSUM.

Variety.

ROCK SALT. Massive.

Variety.

CHALK FORMATION. With marine remains.

Variety.

With pyrites.

BITUMINOUS SHALE.

Variety.

Variety, with impressions.

COARSE GRIT. With feldspar.

Variety.

COAL FORMATION. Coal, with pyrites.

Variety, with sandstone.

Variety.

NEWEST TRAP.

Whinstone, Jew Stone, and Dyke Stones.

Variety.

Variety.

Basalt.

Variety.

Variety, with Olivine.

Trap Tuff. Fragments of trap, coarsely cemented.

TUFFA is a calcareous substance, coating vegetables, &c. which is deposited by water issuing from limestone rocks, and containing lime in solution. These waters have received the name of Petrifying Wells; and if twigs, straw, nests, &c. be laid in them for a year, they will become coated with this deposit*.

* For more particulars upon this subject, see the Author's Familiar Lessons on Mineralogy.

The Author meets with such difficulty, in finding words to distinguish many of the characters which minerals assume, that he feels obliged to make use of the term *Variety*, as a reference to undecided specimens.

ALLUVIAL DEPOSITS.

CASCALHAO is an alluvial deposit, consisting chiefly of rounded and angular pebbles, with sand and clay; this formation is immediately incumbent on the rock in the gold district of Brazil: and amongst the loose stones are found diamonds, gold, topaz, amethysts, &c. This stratum is often covered many feet deep by vegetable earth, forming the richest soil*.

GRAVEL generally consists of pebbles of quartz, chalcidony, jaspers in great variety, &c. which have been rounded by attrition. Pieces of limestone, rounded tin ore†, sometimes galena, fossil wood, iron-stone, animal remains apparently passing into a fossil state, and various other substances, are often found in gravel. In some places this deposit is above forty feet thick.

Gravel and sand sometimes become cemented, and form an aggregate, called pudding stone.

SAND and MARL are too well known to require any description.

* For a more particular description, see the Author's Travels through the Diamond District in Brazil.

† In the island of Banca and other isles in the Indian Archipelago, an enormous quantity of rounded tin pebbles is found. They are also abundant in Cornwall.

VOLCANIC ROCKS.

ROCKS, in which volcanoes are situated, are called volcanic, and are so, if altered by fire; they form a very extensive class, and present considerable varieties, which generally contain olivine, augite, &c. They are commonly black, and more or less cellular; and often, (more particularly the lavas, cinders, and ashes), form whole islands, or extend over considerable territories.

Rocks, altered by fire, have a peculiar vitreous appearance, and may generally be easily distinguished; it frequently occurs that great varieties of crystallized substances are embedded in them.

VOLCANIC ROCK.

Variety.

Variety, containing Olivine.

Variety, containing Augite.

Lava, compact.

Variety.

Variety, cellular.

Variety.

Variety, spongy.

Variety.

Ashes, volcanic.

THE END.

INDEX.

Abrazite	145	Amber	142
Acanticonite	155	Amethyst	79
Actynolite	114	Amianthus	112
asbestos	114	Amianthoide	112
common	114	Amygdaloid	120, 155
glassy	114	Andalusite	98
Adhesive Slate	103	Anhydrite	134
Adularia	98	Antimonial Lead Ore ..	36
Agalmatolite	110	Silver	4
Agaphite	155	Antimony	50
Agaric Mineral	123	ochre	50
Agate	89	Anthophyllite	115
blue	89	Apatite	130
breccia	89	Aphrizite	155
fortification	89	Aphrite	123
landscape	89	Aplome	77
moss	89	Apophyllite	92
oriental	89	Aqua-marine	72
petrification	89	Arfwedsonite	145
striped	89	Arragonite	125
Albin	155	Arsenate of Cobalt	53
Albite	99	Copper	19
Allagite	146	tabular	20
Allanite	62	octahedral	20
Allochroite	77	prismatic	20
Allophane	93	Iron	29
Alluvial Deposits	169	cupreous	29
Almandine	155	Lead	38
Alum Earth	141	Arsenic	57
Alum, native	143	Arsenical Cobalt	53
Slate	104	Copper	16
Stone	103	Nickel	56
Aluminite	103	Pyrites	57
Amblygonite	145	Silver	4

Asbestos	112	Blue Spar	97
Asparagus Stone	130	Vitriol	64
Asteria	70	Bog Iron Ore	26
Atacamite	20	Bole ..	109
Avanturine	79	Boracite	136
Augite	118	Borax ..	144
Automalite	68	Botryolite	136
Axe stone	111	Bovey Coal	141
Axinite	73	Bournonite	36
Azurite	97	Brazilianite	96
Baikalite	115	Brewsterite	145
Barystronianite	153	Bronzite	115
Barytes	137	Brown Coal	141
Basalt	120, 157	earthy	141
Basanite	80	Cobalt Ochre ..	55
Bergmanite	99	Hematites	25
Beryl	72	Iron Ore	25
Bismuth	47	scaly	25
ochre	47	compact	25
Bismuthic Silver	4	Ochre	25
Bitter Spar	121	Spar	121
Bitumen, elastic	140	Bucholzite	146
indurated	140	Byssolite	112
Bituminous Shale ..	141, 165	Cacholong	86
Slate	104	Cadmiferous Blende ..	63
Wood	141	Calamine	63
Black Coal	141	Cadmium	63
Cobalt Ochre ..	53	Cairn gorm ..	81
Copper	15	Calaita	155
Hematites	25	Calamine	42
Iron Ore	25	cupriferous	42
Marble	125	Calamite	146
Manganese	31	Calcareous Spar	124
Ore of Nickel	56	Tuffa	125, 168
Sulphuret of Silver ..	6	Cannel Coal	141
Blende, black	43	Carbonate of Barytes ..	137
brown	43	Copper, hydro blue ..	17
green	43	green	17
yellow	43	Iron	29
Bloedit	145	Lead	38
Blood-Stone	84	black	38
Blue Iron Ore, earthy ..	22	compact	38
Lead Ore	36	earthy	38

Carbonate of Lime	126	Clay Iron ore	25
granular	124	columnar	25
crystallized	124	Cleavelandite	155
fibrous	124	Clink Stone	120
Magnesia	109	Coal	141
Manganese	33	black	141
Silver	8	brown	141
Soda	143	Cannel	141
Strontian	138	foliated	141
Carinthin	146	Glance	141
Carnelian	83	Newcastle	141
Cascalhao	169	shining	141
Cat's eye	80	slaty	141
Celestine	138	Coarse Grit	165
Cereolite	146	Cobalt	53
Cerite	62	Glance	53
Cerium	62	Coccolite	118
Ceylonite	68	Colophonite	77
Chabasite	93	Comptonite	146
Chalcedony	83	Condroidite	146
Chalk	123	Conite	146
Chert	80	Copper	12
Cheveux de Venus	34	azure	17
Chiastolite	98	black	15
Chlorite	105	Glance	13
common	105	Nickel	56
foliated	105	Pyrites	14
Slate	105	Copperas	64
Chloropal	146	Coralloid Limestone	124
Chlorophane	131	Cordierite	155
Chlorophæite	146	Corundum	68
Chromate of Iron	29	Couzeranite	147
Lead	38	Crichtonite	34
green	38	Cronstedite	147
Chrome	62	Cross-Stone	93
Chrysoberyl	69	Cryolite	139
Chrysolite	118	Crysocolla	18
Chrysoprase	84	Cube Spar	134
Cimolite	107	Cubicite	92
Cinnamon Stone	77	Cupreous Bismuth Ore	47
Cinnabar	10	Manganese	33
Clay, common	103	Cymophane	69
Slate	103, 158	Datholite	136
variegated	103	Desmine	147

Diallage.....	115	Flint	80
Diamond	66	Flinty Slate	80, 158
Diaspore	147	transition	163
Dichroite	72	Float Stone	104
Diopside	118	Floetz Formation	164
Diopase	19	Fluor	131
Dipyre	93	argillaceous	131
Dolomite	121, 160	compact.....	131
Domite	147	earthy	131
Drawing-Slate	104	foliated	131
Dyke Stone	168	Fontainebleau Sandstone	79
Dysodile	147	Fossil Copal	142
Earthy Talc	112	Fullers' Earth	109
Endellione	155	Fuscite	148
Egeran	76	Gabronite	155
Elaolite	90	Gadolinite	61
Electric Calamine	42	Gahnite	155
Electrum	3	Galena	36
Emerald	72	Garnet, common	76
Emery	68	precious	76
Epidote	73	Gehlenite	76
Epsom-Salt	148	Geology	157
Euclase	72	Gibbsite	148
Euchysiderite	147	Gieseckite	148
Eudialite	147	Girasol	85
Eukairite	155	Glauberite	134
Fahlertz Grey Copper ..	13	Glauber-Salt	143
Fahlunite	148	Gneiss	158
Fassaite	155	Gold	2
Feldspar	98	Granite	157
common	98	graphic	159
compact	99	orbicular	159
disintegrated	99	porphyritic	159
glassy	98	slaty	158
Labrador	98	Graphite	142
radiated	99	Gravel	169
Fettstein	99	Green Earth	107
Fibrolite	148	Stone	162
Figure-Stone	109	Grenatite	77
Fiorite Pearl Sinter	84	Grey Antimony	50
Fire Marble	124	nickeliferous	50
Flexible Limestone	121	Cobalt	58
Sandstone	79	Copper	13

Grey Manganese	31	Indianite	98
compact	31	Indicolite	72
earthy	31	Iohnite	155
foliated	31	Iolite	72
radiated	31	Iridium	2
Silver ore	6	Iron	22
Wacke	162	Iron Froth	24
Slate	163	Glance	24
Grossular	76	micaceous	24
Gurhofite	121	Pyrites	23
Gypsum	133	hepatic	23
compact	134	magnetic	23
earthy	134	Iserine	34
fibrous	134	Jade	111
first	164	Jasper	87
foliated	134	Jasper agate	89
plumose	134	common	88
primitive	158	Egyptian	87
second	165	porcelain	88
Harmatome	93	striped	88
Hatchetine	148	Jeffersonite	149
Haüyne	97	Jet	141
Heavy Spar	137	Jew Stone	168
Hedenbergite	148	Kaolin	155
Heliotrope	84	Karpholite	149
Helvin	148	Keffekil	155
Hepatic Cinnabar	10	Killinite	149
Hisingerite	149	Kuebelite	150
Holmite	149	Kollyrite	107
Honey Stone	142	Konilite	150
Hornblende	114	Koupholite	155
Slate	114	Kyanite	115
basaltic	114	Lapis Lazuli	96
Horn Mercury	10	Latialite	155
Mangan	149	Latrobite	150
Hornstone	80	Launonite	93
Humboldtine	149	Lazulite	97
Humite	149	Lead	36
Hyacinth	67	Leelite	80
Hyalite	84	Leivrite	118
Hydrophane	85	Lemnian Earth	109
Hyperstene	115	Lenticular Iron Ore	25
Ice Spar	100	Lenzinite	150
Idocrase	76	Lepidolite	105

Leucite	76	Meionite	99
Leuttrite	150	Melanite ..	77
Lhergolite	155	Menachanite	34
Ligurite	150	Menilite	87
Liehuities	155	Mercury	9
Limbilite	150	Mesolite	155
Limestone	123	Meteoric Native Iron ..	22
common	123	Miarcite	156
compact	123	Mica	105
coralloid	124	slate	158
fibrous	124	Miemite	121
floetz	164	Mineral Charcoal	142
granular	124	Mocha Stone	83
magnesian	121	Molybdate of Lead ..	38
primitive	158	Molybdena	52
shell	124, 165	Ochre	52
transition	162	Mona Marble	124
Lithomarge	107	Montmartrite	151
Loam	103	Moon Stone	98
Loboite	155	Morass Iron Ore	26
Lucullite	125	Moroxite	130
Lumachella	124	Mountain Soap	107
Lydian Stone ..	83	Muriate of Copper ..	19
Lythrodes	150	Lead	38
Macle	98	Mercury	10
Magnesia, native	109	Silver	8
Magnesian Limestone ..	121	Soda	144
Magnesite	109	Mussite	118
Malachite	17	Nacrite	112
Malacolite	155	Naptha	140
Manganese	31	Native Alum	143
Maltha	155	Amalgam	10
Marble	124	Antimony	50
black	125	Arsenic	57
Carrara	125, 160	Bismuth	47
green antique	125	Copper	12
Lumachella	124	Gold	3
Mona	124	Iron	22
Tiree	124, 149	meteoric	22
Marekanite	90	terrestrial	22
Margarite	150	Loadstone	24
Marl	125, 158	Magnesia	109
Meadow Iron Ore ..	26	Mercury	10
Meerschaum	109	Nickel	56


Native Platina	1	Oxide of Iridium	2
Silver	4	Nickel	56
auriferous	4	Tin	45
Sulphur	140	Uranium, black	60
Tellurium	49	green	60
Natrolite	93	Zink, red	42
Natron	143	Palladium	1
Necronite	151	Paranthine	99
Needle Ore	47	Pargasite	114
Nepheline	99	Peacock Coal	141
Nephrite	111	Pea Iron Ore	25
Newcastle Coal	141	Pea Stone	125
Nickel	56	Pearl Spar	121
Ochre	56	Stone	90
Nigrine	34	Peliome	156
Nitre	144	Pelium	156
Nodular Iron Ore	25	Petalite	99
Novaculite	156	Petroleum	140
Obsidian	90	Pharmacolite	58
Octahedrite	34	Phosphate of Copper	19
Olivine	113	Iron	29
Omphacite	151	earthy	29
Onyx	83	Phosphate of Lead	38
Oolite	123	Lime	130
Opal	85	Manganese	32
common	85	Phosphorite	130
fire	85	Photizite	151
golden	86	Physalite	156
harlequin	86	Picolite	151
jasper	86	Picrolite	151
precious	85	Picropharmacolite	156
semi	86	Pimelite	107
wood	86	Pinite	105
Oriental Amethyst	70	Pistazite	73
Emerald	70	Pitch Ore	60
Ruby	68	Stone	90
Topaz	70	Porphyry	90
Orpiment	57	Pitchy Iron Ore	26
Orthite	151	Plasma	84
Oxide of Antimony	50	Platina	1
Arsenic	53	Pleonaste	68
Oxide of Cobalt	53	Plumbago, compact	142
Copper, red	16	scaly	142

Polishing Slate	104	Red Iron Ore	24
Polyhallite	151	Froth	24
Porcelain Earth	103	Glance	24
Porphyry	158	scaly	24
Pot-Stone	111	Lead	38
Potters' Clay	103	Manganese ..	38
Prase	79	Ochre	24
Prehnite	92	Oxide of Copper ..	16
Pudding-Stone	88, 169	ferruginous ..	16
Pumice	90	Zinc	42
Purple Copper Ore .. .	13	Silver	6
Pycnite	72	Reddle	25
Pygrom	155	Retin Asphalt	142
Pyrope	76	Rhätizite	115
Pyralloite	151	Rhodonite	152
Pyrophyllite	72	Rhomb Spar	121
Pyrenite	152	Rock Cork	112
Pyrosmalite	152	Crystal	79
Quartz	79	Milk	123
Aventurine	79	Salt	144, 165
Babel-tower	81	Wood	112
black	81	Rocks, primitive	157
blue	81	secondary, transition ..	162
cap	81	volcanic	170
common	79	Roe Stone	123
ferruginous	80	Romanzovite	156
float	79	Rotten Stone	104
milky	81	Rubellite	72
paper	81	Ruby	68
pseudamorphous .. .	79	Rutile	34
red	81	Sahlite	118
rhombic	79	Sand	169
rock	158	Sandstone, flexible	79
rose	81	Fontainebleau	79
yellow	81	old red	164
Rathoffite	152	third	165
Realgar	57	variegated	164
Red Antimony	50	Sappare	115
Cinnabar	10	Sapphire	68
Cobalt	53	blue	68
radiated	53	red	68
slaggy	54	Sarcolite	156
Hematites	24	Sardonyx	83


Sassolin	144	Sphærolite	90
Satin Spar	124	Sphene	34
Sausurite	98	Spinelle	68
Scapolite	99	Spodumene	99
Schiefer Spar	123	Stalactite	124
Schiller Spar	115	Staurotide	77
Schorl	72	Steatite	109
rock	161	Steinheilite	156
Schorlite	72	Stilbite	92
Scorza	156	Stilpnosiderite	153
Selenite	134	Stream Tin	45
Seleniuret of Copper ...	63	Stromuite	153
Selenium	63	Strontianite	138
Serpentine	111, 158	Succinite	153
precious	111	Sulphate of Ammonia ..	144
Shale	165	Barytes	137
Shell Limestone	124	Cobalt	64
Shining Coal	141	Copper ..	64
Siberian Topaz	73	Iron	64
Siberite	156	Lead	38
Siderite	152	Soda	143
Sideroclepte	152	Strontian	138
Sienite	158	Zinc	64
Silicious Copper	18	Sulphur, native	140
Oxide of Chrome ...	62	volcanic	140
Tuffa	84	Sulphuret of Arsenic ...	57
Silver	4	Bismuth	47
Sinopal	88	Cobalt	53
Skolezite	152	Lead	36
Skorodite	153	antimonial ...	36
Slaggy red Cobalt Ochre	54	triple	36
Slate Clay	103	cobaltic	37
Coal	141	Manganese	32
common	160	Silver	5
Spar	123	Tin	45
Slaty Coal	141	Zinc	42
Slickenside	37	Surturbrand ..	156
Smaragdite	117	Swine Stone	125
Soap Stone	109	Tabular Spar	123
Sodalite	99	Talc	111
Sordawalite	153	columnar	112
Spak	153	compact	112
Sparry Anhydrite	134	Venetian	111
Iron Ore	29	Tantalum	61
Lead Ore	38	Tantalite	61

Tellurium	49	Uranium Ochre	60
Tellurium, graphic	49	Valentianite	109
yellow	49	Variegated Copper ore ..	13
black	49	Vauquelinite	154
Tennantite	13	Venetian Talc	111
Terrestrial Native Iron ..	22	Vesuvian	76
Thomsonite	153	Vitriol, blue	64
Thulite	154	white	64
Tin	45	Volcanic Glass	90
Tinder ore	50	Sal Ammoniac	144
Tiree Marble	124	Sulphur	140
Titanious Iron Ore	26	Vulpinite	134
Titanium	34	Wacke	120 155/162 166
Toadstone	164	Wad	31
Topaz	71	Wavellite	96
Brazilian	71	Wernerite	99
rock	158	Whet Slate	104
Saxon	71	Whinstone	165
Siberian	73	White Antimony	50
Topazolite	76	Copper	16
Tourmaline, precious	72	Silver Ore	6
common	72	Stone	158
Trap, primitive	158	Vitriol	64
transition	162	Wodan Pyrites	154
flötz	164	Wolfram	59
newest	165	Wollastonite	154
tuff	168	Woodstone	80
Tremolite	114	Wood Tin	45
asbestos	115	Yellow Cobalt Ochre ..	55
common	115	Earth	107
glassy	115	Yenite	118
Tricklarite	148	Yttrio Tantalite	61
Triphane	99	Zeolite	92
Tripoli	104	earthy	92
Trona	156	fibrous	92
Tuffa	168	foliated	92
Tungsten	59	needle	92
Turnerite	154	radiated	92
Turquoise	87	Zinc	42
Umber	26	Zircon	67
Uranite	60	Zoisite	73
Uranium	60	Zrinite	154














by
CD.

