


1902

USNM

EX LIBRIS

William Healey Dall

Division of Mollusks
Sectional Library







CONCHOLOGIA ICONICA:

OR,

ILLUSTRATIONS

OF THE

SHELLS OF MOLLUSCOUS ANIMALS.

VOL. V.

CONTAINING MONOGRAPHS OF THE GENERA

BULIMUS.

CASSIS.

CASSIDARIA.

ACHATINA.

TURRITELLA.

ONISClA.

DOLIUM.

MESALIA.

EBURNA.

EGLISIA.

BY

LOVELL AUGUSTUS REEVE, F.L.S., F.Z.S.,

CORRESPONDING MEMBER OF THE NATURAL HISTORY SOCIETY OF WURTEMBERG

HONORARY MEMBER OF THE IPSWICH MUSEUM.

" As the Snail whose tender horns being hit,
Shrinks hockwards in his shelly cave with pain,
And there, all smothered up, in shade doth sit,
Long after fearing to creep forth again."—*Shakespeare*.

LONDON:

REEVE, BENHAM, AND REEVE, KING WILLIAM STREET, STRAND.

1849.



MONOGRAPH

OF THE GENUS

B U L I M U S.

"The fertile bosom of the earth gives suck
To myriads, who still grow beneath her care,
Rewarding her with their pure perfectness."—*Shelley*.





B U L I M U S.

PLATE I.

Genus BULIMUS, Lamarck.

*Testa ovata, vel oblonga, vel torrita, angustibasis unus per-
plicis, ventricosis, nus plurimi, contractis, aut fra-
gilibus et sepe translucidis, aut crassiusculis, epidermae
interdum dupliciti undet; columella recta, interdum uni-
plicata, basi unquam truncata; aperta nonnunquam
dentata, plerunque integrâ, marginibus disjunctis, labo
vel simplici, vel reflexo. Operculum nullum.*

Shell ovate, oblong, or turreted, whorls sometimes very few and ventricose, sometimes many and contracted, either fragile and often transparent, or rather thick, covered with, sometimes, a double diaphanous epidermis; columella straight, sometimes one-plaited, never truncated at the base; aperture in some instances toothed, generally entire, with the margins disjoined, lip either simple or reflected.

The beautiful forms and varieties of shells produced by those air-breathing mollusks, which, under the generic appellation of *Bulinus*, constitute an important division of the great tribe of SNAILS, have become objects of especial interest to the conchologist, owing to the zeal with which a few enterprising scientific travellers have lately penetrated into tropical countries in pursuit of them. It is, however, to the productive exertions of Mr. Cuming that we are mainly indebted for the newer and more attractive subjects of the present monograph. The researches of this ardent naturalist in the arid plains on the west side of the Andes, in the dense woods of West Columbia and Central America, and, more recently, in the luxuriant open forests of the Philippine Islands, whilst they present an instructive contrast, exceed any result the most sanguine collector could have anticipated. In the dry and barren regions of Western Chili and Peru, the *Bulinai* are mostly small and of comparatively fragile structure; but in the beautiful islands of the Eastern Archipelago, where climate and vegetation combine to favour the growth of arboreal species, the genus is represented with prolific splendour. Mr. Cuming must have truly felt like one transported to the fabled garden of the Hesperides, when beholding the lofty trees of these sunny isles laden with snails of such magnificent proportions. Aladdin, in the Arabian tale, could not, surely, have contemplated the rich clusters of vari-coloured fruit in the garden of the African Magician with more astonishment, nor probably gathered it with more avidity.

"It was in 1836," relates Mr. Broderip, "that Mr. Cuming proceeded to the Philippine Islands by permission of the Queen Regent of Spain, and aided by powerful recommendations from her government, which opened to him the interior of the islands, and caused him to be received with a noble hospitality, equalled only by the warm interest which facilitated his pursuits wherever he arrived and made himself known." Species of which we had but an imperfect knowledge, in consequence of the bad condition in which a stray individual chanced to reach our cabinets, were found in luxuriant plenty, and many new kinds were discovered in their airy solitude in equal abundance. Had De Féussac, the enthusiastic admirer of this tribe, lived to see the glorious series of *Bulinii* accumulated in the Cumingian collection in different stages of growth, and in the finest state of preservation, from the egg to the adult, he would have been indeed amazed. Sorry am I that the limits of the present work will not allow me to illustrate more than two, or occasionally three, of the typical varieties of each species, and of these but a single view.

The genus *Bulinus*, restricted by Lamarck, comprehends an extremely natural group, though presenting important differences of growth and texture; and these variations are peculiarly local. In the Philippine Islands, as just described, the species are of large and rather solid growth with a remarkable hydrophanous epidermis, that is, one permeable by water or other evaporable fluid. On the barren hills of Lima and the sandy plains of Chili and Peru, they are mostly small and delicately formed; in Brazil, the species are remarkable for having the aperture in frequent instances denticulated; and in New Caledonia, Venezuela, New Grenada, and New Hebrides, they not uncommonly exhibit, with equal peculiarity, a plaited *Auricula*-like columella.

It is a curious feature in the Philippine species that the varieties of pattern, which constitute their chief ornament, reside only in the epidermis. The colours of the shell rarely describe any sort of configuration: they are mostly blended into a uniform tint, over which a fineifl pattern is produced by the epidermis forming a double porous membrane in some places, and a single one only in others, developed, moreover, with the same continuous regularity as the textile marking of a Volute or Cone. This phenomenon is easily detected by immersing the shell in water, when the light portion or upper porous

layer of epidermis becomes saturated, and the ground colour of the shell is seen through it; as the moisture evaporates, the epidermis resumes its light appearance. Sir David Brewster, in reply to a letter from Mr. Broderip on this subject, says: "It appears to me, from very careful observations, that the epidermis consists of two layers, and that it is only the upper layer which is porous wherever the pattern is white. These white or porous portions of the epidermis differ from the other parts of the upper layer only in having been deprived of, or in never having possessed, the element which gives transparency to the membrane; in the same manner as hydrophanous opal has become white, from the expulsion of its water of crystallization."

There is little variety in the animal of *Bulimus*: the Chilian species are mostly of a light colour, and a few in this and the Columbian district are spotted, some having a transparent shell through which the spots are visible. The Philippine species are, without exception, of a sombre oliveaceous brown, and dwell in family groups, as it were, among the shady foliage of the branches. Out of a group of some dozen living specimens not more than three or four may be found in an adult state with the lip of the shell reflected. They may be dislodged by shaking the branches, but are easily disturbed by the heavy rains with which those islands are at times visited. Mr. Cuming preferred, however, to collect them in dry sunny weather, because he was sure of finding the objects of his search in their shady places of retreat. In the immense sandy tract on the west side of the Andes, the reverse of this condition of nature prevails. The *Bulimai* are here physically very distinct; some reside all the year round upon the numerous *Cacti*, but during the dry season, which lasts for several months, they mostly live in a state of torpor, enclosed within their shell by an epiphram, and buried in the sand or under stones. On the approach of the showers, they revive to a state of animation, and crawl about at night in quest of food.

In illustration of the remarkable drought that prevails in Northern Chili, and of its effect upon molluscous life, I am tempted to repeat, in brief, an anecdote related to me by Mr. Cuming. On the arrival of our friend at the Port of Copiapo in 1829, he discovered the beautiful *Bulimus Broderippii* in considerable numbers, in the fissures of the rocks that may be seen here and there in the sandy plains of that country. Finding a large proportion of them dead, with the soft parts entirely decomposed, he requested a solitary inhabitant of the place to collect as many specimens as he could pick out alive whilst he occupied himself with botanizing. Returning from his

excursion, Mr. Cuming was greatly disappointed to find that among the quantity his Chilian collector had accumulated, there was scarcely one in a living state. Upon remonstrating with him for his inattention, the native replied: "Only wait till the dews come, and they will be all alive again." Mr. Cuming rejoined: "I suppose you mean when it rains." The man, however, in perfect astonishment inquired what he meant; though a sexagenarian, he had never heard of such a thing as rain.

The *Bulimai* vary in their mode of propagation: fragile species, with the lip of the shell simple, are mostly viviparous, whilst those with a reflected lip are oviparous. The arboreal species of the Philippines deposit their eggs in little clusters on the trees, between two leaves which the animal manages to curl up one upon the other, so as to form a receptacle for their protection; and, so far as Mr. Cuming's observations go, they are all soft like snake's eggs, with the single exception of the *B. Marmorosus*, in which instance the eggs are calcareous, deposited upon the leaf in parallel rows, each standing perpendicularly on end, attached at the base by a glutinous substance.

The habits of the *Bulimai* in the two widely remote countries explored by Mr. Cuming, having been treated of in the foregoing remarks, it only remains to speak of them in other parts. Turning to New Holland we are unexpectedly surprised to find that the genus is represented to an extremely limited extent. I am not aware of more than three species having been found in this wide expanse of country, although several fine *Helices* have been discovered; and in a region of which the Fauna and Flora exhibit so luxuriant and distinctive a character, the scarcity of a genus of so much importance in the Eastern Isles is remarkable. The same observation applies to New Zealand, from whence, so far as the interior of the islands of that group has been visited, no more than one or two species have been received. In Africa the *Bulimai* are almost as great strangers as in the localities just spoken of; throughout the whole extent of land yet explored of this vast continent, scarcely a dozen species have been obtained. The *Bulimai* are here replaced by *Achatinae*. Such a phenomenon may also be observed in some of the islands of the Pacific: in the Sandwich Islands the *Bulimai* are replaced by the genus *Achatinella*, and in the Society Islands their place is occupied by the *Partula*. In the West Indies the genera *Achatina* and *Glandina* seem to prevail. Howsoever abundant is the genus *Bulimus* in most of the islands of the Eastern Archipelago, few species appear to inhabit the great territories of India and China. On the coast of Borneo a beautiful species was

recently discovered by Mr. Adams, of H.M.S. Samarang, by the accidental falling of a huge tree, in a woody islet situated between Banguey and Balambangan, but they are of rare occurrence in that locality. In Europe, where nature is exposed to the vicissitudes of a colder climate, the *Bulim* are mostly small and exhibit no brilliancy of colour. So also, in the extensive region of North America, where no more than a few insignificant species are known to exist. It is in the richly fertile and woody district of Columbia, that the genus *Bulimus* is represented with a magnificence little inferior to that of the Philippine Islands; here they are large enough and sufficiently abundant to be roasted and eaten by the aborigines as a frequent article of food. Several fine species, entirely new to science, have been collected in Venezuela and New Granada by Mr. Linden, an assiduous Botanical Traveller, only within the last twelvemonth, at an altitude of from 5000 to 8000 feet, and many more, no doubt, dwell in undisturbed solitude in the vast interior of this immense continent. It is extremely probable that a large portion of South America yet remains to be explored by the adventurous naturalist, where there is no doubt a fine expanse of forest country, grand in extent, rich in foliage, and possessing all the elements favourable to the growth and beauty of arboreal mollusks.

Species 1. (Mus. Cuming.)

BULIMUS MACROSTOMA. *Bul. testá ovali, ventricosâ, spirâ subobtusevata, anfractibus seis, convexo-tumidis, concentricâ striatis, columellâ basi subcontortâ; purpureofuscâ, epidermide hydroporâ, in anfractu ultimo unzonatâ, indutâ; -aperturâ cærulescente-albâ, lobro reflexo.*

THE WIDE-MOUTH BULIMUS. Shell oval, ventricose, spine slightly abbreviated, whorls six in number, convexly tumid, concentrically striated, columella slightly twisted at the base; purple-brown, covered with a hydrophanous epidermis, marked with a single zone on the last whorl; aperture bluish white, lip reflected.

PFEIFFER, Pro. Zool. Soc., 1842, p. 152.

Hub. Sual, Province of Pangasinan, Island of Luzon, Philippines (on leaves of trees); Cuming.

This species approaches the *B. rufogaster*, it partakes of the same shades of colour, and the epidermis is characterised by a similar zone round the last whorl; it does not however exhibit the vacant patches beneath the sutures. The columella of *B. macrostoma* is slightly twisted, and

the general form of the shell is eminently distinguished by its swollen egg-shaped growth.

Species 2. (Fig. a and b. Mus. Cuming.)

BULIMUS PYTHOGASTER. *Bul. testâ subpyramidalis-ovata, anfractibus seis, plano-concisis, ultimo ventricoso, columellâ rectâ; purpureo-nigricante, epidermide interdum simplici, interdum longitudinaliter strigata, aperturâ cærulescente-albâ, columellâ rosaceo-purpurascente.*

THE ORIENTAL BULIMUS. Shell somewhat pyramidal ovate, whorls six in number, flatly convex, the last ventricose, columella straight; purple-black, epidermis sometimes simple, sometimes arranged in longitudinal streaks, aperture bluish white, columella pinkish purple.

FERUSSAC, Lamarcq., Anim. sans vert. vol. viii. p. 226.

Hub. Fig. a, Island of Ticao; Fig. b, Mount Isarog, Island of Luzon, Philippines (on leaves of trees); Cuming.

The examples of *B. pythogaster* selected for representation vary so materially in form, as well as in the arrangement of the epidermis, that one might consider them to belong to distinct species, were it not for the similarity in the structure and colouring of the aperture. In the specimen from Ticao, Fig. a, the shell is broader and more acutely ventricose at the base, and the epidermis is arranged in longitudinal streaks; in that from Luzon, Fig. b, the base is somewhat contracted, imparting a more pyramidal form to the shell, the apex is dark purple, and the epidermis is soft and velvety, uniformly distributed throughout without any description of pattern.

Species 3. (Mus. Cuming.)

BULIMUS LIGNARIUS. *Bul. testâ ovo-conoidâ, ventricosissimâ, anfractibus seis, tenuido-converris; columellâ rectâ; purpureo-nigricante, epidermide lineis fasciisque in anfractu ultimo, et interdum penultimo, notatâ; aperturâ cærulescente-albâ, columellâ livido-parparascente.*

THE WOODEN BULIMUS. Shell ovately conoid, very ventricose, whorls six in number, timidly convex; columella straight; purple-black, epidermis marked with bands or lines on the last and sometimes the penultimate whorl; aperture bluish-white, columella livid purple.

PFEIFFER, Pro. Zool. Soc., 1842.

Hub. Gattaranq, Province of Cagayan, Island of Luzon, Philippines (on leaves of trees); Cuming.

BULIMUS.—PLATE I.

This is the largest of the Philippine species, and that of the most ventricose proportions. The columella is of a livid blueish purple, and the epidermal pattern consists of a number of circular bands or lines, frequently extending around the last and penultimate whorls; the specimen figured, in which the bands are confined to the last whorl, is rather an exceptional variety.

Species 4. (Mus. Cuming.)

BULIMUS RUGOGASTER. *Bul. testā ovato-conoidā, sub-ventricosā, anfractibus senis, convexis, columellā sub-intortā; nigricante-purpureā, interdum rufescente, epidermide basin versus unicolor, strigis longitudinalibus nigricantibus, prope ad suturas latioribus, notata, aperturā columellā carunculata-albitā.*

THE RED BULIMUS. Shell ovately conoid, somewhat ventricose, whorls six in number, convex, columella somewhat twisted; blackish purple, sometimes reddish, epidermis marked towards the base with a single band, and with dark longitudinal streaks, which are broader near the sutures, aperture and columella bluish white.

Hab. Province of Baie, Island of Luzon (in dark and lofty forests); Cuming.

In this species the columella is slightly twisted, and of the same blueish white hue as the interior of the shell. The pattern of the epidermis exhibits a single band round the lower part of the last whorl, and a row of broad, short, interrupted streaks around the sutures. This last mentioned design I have not observed in any other species.

Species 5. (Mus. Cuming.)

BULIMUS PHILIPPINENSIS. *Bul. testā ovato-conoidā, basi subplanatā, anfractibus senis, convexis, ultimo basin versus subobscurè angulato; columellā leviter intortā; nigricante-fusca, epidermide in strigis obliquè concentricis, aperturā carunculata-albitā, columellā palea rosaceo-purpurascente.*

THE PHILIPPINE BULIMUS. Shell ovately conoid, somewhat flattened at the base, whorls six in number,

convex, last whorl somewhat obscurely angled towards the base; columella slightly twisted; blackish-brown, epidermis disposed in obliquely concentric streaks, aperture blueish white, columella pale pinkish purple.

PFEIFFER, Pro. Zool. Soc., 1846.

Hab. Islands of Luzon and Marinduque, Philippines (on the leaves of trees); Cuming.

There is little to distinguish this species from the *B. pythogaster*; it has a somewhat angular inclination of growth around the lower part, but this is not a character to which much importance can be attached, seeing that the typical varieties above figured of that species present almost as great a difference of form.

Species 6. (Mus. Cuming.)

BULIMUS REEVII. *Bul. testā oblongo-ovalā, anfractibus senis, tumidusculis, ultima ventricosa, effuso; intensè ustulato-fusca, epidermide in fasciis insignibus latusculis renatis lineisque subtilibus, aperturā carunculata-albitā, columellā viz intortā, peritremate livido-purpurascens.*

REEVE'S BULIMUS. Shell oblong-ovate, whorls six in number, rather swollen, last whorl ventricose and effused; rich burnt-brown, with the epidermis in remarkable rather broad bands and fine lines, aperture bluish white, columella scarcely twisted, and, together with the peristome, of a livid purple hue.

CUMING MSS., Mus. Cuming.

Helix Reevi, Broderip, Pro. Zool. Soc., 1841.

Hab. Lubban, Province of Tayabas, Island of Luzon, Philippines (on the leaves of trees); Cuming.

This is the first species of *Bulimus*, and I believe the first shell, which Mr. Cuming unpacked from his vast collection on his return from the Philippines, and, being present at this exciting moment, I shall not easily forget the impression caused by the sight of so superb an example of an arboreal Snail. The remarkable disposition of the epidermis was quite a matter of astonishment, nothing of the kind had been observed before, nor had any specimen of the *B. pythogaster* been collected in a state of preservation by which this phenomenon could be understood.



BULIMUS.

PLATE II.

Species 7. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS GILVUS. *Bul. testā subglobosā, apice obtusā, angustiōnis senis, rotundatā, concentricē striatis, apertura subcontractā, columellā vix curvatā; gileā vel ustulatōrū, aufractū ultimo obscurē nigrificati, epidermide simplici indutā, peritreme colunselldāque livido-albā.*

THE BRICK-COLOURED BULIMUS. Shell somewhat globose, obtuse at the apex, whorls six in number, rounded, concentrically striated, aperture rather contracted, columella slightly curved; brick-coloured or burnt-red, last whorl encircled with an obscure band, covered with a simple epidermis, peritreme and columella livid-white.

SOWERBY, Pro. Zool. Soc., 1845, p. 45.

Hab. Island of Bohol, Philippines (on the leaves of trees); Cuming.

The specimens of *B. gilvus* figured give a tolerable idea of the varieties of the species, which only affect the shell so far as the depth of colour and thickness of the epidermis are concerned. The last whorl exhibits an obscure band round the lower part, and the concentric striae are rather prominently developed, but there is no epidermal pattern or configuration. The livid-white peritreme and columella are rather a characteristic feature in the species. It appears to be confined to the Island of Bohol.

The *B. gilvus*, it will be observed, partakes much more of the character of *Helix* than the following, which Mr. Broderip refers to that genus according to the method of De Féussac.

Species 8. (Fig. *a*, *b*, and *c*. Mus. Cuming.)

BULIMUS DAPHNIS. *Bul. testā pyramidalē ovatā, aufractū senis, convexis, plus siuis ventricosis, apertura subcontractā, columellā subrectā; nigricante-purpureā, spadicēaque, epidermide vel simplici, vel in striis longitudinalibus, vel in fasciis lineisce transversis; columellā roseo-purpureā, apertura cārnelente-albā, peritreme nigricante-purpureā.*

THE DAPHNIS BULIMUS. Shell pyramidal, ovate, whorls six in number, convex, more or less ventricose, aperture somewhat contracted, columella nearly straight; blackish purple or fawn colour, epidermis either simple or disposed in longitudinal streaks or transverse bands and lines, columella pinkish purple, aperture blueish white, peritreme blackish purple.

Helix (Cochlostyla) Daphnis, Broderip, Pro. Zool. Soc., 1840, p. 180.

Hab. Fig. *a*, Island of Siquijor; Fig. *b*, Argao, Island of Zebu; Fig. *c*, San Nicolas, Island of Zebu, Philippines (on the trees); Cuming.

Of the varieties of this charming species it may be said their name is legion, all remarkable for their neatness and alike worthy to sustain the rural dignity of the pastoral shepherd and poet whose name they bear. Of the specimens selected for representation, one has the epidermis disposed in transverse bands, another in concentric streaks, whilst in the third it is without any indication of pattern; and it may be observed that they vary even in form. In the most important part of the shell, the aperture and columella, they resemble each other perfectly, and not the least permanent character of the species resides in the pinky colouring of those parts.

Species 9. (Mus. Cuming.)

BULIMUS SOLIDUS. *Bul. testā oblongā, subelongatā, angustiōnis septensis, ultimo basia versis obsoletissimē angulatā; columellā subintortā; nigricante-purpureā, epidermide in striis longitudinalibus, confertis, irregulāribus, apertura cārnelente, peritreme nigricante-purpurea.*

THE SOLID BULIMUS. Shell oblong, somewhat elongated, whorls seven in number, the last very obsoletely angled towards the base; columella slightly twisted; blackish-purple, epidermis in close-set irregular longitudinal streaks, aperture bluish, peritreme blackish-purple.

PEIFFER, Pro. Zool. Soc., 1842, p. 152.

Hab. San Juan, Province of Cagayan, Island of Luzon, Philippines (on leaves of trees); Cuming.

This species, of which Mr. Cuming collected only a few specimens, is of an elongately oblong form, and composed of one more whorl than any of the preceding; it is of rather dark colour, and the longitudinal streaks of epidermis are very irregular and close-set. There is a variety in which the last whorl is more inflated and has a whitish band.

The name devoted to this species is rather an inappropriate one, the shell being of no more solid growth than its congeners.

BULIMUS.—PLATE II.

Species 10. (Mus. Cuming.)

BULIMUS MUS. *Bul. testā ovatā, crassiusculā, spirā obtusa, anfractibus quinīs, subrotundetis, columellā sub-intortā; intensē castaneā, epidermide albicant, lineis subtilibus concentricis hic ilic' interruptō; columellā lacteā, peritreme intensē castaneā.*

THE MOUSE BULIMUS. Shell ovate, rather thick, spine obtuse, whorls five in number, somewhat rounded, columnella slightly twisted; deep chesnut, epidermis whitish, interrupted here and there with fine lines, columnella milk-white, peritreme deep chesnut.

Hab. Bamang, Province of Pangasinan, Island of Luzon (on trees); Cuming.

The preceding species, it may be observed, consists of a whorl more than its congeners from the Philippine Islands: this, on the contrary, is shorter and composed of a whorl less. It is of an extremely rich chesnut colour, covered by an epidermis delicately white, and only interrupted here and there by fine lines. The columnella does not partake of the colour of the peritreme, being of a clear milk-white.



BULIMUS.

PLATE III.

Species 11. (Fig. a and b. Mus. Cuming.)

BULIMUS WOODIANUS. *Bul. testā ovato-conoidā, basi peculiariter concavā, anfractibus quinque subrotundis, suturis impressis, columellā vix intortā; albā, epidermide latē tenuissimā nitidā indutā, anfractus ultimo ustulato-fusco tincto, conspicue unifasciati, columellā rosaceo-purpurascente, peritreme et circa columellanam intusē nigricante-castaneis.*

WOOD'S BULIMUS. Shell ovately conoid, peculiarly concave at the base, whorls five in number, somewhat round, sutures impressed, columella slightly twisted; white, covered with a very thin yellow shining epidermis, last whorl stained with burnt brown and conspicuously single banded, columella pinkish purple, peritreme and around the columella deep blackish chestnut.

LEA. Trans. Acad. Nat. Sc. Philadelphia, 1840, pl. xi. f. 5.
Hab. Island of Romblon, Philippines (on the trees); Cuming.

This species is of a character singularly distinct from the rest of its congeners of the Philippine Isles. No pattern is discernible in the epidermis, which is merely a thin transparent horny layer, nor is there any colour in the shell until it approaches maturity. The specimen represented at Fig. 11. a is entirely colourless; that at 11. b has a burnt brown hue upon the last whorl, which is further distinguished by a dark band and intensely dark chestnut-stained peritreme.

Species 12. (Mus. Cuming.)

BULIMUS JUGLANS. *Bul. testā ovatā, crassissimulā, anfractibus quinque, rotundato-tumidis, obliquè striatis, operculari subcontractā, columellā intorsum arcuatā; intensē rufo-castaneā, epidermide tenuiculā, spiraliiter lineatā, aperturā peritreme et columellā livido-cerulescentibus.*

THE WALNUT BULIMUS. Shell ovate, rather thick, whorls five in number, rotundately tumid, obliquely striated, aperture slightly contracted, columella curved inwards; deep reddish chestnut, epidermis rather thin, spirally lined, aperture peritreme and columella pale livid blue.

PFEIFFER, Pro. Zool. Soc., 1842, p. 89.

Hab. Mountains of Igarrotes, northern extremity of the Island of Luzon, Philippines (on leaves of trees); Cuming.

A rather scarce species of solid growth, and of a deep crimson brown colour. The aperture is somewhat contracted, and there is a peculiar twist in the columella. The epidermis is rather slight, disposed in spiral lines.

Species 13. (Mus. Cuming.)

BULIMUS FAUNUS. *Bul. testā subpyramidalis-ovatā, crassissimulā, anfractibus sexis, subconvergis, columellā circūlectā; castaneo-fusca, epidermide simplici, fusco-fusca, maculis paucis nigricantibus prope suturas exhibentibus, aperturā cæruleo-roseata-albā, columellā rosaceo-purpurascente, peritreme nigricante.*

THE FAWN BULIMUS. Shell somewhat pyramidal, ovate, rather thick, whorls six in number, slightly convex, columella nearly straight; chestnut-brown, epidermis simple, smoky-brown, exhibiting a few blackish spots next the sutures, aperture bluish-white, columella pinkish-purple, peritreme blackish.

Helix (Cochlostyla) Faunus, Broderip, Pro. Zool. Soc., 1840, p. 180.

Hab. San Nicolas, Island of Zebu (on the trees); Cuming.

Of a firm pyramidal growth, singularly characterized by a row of irregular small spots around the sutures.

Species 14. (Fig. a, b, c. Mus. Cuming.)

BULIMUS TICAONICUS. *Bul. testā subgloboso-ovatā, apice obtusa, anfractibus quinque, rotundatis, columellā subarcuatā; castaneo-rufa ant floridante, epidermide in floccis strigisve subdistansibus conspicua, ant simplici ant spiraliiter lineatā, columellā rosaceo-purpurascente, peritreme purpureo-castaneā, nonnigra pallida.*

THE TICAO BULIMUS. Shell somewhat globosely ovate, obtuse at the apex, whorls five in number, rounded, columella slightly arched; chestnut-red or pale yellow, with the epidermis in conspicuous, rather distinct flakes or streaks, or simple or spirally lined, columella pinkish-purple, peritreme purple-chestnut, sometimes pale.

Helix (Cochlostyla) Ticaonica, Broderip, Pro. Zool. Soc. 1840.

Hab. Island of Ticão, Philippines (in the woods); Cuming.

This species, like the *B. Daphnis* of the Island of Zebu, is an extremely variable one, assuming numerous styles of pattern and colour, as may be noticed by the contrast afforded in the specimens selected for representation. All are, however, distinguished by their globose form, and by the important delicate pink colouring of the columella.



BULIMUS.

PLATE IV.

Species 15. (Mus. Cuming.)

BULIMUS MINDORENSIS. *Bul. testā cylindraceo-oblongā, anfractibus sexis, convexis, columellā subarcuatā; fasciū nigricante ant. rufescente, epidermide in strigis conspicuis subdistans, aperturā cārulēscētē-albā, peritreme effusā, intēsē fuscō-nigricante.*

THE MINDORO BULIMUS. Shell cylindrically oblong, whorls six in number, convex, columella slightly arched; brownish black or reddish, with the epidermis in conspicuous, rather distant, longitudinal streaks, aperture bluish white, peritreme effused, deeply stained with brown-black.

Hab. Broderip, Pro. Zool. Soc., 1840, p. 84.

Hab. Puerto Galero, Philippine Islands (on the leaves of trees); Cuming.

The epidermal pattern of this species is characteristic, consisting of numerous conspicuously marked longitudinal streaks, which in parts are somewhat distant from each other. The aperture in most of the species is comparatively rather small, and the lip in all is effused rather than reflected back.

I cannot quite agree in the varieties noted by Mr. Broderip, some of which are too nearly alike to make it worth while to mark a distinction, whilst there is one which appears to me of so different a character, that it should be regarded as another species, see Plate V. Sp. 20.

Species 16. (Fig. a and b. Mus. Cuming.)

BULIMUS CHRYSALIDIFORMIS. *Bul. testā cylindraceo-oblongā, medio ventricosiore, anfractibus septenis, sub-plano-convexis, columellā subincurvā, aperturā parencyclā; intēsē rufō-castaneā, epidermide in fasciis latissimis, longitudinalibus, undatis, vel flavescente, epidermide sub fasciā rufō-fuscā infra suturas eximie tessellatā, peritremate intēsē castaneā.*

THE CHRYSALIS-SHAPED BULIMUS. Shell cylindrically oblong, more ventricose in the middle, whorls seven in number, rather flatly convex, columella slightly twisted inwards, aperture somewhat small; deep reddish chestnut colour, with the epidermis in rather broad longitudinal wave streaks, or yellowish, with the epidermis delicately tessellated upon a reddish brown band beneath the sutures, peritreme deep chestnut brown.

SOWERBY. Pro. Zool. Soc., 1833, f. 37. Conch. Illus.

Bulimus, f. 28.

Hab. Fig. 16 a, Mansala, Island of Mindoro. Fig. 16 b,

Puerto Galero, Philippines (in dark forests of thick foliage, upon and beneath the leaves); Cuming.

The *Bulimus chrysalidiformis* described by Mr. Sowerby, prior to Mr. Cuming's voyage to the Philippine Islands, from a pale uncoated specimen, is mainly distinguished by its elongately cylindrical form, the aperture being comparatively rather small, and a little distorted from the twisting of the columella. The yellow variety represented at Fig. 16 a, is very remarkable, and appears, at first sight, to differ materially from the type of the species, which approaches rather to *B. Mindoreensis*.

Species 17. (Mus. Cuming.)

BULIMUS NIMBUS. *Bul. testā pyramidalē-oblongā, spirō acuminatā, anfractibus septenis, convexis, columellā arcuatā; intēsē brunnēā, epidermide in floccis latissimis, subdistans, undatis, aperturā cārulēscētē-albā, peritremate brunnēa.*

THE RAINY BULIMUS. Shell pyramidal oblong, spire acuminated, whorls seven in number, convex, columella arched; deep brown, with the epidermis in broad, rather distant, wavy flakes, aperture bluish white, peritreme brown.

Hab. Broderip, Pro. Zool. Soc., 1840, p. 121.

Hab. Island of Negros (on the summits of the mountains); Cuming.

A fine dark brown species of a pyramidal acuminate form, with the epidermis disposed in light longitudinally waved flakes. Mr. Broderip's variety d has always appeared to me so constantly distinct from this, that I propose to elevate it to the rank of a species, in compliment to my friend, Dr. Pfeiffer.

Species 18. (Mus. Cuming.)

BULIMUS PFEIFFERIANUS. *Bul. testā oblongo-ovatā, sub-ventricosā, anfractibus sexis, tumidinscis, columellā subarcuatā; flavo-viridescente, epidermide in strigis longitudinalibus undatis, columellā et aperturā cārulēscētē-albā, peritremate et parte circa columellam, nigricante-fuscis.*

PFEIFFER'S BULIMUS. Shell oblong-ovate, rather ventricose, whorls six in number, somewhat swollen, columella slightly arched; pale yellowish green, with the epidermis in longitudinally waved streaks, columella and aperture bluish white, peritreme and the part round the columella blackish brown.

Bulimus sinuosus, var. *d*, Broderip.

Hab. Tankay, Island of Negros (on the branches); Cuming.

I have long regarded this shell in my own mind as a species distinct from *B. sinuosus*; it is composed of one whorl less, is constantly more ventricose, less pyramidal, acuminate, and always distinguished by a pale yellowish-green cast of colour with a dark chestnut blotch, nearly black, around the columella, and there is a peculiarity also in the arrangement of the epidermis.

To Dr. Pfeiffer, of Cassel, I dedicate this beautiful species, trusting he will appreciate my desire to do him honour for the service he has rendered to conchological science by his laborious investigation of the great tribe of *Helicidae* of which the present genus forms a part.

Species 19. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS PICTOR. *Bul. testa acuminato-oblonga, anfractuosa*.

tibus senis, rotundatis, columellâ subrectâ; intus brunnescens, epidermae plus minus abâ, fasciis angustis longitudinalibus undulatis, interdum interruptis, prætermitente, aperturâ caroalente, peritreme nigricante-castaneo marginatâ.

THE PAINTER BULIMUS. Shell acuminate oblong, whorls six in number, convex, columella nearly straight: rich brown, epidermis more or less white, leaving narrow longitudinally waved dark bands, which are sometimes interrupted, aperture bluish white, peritreme edged with blackish chestnut.

BRODERIP, Pro. Zool. Soc., 1849, p. 120.

Hab. Dingle, Province of Ilo Ilo, Island of Penang, Philippines (on the trees); Cuming.

A richly coloured species, with the epidermis so deposited as to leave a dark, longitudinal, zig-zag stripe at intervals, which is occasionally broken or otherwise interrupted, and has a striking appearance from the epidermis being of a thick, opaque-white, satiny hue down one side. The aperture is of clear milky white, and the peritreme is peculiar in having the usual dark colouring matter quite at the edge.



20. b.



20. a.



21.



22.



23. a.



23. c.



23. b.

BULIMUS.

PLATE V.

Species 20. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS NOBILIS. *Bal. testá acuminato-oblongá, anfractibus quinque seweise, subplano-concexis, suturis impressis, columellá rectilincolá; virescote-flavida, epidermide tenui, zonulas albas sinuatas, longitudinalis, nunc distantes, nunc interruptas, exhibente, anfracta ultimo fascia angustâ nigricante-fusca parte inferiore cingulato, circa columellam nigrante, peritreme albâ.*

THE NOBLE BULIMUS. Shell acuminate oblong, whorls five or six in number, somewhat flatly convex, sutures impressed, columella rather straight; greenish yellow; epidermis thin, exhibiting longitudinally situated opaque white zones, sometimes interrupted, sometimes rather distant, last whorl with a narrow dark brown band round the lower part, blackish around the columella, peritreme white.

Hab. Mountains of Igbaras, Island of Luzon, Philippines (on the trees); Cuming.

The markings of this beautiful species are similar to those of the *B. fulgetrum*, the epidermis being thin, excepting where it is disposed in striking lightning-like streaks, darting, so to speak, out of the sutures; in form the shell is constantly distinct, and it is further characterized by a conspicuous dark band.

Species 21. (Mus. Cuming.)

BULIMUS ELECTRICUS. *Bal. testá subelongato-ovalá, medio ventricosore, spirá acuminatá, anfractibus senis, subplano-concexis, columellá leviter intortá; lividofusca, epidermide tenuicula, strias pallidus obliquè undatus, valde irregulares, exhibente, aperturá albida, peritreme rufo-fusca.*

THE ELECTRIC BULIMUS. Shell somewhat elongately ovate, more ventricose in the middle, spire acuminate, whorls six in number, rather flatly convex, columella slightly twisted; livid red, epidermis rather thin, exhibiting very irregular obliquely waved pale streaks, aperture whitish, peritreme reddish brown.

Bulimus Mindoreensis, var. *k*, Broderip.

Hab. Puerto Galero, Island of Mindoro, Philippines (on the leaves of trees); Cuming.

If any importance is to be attached to the character and pattern of the epidermis of these shells, there is certainly enough to distinguish this from the *B. Mindoreensis*.

The epidermis is of a soft, very slight texture, of a uniform reddish brown tint in which light streaks descend here and there from the sutures in very zigzag course, somewhat as in *B. fulgetrum* but fainter; and, besides this, the shell is of a more acuminate growth, whilst the last whorl is larger and more effused.

Species 22. (Mus. Cuming.)

BULIMUS BICOLORATUS. *Bal. testá pyramidali, anfractibus septenis, leviter convezis, columellá subintortá; rufo-brunneá, subbus viridi, epidermide in striis longitudinalibus medio peculiariter interruptis, et inferne bifasciatá, peritreme columelláque intense nigro-fusca.*

THE TWO-COLOURED BULIMUS. Shell pyramidal, whorls seven in number, slightly convex, columella slightly twisted; reddish-brown, green beneath, epidermis in rather broad longitudinal bands, peculiarly interrupted in the middle, and two-banded round the lower part, peritreme and columella deep black-brown.

LEA, Trans. Amer. Phil. Soc., vol. vii. p. 459, pl. 11. f. 8.
Bulimus Alberti, Broderip.

Hab. Mount Isarog, Province of Camarinas; and Calanang, Province of Baie, Island of Luzon, Philippines (on the trees); Cuming.

A truly beautiful species, remarkable for its pyramidal growth, and original display of colouring. None of the preceding species have any indication of the bright green which distinguishes the under portion of the body-whorl in this, and the epidermal pattern is equally peculiar.

The shell described by Mr. Broderip under the name of *B. oxyx* appears to be only another state of this, inhabiting the same locality in more lofty situations in the trees. It should be observed that Mr. Cuming collected but two specimens, one in bad condition, the other of immature growth. They are of a singular pupiform shape, but are so nearly allied to the *B. bicoloratus* in other respects, that the species must be received with caution until further and better means of comparison can be obtained.

The honour intended by the dedication of this finest of the Philippine species, to be conferred on H.R.H. Prince Albert, whose interest in Conchology is handsomely testified by his patronage of the present work, was uncon-

sciently intercepted in the present instance by Mr. Lea of Philadelphia.

Species 23 (Fig. *a, b*, and *c*. Mus. Cuming.)

BULIMUS FULGETRUM. *Bal. testa oblongo-ovalis, subcuneatis, anfractibus quinque, ultimo majusculo, columellam ex arcuata; ustulato-fusca et flavicante, epidermide in strigis albis oparis rariè undatis, subdistans, peritremate albâ.*

THE LIGHTNING-MARKED BULIMUS. Shell oblong-ovate, somewhat ventricose, whorls five in number, last whorl rather large, columella slightly curved; burnt-

brown or yellowish, with the epidermis in rather distant variously waved opaque white streaks, peri-treme white.

BRODERIP, Pro. Zool. Soc., 1840., p. 119.

Hab. Island of Negros, Philippines (on the trees); Cuming.

Of all the species collected by Mr. Cuming at the Philippine Islands, this has, perhaps, most excited the admiration of amateurs. The electrical flashes, so to speak, of light satiny epidermis which descend from the sutures are peculiarly striking, and unlike anything that had been seen before. The ground colour of the shell varies from a rich burnt-brown to a pale lemon colour.



BULIMUS.

PLATE VI.

Species 24. (Fig. a and b. Mus. Cuming.)

BULIMUS GUIMARASENSIS. *Bul. testā subacuminato-oblongā, anfractibus quinīs, concavis; albā, glabrā, anfractibus fasciā fasciā conspicuā supra infringe cingulatis, epidermide tenui nitidā latè olivaceā indutā.*

THE GUIMARAS BULIMUS. Shell somewhat acuminate oblong, whorls five in number, convex; white, smooth, whorls encircled above and below by a conspicuous brown band, and covered by a thin shining yellowish olive epidermis.

PRODRERIP, Pro. Zool. Soc., 1840, p. 156.

Hab. Island of Guimaras, Philippines (on the trees); Cuming.

A well-defined though not a particularly striking shell, partaking of the character of *B. ovoides* and *citrinus*. Fig. 24 a is the ordinary state of the species; the variety represented at Fig. 24 b is one of rare occurrence, in which the bands are unusually broad and dark coloured.

Species 25. (Mus. Cuming.)

BULIMUS FRATER. *Bul. testā subacuminato-oblongā, anfractibus quinīs, concavis; rosaceo-albā, anfractum parte inferiori zonā angustā refescente cingulatā, epidermide tenui nitidā rufescere-testā indutā.*

THE BROTHER BULIMUS. Shell somewhat acuminate oblong, whorls five in number, convex; pinkish-white, lower part of the whorls encircled by a narrow reddish zone, covered by a thin shining reddish-yellow epidermis.

DE FÉRUSSAC, Hist. Nat. Moll. terr. et fluv. pl. 112, f. 3 and 4.

Hab. Island of Leyte, Philippines (on the leaves of trees); Cuming.

Dr. Pfeiffer is of opinion that this shell is the *B. frater* of De Féruccac, which is highly probable, although his figure of that species represents one of larger dimensions and devoid of the delicate rose colouring.

Species 26. (Fig. a and b. Mus. Cuming.)

BULIMUS MACULIFERUS. *Bul. testā acuminato-oblongā, plerumque sinistrorsā, anfractibus senis, concavis, concentricè striatis, ultimo longissimo; flavescente-spatulatā, interdum brunnecō maculatā et nebulatā, interdum*

immaculatā, lineā roseā longitudinali hic illuc ornatā sutureis albidis, periremate albā, apice nitidè opacobrunnea.

THE BLOTTCHED BULIMUS. Shell acuminate oblong, mostly sinistral, whorls six in number, concentrically striated, last whorl rather long; yellowish fawn colour, sometimes blotched and clouded with brown, sometimes unspotted, ornamented here and there with a pink longitudinal line, sutures whitish, periremate white, apex shining opaque white.

PRODRERIP, Pro. Zool. Soc., 1841, p. 14.

Hab. Province of Misamis, Philippines (on the trunks of trees); Cuming.

Dr. Pfeiffer has intimated his opinion that the *B. maculiferus* is a variety of the *B. citrinus*. I cannot, however, agree with him in this; the species are very nearly allied but always distinct. The present is of a more elongated growth, and in all its varieties is characterized by the same pale ferruginous fawn colouring. The variety represented at Fig. 26 b is prettily distinguished by three or four narrow longitudinal zones of a delicate rose colour.

Species 27. (Fig. a and b. Mus. Cuming.)

BULIMUS SMARAGDINUS. *Bul. testā oblongo-ovatā, plus minus ventricosā, anfractibus concavis, subdissimilē obliquè striatis, suturis superficialiter canaliculatis; albā, epidermide tenui cornē nitidā infra saturos indutā; in anfractu ultimo smaragdinā vel olivaceo-virescere, purpureo apicem versus evanescente; periremate interdum albā, interdum pallide rosacea.*

THE EMERALD-GREEN BULIMUS. Shell oblong ovate, more or less ventricose, whorls convex, very finely obliquely striated, sutures superficially channeled, white, covered beneath the sutures with a thin shining horny epidermis; bright emerald-green or olive-green on the last whorl, fading into purple towards the apex, perireme sometimes white, sometimes light pink.

REEVE, Pro. Zool. Soc., 1842, p. 49. Conch. Syst. vol. ii. pl. 173, f. 6.

Hab. Surigao, Island of Mindanao, Philippines (on the leaves of trees); Cuming.

The accompanying figures of this bright coloured and delicate species represent extreme states of it, one being much more ventricose and of a lighter colour than the

other. The chief peculiarity of the *B. suaragininus* consists in the nature of its epidermis, which, unlike that in most of the preceding Philippine species, exhibits no pattern, but contains all the colouring matter. The shell is pure white, and the beautiful emerald-green colour, so much to be admired, resides within the epidermis, which is of a thin shining horny character, commencing in a singular manner just beneath the sutures, and may be peeled away.

Species 28. (Mus. Cuming.)

BULIMUS CINCINNIFORMIS. *Bul. testā subpyramidalis oratā, tenuiculā, atractibus senis, plano-convexis; albā, atractibus fasciis duabus distantibus, nigricanib; epidermīs particulis notatis cingulatis, labro tenui, vix reflexo.*

THE CURL-SHAPED BULIMUS. Shell somewhat pyramidal, ovate, rather thin, whorls six in number, flatly convex: white, whorls encircled with two distant black bands marked with particles of epidermis, lip thin, scarcely reflected.

Hab. Island of Luban, Philippines (on the trees); Cuming.

A thin pyramidal shell, in which the black bands are curiously studded with light particles of epidermis.

Species 29. (Mus. Cuming.)

BULIMUS SATYRUS. *Bul. testā subpupiformi, atractibus quinque, convexis, columellā subintortā; purpureo-nigrā, epidermide fasciā simplici iudā, aperturā crenulente-albā, columellā carneo-purpurascente, peristreme purpleo-nigrā.*

THE SATYR BULIMUS. Shell somewhat pupiform, whorls five in number, convex, columella slightly twisted; purple-black, covered with a simple brown epidermis, aperture bluish white, columella light flesh-purple, peristreme purple-black.

BRODERIP, PRO. ZOOL. SOC., 1842. p. 181.

Hab. Island of Tablas, Philippines (on the leaves of trees); Cuming.

Mr. Cuming only obtained a few specimens of this species; it is of a dark purple-black hue without any epidermal pattern to attract particular attention.



30 a



31



30 b



30 c



32



30 d



33



34



35

BULIMUS.

PLATE VII.

Species 30. (Fig. *a*, *b*, *c*, and *d*. Mus Cuming.)

BULIMUS CINCINNUS. *Bul. testá ovato-pyramidalis, tenui,
subhyalino, anfractibus sexis subrotundatis, columellá
subintortá, labro simplici, vix reflexo; pallide rosacea;
albidus aut virescente, epidermide fuscá basin versus
sparsim maculat.*

THE CURL BULIMUS. Shell ovately pyramidal, thin, rather transparent, whorls six in number, somewhat rounded, columella slightly twisted, lip simple, scarcely reflected; pale pink, whitish or greenish, sparingly bespotted towards the base with dark brown epidermis.

SOWERBY, Pro. Zool. Soc., 1840, p. 98.

Hab. Islands of Temple and Burias, Philippines (on the leaves of trees); Cuming.

A very delicate and extremely variable species, the epidermis of which is of so tenacious a character that it is rare to meet with specimens having any portion adhering. The colour of some is purple, of others light pink or pale green, and the shell has not unfrequently a pink columella.

Species 31. (Mus. Cuming.)

BULIMUS CALYPSO. *Bul. testá superpyramidalis-ovatá, dia-
phaná, anfractibus sexis, convexis; columellá pecu-
liariter contortá et truncatá, labro tenui, vix reflexo;
albidus, fasciis pallidis oliveaceo-luteis, precipue in-
fractu ultimo, cingulatá; columellá apice et peritre-
mata roseo-purpureis.*

THE CALYPSO BULIMUS. Shell somewhat pyramidal, ovate, diaphanous, whorls six in number, convex; columella peculiarly twisted and truncated, lip thin, but little reflected; whitish, encircled with pale olive yellow bands, especially round the last whorl, columella apex and peristome rose-purple.

BRODERIP, Pro. Zool. Soc., 1840, p. 182.

Hab. Tanhay, Isle of Negros, Philippines (in the woods); Cuming.

Only two specimens of this very delicate diaphanous species were collected; it is encircled by numerous faint olive-yellow bands, somewhat crowded towards the base, and the columella, peristome, and first two whorls of the shell are of a deep rose-purple; there is, moreover, a dark purplish stripe around the columella.

It is a peculiarity in this species that the banded pattern does not reside in the epidermis, as in most of its congeners, but in the shell, where it may be observed winding round the interior.

Species 32. (Mus. Cuming.)

BULIMUS VIRENS. *Bul. testá pyramidalis-ovatá, laeviusculá,
nitidá, anfractibus septenis, subdolano-converxis, ultimo
infractu obscuré angulato; columellá vix intortá, labro
subreflexo; albá, epidermide tenui virescente basin versus
indulat.*

THE PALE GREEN BULIMUS. Shell pyramidal, rather smooth, shining, whorls seven in number, rather flatly convex, the last obscurely angled round the lower part; columella slightly twisted, lip but little reflected; white, covered with a thin epidermis which is of a pale green colour towards the base.

PFEIFFER, Pro. Zool. Soc., 1840, p. 153.

Hab. Island of Burias, Philippines (on leaves of trees); Cuming.

I cannot trace the resemblance which Dr. Pfeiffer notices between this shell and *B. carinatus*, Lea; the figure of that species in the Transactions of the Academy of Natural Sciences of Philadelphia, represents a shell belonging rather to that division of the genus which is characterised by a hydrophobic epidermis.

Species 33. (Mus. Cuming.)

BULIMUS HINDSI. *Bul. testá oblongo-ovatá, anfractibus
quini converxis, columellá subintortá, labro reflexo;
albá, anfractibus fasciá castaneá supra et infra cingula-
tis, epidermide tenui luteá, parte circa columellam
et peristremate purpureo-fasciis*

HIND'S BULIMUS. Shell oblong-ovate, whorls convex, columella slightly twisted, lip reflected; white, whorls encircled above and below with a chestnut band, epidermis thin, yellow, peristome and part around the columella purple-brown.

PFEIFFER, MSS. Mus. Cuming.

Hab. Tayabas, Island of Luzon, Philippines (on the leaves of trees); Cuming.

The general aspect of this shell is very simple, but it is not a species of common occurrence.

Species 34. (Mus. Cuming.)

BULIMUS ROMBLONENSIS. *Bul. testá pyramidalí-ovatá,* *afractibus sexis leviter convexis, striis transversis et obliquis subtilissimè obscure decussatis, columellā intortā; virescente-albā, glabrá, nitidā, afractibus fasciē castanē infernē cingulatā, epidermide tenui viriditatem, parte circa columellam nigrante-castanē, columellā et peristriate albīs.*

THE ROMBLON BULIMUS. Shell pyramidaly ovate, whorls slightly convex, obscurely decussated with very fine transverse and oblique striae, columella twisted; greenish-white, smooth, shining, whorls encircled round the lower part with a chestnut band, epidermis thin, greenish yellow, blackish chestnut around the columella, columella and peristriate white.

PFEIFFER, Pro. Zool. Soc., 1842, p. 152.

Hab. Island of Romblon, Philippines (on the trees). Cuming.

A bright shining species closely resembling a variety of *B. cinctinus*, not figured in the accompanying plate, which is sufficiently distinct in form to prevent their being confounded, the one for the other.

Species 35. (Mus. Cuming.)

BULIMUS SUBCARINATUS. *Bul. testá conico-ovatā, sub-*

pellucídā, afractibus sexis convexis, striis transversis et obliquis subtilissimè decussatis, ultimo infernē obscurè subcarinato, columellā viz intortā, labro reflexo, subeffuso; cornuescente-albā, epidermide hydrophanā fuscrescente ad suturas peculiariter auratā, sub alterd corneā.

THE SLIGHTLY-KEELED BULIMUS. Shell conically ovate, somewhat transparent, whorls six in number, convex, very finely decussated with fine transverse and oblique striae, last whorl obscurely slightly keeled, columella but little twisted, lip reflected, somewhat effused; bluish white, epidermis hydrophanous, light brown, peculiarly golden tinged at the sutures, over another one of a thin horny substance.

PFEIFFER, Pro. Zool. Soc., 1842, p. 153.

Hab. Island of Romblon, Philippines (on the leaves of trees); Cuming.

This delicate semi-transparent species serves to illustrate in an interesting manner the phenomenon, observed in many *Bulinæ*, of a double epidermis. The shell under consideration is a clear opal white, it is then covered with a thin, shining, greenish, horny epidermis, over which is a soft, tenacious, porous, hydrophanous one, curiously puckered and tinged with a bronze or golden hue around the sutures.



36



37 α



37 δ



38



39



40



41 α



42 α



42 δ

BULIMUS.

PLATE VIII.

Species 36. (Mus. Cuming.)

BULIMUS CAMELEOPARDALIS. *Bul. testā elongato-oblongā, spirāe valde productā, anfractibus septenīs, planō-concresīs, ultīmō obscurē carinatō, columellā ornatā, aperturā parvissimā; castaneo-flavante, epidermide virgō longitudinalē lotusculas conspicuas, cīx nudatas, prætermittente; peritreme purpureo-nigricante.*

THE CAMELEOPARD BULIMUS. Shell elongately oblong, spire much produced, whorls seven in number, flatly convex, the last obscurely keeled, columella arched, aperture rather small; chestnut yellow, epidermis leaving rather broad conspicuous longitudinal stripes, scarcely waved, peritreme purple-black.

Broderip, Pro. Zool. Soc., 1840, p. 157.

Hab. Sibonga, Island of Zebu, Philippines (on the leaves of bushes); Cuming.

Distinguished by its elongated dactylose form, and by the bold longitudinally striped pattern of the epidermis.

Species 37. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS BOHOLENSIS. *Bul. testā subelongato-oblongā, anfractibus seuis, leviter concresīs, columellā vis intortā; polidē castaneā, apicem versus flavicante, epidermide nuculos sparsas strigasque acutē nudatas, sēpē angustas, numerosas, prætermittente, peritreme purpureo-nigricante.*

THE BOHOL BULIMUS. Shell somewhat elongately oblong, whorls six in number, slightly convex, columella a little twisted; pale chestnut, yellowish towards the apex, epidermis leaving sharply zig-zag streaks, sometimes numerous and narrow, with scattered spots, peritreme purple-black.

Broderip, Pro. Zool. Soc., 1840, p. 158.

Hab. Island of Bohol, Philippines (on leaves of trees); Cuming.

In general aspect this species approximates closely to the preceding; it is, however, less elongated in form, and the epidermal pattern is of an oblique zig-zag character, which is sometimes developed with peculiar neatness.

Species 38. (Mus. Cuming.)

BULIMUS CALISTA. *Bul. testā elongato-ovatā, anfractibus septenīs, concresīs, columellā subintortā; stramineā,*

epidermide tenuiculā, in strigis longitudinalibns nudatis, peritreme purpureo-nigricante, apice rufo-castaneo.

THE CALISTA BULIMUS. Shell elongately ovate, whorls seven in number, convex, columella slightly twisted; straw-colour, epidermis rather thin, in waved longitudinal streaks, peritreme purple-black, apex reddish chestnut.

Broderip, Pro. Zool. Soc., 1842, p. 152.

Hab. Tanhay, Island of Negros, Philippines (on the leaves of bushes); Cuming.

The *B. Calista* is of a delicate straw-colour, with a very light epidermis; in form it approaches the *B. Boholensis*, and the colouring of the peritreme and columella is nearly the same.

Species 39. (Mus. Cuming.)

BULIMUS DACTYLUS. *Bul. testā pyramidali-oblongā, basin versis peculiariter effusā; anfractibus septenīs, concresīs, ultīmō obscurē carinatō; columellā subintortā; castaneo-fuscescēte, epidermide longitudinaliter nebulae nudatā, lineam spiralem inferī prætermissente, peritreme castaneo-nigricante.*

THE FINGER BULIMUS. Shell pyramidal oblong, peculiarly effused towards the base; whorls seven in number, convex, the last obscurely keeled; columella slightly twisted; light chestnut-brown, epidermis longitudinally nebulously waved, leaving a spiral linear mark round the lower part, peritreme chestnut-black.

Broderip, Pro. Zool. Soc., 1842.

Hab. Mountains of Tayabas, Island of Luzon, Philippines (on the leaves of trees); Cuming.

This species, of which Mr. Cuming collected but few specimens, may be recognised by its acuminate growth and effused development of the aperture. It appears to come nearer the *B. coriatus* of Lea in these respects, than any that has been attributed to that species.

Species 40. (Mus. Cuming.)

BULIMUS DIANA. *Bul. testā elongato-oblongā, anfractibus septenīs, concresīs, columellā subintortā; stramineā, apicem versus eburneā, epidermide tenui longitudinaliter nebulae et strigata, peritreme eburneā.*

April, 1848.

THE DIANA BULIMUS. Shell elongately oblong, whorls seven in number, convex, columella slightly twisted; straw-colour, ivory-white towards the apex, epidermis thin, longitudinally streaked and clouded, peritreme very white.

Broderip, Pro. Zool. Soc., 1840, p. 157.

Hab. Tamay, Island of Negros, Philippines (on the leaves of bushes); Cuming.

The chief point of difference between this species and the *B. Culista* appears to consist in the shining ivory-white peritreme and apex, and it is not unlikely, as Mr. Broderip suggests, that they may merge into one and the same species.

Species 41. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS NYMPHA. *Bul. testā subpyramidalē-oblongā, anfractibus senis septenīs, convexis, ultimo obscurē carinato; columellā eix arcuatā; rufō-castaneā, epidermē in strigis longitudinalibus, latis, rectiusculis, peritremate purpureo-nigrante.*

NYMPH BULIMUS. Shell somewhat pyramidal oblong, whorls six or seven in number, convex, the last obscurely keeled; columella scarcely curved; reddish brown, epidermis in rather straight broad longitudinal streaks, peritreme purple-black.

Pfeiffer, Pro. Zool. Soc., 1842, p. 89.

Hab. Fig. *a*, Mount Isarog; Fig. *b*, Catanauan, Island of Luzon, Philippines (on the leaves of trees); Cuming.

The variety represented at Fig. *b* is composed of a whorl less than the other, and is of more ventricose proportions. From *B. Boholeensis*, to which this species is perhaps the nearest allied, it differs in being of a more acuminate form, whilst the pretty zig-zag pattern is wanting in the epidermis.

Species 42. (Mus. Cuming.)

BULIMUS SQUIJOORENSIS. *Bul. testā oblongo-ovatā, tenuiculā, anfractibus senis, subcetratōcosis, ultimo obscurē carinato; columellā rectiusculā, aperturā paululum effusā; flavicante, epidermē tenui, pulcherrimē marmoratā et flammatā, peritremate purpureo-nigrante, apice rufescente-castaneo.*

THE SQUIJOR BULIMUS. Shell oblong-ovate, rather thin, whorls six in number, rather ventricose, the last obscurely keeled; columella nearly straight, aperture a little effused; yellowish, prettily flamed and marbled, peritreme purple-black, apex reddish chesnut.

Pfeiffer, Pro. Zool. Soc., 1845, p. 74.

Hab. Island of Siquijor, Philippines (on the leaves of trees); Cuming.

A light ventricose shell, in which the epidermis is disposed in light sharply zig-zag flames, and beneath the keel, which is obscurely developed on the last whorl, there is an interrupted band of indistinct arrow-headed marks.



43. a



44



43. b



45. a



45. b



45. c



46. a



46. b



46. c

B U L I M U S.

PLATE IX.

Species 43. (Fig. a and b. Mus. Cuming.)

BULIMUS CUYOENSIS. *Bul.* testă pyramidalis-ovata, umbilicata, unfractibus quinque, subrotundata, ventricosimarginata, columella subarcuata; sutulato-fusca vel strawina, nigra unifasciata, epidermide striata conspicua longitudinaliter, vir undatus, exhibente, peritreme albo.

THE CUYO BULIMUS. Shell pyramidally ovate, umbilicated, whorls five in number, somewhat rounded, rather ventricose, columella slightly arched; burnt brown or straw-colour, encircled by a black band, epidermis exhibiting conspicuous longitudinal slightly waved streaks, peritreme white.

PFEIFFER, Pro. Zool. Soc., 1842, p. 252.

Bulimus cahaynas. Jonas.

Hab. Islands of Cuyo and Mindoro, Philippines (on the trees); Cuming.

This very pretty species was first described by Dr. Pfeiffer, from a single indifferent specimen collected by Mr. Cuming at the Island of Cuyo; others subsequently arrived from the Island of Mindoro, exceeding the former so greatly in size and beauty, that their identity was not recognised.

Species 44. (Mus. Cuming.)

BULIMUS FESTILIS. *Bul.* testă cylindraceo-oblonga, subumbilicata, unfractibus sexis, concreta, columella lobatoque latè reflexa; reflo-castanea, epidermide lactea opaca, strias paucas longitudinaliter irregulares praetermittente, peritreme albo.

THE EARTHEN BULIMUS. Shell cylindrically oblong, slightly umbilicated, whorls six in number, convex, lip and columella widely reflected; reddish chesnut, covered with an opaque cream-coloured epidermis, leaving a few irregular longitudinal streaks, peritreme white.

Broderip, Pro. Zool. Soc., 1840.

Hab. Island of Cuyo, Philippines (on the leaves of trees); Cuming.

I much doubt if the *Bulimus lorenzus* of Broderip, (Sowerby, Conch. Illus. f. 117) is distinct from this; it was described, with some hesitation, from a single specimen in rather indifferent condition, and its specific value has not been confirmed by the receipt of any specimens showing more distinctly the characters which it served to indicate.

Species 45. (Fig. a, b, c. Mus. Cuming.)

BULIMUS DRYAS. *Bul.* testă elongato-ovata, crassissimata, subumbilicata, unfractibus sexis, convexa, apertura subcontracta, labro crassissculo, latè reflexo; etiamque plerumque castaneo vel fuscoc-nigricante conspicue fasciata, epidermide ciz nullâ.

THE DRYAS BULIMUS. Shell elongately ovate, rather thick, slightly umbilicated, whorls six in number, convex, aperture somewhat contracted, lip rather thick, broadly reflected; ivory-white, mostly conspicuously banded with chesnut and brownish-black, with scarcely any epidermis.

Broderip, Pro. Zool. Soc., 1840.

Bulimus porraceus. Jay.

Testa juvenilis, *Bulimus paradoxus*, Pfeiffer.

Hab. Mansalai, Island of Mindoro, Philippines (in the woods); Cuming.

Of a peculiarly thickened growth compared with other species, giving a widely reflected lip to the aperture, with an extremely slight epidermis. The rich dark chesnut bands with which it is mostly encircled are deeply stained in the shell.

Species 46. (Fig. a, b, c. Mus. Cuming.)

BULIMUS SYLVANUS. *Bul.* testă elongato-ovata, subpyramidalis, unfractibus sexis, subventricosa, apertura subeffusa, labro latè reflexo; albida, castaneo-fusco plerumque latifasciata, epidermide lateat cornet, strias remota longitudinaliter plerumque exhibente, peritreme albo, sepe castaneo vel roseo marginato.

THE SYLVANUS BULIMUS. Shell elongately ovate, somewhat pyramidal, whorls six in number, rather ventricose, aperture slightly effused, lip widely reflected; whitish, generally broadly banded with chesnut-brown and covered with a thin yellow horny epidermis, frequently exhibiting remote longitudinal streaks, peritreme white, often edged with rose or chesnut.

Broderip, Pro. Zool. Soc., 1840.

Hab. Puerto Portero, Island of Mindoro (in the woods); Cuming.

This species, it will be observed, is distinguished from the preceding by its lighter growth, deeper colour, and more developed epidermis.



BULIMUS.

PLATE X.

Species 47. (Mus. Cuming.)

BULIMUS APLOMORPHUS. *Bul. testá conico-ovatá, axi-
fractibus sexis, convexis, ultimo superne obscure angu-
lato, aperturá parviscalá, labro fortiter reflexo; eas-
taneo-lutescente, zonis rufofusca duabus cingulata,
epidermide tenuissimá, peritreme albo.*

THE SIMPLE-FORM **BULIMUS.** Shell conically ovate, whorls six in number, convex, last whorl obscurely angled at the lower part, aperture rather small, lip strongly reflected; chestnut-yellow, encircled with two red-brown zones, epidermis very slight, peritreme white.

JONAS, Pro. Zool. Soc., 1842, p. 189.

Hab. Philippine Islands.

Allied to *A. Guimarasensis* but of another form, differing principally in being more conical and in having a second thin white hydrophanous epidermis, which, however, in the specimen figured, is much rubbed away. It was not collected by Mr. Cuming.

Fig. 19 c and d. (Mus. Cuming.)

The shells here represented are without doubt varieties of the *B. pictor* described at Plate IV, although so much smaller in size, and not characterized by the zigzag shadowed flames depicted in the specimens there figured: in these the outer hydrophanous epidermis is either arranged in slightly waved streaks, or it covers the shell leaving streaked spaces. Where the varieties chiefly simulate is in the livid purple-black colouring of the peritreme.

Species 48. (Mus. Cuming.)

BULIMUS UBER. *Bul. testá subabbreviato-ovatá, tenuiculá,
diaphana, anfractibus quaternis, ventricosus, longitu-
dinaliter confertia regulariter striatus, ultimo spira
magnitudine exaltata superante; pelliculo-albá, epider-
mide tenui intescente indutá.*

THE NIPPLE **BULIMUS.** Shell somewhat shortly ovate, rather thin, diaphanous, whorls four in number, ventricose, longitudinally closely regularly striated, last whorl far exceeding the size of the spire; transparent white, covered by a thin yellowish epidermis.

PFEIFFER, Pro. Zool. Soc., 1842, p. 153.

Hab. Island of Guimaras, Philippines, (on the leaves of trees); Cuming.

The spire of this shell is of a peculiarly swollen mamillary structure, whilst the shell is composed of no more than four whorls; it is of a semitransparent diaphanous texture with only a very simple shining epidermis.

Species 49. (Mus. Cuming.)

BULIMUS EGLE. *Bul. testá oblongo-ovatá, eix pyramidalis,
anfractibus quinque, convexis, oblique creberius striatis,
ultimo subfuso, inferne obscure obtuso-carinato; fuléa,
basí indistincte fasciata, suturis subtusissimè albiliucentis,
peritreme conspicue castaneo-fusco.*

THE EGLE **BULIMUS.** Shell oblong-ovate, scarcely pyramidal, whorls five in number, convex, obliquely very closely striated, last whorl somewhat effused, obscurely obtusely angled round the lower part; fulvous, indistinctly banded at the base, sutures marked with a very fine white line, peritreme conspicuously stained with chestnut-brown.

BRODERIP, Pro. Zool. Soc., 1840, p. 181.

Hab. Casan, Province of Misamis, Island of Mindanao, Philippines (in a dense forest on the leaves of trees); Cuming.

Peculiar in form but of a dull fulvous hue, not distinguished by any ornamental epidermis, nor of an aspect at all complimentary to the fair nymph to whom it is dedicated.

Species 50. (Mus. Cuming.)

BULIMUS LEUCOPHEUS. *Bul. testá subpyramidalis-ovatá,
crassiusculá, plus minus ventricosa, anfractibus sexis,
convexis, columella recta, apertura subcontráta, labro
reflexo; castaneo-fusca, epidermide leucophaea, sepe
albiflumata, inferne fasciá angusta et lineá nigricante
cingulata, interdum zoná, infra suturas, albida, hic illuc
peculiariter abrasa, columellá et apertura fance ca-
runculente-albis, peritreme castaneo-nigricante mar-
ginato.*

THE RUSSET-GREY **BULIMUS.** Shell somewhat pyramidal-ovate, rather thick, more or less ventricose, whorls six in number, convex, columella straight, aperture slightly contracted, lip reflected; chestnut-brown, covered with a russet-grey epidermis, often

tan of with white, encircled round the lower part with a dark line or narrow band, sometimes with a white band beneath the sutures peculiarly abraded here and there, columella and interior of the aperture bluish white, peristome edged with chestnut-black.

Helix leucophaea, Sowerby, Pro. Zool. Soc., 1841, p. 19.

Hab. Dolores, Province of Pampanga, and Mountains of Igarrotes, Province of Cagayan, Island of Luzon, Philippines (on the trees); Cuming.

There is a singular peculiarity attending this species in the manner in which the epidermis, especially on the band which is sometimes seen beneath the sutures, is abraded, denuding the shell in patches.

Species 51. (Mus. Cuming.)

BULIMUS LEOPARDUS. *Bul. testā subabbreviato-ovatā, ventricosā, anfractibus quinque, convexis, longitudinaliter restringit costulato-striatā, ultima spira longitudine suprae, columellā arcuata, aperturā amplissimā; lateo-castaneā, epidermide albida sparsā, sub alternaā tenui cornēa, variegata, peristremae purpuro-castaneo.*

THE LEOPARD BULIMUS. Shell somewhat shortly ovate, ventricose, whorls five in number, convex, very closely sculptured longitudinally with fine rib-like striae, last whorl exceeding the length of the spire, columella arched, aperture very large; yellowish-chestnut, variegated with a scattered whitish epidermis upon a thin horny one, peristome purple-chestnut.

PFEIFFER, Pro. Zool. Soc., 1845, p. 67.

Hab. Island of Mindoro, Philippines.

This remarkable species, of which Mr. Cuming has only the single specimen here represented, is distinguished from all others by the effused and suddenly enlarged development of the last whorl. It is also characterized by a regular succession of longitudinal rib-like striae of a more defined growth than the ordinary lines of structure, which in a genus so devoid of sculpture is an important feature.

The name is suggested by the scattered interrupted stripes in which the outer hydrophanous epidermis is deposited. The columella, it should be observed, has a pioky hue, and the impress of the striae imparts a delicate satiny iridescence to the interior of the shell.

Species 52. (Mus. Cuming.)

BULIMUS CONCINNUS. *Bul. testā pyramidali-oblongā, anfractibus sexis planō-concēxis, aperturā contractā, labro reflexo; castaneo-fusca, epidermide concinna albido-marmoratā et flammatā, columellā et aperturā fauce overulescente-albīs, peristrema nigricante-castaneo*

THE PRETTY BULIMUS. Shell pyramidal oblong, whorls six in number, flatly convex, aperture contracted, lip reflected; chestnut-brown epidermis prettily flamed and marbled with white, columella and interior of the aperture bluish white, peristome dark chestnut.

Helix concinna, Sowerby, Pro. Zool. Soc., 1841, p. 20.

Hab. St. Juan, Province of Cagayan, Island of Luzon, Philippines (on the leaves of trees); Cuming.

I much doubt whether this is anything more than a variety of the *B. leucophaea*; it is, however, a well-marked and extremely pretty one.



54. a



55. a



55. b



b



c



d



e



f



g



h

BULIMUS.

PLATE XI.

Species 53. (Mus. Cuming.)

BULIMUS INCOMPTUS. *Bal. testā oblongo-ovalā, obtuso-pyramidalī, anfractibus quinīs, rotundatī, columellā rectā, labro vix reflexo; lutescēte-brunādā, zonulā nigricante-fuscā cingulatā, epidermide tenī indutā.*

THE UNADORNED BULIMUS. Shell oblong ovate, obtusely pyramidal, whorls five in number, rounded, columella straight, lip but slightly reflexed; yellowish brown, encircled by a narrow blackish brown zone, covered by a thin epidermis.

Helix incuncta, Sowerby, Pro. Zool. Soc., 1840.

Hab. Island of Tablas, Philippines.

Only a single specimen of this species was collected, and it has very much the appearance of being an abortive growth of one subsequently named by Mr. Sowerby, *Helix acuminata* (Sp. 60); they are identical in colour and pattern, but vary materially in form.

Species 54. (Fig. a and b. Mus. Cuming.)

BULIMUS PARTULOIDES. *Bal. testā pyramidalī, spirā acuminatā, anfractibus quinīs, convexis, aperturā parvissentā, subquadrate-ovali, labro latissimō reflexo; alba, zonulā castaneo-nigricante cingulatā, epidermide testē indutā, interdūa omnino intenſe castaneā, peritreme albo.*

THE PARTULA-LIKE BULIMUS. Shell pyramidal, spire acuminate, whorls five in number, convex, aperture rather small, somewhat square-ovate, lip very broadly reflexed; whitish, encircled by a narrow chestnut black zone, covered by a yellow epidermis, sometimes dark chestnut throughout, peritreme white.

BRODERIP, Pro. Zool. Soc., 1840, p. 181.

Hab. Island of Tablas, Philippines (on the leaves of trees); Cuming.

This is the nearest approach to *Partula* of any species of the genus, the aperture having the same square-like form and widely reflexed lip.

Species 55. (Mus. Cuming.)

BULIMUS LUZONICUS. *Bal. testā subelongato-ovalā, apice obusa, anfractibus sensīs, convexis, columellā intortā, aperturā subcontractā, labro reflexo; eburnēdā, zonulā angustā nigricante conspicue cingulatā.*

May, 1848.

THE LUZON BULIMUS. Shell somewhat elongately ovate, obtuse at the apex, whorls six in number, convex, columella twisted, aperture somewhat contracted, lip reflexed; ivory white, conspicuously encircled by a narrow black zone.

Hab. Island of Masbate, Philippines (on the trees); Cuming.

I have always considered this shell to represent a species distinct from *B. ovoideus*, and propose to re-establish Mr. Sowerby's name, although he himself abandoned it upon the grounds of its relationship. It is of a more elongately oblong form, is composed of a whorl more, and the narrow black zone with which it is encircled never varies. These characters may be regarded as of trifling importance, but they are constant. In a parcel of upwards of a hundred of each species which came into my possession some time since, there was no variation or the least tendency to merge one into the other. The name is rather an inappropriate one, for Mr. Cuming did not collect a single species at Luzon; there is, however, no reason why it might not become naturalized in that island.

Species 56. (Fig. a and b. Mus. Cuming.)

BULIMUS OVOIDEUS. *Bal. testā oblongo-ovalā, subeven-tricata, apice obtusa, anfractibus quinīs, convexis, columellā vix intortā, labro reflexo; albā, fasciis dubiis tribus latissimis aurantio-vel castaneo-fuscis cingulatā, epidermide tenuissimā.*

THE OVOID BULIMUS. Shell oblong ovate, rather ventricose, obtuse at the apex, whorls five in number, convex, columella scarcely twisted, lip reflexed; white, encircled with two or three rather broad orange-brown or chestnut bands, epidermis very thin.

BRUGUIERE, Enc. Méth. Dict., No. 64.

Hab. Islands of Ticao and Masbate, Philippines (on the trees); Cuming.

Distinguished by its oval ventricose growth and broad spiral bands of colour.

Species 57. (Mus. Cuming.)

BULIMUS VELATUS. *Bal. testā oblongā, sub-Pupaformi, anfractibus sensīs, subrotundatis, aperturā parvisculptā, labro simplici; lutescente-albā, zonulā rufo-fuscescente*

cingulatā, epidermē *tenui*, *albicante*, *strigas longitū-*
dinates subindistincte prætermittente, *peritremate rufof-*
fusco.

THE CLOTHED BULIMUS. Shell oblong, somewhat pupa-shaped, whorls six in number, subrotund, aperture rather small, lip simple; yellowish white, encircled with a narrow red-brown zone, epidermis thin, whitish, leaving somewhat indistinctly longitudinal streaks, peritreme red-brown.

BRODERIP, Pro. Zool. Soc., 1841, p. 15.

Hab. Island of Camote, Philippines (on the leaves of trees); Cuming.

Of this species Mr. Cuming only collected three specimens, and these appear to be of somewhat immature growth. It is not distinguished by any marked peculiarity of character.

Fig. 58. (Mus. Cuming.)

This shell was named in Mr. Cuming's collection *B. nebulosus*, but as I am doubtful of its being a variety of the *B. calista*, it may for the present be referred to that species. For sp. 58, see Pl. XIII.

Species 59. (Mus. Cuming.)

BULIMUS PYRAMIDALIS. *Bul. testā oblongo-pyramidalī,*
apice obtusā, aequaliter rotundata, subrotundis, apertura
parvisepta, labro simplici; fulcā, basin versus sutu-
risque nigrante-costatae, epidermē *tenui indutā*,
columnellā albā, *peritremate fusco*.

THE PYRAMIDAL BULIMUS. Shell oblong-pyramidal, obtuse at the apex, whorls six in number, somewhat

round, aperture rather small, lip simple; yellowish, blackish chestnut towards the base and in the sutures, covered with a thin epidermis, columella white, peritreme brown.

Helix pyramidalis, Sowerby, Pro. Zool. Soc., 1841, p. 39.
 Hab. Island of Cuyo, Philippines (upon the leaves of trees); Cuming.

The colour of this shell is a dark burnt brown, with traces of a light puckered epidermis beneath the sutures. It may be readily distinguished by this and its pyramidal growth.

Species 60. (Mus. Cuming.)

BULIMUS ACUMINATUS. *Bul. testā subpyramidalī-ovatā,*
apice acuminatā, aequaliter rotundata, convexis, ultimo
obscure subangulatō, labro tenui, vix reflexo; lutescente-
brunneā, zonatā castaneā cingulatā, epidermē *tenui*
indutā, columnellā *albā*, *peritremate rufo-fusco*.

THE ACUMINATED BULIMUS. Shell somewhat pyramidal-ovate, spire acuminated, whorls five in number, convex, the last obscurely slightly angled, lip thin, but slightly reflexed; yellowish-brown, encircled by a narrow chestnut zone, covered with a thin epidermis, columella white, peritreme red-brown.

Helix acuminata, Sowerby, Pro. Zool. Soc., 1841, p. 39.
 Hab. Island of Cuyo, Philippines (upon the leaves of trees); Cuming.

Of a peculiar yellowish-brown colour encircled by a narrow zone of red-brown and stained with very dark chestnut round the columella. It differs from *B. incomptus* only in form, its general aspect and colouring are the same.



BULIMUS.

PLATE XII.

Species 61. (Mus. Cuming.)

BULIMUS CUMINGII. *Bul. testá ovalá, spirá subobtusa, anfractibus quinque, oblique tenuistriatis, ultimis infraeum indistincte obtuso angulato, columellá planá, intortá; pellucido-albá, nitidá, basí versus virescente, linea castanea conspicua infra suturas circumdatá, columellá purpureo-fusca, peritreme alba.*

CUMING'S BULIMUS. Shell ovate, spire rather obtuse, whorls five in number, finely obliquely striated, the last indistinctly obtusely angled round the lower part, columella flat, twisted; transparent white, shining, greenish towards the base, surrounded by a conspicuous chestnut line beneath the sutures, columella purple-brown, peritreme white.

PFEIFFER, Pro. Zool. Soc., 1842, p. 88.

Hab. Island of Cauning, Philippines (on leaves of trees); Cuming.

A transparent shining opal-like shell, peculiarly characterized by a dark chestnut line running beneath the sutures.

Species 62. (Mus. Cuming.)

BULIMUS VERECUNDUS. *Bul. testá ovalá, tenuicula, ventricosa, anfractibus quinque, convexis, tuwidis, columellá recta, labro tenui, eix reflexo; albá, fusca, carno-fuscescentibus latis interruptis cingulata, peritreme carneo-fuscescente.*

THE MODEST BULIMUS. Shell ovate, rather thin, ventricose, whorls five in number, convex, swollen, columella straight, lip thin, very slightly reflected; white, encircled by broad interrupted light flesh-brown bands, peritreme flesh-brown.

Helix modesta, Sowerby, Pro. Zool. Soc., 1841, p. 39.

Hab. Catanauan, Province of Tayabas, Island of Luzon (on the leaves of trees); Cuming.

The adoption of this delicate species under the type of *Bulimus* imposes the necessity of substituting a new specific name, that of *modestus* being already occupied.

Species 63. (Mus. Cuming.)

BULIMUS SIMPLEX. *Bul. testá ovato-conoidea, anfractibus quinque, convexis, columellá subintortá, labro subeffuso, reflexo; albá, basí versus viridescente, epidermide tenui, cornéa.*

THE SIMPLE BULIMUS. Shell ovately conoid, whorls five in number, convex, columella slightly twisted, lip somewhat effused, reflected; white, greenish towards the base, with a very thin horny epidermis.

JONAS, Pro. Zool. Soc., 1842, p. 189.

Hab. Island of Mindoro, Philippines.

The green colouring of this species seems to reside in great measure in its thin horny epidermis.

Species 64. (Mus. Cuming.)

BULIMUS EFFUSUS. *Bul. testá ovalá, basí versus ventricosa, effusa, anfractibus quinque, convexis, columellá vis arcuata, labro sublate reflexo; albá, nitidá, zonulis duabus castaneis conspicue cingulata, epidermide tenui cornéa lutescente inferne intota.*

THE EFFUSED BULIMUS. Shell ovate, ventricose and effused towards the base, whorls five in number, convex, columella scarcely arched, lip somewhat broadly reflected; white, shining, sometimes conspicuously encircled with two narrow chestnut zones, covered at the lower part with a thin yellowish horny epidermis.

PFEIFFER, Pro. Zool. Soc., 1842, p. 152.

Hab. Island of Tablas, Philippines (on the leaves of trees); Cuming.

This shell approaches rather to the *Helix decipiens* in form and general aspect than to the *B. Cumingii*, with which Dr. Pfeiffer compares it, and might be referred to that genus with almost equal propriety.

Species 65. (Mus. Cuming.)

BULIMUS BREVICULUS. *Bul. testá crassiuscula, peculiariter abbreviato-pyramidalis, apice obtusa, anfractibus sex, angustis, aperturá parci, labro sublate reflexo; albá, epidermide luteo-fuscescente.*

THE SHORTENED BULIMUS. Shell rather thick, peculiarly abbreviately pyramidal, obtuse at the apex, whorls six in number, narrow, aperture small, lip rather widely reflected; white, epidermis yellowish brown.

PFEIFFER, Pro. Zool. Soc., 1842, p. 88.

Hab. Island of Romblon, Philippines (on bushes); Cuming.

The shell is now in very good condition and only a vestige of the epidermis is visible near the aperture.

Species 66. (Mus. Cuming.)

BULIMUS LIMA. *Bul. testá orato-conoideá, solidimisculá, sp. in sublonga, anfractibus quinque, rotundatis, columella oblonga, labro reflexo; albá, epidermide tenui strati, opacá.*

LEVI'S BULIMUS. Shell ovately conoid, rather solid, spire somewhat obtuse, whorls five in number, rounded, columnella rather flattened, lip reflected; white, covered with a thin straw-coloured epidermis.

PFEIFFER, Pro. Zool. Soc., 1846, p. 29.

Hab. Philippines Islands.

A shell of simple character though not exactly agreeing with any other species.

Species 67. (Mus. Cuming.)

BULIMUS STABILIS. *Bul. testá cylindraceo-ovatá, crassissimá, basi effusa, anfractibus sex, plano-euariis, angustis, aperturá obliqua, labro forte reflexo; albá, anfractuosa parte inferiori intus ustulato-castanó, peritreme alba.*

THE FIRM BULIMUS. Shell cylindrically ovate, rather thick, effused at the base, whorls six in number, flatly convex, narrow, aperture oblique, lip strongly reflected; white, lower part of the whorls deep burnt chestnut, peritreme white.

Hab. stabili. Sowerby, Pro. Zool. Soc., 1846, p. 104.

Hab. Isle of Temple, Philippines; Cuming.

A solid pupiform shell, approximating in some degree to the *B. oroides* but of more solid structure and very peculiar in its style of colouring.

Species 68. (Mus. Cuming.)

BULIMUS BULLULA. *Bul. testá ovo-conoideá, anfractibus quatuor, convexis, apertura subeffusa, labro vis reflexo; cerascente-albá diaplaná, aut flavicante, apice pallidé rubido.*

THE LITTLE BUBBLE BULIMUS. Shell ovately conoid, whorls four in number, convex, aperture somewhat effused, lip slightly reflected; transparent bluish white or yellowish, pale ruddy colour at the apex.

BRODERIP, Pro. Zool. Soc., 1840, p. 159.

Hab. Puerto Galero, Island of Mindoro, Philippines; Cuming.

A most delicately coloured species of a transparent opal white tinged with red at the apex.

Species 69. (Mus. Cuming.)

BULIMUS DILATATUS. *Bul. testá crassissimá, abbreviatopiramidalis, subcentricosá, anfractibus sex, angustis, columellá plano-dilatátá, apertura parceuseulá; nigricante-purpureá, apicem versus rufescente, epidermide fusca, hydrophaná, columellá et apertura fave albís, peritreme nigricante-purpureo marginata.*

THE DILATED BULIMUS. Shell rather thick, abbreviately pyramidal, somewhat ventricose, whorls six in number, narrow, columnella flatly dilated, aperture rather small; blackish-purple, reddish towards the apex, with a brown hydrophanous epidermis, columnella and interior white, peritreme edged with purple-black.

PFEIFFER, Pro. Zool. Soc., 1846, p. 42.

A solid contracted pyramidal shell approaching somewhat in form to the *B. leucophaea*; and the hydrophanous epidermis, of which this specimen exhibits the remains, appears to be abraded in patches beneath the sutures as in that species.

Singularly allied in form to the *B. dilatus*, which is of a dark purple-brown colour, but somewhat less ventricose.



BULIMUS.

PLATE XIII.

Species 58. (Mus. Cuming.)

BULIMUS ZONULATUS. *Bul. testā conico-oblongā, tenui, umbilicatā, anfractibus concavis, levibus, labro columellarī reflexo, umbilicūm frē occultante, externo simplici; pallide opaco-strawīvā, macularū pellucidarū serierū duabus cinctā, anfractū ultimo zonulā daubas castaneā ornato.*

THE FINELY ZONED BULIMUS. Shell conically oblong, thin, umbilicated, whorls convex, smooth, columellar lip reflected so as almost to conceal the umbilicus, outer lip simple; pale opaque straw-colour, encircled by two rows of transparent spots, last whorl ornamented with two narrow chestnut zones.

PFEIFFER, Pro. Zool. Soc., 1846, p. 41.

Hab. Cabanatuan, Province of Nueva Eaja, Island of Luzon; Cuming.

There is a very singular character attending this species: the shell is of a thin transparent horny substance, covered by an opaque straw-coloured layer except in those places by which the pattern is produced, the spots thus have an appearance, so to speak, of little windows in the shell.

Species 70. (Mus. Cuming.)

BULIMUS PANAMENSIS. *Bul. testā oblongo-ovatā, tenui, vix umbilicatā, anfractibus quinque concavis, levibus, aperturā parviusculā, labro simplici; pallide fulva, diaphana.*

THE PANAMA BULIMUS. Shell oblong-ovate, thin, slightly umbilicated, whorls five in number, smooth, aperture rather small, lip simple; pale fulvous colour, dia-phanaous.

BRODERIP, Pro. Zool. Soc., 1832, p. 105.

Hab. King's and Saboga Islands, Bay of Panama (on the trunks of large trees); Cuming.

The form of this shell is extremely simple, and its colour a semi-transparent nankeen.

Species 71. (Mus. Cuming.)

BULIMUS TRANSLUCENS. *Bul. testā oblongo-ovatā, tenui, valde pellucidā, vix umbilicatā, anfractibus quinque, subcentricosis, levibus aut transversis et longitudinae-liter minime striatis, labro simplici; corneo-lutescente.*

THE TRANSPARENT BULIMUS. Shell oblong-ovate, thin,

very pellucid, slightly umbilicated, whorls five in number, smooth or transversely and longitudinally minutely striated, lip simple; pale horny yellow.

BRODERIP, Pro. Zool. Soc., 1832, p. 31.

Hab. King's and Saboga Islands, Panama (upon the trees); Cuming.

Of a beautiful transparent glassy structure, through which the internal winding of the columella may be plainly seen. It is very similar in form to the *B. Panamensis*, and is from the same locality.

Species 72. (Mus. Cuming.)

BULIMUS EVANESCENTS. *Bul. testā subobtuso-pyramidalī, opice obtusa, anfractibus quinque, concavis, levibus, nitidusculis, aperturā parvissimā, labro reflexo; virginate-alba, subdiaphana, linea nigra infra suturas, parte circa columellam conspicue nigra.*

THE EVANESCENT BULIMUS. Shell somewhat abbreviately pyramidal, obtuse at the apex, whorls five in number, convex, smooth, rather shining, aperture somewhat small, lip reflected; greenish white, subdiaphanous, with a black line beneath the sutures, and the part around the columella conspicuously black.

PFEIFFER, (inedit?).

Hab. Philippine Islands; Cuming.

A delicate semi-transparent shell easily recognised by the black line between the sutures, and broader band around the columella.

Species 73. (Mus. Cuming.)

BULIMUS ADAMSII. *Bul. testā ovato-turrītā, sinistrali, anfractibus concavis, levibus, aperturā parvissimā, labro reflexo; virescente aut latè, maculis irregularibus subquadris, nigricante-viridibus medio confertim uniseriatim cinctā, interdum immaculatā, parte circa columellam purpureo-roseā.*

ADAMS' BULIMUS. Shell ovately turreted, sinistral, whorls convex, smooth, aperture rather small, lip reflected; greenish or yellow, encircled round the middle with a row of close-set irregular somewhat square blackish green spots, sometimes unspotted, the part around the columella purple-rose.

Hab. Eastern Coast of Borneo (on a tall tree in an islet between Banguey and Balambangan); Adams.

An interesting new species discovered by Mr. Arthur Adams, at the above mentioned locality, during the recent expedition of H.M.S. Samarang. "A tree, partially cut through at the base with axes," says the author of the valuable summary of Natural History attached to the Narrative of that Voyage, "fell upon one of the carpenters wooding on the island, depriving him at the time of sensation. What proved a misfortune to the man, however, presented to science a new and beautiful species of *Bulimus*, which I discovered in considerable numbers and variety adhering to the prostrate tree." All the specimens were reversed, of a yellow or yellowish green colour, stained with purple rose about the columella, and mostly encircled round the middle by a chain of square greenish spots.

Species 74. (Mus. Cuming.)

BULIMUS EBURNEUS. *Bul. testá pyramidali-oblonga, an-*
fractibus sex, rotundatis, tarcios, politis, labro reflexo;
eburnea, epidermide tenuissimè straminea partim indutæ.

THE IVORY BULIMUS. Shell pyramidal oblong, whorls six in number, rounded, smooth, polished, lip reflected; ivory white, partially covered with a very slight straw-coloured epidermis.

Bulimus Diana var., Broderip.
Hab. —?

Mr. Broderip has, I think erroneously, considered this shell a variety of *B. Diana*; in form it differs in the whorls being more rounded, whilst it is of thinner texture and of a pure diaphanous white, the surface being characterized by an extremely delicate natural polish.

Species 75. (Mus. Cuming.)

BULIMUS ERYTHROSTOMA. *Bul. testá subgloboso-ovatæ,*
umbilicatæ, anfractibus quinque rotundatis, concentricè
rugè striatis, columellæ sublatæ reflexæ, labro simplici;
opaco-albâ, columellæ apice et aperture fuscæ rosæ.

THE RED-MOUTH BULIMUS. Shell somewhat globosely ovate, umbilicated, whorls five in number, rounded, concentrically rudely striated, columella rather widely reflected, lip simple; opake-white, columella, apex, and interior of the aperture rose-pink.

SOWERBY, Pro. Zool. Soc., 1833, p. 37.

Hab. Huasco, Chili (under bushes); Cuming.

Of a peculiar bluish opake white having the columella, apex, and aperture coloured with a delicate rose-pink.



77



76



78



79



80



81



82



83



84



85

BULIMUS.

PLATE XIV.

Species 76. (Mus. Cuming.)

BULIMUS PANAYENSIS. *Bul. testā elongato-turrītā, levigatā, pellucidā, apice obtusā, aufractibus octo, apertura minimā, labro simplici; cereo-albida.*

THE PANAY BULIMUS. Shell elongately turreted, smooth, pellucid, obtuse at the apex, whorls eight in number, aperture very small, lip simple : waxes white.

PFEIFFER, Pro. Zool. Soc., 1846, p. 33.

Hab. Dingle, Isle of Panay, Philippines; Cuming.

This is the smallest species collected by Mr. Cuming at the Philippines, it is of an elongately turreted form and of a hyaline waxes white.

Species 77. (Mus. Cuming.)

BULIMUS RHODACME. *Bul. testā oblongo-ovatā, umbilicatā, aufractibus sex, minutiè autem conspicue striatis, striis transversis subsoletis vittulissimè decussato-granulatis, labro simplici; opaco-albida, maculis floridisque roseis subpellucidis versus apicem.*

THE RED-TIPPED BULIMUS. Shell oblong-ovate, umbilicated, whorls six in number, minutely but conspicuously striated, and very minutely decussated with somewhat obsolete transverse striae, lip simple ; opake-white, with semitransparent rose-tinted spots and flaves towards the apex.

PFEIFFER, Pro. Zool. Soc., 1842, p. 187.

Hab. Frierina, Huasco, Chili (under small *Cacti*) ; Bridges.

A delicate species of a seemingly transparent texture, enveloped, save about the apex, with an opake-white coating.

Species 78. (Mus. Cuming.)

BULIMUS MICRA. *Bul. testā subcongatā, vix umbilicatā, senipellucidā, aufractibus subrotundatis, creberrime concentricè striatis, spira suturis peculiariter impressis, apertura parvā, labro simplici; corneo-albida.*

THE SMALL BULIMUS. Shell rather elongated, slightly umbilicated, semipellucid, whorls somewhat rounded, very closely concentrically striated, sutures of the spire peculiarly impressed, aperture small, lip simple ; horny white.

D'ORBIGNY, Voy. dans l'Amér. mérid. Moll. p. 262, pl. 41. f. 18 to 20.

Hab. Eastern side of the Bolivian Andes, near Santa Cruz (under stones), D'Orbigny ; near Chilon, Bolivia (on the dead trunks of *Cacti*) ; Bridges.

A very small semitransparent shell, thickly sculptured with rude concentric striae.

Species 79. (Mus. Cuming.)

BULIMUS TEREBRATIS. *Bul. testā elongatā, subcyindrica, latè angulato-umbilicatā, spira suturis impressis, aufractibus deinceps ad undecim, plano-concavis, obliquè crebre corrugato-striatis, columnellā subconcauo-effusa, apertura minima, peririmata dilatata, fuscescente-alba, apice versus nigricante.*

THE AUGER-LIKE BULIMUS. Shell elongated, somewhat cylindrical, broadly angularly umbilicated, sutures of the spire impressed, whorls ten to eleven in number, flatly convex, obliquely closely striated in a wrinkled manner, columnella concavely effused, aperture very small, peristome dilated; brownish-white, blackish towards the apex.

PFEIFFER, Pro. Zool. Soc., 1842, p. 187.

Hab. Coquimbo, Chili (on rocks and plants near the sea) ; Bridges.

Distinguished from the *B. Papiformalis* to which it is most nearly allied, by its obliquely corrugated striae, widely angulated umbilicus, and neat dilated aperture.

Species 80. (Mus. Cuming.)

BULIMUS ELONGATUS. *Bul. testā elongatā, aufractibus octo, latiusculis, plano-concavis, sub epidermide tenuissimè et creberrimè striatis, apertura parvā, superè subcontracta, labro simplici; stramineo-albida, subpellucida.*

THE ELONGATED BULIMUS. Shell elongated, whorls eight in number, rather broad, flatly convex, very finely and closely striated, aperture small, somewhat contracted towards the upper part ; pale straw-colour. semitransparent.

PFEIFFER, Pro. Zool. Soc., 1846, p. 42.

Hab. Catanaun, Province of Baie, Island of Luzon, Philippines (at the roots of trees) ; Cuming.

The whorls of this species are each slightly contracted round the upper part.

BULIMUS.—PLATE XIV.

Species 81. (Mus. Cuming.)

BULIMUS GRATELOUPI. *Bul. testa elongato-oratā, tenui,
pellucida, anfractibus sex ad septem, levibus, nitidis,
et concentricè subtiliissimè striatis, columellā reflexā,
labro simplici, arcuato, superne sinuato; cornueo-
straminea.*

GRATELOUP'S BULIMUS. Shell elongately ovate, thin, pellucid, whorls six to seven in number, smooth, shining, or concentrically very finely striated, columella reflected, lip simple, curved, sinuated at the upper part; horny straw-colour.

PFEIFFER, Pro. Zool. Soc., 1846, p. 42.

Hab. Islands of Luzon and Panay, Philippines (at the roots of trees); Cuming.

A transparent horny shell very finely marked throughout with concentric striae of growth, resulting from the curved structure of the lip.

Species 82. (Mus. Cuming.)

BULIMUS COCHLEADES. *Bul. testa elongata, anfractibus
nove, latiusculis, levibus, convexis, apertura parvā,
superne contractā, labro simplici; albida.*

THE SPOON-LIKE BULIMUS. Shell elongated, whorls nine in number, rather broad, smooth, convex, aperture small, slightly contracted at the upper part, lip simple; whitish.

Hab. Island of Cuyo, Philippines; Cuming.

This singular species, it may be observed, is represented of the natural size, whilst the rest of the plate are all more or less magnified. Mr. Cuming only collected this solitary specimen, which is quite unapproached by any other. It is of extremely simple structure.

Species 83. (Mus. Cuming.)

BULIMUS LICHENORUM. *Bul. testa oblongo-turrīta, um-
bilicata, anfractibus septem, subrotundatis, levibus aut
tenuisstriatis, columellā reflexā, apertura parriculata,
labro simplici; carulecente-alba, striis perparvis
cinereis obliquis, apice nigricante.*

THE LICHEN BULIMUS. Shell oblong-turreted, whorls seven in number, somewhat rounded, smooth or finely striated, columella reflected, aperture rather small, lip simple; bluish-white, with a few oblique ashy streaks, apex blackish.

D'ORBIGNY, Voyage dans l'Amér. mérid. Moll. p. 264.

pl. 41, f. 9 to 11.

Hab. Cobija (on lichens); D'Orbigny. Chilou, Bolivia (under dead trunks of *Cacti*); Bridges.

It is not without some doubt that I venture to refer this shell to the above named species of D'Orbigny; yet it accords sufficiently with his description and figure.

Species 84. (Mus. Cuming.)

BULIMUS SCABIOSUS. *Bul. testa elongato-oblonga, anfrac-
tibus septem, convexis, peculiariter interrupte sulcatis
et striatis, columellā reflexā, apertura parvā, labro
simplici; albida, rufo-fuscescente striigata et marmo-
ratā, apicem versus nigricante.*

THE SCRUB BULIMUS. Shell elongately oblong, whorls seven in number, convex, peculiarly grooved and striated in an interrupted manner, columella reflected, aperture small, lip simple; whitish, streaked and marked with light reddish-brown, blackish towards the apex.

SOWERBY, Pro. Zool. Soc., 1833, p. 74.

Hab. Cobija, Peru (under stones); Cuming.

Mainly distinguished by a roughened peculiarity of sculpture which appears to have suggested the above name to Mr. Sowerby, though no mention is made of it in his description.

Species 85. (Mus. Cuming.)

BULIMUS PUPIFORMIS. *Bul. testa cylindraceo-elongata,
anfractibus decem, convexis, levibus et tenuistratiis,
ad suturas nivale crenulatis, columellā reflexā, apertura
parvā, labro simplici, subdilatato; cinereo-albida,
apicem versus nigricante.*

THE PUPA-SHAPED BULIMUS. Shell cylindrically elongated, whorls ten in number, convex, smooth or finely striated, minutely crenulated at the sutures, columella reflected, aperture small, lip simple, slightly dilated; ashy white, blackish towards the apex.

BRODERIP, Pro. Zool. Soc., 1832, p. 105.

Hab. Huasco, Chili (under stones in shady places); Cuming.

It is important to observe the minute sutural crenulations which characterize this species, as Mr. Broderip appears to have overlooked them.



BULIMUS.

PLATE XV.

Species 86. (Fig. *a*, *b*, *c*, Mus. Cuming.)

BULIMUS TUPACI. *Bul. testā oblongo-vel elongata-ovalā, subumbilicata, ventricosa, aufractibus septem, crenatō fortiter et irregulariter striatis, per marginem superiorē plus minuscre crenatā, labro vir reflexo; olivaceo-brunnea, interdum quinquefasciatā, raro flavo-caerulea, columellā labroque albī,*

TUPAC'S BULIMUS. Shell oblong or elongately ovate, slightly umbilicate, ventricose, whorls seven in number, concentrically strongly and irregularly striated, more or less crenulated along the upper margin, lip scarcely reflected; olive-brown, sometimes five-banded, rarely pale yellow, lip and columella white.
D'ORBIGNY, Voy. dans l'Amer. Mérid. p. 292. pl. 38. f. 1 to 5.

Hab. Provinces of Yungas, Sicasica, and Ayupaya, Bolivia (found on garden walls and on bushes at an elevation of more than 9000 feet); D'Orbigny, Bridges.

The growth of this species varies, as represented by the specimens figured, from a swollen oblong shape to a more elongated form. Sometimes the shell is banded, and it is always more or less crenulated along the margin next the sutures. The colour is mostly a dull livid olive brown, but the specimen Fig. *c*, collected by Mr. Bridges, is a pale yellow.

Species 87. (Fig. *a* and *b*, Mus. Cuming.)

BULIMUS ROSACEUS. *Bul. testā oblongo-ovatā, crassiusculā, ventricosa, striatā aut sub lente minutiissimā granulatā, spirā obtuso-depressā, aufractibus quinque, per marginem superiorē plus minuscre crenatā, labro latissulo, reflexo; olivaceo-viridi, apice versus rosacē, epidermide tenuissimā sepe aurantiorā, columellā labroque albīdis, interdum rosaceo-aureolatis.*

THE ROSE-COLOURED BULIMUS. Shell oblong ovate, rather thick, ventricose, striated or very minutely granulated under the lens, spire obtusely depressed, whorls five in number, more or less crenulated along the upper margin, lip rather broad, reflected; olive-green, rose-coloured towards the apex, frequently marbled with a very light epidermis; lip and columella white, sometimes rose-orange.

KING, Zool. Journ. vol. v. p. 341.

Varietates.

Bulimus Chilensis, Broderip (not of Lesson).

Bulimus pachycheilus, Pfeiffer.

Hab. Chili (under stones and in the sand); King, Cuming, D'Orbigny, Bolivia (under trunks of trees); Bridges.

The *B. rosaceus* inhabits a wide range of country and is found in very different states. The variety Fig. *a*, which may be regarded as the typical form, inhabits the environs of Valparaiso; that represented at Fig. *b*, distinguished by its rude concentric striae, strong crenulations, and warm orange colouring of the lip and interior, was collected in great plenty and with little or no variation at Huasco, Chili; and at Cocapata, Bolivia, a variety of still larger size occurs of thin light growth with scarcely any indication of crenulations or striae. At Huasco, the species was also collected by Mr. Bridges in a state quite distinct from these, of much smaller dimensions, paler in colour, and of a minutely granulated satiny hue. This is the *B. pachycheilus* of Pfeiffer, and apparently the shell from Bahia Blanca taken by D'Orbigny for the *B. nucleus* with which he does not seem to be acquainted. Another condition of the *B. rosaceus* is that of similar size from Coquiubo, described by Mr. Sowerby as a distinct species under the name *B. chilensis*. It should be observed that the eggs of these smaller species are of a more oblong form than those of the first mentioned.

Species 88. (Mus. Cuming.)

BULIMUS PHASIANELLUS. *Bul. oblongo-ovalā, crassissimā, aufractibus quinque, ventricosa, leviter striatā, per marginem superiorē in aufracta ultimo et penultima crenatā, columellā subiactorā, labro incrassato, reflexo; violaceo-griseā, maculis transversis parvis numerosis irregulāribus violaceo-fuscescentib⁹ pictā, columellā labroque intus violaceis.*

THE PHEASANT-MARKED BULIMUS. Shell oblong-ovate, rather thick, whorls five in number, ventricose, slightly striated, crenulated along the upper margin in the last and penultimate whorls, columella a little twisted, lip thickened, reflected; violet-grey, painted with numerous irregular small transverse blotches of violet-brown, lip and columella deep violet.

BULIMUS.—PLATE XV

VALLSEIENES, *Hedwighi*. Zool. v. 6, p. 244, pl. 55, f. 4.
Bulinus zebra, Sowerby.
Hab. Chile and Peru (on trees). Cuming.

At the Isle of Plata Mr. Cuming found this beautiful and well known species in such abundance upon trees stripped of their leaves, that they looked as if laden with fruit. The animal yields a rich purple juice.

Species 80. (Fig. 4 and 6. Mus. Cuming.)

BULIMUS PORPHYRICUS. *Bol.* testū subcylindrica-ovata, apicē acuminata, infractibus sex concentricis umbilicis et regulariter angustostriatis, per marginem superiore striis minute crenulatis; fauoris-fusca, striis concentricis fere oblique.

THE PORPHYRY BULIMUS. Shell somewhat elongately ovate, spire acuminate, whorls six in number, concretely, rudely, and irregularly elevatedly striated, and minutely crenulated here and there along the upper margin; smoky brown, concentric striae here and there white.

PETTERIK. Zool. Pro. Soc. 1816, p. 114
Hab. ——

The growth of this shell not being completed, it is impossible to say whether the lip would remain simple or be reflected at maturity. It is a very characteristic species and cannot easily be confounded with any other; the white ridge-like striae have a rubbed appearance and being on a dark ground have a very peculiar aspect. Mr. Cuming has no information as to its locality.

Species 90. (Mus. Cuming.)

BULIMUS ZEBRA. *Bol.* testū oratā, tenaculā, imperforatā, ventricosa, spirā acuminata, infractibus subrotundatis heribus, columellā rectā, labro simplici: viridi, flammæ strigisque minoribus, peculiariter undatis flavicarneis uratā, columella alba, labro intus castaneo-nigra.

THE ZEBRA BULIMUS. Shell ovate, rather thin, imperforated, ventricose, spire acuminate, whorls somewhat rounded, smooth, columella straight, lip simple, green, ornamented with peculiarly waved flames and lesser streaks of yellow, columella white, lip edged internally with chestnut.

Buccinum zebra, Muller, Verm. Terr. et Fluv. Hist. p. 138

Bulinus undatus, Bruguière.

Varicataes.

Bulinus zigzag, Lamarek.

Bulinus princeps, Sowerby.

Hab. Nicaragua, Panama, Costa Rica, Veragua, Pernambuco, West Indies.

Few species of the genus affect a wider range of country than the present, and it is owing to the different aspects under which it is found, that the above-recorded synomyms have arisen. The beautiful variety figured in the Conchological Illustrations under the name *B. princeps* was collected by Mr. Cuming in the woods of Nearagua where it inhabits the hollows of trees; the more common variety, known as the *B. undatus*, is found in the West Indies and Brazil, and, extremely rich in colour, at Panama and Costa Rica.



BULIMUS.

PLATE XVI.

Species 91. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS PECULIES. *Bul. testă elongato-ovată, tenui, ventricosă, subumbilicată, spiră acuminată, anfractibus sex aut septem, longitudinaliter striatus, labro tenui, simplici; albida, fusco-ventre lineata, caro-nigricante carū fasciată aut maculata aut punctată, apice peculariter nigricante.*

THE VARI-PICTURED BULIMUS. Shell elongately ovate, thin, ventricose, slightly umbilicated, spine acuminated, whorls six or seven in number, longitudinally striated, lip thin, simple; whitish, stained with pale brown and variously banded, or blotched, or dotted with bluish black, apex peculiarly blackish.

D'ORBIGNY, Voyage dans l'Amer. Mérid. p. 286. pl. 31. f. 1 to 10.

Hab. Bolivian Andes, South America (at the roots of shrubs); D'Orbigny, Bridges.

An extremely variable species of which D'Orbigny has considered it necessary to give ten figures, mainly distinguished by its acuminated growth and general disposition of colour.

Species 92. (Mus. Cuming.)

BULIMUS PUNCTULIFER. *Bul. testă oblongo-ovată, tenuis, umbilicata, anfractibus sex, longitudinaliter striatis, sub lente undique creberrimè pleno-granulosis, per marginem superioreum hic illuc minuti ruditē crenulatis, labro tenui simplici, columellari reflexo; albida, fusco irregulariter lineata et punctata, epidermide floridante fulvata.*

THE DOTTED BULIMUS. Shell oblong-ovate, very thin, umbilicated, whorls six in number, longitudinally striated, under the lens very closely flatly granulated, minutely rudely crenulated here and there along the upper margin, lip thin, simple, the columellar lip reflected back; whitish, irregularly lineated and dotted with brown, covered with a yellowish epidermis.

SOWERBY, Pro. Zool. Soc., 1833, p. 39.

Hab. Chili, La Questa Prada, Cuming; La Questa de Arenos, Huasco, Bridges.

Approaching the *B. granulosus* in form and sculpture, the painting of this shell resembles that of *B. Broderipii*, though easily distinguished by its dotted indefinite style of pattern.

Species 93. (Mus. Cuming.)

BULIMUS RUPICOLUS. *Bul. testă fusiformis-ovata, tenui, spiră subconica, anfractibus quinque ultimo maximum, longitudinaliter ruditē striatis, undique minute granulato-rugosis, labro simplici; albida, maculis longitudinalibus punctisque nigricantibus seriatim picta, epidermide floridante iuncta.*

THE ROCK-INHABITING BULIMUS. Shell fusiformly ovate, thin, spine rather sharp, whorls five in number, the last very large, longitudinally rudely striated and everywhere minutely granulately rough, lip simple; whitish, painted with rows of black longitudinal blotches and dots, covered with a yellowish epidermis. *Succinea variegata*, Pfeiffer, Pro. Zool. Soc., 1842, p. 187. *Hab.* Valleys in the north of Coquimbo (in the crevices of rocks); Bridges.

I think of *Bulimus* to the genus *Succinea*, the type of which is an amphibious animal, like the *Lygaea*, affecting damp and swampy situations, such as the banks of stagnant pools, in marshes, or on floating vegetable matter. The habits of these species are much opposed to this: dwelling in the dry crevices of the rocks which appear in the sandy plains west of the Andes, where there is little or no moisture beyond that which arises from the dews.

The name *variegatus* cannot be adopted, in consequence of its having been used under the head of *Bulimus* by Bruguière in reference to the *Tornatella flava*.

Species 94. (Mus. Cuming.)

BULIMUS THAMNOICUS. *Bul. testă oblongo-ovată, tumidinosa, umbilicata, anfractibus sex, convexa, undique minute seriatim granulata, per marginem superioreum crenulata; fusco-ventre, lineis nigricantibus nebulae irregularibus cingulata.*

THE SHRUB-INHABITING BULIMUS. Shell oblong-ovate, rather swollen, umbilicated, whorls six in number, convex, minutely granulated throughout in rows, crenulated along the upper margin; light brown encircled with irregular clouded blackish lines.

D'ORBIGNY, Voyage dans l'Amer. Mérid. p. 290. pl. 37. f. 4 to 9.

Hab. Eastern side of the Bolivian Andes, (at the roots of bushes and, during the dry season, in the crevices of rocks); D'Orbigny, Bridges.

The specimen selected for representation is the Var. *D. carinatus* of D'Orbigny, the larger variety in which there are none of the transverse lines serving to characterize this, approximate rather to the *B. Tupavii* of the same author.

Species 95. (Mus. Cuming.)

BULIMUS COQUIMBOENSIS. *Bul. testá subfusiforme-oratá, tenuissimá, ventricosá, aufractibus sex, coarctis, subobliquis, longitudinaliter rudité striatis, aufracta ultimam maxima, aperturá subangulá, labro simplici; albida, fuscescente tincta, punctis maculis strigisque nigrovittatis aspersa*

THE COQUIMBO BULIMUS. Shell somewhat fusiformly ovate, very thin, ventricose, whorls six in number, rather oblique, longitudinally rudely striated, last whorl very large, aperture rather large, lip simple; whitish, stained with light brown, and sprinkled with blackish dots, blotches and streaks.

SOWERBY, Pro. Zool. Soc., 1832, p. 30.

Succinea Coquimbensis, Pfeiffer.

Hab. Mountains of Coquimbo; Cuming.

The painting of this species is also not much unlike that of the *B. Broderipii* from which it differs most essentially in form.

Species 96. (Mus. Cuming.)

BULIMUS PIPERATUS. *Bul. testá oratá, tenui, callo ianflatá, spirá breviusculá, acuminatá, aufractibus quinque, ultimo maximo, obscurè rugoso-granulatis et striatis, aperturá perangulá, labro vix reflexo; albida, punctis fuscis perpaucae aspersa, epidermide flaccidá indutá.*

THE PEPPED BULIMUS. Shell ovate, thin, greatly inflated, spire rather short, acuminate, whorls five in number, the last very large, obscurely roughly granulated and striated, aperture very large, lip scarcely reflected; whitish, sprinkled with a yellowish epidermis.

SOWERBY, Conch. Illus. f. 93.

Hab. Hualaga, Peru.

The most inflated of the Succinea-form species, distinguished by a more economical style of painting than its congeners.

Species 97. (Mus. Cuming.)

BULIMUS BRODERIPII. *Bul. testá oratá, abbreviato-fusiforme, tenuissimá, valde inflatá, spirá brevi, aufractibus quinque, ultimo maximo, sub lente minutè granulato-rugosis, aperturá perangulá, labro simplici; albida, nigro fulvoque conspicuè et elegantissimè maculata et variegata.*

BRÖDERIPI'S BULIMUS. Shell ovate, abbreviately fusiform, very thin, greatly inflated, spire short, whorls five in number, the last very large, minutely granulately rugose under the lens, aperture very large, lip simple; whitish, conspicuously and very elegantly blotched and variegated with black and fulvous brown.

SOWERBY, Pro. Zool. Soc., 1832, p. 30.

Succinea Broderipii, Pfeiffer.

Hab. Copiapo, Chili (in the fissures of rocks); Cuming.

Much smaller varieties of this species occur than that selected for figuring and more richly coloured; it is an extremely thin and fragile shell.



BULIMUS.

PLATE XVII.

Species 98. (Mus. Cuming.)

BULIMUS CULMINANS. *Bul. testá oblongo-ovatá, tenuiculá, anfractibus sex, subrotundatis, longitudinaliter creberis et subrude striatis, per marginem superioriem subtiliter crenulatis, labro simplici; flavido-olivaceá, apicea versus cinerascente.*

THE LOFTY-DWELLING BULIMUS. Shell oblong-ovate, rather thin, whorls six in number, somewhat rounded, longitudinally very closely and rather rudely striated, finely crenulated along the upper margin, lip simple; yellowish, faintly olive, pale ash-coloured towards the apex.

D'ORBIGNY, Voy. dans l'Amer. Mérid. p. 289.

Hab. Lake of Titicaca, Bolivia (under stones and at the roots of shrubs upon the mountains); D'Orbigny.

M. D'Orbigny describes this species as inhabiting the most elevated regions of the Bolivian Andes. It is of simple structure and not distinguished by any marked peculiarity of colouring.

Species 99. (Mus. Cuming.)

BULIMUS NUCLEUS. *Bul. testá globoso-ovatá, tenuiculá, ventricosá, vis umbilicatá, sub tente minutissimis granulata, anfractibus quatuor, rotundatis, labro reflexo; tenuo-olivaceá, peritreme rufidé nauratio.*

THE KERNEL BULIMUS. Shell globosely ovate, rather thin, ventricose, scarcely umbilicated, very minutely granulated under the lens, whorls four in number, rounded, lip reflected; yellowish olive, peritreme bright orange.

SOWERBY, Conch. Illus., f. 33.

Hab. Maldanao, vicinity of Monte Video.

I doubt very much if the shell which D'Orbigny refers to this species (Voy. dans l'Amer. Mérid. p. 305) is anything more than a variety of *B. rosaceus*. It will be seen by the accompanying figure how much this shell resembles that species, though readily distinguished by its light globose structure and bright orange peritreme.

Species 100. (Fig. a, b, c. Mus. Cuming.)

BULIMUS SORDIDUS. *Bul. testá ovatá, ventricosá, ampliter umbilicatá, spirá breviusculá, acutá, anfractibus sex, convexis, ultimo maximo, inflato, crebellinæ gra-*

nulatis, labro tenui latè effuso; sordidus albæ, aurantio-castaneo plus sinuoso tintet et fasciat.

THE DULL BULIMUS. Shell ovate, ventricose, largely umbilicated, spire rather short, acute, whorls six in number, the last very large and inflated, very closely granulated, lip thin, widely effused; dull white, more or less stained or variegated with orange chestnut.

LESSON. Zool. Voy. Coquille, pl. 13. f. 3.

Bulimus Protens and mutabilis, Broderip.

Bulimus Cora, D' Orbigny.

Hab. Mountains of Peru (under stones); Cuming.

The name given to this species by Mr. Broderip is a far more appropriate one on account of its Protean aspect and pretty varieties of painting, than that to which it is entitled. First collected by M. Lesson during the voyage of the Coquille, it is to Mr. Cuming we are mainly indebted for the beautiful varieties obtained in different parts of Peru, St. Jacinta, Pacosmayo, Truxillo, Santos, &c., out of which Mr. Broderip proposed a second species under the name *B. mutabilis*. M. D'Orbigny collected the *B. sordidus* on the arid ridges of the mountain of San-Cristobal near Lima, and I believe his *B. Cora* from the interior of Peru, of which Mr. Cuming possesses authentic specimens, to be only another state of the same species.

Species 101. (Fig. a, b, c. Mus. Cuming.)

BULIMUS PERUVIANUS. *Bul. testá oblongo-ovatá, tenui, imperforatá, spirá subacutá, anfractibus quaque longitudinaliter rude et irregulariter rugoso-striatis, interdum transversum obscurè punctato-suturatis, labro simplici, tenuissimo; grisœ aut flavidæ aut fuscenscente, sepe roseo et carmineante variegatæ.*

THE PERUVIAN BULIMUS. Shell oblong-ovate, thin, imperforated, spire rather sharp, whorls five in number, longitudinally rudely and irregularly rugosely striated, sometimes encircled with obscure punctured grooves, lip simple, very thin; greyish or pale yellow or light brown, frequently variegated with pink or blue.

BRUGUIERE. Dict. Enc. Méth. no. 37.

Bulimus Gravesii and corrugatus, King.

Hab. Chili (at the roots of shrubs); Cuming.

There is not much variation in the form of this species, but it differs considerably in colour. The animal, unlike

BULIMUS.—PLATE XVII.

most of its congeners, brings forth its young alive, and during the dry season incloses itself within the shell by forming an epiphragm.

Species 102. (Mus. Cuming.)

BULIMUS CHILENSIS. *Bul. testa oblongo-ovata, tenui,
subligeriter cerebellinæ granulatâ, spirâ subacutâ, aprofundata,
tibus sex longitudinaliter rufè striatis, labro levissimo,
simplici, fuscocerato, strigis sordide castaneis longitudinaliter
dispositis hic illuc truetâ.*

THE CHILI BULIMUS. Shell oblong-ovate, thin, finely very closely granulated, spire rather sharp, whorls


six in number, longitudinally rudely striated, lip very thin, simple; light brown, stained here and there with longitudinal dull chesnut streaks.

LESSON, Voy. de la Coquille, pl. 8, f. 3.

Bulimus granulosus, Broderip.

Hab. Valparaiso and Conception, Chili; Cuming.

Although this shell is exactly similar in structure to that of the preceding species, the animal is not viviparous: it is, however, singularly distinguished by the smallness of its eggs, which are like little peas. It inhabits the mountains of Conception, and is found in ravines near the sea-shore as far as Valparaiso. During the dry season it buries itself in the earth, inclosed by an epiphragm. The shell exhibits little variation of form or colouring.



BULIMUS.

PLATE XVIII.

Species 103. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS NITIDUS. *Bul. testā pyramidali-oratā, tenui,
subpellucido, anfractiōes sex, convexis, longitudinaliter
fortiter elevato-striatis, aperturā pareincolū, labro
simplici; opalescente-luteo, strigis cerasulente-fusci-
longitudinalibus ornata, apice nigricante.*

THE SHINING BULIMUS. Shell pyramidal ovate, thin, somewhat transparent, whorls six in number, convex, longitudinally strongly elevatedly striated, aperture rather small, lip simple; opalescent cream-colour, ornamented with longitudinal bluish-brown streaks, apex blackish.

BRODERIP, Pro. Zool. Soc., 1832, f. 31.

Ejusdem var. laeteca.

Bulimus cacticornis, Broderip.

Hab. Columbia and Peru; Cuming.

Of a delicate opalescent cream-colour with a black apex, and longitudinally streaked with blue-brown, a neat finely corded appearance being imparted to the shell in consequence of the striae which cross the dark streaks remaining cream-colour. The *B. cacticornis* of Broderip is merely a variety having no streaks.

Species 104. (Fig. *a*, *b*, *c*. Mus. Cuming.)

BULIMUS VARIANS. *Bul. testā elongatā, tenui, umbilicatā,
anfractiōes septem, convexis, longitudinaliter hic illuc
rude stratiis, aperturā parē, labro simplici, albō aut
rubellā, costaneo sepiissimè aut zonulatā aut longitudi-
nialiter variegatā et strigatā.*

THE VARIABLE BULIMUS. Shell elongated, thin, umbilicated, whorls seven in number, convex, longitudinally here and there rudely striated, aperture small, lip simple; white or reddish, most frequently zoned or longitudinally streaked and variegated with chesnut.

BRODERIP, Pro. Zool. Soc., 1832, p. 106.

Ejusdem varietates.

B. pulchellus and *rubellus*.

Hab. On the mountains about Truxillo, Peru; Cuming.

The present is a remarkable instance of the variation of colour and pattern in one species in the same locality. The variety Fig. *a*, (*B. rubellus*, Brod.) is of a simple reddish nankeen colour; Fig. *b*, (*B. pulchellus*, Brod.) is white, reddish towards the apex, encircled by a regular

series of dark brown ~~zona~~ and Fig. *c*, (*B. variegatus*, Brod.) is of the same ground tint as Fig. *a*, with the addition of numerous irregular mottled longitudinal purple-brown streaks. All are characterized by a dark band in the umbilicus which is rather large.

Species 105. (Mus. Cuming.)

BULIMUS LEMNISCATUS. *Bul. testā ovoatā, ventricosā ant-
plūde umbilicatā, spirā subpyramidalis, acutā, anfracti-
bus septem, rotundatis, levibus, labro simplici, albīdā,
vittis fusci pseudoridere interimplis circumdatā, colum-
nellā et apertura flave roseo.*

THE RIBANDED BULIMUS. Shell ovate, largely umbilicated, spire somewhat pyramidal, sharp, whorls seven in number, rounded, smooth, lip simple; whitish, surrounded by peculiarly interrupted brown fillets, columella and interior of the aperture pink.

DESHAYES, Anim. sans vert. vol. viii. p. 271.

Bulimus vittatus, Broderip (not of Spix).

Hab. Ilo, Peru; Cuming.

The brown bands or fillets which characterize this species are broken up in a peculiar manner into square blotches divided in part by a light transverse dash of white; the columella and interior are pink.

M. D'Orbigny appears to have confounded this and the *B. Henndeli* together; his Fig. 8 (Voy. Amer. Mérid. Pl. 30.) is a variety of the former species, whilst his Fig. 3 and 4 are of the latter.

Species 106. (Mus. Cuming.)

BULIMUS FUSOIDES. *Bul. testā elongatā, basi effusa, an-
fractiōes sex, levibus, nitidis, columellā oblique contortā,
labro via reflexo; albīdā, strigis cerasulente-fusci
angustis undatis longitudinaliter ornata, columellā in-
tensè roseo-purpurea, labro albo.*

THE FUSOID BULIMUS. Shell elongated, effused at the base, whorls six in number, smooth, shining, columella obliquely twisted, lip scarcely reflected; whitish, ornamented longitudinally with narrow waved streaks of bluish-brown, columella deep rose-purple, lip white.

D'ORBIGNY, Voyage dans l'Amer. Mérid. p. 215.

Hab. Cacopata, Bolivia (in the woods); Bridges.

Of this very characteristic and pretty species, collected to perfection by Mr. Bridges, M. D'Orbigny found but two dead specimens in a damp ravine near the banks of the Rio San Mateo, Cochabamba, Bolivia, on the North-East side of the Eastern Cordilleras. It is chiefly remarkable on account of the patulate effused formation of the aperture and deep rose-purple stained columella.

Species 107. (Mus. Cuming.)

BULIMUS TIGRIS. *Bul. testā cyprinaceo-angustā, ampliatae undulatā, unfractibus septem, lacinibus aut subtiliter striatis, latera cincta, latus subtilē expanso, exterior sanguineus, subeffusus, albida, purpureo-fuscoseculo latè obscurata.*

THE TIGER BULIMUS. Shell cylindrically elongated, largely undulated, whorls seven in number, smooth or finely striated, exemplar lip widely expanded, outer lip simple, somewhat effused; white, broadly streaked with purple-brown.

BRODERIP, Proc. Zool. Soc., 1832, p. 107.

Hab. Truxuy, Peru (on bushes on the mountains); Cuming.

Though approximating very closely to the *B. carinus*, there is a sufficient variation in the form of this species to render it distinct: the body-whorl is of greater length and the aperture more effused.

Species 108. (Mus. Cuming.)

BULIMUS ONCA. *Bul. testē conigalo-oblongā, levū, subangustā, aperturā granulata, ultimo maxima, obliquis, longitudinaliter striatis, per marginem superiore leviter exangulatis, latera cincta, cinctores fuligineos, guttis sublineares, per dicitus asperis, columellā et aperture fuscō-rubore.*

THE SPOTTED BULIMUS. Shell elongately oblong, thin, colour brownish, whorls five in number, the last two larger, oblique, longitudinally striated, slightly rounded above the upper margin, lip scarcely retracted, whitish, fulvous colour, sprinkled here and

there with large chestnut dots; columella and interior of the aperture pink.

D'ORBIIGNY, Voyage dans l'Amer. Mérid. p. 295. pl. 30. f. 1 and 2.

Hab. Tutuilima, Bolivia.

It is entirely owing to the enterprising spirit of M. Alcide D'Orbigny that we are indebted for the knowledge of this beautiful species. "Desiring to find a more immediate communication," says that intelligent traveller, "between Cochabamba and Moxas, Bolivia, we made a proposition to the Government of that Republic to make the attempt. The President, General Santa Cruz, furnished us with the means to achieve this undertaking. We set out on foot with a party of Indians to carry provisions, and after passing forty days amid the precipices of uninhabited mountains on the north-east side of the Eastern Cordilleras, and enduring a thousand difficulties and privations we accomplished our journey. It was in this excursion not far from Tutuilima that we observed the *Bulimus onca* at the bottom of almost inaccessible ravines in damp woody situations, where it is of rare occurrence and difficult to procure, from its habit of burying itself in the earth."

Species 109. (Mus. Cuming.)

BULIMUS MODESTUS. *Bul. testē pyramidali, turreta, ampliatae undulatā, unfractibus septem, angustis, rotundatis, fortiter et erberriacē concentricē elevato-striatis, lobis simpliciē fuscoseculo, apicea versus albida, striis concentricis obtusis.*

THE MODEST BULIMUS. Shell pyramidal, turreted, largely undulated, whorls seven in number, narrow, rounded, strongly and very closely concentrically elevatedly striated; light brown, whitish towards the apex stria whitish.

BRODERIP, Proc. Zool. Soc., 1832, p. 106.

Hab. Huacho, Peru, (on small bushes on the mountains); Cuming.

The whorls of this species have rather a tubercular style of growth.



Fig. 2.



Fig. 3.



Fig. 4.



Fig. 5.



Fig. 6.



Fig. 7.



Fig. 8.



Fig. 9.



Fig. 10.



Fig. 11.

BULIMUS.

PLATE XIX.

Species 110. (Mus. Cuming.)

BULIMUS SCUTULATUS. *Bul. testā pyramidali, turrītā, umbilicatā, aufractibus octo, rotundatis, concentricē rudo striatis, aperturā pareinclusā, labro simplici; albida, castaneo plus minusve undique fasciata.*

THE COBWEB-MARKED BULIMUS. Shell pyramidal, turrid, umbilicated, whorls eight in number, rounded, concentrically rudely striated, aperture rather small, lip simple; whitish, more or less banded throughout with chestnut.

Broderip, Pro. Zool. Soc., 1832, p. 106.

Hab. Islay, Port Aréquipa, Peru, on the hills, Cuming; on dry rocks near the shore, D'Orbigny.

M. D'Orbigny speaks of this species as existing in situations where there is not the slightest vegetation or moisture even from the dews.

Species 111. (Mus. Cuming.)

BULIMUS TUMIDULUS. *Bul. testā oblongo-ovatā, tenui, subumbilicatā, spirā breviusculā, acentrā, aufractibus septem, saperāe plano-convergis, infernū peculiariter rotundato-effusis, labro simplici; albida, flavo-fuscescente tinctā, lînis castaneis fuscisqne singulariter nubatis circinatā, columellā roseo-purpureā, apice purpureo-fusco.*

THE SWOLLEN BULIMUS. Shell oblong-ovate, thin, slightly umbilicated, spire rather short, acute, whorls seven in number, flatly concave round the upper part, peculiarly rotundately effused round the lower, lip simple; whitish, stained with yellowish brown, encircled with chestnut lines and singularly shaded bands, columnella rose-purple, apex purple-brown.

Pfeiffer, MSS., Mus. Cuming.

Bulinus inflatus, Broderip (not of Lamarck), Pro. Zool. Soc., 1832, p. 45.

Hab. Ambo, Peru; Cuming.

An extremely pretty shell differing both in its growth and texture from most of the Peruvian species.

Species 112. (Mus. Cuming.)

BULIMUS ELEGANS. *Bul. testā ovatā, tenuissimā, pellucida, inflata, aufractibus quinque, concaisis, ultimo*

maximo, labro simplici; rubello-cornēd, strigis albis opacis livis que nigris undatis longitudinaliter pictā.

THE ELEGANT BULIMUS. Shell ovate, very thin, transparent, inflated, whorls five in number, convex, the last very large, lip simple; reddish horny colour, longitudinally painted with waved black lines and opaque white streaks.

Succinea elegans, Pfeiffer, Pro. Zool. Soc., 1842, p. 187.

Hab. La Questa de Arenas, Ilustre, Chili (found on a small cryptogamic plant); Bridges.

An extremely delicate shell of a thin fragile Succinea-like texture, strikingly painted with opaque white streaks and black lines.

Species 113. (Mus. Cuming.)

BULIMUS VERSICOLOR. *Bul. testā pyramidali-ovatā, ampliè umbilicatā, aufractibus sex, convergis, minutè depresso-granulatis, labro tenui, sublate reflexo: castanā, nuculis albido variegatā,*

THE CHANGEABLE-COLOURED BULIMUS. Shell pyramidal-oval, largely umbilicated, whorls six in number, convex, minutely depresso granulated, lip thin, rather widely reflexed; chestnut, variegated with whitish spots.

Broderip, Pro. Zool. Soc., 1832, p. 108.

Hab. Mongon near Casma, Peru (found on bushes on the mountains); Cuming.

Although this shell varies in its shade and pattern of colour, the specimens all partake more or less of the same red-brown tint which characterizes the *B. Proteus*. In form and sculpture the variation is trifling, and I doubt very much if it should not also be arranged, together with *B. mutabilis*, as a variety of that species.

Species 114. (Mus. Cuming.)

BULIMUS MONTE-VIDENSIS. *Bul. testā pyramidali-ovatā, tenui, pellucida, umbilicatā, aufractibus septem, concaisis, obliquè irregulariter striatis, labro simplici; pallide straminea, cornēd, strigis opaco-albīs.*

THE MONTE-VIDEO BULIMUS. Shell pyramidal-oval, thin, pellucid, umbilicated; whorls seven in number, convex, obliquely irregularly striated, lip simple; pale straw colour, horny, streaks opaque-white.

May, 1848.

PFEIFFER, Pro. Zool. Soc., 1846, p. 93.

Hab. Monte Video, Buenos Ayres (on small trees in swampy places); Cuming.

A very delicate transparent species in which the whorls are concentrically marked with fine opaque-white striae.

Species 115. (Mus. Cuming.)

BULIMUS COTURNIX. *Bul. testā glōbo-ovalā, tenui, in-*
flatā, rīz umbilicatā, anfractibus quinque, ultimo mar-
teo; labro simplici; fluorescente-albā, maculis grandi-
tus unicoloris undulato-custanicis fuscā.

THE QUAIL BULIMUS. Shell globosely ovate, thin, inflated, scarcely umbilicated, whorls five in number, the last very large, lip simple; brownish-white, stained with numerous large blotches of burnt-chestnut.

SOWERBY, Pro. Zool. Soc., 1832, p. 33.

Hab. Huasco, Chili (on dry stones); Cuming, D'Orbigny.

This is certainly not a variety of *B. Broderipii* as M. D'Orbigny intimates, it is, with little or no variation, of a more globose form and another style of painting.

Species 116. (Mus. Cuming.)

BULIMUS DRAPARNAUDI. *Bul. testā subpyramidalē-ovatā,*
anfractibus septem convexis, labro simplici; albā,
strigis latiscutis cœruleis, roseis, et opaco-albīs, lineis
nigra interdau subtiliter undulatis interspersis, pulcher-
rimē longitudinaliter pictā.

DRAPARNAUD'S BULIMUS. Shell somewhat pyramidalily ovate, whorls seven in number, convex, lip simple;

white, very beautifully painted longitudinally with rather broad streaks of blue, pink, and opaque white, interspersed with black lines which are sometimes finely waved.

PFEIFFER, Pro. Zool. Soc., 1846, p. 113.

Hab. Chilon, Bolivia (on the trunks of *Cacti*); Bridges.

This is truly a gem, exhibiting the gayest display of colours of any of the ectivorous species.

Species 117. (Mus. Cuming.)

BULIMUS BRIDGESII. *Bul. testā ovalā, umbilicatā, pel-*
lucidā, apice subobtusa, anfractibus quatuor, striis con-
fertissimis et subtilissimè decussatis, aperturā parrius-
cūlā, labro latè reflexo, frē integro; serico-flavescente,
perileucate albo.

BRIDGES' BULIMUS. Shell ovate, umbilicated, pellucid, rather obtuse at the apex, whorls four in number, very closely and finely decussated with striae, aperture rather small, lip broadly reflected, almost entire; silken yellow, peristome white.

PFEIFFER, Pro. Zool. Soc., 1842, p. 186.

Hab. Frierina, Huaseo, Chili (under stones); Bridges.

A delicate transparent shell, resembling *B. rosaceus* in form, but certainly distinct from that species. The whorls are not crenulated along the upper edge, nor is there the least tinge of rose-colouring; it has more the transparency of a silkworm cocoon.



123



124



125



126



127



128



129



130



131



132

BULIMUS.

PLATE XX.

Species 118. (Mus. Cuming.)

BULIMUS RUGIFERUS. *Bul. testā pyramidali-turritā, umbilicatā, aufractibus octo, transversa tenuissimè striatā, longitudinaliter peculiariter rugosā, aperturā parvā, labro simplici; intus extusque flichti-brunneā.*

THE WRINKLED BULIMUS. Shell pyramidal turreted, umbilicated, whorls eight in number, transversely very finely striated, longitudinally peculiarly wrinkled, aperture small, lip simple; earthy brown within and without.

SOWERBY, Pro. Zool. Soc., 1833, p. 36.

Hab. Jacob Island, Gallapagos; Cuming.

Curiously longitudinally wrinkled, whilst the interstices are finely transversely striated.

Species 119. (Mus. Cuming.)

BULIMUS LAURENTII. *Bul. testā acuminato-octoū, sub-aupitē umbilicatā, aufractibus quinque, subrotundis, lœvatis, aperturā suborbiculari, labro simplici; albida, rubido-fusco concinna fasciatā.*

THE ST. LAWRENCE BULIMUS. Shell acuminately ovate, rather largely umbilicated, whorls five in number, rather rounded, smooth, aperture somewhat orbicular, lip simple; whitish, prettily banded with reddish brown.

SOWERBY, Pro. Zool. Soc., 1833, p. 37.

Hab. Island of San Lorenzo, Bay of Callao, Peru (on stones at the summits of the mountains); Cuming.

An interesting small species existing in great abundance in the localities it inhabits.

Species 120. (Mus. Cuming.)

BULIMUS PRUINOSUS. *Bul. testā oblongo-ovata, tenui, pellucida, subaupitē umbilicatā, aufractibus quinque, lœvatis, apice subpapillari, labro simplici; corneo-albida, opaco-albo varia.*

THE SNOWY BULIMUS. Shell oblong-ovate, thin, pellucid, rather largely umbilicated, whorls five in number, smooth, apex somewhat papillary, lip simple; horny white, variegated with opaque white.

SOWERBY, Pro. Zool. Soc., 1833, p. 36.

Hab. Peru; Cuming.

A light transparent horny shell, delicately variegated with irregular opaque white spots and streaks.

Species 121. (Mus. Cuming.)

BULIMUS ESCHARITERUS. *Bul. testā sublongato-turrīta, umbilicatā, spira solaris impressa, aufractibus septem, depresso-concavis, undique subtiliter seriatim granulatis, aperturā subquadrate-ovali, labro vir reflexo; ustulata-fusca, fascia angustis rufo-fuscis medio cingulata.*

THE TOASTED BULIMUS. Shell somewhat elongately turreted, umbilicated, sutures of the spire impressed, whorls seven in number, depresso convex, finely granulated throughout in rows, aperture somewhat squarely ovate, lip scarcely reflected; scorched-brown, encircled round the middle with two narrow red-brown bands.

SOWERBY, Conch. Illus. Bul. f. 87.

Hab. Chatham Island, Gallapagos; Darwin.

The granulations of this species are somewhat different from those of the *B. Proleus*; they are more isolated and more distinctly set in transverse rows.

Species 122. (Mus. Cuming.)

BULIMUS RIVASI. *Bul. testā acuminato-oblonga, tenui, aufractibus octo, concaesis, aperturā parvissimā, labro simplici; opaco-alba, strigis lineolisque longitudinalibus varieis pellucidis.*

RIVAS' BULIMUS. Shell acuminately oblong, thin, whorls eight in number, convex, aperture rather small, lip simple; opaque-white, with longitudinal transparent horny streaks and lines.

D'ORBIGNY, Voyage dans l'Amér. Mérid. p. 276. pl. 34. f. 8 to 10.

Hab. near Chilon, Bolivia (under dead trunks of *Cueti*); Bridges.

I have not much doubt that this shell is the *B. Rivasi* of D'Orbigny, but it is proper to mention that I have no other authority for it than that obtained by a comparison with the description and figure.

Species 123. (Mus. Cuming.)

BULIMUS RUGELLOSUS. *Bul. testā cylindraceo-elongata, sub-Papiformi, umbilicata, apice subpapillari, aufractibus octo, leviter concentricè rugulosis, columella supra umbilicata reflexa, aperturā parvā, labro vir dilatato; olivaceo-fusca, fascia unica pallidā medio cingulata.*

THE SLIGHTLY WRINKLED BULIMUS. Shell cylindrically elongated, somewhat Pupa-shaped, umbilicated, apex subpapillary, whorls eight in number, slightly concentrically wrinkled, columella reflected over the umbilicus, aperture small, lip very slightly dilated; olive-brown, encircled round the middle with a single pale band.

SOWERBY, Conch. Illus. *Bul.* £ 85.

Hab. Gallapagos Islands; Cuming.

Of a peculiar red-tinged olive brown-colour and chrysanthemum-like form.

Species 124. (Mus. Cuming.)

BULIMUS TURBITUS. *Bul. testā elongato-turritā, imperforatā, spirae suturis impressis, anfractibus decem ad undevit, laevibus, subdepresso-concisis, aperturā pareā, labro simplici; albida, rufo-fluorescente trifasciata.*

THE TURRETED BULIMUS. Shell elongately turreted, imperforated, sutures of the spire impressed, whorls ten to eleven in number, smooth, depresso convex, aperture small, lip simple; whitish, encircled with four reddish-brown bands.

BRODERIP, Pro. Zool. Soc., 1832, p. 106.

Hab. Truxillo, Peru (on the mountains); Cuming.

The uppermost of the bands is next the suture of each whorl, the lowest is only shown in the last whorl.

Species 125. (Mus. Cuming.)

BULIMUS SCULPTRATUS. *Bul. testā oralā, lenientiā, vix umbilicatā, anfractibus septem, subcentrōsis, conspicue reticulatis rugatis, interstītīis transversim corrugatis, columellā simplicē reflectā, labro tenui; intus subsope fimbriis.*

THE SCULPTURED BULIMUS. Shell ovate, rather thin, scarcely umbilicated, whorls seven in number, somewhat ventricose, conspicuously reticulately wrinkled, the interstices being corrugately striated, columella simply reflected, lip thin; earthy brown within and without.

FILIBER, Pro. Zool. Soc., 1846, p. 29.

Hab. Gallapagos Islands (on bushes); Darwin.

The sculpture of this species is precisely that already described in the *B. rugiferus* and *Darwinii*, from both of which it differs in form, and from the latter more especially by the absence of any callous tubercles in the aperture.

Species 126. (Mus. Cuming.)

BULIMUS CALVUS. *Bul. testā oblongā, subturrīta, umbilicatā, anfractibus septem rotundatis, levibus ant minūte concentricē striatis, aperturā pareā, labro cœ dilatato; pallente brunnēa, ad apicem nigricante.*

THE BALD BULIMUS. Shell oblong, somewhat turreted, umbilicated, whorls seven in number, rounded, smooth or minutely concentrically striated, aperture small, lip slightly dilated; pale brown, blackish at the base.

SOWERBY, Pro. Zool. Soc., 1833, p. 72.

Hab. James' Island, Gallapagos (on dried tufts of grass); Cuming.

A smooth pale brown shell having somewhat an *Achatinella*-like appearance.

Species 127. (Mus. Cuming.)

BULIMUS PUSTULOSUS. *Bul. testā subfusiforme-oralā, ampliter umbilicatā, anfractibus sex, subrugosis, striis cœndibis granulatis, moniliformibus, longitudinaliter exsculptis, columellā supra umbilicūm reflectā, labro simplici; intus extusque sordidè albā.*

THE PUSTULOSE BULIMUS. Shell somewhat fusiformly ovate, largely umbilicated, whorls six in number, rather rough, longitudinally engraved with rude granulated necklace-like striae, columella reflected over the umbilicus, lip simple; dead white within and without.

BRODERIP, Pro. Zool. Soc., 1832, p. 105.

Hab. Huasco, Chili (under stones on elevated ground); Cuming.

The necklace-like granulated striae with which this species is longitudinally sculptured are extremely characteristic.



B U L I M U S.

PLATE XXI.

Species 128. (Mus. Cuming.)

BULIMUS DECOLORATUS. *Bul. testā oblongo-ovatā, anfractibus quinque, medio subangulato-ventricosis, infra suturas peculiariter longitudinaliter crenulatis, labro simplici; albida, fuscescente hic illie tenueri strigata, parte circa columellam castaneo-fusca.*

THE DISCOLOURED BULIMUS. Shell oblong-ovate, whorls five in number, somewhat angularly ventricose in the middle, peculiarly longitudinally crenulated beneath the sutures, lip simple; whitish, finely streaked here and there with light brown, chesnut brown round the columella.

SOWERBY, Pro. Zool. Soc., 1833, p. 73.

Hab. Vicinity of Lima, Peru (found buried in the earth under bushes on the hills); Cuming.

The first two whorls of this shell are characterized by a very minute, shagreen appearance, discernible only with the lens, but it may be doubted whether it is of any specific value.

Species 129. (Fig. a and b. Mus. Cuming.)

BULIMUS SCALARIFORMIS. *Bul. testā pyramidalis-oblonga, amplius umbilicata, anfractibus sex, rotundatis, crenulariē concentricē tenuicostatis, aperturā parvā, labro simplici; fuscescente, costis albidis.*

THE SCALARIA-SHAPED BULIMUS. Shell pyramidalily oblong, largely umbilicated, whorls six in number, round, very closely concentrically finely ribbed, aperture small, lip simple; light brown, ribs whitish.

BRODERIP, Pro. Zool. Soc., 1832, p. 31.

Hab. Ancon, Peru; Cuming. Vicinity of Lima; D'Orbigny.

An extremely interesting species of which two specimens are represented, illustrative of its variation of growth. M. D'Orbigny relates having found it in great abundance among the ruins of a temple erected by the ancient Incas in honour of their invisible god, Pachacamac.

Species 130. (Mus. Cuming.)

BULIMUS USTULATUS. *Bul. testā oblongo-ovatā, subamplicher umbilicata, anfractibus septem, convexis, lœvibus, aperturā parvā, labro simplici; ustulato-fusca, spiruliter varia linearū.*

THE SCORCHED BULIMUS. Shell oblong-ovate, rather

largely umbilicated, whorls six in number, somewhat rounded, concentrically rudely striated, lip simple; burnt brown, variously spirally lined.

SOWERBY, Pro. Zool. Soc., 1833, p. 72.

Hab. Charles' Island, Galapagos (under detached pieces of lava); Cuming.

The lined pattern of this species is somewhat variable, but it is always characterized by the same scorched brown hue.

Species 131. (Mus. Cuming.)

BULIMUS MERIDIONALIS. *Bul. testā cylindraceo-ovatā, temniciā, rix umbilicata, anfractibus sex, pecuniariter sebarsco-incisis, prope suturas irregulariter linearisaleatis, columellā angulariter contortā, aperturā oblongā, leviter compressā; sordidè albā, apertura fovea intensè purpureo-rufā.*

THE SOUTHERN BULIMUS. Shell cylindrically ovate, rather thin, scarcely umbilicated, whorls six in number, peculiarly sebarsco engraved, irregularly linearly grooved near the sutures; columella angularly twisted, aperture oblong, slightly compressed; dead white, interior of the aperture rich purple red.

Hab. Chili; Bridges.

A very singular species, which appears to have escaped the vigilant search of both Cuming and D'Orbigny. It is very unlike any other Chilian species. Of a cylindrically oblong form, its dead white surface is curiously engraved with obtuse or obsolete scales, and the interior is stained with a deep purple-red colour. The columella has a curious angular prominence, which does not, however, form an inner fold as in the Auricula-like species.

Species 132. (Mus. Cuming.)

BULIMUS BILINEATUS. *Bul. testā pyramidalis-oblonga, umbilicata, anfractibus septem, convexis, lœvibus, aperturā parvā, labro simplici; fuscescente-albe, lineis dubiis iridescentibus cingulata.*

THE TWO-LINED BULIMUS. Shell pyramidalily oblong, umbilicated, whorls seven in number, convex, smooth, aperture small, lip simple; brownish-white, encircled by two reddish lines.

SOWERBY, Pro. Zool. Soc., 1833, p. 37.

Hab. St. Elmo and Isle of Pinta, West Columbia (under stones, and buried at the roots of bushes); Cuming.
Extremely simple in form and colouring.

Species 133. (Mus. Cuming.)

BULIMUS UNICOLOR. *Bul. testā oblongā, subpyramidalē, umbilicatā, anfractibus sex, subrotundis, concentricis et minutissimè ruditè striatis, labro simplici; cornuā, petiolaridē.*

THE UNICOLOURED BULIMUS. Shell oblong, somewhat pyramidal, umbilicated, whorls six in number, somewhat rounded, concentrically very minutely rudely striated, lip simple; horny, transparent.

SOWERBY, Pro. Zool. Soc., 1833, p. 73.

Hab. Island of Perico, Bay of Panama (on dead leaves); Cuming.

Very similar in form and general character to the *B. transversus* and *Panamensis*, and found in nearly the same locality.

Species 134. (Mus. Cuming.)

BULIMUS CORNEUS. *Bul. testā ovatā, subcentricosā, umbilicatā, anfractibus quinque ad sex, rotundatis, lobi- bus, aut minutissimè striatis, labro simplici; cornuā, petiolaridē.*

THE HORNED BULIMUS. Shell ovate, somewhat ventricose, umbilicated, whorls five to six in number, rounded, smooth or very minutely striated, lip simple; horny, transparent.

SOWERBY, Pro. Zool. Soc., 1833, p. 37.

Hab. Real Llegos, Central America (under decayed grass); Cuming.

A simple transparent horny shell, the whorls being somewhat rounded.

Species 135. (Mus. Cuming.)

BULIMUS JACOBI. *Bul. testā ovatā, subacuminatā, aperturā umbilicatā, anfractibus subrotundis, lobi- bus, columellā supra umbilicū partim expansā, aperturā parvissimā, labro simplici; cornueo-fuscescente.*

THE JACOB BULIMUS. Shell ovate, somewhat acuminate, largely umbilicated, whorls rather rounded, smooth, columella partially expanded over the umbilicus, aperture rather small, lip simple; pale horny brown.

SOWERBY, Pro. Zool. Soc., 1833, p. 74.

Hab. Jacob Island, Gallapagos; Cuming.

Chiefly distinguished from the last species by its large umbilicus, which is not seen in the figure on account of its being partially overlaid by the columellar lip.

Species 136. (Mus. Cuming.)


BULIMUS DARWINII. *Bul. testā subgloboso-ovatā, tenui- catā, inflatā, umbilicatā, anfractibus sex, supernè sub- angulatis, longitudinaliter conspicue rugatis, interstictis corrugato-striatis, columellā subtorsatā, tuberculis duobus cattosis proximatebus armatā, aperturā subquadrate- ovali, labro simplici; intus extusque fictili-brunnea.*

DARWIN'S BULIMUS. Shell somewhat globosely ovate, rather thin, inflated, umbilicated, whorls six in number, slightly angulated round the upper part, longitudinally conspicuously wrinkled, the interstices being transversely corrugately striated, columella slightly twisted, and armed with two prominent callosus tubercles, aperture somewhat squarely oval, lip simple, earthy brown within and without.

PREIFFER, Pro. Zool. Soc., 1846, p. 29.

Hab. Galapagos Islands (on bushes); Darwin.

It will be observed on comparing this shell with the *B. rugiferus* and *sculpturatus*, that a uniform peculiarity of wrinkled sculpture prevails in each species; the present is chiefly distinguished by its globose form and by the calllosities on the columella.



B U L I M U S.

PLATE XXII.

Species 137. (Mus. Cuming.)

BULIMUS CONSPERSUS. *Bul. testá ovatá, tenui, sordide pellucidá, subaustrilicatá, anfractibus quinque, rotundatis, concentricè obsolete ruditè striatis, ad suturas crenulatis, labro simplici; flavante-corneá, lineis guttulisque opaco-albis profusè conspersá, peritremate pallide rosacea.*

THE BESPRINKLED BULIMUS. Shell ovate, thin, semi-transparent, slightly umbilicated, whorls five in number, rounded, concentrically obsoletely rudely striated, crenulated at the sutures, lip simple; yellowish horny colour, profusely sprinkled with opaque white lines and dots, peritreme tinged with pink.

SOWERBY, Pro. Zool. Soc., 1833, p. 73.

Hab. Lima (buried in the earth under bushes on the hills); Cuming.

A beautifully delicate horny species, crowded with effective opaque-white lines and dots, with a pale rose peritreme to the aperture.

Species 138. (Mus. Cuming.)

BULIMUS AQUILUS. *Bul. testá ovatá, ventricosá, umbilicatá, anfractibus quinque, longitudinaliter creberissimè et subtiliter striatis, interstítios reticulato-punctatis, labro simplici; pellucido-castaneo alboque longitudinaliter undatá et caregatá.*

TRE SWARTHY BULIMUS. Shell ovate, ventricose, umbilicated, whorls five in number, longitudinally very closely and finely striated, the interstices being reticulately punctured, lip simple; longitudinally waved and variegated with transparent chestnut and white,

Hab. Tacna, Peru.

A prettily marked species approaching somewhat in character to the *B. versicolor*, but differing materially in form.

Species 139. (Mus. Cuming.)

BULIMUS STRIATUS. *Bul. testá subpyramidalis-ovatá, tenuicatá, subampliter umbilicatá, anfractibus sex, usque concentricè elevato-striatis, columella sublatè expansa, labro simplici; fasciente-alba, lineis strigis concentricis castaneis irregulariter notatá.*

THE STRIATED BULIMUS. Shell somewhat pyramidal, ovate, rather thin, somewhat largely umbilicated, whorls six in number, concentrically elevatedly striated throughout, columella rather widely expanded, lip simple; brownish white, irregularly marked with concentric chestnut-brown lines and streaks.

KING, Zool. Journal.

Hab. Lima (upon the hills); Cuming.

Peculiarly barred here and there in the most irregular manner with dark chestnut brown.

Species 140. (Mus. Cuming.)

BULIMUS EROSUS. *Bul. testá ovatá, subaustriliter umbilicatá, spirá breviusculá, acuminatá, anfractibus sex, ventricosis, longitudinaliter subrugoso-striatis; diaphano fuscenscente, strigis maculatisque opaco-albis aspersa, apice pallide rosacea.*

THE ERODED BULIMUS. Shell ovate, somewhat largely umbilicated, spire rather short, acuminated, whorls six in number, ventricose, longitudinally somewhat roughly striated; light diaphanous brown, sprinkled with opaque white streaks and spots, apex pale pink.

BRODERIP, Pro. Zool. Soc., 1832, p. 106.

Hab. Huantajaya, near Iquique, Peru (under stones on the hills, 2,500 feet above the level of the sea); Cuming.

The detail of character is not much unlike that of *B. conspersus*, but it is a shell of larger size and much more solid growth.

Species 141. (Mus. Cuming.)

BULIMUS ALBICANS. *Bul. testá ocatá, umbilicatá, anfractibus quinque, ventricosis, subrugoso-striatis, columella subexpansá, opaco-albicante, hic illuc diaphaná, apertura favea fulvescente.*

THE WHITE BULIMUS. Shell ovate, umbilicated, whorls five in number, ventricose, somewhat roughly striated, columella rather expanded; opaque-white, here and there diaphanous, interior of the aperture fulvous.

BRODERIP, Pro. Zool. Soc., 1832, p. 105.

Hab. Copiapo, Chili (in the dry sand on elevated ground near the Port); Cuming.

This species partakes in great measure of the characters

of *B. conspersus* and *erosus*, but is, I believe, sufficiently distinct.

Species 142 (Mus. Cuming.)

BULIMUS APODEMETES. *Bul. testā ovalā, tenui, inflatā, virū umbilicatā, anfractibus quinque, subrugoso-striatis, ultimo calde maximo, aperturā subampullā, labro tenui; albidā.*

THE EXILE BULIMUS. Shell ovate, thin, inflated, scarcely umbilicated, whorls five in number, rather roughly striated, last whorl very much the largest, aperture somewhat large, lip thin; whitish.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 279.

Hab. Bolivia, &c.; D'Orbigny.

I must refer the reader for a better illustration of this species to the above-mentioned magnificent work by M. D'Orbigny, the present uncoloured specimen giving no other idea of it than that of form. In reference to colour M. D'Orbigny says, "the tints are extremely variable according to the localities; at Parana the shell is pale-yellow or dotted, often marked with transverse lines of marbled white and clear brown, with indications of distant clear brown longitudinal lines; at Chiquitos, on the other hand, it is distinguished by longitudinal brown bands only, which are then more conspicuous."

Species 143. (Fig. a and b. Mus. Cuming.)

BULIMUS STRIATUS. *Bul. testā pyramidali-ovalā, tenui, diaphanā, conspicue umbilicatā, anfractibus rotundatis, pulcherrimè creberrimè concentricè elevato-striatis, columellā reflexā, labro simplici; pallidè cornē, translucida, striis opaco-albis, fasciis plurimiis albidiis interrumpitis cingulata.*

THE FINELY STRIATED BULIMUS. Shell pyramidal, thin, diaphanous, conspicuously umbilicated, whorls rounded, very beautifully closely concentrically elevatedly striated, columella reflected, lip simple; pale horny colour, transparent, the striae opaque-white, encircled with numerous interrupted whitish bands.

SOWERBY, Pro. Zool. Soc., 1833, p. 73.

Bulimus D'Orbignii, Pfeiffer,

Hab. Lima (under stones on the hills); Cuming.

A beautiful species, distinguished by the delicate aspect of its fine opaque-white striae upon a transparent horny ground.

Fig. a is a representation of the specimen upon which Mr. Sowerby founded the species, and Fig. b is the shell described by Dr. Pfeiffer under the name *B. D'Orbignii*.

Species 144. (Mus. Cuming.)

BULIMUS GUTTATUS. *Bul. testā pyramidalis-oblongā, subfusiformi, umbilicatā, apice papillari, anfractibus sex, longitudinaliter rugoso-striatis, ad suturas crenulatis: fulvo-fuscenscente, guttis apaco-albis longitudinaliter notatā, apice rosaceo-fuscā.*

THE MOTTLED BULIMUS. Shell pyramidal-oblong, somewhat fusiform, umbilicated, apex papillary, whorls six in number, longitudinally rugosely striated, crenulated at the sutures; pale fulvous-brown, longitudinally marked with opaque-white speckles; apex pinkish-brown.

BRODERIP, Pro. Zool. Soc., 1832, p. 31.

Hab. Cobija; Cuming.

This species like others of the plate is characterized by a sprinkling of opaque-white spots, and it is distinguished by its papillary apex.



138



139



140



141



142



143



144



145



146



147

B U L I M U S.

PLATE XXIII.

Species 145. (Mus. Cuming.)

BULIMUS DISCREPANS. *Bul. testā acuminato-oblongā, tenui, anfractibus sex, lœvibus, aut sub leute minutissimè transversim impresso-striatis, labro simplici; lacteā, lineis fuscis subdistans obliquè pictā.*

THE DIFFERENT BULIMUS. Shell acuminate oblong, thin, whorls six in number, smooth, or, under the lens, very minutely transversely impressively striated, lip simple; cream colour, obliquely painted with rather distant brown lines.

SOWERBY, Pro. Zool. Soc., 1833, p. 72.

Hab. Couchagua, Central America (found under the bark of a tree); Cuming.

Distinguished by its painting of delicate brown lines, which are placed, for the most part, at equal distances from each other.

Species 146. (Mus. Cuming.)

BULIMUS MONTAGNEI. *Bul. testā acuminato-oblongā, tenuiculā, vis nubilatā, anfractibus septem, levibus, subrotundis, labro simplici; albida, strigis punctisque cervuleo-nigricandibus longitudinaliter pictā, apertura fauce lutescente.*

MONTAGNE'S BULIMUS. Shell acuminate oblong, rather thin, scarcely umbilicated, whorls seven in number, smooth, somewhat rounded, lip simple; whitish, painted longitudinally with bluish black dots and streaks, interior of the aperture yellowish.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 286. pl. 32. f. 7.

Hab. Chilon, Bolivia (under dead trunks of *Cacti*); Bridges.

I take this shell to be the *B. Montagnei* of D'Orbigny from its very accurate resemblance with his Fig. 7 of that species.

Species 147. (Mus. Cuming.)

BULIMUS HUASCENSIS. *Bul. testā acuminato-oblongā, subampliē umbilicatā, anfractibus sex, rotundatis, concentricè irregulariter rugoso-striatis, suturis subprofundis, labro simplici; cerulescente-albā.*

THE HUASCO BULIMUS. Shell acuminate oblong, rather

largely umbilicated, whorls six in number, rounded, concentrically, irregularly, roughly striated, sutures rather deep, lip simple; bluish-white.

Hab. Huasco, Chili; Bridges.

A small species collected by Mr. Bridges which is not exactly referable to any of those discovered by Mr. Cuming or M. D'Orbigny.

Species 148. (Mus. Cuming.)

BULIMUS RUBESCENS. *Bul. testā subacuminato-ovali, umbilicatā, anfractibus sex, rotundatis, concentricè rugoso-striatis, labro simplici; albida, lineis fusciscentibus longitudinaliter irregulariter notatā, columellā et operculari fauce purpureo-roseis.*

THE BLUSHING BULIMUS. Shell somewhat acuminate ovate, umbilicated, whorls six in number, rounded, concentrically roughly striated, lip simple; whitish, irregularly marked with light brown longitudinal lines, columella and interior of the aperture purple-rose.

Hab. South America.

A characteristic new species, distinguished by the delicate purple-rose bloom which lines the interior.

Species 149. (Mus. Cuming.)

BULIMUS UNIFASCIATUS. *Bul. testā subpyramidalē-ovata, tenui, cornēa, umbilicatā, anfractibus sex, longitudinaliter subtiliter striatis, labro simplici; luteo-fuscescente, fasciā angustā, unicā, stramineā medio cingulatā.*

THE ONE-BANDED BULIMUS. Shell somewhat pyramidal ovate, thin, horny, umbilicated, whorls six in number, longitudinally finely striated, lip simple; yellowish-brown, encircled round the middle by a single narrow straw-coloured band.

SOWERBY, Pro. Zool. Soc., 1833, p. 37.

Hab. Charles's Island, Galapagos (found under detached pieces of lava); Cuming.

A semi-transparent horny shell, mainly distinguished by a narrow pale straw-coloured band by which it is encircled throughout.

Species 150. (Mus. Cuming.)

BULIMUS SUX. *Bal. testā pyramidalē, umbilicatā, anfractibus septem, concentricis, rotundatis, subfuscis, aperturā subangustiori, pallidissime effusa; olivaceofusca, rufosubrosea, necto, apice erigente, apertura fuscō-erubescens-alba, prope marginam fuscā.*

THE SUX BULIMUS. Shell pyramidal ovate, umbilicated, whorls seven in number, concentrically roughly striated in a wrinkled manner, aperture somewhat squarely ovate, slightly effused; olive-brown, stained with rusty red, blackish at the apex, interior of the aperture bluish-white, brown near the edge.

Broderip, Pro. Zool. Soc., 1832, p. 125.

Hab. Charles's Island, Galapagos (on bushes); Cuming.

The aperture of this species is frequently compressed at the sides so as to give it a somewhat square aspect.

Species 151. (Mus. Cuming.)

BULIMUS DERELICTUS. *Bal. testā acuminato-ovatā, angulatā, umbilicatā, anfractibus sex, subrotundatis, subtilissimā granulo-striatis, columellā labro expansis, spire papillari; cornu-albidā, opaco-albo tunc illuc emporso.*

THE FORSAKEN BULIMUS. Shell acuminate ovate, largely umbilicated, whorls six in number, somewhat rounded, very finely granosely striated, lip and columella expanded, apex papillary; flesh-white, sprinkled here and there with opaque-white.

Broderip, Pro. Zool. Soc., 1832, p. 107.

Hab. Cobija; Cuming.

The name attached to this shell is intended to denote the solitary habitat in which it was found by Mr. Cuming, on flat rocks without soil or verdure, all being desolate for a league around. M. D'Orbigny notices having collected it of a punky hue, at a greater elevation upon some lichens and cactuses.

Species 152. (Mus. Cuming.)

RICUS VEXILLUM. *Bal. testā oblongo-ovatā, cornuā, subpellucida, anfractibus quinque, longitudo-nitibus rugoso-striatis, labro simplici; pallide cornē, maculis strigisque undatis opaco-albis profuse notata, apice roseā.*

THE RICUS BULIMUS. Shell oblong-ovate, horny, semi-transparent, whorls five in number, smooth, shining; yellowish-white, encircled by numerous chestnut bands.

Broderip, Pro. Zool. Soc., 1832, p. 105.

Hab. King's and Saboga Islands, Bay of Panama (on the trunks of large trees); Cuming.

This comparatively well-known species has, I believe, been called *B. altervatus* by Dr. Beck, in consequence of Mr. Broderip's name having been formerly applied by Bruguière to the shell commonly known as *Achatina exilis*. M. Deshayes does not, however, think there is any necessity for this change, and lest it might add to the confusion I have avoided it.

Species 153. (Mus. Cuming.)

BULIMUS PESSULATUS. *Bal. testā ovalā, tenuī, inflatā, anfractibus quinque, rotundatis, ventricosis, lacertos, labro simplici; pellucido-cornē, epiderme ferrugineo-albo opaco, castaneo longitudinaliter strigatā, strigis distantibus, conspicuis.*

THE CROSS-BARRED BULIMUS. Shell ovate, thin, inflated, whorls five in number, rounded, ventricose, smooth, lip simple; transparent horny, covered by a rusty white opaque epidermis, crossed with conspicuous distant longitudinal streaks.

Hab. Santa Cruz de la Sierra, Bolivia (on bushes and high grass); Bridges.

This has much the appearance of being an immature shell, but the specimens collected by Mr. Bridges are all alike, and are without doubt of adult growth.

Species 154. (Mus. Cuming.)

BULIMUS AFFINIS. *Bal. testā elongato-ovalā, subfusiforme, umbilicatā, anfractibus septem, convexis, longitudinaliter rugoso-striatis, labro simplici; pallide cornē, maculis strigisque undatis opaco-albis profuse notata, apice roseā.*

THE ALLIED BULIMUS. Shell elongately ovate, somewhat fusiform, umbilicated, whorls seven in number convex, longitudinally roughly striated, lip simple; pale horny, profusely marked with opaque-white waved streaks and spots, pink at the apex.

Broderip, Pro. Zool. Soc., 1832, p. 105.

Hab. Mexillones, desert of Atacama, Peru (in small crevices of the dry earth at an elevation of 2,000 feet above the sea); Cuming.

A light transparent shell covered with delicate flakes of opaque-white.



B U L I M U S.

PLATE XXIV.

Species 155. (Mus. Cuming.)

BULIMUS COLORATUS. *Bul. testá ovalá, ventricosissimá, vix umbilicatá, spirá brevi, anfractibus quinque, tumido-concexis, pulcherrimè minuti promiscuó granulatis, ad suturas subtilissimè crenulatis, columellá plicá unicá ascende, aperturá subamplá, labris lamíná callosa junctis, peritremate conspicuè reflexo; albida, flammis luteis longitudinalibus, maculis strigise nigricantibus umbratis, epidermide olivacea indutá, apicem versus castaneá, columellá labroque roseis.*

THE COLOURED BULIMUS. Shell ovate, very ventricose, scarcely umbilicated, spire short, whorls five in number, tumidly convex, very beautifully minutely promiscuously granulated, very finely crenulated at the sutures, columella with a single ascending fold, aperture rather large, outer and columellar lips connected by a callous lamina on the body whorl, peritreme conspicuously reflected; whitish, with longitudinal yellow flames shaded with blackish spots or streaks, covered by an olive epidermis, chestnut-coloured towards the apex, lip and columella deep rose.

NYST, Bull. de l'Acad. Royale de Bruxelles, vol. xii. 1^{re} partie, pl. I. f. 2 a, b.

Hab. Province of Cumana, Colombia; Funek.

A truly interesting species collected by M. Funek, the South American traveller, in the interior of Colombia, and first described by M. Nyst in the Transactions of the Royal Academy of Brussels, from a specimen of much younger and more pelleted growth than those before me. The most characteristic features of the species are its beautifully granulated surface and richly shadowed undulated painting, whilst the lip is of a deep rose colour, and the interior lined with a delicate pinkish bloom, through which the outer pattern is discernible.

Species 156. (Mus. Cuming.)

BULIMUS LAMARCKIANUS. *Bul. testá ovalá, ventricosissimá, vix umbilicatá, spirá brevi, anfractibus quinque, subplano-concexis, minuti promiscuó obtuso-granulatis, ad suturas subtilissimè crenulatis, columellá plicá unicá ascende, aperturá subamplá, peritremate conspicuè reflexo; pallide livido-fuscescente, punctis grandibus castaneis hic illuc piperatá, epidermide olivacea indutá, peritremate livido-fusco.*

LAMARCK'S BULIMUS. Shell ovate, very ventricose, scarcely umbilicated, spire short, whorls five in number, rather flatly convex, minutely promiscuously obtusely granulated, very finely crenulated at the sutures, columella with a single ascending fold, aperture rather large, peritreme conspicuously reflected; pale livid brown, peppered here and there with large chestnut dots, covered with an olive-brown epidermis, peritreme livid brown.

PFEIFFER, Monog. *Bul.*

Hab. Andes of New Granada (at an elevation of about 8,000 feet); Funek.

So closely does this species approximate to the preceding, that I feel somewhat doubtful of the propriety of retaining it; they appear to differ, so far as may be judged by a comparison of the specimens here figured, but it is easy to conceive that a larger series would reveal a much closer affinity between them.

Species 157. (Mus. Dennison.)

BULIMUS PARDALIS. *Bul. testá oblongo-ovali, subventricosa, crassiuscula, anfractibus quinque, suboblique convoluta, apicem versus minutissimè creberrimè obtuso-granulatis, per suturas undique longitudinaliter crenulatis, columellá contortá, fortiter uniplicatá, labro conspicuè reflexo; purpurascente-fusca, longitudinaliter albiflammata, epidermide lutescente-olivacea indutá, columellá et peritremate albis.*

THE PANTHER BULIMUS. Shell oblong-oval, rather ventricose, somewhat thick, whorls five in number, rather obliquely convoluted, very minutely and closely obtusely granulated towards the apex, longitudinally crenulated throughout along the sutures, columella twisted, with a strong winding plait, lip conspicuously reflected; purplish-brown, longitudinally flamed with white, and covered by a yellowish olive epidermis.

DE FERUSSAC, Hist. Moll. terr. et fluv. pl. 112. f. 7. 8.

Bulimus venezuelensis, Nyst.

Bulimus atrapoides, Jonas.

Hab. Cumana, Venezuela; Funek. Varinas, Venezuela; Dyson.

It is somewhat difficult to recognise this species by the figure of M. De Féussac, it being only known to the illustrious author of the 'Histoire des Coquilles Terrestres'

in a smaller state deprived of epidermis. The specimen here represented, is from the choice collection of Mr. Dennison, and Mr. Cuming possesses it almost equally fine, selected from the stores of Mr. Dyson obtained during a recent expedition to Venezuela.

The journey performed by Mr. Dyson in this fertile district of South America, in search of objects of natural history, was undertaken through the generosity of a limited number of zoological friends, who, with the hope of enriching their collections, were at the same time desirous of encouraging the zeal of a young naturalist distinguished by his love of adventure. Though not perhaps a very profitable excursion, he collected many interesting species, and has no doubt gratefully responded to the patronage of his friends.

Species 155. (Mus. Cuming.)

BULIMUS THOMPSONI. *Bul. testá subelongato-oblongá, imperforatá, apicē obtusā, anfractibus sex, longitudina-
liter striatis, per suturas plano-crenulatis, columellá
rectá vel paululum arcuatá, labro vix reflexo; livido-
cinerá, epidermide cornéa, nítidá, olivaceo-viridi
indutá, apice refescente, anfractuum marginē superiori
crenulato albante, aperture fauce crenulente, juxta
peritreme purpureo-nigricante, labris laminá nigricante-
tividá callosá junctis.*

THOMPSON'S BULIMUS. Shell somewhat elongately oblong, imperforated, obtuse at the apex, whorls six in number, longitudinally striated, flatly crenulated along the sutures, columella straight or a little curved, lip but slightly reflected; livid ash colour, covered with a shining horny olive-green epidermis, reddish towards the apex, upper crenulated edge of the whorls whitish, interior of the aperture bluish, purple-black towards the peritreme, lips joined by a callous blackish livid lamina.

PFEIFFER, Pro Zool. Soc., 1845, p. 74.
Hab. Andes of Quito.

A very distinct species remarkable for the same shining livery green epidermis which characterizes the *Helix Bushii*. (Couch, Syst. vol. ii. pl. 164. fig. 11.)

Species 159. (Mus. Cuming.)

BULIMUS SHONGI. *Bul. testá pyramidati-oblongá, crassá,
imperforatá, anfractibus sex, levibus, per suturas sub-
crenulatis, labro reflexo; fluorescente, epidermide cornéa
rufo-castaneá indutá, anfractuum marginē superiori*

*albicante, columellá et aperture fauce intenſe aurantio-
rufis, peritremate albante.*

SHONG'S BULIMUS. Shell pyramidal oblong, thick, imperforated, whorls six in number, smooth, slightly crenulated along the sutures, lip reflected; light brown, covered with a horny reddish chestnut epidermis, upper edge of the whorls whitish, columella and interior of the aperture deep orange red, peritreme whitish.

LESSON, Voy. de la Coquille, Moll. p. 321. pl. 7. f. 4, 5.
Hab. New Zealand.

A shell of not uncommon occurrence, always distinguished by the deep orange red colouring of the aperture and rich chestnut epidermis, whilst there is no plait on the columella.

It was first discovered by M. Lesson, during the Voyage of the Coquille, at the Bay of Islands, New Zealand, and named after one of the native Chiefs of that district. Subsequently M. Lesson thought to have recognised his species in Chemnitz's figure, Conchyl. Cab. vol. ix. pl. 121. f. 1039-40, under the title *Helix auris-Mida* and in De Féüssac's figure of the *B. auris-bovinus*, Hist. Moll. terr. et fluv. pl. 159. f. 1, 2, but it is perfectly distinct from these, which (*cide* Pl. XXX.) are also distinct from each other, and the honoured name of the New Zealand Chief remains to posterity.

Species 160. (Mus. Cuming.)

BULIMUS FULMINANS. *Bul. testá oblongá, apicem versus
peculiariter acuminatá, anfractibus quinque, oblique
convolutis, undique longitudinaliter et reticulatè cor-
regatis, columellá fortiter uniplicatá labro talè reflexo;
fumefuscá, subpellucídá, strigis longitudinalibus acutè
angulatis intenſe castaneis conspicuè ornatá.*

THE LIGHTNING BULIMUS. Shell oblong, peculiarly acuminate towards the apex, whorls five in number, obliquely convoluted, longitudinally and reticulately wrinkled throughout, columella strongly one-plaited, lip widely reflected; smoky brown, somewhat transparent, conspicuously ornamented with sharply angled, dark chestnut longitudinal streaks.

NYST, Bull. de l'Acad. Royale de Bruxelles.

Bulimus bellulus, Jonas.

Bulimus Loceeni, Pfeiffer.

Hab. Colonia de Tovar, Venezuela; Dyson.

A somewhat rare species rather sharply acuminate towards the apex, of a peculiar semitransparent smoky brown colour, marked with sharply zigzag longitudinal chestnut streaks.



BULIMUS.

PLATE XXV.

Species 161. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS BLAINVILLEANUS. *Bul. testā ovalā, spirā brevissimā, anfractibus quinque, inflatis, suboblique convolutis, undique pulcherrimā longitudinaliter et reticulatē corrugatis, columellā fortiter uniplicatā, labo latē reflexo; fuscō-fuscā, strigis castaneis acute angulatis longitudinaliter pictā, epidermide subtilissimā corrugatā viridi luteo-punctulatā indutā.*

DE BLAINVILLE'S BULIMUS. Shell ovate, spire very short, whorls five in number, inflated, somewhat obliquely convoluted, very beautifully longitudinally and reticulately wrinkled throughout, columella strongly one-plaited, lip widely reflected; smoky-brown, longitudinally painted with sharply zigzag chestnut streaks, covered with a very finely wrinkled epidermis, of a green colour sometimes dotted with yellow.

PFEIFFER, Monog. *Bul.*

Hab. Zaji, Province of Merida, New Granada; Funck, Linden.

This very beautiful species has precisely the sculpture, colour, and marking of the *B. fuluinanus*, overlaid by a remarkable epidermis of a dark green colour, sometimes minutely yellow-dotted, deposited in fine wrinkles resembling those which may frequently be observed on oil-painting that has been exposed to the heat of the sun. It differs in form, being shorter and of a more inflated growth, but is a curious approximation to it in other respects.

Species 162. (Mus. Cuming.)

BULIMUS MORITZIANUS. *Bul. testā oblongo-ovalā, ventricosā, anfractibus sex, convexis, levibus, ad suturas exiliter crenulatis, columellā fortiter uniplicatā, labo reflexo; fuscā, lutescente-albo scēp̄ longitudinaliter undato-striatā, epidermide olivaceo-viridi, nunc levigatā, unne corrugatā, indutā, columellā labroque interdū viridē rufo-aureantia, interdū fuscō-fuscā.*

MORITZ'S BULIMUS. Shell oblong-ovate, ventricose, whorls six in number, convex, smooth, faintly crenulated at the sutures, columella strongly one-plaited, lip reflected; brown, often longitudinally streaked with yellowish white in zigzag pattern,

covered with an olive green epidermis, sometimes smooth, sometimes wrinkled, lip and columella bright reddish-orange or smoky brown.

PFEIFFER, Monog. *Bul.*

Hab. Chachopo, Province of Merida, New Granada (in the woods); Funck, Linden.

A fine new species of which Mr. Cuming possesses several examples varying in colour and painting between those selected for illustration. The rich dark green epidermis and deep orange lip and columella present an effective contrast.

Species 163. (Mus. Cuming.)

BULIMUS CALEDONICUS. *Bul. testā subfusiforme-oblongā, crassissimā, anfractibus sex, subdepressis, lœvibus, columellā valde incrassatā, medio depresso, deinde productā et tenuitā, parte superiori tuberculo calloso munitā, apertura auriculatā, contractā, labo incrassato, non reflexo, superne late sinuato; fuscescente-albā, epidermide castaneā nitidā indutā, columellā labroque albidis, apertura fave intenſe sanguineo-rufā.*

THE NEW CALEDONIA BULIMUS. Shell somewhat fusiformly oblong, very thick, whorls six in number, rather depressed, smooth, columella very much thickened, depressed in the middle, then produced and swollen, upper part furnished with a callous tubercle, aperture auriculated, contracted, lip thickened, not reflected, broadly sinuated above; brownish-white, covered with a shining chestnut epidermis, lip and columella whitish, interior deep blood red.

PETIT, Magasin de Zoologie.

Hab. New Caledonia.

A remarkable species, chiefly distinguished by the curious auriculated structure of the aperture; there is no plait on the columella, but a prominent callous wart or tubercle. The lip is not reflected but pressed over the aperture, and towards the upper part is characterized by a deep sinus. Were this singularity of growth not proved to be constant in the species, it might have been regarded as an unnatural distortion. We are indebted to M. Petit, a zealous conchologist of Paris, for having described this species, and I have to thank him for sending me a specimen which accords in all respects with those in Mr. Cuming's collection.

Species 164. (Mus. Cuming.)

BULIMUS AURIS-SILENI. *Bul.* testē ovato-oblongā, apicem versus abbreviato-tumidā, anfractibus quatuor, corrugato-striatis, subrotundatis, obliquē convolutis, columellā fortissimā uniplicatā, plicis valde productā, lobis latē reflexo : suturā, striis nigricante-viridibus longitudinaliter undatā, epidermide luteo-olivaceā indutā, columellā tubroque albī.

THE SILENUS' EAR *BULIMUS*. Shell ovately oblong, abbreviately swollen towards the apex, whorls four in number, striated in a wrinkled manner, somewhat rounded, obliquely convoluted, columella very strongly plaited, extremely produced, lip widely reflexed, fulvous, longitudinally waved with blackish green stripes, and covered with a yellowish-olive epidermis, lip and columella white.

BRUGUIERE, Enc. Méth. vol. i. p. 345.

Auris Sileni, Born.

Helix Auris-Silenei, De Féüssac.

Annicula caprella, Lamarck.

Carychium undulatum, Leach.

Plekocheilus undulatus, Goulding.

Hab. —?

This is without doubt the original *Auris Sileni* of the Vienna Museum, of which a characteristic representation is given by Born and Chemnitz, and not the shell described by Lamarck under the name *Annicula Sileni*. It is a short thick shell of tumid growth, distinguished by a remarkably prominent plait on the columella and bold oval aperture with rudely reflected lip.



BULIMUS.

PLATE XXVI.

Species 165. (Mus. White.)

BULIMUS ADAMSONI. *Bal. testá subfusiformi-oratá, ventri-*
cosá, spírá acuminatá, pyramidali, aufractiōis sex ad
septem, levibus vel obtigné tenuistrati; columellá sub-
recta, aperturá paulatim effusa, labro vix refexo;
violetaceo-cinerá, basi versus subolivaceá, fuso hic
ilic sparsu nubilatá, vittá conspicuá carvescente-
alba, olivaceo-nigrante subroseo articulata cingulatá;
ritá alterá politid obcurá supra, columellá purpureo-
nigrante, peritreme violato, intus carvescente-albo.

ADAMSON'S BULIMUS. Shell somewhat fusiform oval, ventricose, spire acuminate, pyramidal, whorls six to seven in number, smooth or obliquely finely striated; columella nearly straight, aperture a little effused, lip scarcely reflected; violet ash colour, olive-tinted towards the base, sparingly clouded here and there with brown; encircled by a conspicuous blue-white fillet, rather remotely articulated with olive-black, there being another pale obscure fillet above, columella purple-black, peritreme violet, bluish white within.

Hab. Base of the Purremo, near the volcano of Tolyma, on the eastern slope of the Andes.

This magnificent species was purchased many years since, with the above locality, by John Adamson, Esq., of Newcastle-on-Tyne, and no other specimen has since been heard of. The collection of that gentleman having been recently dispersed, it has fallen into the possession of another eminent amateur conchologist, Henry Hopley White, Esq., through whose kindness and liberality I am enabled to publish it after a long period of obscurity.

It is of an elegant pyramidal fusiform growth, rather ventricose and a little effused towards the base, of a rich olivaceous violet colour, distinguished by a conspicuous white belt remotely articulated with purple-black, and the peritreme of the aperture is of a rich violet.

Species 166. (Mus. Dennison.)

BULIMUS DENNISONI. *Bal. testá acuminato-oratá, ven-*
tricosá, aufructibus quinque ad sex, supernē subconcurvo-
depressa, superficie striis rugosis modique subtilissimè

decussatá, aufractu ultimo peramplo, inflato, columella
subrecta, apertura oblonga, labro vix refexo: supra
ceruleo-albicans, nigricante-virido variegata et punctata,
infra latro-olivacea, panicis strigisque brevibus
undatis nigricante-viridiobus, lateo in uno latere illumi-
natus, ptilcherioe picti, zonis albidis cingulata, quarum
inferiori subobscurá, peritremate fuscó, intus ceruleo-
albicans.

DENNISON'S BULIMUS. Shell acuminately ovate, ventricose, whorls five to six in number, slightly concavely depressed round the upper part, the entire surface being very finely decussated with rough striae, reticulated, last whorl very large, inflated; columella nearly straight, aperture oblong, lip scarcely reflected; bluish white above, variegated and dotted with blackish green, beneath yellowish olive, very beautifully painted with dots and short-waved streaks of blackish green, illuminated along one side with yellow, and encircled by two whitish zones of which the lower is somewhat obscure, peritreme brown, inferior bluish white.

Hab. —?

Another fine species, from probably the same fertile region, of which, I believe, only two specimens are known: one, here figured, in the collection of J. Dennison, Esq.; the other, very much worn and discoloured, in the British Museum. It approximates closely to the preceding species in general appearance, but differs materially in its details. In the *B. Adamsoni* the whorls form a nearly straight pyramidal cone, in the *B. Dennisoni* they are more ventricose and less pyramidal disposed, whilst the last is proportionably larger, more oblong, and more inflated. The texture of the two shells is dissimilar; the former is smooth or obliquely striated and appears to have a thin horny stratum of semi-transparent epidermis, the latter is finely decussated with rough uneven striae in a manner which is rarely if ever observed beneath a horny epidermis. The difference of colouring is best described in the accompanying drawings; they are two elegantly painted shells and, as species quite unapproached by any hitherto described, constitute a most interesting and valuable addition to the genus.



B U L I M U S.

PLATE XXVII.

Species 167. (Fig. 167 *a*, Mus. Dennison. Fig. 167 *b*,
Mus. Cuming.)

BULIMUS POWISIANUS. *Bul. testā acuminato-oblongā,
crassā, apice obtusā, aufractibus levibus, nitidis, vix
striatis; columellā subarcuatā, aperturā pareiusculā,
paulatim effusā, labro incrassato, vix reflexo; futes-
cente-spadicōcā, strigis irregularris obliquis, nigricante-
viridis, cingulo unico in aufractu ultimo, conspicu-
picta, interdum innuculatā, cingulo tantum, columellā
labroque intensa purpureo-violaceis, peritremate nigri-
cante, apice rufo-castaneo.*

POWIS' BULIMUS. Shell acuminate oblong, thick, ob-
tuse at the apex, whorls smooth, shining, scarcely
striated, columnella slightly arched, aperture rather
small, a little effused, lip thickened, but slightly
reflected; fulvous bay, conspicuously painted with
irregular, oblique, blackish-green streaks, and a single
belt round the last whorl, sometimes unspotted, with
the belt only, lip, and columnella deep purple violet,
peritreme blackish, apex reddish chestnut.

PETIT, Magasin de Zool.

Hab. Valle de Canea, New Granada

This very beautiful and rare species approximates in a
manner to the *B. regina* which follows; it is of a less
acuminated form, of extremely solid structure, and, ex-
cepting the lip and columnella, the pattern of colouring is
dissimilar. Of the specimens here represented, that from
the collection of Mr. Dennison is distinguished by its rich
painting of irregular black-green stripes upon a fulvous
bay ground; in Mr. Cuming's specimen the ground colour
is unspotted, but in both examples the last whorl is en-
circled by a characteristic narrow belt of the same dark
colour as the stripes.

Species 168. (Fig. *a*, *b*, *c*. Mus. Cuming.)

BULIMUS REGINA. *Bul. testā subelongato-oblongā, sarpē
sinistrali, spirā pyramidalis-acuminatā, apice obtusā;
aufractibus septem ad octo, levibus vel obliquē tenui-
striatis, columellā subcontortā, labro simplici; futes-
cente-albā, aufractibus superne fusco viridiisque variegatis
et articulatis, infernē rufescensibus aut purpureo-nigri-
cantis, columellā et apertura fauce interdum albis,
interdum intense violaceis, nigro marginatis.*

THE QUEEN BULIMUS. Shell somewhat elongately oblong,
often sinistral, spire pyramidaly acuminate, obtuse
at the apex; whorls seven to eight in number, smooth
or obliquely finely striated, columella slightly twisted,
lip simple; fulvous white, whorls variegated and
articulated at the upper part with brown or green,
greenish, or purple-black at the lower, aperture and
columella sometimes white, sometimes deep violet,
edged with black.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 257. pl. 29.
f. 4, 5.

Helix regina, De Férussac.

Achatina perversa, Swainson.

Achatina melanostoma, Swainson.

Eadem? *Bulinus phlegrus*, D'Orbigny.

Hab. Fig. 168 *a*. Interior of Peru; Poeppig. Fig. 168 *b*.
Chopo, Province of Pauplona, New Granada; Funck,
Linden. Fig. 168 *c*. British Guyana, Brazils. Bo-
livia; D'Orbigny.

It will be seen by the above-mentioned localities that the *Bulinus regina* inhabits a wide range of country, and, by the figures, that it is found under very different aspects, having in some localities a dextral, in others a sinistral growth.

I quite agree with M. D'Orbigny in the propriety of referring this species to the genus under consideration rather than to *Achatina*; the columnella is never strictly truncated, and the shell has all the texture, form, and colouring of a *Bulimus*.

Figure 90 *b*. (Mus. Dennison.)

BULIMUS ZEBRA. This species is also found under so
many different aspects, that I thought it desirable to figure
a singular variety from the collection of Mr. Dennison, of
which that gentleman possesses several examples exactly
similar, brought by some vessel into Liverpool, but he
is not aware from whence. It is of a remarkably solid
growth, heavy as porcelain, and has no indication of the
green which predominates more or less in most varieties.
The upper part of the whorls is white, the lower encircled
by a broad band of light rust colour, and towards the
aperture the whorl is crossed by numerous concentric dark
streaks.



B U L I M U S.

PLATE XXVIII.

Species 160. (Fig. 160 b. Mus. Cuming.)

BULIMUS FULMINANS. The shell here represented is that in Mr. Cuming's collection, described by Dr. Pfeiffer under the name of *B. Loveti*, Pro. Zool. Soc., 1847, p. 229. It is without doubt identical with the *B. fulminans* described at Pl. XXIV., though lighter in structure and of less acuminate growth.

Species 169. (Mus. Cuming.)

BULIMUS MILLERI. *Bul. testā ovato-oblongā, spirā sub-oblītā, atractibūs quinque, tumidisculis, oblique convolutis, aperturā subcontractā, labro reflexo; castaneofuscā, epidermide sericeā, transversim minutè granulosocrispatā, indutā, peritremate roseo.*

MILLER'S BULIMUS. Shell ovately oblong, spire rather obtuse, whorls five in number, rather swollen, obliquely convoluted, aperture somewhat contracted, lip reflected; chestnut-brown, covered with a silken epidermis, which is transversely minutely granulously crisped, peritreme rose-coloured.

SOWERBY, Conch. Illus. f. 94.

Hab. Brazil.

This beautiful species is mainly distinguished by its exquisite silken epidermis, which exhibits two or three bands towards the base, not of colour, but of light and shade, as in a shot-silk.

Species 170. (Mus. Cuming.)

BULIMUS AURIS-MID.E. *Bul. testā oblongo-vel elongato-ovalā, subcompressā, atractibūs sex, levibus aut irregulariter rugosō-striatis, ultimo calde majorē, columellā callōsa pliciformi, interdum duabus, nūridā, aperturā amplā; fulvescente-ablō, epidermide olivaceo-castaneā indutā, apertura fauce rufescente-croceā, peritremate lacteā.*

THE MIDAS' EAR BULIMUS. Shell oblong or elongately ovate, a little compressed, whorls six in number, smooth or irregularly roughly striated, last whorl very much the larger, columella furnished with a plait-like callousity, sometimes two, aperture large; fulvous white, covered with an olive chesnut epidermis, interior of the aperture reddish chesnut, peritreme cream colour.

Auris-Midae terre australis, Chemnitz, Coneh. Cab. vol. ix.

p. 42. pl. 121. f. 1039, 1040.

Bulimus fibrosus, Gray.

Hab. New Caledonia.

We are informed by Chemnitz in the ninth volume of the Conchylian Cabinet, published in 1786, that this shell was known to Favanne as "a very rare *Buccinum* from New Holland bearing some resemblance to the false Midas' Ear," meaning the *Aaricula Midæ*, which Bruguière referred to his genus *Bulimus* under the same specific title as that used by Chemnitz in reference to this. We also learn from the same source that it was known to English collectors upwards of half a century ago, as 'The Midas' Ear from New Caledonia.' Lamarck, however, and his followers appear to have confounded it with the *B. auris-bovina*, from which, as described at Pl. XXX., it is amply distinguished.

The chief variation in the growth of the present species consists in some specimens being of an elongated form as represented in Chemnitz, and of which a specimen exactly agreeing with that figure is before me, and in others being of a compressed oblong tumid growth, as in the shell represented in the accompanying plate. Both varieties are distinguished by a plait-like callosity on the columella and the rich orange-red painting of the interior, but are equally distinct from the *B. Shongi* or *auris-bovina*.

Species 171. (Mus. Cuming.)

BULIMUS FUNCKII. *Bul. testā elongato-oblongā, subcylin-draceā, crassissimā, atractibūs quinque ad sex, longitudinaliter irregulariter rufē striatis, levibus, apicem versus subtilissimē corrugato-reticulatis, columellā subarcuatā, labro fortiter reflexo; rufō-castaneā, epidermide olivaceā crassā indutā, aperturā livido-fuscā, peritremate funeo-aurantio.*

FUNCK'S BULIMUS. Shell elongately oblong, somewhat cylindrical, rather thick, whorls five to six in number, longitudinally irregularly rudely striated, smooth, very finely corrugately reticulated towards the apex; columella slightly arched, lip strongly reflected; reddish chesnut, covered with a thick olive epidermis, interior of the aperture livid brown, peritreme smoky orange.

NYST, Mém. Société Royale de Liège, vol. i.

Bulimus superbus, Jonas.

Bulimus Caripeus, Valenciennes.

Hab. Caripe, Venezuela (in the crevices of high mountains); Funk, Dyson.

First discovered by M. Funk, and subsequently by Mr. Dyson, who found the species in great abundance at Caripe, Venezuela, where the natives were observed to collect and roast the animals for food. It is a rather solid shell of a cylindrically oblong form, and of a peculiar smoky tint of colour.

At the time M. Nyst was engaged in describing this species, it happened that Mr. Cuming, whilst passing through Belgium, observed to him that he had some recollection of its being already described in the Zoological Journal, under the name *B. labeo*. Upon this M. Nyst, in a subsequent notice of the species, took some considerable pains to exhibit the difference between the *B. labeo* and *Funkii*, reproducing the description and figure of the former for comparison, side by side, with the latter. M. Nyst is no doubt to be complimented for his ability and zeal in the matter, but in so doing he unconsciously attaches an importance to a passing conversational remark, a mere *lepsus-lingue*, or rather *mémoria*, which the speaker never intended. The species are too far removed from each other to admit of two opinions.

Species 172. (Mus. Cuming.)

BULIMUS MELANOCHEILUS. *Bul. testá elongato-oblongá, subnudicatá, anfractuós sex, longitudinaliter rudité striatis, per marginem superiorē crenatis, apicem versus subtilissimè corrugatis; columellá subcontortá, aperturā parviusculá, labro reflexo; castaneo-brunneá, medio et infra suturas indistincte fasciatá, epidermide luteo-olivaceá indutá, aperturæ fauce et peritremate scalo-nigricantibus.*

THE BLACK-LIP BULIMUS. Shell elongately oblong, slightly umbilicated, whorls six in number, longitudinally rudely striated, crenated along the upper margin, very finely corrugated; columella a little twisted, aperture rather small, lip reflected; chestnut brown, indistinctly banded round the middle and beneath the sutures, covered with a yellowish olive epidermis, interior and peritreme of the aperture livid-black.

NYST, Bul. Acad. Royale de Bruxelles, vol. xii.—Extrait, p. 8. pl. 3. f. 3 a, b.

Hab. Paraguay.

Chiefly remarkable for the peculiar livid black-lead hue of the enamel, which lines the interior and peritreme of the aperture.

Species 173. (Mus. Cuming.)

BULIMUS RHODOCHEILUS. *Bul. testá oblongo-ovatá, spirá breviusculá, anfractuós quinque, tenuidiusculis, undique minutissimè granulatis; columellá subcontortá, aperiturā suprē contractá, labro expanso; castanéa, strigis albis longitudinalibus, peculiariter undatis et reticulatis, notatá, epidermide cornu latescente indutá, peritremate roseo.*

THE ROSE-LIP BULIMUS. Shell oblong ovate, spire rather short, whorls five in number, somewhat swollen, very minutely granulated throughout; columella slightly twisted, aperture contracted round the upper part, lip expanded; chestnut brown, marked with longitudinal white streaks which are peculiarly waved and reticulated, covered with a yellowish horny epidermis, peritreme rose-coloured.

Hab. Brazil.

The light waved marks which distinguish this species have a singular net-work-like appearance, and the peritreme of the aperture is a particularly deep crimson rose.



175



176



177



177 b



177 a



179



178

BULIMUS.

PLATE XXIX.

Species 174. (Mus. Cuming.)

BULIMUS MALLEATUS. *Bul. testā oblongo-ovatā, inflatā, spirā breviusculā, acuminatā, aufractibus quinque longitudinaliter striatis, transversius peculiariter corrugatis, aufractu ultivo perawplo; columellā fortiter nūplicatā, labro latē reflexo; albā, castaneo hic illuc sparsim irregulariter maculatā, intus lutescente.*

THE BEATEN BULIMUS. Shell oblong-ovate, inflated, spire rather short, acuminate, whorls five in number, longitudinally striated, transversely peculiarly wrinkled, last whorl very large; columella strongly one-plaited, lip broadly reflected; white, sparingly and irregularly spotted here and there with chesnut, inferior yellowish.

JAY, Catalogue of Shells.

Hab.

The chesnut spots of this shell, which is now comparatively well known to collectors, have very much the appearance of scattered fragments of horny epidermis.

Species 175. (Mus. Cuming.)

BULIMUS FULGURATUS. *Bul. testā oblongo-ovatā, mabili-
cicatā, subobtusa, aufractibus quinque longitudinaliter
excissimè striatis, transversius peculiariter corrugatis,
columellā fortiter nūplicatā, labro latē reflexo; albā,
fuscō longitudinaliter undato-strigatā, epidermae luteo-
olivaceā indutā, apice versas rosacē.*

THE LIGHTNING-STRUCK BULIMUS. Shell oblong-ovate, umbilicated, somewhat inflated, whorls five in number, longitudinally very faintly striated, transversely peculiarly wrinkled, columella strongly one-plaited, lip widely reflected; white, longitudinally streaked in a waved manner with brown, and covered with a yellowish olive epidermis, pinkish towards the apex.

JAY, Catalogue of Shells.

Plekocheilus gracilis, Broderip.

Bulinus exsiccans, nudi, in Conch. Syst.

Hab. Feejee Islands.

Although the *B. fulguratus* is of a more cylindrically oblong form than the preceding species, and of a different colour and pattern, it is very similar in regard to sculpture, and in the strongly plaited columella. The transversely corrugated surface of these is rarely met with in the genus.

Species 176. (Mus. Cuming.)

BULIMUS HARTWEGII. *Bul. testā elongato-oblongā, sub-
cylindrōcā, aufractibus sex, convexis, longitudinaliter
radē striatis, per marginem crenulatis, columellā an-
guinidā, rectā, aperturā parvissimā, labro simplici, vix
reflexo; ustulato-spadicē, nigricante-fusco hic illuc
irregulariter anguste strigatā, intus violascente-albā.*

HARTWEG'S BULIMUS. Shell elongately oblong, somewhat cylindrical, whorls six in number, convex, longitudinally rudely striated, crenulated along the margin, columella narrow, straight, aperture rather small, lip simple, scarcely reflected; scorched bay colour, irregularly marked here and there with narrow streaks, interior violet-tinged white.

PFEIFFER, Monog. *Bul.*

Hab. El Catamajia Loxa, Quito; Hartweg.

The columella, though solidified, seems inclined to roll back, after the manner of those species approaching to *Achatina*.

Species 177. (Fig. a. Mus. Cuming. Fig. b. Mus.

Dennison.)

BULIMUS DOWNESII. *Bul. testā pyramidali-oblongā, spirā
acuminatā, apice obtuso; aufractibus septem convexis,
marginē crevatis, concentricē peculiariter plicato-rugati,
rugis hic illuc plus minuscē transversis: columellā
contortā et retrorsum curvata, aperturā subrotundatā,
labro reflexo; aufractuu parte superiori purpurisoerule
ad ustulato-rufescente, rugis hic illuc albī, inferiori
sepissimè castaneo-nigricante, columellā et periremante
roseis, vel pallide coccineis, intus carmineante-albā.*

DOWNES' BULIMUS. Shell pyramidal oblong, spire acuminate, obtuse at the apex; whorls seven in number, convex, crenulated at the edge, concentrically peculiarly plicately wrinkled, wrinkles here and there swollen; columella twisted and curved back, aperture somewhat rounded, lip reflected; upper part of the whorls purplish or of a burnt red colour, wrinkles here and there white, lower part of the whorls most frequently chesnut black, columella and perireme rose or scarlet-tinged, interior bluish white.

GRAY, Sowerby, Conch. Illus. f. 99.

Hab. Prince's Island, West Coast of Africa; Fraser.

There are apparently two well-marked varieties of this species, one of a deep burnt red colour, dark chesnut-black round the lower part, with a rose lip as in the richly coloured specimen here figured from the collection of Mr. Deminson; the other of a purplish colour, merely brownish towards the lower part and having the lip tinged with scarlet, as in Mr. Cuming's specimen.

Several fine examples of *B. Doxenii* were collected by Mr. Fraser at the above-mentioned island, during his adventure in the unfortunate Niger Expedition.

Species 178. (Mus. Deminson.)

BULIMUS AURIPIGMENTUM. *Bul. testá pyramidalí-elongatá, spirá acuminatá, anfractibus septem, levibus, plano-convervis; columellá angustá, rectá, quasi revolutá, aperturá parviusculá, labro simplici, acuto; peculiari refescente-spicicá, anfractibus flammis brevibus irregularibus erectis, costaneis et albis, bosi cingulatis, anfractu ultimo fasciá castaneá latá.*

THE ORPIMENT BULIMUS. Shell pyramidal elongated, spire acuminated, whorls seven in number, smooth, flatly convex, columella narrow, straight, as if rolled back, aperture rather small, lip simple, sharp; of a peculiar reddish bay, the whorls being encircled round the base with irregular short erect chesnut and white flames, last whorl with a broad chesnut band.

Hab. —?

A very handsome characteristic new species totally distinct from any hitherto known, for which I am indebted to the liberality of J. Deminson, Esq., who possesses two examples agreeing with each other in the minutest detail.

Species 179. (Mus. Cuming.)

BULIMUS FAVANNII. *Bul. testá ovatá ventricosá, umbilicata, spirá breviusculá, acuminatá, suturis impressis; anfractibus septem, concentricis striatis, levibus, columellá latá, aperturá subcontractá, labro vir reflexo; albá, maculis grandibus quadratis, rufescente-fuscis superne et inferne nebulatá.*

FAVANNE'S BULIMUS. Shell ovate, ventricose, umbilicated, spire rather short, acuminated, sutures impressed; whorls seven in number, concentrically striated, smooth, columella broad, aperture a little contracted, lip scarcely reflected; white, clouded above and below with large square reddish brown spots.

LAMARCK, *Anim. sans vert.* vol. viii. p. 227.

Helix Favannii. FéruSSAC.

Bulimus cinctus, Jay.

Hab. Eastern Islands.

Easily distinguished by its style of painting, the colour being arranged round the upper and lower parts of the whorls in large oblong-square blotches, generally flowing irregularly one into the other.



181



180



182



183



184



185



186

BULIMUS.

PLATE XXX.

Species 180. (Mus. Cuming.)

BULIMUS AURIS-VULPINUS. *Bul. testā ocaū, ventricosā, subumbilicatā, spirā subacuminatā, anfractibus septem, transversim et longitudinaliter rudo rugoso-striatis, superne depresso angulato-rotundatis et corrugato-crenatis; columellā profundi excavatā, basi contortā, labro incrassato, rudo reflexo, aperturā sinuatā et contractā; sordide alba, columellā croco-rufā, apice et aperture fance pallide crocatis.*

THE FOX-EAR BULIMUS. Shell ovate, ventricose, slightly umbilicated, spire somewhat acuminate, whorls seven in number, transversely and longitudinally roughly striated, depressed angularly rounded and pucker'd above; columella deeply excavated, twisted at the base, lip thickened, rudely reflected, aperture sinuate and contracted; dead white, columella saffron-red, apex and interior of the aperture faintly tinged with saffron colour.

DESHAXES, Anūm. sans vert. vol. viii. p. 257.

Auris vulpina, Chemnitz.

Toluta auris-vulpina, Dillwyn.

Helix auris-vulpina, Féussac.

Hab. St. Helena.

A curious species of which several specimens have been found in a semi-fossil state at St. Helena, supposed to have belonged to a mollusk now extinct; none having been collected in a living state.

Species 181. (Mus. Cuming.)

BULIMUS HANLEYI. *Bul. testā elongato-oblongā, sub-turrītā, tenui, semipellucidā, spirā acuminatā, suturis impressis; anfractibus septem, levibus, nitentibus, plano-concisis, per marginem superiorē subtilliter et creberriōe crenulatis, crenulis in sulcis minutis descenditibus, columellā labroque tenuibus, simplicibus; rufo-castaneā, intus iridescente.*

HANLEY'S BULIMUS. Shell elongately oblong, somewhat turreted, thin, semipellucid, spire acuminate, sutures impressed; whorls seven in number, smooth, shining, flatly convex, very finely and closely crenulated along the upper margin, the crenules descending in fine grooves, lip and columella thin, simple; reddish-chestnut, iridescent within.

PFEIFFER, Monog. *Bul.*

Hab. Brazil; Miller.

A light turriculated shell of a peculiar reddish mahogany brown-colour, the sutures of which are somewhat deeply impressed, whilst the upper edge of the whorls is very finely crenulated, the crenules being delicately prolonged.

Species 182. (Mus. Cuming.)

BULIMUS SPIXII. *Bul. testā elongato-turrītā, subcylin-dracea, spirā acuminatā, apice concavo-depressā; au-fractibus septem, plano-concisis, longitudinaliter subtilissimā corrugato-striatis, per marginem crenulatis; columellā peculiariter incisa, aperturā portā, labro subincrasato, viz recurvo; corneo-fusca, obscurè fasciatā, epidermide lateo-olivaceā iunctā, labro et apertura fance pallide carneo-rosaceis, subridescensibus.*

SPIX'S BULIMUS. Shell elongately turreted, somewhat cylindrical, spire acuminate, concavely depressed at the apex; whorls seven in number, flatly convex, longitudinally very finely striated in a wrinkled manner, crenulated along the edge; columella peculiarly notched, aperture small, lip somewhat thickened, but slightly recurved; flesh-tinted brown, obscurely banded, covered with a yellowish-olive epidermis, lip and interior of the aperture flesh-tinged rose, slightly iridescent.

WAGNER, Test. Bras. p. 11. pl. 10. f. 3.

Columba boliviæ, Spix.

An eadem junior, *Bulimus hyalinus*, Wagner.

Hab. Brazil.

The notched columella is a very conspicuous feature in this species, which, in fine condition, is not white as hitherto described, but of a flesh-tinted rose hue, banded with brown and covered by a yellowish-olive epidermis.

Species 183. (Mus. Cuming.)

BULIMUS IMPERATOR. *Bul. testā pyramidali-ovalē, au-fractibus sex, levibus, ultimo basis versus subangulato, columellā rectā; purpuro-nigricante, epidermide spadiceo-alba, undato-striatā iunctā, aperturā carvescente-alba, columellā rosaceo-purpurascente, peritremate nigricante.*

THE EMPEROR BULIMUS. Shell pyramidal-oval, whorls six in number, smooth, last whorl slightly angled

towards the base, columella straight; purple-black, covered with an epidermis marked with zigzag streaks of fawn-white, aperture bluish-white, columella pinkish-purple, peristome black.

PFEIFFER, Pro Zool. Soc., 1848.

Hab. Philippine Islands: Kierulf.

I doubt if 'The Pretender' would not have been a more appropriate name for this shell than 'The Emperor,' for it appears to be simply a variety of *B. pythagaster*. It would, however, be unfair to judge of its claim to rank as a species by a single specimen not in the best condition. Mr. Cuming informs me that it was collected at the Philippines by Dr. Kierulf of Copenhagen, and that there is another exactly similar to it in the collection of M. Janelle of Paris; and further, that he never collected any *B. pythagaster* with this peculiar zigzag style of marking.

Species 184. (Mus. Cuming.)

BULIMUS ATOMATUS. *Bul. testā oblongo-ovatā, Achatinae-formi, tenui, anfractibus sex, levibus, inflatis, per marginem leviter crenulatis; columellā rectā, reflexā, labro externo, tenui, simplici; flavescente-brunneā, castaneo-rafū minuti undato-lineatā et punctatā, intus cernucente-albā.*

THE MINUTELY DOTTED BULIMUS. Shell oblong-ovate, Achatina-shaped, thin, whorls six in number, smooth, inflated, slightly crenulated along the edge; columella straight, reflected, outer lip thin, simple; yellowish-brown, minutely lined and dotted in a waved manner with chestnut-red, interior bluish-white.

GRAY, Pro Zool. Soc., 1834, p. 64.

Hab. Near Fort Macquarie, New Holland.

A singularly marked species having somewhat the form of the large African *Achatinae*.

Species 185. (Mus. Dennison.)

BULIMUS AURIS-BOVINA. *Bul. testā elongato-oblongā, crassā, spirā subacuminiatā, anfractibus sex, engosistratis; columellā incrassatā, callositatis deabus multis, quarum inferiori peramplo, solidā; apertura*

pareā, peculiariter contractā, labro valde incrassato; rufescente-albō, aperture fauce vivide rufā, columellā labroque chlarneis.

THE OX-EAR BULIMUS. Shell elongately oblong, thick, spire somewhat acuminate, whorls six in number, roughly striated, columella thickened, furnished with two callosities, of which the lower is very large and solid, aperture small, peculiarly contracted, lip very much thickened; reddish-white, interior of the aperture deep red, lip and columella ivory white.

BRUGUIERE, Enc. Méth. dict. no. 80.

Auricula bovinā, Lamarek.

Helix auris-bovina, Féussac.

Hab. Eastern Islands.

The *Bulimū Shougi* and *auris-Mido* have each been confounded in its turn with the present species, which is eminently distinguished from them; first, in being of a natural white, which obtains a reddish hue, owing to the interior being very deeply stained with that colour; secondly, in the aperture being small and much contracted; thirdly, in being of more solid growth, and fourthly, in the lip and columella being like white porcelain or polished ivory.

Species 186. (Mus. Cuming.)

BULIMUS DOMBEYANUS. *Bul. testā ovatā, umbilicatā, spirā brevissimā, mucronulā, anfractibus sex ad septem, subrotundatā, concentricē rugatis; columellā subcontortā, reflexā, labro latissimē expanso et reflexo; albā, columellā et apertura fauce intense laco-purpureis.*

DOMBEY'S BULIMUS. Shell ovate, umbilicated, spire rather short, pointed, whorls six to seven in number, somewhat rounded, concentrically wrinkled; columella a little twisted, reflected, lip very widely expanded and reflected; white, interior deeply coloured with purple-lake.

Helix Dombeiana, Féussac, Hist. Moll.

Hab. Mexico.

This beautiful and very rare species is chiefly remarkable on account of the trumpet-like expansion of the aperture, which is of a deep purple-bronze colour.



BULIMUS.

PLATE XXXI.

Species 187. (Fig. *a*, Mus. Hudson : Fig. *b*, Mus. Cuming.)

BULIMUS CITRINUS. *Bul.* testā subelongato-ovatā, crassissimā, ant. dextrā aut sinistrā, aufractibus sex ad septem, subrotundatis, levibus, nitidis; rieide citrino-flavā, interdum rufescere-fusco-flammatā, prope suturas albā, cīttā longitudinaliō castaneo-nigrā interdum pietā, labro et periremata albus.

THE CITRON BULIMUS. Shell somewhat elongately ovate, rather thick, either dextral or sinistral, whorls six to seven in number, somewhat rounded, smooth, shining; bright citron yellow, sometimes flushed with reddish brown, white near the sutures, sometimes painted with a chestnut-black longitudinal fillet, lip and perireme white.

BRUGUERIE, Enc. Méth. dict. no. 27.

Helix perversa (pars?), Linneus.

Helices dextra, and *sinistra*, Muller.

Helix aurata, Férrussac.

Hab. Moluccas.

This species is so well known by its long-established name, *B. citrinus*, that I have hesitated to adopt the Linnean title *Helix perversa*, believing it probable that the author of the *Systema Naturae*, included more than one reversed species under that appellation.

Species 188. (Mus. Cuming.)

BULIMUS ATRICALLOSOUS. *Bul.* testā oblongo-ovatā, basi versus subinflatā, spira acuminatā: aufractibus sex ad septem, subrotundatis, levibus, nitidis, labro expenso, late reflexo; albida, cirridescente-lato tintā, parte supra columellam laniā collosa intus purpureo-nigrā inductā, aufracta ultimā cīttā simili longitudinaliō pictā, aperturā albā.

THE BLACK-CALLOUS BULIMUS. Shell oblong-ovate, somewhat inflated towards the base, spire acuminated, whorls six or seven in number, somewhat rounded, smooth, shining, lip expanded, widely reflected; white, stained with greenish yellow, the part above the columella covered with a callous deep purple-black lamina, last whorl painted with a longitudinal fillet of the same colour, aperture white.

GOLD, Boston Journ. Nat. Hist.

Hab. Tavoy, Burmah.

I fear there is not much difference between this and the

preceding species, beyond the callous purple-black lamina which lines the interior of the body-whorl, and is of somewhat doubtful importance.

Species 189. (Mus. Cuming.)

BULIMUS LINDENI. *Bul.* testā cylindraceo-oblongā, crassissimā, apice obtuso depresso, aufractibus quinque ad sex, decessante raro striatis et subtilliter indentatis, per marginem exiliter crenulatis; columella saepe platicata, apertura parvisepta, labro subincrustata, reflexo: lac-tea, nigra sparsio striata, apicem versus purporascente, epidermide olivaceo-viridi indutā, apertura fave cæ-reolente-alba, labro pallide crocato.

LINDEN'S BULIMUS. Shell cylindrically oblong, rather thick, apex obtusely depressed, whorls five to six in number, decessantly rudely striated and finely indented, faintly crenulated along the margin; columella one-plaited, aperture rather small, lip somewhat thickened, reflected; cream colour, sparingly streaked with black, purplish towards the apex, and covered with an olive-green epidermis, interior of the aperture bluish white, lip pale saffron colour.

Bulinus leptochelus, Pfeiffer.

Hab. La Baja, Province of Pamplona, New Granada; Funek, Linden.

Notwithstanding the importance which is very properly attached to the priority of names, there are exceptions, as in the instance before us, in which a name may require to be changed. The term *leptocheilus*, or *thin-lipped*, denotes a character which is not found in the adult species: the shells of all *Bulinii* are "thin-lipped" in an immature state, and many remain so at maturity; but in the species under consideration, the lip is thickened and rather widely reflected. I therefore venture to name it anew, in honour of the indefatigable botanist and shell-collector, to whom we are indebted for so many beautiful species of *Bulinii* from the luxuriant di-trict of New Granada.

Species 190. (Mus. Cuming.)

BULIMUS PLAXIDENS. *Bul.* testā oblongo-oriformi, tenuis, spirā brevi, neminatā, aufractibus quinque, tenuidis, longitudinaliter crenberinā corrugato-striatis, striis minutissimis transversis sub lete decussatis, aper-

turrid *oblongo-quadrati*, *parietiscula*, *contracta*, *labro*
dente unico planulato instruto; *oliveaceo-virescente*,
aperturā caralescere-alba, *peritremate intensē par-*
pureo-roseo.

THE FLAT-TOOTHED BULIMUS. Shell oblong egg-shaped, rather thin, spire short, acuminate, whorls five in number, swollen, longitudinally very closely striated in a wrinkled manner, and, under the lens, crossed with very minute transverse striae, aperture oblong-square, rather small, contracted, furnished with a single flattened tooth; light olive-green, aperture bluish white, peritreme deep purple-rose.

MICHELIN, Magasin de Zool. pl. 25.

Hab. Brazil.

The surface of this species, in fine condition, has a delicate silken aspect, occasioned by a multitude of extremely fine transverse striae, which are scarcely discernible without the aid of a lens. The epidermis is of a peculiar plum-green colour, and the lip of an intense purple-rose, distinguished by an oblong flattened tooth.

Species 191. (Mus. Cuming.)

BULIMUS PUDICUS. *Bul. testā oblongo-ovalā, spirā brevi-*
acuminatā, aufractibus quinque tumidiusculis, undique
striato-corruigatis, columellā nuplicatā, labro expanso,
latē reflexo; oliveaceo-rufā, peritremate nunc albo, nunc
vividē roseo.

THE CHASTE BULIMUS. Shell oblong-ovate, spire short, acuminate, whorls five in number, rather swollen, striately wrinkled throughout, columella one-plaited, lip expanded, widely reflected; olive-red, peritreme sometimes white, sometimes deep rose.

DESHAYES, Anim. sans vert., vol. viii. p. 253.

Helix pudica, Linnaeus, Syst. Nat. (Gmelin's edit.)
 p. 3645.

Partula pudica, De Ferussac.

Helix pudica, Moricand.

Auris virginaea, Chemnitz.

Voluta auris-virginis, Dillwyn.

Bulimus virginicus, Bruguière.

Strophochelus Almeida, Spix.

An eadem var. *Bulimus perplexus*, Sowerby.

Hab. Brazil.

M. Deshayes has already observed that the shell figured by Chemnitz under the name of *Auris virginaea* (Coneh. Cab. vol. ix. pl. 121. f. 1042) has a different aspect from that commonly known to us as the *B. pudicus*, figured in the accompanying plate, and among the shells collected by Dr. Spix in Brazil. The *Auris virginaea* has more the form of a *Partula*, but I find no shell to correspond with it, and take it to be merely another state of the species under consideration.

(Species 192. Mus. Cuming.)

BULIMUS UNIDENTATUS. *Bul. testā oblongo-ovalā, spirā*
breviusculā, aufractibus quatuor ad quinque, tumidis,
subrugosis, aperturā pariescentia, labro reflexo, deute-
re uno obtuso instruto; albida, roseo tintā, columellā
labroque purpureo-castaneis.

THE ONE-TOOTHED BULIMUS. Shell oblong-ovate, spire rather short, whorls four to five in number, swollen, rather rough, aperture somewhat small, lip reflected, furnished with a single blunt tooth; whitish, tinged with rose, lip and columella purple-chesnut.

Partula unidentata, Sowerby, Tankerville Catalogue, App. p. vii.

Hab. Brazil.

I am not aware that any example of this species is known in fine condition with its epidermis. It is of more solid growth than the *B. planidens*, and of a less oblong form, whilst the tooth is short and blunt, not oblong and flattened. The species differ essentially also in colour.



194.



195.



196.



197.



198.



199.

BULIMUS.

PLATE XXXII.

Species 193. (Mus. Cuming.)

BULIMUS GUERINI. *Bal. testā oblongo-ovatā, tenuiculā, apice subobtusa, anfractibus quinque, levibus, subinflatā, columellā subcontortā, labro vix reflexo; intensē olivaceā, lateo longitudinaliter striatā, apertura periretate aurantio.*

GUERIN'S BULIMUS. Shell oblong-ovate, rather thin, somewhat obtuse at the apex, whorls five in number, smooth, slightly inflated, columella somewhat twisted, lip scarcely reflected; very dark olive, longitudinally streaked with yellow, peritreme of the aperture orange.

PFEIFFER, Pro. Zool. Soc., 1846, p. 40.

Hab. New Granada.

Although much smaller in size and of comparatively lighter growth, this species approximates so closely to the *B. Moritzianus*, which is found in the same locality under a multitude of aspects, that I fear it may prove to be only a dwarf variety.

Species 194. (Mus. Cuming.)

BULIMUS CRYSTALLINUS. *Bal. testā conico-ovatā, tenui, pellucida, valde inflata, anfractibus quinque ad sex, transversim undique subditissimè impresso-striatis, anfractū ultimo perauplo, basin versus obtuse cariato, columellā concavo-depressā, vix reflexā, labro tenui, simplici; pellucido-olivā, columellā brunnescente.*

THE CRYSTALLINE BULIMUS. Shell conically ovate, thin, pellucid, very much inflated, whorls five to six in number, transversely very finely impressly striated throughout, last whorl very large, obtusely keeled towards the base, columella concavely depressed, slightly reflected, lip thin, simple; transparent white, columella pale brown.

Hab. —?

Of very peculiar form, and differing in all respects from any species hitherto described. It is a light transparent shell, transversely engraved with fine striae, very much inflated, and indistinctly angled towards the base.

Species 195. (Mus. Cuming.)

BULIMUS MONOZONUS. *Bal. testā subpyramidalis-ovatā, apice obtusa, anfractibus quinque, rotundatis, oblique striatis, columellā rectā, labro vix reflexo; rufescente-*

brunnea, zonā latiusculā albā, in anfractu ultimo tantum conspicuā, cingulatā; epidermide fusca indutā, periretate nigricante,

THE ONE-ZONE BULIMUS. Shell somewhat pyramidal in shape, obtuse at the apex, whorls five in number, rounded, obliquely striated, columella straight, lip but slightly reflected; reddish brown, encircled by a rather wide white zone, conspicuous only on the last whorl, and covered with a brown epidermis, peritreme blackish.

PFEIFFER, Monog. *Bal.*

Hab. Philippines; Kierulf.

Collected in the Philippine Islands by Dr. Kierulf of Copenhagen, and apparently distinct from any of Mr. Cuming's species.

Species 196. (Mus. Cuming.)

BULIMUS GIBBONIUS. *Bal. testā ovatā, umbilicata, crassisculā, ventricosissimā, anfractibus quatuor ad quinque, longitudinaliter radè striatis, undique minutiè granulatis, anfractū ultimo perauplo, inflato, columellā subcontortā, labro reflexo; castaneo-rufescente, fuscō sparsissimā punctatā, aperturā albida.*

GIBBON'S BULIMUS. Shell ovate, umbilicated, rather thick, extremely ventricose, whorls four to five in number, longitudinally rudely striated, minutely granulated throughout, last whorl very large, inflated, columella slightly twisted, lip reflected; chestnut red, sparingly dotted with brown, aperture whitish.

LEA, Trans. Amer. Phil. Soc., 1836.

Hab. Between La Plata and Tocaima, New Granada; Dr. Gibbon.

Distinguished by the widely-inflated and rapid enlargement of the last whorl; of a chestnut colour, besprinkled with dashes of brown.

Species 197. (Mus. Cuming.)

BULIMUS CASTANEUS. *Bal. testā ovatā, vix umbilicata, ventricosissimā, anfractibus quatuor ad quinque, longitudinaliter radè striatis, granulis rufibus oblongis propagatis seriatim cingulatis, anfractū ultimo perauplo, inflato, labro vix reflexo; intus exlusque fuscō-castanea.*

THE CHESTNUT BULIMUS. Shell ovate, scarcely umbilicated, extremely ventricose, whorls four to five in

number, longitudinally rude striated, encircled near the sutures with rows of rude oblong granules, last whorl very large, inflated, lip but slightly reflected, smoky chestnut-brown within and without.

PFEIFFER, Pro. Zool. Soc., 1845, p. 68.

Hab. Vegas, on the river Quendu, New Granada.

It may be observed that this shell is almost equally ventricose and inflated with the preceding species, though not exactly of the same form. It is, moreover, of lighter structure, and, instead of being covered with granules, it has merely a few near the sutures of a rude oblong shape, set in exact rows more or less distant from each other.

Species 198. (Mus. Cuming.)

BULIMUS GALLINA-SULTANA. *Bul. testá conico-ovatá, tenuí, ventrisciosissimá, anfractibus sex, transversam creberimè striatis, striis regosis et minime undulatis, anfractu ultimo peramplo, inflato, labro simplici; albicans, olivaceo-viridi maculata et variegata, epidermide floccidá indutá.*

THE SULTAN-FOWL BULIMUS. Shell conically ovate, thin, extremely ventricose, whorls six in number, transversely very closely striated, striae rough and minutely undulated, last whorl very large, inflated, lip simple; whitish, blotched and variegated with olive green, and covered with a yellowish epidermis.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. viii. p. 222.

La poule sultane, Favanne.

Helix gallina-sultana, Chemnitz.

Achatina pavonina, Spix.

Hab. Bolivia; D'Orbigny.

"We met with the *B. gallina-sultana*," says M. D'Orbigny, "in the hot damp forests, inhabited by the savage Guarayos, in the centre of the republic of Bolivia. There, they bury themselves among the roots of trees and only come forth during the rainy season; they then crawl upon the trunk and stems to inhale the moisture, and as soon as the rains begin to cease, they descend and again bury themselves in the earth, where they remain concealed until the following season."

Species 199. (Mus. Cuming.)

BULIMUS HETEROTRICHUS. *Bul. testá conico-ovatá, subglobosá, umbilicatá, anfractibus sex ad septem, rotundatis, striis undique sub lente minime decussatis, columellá latissimá, labro exiliiter reflexo; olivaceo-flavidá, epidermide setosa indutá, setis subtilibus, in seriebus spiralibus subdistantibus.*

THE DIFFERENTIALLY-BRISTLED BULIMUS. Shell conically ovate, somewhat globose, umbilicated, whorls six to seven in number, rounded, under the lens minutely decussated throughout with striae, columella rather broad, lip delicately reflected; olive yellow, covered with a bristly epidermis, of which the bristles are fine and arranged in rather distant spiral rows.

MORICAND, Mém. de Genève, vol. 2, part. 2, p. 430, pl. 2. f. 5, 6.

Hab. Brazil,

Well characterized by its delicate bristly epidermis, deprived of which, the species may be equally recognised by its peculiar form and minute decussated sculpture, looking, as it were, like a fine web of punctures.



BULIMUS.

PLATE XXXIII.

Species 200. (Mus. Cuming.)

BULIMUS SIGNATUS. *Bul.* testā oblongo-ovatā, umbilicatā, spirā brevi, acuminatā, anfractibus quinque, rugositas, columellā basi callotā, rūdē plicatā, aperturā angustā, contractā, auriformi, labro reflexo et latissimè concavocurvato; albida; fasciis tribus tatis nigricantibus cingulatā, fuscoque nebulatā, apice nigro, labro chunneo.

THE MARKED BULIMUS. Shell oblong-ovate, umbilicated, spire short, acuminated, whorls five in number, rough, columella callous at the base and obsoletely pliated, aperture narrow, contracted, car-shaped, lip reflected and very widely concavely expanded; whitish, encircled with three broad blackish bands and clouded with brown, apex black, lip ivory white.

Auricula signata, Wagner, Moll. du Brésil, p. 17. pl. 12. f. 3.
Helix signata, Moricand.

Hab. Brazil.

The wide-spread developement of the lip is rather a peculiar feature in this species. It is not uncommon.

Species 201. (Mus. Cuming.)

BULIMUS BILABIATUS. *Bul.* testā ovatā, spirā acuminatā, anfractibus quinque, subrotundatis, rugositas, sub tenui minute rūde reticulatis et foratis, peculiariter oblique plicato-costatis, costis compressis; columellā subcontortā, aperturā auriformi, labro undique bisecto; sordide albā, fusco hic illuc lineatā, parte suprā columellam nunc auratā, nunc purpureo-nigrā.

THE TWO-LIPPED BULIMUS. Shell ovate, spire acuminated, whorls five in number, somewhat rounded, rough, under the lens minutely reticulated and punctured, peculiarly obliquely plicately ribbed, ribs compressed; columella rather twisted, aperture car-shaped, lip divided in twain all round; dead white, stained here and there with brown, the part above the columella sometimes orange, sometimes purple-white.

BRODERIP and SOWERBY, Zool. Journ. vol. v. p. 49.
pl. supp. 40. f. 1, 2.

Helix maximiliana, Féussae, MSS.

Hab. Brazil.

The name *B. maximiliana* of De Féussae does not appear to have been published. There is a very characteristic dwarf state of the species exhibiting the same varieties of colouring as those of a larger growth.

Species 202. (Mus. Cuming.)

BULIMUS TAUNAYSI. *Bul.* testā elongato-oblongā, subcylindracea, anfractibus sex ad septem, convexis, lăevis, longitudinaliter striatis, columellā subcontortā, labro tenuicolo, eis reflexo; fulvo vel nigricante-fuscā, zonulā pallidā angustā cingulatā, columellā albā.

TAUNAY'S BULIMUS. Shell elongately oblong, somewhat cylindrical, whorls six to seven in number, convex, smooth, longitudinally striated, columella somewhat twisted, lip rather thin, scarcely reflected; fulvous or blackish brown, encircled by a pale narrow zone, columella white.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 294.

Helix Taunaysi, Féussac.

Eadem. *Bulimus magnificus*, Grateloup.

Hab. Brazil.

The shell named *B. magnificus* by M. Grateloup is, according to the specimens so marked in Mr. Cuming's collection, a variety of *B. Taunaysi* of a lighter brown colour.

Species 203. (Fig. a. Mus. Demmison. Fig. b, and c. Mus. Cuming.)

BULIMUS MELANOSTOMA. *Bul.* testā oblongo-ovatā, spirā subacuminatā, anfractibus quinque ad sex, undique peculiariter oblongo-granulatis, per marginem minime crevatis et plicatis, platis subamplo distans, in anfractu ultimo evanidis; columellā contortā, labro latè reflexo, interdum partim bisecto; roseo-albicans, fuscescente obscurè nebulatā, spirā apicem versus saepe rosacea, inter platis nigricante, aperturā partim vel omnino purpureo-nigrā, interdum croco marginatā.

THE BLACK-MOUTH BULIMUS. Shell oblong-ovate, spire somewhat acuminated, whorls five to six in number, peculiarly oblong-granulated throughout, minutely crenulated along the margin, and plicated, the plait being rather large and distant, fading away in the last whorl; columella twisted, lip widely reflected, sometimes partially bisected; pinkish-white, obscurely clouded with pale brown, spire frequently pinkish towards the apex, blackish between the folds, aperture partially or entirely purple-black, sometimes edged with saffron yellow.

SWAINSON, Zool. Illustrations, vol. i. pl. 4.

Helix melanostoma, Féussac.

Auris melanostoma, Spix.

Eadem. *Helix rhodospira*, Moricand.

Hab. Brazil.

The varieties of this species are strongly indicated in the distribution of the colouring matter of the aperture.

In most specimens the aperture is wholly lined with rich purple black, as in fig. 203 c.; in some, the lip is produced a second time outwardly, as in fig. 203 a, bordered with saffron yellow; whilst in others, as in fig. 203 b, the aperture is white, and no purple-black is seen, excepting on the body whorl above the columella.

— — — — —



BULIMUS.

PLATE XXXIV.

Species 204. (Mus. Denison.)

BULIMUS POPELAIRIANS. *Bul. testā ovatā, ventricosissimā, crassiusculā, subcompressā, anfractibus tuvidis, superne crenulatis, apicem versus concentricē subdistanter tiratis et minutissimē granulatis; columellā arcuatā, alata superiorē callōsa, labro incrassato, reflexo: castaneo-brunnecente, epidermaide crassā cornē brunneā, in strigis longitudinalibus acutē undatis interdum pecudioriter dissolida, indatā; aperturā lividā, labro albo.*

POPELAIR'S BULIMUS. Shell ovate, extremely ventricose, rather thick, somewhat compressed, whorls swollen, crenulated round the upper part, concentrically somewhat distantly ridged and minutely granulated towards the apex; columella arched, callous at the upper part with age, lip thickened, reflected; light chesnut-brown, covered with a thick brown epidermis, which is sometimes peculiarly dissolved, as it were, in longitudinal zigzag streaks; aperture livid, lip white.

NYST, Bull. Acad. Roy. de Bruxelles, vol. xii. Extr. p. 10.
pl. 4. f. 5.

Eadem. *Bulimus fulgoratus*, Valenciennes.

Bulimus Valenciennesii, Pfeiffer.

Hab. Brazil.

This fine zigzag-marked species, distinguished under the above names by Pfeiffer and Valenciennes, is, without doubt, the same described by M. Nyst in the Transactions of the Royal Academy of Brussels in honour of the Baron de Popelaïr. Mr. Cuming possesses a specimen, like M. Nyst's, quite destitute of the zigzag marking, which is not a distribution of colour, nor the result of a double-membrane arrangement of the epidermis, as in the Philippine species; it has rather the appearance of being partially dissolved, as though it were occasioned by the percolation of some solvent out of the sutures, though incapable of such an explanation.

Species 205. (Mus. Cuming.)

BULIMUS EGREGIUS. *Bul. elongato-oblongā, subfusiformi, umbilicatā, anfractibus concavusculis, superne concavis, undique crenatimē placo-granulatis, granulis in series transversis ad intervalla subdistantia, majoribus, anfractu ultimo basi angulato-compresso; aperturā oblongo-ovali, basi angulatā et canaliculatā, labro sub-expanso et reflexo, marginibus lirā agrestē callōsa juncis: castaneo-fuscā, latro flammatā, labro intense roseo.*

THE CHOSEN BULIMUS. Shell elongately oblong, somewhat fusiform, umbilicated, whorls rather convex, concave round the upper part, very closely flatly granulated throughout, granules in transverse rows, at rather distant intervals, larger, last whorl angularly compressed at the base; aperture oblong-oval, angulated and channelled at the base, lip slightly expanded and reflected, margins joined by a narrow callous ridge; chesnut-brown, flamed with yellow, lip deep rose.

PFEIFFER, Pro. Zool. Soc., 1845, p. 67.

Bulimus hybridus, Gould (fide Pfeiffer).

Hab. Brazil, in the vicinity of Rio Janeiro; Gould.

I much doubt whether this is anything more than a variety of the *B. goniostoma*. I have not met with it in a state intermediate between the specimens here represented, but it is certain that the sculpture of *B. egregius* is precisely that of *B. goniostoma* in an exaggerated and more luxuriant condition, whilst the species are alike in form.

Species 206. (Mus. Cuming.)

BULIMUS GONIOSTOMA. *Bul. testā elongato-oblongā, fusiformi, umbilicatā, spirā acuminatā, anfractibus convexusculis, superne concavis, undique minutissimē et crenatimē granulatis, granulis, in series transversis ad intervalla subdistantia, paulo majoribus, anfractu ultimo basi compresso; aperturā oblongo-ovali, basi caudatoculatā, labro subexpanso et reflexo, marginibus lirā callōsa juncis: basidā, labro intense roseo.*

THE ANGLED-MOUTH BULIMUS. Shell elongately oblong, fusiform, umbilicated, spire acuminated, whorls slightly convex, concave round the upper part, very minutely and closely granulated throughout, the granules in transverse rows at rather distant intervals a little larger, last whorl compressed at the base; aperture oblong oval, channelled at the base, lip somewhat expanded and reflected, margins joined by a callous ridge; brown, lip deep rose.

Ochlogaea goniostoma, Féussac, Prodrome, No. 441.

Pupa goniostoma, Gray.

Goniostoma erubescens, Swainson.

Hab. Brazil.

Quite an isolated form of the genus, few species presenting much approximation to this in respect of its angular oblong channelled aperture, besides the preceding and one or two of smaller size, to be described hereafter, from the Province of Merida, New Granada.





BULIMUS.

PLATE XXXV.

Species 207. (Mus. (?) olim Soc. Zool. Lond.)

The accompanying figure, copied from the Zoological Journal, Tab. Supp. xxxi, represents a very remarkable species of *Bulimus*, deposited some years since in the Zoological Society's Museum. The shell, however, disappeared shortly afterwards from the collection, together with some other rarities, and the species has never since been obtained or heard of. It only remains therefore to copy the description also.

BULIMUS LABEO. *Bul. testá ovato-producta, fusco-castanea, apice versus rubrā, fusco variá, aufractibüs sex, ventricosis, ultimo fascis duabus nigris, hinc mediá, illid suturali, penultimo fascis duabus nigris, suturalibus; columellā dente obtuso insigni; labro crassissimo, reflexo, sapro pallide castaneo, infra nigro; apertura iutus obtusa.*

THE BLUBBER-LIP BULIMUS. Shell ovately produced, brown-chesnut, red towards the apex, variegated with brown, whorls six in number, ventricose, the last with two black bands, one round the middle, the other next the suture, penultimate whorl with two bands, both sutural; columella with a remarkable obtuse tooth, lip very thick, reflected, pale chesnut above, black beneath, interior of the aperture whitish.

BRODERIP, Zool. Journal, vol. iv. p. 222. pl. supp. xxxi.
Hab. Peru.

To quote Mr. Broderip's more minute description of this remarkable lost species:— “ Shell stout, long oval, of a brown chesnut colour, changing to red at the apex and on the upper whorls, which last are longitudinally striped with reddish brown towards the suture. Whorls six, ventricose; the last has a narrow band across its middle, and another of the same colour close to the suture, which is white; the last whorl but one has two narrow black bands, both near the sutures; the lower bands on the last and penultimate whorls are each thrice interrupted. On the last whorl, near the base, which is very dark, is a faint, broad, lighter coloured band. The columella is remarkable for its obtuse white tooth, surrounded by the rich dark colouring of the aperture. The right lip is of huge thickness, and much reflected: above, it is of a light chesnut colour; below, of a rich brownish japan-like black, which, particularly where it is shading off into the chesnut, gives, when the light is thrown full on it, the same kind of iridescent appearance as is seen in the Limachella, or Fire-marble. The lower edge of this rich lip is punctured

pretty thickly with dots, resembling those of *Cypraea testudinaria*, which seem filled with a whitish opaque substance, and the formation of which has given an irregular, and almost fungus-like appearance to the reflected border of the lip, on its upper side. The interior of the aperture is white. Length three inches; breadth, measured across the body whorl, and including the lip, one inch and six-eighths.

“ This fine shell, of a much firmer fabric than most land-shells, was obtained by Lieut. Maw, R.N., in a rancho (farm-house) at Toulea, about nine leagues to the eastward of Chachapoyas, in Peru, on Christmas day, 1827. It had been taken on the chaera (farm) to which the rancho belonged. The chaera is situated at the highest limit of the montana, or woody district, on the eastern side of the Andes, at an elevation of about 8000 feet above the level of the sea. The species was frequently seen by Lieut. Maw in the woods between Chachapoyas and Moyabamba. Two of a much larger size than the individual here described, were found by him alive, and given in charge to the arriero (muleteer), with a special injunction to take care of them. The arriero, supposing that they were to be prepared for breakfast, roasted them, and thus destroyed the shells.

“ Spix has not mentioned this shell, and I have reason to believe that there is no other specimen in this country. Whether the species is confined within narrow limits in the almost untrdden district traversed by Lieut. Maw, will be an interesting subject for future inquiry. This enterprising traveller is the first European who has performed a journey across the continent of South America in the line of the Maranon, or Amazon River. After passing from Truxillo across the Andes to Chachapoyas he embarked upon the river in Peru, and proceeded throughout its course to Para. In the course of this expedition he collected several animals, which he has liberally presented, together with the subject of this memoir, to the Zoological Society.”

Species 208. (Mus. Cuming.)

BULIMUS LACUNOSUS. *Bul. testá oblongo-ovata, aufractibus subzentricosis, longitudinaliter striatis, striis brevibus impressis peculiariter subtilissimè notatis, apice versus nuncù granulatis et radiatim liratis, columellā vix contortā, labro reflexo; livido-brunneā, longitudinaliter obscurè fasciatā, epidermine tenuicollā coram induitā, labro albo.*

THE RAVINE-INHABITING BULIMUS. Shell oblong-ovate, whorls somewhat ventricose, longitudinally striated, and peculiarly very finely marked besides with short impressed striae, minutely granulated and radiately ridged towards the apex, columnella scarcely twisted, lip reflected; livid brown, longitudinally obscurely banded, covered with a rather thin horny epidermis, lip white.

D'OREIGNY, Voy. dans l'Amér. Mérid. Moll. p. 302.

Hab. Cochabamba, Bolivia (in deep ravines at the base of the snowy peaks of the Cordilleras); D'Orbigny.

Intermediate, in a manner, between the *B. maximus* and *ovatus*, of smaller size, and apparently distinct. In the specimen here figured from Mr. Cuming's collection, the surface is curiously marked by numerous rows of short longitudinal hollowed striae; it is also incircled by a couple of scratched grooves, but these are evidently the result of accident.

Species 209. (Mus. Cuming.)

BULIMUS GRANULOSUS. *Bul. testa oblongo-ovata, au-*
fractibus centricosis, longitudinaliter rude striatis, nu-
disque minutè granulata, prope apicem subobscure
radiatum liratis, columellâ subcoartatâ, labro reflexo;
tuteo-brunnecente, epidermide levicentâ indutâ, colu-
mellâ labraque pallidè roseis, apertura fuscæ carde-
albicans.

THE GRANULOUS BULIMUS. Shell oblong-ovate, whorls ventricose, longitudinally rudely striated, minutely granulated throughout, somewhat obscurely radiately ridged near the apex, columella somewhat twisted, lip reflected; yellowish brown, covered with a thin epidermis, lip and columella pale rose, interior of the aperture bluish white.

Helix granulosa, Rang, Deser. coquilles terrestres, p. 49.
pl. 2.

Hab. Brazil.

It is much to be feared whether this is not truly a variety of the *B. ovatus*. The typical differences are as follows.—in form the *B. granulosus* is of a less shortened umid growth than the *B. ovatus*, and in sculpture, it is more prominently granulated throughout, whilst the radiated ridges about the apex are more obscure: it is,

moreover, of a lighter tint of colour, and varies considerably in the nature of the epidermis, which in this species is slight and fibrous, but in the *B. ovatus*, is thick, horny, and shining.

Species 210. (Mus. Cuming.)

BULIMUS OBLONGUS. *Bul. testa oblongo-ovata, vix umbi-*
licalata, nunc subampita, ventricosa, nunc par-
viore, crassâ, ponderosa, anfractibus convexis, plus
minusve costulato-striatis, labro reflexo; fulvo-spadicea,
columellâ labroque intenſe roseis.

THE OBLONG BULIMUS. Shell oblong-ovate, scarcely umbilicated, sometimes rather large, thin, and ventricose, sometimes smaller, thick, and ponderous, whorls convex, more or less sculptured with rib-like striae, lip reflected; fulvous-bay, lip and columella striae rose.

Helix oblonga, Müller, Verm. vol. ii. p. 86.

Bulinus hemastoma, Scopoli, Lamarek.

Hab. Eastern coast of South America.

This well-known species inhabits a wide range of country, and varies greatly in size and substance according to the local conditions of climate and vegetation, at one time large and ventricose like the specimen selected for illustration, at another smaller, thick, and ponderous, with the lip not unfrequently thickened to an extent truly remarkable.

M. D'Orbigny has some curious remarks on the habits and age of this species. "The *Bulinus oblongus*," says this intelligent traveller and naturalist, "buries itself deeply in the earth during the dry season, and comes forth at the time of the rains. Towards the close of the rainy season, it deposits its cretaceous eggs in the earth, and the young escape from them during the first rains which follow." The author then goes on to describe that about two or three years is occupied in the formation of the shell, when, having arrived at the age of maturity, the lip is reflected, and at a more advanced stage becomes more or less thickened according to circumstances. After an examination of what he considers to be annual marks of increase of growth, Mr. D'Orbigny arrives at the conclusion that the *Bulinus oblongus* lives for a period of about ten years.



211. A.



212. B.



212. C.



212. D.



213. A.



213. E.



214.

— 212. —

— 213. —

B U L I M U S.

PLATE XXXVI

Species 211. (Fig. a, Mus. Cuming, Fig. b and c,
Mus. Dennison.)

BULIMUS FOCILLATUS. *Bal. testā oblongo-ovatā, aufrac-
tibus sex, striis longitudinalibus arcuatis peculiariter
notatis, per marginem superiore crenulatis, crenulis
sudostriatis; columellā revolutā; apertura ampliā,
effusā, labro expanso; pallidē stramineā, vel purporas-
cente-spadicē, maculis fuscisque carneo-aigricantibus
interdum tinctis, columellā et apertura fave roseo-
purpureis.*

THE CHERISHED BULIMUS. Shell oblong-ovate, whorls
six in number, peculiarly marked with arched longi-
tudinal striae, crenulated along the upper margin,
crenules rather distant; columella rolled back;
aperture large, effused, lip expanded; pale straw
colour, or light salmon colour, sometimes painted
with bluish-black bands and blotches, columella and
interior of the aperture rose-purple.

Hab. Vera Cruz, South America.

A delicate and graceful species, extremely variable in its
external colouring, but constant so far as regards the rose-
purple painting of the aperture and columella.

Species 212. (Mus. Dennison.)

BULIMUS OVATUS. *Bal. testā subabbriviatō-ovatā, tumidā,
apice subobtusa; aufractibus quinque, longitudinaliter
rude striatis, apicem versus granulatis et subtiliter
radiatim liratis, labro fortiter reflexo; roseo-fusces-
cente, epidermali cornea olivacea nitente inducū,
apertura cærdescente-alba, labro vivido roseo.*

THE OVATE BULIMUS. Shell somewhat abbreviately
ovate, swollen, rather obtuse at the apex; whorls five
in number, longitudinally rudely striated, granulated
and finely radiately ridged towards the apex, lip
strongly reflected; rose-brown, covered with a shining
olive horny epidermis, aperture bluish-white, lip
bright rose.

BRUGUIERE, Enc. Mith. Diet. no. 33.

Holix ovata, Muller.

Bulla ovata, Chemnitz.

Hab. Brazil.

This fine species is remarkable for its curious abbreviately turgid growth, whilst it has a peculiar shining

horny epidermis, and the lip is mostly of an unusually deep
crimson rose.

Species 213. (Mus. Cuming.)

BULIMUS ABYSSORUM. *Bal. testā oblongo-ovatā, tenuiculā,
subfusiformi, umbilicatā, aufractibus sex, rugoso-striatis,
columellā revolutā, labro subterpanso; albida, rufo-
brunneae longitudinaliter striatā et maculata.*

THE ABYSS-INHABITING BULIMUS. Shell oblong-ovate,
rather thin, somewhat inflated, umbilicated, whorls
six in number, roughly striated, columella rolled back,
lip somewhat expanded; whitish, longitudinally
streaked and blotched with reddish-brown.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 308, pl.
39, f. 7, 8.

Hab. Pampa Ruis, eastern declivity of the Bolivian Andes
(amongst stones at the roots of Cacti); D'Orbigny.

Towards the base of the shell the roughly striated sculpture
produces almost the effect of obtuse granules.

Species 214. (Mus. Cuming.)

BULIMUS FENESTRATUS. *Bal. testā oblongo-ovatā, sub-
fusiformi, umbilicatā, aufractibus sex ad septem, pecu-
liariter undulatè corrugatā et sulcatā; columellā sub-
plicatā, obliquè recedente, plano-revolutā; albida, ma-
culis fuscisque fusco-nigricantibus conspicue fenestrata,
columellā et apertura fave roseo-purparascensibus.*

THE WINDOW-PAINTED BULIMUS. Shell oblong-ovate,
somewhat fusiform, umbilicated, whorls six to seven in
number, peculiarly undulately wrinkled and grooved;
columella slightly plaited, obliquely receding, flatly
rolled back; whitish, conspicuously latticed with
brownish-black blotches and bands, columella and
interior of the aperture like.

PFEIFFER, Pro. Zool. Soc. 1846. p. 29.

Hab. Mexico.

Perfectly distinguished from the preceding species in
form, whilst the undulating grooved sculpture is peculiar.

Species 215. (Mus. Cuming.)

BULIMUS HYGROHYLEUS. *Bal. testā subpyramidalis-
oblongā, Puerulaformi, ampliter umbilicatā, aufractibus*

BULIMUS.—PLATE XXXVI.

septem, longitudinaliter rugoso-striatis; columellā latā, subrecedente, labro latè plano-reflexo; albida, cæruleo-nigricante quinquefasciatā et maculatā, labro nitiōe albo.

THE DAMP-WOODS BULIMUS. Shell somewhat pyramidaly oblong, Partula-shaped, largely umbilicated, whorls seven in number, longitudinally roughly striated: columella broad, slightly receding, lip

broadly and flatly reflexed: whitish, encircled by five bands of bluish-black blotches, lip shining white. D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 311, pl. 40, f. 3—5.

Hab. Province of Santa Cruz, Bolivia (in damp forests): D'Orbiguy.

A very characteristic species readily distinguished from the preceding by the Partula-like growth of the aperture.



215 a



216 a



217 b



216



216 b



219



220



220



222



221

BULIMUS.

PLATE XXXVII.

Species 216. (Fig. *a* and *b*, Mus. Cuming.)

BULIMUS LEVUS. *Bul. testā pyramidalē-oblongā, subcylindrācā, crassissimulā, interdūm umbilicatā, anfractibūs septem, subrotundis, lēvibus; aperturā parvissimulā, subobtusā, labro fortiter reflexo; rufo-aureastrā aut luteā, fasciis rufo-olivaceis aut viridiōbus cingulatā, interdūm longitudinaliter maculatā aut strigatā, labro nitidō albo.*

THE LEFT **BULIMUS.** Shell pyramidaly oblong, somewhat cylindrical, rather thick, sometimes umbilicated, whorls seven in number, somewhat rounded, smooth; aperture rather small, nearly orbicular, lip strongly reflected; reddish orange or yellow, encircled with red-olive or green bands, sometimes longitudinally blotched or streaked, lip shining white.

BRUGUIERE, Enc. Mēth. Vcrs, vol. 1. p. 317.

Helix lava, Muller.

Hab. Amboina and the Moluccas.

Little need be added to assist in the recognition of this well-known species, which is of a more pyramidal growth than *B. citrinus*, and distinguished by a greater variety of colouring. It is mostly sinistral.

Species 217. (Mus. Cuming.)

BULIMUS LACTARIUS. *Bul. testā ocatā, ventricosimaculā, umbilicatā, spirā subacutissimā, anfractibūs sex ad septem, convexis, lēvibus, aut longitudinaliter tenuis-striatis; columellā latiusculā, labro simplici; cerino-albā, fuscō-rufescēte interdūm tinctā, floccis opaco-albīs irregulariōnis aspersā.*

THE MILK-SPOTTED **BULIMUS.** Shell ovate, rather ventricose, umbilicated, spire somewhat attenuated, whorls six to seven in number, convex, smooth, or longitudinally finely striated; columella rather broad, lip simple; waxy-white, sometimes tinged with smoky red, sprinkled with irregular flakes of opaque white.

MENKE, MSS.

Hab. Mexico.

The curious opaque-white flakes with which this shell is marked, constitute an important specific character.

Species 218. (Mus. Cuming.)

BULIMUS GOUDOTI. *Bul. testā ovalā, tenuissimā, pellucidā,*

September, 1848.

ventricosā, anfractibūs quinque, longitudinaliter vnde striatis, labro tenui, simplici; rufo-fuscescente, epidermide cornu olivaceā nitente inolutā, aperture fauce iridescente.

GOUDOT'S BULIMUS. Shell ovate, very thin, transparent, ventricose, whorls five in number, longitudinally rarely striated, lip thin, simple; pale reddish brown, covered with a shining olive horny epidermis, interior of the aperture iridescent.

PETIT, Mag. de Zool.

Hab. Eastern Cordilleras, New Granada.

An extremely delicate shell of little more substance than a glassy film, protected by a thick shining horny epidermis.

Species 219. (Mus. Cuming.)

BULIMUS DUFRESNII. *Bul. testā cylindraceo-ovalā, oleiforā, opice obtusā; anfractibūs quinque, apicea versus minute granulosā, per marginem crenulatā, columellā subrecedente, labro simplici; luteo-viridescēte, olivaceo-fusco fasciatā et lineatā.*

DUFRESNE'S BULIMUS. Shell cylindrically ovate, olive-shaped, obtuse at the apex; whorls five to six in number, minutely granulated towards the apex, crenulated along the margin, columella slightly receding, lip simple; yellowish green, banded and lined with dark olive brown.

LEACH, Zool. Misc. vol. 2. p. 153. pl. 120.

Hab. Van Dieman's Land.

The painting of this species is mostly represented by a conspicuous central band, with five longitudinally waved lines above and below it.

Species 220. (Mus. Cuming.)

BULIMUS CONTUSUS. *Bul. testā oblongo-ovatā, crassissimulā, sepiissimā sinistrali, anfractibūs septem, convexis, lēvibus, labro reflexo; albida, ustulato-fusco tinctā et hic illuc irregulariter longitudinaliter strigatā, labro albo, columellā et peritremate labri porparascente-fusca.*

THE BRUISED **BULIMUS.** Shell oblong-ovate, rather thick, most frequently sinistral, whorls seven in number, convex, smooth, lip reflected; whitish, stained and irregularly longitudinally streaked here

and there with burnt brown, lip white, columnella and outer edge of the lip purplish brown.

Hab. Eastern Islands.

This is unquestionably distinct from the *B. citrinus*, with which it has been hitherto confounded.

Species 221. (Mus. Cuming.)

BULIMUS CRASSILABRIS. *Bul. testā ovatā, cylindraceā, umbilicatā, anfractibus septem, convexis, longitudinālē crenellatā salcato-striatis, columellā latā, aperturā parviusculā, rotundatā, labro simplici; albā, anfractu-fusco late bifasciatā, versus apicem rosacea.*

THE THICKENED-LIP BULIMUS. Shell ovate, cylindrical, umbilicated, whorls seven in number, convex, longitudinally very closely sulyately striated, columnella broad, aperture rather small, rounded, lip simple; white, encircled by two broad bands of orange-brown, pink towards the apex.

GRAY, Pro. Zool. Soc. 1834, p. 66.

Hab. Madagascar.

The name assigned to this species is rather an inappropriate one, the lip being only slightly thickened and not reflected.

Species 222. (Mus. Cuming.)

BULIMUS PETITI. *Bul. testā conico-ovatā, glandiformi, umbilicatā, anfractibus sex, plano-convexis, longitu-*

disaliter rugoso-striatis, suturā subtiliter crenulatis, columellā latiusculā, labro tenui, simplici; albida, ustulato-fusco plus minusce tincta.

PETIT'S BULIMUS. Shell conically ovate, acorn-shaped, umbilicated, whorls six in number, flatly convex, longitudinally roughly striated, finely crenulated at the suture, columnella rather broad, lip thin, simple: whitish, more or less stained with burnt brown.

PFEIFFER, Pro. Zool. Soc. 1846, p. 31.

Hab. Chacopo, Bolivia.

Of rather light growth, crenulated at the suture in a manner extremely characteristic.

Species 223. (Mus. Cuming.)

BULIMUS CHLORIS. *Bul. testā cylindraceo-oblongā, subfusiformi, sinistrali, anfractibus sex ad septem, convexis, lăevibus aut longitudinaliter minutissimè striatis, columellā latā, subcallosā, aperturā parviusculā, labro reflexo; intense citrina-flavā, columellā labroque albī.*

THE CANARY-BIRD BULIMUS. Shell cylindrically oblong, somewhat fusiform, sinistral, whorls six to seven in number, convex, smooth or longitudinally very minutely striated, columnella broad, somewhat callous, aperture rather small, lip reflected; deep citron yellow, lip and columnella white.

Hab. Eastern Islands.

Though approximating very closely in general aspect to the *B. citrinus*, I feel strongly convinced that this is a distinct species: the difference of form is permanent.



BULIMUS.

PLATE XXXVIII.

Species 224. (Mus. Cuming.)

BULIMUS SEX-DENTATUS. *Bul. testā cylindraceo-oblongā,*

Prupiformis, basi carinatā et umbilicatā; anfractibus octo, piano-concisis, longitudinaliter corrugato-striatis et sulcatis, ultimo pone labrum biseriobimaculatum, labro subexpanso, aperturā subquadratā, sex-dentatā; cornucō-albida, apertura fance his illis purpureo-fusco tincta.

THE SIX-TOOTHED BULIMUS. Shell cylindrically oblong,

Pupa-shaped, keeled and umbilicated at the base; whorls eight in number, flatly convex, longitudinally striated and grooved in a wrinkled manner, last whorl with two indentations behind the lip, lip somewhat expanded, aperture rather square, six-toothed; horny white, interior of the aperture stained here and there with purple-brown.

Clavisilia sex-dentata, Spix, Test. Bras., pl. 14. f. 3.

Pupa sex-dentata, Wagner.

Hab. Brazil.

Of a peculiarly cylindrical form, grooved and striated in a wrinkled manner much after the fashion of a Pupa.

Species 225. (Mus. Cuming.)

BULIMUS PUNCTATISSIMUS. *Bul. testā cylindraceo-fusiformi, basi subcarinatā et umbilicatā, spirae suturis subimpressis; anfractibus octo ad noeum, transversis subtilissimis striatis et nivali superficialiter indentatis, labro latè expanso, aperturā subconstrictā, septendentatā; semipellucidus-albus, opaco-albus ad suturas, apertura fance purpureo-fusca.*

THE FINELY PUNCTURED BULIMUS. Shell cylindrically fusiform, slightly keeled and umbilicated at the base, suture of the spire somewhat impressed; whorls eight to nine in number, transversely very finely striated and everywhere superficially indented, lip broadly expanded, aperture somewhat contracted, seven-toothed; semitransparent white, opaque white at the sutures, interior of the aperture purple-brown.

Clavisilia punctatissima, Lesson, Voyage de la Coquille, pl. 15. f. 3.

Pupa septem-plicata, Rossmäler.

Hab. Brazil.

It is incorrect to say that this shell is punctured; the surface is covered with superficial indentations, and it is encircled throughout with very fine impressed striae.

Species 226. (Mus. Cuming.)

BULIMUS JANEIRENSIS. *Bul. testā cylindraceo-fusiformi, umbilicatā, anfractibus octo ad noeum, cornicis, nitidulis, striis obliquis, elevatis scutis, opaco-albis, fasciis longitudinali-*

dinalibus ornatis, evanescitā anti-plicatā; aperturā subquadratā, labro expanso, rosacea.

THE JANEIRO BULIMUS. Shell cylindrically fusiform, umbilicated, whorls eight to nine in number, horny, shining, ornamented with oblique opaque white striae and longitudinal waved bands, columella plaited; aperture somewhat square, lip expanded, pink.

SOWERBY, Conchological Illustrations, f. 97.

Hab. Janeiro, Brazil.

A pretty delicately coloured shell with merely a simple plait upon the columella.

Species 227. (Mus. Cuming.)

BULIMUS EXESUS. *Bul. testā oblongo-fusiformi, umbilicatā, anfractibus octo, subcentricosis, peculiariter corrugatis et longitudinaliter indentatis: labro latissime expanso, aperturā subquadrato-ovali, quadridecatā, dentibus lanuginosis, griseo-albicans, labro purpureo-rufescente.*

THE WORM-EATEN BULIMUS. Shell oblong-fusiform, umbilicated, whorls eight in number, somewhat ventricose, peculiarly wrinkled and longitudinally indented; lip very widely expanded, aperture somewhat squarely ovate, four-toothed, teeth laminated; greyish white, lip purple-red.

Clavisilia exesa, Spix, Test. Bras., pl. 14. f. 1.

Pupa exesa, Wagner.

Pupa rufous, Jay.

Odontostoma exesum, Beck.

Hab. Brazil.

A fine species in which the surface is peculiarly longitudinally indented, the lip widely expanded, and richly coloured, and the teeth of the aperture prominent.

Species 228. (Mus. Cuming.)

BULIMUS ODONTOSTOMA. *Bul. testā cylindraceo-elongatā, basi bicarinatā et subanguliter umbilicatā, anfractibus septem, concisis, obliquè erubescitā subtiliter striatis, striis elevatis, minutissimè granulatis; aperturā productā, septem-decatā, marginibus conjunctis: emarginatā, aperturā rubente.*

THE TOOTHED-MOUTH BULIMUS. Shell cylindrically elongated, double-keeled and rather largely umbilicated at the base, whorls seven in number, convex, obliquely very closely finely striated, striae raised, very minutely granulated; aperture produced, seven-toothed; coppery-brown, aperture reddish.

SOWERBY, Zool. Journ., vol. i. p. 59. pl. 5. f. 3.

Pupa odontostoma, Gray.

Cochlidion Gargantua, De Férussac.

Clausilia Gargantua, Deshayes.

Pupa erythrostoma, Menke.

Odontostoma Gargantua, Beck.

Macrodontes Sowerbyi, Swainson.

Hab. Brazil.

Distinguished by its delicate coppery brown colour, which obtains a satiny hue from the striae being granulated.

Species 229. (Mus. Cuming.)

BULIMUS GRAYANUS. *Bul. testá cylindraceo-elongatá, lateráliter compressá, subconcurvata, basi conspicuè carinatá et umbilicatá, anfractibus sex, striis spiralibus et obliquis undique decussatis, aperturá valde productatá, octo-dentatá, marginibus conjunctis; fuscá, aperturá livido-purpurascens.*

GRAY'S BULIMUS. Shell cylindrically elongated, laterally compressed, somewhat twisted, conspicuously keeled and umbilicated at the base, whorls six in number, decussated throughout with spiral and oblique striae, aperture very much produced, eight-toothed, margins joined: brown, aperture livid-purple.

PFEIFFER, Pro. Zool. Soc., 1845, p. 73.

Hab. Brazil.

This species has very much the appearance of being a distorted example of the preceding; it may prove distinct.

Species 230. (Mus. Cuming.)

BULIMUS PANTAGRELINUS. *Bul. testá cylindraceo-oblóngá, umbilicatá, anfractibus novem, subelevatis, longitudinaliter peculiariter corrugatá et indecatatá; aperturá subproductá, labro latissimè incrassato et expanso, iatus profusè lamello-dentato; cærulescente-albá, labro carneo-antracitá.*

PANTAGRELINUS' BULIMUS. Shell cylindrically oblong, umbilicated, whorls nine in number, somewhat ventricose, longitudinally peculiarly wrinkled and indented; aperture rather produced, lip very widely thickened and expanded, profusely lamellarly toothed; bluish-white, lip flesh-tinted orange.

Cochlodina Pantagrelina, Moricand, Mém. Genève, vol. vi. p. 542, pl. I, f. 7, 8.

Odontostoma Pantagrelinum, Beck.

Odontostoma leucotremum, Beck.

Hab. Brazil.

The largest species of this curious Brazilian group, distinguished by the bold reflection of the lip and singular development of teeth.

Species 231. (Mus. Cuming.)

BULIMUS PUPOIDES. *Bul. testá cylindraceo-oblóngá, umbilicatá, vix carinatá, anfractibus novem, plano-concavis, obliquè levissimè striatis; aperturá subproductá, quadrato-oblóngá, quadri-vel quinque-dentatá, superne bicanaliculatá, labro incrassato, expanso; corneo-albida, strigis castanis perplicatis obliquè picta.*

THE PUPA-LIKE BULIMUS. Shell cylindrically oblong, umbilicated, scarcely keeled, whorls nine in number, flatly convex, obliquely very finely striated; aperture somewhat produced, squarely oblong, four- or five-toothed, with two narrow channels at the upper part, lip thickened, expanded; horny white, obliquely painted with a very few chestnut-brown stripes.

Clausilia Pupoides, Spix, Test. Bras., pl. 14, f. 4.

Pupa infusa, Wagner.

Pupa fasciata, Potiez et Michaud.

Cochlodonta Listeri, De Féüssac.

Pupa Listeri, Gray.

Cyclodontina Pupoides and *Listeri*, Beck.

Bulimus vibrans, Spix.

Hab. Brazil.

A solid shell with a somewhat transparent aspect,

Species 232. (Mus. Cuming.)

BULIMUS WAGNERI. *Bul. testá cylindraceo-pyramidalis, umbilicatá, anfractibus undecim, levissimè rugosostriatis, suturis subrugibus; aperturá quadridentatá, labro simplici; corneo-albida, opaco-alba maculatá et flammatá.*

WAGNER'S BULIMUS. Shell cylindrically pyramidal, umbilicated, whorls eleven in number, very finely roughly striated, sutures somewhat rude, aperture four-toothed, lip simple; horny white, spotted and flamed with opaque white.

PFEIFFER, Monog. Hel. viv., vol. ii. p. 85.

Clausilia striata, Spix.

Pupa striata, Wagner.

Pupa turrula, Anton.

Pupa conspersa, Potiez and Michaud.

Pupa Spixii, D'Orbigny.

Cyclodontina striata, Beck.

Hab. Brazil.

I quote the *Clausilia striata*, Spix, (*Pupa Spixii*, D'Orbigny,) as synonymous with this species on the authority of Dr. Pfeiffer, but with some doubt, for D'Orbigny describes it as having the lip thickened and reflected.

Species 233. (Mus. Cuming.)

BULIMUS DENTATUS. *Bul. testá subobso-cylindracea, umbilicatá, anfractibus septem, concavis, obliquè exiliter striatis, aperturá dentis sex-dentatá, labro producto, vix reflexo; albida.*

THE TOOTHED BULIMUS. Shell somewhat stoutly cylindrical, umbilicated, whorls seven in number, convex, obliquely faintly striated, aperture thickly six-toothed, lip produced, scarcely reflected: whitish.

KING, Zool. Journal, vol. v. p. 310.

Hab. Brazil.

It is only to be presumed that this is the true *B. dentatus* of King.



235



236



237



238



239



240



241



242



243

B U L I M U S.

PLATE XXXIX.

Species 234. (Mus. Cuming.)

BULIMUS GRANADENSIS. *Bul. testá fusiformi-ovalá, tenui, vix umbilicata, anfractibus sex, ventricosis, irregulariter striatis; aperturá ovali, labro tenuissimo, acuto; cæruleo-olivante, luteo-aureario lincta, maculis oblongis irregularibus seriatim ornata.*

THE GRANADA BULIMUS. Shell somewhat fusiformly ovate, thin, scarcely umbilicated, whorls six in number, ventricose, irregularly striated; aperture oval, lip very thin, acute; bluish white, stained with yellowish orange, and ornamented with rows of irregular oblong blue-black spots.

PFEIFFER, Pro. Zool. Soc., 1847. p. 231.

Hab. Merida, New Granada; Funk.

This specimen described by Dr. Pfeiffer is apparently of immature growth, though it exhibits all the characters of a distinct species

Species 235. (Mus. Cuming.)

BULIMUS BADIUS. *Bul. testá ovatá, ventricosa, vix umbilicata, anfractibus sex, ventricosis, rugoso-striatis, ad suturas tenui-crenulatis; columellá revolutá, aperturá ovalá, labro tenui; fuligineo-badius fuscó fasciatis, fasciis interruptis.*

THE BAY BULIMUS. Shell ovate, ventricose, scarcely umbilicated, whorls six in number, ventricose, roughly striated, finely crenulated at the sutures; columella rolled back, aperture ovate, lip thin; light fulvous bay, encircled with interrupted brown bands.

SOWERBY, Pro. Zool. Soc., 1834, p. 141.

Hab. Province of Xagua, Peru; Matthews.

There is a considerable resemblance in the colouring of this species to that of the *B. thomasiensis*.

Species 236. (Mus. Cuming.)

BULIMUS PAPYRACEUS. *Bul. testá fusiformi-ovatá, tenui, vix umbilicata, anfractibus sex, subcentricosis, longitudinaliter subtilissimè plicato-striatis; columellá revolutá, aperturá ovali, labro subexpanso, tenui, acuto; albá, cæruleo-nigrante undato-umbelata, striis opacis albis.*

THE PAPER BULIMUS. Shell fusiformly ovate, thin, scarcely umbilicated, whorls six in number, somewhat ventricose, longitudinally very finely plicated; striated; columella rolled back, aperture oval, lip somewhat expanded, thin, sharp; white, clouded in a waved manner with blue-black.

Helix papryacea, Mawe.

Cochlogena litu, De Féussac.

Bulimus littoralis, Spix.

Bulimus lita, Deshayes.

Bulimus inflatus, Spix.

Bulimus magus, Wagner.

Hab. Brazil.

The shell named *B. magus* and *inflatus*, as above recorded, is another state of the species of larger size.

Species 237. (Mus. Cuming.)

BULIMUS TRI-FASCIATUS. *Bul. testá subpyramidalis-ovalá, solidulá, umbilicata, anfractibus sex, lacibis ant obsoletè striatis, aperturá suborbiculari, labro conspicuè reflexo; albâ, rufo-fusco basia versus fasciata, apicem versus cinereo punctulata, labro albo.*

THE THREE-BANDED BULIMUS. Shell somewhat pyramidalis-oval, rather solid, umbilicated, whorls six in number, smooth or obsoletely striated, aperture rather orbicular, lip conspicuously reflexed; whitish, banded with reddish brown towards the base, finely dotted with ash colour towards the apex, lip white.

Helix trifasciata, Chemnitzi, vol. ix. p. 153. pl. 134. f. 1215
Cochlogena trizonalis, De Féussac.

Bulimus zonatus, Swainson.

Hab. Ceylon; Gardner.

The shell here represented agrees much better with Mr. Swainson's figure, *B. zonatus*, Zool. Illus., pl. 17, than with Chemnitzi's above quoted. I believe, however, with Dr. Pfeiffer, they are varieties of the same species.

Species 238. (Mus. Cuming.)

BULIMUS MULTICOLOR. *Bul. testá conico-ovatá, suboblique convoluta, umbilicata, anfractibus quinque, sub lete minutiissimè granulatis, ultimè ventricosa; aperturá subobliquè oblonga, labro tenuicolo, vix reflexo; opaco-albâ, olivaceo-tinctâ, flammis purpureo-fusca variegata, apertura favea violascens, labro cividè rosea.*

THE MANY-COLOURED BULIMUS. Shell conically ovate, somewhat obliquely convoluted, umbilicated, whorls five in number, very minutely granulated under the lens, last whorl ventricose; aperture rather obliquely oblong, lip rather thin but slightly reflexed; opaque white, olive-stained, variegated with purple-brown flames, interior of the aperture violet, lip bright rose.

Helix multicolor, Rang, Annales des Sci. Nat., 1831, p. 55
pl. 3. f. 1.

Gonyostoma multicolor, Beck.
Hab. Brazil.

The last whorl of this pretty species is distinguished by a peculiar swollen oblique growth.

Species 239. (Mus. Cuming.)

BULIMUS MIERSI. *Bul. testā fusiformi-oblongā, latē umbilicatā, spirā acuminatā, anfractibus sex, lēvibus, sub lente seriatim minutiissimè punctulatis; aperturā oblongo-ovali, labro conspicuā reflexo; pallidē castaneā, opaco albo variegatā, intus purpureo-violascente, labro rividō roseo.*

MIER'S BULIMUS. Shell fusiformly oblong, broadly umbilicated, spire acuminated, whorls six in number, smooth, very minutely punctured in rows beneath the lens; aperture oblong-oval, lip conspicuously reflected; light chestnut, variegated with opaque white, interior tinged with purple-white, lip bright rose.

SOWERBY, Conch. Illus., f. 90.

Hab. Brazil.

Dr. Pfeiffer errs greatly in regarding this as a variety of the preceding species; it has certainly a general resemblance to it, but, as an attentive comparison of the foregoing descriptions will show, it is specifically distinct throughout.

Species 240. (Mus. Cuming.)

BULIMUS CRICHTONI. *Bul. testā sulcissimā, crassisculā, anfractibus ser ad septem, longitudinaliter cerebrimē rūdē plicato-liratis; columellā callosā, basi retrorsum contortā, aperturā oblongā, basi canaliculatā, labro incrassato, expanso, vix reflexo; flavicante-albā, fuscō prope apicem exiliiter flammatā, olivaceo-viridi inter tiras fasciātā et irregulariter tinctā, columellā vivide purpureo-roseā.*

CRICHTON'S BULIMUS. Shell somewhat fusiform, rather thick, whorls six to seven in number, longitudinally very closely rudely plicately ridged; columella callous, twisted backwards at the base, aperture oblong, channelled at the base, lip thickened, expanded, scarcely reflected; yellowish white, faintly flamed with brown near the apex, banded and irregularly stained with olive-green between the ridges, columella deep purple-rose.

BRODERIP, Pro. Zool. Soc., 1836, p. 44.

Hab. Ambu, near Huamueo, Peru, Sir Alexander Crichton.

The channel of the base of this remarkable species, of which this is, I believe, the only specimen known, is thought by Mr. Broderip to be a distortion.

Species 211. (Mus. Cuming.)

BULIMUS FULVICANS. *Bul. testā conico-ovatā, tenui, um-*

bilicatā, anfractibus sex ad septem, subrotundatā, cerebrimē spiraliter elevato-striatis, striis aliis minutis obliquè decussatis; columellā plano-revolutā, labro simplici, acuto; undique rufescente-fulvā.

THE FULVOUS BULIMUS. Shell conically ovate, thin, umbilicated, whorls six to seven in number, somewhat rounded, very closely spirally elevatedly striated, obliquely crossed with other minute striae; columella flatly rolled back, lip simple, sharp; reddish fulvous throughout.

PFEIFFER, Monog. Hel. viv., vol. 2, p. 99.

Hab. Seychelle Islands.

Of a uniform reddish fulvous colour, closely spirally striated in a very characteristic manner.

Species 242. (Mus. Cuming.)

BULIMUS JUSSIEUI. *Bul. testā oblongo-ovatā, tenuiculā, vix umbilicatā, anfractibus sex, longitudinaliter rūdē striatis, transversiū sub lente obscurè seriatim pauculatis; columellā revolutā, labro simplici, acuto; corneo-fuscī, hic illuc saturatim strigata.*

JUSSIEU'S BULIMUS. Shell oblong-ovate, rather thin, scarcely umbilicated, whorls six in number, longitudinally rudely striated, transversely obscurely finely punctured under the lens in rows; columella rolled back, lip simple, sharp, flesh-tinted brown, streaked here and there of a darker colour.

VALENCIENNES, MSS. Pfeiffer, Pro. Zool. Soc., 1846, p. 33.
Hab. Cusoo.

Resembling *B. granulosus* somewhat in general aspect.

Species 243. (Mus. Cuming.)

BULIMUS MELO. *Bul. testā ovatā, tenuiculā subconcrecā, vix umbilicatā, anfractibus sex, longitudinaliter rūdē striatis, ad suturas subuplicatis; columellā revolutā, labro simplici, tenui; intus exlusus nigriceunte-purpureus, extus fulvo longitudinaliter variè flammatus et striatus.*

THE MELON BULIMUS. Shell ovate, rather thin, ventricose, scarcely umbilicated, whorls six in number, longitudinally rudely striated, slightly plicated at the sutures; columella rolled back, lip simple, thin; blackish-purple within and without, variously longitudinally flamed and striated without with fulvous colour.

QUOY and GAIMARD, Voy. de l'Astrol. Zool., vol. ii, p. iii.
pl. 9, f. 8, 9.

Hab. New Holland.

Varied externally with dark brown and yellowish flames, and tinged with rich purple brown in the interior.



BULIMUS.

PLATE XL.

Species 244. (Mus. Cuming.)

BULIMUS MEXICANUS. *Bul. testā acuminatō-ovatā, tenui, vix umbilicatā, anfractib⁹ sex ad septem, longitudinaliter tenuistriotis; columellā reflexā, labro tenui, simplici; albidā, nigricante-fusco tri-quadrī-zonatā.*

THE MEXICAN BULIMUS. Shell acuminate-ovate, thin, but slightly umbilicated, whorls six to seven in number, longitudinally finely striated; columella reflected, lip thin, simple; whitish, encircled by three or four zones of blackish brown.

LAMARCK, Anūm. sans vert., (Deshayes' edit.) vol. vii. p. 232.

Cochlogena vittata, De Férrussac.

Hab. Mexico.

Approximating in general aspect to the *B. Caribaeorum* of the West Indies.

Species 245. (Mus. Cuming.)

BULIMUS LUCIDUS. *Bul. testā oblongo-ovatā, tenui, cornuā, pellucida, vix umbilicatā, anfractib⁹ sex, concavis, levibus, nitentibus, sub teste crenulariā spiraliā impresso-striatis; columellā reflexā, labro simplici, acuto; pallide stramineā.*

THE CLEAR BULIMUS. Shell oblong ovate, thin, horny, transparent, scarcely umbilicated, whorls six in number, convex, smooth, shining, very closely spirally impressed striae.

Hab. St. Vincents, Guilding.

This shell has the name *fragilis* attached to it in Mr. Cuming's collection, but it is not the *B. fragilis* of Lamarck, which, according to M. Deshayes, is the young of the British *Lymnaea stagnalis*.

Species 246. (Mus. Cuming.)

BULIMUS VIMINEUS. *Bul. subpyramidalis-ovatā, basi subobliquē productā, vix umbilicatā, anfractib⁹ septem, convexis, glabris; columellā reflexā, aperturā obliquā angulatā-fusā, labro subexpanso, non reflexo; albā, late et cinereo-carulescente conspicue longitudinaliter vittatā, aperturā fince castaneo-fusca, labro albo.*

THE WICKER BULIMUS. Shell somewhat pyramidal, obliquely produced at the base, scarcely umbilicated, whorls seven in number, convex, smooth,

columella reflected, aperture obliquely angularly effused, lip somewhat expanded, not reflected; white, conspicuously longitudinally filleted with stripes of ashy blue and yellow, interior of the aperture chestnut-brown, lip white.

Helix virescens, Moricand, Mém. de Genève, vol. vi. p. 540 pl. 1. f. 5.

Hab. Brazil.

A very characteristic species both in respect of form and colouring.

Species 247. (Mus. Cuming.)

BULIMUS UNDULATUS. *Bul. testā subfusiformi-ovatā, subumbilicatā, anfractib⁹ sex ad septem, convexisculis, glabris, ultimo basi obtusè angulatis; columellā reflexā, aperturā oblongo-ovali, labro acuto, vix reflexo; albida, late et livido-brunnea conspicue undulata.*

THE WAVED BULIMUS. Shell somewhat fusiformly ovate, umbilicated, whorls six to seven in number, rather convex, smooth, last whorl obtusely angled at the base; columella reflected, aperture oblong-oval, lip sharp, but slightly reflected; whitish, conspicuously waved with yellow and livid brown.

Bulinulus undulatus, Guilding, Zool. Journ., vol. iv. p. 169. Hab. St. Vincents, West Indies.

A prettily striped species described some years since by the Rev. Lansdowne Guilding, but still rare.

Species 248. (Mus. Cuming.)

BULIMUS ANGULATUS. *Bul. testā oendo-fusiformi, subumbilicatā, basi attenuatā, anfractib⁹ octo, concavis, levibus; columellā latè reflexā, aperturā infernè canaliculatā, labro subexpanso, non reflexo; cernilescente-albā, striis fusco longitudinalibus hic illie irregulariter pictā, strigā saturatiore pone labrum, labro albo, apice nigro.*

THE ANGLED BULIMUS. Shell ovately fusiform, umbilicated, attenuated at the base, whorls eight in number, convex, smooth; columella broadly reflected, aperture channeled at the lower part, lip somewhat expanded, not reflected; bluish-white, painted irregularly here and there with brown streaks, with a darker streak behind the lip, lip white, apex black.

WAGNER, Spix, Test. Bras., p. 14. pl. 13. f. 3.

Hab. Brazil.

Distinguished by the aperture being attenuately prolonged at the base, so as to form a channel.

Species 249. (Mus. Cuming.)

BULIMUS FUSIFORMIS. *Bul. testā elongato-fusiformi, umbilicatā, basi attenuatā, compressā et retrosum subcontortā, anfractibus novem ad decem, plano-convergis, longitudinaliter obscure striatis; columellā latē reflexā aperturā subelongatā, basi latē canaliculatā, labro expanso non reflexo; fuscostrigata, strigis fuscis longitudinalibus hic illuc irregulariter pictā, strigā saturatione pone labrum.*

THE FUSIFORM BULIMUS. Shell elongately fusiform, umbilicated, attenuated at the base, compressed and slightly twisted backwards, whorls nine to ten in number, flatly convex, longitudinally obscurely striated; columella broadly reflected, aperture somewhat elongated, broadly channelled at the lower part, lip expanded, not reflected; pale brown, painted irregularly here and there with brown streaks, with a darker streak behind the lip.

Helix fusiformis, Rang, Ann. des Sci. Nat., p. 60. pl. 3. f. 2.
Bulimus dubiosus, Jay.

Bulimus angulatus, var? Deshayes.
Hab. Brazil

There is a remarkable affinity between this and the preceding species, which is not shared by any other; they are nevertheless permanently distinct.

Species 250. (Mus. Cuming.)

BULIMUS DESHAYESI. *Bul. testā elongato-turrīta, umbilicatā, anfractibus novem, rotundatis, subtiliter impresso-striatis, ad suturas obscure crenulatis, columellā reflexā, aperturā pareā, simplici; violaceo-albā, maculis strigisque viridi-fusca irregulariter pictā, apertura fauce roseo-purpurea.*

DESHAYES' BULIMUS. Shell elongately turreted, umbilicated, whorls nine in number, rounded, finely impressed striated, obscurely crenulated at the sutures, columella reflected, aperture small, simple; violetting white, irregularly painted with greenish-brown streaks and blotches, interior of the aperture rose-purple.

PFEIFFER, Pro. Zool. Soc., 1845, p. 73.
Hab. Venezuela; Dyson.

Quite unlike any other species at present known.

Species 251. (Mus. Cuming.)

BULIMUS LOXENSIS. *Bul. testā cylindraceo-oblongā, una-*

bilicatā, anfractibus septem, convexis, lēvibus, aut obscure striatis; columellā latē reflexā, aperturā parvissula, labro simplici, acuto; cōrvescente-albā, nigricante-castaneo hic illic longitudinaliter maculatā aut strigatā.

THE LOXA BULIMUS. Shell cylindrically oblong, umbilicated, whorls seven in number, convex, smooth, or obscurely striated; columella broadly reflected, aperture rather small, lip simple, sharp; bluish-white, here and there longitudinally streaked and blotched with blackish chestnut.

PFEIFFER, MSS., Mus. Cuming.
Hab. El Catamajá, Loxa, Quito; Hartweg.

Of a peculiarly bluish-white, painted in a very irregular manner with dark more or less interrupted streaks.

Species 252. (Mus. Cuming.)

BULIMUS SERPERASTRUS. *Bul. testā conico-oblongā, umbilicatā, anfractibus septem, convexiusculis, lēvibus; columellā reflexā, labro subexpanso, viz reflexo; albida, fasciis ad sex cōrveo-fascis irregularibus, interruptis, cingulatā.*

THE BANDAGED BULIMUS. Shell conically oblong, umbilicated, whorls seven in number, slightly convex, smooth; columella reflected, lip somewhat expanded, scarcely reflected; whitish, encircled with about six irregular interrupted bluish-brown bands.

SAY, Descr. New Terr. Shells, p. 25.
Hab. Mexico.

Banded in a particular manner, whilst the species is also peculiar in form.

Species 253. (Mus. Cuming.)

BULIMUS DURUS. *Bul. testā subgloboso-ovatā, umbilicatā, anfractibus septem, rotundatis, striis longitudinalibus et transversis subtilissimè decussatā; columellā reflexā, labro simplici, viz reflexo; fulcicante, medio albizonatā.*

THE HARD BULIMUS. Shell somewhat globosely ovate, umbilicated, whorls seven in number, rounded, very finely decussated with longitudinal and transverse striae; columella reflected, lip simple, scarcely reflected; pale fulvous colour with a white zone round the middle.

SPIX, Test. Bras., p. 5. pl. 6. f. 2.
Hab. Brazil.

Very closely allied in form to dwarf specimens of *B. heterotrichus*.



BULIMUS.

PLATE XLI.

Species 254. (Mus. Dennis.)

BULIMUS SUBSEMICLAUSUS. *Bul. testā suboblongato-ovatā, tenuiculā, ventricosā, anfractibus quinque, levibus, vel sub teste minutissimè impresso-striatis, anfractu ultimo, subangulato; columellā depressā, labro peculiariter plano-reflexo; lacteo-albidā, obscurè fuscocingulatā, punctis opaco-albis fuscocubratis notatā, castanā circa columellam.*

THE PARTIALLY-CLOSED BULIMUS. Shell somewhat abbreviate ovate, rather thin, ventricose, whorls five in number, smooth or very minutely impressively striated under the lens, last whorl slightly angulated; columella depressed, lip peculiarly flatly turned inwards; yellowish-white, obscurely belted with brown, marked with brown-shaded opaque white dots, chesnut around the columella.

PETIT, Magasin de Zoologie, 1843, Moll. pl. 66.

Hab. Santa Fé di Bogota.

This singular species may at once be recognised by the curious inwardly reflected growth of the lip.

Species 255. (Mus. Cuming.)

BULIMUS MARMARINUS. *Bul. testā subcylindraceo-oblongā, umbilicatā, anfractibus sex ad septem, concentricā subrade striatis; columellā subuplicatā et latè plano-reflexā, labro vix reflexo; albidā, castaneo-fasciata muciculatā et longitudinaliter undulata.*

THE MARBLED BULIMUS. Shell somewhat cylindrically oblong, umbilicated, whorls six to seven in number, concomitantly somewhat rudely striated; columella slightly plaited, and broadly flatly reflected, lip scarcely reflected; whitish, spotted in a banded manner and longitudinally waved with chesnut.

D'ORBIGNY, Voy. dans l'Amér. Mérid., p. 310. pl. 39. f. 11, 12.

Hab. Yungas, Bolivia (under dead leaves and stones in woods near the river Meguilla); D'Orbigny.

Characterized by a waved pattern in the painting as well as by a peculiar roundness of form.

Species 256. (Mus. Cuming.)

BULIMUS CANALICULATUS. *Bul. testā subfusiforme-ovatā, basi peculiariter compresso-carinatā et umbili-*

cata, anfractibus sex ad septem, convexis, longitudinaliter rugoso-striatis; columellā reflexā, apertura oblongā, inferne conspicue canaliculatā; albidā, purpurascente-spadiceo strigata et marmorata, intus purpurascens.

THE CHANNELLED BULIMUS. Shell somewhat fusiformly ovate, peculiarly compressly keeled and umbilicated at the base, whorls six to seven in number, convex, longitudinally roughly striated; columella reflected, aperture oblong, conspicuously channelled at the lower part; whitish, streaked and marbled with purplish bay, interior purplish.

PFEIFFER, Pro. Zool. Soc., 1845. p. 68.

Hab. New Granada.

Chiefly remarkable on account of its abruptly contracted growth at the base, which looks pushed inwards as it were.

Species 257. (Mus. Adamson.)

BULIMUS LYONNETIANUS. *Bul. testā pyramidalis-conicā, basi latissimè et abruptè angulatā, profunde umbilicatā, anfractibus octo, longitudinaliter rugoso-striatis, striis apice versus elevatioribus et regulariōbus, anfractu ultimo sinistrorsum gibbosso-producto; apertura ovata, labro reflexo; albidā, epidermide flavicante indutā.*

LYONNET'S BULIMUS. Shell pyramidalis conical, very broadly and abruptly angled at the base, deeply umbilicated, whorls eight in number, longitudinally roughly striated, stria more elevated and regular towards the apex, last whorl gibbously produced on the left side; aperture ovate, lip reflected; whitish, covered with a yellowish epidermis.

Helix Lyonnetiana, Pallas, Spicilegia Zoologica, vol. x. pl. 3. f. 7, 8.

Trochus Lyonnetianus, Chemnitz.

Trochus distortus, Guenlin.

Helix distorta, Burrows.

Pupa nodiolinus, Bowditch.

Hab. Mauritius.

This remarkable species, which partakes very much of the character of certain *Pupa*, was originally named by Pallas, in honour of M. Lyonnet, an eminent anatomist and anatomical engraver of the last century, resident at the Hague.

Species 258. (Fig. *a* and *b*, Mus. Cuming.)

BULIMUS NAVICULA. *Bul. testā abbreviato-conicā, basi peculiariter plano-truncatā, subobscnē umbilicatā, anfractibus sex, subtiliter striatis et corrugato-malleatis, ultimo dextrorsum valde producto; aperturā infernū canaliculatā, labro subreflexo; albida, carneo-fusco marmoratā, serpē intense roseo-castaneo latifasciatā, labro churvo.*

THE LITTLE BARK BULIMUS. Shell abbreviately conical, peculiarly flatly truncated at the base, somewhat obscurely umbilicated, whorls six in number, finely striated and indented in a wrinkled manner, last whorl much produced towards the right; aperture channelled at the lower part, lip slightly reflexed; whitish, marbled with bluish brown, often broadly banded with deep rose-chesnut, lip ivory white.

HELIX NAVICULA, Wagner, Spix, Test. Bras., pl. 15. f. 2, 3.
Bulinus auris-leporis, Féussac.

Bulinus lateralis, Menke.

Otosoma navicula, Beck.

Hab. Bahia, Brazil; Spix, Moricand.

Amply distinguished by its extraordinary mode of growth, of which an approximation is offered in the species which follows.

Species 259. (Fig. *a* and *b*, Mus. Cuming.)

BULIMUS AURIS-LEPORIS. *Bul. testā subpyramidalis-ovatā, basi depresso-convexā, umbilicatā, anfractibus sex, subtiliter striatis et corrugato-malleatis, anfractū ultimo dextrorsum subproducto, aperturā descendente, labro expanso, vis reflexo; albā, nigro vel fusco marmoratā aut striatā, infernū nufasciatā, labro churvo.*

THE HARE'S-EAR BULIMUS. Shell somewhat pyramidal ovate, depresso convex, umbilicated, whorls six in number, finely striated and indented in a wrinkled manner, last whorl slightly produced towards the right, with the aperture descending, lip expanded, scarcely reflexed; white, marbled or streaked with black or brown, with a single band round the lower part, lip ivory white.

BRUGUIERE, Enc. Méth., vol. i. p. 346.

Auricula leporis, Lamarck.

Cochlogena auris-leporis, Féussac.

Pupa auris-leporis, Gray.

Bulinus lagotis, Menke.

Otosoma lagotis, Beck.

Hab. Brazil.

The plan of growth and detail of sculpture are the same

as in the *B. navicula*, in which species they may be said to be fancifully exaggerated.

Species 260. (Mus. Cuming.)

BULIMUS COARCTATUS. *Bul. testā conico-ovatā, solidulā, umbilicatā, anfractibus sex ad septem, levibus, ant sub lente subtiliter decussatim striatis; columellā incrassatā, reflexā, aperturā coarctatā, labro planè expanso, latissimo; albā, superne lineis interruptis, infernū fasciis, fusco-nigricantibus cingulatā, apice nigro, labro churvo.*

THE CONTRACTED BULIMUS. Shell conically ovate, rather solid, umbilicated, whorls six to seven in number, smooth under the lens, finely decussately striated; columella thickened, reflected, aperture contracted, lip flatly expanded, rather broad; white, encircled round the upper part with brown-black interrupted lines, round the lower with bands, lip ivory white.

PEIFFER, Pro. Zool. Soc., 1845, p. 73.

Hab. —?

Somewhat *Partula*-like in aspect, though not possessing the characters of that genus.

Species 261. (Mus. Cuming.)

BULIMUS AURIS-MURIS. *Bul. testā conico-ovatā, basi depresso-planā, subobscnē umbilicatā, anfractibus sex ad septem, planiscaulis, levibus, sub tenui subtiliter malleatis; columellā latē reflexā, aperturā lateraliter planō-contortā, subtrigonā, labro reflexo; albā, lineis carneo-nigricantibus cingulatā, labeculis pallidioribus obliquis.*

THE MOUSE'S-EAR BULIMUS. Shell conically ovate, depresso flattened at the base, somewhat obscurely umbilicated, whorls six to seven in number, rather flattened, smooth, finely indented under the lens; columella broadly reflected, aperture flatly contracted at the side, somewhat triangular, lip reflexed; white, encircled by narrow bluish-black bands, and oblique dashes of a lighter hue.

Helix auris-muris, Moricand, Mém. de Genève, vol. viii. p. 140. pl. 3. f. 1-3.

Otosoma myotis, Beck.

Hab. Bahia, Brazil.

Although partaking of the form and minute detail of sculpture presented in the *B. navicula* and *auris-leporis*, this is a very distinct and characteristic species.



BULIMUS.

PLATE XLII.

Species 262. (Mus. Cuming.)

BULIMUS VERANYI. *Bul. testā conico-globosā, spirā brevi, asfractibus quatuor ad quinque, ventricosā, cerebrimē minutē granulatis; columellā subuplicatā, labro reflexo; fulvā, fuso subindistincte punctatā, striis albidi sanguinis, epidermide tenui indutā, labro corneo-albido.*

VERANY'S BULIMUS. Shell conically globose, spire short, whorls four to five in number, ventricose, very closely minutely granulated; columella slightly plaited, lip reflected; yellowish, somewhat indistinctly dotted with brown, marked with white lightning-like streaks, and covered with a thin epidermis, lip flesh-tinted white.

PFEIFFER, Pro. Zool. Soc., 1847, p. 230.

Hab. Chacopo, Province of Merida, New Granada.

In some specimens the white zigzag streaks are not present.

Species 263. (Mus. Cuming.)

BULIMUS QUADRICOLOR. *Bul. testā tenui, sub-depresso-ovalā, Succineiformi, spirā brevi, asfractibus quatuor, obliquè elevato-striatis; columellā tortuosā, apertura ampliā, labro tenuiter reflexo; pellucido-cornueā, luteā, fuso nuce undulatā, nuce punctatā, epidermide tenui luteo-punctatā indutā, epidermide sericeā secundā in strijs longitudinalibus antorrum peculiariter serratis, labro purpureo-roseo.*

THE FOUR-COLOURED BULIMUS. Shell somewhat depressed-oval, Succinea-shaped, spire short, whorls four in number, obliquely elevatedly striated; columella twisted, aperture large, lip thinly reflected; transparent-horny, yellow, sometimes waved, sometimes dotted with brown, covered with a thin yellow-dotted epidermis, with a second light silken epidermis in longitudinal streaks peculiarly serrated along the anterior edge, lip purple-rose.

PFEIFFER, Pro. Zool. Soc., 1847, p. 229.

Hab. Chacopo, Province of Merida, New Granada.

A delicate and beautiful species collected in the same locality with the preceding by M M. Funck and Linden.

Species 264. (Mus. Cuming.)

BULIMUS SUCCINOIDES. *Bul. testā depresso-ovalā, Sue-*

cineiformi, spirā brevissimā, asfractibus quatuor, obliquè striatis et cerebrimē minutē granulatis; columellā tortuosā, apertura perampiā, labro reflexo; subcornueā, lutescente, fasciā suturali castaneā, epidermide olivaceo-fusca indutā, epidermide secundā sericeā in strijs brevibus obliquis e suturis descendentibus, labro albido.

THE SUCCINEA-LIKE BULIMUS. Shell depressedly ovate, Succinea-shaped, spire very short, whorls four in number, obliquely striated and very closely minutely granulated; columella tortuous, aperture very large, lip reflected; somewhat horny, yellowish, with a dark chestnut sutural band, covered with an olive brown epidermis, and a second silken one descending from the suture in short oblique streaks, lip whitish,

PETIT, Magasin de Zoologie, 1841, Moll. pl. 31.
Hab. Santa Fé di Bogota.

The form of this species, contrasted with that of *B. Lyoni-*
nianus and *navicula*, is remarkable.

Species 265. (Fig. *a* and *b*, Mus. Sowerby; Fig. *c*,
Mus. Cuming.)

BULIMUS CATHCARTIÆ. *Bul. testā ovalā, ventricosā, spirā brevi, asfractibus quinque, obliquè subtilissimè striatis, asfractū ultimo maximo, apertura patulā, labro sublatè reflexo; albida vel purpurascente-castaneā, epidermide luteo-olivacea indutā, epidermide sericeā secundā, plus minusculæ pulcherrimæ et variè configuratā, intus purpu-*

rascente-castaneā, labro rosso.

CATHCART'S BULIMUS. Shell ovate, ventricose, spire short, whorls five in number, obliquely very finely striated, last whorl very large, aperture wide open, lip somewhat broadly reflected; whitish, or purplish chestnut, covered with a yellowish-olive epidermis, and a second silken one which is more or less beautifully and variously configured, interior purple-chestnut, lip pink.

Hab. Province of Merida, New Granada; Linden.

In addition to the beautiful association of form and colour in this species, it is further distinguished by a remarkable configuration of the outer membrane of the epidermis, which is variously disposed in different specimens. The *B. Cathcartiæ* resembles in some degree the *B. pintoiensis*, D'Orbigny, Moll. Voy. dans l'Amér. Mérid.

of which the author informs us he collected a single dead specimen in a damp forest at the base of the Eastern Cordilleras of Bolivia, but possesses neither the form nor marking of that species.

I am indebted to Mr. Linden, the botanical traveller in New Granada, for the possession of several examples of this chaste and beautiful shell, characterized by different patterns of epidermis, and have thought it a subject not unworthy of being dedicated to an eminent collector in North Britain, the Honourable Mrs. Catheart, distinguished by her zeal and affability.

Species 266. (Mus. Dennison.)

BULIMUS GLANDIFORMIS. *Bul. testā ovatā, ventricosō, vix umbilicatā, anfractibus quatuor ad quinque, rugoso-striatis et granulatis, columellā subtortuōs, labro sublatē reflexo; fumeo-albidā, strigis brevibus fuscis e suturis fulgaratā, epidermide olivaceā, luteo-punctatā et marmoratā, indutā.*

THE ACORN-SHAPED BULIMUS. Shell ovate, ventricose, scarcely umbilicated, whorls four to five in number, roughly striated and granulated, columella slightly twisted, lip rather broadly reflected; smoky-white, marked with short zigzag brown streaks out of the sutures, covered with an olive epidermis, dotted and mottled with yellow.

LEA. Extr. Trans. Nat. Hist. Soc. Philadelphia, pl. 23. f. 92.

Hab. New Granada.

It is not without some doubt that I attribute this shell to Mr. Lea's *B. glandiformis*, whose figure is very obscure.

Species 267. (Mus. Dennison.)

BULIMUS PULICARIUS. *Bul. testā ovatā, tenuiculā, vix umbilicatā, anfractibus quatuor ad quinque, inflatis, subtiliter granulatis, longitudinaliter rugoso-striatis, transversim obscurè sulcatis; columellā latē reflexa,*

aperturā subamplyā, labro reflexo; albidā, castaneo-irregulariter punctatā, labro et interdum fauce aper-turā purpurascente-rosis, epidermide olivaceo-fusciaut viridescente indutā.

THE FLEA-SPOOTTED BULIMUS. Shell ovate, rather thin, scarcely umbilicated, whorls four to five in number, inflated, finely granulated, longitudinally roughly striated, transversely obscurely grooved; columella broadly reflected; whitish, irregularly dotted with chestnut, lip and sometimes the interior of the aperture purple-rose, covered with an olive-brown or greenish epidermis.

Hab. New Granada.

A delicate species more or less irregularly dotted with black, and richly coloured with rose-purple about the lip and aperture.

Species 268. (Mus. Cuming.)

BULIMUS PLECTOSTYLOS. *Bul. testā ovatā, crassiusculā, vix umbilicatā, anfractibus quatuor ad quinque, rugoso-striatis et minuti granulatis, columellā tortuosa et fortiter plicatā, labro incrassato, reflexo; fumeo-castaneā, punctis perpaucis nigris obscurè notata, epidermide fusca indutā, epidermide secundā albā in strigis brevibus e suturis fulgaratā.*

THE TWISTED-COLUMN BULIMUS. Shell ovate, rather thick, scarcely umbilicated, whorls four to five in number, roughly striated and minutely granulated, columella twisted and strongly plaited, lip thickened, reflected; smoky chestnut, obscurely marked with a few black dots, covered with a brown epidermis, and a second white epidermis emerging from the sutures in short zigzag streaks.

PFEIFFER, Pro. Zool. Soc., 1847, p. 230.

Hab. Chacopo, Province of Merida, New Granada; Funec.

From the same locality as the preceding species, remarkable for its twisted columella.



273 b.

BULIMUS.

PLATE XLIII.

Species 269. (Mus. Dennison.)

BULIMUS TRI-BALTEATUS. *Bul. testā fusiformi-ovatā, vix umbilicata, anfractibus sex, convexis, levigatis, vel minutè longitudinaliter striatis; aperturā parviusculā, labro subreflexo; pallide stramineā vel aurantiā, apice roseā, labro vivide croceo.*

THE THREE-BELTED BULIMUS. Shell somewhat fusiformly ovate, slightly umbilicated, whorls six in number, convex, smooth, or minutely longitudinally striated; aperture rather small, lip slightly reflected, pale straw colour, conspicuously belted with three chestnut zones, lip bright pink.

Hab. Santa Fé di Bogota.

A chaste and delicately coloured species, not hitherto described, in the collection of J. Dennison Esq.

Species 270. (Fig. *a* and *b*, Mus. Dennison;
Fig. *c*, Mus. Cuming.)

BULIMUS KNORRI. *Bul. testā fusiformi-ovatā, basi subangulo-producēta, ampliter umbilicata, anfractibus septem, levibus, sub lente transversim minute striatis et indentatis; columellā latè reflexā, aperturā oblongo-ovali, basi attenuatā, quasi sinuatā, labro expanso, vix reflexo; stramineā, aut purpureo-fusca, aut ceraso-fusco obscurè fasciatā, labro intus aut roseo aut vivide croceo.*

KNORR'S BULIMUS. Shell fusiformly ovate, somewhat angularly produced at the base, largely umbilicated, whorls seven in number, smooth, beneath the lens minutely transversely striated and indented; columella broadly reflected; straw colour, or purple-brown, or obscurely banded with bluish-brown, inner edge of the lip pink or deep saffron yellow.

Pfeiffer, Monog. Hel. viv., p. 95.

Hab. La Guayra.

Chiefly distinguished by a peculiar attenuation of the aperture, and by the deep pink or saffron colouring of the lip. The outer colouring of the shell is extremely variable.

Species 271. (Mus. Dennison.)

BULIMUS XANTHOSTOMA. *Bul. testā fusiformi-ovatā, ampliter umbilicata, spirā subacuminatā, anfractibus*

octo, levibus, sub lente transversim minute striatis et indentatis, columellā latè reflexā, aperturā ovatā, labro latè expanso, vix reflexo; pallide stramineā vel aurantiā, apice roseā, labro vivide croceo.

THE YELLOW-MOUTH BULIMUS. Shell fusiformly ovate, largely umbilicated, spire somewhat acuminate, whorls eight in number, smooth, transversely minutely striated and indented under the lens, columella broadly reflected, aperture ovate, lip broadly expanded, scarcely reflected; pale straw colour or orange, pink at the apex and bright saffron yellow at the lip.

D'Orbigny, Voy. dans l'Amér. Mérid. p. 312. pl. 40. f. 1, 2.

Hab. Bolivia and Santa Fé di Bogota.

Mainly distinguished from the preceding species by its more acuminate growth, comprising a whorl more, and less oblong form of the aperture.

Species 272. (Mus. Brit.)

BULIMUS BAIRDII. *Bul. testā cylindraceo-oblongā, crassa, anfractibus septem, piano-convexis, columellā callositatibus grandibus duabus munīta, aperturā parviusculā, labro incrassato, reflexo; albā fusco-linetā, columellā labroque eburneis, apertura fauce crocea.*

BAIRD'S BULIMUS. Shell cylindrically oblong, thick, whorls seven in number, flatly convex, columella furnished with two large callosities, aperture rather small, lip thickened, reflected; white, stained with brown, lip and columella ivory white, interior of the aperture saffron-yellow.

Hab. —?

This very interesting shell partakes of the characters of the *B. auris-bovia* and *auris-Midae*, the latter of which, as may be seen by a comparison of the figure at Pl. 28, with one in Martyn's Universal Conchologist (sub nom. *Helix fibrosa*), assumes very different aspects, but none with which the present agrees.

I have the pleasure to name this species in honour of Dr. Baird of the British Museum, as a trifling acknowledgement of his great diligence in sorting the specimens of the national collection, preparatory to their being critically arranged and named.

BULIMUS.—PLATE XLIII.

Species 273. (Fig. *a* and *b*, Mus. Dennison.)

BULIMUS MURRINUS. *Bul. testa fusiformi-ovata, vix umbilicata, anfractibus sex, convexis, levibus, politis, ultimo basi subproduciā et attenuatā; aperturā suboblongā, infernā leviter sinuatā, labro expanso, vix reflexo; abdita, livido-olivaceo ant. carneo-nigro longitudinaliter strigata et transversim fasciatā, apertura fance viede purpureā, labro alto.*

THE PORCELAIN BULIMUS. Shell fusiformly ovate, scarcely umbilicated, whorls six in number, convex, smooth, polished, last whorl somewhat produced and attenuated at the base; aperture rather oblong, slightly sinuated at the lower part, lip expanded, scarcely reflected; whitish, longitudinally streaked and transversely banded with vivid olive or blue-black, interior of the aperture rich purple, lip white.

Hab. Santa Fé di Bogota.

A beautifully painted species from the collection of

J. Dennison, Esq., nearest allied to the *B. linostoma* of D'Orbigny.

Species 274. (Mus. Cuming.)

BULIMUS CEYLANICUS. *Bul. testa conico-ovata, solidiuscula, spira suturis impressis; anfractibus sex, plano-convexis, obliquè striatis, columellā latè reflexā, aperturā parvā, labro expanso, vix reflexo; intus extusque albā.*

THE CEYLON BULIMUS. Shell conically ovate, rather solid, sutures of the spire impressed; whorls six in number, flatly convex, obliquely striated, columella broadly reflected, aperture small, lip expanded, but slightly reflected; white within and distant.

PFEIFFER, Monog. Hel. viv., vol. ii. p. 59.

Hab. Ceylon.

A rather stout shell of a peculiar conical form.



BULIMUS.

PLATE XLIV.

Species 275. (Mus. Demison.)

BULIMUS CHIMBORASENSIS. *Bul. testā fusiformi, ad basin subattenuatā, vix umbilicatā, aufractibus sex, convexis, longitudinaliter creberrimē elevato-striatis; columellā depresso-reflexā, labro subexpanso, non reflexo; roseofuscescente, albifasciatā, columellā labroque vivida purpureo-roseis.*

THE CHIMBORAZO BULIMUS. Shell fusiform, somewhat attenuated at the base, scarcely umbilicated, whorls six in number, convex, longitudinally very closely elevatedly striated; columella depresso reflected, lip slightly expanded, not reflected; rose-brown, banded with white, lip and columella deep purple-rose.

Bulimus decoratus, Lea (not of Féussac), Obs. vol. ii. pl. 23. f. 108.

Hab. Chimborazo, Columbia, New Granada.

First described by Mr. Lea of Philadelphia from a specimen collected near Carthagena, New Granada, by Dr. Gibbon.

Species 276. (Mus. Cuming.)

BULIMUS BICOLOR. *Bul. testā oblongo-ovatā, umbilicatā, ad basin rotundatā, aufractibus sex, longitudinaliter oblique fortiter rugoso-striatis; carnellescente-albō, fasciis fuscis interruptis cingulatā.*

THE TWO-COLOURED BULIMUS. Shell oblong-ovate, umbilicated, rounded at the base, whorls six in number, longitudinally obliquely strongly roughly striated; bluish-white, encircled by interrupted brown bands.

SOWERBY, Pro. Zool. Soc., 1834, p. 141.

Hab. Xagua, Peru; Miller.

The lower edge of the bands is peculiarly broken or serrated.

Species 277. (Mus. Cuming.)

BULIMUS RUBIFASCIATUS. *Bul. testā cylindraceo-ovatā, vix umbilicatā, aufractibus septem, convexis, levibus, aperturā parvā, labro simplici; albida, rufo-fusco trifasciatā, fasciā medianā angustā, linearī.*

THE RED-BANDED BULIMUS. Shell cylindrically ovate, scarcely umbilicated, whorls seven in number, convex, smooth, aperture small, lip simple; whitish, painted with three red-brown bands, of which the central one is narrow and linear.

Hab. —?

M. Deshayes has, I believe, proposed the name *B. exilis* for this species, already occupied by Gmelin.

Species 278. (Fig. a, Mus. Cuming; Fig. b, Mus. Demison.)

BULIMUS GEOMETRICUS. *Bul. testā subcylindraceo-turritā, umbilicatā, spira suturis impressis; aufractibus septem, subrotundatis, levibus, ultimo peculiariter obliquè producto; columellā latè reflexā, aperturā obliquè ovalatā, labro reflexo; albida, fusco irregulariter tenuiatā et undulatā, vel castanea, albifasciatā, columellā interdum purpuraseante, labro pallide luteo.*

THE GEOMETRIC BULIMUS. Shell somewhat cylindrically turreted, umbilicated, sutures of the spire impressed; whorls seven in number, slightly rounded, smooth, last whorl peculiarly obliquely produced; columella broadly reflected; aperture obliquely ovate, lip reflected; whitish, irregularly filleted, and waved with brown, sometimes chestnut-brown, banded with white, columella sometimes purple, lip pale yellow.

PFEIFFER, Monog. Hel. viv., vol. ii. p. 59.

Hab. New Granada.

An extremely interesting species distinguished in great measure by the obliquely produced growth of the last whorl.

Species 279. (Fig. a and b, Mus. Dennison; Fig. c, Mus. Cuming.)

BULIMUS VITTATUS. *Bul. testā oblongo-turritā, subpellucidā, subaustriter umbilicatā, aufractibus septem ad octo, levibus; columellā reflexā, labro tenui, simplici; flavi-caerulea, purpureo-nigricante sepissimā varie fasciatā, interdum fuscescente longitudinaliter strigata, interdum undique straminea.*

THE BIBAND-BOUND BULIMUS. Shell oblong-turreted, somewhat transparent, rather largely umbilicated, whorls seven to eight in number, smooth; columella reflected, lip thin, simple; yellowish, most frequently variously banded with purple-black, sometimes streaked longitudinally with light brown, sometimes straw-colour throughout.

Spix, Test. Bras., p. 7. pl. 7. f. 4.

Eadem var.

Bulimus Coxeiranus, Moricand.

Hab. Bahia and Pernambuco, Brazil.

The dark purple-black painting of this species is most variable.

Species 280. (Mus. Cuming.)

BULIMUS STRIGATUS. *Bul. testā oblongo-ovatā, subobampliter umbilicatā, anfractibus sex, convexis, longitudinaliter creberrimè elevato-striatis, anfracta ultimo subobliquè producto; columellā latè reflexā, labro expanso; cernu-lescente-albā, livido-nigricante undatā et partim fasciatā, apice nigro.*

THE STREAKED BULIMUS. Shell oblong-ovate, somewhat largely umbilicated, whorls six in number, convex, longitudinally very closely elevately striated, last whorl somewhat obliquely produced; columella broadly reflected, lip expanded; bluish-white, waved and partially banded with livid-black, apex black.

SOWERBY, Conch. Illus., f. 95, 96.

Hab. Huallaga.

The dark painting of this species is partly in bands, and partly in longitudinally waved streaks.

Species 281. (Mus. Dennison.)

BULIMUS BOLIVIANUS. *Bul. testā cylindraceo-oblongā, subampliter umbilicatā, anfractibus septem, planō-convergis, levibus aut obscurè striatis; columellā latè reflexā, aperturā pareiusculā, labro simplici; albido, roseo-fusco crebrifasciatā, fasciis irregularibus, subinterruptis, apice roseo-fusco.*

THE BOLIVIAN BULIMUS. Shell cylindrically-oblong, somewhat largely umbilicated, whorls seven in number, flatly convex, smooth or obscurely striated: columella broadly reflected, aperture rather small, lip simple; whitish, closely banded throughout with rose-brown, bands irregular, somewhat interrupted, apex rose-brown.

PFEIFFER, Pro. Zool. Soc., 1846. p. 34.

Hab. Andes of Bolivia.

The bands are less clearly defined in this species than is usual, and in some specimens run into one another.



BULIMUS.

PLATE XLV.

Species 282. (Fig. *a* and *b*, Mus. Demaison.)

BULIMUS CASTUS. *Bul. testā subfusiformi-oratā, tenui, gracili, anfractibus quinque ad sex, longitudinaliter teniūlē irregulariter striatis, sub lente striis minutis impressis transversim exsculptis; columellā angustā reflexā, labro expanso, vix refexo; albā vel pallidē stramineā, fusco interdum sparsim maculatā, columellā labroque pallidē roseis.*

THE CHASTE BULIMUS. Shell somewhat fusiformly-ovate, thin, slender, whorls five to six in number, longitudinally finely irregularly striated, beneath the lens sculptured transversely with minute impressed striae; columella narrowly reflected; white or pale straw colour, sometimes sparingly spotted with brown, lip and columella pink.

PFEIFFER, Pro. Zool. Soc., 1846, p. 112.

Hab. Central America?

Truly a chaste and delicately coloured species.

Species 283. (Mus. Cuming.)

BULIMUS MORICANDI. *Bul. testā ovalā, tenuiculā, dia-phānā, vix umbilicatā, anfractibus sex, levibus, tumiduscūla, sub lente transversim minitissimè impresso-striatis; columellā reflexā, labro tenui, acuto; indus extrusè crocē.*

MORICAND'S BULIMUS. Shell ovate, rather thin, diaphanous, scarcely umbilicated, whorls six in number, smooth, rather swollen, transversely very minutely impressively striated beneath the lens; columella reflected, lip thin, sharp; saffron yellow within and without.

PFEIFFER, Pro. Zool. Soc., 1846, p. 113.

Hab. Central America.

A bright semitransparent yellowish shell, without any trace of marking.

Species 284. (Mus. Cuming.)

BULIMUS ONAGER. *Bul. testā subpyramidalis-turritā, vix umbilicatā, anfractibus septem ad octo, ad basiu subangula-tis, levibus; columellā subreflexā, labro expanso; albida aut lutescente, cæruleo-viridi varie tinctā et longitudinaliter strigata, aperturā fonce fusco-rufescente.*

THE WILD-ASS BULIMUS. Shell somewhat pyramidaliter turreted, scarcely umbilicated, whorls seven to eight in number, slightly angled at the base, smooth: columella but little reflected, lip expanded; whitish or yellowish, variously mottled and streaked with bluish-green, interior of the aperture brownish red.

Bulinatus onager, Beck, Ind. p. 64.

Bulimus zebra, Spix.

Hab. Bahia, Brazil.

A prettily painted species closely allied to *B. vimineus*. but apparently distinct.

Species 285. (Mus. Cuming.)

BULIMUS VENOSUS. *Bul. testā oblongo-ovalā, vix umbi-licatā, anfractibus sex ad septem, convexis, levibus; columellā reflexā, labro tenui, acuto; albida, lineā castaneā suturali, lineisque castaneis obliquis hic illuc longitudinaliter pictā, apice nigra.*

THE VEINED BULIMUS. Shell oblong-ovate, scarcely umbilicated, whorls six to seven in number, convex, smooth; columella reflected, lip thin, sharp; whitish with a dark chestnut sutural line, and longitudinally painted here and there with chestnut lines, apex black.

Bulinus Menkei, Gruner, Wiegmann's Archives, 1841, (not *B. Menkeana*, Féussac.)

Hab. Angosturi, Banks of the Oronoko.

The painted lines are at very unequal distances throughout.

Species 216. (Mus. Cuming.)

BULIMUS RUDIS. *Bul. testā oratā, ventricosā, vix umbi-licatā anfractibus sex, rugosis, longitudinaliter rudi-plicato-liratis, columellā reflexā, labro simplici; roseo-albida, cinereo-fusco hic illic longitudinaliter strigata, aperturā fonce roseo-purpureo-scente.*

THE RUDE BULIMUS. Shell ovate, ventricose, scarcely umbilicated, whorls six in number, rough, longitudinally rudely plicately ridged, columella reflected, lip simple; pinkish-white, longitudinally streaked here and there with ash-brown, interior of the aperture light rose-purple.

ANTON, MSS.?

Hab. Mexico.

A very characteristic delicately coloured species, of roughly plicated growth.

Species 287. (Mus. Cuming.)

BULIMUS LILIACEUS. *Bul. testā acuminato-oratā, tenui, diaphanā, anfractibus sex, convexis, levibus, vel sub lente minutissimè impresso-striatis; columellā angustè reflexā, labro tenui, acuto; pallidē crocato-aurantia, roseo basin versus tinctā.*

THE LILY-COLOURED BULIMUS. Shell acuminate ovate, thin, diaphanous, whorls six in number, convex, smooth, or very minutely impressly striated beneath the lens; columella narrowly reflected, lip thin, sharp; pale saffron-orange, tinged towards the base with rose.

GULDING, MSS.?

Hab. St. Vincents, West Indies?

Mr. Cuming possesses this shell as an inhabitant of St. Vincents, accompanied with the above name, as having been given to it by the Rev. Lansdowne Gulding. It is but right, however, to add that the authenticity of this name and locality requires confirmation.

Species 288. (Mus. Cuming.)

BULIMUS TENUISSIMUS. *Bul. testā ovatā, tenui, vix umbilicata, anfractibus quinque ad sex, obscurè rugoso-striatis; columellā subdepresso-reflexā, labro tenui, acuto; undique sordide fuscescente.*

THE VERY THIN BULIMUS. Shell ovate, thin, scarcely

umbilicated, whorls five to six in number, obscurely roughly striated; columella somewhat depressedly reflected, lip thin, sharp; pale dirty brown throughout.

Helix tenuissima, De Féussac, MSS.

Hab. Brazil.

One of the most common species, says M. Moricand, on the walls, in Brazil.

Species 289. (Mus. Cuming.)

BULIMUS BENGALENSIS. *Bul. testā conico-ovatā, tenuiculā, umbilicata, anfractibus sex, levibus, basin versus subangulato-productis; columellā latè reflexā, labro tenui, acuto; allidā, lineā castaneā suturali, anfractū ultimo lineis dubiis cingulato, columellā pallidē rosacea, apice nigra.*

THE BENGAL BULIMUS. Shell conically ovate, rather thin, umbilicated, whorls six in number, smooth, slightly angularly produced towards the base; columella broadly reflected, lip thin, sharp; whitish, with a sutural chestnut line, last whorl encircled round the lower part with two chestnut lines, columella tinged with pink, apex black.

LAMARCK, Anim. sans vert. (Deshayes' edit.), vol. viii. p. 233.

Hab. Bengal.

Easily recognised by its black apex and characteristic linear painting, the lower part of the whorls being encircled by two chestnut lines, one of which is concealed in all but the last by the superposition of one whorl upon the other.

Bulimus, Pl. XLVII.



290



291



292



293



294 ♂



294 ♂



295



296



297



298

1000 m - hab

R. E. & R. 1900

BULIMUS.

PLATE XLVI.

Species 290. (Mus. Cuming.)

BULIMUS VITRINOIDES. *Bul. testā subglobosa-ovalā, tenuissimā, hyalinā, apice brevi, acuto; anfractibus quinque, subrude striatis, ultima ventricoso; columellā labroque simplicibus; pellucido-corneā.*

THE VITRINA-LIKE BULIMUS. Shell somewhat globosely ovate, rather thin, hyaline, apex short, sharp; whorls five in number, somewhat rudely striated, last whorl ventricose; lip and columella simple; transparent horny.

Hab. ——?

This shell looks as if it were composed of fine glass, and is so delicately transparent, that the internal winding of the columella is discernible throughout.

Species 291. (Mus. Benson.)

BULIMUS NILAGARICUS. *Bul. testā cylindraceo-turrītā, unibilicatā, anfractibus octo, convexis, concavisè substriatis, et sub lente minutissimā transversa sulcatis, suturis subtilliter crenulatis, columellā labroque expansis, vir reflexis; castaneo-fusca, nitente, suturis albida, labro albido.*

THE NEILGHERRY BULIMUS. Shell cylindrically turrited, umbilicated, whorls eight in number, convex, and, under the lens, very minutely grooved transversely, sutures finely crenulated, lip and columella expanded, scarcely reflected; shining chesnut-brown, whitish at the sutures, lip whitish.

PFEIFFER, Pro. Zool. Soc., 1846, p. 41.

Hab. Neilgherry Hills, India; Jerdon.

Of a silken chesnut hue, whitish at the sutures.

Species 292. (Mus. Cuming.)

BULIMUS EXILIS. *Bul. testā subcylindraceo-oblongā, anfractibus sex ad septem, longitudinaliter rudo striatis, columellā labroque simplicibus, aperturā parenscaulis; flacio-fuscescente, subhyalinā, aperturā fucee violaceo-albicans.*

THE DELICATE BULIMUS. Shell somewhat cylindrically oblong, whorls six to seven in number, longitudinally rudely striated, lip and columella simple, aperture

rather small; yellowish-brown, somewhat hyaline, interior of the aperture violet-tinged.

GMELIN, Syst. Nat. p. 3668.

Hab. ——?

The sutures of this species are inclined to be white as in the preceding. It is not unlike the *B. Guadaloupensis*.

Species 293. (Mus. Cuming.)

BULIMUS CINNAMOMEO-LINEATUS. *Bul. testā subpyramidalis-ovalā, umbilicatā, umbilico parvo; anfractibus septem, convexis, levibus; columellā plano-reflexā, labro paululum expanso; albida, lineis cinnamonaeis obliquè striatā, anfractus ultimi basi innasulatā.*

THE CINNAMON-LINED BULIMUS. Shell somewhat pyramidalis-ovalis, umbilicated, umbilicus small; whorls seven in number, convex, smooth; columella flatly reflected, lip a little expanded; whitish, obliquely streaked with cinnamon lines, last whorl unspotted at the base.

Helix cinnamonomeo-lineata, Moricand, Mém. Genève, vol. ix. p. 46, pl. 4. f. 6, 7.

Hab. Bahia, Brazil.

The cinnamon lines are suddenly arrested at the base of the last whorl.

Species 294. (Fig. a, Mus. Dennison;

Fig. b, Mus. Cuming.)

BULIMUS GUADALOUPENSIS. *Bul. testā cylindraceo-oblongā, solidusnā, anfractibus sex, convexis, longitudinaliter rudo et tenuiter striatis, apertura peculiariter quadrato-ovalā, columellā labroque subexpansis, non reflexis; flacio-fuscescente, nigricante-castaneo interdūs conspicue fasciata.*

THE GUADALOPE BULIMUS. Shell cylindrically oblong, rather solid, whorls six in number, convex, longitudinally rudely and finely striated, aperture peculiarly squarely oval, lip and columella somewhat expanded, not reflected; light yellowish-brown, sometimes conspicuously banded with very dark chesnut.

BRUGUIERE, Ene. Méth. Dict. no. 26.

Hab. West Indies.

Distinguished by the peculiar quadrate form of the aperture.

BULIMUS.—PLATE XLVI.

Species 295. (Mus. Cuming.)

BULIMUS MULTIFASCIATUS. *Bul. testā oblongo-ovatā, vix umbilicatā, anfractibus sex, ləvibus, nitidis, columellā reflexā, labro simplici, acuto; lutescente-albā, zonis roseis, violaceo-nigrante marginatis, cingulatā, apice nigrante.*

THE MANY-BANDED BULIMUS. Shell oblong-ovate, but slightly umbilicated, whorls six in number, smooth, shining, columella reflected, lip simple, sharp; yellowish-white, encircled with pink zones, edged with violet black, apex blackish.

LAMARCK, Ann. sans. vert. (Deshayes' edit.) vol. viii. p. 233.

Hélis picturata, Fénnssae.

Hab. West Indies.

Easily recognised by its pretty pink zones, which are brighter as seen in the interior.

Species 296. (Mus. Benson.)

BULIMUS ABYSSINICUS. *Bul. testā conico-ovatā, subampliter umbilicatā, anfractibus sex, ventricosis, subrotundatis, longitudināller creberrimè tenuicostatis; columellā latē expansā, aperturā suborbiculari, labro expanso, non reflexo; sordidè albā.*

THE ABYSSINIAN BULIMUS. Shell conically ovate, rather largely umbilicated, whorls six in number, ventricose, somewhat rounded, longitudinally very closely finely ribbed; columella broadly expanded, aperture nearly orbicular, lip expanded, not reflected; dull white.

RUPPELL, MS.

Hab. Abyssinia; Malwah, Central India; Benson.

Finely ribbed somewhat after the manner of the *B. Scalariformis*, but of a more *Pupa*-like character. Published

with the above name on the authority of Mr. Benson, to whose kindness I am indebted for the species.

Species 297. (Mus. Benson.)

BULIMUS JERDONI. *Bul. testā subabbreviato-ovatā, ampliter umbilicatā, spirā breviusculā, acuminatā, anfractibus sex, tumidis, rotundatis, columellā latē reflexā, aperturā suborbiculari, labro expanso, vix reflexo; sordidè olivaceā.*

JERDON'S BULIMUS. Shell somewhat shortly oval, largely umbilicated, spire rather short, acuminate, whorls six in number, swollen, rounded, columella broadly reflected, aperture nearly orbicular, lip expanded, scarcely reflected; dirty olive-brown.

BENSON, MS.

Hab. Deccan, Hindostan.

A short swollen species with somewhat of the *Pupa* character.

Species 298. (Mus. Cuming.)

BULIMUS BENGUELENSIS. *Bul. testā oblongā, solidiusculā, vix umbilicatā, anfractibus sex, convexis, subtilissimè rvdè striatis, columellā reflexā, labro ex expanso, non reflexo; albā, ad umbilicum et suturas pallidè aurantio-fascescente.*

THE BENGUELA BULIMUS. Shell oblong, rather solid, scarcely umbilicated, whorls six in number, convex, very finely rudely striated; columella reflected, lip but slightly expanded, not reflected; white, light orange-brown at the sutures and about the umbilicus.

Hab. Benguela.

Allied to the *B. Guadaloupensis*, though from a very different locality.



309



300



301



302



303



304



305



306



307.



308

10 mic.

10 mic.

BULIMUS.

PLATE XLVII.

Species 299. (Mus. Benson.)

BULIMUS VIREX. *Bul. testá cylindraceo-turrítá, sinistrál,*
smb-Pupíformi, diapháná, umbilicatá, anfractibus no-
vem, convexis, obliquí striatis, aperturá parvá, rotundatá,
labro plano-reflexo; pellucido-fuscescente, strigis
opaco-albis obliqué notatá.

THE STRIPED BULIMUS. Shell cylindrically turreted, sinistral, somewhat Pupa-shaped, diaphanous, umbilicated, whorls nine in number, convex, obliquely striated, aperture small, rounded, lip flatly reflected; light transparent brown, obliquely marked with opake white stripes.

HUTTON, Ms., Pfeiffer, Monog. *Bul.* p. 118.

Hab. Simla and Landour, Western Himalaya, India (among moss); Benson.

A reversed cylindrical shell partaking very much of the character of *Pupa*.

Species 300. (Mus. Cuming.)

BULIMUS CREPUNDIA. *Bul. testá subpyramidalis-ovalá,*
tenui, diapháná, umbilicatá, anfractibus septem, sub-
tilissimè rugoso-striatis, columellá reflexá, labro sin-
plici; pellucido-corneá, opaco-albo hic illuc obliqué
notatá.

THE PLAYTHING BULIMUS. Shell somewhat pyramidaly ovate, thin, diaphanous, umbilicated, whorls seven in number, very finely roughly striated, columella reflected, lip simple; transparent horny, obliquely marked here and there with opake white.

D'ORBIGNY, Voy. dans l'Amér. Mérid. p. 275. pl. 33. f. 18,
19.

Hab. Chiquitos, Bolivia.

It is probably not without some doubt that Dr. Pfeiffer assigns this shell to the *B. crepundia*.

Species 301. (Mus. Benson.)

BULIMUS COLEBS. *Bul. testá cylindracea, Pupíformi,*
tenui, diapháná, latè umbilicatá, anfractibus octo, obliquí
striatis, columellá latè reflexá, labro expanso non re-
flecto; pellucido-corneá, strigis opaco-albis obliqué
notatá.

THE BACHELOR BULIMUS. Shell cylindrical, Pupa shaped, thin, diaphanous, broadly umbilicated, whorls

eight in number, obliquely striated, columella broadly reflected, lip expanded, not reflected, transparent horny, obliquely marked with opake-white streaks.

BENSON, Ms., Pfeiffer, Monog. *Bul.* p. 119.

Hab. Landour and Kemaon, Western Himalaya, India, (on dead stems of the wild Indigo); Benson.

This and the two preceding species are of the same texture, a transparent horny substance, obliquely marked with opake white streaks.

Species 302. (Mus. Benson.)

BULIMUS GRIFFITHSII. *Bul. testá elongatá, subcylindrá-*
ced, umbilicatá, anfractibus octo, subplanulatis, levibus,
columella latè reflexá, aperturá suboblique productá,
labro expanso, non reflexo; peculiaritatem calcareo-albá,
apertura fauce pallidè aurantio-fuscescente.

GRIFFITH'S BULIMUS. Shell elongated, somewhat cylindrical, umbilicated, whorls eight in number, rather flattened, smooth, columella reflected, aperture somewhat obliquely produced, lip expanded, not reflected; of a peculiar chalk-white, tinged in the interior with pale orange-brown.

BENSON, Ms.

Hab. Afghanistan, India; Griffith.

Named by Mr. Benson after the lamented botanist Dr. Griffith, by whom the specimen here figured was collected.

Species 303. (Mus. Benson.)

BULIMUS SINDICUS. *Bul. testá cylindraceo-elongatá, latè*
umbilicatá, anfractibus novem, plano-convexis, levibus
aut concentrico striatis, aperturá pareá, columellá la-
brouque expansa, vis reflexis; griseo-albá, strigis pel-
lucidis hic illuc obliqué notatá.

THE SINDE BULIMUS. Shell cylindrically elongated, broadly umbilicated, whorls nine in number, flatly convex, smooth or concentrically striated, aperture small, lip and columella expanded, but slightly reflected; greyish white, here and there obliquely marked with pellucid streaks.

BENSON, Ms.

Hab. Near the head waters of the river Sind, India; Benson.

Mr. Bensou informs me that this species was first collected by Capt. Hutton in the warmer vallies of the sub-Himalayan range, on the left bank of the Sutlej; and that specimens were transmitted to him from Khyber Pass by Capt. Barr.

Species 304. (Mus. Cumug.)

BULIMUS LIVESCENS. *Bal. testā acuminato-ovatā, sub-turrīdā, tenui, anfractibus septem, subtrotundatis, levibus, columellā simplici, labro tenui, acuto; livescente-albā; interdum zonulis fascia angustis tribus cingulatā.*

THE BLUISH BULIMUS. Shell acuminate ovate, rather turreted, thin, whorls seven in number, somewhat rounded, smooth, columnella simple, lip thin, sharp; bluish-white, encircled with three narrow light-brown zones.

PFEIFFER, Monog. *Bul.* p. 175.

Hab. Tehuacan, Mexico.

It should be observed that in the banded variety the two lower bands, which are frequently broader than the uppermost, are only seen in the last whorl.

Species 305. (Mus. Cuming.)

BULIMUS FUSCAGULA. *Bul. testā cylindraceo-elongatā, basi compressā, subumbilicatā, pupiformi, anfractibus novem, obtuse striatis et peculiariter indentatis, columellā bilamellatis, aperturā parvā, contractā, expansā; subcoriē, opaco-intensente-albo strigatā et subtiliter reticulatā, labro roseo.*

THE FUSCOUS-THROAT BULIMUS. Shell cylindrically elongated, compressed at the base, slightly umbilicated, pupa shaped; whorls nine in number, obtusely striated and peculiarly indented, columnella with two tooth-like laminae, aperture small, contracted, expanded; somewhat horny, streaked and finely reticulated with opaque yellowish white, lip rose.

D'ORIGNY, Voy. dans l'Amér. Mérid. p. 310. pl. 39. f. 1, 2.

Hab. Rio Janeiro.

Very closely allied to *B. Janeiroensis*, and from the same locality; it seems, however, to differ in the columnar laminae and reticulated arrangement of the opaque pattern.

Species 306. (Mus. Cuming.)

BULIMUS HELENA. *Bul. testā subobso-cylindraceā, tenui,*

umbilicatā, anfractibus sex, rotundatis, striis minutis crebrerrimè decussatis, suturis impressis, columellā labrovoe ex reflexis; squamidē olivaceā, semipellucida.

THE HELENA BULIMUS. Shell somewhat stoutly cylindric, thin, umbilicated, whorls six in number, rounded, very closely decussated with minute striae, sutures impressed, lip and columnella scarcely reflected; dirty olive, semitransparent.

QUOY and GAIMARD, Voy. dans l'Astrol. vol. 2. p. iii. pl. 9. f. 82.

Hab. St. Helena.

The minute decussated striae of this species are scarcely discernible without the aid of a lens.

Species 307. (Mus. Cumug.)

BULIMUS CONSTRICTUS. *Bul. testā pyramidalis-conicā, umbilicatā, anfractibus septem, rotundatis, striis transversi et longitudinalibus minutissimis decussatis, suturis peculiariter impressis; albida, semipellucida, epidermide tenui olivaceā indutā.*

THE TIED BULIMUS. Shell pyramidal conical, umbilicated, whorls seven in number, rounded, very minutely decussated with longitudinal and transverse striae, sutures peculiarly impressed; whitish, semi-transparent, covered with a thin olive epidermis.

PFEIFFER, Symbolae, vol. 1. p. 43.

Bulimus Angosturensis, Gruner.
Hab. Angostura; Gruner.

Mainly characterized by the impressed growth of the sutures.

Species 308. (Mus. Cuming.)

BULIMUS DIGITALE. *Bul. testā obeso-ovatā, umbilicata, spiræ suturis peculiariter profundè impressis, anfractibus quinque ad sex, subtiliter granuloso-reticulatis, columellā reflexō, labro simplici; olivaceo-fuscā.*

THE THIMBLE BULIMUS. Shell stoutly ovate, umbilicated, sutures of the spire peculiarly deeply impressed, whorls five to six in number, finely granularly reticulated, columnella reflected, lip simple; olive-brown.

Hab. ——?

The surface of this shell is reticulated in a delicate wrinkled manner, covered by a silken olive epidermis, and the sutures are universally deeply impressed.



BULIMUS.

PLATE XLVIII.

Species 309. (Mus. Cuming.)

BULIMUS MACULATUS. *Bul. testā sub-pyramidalē-ovatā, vix umbilicatā, anfractibus convexis, levibus, transversim sub lete minutissimē impresso-striatis, columellā reflexā, labro simplici, acuto; viridescente-albidā, maculis fuscō-olivaceis, quadratis hic illicē notatis.*

THE SPOTTED BULIMUS. Shell somewhat pyramidalily ovate, scarcely umbilicated, whorls convex, smooth, transversely very minutely impressively striated, columella reflected, lip simple, acute; greenish-white, marked here and there with square brownish olive spots.

LEA, Obs. vol. ii. p. 84. p. 23. f. 97.

Hab. Carthagena, New Granada.

The square spots though irregular and scattered are mostly situated in the same line longitudinally.

Species 310. (Mus. Cuming.)

BULIMUS TRILINEATUS. *Bul. testā oblongo-ovatā, tenuiculā, anfractibus sex, subrugoso-striatis, ad suturas crenulatis, labro tenui, acuto; albida, strigis rufo-fuscis longitudinalibus plus minusve densè notata.*

THE TRILINEATED BULIMUS. Shell oblong ovate, rather thin, whorls six in number, somewhat roughly striated, crenulated at the sutures, lip thin, sharp; whitish, more or less thickly marked with red-brown longitudinal stripes.

QUOY and GAIMARD, Voy. dans l'Astrol. vol. ii. p. 107, pl. 9. f. 1 to 3.

Hab. New Holland.

It is necessary to observe that this species is named after a character in the animal, whose neck and back are marked with three lines.

Species 311. (Mus. Cuming.)

BULIMUS MANOELII. *Bul. testā acuminato-oblongā, subumbilicatā, anfractibus septem, levibus, ant obliquē indistincte striatis, columellā latē reflexā, labro tenuiculā, vix reflexo; albida, anfractu ultimo infernē pureo-olivaceo bifasciato.*

MANOEL'S BULIMUS. Shell acuminate ovate, slightly umbilicated, whorls seven in number, smooth, or obliquely indistinctly striated, columella broadly re-

flected, lip rather thin, scarcely reflected; whitish, last whorl encircled round the lower part with two purple olive bands.

Helix Manoelli, Moricand, Mém. de Genève, vol. ix. p. 45.

pl. 4. f. 4. 5.

Hab. Bahia, Brazil.

The upper band, being exactly on the sutural line, is nearly concealed in all but the last whorl.

Species 312. (Mus. Cuming.)

BULIMUS ANGIOSTOMA. *Bul. testā sub-cylindraceo-ovatā, vix umbilicatā, anfractibus octo, tumidiusculis, ultimo ad aperturam peculiariter compresso; aperturā oblongā, angustā, labro exiūne plano-reflexo; cinereo-albida, lineis nigricantibus et fuscis longitudinaliter pictā, apertura fance fuscā.*

THE NARROW-MOUTH BULIMUS. Shell somewhat cylindrically ovate, scarcely umbilicated, whorls eight in number, rather swollen, last whorl peculiarly compressed at the aperture; aperture oblong, narrow, lip delicately flatly reflected; ashy white, longitudinally painted with black and brown lines, interior of the aperture brown.

WAGNER, Spix. Test. Bras. p. 14. pl. 13. f. 4.

Bulimus Capueira, Spix.

Testa junior. *Bulimus virgatus*, Spix.

Hab. Brazil.

The singular compression of the aperture forms a very characteristic feature in this species.

Species 313. (Mus. Cuming.)

BULIMUS OREADES. *Bul. testā acuminato-oblongā, umbilicatā, anfractibus sex ad septem, levibus, vel minutissimē impresso-striatis, columellā reflexā, labro tenui, acuto; flaccidente-albida, strigis olivaceo-fuscis, anfractu medio desincentibus, peculiariter notatis.*

THE OREADES BULIMUS. Shell acuminate oblong, umbilicated, whorls six to seven in number, smooth, or very minutely impressively striated, columella reflected, lip thin, sharp; yellowish-white, peculiarly marked with olive-brown streaks, ending in the middle of the whorls.

D'ORBIGNY, Voy. dans l'Amér. Mérid. p. 270. pl. 31. f. 11.

Hab. Province of Corrientes, Argentine Republic, Bolivia
(under dead leaves); D'Orbigny.

The chief specific peculiarity of this species resides in the painting, which does not descend below the sutural line of the whorls.

Species 314. (Mus. Cuming.)

BULIMUS FERIATUS. *Bul. testā subfusiformi, eix umbilicatā, aufractibus sex, subtiliter rugoso-striatis, saturis impressis, columellā reflexā, aperturā parvissimā, labro expanso; fuscescente, fusco striatā et variegatā, punctis allis aspera.*

THE UNEMPLOYED BULIMUS. Shell somewhat fusiform, scarcely umbilicated, whorls six in number, finely roughly striated, sutures impressed, columella reflected, aperture rather small, lip expanded, light brown, streaked and variegated with bright brown, sprinkled with white dots.

Hab. —?

Dr. Pfeiffer, when in London, attributed this shell to the *B. lophicus* of D'Orbigny, but it fails to agree with the description and figure of that species, and appears to be equally distinct from all others.

Species 315. (Mus. Benson.)

BULIMUS BONTLE. *Bul. testā conico-ovatā, tenui, ventricosā, subtiliter umbilicatā, aufractiō quinque, minutiō obliquē striatis, medio tumidis, inflatis, columellā latē reflexā, labro tenui, expanso, non reflexo.*

THE BONTIA BULIMUS. Shell conically ovate, thin, ventricose, finely umbilicated, whorls five in number, minutely obliquely striated, swollen and inflated round the middle, columella broadly reflected, lip thin, expanded, not reflected.

Helix Bontia, Chemnitz, Conch. Cab. vol. ix. p. 156.
pl. 134. f. 1216, 1217.

Hab. Southern India; Jerdon.

An extremely rare species, though described upwards of sixty years since by Chemnitz in his Conchylion Cabinet. The specimen here rerepresented, for which I am indebted to the kindness of Mr. Benson, was collected in Southern India by Mr. Jerdon of Madras.

Species 316. (Mus. Cuming.)

BULIMUS CONFUSUS. *Bul. testā oblongo-ovalatā, tenuicollatā,*

vix umbilicatā, spire suturis impressis, aufractibus sex, convexis, longitudinaliter crenatim plicato-striatis, et hic illis indentatis; fuscescente-albida.

THE PUZZLING BULIMUS. Shell oblong-ovate, rather thin, scarcely umbilicated, sutures of the spire impressed, whorls six in number, convex, longitudinally very closely plicately striated, and here and there indented; light brownish white.

Hab. —?

This shell resembles M. D'Orbigny's *B. Torallyi* in form, but does not agree with it in the detail of character.

Species 317. (Mus. Cuming.)

BULIMUS QUITENSIS. *Bul. testā conico-ovatā, subangustata, umbilicatā, aufractibus septem, convexis, concentricē subrugoso-striatis; columellā labroque eix reflexis; fuscescente castaneo-fusco hic illis striatā, umbilico nigricante, columellā labroque purpureo-castaneis, nitentibus.*

THE QUITO BULIMUS. Shell conically ovate, rather largely umbilicated, whorls seven in number, convex, concentrically rather roughly striated, lip and columella scarcely reflected; light-brown, streaked here and there with chesnut-brown, umbilicus blackish, lip and columella shining purple chesnut.

PFEIFFER. Pro. Zool. Soc., 1847.

Hab. Quito, Republic of Ecuador.

The general aspect of this shell is not much unlike that of an immature *Partula faba*.

Species 318. (Mus. Cuming.)

BULIMUS MONILIFER. *Bul. testā subpyramidalis-ovatā, umbilicatā, aufractibus convexis, levibus, columellā reflexā, aperturā subquadratā, labro simplici; opacobalbā, fasciis duabus interruptis aurantio-fuscis, valde indistinctis, cingulatā, aperturā aurantio-fuscā.*

THE NECKLACE-BOUND BULIMUS. Shell somewhat pyramidal, umbilicated, whorls convex, smooth, columella reflected, aperture somewhat square, lip simple; opake-white, encircled with two very indistinct interrupted orange-brown bands, aperture tinged with orange.

Hab. —?

Indistinctly encircled by two characteristic interrupted bands. The aperture is remarkable for the same lateral compression as in the *B. Caudatolamprinus*.



320a



320



321c



321



322



323



324



324



325

C. 100 m. S.

C. 100 m. S.

BULIMUS.

PLATE XLIX.

Species 319. (Mus. Cuming.)

BULIMUS FABREFACTUS. *Bul. testā pyramidalis, ad basin angulato-compressā et umbilicatā, anfractibus septem, longitudinaliter rugoso-striatis et inequaliter indentatis, columellā reflexā, aperturā ad basim canaliculatā, labro simplici; albida, fuso subtiliter et crebrissimè striatā, strigis ad angulum basalem tantum descendentibus, columellā pallidè rosaceā, apertura fauce cernuolente.*

THE CURIOUSLY-FORMED BULIMUS. Shell pyramidal, angularly compressed and umbilicated at the base, whorls seven in number, longitudinally roughly striated and irregularly indented, columella reflected, aperture channelled at the base, lip simple; whitish, finely and closely streaked with brown, the streaks descending only to the basal angle, columella tinged with pink, interior of the aperture bluish.

Hab. Province of Merida, New Grauada; Linden.

Of the same peculiarly angled structure at the base as the *B. caudiculatus*, from which it is amply distinguished by its more elongated form, solid growth, and different style of colouring.

Species 320. (Fig. a and b, Mus. Cuming.)

BULIMUS VIRGULATUS. *Bul. testā acuminato-oblongā, vix umbilicatā, anfractibus septem, levibus, aperturā ovatā, labro simplici; albā, fasciis castaneis interruptis variè dispositis, interdum rufescēte-crocā, innuculatā.*

THE STRIPED BULIMUS. Shell acuminately oblong, scarcely umbilicated, whorls seven in number, smooth, aperture ovate, lip simple: white, with interrupted chestnut bands, variously disposed, sometimes of a rich reddish saffron, unspotted.

Cochlogena virgulata, Féussac, Hist. Moll. pl. 142. B. f. 1-7.

Bulimus Caribbeorum, Lamarck.

Hab. West Indies.

Extremely variable both in colour and pattern.

Species 321. (Mus. Cuming.)

BULIMUS LITUS. *Bul. testā oblongo-ovatā, tenui, ventricosā, vix umbilicatā, anfractibus sex, obliquè crebrissimè striatis, labro tenui simplici; albida, nigrante-carulō, longitudinaliter maculatā et undulatā.*

THE MOTTLED BULIMUS. Shell ovate, thin, ventricose, scarcely umbilicated, whorls six in number, obliquely very closely striated, lip thin, simple; whitish, longitudinally blotched and waved with blackish-blue.

Helix lila, Féussac, Prodrome, p. 54.

Bulimus littoralis, Spix.

Bulimus magus, Wagner.

Hab. Brazil.

Very closely allied to the *B. papryraceus*, which it is not improbable may be another state of the species.

Species 322. (Mus. Cuming.)

BULIMUS MILTOCHEILUS. *Bul. testā acuminato-oblongā, subcylindraccā, umbilicatā, anfractibus sex, lăevigatis, semipellucidis, nitentibus, longitudinaliter peculiariter plicatis, columellā latē reflexā, aperturā oblongā, labro reflexo; cereo-albā, labro intenſe coccineo-annulato.*

THE VERMILION-LIP BULIMUS. Shell acuminately oblong, somewhat cylindrical, umbilicated, whorls six in number, smooth, semi-transparent, shining, longitudinally peculiarly plaited, columella broadly reflected, aperture oblong, lip reflected; waxy-white, lip deep scarlet-vermilion.

Hab. Sau Christoval, south-eastern island of Solomon's Group, north-east coast of New Holland.

A shell of curiously plaited growth, remarkable for its semipellucid wax-like appearance, with which its rich vermilion lip presents a notable contrast. Mr. Cuming possesses several specimens from the above named locality, all in the same condition, which seems perfectly natural and not the result of any artificial clearing.

Species 323. (Mus. Cuming.)

BULIMUS RHODOSTOMA. *Bul. testā ovatā, subventricosā, umbilicatā, anfractibus sex, longitudinaliter subrude striatis, ad suturas crenulatis, columellā reflexā, labro simplici; roseo-brunneoscente.*

THE ROSE-MOUTH BULIMUS. Shell ovate, rather ventricose, umbilicated, whorls six in number, longitudinally somewhat rudely striated, crenulated at the sutures: columella reflected, lip simple; rose-brown.

Gray, Pro. Zool. Soc., 1834, p. 65.

Hab. New Holland?

This shell has been ascribed to the above-recorded species in Mr. Cuming's collection, and is so published here, but, it should be added, with considerable doubt.

Species 324. (Mus. Cuming.)

BULIMUS HYEMATUS. *Bul. testā acuminato-ovatā, anfractibus septem, peculiariter plicato-corrugatis, apertura parviusculā, labro simplici; cinerascente-albā, apertura fonce roseo-purpureā.*

THE FROZEN BULIMUS. Shell acuminate ovate, whorls seven in number, peculiarly plicately wrinkled, aperture rather small, lip simple; pale ash-white, rose-purple in the interior.

Hab. ——?

The surface of this shell is plicately wrinkled in a manner different from any other species. Externally it is of French-white hue, purplish towards the apex, and the mouth is tinged with deep purple-rose throughout.

Species 325. (Mus. Cuming.)

BULIMUS SPORADICUS. *Bul. testā elongato-ovatā, pyramidalī, viz umbilicatā, anfractibus octo, sublevibus, aperturā parvā, labro tenui, acuto; albida, nigricante-fusco varie lineatā et undatā.*

THE SCATTERED BULIMUS. Shell elongately ovate, pyramidal, scarcely umbilicated, whorls eight in number, more or less smooth, aperture small, lip thin, sharp: whitish, variously lined and waved with bluish-brown.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 271.
pl. 32, f. 12 to 15.

Hab. Patagonia, Bolivia; D'Orbigny.

M. D'Orbigny describes this species to be widely distributed over South America, and very abundant.

Species 326. (Mus. Cuming.)

BULIMUS TENIOLUS. *Bul. testā cylindraceo-ovatā, ad basin rotundatā, anfractibus septem, minutè granulatis, ad suturas crenulatis, columellā reflexā, labro vix reflexo; fulvo-castaneā, ad suturas albizonatā, aperturā fonce lividā, peritreme albido.*

THE FILLETED BULIMUS. Shell cylindrically ovate, rounded at the base, whorls seven in number, minutely granulated, crenulated at the sutures, columella reflected, lip scarcely reflected, fulvous chestnut, zoned with white at the sutures, interior of the aperture livid, peritreme white.

NYST, Ext. Acad. Royale de Bruxelles, p. 9. pl. 3. f. 4. *a, b.*

Hab. South America.

Of a peculiar obtusely rounded growth at the base.



327



328



329



330

327-330

Fig. 327-330

BULIMUS.

PLATE L.

Species 327. (Fig. 327, 328. Mus. Cuming.)

BULIMUS ADANSONI. *Bul. testā oblongo-ovatā, tenuicellā, vix umbilicatā, anfractibus novem, subtilissimè granulato-striatis, ad suturas subtiliter crenulatis, columellā rectā, revolutā, labro simplici, acuto; albida, castaneo-fusco conspicuè longitudinaliter flammatā, epidermide tenui stramineā induitā, columellā et aperture fauce purascente-ceruleis.*

ADANSON'S BULIMUS. Shell oblong-ovate, rather thin, scarcely umbilicated, whorls nine in number, very finely granulately striated, finely crenulated at the sutures, columella straight, rolled back, lip simple, sharp; whitish, conspicuously longitudinally flamed with chesnut-brown, covered with a thin straw-coloured epidermis, columella and interior of the aperture purplish-blue.

PFEIFFER, Monog. *Bul.* p. 179.

Le Kambeul, Adanson.

Bulimus Kambeul, Bruguïère.

Cochlogena adilis, Féruſſac.

Fide Pfeiffer.

Bulimus speciosus, Parreys.

— *Kordofanus*, Parreys.

— *xantholinus*, Ziegler.

— *Achatinoides*, Ziegler.

Hab. West coast of Africa.

Dr. Pfeiffer's name for this species is to be preferred to that adopted by Bruguïère and Lamarek, which does not enter strictly in the Latin form of nomenclature.

For Species 328, see Pl. LVIII.

Species 329. (Mus. Cuming.)

BULIMUS RUPPELLIANUS. *Bul. testā subabbreviato-ovatā,*

ventricosā, vix umbilicatā, anfractibus septem, creberrimè plano-granulatis, columellā rectā, revolutā, labro simplici, acuto; albida, flammis subangustis castaneo-fusco longitudinaliter pictā, epidermide tenui stramineā induitā.

RUPPELL'S BULIMUS. Shell somewhat shortly ovate, ventricose, scarcely umbilicated, whorls seven in number, very closely flatly granulated, columella straight, rolled back, lip simple, sharp; whitish, longitudinally painted with rather narrow chesnut-brown flames, covered with a thin straw-coloured epidermis.

PFEIFFER, Monog. *Bul.* p. 190.

Hab. Abyssinia; Ruppell.

Constantly distinguished from the preceding species by its shortened ventricose growth.


Species 330. (Mus. Cuming.)

BULIMUS AFRICANUS. *Bul. testā acuminato-ovatā, profunde umbilicatā, anfractibus novem, subtundatis, reticulato-striatis et subtilissimè concentricè corrugatis, ad suturas crenatis, columellā rectā, revolutā; livido-albida, castaneo-fusco pallidè flammatā, flammis subdistansibus.*

THE AFRICAN BULIMUS. Shell acuminately ovate, deeply umbilicated, whorls nine in number, somewhat rounded, reticulately striated and very finely concentrically wrinkled, crenated at the sutures, columella straight, rolled back; livid white, faintly marked with rather distant longitudinal chesnut-brown flames.

Hab. West coast of Africa.

This seems constantly distinguished from the *B. Adansonii* in size, in form, and in pattern of colouring.



B U L I M U S .

PLATE LI.

Species 331. (Mus. Cuming.)

BULIMUS MUTILATUS. *Bul. testā attenuato-pyramidalis, crassiusculā, umbilicatā, anfractibus quatuor ad quinque, deinde latè decollatis, obliquè ruditè striatis, anfractu ultimo basi subangulato-tumido, aperturā parvā, rotundā; squamidē albā.*

THE MUTILATED BULIMUS. Shell attenuately pyramidal, rather thin, umbilicated, whorls four to five in number, then broadly decollated, obliquely rudely striated, last whorl somewhat angularly swollen at the base, aperture small, rounded, dirty-white.

Hab. —?

Distinguished from the *B. decollatus* by its more attenuately pyramidal form and contracted aperture.

Species 332. (Mus. Cuming.)

BULIMUS GRUNERI. *Bul. testā cylindraceo-turrītā, subumbilicatā, anfractibus septem, plani-convexis, lœvibus, nitidis, columellā depresso-reflexā, aperturā parvissimā, labro tenui, simplici; cærudescente-albā, maculis fuscocentribus subquadratis hūi illic irreguliter fusciatim notatā, apice purpureo-nigricante.*

GRUNER'S BULIMUS. Shell cylindrically turreted, slightly umbilicated, whorls seven in number, flatly convex, smooth, shining, columella depressly reflected, aperture rather small, lip thin, simple; bluish-white, marked here and there with irregular bands of light brown square-like spots. apex purple-black.

PFEIFFER, Pro. Zool. Soc. 1846. p. 30.

Hab. Mexico.

Named in honour of Mr. Gruner, an eminent collector at Bremen.

Species 333. (Mus. Cuming.)

BULIMUS ALBATUS. *Bul. testā acuminato-oblongā, subumbilicatā, anfractibus octo, convexis, lœvibus, columellā reflexā, labro simplici; albā, interdum fasciis interrupitis brunnēis duabus vel tribus cingulatā, apertura fance subaurantia.*

THE WHITE BULIMUS. Shell acuminate oblong, slightly umbilicated, whorls eight in number, convex, smooth, columella reflected, lip simple; white, sometimes encircled with two or three interrupted brown bands, interior of the aperture tinged with orange.

PFEIFFER, Symbole, vol. ii. p. 42.

Helix abalta, De Féussac.

Bulinus bicinctus, Recluz.

Hab. Yemen, Arabia.

Rarely more than a single indistinct band is visible on the upper whorls.

Species 334. (Mus. Cuming.)

BULIMUS DECOLLATUS. *Bul. testā cylindraceo-elongatā, anfractibus ad septem, deinde decollatis, plani-convexis, striis spiraliis et oblique subtiliter decussatis, marginē crenulatis, columellā reflexā, aperturā parvissimā, labro simplici; albā, aut pallide spadiceā,*

THE DECOLLATED BULIMUS. Shell cylindrically elongated, whorls about seven in number, then decollated, flatly convex, finely decussated with spiral and oblique striae, crenulated at the margin, columella reflected, aperture rather small, lip simple; whitish, or pale fawn-colour.

BRUGUÈRE, Enc. Méth. Dict. no. 49.

Helix decollata, Linneus.

Hab. South of Europe. Sicily and the Canary Islands.

This well-known species occurs in a larger state in some parts of Northern Africa.

Species 335. (Mus. Cuming.)

BULIMUS AURATUS. *Bul. testā oblongo-turrītā, tenuissimā, pellucidā, anfractibus septem, subventricosis, longitudinaliter crebrèrè elevato-striatis, columellā simplici, labro tenui, acuto; aureo-olivaceā, vitredē, lineis minutis fasciis spiraliis obsolete notatā.*

THE GILDED BULIMUS. Shell oblong-turreted, very thin, transparent, whorls seven in number, rather ventricose, longitudinally very closely elevately striated, columella simple, lip thin, sharp; golden-olive, glassy, obsoletely marked with minute spiral brown lines.

PFEIFFER, Pro. Zool. Soc. 1846. p. 32.

Hab. —?

A delicate glassy species of the substance and colour of a *Fritilia*.

Species 336. (Mus. Cuming.)

BULIMUS KINGII. *Bul. testā acuminato-ovatā, crassiusculā, vix umbilicata, anfractibus septem, levibus aut obscurè striatis, columellā latè reflexā, labro simplici; albā, lineis fasciis frequentibus longitudinaliter pictā, columellā fuscā, aperturae fauce fuscescente.*

KING'S BULIMUS. Shell acuminate-ovate, rather thick, scarcely umbilicated, whorls seven in number, smooth or obscurely striated, columella broadly reflected, lip simple; white, longitudinally painted with numerous brown lines, columella brown, interior of the aperture light brown.

GRAY, Annals of Philosophy, New Series, vol. ix. p. 414.

Wood, Ind. Test. Supp. pl. 7. f. 27.

Hab. New Holland; King.

The brown colouring of the columella has a kind of metallic hue.

Species 337. (Mus. Metcalfe.)

BULIMUS MAXIMUS. *Bul. testā oblongā, ventricosā, anfractibus sex, tenuidis, subrotundatis, longitudinaliter ruvē striatis, penultimo granulato, prope apicem subtilius radiatum costellatis, columellā subrectā, crassiusculā, aperturā parvissimā, labro solidulo, reflexo; sordide albā, purpureo-brunnecente tintetā, epidermide luteo-oliveacea nitente indutā, columellā et peritremate albīs.*

THE VERY LARGE BULIMUS. Shell oblong, ventricose, whorls six in number, swollen, somewhat rounded, longitudinally rudely striated, last whorl but one granulated, finely radiately ribbed near the apex, columella nearly straight, somewhat thick, aperture rather small, lip solid, reflected; dirty white, stained with light purple-brown, covered with a shining yellowish-olive epidermis, columella and peritreme white.

Cochlogena maximas, Sowerby, Tankerville Catalogue, Appendix, p. vii.

Bulimus Kremnoicus, D'Orbigny.

Hab. Santa Cruz, Bolivia; D'Orbigny.

Although a large species, is this inferior in size to the *B. Popelairianus*, from which it differs in its more oblong proportions and smaller aperture.

Species 338. (Mus. Cuming.)

BULIMUS EFFEMINATUS. *Bul. testā acuminato-oblongā, subumbilicata, tenuicnlā, anfractibus octo, subplanulatis, longitudinaliter exiliter impresso-striatis, columellā reflexā, labro simplici, aente; carulecent-albā, apicem versus purpurascētē-fuscis, strigis angustis fasciis distantibus.*

THE EFFEMINATE BULIMUS. Shell acuminate-oblong, slightly umbilicated, rather thin, whorls eight in number, somewhat flattened, longitudinally delicately impressively striated, columella reflected, lip simple, sharp; bluish white, purple-brown towards the apex, with distinct narrow stripes of the same colour.

Hab. —?

A delicate opake-white shell gradually more deeply tinged with purple-brown towards the apex.

Species 339. (Mus. Cuming.)

BULIMUS OLIVIERI. *Bul. testā ovatā, ventricosā, anfractibus sex ad septem, longitudinaliter exiliter plicato-costellatis, intersticiis subtilissimè decussatis et punctatis, columellā reflexā, margine incrassatā, labro incrassato, reflexo; sericeo-brunneā, suboliveacea, peritreme flavicante.*

OLIVIER'S BULIMUS. Shell ovate, ventricose, whorls six to seven in number, somewhat flattened, longitudinally delicately plicately ribbed, with the interstices very finely decussated and punctured, columella reflected, thickened at the edge, lip thickened and reflected; silken-brown, of an olive hue, peritreme yellowish.

PFEIFFER, Monog. *Bul.* p. 117.

Hab. Abyssinia.

Au exceedingly beautiful species in fine condition, when examined with the lens.



340



341



342



343



344



345

B U L I M U S.

PLATE LII.

Species 340. (Mus. Cuming.)

BULIMUS SYLVATICUS. *Bul. testā elongato-turrītā, gracilimā, solidinsculā, anfractibus undecim, plano-concavis, levibus, aut minutissimā striatis, politis, apice sub-papillaribus, aperturā parrā, columellā labroque simplicibus; eburneā, epidermide tenui stramineā partim induitā.*

THE SYLVAN BULIMUS. Shell elongately turreted, very slender, rather solid, whorls eleven in number, flatly convex, smooth, polished, somewhat papillary at the apex, aperture small, lip and columella simple; ivory white, partially covered with a light straw-coloured epidermis.

WAGNER, Spix. Test. Bras. p. 11. pl. 10. f. 4.

Hab. Mexico and Brazil (in the woods); Spix.

A semitransparent ivory-white shining shell, partially covered with a slight epidermis.

Species 341. (Mus. Cuming.)

BULIMUS TEREBRATOR. *Bul. testā elongato-turrītā, tenui, pellucidā, apice tumidā, subtruncatā, anfractibus novem nudique creberim concentricē elevato-striatis, columellā tenui, simplicissimā, introrsum arcuatā, labro tenuissimo, acuto; cornuo-luteo.*

THE TERETE BULIMUS. Shell elongately turreted, thin, transparent, whorls nine in number, swollen at the apex, somewhat truncated, very closely concentrically sculptured throughout with raised striae, columella thin, very simple, arched inwardly, lip very thin and sharp; horny yellow.

FERUSSAC, Lamarek, Anim. sans vert. (Deshayes' edit.) vol. 8. p. 234.

Bulimus haplostylus, Pfeiffer.

Hab. El Catamaja, near Loxa, Republic of Equador, South America; Hartweg.

Chiefly distinguished by the very simple structure of the columella and the raised concentric striae which have almost the appearance of fine ribs.

I am indebted to a visit to the collection of the Rev. S. C. E. Neville Rolfe of Heacham Hall, for having had my attention directed to the above recorded species of De Féüssac, with which Dr. Pfeiffer does not seem to have been acquainted.

Species 342. (Mus. Cuming.)

BULIMUS CALCAREUS. *Bul. testā elongato-turrītā, apicem versus solidinsculā, tenuidā, anfractibus duodecim, longitudinaliter subtiliter striatis, striis minutis impressis spiralibus decussatis, aperturā pareā, columellā labroque simplicibus; calcarea-allā, epidermide castaneo-brunneō nitente induitā.*

THE LIME-WHITE BULIMUS. Shell elongately turreted, rather solid and swollen towards the apex, whorls twelve in number, longitudinally finely striated, decussated with minute spiral impressed striae, aperture small, lip and columella simple; chalk-white, covered with a shining chesnut brown epidermis.

WAGNER, Spix. Test. Bras. p. 10.

Helix calcarea, Born.

Bulimus maritimus, Spix, Deshayes.

Hab. Brazil,

M. Deshayes is correct in assigning this species to the *B. maritimus* of Spix, but it will be found to agree also with the *Helix calcarea* of Born, represented in the Mus. Ces. Vindobonense, stripped of its rich brown chesnut epidermis, beneath which it is of a peculiar opaque lime-white character. It is the largest and most solid of the turreted *Bulimi*, and has not a reflected lip.

Species 343. (Mus. Cuming.)

BULIMUS OBELISCUS. *Bul. testā elongato-turrītā, acuminatā, anfractibus sedecim ad octodecim, plano-concavis, subtiliter obliquā striatis, striis minutissimis spiralibus impressis decussatis, columellā labroque simplicissimis; albā, epidermide tenuicula luteo-olivaceā induitā.*

THE OBELEISK BULIMUS. Shell elongately turreted, acuminate, whorls eighteen in number, flatly convex, finely obliquely striated, and decussated with very minute spiral impressed striae, lip and columella very simple; whitish, covered with a rather thin yellowish epidermis.

Helix obeliscus, Moricand, Mémo. Genève, vol. vi. p. 540. pl. 1. f. 4.

Hab. Near Caravalhas, Brazil; Moricand.

Distinguished by having the largest number of whorls and most sharply terete growth, whilst the epidermis is very thin, and of a light yellowish olive-colour. It should

be observed that the shell grows to a more acuminated point than in the specimen here represented, which has the first whorls broken off.

Species 344. (Mus. Cuming.)

BULIMUS OBTUSATUS. *Bul. testá elongato-turrítá, basi subobliquè productá, umbilicatá, apice tumidá, obtusá, anfractibus decem ad undecim, plano-conexis, obliquè creberrimè costulato-striatis, aperturá ovali, labro incrassato, paululum reflexa; tutescente-albá, rufo-fusco bie illiè irregulariter tinté et strigatá.*

THE BLUNTED BULIMUS. Shell elongately turreted, rather obliquely produced at the base, umbilicated, swollen and obtuse at the apex, whorls ten to eleven in number, flatly convex, obliquely very closely sculptured with rib-like striae, aperture oval, lip thickened, a little reflected; yellowish-white, irregularly stained and streaked here and there with red-brown.

PFEIFFER, Monog. *Bul.* p. 151.

Helix obtusa, Gmelin.

Bulinus calcareus, Bruguière, Lamarck, Deshayes.

Hab. Madagascar; Beck.

This shell, distinguished by Lamarek as having a reflected lip, has been confounded with the *Helix calcarea* of Born, an examination of whose figure will show that it represents a worn specimen of Sp. 342, deprived of its epidermis.

Species 345. (Mus. Cuming.)

BULIMUS CLAVATOR. *Bul. testá elongato-oblongá, basi tumidá et subobliquè productá, ex umbilicato, anfractibus octo, longitudinaliter rudi elevato-striatis, ad suturas crenulatis, columellá subdepresso-reflexá, labro simplici, aut paululum reflexo, superne compresso; pallidè livido-olivacea.*

THE CLUB-BEARER BULIMUS. Shell elongately oblong, swollen and somewhat obliquely produced at the base, scarcely umbilicated, whorls eight in number, longitudinally rudely striated, crenulated at the sutures, columella somewhat depresso reflected, lip simple or a little reflected, compressed at the upper part; pale livid olive.

PETIT, Mag. de Zool. 1844, pl. 94.

Hab. Madagascar.

The sutures, as in many species of this form, are white.



34.7



34.8



34.8



34.9



35.0



35.2



35.2

BULIMUS.

PLATE LIII.

Species 346. (Mus. Cuming.)

BULIMUS CONSIMILIS. *Bul. testā oblongā, subcylindraceā, basi rotundatā, tenui, anfractibus sex ad septem, longitudinaliter tenui-striatis, ultimo ventricosiusculo, columellā simplicissimā, labro simplici; brunnescente, nitente, striis saturatioribus hic illic notatā, aperture fauce carulescente.*

THE SIMILAR **BULIMUS.** Shell oblong, somewhat cylindrical, rounded at the base, thin, whorls six to seven in number, longitudinally finely striated, last whorl rather ventricose, columella very simple, lip simple; light brown, marked here and there with streaks of darker brown, interior of the aperture bluish.

Hab. —?

This shell approaches nearest to the *B. Tannaisii*, but is certainly distinct.

Species 347. (Mus. Cuming.)

BULIMUS TENEBRICUS. *Bul. testā cylindraceo-oblongā, anfractibus septem ad octo, subrotundatis, prope suturas subtilissimè reticulatis, margine crenulatis, columellā revolutā, labro simplici, acuto; albida, purpureo-roseo tintata, purpureo-nigricante conspicue grandimaculata.*

THE DARKENED **BULIMUS.** Shell cylindrically oblong, whorls seven to eight in number, rather rounded, very finely reticulated near the sutures, crenulated along the edge, columella rolled back, lip simple, sharp; whitish, stained with purple-rose and conspicuously painted with large blotches of purple-black.

Hab. Ibu, West Africa; Fraser.

Distinguished from *B. flammatus*, by its dwarf-like growth and rich purple-rose hue. The specimen here represented, was collected by Mr. Fraser during the unfortunate Expedition to explore the source of the Niger.

Species 348. (Mus. Cuming.)

BULIMUS SPECTRALIS. *Bul. testā cylindraceo-oblongā, vix umbilicatā, anfractibus septem, locis, columellā revolutā, labro tenui, acuto; albida, epidermide tenui straminei indutā.*

THE SPECTRAL **BULIMUS.** Shell cylindrically oblong,

slightly umbilicated, whorls seven in number, smooth, columella rolled back, lip thin, sharp; whitish, covered with a thin straw-coloured epidermis.

Hab. —?

Care must be taken not to confound this species with the uncoloured variety of *B. flammatus*, from which it differs in form, and in not being reticulated.

Species 349. (Mus. Metcalfe.)

BULIMUS INCA. *Bul. testā cylindraceo-oblongā, umbilicatā, anfractibus septem ad octo, longitudinaliter plicato-striatis, suturis subprofundis, columellā latè reflexa, labro paululum reflexo; nigricante-brunneā, sublividacē, columellā albā, aperture fauce purpureo-cinereā.*

THE INCA **BULIMUS.** Shell cylindrically oblong, umbilicated, whorls seven to eight in number, longitudinally plicately striated, sutures rather deep, columella broadly reflected, lip a little reflected; blackish-brown, tinged with olive, columella white, interior of the aperture purple-ash.

D'ORBIGNY, Voy. dans l'Amér. Mérid. p. 292. pl. 38. f. 6, 7.

Hab. Yungas and Cochabamba, Bolivia; D'Orbigny.

The specimen here represented from the collection of Mr. Metcalfe, is the only one that I am aware of in this country.

Species 350. (Mus. Dennison.)

BULIMUS SUFFUSUS. *Bul. testā elongato-cylindraceā, umbilicatā, anfractibus novem, subobsoletè plicato-striatis, marginis crenulatis, locis, utidis, suturis profundis, apertura parvā, oblongā, columellā revolutā, labro simplici, acuto; albā, roseo suffusā, epidermide tenui lutescente indutā.*

THE BLUSHING **BULIMUS.** Shell elongately cylindrical, umbilicated, whorls nine in number, rather obsoletely plicately striated, crenulated at the margin, smooth, shining, sutures deep, aperture small, oblong, columella rolled back, lip simple, acute; whitish, suffused with rose, and covered with a thin yellowish epidermis.

Hab. —?

Of a peculiarly swollen cylindrical form, approximating to *B. chrysallidiformis* in an immature state, but of a totally different texture and detail of character.

Species 351. (Mus. Cuming.)

BULIMUS NUMIDICUS. *Bul. testā pyramidali-turritā, anfractibus novem, subrotundatis, subtilissimè reticulato-striatis, margine crenulatis, columellā revolutā, labro simplici, acuto; lutescente-albā, rufo-castaneo peculiariter flammēd.*

THE NUMIDIAN BULIMUS. Shell pyramidal turreted, whorls nine in number, somewhat rounded, very finely reticulately striated, crenulated at the edge, columella rolled back, lip simple, sharp; yellowish white, peculiarly flamed with reddish chesnut.

Hab. Prince's Island, West Africa.

This species has been probably confounded by many with the *B. flammatus*, from which it constantly differs in form, sculpture, and colouring.

Species 352. (Mus. Cuming.)

BULIMUS FLAMMEUS. *Bul. testā oblongo-turritā, anfractibus octo, subrotundatis, reticulato-striatis, columellā revolutā, labro simplici, acuto; lutescente-albā, apicem versus purpureo-rubente, castaneo-brunneo latiflammēd, aperture fauce corruſcente.*


THE FLAMED BULIMUS. Shell oblong-turreted, whorls eight in number, somewhat rounded, reticulately striated, columella rolled back, lip simple, sharp; yellowish-white, tinged with purple-red towards the apex; and broadly flamed with dark chesnut-brown, interior of the aperture bluish.

BRUGUIERE, Enc. Méth. vol. i. p. 322.

Helix flammata, Muller.

Hab. Banks of the Niger, Africa; Fraser.

Distinguished from the preceding species by its more oblong form, absence of sutural crenulations, purple-rose colouring, and broader flames.



BULIMUS.

PLATE LIV.

Species 353. (Fig. 34 and 35.) Mus. Cuming.)

BULIMUS ROSEATUS. *Bul. testā oblongo-ovato, levicula, vix umbilicata, austrorubris sex ad septem, subcentrōcosis, transversis minutis interpresso-striatis, columellā reflexā, labro sollempniter lani, acuto; albido, roseo exiliter tincto, striis fuscocinereis longitudinalibus interdina superne notata.*

THE ROSEATE BULIMUS. Shell oblong-ovate, rather thin, scarcely umbilicated, whorls six to seven in number, rather ventricose, transversely very minutely impressly striated, columella reflected, lip somewhat expanded, thin, sharp; whitish, faintly tinged with rose, and sometimes marked round the upper part with brownish longitudinal streaks.

Hab. Venezuela.

A delicately coloured species, approximating closely in form and style of painting to the *B. oreades*.

Species 354. (Mus. Cuming.)

BULIMUS FERIATUS. *Bul. testā acuminato-oblongā, vix umbilicata, austrorubris sex, transversis sub lente minutiissimè interpresso-striatis, columellā reflexā, labro expanso, simplici, vix reflexo; virescente-nigra, maculis panceo-fuscis quadratis prope apicem notatis, fasciā conspicuā circa umbilicum.*

THE UNEMPLOYED BULIMUS. Shell acuminately oblong, scarcely umbilicated, whorls six in number, transversely very minutely impressly striated beneath the lens, columella reflected, lip expanded, simple, slightly reflected; greenish-white, marked near the apex with a few square olive-brown spots, and a conspicuous band round the umbilicus.

Hab. Venezuela.

Another interesting new species from the same locality as the preceding.

Species 355. (Mus. Cuming.)

BULIMUS PERDIX. *Bul. testā octato-oblongā, solidissimcula, umbilicata, austrorubris quinque ad ser, longitudinaliter creberrimē elevato-striatis, gibbosascutis, columellā obscure plicata, labro reflexo; albido, flammis nunculis punctisque cordeco-nigricantibus variegata, columellā superne purpurascente.*

THE PARTRIDGE BULIMUS. Shell ovately oblong, rather solid, umbilicated, whorls five to six in number, longitudinally very closely elevatedly striated, rather gibbous, columella obscurely plaited, lip reflected; whitish, variegated with blue-black flame blotches and dots, columella purplish at the upper part.

PFEIFFER, Pro. Zool. Soc., 1847.

Hab. New Granada; Funek.

This and the three following having been described as separate species, and not to differ in a case which may be regarded a matter of opinion, they are here published as such; I cannot however forbear expressing my conviction that notwithstanding their apparent differences of form and detail of growth, they are but so many modified conditions of the same species. In the state described as *B. perdix*, the longitudinal striae are extremely regular, the aperture is rounded, and the plait of the columella is only obscurely developed at the upper part.

Species 356. (Mus. Cuming.)

BULIMUS EURYOMPHALUS. *Bul. testā fusiforme-oblongā, solidissimcula, umbilicata, basi angulato-compressa, australibus quinque ad sex, creberrimē longitudinaliter striatis, columellā plicata, aperturā trigono-contracta, labro reflexo; cerasescate-alba, punctis maculisque fuligineantibus nigricantibus variegata, columellā intense purpurea.*

THE BROAD-PLAITED BULIMUS. Shell fusiformly oblong, rather solid, umbilicated, angularly compressed at the base, whorls five to six in number, very closely longitudinally striated, columella plaited, aperture triangularly contracted, lip reflected; bluish-white, clouded with black zigzag blotches and dots, columella deep purple.

Hab. Venezuela; Dyson.

Chiefly distinguished by its clouded zigzag style of painting.

Species 357. (Mus. Cuming.)

BULIMUS GLABER. *Bul. testā subfusiforme-ovato, umbilicata, crassā, solidā, basi subcontracta, australibus quinque ad ser, longitudinaliter corrugato-striatis, columellā plicata, labro invassato, reflexo; albido, striis griseorubentibus nebulosatā et variegata.*

THE SMOOTH BULIMUS. Shell somewhat fusiformly ovate, umbilicated, thick, solid, slightly contracted at the base, whorls five to six in number, longitudinally striated in a wrinkled manner, columella plaited, lip thickened, reflected; whitish, clouded and variegated with greyish-red streaks.

GMELIN, Syst. Nat. p. 3436.

Cochlogena auris-caprinus, De Férrussac.

Auricula Sileni, Lamarck.

Bulinus agotis, Menke.

Hab. West Indies.

This has a somewhat closer affinity with the two following species than with the two which precede.

Species 358. (Mus. Dennison.)

BULIMUS DISTORTUS. *Bul. testā fusiformi-oblongā, crassā, solidā, basi angulato-compressā, anfractibus quinque ad sex, longitudinaliter corrugato-striatis, ad suturas interdum crenulatis, columellā fortiter plicatā, aperturā trigono-contractā, labro incassato, reflexo; albida, cinereo-rubente variegatā et partim strigatā, columellā intensē purpureā.*

THE DISTORTED BULIMUS. Shell fusiformly oblong, thick, solid, angularly compressed at the base, whorls five to six in number, longitudinally striated in a wrinkled manner, sometimes crenulated at the sutures, columella strongly plaited, aperture triangularly contracted, lip thickened, reflected; whitish, variegated and partially streaked with ash-red, columella deep purple.

BRUGUIERE, Enc. Méth. vol. i. p. 344.

Arca Mida distorta, Chemnitz.

Cochlogena distorta, De Férrussac.

Plekocheilos distortus, Beck.

Hab. Venezuela.

This is the type of the group, Sp. 355 to 358, which, as before mentioned, I am inclined to think are but one species in different states.

Species 359. (Mus. Cunning.)

BULIMUS RIMATUS. *Bul. testā cylindraceo-elongatā, tenui, subcornutā, latè umbilicatā, anfractibus sex, lāribus, nitidiusculis, columellā expansā, aperturā suborbiculari et productā; fuscescente.*

THE CLEFTED BULIMUS. Shell cylindrically elongated, thin, somewhat horny, broadly umbilicated, whorls six in number, smooth, rather shining, columella expanded, aperture nearly orbicular and produced; light brown.

PFEIFFER, Pro. Zool. Soc., 1846. p. 112.

Hab. —?

Mainly distinguished by the rounded *Pupa*-like growth of the aperture.

Species 360. (Mus. Cunning.)

BULIMUS CULMINEUS. *Bul. testā oblongo-ovatā, crassissimā, vix umbilicatā, anfractibus septem, minutissimè placo-granulatis, longitudinaliter ruditè elevato-striatis, columellā reflexā, subcontorta, labro simplici, subexpanso; albida, apicem versus cinereo-fuscescente tincta, lineolis rufis hic illuc notata.*

THE LOFTY-DWELLING BULIMUS. Shell oblong-ovate, rather thick, scarcely umbilicated, whorls seven in number, very minutely flatly granulated, longitudinally rudely elevately striated, columella reflected, somewhat twisted, lip simple, slightly expanded; whitish, stained with pale brownish ash-colour towards the apex, and marked here and there with longitudinal red lines.

For authority, locality, and remarks on this species see Pl. XVII. Sp. 98, in which place a shell is represented agreeing better with the *B. lithocicus* of D'Orbigny.

Species 361. (Mus. Dennison.)

BULIMUS SCHIEDEANUS. *Bul. testā ovatā, crassā, subampliè umbilicatā, anfractibus sex, concisis, longitudinaliter rugoso-striatis, columellā reflexā, intercrassatā, labro incassato, non reflexo; calcareo-albā, apertura saepe croceo-aureata.*

SCHIEDE'S BULIMUS. Shell ovate, rather largely umbilicated, thick, whorls six in number, convex, longitudinally roughly striated, columella reflected, thickened, lip thickened, not reflected; chalk-white, interior of the aperture tinged with saffron-orange.

PFEIFFER, Monog. Bul. p. 187.

Hab. Mexico.

A very solid opaque chalk-white shell, tinged with saffron orange in the interior.



365



366



367



368



369



366 A



BULIMUS.

PLATE LV.

Species 362. (Mus. Dennison.)

BULIMUS VELUTINUS. *Bul. testā conico-ovatā, inflatā umbilicatā, anfractibus sex, ultimo marino, striis obtusis longitudinalibus et transversis cerebrinē decussatis; columellā peculiariter reflexā, labro tenui, paululum reflexā; intus exustus rievidē brunnēa.*

THE VELVETTY BULIMUS. Shell conically ovate, inflated, umbilicated, whorls six in number, the last very large, very closely decussated with obtuse longitudinal and transverse striae; columella peculiarly reflected, lip thin, a little reflected; bright brown within and without.

PFEIFFER, *Symbola*, vol. i. p. 42.

Bulimus strigilosis, Dufo.

Hab. Seychelle Islands.

A very delicately formed, largely inflated shell of a rich brown-colour, the surface being decussated in a manner that gives it a soft velvety appearance.

Species 363. (Mus. Dennison.)

BULIMUS JONASI. *Bul. testā subfusiformi oblongā, vix umbilicatā, anfractibus sex, longitudinaliter rugosoplicatis, columellā reflexā, labro simplici; diaphanā, plūcī opaco-lacteis, columellā purpurascente.*

JONAS' BULIMUS. Shell somewhat fusiformly oblong, scarcely umbilicated, whorls six in number, longitudinally roughly plicated, columella reflected, lip simple; diaphanous, folds opaque cream-colour, columella light purple.

PFEIFFER, *Philippi*, Abild. *Bul.* pl. 5. f. 4.

Hab. Vera Cruz, Central America.

A very delicate and graceful shell, having the longitudinal folds of an opaque cream-colour, whilst the interstices are transparent, and the columella tinged with purple.

Species 364. (Mus. Cuming.)

BULIMUS ANDICOLA. *Bul. testā pyramidali-oblongā, subumbilicatā, anfractibus septem, levibus, columellā reflexā, aperturā parvissimā, labro simplici, acuto; albida, strigis perpansis rufo-fuscescentibus hic illuc longitudinaliter pictā.*

THE ANDES-INHABITING BULIMUS. Shell pyramidalily oblong, slightly umbilicated, whorls seven in number, smooth, columella reflected, aperture rather small, lip simple, acute; white, longitudinally painted here and there with a very few reddish-brown streaks.

PFEIFFER, *Pro. Zool. Soc.*, 1846, p. 115.
Hab. Columbian Andes.

Of somewhat doubtful character and apparently of immature growth.

Species 365. (Mus. Cuming.)

BULIMUS PALLIDIOR. *Bul. testā acuminato-ovatā, subventricosa, ampliè umbilicatā, anfractibus septem, levibus, vel obscurè striatis, columellā labroque expansa; albā, intus aurantio tincta.*

THE PALE BULIMUS. Shell acuminately ovate, rather ventricosely, largely umbilicated, whorls seven in number, smooth, or obscurely striated, lip and columella expanded; white, tinged in the interior with orange.

SOWERBY, *Conch. Illus. Bulimus*, Fig. 39 and 44.

Hab. — ?

Very like *B. proteus* in general aspect, but destitute of granules or any similar sculpture.

Species 366. (Fig. a and b, Mus. Cuming.)

BULIMUS VINCENTINUS. *Bul. testā fusiformi-ovatā, tenue culā, subpellucidā, anfractibus sex, levibus, nitentibus, sub lente transversi minutissimè inciso-striatis, columellā plano-reflexā, aperturā oblongā, labro subexpanso. Intercostæ albā, interdum castaneo fasciata.*

THE ST. VINCENT BULIMUS. shell fusiformly ovate, rather thin, somewhat transparent, whorls six in number, smooth, shining, transversely very minutely striated in an engraved manner under the lens, columella flatly reflected, aperture oblong, lip somewhat expanded; yellowish white, sometimes banded with chestnut.

PFEIFFER, *Pro. Zool. Soc.*, 1846, p. 30.

Hab. Venezuela; Linden. Island of St. Vincent?

I fear the locality assigned to this species in the name given to it by Dr. Pfeiffer, is somewhat doubtful.

Species 367. (Mus. Cuming.)

BULIMUS INTERSTINCTUS. *Bul. testā cylindraceo-pyramidalis, non umbilicatā, anfractibus septem, subrotundatis, obliquè ruditè striatis, infra suturas peculiariter crevulatis, columellā subcontortā, aperturā parvā, labro tenui, simplici; peltincedo albā, strigis perpansis fuscis hic illuc obliquè pictā, epidermide tenui cornēa iudicata. columellā rosata.*

BULIMUS.—PLATE LV.

THE DIVIDED BULIMUS. Shell cylindrically pyramidal, not umbilicated, whorls seven in number, somewhat rounded, obliquely rudely striated, peculiarly crenulated beneath the sutures, columella slightly twisted, aperture small, lip thin, simple; transparent white, obliquely painted here and there with a very few brown stripes, covered with a thin horny epidermis, columella pink.

GOULD, Pfeiffer, Monog. *Bul.* p. 169.

Eadem var.

Bulimus rubicundulus, Gould.

Hab. Cape Palmas, West Africa.

An interesting species distinguished by a peculiar beaded row of fine crenulations beneath the sutures, and a delicate pink columella.

able character, but certainly distinct from any other that I am acquainted with.

Species 369. (Mus. Cuming.)

BULIMUS MANUPICTUS. *Bul. testā acuminato-ovatā, basi subattenuatā, umbilicatā, anfractibus sex, lóbibus vel obscurè striatis, columellā reflexā, labro simplici; albocaucaute, purpureo-fusco conspicue longitudinaliter virgatā.*

THE HAND-PAINTED BULIMUS. Shell acuminate ovate, somewhat attenuated at the base, whorls six in number, smooth or obscurely striated, columella reflected, conspicuously longitudinally striped with purple-brown.

Hab. Andes of Columbia.

Of a somewhat diamond-shaped form.

Species 370. (Mus. Cuming.)

BULIMUS NIGROFASCIATUS. *Bul. testā ovatā, subventricosa, viz umbilicatā, anfractibus sex, subtiliter rudē striatis, columellā reflexā, labro tenui, acuto; pellucido-albā, fasciis latis fusco-nigricantibus conspicue pictā.*

THE BLACK-BANDED BULIMUS. Shell ovate, somewhat ventricose, slightly umbilicated, whorls six in number, finely rudely striated, columella reflected, lip thin, sharp; transparent white, conspicuously painted with broad brownish-black bands.

PFEIFFER, MSS. Mus. Cuming.

Hab. Valley of the Magdalina, New Granada.

Distinguished by its broad dark bands.

THE UNIMPORTANT BULIMUS. Shell somewhat acuminate oblong, rather thin, slightly umbilicated, whorls six in number, roughly striated and obscurely rudely flatly granulated, columella broadly reflected, lip thin and simple; fawn-white, irregularly painted with a few white-mottled brown streaks.

Hab. —?

A delicately painted species not possessing any remark-

Bulinus, Pl. LXI.



1 in. = 25 mm. = 100

1 in. = 25 mm. = 100

1 measure = one half "

BULIMUS.

PLATE LVI.

Species 370. (Mus. Cuming.)

BULIMUS MERIDIONALIS. *Bul. testā subtriangularē, umbilicatā, tenui, diaphanā, anfractibus septem, obliquē minutiō rūde striatis, columellā posterioriter late reflexā, labro tenui, acuto; pellucido-albā, nō nācā fuscescēt media cingulatā.*

THE MERIDIONAL BULIMUS. Shell ovately conical, umbilicated, thin, diaphanous, whorls seven in number, obliquely minutely rudely striated, columnella peculiarly broadly reflected, lip thin, sharp; transparent white, encircled round the middle with a pale brown zone.

PFEIFFER, Pro. Zool. Soc., 1847. p. 231.

Hab. South Africa.

The general aspect of this shell reminds one somewhat of an immature specimen of the *Achatina alabaster*.

Species 371. (Mus. Cuming.)

BULIMUS LIMONICUS. *Bul. testā elongatā, pyramidalē, subumbilicatā, anfractibus septem, subtundatā, concentricē subradē rugoso-striatis, columellā reflexā, apertura parvissimā, labro simplici; rufescēt-corneā, lineolis albīs longitudinalē plus minusse profusē notatā.*

THE MEADOW BULIMUS. Shell elongated, pyramidal, slightly umbilicated, whorls seven in number, somewhat rounded, concentrically rather rudely roughly striated, columnella reflected, aperture rather small, lip simple; reddish horny, more or less profusely marked with whitish longitudinal lines.

D'ORBIGNY, Moll. Voy. dans l'Amér. Mérid. p. 284.
Hab. Bolivia; Bridges.

I have every reason to believe this to be the *B. limonicus* of D'Orbigny.

Species 372. (Mus. Cuming.)

BULIMUS CINEREUS. *Bul. testā elongato-oratā, umbilicatā, anfractibus septem ad octo, subtundatā, lacerbā ad concentricē striatis, columellā reflexā, apertura parvā, labro simplici; pallidē cinereā, hic illie obliquē striatā, apice versus pellucido-corneā.*

THE ASH-COLOURED BULIMUS. Shell elongately ovate, umbilicated, whorls seven to eight in number, some-

what rounded, smooth or concentrically striated, columella reflected, aperture small, lip simple; pale ash-coloured, obliquely streaked here and there, transparent horny towards the apex.

Hab. Bolivia.

From a locality well examined by M. Alcide D'Orbigny, but not agreeing with any of the species described in his 'Voyage.'

Species 373. (Mus. Cuming.)

BULIMUS ELECTRUM. *Bul. testā oblongo-ovatā, tenui, pellucida, subitreā, viz umbilicatā, anfractibus sex, subangulatis, transversim minutiō inciso-striatis, columellā reflexā, labro simplici, expanso; pellucido-lataē.*

THE AMBER BULIMUS. Shell oblong-ovate, thin, transparent, somewhat glassy, scarcely umbilicated, whorls six in number, slightly inflated, transversely minutely engraved with striae, columnella reflected, lip simple, expanded; transparent yellow.

Hab. Venezuela.

A delicate transparent shell of somewhat glassy texture, distinguished by its bright amber colour.

Species 374. (Mus. Cuming.)

BULIMUS TUMEFACTUS. *Bul. testā conico-ovatā, latē inflatā, peculiariter umbilicatā, anfractibus quinque ad sex, ultimo maximo, tumido, obliquē plicato-striatis, columellā peculiariter reflexā, labro tenui; pellucido-corneā, epidermide fuscescēt hydrophanā, cingulum angustum punctaque perpanca intermitente, indistā.*

THE PUFFED BULIMUS. Shell conically ovate, broadly inflated, peculiarly umbilicated, whorls five to six in number, the last very large and swollen, obliquely plicately striated, columnella peculiarly reflected, lip thin; transparent horny, covered with a light brown hydrophanous epidermis, leaving a narrow belt and a few small spots.

Hab. Great Bassam, West Africa.

A light horny shell of very peculiar form, distinguished by a second hydrophanous epidermis.

Species 375. (Mus. Dennison.)

BULIMUS CANTATUS. *Bul. testā subfusiformē-oblongā, basi subobliquē productā, umbilicatā, axfractibus sex ad septem, levibus vel subobscure ruditē striatis, columellā medio incurvata, aperturā basi sinuata, labro expanso, subreflexo; pallidē spadiceo-albā, maculis parvis subquadratis distantibus indistincte notata, columellā labroque intus roseo-purpureis.*

THE CHARMED BULIMUS. Shell somewhat fusiformly oblong, somewhat obliquely produced at the base, umbilicated, whorls six to seven in number, smooth or rather obscurely rudely striated, columella curved inwards in the middle, aperture situated at the base, lip expanded, partially reflected: light fawn-white, indistinctly marked with small squarish distant spots, lip and columella coloured interiorly with rose-purple.

Hab. —?

A very characteristic new species from the collection of J. Deunison Esq.

Fig. 376. (Mus. Cuming.)

BULIMUS OBLITUS. *Bul. testā cylindraceo-elongatā, umbilicatā, axfractibus novem, ad suturas peculiariter crenulatis, deinde obscurè obliquē corrugato-striatis, columellā latè reflexā, uniplicatā, aperturā parvā, labro pavulū expenso; pellucido-albā, olivaceo-luteo tinctā.*

THE FORGOTTEN BULIMUS. Shell cylindrically elongated, umbilicated, whorls nine in number, peculiarly crenulated beneath the sutures, then obscurely obliquely striated in a wrinkled manner, columella broadly reflected, with a single plait, aperture small, lip a little expanded; transparent white, tinged with olive-yellow.

Hab. Brazil.

Allied in form to the *B. Joueirensis*, but distinguished amongst other characters by the finely beaded crenulations beneath the sutures.

Species 377. (Mus. Cuming.)

BULIMUS PAZIANUS. *Bul. testā pyramidalē-oblongā tenui, diaphanā, vix umbilicatā, axfractibus sex, levibus, columellā reflexā, labro simplici, acuto; corneo-albā, lineis strigisque fuscis longitudinaliter pictā.*

Paz' BULIMUS. Shell pyramidal oblong, thin, diaphanous, scarcely umbilicated, whorls six in number, smooth, columella reflected, lip simple, sharp; horny white, longitudinally painted with brown lines and streaks.

D'ORBIGNY, Voy. dans l'Amér., Mérid. Moll. p. 286. pl. 32. f. 10, 11.

Hab. Bolivia; D'Orbigny. Guatemala; Sallé.

A delicate transparent shell, well characterized by its linear painting.

Species 378. (Mus. Cuming.)

BULIMUS CALIFORNICUS. *Bul. testā subacuminato-ovatā, tenuiculā, vix umbilicatā, axfractibus sex, levibus, columellā reflexā, labro simplici; lacteā, zonulis interruptis transversis carmineo-nigracantibus cingulatā.*

THE CALIFORNIAN BULIMUS. Shell somewhat acuminate ovate, rather thin, scarcely umbilicated, whorls six in number, smooth, columella reflected, lip simple: cream colour, encircled with interrupted transverse blue-black zones.

Hab. California: Hartweg.

There is little novelty in the character or general aspect of this species, but it is certainly distinct.



380



381



382



383



384



385



386



387 a



387 b



388

1126

BULIMUS.

PLATE LVII.

Species 350. (Mus. Cuming.)

BULIMUS TRICINCTUS. *Bul.* testā acuminato-ovalā, umbilicatā, anfractibus septem, tæribus vel obscurè rudo striatis, aperturā parvā, labro simplici; caro-albidā, usculato-fuscescente longitudinaliter lineatā et strigatā, zonulis tribus nigricantibus basi conspicuè pictā.

THE THREE-GIRT BULIMUS. Shell acuminate ovate, umbilicated, whorls seven in number, smooth or obscurely rudely striated, aperture small, lip simple; bluish-white, longitudinally lined and streaked with burnt-brown, painted in a conspicuous manner at the base with three blackish zones.

Hab. —?

The three dark zones which encircle the lower portion of the whorls are concealed in all but the last.

Species 381. (Mus. Cuming.)

BULIMUS ERUBESCENS. *Bul.* testā pyramidali-oblongā, eis umbilicatā, anfractibus sex, tæribus, transversim sub lente minutiè inciso-striatis, columellā tenuiter reflexā, labro simplici, aente; roseo-lutescente, apice rosaceo.

THE BLUSHING BULIMUS. Shell pyramidal oblong, scarcely umbilicated, whorls six in number, smooth, transversely minutely striated in an engraved manner beneath the lens, columella finely reflected, lip simple, sharp; pinkish yellow, apex pink.

PFEIFFER, Pro. Zool. Soc., 1847. p. 112.

Hab. —?

THE name *erubescens* has been used by Swainson in reference to the *B. goniostoma*, associated with *Goniostoma* as a generic word, but never with *Bulimus*.

Species 382. (Mus. Cuming.)

BULIMUS CHRYSALIS. *Bul.* testā ovato-turrītā, subcylindracea, sub-umbilicata, anfractibus rotundatis, obliquè rugoso-striatis, columellā latè reflexā, aperturā ferè rotundā, labro tenui; interiore purpureo-fusca.

THE CHRYSALIS BULIMUS. Shell ovately turreted, somewhat cylindrical, slightly umbilicated, whorls rounded, obliquely finely roughly striated, columella broadly reflected, aperture nearly round, lip thin; deep purple-brown.

Hab. Guadalupe.

Of a cylindrically turreted form and deep purplish brown hue.

Species 383. (Mus. Cuming.)

BULIMUS SOWERBYI. *Bul.* testā ovato-conicā, tenui, sub-inflatā, umbilicatā, anfractibus sex, tæribus, infra medium subattenuatis, columellā latè reflexā, labro tenui, simplici; albida, strigis fascis oblongis pictā, strigis aliis fascis rectis, medio terminatibus decussatā, zonulis nigricante unicō prope basin.

SOWERBY'S BULIMUS. Shell ovately conical, thin, somewhat inflated, umbilicated, whorls six in number, smooth, slightly attenuated below the middle, columella broadly reflected, lip thin, simple; whitish, painted with oblique brown streaks, crossed with other straight brown streaks terminating in the middle, near the base in a single blackish zone.

PFEIFFER, Pro. Zool. Soc., 1846. p. 112.

Hab. Andes of Columbia.

The double painting of straight lines crossing the oblique on the upper half of the whorls is a very peculiar feature in this shell.

Species 384. (Mus. Cuming.)

BULIMUS STUDERI. *Bul.* testā oblongā, subcylindracea, umbilicata, anfractibus sex, tæribus, columellā reflexā, labro simplici, pantalone expanso; flavigeante-albā, fascis rosae cingulata.

STUDER'S BULIMUS. Shell oblong, somewhat cylindrical, umbilicated, whorls six in number, smooth, columnella reflected, lip simple, a little expanded; yellowish-white, encircled with two pink bands.

PFEIFFER, Pro. Zool. Soc., 1846. p. 112.

Hab. Merida, New Granada.

The two pink bands which encircle the whorls of this shell upon a shining white ground, are very characteristic.

Species 385. (Mus. Cuming.)

BULIMUS PRIMULA. *Bul.* testā oblongā, subcylindracea, umbilicata, anfractibus sex, tæribus vel obliquè subtiliter striatis, columellā latè reflexā, labro simplici, albida, inter hic illuc cincte lineatā, zonis lotis permodere, fascis cingulata, apice purpureo-fusca.

BULIMUS.—PLATE LVII.

THE PRIMROSE BULIMUS. Shell oblong, somewhat cylindric, umbilicated, whorls six in number, smooth or obliquely finely striated, columella broadly reflected, lip simple; whitish, here and there brightly coloured with yellow, encircled with broad pink zones, apex purple-brown.

Hab. Merida, New Granada.

A prettily painted species from the same locality as the preceding, differing slightly, but importantly, in form. The whorls are somewhat more ventricose, and the reflected portion of the columella more appressed.

Species 386. (Mus. Cuming.)

BULIMUS MERIDANUS. *Bul. testá fusiformi-oblongá, subumbilicatá, spirá acuminatá, anfractibus septem, laxeibus, nitensibus, sub tente transversim minute impreso-striatis, columellá reflexá, labro simplici; albida, luteo et roseo eximie tinctá, fasciis pallide purpureis interrruptis cingulatá.*

THE MERIDA BULIMUS. Shell fusiformly oblong, slightly umbilicated, spire acuminated, whorls seven in number, smooth, shining, engraved beneath the lens with minute transverse striae, columella reflected, lip simple; whitish, delicately tinged with rose and yellow, and encircled with interrupted pale purple bands.

PFEIFFER, Pro. Zool. Soc., 1846. p. 33.

Hab. Merida, New Granada.

Similarly delicately painted to the last two species from the same locality, but of very different form.

Species 387. (Fig. a and b, Mus. Cuming.)

BULIMUS LIQUABILIS. *Bul. testá ovato-conicá, ventricosá, umbilicatá, anfractibus sex ad septem, subrotundotis, terebris vel minutissimè striatis, columellá late reflexá, labro tenui, paululum expanso; pellucido-cornéa, strigis longitudinalibus opacis notata.*

THE DISSOLVABLE BULIMUS. Shell ovately conical, ventricose, umbilicated, whorls six to seven in number, somewhat rounded, smooth or very minutely striated, columella broadly reflected, lip thin, a little expanded; transparent horny, marked with opaque streaks.

Hab. Texas; Sallé.

An inflated ovate shell, quite transparent except where ornamented with longitudinal opaque streaks.

Species 388. (Mus. Cuming.)

BULIMUS CONNIVENS. *Bul. testá pyramidalis-conicá, solidiusculá, ampliter umbilicatá, anfractibus septem, oblique creberimè plicato-striatis, columellá late expansa, aperturā rotundatā, labro expanso, paululum reflexo; opaco-albá, nitente, apertura fuce aurantiá.*

THE DISSEMBLING BULIMUS. Shell pyramidal conical, rather solid, largely umbilicated, whorls seven in number, obliquely closely plicately striated, columella broadly expanded, aperture rounded, lip expanded, a little reflected; opaque white, shining, interior of the aperture saffron-orange.

Hab. Senegal.

This shell obtains a pyramidal form from the last whorl being widely convoluted and leaving a large umbilicus.



.123



389 a



389 f



390 a



390 b



396 e



397



393



392



394

B U L I M U S.

PLATE LVIII.

Species 328. (Mus. Cuming.)

BULIMUS MOZAMBIENSIS. *Bul. testā ovoato-conicā, sub-turrītā, subumbilicatā, aufractibūs sex ad septem, lēvibus, columellā latē reflexā, aperturā parvissimā, labo simplici; albā, aufractum parte superiori lineis fasciā brevibus oblique pīctā, medio linea nigra cingulatā, parte inferiori immaculatā.*

THE MOZAMBIQUE BULIMUS. Shell ovately conic, somewhat turreted, slightly umbilicated, whorls six to seven in number, smooth, columella broadly reflected, aperture rather small, lip simple; white, upper part of the whorls painted with short oblique brown lines, encircled round the middle with a black line, lower part unspotted.

PFEIFFER, *Symbolæ*, vol. iii. p. 55.

Hab. Mozambique.

The painting of this species affords a character in being marked only above the line which encircles the whorl at the sutures.

Species 339. (Mus. Cuming.)

BULIMUS ZIEGLERI. *Bul. testā oblongo-conicā, semi-pelucida, rīx umbilicatā, aufractibūs ser., obscurè tenuissimè decessatim striatis, labo simplici; lateo-albida, sapè fasciis castaneis maculoso-interrupitis cingulatā.*

ZIEGLER'S BULIMUS. Shell oblong conical, semitransparent, slightly umbilicated, whorls six in number, obscurely very finely decessately striated, lip simple; yellowish white, frequently encircled with chestnut bands broken up into square spots.

PFEIFFER, *Pro. Zool. Soc.*, 1846. p. 113.

Hab. Central America.

The spots arising out of the interrupted character of the bands are disposed obliquely.

Species 390. (Fig. 390 a, b, c, Mus. Dyson.)

BULIMUS CURIANENSIS. *Bul. testā elongato-oblongā, subcylindraceā, basi obtusè angulatā et ampliter umbilicatā, aufractibūs septem ad octo, plano-convexis, lēvibus vel minutissimā striatis, columellā latē reflexā, aperturā latē expansā, non reflexā, basi attenuatā, subcaualentatā; griseo-albida, fasciis cœruleo-nigricantibus interdum cingulatā, umbilico interdum intenue castaneo, labo intus roseo.*

THE CURIANA BULIMUS. Shell elongately oblong, somewhat cylindrical, obtusely angled at the base and largely umbilicated, whorls seven to eight in number, flatly convex, smooth or very minutely striated, columella broadly reflected, aperture widely expanded, not reflected, attenuated and slightly channelled at the base; greyish-white, sometimes encircled with blue-black bands, umbilicus sometimes very dark chestnut, lip pink within.

Hab. Curiana, Venezuela (on the leaves of Palmas); Dyson.

One of the most interesting of the many beautiful species of shells collected by Mr. Dyson during his recent expedition to Venezuela. It is very closely allied to *B. Knorrī*, but may be readily distinguished by its abrupt angular growth at the base, and large umbilicus.

Species 391. (Mus. Cuming.)

BULIMUS HUMBOLDTI. *Bul. oblongo-conicā, subcetrato-cosā, umbilicatā, suturis impressis, aufractibūs septem, subrotundatā, lēvibus, politis, columellā latē reflexā, labo simplici; porcellaneo-albā, fasciis castaneis tribus, peculiariter tenuē interruptis, infernē cingulatā.*

HUMBOLDT'S BULIMUS. Shell oblong-conic, somewhat ventricose, umbilicated, sutures impressed, whorls seven in number, rather rounded, smooth, polished, columella broadly reflected, lip simple; porcelain white, encircled round the lower part with three peculiarly finely interrupted chestnut bands.

Bolimus Mexicanus, Valenciennes (not of Lamarck) in Humb. *Voy. Zool.* vol. ii. p. 247. pl. 56. f. 1.

Hab. Mexico; Humboldt.

This shell, an authentic specimen of Humboldt's species, received by Mr. Cuming from Paris, is essentially distinct from Lamarck's *B. Mexicanus* (Pl. XL. Sp. 244), though resembling it in general aspect.

Species 392. (Mus. Dyson.)

BULIMUS ERECTUS. *Bul. testā pyramidali-oblongā, umbilicatā, spiri acuminatā, suturis impressis, aufractibūs subrotundatā, longitudinaliter rūde striatis; columellā latē reflexā, labo tenui, simplici; perlucido-corneā.*

THE ERECT BULIMUS. Shell pyramidal oblong, umbilicated, spire acuminated with the sutures impressed,

whorls somewhat rounded, longitudinally rudely striated, columella broadly reflected, lip thin, simple; transparent horny.

Hab. Curiana, Venezuela (on *Cacti*); Dyson.

Of a uniform horn colour with a peculiarly erect spire.

Species 393. (Mus. Dyson.)

BULIMUS CACTICOLUS. *Bal. testá acuminato-oblongá, subventricosá, tenuissimá, umbilicatá, anfractibus septem, subrotundis, obscurè decussatis striatis, columellá reflexá, labro tenui, simplici; pellucido-corneá.*

THE CACTUS-INHABITING BULIMUS. Shell acuminate oblong, somewhat ventricose, very thin, umbilicated, whorls seven in number, somewhat rounded, obscurely decussately striated, columella reflected, lip thin, simple; transparent horny.

Hab. Curiana, Venezuela (on *Cacti*); Dyson.

Very similar to the last in colour and substance, but sufficiently distinguished in form.

Species 394. (Mus. Cuming.)

BULIMUS OPALINUS. *Bal. testá pyramidalí, bass angulatá et subplanatá, vix umbilicata, anfractibus octo, planoconvexis, minitissimè decussatis striatis, columellá reflexá, labro paululum reflexo; opalino-alba, immaculata.*

THE OPAL BULIMUS. Shell pyramidal, angulated and slightly flattened at the base, scarcely umbilicated, whorls eight in number, flatly convex, very minutely decussately striated, columella reflected, lip a little reflected; opal white, unspotted.

SOWERBY, Beechey's Voyage, p. 144. pl. 138. f. 8.

Hab. Brazil.

Distinguished by its acuminate Trochiform growth and pure opal white substance.



395



396a



396



397



398



399



400



401



402



403

BULIMUS.

PLATE LIX.

Species 395. (Mus. Cuming.)

BULIMUS RUFO-NIGER. *Bul. testā conico-turritā, umbili-
catā, aufractibus septem, rotundatis, obliquū irregulariter
striatis, aperturā sabrotundā, labro simplici; nigricante-
castaneū.*

THE RED-BLACK BULIMUS. Shell conically turreted, um-
bilicated, whorls seven in number, rounded, obliquely
irregularly striated, aperture nearly round, lip simple;
blackish chestnut.

FERUSSAC (?)

Hab. Madagascar.

Of a peculiar dark chestnut reddish hue.

Species 396. (Mus. Cuming.)

BULIMUS CASTRENsis. *Bul. testā conico-oblongā, umbili-
catā, aufractibus septem, læcibus, columellā reflexā, labro
simplici; albida, strigis carneo-viridibus albi-denticu-
lati, maculis oblongis opaco-albis peculiariiter notatā.*

THE CAMP-MARKED BULIMUS. Shell conically oblong,
umbilicated, whorls seven in number, smooth, colum-
ella reflected, lip simple; whitish, peculiarly
marked with bluish green streaks denticulated with
white, and opaque white oblong spots.

PFEIFFER, Pro. Zool. Soc., 1846, p. 115.

Hab. —?

It is only by reference to the figure that it is possible to
form an adequate idea of the singular painting of this
species.

Species 397. (Mus. Cuming.)

BULIMUS TRILINEATUS. *Bul. testā cylindraceo-oblongā,
imperforatā, aufractibus sex, longitudinaliter subtiliter
plicato-rugatā, columellā angustā, labro simplici; gri-
geo-albicans, strigis teniabis fuscis longitudinalibus plus
minus densē ornatā, apertura fauce castaneo-fusca.*

THE TRILINEATED BULIMUS. Shell cylindrically oblong,
not umbilicated, whorls six in number, longitudinally
finely plicately wrinkled, columnella narrow, lip simple;
greyish white, more or less thickly ornamented with
fine brown longitudinal streaks, interior of the ap-
erture chestnut-brown.

Helix trilineata, Quoy, Voy. de l'Astrol. vol. ii. p. 107. pl.
9. f. 1 to 3.

Hab. Port George, New Holland.

The painting of this species is probably variable in its
character, as the specific name given to it by M. Quoy does
not strictly apply to the specimen here figured.

Species 398. (Mus. Cuming.)

BULIMUS NITELINUS. *Bul. testā oblongo-oratā, ad basin
subattenuatā, vix umbilicatā, aufractibus quinque ad
sex, locibus, medio transversis, columellā paululum
reflexā, labro simplici; albida, roseo-aureantio tinti,
fusciis vacuulis nigricaudib; remolte interrumpit,
ornatā, aperturā roseā.*

THE GOLD-TINGED BULIMUS. Shell oblong-ovate, slightly
attenuated at the base, scarcely umbilicated, whorls
five to six in number, smooth, rather swollen in the
middle, columella a little reflected, lip simple;
whitish, tinged with rose-orange, ornamented with
remotely interrupted spots and bands, aperture pink.

Hab. Mexico.

Of a warm rose-orange tint, peculiarly marked, though
not of very novel aspect.

Species 399. (Mus. Cuming.)

BULIMUS DETRITUS. *Bul. testā cylindraceo-oblongā, soli-
disculā, vix umbilicatā, aufractibus septem, convexius
calis, ad suturas subtiliter crenulatis, columellā latè
reflexā, aperturā porvissutā, labro subexpanso; sor-
ditate albā, sepe cinerascens longitudinaliter strigatā.*

THE WORN BULIMUS. Shell cylindrically oblong, rather
solid, slightly umbilicated, whorls seven in number,
rather convex, finely crenulated at the sutures,
columella broadly reflected, aperture rather small,
lip somewhat expanded; dead white, frequently lon-
gitudinally streaked with pale ash.

Helix detrita, Müller, Verm. vol. ii. p. 101.

Helix sepiaria, Gmelin.

Cochlogea radula, Féussac.

Cochlogea dealbata, Féussac.

Bulimus melanostriatus, Jan.

Bulimus abbreviatus, Jan.

Bulimus latens, Ziegler.

Hab. Central Europe.

This well-known species is very generally distributed throughout Germany and other parts of Central Europe.

Species 400. (Mus. Cuming.)

BULIMUS HONDURASANUS. *Bul. testā subacuminateo-oblóngā, tenuiculā, umbilicatā, austractibus sex, convexisculis, lèvibus, columellā latē reflexā, labro simplici; pellucido-flavescens, roseo-fusco trifasciatā.*

THE HONDURAS BULIMUS. Shell somewhat acuminate oblong, rather thin, umbilicated, whorls six in number, rather convex, smooth, columella broadly reflected, lip simple; transparent yellowish-white, encircled by three rose-brown bands.

PFEIFFER, Pro. Zool. Soc., 1846, p. 29.

Hab. Honduras; Dyson.

A very delicate transparent rose-tinged shell.

Species 401. (Mus. Cuming.)

BULIMUS SORDIDUS. *Bul. testā conico-ovatā, tenui, sub-amplyter umbilicatā, austractibus sex ad septem, rotundatis, creberimè fortiter elevato-striatis, columellā latē expansā, vix reflexā, labro tenui, simplici; sordidè fusca.*

THE SORDID BULIMUS. Shell conically ovate, thin, rather largely umbilicated, whorls six to seven in number, rounded, very closely strongly elevatedly striated, columella broadly expanded, scarcely reflected, lip thin, simple; dull brown.

Helix sordida, Lesson, Voyage de la Coquille, p. 315. pl. 13. f. 3.

Hab. Environs of Lima.

It has been supposed that Mr. Broderip's *Bulimus Proteus* is the *B. sordidus* of Lesson; but Dr. Pfeiffer has shown it to be a distinct species, as the authentic specimen here figured, obtained by Mr. Cuming from those collected

by M. Lesson, sufficiently verifies. The name *sordidus* applied to Sp. 100, must therefore be cancelled in favour of Mr. Broderip's very significant title of *Proteus*.

Species 402. (Mus. Cuming.)

BULIMUS CONSTRICTUS. *Bul. testā subcylindraceo-ovatā, solidiusculā, profundiè umbilicatā, spiræ suturæ impressæ, austractibus septem, concisis, tenuissimæ striatis, columellā latē expansā, aperturā parviusculā, sub-reflexa; opaco-alba.*

THE CONSTRICTED BULIMUS. Shell somewhat cylindrically ovate, rather solid, deeply umbilicated, sutures of the spire impressed, whorls seven in number, convex, very finely striated, columella broadly expanded, aperture rather small, with the lip a little reflected; opaque white.

PFEIFFER, Symbolic. vol. i. p. 43.

Bulimus Angosturensis, Gruner.

Hab. Angostura.

Dr. Pfeiffer describes this species as having a horny epidermis.

Species 403. (Mus. Cuming.)


BULIMUS SPRATTII. *Bul. testā subgloboso-ovatā, solidiusculā, turgidā, umbilicatā, austractibus sex, longitudin-aliter ruguloso-platicatis, columellā latē expansā, labro subreflexo; albidā, epidermide tenui cornea olivaceo-partim indutā.*

SPRATT'S BULIMUS. Shell somewhat globosely ovate, rather solid, swollen, umbilicated, whorls six in number, longitudinally rather roughly plicated, columella broadly expanded, lip slightly reflected; whitish, partially covered with a thin horny olive epidermis.

PFEIFFER, in Phil. Abbild., *Bulimus*, pl. 4. f. 6.

Hab. Asia Minor, Pisidia; Forbes. Lycia; Spratt.

A turgid delicately plicated species of very distinct form.



BULIMUS.

PLATE LX.

Species 404. (Mus. Cuming.)

BULIMUS CANTORII. *Bul. testā oblongo-cylindracea, soli-
diuscūla, compresē umbilicatā, anfractibus octo, per-
obliquū tenuissimū striatis, columellā concavo-expansā,
aperturā suborbiculari, labro plano-reflexo; pellucido-
cornē, labro albido.*

CANTOR'S BULIMUS. Shell obliquely cylindrical, rather solid, compressly umbilicated, whorls eight in number, very obliquely finely striated, columella concavely expanded, aperture nearly orbicular, lip flatly reflected; transparent horny, lip whitish.

PHILIPPI, Zeitschr. für Mal. 1844. p. 165.

Hab. Environs of Nanking, China.

Although of a subtransparent horny substance, it is rather a stout shell.

Species 405. (Mus. Cuming.)

BULIMUS ROSSMALLERI. *Bul. testā acuminato-oblongā,
subcylindracea, compresē umbilicatā, anfractibus octo,
obliquū crenulariū plicato-striatis, columellā concavo-
expansā, callositate parvā superne umbratā, aperturā
parvā, labro subexpanso; superne fusco, inferne sordide
albā.*

ROSSMALLER'S BULIMUS. Shell acuminate oblong, slightly cylindrical, compressly umbilicated, whorls eight in number, obliquely very closely plicately striated, columella concavely expanded, furnished at the upper part with a small callosity, aperture small, lip slightly expanded; brown at the upper part, dull white about the lower.

PFEIFFER, Pro. Zool. Soc., 1846. p. 113.

Hab. Asia Minor.

It appears to be a general character in the species, to present the above recorded contrast of colour.

Species 406. (Mus. Cuming.)

BULIMUS SYRIACUS. *Bul. testā oblongo-cylindracea, compresē umbilicata, spira suturis linearis sulcatis, anfractibus septem vel octo, planulatis, parum concisis, columellā indistincte uniplicata, aperturā orbiculari, labro plano-reflexo; griseo-albicante, apice fuso, labro albo.*

THE SYRIAN BULIMUS. Shell oblong-cylindrical, compressly umbilicated, sutures of the spire linearly

grooved, whorls seven to eight in number, flattened, but little convex, columella indistinctly one-plaited, aperture orbicular, lip flatly reflected; greyish white, apex brown, lip white.

PFEIFFER, Symbolæ, vol. iii. p. 88.

Hab. Syria.

Distinguished among other characters by the flattened and closely appressed volution of the whorls.

Species 407. (Mus. Cuming.)

BULIMUS LYCICUS. *Bul. testā cylindraceo-ovalā, subven-
tricosā, vis umbilicatā, anfractibus sex ad septem, lāci-
bus, utidissimis, columellā obscurē uniplicata, labro sub-
incassato, reflexo; corneo-lutescente, pellucido, labro
albido.*

THE LYCIAN BULIMUS. Shell cylindrically ovate, rather ventricose, scarcely umbilicated, whorls six to seven in number, smooth, very shining, columella obscurely one-plaited, lip somewhat thickened, reflected; horny yellow, transparent, lip whitish.

PFEIFFER, Phil. Abbild. vol. ii. p. 113. pl. 4. f. 4.

Hab. Lycia; Spratt.

A comparatively stout shell, yet of such clear transparent substance that the axis of the columella may be seen throughout.

Species 408. (Mus. Cuming.)

BULIMUS CANDELARIS. *Bul. testā elongato-oblongā, sinis-
trali, subcylindracea, compresē umbilicata, anfractibus
nove, plano-concisis, subradē tenuisstriatis, columellā
concavo-expansā, aperturā parvā, labro expanso; vor-
dile albā, labro cibreo.*

THE CANDLE-LIKE BULIMUS. Shell elongately oblong, sinistral, somewhat cylindrical, compressly umbilicated, whorls nine in number, flatly convex, rather rudely finely striated, columella concavely expanded, aperture small, lip expanded; dead white, lip ivory white.

PFEIFFER, Pro. Zool. Soc., 1846. p. 40.

Hab. —?

Locality unknown, but probably on the confines of Europe and Asia.

BULIMUS.—PLATE LX.

Species 409. (Mus. Cuming.)

BULIMUS CARNEUS. *Bul. testá cylindraceá, vix umbilicatá, apice obtusá, anfractibus novem, plano-convexis, obliquè subobscure striatis, columella late expansa, subplaticatá, aperturá ferè semiorbiculari, labro subincassato, convexu reflexo; cornéa, pelticeidá, nitidá, labro albo.*

THE FLESHY BULIMUS. Shell cylindrical, scarcely umbilicated, obtuse at the apex, whorls nine in number, flatly convex, obliquely somewhat obscurely striated, columella broadly expanded, slightly plaited, aperture nearly semiorbicular, lip rather thickened, conspicuously reflected; horny, transparent, shining, lip white.

PFEIFFER, Philippi, Abbild. vol. ii. pl. 4. f. 5.

Hab. Lycia; Spratt.

This, like the *B. Lycicus* from the same locality, is a stout shell, although sufficiently transparent to show the axis of the columella.

Species 410. (Mus. Cuming.)

BULIMUS LABROSUS. *Bul. testá cylindraceá, tumidá, solidiusculá, compressè umbilicatá, anfractibus sex ad septem, obscurè oblique tenuistratis, nitidis, columellá concavo-expansá, subplaticatá, aperturá ferè orbiculari, labro incassato, late reflexo, lirá callositate continuo; livido-fuscescente, subpellucidá, labro intenso lacteo.*

THE BROAD-BRIMMED BULIMUS. Shell cylindrical, swollen, rather solid, compressly umbilicated, whorls six to seven in number, obscurely obliquely finely striated, shining, columella concavely expanded, slightly plaited, aperture nearly orbicular, lip thickened, broadly reflected, continuous with a callous ridge; pale livid brown, somewhat transparent, lip dark cream-colour.

OLIVIER, Voy. Levant, vol. ii. p. 222. pl. 31. f. 10.

Cochlogena labrosa, Férrussac.

Pupa labrosa, Lamarck.

Hab. Syria.

This fine species is remarkably characteristic of the semitransparent substance, cylindrical form, and dilated lip typical of the *Bulimii* of Syria and Asia Minor.

Species 411. (Mus. Cuming.)

BULIMUS EHRENCBERGI. *Bul. testá abbreviato-cylindraceá, tumidiusculá, compressè umbilicatá, apice erecto-papillari, anfractibus septem ad octo, levibus, vix striatis, columellá labroque reflexis; sordidè carulescente-alba.*

EHRENCBERG'S BULIMUS. Shell abbreviately cylindrical, rather swollen, compressly umbilicated, apex erectly papillary, whorls seven to eight in number, smooth, slightly striated, lip and columella reflected; dull bluish-white.

PFEIFFER, Pro. Zool. Soc., 1846. p. 113.

Hab. Greece.

Of a stout cylindrical growth with the apex peculiarly erect and papillary.

Species 412. (Mus. Cuming.)

BULIMUS LABIOSUS. *Bul. testá acuminato-cylindraceá, solidiusculá, anfractibus octo, laceribus, politis, columellá latè expansá, subplaticatá, aperturá ferè orbiculari, labro convexo-incassato, latissimè reflexo, laminá callosa continuo; opalino-alba.*

THE WIDE-LIPPED BULIMUS. Shell acuminately cylindrical, rather solid, whorls eight in number, smooth, polished, columella broadly expanded, somewhat plaited, aperture nearly orbicular, lip convexly thickened, very broadly reflected, continuous by a callous lamina; clear opal white.

BRUGUIERE, Eucyclopédie Méthodique, p. 347.

Helix labiosa, Müller.

Pupa Jeheznei, Récluz.

Hab. Island of Socotra, Coast of Africa; Jeheznei.

Distinguished by its solid clear opal-like substance.

Species 413. (Mus. Cuming.)

BULIMUS ALEPI. *Bul. testá oblongo-cylindraceá, compressè umbilicatá, anfractibus septem, obliquè tenuissimè striatis, columellá plano-expansá, subplaticatá, aperturá ferè orbiculari, labro reflexo, callositatibus ferè continuo; corneo-albicante, subpellucidá, suturis opaco-albo marginatis, labro albo.*

THE ALEPO BULIMUS. Shell oblong-cylindrical, compressly umbilicated, whorls seven in number, obliquely very finely striated, columella flatly expanded, somewhat plaited, aperture nearly orbicular, lip reflected, almost continuous with callosities; horny white, transparent, sutures edged with opaque white, lip white.

Cochlogena Alepi, Férrussac, Prodrome, p. 418.

Bulimus halepensis, Pfeiffer.

Hab. Environs of Aleppo, Syria.

Resembling the large *B. labrosus*, both in aspect and general detail of character.



415 a



415 b



416



417



418



419



420



421



422

BULIMUS.

PLATE LXI.

Species 414. (Mus. Cuming.)

BULIMUS SARCODES. *Bul. acuminato-oblonga, tenui, subprofundè umbilicata, aufractibus sex ad septem, tennissimè ruguloso-striatis, striis spirabilis tenuissimè impressis sub leute decussatæ, columellæ reflexæ, labro simplici; pellucido-cornæ, incarnato tinctæ.*

THE FLESH-TINTED BULIMUS. Shell acuminate oblong, thin, somewhat deeply umbilicated, whorls six to seven in number, very finely rugosely striated, and decussated, beneath the lens, with delicate impressed striae, columella reflected, lip simple; transparent, horny, tinged with flesh colour.

PFEIFFER, Pro. Zool. Soc., 1846. p. 30.

Hab. Honduras; Dyson.

An extremely delicate transparent shell.

Species 415. (Mus. Cuming.)

BULIMUS FASCIOLATUS. *Bul. oblongo-cylindracea, solidiuscula, vix umbilicata, apice obtusa, aufractibus septem, plano-convexis, striatulis, prope suturas minutiissimè granulatis, columellæ breviter reflexæ, aperturæ parviusculæ, labro subexpansæ; albida, fasciis longitudinalibus nigricante-fasciis interdum conspicue picta, aperturæ fance intenè castaneo-fasciæ.*

THE LITTLE-BANDED BULIMUS. Shell oblong-cylindrical, rather solid, scarcely umbilicated, obtuse at the apex, whorls seven in number, flatly convex, finely striated, very minutely granulated near the sutures, columella shortly reflected, aperture rather small, lip slightly expanded; whitish, conspicuously painted with longitudinal blackish-brown bands, interior of the aperture deep chestnut-brown.

OLIVIER, Voy. Levant, vol. i. p. 416. pl. 17. f. 5.

Cochlodena fasciolata, Féussac.

Pupa fasciolata, Lamarck.

Hab. Syria, Crete, Rhodes, &c.

The white variety, represented at Fig. 415 b, is still marked with faint longitudinal bands.

Species 416. (Mus. Cuming.)

BULIMUS HOHENACKERI. *Bul. testa oblongo-cylindracea, solidiuscula, vix umbilicata, aufractibus octo, plano-con-*

veris, tennissimè subgranuloso-striatis, columellæ labroque subexpansis; albâ, intus intenè nigricante-fasciâ.

HOHENACKER'S BULIMUS. Shell oblong-cylindrical, rather solid, scarcely umbilicated, whorls eight in number, flatly convex, very finely somewhat granulesly striated, lip and columella a little expanded; white, interior deep chesnut-brown.

KRYNICKI, in Bull. Moscow, vol. ix.

Bulimus xanthostoma, Hohenacker.

Hab. Georgia.

Closely resembling the white variety of the preceding species, especially in regard to the intense dark colouring of the interior.

Species 417. (Mus. Cuming.)

BULIMUS TOURNEFORTIANUS. *Bul. testa elongato-turrita, sinistrali, vix umbilicata, aufractibus duodecim, plano-convexis, lœvibus vel obliquè minitè striatis, columellæ labroque subexpansis, tridentatis; albâ, aperturæ fance aurantio-luteæ.*

TOURNEFORT'S BULIMUS. Shell elongately turreted, sinistral, scarcely umbilicated, whorls twelve in number, flatly convex, smooth or obliquely minutely striated, lip and columella a little expanded, three-toothed; white, interior of the aperture orange-yellow.

Cochlodena Tournefortiana, Féussac in Tournefort, Voy. Levant, vol. iii. p. 308.

Pupa tridentata, Lamarck.

Hab. Turkey in Europe.

The *Clansilia lœvis* of De Blainville, quoted by Dr. Pfeiffer as synonymous with this species, differs in being less elongated and composed of fewer whorls.

Species 418. (Mus. Cuming.)

BULIMUS EBURNEUS. *Bul. testa elongato-turrita, non umbilicata, aufractibus sex, lœvibus, politis, columellæ subreflexæ, labro simplici; eburnea, intus luteo pallide tintata.*

THE IVORY BULIMUS. Shell elongately turreted, not umbilicated, whorls nine in number, smooth, polished, columella a little reflected, lip simple; ivory white, interior tinged with yellow.

PFEIFFER, Symbole, vol. II. p. 44.

Hab. Turkey in Europe.

Of a smooth polished ivory aspect.

Species 419. (Mus. Cuming.)

BULIMUS FORSKALII. *Bul. testā ventricoso-ovalā, cylindraceo-Pupiformi, latē umbilicatā, anfractibus octo, obliquè creberrius elevato-striatis, columellā uniplicatā, aperturā suborbiculari, labro latissimè expanso; pellucido-albā, labo intus fulvo-sanguineo tincto.*

FORSKAL'S BULIMUS. Shell ventricosely ovate, cylindrically Pupa-shaped, widely umbilicated, whorls eight in number, obliquely very closely elevated striated, columella one-plaited, aperture nearly orbicular, lip very much expanded; transparent white, lip tinged within with fulvous-yellow colour.

Bulimus Forskallii, Beck, Ind. p. 68.

Helix sulcata, Chemnitz.

Pupa candida, Lamarek.

Pupa arata, Réchuz.

Hab. Arabia.

This species partakes very much of the character of *Pupa*.

Species 420. (Mus. Cuming.)

BULIMUS TAURICUS. *Bul. testā cylindraceo-turritatā, solidiusculā, compresso-umbilicatā, anfractibus novem ad decem, planiusculis, subobscure obliquè striatis, columellā labroque dilatatis, aperturā parviusculā; albā, apertura fance aurantio-luteā.*

THE TARTARY BULIMUS. Shell cylindrically turreted, rather solid, compressly umbilicated, whorls nine to ten in number, rather flattened, somewhat obscurely obliquely striated, lip and columella dilated, aperture rather small; white, interior of the aperture tinged with orange-yellow.

LANG, Pfeiffer, Monog. *Bul.* p. 226.

Bulimus lineatus, Rossmassler.

Bulimus obsoletus, Ziegler.

Bulimus fusiformis, Menke.

Hab. Tartary.

Very minutely striated beneath the lens.

Species 421. (Mus. Cuming.)

BULIMUS CYLINDRICUS. *Bul. testā cylindraceo-oblongā, sinistrali, tenuiculā, visu umbilicatā, anfractibus octo, convexis, obliquè striatis, columellā expansa, labro simplici; cerasescente-albā, aperturæ fance lutescente tincta.*

THE CYLINDRICAL BULIMUS. Shell cylindrically oblong, sinistral, rather thin, scarcely umbilicated, whorls eight in number, convex, obliquely striated, columella expanded, lip simple; bluish white, interior of the aperture faintly tinged with yellow.

MENKE, MSS.

Hab. —?

The apex of this reversed species has a somewhat papillary aspect.


Species 422. (Mus. Cuming.)

BULIMUS DARDANUS. *Bul. testā acuminato-turritatā, visu umbilicatā, anfractibus octo ad novem, convexis, obliquè striatis, columellā reflexā, aperturā obliquè semiovali, labro simplici; albida, incarnato-fusco obscurè tincta et striata, apertura fance saturatiore.*

THE TROJAN BULIMUS. Shell acuminately turreted, scarcely umbilicated, whorls eight to nine in number, convex, obliquely striated, columella reflected, aperture obliquely semi-oval, lip simple; whitish, obscurely stained and streaked with flesh-tinted brown, interior of a rather darker colour.

Hab. Asia Minor, Turkey in Europe.

Of a more acuminated, less cylindrical form than most of the allied species.



B U L I M U S.

Plate LXII.

Species 423. (Mus. Cuming.)

BULIMUS PALLENS. *Bul. testā conico-ovatā, tenuiculā, anfractibus quinque ad sex, sub lente tenuissima oblique striatis, striis subtilissimis impressis spiralibus exsculptis, anfractū ultimo medio subangulato, columellā tenū, angustā, labro simplici; pallidē stramineā.*

THE PALE BULIMUS. Shell conically ovate, rather thin, whorls five to six in number, beneath the lens very finely obliquely striated, and sculptured with very fine impressed spiral striae, last whorl slightly angled in the middle, columella thin, narrow, lip simple; pale straw-colour.

JONAS, MSS. Mus. Cuming.

Hab. —?

This has very much the appearance of an immature shell, and is published with some doubt.

Species 424. (Mus. Cuming.)

BULIMUS FERRUGINEUS. *Bul. testā conico-ovatā, umbilicata, anfractibus rotundatis, tæcibus, columellā labroque simplicibus; albida, rufo-fusco concentricè strigata, et ad basin spiraliiter interruptè fasciatā.*

THE RUSTY BULIMUS. Shell conically ovate, umbilicated, whorls rounded, smooth, lip and columella simple; whitish, concentrically streaked with reddish-brown, and spirally banded at the base in an interrupted manner.

Hab. Peru.

An interesting little species, distinguished by a very characteristic style of painting.

Species 425. (Mus. Cuming.)

BULIMUS DYSONI. *Bul. testā ovato-turritatā, minutiè umbilicata, anfractibus sex ad septem, rotundatis, irregulariter tenuistriatis, columellā reflexā, labro simplici; fusco-cornē, semipeltucida.*

DYSON'S BULIMUS. Shell ovately turreted, minutely umbilicated, whorls six to seven in number, rounded, irregularly finely striated, columella reflected, lip simple; brown horn colour, semitransparent.

PFEIFFER, Pro. Zool. Soc., 1846, p. 39.

Hab. Honduras.

A bright brown pellucid shell in which the columella is visible throughout.

Species 426. (Mus. Benson.)

BULIMUS KUNAWURENSIS. *Bul. testā elongato-turritatā, sinistrorsā, comprèsè umbilicata, anfractibus duodecim, tæcibus, ant obscurè ruguloso-striatis, ad suturas minutè crenulatis, aperturā pareā, columellā labroque expansis; fuscescente-cornē, strigis opacis hic illuc rotatā, labro intus niveo.*

THE KUNAWUR BULIMUS. Shell elongately turreted, sinistral, compressly umbilicated, whorls twelve in number, smooth, or obscurely roughly striated, minutely crenulated at the sutures, aperture small, lip and columella expanded; brownish horny colour, marked here and there with opaque white streaks, lip snow-white within.

HUTTON, MSS.

Hab. Kunawur, India.

I am indebted for this characteristic Indian species to Mr. Benson, and for the information that it was collected by Capt. Hutton at Chini in Kunawur on the bank of the river Sutlej, before it passes through the snowy range of the Himalaya.

Species 427. (Fig. a and b, Mus. Cuming.)

BULIMUS IRRORATUS. *Bul. testā acuminato-oblongā, media ventricosa, anfractibus sex, subrotundatis, striis tenuidis elevatis interruptis oblique exsculptis, infra suturas peculiariter concentrica crenulatis, columellā strictè uniplicata; rifescente-purpurea, anfractū ultimo epiderma tenui cinerascente, maculis fulvo-albidis, fusco-umbritis aspersa, induto, columellā carnelascente-alba, labro incarnato-roso.*

THE BEDEWED BULIMUS. Shell acuminately oblong, ventricose in the middle, whorls six in number, somewhat rounded, obliquely sculptured with swollen raised interrupted striae, and peculiarly concentrically crenulated beneath the sutures, columella strictly one-plaited; reddish purple, last whorl covered with a pale ash epidermis, sprinkled with brown-shaded fulvous white spots, columella bluish-white, lip flesh-pink.

REEVE, Pro. Zool. Soc., 1849.

Hab. Brazil? New Granada?

Mr. Cuming is indebted for this beautiful new species

to the liberality of A. L. Gubba, Esq., of Havre. It is of a swollen ovate form with the spire rather sharply acuminate, and the columella is distinguished by a sharp winding plait. The ground colour of the shell is a reddish purple, the last whorl being particularly characterized by a thin ash-coloured epidermis, sprinkled with light fulvous spots, all inclining towards the lip which is of a delicate flesh-pink.

Species 428. (Mus. Cuming.)

BULIMUS LINEATUS. *Bul. testá acuminato-conicá, subumbilicatá, anfractibus septem, lèvibus, sub lente minutè striatis, columellá latè reflexá, aperturá subobliquá, labro paululum reflexo; albida, cinnamomeo-fuscescente obliquè lineatá et strigatá, labro niveo.*

THE LINEATED BULIMUS. Shell acuminate conical, slightly umbilicated, whorls seven in number, smooth, minutely striated beneath the lens, columella broadly reflected, aperture rather oblique, lip a little reflected; whitish, obliquely lineated and streaked with light cinnamon brown, lip snow-white.

SPIX, Test. Bras. p. 8. pl. 7. f. 6.

Hab. Brazil.

Not much unlike the *B. cinnamomeo-lineatus* in general appearance.

Species 429. (Mus. Cuming.)

BULIMUS PSEUDO-SUCCINEA. *Bul. testá subfusiformi-oblonga, tenuissimá, vitrea, pellucidá, anfractibus quin-*

que ad sex, subtilissimè striatulis, ultimo inflato, patulo, columellá labroque tenuibus, simplicibus; stramineo-corneá.

THE FALSE-SUCCINEA BULIMUS. Shell somewhat fusiformly oblong, very thin, glassy, transparent, whorls five to six in number, very finely and delicately striated, last whorl inflated, wide open, lip and columella thin, simple; pale straw horn-colour.

POTIEZ and MICHAUD, Gal. de Douai, vol. i. p. 140. pl. 13. f. 5. 6.

Helix pseudo-Succinea, Moricand.

Hab. Bahia, Brazil.

An extremely transparent glassy shell of a delicate pale straw-colour.

Species 430. (Mus. Cuming.)

BULIMUS NATALENSIS. *Bul. testá angulato-conicá, Trochiformi, anfractibus sex, obliquè striatis, ultimo angulato-productis, columellá subincrassatá, labro simplici: fulvæ, nitente, nigro-fusco conspicue fasciatâ.*

THE NATAL BULIMUS. Shell angularly conical, Trochus-shaped, whorls six in number, obliquely striated, last whorl angularly produced, columella rather thickened, lip simple; fulvous, shining, conspicuously banded with black-brown.

KRAUSS, Sudafrikanischen Moll. p. 71. pl. 5. f. 1.

Hab. Port Natal, South Africa.

A light semitransparent trochiform shell, conspicuously dark-bauded at the sutures and round the middle of the last whorl.



BULIMUS.

PLATE LXIII.

Species 431. (Mus. Cuming.)

BULIMUS PUPA. *Bul. testā cylindraceo-oblongā, vix umbilicatā, austractibus septem, obscurē ruditē striatis, aperturā parciusculā, labro subexpanso; pellucido-cornēd, labro albo.*

THE PUPA BULIMUS. Shell cylindrically oblong, scarcely umbilicated, whorls seven in number, obscurely rudely striated, aperture rather small, lip a little expanded; transparent horny, lip white.

BRUGUIERE, Enc. Mith. vol. i. p. 349.

Helix pupa, Linnaeus.

Varietates (vide Pfeiffer).

Bulimus tuberculatus, Turton.

Bulimus evanescens, Deshayes.

Bulimus Babanchii, Anton.

Bulimus labiatus, Ziegler.

Pupa primativa, Menke.

Hab. Sicily, the Morea, Algeria, the Canary Islands, &c.

This species is not found in Britain or Central Europe.

Species 432. (Mus. Cuming.)

BULIMUS LEUCODON. *Bul. testā subelongato-turritatā, vix umbilicatā, austractibus novem, obliquē creberrimè impresso-striatis, aperturā parvā, subquadratā, marginibus jucutis, labro dente valido interno, extus scrobiculatus formante, medio conspicuē munito; olivaceo-cornēd, dente labroque albidis.*

THE WHITE TOOTH BULIMUS. Shell somewhat elongately turreted, scarcely umbilicated, whorls nine in number, obliquely very closely impressively striated, aperture small, somewhat square, margins joined, lip conspicuously furnished in the middle with a strong internal tooth, forming a depression on the outer surface; olive-horny, tooth and lip whitish.

PFEIFFER, Phil. Abbild. Conch. vol. ii. p. 114. *Bulimus*, pl. 4. f. 7.

Hab. Near Trebizond.

Easily distinguished by a prominent internal tooth on the outer lip, whilst there is no indication of one on the columella.

Species 433. (Mus. Cuming.)

BULIMUS SIDONIENSIS. *Bul. testā cylindraceo-turritatā, umbilicatā, austractibus novem, plano-convexis, obliquē*

striatis, aperturā subquadratā, prominulā, labro reflexo. pallidē ceruleo-cinerascente, apice cornēd, aperture fulvo-fuscā.

THE SIDON BULIMUS. Shell cylindrically turreted, umbilicated, whorls nine in number, flatly convex, obliquely striated, aperture somewhat square, prominent, lip reflected; pale bluish ash, apex horny, interior of the aperture fulvous.

Helix Sidoniensis, Fénnec, Prodrome.

Pupa Bulimoides, Pfeiffer.

Hab. Syria.

On the confines of the genera *Bulimus* and *Pupa*, but partaking rather more of the characters of the former.

Species 434. (Mus. Cuming.)

BULIMUS MARTINICENSIS. *Bul. testā acuminate-turritatā, umbilicatā, spirae suturis subimpressis, austractibus septem ad octo, subtiliter obliquē striatis, aperturā subquadrato-ovatā, labro expanso; cornēd, subpellucidā, labro albo.*

THE MARTINIQUE BULIMUS. Shell acuminately turreted, umbilicated, sutures of the spire impressed, whorls seven to eight in number, finely obliquely striated, aperture somewhat squarely ovate, lip expanded; horny, rather transparent, lip white.

PFEIFFER, Pro. Zool. Soc., 1846, p. 40.

Hab. Martinique.

Chiefly distinguished by its acuminated form.

Species 435. (Mus. Cuming.)

BULIMUS PERSPECTIVUS. *Bul. oblongo-turritatā, tenuiculā, ampliē et profunde umbilicatā, austractibus septem, brevis, subtilissimā obliquē striatis, aperturā subquadrato-oblongā; labro parvus expanso; intus extusque intensē castaneā.*

THE PERSPECTIVE BULIMUS. Shell oblong-turreted, rather thin, largely and deeply umbilicated, whorls seven in number, smooth, very finely obliquely striated, aperture somewhat squarely oblong, lip but little expanded; deep chestnut brown within and without.

PFEIFFER, Pro. Zool. Soc., 1846, p. 33.

Hab. —?

Of a uniform transparent deep reddish-chestnut colour.

Species 436. (Mus. Cuming.)

BULIMUS KRAUSSI. *Bul. testā ovato-ventricosā, anfractibus septem, undique pulcherrimè minutiè granulatis, longitudinaliter plicato-striatis, columellā strictè reflexā, labro subincurvato, non reflexo; albida, rufo-fusco apicem versus conspicuè flammatā, et supra suturas peculiariter articulatā, columellā labique limbo interno intenso castaneo-purpureis.*

KRAUSS'S BULIMUS. Shell ovately ventricose, whorls seven in number, very beautifully minutely granulated throughout, longitudinally plicately striated, columella closely reflected, lip rather thickened, not reflected; whitish, conspicuously flamed with reddish brown towards the apex, and peculiarly articulated above the sutures with the same colour, columella and inner edge of the lip deep chestnut-purple.

PFEIFFER, Symbolae, vol. iii. p. 85. Krauss, Die Sudafrikanischen Mollusken, p. 78. pl. 5. f. 4.

Hab. Natal, South Africa (in the woods); Krauss.

This magnificent species, from a locality which has contributed little of particular interest to this genus, partakes very much of the form of *B. oblongus* and its congeners from South Eastern America. In colouring it is characterized by a singular articulated band around the sutures, and the lip and columella are coated with an enamel of deep chestnut-purple. In sculpture the entire surface of the shell is beautifully minutely granulated, besides being plicately striated, a character to which it is proper to direct attention from the difficulty of representing it in a figure.

The discovery of this fine species by Dr. Krauss in the woods of South Africa, gives us some idea of the richly painted snails and other animal forms that may be dwelling at this moment in the vast unexplored regions of that immense continent so fatal to European health and enterprise.

Species 437. (Mus. Cuming.)

BULIMUS PUBESCENS. *Bul. testā oblongo-ovatā, umbilicatā, anfractibus sex ad septem, subrotundatis, sub lente minutissimè reticulatis, columellā uniplicatā, reflexā,*

labro reflexo; pellucido-corneā, epidermide tenui pubescente indutā, columellā labroque albida.

THE Pubescent BULIMUS. Shell oblong-ovate, umbilicated, whorls six to seven in number, somewhat rounded, very minutely reticulated beneath the lens, columella once-plaited, reflected, lip reflected; transparent horny, covered with a thin pubescent epidermis, lip and columella whitish.

MORICAND, Mém. 3rd Supp. p. 63. pl. 5. f. 21-23.

Hab. Bahia, Brazil.

Distinguished by its minutely reticulated surface, and pubescent epidermis.

Species 438. (Mus. Cuming.)

BULIMUS FRATERCULUS. *Bul. testā subacuminato-ovatā, umbilicatā, tenuiculā, axfrictibus sex, obliquè tenuissimè plicato-striatis, striis impressis spiralibus parallelis remotis decussatis, columellā reflexā, labro simplici; pellucido-corneā.*

THE ALLIED BULIMUS. Shell somewhat acuminately ovate, umbilicated, rather thin, whorls six in number, obliquely very finely plicately striated, decussated with remote parallel spiral impressed striae, columella reflected, lip simple; transparent horny.

POTIEZ and MICHAUD, Gal. de Douai, p. 141. pl. 13. f. 7, 8. *Cochlogena fraterculus*, Féussac.

Hab. Guadalupe.

The remote spiral striae which characterize this shell, are totally different in appearance from those which traverse it obliquely.

Species 439. (Mus. Cuming.)

BULIMUS ORBIGNI. *Bul. testā acuminato-oblongā, umbilicatā, anfractibus septem, concentricè creberrimè tenuiliratis, columellā subreflexā, labro simplici; albā.*

ORBIGNY'S BULIMUS. Shell acuminately oblong, whorls seven in number, concentrically very closely finely ridged, columella slightly reflected, lip simple; white.

PFEIFFER, Pro. Zool. Soc., 1846, p. 31.

Hab. Lima.

Very like the *B. scalariformis* in general aspect, but sufficiently distinct on comparison.

Bulinus, Pl. LXIV.



44.6



44.7



44.8



44.9



44.10



44.11



44.12



44.13



44.14



44.15

BULIMUS.

PLATE LXIV.

Species 440. (Mus. Cuming.)

BULIMUS SOCOTRENSIS. *Bul. testá globosá, crassiusculá, umbilicatá, anfractibus quinque ad sex, conspicue concentricè plicato-striatis, aperturá subrotundatá; albida, fusco peculiariter obliquè spiraliter undato-striatá, apertura fuscá fuscá.*

THE SOCOTRA BULIMUS. Shell globose, rather thick, umbilicated, whorls five to six in number, conspicuously concentrically plicately striated, aperture somewhat rounded; whitish, peculiarly obliquely spirally streaked in a waved manner with brown, interior of the aperture brown.

Hab. Island of Socotra.

About the size of a large pea, curiously painted with spirally oblique streaks.

Species 441. (Mus. Cuming.)

BULIMUS FERUSSACI. *Bul. testá oblongo-turrítá, umbilicatá, anfractibus octo, subrotundatá, levibus, columellá reflexá, aperturá puró; opaco-albá, nigripunctatá, anfractu ultimo fasciá nigra cingulata.*

FERUSSAC'S BULIMUS. Shell oblong-turreted, umbilicated, whorls eight in number, somewhat rounded, smooth, columella reflected, aperture small; opake-white, dotted with black, last whorl encircled with a black band.

DUNKER, Zeitschr. fur Malac. 1845, p. 164.

Hab. Loanda, West Africa.

Sprinkled with minute dots upon an opake-white ground, the last whorl being encircled with a conspicuous band.

Species 442. (Mus. Cuming.)

BULIMUS BAHIENSIS. *Bul. testá fusiformi-oblongá, umbilicatá, anfractibus octo, plano-convexis, levibus, columellá reflexá, uniplicatá, aperturá oblongá, labro expansá; pelmido-albá, opaco-albo bic illic peculiariter undatá.*

THE BAHIA BULIMUS. Shell fusiformly oblong, umbilicated, whorls eight in number, flatly convex, smooth, columella reflected, one-plaited, aperture oblong, lip expanded; transparent-white, peculiarly wavy here and there with opake-white.

MORICAND, Mém. de Genève, vol. vi. p. 541. pl. 1. f. 6.
Hab. Bahia, Brazil.

The opake-white streaks on the surface of this transparent shell, have somewhat the character of an epidermis.

Species 443. (Mus. Cuming.)

BULIMUS RETICULATUS. *Bul. testá obeso-cylindraceá, solidusculá, apice mucronato, anfractibus septem ad octo, plano-convexis, sub lete minuté grauso-reticulatis, suturis peculiariter sulcatis, columellá latè reflexá, aperturá parvoscíula, labro conspicue reflexo; albido-fuscescente, labro eburneo.*

THE RETICULATED BULIMUS. Shell stoutly cylindrical, rather solid, apex pointed, whorls seven to eight in number, flatly convex, minutely granously reticulated beneath the lens, sutures peculiarly grooved, columella broadly reflected, aperture rather small, lip conspicuously reflected; whitish-brown, lip ivory-white.

Hab. West Africa.

Distinguished by its delicate minutely grained surface.

Species 444. (Mus. Cuming.)

BULIMUS CYLINDRICUS. *Bul. testá pyramidalis-turrítá, minutè umbilicatá, anfractibus decem ad undecim, levibus, politis, columellá plano-reflexá; albida, fuscescente fasciatá, striis purpureo-nigris peculiariter reticulatis longitudinaliter obliquè pictá, opere nigrante.*

THE CYLINDRICAL BULIMUS. Shell pyramidalily turreted, minutely umbilicated, whorls ten to eleven in number, smooth, polished, columella flatly reflected; whitish, banded with light brown, longitudinally obliquely painted with peculiarly reticulated purple-black streaks, apex blackish.

GRAY, Annals of Philosophy, 1825, vol. ix. p. 414.

Bulimus articulatus, Turton.

Macroceramus signatus, Gmelin.

Bulimus signatus, Sowerby.

Hab. Island of Tortola, West Indies.

A singularly-painted porcelain-like shell, distinguished by its pyramidal growth.

Species 445. (Mus. Cuming.)

BULIMUS GULDINGII. *Bul. testá pyramidalis-turrítá,*

March, 1849.

minutè umbilicatā, anfractibus decem ad undecim, subrotundatis, levibus, columellā parum reflexā, aperturā parā, rotundatā, labro subexpanso; albā, maculis nigricante-fusciis reticulatō obliquē pictā, anfracta ultima fasciata, apice nigricante.

GULDING'S BULIMUS. Shell pyramidal turreted, minutely umbilicated, whorls ten to eleven in number, rather rounded, smooth, columnella but little reflected, aperture small, rounded, lip slightly expanded ; white, obliquely painted with blackish-brown reticulated patches, last whorl banded, apex blackish.

PFEIFFER, Symbolæ, vol. i. p. 82, and vol. ii. p. 115.
Hab. West Indies.

Distinguished from the preceding species, by the whorls being more rounded, and the shell less sharply pyramidal.

Species 446. (Mus. Cuming.)

BULIMUS FRAGOSUS. *Bul. testā obeso-cylindraceā, latè umbilicatā, apice nucronato, anfractibus septem, planocanexis, concentricè crebrinè elevato-striatis, suturis quasi sulcatis, columellā latè reflexā, obscurè uniplicatā, aperturā subrotundatā, labro expanso; sordide albā.*

THE ROUGH BULIMUS. Shell stoutly cylindreal, broadly umbilicated, apex painted, whorls seven in number, flatly convex, concentrically very closely elevatedly striated, sutures as though grooved, columnella broadly reflected, obscurely one-plaited, aperture nearly round, lip expanded ; dead white.

PFEIFFER, Symbolæ, vol. ii. p. 45.
Cochliogena fragosa, Ferussac.

Hab. Yemen, Arabia.

The margins of the aperture almost unite on the body whorl.

Species 447. (Mus. Cuming.)

BULIMUS CORONATUS. *Bul. testā oblongo-turritā, vix umbilicatā, anfractibus levibus vel obscurè rude striatis, tuberculis parvis compressis ad suturas crenulatis, columellā tenui, angustā; pellucido-brunneā.*

THE CORONATED BULIMUS. Shell oblong-turreted, scarcely umbilicated, whorls smooth, or obscurely

rudely striated, crenulated at the sutures with a row of small compressed tubercles, columnella thin, narrow ; transparent straw-colour.

PFEIFFER, Symbolæ, vol. iii. p. 83.
Hab. Brazil.

Each whorl is coronated at the sutures by a row of small compressed tubercles.

Species 448. (Mus. Cuming.)

BULIMUS FORMOSUS. *Bul. testā cylindraceo-oblongā, subumbilicatā, apice nucronato, anfractibus decem, rotundatis, concentricè subtiliter striatis, aperturā pareā, rotundatā, labro subexpanso; albida, nigricante-fusco transversim lineatā et obliquè remotè strigatā, apice nigricante.*

THE BEAUTIFUL BULIMUS. Shell cylindrically oblong, slightly umbilicated, apex pointed, whorls ten in number, rounded, concentrically finely striated, aperture small, rounded, lip a little expanded ; whitish, transversely lined and obliquely remotely streaked with blackish brown, apex blackish.

GRAY, Wood, Index Test. Supp. pl. 6. f. 24.
Hab. Island of St. Domingo, West Indies.

Closely allied to the pretty West Indian species above described, *B. cylindricus* and *Guildingii*.

Species 449. (Mus. Cuming.)

BULIMUS BOISSIERI. *Bul. testā globosā, inflatā, fragili, pellucidā, epidermide tenui cornē nitente indutā, spirae brevi, suturis impressis, anfractibus quatuor ad quinque, rotundatis, columellā tenui, simplici, aperturā subampitā; pellucido-brunneā.*

BOISSIER'S BULIMUS. Shell globose, inflated, fragile, transparent, covered with a thin shining horny epidermis, spire short, sutures impressed, whorls four to five in number, rounded, columnella thin, simple, aperture rather large ; transparent brown.

Hab. Bahia, Brazil.

A curious little *Physa*-like shell, of transparent horny substance, extremely delicate to the touch.



451



450



452



453



454



455



456



457



458



459

R. J. T. 1860.

200 x 40 mm.

[Increased one fourth.]

BULIMUS.

PLATE LXV.

Species 450. (Mus. Cuming.)

BULIMUS COSTATUS. *Bal. testá cylindraceo-turrítá, soli-disculátá, compressa umbilicatá, anfractibus octo, con-vezis, nitidé striato-costulatis, columellá plano-reflexá, plicato-dentalá, aperturá parvá, labro paululum ex-panso; cinerascente-carnéa, nitente, costulis albi-caulis, apertura fuscæ fasciæ.*

THE RIBBED BULIMUS. Shell cylindrically rounded, rather solid, compressly umbilicated, whorls eight in number, convex, neatly sculptured with fine striate ribs, columella flatly reflected, plicately toothed, aperture small, lip a little expanded; pale ashy flesh-colour, shining, ribs whitish, interior of the aperture brown.

PFEIFFER, Pro. Zool. Soc., 1848.

Hab. Brazil.

Very delicately ribbed after the manner of a *Pupa*.

Species 451. (Mus. Cuming.)

BULIMUS CONIFORMIS. *Bal. testá oculo-conicá, minutè umbilicatá, anfractibus quinque, levibns, ultimo ventri-co-so, medio obtusè angulato, columellá brevissimè reflexá, aperturá oblongá, labro simplici, acuto; albida, striis fasciis obliquis irregulariter notatá.*

THE CONE-SHAPED BULIMUS. Shell ovately conical, minutely umbilicated, whorls five in number, smooth, last whorl ventricose, obtusely angled in the middle, columella very shortly reflected, aperture oblong, lip simple, sharp; whitish, obliquely irregularly marked with brown streaks.

PFEIFFER, Pro. Zool. Soc., 1846. p. 114.

Hab. Andes of Columbia, near Merida.

The peculiar conical form of this shell gives it the appearance of immature growth.

Species 452. (Mus. Benson.)

BULIMUS PUNCTATUS. *Bal. testá oculo-conicá, subpro-funde umbilicatá, anfractibus sex, levibns, columellá latè reflexá, appressá, aperturá ovalá, labro simplici; ustulato-albó, zonulá angustá punctisque numerosis sparsis, nigricantibus, ornata, apice nigricante.*

THE DOTTED BULIMUS. Shell ovately conical, rather deeply umbilicated, whorls six in number, smooth,

columella broadly reflected, appressed, aperture ovate, lip simple; scorched white, ornamented with a narrow blackish spiral zone, and numerous scattered spots, apex blackish.

ANTON, Verz. p. 62.

Hab. Bundelkund and Southern India (on plants).
Benson, Jerdon.

The name given to this species in manuscript by Mr. Benson was *solutus*, signifying freckled, not *solutus* as printed by Dr. Pfeiffer. Manuscript names ought not, however, to be quoted as synomyms; a private name is scarcely open to public criticism.

Species 453. (Mus. Cuming.)

BULIMUS PEMPHIGODES. *Bal. testá globoso-conicá, ren-tricoso-inflatá, tenüssimá, minutè umbilicatá, spirá brevi, anfractibus quinque, convexis, obliquis subtilissimis striatis, columellá reflexá, appressá, aperturá ampliá, labro simplici, acuto; pallide cirescente-straminea, dia-planá, epidermide tenui opacá fasciatis indutá.*

THE BLADDER-LIKE BULIMUS. Shell globosely conical, ventricosely inflated, very thin, minutely umbilicated, spire short, whorls five in number, convex, obliquely very finely striated, columella reflected, appressed, aperture large, lip simple, sharp; pale greenish straw-colour, diaphanous, covered with a thin opaque epidermis.

JONAS, Zeitschr. für Malac. 1846. p. 12.

Hab. Guinea.

An extremely delicate inflated species named originally *B. vesiculosus* by Dr. Jonas in manuscript.

Species 454. (Mus. Cuming.)

BULIMUS IRREGULARIS. *Bal. testá ovato-oblongá, subam-pliter umbilicatá, anfractibus sex, ruguloso-striatis, suturis impressis, columellá latè reflexá, aperturá oblongá, labro simplici; carneo-fuscoceerulea.*

THE IRREGULAR BULIMUS. Shell ovately oblong, rather largely umbilicated, whorls six in number, rather roughly striated, sutures impressed, columella broadly reflected, aperture oblong, lip simple; light fleshy brown.

PFEIFFER, Pro. Zool. Soc., 1847. p. 231.

BULIMUS.—PLATE LXV.

Hab. Quito, Ecuador; Latre.

This species is more fully described by Dr. Pfeiffer, but its characters are very simple, approaching those of the Chilian group.

Species 455. (Mus. Cuming.)

BULIMUS DEALBATUS. *Bul. testā ovato-conicā, tenuiculā, umbilicatā, anfractibus sex, apicem versus subtilissimē striatis, columellā reflexā, aperturā ovatā, simplici; virescente-albā, subpellucidā, maculis punctisque opacocolorib; irregulariter striatā et aspersā.*

THE WHITE BULIMUS. Shell ovately conical, rather thin, umbilicated, whorls six in number, very finely striated towards the apex, columella reflected, aperture ovate, simple; pale greenish transparent white, irregularly streaked and sprinkled with opaque dots and blotches.

Say, Journ. Nat. Hist. Soc. Philadelphia, vol. ii. p. 159.

Hab. Alabama, North America.

The surface marking of this shell is not unlike that of the *B. conspersus*.

Species 456. (Mus. Cuming.)

BULIMUS YUNGASENSIS. *Bul. testā subfusiformi-oblongā, tenuiculā, viz umbilicatā, anfractibus sex, subrugosis, obscurè indentatis, columellā recedente, reflexā, planō-appressā; violaceente-albā, maculis violaceo-brunneis inaequilibre triseriatin cinctā.*

THE YUNGAS BULIMUS. Shell somewhat fusiformly oblong, rather thin, scarcely umbilicated, whorls six in number, rather rough, obscurely indented, columella receding, reflected, flatly appressed; violettinged white, encircled with three rows of irregular violet-brown spots.

D'ORBIGNY, Voy. dans l'Amér. Mérid. p. 316. pl. 40. f. 1.

Hab. Eastern side of the Andes of Bolivia, in the woods near the Meguilla river; D'Orbigny.

This species is unknown to me in good condition.

Species 457. (Mus. Cuming.)

BULIMUS DUSSUMIERI. *Bul. testā obeso-cylindracea, unu-*

bilicatā ad apicem subobtusa, anfractibus octo, angustis, obliquè striatis, columellā reflexā, aperturā subquadratā; pellucido-corneā.

DUSSUMIER'S BULIMUS. Shell stoutly cylindrical, unbilicated, rather obtuse at the apex, whorls eight in number, narrow, obliquely striated, columella reflected, aperture somewhat square; transparent horny.

Hab. India.

Mr. Cuming received this shell from the continent with the above name, but I cannot learn that it has been published.

Species 458. (Mus. Cuming.)

BULIMUS SAYI. *Bul. testā conico-oblongā, minutè umbilicatā, anfractibus sex, confertim rugoso-plicatis, columellā breviter reflexā, appressā; opaco-albā, strigis rufescente-fuscis pellucidis longitudinaliter notatā, circa umbilicum rufescente-fuscā.*

SAY'S BULIMUS. Shell conically oblong, minutely umbilicated, whorls six in number, closely rugosely pliated, columella shortly reflected, appressed; opaque white, longitudinally marked with transparent reddish-white streaks, reddish-brown around the umbilicus.

PFEIFFER, Pro. Zool. Soc., 1846, p. 114.

Hab. —?

The transparency of the brown stripes that distinguish this species, is very peculiar.

Species 459. (Mus. Cuming.)

BULIMUS CYCLOSTOMA. *Bul. testā conicā, profundè et subapertile umbilicatā, anfractibus rotundatis, concentricè striatis, columellā latè reflexā, aperturā circulari, labro simplici; caerulecente-albā.*

THE ROUND-MOUTH BULIMUS. Shell conical, deeply and rather largely umbilicated, whorls rounded, concentrically striated, columella broadly reflected, aperture circular, lip simple; bluish-white.

Hab. —?

Of a delicate white texture, deeply unablicated, and approaching the form of a *Cyclostoma*.



-f. 1



-d. 1



-e. 2



463



-f. 2



-g. 1



-g. 2



-h. 1



-i. 1



1 mm. long adult

1 1/8 in. imp.

B U L I M U S.

PLATE LXVI.

Species 460. (Mus. Cuming.)

BULIMUS UMBILICARIS. *Bul. testá pyramidalis-conicá,* profundissimè et amplissimè umbilicatá, anfractibus sex ad septem, compressissimis, concentricè striatis, suturis impressis, aperturá angustè ovali, labro subexpanso; pallidè corneo-cinereá.

THE UMBILICAL BULIMUS. Shell pyramidal and conical, very deeply and largely umbilicated, whorls six to seven in number, very compressed, concentrically striated, sutures impressed, aperture narrow-oval, lip slightly expanded; pale flesh-tinted ash.

SOULEYET, Revue Cuvierienne, 1842, p. 102.
Hab. Cobija, Bolivia.

This species is remarkable for its very capacious umbilicus, the whorls being so much compressed as to form a hollow cone.

Species 461. (Mus. Cuming.)

BULIMUS VARIATUS. *Bul. testá cylindraceo-oblongá,* crassiusculá, compressè umbilicatá, anfractibus septem, obliquè striatis, aperturá pareá, columellá labroque reflexis; brunnéa, strigis interseptis obliquè variegatá.

THE VARIED BULIMUS. Shell cylindrically oblong, rather thick, compressly umbilicated, whorls seven in number, obliquely striated, aperture small, lip and columella reflected; dark brown, obliquely variegated with yellowish streaks.

WEBB and BERTHELOT, Syn. p. 326.
Hab. Canary Islands.

There appears to be a slight yellowish epidermis.

Species 462. (Mus. Cuming.)

BULIMUS GOSSEI. *Bul. testá cylindraceo-turrítá,* compressè umbilicatá, apex attenuato, anfractibus undecim, angustis, subrotundatis, perobliquè costulato-striatis, suturis crenulatis, aperturá circulari, columellá labroque paulidam expansis; griseo-cinerascente, strigis semilunariibus albis, fusco-umbritis, punctisque fuscis, ornatá, fusco peltacido.

GOSSE'S BULIMUS. Shell cylindrically turreted, compressly umbilicated, apex attenuated, whorls eleven in number, narrow, somewhat rounded, very obliquely sculptured with fine rib-like striae, sutures

crenulated, aperture circular, lip and columella a little expanded; greyish-ash, ornamented with brown dots and white semilunar streaks shaded with brown, the brown being transparent.

PFEIFFER, Pro. Zool. Soc., 1843, p. 137.
Hab. Jamaica; Gosse.

The interesting new species of *Bulimus* and *Achatina* collected by Mr. Gosse in the Island of Jamaica, whilst engaged in ornithological pursuits, present an honourable testimony of his zeal in natural science, and furnish an example of what may be done by an active collector, with eyes to observe, and a mind to comprehend, the minute forms of animal life.

— — — — —
Species 463. (Mus. Cuming.)

BULIMUS KIENERI. *Bul. testá cylindraceo-turrítá,* compressè umbilicatá, apice attenuato, anfractibus tredecim, subrotundatis, perobliquè costulato-striatis, suturis crenulatis, aperturá circulari, columellá labroque subexpansis; fusco-cornéa, strigis semilunariibus albis, suturale umbritis, ornatá.

KIENER'S BULIMUS. Shell cylindrically turreted, compressly umbilicated, apex attenuated, whorls thirteen in number, somewhat rounded, very obliquely striated in a rib-like manner, sutures crenulated, aperture circular, lip and columella a little expanded; brownish-horny, ornamented with semilunar white spots shaded with darker brown.

PFEIFFER, Pro. Zool. Soc., 1846, p. 40.
Hab. Honduras.

It is somewhat doubtful whether this is anything more than a local variety of the *B. Gossei* of Jamaica.

Species 464. (Mus. Cuming.)

BULIMUS ZEBRIOLUS. *Bul. testá cylindraceo-elongatá,* Pupa-formá, latè umbilicatá, anfractibus decem, obliquè subobscure striatis, aperturá subquadratá, tridentata, marginibus subexpansis, collo conjunctis; cretuceo-alba, strigis angustis fuscis obliquè notata.

THE STRIPED BULIMUS. Shell cylindrically elongated, Pupa-shaped, broadly umbilicated, whorls ten in number, obliquely rather obscurely striated, aperture somewhat square, three-toothed, margins a little ex-

pounded, joined by a callosity; chalk-white, obliquely marked with narrow brown streaks.

Helix (Cochlogena) zebraiola, Féussac, Prodrome, No. 455.
Bulinus zebra, Olivier.

Var. immaculata. *Bulinus calcareus*, Parreyss.
Hab. Greece; Capt. Spratt.

Of an opake chalk-white obliquely marked with numerous streaks of light brown.

Species 465. (Mus. Cuming.)

BULIMUS RIDENS. *Bul. testa cylindraceo-elongata, crassiuscula, compressè umbilicata, aufractibus novem, planoverexis, obliquè obscurè striatis, columellâ reflexâ, conspicuo dentato-plicata, apertura parvâ; carlescentia alba, stigis angustis perpaucis fascis notata.*

THE DOUBLE-TOOTH BULIMUS. Shell cylindrically elongated, rather thick, compressly umbilicated, whorls nine in number, flatly convex, obliquely obscurely striated, columella reflected, conspicuously dentately plaited, aperture small; bluish-white, marked with a very few brown streaks.

KRYNICKI, Bull. Acad. Moscow, vol. vi. p. 401. pl. 1. f. 3.
Bulinus Clausiliiformis, Menke.

Bulinus appendiculatus, Potiez and Michaud.
Hab. Syria.

Potiez and Michaud regarded this species as the *Helix (Cochlogena) appendiculatus* of Dc Féussac, of which Dr. Pfeiffer is doubtful.

Species 466. (Mus. Cuming.)

BULIMUS CRETENSIS. *Bul. testa cylindraceo-oblonga, apice versus subattenuata, vix umbilicata, aufractibus septem, rugoso-striatis, columellâ subplanata, apertura quadrato-ovata, labro subexpanso, marginibus callo tenueo-conjunctis; rosaceo-tutea, suturis albimarginatis, labro albo.*

THE CRETEAN BULIMUS. Shell cylindrically oblong, rather attenuated towards the apex, scarcely umbilicated, whorls seven in number, delicately roughly striated, columella somewhat flattened, aperture squarely ovate, lip rather expanded, margins joined by a thin callosity; pale pinkish-yellow, sutures edged with white, lip white.

PFEIFFER, Philipp., Abbild. und besch. *Bulinus*, pl. 5. f. 8.
Hab. Island of Candia; Capt. Spratt.

A very delicate semitransparent shell, though of rather solid growth.

Species 467. (Mus. Cuming.)

BULIMUS ACUTUS. *Bul. testa pyramidalis-conica, minime umbilicata, aufractibus novem, rotundata, leviter corrugato-striatis, ad suturas rudi crenulatis, columellâ tenui, reflexâ, apertura circinari, labro simplici, acuto;*

pallide cornuta, maculis striisque opaco-lacteis marmorata, aufractu ultimo zonula nigra cingulata.

THE ACUTE BULIMUS. Shell pyramidalis conical, minutely umbilicated, whorls nine in number, rounded, slightly striated in a wrinkled manner, rudely crenulated at the sutures, columella thin, reflected, aperture circular, lip simple, acute; light horny, marbled with opake cream-coloured streaks and blotches, last whorl encircled by a narrow black zone.

BRUGUIERE, Enc. Méth., vol. i. p. 323.

Helix acuta, Müller,

Helix bifasciata, Pulteney.

Helix fasciata, Pennant.

Turbo turricula maroccaea, Chemnitz.

Hab. Spain, Portugal, West of England and Ireland.

The *B. articulatus* of Lamarek is probably a variety.

Species 468. (Mus. Cuming.)

BULIMUS UNICARINATUS. *Bul. testa cylindraceo-turrato, minute umbilicata, apice attenuata, aufractibus decem, obliquè subtilissimè striatis, ultimo basi acutè unicarinato, apertura parvâ, circinari, labro subreflexo; olivaceo-cornuta, stigis longitudinalibus numerosis opaco-albis ornata.*

THE ONE-KEELED BULIMUS. Shell cylindrically turreted minutely umbilicated, apex attenuated, whorls ten in number, obliquely very finely striated, last whorl encircled by a single sharp keel at the base, aperture small, circular, lip a little reflected; olive horny, ornamented with numerous opake white longitudinal streaks.

PFEIFFER, Monog. Helic. vol. ii. p. 80.

Popa unicarinata, Lamarek.

Bulinus Canimarensis, Pfeiffer.

Hab. Cuba, Guadaloupe.

Belongs to the same peculiar type as *B. Gossei*.

Species 469. (Mus. Cuming.)

BULIMUS BADIOSA. *Bul. testa ovato-conica, compressè umbilicata, aufractibus sex, subtilissimè granulatis, ultimo subventricoso, columellâ reflexâ, appressâ, apertura rotundato-ovali, labro subexpanso, marginibus callo subtiliter folio conjunctis, castaneo-brunneâ, labro albido.*

THE BROWN BULIMUS. Shell ovately conical, compressly umbilicated, whorls six in number, very finely granulated, last whorl rather ventricose, columella reflected, appressed, aperture rotundately oval, lip slightly expanded, margins joined by a somewhat tubercular callosity.

WEBB and BERTHELLOT, Syn. p. 318.

Helix (Cochlogena) badiosa, Féussac.

Hab. Island of St. Thomas, Féussac; Teneriffe, Webb and Berthelot.

Chiefly characterized by its delicate granulated sculpture.



470



471



472



473



474



475



476



477



478



479

100

[Increased one-half.]

B U L I M U S.

PLATE LXVII.

Species 470. (Mus. Cuming.)

BULIMUS ACHATINACEUS. *Bul. testā elongato-turritā, solidiusculā, anfractibus octo, cereberrimè rude striatis, columellā brevissimè reflexā, aperturā parvā, labro acuto; stramineū, cereā.*

THE ACHATINA-LIKE BULIMUS. Shell elongately turreted, rather solid, whorls eight in number, very closely rudely striated, columella very shortly reflected, aperture small, lip acute; straw-colour, waxy.

PFEIFFER, Symbol., vol. iii. p. 52.

Hab. Java.

Very closely allied to some of the small subulate species of *Achatina*.

Species 471. (Mus. Cuming.)

BULIMUS SUBTILIS. *Bul. testā cylindraceo-turritā, rix umbilicatā, anfractibus novem, obliquè subobscure rude striatis, columellā parvā reflexā, aperturā parvā; pelticido-corneā, suturis albi-marginatis.*

THE DELICATE BULIMUS. Shell cylindrically turreted, scarcely umbilicated, whorls nine in number, obliquely rather obscurely rudely striated, columella but little reflected, aperture small; transparent horny, sutures edged with white.

ROSSMASLER, vol. vi. p. 47. f. 392.

Bulimus noctivagus, Parreyss.

Hab. Dalmatia.

The white edge of the sutures is only faintly indicated, but it is a character not to be over-looked.

Species 472. (Mus. Cuming.)

BULIMUS SANDWICENSIS. *Bul. testā cylindraceo-turritā, umbilicatā, anfractibus decem, subplanatis, columellā latiusculā, parvā reflexā, aperturā parvā, labro acuto; rufo-brunneā, strigis undulatis opaco-albis notatā.*

THE SANDWICH BULIMUS. Shell cylindrically turreted, umbilicated, whorls ten in number, somewhat flattened, columella rather broad, but little reflected, aperture small, lip sharp; red-brown, marked with waved opaque-white streaks.

PFEIFFER, Pro. Zool. Soc., 1846, p. 31.

Hab. Sandwich Islands.

Of a bright dark red-brown colour, waved obliquely with delicate opaque-white streaks.

Species 473. (Mus. Cuming.)

BULIMUS MERDUENIANUS. *Bul. testā oblongo-conicā, umbilicatā, anfractibus octo, obscurè striatis, columellā latiusculā, parvā reflexā, aperturā rotundato-ovali, marginib[us] fere conjunctis, sordidè cornēa.*

THE MERDUENI BULIMUS. Shell oblong conical, umbilicated, whorls eight in number, obscurely striated, columella rather broad, but little reflected, aperture rotundately oval, margins almost joined; dull horny.

KRYSICKI, Bull. Acad. Sci. Moscow, 1837, p. 53.

Bulimus tener, Ziegler.

Bulimus fragilis, Parreyss.

Hab. Merdwen; Asia Minor.

Distinguished by its conical form.

Species 474. (Mus. Cuming.)

BULIMUS COSENSIUS *Bul. testā cylindraceo-oblongā, compressè umbilicatā, anfractibus septem, obliquè subrude striatis, columellā latā, concavo-appressā, aperturā subquadrato-ovatā, labro paulatim expanso; cernescente-albā, opacā.*

THE COS BULIMUS. Shell cylindrically oblong, compressly umbilicated, whorls seven in number, obliquely rather rudely striated, columella broad, concavely appressed, aperture rather squarely ovate, lip a little expanded; bluish-white, opaque.

Hab. Island of Cos, Grecian Archipelago; Capt. Spratt.

The lip is slightly expanded within, after the manner of *Partula*.

Species 475. (Mus. Cuming.)

BULIMUS MONTIVAGUS. *Bul. testā pyramidalī, compressè umbilicatā, anfractibus novem, obscurè tenuistriatis, columellā reflexā, aperturā partiuisculā, labro subcylindrico; pallide virescente-albā, pellucida, fuscescente hic illuc obscurè striatā.*

THE MOUNTAIN BULIMUS. Shell pyramidal, compressly umbilicated, whorls nine in number, obscurely finely striated, columella reflected, aperture rather small, lip

somewhat expanded; pale greenish-white, transparent, obscurely streaked here and there with light brown.

D'ORBIGNY, Moll. Voy. dans l'Amér. Mérid. p. 275. pl. 84. f. 1 to 3.

Hab. Parana, Argentine Republic, (under dead leaves); D'Orbigny.

Has sometimes the appearance of being marked with white streaks.

Species 476. (Mus. Benson.)

BULIMUS PULLUS. *Bul. testā elongato-cylindraceā, solidusculā, vix umbilicatā, anfractibus novem, subtilissimè plicato-striatis, infra suturas arcuatis crenulatis, aperturā parvā, circulari, columellā labro paululum expansis; fuscenscente-lacteā.*

THE BROWNISH BULIMUS. Shell elongately cylindrical, rather solid, scarcely umbilicated, whorls nine in number, very finely plicately striated, arcuately crenulated beneath the sutures, aperture small, circular, lip and columella a little expanded; brownish cream-colour.

GRAY, Pro. Zool. Soc., 1834, p. 66.

Pupa cylindrica, Hutton.

Hab. India. At Delhi (under stones); at Bundelkund (common among rocks and brushwood in the rainy season); Benson.

Mr. Benson collected a light variety of this species at Aden, Arabia, among loose stones on the skirts of the barren hills.

Species 477. (Mus. Cuming.)

BULIMUS LOEWI. *Bul. testā cylindraceo-turrītā, sinistrā, compressè umbilicatā, apice obtusè attenuato, anfractibus novem, angustis, planulatis, levibus, aperturā subquadrate-ovali, quadrivalvata; peltucido-corneā, dentibus labro opaco-albis, suturis albidae.*

LOEWE'S BULIMUS. Shell cylindrically turreted, sinistral, compressly umbilicated, apex obtusely attenuated, whorls nine in number, narrow, rather flattened,

smooth, aperture somewhat squarely oval, four-toothed; transparent horny, teeth and lip opake-white, sutures whitish.

PHILIPPI, Zeitschr. fur Malac. 1844. p. 106

Pupa Loewii, Kuster.

Hab. Lycia; Capt. Spratt.

This partakes much of the character of *Pupa*.

Species 478. (Mus. Benson.)

BULIMUS ARCUATUS. *Bul. testā oblongā, sinistrā, subventricosā, compressè umbilicatā, anfractibus septem, ruguloso-striatis, columellā latè appressā, labro subexpanso; suturā corneo-fusca, strigis lutescentibus angustis nitidè pictā, labro albido.*

THE ARCHED BULIMUS. Shell oblong, sinistral, somewhat ventricose, compressly umbilicated, whorls seven in number, finely roughly striated, columella broadly appressed, lip a little expanded; dark horny brown, neatly painted with narrow yellowish streaks, lip whitish.

PFEIFFER, Monog. Helic. viv. vol. ii. p. 118.

Bulimus ornatus, Hutton, MSS.

Hab. Mahassoo, Western Himalaya; Hutton.

I learn on the authority of Mr. Benson, that *ornatus* was the name originally given to this species by Capt. Hutton in manuscript, not *arcuatus* for which Dr. Pfeiffer mistook it. The error proves, however, to be a fortunate one, Hutton's name being already appropriated by Dufo.

Species 479. (Mus. Cuming.)

BULIMUS OLIVACEUS. *Bul. testā oblongā, compressè umbilicatā, anfractibus sex, obliquè striatulis, columellā appressā, labro paululum reflexo; corneo-oliveā, sericeinā, labro albo.*

THE OLIVE BULIMUS. Shell oblong, compressly umbilicated, whorls six in number, obliquely finely striated, columella appressed, lip a little reflected; horny olive, silkcn, lip white.

PFEIFFER, Abbild. und Besch. *Bulimus*, Pl. 5. f. 6.

Hab. Island of Candia; Capt. Spratt.

Of very simple form, with little or no sculpture.

Bulimus. Pl. LXVIII.



480



481




482



483



484



485



486



487



488



489

BULIMUS.

PLATE LXVIII.

Species 480. (Mus. Cuming.)

BULIMUS ORYZA. *Bul. testā cylindraceo-turritā, minūtē umbilicatā, anfractibus octo, rotundatis, longitudinaliter striatis, spira suturis profundis, columellā subreflexā, aperturā minūmā, suborbiculari, labro simplici; vires-cente-albā, vitrēa.*

THE RICE-GRAIN BULIMUS. Shell cylindrically turreted, minutely umbilicated, whorls eight in number, rounded, longitudinally striated, sutures of the spire deep, columella slightly reflected, aperture very small, nearly orbicular, lip simple; greenish white, glassy.
Hab. Bruguier, Encyc. Mél. vers, vol. i. p. 332.

Hab. Brazil.

The shell here figured agrees with the description of this species given by Deshayes (Lamarc, Anima. sans vert., vol. viii. p. 263), but not with Dr. Pfeiffer's diagnosis, which describes a shell of cancellated sculpture, composed of five whorls only.

Species 481. (Mus. Cuming.)

BULIMUS TUCKERI. *Bul. testā subulato-turritā, minūtē umbilicatā, anfractibus novem, convexiusculis, longitudinaliter striatis, spira suturis impressis, columellā subobliquè recedente, tenuiter reflexā, aperturā minūtā, labro simplici; pallidè cornēa, pellucida.*

TUCKER'S BULIMUS. Shell subulately turreted, minutely umbilicated, whorls nine in number, rather convex, longitudinally striated, sutures of the spire impressed, columella somewhat obliquely receding, finely reflected, aperture minute, lip simple; pale horn, transparent.

PFEIFFER, Pro. Zool. Soc., 1846, p. 30.

Hab. Hardy's Island, Pacific Ocean; Tucker.

Chiefly distinguished by its subulate form.

Species 482. (Mus. Cuming.)

BULIMUS BACTERIONIDES. *Bul. testā elongato-turritā, cylindraceō, ad apicem obtusā, anfractibus octo ad novem, subplano-convexis, levibus, nitidis, suturis impressis, columellā rectā, interdum leviter truncatā, aperturā parvā; lutescente-cornēa, pellucidā.*

THE STAFF-LIKE BULIMUS. Shell elongately turreted, cylindrical, obtuse at the apex, whorls eight to nine

in number, rather flatly convex, smooth, shining, sutures impressed, columella straight, sometimes slightly truncated, aperture small; yellowish-horn, transparent.

D'ORBIGNY, Voyage dans l'Amér. Mérid. Moll., p. 260. pl. 29. f. 1 to 3.

Hab. Bolivia; D'Orbigny.

As an example of the little importance of the truncature of the columella as a generic character in the small *Achatinae*, it may be remarked that in this species the columella is sometimes truncated and sometimes not.

Species 483. (Mus. Cuming.)

BULIMUS IMPRESSUS. *Bul. testā subulatā, anfractibus octo, plano-convexis, politis, nitidis, lineis impressis obscuris distantibus longitudinaliter notatis, columellā tenuiter reflexā, subobliquè recedente, aperturā parvā, oblongā; pellucido-cornēa, solidinsculpta.*

THE IMPRESSED BULIMUS. Shell subulate, whorls eight in number, flatly convex, polished, shining, longitudinally marked with distant obscure impressed lines, columella thinly reflected, obliquely receding, aperture small, oblong; transparent horny, rather solid.

Hab. —?

A rather stout polished horny shell, marked here and there with faintly impressed lines, distinct from *B. subula*, in which the whorls are more rounded.

Species 484. (Mus. Cuming.)

BULIMUS MIMOSARUM. *Bul. testā cylindraceo-turritā, minūtē umbilicatā, apice acutā, anfractibus decem, angustis, subrotundatis, levibus, nitidis, suturis impressis, columellā tenui, latè reflexā, appressā, aperturā parvā, labro simplici; sordide albā.*

THE MIMOSA-INHABITING BULIMUS. Shell cylindrically turreted, minutely umbilicated, apex sharp, whorls ten in number, narrow, somewhat rounded, smooth, or finely striated, columella thin, broadly reflected, appressed, aperture small, lip simple; dull white.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll., p. 262. pl. 41. f. 12-14.

Hab. Bolivia; D'Orbigny.

Found principally at the roots of thorny *Mimosa* trees.

BULIMUS.—PLATE LXVIII.

Species 485. (Mus. Cuming.)

BULIMUS SPOLIATUS. *Bul. testā cylindraceo-elongatā, minutè umbilicatā, anfractibus octo, planulatis, punctis obscurè indentatis, columellā subexpansā, lirā denti-formū aperturam intrante superne innatā, aperturā subobliquā, labro medio obsoletè undentata; fuscescente-lacēta, fuso supra suturas spiraliter punctatā.*

THE SPOILED BULIMUS. Shell cylindrically elongated, minutely umbilicated, whorls eight in number, flattened, obscurely indented with punctures, columella rather expanded, furnished above with a tooth-like ridge entering the aperture, aperture rather oblique, lip with an obsolete tooth about the middle; brownish cream-colour, spirally dotted above the sutures with brown.

PARREYSS, Pfeiffer, Symbolæ, vol. iii. p. 87.
Hab. Greece.

Dr. Pfeiffer quotes the *Pupa fusiformis* of Deshayes as a synonymy of this species, but it does not agree with the description.

Species 486. (Mus. Cuming.)

BULIMUS SUBLIFORMIS. *Bul. testā acuminato-elongatā, gracili, apice subpapillari, anfractibus ad quatuordecim, planulatis, obliquè subtilissimè striatis, politis, nitidis, anfractu ultimo ad basin angulato, columellā tenui, aperturā minima; tricemente-cornēa, pellucida.*

THE AWL-SHAPED BULIMUS. Shell acuminate elongated, slender, apex somewhat papillary, whorls about fourteen in number, flattened, obliquely very finely striated, polished, shining, last whorl angulated at the base, columella thin, aperture very small; yellowish-horny, transparent.

Helix subuliformis, Moricand, Mém. Genève, vol. vii. pt. 2. p. 127. pl. 2. f. 3.
Hab. Bahia, Brazil.

A delicate, bright yellow, horny species, distinguished by its prolonged spire of flattened whorls.

Species 487. (Mus. Cuming.)

BULIMUS PELLUCIDUS. *Bul. testā turritā, minutè umbilicatā, tenuissimā, anfractibus septem, rotundatis, longitudinaliter crebristriatis, columellā tenui, reflexā, aperturā parvā, labro simplici; virescente-corneā, pellucida.*

THE TRANSPARENT BULIMUS. Shell turreted, minutely umbilicated, very thin, whorls seven in number, rounded, longitudinally closely striated, columella thin, reflected, aperture small, lip simple; greenish-horny, transparent.

PFEIFFER, Pro. Zool. Soc., 1847.

Hab. Province of Merida, New Granada; Funck.

An exceedingly delicate shell, having the appearance of light green glass.

Species 488. (Mus. Cuming.)

BULIMUS DECEPTUS. *Bul. testā sulfusiformi-conicā, anfractibus septem, lœvibus, nitidis, e suturis obscurè concentricè striatis, columellā tortuosa, recedente, ad marginem collosā, aperturā oblongā, labro simplici; virescente-vitrea, pellucida.*

THE MISTAKEN BULIMUS. Shell somewhat fusiformly conical, whorls seven in number, smooth, shining, obscurely concentrically striated from the sutures, columella tortuous, receding, callous at the edge, aperture oblong, lip simple; greenish-glassy, transparent.

Bulimus interstinctus, Pfeiffer (not of Gould) Monog. Helic. p. 169.
Hab. —?

Dr. Pfeiffer's description of this species is inadvertently referred to at Pl. LV. Sp. 367, owing to his having taken it to be the *B. interstinctus* of Gould; arising probably from the displacement of a label in Mr. Cuming's cabinet. Dr. Pfeiffer does not appear to have seen Dr. Gould's description of that species, and I have been equally unsuccessful myself in finding it.

The shell described and figured at Pl. LV. Sp. 367, was received from Dr. Gould by Mr. Cuming with the name *B. interstinctus* attached to it, and he has also received specimens from Mr. Redfield of the United States similarly named.

Species 489. (Mus. Cuming.)

BULIMUS BERGERI. *Bul. testā cylindraceo-turritā, minutè umbilicatā, anfractibus novem, planicostatis, obliquè striatis, columellā latiusculā, aperturā parvā, tridentata; fuscescente-cornēa, dentibus albidis.*

BERGER'S BULIMUS. Shell cylindrically turreted, minutely umbilicated, whorls nine in number, flatly convex, obliquely striated, columella rather broad, aperture small, three-toothed; brownish horn-colour, teeth white.

PFEIFFER, Symbolæ, vol. iii. p. 52.

Pupa Bergeri, Roth.

Pupa tridentata, Anton.

Pupa tricuspis, Rossmaëslér.

Bulimus tridentatus, Parryss.

Hab. Greece, Egypt, &c.

I follow Dr. Pfeiffer in referring this species to *Bulimus*, but it seems really a matter of indifference whether it is included in this genus or in *Pupa*.




s.9c



s.9d



s.9e



s.9f



s.9g



s.9h



s.9i



s.9j



s.9k



s.9l

BULIMUS.

PLATE LXIX.

Species 490. (Mus. Cuming.)

BULIMUS HOLOSTOMA. *Bal. testā turritā, profundē umbilicatā, anfractibus septem, subquadralis, corrugato-pli-catis, peculiariter laevē convolutis, suturis profundis, aperturā integrā, tubulari; carneo-ciuerē.*

THE ENTIRE-MOUTH BULIMUS. Shell turreted, deeply umbilicated, whorls seven in number, somewhat square, plicated in a wrinkled manner, peculiarly loosely convoluted, sutures deep, aperture entire, tubular; fleshy ash-colour.

PFEIFFER, Pro. Zool. Soc., 1846. p. 28.

Hab. Cobija, Bolivia.

An extremely interesting species in which the shell, forming a loosely convoluted spiral tube, has no axial pillar.

Species 491. (Mus. Cuming.)

BULIMUS RUPESTRIS. *Bal. testā turritā, misutē umbili-catā, anfractibus quinque ad sex, rotundatis, obliquē rosulato-striatis, aperturā ovali, ferè integrā; fusco-corneā.*

THE ROCKY BULIMUS. Shell turreted, minutely umbilicated, whorls five to six in number, rounded, obliquely sculptured with fine rib-like striae, aperture oval, almost entire; brownish-horny.

PHILIPPI, Ennus. Moll. Sicil. p. 141. pl. 8. f. 18.
Pupa rupestris, Rossmaister.

Hab. Siéli; Philippi; Algeria; Tervér.

A delicate *Scalaria*-like species in which the aperture is almost entire.

Species 492. (Mus. Cuming.)

BULIMUS CENOPICTUS. *Bal. testā oblongā, compressē umbilicatā, anfractibus quinque, subrotundatis, levibus, columellā vix reflexā, aperturā parvā; peltucido-corneā.*

THE DIRTY-PAINTED BULIMUS. Shell oblong, compressly umbilicated, whorls five in number, somewhat rounded, smooth, columella scarcely reflected, aperture small; transparent horny.

Pupa cenopicta, Hutton, Journ. Asiatic Soc., vol. iii. p. 85.
Hab. India.

Dr. Pfeiffer remarks that this may be a variety of *B. folliculus*, but that is unlikely.

Species 493. (Mus. Cuming.)

BULIMUS SUBDIAPHANUS. *Bal. testā cylindraceo-turritā, vix umbilicatā, anfractibus septem, convexis, subtilis-simē obliquē striatis, columellā subreflexā, appressā, aperturā minima, latiusculā; eburnea, subdiaphana, solidiuscula.*

THE SEMITRSPARENT BULIMUS. Shell cylindrically turreted, scarcely umbilicated, whorls seven in number, convex, very finely obliquely striated, columella slightly reflected, appressed, aperture very small, rather broad; ivory-white, semitransparent, rather solid.

PFEIFFER, Symbolae, vol. ii. p. 122.

Pupa subdiaphana.

Bulimus Barbacocha, Webb and Berthelot.

Hab. Cape De Verd Islands.

A long *Pupa*-shaped semitransparent ivory-white shell.

Species 494. (Mus. Cuming.)

BULIMUS SUBULA. *Bal. testā subulato-turritā, spira-suturis impressis, anfractibus octo, longitudinaliter striatis, columellā rectā, brevissime reflexā, aperturā suboblongā, labro simplici; pellucido-corneā, tenuissimā.*

THE AWL BULIMUS. Shell subulately turreted, sutures of the spire impressed, whorls eight in number, longitudinally striated, columella straight, very shortly reflected, aperture rather oblong, lip simple; transparent horny, very thin.

PFEIFFER, Symbolae, vol. i. p. 85.

Bulimus octonoides, D'Orbiguy.

Bulimus procerus, Adams.

Hab. Cuba.

A perfectly transparent shell, of a more acuminate subulate growth than the *B. octona*, for which it might be mistaken.

Species 495. (Mus. Benson.)

BULIMUS GRACILIS. *Bal. testā gracili-subulata, vix um-bilicata, suturis impressis, anfractibus decem ad undeci, creberrimē tenuissimē concentricē striatis, columellā tenui, breviter reflexā, aperturā supernā leviter con-tractā, labro simplici; sordidē fuscescere-corneā.*

THE SLENDER BULIMUS. Shell slenderly subulate,

BULIMUS.—PLATE LXIX.

scarcely umbilicated, sutures impressed, whorls ten to eleven in number, very closely and finely concentrically striated, columnella thin, shortly reflected, aperture slightly contracted at the upper part, lip simple; dull brownish horn colour.

HUTTON, Journ. Asiatic Soc., 1834, vol. iii. p. 84.

An eadem, *Bulimus Indicus*, Pfeiffer.

Hab. Bundelkund, and all the plain Provinces of the Bengal Presidency; Benson. Point de Galle, Ceylon; Benson.

Dr. Pfeiffer's *B. Indicus* appears to be merely an immature state of this species.

Species 496. (Mus. Benson.)

BULIMUS NIVICOLA. *Bul. testā cylindraceo-ovatā, obesā, ventricosā, sinistrā, compressa umbilicatā, anfractibus septem, concexis, obliquè striatis, ad suturas subobscure crenulatis, columellā latiusculā, appressā, apertura parvā, labo leviter expanso; rufo-fuscā, strigis albīs obliquis hic illīs irregulariter interruptis notatā, epidermide tenui lutescente intitulā.*

THE SNOW-INHABITING *BULIMUS*. Shell cylindrically ovate, stout, ventricose, sinistral, compressly umbilicated, whorls seven in number, convex, obliquely striated, somewhat obscurely crenulated at the sutures, columnella rather broad, appressed, aperture small, lip slightly expanded; red-brown, marked here and there with irregularly interrupted oblique white streaks, covered with a thin yellowish epidermis.

BENSON, MSS.

Pupa sylvestris, Hutton (fide Benson.)

Hab. Liti Pass, Western Himalaya (among juniper bushes and patches of snow at nearly 14,000 feet above the level of the sea); Capt. Hutton.

Mr. Benson is of opinion that this species, distinguished by its stout cylindrical form and reversed growth, which has been upwards of twelve years in his collection, is the *Pupa sylvestris* of Hutton, described in McClelland's Journal for January, 1841, as being found among juniper bushes at Burrenda Pass, and of which the name is pre-occupied in the present genus to which this species strictly belongs. It is remarkable for its stout cylindrical form and reversed growth; it should, however, be remembered when referring to the plate, that the figure is very considerably magnified.

Species 497. (Mus. Cuming.)

BULIMUS TURRICULA. *Bul. testā cylindraceo-turritatē, minutè umbilicatā, prope apicem peculiariter attenuatā, anfractibus novem, rotundatis, perobliquè creberrimè costulato-striatis, columellā latè appressā, apertura ferè circulari; fuscescente, strigis arcuatis albīs fuso-umbrales hic illīs marmoratae.*

THE LITTLE TURRET *BULIMUS*. Shell cylindrically turrited, minutely umbilicated, peculiarly attenuated near the apex, whorls nine in number, rounded, very obliquely closely striated in a fine rib-like manner, columnella broadly appressed, aperture nearly circular; light brown, marbled here and there with arched brown-shaded white streaks.

PFEIFFER, Wiegman, Archiv., 1839, vol. i. p. 351.

Hab. Cuba.

An interesting little species allied in form to the *B. Gossei*.

Fig. 498. (Mus. Cuming.)

This little triangular shell, of which the figure is very highly magnified, proves to be the extreme young of *B. calebs* (*B. solitarius*, Hutton) Pl. XLVII. Sp. 301, from Western Himalaya.

For Sp. 498. see Pl. LXX.

Species 499. (Mus. Cuming.)

BULIMUS VENTROSUS. *Bul. testā subpyramidaliter-conicā, minutè et profundiè umbilicatā, anfractibus septem, planulato-conexis, peculiariter obscurè plicato-corrugatis, columellā reflexā, apertura ferè circulari, labo simplici; abiebat, basi nigricante unifasciota.*

THE VENTRICOSE *BULIMUS*. Shell somewhat pyramidal conical, minutely and deeply umbilicated, whorls seven in number, rather flatly convex, peculiarly obscurely plicately wrinkled, columnella reflected, aperture nearly circular, lip simple; whitish, encircled with a narrow black band at the base.

Helix ventrosa, Férussac, Prodrome, 377.

Bulimus ventricosus, Draparnaud.

Bulimus variabilis, Hartman.

Helix acuta, Webb and Berthelot.

The narrow black band which encircles the base of this shell is not an unimportant character.



503.



504.



505.



506.



507.



508.



509.



510.



511.



512.

BULIMUS.

PLATE LXX.

Species 498. (Mus. Pfeiffer.)

BULIMUS TRUNCATUS. *Bul. testā attenuato-pyramidalī, basi angulatā, ad apicem decollatā, anfractibus octo, angustis, convexis, obliquē concentricē creberrimā elevato-striatis, columellā rectā, tenuiter expansā, aperturā sub-quadratā, labro tenui; virescuto-cornēā.*

THE TRUNCATE BULIMUS. Shell attenuately pyramidal, angled at the base, decollated at the apex, whorls eight in number, narrow, convex, obliquely concentrically very closely elevated striated, columella straight, thinly expanded, aperture somewhat square, lip thin; greenish horny.

PFEIFFER, *Symbola*, vol. i. p. 43.

Hab. Mexico.

This remarkable species, for the loan of which I am indebted to Dr. Pfeiffer, does not appear to have reached maturity. It is not known to English collectors.

Species 500. (Pl. *Achatina*, XVI. Fig. 77,
Mus. Cuming.)

BULIMUS CLAVA. *Bul. testā subulatā, apice papillari, haud umbilicata, anfractibus decem, levibus, politis, hic illū longitudinaliter impresso-striatis, suturis marginatis, columellā tenui, parua reflexā, aperturā parviusculā, labro simplici; pellucido-cornēā.*

THE CLUB BULIMUS. Shell subulate, papillary at the apex, not umbilicated, whorls ten in number, smooth, polished, here and there longitudinally impressly striated, sutures margined, columella thin, but little reflected, aperture rather small, lip simple; transparent horny.

Hab. —?

The columella of this species is extremely thinly reflected, and the outer lip is rather inclined to be sinuated at the upper part.

Species 501. (Pl. *Achatina*, XVII. Fig. 81.
Mus. Cuming.)

BULIMUS CEREUS. *Bul. testā subulatā, umbilicata, tenui, anfractibus decem, convexis, arcuatum concentricē striatis, columellā reflexā, aperturā pareā, labro simplici; pellucido-stramineā.*

THE WAXEN BULIMUS. Shell subulate, umhilicated, thin,

whorls ten in number, convex, areately concentrically striated, columella reflected, aperture small, lip simple; transparent straw-colour.

Hab. Moradabad, India.

I venture to distinguish this from the *B. gracilis*, from which, in comparison with authentic specimens collected by Mr. Renson, it varies in form, the whorls being of a more bubble-like inflated growth.

Species 502. (Pl. *Achatina*, XVIII, Fig. 95.
Mus. Cuming.)

BULIMUS COLUMELLA. *Bul. testā subulatā, haud umbilicata, apice papillari, anfractibus undecim ad duodecim, minutissimē obliquē striatis, ad suturas subtler crevulatā, columellā vix reflexā, aperturā pareā, labro simplici; pellucido-cornēā, nitente.*

THE LITTLE COLUMN BULIMUS. Shell subulate, not umbilicated, papillary at the apex, whorls eleven to twelve in number, very minutely obliquely striated, finely crenulated at the sutures, columella but little reflected, aperture small, lip simple; transparent horny, shining.

PHILIPPI, Abbild. und Besch. Bul. pl. 2. f. 7.
Hab. Brazil.

This approaches very closely to the *B. sylvaticus*. It is a transparent turreted shell, very finely striated and delicately puckered at the sutures.

Species 503. (Pl. *Achatina*, XX. Fig. 113. Mus. Cuming.)

BULIMUS JAMAICENSIS. *Bul. testā subulatā, angustā, haud umbilicata, anfractibus octo, planice convexis, hic illū impresso-striatis, columellā tenuiter reflexā, aperturā minuta, labro simplici; pellucido-cornēā.*

THE JAMAICA BULIMUS. Shell subulate, narrow, not umbilicated, whorls eight in number, flatly convex, here and there impressly striated, columella thinly reflected, aperture very small, lip simple; transparent horny.

Hab. Jamaica.

An extremely bright transparent shell in which the columellar axis of the whorls is visible throughout.

Species 504. (Mus. Pfeiffer.)

BULIMUS OBESATUS. *Bul. testa cylindraceo-oblonga, subobesa, compressa umbilicata, anfractibus septem, (Quaque corrugato-striatis, columellâ labroque subincrustatis, late expansis, apertura suborbiculari; albida, epidermide olivaceo-cornuta, nitente, inducta).*

THE STOUT BULIMUS. Shell cylindrically oblong, rather stout, compressly umbilicated, whorls seven in number, obliquely corrugately striated, lip and columella somewhat thickened, broadly expanded, aperture nearly orbicular; whitish, covered with a shining olive-horny epidermis.

WEBB and BERTHELOT, Syn. p. 318.

Helix obesata, Féussac.

Hab. Canary Islands.

Of a cylindrical Pupiform shape like the Syrian and Lycian species, but covered with a dark shining olive epidermis.

Species 505. (Mus. Pfeiffer.)

BULIMUS EBTICATUS. *Bul. testa ovato-conica, tenuicula, compressa umbilicata, anfractibus seplex, convexis, peculiariter corrugato-granulatis, apertura suborbiculari, columellâ labroque expansa; sordida castanea.*

THE TAWNY BULIMUS. Shell ovately conical, rather thin, compressly umbilicated, whorls seven in number, convex, peculiarly corrugately granulated, aperture nearly orbicular, lip and columella expanded; dull chestnut.

WEBB and BERTHELOT, Syn. p. 318.

Helix (Coccygea) beticata, Féussac.

Bulimus cornutus, Menke.

Hab. Island of St. Thomas, West Indies; Féussac. Canary Islands; Webb.

Curiously granulated throughout, whilst the apex rises with a papillary appearance.

Species 506. (Mus. Pfeiffer.)

BULIMUS ZIEBMANNI. *Bul. testa fusiformi-oblonga, compressa umbilicata, anfractibus seplex, levibus, columellâ reflexa, apertura parviuscula, labro effuso, vix reflexa; albâ, fasciis carneeo-nigricantibus brevibus, multo interrupitis, cingulata.*

ZIEBMAN'S BULIMUS. Shell fusiformly oblong, compressly umbilicated, whorls seven in number, smooth, columella reflected, aperture rather small, lip effused, scarcely reflected; white, encircled with short blue-black bands, much interrupted.

PFEIFFER, MSS.

Hab. —?

Dr. Pfeiffer has favoured me with this shell for publication, and I am not aware that he has described it.

Species 507. (Mus. Pfeiffer.)

BULIMUS PHYSOIDES. *Bul. testa ventricoso-ovata, tenui, inflata, vix umbilicata, anfractibus quinque, longitudo later rugoso-striatis, ad suturas plicato-crenulatis, columellâ reflexâ, labro simplici; flaccidæ, albivirgata.*

THE PHYSA-LIKE BULIMUS. Shell ventricosely ovate, thin, inflated, scarcely umbilicated, whorls five in number, longitudinally roughly striated, plicately crenulated at the sutures, columella reflected, lip simple; yellowish, mottled with white.

MENKE, MSS.

Hab. —?

Received from Dr. Pfeiffer with the above name. I fear it is an immature shell.

Species 508. (Mus. Pfeiffer.)

BULIMUS HEGEWISCHI. *Bul. testa subacutuato-ovata, tenuicula, vix umbilicata, anfractibus quinque ad sex, irregulariter impresso-striatis, columellâ reflexâ, labro simplici; carneo-albicante, fuscâ hic illuc sparsim striata, intus pallide aurantiâ.*

HEGEWISCH'S BULIMUS. Shell somewhat acuminately ovate, rather thin, scarcely umbilicated, whorls five to six in number, irregularly impressly striated, columella reflected, lip simple; flesh-white, sparingly streaked here and there with brown, pale orange within.

PFEIFFER, MSS.

Hab. —?

For this species I am also indebted to Dr. Pfeiffer.

Species 509. (Mus. Pfeiffer.)

BULIMUS ZOOGRAPHICUS. *Bul. testa subfusiformi-oblonga, tenui, vix umbilicata, anfractibus sex, obsoletè striatis et indentatis, columellâ reflexâ, apertura subeffusa, labro simplici; sordida albâ, carneo-nigricante maculata et variegata.*

THE ZOOGRAPHIC BULIMUS. Shell somewhat fusiformly oblong, thin, scarcely umbilicated, whorls six in number, obsoletely striated and indented, columella reflected, aperture somewhat effused, lip simple; dull white, blotched and variegated with ashy black.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 313. pl. 40, f. 6 to 8.

Hab. Eastern Cordilleras of Bolivia (in a ravine between the Provinces of Mojos and Cochabamba); D'Orbigny.

This appears to be a species of extremely rare occurrence. The specimen here represented from Dr. Pfeiffer's cabinet is one of those collected by D'Orbigny.

Species 510. (Mus. Pfeiffer.)

BULIMUS LYMNOIDES. *Bul. testā subfusiformi-ovalā, solidiscutā, vix umbilicatā, aufractibus sex, subdilatissimē rugoso-striatis, columellā reflexā, aperturā subeffusā; olivaceo-fusca, nitente, saturis albinaarginatis.*

THE LYMNEA-LIKE BULIMUS. Shell somewhat fusiformly ovate, rather solid, scarcely umbilicated, whorls six in number, very finely roughly striated, columella reflected, aperture slightly effused; olive-brown, shining, sutures edged with white.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. viii. p. 260.
Hab. Guadeloupe, St. Vincent.

Externally, this shell has a very delicate satiny appearance.

Species 511. (Mus. Pfeiffer.)

BULIMUS TURRITELLA. *Bul. testā oblongo-ovalā, tenui, leviter ventricosā, umbilicatā, aufractibus sex, rugoso-striatis, columellā latē reflexā, labro simplici; pellicula-albā, lineis opaco-albis profnō longitudinaliter rotatā.*

THE LITTLE TOWER BULIMUS. Shell oblong-ovate, thin, slightly ventricose, umbilicated, whorls six in number, roughly striated, columella broadly reflected, lip simple; transparent white, profusely longitudinally marked with opaque white lines.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 274. pl. 33. f. 12 to 14.

Hab. Province of Chiquitos, Bolivia (in a small plain near the ancient Mission of St. Juan); D'Orbigny.

This species is also illustrated from one of M. D'Orbigny's specimens in the collection of Dr. Pfeiffer.

Species 512. (Mus. Pfeiffer.)

BULIMUS INFLATUS. *Bul. testā subgloboso-oculatā, solidiscutā, umbilicatā, aufractibus quinque, rotundatis, rugoso-striatis, columellā subarcuata, latē reflexā, aperturā ovali, labro simplici; sordidè albā.*

THE INFLATED BULIMUS. Shell somewhat globosely ovate, rather solid, umbilicated, whorls five in number, rounded, roughly striated, columella slightly arched, broadly reflected, aperture oval, lip simple; dead-white.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. viii. p. 230.

Helix (Cochlogena) costulata, Féussac.
Bulimus costulatus, Potiez and Michaud.

Hab. New Holland; Lamarck.

I follow Dr. Pfeiffer in adopting Lamarck's name for this species, in consequence of De Féussac's being inappropriate.



514.



513.



515.



207. b



516. a.



516. b



517.

BULIMUS.

PLATE LXXI.

Species 513. (Mus. Cuming.)

BULIMUS SCITULUS. *Bul. testā sulciformalē-oblongā, vix umbilicatā, anfractibus octo, leviter convexis, lacūbus, columellā parum reflexā, aperturā subangustā, labro simplici; albida, purpureo-ceruleo tinctā, basin versus ferrugineo-rufā, lineis subtilibus albis, irregulariter undulatis, cerebrimē longitudinaliter notatā.*

THE TRIN. BULIMUS. Shell somewhat fusiformly oblong, scarcely umbilicated, whorls eight in number, slightly convex, smooth, columella but little reflected, aperture rather narrow, lip simple; whitish, stained with purple-blue, rust-red towards the base, very closely longitudinally marked with fine, irregularly waved, white lines.

Hab. Chachapoyas, Alto-Peru.

Neatly marked with fine white waved lines upon a purple-blue ground, tinged towards the base with a bright rust-red.

Species 514. (Mus. Cuming.)

BULIMUS CUZCOENSIS. *Bul. testā acuminato-oblongā, subcylindracea, subcompressē umbilicata, anfractibus octo, leviter concavis, sub lute stratis et corrugato-indentatis, columellā reflexā, aperturā parenscaida, labro simplici; fulvescente-spodicēa.*

THE CUZCO BULIMUS. Shell acuminate oblong, somewhat cylindrical, rather compressly umbilicated, whorls eight in number, slightly convex, beneath the lens striated and indented in a wrinkled manner, columella reflected, aperture rather small, lip simple; light fulvous bay.

Hab. Cuzco, Bolivia; W. Lobb.

Of a delicate saffron colour throughout.

Species 515. (Mus. Cuming.)

BULIMUS PRETEXTUS. *Bul. testā acuminato-oblongā, subcylindracea, subampullē umbilicata, anfractibus octo, leviter concavis, lacūbus vel obscurē indentatis, columellā latissimā reflexā, aperturā parenscaida, labro simplici, pungulum reflexo; tincta, ceruleo nebula, unicolor rotundatus albido prouincie floccatā, lineis sinuatis albido undulatis longitudinaliter cerebrimē notata, opice fulvescente.*

THE LACED BULIMUS. Shell acuminate oblong, some-

what cylindrical, rather largely umbilicated, whorls eight in number, slightly convex, smooth or obscurely indented, columella very broadly reflected, aperture rather small, lip simple, a little reflexed; cream-colour, clouded with blue, promiscuously flaked with round whitish spots, and very closely longitudinally marked with minute waved whitish lines, apex tinged with light rusty-brown.

Hab. Andes of Caxamareca, Peru; W. Lobb.

A delicate blue-clouded shell, sprinkled with a few white lines, which are irregularly waved and sometimes ramified like veins.

Species 207. (Pl. LXXI. Fig. 207, b, and Pl. LXXII. Fig. 207, c. Mus. Cuming.)

BULIMUS LABEO. It is with much gratification that I am enabled to give an original figure of the *Bulimus labeo*, illustrated at Pl. XXXV, from a figure in the Zoological Journal. This shell, from the Cumingian collection, which I take to be identical with the lost specimen, is of a lighter olive colour, but has a similar narrow dark band beneath the sutures, the columella is similarly plicately twisted, the lip is less thickened but of the same rich purple-black enamel, destitute, however, of the opaque bushy-like dots, which are of seldom specific importance.

It was collected by Mr. Lobb at Limabamba, Peru; a district seldom visited by travellers, and the same in which Lieut. Mawre obtained the original specimen.

Species 516. (Mus. Cuming.)

BULIMUS LOBBII. *Bul. testā subcylindracea-oblongā, compressē umbilicata, aperturā versus suboblique tumida, anfractibus octo, leviter concavis, lacūbus vel obscurē indentatis, columellā latē expansa, aperturā obliquē effusa, labro reflexo; albā, ciliis longitudinalibus fuscovestibus et purpureo-castaneis irregulariter conspicie pectī, pone labrum et apertura fauce purpureo-nigrinante.*

LOBE'S BULIMUS. Shell somewhat cylindrically oblong, compressly umbilicated, rather obliquely swollen towards the aperture, whorls eight in number, slightly convex, smooth or obscurely indented, columella broadly expanded, aperture obliquely effused,

BULIMUS.—PLATE LXXI.

lip reflected; white, conspicuously irregularly painted with longitudinal light-brown and purple-chestnut ribands, interior of the aperture and behind the lip purple-black.

Hab. Banks of the Maranon near Balsas, Peru (on branches of a species of *Jatropha*); W. Lobb.

This fine species is of a delicate cream-white, striped longitudinally by distinct ribands of light brown and dark purple chestnut, without any of intermediate tint. Immediately behind the lip there is more of the dark purple-chestnut, approaching to black, and the interior of the aperture is coloured with the same, having a somewhat metallic hue.

I have the pleasure to name it in honour of Mr. Lobb, botanical collector of Messrs. Veitch & Son, the eminent nurserymen of Exeter, to whose zeal in the pursuit of natural history, the species of this, and the two following plates, bear honourable testimony.

Species 517. (Mus. Cuming.)

BULIMUS PURPURATUS. *Bul. testō subacuminate-ovatā, compressē umbilicatā, anfractibus sex, convexis, longitudinaliter rugoso-corrugatis, ad suturas plicato-crenatis, anfractu ultima ventricosiusculo, columellā reflexā, labro simplici; purpureo-fuscā, lineis albidis hic illic longitudinaliter interruptā, basi et apertura fave albida.*

THE PURPLE-STAINED BULIMUS. Shell somewhat acuminately ovate, compressly umbilicated, whorls six in number, convex, longitudinally rugosely wrinkled, plicately crenated at the sutures, last whorl rather ventricose, columella reflected, lip simple; purple-brown, longitudinally interrupted here and there with whitish lines, white at the base and within the aperture.

Hab. Andes of Caxamarea, Peru; Mr. W. Lobb.

A rather stout rough shell, stained with dark purple-brown.




519.



518.



520.



267 c.



521 a.



522.



521 b.

BULIMUS.

PLATE LXXII.

Species 518. (Mus. Cuming.)

BULIMUS RHODOLARYNX. *Bul.* testā acuminato-ovatā, basin versus obliquē ventricosā, ampliter umbilicatā, anfractibus septem ad octo, subrotundatis, levibus, sub lente striis obliquis elevatisculis et spiralibus incisis minutè decussatis, aperturā suborbiculari, columellā labroque latē reflexis; rosco-albicante, intus purpureo-roseā.

THE RED-THROAT BULIMUS. Shell acuminately ovate, obliquely ventricose towards the base, largely umbilicated, whorls seven to eight in number, somewhat rounded, smooth, minutely decussated, beneath the lens, with slightly raised oblique striae and engraved spiral ones, aperture nearly orbicular, lip and columella broadly reflected; rose-white, interior purple-rose.

Hab. Banks of the Aparimao, Alto-Peru; W. Lobb.

Distinguished by its very delicate purple-rose interior, the colour of which is seen through the substance of the shell.

Species 519. (Mus. Cuming.)

BULIMUS DECUSSATUS. *Bul.* testā acuminato-oblongā, basi rotundata, subcylindravō, compressè umbilicatā, anfractibus octo, leviter convexis, longitudinaliter corrugato-striatis, ad suturas subcrenulatis, columellā latē reflexā, labro tenui, simplici; pallidè stramineā, strigis brevibus rufo-fuscis, longitudinalibus et obliquis, fasciatis decussatis, peculiariter notatā.

THE DECUSSATED BULIMUS. Shell acuminately oblong, rounded at the base, somewhat cylindrical, compressly umbilicated, whorls eight in number, slightly convex, longitudinally striated in a wrinkled manner, slightly wrinkled at the sutures, columella broadly reflected, lip thin, simple; light straw-colour, peculiarly marked with short red-brown longitudinal and oblique streaks crossing over each other in bands.

Hab. Andes of Caxamarca, Peru; W. Lobb.

Singularly characterized by the bands of short brown streaks, ranging obliquely in the direction opposed to the lines of growth.

Species 520. (Mus. Cuming.)

BULIMUS MYRISTICUS. *Bul.* testā acuminato-oblongā, basi subobliquā rotundata, compressè umbilicatā, anfractibus octo ad novem, planulato-convexis, longitudinaliter striatis, infra suturas suborenulatis, columellā latē reflexā, labro simplici; albida, vittis irregularibus castaneis et fuscocastaneis confertim longitudinaliter pictā.

THE NUTMEG BULIMUS. Shell acuminately oblong, somewhat obliquely rounded at the base, compressly umbilicated, whorls eight to nine in number, rather flatly convex, longitudinally striated, slightly crenulated beneath the sutures, columella broadly reflected, lip simple; whitish, closely longitudinally painted with irregular chestnut and pale brown ribands.

Hab. Andes of Caxamarca, Peru; W. Lobb.

This differs but little from the preceding species in form and detail of sculpture; yet there is a marked distinction in the style of painting.

Species 521. (Mus. Cuming.)

BULIMUS ALTO-PERUVIANUS. *Bul.* testā acuminato-ovatā, tenuiculā, ampliter compressè umbilicatā, aperturam versus obliquē ventricosā, inflatā, anfractibus septem, convexis, apicem versus cerebrinē elevato-striatis, striis spiralibus incisis sub lente minutè decussatis, columellā latē verticaliter reflexā, aperturā obliquē suborbiculari, labro effuso, non reflexo; fulvescentia lactea, apicem versus cerulescente, lineis castaneis subdistantibus irregulariter longitudinaliter notata, maculis brevibus contrariè obliquis bifasciatim pictā, macularum serie unicā infra suturas.

THE ALTO-PERUVIAN BULIMUS. Shell acuminately ovate rather thin, largely compressly umbilicated, obliquely ventricose towards the aperture, inflated, whorls seven in number, convex, very closely elevately striated towards the apex, minutely decussated, beneath the lens, with engraved spiral striae, columella broadly vertically reflected, aperture obliquely suborbicular, lip effused, not reflected; fulvous cream-colour, bluish towards the apex, irregularly longitudinally marked with rather distant chestnut lines, and painted with two bands of spots arranged contrariwise obliquely, with a single row of spots also beneath the sutures.

BULIMUS.—PLATE LXXII.

Hab. Chachapoyas, Alto-Peru; W. Lobb.

The painting of this delicate and boldly convoluted shell is characterized by two bands of short oblong chestnut spots or dashes, ranging obliquely in a direction contrary to that of the painted lines, as in *B. Sowerbyi*.

Species 522. (Mus. Cuming.)

BULIMUS ALUTACEUS. *Bul. testā oblongo-ovalā, umbilicatā, anfractibus septem, convexis, creberrimè corrugato-striatis, aperturā ovali, columellā reflexā, labro vix reflexo;*

intensē ustulato-castaneā, zonulā unicō albā medio cingulatā, labro albo.

THE LEATHERY BULIMUS. Shell oblong ovate, umbilicated, whorls seven in number, convex, very closely corrugately striated, aperture oval, columella reflected, lip but slightly reflexed; very dark burnt chestnut, encircled round the middle with a single white zone, lip white.

Hab. Cuzco, Bolivia; W. Lobb.

The entire surface of this shell is sculptured longitudinally with very closely packed crinkled striae.



523.



524.



525.



526.



527. a



527. b



528.

BULIMUS.

PLATE LXXIII.

Species 523. (Mus. Cuming.)

BULIMUS CLAUSILIOIDES. *Bul. testā elongato-turrītā, subcylindraceā, sinistrā, compressā umbilicatā, onfractib⁹ novem, supernē obscurē costatis, longitudinātiler creberrimē et minutissimō rugoso-striatis, columellā verticaliter reflexā, aperturā subquadratā, labro tenui, simplici; colore murino.*

THE CLAUSILIA-LIKE BULIMUS. Shell elongately turreted, somewhat cylindrical, sinistral, compressly umbilicated, whorls nine in number, obscurely ribbed round the upper part, longitudinally very closely and very minutely roughly striated, columella vertically reflected, aperture somewhat square, lip simple; mouse-colour.

Hab. Andes of Caxamarca, Peru; W. Lobb.

Very like a *Clausilia* in form, and of a silken aspect, arising out of the very close and minute development of longitudinal striae.

Species 524. (Mus. Cuming.)

BULIMUS DEPTUS. *Bul. testā subacuminato-ovalē, compressā umbilicatā, onfractib⁹ septem, rotundatis, levibus, supernē depressissimis, minuti plicato-crenulatis, columellā reflexā, aperturā pareiangularis, labro simplici; intus exusque ustulato-fuscā, hic illuc saturatiore strigata.*

THE TAXED BULIMUS. Shell subacuminately ovate, compressly umbilicated, whorls seven in number, slightly depressed round the upper part, minutely plicately crenulated, columella reflected, aperture rather small, lip simple; light scorched brown within and without, streaked here and there of a darker colour.

Hab. Chachapoyas, Alto-Peru; W. Lobb.

A thin shell approaching in form and colouring to the *Bulimus aux*, from which it differs in being of a lighter structure and having a more rounded aperture.

Species 525. (Mus. Cuming.)

BULIMUS NIGROPILEATUS. *Bul. testā acuminato-ovalā, sublongo-ovalē, umbilicatā, onfractib⁹ septem, convexis, obtuse subrugoso-striatis, columellā verticaliter reflexā.*

apertura ovali, labro simplici; nimbā, basi versu obsoletè fuscō-fasciatā, apice nigro.

THE BLACK-TIPPED BULIMUS. Shell acuminate ovate, rather largely umbilicated, whorls seven in number, convex, obtusely somewhat roughly striated, columella vertically reflected, aperture oval, lip simple; whitish, obsoletely banded with brown towards the base, apex black.

Hab. Chachapoyas, Alto-Peru; W. Lobb.

It is probable from the faintly banded appearance of this shell, that this is but the pale variety of a darker type.

Species 526. (Mus. Cuming.)

BULIMUS FOVEOLATUS. *Bul. testā oblongo-ovalē, tenuicula, subventricosā, bandū umbilicatā, ad apicem obtusa, onfractib⁹ quinque ad sex, convexis, longitudinaliter obtuse plicato-striatis, punctis oblongis spiraliiter lineariter excutis, infra suturas plicato-crenulatis, apicem versus peculiariter foreolatis, suturis radiis, onfractū ultimō obliquū descendente, columellā latā, depresso-estutā, obliquū recedente, aperturā oblongo-ovali, labro subincassato, cir reflexo; intensē oliveo-brunneā, infra suturas pallide unifasciatā, suturis albidi, columellā labroque coralescente-albīs, apertura fave iridescente-filacea.*

THE FITTED BULIMUS. Shell oblong-ovate, rather thin, slightly ventricose, not umbilicated, obtuse at the apex, whorls five to six in number, convex, longitudinally obtusely plicately striated, spirally engraved with lines of oblong punctures, plicately crenulated beneath the sutures, peculiarly pitted towards the apex, sutures rude, last whorl descending obliquely, columella broad, rather depressed, obliquely receding, aperture oblong-oval, lip somewhat thickened, but little reflected; very dark olive-brown, with a single pale band beneath the sutures, sutures whitish, lip and columella bluish-white, interior of the aperture iridescent lilac.

Hab. Vitoc, near Sarma, Alto-Peru; W. Lobb.

This is the species which Dr. Pfeiffer has assigned to the *Bulimus Madagaci*, of Sowerby, Conch. Illus. f. 59, of which I can find no description or tidings.

BULIMUS.—PLATE LXXXIII.

On comparing the shell under consideration with Mr. Sowerby's figure, it will be found to be of a more oblong form, more acuminate at the apex, and very peculiarly pitted round the upper sutures, reminding one very much of the indentations in the shells of *Phorus*.

Species 527. (Fig. a, b, Mus. Cuming.)

BULIMUS PRIMULARIS. *Bol. testâ acuminato-ovatâ, tenuiculâ, subcentricosâ, umbilicatâ, anfractibus septem, levibus, columellâ tenui, reflexâ, labro simplici; albida, basin versus vivide luteâ, fasciis castaneo-nigris quatuor aut pluribus, nonnullis nullo interruptis, cingulatâ.*

THE PRIMROSE-COLOURED BULIMUS. Shell acuminate ovate, rather thin, somewhat ventricose, umbilicated, whorls seven in number, smooth, columella thin, reflected, lip simple; whitish, bright yellow towards the base, encircled towards the base, with four or more chestnut-black bands, some of which are much interrupted.

Hab. Chachapoyas, Alto-Peru; W. Lobb.

Of simple structure, but abundantly characterized by its bright primrose colour and dark basal bands.

Species 528. (Mus. Cuming.)

BULIMUS COLUMELLARIS. *Bul. testâ cylindraceo-elongatâ, haud umbilicata, anfractibus septendecim ad octodecim, planis, angustis, obliquè subsoletè striatis, anfractu ultimo ad basin subangulato, columellâ tortuosa, leviter recedente, operaturâ subquadratâ, ad basin effusa; roseo-albicante, apice versus livido-carulescente et rufescente.*

THE LITTLE-COLUMN BULIMUS. Shell cylindrically elongated, not umbilicated, whorls seventeen to eighteen in number, flattened, narrow, obliquely rather obsoletely striated, last whorl somewhat angled at the base, columella twisted, slightly receding, aperture nearly square, effused at the base; pinkish-white, livid blue and red towards the apex.

Hab. Andes of Caxamarca, Peru, (under stones at an elevation of 12,000 feet); W. Lobb.

An interesting *Pupa*-like species, distinguished by its square effuse aperture and erect columnar form.



529



530 a



531



532



533



535



536

BULIMUS.

PLATE LXXIV.

Species 529. (Mus. Cuming.)

BULIMUS DEPICTUS. *Bul. testā subfusiformi-oblongā, vix umbilicatā, anfractibus sex, levibus, vel, sub leate, longitudinaliter tenuistriatis, columellā reflexā, aperturā sub-oblongā, labro simplici; lateo-albā, cernueo-virescente longitudinaliter variegatā, fasciis duabus angustis in anfractu ultimo pratermittente.*

THE PAINTED BULIMUS. Shell somewhat fusiformly oblong, scarcely umbilicated, whorls six in number, smooth, or beneath the lens, finely striated, columella reflected, aperture rather oblong, lip simple; yellowish white, longitudinally variegated with blue-green, leaving two narrow bands round the last whorl.

Hab. New Granada.

A prettily painted species collected in the above named locality by Mr. Linden.

Species 530. (Fig a and b, Mus. Philippi.)

BULIMUS RECLUZIANUS. *Bul. testā subfusiformi-oblongā, vix umbilicatā, anfractibus sex, convexis, lineis subtilibus confertim spiraliter incisis, columellā tenuē reflexā, labro simplici; corneo-albidi, strigis latīs irregulāribus castaneis et griseo-violaceis ornata.*

RECLUZ'S BULIMUS. Shell somewhat fusiformly oblong, scarcely umbilicated, whorls six in number, convex, closely spirally engraved with fine lines, columella thinly reflected, lip simple; flesh-white, ornamented with irregular chestnut and grey-violet streaks.

PFEIFFER, Zeitschr. fur Malac. 1847. p. 82.

Hab. —?

A very distinct and characteristic species, for the loan of which I am indebted to Dr. Philippi of Cassel.

Species 531. (Mus. Beleher.)

BULIMUS SOLIVAGUS. *Bul. subpyramidalis-ovatā, crassissimulā, basi subventricosā, hanc umbilicatā, anfractibus sex, levibus, columellā paululum incrassatā, arcuatā, labro reflexo; intus castaneo-purpureā, epidermide albida hydrophanā undique indutā.*

THE WANDERING BULIMUS. Shell somewhat pyramidal-ovate, rather thick, slightly ventricose, not umbilicated, whorls six in number, smooth, columella a little

thickened, arched, lip reflected; deep chestnut-purple, covered throughout with a whitish hydrophanous epidermis.

Hab. Philippine Islands; Belcher.

Distinct from any of the species collected in the same locality by Mr. Cuming.

Species 532. (Mus. Cuming.)

BULIMUS LILACINUS. *Bul. testā fusiformi-oblongā, compressè umbilicatā, anfractibus septem, convexis, longitudinaliter plicato-striatis, columellā reflexā, paululum appressā, labro vix reflexo; albidi, purpureis intus undique fasciatā, columellā viciè violaceo-purpureā.*

THE LILAC BULIMUS. Shell fusiformly oblong, compressly umbilicated, whorls seven in number, convex, longitudinally plicately striated, columella reflected, a little appressed, lip slightly reflected; whitish, banded throughout with violet-purple, columella deep violet-purple.

Hab.

There may be other varieties of this very interesting species. The present is lined interiorly with deep purple bands which have an obscure blackish appearance on the outside.

Species 533. (Mus. Pfeiffer.)

BULIMUS DUNKERI. *Bul. testā acuminato-oblongā, subinflatā, umbilicatā, anfractibus sex, longitudinaliter rugoso-striatis, lineis spiralibus impressis obsolete reticulatā, columellā latē reflexā, aperturā ovali, labro tenui, vix reflexo; albidi, spadiceo et cinnereo-fusco variegatā et nebulatā.*

DUNKER'S BULIMUS. Shell acuminate oblong, slightly inflated, umbilicated, whorls six in number, longitudinally roughly striated, obsolete reticulated with spiral impressed lines, columella broadly reflected, aperture oval, lip thin, scarcely reflected; whitish, variegated and clouded with bay and ash-brown.

PFEIFFER, Abbild. und Besch. Bulimus, Pl. 4. f. 10.

Hab. Mexico.

Mottled and streaked with ash-brown upon a whitish ground, leaving a few light spots or dashes upon the darker parts.

August, 1849.

Species 534. (Mus. Cuming.)

BULIMUS SUCCINCTUS. *Bul. testā pyramidalī-conicā, crassiusculā, hand umbilicatā, anfractibus septem, convervis, levibus; albā, anfractibus fasciā angustā castaneā supra et infra sutras cingulatā, anfractū ultimo infra fasciā inferiori ustulato-castaneo tinctō, circa columellam intensoire.*

THE UNDER-GIRT BULIMUS. Shell pyramidally conical, rather thick, not umbilicated, whorls seven in number, convex, smooth; white, whorls encircled above and below the sutures with a narrow chestnut band, last whorl stained below the lower band with burnt chestnut, darker round the umbilicus.

Hab. Philippines.

The loss of its epidermis prevents me from giving a complete description of this species; but the peculiarity of form and colouring abundantly testifies that it is distinct from any hitherto known.

Species 535. (Mus. Cuming.)

BULIMUS AMBUSTUS. *Bul. testā oblongo-ovatā, vix umbilicatā, anfractibus sex, levibus, vel, sub leute, cerebrinū striatis, columellā reflexā, appressā, recedente, aperturā subtilatatā, labro simplici; albā, strigis ustulato-fasciis, nunc latiusculis, nunc angustis, irregulärer pictā.*

THE SCORCHED BULIMUS. Shell oblong-ovate, scarcely umbilicated, whorls six in number, smooth, or, beneath the lens, very closely striated, columella reflected, appressed, receding, aperture slightly dilated, lip simple; whitish, irregularly painted with burnt-brown streaks, some narrow, some rather broad.

Hab. ——?

The base of the aperture obtains a slightly dilated form, the receding growth of the columella.

Species 537. (Mus. Cuming.)

BULIMUS CERUSSATUS. *Bul. testā acuminato-oblongā, sub-ampliter compressē umbilicatā, anfractibus septem, convexis, levibus, columellā latissimā, vix reflexā, aperturā subobliquā productā, labro simplici; crelaceo-albā, strigis perplicis nudatatis fasciis intus pictā.*

THE WHITE-LEAD BULIMUS. Shell acuminately oblong, rather largely compressly umbilicated, whorls seven in number, convex, smooth, columella very broad, scarcely reflected, aperture rather obliquely produced, lip simple; chalk-white, painted interiorly with a few brown waved streaks.

Hab. ——?

Painted interiorly with scattered brown waved streaks, which show somewhat obscurely through the substance of the shell.

Species 537. (Mus. Pfeiffer.)

BULIMUS SPADICEUS. *Bul. testā globose-conicā, tenuiculā, umbilicatā, basi subplanata, anfractibus sex, convexo-declivibus, sulcari sebimpresso, subtilissimē rugosostriatis, anfractū ultimo obsolete carinato, columellā brevi, latē reflexā, labro simplici; intensē rufō-castanē, apertura fance iridescente.*

THE DARK-RAY BULIMUS. Shell globose conical, rather thin, umbilicated, somewhat flat at the base, whorls six in number, convexly slanting, sutures rather impressed, very finely roughly striated, last whorl obsoletely keeled, columella short, broadly reflected, lip simple; deep reddish chestnut, interior of the aperture iridescent.

MENKE, Pfeiffer, Symbolae, vol. iii. p. 17.

Hab. Natal, South Africa (in the woods near the river Umlass); Krauss.

One of the many interesting new species of shells collected in the vicinity of Port Natal, by the indefatigable Dr. Krauss of Stuttgart.



B U L I M U S.

PLATE LXXV.

Species 538. (Mus. Pfeiffer.)

BULIMUS GASTRUM. *Bul. testā obeso-ovatā, subcylindraceā, compressè umbilicatā, anfractibus sex, concavisculis, obliquè striatis, columella brevis, subdilatata, aperturā semiovali, labro plano-reflexo; pellacido-albā, epidermide latescente indutā, suturis opaco-albis.*

THE SWOLLEN BULIMUS. Shell stoutly ovate, somewhat cylindrical, compressly umbilicated, whorls six in number, rather convex, obliquely striated, columella short, rather dilated, aperture semioval, lip flatly reflected; transparent white, covered with a yellowish epidermis, sutures opaque-white.

EHRENBURG, Symb. phys., Pfeiffer Monog. Helic. vol. ii. p. 122.

Hab. Syria.

A swollen Pupa-like species with a small obtuse apex.

Species 539. (Mus. Pfeiffer.)

BULIMUS BARBARUS. *Bul. testā acuminato-oblongā, compressè umbilicatā, anfractibus sex, concavis, obliquè striatis, columella labroque via reflexis; cornēa, flaminis albōdis variegatā.*

THE FOREIGN BULIMUS. Shell acuminately oblong, compressly umbilicated, whorls six in number, convex, obliquely striated, lip and columella but slightly reflected; horny, variegated with whitish flames.

PFEIFFER, Monog. Helicidae, vol. ii. p. 124.

Helix barbarus, Linnaeus.

Bulimus Jeanoli, Tervér.

Bulimus Terevii, Forbes.

Hab. Algeria.

Although this shell is of a semitransparent horny substance, the opaque grey-white colouring occupies the larger surface.

Species 540. (Mus. Dunker.)

BULIMUS SOLUTUS. *Bul. testā turrītā, laevē convolutū, anfractibus sex, primis quatuor contiguis, ultimis duabus omnino solutis, longitudinaliter rugoso-striatis, supra et infra angulato-cornutis, carinis lineā impressā utrinque marginatis, aperturā subquadrato-ovatā, labro simplici; griseo-albatico.*

THE UNLOOSED BULIMUS. Shell turreted, loosely con-

volute, whorls six in number, the first four contiguous, the last two quite unrolled, longitudinally roughly striated, angularly keeled above and below, keels edged on both sides with an impressed line, aperture somewhat squarely oval, lip simple; greyish-white.

TROSCHEI, Zeitschr. für Malac. 1847, p. 49.

Hab. Peru.

A curious species for which I am indebted to Dr. Dunker, in which the last two whorls are unrolled after the manner of a *Vermelus.*

Species 541. (Mus. Pfeiffer.)

BULIMUS BERTHELOTI. *Bul. testā cylindraceo-oblongā, compressè umbilicatā, apice obtusè eleutatā, anfractibus octo, concavo-planis, obliquè striatis, columella latè cervicaliter dilatata, aperturā latiuscula, labro latè reflexo, sordide albā.*

BERTHELOT'S BULIMUS. Shell cylindrically oblong, compressly umbilicated, apex obtusely elevated, whorls eight in number, convexly flattened, obliquely striated, columella broadly vertically dilated, aperture rather broad, lip broadly reflected; dead white.

PFEIFFER, Symbolae, vol. iii. p. 83.

Hab. Canary Islands.

Very closely allied to the *B. Ehrenbergii* which, probably owing to some misarrangement, is far removed from this species in Dr. Pfeiffer's monograph.

Species 542. (Mus. Dunker.)

BULIMUS RINGENS. *Bul. testā fusiformi-cylindracea, compressè umbilicatā, anfractibus octo, levibus, aperturā oblongo-quadratā, effusa, conspicue quadridentata: corneo-fuscā, cinereo-albo strigata et variegata, aperturā violaceo-nigrigante tinctā.*

THE GRINNING BULIMUS. Shell fusiformly cylindrical, compressly umbilicated, whorls eight in number, smooth, aperture oblong square, effused, conspicuously four-toothed; horny brown, streaked and variegated with ash white, aperture stained with violet black.

DUNKER, Zeitschr. für Malac. 1847, p. 83.

Hab. Macahe, Brazil.

The aperture is characterized by having four conspi-

euos flattened teeth, of which the interstices are variously stained with violet-black.

Species 543. (Mus. Pfeiffer.)

BULIMUS VARNENSIS. *Bul. testá cylindraceo-elongatā, solidiusculā, compressè umbilicatā, anfractibus decem, planō-convexitā, obliquē irregulariter striatis, aperturā poriuseculā, columellā labroque dilatatis, non reflexis; albā, intus fuscescente, apice nigricante.*

THE VARNA BULIMUS. Shell cylindrically elongated, rather solid, compressly umbilicated, whorls ten in number, flatly convex, obliquely irregularly striated, aperture rather small, lip and columella dilated, not reflected; white, brownish within, apex blackish.

FRIVALDSKY, Pfeiffer, Monog. Helicidae, vol. ii. p. 227.
Hab. Near Varna, Turkey.

Of a clear white, without any indication of marking.

Species 544. (Mus. Pfeiffer.)

BULIMUS MEMBRANACEUS. *Bul. testá subfusiforme-oblóngā, tenui, hyalinā, compressè umbilicatā, anfractibus septem, planō-convexitā, lăvibūs, sub lente minnī striatis, columellā latē reflexā, depresso, aperturā subobliquā, labro simplici, paululum effusā; pellucido-albicans.*

THE MEMBRANACEOUS BULIMUS. Shell somewhat fusiformly oblong, thin, hyaline, compressly umbilicated, whorls seven in number, flatly convex, smooth, minutely striated beneath the lens, columella broadly reflected, depressed, aperture rather oblique, lip simple, a little effused; transparent white.

PHILIPPI, Abbild. und Besch. Conch. *Bul. Pl. 5. f. 1, 2.*
Hab. —?

A delicate transparent white species, probably inhabiting Central America.

Species 545. (Mus. Pfeiffer.)

BULIMUS FRIVALDSKYI. *Bul. testá cylindraceo-oblóngā, vix compressè umbilicatā, anfractibus novem, planiusculis, subtilliter obliquē striatis, aperturā parvā, obliquā, seniorali, columellā labroque appressis; pellucido-corneā, crassiusculā.*

FRIVALDSKY'S BULIMUS. Shell cylindrically oblong, but slightly compressly umbilicated, whorls nine in number, rather flattened, finely obliquely striated, aperture small, oblique, semi-oval, lip and columella appressed; transparent horny, rather thick.

PFEIFFER, Zeitschr. für Malac. 1847, p. 191.

Hab. Natolia, Turkey in Asia.

A small stout horny cylindrical shell of which the surface is rather flattened.

Species 546. (Mus. Pfeiffer.)

BULIMUS COMPACTUS. *Bul. cylindraceo-oblóngā, solidiusculā, subcompressè umbilicatā, anfractibus novem ad decem, obliquē ruguloso-striatis, aperturā truncato-ovali, tridentalē; opaco-albā, fuscescente pallidē striatā, intus subaurantia.*

THE COMPACT BULIMUS. Shell cylindrically oblong, rather solid, somewhat compressly umbilicated, whorls nine to ten in number, obliquely roughly striated, aperture truncately oval, three-toothed; opake-white, faintly streaked with light brown, interior tinged with orange. FRIVALDSKY, Pfeiffer, Zeitschr. für Malac. 1848, p. 6.
Hab. Island of Crete.

A slight contraction of the aperture forms a noticeable feature in this species.

Species 547. (Mus. Pfeiffer.)

BULIMUS TRICHODES. *Bul. testá acuminato-oblóngā, tenui, umbilicatā, anfractibus octo, subtilissimè granulato-striatis, aperturā ovali, columellā reflexā, labro simplici; olivaceo-corneā, lineis opaco-albis longitudinaliter notatā, pilis brevissimis obsitā.*

THE HAIRY BULIMUS. Shell acuminate-oblong, thin, umbilicated, whorls eight in number, very finely granulately striated, aperture oval, columella reflected, lip simple; olive-horny, longitudinally marked with opake white lines, covered with very short hair.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 277. pl. 33. f. 1-5.

Hab. Province of Santa Cruz, Bolivia; D'Orbigny.

Distinguished by its peculiar hairy covering.



548



549



550



551



552



553



554



555



556



557

— 10 — 20 —

— 10 — 20 —

B U L I M U S.

PLATE LXXVI.

Species 548. (Mus. Cuming.)

BULIMUS BURCHELLII. *Bul. testā acuminato-ovatā, vir umbilicatā, anfractibus octo, levibus vel ruditē obsoletē striatis, columellā breviter reflexā, labro simplici; albidā, fuscescente pallido strigatā.*

BURCHELL'S BULIMUS. Shell acuminate ovate, scarcely umbilicate, whorls eight in number, smooth or rudely obsoletely striated, columella shortly reflected, lip simple; whitish, faintly streaked with brown.

GRAY, Pro. Zool. Soc., 1834, p. 66.

Hab. Cape Natal, South Africa.

I am not very sure that this is the true *B. Burchellii*, of which Mr. Gray describes having seen a number strung together as a necklace. Mr. Cuming has the name attached to it in his cabinet with a mark of doubt, which I am unable to clear up.

Species 549. (Mus. Cuming.)

BULIMUS RECONDITUS. *Bul. testā acuminato-oblongā, tenuicollā, umbilicatā, anfractibus octo, levibus, vel, sub lete, minutissimè striatis, columellā breviter reflexā, labro simplici; cinereo-albicante, carnea-fuso longitudinaliter punctatā et partim strigatā, aperture fave pallidē aurantiā.*

THE CONCEALED BULIMUS. Shell acuminate oblong, rather thin, umbilicate, whorls eight in number, smooth, or, beneath the lens, very minutely striated, columella shortly reflected, lip simple; ashy white, longitudinally dashed and dotted with blue-brown, interior of the aperture tinged with orange.

Hab. —?

Very prettily marked with dark longitudinal dashes and finely dotted lines.

Species 550. (Mus. Cuming.)

BULIMUS NEURICUS. *Bul. testā acuminato-oblongā, vir umbilicatā, anfractibus septem, levibus vel, sub lente, minutē striatis, columellā reflexā, labro simplici; pallidē straminea, fasciā augsā nigricante-costatae lineice interrupitā cingulatā, anfractū ultimo fasciā duabus cingulato.*

THE GOUTY BULIMUS. Shell acuminate oblong, scarcely umbilicate, whorls seven in number, smooth, or

beneath the lens, minutely striated, columella reflected, lip simple; pale straw colour, encircled with a narrow blackish chestnut band and an interrupted line, last whorl with two bands.

Hab. Taboo, Cape Palmas (near the sea-shore).

The interrupted line encircles the whorls at about the middle, the two bands are round the lower part, the lower band being concealed in all but the last whorl.

Species 551. (Mus. Cuming.)

BULIMUS OBLIQUEUS. *Bul. testā subpyramidalis-oratā, umbilicatā, crossincolata, ad basim oblique producita, anfractibus septem ad octo, levibus, aperturā obliquā, columellā labroque latè dilatata; pallide rosaceā, anfractū ultimo fasciā costatā cincta cingulato.*

THE OBLIQUE BULIMUS. Shell somewhat pyramidal, ovate, umbilicate, rather thick, obliquely produced at the base, whorls seven to eight in number, smooth, aperture oblique, lip and columella broadly dilated; light pink, with a single chestnut band round the last whorl.

Hab. Bahia,

A pink shell of firm structure, encircled with a conspicuous chestnut band round the last whorl.

Species 552. (Mus. Cuming.)

BULIMUS RUGATUS. *Bul. testā subacuminato-ovatā, subbilicatā, anfractibus septem, subventricosis, subtiliter et creberius longitudinaliter rugatis, aperturā subquadrato-sinuatā, columellā reflexā, labro paulatim dilatato; olivaceā.*

THE SHRIVELLED BULIMUS. Shell somewhat acuminate ovate, umbilicate, whorls seven in number, rather ventricose, finely and very closely longitudinally shrivelled, aperture somewhat squarely sinuate, columella reflected, lip a little dilated; olive.

Hab. —?

Chiefly distinguished by its finely shrivelled surface.

Species 553. (Mus. Cuming.)

BULIMUS GELIDUS. *Bul. testā acuminato-ovatā, subampliter umbilicatā, anfractibus octo, convexis, subtilis-*

sinuē striatis, columellā latē reflexā, labo simplici; pellucidō-albā, līneis numerosis opaco-albis longitudinaliter notatā.

THE ICY BULIMUS. Shell acuminate ovate, rather largely umbilicated, whorls eight in number, convex, very finely striated, columella broadly reflected, lip simple; transparent white, longitudinally marked with numerous lines of opaque white.

Hab. Central America?

Somewhat of a bluish-white tinge, characterized with opaque white longitudinal lines.

Species 554. (Mus. Cuming.)

BULIMUS CUNCTATOR. *Bul. testā subabbreviato-ovatā, hand umbilicatā, inferū ventricosā, anfractibus quinque, convexis, lēvibus, columellā rectā, labo sub-incassatō, non reflexo; purpureo-castaneā, epidermide albida hydrophanā longitudinaliter strigatā, intus cerasulente.*

THE LOITERER BULIMUS. Shell somewhat abbreviate ovate, not umbilicated, ventricose round the lower part, whorls five in number, convex, smooth, columella straight, lip slightly thickened, not reflected; purple-chestnut, streaked longitudinally with a white hydrophanous epidermis, interior bluish.

Hab. Philippine Islands; Belcher.

This species, collected by Capt. Sir Edward Belcher during the voyage of the Samarang, is quite distinct from any of the same group discovered by Mr. Cuming.

Species 555. (Mus. Cuming.)

BULIMUS DELUMBIS. *Bul. testā subgloboso-ovatā, umbilicatā, anfractibus sex, rotundatis, longitudinaliter rugoso-striatis, columellā reflexā, labo simplici; cretaceo-albā, līneis brevibus ferrugineo-fuscis irregulariter fasciatā pictā, aperture fauce violascente.*

THE FEEBLE BULIMUS. Shell somewhat globosely ovate, umbilicated, whorls six in number, rounded, longitudi-

nally rugosely striated, columella reflected, lip simple; chalk-white, irregularly painted with bands of short rusty-brown lines, interior of the aperture violet.

Hab. —?

Very delicately tinged with orange and purple violet in the aperture

Species 556. (Mus. Cuming.)

BULIMUS OVULUM. *Bul. testā ovatā, crassiusculā, umbilicatā, anfractibus sex, convexis, lēvibus, vel obscurē striatis, columellā incassatā, latē reflexā, labo reflexo; intus extusque eburneā.*

THE LITTLE EGG BULIMUS. Shell ovate, rather thick, umbilicated, whorls six in number, convex, smooth, or obscurely striated, columella thickened, broadly reflected, lip reflected; ivory white within and without.

Hab. Philippine Islands; Cuming.

A shell of rather solid growth, without any appearance of colouring.

Species 557. (Mus. Cuming.)

BULIMUS FIDUSTUS. *Bul. testā acuminato-ovatā, tenuiculā, eis umbilicatā, anfractibus septem, longitudinaliter minutiē et crebrimē striatis, columellā breviter reflexā, labo tenui, simplici; lutescente-albā, castaneo latifasciatā, epidermide tenui cornue politā indata.*

THE FAITHFUL BULIMUS. Shell acuminate ovate, rather thin, scarcely umbilicated, whorls seven in number, longitudinally minutely and very closely striated, columella shortly reflected, lip thin, simple; yellowish white, broadly banded with chestnut, covered with a thin polished horny epidermis.

Hab. Sebundo, New Granada; Funck.

An interesting species from New Granada, which does not appear to have been described



558



559



560



561



562



563.



564



565.



566



567

BULIMUS.

PLATE LXXVII.

Species 558. (Mus. Cuming.)

BULIMUS COAGULATUS. *Bul. testā ovalā, tenui, ventricosā, vix umbilicatā, aufractibus quinque, obscurè longitudinaliter plicato-striatis, columellā reflexā, labro simplici; corneo-fuscescente, pellucidā, opaco-albo peculiariter undato-reticulatā.*

THE CURDLED BULIMUS. Shell ovate, thin, ventricose, scarcely umbilicated, whorls five in number, obscurely longitudinally plicately striated, columella reflected, lip simple; transparent horny brown, peculiarly reticulated in a waved manner with opaque white.

Hab. Peru.

A light inflated shell distinguished by a peculiarly mottled opaque net-work.

Species 559. (Mus. Cuming.)

BULIMUS UMBRATICUS. *Bul. testā acuminato-ovatā, vix umbilicatā, aufractibus sex, longitudinaliter subtilissimè creberrimè elevato-striatis, columellā breviter reflexā, labro tenui, simplici; pellucido-albā, prope apicem fuscescente strigatā.*

THE SLIGHT BULIMUS. Shell acuminately ovate, scarcely umbilicated, whorls six in number, longitudinally very finely elevatedly striated, columella shortly reflected, lip simple; transparent white, streaked with pale brown near the apex.

Hab. Central America.

It is possible that this may be a light variety of a species of more colour from the indications of marking near the apex.

Species 560. (Mus. Cuming.)

BULIMUS EXORNATUS. *Bul. testā acuminato-ovatā, tenui, vix umbilicatā, aufractibus octo, convexis, levibus, columellā breviter reflexā, aperturā pareinclusā, labro simplici; pellucido-fusca, lineis longitudinalibus opaco-albis profuse strigatā.*

THE WELL-TRIMMED BULIMUS. Shell acuminately ovate, thin, scarcely umbilicated, whorls eight in number, convex, smooth, columella shortly reflected, aperture rather small, lip simple; transparent brown, profusely streaked with longitudinal opaque white lines.

Hab. Chilon, Bolivia (under dead trunks of Cacti); Bridges.

Of transparent horny texture, neatly marked with longitudinal opaque-white lines.

Species 561. (Mus. Benson.)

BULIMUS MAVORTIUS. *Bul. testā acuminato-ovatā, vix umbilicatā, aufractibus septem, convexo-declivibus, levibus, vel, sub lente, tenuistriatis, columellā breviter reflexā, labro simplici; pallidè stramineā, sparsim nigripunctatā, lineis purpurascēt ad suturas, aufractū ultimo zonulis dubias rosaceis cingulato, columellā indensè rosacē, apicē purpureo-vigricante.*

THE MARTIAL BULIMUS. Shell acuminately ovate, scarcely umbilicated, whorls seven in number, convexly slanting, smooth, or, beneath the lens, finely striated, columella shortly reflected, lip simple; pale straw-colour, sprinkled here and there with black dots, and with a purple line round the sutures, last whorl encircled with two narrow pinkish zones, columella deep pink, apex purple-black.

Hab. Ceylon?

An extremely characteristic species, of which I have been favoured with two specimens from the collection of Mr. Benson.

Species 562. (Mus. Cuming.)

BULIMUS IGNAVUS. *Bul. testā oblongo-ovatā, subprofunde umbilicatā, aufractibus septem, rotundatis, longitudinaliter subtilissimè striatis, columellā latā, subreflexā, labro simplici; sordidè cornēt.*

THE SLUGGISH BULIMUS. Shell oblong-ovate, rather deeply umbilicated, whorls seven in number, rounded, longitudinally very finely striated, columella broad, slightly reflected, lip simple; dirty horn colour.

Hab. Central America.

A thin horny shell with the whorls somewhat rounded and rather deeply umbilicated.

Species 563. (Mus. Cuming.)

BULIMUS LIMENSIS. *Bul. testā acuminato-ovatā, subprofunde umbilicatā, aufractibus septem, longitudinaliter creberrimè elevato-striatis, striis irregularibus, subplicatis, columellā reflexā, labro simplici; pallidè ferrugineo-fusca.*

BULIMUS.—PLATE LXXVII.

THE LIMA BULIMUS. Shell acuminate ovate, rather deeply umbilicated, whorls seven in number, longitudinally very closely elevately striated, striae irregular, subPLICATE, columella reflected, lip simple; pale rusty brown.

Hab. Lima and Quito, South America.

This is allied to the *B. scalariformis*, but does not agree with any of the varieties of that species.

Species 564. (Mus. Benson.)

BULIMUS SYLHETICUS. *Bul. subacuminato-oblonga, solidus, sinistrali, umbilicata, anfractibus sex, levibus, columellâ labroque subincrassatis, dilatato-reflexis; vivido luteo, columellâ labroque pallidioribus.*

THE SYLHET BULIMUS. Shell somewhat acuminate oblong, rather solid, sinistral, umbilicated, whorls six in number, smooth, lip and columella rather thickened, dilately reflected; bright yellow, lip and columella paler.

Hab. Sylhet, Eastern Himalaya; Benson.

Having very much the aspect of a young or dwarf specimen of *B. citrinus*, it might easily be confounded with that species.

Species 565. (Mus. Cuming.)

BULIMUS MOLLICELLUS. *Bul. testa subcylindraceo-oblonga, vix umbilicata, suturis subconstrictis, anfractibus sex, subtilissimè irregulariter striatis, ad suturas leviter crenulatis, columellâ reflexâ, appressâ, aperturâ pauculum contractâ, labro simplici; pellucido-albâ, epidermide tenui stramineâ inducta.*

THE DELICATE BULIMUS. Shell somewhat cylindrically oblong, scarcely umbilicated, sutures rather constricted, whorls six in number, very finely irregularly

striated, slightly crenulated at the sutures, columella reflected, appressed, aperture a little contracted, lip simple; transparent white, covered with a thin straw coloured epidermis.

Hab. —?

A delicate small species certainly distinct from any hitherto described.

Species 566. (Mus. Cuming.)

BULIMUS TRANSPARENTE. *Bul. testa acuminato-oblonga, tenuissima, haud umbilicata, anfractibus sex, subtilissimè irregulariter striatis, columellâ tenuissimè reflexâ, labro simplici; pellucido-cornea.*

THE TRANSPARENT BULIMUS. Shell acuminate oblong, very thin, not umbilicated, whorls six in number, very finely irregularly striated, columella very thinly reflected, lip simple; transparent horny.

Hab. —?

Of as light and delicate a structure as any *Lymnea*.

Species 567. (Mus. Cuming.)

BULIMUS NIGRILINEATUS. *Bul. testa conico-ovalata, ventricosa, tenicula, haud umbilicata, anfractibus sex, levibus, vel, sub lente, tenuissimè striatis, columellâ tenuiter reflexâ, labro simplici; stramineâ, lineis nigrofuscis fasciatâ.*

THE BLACK-LINED BULIMUS. Shell conically ovate, ventricose, rather thin, not umbilicated, whorls six in number, smooth, or, beneath the lens, very finely striated, columella thinly reflected, lip simple; pale straw-colour, banded with black-brown lines.

Hab. —?

The bands round the upper part of the whorls are somewhat lighter than those round the lower.



BULIMUS.

PLATE LXXVIII.

Species 568. (Mus. Cuming.)

BULIMUS LATIREFLEXUS. *Bul. testā acuminato-cylindraceā, subobsoletā, semipellucida, compressē umbilicata, anfractibus octo, obliquē creberriū plicato-striatis, infra suturas lineā marginatis, columellā latā, verticali, aperturā suborbiculari, labro late reflexo; fulvescente-cornetā, labro opaco-albidā.*

THE BROADLY-REFLECTED BULIMUS. Shell acuminate cylindric, rather stout, semitransparent, compressly umblicated, whorls eight in number, obliquely very closely plicately striated, edged with a line beneath the sutures, columella broad, vertical, aperture nearly orbicular, lip broadly reflected; fulvous horny, lip opake fulvous-white.

Hab. Muscat ?

Mr. Cuming has every reason to believe that this fine species is from Muscat, Arabia.

Species 569. (Mus. Benson.)

BULIMUS CERATINUS. *Bul. testā acuminato-oblongā, tenui, compressē umbilicata, anfractibus sex ad septem, convexis, subobsoletā irregulariter sulcato-striatis, columellā late verticaliter reflexa, aperturā parviusculā, suborbiculari, pellucido-corneā.*

THE INTRICATE BULIMUS. Shell acuminate oblong, thin, compressly umblicated, whorls six to seven in number, convex, somewhat obscurely irregularly striated in a grooved manner, columella broadly vertically reflected, aperture rather small, nearly orbicular; transparent horny.

BENSON, MSS.

Hab. Almorah, Kemaon, Western Himalaya; Capt. Boys.

Distinguished from *B. moulensis*, to which it is next allied, by its delicate transparent texture and absence of granules.

Species 570. (Mus. Benson.)

BULIMUS RUFISTRIGATUS. *Bul. testā cylindraco-oblongā, compressē umbilicata, anfractibus octo, plano-convexis, levibus, ad suturas minutē crenulatis, aperturā parviusculā, suborbiculari; fulvo-corneā, opaco-albo irregulariter tenuistrigata.*

THE RED-STREAKED BULIMUS. Shell cylindrically oblong, compressly umblicated, whorls eight in number, flatly convex, smooth, minutely crenulated at the sutures, aperture rather small, nearly orbicular; fulvous horny, irregularly finely streaked with opaque-white.

BENSON, MSS.

Hab. At the base of the lower range of the Himalaya, between the rivers Jumna and Sutlej; Hutton.

Communicated to Mr. Benson by Capt. Hutton with the title *Pupa strigata*, of which the specific name is occupied in the present genus.

Species 571. (Mus. Benson.)

BULIMUS BENSONI. *Bul. testā acuminato-ovata, hand umbilicata, anfractibus octo, transversi minute et creberriū striatis, striis subtilissimè granulatis, peculiariter corrugato-andulatis, columellā tenuioidi, leviter contortā, labro vir reflexo; violaceo-alba, purpurascē-fusco variegatā, fasciis albis duabus tribusve, maculis intenſe purpureis sagittalibus notatis, cingulatā, columellā castaneo-fusca.*

BENSON'S BULIMUS. Shell acuminate ovate, not umblicated, whorls eight in number, transversely minutely and very closely striated, striae very finely granulated, peculiarly corrugately waved, columella rather thin, slightly twisted, lip scarcely reflected; violet-tinged white, variegated with purple-brown, encircled with two or three white bands marked with deep purple arrow-head spots, columella chestnut brown.

Hab. Banks of the river Amazon.

This fine species, which I have the honour to name after a gentleman who has afforded me much valuable information on the *Bulinia* of Bengal, Afghanistan and North Western India, is closely allied in general aspect to the *B. zebra*. Its chief point of difference consists in the minute waved-striated semigranular sculpture which characterises the entire surface of the shell, whilst the design of painting is marked by the presence of two white bands, painted with dark purple arrow-head spots, pointing towards the lip.

Fig. 572.

As this species belongs more properly to *Achatina*, the reader is referred to Plate XXI. of that genus for description. For *Bulimus* Sp. 572, see Pl. LXXX.

Species 573. (Mus. Benson.)

BULIMUS EREMITA. *Bul. testá cylindraceo-oblongá, subprofundè umbilicatá, anfractibus octo, convexis, obscurè subirregulariter striatis, aperturá obliquè productá, columellá latá, labro paululum expanso; carulescente-albá.*

THE HERMIT BULIMUS. Shell cylindrically oblong, rather deeply umbilicated, whorls eight in number, convex, obscurely rather irregularly striated, aperture obliquely produced, columella broad, lip a little expanded: bluish-white.

BENSON. MSS.

Hab. Afghanistan: Hutton.

Collected by Capt. Hutton on his march with Shah Shuja's force from the Bolan Pass to Cabul, during the recent operations of the British Army in India.

Species 574. (Mus. Benson.)

BULIMUS RUSTICUS. *Bul. testá ovatá, solidiusculá, haud umbilicata, spirá obtusá, anfractibus quinque ad sex, convexis, levibus, columellá plavo-increassatá, subcallosá, quasi dentalá, aperturá obliquá, labro reflexo; castaneo-brunneá, circa columellam obscurè fasciatá, albida versus apicem, columellá labroque albis.*

THE SIMPLE BULIMUS. Shell ovate, rather solid, not umbilicated, spire obtuse, whorls five to six in number, convex, smooth, columella flatly thickened, rather callous, as if toothed, aperture oblique, lip reflected; chestnut-brown, obscurely banded around the columella, white towards the apex, lip and columella white.

Mousson, Land and Suss. Moll. Java, p. 114. pl. 22 f. 1.
Hab. Eastern Java.

I am indebted to Mr. Benson for an authentic example of this species, received from M. Mousson. It is, however, impossible to describe it satisfactorily, owing to the shell having lost its epidermis, which, to judge by analogy, would afford important characters.

The author notices having a number of dead specimens in his possession, collected in Eastern Java, and remarks, that it is the only representative hitherto found in that island, of the richly clad group, comprising *B. pythagaster* and its congeners, inhabiting the Philippines.

Species 575. (Mus. Benson.)

BULIMUS BOYSIANUS. *Bul. testá cylindraceo-oblongá, sinistrali, vix umbilicata, anfractibus octo, convexis, subrugoso-striatis, ad suturas subtilissimè plicatis, lineis spiralibus, sub lete minutè decussatis, columellá reflexa; corneo-fusca, lineis undulatis opaco-albis variegata.*

BOYS' BULIMUS. Shell cylindrically oblong, sinistral, scarcely umbilicated, whorls eight in number, convex, rather coarsely striated, very finely plicated at the sutures, minutely decussated beneath the lens with spiral lines; horny brown, variegated with waved opaque-white lines.

BENSON. MSS.

Hab. Kémaon, Western Himalaya; Boys.

Named by Mr. Benson in honour of Capt. Boys, to whom we are indebted for this and many interesting species collected in North Western India.

Species 576. (Mus. Benson.)

BULIMUS CHERSONESICUS. *Bul. testá acuminato-cylindrácei, sinistrali, compresè umbilicata, anfractibus octo, convexis, subobscure rugoso-striatis, columellá latá, verticali; carulescente-albá, fulvescente hic illuc pallidè tincta, apertura fave pallidi aurantiá.*

THE CRIMEA BULIMUS. Shell acuminately cylindrical, sinistral, compressly umbilicated, whorls eight in number, convex, somewhat obscurely coarsely striated, columella broad, vertical; bluish-white, faintly tinged here and there with light brown, interior of the aperture light orange.

SOWERBY, MSS. Jay, Catalogue, 1839, p. 55.

Hab. Crimea, Southern Russia.

This is one of the list of species unknown to Dr. Pfeiffer. The specimen here figured from the collection of Mr. Benson was received from Mr. Sowerby with the above name in 1834.

Species 577. (Mus. Cuming.)

BULIMUS CONULUS. *Bul. testá conico-ovatá, vix umbilicata, anfractibus septem, rotundatis, obliquè subtilissimè striatis; pelticeudo-cornetá.*

THE LITTLE CONE BULIMUS. Shell conically ovate, scarcely umbilicated, whorls seven in number, rounded, obliquely very finely striated; transparent horny.

Hab. Port Natal, South Africa.

Very thin and transparcuit, of a conical form rather broad at the base.



578



579



580



581



582



583



584



585



586



587

BULIMUS.

PLATE LXXIX.

Species 578. (Mus. Cuming.)

BULIMUS POLYGYRATUS. *Bul. testā pyramidalis-turritā, minutē umbilicatā, anfractibus novem ad decem, sub-augustis, rotundatis, subtiliter plicato-striatis, columellā reflexā, aperturā parvā, labro simplici; cerulescente-alba.*

THE MANY-WHORLED BULIMUS. Shell pyramidalis turreted, minutely umbilicated, whorls nine to ten in number, rather narrow, rounded, finely plicately striated, columella reflected, aperture small, lip simple; bluish-white.

Hab. — ?

An erect closely convoluted shell, of which the sutures are rather impressed.

Species 579. (Mus. Cuming.)

BULIMUS MICRA. *Bul. testā acuminato-turritā, tenuissimā, eix umbilicatā, anfractibus acto, supernè tunidiusculā, rotundatis, longitudinaliter elevato-striatis, quasi costellatis, columellā reflexā, aperturā parvā, labro tenui; pellucidō-stramineā.*

THE SMALL BULIMUS. Shell acuminately turreted, very thin, scarcely umbilicated, whorls eight in number, rather swollen round the upper part, rounded, longitudinally elevately striated, as if finely ribbed, columella reflected, aperture small, lip thin; transparent straw-colour.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 262. pl. 41. f. 18 to 19.

Hab. Chilon, Bolivia (under trunks of Cacti): Bridges.

This little species seems to obtain rather a wide range. According to D'Orbiguy it is found in the environs of Rio Janeiro as well as on the eastern side of the Bolivian Andes.

Species 580. (Mus. Cuming.)

BULIMUS CARACCASENSIS. *Bul. testā acuminato-turritā, umbilicatā, anfractibus novem angustis, rotundatis, longitudinaliter elevato-striatis, infra suturas quasi costellatis, suturas impressis, columellā reflexā, aperturā parvā, labro tenui; sordidō-fulvescente.*

THE CARACAS BULIMUS. Shell acuminately turreted,

umbilicated, whorls nine in number, narrow, rounded, longitudinally elevately striated, beneath the sutures as if finely ribbed, sutures impressed, columella reflected, aperture small, lip thin; dull fulvous-colour.

Hab. Caracas; Sallé.

Very like the first species of this plate, *B. polygyratus*, though smaller and differing in form.

Species 581. (Mus. Cuming.)

BULIMUS TUNETANUS. *Bul. testā oblongo-ovatā, hyalinā, sub-Partula-formā, compressē umbilicatā, anfractibus sex, tunidiusculis, obliquē subtilissimē striatis, columellā latā, expansā, aperturā ovatā, supernè callositate parvā, labro reflexo; hyalino-alba, labro opaco-alba.*

THE TUNIS BULIMUS. Shell oblong-ovate, hyaline, somewhat Partula-shaped, compressly umbilicated, whorls six in number, rather swollen, obliquely very finely striated, columella broad, expanded, aperture ovate, with a small callosity at the upper part, lip reflected; hyaline white, lip opaque-white.

Hab. Tunis; Capt. Spratt.

A delicate transparent shell, though of comparatively solid growth, and with the lip reflected, reminding one of the *Partula hyalina*.

Species 582. (Mus. Cuming.)

BULIMUS CONTIGUUS. *Bul. testā elongato-cylindracea, haud umbilicata, anfractibus novem, plani-convervis, tævis, politis, ad suturas exiliter impresso-crenulatis, columellā latā, verticali, aperturā minuta, marginibus incrassatis, reflexis, supra callo junctis; eburneā.*

THE CLOSELY-CONVOLUTED BULIMUS. Shell elongately cylindrical, not umbilicated, whorls nine in number, flatly convex, smooth, polished, faintly impressively crenulated at the sutures, columella broad, vertical, aperture minute, margins thickened, reflected, joined above by a callosity; ivory-white.

Hab. Island of Socotra.

A very characteristic closely convoluted polished shell with a peculiarly small aperture.

BULIMUS.—PLATE LXXIX.

Species 583. (Mus. Benson.)

BULIMUS OBTUSUS. *Bul. testā cylindraceo-oblongā, basi umbilicatā, apice peculiariter abbreviatā, anfractibus septem ad octo, subplano-convexus, obliquè striatis, columellā arcuata, aperturā parvā, suborbiculari, labro tenui reflexo; coralescente-albā.*

THE OBTUSE BULIMUS. Shell cylindrically oblong, not umbilicated, peculiarly abbreviated at the apex, whorls seven to eight in number, rather flatly convex, obliquely striated, columella arched, aperture small, nearly orbicular, thinly reflected; bluish-white.

PFEIFFER, Symbolæ, vol. i. p. 84. and vol. ii. p. 119.

Pupa obtusa, Draparnaud.

Helix (Cochlostyla) obtusa, Féussac.

Pupa Germannica, Lamarck.

Hab. Australia.

Chiefly distinguished by the peculiar sinking of the first few whorls.

Species 584. (Mus. Cuming.)

BULIMUS EUBOICUS. *Bul. testā acuminato-oblongā, chrysalidiformi, compressè umbilicatā, anfractibus novem, plano-convexus, minutissimè striatus, superne marginatis, aperturā parvā, fortiter tridentalata, supra callositate, labro subreflexo; fulvescente-corneū, dentibus fabroque opaco-albīs.*

THE EUBEAN BULIMUS. Shell acuminate-oblong, chrysallis-shaped, compressly umbilicated, whorls nine in number, flatly convex, very minutely striated, edged round the upper part, aperture small, strongly three-toothed, with a callosity above, lip slightly reflected; fulvous horny, teeth and lip opaque-white.

Hab. Eubea; Capt. Spratt.

Unusually strongly toothed for so small a species.

Species 585. (Mus. Cuming.)

BULIMUS NANUS. *Bul. testā subelongatā, umbilicatā, anfractibus octo, convexis, subtilissimè striatis, striis infra suturas subuplicatis, columellā verticali, aperturā parvā, labro simplici; cinerascente, striis fuscouscentibus hic illuc obscurè notata, apice fuso.*

THE DWARF BULIMUS. Shell rather elongated, umbilicated, whorls eight in number, convex, very finely striated, striae slightly plicated beneath the sutures, columella vertical, aperture small, lip simple; pale ash-colour, obscurely marked here and there with light brown streaks, apex brown.

Hab. Chili.

A small turreted shell very faintly marked with brownish streaks and with a dark-brown apex.

Species 586. (Mus. Cuming.)

BULIMUS MACILENTUS. *Bul. testā acuminatā, basi suboblongā, basi umbilicatā, tenissimā, anfractibus sex, tævibus, politis, vel sub lente subtilissimè concentricè striatis, superne minutè marginatis, columellā vix reflexā, aperturā oblongo-ovalatā, labro tenui; virescente, vitreā.*

THE LEAN BULIMUS. Shell acuminate, rather oblong at the base, not umbilicated, very thin, whorls six in number, smooth, polished, or, beneath the lens, very finely concentrically striated, minutely edged round the upper part, columella only slightly reflected, aperture oblong-ovate, lip thin; greenish, glassy.

Hab. Philippine Islands; Cuming.

A very transparent, greenish, glassy shell, of which the last whorl is rather oblong-inflated.

Species 587. (Mus. Benson.)

BULIMUS SEGREGATUS. *Bul. testā ovato-conicā, subpyramidalis, basi tauridinuscultā, umbilicatā, anfractibus septem, convexis, obliquè striatis, columellā latā, verticali, aperturā parvā, rotundatā, labro tenui; sordide cornēd.*

THE ALIEN BULIMUS. Shell ovately conical, somewhat pyramidal, rather swollen at the base, umbilicated, whorls seven in number, convex, obliquely striated, columella broad, vertical, aperture small, rounded, lip thin; dull horny.

BENSON, MSS.

Hab. Simla, Western Himalaya.

Distinguished from most of the Himalayan species by its conical form.



596



597



598



599



600



601



602



603



604

BULIMUS.

PLATE LXXX.

Species 572. (Mus. Benson.)

BULIMUS LATEBRICOLO. *Bul. testā acuminato-oblongā, compressē umbilicatā, unfractib; sex, subrotundatis, levibus, vel concentricē striatis; pellucido-stramineis, politā.*

THE LURCHER BULIMUS. Shell acuminate oblong, compressly umbilicated, whorls six in number, somewhat rounded, smooth, or concentrically striated; transparent straw-colour, polished.

BENSON, MS.

Hab. Landour, Western Himalaya.

A very delicately coloured transparent species, obtained by Mr. Benson in the lower Himalaya range, at elevations varying from 4,000 to 7,000 feet.

Species 588. (Mus. Pfeiffer.)

BULIMUS NITIDULUS. *Bul. testā subacuminato-oblongā, solidiusculā, compressē umbilicatā, unfractib; quinque ad sex, rotundatis, striatalis, politis, suturis constrictis, columellā latiusculā, expansā; cinnamonaceo-cornē.*

THE SHINING BULIMUS. Shell somewhat acuminate oblong, rather solid, compressly umbilicated, whorls five to six in number, rounded, finely striated, polished, sutures constricted, columella rather wide, expanded; cinnamon horn-colour.

PFEIFFER, Wiegman's Archives, 1839, vol. i. p. 352.

Hab. Matanzas, Island of Cuba.

A small shining cinnamon-brown shell.

Species 589. (Mus. Pfeiffer.)

BULIMUS ANTONT. *Bul. testā ovato-conicā, compressē humiliatā, unfractib; quinque ad sex, convexisculis, levibus, columellā subreflexā; pellucido-cornē.*

ANTON'S BULIMUS. Shell ovately conical, compressly umbilicated, whorls five to six in number, rather convex, smooth, columella slightly reflected; transparent horny.

PFEIFFER, Symbolæ, vol. i. p. 24.

Strobilus oblongus, Anton.

Hab. Island of Oparu.

An extremely small species, of a dull transparent horny substance.

Species 590. (Mus. Cuming.)

BULIMUS SCALARIOIDES. *Bul. testā subulatā, vix suboblique, unfractib; undecim, rotundatis, anguste crenulatissimis plicato-costatis, costis peculiariter acute compressis, aperturā parvā, rotundatā, labro simplici. sordide alba.*

THE SCALARIA-LIKE BULIMUS. Shell subulate, scarcely umbilicated, whorls eleven in number, rounded, very closely plicately ribbed throughout, the ribs being peculiarly sharply compressed, aperture small, rounded, lip simple; dull white.

Hab. Canary Islands?

Distinguished by the very sharply compressed structure of the ribs, which range longitudinally as in *Scalaria*.

Species 591. (Mus. Cuming.)

BULIMUS GLOMERATUS. *Bul. testā conico-subulatā, bandū umbilicatā, unfractib; nuc ea, rotundatis, levibus, suturis subconstrictis, aperturā suborbiculari, labro simplici; pellucido-cornē.*

THE COILED BULIMUS. Shell conically subulate, not umbilicated, whorls nine in number, rounded, smooth, sutures rather constricted, aperture nearly orbicular, lip simple; transparent horny.

Hab. — ?

Of a peculiar conical subulate form, of many whorls.

Species 592. (Mus. Benson.)

BULIMUS DECORTICATUS. *Bul. testā subulatā, solidiusculā, vix umbilicatā, unfractib; octo, concentricē subrude plicato-striatis, columellā tenui, reflexā, apertura parricula, oblongo-ovata, labro tenui; sordide cirecentie-cornē.*

THE DECORTICATED BULIMUS. Shell subulate, rather solid, scarcely umbilicated, whorls eight in number, concentrically rather rudely plicately striated, columella thin, reflected, aperture rather small, oblong-ovate, lip thin; dull greenish horn.

Hab. Macao, China; Dr. Cantor.

A shell of rather solid growth, longitudinally sculptured throughout with rude plicate striae.

Fig. 593.

The shell here figured, inadvertently, is the *B. Papoides*, already published at Pl. XXXVIII. Sp. 231.

For Sp. 593, see Pl. LXXXIV.

Species 594. (Mus. Pfeiffer.)

BULIMUS GRACILLIMUS. *Bal. testá subulatá, gracilis, haud umbilicatá, anfractibus septem ad octo, planulatis, medio leviter coarctatis, longitudinaliter acutè costatis et striatis, suturis impressis, columella subincrassatá, aperturá parvá, labro simplici; sordidè albá.*

THE VERY SLENDER BULIMUS. Shell subulate, slender, not umbilicated, whorls seven to eight in number, rather flattened, slightly contracted in the middle, longitudinally sharply ribbed and striated, sutures impressed, columella a little thickened, aperture small, lip simple: dead white.

PFEIFFER, Symbolae, vol. iii. p. 64.

Bulimus striato-costatus, D'Orbigny.

Hab. Cuba.

Originally described by Dr. Pfeiffer in Wiegman's Archives as an *Achatina*.

Species 595. (Mus. Benson.)

BULIMUS CLAVULINUS. *Bal. testá acuminato-oblongá, virginaliter umbilicatá, basia versus subventricosá, anfractibus sex,*

convexis, levibus, columellá tenué reflexá, apertura ovatá, labro tenui; pellucido-stramineá, vitroá.

THE LITTLE NAIL-LIKE BULIMUS. Shell acuminate oblong, scarcely umbilicated, rather ventricose towards the base, whorls six in number, convex, smooth, columella thinly reflected, aperture ovate, lip thin; transparent straw-colour, glassy.

POTIEZ and MICHAUD, Gal. de Douai, p. 136, pl. 14. f. 9, 10.
Hab. Moka, Isle of France (on woody banks); Benson.

If this shell is Potiez and Michaud's *B. clavulinus*, which I believe it is, Dr. Pfeiffer is wrong in quoting it as synonymous with *B. Goodalli*.

Species 596. (Mus. Pfeiffer.)


BULIMUS HARPA. *Bal. testá ovato-conicá, tenui, subventricosá, via umbilicatá, anfractibus quatuor, rotundatis, concentrici nitide costellato-striatis, aperturá rotundá, labro simplici; pellucido-brunneá.*

THE HARP BULIMUS. Shell ovately conical, thin, rather ventricose, slightly umbilicated, whorls four in number, rounded, concentrically neatly sculptured with rib-like striae, aperture rounded, lip simple: transparent brown.

PFEIFFER, Zeitschr. für Malac., 1847, p. 147.

Hab. ——?

A minute but extremely interesting species, partaking somewhat of the typical form of *Cyclostoma*.



BULIMUS.

PLATE LXXXI.

Species 597. (Mus. Taylor.)

BULIMUS RIVITATUS. *Bul. testā pyramidali-ovatā, basiū versus obliquā, umbilicatā, anfractibus octo, leevibus, ad suturas peculiariter crenulatis, aperturā obliquā ovatā, columellā labroque latē reflexis et effusis; albā, anfracta ultimā castaneo pallidō fasciata.*

THE TWO-FILLETED BULIMUS. Shell pyramidal-oval, oblique towards the base, umbilicated, whorls eight in number, smooth, peculiarly crenulated at the sutures, aperture obliquely oval, lip and columella broadly reflected and effused; white, last whorl faintly two-banded with chestnut.

SOWERBY, Conchological Illustrations, f. 46.
Hab. Brazil.

Effused in an oblique direction at the base, after the manner of the *B. auris-leporis*.

Species 598. (Mus. Taylor.)

BULIMUS MULIERIS. *Bul. testā acuminato-oblongā, um- bilicata, tenuicula, anfractibus septem, subrude sub-tilissimè striata, columellā reflexa, aperturā ovatā, labro simplici; albida, strigis angustis fasciis, hic illuc saturioribus, undique longitudinaliter picta.*

THE EFFEMINATE BULIMUS. Shell acuminate-oblong, umbilicated, rather thin, whorls seven in number, somewhat rudely very finely striated, columella reflected, aperture oval, lip simple; whitish, longitudinally painted throughout with narrow brown streaks which are here and there darker.

Hab. New Granada.

A light delicately painted species, finely rudely striated.

Species 599. (Mus. Cuming.)

BULIMUS BOLIVIANUS. *Bul. testā acuminato-oblongā, compressè umbilicatā, anfractibus septem, leevibus ant inanitissimè decussatis impresso-striatis, columellā contorto-plicata, talissimè depresso-dilatata, aperturā ovatā, labro simplici; carneo-albidā, purpureo-fusco fasciata, fasciis latīs, confertis, hic illuc interrupitis, columellā rubellā, apice rubro.*

THE BOLIVIAN BULIMUS. Shell acuminate-oblong, compressly umbilicated, whorls seven in number, smooth or very minutely decussated with impressed

striae, columella plicately twisted, very broadly depressedly dilated, aperture ovate, lip simple; flesh-white, banded with purple-brown, bands broad, close-set, here and there interrupted, columella reddish, apex red.

PFEIFFER, Pro. Zool. Soc., 1846, p. 34.

Hab. Near Merida, Andes of Bolivia.

The shell here represented is the original type of the species described by Dr. Pfeiffer. To that which I have mistaken for it at Pl. XLIV. Sp. 281, the name *B. annulatus* may be given. The species are truly distinct, as may readily be seen by an accurate comparison of the figures.

Species 600. (Mus. Taylor.)

BULIMUS PATRICIUS. *Bul. testā pyramidali-oblongā, sub- ampliter compressè umbilicata, anfractibus octo, obliquè subrude striatis, aperturā parvissimā, ovatā, columellā labroque latē reflexis: albida, rufescatè-purpureo sub- obscurè interrupitè fasciata, apertura fave roseo-pur- purea.*

THE PATRICIAN BULIMUS. Shell pyramidal-oblong, rather largely compressly umbilicated, whorls obliquely somewhat rudely striated, aperture rather small, oval, lip and columella broadly reflected; whitish, rather obscurely interruptedly banded with reddish purple, interior of the aperture rose-purple.

Hab. —?

This fine species is probably from New Granada, but I have no authority for stating it to be the locality.

Species 601. (Fig. a, b, c, Mus. Taylor.)

BULIMUS SPECTATU.S. *Bul. testā fusiformi-ovatā, eis nu- bilatā, tenuicula, anfractibus sex ad septem, longitu- dimulter subrude tenuistriatis, columellā reflexā, de- pressissimulā, aperturā ovatā, labro peculiariter effuso: albida vel cinereo-fuscenti varii striatā ant fasciatā, columellā et apertura fave roseo-purpureis.*

THE CHOICE BULIMUS. Shell fusiformly oval, scarcely umbilicated, rather thin, whorls six to seven in number, longitudinally somewhat rudely finely striated, columella reflected, rather depressed, aperture ovate, lip peculiarly effused; whitish or ash-brown, vari-

ously streaked or banded, columella and interior of the aperture rose-purple.

Hab. New Granada.

Except in the rose-purple hue of the columella and aperture, the pattern of colouring is too variable in this beautiful species to particularize. Its graceful effused growth is characteristic.

Species 602. (Mus. Taylor.)

BULIMUS TAYLORIANUS. *Bul. testā ventricoso-oblongā, tenui, inflatā, hand umbilicatā, aufractibus quinque, longitudinaliter obscurè plicato-striatis, ad sularus leviter crenulatis, sub leute undique minutissimè granulatis, aufractu ultimo radice maximo, peculiariter descendente, columellā transversā, reflexa, labro cix reflexo; castaneo-fuscā, semipellucidā, liris perparvis nigris apicem versus fulgoratā, columellā labroque roseo tingit.*

FAYLOE'S BULIMUS. Shell ventricosely oblong, thin, inflated, not unibiled, whorls five in number, longitudinally obscurely plicately striated, slightly crenulated at the sutures, very minutely granulated throughout beneath the lens, last whorl very much the largest, peculiarly descending, columella very thin, reflected, lip scarcely reflected; chestnut-brown, semipellucid, lightning-marked towards the apex with a few black lines, lip and columella tinged with rose.

Hab. Environs of Quito.

A delicate inflated species of a dark transparent brown colour, marked with dark zigzag lines near the apex, and having the thin lip and columellar edge tinged with rose.

Being indebted to Thomas Lombe Taylor, Esq., for the species, I have the pleasure to name it in honour of that distinguished collector.

Species 603. (Mus. Cuming.)

BULIMUS SINISTRALIS. *Bul. testā subacuminato-oblongā, crassiunculā, sinistrali, cix umbilicatā, aufracti-*

tibus septem, subrotundatis, obliquè striatis, columellā reflexā, aperturā parvinsculā, labro reflexo; luteā, olivaceo-viridi marmoratā, apertura fauce intensè fuscā, labro albo.

THE SINISTRAL BULIMUS. Shell somewhat acuminately oblong, rather thick, sinistral, scarcely unibiled, whorls seven in number, somewhat rounded, obliquely striated, columella reflected, aperture rather small, lip reflected; yellow, marbled with olive-green, interior of the aperture very dark brown, lip white.

Hab. Java.

Care should be taken not to confound this species with the green variety of *B. latus*, elevated to the rank of a species by Mousson, under the name *B. furcillatus*, from which it is certainly distinct.


Species 604. (Mus. Taylor.)

BULIMUS ALBIZONATUS. *Bul. testā subpyramidalis-conicā, basi depresso-angulatā, virg umbilicatā, aufractibus septem, plano-convexis, obliquè tenuestriatis, suturis peculiariter linearī-irregulari, aufractu ultimo basi angulato, columellā latē reflexa, aperturā parvinsculā, labro reflexo; intus extusque vărlescente-alba, zonulā augustā opaco-alba ad angulum aufractis ultimi.*

THE WHITE-ZONED BULIMUS. Shell somewhat pyramidal conical, rather depressed at the base, scarcely umbilicated, whorls seven in number, flatly convex, obliquely finely striated, sutures peculiarly linearly engraved, last whorl angled at the base, columella broadly reflected; bluish white within and without, having a narrow opaque white zone at the angle of the last whorl.

Hab. Ceylon.

A pale blue-white shell, remarkably characterized by a narrow opaque-white zone round the middle of the whorls, exactly in the place of the suture, concealed in all but the last whorl by the superposition of one whorl upon the other.



B U L I M U S.

PLATE LXXXII.

Species 605. (Mus. Taylor.)

BULIMUS TURBINATUS. *Bul. testā cylindraco-oblongā, vix umbilicatā, anfractibus septem, subrotundatis, longitudinaliter obscurè impresso-striatis, ad suturas subtilliter crenulatis et marginatis, columellā contortā, basi recedente, aperturā parvissimā, labro simplici; lutescente, castaneo maculatā et striatā, columellā livido-purpurascente.*

THE TURBINATED BULIMUS. Shell cylindrically oblong, scarcely umbilicated, whorls seven in number, somewhat rounded, longitudinally obscurely impressly striated, finely crenulated and margined at the sutures, columella twisted, receding at the base, aperture rather small, lip simple; yellowish, spotted and streaked with chestnut, columella livid-purple.

Achatina turbinata, Lca, Trans. Am. Phil. Soc., vol. ii. p. 31.
Hab. West Coast of Africa

I find this shell with the above name attached to it, but am not at all sure that it is the species. It is certainly distinct from the allied African forms.

Species 606. (Mus. Cuming.)

BULIMUS BIFLUGRATUS. *Bul. testā ventricoso-ovatā, tenuiculā, anfractibus sex, obliquè plicato-striatis, striis regularibus, labrum versus evanidis, columellā tenui, subcontortā, aperturā ovatā, labro simplici; fuscescente, intus cæruleo-viridi tincta, luceis flavidis, binis, oculi nudatis, undique fulgaratā.*

THE DOUBLE-WAVED BULIMUS. Shell ventricously ovate, rather thin, obliquely plicately striated, striae regular, fading towards the lip, columella thin, slightly twisted, aperture ovate, lip simple; light brown, stained with deep blue-green, lightning-marked throughout with yellow zigzag lines in pairs.

Hab. Andes of Columbia.

An interesting new species allied to *B. zebra*, but very distinctly characterized in its varieties of painting.

Species 607. (Mus. Cuming.)

BULIMUS PROXIMUS. *Bul. testā globoso-ovatā, ventricosissimā, haud umbilicatā, anfractibus sex, undique minutissimè granulosis, apicea versus obscurè radiativa costulatis, columellā subcontortā, basi recedente, labro sabinrassato, reflexo; fuscā, infra suturas pallide fasciatā, epidermide olivaceo-cornēa indutā, columellā labroque albī.*

THE ALLIED BULIMUS. Shell globosely ovate, very ventricose, not umbilicated, whorls six in number, very minutely granulated throughout, obscurely radially finely ribbed towards the apex, columella slightly twisted, receding at the base, lip rather thickened, reflected; brown, with a pale band beneath the sutures, covered with an olive horny epidermis, lip and columella white.

SOWERBY, Conch. Illustrations, f. 102.

Hab. Brazil.

Constantly distinguished by its broad rudely swollen form, and light band beneath the sutures.

Species 608. (Mus. Cuming.)

BULIMUS ADOPTUS. *Bul. testā subelongato-oblongā, crassiusculā, haud umbilicatā, anfractibus quinque ad sex, longitudinaliter subrude striatis, ad suturas subtilliter plicato-crenulatis, columellā incrassatā, peculiariter latè depressa, aperturā ovatā, labro crassatius reflexo; olivaceo-fuscā, apicea versus pallide flammatā.*

THE ADOPTED BULIMUS. Shell somewhat elongately oblong, rather thick, not umbilicated, whorls five to six in number, longitudinally somewhat rudely striated, finely plicately crenulated at the sutures, columella thickened, peculiarly broadly depressed, aperture ovate, lip thickly reflected; olive-brown, palely flamed towards the apex.

Hab. Banks of the Orinoco.

Allied in form to *B. Fauckii*, from which it may be recognised by its colouring and pattern of light flames.

Species 609. (Mus. Cuming.)

BULIMUS CALIGINOSUS. *Bul. testā oblongo-ovatā, haud umbilicatā, anfractibus sex, obliquè rude striatis, columellā latè reflexā, appressā, aperturā ovatā, labro simplici; cuneata vel albicans, ferrugineo-fusco irreguliter striato-lincta.*

THE DARKENED BULIMUS. Shell oblong-ovate, not umbilicated, whorls six in number, obliquely rudely striated, columella broadly reflected, appressed, aperture ovate, lip simple; ash-colour or whitish, irregularly stained in a streaked manner with rusty brown.

Hab. — ?

There is little of character to distinguish this species, although it is not referable to any hitherto described.



610



611.



612.



613.



614.



615.



616.



617.



618.



619.

BULIMUS.

PLATE LXXXIII.

Species 610. (Mus. Hamilton.)

BULIMUS HAMILTONI. *Bul. testá acuminato-turrítá, umbilicatá, spire suturis profunde impressis, anfractibus septem ad octo, subrotundatis, rudi plicato-striatis, aperturá pareá, columellá labroque simplicib; sordide albá, apice rubro.*

HAMILTON'S BULIMUS. Shell acuminately turreted, umbilicated, sutures of the spire deeply impressed, whorls seven to eight in number, somewhat rounded, rudely plicately striated, aperture small, lip and columella simple; dead white, apex red.

Hab. Near the Lake of Titicaca, Bolivia.

This interesting little species was collected by Mr. Pentland on the mountains in the environs of the Lake of Titicaca, which is 14,000 feet above the level of the sea. It may be distinguished by its characteristic red apex.

Species 611. (Mus. Hamilton.)

BULIMUS GIBBER. *Bul. testá oblongo-turrítá, sinistrali, compressé umbilicatá, anfractibus septem ad octo, convexus, rugoso-striatis, ad suturas subtiliter marginatis, columellá dilatata, aperturá pareá, labro pouatum reflexo; lacteá.*

THE GIBBOS BULIMUS. Shell oblong-turreted, sinistral, compressly umbilicated, whorls seven to eight in number, slightly convex, roughly striated, finely marginated at the sutures, columella dilated, aperture small, lip a little reflected; cream-colour.

KRYNICKI, Bull. Moscow, vol. vi. p. 416. pl. 3. f. 6.

Bulimus revolutus, Ziegler.

Hab. European Tartary.

The first three species of this plate are somewhat enlarged.

Species 612. (Mus. Bleher.)

BULIMUS GREGARIUS. *Bul. testá cylindraceo-turrítá, compressé umbilicatá, anfractibus octo, obliquè impresso-striatis, suturis impressis, columellá verticaliter dilatata, aperturá pareá, subquadrate-ovatá, labro sub-reflexo; pellucido-cornéa.*

THE ORDINARY BULIMUS. Shell cylindrically turreted, compressly umbilicated, whorls eight in number, ob-

liquely impressly striated, sutures impressed, columella vertically dilated, aperture small, somewhat squarely ovate, lip a little reflected; transparent horny.

ADAMS and REEVE, Moll. Voy. Samarang, pl. xiii. f. 12.
Hab. Sarawak, Borneo.

The oblique striae are extremely superficial, and the shell is of a very thin horny substance.

Fig. 613. (Mus. Hamilton.)

This interesting shell, with a delicate purple-coloured aperture, proves to be a variety of *B. strigatus*, Sp. 280. For Sp. 613, see Pl. LXXXV.

Species 614. (Mus. Hamilton.)

BULIMUS PENTLANDI. *Bul. testá acuminato-ovatá, basi subinflatá, vix umbilicatá, anfractibus sex, irregulariter rudi striatis, columellá reflexi, aperturá orbiculari-ovatá, labro simplici; sordide olivaceá, epidermide inductá.*

PENTLAND'S BULIMUS. Shell acuminately ovate, rather inflated at the base, scarcely umbilicated, whorls six in number, irregularly rudely striated, columella reflected, aperture orbicularly ovate, lip simple; dull olive, covered with an epidermis.

Hab. Near the Lake of Titicaca, Bolivia.

This and the *B. rubricatus* were the only species found by Mr. Pentland on the mountains in the vicinity of the Lake of Titicaca, at the same lofty elevation.

Species 615. (Mus. Taylor.)

BULIMUS FUCATUS. *Bul. testá acuminato-ovatá, basi subobliquè effuso, vix umbilicatá, anfractibus sex, obscurè striatis, ad suturas subcrenulatis, columellá reflexa, aperturá ovatá, labro effuso, vix reflexo; fuscescente, carmine-nigricante, subobscurè strigato-dimicti, columellá labroque pallidò rosæis.*

THE DYED BULIMUS. Shell acuminately ovate, rather obliquely effused at the base, scarcely umbilicated, whorls six in number, obscurely striated, slightly crenulated at the sutures, columella reflected, aperture ovate, lip effused, scarcely reflected; light brown.

somewhat obscurely stained with streaks of blue-black, lip and columella pale rose.

Hab. Sebundoi, New Granada.

Distinguished by its elegantly effused growth. The dark painting is somewhat obscure and has a mottled appearance.

Species 616. (Mus. Cuming.)

BULIMUS LETUS. *Bul. testā acuminato-oblongā, vix umbilicatā, anfractibus sex, levibus vel tenuistratis, columellā reflexā, appressā, aperturā ocatā, labro vix reflexā; castaneo-spadiceā, zonulis duabus angustis lateribus cingulatā, columellā labroque roseis.*

THE CHEERFUL BULIMUS. Shell acuminate oblong, scarcely umbilicated, whorls six in number, smooth or finely striated, columella reflected, appressed, aperture ovate, lip scarcely reflected; chestnut-fawn colour, encircled by two narrow yellowish-white zones, lip and columella rose-pink.

Hab. Sebundoi, New Granada.

From the same locality as the preceding species, and very closely allied; it differs, however, in form. Mr. Taylor has a specimen collected independently of this, of exactly similar pattern and colour.

Species 617. (Mus. Cuming.)

BULIMUS OCCULTUS. *Bul. testā cylindraceo-fusiformi, basi angustè compressā, subumbilicatā, anfractibus novem, plano-converxis, ad suturas superficialiter canaliculatis, aperturā quadrato-oblongā, tridentatā, labro tenui, effuso; pellucido-alba, ad umbilicatum atque pone labrum nigricante-castaneo-tinctā.*

THE CONCEALED BULIMUS. Shell cylindrically fusiform, narrowly compressed at the base, slightly umbilicated, whorls nine in number, flatly convex, superficially channelled at the sutures, aperture squarely oblong, three-toothed, lip thin, effused; transparent white, stained with dark chestnut at the umbilicus, and behind the lip.

Hab. Brazil.

Allied in form and general aspect to *B. Bahiensis*, but differing materially in the character of the aperture.

Species 618. (Mus. Taylor.)

BULIMUS QUINQUEDENTATUS. *Bul. testā cylindraceā, erassiusculā, basi compressā umbilicatā, anfractibus octo, plano-converxis, obliquè tenuistratis, aperturā parvā, coarctatā, quinque-vel sex-dentatā, labro plano-incassato; corneo-olivaceō, dentibus labroque albīs.*

THE FIVE-TOOTHED BULIMUS. Shell cylindric, rather thick, base compressly umbilicated, whorls eight in number, flatly convex, obliquely finely striated, aperture small, contracted, five- or six-toothed, lip flatly thickened; horny olive, teeth and lip white.

PFEIFFER, Symbolae, vol. i. p. 84.

Pupa quinquedentata, Mühlfeldt.

Hab. Dalmatia.

A shell of rather solid cylindrical growth, with the aperture much contracted.

Species 619. (Mus. Benson.)

BULIMUS PRETIOSUS. *Bul. testā cylindraceo-ovato, compressā umbilicatā, anfractibus levibus aut tenuistratis, columellā latē verticaliter dilatatā, aperturā parvissimā, suborbiculari, labro tenuē reflexā; pallide cornē, pellucida, striis opaco-albis numerosis nudata.*

THE PRECIOUS BULIMUS. Shell cylindrically ovate, compressly umbilicated, whorls smooth or finely striated, columella broadly vertically dilated, aperture rather small, nearly orbicular, lip thinly reflected; pale horny, transparent, waved with numerous opaque-white streaks.

CANTOR, MS.

Hab. Field of Chillianwalla, India (found on *Mimosa* bushes on the banks of the Jhelum); Cantor.

An extremely delicate and interesting species, allied to *B. rufistriatus*, from which it differs slightly in form, but most distinctly in pattern. Mr. Benson, to whom it was communicated by Dr. Cantor, informs me that it was collected during the late campaign of the British army in the Punjab, on the plain of Chillianwalla, the locality of Lord Gough's dearly purchased victory.



593



620



621



622



119



120



121



623



624



625

Small species.

Small species.

[All exceedingly minute.]

BULIMUS.

PLATE LXXXIV.

Species 593. (Mus. Adams.)

BULIMUS OCTONOIDES. *Bal. testā subelongatā, spirae suturis impressis, anfractibus septem, superne rotundatis et concentricè impresso-striatis, columellā tenuē reflexā, aperturā parvā, labro simplici; pallide stramineis, diaphanis.*

THE OCTONOIDS BULIMUS. Shell somewhat elongated, sutures of the spire impressed, whorls seven in number, rounded at the upper part and concentrically impressly striated, columella thinly reflected, aperture small, lip simple; light straw-colour, diaphanous.

ADAMS, Pro. Nat. Hist. Soc. Boston, 1845, p. 12.
Hab. Jamaica.

Named to denote its general resemblance with the *B. octonus*, which is referred to *Achatina*.

I am indebted for eight of the species of this plate to the kindness of Mr. C. B. Adams, Professor of Zoology, &c., in Amherst College, Massachusetts. They were collected during an excursion in Jamaica, in company with the Honourable Mr. Chitty, Chairman of Quarter Sessions in that Island, together with a number of interesting land shells of other genera, which Mr. Adams is preparing for publication in a periodical pamphlet form, under the title of "Contributions to Conchology."

Species 620. (Mus. Adams.)

BULIMUS TEREBELLA. *Bal. testā subulatā, anfractibus octo, convexiusculis, concentricè impresso-striatis, columellā tenuē reflexā, aperturā parvā, labro simplici; sordidè fuscescente.*

THE LITTLE AUGER BULIMUS. Shell subulate, whorls eight in number, rather convex, concentrically impressly striated, columella thinly reflected, aperture small, lip simple; dull brownish.

ADAMS, MS.
Hab. Jamaica.

This is the most elongated form of the Jamaica type, of light diaphanous growth, concentrically impressly striated. It is very minute, and, with the rest of the species of this plate, considerably magnified.

Species 621. (Mus. Cuming.)

BULIMUS GOODALLII. *Bal. testā conico-elongatā, vix umbilicatā, anfractibus septem, convexiusculis, levibus,*

December, 1849.

columellā salientiā, tenuē reflexā, apertura parvā, labro simplici; pallido-cornuei.

GOODALL'S BULIMUS. Shell conically elongated, scarcely umbilicated, whorls seven in number, slightly convex, smooth, columella somewhat twisted, thinly reflected, aperture small, lip simple; transparent horny.

Helix Goodallii, Miller, Ann. Phil. 1822, p. 381.

Bulinus Goodallii, Gray.

Helix (Cochlicella) clavulus, Féussac.

Bulinus clavulus, Potiez and Michaud.

Bulinus penitus, Pfeiffer.

Hab. West Indies.

This little species, being transported from Jamaica, has become partially naturalized in this country, and is included by some authors in the British fauna. It is found abundantly in the pine-pits at Bristol, especially on the sides of the pots, where they are immersed in the decayed bark.

Species 622. (Mus. Adams.)

BULIMUS MIRABILIS. *Bal. testā pyramidalis-conicā, anfractibus octo, rotundatis, superne squamis grandibus erectis nudique peculiariter coronatis, columellā contractā, aperturā parvā, medio contractā; carneo-fuscescente.*

THE WONDERFUL BULIMUS. Shell pyramidal conical, whorls eight in number, rounded, peculiarly coronated throughout with large erect scales, columella twisted, aperture small, contracted in the middle; light flesh-brown.

ADAMS, MS.
Hab. Jamaica.

The whorls in this remarkable species are contracted in the middle, and the upper portion is prominently coronated throughout with broad vaulted scales.

The three following *Bulinii* of Adams are numbered to belong to *Achatina*, under which genus the descriptions are given at Pl. XXII.

Species 623. (Mus. Adams.)

BULIMUS MINIMUS. *Bal. testā subulatā, spirae suturis subimpressis, anfractibus septem, subrotundatis, con-*

ventricē impresso-striatis, columellā tenuē reflexā, aper-turā parvissimulā, labro simplici; pallidē stramineā.

THE VERY SMALL BULIMUS. Shell subulate, sutures of the spire somewhat impressed, whorls seven in number, somewhat rounded, concentrically impressly striated, columella thinly reflected, aperture rather small, lip simple; light straw-colour.

ADAMS, MS.

Hab. Jamaica.

It may be as well to repeat that all the species of this plate are very considerably magnified. The *B. minimus* under consideration, the smallest of the genus at present known, is not half the size of a grain of corn.

Species 624. (Mns. Adams.)

BULIMUS PAUPERCULUS. *Bul. testā acuminato-oblongā, anfractibus quinque, levibus, supernē vix impresso-striatis, columellā tenuē reflexā, aperturā ovalā, labro simplici; sordide stramineā.*

THE MEAN-LOOKING BULIMUS. Shell acuminate oblong, whorls five in number, smooth, slightly impressly striated round the upper part, columella thinly reflected, aperture ovate, lip simple; dull straw-colour.

ADAMS, MS.

Hab. Jamaica.

It may be observed that there is a great similarity in the majority of these small *Bulimus* and *Achatinae* from Jamaica. Though specifically distinct, they are but slight modifications of one typical character,—a diaphanous shell of more or less elongated convolution, with the whorls concentrically impressly striated.

Species 625. (Mus. Benson.)

BULIMUS TUTULUS. *Bul. testā conicā, basi latiusculā, compressē umbilicatā, anfractibus sex, rotundatis, mi-nutissimā striatis, columellā latē verticaliter dilatatā, callositate parvā supra, aperturā ferè rotundā, labro subreflexo; pallidē castaneā, subcorneā, callositate albā.*

THE LITTLE TUFT BULIMUS. Shell conical, rather broad at the base, compressly umbilicated, whorls six in number, rounded, very minutely striated, columella broadly vertically dilated, with a small callosity above, aperture nearly round, lip slightly reflected; light chestnut, somewhat horny, callosity white.

BENSON, MS.

Hab. Humeerpore, Bundelkund, India; Benson.

This interesting little species is well distinguished by the broad convolution of the last whorl. Only two specimens were found.



BULIMUS.

PLATE LXXXV.

Species 613. (Mus. Cuming.)

BULIMUS LINOSTOMA. *Bul. testá fusiforme-ovatá, compressé umbilicata, tenuicula, unfractibus sex, levibus, vel sub lente minime striatis, columellá contortá, subplicata et appressa; aperturá obliqua dilatata, labro effuso; albida, flammis cerasescensibus et cinereis ornata, aperturá vixide violaceo-purpurea.*

THE THREADED BULIMUS. Shell fusiformly ovate, compressly umbilicated, rather thin, whorls six in number, smooth, or minutely striated beneath the lens, columella twisted, subplicate and appressed, aperture obliquely dilated, lip effused; whitish, ornamented with bluish and ash flames, aperture deep violet-purple.

Hab. Province of Santa Cruz, Bolivia.

M. D'Orbigny notices having found this delicately-coloured species in the woods of Santa Cruz, after rain, upon the ground as well as upon trees and shrubs, but in comparative scarcity.

Species 626. (Mus. Cuming.)

BULIMUS LIVIDUS. *Bul. testá acuminato-oblonga, tenui, via umbilicata, unfractibus sex ad septem, subrotundatis, sub lente creberriam inciso-striatis, columellá tenui reflexa, aperturá parviscalá, labro simplici; livido-olivaceá, nuculis perpansis subquadratris suturariibus infra suturas.*

THE LIVID BULIMUS. Shell acuminately oblong, thin, scarcely umbilicated, whorls six to seven in number, somewhat rounded, very closely engraved with striae, columella thinly reflected, aperture rather small, lip simple; livid olive, with a few somewhat square spots of a darker colour beneath the sutures.

Hab. Venezuela.

A thin shell, of a peculiar mottled livid colouring, marked somewhat irregularly with square spots beneath the sutures.

Species 627. (Mus. Cuming.)

BULIMUS HACHENSIS. *Bul. testá acuminato-oblonga, subpyramidalis, subumbilicata, unfractibus septem ad octo, concisis, levibus, nitidis, columellá tenui reflexa, ap-*

pressa, aperturá parviscalá, labro simplici; intus extusque roseo-intecta.

THE HACHA BULIMUS. Shell acuminately oblong, somewhat pyramidal, slightly umbilicated, whorls seven to eight in number, convex, smooth, shining, columella thinly reflected, appressed, aperture rather small, lip simple; pinkish yellow within and without.

Hab. Banks of the Rio Hacha, Guatemala.

A delicate yellow shell, of a peculiarly pyramidal form.

Species 628. (Mus. Cuming.)

BULIMUS REFLEXUS. *Bul. testá attenuato-ovalis, subinflatá, tenui, pellucida, unfractibus quinque ad sex, planiusculis, substriatis, columellá tenui, parum reflexa, appressa, aperturá oblonga, labro simplici; pallide cornéa, strigis irregularibus castaneis, interdum ramosis, ornata.*

THE REFLECTED BULIMUS. Shell attenuately ovate, somewhat inflated, thin, transparent, whorls five to six in number, rather flat, slightly striated, columella thin, but little reflected, appressed, aperture oblong, lip simple; pale horny, ornamented with chestnut streaks, sometimes branched.

PFEIFFER, PRO. ZOOL. SOC. 1842, p. 187.

Hab. Vicinity of Coquimbo (upon leaves of plants); Cuming.

An extremely thin, horny, inflated shell, belonging to the group of which *B. Broderipii* and *elegans* are typical examples.

Species 629. (Mus. Cuming.)

BULIMUS NUCINUS. *Bul. testá oblongo-ovata, vix umbilicata, crassiuscula, unfractibus quinque ad sex, convexis, minime granulosis, columellá subincrustatissima reflexa, appressa, aperturá ovata, labro crasso, simplici; sordidè albá, strigis perpansis rufo-fuscis ornata.*

THE NUT-TREE BULIMUS. Shell oblong-ovate, scarcely umbilicated, rather thick, whorls five to six in number, convex, minutely granulated, columella rather thickly reflected, appressed, aperture ovate, lip thick, simple; dull white, ornamented with a few red-brown streaks.

Hab. —?

A stout solid, dull-whitish shell, marked with red-brown longitudinal streaks, having a finely granulated surface. Distinct from *B. granulosus*, though agreeing with it in general aspect.

Species 630. (Mus. Cumiu.)

BULIMUS BALANOIDES. *Bul. testá cylindraceo-ovatá, solidiusculá, opice obtusá, anfractibus sex, plano-convexis, levibus, columellá incrassatá, reflexá, aperturá parviusculá, labro incrassato, reflexo; albida, fasciis castaneis duabus cingulatá, epidermide tenui hydrophaná induitá.*

THE DATE-LIKE BULIMUS. Shell cylindrically ovate, rather solid, obtuse at the apex, whorls six in number, flatly convex, smooth, columella thickened, reflected, aperture rather small, lip thickened, reflected; whitish, encircled by two chestnut bands, covered with a thin hydrophanous epidermis.

JONAS, Pro. Zool. Soc. 1842, p. 188. Phil. Abbild. und Besch. *Bul. pl. 6. f. 2.*

Hab. Island of Mindoro, Philippines.

Of solid growth, allied in form to *B. ovoides*.

Species 631. (Mus. Adams.)

BULIMUS IMMACULATUS. *Bul. testá acuminato-ovatá, vix umbilicata, anfractibus sex ad septem, planiusculis, sub lente minutè et creberrimè inciso-striatis, columellá incrassatá, aperturá parviusculá, labro vix reflexo; pallide straminea, castaneo-nigricante circa columellanam et aperturam marginem, epidermide tenui adustá basin versus indutá.*

THE UNSPOTTED BULIMUS. Shell acuminate-ovate, scarcely umbilicated, whorls six to seven in number, rather flat, beneath the lens minutely and very closely engraved with striae, columella thinly reflected, appressed, aperture ovate, lip simple; pale straw-colour, delicate rose about the umbilicus.

ADAMS, MSS.

Hab. Jamaica.

An interesting species, for which I am indebted to Mr. Adams, of Jamaica, allied to *B. stramineus*, but of rather stouter substance and more acuminate growth.

Species 632. (Mus. Metcalf.)

BULIMUS STRAMINEUS. *Bul. testá acuminato-ovatá, sub-inflatá, vix umbilicata, anfractibus sex, convexis, sub lente minutissimè et creberrimè inciso-striatis, columellá reflexa, appressa, aperturá ovatá, labro vix reflexo; pallidido-straminea, suturis indistinctè albinarginatis, apice nigra.*

THE STRAW-COLOURED BULIMUS. Shell acuminate-ovate, rather inflated, scarcely umbilicated, whorls six in number, convex, beneath the lens very minutely and closely engraved with striae, columella reflected, appressed, aperture ovate, lip slightly reflected; transparent straw-colour, sutures indistinctly edged with white, apex black.

GULDING, Trans. Liun. Soc. vol. xiv. pt. 2. p. 340.

Bulimus facetus, Menke.

Hab. Island of St. Vincent, West Indies.

A more transparent shell than the preceding, and of more inflated growth.

Species 633. (Mus. Cuming.)

BULIMUS SPRETUS. *Bul. testá pyramidali-conicá, solidiusculá, anfractibus sex ad septem, planio-convexis, sub lente minutissimè decussatim striatis, columellá incrassatá, aperturá parviusculá, labro vix reflexo; pallide straminea, castaneo-nigricante circa columellanam et aperturam marginem, epidermide tenui adustá basin versus indutá.*

THE DISREGARDED BULIMUS. Shell pyramidal-conical, rather solid, whorls six to seven in number, flatly convex, very minutely decussately striated beneath the lens, columella thickened, aperture rather small, lip slightly reflected; pale straw-colour, chestnut black round the umbilicus and edge of the aperture, covered towards the base with a thin burnt-brown epidermis.

Hab. Island of Romblon, Philippines; Cuming.

This species, of which Mr. Cuming collected several species, has been confounded with *B. cincinus*, from which it differs principally in being of a more truly conical form.



634.



635.



636.



637.



638.



639.



640.



641.



642.



643.

BULIMUS.

PLATE LXXVI.

Species 634. (Mus. Cuming.)

BULIMUS MONTANUS. *Bul. testā oblongo-cylindraceō, tenuiculā, compressē umbilicatā, anfractibus septem, obliquē subrude striatis et minutissimā granulatis, ad suturas obscurē crenulatis, columellā verticali, sublatē dilatatā, aperturā pareinclusa, labro subreflexo: pallidē castanea, semipelticida, suturis et apice albidis.*

THE MOUNTAIN BULIMUS. Shell oblong-cylindrical, rather thin, compressly umbilicated, whorls seven in number, obliquely somewhat rudely striated and very minutely granulated, obscurely crenulated at the sutures, columella vertical, rather broadly dilated, aperture rather small, lip slightly reflected; pale chestnut, semitransparent, sutures and apex whitish.

DRAPARNAUD, Hist. Moll. p. 74. pl. 4. f. 22.

Bulinus Lackhausenensis, Fleming.

Helix Lockhausenensis, Montagu.

Bulinus Montacuti, Jeffreys.

Bulinus Collini, Michaud.

**Helix buccinota*, Alten.

Hab. Britain, Germany, France, Spain.

A minutely granulated, delicate brown shell, varying a little in size in different localities.

Species 635. (Mus. Metcalfe.)

BULIMUS SEDUCTILIS. *Bul. testā oblongo-cylindraceō, sinistrali, compressē umbilicata, anfractibus decem, planisulcatis, densē convolutis, columellā verticali, dilatatā, aperturā parvā, bidentatā: carneō-albida.*

THE REMOTE BULIMUS. Shell oblong-cylindrical, sinistral, compressly umbilicated, whorls ten in number, rather flattened, thickly convoluted, columella vertical, dilated, aperture small, two-toothed; horn-white.

PFEIFFER, Symbolae, vol. i. p. 85.

Pupa seductilis, Ziegler.

Bulinus Niso, Pfeiffer.

Hab. Dalmatia, Austria.

A small, cylindrical, sinistral shell, remarkably closely convoluted.

Species 636. (Mus. Metcalfe.)

BULIMUS TRIDENS. *Bul. testā oblongā, compressē umbilicatā, anfractibus septem, planisulcatis, subtiliter stri-*

atis, columellā verticali, dilatatā, aperturā parvā, subquadratā, tridentatā, labro incrassato, subreflexo: carneō-fuscā.

THE THREE-TOOTHED BULIMUS. Shell oblong, compressly umbilicated, whorls seven in number, rather flattened, finely striated, columella vertical, dilated, aperture small, somewhat square, three-toothed, lip thickened, slightly reflexed; horny-brown.

BRUGUIERE, Enc. Méth. vol. i. p. 350.

Helix tridens, Müller.

Turbo tridens, Gmelin.

Pupa tridens, Draparnaud.

Chondrus tridens, Cuvier.

Jaanius tridens, Risso.

Bulinus variegatus, Hartman.

Hab. Germany, France, Austria, Sardinia.

Of a less cylindrical form than the preceding, of fewer whorls, and more horny substance.

Species 637. (Mus. Cuming.)

BULIMUS PUELLARIS. *Bul. testā acuminato-ovatā, tenuiculā, subdiaphanā, vix umbilicatā, anfractibus sex, convexis, levibus, sub lete crenatim et minutissimā inciso-striatis, columellā tenuē reflexā, aperturā ovatā, labro siuplici; extimū albā.*

THE MAIDEN BULIMUS. Shell acuminate-ovate, rather thin, somewhat diaphanous, scarcely umbilicated, whorls six in number, convex, smooth, beneath the lens very closely and minutely engraved with striae, columella thinly reflected, aperture ovate, lip simple: delicate white.

Hab. Brazil.

An elegantly convoluted semitransparent shell, not referable to any hitherto described species.

Species 638. (Mus. East India Company.)

BULIMUS HARRISII. *Bul. testā ovalā, subcentricosā, compressē umbilicatā, anfractibus septem, obliquē crenatim plicato-striatis, suturis subinpressis, columellā latē dilatatā, verticali, aperturā suborbiculari, labro simplici, vix reflexo; sordido albā, nitente.*

HARRIS'S BULIMUS. Shell ovate, somewhat ventricose, compressly umbilicated, whorls seven in number, obliquely very closely plicately striated, sutures

rather impressed, columella broadly dilated, vertical, aperture nearly orbicular, lip simple, scarcely reflected; dull white, shining.

Hab. Ankobar, Abyssinia.

Dedicated, at the request of Mr. Benson, to whom I am indebted for calling my attention to this shell in the museum of the East India Company, to the memory of its enterprising discoverer, Sir C. W. Harris.

Species 639. (Mus. Cuming.)

BULIMUS INUTILIS. *Bal. testā ovatā, tenuiculā, umbilicatā, aufractibus sex, subcentricosis, ruditē tenuistriatis, columellā dilatatā, verticali, aperturā ovatā, labro simplici; fuscescente.*

THE INSIGNIFICANT BULIMUS. Shell ovate, rather thin, umbilicated, whorls six in number, rather ventricose, rudely finely striated, columella dilated, vertical, aperture ovate, lip simple; light brown.

Hab. ——?

There is little to characterize this species, of which Mr. Cuming has several specimens, yet it is distinct from any previously described.

Species 640. (Mus. East India Company.)

BULIMUS EOUS. *Bal. testā cylindraceo-oblongā, obesiuscultā, compressè umbilicata, aufractibus sex, convexo-planis, columellā verticali, latè dilatatā, aperturā suborbiculari, productā, marginib⁹ lamina conjunctis; albā.*

THE EASTERN BULIMUS. Shell cylindrically oblong, rather stout, compressly umbilicated, whorls six in number, convexly flattened, columella vertical, broadly dilated, aperture nearly orbicular, produced, margin joined together by a lamina; white.

Hab. Hindoostan.

Collected by Dr. Griffiths, and very closely allied to the species named in honour of that gentleman represented at Sp. 362.

Species 641. (Mus. Cuming.)

BULIMUS GUINEENSIS. *Bal. testā fusiformi-conicā, tenui, vix umbilicatā, aufractibus septem, subtilissimè concen-trice striatis, striis peculiariter arcuatis, columellā ruditē contortā, tenuissimè reflexā, aperturā parvius-*

cūlā, labro simplici, supernē sinuato; pallidè vires-cente-stramineā.

THE GUINEA BULIMUS. Shell fusiformly conical, thin, scarcely umbilicated, whorls seven in number, very finely concentrically striated, striae peculiarly arched, columella much twisted, very thinly reflected, aperture rather small, lip simple, situated at the upper part: pale greenish straw-colour.

JONAS. Phil. Abbild. und Besch. Conch. vol. i. p. 54. *Bal.* pl. 1. f. 4.

Hab. Guinea.

A very delicate, thin shell, chiefly remarkable for the twisted growth of the columella, and for the sinuated lip, which causes the striae to be formed in curves.

Species 642. (Mus. Cuming.)

BULIMUS TEPIDULUS. *Bal. testā subpyramidalis, compresse umbilicata, aufractibus septem, convexis, levibus, suturis impressis, columellā verticali, aperturā parvā, labro simplici; sordide albā, apicea versus fuscescente.*

THE COOL BULIMUS. Shell somewhat pyramidal, compressly umbilicated, whorls seven in number, convex, smooth, sutures impressed, columella vertical, aperture small, lip simple; dull white, light brown towards the apex.

Hab. Canary Islands.

This does not agree with any of the species from the Canary Islands described by Webb and Berthelot.

Species 643. (Mus. Cuming.)

BULIMUS CONFINIS. *Bal. testā ovatā, subcentrica, profunde umbilicata, aufractibus sex, rotundatis, tenui striatis, columellā latè dilatatā, subreflexā, aperturā orbiculari, labro simplici; pellucido-corneā, opaco-albidō variegata.*

THE BORDERER BULIMUS. Shell ovate, rather ventricose, deeply umbilicated, whorls six in number, rounded, finely striated, columella broadly dilated, a little reflected, aperture orbicular, lip simple; transparent horny, variegated with opaque whitish.

Hab. Texas, United States.

The locality of this species may be regarded as about the northern limit of the genus in the new world.



644



645



646



647



648



649



650



651



652



653

BULIMUS.

PLATE LXXXVII.

Species 644. (Mus. Metcalfe.)

BULIMUS FOLLICULUS. *Bul. testā abbreviato-cylindraceā, compressē umbilicatā, apice acuminatā, aufractibus rotundatis, tumidis, subtilissimè costatis, suturis impressis, columellā verticali, dilatatā, aperturā subcirculari, marginibus lamina callosa conjunctis; cornue-fuscescente.*

THE WHEAT-HUSK BULIMUS. Shell abbreviate cylindric, compressly umbilicate, acuminate at the apex, whorls rounded, swollen, very finely ribbed, suture impressed, columella vertical, dilated, aperture nearly round, with the margins joined by a callous lamina; horny brown.

PFEIFFER, *Symbola*, vol. iii. p. 83.
Carychium costatum, Hutton.

Hab. Bengal.

Partaking somewhat of the typical form of *B. Gossei* of the West Indies.

Species 645. (Mus. Metcalfe.)

BULIMUS LUBRICUS. *Bul. testā oblonga-ovatā, apice oblusincurvā, aufractibus sex, convexis, levibus, nitetibus, columellā arcuatā, callosā, vix obsoletē truncatā, aperturā parvā, labro simplici, intus subincrassato; fuscescente, vitrea.*

THE SLIPPERY BULIMUS. Shell oblong-ovate, rather obtuse at the apex, whorls six in number, convex, smooth, shining, columella arched, callous, slightly obsoletely truncated, aperture small, lip simple, slightly thickened within; brownish, glassy.

BRUGUIERE, Enc. Méth. vol. i. p. 311. no. 23.

Helix lubrica, Müller.
Helix subcylinndrica, Chemnitz.
Turbo glaber, Da Costa.
Achatina lubrica, Michaud.
Zoa lubrica, Leach.
Cionella lubrica, Jeffreys.
Columna lubrica, Jan.
Styloides lubricus, Fizing.
Bulinus nitens, Schmidt.

Hab. Europe and the British Isles. Massachusetts, North America: Gould.

This is the most abnormal species of the genus, and affects a colder latitude by several degrees than any other. The

shell is distinguished by its shining glassy composition and by an inclination in the columella to be truncated, partaking in these respects of the character of the small glassy *Achatinae*. Dr. Gould describes having found it in great abundance in the woods of Oak Island, Massachusetts, after a warm October rain, clinging to the fallen leaves, from which they disappeared as the moisture evaporated. It inhabits the British Isles throughout, and has a wide range in Spain, Portugal, Italy, Switzerland, France, and Germany, reaching as far north as Sweden.

Species 646. (Mus. Cuming.)

BULIMUS OPARANUS. *Bul. testā subulatā, vix umbilicatā, spire suturis impressis, aufractibus septem, rotundatis, concentricè striatis, columellā verticaliter tenuè dilatatā, aperturā parvā, superne sinnatā, labro simplici: pellucido-vitrea, straminā.*

THE OPARA BULIMUS. Shell subulate, scarcely umbilicate, suture of the spire impressed, whorls seven in number, rounded, concentrically striated, columella vertically thinly dilated, aperture small, situated at the upper part, lip simple; transparent glassy, straw-colour.

PFEIFFER, Pro. Zool. Soc., 1846, p. 34.

Hab. Island of Opara, Society group (at the roots of plants); Cuming.

A thin, glassy, subulate shell, delicately concentrically striated after the manner of those from Jamaica represented in Pl. 84.

Species 647. (Mus. Metcalfe.)

BULIMUS OBSCURUS. *Bul. testā oblongā, compressē umbilicatā, aufractibus septem, convexis, transversaliter, oblique striatis, columellā verticaliter dilatatā, aperturā parvā, subquadrate-ovatā, obliquā, labro subeffuso: castaneo-fuscescente.*

THE OBSCURE BULIMUS. Shell oblong, compressly umbilicate, whorls seven in number, convex, rather swollen, obliquely striated, columella vertically dilated, aperture small, somewhat squarely ovate, oblique, lip rather effused; chestnut-brown.

DRAPARNAUD, Hist. Moll. p. 74. pl. 4. f. 23.

Helix obscura, Müller.
Bulinus hordeaceus, Bruguiere.

- Turbo rupina*, Da Costa.
Eua obscura, Leach.
Merdigera obscura, Held.
Hab. Europe and the British Isles.

Allied to *B. montanus*, from which it differs in form, and is uniformly of smaller size.

Species 648. (Mus. Cuming.)

BULIMUS LINEARIS. *Bul. testā subulatā, minutiū compresē umbilicatā, anfractibus decem, plano-converxis, subtilissimè arcuatim striatis, lineaē elevatā superiē marginatis, columellā brevi, tenuē reflexā, aperturā parvā, labro simplici; pellucido-corneā, stramineā.*

THE LINEAR BULIMUS. Shell subulate, minutely compressly umbilicated, whorls ten in number, flatly convex, very finely arcuately striated, margined round the upper part with a raised line, columella short, thinly reflected, aperture small, lip simple; transparent horny, straw-colour.

KRAUSS, Sudafrik. Moll. p. 78. pl. 5. f. 3.
Hab. Port Natal, South Africa (near the river Limpopo); Wahlberg.

A delicate glassy shell, in which the suture is distinguished throughout by a fine elevated ridge.

Fig. 649 proves to be a variety of *B. subula*, Sp. 494.

Species 650. (Mus. Metcalfe.)

BULIMUS QUADRIDIENS. *Bul. testā cylindraceo-oblongā, sinistrati, compresē umbilicatā, anfractibus octo, plano-converxis, obliquè striatis, aperturā subquadratā, quadridentata, columellā labroco effusis; olivaceo-corneā, labro albido.*

THE FOUR-TOOTHED BULIMUS. Shell cylindrically oblong, sinistral, compressly umbilicated, whorls eight in number, flatly convex, obliquely striated, aperture somewhat square, four-toothed, lip and columella effused; olive horny, lip whitish.

BRUGUIERE, Enc. Méth. vol. i. p. 351. no. 91.

Helix quadridens, Müller.
Pupa quadridens, Draparnaud.
Chondrus quadridens, Cuvier.
Janilia heterostrophia, Risso.
Gonodon quadridens, Held.
Torquula quadridens, Villa.
Hab. Frauec, Switzerland, Italy, Spain.

This species partakes very much of the character of *Pupa*, and may, with equal propriety, be referred to that genus.

Species 651. (Mus. Benson.)

BULIMUS SIKKIMENSIS. *Bul. testā ovatā, subinflatā, spirā breviusculā, apice obtusa, anfractibus quatuor, ultimo multo maximo, concentricè tenuē striatis, suturis impressis, columellā subcontortā, tenuē reflexā, aperturā oblongā, superne leviter sinuatā; pallide olivaceā, semi-pellucida.*

THE SIKKIM BULIMUS. Shell ovate, somewhat inflated, spine rather short, obtuse at the apex, whorls four in number, the last much the largest, concentrically finely striated, sutures impressed, columella slightly twisted, thinly reflected, aperture oblong, slightly sinuated at the upper part; pale olive, semitransparent.

Hab. Darjeeling, Sikkim-Himalaya.

An interesting species from the collection of Mr. Benson, of a different type from those inhabiting the western Himalaya range.

Species 652. (Mus. Cuming.)

BULIMUS TERRIFORMIS. *Bul. testā subulatā, subumbilicatā, anfractibus octo, converxis, subtilissimè concentricè striatis, columellā tenuē reflexā, aperturā parvā, labro simplici; pellucido-corneā, stramineā.*

THE TURRETED BULIMUS. Shell subulate, slightly umbilicated, whorls eight in number, convex, very finely concentrically striated, columella thinly reflected, aperture small, lip simple; transparent horny, straw-colour.

KRAUSS, Sudafrik. Moll. p. 78. pl. 5. f. 2.
Hab. Natal; Wahlberg.

Very closely allied to *B. Oparanus* from the Society Islands.

Species 653. (Mus. Benson.)

BULIMUS SOLITARIUS. *Bul. testā globose-conicā, umbilicatā, anfractibus sex, rotundatis, obliquè striatis, columellā brevi, parvā dilatatis, aperturā suborbiculari, labro simplici; albido, fusco, fasciatā ant variegatā.*

THE SOLITARY BULIMUS. Shell globosely conical, umbilicated, whorls six in number, rounded, obliquely striated, columella short, but little dilated, aperture nearly orbicular, lip simple; whitish, banded or variegated with brown.

PFEIFFER, Symbolae, vol. ii. p. 122.

Helix solitarius, Poiret.

Helix couoidea, Draparnaud.

Bulimus couoidens, Jan.

Hab. Southern Europe.

The most trochiform of any species of the genus.



B U L I M U S .

PLATE LXXXVIII.

Species 654. (Mus. Cuming.)

BULIMUS EXIGUUS. *Bul. testá oblongá, compressè umbilicatá, anfractibus quinque, subrotundatis, columellá verticaliter dilatatá, aperturá parvá, rotundatá, labro subreflexo; castaneo-cornéa.*

THE PUNY BULIMUS. Shell oblong, compressly umbilicated, whorls five in number, somewhat rounded, columella vertically dilated, aperture small, rounded, lip slightly reflexed; chestnut horny.

Hab. St. Domingo; Sallé.

A little dark horny species, approaching somewhat in general aspect to *B. crenipictus* of Hindooostan.

Species 655. (Mus. Beleher.)

BULIMUS MEIACOSHIMENSIS. *Bul. testá subpyramidalis oblongi, vix umbilicata, anfractibus decem, subangustis, columellá verticaliter reflexa, aperturá rotundatá; pellucido-cornéa.*

THE MEIACOSHIMA BULIMUS. Shell somewhat pyramidal oblong, scarcely umbilicated, whorls ten in number, rather narrow, columella vertically reflected, aperture rounded; transparent horny.

ADAMS and REEVE, Moll. Voy. Samarang, PL 14. f. 5.

Hab. Islands of Ty-pin-sau and Koo-Kien-san of the Meiacoshima group, Yellow Sea.

A small thin horny species, found by Mr. Arthur Adams on decayed leaves among the loose stones surrounding the tombs in the above-named islands.

Species 656. (Mus. Cuming.)

BULIMUS CONCENTRICUS. *Bul. testá subacuminato-oblonga, vix umbilicata, anfractibus septem, subcentricosis, concentricè tenuistratis, columellá tenué reflexa, aperturá subangulata, supernè sinuata; pellucido-straminea.*

THE CONCENTRIC BULIMUS. Shell somewhat acuminately oblong, scarcely umbilicated, whorls seven in number, rather ventricose, concentrically finely striated, columella thinly reflected, aperture rather large, situated at the upper part; transparent straw-colour.

Hab. Liberia, West Africa.

An extremely delicate inflated species, of the *Glandina* type, allied to *B. Guineensis*.

Species 148. (Mus. Metcalfe.)

BULIMUS HENNAHII. *Bul. testá acuminato-ovatá, umbilicata, anfractibus sex ad septem, rotundatis, rugoso-striatis, columellá reflexa, aperturá ovatá, labro simplici; albida, linea purpureo-nigricantibus longitudinaliter irregulariter notata, aperturæ fauce purpureo-nigricante, apice roseo.*

HENNAH'S BULIMUS. Shell acuminately ovate, umbilicated, whorls six to seven in number, rounded, roughly striated, columella reflected, aperture ovate, lip simple; whitish, irregularly longitudinally marked with purple-black lines, apex pink.

GRAY, Spicilegia Zoologica, part 1. p. 5. pl. 5. f. 1.
Bulinus lychnorum, Sowerby.
Bulinus eactornus, D'Orbigny.

Hab. Peru.

To this species, which is extremely variable in colour, must be added *B. erubescens*, mihi, Sp. 148, of which it is a variety.

Species 657. (Fig. a, b, c, Mus. Cuming.)

BULIMUS SALLEANUS. *Bul. testá subulata, solidiuscula, anfractibus undecim, planulatis, concentricè tenuissimè striatis, columellá reflexa, aperturá parvá, basi subflexuosa, labro simplici; albicans, epidermide tenui indutâ.*

SALLE BULIMUS. Shell subulate, rather solid, whorls eleven in number, flattened, concentrically very finely striated, columella reflected, aperture small, slightly flexuous at the base, lip simple; transparent white, covered with a thin epidermis.

Hab. St. Domingo; Sallé.

An interesting, sharply subulate, solid, ground species.

Species 658. (Mus. Miers.)

BULIMUS CONTORTPLICATUS. *Bul. testá oblongo-ovata, subcentricos, anfractibus quinque, convexis, minutissimè indentatis, columellá subincrassata, uniplicata, aperturæ oblongo-ovata, labro subincrassato, reflexo; albida, partis rosaceæ, epidermide tenui cornæ tute-olivaceæ indutâ, aperturæ fauce alba.*

THE TWISTED-PLAITED BULIMUS. Shell oblong-ovate, rather ventricose, whorls five in number, convex.

very minutely indented, columella rather thickened, reflected; whitish, partially tinged with pink, covered with a thin horny olive-yellow epidermis, interior of the aperture white.

Hab. Minas Geraes, Brazil; Miers.

This fine species is nearest allied to *B. pudicus*, from which the plicate growth of the columella differs materially in form; whilst the shell is of a more oblong growth.

Species 659. (Mus. Cuming.)

BULIMUS DOMINICUS. *Bul. testá ovatá, tenui, inflatá, anfractibus quinque, sub lente miuntissimè decussatim striatis, columellá tenui reflexá et appressá, aperturá ovatá, labro simplici; peltacido-corneá, fasciá castaneá interruptá cingulatá, anfractu ultimo trifasciato.*

THE DOMINGO BULIMUS. Shell ovate, thin, inflated, whorls five in number, very minutely decussately striated beneath the lens, columella thinly reflected and appressed, aperture ovate, lip simple; transparent horny, encircled with an interrupted chestnut band, last whorl three-banded.

Hab. St. Domingo; Sallé.

Resembling *B. Humboldtii*, but much smaller and more transparent, and of a different form.

Species 660. (Mus. Cuming.)

BULIMUS LIBERIANUS. *Bul. testá oblongo-ovatá, tumiduscílata, compressè umbilicatá, apice obtusá, anfractibus sex, convexis, ad suturas tenuimarginatis, obliquè subtíliter et creberrimè elevato-striatis, columellá latè verticaliter reflexá, aperturá obliquè subquadratá: pelliculo-albá, innaculatá.*

THE LIBERIAN BULIMUS. Shell oblong-ovate, rather swollen, compressly umbilicated, obtuse at the apex, whorls six in number, convex, thinly margined at the sutures, obliquely finely and very closely sculptured with raised striae, columella broadly rather vertically reflected, aperture obliquely square-formed; transparent white, unspotted.

GOULD, MSS.

Hab. Liberia, West Africa.

This very interesting species belongs to the same type as the *Pupa grandis* of the islands of Socotra and Madagascar, and, bordering on the confines of the two genera, may be referred with equal propriety to either.



B U L I M U S.

PLATE LXXXIX.

Sp. 367. (Mus. Cuming.)

BULIMUS INTERSTINCTUS. Having been favoured with finer specimens of this very interesting species, collected by Dr. Perkins at Cape Palmas, West Africa, I avail myself of the opportunity of figuring two characteristic varieties.

Species 661. (Mus. Cuming.)

BULIMUS KELLETTII. *Bul. testā ovatā, subventricosā, anfractibus quinque, conexis, rugoso-striatis, ad suturas subplicato-crenulatis, anfractu ultimo multo maximo, columellā incrassatā, peculiariter contortā, aperturā subeffusā, labro incrassato, reflexo; carulescente-griseā saturata quadrisciatā, strigis angustis albis interrumpitis et undatis longitudinaliter fulguratis, columellā ei apertura fuce iridescente-purpureis.*

KELLETT'S BULIMUS. Shell ovate, rather ventricose, whorls five in number, convex, roughly striated, somewhat plicately crenulated at the sutures, last whorl much the largest, columella thickened, peculiarly twisted, aperture rather effused, lip thickened, reflected: bluish-grey, encircled with four bands of a darker colour, and longitudinally lightning-marked with narrow waved and interrupted white streaks.

Hab. Ecuador?

I have much pleasure in naming this new and very beautiful species after the zealous naturalist, Capt. Kellett, of H. M. S. Herald, by whom it was procured in Central America, and is probably from Ecuador.

Species 212. (Mus. Miers.)

BULIMUS OVATUS. An interesting white-lip dwarf variety, from Bahia, described by M. Sowerby in his 'Conchological Illustrations' under the name *B. auritus*.

Species 662. (Fig. *a* and *b*. Mus. Cuming.)

BULIMUS TORRIDUS. *Bul. testā oblongo-tumidā, apice subpapillari, anfractibus septem, convexis, obliquè elevato-striatis, ad suturas subtiliter et creberrimè crenulatis, columellā simplici, aperturā ovatā, labro porām reflexo; ustulato-fuscā, epidermide olivaceō indutā, aperturā fuce intense fuscā.*

THE TORRID BULIMUS. Shell oblong-tumid, somewhat papillary at the apex, whorls seven in number, convex, obliquely elevately striated, finely and very closely crenulated at the sutures, columella simple, aperture ovate, lip but little reflected; scorched-brown, covered with an olive epidermis, interior of the aperture dark brown.

GOULD, MSS.

Hab. Liberia, West Africa.

The African species received from Dr. Gould, of Boston, United States, just at the close of this monograph, will probably be described, if not so already, in the Proceedings of the Natural History Society of that city. They are of an extremely interesting character, and were collected by Dr. George A. Perkins, a zealous American missionary, for many years resident at Cape Palmas.

February, 1850.

B U L I M U S.

	Plate.	Species.		Plate.	Species
<i>abbreviatus</i> , Jan.	LIX.	399	<i>auratus</i> , <i>Pfr.</i>	L.I.	335
<i>Abyssinicus</i> , Ruppell	XLVI.	296	<i>aureus</i> , Swainson	XXIX.	187
<i>abyssorum</i> , <i>D'Orb.</i>	XXXVI.	213	<i>auripigmentum</i> , <i>Reeve</i>	XXIX.	178
<i>Achatinaeus</i> , <i>Pfr.</i>	LXVII.	470	<i>auritus</i> , Sow.	XXXVI.	212
<i>Achatinoides</i> , Ziegler	L.	327	<i>auris-bovina</i> (<i>Auric.</i>), Lam.	XXX.	185
<i>acuminatus</i> (<i>Helix</i>), Sow.	XI.	60	<i>auris-leporis</i> (<i>Auric.</i>), Lam.	XLI.	259
<i>acutus</i> (<i>Helix</i>), Muller	LXVI.	467	<i>auris-Midae</i> , <i>Chenn.</i>	XXVIII.	170
<i>Adamsii</i> , <i>Reeve</i>	XIII.	73	<i>auris-muris</i> (<i>Helix</i>), Mor.	XLI.	261
<i>Adamsoni</i> , Gray	XXVI.	165	<i>auris-Silene</i> , <i>Born.</i>	XXV.	164
<i>Adamsoni</i> , <i>Pfr.</i>	L.	327	<i>auris-vulpina</i> , <i>Chenn.</i>	XXX.	180
<i>adoptus</i> , <i>Reeve</i>	LXXXII.	608	<i>Babauchi</i> , Anton.	LXIII.	431
<i>Ægle</i> , Brod.	X.	49	<i>baeterionides</i> , <i>D'Orb.</i>	LXVIII.	482
<i>ægots</i> , Menke	LIV.	357	<i>badiosa</i> (<i>Helix</i>), Fér.	LXVI.	469
<i>affinis</i> , Brod.	XXIII.	154	<i>badius</i> , Sow.	XXXIX.	235
<i>Africanus</i> , <i>Reeve</i>	L.	330	<i>baraticatus</i> (<i>Helix</i>), Fér.	LXX.	505
<i>albatus</i> (<i>Helix</i>), Fér.	LI.	333	<i>Bahiensis</i> , <i>Moric.</i>	LXIV.	442
<i>Alberti</i> , Brod.	V.	22	<i>Bairdii</i> , <i>Reeve</i>	XLIII.	272
<i>albicans</i> , Brod.	XXII.	141	<i>balanoides</i> , <i>Jonas</i>	LXXXV.	630
<i>albizonatus</i> , <i>Reeve</i>	LXXXI.	604	<i>Bomboncha</i> , W. & B.	LXIX.	493
<i>allus</i> , Sow.	XIII.	75	<i>barbarus</i> (<i>Helix</i>), Linn.	LXXV.	539
<i>Alipi</i> (<i>Helix</i>), Féussac	LX.	413	<i>Beckianus</i> , <i>Pfr.</i>	LXXVIII.	480
<i>Alto-Peruvianus</i> , <i>Reeve</i>	LXXII.	521	<i>bellulus</i> , <i>Jonas</i>	XXIV.	160
<i>alternans</i> , Beck.	XXIII.	152	<i>Bengaleensis</i> , <i>Lam.</i>	XLV.	289
<i>alutaceus</i> , <i>Reeve</i>	LXXII.	522	<i>Benguelensis</i> , <i>Pfr.</i>	XLVI.	298
<i>ambustus</i> , <i>Reeve</i>	LXXIV.	535	<i>Bensonii</i> , <i>Reeve</i>	LXXVIII.	571
<i>Andicola</i> , <i>Pfr.</i>	LV.	364	<i>Bergeri</i> , <i>Pfr.</i>	LXVIII.	489
<i>angistoma</i> , Wagner	LXVIII.	312	<i>Bertheloti</i> , <i>ib.</i>	LXXV.	541
<i>Angosturensis</i> , Grinner	XLVII.	307	<i>biciinctus</i> , Recluz	LI.	333
<i>Angosturensis</i> , Grinner	LIX.	402	<i>bicolor</i> , <i>Sow.</i>	XLIV.	276
<i>angulatus</i> , Wagn.	XL.	248	<i>bidens</i> , Krynicki	LXVI.	465
Antoni, <i>Pfr.</i>	LXXX.	589	<i>bicoloratus</i> , <i>Lea</i>	V.	22
<i>aplomorphus</i> , <i>Jonas</i>	X.	47	<i>bifasciatus</i> , Philippi	XXX.	182
<i>apodemetes</i> , <i>D'Orb.</i>	XXII.	142	<i>bilabiatus</i> , Brod. & Sow.	XXXIII.	201
<i>appendiculatus</i> , P. & M.	XLVI.	465	<i>bifulgaratus</i> , <i>Reeve</i>	LXXXII.	606
<i>aquilus</i> , <i>Reeve</i>	XXII.	138	<i>bilineatus</i> , <i>Sow.</i>	XXI.	132
<i>arcuatus</i> , <i>Pfr.</i>	LXVII.	478	<i>bivittatus</i> , <i>Sow.</i>	LXXXI.	597
<i>articulatus</i> , Turton	LXIV.	444	<i>Blainvilleanus</i> , <i>Pfr.</i>	XXV.	161
<i>astrapoides</i> , <i>Jonas</i>	XXXI.	159	<i>Boholensis</i> , Brod.	VIII.	37
<i>atomatus</i> , Gray	XXX.	184	<i>Boissieri</i> , <i>Moric.</i>	LXIV.	449
<i>atrapoides</i> , <i>Jonas</i>	XXIV.	157	<i>Boliviensis</i> , <i>Pfr.</i>	XLIV.	281
<i>atricallosus</i> , Gould	XXXI.	188	<i>Boliviensis</i> , <i>Pfr.</i>	LXXXI.	599

INDEX.

	Plate.	Species.		Plate.	Species.
Boutre (<i>Helix</i>), Chemn.	XLVIII.	315	<i>ciactus</i> , Jay	XXIX.	179
Boysiana, <i>Benson</i>	LXXVIII.	575	<i>cinerous</i> , <i>Reeve</i>	LVI.	372
breviculus, <i>Pfr.</i>	XII.	65	<i>cinnamomeo-lucens</i> , <i>Moric</i>	XLVI.	293
Bridgei, <i>Pfr.</i>	XIX.	117	<i>citrino-vitrea</i> , <i>Moric</i>	XLVI.	290
Broderipii, <i>Sow.</i>	XVI.	97	<i>citrinus</i> , <i>Brug.</i>	XXXI.	187
Browni, <i>Pfr.</i>	XXXVI.	212	<i>Clausiliaforata</i> , <i>Menke</i>	LXVI.	465
bulbula, <i>Brod.</i>	XII.	68	<i>Clausilioides</i> , <i>Reeve</i>	LXXIII.	533
Burchellii, <i>Gray</i>	LXXVI.	548	<i>clava</i> , <i>Reeve</i>	LXX.	500
Buschii, <i>Pfr.</i>	XIX.	112	<i>elevator</i> , <i>Petit</i>	LII.	345
Cacticulus, <i>Reeve</i>	LVIII.	393	<i>clavulinus</i> , <i>Potiez</i>	LXXX.	595
cacticinus, <i>Brod.</i>	XVIII.	103	<i>clavulinus</i> , <i>Pot. & Mich.</i>	LXXXIV.	621
cactorius, D'Orb.	LXXXVIII.	148	<i>coagulatus</i> , <i>Reeve</i>	LXXVII.	558
calcareus, <i>Brug.</i>	LII.	344	<i>coarctatus</i> , <i>Pfr.</i>	XLI.	260
calcareus (<i>Helix</i>), Born.	LII.	342	<i>cochleatus</i> , <i>Pfr.</i>	XIV.	82
calcareus, Parryss.	LXVI.	464	<i>colebs</i> , <i>Benson</i>	XLVII.	301
Caledonicus, <i>Petit</i>	XXV.	163	<i>conopictus</i> (<i>Pupa</i>), <i>Hutton</i>	LXIX.	492
Californicus, <i>Reeve</i>	LVI.	378	<i>Collini</i> , <i>Michaud</i>	LXXXVI.	634
caliginosus, <i>Reeve</i>	LXXXII.	609	<i>coloratus</i> , <i>Nyst.</i>	XXIV.	155
Calista, <i>Brod.</i>	VIII.	38	<i>columella</i> , <i>Philippi</i>	LXX.	502
calobaptus, Jonas	IX.	43	<i>columellaris</i> , <i>Reeve</i>	LXXXIII.	528
calvus, <i>Sow.</i>	XX.	126	<i>compactus</i> , <i>Friedelsky</i>	LXXV.	546
Calypso, <i>Brod.</i>	VII.	31	<i>concentricus</i> , <i>Reeve</i>	LXXXVIII.	656
camelopardalis, <i>Brod.</i>	VIII.	36	<i>concininus</i> (<i>Helix</i>), <i>Sow.</i>	X.	52
canaliculatus, <i>Pfr.</i>	XII.	256	<i>confusus</i> , <i>Reeve</i>	LXXXVI.	643
candeleris, <i>Pfr.</i>	LX.	408	<i>confusus</i> , <i>Reeve</i>	LXVIII.	316
Canimarensis, <i>Pfr.</i>	LX.	468	<i>coniformis</i> , <i>Pfr.</i>	LXV.	451
cantatus, <i>Reeve</i>	LVII.	375	<i>connivens</i> , <i>Reeve</i>	LVIII.	388
Cantorii, <i>Philippi</i>	LX.	404	<i>conoidens</i> , Jan.	LXXXVII.	653
Capuira, Spix	LXVIII.	312	<i>copersius</i> , <i>Sow.</i>	XXII.	137
Caraceasensis, <i>Reeve</i>	LXXIX.	580	<i>constrictus</i> , <i>Pfr.</i>	LIX.	402
Caribbeorum, Lam.	XLIX.	320	<i>contiguus</i> , <i>Reeve</i>	LXXIX.	552
Coripeusia, Val.	XXVIII.	171	<i>contortuplicatus</i> , <i>Reeve</i>	LXXXVIII.	658
carneus, <i>Pfr.</i>	LX.	409	<i>contusus</i> , <i>Reeve</i>	XXXVII.	220
castaneus, <i>Pfr.</i>	XXXII.	197	<i>conulus</i> , <i>Reeve</i>	LXXVIII.	577
castrensis, <i>Pfr.</i>	LIX.	396	<i>Coquimbensis</i> , <i>Sow.</i>	XVI.	95
castus, <i>Pfr.</i>	XLV.	282	<i>Cora</i> , D'Orb.	XVII.	100
Cuthartiae, <i>Reeve</i>	XLII.	265	<i>conicus</i> , <i>Sow.</i>	XXI.	134
ceratinus, <i>Benson</i>	LXXVIII.	569	<i>corneus</i> , <i>Menke</i>	LXX.	505
cereus, <i>Reeve</i>	LXX.	501	<i>coronatus</i> , <i>Pfr.</i>	LXIV.	447
cerussatus, <i>Reeve</i>	LXXIV.	536	<i>corrugatus</i> , King	XVII.	101
Ceylanicus, <i>Pfr.</i>	XLIII.	274	<i>Cosenus</i> , <i>Reeve</i>	LXVII.	474
Chersoneicus, <i>Sow.</i>	LXXVIII.	576	<i>costatus</i> , <i>Pfr.</i>	LXV.	450
Chileensis, <i>Brod.</i>	XV.	87	<i>Costerii</i> , Eydoux	XI.	55
Chilensis, Lesson	XVII.	102	<i>costulatus</i> , <i>Pot. & Mich.</i>	LXX.	512
Chimborascensis, <i>Reeve</i>	XLIV.	275	<i>coturnix</i> , <i>Sow.</i>	XIX.	115
chloris, <i>Reeve</i>	XXXVIII.	223	<i>Coxerians</i> , <i>Mor.</i>	XLIV.	279
chrysalidiformis, <i>Sow.</i>	IV.	16	<i>crassilabris</i> , <i>Gray</i>	XXXVII.	221
chrysalis, <i>Pfr.</i>	LVIII.	382	<i>crepundia</i> , D'Orbigny	XLVII.	300
cincinniformis (<i>Helix</i>), <i>Sow.</i>	VI.	28	<i>Cretensis</i> , <i>Pfr.</i>	LXVI.	466
cincinninus, <i>Sow.</i>	VII.	30	<i>Crichtoni</i> , <i>Brod.</i>	XXXIX.	240

Plate.	Species.	Plate.	Species.
crystallinus, <i>Reeve</i>	XXXII.	194 effeminatus, <i>Reeve</i>	LII. 338
eulmineus, <i>D'Orb.</i>	LIV.	360 effusus, <i>Pfr.</i>	XII. 64
Cumingii, <i>Pfr.</i>	XII.	61 egregius, <i>Pfr.</i>	XXXIV. 205
cunctator, <i>Reeve</i>	LXXVI.	554 Ehrebergi, <i>Pfr.</i>	LX. 411
Curiensis, <i>Reeve</i>	LVIII.	399 electricus, <i>Reeve</i>	V. 21
curtus, Koch	XXIII.	151 electrum, <i>Reeve</i>	LVI. 373
Cuyensis, <i>Pfr.</i>	IX.	43 elegans (<i>Succinea</i>), <i>Pfr.</i>	XIX. 112
Cuzeoensis, <i>Reeve</i>	LXXI.	514 elongatus, <i>Pfr.</i>	XIV. 80
cyclostoma, <i>Reeve</i>	LXV.	459 emarginatus, Deshayes	LXIII. 431
cylindricus, <i>Gray</i>	LXIV.	444 eous, <i>Reeve</i>	LXXXVI. 640
cylindricus, Menke	LXI.	421 erectus, <i>Reeve</i>	LVIII. 392
eymatilis, <i>Reeve</i>	LXI.	421 eremita, <i>Benson</i>	LXXVIII. 573
dauctylus, <i>Brod.</i>	VIII.	39 erosus, <i>Brod.</i>	XXII. 149
Daphnus (<i>Helix</i>), Brod.	II.	8 erubescens, <i>Pfr.</i>	LVII. 381
Dardanus, <i>Fris.</i>	LXI.	422 erythrostoma, <i>Sow.</i>	XIII. 75
Darwinii, <i>Pfr.</i>	XXI.	136 eschariferus, <i>Sow.</i>	XX. 121
dealbatus, <i>Sny.</i>	LXV.	455 Eubioicus, <i>Reeve</i>	LXXIX. 584
deceptus, <i>Reeve</i>	LXVIII.	488 euryonophalus, <i>Jouas</i>	LIV. 356
decoratus, <i>Dec.</i>	LI.	334 euryzonatus, <i>Pfr.</i>	XI. 56
decoratus, <i>Sow.</i>	XXI.	128 evanescens, <i>Brod.</i>	XIII. 72
decoratus, Lea	XLIV.	275 exesus (<i>Clau.</i>), <i>Spix</i>	XXXVIII. 227
decorticatus, <i>Reeve</i>	XXX.	592 exiguis, <i>Reeve</i>	LXXXVIII. 654
decussatus, <i>Reeve</i>	LXXII.	519 exilis, <i>Gmelin</i>	XLVI. 292
delumbis, <i>Reeve</i>	LXXVI.	555 eximius, <i>Reeve</i>	XXIX. 175
demotus, <i>Reeve</i>	LIV.	354 exornatus, <i>Reeve</i>	LXXVII. 560
Dennisoni, <i>Reeve</i>	XXVI.	166 fabrefactus, <i>Reeve</i>	XLIX. 319
dentatus, King	XXXVIII.	233 fasciolatus, <i>Oliv.</i>	LXI. 415
depstus, <i>Reeve</i>	LXXXIII.	524 Faunus (<i>Helix</i>), Brod.	III. 13
depictus, <i>Reeve</i>	LXXXIV.	529 Favanius, <i>Lam.</i>	XXIX. 179
derelectus, <i>Brod.</i>	XXIII.	151 fenestratus, <i>Pfr.</i>	XXXVI. 214
Deshayesii, <i>Pfr.</i>	XL.	250 feriatus, <i>Reeve</i>	XLVIII. 314
detritus (<i>Helix</i>), Mull.	LIX.	399 ferrugineus, <i>Reeve</i>	LXII. 424
Diana, <i>Brod.</i>	VIII.	40 Ferussaci, <i>Dunker</i>	LXIV. 411
digitele, <i>Reeve</i>	XLVII.	308 fibratulus, <i>Martyn</i>	XXVIII. 170
dilatatus, <i>Pfr.</i>	XII.	69 fictilis, <i>Brod.</i>	IX. 44
discrepans, <i>Sow.</i>	XXIII.	145 fidustus, <i>Reeve</i>	LXXVI. 557
distortus, <i>Brug.</i>	LIV.	355 flammeus (<i>Helix</i>), Mull.	LIII. 352
Dombeyanus (<i>Helix</i>), Fév.	XXX.	186 foliatus, <i>Reeve</i>	XXXVI. 211
Dominiucus, <i>Reeve</i>	LXXXVIII.	659 follieulus, <i>Pfr.</i>	LXXXVII. 644
D'Orbignii, <i>Pfr.</i>	XXII.	143 formosus, <i>Gray</i>	LXIV. 448
Downesii, <i>Gray</i>	XXIX.	177 Forskalii, <i>Beck</i>	LXI. 419
Draparnaudi, <i>Pfr.</i>	XIX.	116 foveolatus, <i>Reeve</i>	LXXXIII. 526
Dryas, <i>Brod.</i>	IX.	45 fragilis, Parreyss	LXXVII. 473
dubiosus, Jay	XL.	249 fragosus (<i>Helix</i>), Fév.	LXIV. 446
Dufresnii, <i>Leach</i>	XXXVII.	219 frater, <i>Fév.</i>	VI. 25
Dunkeri, <i>Pfr.</i>	LXXXIV.	533 fraterculus (<i>Helix</i>), Fév.	LXIII. 438
durus, <i>Spix</i>	XL.	253 Frivaldszkyi, <i>Pfr.</i>	LXXV. 545
Dussumieri, —?	LXV.	457 fucatus, <i>Reeve</i>	LXXXIII. 615
Dysoui, <i>Pfr.</i>	LXII.	425 fulgetrum, <i>Brod.</i>	V. 23
eburneus, <i>Pfr.</i>	LXI.	418 fulguratus, <i>Jay</i>	XXIX. 175

	Plate.	Species.	Plate.	Species.		
<i>fulguratus</i> , Val.	XXXIV.	204	<i>Hegewischii</i> , <i>Pfr.</i>	LXX.	508	
<i>fulminans</i> , <i>Nyst</i>	XXIV.	160	<i>Helema</i> , <i>Quoy</i>	XLVII.	306	
<i>fulvicanus</i> , <i>Pfr.</i>	XXXIX.	241	<i>Hennabi</i> , <i>Gray</i>	XXIII., LXXXVIII.	148	
<i>Funckii</i> , <i>Nyst</i>	XXVIII.	171	<i>heterotrichus</i> , <i>Mor.</i>	XXXII.	199	
<i>fuseagula</i> , <i>D'Orbigny</i>	XLVII.	305	<i>Hindsii</i> , <i>Pfr.</i>	VII.	33	
<i>fusiformis</i> , Menke	LXI.	420	<i>Hohenackeri</i> , <i>Kryn.</i>	LXI.	416	
<i>fusiformis</i> (<i>Helix</i>), Rang	XL.	249	<i>holostoma</i> , <i>Pfr.</i>	LIX.	490	
<i>fusoides</i> , <i>D'Orb.</i>	XVIII.	106	<i>Hondurasinus</i> , <i>Brug.</i>	LXXXVII.	647	
<i>Galeottii</i> , <i>Nyst</i>	XXXVII.	217	<i>hordeaceus</i> , <i>Brug.</i>	XXIII.	147	
<i>gallina-sultana</i> , <i>Fav.</i>	XXXII.	198	<i>huancensis</i> , <i>Reeve</i>	LVIII.	391	
<i>gastrum</i> , <i>Ehrenberg</i>	LXXV.	538	<i>Humboldtii</i> , <i>Reeve</i>	XXX.	182	
<i>gelidus</i> , <i>Reeve</i>	LXXVI.	553	<i>hyalinus</i> , <i>Wagn.</i>	XXXIV.	205	
<i>geometricus</i> , <i>Pfr.</i>	XLIV.	278	<i>hybridus</i> , <i>Gould</i>	CLXIX.	324	
<i>gibber</i> , <i>Krynicki</i>	LXXXIII.	611	<i>hyenatus</i> , <i>Reeve</i>	XXXVI.	215	
<i>Gibbosius</i> , <i>Lea</i>	XXXII.	196	<i>hygrohykeus</i> , <i>D'Orb.</i>	ignavus, <i>Reeve</i>	LXXXVII.	562
<i>gilvus</i> , <i>Sow.</i>	II.	7	<i>immaculatus</i> , <i>Adams</i>	XXXIII.	203	
<i>glaber</i> , <i>Gmelin</i>	LIV.	357	<i>imperator</i> , <i>Pfr.</i>	XXXV.	631	
<i>glandiformis</i> , <i>Lea</i>	XLII.	266	<i>impressus</i> , <i>Reeve</i>	XXX.	183	
<i>glomeratus</i> , <i>Reeve</i>	LXXX.	591	<i>Inca</i> , <i>D'Orb.</i>	LIII.	349	
<i>goniostoma</i> (<i>Helix</i>), Fér.	XXXIV.	206	<i>incomptus</i> (<i>Helix</i>), <i>Sow.</i>	XL.	53	
<i>Goodallii</i> (<i>Helix</i>), Müller	LXXXIV.	621	<i>Indicus</i> , <i>Pfr.</i>	XLIX.	495	
<i>Gossei</i> , <i>Pfr.</i>	XLVI.	462	<i>indutus</i> , Menke	XV.	86	
<i>Goudotii</i> , <i>Pétît</i>	XXXVII.	218	<i>inflatus</i> , Brod.	XIX.	111	
<i>gracilis</i> , <i>Watton</i>	LXIX.	495	<i>inflatus</i> , <i>Spix</i>	XXXIX.	236	
<i>gracillimus</i> , <i>Pfr.</i>	LXXX.	594	<i>inflatus</i> , <i>Laurek</i>	LXX.	512	
<i>Granadensis</i> , <i>Pfr.</i>	XXXIX.	234	<i>infrafuscatus</i> , <i>Gould</i>	LXXV.	550	
<i>granulosus</i> , Brod.	XVII.	102	<i>inglorius</i> , <i>Reeve</i>	LV.	368	
<i>granulosus</i> (<i>Helix</i>), Rang	XXXV.	209	<i>interstinctus</i> , <i>Gould</i>	LXXXIX.	367	
<i>Grateloupii</i> , <i>Pfr.</i>	XIV.	81	<i>imutilus</i> , <i>Reeve</i>	LXXXVI.	639	
<i>Gravesii</i> , King	XVII.	101	<i>iostoma</i> , <i>Sow.</i>	XV.	88	
<i>Grayanus</i> , <i>Pfr.</i>	XXXVIII.	229	<i>irregularis</i> , <i>Pfr.</i>	LXV.	454	
<i>gregarius</i> , <i>Adams</i> & <i>Reeve</i>	LXXXIII.	612	<i>irroratus</i> , <i>Reeve</i>	LXII.	427	
<i>Griffithsi</i> , <i>Benson</i>	XLVII.	302	<i>interruptus</i> , <i>Brug.</i>	XXXI.	187	
<i>Gruneri</i> , <i>Pfr.</i>	LI.	332	<i>inversus</i> , <i>Brug.</i>	XXXI.	187	
<i>Guadaloupeusis</i> , <i>Brug.</i>	XLVI.	294	<i>Jacobi</i> , <i>Sow.</i>	XXI.	135	
<i>Guerini</i> , <i>Pfr.</i>	XXXII.	193	<i>Jamaicensis</i> , <i>Reeve</i>	LXX.	503	
<i>Guildingii</i> , <i>Pfr.</i>	LXIV.	445	<i>Jancirensis</i> , <i>Sow.</i>	XXXVIII.	226	
<i>Guimaraensis</i> , Brod.	VI.	24	<i>Jayanus</i> , <i>Lea</i>	XXXVII.	220	
<i>Guineensis</i> , <i>Jous</i>	LXXXVI.	641	<i>Jeanoli</i> , <i>Terver.</i>	LXXV.	539	
<i>guttatus</i> , Brod.	XXII.	144	<i>Jerdoni</i> , <i>Benson</i>	XLVI.	297	
<i>Hachecusis</i> , <i>Reeve</i>	LXXXV.	627	<i>Jonasi</i> , <i>Pfr.</i>	LV.	363	
<i>laevastoma</i> , <i>Scop.</i>	XXXV.	210	<i>juglans</i> , <i>Pfr.</i>	III.	12	
<i>halepeusis</i> , <i>Pfr.</i>	LX.	413	<i>Jussieui</i> , <i>Val.</i>	XXXIX.	242	
<i>Hamiltoni</i> , <i>Reeve</i>	LXXXIII.	610	<i>Kauboul</i> , <i>Brug.</i>	L.	327	
<i>Hankeyi</i> , <i>Pfr.</i>	XXX.	181	<i>Kelletii</i> , <i>Reeve</i>	LXXXIX.	661	
<i>haplostylus</i> , <i>Pfr.</i>	III.	341	<i>Kieneri</i> , <i>Pfr.</i>	LXVI.	463	
<i>arpa</i> , <i>Pfr.</i>	LXXX.	596	<i>Kingii</i> , <i>Gray</i>	LI.	336	
<i>arpa</i> , <i>Pfr.</i>	LXXX.	596	<i>Knorr</i> , <i>Pfr.</i>	XLIII.	270	
<i>Harrisii</i> , <i>Reeve</i>	LXXXVI.	638				
<i>Hartwegii</i> , <i>Pfr.</i>	XXIX.	176				

	Plate.	Species.		Plate.	Species.
<i>Kordofanus</i> , Parreyss	L.	327	<i>Loxensis</i> , Pfr.	XL.	251
<i>Kraussi</i> , Pfr.	LXIII.	436	<i>lucidus</i> , Reeve	XL.	245
<i>Kremnouensis</i> , D'Orb.	LI.	337	<i>lubricus</i> (<i>Helix</i>), Müller	LXXXVII.	645
<i>Kunawurensis</i> , Hutton	LXII.	426	<i>latens</i> , Ziegler	LIX.	399
<i>labeo</i> , Brod.	XXXV., LXXI., LXXII.	207	<i>Luzonicus</i> , Sow.	XI.	55
<i>Iabiatius</i> , Ziegler	LXIII.	431	<i>Lycicus</i> , Pfr.	LX.	407
<i>labiosus</i> (<i>Helix</i>), Mull.	LX.	412	<i>Lymnoëdes</i> , Lamarck	LXX.	510
<i>labrosus</i> , Oliv.	LX.	410	<i>Lyonuctianus</i> (<i>Helix</i>), Pallas	XLI.	257
<i>Lackhamensis</i> , Fleming.	LXXXVI.	634	<i>maculentus</i> , Reeve	LXXIX.	586
<i>lectarius</i> , Menke	XXXVII.	217	<i>macrostoma</i> , Pfr.	I.	1
<i>lacunosus</i> , D'Orb.	XXXV.	208	<i>maculatus</i> , Lea	XLVIII.	309
<i>ketus</i> , Reeve	LXXXIII.	616	<i>maculiferus</i> , Brod.	VI.	26
<i>levus</i> (<i>Helix</i>), Mull.	XXXVII.	216	<i>naufragicus</i> , Grat.	XXXIII.	202
<i>lagotis</i> , Menke	XLI.	259	<i>magnus</i> , Wagner	XLIX.	321
<i>Lamarckianus</i> , Pfr.	XXIV.	156	<i>mauleatus</i> , Jay	XXIX.	174
<i>Largilliertii</i> , Philippi	LI.	346	<i>Manoelii</i> (<i>Helix</i>), Mor.	XLVIII.	311
<i>larvatus</i> , Brod.	IX.	44	<i>mampticus</i> , Reeve	LV.	369
<i>latebricola</i> , Benson	LXXX.	572	<i>maritimus</i> , Spix	LIL.	342
<i>lateralis</i> , Menke	XLI.	258	<i>marmorarius</i> , D'Orb.	XLI.	255
<i>latireflexus</i> , Reeve	LXXXVIII.	568	<i>marmoratus</i> , Phil.	XXIV.	157
<i>Latrirei</i> , Pfr.	XXXVI.	211	<i>Martinicensis</i> , Pfr.	LXIII.	434
<i>Laurettii</i> , Sow.	XX.	119	<i>mauviorius</i> , Reeve	LXXVII.	561
<i>Levi</i> , Pfr.	XII.	66	<i>maximus</i> , Sow.	LI.	337
<i>lemniscatus</i> , Desh.	XVIII.	105	<i>Meiacoshimensis</i> , Ad. & Reeve	LXXXVIII.	655
<i>leopardus</i> , Pfr.	X.	51	<i>melanocheilus</i> , Nyst.	XXVIII.	172
<i>leptocheilus</i> , Pfr.	XXXI.	189	<i>melanorhinus</i> , Jan.	LIX.	399
<i>leucodou</i> , Pfr.	LXIII.	432	<i>melanostoma</i> , Swain.	XXXIII.	203
<i>leucophoeus</i> (<i>Helix</i>), Sow. . .	X.	50	<i>melo</i> , Quoy	XXXIX.	243
<i>Liberianus</i> , Reeve	LXXXVIII.	660	<i>membranaceus</i> , Phil.	LXXV.	544
<i>lichenorum</i> , D'Orb.	XIV.	83	<i>Menkei</i> , Gruner	XLV.	285
<i>lignarius</i> , Pfr.	I.	3	<i>Merdueianus</i> , Kryn.	LXVII.	473
<i>lilacinus</i> , Reeve	LXXIV.	532	<i>Meridanus</i> , Pfr.	LVII.	386
<i>iliacinus</i> , Gmeling	XLV.	287	<i>meridionalis</i> , Pfr.	LVI.	370
<i>Limensis</i> , Reeve	LXXVII.	563	<i>Mexicanus</i> , Linn.	XL.	244
<i>limonocoicus</i> , D'Orb.	LVI.	371	<i>Mexicanus</i> , Val.	LVIII.	391
<i>Lindei</i> , Reeve	XXXI.	189	<i>micra</i> , D'Orb.	XIV.	78
<i>linearis</i> , Krauss	LXXXVII.	648	<i>micra</i> , D'Orb.	LXXIX.	579
<i>lineatus</i> , Rossmässler	LXI.	420	<i>Miersii</i> , Sow.	XXXIX.	239
<i>lineatus</i> , Spix	LXII.	428	<i>Milleri</i> , Sow.	XXVIII.	169
<i>linostoma</i> , D'Orb.	LXXXV.	613	<i>mittochelus</i> , Reeve	XLIX.	322
<i>liquabilis</i> , Reeve	LVII.	387	<i>Minosarum</i> , D'Orb.	LXVII.	484
<i>lita</i> , Desh.	XXXIX.	236	<i>Mindorensis</i> , Brod.	IV.	15
<i>lithocicus</i> , D'Orb.	XVII.	98	<i>minimus</i> , Adams	LXXXIV.	623
<i>lituratus</i> , Spix	XLIX.	321	<i>mirabilis</i> , Adams	LXXXIV.	622
<i>litus</i> (<i>Helix</i>), Fé.	XLIX.	321	<i>modestus</i> , Brod.	XVIII.	109
<i>livescens</i> , Br.	XLVII.	304	<i>modestus</i> , Sow.	XII.	62
<i>lividus</i> , Reeve	LXXXV.	626	<i>mollicellus</i> , Reeve	LXXVII.	565
<i>Lobbi</i> , Reeve	LXXI.	516	<i>monilifer</i> , Pfr.	XLVIII.	318
<i>Loevei</i> , Pfr.	XXIV.	160	<i>moniliferus</i> , Gould	XIII.	73
<i>Loewii</i> , Philippi	LXXVII.	477	<i>monozonus</i> , Pfr.	XXXII.	195

INDEX.

	Plate	Species		Plate	Species
Montagnei, D'Orb.	XXIII.	146	<i>octonoides</i> , D'Orb.	LIX.	494
montanus, Drap.	LXXXVI.	634	<i>odontostoma (Pupa)</i> , Gray	XXXVIII.	228
<i>Montacuti</i> , Jeffreys	LXXXVI.	634	<i>olivaceus</i> , Pfr.	LXVII.	479
Monte-Videusis, Pfr.	XIX.	114	<i>Olivieri</i> , Pfr.	LI.	339
montivagus, D'Orb.	LXVII.	475	<i>olorinus</i> , Duclos	XIII.	75
Moricandi, Pfr.	XLV.	283	<i>omager</i> , Beck	XLV.	284
Moritzianus, Pfr.	XXV.	162	<i>onça</i> , D'Orb.	XVIII.	108
Mozambicensis, Pfr.	LVIII.	328	<i>oryza</i> , Brodrip	V.	22
multicribris, Reeve	LXXXI.	598	<i>opalinus</i> , Sow.	LVIII.	394
multicolor (<i>Helix</i>), Rang	XXXIX.	238	<i>Oparanus</i> , Pfr.	LXXXVII.	646
multifasciatus, Lam.	XLVI.	295	<i>Orbignyi</i> , Pfr.	LXIII.	439
murrinus, Reeve	XLIII.	273	<i>Oreades</i> , D'Orbigny	XLVIII.	313
mus, Brod.	II.	10	<i>ornatus</i> , Hutton	LXVII.	478
mutabilis, Brod.	XVII.	100	<i>oryza</i> , Brug.	LXVIII.	480
multilatus, Reeve	LI.	331	<i>ovatus</i> (<i>Helix</i>), Mull.	XXXVI., LXXXIX.	212
myristicus, Reeve	LXXII.	520	<i>ovoideus</i> , Brug.	XI.	56
nanus, Reeve	LXXIX.	585	<i>ovulum</i> , Reeve	LXXVI.	556
Natalensis, Krauss	LXII.	439	<i>pachychelus</i> , Pfr.	V.	87
navicula (<i>Helix</i>), Wagn.	XLI.	258	<i>palaceus</i> , Busch	XXXI.	187
neglectus, Pfr.	LVI.	376	<i>pallaeus</i> , Jonas	LXII.	423
neuricus, Reeve	LXXVI.	550	<i>pallidior</i> , Sow.	LV.	365
nigrofasciatus, Pfr.	LV.	379	<i>Panameensis</i> , Brod.	XIII.	70
nigropileatus, Reeve	LXXXIII.	525	<i>Panayensis</i> , Pfr.	XIV.	76
nigriluteatus, Reeve	LXXXVII.	567	<i>Pantagruelius</i> , Mor.	XXXVIII.	230
Nilagaricus, Pfr.	XLVI.	291	<i>papyraceus</i> (<i>Helix</i>), Mawe	XXXIX.	236
nimbosus, Brod.	IV.	17	<i>paradoxus</i> , Pfr.	IX.	45
Niso, Pfr.	LXXVI.	635	<i>pardalis</i> , Fr.	XXIV.	157
nitelinus, Reeve	LIX.	398	<i>Partuloides</i> , Brod.	XI.	54
nitens, Schmidt	LXXXVII.	645	<i>patricius</i> , Reeve	LXXXI.	600
nitidulus, Pfr.	LXXX.	588	<i>pauperculus</i> , Adams	LXXXIV.	624
nitidus, Brod.	XVIII.	103	<i>Pazianus</i> , D'Orb.	LVI.	377
nivicola, Bous	LXIX.	496	<i>pellucidus</i> , Pfr.	LXVIII.	487
nobilis, Reeve	V.	20	<i>pemphigodes</i> , Jonas	LXV.	453
noctivagus, Parryss	LXVII.	471	<i>Peutlandi</i> , Reeve	LXXXIII.	614
nucinus, Reeve	LXXXV.	629	<i>perdix</i> , Pfr.	LIV.	355
nucleus, Sow.	XVII.	99	<i>perplexus</i> , Sow.	XXXI.	191
Numidicus, Reeve	LIII.	351	<i>perspectivus</i> , Pfr.	LXIII.	435
nux, Brod.	XXIII.	150	<i>Peruvianus</i> , Brug.	XVII.	101
Nympha, Pfr.	VIII.	41	<i>pessulatus</i> , Reeve	XXIII.	153
obeliscus (<i>Helix</i>), Mor.	LII.	343	<i>Petiti</i> , Pfr.	XXXVII.	222
obesus (<i>Helix</i>), Féussac	LXX.	504	<i>Pfeifferianus</i> , Reeve	IV.	18
oblitus, Reeve	LVI.	376	<i>phasianellus</i> , Val.	XV.	88
obliquus, Reeve	LXXVI.	551	<i>Philippinensis</i> , Pfr.	I.	5
oblungus (<i>Helix</i>), Mull.	XXXV.	210	<i>Philippi</i> , Pfr.	XXII.	143
obscurus (<i>Helix</i>), Mull.	LXXXVII.	647	<i>phlogerus</i> , D'Orb.	XXVII.	168
obsolete, Ziegler	LXI.	420	<i>pictor</i> , Brod.	IV.	19
obtusatus (<i>Helix</i>), Gmel.	LII.	344	<i>Physoides</i> , Menke	LXX.	507
obtusus (<i>Pupa</i>), Drap.	LXXIX.	583	<i>picturata</i> , Féussac	XLVI.	295
occultus, Reeve	LXXXIII.	617	<i>pintadinus</i> , D'Orb.	XLII.	265
octonoides, Adams	LXXXIV.	593	<i>piperatus</i> , Sow.	XVI.	96

	Plate.	Species.		Plate.	Species.
planidens, <i>Mich.</i>	XXXI.	190	rhodocheilus, <i>Reeve</i>	XXVIII.	173
plectostylus, <i>Pfr.</i>	XLII.	268	rholodarynx, <i>Reeve</i>	LXXII.	518
poculus, <i>D'Orb.</i>	XVI.	91	rhodospira, <i>Mor.</i>	XXXIII.	203
politus, <i>Reeve</i>	XIII.	74	rhodostoma, <i>Gray</i>	XLIX.	323
polygyrus, <i>Reeve</i>	LXXIX.	578	rimatus, <i>Pfr.</i>	LIV.	359
Popelairianus, <i>Nyst.</i>	XXXIV.	204	ringens, <i>Dunker</i>	LXXV.	542
porphyrius, <i>Pfr.</i>	XV.	89	Rivasi, <i>D'Orb.</i>	XX.	122
porraceus, <i>Jay</i>	IX.	45	Romblorensis, <i>Pfr.</i>	VII.	34
Powisianus, <i>Pelt</i>	XXVII.	167	rosaceus, <i>King</i>	XV.	87
prætextus, <i>Reeve</i>	LXXI.	515	rosatus, <i>Reeve</i>	LIV.	353
pretiosus, <i>Caudor</i>	LXXXIII.	619	Rossmässleri, <i>Pfr.</i>	LX.	405
primula, <i>Reeve</i>	LVII.	385	rubellus, <i>Brod.</i>	XVIII.	104
primirularis, <i>Reeve</i>	LXXXIII.	527	rubescens, <i>Reeve</i>	XXIII.	148
princeps, <i>Sow.</i>	XV.	90	rubicundulus, <i>Gould</i>	LV.	367
procerus, <i>Adams</i>	LXIX.	494	rubrifasciatus, <i>Reeve</i>	XLIV.	277
Proteus, <i>Brod.</i>	XVII.	100	rudis, <i>Anton</i>	XLV.	286
proximus, <i>Sow.</i>	LXXXII.	607	rufistrigatus, <i>Benson</i>	LXXVIII.	570
pruinosus, <i>Sow.</i>	XX.	120	rufogaster, <i>Lesson</i>	I.	4
pseudo-Succinea (<i>Helix</i>), <i>Mor.</i>	LXII.	429	rufo-niger, <i>Fér.</i>	LIX.	395
pubescens, <i>Mor.</i>	LXIII.	437	rugatus, <i>Reeve</i>	LXXVI.	552
pubicus (<i>Helix</i>), <i>Linu.</i>	XXXI.	191	rugiferus, <i>Sow.</i>	XX.	118
pulchellus, <i>Brod.</i>	XVIII.	104	rugulosus, <i>Sow.</i>	XX.	123
puellaris, <i>Reeve</i>	LXXXVI.	637	rupestrus, <i>Phil.</i>	LIX.	491
puicularis, <i>Reeve</i>	XLII.	267	rupicolus, <i>Reeve</i>	XVI.	93
pullus, <i>Gray</i>	LXVII.	476	Ruppellianus, <i>Pfr.</i>	L.	329
pulverulculus, <i>Pfr.</i>	XXXIX.	241	rusticus, <i>Mousson</i>	LXXVIII.	574
punctatus, <i>Anton</i>	LXV.	452	Salicarus, <i>Reeve</i>	LXXXVIII.	657
punctatissimus (<i>Clau.</i>), <i>Less.</i>	XXXVIII.	225	Sandwicensis, <i>Pfr.</i>	LXVII.	472
punctulifer, <i>Sow.</i>	XVI.	92	sarcodes, <i>Pfr.</i>	LXI.	414
pupa (<i>Helix</i>), <i>Linu.</i>	LXIII.	431	Satyrus, <i>Brod.</i>	VI.	29
Pupiformis, <i>Brod.</i>	XIV.	85	Sayi, <i>Pfr.</i>	LXV.	458
Pupoidea (<i>Clau.</i>), <i>Spix.</i>	XXXVIII.	231	seabious, <i>Sow.</i>	XIV.	84
purpuratus, <i>Reeve</i>	LXXI.	517	Scalariformis, <i>Brod.</i>	XXI.	129
pustulosus, <i>Brod.</i>	XX.	127	Scalarioides, <i>Reeve</i>	LXXX.	590
purus, <i>Mousson</i>	XXI.	187	Schiedeana, <i>Pfr.</i>	LIV.	361
pyramidalis (<i>Helix</i>), <i>Sow.</i>	XI.	59	scitulus, <i>Reeve</i>	LXXI.	513
pythagaster, <i>Fér.</i>	I.	2	scobinatus, <i>Wood</i>	XXXII.	199
quadricolor, <i>Pfr.</i>	XLII.	263	sculpturatus, <i>Pfr.</i>	XX.	125
quadridens (<i>Helix</i>), <i>Muller</i>	LXXXVII.	650	scutulatus, <i>Brod.</i>	XIX.	110
quinquedentatus, <i>Pfr.</i>	LXXXIII.	618	seductilis, <i>Pfr.</i>	LXXXVI.	635
Quiteus, <i>Pfr.</i>	XLVIII.	317	segregatus, <i>Benson</i>	LXXIX.	587
Recluzianus, <i>Pfr.</i>	LXXIV.	530	serperastrus, <i>Say</i>	XL.	252
recouditus, <i>Reeve</i>	LXXVI.	549	sex-dentatus (<i>Clau.</i>), <i>Spix.</i>	XXXVIII.	224
redditus, <i>Reeve</i>	XLVII.	307	Shongi, <i>Lesson</i>	XXIV.	159
Reevii (<i>Helix</i>), <i>Brod.</i>	I.	6	Sidonensis (<i>Helix</i>), <i>Fér.</i>	LXIII.	433
reflexus, <i>Pfr.</i>	LXXXV.	628	signatus (<i>Auric.</i>), <i>Wagn.</i>	XXXIII.	200
regina (<i>Helix</i>), <i>Fér.</i>	XXVII.	168	signatus, <i>Sow.</i>	LXIV.	444
reticulatus, <i>Reeve</i>	XXIV.	443	Sikkimeus, <i>Reeve</i>	LXXXVII.	651
revolutus, <i>Ziegler</i>	LXXXIII.	611	simplex, <i>Jonas</i>	XII.	63
rhodaemus, <i>Pfr.</i>	XIV.	77	Siudicus, <i>Benson</i>	XLVII.	303

	Date.	Species.		Date.	Species.
sinistralis, <i>Reeve</i>	LXXXI.	603	tener, <i>Ziegler</i>	LXVII.	473
Siquijorensis, <i>Pfr.</i>	VIII.	42	tenuissimus (<i>Helix</i>), <i>Fér.</i>	XLV.	288
smaragdinus, <i>Reeve</i>	VI.	27	tepidulus, <i>Reeve</i>	LXXXVI.	642
Socotrensis, <i>Pfr.</i>	LXIV.	440	terebella, <i>Adams</i>	LXXXIV.	620
solidus, <i>Pfr.</i>	II.	9	terebialis, <i>Pfr.</i>	XIV.	79
solitarius (<i>Helix</i>), <i>Poiret</i> . . .	LXXXVII.	653	terebrastris, <i>Férussas</i>	LII.	341
solivagus, <i>Reeve</i>	LXXIV.	531	Terterii, <i>Forbes</i>	LXXV.	539
solutus, <i>Troschel</i>	LXXV.	540	thamnoicus, <i>D'Orb.</i>	XVI.	94
sordidus, <i>Lesson</i>	XVII.	100	Thompsoni, <i>Pfr.</i>	XXIV.	158
sordidus (<i>Helix</i>), <i>Lesson</i>	LIX.	401	Ticaonicus (<i>Helix</i>), <i>Brod.</i>	III.	14
Sowerbyi, <i>Pfr.</i>	LVII.	383	tigris, <i>Brod.</i>	XVIII.	107
spadicetus, <i>Menke</i>	LXXXIV.	537	torridus, <i>Gould</i>	LXXXIX.	662
speciosus, <i>Parryss</i>	L.	327	Tournefortianus, <i>Fér.</i>	LXI.	417
spectatus, <i>Reeve</i>	LXXXI.	601	translucens, <i>Brod.</i>	XIII.	71
spectralis, <i>Reeve</i>	LIII.	348	transparcus, <i>Reeve</i>	LXXVII.	566
Spixii, <i>Wagn.</i>	XXX.	182	tribalteatus, <i>Reeve</i>	XLI.	269
spoliatus, <i>Parryss</i>	L XVIII.	485	trichodes, <i>D'Orbigay</i>	LXXV.	547
sporadicus, <i>D'Orbigy</i>	XLIX.	325	tricinctus, <i>Reeve</i>	LVII.	380
Spratii, <i>Pfr.</i>	LIX.	403	tridens, <i>Brug.</i>	LXXXVI.	636
spretus, <i>Reeve</i>	LXXXV.	633	tridentatus, <i>Parryss</i>	LXVIII.	489
stabili, (<i>Helix</i>), <i>Sow.</i>	XII.	67	trifasciatus (<i>Helix</i>), <i>Chemn.</i>	XXXIX.	237
stramineus, <i>Goulding</i>	LXXXV.	632	trilineatus, <i>Quoy</i>	XLVIII.	310
striato-costatus, <i>D'Orbigy</i>	LXXX.	594	trilineatus (<i>Helix</i>), <i>Quoy</i>	LIX.	397
striatulus, <i>Sow.</i>	XXII.	143	truncatus, <i>Pfr.</i>	LXX.	498
striatus, <i>King</i>	XXII.	139	tuberculatus, <i>Turton</i>	LXIII.	431
strigatus, <i>Sow.</i>	XLIV.	280	Tuckeri, <i>Pfr.</i>	LXVIII.	481
strigilosis, <i>Dufo</i>	LV.	362	tumefactus, <i>Reeve</i>	LVI.	374
Studerii, <i>Pfr.</i>	LVII.	384	tumidulus, <i>Pfr.</i>	XIX.	111
subcarinatus, <i>Pfr.</i>	VII.	35	Tunetanus, <i>Reeve</i>	LXXIX.	581
subdiaphanus (<i>Pupa</i>), <i>King</i> . . .	LXIX.	493	Tupacii, <i>D'Orb.</i>	XV.	86
subsemiclausus, <i>Petit</i>	XLI.	254	turbatinus, <i>Lea</i>	LXXXII.	605
subtilis, <i>Rossaußler</i>	LXVII.	471	turricola, <i>Pfr.</i>	LIX.	497
subula, <i>Pfr.</i>	LXIX.	494	turriformis, <i>Krauss</i>	LXXXVII.	652
subuliformis (<i>Helix</i>), <i>Moric.</i> . . .	LXVIII.	486	turritella, <i>D'Orbigy</i>	LXX.	511
succinctus, <i>Reeve</i>	LXXIV.	534	turritus, <i>Brod.</i>	XX.	124
Suecinoïdes, <i>Petit</i>	XLII.	264	tutulus, <i>Benson</i>	LXXXIV.	625
suffusus, <i>Reeve</i>	LIII.	350	uber, <i>Pfr.</i>	X.	48
sulcosus, <i>Pfr.</i>	XLIX.	324	umbilicaris, <i>Soul.</i>	LXVI.	460
sultanus, <i>Lamarck</i>	XXXI.	187	umbomaticus, <i>Reeve</i>	LXVII.	559
superbus, <i>Jonas</i>	XXVIII.	171	undatus, <i>Brug.</i>	XV.	90
Swainsonii, <i>Pfr.</i>	XXXIII.	203	undulatus, <i>Guild</i>	XL.	247
Sylheticus, <i>Reeve</i>	LXXVII.	564	unicarinatus (<i>Pupa</i>), <i>Lam.</i>	LXVI.	468
Sylvanus, <i>Brod.</i>	IX.	46	unicolor, <i>Sow.</i>	XXI.	133
sylvaticus, <i>Wagner</i>	LII.	340	uvidentatus (<i>Part.</i>), <i>Sow.</i>	XXXI.	192
Syriacus, <i>Pfr.</i>	LX.	406	unifasciatus, <i>Sow.</i>	XXIII.	149
teniolus, <i>Nyst</i>	XLIX.	326	ustulatus, <i>Sow.</i>	XXI.	130
Taunaysii (<i>Helix</i>), <i>Fér.</i>	XXXIII.	202	Valencienensis, <i>Pfr.</i>	XXXIV.	204
Tauricus, <i>Lang</i>	LXI.	420	variabilis, <i>Hartm.</i>	LNIX.	499
Taylorianus, <i>Reeve</i>	LXXXI.	602	variatus, <i>Webb & Berth.</i>	LVII.	461
tencibricus, <i>Reeve</i>	LIII.	347	varians, <i>Brod.</i>	XVIII.	104

	Plate.	Species.		Plate.	Species.
<i>variedentatus</i> , Hartman	LXXXVI.	636	<i>vitreus</i> , Spix	XXXVIII.	231
<i>variegatus</i> , Pfr.	XVI.	93	<i>Vitrinoides</i> , Reeve	XLVI.	290
<i>Varneensis</i> , Fricaldsky	LXXV.	543	<i>vittatus</i> , Spix	XLIV.	279
<i>velatus</i> , Brod.	XI.	57	<i>vittatus</i> , Brod.	XVIII.	105
<i>velutinus</i> , Pfr.	LV.	362	<i>voithianus</i> , Pfr.	XXI.	131
<i>velutino-hispidus</i> , Moric.	XXXII.	199	<i>Wagneri</i> , Pfr.	XXXVIII.	232
<i>Venezuelensis</i> , Nyst.	XXIV.	157	<i>Woodiana</i> , Lea	III.	11
<i>venosus</i> , Reeve	XLV.	285	<i>xantholinus</i> , Ziegler	L.	327
<i>ventricosus</i> , Drap.	LIX.	499	<i>xanthostoma</i> , Hohen.	LXI.	416
<i>ventrosus</i> (<i>Helix</i>), Fér.	LXIX.	499	<i>xanthostoma</i> , D'Orb.	XLIII.	271
<i>Veranyi</i> , Pfr.	XLII.	262	<i>Yungasensis</i> , D'Orb.	LNV.	456
<i>vereundus</i> , Reeve	II.	62	<i>zebra</i> (<i>Bucc.</i>), Mull.	XV., XXVII.	90
<i>versicolor</i> , Brod.	XIX.	113	<i>zebra</i> , Olivier	XLVI.	464
<i>vezillum</i> , Brod.	XXIII.	152	<i>zebra</i> , Spix	XXX.	284
<i>vibex</i> , Hutton	XLVII.	299	<i>zebriolus</i> (<i>Helix</i>), Fér.	LXVI.	464
<i>vinineus</i> (<i>Helix</i>), Mor.	XL.	246	<i>Ziebmanni</i> , Pfr.	LXX.	506
<i>Vincentinus</i> , Pfr.	LV.	366	<i>Ziegleri</i> , Pfr.	LVIII.	389
<i>virens</i> , Pfr.	VII.	32	<i>zigzag</i> , Lam.	XV.	90
<i>virescens</i> , Swainson	XXXI.	187	<i>zonatus</i> , Swain.	XXXIX.	237
<i>virgatus</i> , Spix	XLVIII.	312	<i>zonulatus</i> , Pfr.	XIII.	58
<i>virginicus</i> , Brug.	XXXI.	191	<i>zoographicus</i> , D'Orbigny	LXX.	509
<i>virgulatus</i> (<i>Helix</i>), Fér.	XLIX.	320			

- Sp. 4. *B. rufogaster*. Add authority, Lesson, Illus. de Zool. pl. 42.
- Sp. 10. *B. mus*. Add authority, Broderip, MS.
- Sp. 15. *B. Mindorensis*. Add as synonymes, *B. aspersus*, Grataloup, and *B. amoenus*, Pfeiffer.
- Sp. 28. *B. cincinniformis*. Add authority, *Helix cincinniformis*, Sowerby, Pro. Zool. Soc. 1841, p. 17.
- Sp. 53. *B. incomplata*. Having expressed a doubt of this species, it may be as well to add that specimens similar to the original one have been received by Mr. Cuming from the Philippine Islands, and others were collected by Capt. Belcher in the same locality during the voyage of the Samarang.
- Sp. 55. *B. Luzonicus*. Add as synonyme, *B. Costerii*, Eydoux, Mag. de Zool. 1838, Moll. pl. 116.
- Sp. 56. *B. ovoideus*. Add as variety *B. euryzonus*, Pfeiffer.
- Sp. 69. *B. dilatatus*. Insert Hab. Island of Luzon, Philippines.
- Sp. 72. *B. evanescens*. Add authority, Broderip, Pro. Zool. Soc. 1841, p. 15.
- Sp. 73. *B. Adamsii*. Dr. Gould informs me that this species is identical with his *B. moniliferus*, from Savoy, Pro. Bost. Soc. Nat. Hist. 1846.
- Sp. 74. For *B. eburneus* read *B. politus*, the name being pre-occupied by Pfeiffer, Sp. 41.
- Sp. 75. *B. erythrostoma*. Add as variety, *B. albus*, Sow., Pro. Zool. Soc. 1833, p. 73; and *B. olorinus*, Duelos, Guerin's Mag. de Zool. 1833, Moll. pl. 24.
- Sp. 82. *B. cochleatus*. Add authority, Pfeiffer, Pro. Zool. Soc. 1842, p. 151.
- Sp. 86. *B. Tupacii*. Add as synonyme *B. indutus*, Menke.
- Sp. 98. For *B. culminans* read *B. lithoicus*, D'Orbigny, Voy. dans l'Amér. Mérid. Moll. pl. 33, f. 10, 11.
- Sp. 100. For *B. sordidus*, Lesson, read *B. Proteus*, Broderip, the former species being correctly represented at Fig. 401.
- Sp. 112. *B. elegans*. Add as synonyme *B. Buschii*, Pfeiffer.
- Sp. 131. For *B. meridionalis*, Reeve, read *B. Voithianus*, Pfeiffer, Pro. Zool. Soc. 1847, p. 114.
- Sp. 143. For *B. striatulus*, Sow., read *B. Philippii*, Pfeiffer, the first name being pre-occupied by Bruguière in reference to *Buccinum striatum*, Müller, unidentified. For the true *B. D'Orbignii*, under which name this species was communicated to me, see Sp. 439.
- Sp. 148. For *B. rubescens*, Reeve, read *B. Hennahi*, Gray, for which see Pl. LXXXVIII.
- Sp. 151. *B. derelictus*. Add as synonyme *B. curtus*, Koch, Philippi, Abbild. vol. i. p. 155. *Bul. pl. 2. f. 8.*
- Sp. 152. For *B. vexillum*, Brod., read *B. alternans*, Beck, Pfeiffer, Mon. Helic. vol. ii. p. 207, the former name being pre-occupied by Bruguière and by Wood.
- Sp. 157. *B. pardalis*. Add as synonyme *B. marmoratus*, Philippi, Abbild. vol. i. p. 157. *Bul. pl. 2. f. 1 and 2*, and *B. Venezuelensis* and *Venetiolensis*, Nyst, Bull. de l'Acad. Royale de Bruxelles, vol. xii.
- Sp. 164. *B. auris-Sileni*. Insert Hab. Island of St. Vincent, West Indies; Guilding.
- Sp. 165. *B. Adamsoni*. Add authority, Gray, Pro. Zool. Soc. 1833, p. 123.
- Sp. 170. For *B. auris-Midae*, read *B. fibratus*, Martyn, Univ. Conch. pl. 25, published anterior to Chemnitz's *Auris Midae*.
- Sp. 182. *B. Spixii*. Add as synonyme *B. bifasciatus*, Philippi. The fold represented upon the columella in this figure is merely casual.
- Sp. 187. *B. citrinus*. Add as synonymes *B. inversus* and *interruptus*, Bruguière, *B. sultanus*, Lamarck, *B. aureus* and *virescens*, Swainson, *B. palaceus*, Busch, and *B. purus*, Mousson.
- Sp. 189. For *B. Liudenii*, Reeve, read *B. astrapoides*, Jonas, Philippi, Abbild. vol. i. p. 157. *Bul. pl. 2. f. 3 and 4*, quoted erroneously by Dr. Pfeiffer, as a variety of *B. pardalis*.
- Sp. 199. *B. heterotrichus*. Add as varieties *B. velutino-hispidus*, Moricand, and *B. scobinatus*, Wood.
- Sp. 203. *B. melanostoma*. Add as synonymes *B. Swainsoni*, Pfeiffer, and *B. Illeoccola*, Moricand.
- Sp. 204. For *B. Popelairianus*, Nyst, read *B. Valencienensis*, Pfeiffer, which has priority, and include the former name as synonyme.

MONOGRAPH
OF THE GENUS
A C H A T I N A.

.. Each shell, each crawling insect, holds a rank
Important in the scale of Him who framed
This scale of beings; holds a rank, which lost,
Would break the chain, and leave a gap behind,
Which Nature's self would rue!"—*Stillingfleet*



ACHATINA.

PLATE I.

Genus ACHATINA, Lamarck.

Testa ovata vel fusiformi-oblonga, anfractibus plerumque striatis vel granatis, columellā involuta, basi pecudariata truncata, apertura labro simplici, nunquam incassato art refexo. Operculum nullum.

Shell ovate or fusiformly oblong, whorls mostly striated or grained, columella rolled inwards, peculiarly truncated at the base, lip of the aperture simple, never thickened or reflected. No operculum.

The general aspect of the *Achatinae* is very similar to that of the *Bulinus*, but the particulars of the shell in detail present a sufficient typical difference throughout. The chief peculiarity of structure which characterises this genus resides in the columella, it being always involuted towards the aperture, and more or less abruptly truncated at the base. A negative character is also afforded in the outer lip being never thickened or reflected. The *Achatinae* are not of the same arboreal habits as the *Bulinus*, but dwell in places where there is more moisture, in swamps or in the vicinity of lakes and rivers; and are more confined to particular localities. They comprehend three very distinct groups, one consisting of large, richly coloured, ventricose species, chiefly inhabiting West Africa, where they may be said to take the place of the *Bulinus*, another comprising numerous delicate species, some of a semi-transparent, glassy polish, others finely sculptured with granulated or raised striae, inhabiting Central America, the West Indies, and Southern Europe, and constituting the genus *Glandina* of authors. The prettily painted and porcelain-like species *A. virginea* and *vexillaria*, are also representatives of another type, the genus *Liguus* of De Montford.

Species 1. (Mus. Cuming.)

ACHATINA PFEIFFERI. *Achat. testa oblongo-turritā, anfractibus octo, convexis, longitudinaliter obscure plicato-striatis, apicem versus subtler decussatā, ad suturas indistincte crenulatis, columellā vnde arcuatā et contortā, abrupte truncatā; fulvescente-alba, rufo-fusco, praecipue basin versus, flammatā.*

PFEIFFER'S ACHATINA. Shell oblong-turreted, whorls eight in number, convex, longitudinally obscurely plicately striated, finely decussated towards the apex, indistinctly crenulated at the sutures, columella much arched and twisted, abruptly truncated; pale fulvous

white, flamed especially towards the base with reddish brown.

DUNKER, Zeitschr. für Malac. 1845, p. 163.

Hab. Near Loanda, West Africa (on sandy mountains).

As in most species of the genus the colouring is darker and more thickly displayed towards the base.

Species 2. (Mus. Cumiuig.)

ACHATINA EQUATORIA. *Achat. testa acuminato-oblonga, anfractibus septem, tumidis, rotundatis, lēvibus, obliquè teviestratis, ad suturas lirā parvā spirali crenulata cinctis, columellā vix truncatā, labro simplici; ciuere carvalescente, apicem versus albicans, strigis maculatisque fuscis irregulariter aspersa.*

THE EQUATORIAL ACHATINA. Shell acuminate oblong whorls seven in number, swollen, rounded, smooth, obliquely finely striated, encircled with a small spiral crenulated ridge at the sutures, columella scarcely truncated, lip simple; ash-blue, whitish towards the apex, sprinkled irregularly with brown streaks and spots.

Hab. Banks of the river Gaboon, West Africa.

The marking of this species, from a locality nearly on the equator, is very peculiar, not disposed in regular stripes, but wavy and scattered; the whorls are more than usually rounded, and the apex has a papillary aspect.

Species 3. (Mus. Cumiuig.)

ACHATINA VARIEGATA. *Achat. testa conico-oblonga, basin versus ventricosā, anfractibus septem ad octo, plus minus subtler decussatis, ad suturas crenulatis, linea impressā marginatis, columellā contortā, oblique truncatā, labro simplici; fulvo-albidā, strigis nigro-custaneis nudatis, hic illic fulgoratis, sepiissimè pictā, columellā vivide roseā.*

THE VARIEGATED ACHATINA. Shell conically oblong, ventricose towards the base, whorls seven to eight in number, more or less finely decussated, crenulated at the sutures, margined with an impressed line, columella twisted, obliquely truncated, lip simple; fulvous-white, most frequently painted with dark waved chestnut streaks here and there zig-zag, columella very deep rose.

ACHATINA.—PLATE I.

ROISSY, Buffon de Sonnini, Hist. Moll. p. 354.

Bulla achatina, Linnaeus.

Buccinum achatinum, Muller.

Buccinum variegatum, Fabius Columna.

Bulinus achatinus, Bruguière.

Cochlitoma achatina, Féussac.

Achatina perdix, Lamarck.

Hab. Sierra Leone &c., West Africa (in the woods).

This fine large species exists in great abundance in the woods of West Africa, with little variation except that the striped painting is sometimes absent, as in Pl. V.

Species 4. (Mus. Cuming.)

ACHATINA BICOLOR. *Achat. testā subelongato-turritā, onfractibus septem, subirregulariter striatis, ad suturas indistinctè plicato-crenulatis, columellā arcuatā et contortā, fortiter truncatā, aperturā parvō; intensè pureo-nigrā, apicem versus castanē, columellā cæruleo-centre-albā.*

THE TWO-COLOURED ACHATINA. Shell somewhat elongately turreted, whorls seven in number, rather irregularly striated, indistinctly plicately crenulated at the sutures, columella arched and twisted, strongly truncated, aperture small; intense purple-black, chesnut towards the apex, columella bluish-white.

JAY, Cat. Mus. p. 119.

Achatinella bicolor, Pfeiffer.

Achatina adusta, Gould (fide Pfeiffer).

Hab. Sandwich Islands.

This shell has not the generic character of *Achatinella*, the species of which are more or less polished, and distinguished by a peculiar screw-like twist of the columella.

Species 5. (Mus. Cuming.)

ACHATINA COCHLEA. *Achat. testā elongato-turritā, onfractibus octo, spiraliiter costatis, supernè depresso, infra convexo-planis, basi subangulatis, columellā profundi arcuata et contorta; ustulato-castaneā, zonula albā infra suturas.*

THE SCREW ACHATINA. Shell elongately turreted, whorls eight in number, spirally ribbed, depressed round the upper part, convexly flattened beneath, slightly angled at the base, columella deeply arched and twisted; burnt chestnut colour, with a narrow white zone beneath the sutures.

Hab. Peru.

This species is chiefly remarkable on account of its elongated form and spiral ribbed structure, having much the aspect of a *Turritella*.



ACHATINA.

PLATE II.

Species 6. (Mus. Cuming.)

ACHATINA PAPYRACEA. *Achat. testā oblongo-ovatā, tenuis, anfractibus sex, windis decussatis granulatis, ad suturas maryinatis, columellā leviter contortā, parum truncatā; fuscescente, fuscō nebulatā, columellā interdūa subrosacea.*

THE PAPER ACHATINA. Shell oblong-ovate, thin, whorls six in number, minutely decussately granulated, marbled at the sutures, columella slightly twisted, but little truncated; light brown, clouded with darker brown, columella sometimes pinkish.

PFEIFFER, Pro. Zool. Soc. 1845, p. 74.

Hab. Banks of the river Nun, Africa.

A light semi-transparent shell, of which the interior is slightly iridescent.

Species 7. (Mus. Cuming.)

ACHATINA BALTEATA. *Achat. testā acuminato-oblongā, subfusiformi, crassissimula, anfractibus octo ad novem, subplicato-striatis, nudique creberinē minutē granulatis, columellā peculiariter arcuatā, et contortā, abrupte truncatā; fuscescente, fasciis castaneis longitudinalibus apicem versus tinctō, fasciā spirali interrupta, subindistinctā, medio cingulatis, columellā et aperture fuscō carunculacente-albā.*

THE BELTED ACHATINA. Shell acuminately oblong, somewhat fusiform, rather thick, whorls eight to nine in number, slightly plicately striated, very closely minutely granulated, columella peculiarly arched and twisted, abruptly truncated; yellowish, stained towards the apex with chestnut longitudinal bands, and encircled round the middle with a rather indistinct interrupted spiral band, columella and interior of the aperture bluish-white.

Hab. Banks of the river Gambia, Africa.

Very beautifully minutely granulated throughout, and characterized by a faint spiral band, in a manner which obtains some specific importance on account of its singularity.

Species 8. (Mus. Cuming.)

ACHATINA FULICA. *Achat. testā fusiformi-oblongā, subventricosā, nunc tenui, nunc crassissimula, anfractibus*

octo, lœvibus, apicem versus obscurè granulato-striatis, ad suturas vix crenulatis, columellā subcontortā, angustè truncatā; lutescente-albā, fuscō profuse latistriata et nebulatā, columellā et apertura fuscō albīs.

THE SMOKED ACHATINA. Shell fusiformly oblong, rather ventricose, sometimes thin, sometimes thick, whorls eight in number, smooth, obscurely granules striated towards the apex, but slightly crenulated at the sutures, columella rather twisted, narrowly truncated; yellowish-white, profusely broadly streaked and clouded with brown, columella and interior of the aperture bluish-white.

DE FéRUSSAC, Hist. Moll. pl. 124 A. f. 1.

Eadem varietates.

Helix borbonica, De FéruSSAC.

Helix zebra, De FéruSSAC.

Helix Mauritiana, Quoy and Gaimard.

Achatina Conroupa, Lesson.

Hab. Mauritius.

An extremely variable species; in addition to the typical specimen here figured it has been necessary to represent (Pl. III.) a variety of smaller size which is of frequent occurrence, and one (Pl. XI.) of singular umbilicated growth. The variety Fig. 8, b. is of rather light structure, and extremely constant in this peculiarity. MM. Quoy and Gaimard distinguished it as a separate species with the name *B. Mauritiana*, but it is found associated with the larger kind and differs immaterially in form or sculpture. This variety is, however, a very characteristic one, and is found abundantly in some parts of Mauritius. When the Samarang touched at that island on its return to England, Sir David Barclay, an eminent collector residing at Port Luis, presented Sir Edward Belcher with some thirty or forty of them alive from his garden, scarcely differing from one another in size or painting.

In the variety figured at Pl. XI, a very remarkable one, the shell is pressed down, as it were, and the whorls are turned in a manner to form a large umbilicus. Specimens of this variety are not very uncommon, and may be seen in different stages of growth.

Species 9. (Mus. Cuming.)

ACHATINA RETICULATA. *Achat. testā elongato-ovatā, subfusiformi, crassā, anfractibus novem, plicato-rugosis, et striatis, ad suturas concentricē crenatis, columellā*

ACHATINA.—PLATE II.

variolosus, subangustè truncatò; *nivei*, purpureo-castaneo
variegatà, columellà et aperture fauce niveis.

THE RETICULATED ACHATINA. Shell elongately ovate, somewhat fusiform, thick, whorls nine in number, plicately wrinkled and striated, concentrically crenated at the sutures, columella callous, rather narrowly truncated; snowy-white, variegated with purple-chesnut, columella and interior of the aperture snowy-white.

PFEIFFER, Pro. Zool. Soc. 1845, p. 74.
Hab. Africa.

This fine species is remarkably distinct from any other, unless a comparison be made with the *A. lactea*, which is smaller and unspotted.



ACHATINA.

PLATE III.

Species 10. (Mus. Cuming.)

ACHATINA FULVA. *Achat. testā sub-acuminato-oblongā, solidiusculā, anfractibus octo ad novem, convexis, irregulariter striatis, apicem versus minutē decussatis, ad suturas rude crenatis, columellā profundē arcuatā, subabruptè truncatā; griseo-albā, castaneo-fusco intenſe nebulato-strigatā, columellā et aperture fave carmineo-albicantibus.*

THE FULVOUS ACHATINA. Shell acuminately oblong, rather solid, whorls eight to nine in number, convex, irregularly striated, minutely decussated towards the apex, rudely crenulated at the sutures, columella deeply arched, not very abruptly truncated; greyish white, deeply striped in a clouded manner with chestnut brown, columella and interior of the aperture bluish-white.

DESHAYES, *Auim. sans vert. vol. viii. p. 309.*

Bulinus faleus, Bruguière.

Hab. East Africa.

The columella of this species is remarkable for its deeply arched curve.

Species 11. (Mus. Cuming.)

ACHATINA ACUTA. *Achat. testā subfusiformi-oblongā, solidiusculā, anfractibus octo ad novem, apicem versus minuto decussatis, ad suturas subdistincte crenulatis, columellā arcuatā et contortā, elongato-productā, vix truncatā; albida, fusco profuse nebulatā et strigatā, columellā et aperture fave carmineo-albītibus.*

THE SHARP ACHATINA. Shell somewhat fusiformly oblong, rather solid, whorls eight to nine in number, minutely decussated towards the apex, rather indis-

tinctly crenulated at the sutures, columella arched and twisted, elongately produced, but slightly truncated; whitish, profusely clouded and streaked with brown, columella and interior of the aperture bluish-white.

FERUSSAC, *Hist. Moll. pl. 124. A. f. 2.*

Hab. Mauritius.

This species is mainly distinguished from the preceding by its more elongated fusiform growth, by which the columella is more produced and less truncated.

Species 12. (Mus. Cuming.)

ACHATINA PANTHERA. *Achat. testā conico-oblongā, ventricosā, anfractibus octo, superne planō-declinibus, medio truidis, longitudinaliter plicato-striatis, ad suturas rude crenulatis, apicem versus minuto decussatis; columellā contortā, abruptè truncatā; albida, maculis striisque fusca carmineo-nebulatis pictā, columellā intenſe purpureo-rosā.*


THE PANTHER ACHATINA. Shell conically oblong, ventricose, whorls eight in number, flatly slanting round the upper part, swollen in the middle, longitudinally plicately striated, rudely crenulated at the sutures, minutely decussated towards the apex, columella twisted, abruptly truncated; whitish, painted with blue-clouded spots and streaks, columella deep purple-rose.

FERUSSAC, *Prod. p. 49. Hist. Moll. pl. 126.*

Achatina Lamarchiana, Pfeiffer.

Hab. Interior of Madagascar.

A beautifully painted species from a locality in which many fine *Colimacea* doubtless remain to be discovered.



ACHATINA.

PLATE IV.

Species 13. (Mus. Cuming.)

ACHATINA PRUXEM. *Achat. testis ovalis, tenui, ventricosius, spirid apicem versus obscurâ, aufrictibus quinque, infra suturas appressis, lacerusculis, apicem versus minutissimè decussatis, columellâ arcuata, vix truncatâ; aureo-lutescente, castaneo latistrigatâ, superne purpureopurpura.*

(a). PLUM ACHATINA. Shell ovate, thin, rather ventricle, obtuse towards the apex, whorls five in number, appressed beneath the sutures, somewhat smooth, very minutely decussated towards the apex, columella arched, but slightly truncated; pale golden-yellow, broadly streaked with chesnut, purple-rose towards the upper part.

Hab. West Africa.

A striking species, distinguished by its obtuse growth and peculiar colouring. The last whorl is of a delicate golden-yellow, marked with dark chesnut streaks converging together at the base, whilst the upper whorls are of a purple-rose, indistinctly spotted at the sutures.

This species may possibly have been taken for the young of *A. marginata*, but it is certainly distinct.

Species 14. (Mus. Dennison.)

ACHATINA MARGINATA. *Achat. testis conico-ovatâ, ventricosa, aufrictibus sex ad septem, levibus, apicem versus minutissimè decussatis, infra suturas peculiariter marginatis, columellâ subprofunda arcuata et contorta; lutescente, nigricante-castaneo profuse undulatâ et striata, apice intenso roseo, columellâ unice albâ, nunc roseâ, apertura fave cerulescente-alba.*

THE MARGINED ACHATINA. Shell conically ovate, ventricle, whorls six to seven in number, smooth, very minutely decussated towards the apex, peculiarly margined beneath the sutures, columella rather deeply

arched and twisted. yellowish, profusely waved and streaked with blackish chesnut, apex deep rose, columella sometimes white, sometimes rose, interior of the aperture bluish-white.

SWAINSON, Zool. Illus. pl. 30.

Achatina amphora, Jay.

Hab. West Africa.

Mostly very richly marked externally, but not coloured in the aperture as in the species which follows.

Species 15. (Fig. a, Mus. Cuming; Fig. b, Mus. Dennison.)

ACHATINA PURPUREA. *Achat. testis ovalis, solidiscutâ, aufrictibus sex, convexis, infra suturas marginatis et eximie crenulatis, longitudinaliter striatis, apicem versus decussatis, columellâ subprofunda arcuata et contorta; crenulcente-olivaceâ, pallidi fusco raro striata, apice roseo, columellâ et apertura fave intenso roseâ, interdum nigro-purpureo marginatis.*


THE PURPLE ACHATINA. Shell ovate, rather solid, whorls six in number, convex, margined and delicately crenulated beneath the sutures, longitudinally striated, decussated towards the apex, columella somewhat deeply arched and twisted; bluish-olive, faintly streaked at distant intervals with pale brown, pink at the apex, columella and interior of the aperture deep rose, sometimes edged with black-purple.

LAMARCK, Ann. sans vert. (Deshayes' edit.) vol. viii. p. 296.
Bulla purpurea, Chemnitz.

Bulinus purpurascens, Bruguière.

Hab. Cape Palmas, West Africa.

This species may generally be recognised by the deep rose colouring of the aperture and columella, which, as in the beautiful specimen figured from Mr. Dennison's collection, is sometimes edged with very dark purple, approaching to black.



A C H A T I N A.

PLATE V.

Species 16. (Mus. Cuming.)

ACHATINA ALLISA. *Achat. testā elongato-oblongā, subcylindraceā, anfractibus septem, obliquè subtilliter granato-striatis, infra suturas tenuē crenulatis, apertura parvissimā; columellā arcuatā, subcontortā, abrupte truncatā; olivaceo-lutescente, maculis rufo-castaneis oblongis hic illuc irregulariter inquinatā.*

THE BRUISED ACHATINA. Shell elongately oblong, somewhat cylindrical, whorls seven in number, obliquely finely granosely striated, finely crenulated beneath the sutures, aperture rather small; columella arched, slightly twisted, abruptly truncated; olive-yellow, here and there irregularly bedaubed with reddish chesnut blotches.

Hab. Cape Palmas, West Africa.

This species may be readily distinguished from any of those hitherto described, by its oblong cylindrical form and peculiar style of irregular blotching. The aperture is proportionally smaller and narrower than in the allied species.

Species 17. (Mus. Cuming.)

ACHATINA BICARINATA. *Achat. testā oblongo-ovata, cras-*

sinsculā, anfractibus sex ad septem, concentricè plicato-striatis, ultimo inferne obscurè bicarinato; columellā subarcuatā, fortiter truncatā; intense purpureā, apicē albicantē, fusco obscurè fulgaratā, apertura fauce corrucenti-albā.

THE TWO-KEELED ACHATINA. Shell oblong-ovate, rather thick, whorls six to seven in number, concentrically plicately striated, last whorl obscurely two-keeled towards the lower part, columella arched, strongly truncated; deep purple, whitish towards the apex, marked with obscure brown waves, interior of the aperture bluish-white.

LAMARCK, Annu. sans vert. (Deshayes' edit.) vol. viii. p. 296.

Bulla bicarinata, Dillwyn.

Bulinus bicarinatus, Bruguière.

Cochlitoma bicarinata, Fréussac.

Bulla achatina sinistrorsa maxima, Chemnitz.

Achatina sinistrorsa, Pfeiffer.

Hab. Prince's Islands, West Africa.

The keeled growth of this shell is very obscure and not always perceptible. The name *sinistrorsa* was not given to the species by Chemnitz as a technical specific appellation; it merely formed part of his diagnose.

February, 1849.



23



24



25

ACHATINA.

PLATE VI.

Species 18. (Mus. Cuming.)

ACHATINA INDOTATA. *Achat. testā oblongo-ovatā, tenui, ventricosā, anfractibus septem, infernē levibus, superne granoso-striatis, columellā arcuatā; lutescente-albā, strigis latī castaneis fulgarotis.*

THE UNXENODED ACHATINA. Shell oblong-ovate, thin, ventricose, whorls seven in number, smooth round the lower part, granosely striated round the upper part, columella arched; yellowish-white, marked with broad zigzag chestnut-brown streaks.

Hab. West Africa.

This shell, of which I have examined many specimens, is frequently confounded with *A. zebra*, from which it differs constantly in its proportions and the contour of the columella. The painting, though it assimilates in a remarkable degree with some varieties of *A. zebra*, has nevertheless a peculiarity of character about it.

Species 19. (Mus. Cuming.)

ACHATINA STRIATA. *Achat. testā fusiformi-oblongā, anfractibus octo, longitudinaliter conspicuè plicato-striatis, ad suturas crenulatis, columellā contortā, subelongato-protrudē, apertura parvissimā, angustā; opaco-albā.*

THE STRIATED ACHATINA. Shell fusiformly oblong, whorls eight in number, longitudinally conspicuously plicately striated, crenulated at the sutures, columella twisted, somewhat elongately produced, aperture rather small, narrow; opake white.

Buccinum striatum, Müller, Verm. p. 149.

Chemnitz, Conch. Cab. vol. ix, p. 36, pl. 120, f. 1028-9.

Hab. Mexico.

M. Deshayes assigns the *A. rosea*, De Férrussac, and Dr. Pfeiffer the *A. Mulleri* of the same author, to the old *Buccinum striatum* of Müller; the species under consideration seems, however, to agree far better both with Müller's description and Chemnitz's figures, excluding f. 1030, which is *A. Mulleri*.

It is a shell of rather solid growth, opake snowy white, with the striae raised in prominent longitudinal folds.

Species 20. (Mus. Cuming.)

ACHATINA ORNATA. *Achat. testā sulciformi-oblongā, ventricosiusculā, anfractibus octo, longitudinaliter crebrè elevato-striatis, ad suturas anguste marginatis, columellā breviusculā, abruptè truncatā, aperturā pareō, oblongā; fulvescente, strigis brevibus fasciis irreguläribus infra suturas obscurè notatā.*

THE ADORNED ACHATINA. Shell somewhat fusiformly oblong, rather ventricose, whorls eight in number, longitudinally very closely elevated striated, narrowly marginated at the sutures, columella rather short, abruptly truncated, aperture small, oblong; light fulvous-brown, obscurely marked beneath the sutures with short, irregular, brown streaks.

PFEIFFER, Ztschr. fur. Malac. 1846, p. 117.

Achatina costalata, Pfeiffer (not of Adams).

Hab. New Granada.

Curiously, but obscurely, marked with short scattered streaks striking out of the sutures.

Species 21. (Mus. Beuson.)

ACHATINA KRAUSSI. *Achat. testā oblongo-ovatā, anfractibus septem, parua striatis aut decussatis, columellā arcuatā et contortā; nigricastanā, strigis angustis albidiis remotè fulgaratis, columellā et apertura fance curvato-albīs.*

KRAUSS'S ACHATINA. Shell oblong-ovate, whorls seven in number, but little striated or decussated, columella arched and twisted; blackish-chestnut, marked with distant narrow white zigzag streaks, columella and interior of the aperture bluish-white.

REEVE, Pro. Zool. Soc., 1842, p. 55. *Couch. Syst. vol. ii. pl. 179. f. 19.*

Hab. Port Natal, Coast of Africa.

Named in honour of Dr. Krauss of Stutgard, whose researches in South Africa have been productive of a valuable work on the Mollusca of that district.



278.



279.



277.



280.



281.

ACCHATINA.

PLATE VII.

Species 22. (Fig. *a* and *b*, Mus. Cuming.)

ACCHATINA MUREA. *Achat. testá pyramidali-conicā, apice truncatā, basi subplanatā et angulatā, aufractibus sex, laevibus, politis, ad suturas minutissimā marginatis, columellā vix truncatā; pallide luteā, maculis ceruleo-viridibus oblongo-quadratis undatis pulcherrimè tessellatā, interdum virente luteā, immaculatā, apice roseo-incarnata.*

THE PORCELAIN ACHATINA. Shell pyramidal conical, swollen at the apex, somewhat flattened and angular at the base, whorls six in number, smooth, polished, very minutely marginated at the sutures, columella slightly truncated; pale yellow, very prettily tessellated with waved oblong-square blue-green spots, sometimes bright yellow, unspotted, tinged with flesh-pink at the apex.

Hab. —?

A pretty species, clearly distinct from the *A. fasciata*.

Species 23. (Mus. Cuming.)

ACCHATINA ZEBRA. *Achat. testá subelongato-oblongā, crassisuscida, aufractibus octo, granoso-striatis, granis in aufractu ultimo exauditis, columellā subcontortā; intensē castaneā, striis angustis albīs crebrè fulgoratā, aertura fance cārnelescente-albā.*

THE ZEBRA ACHATINA. Shell elongately oblong, rather thick, whorls eight in number, granosely striated, grains disappearing on the last whorl, columella slightly twisted; very dark chesnut, marked with close narrow zigzag white streaks, interior of the aperture bluish-white.

LAMARCK, Anni. sans vert. (Deshayes' edit.) vol. viii. p. 295.

Buccinum ochotina, pars, Muller.

Bulla zebra, Chemnitz.

Bulinus zebra, Bruguière.

Cochlitoma zebra, Féussac.

Hab. Caffaria. South Africa.

It is curious to observe in this species, that the longitudinal streaks become gradually narrower and more frequent as the shell increases in growth. For a species commonly placed with this in collections, in which the stripes are broader and much fewer in number, and which differs manifestly in form, see the preceding Plate, Sp. 18.

Species 24. (Mus. Cuming.)

ACCHATINA CARINATA. *Achat. testá pyramidali-conicā, basi plano-augulatā, aufractibus sex ad septem, plano-convexis, sub lente minutissimā decussatis, suturis sub-excavatis, columellā vix truncatā; petalido-albā, fascia opaco-albō ad suturas et, in aufractu ultimo, ad angulum.*

THE KEELED ACHATINA. Shell pyramidal conical, flatly angled at the base, whorls six to seven in number, flatly convex, under the lens very minutely decussated, sutures somewhat hollowed, columella scarcely truncated; transparent white, encircled at the sutures with a band of opaque white, continued round the angle of the last whorl.

PFEIFFER, Symbole, vol. iii. p. 90.

Hab. —?

An interesting species, distinguished by its angular structure and delicate milk-white sutural band.



ACHATINA.

Plate VIII.

Species 25. (Mus. Dennison.)

ACHATINA MULLERI. *Achat. testá fusiformi, spirá elongatō, anfractibus octo, longitudinaliter creberrimè plicato-striatis, ad suturas crenulatis, columellā subconforta, conspicuè truncatā; pellucido-fuscescente, strigis castaneis remotis irregularibus picta.*

MULLER'S ACHATINA. Shell fusiform, spire elongated, whorls eight in number, longitudinally very closely plicately striated, crenulated at the sutures, columella slightly twisted, conspicuously truncated; light transparent brown, painted with irregular distant chestnut streaks.

FERUSSAC, Prodrome, p. 50. No. 357.

Achatina daedylas, Broderip.

Hab. Tumaco, West Coast of South America.

Dr. Pfeiffer refers this shell to the *Buccinum striatum* of Muller, but there is no mention of the characteristic striped painting in the description of that author; neither is there any account of the delicate tinge of yellow or rose colouring which characterizes the species referred to it by M. Deshayes, namely, the *Achatina rosea*. There is no doubt, however, of the shell under consideration being identical with that figured by Chemnitz under the head of *B. striatum* in Conch. Cab. vol. ix. pl. 120, f. 1030.

Species 26. (Fig. a and b, Mus. Cuming.)

ACHATINA SOWERBYANA. *Achat. testá oœato-fusiformi, tenuiculá, subcentricosá, anfractibus sex ad septem, undique pulcherrimè granoso-decussatis, ad suturas crenulatis, columellā valde arcuatā, leviter conforta, conspicuè truncatā; diaphano-fuscescente, nitente, strigis longitudinalibus castaneis angustis frequenter irregulariter picta.*

SOWERBY'S ACHATINA. Shell ovately fusiform, somewhat thin, rather ventricose, whorls six to seven in number, everywhere very beautifully granosely decussated, crenulated at the sutures, columella much arched, slightly twisted, conspicuously truncated; bright transparent brown, shining, irregularly painted with frequent narrow chestnut streaks.

PFEIFFER, Pro. Zool. Soc., 1846. p. 32.

Hab. Mexico.

A fine large delicately bright transparent species of the *Glandina* section of the genus, prettily granulated throughout.

Species 27. (Mus. Dennison.)

ACHATINA LIGNARIA. *Achat. testá oblongo-cylindracei, anfractibus octo, granulis obtongis undique excutatis, ad suturas rudē crenulatis, columellā abrupte truncatā, aperturā parciuersulā; fuscescente-spindicea, strigis angustis saturationibus subobscurè et irregulariter notata.*

THE WOODEN ACHATINA. Shell oblong-cylindrical, whorls eight in number, sculptured throughout with oblong granules, rudely crenulated at the sutures, columella abruptly truncated, aperture rather small; light brown bay, somewhat obscurely and irregularly marked with streaks of a darker colour.

Hab. —?

This fine species from the collection of J. Dennison, Esq., approaches nearer to *A. Sowerbyana* than to any hitherto described.



ACHATINA.

PLATE IX.

Species 28. (Mus. Cuming.)

ACHATINA ALABASTER. *Achat. testă conico-ovată, crassi-
usculă, anfractibus septem, sub lente minutissimè spi-
raliter striatis, suturis subimpresis, columellă vix
truncată, apertură pareă; pelticido-albă, fasciă
aurantio-castaneă media cingulată.*

THE ALABASTER ACHATINA. Shell conically ovate, whorls seven in number, very minutely spirally striated beneath the lens, sutures slightly impressed, columnella scarcely truncated, aperture small; transparent white, encircled round the middle with an orange-chestnut band.

DESHAYES. Anim. sans vert. vol. viii. p. 312.

Helix alabaster, Rang.

Hab. Prince's Islands, West Africa.

The bright chestnut band which encircles this species, is constant in all the examples I have seen.

Fig. 29. (Mus. Cuming.)

For this species see Plate X. Sp. 35, *A. fasciata*, of which the shell here figured (*A. cretata*, Swainson) is a variety.

Species 30. (Mus. Cuming.)

ACHATINA REEVEANA. *Achat. testă conico-ovată, tenui-
colă, anfractibus septem, sub lente minutissimè spiraliter
striatis, obliquè subtiliter plicato-striatis, ad suturas
concentricè crenulatis, columellă vix truncată, apertură
parcinsulă; pelticido-albă, fasciă obscură luteo-fuscă
media cingulată.*

REEVE'S ACHATINA. Shell conically ovate, rather thin, whorls seven in number, very minutely spirally striated beneath the lens, obliquely finely plicately striated, concentrically crenulated at the sutures, columnella scarcely truncated, aperture rather small; transparent white, encircled round the middle with an obscure yellowish-brown band.

PFEIFFER, Monog. Helie. viv. vol. ii. p. 247.

Hab. Banks of the river Gaboon, West Africa.

Distinguished from *A. alabaster*, to which it is next allied, by its larger and lighter structure, sutural crenulations, and faint colouring of the band.

Species 31. (Fig. a and b, Mus. Dennison.)

ACHATINA FUSIFORMIS. *Achat. testă fusiformi-oblongă
tenui, ventricosă, anfractibus septem, superne obscure
subangulato-productis, longitudinaliter creberimè sub-
irregulariter plicato-striatis, striis minutis impressis
spiraliindentatis, ad suturas crispato-marginatis.
columellă abrupte truncată, apertură angustă; strami-
nata vel fuscescens, extimè miniată, strijs angustis huc
illuc saturatè pictă, intus pallidè rosacea.*

THE FUSIFORM ACHATINA. Shell fusiformly oblong, thin, ventricose, whorls seven in number, obscurely slightly angulated round the upper part, longitudinally very closely but rather irregularly plicately striated, spirally indented with minute impressed striae crisply margined at the sutures, columnella abruptly truncated, aperture narrow; pale straw-colour, or brownish, delicately tinged with vermillion, and painted here and there with narrow streaks of darker colour, pale rose within.

PFEIFFER, Pro. Zool. Soc., 1845, p. 75.

Hab. Mountain of Coban, Vera Cruz, Central America.

It is scarcely possible to do justice to the delicate colouring and plicated sculpture of this beautiful species, the inner edge of the lip of which is stained with a peculiar saturnine vermillion, whilst the entire shell is more or less tinged with a warm hue of the same, excepting in the variety represented at Fig. 31. b, which is of a bright silky straw-colour.

Species 32. (Mus. Dennison.)

ACHATINA DENNISONI. *Achat. testă fusiformi-ovată,
spira subacuminată, anfractibus sex, subrude exiliter
plicato-striatis, striis subtilibus impressis decussatis,
apice subpapillari, columella arcuată et contortă,
attenuatè truncată; fuscescens, fuso apicra versus
variegata, anfractu ultimo insinuata, fasciă pallida
obscură cingulata, columellă pallide rosacea, supra
livido-cerulescata, fuso marginata.*

DENNISON'S ACHATINA. Shell fusiformly ovate, spire rather acuminated, whorls six in number, somewhat rudely faintly plicately striated, decussated with fine impressed striae, apex somewhat papillary, columnella arched and twisted, attenuately truncated; pale brown, variegated towards the apex with darker

ACHATINA—PLATE IX.

brown, last whorl unspotted, encircled with a pale obscure band, columella pinkish.

Hab. Bogota, Central America.

A new and very distinct species for which I am indebted to the magnificent collection of J. Dennison, Esq., who received it from the above-named locality.

Species 33. (Mus. Dennison.)

ACHATINA MAGNIFICA. *Achat. testā subfusciformi-ovalā, anfractibus quinque ad ser, subrugosis, obscurē tenuiplicatis, striis minutis spiralibus obscurē decussatis, columellā arcuatā; cæruleo-viridi, luteo fasciatā et flammatā, columellā cæruleo-fuliginea, supra intēse pur-*

pureo-castaneā, labro intus nigricante-castanō, margine minato.

THE MAGNIFICENT ACHATINA. Shell somewhat fusiformly ovate, whorls five to six in number, rather rough, obscurely finely plicated, obscurely decussated with minute spiral striae, columella arched; dark bluish-green, banded and flamed with yellow, columella bluish, purple chestnut above, lip blackish chestnut within, edged with vermilion.

PFEIFFER, Pro. Zool. Soc., 1847, p. 232.

Hab. Quito, Ecuador, Central America.

Another very beautiful species from the collection of J. Dennison, Esq., of which there is also a specimen in the possession of Mr. Cuming.



A C H A T I N A.

Plate X.

Species 34. (Mus. Cuming.)

ACHATINA PICTA. *Achat. testā conico-ovatā, spira suturis impressis, anfractibus septem, levibus, politis, vel obliquè tenuissimè striatis, columellā brevi, vix truncatā, aperturā parvā, luteā, maculis castaneis nebulatis, tenuiā duplicitā, medio cinctā, maculis viridiibus grandibus hic illuc ornatā.*

THE PAINTED ACHATINA. Shell conically ovate, sutures of the spire impressed, whorls seven in number, smooth, polished, or obliquely very finely striated, columella short, scarcely truncated, aperture small; yellow, encircled round the middle with a double fillet of chestnut clouded spots, and ornamented here and there with large green blotches.

REEVE, Pro. Zool. Soc., 1812, p. 56. Conch. Syst. vol. ii. p. 87. pl. 177. f. 5.

Hab. Island of Cuba.

Very richly painted, but in a different style from either of the following species.

Species 35. (Fig. a, b, c; and Pl. IX. Fig. 29, Mus. Cuming.)

ACHATINA FASCIATA. *Achat. testā subpyramidalē-ovatā, solidiusculā, anfractibus octo, plano-converx, levibus, politis, columellā brevissimā, callosā, vix truncatā, aperturā parvā, labro crenato; albā, fasciis latis luteis, maculis viridiibus quadrato-oblongis obliquis ornatā, lineis viridiibus cingulatā; interdum albā, viridi lineatā, apice roseo.*

THE BANDED ACHATINA. Shell somewhat pyramidal ovate, rather solid, whorls eight in number, flatly convex, smooth, polished, columella rather short, callous, scarcely truncated, aperture small, lip crenated; white, ornamented with broad yellow bands painted with large oblique square-oblong green blotches, encircled with green lines; sometimes white, with the green lines only, apex pink.

SWAINSON, Zool. Illus., vol. iii. pl. 162.

Buccinum fasciatum, Muller.

Bulla fasciata, Chemnitz.

Bulinus vexillum, Bruguière.

Achatina vexillum, Lamarck.

Cochlitoma vexillum, Féussac.

Achatina lineata, Valenciennes.

Varietates.

Achatina crenata, Swainson.

Achatina pallida, Swainson.

Achatina Anais, Lesson.

Bulinus zebra, D'Orbigny.

Hab. Island of Cuba, West Indies.

It will be seen by the accompanying figures and the one referred to in the preceding plate, that the painting of this species is extremely variable; one portion of the colour, however, the fine green lines which encircle the shell, is rarely absent, and it is curious to remark that where these reach the lip, there is a notch corresponding to each line.

Species 36. (Fig. a, b, c, Mus. Cuming.)

ACHATINA VIRGINEA. *Achat. testā pyramidalē-ovatā, solidiusculā, anfractibus septem ad octo, levibus, politis, columellā arenatā, brevi, truncatā: aperturā parvā; carneliente-albā, lineis fasciisque angustis miniatis viridiibus et nigris variè cingulatā, columellā nunc albā, nunc intenſe roſatā.*

THE VIRGIN ACHATINA. Shell pyramidal ovate, rather solid, whorls seven to eight in number, smooth, polished, columella small, short, truncated; aperture small; bluish-white, variously encircled with vermilion green and black lines and narrow bands, columella sometimes white, sometimes deep rose.

LAMARCK, Anim. sans vert. (Deshaye's edit.) vol. viii. p. 299.

Bulla virginaea, Linnæus.

Buccinum virginicum, Muller.

Liguus virginicus, De Montford.

Bulinus virginicus, Bruguière.

Cochlitoma virginica, Féussac.

Achatina vittata, Swainson.

Hab. Cuba &c., West Indies.

There is little variation in the style of painting of this well-known species although the colours occasionally differ. They are mostly very brilliant and the shell has all the appearance of painted porcelain.



29



37.



8c



33

ACHATINA.

PLATE XI

Species 29. (Mus. Brit.)

ACHATINA TINTA. *Achat. testa acuminato-oblonga, crassiuscula, anfractibus septem ad octo, levibus, ad suturas marginatis, columellam contortam, subanguste truncatam; flavidam, castaneo profusa tinctam, apice purpurascens, columellam et apertura fance cerascente-albitam.*

THE STAINED ACHATINA. Shell acuminate oblong, rather thick, whorls seven to eight in number, smooth, marginated at the sutures, columella twisted, rather narrowly truncated; yellow, profusely stained with chestnut, columella and interior of the aperture bluish-white.

REEVE, Conch. Syst. vol. ii. pl. 179. f. 18.

Hab. —?

The painting of this shell is very characteristic, and the outer surface of it is further distinguished by its smooth porcelain appearance. It is an extremely rare species; the specimen represented in the 'Conchologia Systematica' is in Mr. Cuming's collection, and that here figured from the British Museum was formerly in the Tankerville collection, afterwards in that of W. J. Broderip, Esq.

Species 37. (Mus. Cuming.)

ACHATINA SEMISULCATA. *Achat. testa subfusiformis oblonga, anfractibus septem, superna granato-striatis, inferne levibus, columellam arcuatam, subdepresso-truncatam; albidam, castaneo interrupto fulguratum.*

THE HALF-SCULPTURED ACHATINA. Shell somewhat fusiformly oblong, whorls seven in number, granosely striated round the upper part, smooth round the lower, columella arched, rather depresso truncated;

whitish, streaked in an interrupted manner with chestnut.

PFEIFFER, Pro. Zool. Soc., 1845, p. 74.

Hab. Benguela, West Africa.

All Mr. Cuming's examples of this species are distinguished by the same broken interrupted style of painting with which this is characterized.

Species 38. (Mus. Cuming.)

ACHATINA COLUMNA. *Achat. testa elongato-turrata, anfractibus septem, subrade granoso-striatis, suturis depresso-concavis, columellam contortam, callosam, plicato-productam, aperturam parvam; flavidam, castaneo undique fulguratum, columellam et apertura fance cerascente-albitam.*

THE COLUMN ACHATINA. Shell elongately turreted, whorls seven in number, rather rudely granosely striated, sutures depresso concave, columella twisted, callous, produced in the form of a plait, aperture small; yellow, streaked throughout with narrow chestnut flames, columella and interior of the aperture bluish-white.

BRUGUIERE, Encyclopédie Méthodique, vol. i. f. 332.

Buccinum columnum, Muller.

Limax flammatus, Martyn.

Helix columnaria, Chemnitz.

Hab. Prince's Islands, West Africa.

It is somewhat remarkable that no species has been discovered to approximate to this. Lamarck, believing it to be an inhabitant of freshwater, placed it originally in the genus *Lymnaea*.



ACHATINA.

PLATE XII.

Species 42. (Fig. 38. Mus. Cuming.)

ACHATINA SIDERATUS. *Achat. testā oratā, ventricosā, tenuestrigata, anfractibus quinque, cerebrinē minutiē granulatis, columellā contortā; fuscā, castano acutē et subtiliter fulgaratā, apertura fance pallidē cœrulecentē.*

THE THUNDER-STRICKEN ACHATINA. Shell ovate, ventricose, rather thin, whorls five in number, very closely minutely granulated, columella twisted; brown, marked with fine sharply zigzag chestnut streaks, interior of the aperture faintly blue.

Hab. ——?

This is apparently a shell of immature growth, but so peculiar in its colour and marking, that I venture to describe it as new. Of a warm brown silken hue, painted with fine sharply zigzag streaks somewhat broken, much more delicate and numerous towards the apex.

Species 39. (Mus. Cuming.)

ACHATINA FLAMMIGERA. *Achat. testā subpyramidaliter ovalā, anfractibus septem ad octo, obliquè obscurè pli-cato-rugosis, columellā rectā, vix truncatā, apertura parvissimā; latescente, flavous subdilatans frequentibus ustulato-fuscis obliquè pictā, basi nigrante-castanā.*

THE FLAMED ACHATINA. Shell somewhat pyramidaliter ovate, whorls seven to eight in number, obliquely obscurely plicately wrinkled, columella straight, scarcely truncated, aperture rather small; yellowish, obliquely painted with frequent fine burnt-brown flames, base blackish chestnut.

DESHAYES, Enc. Méth. vol. ii. p. 10.

Cochlitoma flammigera, Féussac.

Bulinus Richii, Lamarck.

Achatina Richii, mihi, in Conch. Syst.

Hab. Peru.

The columella being very slightly truncated in this species, partakes very much of the character of *Bulinus*.

Species 40. (Mus. Cuming.)

ACHATINA USTULATA. *Achat. testā cylindraceo-oblongā,*

spira exserta, anfractibus septem, granato-striatis, striis in anfractu ultimo evanidis, columellā subrotundatā, abrupte truncatā, apertura parvā; nigrante-castanā, strigis latescientibus longitudinaliter fulguratis, apertura fave cœrulecente.

THE BURNT ACHATINA. Shell cylindrically oblong, spire stretched out, whorls seven in number, granosly striated, striae fading away in the last whorl, columella slightly twisted, abruptly truncated, aperture small; blackish chestnut, marked with longitudinal yellowish zigzag streaks, interior of the aperture bluish.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. viii. p. 297.

Cochlitoma ustulata, Féussac.

Hab. Cape of Good Hope; Krauss.

Well characterized by its cylindrically drawn out form and style of painting.

Species 41. (Mus. Rolfe.)

ACHATINA LACTEA. *Achat. testā oratā, crassiusculā, anfractibus tumidiusculis, audique plicato-striatis et granatis, columellā arcuatā, callous, attenuatè truncatā: intus extusque lactea, immaculatā.*

THE CREAM-COLOURED ACHATINA. Shell ovate, rather thick, whorls somewhat swollen, plicately striated and graned, columella arched, callous, attenuately truncated, cream-coloured within and without, unspotted.

REEVE, Pro. Zool. Soc., 1842. p. 55. Coneh. Syst. vol. ii. p. 86. pl. 177. f. 6.

Hab. Zanzibar.

An extremely characteristic species of a pure uniform cream colour without any indication of pattern.

The specimen originally described, and from which the accompanying figure from the *Conchologia Systematica* was engraved, is in the possession of the Rev. S. C. E. Neville Rolfe of Headlam Hall. Mr. Cuming has since obtained similar specimens from Zanzibar.



ACHATINA.

PLATE XIII.

Species 43. (Mus. Cuming.)

ACHATINA ALGIRA. *Achat. testā acuminato-oblongā, tenui, pellucidā, epidermide pallidē stramineā, anfractibus sex ad septem, longitudinaliter subrude tenuistratis, columellā arcuatā.*

THE ALGERINE ACHATINA. Shell acuminate oblong, thin, transparent, with a pale straw-coloured epidermis, whorls six to seven in number, longitudinally rather rudely finely striated, columella arched.

DESHAYES, *Expedition de Morée*, p. 165.

Batina Algirus, Bruguière.

Cochlicopa Poireti, Féussac.

Achatina Poireti, Rossmaesler.

Hab. Southern Europe and Algeria.

This delicate and well-known species inhabits the provinces of Austria and Italy, the Morea, some of the Islands of the Mediterranean, and the Algerine district of North Africa.

Species 44. (Mus. Cuming.)

ACHATINA CYANOSTOMA. *Achat. testā elongato-turritā, teani, subpellucidā, anfractibus novem ad decem, obscure irregulariter tenuistratis, columellā angustā, arcuatā; albidō-cornuā, cincto-fusco nitente longitudinaliter tinctā, aperturā fauce cinnamomeo-purpurea.*

THE BLUE-MOUTH ACHATINA. Shell elongately turreted, thin, semi-transparent, whorls nine to ten in number, obscurely irregularly finely striated, columella narrow, arched; whitish horn-colour, longitudinally stained with shining chestnut-brown, interior of the aperture bluish.

RUPPELL, Pfeiffer, *Symbolæ*, vol. ii. f. 58.

Hab. Abyssinia; Ruppell.

The rich chestnut-brown colouring of this species resides only in a thin horny epidermal coating with which it is covered.

Species 45. (Mus. Cuming.)

ACHATINA TURRIS. *Achat. testā oblongo-turritā, tenuiculā, diaphanidā, anfractibus septem, longitudinaliter creberrimā et tenuissimā costulato-striatis, columellā rectā, abrupte truncatā, aperturā parvā; pallidē carnaea.*

THE TURRET ACHATINA. Shell oblong-turreted, rather thin, diaphanous, whorls seven in number, longitudinally very closely and finely rib-like striated, columella straight, abruptly truncated, aperture small; pale flesh-colour.

PFEIFFER, *Symbolæ*, vol. iii. f. 91.

Hab. —?

Distinguished from the following species, to which it is nearly allied, by its more elongated structure and smaller aperture, and more delicate striae.

Species 46. (Fig. a, Mus. Cuming; Fig. b, Mus. Dennisou.)

ACHATINA ROSEA. *Achat. cylindracro-oblongā, interdum subfusiformā, anfractibus septem, ad suturas marginatis, longitudinaliter creberrimā costulato-striatis, striis interstiliisque punctato-indentatis, columellā subprofunde arcuatā et contortā, abrupte truncatā; stramineā, vel, roseā, vel pallidē incarnato spadiceā.*

THE ROSE ACHATINA. Shell cylindrically oblong, sometimes inclined to fusiform, whorls seven in number, margined at the sutures, longitudinally very closely rib-like striated, striae and interstices indented with punctures, columella rather deeply arched and twisted, abruptly truncated; straw-colour, or rose, or light flesh-bay.

GRAY, *Annals of Philosophy*, vol. ix. p. 414.

Cochlicopa rosea, Féussac.

Glandina rosea, Pfeiffer.

Hab. Central America (at the roots of trees): Cuming. Honduras; Dyson. Vera Cruz, &c.

Distinguished from its congeners by the minute indented punctures which characterize the surface of the shell in all its varieties.

Species 47. (Mus. Cuming.)

ACHATINA TRUNCATA. *Achat. testā fusiformi-oblongā, subangustā, anfractibus sex, longitudinaliter creberrimā pliato-costulato-striatis, columellā arcuatā et contortā; cinerascente-rosacea, aperturā fauce rosacea.*

THE TRUNCATED ACHATINA. Shell fusiformly oblong, rather narrow, whorls six in number, longitudinally very closely plicately rib-like striated, columella arched and twisted; ashy-pink, interior of the aperture pink.

Glandina truncata, Say, Amer. Couch. pl. 20.
Hab. Florida.

This very interesting shell is, without doubt, the *Glandina truncata* of Say, according to his figure, but whether it is the original *Bulla truncata* of Gmelin, it is impossible to tell; for Gmelin, like Say in the instance before us, may have confounded two or three species as synonyms under one specific name. A close examination of the characters of the shell under consideration, which, as being the species figured by Say, may as well have the name *truncata* assigned to it, will show that it is perfectly distinct. The rib-like striae have a more elevated plicate structure, and they differ materially in having no indented punctures; the colouring is also peculiar, the aperture being tinged with a rich flesh-pink.

Species 48. (Mus. Cuming.)

ACHATINA VANUXEMENSIS. *Achat. testá fusiformi-ovatá, tenui, inflatá, anfractibus sex, longitudinaliter costulato-striatis, ad suturas marginatis, columellá arcuatá; fulvá, fuscó hic illi pallidé strigatá, maculis subobscureis albidis sparsim floccatá.*

VANUXEMEN'S ACHATINA. Shell fusiformly ovate, thin, inflated, whorls six in number, longitudinally rib-like striated, marginated at the sutures, columella arched; fulvous, palely streaked here and there with brown, sparingly flaked with rather obscure white spots.

LEA, Trans. Amer. Phil. Soc. (Extr.) p. 196, pl. 19. f. 78.
Hab. Mexico.

This species of Mr. Lea's approaches very closely to that subsequently named *A. Sonerbyana*, by Dr. Pfeiffer, represented at Pl. VIII. The white flakes serve nevertheless to distinguish it.



50.



52.



51.



53.



52.



54.



56.



55.



57.



58.

A C H A T I N A .

PLATE XIV.

Species 49. (Mus. Cuming.)

ACHATINA PHILIPPINA. *Achat. testá fusiformi-ovatá, subpellucídá, anfractibus octo, supernú rotundatis, undique longitudinaliter creberrimè costulatis, columellá contortá, attenuatá truncatá, aperturá parvissimá, medio subdepressa; sericeo-lutescente, strigis latis intenſe purpureo-brunneis, e suturis hic illic decurrentibus, conspicuā picta.*

PHILIPPI'S ACHATINA. Shell fusiformly ovate, somewhat transparent, whorls eight in number, rounded at the upper part, longitudinally very closely finely ribbed throughout, columella twisted, attenuately truncated, aperture rather small, slightly depressed in the middle; pale silken yellow, conspicuously painted with broad deep purple-brown streaks, descending here and there from the sutures.

PFEIFFER, (*Glandina*,) *Symbolæ*, vol. iii. p. 90.

Hab. Jamaica, Gosse.

The fine ribs with which the surface of this shell is excessively closely sculptured, give it a delicate silken aspect. It is very richly painted in the manner above described.

Species 50. (Mus. Dennison.)

ACHATINA MONILIFERA. *Achat. testá oblongo-ovatá, ven-tricosa, tenui, pellucídá, anfractibus septem, longitudi-naliter creberrimè costulato-striatis, striis basin versus evanidis, columellá contortá, abrupte truncatá; fulvo-roseá, cornéa, strigis angustis remotis rufo-brunneis irreguloriter picta.*

THE NECKLACE-BOUNDED ACHATINA. Shell oblong-ovate, ventricose, thin, transparent, whorls seven in number, longitudinally very closely striated after the manner of fine ribs, stripe fading towards the base, columella twisted, abruptly truncated; fulvous-rose, horny, irregularly painted with distant narrow red-brown streaks.

PFEIFFER, (*Glandina*,) *Pro. Zool. Soc.*, 1845, p. 75.

Hab. Coban, Vera Cruz, Central America; Latre.

An extremely delicate transparent swollen shell in which the columella is not attenuated as in the last species, but abruptly truncated.

Species 51. (Mus. Cuming.)

ACHATINA OLEACEA. *Achat. testá cylindraceo-oblongá,*

pellucido-corneá, spirá brevi, anfractibus septem, levigatis, columellá subprofundè arcuatá, aperturá oblongá, supernú angustá, labro medio dilatato; fulvæ, nitide.

THE OLIVE-SHAPED ACHATINA. Shell cylindrically oblong, transparent-horny, spire short, whorls seven in number, smooth, columella rather deeply arched, aperture oblong, narrow at the upper part, lip dilated in the middle; fulvous, shining.

Helix (Cochlicopa) oleacea, Féussac, Prodrome, No. 360.
Glandina oleacea, Beck.

Hab. Cuba.

A smooth transparent fulvous horny shell, of cylindric oblong form, without any design of painting.

Species 52. (Mus. Cuming.)

ACHATINA LEUCOZOONIAS. *Achat. testá fusiformi-oblongá, spirá subacutainatá, anfractibus octo, plano-convercis, longitudinaliter subtiliter et creberrimè inciso-striatis, columellá contortá, aperturá oblongá, labro angulato-dilatato; intensè purpureo-brunneá, zonulis longitudinalibus angustis albis notata, labro albimarginato.*

THE WHITE-ZONED ACHATINA. Shell fusiformly oblong, spire somewhat acuminate, whorls eight in number, flatly convex, longitudinally finely and very closely striated in a grooved manner, columella twisted, aperture oblong, lip angularly dilated; deep purple-brown, marked with narrow longitudinal white zones, lip edged with white.

SOWERBY, Genera of Shells, f. 3.

Voluta leucozonias, Walch, Naturforsch, vol. iv. p. 40. pl. 1. f. 3, 4.

Achatina albolineata, Lamarck.

Helix (Cochlicopa) leucozonias, Féussac.

Polyphemus leucozonias, Jan.

Glandina leucozonias, Beck.

Hab. Martinique.

This species is remarkable for the angularly dilated growth of the lip, which forms a slightly elevated ridge or varix at intervals of from four to five in a whorl. The shell being of a uniform dark purple-brown colour with the lip edged with white, these delicate concentric varices appear as stripes, and turning inwards over the columella have the appearance of false plaits. Walch, Gimel, and Dillwyn, misled by this varicose peculiarity of growth,

and regarding the columella to be truly plaited, placed the species among the Mitres in their genus *Voluta*.

Species 53. (Mus. Cuming.)

ACHATINA LATREI. *Achat. testā cylindraceo-oblongā, Volvariaformi, spirā brevi, aufRACTIBUS septem, convexo-planis, levibus, cuticulam carneam politam indutā, columellā contortā, lirā callosā marginatā, apertura elongatā, angustā; fulvā, rufō-brunneo longitudinaliter subremotē strigatā.*

LATRE'S ACHATINA. Shell cylindrically oblong, Volvaria-shaped, spire short, whorls seven in number, convexly flattened, smooth, covered with a polished horny cuticle, columella twisted, edged with a callous ridge, aperture elongated and narrow; fulvous, longitudinally rather remotely streaked with red-brown.

PFEIFFER, Pro. Zool. Soc., 1845, p. 138.

Hab. Central America; Latre.

The shell of this species approaches the form of *Volvaria* or *Tornatella*, though totally unconnected with those genera; the columella is not, however, plaited, but elevated at the base into a callous ridge, as in the *Achatina columnata*. The entire surface of the shell is covered with a perfectly transparent polished horny cuticle, through which the dark stripes show with additional lustre.

It is a rare and extremely interesting species, collected by M. Latre in Central America, probably in the province of Vera Cruz, and named by Dr. Pfeiffer in honour of that eminent traveller.

Species 54. (Mus. Dennison.)

ACHATINA NIGRICANS. *Achat. testā oblongo-ovatā, solidissimā, aufRACTIBUS septem, convexis, superne subtilliter concentricē striatis, deinde levibus, cuticulam carneam politam indutā, columellā brevi, contortā, attenuatē truncatā, apertura oblongā, labro medio subdilatato; livido-nigrā, suturis basique fulvis, strigis fulvis per-pans longitudinaliter notatā.*

THE BLACKISH ACHATINA. Shell oblong-ovate, rather solid, whorls seven in number, convex, finely concentrically striated round the upper part, then smooth, covered with a polished horny cuticle, columella short, twisted, attenuately truncated, aperture oblong, lip slightly dilated in the middle; livid-black, sutures and base fulvous, marked with a very few fulvous streaks.

PFEIFFER, (*Glandina*) Pro. Zool. Soc., 1845, p. 75.

Hab. Vera Cruz, Central America; Latre.

An interesting species, covered with a transparent

cuticle of the same highly glazed character as the preceding, from the same locality, more swollen in form, and of a peculiar dark beetle brown or black.

Species 55. (Mus. Cuming.)

ACHATINA VOLUTA. *Achat. testā subcylindraceo-oblongā, oleaceā, solidissimā, spirā brevi, aufRACTIBUS septem, subplanō-convexis, levibus, cuticulam carneam politam indutā, columellā brevi, arcuatā, abrupte truncatā, apertura elongatā, angustā, labro medio subdilatato; olivaceo-brunnescente, subindistinctē longitudinaliter strigata.*

THE VOLUTE ACHATINA. Shell somewhat cylindrically oblong, olive-shaped, rather solid, spire short, whorls seven in number, somewhat flatly convex, smooth, covered with a polished horny cuticle, columella short, arched, abruptly truncated, aperture elongated, narrow, lip slightly dilated in the middle; light olive-brown, somewhat indistinctly longitudinally streaked.

DESHAYES, Anim. sans vert. vol. viii. p. 300, note.

Bulla voluta, Chemnitz, Conch. Cab. vol. ix. pt. 2. p. 16. pl. 117. f. 1009, 1010.

Bulimus glans, Bruguière.

Helix (Cochlicopa) glans, Féussac.

Achatina glans, Lamarck.

Polypheus glans, De Montford.

Glandina glans, Beck.

Glandina olivacea, Schumacher.

Hab. West Indies.

The above species may be regarded as the type of this shining horn-clad group. Though of rare occurrence, it is admirably figured in vol. ix. of the Conchylien Cabinet, published in 1786.

Species 56. (Mus. Cuming.)

ACHATINA DOMINICENSIS. *Achat. testā fusiformi-oblongā, aufRACTIBUS septem, convexis, superne subrotundatis, longitudinaliter subtilissimē et creberimē inciso-striatis, columellā subcontortā, labro medio dilatato; fulvo-aureantā, strigis latiusculis intensē castaneis, remotis, e suturis descendentibus, notatis.*

THE ST. DOMINGO ACHATINA. Shell fusiformly oblong, whorls seven in number, convex, somewhat rounded at the upper part, longitudinally very finely and closely deeply striated, columella rather twisted, lip dilated in the middle; fulvous-orange, marked with distant rather broad dark chestnut streaks descending from the sutures.

DESHAYES, Anim. sans vert. vol. viii. p. 302, note.

Bulla Dominicensis, Gmelin.

ACHATINA.—PLATE XIV.

Helix (Cochlicopa) Dominicensis, Féussac.*Bulinus maculatus*, Bruguière.*Achatina fusco-lineata*, Lamarck.*Polypleura Bruguereus*, Bowditch.*Glandina maculata*, Beck.*Hab.* West Indies.

The varieose peculiarity of growth which characterizes the *A. leucozonias*, is developed in a less prominent degree in this species, the contrast of painting being reversed. The *A. leucozonias* is of a dark chestnut-brown marked with longitudinal white stripes, in places where the white-edged lip has been matured after the manner of a varix. In the present species the shell is of a uniform orange-yellow, and the periodical maturity of the lip is indicated in its course of growth, about four to five times in a whorl, by a broad chestnut stripe.

Species 57. (Mus. Cuming.)

ACHATINA PERUVIANA. *Achat. testá fusiformi-oblongā, spirae suturis peculiariter profundi excavatis, anfractibus subplano-concavis, longitudinaliter crenatim costulatis, intersticiis transversis sulcato-striatis, columellā contortā, et abrupte truncatā, aperturā oblongā; albā, flauoris undatis longitudinalibus, linearisque obliquis rufo-fuscis, conspicuē pictā.*

THE PERUVIAN ACHATINA. Shell fusiformly oblong, sutures of the spire peculiarly deeply excavated, whorls rather flatly convex, longitudinally very closely finely ribbed, interstices marked transversely with grooved striae, columella twisted and abruptly truncated, aperture oblong; white, conspicuously painted with red-brown longitudinal flames and oblique lines.

LAMARCK. Anim. s. vert. (Deshayes' ed.) vol. viii. p. 301.*Helix (Cochlicopa) pretiosa*, Féussac, Hist. Moll. pl. 135 f. 4.*Hab.* Peru; Dombey (fide Lamarck).

This species belongs apparently to a distinct type, the chief peculiarity of which resides in the deeply grooved formation of the sutures. It is of rather solid texture with somewhat the aspect of a marine shell, finely ribbed longitudinally, the ribs turning over the upper edge of the whorls into the sutural depression. In painting it is richly marked with red brown stripes, crossed obliquely with fine lines of the same colour.

Species 58. (Mus. Cuming.)

ACHATINA GLabra. *Achat. testá ovaínula-ovata, tenuiculá, anfractibus septem ad octo, subtilissimè longitudinaliter striatis, glabris, nitidis, columellá brevi, arcuata et contortā; fulvescente cornu, strigis fascis renatis, suturas anfractuum medio descendentibus, ornata.*

THE SMOOTH ACHATINA. Shell subminutely ovate, rather thin, whorls seven to eight in number, very finely longitudinally striated, smooth, shining, columella short, arched and twisted; light fulvous horn-colour, ornamented with distant brown streaks, descending from the sutures to the middle of the whorls.

PFEIFFER. (*Glandina*) Symbolæ, vol. iii. p. 90.*Hab.* Island of Porto Rico.

This species has a smooth glazed appearance, but not arising from the lustre of a horny cuticle, such as that which characterizes the *A. Latrei* and its congeners.



ACHATINA.

Plate XV.

Species 59. (Mus. Benson.)

ACHATINA CEYLANICA. *Achat. testā ovato-oblongā, spirā subpyramidalē, anfractibus septem, rotundelis, levibus, politis, columellā brevi, arcuatā, abruptē truncatā; fulvescente-cornēd.*

THE CEYLON ACHATINA. Shell ovately oblong, spire somewhat pyramidal, whorls seven in number, rounded, smooth, polished, columella short, arched, abruptly truncated; pale fulvous horny.

PFEIFFER, Zeitschr. für Malac., 1845, p. 157.

Hab. Ceylon; Benson, Templeton.

Chiefly distinguished by the confined pyramidal growth of the spire, and short arched columella. The specimen here figured, for which I am indebted to Mr. Benson, was found under some fallen leaves in a grove near Point de Galle.

Species 60. (Mus. Cuming.)

ACHATINA PHILLIPSII. *Achat. testā acuminata-fusiformi, tenui, pellucidā, anfractibus octo, longitudinaliter striatis, columellā vix arcuatā, attenuatē truncatā, operariū parvā; fulvescente-cornēa, strigis castaneis hic illī concinē pictā.*

PHILLIPS' ACHATINA. Shell acuminately fusiform, thin, transparent, whorls eight in number, longitudinally faintly striated, columella slightly arched, attenuately truncated, aperture small; light fulvous horny, neatly painted here and there with chestnut stripes.

ADAMS, Proc. Boston Nat. Hist. Soc., 1845, p. 13.

Hab. Jamaica; Gosse.

A very delicate transparent horny species, neatly painted at irregular intervals with longitudinal chestnut stripes.

Species 61. (Mus. Cuming.)

ACHATINA JAMAICENSIS. *Achat. testā fusiformi-turritatā, crassiusculā, anfractibus decem concrexis, longitudinaliter cerebellinū multicostatis, ad suturas subtiliter crevatis, columellā subrectā, attenuatē truncatā, apertura parvā; pellucido-oblā, strigis rufi-fuscis raris oblique pictā.*

THE JAMAICA ACHATINA. Shell fusiformly turreted, rather thick, whorls ten in number, longitudinally very closely many-ribbed, finely crenulated at the sutures, columella nearly straight, attenuately truncated, aperture small; transparent white, obliquely painted with a few red-brown streaks.

PFEIFFER, Pro. Zool. Soc., 1845, p. 137.
Hab. Jamaica; Gosse.

This species is finely ribbed after the manner of a *Pupa*, where others are simply striated. The painting is arranged obliquely.

Species 62. (Mus. Cuming.)

ACHATINA OBTUSA. *Achat. testā subfusiformi-oblongi, crassiusculā, pellucidā, spirā subobtusa, anfractibus septem, levibus, ad suturas marginatis, columellā ex contortā, abruptē truncatā; fulvescente-cornēa.*

THE OBTUSE ACHATINA. Shell somewhat fusiformly oblong, rather thick, transparent, spire somewhat obtuse, whorls seven in number, smooth, margined at the sutures, columella slightly twisted, abruptly truncated; light fulvous horny.

PFEIFFER, (Glandina,) Pro. Zool. Soc., 1845, p. 42.

Hab. Real Llejós, Province of Nicaragua, Central America, Cuming.

Rather a stout shell, but still of so transparent a substance that the columella of the interior is visible.

Species 63. (Mus. Cuming.)

ACHATINA VENUSTA. *Achat. testā subfusiformi-oblongi, anfractibus septem, longitudinaliter cerebellinū costulato-striatis, columellā vix contortā, attenuatē truncatā; pellucido-rosā, strigis castaneis, albi-marginatis, hic illī e suturis medio descendentibus.*

THE GRACEFUL ACHATINA. Shell somewhat fusiformly oblong, whorls seven in number, longitudinally very closely sculptured with fine rib-like striae; columella scarcely twisted, attenuately truncated; transparent rose-coloured, with white edged chestnut streaks descending here and there out of the sutures to the middle.

PFEIFFER, (Glandina,) Symb. vol. i. p. 46.

Hab. Jamaica.

A delicately rose-tinged species, painted with chestnut streaks edged with white.

Species 64. (Mus. Cuming.)

ACHATINA GRIFFITHII. *Achat. testā subfusiformi, spirā acuminatā, anfractibus novet, cerebrimē et subtilissimē costulato-striatis, ultimo medio impresso, basi expanso, columellā contortā, attenuatè truncatū : pellucido-albā, strigis castaneis sparsis obliquè pictā.*

GRIFFITH'S ACHATINA. Shell somewhat fusiform, spire acuminate, whorls nine in number, very closely and finely sculptured with rib-like striae, last whorl impressed in the middle and expanded at the base, columella twisted, attenuately truncated; transparent white, obliquely painted with scattered chestnut streaks.

ADAMS, Pro. Boston Nat. Hist. Soc., 1845, p. 14.

Hab. Jamaica.

The compressed growth of the last whorl has very much the appearance of an accidental distortion.

Species 65. (Mus. Cuming.)

ACHATINA RANGIANA. *Achat. testā elongato-turritatā, crassā, solidā, anfractibus undecim, plano-convexis, subtilissimā areolatim striatis, lineis impressis spiralibus subdistantibus obscure notatis, columellā simplici, callousā, vis truncatā, aperturā parvā ; albido-stramineā.*

RANG'S ACHATINA. Shell elongately turreted, thick, solid, whorls eleven in number, flatly convex, very finely areately striated, obscurely marked with rather distant impressed spiral lines, columella simple, callous, scarcely truncated, aperture small; whitish straw-colour.

PFEIFFER, Pro. Zool. Soc., 1846, p. 115.

Hab. Republic of Mexico; Linden.

A shell of peculiarly solid growth verging on *Bulimus*, allied to the elongately turreted group of that genus of which *B. calcareus* is the type.

Species 66. (Mus. Cuming.)

ACHATINA TORTILLANA. *Achat. testā subfusiformi-oblongā, subcavatō, anfractibus septem, subtilissimā et cerebrimē costulato-striatis, columellā contortā et attenuatè truncatā ; lutescente-corneā, pellucidā, quasi vitrā.*

THE TORTILLA ACHATINA. Shell somewhat fusiformly

oblong, rather ventricose, whorls seven in number, very finely and closely rib-like striated, columella twisted and attenuately truncated; yellowish horny, transparent, as though glassy.

PFEIFFER, Pro. Zool. Soc., 1846, p. 32.

Hab. Tortilla, Central America.

The opaque blotches and streaks described by Dr. Pfeiffer, appear to be merely accidental in the individual specimen.

Species 67. (Mus. Cuming.)

ACHATINA ASSIMILIS. *Achat. testā oblongā, apice subodosus, anfractibus sex, levibus, politis, e suturis subtilissimā impresso-striatis, superne tenuimarginatis, columellā breviter reflexā, aperturā oblongā ; vividē pellucido-cornēa.*

THE ALLIED ACHATINA. Shell oblong, rather obtuse at the apex, whorls six in number, smooth, polished, very finely impressively striated at the sutures, thinly margined round the upper part, columella shortly reflected, aperture oblong; bright transparent horny.

Hab. —?

This approaches very closely to the *A. obtusa*, Sp. 62, but is of more transparent texture, whilst the columella is less abruptly truncated. The shell is composed of a whorl less in number, and the spire is shorter.

Species 68. (Mus. Cuming.)

ACHATINA HUGELI. *Achat. testā pyramidalis-oblongā, anfractibus decem, rotundatis, superne subtilissimā areolatim striatis, ad suturas sinuulē marginatis, columellā valde arcuatā et contortā, aperturā parvā ; pellucido-albante, hyalina, epidermē tenui cornēa lutescente.*

HUGEL'S ACHATINA. Shell pyramidal oblong, whorls ten in number, rounded, very finely areately striated at the upper part, minutely marginated at the sutures, columella much arched and twisted, aperture small; transparent whitish, hyaline, with a thin yellowish horny epidermis.

PFEIFFER, Symbolae, vol. ii. f. 58.

Hab. —?

A neat hyaline species of pyramidal growth, marked with delicate covered striae round the upper part of the whorls, which become fainter as the shell approaches maturity.



73



76



77

78



ACHATINA.

PLATE XVI.

Species 69. (Mus. Cuming.)

ACHATINA ERECTA. *Achat. testā sublongato-turritā, solitinsculā, apice obtusā, anfractibus octo, convexis, arcuatis striatōis, suturis impressis, columellā contortā, attenuatā truncatā, aperturā parvā, subrotundatā; sordidē fulcescente.*

THE ERECT ACHATINA. Shell somewhat elongately turreted, rather solid, obtuse at the apex; whorls eight in number, convex, finely areately striated, sutures impressed, columella twisted, attenuately truncated, aperture small, rather rounded; dull fulvous colour.

BENSON, Ann. and Mag. Nat. Hist., vol. ix. p. 487.

Hab. China.

Mr. Benson notices this species as inhabiting the island of Chusan and environs of Macao, and M. Largiliert has received it from Nanking.

Species 70. (Mus. Cuming.)

ACHATINA VIVIPARA. *Achat. testā elongato-turritā, suturis impressis, anfractibus decem, cerebrinā subtiliter costulatis, costulis apicem versus remotioribus, columellā subcontortā, abrupte truncatā; albida, epidermide sordidē stramineā indutā.*

THE VIVIPAROUS ACHATINA. Shell elongately turreted, sutures impressed, whorls ten in number, very closely sculptured with fine rib-like striae, last whorl slightly angled at the base, columella rather thin, twisted, attenuately truncated, aperture small; transparent straw-colour.

SOWERBY, Malac. and Conch. Mag., vol. i. p. 27.

Hab. —?

Though of delicate transparent texture, this shell is of rather solid growth, the whorls rounded and aperture small.

Species 71. (Mus. Cuming.)

ACHATINA FRASERI. *Achat. testā subulato-turritā, apice obtusā, subpapillari, anfractibus decem ad undecim, piano-convexis, levibus, ad suturas subtilissimè plicato-crenulatis, columellā arcuatā, abrupte truncatā; fulvescente-albā, epidermide cornea, olateaceā, obscurè fuscostriatā indutā.*

FRASER'S ACHATINA. Shell subulately-turreted, apex obtuse, rather papillary, whorls ten to eleven in

number, flatly convex, smooth, very finely plicately crenulated at the sutures, columella arched, abruptly truncated; pale fulvous white, covered with an olive horny epidermis, obscurely streaked with brown.

PFEIFFER, Symbolae, vol. iii. p. 90.

Hab. Banks of the river Nun, Africa.

Dr. Pfeiffer quotes as synonymous with this species, on the authority of a label in Mr. Cuming's cabinet, *A. iheringi*, Gould, from Cape Palmas.

Species 72. (Mus. Cuming.)

ACHATINA CLAVATA. *Achat. testā acuminato-turritā, anfractibus decem, cerebrinā subtiliter costulatis, costulis apicem versus remotioribus, columellā subcontortā, abrupte truncatā; albida, epidermide sordidē stramineā indutā.*

THE CLAVATE ACHATINA. Shell acuminately turreted, whorls ten in number, very closely finely ribbed, ribs more distant towards the apex, columella slightly twisted, abruptly truncated; whitish, covered with a dull straw-coloured epidermis.

GRAY, Loudon's Magazine, New Series, vol. i. p. 487.

Hab. Sierra Leone.

It is a character to be observed in this species, that the delicate longitudinal ribs are rather distant on the first few whorls of the shell, and become gradually finer and closer towards the aperture.

Species 73. (Mus. Cuming.)

ACHATINA SEMITARUM. *Achat. testā subulata, tenuissimā, apice subpapillari, anfractibus novem ad decem, planulata, subtilissimè superficiatiter striatis, nitidis superne marginatis, columellā arcuatā, attenuatā truncatā, opercularis suboblonga; pelliculo-cornēa.*

THE WINDING-PATH ACHATINA. Shell subulate, very thin, somewhat papillary at the apex, whorls nine to ten in number, flattened, very finely superficially striated, shining, margined round the upper part, columella arched, attenuately truncated, aperture rather oblong; transparent horny.

Helix semitarum, Rang, Pfeiffer, Symbolae, vol. ii. p. 59.

Hab. West Indies.

The sutural margin of the whorls is characterized by a

delicate spiral cord, which Dr. Pfeiffer does not appear to have remarked.

Species 74. (Mus. Cuming.)

ACHATINA STRIATELLA. *Achat. testa subulato-turrata, tenui, anfractibus decem, creberrimè et sublississime costulato-striatis, ultimo obscurè angulato, subproducto, columellâ tenui, contortâ, attenuè truncatâ, aperturâ parvâ, oblongo-ovata; pellucido-corned.*

THE FINELY STRIATED ACHATINA. Shell subulately turrted, thin, whorls ten in number, very closely sculptured with fine ribs, last whorl obscurely angled and slightly produced, columella thin, twisted, attenuately truncated, aperture small, oblong-ovate; transparent horny.

Helix striatella, Rang, Annales Sci. Nat. vol. xxiv. p. 38. pl. 3. f. 7.

Hab. Prince's Island, West Africa; Rang.

The sculpture of this shell is not unlike the *Achatina vi-vipara*, which is of a stouter texture and less subulate form.

Species 75. (Mus. Cuming.)

ACHATINA CLAVUS. *Achat. testa acuminato-oblonga, sub-Pupaformi, solidiusculâ, anfractibus novem ad decem, piano-concavis, exilissime arcuatè striatis, columellâ arcuatâ, aperturâ parvissimâ; sordidè olivaceâ, epidermide tenui cornæ brunnescente induitâ.*

THE NAIL ACHATINA. Shell acuminately oblong, somewhat Pupa-shaped, rather solid, whorls nine to ten in number, flatly convex, very delicately striated in a curved direction, columella arched, aperture rather small; dull olive, covered with a thin brownish epidermis.

PFEIFFER, Symbolae, vol. iii. p. 90.

Hab. —?

This species has a much more solid appearance than the rest of the group.

Species 76. (Mus. Cuming.)

ACHATINA TENUISPIRA. *Achat. testa elongato-turrata, suturis subprofundè impressis, anfractibus undecim, tenuissimâ elevato striatis, columellâ valde arcuatâ; pellucido-cornueâ.*

THE THIN SPIRE ACHATINA. Shell elongately-turreted, sutures rather deeply impressed, whorls eleven in number, very finely elevately striated, columella much arched; transparent horny.

BENSON, Journ. Asiatic Soc., 1836, vol. v. p. 353.
Hab. Darjeeling, Sikkim-Himalaya.

A delicate transparent species from the locality in which Dr. Hooker is at this moment occupied in his natural history researches.

Fig. 77. (Mus. Cuming.)

For description of this species, see *Bulinus*, Pl. LXX.
For *Achatina*, Sp. 77, see Pl. XXI.

Species 78. (Mus. Cuming.)

ACHATINA PAXILLUS. *Achat. testa subulata, subfusiformis, solidiusculâ, basi rotundata, anfractibus novem, sub-plano-concavis, obliquè tenuistriatis, columellâ arcuatâ, subcontortâ, aperturâ parvâ; albidâ, epidermide tenui cornæ luteo-olivaceâ induitâ.*

THE PEG ACHATINA. Shell subulate, somewhat fusiform, rather solid, rounded at the base, whorls nine in number, rather flatly convex, obliquely finely striated, columella arched, slightly twisted, aperture small; whitish, covered with a thin horny yellowish-green epidermis.

Hab. —?

This is rather a stout shell, peculiarly rounded at the base. In other respects it approaches very closely to *A. clavus*.

Achatina. PL. VII.



S1



S2



S3



S4



S5



S6



S7



S8



S9



S10

A C H A T I N A.

PLATE XVII.

Species 79. (Mus. Cuming.)

ACHATINA JAVANICA. *Achat. testa oblongo-turrata, anfractibus octo, rotundatis, obscurè subtilissimè rugè striatis, columellâ tenui, attenuatè truncatâ, apertura parvâ; pallide stramineâ.*

THE JAVA ACHATINA. Shell oblong-turreted, whorls eight in number, rounded, obscurely very finely rudely striated, columella thin, attenuately truncated, aperture small; pale straw-colour.

Hab. Java.

Characterized in some measure by a delicate waxy appearance.

Fig. 80. (Mus. Cuming.)

The shell here represented proves to be a variety of *A. striatella*, a species varying much in growth, but always well characterized by its delicate close-set rib-like striae.

For *Achatina*, Sp. 80, see Pl. XXI.

Fig. 81. (Mus. Cuming.)

For description of this species, see *Bulinus*. Pl. LXX.
For *Achatina*, Sp. 81, see Pl. XXI.

Species 82. (Mus. Benson.)

ACHATINA AMENTUM. *Achat. testa cylindraceo-turrata, tenuissimâ, anfractibus novem, rotundatis, obscurè subtilissimâ plicato-striatis, columellâ arcuatâ et contortâ, attenuatè truncatâ; pellucido-cornicata.*

THE THONG ACHATINA. Shell cylindrically turreted, very thin, whorls nine in number, rounded, obscurely very finely plicately striated, columella arched and twisted, attenuately truncated; transparent horny.

BENSON, MSS.

Hab. Howrah, near Calcutta.

For this species I am indebted to Mr. Benson, who relates having taken three specimens alive in 1835, from beneath a fallen palm tree at Howrah, near Calcutta; and he is not aware that it has been collected since.

Species 83. (Mus. Cuming.)

ACHATINA ARCUATA. *Achat. testa subulata, anfractibus*

octo, planulatis, levibus, politis, striis subtilibus removetis impressis longitudinaliter excutplis, columellâ arcuatâ, abruptè truncatâ; pellucido-lactea, strigis cascarie perspicuis angustis ornata.

THE ARCHED ACHATINA. Shell subulate, whorls eight in number, flattened, smooth, polished, engraved with rather distant fine impressed longitudinal striae, columella arched, abruptly truncated; transparent cream-colour, ornamented with a few narrow chestnut streaks.

PFEIFFER, Pro. Zool. Soc., 1835. p. 138.

Hab. Jamaica; Gosse.

A bright polished species, delicately grooved here and there longitudinally.

Species 84. (Mus. Cuming.)

ACHATINA OCTONA. *Achat. testa subulata-turrata, suturis impressis, anfractibus octo ad decem subrotundatis, minutissimâ striatis, nileantibus, columellâ arcuatâ, contortâ, parva truncatâ, apertura minima; pellucido-cereâ, pallidè stramineâ.*

THE EIGHT-WHORLED ACHATINA. Shell subulately turreted, sutures impressed, whorls eight to ten in number, somewhat rounded, very minutely striated, shining, columella arched, twisted, but little truncated, aperture very small; transparent waxy, pale straw-colour.

GRAY, Annals of Philosophy, New Series, vol. ix. p. 414.
Helix octona, Chemnitz.

Bulinus octonus, Lamarck.

Hab. West Indies, South America, Pacific Islands.

This species, for which others have frequently been mistaken, may be distinguished by its delicate shining surface and very transparent texture; the sutures of the spire are also peculiarly impressed, giving a globose form to the whorls. It has a very wide range of habitation, and is composed sometimes of eight, sometimes of ten whorls.

Species 85. (Mus. Beuson.)

ACHATINA CASSIACA. *Achat. testa elongato-subulata, apice subpapillata, anfractibus quatordecim ad quindecim, planulatis, obliquè crebrè subrude costulato-striatis, suturis impressis, columellâ arcuatâ, abruptè truncatâ; albida, epidermide tenui olivaceo-fuscâ indulda.*

THE CASSIA ACHATINA. Shell elongately subulate, apex somewhat papillary, whorls fourteen to fifteen in number, flat, obliquely very closely and somewhat rudely sculptured with rib-like striae, sutures impressed, columella arched, abruptly truncated; whitish, covered with a thin olive-brown epidermis.

BENSON, MSS.

Hab. Cassia Hills, North East Frontier of Bengal;

Dr. Griffith.

The rib-like striae of this fine species, contributed by Mr. Benson, have an oblique plicate growth.

Species 86. (Mus. Cuming.)

ACHATINA SULCATA. *Achat. testâ elongato-turrilitâ, anfractibus decem, obliquè creberrimè costulato-striatis, anfractu ultimo basi angulato, infra angulum striis minus prominentibus, columellâ arcuatâ et contortâ, parum truncatâ; virente-stramineâ,*

THE GROOVED ACHATINA. Shell elongately turreted, whorls ten in number, obliquely very closely striated after the manner of fine ribs, last whorl angled at the base, striae beneath the angle less prominent, colu-

mella arched and twisted, but little truncated; greenish straw-colour.

GRAY, Annals of Philosophy, New Series, vol. ix. p. 415.
Hab. West Indies.

I much doubt if the *A. striatella* of Rang is not a variety of this species, although the striae are finer and closer together.

Fig. 87. (Mus. Cuming.)

The shell here represented illustrates another form of *A. striatella*, in which the columnella is shorter and of more callous substance.

Fig. 88. (Mus. Cuming.)

This shell, an authentic specimen of Dr. Gould's *A. involuta*, from Cape Palmas, proves to be identical with *A. Fraseri*.

For Achatina Sp. 87 and 88, see Pl. XXI.

Achatina. Pl. XVIII.



ACHATINA.

PLATE XVIII.

Species 89. (Mus. Cuming.)

ACHATINA SUBULATA. *Achat. testá cylindracco-oblongá, spirá subexsertá, anfractibus sex ad septem, lœvibus, nitidis, supremis convexisculis, reliquis sub-planulatis, columellá angustá truncatá, aperturá parviangusta, oblongá; pellucido-corneá, lineá spirali opacá sub-obscurá infra suturas.*

THE SUBULATE ACHATINA. Shell cylindrically oblong, spire rather exserted, whorls six to seven in number, smooth, shining, the uppermost whorls rather convex, the lower a little flattened, columella narrowly truncated, aperture rather small, oblong; transparent horny, with a somewhat obscure spiral opaque line beneath the sutures.

PFEIFFER, (*Polyphemus*) Wiegmann's Archiv. 1839, vol. i. p. 352.

Glandina subulata, Philippi.

Helix orysacea, Rang.

Achatina orysacea, D'Orbigny.

Hab. Cuba.

Distinguished from its allied species by a more oblong-cylindrical growth.

Species 90. (Mus. Cuming.)

ACHATINA INCERTA. *Achat. testá oblongó, spirá obtusissim-
culá, anfractibus quinque ad sex, subplanulatis, lœvibus,
nitidis, superne exiliter marginatis, columellá con-
spicue truncatá, aperturá oblongá, superne angustatá;
pellucido-corneá.*

THE DOUBTFUL ACHATINA. Shell oblong, spire somewhat obtuse, whorls six in number, rather flattened, smooth, shining, faintly margined round the upper part, columella conspicuously truncated, aperture oblong, narrowed at the upper part; transparent horny.

Hab. —?

Of a transparent horny substance, very like *A. solidula*, from which it differs in being more ventricose towards the base.

Species 91. (Mus. Cuming.)

ACHATINA CYLINDRACEA. *Achat. testá cylindracco-ovalá,
tenuiculá, spirá brevissimá, conoidea, anfractibus quin-*

*que, convexo-planis, lœvibus, politis, columellá excavata.
lirá callosa angustá basi conspicue marginatá, apertura
superne elongatá et angustatá, pellucido-corneá.*

THE CYLINDRICAL ACHATINA. Shell cylindrically ovate, rather thin, spire very short, conoid, whorls five in number, convexly flattened, smooth, polished, columella excavated, conspicuously edged at the base with a narrow callous ridge, aperture elongate and narrowed at the upper part; transparent horny.

PFEIFFER, Pro. Zool. Soc. 1846, p. 31.

Hab. Tortilla, Central America.

The structure of this species is similar to that of the *A. Lattevii*, in which the columella instead of being truncated turns up at the edge, as in the present species, into a callous ridge.

Species 92. (Mus. Cuming.)

ACHATINA SOLIDULA. *Achat. testá cylindracco-oblongá,
spirá brevi, anfractibus quinque ad sex, subplanulatis,
lœvibus, politis, superne exiliter marginatis, columellá
brevis, arcuatá, abrupte truncatá, aperturá oblongá,
superne angustatá; pellucido-corneá.*

THE RATHER-SOLID ACHATINA. Shell cylindrically oblong, spire short, whorls five to six in number, rather flattened, smooth, polished, faintly margined at the upper part, columella short, arched, abruptly truncated, aperture oblong, narrowed at the upper part; transparent horny.

PFEIFFER, Wiegmann's Archiv. 1840, vol. i. p. 252.

Glandina solidula, Philippi.

Hab. Cuba.

A bright transparent shell of rather stouter growth than most of this group.

Species 93. (Mus. Cuming.)

ACRATINA PORTO-RICENSESIS. *Achat. testá oblongo-tar-
ritá, anfractibus octo, convexis, lœvibus, politis, lineis
impressis longitudinalibus subobscure exsculptis, colu-
mellá arcuatá, abrupte truncatá, aperturá parviangusta :
cinerascens-corneá, strigis saturatioribus ornatá.*

THE PORTO-RICO ACHATINA. Shell oblong-turreted, whorls eight in number, convex, smooth, polished, rather obscurely sculptured with impressed longitudinal lines, columella arched, abruptly truncated,

aperture rather small; pale ash horn-colour, ornamented with darker stripes.

PFEIFFER, Monog. Helic. viv. vol. ii. p. 258.
Hab. Island of St. John, Porto Rico.

There is a peculiarity of form, as well as of colour, in this species.

Species 94. (Mus. Cuming.)

ACHATINA CURVILABRIS. *Achat. testá fusiformi, spiri acuminati, anfractibus octo, creberrimè costulato-striatis, ad suturas marginatis, columellâ contortâ, basi subproductâ et truncatâ, labro medio impresso; peltacido-cornatâ, strigis angustis albidis obscurè notatâ, labii limbo albido.*

THE CURVED-LIP ACHATINA. Shell fusiform, spire acuminated, whorls eight in number, very closely sculptured with rib-like striae, margined at the sutures, columnella twisted, somewhat produced and truncated at the base; transparent horny, obscurely marked with narrow whitish streaks, edge of the lip whitish.

PFEIFFER, (*Glandina*) Pro. Zool. Soc., 1845, p. 137.
Hab. Jamaica; Goss.

Belonging to the group typified by the *A. lencozonias*, the shell is marked in like manner by white streaks deposited at intervals by the lip after the fashion of variees.

Fig. 95. (Mus. Cuming.)

For description of this species see *Balimus*, Pl. LXX.

For *Achatina*, Sp. 95. see Pl. XXI.

Species 96. (Mus. Cuming.)

ACHATINA FUNCKI. *Achat. testá ovato-conicâ, subumbilicato, tenui, anfractibus sex, sabrotundatis, striatis, nitidis, anfracta ultimo firâ callosa, apertura intranitâ, prope mediuma manita, columellâ tanellato-truncatâ; peltacido-albâ, diaphana.*

FUNCK'S ACHATINA. Shell ovately conical, slightly umbilicated, thin, whorls six in number, rather rounded, very delicately striated, shining, last whorl furnished about the middle with a callous plait entering the aperture; columnella flatly truncated; transparent white, diaphanous.

PFEIFFER, Pro. Zool. Soc., 1847, p. 232.

Hab. Province of Merida, New Granada.

A very transparent wax-like shell, intermediate between *Bulinus* and *Achatina*; the columella, although truncated, is slightly reflected back forming a small umbilicus. On the middle of the body-whorl is a very characteristic callous ridge, entering the aperture.

Species 97. (Mus. Cuming.)

ACHATINA LAMELLATA. *Achat. testá ovato-conicâ, hic subcentrica, minutiè umbilicata, anfractibus sex, hic illuc sublamellato-striatis, columellâ subreflexâ et truncatâ, firâ parvâ callosâ, obscurâ, prope mediuma intranitâ, manita; peltacido-albâ, diaphana.*

THE LAMELLATED ACHATINA. Shell ovately conic, thin, rather ventricose, minutely umbilicated, whorls six in number, here and there slightly lamellately striated; columnella a little reflected and truncated, furnished with an obscure callous ridge entering near the middle; transparent white, diaphanous.

POTIEZ and MICHAUD, Gal. de Douai, p. 128. pl. 11. f. 7 and 8.

Hab. New Lima; Latre.

This species, it will be seen, differs little from the preceding except in being rather more ventricose. The locality, to judge from analogy, is equally doubtful.

Species 98. (Mus. Cuming.)


ACHATINA CONIFERA. *Achat. testá pyramidalis-conicâ, anfractibus octo, subangulis, rotundatis, longitudinaliter obscure striatis, nitidis, columellâ brevi, arcuatâ, contortâ, attenuatâ truncatâ, aperturâ parvâ; fusco-cornatâ.*

THE CONICAL ACHATINA. Shell pyramidalis conical, whorls eight in number, rather narrow, rounded, longitudinally obscurely striated, shining, columella short, arched, twisted, attenuately truncated, aperture small; brown-horny.

FERUSSAC, —?

Hab. Juan Fernandez.

Distinguished by its small aperture and by the elevated growth of the spire.



A C H A T I N A .

PLATE XIX.

(Figures magnified to twice the natural size.)

Species 99. (Mus. Cuming.)

ACHATINA DYSONI. *Achat. testā cylindraceo-oblongā, tenui, basin versus subdilatatā, spirā brevi, anfractibus quinque, lœvibus, politis, columellā contortā, lirā callosā marginatā, aperturā oblongā, medio subconstrictā: peltūcida-cornētā.*

DYSON'S ACHATINA. Shell cylindrically oblong, thin, slightly dilated towards the base, spire short, whorls five in number, smooth, polished, columella twisted, edged with a callous ridge, aperture oblong, somewhat contracted in the middle; transparent horny.

PFEIFFER, Pro. Zool. Soc. 1846, p. 32.

Hab. Honduras, (under decayed leaves): Dyson.

The columella of this delicate transparent species is not truncated at the base, but edged with a callous ridge.

Species 100. (Mus. Cuming.)

ACHATINA SPLENDIDA. *Achat. testā acuminate-ovatā, tenui, striatā, nūtidā, anfractibus sex ad septem, subrotundatis, columellā brevi, contortā, subcallosā, vir reflexi; peltūcida-cornētā.*

THE SPLENDID ACHATINA. Shell acuminate-ovate, thin, finely striated, shining, whorls six to seven in number, subrotund, columella short, twisted, rather callous, scarcely reflected; transparent horny.

ANTON, Verz. p. 41. Philippi, Abbild. *Achat.* p. 1. f. 11.
Hab. Island of Juan Fernandez.

A small bright transparent horny shell, in which the columella is frequently turned up at the edge into a callous ridge.

Species 101. (Mus. Cuming.)

ACHATINA ACCINCTA. *Achat. testā conico-ovatā, tenui, anfractibus sex, subplanu-coriexis, e suturis arcuatis striatis, columellā contortā, plicato-truncatā, aperturā porrexiscula; peltūcida-cornētā.*

THE GIRDLED ACHATINA. Shell conically ovate, thin, whorls six in number, somewhat flatly convex, arcuately striated from the sutures, columella twisted, plicately truncated, aperture rather small; transparent horny.

MIGHELS, Pro. Nat. Hist. Soc. Boston, 1845, p. 20.

Achatina Sandwicensis, Pfeiffer.
Hab. Sandwich Islands.

Dr. Pfeiffer notices that this group of *Achatinae* is allied to the *Achatinellae*; it should be observed that they are of very different substance and texture.

Fig. 102. (Mus. Cuming.)

For description, see Genus *Tornatellina*.
For *Achatina*, Sp. 102, see Pl. XXI.

Species 103. (Mus. Cuming.)

ACHATINA BULIMOIDES. *Achat. testā ventricoso-ovatā, spirā subacuminatā, anfractibus quinque ad sex, subrotundatis, columellā tenui, contortā, aperturā suborbiculari; cornētā, epidermide lateo-olivacea indutā.*

THE BULIMUS-LIKE ACHATINA. Shell ventricosely ovate, spire somewhat acuminate, whorls five to seven in number, rather rounded, columella thin, twisted, aperture nearly orbicular; horny, covered with a yellowish-olive epidermis.

PFEIFFER, Pro. Zool. Soc. 1846, p. 116.

Hab. Island of Juan Fernandez.

A delicately formed species, covered with a thin yellowish-olive horny epidermis.

Species 104. (Mus. Cuming.)

ACHATINA CONSIMILIS. *Achat. testā conico-ovatā, subventricoso, anfractibus quinque ad sex, subplanu-coriexis, striatis, nūtidis, columellā brevi, rectā, subcallosā; lateo-olivacea, yellucida.*

THE VERY SIMILAR ACHATINA. Shell conically ovate, rather ventricose, whorls five to six in number, rather flatly convex, finely striated, shining, columella short, straight, rather callous; yellowish-olive, transparent.

Hab. Juan Fernandez: Müller.

Very similar in general aspect to the preceding species, but clearly distinct; it is less ventricose and the columella is nearly straight.

ACHATINA.—PLATE XIX.

Species 105. (Mus. Cuming.)

ACHATINA OROPHILA. *Achat. testā conico-ovalā, spirā acuminatā, suturis impressis, aufractibus septem, rotundatis, peculiariter obscurè indentatis, columellā brevi, valde arcuatā et truncatā, aperturā parvusculā; oliveo-cornēā.*

THE THATCHED ACHATINA. Shell conically ovate, spire acuminate, sutures impressed, whorls seven in number, rounded, peculiarly obscurely indented, columella short, deeply arched and truncated, aperture rather small; olive-horn.

BENSON, MSS.

Hab. Neilgherry Hills, India; Jerdon. Colombo, Ceylon; Templeton.

Distinguished by the deeply arched curvature of the columella, and by the whorls being rather more numerous than is usual in species of this form.

Species 106. (Mus. Cuming.)

ACHATINA CONCENTRICA. *Achat. testā acuminato-ovalā, aufractibus sex, concentricè plicato-striatis, columellā brevi, contortā, abrupte truncatā, superne lirā callosā muricatā, aperturā parvusculā; cornēā, sordidè oliveacēā.*

THE CONCENTRIC ACHATINA. Shell acuminate-ovate, whorls six in number, concentrically plicately striated, columella short, twisted, abruptly truncated, furnished at the upper part with a callous ridge, aperture rather small; horny, dull olive.

Hab. Bolivia, (under dead Cacti); Bridges.

Finely sculptured by numerous arched concentric striae, whilst the body-whorl is furnished with a winding callous ridge.

Species 107. (Mus. Cuming.)

ACHATINA LAMELLOSA. *Achat. testā acuminato-ovalā, aufractibus sex, subrotundatis, subtilissimè concentricè plicato-lamellatis, columellā contortā, subtruncatā, lirā parvā superne muricatā; pellucido-cornēā.*

THE LAMELLOSE ACHATINA. Shell acuminate-ovate, whorls six in number, somewhat rounded, very finely slightly plicately lamellated, columella twisted, slightly truncated, furnished at the upper part with a small ridge; transparent horny.

MORICAND, —?

Hab. —?

A very transparent shell, most delicately concentrically lamellated.

Species 108. (Mus. Cuming.)


ACHATINA ANOMALA. *Achat. testā turritā, minulā, umbilicatā, spire suturis impressis, aufractibus septem ad octo, rotundatis, striatulis, columellā brevi, callosā, reflexā, aperturā parvā; sordidè albicans.*

THE ANOMALOUS ACHATINA. Shell turreted, minutely umbilicated, sutures of the spire impressed, whorls seven to eight in number, rounded, finely striated, columella short, callous, reflected, aperture small: dull whitish.

PFEIFFER, Symbole, vol. iii. p. 89. Philippi, Abbild. *Achat.* pl. 1. f. 12.

Hab. Peru.

A small turreted shell, with the sutures of the spire rather deeply impressed, verging very closely on *Bulimus* by reason of the reflected growth of the columella.



ACHATINA.

PLATE XX.

Species 109. (Mus. Benson.)

ACHATINA BALANUS. *Achat. testā cylindraceo-oblongā, subfusiformi, anfractib⁹ quatuor, lævibus, apice obtuso, columellā arcuatā, truncatā, aperturā parvā; sordidè albā.*

THE BARNACLE ACHATINA. Shell cylindrically oblong, somewhat fusiform, whorls four in number, smooth, shining, apex obtuse, columella arched, truncated, aperture small; dull white.

BENSON, MSS.

Hab. Banks of the Jumna, India.

A very minute species, allied to the following, but composed of fewer whorls. First collected by Mr. Benson in 1825 on the banks of the river Jumna near Hameerpore, Bundelkund, among the roots of a large fallen tree; subsequently on a porphyritic hill on the border of the desert south of Hawee. It occurs also at Agra, on the right bank of the Jumna.

Species 110. (Mus. Cuming.)

ACHATINA ABERRANS. *Achat. testā subulatā, tenui, anfractibus sex, longitudinaliter impresso-striatis, columellā tortuosa, callosa-marginatā, aperturā parvā; pellucida, vitrea, striis rufulis angustis obliquè pictā.*

THE ABERRANT ACHATINA. Shell subulate, thin, whorls six in number, longitudinally impressly striated, columnella twisted, callous at the edge, aperture small; transparent, glassy, obliquely painted with narrow reddish streaks.

PFEIFFER, Pro. Zool. Soc. 1845, p. 138.

Hab. Jamaica; Gosse.

A pretty little glassy shell, striped at rather distant intervals with red-brown lines.

Species 111. (Mus. Cuming.)

ACHATINA ACICULA. *Achat. testā cylindraceo-subulatā, subfusiformi, anfractibus sex, lævibus, politis, apice obtuso, columellā arcuatā, angustè truncatā; albē, diaphana.*

THE LITTLE NEEDLE ACHATINA. Shell cylindrically subulate, somewhat fusiform, whorls six in number, smooth, polished, apex obtuse, columella arched, narrowly truncated; white, diaphanous.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. ix. p. 304

Buccinum acicula, Müller.

Buccinum terrestre, Montagu.

Helix (Cochlicopa) acicula, De Féussac.

Bulinus acicula, Bruguière.

Achatina alba, Brown.

Cionella acicula, Jeffreys.

Acicula ebarnea, Risso.

Hab. Europe, including the British Isles.

A light, shining, glassy shell, with a distinctly truncated columnella.

Species 112. (Mus. Cuming.)

ACHATINA COSTULATA. *Achat. testā turritā, anfractibus octo, rotundatā, longitudinaliter costulatā, columellā subrecti, truncatā, aperturā parvā; pellucido-cornēa, striis rufo-fuscis irregulariter pictā.*

THE DELICATELY-RIBBED ACHATINA. Shell turreted, whorls eight in number, rounded, longitudinally finely ribbed, columnella nearly straight, truncated, aperture small; transparent horny, irregularly painted with red-brown streaks.

ADAMS, Pro. Nat. Hist. Soc. Boston, 1845, p. 13.

Achatina Adamsi, Pfeiffer.

Hab. Jamaica.

A finely ribbed, truncated shell, of a brownish horn-colour, streaked irregularly with darker brown.

Fig. 113. (Mus. Cuming.)

For description of this species, see *Bulinus*, Pl. LXX.
For *Achatina*, Sp. 113, see Pl. XXI.

Species 114. (Mus. Cuming.)

ACHATINA GOSSEI. *Achat. testā subulatā, aciculari, anfractibus decen, subrenatō costatis, costis basia versus gradatim evanidis, juxta suturas fortioribus, columellā obliquè arcuatā, vix truncatā, aperturā minatā, subdilatotā; sordidè albā.*

GOSSE'S ACHATINA. Shell subulate, finely pointed, whorls ten in number, somewhat remotely ribbed, ribs gradually fading towards the base, stronger next the sutures, columnella obliquely arched, scarcely truncated, aperture very small, rather dilated; dead white.

ACHATINA.—PLATE XX.

PFEIFFER, Pro. Zool. Soc. 1845, p. 138.

Hab. Jamaica; Gosse.

An interesting delicately ribbed shell, distinguished by its sharp aciculate form.

Species 115. (Mus. Cuming.)

ACHATINA CALIFORNICA. *Achat. testā cylindraceo-elongatā, anfractibus tredecim, obliquè creberimè striatis, basi versus subangulato-planatis, ultimo basi acutè carinato, infra carinam peculiariter excavato, columellā valde arcuata, callosa-marginatā, aperturā parvā, subquadratā; cereo-albicante.*

THE CALIFORNIAN ACHATINA. Shell cylindrically elongated, whorls thirteen in number, obliquely very closely striated, rather angularly flattened towards the base, last whorl sharply keeled at the base, peculiarly excavated beneath the keel, columella very much arched, callous at the edge, aperture small, somewhat square; waxen white.

PFEIFFER, Symbolæ, vol. iii. p. 89.

Hab. Monterey, California.

This very remarkable species is mainly distinguished by the excavated and keeled growth of each whorl at the base.

Fig. 116. (Mus. Cuming.)

ACHATINA ACCINCTA. Already described and figured from a lighter specimen in the preceding plate, Sp. 101. For Sp. 116, see Pl. XXIII.

Fig. 117. (Mus. Cuming.)

ACHATINA LUBRICA. Referred to the genus *Bulimus*, Sp. 645. For Sp. 117, see Pl. XXIII.

Species 118. (Mus. Cuming.)

ACHATINA SUTURALIS. *Achat. testā oblongo-ovatā, spirā breviusculā, conicā, suturis impressis, anfractibus quinque, convexis, laevis, politis, columellā subrectā, truncatā; cornēd, solidiusculā, pellucida.*

THE SUTURAL ACHATINA. Shell oblong-ovate, spire rather short, conical, sutures impressed, whorls five in number, convex, smooth, polished, columella nearly straight, truncated; horny, rather solid, transparent.

Polypheus suturalis, Pfeiffer, in Wiegman Archiv. 1839, p. 353.

Glandina suturalis, Philippi.

Hab. Cuba.

An oblong transparent polished shell, belonging to the group *Glandina* of Dr. Philippi.



26



27



28



29



30



31



32



33



ACHATINA.

PLATE XXI.

Species 77. (*Bulimus*, Fig. 572. Mus. Benson.)

ACHATINA EXULATA. *Achat. testā oblongā, subcylindracē, anfractibus sex ad septem, convexis, obliquū subtilliter striatis, columellā arcuatā, subcontortā, obtuse truncatā, aperturā porciusculā, peculiariter effusā; pellucido-albā, opaco-albo sparsim variegatā.*

THE EXILED ACHATINA. Shell oblong, somewhat cylindrical, whorls six to seven in number, convex, obliquely finely striated, columella arched, somewhat twisted, obtusely truncated, aperture rather small, peculiarly effused; transparent white, sparingly mottled with opaque white.

BENSON, MS.

Hab. St. Helena; Lieut. Lefroy.

Of a very distinct type from any other species of the genus.

Species 80. (Mus. Benson.)

ACHATINA JERDONI. *Achat. testā subpyramidalē-oblongā, anfractibus septem ad octo, convexis, levibus, poitis, ad suturas subtilliter marginatis, columellā arcuatā, conspicue truncatā, aperturā parvā; fuso-cornēd.*

JERDON'S ACHATINA. Shell somewhat pyramidal oblong, whorls seven to eight in number, convex, smooth, polished, finely margined at the sutures, columella arched, conspicuously truncated, aperture small; brown horny.

BENSON, MS.

Hab. Nilgherries, Hindooostan; Jerdon.

A polished horny species, of rather narrow, pyramidal growth.

Species 81. (Mus. Benson.)

ACHATINA CRASSILABRIS. *Achat. testā subpyramidalē-conicē, basia versus ventricosā, anfractibus octo, subrotundatis, ad suturas crenulatis, crenulis numerosis, confertis, in striis impressis descendentibus, columellā profunde arcuatā, conspicue truncatā, aperturā parvissimā, labro subincurvato; cornē-fuscā.*

THE THICK-LIP ACHATINA. Shell somewhat pyramidal conical, ventricle towards the base, whorls eight in number, rather rounded, crenulated at the sutures, crenules numerous, close-set, descending in impressed striae, columella deeply arched, conspicuously trun-

cated, aperture rather small, lip slightly thickened; horny brown.

BENSON, Journ. Asiatic Soc. 1836, vol. v. p. 353.
Hab. North-east Frontier of Bengal; Benson.

Distinguished by its rounded, conical form, finely impressed crenulated sculpture, and slightly thickened lip.

Species 87. (Mus. Benson.)

ACHATINA NILGARICA. *Achat. testā pyramidali-turritā, anfractibus decem, convexis, ad suturas subtillissimē crenulatis, deinde striatis, columellā subprofundē arcuatā, aperturā parvā; fuso-cornēd.*

THE NILGHERRY ACHATINA. Shell pyramidal turreted, whorls ten in number, convex, very finely crenulated at the sutures, then striated, columella rather deeply arched, aperture small; brown-horny.

BENSON, MS.

Hab. Nilgherries, Hindooostan; Jerdon.

Allied to the preceding species in detail of character, but amply distinguished by its more elongated pyramidal form.

Species 88. (Mus. Cuming.)

ACHATINA LINDONI. *Achat. testā cylindracē-oblongā, utrinque attenuatā, solidatā, anfractibus octo, planulatis, levigatis, lineis crenatis incrementi leviter impressis, ad suturas marginatis, duobus ultimis obliquē descendantibus, columellā basi arcuatā, suprā gibbosā, aperturā angustā, suprē sinuata; fulvo-cornēd, pallidā, strigis castaneis arcatis obscurē lineata.*

LINDON'S ACHATINA. Shell cylindrically oblong, attenuated at each end, rather solid, whorls eight in number, rather flattened, smooth, slightly impressed with curved lines of growth, margined at the sutures, the last two whorls descending obliquely, columella arched, gibbosus above, aperture narrow, situated at the upper part; fulvous horny, polished, obscurely stained with arched chestnut streaks.

PFEIFFER, Pro. Zool. Soc. 1846, p. 116.

Hab. Cuba; Lindon.

Of a compressed cylindrical form, belonging to the *Glaucina* type.

Species 95. (Mus. Cumig.)

ACHATINA ISABELLINA. *Achat. testā fusiformi-oblongā, anfractibus sex, convexis, sub lente minutissimè decussatis striatis, ad suturas marginatis, columellā obliquā, sublatē truncatā, aperturā ovaē; pellucidō cornēā, sordidē albicante.*

THE SOILED ACHATINA. Shell fusiformly oblong, whorls six in number, convex, very minutely decussately striated beneath the lens, marginated at the sutures, columella oblique, rather broadly truncated, aperture ovoid; transparent horny, dull whitish.

PFEIFFER, Pro. Zool. Soc. 1846, p. 31.
Hab. Mexico.

A shell of rather firm growth, though of transparent substance.

Species 102. (Mus. Benson.)

ACHATINA PERROTTETI. *Achat. testā pyramidalī-conicā, anfractibus octo, rotundatis, levibus, columellā subpropende arcuatā, aperturā parvā; fuso-corneā.*

PERBOTTE'S ACHATINA. Shell pyramidaly conical, whorls eight in number, rounded, smooth, columella

rather deeply arched, aperture small; brownish horny.

PFEIFFER, Revue Zool. Cuv., 1842, p. 305.
Hab. Nilgherries, Hindooostan; Jerdon.

Very closely allied to *A. crassilabris*, from which it mainly differs in being of a rather more elongated form, and not impressly crenulated.

Species 113. (Mus. Benson.)

ACHATINA OREAS. *Achat. testā oblongo-conicā, anfractibus septem, convexis, ad suturas subtiliter marginatis, longitudinaliter obscurè impresso-striatis, columellā arcuatā, aperturā parvā; fuso-corneā.*

THE MOUNTAIN-NYMPH ACHATINA. Shell oblong-conical, whorls seven in number, convex, finely marginated at the sutures, longitudinally obscurely impressly striated, columella arched, aperture small; brown horny.

BENSON, MS.

Hab. Nilgherries, Hindooostan; Jerdon.

Another interesting species of the Nilgherry type, approaching to the form of *A. Jerdoni*.

Achatina. Pl. XXXII.



125.



127.



129.



126.



128a.



128b.



122.



125.



123b.



124.

ACHATINA.

PLATE XXII.

[Figures considerably magnified.]

Species 116. (Mus. Cuming.)

ACHATINA TRITICEA. *Achat. testā ovo-cylindracea, solidā, anfractibus sex, levibus, politis, infra suturas lineā circumdatā, columellā plerunque bīplicatā, vix truncatā, aperturā basi rotundata et effusa, superne leviter sinuatā; fulvescente-cornēa.*

THE WHEAT-GRAIN ACHATINA. Shell ovately cylindrical, rather solid, whorls six in number, smooth, polished, surrounded with a line beneath the sutures, columella generally two-plaited, scarcely truncated, aperture rounded and effused at the base, slightly sinuated at the upper part; fulvous horny.

Helix triticea, Lowe, Faun. Mader. p. 90. pl. 6. f. 26.

Glandina triticea, Pfeiffer.

Hab. Porto Sancto, Madeira; Lowe.

This species, in which the columellar plaits are sometimes wanting, partakes of the character of the genus *Tornatellina*.

Species 117. (Mus. Cuming.)

ACHATINA GRACILIS. *Achat. testā acuminato-oblongā, anfractibus quinque, levibus, politis, ad suturas subtilliter marginatis, columellā levigatā, vix truncatā; pellucido-vitrēa, virescens.*

THE SLENDER ACHATINA. Shell acuminately oblong, whorls five in number, smooth, polished, finely margined at the sutures, columella smooth, scarcely truncated; transparent glassy, greenish.

Helix gracilis, Lowe, Faun. Mader. p. 61. pl. 6. f. 28.

Hab. Porto Sancto, Madeira; Lowe.

A small transparent glassy species, very simply characterized.

Species 119. (Mus. Cuming.)

ACHATINA OVALIFORMIS. *Achat. testā oblongo-ovatā, anfractibus quatuor, levibus, politis, suturis subimpressis, columellā tenuē bīplicatā, obliquē truncatā; pellucido-vitrēa, fuscenscente.*

THE LITTLE EGG-SHAPED ACHATINA. Shell oblong-ovate, whorls four in number, smooth, polished, sutures rather impressed, columella thinly two-

plaited, obliquely truncated; transparent glassy, brownish.

Helix ovaliformis, Lowe, Faun. Mader. p. 61. pl. 6. f. 27.
Hab. Porto Sancto, Madeira; Lowe.

Less acuminate in form and browner in colour, whilst the columella is plaited, after the manner of *Tornatellina*.

Species 120. (Mus. Benson.)

ACHATINA CRASSULA. *Achat. testā pyramidali-conica, anfractibus septem, plano-convexis, longitudinaliter impresso-striatis, suturis excavatis, columellā arcuata, conspicue truncatā, aperturā pareā; albida, epidermide olivaceo-cornēa indutā.*

THE THICKENED ACHATINA. Shell pyramidal conical, whorls seven in number, flatly convex, longitudinally impressively striated, sutures excavated, columella arched, conspicuously truncated, aperture small; whitish, covered with an olive horny epidermis.

BENSON, MS.

Hab. Himalaya.

Of more solid growth than the Nilgherry species of allied form, represented in the preceding plate.

Species 121. (Mus. Cuming.)

ACHATINA CORUSCA. *Achat. testā pyramidalis, subfusiformi, apice obtusā, anfractibus septem, convexis, levibus, politis, ad suturas subtilliter marginatis, columellā arcuata et contortā, aperturā ovatā; pellucido-fuscenscente, vitrēa, strigis castaneis flexuosis obscurè tinctā.*

THE GLITTERING ACHATINA. Shell pyramidal, somewhat fusiform, obtuse at the apex, whorls seven in number, convex, smooth, polished, finely margined at the sutures, columella arched and twisted, aperture ovate; transparent brown, glassy, obscurely stained with flexuous chestnut streaks.

Hab. — ?

A shell of firm growth, though transparent and glassy.

Species 122. (Mus. Cuming.)

ACHATINA MELAMPOIDES. *Achat. testā oblongo-ovata,*

solidiusculā, spirā brevi, anfractibus sex, lēvibus, suturis indistinctis, līneis circumdati, columellā excavatā et contortā, parū truncatā, aperturā oblongā, supernē leviter sinuatā, basi peculiariter effusā; ciuere-fuscā, subobliqueā, columellā albā.

THE MELAMPUS-LIKE ACHATINA. Shell oblong-ovate, rather solid, spire short, whorls six in number, smooth, sutures indistinct, surrounded by a line, columella excavated and twisted, but little truncated, aperture oblong, slightly sinuated at the upper part, peculiarly effused at the base; ash-brown, olive tinged, columella white.

LOWE, MS.

Hab. Madeira.

A shell of rather solid growth, having the form of a *Melampus* or *Auricula*.

Species 123. (Mus. Benson.)

ACHATINA GEMMA. *Achat. testā oblongo-conicā, solidiusculā, anfractibus sex, rotundatis, lēvibus, columellā arcuatā, abbreviatā, aperturā subrotundā; purpureo-nigricante, nitente.*

THE BUD ACHATINA. Shell oblong-conical, rather solid, whorls six in number, rounded, smooth, columella

arched, abbreviated, aperture nearly round; purple-black, shining.

BENSON, MS.

Hab. Barrackpore, Bengal; Benson, Bacon.

Of a much darker purple-black colour than any other of the Indian species, and of shorter growth, with a shining, polished surface.

Species 124. (Mus. Cuming.)

ACHATINA FRUMENTUM. *Achat. testā oblongo-conicā, solidiusculā, anfractibus sex, convexis, lēvibus, columellā arcuatā et contortā, aperturā subrotundā; fulvo-spaticeā.*

THE RYE-GRAIN ACHATINA. Shell oblong-conical, rather solid, whorls six in number, convex, smooth, columella arched and twisted, aperture nearly round; fulvous bay.

Hab. Chandpore, Bengal; Bacon.

This approaches very closely to the preceding species, yet there is a characteristic difference in the appearance of the shell, of which Mr. Cuming possesses many specimens. The whorls are less rounded, less polished, and of a uniform lighter colour.



ACHATINA.

PLATE XXIII.

Species 125. (Mus. Taylor.)

ACHATINA MARMOREA. *Achat. testá elongato-turrítá, au-*
gustá, subcylindráceá, solidiusculá, anfractibús plano-
convéxis, obliquè subrude stríatís, columellá tenuē trun-
cátá, aperturá parvá; albá, epidermide cornéa cas-
tanea inadúta.

THE MARBLE ACHATINA. Shell elongately turreted, narrow, somewhat cylindrical, rather solid, whorls flatly convex, obliquely rather rudely striated, columella thinly truncated, aperture small; white, covered with a chestnut horny epidermis.

Hab. —?

This species resembles the *Bulinus calcar* in form. It is of more solid growth, and has the columella distinctly truncated.

Species 126. (Mus. Cuming.)

ACHATINA MUCIDA. *Achat. testá ovalá, tenui, subven-*
tricosá, anfractibús sex, decussati subtilissimè granu-
losso-stríatás, superne marginatás, et obscurè crenulatás,
columellá tenuissimè truncatá; opaco-albo et pur-
pureo-fusco strigatá et variegatá, epidermide tenui-
inadúta.

THE MOULDY ACHATINA. Shell ovate, thin, rather ventricose, whorls six in number, decussately very finely granulately striated, marginated round the upper part, and obscurely crenulated, columella very thinly truncated; streaked and variegated with opaque white and purple-brown, covered with a thin epidermis.

Bulinus mucidus, Gould.

Hab. Liberia, West Africa.

A light, strongly painted shell, of which the colouring is seen through in the interior.

Species 127. (Fig. Guérin.)

ACHATINA SAULCYDI. *Achat. testá oblongo-ovalá, ven-*
tricosá, sinistrá, apice subpapillari, anfractibús con-
vexis, irregulariter rugoso-stríatás, ad suturas crenula-
tis, columellá subexcavatá et truncatá; violaceo-griseá,
nigrigante-fusco longitudinaliter lineatá et strigatá,
aperture fance ferrugineo-falcá.

SAULCYD'S ACHATINA. Shell oblong-ovate, ventricose, sinistral, somewhat papillary at the apex, whorls convex, irregularly roughly striated, crenulated at the sutures, columella slightly excavated and truncated; violet-grey, longitudinally streaked and lined with blackish brown, aperture rusty-fuscous.

DE JOANNIS, Guérin's Mag. de Zool. 1834, Moll. pl. 50.
Hab. Prince's Island, West Africa.

I have not seen this species, but have ventured to copy the excellent figure of it in the work above referred to, on account of its characteristic appearance. The papillary type of the apex appears also in *Bulinus torridus*, from the neighbouring locality of Liberia.

Species 128. (Mus. Cuming.)

ACHATINA GOULDII. *Achat. testá ovalo-conicá, tenui,*
ventricosá, apice subpapillari, anfractibús plano-con-
vexis, obliquè tenuissimè striatás, ad suturas subdilatér
crenulatás, anfractu ultime basi leciter angulato; ful-
vescente-olivaceá, epidermide tenui inadúta, epidermide
secundá hydrophaná prope apicem radiatá et infre
angulum anfractiú ultimi unifasciatá.

GOULD'S ACHATINA. Shell ovately conical, thin, ventricose, somewhat papillary at the apex, whorls flatly convex, obliquely very finely striated, finely crenulated at the sutures, last whorl slightly angled at the base; fulvous-olive, covered with a thin epidermis, rayed at the apex with a second hydrophanous epidermis, and encircled with a single band of the same beneath the angle of the last whorl.

Bulinus balteatus, Gould.

Hab. Liberia, West Africa.

The name *balteatus* being already occupied in this genus, I have the pleasure of naming the species after the accomplished naturalist to whom I am indebted for it,— Dr. Gould, of Boston, United States.

Species 129. (Mus. Taylor.)

ACHATINA OBELISCUS. *Achat. testá elongato-turrítá, an-*
fractibus novem ad drem, planis, medio subconcavis,
transversim obscurè et irregulariter liratis, anfractu
ultimo basi obtuse angulato, columellá arcuatá et

ACHATINA.—PLATE XXIII.

truncatā; sordidē fusco-albā, epidermide fulvescente-fusca cornēā indutā, anfractū ultimo circa columellam castaneo-nigricante fasciato.

THE OBELISK ACHATINA. Shell elongately turreted, whorls nine to ten in number, flattened, slightly concave in the middle, transversely obscurely and irregularly ridged, last whorl obtusely angled at the base, columella arched and truncated; dirty

brownish-white, covered with a horny fulvous-brown epidermis, last whorl banded with chestnut-black around the columella.

Hab. ——?

For this remarkable and very characteristic species I am indebted to the collection of Thomas Lombe Taylor, Esq.

ACHATINA.

Plate.	Species.	Plate.	Species.		
aberrans, <i>Pfr.</i>	XX.	110	cylindracea, <i>Pfr.</i>	XVIII.	91
accincta, <i>Mighels</i>	XIX.	101	<i>dactylus</i> , Brod.	VIII.	26
acicula, <i>Lamarek</i>	XX.	111	Dennisoni, <i>Reeve</i>	IX.	32
acuta, <i>Férussac</i>	III.	11	Dominicensis (<i>Helix</i>), <i>Fér.</i>	XIV.	56
<i>Adamsi</i> , <i>Pfr.</i>	XX.	112	Dysoni, <i>Pfr.</i>	XIX.	99
<i>adusta</i> , <i>Gould</i>	I.	4	erecta, <i>Benson</i>	XVI.	69
aquatoria, <i>Reeve</i>	I.	2	exulata, <i>Benson</i>	XXI.	77
alabaster (<i>Helix</i>), <i>Rang</i>	IX.	28	fasciata (<i>Bucc.</i>), <i>Müller</i>	X.	35
alba, <i>Brown</i>	XX.	111	flammea, <i>Férussac</i>	XII.	39
abolineata, <i>Lamarek</i>	XIV.	52	Fraseri, <i>Pfr.</i>	XVI.	71
Algira, <i>Brug.</i>	XIII.	43	frumentum, <i>Reeve</i>	XXII.	124
allisa, <i>Reeve</i>	V.	16	fulica, <i>Férussac</i>	II.	8
amentum, <i>Benson</i>	XVII.	82	fulva (<i>Bulinus</i>), <i>Brug.</i>	III.	10
amphora, <i>Jay</i>	IV.	14	Funeki, <i>Pfr.</i>	XVIII.	96
Annis, <i>Lesson</i>	X.	35	fusco-lineata, <i>Lamarek</i>	XIV.	56
anomala, <i>Pfr.</i>	XIX.	108	fusiformis, <i>Pfr.</i>	IX.	31
arcuata, <i>Pfr.</i>	XVII.	83	gemma, <i>Benson</i>	XXII.	123
assimilis, <i>Reeve</i>	XV.	67	glabra (<i>Gland.</i>), <i>Pfr.</i>	XIV.	58
balanus, <i>Benson</i>	XX.	109	glaea, <i>Lamarek</i>	XIV.	55
balteata, <i>Reeve</i>	II.	7	Gosschi, <i>Pfr.</i>	XX.	114
bicarinata (<i>Bulle</i>), <i>Dillw.</i>	V.	17	Gouldii, <i>Reeve</i>	XXIII.	128
bicolor, <i>Jay</i>	I.	4	gracilis, <i>Loeve</i>	XXII.	117
Bulimoides, <i>Pfr.</i>	XIX.	103	Griffithsii, <i>Adams</i>	XV.	64
Californica, <i>Pfr.</i>	XX.	115	Hugeli, <i>Pfr.</i>	XV.	68
carinata, <i>Pfr.</i>	VII.	24	incerta, <i>Reeve</i>	XVIII.	90
Cassica, <i>Benson</i>	XVII.	85	indotata, <i>Reeve</i>	VI.	18
Ceylanica, <i>Pfr.</i>	XV.	59	isabellina, <i>Pfr.</i>	XXI.	95
elavata, <i>Gray</i>	XVI.	72	Jamaicensis, <i>Pfr.</i>	XV.	61
elavus, <i>Pfr.</i>	XVI.	75	Javanica, <i>Reeve</i>	XVII.	79
cochlea, <i>Reeve</i>	I.	5	Jerdoni, <i>Benson</i>	XXI.	80
columna (<i>Bucc.</i>), <i>Müller</i>	XI.	38	Kraussi, <i>Reeve</i>	VI.	21
concentrica, <i>Reeve</i>	XIX.	106	Iactea, <i>Reeve</i>	XII.	41
conifera, <i>Férussac</i>	XVIII.	98	Lamarekiana, <i>Pfr.</i>	III.	12
consimilis, <i>Reeve</i>	XIX.	104	lamellata, <i>Pöliez</i> and <i>Michaud</i>	XVIII.	97
corusca, <i>Reeve</i>	XXII.	121	lamellosa, <i>Moricand</i>	XIX.	107
costulata, <i>Adams</i>	XX.	112	Lattrei, <i>Pfr.</i>	XIV.	53
costulata, <i>Pfr.</i>	VI.	20	leucozonias (<i>Voluta</i>), <i>Waleh</i>	XIV.	52
Couroupa, <i>Lesson</i>	II.	8	lignaria, <i>Reeve</i>	VIII.	27
crassilabris, <i>Benson</i>	XXI.	81	Lindomi, <i>Pfr.</i>	XXI.	88
crassula, <i>Benson</i>	XXII.	120	lineata, <i>Valenciennes</i>	X.	35
crenata, <i>Swainson</i>	X.	35	magnifica, <i>Pfr.</i>	IX.	33
curlabilis, <i>Pfr.</i>	XVIII.	94	marginata, <i>Schaubon</i>	IV.	14
cyanostoma, <i>Rüppell</i>	XIII.	44	marmorea, <i>Reeve</i>	XXIII.	125

INDEX.

	Plate.	Species.		Plate.	Species.
Melampoides, <i>Love</i>	XXII.	122	reticulata, <i>Pfr.</i>	II.	9
monilifera (<i>Gland.</i>), <i>Pfr.</i>	XIV.	50	<i>Rickii</i> , <i>Reeve</i>	X.	39
mucida (<i>Bulimus</i>), <i>Gould</i>	XXIII.	126	<i>rosea</i> , <i>Férussac</i>	XIII.	46
Mülleria, <i>Férussac</i>	VIII.	25	<i>Sandwicensis</i> , <i>Pfr.</i>	XIX.	101
murrea, <i>Reeve</i>	VII.	22	<i>Saulcydi</i> , <i>De Joannis</i>	XXIII.	127
nigricans (<i>Gland.</i>), <i>Pfr.</i>	XIV.	54	semisculpta, <i>Pfr.</i>	XI.	37
Nilagericus, <i>Benson</i>	XXI.	87	semistarum (<i>Helix</i>), <i>Rang</i>	XVI.	73
obeliscus, <i>Reeve</i>	XXIII.	129	sideratus, <i>Reeve</i>	XII.	38
obtusa (<i>Gland.</i>), <i>Pfr.</i>	XV.	62	<i>sinistrorsa</i> , <i>Pfr.</i>	V.	17
oetona, <i>Gray</i>	XVII.	84	<i>solidula</i> , <i>Pfr.</i>	XVIII.	92
oleacea (<i>Helix</i>), <i>Férussac</i>	XIV.	51	<i>Sowerbyana</i> , <i>Pfr.</i>	VIII.	26
Orcas, <i>Benson</i>	XXI.	113	<i>splendida</i> , <i>Audou</i>	XIX.	100
ornata, <i>Pfr.</i>	VI.	20	<i>striata</i> (<i>Bucc.</i>), <i>Müller</i>	VI.	19
orophila, <i>Benson</i>	XIX.	105	<i>striatella</i> (<i>Helix</i>), <i>Rang</i>	XVI.	74
orysacea, <i>D'Orbigny</i>	XVIII.	89	<i>subulata</i> , <i>Pfr.</i>	XVIII.	89
ovuliformis, <i>Love</i>	XXII.	119	<i>sulecta</i> , <i>Gray</i>	XVII.	86
pallida, <i>Swainson</i>	X.	35	<i>suturalis</i> , <i>Pfr.</i>	XX.	118
panthera, <i>Férussac</i>	III.	12	<i>tenuispira</i> , <i>Benson</i>	XVI.	76
papyracca, <i>Pfr.</i>	II.	6	<i>tineta</i> , <i>Reeve</i>	XI.	29
pixillus, <i>Reeve</i>	XVI.	77	<i>Tortillana</i> , <i>Pfr.</i>	XV.	66
perdix, <i>Lamarek</i>	I.	3	<i>triticea</i> , <i>Love</i>	XXII.	116
Perroteti, <i>Pfr.</i>	XXI.	102	<i>truncata</i> (<i>Gland.</i>), <i>Say</i>	XIII.	47
Peruviana, <i>Lamarek</i>	XIV.	57	<i>turris</i> , <i>Pfr.</i>	XIII.	45
Pfeifferi, <i>Dauker</i>	I.	1	<i>ustulata</i> , <i>Lamarek</i>	XII.	40
Philippiana (<i>Gland.</i>), <i>Pfr.</i>	XIV.	49	<i>Vanuxemensis</i> , <i>Lea</i>	XIII.	48
Phillipsii, <i>Adams</i>	XV.	60	<i>variegata</i> , <i>Roissy</i>	I.	3
pieta, <i>Reeve</i>	X.	34	<i>venusta</i> (<i>Gland.</i>), <i>Pfr.</i>	XV.	63
Poirieri, <i>Rossmoser</i>	XIII.	43	<i>vezillum</i> , <i>Lamarek</i>	X.	35
Porto-Ricensis, <i>Pfr.</i>	XVIII.	93	<i>virginea</i> (<i>Bulla</i>), <i>Limn</i>	X.	36
prunum, <i>Reeve</i>	IV.	13	<i>vittata</i> , <i>Swainson</i>	X.	36
purpurea (<i>Bulla</i>), <i>Chemn</i>	IV.	15	<i>vivipara</i> , <i>Sowerby</i>	XVI.	70
Ranguana, <i>Pfr.</i>	XV.	65	<i>voluta</i> (<i>Bulla</i>), <i>Chemn</i>	XIV.	55
Reeveana, <i>Pfr.</i>	IX.	29	<i>zebra</i> (<i>Bulla</i>), <i>Chemn</i>	VII.	23

MONOGRAPH
OF THE GENUS
DOLIUM.

Ocean, in thy unfathomable depths
What beauties lie concealed from wondering eyes?
Pearls that adorn the neck of beauty; shells
On which the sea-born Venus finds a couch -- *Bryant*



DOLIUM.

PLATE I.

Genus DOLIUM, Lamarck

Test. tenuis, globoso-omnida, ventricosissima, transversim costata, unquam longitudinaliter exculpta, aperturā angusta, labro simplici, crenato, raro fimbriato, labro columellarī lato expanso. Operculum nullum.

Shell thin, globosely swollen, extremely ventricose, transversely ribbed, never longitudinally sculptured, aperture large, lip simple, crenated, rarely fimbriated, columellar lip broadly expanded. No operculum.

The *Dolia* or 'Tuns' are mainly distinguished from other genera of the great family of *Parupifera* to which they belong, by their large globosely inflated growth, transverse ribs, and absence of any sort of longitudinal sculpture, presenting a singular contrast in this respect with their allies, the 'Harps,' in which ribs only occur in a longitudinal direction.

As may be gathered from the very light structure and ample evolution of the shell, its animal inhabitant is distinguished for its large size and muscular strength, as well as by its voracity and comparative activity of habit. The head is furnished with a long retractile proboscis which the animal is capable of rapidly protruding or withdrawing, furnished at the extremity with a kind of rosette, as represented in the beautiful figure of *D. perdis* in the Voyage de l'Astrolabe by MM. Quoy and Gaimard. The disk or foot of *Dolium* is large and fleshy and, according to M. Deshayes, becomes inflated with water, when the animal desires to swim, imbibing the fluid through certain pores in the skin. The colours of the animal present a brilliant contrast with those depicted in the shell; the *D. perdis* is striped with light azure blue, the *D. pomaia* is also light blue, and the *D. olearina* is of a rich coppery green.

The *Dolia* are partially distributed, although the *D. perdis* occurs equally on the shores of the Philippine Islands and in the West Indies. The *D. pomaia* is found also both in the Philippine and Society Islands. This and the *D. ringens* from Peru are of more solid growth than the rest, and peculiar in having a hollow excavated in the columella, on which account they have been separated

from the genus by M. Valenciennes under the title of *Malea*. The *D. galea* of the Mediterranean is remarkable for its large size, but the *D. variegatum* has been recently collected at Torres Straits, North Australia, of even more gigantic proportions.

The shells of this genus, though not much esteemed by collectors, are by no means common; only fourteen species are known, and of these, but one new one has resulted from the researches of Mr. Cuming or Sir E. Belcher.

Species 1. (Mus. Cuming.)

DOLIUM GALEA. *Dol. testā maximā, eucratosissimā, umbilicatā, spirā brevi, paullatū subsidente, suturis profundè impressis, aufrictibus septem, superne tenuidis, transversim costatis, costis convexis, confertis, superioribus lirā sepiissimè interveniente, columellā subcavatā; albida aut pallidè fulvā, apice nigricante, labro columellarī albā, labro intus fuso tincto.*

THE HELMET TUN. Shell very large, extremely ventricose, umbilicated, spire short, a little sunk in, sutures deeply impressed, whorls seven in number, swollen round the upper part, transversely ribbed, ribs convex, close-set, upper ones with most frequently an intervening ridge, columella somewhat twisted; whitish or pale fulvous colour, apex blackish, columellar lip white, lip stained within with brown.

LAMARCK. Anim. sans vert. (Deshayes' edit.) vol. x. p. 139.
Buccinum galea, Linnaeus.

An eadem var.?

Dolium ampullaceum, Philippi.

Testa juvenis.

Dolines tenuis, Menke.

Hab. Mediterranean.

The above recorded synonymes of this well-known species have arisen out of the circumstance of its having a more oblong form in an early stage of growth, and an occasional irregularity in the development of the ribs.



DOLIUM.

PLATE II.

Species 2. (Mus. Brit.)

DOLIUM (ELANOSTOMA). *Dol. testa subpyriforme-turbinata, nubilatim, ventricosissimam, spiris subpropria-anam, unfractibus septem, transversum costatis, costis subdepresso-convexis, interstitiis liri paré subsidente undique rugulis, columellâ latè expansâ, aperturâ ampliâ, basi subprofundiè emarginatâ; latescere-albidâ, levigata, sileante, columellâ intense nigricante-brunneâ.*

THE BLACK-MOUTH TUN. Shell somewhat globosely turbinated, umbilicate, very ventricose, spire rather prominent, whorls seven in number, transversely ribbed, ribs somewhat depresso convex, interstices filled throughout with a small ridge, columella widely expanded, aperture large, rather deeply emarginated at the base; yellowish white, smooth, shining, columella deep blackish brown.


JAY, Cat. Mus. p. 124. pl. 8-9.
Hub. Friendly Islands; Jay.

The shell here represented, from the British Museum,

belonged to the late Earl of Tankerville, and is the individual specimen noted by Mr. Sowerby in his well-known Catalogue of that nobleman's collection, as *Dolium galba* var. *d.* I believe it to be distinct from that species, which is an inhabitant of the Mediterranean, and to be identical with a shell from the Friendly Islands in the collection of Dr. Jay of New York, described by that gentleman, under the name above recorded, in his published 'Catalogue' of 1839. "This splendid shell," says Dr. Jay, "was brought to Nantucket by a whaler from the Friendly Islands, where it was obtained by the natives in diving for shells."

The *D. melanostoma* is chiefly distinguished from the *D. galba* in having a more prominent spire and a regularly developed ridge throughout the interstices between the ribs, which have a lighter and more polished surface. The base of the shell beneath the umbiliens is characterized by its numerous concentric striae, whilst in the *D. galba* it is ridged in a longitudinal direction; and the rich dark colouring of the columellar lip appears also to be a character of the species.

December, 1848.



DOLIUM.

PLATE III.

Species 3. (Fig. *a* and *b*, Mus. Cuming.)

DOLIUM FIMBRIATUM. *Dol. testá globosá, ventricosissimá, vir umbilicatá, spirá suturis impressa, anfractibus sex, conspicuè costatis, costis plus minusve numerosis, quarum præcipuus quatuordecim ad quindecim, columellá contortá, labro externo fimbriato et dentato; fulvo-cente-albá, costis rufescens-fusco maculatis, apertura fance fusca.*

THE FIMBRIATED TUN. Shell globose, very ventricose, scarcely umbilicated, sutures of the spire impressed, whorls six in number, conspicuously ribbed, ribs more or less numerous, the chief of which are about fourteen in number, columella twisted, outer lip fimbriated and toothed; fulvous white, ribs spotted with reddish-brown, interior of the aperture brown.

SOWERBY, Genera of Shells, f. 2.

Lr Minjac, Adanson.

Dolium tessellatum (*pars*), Enc. Méth.

An eadem var.?

Dolium marginatum, Philippi.

Hab. Bay of Manila, Island of Luzon, Philippines; Cuming.

M. Deshayes notices as a character by which to distinguish this from the following species, that it has about fourteen ribs, whilst the *D. maculatum* has mostly nine. I find this observation may be mainly relied on, for when the ribs are really more numerous, as in the magnificent specimen represented at Fig. 3 *a*, there are still from fourteen to fifteen principal ribs, the rest being manifestly inferior both in colour and development. The species is further distinguished by the columella being abruptly

twisted and by the outer lip becoming strongly fimbriated on arriving at maturity.

M. Adanson named this shell after the Minjac or Muntjak, a spotted animal of the deer tribe, and M. Deshayes proposes to use this for the specific title. I do not, however, see how a common name of this kind can be adopted with propriety in the Latin.

Species 4. (Mus. Cuming.)

DOLIUM MACULATUM. *Dol. testá ovalá, ventricosissimá, vir umbilicatá, spirá suturis subimpressa, anfractibus sex, transversa costatis, costis ad novem, distantibus, lirá subobscura interveniente, columellá subrectá, vir contortá, labro externo simplici; cærulescente-albá, costis rufescens-fusco maculatis, apertura fance fusca.*

THE SPOTTED TUN. Shell ovate, very ventricose, scarcely umbilicated, sutures of the spire somewhat impressed, whorls six in number, encircled by about nine distant ribs having a somewhat obscure ridge running between them, columella nearly straight, scarcely twisted, outer lip simple; bluish-white, ribs spotted with reddish-brown, interior of the aperture brown.

LANARCK, Anim. sam. vert. (Deshayes' edit.) vol. x. p. 140.

Buccinum dolium, Linnaeus.

Dolium tessellatum (*pars*), Enc. Méth.

Hab. Eastern Seas.

This is a shell of lighter growth than the preceding species, the ribs are less numerous and consequently more distant from each other, the columella is straighter, and the lip simple.

December, 1848.



DOLIUM.

PLATE IV.

Species 5. (Mus. Cuming.)

DOLIUM RINGENS. *Dol.* testá globosá, crossiusculá, spirá brevi, acutá, anfractibüs septem, transversim costatis, costis ad sedecim, obscurè superficialiter bisectis, columellá medio peculiariter excavatá, callositate magná costata supra et infra munidá, labro columellarí latè expanso, aperturá contractá, labro externo latissimè plano-reflexo et fortiter dentato; fidescente vel aurantió-athidá.

THE GRINNING TUN. Shell globose, rather thick, spire short, sharp, whorls seven in number, transversely ribbed, ribs about sixteen in number, obscurely superficially bisected, columella distinguished by a peculiar excavation in the middle, having a large ribbed callosity above and below it, columellar lip widely expanded, aperture contracted, outer lip very broadly flatly reflected and strongly toothed; light fulvous or orange-white.

Cassis ringens, Swainson in Bligh Catalogue; Sowerby in Tankerville Catalogue, App. p. xxi.

Malea latilabris, Valenciennes.

Dolium latilabre, Kiener.

Hab. Payta, Peru; Cuming.

This remarkable species and the one which follows, possess an association of character intermediate between *Dolium* and *Cassis*, which gives them a strong claim to rank as a separate genus, as proposed by M. Valenciennes under the title of *Malea*. They are of more solid growth than any other species of *Dolium*, and are singularly characterized by an excavation of the columella, which has

some obtuse wrinkles or rills above and below it. The outer lip is moreover distinguished by the mode in which it is flatly reflected and toothed.

Species 6. (Mus. Cuming.)

DOLIUM POMUM. *Dol.* testá oblongo-ovatá, turgidá, solidá, subumbilicatá, spirá brevi, anfractibüs sex, transversim costatis, costis ad duodecim, obtusis, medianis, rix elevatis, columellarí rinde corrugato-costatá, callósá, non expansá, basin versus excavatá, aperturá subcontractá, labro plano-reflexo, intus dentato; fido-spedicat, albimaculatá, apertura fince arroñatá.

THE APPLE TUN. Shell oblong-ovate, swollen, solid, slightly umbilicated, spire short, whorls six in number, transversely ribbed, ribs about twelve in number, obtuse, the middle ones scarcely elevated, columella rudely ribbed in a wrinkled manner, callous, not expanded, excavated towards the base, aperture somewhat contracted, lip flatly reflected, toothed within; light fulvous bay, sprinkled with white spots, interior of the aperture orange.

LAMARCK, Aun. sans vert. (Deshayes' edit.) vol. x. p. 142.
Buccinum pomum, Linnaeus.

Cassis labrosa, Martini.

Hab. Society and Philippine Islands; Cuming.

The excavation of the columella is much less defined than in the preceding species, and the columella also differs in not having an expanded laminar lip, its place being occupied by a callous deposit of white enamel.

December, 1848.



DOLIUM.

PLATE V.

Species 7. (Fig. a, Mus. Brit. Fig. b, Mus. Cuming.)

DOLIUM VARIEGATUM. *Dol. testā ovato-globosā, interdum maximā, umbilicatā, unfractib⁹ septem, transversum costatis, costis decim⁹ ad septendecim, subpronuntialis, interst̄itīs excavatis, superiorib⁹ lirā pareā interveniente; albida, fulvo-rufescente tincta, costis aliis rufofusco maculatis, aliis innacutatis, aperturā fave plns attus intēs aurantid.*

THE VARIEGATED TUN. Shell ovately globose, sometimes very large, umbilicated, whorls seven in number, transversely ribbed, ribs from sixteen to seventeen in number, rather prominent, interstices hollowed, upper ones with a small intervening ridge; whitish, stained with fulvous red, some ribs spotted with reddish-brown, some unsotted, interior of the aperture more or less deeply stained with orange.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 143.
Dolium Kieseri, Philippi.

Hab. North Coast of New Holland: Jukes.

There is little doubt of this species being the true *D. variegatum* of Lamarck, who notifies the locality New Holland, from whence a magnificent series may be seen in the British Museum, including specimens even larger than any to be found of the *D. gaea*. They were collected by Mr. Jukes during his recent expedition to that country, and are from Torres Straits.

The *D. Chiaueuse*, which Dr. Philippi refers to Lamarck's *D. variegatum*, is no doubt the Chinese analogue of this species: its plan of growth and style of colouring are the

same, and it differs only in being more globose and of lighter structure; still it is not Lamarck's type, and when named by Chemnitz *Buccinum Australē seu Chiaueuse*, he may have had specimens from both localities before him.

Species 8. (Mus. Cuming.)

DOLIUM COSTATUM. *Dol. testā ovato-oblongā, ventricosā, vix umbilicatā, unfractib⁹ septem, superne plano-depressis, subcaudalib⁹ transversis costatis, costis tredecim ad quatuordecim, prominentis, distantib⁹, interst̄itīs lati, concavis, levibus; aperturā oblongā, labo fimbriato et dentato; carinae-umbilicante, costis fulvo-fuscentibus, apicem versus obscure maculatis, apice nigricante.*


THE RIBBED TUN. Shell ovately oblong, ventricose, scarcely umbilicated, whorls seven in number, flatly depressed and slightly channelled round the upper part, transversely ribbed, ribs thirteen to fourteen in number, rather prominent, distant, interstices broad, concave, smooth; aperture oblong, lip fimbriated and toothed; bluish-white, ribs light fulvous brown, and obscurely spotted towards the apex, which is blackish.

DESHAYES, Anim. sans vert. vol. x. p. 144.

Dolium fasciatum, var., Kiener.

Hab. Philippine Islands; Cuming.

This bold species is amply distinguished from the *D. fasciatum*, by the prominent and remote development of the ribs, and absence of banded colouring.



g



g

DOLIUM.

PLATE VI.

Species 9. (Mus. Brit.)

DOLIUM PERDIX. *Dol. testā ovato-oblongā, tenui, inflatā, basin ceras obliquè effusā, umbilicatā, spirā subexsertā, aufractibus sex, transversim sulcatis et costatis, costis ad cingulū, contiguous, plano-concavis, vix elevatis, columellā arcuatā, labro simplici; fulvo-brunneā aut rufescente-spadiceā, maculis lunatis albī plus minusve frequentiōnis, rotatā, aperturæ fance caroescente-albā, ant brunnescute.*

THE PARTRIDGE TUN. Shell ovately oblong, thin, inflated, obliquely effused towards the base, umbilicated, spire somewhat exserted, whorls six in number, transversely grooved and ribbed, ribs about twenty in number, contiguous, flatly convex, scarcely raised, columella arched, lip simple; fulvous brown or light reddish bay, marked, more or less plentifully, with white lunate spots, interior of the aperture bluish-white or brownish.

LAMARCK. Amin. sans vert. (Deshayes' edit.) vol. x. p. 144.
Buccinum perdix, Linnaeus.

Eadem var. *Dolium plumatum*, Green.

Hab. Eastern, Western, and Pacific Oceans.

It is not often that in genera so limited in kind as the present, a species occurs with so wide a range of habitatio as the *Dolium perdix*. Inhabiting both hemispheres, it differs in the ribs being more or less elevated, but in no feature, sufficiently, to allow of its being made the subject of another species, as introduced by Dr. Jacob Green in the Transactions of the Albany Institute.

The *D. perdix*, it may be observed, is modelled on a different type from the rest of the genus, the spire being more elevated and the aperture more obliquely effused.

Species 10. (Mus. Rolfe.)

DOLIUM CHINENSE. *Dol. testā ovato-globosā, tenuissimā, ventricosā, basi subtruncatā, vix umbilicatā, aufractibus sex, transversim multicostatis, costis tenuibus depresso-sinuatis, liridē parvā intercrenante, columellā contortā, labro simplici; albido fulvoque alternatione zonatā, zonis albidiis rufo-fusco maculatis, aperture fance pallidē arrantio tintatā.*

THE CHINA TUN. Shell ovately globose, very thin, ventricose, somewhat truncated at the base, scarcely umbilicated, whorls six in number, transversely many-ribbed, ribs rather depressed, with a small ridge running between them, columella twisted, lip simple; painted alternately with fulvous and white zones, white zones spotted with red-brown, interior of the aperture faintly tinged with orange.

PESHAYES, Anim. sans vert. vol. x. p. 146.
Buccinum Australe seu Chinense, Chemnitz.
Dolium variegatum, Philippi (not of Lamarck).

Hab. China Seas.

If Dr. Philippi had proclaimed the *D. Chinense* and *variegatum* to be one and the same species in different states, instead of assigning the shell under consideration to the Lamarckian species, and distinguishing that from Torres Straits by a new name, his opinion would have been less open to objection; for the characters of the New Holland specimens are so truly represented in this, under certain modifications, that they might almost be regarded as local varieties. They agree even in colour and pattern, and the only difference between them consists in the *D. Chinense* being of much smaller and lighter structure, of more contracted growth, and more numerously and finely ribbed.



DOLIUM.

PLATE VII.

Species 11. (Mus. Cuming.)

DOLIUM FASCIATUM. *Dol. testā ovato-ventricosā, vix umbilicatā, spirā canaliculatā, aufractibūs sex ad septēm, transversis costatis, costis sedecim ad septēdecim, planō-concavis, levibus, interstītiā suprēmā lati-nsuītā, columellā subcōvolutā, labro reflexo, fimbriato-deutato; albā, fasciis quatnor latiunculis aurantiō-nigris, pone labrum evanidis, cingulatā, apice ceruleo-nigricante.*

THE BANDED TUN. Shell ovately ventricose, scarcely umbilicated, channelled at the spire, whorls six to seven in number, transversely ribbed, ribs sixteen to seventeen, flatly convex, smooth, uppermost interstice rather wide, columnella a little twisted, lip reflected, fimbriately toothed; white, encircled with four rather broad orange-brown bands, fading behind the lip, apex bluc-black.

LAMARCK, Anim. sans vert. vol. x. p. 142.

Buccinum fasciatum, Bruguière.

Hab. Philippine Islands; Cuming.

The orange-brown bands which encircle this shell constitute a marked character of the species, whilst the manner in which they gradually fade away before reaching the lip, neatly illustrates a peculiarity which may be observed throughout the species, of a certain deficiency of pattern towards the completion of the shell. Before arriving at maturity, the animal appears to lose its powers of design in the display of ornamental colouring, and it not unfrequently happens, that after any sudden accident or concussion, sufficient to cause a sea-break, the shell is completed in a coarse rugged manner, with little indication of the design and variety of colouring which has characterized its previous growth.

It is hardly necessary to draw a comparison between this species and the *D. costatum*, confounded with it by M. Kiener, for the ribs in that species are very prominent,

and widely separated, whilst in the *D. fasciatum* they are very depressed and almost contiguous; the bands, moreover, wanting in *D. costatum*, are not to be lightly regarded.

Species 12. (Fig. a, Mus. Hanley; Fig. b, Ms. Cuming.)

DOLIUM ZONATUM. *Dol. testā globoso-ovatā, tenuicollā, vix nabilicatā, spirae suturis canaliculatis, anfractibūs sex, transversis costatis, costis depresso-concavis, qua-tuordecim ad sedecim, subdistantibus, lirā tenui, superne duobus tribus, intercavente; columellā tortuosā, labro simplici aut subtili crenulato; castaneo-rufescente, concolori, anfractuā parte suprēmā albida, apice nigricante.*

THE ZONED TUN. Shell globosely ovate, rather thin, scarcely umbilicated, sutures of the spire channelled, whorls six in number, transversely ribbed, ribs depresso convex, fourteen to sixteen in number, rather distant, with a fine ridge, two or three towards the upper part, running between them; columnella twisted, lip simple or finely crenulated; chestnut-red, of one uniform colour, uppermost portion of the whorls whitish, apex blackish.

GREEN, Transactions of the Albany Institute, vol. i. part 1. p. 131.

Dolium crenulatum, Philippi.

Hab. China.

This is the only species of the genus in which there is an isolated raised line between the interstices of the lower ribs of the shell; the *D. galea* has an elevated line between the upper ribs in an early stage of growth, but in this portion of the shell of *D. zonatum*, there are three. It is accurately described and figured by Dr. Jacob Green of the United States in the volume above referred to, and a magnificent specimen has been recently described and figured by Dr. Philippi in the third volume of his 'Abbildung' as a new species with the name above recorded.



DOLIUM.

PLATE VIII.

Species 13. (Fig. 13. b, and 13. c, Mus. Cuming.)

DOLIUM CUMINGI. *Dol. testa globoso-ovata, tenui, ventricosa, umbilicata, anfractibus septem, transversioris multicotatis, costis depresso-planis, confertis, interstitiis linearis-sulcatis, columella subcontorta, labro simplici: fulvo-caeruleo, mucalis fuscis et albis fasciatus pecuniariter nebulosus.*

CUMING'S TUN. Shell globosely ovate, thin, ventricose, umbilicated, whorls seven in number, transversely many-ribbed, ribs depressedly flattened, close-set, interstices linearly grooved, columella slightly twisted, lip simple; fulvous chestnut, peculiarly clouded in a banded manner with brown and white spots.

HANLEY, MSS. Mus. Cuming.
Hab. Philippine Islands; Cuming.

This is unquestionably distinct both from the *D. olearium* and *Deshayesi*. Characterized in great measure by its globose form, it is also distinguished by its style of painting which is very peculiar, and, as in all the species of the genus, a constant and not unimportant feature.

Species 14. (Mus. Cuming.)

DOLIUM OLEARIUM. *Dol. testa orata, ventricosissima, umbilicata, spira sutaris canaliculatis, anfra tibus septem, transversioris costatis, costis plano-depressis, latiusculis, confertis, interstitiis linearis-sulcatis, labro simplici: fulvo-caeruleo, mucalis conspicuis albis aspersa.*

THE OIL-JAR TUN. Shell ovate, extremely ventricose, umbilicated, sutures of the spire channelled, whorls seven in number, transversely ribbed, ribs flatly depressed, rather broad, close set, interstices linearly grooved, lip simple; fulvous chestnut, sprinkled with conspicuous white spots.

LAMARCK, Anin. sans vert. vol. x. p. 140.

Buccinum olearium, Bruguière.
Hab. Ceylon.

This is the *Buccinum olearium* of Bruguière, but not of Linnaeus, which accounts for his remark in the Encyclopédie Méthodique: "J'ignore ce que Linnaé a entendu par les mots 'salcis obtusis liacolâ elevata interstictis,' qu'il a

employé dans la différence spécifique de ce Buccin, puisqu'il ses côtes, de même que ses sillons, n'offrent pas des lignes élevées." The shell upon which Linnaeus founded his *Buccinum olearium*, preserved in the museum of the Linnean Society of London, proves on examination to be nothing more than a very young specimen of the *D. galea*, in the early growth of which species there is always a fine elevated line in the interstices of the upper ribs.

From an observation of two young individuals of the Lamarcian *Dolium olearium* in the same collection, I am inclined to think Linnaeus' *Bulla caeruleolata*, which has never been identified, is this shell; there is, however, one point in which the description of that species in the 'Systema Naturae' does not agree, namely, in respect of form, "cylindrico." For this reason the name *olearium*, with the above explanation, may be retained.

Species 15. (Fig. 13. a, Mus. Cuming.)

DOLIUM DESHAYESII. *Dol. testa subgloboso-oblonga, ex umbilicata, spirâ uniuersâ canaliculata, anfractibus sec spiraliter linearis-sulcatis, labro simplici: pallido fulvo-elevata, mucalis conspicua albis, fuscocinnabrotis, plerunque binis, subdistinctis fasciatus picta.*

DESHAYES' TUN. Shell somewhat globosely oblong, scarcely umbilicated, spire never channelled, whorls six in number, spirally linearly grooved, lip simple pale fulvous chestnut, painted with bands of rather distant conspicuous white spots, shaded with brown and ranging mostly in pairs.

Hab. Philippine Islands; Cuming

A comparison of this shell with the Lamarcian *Dolium olearium* enables me to confirm the accuracy of M. Deshayes' observation (Anin. sans vert. nov. edit. vol. x. p. 140, note) to the effect, that the specimen figured for that species by Mr. Solley in his 'Genera of Shells,' and by myself, from the same engraving, in 'Couchologia Systematica,' is distinct from it. The sutures of the spire are not channelled, and there is a singular peculiarity in the style of colouring, being painted with white blotches, shaded with brown, ranging two and two in distant bands, or in bands longitudinally. The figure in 'The Genera' gives a characteristic view of the back of the species.

DOLIUM.


	Plate	Species		Plate	Species
<i>anapulaceum</i> , Philippi	I.	1 melanostoma, Jay		II.	2
<i>Chinense</i> (<i>Bucc.</i>), Chemnitz	VI.	10 <i>Minjac</i> , Adanson	III.	3	
<i>costatum</i> , Deshayes	V.	8 <i>olearium</i> (<i>Bucc.</i>), Bruguière	VIII.	14	
<i>crenulatum</i> , Philippi	VII.	12 <i>perdix</i> (<i>Bucc.</i>), Linnaeus	VI.	9	
<i>Cumingii</i> , Houley	VIII.	13 <i>plumatum</i> , Green	VI.	9	
<i>Deshayesii</i> , Reeve	VIII.	15 <i>pomnum</i> (<i>Bucc.</i>), Linnaeus	IV.	6	
<i>fasciatum</i> (<i>Bucc.</i>), Bruguière	VII.	11 <i>ringens</i> (<i>Cassis</i>), Swainson	IV.	5	
<i>tumbratum</i> , Sowerby	III.	3 <i>teue</i> , Menke	I.	1	
<i>galca</i> (<i>Bucc.</i>), Linnaeus	I.	1 <i>tesselatum</i> , Eie. Mdh.	III.	3	
<i>Kieueri</i> , Philippi	V.	7 <i>variegatum</i> , Lamarck	V.	7	
<i>latilabre</i> , Kienner	IV.	5 <i>variegatum</i> , Philippi	VI.	10	
<i>maculatum</i> , Lamarck	III.	4 <i>zonatum</i> , Green	VII.	12	
<i>marginatum</i> , Philippi	III.	3			

MONOGRAPH

OF THE GENUS

C A S S I S.

" Yet this obtains in all,
That all discern a beauty in His works,
And all can taste them. Minds have been formed
And tutor'd, with a relish more exact,
But none without some relish, none unmoved."—*Cowper*



CASSIS.

PLATE I.

Genus CASSIS, Lamarck.

*Testa ovata vel trigono-ovata, ad basin canali recurvo ascende-
dente traiunata, spirá brevi, aufractibus interdum varico-
sis, aufractu ultimo valde maximo, inflato; colum-
nella sepsiisimè plicato-rugatá vel granosá et plerunque
latè expausá, aperturá elongatá plus minusve angustá,
labro incassato, dentato.*

Shell ovate or triangularly ovate, terminating at the base with a recurved ascending canal, spire short, whorls sometimes varicose, last whorl very large, inflated; columella most frequently plicately wrinkled or grained, and generally widely expanded, aperture elongated, more or less narrow, lip thickened, toothed.

The *Cassides*, or Helmets, are a strong, muscular, but inactive group of mollusks, whose shell is mainly distinguished by its solid or inflated growth and short spire, and by an abruptly ascending recurved canal through which the siphon conveying the water to the breathing organs passes, and is of more than ordinary dimensions. The head is stout and fleshy, prolonged into the tentacles after the manner of *Purpura* and *Buccinum*, the eyes being situated on the outer surface, near the base; M. Deshayes has, however, observed that in the *C. sulcosa* of the Mediterranean they are elevated on a short pedicle.

The calcifying functions of the *Cassides* are somewhat vigorously exercised, some of the shells being of large size and often of extremely solid growth; the mantle which lines the interior of the shell, extends in ample folds about the aperture, and is freely occupied in the secretion of that rich display of enamel which, through being deposited in layers of different tints, affords so beautiful a material for the engraving of Cameos. The *C. Madagascariensis*, on account of its finely expanded lip and purity of colour, is particularly adapted for this singular art of gem-engraving, practised by the ancients; and the well-known 'Bull's Mouth,' *C. rugosa*, is also much used. The operculum of *Cassis* is very peculiar; it is of a horny composition of an elongated crescent form, crenulated along the edge, and rayed with deep striae.

We have no representative of this genus on our own coast, and only one species, *C. sulcosa*, ranges so far north as the Mediterranean. According to the observations of M. Deshayes, it lives upon the juices of different Bivalves, whose shells it penetrates with its proboscis, like the *Murexes*.

The chief localities noticed in the following monograph are China, the Molucca and Philippine Islands, Algoa Bay and the islands of the Pacific, the West Indies, New Holland, and New Zealand. It should be observed, however, that the species are but sparingly distributed, and very few new ones have been collected during the last few years, amongst the vast numbers that have been discovered of other genera. The shells present a lively display of colours, and, from their singular plan of growth, constitute a very characteristic type in the great series of Gastropods.

Species 1. (Fig. a, b, c, Mus. Cuming.)

CASSIS TORQUATA. *Cass. testá ovatá, spirá subacuminatá,
aufractibus varicosis, convexis, levibus, columella basi
uniplacatá, labro quinque- vel sex-spinoso; fidecescente-
laetá, aufractibus maculis quadratis intensè castanis
equidistantibus supernè et infernè unicoloria pictis.
labro castaneo maculato.*

THE ENCHAINED HELMET. Shell ovate, spire somewhat acuminated, whorls destitute of varices, convex, smooth, columella one-plaited at the base, lip five- or six-spinose; fulvous umber-colour, whorls painted round the upper and lower parts with a row of equidistant, square, dark chestnut spots, lip spotted with chestnut.

Variety β . (Fig. 1. c.)

*Testa incrassata, aufractibus supernè nodoso-tuberculatis.
Shell thickened, with the whorls nodosely tubercled round
the upper part.*

Hab. New Holland.

This species, of which four excellent figures were published in 1773 by Martini (Conch. Cab. vol. ii. pl. 38. f. 333 to 336), has been confounded hitherto with that commonly known to collectors under its two-fold aspect of *C. vibex* and *erinnaceus*. It is uniformly of smaller size and eminently distinguished by its style of painting, the pattern of which does not consist of longitudinal flames or dot-lines, but of two conspicuous chains of square chestnut spots, one encircling the whorls immediately beneath the sutures, the other at the base; the large intermediate surface being clear and unspotted, beyond three faint indications of bands.

Species 2. (Mus. Brit.)

CASSIS CORNUA. *Cass. testā maximā, ovalā, ventricosā, spirā depressā, apice acutā; anfractibus uniaricosis, s' spira angulatis, fortiter tuberculatis, tuberulus crebro-coarctatis, infra cingulis duobus callous obsoletè tuberculatis, tuberulus ectate crescentibus, cingulum intersticis scrobiculis parvis, hexagonis, una regularibus, una irreguläribus, crebrevisim reticulatis; columellā conicae-depressā, labro externo rude et fortiter rugosā, labris plus minus pleno-expansae et superne levigata; luctuosa, ferrugineo-rufescente hic illuc pallidior; cinctis, cingulisque fuso grandinaculatis, columellā et apertura fulvo-vitrinīs, rugis albocanulatis, scrobiculis sub encanosis peculiariter opaco-albis.*

THE HORNED HELMET. Shell very large, ovate, ventricose, spire depressed, sharp at the apex; whorls one-varieose, angulated round the upper part, strongly tubercled, tubercles eructly conical, two callous obsoletely tubercled belts beneath, the tubercles increasing with age, interstices between the belts closely reticulated with small hexagonal pits, sometimes regular, sometimes irregular; columella concavely depressed, and, with the outer lip, rudely and strongly wrinkled, lips more or less flatly expanded, joined at the upper part: cream-coloured, palely tinged here and there with rusty red, varices and belts stained with large brown blotches, columella and aperture fulvous citron, wrinkles whitish, pits beneath the enamel peculiarly opake-white.

(VANCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 20.
Buccinum cornutum, Linnaeus.

Cassis cornuta, Bruguière.

Eadem, *Cassis tabula*, Chemnitz.

Hab. Moluccas, West Indies, Pacific Islands.

This fine species, the largest of the genus, grows to an immense size, yet examples are not unfrequently met with, of comparatively small dimensions, bearing all the characters of the adult shell, except in having little indication of tubercles on the two central callous belts. The tubercles between these, and other parts of the surface, are distinguished in all stages of growth by a beautiful hexagon coumed structure, consisting of a number of hexa-

gonal cells, which in that part of the body-whorl overlaid by the columellar lip, become filled up with opake-white matter. The tubercles along the upper angle of the whorls are always very prominent, and sometimes rather compressed.

The *C. cornuta* assumes two very distinct varieties of form: one extremely broad and ventriose at the base of the spire, attaining rather the larger size; the other of a more oblong form, in which the lip is more widely expanded into a flattened plane. It is the latter variety of which Chemnitz has given two admirable figures in the Conchylion Cabinet, vol. ix. pl. 184 and 185, under the name *Cassis tabula*.

Species 3. (Mus. Cuming.)

CASSIS SEMIGRANOSA. *Cass. testā ovalā, basi contractā, spirā acuminatā, apice subpapillari, anfractibus varicosis, rotundatis, laceris, basi striatis, superne granularis sericulus quinque vel sex conspicuū cingulatis, granis sub plicas descendentibus; columellā teniplicatā, basi truncatā, labro reflexo, levigato; albidā, fulvo-ferrugineo plus minus tincta.*

THE HALF-GRAINED CASSIS. Shell ovate, contracted at the base, spire acuminate, somewhat papillary at the apex; whorls destitute of varices, rounded, smooth, striated at the base, circled round the upper part with five or six conspicuous rows of grains, which are upon descending folds; columella finely plicated, truncated at the base, lip reflected, smooth; whitish, more or less stained with fulvous rust colour.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 37
Hab. Van Dieman's Land.

The rows of grains which are ranged along the summit of the descending folds round the upper portion of the whorls are extremely characteristic, the columella is peculiarly short and truncated, and the smoothness of the outer lip is an unusual feature. The shell is usually stained with a kind of fulvous rust colour, whilst the apex is of ivory whiteness with a somewhat papillary aspect.



CASSIS.

PLATE II.

Species 4. (Figures, b, c, d, Mus. Cuming.)

CASSIS DECUSATA. *Cass. testá ovatá compressiuscula, spirá subacuminatá, apice acutá; anfractibus mucaricosis, striis elevatiscenis longitudinalibus, incisisque transversis nudique decussatis, praecipue ad spiram; varicibus superne peculiariter bi-tri-deutato-spinosis, columellá fusciter plicato-rugata; livido-caeruleo-roseata, maculis quadratis vel striatis fulcis longitudinaliter undata, naviatis strigata super varices et apertura fauce intensè castaneo-rufa.*

THE DECUSATED HELMET. Shell ovate, rather compressed, spire somewhat acuminate, sharp at the apex; whorls one-varicose, decussated throughout with rather elevated longitudinal and engraved transverse striae, especially on the spire; varices peculiarly two-three-dentately spinous at the upper part; columella strongly plicately wrinkled; livid blue, longitudinally waved with fulvous streaks or square spots, interior of the aperture and spots or streaks upon the varices deep chestnut-red.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 29.
Buccinum decussatum, Linnaeus.

Cassis decussata, Bruguière,
Hab. —?

In this species the striped and tessellated designs of painting which characterize the *C. areola* and *nudata* are united, the varieties being unaccompanied by any difference of form or sculpture. The chief peculiarity in the sculpture of *C. decussata*, the same in both varieties, consists in the longitudinal striae being raised, whilst the transverse striae are engraved; in other words, the former are ridges, the latter grooves. Another feature to be remarked is that the varices, including that which forms the reflected margin of the aperture, are armed at the summit, next the spire, with two or three dentate spines which in all other species, where they occur, are at the base.

M. Deshayes remarks in a note on *C. decussata* (Anim. sans vert. vol. x. p. 9.) that Lamarck in referring to

Martini's figures (Couch. Cab. vol. 2. f. 360-1, and 367-8.) has confounded a young specimen of his *C. zebra* (*C. nudata*) with it. This is a mistake: Lamarck was perfectly correct, and so was Martini, who, although he describes each of these shells separately, refers them both to the *Buccinum decussatum* of Linnaeus. They represent truly the species in different states, and perhaps the most extreme.

Species 5. (Mus. Brit.)

CASSIS MADAGASCARIENSIS. *Cass. testá sub-trigonovatá, ventricosa, elevato-rotundata, anfractibus longitudinaliter subirregulariter striatis, transversim superficialiter sulcatis, triseriatim tuberculatis, tuberculis supraeis granulatis, prominulis profusè plicato-rugatis, ringis hic illuc bifurcatis, labro latissimè expanso, marginibus conjunctis, labro externo intus conspicuè dentato; varicello-oblivate, facie ecausticā rufente-incipiatō, dentibus rugisque albis, interstitiis intense purpureo-costaneis.*

THE MADAGASCAR HELMET. Shell somewhat triangularly ovate, ventricose, elevately rotundate, whorls longitudinally somewhat irregularly striated, transversely superficially grooved, encircled with three rows of tubercles, of which the uppermost are large and prominent; columella profusely plicately wrinkled, wrinkles here and there bifurcated, lip very widely expanded, margins joined, outer lip conspicuously toothed within; bluish white, enamelled face reddish flesh-coloured, teeth and wrinkles white, interstices intense purple-chestnut.

LAMARCK, Anim. sans vert. vol. x. p. 20.
Hab. Madagascar.

This fine species may be recognised by the delicate flesh tint of its enamelled face and deep purple-chestnut colouring of the lip and columella; it attains almost as large a size as the *C. corallinus*. In sculpture it differs from that species in having the surface transversely rather superficially grooved, not cancellated.



CASSIS.

PLATE III.

Species 6. (Mus. Cuming.)

CASSIS BISULCATA. *Cass. testa globosa, tenuicula, spirá brevi, anfractibus evaricosis, rotundatis, transversis impresso-striatis, sulcis ducibus subobscureis infra suturas; columellá fortissimè rude rugoso-plicatá, labro dentato-lirato; céruleo-albicans, muculis fideo-rufescens; tuberculis parvis quadratis distansibus ornata, apice lido-nigricante.*

THE DOUBLE-GROOVED HELMET. Shell globose, rather thin, spire short, whorls destitute of varices, rounded, transversely impressively striated, with two somewhat obscure grooves beneath the sutures; columella very strongly roughly plaited, lip dentately ridged; bluish white, ornamented with small fulvous red distant square spots, apex pale livid black.

SCHUBERT and WAGNER, Conch. Cab. Supp. p. 68, pl. 223. f. 3051, 3052.

Hab. Manila, Island of Luzon (on the reefs); Cuming.

The general aspect and sculpture of this shell approaches nearest to that of *C. canaliculata*, from which it differs in not having the deep broad channelled groove around the sutures, whilst the coloured squares are more distant, darker, and more defined.

Species 7. (Mus. Brit.)

CASSIS TUBEROSA. *Cass. testa magiá, trigono-ovata, gibbosissimá, spirá plano-convexa, apice acutá; anfractibus unicolorosis, superè angustatis, tuberculatis, tubero centro ventrali dorsali prominentiore, infra cingulos tribus callosis, obsoletè tuberculatis, interstítios transversis sulcatis, longitudináliter crebre rufescens; columellá planiuscula, leviter concava, rudi rugata, rugis nonnullis seleni divisis, labro externo fortiter dentato; aquilo-alba, dorso rufo-fuscis ocreo-locque nebulatá, castaneo conspicè longitudinaliter flammantá, varicibus maculis perporatis nigricantibus notatis, columellá vixidè castanæ, rugis albidis.*

THE TUBEROUS HELMET. Shell large, triangularly ovate, very hump-backed, spire flatly convex, sharp at the apex; whorls one-varicose, angulated round the

upper part, tubercled, ventral dorsal tubercle more prominent than the rest, with three obsoletely tubercled callous belts, interstices transversely grooved and longitudinally very closely lineated in a somewhat wrinkled manner; columella rather flattened, slightly concave, rudely wrinkled, some of the wrinkles divided by a groove, outer lip strongly dentated; swarthy-white, clouded on the back with reddish brown or blue, varices marked with a very few blackish blotches, columella bright chestnut, wrinkles whitish.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 21.
Cassidea tuberosa, Bruguière.

Testa junior. Boeckiana striatula. Gronovius.
Hab. Philippine Islands; Cuming.

This species is mainly distinguished by its swarthy hue and solid triangular growth, the back of the shell being peculiarly gibbous and strongly tubercled.

Species 8. (Mus. Metcalfe.)


CASSIS CANALICULATA. *Cass. testa globosa-ovata, tenuicula, subangustata, spirá brevissimè, suturis peculiariter late et profunde evanescientib; evaricosis, transversis inciso-striatis; columellá rude plicato-lirata, labro dentato; lido-albicans, muculis oblongo-quadratis inaequalibus rufescente-fuscis subindistinctè quadrifasciatis.*

THE CHANNELLED HELMET. Shell globosely ovate, rather thin, somewhat inflated, spire rather short, sutures peculiarly broadly and deeply channelled; whorls destitute of varices, encircled with engraved striae; columella rudely plicately ridged, lip dentated; livid white, somewhat indistinctly four-banded with unequal oblong-square reddish brown spots.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 37.
Cassidea canaliculata, Bruguière.

Hab. Philippine Islands; Cuming.

Distinguished from all other species by the curious deep broad channel around the sutures.



C A S S I S.

PLATE IV.

Species 9. (Mus. Cuming.)

CASSIS SPINOSA. *Cass. testā oblongo-ovatā, leviculā, reutricosā, basi alteauatā, spirā depresso-concavā, apice acutā, aufractibus unicoloricos, transversim obsoletē sulcatis, spirā basique longitudinaliter tenuiplicatis, superē nodularum series tribus, quarum superiorē minuta moniliformi, cingulata; columellā concavā, plicato-rugatā, rugis brevibus irregularibus, labro exterō rudi dentato, serie nodularum; lactea, fulvo-spadicis pallio tinctā, fasciis quinque ant ploribus, maculis semilunaris satartioribus abso articulatis, cingulata.*

THE SPINOUS CASSIS. Shell oblong-ovate, rather thin, ventricose, attenuated at the base, spire very short, sharp at the apex, whorls one-varicose, transversely obsoletely grooved, longitudinally finely plicated on the spire and at the base, encircled with three rows of nodules round the upper part of which the uppermost row is minute and necklace-like; columella concave, plicately wrinkled, wrinkles short and irregular, outer lip rudely dentated, with the addition of a row of nodules; cream-coloured, palely stained with fulvous bay, and encircled by four or more bands of semilunar darker bay spots articulated with white.

DESHAYES, Anim. sans vert. vol. x. p. 23, note.

Buccinum spinosum, Gronovius, Zooph. p. 302, pl. 19. f. 9.

Buccinum Rupicollis, Gmelin,

Buccinum tessellatum, Gmelin.

Buccinum maculosum, Gmelin.

Cassidea fasciata, Bruguière.

Cassis fasciata, Lamarck.

Hab. West Indies.

There are three peculiarities of character in this species which distinguish it in an eminent degree from any other. The first consists of a necklace-like row of small spinous nodules just within the base of the spire; the second feature of interest is the dorsal bands of articulated semilunar spots, which are presented in all stages of growth; and the third is that of a row of spinous nodules upon the flattened surface of the outer lip. In form the *C. spinosa* has some general resemblance with the *C. flavanea*, but it is a shell of much lighter growth and has no indication of any tubercles.

Species 10. (Mus. Cuming.)

CASSIS TESTICULUS. *Cass. testā suboblongo-ovatā, crassā, ponderosā, Cypræformis, spirā brevissimā, apice acutā, aufractibus unicoloricos, longitudinaliter tenuē et creberū plicato-liratis, superē rotundatis, basia versus sulcatis; columellā cattosa, undique plicato-rugatā, labro crasso reflexo, inter dentato, aperturā elongatā, angustā; lividio-violaceata, purpareo-fusco longitudinaliter undulatā et muculatā, liris dorsalis albida, columellarū labroque rufescensibus, posteriore poenā nigromaculata.*

THE PURSE HELMET. Shell somewhat oblong-ovate, thick, ponderous. Cowry-shaped, spire very short, sharp at the apex; whorls destitute of varices, longitudinally finely and very closely plicately ridged, rounded at the upper part, grooved towards the base; columella callous, plicately wrinkled throughout, lip thickly reflected, toothed within, aperture elongated and narrow; livid violet, longitudinally waved and spotted with purple brown, dorsal ridges whitish, columella and lip reddish, the latter spotted behind with black.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 32.

Buccinum testiculus, Linnaeus.

Cassidea testiculus, Bruguière.

Hab. West Indies.

Variety β . (Fig. 10. a.)

Testa subpyriformis, superē crassissima, turgida, plicato-tuberculata.

Shell somewhat pear-shaped, very thick, swollen, and plicately tubercled round the upper part.

Cassis plicata, Martini.

Buccinum plicatum, Dillwyn.

Cassidea crassa, Bruguière.

Cassis crassa, Lamarck.

Hab. West Indies.

M. Kiener has very truly indicated these to be varieties of the same species, the latter being of a very thick swollen callous growth, plicately tubercled round the upper part, but agreeing in the minutest detail with the former. Both are from the same locality.



CASSIS.

PLATE V.

Species 11. (Mus. Cuming.)

Cassis Saburon. *Cass. testá ovalá, ventricosá, spirá breviusculá, apice subpapillari; anfractibus evanescis rotundatis, transversas undique creberrimè sulcatis, marginè superiori crenulatis; columella callosa, rugis brevibus perplicatis obliquis, labris subirregulariter dentato-irato; pallidè ciucrea ant fulvescente, obscurè quinquefasciatá, fasciis indistinctè quadro-maculatis, maculis ponit varices intense rufo-caustaeis.*

THE SABURON HELMET. Shell ovate, ventricose, spire rather short, apex slightly papillary; whorls destitute of varices, rounded, transversely very closely grooved throughout, crenulated at the upper margin; columella callous, with a very few short oblique wrinkles, lip somewhat irregularly dentately ridged; pale ash or light fulvous colour, obscurely five-banded, bands indistinctly square-spotted, spots behind the varices dark reddish chestnut.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 36.

Le Saburon, Adanson.

Cussiaea Saburon, Bruguière.

Buccinum Saburon, Dillwyn.

Eadem. *Cassis posanae,* Schubert and Wagner.

Hab. Japan (on the sands); Dr. Siebold.

Mainly distinguished by its close transverse grooves and rounded form; the spots are often much obscured, excepting behind the marginal varix, where in comparison they are unusually dark.

Martini has a very excellent figure of this species (Conch. Cab. vol. 2. pl. 34. f. 350) referred to by Lamarck under the head of *C. granulosa* (*C. inflata*), and the figure in a vignette, f. 1-2 at p. 10, of the same work to which M. Deshayes refers for *C. Saburon*, appears to correspond with one not hitherto described, which I propose to distinguish by the name *C. pila*.

Species 12. (Mus. Cuming.)

Cassis flamma. *Cass. testá oblongo-ovalá, vis. levigatá, spirá breviusculá, apice acutá; anfractibus evanescis longitudinaliter teninder plicato-rugatis, rugis quadrum versus etate obsolete, anfractuum parte superiori ex angulis, tuberculorum seribus trilobis aut pluribus armatis, tuberculis lateraliiter compressis, labrum versus obsoletioribus; columella concavusculá, ruditè rugata, labro externo fortiter dentato; laetá, dorso griseo-carmesinante nebulosá, castaneo longitudinaliter flammatá, varicibus maculis granulibus nigris tinctis.*

THE FLAMED CASSIS. Shell oblong-ovate, scarcely triangular, spire rather short, sharp at the apex; whorls one-variegate, longitudinally finely plicately wrinkled, the wrinkles becoming obsolete towards the middle with age, upper part of the whorls but slightly angulated, armed with three or more rows of tubercles, which are laterally compressed and more obsolete towards the lip; columella rather concave, rudely wrinkled, outer lip strongly toothed; cream-coloured, clouded at the back with greyish blue, longitudinally flamed with chestnut, varices stained with large black blotches.


LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 22.

Buccinum flammatum, Linneus.

Cassidea flammata, Bruguière.

Hab. —?

The peculiar oblong oval form of this species, which is of constant occurrence, fully distinguishes it from *C. tuberosa*, of which M. Kiener intimates his opinion that it is a variety. There is moreover a uniform variation in the sculpture, which, in the species under consideration, rather assumes the nature of plicated wrinkles; the tubercles are more numerous, smaller, and more compressed



CASSIS.

PLATE VI.

Species 13. (Fig. *a*, Mus. Saul; Fig. *b* and *c*, Mus. Cuming.)

CASSIS TENUIS. *Cass. testá ovato-oblongá, cylindracea,* *Cypraeiformis, teniculá, spirá brevissimá, anfractibus* *locibus, transversum regulariter sulcatis, sulcis binis,* *interstitti longitudinaliter obtuse liratis, tuberculorum* *series quatuor arcuatis; columellá eis concavá, su-* *perré et inferná rugatá, labro exterio peculiariter* *reflexo, fortiter dentato-lirato, liris binis; apertura* *elongatá, extremitatibus ambabus conicalitatis; pur-* *pureo-fuscá, livido-ceruleo nebulatá, sulcis lirisque dor-* *salibus ceruleo-albidis, columellá purpureo-nigrá, rugis* *albis, tubo columellarí basin versus raso vixide tinctá,* *arcá pone labrum exterum purpureo-nigro conspicuá* *maculatá.*

THE THIN CASSIS. Shell ovately-oblong, cylindrical, Cowry-shaped, rather thin, spire very short, whorls smooth, transversely regularly grooved, grooves two and two, interstices longitudinally obtusely ridged, armed with four rows of tubercles; columella slightly concave, wrinkled above and below, outer lip peculiarly reflected, strongly dentately ridged, ridges in pairs, aperture elongated, both extremities channelled; purple-brown, clouded with livid blue, dorsal grooves and ridges white, columella stained towards the base with bright red, area behind the outer lip conspicuously blotched with purple-black.

GRAY, Wood Index Test. Supp. pl. 4. f. 4.

Cassis Masseæ, Kiener.

Hab. Galapagos Islands (in sandy mud at the depth of six fathoms); Cuming.

M. Deshayes in his notice of this species likens it to the *C. testiculus*; the accompanying figure of a mature specimen shows, however, that it is much more nearly allied to the *C. rufa*. In its most advanced stage of growth the shell is comparatively thin, the lip being curled back without any great deposit of enamel; the general form is more

cylindrically oblong than that of *C. rufa*, and the aperture is wider. The colour and marking are extremely pretty.

The *C. tenuis*, of which a very small but accurate figure was published by Mr. Wood in his supplement to the "Index Testaceologicus," has been very generally mistaken by English collectors for Lamarek's *C. penuata*, which is merely the young of *C. rufa*.

Species 14. (Mus. Saul.)

CASSIS COARCTATA. *Cass. testá cylindraceo-ovatá, crassi-* *sculpá, spirá brevi, anfractibus ericicis, transversum* *tevissimè striatá, costis nodulosis cingulatá, quarum* *quatuor exteris majoribus; columellá brevissimè et* *obliquè subtilè rugatá, labro intus dentato-lirato,* *superne corrugato, non reflexo, aperturá elongatá, angustá;* *cirreto-albida, purpureo-fusco tinctá, nodulis* *nonnullis albicantibus, columellá labroque refuscensibus.*

THE CONTRACTED HELMET. Shell cylindrically ovate, rather thick, spire short, whorls destitute of varices, transversely very finely striated, encircled by nodulous ribs, four of which are larger than the rest; columella shortly plicated, and obliquely finely wrinkled, lip dentately ridged within, contracted at the upper part, not reflected, aperture elongated and narrow; ash-white, stained with purple-brown, some of the nodules whitish, lip and columella reddish.

GRAY, Wood Index Test. Supp. pl. 4. f. 5.

Cypraeacassis coarctata, Stutchbury.

Hab. Galapagos Islands, Pacific Ocean (in crevices of rocks); Cuming.

The contraction of the outer lip, which is always constant, is a singular feature in this species. For the magnificent specimen here represented, which is of unusually large size, I am indebted to the choice collection of Miss Saul.



CASSIS.

PLATE VII.

Species 15. (Mus. Cuming.)

CASSIS VIBEX. *Cass. testā elongato-ovatā, basi peculiariiter effusā, spiri' subacuminotā; anfractibus evanescis, convexis, subobliquā convolutis, levigatis, politis, columellā incrassatā, basi uniplicatā, labro basi quadrispinoso; cornu-euercante, obscurè fasciatā, flammis fulvescentibus longitudinaliter pīctā, lineisque indistincte fuscō-punctatis cingulatā; columellā eburinā, labro castaneo intēnsū maculato.*

THE STRIPED HELMET. Shell elongately ovate, peculiarly effused at the base, spire somewhat acuminate; whorls destitute of varices, convex, somewhat obliquely convoluted, smooth, polished, columella thickened, one-plaited at the base, lip four-spined at the base; fleshy ash-colour, obscurely banded, painted longitudinally with light fulvous flames and encircled with indistinctly brown-dotted lines; columella ivory-white, lip intensely spotted with chestnut.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 38.

Buccinum riberi, Linneus.

Cassidea riberi, Bruguière.

Variety β . (Fig. 15. a.)

Testa minor, valde incrassata, anfractibus superne nodosis, nodis in plicis sep̄e descendantibus.

Shell smaller, very much thickened, with the whorls noduled round the upper part, nodules frequently descending in folds.

Buccinum erinaceus, Linnaeus.

Buccinum nodulosum, Gmelin.

Buccinum hirsutum, Dillwyn.

Cassidea erinaceus, Bruguière.

Cassis erinaceus, Lamarck.

Hab. Philippine and Pacific Islands; Cuming.

The species *C. vibex* and *torquata* were confounded indiscriminately by Linneus, Lamarck, and others, to judge by the reference made to Martin's figures, the smooth specimens being set apart under the specific title of *riberi*, the noduled variety under that of *erinaceus*: it will, however, be seen that there are two very distinct species, both equally distinguished in having the upper portion of the whorls sometimes rounded, sometimes strongly noduled or plicately tubercled. The name *erinaceus* is therefore suppressed, as being founded on the

thickened noduled variety of either species, and that which had not been recognised is named *C. torquata*.

Species 16. (Mus. Cuming.)

CASSIS RECURVIROSTRUM. *Cass. testā ovatā, tenuiculā, spirā subacuminatā, anfractibus evanescis, superne concavo-declivibus, pleis tenibus interdum obsolete s et suturis descendantibus; columellā subampullā uniplicatā et obtusā granosā, labro subtiliter dentatolitato; pallidē fulvo-spadiceā, maculis parvis rufescente-fuscis subquadratris distantibus seriatim rotundatā.*

THE RECURVED-BEAK HELMET. Shell ovate, rather thin, spire somewhat acuminate, whorls destitute of varices, smooth, concavely slanting round the upper part, with thin plait, sometimes obsolete, descending from the sutures; columella somewhat largely one-plaited and obtusely grained, lip finely dentately ridged; light fulvous bay, marked with rows of small rather square distant reddish brown spots.

Buccinum recurvirostrum, Wood, Index Test. pl. 22. f. 30. Hab. Raines' Island, Torres Straits; Capt. Ince.

M. Kiener figures this shell as the young of *C. inflata* (*C. granulosa*, Lamarck), from which, it is scarcely necessary to add, it is perfectly distinct. The shell here represented, from Mr. Cuming's collection, is one of mature growth; and I take it to be the *Buccinum recurvirostrum* of Wood, founded on Lister's figure, Hist. Conch. t. 1016. f. 75. It may be observed that this species is from New Holland, whilst the *C. inflata* is a native of the West Indies.

Species 17. (Mus. Cuming.)

CASSIS FIMBRITA. *Cass. testā oblongo-ovatā, plerunque tenaculā, spirā subprodnetā, apice papillari, anfractibus evanescis, longitudinaliter irreguliter levigatis, striis transversis prope apicem basiique decussatis, parte superiori dorsali cingulis nodosis tribus armatā, nodis posticè compressis; columellā arcuatā, leviter rugulā, labro externo eis obsoletè dentato; pallidē luteā, fasciis sc̄r. fulvescentibus, lineisque fuscis interrupitis, octo aut pluribus cingulatā, variebus fuscō-nodulatis.*

THE FIMBRIATED HELMET. Shell oblong-ovate, generally rather thin, spire somewhat produced, papillary at the apex, whorls one-varicose, longitudinally irregularly finely plicated, crossed near the base and apex with transverse striae; upper dorsal part of the shell armed with three rows of posteriorly compressed nodules; columella curved, but slightly wrinkled, outer lip scarcely obsolete dentated; pale cream-colour, encircled by six light fulvous bands and eight or more interrupted brown lines, varices stained with large brown blotches.

QUOY and GAIMARD, Voy. de l'Astrolabe, Zool. vol. II.
p. 596, pl. 43, f. 7, 8.
Hab. New Holland.

The form of this species approaches nearest to that of the *C. fasciata* and *flammeata*; its principal features of distinction consist in the lip and columella being only very slightly wrinkled and dentated, and in the shell being encircled with brown interrupted lines.

The name *C. fimbriata* was applied in general terms by Martini in reference to the *C. villex* and *torquata*, but not upon the binomial principle of nomenclature introduced by Linnaeus.



CASSIS.

PLATE VIII.

Species 18. (Mus. Hamilton.)

CASSIS ABBREVIATA. *Cass. testá subgloboso-oratá, crissá, spirá breviscutá, acuminatá, apice subpapillari; anfractibus, ultimo interdum varicosis, subrotundis, striis longitudinalibus elevatis, sulcis striis incisis transversis decussatis, liris apicea versus conspicuè granosis; columellá fortiter granosá, labro calde incrassato, propria basi leviter effuso, utes dentato-lirato; eburnea, maculis parvis aurantio-fuscis quadratis vel inaequalibus distantibus plus minus obscurè pictâ, interdum innuculata, labri marginé retrorsò fuso radiato*

THE ABBREVIATED HELMET. Shell somewhat globosely ovate, thick, spire rather short, acuminate, apex rather papillary; whorls, the last being sometimes one-varicose, somewhat rounded, decussated with longitudinal raised striae and transverse engraved striae or grooves, ridges conspicuously granose towards the apex; columella strongly granose, lip excessively thickened, slightly effused towards the base, dentately ridged within; ivory white, more or less obscurely painted with small orange-brown spots which are square or unequal and distant, sometimes unspotted, reflected edge of the lip rayed with brown.

LAMARCK, Anim. sans vert. (Peshayes' edit.) vol. x. p. 30.

Eadem. *Cassis lactea*, Kiener.

Hab. Acapudo.

When Lamarck introduced this species, it escaped his notice, as subsequently that of his commentators, that he had already referred to a characteristic figure of it in Martini (Conch. Cab. vol. ii. pl 34. f. 351, 2) among the synonymy of his *C. granulosa* (*C. inflata*). It presents itself under so many different aspects, that it is somewhat difficult to recognise the species in all its varieties without a comparison of numerous specimens. M. Kiener's *C. lactea* is undoubtedly a variety of it, and the same may be said, with less certainty in the absence of figures, of the *C. cestigondrata* and *dolata* described in 1811 by Valentiniennes in the Zoology of the Voyage of Humboldt and Bonpland.

Species 19. (Mus. Metcalfe.)

CASSIS PAUCIRUGIS. *Cass. testá oratá, spirá subacuminatá, apice acutá; anfractibus varicosis, lacibus, nitidis, longitudinaliter irregulariter tenuissimè inciso-striatis,*

apicem versus decussatis, superne angulato-declivibus, ad angulum obsoletè nodosis; columellá vix plicatá, labro utes dentato-lirato; eburnea, polita, maculis quadratis fulcescentibus distantibus exilissimè tincta, maculis saturatioribus pone labrum; apertura fauce fulvescente-crocata.

THE FEW-WRINKLED HELMET. Shell ovate, spire somewhat acuminate, sharp at the apex; whorls destitute of varices, smooth, shining, longitudinally irregularly very finely marked with engraved striae, decussated towards the apex, angularly slanting round the upper part, obsoletely nodose at the angle; columella scarcely plicated, tip dentately ridged within; ivory white, polished, very faintly stained with distant square fulvous spots, which are darker behind the lip; interior of the aperture fulvous saffron colour.

MENKE, Moll. Nov. Hollandiae, p. 23. no. 107.

Hab. West coast of New Holland; Menke.

The *Cassis paucirugis* was described about five years ago by Dr. Menke of Pyrmont, Germany, in his Synopsis of Shells from New Holland, and kindly made known to me by the transmission of his original specimen. From its very discoloured and bleached appearance the species was regarded with some doubt, but a comparison of further specimens from the collections of Mr. Cuming, Mr. Metcalfe, and Mr. Hanley enables me to add satisfactorily to the characters already adduced by Dr. Menke. In the specimen here presented there is a perceptible indication of square fulvous spots beneath its ivory enamelled surface, the columellar lip remaining of as clear a porcelain white as that of the *Oenida ovaum*, and the interior is distinguished by a warm, fulvous saffron hue.

The *C. paucirugis* has not hitherto been illustrated, and is but little known. It is impossible to identify a species with any degree of certainty without the aid of a figure, and this, like many more whose existence has only been recorded by the simple technicalities of a diagnose, has been waiting in obscurity to be published under a more intelligible aspect.

Species 20. (Mus. Brit.)

CASSIS RUTA. *Cass. testá oratá, Cypraeiformis, crassissimá, ponderosá, spirá brevissimá, anfractibus varic-*

Cassis. *longitudinaliter subtiliter et crenatissimè plicato-corrugatis, tubercularum series tribus aut pluribus cingulatis; columnellā concaviusculā, undique tenuē rugatā, lobis valde incrassatis externo fortiter rugato, aperturā elongatā et angustā, extremitatibus amboibus profundè canaliculatis; livido-rufescente, fuso alboque maculata, aperturā rufide coccineo-rufā, hic illuc nigri-cante tructā, rugis alludis.*

THE RED HELMET. Shell ovate, Cowry-shaped, very thick, ponderous, spire very short, whorls destitute of varices, longitudinally finely and very closely plicately wrinkled, encircled with three or more rows of tubercles; columnella slightly concave, finely wrinkled throughout, lips extremely thickened, the outer lip strongly wrinkled, aperture elongated and narrow, with both the extremities deeply channelled; livid red, spotted with brown and white, aperture deep scarlet-red, tinged in parts with black, wrinkles whitish.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 30.

Buccinum rufum, Linnaeus.

Cassidea rufa, Bruguière.

Testa juvenis.

Buccinum pullum, Born.

Buccinum pennatum, Gmelin.

Cassidea pennata, Bruguière.

Cassis pennata, Lamarck.

Hab. Moluccas, &c. Isle of Annaa, Pacific Ocean (on the reefs); Cuming.

The *Cassis rufa* appears to be the species most in request for the manufacture of ordinary cameos, owing to the great thickness of the enamel, which is frequently deposited in a dozen to twenty layers. Large quantities of these shells are imported for the purpose, and the species has become a common article of merchandise.

The *Cassis pennata* of Lamarck, for the illustration of which the author refers to Martini's Couch. Cab. vol. ii. pl. 36. f. 372-3, is undoubtedly the young of *C. rufa*.



CASSIS.

PLATE IX.

Species 21. (Mus. Cuming.)

CASSIS PILA. *Cass. testā subgloboso-ovatā, spirā brevisculā, apice acutā; anfractibus evanicos, transversim undique crebrimē inciso-striatis; columellā densē plicato-liratā, labro dentato-brato; fulvo-albicante, maculis numerosis parvis quadratis rufo-fuscis seriatim pictā.*

THE BALL HELMET. Shell somewhat globosely ovate, spire rather short, sharp at the apex; whorls destitute of varices, very closely encircled throughout with engraved striae; columella thickly plicately ridged, lip dentately ridged; fulvous white, painted with rows of numerous small square red-brown spots.

Hab. China.

A shell of rather light growth, globose in form, painted with neatly defined distant spots.

Species 22. (Fig. a and b. Mus. Hanley. Fig. c. Mus. Cuming.)

CASSIS INFELA. *Cass. testā subgloboso-ovatā, tenuiculā, centricissimā, spirā acutā; anfractibus rotundatis, inflatis, levigatis, transversim obsoletē sulcatis, epicenos versus reticulatis striatis; columellā leviter plicatā et conspicue granosā, labro reflexo, intus acutē brevibrato; cōrdelescente-albidā, fasciis griseis macularum rufofuscescentium cingulatā, maculis subquadratis, irregulare interrupitis.*

THE INFLATED HELMET. Shell somewhat globosely ovate, rather thin, extremely ventricose, spire sharp; whorls rounded, inflated, smooth, transversely obsoletely grooved, reticulately striated towards the apex; columella smoothly plicated and conspicuously granose, lip reflected, sharply shortly ridged within; bluish white, encircled by five bands of somewhat square and irregularly interrupted reddish-brown spots.

Variety β (Fig. a and b. Mus. Hanley.)

Testa multo raior, anfractibus superne crebrè nodosis, non nullis unicoloriosis.

Shell much smaller, with the whorls closely noduled round the upper part, some of them distinguished by a varix. *Buccinum inflatum*, Shaw, Nat. Misc. v. 22. pl. 959.

Buccinum tessellatum, Chemnit.

Cassidea granulosa, Bruguière.

Cassis granulosa, Lamarck.

Hab. Barbados.

The first authentic illustration of this species occurs in Shaw's Naturalist's Miscellany, 1789, with the name *Buccinum inflatum*. Ten years subsequently an equally characteristic figure of it was published by Chemnit in his last volume of the Conchylien Cabinet under the title *Buccinum cassidiforme tessellatum*. These, however, were overlooked by Bruguière and Lamarck, who introduced the species with the name *granulosa*, and referred for illustration to Martini's figures, Conch. Cab. v. 2. f. 344—5 and 350—52, which also comprehend the *C. abbreviata* and *sabroui*. Kiener, following in the steps of his predecessors in adopting the name *granulosa*, added much to the confusion by representing the *C. recurvirostrum* to be the young of it, and that both were merely varieties of the *C. subcosta* of the Mediterranean, arising out of differences of age or sex. It does not appear that this crochet of M. Kiener was noticed in the new edition of Lamarck, and M. Deshayes errs in stating that the figures above alluded to in Martini, "can only be referred to this species." The shell Fig. 344—5 is without doubt a *C. inflata*, though presenting the unusual feature of a strong lateral varix, but fig. 350, is a variety of *C. sabroui*, distinguished by the dark spots behind the reflected lip, and the shell, Fig. 351—2 is a pale variety of Lamarck's *C. abbreviata*.

The coronated variety of *C. inflatum* is very uniform and I am not aware that it exceeds the size of the specimen here figured.

Species 23. (Mus. Cuming.)

CASSIS JAPONICA. *Cass. testā ovatā, tenuiculā, ventricosa, inflatā, spirā brevissimā, acutā; anfractibus suprae-rotundatis, transversim acutē sulcatis, sulcis supremis ceteris latioribus, striis subtilibus elevatis sulculis longitudinalibus obliquis decussatis; columellā plicato-rugatā, rugis partim obsoletis, interrupitis et obscurè granulatis; labro dentato-brato; albidā, maculis oblongo-quadratis inaequalibus castaneo-rufis quaque fasciatis.*

THE JAPAN HELMET. Shell ovate, rather thin, ventricose, inflated, spire rather short, sharp; whorls rounded at the upper part, transversely sharply grooved, uppermost grooves broader than the rest, decussated with very fine slightly elevated longitudinal oblique striae; columella plicately wrinkled, wrinkles partially obsolete, interrupted and obscurely granulated; lip dentately toothed; whitish, encircled

with five bands of unequal oblong-square chesnut-red spots.

Hab. China and Japan (on the sands); Siebold.

A beautiful species strongly characterized by its close grooving and deeply excavated trench behind the varices; it is moreover distinguished by its light purple tinge of colouring and by the oblique disposition of the spots.

Species 24. (Mus. Cuming.)

CASSIS AREOLA. *Cass.*, testā ovatā, spira acuminatā, apice acutā; anfractibus univaricosis, supernè concaviusculis, iugrā gibbosis, levibus, nitidis, apicem versus decussatis; columellā plicatā et interrupte rugatā, labro acutē dentato-lirato, medio transversulo; ceruleo-albicans, marginis grandibus quadratis fulvescente-fuscis quinquefasciatā.

THE DRAUGHT-BOARD HELMET. Shell ovate, spire acu-

minated, sharp at the apex; whorls one-varicose, rather concave round the upper part, gibbous beneath, smooth, shining, decussated towards the apex; columnella painted and interruptedly wrinkled, lip sharply dentately ridged, rather swollen in the middle: bluish white, encircled by five bands of large square fulvous-brown spots.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. pl. 27.

Buccinum areola, Linnaeus.

Cassidea areola, Bruguière.

Bezoardica areola, Schumacher.

Hab. Island of Negros, Philippines (on mud banks). Cuming.

The tessellated painting of this well-known species is so peculiar, that little need be added to the characters above detailed. In form it is mainly distinguished by its elevated gibbous growth.



CASSIS.

PLATE X.

Species 25. (Fig. a. Mus. Cuming.
Fig. b. Mus. Saul.)

CASSIS TURGIDA. *Cass. testā ovatā, tenuicūlā, ventricosā, spirā subacuminatā, anfractibus univaricosis, lăvibus, turridis, supernē rotundatis; columellā valde confortatā, basi uniplicatā, labro undique dentato-spinoso; lactea, pallidē castaneo longitudinaliter flammatā, labro intēse castaneo maculatō.*

THE SWOLLEN HELMET. Shell ovate, rather thin, ventricose, spire somewhat acuminate, whorls destitute of varices, smooth, swollen, rounded at the upper part; columella much twisted, one-plaited at the base, lip dentately spinous throughout; cream-coloured, longitudinally flamed with pale chesnut, lip spotted with very dark chesnut.

Hab. Boljoon, Island of Zebu, Philippines (on sand banks); Cuming.

A fine species belonging to the same group as *C. aecho-tina*, *vibex*, and *torquata*; the whorls are distinguished for their swollen rotundity, and the columella is remarkably twisted. The lip is dentately spinous throughout. I have not seen any noduled variety of this species.

Species 26. (Mus. Cuming.)

CASSIS UNDATA. *Cass. testā ocatā, gibbosissimolā, spirā acuminatā, apice ventā; anfractibus univaricosis, lăvibus, basi conspicuē saepe, superne depresso-sculptis et striatis, apice versus granulosō-decussatis; columellā plicatā et interrupē rugatā, medio tuberculatā, labro fortissimō dentato-lirato, liris elongatis; veruculo-albicante, fulvescente-fusco longitudinaliter striatā.*

THE WAVED HELMET. Shell ovate, rather gibbose, spire acuminate, sharp at the apex; whorls one-varicose, smooth, conspicuously grooved at the base, a little depressed and striated round the upper part, granulously decussated towards the apex; columella plicated and interruptedly wrinkled, tuberculated in the middle, lip very strongly dentately ridged, ridges elongated; bluish-white, longitudinally streaked with fulvous brown.

DESHAYES, Anim. sans vert. vol. x. p. 28, note.

Cassis levis undata, Martini.

Buccinum striatum, Gmelin.

Cassidea areola, var., Bruguière.

Cassis zebra, Lamarck.

Hab. Philippine Islands; Cuming.

Although the patterns of this species and the *C. areola* are united in the *C. decussata*, they are not varieties of the same as assumed by Bruguière and Kiener. The *C. areola* is a narrower and more gibbose shell than the *C. undata*, which is eminently distinguished from the former by the strong linear grooves around the base. The dentate ridges on the outer lip are longer and more strongly developed.

Species 27. (Mus. Cuming.)

CASSIS PLICATA. *Cass. testā subelongato-ovatā, peculiariter compressā, spirā subacuminatā, apice acutā; anfractibus univaricosis, transversis subtilissimis linearē-sulcatis, supernē concavo-declinibus, margine nodulosō-crenatis, infra subtiliter bi-seriatim gemmatis, ad angulum conspicuē muricato-nodosis, nodis in pliis prononiis subarcuatis descendentibus; columellā latē-expansā, plicato-rugolatā, rugis brevibus, labro acutē dentato-liratis, basi obsoletē trispinoso; livido-albante, fulvescente longitudinaliter undulatā strigatā, variculis fusco-maculatis, columellā labroque albīs, aperture fruce fulvo-castanēa.*

THE PLICATED HELMET. Shell somewhat elongately ovate, peculiarly compressed, spire somewhat acuminate, sharp at the apex; whorls one-varicose, transversely finely linearly grooved, concavely slanting at the upper part, nodulously crenated at the margin, encircled by two finely beaded ridges beneath, conspicuously prickly noduled at the angle, nodules descending in prominent slightly curved plait; columella widely expanded, plicately wrinkled, wrinkles short, lip sharply dentately ridged, obsoletely three-spinose at the base; livid-white, longitudinally unduly streaked with pale fulvous-colour, varices spotted with brown, lip and columella whitish, interior of the aperture fulvous chesnut.

DESHAYES, Anim. sans vert. vol. x. p. 26, note.

Buccinum plicatum, Linnaeus.

Buccinum fibriatū, Gmelin.

Cassis plicaria, Lamarck.

Hab. Moluccas.

We are indebted to M. Deshayes for having identified Lamarck's *Cassis plicaria* with the *Buccinum plicatum* of

Linnaeus . . . are must, however, be taken not to confound it with the *Cassis plicata* of Martini, which is the *C. cruentata* of Lamarek, and without doubt a variety of *C. testiculus*. The plicated sculpture of this species is constant, so also is its singularly compressed growth, and dark brown blotches behind the varices.

Species 25. (Fig. a. Mus. Saul.

Fig. b. Mus. Cuming.)

CASSIS ACHATINA. *Cass. testá ovalá, ventricosá, spirá brevi, anfractibus ovaricosis, levibus, columellá vix rugatá, basi truncatá, labro tenué reflexo, basin versus leviter denticulato: cævereō aut cœrulecente, rufo purpureo longitudinaliter flammatá et arcuatim articulatá: columellá lacteá aut rufescente, labro fuscō nigricante maculato.*

THE AGATE HELMET. Shell ovate, ventricose, spire short, whorls destitute of varices, smooth, columella scarcely wrinkled, truncated at the base, lip thinly reflected, slightly denticulated towards the base; ash or bluish colour, longitudinally flamed and arcuately articulated with red or purple-brown; columella cream-colour or reddish, lip spotted with brownish-black.

LAMARCK. Anim. sans vert. (Deshayes' edit.) vol. x. p. 33.
Hab. Algoa Bay: Hennah.

A well-known species, not of very common occurrence, mainly distinguished by its richly variegated style of painting. Lamarek cites New Holland as its place of habitation, but Mr. Cuming's specimens were collected in Algoa Bay, Cape of Good Hope, by the Rev. Mr. Hennah.



29 - a



29 - b



29 - c



29 - d

CASSIS.

PLATE XI.

Species 29. (Mus. Cuming.)

CASSIS PYRUM. *Cass. testá orato-globosá, pleruque crassi-
sissimá, anfractibus concérvis, levigatis, superná plus
minusve angulatis, interdum subnodosis; columellá
crassá, levigatá, basi obscurè sulcatá, labro reflexo,
incrassato; aurantió-lactéa, nigricante-fusco pone va-
rices maculatá.*

THE PEAR HELMET. Shell ovately globose, generally very thick, whorls convex, smooth, more or less angulated round the upper part, sometimes slightly nodose; columella thickened, smooth, obscurely grooved at the base, lip reflected, thickened; orange cream-colour, spotted with blackish-brown behind the varices.

LAMARCK, Auct. sans vert. (Deshayes' edit.) vol. x. p. 33.
Eadem. *Cassis Zeylanica*, Lamarck.

Hab. New Holland.

I cannot observe any character of sufficient permanence to distinguish Lamarck's *Zeylanica* from *C. pyrum*. The species is sometimes comparatively thin and lightly noduled, but more frequently very much thickened, with the enamel sometimes deposited in rude callous layers.

Species 30. (Mus. Cuming.)

CASSIS SULCOSA. *Cass. testá ovalá, spirá subproducta,
acutá, anfractibus evarciosis, superná concavo-depressis,
sulcis regularibus equidistantibus andique cingulatis;
columellá plicato-rugatá et granosá, labro sapissimè
valde incrassato, intus acide fortiter lirato; griseo-
fulvá, ferrugineo-fusco maculatá et flammatá.*

THE GROOVED HELMET. Shell ovate, spire somewhat produced, sharp, whorls destitute of varices, concavely depressed round the upper part, encircled throughout by regular, equidistant grooves; columella plicately wrinkled and granose, lip frequently very much thickened, sharply and strongly ridged within; greyish fulvous colour, blotched and flamed with rusty brown.

LAMARCK, Auct. sans vert. (Deshayes' edit.) vol. x. p. 34.
Buccinum sulcosum, Born.


Cassidea sulcosa, Bruguire.

Buccinum undulatum, Gmelin.

Hab. Mediterranean.

The common species of the Mediterranean, perfectly distinct from the *C. inflata* and others with which M. Kiener seems to confound it. Owing to its ventricose growth and the regularity with which it is spirally grooved throughout, it has somewhat the aspect of a *Dolium*.

September, 1848.



1000

1000

CASSIS.

PLATE XII.

Species 31. (Mus. Cuming.)

CASSIS CORONULATA. *Cass. testâ suboblongo-ovatâ, spirâ subacuminatâ, apicé acutâ; aufractibâ lœvigatis, undis, apicem versus decussatis, supernè depresso-concavis, ad angulum tenuè muricato-nodosis, aufractu ultimo unicoloris; columellâ expansâ, undulatâ plicato-rugatâ, labro externo acutâ dentato-lirato, basi obsoletè tri-spinoso; livido-albante, fasciis transversis quinque fulvo-citris pallide fuscescitibus, fasciis pallidioribus longitudinalibus nudatis decussatis, nitide picta, fasciis saturatioribus pone varices, labro citriuo-fulvocente tessellata.*

THE CORONATED HELMET. Shell somewhat oblong-ovate, spire somewhat acuminate, sharp at the apex; whorls smooth, shining, decussated towards the apex, depresso concave round the upper part, finely prickly noduled at the angle, last whorl one-varicose; columella expanded, undulately plicately wrinkled, outer lip sharply dentately ridged, obsoletely trispinose at the base; livid white, neatly painted with five rather broad transverse pale fulvous bands, decussated with paler longitudinal waved bands, bands darker behind the varices, lip tessellated with citron fulvous brown.

SOWERBY, TANKERVILLE Catalogue, App. p. xx.

Hab. ——
Philippine Islands; Cuming.

This is certainly not the young of *Cassis glauca*, as stated by M. Kiener; whether it should be regarded as a variety of that species may remain a matter of opinion, but I believe it to be distinct. Though not of very common occurrence it is not rare, and there appears to be no nearer approximation between the species than is represented by the accompanying figures. The *C. corona* is of a more oblong-oval form and perfectly smooth, without the least indication of those curious flattened indentations which characterize the *C. glauca*; the tessellated painting is peculiar and constant, and the spinous processes at the base of the lip are but three in number and very obscure.

Species 32. (Mus. Cuming.)

CASSIS EXARATA. *Cass. testâ ovatâ, spirâ subacuminatâ, apicé acutâ; aufractibâ naiacaricosis, levibus, transversim regulariter acutè soleatis, supernè declivibus, spiraliter costatis, costis creberriam nodoso-gemmatis; columellâ latè concavo-expansâ, profuse et subtilliter*

plicato-rugatâ, labro externo fortiter dentato-lirato; eburnea, maculis quadratis distantibus pallide fulvescentibus tessellata.

THE PLOUGHED HELMET. Shell ovate, spire somewhat acuminate, sharp at the apex, whorls one-varicose, smooth, transversely regularly sharply grooved, slanting at the upper part, spirally ribbed, ribs very closely beaded; columella broadly concavely expanded, profusely and very finely plicately wrinkled, outer lip strongly dentately ridged; ivory white, tessellated with distant pale fulvous square spots.

Hab. —— ?

A new and very distinct species allied in form to the *C. glauca* and *coronata*. Its chief peculiarities consist in the surface being encircled with narrow equi-distant grooves, and in the spire being strongly beaded. The columella is very widely expanded, and profusely sculptured with fine divercating wrinkles.

Species 33. (Mus. Hudson.)

CASSIS GLAUCÀ. *Cass. testâ ovalâ, turgidâ, spirâ subacuminatâ, apicé acutâ; aufractibâ lœvigatis, undique peculiariter indentatis, basi superficialiter sulcatâ, supernè angulatis, ad angulum acutè muricato-nodulosis, apicem versus decussatis, aufractu ultimo unicoloris; columellâ expansâ, exiliter plicato-rugatâ, labro externo dentato-lirato, basi quadrispinoso; livido-glauca, indistinctè fasciatâ, lobris vicinè falcato-citrinâ, apertura fauce purpureo-castanea.*

THE GLAUCOUS HELMET. Shell ovate, swollen, spire somewhat acuminate, sharp at the apex; whorls smooth, everywhere peculiarly indented, superficially grooved at the base, angulated round the upper part, sharply prickly noduled at the angle, decussated towards the apex, last whorl one-varicose; columella expanded, faintly plicately wrinkled, outer lip dentately ridged, four-spined at the base; livid glaucous colour, indistinctly beaded, lips bright fulvous citron colour, interior of the aperture purple-chesnut.

LAMARCK. Anni. sans. vert. (Deshayes' edit.) vol. x. p. 24.
Buccinum glaucum, Linneus.

Cassidea glauca, Bruguière.
Bezzardica vulgaris, Schumacher.

Hab. Philippine Islands; Cuming.

A species of common occurrence, exhibiting little variation of colour or sculpture.

C A S S I S.

	<i>Plate.</i>	<i>Species.</i>		<i>Plate.</i>	<i>Species.</i>
<i>abbreviata</i> , Lamarck	VIII.	18	<i>paucirugis</i> , Menke	VIII.	19
<i>achatina</i> , Lamarck	X	28	<i>pennata</i> , Lamarck	VIII.	20
<i>arcola</i> , (Bucc.) Linn.	IX	24	<i>pila</i> , Reeve	IX.	21
<i>bisulcata</i> , Schubert & Wagner	III.	6	<i>plicaria</i> , Lamarck	X.	27
<i>canaliculata</i> , (<i>Cassidea</i>) Brug.	III.	8	<i>plicata</i> , (Bucc.) Martini	X.	27
<i>coarctata</i> , Gray	VI.	14	<i>plicata</i> , Martini	IV.	10
<i>cornuta</i> , (Bucc.) Linn.	I	2	<i>pomum</i> , Schubert & Wagner	V.	11
<i>coronulata</i> , Sowerby	XII.	31	<i>pyrum</i> , Lamarck	XI.	29
<i>crusacea</i> , Lamarck	IV.	10	<i>recurvirostrum</i> , (Bucc.) Wood	VII.	16
<i>decussata</i> , (Bucc.) Linn.	II.	4	<i>rufa</i> , (Bucc.) Linn.	VIII.	20
<i>erinaceus</i> , Lamarck	VII.	15	<i>Saburon</i> , Adanson	V.	11
<i>exarata</i> , Reeve	XII.	32	<i>semigranosa</i> , Lamarck	I.	3
<i>fasciata</i> , Lamarck	IV.	9	<i>spinosa</i> , (Bucc.) Gronov.	IV.	9
<i>fimbriata</i> , Quoy & Gaimard	VII.	17	<i>sulecosa</i> , (Bucc.) Born	XI.	30
<i>flammea</i> , (Bucc.) Linn.	V.	12	<i>tenuis</i> , Gray	VI.	13
<i>glauea</i> , (Bucc.) Linn.	XII.	33	<i>testiculus</i> , (Bucc.) Linn.	IV.	10
<i>granulosa</i> , Lamarck	IX	22	<i>tortuata</i> , Reeve	I.	1
<i>inflata</i> , (Bucc.) Shaw	IX.	22	<i>tuberosa</i> , (<i>Cassidea</i>) Brug.	III.	7
<i>Japonica</i> , Reeve	IX.	23	<i>turgida</i> , Reeve	X.	25
<i>labiata</i> , Chemnitz	I.	2	<i>undata</i> , Deshayes	X.	26
<i>loefia</i> , Kiener	VIII.	18	<i>vibex</i> , (Bucc.) Linn.	VII.	15
<i>Madagascariensis</i> , Lamarck	II.	5	<i>zebra</i> , Lamarck	X.	26
<i>Muscica</i> , Kiener	VI.	13	<i>Zeylanica</i> , Lamarck	XI.	29

MONOGRAPH
OF THE GENUS
T U R R I T E L L A.

He compelled
The Proteus shape of Nature, as it slept,
To wake, and lead him to the caves that held
The treasures of the secrets of its reign.—*Shelley*.



TURRITELLA.

PLATE I.

Genus TURRITELLA. Lamarck.

Testa elongato-turrata, lanceolato-acuminata, unguinam umbilicata, anfractibus unnerosis, sepsissimè transversum costatis vel striatis, unguinum longitudinaliter exsculptis, evanescens, unguinum sponiosis aut unicalis; apertura parvissimata, plus minusse rotundata, marginibus disjunctis, labro septè medio sinuato.

Shell elongately turreted, lanceolate acuminate, never umbilicated, whorls numerous, mostly transversely ribbed or striated, never longitudinally sculptured, destitute of varices, never spined or prickly; aperture rather small, more or less rounded, margins disjoined, lip often sinuated in the middle.

The genus *Turritella* affords excellent characters for the distinction of species. Constructed upon the simple type of an enlarging tube, the shell is developed with geometrical symmetry in short convolutions, varying from fifteen to thirty in number, the same number in each species, forming a sharply acuminate screw. The whorls are sometimes flattened or a little concave, and being always contiguous there is no indication of any umbilicus. No varices or tubercles are formed, as in *Cerithium*, nor is there any periodical development of sculpture on the margin of the aperture. The ribs and striae of *Turritella* range transversely, that is, spirally, never longitudinally, and are rarely granulated or nodulated. The best characters for the discrimination of species reside in the first eight or ten whorls from the apex; beyond this point the sculpture varies slightly or becomes obsolete, as the shell approaches maturity. In colour the reverse of this condition is presented, the pattern becoming gradually darker and more defined towards the aperture,

The genus *Turritella* has been somewhat neglected by conchologists. Only eleven species were described by Lamarck, and a monograph lately published in Paris by M. Kiener includes less than half the number now before me from the cabinets of Mr. Cuming, Mr. Metcalfe, Mr. Hanley, Sir. Edward Belcher, the British Museum, and the Banksian Collection in the museum of the Linnean Society; amounting to sixty-five all distinctly characterized in respect of form, sculpture, and painting.

The Linnean species have not been correctly interpreted by Lamarck, Deshayes, or Kiener, for want of access to the original types of the 'Systema Naturae' preserved in the museum of the Linnean Society of London. The *Turbo acutangulus* of Linnaeus proves to be a simple variety of the *T. duplicatus*, and not the shell assigned to it by M. Deshayes. The shell described as *Turbo variegatus* in the 'Systema Naturae,' though differing materially in

general aspect from that which served for the description of the *Turbo inbriatus*, may, nevertheless, be traced to belong to the same species; and the *Turbo angulatus* of Linnaeus proves to be identical with Lamarck's *Turritella fuscosta*. The British species *Turbo terebra*, Linn. in 'Fauna Suecia' (not of Syst. Nat.) and Montagu, must be referred to the *T. cornuta* of Lamarck,

The *Turritella brevialis* of Lamarck (*Le Mesal* of Adanson) constitutes the type of a new genus, founded by Mr. Gray, under the name *MESALIA*, to which I venture to add the *Turritella laetea* of Möller, and a very characteristic species from the Cumingian collection. A genus has been also introduced by Mr. Gray, with equal judgment, under the name *EGLISIA*, founded on the *Turritella spirata* of the Tamkerville Catalogue; and to this is added a species collected by Mr. Cuming at one of the Philippine Islands.

The *Turritelle* have a wide range of habitation and dwell at a considerable depth as well as near the surface. The *T. cornuta* has been recently dredged off the Shetland Islands by Mr. M'Andrew at a depth of many fathoms, beyond the influence of light and colourless. Only two comparatively small species besides this, extend so far from north of the equator as the Mediterranean. The greater number are from the Eastern Seas and the shores of Central America, and there are a few known from New Zealand, Australia, and Van Dieman's Land.

Species 1. (Mus. Cuming.)

TURRITELLA UNGULINA. Turr. testa acuminato-turrata, anfractibus quindecim, convexis, levibus, regulariter decem-striatis, intersticibus superficie ad latera sulcatis, apertura suboblongo-ovali; castaneo-rufâ, aperturam versus nudatato-nigrâ.

THE NAIL-BRUISED TURRITELLA. Shell acuminate turreted, whorls fifteen in number, convex, smooth, regularly ten striated, interstices superficially grooved, aperture rather oblong-oval; chestnut-red, burnt black towards the aperture.

Turbo angulatus, Linnaeus, Syst. Nat. (12th. edit.) p. 1240.
Turritella fuscosta, Lamarck.

Hab. Senegal, Mouth of the Gambia, West Africa.

The original specimen which served Linnaeus for the type of his *Turbo angulatus*, still preserved in the museum of the Linnean Society of London, proves, as Mr. Hanley first pointed out to me, to be Lamarck's *T. fuscosta*. The locality "Seas of Europe" erroneously given in the 'Systema Naturae' led Dr. Pulteney, and recently M. Deshayes, to assign the Linnean species to the only *Turritella* existing north of the Mediterranean; on the

authority of M. Kiener, however, who has the same access to the types of the Lamarckian species as we have to those of Linnaeus, the European representative of the genus is the *T. cornea*, Lamarck. M. Deshayes' *T. ungulina* (Anim. sans vert. vol. ix. p. 260) must therefore be cancelled.

Species 2. (Mus. Cuming.)

TURRITELLA DUPLICATA. *Turr. testā obeso-turbinatā, crassā, ponderosā, anfractibus ad sedecim, primis per paucis multi-carinatis, carinū unicā, deinde duabus multo eminentioribus, carinis parris cito evanidis, majoribus gradatim minus elatis, aperturā valde sinuatā; fulvescente-lacteā, anfractuum parte supremā saturnato.*

THE DUPLICATE TURRITELLA. Shell stoutly turbinated, thick, ponderous, whorls about sixteen in number, the first few many-keeled, one keel and afterwards two to three very conspicuously elevated, small keels quickly disappearing, larger keels gradually less raised, aperture considerably sinuated; fulvous cream-colour, uppermast part of the whorls of rather darker colour.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. ix. p. 251.
Turbo duplicatus, Linnaeus.

Eadem var. *Turbo acutangulus*, Linnaeus.
An eadem var.? *Turbo replicatus*, Linnaeus.

Hab. Eastern Seas.

The *Turbo acutangulus* of Linnaeus is truly a variety of his *Turbo duplicatus*, as may be seen on reference to the original specimen in the Linnaean collection; but not so the species to which M. Deshayes has assigned that title, figured by M. Kiener at Pl. 2. Fig. 26, and in the accompanying plate at Fig. 4.

The first few whorls of *Turritella duplicata* are encircled by numerous keels, one of which is particularly sharp and prominent. The smaller keels quickly disappear, and a second of equal prominence with the principal keel makes its appearance in most specimens at about the tenth whorl. Linnaeus's *Turbo acutangulus* is a specimen in which the second prominent keel is not developed. The first sharp-angled keel is equally prominent in all varieties of the species during the first eight or ten whorls.

Species 3. (Mus. Cuming.)

TURRITELLA TEREbra. *Turr. testā lanceolato-turritā, acutissimè attenuatā, anfractibus ad quinque et viginti, medio tumidiusculis, spiraliter acutè sex-carinatis, suturis lacibas, excavatis; castaneo-brunneā aut fulvescente.*

THE AUGUR TURRITELLA. Shell lanceolately turreted,

very sharply attenuated, whorls about twenty-five in number, rather swollen in the middle, spirally sharply six-keeled, sutures smooth, excavated; chestnut-brown or fulvous.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. ix. p. 253.
Turbo terebra, Linnaeus, Syst. Nat. (not of Fauna Suecica.)

Turritella Archimedis, Dillwyn, Mawe.
Hab. Philippine Islands; Cuming.

The magnificent specimen here represented, is very similar to that selected by Martyn for the frontispiece of his 'Universal Conchologist.' Struck by the perfect symmetry of its convolutions, he distinguished it by the title of *Αφροδίτη*, significant of its graceful form and marine origin, and truly the geometrical proportions of this species are the perfection of beauty. The specimen figured was dredged by Mr. Cuming in fine sand at low water at the Island of Negros, one of the Philippines. The *T. terebra* of the 'Fauna Suecica' and of British authors (*T. Linnei* and *ungulinus*, Deshayes, *T. communis*, Philippi) is the *T. cornea* of Lamarck.

Species 4. (Mus. Cuming.)

TURRITELLA ATTENUATA. *Turr. testā acutissimè attenuatā, anfractibus ad viginti, spiraliter striatis, anterioribus medio acutè et tenuè mucarinatis striis evanidis, suturis subexcavatis; fuscoceruleo-albâ, livida, anfractuum parte superiori saturatore, aperturā sinuatā.*

THE ATTENUATED TURRITELLA. Shell very sharply attenuated, whorls about twenty in number, spirally striated, latter whorls sharply and finely onc-keeled in the middle, striae disappearing, sutures rather excavated; light fulvous white, livid, upper part of the whorls darker, aperture sinuated.

Turritella acutangulus, Deshayes, Anim. sans vert. vol. ix. p. 259 (not *Turbo acutangulus*, Linn.).

Hab. —?

The "carinā majore acutā" by which Linnaeus characterizes his *Turbo acutangulus*, constitutes a very prominent feature in that and the *T. duplicatus*, and proves them, beyond doubt, to be one and the same species. In the species under consideration there is no indication of this character, the whorls being merely encircled from the apex with ten elevated striae or fine ridges which become more numerous, fainter, and almost obsolete. The species is further distinguished by its more attenuated growth, and a central fine keel or cord which is not developed until after the tenth or twelfth whorl, and the keel-like striae become obsolete.



5. ♂



5.



6. ♂



7.

TURRITELLA.

PLATE II.

Species 5. (Mus. Cuming.)

TURRITELLA LEUCOSTOMA. *Turr. testā subpyramidalis acuminate, angustā, acutissimē lanceolatā, solidissimā, anfractibus ad viginti, superne contractis, suturis excavatis, infernō extorsuō projectis, planulatis, spiraliiter sex-liratis, liris angustis, subdistantib⁹, interstatis subtilissimē striatis; fulvescente-albā, liris nitide aurantio-rufo articulatis.*

THE WHITE-MOUTH TURRITELLA. Shell somewhat pyramidalis acuminate, narrow, very sharply lanceolate, rather solid, whorls about twenty in number, contracted round the upper part, sutures excavated, projected outwardly round the lower part, flattened, spirally six-ridged, ridges narrow, rather distant, interstices very finely striated; fulvous white, ridges neatly articulated with orange-brown.

VALENCIENNES, Zool. Voy. Humboldt et Bonpland, vol. ii. p. 275.

Hab. Gulf of Nicoya (in sandy mud at the depth of eleven fathoms); Cuming.

Distinguished by the truly regular style in which the shell is turreted. The whorls contracted round the upper part and projecting outwards gradually towards the base, look like an attenuated pyramid of thimbles; and they are spirally encircled throughout by a row of neatly articulated ridges.

Species 6. (Fig. a and b, Mus. Cuming.)

TURRITELLA BRODERPIANA. *Turr. testā elongato-acuminate, subcylindraceo-tunnidā, anfractibus octodecim, undique crebellinē impresso-striatis, perpans primis unicarinatis, medianis planulatis, ceteris media depresso-concavis, suturis interdūa indistinctis; fulvescente, ceruleo-fuscente nebulatā, lineis rubido-fusca transversis brevibus interruptis, in strigis undulatis frequenter dispositis, densē testigivosis.*

BRODERIP'S TURRITELLA. Shell elongately turreted, somewhat cylindrically swollen, whorls eighteen in

number, very closely impressly striated throughout, the first few whorls one-keeled, the next flattened, then depressly concave in the middle, sutures sometimes indistinct; light fulvous colour, clouded with blue-brown and thickly freckled with short interrupted red-brown transverse lines, mostly disposed in waved streaks.

D'ORBIGNY, Voy. dans l'Amér. Mérid. Moll. p. 388.

Turritella marmorata, Kiener.

Hab. Payta, Peru (in fine sand at a depth of about eight fathoms); Cuming.

The *T. Broderipiana* has a very characteristic style of painting, consisting of a profusion of short irregular linear dashes, ranged transversely, but mostly disposed in confused longitudinally waved streaks; and over these, there is generally a scorched blue-brown stain of colouring. The shell is of an uncouth subcylindrical growth, spirally encircled throughout with fine grooved striae.

Species 7. (Mus. Cuming.)

TURRITELLA BACILLUM. *Turr. testā elongato-acuminate, anfractibus octodecim ad viginti, convexis, superne subplanulatis, spiraliiter quinque- vel sex-carinatis, carinis angustis, subirregulariter distantibus, aperturam versus ferè evanidis; livido-futescente, carinis saturatioribus.*

THE STAFF TURRITELLA. Shell elongately acuminate, whorls eighteen to twenty in number, convex, rather flattened round the upper part, spirally five- to six-keeled, keels narrow, somewhat irregularly distant, almost disappearing towards the aperture; livid fulvous colour, the keels being of darker hue.

KIENER, Icon. coq. viv., p. 5, pl. 4, f. 1.

Hab. Ceylon.

The spiral keels of this species are not developed in the sharp prominent style of the *T. terebra*. Towards the aperture they nearly disappear, leaving their colour on the surface as though they had sunk, so to speak, into the substance of the shell.



TURRITELLA.

PLATE III.

Species 8. (Mus. Cuming.)

TURRITELLA TIGRINA. *Turr. testá pyramidal-turritá, subangustá, solidiusculá, anfractibus octodecim ad viginti, spiraliter quinque- vel sex-liratis, costá tumidá declivi ad basim angulatis; albida, flammis oblique purpureo-nigrae profuse pictá.*

THE TIGER-STRIPED TURRITELLA. Shell pyramidal, turreted, somewhat narrow, rather solid, whorls eighteen to twenty in number, spirally five- or six-ridged, angled at the base with a slanting swollen rib; whitish, profusely painted with oblique purple-black flames.

KIENER, Icon. coq. viv., p. 29. pl. 4. f. 2.

Hab. Gulf of California.

The peculiarities of this species, which approaches the *T. lentiginosa* and *goniostoma*, consist in its more solid structure and narrow pyramidal growth, whilst the whorls are swollen to an obtuse angle round the base. The colouring is disposed in more definite stripes, especially towards the apex. In the species just mentioned the first few whorls are conspicuously keeled round the middle, but in the *T. tigrina* there is little variation from the general pattern.

Species 9. (Mus. Cuming.)

TURRITELLA LENTIGINOSA. *Turr. testá subobeso-turritá, anfractibus od viginti, primis perpaucis carinatis, medianis planulatis, ceteris medio convexis, basi tumidis, omnibus creberrime spiraliter striatis et salentis, aperturam versus laniis irregularibus iubricatis; albida, lineis subtilibus fusca, brevibus, interrupitis, spiraliter lentiginosa, maculis strigisve nigricantibus subindistinctis oblique nebulatá, apice interdum violaceo.*

THE FRECKLED TURRITELLA. Shell rather stoutly turreted, whorls about twenty in number, the first few carinated, the middle whorls rather flattened, the rest convex in the middle and swollen round the base; the whole very finely spirally grooved and striated, and imbricated with irregular laminae towards the aperture; whitish, spirally freckled with short interrupted fine brown lines, and obliquely clouded with rather indistinct blackish streaks and blotches, apex sometimes tinged with violet.

Hab. Payta, Peru (in coarse sand at a depth of about five fathoms); Cuming.

A shell of rather stout growth, generally more or less imbricated with rude irregular laminae near the outer lip. The first seven or eight whorls are distinctly keeled round the middle, the next seven or eight are flattened in a manner almost to obliterate the suture, and the remainder are sunk in the middle and swollen above and below, showing the sutures more distinctly.

The painting of this species consists of a profusion of extremely fine, short, brown, transverse lines or dashes, over which is an obscure confused marbling of oblique blackish blotches; in some specimens this is wanting and its place is supplied by a dull greyish tinting. Care should be taken not to confound the *T. lentiginosa* with the following species which is characterized by a different style of painting and a slight difference of form.

Species 10. (Fig. a and b, Mus. Cuming.)

TURRITELLA GONIOSTOMA. *Turr. testá subcylindraceo-acuminatá, anfractibus octodecim ad viginti, spiraliter tenuistratis, primis perpaucis carinatis, medianis planulatis, ceteris medio concavis, costá interdum prominulá et crenulá, supra et infra angulatis, anfractu ultimo sapé prope aperturam laniis subimbricatis; albá, griseo- vel rufescente-nigro dñe maroratá.*

THE ANGULAR-NOUTH TURRITELLA. Shell somewhat cylindrically acuminate, whorls eighteen to twenty in number, spirally finely striated, the first very few whorls keeled, the middle whorls smooth, the rest concave in the middle, angled above and below with a rib, which is sometimes prominent and crenulated, last whorl often slightly imbricated near the aperture with laminae; white, thickly marbled with grey or reddish-black.

VALENCIENNES, Zool. Voy. Humboldt et Bonpland, vol. ii. p. 275.

Hab. Payta; Salango, West Columbia; Guacomayo, Central America (in sandy mud at depths of from five to ten fathoms); Cuming.

The variation of sculpture in the several stages of the growth of this species is very similar to that of the pre-

TURRITELLA.—PLATE III.

ceding, save that the latter whorls are more sharply angled above and below. The ribs which impart this angular structure to the whorls just before the growth of the shell is matured, are much more strongly developed in some specimens than in others; they are nevertheless sharply defined, and never have the obtuse swollen appearance presented in the *T. lentiginosa*. The sutural

grooving of the middle whorls is nearly obliterated, as in that species, by their flattened and close contact. In colouring, the *T. goniostoma* is characterized by a very dense marbled or mottled pattern of grey-black, sometimes a little rusty; and exhibits no indication of the short transverse linear dashes which serve to distinguish the *T. lentiginosa*.



11 b.



11 a.



12 a.



12 b.



13.



15.



14.

TURRITELLA.

PLATE IV.

Species 11. (Fig. *a*, Mus. Belcher. Fig. *b*, Mus. Cuming.)

TURRITELLA NODULOSA. *Turr. testā acuminato-turritā, unfractibus quatuordecim ad quindecim, transversio fortiter impresso-striatis, obscure uni- bi-angulatis, ad angulos nodulosos, medio concavis; sortidē griseo-albā, flammulis fasciis inter nodulos longitudinaliter tinctā.*

THE NODULOUS TURRITELLA. Shell acuminate turreted, whorls fourteen to fifteen in number, transversely strongly impressively striated, obscurely one- two-angled, nodulous at the angles, concave in the middle; dull greyish-white, longitudinally stained with brown flames between the nodules.

KING, Zool. Journ. vol. v. p. 347.

Turridella papillosa, Kiener.

Hab. Gulf of Dulce, Central America (found in sandy mud at the depth of six to ten fathoms); Cuming.

Distinguished by its nodulous structure, which is a character of rare occurrence in this genus.

Species 12. (Fig. *a* and *b*, Mus. Cuming.)

TURRITELLA CARINIFERA. *Turr. testā pyramidalis-turritā, basi tumida et acutè angulata, unfractibus quatuordecim ad quindecim, superne decoloratis, inferne acutè bicarinatis, undique spiratiliter impresso-striatis, striis subtilissimè nodulato-corrigatis; nivea aut pallide purpureo-roseata.*

THE KEELED TURRITELLA. Shell pyramidal turreted, swollen and sharply angled at the base, whorls fourteen to fifteen in number, slanting round the upper part, sharply two-keeled round the lower, spirally impressively striated, striae very finely undulately wrinkled; snowy white or pale purple-pink.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. ix. p. 258.
Hab. —?

This remarkable shell has been thought to belong to a *Fervetulus*, but its growth is perfectly regular.

Species 13. (Mus. Cuming.)

TURRITELLA CUMINGII. *Terr. testā sub-luncolato-pyramidalis, crassissimā, unfractibus ad cinqū, planodecībus, inferne tumidis et obtuse angulatis, spiratiter quinqūlateralis, liris angustis, acutincis, regularibus; fluorescente-albā, strigis maculatisque purpureofuscis undique obliquè nebulatā.*

CUMING'S TURRITELLA. Shell rather lanceolately pyramidal, rather thick, whorls about twenty in number, flatly slanting, swollen and obtusely angled round the lower part, spirally five-ridged, ridges narrow, rather sharp, regular; brownish white, obliquely clouded throughout with purple-brown streaks and blotches.

Hab. Panama (in mud at depths from eleven to sixteen fathoms); Cuming. Conchagua; Belcher.

Several specimens of this fine species, which I have the pleasure to dedicate to Mr. Cuming, were dredged by him at Panama. It was also collected by Sir E. Belcher at Conehagua during the voyage of the Sulphur. Each whorl is swollen outwardly at the base so as to form an obtuse angle, and the five ridges which encircle them are continued without change throughout the spire.

Species 14. (Mus. Cuming.)

TURRITELLA COLUMNARIS. *Turr. testā acutissimè lanceolato-turritā, ad basin acutè angulatā, unfractibus ad triginta, priuia bicarinatis, carinis approximatis, ceteris planulatis, decoloratis, liris obsolete granulatis, suturis excavatis; griseo-albā, strigis purpureo-spadiceis oblique undulatis nebulatā, basi purpurascens tinctā.*

THE COLUMNAR TURRITELLA. Shell very sharply lanceolately turreted, sharply angled at the base, whorls about thirty in number, the first two-keeled, keels approximated, the rest rather flattened, ten-ridged, ridges obsolete granulated, sutures excavated; greyish-white, clouded with purple-fawn obliquely waved streaks, base tinged with purple.

KIENER, Icon. coq. viv. p. 10. pl. 7. f. 1.

Hab. Ceylon.

This species has the largest number of whorls, and is the most elongated form of the genus.

Species 15. (Mus. Cuming.)

TURRITELLA BANKSII. *Tarr. testā pyramidalis-turritā, crassā, ad basin tumidatā et obtuse angulatā, unfractibus quindecim, undique minitissimā granuloso-reticulatis, concavis, liris medianā aliisque minoribus irregularris cingulatis, marginis inferiori rotundatis, suturis indistinctis, columellā peculiariter contorto-productā, aperturā subquadratā: cinereo-olivaceā, nigro alboque cavigalā, liris siagro articulatis.*

TURRITELLA.—PLATE IV.

BANKS' TURRITELLA. Shell pyramidal, thick, rather swollen and obtusely angled at the base, whorls fifteen in number, very minutely granulously reticulated throughout, concave, encircled with a middle ridge and other irregular smaller ones, rounded at the lower edge, sutures indistinct, columella peculiarly produced in a twisted manner, aperture somewhat square; ash-olive, mottled with black and white, ridges articulated with black.

GRAY, MSS. Brit. Mus.

Hab. Panama (dredged from sandy mud at the depth of ten fathoms): Cuniling.

A solid pyramidal shell in which the whorls are convex in the middle, rounded at the lower edge, and encircled with neatly articulated ridges, one of which is central and prominent.



3 - 1a]



16



19



18



21



22

TURRITELLA.

PLATE V.

Species 16. (Fig. *a* and *b*, Mus. Cuming.)

TURRITELLA BREVIALIS.

For description of this Lamarekian species see Genus MESALIA, Gray, immediately following the present.

For Sp. 16, see Pl. X.

Species 17. (Mus. Cuming.)

TURRITELLA TRISULCATA. *Terr. testá acuminato-turrítá, anfractibus ad octodecim, transversis conspicue tricostatis, costis distantibus, sulcis intermetatis excavatis; albidá, apicem versus violaceat, flammulis perpansis irregularibus aurantio-fusco oblique picta.*

THE THREE-GROOVED TURRITELLA. Shell acuminately turreted, whorls about eighteen in number, transversely conspicuously three-ribbed, ribs distant, intermediate grooves hollowed; whitish, tinged with violet towards the apex, obliquely painted with a very few irregular orange-brown flames.

LAMARCK, Anni. sans vert. (Deshayes's edit.) vol. ix. p. 256.
Hab. Red Sea; Kiener.

The only species which is prominently three-ribbed. The painting is well characterized by Lamarck in the words, "ses flammules sont éparses."

Species 18. (Mus. Cuming.)

TURRITELLA ANNULATA. *Terr. testá lanceolato-turrítá, anfractibus viginti, spiraliiter striatis, superne declivibus, deinde concavis et bicarinatis, carinis aperturas versus distantiioribus, evanescuntioribus, et interdum duplicitis; fulvo-spodicæ, longitudinaliter obscurè albinebulata.*

THE RINGED TURRITELLA. Shell lanceolately turreted, whorls twenty in number, spirally striated, slanting round the upper part, then concave and two-keeled, keels becoming gradually more distant, towards the aperture, fainter, and sometimes duplicate; fulvous-brown, longitudinally obscurely clouded with white.

KIENER, Icon. coq. viv. p. 20. pl. 13. f. 1.
Hab. Mouth of the Gambia, West Africa.

In the first ten whorls of this species, according to the specimen before me, the whorls are prominently developed and contiguous; after this period of its growth they become

gradually fainter and more distant, separating at length into two faint ridges.

Species 19. (Mus. Cuming.)

TURRITELLA VARIEGATA. *Terr. testá pyramidalis-turrítá, anfractibus quindecim ad sedecim, perpansis primis superne declivibus, inferne excavatis, cæteris plano-concavis, supra suturas transversulis, striis confertis linearisque quadratæ vel quinque subtilibus, distantiolis, interduo obsolete granulosis, cingulatis; albidâ vel purpurascente-spodicæ, rufo-fusco punctatâ et longitudinaliter variegatâ, liris interdum fusco alboque articulatis, aperture fauce purparascente.*

THE VARIEGATED TURRITELLA. Shell pyramidaly turreted, whorls fifteen to sixteen in number, the first few slanting round the upper part, excavated beneath, the rest flatly concave, rather swollen above the sutures, encircled with four or five distant sometimes obsolete granulated ridges; whitish or purple-tinged fawn, dotted and longitudinally variegated with reddish-brown, ridges sometimes articulated with brown and white, interior of the aperture purplish.

Turbo variegatus, Linn. Syst. Nat. (12th edit.) p. 1240.
Turritella imbricata, Lamarek.

Var. elongata. *Turbo imbricatus*, Linnaeus.

Hab. Island of Margarita, West Indies; Dr. Hornbeck.

The original type of the *Turbo variegatus* in the Linnaean collection is a shell of immature growth, but extremely well characterized and highly coloured. The shell which served for Linnaeus's *Turbo imbricatus* is, on the contrary, in the worst possible condition, perfectly bleached, and only retaining a sufficient amount of sculpture to identify it with the elongated variety represented by Kiener at Pl. IX. fig. 2 *a*. A comparison of these shells with specimens in Mr. Cuming's collection and in the British Museum (for one of which illustrative of the elongated variety, see Pl. IX. fig. 19 *b*) has enabled me to determine their affinity with satisfaction; and the name *variegatus* should be used in preference to that adopted by Lamarck, from the fact of its being a more appropriate one, and that which was assigned by Linnaeus to the species in its normal form and condition.

Species 20. (Mus. Cuming.)

TURRITELLA BICINGULATA. *Terr. testá acuminato-turrítá, anfractibus octodentaria, subtiliter striatis, superne de-*

TURRITELLA.—PLATE V.

presso-excurrentis, deinde bicostatis, costis prominentibus, tumidis, planisculis, distantibus, interstilio excavato, sutura subprofundè excavatis; fulvo-albidâ, rufo-fusco profuse flammatâ et variegatâ, areâ inter costas parum pictâ, basi et apertura fauce purpureo-roseis.

THE TWO-GIRDLED TURRITELLA. Shell acuminate turreted, whorls eighteen in number, finely striated, deeply excavated round the upper part, then two-ribbed, ribs prominent, swollen, rather flattened, distant, with the interstice excavated, sutures rather deeply excavated; fulvous white, profusely flamed and variegated with red-brown, area between the ribs but little painted, base and interior of the aperture purple-rose.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. ix. p. 256.

Turritella biangulata, De Blainville.

Hab. Porto Praya, Cape de Verd Islands.

This fine species from the Cape de Verd Islands is distinguished by a pair of conspicuous swollen keels and a bold profuse style of painting; the base and aperture are of a peculiar purplish-hue.

At Pl. VII. Sp. 33, is another species very similar to

this, collected by Capt. Sir E. Belcher during the Voyage of the Sulphur.

Species 21. (Mus. Cuming.)

TURRITELLA TORULOSA. *Turr. testâ acuminato-turritâ, crassiusculâ, anfractibus quindecim, rotundatis, striatis, primis bicostatis, ceteris granatis quadricostatis, costis obsoletè granatis, duabus inferioribus fortioribus; fulvescente-albâ, rufo-fusco punctatâ et eximie flammulatâ.*

THE CORDED TURRITELLA. Shell acuminate turreted, rather thick, whorls fifteen in number, rounded, striated, first whorls two ribbed, the rest gradually four-ribbed, ribs obsoletely grained, the two lower ribs always the stronger; light fulvous white, dotted and delicately flamed with red-brown.

KIENER, Icon. coq. viv. p. 18. pl. 6. f. 3.

Hab. —?

The whorls are more than usually rounded in this species, and they are corded with four rather prominent ribs, the upper two of which are not developed until the shell has advanced to its ninth or tenth convolution. The painting has a peculiarly dotted character.



24.



25.



26.



27.



28.



29.



30.

TURRITELLA.

PLATE VI.

Species 22. (Mus. Walton.)

TURRITELLA EXOLETA. *Turr. testá acuminato-turritá, crassi-serrulá, ad basin obtuse angulatá, anfractibus sedecia ad septendecim, superne et inferne rudi costatis, medio concavis, interquatiter excavatis, septis levibus lamelli-formibus interdum longitudinaliter iubricatis, costá inferiori duplice, apertura marginé medio sinuata; sordidó ceraso-albá, rufo-aurantió obscurè flammatá, costis maculatis.*

THE DECAYED TURRITELLA. Shell acuminate turreted, rather thick, obtusely angled at the base, whorls sixteen to seventeen in number, rudely ribbed above and below, concave in the middle, unevenly excavated, sometimes longitudinally imbricated with thin lamelliform septa, lower rib duplex, margin of the aperture sinuated in the middle; dull waxen white, obscurely flamed with orange-brown, ribs spotted with the same.

Turbo croesus, Linneus, Gmelin's edit. p. 3607.

Turbo torcularius, Born.

Hub. Martinique, Guînea; Kiener.

The chief peculiarity of this remarkable species consists in the whorls being rudely excavated, and imbricated across the hollow at frequent intervals with thin lamelliform septa. Specimens, however, are more commonly seen without the septa; not merely worn, but in which they apparently never existed.

The shell in the Linnaean collection, numbered to refer to the 'Systema,' is a worn *T. bicarinata*, and to that species the present has been assigned by Dillwyn. I believe, however, that Linnaeus's description refers to the shell under consideration, and the figure of Bonani, to which he refers, goes far to support that opinion.

Species 23. (Mus. Cuming.)

TURRITELLA CINGULATA. *Turr. testá pyramidalis-turritá, crassá, subponderosa, anfractibus ad septendecim, planulatis, spiraliiter fortiter tricostatis, costis planatis, gravo-crenatis; ferrugineo-albida, costis castaneo-nigris.*

THE GIRDLED TURRITELLA. Shell pyramidal turreted, thick, rather ponderous, whorls about seventeen in number, rather flattened, spirally strongly three-ribbed, ribs flattened and granosely crenated; rusty white, ribs chestnut-black.

SOWERBY, Taurerville Catalogue, App. p. xiii.

Turritella tricarinata, King.

Hub. Valparaiso (in coarse gravel at the depth of six fathoms); Cuming.

Mr. Cuming speaks of this very characteristic species as being very abundant at Valparaiso, having frequently brought up two bushels of living specimens at a time, mingled with coarse gravel.

Species 24. (Mus. Cuming.)

TURRITELLA FLAMMULATA. *Turr. testá acuminato-turritá, anfractibus octodecim, perpansis primis bicostatis, ceteris rotundatis, septem-costatis, costis medianis latiusculis, ceteris striisque gradatim angustioribus; fuligineo-albá, roseo tinctá, purpureo-fusco punctatá et marmoratá, basi roseá.*

THE FLAMED TURRITELLA. Shell acuminate turreted, whorls eighteen in number, the few first two-ribbed, the rest rounded, seven-ribbed, middle ribs rather broad, the remainder on either side gradually narrower; fulvous-white, stained with rose, dotted and marbled with purple-brown, base rose.

KIENER, Icon. coq. viv. p. 7. pl. 5. f. 1.

Le Ligar, Adanson.

Turritella Ligar, Deshayes.

Hub. Muscat, Red Sea.

The shell here represented from the above-mentioned locality is, no doubt, *Le Ligar* of Adanson, a name which cannot correctly be adopted in the Latin. It will be observed that both the sculpture and colouring of the species are peculiar.

Species 25. (Mus. Belcher.)

TURRITELLA CEREA. *Turr. testá elongato-turritá, crassissimul, anfractibus octodecim ad viginti, superne subdeclinibus, inferne transversalibus, spiraliiter sexcarinatis, carinis angustis, acentis, apertoram versus evanidis; fuligineo-albá, anfractuum parte superiori falco-rufescente tinctá.*

THE WAXEN TURRITELLA. Shell elongately turreted, rather thick, whorls eighteen to twenty in number, somewhat inclined round the upper part, rather swollen round the lower, spirally six-keeled, keels narrow, sharp, fading towards the aperture; fulvous white, upper portion of the whorls tinged with fulvous-red.

Hub. —?

The characters by which this species differs from *T. bacillum* are apparently of trivial importance, but they are constant in several specimens that have come under my observation. The whorls are more swollen round the lower part, and the keels are finer, more numerous, and not distinguished by any darker colouring.

Species 26. (Mus. Cuming.)

TURRITELLA CROCEA. *Turr. testā pyramidalī-turritā, solidiusculā, anfractibus viginti aut pluribus, convexo-planulatis, spiraliter quinque-ad decem-carinalis, carinis subtilibus, inaequidistantibus; croceo-brunneā, anfractuum parte superiori saturatiore, suturis pallidioribus.*

THE SAFFRON-TINGED TURRITELLA. Shell pyramidalily turreted, rather solid, whorls twenty or more in number, convexly flattened, spirally five to ten keeled, keels fine, inequidistant; saffron-brown, upper part of the whorls darker, sutures paler.

KIENER, Icon. coq. viv. p. 24. pl. 11. f. 2.
Hab. China.

The flattened surface of the whorls and light colouring of the sutures constitute the characteristic features of this species. The spiral keels are unusually numerous, but the alternate ones are sometimes only faintly developed, leaving five only of a prominent character.

Species 27. (Mus. Belcher.)

TURRITELLA SANGUINEA. *Turr. testā subelongato-turritā, anfractibus octodecim ad viginti, convexis, spiraliter impresso-sulcatis, sulcis irregularibus, liris intermediis*

planiusculis; fulvo-albicante, liris maculis sanguineis transversis, oblongo-quadratis, profusè pictis.

THE BLOOD-STAINED TURRITELLA. Shell somewhat elongately turreted, whorls eighteen to twenty in number, convex, spirally impressly grooved, grooves irregular, intermediate ridges rather flattened; fulvous white, ridges profusely painted with transverse oblong-square blood-stained spots.

Hab. California.

A very characteristic species in which the entire surface is closely painted with transverse oblong-square dull blood-stained spots.

Species 28. (Mus. Cuming.)

TURRITELLA GEMMATA. *Turr. testā pyramidalī-turritā, crassiusculā, ad basin angulatā, anfractibus ad quindecim, supernē bicostatis, medio concavis, infernū unicostatis, costis superioribus granatis, inferiori levē; albā, rufescente-fusco hic illuc pallide tinctā.*

THE GEMMED TURRITELLA. Shell pyramidalily turreted, rather thick, angled at the base, whorls about fifteen in number, encircled with two granulated ribs round the upper part, and one smooth one round the lower, concave in the middle; white, faintly stained here and there with light orange brown.

Hab. — ?

The structure of this species approaches somewhat in pattern to that of the *T. radula*, though it will be found to differ on comparison in the lower spiral rib being smooth except on the last whorl. The species differ materially in form, the *T. gemmata* being shorter and composed of much fewer whorls.



200.



201.



202.



203.



204.



205.



206.

TURRITELLA.

PLATE VII.

Species 29. (Mus. Cuming.)

TURRITELLA COCHLEA. *Turr. testā subulato-turrīdā, crassiusculā, anfractibus quindecim, spiraliter striatis, superē depresso-excavatis, medio bicarinatis, carinis angustis, elevatis, acutis, distantibus, intersticio excavato, aperturā parvā, albida.*

THE SCREW TURRITELLA. Shell subulately turreted, rather thick, whorls fifteen in number, spirally striated, depressedly excavated round the upper part, two-keeled in the middle, keels narrow, elevated, sharp, distant, with the interstices between them excavated, aperture small; white.

Hab. —?

The form of this species is remarkably characterized by its very prominent bicarinate structure, resembling the thread of a large flattened press-screw. The specimen here represented from Mr. Cuming's collection is the only one I have seen; and he possesses no information as to its locality. It exhibits no indication of colour.

Species 30. (Mus. Cuming.)

TURRITELLA RADULA. *Turr. testā lanceolato-acuminatā, gracili, anfractibus ad duobus et viginti, supra infrāque biliaris, medio concavis, liris, nisi infimā, granoso-crenatis, anfractu ultimo laevibus septiformibus interdum imbricato, granulis subsoletis; violascente-ashida, rufofuscente flammatū et variegatū.*

THE GRATER TURRITELLA. Shell lanceolately acuminate, slender, whorls about two and twenty in number, double-ridged above and below, concave in the middle, ridges, excepting the lowest, granosely crenated, last whorl sometimes imbricated with septiform lamellae, granules rather obsolete; violet-white, flamed and variegated with reddish-brown.

KIENER, Icon. coq. viv. p. 13. pl. 2. f. 1.

Hab. Isle of Muerte, Bay of Guayaquil (found at the depth of eleven fathoms in sandy mud); Cuming.

The form of the whorls of this elongated lanceolate species is very similar to that of *T. exoleta*; the middle is however less concave and the upper and lower ribs are represented each by two smaller ridges which are granosely crenulated.

Species 31. (Mus. Metcalfe.)

TURRITELLA HASTULA. *Turr. testā acutē lanceolata, gracili, anfractibus duobus vel tribus et viginti, convexis, concentricè plicato-corrugatis, anterioribus planatis, basi tumidiscalis, rugis levidis, anfractibus omnibus spiraliter inciso-striatis, liris intermediis obsoletē granulatis; cinereo-albicans, purpureo-brunneu, arcuatim strigatā et variegatā.*

THE LITTLE LANCE TURRITELLA. Shell sharply lanceolate, slender, whorls two or three and twenty in number, rather convex, concentrically plicately wrinkled, the latter flattened, rather swollen at the base, the wrinkles faded away, all the whorls spirally impressly striated, the intermediate ridges being obsoletely granulated; ash-white, arcuately streaked and variegated with purple-brown.

Hab. —?

A very slender sharply lanceolate shell, fully distinguished by the peculiarities of sculpture above described from any other species of this monograph.

Species 32. (Mus. Cuming.)

TURRITELLA FERRUGINEA. *Turr. testā subelongato-turrīdā, ad basim angulatā et subcarinatā, anfractibus se-decim, convexis, spiraliter subtiliter liratis, primis bicanalis, carinis gradatim evanidis, lirarum intersticiis striis minutis longitudinalibus sub lete cerebrinæ decussatis; albida, basi versus ferrugineo-castaneo tincta.*

THE RUSTY TURRITELLA. Shell somewhat elongately turreted, angulated and slightly keeled at the base, whorls sixteen in number, convex, spirally finely ridged, the first whorls two-keeled, keels gradually fading, interstices between the ridges very closely decussated beneath the lens with minute longitudinal striae; whitish, stained towards the base with rusty chestnut.

Hab. —?

The whorls of this species have a peculiar angular carinate structure at the base, in the place of the suture.

Species 33. (Mus. Belcher.)

TURRITELLA MACULATA. *Turr. testā acuminato-turrīdā, basi concavā et nitide striatā, anfractibus octodecim spira-*

siter acutè elevato-striatis, superiòr excavatis, deinde bicostatis, costis tumidinascens, interstitio concavo, anfractibus primis parum costatis; allidà, aut fulvescente, striis spiraliis purpureo-fusco tinctis, anfractibus infra suturas purpureo-fusco maculatis et interdum linearibus, basi violaceo.

THE SPOTTED TURRITELLA. Shell acuminate turreted, concave and neatly striated at the base, whorls eighteen in number, spirally sharply elevately striated, excavated round the upper part, then two-ribbed, ribs rather swollen, with the interstices concave, the first whorls but slightly ribbed; whitish or fulvous, spiral striae stained with purple-brown, whorls spotted beneath the sutures and sometimes lineated with the same colour, base violet.

Hab. China Seas; Belcher.

Although this species approximates so closely to the *T. bicingulata* in general appearance, it will be found to differ on examination. In the earlier whorls of *T. bicingulata* the keels are very prominently developed even to the apex; in the species under consideration they are comparatively obsolete, the base of the shell is more concave, and is marked by numerous coloured linear striae, which are continued throughout the shell. There is none of the bold blotchy painting which characterizes the *T. bicingulata*, but the upper margin of the whorls is always distinguished by a row of irregular red-brown spots immediately beneath the sutures.

Species 34. (Mus. Hanley.)

TURRITELLA META. *Turr. testá pyramidalí-conicā, crassiusculā, anfractibus quindecim, plano-declinib. basi tumidinascens, spiraliter exeribris, anfractua diuidio superiori eximia subirregulariter lirato, liris subobsoletè granosis, fulvescente-albidè, purpureo-rufo maculatè et variegatâ, liris striisque purpureo-rufo alboque nitide articulatâ.*

THE CONICAL PILLAR TURRITELLA. Shell pyramidal conical, rather thick, whorls fifteen in number, flatly slanting, rather swollen at the base, spirally closely striated, upper half of the whorls delicately and somewhat irregularly ridged, ridges somewhat obsoletely granose; fulvous-white, blotched and variegated with purple-red, ridges and striae neatly articulated with purple-red and white.

Hab. —?

Unacquainted with the *T. nebula* of M. Kiener, I have seen somewhat inclined to assign the present species to it, save that it is a stouter shell, the ridges appear on the upper half of the whorls only, and he makes no mention of their very delicate articulated painting.

Species 35. (Mus. Metcalfe.)

TURRITELLA CORNEA. *Turr. testá acuminato-turrítâ, interdum crassiusculâ, ad basin obtuse angulatâ, anfractibus quindecim, spiraliter tricarinatis et striatis, anfractibus ultimi, et interdum penultimi, carinis evanidioribus, lirâ pareâ in interstitiis; ferrugineo-castaneâ, aperturae parte purpureo tinctâ.*

THE HORNY TURRITELLA. Shell acuminate-turreted, sometimes rather thick, obtusely angled at the base, whorls fifteen in number, spirally three-keeled and striated, keels of the last whorl, and sometimes of the last but one, fainter, having a small ridge in the interstices; rusty-chestnut, interior of the aperture tinged with purple.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. ix. p. 255.

Turbo terebra, Linnaeus, (of Fauna Suecia, not of Syst. Nat.)

Territella ungulina, Deshayes, (not *Turbo ungulinus*, Linna.)

Territella Linnæi, Deshayes.

Turritella communis, Risso.

Hab. Seas of Europe.

Described by Linnaeus in his 'Fauna Suecia' under the name *T. terebra*, which has been adopted by Montagu and other British authors, without considering that the name had already been appropriated in the 'Systema Naturae' to a foreign species of much larger dimensions. Lamarck described it under the name given above, but from a worn and very obscure specimen, and with so brief a description, and no information as to its locality, that until M. Kiener, who held the keys of Lamarck's collection, published the *T. cornuta*, it was impossible correctly to identify the species. M. Deshayes not having obtained access either to Linnaeus's collection for the type of *Turbo ungulinus*, or to Lamarck's for that of *Territella cornuta*, assigned the species under consideration, agreeably with the views of Dr. Pulteney, to the former of these, which is Lamarck's *T. fuscata*; and did not observe it to be identical with the *T. terebra* of the 'Fauna Suecia,' which he proposes to call *T. Linnæi*.

The figure to which Lamarck refers in the 'Encyclopédie Méthodique' in illustration of his *T. cornuta*, is the Mediterranean form of the species, which is more elongated and has the sutures more excavated than that found on our own coast. The accompanying figure represents a specimen of the British state of the species dredged at Torbay by Mr. Metcalfe; and Mr. Hanley possesses a transparent white variety, with the keels more prominently developed, dredged off the Shetland Isles at a considerable depth by Mr. M'Andrew. Professor E. Forbes informs me, however, that this variety is not uncommon, and occurs in shallows as well as deeps.



28. *a*



28. *b*



28. *c*



29. *a*



29. *b*



— 2 —

TURRITELLA.

PLATE VIII.

Species 36. (Mus. Cumming.)

TURRITELLA HANLEYANA. *Turr. testá acuminato-pyramidalis, ad basim angulatá et plano-concavá, aufractibus quatuordecim, plano-declivibus, spiraleriter quadriliratis et striatis, basi marginatis; sordidé lacteá, maculis nigricantibus indistinctis obliquè tincta.*

HANLEY'S TURRITELLA. Shell acuminate pyramidal, angulated and flatly concave at the base, whorls fourteen in number, flatly slanting, spirally four-ringed and striated, margined round the base; dull cream-colour, obliquely stained with indistinct blackish spots.

Hab. ——?

An extremely characteristic species both in respect of form and sculpture. Mr. Metcalfe also possesses a specimen.

Species 37. (Mus. Metcalfe.)

TURRITELLA CLATHRATA. *Turr. testá lanceolato-acuminata, angustá, solidiusculá, aufractibus octadecim, plurimis levibus, dividens inferiori fortílier bicarinatis; fulvo-costata, aurantio-frisco obscurè maculata, carinis pallidioribus.*

THE CROSS-BARRED TURRITELLA. Shell lanceolately acuminate, narrow, rather solid, whorls eighteen in number, flattened, smooth, strongly two-keeled upon the lower half; fulvous chestnut, obscurely spotted with orange-brown, keels paler.

KIENER, Icon. coq. viv. p. 38. pl. 14. f. 1.

Hab. ——?

Distinguished by its very narrow acuminate growth, and smooth bold pair of keels.

Species 38. (Mus. Banks. Soc. Linn.)

TURRITELLA CANDIDA. *Turr. testó acuminato-turrítá, aufractibus octadecim, spiraleriter striatis, superne bicarinatis; medio concavis, deinde unicarinatis, aufractu ultimo et penultimo quadricarinatis, suturis excavatis; niveá, immaculata.*

THE WHITE TURRITELLA. Shell acuminate turreted, whorls eighteen in number, spirally striated, two-keeled round the upper part, concave in the middle, then encircled with a single keel, last whorl,

and last but one, four-keeled, sutures excavated; snowy white, unspotted.

Hab. ——?

A delicate semi-transparent white species of which I have seen two very characteristic specimens; one in Mr. Cumming's collection and the other in the Banksian cabinet of the Linnaean Society. Amongst the shells collected by Sir Joseph Banks, preserved in this Museum, are many interesting species.

Species 39. (Fig. a, Mus. Cumming. Fig. b, Mus. Walton.)

TURRITELLA PUNCTATA. *Turr. testá subpyramidalis-ocuminata, crassiusculá, basi fortiter costata, aufractibus quindecim, spiraleriter striatis, carinatis et concavis, infernè tuvidis, carinis granatis; albida, purpureo-fusco profuse punctata, maculata, et variegata.*

THE DOTTED TURRITELLA. Shell somewhat pyramidaly acuminate, rather thick, strongly ribbed at the base, whorls fifteen in number, spirally striated, keeled and concave, swollen round the lower part, keels grained; whitish, profusely dotted, blotched, and variegated with purple-brown.

KIENER, Icon. coq. viv., p. 19. pl. xi. f. 3.

Hab. ——?

The keels are somewhat irregular in this species, but it is constantly distinguished by a slight swelling out of the base of each whorl. The upper part of the whorl is always more or less concave with a smaller keel or ridge in the concavity.

Species 40. (Mus. Cumming.)

TURRITELLA SPECTRUM. *Turr. elongato-turrítá, tenuiculá, aufractibus octodécim, rotundatis, primis sex-carinatis, deinde duodecim-carinatis, carinis evanescib; gradatim subtilioribus; semipellucido-olivaceo, aufractu parte superiori fuscavite tincta.*

THE SPECTRE TURRITELLA. Shell elongately turreted, rather thin, whorls eighteen in number, rounded, the first six-keeled, then twelve-keeled, keels delicate, gradually finer; semitransparent white, upper part of the whorls tinged with brown.

Hab. ——?

The gradual doubling of the number of keels in this species is curious. The first ten whorls are encircled by six keels, an intermediate keel then gradually appears in each of the interstices, increasing in size until they obtain the same degree of development as the original keels.

Species 41. (Mus. Cuming.)

TURRITELLA ROSEA. *Turr. testá pyramidali-conicā, basi plano-angulatā, anfractibus quindecim, planatis, spirali ter quinque-liratis, liris inaequidistantibus, striis elevatisculis in interstitiis; albidā, basin versus ferrugineo-fusco, liris striisque saturatiōribus, apertura fūce livido-purpurascēte tinctā.*

THE ROSE TURRITELLA. Shell pyramidal conical, flatly angled at the base, whorls fifteen in number, flattened, spirally six-ridged, ridges inequidistant, with

rather elevated striae in the interstices; whitish, rusty brown towards the base, lines and striae darker brown, interior of the aperture tinged with livid purple.

QUOY and GAIMARD, Zool. Voy. de l'Astrolabe, vol. iii. p. 136, pl. 55, f. 24—26.

Au cedem? *Turritella lineolata*, Kiener.
Hab. New Zealand.

The truly conical form of this species is very characteristic, and it is neatly encircled throughout with brown lines, consisting of ridges and intermediate striae at unequal distances. I can discover little of the rose-colouring observed by M. Quoy; and I incline to suspect the rarity of this has led M. Kiener to describe the yellower and lighter variety of the species under another name, *T. lineolata*.



43. a



44.



b



44.



c



d

TURRITELLA.

PLATE IX.

Species 42. (Mus. Brit.)

TURRITELLA TASMANICA. *Turr. testá angusto-acuminatá, solidiusculá, anfractibus quindecim, planiusculis, ad marginem inferioriem liratis, liris dubiis striisque intermediiis supra cingulatis, interstictiis striis minutis elevatis obliqui creberimè decessatis; sordide albida.*

THE TASMANIAN TURRITELLA. Shell narrowly acuminate, rather solid, whorls fifteen in number, rather flattened, ridged at the lower edge, encircled with two ridges and intermediate striae above, interstices very closely decessated with minute oblique raised striae; dead white.

Hab. Van Dieman's Land; Dr. Sinclair.

The only specimen of this, in the British Museum, is in bad condition and colourless, but affords strong characters for the foundation of the species. It approximates nearest to the *T. clathrata* in which the spiral ridges are more strongly and conspicuously developed, and the interstices smooth.

Species 43. (Fig. *a* and *b*, Mus. Cuming.)

TURRITELLA TRIPLOCATA. *Turr. testá acuminato-turritá, crassiussentia, anfractibus quindecim, spiraliiter striatis, et tricostatis, costis planiusculis, medianá multo maiore, superiori parvá, duplicitá, apicem versus interdum obsoletá; nunc albida, castaneo variegatá et flammatá, nunc castaneá, unicolor.*

THE THREE-PLAITED TURRITELLA. Shell acuminate turreted, rather thick, whorls fifteen in number, spirally striated, and three-ribbed, ribs flattened, the middle rib much the largest, the upper one small, duplicate, and sometimes obsolete towards the apex; whitish, variegated and flamed with chestnut, sometimes entirely chestnut.

STUDER, Philippi, Enum. Moll. Sicil. p. 190.

Hab. Mediterranean and West Coast of Africa.

The specimens here figured of this species are from the mouth of the Gambia; in the Mediterranean it is of somewhat smaller dimensions and of lighter growth.

Species 44. (Mus. Brit.)

TURRITELLA NIVEA. *Turr. testá elongato-turritá, crassi-usculá, anfractibus viginti, superne contractis, deinde tumidiusculis, transversis septen-costatis, costis interstitiisque striatis, aequalibus, striis minutis incrementi creberimè decessatis, anfractibus perpacuis primis bicarinatis, carinis cito evanidis; nivea.*

THE SNOW-WHITE TURRITELLA. Shell elongately turreted, rather thick, whorls twenty in number, contracted round the upper part, then rather swollen, transversely seven-ribbed, ribs and interstices striated, equal, very closely decessated with minute lines of increase, first few whorls two-keeled, keels quickly disappearing; snowy-white.

GRAY, MSS. British Museum.

Hab. East Coast of Africa; Capt. Owen, R.N.

A genuine snow-white species, rather closely encircled with neat ridges of equal width with the interstices.

Species 45. (Mus. Brit.)

TURRITELLA GUNNII. *Turr. testá elongato-acuminatá, solidiusculá, angusti, anfractibus octodecim, plano-concavis, marginibus tumidiusculis, medio obscurè et subirregulariter liratis et striatis, suturis excavatis; albida, flammis undulatis pallidè fulvo-centre-fuscis obliquè picta, basi pallide rosá.*

GUNN'S TURRITELLA. Shell elongately acuminate, rather solid, narrow, whorls eighteen in number, flatly concave, margins rather swollen, obscurely and somewhat irregularly ridged and striated in the middle, sutures excavated; whitish, obliquely painted with light fulvous-brown waved flames, base pinkish.

Hab. Van Dieman's Land; Gunn.

An interesting new species collected by Mr. Ronald Gunn, an eminent naturalist of Hobart Town. It is a narrow, sharply acuminate shell, with the whorls flatly concave, and margined at the edges so as to give an excavated appearance to the sutures.

Species 46. (Mus. Belcher.)

TURRITELLA AQUILA. *Turr. testá pyramidali-turritá, anfractibus quatuordecim aut quindecim, superne concavo-declicibus, deinde subobscurè latè bicosostatis, undique con-*

TURRITELLA.—PLATE IX.

*spiculæ inciso-striatis, anfractu ultimo infra nitide tirato
et striato; ustulato-fuscescente, rufo-castaneo oblique
maculata vel strigata.*

THE SWARTHY TURRITELLA. Shell pyramidal turreted, whorls fourteen to fifteen in number, concavely slanting round the upper part, then somewhat obscurely broadly two-ribbed, conspicuously engraved throughout with grooved striae, last whorl neatly

ridged and striated beneath; swarthy brown, obliquely spotted and streaked with reddish chestnut. ADAMS and REEVE, Moll. Voy. Samarang.
Hab. Nangasaki Bay, Japan.

The painting of this species has rather a dotted appearance from its being every where cut through by the striate grooves.



47



46



48



49



50



51



52



53



55



54

TURRITELLA.

PLATE X.

Species 16. (Mus. Banks).

TURRITELLA CONSTRICTA. *Turr. testá turritá, sulcatis profundis, anfractibus duodecim ad quattuordecim, bicarinatis, carinis remoto distantibus, intersticio concavo; sordida albá.*

THE TIED TURRITELLA. Shell turreted, sutures deep, whorls twelve to fourteen in number, two-keeled, keels widely distant, the interstices being concave; dead white.

Hab. —?

Three specimens of this species, not hitherto described, are preserved in the Banksian collection of the Linnaean Society.

Species 47. (Mus. Hanley.)

TURRITELLA FASCIALIS. *Turr. testá luscolato-acuminata, gravillima, anfractibus octodecim, concavis, exiliter quadrangularibus, sulcatis subcontractis; intescente, anfractuum parte superiori rubido-fuscata.*

THE BANDED TURRITELLA. Shell lanceolately acuminate, very slender, whorls eighteen in number, convex, faintly four-ridged, sutures a little contracted; yellowish, upper part of the whorls banded with dull red.

MENKE, —?

Hab. Bay of Montija, Central America (in coarse sand at the depth of seven fathoms); Cuming.

Distinguished by its slender erect form and ruddy band beneath the sutures.

Species 48. (Mus. Belcher.)

TURRITELLA FASTIGIATA. *Terr. testá graciliuso-subulata, tenacula, anfractibus octodecim ad viginti, superne contractis, declivibus, deinde rotundatis, uniques subtilissime levatis et striatis, apicem versus bicarinatis; rigilaceo alboque pallide variegatis, striis fuscovenustibus oblongis, levis obscure fuscō punctatis vel articulatis.*

THE MINARET TURRITELLA. Shell very slenderly subulate, rather thin, whorls eighteen to twenty in number, contracted round the upper part, slanting, then rounded, very finely ridged and striated, two-keeled towards the apex: variegated with pale violet

and white, and oblique brownish streaks, ridges dotted or articulated with light brown.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas; Belcher.

The oblique clouded streaks of painting are characteristic in this species.

Species 49. (Mus. Cuming.)

TURRITELLA FROSA. *Turr. testá turritá, anfractibus decem ad duodecim, subplano-concavis, spiritaliter quadrangularibus, sulcis angustis, subprofundis; fuscescens, infra suturas albivante.*

THE ERODED TURRITELLA. Shell turreted, whorls ten to twelve in number, rather flatly convex, spirally four-grooved, grooves narrow, rather deep; light brown, whitish beneath the sutures.

COUTHOUY, Bost. Journ. Nat. Hist., vol. ii. p. 103. pl. 3. f. 1.

Hab. Massachusetts Bay, United States (found in stomachs of fishes); Gould, De Kay.

Generally more or less eroded, the apex being semi-transparent and somewhat papillary.

Species 50. (Mus. Belcher.)

TURRITELLA MONILIFERA. *Turr. testá acuminato-pyramidalis, basi depresso-concavā, et aculeo angulatō, anfractibus quindecim, carinā latissimā infra suturas medio concavā, deinde bi-liratis; rosaceo-albā, carinā punctis distantibus obliquis rufescente-fuscis ornatā.*

THE NECKLACE TURRITELLA. Shell acuminately pyramidal, depresso concave and sharply angled at the base, whorls fifteen in number, with a broad keel beneath the sutures, concave in the middle, then two-ridged; pinkish white, keel ornamented with distant oblique red-brown dots.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas; Belcher. *

The only painting in this species, beyond its delicate violet-pink hue, consists in the necklace-like row of red-brown dots upon the keel.

Species 51. (Mus. Beleher.)

TURRITELLA OPALINA. *Turr. testá subventricoso-turritá, tenuílata, anfractibus duodecim, rotundatis, supernè depresso-caualiculatis, sub lente minutissimè creberrimè iaciso-striatis; pellucido-albá, fuscescente pallidè con-centricaè flammati.*

THE OPAL TURRITELLA. Shell somewhat ventricosely turreted, rather thin, whorls twelve in number, rounded, depressedly channelled round the upper part, beneath the lens very closely minutely engraved with striae; opal-white, faintly concentrically flamed with light brown.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas; Belcher.

An extremely interesting species, quite unlike any other.

Species 52. (Mus. Beleher.)

TURRITELLA DECLIVIS. *Turr. testá pyramidalis-turritá, basi plano-angulata, anfractibus octodecim, plano-de-clivibus, basin versus gradatim latioribus, undique cre-berrimè subtilissimè undulato-striatis, prope apicem media plicatis; lutescente-albá, livido-fusco tincta et apicem versus peculiariter maculata.*

THE SLANTING TURRITELLA. Shell pyramidaly turreted, flatly angled at the base, whorls eighteen in number, flatly slanting, gradually wider towards the base, very closely and finely undulately striated, plicated in the middle towards the apex; yellowish-white, stained with livid brown, and peculiarly spotted towards the apex.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas; Belcher.

In this very interesting species, the livid brown appears in the first few whorls in a necklace of spots beneath the sutures.

Fig. 53. (Mus. Beleher.)

For description of this species, see Genus *Eglisia*, Sp. 3.

Species 54. (Mus. Belcher.)

TURRITELLA MULTILIRATA. *Turr. testá acuminato-turritá, anfractibus supernè contractis, inférâ leviter declivibus, inferiore caualiculatis, spiraliter multiliratis, liris sub-tiliissimè granulatis; pellucido-albá.*

THE MANY-RIDGED TURRITELLA. Shell acuminately turreted, whorls contracted round the upper part, gently slanting below, channelled round the lower part, spirally many-ridged, ridges very finely granulated; transparent white.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas; Belcher.

An extremely delicate transparent shell.

Species 55. (Mus. Belcher.)

TURRITELLA CONSPERSA. *Turr. testó turritá, anfractibus duodecim, supernè declivibus, deinde tuwidis et con-spicie bicanalatis et liratis; lutescente-albá, fuscescente longitudinaliter undulati et punctatá.*

THE SPRINKLED TURRITELLA. Shell turreted, whorls twelve in number, slanting round the upper part, then swollen and conspicuously two-keeled and ridged; yellowish white, longitudinally waved and dotted with light brown.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas; Belcher.

Delicately mottled throughout with irregular, light brown waved streaks, forming dots here and there next the sutures.



TURRITELLA.

PLATE XI.

Species 56. (Mus. Belcher.)

TURRITELLA BICOLOR. *Turr. testā acuminato-turritā, aufractibus decea ad duodecim, convexis, subtilissimè quadriliratis et striatis, liris distantibus obscurè granulatis; aureo-lutea, sutaris lirisque nigrescente-purpureis.*

THE TWO-COLOURED TURRITELLA. Shell acuminate turreted, whorls ten to twelve in number, convex, very finely four-ridged and striated, ridges distant, obscurely granulated; rich yellow, sutures and ridges blackish purple.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas.

In addition to the above characters there are a few pockered obliquely wrinkled striae next the sutures.

Species 57. (Mus. Belcher.)

TURRITELLA CANALICULATA. *Turr. testā acuminato-turritā, aufractibus duodecim, spiraliiter acutē sex-liratis et striatis, interstictis striis obliquis cancellatis, liris tenuibus, duabus inferioribus prominentibus; sordidè alba.*

THE CHANNELLED TURRITELLA. Shell acuminate turreted, whorls twelve in number, spirally sharply six-ridged and striated, interstices cancellated with oblique striae, ridges thin, the two lower ones rather prominent; dull white.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas.

The two lower ridges, more prominent than the rest, form a spiral channel immediately above the suture.

Species 58. (Mus. Belcher.)

TURRITELLA VITTULATA. *Turr. testā acuminato-turritā, basi subconcavā, aufractibus duodecim, ad suturas contractis, creberrimè spiraliiter striatis, striis elevatis, subirregularibus, aufractibus perpansis primis bicarinatis; fuscescente, striis interrupè castaneis.*

THE FINELY-FILLETED TURRITELLA. Shell acuminate turreted, slightly concave at the base, whorls twelve in number, contracted at the sutures, very closely spirally striated, striae raised, somewhat irregular, first few whorls two-keeled; light-brown, striae stained with chestnut here and there in an interrupted manner.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas.

The striae of this species are not articulated with chestnut, but merely coloured in an irregularly interrupted manner.

Species 59. (Mus. Belcher.)

TURRITELLA CONGELATA. *Turr. testā acutē subulata, basi angulata, aufractibus sedecim, convexo-planis, liris, obscurè triliratis, liris tenuibus, distantibus, pellucido-alba.*

THE CONGEALED TURRITELLA. Shell sharply subulate, angled at the base, whorls sixteen in number, convexly flattened, smooth, obscurely three-ridged, ridges fine, distant; transparent white.

ADAMS and REEVE, Moll. Voy. Samarang.
Hab. China Seas.

Transparent at the base, but more opake towards the apex.

Species 60. (Mus. Brit.)

TURRITELLA PAGODA. *Turr. testā subpyramidalis-turritā, aufractibus quatuordecim, spiraliiter acutē liratis, liris versus conspicuè acutē unicostatis, aufractibus perpansis priuis bicaratis, costæ superiori cito evanida; albida, fuligineo obscurè flammata.*

THE PAGODE TURRITELLA. Shell somewhat pyramidal turreted, whorls fourteen to fifteen in number, spirally sharply ridged, conspicuously encircled with a single sharp rib towards the base, first few whorls two-ribbed, the upper one quickly disappearing; whitish, obscurely flamed with light fulvous colour.

Hab. New Zealand; Dr. Sinclair.

Distinctly characterized by its single spiral keel and elegant form; yet intimately allied with the double-keeled *T. conspersa*.

Species 61. (Mus. Brit.)

TURRITELLA HOOKERI. *Turr. testā acuminato-turritā, tenuiculā, aufractibus quindecim, medio concurvo-turoidis, carinis duabus acutis, rotatis, undique conspicue cingulatis, aufracta ultimo et penultimo subtiliter elevato-striatis; pellucido-alba.*

HOOKE'S TURRITELLA. Shell acuminate turreted, rather thin, whorls fifteen in number, concavely swollen in the middle, and conspicuously encircled throughout with two sharp remote keels, last and penultimate whorls finely elevately striated; transparent-white.

Hab. ——?

An extremely interesting species, collected during the Antarctic Expedition by Capt. Sir James Clark Ross, which I have the pleasure to name in honour of Dr. Joseph Hooker, the eminent naturalist of that productive voyage, and author of the 'Flora Antarctica.'

Species 62. (Mus. Cuming.)

TURRITELLA SINUATA. *Turr. testá pyramidalí-acuminatá, anfractibus quatuordecim, planulatis, spiraliter tenui-liratis et striatis, aperturá parviusculá, labro ampliter sinuato: fulvescente, zonulá albida rufo-punctatá infra suturas.*

(I) **SINUATED TURRITELLA.** Shell pyramidaly acuminate, whorls fourteen in number, flattened, spirally finely ridged and striated, aperture rather small, lip largely sinuated; light fulvous colour, having a red-dotted white zone beneath the sutures.

Hab. ——?

Readily distinguished by its dotted painting, and by the broad sinus of the outer lip.

Species 63. (Mus. Cuming.)

TURRITELLA RUBESCENS. *Turr. testá acuminato-turrítá, anfractibus tredecim, spiraliter conformati liratis et striatis, primis paucis medio carinatis, cariná cito evanidá: rubescens.*

(II) **LIGHT-RED TURRITELLA.** Shell acuminate turreted,

whorls thirteen in number, spirally closely ridged and striated, the first few whorls keeled in the middle, keel quickly vanishing; light-red.

Hab. Bay of Montijo, West Columbia (in coarse sand at the depth of seven fathoms); Cuming.

All the whorls are somewhat prominent in the middle, occupied by the keel in the first few whorls.

Species 64. (Mus. Cuming.)

TURRITELLA CINGULIFERA. *Turr. testá gracili, turritá, anfractibus ad duodecim, superne contractis, deinde rotundatis, spiraliter elevato-striatis; albá, infra suturas fuscescere.*

THE ENCIRCLED TURRITELLA. Shell slenderly turreted, whorls about twelve in number, contracted round the upper part, then rounded, spirally elevately striated; white, light brown beneath the sutures.

SOWERBY, Tankerville Cat. App. p. 14.

Turritella fragilis, Kiener.

Hab. Port Essington; Jukes.

The light brown colouring immediately beneath the sutures is a prominent character of this species.

Species 65. (Mus. Cuming.)

TURRITELLA INCISA. *Turr. testá subpyramidalí-acuminatá, basi concavo-angulata, anfractibus duodecim, plano-convervis, spiraliter undique creberimè inciso-striatis: cinereo-fusca.*

THE ENGRAVED TURRITELLA. Shell somewhat pyramidaly acuminate, concavely angled at the base, whorls twelve in number, flatly convex, spirally very closely marked with engraved striae; ash-brown.

Hab. Sidney (in deep water); Strange.

Closely but irregularly engraved throughout.

TURRITELLA.

	Plate.	Species.		Plate.	Species.
annulata, Kiener	V.	18	Hanleyana, Reeve	VIII.	36
aquilus, Reeve	IX.	46	hastula, Reeve	VII.	31
Archimedis, Mawe	I.	3	Hookeri, Reeve	XI.	61
attenuata, Reeve	I.	4	imbricata, Lam.	V.	19
baeumii, Kiener	II.	7	imbricata (<i>Turbo</i>), Linn.	V.	19
Banksii, Gray	IV.	15	inceisa, Reeve	XI.	63
biangulata, De Blaive	V.	20	lentigiosa, Reeve	III.	9
bicingulata, Lam.	V.	20	leucostoma, Valenc.	II.	5
bicolor, Adams & Reeve	XI.	56	Liger, Deshayes	VI.	24
brevialis, Lam.	V.	16	lineolata, Kiener	VIII.	41
Broderipiana, D'Orb.	II.	6	linnei, Deshayes	VII.	35
canaliculata, Adams & Reeve	XI.	57	maculata, Reeve	VII.	33
candida, Reeve	VIII.	38	marmorata, Kiener	II.	6
carinifera, Lao.	IV.	12	meta, Reeve	VII.	34
cerea, Reeve	VI.	25	monilifera, Adams & Reeve	X.	50
cingulata, Sowerby	VL	23	multilirata, Adams & Reeve	X.	54
cingulifera, Sowerby	XI.	62	nivea, Gray	IX.	44
clathrata, Kiener	VIII.	37	nodulosa, King	IV.	11
cochlea, Reeve	VII.	29	opalina, Adams & Reeve	X.	51
columnaris, Kiener	IV.	14	pagodina, Reeve	XI.	60
communis, Risso	VII.	35	papillosa, Kiener	IV.	11
congelata, Adams & Reeve	XI.	59	punctata, Kiener	VIII.	39
conspersa, Adams & Reeve	X.	55	radula, Kiener	VII.	30
constricta, Reeve	X.	16	rosea, Quoy	VIII.	41
cornea, Lamarck	VII.	35	rubescens, Reeve	XI.	61
crocea, Kiener	VI.	26	sanguinea, Reeve	VI.	27
Cumingii, Reeve	IV.	13	sinnata, Reeve	XI.	60
deglivis, Adams & Reeve	X.	52	spectrum, Reeve	VIII.	40
duplicata (<i>Turbo</i>), Linn.	I.	2	Tasmanica, Reeve	IX.	42
erosa, Couthouy	X.	49	terebra (<i>Turbo</i>), Linn.	I.	3
exoleta (<i>Turbo</i>), Linn.	VI.	22	tigrina, Kiener	III.	8
fascialis, Menke	X.	47	torulosa, Kiener	V.	21
fastigiata, Adams & Reeve	X.	48	tricarinata, King	VI.	23
ferruginea, Reeve	VII.	32	triplicata, Studer	IX.	43
flamminata, Kiener	VI.	24	trisulata, Lam.	V.	17
fragilis, Kiener	XI.	62	ungulina, Deshayes	VII.	35
fuscata, Lam.	I.	1.	ungulina (<i>Turbo</i>), Linn.	I.	1
gemmata, Reeve	VI.	28	varia, Kiener	V.	16
goniostoma, Valenc.	III.	10	variegata (<i>Turbo</i>), Linn.	V.	19
Gunnii, Reeve	IX.	45	vittulata, Adams & Reeve	XI.	58

MONOGRAPHS
OF THE GENERA
MESALIA AND EGLISIA.

There is a rapture on the lonely shore,
There is society where none intrudes,
By the deep sea, and music in its roar.—*Byron*.



2. a



2. b



Eglisia. Pl. I.



2. c



2. d



2. e

MESALIA.

PLATE I.

Genus MESALIA, Gray.

Testa acuminato-turrata, basi rotundata, columellā plano-contortā, recedente, aperture marginē infra columellam leviter sinuata et reflexa.

Shell acuminate turreted, rounded at the base, columella flatly twisted, receding, margin of the aperture below the columella sinuated and reflected.

The flattened receding structure of the columella of the well-known Senegal *Turritella* (*T. brevialis*, Lamarck), *Le Mesal* of Adanson, induced Mr. Gray to found a genus for its reception with the name *Mesalia*, which it is desirable should be adopted.

It is materially distinguished in this respect from the rest of the *Turritellae*, excepting the little Greenland *T. lactea* of Möller, also a *Mesalia*; and to these may be added a species (*M. Melanoides*) from the Cummingian collection, in which the typical character of the genus is yet more strongly developed, and which Mr. Cumming had already put aside in his cabinet as a new form.

Species 1. (Mus. Brit.)

MESALIA LACTEA. *Mes. testā turrīta, anfractibus tredecim, longitudinatiliter sub-oblique et obèse plicato-costatis, lineis impressis inter costas, lirā angustā inferne funiculatis, apertura parvī; opaco-albā.*

THE MILK-WHITE MESALIA. Shell turreted, whorls thirteen in number, longitudinally rather obliquely and stoutly plicately ribbed, with impressed lines in the interstices, corded round the lower part with a narrow ridge, aperture small; opaque-white.

Turritella lactea, Möller, Index Moll. Greenlandie, p. 9.
Hab. Greenland.

Möller describes this shell as being a diaphanous white, but the specimens in the British Museum are rather opaque.

Species 2. (Fig. 2 a, b, Mus. Hanley; and Pl. V. Fig. 16 a, b, of Turritella, Mus. Cuming.)

MESALIA BREVIALIS. *Mes. testā subabbreviato-turrīta, crassā, ponderosā, anfractibus quindecim, rotundatis, supernā obcurè canaliculatis, spiraliter quinqüeliratis, liris superioribus fortioribus, inferioribus interdum obsoletis, intersticiis undique subtilissime et creberissimè striatis, columellā plano-contortā, recedente, marginē*

subreflexo; albā, vel cinereo-albā, vel sordidè olivaceā, interdum strigis fuscescensibus indistinctè pallide tinctā.

THE SHORT MESALIA. Shell somewhat abbreviate turreted, thick, ponderous, whorls fifteen in number, rounded, obscurely channelled round the upper part, spirally five-ridged, upper ridges stronger, lower sometimes obsolete, interstices very finely and closely striated throughout, columella flatly twisted, receding and a little reflected at the edge; white, or ashy-white, or dirty olive, sometimes faintly stained with brownish streaks.

Turritella brevialis, Lamarck, Anim. sans vert. (Deshayes' edit.) vol. ix. p. 256.

Le Mesal, Adanson.

Turritella Mesal, Deshayes.

Mesalia sinuata, Gray.

Eadem var. *Turritella varia*, Kiener.

Hab. West Coast of Africa.

An extremely variable species in size and general aspect, but always distinctly characterized throughout by its one idea of sculpture. Each whorl is encircled by nine thread-like ridges, five of which remain exposed to view; of these the two uppermost are more prominently developed than the rest and form a slight channel. The interstices are sculptured throughout with exceedingly fine, close-set, sharply defined striae. In some specimens the shell is of a dark dirty olive colour and the ridges lighter, others are of a light stone colour without any distinction of tone in the ridges. Of the smaller varieties one is of a dark lead colour, another snowy white and more elongated, with the ridges almost obsolete; and a third equally white with the ridges like so many fine opaque threads.

Lamarck, when first describing this species according to the present system of nomenclature, apparently from a worn specimen, did not recognise in it *Le Mesal* of Adanson (*Voyage en Scincégal*, p. 159. pl. 10. f. 7); and M. Deshayes was unable to identify *Turritella brevialis* in his edition of the 'Animaux sans vertèbres' for want of access to the Lamarckian collection, reserved by M. Kiener for the use of his 'Iconographie des coquilles vivantes'; in which work the identification of *T. brevialis* with Adanson's *Mesal* has been recently made known.

Unacquainted with Lamarck's *T. brevialis*, from his not referring to Adanson's figure, M. Deshayes described *Le Mesal* anew. It is much to be regretted that the learned editor of the 'Anim. sans vert.' was not allowed

access to the Lamarekian types, when engaged on his important edition of that work; either whilst the collection was in the possession of the Due de Rivoli, under the curatorship of M. Kiener, or subsequently in that of Baron Delessert under the care of M. Cléén. That such was the case M. Deshayes has related to me more than once with much concern.

Species 3. (Mus. Cuming.)

MESALIA MELANOIDES. *Mes. testā acuminato-turrītā, basi rotundatā, aufractib⁹ decem, supernē leviter con-*

cavis, obliquè longitudinaliter plicato-costellatis, striis spiralibus numerosis lirisque subdistantibus decussatis, suturis impressis; albā, punctis varis rufo-fuscis ornatā.

THE MELANIA-LIKE MESALIA. Shell acuminate turreted, rounded at the base, whorls ten in number, slightly concave at the upper part, obliquely longitudinally plicately ribbed, decussated with numerous spiral striae and distant ridges, sutures impressed; white, ornamented with one or two red-brown dots.
Hab. — ?

The generic peculiarities of *Mesalia* are well-illustrated in this species.

EGLISIA.

PLATE I.

Genus EGLISIA, Gray.

Testa elongato-turrata, anfractibus numerosis, suturis depresso-serratis, columellā plano-increassatā, ūsi quadrate-angularatā, aperturā parē, rotundatā, marginibus ferè conjunctis, non reflexis.

Shell elongately turreted, whorls numerous, sutures depressed, columella flatly thickened, squarely angled at the base, aperture small, rounded, margins almost joined, not reflected.

The necessity for instituting a genus for the reception of the *Turritella spirata* of the Tankerville Catalogue, was first made known by Mr. Sowerby in his description of that remarkable shell. "Two circumstances," says the author of the Appendix, Tank. Cat. p.xiv., "are remarkable in this shell, the broad flat depression in the upper part of the volutions, and the straight edge of the outer lip. Aware that it is generically distinguished by this latter circumstance from *Turritella*, we should not have added it to this genus, had there been any more convenient place for it."

Mr. Gray proposed the name *Eglisia*, and I am able to confirm the value of the genus by the addition of two more equally characteristic species, one collected by Mr. Cuming at the Philippine Islands, another dredged by Sir Edward Belcher in the China Seas.

Species 1. (Mus. Cuming.)

EGLISIA SPIRATA. *Egl. testā pyramidalī-turritatā, anfractibus ad duodecim, superē peculiariter plano-depressis, ad suturas concavo-excavatis, spiraliter subobtuse liratis; albida; fulvescente-fusca; obliquè inequinatā et variegatā.*

THE SPIRAL EGLISIA. Shell pyramidalily turreted, whorls about twelve in number, peculiarly flatly depressed round the upper part, concavely excavated at the sutures, spirally somewhat obtusely ridged; whitish, obliquely stained and variegated with fulvous brown. *Turritella spirata*, Sowerby, Tankerville Catalogue, App. p. xiv.

Turbo suturalis, Gray (in Wood Index Test.)

Eglisia suturalis, Gray (in Brit. Mus.)

Turritella suturalis, Kiener.

Hab. Island of St. Thomas, West Indies; Sowerby.

I am not aware of any other specimens of this remarkable species, than those spoken of by Mr. Sowerby in the Tankerville Catalogue, as having been purchased in the old collection of Mr. Humphrey. Of these the one here figured is in the Cumingian collection, and another is in the British Museum.

Species 2. (Mus. Cuming.)

EGLISIA LANCEOLATA. *Egl. testā lanceolato-turritatā, anfractibus ad octodecim, rotundatis, subobscure octoliratis, interstices sub lente tenuissimè caeculatī, suturis profundè excavatis, aperturā parē, rotundā; fulvescente-spadicēa.*

THE LANCEOLATE EGLISIA. Shell lanceolately turreted, whorls about eighteen in number, rounded, rather obscurely eight-ridged, interstices, beneath the lens, very finely caeculated, sutures deeply excavated, aperture small, rounded; light fulvous bay.

Hab. Pasacao, Island of Luzon (in sandy mud at the depth of ten fathoms); Cuming.

An interesting species collected by Mr. Cuming at the above-named locality, of the same generic type as the preceding, the sutures having the same excavated structure, though in an inferior degree.

Species 3. (Mus. Belcher.)

EGLISIA TRICARINATA. *Egl. testā acutē turritatā, anfractibus ad octodecim, superē contractis et declivibus, deinde tricarinatis, sub lente longitudinaliter creberius et tenuissime caeculato-striatis, aperturā parē, rotundā; sordide cinereo-fusca.*

THE THREE-KEELED EGLISIA. Shell sharply turreted, whorls about eighteen in number, contracted and slanting round the upper part, then three-keeled, beneath the lens very finely longitudinally cancellately striated, aperture small, round; dull ashy-brown.

ADAMS and REEVE, Moll. Voy. Samarang.

Hab. China Seas; Belcher.

The cancellated structure of this species is similar to that of the preceding.

MONOGRAPH
OF THE GENUS
CASSIDARIA.

Not lost the time in sea-side ramble spent ;
Braced is the frame, and mental health is gained ;
Knowledge is gained of Him who made the deep,
And blissful love acquired of Nature's works.
Each rock-pool has its treasure, every tide
Strews on the yellow sand from Ocean's lap.—*Leedsborough.*



3 a



2



3 b

CASSIDARIA.

PLATE I.

GENUS CASSIDARIA, Lamarck.

Testa ovoidea, ventricosa, ad basin attenuata, in canalem curvum, postice ascendente, desinens, apertura longitudinali, labro columellarí laxe, latissimè effuso, labro externo reflexo, intus crenato. Operculum cornuum.

Shell ovoid, ventriose, attenuated at the base, ending in a posteriorly ascending recurved canal, aperture longitudinal, columellar lip smooth, very widely effused, outer lip reflected, crenated within. Operculum horny.

The genus *Cassidaria* is one of the very few to which no species have been added since the time of Lamarck. It is, indeed, reduced within narrower limits; for whilst, on the one hand, the researches of Cuming, Quoy, Béleher, D'Orbigny and others, compassing the four quarters of the globe, have failed to discover any new species; the Lamarckian genus has been reduced, on the other hand, by the distribution of two to form the genus *Oviscia*, and of another to the genus *Triton*. Three recent species only remain to the genus, and it is a matter of opinion amongst authors whether two of these, both inhabiting the shores of the Mediterranean, are not varieties of the same specific type.

The *Cassidaria* differs from *Cassis*, to which it is most nearly allied, by its oval inflated growth, projecting spire, and effused aperture, ending in a slightly recurved canal.

The locality of the third species is not known.

Species 1. (Mus. Cuming.)

CASSIDARIA TYRRHENIA. *Cass. testá subpyriformi-ovalá, superne ventricosá, anfractibus liris regularibus undique crebre fuscatis, ultimo superne angulato, ad angulum compressò nodoso, supra exiliter oblique plicato, apertura elongato-oblongá, labris obsolete crenatis; fulvescente-spadicé, apertura albá.*

THE TUSCAN CASSIDARIA. Shell somewhat pyramidal ovate, ventriose round the upper part, whorls closely corded throughout with regular ridges, last whorl angled at the upper part, compressly noduled at the angle, faintly obliquely plicated above, aperture elongately oblong, lips obsolete crenulated; light fulvous bay, aperture white.

LAMARCK, Ainsi, sans vert. (Deshayes' edit.) vol. vii. p. 8.
Buccinum Tyrrhenum, Chemnitz.

Cassidea Tyrrhenia, Bruguière.

Buccinum echinophorum, var. Gmelin.

Buccinum ochroleucum, Gmelin.

Cassidaria echinophora, Deshayes.

Hab. Mediterranean.

Dr. Philippi and M. Deshayes are of opinion that this shell, of which there is an accurate representation in Chemnitz 'Conchylien Cabinet,' vol. x. p. 153. f. 1461-2, is a smooth variety of the well-known *C. echinophora* which follows. It is remarkably distinguished by its angular pyriform growth, and I have not seen any specimen which may be said to represent a state intermediate between this and the commoner form, of which specimens with few or no nodules are frequently mistaken for it.

Species 2. (Fig. a and b, Mus. Cuming.)

CASSIDARIA ECHINOPHORA. *Cass. testá oviformi, vel globoso-ovatá, transversim medio costatá, supra et infra tenuiratá, costis planulatis, plus minusve nodosis, liris parvæ intermediarí, labro externo subincassato, intus crenato; fulvescente-albá, ferrugineo-fusco plus minusve tinctuá.*

THE PRICKLY CASSIDARIA. Shell egg-shaped or globosely ovate, transversely ribbed in the middle, finely ridged above and below, ribs rather flattened, more or less nodose, with a small intermediate ridge, outer lip somewhat thickened, crenated within; light fulvous white, more or less stained with rusty brown.

LAMARCK, Animi. sans vert. (Deshayes' edit.) vol. x. p. 7.

Buccinum echinophorum, Linnaeus.

Cassidea echinophora, Bruguière.

Echinora tuberculosa, Schumacher.

Buccinum strigosam, Gmelin.

Buccinum nodosum, Dillwyn.

Cassidaria Tyrrhenia, var., Philippi.

Hab. Mediterranean.

The large noduled specimen represented at Fig. 2 b, sufficiently shows that the growth of this species is extremely variable; still, I have not observed in any variety the peculiar angulated pear-shape which distinguishes the *C. Tyrrhenia*.

Species 3. (Mus. Brit.)

CASSIDARIA STRIATA. *Cass. testá subpyriformi-ovalá,*

August, 1849.

CASSIDARIA.—PLATE I.

basin versus præcipue attenuatâ, vix recurvâ, transversim creberrimè elevato-striatâ, striis longitudinibus incisis versus apicem decussatis, spiræ saturis excavatis, labris incrassatis, intus fortiter crenato-liratis, parum effusis; albâ, interduu fulcescente, supra columellam pallidâ fulcescente areolatâ.

THE STRIATED CASSIDARIA. Shell somewhat pyriformly ovate, particularly attenuated towards the base, scarcely recurved, transversely very closely elevated striated, decussated towards the apex with longitu-

dinal engraved striae, sutures of the spire excavated, lips thickened, strongly crenately ridged within, but little effused: white, sometimes brownish, marked with faint brownish squares above the columella,

LAMARCK, *Anim. sans vert.* (Deshayes' edit.) vol. x. p. 10.
Hab. —?

An extremely rare shell, concerning the habitat of which I am unable to procure any information.

MONOGRAPH
OF THE GENUS
ONISCI A.

Under the salt waves bright and clear,
On moss and sand and soft sea-weed,
Lies many a gift of ocean rare.—*Freiligrath.*



ONIS CIA.

PLATE I.

Genus ONIS CIA, Sowerby.

Testa oblongo-ovalata, basi emarginata, sub-recurea, spirâ brevi, unuc acuminatâ, unuc depresso, aperturâ elongatâ, angustâ, usque ad spiram extensâ, labro columellarî latissimâ dilatato-reflexo et granulato, labro externo denticulato, incassato, plerunque medio subeortato.

Shell oblong-ovate, emarginated and a little recurved at the base, spire sometimes acuminated, sometimes depressed, aperture elongated, narrow, extending as far as the spire, columellar lip very broadly dilately reflexed and granulated, outer lip denticulated, thickened, generally slightly contracted in the middle.

The genus *Oniscia* founded by Mr. Sowerby in his "Genera of Shells" on the Linnean *Strombus oniscus* (*Cassidaria oniscus*, Lamarek) has been universally adopted, as representing a very characteristic and well-defined assemblage of characters. They were first remarked on the appearance of the well-known *Oniscia cancellata* of the China Seas, and their importance has been fully confirmed by the discovery of several interesting new species.

The *Onisciae* are closely related to the *Cassidariae*, but are uniformly of smaller size, of more solid growth, with more elaborate sculpture, and remarkably distinguished by a wide-spread dilation of the columellar. The outer lip is also peculiar in being denticulated within.

Of the localities known of the *Onisciae* three are from the Western, two from the Eastern world.

Species 1. (Mus. Cuming.)

ONIS CIA ONISCUS. *Onisc. testâ oblongo-ovalâ, crassâ, apice unuc obtusâ, unuc mucronatâ, costis tribus vel quatuor tuberculatis cinctâ, tuberculis interdum bipartitis, labro incassato, tenuido, lîris interruptis irregularibus intus denticulato; albida, maculis perpuscuis penitus numerosis carneo-nigris aut fuscis asperâ, labris interdum albis, interdum eximie rosaceis.*

THE WOOD-LOUSE ONIS CIA. Shell oblong-ovate, thick, sometimes obtuse, sometimes pointed at the apex, encircled with three or four tuberculated ribs, tubercles sometimes divided into two, lip thickened, swollen, denticulated within with irregular interrupted ridges; whitish, sprinkled with a few spots and numerous dots of brown or blue-black, lips sometimes white, sometimes delicate pink.

SOWERBY, Genera of Shells, *Oniscia*, f. 4.

Strombus oniscus, Linnaeus.

Eadem var. *Strombus Lamarcii*, Deshayes.
Hab. Jamaica, Gosse.

The shell selected for illustration is the pink-lip variety to which M. Deshayes has given the name *O. Lamarcii*. The white varieties are usually of smaller dimensions, but the characters are in all respects the same.

Species 2. (Mus. Cuming.)

ONIS CIA STROMBIIFORMIS. *Onisc. testâ subtrigono-pyramidalis, spirâ elevatâ, apice mucronatâ, anfractibus superne angulatis, costis numerosis angustis nodosis cinctis, labro columellarî minutè granulato, labro externo tirato-denticulato, lîris brevi interruptâ interveniente: albâ, rufo-fusco maculatâ et punctatâ, labris albis.*

THE STROMBUS-SHAPED ONIS CIA. Shell somewhat triangularly pear-shaped, spire elevated, pointed at the apex, whorls angulated round the upper part, encircled with many narrow nodulous ridges, columellar lip minutely granulated, outer lip denticulated with ridges, with a short interrupted ridge in the interstices; white, spotted and dotted with red brown, lips white.

REEVE, Pro Zool. Soc., 1842. p. 91. Conch. Syst. vol. ii. p. 210. pl. 253. f. 1.

Hab. Honduras; Dyson.

Distinguished from the preceding species by its slighter growth, more elevated spire, and finely noduled narrow ribs.

Species 3. (Mus. Dennison.)

ONIS CIA EXQUISITA. *Onisc. testâ subtrigono-ovalâ, spirâ brevi, acentâ, anfractibus supernâ concavo-depressis, tuberculis papillaribus undique cingulatis, lîris obscurâ interveniente, labro columellarî latè expanso, granulis valde irregularibus, labro externo incassatius reflexo, lîris brevibus dentiformibus irregulariter munito; albida, anrantio-fusco hic illic sparsius punctatâ et pone labrum triangulatâ, labris pallide purpureo-rosaceis, apice reflo.*

THE EXQUISITE ONIS CIA. Shell somewhat triangularly ovate, spire short, sharp, whorls concavely depressed,

encircled throughout with papillary tubercles, with an obscure ridge intervening; columellar lip broadly expanded, granules very irregular, outer lip thickly reflected, furnished with short dentiform ridges; whitish, sparingly dotted, and, behind the lip, three-spotted with orange-brown, lips light purple-pink, apex red.

ADAMS and REEVE, Zool. Voy. Samarang, Moll. pl. 5, fig. 3, a and b.

Hab. Sooloo Archipelago (outside a coral reef near the city of Sooloo, in sandy mud at the depth of from sixteen to twenty fathoms); Belcher.

The surface of this very characteristic species, collected by Captain Sir Edward Belcher during the voyage of the Samarang, is covered with papillary tubercles in rows of about ten, the same number longitudinally as transversely. The columellar and outer lips are delicately enamelled of a livid purplish pink colour.

Species 4. (Mus. Cuming.)

ONISCA CANCELLOTA. *Onisc. testa oblongo-ovata, basi attenuata, plus minusve pyriformi, spirò elevatiscula, aenà, anfractibus superne rotundatis, longitudinaliter compressè plicatis, costis numerosis angustis crenulatis transversis cancellatis, costis ad juncturas squamatis, labro columellarì fortiter granulato, externo fortiter dentato; albida aut fuscescente-albida, fusco trifasciata.*

THE CANCELLED ONISCA. Shell oblong-ovate, attenuated at the base, more or less pear-shaped, spire rather elevated, sharp, whorls rounded at the upper part, longitudinally compressly plicated, cancellated transversely with numerous narrow crenulated ribs, ribs squamate at the crossing, columellar lip strongly granulated, outer lip strongly toothed; whitish or brownish white, encircled with three brown bands.

SOWERBY, Genera of Shells, *Oniscia*, f. 1 to 3.

Cassidaria cancellata, Kiener (not of Lamarck).

Hab. China Seas.

According to the testimony of M. Deshayes, the fossil *Cassidaria cancellata* of Lamarck is a different species from this, with which it has been usually identified.

Species 5. (Fig. a and b, Mus. Cuming.)

ONISCA TUBERCULOSA. *Onisc. testa cylindraceo-oblonga, crassa, spirò nixa concezà, nunc depresso-planà, apice aenè mncronotata, anfractibus quinquecostatis, costis subdistantibus, tuberculatis, interstitiis subtiliter liratis,*

labro columellarì exiliter granulato, externo medio contractato, intus denticulato; intensè nigra, irregulärer albipunctatà, aperturæ fauce abù, interdum viciè croceâ.

THE TUBERCLED ONISCA. Shell cylindrically oblong, thick, spire sometimes convex, sometimes depressedly flattened, pointed at the apex, whorls five-ribbed, ribs rather distant, tuberculated, interstices finely ridged, columellar lip faintly granulated, outer lip contracted in the middle, denticulated within; deep black, dotted irregularly with white, interior of the aperture white, sometimes bright saffron.

SOWERBY, Genera of Shells, *Oniscia*, p. 2. Reeve, Conch. Syst. vol. ii. p. 211, pl. 253, f. 2 to 4.

Hab. Fig. a. Galapagos Island (in clefts of rocks at low water); Cuming. Fig. 2 b, Gulf of California.

Of the two specimens here figured from Mr. Cuming's collection, that represented at Fig. 5 a is but indistinctly granulated, the spire is depressedly flattened, and the aperture is of bright saffron colour. In the specimen represented at Fig. 5 b, the spire is convexly raised, and there is no indication of colour in the aperture. The granules are numerous, but extremely superficial.

Species 6. (Mus. Dennison.)

ONISCA DENNISONI. *Onisc. testa trigono-ovata, spirò acuta, anfractibus superne angulatis, longitudinaliter compressè plicatis, costis numerosis angustis cancellatis, costis ad juncturas squamatis, labro columellarì latè expanso, minutè granulato, externo liris subirregularibus denticulato; albida, fusco subobscurè tri-quadrifasciata, labro columellarì intensè sanguineo-rutilo, granulis albidis, labro externo rufescente, liris, albida.*

DENNISON'S ONISCA. Shell triangularly ovate, spire sharp, whorls angled round the upper part, longitudinally compressly plicated, cancellated with numerous narrow ribs, which are squamate at the crossing, columellar lip broadly expanded, minutely granulated, outer lip denticulated with somewhat irregular ridges; whitish, rather obscurely three- or four-banded, columellar lip very deep blood-red, granules whitish, outer lip light red, with the ridges whitish.

REEVE, Pro. Zool. Soc., 1842, p. 21. Conch. Syst. vol. ii. p. 211, pl. 253, f. 5 and 6.

Hab. —?

This species approaches very closely in form and detail of sculpture to the *O. cancellata*, but is especially remarkable on account of the intensely red colouring of the columella.

MONOGRAPH

OF THE GENUS

E B U R N A .

Now is it pleasant in the summer eve,
When a broad shore retiring waters leave,
Awhile to wait upon the firm fair sand,
When all is calm at sea, all still on land,
And there the ocean's produce to explore.—*Crabbé.*



E B U R N A.

PLATE I.

Genus EBURNA, Lamarck.

Testa ovata vel ovato-conoidea, umbilicata, umbilico ampio, infra caudiculato, interdum impletio, spiræ suturis nunc simplicibus, nunc profundiæ canaliculatis, aperturâ basi emarginatâ, plerunque superne canaliculatâ, tubo columellarî incrassato, calloso, exteriori simplici. Operculum parvum, acuminatum.

Shell ovate or ovately conoid, umbilicated, umbilicus large, channelled beneath, sometimes filled up, sutures of the spire simple or very deeply channelled, columellar lip thickened, callous, outer lip simple. Operculum small, acuminated.

The name *Eburna*, signifying *ivory*, is not very appropriate for the present genus, inasmuch as it was founded on the polished *E. glabrata* of Lamarck, which proves to be an *Ancillaria*, almost entirely enveloped by the soft parts, whilst the species that remain are distinguished for having a dark thick epidermis. The animal, of which drawings have been published in the Mollusca of the Voyage of the Saumurang and of Le Bonite, approaches so near to *Eucrinium*, that M. Deshayes is disposed to return rather to the views of Bruguière and Linnaeus; it will, however, be readily acknowledged that there is a marked association of generic character in the objects under consideration. They are all largely umbilicated, the umbilicus being sometimes filled up, and are devoid of sculpture. The aperture is alike in all, and the colouring and pattern are peculiar, very similar throughout, yet varying in a manner sufficiently constant in each species to be regarded of considerable specific importance.

It is perhaps scarcely necessary to refer to M. Kiener's remarks that *E. papillosa* and *Valeantiana* are varieties of *E. spirata*, and that *E. ambulacrum* is a variety of *E. lutosa* (figured as *E. spirata* var.). This is a grave mistake; the author is not, unfortunately, acquainted with the species. It will also be found on reference to Sp. 4, that the *E. australis* is not identical with Lamarck's *Cancellaria spirata* as generally supposed; whilst *E. plumbea* is referred to the genus *Monoceros*.

The *Eburnæ* are all inhabitants of the Eastern hemisphere.

Species 1. (Mus. Brit.)

EBURNA PAPILLARIS. *Eburna. testâ oblongo-conoideâ, imperforatâ, spirâ elevatâ, apice papillari, suturis sim-*

plicibus, anfractibus subrotundatis, superne obtuse depresso, vix angulatis; albâ, fuso undique punctata.

THE PAPILLARY EBURNA. Shell oblong-conoid, imperforated, spire raised, papillary at the apex, sutures simple, whorls somewhat rounded, obtusely depressed at the upper part, scarcely angulated; white, dotted throughout with brown.

SOWERBY, Tankerville Catalogue, App. p. xxii.
Hab. Cape of Good Hope.

I am only aware of four specimens of this very interesting species; one in the British Museum, one in the Cumingian collection, one in the collection of Mr. Gaskoin, and one in some continental cabinet.

Species 2. (Mus. Cuming.)

EBURNA LUTOSA. *Eburna. testâ ovaliâ, vix umbilicata, spirâ elevata, acutâ, anfractibus concavis, tævibus, superne obtuse depressa, vix angulatis; albâ, pallidè spadiceo prouincia nebuloso-nudato-nebulatâ.*

THE MUDDY EBURNA. Shell ovate, scarcely umbilicated, spire raised, sharp, whorls convex, smooth, obtusely depressed above, scarcely angulated; white, promiscuously clouded in a waved manner with light fawn colour.

LAMARCK, Ainsi. sans vert. (Deshayes' edit.) vol. x. p. 235.
Eburna Pacifica, Swainson.

Hab. China.

The colouring obtains a less defined pattern in this species than in any other, and the whorls are more obscurely depressed rounded at the upper part.

Species 3. (Mus. Cuming.)

EBURNA JAPONICA. *Eburna. testâ ovato-conoideâ, umbilicata, apice subpapillari, suturis simplicibus, anfractibus concavis, tævibus, superne vix depresso; albida, maculis semilunaris vel oblongis fulvo-spadiceis, confertis, biseriatim cinctâ, intersitis maculis parvis numerosis. diagonaliiter dispositis, notatis.*

THE JAPAN EBURNA. Shell ovately conoid, umbilicated, somewhat papillary at the apex, sutures simple, whorls convex, smooth, very slightly depressed at the upper part; whitish, encircled with two rows of close-set semilunar or oblong fulvous bay spots, the interspaces between which are marked with numerous small spots ranging diagonally.

REEVE, Proc. Zool. Soc. 1842, p. 200 : Conch. Syst. pl. 271. f. 1.

Hab. Japan; Dr. Siebold.

Allied in form to *E. Zeylanica*, but less umbilicated, and less sharply acuminate; whilst the design of the painting is very characteristic.

Species 4. (Mus. Cuming.)

EBURNA AUSTRALIS. *Eburn. testá ovalá, imperforatá, spirá brevisealá, suturis laté depresso-canaliculatis, basi leviter subcavata, siccis quinque vel sex, deinde salvo unico subprofundo, supra punctato-striatá, anfractibus quinque; latescente-albidá, maculis grandibus rufi-fuscis juxta suturas, perplicatis minoribus iufra.*

THE SOUTHERN EBURNA. Shell ovate, imperforated, spire rather short, sutures broadly depresso channelled, sculptured at the base with five or six light grooves, then with a single rather deep one, above striated with fine punctures, whorls five in number; yellowish white, with a row of red-brown spots next the sutures, and a very few smaller ones beneath.

SOWERBY, Conch. Illustrations, f. 5.

Hab. Australia.

The *Eburna australis* has been confounded by many authors with the *Cancellaria spirata* of Lamarck, believing them to be identical. M. Deshayes goes so far as to say that Mr. Sowerby has represented the same shell in two places (Conch. Illus. *Cancellaria*, f. 25 and *Eburna*, f. 5.), suiting the configuration of the columella in each instance to the character of the genus. Such is not the case. The three-plaited *Cancellaria spirata* and the *Eburna australis* are two very distinct shells, the latter being of more solid growth, smooth towards the apex, and furnished with an arched callous columella, characteristic of the genus to which it has been referred in the absence of any more suitable place.

Species 5. (Mus. Cuming.)

EBURNA AMBULACRUM. *Eburn. testá ocatá, umbilicatá, spira suturis profunde canaliculatis, anfractibus subplanó-convexis, lœvibus; albidá, fuscescente-spadiceo maculatá, creberimē maculatá et nebulae-lineatá, apice nigritá.*

THE GALLERY EBURNA. Shell ovate, umbilicated, sutures of the spire deeply channelled, whorls rather flatly convex, smooth; whitish, very closely spotted and clouded with brownish bay, apex blackish.

SOWERBY, Tankerville Catalogue, App. p. xxii.

Hab. Isle of Mindanao, Philippines (in sandy mud at the depth of twelve fathoms); Cuming.

Very closely allied to the *E. spirata*, but always readily distinguished by its smaller size, darker colour, and more closely clouded spots.

Species 6. (Mus. Cuming.)

EBURNA AREOLATA. *Eburn. testá ovato-ventricosá, spirá elevatá, profunde et ampliter umbilicatá, convexis, lœvibus, superne obtusè depressis, subconicalatis, viz angulatis; albá, maculis rufo-castaneis grandibus oblongo-quadratis subundatis seriatim ciuctá.*

THE SQUARE-SPOOTTED EBURNA. Shell ovate ventricose, spire elevated, deeply and largely umbilicated, convex, smooth, obtusely depressed round the upper part, slightly channelled, scarcely angulated; white, encircled with rows of large oblong-square slightly waved reddish-chestnut spots.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 235.
Buccinum spiratum var., Bruguière.

Eburna tessellata, Swainson.

Hab. Ceylon.

The spots are disposed in three rows, the lowest being concealed in all but the last whorl, by the superposition of one whorl upon the other.

Species 7. (Mus. Cuming.)

EBURNA SPIRATA. *Eburn. testá ovalá, parum umbilicatá, spirá acutá, suturis profunde canaliculatis, anfractibus subplanó-convexis; albá, fuscescente-spadiceo maculatá, maculis medianis minoribus, apice cérulo-nigricante.*

THE SPIRAL EBURNA. Shell ovate, but little umbilicated, spire sharp, sutures deeply channelled, whorls rather flatly convex; white, spotted with fulvous bay, middle spots the smaller, apex bluish black.

LAMARCK, Anim. sans vert. (Deshayes' edit.) vol. x. p. 234.
Buccinum spiratum, Linnaeus.

Hab. Ceylon. Philippines.

An interesting variety of this well-known species was collected during the voyage of the Samarang, in which the spots were smaller and of much darker colour.

Species 8. (Mus. Cuming.)

EBURNA ZEYLANICA. *Eburn. testá ovato-conicá, profunde et ampliter umbilicatá, spirá acuminatá, suturis non quam canaliculatis, anfractibus convexo-declicibus, lœvatis; albá, maculis grandibus sinnatis fulvo-spadiceis*

EBURNA.—PLATE I.

*spira suturas, umbilis ovatis minoribus medio picta,
umbilico violente, apice nigricante.*

THE CEYLON EBURNA. Shell ovately conic, deeply and largely umbilicated, spire acuminate, sutures never channelled, whorls convexly slanting, smooth : white, painted beneath the sutures with large sinuated fulvous-bay blotches, in the middle with smaller oval spots of the same colour, umbilicus tinged with violet, apex blackish.

LAMARCK, *Anni. sans vrt.* (Deshayes' edit.) vol. x. p. 233.
Buccinum Zeylanicum, Bruguière.

Hab. Ceylon.

Distinguished by its regular conical growth, without the sutures being in any degree canaliculated.

Species 9. (Mus. Cuming.)

EBURNA VALENTIANA. *Ebura testa globosa, crassusculta,
imperforata, spirá peculiariter demersa, suturis pro-
funde canaliculatis, apice acuto, aufrictibus rotundatis,
loribus : albá, maculis fulvo-spadicis, promiscue longi-
tudinaliter sinuatis et undatis, picta.*

VALENTIA'S EBURNA. Shell globose, rather thick, imperforated, spire peculiarly sunk, sutures deeply channelled, apex sharp, whorls rounded, smooth : white, painted with fulvous-bay spots, promiscuously longitudinally situated and waved.

SWAINSON, *Zoological Illustrations*, vol. iii. pl. 144.
Hab. Red Sea.

A very distinct species, named in honour of Lord Valentia, afterwards Earl of Mount Norris.

1681

