

Crystal Data: Monoclinic. *Point Group:* 2/m. As prismatic crystals to 1 cm in bunches, sheaf-like or radial aggregates, typically in crusts.

Physical Properties: *Cleavage:* Imperfect. *Tenacity:* Brittle. *Fracture:* Uneven. Hardness = ~3.5 D(meas.) = n.d. D(calc.) = 3.915

Optical Properties: Transparent. *Color:* Colorless, pale green, pale yellow, light blue, pale lilac or pink. *Streak:* White. *Luster:* Vitreous.

Optical Class: Biaxial (-). $\alpha = 1.719(3)$ $\beta = \gamma = 1.732(3)$ $2V(\text{meas.}) = 15(10)^\circ$ $2V(\text{calc.}) = 0^\circ$
Dispersion: Medium, $r > v$. *Orientation:* $Y = b$.

Cell Data: *Space Group:* C2/c. $a = 11.8405(3)$ $b = 12.7836(2)$ $c = 6.69165(16)$ $\beta = 112.425(3)^\circ$
 $Z = 4$

X-Ray Diffraction Pattern: Arsenatnaya fumarole, Tolbachik Volcano, Russia.
 2.758 (100), 2.735 (25), 3.220 (19), 2.910 (17), 3.509 (16), 1.661 (16), 2.620 (12)

| Chemistry: | (1) | (2) | | (1) | (2) |
|--------------------------------|-------|-------|--------------------------------|-------|--------|
| Na ₂ O | 7.32 | 5.61 | Fe ₂ O ₃ | 3.53 | |
| K ₂ O | 0.10 | | TiO ₂ | 0.01 | |
| CaO | 6.82 | 10.15 | SiO ₂ | 0.03 | |
| MgO | 20.31 | 21.87 | P ₂ O ₅ | 1.25 | |
| MnO | 0.68 | | V ₂ O ₅ | 0.10 | |
| CuO | 0.27 | | As ₂ O ₅ | 58.77 | 62.37 |
| ZnO | 0.02 | | SO ₃ | 0.13 | |
| Al ₂ O ₃ | 0.56 | | Total | 99.90 | 100.00 |

(1) Arsenatnaya fumarole, Tolbachik Volcano, Kamchatka, Russia; average electron microprobe analysis supplemented by Raman spectroscopy; corresponds to $(\text{Na}_{1.30}\text{K}_{0.01}\text{Ca}_{0.67}\text{Mg}_{2.78}\text{Mn}_{0.05}\text{Cu}_{0.02}\text{Al}_{0.06}\text{Fe}^{3+}_{0.24})_{\Sigma=5.13}(\text{As}_{2.83}\text{P}_{0.10}\text{S}_{0.01}\text{V}_{0.01})_{\Sigma=2.95}\text{O}_{12}$. (2) NaCaMgMg₂(AsO₄)₃.

Mineral Group: Alluaudite supergroup, alluaudite group - arsenates.

Occurrence: A sublimate at an active volcanic fumarole.

Association: Hematite, tenorite, johillerite, nickenichite, bradaczekite, hatertite, magnesiohatertite, badalovite, aphthitalite-group sulfates, langbeinite, calciolangbeinite, sanidine (As-bearing variety), fluorophlogopite, fluoborite, tilasite, anhydrite, pseudobrookite, rutile, sylvite, halite, lammerite, lammerite-β, urusovite, ericlaxmanite, kozyrevskite, arsmirandite, svabite, popovite, dmisokolovite, shchurovskyite, yurmarinite, krasheninnikovite, euchlorine, wulffite, alumoklyuchevskite, sellaite, gahnite, corundum (shallow zone); badalovite, fluorophlogopite, sanidine, diopsidite, hematite, cassiterite, anhydrite, metathénardite, belomarinaite, sylvite, tilasite, svabite, arsenowagnerite (medium zone); anhydrite, diopsidite, hematite, svabite, berzeliite, schäferite, forsterite, magnesioferrite, ludwigite, rhabdobarite-group fluoroborates, powellite, baryte, fluorapatite, udinaite, arsenudinaite, paraberzeliite (deep zone).

Distribution: From the Arsenatnaya fumarole, Second scoria cone, Northern Breakthrough of the Great Tolbachik Fissure Eruption, Tolbachik Volcano, Kamchatka, Russia.

Name: The prefix, *calcio*, identifies the calcium analogue of *johillerite*.

Type Material: A.E. Fersman Mineralogical Museum, RAS, Moscow, Russia (95621).

References: (1) Pekov, I.V., N.N. Koshlyakova, A.A. Agakhanov, N.V. Zubkova, D.I. Belakovskiy, M.F. Vigasina, A.G. Turchkova, E.G. Sidorov, and D.Y. Pushcharovsky (2021) New arsenate minerals from the Arsenatnaya fumarole, Tolbachik volcano, Kamchatka, Russia. XV. Calciojohillerite, NaCaMgMg₂(AsO₄)₃, a member of the alluaudite group. *Mineral. Mag.*, 85, 215–223.