

**Calciosamarskite****(Ca, Fe, Y)(Nb, Ta, Ti)O<sub>4</sub>**

**Crystal Data:** Orthorhombic; metamict. *Point Group:* 2/m 2/m 2/m. As anhedral masses.

**Physical Properties:** *Cleavage:* None. *Fracture:* Conchoidal. *Tenacity:* Brittle. Hardness = 5-6 D(meas.) = 5.8(1) D(calc.) = n.d. Radioactive.

**Optical Properties:** Opaque. *Color:* Black. *Streak:* Brown to black. *Luster:* Vitreous. *Optical Class:* Isotropic.  $n > 2.0$

**Cell Data:** *Space Group:* n.d.  $a = 5.632$   $b = 9.912$   $c = 5.221$   $\beta = 93.87^\circ$

**X-ray Powder Pattern:** Mitchell Co., North Carolina, USA. (After heating at 1100 °C for 12 hours) 2.952 (100), 3.079 (90), 2.599 (35), 3.867 (30), 2.824 (30), 3.614 (25), 1.839 (20)

<b>Chemistry:</b>	(1)	(1)	(1)	(1)	
UO <sub>2</sub>	17.9	Sc <sub>2</sub> O <sub>3</sub>	1.09	Er <sub>2</sub> O <sub>3</sub>	0.38
WO <sub>3</sub>	1.83	Tb <sub>2</sub> O <sub>3</sub>	0.11	Yb <sub>2</sub> O <sub>3</sub>	0.48
Nb <sub>2</sub> O <sub>5</sub>	37.4	Y <sub>2</sub> O <sub>3</sub>	6.90	Lu <sub>2</sub> O <sub>3</sub>	0.12
Ta <sub>2</sub> O <sub>5</sub>	5.57	Pr <sub>2</sub> O <sub>3</sub>	0.03	MnO	0.79
SiO <sub>2</sub>	0.21	Ce <sub>2</sub> O <sub>3</sub>	0.05	PbO	0.70
TiO <sub>2</sub>	1.17	Dy <sub>2</sub> O <sub>3</sub>	1.38	MgO	0.01
ThO <sub>2</sub>	3.75	Nd <sub>2</sub> O <sub>3</sub>	0.50	CaO	6.17
SnO <sub>2</sub>	0.10	Fe <sub>2</sub> O <sub>3</sub>	7.40	Na <sub>2</sub> O	0.84
Al <sub>2</sub> O <sub>3</sub>	0.15	Sm <sub>2</sub> O <sub>3</sub>	0.29	H <sub>2</sub> O	n.d.
Gd <sub>2</sub> O <sub>3</sub>	0.97	Ho <sub>2</sub> O <sub>3</sub>	0.21	Total	96.5

(1) Mitchell Co., North Carolina, USA; average electron microprobe analysis; corresponds to (Ca<sub>0.309</sub>Fe<sub>0.261</sub>U<sub>0.186</sub>Y<sub>0.172</sub>REE<sub>0.114</sub>Na<sub>0.077</sub>Th<sub>0.04</sub>Mn<sub>0.031</sub>)<sub>Σ=1.19</sub>(Nb<sub>0.791</sub>Ta<sub>0.071</sub>Ti<sub>0.041</sub>)<sub>Σ=0.903</sub>O<sub>4</sub>.

**Mineral Group:** Samarskite group.

**Occurrence:** An accessory mineral in RE-rich granite pegmatites, common but not abundant.

**Association:** Columbite, zircon, monazite, uraninite, aeschynite, magnetite, albite, topaz, beryl, garnet, muscovite, biotite.

**Distribution:** From Mitchell Co., North Carolina, USA. At the Woodcox mine, Hybla, Monteagle township, Hastings county, Ontario, Canada.

**Name:** Honors Colonel Vasilii Evgrafovich von Samarskii-Bykhovets (1803-1870), Chief of Staff of the Russian Corps of Mining Engineers. The prefix, *calcio*, indicates Ca is dominant in the A-site.

**Type Material:** n.d.

**References:** (1) Hanson, S.L., W.B. Simmons, A.U. Falster, E.E. Foord, and F.E. Lichte (1999) Proposed nomenclature for samarskite-group minerals: new data on ishikawaite and calciosamarskite. *Mineral. Mag.*, 63(1), 27-36. (2) Ellsworth H.V. (1928) A mineral related to samarskite from the Woodcox Mine, Hybla Ontario. *Amer. Mineral.*, 13, 63-65. (3) Palache, C., H. Berman, and C. Frondel (1944) Dana's system of mineralogy, (7th edition), v. I, 797-801, 806-807. (4) Sugitani, Y., Y. Suzuki, and K. Nagashima (1985) Polymorphism of samarskite and its relationship to other structurally related Nb-Ta oxides with the  $\alpha$ -PbO<sub>2</sub> structure. *Amer. Mineral.*, 70, 856-866.