

## Laverovite

## $K_2NaMn_7Zr_2(Si_4O_{12})_2O_2(OH)_4F$

**Crystal Data:** Triclinic. *Point Group:*  $\bar{1}$ . As an elongated zone,  $\sim 500 \mu m$  long, on the rim of a lath-like intergrowth of four minerals: laverovite, zircophyllite, astrophyllite, and kupletsksite.

**Physical Properties:** *Cleavage:* Perfect on {001}. *Tenacity:* Brittle. *Fracture:* Hackly.  
Hardness = 3 D(meas.) = n.d. D(calc.) = 3.367

**Optical Properties:** Transparent. *Color:* Pale brown to dark brown. *Streak:* Light brown.  
*Luster:* Vitreous.

*Optical Class:* Biaxial (-).  $\alpha = 1.670(2)$   $\beta = 1.710(5)$   $\gamma = 1.740(5)$   $2V(\text{meas.}) = 82(2)^\circ$   
 $2V(\text{calc.}) = 80^\circ$ . *Pleochroism:*  $X =$  yellowish brown,  $Y =$  brownish yellow,  $Z =$  pale yellow.  
*Absorption:*  $X > Y > Z$ . *Dispersion:* Strong,  $r > v$ .

**Cell Data:** *Space Group:*  $P\bar{1}$ .  $a = 5.4329(1)$   $b = 11.9232(3)$   $c = 11.7491(3)$   $\alpha = 112.905(2)^\circ$   
 $\beta = 94.696(1)^\circ$   $\gamma = 103.178(1)^\circ$   $Z = 1$

**X-ray Powder Pattern:** Desourdy quarry, Mont Saint-Hilaire, Montérégie, Québec, Canada.  
2.589 (100), 2.788 (97), 3.452 (92), 2.680 (68), 1.590 (50), 2.504 (44), 1.776 (39)

| Chemistry: | (1)   | (1)                  |        |
|------------|-------|----------------------|--------|
| $Nb_2O_5$  | 0.56  | $CaO$                | 0.48   |
| $ZrO_2$    | 9.78  | $MgO$                | 0.76   |
| $TiO_2$    | 4.69  | $Cs_2O$              | 0.05   |
| $SiO_2$    | 33.52 | $K_2O$               | 6.00   |
| $Al_2O_3$  | 0.94  | $Na_2O$              | 2.28   |
| $SrO$      | 0.13  | F                    | 1.80   |
| $ZnO$      | 0.07  | $H_2O$               | [2.57] |
| $FeO$      | 13.94 | $\underline{-O=F_2}$ | 0.76   |
| $MnO$      | 20.51 | Total                | 97.32  |

(1) Desourdy quarry, Mont Saint-Hilaire, Québec, Canada; average electron microprobe analysis supplemented by FTIR spectroscopy,  $H_2O$  calculated from structure; corresponds to  
 $(K_{1.78}Sr_{0.02}Cs_{0.01}\square_{0.19})_{\Sigma=2}(\square_{1.85}Na_{0.15})_{\Sigma=2}(Na_{0.88}Ca_{0.12})_{\Sigma=1}(Mn_{4.03}Fe^{2+}_{2.71}Mg_{0.25}Zn_{0.01})_{\Sigma=7}$   
 $(Zr_{1.11}Ti_{0.82}Nb_{0.06}Mg_{0.01})_{\Sigma=2}[(Si_{7.78}Al_{0.26})_{\Sigma=8.04}O_{24}]O_2[(OH)_{3.68}F_{0.32}]_{\Sigma=4}[\square_{1.85}(H_2O)_{0.15}]_{\Sigma=2}$ .

**Mineral Group:** Astrophyllite supergroup, kupletsksite group.

**Occurrence:** A late-stage hydrothermal mineral in a nepheline-syenite pegmatite.

**Association:** Zircophyllite, kupletsksite, astrophyllite, aegirine, analcime, orthoclase, albite.

**Distribution:** From the Desourdy quarry (which became part of the Demix and later the Poudrette quarry), Mont Saint-Hilaire, La Vallée-du-Richelieu RCM, Montérégie, Québec, Canada.

**Name:** Honors Professor Nikolay Pavlovich Laverov (1930-2016) Academician of the Russian Academy of Sciences, a prominent Russian ore geologist.

**Type Material:** RRUFF Database (R060216), University of Arizona, Tucson, USA. Acquired from the Royal Ontario Museum, Toronto, Ontario, Canada (M57542).

**References:** (1) Sokolova, E., M.C. Day, F.C. Hawthorne, A.V. Kasatkin, R.T. Downs, L. Horváth, and E. Pfenninger-Horváth (2019) Laverovite,  $K_2NaMn_7Zr_2(Si_4O_{12})_2O_2(OH)_4F$ , a new astrophyllite-supergroup mineral from Mont Saint-Hilaire, Québec, Canada. *Can. Mineral.*, 57(2), 201-213.  
(2) (2021) Amer. Mineral., 106, 1186-1187 (abs. ref. 1).