

Crystal Data: Triclinic. *Point Group:* $\bar{1}$. Lath-shaped to 150 μm ; as thin-lamellar pseudo-hexagonal rhomb-like and band-shaped crystals to 2 mm, usually curved.

Physical Properties: *Cleavage:* Perfect on {001}. *Tenacity:* Elastic flakes. *Fracture:* n.d. Hardness = n.d. D(meas.) = n.d. D(calc.) = 7.598

Optical Properties: Semi-transparent. *Color:* Silvery, brown-red inner reflections; white with slight pink hue in reflected light. *Streak:* Gray-black. *Luster:* Metallic.

Optical Class: Anisotropy: Strong.

R₁-R₂: (470) 44.1-39.1, (546) 42.8-37.6, (589) 42.2-37.6, (650) 42.4-37.3

Cell Data: *Space Group:* P $\bar{1}$. $a = 6.470(5)$ $b = 6.368(5)$ $c = 6.401(7)$ $\alpha = 105.0(1)^\circ$ $\beta = 91.59(9)^\circ$ $\gamma = 118.90(6)^\circ$ Z = 4

X-ray Powder Pattern: Kudryavy volcano, Kurile volcanic arc, Russia.

2.7834 (10), 2.764 (10), 2.733 (10), 1.6156 (10), 1.5938 (10), 2.371 (9), 2.0914 (9),

Chemistry:	(1)	(2)
Mo	1.58	0.13
Re	74.77	74.30
Fe		
Cu	0.42	
S	23.43	25.46
Total	100.20	99.89

(1) Pagoni Rachi Mo-Cu-Te-Ag-Au prospect, northern Greece; electron microprobe analysis; corresponds to Re_{1.043}Mo_{0.043}Cu_{0.017}S_{1.897}. (2) Kudryavy volcano, Kurile volcanic arc, Russia; electron microprobe analysis; corresponds to Re_{1.002}Mo_{0.005}Cu_{0.017}S_{1.993}.

Occurrence: As sublimes in high-temperature fumaroles and in porphyry-type Mo-Cu-Te-Ag-Au hydrothermal mineralization.

Association: Molybdenite, pyrite, quartz (Pagoni Rachi); magnetite, corundum, wollastonite, andradite-grossular garnet, wurtzite, greenockite, cadmoindite, halite (Kudryavy volcano).

Distribution: At the Kudryavy volcano, Kurile volcanic arc, Russia [TL] and in the Pagoni Rachi and Konos prospects, in northeastern Greece.

Name: For its chemical composition as a unique mineral of *rhenium*.

Type Material: A.E. Fersman Mineralogical Museum, Moscow, Russia.

References: (1) Korzhinsky, M.A., S.I. Tkachenko, K.I. Shmulovich, Y.A. Taran, and G.S. Steinberg (1994) Discovery of a pure rhenium mineral at Kudriavy volcano. *Nature* 1994, 369, 51-52. (2) Znamensky, V.S., M.A. Korzhinsky, G.S. Steinberg, S.I. Tkachenko, A.I. Yakushev, I.P. Laputina, I.A. Bryzgalov, N.D. Samotoin, L.O. Magazina, O.V. Kuzmina, N.I. Organova, V.A. Rassulov, and I.V. Chaplygin (2005) Rheniite, ReS₂ - natural rhenium disulfide from fumaroles of Kudriavy volcano (Iturup Island, Kurile islands). *Zap. Vses. Mineral. Obshchest.* 134(5), 32-40 (in Russian, English abstract). (4) Ibáñez-Insa, J., T. Woźniak, R. Oliva, C. Popescu, S. Hernández, and J. López-Vidrier (2021) Structural and High-Pressure Properties of Rheniite (ReS₂) and (Re,Mo)S₂. *Minerals*, 11, 207. (5) Voudouris, P.C., V. Melfos, P.G. Spry, L. Bindi, T. Kartal, K. Arikas, R. Moritz, M. Ortelli (2009) Rhenium-rich molybdenite and rheniite in the Pagoni Rachi Mo-Cu-Te-Ag-Au prospect, northern Greece: implications for the Re geochemistry of porphyry-style Cu-Mo and Mo mineralization. *Can. Mineral.*, 47(5), 1013-1036.