

**Crystal Data:** Isometric. **Point Group:**  $4/m\bar{3}2/m$ . As grains to 0.25 mm.

**Physical Properties:** **Cleavage:** None. **Fracture:** Conchoidal. **Tenacity:** Brittle.  
Hardness = 4-4.5 D(meas.) = n.d. D(calc.) = 5.275

**Optical Properties:** Translucent to transparent. **Color:** Brownish yellow to reddish orange.  
**Streak:** Light yellow. **Luster:** Adamantine.  
**Optical Class:** Isotropic.  $n = 2.10(5)$

**Cell Data:** **Space Group:**  $Fd\bar{3}m$ .  $a = 10.5053(10)$  Z = 8

**X-ray Powder Pattern:** Boziguuer REE deposit, Akesu, Xinjiang Autonomous Region, China.  
3.042 (100), 2.628 (38), 1.859 (34), 1.582 (15), 1.515 (4), 6.074 (3), 1.2045 (3)

Chemistry:	(1)		(1)
Na <sub>2</sub> O	6.80	UO <sub>2</sub>	5.81
K <sub>2</sub> O	0.01	Ta <sub>2</sub> O <sub>5</sub>	3.00
CaO	2.01	Nb <sub>2</sub> O <sub>5</sub>	53.42
MgO	0.01	F	3.19
FeO	0.05	Cl	0.02
SrO	0.03	ThO <sub>2</sub>	0.48
PbO	16.17	Sb <sub>2</sub> O <sub>5</sub>	0.01
Ce <sub>2</sub> O <sub>3</sub>	4.29	ZrO <sub>2</sub>	0.01
La <sub>2</sub> O <sub>3</sub>	1.65	MnO	0.04
Nd <sub>2</sub> O <sub>3</sub>	0.41	SnO <sub>2</sub>	0.34
Y <sub>2</sub> O <sub>3</sub>	0.42	<u>-O = (F,Cl)<sub>2</sub></u>	1.35
SiO <sub>2</sub>	0.03	Total	98.21
TiO <sub>2</sub>	1.36		

(1) Boziguuer REE deposit, Akesu, Xinjiang Autonomous Region, China; average of 10 electron microprobe analyses supplemented by FTIR spectroscopy; corresponding to  $(Na_{1.03}Pb_{0.34}Ca_{0.17}U_{0.10}Th_{0.01}Ce_{0.12}La_{0.05}Y_{0.02}Nd_{0.01})_{\Sigma=1.85}(Nb_{1.88}Ti_{0.08}Ta_{0.06}Sn_{0.01})_{\Sigma=2.03}O_{6.21}F_{0.79}$ .

**Mineral Group:** Pyrochlore supergroup.

**Occurrence:** An accessory phase in niobium-tantalum-rich alkali syenite.

**Association:** Microcline, albite, aegirine, sodic amphibole, biotite, zircon, rutile, thorite, fluorite, fluocerite-(Ce), columbite-(Fe), xenotime-(Y), astrophyllite, chevkinite-(Ce), fergusonite-(Y).

**Distribution:** From the Boziguuer REE deposit, 43 km north of Baicheng County City, Baicheng County, Akesu, Xinjiang Autonomous Region, China.

**Name:** As a member of the pyrochlore supergroup with dominant fluorine in the Y site and sodium in the A site.

**Type Material:** Geological Museum of China, Beijing, China (M12183).

**References:** (1) Yin, J., G. Li, G. Yang, X. Ge, H. Xu, and J. Wang (2015) Fluornatropyrochlore, a new pyrochlore supergroup mineral from the Boziguuer rare earth element deposit, Baicheng County, Akesu, Xinjiang, China. Can. Mineral., 53(3), 455-460. (2) (2016) Amer. Mineral., 101, 2357-2358 (abs. ref. 1).