

Morinite

NaCa₂Al₂(PO₄)₂(F, OH)₅·2H₂O

©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Monoclinic. *Point Group:* 2/m. As tabular to blocky, prismatic crystals, to 1 cm; flattened on {100}, elongated along [010]; {hk0} faces striated || [001]; subparallel aggregates of crystals terminated by $\{\bar{1}11\}$, {110} yield groups with serrated upper edges; as botryoidal crusts and radial fibrous aggregates.

Physical Properties: *Cleavage:* Perfect on {100}; imperfect on {001}. *Hardness* = 4–4.5
D(meas.) = 2.94–2.96 D(calc.) = 2.98

Optical Properties: Transparent to translucent. *Color:* Colorless to white, pale pink, wine-red. *Streak:* White. *Luster:* Vitreous to pearly on cleavages.
Optical Class: Biaxial (-). *Orientation:* Y = b; X ∧ a = 45°. *Dispersion:* r < v, weak.
α = 1.553 β = 1.565 γ = 1.567 2V(meas.) = 40°

Cell Data: *Space Group:* P2₁/m. a = 9.454(3) b = 10.692(4) c = 5.444(2)
β = 105.46(2)° Z = 2

X-ray Powder Pattern: Hugo mine, South Dakota, USA.
2.94 (10), 3.47 (8), 1.783 (8), 4.70 (7), 2.63 (7), 9.11 (6), 3.73 (6)

Chemistry:

	(1)	(2)
P ₂ O ₅	29.31	30.04
SiO ₂	2.40	
Al ₂ O ₃	22.46	22.33
MnO	0.02	
CaO	22.49	23.60
Na ₂ O	5.85	6.62
F	14.58	13.31
H ₂ O	9.80	9.98
insol.	0.14	
-O = F ₂	6.12	5.60
Total	100.93	100.28

(1) Montebbras, France. (2) Hugo mine, South Dakota, USA; corresponds to Na_{1.00}Ca_{1.97}Al_{2.05}(P_{0.99}O₄)₂[F_{3.52}(OH)_{1.22}]_{Σ=4.74}·2H₂O.

Occurrence: An uncommon late-stage mineral in complex granite pegmatites.

Association: Montebbrasite, apatite, augelite, wardite, wavellite, cassiterite.

Distribution: From Montebbras, Creuse, France. On the Greifensteine, near Ehrenfriedersdorf, Saxony, and at Hagendorf, Bavaria, Germany. In the Gunheath china clay pit, St. Austell, Cornwall, England. From the Viitaniemi pegmatite, near Eräjärvi, Finland. In the USA, in the Hugo mine, 1.5 km south of Keystone, Pennington Co., South Dakota; at the Stewart mine, Pala, San Diego Co., California; from the Silver Coin mine, near Valmy, Iron Point district, Humboldt Co., Nevada. In a pegmatite nearby Broken Hill, New South Wales, and at the Mt. Cleveland tin mine, 14 km southwest of Waratah, Tasmania, Australia.

Name: Honors Mr. E.A. Morineau, Director of the tin mine, Montebbras, France, who supplied the first specimens.

References: (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 783–784, 784–785 [ježekite = morinite]. (2) Fisher, D.J. and J.J. Runner (1958) Morinite from the Black Hills. *Amer. Mineral.*, 43, 585–594. (3) Hawthorne, F.C. (1979) The crystal structure of morinite. *Can. Mineral.*, 17, 93–102. (4) Čech, F. and P. Povondra (1985) Identity of ježekite with morinite. *Bull. Minéral.*, 108, 533–539.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.