

Crystal Data: Triclinic. **Point Group:** 1. As cores of equant, well-formed pseudocubes to 1 mm; intimately intergrown with peatite-(Y).

Physical Properties: *Cleavage:* Weak to poor on {100}, {010}, {001} probable. *Fracture:* Splintery. *Tenacity:* Brittle. Hardness = ~3 D(meas.) = n.d. D(calc.) = 3.60(1)

Optical Properties: Translucent. *Color:* Yellowish white. *Streak:* White. *Luster:* Sub-vitreous to dull.

Optical Class: Biaxial (n.d.). α = n.d. β = 1.636(1) γ = n.d. 2V(meas.) = n.d. 2V(calc.) = n.d. *Pleochroism:* None. *Dispersion:* None.

Cell Data: *Space Group:* P1. a = 10.9977(6) b = 10.9985(6) c = 10.9966(6)
 α = 90.075(4) $^\circ$ β = 89.984(4) $^\circ$ γ = 89.969(4) $^\circ$ Z = 1

X-ray Powder Pattern: Poudrette pegmatite, Mont Saint-Hilaire, Canada.
 3.89 (100), 2.94 (98), 2.59 (98), 7.80 (79), 11.04 (76), 6.36 (75), 3.48 (68)

Chemistry:	(1)	(2)	(1)	(2)
Na ₂ O	11.25	12.77	ZrO ₂	23.40
CaO	4.15		ThO ₂	0.49
La ₂ O ₃	0.11		HfO ₂	0.69
Y ₂ O ₃	16.48	23.26	P ₂ O ₅	28.10
Nd ₂ O ₃	0.08		F	0.62
Dy ₂ O ₃	1.11		-O=F ₂	0.26
Ce ₂ O ₃	0.10		CO ₂	[5.92]
Er ₂ O ₃	1.18		H ₂ O	[0.92]
Tm ₂ O ₃	0.28		<u>Li₂O</u>	[2.01]
Yb ₂ O ₃	0.57		Total	99.75
Al ₂ O ₃	0.14			100.00

(1) Poudrette pegmatite, Mont Saint-Hilaire, Canada; average of 22 electron microprobe analyses, H₂O, CO₂ and Li₂O calculated from stoichiometry and their presence confirmed by LA-ICP-MS and Raman analyses; corresponding to $\text{Li}_4(\text{Na}_{10.79}\text{Ca}_{1.21})_{\Sigma=12}(\text{Y}_{4.34}\text{Ca}_{0.99}\text{Dy}_{0.18}\text{Er}_{0.18}\text{Yb}_{0.09}\text{La}_{0.02}\text{Ce}_{0.02}\text{Nd}_{0.01})_{\Sigma=5.83}(\text{Zr}_{5.65}\text{Hf}_{0.10}\text{Th}_{0.06})_{\Sigma=5.81}[(\text{P}_{0.98}\text{Al}_{0.01})_{\Sigma=0.99}\text{O}_4]_{12}(\text{CO}_3)_4\text{O}_4[(\text{OH})_{3.03}\text{F}_{0.97}]_{\Sigma=4}$.

(2) $\text{Li}_4\text{Na}_{12}\text{Y}_6\text{Zr}_6(\text{PO}_4)_{12}(\text{CO}_3)_4\text{O}_4(\text{OH})_4$.

Occurrence: A late-stage product possibly related to the in situ alteration of the pre-existing mineral assemblage present in the core of a zoned peralkaline pegmatite dike encased in a hornfels xenolith.

Association: Peatite-(Y), albite, rhodochrosite, siderite, chabazite-Na, synchysite-(Ce), sabinaite.

Distribution: From the Poudrette pegmatite, Mont Saint-Hilaire, La Vallée-du-Richelieu, Montérégie (formerly Rouville County), Québec, Canada.

Name: Honors Robert A.J. Ramik (*b.*1951), an analytical technician and mineralogist in the Department of Natural History, Royal Ontario Museum, Toronto, Ontario, Canada.

Type Material: The Royal Ontario Museum, Toronto, Ontario, Canada (M53893).

References: (1) McDonald, A.M., M.E. Back, R.A. Gault, and L. Horváth (2013) Peatite-(Y) and ramikite-(Y), two new Na-Li-Y±Zr phosphate-carbonate minerals from the Poudrette pegmatite, Mont Saint-Hilaire, Quebec. *Can. Mineral.*, 51, 569-596. (2) (2014) Amer. Mineral., 99, 2441 (abs. ref. 1).