


Sodium-zippeite



©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Orthorhombic. *Point Group:* n.d. Rare in crystals, to 100 μm , rhomboidal to slightly elongated plates, extremely thin, may be petalike, lathlike, spindle-shaped, in vermiform aggregates; as cross-fiber veinlets, warty crusts, very fine-grained powdery to massive; commonly in efflorescences. *Twinning:* Common on $\{h0l\}$, contact twins, with another law, polysynthetic.

Physical Properties: *Cleavage:* Perfect on $\{010\}$; perhaps another perpendicular. Hardness = ~ 2 D(meas.) = > 3.3 D(calc.) = n.d. Radioactive; fluoresces bright yellow under SW and LW UV.

Optical Properties: Semitransparent. *Color:* Yellow, lemon-yellow, yellow-orange. *Optical Class:* Biaxial (-). *Pleochroism:* X = colorless to pale yellow; Z = dark yellow to golden yellow. *Orientation:* X \perp plane of flattening; Z = c. $\alpha = 1.630\text{--}1.637$ $\beta = 1.685\text{--}1.689$ $\gamma = 1.732\text{--}1.739$ 2V(meas.) = Moderate to large.

Cell Data: *Space Group:* n.d. $a = 8.82$ $b = 17.12$ $c = 7.32$ $Z = \text{n.d.}$

X-ray Powder Pattern: Synthetic.

7.34 (100), 3.663 (54), 3.490 (44), 3.153 (35), 2.858 (15), 3.754 (14), 2.118 (11)

Chemistry:

	(1)	(2)
SO ₃	10.43	10.71
UO ₃	75.86	76.53
Na ₂ O	5.25	5.53
K ₂ O	0.47	
H ₂ O	7.99	7.23
Total	[100.00]	100.00

(1) Delta mine, Utah, USA; recalculated to 100% after deduction of gypsum 0.1% and insoluble SiO₂ + Al₂O₃ + Fe₂O₃ 4.39%. (2) Na₄(UO₂)₆(SO₄)₃(OH)₁₀ · 4H₂O.

Occurrence: A secondary alteration product of uraninite and base-metal sulfides in the oxidized zone of uranium-rich deposits, typically in sandstone-type uranium deposits; of post-mine origin, where the sodium may be supplied by percolating saline water.

Association: Andersonite, uranopilite, johannite, schröckingerite, uranopilite, other “zippeite-group” minerals.

Distribution: In the USA, in Utah, from the Happy Jack mine, White Canyon, and the W.N. mine, Deer Flat, San Juan Co.; in the Delta and Lucky Strike No. 2 mines, San Rafael district, and at the Sodaroll mine, Green River district, Emery Co.; from the Parco No. 23 mine, Thompsons district, Grand Co.; at the Atomic King mine, Cane Springs, Moab Co.; in the Oyler mine, Henry Mountains district, Wayne Co.; in Arizona, from the Sue mine, Cherry Creek district, Gila Co., and in the Hillside mine, about 5.5 km north of Bagdad, Eureka district, Yavapai Co.; at the Jackpile mine, Grants district, Socorro Co., New Mexico. From Jáchymov (Joachimsthal), Czech Republic. In the Geevor mine, St. Just, Cornwall, England.

Name: For its content of *sodium* and relation to the other *zippeite* group species.

Type Material: n.d.

References: (1) Frondel, C., J. Ito, R.M. Honea, and A.M. Weeks (1976) Mineralogy of the zippeite group. *Can. Mineral.*, 14, 429–436.