

$$43. r = \frac{v_0^2}{32} (\sin 2\theta)$$

$$v_0 = 2200 \text{ ft/sec}$$

$\theta$  changes from  $10^\circ$  to  $11^\circ$

$$dr = \frac{(2200)^2}{16} (\cos 2\theta) d\theta$$

$$\theta = 10 \left( \frac{\pi}{180} \right)$$

$$d\theta = (11 - 10) \frac{\pi}{180}$$

$$\Delta r \approx dr$$

$$= \frac{(2200)^2}{16} \cos \left( \frac{20\pi}{180} \right) \left( \frac{\pi}{180} \right) \approx 4961 \text{ feet}$$

$$\approx 4961 \text{ feet}$$

$$47. \text{ Let } f(x) = \sqrt[4]{x}, x = 625, dx = -1.$$

$$f(x + \Delta x) \approx f(x) + f'(x) dx = \sqrt[4]{x} + \frac{1}{4^4 \sqrt{x^3}} dx$$

$$f(x + \Delta x) = \sqrt[4]{624} \approx \sqrt[4]{625} + \frac{1}{4(\sqrt[4]{625})^3} (-1)$$

$$= 5 - \frac{1}{500} = 4.998$$

Using a calculator,  $\sqrt[4]{624} \approx 4.9980$ .

51. In general, when  $\Delta x \rightarrow 0$ ,  $dy$  approaches  $\Delta y$ .

53. True

$$45. \text{ Let } f(x) = \sqrt{x}, x = 100, dx = -0.6.$$

$$f(x + \Delta x) \approx f(x) + f'(x) dx$$

$$= \sqrt{x} + \frac{1}{2\sqrt{x}} dx$$

$$f(x + \Delta x) = \sqrt{99.4}$$

$$\approx \sqrt{100} + \frac{1}{2\sqrt{100}} (-0.6) = 9.97$$

Using a calculator:  $\sqrt{99.4} \approx 9.96995$

$$49. \text{ Let } f(x) = \sqrt{x}, x = 4, dx = 0.02, f'(x) = 1/(2\sqrt{x}).$$

Then


$$f(4.02) \approx f(4) + f'(4) dx$$

$$\sqrt{4.02} \approx \sqrt{4} + \frac{1}{2\sqrt{4}} (0.02) = 2 + \frac{1}{4} (0.02).$$

55. True

## Review Exercises for Chapter 3

1. A number  $c$  in the domain of  $f$  is a critical number if  $f'(c) = 0$  or  $f'$  is undefined at  $c$ .



$$3. g(x) = 2x + 5 \cos x, [0, 2\pi]$$

$$g'(x) = 2 - 5 \sin x$$

$$= 0 \text{ when } \sin x = \frac{2}{5}.$$


Critical numbers:  $x \approx 0.41, x \approx 2.73$

Left endpoint:  $(0, 5)$

Critical number:  $(0.41, 5.41)$

Critical number:  $(2.73, 0.88)$  Minimum

Right endpoint:  $(2\pi, 17.57)$  Maximum



5. Yes.  $f(-3) = f(2) = 0$ .  $f$  is continuous on  $[-3, 2]$ , differentiable on  $(-3, 2)$ .

$$f'(x) = (x + 3)(3x - 1) = 0 \text{ for } x = \frac{1}{3}.$$

$$c = \frac{1}{3} \text{ satisfies } f'(c) = 0.$$

9.  $f(x) = x^{2/3}, 1 \leq x \leq 8$

$$f'(x) = \frac{2}{3}x^{-1/3}$$

$$\frac{f(b) - f(a)}{b - a} = \frac{4 - 1}{8 - 1} = \frac{3}{7}$$

$$f'(c) = \frac{2}{3}c^{-1/3} = \frac{3}{7}$$

$$c = \left(\frac{14}{9}\right)^3 = \frac{2744}{729} \approx 3.764$$

13.  $f(x) = Ax^2 + Bx + C$

$$f'(x) = 2Ax + B$$

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = \frac{A(x_2^2 - x_1^2) + B(x_2 - x_1)}{x_2 - x_1}$$

$$= A(x_1 + x_2) + B$$

$$f'(c) = 2Ac + B = A(x_1 + x_2) + B$$

$$2Ac = A(x_1 + x_2)$$

$$c = \frac{x_1 + x_2}{2} = \text{Midpoint of } [x_1, x_2]$$

15.  $f(x) = (x - 1)^2(x - 3)$

$$f'(x) = (x - 1)^2(1) + (x - 3)(2)(x - 1)$$

$$= (x - 1)(3x - 7)$$

$$\text{Critical numbers: } x = 1 \text{ and } x = \frac{7}{3}$$

17.  $h(x) = \sqrt{x}(x - 3) = x^{3/2} - 3x^{1/2}$


$$\text{Domain: } (0, \infty)$$

$$h'(x) = \frac{3}{2}x^{1/2} - \frac{3}{2}x^{-1/2}$$

$$= \frac{3}{2}x^{-1/2}(x - 1) = \frac{3(x - 1)}{2\sqrt{x}}$$

$$\text{Critical number: } x = 1$$

7.  $f(x) = 3 - |x - 4|$



$$f(1) = f(7) = 0$$

- (b)  $f$  is not differentiable at  $x = 4$ .

11.  $f(x) = x - \cos x, -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$

$$f'(x) = 1 + \sin x$$

$$\frac{f(b) - f(a)}{b - a} = \frac{(\pi/2) - (-\pi/2)}{(\pi/2) - (-\pi/2)} = 1$$

$$f'(c) = 1 + \sin c = 1$$

$$c = 0$$

|                   |                   |                       |                            |
|-------------------|-------------------|-----------------------|----------------------------|
| Interval:         | $-\infty < x < 1$ | $1 < x < \frac{7}{3}$ | $\frac{7}{3} < x < \infty$ |
| Sign of $f'(x)$ : | $f'(x) > 0$       | $f'(x) < 0$           | $f'(x) > 0$                |
| Conclusion:       | Increasing        | Decreasing            | Increasing                 |

|                   |             |                  |
|-------------------|-------------|------------------|
| Interval:         | $0 < x < 1$ | $1 < x < \infty$ |
| Sign of $h'(x)$ : | $h'(x) < 0$ | $h'(x) > 0$      |
| Conclusion:       | Decreasing  | Increasing       |

19.  $h(t) = \frac{1}{4}t^4 - 8t$

$h'(t) = t^3 - 8 = 0$  when  $t = 2$ .

 Relative minimum:  $(2, -12)$ 

|                   |                   |                  |
|-------------------|-------------------|------------------|
| Test Interval:    | $-\infty < t < 2$ | $2 < t < \infty$ |
| Sign of $h'(t)$ : | $h'(t) < 0$       | $h'(t) > 0$      |
| Conclusion:       | Decreasing        | Increasing       |

21.  $y = \frac{1}{3} \cos(12t) - \frac{1}{4} \sin(12t)$

$v = y' = -4 \sin(12t) - 3 \cos(12t)$

 (a) When  $t = \frac{\pi}{8}$ ,  $y = \frac{1}{4}$  inch and  $v = y' = 4$  inches/second.

 (b)  $y' = -4 \sin(12t) - 3 \cos(12t) = 0$  when  $\frac{\sin(12t)}{\cos(12t)} = -\frac{3}{4} \Rightarrow \tan(12t) = -\frac{3}{4}$ .

 Therefore,  $\sin(12t) = -\frac{3}{5}$  and  $\cos(12t) = \frac{4}{5}$ . The maximum displacement is

$$y = \left(\frac{1}{3}\right)\left(\frac{4}{5}\right) - \frac{1}{4}\left(-\frac{3}{5}\right) = \frac{5}{12} \text{ inch.}$$

 (c) Period:  $\frac{2\pi}{12} = \frac{\pi}{6}$ 

Frequency:  $\frac{1}{\pi/6} = \frac{6}{\pi}$

23.  $f(x) = x + \cos x$ ,  $0 \leq x \leq 2\pi$

$f'(x) = 1 - \sin x$

$f''(x) = -\cos x = 0$  when  $x = \frac{\pi}{2}, \frac{3\pi}{2}$ .

 Points of inflection:  $\left(\frac{\pi}{2}, \frac{\pi}{2}\right), \left(\frac{3\pi}{2}, \frac{3\pi}{2}\right)$ 

|                    |                         |                                      |                             |
|--------------------|-------------------------|--------------------------------------|-----------------------------|
| Test Interval:     | $0 < x < \frac{\pi}{2}$ | $\frac{\pi}{2} < x < \frac{3\pi}{2}$ | $\frac{3\pi}{2} < x < 2\pi$ |
| Sign of $f''(x)$ : | $f''(x) < 0$            | $f''(x) > 0$                         | $f''(x) < 0$                |
| Conclusion:        | Concave downward        | Concave upward                       | Concave downward            |


25.  $g(x) = 2x^2(1 - x^2)$

$g'(x) = -4x(2x^2 - 1)$  Critical numbers:  $x = 0, \pm\frac{1}{\sqrt{2}}$


$g''(x) = 4 - 24x^2$

$g''(0) = 4 > 0$  Relative minimum at  $(0, 0)$

$g''\left(\pm\frac{1}{\sqrt{2}}\right) = -8 < 0$  Relative maximums at  $\left(\pm\frac{1}{\sqrt{2}}, \frac{1}{2}\right)$




27.



29. The first derivative is positive and the second derivative is negative. The graph is increasing and is concave down.

31. (a)  $D = 0.0034t^4 - 0.2352t^3 + 4.9423t^2 - 20.8641t + 94.4025$



(c) Maximum at (21.9, 319.5) ( $\approx 1992$ )

Minimum at (2.6, 69.6) ( $\approx 1972$ )

(d) Outlays increasing at greatest rate at the point of inflection (9.8, 173.7) ( $\approx 1979$ )

33.  $\lim_{x \rightarrow \infty} \frac{2x^2}{3x^2 + 5} = \lim_{x \rightarrow \infty} \frac{2}{3 + 5/x^2} = \frac{2}{3}$

35.  $\lim_{x \rightarrow \infty} \frac{5 \cos x}{x} = 0$ , since  $|5 \cos x| \leq 5$ .

37.  $h(x) = \frac{2x + 3}{x - 4}$

Discontinuity:  $x = 4$

$$\lim_{x \rightarrow \infty} \frac{2x + 3}{x - 4} = \lim_{x \rightarrow \infty} \frac{2 + (3/x)}{1 - (4/x)} = 2$$

Vertical asymptote:  $x = 4$

Horizontal asymptote:  $y = 2$

39.  $f(x) = \frac{3}{x} - 2$

Discontinuity:  $x = 0$

$$\lim_{x \rightarrow \infty} \left( \frac{3}{x} - 2 \right) = -2$$


Vertical asymptote:  $x = 0$

Horizontal asymptote:  $y = -2$

41.  $f(x) = x^3 + \frac{243}{x}$

Relative minimum: (3, 108)

Relative maximum: (-3, -108)




Vertical asymptote:  $x = 0$

43.  $f(x) = \frac{x - 1}{1 + 3x^2}$

Relative minimum: (-0.155, -1.077)

Relative maximum: (2.155, 0.077)



Horizontal asymptote:  $y = 0$

45.  $f(x) = 4x - x^2 = x(4 - x)$


Domain:  $(-\infty, \infty)$ ; Range:  $(-\infty, 4)$

$$f'(x) = 4 - 2x = 0 \text{ when } x = 2.$$

$$f''(x) = -2$$

Therefore, (2, 4) is a relative maximum.

Intercepts: (0, 0), (4, 0)



47.  $f(x) = x\sqrt{16 - x^2}$ , Domain:  $[-4, 4]$ , Range:  $[-8, 8]$

Domain:  $[-4, 4]$ ; Range:  $[-8, 8]$

$$f'(x) = \frac{16 - 2x^2}{\sqrt{16 - x^2}} = 0 \text{ when } x = \pm 2\sqrt{2} \text{ and undefined when } x = \pm 4.$$

$$f''(x) = \frac{2x(x^2 - 24)}{(16 - x^2)^{3/2}}$$

$$f'(-2\sqrt{2}) > 0$$

Therefore,  $(-2\sqrt{2}, -8)$  is a relative minimum.


$$f'(2\sqrt{2}) < 0$$

Therefore,  $(2\sqrt{2}, 8)$  is a relative maximum.

Point of inflection:  $(0, 0)$

Intercepts:  $(-4, 0)$ ,  $(0, 0)$ ,  $(4, 0)$

Symmetry with respect to origin



49.  $f(x) = (x - 1)^3(x - 3)^2$

Domain:  $(-\infty, \infty)$ ; Range:  $(-\infty, \infty)$

$$f'(x) = (x - 1)^2(x - 3)(5x - 11) = 0 \text{ when } x = 1, \frac{11}{5}, 3.$$

$$f''(x) = 4(x - 1)(5x^2 - 22x + 23) = 0 \text{ when } x = 1, \frac{11 \pm \sqrt{6}}{5}.$$

$$f''(3) > 0$$


Therefore,  $(3, 0)$  is a relative minimum.

$$f''\left(\frac{11}{5}\right) < 0$$

Therefore,  $\left(\frac{11}{5}, \frac{3456}{3125}\right)$  is a relative maximum.

Points of inflection:  $(1, 0)$ ,  $\left(\frac{11 - \sqrt{6}}{5}, 0.60\right)$ ,  $\left(\frac{11 + \sqrt{6}}{5}, 0.46\right)$

Intercepts:  $(0, -9)$ ,  $(1, 0)$ ,  $(3, 0)$



51.  $f(x) = x^{1/3}(x + 3)^{2/3}$


Domain:  $(-\infty, \infty)$ ; Range:  $(-\infty, \infty)$

$$f'(x) = \frac{x + 1}{(x + 3)^{1/3}x^{2/3}} = 0 \text{ when } x = -1 \text{ and undefined when } x = -3, 0.$$

$$f''(x) = \frac{-2}{x^{5/3}(x + 3)^{4/3}} \text{ is undefined when } x = 0, -3.$$

By the First Derivative Test  $(-3, 0)$  is a relative maximum and  $(-1, -\sqrt[3]{4})$  is a relative minimum.  $(0, 0)$  is a point of inflection.

Intercepts:  $(-3, 0)$ ,  $(0, 0)$




53.  $f(x) = \frac{x+1}{x-1}$

Domain:  $(-\infty, 1), (1, \infty)$ ; Range:  $(-\infty, 1), (1, \infty)$ 

$$f'(x) = \frac{-2}{(x-1)^2} < 0 \text{ if } x \neq 1.$$

$$f''(x) = \frac{4}{(x-1)^3}$$

Horizontal asymptote:  $y = 1$ Vertical asymptote:  $x = 1$ Intercepts:  $(-1, 0), (0, -1)$ 

55.  $f(x) = \frac{4}{1+x^2}$

Domain:  $(-\infty, \infty)$ ; Range:  $(0, 4]$ 


$$f'(x) = \frac{-8x}{(1+x^2)^2} = 0 \text{ when } x = 0.$$

$$f''(x) = \frac{-8(1-3x^2)}{(1+x^2)^3} = 0 \text{ when } x = \pm \frac{\sqrt{3}}{3}.$$

$$f''(0) < 0$$

Therefore,  $(0, 4)$  is a relative maximum.Points of inflection:  $(\pm\sqrt{3}/3, 3)$ Intercept:  $(0, 4)$ 

Symmetric to the y-axis

Horizontal asymptote:  $y = 0$ 

57.  $f(x) = x^3 + x + \frac{4}{x}$

Domain:  $(-\infty, 0), (0, \infty)$ ; Range:  $(-\infty, -6], [6, \infty)$ 

$$f'(x) = 3x^2 + 1 - \frac{4}{x^2} = \frac{3x^4 + x^2 - 4}{x^2} = 0 \text{ when } x = \pm 1.$$

$$f''(x) = 6x + \frac{8}{x^3} = \frac{6x^4 + 8}{x^3} \neq 0$$


$$f''(-1) < 0$$

Therefore,  $(-1, -6)$  is a relative maximum.

$$f''(1) > 0$$

Therefore,  $(1, 6)$  is a relative minimum.Vertical asymptote:  $x = 0$ 

Symmetric with respect to origin



59.  $f(x) = |x^2 - 9|$

Domain:  $(-\infty, \infty)$ ; Range:  $[0, \infty)$

$$f'(x) = \frac{2x(x^2 - 9)}{|x^2 - 9|} = 0 \text{ when } x = 0 \text{ and is undefined when } x = \pm 3.$$

$$f''(x) = \frac{2(x^2 - 9)}{|x^2 - 9|} \text{ is undefined at } x = \pm 3.$$

$$f''(0) < 0$$


Therefore,  $(0, 9)$  is a relative maximum.

Relative minima:  $(\pm 3, 0)$

Points of inflection:  $(\pm 3, 0)$

Intercepts:  $(\pm 3, 0), (0, 9)$

Symmetric to the y-axis



61.  $f(x) = x + \cos x$


Domain:  $[0, 2\pi]$ ; Range:  $[1, 1 + 2\pi]$

$$f'(x) = 1 - \sin x \geq 0, \text{ } f \text{ is increasing.}$$

$$f''(x) = -\cos x = 0 \text{ when } x = \frac{\pi}{2}, \frac{3\pi}{2}.$$

$$\text{Points of inflection: } \left(\frac{\pi}{2}, \frac{\pi}{2}\right), \left(\frac{3\pi}{2}, \frac{3\pi}{2}\right)$$

Intercept:  $(0, 1)$



63.  $x^2 + 4y^2 - 2x - 16y + 13 = 0$

(a)  $(x^2 - 2x + 1) + 4(y^2 - 4y + 4) = -13 + 1 + 16$


$$(x - 1)^2 + 4(y - 2)^2 = 4$$

$$\frac{(x - 1)^2}{4} + \frac{(y - 2)^2}{1} = 1$$

The graph is an ellipse:

Maximum:  $(1, 3)$

Minimum:  $(1, 1)$



(b)  $x^2 + 4y^2 - 2x - 16y + 13 = 0$

$$2x + 8y \frac{dy}{dx} - 2 - 16 \frac{dy}{dx} = 0$$

$$\frac{dy}{dx}(8y - 16) = 2 - 2x$$

$$\frac{dy}{dx} = \frac{2 - 2x}{8y - 16} = \frac{1 - x}{4y - 8}$$

The critical numbers are  $x = 1$  and  $y = 2$ . These correspond to the points  $(1, 1)$ ,  $(1, 3)$ ,  $(2, -1)$ , and  $(2, 3)$ . Hence, the maximum is  $(1, 3)$  and the minimum is  $(1, 1)$ .

65. Let  $t = 0$  at noon.

$$L = d^2 = (100 - 12t)^2 + (-10t)^2 = 10,000 - 2400t + 244t^2$$


$$\frac{dL}{dt} = -2400 + 488t = 0 \text{ when } t = \frac{300}{61} \approx 4.92 \text{ hr.}$$

Ship A at (40.98, 0); Ship B at (0, -49.18)

$$d^2 = 10,000 - 2400t + 244t^2$$

$$\approx 4098.36 \text{ when } t \approx 4.92 \approx 4:55 \text{ P.M.}$$

$$d \approx 64 \text{ km}$$



67. We have points  $(0, y)$ ,  $(x, 0)$ , and  $(1, 8)$ . Thus,

$$m = \frac{y - 8}{0 - 1} = \frac{0 - 8}{x - 1} \text{ or } y = \frac{8x}{x - 1}.$$


$$\text{Let } f(x) = L^2 = x^2 + \left(\frac{8x}{x - 1}\right)^2.$$

$$f'(x) = 2x + 128\left(\frac{x}{x - 1}\right)\left[\frac{(x - 1) - x}{(x - 1)^2}\right] = 0$$

$$x - \frac{64x}{(x - 1)^3} = 0$$

$$x[(x - 1)^3 - 64] = 0 \text{ when } x = 0, 5 \text{ (minimum).}$$

Vertices of triangle:  $(0, 0)$ ,  $(5, 0)$ ,  $(0, 10)$




69.  $A = (\text{Average of bases})(\text{Height})$

$$= \left(\frac{x + s}{2}\right) \frac{\sqrt{3s^2 + 2sx - x^2}}{2} \text{ (see figure)}$$

$$\frac{dA}{dx} = \frac{1}{4} \left[ \frac{(s - x)(s + x)}{\sqrt{3s^2 + 2sx - x^2}} + \sqrt{3s^2 + 2sx - x^2} \right]$$

$$= \frac{2(2s - x)(s + x)}{4\sqrt{3s^2 + 2sx - x^2}} = 0 \text{ when } x = 2s.$$

$A$  is a maximum when  $x = 2s$ .



71. You can form a right triangle with vertices  $(0, 0)$ ,  $(x, 0)$  and  $(0, y)$ .

Assume that the hypotenuse of length  $L$  passes through  $(4, 6)$ .

$$m = \frac{y - 6}{0 - 4} = \frac{6 - 0}{4 - x} \text{ or } y = \frac{6x}{x - 4}$$

$$\text{Let } f(x) = L^2 = x^2 + y^2 = x^2 + \left(\frac{6x}{x - 4}\right)^2.$$

$$f'(x) = 2x + 72\left(\frac{x}{x - 4}\right)\left[\frac{-4}{(x - 4)^2}\right] = 0$$

$$x[(x - 4)^3 - 144] = 0 \text{ when } x = 0 \text{ or } x = 4 + \sqrt[3]{144}.$$

$$L \approx 14.05 \text{ feet}$$



73.  $\csc \theta = \frac{L_1}{6}$  or  $L_1 = 6 \csc \theta$  (see figure)

$$\csc\left(\frac{\pi}{2} - \theta\right) = \frac{L_2}{9} \text{ or } L_2 = 9 \csc\left(\frac{\pi}{2} - \theta\right)$$

$$L = L_1 + L_2 = 6 \csc \theta + 9 \csc\left(\frac{\pi}{2} - \theta\right) = 6 \csc \theta + 9 \sec \theta$$


$$\frac{dL}{d\theta} = -6 \csc \theta \cot \theta + 9 \sec \theta \tan \theta = 0$$

$$\tan^3 \theta = \frac{2}{3} \Rightarrow \tan \theta = \frac{\sqrt[3]{2}}{\sqrt[3]{3}}$$

$$\sec \theta = \sqrt{1 + \tan^2 \theta} = \sqrt{1 + \left(\frac{2}{3}\right)^{2/3}} = \frac{\sqrt{3^{2/3} + 2^{2/3}}}{3^{1/3}}$$

$$\csc \theta = \frac{\sec \theta}{\tan \theta} = \frac{\sqrt{3^{2/3} + 2^{2/3}}}{2^{1/3}}$$

$$L = 6 \frac{(3^{2/3} + 2^{2/3})^{1/2}}{2^{1/3}} + 9 \frac{(3^{2/3} + 2^{2/3})^{1/2}}{3^{1/3}} = 3(3^{2/3} + 2^{2/3})^{3/2} \text{ ft} \approx 21.07 \text{ ft (Compare to Exercise 72 using } a = 9 \text{ and } b = 6.)$$



75. Total cost = (Cost per hour)(Number of hours)

$$T = \left(\frac{v^2}{600} + 5\right)\left(\frac{110}{v}\right) = \frac{11v}{60} + \frac{550}{v}$$

$$\frac{dT}{dv} = \frac{11}{60} - \frac{550}{v^2} = \frac{11v^2 - 33,000}{60v^2}$$

$$= 0 \text{ when } v = \sqrt{3000} = 10\sqrt{30} \approx 54.8 \text{ mph.}$$

$$\frac{d^2T}{dv^2} = \frac{1100}{v^3} > 0 \text{ when } v = 10\sqrt{30} \text{ so this value yields a minimum.}$$

77.  $f(x) = x^3 - 3x - 1$

From the graph you can see that  $f(x)$  has three real zeros.

$$f'(x) = 3x^2 - 3$$

| $n$ | $x_n$   | $f(x_n)$ | $f'(x_n)$ | $\frac{f(x_n)}{f'(x_n)}$ | $x_n - \frac{f(x_n)}{f'(x_n)}$ |
|-----|---------|----------|-----------|--------------------------|--------------------------------|
| 1   | -1.5000 | 0.1250   | 3.7500    | 0.0333                   | -1.5333                        |
| 2   | -1.5333 | -0.0049  | 4.0530    | -0.0012                  | -1.5321                        |

| $n$ | $x_n$   | $f(x_n)$ | $f'(x_n)$ | $\frac{f(x_n)}{f'(x_n)}$ | $x_n - \frac{f(x_n)}{f'(x_n)}$ |
|-----|---------|----------|-----------|--------------------------|--------------------------------|
| 1   | -0.5000 | 0.3750   | -2.2500   | -0.1667                  | -0.3333                        |
| 2   | -0.3333 | -0.0371  | -2.6667   | 0.0139                   | -0.3472                        |
| 3   | -0.3472 | -0.0003  | -2.6384   | 0.0001                   | -0.3473                        |

| $n$ | $x_n$   | $f(x_n)$ | $f'(x_n)$ | $\frac{f(x_n)}{f'(x_n)}$ | $x_n - \frac{f(x_n)}{f'(x_n)}$ |
|-----|---------|----------|-----------|--------------------------|--------------------------------|
| 1   | -1.9000 | 0.1590   | 7.8300    | 0.0203                   | 1.8797                         |
| 2   | 1.8797  | 0.0024   | 7.5998    | 0.0003                   | 1.8794                         |

The three real zeros of  $f(x)$  are  $x \approx -1.532$ ,  $x \approx -0.347$ , and  $x \approx 1.879$ .

79. Find the zeros of  $f(x) = x^4 - x - 3$ .

$$f'(x) = 4x^3 - 1$$

From the graph you can see that  $f(x)$  has two real zeros.

$f$  changes sign in  $[-2, -1]$ .

| $n$ | $x_n$   | $f(x_n)$ | $f'(x_n)$ | $\frac{f(x_n)}{f'(x_n)}$ | $x_n - \frac{f(x_n)}{f'(x_n)}$ |
|-----|---------|----------|-----------|--------------------------|--------------------------------|
| 1   | -1.2000 | 0.2736   | -7.9120   | -0.0346                  | -1.1654                        |
| 2   | -1.1654 | 0.0100   | -7.3312   | -0.0014                  | -1.1640                        |

On the interval  $[-2, -1]$ :  $x \approx -1.164$ .

$f$  changes sign in  $[1, 2]$ .

| $n$ | $x_n$  | $f(x_n)$ | $f'(x_n)$ | $\frac{f(x_n)}{f'(x_n)}$ | $x_n - \frac{f(x_n)}{f'(x_n)}$ |
|-----|--------|----------|-----------|--------------------------|--------------------------------|
| 1   | 1.5000 | 0.5625   | 12.5000   | 0.0450                   | 1.4550                         |
| 2   | 1.4550 | 0.0268   | 11.3211   | 0.0024                   | 1.4526                         |
| 3   | 1.4526 | -0.0003  | 11.2602   | 0.0000                   | 1.4526                         |

On the interval  $[1, 2]$ :  $x \approx 1.453$ .

81.  $y = x(1 - \cos x) = x - x \cos x$

$$\frac{dy}{dx} = 1 + x \sin x - \cos x$$

$$dy = (1 + x \sin x - \cos x) dx$$

83.  $S = 4\pi r^2$ .  $dr = \Delta r = \pm 0.025$

$$\begin{aligned} dS &= 8\pi r dr = 8\pi(9)(\pm 0.025) \\ &= \pm 1.8\pi \text{ square cm} \end{aligned}$$

$$\begin{aligned} \frac{dS}{S}(100) &= \frac{8\pi r dr}{4\pi r^2}(100) = \frac{2 dr}{r}(100) \\ &= \frac{2(\pm 0.025)}{9}(100) \approx \pm 0.56\% \end{aligned}$$

$$V = \frac{4}{3}\pi r^3$$

$$\begin{aligned} dV &= 4\pi r^2 dr = 4\pi(9)^2(\pm 0.025) \\ &= \pm 8.1\pi \text{ cubic cm} \end{aligned}$$

$$\begin{aligned} \frac{dV}{V}(100) &= \frac{4\pi r^2 dr}{(4/3)\pi r^3}(100) = \frac{3 dr}{r}(100) \\ &= \frac{3(\pm 0.025)}{9}(100) \approx \pm 0.83\% \end{aligned}$$

## Problem Solving for Chapter 3

1. Assume  $y_1 < d < y_2$ . Let  $g(x) = f(x) - d(x - a)$ .  $g$  is continuous on  $[a, b]$  and therefore has a minimum  $(c, g(c))$  on  $[a, b]$ . The point  $c$  cannot be an endpoint of  $[a, b]$  because

$$g'(a) = f'(a) - d = y_1 - d < 0$$

$$g'(b) = f'(b) - d = y_2 - d > 0$$

Hence,  $a < c < b$  and  $g'(c) = 0 \Rightarrow f'(c) = d$ .

3. (a) For  $a = -3, -2, -1, 0$ ,  $p$  has a relative maximum at  $(0, 0)$ .

For  $a = 1, 2, 3$ ,  $p$  has a relative maximum at  $(0, 0)$  and 2 relative minima.

(b)  $p'(x) = 4ax^3 - 12x = 4x(ax^2 - 3) = 0 \Rightarrow x = 0, \pm\sqrt{\frac{3}{a}}$

$p''(x) = 12ax^2 - 12 = 12(ax^2 - 1)$

For  $x = 0, p''(0) = -12 < 0 \Rightarrow p$  has a relative maximum at  $(0, 0)$ .

(c) If  $a > 0, x = \pm\sqrt{\frac{3}{a}}$  are the remaining critical numbers.


$p''\left(\pm\sqrt{\frac{3}{a}}\right) = 12\left(\frac{3}{a}\right) - 12 = 24 > 0 \Rightarrow p$  has relative minima for  $a > 0$ .

(d)  $(0, 0)$  lies on  $y = -3x^2$ .

Let  $x = \pm\sqrt{\frac{3}{a}}$ . Then

$p(x) = a\left(\frac{3}{a}\right)^2 - 6\left(\frac{3}{a}\right) = \frac{9}{a} - \frac{18}{a} = -\frac{9}{a}$ .

Thus,  $y = -\frac{9}{a} = -3\left(\pm\sqrt{\frac{3}{a}}\right)^2 = -3x^2$  is satisfied by all the relative extrema of  $p$ .



5.  $p(x) = x^4 + ax^2 + 1$

(a)  $p'(x) = 4x^3 + 2ax = 2x(2x^2 + a)$


$p''(x) = 16x^2 + 2a$

For  $a \geq 0$ , there is one relative minimum at  $(0, 0)$ .

(b) For  $a < 0$ , there is a relative maximum at  $(0, 1)$ .

(c) For  $a < 0$ , there are two relative minima at  $x = \pm\sqrt{-\frac{a}{2}}$ .

(d) There are either 1 or 3 critical points. The above analysis shows that there cannot be exactly two relative extrema.



7.  $f(x) = \frac{c}{x} + x^2$

$f'(x) = -\frac{c}{x^2} + 2x = 0 \Rightarrow \frac{c}{x^2} = 2x \Rightarrow x^3 = \frac{c}{2} \Rightarrow x = \sqrt[3]{\frac{c}{2}}$

$f''(x) = \frac{2c}{x^3} + 2$

If  $c = 0, f(x) = x^2$  has a relative minimum, but no relative maximum.

If  $c > 0, x = \sqrt[3]{\frac{c}{2}}$  is a relative minimum, because  $f''\left(\sqrt[3]{\frac{c}{2}}\right) > 0$ .

If  $c < 0, x = \sqrt[3]{\frac{c}{2}}$  is a relative minimum too.

Answer: all  $c$ .

9. Set  $\frac{f(b) - f(a) - f'(a)(b - a)}{(b - a)^2} = k$ .

Define  $F(x) = f(x) - f(a) - f'(a)(x - a) - k(x - a)^2$ .

$F(a) = 0, F(b) = f(b) - f(a) - f'(a)(b - a) - k(b - a)^2 = 0$

$F$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ .

There exists  $c_1, a < c_1 < b$ , satisfying  $F'(c_1) = 0$ .

$F'(x) = f'(x) - f'(a) - 2k(x - a)$  satisfies the hypothesis of Rolle's Theorem on  $[a, c_1]$ :

$F'(a) = 0, F'(c_1) = 0$ .

There exists  $c_2, a < c_2 < c_1$  satisfying  $F''(c_2) = 0$ .

Finally,  $F''(x) = f''(x) - 2k$  and  $F''(c_2) = 0$  implies that

$k = \frac{f''(c_2)}{2}$ .

Thus,  $k = \frac{f(b) - f(a) - f'(a)(b - a)}{(b - a)^2} = \frac{f''(c_2)}{2} \Rightarrow f(b) = f(a) + f'(a)(b - a) + \frac{1}{2}f''(c_2)(b - a)^2$ .

11.  $E(\phi) = \frac{\tan \phi(1 - 0.1 \tan \phi)}{0.1 + \tan \phi} = \frac{10 \tan \phi - \tan^2 \phi}{1 + 10 \tan \phi}$

$E'(\phi) = \frac{(1 + 10 \tan \phi)(10 \sec^2 \phi - 2 \tan \phi \sec^2 \phi) - (10 \tan \phi - \tan^2 \phi)10 \sec^2 \phi}{(1 + 10 \tan \phi)^2} = 0$

$\Rightarrow (1 + 10 \tan \phi)(10 \sec^2 \phi - 2 \tan \phi \sec^2 \phi) = (10 \tan \phi - \tan^2 \phi)10 \sec^2 \phi$

$\Rightarrow 10 \sec^2 \phi - 2 \tan \phi \sec^2 \phi + 100 \tan \phi \sec^2 \phi - 20 \tan^2 \phi \sec^2 \phi$

$= 100 \tan \phi \sec^2 \phi - 10 \tan^2 \phi \sec^2 \phi$

$\Rightarrow 10 - 2 \tan \phi = 10 \tan^2 \phi$

$\Rightarrow 10 \tan^2 \phi + 2 \tan \phi - 10 = 0$

$\tan \phi = \frac{-2 \pm \sqrt{4 + 400}}{20} \approx 0.90499, -1.10499$

Using the positive value,  $\phi \approx 0.7356$ , or  $42.14^\circ$ .

13.  $v = -2400\pi \sin \theta$

$v' = -2400\pi \cos \theta = 0$

$\theta = \frac{\pi}{2} + 2n\pi, \frac{3\pi}{2} + 2n\pi, n$  an integer

15. The line has equation  $\frac{x}{3} + \frac{y}{4} = 1$  or  $y = -\frac{4}{3}x + 4$ .

Rectangle:

$$\text{Area} = A = xy = x\left(-\frac{4}{3}x + 4\right) = -\frac{4}{3}x^2 + 4x.$$

$$A'(x) = -\frac{8}{3}x + 4 = 0 \Rightarrow \frac{8}{3}x = 4 \Rightarrow x = \frac{3}{2}$$

Dimensions:  $\frac{3}{2} \times 2$  Calculus was helpful.

Circle: The distance from the center  $(r, r)$  to the line  $\frac{x}{3} + \frac{y}{4} - 1 = 0$  must be  $r$ :

$$r = \frac{\left|\frac{r}{3} + \frac{r}{4} - 1\right|}{\sqrt{\frac{1}{9} + \frac{1}{16}}} = \frac{12|7r - 12|}{5|12|} = \frac{|7r - 12|}{5}$$

$$5r = |7r - 12| \Rightarrow r = 1 \text{ or } r = 6.$$

Clearly,  $r = 1$ .

Semicircle: The center lies on the line  $\frac{x}{3} + \frac{y}{4} = 1$  and satisfies  $x = y = r$ .

Thus  $\frac{r}{3} + \frac{r}{4} = 1 \Rightarrow \frac{7}{12}r = 1 \Rightarrow r = \frac{12}{7}$ . No calculus necessary.

17.  $y = (1 + x^2)^{-1}$

$$y' = \frac{-2x}{(1 + x^2)^2}$$

$$y'' = \frac{2(3x^2 - 1)}{(x^2 + 1)^3} = 0 \Rightarrow x = \pm \frac{1}{\sqrt{3}} = \pm \frac{\sqrt{3}}{3}$$

$$y'': \begin{array}{c} \text{+++} \text{---} \text{---} \text{---} \text{+++} \\ \frac{1}{-\frac{\sqrt{3}}{3}} \quad 0 \quad \frac{\sqrt{3}}{3} \end{array}$$

The tangent line has greatest slope at  $\left(-\frac{\sqrt{3}}{3}, \frac{3}{4}\right)$  and least slope at  $\left(\frac{\sqrt{3}}{3}, \frac{3}{4}\right)$ .

19. (a)

|          |         |         |         |         |         |         |
|----------|---------|---------|---------|---------|---------|---------|
| $x$      | 0.1     | 0.2     | 0.3     | 0.4     | 0.5     | 1.0     |
| $\sin x$ | 0.09983 | 0.19867 | 0.29552 | 0.38942 | 0.47943 | 0.84147 |

$$\sin x \leq x$$

(b) Let  $f(x) = \sin x$ . Then  $f'(x) = \cos x$  and on  $[0, x]$  you have by the Mean Value Theorem,

$$f'(c) = \frac{f(x) - f(0)}{x - 0}, \quad 0 < c < x$$

$$\cos(c) = \frac{\sin x}{x}$$

$$\text{Hence, } \left|\frac{\sin x}{x}\right| = |\cos(c)| \leq 1$$

$$\Rightarrow |\sin x| \leq |x|$$

$$\Rightarrow \sin x \leq x$$