

85. As k increases, the time required for the object to reach the ground increases.

$$87. y = \cosh x = \frac{e^x + e^{-x}}{2}$$

$$y' = \frac{e^x - e^{-x}}{2} = \sinh x$$

$$89. y = \cosh^{-1} x$$

$$\cosh y = x$$

$$(\sinh y)(y') = 1$$

$$y' = \frac{1}{\sinh y} = \frac{1}{\sqrt{\cosh^2 y - 1}} = \frac{1}{\sqrt{x^2 - 1}}$$

$$91. y = \operatorname{sech} x = \frac{2}{e^x + e^{-x}}$$


$$y' = -2(e^x + e^{-x})^{-2}(e^x - e^{-x}) = \left(\frac{-2}{e^x + e^{-x}}\right)\left(\frac{e^x - e^{-x}}{e^x + e^{-x}}\right) = -\operatorname{sech} x \tanh x$$

Review Exercises for Chapter 5

1. $f(x) = \ln x + 3$

Vertical shift 3 units upward

Vertical asymptote: $x = 0$



$$3. \ln \sqrt[5]{\frac{4x^2 - 1}{4x^2 + 1}} = \frac{1}{5} \ln \frac{(2x - 1)(2x + 1)}{4x^2 + 1} = \frac{1}{5} [\ln(2x - 1) + \ln(2x + 1) - \ln(4x^2 + 1)]$$

$$5. \ln 3 + \frac{1}{3} \ln(4 - x^2) - \ln x = \ln 3 + \ln \sqrt[3]{4 - x^2} - \ln x = \ln \left(\frac{3\sqrt[3]{4 - x^2}}{x} \right)$$

$$7. \ln \sqrt{x + 1} = 2$$

$$\sqrt{x + 1} = e^2$$

$$x + 1 = e^4$$

$$x = e^4 - 1 \approx 53.598$$

$$9. g(x) = \ln \sqrt{x} = \frac{1}{2} \ln x$$

$$g'(x) = \frac{1}{2x}$$

$$11. f(x) = x\sqrt{\ln x}$$

$$f'(x) = \left(\frac{x}{2}\right)(\ln x)^{-1/2}\left(\frac{1}{x}\right) + \sqrt{\ln x}$$

$$= \frac{1}{2\sqrt{\ln x}} + \sqrt{\ln x} = \frac{1 + 2\ln x}{2\sqrt{\ln x}}$$

$$13. y = \frac{1}{b^2} \left[\ln(a + bx) + \frac{a}{a + bx} \right]$$

$$\frac{dy}{dx} = \frac{1}{b^2} \left[\frac{b}{a + bx} - \frac{ab}{(a + bx)^2} \right] = \frac{x}{(a + bx)^2}$$

$$15. y = -\frac{1}{a} \ln \left(\frac{a + bx}{x} \right) = -\frac{1}{a} [\ln(a + bx) - \ln x]$$

$$\frac{dy}{dx} = -\frac{1}{a} \left(\frac{b}{a + bx} - \frac{1}{x} \right) = \frac{1}{x(a + bx)}$$

$$17. u = 7x - 2, du = 7dx$$

$$\int \frac{1}{7x - 2} dx = \frac{1}{7} \int \frac{1}{7x - 2} (7) dx = \frac{1}{7} \ln|7x - 2| + C$$

$$\begin{aligned}
 19. \int \frac{\sin x}{1 + \cos x} dx &= -\int \frac{-\sin x}{1 + \cos x} dx \\
 &= -\ln|1 + \cos x| + C
 \end{aligned}$$

$$23. \int_0^{\pi/3} \sec \theta d\theta = \left[\ln|\sec \theta + \tan \theta| \right]_0^{\pi/3} = \ln(2 + \sqrt{3})$$


$$\begin{aligned}
 25. (a) \quad f(x) &= \frac{1}{2}x - 3 \\
 y &= \frac{1}{2}x - 3 \\
 2(y + 3) &= x \\
 2(x + 3) &= y \\
 f^{-1}(x) &= 2x + 6
 \end{aligned}$$

$$\begin{aligned}
 27. (a) \quad f(x) &= \sqrt{x + 1} \\
 y &= \sqrt{x + 1} \\
 y^2 - 1 &= x \\
 x^2 - 1 &= y \\
 f^{-1}(x) &= x^2 - 1, x \geq 0
 \end{aligned}$$


$$\begin{aligned}
 29. (a) \quad f(x) &= \sqrt[3]{x + 1} \\
 y &= \sqrt[3]{x + 1} \\
 y^3 - 1 &= x \\
 x^3 - 1 &= y \\
 f^{-1}(x) &= x^3 - 1
 \end{aligned}$$

$$\begin{aligned}
 31. \quad f(x) &= x^3 + 2 \\
 f^{-1}(x) &= (x - 2)^{1/3} \\
 (f^{-1})'(x) &= \frac{1}{3}(x - 2)^{-2/3} \\
 (f^{-1})'(-1) &= \frac{1}{3}(-1 - 2)^{-2/3} = \frac{1}{3(-3)^{2/3}} \\
 &= \frac{1}{3^{5/3}} \approx 0.160
 \end{aligned}$$


$$21. \int_1^4 \frac{x + 1}{x} dx = \int_1^4 \left(1 + \frac{1}{x}\right) dx = \left[x + \ln|x| \right]_1^4 = 3 + \ln 4$$



$$\begin{aligned}
 (c) \quad f^{-1}(f(x)) &= f^{-1}\left(\frac{1}{2}x - 3\right) = 2\left(\frac{1}{2}x - 3\right) + 6 = x \\
 f(f^{-1}(x)) &= f(2x + 6) = \frac{1}{2}(2x + 6) - 3 = x
 \end{aligned}$$



$$\begin{aligned}
 (c) \quad f^{-1}(f(x)) &= f^{-1}(\sqrt{x + 1}) = \sqrt{(x^2 - 1)^2} - 1 = x \\
 f(f^{-1}(x)) &= f(x^2 - 1) = \sqrt{(x^2 - 1) + 1} \\
 &= \sqrt{x^2} = x \text{ for } x \geq 0.
 \end{aligned}$$



$$\begin{aligned}
 (c) \quad f^{-1}(f(x)) &= f^{-1}(\sqrt[3]{x + 1}) = (\sqrt[3]{x + 1})^3 - 1 = x \\
 f(f^{-1}(x)) &= f(x^3 - 1) = \sqrt[3]{(x^3 - 1) + 1} = x
 \end{aligned}$$

$$\begin{aligned}
 33. \quad f(x) &= \tan x \\
 f\left(\frac{\pi}{6}\right) &= \frac{\sqrt{3}}{3} \\
 f'(x) &= \sec^2 x \\
 f'\left(\frac{\pi}{6}\right) &= \frac{4}{3} \\
 (f^{-1})'\left(\frac{\sqrt{3}}{3}\right) &= \frac{1}{f'(\pi/6)} = \frac{3}{4}
 \end{aligned}$$

35. (a) $f(x) = \ln \sqrt{x}$


$$y = \ln \sqrt{x}$$

$$e^y = \sqrt{x}$$

$$e^{2y} = x$$

$$e^{2x} = y$$


$$f^{-1}(x) = e^{2x}$$



(c) $f^{-1}(f(x)) = f^{-1}(\ln \sqrt{x}) = e^{2 \ln \sqrt{x}} = e^{\ln x} = x$

$$f(f^{-1}(x)) = f(e^{2x}) = \ln \sqrt{e^{2x}} = \ln e^x = x$$

37. $y = e^{-x/2}$



41. $g(t) = t^2 e^t$

$$g'(t) = t^2 e^t + 2te^t = te^t(t + 2)$$

45. $g(x) = \frac{x^2}{e^x}$

$$g'(x) = \frac{e^x(2x) - x^2 e^x}{e^{2x}} = \frac{x(2-x)}{e^x}$$

49. Let $u = -3x^2$, $du = -6x dx$.

$$\int x e^{-3x^2} dx = -\frac{1}{6} \int e^{-3x^2} (-6x) dx = -\frac{1}{6} e^{-3x^2} + C$$

53. $\int x e^{1-x^2} dx = -\frac{1}{2} \int e^{1-x^2} (-2x) dx$

$$= -\frac{1}{2} e^{1-x^2} + C$$

57. $y = e^x(a \cos 3x + b \sin 3x)$

$$y' = e^x(-3a \sin 3x + 3b \cos 3x) + e^x(a \cos 3x + b \sin 3x)$$

$$= e^x[(-3a + b) \sin 3x + (a + 3b) \cos 3x]$$

$$y'' = e^x[3(-3a + b) \cos 3x - 3(a + 3b) \sin 3x] + e^x[(-3a + b) \sin 3x + (a + 3b) \cos 3x]$$

$$= e^x[(-6a - 8b) \sin 3x + (-8a + 6b) \cos 3x]$$

$$y'' - 2y' + 10y = e^x[(-6a - 8b) - 2(-3a + b) + 10b] \sin 3x + [(-8a + 6b) - 2(a + 3b) + 10a] \cos 3x = 0$$

39. $f(x) = \ln(e^{-x^2}) = -x^2$

$$f'(x) = -2x$$

43. $y = \sqrt{e^{2x} + e^{-2x}}$

$$y' = \frac{1}{2}(e^{2x} + e^{-2x})^{-1/2}(2e^{2x} - 2e^{-2x}) = \frac{e^{2x} - e^{-2x}}{\sqrt{e^{2x} + e^{-2x}}}$$

47. $y(\ln x) + y^2 = 0$

$$y\left(\frac{1}{x}\right) + (\ln x)\left(\frac{dy}{dx}\right) + 2y\left(\frac{dy}{dx}\right) = 0$$

$$(2y + \ln x)\frac{dy}{dx} = \frac{-y}{x}$$

$$\frac{dy}{dx} = \frac{-y}{x(2y + \ln x)}$$

51. $\int \frac{e^{4x} - e^{2x} + 1}{e^x} dx = \int (e^{3x} - e^x + e^{-x}) dx$

$$= \frac{1}{3} e^{3x} - e^x - e^{-x} + C$$


$$= \frac{e^{4x} - 3e^{2x} - 3}{3e^x} + C$$

55. Let $u = e^x - 1$, $du = e^x dx$.

$$\int \frac{e^x}{e^x - 1} dx = \ln|e^x - 1| + C$$

$$59. \text{Area} = \int_0^4 x e^{-x^2} dx = \left[-\frac{1}{2} e^{-x^2} \right]_0^4 = -\frac{1}{2}(e^{-16} - 1) \approx 0.500$$

$$61. y = 3^{3/2}$$



$$65. f(x) = 3^{x-1}$$

$$f'(x) = 3^{x-1} \ln 3$$

$$69. g(x) = \log_3 \sqrt{1-x} = \frac{1}{2} \log_3(1-x)$$

$$g'(x) = \frac{1}{2} \frac{-1}{(1-x)\ln 3} = \frac{1}{2(x-1)\ln 3}$$


$$73. (a) y = x^a$$

$$y' = ax^{a-1}$$

$$(b) y = a^x$$

$$y' = (\ln a)a^x$$

$$63. y = \log_2(x-1)$$



$$67. y = x^{2x+1}$$

$$\ln y = (2x+1) \ln x$$

$$\frac{y'}{y} = \frac{2x+1}{x} + 2 \ln x$$

$$y' = y \left(\frac{2x+1}{x} + 2 \ln x \right) = x^{2x+1} \left(\frac{2x+1}{x} + 2 \ln x \right)$$

$$71. \int (x+1)5^{(x+1)^2} dx = \frac{1}{2} \frac{1}{\ln 5} 5^{(x+1)^2} + C$$

$$(c) y = x^x$$

$$\ln y = x \ln x$$

$$\frac{1}{y} y' = x \cdot \frac{1}{x} + (1) \ln x$$

$$y' = y(1 + \ln x)$$

$$y' = x^x(1 + \ln x)$$

$$(d) y = a^a$$

$$y' = 0$$

$$75. 10,000 = P e^{(0.07)(15)}$$

$$P = \frac{10,000}{e^{1.05}} \approx \$3499.38$$

$$77. P(h) = 30e^{kh}$$

$$P(18,000) = 30e^{18,000k} = 15$$

$$k = \frac{\ln(1/2)}{18,000} = \frac{-\ln 2}{18,000}$$

$$P(h) = 30e^{-(h \ln 2)/18,000}$$

$$P(35,000) = 30e^{-(35,000 \ln 2)/18,000} \approx 7.79 \text{ inches}$$

$$79. P = C e^{0.015t}$$

$$2C = C e^{0.015t}$$

$$2 = e^{0.015t}$$

$$\ln 2 = 0.015t$$

$$t = \frac{\ln 2}{0.015} \approx 46.21 \text{ years}$$

$$81. \frac{dy}{dx} = \frac{x^2 + 3}{x}$$

$$\int dy = \int \left(x + \frac{3}{x} \right) dx$$

$$y = \frac{x^2}{2} + 3 \ln|x| + C$$

83. $y' - 2xy = 0$

$$\frac{dy}{dx} = 2xy$$

$$\int \frac{1}{y} dy = \int 2x dx$$

$$\ln|y| = x^2 + C_1$$

$$e^{x^2+C_1} = y$$

$$y = Ce^{x^2}$$

85. $\frac{dy}{dx} = \frac{x^2 + y^2}{2xy}$ (homogeneous differential equation)

$$(x^2 + y^2) dx - 2xy dy = 0$$

Let $y = vx$, $dy = x dv + v dx$.

$$(x^2 + v^2x^2) dx - 2x(vx)(x dv + v dx) = 0$$

$$(x^2 + v^2x^2 - 2x^2v^2) dx - 2x^3v dv = 0$$

$$(x^2 - x^2v^2) dx = 2x^3v dv$$

$$(1 - v^2) dx = 2x dv$$

$$\int \frac{dx}{x} = \int \frac{2v}{1 - v^2} dv$$

$$\ln|x| = -\ln|1 - v^2| + C_1 = -\ln|1 - v^2| + \ln C$$

$$x = \frac{C}{1 - v^2} = \frac{C}{1 - (y/x)^2} = \frac{Cx^2}{x^2 - y^2}$$

$$1 = \frac{Cx}{x^2 - y^2} \quad \text{or} \quad C_1 = \frac{x}{x^2 - y^2}$$

87. $y = C_1x + C_2x^3$

$$y' = C_1 + 3C_2x^2$$

$$y'' = 6C_2x$$

$$\begin{aligned} x^2y'' - 3xy' + 3y &= x^2(6C_2x) - 3x(C_1 + 3C_2x^2) + (C_1x + C_2x^3) \\ &= 6C_2x^3 - 3C_1x - 9C_2x^3 + 3C_1x + 3C_2x^3 = 0 \end{aligned}$$


$$x = 2, y = 0: 0 = 2C_1 + 8C_2 \Rightarrow C_1 = -4C_2$$

$$x = 2, y' = 4: 4 = C_1 + 12C_2$$

$$4 = (-4C_2) + 12C_2 = 8C_2 \Rightarrow C_2 = \frac{1}{2}, C_1 = -2$$

$$y = -2x + \frac{1}{2}x^3$$


89. $f(x) = 2 \arctan(x + 3)$



91. (a) Let $\theta = \arcsin \frac{1}{2}$

$$\sin \theta = \frac{1}{2}$$

$$\sin\left(\arcsin \frac{1}{2}\right) = \sin \theta = \frac{1}{2}.$$



(b) Let $\theta = \arcsin \frac{1}{2}$

$$\sin \theta = \frac{1}{2}$$

$$\cos\left(\arcsin \frac{1}{2}\right) = \cos \theta = \frac{\sqrt{3}}{2}.$$

93. $y = \tan(\arcsin x) = \frac{x}{\sqrt{1-x^2}}$
 $y' = \frac{(1-x^2)^{1/2} + x^2(1-x^2)^{-1/2}}{1-x^2} = (1-x^2)^{-3/2}$

95. $y = x \operatorname{arcsec} x$

$$y' = \frac{x}{|x|\sqrt{x^2-1}} + \operatorname{arcsec} x$$

97. $y = x(\arcsin x)^2 - 2x + 2\sqrt{1-x^2} \arcsin x$
 $y' = \frac{2x \arcsin x}{\sqrt{1-x^2}} + (\arcsin x)^2 - 2 + \frac{2\sqrt{1-x^2}}{\sqrt{1-x^2}} - \frac{2x}{\sqrt{1-x^2}} \arcsin x = (\arcsin x)^2$

99. Let $u = e^{2x}$, $du = 2e^{2x} dx$.

$$\int \frac{1}{e^{2x} + e^{-2x}} dx = \int \frac{e^{2x}}{1 + e^{4x}} dx = \frac{1}{2} \int \frac{1}{1 + (e^{2x})^2} (2e^{2x}) dx = \frac{1}{2} \arctan(e^{2x}) + C$$

101. Let $u = x^2$, $du = 2x dx$.

$$\int \frac{x}{\sqrt{1-x^4}} dx = \frac{1}{2} \int \frac{1}{\sqrt{1-(x^2)^2}} (2x) dx = \frac{1}{2} \arcsin x^2 + C$$

103. Let $u = 16 + x^2$, $du = 2x dx$.

$$\int \frac{x}{16+x^2} dx = \frac{1}{2} \int \frac{1}{16+x^2} (2x) dx = \frac{1}{2} \ln(16+x^2) + C$$

105. Let $u = \arctan\left(\frac{x}{2}\right)$, $du = \frac{2}{4+x^2} dx$.

$$\int \frac{\arctan(x/2)}{4+x^2} dx = \frac{1}{2} \int \left(\arctan \frac{x}{2}\right) \left(\frac{2}{4+x^2}\right) dx = \frac{1}{4} \left(\arctan \frac{x}{2}\right)^2 + C$$

107. $\int \frac{dy}{\sqrt{A^2-y^2}} = \int \sqrt{\frac{k}{m}} dt$
 $\arcsin\left(\frac{y}{A}\right) = \sqrt{\frac{k}{m}} t + C$

109. $y = 2x - \cosh \sqrt{x}$

$$y' = 2 - \frac{1}{2\sqrt{x}} (\sinh \sqrt{x}) = 2 - \frac{\sinh \sqrt{x}}{2\sqrt{x}}$$

Since $y = 0$ when $t = 0$, you have $C = 0$. Thus,

$$\sin\left(\sqrt{\frac{k}{m}} t\right) = \frac{y}{A}$$

$$y = A \sin\left(\sqrt{\frac{k}{m}} t\right)$$

111. Let $u = x^2$, $du = 2x dx$.

$$\int \frac{x}{\sqrt{x^4-1}} dx = \frac{1}{2} \int \frac{1}{\sqrt{(x^2)^2-1}} (2x) dx = \frac{1}{2} \ln(x^2 + \sqrt{x^4-1}) + C$$

Problem Solving for Chapter 5

1. $\tan \theta_1 = \frac{3}{x}$

$$\tan \theta_2 = \frac{6}{10-x}$$

Minimize $\theta_1 + \theta_2$:

$$f(x) = \theta_1 + \theta_2 = \arctan\left(\frac{3}{x}\right) + \arctan\left(\frac{6}{10-x}\right)$$

$$f'(x) = \frac{1}{1 + \frac{9}{x^2}} \left(\frac{-3}{x^2}\right) + \frac{1}{1 + \frac{36}{(10-x)^2}} \left(\frac{6}{(10-x)^2}\right) = 0$$

$$\frac{3}{x^2 + 9} = \frac{6}{(10-x)^2 + 36}$$

$$(10-x)^2 + 36 = 2(x^2 + 9)$$

$$100 - 20x + x^2 + 36 = 2x^2 + 18$$

$$x^2 + 20x - 118 = 0$$

$$x = \frac{-20 \pm \sqrt{20^2 - 4(-118)}}{2} = -10 \pm \sqrt{218}$$


$$a = -10 + \sqrt{218} \approx 4.7648 \quad f(a) \approx 1.4153$$

$$\theta = \pi - (\theta_1 + \theta_2) \approx 1.7263 \quad \text{or} \quad 98.9^\circ$$

Endpoints: $a = 0$: $\theta \approx 1.0304$

$$a = 10$$
: $\theta \approx 1.2793$

Maximum is 1.7263 at $a = -10 + \sqrt{218} \approx 4.7648$.



3. $f(x) = \sin(\ln x)$

(a) Domain: $x > 0$ or $(0, \infty)$

(b) $f(x) = 1 = \sin(\ln x) \Rightarrow \ln x = \frac{\pi}{2} + 2k\pi$

Two values are $x = e^{\pi/2}, e^{(\pi/2)+2\pi}$.

(c) $f(x) = -1 = \sin(\ln x) \Rightarrow \ln x = \frac{3\pi}{2} + 2k\pi$

Two values are $x = e^{-\pi/2}, e^{3\pi/2}$.


(d) Since the range of the sine function is $[-1, 1]$, parts (b) and (c) show that the range of f is $[-1, 1]$.

(e) $f'(x) = \frac{1}{x} \cos(\ln x)$

$$f'(x) = 0 \Rightarrow \cos(\ln x) = 0 \Rightarrow \ln x = \frac{\pi}{2} + k\pi \Rightarrow$$

$$x = e^{\pi/2} \text{ on } [1, 10]$$

$$\left. \begin{array}{l} f(e^{\pi/2}) = 1 \\ f(1) = 0 \\ f(10) \approx 0.7440 \end{array} \right\} \text{Maximum is 1 at } x = e^{\pi/2} \approx 4.8105$$



$\lim_{x \rightarrow 0^+} f(x)$ seems to be $-\frac{1}{2}$. (This is incorrect.)

(g) For the points $x = e^{\pi/2}, e^{-3\pi/2}, e^{-7\pi/2}, \dots$

we have $f(x) = 1$.

For the points $x = e^{-\pi/2}, e^{-5\pi/2}, e^{-9\pi/2}, \dots$

we have $f(x) = -1$.

That is, as $x \rightarrow 0^+$, there is an infinite number of points where $f(x) = 1$, and an infinite number where $f(x) = -1$. Thus $\lim_{x \rightarrow 0^+} \sin(\ln x)$ does not exist.

You can verify this by graphing $f(x)$ on small intervals close to the origin.

5. (a) $\frac{\text{Area sector}}{\text{Area circle}} = \frac{t}{2\pi} \Rightarrow \text{Area sector} = \frac{t}{2\pi}(\pi) = \frac{t}{2}$

(b) $\text{Area } AOP = \frac{1}{2}(\text{base})(\text{height}) - \int_1^{\cosh t} \sqrt{x^2 - 1} dx$

$$A(t) = \frac{1}{2} \cosh t \cdot \sinh t - \int_1^{\cosh t} \sqrt{x^2 - 1} dx$$

$$\begin{aligned} A'(t) &= \frac{1}{2}[\cosh^2 t + \sinh^2 t] - \sqrt{\cosh^2 t - 1} \sinh t \\ &= \frac{1}{2}[\cosh^2 t + \sinh^2 t] - \sinh^2 t \\ &= \frac{1}{2}[\cosh^2 t - \sinh^2 t] = \frac{1}{2} \end{aligned}$$

$A(t) = \frac{1}{2}t + C$. But, $A(0) = C = 0 \Rightarrow C = 0$

Thus, $A(t) = \frac{1}{2}t$ or $t = 2A(t)$.

7. $y = \ln x$

$$y' = \frac{1}{x}$$

$$y - b = \frac{1}{a}(x - a)$$

$$y = \frac{1}{a}x + b - 1 \quad \text{Tangent line}$$

If $x = 0, c = b - 1$. Thus, $b - c = b - (b - 1) = 1$.

9. Let $u = 1 + \sqrt{x}, \sqrt{x} = u - 1, x = u^2 - 2u + 1,$
 $dx = (2u - 2)du.$

$$\begin{aligned} \text{Area} &= \int_1^4 \frac{1}{\sqrt{x} + x} dx = \int_2^3 \frac{2u - 2}{(u - 1) + (u^2 - 2u + 1)} du \\ &= \int_2^3 \frac{2(u - 1)}{u^2 - u} du \\ &= \int_2^3 \frac{2}{u} du \\ &= \left[2 \ln u \right]_2^3 \\ &= 2 \ln 3 - 2 \ln 2 = 2 \ln \left(\frac{3}{2} \right) \\ &\approx 0.8109 \end{aligned}$$

11. (a) $\frac{dy}{dt} = y^{1.01}$

$$\int y^{-1.01} dy = \int dt$$

$$\frac{y^{-0.01}}{-0.01} = t + C_1$$

$$\frac{1}{y^{0.01}} = -0.01t + C$$

$$y^{0.01} = \frac{1}{C - 0.01t}$$

$$y = \frac{1}{(C - 0.01t)^{100}}$$

$y(0) = 1: 1 = \frac{1}{C^{100}} \Rightarrow C = 1$

Hence, $y = \frac{1}{(1 - 0.01t)^{100}}$.

For $T = 100, \lim_{t \rightarrow T^-} y = \infty$.

(b) $\int y^{-(1+\varepsilon)} dy = \int k dt$

$$\frac{y^{-\varepsilon}}{-\varepsilon} = kt + C_1$$

$$y^{-\varepsilon} = -\varepsilon kt + C$$

$$y = \frac{1}{(C - \varepsilon kt)^{1/\varepsilon}}$$

$y(0) = y_0 = \frac{1}{C^{1/\varepsilon}} \Rightarrow C^{1/\varepsilon} = \frac{1}{y_0} \Rightarrow C = \left(\frac{1}{y_0} \right)^\varepsilon$

Hence, $y = \frac{1}{\left(\frac{1}{y_0^\varepsilon} - \varepsilon kt \right)^{1/\varepsilon}}$.

For $t \rightarrow \frac{1}{y_0^\varepsilon \varepsilon k}, y \rightarrow \infty$.

13. Since $\frac{dy}{dt} = k(y - 20)$,

$$\int \frac{1}{y - 20} dy = \int k dt$$

$$\ln(y - 20) = kt + C$$

$$y = Ce^{kt} + 20.$$

When $t = 0$, $y = 72$. Therefore, $C = 52$.

When $t = 1$, $y = 48$. Therefore, $48 = 52e^k + 20$, $e^k = (28/52) = (7/13)$, and $k = \ln(7/13)$.

Thus, $y = 52e^{\ln(7/13)t} + 20$.

When $t = 5$, $y = 52e^{5 \ln(7/13)} + 20 \approx 22.35^\circ$.

15. (a) $\frac{dS}{dt} = k_1 S(L - S)$

$S = \frac{L}{1 + Ce^{-kt}}$ is a solution because

$$\frac{dS}{dt} = -L(1 + Ce^{-kt})^{-2}(-Cke^{-kt})$$

$$= \frac{LCke^{-kt}}{(1 + Ce^{-kt})^2}$$

$$= \left(\frac{k}{L}\right) \frac{L}{1 + Ce^{-kt}} \cdot \frac{CLe^{-kt}}{1 + Ce^{-kt}}$$

$$= \left(\frac{k}{L}\right) \frac{L}{1 + Ce^{-kt}} \cdot \left(L - \frac{L}{1 + Ce^{-kt}}\right)$$


$$= k_1 S(L - S), \text{ where } k_1 = \frac{k}{L}.$$

$L = 100$. Also, $S = 10$ when $t = 0 \Rightarrow C = 9$. And,

$S = 20$ when $t = 1 \Rightarrow k = -\ln(4/9)$.

$$\text{Particular Solution. } S = \frac{100}{1 + 9e^{\ln(4/9)t}}$$

$$= \frac{100}{1 + 9e^{-0.8109t}}$$



(b) $\frac{dS}{dt} = \ln\left(\frac{4}{9}\right) S(100 - S)$

$$\frac{d^2S}{dt^2} = \ln\left(\frac{4}{9}\right) \left[S\left(-\frac{dS}{dt}\right) + (100 - S) \frac{dS}{dt} \right]$$

$$= \ln\left(\frac{4}{9}\right) (100 - 2S) \frac{dS}{dt}$$

$$= 0 \text{ when } S = 50 \text{ or } \frac{dS}{dt} = 0.$$


Choosing $S = 50$, we have:

$$50 = \frac{100}{1 + 9e^{\ln(4/9)t}}$$

$$2 = 1 + 9e^{\ln(4/9)t}$$

$$\frac{\ln(1/9)}{\ln(4/9)} = t$$

$$t \approx 2.7 \text{ months}$$



(e) Sales will decrease toward the line $S = L$.