


## Chapter Two

### Section 2.1


1(a).



(b). Based on the direction field, all solutions seem to converge to a specific increasing function.

(c). The integrating factor is  $\mu(t) = e^{3t}$ , and hence  $y(t) = t/3 - 1/9 + e^{-2t} + c e^{-3t}$ . It follows that all solutions converge to the function  $y_1(t) = t/3 - 1/9$ .


2(a).



(b). All slopes *eventually* become positive, hence all solutions will increase without bound.

(c). The integrating factor is  $\mu(t) = e^{-2t}$ , and hence  $y(t) = t^3 e^{2t}/3 + c e^{2t}$ . It is evident that all solutions increase at an exponential rate.


3(a)



(b). All solutions seem to converge to the function  $y_0(t) = 1$ .

(c). The integrating factor is  $\mu(t) = e^{2t}$ , and hence  $y(t) = t^2 e^{-t}/2 + 1 + c e^{-t}$ . It is clear that all solutions converge to the specific solution  $y_0(t) = 1$ .

4(a).




(b). Based on the direction field, the solutions eventually become oscillatory.

(c). The integrating factor is  $\mu(t) = t$ , and hence the general solution is

$$y(t) = \frac{3\cos(2t)}{4t} + \frac{3}{2}\sin(2t) + \frac{c}{t}$$

in which  $c$  is an arbitrary constant. As  $t$  becomes large, all solutions converge to the function  $y_1(t) = 3\sin(2t)/2$ .


5(a).



(b). All slopes *eventually* become positive, hence all solutions will increase without bound.

(c). The integrating factor is  $\mu(t) = \exp(-\int 2dt) = e^{-2t}$ . The differential equation can be written as  $e^{-2t}y' - 2e^{-2t}y = 3e^{-t}$ , that is,  $(e^{-2t}y)' = 3e^{-t}$ . Integration of both sides of the equation results in the general solution  $y(t) = -3e^t + ce^{2t}$ . It follows that all solutions will increase exponentially.

6(a)




(b). All solutions seem to converge to the function  $y_0(t) = 0$ .

(c). The integrating factor is  $\mu(t) = t^2$ , and hence the general solution is

$$y(t) = -\frac{\cos(t)}{t} + \frac{\sin(2t)}{t^2} + \frac{c}{t^2}$$

in which  $c$  is an arbitrary constant. As  $t$  becomes large, all solutions converge to the function  $y_0(t) = 0$ .


7(a).



(b). All solutions seem to converge to the function  $y_0(t) = 0$ .

(c). The integrating factor is  $\mu(t) = \exp(t^2)$ , and hence  $y(t) = t^2 e^{-t^2} + c e^{-t^2}$ . It is clear that all solutions converge to the function  $y_0(t) = 0$ .


8(a)



(b). All solutions seem to converge to the function  $y_0(t) = 0$ .

(c). Since  $\mu(t) = (1 + t^2)^2$ , the general solution is  $y(t) = [\tan^{-1}(t) + C]/(1 + t^2)^2$ . It follows that all solutions converge to the function  $y_0(t) = 0$ .


9(a).



(b). All slopes *eventually* become positive, hence all solutions will increase without bound.

(c). The integrating factor is  $\mu(t) = \exp(\int \frac{1}{2} dt) = e^{t/2}$ . The differential equation can be written as  $e^{t/2}y' + e^{t/2}y/2 = 3t e^{t/2}/2$ , that is,  $(e^{t/2}y/2)' = 3t e^{t/2}/2$ . Integration of both sides of the equation results in the general solution  $y(t) = 3t - 6 + c e^{-t/2}$ . All solutions approach the specific solution  $y_0(t) = 3t - 6$ .


10(a).



(b). For  $y > 0$ , the slopes are *all* positive, and hence the corresponding solutions increase without bound. For  $y < 0$ , almost all solutions have negative slopes, and hence solutions tend to decrease without bound.

(c). First divide both sides of the equation by  $t$ . From the resulting *standard form*, the integrating factor is  $\mu(t) = \exp(-\int \frac{1}{t} dt) = 1/t$ . The differential equation can be written as  $y'/t - y/t^2 = t e^{-t}$ , that is,  $(y/t)' = t e^{-t}$ . Integration leads to the general solution  $y(t) = -t e^{-t} + c t$ . For  $c \neq 0$ , solutions *diverge*, as implied by the direction field. For the case  $c = 0$ , the specific solution is  $y(t) = -t e^{-t}$ , which evidently approaches *zero* as  $t \rightarrow \infty$ .


11(a).



(b). The solutions appear to be oscillatory.

(c). The integrating factor is  $\mu(t) = e^t$ , and hence  $y(t) = \sin(2t) - 2 \cos(2t) + c e^{-t}$ . It is evident that all solutions converge to the specific solution  $y_0(t) = \sin(2t) - 2 \cos(2t)$ .

12(a).



(b). All solutions *eventually* have positive slopes, and hence increase without bound.

(c). The integrating factor is  $\mu(t) = e^{2t}$ . The differential equation can be written as  $e^{t/2}y' + e^{t/2}y/2 = 3t^2/2$ , that is,  $(e^{t/2}y/2)' = 3t^2/2$ . Integration of both sides of the equation results in the general solution  $y(t) = 3t^2 - 12t + 24 + c e^{-t/2}$ . It follows that all solutions converge to the specific solution  $y_0(t) = 3t^2 - 12t + 24$ .


14. The integrating factor is  $\mu(t) = e^{2t}$ . After multiplying both sides by  $\mu(t)$ , the equation can be written as  $(e^{2t}y)' = t$ . Integrating both sides of the equation results in the general solution  $y(t) = t^2 e^{-2t}/2 + c e^{-2t}$ . Invoking the specified condition, we require that  $e^{-2}/2 + c e^{-2} = 0$ . Hence  $c = -1/2$ , and the solution to the initial value problem is  $y(t) = (t^2 - 1)e^{-2t}/2$ .

16. The integrating factor is  $\mu(t) = \exp(\int \frac{2}{t} dt) = t^2$ . Multiplying both sides by  $\mu(t)$ , the equation can be written as  $(t^2 y)' = \cos(t)$ . Integrating both sides of the equation results in the general solution  $y(t) = \sin(t)/t^2 + c t^{-2}$ . Substituting  $t = \pi$  and setting the value equal to *zero* gives  $c = 0$ . Hence the specific solution is  $y(t) = \sin(t)/t^2$ .

17. The integrating factor is  $\mu(t) = e^{-2t}$ , and the differential equation can be written as  $(e^{-2t}y)' = 1$ . Integrating, we obtain  $e^{-2t}y(t) = t + c$ . Invoking the specified initial condition results in the solution  $y(t) = (t + 2)e^{2t}$ .

19. After writing the equation in *standard form*, we find that the integrating factor is  $\mu(t) = \exp(\int \frac{4}{t} dt) = t^4$ . Multiplying both sides by  $\mu(t)$ , the equation can be written as  $(t^4 y)' = t e^{-t}$ . Integrating both sides results in  $t^4 y(t) = -(t + 1)e^{-t} + c$ . Letting  $t = -1$  and setting the value equal to *zero* gives  $c = 0$ . Hence the specific solution of the initial value problem is  $y(t) = -(t^{-3} + t^{-4})e^{-t}$ .

21(a).




The solutions appear to diverge from an *apparent* oscillatory solution. From the direction field, the critical value of the initial condition seems to be  $a_0 = -1$ . For  $a > -1$ , the solutions increase without bound. For  $a < -1$ , solutions decrease without bound.

(b). The integrating factor is  $\mu(t) = e^{-t/2}$ . The general solution of the differential equation is  $y(t) = (8\sin(t) - 4\cos(t))/5 + c e^{t/2}$ . The solution is sinusoidal as long as  $c = 0$ . The *initial value* of this sinusoidal solution is  $a_0 = (8\sin(0) - 4\cos(0))/5 = -4/5$ .

(c). See part (b).

22(a).




All solutions appear to *eventually* increase without bound. The solutions *initially* increase or decrease, depending on the initial value  $a$ . The critical value seems to be  $a_0 = -1$ .

(b). The integrating factor is  $\mu(t) = e^{-t/2}$ , and the general solution of the differential equation is  $y(t) = -3e^{t/3} + c e^{t/2}$ . Invoking the initial condition  $y(0) = a$ , the solution may also be expressed as  $y(t) = -3e^{t/3} + (a + 3) e^{t/2}$ . Differentiating, follows that  $y'(0) = -1 + (a + 3)/2 = (a + 1)/2$ . The critical value is evidently  $a_0 = -1$ .

(c). For  $a_0 = -1$ , the solution is  $y(t) = -3e^{t/3} + 2e^{t/2}$ , which (for large  $t$ ) is dominated by the term containing  $e^{t/2}$ .

is  $y(t) = (8\sin(t) - 4\cos(t))/5 + ce^{t/2}$ .

23(a).




As  $t \rightarrow 0$ , solutions increase without bound if  $y(1) = a > .4$ , and solutions decrease without bound if  $y(1) = a < .4$ .

(b). The integrating factor is  $\mu(t) = \exp\left(\int \frac{t+1}{t} dt\right) = te^t$ . The general solution of the differential equation is  $y(t) = te^{-t} + ce^{-t}/t$ . Invoking the specified value  $y(1) = a$ , we have  $1 + c = ae$ . That is,  $c = ae - 1$ . Hence the solution can also be expressed as  $y(t) = te^{-t} + (ae - 1)e^{-t}/t$ . For *small* values of  $t$ , the second term is dominant. Setting  $ae - 1 = 0$ , critical value of the parameter is  $a_0 = 1/e$ .

(c). For  $a > 1/e$ , solutions increase without bound. For  $a < 1/e$ , solutions decrease without bound. When  $a = 1/e$ , the solution is  $y(t) = te^{-t}$ , which approaches 0 as  $t \rightarrow 0$ .

24(a).



As  $t \rightarrow 0$ , solutions increase without bound if  $y(1) = a > .4$ , and solutions decrease without bound if  $y(1) = a < .4$ .



(b). Given the initial condition,  $y(-\pi/2) = a$ , the solution is  $y(t) = (a\pi^2/4 - \cos t)/t$ .

Since  $\lim_{t \rightarrow 0} \cos t = 1$ , solutions increase without bound if  $a > 4/\pi^2$ , and solutions decrease without bound if  $a < 4/\pi^2$ . Hence the critical value is  $a_0 = 4/\pi^2 = 0.452847\dots$

(c). For  $a = 4/\pi^2$ , the solution is  $y(t) = (1 - \cos t)/t$ , and  $\lim_{t \rightarrow 0} y(t) = 1/2$ . Hence the solution is bounded.

25. The integrating factor is  $\mu(t) = \exp(\int \frac{1}{2} dt) = e^{t/2}$ . Therefore general solution is  $y(t) = [4\cos(t) + 8\sin(t)]/5 + c e^{-t/2}$ . Invoking the initial condition, the specific solution is  $y(t) = [4\cos(t) + 8\sin(t) - 9 e^{t/2}]/5$ . Differentiating, it follows that

$$\begin{aligned} y'(t) &= [-4\sin(t) + 8\cos(t) + 4.5 e^{-t/2}]/5 \\ y''(t) &= [-4\cos(t) - 8\sin(t) - 2.25 e^{-t/2}]/5 \end{aligned}$$

Setting  $y'(t) = 0$ , the first solution is  $t_1 = 1.3643$ , which gives the location of the *first* stationary point. Since  $y''(t_1) < 0$ , the first stationary point is a local *maximum*. The coordinates of the point are  $(1.3643, .82008)$ .

26. The integrating factor is  $\mu(t) = \exp(\int \frac{2}{3} dt) = e^{2t/3}$ , and the differential equation can

be written as  $(e^{2t/3} y)' = e^{2t/3} - t e^{2t/3}/2$ . The general solution is  $y(t) = (21 - 6t)/8 + c e^{-2t/3}$ . Imposing the initial condition, we have  $y(t) = (21 - 6t)/8 + (y_0 - 21/8)e^{-2t/3}$ . Since the solution is smooth, the desired intersection will be a point of tangency. Taking the derivative,  $y'(t) = -3/4 - (2y_0 - 21/4)e^{-2t/3}/3$ . Setting  $y'(t) = 0$ , the solution is  $t_1 = \frac{3}{2} \ln[(21 - 8y_0)/9]$ . Substituting into the solution, the respective *value* at the stationary point is  $y(t_1) = \frac{3}{2} + \frac{9}{4} \ln 3 - \frac{9}{8} \ln(21 - 8y_0)$ . Setting this result equal to *zero*, we obtain the required initial value  $y_0 = (21 - 9 e^{4/3})/8 = -1.643$ .

27. The integrating factor is  $\mu(t) = e^{t/4}$ , and the differential equation can be written as  $(e^{t/4} y)' = 3 e^{t/4} + 2 e^{t/4} \cos(2t)$ . The general solution is

$$y(t) = 12 + [8\cos(2t) + 64\sin(2t)]/65 + c e^{-t/4}.$$

Invoking the initial condition,  $y(0) = 0$ , the specific solution is

$$y(t) = 12 + [8\cos(2t) + 64\sin(2t) - 788 e^{-t/4}]/65.$$

As  $t \rightarrow \infty$ , the exponential term will decay, and the solution will oscillate about an *average value* of 12, with an *amplitude* of  $8/\sqrt{65}$ .

29. The integrating factor is  $\mu(t) = e^{-3t/2}$ , and the differential equation can be written as  $(e^{-3t/2} y)' = 3t e^{-3t/2} + 2 e^{-t/2}$ . The general solution is  $y(t) = -2t - 4/3 - 4e^t + c e^{3t/2}$ . Imposing the initial condition,  $y(t) = -2t - 4/3 - 4e^t + (y_0 + 16/3) e^{3t/2}$ . As  $t \rightarrow \infty$ , the term containing  $e^{3t/2}$  will *dominate* the solution. Its *sign* will determine the divergence properties. Hence the critical value of the initial condition is

$$y_0 = -16/3.$$

The corresponding solution,  $y(t) = -2t - 4/3 - 4e^t$ , will also decrease without bound.

Note on Problems 31-34 :

Let  $g(t)$  be *given*, and consider the function  $y(t) = y_1(t) + g(t)$ , in which  $y_1(t) \rightarrow \infty$  as  $t \rightarrow \infty$ . Differentiating,  $y'(t) = y_1'(t) + g'(t)$ . Letting  $a$  be a *constant*, it follows that  $y'(t) + ay(t) = y_1'(t) + ay_1(t) + g'(t) + ag(t)$ . Note that the hypothesis on the function  $y_1(t)$  will be satisfied, if  $y_1'(t) + ay_1(t) = 0$ . That is,  $y_1(t) = c e^{-at}$ . Hence  $y(t) = c e^{-at} + g(t)$ , which is a solution of the equation  $y' + ay = g'(t) + ag(t)$ . For convenience, choose  $a = 1$ .

31. Here  $g(t) = 3$ , and we consider the linear equation  $y' + y = 3$ . The integrating factor is  $\mu(t) = e^t$ , and the differential equation can be written as  $(e^t y)' = 3e^t$ . The general solution is  $y(t) = 3 + c e^{-t}$ .

33.  $g(t) = 3 - t$ . Consider the linear equation  $y' + y = -1 + 3 - t$ . The integrating factor is  $\mu(t) = e^t$ , and the differential equation can be written as  $(e^t y)' = (2 - t)e^t$ . The general solution is  $y(t) = 3 - t + c e^{-t}$ .

34.  $g(t) = 4 - t^2$ . Consider the linear equation  $y' + y = 4 - 2t - t^2$ . The integrating factor is  $\mu(t) = e^t$ , and the equation can be written as  $(e^t y)' = (4 - 2t - t^2)e^t$ . The general solution is  $y(t) = 4 - t^2 + c e^{-t}$ .

**Section 2.2**

2. For  $x \neq -1$ , the differential equation may be written as  $y dy = [x^2/(1+x^3)]dx$ . Integrating both sides, with respect to the appropriate variables, we obtain the relation

$$y^2/2 = \frac{1}{3} \ln|1+x^3| + c. \text{ That is, } y(x) = \pm \sqrt{\frac{2}{3} \ln|1+x^3| + c}.$$

3. The differential equation may be written as  $y^{-2}dy = -\sin x dx$ . Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation  $-y^{-1} = \cos x + c$ . That is,  $(C - \cos x)y = 1$ , in which  $C$  is an arbitrary constant. Solving for the dependent variable, explicitly,  $y(x) = 1/(C - \cos x)$ .

5. Write the differential equation as  $\cos^{-2} 2y dy = \cos^2 x dx$ , or  $\sec^2 2y dy = \cos^2 x dx$ . Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation  $\tan 2y = \sin x \cos x + x + c$ .


7. The differential equation may be written as  $(y + e^y)dy = (x - e^{-x})dx$ . Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation

$$y^2 + 2e^y = x^2 + 2e^{-x} + c.$$

8. Write the differential equation as  $(1+y^2)dy = x^2 dx$ . Integrating both sides of the equation, we obtain the relation  $y + y^3/3 = x^3/3 + c$ , that is,  $3y + y^3 = x^3 + C$ .

9(a). The differential equation is separable, with  $y^{-2}dy = (1 - 2x)dx$ . Integration yields  $-y^{-1} = x - x^2 + c$ . Substituting  $x = 0$  and  $y = -1/6$ , we find that  $c = 6$ . Hence the specific solution is  $y^{-1} = x^2 - x - 6$ . The *explicit form* is  $y(x) = 1/(x^2 - x - 6)$ .


(b)



(c). Note that  $x^2 - x - 6 = (x + 2)(x - 3)$ . Hence the solution becomes *singular* at  $x = -2$  and  $x = 3$ .


10(a).  $y(x) = -\sqrt{2x - 2x^2 + 4}$ .

10(b).



11(a). Rewrite the differential equation as  $x e^x dx = -y dy$ . Integrating both sides of the equation results in  $x e^x - e^x = -y^2/2 + c$ . Invoking the initial condition, we obtain  $c = -1/2$ . Hence  $y^2 = 2e^x - 2x e^x - 1$ . The *explicit form* of the solution is  $y(x) = \sqrt{2e^x - 2x e^x - 1}$ . The *positive sign* is chosen, since  $y(0) = 1$ .


(b).



(c). The function under the radical becomes *negative* near  $x = -1.7$  and  $x = 0.76$ .

11(a). Write the differential equation as  $r^{-2} dr = \theta^{-1} d\theta$ . Integrating both sides of the equation results in the relation  $-r^{-1} = \ln \theta + c$ . Imposing the condition  $r(1) = 2$ , we obtain  $c = -1/2$ . The *explicit form* of the solution is  $r(\theta) = 2/(1 - 2 \ln \theta)$ .


(b).



(c). Clearly, the solution makes sense only if  $\theta > 0$ . Furthermore, the solution becomes singular when  $\ln \theta = 1/2$ , that is,  $\theta = \sqrt{e}$ .


13(a).  $y(x) = -\sqrt{2\ln(1+x^2)+4}$ .

(b).



14(a). Write the differential equation as  $y^{-3}dy = x(1+x^2)^{-1/2} dx$ . Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation  $-y^{-2}/2 = \sqrt{1+x^2} + c$ . Imposing the initial condition, we obtain  $c = -3/2$ . Hence the specific solution can be expressed as  $y^{-2} = 3 - 2\sqrt{1+x^2}$ . The *explicit form* of the solution is  $y(x) = 1/\sqrt{3 - 2\sqrt{1+x^2}}$ . The *positive* sign is chosen to satisfy the initial condition.


(b).



(c). The solution becomes singular when  $2\sqrt{1+x^2} = 3$ . That is, at  $x = \pm\sqrt{5}/2$ .


15(a).  $y(x) = -1/2 + \sqrt{x^2 - 15/4}$ .

(b).



16(a). Rewrite the differential equation as  $4y^3 dy = x(x^2 + 1)dx$ . Integrating both sides of the equation results in  $y^4 = (x^2 + 1)^2/4 + c$ . Imposing the initial condition, we obtain  $c = 0$ . Hence the solution may be expressed as  $(x^2 + 1)^2 - 4y^4 = 0$ . The *explicit* form of the solution is  $y(x) = -\sqrt{(x^2 + 1)/2}$ . The *sign* is chosen based on  $y(0) = -1/\sqrt{2}$ .


(b).



(c). The solution is valid for all  $x \in \mathbb{R}$ .

17(a).  $y(x) = -5/2 - \sqrt{x^3 - e^x + 13/4}$ .


(b).



(c). The solution is valid for  $x > -1.45$ . This value is found by estimating the root of  $4x^3 - 4e^x + 13 = 0$ .

18(a). Write the differential equation as  $(3 + 4y)dy = (e^{-x} - e^x)dx$ . Integrating both sides of the equation, with respect to the appropriate variables, we obtain the relation  $3y + 2y^2 = -(e^x + e^{-x}) + c$ . Imposing the initial condition,  $y(0) = 1$ , we obtain  $c = 7$ . Thus, the solution can be expressed as  $3y + 2y^2 = -(e^x + e^{-x}) + 7$ . Now by *completing the square* on the left hand side,  $2(y + 3/4)^2 = -(e^x + e^{-x}) + 65/8$ . Hence the *explicit* form of the solution is  $y(x) = -3/4 + \sqrt{65/16 - \cosh x}$ .


(b).



(c). Note the  $65 - 16 \cosh x \geq 0$ , as long as  $|x| > 2.1$ . Hence the solution is valid on the interval  $-2.1 < x < 2.1$ .

19(a).  $y(x) = -\pi/3 + \frac{1}{3} \sin^{-1}(3 \cos^2 x)$ .


(b).



20(a). Rewrite the differential equation as  $y^2 dy = \arcsin x / \sqrt{1 - x^2} dx$ . Integrating both sides of the equation results in  $y^3/3 = (\arcsin x)^2/2 + c$ . Imposing the condition  $y(0) = 0$ , we obtain  $c = 0$ . The *explicit* form of the solution is  $y(x) = \sqrt[3]{\frac{3}{2} (\arcsin x)^2}$ .




(b).



(c). Evidently, the solution is defined for  $-1 \leq x \leq 1$ .

22. The differential equation can be written as  $(3y^2 - 4)dy = 3x^2dx$ . Integrating both sides, we obtain  $y^3 - 4y = x^3 + c$ . Imposing the initial condition, the specific solution is  $y^3 - 4y = x^3 - 1$ . Referring back to the differential equation, we find that  $y' \rightarrow \infty$  as  $y \rightarrow \pm 2/\sqrt{3}$ . The respective values of the abscissas are  $x = -1.276, 1.598$ .



Hence the solution is valid for  $-1.276 < x < 1.598$ .

24. Write the differential equation as  $(3 + 2y)dy = (2 - e^x)dx$ . Integrating both sides, we obtain  $3y + y^2 = 2x - e^x + c$ . Based on the specified initial condition, the solution can be written as  $3y + y^2 = 2x - e^x + 1$ . *Completing the square*, it follows that  $y(x) = -3/2 + \sqrt{2x - e^x + 13/4}$ . The solution is defined if  $2x - e^x + 13/4 \geq 0$ , that is,  $-1.5 \leq x \leq 2$  (*approximately*). In that interval,  $y' = 0$ , for  $x = \ln 2$ . It can be verified that  $y''(\ln 2) < 0$ . In fact,  $y''(x) < 0$  on the interval of definition. Hence the solution attains a global maximum at  $x = \ln 2$ .


26. The differential equation can be written as  $(1 + y^2)^{-1}dy = 2(1 + x)dx$ . Integrating both sides of the equation, we obtain  $\arctan y = 2x + x^2 + c$ . Imposing the given initial condition, the specific solution is  $\arctan y = 2x + x^2$ . Therefore,  $y(x) = \tan(2x + x^2)$ . Observe that the solution is defined as long as  $-\pi/2 < 2x + x^2 < \pi/2$ . It is easy to see that  $2x + x^2 \geq -1$ . Furthermore,  $2x + x^2 = \pi/2$  for  $x = -2.6$  and  $0.6$ . Hence the solution is valid on the interval  $-2.6 < x < 0.6$ . Referring back to the differential

equation, the solution is *stationary* at  $x = -1$ . Since  $y''(x) > 0$  on the entire interval of definition, the solution attains a global minimum at  $x = -1$ .

28(a). Write the differential equation as  $y^{-1}(4 - y)^{-1}dy = t(1 + t)^{-1}dt$ . Integrating both sides of the equation, we obtain  $\ln|y| - \ln|y - 4| = 4t - 4\ln|1 + t| + c$ . Taking the *exponential* of both sides, it follows that  $|y/(y - 4)| = C e^{4t}/(1 + t)^4$ . It follows that as  $t \rightarrow \infty$ ,  $|y/(y - 4)| = |1 + 4/(y - 4)| \rightarrow \infty$ . That is,  $y(t) \rightarrow 4$ .


(b). Setting  $y(0) = 2$ , we obtain that  $C = 1$ . Based on the initial condition, the solution may be expressed as  $y/(y - 4) = -e^{4t}/(1 + t)^4$ . Note that  $y/(y - 4) < 0$ , for all  $t \geq 0$ . Hence  $y < 4$  for all  $t \geq 0$ . Referring back to the differential equation, it follows that  $y'$  is always *positive*. This means that the solution is *monotone increasing*. We find that the root of the equation  $e^{4t}/(1 + t)^4 = 399$  is near  $t = 2.844$ .

(c). Note the  $y(t) = 4$  is an equilibrium solution. Examining the local direction field,




we see that if  $y(0) > 0$ , then the corresponding solutions converge to  $y = 4$ . Referring back to part (a), we have  $y/(y - 4) = [y_0/(y_0 - 4)]e^{4t}/(1 + t)^4$ , for  $y_0 \neq 4$ . Setting  $t = 2$ , we obtain  $y_0/(y_0 - 4) = (3/e^2)^4 y(2)/(y(2) - 4)$ . Now since the function  $f(y) = y/(y - 4)$  is *monotone* for  $y < 4$  and  $y > 4$ , we need only solve the equations  $y_0/(y_0 - 4) = -399(3/e^2)^4$  and  $y_0/(y_0 - 4) = 401(3/e^2)^4$ . The respective solutions are  $y_0 = 3.6622$  and  $y_0 = 4.4042$ .

30(f).




31(c)




32(a). Observe that  $(x^2 + 3y^2)/2xy = \frac{1}{2}\left(\frac{y}{x}\right)^{-1} + \frac{3}{2}\frac{y}{x}$ . Hence the differential equation is *homogeneous*.

(b). The substitution  $y = xv$  results in  $v + xv' = (x^2 + 3x^2v^2)/2x^2v$ . The transformed equation is  $v' = (1 + v^2)/2xv$ . This equation is *separable*, with general solution  $v^2 + 1 = cx$ . In terms of the original dependent variable, the solution is  $x^2 + y^2 = cx^3$ .

(c).




33(c).




34(a). Observe that  $-(4x + 3y)/(2x + y) = -2 - \frac{y}{x} \left[2 + \frac{y}{x}\right]^{-1}$ . Hence the differential equation is *homogeneous*.

(b). The substitution  $y = xv$  results in  $v + xv' = -2 - v/(2 + v)$ . The transformed equation is  $v' = -(v^2 + 5v + 4)/(2 + v)x$ . This equation is *separable*, with general solution  $(v+4)^2|v+1| = C/x^3$ . In terms of the original dependent variable, the solution is  $(4x + y)^2|x+y| = C$ .

(c).




35(c).



36(a). Divide by  $x^2$  to see that the equation is homogeneous. Substituting  $y = xv$ , we obtain  $xv' = (1 + v)^2$ . The resulting differential equation is separable.

(b). Write the equation as  $(1 + v)^{-2}dv = x^{-1}dx$ . Integrating both sides of the equation, we obtain the general solution  $-1/(1 + v) = \ln|x| + c$ . In terms of the original dependent variable, the solution is  $y = x [C - \ln|x|]^{-1} - x$ .

(c).




37(a). The differential equation can be expressed as  $y' = \frac{1}{2} \left(\frac{y}{x}\right)^{-1} - \frac{3}{2} \frac{y}{x}$ . Hence the equation is homogeneous. The substitution  $y = xv$  results in  $xv' = (1 - 5v^2)/2v$ . Separating variables, we have  $\frac{2v}{1-5v^2} dv = \frac{1}{x} dx$ .

(b). Integrating both sides of the transformed equation yields  $-\frac{1}{5}$

$$\ln|1 - 5v^2| = \ln|x| + c,$$

that is,  $1 - 5v^2 = C/|x|^5$ . In terms of the original dependent variable, the general solution is  $5y^2 = x^2 - C/|x|^3$ .


(c).



38(a). The differential equation can be expressed as  $y' = \frac{3}{2} \frac{y}{x} - \frac{1}{2} \left(\frac{y}{x}\right)^{-1}$ . Hence the equation is homogeneous. The substitution  $y = xv$  results in  $xv' = (v^2 - 1)/2v$ , that is,  $\frac{2v}{v^2-1} dv = \frac{1}{x} dx$ .

(b). Integrating both sides of the transformed equation yields  $\ln|v^2 - 1| = \ln|x| + c$ , that is,  $v^2 - 1 = C|x|$ . In terms of the original dependent variable, the general solution is  $y^2 = Cx^2|x| + x^2$ .

(c).




**Section 2.3**

5(a). Let  $Q$  be the amount of salt in the tank. Salt enters the tank of water at a rate of  $2\frac{1}{4}(1 + \frac{1}{2}\sin t) = \frac{1}{2} + \frac{1}{4}\sin t$  oz/min. It leaves the tank at a rate of  $2Q/100$  oz/min. Hence the differential equation governing the amount of salt at any time is

$$\frac{dQ}{dt} = \frac{1}{2} + \frac{1}{4}\sin t - Q/50.$$

The initial amount of salt is  $Q_0 = 50$  oz. The governing ODE is *linear*, with integrating factor  $\mu(t) = e^{t/50}$ . Write the equation as  $(e^{t/50}Q)' = e^{t/50}(\frac{1}{2} + \frac{1}{4}\sin t)$ . The specific solution is  $Q(t) = 25 + [12.5\sin t - 625\cos t + 63150 e^{-t/50}]/2501$  oz.

(b).



(c). The amount of salt approaches a *steady state*, which is an oscillation of amplitude  $1/4$  about a level of  $25$  oz.

6(a). The equation governing the value of the investment is  $dS/dt = rS$ . The value of the investment, at any time, is given by  $S(t) = S_0e^{rt}$ . Setting  $S(T) = 2S_0$ , the required time is  $T = \ln(2)/r$ .


(b). For the case  $r = 7\% = .07$ ,  $T \approx 9.9$  yrs.

(c). Referring to Part(a),  $r = \ln(2)/T$ . Setting  $T = 8$ , the required interest rate is to be approximately  $r = 8.66\%$ .

8(a). Based on the solution in Eq.(16), with  $S_0 = 0$ , the value of the investments *with* contributions is given by  $S(t) = 25,000(e^{rt} - 1)$ . After *ten* years, person A has  $S_A = \$25,000(1.226) = \$30,640$ . Beginning at age 35, the investments can now be analyzed using the equations  $S_A = 30,640 e^{.08t}$  and  $S_B = 25,000(e^{.08t} - 1)$ . After *thirty* years, the balances are  $S_A = \$337,734$  and  $S_B = \$250,579$ .

(b). For an *unspecified* rate  $r$ , the balances after *thirty* years are  $S_A = 30,640 e^{30r}$  and  $S_B = 25,000(e^{30r} - 1)$ .

(c).



(d). The two balances can *never* be equal.

11(a). Let  $S$  be the value of the mortgage. The debt accumulates at a rate of  $rS$ , in which  $r = .09$  is the *annual* interest rate. Monthly payments of \$ 800 are equivalent to \$ 9,600 *per year*. The differential equation governing the value of the mortgage is  $dS/dt = .09S - 9,600$ . Given that  $S_0$  is the original amount borrowed, the debt is  $S(t) = S_0e^{.09t} - 106,667(e^{.09t} - 1)$ . Setting  $S(30) = 0$ , it follows that  $S_0 = \$99,500$ .

(b). The *total* payment, over 30 years, becomes \$ 288,000. The interest paid on this purchase is \$ 188,500.

13(a). The balance *increases* at a rate of  $rS$  \$/yr, and *decreases* at a constant rate of  $k$  \$ *per year*. Hence the balance is modeled by the differential equation  $dS/dt = rS - k$ . The balance at any time is given by  $S(t) = S_0e^{rt} - \frac{k}{r}(e^{rt} - 1)$ .

(b). The solution may also be expressed as  $S(t) = (S_0 - \frac{k}{r})e^{rt} + \frac{k}{r}$ . Note that if the withdrawal rate is  $k_0 = rS_0$ , the balance will remain at a constant level  $S_0$ .

(c). Assuming that  $k > k_0$ ,  $S(T_0) = 0$  for  $T_0 = \frac{1}{r} \ln \left[ \frac{k}{k - k_0} \right]$ .

(d). If  $r = .08$  and  $k = 2k_0$ , then  $T_0 = 8.66$  *years*.

(e). Setting  $S(t) = 0$  and solving for  $e^{rt}$  in Part(b),  $e^{rt} = \frac{k}{k - rS_0}$ . Now setting  $t = T$  results in  $k = rS_0e^{rT} / (e^{rT} - 1)$ .

(f). In part(e), let  $k = 12,000$ ,  $r = .08$ , and  $T = 20$ . The required investment becomes  $S_0 = \$119,715$ .

14(a). Let  $Q' = -rQ$ . The general solution is  $Q(t) = Q_0e^{-rt}$ . Based on the definition of *half-life*, consider the equation  $Q_0/2 = Q_0e^{-5730r}$ . It follows that



$-5730r = \ln(1/2)$ , that is,  $r = 1.2097 \times 10^{-4}$  per year.

(b). Hence the amount of carbon-14 is given by  $Q(t) = Q_0 e^{-1.2097 \times 10^{-4}t}$ .

(c). Given that  $Q(T) = Q_0/5$ , we have the equation  $1/5 = e^{-1.2097 \times 10^{-4}T}$ . Solving for the *decay time*, the apparent age of the remains is approximately  $T = 13,304.65$  years.


15. Let  $P(t)$  be the population of mosquitoes at any time  $t$ . The rate of *increase* of the mosquito population is  $rP$ . The population *decreases* by 20,000 per day. Hence the equation that models the population is given by  $dP/dt = rP - 20,000$ . Note that the variable  $t$  represents *days*. The solution is  $P(t) = P_0 e^{rt} - \frac{20,000}{r}(e^{rt} - 1)$ . In the absence of predators, the governing equation is  $dP_1/dt = rP_1$ , with solution  $P_1(t) = P_0 e^{rt}$ . Based on the data, set  $P_1(7) = 2P_0$ , that is,  $2P_0 = P_0 e^{7r}$ . The growth rate is determined as  $r = \ln(2)/7 = .09902$  per day. Therefore the population, including the *predation* by birds, is  $P(t) = 2 \times 10^5 e^{.099t} - 201,997(e^{.099t} - 1) = 201,997.3 - 1977.3 e^{.099t}$ .

16(a).  $y(t) = \exp[2/10 + t/10 - 2\cos(t)/10]$ . The *doubling-time* is  $\tau \approx 2.9632$ .

(b). The differential equation is  $dy/dt = y/10$ , with solution  $y(t) = y(0)e^{t/10}$ . The *doubling-time* is given by  $\tau = 10\ln(2) \approx 6.9315$ .


(c). Consider the differential equation  $dy/dt = (0.5 + \sin(2\pi t))y/5$ . The equation is *separable*, with  $\frac{1}{y}dy = (0.1 + \frac{1}{5}\sin(2\pi t))dt$ . Integrating both sides, with respect to the appropriate variable, we obtain  $\ln y = (\pi t - \cos(2\pi t))/10\pi + c$ . Invoking the initial condition, the solution is  $y(t) = \exp[(1 + \pi t - \cos(2\pi t))/10\pi]$ . The *doubling-time* is  $\tau \approx 6.3804$ . The *doubling-time* approaches the value found in part(b).

(d).



17(a). The differential equation  $dy/dt = r(t)y - k$  is *linear*, with integrating factor  $\mu(t) = \exp[-\int r(t)dt]$ . Write the equation as  $(\mu y)' = -k\mu(t)$ . Integration of both

sides yields the general solution  $y = [-k \int \mu(\tau) d\tau + y_0 \mu(0)] / \mu(t)$ . In this problem, the integrating factor is  $\mu(t) = \exp[(\cos t - t)/5]$ .



(b). The population becomes *extinct*, if  $y(t^*) = 0$ , for some  $t = t^*$ . Referring to part(a), we find that  $y(t^*) = 0 \Rightarrow$

$$\int_0^{t^*} \exp[(\cos \tau - \tau)/5] d\tau = 5 e^{1/5} y_c.$$

It can be shown that the integral on the left hand side increases *monotonically*, from zero to a limiting value of approximately 5.0893. Hence extinction can happen *only if*  $5 e^{1/5} y_c < 5.0893$ , that is,  $y_c < 0.8333$ .

(c). Repeating the argument in part(b), it follows that  $y(t^*) = 0 \Rightarrow$

$$\int_0^{t^*} \exp[(\cos \tau - \tau)/5] d\tau = \frac{1}{k} e^{1/5} y_c.$$

Hence extinction can happen *only if*  $e^{1/5} y_c / k < 5.0893$ , that is,  $y_c < 4.1667 k$ .

(d). Evidently,  $y_c$  is a *linear* function of the parameter  $k$ .

19(a). Let  $Q(t)$  be the *volume* of carbon monoxide in the room. The rate of *increase* of CO is  $(.04)(0.1) = 0.004 \text{ ft}^3/\text{min}$ . The amount of CO *leaves the room* at a rate of  $(0.1)Q(t)/1200 = Q(t)/12000 \text{ ft}^3/\text{min}$ . Hence the total rate of change is given by the differential equation  $dQ/dt = 0.004 - Q(t)/12000$ . This equation is *linear* and separable, with solution  $Q(t) = 48 - 48 \exp(-t/12000) \text{ ft}^3$ . Note that  $Q_0 = 0 \text{ ft}^3$ . Hence the *concentration* at any time is given by  $x(t) = Q(t)/1200 = Q(t)/12 \%$ .


(b). The *concentration* of CO in the room is  $x(t) = 4 - 4 \exp(-t/12000) \%$ . A level of 0.00012 corresponds to 0.012%. Setting  $x(\tau) = 0.012$ , the solution of the equation  $4 - 4 \exp(-t/12000) = 0.012$  is  $\tau \approx 36 \text{ minutes}$ .

20(a). The concentration is  $c(t) = k + P/r + (c_0 - k - P/r)e^{-rt/V}$ . It is easy to see that  $c(t \rightarrow \infty) = k + P/r$ .

(b).  $c(t) = c_0 e^{-rt/V}$ . The reduction times are  $T_{50} = \ln(2)V/r$  and  $T_{10} = \ln(10)V/r$ .

(c). The reduction times, in years, are  $T_S = \ln(10)(65.2)/12,200 = 430.85$   
 $T_M = \ln(10)(158)/4,900 = 71.4$ ;  $T_E = \ln(10)(175)/460 = 6.05$   
 $T_O = \ln(10)(209)/16,000 = 17.63$ .


21(c).



22(a). The differential equation for the motion is  $m dv/dt = -v/30 - mg$ . Given the initial condition  $v(0) = 20 \text{ m/s}$ , the solution is  $v(t) = -44.1 + 64.1 \exp(-t/4.5)$ . Setting  $v(t_1) = 0$ , the ball reaches the maximum height at  $t_1 = 1.683 \text{ sec}$ . Integrating  $v(t)$ , the position is given by  $x(t) = 318.45 - 44.1t - 288.45 \exp(-t/4.5)$ . Hence the maximum height is  $x(t_1) = 45.78 \text{ m}$ .

(b). Setting  $x(t_2) = 0$ , the ball hits the ground at  $t_2 = 5.128 \text{ sec}$ .

(c).




23(a). The differential equation for the upward motion is  $m dv/dt = -\mu v^2 - mg$ , in which  $\mu = 1/1325$ . This equation is separable, with  $\frac{m}{\mu v^2 + mg} dv = -dt$ . Integrating

both sides and invoking the initial condition,  $v(t) = 44.133 \tan(.425 - .222t)$ . Setting  $v(t_1) = 0$ , the ball reaches the maximum height at  $t_1 = 1.916 \text{ sec}$ . Integrating  $v(t)$ , the position is given by  $x(t) = 198.75 \ln[\cos(0.222t - 0.425)] + 48.57$ . Therefore the *maximum height* is  $x(t_1) = 48.56 \text{ m}$ .

(b). The differential equation for the *downward* motion is  $m \, dv/dt = +\mu v^2 - mg$ . This equation is also separable, with  $\frac{m}{mg - \mu v^2} dv = -dt$ . For convenience, set  $t = 0$  at the *top* of the trajectory. The new initial condition becomes  $v(0) = 0$ . Integrating both sides and invoking the initial condition, we obtain  $\ln[(44.13 - v)/(44.13 + v)] = t/2.25$ .

Solving for the velocity,  $v(t) = 44.13(1 - e^{t/2.25})/(1 + e^{t/2.25})$ . Integrating  $v(t)$ , the position is given by  $x(t) = 99.29 \ln[e^{t/2.25}/(1 + e^{t/2.25})^2] + 186.2$ . To estimate the *duration* of the downward motion, set  $x(t_2) = 0$ , resulting in  $t_2 = 3.276 \text{ sec}$ . Hence the *total time* that the ball remains in the air is  $t_1 + t_2 = 5.192 \text{ sec}$ .

(c).



24(a). Measure the positive direction of motion *downward*. Based on Newton's 2nd law, the equation of motion is given by

$$m \frac{dv}{dt} = \begin{cases} -0.75v + mg & , 0 < t < 10 \\ -12v + mg & , t > 10 \end{cases} .$$

Note that gravity acts in the *positive* direction, and the drag force is *resistive*. During the first ten seconds of fall, the initial value problem is  $dv/dt = -v/7.5 + 32$ , with initial velocity  $v(0) = 0 \text{ fps}$ . This differential equation is separable and linear, with solution  $v(t) = 240(1 - e^{-t/7.5})$ . Hence  $v(10) = 176.7 \text{ fps}$ .


(b). Integrating the velocity, with  $x(t) = 0$ , the distance fallen is given by

$$x(t) = 240t + 1800 e^{-t/7.5} - 1800 .$$

Hence  $x(10) = 1074.5 \text{ ft}$ .

(c). For computational purposes, reset time to  $t = 0$ . For the remainder of the motion, the initial value problem is  $dv/dt = -32v/15 + 32$ , with specified initial velocity  $v(0) = 176.7 \text{ fps}$ . The solution is given by  $v(t) = 15 + 161.7 e^{-32t/15}$ . As  $t \rightarrow \infty$ ,  $v(t) \rightarrow v_L = 15 \text{ fps}$ . Integrating the velocity, with  $x(0) = 1074.5$ , the distance fallen after the parachute is open is given by  $x(t) = 15t - 75.8 e^{-32t/15} + 1150.3$ . To find the duration of the second part of the motion, estimate the root of the transcendental equation  $15T - 75.8 e^{-32T/15} + 1150.3 = 5000$ . The result is  $T = 256.6 \text{ sec}$ .

(d).



25(a). Measure the positive direction of motion *upward*. The equation of motion is given by  $mdv/dt = -kv - mg$ . The initial value problem is  $dv/dt = -kv/m - g$ , with  $v(0) = v_0$ . The solution is  $v(t) = -mg/k + (v_0 + mg/k)e^{-kt/m}$ . Setting  $v(t_m) = 0$ , the maximum height is reached at time  $t_m = (m/k)\ln[(mg + kv_0)/mg]$ . Integrating the velocity, the position of the body is

$$x(t) = -mgt/k + \left[ \left(\frac{m}{k}\right)^2 g + \frac{m v_0}{k} \right] (1 - e^{-kt/m}).$$

Hence the maximum height reached is

$$x_m = x(t_m) = \frac{m v_0}{k} - g \left(\frac{m}{k}\right)^2 \ln \left[ \frac{mg + k v_0}{mg} \right].$$

(b). Recall that for  $\delta \ll 1$ ,  $\ln(1 + \delta) = \delta - \frac{1}{2}\delta^2 + \frac{1}{3}\delta^3 - \frac{1}{4}\delta^4 + \dots$


26(b).  $\lim_{k \rightarrow 0} \frac{-mg + (k v_0 + mg)e^{-kt/m}}{k} = \lim_{k \rightarrow 0} -\frac{t}{m} (k v_0 + mg)e^{-kt/m} = -gt$ .

(c).  $\lim_{m \rightarrow 0} \left[ -\frac{mg}{k} + \left(\frac{mg}{k} + v_0\right)e^{-kt/m} \right] = 0$ , since  $\lim_{m \rightarrow 0} e^{-kt/m} = 0$ .

28(a). In terms of displacement, the differential equation is  $mv dv/dx = -kv + mg$ . This follows from the *chain rule*:  $\frac{dv}{dt} = \frac{dv}{dx} \frac{dx}{dt} = v \frac{dv}{dx}$ . The differential equation is separable, with

$$x(v) = -\frac{mv}{k} - \frac{m^2g}{k^2} \ln \left| \frac{mg - kv}{mg} \right|.$$

The inverse *exists*, since both  $x$  and  $v$  are monotone increasing. In terms of the given parameters,  $x(v) = -1.25v - 15.31 \ln|0.0816v - 1|$ .



(b).  $x(10) = 13.45$  meters. The required value is  $k = 0.24$ .

(c). In part(a), set  $v = 10$  m/s and  $x = 10$  meters.

29(a). Let  $x$  represent the height above the earth's surface. The equation of motion is given by  $m \frac{dv}{dt} = -G \frac{Mm}{(R+x)^2}$ , in which  $G$  is the universal gravitational constant. The symbols  $M$  and  $R$  are the *mass* and *radius* of the earth, respectively. By the chain rule,

$$mv \frac{dv}{dx} = -G \frac{Mm}{(R+x)^2}.$$

This equation is separable, with  $v dv = -GM(R+x)^{-2} dx$ . Integrating both sides, and

invoking the initial condition  $v(0) = \sqrt{2gR}$ , the solution is  $v^2 = 2GM(R+x)^{-1} + 2gR - 2GM/R$ . From elementary physics, it follows that  $g = GM/R^2$ . Therefore  $v(x) = \sqrt{2g} \left[ R/\sqrt{R+x} \right]$ . (Note that  $g = 78,545$  mi/hr<sup>2</sup>.)


(b). We now consider  $dx/dt = \sqrt{2g} \left[ R/\sqrt{R+x} \right]$ . This equation is also separable, with  $\sqrt{R+x} dx = \sqrt{2g} R dt$ . By definition of the variable  $x$ , the initial condition is  $x(0) = 0$ . Integrating both sides, we obtain  $x(t) = \left[ \frac{3}{2} (\sqrt{2g} R t + \frac{2}{3} R^{3/2}) \right]^{2/3} - R$ . Setting the distance  $x(T) + R = 240,000$ , and solving for  $T$ , the duration of such a flight would be  $T \approx 49$  hours.

32(a). Both equations are linear and separable. The initial conditions are  $v(0) = u \cos A$  and  $w(0) = u \sin A$ . The two solutions are  $v(t) = u \cos A e^{-rt}$  and  $w(t) = -g/r + (u \sin A + g/r) e^{-rt}$ .

(b). Integrating the solutions in part(a), and invoking the initial conditions, the coordinates are  $x(t) = \frac{u}{r} \cos A(1 - e^{-rt})$  and

$$y(t) = -gt/r + (g + ur \sin A + hr^2)/r^2 - \left(\frac{u}{r} \sin A + g/r^2\right)e^{-rt}.$$

(c).



(d). Let  $T$  be the time that it takes the ball to go 350 ft horizontally. Then from above,  $e^{-T/5} = (u \cos A - 70)/u \cos A$ . At the same time, the height of the ball is given by  $y(T) = -160T + 267 + 125u \sin A - (800 + 5u \sin A)[(u \cos A - 70)/u \cos A]$ . Hence  $A$  and  $u$  must satisfy the inequality

$$800 \ln \left[ \frac{u \cos A - 70}{u \cos A} \right] + 267 + 125u \sin A - (800 + 5u \sin A) \left[ \frac{u \cos A - 70}{u \cos A} \right] \geq 10.$$

33(a). Solving equation (i),  $y'(x) = [(k^2 - y)/y]^{1/2}$ . The *positive* answer is chosen, since  $y$  is an *increasing* function of  $x$ .

(b). Let  $y = k^2 \sin^2 t$ . Then  $dy = 2k^2 \sin t \cos t dt$ . Substituting into the equation in part(a), we find that

$$\frac{2k^2 \sin t \cos t dt}{dx} = \frac{\cos t}{\sin t}.$$

Hence  $2k^2 \sin^2 t dt = dx$ .

(c). Letting  $\theta = 2t$ , we further obtain  $k^2 \sin^2 \frac{\theta}{2} d\theta = dx$ . Integrating both sides of the equation and noting that  $t = \theta = 0$  corresponds to the *origin*, we obtain the solutions  $x(\theta) = k^2(\theta - \sin \theta)/2$  and [from part(b)]  $y(\theta) = k^2(1 - \cos \theta)/2$ .

(d). Note that  $y/x = (1 - \cos \theta)/(\theta - \sin \theta)$ . Setting  $x = 1, y = 2$ , the solution of the equation  $(1 - \cos \theta)/(\theta - \sin \theta) = 2$  is  $\theta \approx 1.401$ . Substitution into either of the expressions yields  $k \approx 2.193$ .

## Section 2.4

2. Considering the roots of the coefficient of the leading term, the ODE has unique solutions on intervals *not* containing 0 or 4. Since  $2 \in (0, 4)$ , the initial value problem has a unique solution on the interval  $(0, 4)$ .

3. The function  $\tan t$  is discontinuous at *odd multiples* of  $\frac{\pi}{2}$ . Since  $\frac{\pi}{2} < \pi < \frac{3\pi}{2}$ , the initial value problem has a unique solution on the interval  $(\frac{\pi}{2}, \frac{3\pi}{2})$ .

5.  $p(t) = 2t/(4 - t^2)$  and  $g(t) = 3t^2/(4 - t^2)$ . These functions are discontinuous at  $x = \pm 2$ . The initial value problem has a unique solution on the interval  $(-2, 2)$ .

6. The function  $\ln t$  is defined and continuous on the interval  $(0, \infty)$ . Therefore the initial value problem has a unique solution on the interval  $(0, \infty)$ .

7. The function  $f(t, y)$  is continuous everywhere on the plane, *except* along the straight line  $y = -2t/5$ . The partial derivative  $\partial f/\partial y = -7t/(2t + 5y)^2$  has the *same* region of continuity.

9. The function  $f(t, y)$  is discontinuous along the coordinate axes, and on the hyperbola  $t^2 - y^2 = 1$ . Furthermore,

$$\frac{\partial f}{\partial y} = \frac{\pm 1}{y(1 - t^2 + y^2)} - 2 \frac{y \ln|ty|}{(1 - t^2 + y^2)^2}$$

has the *same* points of discontinuity.

10.  $f(t, y)$  is continuous everywhere on the plane. The partial derivative  $\partial f/\partial y$  is also continuous everywhere.

12. The function  $f(t, y)$  is discontinuous along the lines  $t = \pm k\pi$  and  $y = -1$ . The partial derivative  $\partial f/\partial y = \cot(t)/(1 + y)^2$  has the *same* region of continuity.

14. The equation is separable, with  $dy/y^2 = 2t dt$ . Integrating both sides, the solution is given by  $y(t) = y_0/(1 - y_0 t^2)$ . For  $y_0 > 0$ , solutions exist as long as  $t^2 < 1/y_0$ . For  $y_0 \leq 0$ , solutions are defined for *all*  $t$ .

15. The equation is separable, with  $dy/y^3 = -dt$ . Integrating both sides and invoking the initial condition,  $y(t) = y_0/\sqrt{2y_0 t + 1}$ . Solutions exist as long as  $2y_0 t + 1 > 0$ , that is,  $2y_0 t > -1$ . If  $y_0 > 0$ , solutions exist for  $t > -1/2y_0$ . If  $y_0 = 0$ , then the solution  $y(t) = 0$  exists for all  $t$ . If  $y_0 < 0$ , solutions exist for  $t < -1/2y_0$ .

16. The function  $f(t, y)$  is discontinuous along the straight lines  $t = -1$  and  $y = 0$ . The partial derivative  $\partial f/\partial y$  is discontinuous along the same lines. The equation is




separable, with  $y dy = t^2 dt/(1 + t^3)$ . Integrating and invoking the initial condition, the solution is  $y(t) = [\frac{2}{3} \ln|1 + t^3| + y_0^2]^{1/2}$ . Solutions exist as long as


$$\frac{2}{3} \ln|1 + t^3| + y_0^2 \geq 0,$$

that is,  $y_0^2 \geq -\frac{2}{3} \ln|1 + t^3|$ . For all  $y_0$  (it can be verified that  $y_0 = 0$  yields a valid solution, even though Theorem 2.4.2 does not guarantee one), solutions exist as long as  $|1 + t^3| \geq \exp(-3y_0^2/2)$ . From above, we must have  $t > -1$ . Hence the inequality may be written as  $t^3 \geq \exp(-3y_0^2/2) - 1$ . It follows that the solutions are valid for  $[\exp(-3y_0^2/2) - 1]^{1/3} < t < \infty$ .

17.




18.




Based on the direction field, and the differential equation, for  $y_0 < 0$ , the slopes *eventually* become negative, and hence solutions tend to  $-\infty$ . For  $y_0 > 0$ , solutions increase without bound if  $t_0 < 0$ . Otherwise, the slopes *eventually* become negative, and solutions tend to zero. Furthermore,  $y_0 = 0$  is an *equilibrium solution*. Note that slopes are zero along the curves  $y = 0$  and  $ty = 3$ .

19.



For initial conditions  $(t_0, y_0)$  satisfying  $ty < 3$ , the respective solutions all tend to zero. Solutions with initial conditions above or below the hyperbola  $ty = 3$  eventually tend to  $\pm\infty$ . Also,  $y_0 = 0$  is an equilibrium solution.

20.



Solutions with  $t_0 < 0$  all tend to  $-\infty$ . Solutions with initial conditions  $(t_0, y_0)$  to the right of the parabola  $t = 1 + y^2$  asymptotically approach the parabola as  $t \rightarrow \infty$ . Integral curves with initial conditions above the parabola (and  $y_0 > 0$ ) also approach the curve. The slopes for solutions with initial conditions below the parabola (and  $y_0 < 0$ ) are all negative. These solutions tend to  $-\infty$ .

21. Define  $y_c(t) = \frac{2}{3}(t - c)^{3/2}u(t - c)$ , in which  $u(t)$  is the Heaviside step function. Note that  $y_c(c) = y_c(0) = 0$  and  $y_c(c + (3/2)^{2/3}) = 1$ .

(a). Let  $c = 1 - (3/2)^{2/3}$ .

(b). Let  $c = 2 - (3/2)^{2/3}$ .

(c). Observe that  $y_0(2) = \frac{2}{3}(2)^{3/2}$ ,  $y_c(t) < \frac{2}{3}(2)^{3/2}$  for  $0 < c < 2$ , and that  $y_c(2) = 0$  for  $c \geq 2$ . So for any  $c \geq 0$ ,  $\pm y_c(2) \in [-2, 2]$ .

26(a). Recalling Eq. (35) in Section 2.1,

$$y = \frac{1}{\mu(t)} \int \mu(s)g(s) ds + \frac{c}{\mu(t)}.$$

It is evident that  $y_1(t) = \frac{1}{\mu(t)}$  and  $y_2(t) = \frac{1}{\mu(t)} \int \mu(s)g(s) ds$ .

(b). By definition,  $\frac{1}{\mu(t)} = \exp(-\int p(t)dt)$ . Hence  $y_1' = -p(t) \frac{1}{\mu(t)} = -p(t)y_1$ . That is,  $y_1' + p(t)y_1 = 0$ .

(c).  $y_2' = \left(-p(t) \frac{1}{\mu(t)}\right) \int_0^t \mu(s)g(s) ds + \left(\frac{1}{\mu(t)}\right) \mu(t)g(t) = -p(t)y_2 + g(t)$ . That is,  $y_2' + p(t)y_2 = g(t)$ .

30. Since  $n = 3$ , set  $v = y^{-2}$ . It follows that  $\frac{dv}{dt} = -2y^{-3} \frac{dy}{dt}$  and  $\frac{dy}{dt} = -\frac{y^3}{2} \frac{dv}{dt}$ . Substitution into the differential equation yields  $-\frac{y^3}{2} \frac{dv}{dt} - \varepsilon y = -\sigma y^3$ , which further results in  $v' + 2\varepsilon v = 2\sigma$ . The latter differential equation is linear, and can be written as  $(e^{2\varepsilon t})' = 2\sigma$ . The solution is given by  $v(t) = 2\sigma t e^{-2\varepsilon t} + c e^{-2\varepsilon t}$ . Converting back to the original dependent variable,  $y = \pm v^{-1/2}$ .

31. Since  $n = 3$ , set  $v = y^{-2}$ . It follows that  $\frac{dv}{dt} = -2y^{-3} \frac{dy}{dt}$  and  $\frac{dy}{dt} = -\frac{y^3}{2} \frac{dv}{dt}$ . The differential equation is written as  $-\frac{y^3}{2} \frac{dv}{dt} - (\Gamma \cos t + T)y = \sigma y^3$ , which upon further substitution is  $v' + 2(\Gamma \cos t + T)v = 2$ . This ODE is linear, with integrating factor  $\mu(t) = \exp(2\int(\Gamma \cos t + T)dt) = \exp(-2\Gamma \sin t + 2Tt)$ . The solution is

$$v(t) = 2 \exp(2\Gamma \sin t - 2Tt) \int_0^t \exp(-2\Gamma \sin \tau + 2T\tau) d\tau + c \exp(-2\Gamma \sin t + 2Tt).$$

Converting back to the original dependent variable,  $y = \pm v^{-1/2}$ .

33. The solution of the initial value problem  $y_1' + 2y_1 = 0$ ,  $y_1(0) = 1$  is  $y_1(t) = e^{-2t}$ . Therefore  $y(1^-) = y_1(1) = e^{-2}$ . On the interval  $(1, \infty)$ , the differential equation is  $y_2' + y_2 = 0$ , with  $y_2(t) = ce^{-t}$ . Therefore  $y(1^+) = y_2(1) = ce^{-1}$ . Equating the limits  $y(1^-) = y(1^+)$ , we require that  $c = e^{-1}$ . Hence the global solution of the initial value problem is


$$y(t) = \begin{cases} e^{-2t}, & 0 \leq t \leq 1 \\ e^{-1-t}, & t > 1 \end{cases}.$$

Note the discontinuity of the derivative

$$y(t) = \begin{cases} -2e^{-2t}, & 0 < t < 1 \\ -e^{-1-t}, & t > 1 \end{cases}.$$


Section 2.5

1.




For  $y_0 \geq 0$ , the only equilibrium point is  $y^* = 0$ .  $f'(0) = a > 0$ , hence the equilibrium solution  $\phi(t) = 0$  is *unstable*.

2.




The equilibrium points are  $y^* = -a/b$  and  $y^* = 0$ .  $f'(-a/b) < 0$ , therefore the equilibrium solution  $\phi(t) = -a/b$  is *asymptotically stable*.

3.




4.




The only equilibrium point is  $y^* = 0$ .  $f'(0) > 0$ , hence the equilibrium solution  $\phi(t) = 0$  is *unstable*.

5.




The only equilibrium point is  $y^* = 0$ .  $f'(0) < 0$ , hence the equilibrium solution  $\phi(t) = 0$  is *asymptotically stable*.


6.



7(b).




8.




The only equilibrium point is  $y^* = 1$ . Note that  $f'(1) = 0$ , and that  $y' < 0$  for  $y \neq 1$ . As long as  $y_0 \neq 1$ , the corresponding solution is *monotone decreasing*. Hence the equilibrium solution  $\phi(t) = 1$  is *semistable*.

9.




10.




The equilibrium points are  $y^* = 0, \pm 1$ .  $f'(y) = 1 - 3y^2$ . The equilibrium solution  $\phi(t) = 0$  is *unstable*, and the remaining two are *asymptotically stable*.

11.




12.



The equilibrium points are  $y^* = 0, \pm 2$ .  $f'(y) = 8y - 4y^3$ . The equilibrium solutions  $\phi(t) = -2$  and  $\phi(t) = +2$  are *unstable* and *asymptotically stable*, respectively. The equilibrium solution  $\phi(t) = 0$  is *semistable*.

13.



The equilibrium points are  $y^* = 0$  and  $1$ .  $f'(y) = 2y - 6y^2 + 4y^3$ . Both equilibrium solutions are *semistable*.

15(a). Inverting the Solution (11), Eq. (13) shows  $t$  as a function of the population  $y$  and the carrying capacity  $K$ . With  $y_0 = K/3$ ,

$$t = -\frac{1}{r} \ln \left| \frac{(1/3)[1 - (y/K)]}{(y/K)[1 - (1/3)]} \right|.$$

Setting  $y = 2y_0$ ,

$$\tau = -\frac{1}{r} \ln \left| \frac{(1/3)[1 - (2/3)]}{(2/3)[1 - (1/3)]} \right|.$$


That is,  $\tau = \frac{1}{r} \ln 4$ . If  $r = 0.025$  per year,  $\tau = 55.45$  years.

(b). In Eq. (13), set  $y_0/K = \alpha$  and  $y/K = \beta$ . As a result, we obtain

$$T = -\frac{1}{r} \ln \left| \frac{\alpha[1 - \beta]}{\beta[1 - \alpha]} \right|.$$

Given  $\alpha = 0.1$ ,  $\beta = 0.9$  and  $r = 0.025$  per year,  $\tau = 175.78$  years.

16(a).





17. Consider the change of variable  $u = \ln(y/K)$ . Differentiating both sides with respect

to  $t$ ,  $u' = y'/y$ . Substitution into the Gompertz equation yields  $u' = -ru$ , with solution  $u = u_0 e^{-rt}$ . It follows that  $\ln(y/K) = \ln(y_0/K) e^{-rt}$ . That is,

$$\frac{y}{K} = \exp[\ln(y_0/K) e^{-rt}].$$

(a). Given  $K = 80.5 \times 10^6$ ,  $y_0/K = 0.25$  and  $r = 0.71$  per year,  $y(2) = 57.58 \times 10^6$ .

(b). Solving for  $t$ ,

$$t = -\frac{1}{r} \ln \left[ \frac{\ln(y/K)}{\ln(y_0/K)} \right].$$

Setting  $y(\tau) = 0.75K$ , the corresponding time is  $\tau = 2.21$  years.

19(a). The rate of *increase* of the volume is given by rate of *flow in* – rate of *flow out*. That is,  $dV/dt = k - \alpha a \sqrt{2gh}$ . Since the cross section is *constant*,  $dV/dt = Adh/dt$ . Hence the governing equation is  $dh/dt = (k - \alpha a \sqrt{2gh})/A$ .

(b). Setting  $dh/dt = 0$ , the equilibrium height is  $h_e = \frac{1}{2g} \left( \frac{k}{\alpha a} \right)^2$ . Furthermore, since  $f'(h_e) < 0$ , it follows that the equilibrium height is *asymptotically stable*.

(c). Based on the answer in part(b), the water level will intrinsically tend to approach  $h_e$ . Therefore the height of the tank must be *greater* than  $h_e$ ; that is,  $h_e < V/A$ .

22(a). The equilibrium points are at  $y^* = 0$  and  $y^* = 1$ . Since  $f'(y) = \alpha - 2\alpha y$ , the equilibrium solution  $\phi = 0$  is *unstable* and the equilibrium solution  $\phi = 1$  is *asymptotically stable*.

(b). The ODE is separable, with  $[y(1-y)]^{-1} dy = \alpha dt$ . Integrating both sides and invoking the initial condition, the solution is

$$y(t) = \frac{y_0 e^{\alpha t}}{1 - y_0 + y_0 e^{\alpha t}}.$$

It is evident that (independent of  $y_0$ )  $\lim_{t \rightarrow -\infty} y(t) = 0$  and  $\lim_{t \rightarrow \infty} y(t) = 1$ .

23(a).  $y(t) = y_0 e^{-\beta t}$ .

(b). From part(a),  $dx/dt = \alpha x y_0 e^{-\beta t}$ . Separating variables,  $dx/x = \alpha y_0 e^{-\beta t} dt$ . Integrating both sides, the solution is  $x(t) = x_0 \exp[\alpha y_0 / \beta (1 - e^{-\beta t})]$ .


(c). As  $t \rightarrow \infty$ ,  $y(t) \rightarrow 0$  and  $x(t) \rightarrow x_0 \exp(\alpha y_0 / \beta)$ . Over a *long* period of time, the

proportion of carriers *vanishes*. Therefore the proportion of the population that escapes the epidemic is the proportion of *susceptibles* left at that time,  $x_0 \exp(\alpha y_0/\beta)$ .

25(a). Note that  $f(x) = x[(R - R_c) - ax^2]$ , and  $f'(x) = (R - R_c) - 3ax^2$ . So if  $(R - R_c) < 0$ , the only equilibrium point is  $x^* = 0$ .  $f'(0) < 0$ , and hence the solution  $\phi(t) = 0$  is *asymptotically stable*.

(b). If  $(R - R_c) > 0$ , there are *three* equilibrium points  $x^* = 0, \pm\sqrt{(R - R_c)/a}$ . Now  $f'(0) > 0$ , and  $f'(\pm\sqrt{(R - R_c)/a}) < 0$ . Hence the solution  $\phi = 0$  is *unstable*, and the solutions  $\phi = \pm\sqrt{(R - R_c)/a}$  are *asymptotically stable*.

(c).



## Section 2.6

1.  $M(x, y) = 2x + 3$  and  $N(x, y) = 2y - 2$ . Since  $M_y = N_x = 0$ , the equation is *exact*. Integrating  $M$  with respect to  $x$ , while holding  $y$  constant, yields  $\psi(x, y) = x^2 + 3x + h(y)$ . Now  $\psi_y = h'(y)$ , and equating with  $N$  results in the possible function  $h(y) = y^2 - 2y$ . Hence  $\psi(x, y) = x^2 + 3x + y^2 - 2y$ , and the solution is defined *implicitly* as  $x^2 + 3x + y^2 - 2y = c$ .
2.  $M(x, y) = 2x + 4y$  and  $N(x, y) = 2x - 2y$ . Note that  $M_y \neq N_x$ , and hence the differential equation is *not exact*.
4. First divide both sides by  $(2xy + 2)$ . We now have  $M(x, y) = y$  and  $N(x, y) = x$ . Since  $M_y = N_x = 0$ , the resulting equation is *exact*. Integrating  $M$  with respect to  $x$ , while holding  $y$  constant, results in  $\psi(x, y) = xy + h(y)$ . Differentiating with respect to  $y$ ,  $\psi_y = x + h'(y)$ . Setting  $\psi_y = N$ , we find that  $h'(y) = 0$ , and hence  $h(y) = 0$  is acceptable. Therefore the solution is defined *implicitly* as  $xy = c$ . Note that if  $xy + 1 = 0$ , the equation is trivially satisfied.
6. Write the given equation as  $(ax - by)dx + (bx - cy)dy$ . Now  $M(x, y) = ax - by$  and  $N(x, y) = bx - cy$ . Since  $M_y \neq N_x$ , the differential equation is *not exact*.
8.  $M(x, y) = e^x \sin y + 3y$  and  $N(x, y) = -3x + e^x \sin y$ . Note that  $M_y \neq N_x$ , and hence the differential equation is *not exact*.
10.  $M(x, y) = y/x + 6x$  and  $N(x, y) = \ln x - 2$ . Since  $M_y = N_x = 1/x$ , the given equation is *exact*. Integrating  $N$  with respect to  $y$ , while holding  $x$  constant, results in  $\psi(x, y) = y \ln x - 2y + h(x)$ . Differentiating with respect to  $x$ ,  $\psi_x = y/x + h'(x)$ . Setting  $\psi_x = M$ , we find that  $h'(x) = 6x$ , and hence  $h(x) = 3x^2$ . Therefore the solution is defined *implicitly* as  $3x^2 + y \ln x - 2y = c$ .
11.  $M(x, y) = x \ln y + xy$  and  $N(x, y) = y \ln x + xy$ . Note that  $M_y \neq N_x$ , and hence the differential equation is *not exact*.
13.  $M(x, y) = 2x - y$  and  $N(x, y) = 2y - x$ . Since  $M_y = N_x = -1$ , the equation is *exact*. Integrating  $M$  with respect to  $x$ , while holding  $y$  constant, yields  $\psi(x, y) = x^2 - xy + h(y)$ . Now  $\psi_y = -x + h'(y)$ . Equating  $\psi_y$  with  $N$  results in  $h'(y) = 2y$ , and hence  $h(y) = y^2$ . Thus  $\psi(x, y) = x^2 - xy + y^2$ , and the solution is given *implicitly* as  $x^2 - xy + y^2 = c$ . Invoking the initial condition  $y(1) = 3$ , the specific solution is  $x^2 - xy + y^2 = 7$ . The *explicit* form of the solution is  $y(x) = \frac{1}{2} \left[ x + \sqrt{28 - 3x^2} \right]$ . Hence the solution is valid as long as  $3x^2 \leq 28$ .
16.  $M(x, y) = y e^{2xy} + x$  and  $N(x, y) = bx e^{2xy}$ . Note that  $M_y = e^{2xy} + 2xy e^{2xy}$ , and  $N_x = b e^{2xy} + 2bxy e^{2xy}$ . The given equation is *exact*, as long as  $b = 1$ . Integrating

$N$  with respect to  $y$ , while holding  $x$  constant, results in  $\psi(x, y) = e^{2xy}/2 + h(x)$ . Now differentiating with respect to  $x$ ,  $\psi_x = ye^{2xy} + h'(x)$ . Setting  $\psi_x = M$ , we find that  $h'(x) = x$ , and hence  $h(x) = x^2/2$ . Conclude that  $\psi(x, y) = e^{2xy}/2 + x^2/2$ . Hence the solution is given *implicitly* as  $e^{2xy} + x^2 = c$ .

17. Integrating  $\psi_y = N$ , while holding  $x$  constant, yields

$$\psi(x, y) = \int N(x, y)dy + h(x).$$

Taking the partial derivative with respect to  $x$ ,  $\psi_x = \int \frac{\partial}{\partial x} N(x, y)dy + h'(x)$ . Now set  $\psi_x = M(x, y)$  and therefore  $h'(x) = M(x, y) - \int \frac{\partial}{\partial x} N(x, y)dy$ . Based on the fact that  $M_y = N_x$ , it follows that  $\frac{\partial}{\partial y}[h'(x)] = 0$ . Hence the expression for  $h'(x)$  can be integrated to obtain

$$h(x) = \int M(x, y)dx - \int \left[ \int \frac{\partial}{\partial x} N(x, y)dy \right] dx.$$

18. Observe that  $\frac{\partial}{\partial y}[M(x)] = \frac{\partial}{\partial x}[N(y)] = 0$ .

20.  $M_y = y^{-1}\cos y - y^{-2}\sin y$  and  $N_x = -2e^{-x}(\cos x + \sin x)/y$ . Multiplying both sides by the integrating factor  $\mu(x, y) = ye^x$ , the given equation can be written as  $(e^x \sin y - 2y \sin x)dx + (e^x \cos y + 2\cos x)dy = 0$ . Let  $\overline{M} = \mu M$  and  $\overline{N} = \mu N$ . Observe that  $\overline{M}_y = \overline{N}_x$ , and hence the latter ODE is *exact*. Integrating  $\overline{N}$  with respect to  $y$ , while holding  $x$  constant, results in  $\psi(x, y) = e^x \sin y + 2y \cos x + h(x)$ . Now differentiating with respect to  $x$ ,  $\psi_x = e^x \sin y - 2y \sin x + h'(x)$ . Setting  $\psi_x = \overline{M}$ , we find that  $h'(x) = 0$ , and hence  $h(x) = 0$  is feasible. Hence the solution of the given equation is defined *implicitly* by  $e^x \sin y + 2y \cos x = \beta$ .

21.  $M_y = 1$  and  $N_x = 2$ . Multiply both sides by the integrating factor  $\mu(x, y) = y$  to obtain  $y^2 dx + (2xy - y^2 e^y)dy = 0$ . Let  $\overline{M} = yM$  and  $\overline{N} = yN$ . It is easy to see that  $\overline{M}_y = \overline{N}_x$ , and hence the latter ODE is *exact*. Integrating  $\overline{M}$  with respect to  $x$  yields  $\psi(x, y) = xy^2 + h(y)$ . Equating  $\psi_y$  with  $\overline{N}$  results in  $h'(y) = -y^2 e^y$ , and hence  $h(y) = -e^y(y^2 - 2y + 2)$ . Thus  $\psi(x, y) = xy^2 - e^y(y^2 - 2y + 2)$ , and the solution is defined *implicitly* by  $xy^2 - e^y(y^2 - 2y + 2) = c$ .

24. The equation  $\mu M + \mu N y' = 0$  has an integrating factor if  $(\mu M)_y = (\mu N)_x$ , that is,  $\mu_y M - \mu_x N = \mu N_x - \mu M_y$ . Suppose that  $N_x - M_y = R(xM - yN)$ , in which  $R$  is some function depending *only* on the quantity  $z = xy$ . It follows that the modified form of the equation is *exact*, if  $\mu_y M - \mu_x N = \mu R(xM - yN) = R(\mu xM - \mu yN)$ . This relation is satisfied if  $\mu_y = (\mu x)R$  and  $\mu_x = (\mu y)R$ . Now consider  $\mu = \mu(xy)$ . Then the partial derivatives are  $\mu_x = \mu' y$  and  $\mu_y = \mu' x$ . Note that  $\mu' = d\mu/dz$ . Thus  $\mu$  must satisfy  $\mu'(z) = R(z)$ . The latter equation is *separable*, with  $d\mu = R(z)dz$ , and  $\mu(z) = \int R(z)dz$ . Therefore, given  $R = R(xy)$ , it is possible to determine  $\mu = \mu(xy)$  which becomes an integrating factor of the differential equation.

28. The equation is not exact, since  $N_x - M_y = 2y - 1$ . However,  $(N_x - M_y)/M = (2y - 1)/y$  is a function of  $y$  alone. Hence there exists  $\mu = \mu(y)$ , which is a solution of the differential equation  $\mu' = (2 - 1/y)\mu$ . The latter equation is *separable*, with  $d\mu/\mu = 2 - 1/y$ . One solution is  $\mu(y) = \exp(2y - \ln y) = e^{2y}/y$ . Now rewrite the given ODE as  $e^{2y}dx + (2xe^{2y} - 1/y)dy = 0$ . This equation is *exact*, and it is easy to see that  $\psi(x, y) = xe^{2y} - \ln y$ . Therefore the solution of the given equation is defined implicitly by  $xe^{2y} - \ln y = c$ .

30. The given equation is not exact, since  $N_x - M_y = 8x^3/y^3 + 6/y^2$ . But note that  $(N_x - M_y)/M = 2/y$  is a function of  $y$  alone, and hence there is an integrating factor  $\mu = \mu(y)$ . Solving the equation  $\mu' = (2/y)\mu$ , an integrating factor is  $\mu(y) = y^2$ . Now rewrite the differential equation as  $(4x^3 + 3y)dx + (3x + 4y^3)dy = 0$ . By inspection,  $\psi(x, y) = x^4 + 3xy + y^4$ , and the solution of the given equation is defined implicitly by  $x^4 + 3xy + y^4 = c$ .

32. Multiplying both sides of the ODE by  $\mu = [xy(2x + y)]^{-1}$ , the given equation is equivalent to  $[(3x + y)/(2x^2 + xy)]dx + [(x + y)/(2xy + y^2)]dy = 0$ . Rewrite the differential equation as

$$\left[\frac{2}{x} + \frac{2}{2x + y}\right]dx + \left[\frac{1}{y} + \frac{1}{2x + y}\right]dy = 0.$$

It is easy to see that  $M_y = N_x$ . Integrating  $M$  with respect to  $x$ , while keeping  $y$  constant, results in  $\psi(x, y) = 2\ln|x| + \ln|2x + y| + h(y)$ . Now taking the partial derivative with respect to  $y$ ,  $\psi_y = (2x + y)^{-1} + h'(y)$ . Setting  $\psi_y = N$ , we find that  $h'(y) = 1/y$ , and hence  $h(y) = \ln|y|$ . Therefore

$$\psi(x, y) = 2\ln|x| + \ln|2x + y| + \ln|y|,$$

and the solution of the given equation is defined implicitly by  $2x^3y + x^2y^2 = c$ .

**Section 2.7**


2(a). The Euler formula is  $y_{n+1} = y_n + h(2y_n - 1) = (1 + 2h)y_n - h$ .

(d). The differential equation is *linear*, with solution  $y(t) = (1 + e^{2t})/2$ .

4(a). The Euler formula is  $y_{n+1} = (1 - 2h)y_n + 3h \cos t_n$ .


(d). The exact solution is  $y(t) = (6\cos t + 3\sin t - 6e^{-2t})/5$ .

5.




All solutions seem to converge to  $\phi(t) = 25/9$ .

6.




Solutions with *positive* initial conditions seem to converge to a specific function. On the other hand, solutions with *negative* coefficients decrease without bound.  $\phi(t) = 0$  is an equilibrium solution.

7.




All solutions seem to converge to a specific function.

8.




Solutions with initial conditions to the 'left' of the curve  $t = 0.1y^2$  seem to diverge. On the other hand, solutions to the 'right' of the curve seem to converge to zero. Also,  $\phi(t)$  is an equilibrium solution.

9.



All solutions seem to diverge.

10.



Solutions with *positive* initial conditions increase without bound. Solutions with *negative* initial conditions decrease without bound. Note that  $\phi(t) = 0$  is an equilibrium solution.

11. The Euler formula is  $y_{n+1} = y_n - 3h\sqrt{y_n} + 5h$ . The initial value is  $y_0 = 2$ .

12. The iteration formula is  $y_{n+1} = (1 + 3h)y_n - h t_n y_n^2$ .  $(t_0, y_0) = (0, 0.5)$ .

14. The iteration formula is  $y_{n+1} = (1 - h t_n)y_n + h y_n^3/10$ .  $(t_0, y_0) = (0, 1)$ .


17. The Euler formula is

$$y_{n+1} = y_n + \frac{h(y_n^2 + 2t_n y_n)}{3 + t_n^2}.$$

The initial point is  $(t_0, y_0) = (1, 2)$ .

18(a). See Problem 8.

19(a).



(b). The iteration formula is  $y_{n+1} = y_n + h y_n^2 - h t_n^2$ . The critical value of  $\alpha$  appears to be near  $\alpha_0 \approx 0.6815$ . For  $y_0 > \alpha_0$ , the iterations diverge.



20(a). The ODE is *linear*, with general solution  $y(t) = t + c e^t$ . Invoking the specified initial condition,  $y(t_0) = y_0$ , we have  $y_0 = t_0 + c e^{t_0}$ . Hence  $c = (y_0 - t_0)e^{-t_0}$ . Thus the solution is given by  $\phi(t) = (y_0 - t_0)e^{t-t_0} + t$ .

(b). The Euler formula is  $y_{n+1} = (1 + h)y_n + h - h t_n$ . Now set  $k = n + 1$ .

(c). We have  $y_1 = (1 + h)y_0 + h - h t_0 = (1 + h)y_0 + (t_1 - t_0) - h t_0$ . Rearranging the terms,  $y_1 = (1 + h)(y_0 - t_0) + t_1$ . Now suppose that  $y_k = (1 + h)^k(y_0 - t_0) + t_k$ , for some  $k \geq 1$ . Then  $y_{k+1} = (1 + h)y_k + h - h t_k$ . Substituting for  $y_k$ , we find that  $y_{k+1} = (1 + h)^{k+1}(y_0 - t_0) + (1 + h)t_k + h - h t_k = (1 + h)^{k+1}(y_0 - t_0) + t_k + h$ . Noting that  $t_{k+1} = t_k + h$ , the result is verified.

(d). Substituting  $h = (t - t_0)/n$ , with  $t_n = t$ ,

$$y_n = \left(1 + \frac{t - t_0}{n}\right)^n (y_0 - t_0) + t.$$

Taking the limit of both sides, as  $n \rightarrow \infty$ , and using the fact that  $\lim_{n \rightarrow \infty} (1 + a/n)^n = e^a$ , pointwise convergence is proved.

21. The exact solution is  $\phi(t) = e^t$ . The Euler formula is  $y_{n+1} = (1 + h)y_n$ . It is easy to see that  $y_n = (1 + h)^n y_0 = (1 + h)^n$ . Given  $t > 0$ , set  $h = t/n$ . Taking the limit, we find that  $\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} (1 + t/n)^n = e^t$ .

23. The exact solution is  $\phi(t) = t/2 + e^{2t}$ . The Euler formula is  $y_{n+1} = (1 + 2h)y_n + h/2 - h t_n$ . Since  $y_0 = 1$ ,  $y_1 = (1 + 2h) + h/2 = (1 + 2h) + t_1/2$ . It is easy to show by mathematical induction, that  $y_n = (1 + 2h)^n + t_n/2$ . For  $t > 0$ , set  $h = t/n$  and thus  $t_n = t$ . Taking the limit, we find that  $\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} [(1 + 2t/n)^n + t/2] = e^{2t} + t/2$ . Hence pointwise convergence is proved.

**Section 2.8**


2. Let  $z = y - 3$  and  $\tau = t + 1$ . It follows that  $dz/d\tau = (dz/dt)(dt/d\tau) = dz/dt$ . Furthermore,  $dz/dt = dy/dt = 1 - y^3$ . Hence  $dz/d\tau = 1 - (z + 3)^3$ . The new initial condition is  $z(\tau = 0) = 0$ .

3. The approximating functions are defined recursively by  $\phi_{n+1}(t) = \int_0^t 2[\phi_n(s) + 1]ds$ . Setting  $\phi_0(t) = 0$ ,  $\phi_1(t) = 2t$ . Continuing,  $\phi_2(t) = 2t^2 + 2t$ ,  $\phi_3(t) = \frac{4}{3}t^3 + 2t^2 + 2t$ ,  $\phi_4(t) = \frac{2}{3}t^4 + \frac{4}{3}t^3 + 2t^2 + 2t, \dots$ . Given convergence, set

$$\begin{aligned} \phi(t) &= \phi_1(t) + \sum_{k=1}^{\infty} [\phi_{k+1}(t) - \phi_k(t)] \\ &= 2t + \sum_{k=2}^{\infty} \frac{a_k}{k!} t^k. \end{aligned}$$

Comparing coefficients,  $a_3/3! = 4/3$ ,  $a_4/4! = 2/3, \dots$ . It follows that  $a_3 = 8$ ,  $a_4 = 16$ , and so on. We find that in general, that  $a_k = 2^k$ . Hence

$$\begin{aligned} \phi(t) &= \sum_{k=1}^{\infty} \frac{2^k}{k!} t^k \\ &= e^{2t} - 1. \end{aligned}$$



5. The approximating functions are defined recursively by


$$\phi_{n+1}(t) = \int_0^t [-\phi_n(s)/2 + s]ds.$$

Setting  $\phi_0(t) = 0$ ,  $\phi_1(t) = t^2/2$ . Continuing,  $\phi_2(t) = t^2/2 - t^3/12$ ,  $\phi_3(t) = t^2/2 - t^3/12 + t^4/96$ ,  $\phi_4(t) = t^2/2 - t^3/12 + t^4/96 - t^5/960, \dots$ . Given convergence, set

$$\begin{aligned}\phi(t) &= \phi_1(t) + \sum_{k=1}^{\infty} [\phi_{k+1}(t) - \phi_k(t)] \\ &= t^2/2 + \sum_{k=3}^{\infty} \frac{a_k}{k!} t^k.\end{aligned}$$

Comparing coefficients,  $a_3/3! = -1/12$ ,  $a_4/4! = 1/96$ ,  $a_5/5! = -1/960$ ,  $\dots$ . We find that  $a_3 = -1/2$ ,  $a_4 = 1/4$ ,  $a_5 = -1/8$ ,  $\dots$ . In general,  $a_k = 2^{-k+1}$ . Hence

$$\begin{aligned}\phi(t) &= \sum_{k=2}^{\infty} \frac{2^{-k+2}}{k!} (-t)^k \\ &= 4e^{-t/2} + 2t - 4.\end{aligned}$$




6. The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t [\phi_n(s) + 1 - s] ds.$$

Setting  $\phi_0(t) = 0$ ,  $\phi_1(t) = t - t^2/2$ ,  $\phi_2(t) = t - t^3/6$ ,  $\phi_3(t) = t - t^4/24$ ,  $\phi_4(t) = t - t^5/120$ ,  $\dots$ . Given convergence, set

$$\begin{aligned}\phi(t) &= \phi_1(t) + \sum_{k=1}^{\infty} [\phi_{k+1}(t) - \phi_k(t)] \\ &= t - t^2/2 + [t^2/2 - t^3/6] + [t^3/6 - t^4/24] + \dots \\ &= t + 0 + 0 + \dots.\end{aligned}$$

Note that the terms can be rearranged, as long as the series converges *uniformly*.




8(a). The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t [s^2 \phi_n(s) - s] ds.$$

Set  $\phi_0(t) = 0$ . The iterates are given by  $\phi_1(t) = -t^2/2$ ,  $\phi_2(t) = -t^2/2 - t^5/10$ ,  $\phi_3(t) = -t^2/2 - t^5/10 - t^8/80$ ,  $\phi_4(t) = -t^2/2 - t^5/10 - t^8/80 - t^{11}/880, \dots$ . Upon inspection, it becomes apparent that

$$\begin{aligned} \phi_n(t) &= -t^2 \left[ \frac{1}{2} + \frac{t^3}{2 \cdot 5} + \frac{t^6}{2 \cdot 5 \cdot 8} + \dots + \frac{(t^3)^{n-1}}{2 \cdot 5 \cdot 8 \dots [2 + 3(n-1)]} \right] \\ &= -t^2 \sum_{k=1}^n \frac{(t^3)^{k-1}}{2 \cdot 5 \cdot 8 \dots [2 + 3(k-1)]}. \end{aligned}$$

(b).




The iterates appear to be converging.

9(a). The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t [s^2 + \phi_n^2(s)] ds.$$

Set  $\phi_0(t) = 0$ . The first three iterates are given by  $\phi_1(t) = t^3/3$ ,  $\phi_2(t) = t^3/3 + t^7/63$ ,  $\phi_3(t) = t^3/3 + t^7/63 + 2t^{11}/2079 + t^{15}/59535$ .

(b).




The iterates appear to be converging.

10(a). The approximating functions are defined recursively by

$$\phi_{n+1}(t) = \int_0^t [1 - \phi_n^3(s)] ds.$$

Set  $\phi_0(t) = 0$ . The first three iterates are given by  $\phi_1(t) = t$ ,  $\phi_2(t) = t - t^4/4$ ,  $\phi_3(t) = t - t^4/4 + 3t^7/28 - 3t^{10}/160 + t^{13}/833$ .

(b).



The approximations appear to be diverging.

12(a). The approximating functions are defined recursively by


$$\phi_{n+1}(t) = \int_0^t \left[ \frac{3s^2 + 4s + 2}{2(\phi_n(s) - 1)} \right] ds.$$

Note that  $1/(2y - 2) = -\frac{1}{2} \sum_{k=0}^6 y^k + O(y^7)$ . For computational purposes, replace the above iteration formula by

$$\phi_{n+1}(t) = -\frac{1}{2} \int_0^t \left[ (3s^2 + 4s + 2) \sum_{k=0}^6 \phi_n^k(s) \right] ds.$$

Set  $\phi_0(t) = 0$ . The first four approximations are given by  $\phi_1(t) = -t - t^2 - t^3/2$ ,  
 $\phi_2(t) = -t - t^2/2 + t^3/6 + t^4/4 - t^5/5 - t^6/24 + \dots$ ,  
 $\phi_3(t) = -t - t^2/2 + t^4/12 - 3t^5/20 + 4t^6/45 + \dots$ ,  
 $\phi_4(t) = -t - t^2/2 + t^4/8 - 7t^5/60 + t^6/15 + \dots$

(b).



The approximations appear to be converging to the exact solution,

$$\phi(t) = 1 - \sqrt{1 + 2t + 2t^2 + t^3}.$$

13. Note that  $\phi_n(0) = 0$  and  $\phi_n(1) = 1, \forall n \geq 1$ . Let  $a \in (0, 1)$ . Then  $\phi_n(a) = a^n$ . Clearly,  $\lim_{n \rightarrow \infty} a^n = 0$ . Hence the assertion is true.

14(a).  $\phi_n(0) = 0, \forall n \geq 1$ . Let  $a \in (0, 1]$ . Then  $\phi_n(a) = 2na e^{-na^2} = 2na/e^{na^2}$ . Using l'Hospital's rule,  $\lim_{z \rightarrow \infty} 2az/e^{az^2} = \lim_{z \rightarrow \infty} 1/ze^{az^2} = 0$ . Hence  $\lim_{n \rightarrow \infty} \phi_n(a) = 0$ .

(b).  $\int_0^1 2nx e^{-nx^2} dx = -e^{-nx^2} \Big|_0^1 = 1 - e^{-n}$ . Therefore,

$$\lim_{n \rightarrow \infty} \int_0^1 \phi_n(x) dx \neq \int_0^1 \lim_{n \rightarrow \infty} \phi_n(x) dx.$$

15. Let  $t$  be fixed, such that  $(t, y_1), (t, y_2) \in D$ . Without loss of generality, assume that  $y_1 < y_2$ . Since  $f$  is differentiable with respect to  $y$ , the mean value theorem asserts that  $\exists \xi \in (y_1, y_2)$  such that  $f(t, y_1) - f(t, y_2) = f_y(t, \xi)(y_1 - y_2)$ . Taking the absolute value of both sides,  $|f(t, y_1) - f(t, y_2)| = |f_y(t, \xi)| |y_1 - y_2|$ . Since, by assumption,  $\partial f / \partial y$  is continuous in  $D$ ,  $f_y$  attains a *maximum* on any closed and bounded subset of  $D$ .

Hence  $|f(t, y_1) - f(t, y_2)| \leq K |y_1 - y_2|$ .

16. For a *sufficiently small* interval of  $t$ ,  $\phi_{n-1}(t), \phi_n(t) \in D$ . Since  $f$  satisfies a Lipschitz condition,  $|f(t, \phi_n(t)) - f(t, \phi_{n-1}(t))| \leq K |\phi_n(t) - \phi_{n-1}(t)|$ . Here  $K = \max |f_y|$ .

17(a).  $\phi_1(t) = \int_0^t f(s, 0) ds$ . Hence  $|\phi_1(t)| \leq \int_0^{|t|} |f(s, 0)| ds \leq \int_0^{|t|} M ds = M|t|$ , in which  $M$  is the maximum value of  $|f(t, y)|$  on  $D$ .

(b). By definition,  $\phi_2(t) - \phi_1(t) = \int_0^t [f(s, \phi_1(s)) - f(s, 0)] ds$ . Taking the absolute value of both sides,  $|\phi_2(t) - \phi_1(t)| \leq \int_0^{|t|} |[f(s, \phi_1(s)) - f(s, 0)]| ds$ . Based on the results in Problems 16 and 17,  $|\phi_2(t) - \phi_1(t)| \leq \int_0^{|t|} K |\phi_1(s) - 0| ds \leq KM \int_0^{|t|} |s| ds$ . Evaluating the last integral, we obtain  $|\phi_2(t) - \phi_1(t)| \leq MK|t|^2/2$ .

(c). Suppose that

$$|\phi_i(t) - \phi_{i-1}(t)| \leq \frac{MK^{i-1}|t|^i}{i!}$$

for some  $i \geq 1$ . By definition,  $\phi_{i+1}(t) - \phi_i(t) = \int_0^t [f(t, \phi_i(s)) - f(s, \phi_{i-1}(s))] ds$ . It follows that

$$\begin{aligned} |\phi_{i+1}(t) - \phi_i(t)| &\leq \int_0^{|t|} |f(s, \phi_i(s)) - f(s, \phi_{i-1}(s))| ds \\ &\leq \int_0^{|t|} K |\phi_i(s) - \phi_{i-1}(s)| ds \\ &\leq \int_0^{|t|} K \frac{MK^{i-1}|s|^i}{i!} ds \\ &= \frac{MK^i |t|^{i+1}}{(i+1)!} \leq \frac{MK^i h^{i+1}}{(i+1)!}. \end{aligned}$$

Hence, by mathematical induction, the assertion is true.

18(a). Use the triangle inequality,  $|a + b| \leq |a| + |b|$ .

(b). For  $|t| \leq h$ ,  $|\phi_1(t)| \leq Mh$ , and  $|\phi_n(t) - \phi_{n-1}(t)| \leq MK^{n-1}h^n/(n!)$ . Hence

$$\begin{aligned} |\phi_n(t)| &\leq M \sum_{i=1}^n \frac{K^{i-1}h^i}{i!} \\ &= \frac{M}{K} \sum_{i=1}^n \frac{(Kh)^i}{i!}. \end{aligned}$$

(c). The sequence of partial sums in (b) converges to  $\frac{M}{K}(e^{Kh} - 1)$ . By the *comparison test*, the sums in (a) also converge. Furthermore, the sequence  $|\phi_n(t)|$  is *bounded*, and hence has a convergent subsequence. Finally, since individual terms of the series must tend to zero,  $|\phi_n(t) - \phi_{n-1}(t)| \rightarrow 0$ , and it follows that the sequence  $|\phi_n(t)|$  is convergent.

19(a). Let  $\phi(t) = \int_0^t f(s, \phi(s))ds$  and  $\psi(t) = \int_0^t f(s, \psi(s))ds$ . Then by *linearity of the integral*,  $\phi(t) - \psi(t) = \int_0^t [f(s, \phi(s)) - f(s, \psi(s))]ds$ .

(b). It follows that  $|\phi(t) - \psi(t)| \leq \int_0^t |f(s, \phi(s)) - f(s, \psi(s))|ds$ .

(c). We know that  $f$  satisfies a Lipschitz condition,

$$|f(t, y_1) - f(t, y_2)| \leq K |y_1 - y_2|,$$

based on  $|\partial f / \partial y| \leq K$  in  $D$ . Therefore,

$$\begin{aligned} |\phi(t) - \psi(t)| &\leq \int_0^t |f(s, \phi(s)) - f(s, \psi(s))|ds \\ &\leq \int_0^t K |\phi(s) - \psi(s)|ds. \end{aligned}$$



**Section 2.9**

1. Writing the equation for each  $n \geq 0$ ,  $y_1 = -0.9 y_0$ ,  $y_2 = -0.9 y_1$ ,  $y_3 = -0.9 y_2$  and so on, it is apparent that  $y_n = (-0.9)^n y_0$ . The terms constitute an *alternating series*, which converge to *zero*, regardless of  $y_0$ .

3. Write the equation for each  $n \geq 0$ ,  $y_1 = \sqrt{3} y_0$ ,  $y_2 = \sqrt{4/2} y_1$ ,  $y_3 = \sqrt{5/3} y_2, \dots$ . Upon substitution, we find that  $y_2 = \sqrt{(4 \cdot 3)/2} y_1$ ,  $y_3 = \sqrt{(5 \cdot 4 \cdot 3)/(3 \cdot 2)} y_0, \dots$ . It can be proved by mathematical induction, that

$$\begin{aligned} y_n &= \frac{1}{\sqrt{2}} \sqrt{\frac{(n+2)!}{n!}} y_0 \\ &= \frac{1}{\sqrt{2}} \sqrt{(n+1)(n+2)} y_0. \end{aligned}$$

This sequence is *divergent*, except for  $y_0 = 0$ .

4. Writing the equation for each  $n \geq 0$ ,  $y_1 = -y_0$ ,  $y_2 = y_1$ ,  $y_3 = -y_2$ ,  $y_4 = y_3$ , and so on, it can be shown that

$$y_n = \begin{cases} y_0 & , \text{ for } n = 4k \text{ or } n = 4k - 1 \\ -y_0 & , \text{ for } n = 4k - 2 \text{ or } n = 4k - 3 \end{cases}$$

The sequence is convergent *only* for  $y_0 = 0$ .

6. Writing the equation for each  $n \geq 0$ ,

$$\begin{aligned} y_1 &= 0.5 y_0 + 6 \\ y_2 &= 0.5 y_1 + 6 = 0.5(0.5 y_0 + 6) + 6 = (0.5)^2 y_0 + 6 + (0.5)6 \\ y_3 &= 0.5 y_2 + 6 = 0.5(0.5 y_1 + 6) + 6 = (0.5)^3 y_0 + 6[1 + (0.5) + (0.5)^2] \\ &\vdots \\ y_n &= (0.5)^n y_0 + 12[1 - (0.5)^n] \end{aligned}$$

which can be verified by mathematical induction. The sequence is convergent for all  $y_0$ , and in fact  $y_n \rightarrow 12$ .

7. Let  $y_n$  be the balance at the end of the  $n$ -th day. Then  $y_{n+1} = (1 + r/365) y_n$ . The solution of this difference equation is  $y_n = (1 + r/365)^n y_0$ , in which  $y_0$  is the initial balance. At the end of *one year*, the balance is  $y_{365} = (1 + r/365)^{365} y_0$ . Given that  $r = .07$ ,  $y_{365} = (1 + r/365)^{365} y_0 = 1.0725 y_0$ . Hence the effective annual yield is  $(1.0725 y_0 - y_0)/y_0 = 7.25\%$ .

8. Let  $y_n$  be the balance at the end of the  $n$ -th month. Then  $y_{n+1} = (1 + r/12) y_n + 25$ . As in the previous solutions, we have

$$y_n = \rho^n \left[ y_0 - \frac{25}{1 - \rho} \right] + \frac{25}{1 - \rho},$$

in which  $\rho = (1 + r/12)$ . Here  $r$  is the annual interest rate, given as 8%. Therefore  $y_{36} = (1.0066)^{36} \left[ 1000 + \frac{(12)25}{r} \right] - \frac{(12)25}{r} = 2,283.63$  dollars.

9. Let  $y_n$  be the balance due at the end of the  $n$ -th month. The appropriate difference equation is  $y_{n+1} = (1 + r/12) y_n - P$ . Here  $r$  is the annual interest rate and  $P$  is the monthly payment. The solution, in terms of the amount borrowed, is given by

$$y_n = \rho^n \left[ y_0 + \frac{P}{1 - \rho} \right] - \frac{P}{1 - \rho},$$

in which  $\rho = (1 + r/12)$  and  $y_0 = 8,000$ . To figure out the monthly payment,  $P$ , we require that  $y_{36} = 0$ . That is,

$$\rho^{36} \left[ y_0 + \frac{P}{1 - \rho} \right] = \frac{P}{1 - \rho}.$$

After the specified amounts are substituted, we find the  $P = \$258.14$ .

11. Let  $y_n$  be the balance due at the end of the  $n$ -th month. The appropriate difference equation is  $y_{n+1} = (1 + r/12) y_n - P$ , in which  $r = .09$  and  $P$  is the monthly payment. The initial value of the mortgage is  $y_0 = 100,000$  dollars. Then the balance due at the end of the  $n$ -th month is

$$y_n = \rho^n \left[ y_0 + \frac{P}{1 - \rho} \right] - \frac{P}{1 - \rho}.$$

where  $\rho = (1 + r/12)$ . In terms of the specified values,

$$y_n = (0.0075)^n \left[ 10^5 - \frac{12P}{r} \right] + \frac{12P}{r}.$$

Setting  $n = 30(12) = 360$ , and  $y_{360} = 0$ , we find that  $P = 804.62$  dollars. For the monthly payment corresponding to a 20 year mortgage, set  $n = 240$  and  $y_{240} = 0$ .

12. Let  $y_n$  be the balance due at the end of the  $n$ -th month, with  $y_0$  the initial value of the mortgage. The appropriate difference equation is  $y_{n+1} = (1 + r/12) y_n - P$ , in which  $r = 0.1$  and  $P = 900$  dollars is the *maximum* monthly payment. Given that the life of the mortgage is 20 years, we require that  $y_{240} = 0$ . The balance due at the end of the  $n$ -th month is





$$y_n = \rho^n \left[ y_0 + \frac{P}{1 - \rho} \right] - \frac{P}{1 - \rho}.$$

In terms of the specified values for the parameters, the solution of


$$(.00833)^{240} \left[ y_0 - \frac{12(1000)}{0.1} \right] = - \frac{12(1000)}{0.1}$$

is  $y_0 = 103,624.62$  dollars.

15.



16. For example, take  $\rho = 3.5$  and  $u_0 = 1.1$  :




19(a).  $\delta_2 = (\rho_2 - \rho_1)/(\rho_3 - \rho_2) = (3.449 - 3)/(3.544 - 3.449) = 4.7263$ .

(b).  $\% \text{ diff} = \frac{|\delta - \delta_2|}{\delta} \times 100 = \frac{|4.6692 - 4.7363|}{4.6692} \times 100 \approx 1.22 \%$ .

(c). Assuming  $(\rho_3 - \rho_2)/(\rho_4 - \rho_3) = \delta$ ,  $\rho_4 \approx 3.5643$

(d). A period 16 solutions appears near  $\rho \approx 3.565$ .



(e). Note that  $(\rho_{n+1} - \rho_n) = \delta_n^{-1}(\rho_n - \rho_{n-1})$ . With the assumption that  $\delta_n = \delta$ , we have  $(\rho_{n+1} - \rho_n) = \delta^{-1}(\rho_n - \rho_{n-1})$ , which is of the form  $y_{n+1} = \alpha y_n$ ,  $n \geq 3$ . It follows that  $(\rho_k - \rho_{k-1}) = \delta^{3-k}(\rho_3 - \rho_2)$  for  $k \geq 4$ . Then

$$\begin{aligned} \rho_n &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2) + (\rho_4 - \rho_3) + \dots + (\rho_n - \rho_{n-1}) \\ &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2)[1 + \delta^{-1} + \delta^{-2} + \dots + \delta^{3-n}] \\ &= \rho_1 + (\rho_2 - \rho_1) + (\rho_3 - \rho_2) \left[ \frac{1 - \delta^{4-n}}{1 - \delta^{-1}} \right]. \end{aligned}$$

Hence  $\lim_{n \rightarrow \infty} \rho_n = \rho_2 + (\rho_3 - \rho_2) \left[ \frac{\delta}{\delta - 1} \right]$ . Substitution of the appropriate values yields

$$\lim_{n \rightarrow \infty} \rho_n = 3.5699$$

**Miscellaneous Problems**

1. Linear  $[y = c/x^2 + x^3/5]$ .
2. Homogeneous  $[\arctan(y/x) - \ln\sqrt{x^2 + y^2} = c]$ .
3. Exact  $[x^2 + xy - 3y - y^3 = 0]$ .
4. Linear in  $x(y)$   $[x = ce^y + ye^y]$ .
5. Exact  $[x^2y + xy^2 + x = c]$ .
6. Linear  $[y = x^{-1}(1 - e^{1-x})]$ .
7. Let  $u = x^2$   $[x^2 + y^2 + 1 = ce^{y^2}]$ .
8. Linear  $[y = (4 + \cos 2 - \cos x)/x^2]$ .
9. Exact  $[x^2y + x + y^2 = c]$ .
10.  $\mu = \mu(x)$   $[y^2/x^3 + y/x^2 = c]$ .
11. Exact  $[x^3/3 + xy + e^y = c]$ .
12. Linear  $[y = ce^{-x} + e^{-x}\ln(1 + e^x)]$ .
13. Homogeneous  $[2\sqrt{y/x} - \ln|x| = c]$ .
14. Exact/Homogeneous  $[x^2 + 2xy + 2y^2 = 34]$ .
15. Separable  $[y = c/\cosh^2(x/2)]$ .
16. Homogeneous  $[(2/\sqrt{3})\arctan[(2y - x)/\sqrt{3}x] - \ln|x| = c]$ .
17. Linear  $[y = ce^{3x} - e^{2x}]$ .
18. Linear/Homogeneous  $[y = cx^{-2} - x]$ .
19.  $\mu = \mu(x)$   $[3y - 2xy^3 - 10x = 0]$ .
20. Separable  $[e^x + e^{-y} = c]$ .
21. Homogeneous  $[e^{-y/x} + \ln|x| = c]$ .
22. Separable  $[y^3 + 3y - x^3 + 3x = 2]$ .
23. Bernoulli  $[1/y = -x \int x^{-2}e^{2x} dx + cx]$ .
24. Separable  $[\sin^2x \sin y = c]$ .
25. Exact  $[x^2/y + \arctan(y/x) = c]$ .
26.  $\mu = \mu(x)$   $[x^2 + 2x^2y - y^2 = c]$ .
27.  $\mu = \mu(x)$   $[\sin x \cos 2y - \frac{1}{2}\sin^2x = c]$ .
28. Exact  $[2xy + xy^3 - x^3 = c]$ .
29. Homogeneous  $[\arcsin(y/x) - \ln|x| = c]$ .
30. Linear in  $x(y)$   $[xy^2 - \ln|y| = 0]$ .
31. Separable  $[x + \ln|x| + x^{-1} + y - 2\ln|y| = c]$ .
32.  $\mu = \mu(y)$   $[x^3y^2 + xy^3 = -4]$ .