



INTERMEDIATE ALGEBRA

LIAL / HORNSBY / MCGINNIS




thirteenth edition

Get the Most Out of MyLab Math

When it comes to developmental math, we know one size does not fit all. Pearson’s solutions offer market-leading content written by our author-educators, tightly integrated with the #1 choice in digital learning-MyLab Math. MyLab Math is the teaching and learning platform that empowers instructors to reach every student. By combining trusted author content with digital tools and a flexible platform, MyLab Math personalizes the learning experience and improves results for each student.


- **Flexible Platform—**

Your course is unique. Whether you’d like to build your own assignments, structure students’ work with a learning path, or set prerequisites, you have the flexibility to easily create *your* course to fit *your* needs.



- **Personalized Learning—** Each student learns at a different pace. Personalized learning pinpoints the areas each student needs to practice, giving every student the support they need—when and where they need it—to be successful. A variety of options are available to personalize learning in MyLab Math:

- With **Personalized Homework**, students take a quiz or test and receive a subsequent homework assignment that is personalized based on their performance. This way, students can focus on just the topics they have not yet mastered.



- **Skill Builder** offers adaptive practice that is designed to increase students’ ability to complete their assignments. By monitoring student performance on their homework, Skill Builder adapts to each student’s needs and provides just-in-time, in-assignment practice to help them improve their proficiency of key learning objectives. Available for select MyLab™ courses.

Visit pearson.com/mylab/math and click **Get Trained** to make sure you’re getting the most out of MyLab Math.

EDITION

13

Intermediate Algebra

Margaret L. Lial

American River College

John Hornsby

University of New Orleans

Terry McGinnis

Vice President, Courseware Portfolio Management: Chris Hoag
Director, Courseware Portfolio Management: Michael Hirsch
Courseware Portfolio Manager: Karen Montgomery
Courseware Portfolio Assistant: Kayla Shearns
Managing Producer: Scott Disanno
Content Producer: Lauren Morse
Producers: Stacey Miller and Noelle Saligumba
Managing Producer: Vicki Dreyfus
Associate Content Producer, TestGen: Rajinder Singh
Content Managers, MathXL: Eric Gregg and Dominick Franck
Manager, Courseware QA: Mary Durnwald
Senior Product Marketing Manager: Alicia Frankel
Product Marketing Assistant: Brooke Imbornone
Senior Author Support/Technology Specialist: Joe Vetere
Full Service Vendor, Cover Design, Composition: Pearson CSC
Full Service Project Management: Pearson CSC (Carol Merrigan)
Cover Image: Borchee/E+/Getty Images

Copyright © 2020, 2016, 2012 by Pearson Education, Inc. 221 River Street, Hoboken, NJ 07030. All Rights Reserved. Printed in the United States of America. This publication is protected by copyright, and permission should be obtained from the publisher prior to any prohibited reproduction, storage in a retrieval system, or transmission in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise. For information regarding permissions, request forms, and the appropriate contacts within the Pearson Education Global Rights & Permissions department, please visit www.pearsoned.com/permissions/.

Acknowledgments of third-party content appear on page C-1, which constitutes an extension of this copyright page.

PEARSON, ALWAYS LEARNING, MyLab™ Math, MathXL, and TestGen are exclusive trademarks in the U.S. and/or other countries owned by Pearson Education, Inc. or its affiliates.

Unless otherwise indicated herein, any third-party trademarks that may appear in this work are the property of their respective owners and any references to third-party trademarks, logos or other trade dress are for demonstrative or descriptive purposes only. Such references are not intended to imply any sponsorship, endorsement, authorization, or promotion of Pearson's products by the owners of such marks, or any relationship between the owner and Pearson Education, Inc. or its affiliates, authors, licensees or distributors.

Library of Congress Cataloging-in-Publication Data

Names: Lial, Margaret L., author. | Hornsby, John, 1949- author. | McGinnis, Terry, author.
Title: Intermediate algebra / Margaret L. Lial (American River College), John Hornsby (University of New Orleans), Terry McGinnis.
Description: 13th edition. | Boston : Pearson, [2020] | Includes index.
Identifiers: LCCN 2018048199 | ISBN 9780134895987 (student edition) | ISBN 0134895983 (student edition) | ISBN 9780134996752 (annotated instructor edition) | ISBN 0134996755 (annotated instructor edition)
Subjects: LCSH: Algebra--Textbooks.
Classification: LCC QA152.3 .L534 2020 | DDC 512.9--dc23
LC record available at <https://lccn.loc.gov/2018048199>

Study Skills S-1

- | | |
|---------------------------------------------------|---------------------------------------------------------------|
| STUDY SKILL 1 Using Your Math Text S-1 | STUDY SKILL 6 Managing Your Time S-6 |
| STUDY SKILL 2 Reading Your Math Text S-2 | STUDY SKILL 7 Reviewing a Chapter S-7 |
| STUDY SKILL 3 Taking Lecture Notes S-3 | STUDY SKILL 8 Taking Math Tests S-8 |
| STUDY SKILL 4 Completing Your Homework S-4 | STUDY SKILL 9 Analyzing Your Test Results S-9 |
| STUDY SKILL 5 Using Study Cards S-5 | STUDY SKILL 10 Preparing for Your Math Final Exam S-10 |

R Review of the Real Number System 1

- | | |
|---------------------------------------------------------|------------------------------------------|
| R.1 Fractions, Decimals, and Percents 1 | R.5 Properties of Real Numbers 45 |
| R.2 Basic Concepts from Algebra 14 | Chapter R Summary 52 |
| R.3 Operations on Real Numbers 26 | Chapter R Test 54 |
| R.4 Exponents, Roots, and Order of Operations 36 | |

1 Linear Equations, Inequalities, and Applications 55

- | | |
|------------------------------------------------------------------|-------------------------------------------------------------------------------------------|
| 1.1 Linear Equations in One Variable 56 | 1.7 Absolute Value Equations and Inequalities 125 |
| 1.2 Formulas and Percent 65 | SUMMARY EXERCISES Solving Linear and Absolute Value Equations and Inequalities 136 |
| 1.3 Applications of Linear Equations 78 | Chapter 1 Summary 137 |
| 1.4 Further Applications of Linear Equations 92 | Chapter 1 Review Exercises 142 |
| SUMMARY EXERCISES Applying Problem-Solving Techniques 101 | Chapter 1 Mixed Review Exercises 145 |
| 1.5 Linear Inequalities in One Variable 103 | Chapter 1 Test 146 |
| 1.6 Set Operations and Compound Inequalities 116 | Chapters R and 1 Cumulative Review Exercises 148 |

2 Linear Equations, Graphs, and Functions 149

- | | |
|--------------------------------------------------------------------|--------------------------------------------------------|
| 2.1 Linear Equations in Two Variables 150 | 2.5 Introduction to Relations and Functions 199 |
| 2.2 The Slope of a Line 161 | 2.6 Function Notation and Linear Functions 210 |
| 2.3 Writing Equations of Lines 176 | Chapter 2 Summary 219 |
| SUMMARY EXERCISES Finding Slopes and Equations of Lines 191 | Chapter 2 Review Exercises 222 |
| 2.4 Linear Inequalities in Two Variables 192 | Chapter 2 Mixed Review Exercises 224 |
| | Chapter 2 Test 225 |
| | Chapters R–2 Cumulative Review Exercises 227 |

3 Systems of Linear Equations 229

- 3.1 Systems of Linear Equations in Two Variables 230
- 3.2 Systems of Linear Equations in Three Variables 245
- 3.3 Applications of Systems of Linear Equations 254
- Chapter 3 Summary 270
- Chapter 3 Review Exercises 273
- Chapter 3 Mixed Review Exercises 275
- Chapter 3 Test 276
- Chapters R–3 Cumulative Review Exercises 277

4 Exponents, Polynomials, and Polynomial Functions 279

- 4.1 Integer Exponents 280
- 4.2 Scientific Notation 290
- 4.3 Adding and Subtracting Polynomials 296
- 4.4 Polynomial Functions, Graphs, and Composition 302
- 4.5 Multiplying Polynomials 315
- 4.6 Dividing Polynomials 324
- Chapter 4 Summary 331
- Chapter 4 Review Exercises 334
- Chapter 4 Mixed Review Exercises 337
- Chapter 4 Test 337
- Chapters R–4 Cumulative Review Exercises 338

5 Factoring 341

- 5.1 Greatest Common Factors and Factoring by Grouping 342
- 5.2 Factoring Trinomials 348
- 5.3 Special Factoring 356
- 5.4 A General Approach to Factoring 363
- 5.5 Solving Quadratic Equations Using the Zero-Factor Property 367
- Chapter 5 Summary 376
- Chapter 5 Review Exercises 379
- Chapter 5 Mixed Review Exercises 381
- Chapter 5 Test 382
- Chapters R–5 Cumulative Review Exercises 382

6 Rational Expressions and Functions 385

- 6.1 Rational Expressions and Functions; Multiplying and Dividing 386
- 6.2 Adding and Subtracting Rational Expressions 396
- 6.3 Complex Fractions 405
- 6.4 Equations with Rational Expressions and Graphs 411
- 6.5 Applications of Rational Expressions 421
- 6.6 Variation 433
- Chapter 6 Summary 443
- Chapter 6 Review Exercises 447
- Chapter 6 Mixed Review Exercises 449
- Chapter 6 Test 450
- Chapters R–6 Cumulative Review Exercises 452
- SUMMARY EXERCISES** Simplifying Rational Expressions vs. Solving Rational Equations 419

7 Roots, Radicals, and Root Functions 455

- 7.1 Radical Expressions and Graphs 456
- 7.2 Rational Exponents 464
- 7.3 Simplifying Radicals, the Distance Formula, and Circles 473
- 7.4 Adding and Subtracting Radical Expressions 487
- 7.5 Multiplying and Dividing Radical Expressions 492
- SUMMARY EXERCISES** Performing Operations with Radicals and Rational Exponents 502
- 7.6 Solving Equations with Radicals 503
- 7.7 Complex Numbers 510
- Chapter 7 Summary 518
- Chapter 7 Review Exercises 523
- Chapter 7 Mixed Review Exercises 526
- Chapter 7 Test 527
- Chapters R–7 Cumulative Review Exercises 528

8 Quadratic Equations, Inequalities, and Functions 531

- 8.1 The Square Root Property and Completing the Square 532
- 8.2 The Quadratic Formula 541
- 8.3 Equations That Lead to Quadratic Methods 549
- SUMMARY EXERCISES** Applying Methods for Solving Quadratic Equations 560
- 8.4 Formulas and Further Applications 561
- 8.5 Graphs of Quadratic Functions 570
- 8.6 More about Parabolas and Their Applications 579
- 8.7 Polynomial and Rational Inequalities 590
- Chapter 8 Summary 599
- Chapter 8 Review Exercises 603
- Chapter 8 Mixed Review Exercises 606
- Chapter 8 Test 607
- Chapters R–8 Cumulative Review Exercises 609

9 Inverse, Exponential, and Logarithmic Functions 611

- 9.1 Inverse Functions 612
- 9.2 Exponential Functions 621
- 9.3 Logarithmic Functions 630
- 9.4 Properties of Logarithms 639
- 9.5 Common and Natural Logarithms 646
- 9.6 Exponential and Logarithmic Equations; Further Applications 654
- Chapter 9 Summary 663
- Chapter 9 Review Exercises 667
- Chapter 9 Mixed Review Exercises 670
- Chapter 9 Test 672
- Chapters R–9 Cumulative Review Exercises 673

10 Nonlinear Functions, Conic Sections, and Nonlinear Systems 677

- 10.1 Additional Graphs of Functions 678
- 10.2 Circles Revisited and Ellipses 684
- 10.3 Hyperbolas and Functions Defined by Radicals 694
- 10.4 Nonlinear Systems of Equations 702
- 10.5 Second-Degree Inequalities and Systems of Inequalities 708
- Chapter 10 Summary 715
- Chapter 10 Review Exercises 718
- Chapter 10 Mixed Review Exercises 720
- Chapter 10 Test 721
- Chapters R–10 Cumulative Review Exercises 722

11 Further Topics in Algebra 725

- 11.1** Sequences and Series 726
- 11.2** Arithmetic Sequences 732
- 11.3** Geometric Sequences 740
- 11.4** The Binomial Theorem 750

- Chapter 11 Summary** 756
- Chapter 11 Review Exercises** 759
- Chapter 11 Mixed Review Exercises** 760
- Chapter 11 Test** 761
- Chapters R–11 Cumulative Review Exercises** 762

Appendix Synthetic Division 765

- Answers to Selected Exercises A-1
- Photo Credits C-1
- Index I-1

WELCOME TO THE 13TH EDITION

The first edition of Marge Lial’s *Intermediate Algebra* was published in 1972, and now we are pleased to present the 13th edition—with the same successful, well-rounded framework that was established 50 years ago and updated to meet the needs of today’s students and professors. The names Lial and Miller, two faculty members from American River College in Sacramento, California, have become synonymous with excellence in Developmental Mathematics, Precalculus, Finite Mathematics, and Applications-Based Calculus.

With Chuck Miller’s passing in 1986, Marge Lial was joined by a team of carefully selected coauthors who partnered with her. John Hornsby (University of New Orleans) joined Marge in this capacity in 1992, and in 1999, Terry McGinnis became part of this developmental author team. Since Marge’s passing in 2012, John and Terry have dedicated themselves to carrying on the Lial/Miller legacy.

In the preface to the first edition of *Intermediate Algebra*, Marge Lial wrote

“ . . . the strongest theme . . . is a combination of readability and suitability for the book’s intended audience: students who are not completely self-confident in mathematics as they come to the course, but who must be self-confident and proficient . . . by the end of the course.”

Today’s Lial author team upholds these same standards. With the publication of the 13th edition of *Intermediate Algebra*, we proudly present a complete course program for students who need developmental algebra. Revisions to the core text, working in concert with such innovations in the MyLab Math course as Skill Builder and Learning Catalytics, combine to provide superior learning opportunities appropriate for all types of courses (traditional, hybrid, online).

We hope you enjoy using it as much as we have enjoyed writing it. We welcome any feedback that you have as you review and use this text.

WHAT’S NEW IN THIS EDITION?

We are pleased to offer the following new features and resources in the text and MyLab.

IMPROVED STUDY SKILLS These special activities are now grouped together at the front of the text, prior to Chapter R. **Study Skills Reminders** that refer students to specific Study Skills are found liberally throughout the text. Many Study Skills

now include a *Now Try This* section to help students implement the specific skill.

REVISED EXPOSITION With each edition of the text, we continue to polish and improve discussions and presentations of topics to increase readability and student understanding. This edition is no exception.

NEW FIGURES AND DIAGRAMS For visual learners, we have included more than 50 new mathematical figures, graphs, and diagrams, including several new “hand drawn” style graphs. These are meant to suggest what a student who is graphing with paper and pencil should obtain. We use this style when introducing a particular type of graph for the first time.

ENHANCED USE OF PEDAGOGICAL COLOR We have thoroughly reviewed the use of pedagogical color in discussions and examples and have increased its use whenever doing so would enhance concept development, emphasize important steps, or highlight key procedures.

INCREASED Concept Check AND WHAT WENT WRONG? EXERCISES The number of **Concept Check** exercises, which facilitate students’ mathematical thinking and conceptual understanding, and which begin each exercise set, has been increased. We have also more than doubled the number of **WHAT WENT WRONG?** exercises that highlight common student errors.

INCREASED RELATING CONCEPTS EXERCISES We have doubled the number of these flexible groups of exercises, which are located at the end of many exercise sets. These sets of problems were specifically written to help students tie concepts together, compare and contrast ideas, identify and describe patterns, and extend concepts to new situations. They may be used by individual students or by pairs or small groups working collaboratively. All answers to these exercises appear in the student answer section.

ENHANCED MYLAB MATH RESOURCES MyLab exercise coverage in the revision has been expanded, and video coverage has also been expanded and updated to a modern format for today’s students. **WHAT WENT WRONG?** problems and all **RELATING CONCEPTS** exercise sets (both even- and odd-numbered problems) are now assignable in MyLab Math.

SKILL BUILDER These exercises offer just-in-time additional adaptive practice in MyLab Math. The adaptive engine tracks student performance and delivers, to each individual, questions that adapt to his or her level of understanding. This new feature enables instructors to assign fewer questions for

homework, allowing students to complete as many or as few questions as they need.

LEARNING CATALYTICS This new student response tool uses students' own devices to engage them in the learning process. Problems that draw on prerequisite skills are included at the beginning of each section to gauge student readiness for the section. Accessible through MyLab Math and customizable to instructors' specific needs, these problems can be used to generate class discussion, promote peer-to-peer learning, and provide real-time feedback to instructors. More information can be found via the Learning Catalytics link in MyLab Math. Specific exercises notated in the text can be found by searching LialIntermediate# where the # is the chapter number.

CONTENT CHANGES

Specific content changes include the following:

- **Exercise sets** have been scrutinized and updated with a renewed focus on conceptual understanding and skill development. Even and odd pairing of the exercises, an important feature of the text, has been carefully reviewed.
- **Real world data** in all examples and exercises and their accompanying graphs has been updated.
- **An increased emphasis on fractions, decimals, and percents** appears throughout the text. **Chapter R** begins with a new section that thoroughly reviews these topics. And we have included an **all-new set of Cumulative Review Exercises**, many of which focus on fractions, decimals, and percents, at the end of Chapter 1. Sets of Cumulative Review Exercises in subsequent chapters now begin with new exercises that review skills related to these topics.
- **Solution sets of linear inequalities in Sections 1.5–1.7** are now graphed first before writing them using interval notation.
- **Scientific notation is covered in a separate section in Chapter 4.**
- **Chapters 10 and 11 on Nonlinear Functions, Conic Sections, Nonlinear Systems, and Further Topics in Algebra**, previously available online in MyLab Math, are now included in the text. The material has been fully revised and updated.
- **Presentations of the following topics have been enhanced and expanded**, often including new examples and exercises:
 - Evaluating exponential expressions (Section R.4)
 - Geometric interpretation of slope as rise/run (Section 2.2)
 - Solving systems of linear equations in three variables (Section 3.2)

Identifying functions and domains from equations (Section 2.6)

Determining strategies for factoring polynomials (Section 5.4)

Solving quadratic equations with double solutions (Section 5.5)

Solving rational equations (Section 6.4)

Concepts and relationships among real numbers, nonreal complex numbers, and imaginary numbers; simplifying powers of i (Section 7.7)

Solving quadratic equations using the quadratic formula (Section 8.2)

Solving exponential and logarithmic equations (Sections 9.2, 9.3)

LIAL DEVELOPMENTAL HALLMARK FEATURES

We have enhanced the following popular features, each of which is designed to increase ease of use by students and/or instructors.

- **Emphasis on Problem-Solving** We introduce our six-step problem-solving method in Chapter 2 and integrate it throughout the text. The six steps, *Read, Assign a Variable, Write an Equation, Solve, State the Answer,* and *Check*, are emphasized in boldface type and repeated in examples and exercises to reinforce the problem-solving process for students. We also provide students with **PROBLEM-SOLVING HINT** boxes that feature helpful problem-solving tips and strategies.
- **Helpful Learning Objectives** We begin each section with clearly stated, numbered objectives, and the included material is directly keyed to these objectives so that students and instructors know exactly what is covered in each section.
- **Cautions and Notes** One of the most popular features of previous editions is our inclusion of information marked **CAUTION** and **NOTE** to warn students about common errors and to emphasize important ideas throughout the exposition. The updated text design makes them easy to spot.
- **Comprehensive Examples** The new edition features a multitude of step-by-step, worked-out examples that include pedagogical color, helpful side comments, and special pointers. We give special attention to checking example solutions—more checks, designated using a special **CHECK** tag and ✓, are included than in past editions.
- **More Pointers** There are more pointers in examples and discussions throughout this edition of the text. They

provide students with important on-the-spot reminders, as well as warnings about common pitfalls.

- **Numerous Now Try Problems** These margin exercises, with answers immediately available at the bottom of the page, have been carefully written to correspond to every example in the text. This key feature allows students to immediately practice the material in preparation for the exercise sets.
- **Updated Figures, Photos, and Hand-Drawn Graphs** Today’s students are more visually oriented than ever. As a result, we provide detailed mathematical figures, diagrams, tables, and graphs, including a “hand-drawn” style of graphs, whenever possible. We have incorporated depictions of well-known mathematicians, as well as appealing photos to accompany applications in examples and exercises.
- **Relevant Real-Life Applications** We include many new or updated applications from fields such as business, pop culture, sports, technology, and the health sciences that show the relevance of algebra to daily life.
- **Extensive and Varied Exercise Sets** The text contains a wealth of exercises to provide students with opportunities to practice, apply, connect, review, and extend the skills they are learning. Numerous illustrations, tables, graphs, and photos help students visualize the problems they are solving. Problem types include skill building and writing exercises, as well as applications, matching, true/false, multiple-choice, and fill-in-the-blank problems. Special types of exercises include **Concept Check**, **WHAT WENT WRONG?**, **Extending Skills**, and **RELATING CONCEPTS**.
- **Special Summary Exercises** We include a set of these popular in-chapter exercises in every chapter. They provide students with the all-important *mixed review problems* they need to master topics and often include summaries of solution methods and/or additional examples.
- **Extensive Review Opportunities** We conclude each chapter with the following review components:

A **Chapter Summary** that features a helpful list of **Key Terms** organized by section, **New Symbols**, a **Test Your Word Power** vocabulary quiz (with answers immediately following), and a **Quick Review** of each section’s main concepts, complete with additional examples.

A comprehensive set of **Chapter Review Exercises**, keyed to individual sections for easy student reference.

A set of **Mixed Review Exercises** that helps students further synthesize concepts and skills.

A **Chapter Test** that students can take under test conditions to see how well they have mastered the chapter material.

A set of **Cumulative Review Exercises** for ongoing review that covers material going back to Chapter R.

- **Comprehensive Glossary** The online Glossary includes key terms and definitions (with section references) from throughout the text.

ACKNOWLEDGMENTS

The comments, criticisms, and suggestions of users, non-users, instructors, and students have positively shaped this text over the years, and we are most grateful for the many responses we have received. The feedback gathered for this edition was particularly helpful.

We especially wish to thank the following individuals who provided invaluable suggestions.

Barbara Aaker, *Community College of Denver*
 Kim Bennekin, *Georgia Perimeter College*
 Dixie Blackinton, *Weber State University*
 Eun Cha, *College of Southern Nevada, Charleston*
 Callie Daniels, *St. Charles Community College*
 Cheryl Davids, *Central Carolina Technical College*
 Robert Diaz, *Fullerton College*
 Chris Diorietes, *Fayetteville Technical Community College*
 Sylvia Dreyfus, *Meridian Community College*
 Sabine Eggleston, *Edison State College*
 LaTonya Ellis, *Bishop State Community College*
 Beverly Hall, *Fayetteville Technical Community College*
 Loretta Hart, *NHTI, Concord’s Community College*
 Sandee House, *Georgia Perimeter College*
 Joe Howe, *St. Charles Community College*
 Lynette King, *Gadsden State Community College*
 Linda Kodama, *Windward Community College*
 Carlea McAvoy, *South Puget Sound Community College*
 James Metz, *Kapi’olani Community College*
 Jean Millen, *Georgia Perimeter College*
 Molly Misko, *Gadsden State Community College*
 Charles Patterson, *Louisiana Tech*
 Jane Roads, *Moberly Area Community College*
 Melanie Smith, *Bishop State Community College*
 Erik Stubsten, *Chattanooga State Technical Community College*
 Tong Wagner, *Greenville Technical College*
 Rick Woodmansee, *Sacramento City College*
 Sessia Wyche, *University of Texas at Brownsville*

Over the years, we have come to rely on an extensive team of experienced professionals. Our sincere thanks go to these dedicated individuals at Pearson who worked long and hard to make this revision a success.

We would like to thank Michael Hirsch, Matthew Summers, Karen Montgomery, Alicia Frankel, Lauren Morse, Vicki Dreyfus, Stacey Miller, Noelle Saligumba, Eric Gregg, and all of the Pearson math team for helping with the revision of the text.

We are especially pleased to welcome Callie Daniels, who has taught from our texts for many years, to our team. Her assistance has been invaluable. She thoroughly reviewed all chapters and helped extensively with manuscript preparation.

We are grateful to Carol Merrigan for her excellent production work. We appreciate her positive attitude, responsiveness, and expert skills. We would also like to thank Pearson CSC for their production work; Emily Keaton for her detailed help in updating real data applications; Connie Day for supplying her copyediting expertise; Pearson CSC for their photo research; and Lucie Haskins

for producing another accurate, useful index. Paul Lorczak and Hal Whipple did a thorough, timely job accuracy-checking the page proofs and answers, and Sarah Sponholz checked the index.

We particularly thank the many students and instructors who have used this text over the years. You are the reason we do what we do. It is our hope that we have positively impacted your mathematics journey. We would welcome any comments or suggestions you might have via email to math@pearson.com.

John Hornsby
Terry McGinnis

DEDICATION

To my friend, Carol Merrigan

Johnny

To Andrew and Tyler

Mom

Resources for Success


Get the Most Out of MyLab Math for *Intermediate Algebra*, Thirteenth Edition by Lial, Hornsby, McGinnis

The Lial team has helped thousands of students learn algebra with an approachable, teacherly writing style and balance of skill and concept development. With this revision, the series retains the hallmarks that have helped students succeed in math, and includes new and updated digital tools in the MyLab Math course.

Take advantage of the following resources to get the most out of your MyLab Math course.

Get Students Prepared with Integrated Review

Every student enters class with different levels of preparedness and prerequisite knowledge. To ensure students are caught up on prior skills, every Lial MyLab course now includes Integrated Review.




Integrated Review provides embedded and personalized review of prerequisite topics within relevant chapters. Students can check their prerequisite skills, and receive personalized practice on the topics they need to focus on, with study aids like worksheets and videos also available to help.

Integrated Review assignments are pre-made and available to assign in the Assignment Manager.

Personalize Learning

New! Skill Builder exercises offer just-in-time additional adaptive practice. The adaptive engine tracks student performance and delivers questions to each individual that adapt to his or her level of understanding. This new feature allows instructors to assign fewer questions for homework, allowing students to complete as many or as few questions as they need.




Resources for Success

Support Students Whenever, Wherever

Updated! The complete video program for the Lial series includes:

- Full Section Lecture Videos
- Solution clips for select exercises
- Chapter Test Prep videos
- Short Quick Review videos that recap each section

Full Section Lecture Videos are also available as shorter, objective-level videos. No matter your students' needs—if they missed class, need help solving a problem, or want a short summary of a section's concepts—they can get support whenever they need it, wherever they need it. Much of the video series has been updated in a modern presentation format.




Foster a Growth Mindset

New! A Mindset module is available in the course, with mindset-focused videos and exercises that encourage students to maintain a positive attitude about learning, value their own ability to grow, and view mistakes as a learning opportunity.

Get Students Engaged

New! Learning Catalytics Learning Catalytics is an interactive student response tool that uses students' smartphones, tablets, or laptops to engage them in more sophisticated tasks and thinking.

In addition to a library of developmental math questions, Learning Catalytics questions created specifically for this text are pre-built to make it easy for instructors to begin using this tool! These questions, which cover prerequisite skills before each section, are noted in the margin of the Annotated Instructor's Edition, and can be found in Learning Catalytics by searching for "LialIntermediate#", where # is the chapter number.



Resources for Success

Instructor Resources

Annotated Instructor's Edition

Contains all the content found in the student edition, plus answers to even and odd exercises on the same text page, and Teaching Tips and Classroom Examples throughout the text placed at key points.

The resources below are available through Pearson's Instructor Resource Center, or from MyLab Math.

Instructor's Resource Manual with Tests

Includes mini-lectures for each text section, several forms of tests per chapter—two diagnostic pretests, four free-response and two multiple-choice test forms per chapter, and two final exams.

Instructor's Solutions Manual

Contains detailed, worked-out solutions to all exercises in the text.

TestGen®

Enables instructors to build, edit, print, and administer tests using a computerized bank of questions developed to cover all the objectives of the text. TestGen is algorithmically based, allowing instructors to create multiple but equivalent versions of the same question or test with the click of a button. Instructors can also modify test bank questions or add new questions.

PowerPoint Lecture Slides

Available for download only, these slides present key concepts and definitions from the text. Accessible versions of the PowerPoint slides are also available for students who are vision-impaired.

Student Resources

Guided Notebook

This Guided Notebook helps students keep their work organized as they work through their course. The notebook includes:

- Guided Examples that are worked out for students, plus corresponding Now Try This exercises for each text objective.
- Extra practice exercises for every section of the text, with ample space for students to show their work.
- Learning objectives and key vocabulary terms for every text section, along with vocabulary practice problems.

Student Solutions Manual

Provides completely worked-out solutions to the odd-numbered section exercises and to all exercises in the Now Trys, Relating Concepts, Chapter Reviews, Mixed Reviews, Chapter Tests, and Cumulative Reviews.

This page intentionally left blank

Using Your Math Text

Your text is a *valuable resource*. You will learn more if you make full use of the features it offers.

Now TRY THIS

General Features of This Text

Locate each feature, and complete any blanks.


- **Table of Contents** This is located at the front of the text.
Find it and mark the chapters and sections you will cover, as noted on your course syllabus.
- **Answer Section** This is located at the back of the text.
Tab this section so you can easily refer to it when doing homework or reviewing for tests.
- **List of Formulas** This helpful list of geometric formulas, along with review information on triangles and angles, is found at the back of the text.

The formula for the volume of a cube is _____.

Specific Features of This Text

Look through Chapter 1 and give the number of a page that includes an example of each of the following specific features.

- **Objectives** The objectives are listed at the beginning of each section and again within the section as the corresponding material is presented. Once you finish a section, ask yourself if you have accomplished them.
See page _____.
- **Vocabulary List** Important vocabulary is listed at the beginning of each section. You should be able to define these terms when you finish a section. *See page _____.*
- **Now Try Exercises** These margin exercises allow you to immediately practice the material covered in the examples and prepare you for the exercises. Check your results using the answers at the bottom of the page. *See page _____.*
- **Pointers** These small, shaded balloons provide on-the-spot warnings and reminders, point out key steps, and give other helpful tips. *See page _____.*
- **Cautions** These provide warnings about common errors that students often make or trouble spots to avoid. *See page _____.*
- **Notes** These provide additional explanations or emphasize other important ideas.
See page _____.
- **Problem-Solving Hints** These boxes give helpful tips or strategies to use when you work applications. *See page _____.*



STUDY SKILL 2



Reading Your Math Text

Take time to read each section and its examples before doing your homework. You will learn more and be better prepared to work the exercises your instructor assigns.

Approaches to Reading Your Math Text

Student A learns best by listening to her teacher explain things. She “gets it” when she sees the instructor work problems. She previews the section before the lecture, so she knows generally what to expect. **Student A carefully reads the section in her text AFTER she hears the classroom lecture on the topic.**

Student B learns best by reading on his own. He reads the section and works through the examples before coming to class. That way, he knows what the teacher is going to talk about and what questions he wants to ask. **Student B carefully reads the section in his text BEFORE he hears the classroom lecture on the topic.**

Which of these reading approaches works best for you—that of Student A or Student B?

Tips for Reading Your Math Text

- **Turn off your cell phone and the TV.** You will be able to concentrate more fully on what you are reading.
- **Survey the material.** Glance over the assigned material to get an idea of the “big picture.” Look at the list of objectives to see what you will be learning.
- **Read slowly.** Read only one section—or even part of a section—at a sitting, with paper and pencil in hand.
- **Pay special attention to important information given in colored boxes or set in bold-face type.** Highlight any additional information you find helpful.
- **Study the examples carefully.** Pay particular attention to the blue side comments and any pointer balloons.
- **Do the Now Try exercises in the margin on separate paper as you go.** These problems mirror the examples and prepare you for the exercise set. Check your answers with those given at the bottom of the page.
- **Make study cards as you read.** Make cards for new vocabulary, rules, procedures, formulas, and sample problems.
- **Mark anything you don’t understand. ASK QUESTIONS** in class—everyone will benefit. Follow up with your instructor, as needed.

Now TRY THIS

Think through and answer each question.

1. Which two or three reading tips given above will you try this week?
2. Did the tips you selected improve your ability to read and understand the material? Explain.

Taking Lecture Notes

Come to class prepared.

- Bring paper, pencils, notebook, text, completed homework, and any other materials you need.
- Arrive 10–15 minutes early if possible. Use the time before class to review your notes or study cards from the last class period.
- Select a seat carefully so that you can hear and see what is going on.

Study the set of sample math notes given at the right.

- Use a new page for each day's lecture.
- Include the date and title of the day's lecture topic.
- Skip lines and write neatly to make reading easier.
- Include cautions and warnings to emphasize common errors to avoid.
- Mark important concepts with stars, underlining, etc.
- Use two columns, which allows an example and its explanation to be close together.
- Use brackets and arrows to clearly show steps, related material, etc.
- Highlight any material and/or information that your instructor emphasizes. Instructors often give "clues" about material that will definitely be on an exam.

Sept. 1

Translating Words to Expressions and Equations

Problem solving: key words or phrases translate to algebraic expressions.

Caution *Subtraction is not commutative; the order does matter.*

Examples:	Correct	Wrong
10 less than a number	$x - 10$	$10 - x$
a number subtracted from 10	$10 - x$	$x - 10$
10 minus a number	$10 - x$	$x - 10$

A phrase (part of a sentence)

↓

algebraic expression

A sentence

↓

equation with = sign

Note difference *No equality symbol in an expression. $3x + 2$*
Equation has an equality symbol. $3x + 2 = 14$

★ *Pay close attention to exact wording of the sentence; watch for commas.*

The quotient of (a number) and (the number plus 4) is 28.

$$\frac{x}{x + 4} = 28$$

The quotient of (a number) and (4), plus the number, is 28.

$$\frac{x}{4} + x = 28$$

Commas separate this from division part

Consider using a three-ring binder to organize your notes, class handouts, and completed homework.

Now TRY THIS

With a study partner or in a small group, compare lecture notes. Then answer each question.

1. What are you doing to show main points in your notes (such as boxing, using stars, etc.)?
2. In what ways do you set off explanations from worked problems and subpoints (such as indenting, using arrows, circling, etc.)?
3. What new ideas did you learn by examining your classmates' notes?
4. What new techniques will you try when taking notes in future lectures?

STUDY SKILL 4



Completing Your Homework

You are ready to do your homework **AFTER** you have read the corresponding text section and worked through the examples and Now Try exercises.

Homework Tips

- **Keep distractions and potential interruptions to a minimum.** Turn off your cell phone and the TV. Find a quiet, comfortable place to work, away from a lot of other people, so you can concentrate on what you are doing.
- **Review your class notes.** Pay particular attention to anything your instructor emphasized during the lecture on this material.
- **Survey the exercise set.** Glance over the problems that your instructor has assigned to get a general idea of the types of exercises you will be working. Skim directions, and note any references to section examples.
- **Work problems neatly.** NEVER do your math homework in pen. Use pencil and write legibly, so others can read your work. Skip lines between steps. Clearly separate problems from each other.
- **Show all your work.** It is tempting to take shortcuts. Include ALL steps.
- **Check your work frequently to make sure you are on the right track.** It is hard to unlearn a mistake. For all odd-numbered problems, answers are given in the back of the text.
- **If you have trouble with a problem, refer to the corresponding worked example in the section.** The exercise directions will often reference specific examples to review. Pay attention to every line of the worked example to see how to get from step to step.
- **If you have trouble with an even-numbered problem, work the corresponding odd-numbered problem.** Check your answer in the back of the text, and apply the same steps to work the even-numbered problem.
- **If you have genuinely tried to work a problem but have not been able to complete it in a reasonable amount of time, it's ok to STOP.** Mark these problems. Ask for help at your school's tutor center or from fellow classmates, study partners, or your instructor.
- **Do some homework problems every day.** This is a good habit, even if your math class does not meet each day.

Now TRY THIS

Think through and answer each question.

1. What is your instructor's policy regarding homework?
2. Think about your current approach to doing homework. Be honest in your assessment.
 - (a) What are you doing that is working well?
 - (b) What improvements could you make?
3. Which one or two homework tips will you try this week?
4. In the event that you need help with homework, what resources are available? When does your instructor hold office hours?

Using Study Cards

You may have used “flash cards” in other classes. In math, “study cards” can help you remember terms and definitions, procedures, and concepts. Use study cards to

- Help you understand and learn the material;
- Quickly review when you have a few minutes;
- Review before a quiz or test.

One of the advantages of study cards is that you learn the material while you are making them.

Vocabulary Cards

Put the word and a page reference on the front of the card. On the back, write the definition, an example, any related words, and a sample problem (if appropriate).

Procedure (“Steps”) Cards

Write the name of the procedure on the front of the card. Then write each step in words. On the back of the card, put an example showing each step.


Practice Problem Cards


Write a problem with direction words (like *solve*, *simplify*) on the front of the card, and work the problem on the back. Make one for each type of problem you learn.

Now TRY THIS

Make a vocabulary card, a procedure card, and a practice problem card for material that you are learning or reviewing.

<i>Interval notation</i>	<i>p. 104</i>	Front of Card
<p><i>Definition:</i> Using symbols to describe an interval on a number line.</p> <p><i>symbols:</i> ∞ $-\infty$ () [] (] [)</p> <p>Use interval notation to tell what numbers are in the solution set for an inequality.</p> <p><i>Examples:</i> $(-5, \infty)$ All numbers greater than -5, <u>not including</u> -5</p> <p>$[-5, 5)$ All numbers between -5 and 5, <u>including</u> -5, <u>excluding</u> 5</p>		Back of Card

<i>Solving a Linear Inequality</i>	<i>p. 108</i>	Front of Card
<p>1. Simplify each side separately. (Clear parentheses, fractions, and decimals and combine like terms.)</p> <p>2. Isolate variable terms on one side. (Add or subtract the same number from each side.)</p> <p>3. Isolate the variable. (Divide each side by the same number; if dividing by a <u>negative</u> number, <u>reverse direction</u> of inequality.)</p>		Back of Card
<p><i>Solve</i> $-3(x+4) + 2 \geq 7 - x$ and graph the solution set.</p> <p>$-3(x+4) + 2 \geq 7 - x$ Clear parentheses.</p> <p>$-3x - 12 + 2 \geq 7 - x$ Combine like terms.</p> <p>$-3x - 10 \geq 7 - x$ Both sides are simplified.</p> <p>$-3x - 10 + x \geq 7 - x + x$ Add x to each side.</p> <p>$-2x - 10 \geq 7$ Variable term still not isolated.</p> <p>$-2x - 10 + 10 \geq 7 + 10$ Add 10 to each side.</p> <p>$\frac{-2x}{-2} \geq \frac{17}{-2}$ Divide each side by -2; dividing by negative, <u>reverse</u> direction of inequality symbol.</p> <p>$x \leq -\frac{17}{2}$</p> <p></p>		Back of Card

<i>Solve this inequality. Give the solution set in both interval and graph forms.</i>		Front of Card
$-5x - 4 \geq 11$		
<p>$-5x - 4 \geq 11$</p> <p>$-5x - 4 + 4 \geq 11 + 4$</p> <p>$-5x \geq 15$</p> <p>$\frac{-5x}{-5} \geq \frac{15}{-5}$</p> <p>$x \leq -3$</p> <p>$(-\infty, -3]$</p> <p></p>		Back of Card
<p>Neither side can be simplified.</p> <p>Add 4 to each side.</p> <p>Divide each side by -5.</p> <p>Reverse direction of inequality symbol because dividing by <u>negative</u> number.</p> <p>Solution set in interval form; all numbers less than or equal to -3, <u>including</u> -3.</p> <p>Graph of solution set.</p>		

STUDY SKILL 6

Managing Your Time

Many college students juggle a busy schedule and multiple responsibilities, including school, work, and family demands.



Time Management Tips

- **Read the syllabus for each class.** Understand class policies, such as attendance, late homework, and make-up tests. Find out how you are graded.
 - **Make a semester or quarter calendar.** Put test dates and major due dates for *all* your classes on the *same* calendar. Try using a different color for each class.
 - **Make a weekly schedule.** After you fill in your classes and other regular responsibilities, block off some study periods. Aim for 2 hours of study for each 1 hour in class.
 - **Choose a regular study time and place** (such as the campus library). Routine helps.
 - **Keep distractions to a minimum.** Get the most out of the time you have set aside for studying by limiting interruptions. Turn off your cell phone. Take a break from social media. Avoid studying in front of the TV.
 - **Make “to-do” lists.** Number tasks in order of importance. To see your progress, cross off tasks as you complete them.
- **Break big assignments into smaller chunks.** Don't wait until the last minute to begin big assignments or to study for tests. Make deadlines for each smaller chunk so that you stay on schedule.
 - **Take breaks when studying.** Do not try to study for hours at a time. Take a 10-minute break each hour or so.
 - **Ask for help when you need it.** Talk with your instructor during office hours. Make use of the learning/tutoring center, counseling office, or other resources available at your school.

Now TRY THIS

Work through the following, answering any questions.

1. Evaluate when and where you are currently studying. Are these places quiet and comfortable? Are you studying when you are most alert?
2. Which of the above tips will you try this week to improve your time management?
3. Create a weekly calendar that includes your class times, study times, and other family and/or work obligations.
4. Once the week is over, evaluate how these tips worked. Did you use your calendar and stick to it? What will you do differently next week?
5. Ask classmates, friends, and/or family members for tips on how they manage their time. Try any that you think might work for you.

Reviewing a Chapter

Your text provides extensive material to help you prepare for quizzes or tests in this course. Refer to the **Chapter 1 Summary** as you read through the following techniques.

Techniques for Reviewing a Chapter

- **Review the Key Terms and any New Symbols.** Make a study card for each. Include a definition, an example, a sketch (if appropriate), and a section or page reference.
- **Take the Test Your Word Power quiz** to check your understanding of new vocabulary. The answers immediately follow.
- **Read the Quick Review.** Pay special attention to the headings. Study the explanations and examples given for each concept. Try to think about the whole chapter.
- **Reread your lecture notes.** Focus on what your instructor has emphasized in class, and review that material in your text.
- **Look over your homework.** Pay special attention to any trouble spots.
- **Work the Review Exercises.** They are grouped by section. Answers are included at the back of the text.
 - ▶ Pay attention to direction words, such as *simplify*, *solve*, and *evaluate*.
 - ▶ Are your answers exact and complete? Did you include the correct labels, such as \$, cm², ft, etc.?
 - ▶ Make study cards for difficult problems.
- **Work the Mixed Review Exercises.** They are in random order. Check your answers in the answer section at the back of the text.
- **Take the Chapter Test under test conditions.**
 - ▶ Time yourself.
 - ▶ Use a calculator or notes only if your instructor permits them on tests.
 - ▶ Take the test in one sitting.
 - ▶ Show all your work.
 - ▶ Check your answers in the answer section. Section references are provided.

Reviewing a chapter takes time. Avoid rushing through your review in one night. Use the suggestions over a few days or evenings to better understand and remember the material.

Now TRY THIS

Follow these reviewing techniques to prepare for your next test. Then answer each question.

1. How much time did you spend reviewing for your test? Was it enough?
2. Which reviewing techniques worked best for you?
3. Are you investing enough time and effort to really *know* the material and set yourself up for success? Explain.
4. What will you do differently when reviewing for your next test?

CHAPTER 1 Summary 137

Chapter 1 Summary

STUDY SKILLS REMINDER
How do you best prepare for a test? Review Study Skill 7, Reviewing a Chapter.

Key Terms

<p>1.1 linear (first-degree) equation in one variable solution set equivalent equations conditional equation identity contradiction</p>	<p>1.2 mathematical model formula percent percent increase percent decrease</p>	<p>1.5 inequality interval linear inequality in one variable equivalent inequalities three-part inequality</p>	<p>1.6 intersection compound inequality union</p> <p>1.7 absolute value equation absolute value inequality</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------

1.4 consecutive integers
consecutive even integers
consecutive odd integers

New Symbols

% percent	∞ infinity	(a, b) interval notation for a < x < b	∩ set intersection
1° one degree	-∞ negative infinity	[a, b] interval notation for a ≤ x ≤ b	∪ set union
	(-∞, ∞) the set of real numbers		

Test Your Word Power
See how well you have learned the vocabulary in this chapter.

1. An equation is
 - A. an algebraic expression
 - B. an expression that contains fractions
 - C. an expression that uses any of the four basic operations or the operation of raising to powers or taking roots on any collection of variables and numbers formed according to the rules of algebra
 - D. a statement that two algebraic expressions are equal.
2. A linear equation that is a conditional equation has
 - A. no solution
 - B. one solution
 - C. two solutions
 - D. infinitely many solutions.
3. An inequality is
 - A. a statement that two algebraic expressions are equal
 - B. a point on a number line
 - C. an equation with no solutions
 - D. a statement consisting of algebraic expressions related by <, ≤, >, or ≥.
4. Interval notation is
 - A. a point on a number line
 - B. a special notation for describing a point on a number line
 - C. a way to use symbols to describe an interval on a number line
 - D. a notation to describe unequal quantities.
5. The intersection of two sets A and B is the set of elements that belong
 - A. to both A and B
 - B. to either A or B, or both
 - C. to either A or B, but not both
 - D. to just A.
6. The union of two sets A and B is the set of elements that belong
 - A. to both A and B
 - B. to either A or B, or both
 - C. to either A or B, but not both
 - D. to just B.

ANSWERS

1. D. Examples: $2x + 3 = 7$, $5y = -8$, $x^2 = 4$ 2. B. Example: $x - 4 = 6$ is a conditional equation with one solution, 10.
3. D. Examples: $x < 5$, $7 + 2x \geq 11$, $-5 < 2x - 1 \leq 3$ 4. C. Examples: $(-\infty, 5)$, $(1, \infty)$, $[-3, 3)$
5. A. Example: $H = \{2, 4, 6, 8\}$ and $B = \{1, 2, 3\}$, then $A \cap B = \{2\}$.
6. B. Example: Using the sets A and B from Answer 5, $A \cup B = \{1, 2, 3, 4, 6, 8\}$.

STUDY SKILL 8

Taking Math Tests

Techniques to Improve Your Test Score	Comments
Come prepared with a pencil, eraser, paper, and calculator, if allowed.	Working in pencil lets you erase, keeping your work neat.
Scan the entire test, note the point values of different problems, and plan your time accordingly.	To do 20 problems in 50 minutes, allow $50 \div 20 = 2.5$ minutes per problem. Spend less time on easier problems.
Do a “knowledge dump” when you get the test. Write important notes, such as formulas, in a corner of the test for reference.	Writing down tips and other special information that you’ve learned at the beginning allows you to relax as you take the test.
Read directions carefully, and circle any significant words. When you finish a problem, reread the directions. Did you do what was asked?	Pay attention to any announcements written on the board or made by your instructor. Ask if you don’t understand something.
Show all your work. Many teachers give partial credit if some steps are correct, even if the final answer is wrong. Write neatly.	If your teacher can’t read your writing, you won’t get credit for it. If you need more space to work, ask to use extra paper.
Write down anything that might help solve a problem: a formula, a diagram, etc. If necessary, circle the problem and come back to it later. Do not erase anything you wrote down.	If you know even a little bit about a problem, write it down. The answer may come to you as you work on it, or you may get partial credit. Don’t spend too long on any one problem.
If you can’t solve a problem, make a guess. Do not change it unless you find an obvious mistake.	Have a good reason for changing an answer. Your first guess is usually your best bet.
Check that the answer to an application problem is reasonable and makes sense. Reread the problem to make sure you’ve answered the question.	Use common sense. Can the father really be seven years old? Would a month’s rent be \$32,140? Remember to label your answer if needed: \$, years, inches, etc.
Check for careless errors. Rework each problem without looking at your previous work. Then compare the two answers.	Reworking a problem from the beginning forces you to rethink it. If possible, use a different method to solve the problem.



Now TRY THIS

Think through and answer each question.

1. What two or three tips will you try when you take your next math test?
2. How did the tips you selected work for you when you took your math test?
3. What will you do differently when taking your next math test?
4. Ask several classmates how they prepare for math tests. Did you learn any new preparation ideas?

Analyzing Your Test Results

An exam is a learning opportunity—learn from your mistakes. After a test is returned, do the following:

- **Note what you got wrong and why you had points deducted.**
- **Figure out how to solve the problems you missed.** Check your text or notes, or ask your instructor. Rework the problems correctly.
- **Keep all quizzes and tests that are returned to you.** Use them to study for future tests and the final exam.

Typical Reasons for Errors on Math Tests

1. You read the directions wrong.
2. You read the question wrong or skipped over something.
3. You made a computation error.
4. You made a careless error. (For example, you incorrectly copied a correct answer onto a separate answer sheet.)
5. Your answer was not complete.
6. You labeled your answer wrong. (For example, you labeled an answer “ft” instead of “ft².”)
7. You didn’t show your work.
8. You didn’t understand a concept.
9. You were unable to set up the problem (in an application).
10. You were unable to apply a procedure.

These are test-taking errors. They are easy to correct if you read carefully, show all your work, proofread, and double-check units and labels.

These are test preparation errors. Be sure to practice all the kinds of problems that you will see on tests.

Now TRY THIS

Work through the following, answering any questions.

1. Use the sample charts at the right to track your test-taking progress. Refer to the tests you have taken so far in your course. For each test, check the appropriate box in the charts to indicate that you made an error in a particular category.

▼ Test-Taking Errors

Test	Read directions wrong	Read question wrong	Made computation error	Made careless error	Answer not complete	Answer labeled wrong	Didn't show work
1							
2							
3							

2. What test-taking errors did you make? Do you notice any patterns?
3. What test preparation errors did you make? Do you notice any patterns?
4. What will you do to avoid these kinds of errors on your next test?

▼ Test Preparation Errors

Test	Didn't understand concept	Didn't set up problem correctly	Couldn't apply a procedure
1			
2			
3			

STUDY SKILL 10



Preparing for Your Math Final Exam

Your math final exam is likely to be a comprehensive exam, which means it will cover material from the entire term. **One way to prepare for it now is by working a set of Cumulative Review Exercises** each time your class finishes a chapter. This continual review will help you remember concepts and procedures as you progress through the course.

Final Exam Preparation Suggestions

- 1. Figure out the grade you need to earn on the final exam to get the course grade you want.** Check your course syllabus for grading policies, or ask your instructor if you are not sure.
- 2. Create a final exam week plan.** Set priorities that allow you to spend extra time studying. This may mean making adjustments, in advance, in your work schedule or enlisting extra help with family responsibilities.
- 3. Use the following suggestions to guide your studying.**
 - **Begin reviewing several days before the final exam.** DON'T wait until the last minute.
 - **Know exactly which chapters and sections will be covered on the exam.**
 - **Divide up the chapters.** Decide how much you will review each day.
 - **Keep returned quizzes and tests. Use them to review.**
 - **Practice all types of problems. Use the Cumulative Review Exercises** at the end of each chapter in your text beginning in Chapter 1. All answers, with section references, are given in the answer section at the back of the text.
 - **Review or rewrite your notes** to create summaries of important information.
 - **Make study cards for all types of problems.** Carry the cards with you, and review them whenever you have a few minutes.
 - **Take plenty of short breaks as you study to reduce physical and mental stress.** Exercising, listening to music, and enjoying a favorite activity are effective stress busters.

Finally, **DON'T** stay up all night the night before an exam—*get a good night's sleep.*

Now TRY THIS

Think through and answer each question.

1. How many points do you need to earn on your math final exam to get the grade you want in your course?
2. What adjustments to your usual routine or schedule do you need to make for final exam week? List two or three.
3. Which of the suggestions for studying will you use as you prepare for your math final exam? List two or three.
4. Analyze your final exam results. How will you prepare differently next time?

REVIEW OF THE REAL NUMBER SYSTEM

R

- R.1 Fractions, Decimals, and Percents
- R.2 Basic Concepts from Algebra
- R.3 Operations on Real Numbers
- R.4 Exponents, Roots, and Order of Operations
- R.5 Properties of Real Numbers


R.1 Fractions, Decimals, and Percents

OBJECTIVES

- 1 Write fractions in lowest terms.
- 2 Convert between improper fractions and mixed numbers.
- 3 Perform operations with fractions.
- 4 Write decimals as fractions.
- 5 Perform operations with decimals.
- 6 Write fractions as decimals.
- 7 Write percents as decimals and decimals as percents.
- 8 Write percents as fractions and fractions as percents.

Recall that **fractions** are a way to represent parts of a whole. See **FIGURE 1**. In a fraction, the **numerator** gives the number of parts being represented. The **denominator** gives the total number of equal parts in the whole. The fraction bar represents division ($\frac{a}{b} = a \div b$).

Fraction bar $\rightarrow \frac{3}{8}$
3 ← Numerator
8 ← Denominator



The shaded region represents $\frac{3}{8}$ of the circle.

FIGURE 1

A fraction is classified as either a **proper fraction** or an **improper fraction**.

Proper fractions $\frac{1}{5}$, $\frac{2}{7}$, $\frac{9}{10}$, $\frac{23}{25}$ Numerator is less than denominator. Value is less than 1.

Improper fractions $\frac{3}{2}$, $\frac{5}{5}$, $\frac{11}{7}$, $\frac{28}{4}$ Numerator is greater than or equal to denominator. Value is greater than or equal to 1.

OBJECTIVE 1 Write fractions in lowest terms.

A fraction is in **lowest terms** when the numerator and denominator have no factors in common (other than 1).

Writing a Fraction in Lowest Terms

Step 1 Write the numerator and denominator in factored form.

Step 2 Replace each pair of factors common to the numerator and denominator with 1.

Step 3 Multiply the remaining factors in the numerator and in the denominator.

(This procedure is sometimes called “**simplifying the fraction.**”)

VOCABULARY

- fractions
- numerator
- denominator
- proper fraction
- improper fraction
- lowest terms
- mixed number
- reciprocals
- decimal
- terminating decimal
- repeating decimal
- percent


NOW TRY EXERCISE 1

Write each fraction in lowest terms.

(a) $\frac{30}{42}$ (b) $\frac{10}{70}$ (c) $\frac{72}{120}$


NOW TRY EXERCISE 2

Write $\frac{92}{5}$ as a mixed number.


NOW TRY EXERCISE 3

Write $11\frac{2}{3}$ as an improper fraction.

NOW TRY ANSWERS

1. (a) $\frac{5}{7}$ (b) $\frac{1}{7}$ (c) $\frac{3}{5}$
2. $18\frac{2}{5}$
3. $\frac{35}{3}$

EXAMPLE 1 Writing Fractions in Lowest Terms

Write each fraction in lowest terms.

(a) $\frac{10}{15} = \frac{2 \cdot 5}{3 \cdot 5} = \frac{2}{3} \cdot \frac{5}{5} = \frac{2}{3} \cdot 1 = \frac{2}{3}$ 5 is the greatest common factor of 10 and 15.

(b) $\frac{15}{45} = \frac{15}{3 \cdot 15} = \frac{1}{3 \cdot 1} = \frac{1}{3}$ Remember to write 1 in the numerator.

(c) $\frac{150}{200} = \frac{3 \cdot 50}{4 \cdot 50} = \frac{3}{4} \cdot 1 = \frac{3}{4}$ 50 is the greatest common factor of 150 and 200.

Another strategy is to choose *any* common factor and work in stages.

$$\frac{150}{200} = \frac{15 \cdot 10}{20 \cdot 10} = \frac{3 \cdot 5 \cdot 10}{4 \cdot 5 \cdot 10} = \frac{3}{4} \cdot 1 \cdot 1 = \frac{3}{4}$$

The same answer results.

NOW TRY 

OBJECTIVE 2 Convert between improper fractions and mixed numbers.

A **mixed number** is a single number that represents the sum of a natural (counting) number and a proper fraction.

$$\text{Mixed number} \rightarrow 2\frac{3}{4} = 2 + \frac{3}{4}$$

EXAMPLE 2 Converting an Improper Fraction to a Mixed Number

Write $\frac{59}{8}$ as a mixed number.

Because the fraction bar represents division ($\frac{a}{b} = a \div b$, or $b\overline{)a}$), divide the numerator of the improper fraction by the denominator.

$$\begin{array}{r} 7 \leftarrow \text{Quotient} \\ \text{Denominator of fraction} \rightarrow 8 \overline{)59} \leftarrow \text{Numerator of fraction} \\ \underline{56} \\ 3 \leftarrow \text{Remainder} \end{array} \quad \frac{59}{8} = 7\frac{3}{8}$$

NOW TRY 

EXAMPLE 3 Converting a Mixed Number to an Improper Fraction

Write $6\frac{4}{7}$ as an improper fraction.

Multiply the denominator of the fraction by the natural number, and then add the numerator to obtain the numerator of the improper fraction.

$$7 \cdot 6 = 42 \quad \text{and} \quad 42 + 4 = 46$$

The denominator of the improper fraction is the same as the denominator in the mixed number, which is 7 here.

$$6\frac{4}{7} = \frac{7 \cdot 6 + 4}{7} = \frac{46}{7}$$

NOW TRY 




FIGURE 2

NOW TRY EXERCISE 4

Multiply. Write the answer in lowest terms.

$$\frac{4}{7} \cdot \frac{5}{8}$$

OBJECTIVE 3 Perform operations with fractions.

FIGURE 2 illustrates multiplying fractions.

Multiplying Fractions

$$\text{If } \frac{a}{b} \text{ and } \frac{c}{d} \text{ are fractions, then } \frac{a}{b} \cdot \frac{c}{d} = \frac{a \cdot c}{b \cdot d}.$$

That is, to multiply two fractions, multiply their numerators and then multiply their denominators.

EXAMPLE 4 Multiplying Fractions

Multiply. Write the answer in lowest terms.

$$\frac{3}{8} \cdot \frac{4}{9}$$

$$= \frac{3 \cdot 4}{8 \cdot 9} \quad \begin{array}{l} \text{Multiply numerators.} \\ \text{Multiply denominators.} \end{array}$$

$$= \frac{12}{72} \quad \text{Multiply.}$$

$$= \frac{1 \cdot 12}{6 \cdot 12} \quad \begin{array}{l} \text{The greatest common factor} \\ \text{of 12 and 72 is 12.} \end{array}$$

Make sure the product is in lowest terms.

$$= \frac{1}{6}$$

$$\frac{1 \cdot 12}{6 \cdot 12} = \frac{1}{6} \cdot 1 = \frac{1}{6}$$

NOW TRY

Two numbers are **reciprocals** of each other if their product is 1. For example, $\frac{3}{4} \cdot \frac{4}{3} = \frac{12}{12}$, or 1. Division is the inverse or opposite of multiplication, and as a result, we use reciprocals to divide fractions. FIGURE 3 illustrates dividing fractions.




FIGURE 3

Dividing Fractions

$$\text{If } \frac{a}{b} \text{ and } \frac{c}{d} \text{ are fractions, then } \frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c}.$$

That is, to divide by a fraction, multiply by its reciprocal.

EXAMPLE 5 Dividing Fractions

Divide. Write answers in lowest terms as needed.

(a) $\frac{3}{4} \div \frac{8}{5}$

$$= \frac{3}{4} \cdot \frac{5}{8} \quad \begin{array}{l} \text{Multiply by } \frac{5}{8}, \text{ the} \\ \text{reciprocal of } \frac{8}{5}. \end{array}$$

$$= \frac{3 \cdot 5}{4 \cdot 8} \quad \begin{array}{l} \text{Multiply numerators.} \\ \text{Multiply denominators.} \end{array}$$

$$= \frac{15}{32} \quad \text{Make sure the answer is in lowest terms.}$$

(b) $\frac{5}{8} \div 10$ Think of 10 as $\frac{10}{1}$ here.

$$= \frac{5}{8} \cdot \frac{1}{10} \quad \begin{array}{l} \text{Multiply by } \frac{1}{10}, \text{ the} \\ \text{reciprocal of 10.} \end{array}$$

$$= \frac{5 \cdot 1}{8 \cdot 2 \cdot 5} \quad \text{Multiply and factor.}$$

$$= \frac{1}{16} \quad \text{Remember to write 1 in the numerator.}$$

NOW TRY ANSWER

4. $\frac{5}{14}$

NOW TRY
EXERCISE 5

Divide. Write answers in lowest terms as needed.

(a) $\frac{2}{7} \div \frac{8}{9}$ (b) $3\frac{3}{4} \div 4\frac{2}{7}$

$$\begin{aligned} \text{(c)} \quad & 1\frac{2}{3} \div 4\frac{1}{2} \\ &= \frac{5}{3} \div \frac{9}{2} && \text{Write each mixed number as an improper fraction.} \\ &= \frac{5}{3} \cdot \frac{2}{9} && \text{Multiply by } \frac{2}{9}, \text{ the reciprocal of } \frac{9}{2}. \\ &= \frac{10}{27} && \text{Multiply. The quotient is in lowest terms.} \end{aligned}$$

NOW TRY 

FIGURES 4 and 5 illustrate adding and subtracting fractions.




FIGURE 4




FIGURE 5

Adding and Subtracting Fractions

If $\frac{a}{b}$ and $\frac{c}{b}$ are fractions (where $b \neq 0$), then add or subtract as follows.

$$\frac{a}{b} + \frac{c}{b} = \frac{a + c}{b}$$

$$\frac{a}{b} - \frac{c}{b} = \frac{a - c}{b}$$

That is, to add or subtract two fractions having the same denominator, add or subtract the numerators and keep the same denominator.

If the denominators are different, first find the least common denominator (LCD). Write each fraction as an equivalent fraction with this denominator. Then add or subtract as above.

EXAMPLE 6 Adding and Subtracting Fractions

Add or subtract as indicated. Write answers in lowest terms as needed.

(a) $\frac{2}{10} + \frac{3}{10}$

$$= \frac{2 + 3}{10} \quad \begin{array}{l} \text{Add numerators.} \\ \text{Keep the same} \\ \text{denominator.} \end{array}$$

$$= \frac{5}{10}$$

$$= \frac{1}{2} \quad \begin{array}{l} \text{Write in lowest} \\ \text{terms.} \end{array}$$

(b) $\frac{4}{15} + \frac{5}{9}$

$$= \frac{4}{15} \cdot \frac{3}{3} + \frac{5}{9} \cdot \frac{5}{5}$$

$$= \frac{12}{45} + \frac{25}{45}$$

$$= \frac{37}{45}$$

$15 = 3 \cdot 5$ and
 $45 = 3 \cdot 3 \cdot 5$, so the
LCD is $3 \cdot 3 \cdot 5 = 45$.

Write equivalent
fractions with the
common denominator.

Add numerators.
Keep the same
denominator.

NOW TRY ANSWERS

5. (a) $\frac{9}{28}$ (b) $\frac{7}{8}$


NOW TRY EXERCISE 6

Add or subtract as indicated.
Write answers in lowest terms as needed.

(a) $\frac{1}{8} + \frac{3}{8}$ (b) $\frac{5}{12} + \frac{3}{8}$

(c) $\frac{5}{11} - \frac{2}{9}$ (d) $4\frac{1}{3} - 2\frac{5}{6}$

$$\begin{aligned} \text{(c)} \quad & \frac{15}{16} - \frac{4}{9} \\ & = \frac{15}{16} \cdot \frac{9}{9} - \frac{4}{9} \cdot \frac{16}{16} \\ & = \frac{135}{144} - \frac{64}{144} \\ & = \frac{71}{144} \end{aligned}$$

Because 16 and 9 have no common factors except 1, the LCD is $16 \cdot 9 = 144$.

Write equivalent fractions with the common denominator.

Subtract numerators.

Keep the common denominator.

(d) $4\frac{1}{2} - 1\frac{3}{4}$

$$\begin{aligned} \text{Method 1} \quad & 4\frac{1}{2} - 1\frac{3}{4} \\ & = \frac{9}{2} - \frac{7}{4} \\ & = \frac{18}{4} - \frac{7}{4} \\ & = \frac{11}{4}, \text{ or } 2\frac{3}{4} \end{aligned}$$

Write each mixed number as an improper fraction.

Find a common denominator. The LCD is 4.

Subtract. Write as a mixed number.

Think: $\frac{9}{2} \cdot \frac{2}{2} = \frac{18}{4}$


$$\begin{aligned} \text{Method 2} \quad & 4\frac{1}{2} = 4\frac{2}{4} = 3\frac{6}{4} \\ & -1\frac{3}{4} = 1\frac{3}{4} = 1\frac{3}{4} \\ & \hline & 2\frac{3}{4}, \text{ or } \frac{11}{4} \end{aligned}$$

The LCD is 4.

$$4\frac{2}{4} = 3 + 1 + \frac{2}{4} = 3 + \frac{4}{4} + \frac{2}{4} = 3\frac{6}{4}$$

The same answer results.

NOW TRY 




3 parts of the whole 10 are shaded.
As a fraction, $\frac{3}{10}$ of the figure is shaded. As a decimal, 0.3 is shaded.
Both of these numbers are read “three-tenths.”

FIGURE 6

Fractions are one way to represent parts of a whole. Another way is with a decimal fraction or **decimal**, a number written with a decimal point.

9.25, 14.001, 0.3 **Decimal numbers**

See **FIGURE 6**. Each digit in a decimal number has a place value, as shown below.



NOW TRY ANSWERS

6. (a) $\frac{1}{2}$ (b) $\frac{19}{24}$ (c) $\frac{23}{99}$
(d) $\frac{3}{2}$, or $1\frac{1}{2}$

Each successive place value is ten times greater than the place value to its right and one-tenth as great as the place value to its left.

OBJECTIVE 4 Write decimals as fractions.

Place value is used to write a decimal number as a fraction.

Converting a Decimal to a Fraction

Read the decimal using the correct place value. Write it in fractional form just as it is read.

- The numerator will be the digits to the right of the decimal point.
- The denominator will be a power of 10—that is, 10 for tenths, 100 for hundredths, and so on.

NOW TRY EXERCISE 7

Write each decimal as a fraction. (Do not write in lowest terms.)

- (a) 0.8 (b) 0.431 (c) 2.58

EXAMPLE 7 Writing Decimals as Fractions

Write each decimal as a fraction. (Do not write in lowest terms.)

- (a) 0.95 We read 0.95 as “
- ninety-five hundredths.*
- ”

$$0.95 = \frac{95}{100} \leftarrow \text{For hundredths}$$

- (b) 0.056 We read 0.056 as “
- fifty-six thousandths.*
- ”

Do not confuse 0.056 with 0.56, read “fifty-six hundredths,” which is the fraction $\frac{56}{100}$.

$$0.056 = \frac{56}{1000} \leftarrow \text{For thousandths}$$

- (c) 4.2095 We read this decimal number, which is greater than 1, as “
- Four and two thousand ninety-five ten-thousandths.*
- ”

$$4.2095 = 4 \frac{2095}{10,000} \quad \text{Write the decimal number as a mixed number.}$$

Think: $10,000 \cdot 4 + 2095$

$$= \frac{42,095}{10,000}$$

Write the mixed number as an improper fraction.

NOW TRY **OBJECTIVE 5** Perform operations with decimals.**EXAMPLE 8** Adding and Subtracting Decimals

Add or subtract as indicated.

- (a)
- $6.92 + 14.8 + 3.217$

Place the digits of the decimal numbers in columns by place value. Attach zeros as placeholders so that there are the same number of places to the right of each decimal point.

Be sure to line up decimal points.

$$\begin{array}{r} 6.92 \\ 14.8 \\ + 3.217 \\ \hline \end{array}$$

becomes

$$\begin{array}{r} 6.920 \\ 14.800 \\ + 3.217 \\ \hline 24.937 \end{array}$$

Attach 0s.

6.92 is equivalent to 6.920.
14.8 is equivalent to 14.800.

- (b)
- $47.6 - 32.509$

$$\begin{array}{r} 47.6 \\ - 32.509 \\ \hline \end{array}$$

becomes

$$\begin{array}{r} 47.600 \\ - 32.509 \\ \hline 15.091 \end{array}$$

Write the decimal numbers in columns, attaching 0s to 47.6.

NOW TRY EXERCISE 8

Add or subtract as indicated.

- (a)
- $68.9 + 42.72 + 8.973$
-
- (b)
- $351.8 - 2.706$

NOW TRY ANSWERS

7. (a)
- $\frac{8}{10}$
- (b)
- $\frac{431}{1000}$
- (c)
- $\frac{258}{100}$
-
8. (a) 120.593 (b) 349.094

NOW TRY 


NOW TRY EXERCISE 9

Multiply or divide as indicated.

(a) 9.32×1.4

(b) 0.6×0.004

(c) $7.334 \div 3.1$

(Round the answer to two decimal places.)

EXAMPLE 9 Multiplying and Dividing Decimals

Multiply or divide as indicated.

(a) 29.3×4.52

Multiply normally. Place the decimal point in the answer as shown.

$$\begin{array}{r} 29.3 \\ \times 4.52 \\ \hline 586 \\ 1465 \\ 1172 \\ \hline 132.436 \end{array}$$

1 decimal place
2 decimal places
 $1 + 2 = 3$
3 decimal places

(b) 0.05×0.3

Here $5 \times 3 = 15$. Be careful placing the decimal point.

$$\begin{array}{r} 0.05 \\ \times 0.3 \\ \hline \end{array}$$

2 decimal places 1 decimal place

Do not write 0.150. $= 0.015$

2 + 1 = 3 decimal places
Attach 0 as a placeholder in the tenths place.

(c) $8.949 \div 1.25$ (Round the answer to two decimal places.)

$$\begin{array}{r} 7.159 \\ 1.25 \overline{)8.949} \\ \underline{875} \\ 199 \\ \underline{125} \\ 740 \\ \underline{625} \\ 1150 \\ \underline{1125} \\ 25 \end{array}$$

Move each decimal point two places to the right.

Move the decimal point straight up, and divide as with whole numbers. Attach 0s as placeholders.

We carried out the division to three decimal places so that we could round to two decimal places, obtaining the answer 7.16.

NOW TRY 

NOTE To round 7.159 in **Example 9(c)** to two decimal places (that is, to the nearest hundredth), we look at the digit to the *right* of the hundredths place.

- If this digit is 5 or greater, we round up.
- If this digit is less than 5, we drop the digit(s) beyond the desired place.

Hundredths place

$$\begin{array}{r} 7.159 \\ \approx 7.16 \end{array}$$

9, the digit to the right of the hundredths place, is 5 or greater.
Round 5 up to 6. \approx means *is approximately equal to*.

Multiplying and Dividing by Powers of 10 (Shortcuts)

- To **multiply** by a power of 10, **move the decimal point to the right** as many places as the number of zeros.
- To **divide** by a power of 10, **move the decimal point to the left** as many places as the number of zeros.

In both cases, insert 0s as placeholders if necessary.

NOW TRY ANSWERS

9. (a) 13.048 (b) 0.0024
(c) 5.64

NOW TRY EXERCISE 10

Multiply or divide as indicated.

- (a) 294.72×10
 (b) $4.793 \div 100$

EXAMPLE 10 Multiplying and Dividing by Powers of 10

Multiply or divide as indicated.

- (a) 48.731×10
 $= 48.731$ Move the decimal point one place to the right.
 $= 487.31$
- (b) 48.731×1000
 $= 48.731$ Move the decimal point three places to the right.
 $= 48,731$
- (c) $48.731 \div 10$
 $= 48.731$ Move the decimal point one place to the left.
 $= 4.8731$
- (d) $48.731 \div 1000$
 $= 048.731$ Move the decimal point three places to the left.
 $= 0.048731$

NOW TRY

OBJECTIVE 6 Write fractions as decimals.

Converting a Fraction to a Decimal

Because a fraction bar indicates division, write a fraction as a decimal by dividing the numerator by the denominator.

NOW TRY EXERCISE 11

Write each fraction as a decimal. For repeating decimals, write the answer by first using bar notation and then rounding to the nearest thousandth.


- (a) $\frac{17}{20}$ (b) $\frac{2}{9}$

EXAMPLE 11 Writing Fractions as Decimals

Write each fraction as a decimal.

- (a) $\frac{19}{8}$ $\begin{array}{r} 2.375 \\ 8 \overline{)19.000} \\ \underline{16} \\ 30 \\ \underline{24} \\ 60 \\ \underline{56} \\ 40 \\ \underline{40} \\ 0 \end{array}$ Divide 19 by 8. Add a decimal point and as many 0s as necessary to 19.
- (b) $\frac{2}{3}$ $\begin{array}{r} 0.6666 \dots \\ 3 \overline{)2.0000 \dots} \\ \underline{18} \\ 20 \\ \underline{18} \\ 20 \\ \underline{18} \\ 20 \\ \underline{18} \\ 2 \end{array}$
- $\frac{19}{8} = 2.375$ ← Terminating decimal $\frac{2}{3} = 0.6666 \dots$ ← Repeating decimal
 $= 0.\overline{6}$ A bar is written over the repeating digit(s).
 ≈ 0.667 Nearest thousandth

NOW TRY



35 of the 100 squares are shaded. That is, $\frac{35}{100}$, or 35%, of the figure is shaded.

FIGURE 7

OBJECTIVE 7 Write percents as decimals and decimals as percents.

The word **percent** means “per 100.” Percent is written with the symbol **%**. “One percent” means “one per one hundred,” or “one one-hundredth.” See **FIGURE 7**.

Percent, Fraction, and Decimal Equivalents

$$1\% = \frac{1}{100} = 0.01, \quad 10\% = \frac{10}{100} = 0.10, \quad 100\% = \frac{100}{100} = 1$$

NOW TRY ANSWERS

10. (a) 2947.2 (b) 0.04793
 11. (a) 0.85 (b) $0.\overline{2}$; 0.222

For example, 73% means “73 per one hundred.”

$$73\% = \frac{73}{100} = 0.73$$

Essentially, we are dropping the % symbol from 73% and dividing 73 by 100. Doing this moves the decimal point, which is understood to be after the 3, two places to the *left*.


Writing 0.73 as a percent is the opposite process.

$$0.73 = 0.73 \cdot 100\% = 73\% \quad 100\% = 1$$

Moving the decimal point two places to the *right* and attaching a % symbol give the same result.

Converting Percents and Decimals (Shortcuts)

- To convert a percent to a decimal, move the decimal point two places to the *left* and drop the % symbol.
- To convert a decimal to a percent, move the decimal point two places to the *right* and attach a % symbol.



NOW TRY EXERCISE 12

Convert each percent to a decimal and each decimal to a percent.

- (a) 52% (b) 2%
(c) 0.45 (d) 3.5

EXAMPLE 12 Converting Percents and Decimals by Moving the Decimal Point

Convert each percent to a decimal and each decimal to a percent.

- (a) $45\% = 0.45$ (b) $250\% = 2.50$, or 2.5 (c) $9\% = 0.09$
(d) $0.57 = 57\%$ (e) $1.5 = 150\%$ (f) $0.007 = 0.7\%$

NOW TRY

OBJECTIVE 8 Write percents as fractions and fractions as percents.

EXAMPLE 13 Writing Percents as Fractions

Write each percent as a fraction. Give answers in lowest terms.

- (a) 8%

We use the fact that *percent* means “per one hundred,” and convert as follows.

$$8\% = \frac{8}{100} = \frac{2 \cdot 4}{25 \cdot 4} = \frac{2}{25} \quad \text{As with converting percents to decimals, drop the \% symbol and divide by 100.}$$

Write in lowest terms.

(b) $175\% = \frac{175}{100} = \frac{7 \cdot 25}{4 \cdot 25} = \frac{7}{4}$, or $1\frac{3}{4}$ A number greater than 1 is more than 100%.

Write in lowest terms.

NOW TRY ANSWERS

12. (a) 0.52 (b) 0.02
(c) 45% (d) 350%


NOW TRY EXERCISE 13

Write each percent as a fraction. Give answers in lowest terms.

- (a) 20% (b) 160%

(c) 13.5%

$$= \frac{13.5}{100}$$

Drop the % symbol. Divide by 100.

$$= \frac{13.5}{100} \cdot \frac{10}{10}$$

Multiply by 1 in the form $\frac{10}{10}$ to eliminate the decimal in the numerator.

$$= \frac{135}{1000}$$

Multiply.

$$= \frac{27}{200}$$

Write in lowest terms.

$$\frac{135}{1000} = \frac{27 \cdot 5}{200 \cdot 5} = \frac{27}{200}$$

NOW TRY 

We know that 100% of something is the whole thing. One way to convert a fraction to a percent is to multiply by 100%, which is equivalent to 1. This involves the same steps that are used for converting a decimal to a percent—**attach a % symbol and multiply by 100**.


NOW TRY EXERCISE 14

Write each fraction as a percent.

- (a) $\frac{6}{25}$ (b) $\frac{7}{9}$

EXAMPLE 14 Writing Fractions as Percents

Write each fraction as a percent.

(a) $\frac{2}{5}$

$$= \frac{2}{5} \cdot 100\%$$

Multiply by 1 in the form 100%.

$$= \frac{2}{5} \cdot \frac{100}{1} \%$$

$$= \frac{2 \cdot 5 \cdot 20}{5 \cdot 1} \%$$

Multiply and factor.

$$= \frac{2 \cdot 20}{1} \%$$

Divide out the common factor.

$$= 40\%$$

Simplify.

(b) $\frac{1}{6}$

$$= \frac{1}{6} \cdot 100\%$$

$$= \frac{1}{6} \cdot \frac{100}{1} \%$$

$$= \frac{1 \cdot 2 \cdot 50}{2 \cdot 3 \cdot 1} \%$$

$$= \frac{50}{3} \%$$

$$= 16\frac{2}{3} \%, \text{ or } 16.\bar{6}\%$$

NOW TRY 

NOW TRY ANSWERS

13. (a) $\frac{1}{5}$ (b) $\frac{8}{5}$, or $1\frac{3}{5}$


14. (a) 24% (b) $77.\bar{7}\%$

R.1 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Decide whether each statement is true or false. If it is false, explain why.

- In the fraction $\frac{5}{8}$, 5 is the numerator and 8 is the denominator.
- The mixed number equivalent of the improper fraction $\frac{31}{5}$ is $6\frac{1}{5}$.
- The fraction $\frac{7}{7}$ is proper.
- The reciprocal of $\frac{6}{2}$ is $\frac{3}{1}$.

Concept Check Choose the letter of the correct response.

5. Which choice shows the correct way to write $\frac{16}{24}$ in lowest terms?

A. $\frac{16}{24} = \frac{8+8}{8+16} = \frac{8}{16} = \frac{1}{2}$

B. $\frac{16}{24} = \frac{4 \cdot 4}{4 \cdot 6} = \frac{4}{6}$

C. $\frac{16}{24} = \frac{8 \cdot 2}{8 \cdot 3} = \frac{2}{3}$

6. Which fraction is *not* equal to $\frac{5}{9}$?

- A. $\frac{15}{27}$ B. $\frac{30}{54}$ C. $\frac{40}{74}$ D. $\frac{55}{99}$

Write each fraction in lowest terms. See Example 1.

7. $\frac{8}{16}$ 8. $\frac{4}{12}$ 9. $\frac{15}{18}$ 10. $\frac{16}{20}$ 11. $\frac{90}{150}$
 12. $\frac{100}{140}$ 13. $\frac{18}{90}$ 14. $\frac{16}{64}$ 15. $\frac{144}{120}$ 16. $\frac{132}{77}$

Write each improper fraction as a mixed number. See Example 2.

17. $\frac{12}{7}$ 18. $\frac{16}{9}$ 19. $\frac{77}{12}$ 20. $\frac{67}{13}$

Write each mixed number as an improper fraction. See Example 3.

21. $2\frac{3}{5}$ 22. $5\frac{6}{7}$ 23. $12\frac{2}{3}$ 24. $10\frac{1}{5}$

Multiply or divide as indicated. Write answers in lowest terms as needed. See Examples 4 and 5.


25. $\frac{4}{5} \cdot \frac{6}{7}$ 26. $\frac{5}{9} \cdot \frac{2}{7}$ 27. $\frac{2}{15} \cdot \frac{3}{8}$ 28. $\frac{3}{20} \cdot \frac{5}{21}$
 29. $\frac{1}{10} \cdot \frac{12}{5}$ 30. $\frac{1}{8} \cdot \frac{10}{7}$ 31. $\frac{15}{4} \cdot \frac{8}{25}$ 32. $\frac{21}{8} \cdot \frac{4}{7}$
 33. $21 \cdot \frac{3}{7}$ 34. $36 \cdot \frac{4}{9}$ 35. $3\frac{1}{4} \cdot 1\frac{2}{3}$ 36. $2\frac{2}{3} \cdot 1\frac{3}{5}$
 37. $\frac{7}{9} \div \frac{3}{2}$ 38. $\frac{6}{11} \div \frac{5}{4}$ 39. $\frac{5}{4} \div \frac{3}{8}$ 40. $\frac{7}{5} \div \frac{3}{10}$
 41. $\frac{32}{5} \div \frac{8}{15}$ 42. $\frac{24}{7} \div \frac{6}{21}$ 43. $\frac{3}{4} \div 12$ 44. $\frac{2}{5} \div 30$
 45. $6 \div \frac{3}{5}$ 46. $8 \div \frac{4}{9}$ 47. $6\frac{3}{4} \div \frac{3}{8}$ 48. $5\frac{3}{5} \div \frac{7}{10}$
 49. $2\frac{1}{2} \div 1\frac{5}{7}$ 50. $2\frac{2}{9} \div 1\frac{2}{5}$ 51. $2\frac{5}{8} \div 1\frac{15}{32}$ 52. $2\frac{3}{10} \div 1\frac{4}{5}$

Add or subtract as indicated. Write answers in lowest terms as needed. See Example 6.

53. $\frac{7}{15} + \frac{4}{15}$ 54. $\frac{2}{9} + \frac{5}{9}$ 55. $\frac{7}{12} + \frac{1}{12}$ 56. $\frac{3}{16} + \frac{5}{16}$
 57. $\frac{5}{9} + \frac{1}{3}$ 58. $\frac{4}{15} + \frac{1}{5}$ 59. $\frac{3}{8} + \frac{5}{6}$ 60. $\frac{5}{6} + \frac{2}{9}$
 61. $\frac{5}{9} + \frac{3}{16}$ 62. $\frac{3}{4} + \frac{6}{25}$ 63. $3\frac{1}{8} + 2\frac{1}{4}$ 64. $4\frac{2}{3} + 2\frac{1}{6}$
 65. $\frac{7}{9} - \frac{2}{9}$ 66. $\frac{8}{11} - \frac{3}{11}$ 67. $\frac{13}{15} - \frac{3}{15}$ 68. $\frac{11}{12} - \frac{3}{12}$
 69. $\frac{7}{12} - \frac{1}{3}$ 70. $\frac{5}{6} - \frac{1}{2}$ 71. $\frac{7}{12} - \frac{1}{9}$ 72. $\frac{11}{16} - \frac{1}{12}$
 73. $4\frac{3}{4} - 1\frac{2}{5}$ 74. $3\frac{4}{5} - 1\frac{4}{9}$ 75. $8\frac{2}{9} - 4\frac{2}{3}$ 76. $7\frac{5}{12} - 4\frac{5}{6}$

Work each problem involving fractions.

77. For each description, write a fraction in lowest terms that represents the region described.



- (a) The dots in the rectangle as a part of the dots in the entire figure
- (b) The dots in the triangle as a part of the dots in the entire figure
- (c) The dots in the overlapping region of the triangle and the rectangle as a part of the dots in the triangle alone
- (d) The dots in the overlapping region of the triangle and the rectangle as a part of the dots in the rectangle alone

78. At the conclusion of the Pearson softball league season, batting statistics for five players were as shown in the table.

Player	At-Bats	Hits	Home Runs
Maureen	36	12	3
Christine	40	9	2
Chase	11	5	1
Joe	16	8	0
Greg	20	10	2

Use this information to answer each question. Estimate as necessary.

- (a) Which player got a hit in exactly $\frac{1}{3}$ of his or her at-bats?
- (b) Which player got a home run in just less than $\frac{1}{10}$ of his or her at-bats?
- (c) Which player got a hit in just less than $\frac{1}{4}$ of his or her at-bats?
- (d) Which two players got hits in exactly the same fractional part of their at-bats? What was that fractional part, expressed in lowest terms?

Concept Check Provide the correct response.

79. In the decimal number 367.9412, name the digit that has each place value.

- (a) tens (b) tenths (c) thousandths (d) ones or units (e) hundredths

80. Write a decimal number that has 5 in the thousands place, 0 in the tenths place, and 4 in the ten-thousandths place.

81. For the decimal number 46.249, round to the place value indicated.

- (a) hundredths (b) tenths (c) ones or units (d) tens

82. Round each decimal to the nearest thousandth.

- (a) $0.\bar{8}$ (b) $0.\bar{4}$ (c) 0.9762 (d) 0.8645

Write each decimal as a fraction. (Do not write in lowest terms.) See Example 7.

83. 0.4 84. 0.6 85. 0.64 86. 0.82 87. 0.138
 88. 0.104 89. 0.043 90. 0.087 91. 3.805 92. 5.166

Add or subtract as indicated. See Example 8.

93. $25.32 + 109.2 + 8.574$ 94. $90.527 + 32.43 + 589.8$ 95. $28.73 - 3.12$
 96. $46.88 - 13.45$ 97. $43.5 - 28.17$ 98. $345.1 - 56.31$

99. $3.87 + 15 + 2.9$ 100. $8.2 + 1.09 + 12$ 101. $32.56 + 47.356 + 1.8$
 102. $75.2 + 123.96 + 3.897$ 103. $18 - 2.789$ 104. $29 - 8.582$

Multiply or divide as indicated. See Example 9.

105. 12.8×9.1 106. 34.04×0.56 107. 22.41×33 108. 55.76×72
 109. 0.2×0.03 110. 0.07×0.004 111. $78.65 \div 11$ 112. $73.36 \div 14$
 113. $32.48 \div 11.6$ 114. $85.26 \div 17.4$ 115. $19.967 \div 9.74$ 116. $44.4788 \div 5.27$

Multiply or divide as indicated. See Example 10.

117. 123.26×10 118. 785.91×10 119. 57.116×100 120. 82.053×100
 121. 0.094×1000 122. 0.025×1000 123. $1.62 \div 10$ 124. $8.04 \div 10$
 125. $124.03 \div 100$ 126. $490.35 \div 100$ 127. $23.29 \div 1000$ 128. $59.8 \div 1000$

Concept Check Complete the table of fraction, decimal, and percent equivalents.

	Fraction in Lowest Terms (or Whole Number)	Decimal	Percent
129.	$\frac{1}{100}$	0.01	
130.	$\frac{1}{50}$		2%
131.		0.05	5%
132.	$\frac{1}{10}$		
133.	$\frac{1}{8}$	0.125	
134.			20%
135.	$\frac{1}{4}$		
136.	$\frac{1}{3}$		
137.			50%
138.	$\frac{2}{3}$		$66\frac{2}{3}\%$, or $66.\bar{6}\%$
139.		0.75	
140.	1	1.0	

Write each fraction as a decimal. For repeating decimals, write the answer by first using bar notation and then rounding to the nearest thousandth. See Example 11.

141. $\frac{21}{5}$ 142. $\frac{9}{5}$ 143. $\frac{9}{4}$ 144. $\frac{15}{4}$ 145. $\frac{3}{8}$
 146. $\frac{7}{8}$ 147. $\frac{5}{9}$ 148. $\frac{8}{9}$ 149. $\frac{1}{6}$ 150. $\frac{5}{6}$

Write each percent as a decimal. See Examples 12(a)–12(c).

151. 54% 152. 39% 153. 7% 154. 4% 155. 117%
 156. 189% 157. 2.4% 158. 3.1% 159. $6\frac{1}{4}\%$ 160. $5\frac{1}{2}\%$

Write each decimal as a percent. See Examples 12(d)–12(f).

161. 0.79 162. 0.83 163. 0.02 164. 0.08 165. 0.004
 166. 0.005 167. 1.28 168. 2.35 169. 0.4 170. 0.6

Write each percent as a fraction. Give answers in lowest terms. See Example 13.

171. 51% 172. 47% 173. 15% 174. 35% 175. 2%
 176. 8% 177. 140% 178. 180% 179. 7.5% 180. 2.5%

Write each fraction as a percent. See Example 14.

181. $\frac{4}{5}$ 182. $\frac{3}{25}$ 183. $\frac{7}{50}$ 184. $\frac{9}{20}$ 185. $\frac{2}{11}$
 186. $\frac{4}{9}$ 187. $\frac{9}{4}$ 188. $\frac{8}{5}$ 189. $\frac{13}{6}$ 190. $\frac{31}{9}$

R.2 Basic Concepts from Algebra

OBJECTIVES

- 1 Write sets using set notation.
- 2 Use number lines.
- 3 Classify numbers.
- 4 Find additive inverses.
- 5 Use absolute value.
- 6 Use inequality symbols.

VOCABULARY

- set
- elements (members)
- finite set
- natural (counting) numbers
- infinite set
- whole numbers
- empty (null) set
- variable
- number line
- integers
- coordinate
- graph
- rational numbers
- irrational numbers
- real numbers
- additive inverse
(opposite, negative)
- signed numbers
- absolute value
- equation
- inequality

OBJECTIVE 1 Write sets using set notation.

A **set** is a collection of objects called the **elements**, or **members**, of the set. In algebra, the elements of a set are usually numbers. Set braces, $\{ \}$, are used to enclose the elements. For example,

2 is an element of the set $\{1, 2, 3\}$.

Because we can count the number of elements in the set $\{1, 2, 3\}$ and the counting comes to an end, it is a **finite set**.

In algebra, we refer to certain sets of numbers by name. The set

$$N = \{1, 2, 3, 4, 5, 6, \dots\} \quad \text{Natural (counting) numbers}$$

is the **natural numbers**, or the **counting numbers**. The three dots (*ellipsis* points) show that the list continues in the same pattern indefinitely. We cannot list all of the elements of the set of natural numbers, so it is an **infinite set**.

Including 0 with the set of natural numbers gives the set of **whole numbers**.

$$W = \{0, 1, 2, 3, 4, 5, 6, \dots\} \quad \text{Whole numbers}$$

The set containing no elements is the **empty set**, or **null set**, usually written \emptyset . For example, the set of whole numbers less than 0 is \emptyset .

CAUTION Do not write $\{\emptyset\}$ for the empty set. $\{\emptyset\}$ is a set with one element: \emptyset . Use the notation \emptyset for the empty set.

To indicate that 2 is an element of the set $\{1, 2, 3\}$, we use the symbol \in , which is read “*is an element of*.”

$$2 \in \{1, 2, 3\}$$

The number 2 is also an element of the set of natural numbers N .

$$2 \in N$$

To show that 0 is *not* an element of set N , we draw a slash through the symbol \in .

$$0 \notin N$$

Two sets are equal if they contain exactly the same elements. For example,

$$\{1, 2\} = \{2, 1\}. \quad \text{Order does not matter.}$$

To indicate that two sets are not equal, we use the symbol \neq , which is read “is not equal to.” For example,

$$\{1, 2\} \neq \{0, 1, 2\}$$

because one set contains the element 0 and the other does not.


A **variable** is a symbol, usually a letter, used to represent a number or to define a set of numbers. For example,

$$\{x \mid x \text{ is a natural number between 3 and 15}\}$$

(read “the set of all elements x such that x is a natural number between 3 and 15”) defines the following set.

$$\{4, 5, 6, 7, \dots, 14\}$$

The notation $\{x \mid x \text{ is a natural number between 3 and 15}\}$ is an example of **set-builder notation**.



NOW TRY EXERCISE 1

List the elements in the set.

$$\{p \mid p \text{ is a natural number less than 6}\}$$

EXAMPLE 1 Listing the Elements in Sets

List the elements in each set.

(a) $\{x \mid x \text{ is a natural number less than 4}\}$

The natural numbers less than 4 are 1, 2, and 3. This set is $\{1, 2, 3\}$.

(b) $\{x \mid x \text{ is one of the first five even natural numbers}\}$ is $\{2, 4, 6, 8, 10\}$.

(c) $\{x \mid x \text{ is a natural number greater than or equal to 7}\}$

The set of natural numbers greater than or equal to 7 is an infinite set, which is written with ellipsis points as

$$\{7, 8, 9, 10, \dots\}.$$

NOW TRY

NOW TRY EXERCISE 2

Use set-builder notation to describe the set.

$$\{9, 10, 11, 12\}$$

EXAMPLE 2 Using Set-Builder Notation to Describe Sets

Use set-builder notation to describe each set.

(a) $\{1, 3, 5, 7, 9\}$

There are often several ways to describe a set in set-builder notation. One way to describe the given set is

$$\{x \mid x \text{ is one of the first five odd natural numbers}\}.$$

(b) $\{5, 10, 15, \dots\}$

This set can be described as $\{x \mid x \text{ is a positive multiple of 5}\}$.

NOW TRY

NOW TRY ANSWERS

- $\{1, 2, 3, 4, 5\}$
- One answer is $\{x \mid x \text{ is a natural number between 8 and 13}\}$.

OBJECTIVE 2 Use number lines.

A good way to picture a set of numbers is to use a **number line**. See **FIGURE 8**.




FIGURE 8

To draw a number line, choose any point on the line and label it 0. Then choose any point to the right of 0 and label it 1. Use the distance between 0 and 1 as the scale to locate, and then label, other points.

The set of numbers identified on the number line in **FIGURE 8**, including positive and negative numbers and 0, is part of the set of **integers**.

$$I = \{ \dots, -3, -2, -1, 0, 1, 2, 3, \dots \} \quad \text{Integers}$$

Each number on a number line is the **coordinate** of the point that it labels, while the point is the **graph** of the number. **FIGURE 9** shows a number line with several points graphed on it.




FIGURE 9

The fractions $-\frac{1}{2}$ and $\frac{3}{4}$, graphed on the number line in **FIGURE 9**, are examples of **rational numbers**. A **rational number** can be expressed as the quotient of two integers, with denominator not 0. The set of all rational numbers is written as follows.

$$\left\{ \frac{p}{q} \mid p \text{ and } q \text{ are integers, where } q \neq 0 \right\} \quad \text{Rational numbers}$$

The set of rational numbers includes the natural numbers, whole numbers, and integers because these numbers can be written as fractions.

$$14 = \frac{14}{1}, \quad -3 = \frac{-3}{1}, \quad \text{and} \quad 0 = \frac{0}{1} \quad \text{Rational numbers}$$

A rational number written as a fraction, such as $\frac{1}{8}$ or $\frac{2}{3}$, can also be expressed as a decimal by dividing the numerator by the denominator.

$$\begin{array}{r} 0.125 \leftarrow \text{Terminating decimal} \\ 8 \overline{)1.000} \quad \text{(rational number)} \\ \underline{8} \\ 20 \\ \underline{16} \\ 40 \\ \underline{40} \\ 0 \leftarrow \text{Remainder is 0.} \end{array}$$

$$\frac{1}{8} = 0.125$$

$$\begin{array}{r} 0.666 \dots \leftarrow \text{Repeating decimal} \\ 3 \overline{)2.000} \dots \quad \text{(rational number)} \\ \underline{18} \\ 20 \\ \underline{18} \\ 20 \\ \underline{18} \\ 2 \leftarrow \text{Remainder is never 0.} \end{array}$$

$$\frac{2}{3} = 0.\overline{6} \leftarrow \text{A bar is written over the repeating digit(s).}$$

Thus terminating decimals, such as $0.125 = \frac{1}{8}$, $0.8 = \frac{4}{5}$, and $2.75 = \frac{11}{4}$, and decimals that have a repeating block of digits, such as $0.\overline{6} = \frac{2}{3}$ and $0.\overline{27} = \frac{3}{11}$, are rational numbers.

Decimal numbers that neither terminate nor repeat, which include many square roots, are **irrational numbers**.

$$\sqrt{2} = 1.414213562 \dots \quad \text{and} \quad -\sqrt{7} = -2.6457513 \dots \quad \text{Irrational numbers}$$



NOTE Some square roots, such as $\sqrt{16} = 4$ and $\sqrt{\frac{9}{25}} = \frac{3}{5}$, are rational.

A decimal number such as $0.010010001 \dots$ has a pattern, but it is irrational because there is no fixed block of digits that repeat. Another irrational number is π . See **FIGURE 10**.




FIGURE 10

Some rational and irrational numbers are graphed on the number line in **FIGURE 11**. The rational numbers together with the irrational numbers make up the set of **real numbers**.




FIGURE 11

Every point on a number line corresponds to a real number, and every real number corresponds to a point on the number line.

OBJECTIVE 3 Classify numbers.

Sets of Numbers

Natural numbers, or counting numbers	$\{1, 2, 3, 4, 5, 6, \dots\}$
Whole numbers	$\{0, 1, 2, 3, 4, 5, 6, \dots\}$
Integers	$\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
Rational numbers	$\left\{\frac{p}{q} \mid p \text{ and } q \text{ are integers, where } q \neq 0\right\}$ Examples: $\frac{4}{1}$, 1.3 , $-\frac{9}{2}$ or $-4\frac{1}{2}$, $\frac{16}{8}$ or 2 , $\sqrt{9}$ or 3 , $0.\bar{6}$
Irrational numbers	$\{x \mid x \text{ is a real number that cannot be represented by a terminating or repeating decimal}\}$ Examples: $\sqrt{3}$, $-\sqrt{2}$, π , $0.010010001 \dots$
Real numbers	$\{x \mid x \text{ is a rational or an irrational number}\}^*$

*An example of a number that is not real is $\sqrt{-1}$. This number is part of the *complex number system* and is discussed later in the text.

FIGURE 12 shows the set of real numbers. *Every real number is either rational or irrational.* Notice that the integers are elements of the set of rational numbers and that the whole numbers and natural numbers are elements of the set of integers.




FIGURE 12

NOW TRY EXERCISE 3

List the numbers in the following set that are elements of each set.

$$\{-2.4, -\sqrt{1}, -\frac{1}{2}, 0, 0.\bar{3}, \sqrt{5}, \pi, 5\}$$

- (a) Whole numbers
- (b) Rational numbers

NOW TRY EXERCISE 4

Determine whether each statement is *true* or *false*. If it is false, tell why.

- (a) All integers are irrational numbers.
- (b) Every whole number is an integer.

NOW TRY ANSWERS

- 3. (a) 0, 5
- (b) $-2.4, -\sqrt{1}, -\frac{1}{2}, 0, 0.\bar{3}, 5$
($-\sqrt{1} = -1$)
- 4. (a) false; All integers are rational numbers.
- (b) true

EXAMPLE 3 Identifying Examples of Number Sets

List the numbers in the following set that are elements of each set.

$$\{-8, -\sqrt{2}, -\frac{9}{64}, 0, 0.5, \frac{1}{3}, 1.\bar{1}2, \sqrt{3}, 2, \pi\}$$

- (a) Integers
-8, 0, and 2
- (b) Rational numbers
 $-8, -\frac{9}{64}, 0, 0.5, \frac{1}{3}, 1.\bar{1}2,$ and 2
- (c) Irrational numbers
 $-\sqrt{2}, \sqrt{3},$ and π
- (d) Real numbers
All are real numbers.

NOW TRY

EXAMPLE 4 Determining Relationships between Sets of Numbers

Determine whether each statement is *true* or *false*. If it is false, tell why.

- (a) All irrational numbers are real numbers.
This statement is true. As shown in **FIGURE 12**, the set of real numbers includes all irrational numbers.
- (b) Every rational number is an integer.
This statement is false. Although some rational numbers are integers, other rational numbers, such as $\frac{2}{3}$ and $-\frac{1}{4}$, are not.

NOW TRY

OBJECTIVE 4 Find additive inverses.

In **FIGURE 13**, for each positive number, there is a negative number on the opposite side of 0 that lies the same distance from 0. These pairs of numbers are *additive inverses*, *opposites*, or *negatives* of each other. For example, 3 and -3 are additive inverses.




FIGURE 13

▼ Additive Inverses of Signed Numbers

Number	Additive Inverse
6	-6
-4	4
$\frac{2}{3}$	$-\frac{2}{3}$
-8.7	8.7
0	0

The number 0 is its own additive inverse.

Additive Inverse

For any real number a , the number $-a$ is the **additive inverse** of a .

We change the sign of a number to find its additive inverse. As we shall see later, the sum of a number and its additive inverse is always 0.

Uses of the $-$ Symbol

The $-$ symbol can be used to indicate any of the following.

1. A negative number, as in -9 , read “*negative 9*”
2. The additive inverse of a number, as in “ -4 is the additive inverse of 4”
3. Subtraction, as in $12 - 3$, read “12 *minus* 3”

In the expression $-(-5)$, the $-$ symbol is being used in two ways. The first $-$ symbol indicates the additive inverse (or opposite) of -5 , and the second indicates a negative number, -5 . Because the additive inverse of -5 is 5,

$$-(-5) = 5.$$

$-(-a)$

For any real number a , $-(-a) = a$.

Numbers written with positive or negative signs, such as $+4$, $+8$, -9 , and -5 , are **signed numbers**. A positive number can be called a signed number even though the positive sign is usually left off.

OBJECTIVE 5 Use absolute value.

Geometrically, the **absolute value** of a number a , written $|a|$, is the distance on the number line from 0 to a . For example, the absolute value of 5 is the same as the absolute value of -5 because each number lies five units from 0. See **FIGURE 14**.




FIGURE 14

! CAUTION Because absolute value represents undirected distance, the absolute value of a number is always positive or 0.

The formal definition of absolute value follows.

Absolute Value

For any real number a , $|a| = \begin{cases} a & \text{if } a \text{ is positive or } 0 \\ -a & \text{if } a \text{ is negative.} \end{cases}$

OBJECTIVE 6 Use inequality symbols.

The statement

$$4 + 2 = 6$$

is an **equation**—a statement that two quantities are equal. The statement

$$4 \neq 6 \quad (\text{read “4 is not equal to 6”})$$

is an **inequality**—a statement that two quantities are *not* equal. When two numbers are not equal, one must be less than the other. When reading from left to right, the symbol $<$ means “*is less than*.”

$$8 < 9, \quad -6 < 15, \quad -6 < -1, \quad \text{and} \quad 0 < \frac{4}{3} \quad \text{All are true.}$$

Reading from left to right, the symbol $>$ means “*is greater than*.”

$$12 > 5, \quad 9 > -2, \quad -4 > -6, \quad \text{and} \quad \frac{6}{5} > 0 \quad \text{All are true.}$$

In each case, the symbol “points” toward the lesser number.

The number line in **FIGURE 15** shows the graphs of the numbers 4 and 9. On the graph, 4 is to the *left* of 9, so $4 < 9$. *The lesser of two numbers is always to the left of the other on a number line.*



FIGURE 15

Inequalities on a Number Line

On a number line, the following hold true.

$$a < b \text{ if } a \text{ is to the left of } b. \quad a > b \text{ if } a \text{ is to the right of } b.$$

NOW TRY EXERCISE 7

Use a number line to determine whether each statement is *true* or *false*.

- (a) $-5 > -1$
 (b) $-7 < -6$

EXAMPLE 7 Determining Order on a Number Line

Use a number line to compare -6 and 1 and to compare -5 and -2 .

As shown on the number line in **FIGURE 16**, -6 is located to the left of 1 . For this reason, $-6 < 1$. Also, $1 > -6$. From **FIGURE 16**, $-5 < -2$, or $-2 > -5$. In each case, the symbol points to the lesser number.




FIGURE 16

NOW TRY

The following table summarizes this discussion.

▼ Results about Positive and Negative Numbers

Words	Symbols
Every negative number is less than 0.	If a is negative, then $a < 0$.
Every positive number is greater than 0.	If a is positive, then $a > 0$.
0 is neither positive nor negative.	

NOW TRY ANSWERS

7. (a) false (b) true

In addition to the symbols \neq , $<$, and $>$, the symbols \leq and \geq are often used.

▼ Inequality Symbols

Symbol	Meaning	Example
\neq	is not equal to	$3 \neq 7$
$<$	is less than	$-4 < -1$
$>$	is greater than	$3 > -2$
\leq	is less than or equal to	$6 \leq 6$
\geq	is greater than or equal to	$-8 \geq -10$

NOW TRY EXERCISE 8

Determine whether each statement is *true* or *false*.

- (a) $-5 \leq -6$
 (b) $-2 \geq -10$
 (c) $0.5 \leq 0.5$
 (d) $10 \cdot 6 > 8(7)$

EXAMPLE 8 Using Inequality Symbols

The table shows several uses of inequalities and indicates why each is true.

Inequality	Why It Is True
$6 \leq 8$	$6 < 8$
$-2 \leq -2$	$-2 = -2$
$-9 \geq -12$	$-9 > -12$
$-3 \geq -3$	$-3 = -3$
$6 \cdot 4 \leq 5(5)$	$24 < 25$

Notice the reason why $-2 \leq -2$ is true. *With the symbol \leq , if either the $<$ part or the $=$ part is true, then the inequality is true. This is also the case with the \geq symbol.*

The dot in $6 \cdot 4$ indicates the product 6×4 , or 24, and $5(5)$ means 5×5 , or 25. Thus, $6 \cdot 4 \leq 5(5)$ becomes $24 \leq 25$, which is true.

NOW TRY ANSWERS

8. (a) false (b) true
 (c) true (d) $60 > 56$; true


NOW TRY 

R.2 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

- Concept Check** A student claimed that $\{x \mid x \text{ is a natural number greater than } 3\}$ and $\{y \mid y \text{ is a natural number greater than } 3\}$ actually name the same set, even though different variables are used. Was this student correct?
- Concept Check** Give a real number that satisfies each condition.

(a) An integer between 6.75 and 7.75	(b) A rational number between $\frac{1}{4}$ and $\frac{3}{4}$
(c) A whole number that is not a natural number	(d) An integer that is not a whole number
(e) An irrational number between $\sqrt{4}$ and $\sqrt{9}$	(f) An irrational number that is negative

List the elements in each set. See Example 1.

- $\{x \mid x \text{ is a natural number less than } 6\}$
- $\{m \mid m \text{ is a natural number less than } 9\}$
- $\{z \mid z \text{ is an integer greater than } 4\}$
- $\{y \mid y \text{ is an integer greater than } 8\}$
- $\{z \mid z \text{ is an integer less than or equal to } 4\}$
- $\{p \mid p \text{ is an integer less than } 3\}$
- $\{a \mid a \text{ is an even integer greater than } 8\}$
- $\{k \mid k \text{ is an odd integer less than } 1\}$
- $\{p \mid p \text{ is a number whose absolute value is } 4\}$
- $\{w \mid w \text{ is a number whose absolute value is } 7\}$

55. $-|10 - 9|$

56. $-|12 - 6|$

57. $|-9| - |-3|$

58. $|-10| - |-7|$

59. $|-1| + |-2| - |-3|$

60. $|-7| + |-3| - |-10|$

Solve each problem. See Example 6.

61. The table shows the percent change in population from 2010 through 2016 for selected metropolitan areas.
62. The table gives the net trade balance, in millions of U.S. dollars, for selected U.S. trade partners for October 2017.

Metropolitan Area	Percent Change
Detroit	-5.8
Houston	9.7
Las Vegas	8.3
New Orleans	13.9
Phoenix	11.6
Toledo	-3.0

Data from U.S. Census Bureau.

- (a) Which metropolitan area had the greatest change in population? What was this change? Was it an increase or a decrease?
- (b) Which metropolitan area had the least change in population? What was this change? Was it an increase or a decrease?

Country	Trade Balance (in millions of dollars)
India	-2419
China	-35,230
Netherlands	2148
France	-1993
New Zealand	230

Data from U.S. Census Bureau.

A negative balance means that imports to the United States exceeded exports from the United States. A positive balance means that exports exceeded imports.

- (a) Which country had the greatest discrepancy between exports and imports? Explain.
- (b) Which country had the least discrepancy between exports and imports? Explain.

Sea level refers to the surface of the ocean. The depth of a body of water can be expressed as a negative number, representing average depth in feet below sea level. The altitude of a mountain can be expressed as a positive number, indicating its height in feet above sea level. The table gives selected depths and altitudes.

Body of Water	Average Depth in Feet (as a negative number)	Mountain	Altitude in Feet (as a positive number)
Pacific Ocean	-14,040	Denali	20,310
South China Sea	-4802	Point Success	14,164
Gulf of California	-2375	Matlalcueytl	14,636
Caribbean Sea	-8448	Rainier	14,410
Indian Ocean	-12,800	Steele	16,624

Data from *The World Almanac and Book of Facts*.

63. List the bodies of water in order, starting with the deepest and ending with the shallowest.
64. List the mountains in order, starting with the shortest and ending with the tallest.
65. *True or false:* The absolute value of the depth of the Pacific Ocean is greater than the absolute value of the depth of the Indian Ocean.
66. *True or false:* The absolute value of the depth of the Gulf of California is greater than the absolute value of the depth of the Caribbean Sea.

Use a number line to determine whether each statement is true or false. See Example 7.

67. $-6 < -1$ 68. $-4 < -2$ 69. $-4 > -3$ 70. $-3 > -1$
 71. $3 > -2$ 72. $6 > -3$ 73. $-3 > -3$ 74. $-5 < -5$

Rewrite each statement with $>$ so that it uses $<$ instead. Rewrite each statement with $<$ so that it uses $>$. See Example 7.

75. $6 > 2$ 76. $5 > 1$ 77. $-9 < 4$
 78. $-6 < 1$ 79. $-5 > -10$ 80. $-7 > -12$

Use an inequality symbol to write each statement. See Example 8.


81. 7 is greater than -1 . 82. -4 is less than 10.
 83. 5 is greater than or equal to 5. 84. -6 is less than or equal to -6 .
 85. $13 - 3$ is less than or equal to 10. 86. $5 + 14$ is greater than or equal to 19.
 87. $5 + 0$ is not equal to 0. 88. $6 + 7$ is not equal to -13 .

Simplify each inequality if needed. Then determine whether the statement is true or false. See Example 8.

89. $0 \leq -5$ 90. $-11 \geq 0$ 91. $7 \leq 7$
 92. $10 \geq 10$ 93. $-6 < 7 + 3$ 94. $-7 < 4 + 1$
 95. $2 \cdot 5 \geq 4 + 6$ 96. $8 + 7 \leq 3 \cdot 5$ 97. $-|-3| \geq -3$
 98. $-|-4| \leq -4$ 99. $-8 > -|-6|$ 100. $-10 > -|-4|$

The graph shows egg production in millions of eggs in selected states for 2015 and 2016. Use this graph to work each problem.

U.S. Egg Production



Data from U.S. Department of Agriculture.

101. In 2016, which states had production greater than 6000 million eggs?
 102. In which states was 2016 egg production less than 2015 egg production?
 103. If x represents 2016 egg production for Texas (TX) and y represents 2016 egg production for Ohio (OH), which is true, $x < y$ or $x > y$?
 104. If x represents 2015 egg production for Iowa (IA) and y represents 2015 egg production for Pennsylvania (PA), write two inequalities, one with $>$ and one with $<$, that compare the numbers of eggs produced in these two states.

R.3 Operations on Real Numbers

OBJECTIVES

- 1 Add real numbers.
- 2 Subtract real numbers.
- 3 Find the distance between two points on a number line.
- 4 Multiply real numbers.
- 5 Find reciprocals and divide real numbers.

VOCABULARY

- sum
- difference
- product
- quotient
- reciprocal (multiplicative inverse)
- dividend
- divisor

NOW TRY EXERCISE 1

Find each sum.

(a) $-4 + (-9)$

(b) $-7.25 + (-3.57)$

(c) $-\frac{2}{5} + \left(-\frac{3}{10}\right)$

NOW TRY ANSWERS

1. (a) -13 (b) -10.82
 (c) $-\frac{7}{10}$

OBJECTIVE 1 Add real numbers.

Recall that the answer to an addition problem is a **sum**.

Adding Real Numbers

Same sign To add two numbers with the *same* sign, add their absolute values. The sum has the same sign as the given numbers.

Examples: $2 + 7 = 9$, $-2 + (-7) = -9$

Different signs To add two numbers with *different* signs, find the absolute values of the numbers, and subtract the lesser absolute value from the greater. The sum has the same sign as the number with the greater absolute value.

Examples: $-8 + 3 = -5$, $15 + (-9) = 6$

EXAMPLE 1 Adding Two Negative Real Numbers

Find each sum.

(a) $-12 + (-8)$

First find the absolute values: $|-12| = 12$ and $|-8| = 8$.

Because -12 and -8 have the *same* sign, add their absolute values.

Both numbers are negative, so the sum will be negative.

$$\begin{aligned} & -12 + (-8) \\ &= -(12 + 8) \quad \text{Add the absolute values.} \\ &= -(20) \\ &= -20 \end{aligned}$$

(b) $-6 + (-3)$

$$\begin{aligned} &= -(|-6| + |-3|) \\ &= -(6 + 3) \\ &= -9 \end{aligned}$$

(c) $-1.2 + (-0.4)$

$$\begin{aligned} &= -(1.2 + 0.4) \\ &= -1.6 \end{aligned}$$

(d) $-\frac{5}{6} + \left(-\frac{1}{3}\right)$

$$= -\left(\frac{5}{6} + \frac{1}{3}\right) \quad \text{Add the absolute values. Both numbers are negative, so the sum will be negative.}$$

$$= -\left(\frac{5}{6} + \frac{2}{6}\right) \quad \text{The least common denominator is 6.}$$

$$\frac{1}{3} \cdot \frac{2}{2} = \frac{2}{6}$$

$$= -\frac{7}{6} \quad \begin{array}{l} \text{Add numerators.} \\ \text{Keep the same denominator.} \end{array}$$


NOW TRY EXERCISE 2

Find each sum.

- (a) $-15 + 7$
 (b) $-5 + 12$
 (c) $4.6 + (-2.8)$
 (d) $-\frac{5}{9} + \frac{2}{7}$

EXAMPLE 2 Adding Real Numbers with Different Signs

Find each sum.

(a) $-17 + 11$

First find the absolute values: $|-17| = 17$ and $|11| = 11$.Because -17 and 11 have *different* signs, subtract their absolute values.

$$17 - 11 = 6$$

The number -17 has a greater absolute value than 11 , so the answer is negative.

$$-17 + 11 = -6$$

The sum is negative because $|-17| > |11|$.

(b) $4 + (-1)$

Subtract the absolute values, 4 and 1 . Because 4 has the greater absolute value, the sum must be positive.

$$4 + (-1) = 4 - 1 = 3$$

The sum is positive because $|4| > |-1|$.

(c) $-9 + 17$

$$= 17 - 9$$

$$= 8$$

(d) $-2.3 + 5.6$

$$= 5.6 - 2.3$$

$$= 3.3$$

(e) $-16 + 12$

The absolute values are 16 and 12 . Subtract the absolute values.

$$-16 + 12 = -(16 - 12) = -4$$

The sum is negative because $|-16| > |12|$.

(f) $-\frac{4}{5} + \frac{2}{3}$

$$= -\frac{12}{15} + \frac{10}{15}$$

The least common denominator is 15 .

$$-\frac{4}{5} \cdot \frac{3}{3} = -\frac{12}{15}, \quad \frac{2}{3} \cdot \frac{5}{5} = \frac{10}{15}$$

$$= -\left(\frac{12}{15} - \frac{10}{15}\right)$$

Subtract the absolute values. $-\frac{12}{15}$ has the greater absolute value, so the answer will be negative.

$$= -\frac{2}{15}$$

Subtract numerators.

Keep the same denominator.

NOW TRY 
OBJECTIVE 2 Subtract real numbers.
Recall that the answer to a subtraction problem is a **difference**. Compare the following two statements.

$$6 - 4 = 2$$

$$6 + (-4) = 2$$

Thus,

$$6 - 4 = 6 + (-4).$$

To subtract 4 from 6 , we add the additive inverse of 4 to 6 . This example suggests the following definition of subtraction of real numbers.
NOW TRY ANSWERS

2. (a) -8 (b) 7
 (c) 1.8 (d) $-\frac{17}{63}$

Subtraction

For all real numbers a and b , the following holds.

$$a - b = a + (-b)$$

To subtract b from a , add the additive inverse (or opposite) of b to a .

Examples: $5 - 12 = 5 + (-12) = -7$, $6 - (-3) = 6 + 3 = 9$

NOW TRY EXERCISE 3

Find each difference.

- (a) $9 - 15$
 (b) $-4 - 11$
 (c) $-5.67 - (-2.34)$
 (d) $\frac{4}{9} - \frac{3}{5}$

EXAMPLE 3 Subtracting Real Numbers

Find each difference.

(a) $6 - 8 = 6 + (-8) = -2$

Change to addition.
 The additive inverse of 8 is -8 .

(b) $-12 - 4 = -12 + (-4) = -16$

Change to addition.
 The additive inverse of 4 is -4 .

(c) $-10 - (-7)$
 $= -10 + 7$
 $= -3$

The additive inverse of -7 is 7.

(d) $-2.4 - (-8.1)$
 $= -2.4 + 8.1$
 $= 5.7$

(e) $\frac{5}{6} - \left(-\frac{3}{8}\right)$
 $= \frac{5}{6} + \frac{3}{8}$
 $= \frac{20}{24} + \frac{9}{24}$
 $= \frac{29}{24}$

To subtract $a - b$, add the additive inverse (opposite) of b to a .

The least common denominator is 24.
 $\frac{5}{6} \cdot \frac{4}{4} = \frac{20}{24}$, $\frac{3}{8} \cdot \frac{3}{3} = \frac{9}{24}$

Add numerators.
 Keep the same denominator.

NOW TRY

When working a problem that involves both addition and subtraction, add and subtract in order from left to right. Work inside brackets or parentheses first.

EXAMPLE 4 Adding and Subtracting Real Numbers

Perform the indicated operations.

(a) $15 - (-3) - 5 - 12$
 $= (15 + 3) - 5 - 12$ Work in order from left to right.
 $= 18 - 5 - 12$ Add inside the parentheses.
 $= 13 - 12$ Subtract from left to right.
 $= 1$ Subtract.

NOW TRY ANSWERS

3. (a) -6 (b) -15
 (c) -3.33 (d) $-\frac{7}{45}$

NOW TRY
EXERCISE 4

Perform the indicated operations.

$$-4 - (-2 - 7) - 12$$

$$\begin{aligned} \text{(b)} \quad & -10 - [-8 - (-4)] + 6 \\ & = -10 - [-8 + 4] + 6 && \text{Work inside the brackets.} \\ & = -10 - [-4] + 6 && \text{Add.} \\ & = -10 + 4 + 6 && \text{Add the additive inverse.} \\ & = -6 + 6 && \text{Add from left to right.} \\ & = 0 && \text{Add.} \end{aligned}$$

NOW TRY 

OBJECTIVE 3 Find the distance between two points on a number line.

The number line in **FIGURE 17** shows several points.




FIGURE 17

To find the distance between the points **4** and **7**, we subtract $7 - 4 = 3$. Because distance is positive (or 0), we must be careful to subtract in such a way that the answer is positive (or 0). To avoid this problem altogether, we can find the absolute value of the difference. Then the distance between 4 and 7 is found as follows.

$$|7 - 4| = |3| = 3 \quad \text{or} \quad |4 - 7| = |-3| = 3 \quad \text{Distance between 4 and 7}$$

Distance

The **distance** between two points on a number line is the absolute value of the difference of their coordinates.

EXAMPLE 5 Finding Distance between Points on a Number Line

Find the distance between each pair of points. Refer to **FIGURE 17** as needed.

(a) 8 and -4

Find the absolute value of the difference of the numbers, taken in either order.

$$|8 - (-4)| = 12, \quad \text{or} \quad |-4 - 8| = 12$$

(b) -4 and -6

$$|-4 - (-6)| = 2, \quad \text{or} \quad |-6 - (-4)| = 2$$

NOW TRY 

OBJECTIVE 4 Multiply real numbers.

Recall that the answer to a multiplication problem is a **product**.

Multiplying Real Numbers

Same sign The product of two numbers with the *same* sign is positive.

Examples: $4(8) = 32$, $-4(-8) = 32$

Different signs The product of two numbers with *different* signs is negative.

Examples: $-8(9) = -72$, $6(-7) = -42$

NOW TRY ANSWERS

4. -7
5. 17

The **multiplication property of 0** states that the product of any real number and 0 is 0.

Multiplication Property of 0

For any real number a , the following hold.

$$a \cdot 0 = 0 \quad \text{and} \quad 0 \cdot a = 0$$

Examples: $12 \cdot 0 = 0$, $0 \cdot 12 = 0$, $-12 \cdot 0 = 0$, $0 \cdot (-12) = 0$

NOW TRY EXERCISE 6

Find each product.

(a) $-3(-10)$

(b) $0.7(-1.2)$

(c) $-\frac{8}{11}(33)$

(d) $14(0)$

EXAMPLE 6 Multiplying Real Numbers

Find each product.

(a) $-3(-9) = 27$

The numbers have the same sign, so the product is positive.

(b) $-0.5(-0.4) = 0.2$

(c) $-\frac{3}{4}\left(-\frac{5}{6}\right)$

$$= \frac{15}{24}$$

Multiply numerators.

Multiply denominators.

$$= \frac{5 \cdot 3}{8 \cdot 3}$$

Factor to write in lowest terms.

$$= \frac{5}{8}$$

Divide out the common factor, 3.

(d) $-6(0) = 0$

Multiplication property of 0

(e) $6(-9) = -54$

The numbers have different signs, so the product is negative.

(f) $-0.05(0.3) = -0.015$

(g) $-\frac{3}{4}\left(\frac{2}{9}\right) = -\frac{1}{6}$

(h) $\frac{2}{3}(-6) = -4$

$$-6 = -\frac{6}{1}$$

NOW TRY

OBJECTIVE 5 Find reciprocals and divide real numbers.

The definition of division uses the concept of a **multiplicative inverse**, or *reciprocal*. Recall that two numbers are *reciprocals* if they have a product of 1.

Reciprocal

The **reciprocal** of a nonzero number a is $\frac{1}{a}$.

Reciprocals

Number	Reciprocal
$-\frac{2}{5}$	$-\frac{5}{2}$
-6 , or $-\frac{6}{1}$	$-\frac{1}{6}$
$\frac{7}{11}$	$\frac{11}{7}$
0.05	20
0	None

$$\left. \begin{aligned} -\frac{2}{5}\left(-\frac{5}{2}\right) &= 1 \\ -6\left(-\frac{1}{6}\right) &= 1 \\ \frac{7}{11}\left(\frac{11}{7}\right) &= 1 \\ 0.05(20) &= 1 \end{aligned} \right\}$$

Reciprocals have a product of 1.

There is no reciprocal for 0 because there is no number that can be multiplied by 0 to give a product of 1.

NOW TRY ANSWERS

6. (a) 30 (b) -0.84
(c) -24 (d) 0

! CAUTION Keep the following in mind.

- A number and its additive inverse have opposite signs, such as 3 and -3 .
- A number and its reciprocal always have the same sign, such as 3 and $\frac{1}{3}$.

The result of dividing one number by another is a **quotient**. For example, we can write the quotient of 45 and 3 as $\frac{45}{3}$, which equals 15. We obtain the same answer if we multiply 45 and $\frac{1}{3}$.

$$45 \div 3 = \frac{45}{3} = 15 \quad \text{and} \quad 45 \cdot \frac{1}{3} = 15$$

This suggests the following definition of division of real numbers.

Division

For all real numbers a and b (where $b \neq 0$), the following holds true.

$$a \div b = \frac{a}{b} = a \cdot \frac{1}{b}$$

To divide a by b , multiply a (the **dividend**) by the reciprocal of b (the **divisor**).

There is no reciprocal for the number 0, so **division by 0 is undefined**. For example, $\frac{15}{0}$ is undefined and $-\frac{1}{0}$ is undefined.

! CAUTION Although division by 0 is undefined, dividing 0 by a nonzero number gives the quotient 0.

Example: $\frac{6}{0}$ is undefined, but $\frac{0}{6} = 0$ (because $0 \cdot 6 = 0$).

Be careful when 0 is involved in a division problem.

Because division is defined as multiplication by the reciprocal, the rules for signs of quotients are the same as those for signs of products.

Dividing Real Numbers

Same sign The quotient of two nonzero real numbers with the *same* sign is positive.

Examples: $\frac{24}{6} = 4$, $\frac{-24}{-6} = 4$

Different signs The quotient of two nonzero real numbers with *different* signs is negative.

Examples: $\frac{-36}{3} = -12$, $\frac{36}{-3} = -12$


NOW TRY EXERCISE 7

Find each quotient.

(a) $\frac{-10}{-5}$ (b) $\frac{-1.5}{0.3}$

(c) $\frac{0}{-4}$ (d) $\frac{-\frac{10}{3}}{\frac{3}{8}}$

(e) $-\frac{6}{5} \div \left(-\frac{3}{7}\right)$

EXAMPLE 7 Dividing Real Numbers

Find each quotient.

(a) $\frac{-12}{4} = -3$ The numbers have opposite signs, so the quotient is negative.

(b) $\frac{6}{-3} = -2$

(c) $\frac{-3.6}{-0.4} = 9$ The numbers have the same sign, so the quotient is positive.

(d) $\frac{-9}{0}$ is **undefined**.

(e) $\frac{0}{-12} = 0$ This is true because $0(-12) = 0$.

(f) $\frac{-\frac{2}{3}}{-\frac{5}{9}}$ This is a *complex fraction*. A complex fraction has a fraction in the numerator, the denominator, or both.

$$= -\frac{2}{3} \left(-\frac{9}{5}\right)$$
 $\frac{a}{b} = a \div b$; Multiply by $-\frac{9}{5}$, the reciprocal of the divisor $-\frac{5}{9}$.

$$= \frac{18}{15}$$
 Multiply numerators.
Multiply denominators.

$$= \frac{6}{5}$$
 Write in lowest terms; $\frac{18}{15} = \frac{2 \cdot 3 \cdot 3}{3 \cdot 5} = \frac{6}{5}$

(g) $-\frac{9}{14} \div \frac{3}{7}$

$$= -\frac{9}{14} \cdot \frac{7}{3}$$
 Multiply by $\frac{7}{3}$, the reciprocal of the divisor $\frac{3}{7}$.

$$= -\frac{63}{42}$$
 Multiply numerators.
Multiply denominators.


$$= -\frac{3}{2}$$
 Write in lowest terms; $-\frac{63}{42} = -\frac{3 \cdot 3 \cdot 7}{2 \cdot 3 \cdot 7} = -\frac{3}{2}$

NOW TRY 

Every fraction has three signs: the sign of the numerator, the sign of the denominator, and the sign of the fraction itself.

Equivalent Forms of a FractionThe fractions $\frac{-x}{y}$, $\frac{x}{-y}$, and $-\frac{x}{y}$ are equivalent (where $y \neq 0$).The fractions $\frac{x}{y}$ and $\frac{-x}{-y}$ are equivalent (where $y \neq 0$).Examples: $\frac{-4}{7} = \frac{4}{-7} = -\frac{4}{7}$ and $\frac{4}{7} = \frac{-4}{-7}$ **NOW TRY ANSWERS**

7. (a) 2 (b) -5 (c) 0
-
- (d)
- $-\frac{80}{9}$
- (e)
- $\frac{14}{5}$

R.3 Exercises**FOR EXTRA HELP****MyLab Math** Video solutions for select problems available in MyLab Math**Concept Check** Complete each statement and give an example.

- The sum of two positive numbers is a _____ number.
- The sum of two negative numbers is a _____ number.

3. The sum of a positive number and a negative number is negative if the negative number has the _____ absolute value.
4. The sum of a positive number and a negative number is positive if the positive number has the _____ absolute value.
5. The difference of two positive numbers is negative if _____.
6. The difference of two negative numbers is negative if _____.
7. The product of two numbers with the same sign is _____.
8. The product of two numbers with different signs is _____.
9. The quotient formed by any nonzero number divided by 0 is _____, and the quotient formed by 0 divided by any nonzero number is _____.
10. The sum of a positive number and a negative number is 0 if the numbers are _____.


Find each sum or difference. See Examples 1–3.

- | | | |
|----------------------------------|------------------------------------------------|------------------------------------------------|
| 11. $-6 + (-13)$ | 12. $-8 + (-16)$ | 13. $-15 + 6$ |
| 14. $-17 + 9$ | 15. $13 + (-4)$ | 16. $19 + (-13)$ |
| 17. $-17 + 22$ | 18. $-12 + 16$ | 19. $-\frac{7}{3} + \frac{3}{4}$ |
| 20. $-\frac{5}{6} + \frac{4}{9}$ | 21. $-2.8 + 4.5$ | 22. $-3.8 + 6.2$ |
| 23. $4 - 9$ | 24. $3 - 7$ | 25. $-6 - 5$ |
| 26. $-8 - 17$ | 27. $8 - (-13)$ | 28. $12 - (-22)$ |
| 29. $-16 - (-3)$ | 30. $-21 - (-6)$ | 31. $-12.31 - (-2.13)$ |
| 32. $-15.88 - (-9.42)$ | 33. $\frac{9}{10} - \left(-\frac{4}{3}\right)$ | 34. $\frac{3}{14} - \left(-\frac{3}{4}\right)$ |
| 35. $ -8 - 6 $ | 36. $ -7 - 15 $ | 37. $- -4 + 9 $ |
| 38. $- -5 + 6 $ | 39. $-2 - -4 $ | 40. $16 - -13 $ |

Perform the indicated operations. See Example 4.

- | | |
|-------------------------------------------------------------|--------------------------------------------------------------|
| 41. $-7 + 5 - 9$ | 42. $-12 + 14 - 18$ |
| 43. $6 - (-2) - 8$ | 44. $7 - (-4) - 11$ |
| 45. $8 - (-12) - 2 - 6$ | 46. $3 - (-14) - 6 - 4$ |
| 47. $-9 - 4 - (-3) + 6$ | 48. $-10 - 6 - (-12) + 9$ |
| 49. $-0.38 + 4 - 0.62$ | 50. $-2.95 + 8 - 0.05$ |
| 51. $\left(-\frac{5}{4} - \frac{2}{3}\right) + \frac{1}{6}$ | 52. $\left(-\frac{5}{9} - \frac{1}{6}\right) + \frac{1}{2}$ |
| 53. $-\frac{3}{4} - \left(\frac{1}{2} - \frac{3}{8}\right)$ | 54. $-\frac{1}{2} - \left(\frac{5}{6} - \frac{5}{12}\right)$ |
| 55. $-4 - [-3 - (-6)] + 9$ | 56. $-10 - [-2 - (-5)] + 16$ |
| 57. $ -11 - -5 - 7 + -2 $ | 58. $ -6 + -3 - 4 - -8 $ |

The number line has several points labeled. Find the distance between each pair of points. See Example 5.



59. A and B 60. A and C 61. D and F 62. E and C

63. **Concept Check** Consider the statement “Two negatives give a positive.” When is this true? When is this false? Give a more precise statement that conveys this message.
64. **Concept Check** Why must the reciprocal of a nonzero number have the same sign as the number?

Find each product. See Example 6.

65. $-8(-5)$ 66. $-20(-4)$ 67. $5(-7)$ 68. $6(-9)$
69. $4(0)$ 70. $0(-8)$ 71. $\frac{1}{2}(0)$ 72. $0(-4.5)$
73. $\frac{3}{4}(-16)$ 74. $\frac{3}{5}(-35)$ 75. $-10\left(-\frac{1}{5}\right)$ 76. $-18\left(-\frac{1}{2}\right)$
77. $-\frac{5}{2}\left(-\frac{12}{25}\right)$ 78. $-\frac{9}{7}\left(-\frac{21}{36}\right)$ 79. $-\frac{3}{8}\left(-\frac{24}{9}\right)$ 80. $-\frac{2}{11}\left(-\frac{22}{4}\right)$
81. $-0.8(-0.5)$ 82. $-0.5(-0.6)$ 83. $-0.06(0.4)$ 84. $-0.08(0.7)$

Find each quotient where possible. See Example 7.

85. $\frac{-14}{2}$ 86. $\frac{-39}{13}$ 87. $\frac{-24}{-4}$ 88. $\frac{-45}{-9}$ 89. $\frac{100}{-25}$
90. $\frac{150}{-30}$ 91. $\frac{0}{-8}$ 92. $\frac{0}{-14}$ 93. $\frac{5}{0}$ 94. $\frac{13}{0}$
95. $-\frac{10}{17} \div \left(-\frac{12}{5}\right)$ 96. $-\frac{22}{23} \div \left(-\frac{33}{5}\right)$ 97. $\frac{\frac{12}{13}}{\frac{-4}{-3}}$ 98. $\frac{\frac{7}{6}}{\frac{-2}{-3}}$
99. $\frac{-7.2}{0.8}$ 100. $\frac{-4.5}{0.9}$ 101. $\frac{-1.28}{-0.4}$ 102. $\frac{-1.82}{-0.2}$

The following problems provide more practice on operations with fractions and decimals. Perform the indicated operations.

103. $\frac{1}{6} - \left(-\frac{7}{9}\right)$ 104. $\frac{7}{10} - \left(-\frac{1}{6}\right)$ 105. $-\frac{1}{9} + \frac{7}{12}$
106. $-\frac{1}{12} + \frac{13}{16}$ 107. $-\frac{3}{8} - \frac{5}{12}$ 108. $-\frac{11}{15} - \frac{4}{9}$
109. $-\frac{7}{30} + \frac{2}{45} - \frac{3}{10}$ 110. $-\frac{8}{15} + \frac{7}{6} - \frac{3}{20}$ 111. $\frac{8}{25}\left(-\frac{5}{12}\right)$
112. $\frac{9}{20}\left(-\frac{7}{15}\right)$ 113. $\frac{5}{6}\left(-\frac{9}{10}\right)\left(-\frac{4}{5}\right)$ 114. $\frac{4}{3}\left(-\frac{9}{20}\right)\left(-\frac{5}{12}\right)$
115. $\frac{8}{3} \div \left(-\frac{14}{15}\right)$ 116. $\frac{12}{5} \div \left(-\frac{18}{25}\right)$ 117. $\frac{\frac{2}{3}}{-2}$

118. $\frac{\frac{3}{4}}{-6}$

119. $\frac{-\frac{8}{9}}{\frac{2}{3}}$

120. $\frac{-\frac{15}{16}}{\frac{3}{8}}$

121. $-8.6 - 23.751$

122. $-37.8 - 13.582$

123. $-2.5(0.8)(1.5)$

124. $-1.6(0.5)(2.5)$

125. $-24.84 \div 6$

126. $-32.84 \div 8$

127. $-2496 \div (-0.52)$

128. $-1875 \div (-0.25)$

129. $\frac{-100}{-0.01}$


130. $\frac{-60}{-0.06}$

131. $-14.2 + 9.81$


132. $-89.41 + 21.325$

Solve each problem.

133. On June 27, 1915, a temperature of 100°F was recorded in Fort Yukon, Alaska. On January 23, 1971, Prospect Creek Camp, Alaska, recorded a temperature of -80°F . Express the difference between these two temperatures as a positive number. (Data from *The World Almanac and Book of Facts*.)





134. On August 10, 1936, a temperature of 120°F was recorded in Ozark, Arkansas. On February 13, 1905, Gravette, Arkansas, recorded a temperature of -29°F . Express the difference between these two temperatures as a positive number. (Data from *The World Almanac and Book of Facts*.)



135. Andrew has \$48.35 in his checking account. He uses his debit card to make purchases of \$35.99 and \$20.00, which overdraws his account. His bank charges his account an overdraft fee of \$28.50. He then deposits his paycheck for \$66.27 from his part-time job at Arby's. What is the balance in his account?
136. Kayla has \$37.60 in her checking account. She uses her debit card to make purchases of \$25.99 and \$19.34, which overdraws her account. Her bank charges her account an overdraft fee of \$25.00. She then deposits her paycheck for \$58.66 from her part-time job at Subway. What is the balance in her account?
137. Ahmad owes \$382.45 on his Visa account. He returns two items costing \$25.10 and \$34.50 for credit. Then he makes purchases of \$45.00 and \$98.17.
- (a) How much is his payment if he wants to pay off the balance on the account?
- (b) Instead of paying off the balance, he makes a payment of \$300 and then incurs a finance charge of \$24.66. What is the balance on his account?
138. Charlene owes \$237.59 on her MasterCard account. She returns one item costing \$47.25 for credit and then makes two purchases of \$12.39 and \$20.00.
- (a) How much is her payment if she wants to pay off the balance on the account?
- (b) Instead of paying off the balance, she makes a payment of \$75.00 and incurs a finance charge of \$32.06. What is the balance on her account?



- 139.** The graph shows annual returns in percent for shares of a charter fund.
- Find the sum of the percents for the years shown in the graph.
 - Find the difference between the returns in 2017 and 2016.
 - Find the difference between the returns in 2018 and 2017.
- 140.** The graph shows profits and losses in thousands of dollars for a company.
- What was the total profit or loss for the years 2015 through 2018?
 - Find the difference between the profit or loss in 2018 and that in 2017.
 - Find the difference between the profit or loss in 2016 and that in 2015.



The table shows receipts (income) and outlays (spending) in billions of dollars for the U.S. government in selected years.

Fiscal Year	Receipts	Outlays
2001	1991	1863
2006	2407	2655
2011	2303	3603
2016	3268	3853

Data from Office of Management and Budget.

- 141.** Find the difference between U.S. government receipts and outlays for each year shown in the table.
- 142.** During which years did the budget show a surplus? A deficit? Explain the answers.

R.4 Exponents, Roots, and Order of Operations

OBJECTIVES

- Use exponents.
- Find square roots.
- Use the rules for order of operations.
- Evaluate algebraic expressions for given values of variables.

Two (or more) numbers whose product is a third number are **factors** of that third number. For example, 2 and 6 are factors of 12 because $2 \cdot 6 = 12$.

OBJECTIVE 1 Use exponents.

In algebra, we use *exponents* as a way of writing products of repeated factors. For example, we write the product $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$ as follows.

$$\underbrace{2 \cdot 2 \cdot 2 \cdot 2 \cdot 2}_{5 \text{ factors of } 2} = 2^5$$

The number 5 shows that 2 is used as a factor 5 times. The number 5 is the *exponent*, and 2 is the *base*.

$$2^5 \leftarrow \text{Exponent}$$

$$\uparrow$$

$$\text{Base}$$

Read 2^5 as “2 to the fifth power,” or “2 to the fifth.” Multiplying five 2s gives 32.

$$2^5 \text{ means } 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2, \text{ which equals } 32.$$

VOCABULARY

- factors
- exponent (power)
- base
- exponential expression
- square root
- positive (principal) square root
- negative square root
- constant
- algebraic expression

Exponential Expression

If a is a real number and n is a natural number, then

$$a^n = \underbrace{a \cdot a \cdot a \cdot \dots \cdot a}_{n \text{ factors of } a},$$

where n is the **exponent**, a is the **base**, and a^n is an **exponential expression**. Exponents are also called **powers**.



NOW TRY EXERCISE 1

Write using exponents.

(a) $(-3)(-3)(-3)$

(b) $(0.5)(0.5)$

(c) $t \cdot t \cdot t \cdot t \cdot t$

EXAMPLE 1 Using Exponential Notation

Write using exponents.

(a) $\underbrace{4 \cdot 4 \cdot 4}_{3 \text{ factors of } 4} = 4^3$

Read 4^3 as “4 *cubed*.”

(b) $\frac{3}{5} \cdot \frac{3}{5} = \left(\frac{3}{5}\right)^2$ 2 factors of $\frac{3}{5}$

Read $\left(\frac{3}{5}\right)^2$ as “ $\frac{3}{5}$ *squared*.”

(c) $(-6)(-6)(-6)(-6) = (-6)^4$ 4 factors of -6

Read $(-6)^4$ as “ -6 to the fourth power,” or “ -6 to the fourth.”

(d) $(0.3)(0.3)(0.3)(0.3)(0.3) = (0.3)^5$ (e) $x \cdot x \cdot x \cdot x \cdot x \cdot x = x^6$ NOW TRY

NOTE In **Example 1**, we used the terms *squared* and *cubed* to refer to powers of 2 and 3, respectively. The term *squared* comes from the figure of a square, which has the same measure for both length and width. See **FIGURE 18(a)**. Similarly, the term *cubed* comes from the figure of a cube, where the length, width, and height have the same measure. See **FIGURE 18(b)**.




FIGURE 18

EXAMPLE 2 Evaluating Exponential Expressions

Evaluate.

(a) 5^2 means $5 \cdot 5$, which equals 25. 5 is used as a factor 2 times.

5^2 means $5 \cdot 5$, *not* $5 \cdot 2$.

NOW TRY ANSWERS

1. (a) $(-3)^3$ (b) $(0.5)^2$ (c) t^5


NOW TRY EXERCISE 2

Evaluate.

- (a) 5^3 (b) $\left(\frac{2}{5}\right)^4$
 (c) $(-3)^2$ (d) $(-3)^3$

- (b) $\left(\frac{2}{3}\right)^3$ means $\frac{2}{3} \cdot \frac{2}{3} \cdot \frac{2}{3}$, which equals $\frac{8}{27}$. $\frac{2}{3}$ is used as a factor 3 times.
 (c) $(0.2)^3$ means $(0.2)(0.2)(0.2)$, which equals 0.008.
 (d) $(-2)^4$ means $(-2)(-2)(-2)(-2)$, which equals 16.
 (e) $(-2)^5$ means $(-2)(-2)(-2)(-2)(-2)$, which equals -32 .

NOW TRY 

Examples 2(d) and (e) suggest the following generalizations.

Sign of an Exponential Expression

The product of an *even* number of negative factors is positive.

Example: $(-2)(-2)(-2)(-2) = 16$ 4 factors of -2

The product of an *odd* number of negative factors is negative.

Example: $(-2)(-2)(-2)(-2)(-2) = -32$ 5 factors of -2


NOW TRY EXERCISE 3

Evaluate.

- (a) 7^2 (b) $(-7)^2$ (c) -7^2

EXAMPLE 3 Evaluating Exponential Expressions

Evaluate.

(a) 2^6 means $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$, which equals 64. The base is 2.

(b) $(-2)^6$ means $(-2)(-2)(-2)(-2)(-2)(-2)$, which equals 64. The base is -2 .

(c) -2^6

There are no parentheses. The exponent 6 applies *only* to the number 2, not to -2 .

-2^6 means $-(2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2)$, which equals -64 . The base is 2.

NOW TRY 

! CAUTION It is important to distinguish between $-a^n$ and $(-a)^n$.

$-a^n$ means $-1 \underbrace{(a \cdot a \cdot a \cdot \dots \cdot a)}_{n \text{ factors of } a}$ The base is a .

$(-a)^n$ means $\underbrace{(-a)(-a) \cdot \dots \cdot (-a)}_{n \text{ factors of } -a}$ The base is $-a$.

Be careful when evaluating an exponential expression with a negative sign.

OBJECTIVE 2 Find square roots.

As we saw in **Example 2(a)**, 5 squared (or 5^2) equals 25. The opposite (inverse) of squaring a number is taking its **square root**. For example, a square root of 25 is 5. Another square root of 25 is -5 because $(-5)^2 = 25$. Thus, 25 has two square roots, 5 and -5 .

We write the **positive** or **principal square root** of a number using a **radical symbol** $\sqrt{\quad}$. The positive or principal square root of 25 is written

$$\sqrt{25} = 5.$$

The **negative square root** of 25 is written

$$-\sqrt{25} = -5.$$

NOW TRY ANSWERS

2. (a) 125 (b) $\frac{16}{625}$ (c) 9
 (d) -27
 3. (a) 49 (b) 49 (c) -49

Because the square of any nonzero real number is positive, the square root of a negative number, such as $\sqrt{-25}$, is not a real number.

NOW TRY EXERCISE 4

Find each square root that is a real number.

(a) $\sqrt{121}$ (b) $\sqrt{\frac{100}{9}}$
 (c) $-\sqrt{121}$ (d) $\sqrt{-121}$

EXAMPLE 4 Finding Square Roots

Find each square root that is a real number.

- (a) $\sqrt{36} = 6$ because $6^2 = 36$. (b) $\sqrt{0} = 0$ because $0^2 = 0$.
 (c) $\sqrt{\frac{9}{16}} = \frac{3}{4}$ because $\left(\frac{3}{4}\right)^2 = \frac{9}{16}$. (d) $\sqrt{0.16} = 0.4$ because $(0.4)^2 = 0.16$.
 (e) $\sqrt{100} = 10$ because $10^2 = 100$.
 (f) $-\sqrt{100} = -10$ because the negative sign is outside the radical symbol.
 (g) $\sqrt{-100}$ is not a real number because the negative sign is inside the radical symbol. No *real number* squared equals -100 .

Notice that part (e) is the positive or principal square root of 100, part (f) is the negative square root of 100, and part (g) is the square root of -100 , which is not a real number.

NOW TRY 

OBJECTIVE 3 Use the rules for order of operations.

To simplify the following expression, what should we do first—add 5 and 2, or multiply 2 and 3?

$$5 + 2 \cdot 3$$

When an expression involves more than one operation symbol, we use the following rules for **order of operations**.

Order of Operations

If grouping symbols are present, work within them, innermost first (and above and below fraction bars separately), in the following order.

Step 1 Apply all **exponents**.

Step 2 Do any **multiplications** and **divisions** in order from left to right.

Step 3 Do any **additions** and **subtractions** in order from left to right.

If no grouping symbols are present, start with Step 1.

NOW TRY EXERCISE 5

Simplify.

$$15 - 3 \cdot 4 + 2$$

NOW TRY ANSWERS

4. (a) 11 (b) $\frac{10}{3}$ (c) -11
 (d) not a real number
 5. 5

EXAMPLE 5 Using the Rules for Order of Operations

Simplify.

- | | | |
|----------------------------|-----------------------------|----------------------------------------------------------------------------------------------|
| (a) $5 + 2 \cdot 3$ | (b) $24 \div 3 \cdot 2 + 6$ | Multiplications and divisions are done in order from left to right. So we divide first here. |
| $= 5 + 6$ Multiply. | $= 8 \cdot 2 + 6$ | |
| $= 11$ Add. | $= 16 + 6$ Multiply. | |
| | $= 22$ Add. | |

NOW TRY 


NOW TRY EXERCISE 6

Simplify.

(a) $-5^2 + 10 \div 5 - |3 - 7|$

(b) $6 + \frac{2}{3}(-9) - \frac{5}{8} \cdot 16$

EXAMPLE 6 Using the Rules for Order of Operations

Simplify.

(a) $10 \div 5 + 2|3 - 4|$ *Work inside the absolute value bars first.*

$= 10 \div 5 + 2|-1|$ *Subtract inside the absolute value bars.*

$= 10 \div 5 + 2 \cdot 1$ *Take the absolute value.*

$= 2 + 2$ *Divide first, and then multiply.*

$= 4$ *Add.*

(b) $4 - 3^2 + 7 - (2 + 8)$ *Work inside the parentheses first.*

$= 4 - 3^2 + 7 - 10$ *Add inside the parentheses.*

$= 4 - 9 + 7 - 10$ *Evaluate the power.*

The base is 3. 3^2 means $3 \cdot 3$, *not* $3 \cdot 2$.

$= -5 + 7 - 10$

$= 2 - 10$

$= -8$

Additions and subtractions are done in order from left to right, so subtract $4 - 9$ first here, then add $-5 + 7$.

(c) $\frac{1}{2} \cdot 4 + (6 \div 3 - 7)$ *Work inside the parentheses first.*

$= \frac{1}{2} \cdot 4 + (2 - 7)$ *Divide inside the parentheses.*

$= \frac{1}{2} \cdot 4 + (-5)$ *Subtract inside the parentheses.*

$= 2 + (-5)$ *Multiply.*

$= -3$ *Add.*

NOW TRY 

NOW TRY EXERCISE 7

Simplify.

$$\frac{\sqrt{36} - 4 \cdot 3^2}{-2^2 - 8 \cdot 3 + 28}$$

EXAMPLE 7 Using the Rules for Order of Operations

Simplify.

$$\frac{5 - 2^4}{6 \cdot \sqrt{9} - 9 \cdot 2}$$
Work separately above and below the fraction bar.

$$= \frac{5 - 16}{6 \cdot 3 - 9 \cdot 2}$$
Evaluate the power and the root.

$$= \frac{5 - 16}{18 - 18}$$
Multiply.

$$= \frac{-11}{0}$$
Subtract.

Because division by 0 is undefined, the given expression is undefined. **NOW TRY** 

NOW TRY ANSWERS

6. (a) -27 (b) -10
7. undefined

OBJECTIVE 4 Evaluate algebraic expressions for given values of variables.

A **constant** is a fixed, unchanging number.

$$1, \quad 6, \quad -10, \quad \frac{2}{5}, \quad -3.75 \quad \text{Constants}$$

A collection of constants, variables, operation symbols, and/or grouping symbols is an **algebraic expression**.

$$6ab, \quad 5m - 9n, \quad -2(x^2 + 4y) \quad \text{Algebraic expressions}$$

Algebraic expressions have different numerical values for different values of the variables. We evaluate such expressions by *substituting* given values for the variables.

 **NOW TRY EXERCISE 8**

Evaluate each expression for $x = -4$, $y = 7$, and $z = 36$.

(a) $3y - 2x$ (b) $\frac{x^2 - \sqrt{z}}{-3xy}$

EXAMPLE 8 Evaluating Algebraic Expressions

Evaluate each expression for $m = -4$, $n = 5$, $p = -6$, and $q = 25$.

(a) $5m - 9n$

$$= 5(-4) - 9(5) \quad \text{Substitute } m = -4 \text{ and } n = 5.$$

$$= -20 - 45 \quad \text{Multiply.}$$

$$= -65 \quad \text{Subtract.}$$

Use parentheses around substituted values to avoid errors.

(b) $\frac{m + 2n}{4p}$

Work separately above and below the fraction bar.

$$= \frac{-4 + 2(5)}{4(-6)}$$

Substitute $m = -4$, $n = 5$, and $p = -6$.

$$= \frac{-4 + 10}{-24}$$

Multiply in the numerator.
Multiply in the denominator.

$$= \frac{6}{-24}$$

Add in the numerator.

$$= -\frac{1}{4}$$

Write in lowest terms, and rewrite using $\frac{a}{-b} = -\frac{a}{b}$.

(c) $-3m^3 - n^2(\sqrt{q})$

$$= -3(-4)^3 - (5)^2(\sqrt{25}) \quad \text{Substitute } m = -4, n = 5, \text{ and } q = 25.$$

$$= -3(-64) - 25(5) \quad \text{Evaluate the powers and the root.}$$

$$= 192 - 125 \quad \text{Multiply.}$$

$$= 67 \quad \text{Subtract.}$$

NOW TRY ANSWERS

8. (a) 29 (b) $\frac{5}{42}$


NOW TRY 

R.4 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Determine whether each statement is true or false. If it is false, correct the statement so that it is true.

1. $-7^6 = (-7)^6$

2. $-5^7 = (-5)^7$

3. $\sqrt{25}$ is a positive number. 4. $3 + 5 \cdot 8 = 3 + (5 \cdot 8)$
5. $(-6)^7$ is a negative number. 6. $(-6)^8$ is a positive number.
7. The product of 10 positive factors and 10 negative factors is positive.
8. The product of 5 positive factors and 5 negative factors is positive.
9. In the exponential expression -2^5 , the base is -2 .
10. \sqrt{a} is positive for all positive numbers a .

Write each expression using exponents. See Example 1.

11. $10 \cdot 10 \cdot 10 \cdot 10$ 12. $8 \cdot 8 \cdot 8$
13. $\frac{3}{4} \cdot \frac{3}{4} \cdot \frac{3}{4} \cdot \frac{3}{4} \cdot \frac{3}{4}$ 14. $\frac{1}{2} \cdot \frac{1}{2}$
15. $(-9)(-9)(-9)$ 16. $(-4)(-4)(-4)(-4)$
17. $(0.8)(0.8)$ 18. $(0.1)(0.1)(0.1)(0.1)(0.1)(0.1)$
19. $z \cdot z \cdot z \cdot z \cdot z \cdot z \cdot z$ 20. $a \cdot a \cdot a \cdot a \cdot a$

Concept Check Evaluate each exponential expression.

21. (a) 8^2 (b) -8^2 (c) $(-8)^2$ (d) $-(-8)^2$
22. (a) 4^3 (b) -4^3 (c) $(-4)^3$ (d) $-(-4)^3$

Evaluate each expression. See Examples 2 and 3.

23. 4^2 24. 6^2 25. $(0.3)^3$ 26. $(0.1)^3$
27. $\left(\frac{1}{5}\right)^3$ 28. $\left(\frac{1}{6}\right)^4$ 29. $\left(\frac{4}{5}\right)^4$ 30. $\left(\frac{7}{10}\right)^3$
31. $(-5)^3$ 32. $(-3)^5$ 33. $(-2)^8$ 34. $(-3)^6$
35. -3^6 36. -4^6 37. -8^4 38. -10^3

39. **Concept Check** Determine whether each expression is *positive* or *negative* when evaluated. Do not actually evaluate.

- (a) -7^2 (b) $(-7)^2$
- (c) -7^3 (d) $(-7)^3$
- (e) -7^4 (f) $(-7)^4$

40. **Concept Check** Match each square root in Column I with the appropriate value or description in Column II.

- | I | II |
|-------------------|----------------------|
| (a) $\sqrt{144}$ | A. -12 |
| (b) $\sqrt{-144}$ | B. 12 |
| (c) $-\sqrt{144}$ | C. Not a real number |

Find each square root. If it is not a real number, say so. See Example 4.

41. $\sqrt{81}$ 42. $\sqrt{64}$ 43. $\sqrt{169}$ 44. $\sqrt{225}$
45. $-\sqrt{400}$ 46. $-\sqrt{900}$ 47. $\sqrt{\frac{100}{121}}$ 48. $\sqrt{\frac{225}{169}}$
49. $-\sqrt{0.49}$ 50. $-\sqrt{0.64}$ 51. $\sqrt{-36}$ 52. $\sqrt{-121}$

53. **Concept Check** If a is a positive number, is $-\sqrt{-a}$ positive, negative, or not a real number?

54. **Concept Check** If a is a positive number, is $-\sqrt{a}$ positive, negative, or not a real number?

55. Concept Check Frank's grandson was asked to evaluate the following expression.

$$9 + 15 \div 3$$

Frank gave the answer as 8, but his grandson gave the answer as 14. The grandson explained that the answer is 14 because of the “Order of Process rule,” which says that when evaluating expressions, we proceed from right to left rather than left to right. (*Note:* This is a true story.)

- (a) Whose answer was correct for this expression, Frank's or his grandson's?
 (b) Was the *reasoning* for the correct answer valid? Explain.

56. Concept Check Problems like this one occasionally appear on social media. Simplify this expression.

$$7 + 7 \div 7 + 7 \times 7 - 7$$

Simplify each expression. See Examples 5–7.

57. $12 + 3 \cdot 4$

58. $15 + 5 \cdot 2$

59. $6 \cdot 3 - 12 \div 4$

60. $9 \cdot 4 - 8 \div 2$

61. $10 + 30 \div 2 \cdot 3$

62. $12 + 24 \div 3 \cdot 2$

63. $-3(5)^2 - (-2)(-8)$

64. $-9(2)^2 - (-3)(-2)$

65. $5 - 7 \cdot 3 - (-2)^3$

66. $-4 - 3 \cdot 5 - (-3)^3$

67. $-7(\sqrt{36}) - (-2)(-3)$

68. $-8(\sqrt{64}) - (-3)(-7)$

69. $6|4 - 5| - 24 \div 3$

70. $-4|2 - 4| + 8 \cdot 2$

71. $|-6 - 5|(-8) - 3^2$

72. $|-2 - 3|(-9) - 4^2$

73. $18 - 4^2 + 5 - (3 - 7)$

74. $10 - 2^2 + 9 - (1 - 8)$

75. $6 + \frac{2}{3}(-9) - \frac{5}{8} \cdot 16$

76. $7 - \frac{3}{4}(-8) + 12 \cdot \frac{5}{6}$

77. $-14\left(-\frac{2}{7}\right) \div (2 \cdot 6 - 10)$

78. $-12\left(-\frac{3}{4}\right) - (6 \cdot 5 \div 3)$

79. $\frac{(-5 + \sqrt{4}) - 2^2}{-5 - 2}$

80. $\frac{(-9 + \sqrt{16}) - 3^2}{-6 - 1}$

81. $\frac{2(-5) + (-3)(-2)}{-8 + 3^2 - 1}$

82. $\frac{3(-4) + (-5)(-8)}{2^3 - 2 - 6}$

Evaluate each expression for $a = -3$, $b = 64$, and $c = 6$. See Example 8.

83. $3a + \sqrt{b}$

84. $-2a - \sqrt{b}$

85. $\sqrt{b} + c - a$

86. $\sqrt{b} - c + a$

87. $4a^3 + 2c$

88. $-3a^4 - 3c$

89. $2(a - c)^2 - ac$

90. $-4ac + (c - a)^2$

91. $\frac{\sqrt{b} - 4a}{c^2}$

92. $\frac{a^3 + 2c}{-\sqrt{b}}$

93. $\frac{2c + a^3}{4b + 6a}$

94. $\frac{3c + a^2}{2b - 6c}$

Evaluate each expression for $w = 4$, $x = -\frac{3}{4}$, $y = \frac{1}{2}$, and $z = 1.25$. See Example 8.

95. $wy - 8x$

96. $wz - 12y$

97. $xy + y^4$

98. $xy - x^2$

99. $-w + 2x + 3y + z$

100. $w - 6x + 5y - 3z$

Residents of Linn County, Iowa, in the Cedar Rapids Community School District can use the expression

$$(v \times 0.5485 - 4850) \div 1000 \times 31.44$$

to determine their property taxes, where v is assessed home value. (Source: *The Gazette*.)

Use this expression to calculate, to the nearest dollar, the amount of property taxes that the owner of a home with each of the following values would pay. Follow the rules for order of operations.

101. \$150,000

102. \$200,000



The Blood Alcohol Concentration (BAC) of a person who has been drinking is given by the following expression.

$$\text{number of oz} \times \% \text{ alcohol} \times 0.075 \div \text{body weight in lb} - \text{hr of drinking} \times 0.015$$

(Source: Lawlor, J., *Auto Math Handbook: Calculations, Formulas, Equations and Theory for Automotive Enthusiasts*, Penguin, © 1991.)

103. Suppose a policeman stops a 190-lb man who, in 2 hr, has ingested four 12-oz beers (48 oz), each having a 3.2% alcohol content.

- (a) Substitute the values into the formula, and write the expression for the man's BAC.
- (b) Calculate the man's BAC to the nearest thousandth. Follow the rules for order of operations.

104. Find the BAC to the nearest thousandth for a 135-lb woman who, in 3 hr, has drunk three 12-oz beers (36 oz), each having a 4.0% alcohol content.

105. Refer to Exercises 103 and 104.

- (a) Calculate the BACs if each person weighed 25 lb more, and the rest of the variables stayed the same. How does increased weight affect a person's BAC?
- (b) Predict how decreased weight will affect the BAC of each person. Calculate the BACs if each person weighed 25 lb less and the rest of the variables stayed the same.

106. Calculate the BACs in Exercises 103 and 104 if the time for each person decreases by 1 hr and the other variable values stay the same. How does decreased time affect a person's BAC?

Solve each problem.

107. The amount, in billions of dollars, that Americans spent on their pets from 2003 to 2016 can be approximated by substituting a given year for x in the following expression.

$$2.493x - 4962$$

(Data from American Pet Products Manufacturers Association.) Find the amount spent in each year. Round answers to the nearest tenth.

(a) 2003 (b) 2010 (c) 2016

(d) How did the amount Americans spent on their pets change from 2003 to 2016?



Boots



108. The average price in dollars of a movie ticket in the United States from 2000 to 2016 can be approximated using the expression

$$0.2050x - 404.6,$$

where x represents the year. (Data from National Association of Theater Owners.)

- (a) Use the expression to complete the table.
 (b) How much did the average price of a movie ticket in the United States increase from 2000 to 2016?

Year	Average Price (in dollars)
2000	
2005	6.43
2010	
2016	

R.5


Properties of Real Numbers

OBJECTIVES

- 1 Use the distributive property.
- 2 Use the identity properties.
- 3 Use the inverse properties.
- 4 Use the commutative and associative properties.

VOCABULARY

- identity element for addition (additive identity)
- identity element for multiplication (multiplicative identity)
- term
- coefficient (numerical coefficient)
- like terms
- unlike terms



Area of left part is $2 \cdot 3 = 6$.
 Area of right part is $2 \cdot 5 = 10$.
 Area of total rectangle is $2(3 + 5) = 16$.

FIGURE 19

The basic properties of real numbers studied in this section, along with the multiplication property of zero presented earlier, reflect results that occur consistently in work with numbers. They have been generalized to apply to expressions with variables as well.

OBJECTIVE 1 Use the distributive property.

Notice the following.

$$2(3 + 5) = 2 \cdot 8 = 16$$

and $2 \cdot 3 + 2 \cdot 5 = 6 + 10 = 16,$

so $2(3 + 5) = 2 \cdot 3 + 2 \cdot 5.$

This idea is illustrated by the divided rectangle in **FIGURE 19**.

Similarly, observe these facts.

$$-4[5 + (-3)] = -4(2) = -8$$

and $-4(5) + (-4)(-3) = -20 + 12 = -8,$

so $-4[5 + (-3)] = -4(5) + (-4)(-3).$

These two examples are generalized to the set of *all* real numbers as the **distributive property of multiplication with respect to addition**, or simply the **distributive property**.

Distributive Property

For any real numbers a , b , and c , the following hold true.

$$a(b + c) = ab + ac \quad \text{and} \quad (b + c)a = ba + ca$$

Examples: $12(4 + 2) = 12 \cdot 4 + 12 \cdot 2$

$$(4 + 2)12 = 4 \cdot 12 + 2 \cdot 12$$

The distributive property can also be applied “in reverse.”

$$ab + ac = a(b + c) \quad \text{and} \quad ba + ca = (b + c)a$$

Examples: $6 \cdot 8 + 6 \cdot 9 = 6(8 + 9), \quad 8 \cdot 6 + 9 \cdot 6 = (8 + 9)6$

This property can be extended to more than two numbers as well.

$$a(b + c + d) = ab + ac + ad$$

The distributive property provides a way to rewrite a product $a(b + c)$ as a sum $ab + ac$, or a sum as a product.

NOTE When we use the distributive property to rewrite $a(b + c)$ as $ab + ac$, we refer to the process as “removing” or “clearing” parentheses.

NOW TRY EXERCISE 1

Use the distributive property to rewrite each expression.

- (a) $-2(3x - y)$
 (b) $4k - 12k$
 (c) $3a - 5b$

EXAMPLE 1 Using the Distributive Property

Use the distributive property to rewrite each expression.

(a) $3(x + y)$ Use the first form of the property to rewrite the given product as a sum.
 $= 3x + 3y$

(b) $-2(5 + k)$ Distributive property
 $= -2(5) + (-2)(k)$
 $= -10 - 2k$ Multiply.

(c) $4x + 8x$ Use the distributive property in reverse to rewrite the given sum as a product.
 $= (4 + 8)x$
 $= 12x$ Add inside the parentheses.

(d) $3r - 7r$ Definition of subtraction
 $= 3r + (-7r)$
 $= [3 + (-7)]r$ Distributive property in reverse
 $= -4r$ Add.

(e) $5p + 7q$ This expression *cannot* be rewritten as $12pq$.

Because there is no common number or variable here, we cannot use the distributive property to rewrite the expression.

(f) $6(x + 2y - 3z)$ Distributive property
 $= 6x + 6(2y) + 6(-3z)$
 $= 6x + 12y - 18z$ Multiply.

NOW TRY 

The distributive property can also be used for subtraction as in **Example 1(d)**.

$$a(b - c) = ab - ac \quad \text{Example: } 6(x - y) = 6x - 6y$$

OBJECTIVE 2 Use the identity properties.

The number 0 is the only number that can be added to any number to get that number, leaving the identity of the number unchanged. Thus, 0 is the **identity element for addition**, or the **additive identity**.

In a similar way, multiplying any number by 1 leaves the identity of the number unchanged. Thus, 1 is the **identity element for multiplication**, or the **multiplicative identity**. The **identity properties** summarize this discussion.

NOW TRY ANSWERS

1. (a) $-6x + 2y$ (b) $-8k$
 (c) cannot be rewritten



NOW TRY EXERCISE 2

Simplify each expression.

(a) $7x + x$ (b) $-(5p - 3q)$

Identity Properties

For any real number a , the following hold true.

$$a + 0 = a \quad \text{and} \quad 0 + a = a$$

Examples: $9 + 0 = 9$, $0 + 9 = 9$

$$a \cdot 1 = a \quad \text{and} \quad 1 \cdot a = a$$

Examples: $9 \cdot 1 = 9$, $1 \cdot 9 = 9$

The identity properties leave the identity of a real number unchanged. Think of a child wearing a costume on Halloween. The child's appearance is changed, but his or her identity is unchanged.

EXAMPLE 2 Using the Identity Property $1 \cdot a = a$

Simplify each expression.

(a) $12m + m$

$$= 12m + 1m \quad \text{Identity property}$$

$$= (12 + 1)m \quad \text{Distributive property}$$

$$= 13m \quad \text{Add inside the parentheses.}$$

(b) $y + y$

$$= 1y + 1y$$

$$= (1 + 1)y$$

$$= 2y$$

(c) $-(m - 5n)$

$$= -1(m - 5n) \quad \text{Identity property}$$

$$= -1(m) + (-1)(-5n) \quad \text{Distributive property}$$

$$= -m + 5n \quad \text{Multiply.}$$

Multiply each term by -1 .
Be careful with signs.

NOW TRY

OBJECTIVE 3 Use the inverse properties.

The **additive inverse** (or **opposite**) of a number a is $-a$. Additive inverses have a sum of 0 (the additive identity).

$$5 \text{ and } -5, \quad -\frac{1}{2} \text{ and } \frac{1}{2}, \quad -34 \text{ and } 34 \quad \text{Additive inverses (sum of 0)}$$

The **multiplicative inverse** (or **reciprocal**) of a number a is $\frac{1}{a}$ (where $a \neq 0$). Multiplicative inverses have a product of 1.

$$5 \text{ and } \frac{1}{5}, \quad -\frac{1}{2} \text{ and } -2, \quad \frac{3}{4} \text{ and } \frac{4}{3} \quad \text{Multiplicative inverses (product of 1)}$$

This discussion leads to the **inverse properties**.

Inverse Properties

For any real number a , the following hold true.

$$a + (-a) = 0 \quad \text{and} \quad -a + a = 0$$

Examples: $7 + (-7) = 0$, $-7 + 7 = 0$

$$a \cdot \frac{1}{a} = 1 \quad \text{and} \quad \frac{1}{a} \cdot a = 1 \quad (\text{where } a \neq 0)$$

Examples: $7 \cdot \frac{1}{7} = 1$, $\frac{1}{7} \cdot 7 = 1$

NOW TRY ANSWERS

2. (a) $8x$ (b) $-5p + 3q$

▼ Terms and Their Coefficients

Term	Numerical Coefficient
$-7y$	-7
$34r^3$	34
$-26x^5yz^4$	-26
$-k = -1k$	-1
$r = 1r$	1
$\frac{3x}{8} = \frac{3}{8}x$	$\frac{3}{8}$
$\frac{x}{3} = \frac{1x}{3} = \frac{1}{3}x$	$\frac{1}{3}$

The inverse properties “undo” addition or multiplication. Putting on your shoes when you get up in the morning and then taking them off before you go to bed at night are inverse operations that undo each other.

Expressions such as $12m$ and $5n$ from **Example 2** are examples of *terms*. A **term** is a number or the product of a number and one or more variables raised to powers. The numerical factor in a term is the **numerical coefficient**, or just the **coefficient**.

Terms with exactly the same variables raised to exactly the same powers are **like terms**. Otherwise, they are **unlike terms**.

$$\begin{array}{lll}
 5p \text{ and } -21p & -6x^2 \text{ and } 9x^2 & \text{Like terms} \\
 3m \text{ and } 16x & 7y^3 \text{ and } -3y^2 & \text{Unlike terms} \\
 \uparrow \quad \uparrow & \uparrow \quad \uparrow & \\
 \text{Different variables} & \text{Different exponents} & \\
 & \text{on the same variable} &
 \end{array}$$

OBJECTIVE 4 Use the commutative and associative properties.

Simplifying expressions as in **Examples 2(a) and (b)** is called **combining like terms**. *Only like terms may be combined.* To combine like terms in an expression such as

$$-2m + 5m + 3 - 6m + 8,$$

we need two more properties. From arithmetic, we know that the following are true.

$$\begin{array}{ll}
 3 + 9 = 12 & \text{and} \quad 9 + 3 = 12 \\
 3 \cdot 9 = 27 & \text{and} \quad 9 \cdot 3 = 27
 \end{array}$$

The order of the numbers being added or multiplied does not matter. The same answers result. The following computations are also true.

$$\begin{array}{l}
 (5 + 7) + 2 = 12 + 2 = 14 \\
 5 + (7 + 2) = 5 + 9 = 14 \\
 (5 \cdot 7) \cdot 2 = 35 \cdot 2 = 70 \\
 5 \cdot (7 \cdot 2) = 5 \cdot 14 = 70
 \end{array}$$

The grouping of the numbers being added or multiplied does not matter. The same answers result.

These arithmetic examples can be extended to algebra.

Commutative and Associative Properties

For any real numbers a , b , and c , the following hold true.

$$\left. \begin{array}{l}
 a + b = b + a \\
 ab = ba
 \end{array} \right\} \text{Commutative properties}$$

(The *order* of the two terms or factors changes.)

Examples: $9 + (-3) = -3 + 9$, $9(-3) = (-3)9$

$$\left. \begin{array}{l}
 a + (b + c) = (a + b) + c \\
 a(bc) = (ab)c
 \end{array} \right\} \text{Associative properties}$$

(The *grouping* among the three terms or factors changes, but the order stays the same.)

Examples: $7 + (8 + 9) = (7 + 8) + 9$, $7 \cdot (8 \cdot 9) = (7 \cdot 8) \cdot 9$

The commutative properties are used to change the order of the terms or factors in an expression. Think of *commuting* from home to work and then from work to home. The associative properties are used to regroup the terms or factors of an expression. Think of *associating* the grouped terms or factors.

 **NOW TRY
EXERCISE 3**

Simplify.

$$-7x + 10 - 3x - 4 + x$$

EXAMPLE 3 Using the Commutative and Associative Properties

Simplify.

$$\begin{aligned} & -2m + 5m + 3 - 6m + 8 \\ &= (-2m + 5m) + 3 - 6m + 8 && \text{Associative property} \\ &= (-2 + 5)m + 3 - 6m + 8 && \text{Distributive property} \\ &= 3m + 3 - 6m + 8 && \text{Add inside parentheses.} \end{aligned}$$

The next step would be to add $3m$ and 3 , but they are unlike terms. To combine $3m$ and $-6m$, we use the associative and commutative properties, inserting parentheses and brackets according to the rules for order of operations.

$$\begin{aligned} &= [3m + (3 - 6m)] + 8 && \text{Associative property} \\ &= [3m + (-6m + 3)] + 8 && \text{Commutative property} \\ &= [(3m + [-6m]) + 3] + 8 && \text{Associative property} \\ &= (-3m + 3) + 8 && \text{Combine like terms.} \\ &= -3m + (3 + 8) && \text{Associative property} \\ &= -3m + 11 && \text{Add.} \end{aligned}$$

In practice, many of these steps are not written down, but it is important to realize that the commutative and associative properties are used whenever the terms in an expression are rearranged and regrouped to combine like terms.

NOW TRY 

EXAMPLE 4 Using the Properties of Real Numbers

Simplify each expression.

(a) $5y - 8y - 6y + 11y$

$$\begin{aligned} &= (5 - 8 - 6 + 11)y && \text{Distributive property} \\ &= 2y && \text{Combine like terms.} \end{aligned}$$

(b) $3x + 4 - 5(x + 1) - 8$

$$\begin{aligned} &= 3x + 4 - 5x - 5 - 8 && \text{Distributive property} \\ &= 3x - 5x + 4 - 5 - 8 && \text{Commutative property} \\ &= -2x - 9 && \text{Combine like terms.} \end{aligned}$$

Be careful with signs.

(c) $8 - (3m + 2)$

$$\begin{aligned} &= 8 - 1(3m + 2) && \text{Identity property} \\ &= 8 - 3m - 2 && \text{Distributive property} \\ &= 6 - 3m && \text{Combine like terms.} \end{aligned}$$

NOW TRY ANSWER

3. $-9x + 6$


NOW TRY EXERCISE 4

Simplify each expression.

- (a) $-3(t - 4) - t + 15$
 (b) $7x - (4x - 2)$
 (c) $5x(6y)$
 (d) $3(5x - 7) - 8(x + 4)$

$$\begin{aligned}
 \text{(d)} \quad & 3x(5)(y) \\
 &= [3x(5)]y && \text{Order of operations} \\
 &= [3(x \cdot 5)]y && \text{Associative property} \\
 &= [3(5x)]y && \text{Commutative property} \\
 &= [(3 \cdot 5)x]y && \text{Associative property} \\
 &= (15x)y && \text{Multiply.} \\
 &= 15(xy) && \text{Associative property} \\
 &= 15xy
 \end{aligned}$$

As previously mentioned, many of these steps are not usually written out.

$$\begin{aligned}
 \text{(e)} \quad & 4(3x - 5) - 2(4x + 7) \\
 &= 12x - 20 - 8x - 14 && \text{Distributive property} \\
 &= 12x - 8x - 20 - 14 && \text{Commutative property} \\
 &= 4x - 34 && \text{Combine like terms.}
 \end{aligned}$$

Like terms may be combined by adding or subtracting the coefficients of the terms and keeping the same variable factors.

NOW TRY 

CAUTION Be careful. The distributive property does not apply in **Example 4(d)** because there is no addition or subtraction involved.

$$(3x)(5)(y) \neq (3x)(5) \cdot (3x)(y)$$


NOW TRY ANSWERS

4. (a) $-4t + 27$ (b) $3x + 2$
 (c) $30xy$ (d) $7x - 53$

R.5 Exercises

**FOR
EXTRA
HELP**

MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Choose the correct response.

- | | |
|---------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| 1. The identity element for addition is
A. $-a$ B. 0 C. 1 D. $\frac{1}{a}$. | 2. The identity element for multiplication is
A. $-a$ B. 0 C. 1 D. $\frac{1}{a}$. |
| 3. The additive inverse of a is
A. $-a$ B. 0 C. 1 D. $\frac{1}{a}$. | 4. The multiplicative inverse of a , where $a \neq 0$, is
A. $-a$ B. 0 C. 1 D. $\frac{1}{a}$. |

Concept Check Complete each statement.

- The distributive property provides a way to rewrite a product such as $a(b + c)$ as the sum _____.
- The commutative property is used to change the _____ of two terms or factors.
- The associative property is used to change the _____ of three terms or factors.
- Like terms are terms with the _____ variables raised to the _____ powers.
- When simplifying an expression, only _____ terms can be combined.
- The numerical coefficient in the term $-7yz^2$ is _____.

Simplify each expression. See Examples 1 and 2.

- | | | | |
|-----------------|-------------------|----------------------|----------------------|
| 11. $2(m + p)$ | 12. $3(a + b)$ | 13. $-12(x - y)$ | 14. $-10(p - q)$ |
| 15. $5k + 3k$ | 16. $6a + 5a$ | 17. $7r - 9r$ | 18. $4n - 6n$ |
| 19. $-8z + 4w$ | 20. $-12k + 3r$ | 21. $a + 7a$ | 22. $s + 9s$ |
| 23. $x + x$ | 24. $a + a$ | 25. $-(2d - f)$ | 26. $-(3m - n)$ |
| 27. $-(-x - y)$ | 28. $-(-3x - 4y)$ | 29. $2(x - 3y + 2z)$ | 30. $8(3x + y - 5z)$ |

Simplify each expression. See Examples 1–4.

- | | |
|-------------------------------------|--------------------------------------|
| 31. $-12y + 4y + 3y + 2y$ | 32. $-5r - 9r + 8r - 5r$ |
| 33. $-6p + 5 - 4p + 6 + 11p$ | 34. $-8x - 12 + 3x - 5x + 9$ |
| 35. $3(k + 2) - 5k + 6 + 3$ | 36. $5(r - 3) + 6r - 2r + 4$ |
| 37. $10 - (4y + 8)$ | 38. $6 - (9y + 5)$ |
| 39. $10x(3)(y)$ | 40. $8x(6)(y)$ |
| 41. $-\frac{2}{3}(12w)(7z)$ | 42. $-\frac{5}{6}(18w)(5z)$ |
| 43. $3(m - 4) - 2(m + 1)$ | 44. $6(a - 5) - 4(a + 6)$ |
| 45. $0.25(8 + 4p) - 0.5(6 + 2p)$ | 46. $0.4(10 - 5x) - 0.8(5 + 10x)$ |
| 47. $-(2p + 5) + 3(2p + 4) - 2p$ | 48. $-(7m - 12) + 2(4m + 7) - 6m$ |
| 49. $2 + 3(2z - 5) - 3(4z + 6) - 8$ | 50. $-4 + 4(4k - 3) - 6(2k + 8) + 7$ |

Complete each statement so that the indicated property is illustrated. Simplify each answer if possible. See Examples 1–4.

- | | |
|--------------------------------------------------------------------|---------------------------------------------------------------------------|
| 51. $5x + 8x =$ _____
(distributive property) | 52. $9y - 6y =$ _____
(distributive property) |
| 53. $5(9r) =$ _____
(associative property) | 54. $-4 + (12 + 8) =$ _____
(associative property) |
| 55. $5x + 9y =$ _____
(commutative property) | 56. $-5 \cdot 7 =$ _____
(commutative property) |
| 57. $1 \cdot 7 =$ _____
(identity property) | 58. $-12x + 0 =$ _____
(identity property) |
| 59. $-\frac{1}{4}ty + \frac{1}{4}ty =$ _____
(inverse property) | 60. $-\frac{9}{8}\left(-\frac{8}{9}\right) =$ _____
(inverse property) |
| 61. $8(-4 + x) =$ _____
(distributive property) | 62. $3(x - y + z) =$ _____
(distributive property) |
| 63. $0(0.875x + 9y) =$ _____
(multiplication property of 0) | 64. $0(0.35t + 12u) =$ _____
(multiplication property of 0) |

65. **Concept Check** Give an “everyday” example of a commutative operation.

66. **Concept Check** Give an “everyday” example of inverse operations.

The distributive property can be used to mentally perform calculations.

$$\begin{aligned} & 38 \cdot 17 + 38 \cdot 3 \\ &= 38(17 + 3) && \text{Distributive property} \\ &= 38(20) && \text{Add inside the parentheses.} \\ &= 760 && \text{Multiply.} \end{aligned}$$

Use the distributive property to calculate each value mentally.

$$\begin{array}{lll} 67. 96 \cdot 19 + 4 \cdot 19 & 68. 27 \cdot 60 + 27 \cdot 40 & 69. 58 \cdot \frac{3}{2} - 8 \cdot \frac{3}{2} \\ 70. \frac{8}{5}(17) + \frac{8}{5}(13) & 71. 8.75(15) - 8.75(5) & 72. 4.31(69) + 4.31(31) \end{array}$$

RELATING CONCEPTS For Individual or Group Work (Exercises 73–78)

When simplifying an expression, we usually do some steps mentally. **Work Exercises 73–78 in order**, providing the property that justifies each statement in the given simplification. (These steps could be done in other orders.)

$$\begin{array}{ll} & 3x + 4 + 2x + 7 \\ 73. & = (3x + 4) + (2x + 7) \quad \underline{\hspace{2cm}} \\ 74. & = 3x + (4 + 2x) + 7 \quad \underline{\hspace{2cm}} \\ 75. & = 3x + (2x + 4) + 7 \quad \underline{\hspace{2cm}} \\ 76. & = (3x + 2x) + (4 + 7) \quad \underline{\hspace{2cm}} \\ 77. & = (3 + 2)x + (4 + 7) \quad \underline{\hspace{2cm}} \\ 78. & = 5x + 11 \quad \underline{\hspace{2cm}} \end{array}$$

Chapter R Summary

Key Terms

R.1

fractions
numerator
denominator
proper fraction
improper fraction
lowest terms
mixed number
reciprocals
decimal
terminating decimal
repeating decimal
percent

R.2

set
elements (members)
finite set

natural (counting) numbers
infinite set
whole numbers
empty (null) set
variable
number line
integers
coordinate
graph
rational numbers
irrational numbers
real numbers
additive inverse
(opposite, negative)
signed numbers
absolute value
equation
inequality

R.3

sum
difference
product
quotient
reciprocal
(multiplicative inverse)
dividend
divisor

R.4

factors
exponent (power)
base
exponential expression
square root
positive (principal)
square root

negative square root
constant
algebraic expression

R.5

identity element for
addition
(additive identity)
identity element for
multiplication
(multiplicative
identity)
term
coefficient
(numerical coefficient)
like terms
unlike terms

New Symbols

$0.\overline{6}$	bar notation that signifies repeating digit(s)	\notin	is not an element of	$>$	is greater than
$\%$	percent	\neq	is not equal to	\geq	is greater than or equal to
$\{a, b\}$	set containing the elements a and b	$\{x \mid x \text{ has property } P\}$	set-builder notation	a^m	m factors of a
\emptyset	empty set	$ x $	absolute value of x	$\sqrt{\quad}$	radical symbol
\in	is an element of (a set)	$<$	is less than	\sqrt{a}	positive (principal) square root of a
		\leq	is less than or equal to		

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- The **denominator** of a fraction
 - is the number above the fraction bar
 - gives the total number of equal parts in the whole
 - gives the number of shaded parts in the whole
 - is the smaller number in the fraction.
- A **proper fraction** is a fraction that has
 - numerator greater than denominator
 - numerator equal to denominator
 - numerator less than denominator
 - denominator less than numerator.
- The **empty set** is a set
 - with 0 as its only element
 - with an infinite number of elements
 - with no elements
 - of ideas.
- A **variable** is
 - a symbol used to represent an unknown number
 - a value that makes an equation true
 - a solution of an equation
 - the answer in a division problem.
- An **integer** is
 - a positive or negative number
 - a natural number, its opposite, or zero
 - any number that can be graphed
 - the quotient of two numbers.
- The **absolute value** of a number is
 - the graph of the number
 - the reciprocal of the number
 - the opposite of the number
 - the distance between 0 and the number on a number line.
- The **reciprocal** of a nonzero number a is
 - a
 - $\frac{1}{a}$
 - $-a$
 - 1.
- A **factor** is
 - the answer in an addition problem
 - the answer in a multiplication problem
 - one of two or more numbers that are added to get another number
 - any number that divides evenly into a given number.
- An **exponent** is
 - a symbol that tells how many numbers are being multiplied
 - a number raised to a power
 - a number that tells how many times a factor is repeated
 - a number that is multiplied.
- An **exponential expression** is
 - a number that is a repeated factor in a product
 - a number or a variable written with an exponent
 - a number that tells how many times a factor is repeated in a product
 - an expression that involves addition.
- A **term** is
 - a numerical factor
 - a number or a product of a number and one or more variables raised to powers
 - one of several variables with the same exponents
 - a sum of numbers and variables raised to powers.
- A **numerical coefficient** is
 - the numerical factor in a term
 - the number of terms in an expression
 - a variable raised to a power
 - the variable factor in a term.

ANSWERS

1. B; *Example:* In the fraction $\frac{3}{4}$, the denominator is 4. 2. C; *Examples:* $\frac{1}{2}, \frac{2}{7}, \frac{5}{12}$ 3. C; *Example:* The set of whole numbers less than 0 is the empty set, written \emptyset . 4. A; *Examples:* x, y, z 5. B; *Examples:* $-9, 0, 6$ 6. D; *Examples:* $|2| = 2$ and $|-2| = 2$ 7. B; *Examples:* 3 is the reciprocal of $\frac{1}{3}$; $-\frac{5}{2}$ is the reciprocal of $-\frac{2}{5}$. 8. D; *Example:* 2 and 5 are factors of 10 because both divide evenly (without remainder) into 10. 9. C; *Example:* In 2^3 , the number 3 is the exponent (or power), so 2 is a factor three times, and $2^3 = 2 \cdot 2 \cdot 2 = 8$. 10. B; *Examples:* 3^4 and x^{10} 11. B; *Examples:* $6, \frac{x}{2}, -4ab^2$ 12. A; *Examples:* The term $8z$ has numerical coefficient 8, and the term $-10x^3y$ has numerical coefficient -10 .

Chapter R

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

View the complete solutions to all Chapter Test exercises in MyLab Math.

Perform the indicated operations.

1. $\frac{3}{4} + \frac{1}{6}$

2. $\frac{3}{7} \div \frac{9}{14}$

3. $13.25 - 6.417$

4. 0.7×0.04

Complete the table of fraction, decimal, and percent equivalents.

	Fraction in Lowest Terms	Decimal	Percent
5.			4%
6.	$\frac{5}{6}$		
7.		1.5	

8. Graph $\left\{-3, 0.75, \frac{5}{3}, 5, 6.3\right\}$ on a number line.

Let $A = \{-\sqrt{6}, -1, -0.5, 0, 3, \sqrt{25}, 7.5, \frac{24}{2}, \sqrt{-4}\}$. Simplify the elements of A as necessary, and then list those elements of A that belong to the specified set.

9. Whole numbers

10. Integers

11. Rational numbers

12. Real numbers

Perform the indicated operations.

13. $-6 + 14 + (-11) - (-3)$

14. $-\frac{5}{7} - \left(-\frac{10}{9} + \frac{2}{3}\right)$

15. $10 - 4 \cdot 3 + 6(-4)$

16. $7 - 4^2 + 2(6) + (-4)^2$

17. $\frac{-2[3 - (-1 - 2) + 2]}{\sqrt{9}(-3) - (-2)}$

18. $\frac{8 \cdot 4 - 3^2 \cdot 5 - 2(-1)}{-3 \cdot 2^3 + 24}$

Find each square root. If it is not a real number, say so.

19. $\sqrt{196}$

20. $-\sqrt{225}$

21. $\sqrt{-16}$

22. Evaluate $\frac{8k + 2m^2}{r - 2}$ for $k = -3$, $m = -3$, and $r = 25$.

Simplify each expression.

23. $-3(2k - 4) + 4(3k - 5) - 2 + 4k$

24. $(3r + 8) - (-4r + 6)$

25. Match each statement in Column I with the appropriate property in Column II. Answers may be used more than once.

I

(a) $6 + (-6) = 0$

(b) $-2 + (3 + 6) = (-2 + 3) + 6$

(c) $5x + 15x = (5 + 15)x$

(d) $13 \cdot 0 = 0$

(e) $-9 + 0 = -9$

(f) $4 \cdot 1 = 4$

(g) $(a + b) + c = (b + a) + c$

II

A. Distributive property

B. Inverse property

C. Identity property

D. Associative property

E. Commutative property

F. Multiplication property of 0

STUDY SKILLS REMINDER

You will increase your chance of success in this course if you fully utilize your text. **Review Study Skill 1, Using Your Math Text.**



LINEAR EQUATIONS, INEQUALITIES, AND APPLICATIONS

The concept of balance can be applied to solving *linear equations in one variable*, a topic of this chapter.

1.1 Linear Equations in One Variable

1.2 Formulas and Percent

1.3 Applications of Linear Equations

1.4 Further Applications of Linear Equations

SUMMARY EXERCISES Applying Problem-Solving Techniques

1.5 Linear Inequalities in One Variable

1.6 Set Operations and Compound Inequalities

1.7 Absolute Value Equations and Inequalities

SUMMARY EXERCISES Solving Linear and Absolute Value Equations and Inequalities

1.1 Linear Equations in One Variable

OBJECTIVES

- 1 Distinguish between expressions and equations.
- 2 Identify linear equations.
- 3 Solve linear equations using the addition and multiplication properties of equality.
- 4 Solve linear equations using the distributive property.
- 5 Solve linear equations with fractions or decimals.
- 6 Identify conditional equations, contradictions, and identities.

VOCABULARY

- linear (first-degree) equation in one variable
- solution
- solution set
- equivalent equations
- conditional equation
- identity
- contradiction

NOW TRY EXERCISE 1

Determine whether each of the following is an *expression* or an *equation*.

- (a) $2x + 17 - 3x$
 (b) $2x + 17 = 3x$

NOW TRY ANSWERS

1. (a) expression (b) equation

OBJECTIVE 1 Distinguish between expressions and equations.

In our earlier work, we reviewed *algebraic expressions*.

$$8x + 9, \quad y - 4, \quad \frac{x^3y^8}{z} \quad \text{Algebraic expressions}$$

Equations and inequalities compare algebraic expressions, just as a balance scale compares the weights of two quantities. Recall that an **equation** is a statement that two algebraic expressions are equal. **An equation always contains an equality symbol, while an expression does not.**

EXAMPLE 1 Distinguishing between Expressions and Equations

Determine whether each of the following is an *expression* or an *equation*.

(a) $3x - 7 = 2$

(b) $3x - 7$

In part (a) we have an equation because there is an equality symbol. In part (b), there is no equality symbol, so it is an expression.

Equation

$$\underbrace{3x - 7}_{\text{Left side}} = \underbrace{2}_{\text{Right side}}$$

An equation can be solved.

Expression

$$3x - 7$$

An expression cannot be solved.
It can often be evaluated or simplified.

NOW TRY

OBJECTIVE 2 Identify linear equations.

A **linear equation in one variable** involves only real numbers and one variable raised to the first power.

Linear Equation in One Variable

A **linear equation in one variable** (here x) is an equation that can be written in the form

$$ax + b = 0,$$

where a and b are real numbers and $a \neq 0$.

Examples: $x + 1 = -2, \quad x - 3 = 5, \quad 2x + 5 = 10$

Linear equations
in one variable

A linear equation is a **first-degree equation** because the greatest power on the variable is 1. Some equations that are not linear equations in one variable follow.

$$\left. \begin{array}{l} 4x - 5y = 6 \quad (\text{There is more than one variable.}) \\ x^2 + 4 = 0 \quad (\text{The exponent on the variable is not 1.}) \end{array} \right\} \text{Not linear equations} \\ \text{in one variable}$$

If the variable in an equation can be replaced by a real number that makes the statement true, then that number is a **solution** of the equation. For example, 8 is a solution of the equation $x - 3 = 5$ because replacing x with 8 gives a true statement.

An equation is *solved* by finding its **solution set**, the set of all solutions. The solution set of the equation $x - 3 = 5$ is $\{8\}$.

Equivalent equations are related equations that have the same solution set. To solve an equation, we usually start with the given equation and replace it with a series of simpler equivalent equations. The following are all equivalent because each has solution set $\{3\}$.

$$5x + 2 = 17, \quad 5x = 15, \quad x = 3 \quad \text{Equivalent equations}$$

OBJECTIVE 3 Solve linear equations using the addition and multiplication properties of equality.

We use two important properties of equality to produce equivalent equations.

Addition and Multiplication Properties of Equality

Addition Property of Equality

If a , b , and c represent real numbers, then the equations

$$a = b \quad \text{and} \quad a + c = b + c \quad \text{are equivalent.}$$

That is, the same number may be added to each side of an equation without changing the solution set.

Multiplication Property of Equality

If a , b , and c represent real numbers and $c \neq 0$, then the equations

$$a = b \quad \text{and} \quad ac = bc \quad \text{are equivalent.}$$

That is, each side of an equation may be multiplied by the same nonzero number without changing the solution set.

Because subtraction and division are defined in terms of addition and multiplication, respectively, the preceding properties can be extended.

The same number may be subtracted from each side of an equation, and each side of an equation may be divided by the same nonzero number, without changing the solution set.

EXAMPLE 2 Solving a Linear Equation

Solve $4x - 2x - 5 = 4 + 6x + 3$.

The goal is to isolate x on one side of the equation.

$$4x - 2x - 5 = 4 + 6x + 3$$

$$2x - 5 = 7 + 6x \quad \text{Combine like terms.}$$

$$2x - 5 - 6x = 7 + 6x - 6x \quad \text{Subtract } 6x \text{ from each side.}$$

$$-4x - 5 = 7 \quad \text{Combine like terms.}$$

$$-4x - 5 + 5 = 7 + 5 \quad \text{Add 5 to each side.}$$

$$-4x = 12 \quad \text{Combine like terms.}$$

$$\frac{-4x}{-4} = \frac{12}{-4} \quad \text{Divide each side by } -4.$$

$$x = -3$$


NOW TRY EXERCISE 2

Solve.

$$5x + 11 = 2x - 13 - 3x$$

CHECK Substitute -3 for x in the original equation.

$$4x - 2x - 5 = 4 + 6x + 3 \quad \text{Original equation}$$

$$4(-3) - 2(-3) - 5 \stackrel{?}{=} 4 + 6(-3) + 3 \quad \text{Let } x = -3.$$

$$-12 + 6 - 5 \stackrel{?}{=} 4 - 18 + 3 \quad \text{Multiply.}$$

$$-11 = -11 \quad \checkmark \quad \text{True}$$

Use parentheses around substituted values to avoid errors.

This is *not* the solution.

The true statement indicates that $\{-3\}$ is the solution set.**NOW TRY** 

CAUTION In Example 2, the equality symbols are aligned in a column. *Use only one equality symbol in a horizontal line of work when solving an equation.*

Solving a Linear Equation in One Variable

Step 1 Simplify each side separately. Use the distributive property as needed.

- Clear any parentheses.
- Clear any fractions or decimals.
- Combine like terms.

Step 2 Isolate the variable terms on one side. Use the addition property of equality so that all terms with variables are on one side of the equation and all constants (numbers) are on the other side.

Step 3 Isolate the variable. Use the multiplication property of equality to obtain an equation that has just the variable with coefficient 1 on one side.

Step 4 Check. Substitute the value found into the *original* equation. If a true statement results, write the solution set. If not, rework the problem.

OBJECTIVE 4 Solve linear equations using the distributive property.

EXAMPLE 3 Solving a Linear Equation

Solve $2(x - 5) + 3x = x + 6$.

Step 1 Clear parentheses using the distributive property. Then combine like terms.

Be sure to distribute over *all* terms within the parentheses.

$$2(x - 5) + 3x = x + 6$$

$$2x + 2(-5) + 3x = x + 6 \quad \text{Distributive property}$$

$$2x - 10 + 3x = x + 6 \quad \text{Multiply.}$$

$$5x - 10 = x + 6 \quad \text{Combine like terms.}$$

Step 2 Isolate the variable terms on one side and constants on the other.

$$5x - 10 - x = x + 6 - x \quad \text{Subtract } x.$$

$$4x - 10 = 6 \quad \text{Combine like terms.}$$

$$4x - 10 + 10 = 6 + 10 \quad \text{Add 10.}$$

$$4x = 16 \quad \text{Combine like terms.}$$

NOW TRY ANSWER2. $\{-4\}$


**NOW TRY
EXERCISE 3**

Solve.

$$5(x - 4) - 12 = 3 - 2x$$

Step 3 Use the multiplication property of equality to isolate x on the left.

$$\frac{4x}{4} = \frac{16}{4} \quad \text{Divide by 4.}$$

$$x = 4$$

Step 4 Check by substituting 4 for x in the original equation.

CHECK $2(x - 5) + 3x = x + 6$ Original equation

$2(4 - 5) + 3(4) \stackrel{?}{=} 4 + 6$ Let $x = 4$.

$2(-1) + 12 \stackrel{?}{=} 10$ Simplify.

$10 = 10$ ✓ True

Always check
your work.

A true statement results, so the solution set is $\{4\}$.

NOW TRY 


**NOW TRY
EXERCISE 4**

Solve.

$$2 - 3(2 + 6x) = 4(x + 1) + 36$$

EXAMPLE 4 Solving a Linear Equation

Solve $4(3x - 2) = 38 - 2(2x - 1)$.

$$4(3x - 2) = 38 - 2(2x - 1)$$

Step 1 $4(3x) + 4(-2) = 38 - 2(2x) - 2(-1)$ Clear parentheses using the distributive property.

Be careful with signs
when distributing.

$$12x - 8 = 38 - 4x + 2$$

Multiply.

$$12x - 8 = 40 - 4x$$

Combine like terms.

Step 2 $12x - 8 + 4x = 40 - 4x + 4x$ Add $4x$.

$$16x - 8 = 40$$

Combine like terms.

$$16x - 8 + 8 = 40 + 8$$

Add 8.

$$16x = 48$$

Combine like terms.

Step 3 $\frac{16x}{16} = \frac{48}{16}$ Divide by 16.

$$x = 3$$

Step 4 **CHECK** $4(3x - 2) = 38 - 2(2x - 1)$ Original equation

$4[3(3) - 2] \stackrel{?}{=} 38 - 2[2(3) - 1]$ Let $x = 3$.

$4[7] \stackrel{?}{=} 38 - 2[5]$ Work inside the brackets.

$28 = 28$ ✓ True

A true statement results, so the solution set is $\{3\}$.

NOW TRY 

OBJECTIVE 5 Solve linear equations with fractions or decimals.

When fractions appear as coefficients in an equation, we multiply each side by the least common denominator (LCD) of all the fractions. This is an application of the multiplication property of equality.

NOW TRY ANSWERS

3. $\{5\}$

4. $\{-2\}$


**NOW TRY
EXERCISE 5**

Solve.

$$\frac{x-4}{4} + \frac{2x+4}{8} = 5$$

EXAMPLE 5 Solving a Linear Equation with Fractions

Solve $\frac{x+7}{6} + \frac{2x-8}{2} = -4$.

$$\frac{x+7}{6} + \frac{2x-8}{2} = -4$$

$$\text{Step 1} \quad 6\left(\frac{x+7}{6} + \frac{2x-8}{2}\right) = 6(-4)$$

Eliminate the fractions.
Multiply each side by the LCD, 6.

$$6\left(\frac{x+7}{6}\right) + 6\left(\frac{2x-8}{2}\right) = 6(-4)$$

Distributive property

$$(x+7) + 3(2x-8) = -24$$

Multiply.

$$x+7 + 3(2x) + 3(-8) = -24$$

Distributive property

$$x+7 + 6x - 24 = -24$$

Multiply.

$$7x - 17 = -24$$

Combine like terms.

$$\text{Step 2} \quad 7x - 17 + 17 = -24 + 17$$

Add 17.

$$7x = -7$$

Combine like terms.

$$\text{Step 3} \quad \frac{7x}{7} = \frac{-7}{7}$$

Divide by 7.

$$x = -1$$

$$\text{Step 4 CHECK} \quad \frac{x+7}{6} + \frac{2x-8}{2} = -4$$

Original equation

$$\frac{-1+7}{6} + \frac{2(-1)-8}{2} \stackrel{?}{=} -4$$

Let $x = -1$.

$$1 - 5 \stackrel{?}{=} -4$$

Simplify each fraction.

$$-4 = -4 \quad \checkmark$$

True

A true statement results, so the solution set is $\{-1\}$.
NOW TRY 

When an equation involves decimal coefficients, we can clear the decimals by multiplying by a power of 10, such as $10^1 = 10$, $10^2 = 100$, and so on, to obtain an equivalent equation with integer coefficients.

EXAMPLE 6 Solving a Linear Equation with Decimals

Solve $0.06x + 0.09(15 - x) = 0.07(15)$.

$$0.06x + 0.09(15 - x) = 0.07(15) \quad \leftarrow \text{Clear parentheses first.}$$

$$\text{Step 1} \quad 0.06x + 0.09(15) + 0.09(-x) = 0.07(15) \quad \text{Distributive property}$$

$$0.06x + 1.35 - 0.09x = 1.05 \quad \text{Multiply.}$$

Now clear the decimals by multiplying by a power of 10. Because each decimal number is given in hundredths, multiply each term of the equation by 100. A number can be multiplied by 100 (that is, by 10^2) by moving the decimal point two places to the right.

NOW TRY ANSWER
5. $\{11\}$


NOW TRY EXERCISE 6

Solve.

$$0.08x - 0.12(x - 4) = 0.03(x - 5)$$

$$0.06x + 1.35 - 0.09x = 1.05$$

Multiply each term by 100.

Move decimal points 2 places to the right.

$$6x + 135 - 9x = 105$$

This is an equivalent equation without decimals.

$$-3x + 135 = 105$$

Combine like terms.

Step 2

$$-3x + 135 - 135 = 105 - 135$$

Subtract 135.

$$-3x = -30$$

Combine like terms.

Step 3

$$\frac{-3x}{-3} = \frac{-30}{-3}$$

Divide by -3 .

$$x = 10$$

Step 4 CHECK $0.06x + 0.09(15 - x) = 0.07(15)$

Original equation

$$0.06(10) + 0.09(15 - 10) \stackrel{?}{=} 0.07(15)$$

Let $x = 10$.

$$0.6 + 0.09(5) \stackrel{?}{=} 1.05$$

Multiply and subtract.

$$0.6 + 0.45 \stackrel{?}{=} 1.05$$

Multiply.

$$1.05 = 1.05 \quad \checkmark$$

True

The solution set is $\{10\}$.
NOW TRY 

NOTE Some students prefer to solve an equation with decimal coefficients without clearing the decimals.

$$0.06x + 0.09(15 - x) = 0.07(15)$$

Equation from Example 6

$$0.06x + 0.09(15) + 0.09(-x) = 0.07(15)$$

Distributive property

$$0.06x + 1.35 - 0.09x = 1.05$$

Multiply.

Be careful with decimal points.

$$-0.03x + 1.35 = 1.05$$

Combine like terms.

$$-0.03x + 1.35 - 1.35 = 1.05 - 1.35$$

Subtract 1.35.

$$-0.03x = -0.3$$

Combine like terms.

$$\frac{-0.03x}{-0.03} = \frac{-0.3}{-0.03}$$

Divide by -0.03 .

$$x = 10$$

The same solution results.

OBJECTIVE 6 Identify conditional equations, contradictions, and identities.

Some equations have one solution, some have no solution, and others have an infinite number of solutions. The table gives the names of these types of equations.

▼ Solution Sets of Equations

Type of Linear Equation	Number of Solutions	Indication When Solving
Conditional	One solution; solution set: {a number}	Final line is $x = a$ number. (See Examples 2–6, 7(a).)
Identity	Infinitely many solutions; solution set: {all real numbers}	Final line is true, such as $0 = 0$. (See Example 7(b).)
Contradiction	No solution; solution set: \emptyset	Final line is false, such as $-15 = -20$. (See Example 7(c).)

NOW TRY ANSWER
6. $\{9\}$


**NOW TRY
EXERCISE 7**

Solve each equation.
Determine whether it is a *conditional equation*, an *identity*, or a *contradiction*.

- (a) $9x - 3(x + 4) = 6(x - 2)$
 (b) $-3(2x - 1) - 2x = 3 + x$
 (c) $10x - 21 = 2(x - 5) + 8x$

EXAMPLE 7 Recognizing Conditional Equations, Identities, and Contradictions

Solve each equation. Determine whether it is a *conditional equation*, an *identity*, or a *contradiction*.

(a) $5(2x + 6) - 2 = 7(x + 4)$

$$10x + 30 - 2 = 7x + 28$$

Distributive property

$$10x + 28 = 7x + 28$$

Combine like terms.

$$10x + 28 - 7x - 28 = 7x + 28 - 7x - 28$$

Subtract $7x$. Subtract 28.

$$3x = 0$$

Combine like terms.

$$\frac{3x}{3} = \frac{0}{3}$$

Divide by 3.

$$x = 0$$

$\frac{0}{3} = 0$

The last line has a variable. The number following "=" is a solution.

CHECK

$$5(2x + 6) - 2 = 7(x + 4)$$

$$5[2(0) + 6] - 2 \stackrel{?}{=} 7(0 + 4)$$

$$5(6) - 2 \stackrel{?}{=} 7(4)$$

$$28 = 28 \quad \checkmark$$

Original equation
Let $x = 0$.
Multiply and then add.
True

The value 0 checks, so the solution set is $\{0\}$. Because the solution set has only one element, $5(2x + 6) - 2 = 7(x + 4)$ is a conditional equation.

(b) $5x - 15 = 5(x - 3)$

$$5x - 15 = 5x - 15$$

Distributive property

$$5x - 15 - 5x + 15 = 5x - 15 - 5x + 15$$

Subtract $5x$. Add 15.

$$0 = 0$$

True

The variable has "disappeared."

The *true* statement $0 = 0$ indicates that the solution set is {all real numbers}. The equation $5x - 15 = 5(x - 3)$ is an identity. Notice that the first step yielded

$$5x - 15 = 5x - 15, \quad \text{which is true for all values of } x.$$

We could have identified the equation as an identity at that point.

(c) $5x - 15 = 5(x - 4)$

$$5x - 15 = 5x - 20$$

Distributive property

$$5x - 15 - 5x = 5x - 20 - 5x$$

Subtract $5x$.

$$-15 = -20$$

False

The variable has "disappeared."

Because the result, $-15 = -20$, is *false*, the equation has no solution. The solution set is the empty set \emptyset , so the equation $5x - 15 = 5(x - 4)$ is a contradiction.

NOW TRY 

NOW TRY ANSWERS

7. (a) {all real numbers}; identity
 (b) $\{0\}$; conditional equation
 (c) \emptyset ; contradiction

CAUTION A common error in solving an equation like that in **Example 7(a)** is to think that the equation has no solution and write the solution set as \emptyset . This equation has one solution, the number 0, so it is a conditional equation with solution set $\{0\}$.

1.1 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Be sure to read and work through the section material before working the exercises.

Review Study Skill 2,
Reading Your Math Text.

Concept Check Complete each statement. The following key terms may be used once, more than once, or not at all.

linear equation solution algebraic expression contradiction all real numbers
solution set identity conditional equation first-degree equation empty set \emptyset

- A collection of numbers, variables, operation symbols, and grouping symbols, such as $2(8x - 15)$, is a(n) _____. While an equation (*does/does not*) include an equality symbol, there (*is/is not*) an equality symbol in an algebraic expression.
- A(n) _____ in one variable (here x) can be written in the form $ax + b$ ($=/ > / <$) 0 , with $a \neq 0$. Another name for a linear equation is a(n) _____, because the greatest power on the variable is (*one/two/three*).
- If we let $x = 2$ in the linear equation $2x + 5 = 9$, a (*true/false*) statement results. The number 2 is a(n) _____ of the equation, and $\{2\}$ is the _____.
- A linear equation with one solution in its _____, such as $2x + 5 = 9$, is a(n) _____.
- A linear equation with an infinite number of solutions is a(n) _____. Its solution set is {_____}.
- A linear equation with no solution is a(n) _____. Its solution set is the _____.
- Concept Check** Which equations are linear equations in x ?
A. $3x + x - 2 = 0$ B. $12 = x^2$ C. $9x - 4 = 9$ D. $3x + 2y = 6$
- Concept Check** Which of the equations in **Exercise 7** are nonlinear equations in x ? Explain why.

Determine whether each of the following is an expression or an equation. *See Example 1.*

- $-3x + 2 - 4 = x$
- $4(x + 3) - 2(x + 1) - 10$
- $-10x + 12 - 4x = -3$
- $-3x + 2 - 4 - x = 4$
- $4(x + 3) - 2(x + 1) + 10$
- $-10x + 12 - 4x + 3 = 0$
- Concept Check** This incorrect solution contains a common error.

$$8x - 2(2x - 3) = 3x + 7$$

$$8x - 4x - 6 = 3x + 7 \quad \text{Distributive property}$$

$$4x - 6 = 3x + 7 \quad \text{Combine like terms}$$

$$x = 13 \quad \text{Subtract } 3x. \text{ Add } 6.$$

WHAT WENT WRONG? Give the correct solution.

- Concept Check** This incorrect solution contains a common error.

$$12 - 2(3x + 1) = 11$$

$$10(3x + 1) = 11 \quad \text{Subtract } 12 - 2.$$

$$30x + 10 = 11 \quad \text{Distributive property}$$

$$30x = 1 \quad \text{Subtract } 10.$$

$$x = \frac{1}{30} \quad \text{Divide by } 30.$$

WHAT WENT WRONG? Give the correct solution.

- 17. Concept Check** Suppose we solve a linear equation and obtain, as our final result, an equation in Column I. Match each result with the solution set in Column II for the original equation.

I	II
(a) $7 = 7$	A. $\{0\}$
(b) $x = 0$	B. $\{\text{all real numbers}\}$
(c) $7 = 0$	C. \emptyset

- 18. Concept Check** Which one of the following linear equations does *not* have solution set $\{\text{all real numbers}\}$?

A. $4x = 5x - x$	B. $3(x + 4) = 3x + 12$
C. $4x = 3x$	D. $\frac{3}{4}x = 0.75x$
E. $4(x - 2) = 2(2x - 4)$	F. $2x + 18x = 20x$

Solve each equation, and check the solution. If applicable, tell whether the equation is an identity or a contradiction. See Examples 2–4 and 7.

- | | |
|-----------------------------------------------|---------------------------------------|
| 19. $7x + 8 = 1$ | 20. $5x - 4 = 21$ |
| 21. $5x + 2 = 3x - 6$ | 22. $9x + 1 = 7x - 9$ |
| 23. $7x - 5x + 15 = x + 8$ | 24. $2x + 4 - x = 4x - 5$ |
| 25. $12w + 15w - 9 + 5 = -3w + 5 - 9$ | 26. $-4x + 5x - 8 + 4 = 6x - 4$ |
| 27. $3(2t - 4) = 20 - 2t$ | 28. $2(3 - 2x) = x - 4$ |
| 29. $-5(x + 1) + 3x + 2 = 6x + 4$ | 30. $5(x + 3) + 4x - 5 = 4 - 2x$ |
| 31. $-2x + 5x - 9 = 3(x - 4) - 5$ | 32. $-6x + 2x - 11 = -2(2x - 3) + 4$ |
| 33. $-2(x + 3) = -6(x + 7)$ | 34. $-4(x - 9) = -8(x + 3)$ |
| 35. $3(2x + 1) - 2(x - 2) = 5$ | 36. $4(x - 2) + 2(x + 3) = 6$ |
| 37. $2x + 3(x - 4) = 2(x - 3)$ | 38. $6x - 3(5x + 2) = 4(1 - x)$ |
| 39. $6x - 4(3 - 2x) = 5(x - 4) - 10$ | 40. $-2x - 3(4 - 2x) = 2(x - 3) + 2$ |
| 41. $-2(x + 3) - x - 4 = -3(x + 4) + 2$ | 42. $4(2x + 7) = 2x + 25 + 3(2x + 1)$ |
| 43. $2[x - (2x + 4) + 3] = 2(x + 1)$ | 44. $4[2x - (3 - x) + 5] = -(2 + 7x)$ |
| 45. $-[2x - (5x + 2)] = 2 + (2x + 7)$ | 46. $-[6x - (4x + 8)] = 9 + (6x + 3)$ |
| 47. $-3x + 6 - 5(x - 1) = -5x - (2x - 4) + 5$ | |
| 48. $4(x + 2) - 8x - 5 = -3x + 9 - 2(x + 6)$ | |
| 49. $7[2 - (3 + 4x)] - 2x = -9 + 2(1 - 15x)$ | |
| 50. $4[6 - (1 + 2x)] + 10x = 2(10 - 3x) + 8x$ | |

Concept Check Answer each question.

51. To solve the following linear equation, we multiply each side by the least common denominator of all the fractions in the equation. What is this LCD?

$$\frac{3}{4}x - \frac{1}{3}x = \frac{5}{6}x - 5$$

52. Suppose that in solving the following equation, we multiply each side by 12, rather than the *least* common denominator, 6. Would we obtain the correct solution? Explain.

$$\frac{1}{3}x + \frac{1}{2}x = \frac{1}{6}x$$

53. To solve a linear equation with decimals, we usually begin by multiplying by a power of 10 so that all coefficients are integers. What is the least power of 10 that will accomplish this goal in each equation?

(a) $0.05x + 0.12(x + 5000) = 940$ (b) $0.006(x + 2) = 0.007x + 0.009$

54. The expression $0.06(10 - x)(100)$ is equivalent to which one of the following?

A. $0.06 - 0.06x$ B. $60 - 6x$ C. $6 - 6x$ D. $6 - 0.06x$

Solve each equation, and check the solution. See Examples 5 and 6.

55. $-\frac{5}{9}x = 2$

56. $\frac{3}{11}x = -5$

57. $\frac{6}{5}x = -1$

58. $-\frac{7}{8}x = 6$

59. $\frac{x}{2} + \frac{x}{3} = 5$

60. $\frac{x}{5} - \frac{x}{4} = 1$

61. $\frac{3x}{4} + \frac{5x}{2} = 13$

62. $\frac{8x}{3} - \frac{x}{2} = -13$

63. $\frac{x - 10}{5} + \frac{2}{5} = -\frac{x}{3}$

64. $\frac{5 - x}{6} + \frac{5}{6} = \frac{x}{54}$

65. $\frac{3x - 1}{4} + \frac{x + 3}{6} = 3$

66. $\frac{3x + 2}{7} - \frac{x + 4}{5} = 2$

67. $\frac{4x + 1}{3} = \frac{x + 5}{6} + \frac{x - 3}{6}$

68. $\frac{2x + 5}{5} = \frac{3x + 1}{2} + \frac{-x + 8}{16}$

69. $0.04x + 0.06 + 0.03x = 0.03x + 1.46$

70. $0.05x + 0.08 + 0.06x = 0.07x + 0.68$

71. $0.006x - 0.02x + 0.03 = 0.008x + 0.25$

72. $0.05x - 0.1x + 0.6 = 0.04x + 2.22$

73. $0.05x + 0.12(x + 5000) = 940$

74. $0.09x + 0.13(x + 300) = 61$

75. $0.02(50) + 0.08x = 0.04(50 + x)$

76. $0.04(90) + 0.12x = 0.06(460 + x)$

77. $0.05x + 0.10(200 - x) = 0.45x$

78. $0.08x + 0.12(260 - x) = 0.48x$

79. $0.006(x + 2) = 0.007x + 0.009$

80. $0.006(50 - x) = 0.272 - 0.004x$

1.2 Formulas and Percent

OBJECTIVES

- 1 Solve a formula for a specified variable.
- 2 Solve applied problems using formulas.
- 3 Solve percent problems.
- 4 Solve problems involving percent increase or decrease.
- 5 Solve problems from the health care industry.

A **mathematical model** is an equation or inequality that describes a real situation. Models for many applied problems, called *formulas*, already exist. A **formula** is an equation in which variables are used to describe a relationship. For example, the formula for finding the area \mathcal{A}^* of a triangle is

$$\mathcal{A} = \frac{1}{2}bh.$$

Here, b is the length of the base and h is the height. See **FIGURE 1**. A list of formulas used in algebra is given at the back of this text.




FIGURE 1

*In this text, we use \mathcal{A} to denote area.

OBJECTIVE 1 Solve a formula for a specified variable.

The formula $I = prt$ says that interest on a loan or investment equals principal (amount borrowed or invested) times rate (percent) times time at interest (in years). To determine how long it will take for an investment at a stated interest rate to earn a predetermined amount of interest, it would help to first solve the formula for t . This process is called **solving for a specified variable** or **solving a literal equation**.

When solving for a specified variable, the key is to treat that variable as if it were the only one. Treat all other variables like constants (numbers).

NOW TRY EXERCISE 1

Solve the formula $I = prt$ for p .

VOCABULARY

- mathematical model
- formula
- percent
- percent increase
- percent decrease

STUDY SKILLS REMINDER

Are you getting the most out of your class time?

Review Study Skill 3,
Taking Lecture Notes.

EXAMPLE 1 Solving for a Specified Variable

Solve the formula $I = prt$ for t .

We solve this formula for t by treating I , p , and r as constants (having fixed values) and treating t as the only variable.

$$\begin{aligned}
 prt &= I && \text{Our goal is to isolate } t. \\
 (pr)t &= I && \text{Associative property} \\
 \frac{(pr)t}{pr} &= \frac{I}{pr} && \text{Divide by } pr. \\
 t &= \frac{I}{pr} && \text{The result is a formula for time } t, \text{ in years.}
 \end{aligned}$$


NOW TRY**Solving for a Specified Variable**

- Step 1 Clear any parentheses and fractions.** If the equation contains parentheses, clear them using the distributive property. Clear any fractions by multiplying both sides by the LCD.
- Step 2 Isolate all terms with the specified variable.** Transform so that all terms containing the specified variable are on one side of the equation and all terms without that variable are on the other side.
- Step 3 Isolate the specified variable.** Divide each side by the factor that is the coefficient of the specified variable.

EXAMPLE 2 Solving for a Specified Variable

Solve the formula $P = 2L + 2W$ for W .

This formula gives the relationship between perimeter of a rectangle, P , length of the rectangle, L , and width of the rectangle, W . See **FIGURE 2**.



Perimeter, P , the sum of the lengths of the sides of this rectangle, is given by

$$P = 2L + 2W.$$

FIGURE 2**NOW TRY ANSWER**

1. $p = \frac{I}{rt}$

$$P = 2L + 2W$$

Our goal is to isolate W .


**NOW TRY
EXERCISE 2**
Solve the formula for b .

$$P = a + 2b + c$$

Step 1 is not needed here because there are no parentheses or fractions to clear in this formula.

$$P = 2L + 2W$$

Step 2 $P - 2L = 2L + 2W - 2L$ Subtract $2L$.

$$P - 2L = 2W \quad \text{Combine like terms.}$$

Step 3 $\frac{P - 2L}{2} = \frac{2W}{2}$ Divide by 2.

$$\frac{P - 2L}{2} = W \quad \frac{2W}{2} = \frac{2}{2} \cdot W = 1 \cdot W = W$$

Although W is isolated, we simplify the right side further.

$$W = \frac{P - 2L}{2}$$

Interchange sides.

Be careful here. $\frac{P - 2L}{2} \neq P - L$

$$W = \frac{P}{2} - \frac{2L}{2}$$

$$\frac{a - b}{c} = \frac{a}{c} - \frac{b}{c}$$

$$W = \frac{P}{2} - L$$

$$\frac{2L}{2} = \frac{2}{2} \cdot L = 1 \cdot L = L$$

NOW TRY 

! CAUTION In Step 3 of **Example 2**, we cannot simplify the fraction $\frac{P - 2L}{2}$ by dividing 2 into the term $2L$. The fraction bar serves as a grouping symbol. Thus, the subtraction in the numerator must be done before the division.


**NOW TRY
EXERCISE 3**
Solve $P = 2(L + W)$ for L .
EXAMPLE 3 Solving a Formula Involving Parentheses
Solve the formula $s = \frac{1}{2}(a + b + c)$ for a .

$$s = \frac{1}{2}(a + b + c)$$

Step 1 $s = \frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}c$ Distributive property

$$2s = 2\left(\frac{1}{2}a + \frac{1}{2}b + \frac{1}{2}c\right) \quad \text{Multiply by 2 to clear the fractions.}$$

$$2s = a + b + c \quad \text{Distributive property}$$

Step 2 $2s - b - c = a + b + c - b - c$ Subtract b and c .

Step 3 $a = 2s - b - c$ Combine like terms. Interchange sides.

NOW TRY 
NOW TRY ANSWERS

2. $b = \frac{P - a - c}{2}$

3. $L = \frac{P - 2W}{2}$, or $L = \frac{P}{2} - W$

In **Examples 1–3**, we solved formulas for specified variables. In **Example 4**, we solve an equation with two variables for one of these variables. This process will be useful later when we work with *linear equations in two variables*.


NOW TRY EXERCISE 4
Solve $5x - 6y = 12$ for y .
EXAMPLE 4 Solving an Equation for One of the Variables
Solve $3x - 4y = 12$ for y .

$$3x - 4y = 12$$

Our goal is to isolate y .

$$3x - 4y - 3x = 12 - 3x$$

Subtract $3x$.

$$-4y = -3x + 12$$

Combine like terms; commutative property

$$\frac{-4y}{-4} = \frac{-3x + 12}{-4}$$

Divide by -4 .

$$y = \frac{-3x}{-4} + \frac{12}{-4}$$

 $\frac{a+b}{c} = \frac{a}{c} + \frac{b}{c}$

$$\frac{-3x}{-4} = \frac{-3}{-4} \cdot \frac{x}{1} = \frac{3}{4}x$$

$$y = \frac{3}{4}x - 3$$

Simplify the expression on the right.

NOW TRY 
OBJECTIVE 2 Solve applied problems using formulas.

The distance formula $d = rt$ relates d , the distance traveled, r , the rate or speed, and t , the travel time.

EXAMPLE 5 Finding Average Rate

Phyllis found that on average it took her $\frac{3}{4}$ hr each day to drive a distance of 15 mi to work. What was her average rate (or speed)?

$$d = rt \quad \text{Solve the distance formula for } r.$$

$$\frac{d}{t} = \frac{rt}{t} \quad \text{Divide by } t.$$

$$\frac{d}{t} = r, \quad \text{or} \quad r = \frac{d}{t}$$


We find the desired rate by substituting the given values of d and t into this formula.

$$r = \frac{15}{\frac{3}{4}} \quad \text{Let } d = 15, t = \frac{3}{4}.$$

$$r = 15 \cdot \frac{4}{3} \quad \text{To divide by } \frac{3}{4}, \text{ multiply by its reciprocal, } \frac{4}{3}.$$

$$r = 20$$

Her average rate was 20 mph. (That is, at times she may have traveled a little faster or slower than 20 mph, but overall her rate was 20 mph.)

NOW TRY 
OBJECTIVE 3 Solve percent problems.

A percent is a ratio where the second number is always 100. The percent symbol is %.

50% represents the ratio of 50 to 100—that is, $\frac{50}{100}$, or, as a decimal, 0.50.

The word **percent** means “per 100.” One percent means “one per 100.”

$$1\% = \frac{1}{100} = 0.01, \quad 10\% = \frac{10}{100} = 0.10, \quad 100\% = \frac{100}{100} = 1$$


NOW TRY EXERCISE 5

It takes $\frac{1}{2}$ hr for Dorothy to drive 21 mi to work each day.

What is her average rate?

NOW TRY ANSWERS

4. $y = \frac{5}{6}x - 2$

5. 42 mph

Solving a Percent Problem

Let a represent a partial amount of b , the whole amount (or base). Then the following equation can be used to solve a percent problem.

$$\frac{\text{partial amount } a}{\text{whole amount } b} = \text{decimal value (which is converted to a percent)}$$

For example, if a class consists of 50 students and 32 are males, then the percent of males in the class is found as follows.

$$\begin{aligned} \frac{\text{partial amount } a}{\text{whole amount } b} &= \frac{32}{50} && \text{Let } a = 32, b = 50. \\ &= \frac{64}{100} && \frac{32}{50} \cdot \frac{2}{2} = \frac{64}{100} \\ &= 0.64 && \text{Write as a decimal.} \\ &= 64\% && \text{Convert to a percent.} \end{aligned}$$

NOTE When interpreting the “partial amount” in the formula for solving a percent problem, understand that it may represent a quantity that is actually *larger* than the whole amount. In these cases, the percent will be greater than 100%. For example,

$$25 \text{ is } 250\% \text{ of } 10.$$

Here 25 is the partial amount and 10 is the whole amount.



NOW TRY EXERCISE 6

Solve each problem.

- (a) A 5-L mixture of water and antifreeze contains 2 L of antifreeze. What is the percent of antifreeze in the mixture?
- (b) If a savings account earns 2.5% interest on a balance of \$7500 for one year, how much interest is earned?

EXAMPLE 6 Solving Percent Problems

- (a) A 50-L mixture of acid and water contains 10 L of acid. What is the percent of acid in the mixture?

The given amount of the mixture is 50 L, and the part that is acid is 10 L. Let x represent the percent of acid in the mixture.

$$\begin{aligned} \frac{\text{partial amount}}{\text{whole amount}} &\rightarrow \frac{10}{50} = x \\ x &= 0.20 && \text{Divide to write as a decimal.} \\ x &= 20\% && \text{Interchange sides.} \\ &&& 0.20 = \frac{20}{100} = 20\% \end{aligned}$$

The mixture is 20% acid.

- (b) If \$4780 earns 2% interest in one year, how much interest is earned?

Let x represent the amount of interest earned (that is, the part of the whole amount invested). Because $2\% = 0.02$, the equation is written as follows.

$$\begin{aligned} \frac{x}{4780} &= 0.02 && \frac{\text{partial amount } a}{\text{whole amount } b} = \text{decimal value} \\ 4780 \cdot \frac{x}{4780} &= 4780(0.02) && \text{Multiply by 4780.} \\ x &= 95.6 \end{aligned}$$

The interest earned is \$95.60.

NOW TRY ANSWERS

6. (a) 40% (b) \$187.50


NOW TRY EXERCISE 7

Refer to **FIGURE 3**. How much was spent on vet care? Round the answer to the nearest tenth of a billion dollars.

EXAMPLE 7 Interpreting Percents from a Graph

In 2017, Americans spent about \$69.5 billion on their pets. Use the graph in **FIGURE 3** to determine how much of this amount was spent on pet food.

Spending on Kitty and Rover



Data from American Pet Products Manufacturers Association Inc.



Bert

FIGURE 3

The graph shows that 41.8% was spent on food. Let x = this amount in billions of dollars.

$$\begin{array}{l} \text{partial amount} \rightarrow \frac{x}{69.5} = 0.418 \\ \text{whole amount} \rightarrow 69.5 \end{array} \quad 41.8\% = 0.418$$

$$69.5 \cdot \frac{x}{69.5} = 69.5(0.418) \quad \text{Multiply by 69.5.}$$

$$x \approx 29.1 \quad \text{Nearest tenth}$$

Therefore, about \$29.1 billion was spent on pet food.

NOW TRY

OBJECTIVE 4 Solve problems involving percent increase or decrease.

Percent is often used to express a change in some quantity. Buying an item that has been marked up and getting a raise at a job are applications of **percent increase**. Buying an item on sale and finding population decline are applications of **percent decrease**.

To solve problems of this type, we use the following equation.

$$\text{percent change (as a decimal)} = \frac{\text{amount of change}}{\text{original amount}} \quad \begin{array}{l} \text{Subtract to} \\ \text{find this.} \end{array}$$

EXAMPLE 8 Solving Problems about Percent Increase or Decrease

Use the percent change equation in each application.

- (a) An electronics store marked up a laptop computer from their cost of \$1200 to a selling price of \$1464. What was the percent markup?

“Markup” is a name for an increase.

Let x = the percent increase (as a decimal).

NOW TRY ANSWER

7. \$17.1 billion


NOW TRY EXERCISE 8

Use the percent change equation in each application.

- (a) Jane bought a jacket on sale for \$56. The regular price of the jacket was \$80. What was the percent markdown?
- (b) When it was time for Horatio to renew the lease on his apartment, the landlord raised his monthly rent from \$650 to \$689. What was the percent increase?

$$\text{percent increase} = \frac{\text{amount of increase}}{\text{original amount}}$$

Subtract to find the amount of increase.

$$x = \frac{1464 - 1200}{1200}$$

Substitute the given values.

$$x = \frac{264}{1200}$$

Use the original cost.

$$x = 0.22$$

Use a calculator.

$$x = 22\%$$

Convert to a percent.

The computer was marked up 22%.

- (b) The enrollment at a community college declined from 12,750 during one school year to 11,350 the following year. Find the percent decrease.

Let x = the percent decrease (as a decimal).

$$\text{percent decrease} = \frac{\text{amount of decrease}}{\text{original amount}}$$

Subtract to find the amount of decrease.

$$x = \frac{12,750 - 11,350}{12,750}$$

Substitute the given values.

$$x = \frac{1400}{12,750}$$

Use the original number.

$$x \approx 0.11$$

Use a calculator.

$$x = 11\%$$

Convert to a percent.

The college enrollment decreased by about 11%.

NOW TRY 

! CAUTION When calculating a percent increase or decrease, be sure to use the original number (*before* the increase or decrease) as the denominator of the fraction, **not** the final number (*after* the increase or decrease).

$$\text{percent change} = \frac{\text{amount of change}}{\text{original amount}}$$

OBJECTIVE 5 Solve problems from the health care industry.

EXAMPLE 9 Determining a Child's Body Surface Area

If a child weighs k kilograms, then the child's body surface area S in square meters (m^2) is determined by the formula

$$S = \frac{4k + 7}{k + 90}$$

What is the body surface area of a child who weighs 40 lb? (Data from Hegstad, L., and W. Hayek, *Essential Drug Dosage Calculations*, Fourth Edition, Prentice Hall.)

To convert 40 lb to kilograms, we use the identity property of multiplication and the fact that $1 \text{ lb} \approx 0.4536 \text{ kg}$.

This fraction is approximately equal to 1.

$$40 \text{ lb} \cdot \frac{0.4536 \text{ kg}}{1 \text{ lb}} \approx 18.144 \text{ kg}$$

NOW TRY ANSWERS

8. (a) 30% (b) 6%


NOW TRY EXERCISE 9

Use the formula in **Example 9** to find the body surface area, to the nearest hundredth, of a child who weighs 32 lb.

Use the value 18.144 for k in the given formula.

$$S = \frac{4k + 7}{k + 90}$$

$$S = \frac{4(18.144) + 7}{18.144 + 90} \quad \text{Let } k = 18.144.$$

$$S \approx 0.74 \quad \text{Use a calculator. Round to the nearest hundredth.}$$

The child's body surface area is 0.74 m^2 .

NOW TRY 


NOW TRY EXERCISE 10

Refer to **Now Try Exercise 9**. Determine the appropriate dose, to the nearest unit, for this child if the usual adult dose is 100 mg.

EXAMPLE 10 Determining a Child's Dose of a Medication

If D represents the usual adult dose of a medication in milligrams, the corresponding child's dose C in milligrams is calculated using the formula

$$C = \frac{\text{body surface area (in square meters)}}{1.7} \times D.$$

Determine the appropriate dose for a child weighing 40 lb if the usual adult dose is 50 mg. (Data from Hegstad, L., and W. Hayek, *Essential Drug Dosage Calculations*, Fourth Edition, Prentice Hall.)

From **Example 9**, the body surface area of a child weighing 40 lb is 0.74 m^2 . Use this value in the above formula to find the child's dose.

$$C = \frac{\text{body surface area (in square meters)}}{1.7} \times D$$

$$C = \frac{0.74}{1.7} \times 50 \quad \text{Body surface area} = 0.74, D = 50$$

$$C \approx 22 \quad \text{Use a calculator. Round to the nearest unit.}$$

The child's dose is 22 mg.

NOW TRY 

NOW TRY ANSWERS


9. 0.62 m^2

10. 36 mg

1.2 Exercises

FOR
EXTRA
HELP

 **MyLab Math**

 Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

How are you doing on your homework? **Review Study Skill 4, Completing Your Homework.**

Concept Check Fill in each blank with the correct response.

1. A(n) _____ is an equation in which variables are used to describe a relationship.
2. To solve a formula for a specified variable, treat that _____ as if it were the only one and treat all other variables like _____ (numbers).


Concept Check Work each problem to review converting between decimals and percents.

3. Write each decimal as a percent.
(a) 0.35 (b) 0.18 (c) 0.02 (d) 0.075 (e) 1.5
4. Write each percent as a decimal.
(a) 60% (b) 37% (c) 8% (d) 3.5% (e) 210%


Solve each formula for the specified variable. See Examples 1–3.

5. $I = prt$ for r (simple interest)
6. $d = rt$ for t (distance)


7. $\mathcal{A} = LW$ (area of a rectangle)
 (a) for W (b) for L




9. $P = 2L + 2W$ for L
 (perimeter of a rectangle)




11. $V = LWH$
 (volume of a rectangular solid)
 (a) for W (b) for H



13. $C = 2\pi r$ for r
 (circumference of a circle)



15. $\mathcal{A} = \frac{1}{2}h(b + B)$ (area of a trapezoid)
 (a) for b (b) for B




17. $F = \frac{9}{5}C + 32$ for C
 (Celsius to Fahrenheit)


19. $ax + b = 0$
 (linear equation in x)
 (a) for x (b) for a

21. $A = P(1 + rt)$ for t
 (future value for simple interest)


8. $\mathcal{A} = bh$ (area of a parallelogram)
 (a) for b (b) for h




10. $P = a + b + c$ (perimeter of a triangle)
 (a) for b (b) for c




12. $\mathcal{A} = \frac{1}{2}bh$ (area of a triangle)
 (a) for h (b) for b



14. $V = \pi r^2 h$ for h
 (volume of a right circular cylinder)



16. $V = \frac{1}{3}\pi r^2 h$ for h (volume of a cone)



18. $C = \frac{5}{9}(F - 32)$ for F
 (Fahrenheit to Celsius)

20. $y = mx + b$
 (slope-intercept form of a linear equation)
 (a) for x (b) for m

22. $M = C(1 + r)$ for r
 (markup)

- 23. Concept Check** When a formula is solved for a particular variable, several different equivalent forms may be possible. If we solve $\mathcal{A} = \frac{1}{2}bh$ for h , one correct answer is

$$h = \frac{2\mathcal{A}}{b}.$$

Which one of the following is *not* equivalent to this?

A. $h = 2\left(\frac{\mathcal{A}}{b}\right)$ B. $h = 2\mathcal{A}\left(\frac{1}{b}\right)$ C. $h = \frac{\mathcal{A}}{\frac{1}{2}b}$ D. $h = \frac{\frac{1}{2}\mathcal{A}}{b}$

- 24. Concept Check** The formula $F = \frac{9}{5}C + 32$ can be solved for C to obtain $C = \frac{5}{9}(F - 32)$. Which one of the following is *not* equivalent to this?


A. $C = \frac{5}{9}F - \frac{160}{9}$ B. $C = \frac{5F}{9} - \frac{160}{9}$ C. $C = \frac{5F - 160}{9}$ D. $C = \frac{5}{9}F - 32$

Solve each equation for y . See Example 4.

25. $4x + y = 1$ 26. $3x + y = 9$ 27. $x - 2y = -6$
 28. $x - 5y = -20$ 29. $4x + 9y = 11$ 30. $7x + 8y = 11$
 31. $-3x + 2y = 5$ 32. $-5x + 3y = 12$ 33. $6x - 5y = 7$
 34. $8x - 3y = 4$ 35. $\frac{1}{2}x - \frac{1}{3}y = 1$ 36. $\frac{2}{3}x - \frac{2}{5}y = 2$

Solve each problem. See Example 5.

37. Kurt Busch won the Daytona 500 (mile) race with a rate of 143.187 mph in 2017. Find his time to the nearest thousandth. (Data from *The World Almanac and Book of Facts*.)
38. In 2007, rain shortened the Indianapolis 500 race to 415 mi. It was won by Dario Franchitti, who averaged 151.774 mph. What was his time to the nearest thousandth? (Data from *The World Almanac and Book of Facts*.)
39. Nora traveled from Kansas City to Louisville, a distance of 520 mi, in 10 hr. Find her rate in miles per hour.
40. The distance from Melbourne to London is 10,500 mi. If a jet averages 500 mph between the two cities, what is its travel time in hours?
41. As of 2017, the highest temperature ever recorded in the state of Washington was 42°C . Find the corresponding Fahrenheit temperature to the nearest degree. (Data from National Climatic Data Center.)
42. As of 2017, the lowest temperature ever recorded in South Dakota was -41°F . Find the corresponding Celsius temperature to the nearest degree. (Data from National Climatic Data Center.)
43. The base of the Great Pyramid of Cheops is a square whose perimeter is 920 m. What is the length of each side of this square? (Data from *Atlas of Ancient Archaeology*.)



In baseball, winning percentage (*Pct.*) is commonly expressed as a decimal rounded to the nearest thousandth. To find the winning percentage of a team, divide the number of wins (*W*) by the total number of games played ($W + L$).

59. The final 2017 standings of the American League West division in Major League Baseball are shown in the table. Find the winning percentage of each team.

- (a) L.A. Angels (b) Seattle
(c) Texas (d) Oakland

	W	L	Pct.
Houston	101	61	.623
L.A. Angels	80	82	
Seattle	78	84	
Texas	78	84	
Oakland	75	87	

Data from *The World Almanac and Book of Facts*.

60. The final 2017 standings of the National League Central division in Major League Baseball are shown in the table. Find the winning percentage of each team.

- (a) Milwaukee (b) St. Louis
(c) Pittsburgh (d) Cincinnati

	W	L	Pct.
Chicago Cubs	92	70	.568
Milwaukee	86	76	
St. Louis	83	79	
Pittsburgh	75	87	
Cincinnati	68	94	

Data from *The World Almanac and Book of Facts*.

In June 2017, 69.5 million U.S. households owned at least one Internet-enabled device (such as a smart TV, a video game console, or a multimedia device) that could stream video to a TV set. (Data from *The Nielsen Company*.) Use this information to work each problem. Round answers to the nearest percent or to the nearest tenth of a million, as needed. **See Example 6.**


61. About 13.4 million U.S. households owned only a smart TV in 2017. What percent of those owning at least one Internet-enabled device was this?
62. About 14.5 million U.S. households owned only a video game console in 2017. What percent of those owning at least one Internet-enabled device was this?
63. Of the households owning at least one such device in 2017, 9.4% owned all three types. How many households owned all three?
64. Of the households owning at least one Internet-enabled device in 2017, 15.8% owned both a video game console and a smart TV. How many households owned both of these?



An average middle-income family will spend \$241,080 to raise a child from birth through age 17. The graph shows the percents spent for various categories. Use the graph to answer each question. **See Example 7.**

65. To the nearest dollar, how much will be spent to provide housing for the child?
66. To the nearest dollar, how much will be spent for child care and education?
67. How much will be spent on clothing?

The Cost of Parenthood



Data from U.S. Department of Agriculture.

68. About \$38,500 will be spent for food. To the nearest percent, what percent of the cost of raising a child from birth through age 17 is this? Does the answer agree with the graph?

Solve each problem. See Example 8.

69. After 1 yr on the job, Grady got a raise from \$10.50 per hour to \$11.76 per hour. What was the percent increase in his hourly wage?
70. Clayton bought a ticket to a rock concert at a discount. The regular price of the ticket was \$70.00, but he paid only \$56.00. What was the percent discount?
71. Between 2000 and 2016, the estimated population of metro New Orleans, Louisiana, declined from 1,337,726 to 1,268,883. What was the percent decrease to the nearest tenth? (Data from U.S. Census.)
72. Between 2000 and 2016, the estimated population of metro Chicago, Illinois, grew from 9,098,316 to 9,512,999. What was the percent increase to the nearest tenth? (Data from U.S. Census.)
73. The movie *Transformers: The Last Knight* in DVD was on sale for \$16.17. The list price (full price) of this disc was \$17.99. To the nearest tenth, what was the percent discount? (Data from www.amazon.com)
74. The *Jurassic Park Collection* in Blu-Ray was on sale for \$29.96. The list price (full price) of this three-disc set was \$44.98. To the nearest tenth, what was the percent discount? (Data from www.amazon.com)

Apply the formulas for determining a child's body surface area and a child's dose of a medication. See Examples 9 and 10.

$$S = \frac{4k + 7}{k + 90}$$

Child's body surface area S ,
in square meters;
 k = weight in kilograms

$$C = \frac{\text{body surface area (in square meters)}}{1.7} \times D$$

Child's dose of medication C ,
in milligrams;
 D = usual adult dose, in milligrams

75. Find the body surface area, to the nearest hundredth, of a child with each weight.
- (a) 20 kg (b) 26 kg
76. Find the body surface area, to the nearest hundredth, of a child with each weight.
- (a) 30 lb (b) 26 lb
77. If the usual adult dose of a medication is 250 mg, what is the child's dose, to the nearest unit, for a child with each weight? (Hint: Use the results of Exercise 75.)
- (a) 20 kg (b) 26 kg
78. If the usual adult dose of a medication is 500 mg, what is the child's dose, to the nearest unit, for a child with each weight? (Hint: Use the results of Exercise 76.)
- (a) 30 lb (b) 26 lb

RELATING CONCEPTS For Individual or Group Work (Exercises 79–85)

Consider the following equations.

First Equation

$$\frac{7x + 8}{3} = 12$$

Second Equation

$$\frac{ax + k}{c} = t \quad (c \neq 0)$$

Solving the second equation for x requires the same logic as solving the first equation for x . **Work Exercises 79–85 in order**, to see this “parallel logic.”

- 79.** (a) Clear the first equation of fractions by multiplying each side by 3.
 (b) Clear the second equation of fractions by multiplying each side by c .
- 80.** (a) Transform so that the term involving x is on the left side of the first equation and the constants are on the right by subtracting 8 from each side. (Do not simplify yet.)
 (b) Transform so that the term involving x is on the left side of the second equation by subtracting k from each side. (Do not simplify yet.)
- 81.** (a) Combine like terms in the first equation.
 (b) Combine like terms in the second equation.
- 82.** (a) Divide each side of the first equation by the coefficient of x . Give the solution set.
 (b) Divide each side of the second equation by the coefficient of x .
- 83.** Look at the answer for the second equation in **Exercise 82(b)**. What restriction must be placed on the variables? Why is this necessary?

Now apply the concepts developed in **Exercises 79–83** to work the following problems.

- 84.** Solve the equation. Give the solution set.

$$28 = \frac{7}{2}(a + 13)$$

- 85.** Refer to the steps used to solve the equation in **Exercise 84**, and solve the following formula (from the mathematical topic of series) for a .

$$S = \frac{n}{2}(a + \ell)$$

What restriction must be placed on the variables?

1.3

Applications of Linear Equations

OBJECTIVES

- 1 Translate from words to mathematical expressions.
- 2 Write equations from given information.
- 3 Distinguish between simplifying expressions and solving equations.

OBJECTIVE 1 Translate from words to mathematical expressions.

Producing a mathematical model of a real situation often involves translating verbal statements into mathematical statements.

PROBLEM-SOLVING HINT

Usually there are key words and phrases in a verbal problem that translate into mathematical expressions involving addition, subtraction, multiplication, and division. Translations of some commonly used expressions follow in the table on the next page.

OBJECTIVES (continued)

- 4 Use the six steps in solving an applied problem.
- 5 Solve percent problems.
- 6 Solve investment problems.
- 7 Solve mixture problems.

Use this table for reference when solving applications in this chapter.

▼ **Translating from Words to Mathematical Expressions**

<i>Verbal Expression</i>	<i>Mathematical Expression (where x and y are numbers)</i>
Addition	
The sum of a number and 7	$x + 7$
6 more than a number	$x + 6$
3 plus a number	$3 + x$
24 added to a number	$x + 24$
A number increased by 5	$x + 5$
The sum of two numbers	$x + y$
Subtraction	
2 less than a number	$x - 2$
2 less a number	$2 - x$
12 minus a number	$12 - x$
A number decreased by 12	$x - 12$
A number subtracted from 10	$10 - x$
10 subtracted from a number	$x - 10$
The difference of two numbers	$x - y$
Multiplication	
16 times a number	$16x$
A number multiplied by 6	$6x$
$\frac{2}{3}$ of a number (used with fractions and percent)	$\frac{2}{3}x$
$\frac{3}{4}$ as much as a number	$\frac{3}{4}x$
Twice (2 times) a number	$2x$
Triple (3 times) a number	$3x$
The product of two numbers	xy
Division	
The quotient of 8 and a number	$\frac{8}{x}$ ($x \neq 0$)
A number divided by 13	$\frac{x}{13}$
The ratio of two numbers x and y	$\frac{x}{y}$ ($y \neq 0$)
The quotient of two numbers x and y	$\frac{x}{y}$ ($y \neq 0$)

! CAUTION Because subtraction and division are not commutative operations, it is important to correctly translate expressions involving them.

Examples: “2 less than a number” is translated as $x - 2$, **not** $2 - x$.

“A number subtracted from 10” is translated as $10 - x$, **not** $x - 10$.

For division, the number *by which* we are dividing is the denominator, and the number *into which* we are dividing is the numerator.

Examples: “A number divided by 13” and “13 divided into x ” both translate as $\frac{x}{13}$.

“The quotient of x and y ” translates as $\frac{x}{y}$.

OBJECTIVE 2 Write equations from given information.

Any words that indicate equality or “sameness,” such as *is*, translate as $=$.

NOW TRY EXERCISE 1

Translate each verbal sentence into an equation, using x as the variable.

- (a) The quotient of a number and 10 is twice the number.
 (b) The product of a number and 5, decreased by 7, is 0.

EXAMPLE 1 Translating Words into Equations

Translate each verbal sentence into an equation, using x as the variable.

Verbal Sentence	Equation
Twice a number, decreased by 3, is 42.	$2x - 3 = 42$
The product of a number and 12, decreased by 7, is 105.	$12x - 7 = 105$
The quotient of a number and the number plus 4 is 28.	$\frac{x}{x + 4} = 28$
The quotient of a number and 4, plus the number, is 10.	$\frac{x}{4} + x = 10$

NOW TRY**OBJECTIVE 3** Distinguish between simplifying expressions and solving equations.

Recall the difference between an *expression* and an *equation*.

- An expression translates as a phrase.
- An equation includes the equality symbol ($=$), with expressions on both sides, and translates as a sentence.

NOW TRY EXERCISE 2

Determine whether each is an *expression* or an *equation*. Simplify any expressions, and solve any equations.

- (a) $3(x - 5) + 2x - 1$
 (b) $3(x - 5) + 2x = 1$

EXAMPLE 2 Simplifying Expressions vs. Solving Equations

Determine whether each is an *expression* or an *equation*. Simplify any expressions, and solve any equations.

(a) $2(3 + x) - 4x + 7$

Clear parentheses and combine like terms to **simplify**.

$$2(3 + x) - 4x + 7$$

$$= 6 + 2x - 4x + 7$$

$$= -2x + 13$$

There is no equality symbol.
This is an **expression**.

Distributive property

Simplified expression

(b) $2(3 + x) - 4x + 7 = -1$

Find the value of x to **solve**.

$$2(3 + x) - 4x + 7 = -1$$

$$6 + 2x - 4x + 7 = -1$$

$$-2x + 13 = -1$$

$$-2x = -14$$

$$x = 7$$

There is an equality symbol.
This is an **equation**.

Distributive property

Combine like terms.

Subtract 13.

Divide by -2 .

NOW TRY ANSWERS

1. (a) $\frac{x}{10} = 2x$ (b) $5x - 7 = 0$
 2. (a) expression; $5x - 16$
 (b) equation; $\left\{\frac{16}{5}\right\}$

The solution set is $\{7\}$.

NOW TRY

OBJECTIVE 4 Use the six steps in solving an applied problem.

The following six steps are helpful when solving applied problems.

Solving an Applied Problem

- Step 1** **Read** the problem carefully. *What information is given? What is to be found?*
- Step 2** **Assign a variable** to represent the unknown value. Make a sketch, diagram, or table, as needed. If necessary, express any other unknown values in terms of the variable.
- Step 3** **Write an equation** using the variable expression(s).
- Step 4** **Solve** the equation.
- Step 5** **State the answer.** Label it appropriately. *Does it seem reasonable?*
- Step 6** **Check** the answer in the words of the *original* problem.

NOW TRY EXERCISE 3

The length of a rectangle is 2 ft more than twice the width. The perimeter is 34 ft. Find the length and the width of the rectangle.

EXAMPLE 3 Solving a Perimeter Problem

The length of a rectangle is 1 cm more than twice the width. The perimeter of the rectangle is 110 cm. Find the length and the width of the rectangle.

- Step 1** **Read** the problem. *What must be found? We must find the length and width of a rectangle. What is given? The length is 1 cm more than twice the width and the perimeter is 110 cm.*




FIGURE 4

- Step 2** **Assign a variable.** Make a sketch, as shown in FIGURE 4.

Let W = the width.

Then $2W + 1$ = the length.

- Step 3** **Write an equation.** Use the formula for the perimeter of a rectangle.

$$P = 2L + 2W \quad \text{Perimeter of a rectangle}$$

$$110 = 2(2W + 1) + 2W \quad P = 110 \text{ and } L = 2W + 1$$

- Step 4** **Solve.** $110 = 4W + 2 + 2W$ Distributive property

$$110 = 6W + 2 \quad \text{Combine like terms.}$$

$$110 - 2 = 6W + 2 - 2 \quad \text{Subtract 2.}$$

$$108 = 6W \quad \text{Combine like terms.}$$

$$\frac{108}{6} = \frac{6W}{6} \quad \text{Divide by 6.}$$

$$18 = W$$

We also need to find the length.

- Step 5** **State the answer.** The width of the rectangle is 18 cm and the length is $2(18) + 1 = 37$ cm.

- Step 6** **Check.** The length, 37 cm, is 1 cm more than $2(18)$ cm (twice the width). The perimeter is $2(37) + 2(18) = 110$ cm, as required.

NOW TRY ANSWER

3. width: 5 ft; length: 12 ft



NOW TRY EXERCISE 4

During the 2017 regular NFL season, Russel Wilson of the Seattle Seahawks threw 11 more touchdown passes than Drew Brees of the New Orleans Saints. Together, these two quarterbacks completed a total of 57 touchdown passes. How many passes did each player complete? (Data from www.nfl.com)

EXAMPLE 4 Finding Unknown Numerical Quantities

During the 2017 regular Major League Baseball season, Chris Sale of the Boston Red Sox and Max Scherzer of the Washington Nationals were the top major league pitchers for strikeouts. The two pitchers had a total of 576 strikeouts. Sale had 40 more strikeouts than Scherzer. How many strikeouts did each pitcher have? (Data from www.mlb.com)

Step 1 Read the problem. We are asked to find the number of strikeouts each pitcher had.

Step 2 Assign a variable to represent the number of strikeouts for one of the men.

Let s = the number of strikeouts for Max Scherzer.

We must also find the number of strikeouts for Chris Sale. Because he had 40 more strikeouts than Scherzer,

$s + 40$ = the number of strikeouts for Sale.

Step 3 Write an equation. The sum of the numbers of strikeouts is 576.

$$\begin{array}{rccccccc} \text{Scherzer's strikeouts} & + & \text{Sale's strikeouts} & = & \text{total strikeouts.} & & \\ \downarrow & & \downarrow & & \downarrow & & \\ s & + & (s + 40) & = & 576 & & \end{array}$$

Step 4 Solve.

$$s + (s + 40) = 576$$

$$2s + 40 = 576 \quad \text{Combine like terms.}$$

$$2s + 40 - 40 = 576 - 40 \quad \text{Subtract 40.}$$

$$2s = 536 \quad \text{Combine like terms.}$$

$$\frac{2s}{2} = \frac{536}{2} \quad \text{Divide by 2.}$$

$$s = 268$$

Don't stop here.

Step 5 State the answer. The variable s represents the number of strikeouts for Scherzer, so he had 268. Then Sale had $s + 40$, which is

$$268 + 40 = 308 \text{ strikeouts.}$$

Step 6 Check. 308 is 40 more than 268, and $268 + 308 = 576$. The conditions of the problem are satisfied, and our answer checks.

NOW TRY

CAUTION Be sure to answer all the questions asked in the problem. In Example 4, we were asked for the number of strikeouts for each player, so there was extra work in Step 5 in order to find Sale's number.

PROBLEM-SOLVING HINT In Example 4, we chose to let the variable represent the number of strikeouts for Scherzer. Students often ask, "Can I let the variable represent the other unknown?" The answer is yes. The equation will be different, but in the end the answers will be the same. Let s represent the number of strikeouts for Sale, and confirm this.

$$s + (s - 40) = 576 \quad \text{Alternative equation for Example 4}$$

NOW TRY ANSWER

4. Wilson: 34; Brees: 23

OBJECTIVE 5 Solve percent problems.

Recall that percent means “per 100.”

5% means $\frac{5}{100}$ or $5 \cdot \frac{1}{100}$ or $5(0.01)$, all of which equal 0.05.

14% means $\frac{14}{100}$ or $14 \cdot \frac{1}{100}$ or $14(0.01)$, all of which equal 0.14.


**NOW TRY
EXERCISE 5**

In the fall of 2018, there were 96 Introductory Statistics students at a certain community college, an increase of 700% over the number of Introductory Statistics students in the fall of 2010. How many Introductory Statistics students were there in the fall of 2010?

EXAMPLE 5 Solving a Percent Problem

Total annual health expenditures in the United States were about \$3200 billion (or \$3.2 trillion) in 2015. This is an increase of 345% over the total for 1990. What were the approximate total health expenditures in billions of dollars in the United States in 1990? (Data from U.S. Centers for Medicare & Medicaid Services.)

Step 1 **Read** the problem. We are given that the total health expenditures increased by 345% from 1990 to 2015, and \$3200 billion was spent in 2015. We must find the expenditures in 1990.

Step 2 **Assign a variable.** Let x = the total health expenditures for 1990.

$$345\% = 345(0.01) = 3.45,$$

so $3.45x$ represents the increase in expenditures from 1990 to 2015.

Step 3 **Write an equation** from the given information.

$$\begin{array}{rcccl} \text{The expenditures in 1990} & + & \text{the increase} & = & 3200. \\ \downarrow & & \downarrow & & \downarrow \\ x & + & 3.45x & = & 3200 \end{array}$$

Note the x in $3.45x$.

Step 4 **Solve** the equation. (We do so without clearing the decimal.)

$$1x + 3.45x = 3200 \quad \text{Identity property}$$

$$4.45x = 3200 \quad \text{Combine like terms.}$$

$$x \approx 719 \quad \text{Divide by 4.45.}$$

Step 5 **State the answer.** Total health expenditures in the United States for 1990 were about \$719 billion.

Step 6 **Check** that the increase, which is $\$3200 - \$719 = \$2481$ billion, is about 345% of \$719 billion.

$$3.45 \cdot 719 \approx 2481, \quad \text{as required.}$$

NOW TRY 

CAUTION Avoid two common errors that occur when solving problems like the one in Example 5.

- Do not try to find 345% of 3200 and subtract that amount from 3200. The 345% should be applied to *the amount in 1990, not the amount in 2015*.
- In Step 3, do not write the equation as

$$x + 3.45 = 3200. \quad \text{Incorrect}$$

The percent must be multiplied by some number. In this case, the number is the amount spent in 1990, that is, $3.45x$.

NOW TRY ANSWER

5. 12


NOW TRY EXERCISE 6

Sharon received a \$20,000 inheritance from her grandfather. She invested some of the money in an account earning 3% annual interest and the remaining amount in an account earning 2.5% annual interest. If the total annual interest earned is \$575, how much is invested at each rate?

OBJECTIVE 6 Solve investment problems.

The investment problems in this chapter deal with *simple interest*. In most real-world applications, *compound interest* (covered in a later chapter) is used.

EXAMPLE 6 Solving an Investment Problem

Thomas invested \$40,000. He put part of the money in an account paying 2% interest and the remainder into stocks paying 3% interest. The total annual income from these investments was \$1020. How much did he invest at each rate?

Step 1 Read the problem again. We must find the two amounts.

Step 2 Assign a variable.

Let x = the amount invested at 2%;

$40,000 - x$ = the amount invested at 3%.

The formula for interest is $I = prt$. Here the time t is 1 yr. Use a table to organize the given information.

Principal (in dollars)	Rate (as a decimal)	Interest (in dollars)
x	0.02	$0.02x$
$40,000 - x$	0.03	$0.03(40,000 - x)$
40,000		1020

Multiply principal, rate, and time (here, 1 yr) to find interest.

← Total

Step 3 Write an equation. The last column of the table gives the equation.

$$\begin{array}{rcccl}
 \text{Interest at 2\%} & + & \text{interest at 3\%} & = & \text{total interest.} \\
 \downarrow & & \downarrow & & \downarrow \\
 0.02x & + & 0.03(40,000 - x) & = & 1020
 \end{array}$$

Step 4 Solve the equation. Clear the parentheses and then the decimals.

$$0.02x + 1200 - 0.03x = 1020 \quad \text{Distributive property}$$

$$100(0.02x + 1200 - 0.03x) = 100(1020) \quad \text{Multiply by 100.}$$

$$2x + 120,000 - 3x = 102,000 \quad \text{Distributive property; Multiply.}$$

$$-x + 120,000 = 102,000 \quad \text{Combine like terms.}$$

$$-x = -18,000 \quad \text{Subtract 120,000.}$$

$$x = 18,000 \quad \text{Multiply by } -1.$$

Step 5 State the answer. Thomas invested \$18,000 of the money at 2%. At 3%, he invested

$$\$40,000 - \$18,000 = \$22,000.$$

Step 6 Check. Find the annual interest at each rate. The sum of these two amounts should total \$1020.

$$0.02(\$18,000) = \$360 \quad \text{and} \quad 0.03(\$22,000) = \$660$$

$$\$360 + \$660 = \$1020, \quad \text{as required.}$$

NOW TRY ANSWER

6. \$15,000 at 3%; \$5000 at 2.5%

OBJECTIVE 7 Solve mixture problems.

Mixture problems involving rates of concentration can be solved with linear equations.


NOW TRY EXERCISE 7

How many liters of a 20% acid solution must be mixed with 5 L of a 30% acid solution to make a 24% acid solution?

EXAMPLE 7 Solving a Mixture Problem


A chemist must mix 8 L of a 40% acid solution with some 70% solution to make a 50% solution. How much of the 70% solution should be used?

Step 1 Read the problem. The problem asks for the amount of 70% solution to be used.

Step 2 Assign a variable.

Let x = the number of liters of 70% solution.

The information in the problem is illustrated in **FIGURE 5**. We use it to complete a table.

**FIGURE 5**

Liters of Solution	Percent (as a decimal)	Liters of Pure Acid
8	0.40	$0.40(8) = 3.2$
x	0.70	$0.70x$
$8 + x$	0.50	$0.50(8 + x)$

Sum must equal

The values in the last column were found by multiplying the strengths by the numbers of liters.

Step 3 Write an equation using the values in the last column of the table.

$$\begin{array}{rcccl}
 \text{Liters of pure acid} & & & & \\
 \text{in 40\% solution} & + & \text{liters of pure acid} & = & \text{liters of pure acid} \\
 & & \text{in 70\% solution} & & \text{in 50\% solution.} \\
 \downarrow & & \downarrow & & \downarrow \\
 3.2 & + & 0.70x & = & 0.50(8 + x)
 \end{array}$$

Step 4 Solve. $3.2 + 0.70x = 4 + 0.50x$ Distributive property

Move decimal points
1 place to the right.

$$32 + 7x = 40 + 5x$$

Multiply by 10 to clear the decimals.

$$2x = 8$$

Subtract 32 and $5x$.

$$x = 4$$

Divide by 2.

Step 5 State the answer. The chemist should use 4 L of the 70% solution.

Step 6 Check. 8 L of 40% solution plus 4 L of 70% solution is

$$8(0.40) + 4(0.70) = 6 \text{ L of acid.}$$

Similarly, 8 + 4 or 12 L of 50% solution has

$$12(0.50) = 6 \text{ L of acid.}$$

The total amount of pure acid is 6 L both before and after mixing, so the answer checks.

NOW TRY ANSWER

7. $7\frac{1}{2}$ L

PROBLEM-SOLVING HINT When pure water is added to a solution, water is 0% of the chemical (acid, alcohol, etc.). Similarly, pure chemical is 100% chemical.

NOW TRY EXERCISE 8

How much pure antifreeze must be mixed with 3 gal of 30% antifreeze solution to obtain 40% antifreeze solution?

EXAMPLE 8 Solving a Mixture Problem When One Ingredient Is Pure

The octane rating of gasoline is a measure of its antiknock qualities. For a standard fuel, the octane rating is the percent of isooctane. How many liters of pure isooctane should be mixed with 200 L of 94% isooctane, referred to as 94 octane, to obtain a mixture that is 98% isooctane?

Step 1 **Read** the problem. We must find the amount of pure isooctane.

Step 2 **Assign a variable.** Let x = the number of liters of pure (100%) isooctane. Complete a table. Recall that $100\% = 100(0.01) = 1$.

Liters of Solution	Percent (as a decimal)	Liters of Pure Isooctane
x	1	x
200	0.94	$0.94(200)$
$x + 200$	0.98	$0.98(x + 200)$

Step 3 **Write an equation** using the values in the last column of the table.

$$x + 0.94(200) = 0.98(x + 200) \quad \text{Refer to the table.}$$

Step 4 **Solve.** $x + 188 = 0.98x + 196$ Multiply; distributive property

$$100(x + 188) = 100(0.98x + 196) \quad \text{Multiply by 100 to clear the decimal.}$$

$$100x + 18,800 = 98x + 19,600 \quad \text{Distributive property}$$

$$2x = 800 \quad \text{Subtract } 98x \text{ and } 18,800.$$

$$x = 400 \quad \text{Divide by 2.}$$

Step 5 **State the answer.** 400 L of isooctane is needed.

Step 6 **Check** by substituting 400 for x in the equation from Step 3.

$$400 + 0.94(200) \stackrel{?}{=} 0.98(400 + 200) \quad \text{Let } x = 400.$$

$$400 + 188 \stackrel{?}{=} 0.98(600) \quad \text{Multiply. Add.}$$

$$588 = 588 \quad \checkmark \quad \text{True}$$

A true statement results, so the answer checks.

NOW TRY ANSWER

8. $\frac{1}{2}$ gal


NOW TRY 

1.3 Exercises

FOR EXTRA HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check In each of the following, translate part (a) as an expression and translate part (b) as an equation or inequality. Use x to represent the number.

- | | |
|-------------------------------|---------------------------------|
| 1. (a) 15 more than a number | 2. (a) 5 greater than a number |
| (b) 15 is more than a number. | (b) 5 is greater than a number. |
| 3. (a) 8 less than a number | 4. (a) 6 less than a number |
| (b) 8 is less than a number. | (b) 6 is less than a number. |

STUDY SKILLS REMINDER

You will increase your chance of success in this course if you fully utilize your text.

Review Study Skill 1, Using Your Math Text.

5. **Concept Check** A student translated the phrase “the difference of a number x and 7” incorrectly as $7 - x$. **WHAT WENT WRONG?** Give the correct mathematical expression.
6. **Concept Check** A student translated the phrase “the quotient of a number n and 12” incorrectly as $\frac{12}{n}$. **WHAT WENT WRONG?** Give the correct mathematical expression.

Translate each verbal phrase into a mathematical expression using x as the variable. See *Objective 1*.

- | | |
|-------------------------------------------------|--------------------------------------------------------------------|
| 7. Twice a number, decreased by 13 | 8. Triple a number, decreased by 14 |
| 9. 12 increased by four times a number | 10. 15 more than one-half of a number |
| 11. The product of 8 and 16 less than a number | 12. The product of 8 more than a number and 5 less than the number |
| 13. The quotient of three times a number and 10 | 14. The quotient of 9 and five times a nonzero number |

Translate each verbal sentence into an equation, using x as the variable. Then solve the equation. See *Example 1*.

15. The sum of a number and 6 is -31 . Find the number.
16. The sum of a number and -4 is 18. Find the number.
17. If the product of a number and -4 is subtracted from the number, the result is 9 more than the number. Find the number.
18. If the quotient of a number and 6 is added to twice the number, the result is 8 less than the number. Find the number.
19. When $\frac{2}{3}$ of a number is subtracted from 12, the result is 10. Find the number.
20. When 75% of a number is added to 6, the result is 3 more than the number. Find the number.

Determine whether each is an expression or an equation. Simplify any expressions, and solve any equations. See *Example 2*.

- | | |
|-----------------------------------------------------|-----------------------------------------------------|
| 21. $5(x + 3) - 8(2x - 6)$ | 22. $-7(x + 4) + 13(x - 6)$ |
| 23. $5(x + 3) - 8(2x - 6) = 12$ | 24. $-7(x + 4) + 13(x - 6) = 18$ |
| 25. $\frac{1}{2}x - \frac{1}{6}x + \frac{3}{2} - 8$ | 26. $\frac{1}{3}x + \frac{1}{5}x - \frac{1}{2} + 7$ |

Complete the six problem-solving steps to solve each problem.

27. In 2015, the corporations securing the most U.S. patents were IBM and Samsung. Together, the two corporations secured a total of 12,368 patents, with Samsung receiving 2250 fewer patents than IBM. How many patents did each corporation secure? (Data from U.S. Patent and Trademark Office.)

Step 1 Read the problem carefully. We are asked to find _____.

Step 2 Assign a variable. Let x = the number of patents that IBM secured.
Then $x - 2250$ = the number of _____.

Step 3 Write an equation. _____ + _____ = 12,368

Step 4 Solve the equation. x = _____

Step 5 State the answer. IBM secured _____ patents. Samsung secured _____ patents.

Step 6 Check. The number of Samsung patents was _____ fewer than the number of _____.
The total number of patents was $7309 + \text{_____} = \text{_____}$.

28. In 2015, 16.2 million more U.S. residents traveled to Mexico than to Canada. There was a total of 41.2 million U.S. residents traveling to these two countries. How many traveled to each country? (Data from U.S. Department of Commerce.)

Step 1 Read the problem carefully. We are asked to find _____.

Step 2 Assign a variable. Let x = the number of travelers to Mexico (in millions).
Then $x - 16.2$ = the number of _____.

Step 3 Write an equation. _____ + _____ = 41.2


Step 4 Solve the equation. x = _____

Step 5 State the answer. There were _____ travelers to Mexico and _____ travelers to Canada.


Step 6 Check. The number of _____ was _____ more than the number of travelers to _____.
The total number of travelers was $28.7 +$ _____ = _____.

Solve each problem. See Examples 3 and 4.



29. The John Hancock Center in Chicago has a rectangular base. The length of the base measures 65 ft less than twice the width. The perimeter of the base is 860 ft. What are the dimensions of the base?
30. The John Hancock Center tapers as it rises. The top floor is rectangular and has perimeter 520 ft. The width of the top floor measures 20 ft more than one-half its length. What are the dimensions of the top floor?



31. Grant Wood painted his most famous work, *American Gothic*, in 1930 on composition board with perimeter 108.44 in. If the rectangular painting is 5.54 in. taller than it is wide, find the dimensions of the painting. (Data from *The Gazette*.)
32. The perimeter of a certain rectangle is 16 times the width. The length is 12 cm more than the width. Find the length and width of the rectangle.




33. The Bermuda Triangle supposedly causes trouble for aircraft pilots. It has a perimeter of 3075 mi. The shortest side measures 75 mi less than the middle side, and the longest side measures 375 mi more than the middle side. Find the lengths of the three sides.
34. The Vietnam Veterans Memorial in Washington, DC, is in the shape of two sides of an isosceles triangle. If the two walls of equal length were joined by a straight line of 438 ft, the perimeter of the resulting triangle would be 931.5 ft. Find the lengths of the two walls. (Data from pamphlet obtained at Vietnam Veterans Memorial.)



35. The two companies with top revenues in the Fortune 500 list for 2016 were Walmart and Berkshire Hathaway. Their revenues together totaled \$710.5 billion. Berkshire Hathaway revenues were \$263.3 billion less than those of Walmart. What were the revenues of each corporation? (Data from *Fortune* magazine.)
36. On a particular day in 2017, two of the longest-running Broadway shows were *The Phantom of the Opera* and *Chicago*. Together, there were 20,721 performances of these two shows during their Broadway runs. There were 3675 fewer performances of *Chicago* than of *The Phantom of the Opera*. How many performances were there of each show? (Data from The Broadway League.)
37. Galileo Galilei conducted experiments involving Italy's famous Leaning Tower of Pisa to investigate the relationship between an object's speed of fall and its weight. The Leaning Tower is 804 ft shorter than the Eiffel Tower in Paris, France. The two towers have a total height of 1164 ft. How tall is each tower? (Data from *The World Almanac and Book of Facts*.)



38. In 2017, the Los Angeles Dodgers and the New York Yankees had the highest payrolls in Major League Baseball. The Dodgers' payroll was \$40.6 million more than the Yankees' payroll, and the two payrolls totaled \$443.6 million. What was the payroll for each team? (Data from www.stevetheump.com)
39. In 2017, two popular brands on Instagram were National Geographic and Nike. National Geographic had 8.0 million more followers than Nike. Together, the two brands had 158.8 million followers. How many followers did each brand have? (Data from Iconosquare.)
40. Ted Williams and Rogers Hornsby were two great hitters in Major League Baseball. Together, they had 5584 hits in their careers. Hornsby had 276 more hits than Williams. How many hits did each have? (Data from Neft, D. S., and R. M. Cohen, *The Sports Encyclopedia: Baseball*, St. Martins Griffin, New York.)



Rogers Hornsby

Solve each percent problem. See Example 5.

41. In 2015, the number of graduating seniors taking the ACT exam was 1,924,436. In 2011, a total of 1,617,173 graduating seniors took the exam. By what percent did the number increase over this period of time, to the nearest tenth of a percent? (Data from ACT.)
42. The U.S. population reached 321 million in 2015 and is expected to reach 347 million by the year 2025. What percent increase is expected over this time period, to the nearest tenth of a percent? (Data from www.census.gov)
43. In 2013, mobile messaging service WhatsApp had 300 million monthly active users worldwide. By 2017, this number had increased 333.3%. To the nearest million, what was the number of WhatsApp monthly active users worldwide in 2017? (Data from WhatsApp.)
44. On Major League Baseball Opening Day 2016, the payroll of the Cleveland Indians was \$87.2 million. In 2017, it had increased 44.3%. To the nearest tenth of a million, what was the opening-day payroll of the Indians for 2017? (Data from The Associated Press.)

45. In 2017, the average cost of a traditional Thanksgiving dinner for 10, featuring turkey, stuffing, cranberries, pumpkin pie, and trimmings, was \$49.12, a decrease of 1.48% from the cost in 2016. What was the average cost, to the nearest cent, of a traditional Thanksgiving dinner in 2016? (Data from American Farm Bureau.)



46. In 2016, the average cost of a Thanksgiving turkey was \$22.74. This price had decreased by 1.58% in 2017. What was the average cost of a Thanksgiving turkey in 2017? (Data from American Farm Bureau.)
47. At the end of a day, Lawrence found that the total cash register receipts at the motel where he works were \$2725. This included the 9% sales tax charged. Find the amount of tax.
48. David sold his house for \$159,000. He got this amount knowing that he would have to pay a 6% commission to his agent. What amount did he have after the agent was paid?
49. **Concept Check** Why is it impossible to mix candy worth \$4 per lb and candy worth \$5 per lb to obtain a final mixture worth \$6 per lb?
50. **Concept Check** Write an equation based on the following problem, solve the equation, and explain why the problem has no solution.

How much 30% acid should be mixed with 15 L of 50% acid to obtain a mixture that is 60% acid?

Solve each investment problem. See Example 6.

51. Mario earned \$12,000 last year giving tennis lessons. He invested part of the money at 3% simple interest and the rest at 4%. In one year, he earned a total of \$440 in interest. How much did he invest at each rate?
52. Sheryl won \$60,000 on a slot machine in Las Vegas. She invested part of the money at 2% simple interest and the rest at 3%. In one year, she earned a total of \$1600 in interest. How much was invested at each rate?

Principal (in dollars)	Rate (as a decimal)	Interest (in dollars)
x	0.03	
	0.04	

Principal (in dollars)	Rate (as a decimal)	Interest (in dollars)
x	0.02	

53. Michelle invested some money at 4.5% simple interest and \$1000 less than twice this amount at 3%. Her total annual income from the interest was \$1020. How much was invested at each rate?
54. Piotr invested some money at 3.5% simple interest, and \$5000 more than three times this amount at 4%. He earned \$1440 in annual interest. How much did he invest at each rate?
55. Dan has invested \$12,000 in bonds paying 6%. How much additional money should he invest in a certificate of deposit paying 3% simple interest so that the total return on the two investments will be 4%?
56. Mona received a year-end bonus of \$17,000 from her company and invested the money in an account paying 6.5%. How much additional money should she deposit in an account paying 5% so that the return on the two investments will be 6%?

Solve each problem. See Examples 7 and 8.

57. How many liters of a 10% acid solution must be mixed with 10 L of a 4% solution to obtain a 6% solution?

Liters of Solution	Percent (as a decimal)	Liters of Pure Acid
10	0.04	
x	0.10	
	0.06	

58. How many liters of a 14% alcohol solution must be mixed with 20 L of a 50% solution to obtain a 30% solution?

Liters of Solution	Percent (as a decimal)	Liters of Pure Alcohol
x	0.14	
	0.50	

59. In a chemistry class, 12 L of a 12% alcohol solution must be mixed with a 20% solution to obtain a 14% solution. How many liters of the 20% solution are needed?

60. How many liters of a 10% alcohol solution must be mixed with 40 L of a 50% solution to obtain a 40% solution?

61. How much pure dye must be added to 4 gal of a 25% dye solution to increase the solution to 40%? (Hint: Pure dye is 100% dye.)

62. How much water must be added to 6 gal of a 4% insecticide solution to reduce the concentration to 3%? (Hint: Water is 0% insecticide.)

63. Randall wants to mix 50 lb of nuts worth \$2 per lb with some nuts worth \$6 per lb to make a mixture worth \$5 per lb. How many pounds of \$6 nuts must he use?

Pounds of Nuts	Cost per Pound (in dollars)	Total Cost (in dollars)

64. Lee Ann wants to mix tea worth 2¢ per oz with 100 oz of tea worth 5¢ per oz to make a mixture worth 3¢ per oz. How much 2¢ tea should be used?

Ounces of Tea	Cost per Ounce (in dollars)	Total Cost (in dollars)



RELATING CONCEPTS For Individual or Group Work (Exercises 65–68)

Consider each problem.

Problem A

Jack has \$800 invested in two accounts. One pays 3% interest per year and the other pays 6% interest per year. The amount of yearly interest is the same as he would get if the entire \$800 was invested at 5.25%. How much does he have invested at each rate?

Problem B

Jill has 800 L of acid solution. She obtained it by mixing some 3% acid with some 6% acid. Her final mixture of 800 L is 5.25% acid. How much of each of the 3% and 6% solutions did she use to obtain her final mixture?

In Problem A, let x represent the amount invested at 3% interest, and in Problem B, let y represent the amount of 3% acid used. **Work Exercises 65–68 in order.**

65. (a) Write an expression in x that represents the amount of money Jack invested at 6% in Problem A.
- (b) Write an expression in y that represents the amount of 6% acid solution Jill used in Problem B.

- 66. (a)** Write expressions that represent the amount of interest Jack earns per year at 3% and at 6%.
- (b)** Write expressions that represent the amount of pure acid in Jill’s 3% and 6% acid solutions.
- 67. (a)** The sum of the two expressions in **Exercise 66(a)** must equal the total amount of interest earned in one year. Write an equation representing this fact.
- (b)** The sum of the two expressions in **Exercise 66(b)** must equal the amount of pure acid in the final mixture. Write an equation representing this fact.
- 68. (a)** Solve Problem A.
- (b)** Solve Problem B.
- (c)** Explain the similarities between the processes used in solving Problems A and B.

1.4 Further Applications of Linear Equations

OBJECTIVES

- 1** Solve problems about different denominations of money.
- 2** Solve problems about uniform motion.
- 3** Solve problems about angles.
- 4** Solve problems about consecutive integers.

VOCABULARY

- consecutive integers
- consecutive even integers
- consecutive odd integers

OBJECTIVE 1 Solve problems about different denominations of money.

PROBLEM-SOLVING HINT In problems involving money, use the following basic fact.

$$\begin{array}{l} \text{number of monetary} \\ \text{units of the same kind} \end{array} \times \text{denomination} = \begin{array}{l} \text{total monetary} \\ \text{value} \end{array}$$

Examples: 30 dimes have a monetary value of $30(\$0.10) = \3.00 .
 15 five-dollar bills have a value of $15(\$5) = \75 .

EXAMPLE 1 Solving a Money Denomination Problem

For a bill totaling \$5.65, a cashier received 25 coins consisting of nickels and quarters. How many of each denomination of coin did the cashier receive?

Step 1 Read the problem. We must find the number of nickels and the number of quarters the cashier received.

Step 2 Assign a variable.

Let x = the number of nickels.

Then $25 - x$ = the number of quarters.

	Conect Number of Coins	Denomination (in dollars)	Value (in dollars)
Nickels	x	0.05	$0.05x$
Quarters	$25 - x$	0.25	$0.25(25 - x)$
			5.65

Organize the information in a table.

← Total

Step 3 Write an equation. Use the values from the last column of the table.

$$0.05x + 0.25(25 - x) = 5.65$$


NOW TRY EXERCISE 1

Steven has a collection of 52 coins worth \$3.70. His collection contains only dimes and nickels. How many of each type of coin does he have?

Step 4 Solve. Clear the parentheses and then the decimals.

$$0.05x + 0.25(25 - x) = 5.65$$

Equation from Step 3

$$0.05x + 6.25 - 0.25x = 5.65$$

Distributive property

$$100(0.05x + 6.25 - 0.25x) = 100(5.65)$$

Multiply by 100.

$$5x + 625 - 25x = 565$$

Distributive property; Multiply.

Move decimal points 2 places to the right.

$$-20x = -60$$

Subtract 625. Combine like terms.

$$x = 3$$

Divide by -20 .

Step 5 State the answer. There are 3 nickels and $25 - 3 = 22$ quarters.

Step 6 Check. The cashier has $3 + 22 = 25$ coins. The value is

$$\$0.05(3) + \$0.25(22) = \$5.65, \text{ as required.}$$

NOW TRY 

CAUTION Be sure that your answer is reasonable when working problems like Example 1. Because we are dealing with a number of coins, the correct answer can be neither negative nor a fraction.

OBJECTIVE 2 Solve problems about uniform motion.

PROBLEM-SOLVING HINT Uniform motion problems use the distance formula $d = rt$. When rate (or speed) is given in miles per hour, time must be given in hours. Draw a sketch to illustrate what is happening. Make a table to summarize given information.

EXAMPLE 2 Solving a Motion Problem (Opposite Directions)

Two cars leave the same place at the same time, one going east and the other west. The eastbound car averages 40 mph, while the westbound car averages 50 mph. In how many hours will they be 300 mi apart?

Step 1 Read the problem. We are looking for the time it takes for the two cars to be 300 mi apart.

Step 2 Assign a variable. A sketch shows what is happening in the problem. The cars are going in opposite directions. See FIGURE 6.




FIGURE 6

Let x = the time traveled by each car.

	Rate	Time	Distance
Eastbound Car	40	x	$40x$
Westbound Car	50	x	$50x$
			300

Fill in each distance by multiplying rate by time, using the formula $d = rt$, or $rt = d$. The sum of the two distances is 300.

NOW TRY ANSWER

1. 22 dimes; 30 nickels

NOW TRY EXERCISE 2

Two trains leave a city traveling in opposite directions. One travels at a rate of 80 km per hr and the other at a rate of 75 km per hr. How long will it take before they are 387.5 km apart?

Step 3 Write an equation. $40x + 50x = 300$

Step 4 Solve. $90x = 300$ Combine like terms.

$$x = \frac{300}{90}$$

Divide by 90.

$$x = \frac{10}{3}$$

Lowest terms

Step 5 State the answer. The cars travel $\frac{10}{3} = 3\frac{1}{3}$ hr, or 3 hr, 20 min.

Step 6 Check. The eastbound car traveled $40\left(\frac{10}{3}\right) = \frac{400}{3}$ mi. The westbound car traveled $50\left(\frac{10}{3}\right) = \frac{500}{3}$ mi, for a total distance of $\frac{400}{3} + \frac{500}{3} = \frac{900}{3}$, or 300 mi, as required.

NOW TRY

CAUTION It is a common error to write 300 as the distance traveled by each car in **Example 2**. The **total** distance traveled is 300 mi.

As in **Example 2**, in general, the equation for a problem involving motion in *opposite* directions is of the following form.

$$\text{partial distance} + \text{partial distance} = \text{total distance}$$

EXAMPLE 3 Solving a Motion Problem (Same Direction)

Geoff can bike to work in $\frac{3}{4}$ hr. When he takes the bus, the trip takes $\frac{1}{4}$ hr. If the bus travels 20 mph faster than Geoff rides his bike, how far is it to his workplace?

Step 1 Read the problem. We must find the distance between Geoff's home and his workplace.

Step 2 Assign a variable. Make a sketch to show what is happening. See **FIGURE 7**.



FIGURE 7

The problem asks for a *distance*, but it is easier here to let x be Geoff's *rate* when he rides his bike to work. Then the rate of the bus is $(x + 20)$ mph.

For the trip by bike, $d = rt = x \cdot \frac{3}{4} = \frac{3}{4}x$.

For the trip by bus, $d = rt = (x + 20) \cdot \frac{1}{4} = \frac{1}{4}(x + 20)$.

	Rate	Time	Distance
Bike	x	$\frac{3}{4}$	$\frac{3}{4}x$
Bus	$x + 20$	$\frac{1}{4}$	$\frac{1}{4}(x + 20)$

← Same distance

Step 3 Write an equation.

$$\frac{3}{4}x = \frac{1}{4}(x + 20)$$

The distance is the same in each case.

NOW TRY ANSWER

2. $2\frac{1}{2}$ hr

NOW TRY
EXERCISE 3

Michael can drive to work in $\frac{1}{2}$ hr. When he rides his bicycle, it takes $1\frac{1}{2}$ hours. If his average rate while driving to work is 30 mph faster than his rate while bicycling to work, determine the distance that he lives from work.

Step 4 Solve.

$$\frac{3}{4}x = \frac{1}{4}(x + 20) \quad \text{Equation from Step 3}$$

$$\frac{3}{4}x = \frac{1}{4}x + 5 \quad \text{Distributive property}$$

$$4\left(\frac{3}{4}x\right) = 4\left(\frac{1}{4}x + 5\right) \quad \text{Multiply by 4 to clear the fractions.}$$

$$3x = x + 20 \quad \text{Multiply; } 4\left(\frac{1}{4}\right) = 1 \text{ and } 1x = x$$

$$2x = 20 \quad \text{Subtract } x.$$

$$x = 10 \quad \text{Divide by 2.}$$

Step 5 State the answer. The required distance is

$$d = \frac{3}{4}x = \frac{3}{4}(10) = \frac{30}{4} = 7.5 \text{ mi.}$$

Step 6 Check by finding the distance using

$$d = \frac{1}{4}(x + 20) = \frac{1}{4}(10 + 20) = \frac{30}{4} = 7.5 \text{ mi.}$$

The same distance results.

NOW TRY

As in **Example 3**, the equation for a problem involving motion in the *same* direction is often of the following form.

$$\text{one distance} = \text{other distance}$$

PROBLEM-SOLVING HINT In **Example 3**, it was easier to let the variable represent a quantity other than the one that we were asked to find. It takes practice to learn when this approach works best.

OBJECTIVE 3 Solve problems about angles.

An important result of Euclidean geometry (the geometry of the Greek mathematician Euclid) is as follows.

The sum of the angle measures of any triangle is 180° .

EXAMPLE 4 Finding Angle Measures

Find the value of x , and determine the measure of each angle in **FIGURE 8**.

Step 1 Read the problem. We are asked to find the measure of each angle.

Step 2 Assign a variable.

Let x = the measure of one angle.

Step 3 Write an equation. The sum of the three measures shown in the figure must be 180° .

$$x + (x + 20) + (210 - 3x) = 180$$

Measures are in degrees.

Step 4 Solve.

$$-x + 230 = 180$$

Combine like terms.

$$-x = -50$$

Subtract 230.

$$x = 50$$

Multiply by -1 .





FIGURE 8

NOW TRY ANSWER

3. 22.5 mi


NOW TRY EXERCISE 4

Find the value of x , and determine the measure of each angle.



Step 5 State the answer. One angle measures 50° . The other two angles measure

$$x + 20 = 50 + 20 = 70^\circ$$

and

$$210 - 3x = 210 - 3(50) = 60^\circ.$$

Step 6 Check. Because $50^\circ + 70^\circ + 60^\circ = 180^\circ$ is true, the answer is correct.

NOW TRY 

OBJECTIVE 4 Solve problems about consecutive integers.

Recall that the set of integers is $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$. **Consecutive integers** are integers that are next to each other on a number line. Two consecutive integers differ by 1. Some examples of consecutive integers are 4, 5 and $-6, -5$.

Consecutive even integers always differ by 2. Examples are 4, 6 and $-4, -2$. Similarly, **consecutive odd integers** also differ by 2, such as 1, 3 and $-5, -3$.

PROBLEM-SOLVING HINT If $x =$ the lesser (least) integer in a consecutive integer problem, then the following apply.

- For two consecutive integers, use $x, x + 1$.
- For three consecutive integers, use $x, x + 1, x + 2$.
- For two consecutive even or odd integers, use $x, x + 2$.
- For three consecutive even or odd integers, use $x, x + 2, x + 4$.


NOW TRY EXERCISE 5

Find three consecutive integers such that the sum of the first and second is 43 less than three times the third.

EXAMPLE 5 Solving a Consecutive Integer Problem

Find three consecutive integers such that the sum of the first and third, increased by 3, is 50 more than the second.

Step 1 Read the problem. We are told to find three consecutive integers satisfying the given conditions.

Step 2 Assign a variable.

Let $x =$ the least of the three consecutive integers.

Then $x + 1 =$ the second (middle) integer,

and $x + 2 =$ the greatest of the three consecutive integers.

Step 3 Write an equation.

$$\begin{array}{ccccccc} \text{Sum of the} & & \text{increased} & & \text{is} & & \text{50 more than} \\ \text{first and third,} & & \text{by 3,} & & & & \text{the second.} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ x + (x + 2) & & + 3 & & = & & (x + 1) + 50 \end{array}$$

Step 4 Solve.

$$2x + 5 = x + 51 \quad \text{Combine like terms.}$$

$$x = 46 \quad \text{Subtract } x \text{ and } 5.$$

Step 5 State the answer. The solution is 46, so the first integer is $x = 46$, the second is $46 + 1 = 47$, and the third is $46 + 2 = 48$. The three integers are 46, 47, and 48.

Step 6 Check. The sum of the first and third is $46 + 48 = 94$. If this is increased by 3, the result is $94 + 3 = 97$, which is indeed 50 more than the second (47). The answer is correct.

NOW TRY 

NOW TRY ANSWERS

4. $41^\circ, 52^\circ, 87^\circ$

5. 38, 39, 40

1.4 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Solve each problem.

1. What amount of money is found in a coin hoard containing 12 dimes and 18 quarters?
2. The distance between Cape Town, South Africa, and Miami is 7700 mi. If a jet averages 550 mph between the two cities, what is its travel time in hours?
3. Tri traveled from Chicago to Des Moines, a distance of 300 mi, in 10 hr. What was his rate in miles per hour?
4. A square has perimeter 160 in. What would be the perimeter of an equilateral triangle whose sides each measure the same length as the side of the square?
5. An equilateral triangle has perimeter 27 in. What would be the area of a square whose sides each measure the same length as the side of the equilateral triangle?
6. A circle has area 25π ft². What would be the perimeter of a square whose sides each measure the same length as the radius of the circle?
7. **Concept Check** A student was asked to find the measures of the angles of a triangle with degree measures that are consecutive integers. Letting x represent the measure of the smallest angle, the student wrote the following equation.

$$x + (x + 1) + (x + 2) = 90 \quad \text{Incorrect}$$

WHAT WENT WRONG? Write the correct equation and solve the problem.

8. **Concept Check** A student was asked to find the number of nickels and dimes that have a value of \$1.45. The total number of coins is 21. Letting x represent the number of nickels, the student wrote the following equation.

$$0.05x + 0.10(x - 21) = 1.45 \quad \text{Incorrect}$$

WHAT WENT WRONG? Write the correct equation and solve the problem.

Solve each problem. See Example 1.

9. Otis has a box of coins that he uses when he plays poker with his friends. The box contains 44 coins, consisting of pennies, dimes, and quarters. The number of pennies is equal to the number of dimes. The total value is \$4.37. How many of each denomination of coin does he have in the box?

Number of Coins	Denomination (in dollars)	Value (in dollars)
x	0.01	$0.01x$
x		
	0.25	
XXXXX	XXXXXX	4.37 ← Total

10. Nana found some coins under her sofa pillows. There were equal numbers of nickels and quarters and twice as many half-dollars as quarters. If she found \$2.60 in all, how many of each denomination of coin did she find?

Number of Coins	Denomination (in dollars)	Value (in dollars)
x	0.05	$0.05x$
x		
$2x$	0.50	
XXXXX	XXXXXX	2.60 ← Total

11. In Canada, \$1 coins are called “loonies” because they have a picture of a loon on the reverse, and \$2 coins are called “toonies.” When Marissa returned home to San Francisco from a trip to Vancouver, she found that she had acquired 37 of these coins, with a total value of 51 Canadian dollars. How many coins of each denomination did she have?



12. Ahmad works at an ice cream shop. At the end of his shift, he counted the bills in his cash drawer and found 119 bills with a total value of \$347. If all of the bills are \$5 bills and \$1 bills, how many of each denomination were in his cash drawer?
13. Hussein collects U.S. gold coins. He has a collection of 41 coins. Some are \$10 coins, and the rest are \$20 coins. If the face value of the coins is \$540, how many of each denomination does he have?



14. In the 19th century, the United States minted two-cent and three-cent pieces. Frances has three times as many three-cent pieces as two-cent pieces, and the face value of these coins is \$2.42. How many of each denomination does she have?
15. In 2018, general admission to the Art Institute of Chicago cost \$25 for adults and \$19 for children and seniors. If \$32,972 was collected from the sale of 1460 general admission tickets, how many adult tickets were sold? (Data from www.artic.edu)
16. For a high school production of *Hello, Dolly!*, student tickets cost \$5 each and nonstudent tickets cost \$8 each. If 480 tickets were sold and a total of \$2895 was collected, how many tickets of each type were sold?

For each event, find the rate on the basis of the information provided. Use a calculator and round answers to the nearest hundredth. All events were at the 2016 Summer Olympics in Rio de Janeiro, Brazil. (Data from *The World Almanac and Book of Facts*.)



	Event	Winner	Time
17.	100-m hurdles, women	Brianna Rollins, United States	12.48 sec
18.	400-m hurdles, women	Dalilah Muhammad, United States	53.13 sec
19.	200-m run, men	Usain Bolt, Jamaica	19.78 sec
20.	400-m run, men	Wayde van Niekerk, South Africa	43.03 sec

Solve each problem. See Examples 2 and 3.

21. Two steamers leave a port on a river at the same time, traveling in opposite directions. Each is traveling 22 mph. How long will it take for them to be 110 mi apart?
22. A train leaves Kansas City, Kansas, and travels north at 85 km per hr. Another train leaves at the same time and travels south at 95 km per hr. How long will it take before they are 315 km apart?

	Rate	Time	Distance
First Steamer		t	
Second Steamer	22		
XXXXXXXXXX	XX	XX	110

	Rate	Time	Distance
First Train	85	t	
Second Train			
XXXXXXXXXX	XX	XX	315

23. Mulder and Scully are driving to Georgia to investigate “Big Blue,” a giant reptile reported in one of the local lakes. Mulder leaves the office at 8:30 A.M. averaging 65 mph. Scully leaves at 9:00 A.M., following the same path and averaging 68 mph. At what time will Scully catch up with Mulder?
24. Lois and Clark are covering separate stories and have to travel in opposite directions. Lois leaves the *Daily Planet* building at 8:00 A.M. and travels at 35 mph. Clark leaves at 8:15 A.M. and travels at 40 mph. At what time will they be 140 mi apart?

	Rate	Time	Distance
Mulder			
Scully			

	Rate	Time	Distance
Lois			
Clark			

25. It took Charmaine 3.6 hr to drive to her mother's house on Saturday morning for a weekend visit. On her return trip on Sunday night, traffic was heavier, so the trip took her 4 hr. Her average rate on Sunday was 5 mph slower than on Saturday. What was her average rate on Sunday?

	Rate	Time	Distance
Saturday			
Sunday			


26. Sharon commutes to her office by train. When she walks to the train station, it takes her 40 min. When she rides her bike, it takes her 12 min. Her average walking rate is 7 mph less than her average biking rate. Find the distance from her house to the train station.

	Rate	Time	Distance
Walking			
Biking			


27. Johnny leaves Memphis to visit his cousin Anne, who lives in the town of Hornsby, Tennessee, 80 mi away. He travels at an average rate of 50 mph. One-half hour later, Anne leaves to visit Johnny, traveling at an average rate of 60 mph. How long after Anne leaves will it be before they meet?
28. On an automobile trip, Laura maintained a steady rate for the first two hours. Rush-hour traffic slowed her rate by 25 mph for the last part of the trip. The entire trip, a distance of 125 mi, took $2\frac{1}{2}$ hr. What was her rate during the first part of the trip?

Find the measure of each angle in the triangles shown. See Example 4.


29.




30.




31.




32.





33.





34.





In the following problems, the angles marked with variable expressions are **vertical angles**. It is shown in geometry that vertical angles have equal measures. Find the measure of each angle.



Two angles whose sum is 90° are **complementary angles**. Find the measures of the complementary angles shown in each figure.



Two angles whose sum is 180° are **supplementary angles**. Find the measures of the supplementary angles shown in each figure.



Solve each problem involving consecutive integers. See Example 5.


- 41. Find three consecutive integers such that the sum of the first and twice the second is 17 more than twice the third.
- 42. Find three consecutive integers such that the sum of the first and twice the third is 39 more than twice the second.
- 43. Find four consecutive integers such that the sum of the first three is 54 more than the fourth.
- 44. Find four consecutive integers such that the sum of the last three is 86 more than the first.
- 45. If I add my current age to the age I will be next year on this date, the sum is 103 yr. How old will I be 10 yr from today?
- 46. If I add my current age to the age I will be next year on this date, the sum is 129 yr. How old will I be 5 yr from today?

October						
Sun	Mon	Tue	Wed	Thu	Fri	Sat
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			


- 47. Find three consecutive *even* integers such that the sum of the least integer and the middle integer is 26 more than the greatest integer.
- 48. Find three consecutive *even* integers such that the sum of the least integer and the greatest integer is 12 more than the middle integer.
- 49. Find three consecutive *odd* integers such that the sum of the least integer and the middle integer is 19 more than the greatest integer.
- 50. Find three consecutive *odd* integers such that the sum of the least integer and the greatest integer is 13 more than the middle integer.

RELATING CONCEPTS For Individual or Group Work (Exercises 51–54)

Consider the following two figures. Work Exercises 51–54 in order.





51. Solve for the measures of the unknown angles in **FIGURE A**. (*Hint*: What is the sum of the angle measures of any triangle?)
52. Solve for the measure of the unknown angle marked y° in **FIGURE B**. (*Hint*: The angles are supplementary angles.)
53. Add the measures of the two angles found in **Exercise 51**. How does the sum compare to the measure of the angle found in **Exercise 52**?
54. Based on the answers to **Exercises 51–53**, make a conjecture (an educated guess) about the relationship among the angles marked ①, ②, and ③ in the figure below.



SUMMARY EXERCISES Applying Problem-Solving Techniques

Solve each problem.

1. The length of a rectangle is 3 in. more than its width. If the length were decreased by 2 in. and the width were increased by 1 in., the perimeter of the resulting rectangle would be 24 in. Find the dimensions of the original rectangle.
2. A farmer wishes to enclose a rectangular region with 210 m of fencing in such a way that the length is twice the width and the region is divided into two equal parts, as shown in the figure. What length and width should be used?




3. After a discount of 28.57%, the sale price for a Samsung Galaxy tablet computer was \$199.99. What was the regular price of the tablet computer to the nearest cent? (Data from www.amazon.com)
4. An electronics store offered a new HD television for \$255, which was the sale price after the regular price was discounted 40%. What was the regular price?
5. An amount of money is invested at 4% annual simple interest, and twice that amount is invested at 5%. The total annual interest is \$112. How much is invested at each rate?
6. An amount of money is invested at 3% annual simple interest, and \$2000 more than that amount is invested at 4%. The total annual interest is \$920. How much is invested at each rate?
7. Russell Westbrook of the Oklahoma City Thunder and James Harden of the Houston Rockets were the leading scorers in the NBA for the 2016–2017 regular season. Together they scored 4914 points, with Harden scoring 202 fewer points than Westbrook. How many points did each player score? (Data from stats.nba.com)


8. In 2016, the number of films released in U.S. and Canadian movie theaters was 161 more than the number of films released in 2009. A total of 1275 films were released in both years. How many films were released each year? (Data from comScore—Box Office Essentials.)



9. Atlanta and Cincinnati are 440 mi apart. John leaves Cincinnati, driving toward Atlanta at an average rate of 60 mph. Pat leaves Atlanta at the same time, driving toward Cincinnati in her antique auto, averaging 28 mph. How long will it take them to meet?





10. Joshua has a sheet of tin 12 cm by 16 cm. He plans to make a box by cutting equal squares out of each of the four corners and folding up the remaining edges. How large a square should he cut so that the finished box will have a length that is 5 cm less than twice the width?



11. A pharmacist has 20 L of a 10% drug solution. How many liters of 5% solution must be added to obtain a mixture that is 8%?
12. A certain metal is 20% tin. How many kilograms of this metal must be mixed with 80 kg of a metal that is 70% tin to obtain a metal that is 50% tin?

Kilograms of Metal	Percent (as a decimal)	Kilograms of Tin
	0.20	
	0.50	

13. A cashier has a total of 126 bills in fives and tens. The total value of the money is \$840. How many of each denomination of bill does he have?
14. The top-grossing domestic movie in 2016 was *Finding Dory*. On opening weekend, one theater showing this movie took in \$20,520 by selling a total of 2460 tickets, some at \$9 and the rest at \$7. How many tickets were sold at each price? (Data from comScore, Inc.)
15. Find the measure of each angle.
16. Find the measure of each marked angle.



17. The sum of the least and greatest of three consecutive integers is 32 more than the middle integer. What are the three integers?
18. If the lesser of two consecutive odd integers is doubled, the result is 7 more than the greater of the two integers. Find the two integers.
19. The perimeter of a triangle is 34 in. The middle side is twice as long as the shortest side. The longest side is 2 in. less than three times the shortest side. Find the lengths of the three sides.
20. The perimeter of a rectangle is 43 in. more than the length. The width is 10 in. Find the length of the rectangle.

1.5 Linear Inequalities in One Variable

OBJECTIVES

- 1 Graph intervals on a number line.
- 2 Solve linear inequalities using the addition property.
- 3 Solve linear inequalities using the multiplication property.
- 4 Solve linear inequalities with three parts.
- 5 Solve applied problems using linear inequalities.

VOCABULARY

- inequality
- interval
- linear inequality in one variable
- equivalent inequalities
- three-part inequality

An **inequality** consists of algebraic expressions related by one of the following symbols.

$<$	“is less than”	\leq	“is less than or equal to”
$>$	“is greater than”	\geq	“is greater than or equal to”

These symbols are read as shown when the inequality is read from left to right.

We **solve an inequality** by finding all real number solutions for it. For example, the solution set of $x \leq 2$ includes *all* real numbers that are less than or equal to 2, not just the integers less than or equal to 2.

OBJECTIVE 1 Graph intervals on a number line.

We graph all the real numbers satisfying $x \leq 2$ by placing a square bracket at 2 and drawing an arrow to the left to represent the fact that all numbers less than 2 are also part of the graph. See **FIGURE 9**.




FIGURE 9 The interval $(-\infty, 2]$

The set of numbers less than or equal to 2 is an example of an **interval** on a number line. To write intervals, we use **interval notation**, which includes the **infinity symbols**, ∞ or $-\infty$. We write the interval of *all* numbers less than or equal to 2 as $(-\infty, 2]$. The negative infinity symbol $-\infty$ does not indicate a number, but shows that the interval includes all real numbers less than 2. Both on the number line and in interval notation, the square bracket indicates that 2 is included in the solution set.

Remember the following important concepts regarding interval notation.

- A parenthesis indicates that an endpoint is *not* included.
- A square bracket indicates that an endpoint is included.
- A parenthesis is always used next to an infinity symbol, $-\infty$ or ∞ .
- The set of real numbers is written in interval notation as $(-\infty, \infty)$.

EXAMPLE 1 Using Interval Notation

Graph each inequality, and write it using interval notation.

(a) $x > -5$

This statement says that x can represent any number greater than -5 , but x cannot equal -5 . We graph it by placing a parenthesis at -5 and drawing an arrow to the right, as in **FIGURE 10**. The parenthesis at -5 shows that -5 is *not* part of the graph. This interval is written $(-5, \infty)$.




FIGURE 10 The interval $(-5, \infty)$

NOW TRY EXERCISE 1

Graph each inequality, and write it using interval notation.

- (a) $x < -1$
- (b) $-4 \leq x < 2$

(b) $-1 \leq x < 3$

This statement is read “ -1 is less than or equal to x **and** x is less than 3 .” We want the set of numbers that are *between* -1 and 3 , with -1 included and 3 excluded. We use a bracket at -1 because it is part of the graph and a parenthesis at 3 because it is not part of the graph. See **FIGURE 11**. In interval notation, we write $[-1, 3)$.




FIGURE 11 The interval $[-1, 3)$

NOW TRY

▼ Summary of Types of Intervals

Type of Interval	Set-Builder Notation	Graph	Interval Notation
Open interval	$\{x a < x < b\}$		(a, b)
Closed interval	$\{x a \leq x \leq b\}$		$[a, b]$
Half-open (or half-closed) interval	$\{x a \leq x < b\}$		$[a, b)$
	$\{x a < x \leq b\}$		$(a, b]$
Disjoint interval*	$\{x x < a \text{ or } x > b\}$		$(-\infty, a) \cup (b, \infty)$
Infinite interval	$\{x x > a\}$		(a, ∞)
	$\{x x \geq a\}$		$[a, \infty)$
	$\{x x < a\}$		$(-\infty, a)$
	$\{x x \leq a\}$		$(-\infty, a]$
	$\{x x \text{ is a real number}\}$		$(-\infty, \infty)$

CAUTION A parenthesis is *always* used next to an infinity symbol, $-\infty$ or ∞ .

Solving inequalities is similar to solving equations.

Linear Inequality in One Variable

A **linear inequality in one variable** (here x) is an inequality that can be written in the form


$$ax + b < 0, \quad ax + b \leq 0, \quad ax + b > 0, \quad \text{or} \quad ax + b \geq 0,$$

where a and b are real numbers and $a \neq 0$.

Examples: $x + 5 < 0$, $x - 3 \geq 5$, $2x + 5 \leq 10$, $x > -1$

Linear inequalities in one variable

NOW TRY ANSWERS



*We will work with disjoint intervals later when we study *set operations* and *compound inequalities*.

OBJECTIVE 2 Solve linear inequalities using the addition property.

We solve an inequality by finding all numbers that make the inequality true. Usually, an inequality has an infinite number of solutions. These solutions, like solutions of equations, are found by producing a series of simpler equivalent inequalities. **Equivalent inequalities** are inequalities with the same solution set.

We use two important properties to produce equivalent inequalities.

Addition Property of Inequality

If a , b , and c represent real numbers, then the inequalities

$$a < b \quad \text{and} \quad a + c < b + c \quad \text{are equivalent.}^*$$

That is, the same number may be added to each side of an inequality without changing the solution set.

*This also applies to $a \leq b$, $a > b$, and $a \geq b$.

**NOW TRY EXERCISE 2**

Solve $x - 10 > -7$, and graph the solution set.

EXAMPLE 2 Using the Addition Property of Inequality

Solve $x - 7 < -12$, and graph the solution set.

$$x - 7 < -12$$

$$x - 7 + 7 < -12 + 7 \quad \text{Add 7.}$$

$$x < -5 \quad \text{Combine like terms.}$$

CHECK Substitute -5 for x in the equation $x - 7 = -12$.

$$x - 7 = -12$$

$$-5 - 7 \stackrel{?}{=} -12 \quad \text{Let } x = -5.$$

$$-12 = -12 \quad \checkmark \quad \text{True}$$

The result, a true statement, shows that -5 is the boundary point. Now test a number on each side of -5 to verify that numbers *less than* -5 make the inequality true. We choose -6 and -4 .

$$x - 7 < -12$$

$$-6 - 7 \stackrel{?}{<} -12 \quad \text{Let } x = -6.$$

$$-13 < -12 \quad \text{True } \checkmark$$

-6 is in the solution set.

$$-4 - 7 \stackrel{?}{<} -12 \quad \text{Let } x = -4.$$

$$-11 < -12 \quad \text{False}$$

-4 is *not* in the solution set.

The check confirms that the solution set is the graph shown in **FIGURE 12**.





FIGURE 12

Using interval notation, the solution set is the infinite interval $(-\infty, -5)$.

NOW TRY

NOW TRY ANSWER

$(3, \infty)$

As with equations, the addition property can also be used to *subtract* the same number from each side of an inequality.

NOW TRY EXERCISE 3

Solve $4x + 1 \geq 5x$, and graph the solution set.

EXAMPLE 3 Using the Addition Property of Inequality

Solve $14 + 2x \leq 3x$, and graph the solution set.

$$14 + 2x \leq 3x$$

$$14 + 2x - 2x \leq 3x - 2x \quad \text{Subtract } 2x.$$

$$14 \leq x \quad \text{Combine like terms.}$$

$$\text{Be careful.} \quad x \geq 14 \quad \text{Rewrite.}$$

The inequality $14 \leq x$ (that is, 14 is less than or equal to x) can also be written $x \geq 14$ (that is, x is greater than or equal to 14). **Notice that in each case the inequality symbol points to the lesser number, 14.**

CHECK

$$14 + 2x = 3x$$

$$14 + 2(14) \stackrel{?}{=} 3(14) \quad \text{Let } x = 14.$$

$$42 = 42 \quad \checkmark \quad \text{True}$$

So 14 satisfies the equality part of \leq . Choose 10 and 16 as test values.

$$14 + 2x < 3x$$

$$14 + 2(10) \stackrel{?}{<} 3(10) \quad \text{Let } x = 10.$$

$$34 < 30 \quad \text{False}$$

10 is not in the solution set.

$$14 + 2(16) \stackrel{?}{<} 3(16) \quad \text{Let } x = 16.$$

$$46 < 48 \quad \checkmark \quad \text{True}$$

16 is in the solution set.

The check confirms that the solution set is the graph shown in **FIGURE 13**.




FIGURE 13

Using interval notation, the solution set is the infinite interval $[14, \infty)$. **NOW TRY**

OBJECTIVE 3 Solve linear inequalities using the multiplication property.

Consider the following true statement.

$$-2 < 5$$

Multiply each side by some positive number—for example, 8.

$$-2(8) < 5(8) \quad \text{Multiply by } 8.$$

$$-16 < 40 \quad \text{True}$$

The result is true. Start again with $-2 < 5$, and multiply each side by some negative number—for example, -8 .

$$-2(-8) < 5(-8) \quad \text{Multiply by } -8.$$


$$16 < -40 \quad \text{False}$$

The result is false. To make it true, we must change the direction of the inequality symbol.

$$16 > -40 \quad \text{True}$$

As these examples suggest, multiplying each side of an inequality by a *negative* number requires reversing the direction of the inequality symbol.

NOW TRY ANSWER



Multiplication Property of Inequality

Let a , b , and c represent real numbers, where $c \neq 0$.

(a) If c is **positive**, then the inequalities

$$a < b \quad \text{and} \quad ac < bc \quad \text{are equivalent.}^*$$

(b) If c is **negative**, then the inequalities

$$a < b \quad \text{and} \quad ac > bc \quad \text{are equivalent.}^*$$

That is, each side of an inequality may be multiplied (or divided) by the same *positive* number without changing the direction of the inequality symbol. **If the multiplier is negative, we must reverse the direction of the inequality symbol.**

*This also applies to $a \leq b$, $a > b$, and $a \geq b$.

Because division is defined in terms of multiplication, the multiplication property also applies when dividing each side of an inequality by the same number.

NOW TRY EXERCISE 4

Solve each inequality, and graph the solution set.

(a) $8x \geq -40$

(b) $-20x > -60$

EXAMPLE 4 Using the Multiplication Property of Inequality

Solve each inequality, and graph the solution set.

(a) $5x \leq -30$

Divide each side by 5. **Because $5 > 0$, do not reverse the direction of the inequality symbol.**

$$\begin{aligned} 5x &\leq -30 \\ \frac{5x}{5} &\leq \frac{-30}{5} && \text{Divide by 5.} \\ x &\leq -6 \end{aligned}$$



FIGURE 14

The solution set is graphed in **FIGURE 14** and written $(-\infty, -6]$.

(b) $-4x \leq 32$

Divide each side by -4 . **Because $-4 < 0$, reverse the direction of the inequality symbol.**

$$\begin{aligned} -4x &\leq 32 \\ \frac{-4x}{-4} &\geq \frac{32}{-4} && \begin{array}{l} \text{Divide by } -4. \\ \text{Reverse the direction of the symbol.} \end{array} \\ x &\geq -8 \end{aligned}$$

Reverse the inequality symbol when dividing by a *negative* number.






FIGURE 15

FIGURE 15 shows the graph of the solution set, $[-8, \infty)$.

NOW TRY

NOW TRY ANSWERS



To solve a linear inequality in one variable, use the following steps.

Solving a Linear Inequality in One Variable

Step 1 Simplify each side separately. Use the distributive property as needed.

- Clear any parentheses.
- Clear any fractions or decimals.
- Combine like terms.

Step 2 Isolate the variable terms on one side. Use the addition property of inequality so that all terms with variables are on one side of the inequality and all constants (numbers) are on the other side.

Step 3 Isolate the variable. Use the multiplication property of inequality to obtain an inequality in one of the following forms, where k is a constant (number).

$$\text{variable} < k, \quad \text{variable} \leq k, \quad \text{variable} > k, \quad \text{or} \quad \text{variable} \geq k$$

Remember: Reverse the direction of the inequality symbol only when multiplying or dividing each side of an inequality by a negative number.

NOW TRY EXERCISE 5

Solve and graph the solution set.

$$5 - 2(x - 4) \leq 11 - 4x$$

EXAMPLE 5 Solving a Linear Inequality

Solve $-3(x + 4) + 2 \geq 7 - x$, and graph the solution set.

Step 1

$$-3(x + 4) + 2 \geq 7 - x$$

$$-3x - 3(4) + 2 \geq 7 - x \quad \text{Distributive property}$$

$$-3x - 12 + 2 \geq 7 - x \quad \text{Multiply.}$$

$$-3x - 10 \geq 7 - x \quad (*) \text{ Combine like terms.}$$

Step 2

$$-3x - 10 + x \geq 7 - x + x \quad \text{Add } x.$$

$$-2x - 10 \geq 7 \quad \text{Combine like terms.}$$

$$-2x - 10 + 10 \geq 7 + 10 \quad \text{Add 10.}$$

$$-2x \geq 17 \quad \text{Combine like terms.}$$

Step 3

$$\frac{-2x}{-2} \leq \frac{17}{-2} \quad \begin{array}{l} \text{Divide by } -2. \\ \text{Change } \geq \text{ to } \leq. \end{array}$$

Be sure to reverse the direction of the inequality symbol.


$$x \leq -\frac{17}{2}$$


FIGURE 16

NOW TRY ANSWER

5.
 $(-\infty, -1]$

FIGURE 16 shows the graph of the solution set, $(-\infty, -\frac{17}{2}]$.

NOTE In Step 2 of **Example 5**, if we add $3x$ (instead of x) to both sides of the inequality, we obtain the following sequence of equivalent inequalities.

$$\begin{array}{ll}
 -3x - 10 \geq 7 - x & \text{See (*) of Example 5.} \\
 -3x - 10 + 3x \geq 7 - x + 3x & \text{Add } 3x. \\
 -10 \geq 2x + 7 & \text{Combine like terms.} \\
 -10 - 7 \geq 2x + 7 - 7 & \text{Subtract 7.} \\
 -17 \geq 2x & \text{Combine like terms.} \\
 -\frac{17}{2} \geq x & \text{Divide by 2.} \\
 \text{The symbol points to } x \text{ in each case. } & x \leq -\frac{17}{2} \quad \text{Rewrite. The same solution results.}
 \end{array}$$

NOW TRY EXERCISE 6

Solve and graph the solution set.

$$\frac{3}{4}(x - 2) + \frac{1}{2} > \frac{1}{5}(x - 8)$$

EXAMPLE 6 Solving a Linear Inequality with Fractions

Solve $-\frac{2}{3}(x - 3) - \frac{1}{2} < \frac{1}{2}(5 - x)$, and graph the solution set.

$$-\frac{2}{3}(x - 3) - \frac{1}{2} < \frac{1}{2}(5 - x)$$

Step 1

$$-\frac{2}{3}x + 2 - \frac{1}{2} < \frac{5}{2} - \frac{1}{2}x \quad \text{Clear parentheses.}$$

$$6\left(-\frac{2}{3}x + 2 - \frac{1}{2}\right) < 6\left(\frac{5}{2} - \frac{1}{2}x\right) \quad \text{To clear the fractions multiply by 6, the LCD.}$$

$$6\left(-\frac{2}{3}x\right) + 6(2) + 6\left(-\frac{1}{2}\right) < 6\left(\frac{5}{2}\right) + 6\left(-\frac{1}{2}x\right) \quad \text{Distributive property}$$

$$-4x + 12 - 3 < 15 - 3x \quad \text{Multiply.}$$

$$-4x + 9 < 15 - 3x \quad \text{Combine like terms.}$$

Step 2

$$-4x + 9 + 3x < 15 - 3x + 3x \quad \text{Add } 3x.$$

$$-x + 9 < 15 \quad \text{Combine like terms.}$$

$$-x + 9 - 9 < 15 - 9 \quad \text{Subtract 9.}$$

$$-x < 6 \quad \text{Combine like terms.}$$

Step 3

$$-1(-x) > -1(6) \quad \text{Multiply by } -1. \\ \text{Change } < \text{ to } >.$$

$$x > -6$$

Reverse the inequality symbol when multiplying by a negative number.




FIGURE 17

FIGURE 17 shows the graph of the solution set, $(-6, \infty)$.

NOW TRY ANSWER

6. $(-\frac{12}{11}, \infty)$

OBJECTIVE 4 Solve linear inequalities with three parts.

Some applications involve a **three-part inequality** such as

$$3 < x + 2 < 8, \text{ where } x + 2 \text{ is between } 3 \text{ and } 8.$$

NOW TRY EXERCISE 7

Solve and graph the solution set.

$$-1 < x - 2 < 3$$

EXAMPLE 7 Solving a Three-Part Inequality

Solve $3 < x + 2 < 8$, and graph the solution set.

$$3 < x + 2 < 8$$

$$3 - 2 < x + 2 - 2 < 8 - 2$$

$$1 < x < 6$$

Do not write $8 < x + 2 < 3$, which implies $8 < 3$, a false statement.

Subtract 2 from all three parts.

Thus, x must be between 1 and 6 so that $x + 2$ will be between 3 and 8.



FIGURE 18

The solution set is graphed in **FIGURE 18** and written as the open interval $(1, 6)$.

NOW TRY

CAUTION Write three-part inequalities so that the symbols point in the same direction, and both point toward the lesser number.

NOW TRY EXERCISE 8

Solve and graph the solution set.

$$-2 < -4x - 5 \leq 7$$

EXAMPLE 8 Solving a Three-Part Inequality

Solve $-2 \leq -3x - 1 \leq 5$, and graph the solution set.

$$-2 \leq -3x - 1 \leq 5$$

$$-2 + 1 \leq -3x - 1 + 1 \leq 5 + 1 \quad \text{Add 1 to each part.}$$

$$-1 \leq -3x \leq 6$$

$$\frac{-1}{-3} \geq \frac{-3x}{-3} \geq \frac{6}{-3} \quad \begin{array}{l} \text{Divide each part by } -3. \\ \text{Reverse the direction of the} \\ \text{inequality symbols.} \end{array}$$

$$\frac{1}{3} \geq x \geq -2$$

$$-2 \leq x \leq \frac{1}{3} \quad \begin{array}{l} \text{Rewrite in the} \\ \text{order on the} \\ \text{number line.} \end{array}$$





FIGURE 19





The solution set is graphed in **FIGURE 19** and written as the closed interval $\left[-2, \frac{1}{3}\right]$.

NOW TRY

NOW TRY ANSWERS



▼ Solution Sets of Equations and Inequalities

Equation or Inequality	Typical Solution Set	Graph of Solution Set
Linear equation $5x + 4 = 14$	$\{2\}$	
Linear inequality $5x + 4 < 14$	$(-\infty, 2)$	
$5x + 4 > 14$	$(2, \infty)$	
Three-part inequality $-1 \leq 5x + 4 \leq 14$	$[-1, 2]$	

OBJECTIVE 5 Solve applied problems using linear inequalities.

▼ Words and Phrases That Suggest Inequality

Word Statement	Interpretation	Example	Inequality
a exceeds b .	$a > b$	Juan's age j exceeds 21 yr.	$j > 21$
a is at least b .	$a \geq b$	Juan is at least 21 yr old.	$j \geq 21$
a is no less than b .	$a \geq b$	Juan is no less than 21 yr old.	$j \geq 21$
a is at most b .	$a \leq b$	Mia's age m is at most 10 yr.	$m \leq 10$
a is no more than b .	$a \leq b$	Mia is no more than 10 yr old.	$m \leq 10$

In **Examples 9 and 10**, we use the six problem-solving steps from earlier, changing Step 3 from

“Write an equation” to “Write an inequality.”


NOW TRY EXERCISE 9

A local health club charges a \$40 one-time enrollment fee, plus \$35 per month for a membership. Sara can spend no more than \$355 on this exercise expense. What is the *maximum* number of months that Sara can belong to this health club?

EXAMPLE 9 Using a Linear Inequality to Solve a Rental Problem

A rental company charges \$20 to rent a chain saw, plus \$9 per hr. Tom can spend no more than \$65 to clear some logs from his yard. What is the maximum amount of time he can use the rented saw?

Step 1 Read the problem again.

Step 2 Assign a variable. Let $x =$ the number of hours he can rent the saw.

Step 3 Write an inequality. He must pay \$20, plus \$9 x , to rent the saw for x hours, and this amount must be *no more than* \$65.

$$\underbrace{20 + 9x}_{\text{Cost of renting}} \quad \underbrace{\leq}_{\text{is no more than}} \quad \underbrace{65}_{\text{dollars.}}$$

Step 4 Solve.

$$9x \leq 45 \quad \text{Subtract 20.}$$

$$x \leq 5 \quad \text{Divide by 9.}$$

Step 5 State the answer. He can use the saw for a maximum of 5 hr. (Of course, he may use it for less time, as indicated by the inequality $x \leq 5$.)

Step 6 Check. If Tom uses the saw for 5 hr, he will spend

$$20 + 9(5) = 65 \text{ dollars, the maximum amount.}$$

NOW TRY 

NOW TRY ANSWER

9. 9 months

NOW TRY EXERCISE 10

Joel has scores of 82, 97, and 93 on his first three exams. What score must he earn on the fourth exam to keep an average of at least 90?

EXAMPLE 10 Finding an Average Test Score

Martha has scores of 88, 86, and 90 on her first three algebra tests. An average score of at least 90 will earn an A in the class. What possible scores on her fourth test will earn her an A average?

Let $x =$ the score on the fourth test. Her average score must be at least 90. To find the average of four numbers, add them and then divide by 4.

$$\begin{aligned} \frac{88 + 86 + 90 + x}{4} &\geq 90 && \text{Average is at least } 90. \\ 264 + x &\geq 360 && \text{Add the scores.} \\ 264 + x &\geq 360 && \text{Multiply by 4.} \\ x &\geq 96 && \text{Subtract 264.} \end{aligned}$$

She must score 96 or more on her fourth test.

CHECK $\frac{88 + 86 + 90 + 96}{4} = \frac{360}{4} = 90$, the minimum score. ✓

A score of 96 or more will give an average of at least 90, as required. **NOW TRY**

NOW TRY ANSWER
10. at least 88

1.5 Exercises **FOR EXTRA HELP** **MyLab Math**

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Study cards are a great way to learn vocabulary, procedures, and so on. **Review Study Skill 5, Using Study Cards.**

Concept Check Match each inequality in Column I with the correct graph or interval in Column II.

I	II
1. $x \leq 3$	A.
2. $x > 3$	B.
3. $x < 3$	C. $(3, \infty)$
4. $x \geq 3$	D. $(-\infty, 3]$
5. $-3 \leq x \leq 3$	E. $(-3, 3)$
6. $-3 < x < 3$	F. $[-3, 3]$

Concept Check Work each problem involving inequalities.

7. A high level of LDL cholesterol (“bad cholesterol”) in the blood increases a person’s risk of heart disease. The table shows how LDL levels affect risk.

If x represents the LDL cholesterol number, write a linear inequality or three-part inequality for each category. Use x as the variable.

- (a) Optimal
- (b) Near optimal/above optimal
- (c) Borderline high
- (d) High
- (e) Very high

LDL Cholesterol	Risk Category
Less than 100	Optimal
100–129	Near optimal/ above optimal
130–159	Borderline high
160–189	High
190 and above	Very high

Data from WebMD.

8. A high level of triglycerides in the blood also increases a person's risk of heart disease. The table shows how triglyceride levels affect risk.

If x represents the triglycerides number, write a linear inequality or three-part inequality for each category. Use x as the variable.

- (a) Normal (b) Mildly high
(c) High (d) Very high

Triglycerides	Risk Category
Less than 100	Normal
100–199	Mildly high
200–499	High
500 or higher	Very high

Data from WebMD.

9. **Concept Check** A student solved the following inequality incorrectly as shown.

$$4x \geq -64$$

$$\frac{4x}{4} \leq \frac{-64}{4}$$

$$x \leq -16$$

Solution set:

$$(-\infty, -16]$$

WHAT WENT WRONG? Give the correct solution set.

10. **Concept Check** A student solved the following inequality incorrectly as shown.

$$-2x < -18$$

$$\frac{-2x}{-2} < \frac{-18}{-2}$$

$$x < 9$$

Solution set:

$$(-\infty, 9)$$

WHAT WENT WRONG? Give the correct solution set.

Solve each inequality. Graph the solution set, and write it using interval notation. See Examples 1–6.

11. $x - 4 \geq 12$

12. $x - 3 \geq 7$

13. $3k + 1 > 22$

14. $5x + 6 < 76$

15. $4x < -16$

16. $2x > -10$

17. $-4x < 16$

18. $-5x > 25$

19. $-\frac{3}{4}x \geq 30$

20. $-\frac{2}{3}x \leq 12$

21. $-1.3x \geq -5.2$

22. $-2.5x \leq -1.25$

23. $5x + 2 \leq -48$

24. $4x + 1 \leq -31$

25. $\frac{3k - 1}{4} > 5$

26. $\frac{5x - 6}{8} < 8$

27. $\frac{2x - 5}{-4} > 5$

28. $\frac{3x - 2}{-5} < 6$

29. $3k + 1 < -20$

30. $5z + 6 > -29$

31. $6x - 4 \geq -2x$

32. $2x - 8 \geq -2x$

33. $x - 2(x - 4) \leq 3x$

34. $x - 3(x + 1) \leq 4x$

35. $-(4 + r) + 2 - 3r < -14$

36. $-(9 + x) - 5 + 4x \geq 4$

37. $-3(x - 6) > 2x - 2$

38. $-2(x + 4) \leq 6x + 16$

39. $\frac{2}{3}(3x - 1) \geq \frac{3}{2}(2x - 3)$

40. $\frac{7}{5}(10x - 1) < \frac{2}{3}(6x + 5)$

41. $-\frac{1}{4}(p + 6) + \frac{3}{2}(2p - 5) < 10$

42. $\frac{3}{5}(t - 2) - \frac{1}{4}(2t - 7) \leq 3$

43. $3(2x - 4) - 4x < 2x + 3$

44. $7(4 - x) + 5x < 2(16 - x)$

45. $8\left(\frac{1}{2}x + 3\right) < 8\left(\frac{1}{2}x - 1\right)$

46. $10\left(\frac{1}{5}x + 2\right) < 10\left(\frac{1}{5}x + 1\right)$

Solve each inequality. Graph the solution set, and write it using interval notation. See Examples 7 and 8.

47. $-4 < x - 5 < 6$

48. $-1 < x + 1 < 8$

49. $-9 \leq x + 5 \leq 15$

50. $-4 \leq x + 3 \leq 10$

51. $-6 \leq 2x + 4 \leq 16$

52. $-15 < 3x + 6 < -12$

53. $-19 \leq 3x - 5 \leq 1$

54. $-16 < 3x + 2 < -10$

55. $4 \leq -9x + 5 < 8$

56. $4 \leq -2x + 3 < 8$

57. $-8 \leq -4x + 2 \leq 6$

58. $-12 \leq -6x + 3 \leq 15$

59. $-3 < \frac{3}{4}x < 6$

60. $-4 < \frac{2}{3}x < 12$

61. $-1 \leq \frac{2x - 5}{6} \leq 5$

62. $-3 \leq \frac{3x + 1}{4} \leq 3$

63. $-5 \leq \frac{6 - 5x}{2} \leq 0$

64. $-4 \leq \frac{2 - 4x}{3} \leq 0$

Give, in interval notation, the unknown numbers in each description.

65. A number is between 0 and 1.

66. A number is between -3 and -2 .67. Six times a number is between -12 and 12 .68. Half a number is between -3 and 2 .

69. When 1 is added to twice a number, the result is greater than or equal to 7.

70. If 8 is subtracted from a number, then the result is at least 5.

71. One third of a number is added to 6, giving a result of at least 3.

72. Three times a number, minus 5, is no more than 7.

Solve each problem. See Examples 9 and 10.

73. Bonnie earned scores of 90 and 82 on her first two tests in English literature. What score must she make on her third test to keep an average of 84 or greater?

74. Scott scored 92 and 96 on his first two tests in "Methods in Teaching Mathematics." What score must he make on his third test to keep an average of 90 or greater?


75. To earn a B in an algebra course requires an average of at least 80 on five tests. A student has scores of 75, 91, 82, and 74. What possible scores on the fifth test would guarantee this student a B in the class?

76. To pass Algebra II requires an average of at least 70 on four tests. A student has scores of 80, 62, and 73. What possible scores on the fourth test would guarantee this student a passing score in the class?

77. Latrice is signing up for cell phone service. She must decide between Plan A, which costs \$54.99 per month with a free phone included, and Plan B, which costs \$49.99 per month, but would require her to buy a phone for \$129. Under either plan, she does not expect to go over the included number of monthly minutes. After how many months would Plan B be a better deal?



78. Newlyweds Bryce and Lauren need to move their belongings to their new apartment. They can rent a truck from U-Haul for \$29.95 per day plus 28 cents per mile or from Budget Truck Rentals for \$34.95 per day plus 25 cents per mile. After how many miles (to the nearest mile) would the Budget rental be a better deal than the U-Haul rental?



79. The average monthly precipitation for Dallas, TX, for October, November, and December is 3.47 in. If 2.88 in. falls in October and 3.13 in. falls in November, how many inches must fall in December so that the average monthly precipitation for these months exceeds 3.47 in.? (Data from www.weather.com)
80. The average monthly precipitation for Honolulu, HI, for October, November, and December is 3.11 in. If 2.98 in. falls in October and 3.05 in. falls in November, how many inches must fall in December so that the average monthly precipitation for these months exceeds 3.11 in.? (Data from www.weather.com)
81. According to some medical advisors, a body mass index (BMI) between 19 and 25 suggests a healthful weight. Use the formula

$$\text{BMI} = \frac{704 \times (\text{weight in pounds})}{(\text{height in inches})^2}$$

to find the weight range w , to the nearest pound, that gives a healthful BMI for each height.

- (a) 72 in. (b) 63 in. (c) Your height in inches
82. To achieve the maximum benefit from exercising, the heart rate, in beats per minute, should be in the target heart rate (THR) zone. For a person aged A , the formula is as follows.

$$0.7(220 - A) \leq \text{THR} \leq 0.85(220 - A)$$

Find the THR to the nearest whole number for each age. (Source: Hockey, R. V., *Physical Fitness: The Pathway to Healthful Living*, Times Mirror/Mosby College Publishing.)

- (a) 35 (b) 55 (c) Your age

A product will produce a profit only when the revenue R from selling the product exceeds the cost C of producing it. Find the least whole number of units x that must be sold for the business to show a profit for the item described.

83. Peripheral Visions, Inc., finds that the cost of producing x studio-quality DVDs is $C = 20x + 100$, while the revenue produced from them is $R = 24x$ (C and R in dollars).
84. Speedy Delivery finds that the cost of making x deliveries is $C = 3x + 2300$, while the revenue produced from them is $R = 5.50x$ (C and R in dollars).

RELATING CONCEPTS For Individual or Group Work (Exercises 85–90)

Work Exercises 85–90 in order.

85. Solve the linear equation. Graph the solution set on a number line.

$$5(x + 3) - 2(x - 4) = 2(x + 7)$$

86. Solve the linear inequality. Graph the solution set on a number line.

$$5(x + 3) - 2(x - 4) > 2(x + 7)$$

87. Solve the linear inequality. Graph the solution set on a number line.

$$5(x + 3) - 2(x - 4) < 2(x + 7)$$

88. Graph all the solution sets of the equation and inequalities in **Exercises 85–87** on the same number line. (This is the **union** of the solution sets.) What set do we obtain?

89. Complete the following:

The solution set of $-3(x + 2) = 3x + 12$ is _____.

The solution set of $-3(x + 2) < 3x + 12$ is _____.

Therefore, the solution set of $-3(x + 2) > 3x + 12$ is _____.

90. Describe the union of the three solution sets in **Exercise 89** in words, and write it in interval notation as a single interval.

1.6 Set Operations and Compound Inequalities

OBJECTIVES

- 1 Recognize set intersection and union.
- 2 Find the intersection of two sets.
- 3 Solve compound inequalities with the word *and*.
- 4 Find the union of two sets.
- 5 Solve compound inequalities with the word *or*.

VOCABULARY

- intersection
- compound inequality
- union

NOW TRY EXERCISE 1

Let $A = \{2, 4, 6, 8\}$ and $B = \{0, 2, 6, 8\}$. Find $A \cap B$.

NOW TRY ANSWER

1. $\{2, 6, 8\}$

OBJECTIVE 1 Recognize set intersection and union.

Consider the two sets A and B defined as follows.

$$A = \{1, 2, 3\}, \quad B = \{2, 3, 4\}$$

The set of all elements that belong to both A **and** B , called their *intersection* and symbolized $A \cap B$, is given by

$$A \cap B = \{2, 3\}. \quad \text{Intersection}$$

The set of all elements that belong to either A **or** B , or both, called their *union* and symbolized $A \cup B$, is given by

$$A \cup B = \{1, 2, 3, 4\}. \quad \text{Union}$$


OBJECTIVE 2 Find the intersection of two sets.

The intersection of two sets is defined with the word *and*.

Intersection of Sets

For any two sets A and B , the **intersection** of A and B , symbolized $A \cap B$, is defined as follows.

$$A \cap B = \{x \mid x \text{ is an element of } A \text{ and } x \text{ is an element of } B\}$$



EXAMPLE 1 Finding the Intersection of Two Sets

Let $A = \{1, 2, 3, 4\}$ and $B = \{2, 4, 6\}$. Find $A \cap B$.

The set $A \cap B$ contains those elements that belong to both A *and* B .

$$\begin{aligned} A \cap B &= \{1, 2, 3, 4\} \cap \{2, 4, 6\} \\ &= \{2, 4\} \end{aligned}$$

NOW TRY

OBJECTIVE 3 Solve compound inequalities with the word *and*.

A **compound inequality** consists of two inequalities linked by a connective word.

$$\begin{aligned} x + 1 \leq 9 \quad \text{and} \quad x - 2 \geq 3 \\ 2x > 4 \quad \text{or} \quad 3x - 6 < 5 \end{aligned} \quad \text{Compound inequalities}$$

Solving a Compound Inequality with *and*

Step 1 Solve each inequality individually.

Step 2 The solution set of the compound inequality includes all numbers that satisfy both inequalities in Step 1—that is, the *intersection* of the solution sets.

NOW TRY EXERCISE 2

Solve the compound inequality, and graph the solution set.

$$x - 2 \leq 5 \quad \text{and} \quad x + 5 \geq 9$$

EXAMPLE 2 Solving a Compound Inequality with and

Solve the compound inequality, and graph the solution set.

$$x + 1 \leq 9 \quad \text{and} \quad x - 2 \geq 3$$


Step 1 Solve each inequality individually.

$$x + 1 \leq 9 \quad \text{and} \quad x - 2 \geq 3$$

$$x + 1 - 1 \leq 9 - 1 \quad \text{and} \quad x - 2 + 2 \geq 3 + 2$$

$$x \leq 8 \quad \text{and} \quad x \geq 5$$

Step 2 The solution set will include all numbers that satisfy *both* inequalities in Step 1 at the same time. The compound inequality is true whenever $x \leq 8$ and $x \geq 5$ are both true. See the graphs in **FIGURE 20**.



The set of points where the graphs “overlap” represents the intersection.

FIGURE 20

The intersection of the two graphs in **FIGURE 20** is the solution set. **FIGURE 21** shows the graph of the solution set, written using interval notation as the closed interval $[5, 8]$.




FIGURE 21

NOW TRY

NOW TRY EXERCISE 3

Solve and graph.

$$\begin{aligned} -4x - 1 < 7 \quad \text{and} \\ 3x + 4 \geq -5 \end{aligned}$$

EXAMPLE 3 Solving a Compound Inequality with and

Solve the compound inequality, and graph the solution set.

$$-3x - 2 > 5 \quad \text{and} \quad 5x - 1 \leq -21$$

Step 1 Solve each inequality individually.

$$-3x - 2 > 5 \quad \text{and} \quad 5x - 1 \leq -21$$

$$-3x > 7 \quad \text{and} \quad 5x \leq -20$$

$$x < -\frac{7}{3} \quad \text{and} \quad x \leq -4$$

Remember to reverse the direction of the inequality symbol.

The graphs of $x < -\frac{7}{3}$ and $x \leq -4$ are shown in **FIGURE 22**.




FIGURE 22

Step 2 Now find all values of x that are less than $-\frac{7}{3}$ and also less than or equal to -4 . This is shown in **FIGURE 23**. The solution set is written as the infinite interval $(-\infty, -4]$.






FIGURE 23

NOW TRY

NOW TRY ANSWERS



NOW TRY
EXERCISE 4

Solve and graph.

$$x - 7 < -12 \quad \text{and} \\ 2x + 1 > 5$$

EXAMPLE 4 Solving a Compound Inequality with *and*

Solve the compound inequality, and graph the solution set.

$$x + 2 < 5 \quad \text{and} \quad x - 10 > 2$$

Step 1 Solve each inequality individually.

$$x + 2 < 5 \quad \text{and} \quad x - 10 > 2$$

$$x < 3 \quad \text{and} \quad x > 12$$


The graphs of $x < 3$ and $x > 12$ are shown in **FIGURE 24**.

FIGURE 24


Step 2 There is no number that is both less than 3 *and* greater than 12, so the given compound inequality has no solution. See **FIGURE 25**. The solution set is \emptyset .

FIGURE 25

NOW TRY


OBJECTIVE 4 Find the union of two sets.

The union of two sets is defined with the word *or*.

Union of Sets

For any two sets A and B , the **union** of A and B , symbolized $A \cup B$, is defined as follows.

$$A \cup B = \{x \mid x \text{ is an element of } A \text{ or } x \text{ is an element of } B\}$$



NOW TRY
EXERCISE 5

Let $A = \{5, 10, 15, 20\}$
and $B = \{5, 15, 25\}$.
Find $A \cup B$.

EXAMPLE 5 Finding the Union of Two Sets

Let $A = \{1, 2, 3, 4\}$ and $B = \{2, 4, 6\}$. Find $A \cup B$.Begin by listing all the elements of set A : 1, 2, 3, 4. Then list any additional elements from set B . In this case the elements 2 and 4 are already listed, so the only additional element is 6.

$$A \cup B = \{1, 2, 3, 4\} \cup \{2, 4, 6\} \\ = \{1, 2, 3, 4, 6\}$$

The union consists of all elements in either A *or* B (or both).

NOW TRY

NOW TRY ANSWERS

4. \emptyset
5. $\{5, 10, 15, 20, 25\}$

NOTE Although the elements 2 and 4 appeared in both sets A and B in **Example 5**, they are written only once in $A \cup B$.

OBJECTIVE 5 Solve compound inequalities with the word *or*.**Solving a Compound Inequality with *or*****Step 1** Solve each inequality individually.**Step 2** The solution set of the compound inequality includes all numbers that satisfy either one or the other (or both) of the inequalities in Step 1—that is, the *union* of the solution sets.**NOW TRY EXERCISE 6**

Solve and graph.

$$-12x \leq -24 \quad \text{or} \quad x + 9 < 8$$

EXAMPLE 6 Solving a Compound Inequality with *or*

Solve the compound inequality, and graph the solution set.

$$6x - 4 < 2x \quad \text{or} \quad -3x \leq -9$$


Step 1 Solve each inequality individually.

$$6x - 4 < 2x \quad \text{or} \quad -3x \leq -9$$

$$4x < 4$$

$$x < 1 \quad \text{or} \quad x \geq 3$$

Remember to reverse the inequality symbol.

The graphs of these two inequalities are shown in **FIGURE 26**.The set of points in *either* of the graphs represents the union.**FIGURE 26****Step 2** Because the inequalities are joined with *or*, find the union of the two solution sets. The union is the disjoint interval in **FIGURE 27**.**FIGURE 27**In interval notation, the solution set is $(-\infty, 1) \cup [3, \infty)$. Always pay particular attention to the end points of the solution sets and whether parentheses, brackets, or one of each should be used.**NOW TRY** **CAUTION** When inequalities are used to write the solution set in **Example 6**, it *must* be written using two separate inequalities.

$$x < 1 \quad \text{or} \quad x \geq 3$$

Writing $3 \leq x < 1$, which translates using *and*, would imply that

$$3 \leq 1, \quad \text{which is } \mathbf{FALSE}.$$

NOW TRY ANSWER

6. $(-\infty, -1) \cup [2, \infty)$

NOW TRY EXERCISE 7

Solve and graph.

$$-x + 2 < 6 \quad \text{or} \quad 6x - 8 \geq 10$$

EXAMPLE 7 Solving a Compound Inequality with or

Solve the compound inequality, and graph the solution set.

$$-4x + 1 \geq 9 \quad \text{or} \quad 5x + 3 \leq -12$$

Step 1 Solve each inequality individually.

$$-4x + 1 \geq 9 \quad \text{or} \quad 5x + 3 \leq -12$$

$$-4x \geq 8 \quad \text{or} \quad 5x \leq -15$$

$$x \leq -2 \quad \text{or} \quad x \leq -3$$

The graphs of these two inequalities are shown in **FIGURE 28**.




FIGURE 28

Step 2 **FIGURE 29** shows the union, the infinite interval $(-\infty, -2]$.



FIGURE 29

NOW TRY

NOW TRY EXERCISE 8

Solve and graph.

$$8x - 4 \geq 20 \quad \text{or} \\ -2x + 1 > -9$$

EXAMPLE 8 Solving a Compound Inequality with or

Solve the compound inequality, and graph the solution set.

$$-2x + 5 \geq 11 \quad \text{or} \quad 4x - 7 \geq -27$$

Step 1 Solve each inequality individually.

$$-2x + 5 \geq 11 \quad \text{or} \quad 4x - 7 \geq -27$$

$$-2x \geq 6 \quad \text{or} \quad 4x \geq -20$$

$$x \leq -3 \quad \text{or} \quad x \geq -5$$

The graphs of these two inequalities are shown in **FIGURE 30**.




FIGURE 30

Step 2 By taking the union, we obtain every real number as a solution because every real number satisfies at least one of the two inequalities. The set of all real numbers is graphed in **FIGURE 31**. It is written in interval notation as $(-\infty, \infty)$.






FIGURE 31

NOW TRY

NOW TRY ANSWERS



NOW TRY EXERCISE 9

In **Example 9**, list the elements of each set.

- (a) The set of countries to which exports were greater than \$100,000 million and from which imports were less than \$400,000 million
- (b) The set of countries to which exports were less than \$200,000 million or from which imports were greater than \$250,000 million

EXAMPLE 9 Applying Intersection and Union

The five top U.S. trading partners for 2016 are listed in the table. Amounts are in millions of dollars.

Country	U.S. Exports to Country	U.S. Imports from Country
China	115,602	462,618
Canada	266,797	277,756
Mexico	229,702	294,056
Japan	63,236	132,046
Germany	49,363	114,099



Data from U.S. Census Bureau.

List the elements of the following sets.

- (a) The set of countries to which exports were greater than \$200,000 million *and* from which imports were less than \$300,000 million

The countries that satisfy both conditions are Canada and Mexico, so the set is

$$\{\text{Canada, Mexico}\}.$$

- (b) The set of countries to which exports were less than \$100,000 million *or* from which imports were greater than \$200,000 million

Here, any country that satisfies at least one of the conditions is in the set. This set includes all five countries:

$$\{\text{China, Canada, Mexico, Japan, Germany}\}.$$

NOW TRY

NOW TRY ANSWERS

- 9. (a) {Canada, Mexico}
- (b) {China, Canada, Mexico, Japan, Germany}

1.6 Exercises FOR EXTRA HELP MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Reread your class notes before working the assigned exercises. **Review Study Skill 3, Taking Lecture Notes.**





Concept Check Determine whether each statement is true or false. If it is false, explain why.

1. The union of the solution sets of $x + 1 = 6$, $x + 1 < 6$, and $x + 1 > 6$ is $(-\infty, \infty)$.
2. The intersection of the sets $\{x | x \geq 9\}$ and $\{x | x \leq 9\}$ is \emptyset .
3. The union of the sets $(-\infty, 7)$ and $(7, \infty)$ is $\{7\}$.
4. The intersection of the sets $(-\infty, 7]$ and $[7, \infty)$ is $\{7\}$.

Let $A = \{1, 2, 3, 4, 5, 6\}$, $B = \{1, 3, 5\}$, $C = \{1, 6\}$, and $D = \{4\}$. Find each set. See Examples 1 and 5.

- 5. $A \cap D$
- 6. $B \cap C$
- 7. $B \cap A$
- 8. $C \cap A$
- 9. $B \cap \emptyset$
- 10. $A \cap \emptyset$
- 11. $A \cup B$
- 12. $B \cup D$
- 13. $B \cup C$
- 14. $C \cup D$

Concept Check Two sets are specified by graphs. Graph the intersection of the two sets.



Solve each compound inequality. Graph the solution set, and write it using interval notation. See Examples 2–4.

19. $x < 2$ and $x > -3$

21. $x \leq 2$ and $x \leq 5$

23. $x \leq 3$ and $x \geq 6$

25. $x - 3 \leq 6$ and $x + 2 \geq 7$

27. $-3x > 3$ and $x + 3 > 0$

29. $3x - 4 \leq 8$ and $-4x + 1 \geq -15$

20. $x < 5$ and $x > 0$

22. $x \geq 3$ and $x \geq 6$





24. $x \leq -1$ and $x \geq 3$

26. $x + 5 \leq 11$ and $x - 3 \geq -1$

28. $-3x < 3$ and $x + 2 < 6$

30. $7x + 6 \leq 48$ and $-4x + 3 \geq -21$

Concept Check Two sets are specified by graphs. Graph the union of the two sets.



Solve each compound inequality. Graph the solution set, and write it using interval notation. See Examples 6–8.

35. $x \leq 1$ or $x \leq 8$

37. $x \geq -2$ or $x \geq 5$

39. $x \geq -2$ or $x \leq 4$

41. $x + 2 > 7$ or $1 - x > 6$

43. $x + 1 > 3$ or $-4x + 1 > 5$

45. $4x + 1 \geq -7$ or $-2x + 3 \geq 5$

36. $x \geq 1$ or $x \geq 8$

38. $x \leq -2$ or $x \leq 6$

40. $x \geq 5$ or $x \leq 7$

42. $x + 1 > 3$ or $x + 4 < 2$

44. $-2x + 1 > -11$ or $x + 1 > 10$

46. $3x + 2 \leq -7$ or $-2x + 1 \leq 9$

Concept Check Express each set in simplest interval form. (Hint: Graph each set and look for the intersection or union.)

47. $(-\infty, -1] \cap [-4, \infty)$

49. $(-\infty, -6] \cap [-9, \infty)$

51. $(-\infty, 3) \cup (-\infty, -2)$

53. $[3, 6] \cup (4, 9)$

48. $[-1, \infty) \cap (-\infty, 9]$

50. $(5, 11] \cap [6, \infty)$

52. $[-9, 1] \cup (-\infty, -3)$

54. $[-1, 2] \cup (0, 5)$

Solve each compound inequality. Graph the solution set, and write it using interval notation. See Examples 2–4 and 6–8.

55. $x < -1$ and $x > -5$

57. $x < 4$ or $x < -2$

59. $-3x \leq -6$ or $-3x \geq 0$

61. $x + 1 \geq 5$ and $x - 2 \leq 10$

56. $x > -1$ and $x < 7$

58. $x < 5$ or $x < -3$

60. $-8x \leq -24$ or $-5x \geq 15$

62. $2x - 6 \leq -18$ and $2x \geq -18$

Average expenses for full-time resident college students at 4-year institutions during the 2015–2016 academic year are shown in the table.

▼ College Expenses (in Dollars)

Type of Expense	Public Schools (in-state)	Private Schools
Tuition and fees	8778	27,951
Board rates	4561	5116
Dormitory charges	5850	6463


Data from National Center for Education Statistics.

List the elements of each set. See Example 9.

- 63. The set of expenses that are less than \$9000 for public schools and are greater than \$15,000 for private schools
- 64. The set of expenses that are greater than \$4000 for public schools and are less than \$6000 for private schools
- 65. The set of expenses that are less than \$9000 for public schools or are greater than \$15,000 for private schools
- 66. The set of expenses that are greater than \$15,000 or are between \$7000 and \$8000



The figures represent the backyards of neighbors Luigi, Mario, Than, and Joe. Suppose that each resident has 150 ft of fencing and enough sod to cover 1400 ft² of lawn.



- 67. Determine the perimeters of the four yards.
- 68. Determine the areas of the four yards.

Give the name or names of the residents whose yards satisfy each description. (Hint: Use the perimeters and areas found in Exercises 67 and 68.)

- 69. The yard can be fenced and the yard can be sodded.
- 70. The yard can be fenced and the yard cannot be sodded.
- 71. The yard cannot be fenced and the yard can be sodded.
- 72. The yard cannot be fenced and the yard cannot be sodded.
- 73. The yard can be fenced or the yard can be sodded.
- 74. The yard cannot be fenced or the yard can be sodded.

RELATING CONCEPTS For Individual or Group Work (Exercises 75–78)

An intermediate algebra teacher bases final grades on points earned for activities as given in the Graded Classwork table on the left. To determine final grades, the teacher strictly adheres to the point ranges given in the Grade Distribution table on the right.

▼ Graded Classwork

Activity	Points Available
Homework and vocabulary	45
Daily activities (scaled)	55
Lab participation and completion	100
Major exams (3 at 100 points)	300
Final Exam	150

Total points 650

▼ Grade Distribution

Grade	Points Required
A	585–650
B	520–584
C	455–519
IP*	< 455 and active
F	< 455 and inactive

* In Progress

Use this information to work Exercises 75–78 in order.

75. Suppose Lauren earns all of the homework and vocabulary points, 50 points for daily activities, and 90 points for lab participation and completion.

Let x = points to be earned on exams.

- Write three inequalities to find the minimum number of points she needs on exams to earn grades no lower than A, B, and C.
- Solve each inequality from part (a) to find the minimum number of points she needs for each grade. What “test average” (as a percent) corresponds to this number of points, given that exams account for 450 possible points? (Round up to the nearest whole number.)



76. See Exercise 75. Write and solve a compound inequality to find the range of points Lauren needs in exam scores to earn a B average. What range of “test averages” (as percents) correspond to these scores, given that exams account for 450 possible points? (Round up to the nearest whole number.)
77. Suppose Mark earns only 15 points in homework and vocabulary, 40 points in daily activities, and 50 points in lab participation. Repeat Exercise 75 using these values.
78. Repeat Exercise 76 given that Mark wants to earn a C average. (Use his classwork points given in Exercise 77.)

1.7

Absolute Value Equations and Inequalities

OBJECTIVES

- 1 Use the distance definition of absolute value.
- 2 Solve equations of the form $|ax + b| = k$, for $k > 0$.
- 3 Solve inequalities of the form $|ax + b| < k$ and of the form $|ax + b| > k$, for $k > 0$.
- 4 Solve absolute value equations that involve rewriting.
- 5 Solve equations of the form $|ax + b| = |cx + d|$.
- 6 Solve special cases of absolute value equations and inequalities.
- 7 Solve an application involving relative error.

VOCABULARY

- absolute value equation
- absolute value inequality

Suppose the government of a country decides that it will comply with a restriction on greenhouse gas emissions *within* 3 years of 2025. This means that the *difference* between the year it will comply and 2025 is less than 3, *without regard to sign*. We state this mathematically as follows, where x represents the year in which it complies.

$$|x - 2025| < 3 \quad \text{Absolute value inequality}$$

We can intuitively reason that the year must be between 2022 and 2028, and thus values that satisfy $2022 < x < 2028$ make this inequality true.

OBJECTIVE 1 Use the distance definition of absolute value.

The **absolute value** of a number x , written $|x|$, represents the undirected distance from x to 0 on a number line. For example, the solution set of $|x| = 4$ is $\{-4, 4\}$, which means $x = -4$ or $x = 4$, as shown in FIGURE 32.




FIGURE 32

Because absolute value represents distance from 0, the solution set of $|x| > 4$ consists of all numbers that are *more* than four units from 0 on a number line. The set $(-\infty, -4) \cup (4, \infty)$ fits this description. The graph consists of disjoint intervals, which means $x < -4$ or $x > 4$, as shown in FIGURE 33.




FIGURE 33

The solution set of $|x| < 4$ consists of all numbers that are *less* than 4 units from 0 on a number line. This is represented by all numbers *between* -4 and 4 , which is given by the open interval $(-4, 4)$, as shown in FIGURE 34. Here, $-4 < x < 4$, which means $x > -4$ and $x < 4$.




FIGURE 34

Absolute value equations and inequalities generally take the form

$$|ax + b| = k, \quad |ax + b| > k, \quad \text{or} \quad |ax + b| < k,$$

where k is a positive number. From FIGURES 32–34, we see that

$$|x| = 4 \quad \text{has the same solution set as} \quad x = -4 \quad \text{or} \quad x = 4,$$

$$|x| > 4 \quad \text{has the same solution set as} \quad x < -4 \quad \text{or} \quad x > 4,$$

$$|x| < 4 \quad \text{has the same solution set as} \quad x > -4 \quad \text{and} \quad x < 4.$$

This is equivalent to $-4 < x < 4$.


Solving Absolute Value Equations and Inequalities

Let k be a positive real number, and p and q be real numbers.

Case 1 To solve $|ax + b| = k$, solve the compound equation

$$ax + b = k \quad \text{or} \quad ax + b = -k.$$


The solution set is usually of the form $\{p, q\}$, which includes two numbers.



Case 2 To solve $|ax + b| > k$,* solve the compound inequality

$$ax + b > k \quad \text{or} \quad ax + b < -k.$$

The solution set is of the form $(-\infty, p) \cup (q, \infty)$, which consists of disjoint intervals.



Case 3 To solve $|ax + b| < k$,** solve the three-part inequality

$$-k < ax + b < k.$$

The solution set is of the form (p, q) , which consists of a single interval.



*This also applies to $|ax + b| \geq k$. The solution set *includes* the endpoints, using brackets rather than parentheses.

**This also applies to $|ax + b| \leq k$. The solution set *includes* the endpoints, using brackets rather than parentheses.

NOTE It is acceptable to write the compound statements in Cases 1 and 2 of the preceding box in the following equivalent forms.

$$ax + b = k \quad \text{or} \quad -(ax + b) = k \quad \text{Alternative for Case 1}$$

$$ax + b > k \quad \text{or} \quad -(ax + b) > k \quad \text{Alternative for Case 2}$$

OBJECTIVE 2 Solve equations of the form $|ax + b| = k$, for $k > 0$.

Remember that because absolute value refers to distance from the origin, an absolute value equation (with $k > 0$) will have two parts.

EXAMPLE 1 Solving an Absolute Value Equation (Case 1)

Solve $|2x + 1| = 7$.

For $|2x + 1|$ to equal 7, $2x + 1$ must be 7 units from 0 on a number line. This can happen only when $2x + 1 = 7$ or $2x + 1 = -7$. This is **Case 1** in the box above. Solve this compound equation as follows.

$$2x + 1 = 7 \quad \text{or} \quad 2x + 1 = -7$$

$$2x = 6 \quad \text{or} \quad 2x = -8 \quad \text{Subtract 1.}$$

$$x = 3 \quad \text{or} \quad x = -4 \quad \text{Divide by 2.}$$

NOW TRY
EXERCISE 1

Solve $|4x - 1| = 11$.

CHECK

$$|2x + 1| = 7$$

$ 2(3) + 1 \stackrel{?}{=} 7$ $ 6 + 1 \stackrel{?}{=} 7$ $ 7 \stackrel{?}{=} 7$ $7 = 7 \checkmark$ True	$\left. \begin{array}{l} \text{Let } x = 3. \\ \\ \\ \end{array} \right $	$ 2(-4) + 1 \stackrel{?}{=} 7$ Let $x = -4$. $ -8 + 1 \stackrel{?}{=} 7$ $ -7 \stackrel{?}{=} 7$ $7 = 7 \checkmark$ True
---------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------

The solution set is $\{-4, 3\}$. See FIGURE 35.



FIGURE 35

NOW TRY

OBJECTIVE 3 Solve inequalities of the form $|ax + b| < k$ and of the form $|ax + b| > k$, for $k > 0$.

NOW TRY
EXERCISE 2

Solve $|4x - 1| > 11$.

EXAMPLE 2 Solving an Absolute Value Inequality (Case 2)

Solve $|2x + 1| > 7$.

Because $2x + 1$ must represent a number that is *more* than 7 units from 0 on either side of a number line, this absolute value inequality is rewritten as the following compound inequality. This is **Case 2**.

$$\begin{aligned}
 2x + 1 &> 7 \quad \text{or} \quad 2x + 1 < -7 \\
 2x &> 6 \quad \text{or} \quad 2x < -8 && \text{Subtract 1.} \\
 x &> 3 \quad \text{or} \quad x < -4 && \text{Divide by 2.}
 \end{aligned}$$

The solution set consists of the disjoint intervals shown in FIGURE 36 and is written $(-\infty, -4) \cup (3, \infty)$.



FIGURE 36

CHECK The excluded endpoints -4 and 3 are correct because from **Example 1** we know that -4 and 3 are the solutions of the related equation. Referring to FIGURE 36, we choose a test point in each of the three intervals $(-\infty, -4)$, $(-4, 3)$, and $(3, \infty)$.

For $(-\infty, -4)$, let $x = -5$. $ 2x + 1 > 7$ $ 2(-5) + 1 \stackrel{?}{>} 7$ $ -9 \stackrel{?}{>} 7$ $9 > 7 \checkmark$ True	$\left. \begin{array}{l} \text{For } (-4, 3), \text{ let } x = 0. \\ \\ \\ \end{array} \right $	For $(3, \infty)$, let $x = 4$. $ 2x + 1 > 7$ $ 2(4) + 1 \stackrel{?}{>} 7$ $ 9 \stackrel{?}{>} 7$ $9 > 7 \checkmark$ True
--------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------

The check confirms that the solution set is $(-\infty, -4) \cup (3, \infty)$.

NOW TRY

NOW TRY ANSWERS

1. $\{-\frac{5}{2}, 3\}$
2. $(-\infty, -\frac{5}{2}) \cup (3, \infty)$

NOW TRY
EXERCISE 3

Solve $|4x - 1| < 11$.

EXAMPLE 3 Solving an Absolute Value Inequality (Case 3)

Solve $|2x + 1| < 7$. Graph the solution set.

The expression $2x + 1$ must represent a number that is less than 7 units from 0 on either side of a number line. That is, $2x + 1$ must be between -7 and 7 , which is written as a three-part inequality. This is **Case 3**.

$$-7 < 2x + 1 < 7$$

$$-8 < 2x < 6 \quad \text{Subtract 1 from each part.}$$

$$-4 < x < 3 \quad \text{Divide each part by 2.}$$

The solution set consists of the open interval $(-4, 3)$ shown in **FIGURE 37**.




FIGURE 37

NOW TRY

Look back at **FIGURES 35, 36, AND 37**, with the graphs of

$$|2x + 1| = 7, \quad |2x + 1| > 7, \quad \text{and} \quad |2x + 1| < 7.$$

If we find the union of the three sets, we obtain the set of all real numbers. For any value of x , $|2x + 1|$ will satisfy *one and only one* of the following: It is equal to 7, greater than 7, or less than 7.

CAUTION Remember the following important concepts.

1. The methods described apply when the constant is alone on one side of the absolute value equation or inequality and is *positive*.
2. Absolute value equations $|ax + b| = k$ and inequalities of the form $|ax + b| > k$ translate into “**or**” compound statements.
3. Absolute value inequalities of the form $|ax + b| < k$ translate into “**and**” compound statements. **Only “and” compound statements may be written as three-part inequalities.**

NOW TRY
EXERCISE 4

Solve $|7 - 4x| \geq 7$.

EXAMPLE 4 Solving an Absolute Value Inequality (Case 2, for \geq)

Solve $|5 - 2x| \geq 5$.

Case 2 is applied. Notice that the endpoints are included because equality is part of the symbol \geq .

$$5 - 2x \geq 5 \quad \text{or} \quad 5 - 2x \leq -5$$

$$-2x \geq 0 \quad \text{or} \quad -2x \leq -10 \quad \text{Subtract 5.}$$

$$x \leq 0 \quad \text{or} \quad x \geq 5 \quad \text{Divide by } -2. \text{ Reverse the direction of the inequality symbols.}$$

See the graph of the disjoint intervals in **FIGURE 38**. The solution set is written $(-\infty, 0] \cup [5, \infty)$.




FIGURE 38

NOW TRY

NOW TRY ANSWERS

3. $(-\frac{5}{2}, 3)$

4. $(-\infty, 0] \cup [\frac{7}{2}, \infty)$

OBJECTIVE 4 Solve absolute value equations that involve rewriting. **NOW TRY EXERCISE 5**Solve $|10x - 2| - 2 = 12$.**EXAMPLE 5** Solving an Absolute Value Equation That Requires RewritingSolve $|x + 3| + 5 = 12$.

Isolate the absolute value expression on one side of the equality symbol.

$$|x + 3| + 5 = 12$$

$$|x + 3| + 5 - 5 = 12 - 5 \quad \text{Subtract 5.}$$

$$|x + 3| = 7 \quad \text{Combine like terms.}$$

$$x + 3 = 7 \quad \text{or} \quad x + 3 = -7 \quad \text{Case 1}$$

$$x = 4 \quad \text{or} \quad x = -10 \quad \text{Subtract 3.}$$

CHECK

$$|x + 3| + 5 = 12$$

$ 4 + 3 + 5 \stackrel{?}{=} 12$	Let $x = 4$.	$ -10 + 3 + 5 \stackrel{?}{=} 12$	Let $x = -10$.
$ 7 + 5 \stackrel{?}{=} 12$		$ -7 + 5 \stackrel{?}{=} 12$	
$12 = 12$ ✓ True		$12 = 12$ ✓ True	

The check confirms that the solution set is $\{-10, 4\}$.**NOW TRY**  **NOW TRY EXERCISE 6**

Solve each inequality.

(a) $|x - 1| - 4 \leq 2$

(b) $|x - 1| - 4 \geq 2$

EXAMPLE 6 Solving Absolute Value Inequalities That Require Rewriting

Solve each inequality.

(a) $|x + 3| + 5 \geq 12$

(b) $|x + 3| + 5 \leq 12$

$$|x + 3| \geq 7 \quad \text{Case 2}$$

$$|x + 3| \leq 7 \quad \text{Case 3}$$

$$x + 3 \geq 7 \quad \text{or} \quad x + 3 \leq -7$$

$$-7 \leq x + 3 \leq 7$$

$$x \geq 4 \quad \text{or} \quad x \leq -10$$

$$-10 \leq x \leq 4$$

The solution set is $(-\infty, -10] \cup [4, \infty)$.The solution set is $[-10, 4]$.**NOW TRY** **OBJECTIVE 5** Solve equations of the form $|ax + b| = |cx + d|$.*If two expressions have the same absolute value, they must either be equal or be negatives of each other.***Solving $|ax + b| = |cx + d|$**

To solve an absolute value equation of the form

$$|ax + b| = |cx + d|,$$

solve the compound equation

$$ax + b = cx + d \quad \text{or} \quad ax + b = -(cx + d).$$

NOW TRY ANSWERS

5. $\left\{-\frac{6}{5}, \frac{8}{5}\right\}$

6. (a) $[-5, 7]$

(b) $(-\infty, -5] \cup [7, \infty)$

 **NOW TRY EXERCISE 7**
Solve

$$|3x - 4| = |5x + 12|.$$

EXAMPLE 7 Solving an Equation Involving Two Absolute Values

Solve $|x + 6| = |2x - 3|$.

This equation is satisfied either if $x + 6$ and $2x - 3$ are equal to each other, or if $x + 6$ and $2x - 3$ are negatives of each other.

$$x + 6 = 2x - 3 \quad \text{or} \quad x + 6 = -(2x - 3)$$

$$x + 9 = 2x \quad \text{or} \quad x + 6 = -2x + 3$$

$$9 = x \quad \text{or} \quad 3x = -3$$

$$x = 9 \quad \text{or} \quad x = -1$$

CHECK

$$|x + 6| = |2x - 3|$$

$ 9 + 6 \stackrel{?}{=} 2(9) - 3 $	Let $x = 9$.	$ -1 + 6 \stackrel{?}{=} 2(-1) - 3 $	Let $x = -1$.
$ 15 \stackrel{?}{=} 18 - 3 $		$ 5 \stackrel{?}{=} -2 - 3 $	
$ 15 \stackrel{?}{=} 15 $		$ 5 \stackrel{?}{=} -5 $	
15 = 15 ✓	True	5 = 5 ✓	True

The check confirms that the solution set is $\{-1, 9\}$.

NOW TRY 

OBJECTIVE 6 Solve special cases of absolute value equations and inequalities.

When an absolute value equation or inequality involves a *negative constant* or 0 alone on one side, we use the following properties to solve it.

Special Properties of Absolute Value

Property 1 The absolute value of an expression can never be negative—that is, $|a| \geq 0$ for all real numbers a .

Property 2 The absolute value of an expression equals 0 only when the expression is equal to 0.

 **NOW TRY EXERCISE 8**

Solve each equation.

(a) $|3x - 8| = -2$

(b) $|7x + 12| = 0$

EXAMPLE 8 Solving Special Cases of Absolute Value Equations

Solve each equation.

(a) $|5x - 3| = -4$

See **Property 1** in the preceding box. *The absolute value of an expression can never be negative*, so there are no solutions for this equation. The solution set is \emptyset .

(b) $|7x - 3| = 0$

See **Property 2** in the preceding box. The expression $|7x - 3|$ will equal 0 *only* if $7x - 3 = 0$.

$$7x - 3 = 0$$

$$7x = 3 \quad \text{Add 3.}$$

$$x = \frac{3}{7} \quad \text{Divide by 7.}$$

Check by substituting in the original equation.

NOW TRY ANSWERS

7. $\{-8, -1\}$

8. (a) \emptyset (b) $\left\{-\frac{12}{7}\right\}$

The solution set $\left\{\frac{3}{7}\right\}$ consists of just one element.

NOW TRY 


NOW TRY EXERCISE 9

Solve each inequality.

- (a) $|x| > -10$
 (b) $|4x + 1| + 5 < 4$
 (c) $|x - 2| - 3 \leq -3$

EXAMPLE 9 Solving Special Cases of Absolute Value Inequalities

Solve each inequality.

(a) $|x| \geq -4$

The absolute value of a number is always greater than or equal to 0. (Property 1)
 Thus, $|x| \geq -4$ is true for *all* real numbers. The solution set is $(-\infty, \infty)$.

(b) $|x + 6| - 3 < -5$

$|x + 6| < -2$ Add 3 to each side.

There is no number whose absolute value is less than -2 , so this inequality has no solution. The solution set is \emptyset .

(c) $|x - 7| + 4 \leq 4$

$|x - 7| \leq 0$ Subtract 4 from each side.

The value of $|x - 7|$ will never be less than 0. However, $|x - 7|$ will *equal* 0 when $x = 7$. Therefore, the solution set is $\{7\}$.

NOW TRY 
OBJECTIVE 7 Solve an application involving relative error.

Absolute value is used to find the **relative error**, or **tolerance**, in a measurement. If x represents the actual measurement and x_t represents the expected measurement, then taking the absolute value of the difference of x and x_t , divided by x_t , gives the relative error in x .

$$\text{relative error in } x = \left| \frac{x - x_t}{x_t} \right|$$

In quality control situations, the relative error must often be less than some predetermined amount.



NOW TRY EXERCISE 10

Suppose a machine filling *quart* milk cartons is set for a relative error that *is no greater than* 0.03. How many ounces may a filled carton contain?

EXAMPLE 10 Solving an Application Involving Relative Error

Suppose a machine filling *quart* milk cartons is set for a relative error that *is no greater than* 0.05. How many ounces may a filled carton contain?

Here $x_t = 32$ oz (because 1 qt = 32 oz) and the relative error = 0.05. We must find x , the actual measure of a filled carton.

$$\left| \frac{x - 32}{32} \right| \leq 0.05 \quad \begin{array}{l} \text{Substitute the given values.} \\ \text{Is no greater than translates as } \leq. \end{array}$$

$$-0.05 \leq \frac{x - 32}{32} \leq 0.05 \quad \text{Case 3}$$

$$-1.6 \leq x - 32 \leq 1.6 \quad \text{Multiply by 32.}$$

$$30.4 \leq x \leq 33.6 \quad \text{Add 32.}$$

The filled carton may contain between 30.4 and 33.6 oz, inclusive.

NOW TRY 
NOW TRY ANSWERS

9. (a) $(-\infty, \infty)$ (b) \emptyset (c) $\{2\}$
 10. between 31.04 and 32.96 oz, inclusive

1.7 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Time management can be a challenge for students.

Review Study Skill 6, Managing Your Time.

Concept Check Match each absolute value equation or inequality in Column I with the graph of its solution set in Column II.

I	II	I	II
1. $ x = 5$	A.	2. $ x = 9$	A.
$ x < 5$	B.	$ x > 9$	B.
$ x > 5$	C.	$ x \geq 9$	C.
$ x \leq 5$	D.	$ x < 9$	D.
$ x \geq 5$	E.	$ x \leq 9$	E.

3. **Concept Check** How many solutions will $|ax + b| = k$ have for each situation?

- (a) $k = 0$ (b) $k > 0$ (c) $k < 0$

4. **Concept Check** If $k < 0$, what is the solution set of each of the following?

- (a) $|x - 1| < k$ (b) $|x - 1| > k$ (c) $|x - 1| = k$


Solve each equation. See Example 1.

- | | | |
|------------------------------------------|-----------------------------------------|------------------------------------------|
| 5. $ x = 12$ | 6. $ x = 14$ | 7. $ 4x = 20$ |
| 8. $ 5x = 30$ | 9. $ x - 3 = 9$ | 10. $ x - 5 = 13$ |
| 11. $ 2x - 1 = 11$ | 12. $ 2x + 3 = 19$ | 13. $ 4x - 5 = 17$ |
| 14. $ 5x - 1 = 21$ | 15. $ 2x + 5 = 14$ | 16. $ 2x - 9 = 18$ |
| 17. $ -3x + 8 = 1$ | 18. $ -6x + 5 = 4$ | 19. $\left 12 - \frac{1}{2}x\right = 6$ |
| 20. $\left 14 - \frac{1}{3}x\right = 8$ | 21. $ 0.5x = 6$ | 22. $ 0.3x = 9$ |
| 23. $\left \frac{1}{2}x + 3\right = 2$ | 24. $\left \frac{2}{3}x - 1\right = 5$ | 25. $\left 1 + \frac{3}{4}x\right = 7$ |
| 26. $\left 2 - \frac{5}{2}x\right = 14$ | 27. $ 0.02x - 1 = 2.50$ | 28. $ 0.04x - 3 = 5.96$ |

Solve each inequality. Graph the solution set, and write it using interval notation. See Example 2.

- | | | |
|---------------------|-------------------------|------------------------|
| 29. $ x > 3$ | 30. $ x > 5$ | 31. $ x \geq 4$ |
| 32. $ x \geq 6$ | 33. $ r + 5 \geq 20$ | 34. $ x + 4 \geq 8$ |
| 35. $ 5x + 2 > 10$ | 36. $ 4x + 1 \geq 21$ | 37. $ 3 - x > 5$ |
| 38. $ 5 - x > 3$ | 39. $ -5x + 3 \geq 12$ | 40. $ -2x - 4 \geq 5$ |

41. Concept Check The graph of the solution set of $|2x + 1| = 9$ is given here.



Without doing any algebraic work, graph the solution set of each inequality, referring to the graph shown.

(a) $|2x + 1| < 9$ (b) $|2x + 1| > 9$

42. Concept Check The graph of the solution set of $|3x - 4| < 5$ is given here.



Without doing any algebraic work, graph the solution set of the following, referring to the graph shown.

(a) $|3x - 4| = 5$ (b) $|3x - 4| > 5$

Solve each inequality. Graph the solution set, and write it using interval notation. See Example 3. (Hint: Compare the answers with those in Exercises 29–40.)

43. $|x| \leq 3$

44. $|x| \leq 5$

45. $|x| < 4$

46. $|x| < 6$

47. $|r + 5| < 20$

48. $|x + 4| < 8$

49. $|5x + 2| \leq 10$

50. $|4x + 1| < 21$

51. $|3 - x| \leq 5$

52. $|5 - x| \leq 3$

53. $|-5x + 3| < 12$

54. $|-2x - 4| < 5$

Solve each equation or inequality. In Exercises 55–66, graph the solution set. See Examples 1–4.

55. $|-4 + x| > 9$

56. $|-3 + x| > 8$

57. $|3x + 2| < 11$

58. $|2x - 1| < 7$

59. $|7 + 2x| = 5$

60. $|9 - 3x| = 3$

61. $|3x - 1| \leq 11$

62. $|2x - 6| \leq 6$

63. $|-6x - 6| \leq 1$

64. $|-2x - 6| \leq 5$

65. $|-8 + x| \leq 5$

66. $|-4 + x| \leq 9$

67. $|10 - 12x| \geq 4$

68. $|8 - 10x| \geq 2$

69. $|3(x - 1)| = 8$

70. $|7(x - 2)| = 4$

71. $|0.1x - 1| > 3$

72. $|0.1x + 1| > 2$

73. $|x + 2| = 5 - 2$

74. $|x + 3| = 12 - 2$

75. $3|x - 6| = 9$

76. $5|x - 4| = 5$

77. $|2 - 0.2x| = 2$

78. $|5 - 0.5x| = 4$

Solve each equation or inequality. See Examples 5 and 6.

79. $|x| - 1 = 4$

80. $|x| + 3 = 10$

81. $|x + 4| + 1 = 2$

82. $|x + 5| - 2 = 12$

83. $|2x + 1| + 3 > 8$

84. $|6x - 1| - 2 > 6$

85. $|x + 5| - 6 \leq -1$

86. $|x - 2| - 3 \leq 4$

87. $|0.1x - 2.5| + 0.3 \geq 0.8$

88. $|0.5x - 3.5| + 0.2 \geq 0.6$

89. $\left| \frac{1}{2}x + \frac{1}{3} \right| + \frac{1}{4} = \frac{3}{4}$

90. $\left| \frac{2}{3}x + \frac{1}{6} \right| + \frac{1}{2} = \frac{5}{2}$

Solve each equation. See Example 7.

91. $|3x + 1| = |2x + 4|$

92. $|7x + 12| = |x - 8|$

93. $\left|x - \frac{1}{2}\right| = \left|\frac{1}{2}x - 2\right|$

94. $\left|\frac{2}{3}x - 2\right| = \left|\frac{1}{3}x + 3\right|$

95. $|6x| = |9x + 1|$

96. $|13x| = |2x + 1|$

97. $|2x - 6| = |2x + 11|$

98. $|3x - 1| = |3x + 9|$

Solve each equation or inequality. See Examples 8 and 9.

99. $|x| \geq -10$

100. $|x| \geq -15$

101. $|12t - 3| = -8$

102. $|13x + 1| = -3$

103. $|4x + 1| = 0$

104. $|6x - 2| = 0$

105. $|2x - 1| = -6$

106. $|8x + 4| = -4$

107. $|x + 5| > -9$

108. $|x + 9| > -3$

109. $|7x + 3| \leq 0$

110. $|4x - 1| \leq 0$

111. $|5x - 2| = 0$

112. $|7x + 4| = 0$

113. $|x - 2| + 3 \geq 2$

114. $|x - 4| + 5 \geq 4$

115. $|10x + 7| + 3 < 1$

116. $|4x + 1| - 2 < -5$

Determine the number of ounces a filled carton of the given size may contain for the given relative error. See Example 10.

$$\left|\frac{x - x_t}{x_t}\right| = \text{relative error in } x$$

x represents actual measurement.

x_t represents expected measurement.

117. 64-oz carton; relative error no greater than 0.05

118. 24-oz carton; relative error no greater than 0.05

119. 32-oz carton; relative error no greater than 0.02

120. 36-oz carton; relative error no greater than 0.03

In later courses in mathematics, it is sometimes necessary to find an interval in which x must lie in order to keep y within a given difference of some number. For example, suppose

$$y = 2x + 1$$

and we want y to be within 0.01 unit of 4. This criterion can be written as

$$|y - 4| < 0.01.$$

Solving this inequality shows that x must lie in the interval (1.495, 1.505) to satisfy the requirement.

Find the open interval in which x must lie in order for the given condition to hold.

121. $y = 2x + 1$, and the difference of y and 1 is less than 0.1.

122. $y = 4x - 6$, and the difference of y and 2 is less than 0.02.

123. $y = 4x - 8$, and the difference of y and 3 is less than 0.001.

124. $y = 5x + 12$, and the difference of y and 4 is less than 0.0001.

Work each problem.

125. Dr. Mosely has determined that 99% of the babies he has delivered have weighed x pounds, where

$$|x - 8.3| < 1.5.$$

What range of weights corresponds to this inequality?

126. The Celsius temperatures x on Mars approximately satisfy the inequality

$$|x + 85| \leq 55.$$

What range of temperatures corresponds to this inequality?

127. The recommended daily intake (RDI) of calcium for females aged 19–50 is 1000 mg. Write this statement as an absolute value inequality, with x representing the RDI, to express the RDI plus or minus 100 mg. Solve the inequality. (Data from National Academy of Sciences—Institute of Medicine.)
128. The average clotting time of blood is 7.45 sec, with a variation of plus or minus 3.6 sec. Write this statement as an absolute value inequality, with x representing the time. Solve the inequality.

RELATING CONCEPTS For Individual or Group Work (Exercises 129–132)

The 10 tallest buildings in Houston, Texas, are listed along with their heights.

Building	Height (in feet)
JPMorgan Chase Tower	1002
Wells Fargo Plaza	992
Williams Tower	901
Bank of America Center	780
Texaco Heritage Plaza	762
609 Main at Texas	757
Enterprise Plaza	756
Centerpoint Energy Plaza	741
1600 Smith St.	732
Fulbright Tower	725



Data from *The World Almanac and Book of Facts*.

Use this information to work Exercises 129–132 in order.

129. To find the average of a group of numbers, we add the numbers and then divide by the number of numbers. Use a calculator to find the average of the heights.
130. Let k represent the average height of these buildings. If a height x satisfies the inequality

$$|x - k| < t,$$

then the height is said to be within t feet of the average. Using the result from **Exercise 129**, list the buildings that are within 50 ft of the average.

131. Repeat **Exercise 130**, but list the buildings that are within 95 ft of the average.
132. Work each of the following.
- Write an absolute value inequality that describes the height of a building that is *not* within 95 ft of the average.
 - Solve the inequality from part (a).
 - Use the result of part (b) to list the buildings that are not within 95 ft of the average.
 - Confirm that the answer to part (c) makes sense by comparing it with the answer to **Exercise 131**.

SUMMARY EXERCISES Solving Linear and Absolute Value Equations and Inequalities

Solve each equation or inequality. Give the solution set in set notation for equations and in interval notation for inequalities.

1. $4x + 1 = 49$
2. $|x - 1| = 6$
3. $6x - 9 = 12 + 3x$
4. $3x + 7 = 9 + 8x$
5. $|x + 3| = -4$
6. $2x + 1 \leq x$
7. $8x + 2 \geq 5x$
8. $4(x - 11) + 3x = 20x - 31$
9. $2x - 1 = -7$
10. $|3x - 7| - 4 = 0$
11. $6x - 5 \leq 3x + 10$
12. $|5x - 8| + 9 \geq 7$
13. $9x - 3(x + 1) = 8x - 7$
14. $|x| \geq 8$
15. $9x - 5 \geq 9x + 3$
16. $13x - 5 > 13x - 8$
17. $|x| < 5.5$
18. $4x - 1 = 12 + x$
19. $\frac{2}{3}x + 8 = \frac{1}{4}x$
20. $-\frac{5}{8}x \geq -20$
21. $\frac{1}{4}x < -6$
22. $\frac{1}{2} \leq \frac{2}{3}x \leq \frac{5}{4}$
23. $\frac{3}{5}x - \frac{1}{10} = 2$
24. $\frac{x}{6} - \frac{3x}{5} = x - 86$
25. $x + 9 + 7x = 4(3 + 2x) - 3$
26. $6 - 3(2 - x) < 2(1 + x) + 3$
27. $-6 \leq \frac{3}{2} - x \leq 6$
28. $\frac{x}{4} - \frac{2x}{3} = -10$
29. $|5x + 1| \leq 0$
30. $5x - (3 + x) \geq 2(3x + 1)$
31. $-2 \leq 3x - 1 \leq 8$
32. $-1 \leq 6 - x \leq 5$
33. $|7x - 1| = |5x + 3|$
34. $|x + 2| = |x + 4|$
35. $|1 - 3x| \geq 4$
36. $7x - 3 + 2x = 9x - 8x$
37. $-(x + 4) + 2 = 3x + 8$
38. $|x - 1| < 7$
39. $|2x - 3| > 11$
40. $|5 - x| < 4$
41. $|x - 1| \geq -6$
42. $|2x - 5| = |x + 4|$
43. $8x - (1 - x) = 3(1 + 3x) - 4$
44. $8x - (x + 3) = -(2x + 1) - 12$
45. $|x - 5| = |x + 9|$
46. $|x + 2| < -3$
47. $2x + 1 > 5$ or $3x + 4 < 1$
48. $1 - 2x \geq 5$ and $7 + 3x \geq -2$

Chapter 1

Summary

STUDY SKILLS REMINDER

How do you best prepare for a test? **Review Study Skill 7, Reviewing a Chapter.**

Key Terms

<p>1.1</p> <p>linear (first-degree) equation in one variable solution solution set equivalent equations conditional equation identity contradiction</p>	<p>1.2</p> <p>mathematical model formula percent percent increase percent decrease</p> <p>1.4</p> <p>consecutive integers consecutive even integers consecutive odd integers</p>	<p>1.5</p> <p>inequality interval linear inequality in one variable equivalent inequalities three-part inequality</p>	<p>1.6</p> <p>intersection compound inequality union</p> <p>1.7</p> <p>absolute value equation absolute value inequality</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------

New Symbols

$\%$ percent	∞ infinity	(a, b) interval notation for $a < x < b$	\cap set intersection
1° one degree	$-\infty$ negative infinity	$[a, b]$ interval notation for $a \leq x \leq b$	\cup set union
	$(-\infty, \infty)$ the set of real numbers		

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- An **equation** is
 - an algebraic expression
 - an expression that contains fractions
 - an expression that uses any of the four basic operations or the operation of raising to powers or taking roots on any collection of variables and numbers formed according to the rules of algebra
 - a statement that two algebraic expressions are equal.
- A linear equation that is a **conditional equation** has
 - no solution
 - one solution
 - two solutions
 - infinitely many solutions.
- An **inequality** is
 - a statement that two algebraic expressions are equal
 - a point on a number line
 - an equation with no solutions
 - a statement consisting of algebraic expressions related by $<$, \leq , $>$, or \geq .
- Interval notation** is
 - a point on a number line
 - a special notation for describing a point on a number line
 - a way to use symbols to describe an interval on a number line
 - a notation to describe unequal quantities.
- The **intersection** of two sets A and B is the set of elements that belong
 - to both A and B
 - to either A or B , or both
 - to either A or B , but not both
 - to just A .
- The **union** of two sets A and B is the set of elements that belong
 - to both A and B
 - to either A or B , or both
 - to either A or B , but not both
 - to just B .

ANSWERS

- D; Examples: $2a + 3 = 7$, $3y = -8$, $x^2 = 4$
- B; Example: $x - 4 = 6$ is a conditional equation with one solution, 10.
- D; Examples: $x < 5$, $7 + 2k \geq 11$, $-5 < 2z - 1 \leq 3$
- C; Examples: $(-\infty, 5]$, $(1, \infty)$, $[-3, 3)$
- A; Example: If $A = \{2, 4, 6, 8\}$ and $B = \{1, 2, 3\}$, then $A \cap B = \{2\}$.
- B; Example: Using the sets A and B from Answer 5, $A \cup B = \{1, 2, 3, 4, 6, 8\}$.

Quick Review

CONCEPTS

1.1 Linear Equations in One Variable**Addition and Multiplication Properties of Equality**

The same number may be added to (or subtracted from) each side of an equation without changing the solution set.

Similarly, each side of an equation may be multiplied (or divided) by the same nonzero number without changing the solution set.

Solving a Linear Equation in One Variable

Step 1 Simplify each side separately.

- Clear any parentheses.
- Clear any fractions or decimals.
- Combine like terms.

Step 2 Isolate the variable terms on one side.

Step 3 Isolate the variable.

Step 4 Check.

EXAMPLES

Solve. $x - 4 = 11$

$$x - 4 + 4 = 11 + 4$$

$$x = 15$$

Add 4 to each side by the addition property of equality.

Solution set: $\{15\}$

Solve. $4(8 - 3x) = 32 - 8(x + 2)$

$$32 - 12x = 32 - 8x - 16$$

$$32 - 12x = 16 - 8x$$

$$32 - 12x + 12x = 16 - 8x + 12x$$

$$32 = 16 + 4x$$

$$32 - 16 = 16 + 4x - 16$$

$$16 = 4x$$

$$\frac{16}{4} = \frac{4x}{4}$$

$$4 = x$$

Distributive property
Combine like terms.

Add 12x.

Combine like terms.

Subtract 16.

Combine like terms.

Divide by 4.

CHECK $4(8 - 3x) = 32 - 8(x + 2)$

$$4[8 - 3(4)] \stackrel{?}{=} 32 - 8(4 + 2)$$

$$4[8 - 12] \stackrel{?}{=} 32 - 8(6)$$

$$4[-4] \stackrel{?}{=} 32 - 48$$

$$-16 = -16 \quad \checkmark$$

Original equation

Let $x = 4$.

Multiply. Add.

Subtract. Multiply.

True

Solution set: $\{4\}$

1.2 Formulas and Percent**Solving a Formula for a Specified Variable (Solving a Literal Equation)**

Step 1 Clear any parentheses and fractions.

Step 2 Isolate all terms with the specified variable.

Step 3 Isolate the specified variable.

Solve $\mathcal{A} = \frac{1}{2}bh$ for h .

$$\mathcal{A} = \frac{1}{2}bh$$

$$2\mathcal{A} = 2\left(\frac{1}{2}bh\right)$$

$$2\mathcal{A} = bh$$

$$\frac{2\mathcal{A}}{b} = h, \quad \text{or} \quad h = \frac{2\mathcal{A}}{b}$$

Multiply by 2.

$$2 \cdot \frac{1}{2} = 1$$

Divide by b .

1.3 Applications of Linear Equations**Solving an Applied Problem**

Step 1 Read the problem.

Step 2 Assign a variable.

How many liters of 30% alcohol solution and 80% alcohol solution must be mixed to obtain 100 L of 50% alcohol solution?

Let x = number of liters of 30% solution needed.

Then $100 - x$ = number of liters of 80% solution needed.

Liters of Solution	Percent (as a decimal)	Liters of Pure Alcohol
x	0.30	$0.30x$
$100 - x$	0.80	$0.80(100 - x)$
100	0.50	$0.50(100)$

CONCEPTS

- Step 3** Write an equation.
- Step 4** Solve the equation.
- Step 5** State the answer.
- Step 6** Check.

EXAMPLES

From the last column of the table, write an equation.

$$0.30x + 0.80(100 - x) = 0.50(100)$$

$$0.30x + 80 - 0.80x = 50 \quad \text{Distributive property; Multiply.}$$

$$100(0.30x + 80 - 0.80x) = 100(50) \quad \text{Multiply by 100.}$$

$$30x + 8000 - 80x = 5000 \quad \text{Distributive property; Multiply.}$$

$$-50x + 8000 = 5000 \quad \text{Combine like terms.}$$

$$-50x = -3000 \quad \text{Subtract 8000.}$$

$$x = 60 \quad \text{Divide by -50.}$$

The solution of the equation is 60.

60 L of 30% solution and $100 - 60 = 40$ L of 80% solution are needed.

$$0.30(60) + 0.80(100 - 60) = 50 \text{ is true.}$$

1.4 Further Applications of Linear Equations

To solve a uniform motion problem, draw a sketch and make a table. Use the distance formula.

$$d = rt, \text{ or } rt = d$$

Two cars start from towns 400 mi apart and travel toward each other. They meet after 4 hr. Find the rate of each car if one travels 20 mph faster than the other.


Let x = rate of the slower car in miles per hour.

Then $x + 20$ = rate of the faster car.

Use the information in the problem and $d = rt$ in the form $rt = d$ to complete a table.

	Rate	Time	Distance	
Slower Car	x	4	$4x$	
Faster Car	$x + 20$	4	$4(x + 20)$	
			400	← Total

A sketch shows that the sum of the distances, $4x$ and $4(x + 20)$, must be 400.



$$4x + 4(x + 20) = 400$$

$$4x + 4x + 80 = 400 \quad \text{Distributive property}$$

$$8x + 80 = 400 \quad \text{Combine like terms.}$$

$$8x = 320 \quad \text{Subtract 80.}$$

$$x = 40 \quad \text{Divide by 8.}$$

The slower car travels 40 mph, and the faster car travels $40 + 20 = 60$ mph.

$$4(40) + 4(40 + 20) = 400 \text{ is true.}$$

CONCEPTS

1.5 Linear Inequalities in One Variable**Solving a Linear Inequality in One Variable****Step 1** Simplify each side separately.

- Clear any parentheses.
- Clear any fractions or decimals.
- Combine like terms.

Step 2 Isolate the variable terms on one side.**Step 3** Isolate the variable (here x) to write the inequality in one of these forms.

$$x < k, \quad x \leq k, \quad x > k, \quad \text{or} \quad x \geq k$$

If an inequality is multiplied or divided by a negative number, the direction of the inequality symbol must be reversed.

To solve a three-part inequality, work with all three parts at the same time.

1.6 Set Operations and Compound Inequalities**Solving a Compound Inequality****Step 1** Solve each inequality in the compound inequality individually.**Step 2** If the inequalities are joined with *and*, then the solution set is the intersection of the two individual solution sets.

If the inequalities are joined with *or*, then the solution set is the union of the two individual solution sets.

EXAMPLES

Solve each inequality.

$$3(x + 2) - 5x \leq 12$$

$$3x + 6 - 5x \leq 12 \quad \text{Distributive property}$$


$$-2x + 6 \leq 12 \quad \text{Combine like terms.}$$

$$-2x + 6 - 6 \leq 12 - 6 \quad \text{Subtract 6.}$$

$$-2x \leq 6 \quad \text{Combine like terms.}$$

$$\frac{-2x}{-2} \geq \frac{6}{-2} \quad \text{Divide by } -2.$$

$$x \geq -3 \quad \text{Change } \leq \text{ to } \geq.$$

Solution set:  $[-3, \infty)$


$$-4 < 2x + 3 \leq 7$$

$$-4 - 3 < 2x + 3 - 3 \leq 7 - 3 \quad \text{Subtract 3.}$$

$$-7 < 2x \leq 4$$

$$\frac{-7}{2} < \frac{2x}{2} \leq \frac{4}{2} \quad \text{Divide by 2.}$$


$$-\frac{7}{2} < x \leq 2$$

Solution set:  $(-\frac{7}{2}, 2]$

Solve each compound inequality.

$$x + 1 > 2 \quad \text{and} \quad 2x < 6$$

$$x > 1 \quad \text{and} \quad x < 3$$


Solution set:  $(1, 3)$

$$2(x + 3) - 2 \leq 4 \quad \text{or} \quad -4x \leq -16$$

$$2x + 6 - 2 \leq 4 \quad \text{or} \quad \frac{-4x}{-4} \geq \frac{-16}{-4}$$

$$2x + 4 \leq 4 \quad \text{or} \quad x \geq 4$$

$$x \leq 0$$

Solution set:  $(-\infty, 0] \cup [4, \infty)$

CONCEPTS

1.7 Absolute Value Equations and Inequalities

Solving Absolute Value Equations and Inequalities

Let k be a positive number.

Case 1 To solve $|ax + b| = k$, solve the compound equation

$$ax + b = k \quad \text{or} \quad ax + b = -k.$$

Case 2 To solve $|ax + b| > k$, solve the compound inequality

$$ax + b > k \quad \text{or} \quad ax + b < -k.$$

Case 3 To solve $|ax + b| < k$, solve the compound inequality

$$-k < ax + b < k.$$

To solve an absolute value equation of the form

$$|ax + b| = |cx + d|,$$

solve the compound equation

$$ax + b = cx + d \quad \text{or} \quad ax + b = -(cx + d).$$

Special Properties of Absolute Value

Property 1 The absolute value of an expression can never be negative—that is, $|a| \geq 0$ for all real numbers a .

Property 2 The absolute value of an expression equals 0 only when the expression is equal to 0.

EXAMPLES

Solve each equation or inequality.

$$|x - 7| = 3 \quad \text{Case 1}$$

$$x - 7 = 3 \quad \text{or} \quad x - 7 = -3$$


$$x = 10 \quad \text{or} \quad x = 4 \quad \text{Add 7.}$$



$$|x - 7| > 3 \quad \text{Case 2}$$

$$x - 7 > 3 \quad \text{or} \quad x - 7 < -3$$


$$x > 10 \quad \text{or} \quad x < 4 \quad \text{Add 7.}$$



$$|x - 7| < 3 \quad \text{Case 3}$$

$$-3 < x - 7 < 3$$

$$4 < x < 10 \quad \text{Add 7.}$$




$$|x + 2| = |2x - 6|$$

$$x + 2 = 2x - 6 \quad \text{or} \quad x + 2 = -(2x - 6)$$

$$-x = -8 \quad \text{or} \quad x + 2 = -2x + 6$$

$$x = 8 \quad \text{or} \quad 3x = 4$$

$$x = \frac{4}{3}$$



$$\left. \begin{array}{l} |x - 2| = -4 \quad \text{Solution set: } \emptyset \\ |x| \geq -1 \quad \text{Solution set: } (-\infty, \infty) \\ |2x - 3| < -6 \quad \text{Solution set: } \emptyset \end{array} \right\} \text{Property 1}$$

$$|3x - 9| = 0 \quad \text{Property 2}$$

$$3x - 9 = 0 \quad |a| = 0 \text{ implies } a = 0.$$

$$3x = 9 \quad \text{Add 9.}$$

$$x = 3 \quad \text{Divide by 3.}$$



Chapter 1

Review Exercises

1.1 Solve each equation.

1. $-(8 + 3x) + 5 = 2x + 6$

2. $-\frac{3}{4}x = -12$

3. $\frac{2x + 1}{3} - \frac{x - 1}{4} = 0$

4. $5(2x - 3) = 6(x - 1) + 4x$

Solve each equation. Decide whether the equation is a conditional equation, an identity, or a contradiction.

5. $10x - 3(2x - 4) = 4(x + 3)$

6. $13x - (x + 7) = 12x - 13$

7. $x + 6(x - 1) - (4 - x) = -10$

8. $5 + 3(2x + 6) = 5 + 3(2x + 6)$

1.2 Solve for the specified variable.

9. $V = LWH$ for L


10. $\mathcal{A} = \frac{1}{2}h(b + B)$ for b

11. $P = a + b + c + B$ for c

12. $4x + 7y = 9$ for y


1.2, 1.3 Solve each problem.

13. A rectangular solid has a volume of
- 180 ft^3
- . Its length is 6 ft and its width is 5 ft. Find its height.



14. The number of students attending college in 2015 was 20.2 million. This number is expected to increase to 23.1 million by the year 2024. To the nearest tenth, what is the expected percent increase? (Data from National Center for Education Statistics.)
15. Find the annual simple-interest rate that Halina is earning if the principal of \$30,000 earns \$3900 interest in 4 yr.
16. If the Fahrenheit temperature is 77° , what is the corresponding Celsius temperature?



At the end of 2016, there were 395.9 million wireless subscriptions in the United States. (Data from CTIA.) The circle graph shows market share by carrier during the fourth quarter of 2016.

2016 Wireless Subscriptions Market Share

Data from Strategy Analytics; FierceWireless.

17. To the nearest tenth of a million, about how many Verizon subscriptions were there?
18. To the nearest tenth of a million, about how many T-Mobile subscriptions were there?

34. An 85-mi trip to the beach took Susan 2 hr. During the second hour, a rainstorm caused her to average 7 mph less than she traveled during the first hour. Find her average rate for the first hour.
35. Find the measure of each angle in the triangle.
36. Find the measure of each marked angle.



37. Find three consecutive integers such that the sum of the first and third is 49 less than three times the second.
38. Find three consecutive even integers such that the sum of the second and third is 44 more than the first.

1.5 Solve each inequality. Give the solution set in interval form.

39. $-\frac{2}{3}x < 6$
40. $-5x - 4 \geq 11$
41. $\frac{6x + 3}{-4} < -3$
42. $5 - (6 - 4x) \geq 2x - 7$
43. $8 \leq 3x - 1 < 14$
44. $\frac{5}{3}(x - 2) + \frac{2}{5}(x + 1) > 1$

Solve each problem.

45. Dr. Paul Donohue writes a syndicated column in which readers question him on a variety of health topics. Reader C. J. wrote, “Many people say they can weigh more because they have a large frame. How is frame size determined?” Here is Dr. Donohue’s response:


“For a man, a wrist circumference between 6.75 and 7.25 in. [inclusive] indicates a medium frame. Anything above is a large frame and anything below, a small frame.”

(Source: Dr. Donohue, © 2004. Used by permission of North America Syndicate.)

Using x to represent wrist circumference in inches, write an inequality or a three-part inequality that represents wrist circumference for a male with the following.

- (a) a small frame (b) a medium frame (c) a large frame

46. The perimeter of a rectangular playground must be no greater than 120 m. One dimension of the playground must be 22 m. Find the possible lengths of the other dimension.



47. A group of college students wanted to buy tickets to attend a performance of *Motown the Musical* at the Lunt-Fontanne Theatre in New York City. They could buy student mezzanine seats for a group rate of \$51 per ticket for 15 tickets or more. If they had \$1600 available to spend, how many tickets could they purchase at this price? (Data from www.broadway.com)



48. To pass algebra, a student must have an average of at least 70 on five tests. On the first four tests, a student has scores of 75, 79, 64, and 71. What possible scores on the fifth test will guarantee the student a passing grade in the class?

1.6 Let $A = \{a, b, c, d\}$, $B = \{a, c, e, f\}$, and $C = \{a, e, f, g\}$. Find each set.

49. $A \cap B$ 50. $A \cap C$ 51. $B \cup C$ 52. $A \cup C$

Solve each compound inequality. Graph the solution set, and write it using interval notation.

53. $x > 6$ and $x < 9$ 54. $x + 4 > 12$ and $x - 2 < 12$
 55. $x > 5$ or $x \leq -3$ 56. $x \geq -2$ or $x < 2$
 57. $x - 4 > 6$ and $x + 3 \leq 10$ 58. $-5x + 1 \geq 11$ or $3x + 5 \geq 26$

Express each set in simplest interval form.

59. $(-3, \infty) \cap (-\infty, 4]$ 60. $(-\infty, 6] \cap (-\infty, 2]$
 61. $(4, \infty) \cup (9, \infty)$ 62. $(1, 2) \cup (1, \infty)$

1.7 Solve each equation or inequality.

63. $|x| = 7$ 64. $|x + 2| = 9$
 65. $|3x - 7| = 8$ 66. $|x - 4| = -12$
 67. $|2x - 7| + 4 = 11$ 68. $|4x + 2| - 7 = -3$
 69. $|3x + 1| = |x + 2|$ 70. $|2x - 1| = |2x + 3|$
 71. $|x| < 14$ 72. $|-x + 6| \leq 7$
 73. $|2x + 5| \leq 1$ 74. $|x + 1| \geq -3$
 75. $|3 - 4x| + 7 < -8$ 76. $|-8 - 3x| - 7 > -8$

Determine the number of ounces a filled carton of the given size may contain for the given relative error.

$$\left| \frac{x - x_t}{x_t} \right| = \text{relative error in } x \quad \begin{array}{l} x \text{ represents actual measurement.} \\ x_t \text{ represents expected measurement.} \end{array}$$

77. 48-oz carton; relative error no greater than 0.03
 78. 32-oz carton; relative error no greater than 0.01

Solve.

1. $5 - (6 - 4x) > 2x - 5$ 2. $ak + bt = 6r$ for k
 3. $x + 4 < 7$ and $x + 5 \geq 3$ 4. $\frac{4x + 2}{4} + \frac{3x - 1}{8} = \frac{x + 6}{16}$
 5. $|3x + 6| \geq 0$ 6. $-5x \geq -10$
 7. $|3x + 2| + 4 = 9$ 8. $0.05x + 0.03(1200 - x) = 42$
 9. $|x + 3| \leq 13$ 10. $\frac{3}{4}(x - 2) - \frac{1}{3}(5 - 2x) < -2$

11. $-4 < 3 - 2x < 9$
12. $-0.3x + 2.1(x - 4) \leq -6.6$
13. $|5x - 1| > 14$
14. $x \geq -2$ or $x < 4$
15. $|x - 1| = |2x + 3|$
16. $\frac{3x}{5} - \frac{x}{2} = 3$
17. $|3x - 7| = 4$
18. $5(2x - 7) = 2(5x + 3)$
19. $-5x < -30$ and $-7x > -56$
20. $-5x + 1 \geq 11$ or $3x + 5 \geq 26$

Solve each problem.

21. A newspaper recycling collection bin is in the shape of a box 1.5 ft wide and 5 ft long. If the volume of the bin is 75 ft^3 , find the height.
22. The sum of the first and third of three consecutive integers is 47 more than the second integer. What are the integers?
23. To qualify for a company pension plan, an employee must average at least \$1000 per month in earnings. During the first four months of the year, an employee made \$900, \$1200, \$1040, and \$760. What possible amounts earned during the fifth month will qualify the employee for a pension plan?
24. How many liters of a 20% solution of a chemical should be mixed with 10 L of a 50% solution to obtain a 40% mixture?

Chapter 1

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

► View the complete solutions to all Chapter Test exercises in MyLab Math.

STUDY SKILLS REMINDER

Using test-taking strategies can help you improve your test scores. **Review Study Skill 8, Taking Math Tests.**

Solve each equation.

1. $3(2x - 2) - 4(x + 6) = 3x + 8 + x$
2. $0.08x + 0.06(x + 9) = 1.24$
3. Solve each equation. Then determine whether the equation is a *conditional equation*, an *identity*, or a *contradiction*.
- (a) $3x - (2 - x) + 4x + 2 = 8x + 3$
- (b) $\frac{x}{3} + 7 = \frac{5x}{6} - 2 - \frac{x}{2} + 9$
- (c) $-4(2x - 6) = 5x + 24 - 7x$
- (d) $\frac{x + 6}{10} + \frac{x - 4}{15} = \frac{x + 2}{6}$
4. Solve $V = \frac{1}{3}bh$ for h .
5. Solve $-3x + 2y = 6$ for y .


Solve each problem.

6. The 2017 Brickyard 400 (mile) race was won by Kasey Kahne, who averaged 114.384 mph. What was his time to the nearest thousandth of an hour? (Data from *The World Almanac and Book of Facts*.)
7. A certificate of deposit pays \$456.25 in simple interest for 1 yr on a principal of \$36,500. What is the rate of interest?
8. The final 2017 standings of the American League East division in Major League Baseball are shown in the table. Find the winning percentage of each team.
- (a) NY Yankees (b) Tampa Bay
- (c) Toronto (d) Baltimore


Team	W	L	Pct.
Boston	93	69	.574
NY Yankees	91	71	
Tampa Bay	80	82	
Toronto	76	86	
Baltimore	75	87	

Data from www.MLB.com

9. Tyler invested some money at 1.5% simple interest and some at 2.5% simple interest. The total amount of his investments was \$28,000, and the interest he earned during the first year was \$620. How much did he invest at each rate?
10. After working 1 yr at his new job, Kevin got a raise from \$11.25 per hour to \$12.75 per hour. What was the percent increase in his hourly wage to the nearest tenth?
11. How many liters of a 20% alcohol solution should be added to 40 L of a 50% alcohol solution to make a 30% solution?
12. Two cars leave from the same point at the same time, traveling in opposite directions. One travels 15 mph slower than the other. After 6 hr, they are 630 mi apart. Find the rate of each car.



13. Find the measure of each angle in the triangle.



Solve each inequality. Graph the solution set, and write it using interval notation.

14. $4 - 6(x + 3) \leq -2 - 3(x + 6) + 3x$ 15. $-\frac{4}{7}x > -16$
16. $-1 < 3x - 4 < 2$ 17. $-6 \leq \frac{4}{3}x - 2 \leq 2$
18. Which one of the following inequalities is equivalent to $x < -3$?
- A. $-3x < 9$ B. $-3x > -9$ C. $-3x > 9$ D. $-3x < -9$
19. Justin must have an average of at least 80 on the four tests in a course to get a B. He had scores of 83, 76, and 79 on the first three tests. What possible scores on the fourth test would guarantee him a B in the course?
20. Let $A = \{1, 2, 5, 7\}$ and $B = \{1, 5, 9, 12\}$. Find each set.
- (a) $A \cap B$ (b) $A \cup B$

Solve each compound or absolute value equation or inequality.

21. $3x \geq 6$ and $x < 9$ 22. $-4x \leq -24$ or $4x < 12$
23. $|4x + 3| \leq 7$ 24. $|5 - 6x| > 12$
25. $|3x - 9| = 6$ 26. $|-3x + 4| - 4 < -1$
27. $|7 - x| \leq -1$ 28. $|3x - 2| + 1 = 8$ 29. $|3 - 5x| = |2x + 8|$
30. If $k < 0$, what is the solution set of each of the following?
- (a) $|8x - 5| < k$ (b) $|8x - 5| > k$ (c) $|8x - 5| = k$

STUDY SKILLS REMINDER

We can learn from the mistakes we make.

Review Study Skill 9, Analyzing Your Test Results.

Chapters R and 1

Cumulative Review Exercises

- Write $\frac{7}{20}$ as (a) a decimal and (b) a percent.
- Write 0.66 as (a) a percent and (b) a fraction in lowest terms.

Perform the indicated operations.

- $\frac{5}{6} + \frac{1}{4} - \frac{7}{15}$
- $0.17 + 3.965 - 12$
- $\frac{-4(9)(-2)}{-3^2}$
- $\sqrt{25} - 5(-1)^3$

Evaluate each exponential expression.

- $(-5)^3$
- $\left(\frac{3}{2}\right)^4$

Evaluate each expression for $x = 2$, $y = -3$, and $z = 4$.

- $-2y + 4(x - 3z)$
- $\frac{3x^2 - y^2}{4z}$

Simplify each expression.

- $-7r + 5 - 13r + 12$
- $-(3k + 8) - 2(4k - 7) + 3(8k + 12)$

Solve each equation.


- $\frac{2}{3}x + \frac{3}{4}x = -17$
- $\frac{2x + 3}{5} = \frac{x - 4}{2}$
- $|3m - 5| = |m + 2|$
- $A = P(1 + ni)$ for n

Solve each inequality. Graph the solution set, and write it using interval notation.

- $3 - 2(x + 7) \leq -x + 3$
- $-4 < 5 - 3x \leq 0$
- $2x + 1 > 5$ or $2 - x > 2$
- $|-7k + 3| \geq 4$

Solve each problem.

- Luke invested some money at 7% interest and the same amount at 10%. His total interest for the year was \$150 less than one-tenth of the total amount he invested. How much did he invest at each rate?
- A diet must include three foods, A, B, and C, with twice as many grams of food A as of food C, and 5 g of food B. The three foods must total at most 24 g. What is the largest amount of food C that can be used?
- Zach has scores of 88 and 78 on his first two tests. What score must he make on his third test to earn an average of 80 or greater?
- Telescope Peak, altitude 11,049 ft, is next to Death Valley, 282 ft below sea level. Find the difference between these altitudes. (Data from National Park Service.)
- For a woven hanging, Janette needs three pieces of yarn, which she will cut from a 40 cm piece. The longest piece is to be 3 times as long as the middle-sized piece, and the shortest piece is to be 5 cm shorter than the middle-sized piece. What lengths should she cut?





LINEAR EQUATIONS, GRAPHS, AND FUNCTIONS

The concept of steepness, or grade, is mathematically interpreted using *slope*, one of the topics of this chapter.

2.1 Linear Equations in Two Variables

2.2 The Slope of a Line

2.3 Writing Equations of Lines

SUMMARY EXERCISES Finding Slopes and Equations of Lines

2.4 Linear Inequalities in Two Variables

2.5 Introduction to Relations and Functions

2.6 Function Notation and Linear Functions


2.1 Linear Equations in Two Variables

OBJECTIVES

- 1 Interpret a line graph.
- 2 Plot ordered pairs.
- 3 Find ordered pairs that satisfy a given equation.
- 4 Graph lines.
- 5 Find x - and y -intercepts.
- 6 Graph equations of horizontal and vertical lines.
- 7 Find the midpoint of a line segment.

OBJECTIVE 1 Interpret a line graph.

The line graph in **FIGURE 1** shows personal spending (in billions of dollars) on medical care in the United States from 2010 through 2016. *About how much was spent on medical care in 2015?*




Data from Centers for Medicare & Medicaid Services.

FIGURE 1

VOCABULARY

- ordered pair
- origin
- x -axis
- y -axis
- rectangular (Cartesian) coordinate system
- components
- plot
- coordinate
- quadrant
- table of ordered pairs (table of values)
- graph of an equation
- first-degree equation
- linear equation in two variables
- x -intercept
- y -intercept

The line graph in **FIGURE 1** presents information based on a method for locating a point in a plane developed by René Descartes, a 17th-century French mathematician. According to legend, Descartes was lying in bed ill watching a fly crawl about on the ceiling near a corner of the room. It occurred to him that the location of the fly could be described by determining its distances from the two adjacent walls, as illustrated in the figure at the right.




Locating a fly on a ceiling

We use this insight to plot points and graph linear equations in two variables whose graphs are straight lines.

OBJECTIVE 2 Plot ordered pairs.

Each of the pairs of numbers $(3, 2)$, $(-5, 6)$, and $(4, -1)$ is an example of an **ordered pair**—that is, a pair of numbers written within parentheses. The *order* in which the numbers are written is important. We graph an ordered pair using two perpendicular number lines that intersect at their 0 points. See **FIGURE 2**. The common 0 point is the **origin**.



Rectangular coordinate system

FIGURE 2



René Descartes (1596–1650)

The position of any point in this coordinate plane is determined by referring to the horizontal number line, or **x -axis**, and the vertical number line, or **y -axis**. The x -axis and the y -axis make up a **rectangular** (or **Cartesian**, for Descartes) **coordinate system**.

The numbers in an ordered pair (x, y) , are its **components**. The first component indicates position relative to the x -axis, and the second component indicates position relative to the y -axis. For example, to locate, or **plot**, the point on the graph that corresponds to the ordered pair $(3, 2)$, we move three units from 0 to the right along the x -axis and then two units up parallel to the y -axis. See **FIGURE 3**. The numbers in an ordered pair are the **coordinates** of the corresponding point.




FIGURE 3

We can apply this method of locating ordered pairs to the line graph in **FIGURE 1**. We move along the horizontal axis to a year and then up parallel to the vertical axis to approximate spending for that year. Thus, we can write the ordered pair $(2015, 2716)$ to indicate that in 2015, personal spending on medical care was about \$2716 billion.

CAUTION The parentheses used to represent an ordered pair are also used to represent an open interval. The context of the discussion tells whether ordered pairs or open intervals are being represented.

The four regions of the graph shown in **FIGURE 3** are **quadrants I, II, III, and IV**, reading counterclockwise from the upper right quadrant. *The points on the x -axis and y -axis do not belong to any quadrant.*

OBJECTIVE 3 Find ordered pairs that satisfy a given equation.

Each solution of an equation with two variables, such as

$$2x + 3y = 6, \quad \text{Equation with two variables } x \text{ and } y$$

will include two numbers, one for each variable. To keep track of which number goes with which variable, we write the solutions as ordered pairs. *(If x and y are used as the variables, the x -value is given first.)*

We can show that $(6, -2)$ is a solution of $2x + 3y = 6$ by substitution.

$$2x + 3y = 6 \quad \text{Equation with two variables}$$

$$2(6) + 3(-2) \stackrel{?}{=} 6 \quad \text{Let } x = 6, y = -2.$$

$$12 - 6 \stackrel{?}{=} 6 \quad \text{Multiply.}$$

$$6 = 6 \quad \checkmark \quad \text{True}$$

Use parentheses to avoid errors.

Because the ordered pair $(6, -2)$ makes the equation true, it is a solution.

On the other hand, $(5, 1)$ is *not* a solution of $2x + 3y = 6$.

$$\begin{aligned} 2x + 3y &= 6 && \text{Equation with two variables} \\ 2(5) + 3(1) &\stackrel{?}{=} 6 && \text{Let } x = 5, y = 1. \\ 10 + 3 &\stackrel{?}{=} 6 && \text{Multiply.} \\ 13 &= 6 && \text{False} \end{aligned}$$

To find ordered pairs that satisfy an equation, we select any number for one of the variables, substitute it into the equation for that variable, and then solve for the other variable.

Because any real number could be selected for one variable and would lead to a real number for the other variable, an equation with two variables such as $2x + 3y = 6$ has an infinite number of solutions.

NOW TRY EXERCISE 1

Complete each ordered pair for $2x - y = 4$.

$$\begin{aligned} (0, \underline{\quad}), & \quad (\underline{\quad}, 0), \\ (4, \underline{\quad}), & \quad (\underline{\quad}, 2) \end{aligned}$$

Then write the results as a table of ordered pairs.

EXAMPLE 1 Completing Ordered Pairs and Making a Table

In parts (a)–(d), complete each ordered pair for $2x + 3y = 6$. Then, in part (e), write the results as a table of ordered pairs.

(a) $(0, \underline{\quad})$

$$\begin{aligned} 2x + 3y &= 6 \\ 2(0) + 3y &= 6 && \text{Let } x = 0. \\ 3y &= 6 && \text{Multiply. Add.} \\ y &= 2 && \text{Divide by 3.} \end{aligned}$$

The ordered pair is $(0, 2)$.

(c) $(-3, \underline{\quad})$

$$\begin{aligned} 2x + 3y &= 6 \\ 2(-3) + 3y &= 6 && \text{Let } x = -3. \\ -6 + 3y &= 6 && \text{Multiply.} \\ 3y &= 12 && \text{Add 6.} \\ y &= 4 && \text{Divide by 3.} \end{aligned}$$

The ordered pair is $(-3, 4)$.

(b) $(\underline{\quad}, 0)$

$$\begin{aligned} 2x + 3y &= 6 \\ 2x + 3(0) &= 6 && \text{Let } y = 0. \\ 2x &= 6 && \text{Multiply. Add.} \\ x &= 3 && \text{Divide by 2.} \end{aligned}$$

The ordered pair is $(3, 0)$.

(d) $(\underline{\quad}, -4)$

$$\begin{aligned} 2x + 3y &= 6 \\ 2x + 3(-4) &= 6 && \text{Let } y = -4. \\ 2x - 12 &= 6 && \text{Multiply.} \\ 2x &= 18 && \text{Add 12.} \\ x &= 9 && \text{Divide by 2.} \end{aligned}$$

The ordered pair is $(9, -4)$.

(e)

x	y	Ordered pairs
0	2	← Represents $(0, 2)$ from part (a)
3	0	← Represents $(3, 0)$ from part (b)
-3	4	← Represents $(-3, 4)$ from part (c)
9	-4	← Represents $(9, -4)$ from part (d)

Table of ordered pairs
(or table of values)

NOW TRY

NOW TRY ANSWER

1. $(0, -4), (2, 0), (4, 4), (3, 2)$

x	y
0	-4
2	0
4	4
3	2

OBJECTIVE 4 Graph lines.

The **graph of an equation** is the set of points corresponding to *all* ordered pairs that satisfy the equation. It gives a “picture” of the equation.

To graph $2x + 3y = 6$, we plot the ordered pairs found in **Objective 3** and **Example 1**. See **FIGURE 4(a)** on the next page. If *all* of the ordered pairs that satisfy the equation $2x + 3y = 6$ were graphed, they would form the straight line shown in **FIGURE 4(b)**.




FIGURE 4

The equation $2x + 3y = 6$ is a **first-degree equation** because it has no term with a variable to a power greater than 1.

The graph of any first-degree equation in two variables is a straight line.

Because first-degree equations with two variables have straight-line graphs, they are called **linear equations in two variables**.

Linear Equation in Two Variables

A **linear equation in two variables** (here x and y) is an equation that can be written in the form

$$Ax + By = C,$$

where A , B , and C are real numbers and A and B are not both 0. This form is called **standard form**.

Examples: $3x + 4y = 9$, $x - y = 0$, $x + 2y = -8$

OBJECTIVE 5 Find x - and y -intercepts.

A straight line is determined if any two different points on the line are known. Therefore, finding two different points is sufficient to graph the line.

Two useful points for graphing are the x - and y -intercepts. The **x -intercept** is the point (if any) where the line intersects the x -axis. The **y -intercept** is the point (if any) where the line intersects the y -axis.* See FIGURE 5.

- The y -value of the point where the line intersects the x -axis is always 0.
- The x -value of the point where the line intersects the y -axis is always 0.

This suggests a method for finding the x - and y -intercepts.




FIGURE 5

Finding Intercepts

When graphing the equation of a line, find the intercepts as follows.

Let $y = 0$ to find the x -intercept.

Let $x = 0$ to find the y -intercept.

*Some texts define an intercept as a number, not a point. For example, “ y -intercept $(0, 4)$ ” would be given as “ y -intercept 4.”

NOW TRY EXERCISE 2

Find the x - and y -intercepts, and graph the equation.

$$x - 2y = 4$$

EXAMPLE 2 Finding Intercepts

Find the x - and y -intercepts of $4x - y = -3$, and graph the equation.

To find the x -intercept, let $y = 0$.

$$\begin{aligned} 4x - y &= -3 \\ 4x - 0 &= -3 && \text{Let } y = 0. \\ 4x &= -3 && \text{Subtract.} \\ x &= -\frac{3}{4} && \text{Divide by 4.} \end{aligned}$$

The x -intercept is $(-\frac{3}{4}, 0)$.

To find the y -intercept, let $x = 0$.


$$\begin{aligned} 4x - y &= -3 \\ 4(0) - y &= -3 && \text{Let } x = 0. \\ -y &= -3 && \text{Multiply. Subtract.} \\ y &= 3 && \text{Multiply by } -1. \end{aligned}$$

The y -intercept is $(0, 3)$.

To guard against errors when graphing the equation, it is a good idea to find a third point. We arbitrarily choose $x = -2$, and substitute this value in the equation to find the corresponding value of y .

$$\begin{aligned} 4x - y &= -3 \\ 4(-2) - y &= -3 && \text{Let } x = -2. \\ -8 - y &= -3 && \text{Multiply.} \\ -y &= 5 && \text{Add 8.} \\ y &= -5 && \text{Multiply by } -1. \end{aligned}$$

The ordered pair $(-2, -5)$ lies on the graph. We plot the three ordered pairs and draw a line through them. See **FIGURE 6**.




A linear equation with both x and y variables will have both x - and y -intercepts. Its graph will be a “slanted” line.

FIGURE 6

NOW TRY

NOW TRY ANSWER

2. x -intercept: $(4, 0)$;
 y -intercept: $(0, -2)$



EXAMPLE 3 Graphing a Line That Passes through the Origin

Graph $x + 2y = 0$.

Find the x -intercept.

$$\begin{aligned} x + 2y &= 0 \\ x + 2(0) &= 0 && \text{Let } y = 0. \\ x + 0 &= 0 && \text{Multiply.} \\ x &= 0 && \text{x-intercept is } (0, 0). \end{aligned}$$

Find the y -intercept.

$$\begin{aligned} x + 2y &= 0 \\ 0 + 2y &= 0 && \text{Let } x = 0. \\ 2y &= 0 && \text{Add.} \\ y &= 0 && \text{y-intercept is } (0, 0). \end{aligned}$$

Both intercepts are the *same* point, $(0, 0)$, which means that the graph passes through the origin. To find a second point so we can graph the line, we choose any nonzero number for x or y and solve for the other variable. We arbitrarily choose $x = 4$.

NOW TRY EXERCISE 3
Graph $2x + 3y = 0$.

Choosing a multiple of 2 for x makes dividing by 2 in the last step easier.

$$x + 2y = 0$$

$$4 + 2y = 0 \quad \text{Let } x = 4.$$

$$2y = -4 \quad \text{Subtract 4.}$$

$$y = -2 \quad \text{Divide by 2.}$$

This gives the ordered pair $(4, -2)$. As a final check, substitute 1 for y in the equation and verify that $(-2, 1)$ also lies on the line. The graph is shown in FIGURE 7.




FIGURE 7

NOW TRY

OBJECTIVE 6 Graph equations of horizontal and vertical lines.

A line parallel to the x -axis will not have an x -intercept. Similarly, a line parallel to the y -axis will not have a y -intercept. This is why we included the phrase “if any” when we introduced intercepts.

NOW TRY EXERCISE 4
Graph each equation.

- (a) $y = -2$ (b) $x + 3 = 0$

EXAMPLE 4 Graphing Horizontal and Vertical Lines

Graph each equation.

- (a) $y = 2$ (This equation can be written as $0x + y = 2$.)

Because y *always* equals 2, there is no value of x corresponding to $y = 0$, and the graph has no x -intercept. One value where $y = 2$ is on the y -axis, so the y -intercept is $(0, 2)$. Plot any two other points with y -coordinate 2, such as $(-1, 2)$ and $(3, 2)$.

The graph is shown in FIGURE 8. It is a horizontal line.




FIGURE 8






FIGURE 9

NOW TRY ANSWERS



- (b) $x + 1 = 0$ (This equation can be written as $x = -1$ or $x + 0y = -1$.)

Because x *always* equals -1 , there is no value of y that makes $x = 0$, and the graph has no y -intercept. One value where $x = -1$ is on the x -axis, so the x -intercept is $(-1, 0)$. Plot any two other points with x -coordinate -1 , such as $(-1, -4)$ and $(-1, 5)$.

The graph is shown in FIGURE 9. It is a vertical line.

NOW TRY

CAUTION A linear equation that has only *one* variable x or y will have a vertical or horizontal line as its graph.

1. An equation with only the variable x will always intersect the x -axis and thus will be **vertical**. It has the form $x = a$. **The vertical line with equation $x = 0$ is the y -axis.**
2. An equation with only the variable y will always intersect the y -axis and thus will be **horizontal**. It has the form $y = b$. **The horizontal line with equation $y = 0$ is the x -axis.**

OBJECTIVE 7 Find the midpoint of a line segment.

If the coordinates of the endpoints of a line segment are known, then the coordinates of the *midpoint* of the segment can be found.

FIGURE 10 shows a line segment PQ with endpoints $P(-8, 4)$ and $Q(3, -2)$. R is the point with the same x -coordinate as P and the same y -coordinate as Q . So the coordinates of R are $(-8, -2)$.




FIGURE 10

The x -coordinate of the midpoint M of PQ is the same as the x -coordinate of the midpoint of RQ . Because RQ is horizontal, the x -coordinate of its midpoint is the *average* (or *mean*) of the x -coordinates of its endpoints.

$$\frac{1}{2}(-8 + 3) = -\frac{5}{2}$$

The y -coordinate of M is the average (or *mean*) of the y -coordinates of the midpoint of PR .

$$\frac{1}{2}(4 + (-2)) = 1$$

The midpoint of PQ is $M(-\frac{5}{2}, 1)$.

This discussion leads to the *midpoint formula*.

Midpoint Formula

The midpoint M of a line segment PQ with endpoints $P(x_1, y_1)$ and $Q(x_2, y_2)$ is found as follows.

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

The two nonspecific points (x_1, y_1) and (x_2, y_2) use **subscript notation**. Read (x_1, y_1) as “*x*-sub-one, *y*-sub-one.”

NOW TRY EXERCISE 5

Find the midpoint of line segment PQ with endpoints $P(2, -5)$ and $Q(-4, 7)$.

EXAMPLE 5 Finding the Coordinates of a Midpoint

Find the midpoint of line segment PQ with endpoints $P(4, -3)$ and $Q(6, -1)$.

$$\begin{array}{ccc} (x_1, y_1) & & (x_2, y_2) \\ \downarrow & & \downarrow \\ P(4, -3) & \text{and} & Q(6, -1) \end{array} \quad \text{Label the points.}$$

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) \quad \text{Midpoint formula}$$

We are finding the average of the x -coordinates and the average of the y -coordinates.


$$= \left(\frac{4 + 6}{2}, \frac{-3 + (-1)}{2} \right) \quad \text{Substitute.}$$

$$= \left(\frac{10}{2}, \frac{-4}{2} \right) \quad \text{Add in the numerators.}$$

$$= (5, -2) \leftarrow \text{Midpoint of segment } PQ$$

NOW TRY 

NOTE When graphing with a graphing calculator, we “set up” a rectangular coordinate system. The screen in **FIGURE 11** shows the **standard viewing window**. Minimum x - and y -values are -10 , and maximum x - and y -values are 10 . The scale on each axis, here 1 , determines the distance between the tick marks.



Standard viewing window

FIGURE 11




FIGURE 12

To graph an equation such as $4x - y = -3$, we use the intercepts $(-0.75, 0)$ and $(0, 3)$ to determine an appropriate window. Here, we choose the standard viewing window. We solve the equation for y to obtain $y = 4x + 3$ and enter it into the calculator. The graph in **FIGURE 12** also gives the coordinates of the x -intercept at the bottom of the screen.

NOW TRY ANSWER


5. $(-1, 1)$

2.1 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Are you fully utilizing the features of your text? **Review Study Skill 1, Using Your Math Text.**


Concept Check Fill in each blank with the correct response.

- The point with coordinates $(0, 0)$ is the _____ of a rectangular coordinate system.
- For any value of x , the point $(x, 0)$ lies on the _____-axis. For any value of y , the point $(0, y)$ lies on the _____-axis.
- To find the x -intercept of a line, we let _____ equal 0 and solve for _____. To find the y -intercept, we let _____ equal 0 and solve for _____.
- The equation $y = 4$ has a _____ line as its graph, and $x = 4$ has a _____ line as its graph.

5. To graph a straight line, we must find a minimum of _____ points. The points $(3, \text{_____})$ and $(\text{_____}, 4)$ lie on the graph of $2x - 3y = 0$.
6. The equation of the x -axis is _____. The equation of the y -axis is _____.


Concept Check Solve each problem by locating ordered pairs on the graphs.

7. The graph indicates personal spending in billions of dollars on medical care in the United States.



Data from Centers for Medicare & Medicaid Services.

- (a) If (x, y) represents a point on the graph, what does x represent? What does y represent?
 - (b) Estimate spending in 2016.
 - (c) Write an ordered pair (x, y) that represents approximate spending in 2016.
 - (d) In what year was spending about \$2200 billion?
8. The graph shows the percentage of Americans who moved in selected years.



Data from U.S. Census Bureau.
* Data not collected for 1980.

- (a) If the ordered pair (x, y) represents a point on the graph, what does x represent? What does y represent?
- (b) Estimate the percentage of Americans who moved in 2017.
- (c) Write an ordered pair (x, y) that gives the approximate percentage of Americans who moved in 2017.
- (d) What does the ordered pair $(1960, 20)$ mean in the context of this graph?

Name the quadrant, if any, in which each point is located. See Objective 2.

- | | | | |
|-----------------|-----------------|---------------------|----------------|
| 9. (a) $(1, 6)$ | (b) $(-4, -2)$ | 10. (a) $(-2, -10)$ | (b) $(4, 8)$ |
| (c) $(-3, 6)$ | (d) $(7, -5)$ | (c) $(-9, 12)$ | (d) $(3, -9)$ |
| (e) $(-3, 0)$ | (f) $(0, -0.5)$ | (e) $(0, -8)$ | (f) $(2.5, 0)$ |

11. **Concept Check** Use the given information to determine the quadrants in which the point (x, y) must lie. (*Hint:* Consider the signs of the coordinates in each quadrant, and the signs of their product and quotient.)

- | | | | |
|--------------|--------------|-----------------------|-----------------------|
| (a) $xy > 0$ | (b) $xy < 0$ | (c) $\frac{x}{y} < 0$ | (d) $\frac{x}{y} > 0$ |
|--------------|--------------|-----------------------|-----------------------|

12. **Concept Check** What must be true about the value of at least one of the coordinates of any point that lies along an axis?

- 13. Concept Check** A student plotted the point with coordinates $(-4, 2)$ incorrectly by moving 2 units from 0 to the right along the x -axis and then 4 units down parallel to the y -axis.

WHAT WENT WRONG?

- 14. Concept Check** A student incorrectly claimed that the point $(0, -4)$ lies on the x -axis because the x -coordinate is 0. **WHAT WENT WRONG?**

Plot each point in a rectangular coordinate system. See **Objective 2**.

15. $(2, 3)$ 16. $(-1, 2)$ 17. $(-3, -2)$ 18. $(1, -4)$ 19. $(0, 5)$
 20. $(-2, -4)$ 21. $(-2, 4)$ 22. $(3, 0)$ 23. $(-2, 0)$ 24. $(3, -3)$

Complete the given table for each equation and then graph the equation. See **Example 1** and **FIGURE 4**.

25. $y = x - 4$

x	y
0	
1	
2	
3	
4	

26. $y = x + 3$

x	y
0	
1	
2	
3	
4	

27. $x - y = 3$

x	y
0	
	0
5	
2	

28. $x - y = 5$

x	y
0	
	0
1	
3	

29. $x + 2y = 5$

x	y
0	
	0
2	
	2

30. $x + 3y = -5$

x	y
0	
	0
1	
	-1

31. $4x - 5y = 20$

x	y
0	
	0
2	
	-3

32. $6x - 5y = 30$

x	y
0	
	0
3	
	-2





- 33. Concept Check** Match each equation in parts (a)–(d) with its graph in choices A–D. (Coordinates of the points shown are integers.)

(a) $x + 3y = 3$

(b) $x - 3y = -3$

(c) $x - 3y = 3$

(d) $x + 3y = -3$



- 34. Concept Check** Which of the following equations have a graph that is a horizontal line? A vertical line?

A. $x - 6 = 0$

B. $x + y = 0$

C. $y + 3 = 0$

D. $y = -10$

E. $x + 1 = 5$

Find the x - and y -intercepts. Then graph each equation. See **Examples 2–4**.

35. $2x + 3y = 12$

36. $5x + 2y = 10$

37. $x - 3y = 6$

38. $x - 2y = -4$

39. $5x + 6y = -10$

40. $3x - 7y = 9$

41. $\frac{2}{3}x - 3y = 7$

42. $\frac{5}{7}x + \frac{6}{7}y = -2$

43. $x + 5y = 0$

44. $x - 3y = 0$

45. $2x + y = 0$

46. $4x - y = 0$

47. $2x = 3y$

48. $4y = 3x$

49. $-\frac{2}{3}y = x$

50. $-\frac{3}{4}y = x$

51. $y = 5$

52. $y = -3$

53. $x = 2$

54. $x = -3$

55. $x + 4 = 0$

56. $x - 4 = 0$

57. $y + 2 = 0$

58. $y - 5 = 0$

Each table of values gives several points that lie on a line.

(a) What is the x -intercept of the line? The y -intercept?

(b) Which equation in choices A–D corresponds to the given table of values?

(c) Graph the equation.

x	y
-4	-3
-2	0
0	3
2	6

A. $3x + 2y = 6$

B. $3x - 2y = -6$

C. $3x + 2y = -6$

D. $3x - 2y = 6$

x	y
-1	6
0	4
1	2
2	0

A. $2x - y = 4$

B. $2x + y = -4$

C. $2x + y = 4$

D. $2x - y = -4$

x	y
-2	-1
0	-1
2	-1
4	-1

A. $y = -1$

B. $y = 1$

C. $x = 1$

D. $x = -1$

x	y
6	-1
6	0
6	1
6	2

A. $x = -6$

B. $y = 0$

C. $y = 6$

D. $x = 6$

Find the midpoint of each segment with the given endpoints. See Example 5.

63. $(-8, 4)$ and $(-2, -6)$

64. $(5, 2)$ and $(-1, 8)$

65. $(3, -6)$ and $(6, 3)$

66. $(-10, 4)$ and $(7, 1)$

67. $(-9, 3)$ and $(9, 8)$

68. $(4, -3)$ and $(-1, 3)$

69. $(2.5, 3.1)$ and $(1.7, -1.3)$

70. $(6.2, 5.8)$ and $(1.4, -0.6)$

Extending Skills Find the midpoint of each segment with the given endpoints.

71. $\left(\frac{1}{2}, \frac{1}{3}\right)$ and $\left(\frac{3}{2}, \frac{5}{3}\right)$

72. $\left(\frac{21}{4}, \frac{2}{5}\right)$ and $\left(\frac{7}{4}, \frac{3}{5}\right)$

73. $\left(-\frac{1}{3}, \frac{2}{7}\right)$ and $\left(-\frac{1}{2}, \frac{1}{14}\right)$

74. $\left(\frac{3}{5}, -\frac{1}{3}\right)$ and $\left(\frac{1}{2}, -\frac{7}{2}\right)$

Extending Skills Segment PQ has the given coordinates for one endpoint P and for its midpoint M . Find the coordinates of the other endpoint Q . (Hint: Represent Q by (x, y) and write two equations using the midpoint formula, one involving x and the other involving y . Then solve for x and y .)

75. $P(5, 8), M(8, 2)$

76. $P(7, 10), M(5, 3)$

77. $P(1.5, 1.25), M(3, 1)$

78. $P(2.5, 1.75), M(3, 2)$

2.2 The Slope of a Line

OBJECTIVES

- 1 Find the slope of a line given two points on the line.
- 2 Find the slope of a line given an equation of the line.
- 3 Graph a line given its slope and a point on the line.
- 4 Determine whether two lines are parallel, perpendicular, or neither using slope.
- 5 Solve problems involving average rate of change.

Slope (steepness) is used in many practical ways. The slope (or *grade*) of a highway is often given as a percent. For example, a 10% (or $\frac{10}{100} = \frac{1}{10}$) slope means that the highway rises 1 unit for every 10 horizontal units. Stairs and roofs have slopes too, as shown in **FIGURE 13**.




FIGURE 13

Slope is the ratio of vertical change, or **rise**, to horizontal change, or **run**. A simple way to remember this is to think, “*Slope is rise over run.*”

OBJECTIVE 1 Find the slope of a line given two points on the line.

To obtain a formal definition of the slope of a line, we designate two different points on the line as (x_1, y_1) and (x_2, y_2) . See **FIGURE 14**.




FIGURE 14

As we move along the line in **FIGURE 14** from (x_1, y_1) to (x_2, y_2) , the y -value changes (vertically) from y_1 to y_2 , an amount equal to $y_2 - y_1$. As y changes from y_1 to y_2 , the value of x changes (horizontally) from x_1 to x_2 by the amount $x_2 - x_1$.

NOTE The Greek letter **delta**, Δ , is used in mathematics to denote “change in,” so Δy and Δx represent the change in y and the change in x , respectively.

The ratio of the change in y to the change in x (“rise over run,” or $\frac{\text{rise}}{\text{run}}$) is the *slope* of the line, with the letter m traditionally used for slope.

Slope Formula

The **slope** m of the line passing through the distinct points (x_1, y_1) and (x_2, y_2) is defined as follows.

$$m = \frac{\text{rise}}{\text{run}} = \frac{\text{change in } y}{\text{change in } x} = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1} \quad (\text{where } x_1 \neq x_2)$$

VOCABULARY

- rise
- run
- slope

STUDY SKILLS REMINDER

Are you getting the most out of your class time? **Review Study Skill 3, Taking Lecture Notes.**

NOW TRY
EXERCISE 1

Find the slope of the line passing through the points $(2, -6)$ and $(-3, 5)$.

EXAMPLE 1 Finding the Slope of a Line

Find the slope of the line passing through the points $(2, -1)$ and $(-5, 3)$.
Label the points, and then apply the slope formula.

$$\begin{array}{ccc} (x_1, y_1) & & (x_2, y_2) \\ \downarrow \downarrow & & \downarrow \downarrow \\ (2, -1) & \text{and} & (-5, 3) \end{array}$$

$$\text{slope } m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - (-1)}{-5 - 2} \quad \text{Substitute.}$$

$$= \frac{4}{-7}, \quad \text{or} \quad -\frac{4}{7} \quad \text{Subtract; } \frac{a}{-b} = -\frac{a}{b}$$

The slope is $-\frac{4}{7}$.

The same slope is obtained if we label the points in reverse order. *It makes no difference which point is identified as (x_1, y_1) and which as (x_2, y_2) .*

$$\begin{array}{ccc} (x_2, y_2) & & (x_1, y_1) \\ \downarrow \downarrow & & \downarrow \downarrow \\ (2, -1) & \text{and} & (-5, 3) \end{array}$$


$$\text{slope } m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-1 - 3}{2 - (-5)} \quad \text{Substitute.}$$

$$= \frac{-4}{7}, \quad \text{or} \quad -\frac{4}{7} \quad \text{Subtract; } \frac{-a}{b} = -\frac{a}{b}$$

y-values are in the numerator, x-values in the denominator.

To confirm this slope, see **FIGURE 15**. Using the geometric interpretation of slope (as $\frac{\text{rise}}{\text{run}}$) and beginning at the point $(2, -1)$, we move **up 4** units and to the **left 7** units to reach the point $(-5, 3)$.

$$\text{slope } m = \frac{\text{change in } y \text{ (rise)}}{\text{change in } x \text{ (run)}} = \frac{4}{-7}, \quad \text{or} \quad -\frac{4}{7}$$



Moves *down* or to the *left* are denoted as negative numbers.

FIGURE 15

The same slope results if we begin at the point $(-5, 3)$ and move **down 4** units and to the **right 7** units to reach the point $(2, -1)$. See **FIGURE 15**.

NOW TRY ANSWER

1. $-\frac{11}{5}$

$$\text{slope } m = \frac{\text{change in } y \text{ (rise)}}{\text{change in } x \text{ (run)}} = \frac{-4}{7}, \quad \text{or} \quad -\frac{4}{7}$$

Example 1 suggests the following important ideas regarding slope.

1. The slope is the same no matter which point we consider first.
2. Using similar triangles from geometry, we can show that the slope is the same no matter which two different points on the line we choose.

! CAUTION When calculating slope, remember that the change in y (rise) is the numerator and the change in x (run) is the denominator.

Correct

$$\frac{y_2 - y_1}{x_2 - x_1}$$

Incorrect

$$\frac{x_2 - x_1}{y_2 - y_1} \text{ or } \frac{y_2 - y_1}{x_1 - x_2} \text{ or } \frac{y_1 - y_2}{x_2 - x_1}$$

Be careful to subtract the y -values and the x -values in the same order.

OBJECTIVE 2 Find the slope of a line given an equation of the line.

When an equation of a line is given, one way to find its slope is to use the definition of slope with two different points on the line.

NOW TRY EXERCISE 2

Find the slope of the line $3x - 7y = 21$.




FIGURE 16

EXAMPLE 2 Finding the Slope of a Line

Find the slope of the line $3x - y = -6$.

The intercepts can be used as the two points needed to find the slope.

Let $y = 0$ to find that the x -intercept is $(-2, 0)$.

Let $x = 0$ to find that the y -intercept is $(0, 6)$.

Use these two points in the slope formula.

$$\begin{aligned} \text{slope } m &= \frac{y_2 - y_1}{x_2 - x_1} = \frac{6 - 0}{0 - (-2)} && (x_1, y_1) = (-2, 0) \\ & && (x_2, y_2) = (0, 6) \\ &= \frac{6}{2} && \text{Subtract.} \\ &= 3 && \text{Divide.} \end{aligned}$$

The graph of $3x - y = -6$ in **FIGURE 16** confirms that the slope of the line is 3.

NOW TRY

EXAMPLE 3 Applying the Slope Concept to Horizontal and Vertical Lines

Find the slope of each line.

(a) $y = 2$

The graph of $y = 2$ is a horizontal line. See **FIGURE 17**. To find the slope, select two points on the line, such as $(3, 2)$ and $(-1, 2)$, and apply the slope formula.

$$m = \frac{2 - 2}{-1 - 3} = \frac{0}{-4} = 0 \quad \begin{aligned} (x_1, y_1) &= (3, 2) \\ (x_2, y_2) &= (-1, 2) \end{aligned}$$

In this case, the *rise* is 0, so the slope is 0.




FIGURE 17

NOW TRY ANSWER

2. $\frac{3}{7}$

NOW TRY EXERCISE 3

Find the slope of each line.

- (a) $x = 4$ (b) $y - 6 = 0$

(b) $x + 1 = 0$

The graph of $x + 1 = 0$, or $x = -1$, is a vertical line. See **FIGURE 18**. Two points that satisfy the equation $x = -1$ are $(-1, 5)$ and $(-1, -4)$. Use these two points and the slope formula.

$$m = \frac{-4 - 5}{-1 - (-1)} = \frac{-9}{0} \quad \begin{array}{l} (x_1, y_1) = (-1, 5) \\ (x_2, y_2) = (-1, -4) \end{array}$$

Here the *run* is 0. Because division by 0 is undefined, the slope is undefined. This is why the definition of slope includes the restriction that $x_1 \neq x_2$.




FIGURE 18

NOW TRY

Horizontal and Vertical Lines

- An equation of the form $y = b$ always intersects the y -axis at the point $(0, b)$. A line with this equation is **horizontal** and has **slope 0**.
- An equation of the form $x = a$ always intersects the x -axis at the point $(a, 0)$. A line with this equation is **vertical** and has **undefined slope**.

The slope of a line can also be found directly from its equation. Consider the equation $3x - y = -6$ from **Example 2**. Solve this equation for y .

$$\begin{aligned} 3x - y &= -6 && \text{Equation from Example 2} \\ -y &= -6 - 3x && \text{Subtract } 3x. \\ y &= 3x + 6 && \text{Multiply by } -1; \text{Commutative property} \end{aligned}$$

The slope, 3, found using the slope formula in **Example 2**, is the same number as the coefficient of x in the equation $y = 3x + 6$. We will see in the next section that this is true in general, as long as the equation is solved for y .

EXAMPLE 4 Finding the Slope from an Equation

Find the slope of the graph of $3x - 5y = 8$.

Many of the ordered-pair solutions of this equation have coordinates that are fractions, including the intercepts $(\frac{8}{3}, 0)$ and $(0, -\frac{8}{5})$. See **FIGURE 19**. This makes calculations using the slope formula more difficult.

Instead, we solve the equation for y as discussed above.

$$\begin{aligned} 3x - 5y &= 8 && \text{Solve for } y. \\ -5y &= 8 - 3x && \text{Subtract } 3x. \\ \frac{-5y}{-5} &= \frac{-3x + 8}{-5} && \text{Commutative property;} \\ &&& \text{Divide each side by } -5. \\ \frac{-3x}{-5} = \frac{-3}{-5} \cdot \frac{x}{1} = \frac{3}{5}x &&& \frac{a+b}{c} = \frac{a}{c} + \frac{b}{c} \\ y &= \frac{3}{5}x - \frac{8}{5} \end{aligned}$$

The slope, given by the coefficient of x , is $\frac{3}{5}$.

We can confirm this slope using the intercepts and the slope formula, which involves simplifying a *complex fraction*.

$$m = \frac{0 - (-\frac{8}{5})}{\frac{8}{3} - 0} = \frac{\frac{8}{5}}{\frac{8}{3}} = \frac{8}{5} \div \frac{8}{3} = \frac{8}{5} \cdot \frac{3}{8} = \frac{3}{5} \quad \text{The same slope results.}$$

NOW TRY

NOW TRY EXERCISE 4

Find the slope of the graph of $5x - 4y = 7$.




FIGURE 19

NOW TRY ANSWERS

3. (a) undefined (b) 0
4. $\frac{5}{4}$

NOTE We can solve the standard form of a linear equation $Ax + By = C$ (where $B \neq 0$) for y to show that, in general, **the slope of a line in this form is $-\frac{A}{B}$.**

$$\begin{array}{ll} Ax + By = C & \text{Standard form} \\ By = C - Ax & \text{Subtract } Ax. \\ By = -Ax + C & \text{Commutative property} \\ y = -\frac{A}{B}x + \frac{C}{B} & \text{Divide each term by } B. \end{array}$$

The slope is given by the coefficient of x , $-\frac{A}{B}$. In the equation $3x - 5y = 8$ from **Example 4**, $A = 3$ and $B = -5$, so the slope is

$$-\frac{A}{B} = -\frac{3}{-5} = \frac{3}{5}. \quad \text{The same slope results.}$$

OBJECTIVE 3 Graph a line given its slope and a point on the line.

NOW TRY EXERCISE 5

Graph the line passing through $(-4, 1)$ that has slope $-\frac{2}{3}$.

EXAMPLE 5 Using the Slope and a Point to Graph Lines

Graph each line passing through the given point having the given slope.

(a) $(0, -4)$; slope $\frac{2}{3}$

Begin by plotting the point $P(0, -4)$, as shown in **FIGURE 20**. Then use the geometric interpretation of slope to find a second point.

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{2 \leftarrow \text{rise}}{3 \leftarrow \text{run}}$$

We move *up 2* units from $(0, -4)$ and to the *right 3* units to locate a second point on the graph, $R(3, -2)$. The line through $P(0, -4)$ and R is the required graph.




FIGURE 20

(b) $(3, 1)$; slope -4

Start by plotting the point $P(3, 1)$. See **FIGURE 21**. Find a second point R on the line by writing the slope -4 as $\frac{-4}{1}$ and using the geometric interpretation of slope.

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{-4 \leftarrow \text{rise}}{1 \leftarrow \text{run}}$$

We move *down 4* units from $(3, 1)$ and to the *right 1* unit to locate a second point, $R(4, -3)$. The line through $P(3, 1)$ and R is the required graph.




FIGURE 21


The slope -4 also could be written as

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{4}{-1}$$

In this case, the second point R is located *up 4* units and to the *left 1* unit. Verify that this approach also produces the line in **FIGURE 21**.

NOW TRY

NOW TRY ANSWER



The line in **Example 5(a)** has *positive* slope $\frac{2}{3}$. The graph of the line in **FIGURE 20** *slants up (rises)* from left to right. The line in **Example 5(b)** has *negative* slope -4 . As **FIGURE 21** shows, its graph *slants down (falls)* from left to right. These facts illustrate the following generalization.

Orientation of a Line in the Plane

A *positive slope* indicates that the line *slants up (rises)* from left to right.

A *negative slope* indicates that the line *slants down (falls)* from left to right.

FIGURE 22 summarizes the four cases for slopes of lines.




FIGURE 22

OBJECTIVE 4 Determine whether two lines are parallel, perpendicular, or neither using slope.

The slope of a line measures its steepness, and parallel lines have equal steepness.

Slopes of Parallel Lines

Two nonvertical lines with the same slope are parallel.

Two nonvertical parallel lines have the same slope.

NOW TRY EXERCISE 6


Determine whether the line passing through (2, 5) and (4, 8) and the line passing through (2, 0) and (-1, -2) are parallel.

EXAMPLE 6 Determining Whether Two Lines Are Parallel

Determine whether the lines L_1 , passing through (-2, 1) and (4, 5), and L_2 , passing through (3, 0) and (0, -2), are parallel.

Slope of L_1 :

$$\begin{aligned} m_1 &= \frac{5 - 1}{4 - (-2)} \\ &= \frac{4}{6} \\ &= \frac{2}{3} \end{aligned}$$



Slope of L_2 :

$$\begin{aligned} m_2 &= \frac{0 - (-2)}{3 - 0} \\ &= \frac{2}{3} \end{aligned}$$

FIGURE 23

The slopes are the *same*, both $\frac{2}{3}$, so the two lines are *parallel*. See **FIGURE 23**.

NOW TRY ANSWER

6. not parallel

To see how the slopes of perpendicular lines are related, consider a nonvertical line with slope $\frac{a}{b}$. If this line is rotated 90° , the vertical change and the horizontal change are interchanged, and the slope is $-\frac{b}{a}$ because the horizontal change is now negative. See FIGURE 24. Thus, the slopes of perpendicular lines have product -1 and are negative reciprocals of each other.




FIGURE 24

For example, if the slopes of two lines are $\frac{3}{4}$ and $-\frac{4}{3}$, then the lines are perpendicular because

$$\frac{3}{4} \left(-\frac{4}{3} \right) = -1.$$

Slopes of Perpendicular Lines

If neither is vertical, perpendicular lines have slopes that are negative reciprocals—that is, their product is -1 . Also, lines with slopes that are negative reciprocals are perpendicular.

A line with slope 0 is perpendicular to a line with undefined slope.

EXAMPLE 7 Determining Whether Two Lines Are Parallel, Perpendicular, or Neither

Determine whether each pair of lines is *parallel*, *perpendicular*, or *neither*.

(a) $2y = 3x - 6$ and $2x + 3y = -6$

Find the slope of each line by solving each equation for y .

$$2y = 3x - 6$$

$$y = \frac{3}{2}x - 3 \quad \text{Divide by 2.}$$

↑
Slope

$$2x + 3y = -6$$

$$3y = -2x - 6 \quad \text{Subtract } 2x.$$

$$y = -\frac{2}{3}x - 2 \quad \text{Divide by 3.}$$

↑
Slope

The slopes are *negative reciprocals* because their product is $\frac{3}{2} \left(-\frac{2}{3} \right) = -1$. The two lines are *perpendicular*. See FIGURE 25.




FIGURE 25




FIGURE 26

NOW TRY EXERCISE 7

Determine whether each pair of lines is *parallel*, *perpendicular*, or *neither*.

- (a) $x + 2y = 7$ and $2x = y - 4$
- (b) $2x - y = 4$ and $2x + y = 6$

(b) $2x - 5y = 8$ and $2x + 5y = 8$

$$2x - 5y = 8$$

$$-5y = -2x + 8 \quad \text{Subtract } 2x.$$

$$y = \frac{2}{5}x - \frac{8}{5} \quad \text{Divide by } -5.$$

↑
Slope

$$2x + 5y = 8$$

$$5y = -2x + 8 \quad \text{Subtract } 2x.$$

$$y = -\frac{2}{5}x + \frac{8}{5} \quad \text{Divide by } 5.$$

↑
Slope

The slopes, $\frac{2}{5}$ and $-\frac{2}{5}$, are not the same. They are not negative reciprocals because their product is $-\frac{4}{25}$, *not* -1 . The two lines are *neither* parallel nor perpendicular. See FIGURE 26. **NOW TRY** ↻

NOTE In Example 7(a), we could have found the slope of each line using intercepts and the slope formula. The graph of $2y = 3x - 6$ has intercepts $(2, 0)$ and $(0, -3)$.

$$m = \frac{0 - (-3)}{2 - 0} = \frac{3}{2} \quad \text{The same slope found in Example 7(a) results.}$$

Find the intercepts of the graph of $2x + 3y = -6$ and use them to confirm the slope $-\frac{2}{3}$. Because the slopes are negative reciprocals, the lines are perpendicular.

To avoid fractions when using the slope formula in Example 7(b), we suggest identifying points with integer coordinates, such as $(4, 0)$ and $(-1, -2)$ for the equation $2x - 5y = 8$.

OBJECTIVE 5 Solve problems involving average rate of change.

The slope formula applied to any two points on a line gives the **average rate of change** in y per unit change in x .

For example, suppose the height of a boy increased from 60 to 68 in. between the ages of 12 and 16, as shown in FIGURE 27.




FIGURE 27

$$\begin{aligned} \text{Change in height } y &\longrightarrow \frac{68 - 60}{16 - 12} = \frac{8}{4} = 2 \text{ in. per yr} \\ \text{Change in age } x &\longrightarrow \end{aligned}$$

The boy may actually have grown more (or less) than 2 in. during some years. If we plotted ordered pairs (age, height) for those years and drew a line connecting any two of the points, the average rate of change would likely be slightly different. However, using the data for ages 12 and 16, the boy's *average* change in height was 2 in. per year over these years.

NOW TRY ANSWERS


- 7. (a) perpendicular
- (b) neither

NOW TRY EXERCISE 8

There were approximately 84 million high-speed Internet subscribers in 2013. Using this number for 2013 and the number for 2017 from the graph in **FIGURE 28**, find the average rate of change per year from 2013 to 2017. How does it compare with the average rate of change found in **Example 8**?

EXAMPLE 8 Interpreting Slope as Average Rate of Change

The graph in **FIGURE 28** shows the number of high-speed Internet subscribers in the United States from 2012 to 2017. Find the average rate of change in number of subscribers per year.



Data from Leichtman Research Group.

FIGURE 28

To find the average rate of change, we need two pairs of data. From the graph, we have the ordered pairs (2012, 81) and (2017, 95). We use the slope formula.

$$\text{average rate of change} = \frac{95 - 81}{2017 - 2012} = \frac{14}{5} = 2.8$$

A positive slope indicates an increase.

This means that the number of high-speed Internet subscribers *increased* by an average of 2.8 million subscribers per year from 2012 to 2017.

NOW TRY

NOW TRY EXERCISE 9

In 2011, global sales of digital cameras totaled 33 billion euros. In 2016, sales totaled 17 billion euros. Find the average rate of change in sales of digital cameras per year. (Data from Photoindustrie-Verband.)

EXAMPLE 9 Interpreting Slope as Average Rate of Change

In 2011, there were 105 million households with cable TV in the United States. There were 96 million such households in 2017. Find the average rate of change in the number of households per year. (Data from Nielsen Media Research.)

We must write two ordered pairs of the form (year, number of households) to use to find the average rate of change. From the problem, we have (2011, 105) and (2017, 96). We substitute these values in the slope formula.

$$\text{average rate of change} = \frac{96 - 105}{2017 - 2011} = \frac{-9}{6} = -1.5$$

A negative slope indicates a decrease.

The graph in **FIGURE 29** confirms that the line through the ordered pairs falls from left to right and therefore has negative slope. Thus, the number of households with cable TV *decreased* by an average of 1.5 million households per year from 2011 to 2017.

The negative sign in -1.5 denotes the *decrease*. (We say “The number of customers decreased by 1.5 million per year.” It is *incorrect* to say “The number of customers decreased by -1.5 million per year.”)




FIGURE 29

NOW TRY ANSWERS


8. 2.75 million subscribers per year; It is less than the average rate of change per year from 2012 to 2017.
9. -3.2 billion euros per year

NOW TRY

2.2 Exercises

FOR
EXTRA
HELP

 MyLab Math

 Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER


Be sure to read and work through the section material before working the exercises.

Review Study Skill 2, Reading Your Math Text.

Concept Check Answer each question.


1. A hill rises 30 ft for every horizontal 100 ft. Which of the following express its slope (or grade)? (There are several correct choices.)

- A. 0.3 B. $\frac{3}{10}$ C. $3\frac{1}{3}$
 D. $\frac{30}{100}$ E. $\frac{10}{3}$ F. 30%





2. If a walkway rises 2 ft for every 24 ft on the horizontal, which of the following express its slope (or grade)? (There are several correct choices.)

- A. 12% B. $\frac{2}{24}$ C. $\frac{1}{12}$
 D. 12 E. 8. $\bar{3}$ % F. $\frac{24}{2}$







3. A ladder leaning against a wall has slope 3. How many feet in the horizontal direction correspond to a rise of 15 ft?
4. A hill has slope 0.05. How many feet in the vertical direction correspond to a run of 50 ft?




5. **Concept Check** Match each situation in parts (a)–(d) with the most appropriate graph in choices A–D.

- (a) Sales rose sharply during the first quarter, leveled off during the second quarter, and then rose slowly for the rest of the year.
- (b) Sales fell sharply during the first quarter and then rose slowly during the second and third quarters before leveling off for the rest of the year.
- (c) Sales rose sharply during the first quarter and then fell to the original level during the second quarter before rising steadily for the rest of the year.
- (d) Sales fell during the first two quarters of the year, leveled off during the third quarter, and rose during the fourth quarter.




6. **Concept Check** Using the given axes, draw a graph that illustrates the following description.

Profits for a business were \$10 million in 2013. They rose sharply from 2013 through 2015, remained constant from 2015 through 2017, and then fell slowly from 2017 through 2018.



Concept Check Determine the slope of each line segment in the given figure.


7. AB 8. BC 9. CD
 10. DE 11. EF 12. FG



13. If A and F were joined by a line segment in the figure, what would be its slope?
 14. If B and D were joined by a line segment in the figure, what would be its slope?

15. **Concept Check** Refer to the figure shown here and determine which line satisfies the given description.

- (a) The line has positive slope.
 (b) The line has negative slope.
 (c) The line has slope 0.
 (d) The line has undefined slope.



16. **Concept Check** Which forms of the slope formula are correct? Explain.

A. $m = \frac{y_1 - y_2}{x_2 - x_1}$ B. $m = \frac{y_1 - y_2}{x_1 - x_2}$ C. $m = \frac{x_2 - x_1}{y_2 - y_1}$ D. $m = \frac{y_2 - y_1}{x_2 - x_1}$

17. **Concept Check** A student found the slope of the line through the points $(-4, 5)$ and $(2, 7)$ as follows.

$$m = \frac{7 - 5}{-4 - 2} = \frac{2}{-6} = -\frac{1}{3}$$

This is incorrect. **WHAT WENT WRONG?** Give the correct slope.

18. **Concept Check** A student found the slope of the line through the points $(-3, 4)$ and $(-5, -1)$ as follows.

$$m = \frac{-3 - (-5)}{4 - (-1)} = \frac{2}{5}$$

This is incorrect. **WHAT WENT WRONG?** Give the correct slope.

Evaluate each expression for m , applying the slope formula. See Example 1.

19. $m = \frac{6 - 2}{5 - 3}$

20. $m = \frac{6 - 0}{0 - 3}$

21. $m = \frac{-5 - (-5)}{3 - 2}$

22. $m = \frac{-2 - (-2)}{4 - (-3)}$

23. $m = \frac{3 - 8}{-2 - (-2)}$

24. $m = \frac{5 - 6}{-8 - (-8)}$

In the following exercises, (a) find the slope of the line passing through each pair of points, if possible, and (b) based on the slope, indicate whether the line rises from left to right, falls from left to right, is horizontal, or is vertical. See Examples 1 and 3 and FIGURE 22.

25. $(-2, -3)$ and $(-1, 5)$ 26. $(-4, 1)$ and $(-3, 4)$ 27. $(-4, 1)$ and $(2, 6)$

28. $(-3, -3)$ and $(5, 6)$ 29. $(2, 4)$ and $(-4, 4)$ 30. $(-6, 3)$ and $(2, 3)$

31. $(-2, 2)$ and $(4, -1)$ 32. $(-3, 1)$ and $(6, -2)$ 33. $(5, -3)$ and $(5, 2)$

34. $(4, -1)$ and $(4, 3)$ 35. $(1.5, 2.6)$ and $(0.5, 3.6)$ 36. $(3.4, 4.2)$ and $(1.4, 10.2)$

Extending Skills Find the slope of the line passing through the given pair of points.

(Hint: $\frac{\frac{a}{b}}{\frac{c}{d}} = \frac{a}{b} \div \frac{c}{d}$)

37. $(\frac{1}{6}, \frac{1}{2})$ and $(\frac{5}{6}, \frac{9}{2})$

38. $(\frac{3}{4}, \frac{1}{3})$ and $(\frac{5}{4}, \frac{10}{3})$

39. $(-\frac{2}{9}, \frac{5}{18})$ and $(\frac{1}{18}, -\frac{5}{9})$

40. $(-\frac{4}{5}, \frac{9}{10})$ and $(-\frac{3}{10}, \frac{1}{5})$

Each table of values gives several points that lie on a line. Find the slope of the line.

41.

x	y
-1	8
0	6
2	2
3	0

42.

x	y
-3	6
-1	0
0	-3
2	-9





43.

x	y
-6	-4
-3	0
0	4
3	8

44.

x	y
-5	-4
0	-2
5	0
10	2

Use the geometric interpretation of slope (“rise over run”) to find the slope of each line. (Coordinates of the points shown are integers.)



Find the slope of each line in three ways by doing the following.

(a) Give any two points that lie on the line, and use them to determine the slope. See Example 2.

(b) Solve the equation for y, and identify the slope from the equation. See Example 4.

(c) For the form $Ax + By = C$, calculate $-\frac{A}{B}$. See the Note following Example 4.

49. $2x - y = 8$

50. $3x - y = -6$

51. $3x + 4y = 12$

52. $6x + 5y = 30$

53. $x + y = -3$

54. $x - y = 4$

Find the slope of each line, and sketch its graph. See Examples 1–4.

55. $x + 2y = 4$

56. $x + 3y = -6$

57. $5x - 2y = 10$

58. $4x - y = 4$

59. $y = 4x$

60. $y = -3x$

61. $x - 3 = 0$

62. $x + 2 = 0$

63. $y = -5$

64. $y = -4$

65. $2y = 3$

66. $3x = 4$

Graph each line passing through the given point and having the given slope. See Example 5.

67. $(-4, 2); m = \frac{1}{2}$

68. $(-2, -3); m = \frac{5}{4}$

69. $(0, -2); m = -\frac{2}{3}$

70. $(0, -4); m = -\frac{3}{2}$

71. $(-1, -2); m = 3$

72. $(-2, -4); m = 4$

73. $(0, 0); m = \frac{1}{5}$

74. $(0, 0); m = \frac{5}{3}$

75. $(2, -5); m = 0$

76. $(5, 3); m = 0$

77. $(-3, 1);$ undefined slope

78. $(-4, 1);$ undefined slope


79. **Concept Check** If a line has slope $-\frac{4}{9}$, then any line parallel to it has slope _____, and any line perpendicular to it has slope _____.
80. **Concept Check** If a line has slope 0.2, then any line parallel to it has slope _____, and any line perpendicular to it has slope _____.

Determine whether each pair of lines is parallel, perpendicular, or neither. See Examples 6 and 7.


81. The line passing through (15, 9) and (12, -7) and the line passing through (8, -4) and (5, -20)
82. The line passing through (4, 6) and (-8, 7) and the line passing through (-5, 5) and (7, 4)
83. $x + 4y = 7$ and $4x - y = 3$
84. $2x + 5y = -7$ and $5x - 2y = 1$
85. $4x - 3y = 6$ and $3x - 4y = 2$
86. $2x + y = 6$ and $x - y = 4$
87. $x = 6$ and $6 - x = 8$
88. $3x = y$ and $2y - 6x = 5$
89. $4x + y = 0$ and $5x - 8 = 2y$
90. $2x + 5y = -8$ and $6 + 2x = 5y$
91. $4x - 3y = 8$ and $4y + 3x = 12$
92. $2x = y + 3$ and $2y + x = 3$
93. $x = 6$ and $y = 4$
94. $x + 1 = 0$ and $y - 2 = 0$

Extending Skills Solve each problem.




95. The upper deck at Guaranteed Rate Field in Chicago has produced, among other complaints, displeasure with its steepness. It is 160 ft from home plate to the front of the upper deck and 250 ft from home plate to the back. The top of the upper deck is 63 ft above the bottom. What is its slope?



96. When designing the TD Garden in Boston, architects designed the ramps leading up to the entrances so that circus elephants would be able to walk up the ramps. The maximum grade (or slope) that an elephant will walk on is 13%. Suppose that such a ramp was constructed with a horizontal run of 150 ft. What would be the maximum vertical rise the architects could use?




Find and interpret the average rate of change illustrated in each graph. See Objective 5 and FIGURE 27.



100. **Concept Check** If the graph of a linear equation rises from left to right, then the average rate of change is (positive / negative). If the graph of a linear equation falls from left to right, then the average rate of change is (positive / negative).

Solve each problem. See Examples 8 and 9.


- 101.** The graph shows the number of wireless subscriber connections (that is, active devices, including smartphones, feature phones, tablets, etc.) in millions in the United States for the years 2011 to 2016.



Data from CTIA.

- In the context of this graph, what does the ordered pair $(2016, 396)$ mean?
- Use the given ordered pairs to find the slope of the line.
- Interpret the slope in the context of this problem.


- 102.** The graph shows the percent of households in the United States that were wireless-only households for the years 2011 to 2016.



Data from CTIA.

- In the context of this graph, what does the ordered pair $(2016, 51)$ mean?
- Use the given ordered pairs to find the slope of the line.
- Interpret the slope in the context of this problem.


- 103.** The graph shows the number of drive-in movie theaters in the United States from 2010 through 2017.



Data from www.drive-ins.com

- Use the given ordered pairs to find the average rate of change in the number of drive-in theaters per year during this period. Round the answer to the nearest whole number.
- Explain how a negative slope is interpreted in this situation.

104. The graph shows the number of U.S. travelers to Canada (in thousands) from 2000 through 2016.



Data from U.S. Department of Commerce.

- (a) Use the given ordered pairs to find the average rate of change in the number of U.S. travelers to Canada per year during this period. Round the answer to the nearest thousand.
- (b) Explain how a negative slope is interpreted in this situation.
105. The average price of a gallon of unleaded gasoline in 2000 was \$1.51. In 2016, the average price was \$2.14. Find and interpret the average rate of change in the price of a gallon of gasoline per year to the nearest cent. (Data from Energy Information Administration.)
106. The average price of a movie ticket in 2004 was \$6.21. In 2016, the average price was \$8.65. Find and interpret the average rate of change in the price of a movie ticket per year to the nearest cent. (Data from Motion Picture Association of America.)
107. In 2010, the number of digital cameras shipped worldwide totaled 122 million. There were 24 million shipped in 2016. Find and interpret the average rate of change in the number of digital cameras shipped worldwide per year to the nearest million. (Data from CIPA.)
108. In 2010, worldwide shipments of desktop computers totaled 157 million units. In 2016, 103 million units were shipped. Find and interpret the average rate of change in worldwide shipments of desktop computers per year. (Data from IDC.)

Extending Skills Use your knowledge of the slopes of parallel and perpendicular lines.

109. Show that $(-13, -9)$, $(-11, -1)$, $(2, -2)$, and $(4, 6)$ are the vertices of a parallelogram. (Hint: A parallelogram is a four-sided figure with opposite sides parallel.)
110. Is the figure with vertices at $(-11, -5)$, $(-2, -19)$, $(12, -10)$, and $(3, 4)$ a parallelogram? Is it a rectangle? (Hint: A rectangle is a parallelogram with a right angle.)

RELATING CONCEPTS For Individual or Group Work (Exercises 111–116)

Three points that lie on the same straight line are said to be **collinear**. Consider the points $A(3, 1)$, $B(6, 2)$, and $C(9, 3)$. **Work Exercises 111–116 in order.**

111. Find the slope of segment AB .
112. Find the slope of segment BC .
113. Find the slope of segment AC .
114. If slope of segment $AB =$ slope of segment $BC =$ slope of segment AC , then A , B , and C are collinear. Use the results of **Exercises 111–113** to show that this statement is satisfied.
115. Use the slope formula to determine whether the points $(1, -2)$, $(3, -1)$, and $(5, 0)$ are collinear.
116. Repeat **Exercise 115** for the points $(0, 6)$, $(4, -5)$, and $(-2, 12)$.

2.3 Writing Equations of Lines

OBJECTIVES

- 1 Write an equation of a line given its slope and y-intercept.
- 2 Graph a line using its slope and y-intercept.
- 3 Write an equation of a line given its slope and a point on the line.
- 4 Write an equation of a line given two points on the line.
- 5 Write equations of horizontal and vertical lines.
- 6 Write an equation of a line parallel or perpendicular to a given line.
- 7 Write an equation of a line that models real data.

VOCABULARY

□ scatter diagram

NOW TRY EXERCISE 1

Write an equation of the line with slope $\frac{2}{3}$ and y-intercept $(0, 1)$.

NOW TRY ANSWER

1. $y = \frac{2}{3}x + 1$

OBJECTIVE 1 Write an equation of a line given its slope and y-intercept.

Recall that we can find the slope of a line from its equation by solving the equation for y . For example, the slope of the line with equation

$$y = 3x + 6$$

is 3, the coefficient of x . *What does the number 6 represent?*

To answer this question, suppose a line has slope m and y-intercept $(0, b)$. We can find an equation of this line by choosing another point (x, y) on the line, as shown in FIGURE 30, and applying the slope formula.

$$m = \frac{y - b}{x - 0} \quad \begin{array}{l} \leftarrow \text{Change in } y \\ \leftarrow \text{Change in } x \end{array}$$

$$m = \frac{y - b}{x} \quad \text{Subtract.}$$

$$mx = y - b \quad \text{Multiply by } x.$$

$$mx + b = y \quad \text{Add } b.$$

$$y = mx + b \quad \text{Interchange sides.}$$




FIGURE 30

The last equation is the *slope-intercept form* of the equation of a line, because we can identify the slope m and y-intercept $(0, b)$ at a glance. In the line with equation $y = 3x + 6$, the number 6 indicates that the y-intercept is $(0, 6)$.

Slope-Intercept Form

The **slope-intercept form** of the equation of a line with slope m and y-intercept $(0, b)$ is

$$y = mx + b.$$

Slope \uparrow \uparrow $(0, b)$ is the y-intercept.

EXAMPLE 1 Writing an Equation of a Line

Write an equation of the line with slope $-\frac{4}{5}$ and y-intercept $(0, -2)$.

Here, $m = -\frac{4}{5}$ and $b = -2$. Substitute these values into the slope-intercept form.

$$y = mx + b \quad \text{Slope-intercept form}$$

$$y = -\frac{4}{5}x + (-2) \quad \text{Let } m = -\frac{4}{5} \text{ and } b = -2.$$

$$y = -\frac{4}{5}x - 2 \quad \text{Definition of subtraction}$$

NOW TRY

NOTE Every linear equation (of a nonvertical line) has a *unique* (one and only one) slope-intercept form. *Linear functions* are defined using slope-intercept form. Also, this form is used in graphing a line with a graphing calculator.

OBJECTIVE 2 Graph a line using its slope and y-intercept.

We first saw this approach in the previous section.


NOW TRY EXERCISE 2

Graph the line using its slope and y-intercept.

$$4x + 3y = 6$$

EXAMPLE 2 Graphing Lines Using Slope and y-Intercept

Graph each line using its slope and y-intercept.

(a) $y = 3x - 6$ (In slope-intercept form)

Here, $m = 3$ and $b = -6$. Plot the y-intercept $(0, -6)$. The slope 3 can be interpreted geometrically as follows.

$$m = \frac{\text{rise}}{\text{run}} = \frac{\text{change in } y}{\text{change in } x} = \frac{3}{1}$$

From $(0, -6)$, move *up* 3 units and to the *right* 1 unit, and plot a second point at $(1, -3)$. Join the two points with a straight line. See **FIGURE 31**.




FIGURE 31

(b) $3y + 2x = 9$ (Not in slope-intercept form)

Write the equation in slope-intercept form by solving for y .

$$3y + 2x = 9$$

$$3y = -2x + 9 \quad \text{Subtract } 2x.$$

$$\text{Slope-intercept form} \rightarrow y = -\frac{2}{3}x + 3 \quad \text{Divide by } 3.$$

$$\text{Slope} \rightarrow \uparrow \quad \uparrow \text{ } y\text{-intercept is } (0, 3).$$

To graph this equation, plot the y-intercept $(0, 3)$. The slope can be interpreted as either $-\frac{2}{3}$ or $\frac{2}{-3}$. Using $-\frac{2}{3}$, begin at $(0, 3)$ and move *down* 2 units and to the *right* 3 units to locate the point $(3, 1)$. The line through these two points is the required graph. See **FIGURE 32**. (Verify that the point obtained using $\frac{2}{-3}$ as the slope is also on this line.)





FIGURE 32

NOW TRY ANSWER

OBJECTIVE 3 Write an equation of a line given its slope and a point on the line.

Let m represent the slope of a line and (x_1, y_1) represent a given point on the line. Let (x, y) represent any other point on the line. See **FIGURE 33**.

$$m = \frac{y - y_1}{x - x_1} \quad \text{Slope formula}$$

$$m(x - x_1) = y - y_1 \quad \text{Multiply each side by } x - x_1.$$

$$y - y_1 = m(x - x_1) \quad \text{Interchange sides.}$$




FIGURE 33

This is the *point-slope form* of the equation of a line.

Point-Slope Form

The **point-slope form** of the equation of a line with slope m passing through the point (x_1, y_1) is

$$y - y_1 = m(x - x_1).$$

↓ Slope
↑ Given point ↑

NOW TRY EXERCISE 3

Write an equation of the line with slope $-\frac{1}{5}$ passing through the point $(5, -3)$.

EXAMPLE 3 Writing an Equation of a Line Given Its Slope and a Point

Write an equation of the line with slope $\frac{1}{3}$ passing through the point $(-2, 5)$.

Method 1 Use **point-slope form**. Let $(x_1, y_1) = (-2, 5)$ and $m = \frac{1}{3}$.

$$y - y_1 = m(x - x_1) \quad \text{Point-slope form}$$

Leave x and y as variables. $y - 5 = \frac{1}{3}[x - (-2)]$ Let $y_1 = 5$, $m = \frac{1}{3}$, and $x_1 = -2$.

$$y - 5 = \frac{1}{3}(x + 2) \quad \text{Definition of subtraction}$$

$$y - 5 = \frac{1}{3}x + \frac{2}{3} \quad (*) \text{ Distributive property}$$

$$y = \frac{1}{3}x + \frac{2}{3} + \frac{15}{3} \quad \text{Add } 5 = \frac{15}{3}.$$

Slope-intercept form $\rightarrow y = \frac{1}{3}x + \frac{17}{3}$ $\frac{2}{3} + \frac{15}{3} = \frac{17}{3}$

Method 2 Use **slope-intercept form**. Let $(x, y) = (-2, 5)$ and $m = \frac{1}{3}$.

$$y = mx + b \quad \text{Slope-intercept form}$$

$$5 = \frac{1}{3}(-2) + b \quad \text{Let } y = 5, m = \frac{1}{3}, \text{ and } x = -2.$$

Solve for b . $5 = -\frac{2}{3} + b$ Multiply.

$5 = \frac{15}{3}$ $\frac{17}{3} = b$, or $b = \frac{17}{3}$ Add $\frac{2}{3}$; $\frac{15}{3} + \frac{2}{3} = \frac{17}{3}$

NOW TRY ANSWER

3. $y = -\frac{1}{5}x - 2$

Substitute $m = \frac{1}{3}$ and $b = \frac{17}{3}$ to obtain $y = \frac{1}{3}x + \frac{17}{3}$, as above.

Previously we defined *standard form* for a linear equation.

$$Ax + By = C \quad \text{Standard form}$$

Here A , B , and C are real numbers and A and B are not both 0. **For consistency, we give A , B , and C as integers with greatest common factor 1, and $A \geq 0$.** (If $A = 0$, then we give $B > 0$.) The equation in **Example 3** is written in standard form as follows.

$$y - 5 = \frac{1}{3}x + \frac{2}{3} \quad \text{Equation (*) from Example 3}$$

$$3y - 15 = x + 2 \quad \text{Multiply each term by 3.}$$

$$-x + 3y = 17 \quad \text{Subtract } x. \text{ Add 15.}$$

$$\text{Standard form} \rightarrow x - 3y = -17 \quad \text{Multiply by } -1.$$

NOTE “Standard form” is not standard among texts. A linear equation can be written correctly in many different ways.

Example:

$$\underbrace{2x + 3y = 8,}_{\substack{\text{Standard form} \\ \text{as described above}}} \quad \underbrace{2x = 8 - 3y, \quad 3y = 8 - 2x, \quad x + \frac{3}{2}y = 4, \quad 4x + 6y = 16}_{\text{Equivalent forms of } 2x + 3y = 8}$$

OBJECTIVE 4 Write an equation of a line given two points on the line.

 **NOW TRY EXERCISE 4**

Write an equation of the line passing through the points $(3, -4)$ and $(-2, -1)$. Give the final answer in standard form.

EXAMPLE 4 Writing an Equation of a Line Given Two Points

Write an equation of the line passing through the points $(-4, 3)$ and $(5, -7)$. Give the final answer in standard form.

First find the slope using the slope formula.

$$m = \frac{-7 - 3}{5 - (-4)} = -\frac{10}{9}$$

Use either $(-4, 3)$ or $(5, -7)$ as (x_1, y_1) in the point-slope form of the equation of a line. We choose $(-4, 3)$, so $-4 = x_1$ and $3 = y_1$.

$$y - y_1 = m(x - x_1) \quad \text{Point-slope form}$$

$$y - 3 = -\frac{10}{9}[x - (-4)] \quad \text{Let } y_1 = 3, m = -\frac{10}{9}, \text{ and } x_1 = -4.$$

$$y - 3 = -\frac{10}{9}(x + 4) \quad \text{Definition of subtraction}$$

$$y - 3 = -\frac{10}{9}x - \frac{40}{9} \quad \text{Distributive property}$$

$$9y - 27 = -10x - 40 \quad \text{Multiply each term by 9.}$$

$$\text{Standard form} \rightarrow 10x + 9y = -13 \quad \text{Add } 10x. \text{ Add } 27.$$

Verify that if $(5, -7)$ were used, the same equation would result.

NOW TRY 

NOW TRY ANSWER

4. $3x + 5y = -11$

NOTE Once the slope is found in **Example 4**, the equation of the line could also be determined using slope-intercept form.

OBJECTIVE 5 Write equations of horizontal and vertical lines.

A horizontal line has slope 0. Using point-slope form, we can find the equation of a horizontal line through the point (a, b) .

$$y - y_1 = m(x - x_1) \quad \text{Point-slope form}$$

$$y - b = 0(x - a) \quad y_1 = b, m = 0, x_1 = a$$

$$y - b = 0 \quad \text{Multiplication property of 0}$$

$$\text{Horizontal line} \rightarrow y = b \quad \text{Add } b.$$

Point-slope form does not apply to a vertical line because the slope of a vertical line is undefined. A vertical line through the point (a, b) has equation $x = a$.

Equations of Horizontal and Vertical Lines

A **horizontal line** through the point (a, b) has equation $y = b$.

A **vertical line** through the point (a, b) has equation $x = a$.

NOW TRY EXERCISE 5

Write an equation of the line passing through the point $(4, -4)$ and satisfying the given condition.

- (a) The line has undefined slope.
 (b) The line has slope 0.

EXAMPLE 5 Writing Equations of Horizontal and Vertical Lines

Write an equation of the line passing through the point $(-3, 3)$ and satisfying the given condition.

- (a) The line has slope 0.

Because the slope is 0, this is a horizontal line. A horizontal line through the point (a, b) has equation $y = b$. In $(-3, 3)$, the y -coordinate is 3, so the equation is $y = 3$.

- (b) The line has undefined slope.

This is a vertical line because the slope is undefined. A vertical line through the point (a, b) has equation $x = a$. In $(-3, 3)$, the x -coordinate is -3 , so the equation is $x = -3$.

Both lines are graphed in **FIGURE 34**.




FIGURE 34

NOW TRY

OBJECTIVE 6 Write an equation of a line parallel or perpendicular to a given line.

EXAMPLE 6 Writing Equations of Parallel or Perpendicular Lines

Write an equation of the line passing through the point $(-3, 6)$ and satisfying the given condition. Give final answers in slope-intercept form.

- (a) The line is parallel to the line $2x + 3y = 6$.

We can find the slope of the given line by solving for y .

$$2x + 3y = 6$$

$$3y = -2x + 6 \quad \text{Subtract } 2x.$$

$$y = -\frac{2}{3}x + 2 \quad \text{Divide by 3.}$$

↑ Slope

NOW TRY ANSWERS

5. (a) $x = 4$ (b) $y = -4$

NOW TRY
EXERCISE 6

Write an equation of the line passing through the point $(6, -1)$ and satisfying the given condition. Give final answers in slope-intercept form.

- (a) The line is parallel to the line $3x - 5y = 7$.
- (b) The line is perpendicular to the line $3x - 5y = 7$.

The slope of the line is given by the coefficient of x , so $m = -\frac{2}{3}$. See **FIGURE 35**.

The required equation of the line through $(-3, 6)$ and *parallel* to $2x + 3y = 6$ must have the *same* slope $-\frac{2}{3}$. To find this equation, we use the point-slope form with $(x_1, y_1) = (-3, 6)$ and $m = -\frac{2}{3}$.

$$y - y_1 = m(x - x_1) \quad \text{Point-slope form}$$

$$y - 6 = -\frac{2}{3}[x - (-3)] \quad y_1 = 6, m = -\frac{2}{3}, x_1 = -3$$

$$y - 6 = -\frac{2}{3}(x + 3) \quad \text{Definition of subtraction}$$

$$y - 6 = -\frac{2}{3}x - 2 \quad \text{Distributive property}$$

$$y = -\frac{2}{3}x + 4 \quad \text{Add 6.}$$





FIGURE 35



Both lines have slope $-\frac{2}{3}$. The lines are parallel.

FIGURE 36

We did not clear the fraction because we want the final equation in slope-intercept form—that is, solved for y . Both lines are shown in **FIGURE 36**.

- (b) The line is perpendicular to the line $2x + 3y = 6$.

In part (a), we wrote the equation of the given line in slope-intercept form.

$$2x + 3y = 6$$

$$y = -\frac{2}{3}x + 2 \quad \text{Slope-intercept form}$$

↑
Slope

To be *perpendicular* to the line $2x + 3y = 6$, a line must have slope $\frac{3}{2}$, the *negative reciprocal* of $-\frac{2}{3}$.

We use $(-3, 6)$ and slope $\frac{3}{2}$ in the point-slope form to find the equation of the perpendicular line shown in **FIGURE 37**.

$$y - y_1 = m(x - x_1) \quad \text{Point-slope form}$$


$$y - 6 = \frac{3}{2}[x - (-3)] \quad y_1 = 6, m = \frac{3}{2}, x_1 = -3$$

$$y - 6 = \frac{3}{2}(x + 3) \quad \text{Definition of subtraction}$$

$$y - 6 = \frac{3}{2}x + \frac{9}{2} \quad \text{Distributive property}$$

$$y = \frac{3}{2}x + \frac{9}{2} + \frac{12}{2} \quad \text{Add } 6 = \frac{12}{2}.$$

$$y = \frac{3}{2}x + \frac{21}{2} \quad \frac{9}{2} + \frac{12}{2} = \frac{21}{2}$$



The slopes are negative reciprocals, $-\frac{2}{3}(\frac{3}{2}) = -1$. The lines are perpendicular.

FIGURE 37

NOW TRY ANSWERS

- 6. (a) $y = \frac{3}{5}x - \frac{23}{5}$
- (b) $y = -\frac{5}{3}x + 9$

▼ Summary of Forms of Linear Equations

Equation	Description	When to Use
$y = mx + b$	Slope-Intercept Form Slope is m . y-intercept is $(0, b)$.	The slope and y-intercept can be easily identified and used to quickly graph the equation.
$y - y_1 = m(x - x_1)$	Point-Slope Form Slope is m . Line passes through (x_1, y_1) .	This form is ideal for finding the equation of a line if the slope and a point on the line or two points on the line are known.
$Ax + By = C$	Standard Form (A, B , and C integers, $A \geq 0$) Slope is $-\frac{A}{B}$ ($B \neq 0$). x-intercept is $(\frac{C}{A}, 0)$ ($A \neq 0$). y-intercept is $(0, \frac{C}{B})$ ($B \neq 0$).	The x- and y-intercepts can be found quickly and used to graph the equation. The slope must be calculated.
$y = b$	Horizontal Line Slope is 0. y-intercept is $(0, b)$.	If the graph intersects only the y-axis, then y is the only variable in the equation.
$x = a$	Vertical Line Slope is undefined. x-intercept is $(a, 0)$.	If the graph intersects only the x-axis, then x is the only variable in the equation.

OBJECTIVE 7 Write an equation of a line that models real data.

If a given set of data changes at a fairly constant rate, the data may fit a linear pattern, where the rate of change is the slope of the line.

EXAMPLE 7 Writing a Linear Equation to Describe Real Data

A local gasoline station is selling 89-octane gas for \$3.50 per gal.

(a) Write an equation that describes the cost y to buy x gallons of gas.

The total cost is determined by the number of gallons multiplied by the price per gallon (in this case, \$3.50). As the gas is pumped, two sets of numbers spin by: the number of gallons pumped and the cost of that number of gallons.

The table illustrates this situation.

Number of Gallons Pumped	Cost of This Number of Gallons
0	$0(\$3.50) = \$ 0.00$
1	$1(\$3.50) = \$ 3.50$
2	$2(\$3.50) = \$ 7.00$
3	$3(\$3.50) = \10.50
4	$4(\$3.50) = \14.00



If we let x denote the number of gallons pumped, then the total cost y in dollars can be found using the following linear equation.

$$\begin{array}{ccc} \text{Total cost} & \searrow & \swarrow \text{Number of gallons} \\ & y = 3.50x & \end{array}$$

Theoretically, there are infinitely many ordered pairs (x, y) that satisfy this equation, but here we are limited to nonnegative values for x because we cannot have a negative number of gallons. In this situation, there is also a practical maximum value for x , which varies from one car to another—the size of the gas tank.

NOW TRY
EXERCISE 7

A cell phone plan costs \$100 for the telephone plus \$85 per month for service. Write an equation that gives the cost y in dollars for x months of cell phone service using this plan.

- (b) A car wash at this gas station costs an additional \$3.00. Write an equation that defines the cost of gas and a car wash.

The cost will be $3.50x + 3.00$ dollars for x gallons of gas and a car wash.

$$y = 3.5x + 3 \quad \text{Final 0's need not be included.}$$

- (c) Interpret the ordered pairs $(5, 20.5)$ and $(10, 38)$ in the context of the equation from part (b).

$(5, 20.5)$ indicates that 5 gal of gas and a car wash cost \$20.50.

$(10, 38)$ indicates that 10 gal of gas and a car wash cost \$38.00.

NOW TRY

NOTE In **Example 7(a)**, the ordered pair $(0, 0)$ satisfied the equation $y = 3.50x$, so the linear equation has the form

$$y = mx, \quad \text{where } b = 0.$$

If a realistic situation involves an initial charge b plus a charge per unit m , as in the equation $y = 3.5x + 3$ in **Example 7(b)**, then the equation has the form

$$y = mx + b, \quad \text{where } b \neq 0.$$

EXAMPLE 8 Writing an Equation of a Line That Models Data

Average annual tuition and fees for in-state students at public two-year colleges are shown in the table for selected years and graphed as ordered pairs of points in the **scatter diagram** in **FIGURE 38**, where $x = 0$ represents 2012, $x = 1$ represents 2013, and so on, and y represents the cost in dollars.

Year	Cost (in dollars)
2012 ($x = 0$)	3216
2013 ($x = 1$)	3264
2014 ($x = 2$)	3347
2015 ($x = 3$)	3435
2016 ($x = 4$)	3520
2017 ($x = 5$)	3570

Data from The College Board.




FIGURE 38

- (a) Write an equation that models the data.

Because the points in **FIGURE 38** lie approximately on a straight line, we can write a linear equation that models the relationship between year x and cost y . We choose two data points, $(0, 3216)$ and $(5, 3570)$, to find the slope of the line.

$$m = \frac{3570 - 3216}{5 - 0} = \frac{354}{5} = 70.8$$

The slope 70.8 indicates that the cost of tuition and fees increased by about \$71 per year from 2012 to 2017. We use this slope and the y -intercept $(0, 3216)$ to write an equation of the line in slope-intercept form.

$$y = 70.8x + 3216$$

NOW TRY ANSWER

7. $y = 85x + 100$

NOW TRY EXERCISE 8

Refer to **Example 8**.

- (a) Using the data values for the years 2012 and 2016, write an equation that models the data.
- (b) Use the equation from part (a) to approximate the cost of tuition and fees in 2015.

- (b) Use the equation from part (a) to approximate the cost of tuition and fees in 2018.
The value $x = 6$ corresponds to the year 2018.

$$y = 70.8x + 3216 \quad \text{Equation from part (a)}$$

$$y = 70.8(6) + 3216 \quad \text{Substitute 6 for } x.$$

$$y = 3640.8 \quad \text{Multiply, and then add.}$$


According to the model, average tuition and fees for in-state students at public two-year colleges in 2018 were about \$3641.

NOW TRY

NOTE Choosing different data points in **Example 8** would result in a slightly different line (particularly in regard to its slope) and, hence, a slightly different equation. However, all such equations should yield similar results. See **Now Try Exercise 8**.

EXAMPLE 9 Writing an Equation of a Line That Models Data

Hospital expenditures (in billions of dollars) in the United States are shown in the graph in **FIGURE 39**.



Data from Centers for Medicare & Medicaid Services.

FIGURE 39

- (a) Write an equation that models the data.

The data increase linearly—that is, a straight line through the tops of any two bars in the graph would be close to the top of each bar. To model the relationship between year x and hospital expenditures y , we let $x = 1$ represent 2011, $x = 2$ represent 2012, and so on. The given data for 2011 and 2016 can be written as the ordered pairs $(1, 852)$ and $(6, 1082)$.

$$m = \frac{1082 - 852}{6 - 1} = \frac{230}{5} = 46 \quad \text{Find the slope of the line through } (1, 852) \text{ and } (6, 1082).$$

Thus, hospital expenditures increased by \$46 billion per year. To write an equation, we substitute this slope and the point $(1, 852)$ into the point-slope form.

$$y - y_1 = m(x - x_1) \quad \text{Point-slope form}$$

$$y - 852 = 46(x - 1) \quad (x_1, y_1) = (1, 852); m = 46$$

$$y - 852 = 46x - 46 \quad \text{Distributive property}$$

$$y = 46x + 806 \quad \text{Add 852.}$$

Either point can be used here. $(6, 1082)$ provides the same answer.

NOW TRY ANSWERS

- 8. (a) $y = 76x + 3216$
- (b) \$3444


NOW TRY EXERCISE 9
Refer to **Example 9**.

- (a) Use the ordered pairs (4, 978) and (6, 1082) to write an equation that models the data.
- (b) Use the equation from part (a) to estimate hospital expenditures in 2019.

- (b) Use the equation from part (a) to estimate hospital expenditures in the United States in 2019. (Assume a constant rate of change.)

Because we let $x = 1$ represent 2011, $x = 9$ represents 2019.

$$y = 46x + 806 \quad \text{Equation from part (a)}$$

$$y = 46(9) + 806 \quad \text{Substitute 9 for } x.$$

$$y = 1220 \quad \text{Multiply, and then add.}$$

Hospital expenditures in the United States in 2019 were about \$1220 billion.


NOW TRY 
NOW TRY ANSWERS

9. (a) $y = 52x + 770$
 (b) \$1238 billion

2.3 Exercises

FOR EXTRA HELP

MyLab Math

 Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

How are you doing on your homework? **Review Study Skill 4, Completing Your Homework.**


Concept Check Provide the appropriate response.

- The following equations all represent the same line. Which one is in standard form as specified in this section?
 A. $3x - 2y = 5$ B. $y = \frac{3}{2}x - \frac{5}{2}$ C. $\frac{3}{5}x - \frac{2}{5}y = 1$ D. $3x = 2y + 5$
- Which equation is in point-slope form?
 A. $y = 6x + 2$ B. $4x + y = 9$ C. $y - 3 = 2(x - 1)$ D. $2y = 3x - 7$
- Which equation in **Exercise 2** is in slope-intercept form?
- Write the equation $y + 2 = -3(x - 4)$ in slope-intercept form.
- Write the equation $y + 2 = -3(x - 4)$ in standard form.
- Write the equation $10x - 7y = 70$ in slope-intercept form.


Concept Check Match each equation with the graph that it most closely resembles. (Hint: Determining the signs of m and b will help in each case.)

7. $y = 2x + 3$


A.



B.



C.



8. $y = -2x + 3$

9. $y = -2x - 3$

10. $y = 2x - 3$


11. $y = 2x$

12. $y = -2x$


13. $y = 3$

14. $y = -3$


D.




E.




F.




G.



H.



I.



Write an equation in slope-intercept form of the line that satisfies the given conditions. See Example 1.

15. $m = 5; b = 15$

17. $m = -\frac{2}{3}; b = \frac{4}{5}$

19. Slope 1; y-intercept $(0, -1)$

21. Slope $\frac{2}{5}$; y-intercept $(0, 5)$





16. $m = 2; b = 12$

18. $m = -\frac{5}{8}; b = -\frac{1}{3}$

20. Slope -1 ; y-intercept $(0, -3)$

22. Slope $-\frac{3}{4}$; y-intercept $(0, 7)$

Write an equation in slope-intercept form of the line shown in each graph. (Coordinates of the points shown are integers.)



Each table of values gives several points that lie on a line. Write an equation in slope-intercept form of the line.

 27.

x	y
-2	-8
0	-4
1	-2
3	2

 28.

x	y
-2	-3
0	3
2	9
3	12

 29.

x	y
-5	6
0	3
5	0
10	-3

 30.

x	y
-4	5
-2	0
0	-5
2	-10

For each equation, (a) write it in slope-intercept form, (b) give the slope of the line, (c) give the y-intercept, and (d) graph the line. See Example 2.

31. $-x + y = 4$

33. $6x + 5y = 30$

35. $4x - 5y = 20$

37. $x + 2y = -9$

32. $-x + y = 6$

34. $3x + 4y = 12$

36. $7x - 3y = 3$

38. $x + 3y = -9$

Write an equation of the line passing through the given point and having the given slope. Give the equation (a) in slope-intercept form and (b) in standard form. See Example 3 and the discussion on standard form.

39. $(5, 8)$; slope -2

41. $(-2, 4)$; slope $-\frac{3}{4}$

43. $(-5, 4)$; slope $\frac{1}{2}$

45. $(3, 0)$; slope 4

47. $(2, 6.8)$; slope 1.4

40. $(12, 10)$; slope 1

42. $(-1, 6)$; slope $-\frac{5}{6}$

44. $(7, -2)$; slope $\frac{1}{4}$

46. $(-2, 0)$; slope -5

48. $(6, -1.2)$; slope 0.8

Write an equation of the line passing through the given points. Give the final answer in standard form. See Example 4.

49. $(3, 4)$ and $(5, 8)$

51. $(6, 1)$ and $(-2, 5)$

53. $(2, 5)$ and $(1, 5)$

50. $(5, -2)$ and $(-3, 14)$

52. $(-2, 5)$ and $(-8, 1)$

54. $(-2, 2)$ and $(4, 2)$

55. $(7, 6)$ and $(7, -8)$

56. $(13, 5)$ and $(13, -1)$

57. $\left(\frac{1}{2}, -3\right)$ and $\left(-\frac{2}{3}, -3\right)$

58. $\left(-\frac{4}{9}, -6\right)$ and $\left(\frac{12}{7}, -6\right)$

59. $\left(-\frac{2}{5}, \frac{2}{5}\right)$ and $\left(\frac{4}{3}, \frac{2}{3}\right)$

60. $\left(\frac{3}{4}, \frac{8}{3}\right)$ and $\left(\frac{2}{5}, \frac{2}{3}\right)$

Write an equation of the line passing through the given point and satisfying the given condition. See Example 5.

61. $(9, 5)$; slope 0

62. $(-4, -2)$; slope 0

63. $(9, 10)$; undefined slope

64. $(-2, 8)$; undefined slope

65. $\left(-\frac{3}{4}, -\frac{3}{2}\right)$; slope 0

66. $\left(-\frac{5}{8}, -\frac{9}{2}\right)$; slope 0

67. $(-7, 8)$; horizontal

68. $(2, -7)$; horizontal

69. $(0.5, 0.2)$; vertical

70. $(0.1, 0.4)$; vertical

71. $(0, 0)$; horizontal

72. $(0, 0)$; vertical

Write an equation of the line passing through the given point and satisfying the given condition. Give the equation (a) in slope-intercept form and (b) in standard form. See Example 6.

73. $(7, 2)$; parallel to $3x - y = 8$

74. $(4, 1)$; parallel to $2x + 5y = 10$

75. $(-2, -2)$; parallel to $-x + 2y = 10$

76. $(-1, 3)$; parallel to $-x + 3y = 12$

77. $(8, 5)$; perpendicular to $2x - y = 7$

78. $(2, -7)$; perpendicular to $5x + 2y = 18$

79. $(-2, 7)$; perpendicular to $x = 9$

80. $(8, 4)$; perpendicular to $x = -3$

Write an equation in the form $y = mx$ for each situation. Then give the three ordered pairs associated with the equation for x -values 0, 5, and 10. See Example 7(a).

81. x represents the number of hours traveling at 45 mph, and y represents the distance traveled (in miles).

82. x represents the number of caps sold at \$26 each, and y represents the total cost of the caps (in dollars).



83. x represents the number of gallons of gas sold at \$3.75 per gal, and y represents the total cost of the gasoline (in dollars).

84. x represents the number of hours a bicycle is rented at \$7.50 per hour, and y represents the total charge for the rental (in dollars).

85. x represents the number of credit hours taken at a community college at \$150 per credit hour, and y represents the total tuition paid for the credit hours (in dollars).



86. x represents the number of tickets to a performance of *Hamilton* purchased at \$250 per ticket, and y represents the total paid for the tickets (in dollars).

For each situation, do the following.

- (a) Write an equation in the form $y = mx + b$.
- (b) Find and interpret the ordered pair associated with the equation for $x = 5$.
- (c) Answer the question posed in the problem.

See Examples 7(b) and 7(c).

- 87.** A ticket for Taylor Swift's Reputation Stadium Tour costs \$140. There is a ticket order delivery fee of \$18.50. Let x represent the number of tickets and y represent the cost in dollars. How much does it cost to purchase 2 tickets and have them delivered? (Data from Ticketmaster.)
- 88.** Resident tuition at North Shore Community College is \$206 per credit hour. There is also a \$300 program fee for physical therapy. Let x represent the number of credit hours and y represent the cost in dollars. How much does it cost for a student in physical therapy to take 15 credit hours? (Data from www.northshore.edu)
- 89.** A health club membership costs \$99, plus \$41 per month. Let x represent the number of months and y represent the cost in dollars. How much does the first year's membership cost? (Data from Midwest Athletic Club.)




- 90.** An Executive VIP/Gold membership to a health club costs \$159, plus \$57 per month. Let x represent the number of months and y represent the cost in dollars. How much does a one-year membership cost? (Data from Midwest Athletic Club.)
- 91.** A wireless plan includes unlimited talk, text, and data for \$90 per month. There is a \$25 activation fee. Let x represent the number of months and y represent the cost in dollars. Over two years, how much will this plan cost?
- 92.** A wireless plan includes unlimited talk, text, and data for \$75 per month. There is a \$350 charge for a phone. Let x represent the number of months and y represent the cost in dollars. Over two years, how much will this plan cost?



- 93.** There is a \$30 fee to rent a chain saw, plus \$6 per day. Let x represent the number of days the saw is rented and y represent the charge to the user in dollars. If the total charge is \$138, for how many days is the saw rented?
- 94.** A rental car costs \$99, plus \$0.50 per mile. Let x represent the number of miles driven and y represent the total charge to the renter in dollars. How many miles was the car driven if the renter paid \$174.00?

Solve each problem. Give equations in slope-intercept form. See Examples 8 and 9.


95. Total sales of e-readers in the United States (in millions of dollars) are shown in the graph, where the year 2013 corresponds to $x = 0$.



*Data for this year are unavailable. Data from Consumer Technology Association.

- Use the ordered pairs from the graph to write an equation that models the data. (Round the slope to the nearest tenth.) Interpret the slope in the context of this problem.
- Use the equation from part (a) to approximate sales of e-readers in the United States in 2014, the year data was unavailable.


96. Total sales of smartphones in the United States (in billions of dollars) are shown in the graph, where the year 2013 corresponds to $x = 0$.



*Data for this year are unavailable. Data from Consumer Technology Association.

- Use the ordered pairs from the graph to write an equation that models the data. (Round the slope to the nearest tenth.) Interpret the slope in the context of this problem.
- Use the equation from part (a) to approximate smartphone sales in the United States in 2014, the year data was unavailable.

97. Expenditures for home health care in the United States are shown in the graph.




Data from Centers for Medicare & Medicaid Services.

- If a straight line were drawn through the tops of the bars for 2012 and 2016, would this line have positive or negative slope? Explain.
- Use the information given for the years 2012 and 2016, letting $x = 12$ represent 2012 and $x = 16$ represent 2016, and letting y represent spending (in billions of dollars), to write an equation that models the data.
- Use the equation from part (b) to approximate the amount spent on home health care in 2015. How does the result compare with the actual value, \$89 billion?

98. The number of pieces of first-class mail delivered in the United States is shown in the graph.

- (a) If a straight line were drawn through the tops of the bars for 2011 and 2016, would this line have positive or negative slope? Explain.
- (b) Use the information given for the years 2011 and 2016, letting $x = 11$ represent 2011 and $x = 16$ represent 2016, and letting y represent the number of pieces of mail (in billions), to write an equation that models the data.

First-Class Mail



Data from U.S. Postal Service.

- (c) Use the equation from part (b) to approximate the number of pieces of first-class mail delivered in 2015 to the nearest tenth. How does this result compare to the actual value, 62.6 billion?

RELATING CONCEPTS For Individual or Group Work (Exercises 99–106)

To see how the formula that relates Celsius and Fahrenheit temperatures is derived, work Exercises 99–106 in order.

99. There is a linear relationship between Celsius and Fahrenheit temperatures.

$$\text{When } C = 0^\circ, F = \text{---}^\circ.$$

$$\text{When } C = 100^\circ, F = \text{---}^\circ.$$

100. Think of ordered pairs of temperatures (C, F) , where C and F represent corresponding Celsius and Fahrenheit temperatures. The equation that relates the two scales has a straight-line graph that contains the two points determined in Exercise 99.

- (a) What are these two points?
- (b) Find the slope of the line passing through these two points.

101. Use the slope found in Exercise 100(b) and one of the two points determined earlier, and write an equation that gives F in terms of C . (Hint: Use the point-slope form, where C replaces x and F replaces y .)

102. Use the equation found in Exercise 101 and solve for C in terms of F . For what temperature does $F = C$? (Use the photo to confirm this temperature.)

103. A quick way to estimate Fahrenheit temperature for a given Celsius temperature is to double C and add 30. Use this method to estimate F if $C = 15$.

104. Use the equation found in Exercise 101 to find F if $C = 15$. How does the answer compare with the answer to Exercise 103?

105. Use the method given in Exercise 103 to estimate the Fahrenheit temperature given $C = 30$. Then use the equation from Exercise 101 to find F when $C = 30$. How do the temperatures compare?

106. Explain why the method given in Exercise 103 to estimate Fahrenheit temperature gives a good approximation of $F = \frac{9}{5}C + 32$.



SUMMARY EXERCISES Finding Slopes and Equations of Lines

Find the slope of each line passing through the given points or having the given equation.

1. $(3, -3)$ and $(8, -6)$ 2. $(4, -5)$ and $(-1, -5)$ 3. $y = x - 5$
 4. $3x - 7y = 21$ 5. $x - 4 = 0$ 6. $4x + 7y = 3$

7. **Concept Check** Match the description in Column I with its equation in Column II.

I	II
(a) Slope -0.5 , $b = -2$	A. $y = -\frac{1}{2}x$
(b) x -intercept $(4, 0)$, y -intercept $(0, 2)$	B. $y = -\frac{1}{2}x - 2$
(c) Passes through $(4, -2)$ and $(0, 0)$	C. $x - 2y = 2$
(d) $m = \frac{1}{2}$, passes through $(-2, -2)$	D. $y = 2x$
(e) $m = \frac{1}{2}$, passes through the origin	E. $x = 2y$
(f) Slope 2 , $b = 0$	F. $x + 2y = 4$

8. **Concept Check** Which equation is written in standard form as specified in the text?

- A. $y = -4x - 7$ B. $-3x + 4y = 12$ C. $x - 6y = 3$
 D. $\frac{1}{2}x + y = 0$ E. $6x - 2y = 10$ F. $3y - 5x = -15$

Write the equations not written in standard form in that form.

Write an equation of the line passing through the given point(s) and satisfying the given condition. Give the equation (a) in slope-intercept form and (b) in standard form.

9. $(4, -2)$; slope -3
 10. $(-3, 6)$; slope $\frac{2}{3}$
 11. $(-2, 6)$ and $(4, 1)$
 12. $(4, -8)$ and $(-4, 12)$
 13. $(-2, 5)$ and $(1, 14)$
 14. Origin; slope $-\frac{5}{2}$
 15. $(5, -8)$; parallel to $y = 4$
 16. $(\frac{3}{4}, -\frac{7}{9})$; perpendicular to $x = \frac{2}{3}$
 17. $(4, -2)$; perpendicular to the line passing through $(3, 7)$ and $(5, 6)$
 18. $(-4, 2)$; parallel to the line passing through $(3, 9)$ and $(6, 11)$
 19. $(2, -1)$; parallel to $y = \frac{1}{5}x + \frac{7}{4}$
 20. $(0, -6)$; perpendicular to $y = \frac{4}{3}x + \frac{3}{8}$

2.4 Linear Inequalities in Two Variables

OBJECTIVES

- 1 Graph linear inequalities in two variables.
- 2 Graph the intersection of two linear inequalities.
- 3 Graph the union of two linear inequalities.

VOCABULARY

- linear inequality in two variables
- boundary line

OBJECTIVE 1 Graph linear inequalities in two variables.

Earlier we graphed linear inequalities in one variable on a number line. In this section, we graph linear inequalities in two variables on a rectangular coordinate system.

Linear Inequality in Two Variables

A **linear inequality in two variables** (here x and y) is an inequality that can be written in the form

$$Ax + By < C, \quad Ax + By \leq C, \quad Ax + By > C, \quad \text{or} \quad Ax + By \geq C,$$

where A , B , and C are real numbers and A and B are not both 0.

Examples:

$$2x + 5y < 10, \quad x - y > 0, \quad y \geq 1 \text{ (here } A = 0\text{)}, \quad x \leq -6 \text{ (here } B = 0\text{)}$$

Consider the graph in **FIGURE 40**. The graph of the line $x + y = 5$ divides the points in the rectangular coordinate system into three sets of points.

1. Those points that lie **on** the line itself and satisfy the equation $x + y = 5$, such as $(0, 5)$, $(2, 3)$, and $(5, 0)$
2. Those points that lie in the region **above** the line and satisfy the inequality $x + y > 5$, such as $(5, 3)$ and $(2, 4)$
3. Those points that lie in the region **below** the line and satisfy the inequality $x + y < 5$, such as $(0, 0)$ and $(-3, -1)$




FIGURE 40

The graph of the line $x + y = 5$ is the **boundary line** for the two inequalities

$$x + y > 5 \quad \text{and} \quad x + y < 5.$$

A graph of a linear inequality in two variables is a region in the real number plane that may or may not include the boundary line.

To graph a linear inequality in two variables, follow these steps.

Graphing a Linear Inequality in Two Variables

Step 1 Draw the graph of the straight line that is the boundary.


- Make the line solid if the inequality involves \leq or \geq .
- Make the line dashed if the inequality involves $<$ or $>$.

Step 2 Choose a test point. Choose any point not on the line, and substitute the coordinates of that point in the inequality.

Step 3 Shade the appropriate region. Shade the region that includes the test point if it satisfies the original inequality. Otherwise, shade the region on the other side of the boundary line.


NOW TRY EXERCISE 1
Graph $-x + 2y \geq 4$.**EXAMPLE 1** Graphing a Linear InequalityGraph $3x + 2y \geq 6$.

Step 1 First graph the boundary line $3x + 2y = 6$, which has intercepts $(2, 0)$ and $(0, 3)$, as shown in **FIGURE 41**.

**FIGURE 41**

Step 2 The graph of the inequality $3x + 2y \geq 6$ includes the points of the boundary line $3x + 2y = 6$ (because the inequality symbol \geq includes equality) and either the points *above* that line or the points *below* it. To decide which, select any point *not* on the boundary line to use as a test point. Substitute the values from the test point, here $(0, 0)$, for x and y in the inequality.



$$\begin{array}{l}
 3x + 2y > 6 \quad \text{We are testing the region.} \\
 \text{\textcircled{\scriptsize $(0, 0)$ is a convenient test point.}} \quad 3(0) + 2(0) \stackrel{?}{>} 6 \quad \text{Let } x = 0 \text{ and } y = 0. \\
 0 > 6 \quad \text{False}
 \end{array}$$

Step 3 Because the result is false, $(0, 0)$ does *not* satisfy the inequality. The solution set includes all points in the region on the *other* side of the line. This region is shaded in **FIGURE 42**.

As a further check, select a test point in the shaded region, say $(3, 0)$.

$$\begin{array}{l}
 3x + 2y > 6 \quad \text{Test the region.} \\
 3(3) + 2(0) \stackrel{?}{>} 6 \quad \text{Let } x = 3 \text{ and } y = 0. \\
 9 + 0 \stackrel{?}{>} 6 \quad \text{Multiply.} \\
 9 > 6 \quad \checkmark \quad \text{True}
 \end{array}$$

The check confirms that the correct region is graphed in **FIGURE 42**.

**FIGURE 42****NOW TRY** **NOW TRY ANSWER**

CAUTION When drawing the boundary line in Step 1, be careful to draw a solid line if the inequality includes equality (\leq , \geq) or a dashed line if equality is not included ($<$, $>$).

If an inequality is written in the form $y > mx + b$ or $y < mx + b$, then the inequality symbol indicates which region to shade.

If $y > mx + b$, then shade **above** the boundary line.

If $y < mx + b$, then shade **below** the boundary line.

This method works only if the inequality is solved for y.

NOW TRY EXERCISE 2
Graph $3x - 2y < 0$.

EXAMPLE 2 Graphing a Linear Inequality with Boundary Passing through the Origin

Graph $3x - 4y > 0$.

Graph the boundary line. The x - and y -intercepts are the same point, $(0, 0)$, so this line passes through the origin. Two other points on the line are $(4, 3)$ and $(-4, -3)$. The points of the boundary line do *not* belong to the inequality $3x - 4y > 0$ (because the inequality symbol is $>$, *not* \geq). For this reason, the line is dashed.

To use the method explained above, we solve the inequality for y .

$3x - 4y > 0$ Original inequality

$-4y > -3x$ Subtract $3x$.

Use this equivalent inequality to decide which region to shade.

$y < \frac{3}{4}x$ Divide by -4 .
Change $>$ to $<$.

Because the *is less than* symbol occurs *when the original inequality is solved for y*, shade the region *below* the boundary line. See **FIGURE 43**.

CHECK Choose a test point not on the line, which rules out the origin. We choose $(2, -1)$.

$3x - 4y > 0$ Original inequality

$3(2) - 4(-1) \stackrel{?}{>} 0$ Let $x = 2$ and $y = -1$.

$6 + 4 \stackrel{?}{>} 0$ Multiply.

$10 > 0$ ✓ True




FIGURE 43

This result agrees with the decision to shade below the line. The solution set, graphed in **FIGURE 43**, includes only those points in the shaded region (and *not* those on the line).

NOW TRY

NOW TRY EXERCISE 3
Graph $x + 2 > 0$.

EXAMPLE 3 Graphing a Linear Inequality

Graph $x - 3 < 1$.

We graph $x - 3 = 1$, which is equivalent to $x = 4$, as a dashed vertical line passing through the point $(4, 0)$. To determine which region to shade, we choose $(0, 0)$ as a test point.

$x - 3 < 1$ Original inequality

$0 - 3 \stackrel{?}{<} 1$ Let $x = 0$.

$-3 < 1$ True





FIGURE 44

Because a true statement results, we shade the region containing $(0, 0)$. See **FIGURE 44**.

NOW TRY

NOW TRY ANSWERS



OBJECTIVE 2 Graph the intersection of two linear inequalities.

A pair of inequalities joined with the word *and* is interpreted as the *intersection* of the solution sets of the inequalities.

The graph of the intersection of two or more inequalities is the region of the plane where all points satisfy all of the inequalities at the same time.


NOW TRY EXERCISE 4

Graph $x + y < 3$ and $y \leq 2$.

EXAMPLE 4 Graphing the Intersection of Two Inequalities

Graph $2x + 4y \geq 5$ and $x \geq 1$.

To begin, we graph the solid boundary line $2x + 4y = 5$ through the intercepts $(\frac{5}{2}, 0)$ and $(0, \frac{5}{4})$. Using $(0, 0)$ as a test point in $2x + 4y > 5$ produces a false statement, so we shade the region that does not include $(0, 0)$. See **FIGURE 45(a)**.

For the inequality $x \geq 1$, we graph the solid boundary line $x = 1$. Again, using $(0, 0)$ as a test point produces a false statement, so we shade the region to the right of the boundary line. See **FIGURE 45(b)**.

We use purple shading to identify the intersection of the graphs in **FIGURE 45(c)**.





FIGURE 45

In practice, the graphs in **FIGURES 45(a)** and **(b)** are graphed on the same axes.

CHECK Using **FIGURE 45(c)**, we can choose a test point from each of the four regions formed by the intersection of the boundary lines.




Verify that only ordered pairs in the purple shaded region satisfy *both* inequalities. Ordered pairs in the other regions satisfy only one of the inequalities or neither of them. ✓

NOW TRY 

OBJECTIVE 3 Graph the union of two linear inequalities.

When two inequalities are joined by the word *or*, we must find the *union* of the graphs of the inequalities.

The graph of the union of two inequalities includes all of the points that satisfy either inequality.

NOW TRY ANSWER


NOW TRY EXERCISE 5

Graph $3x - 5y < 15$ or $x > 4$.

EXAMPLE 5 Graphing the Union of Two Inequalities

Graph $2x + 4y \geq 5$ or $x \geq 1$.

The graphs of the two inequalities are shown in FIGURES 45(a) and (b) in Example 4 on the preceding page. The graph of the union includes all points in *either* inequality, as shown in FIGURE 46.




FIGURE 46


CHECK Using FIGURE 46, we can choose a test point from each of the four regions formed by the intersection of the boundary lines.

$(2, 1)$, $(0, 2)$, $(0, 0)$, and $(2, -2)$ Possible test points

Verify that ordered pairs from the shaded region satisfy *either* inequality—that is, they satisfy one or the other or both. (Note also that $(0, 0)$ satisfies *neither* inequality.) ✓

NOW TRY

NOW TRY ANSWER



2.4 Exercises

FOR EXTRA HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Study cards are a great way to learn vocabulary, procedures, and so on.

Review Study Skill 5, Using Study Cards.

Concept Check Determine whether each ordered pair is a solution of the given inequality.

- | | | | | | |
|--------------------|---------------|---------------|----------------|---------------|---------------|
| 1. $x - 2y \leq 4$ | (a) $(0, 0)$ | (b) $(2, -1)$ | 2. $x + y > 0$ | (a) $(0, 0)$ | (b) $(-2, 1)$ |
| | (c) $(7, 1)$ | (d) $(0, 2)$ | | (c) $(2, -1)$ | (d) $(-4, 6)$ |
| 3. $x - 5 > 0$ | (a) $(0, 0)$ | (b) $(5, 0)$ | 4. $y \leq 1$ | (a) $(0, 0)$ | (b) $(3, 1)$ |
| | (c) $(-1, 3)$ | (d) $(6, 2)$ | | (c) $(2, -1)$ | (d) $(-3, 3)$ |

Concept Check In each statement, fill in the first blank with either solid or dashed. Fill in the second blank with either above or below.

- The boundary of the graph of $y \leq -x + 2$ will be a _____ line, and the shading will be _____ the line.
- The boundary of the graph of $y < -x + 2$ will be a _____ line, and the shading will be _____ the line.
- The boundary of the graph of $y > -x + 2$ will be a _____ line, and the shading will be _____ the line.
- The boundary of the graph of $y \geq -x + 2$ will be a _____ line, and the shading will be _____ the line.





Concept Check Refer to the given graph, and complete each statement with the correct inequality symbol $<$, \leq , $>$, or \geq .

9. x _____ 4

10. y _____ -3

11. y _____ $3x - 2$

12. y _____ $-x + 3$



Graph each inequality. See Examples 1–3.

13. $x + y \leq 2$

14. $x + y \leq -3$

15. $4x - y < 4$

16. $3x - y < 3$

17. $x + 3y \geq -2$

18. $x + 4y \geq -3$

19. $y < \frac{1}{2}x + 3$

20. $y < \frac{1}{3}x - 2$

21. $y \geq -\frac{2}{5}x + 2$

22. $y \geq -\frac{3}{2}x + 3$

23. $2x + 3y \geq 6$

24. $3x + 4y \geq 12$

25. $5x - 3y > 15$

26. $4x - 5y > 20$

27. $x + y > 0$

28. $x + 2y > 0$

29. $x - 3y \leq 0$

30. $x - 5y \leq 0$

31. $y < x$

32. $y \leq 4x$

33. $x + 3 \geq 0$

34. $x - 1 \leq 0$

35. $y + 5 < 2$

36. $y - 1 > 3$

Extending Skills Complete each of the following to write an inequality for the graph shown.

37. Determine the following for the boundary line.

Slope: _____


y-intercept: _____

Equation: $y =$ _____

The boundary line here is (solid / dashed), and the region (above / below) it is shaded.

The inequality symbol to indicate this is ($<$ / \leq / $>$ / \geq).

Inequality for the graph: y _____



38. Determine the following for the boundary line.

Slope: _____


y-intercept: _____

Equation: $y =$ _____

The boundary line here is (solid / dashed), and the region (above / below) it is shaded.

The inequality symbol to indicate this is ($<$ / \leq / $>$ / \geq).

Inequality for the graph: y _____



Graph the intersection of each pair of inequalities. See Example 4.

39. $x + y \leq 1$ and $x \geq 1$ 40. $x - y \geq 2$ and $x \geq 3$
 41. $2x - y \geq 2$ and $y < 4$ 42. $3x - y \geq 3$ and $y < 3$
 43. $x + y > -5$ and $y < -2$ 44. $6x - 4y < 10$ and $y > 2$

Extending Skills Use the definition of absolute value to write each inequality as a compound inequality, and graph its solution set in the rectangular coordinate plane.

45. $|x| < 3$ 46. $|y| < 5$ 47. $|x + 1| < 2$ 48. $|y - 3| < 2$

Graph the union of each pair of inequalities. See Example 5.

49. $x - y \geq 1$ or $y \geq 2$ 50. $x + y \leq 2$ or $y \geq 3$
 51. $x - 2 > y$ or $x < 1$ 52. $x + 3 < y$ or $x > 3$
 53. $3x + 2y < 6$ or $x - 2y > 2$ 54. $x - y \geq 1$ or $x + y \leq 4$


RELATING CONCEPTS For Individual or Group Work (Exercises 55–60)

Linear programming is a method for finding the optimal (best possible) solution that meets all the conditions for a problem such as the following.

A factory can have no more than 200 workers on a shift, but must have at least 100 and must manufacture at least 3000 units at minimum cost. How many workers should be on a shift in order to produce the required units at minimal cost?

Let x represent the number of workers and y represent the number of units manufactured. **Work Exercises 55–60 in order.**

55. Write three inequalities expressing the problem conditions.
 56. Graph the inequalities from **Exercise 55** using the axes below, and shade the intersection.



57. The cost per worker is \$50 per day and the cost to manufacture 1 unit is \$100. Write an equation in x , y , and C representing the total daily cost C .
 58. Find values of x and y for several points in or on the boundary of the shaded region. Include any “corner points,” where C is maximized or minimized.
 59. Of the values of x and y found in **Exercise 58**, which ones give the least value when substituted in the cost equation from **Exercise 57**?
 60. What does the answer in **Exercise 59** mean in terms of the given problem?

2.5 Introduction to Relations and Functions

OBJECTIVES

- 1 Define and identify relations and functions.
- 2 Find the domain and range.
- 3 Identify functions defined by graphs and equations.

OBJECTIVE 1 Define and identify relations and functions.

Consider the relationship illustrated in the following table between number of hours worked and paycheck amount for an hourly worker.

Number of Hours Worked	Paycheck Amount (in dollars)	Ordered pairs
5	50	→ (5, 50)
10	100	→ (10, 100)
20	200	→ (20, 200)
40	400	→ (40, 400)

The data from the table can be represented by a set of ordered pairs.

$$\{(5, 50), (10, 100), (20, 200), (40, 400)\}$$

↑ Number of hours worked ↑ Paycheck amount in dollars

VOCABULARY

- relation
- function
- dependent variable
- independent variable
- domain
- range

Each first component of the ordered pairs represents a number of hours worked, and each second component represents the corresponding paycheck amount. Such a set of ordered pairs is a *relation*.

Relation

A **relation** is any set of ordered pairs.

NOW TRY EXERCISE 1

Write the relation as a set of ordered pairs.

Year	Average Gas Price per Gallon (in dollars)
2010	2.79
2012	3.64
2014	3.37
2016	2.14

Data from Energy Information Administration.

EXAMPLE 1 Writing Ordered Pairs for a Relation

Write the relation as a set of ordered pairs.

Number of Gallons of Gas	Cost (in dollars)
0	0
1	3.50
2	7.00
3	10.50
4	14.00

The data in the table defines a relation between number of gallons of gas and cost and can be written as the following set of ordered pairs.

$$\{(0, 0), (1, 3.50), (2, 7.00), (3, 10.50), (4, 14.00)\}$$

↑ Number of gallons of gas ↑ Cost in dollars

NOW TRY ANSWER

1. $\{(2010, 2.79), (2012, 3.64), (2014, 3.37), (2016, 2.14)\}$

A *function* is a special kind of relation.

Function

A **function** is a relation in which, for each distinct value of the first component of the ordered pairs, there is *exactly one* value of the second component.


NOW TRY EXERCISE 2

Determine whether each relation defines a function.

- (a) $\{(1, 5), (3, 5), (5, 5)\}$
 (b) $\{(-1, -3), (0, 2), (-1, 6)\}$

EXAMPLE 2 Determining Whether Relations Are Functions

Determine whether each relation defines a function.

(a) $F = \{(1, 2), (-2, 4), (3, -1)\}$

Look at the ordered pairs that define this relation. For each distinct x -value, there is *exactly one* y -value. We can show this correspondence as follows.

$$\begin{array}{ccc} \{1, & -2, & 3\} & \text{x-values of } F \\ \downarrow & \downarrow & \downarrow & \\ \{2, & 4, & -1\} & \text{y-values of } F \end{array}$$

Therefore, relation F is a function.

(b) $G = \{(-2, -1), (-1, 0), (0, 1), (1, 2), (2, 2)\}$

Relation G is also a function. Although the last two ordered pairs have the same y -value (1 is paired with 2, and 2 is paired with 2), this does not violate the definition of a function.

$$\begin{array}{cccc} \{-2, & -1, & 0, & 1, & 2\} & \text{x-values of } G \\ \downarrow & \downarrow & \downarrow & \downarrow & \swarrow \\ \{-1, & 0, & 1, & 2\} & \text{y-values of } G \end{array}$$

The first components (x -values) are distinct, and each is paired with only one second component (y -value).

(c) $H = \{(-4, 1), (-2, 1), (-2, 0)\}$

In relation H , the last two ordered pairs have the *same* x -value paired with *two different* y -values (-2 is paired with both 1 and 0). H is a relation, but *not* a function.

$$\begin{array}{ccc} \{-4, & -2\} & \text{x-values of } H \\ \downarrow & \swarrow & \downarrow \\ \{1, & 0\} & \text{y-values of } H \end{array}$$

In a function, no two ordered pairs have the same first component and different second components.

$$\begin{array}{c} \text{Different y-values} \\ \downarrow \quad \downarrow \\ \text{Relation } H = \{(-4, 1), (-2, 1), (-2, 0)\} \quad \text{Not a function} \\ \uparrow \quad \uparrow \\ \text{Same x-value} \end{array}$$

NOW TRY 

Relations may be defined in several different ways.

- A relation may be defined as a set of ordered pairs.

Relation $F = \{(1, 2), (-2, 4), (3, -1)\}$ Function (Example 2(a))

Relation $H = \{(-4, 1), (-2, 1), (-2, 0)\}$ Not a function (Example 2(c))

NOW TRY ANSWERS

2. (a) function
 (b) not a function

- A relation may be defined as a correspondence or *mapping*.

See FIGURE 47. In the mapping for relation F , the arrow from 1 to 2 indicates that the ordered pair $(1, 2)$ belongs to F . Also, -2 is mapped to 4, and 3 is mapped to -1 . Thus, F is a function—each first component is paired with exactly one second component.

In the mapping for relation H , which is *not* a function, the first component -2 is paired with two different second components, 1 and 0.




FIGURE 47

- A relation may be defined as a table.
- A relation may be defined as a graph.

FIGURE 48 includes a table and graph for relation F , which is a function.




FIGURE 48

- A relation may be defined as a rule.

The rule may be given in words, such as “ y is twice x .” Usually the rule is an equation, such as

$$y = 2x.$$

The infinite number of ordered-pair solutions (x, y) can be represented by the graph in FIGURE 49.

In the equation $y = 2x$, the value of y *depends* on the value of x . Thus, the variable y is the **dependent variable**. The variable x is the **independent variable**.


Graph of the relation $y = 2x$

FIGURE 49

$$\text{Dependent variable} \rightarrow y = 2x \leftarrow \text{Independent variable}$$

An equation tells how to determine the value of the dependent variable for a specific value of the independent variable.


NOTE An equation that describes the relationship given at the beginning of this section between number of hours worked and paycheck amount is

$$y = 10x. \quad 10 \text{ represents the hourly rate, } \$10.$$

Paycheck amount y *depends* on number of hours worked x . Thus, *paycheck amount* is the dependent variable, and *number of hours worked* is the independent variable.

In a function, there is exactly one value of the dependent variable, the second component, for each value of the independent variable, the first component.

NOTE Another way to think of a function relationship is to visualize the independent variable as an **input** and the dependent variable as an **output**. This **input-output (function) machine** illustrates the relationship between number of hours worked and paycheck amount.



OBJECTIVE 2 Find the domain and range.

Domain and Range

For every relation defined by a set of ordered pairs (x, y) , there are two important sets of elements.

- The set of all values of the independent variable (x) is the **domain**.
- The set of all values of the dependent variable (y) is the **range**.

NOW TRY EXERCISE 3

Give the domain and range of each relation. Determine whether the relation defines a function.

- (a) $\{(2, 2), (2, 5), (4, 8)\}$
 (b) The table from **Objective 1**


Number of Hours Worked	Paycheck Amount (in dollars)
5	50
10	100
20	200
40	400

EXAMPLE 3 Finding Domains and Ranges of Relations

Give the domain and range of each relation. Determine whether the relation defines a function.

- (a) $\{(3, -1), (4, 2), (4, 5), (6, 8)\}$
- List 4 only once.
- Domain: $\{3, 4, 6\}$ Set of x -values
- Range: $\{-1, 2, 5, 8\}$ Set of y -values

This relation is not a function because the same x -value 4 is paired with two different y -values, 2 and 5.



This mapping represents the following set of ordered pairs.

$$\{(95, A), (89, B), (88, B), (78, C)\}$$

Domain: $\{95, 89, 88, 78\}$ Set of first components

Range: $\{A, B, C\}$ Set of second components

The mapping defines a function—each domain value corresponds to exactly one range value.

- (c)
- | x | y |
|-----|-----|
| -5 | 2 |
| 0 | 2 |
| 5 | 2 |
- This table represents the following set of ordered pairs.
- $$\{(-5, 2), (0, 2), (5, 2)\}$$
- Domain: $\{-5, 0, 5\}$ Set of x -values
- Range: $\{2\}$ Set of y -values

The table defines a function—each distinct x -value corresponds to exactly one y -value (even though it is the same y -value).

NOW TRY ANSWERS


3. (a) domain: $\{2, 4\}$;
 range: $\{2, 5, 8\}$;
 not a function
 (b) domain: $\{5, 10, 20, 40\}$;
 range: $\{50, 100, 200, 400\}$;
 function

A graph gives a “picture” of a relation and can be used to determine its domain and range.

NOTE Pay particular attention to the use of color to interpret domain and range in **Example 4**—blue for domain and red for range.


NOW TRY
EXERCISE 4

Give the domain and range of the relation.



EXAMPLE 4 Finding Domains and Ranges from Graphs

Give the domain and range of each relation.




This relation includes the five ordered pairs that are graphed.

$$\{(-1, 1), (0, -1), (1, 2), (4, -3), (5, 2)\}$$

Domain: $\{-1, 0, 1, 4, 5\}$ Set of x -values

Range: $\{1, -1, 2, -3\}$ Set of y -values

List 2 only once.




The x -values of the ordered pairs that form the graph include all numbers between -4 and 4 , inclusive, as shown in blue. The y -values include all numbers between -6 and 6 , inclusive, as shown in red.

Domain: $[-4, 4]$


Range: $[-6, 6]$

Use interval notation.



The arrowheads on the graphed line indicate that the line extends indefinitely left and right, as well as up and down. Therefore, both the domain (set of x -values), shown in blue, and the range (set of y -values), shown in red, include all real numbers.

Domain: $(-\infty, \infty)$ Range: $(-\infty, \infty)$



The graphed curve extends indefinitely left and right, as well as upward. The domain, shown in blue, includes all real numbers. Because there is a least y -value, -3 , the range, shown in red, includes all numbers greater than or equal to -3 .

Domain: $(-\infty, \infty)$ Range: $[-3, \infty)$

NOW TRY

OBJECTIVE 3 Identify functions defined by graphs and equations.

Because each value of x in a function corresponds to only one value of y , any vertical line drawn through the graph of a function must intersect the graph in at most one point. This is the *vertical line test* for a function.

FIGURE 50 on the next page illustrates this test with the graphs of two relations.

NOW TRY ANSWER

4. domain: $(-\infty, \infty)$;
range: $[-2, \infty)$





FIGURE 50

Vertical Line Test

If every vertical line intersects the graph of a relation in no more than one point, then the relation represents a function.


NOW TRY EXERCISE 5

Use the vertical line test to determine whether the relation is a function.



EXAMPLE 5 Using the Vertical Line Test

Use the vertical line test to determine whether each relation graphed in **Example 4** is a function. (We repeat the graphs here.)



The graphs in (a), (c), and (d) satisfy the vertical line test because every vertical line intersects each graph no more than once. These graphs represent functions.

The graph in (b) fails the vertical line test because a vertical line intersects the graph more than once—that is, the same x -value corresponds to two different y -values. This is not the graph of a function.

NOW TRY

NOTE Graphs that do not represent functions are still relations. *All equations and graphs represent relations, and all relations have a domain and range.*

If a relation is defined by an equation involving a fraction or a radical, apply these guidelines when finding its domain.

1. Exclude from the domain any values that make the denominator of a fraction equal to 0.

Example: The function $y = \frac{1}{x}$ has all real numbers *except* 0 as its domain because division by 0 is undefined.

2. Exclude from the domain any values that result in an even root of a negative number.

Example: The function $y = \sqrt{x}$ has all *nonnegative* real numbers as its domain because the square root of a negative number is not real.

NOW TRY ANSWER

5. not a function

Agreement on Domain

Unless specified otherwise, the domain of a relation is assumed to be all real numbers that produce real numbers when substituted for the independent variable.

EXAMPLE 6 Identifying Functions and Domains from Equations

Determine whether each relation defines y as a function of x . Give the domain.

(a) $y = x + 4$

In this equation, y is found by adding 4 to x . Thus, each value of x corresponds to just one value of y , and the relation defines a function. Because x can be any real number, the domain is $(-\infty, \infty)$. The graph in **FIGURE 51** confirms this reasoning.




FIGURE 51




FIGURE 52

(b) $y = \sqrt{2x - 1}$

For any choice of x in the domain, there is exactly one corresponding value for y . (The radical is a single nonnegative number.) This equation defines a function. The quantity under the radical symbol cannot be negative—that is, $2x - 1$ must be greater than or equal to 0.

$$2x - 1 \geq 0$$

$$2x \geq 1 \quad \text{Add 1.}$$

$$x \geq \frac{1}{2} \quad \text{Divide by 2.}$$

The domain of the function is $[\frac{1}{2}, \infty)$. The graph in **FIGURE 52** confirms our work.

(c) $y^2 = x$

The ordered pairs $(4, 2)$ and $(4, -2)$ both satisfy this equation. One value of x , 4, corresponds to two values of y , 2 and -2 , so this equation does not define a function. Because x is equal to the square of y , the values of x must always be nonnegative. The domain of the relation is $[0, \infty)$. See **FIGURE 53**.




FIGURE 53




FIGURE 54

(d) $y \leq x - 1$

By definition, y is a function of x if every value of x leads to exactly one value of y . Here, a particular value of x , such as 1, corresponds to many values of y . The ordered pairs

$$(1, 0), (1, -1), (1, -2), (1, -3), \quad \text{and so on}$$

all satisfy the inequality. This relation does not define a function. Any number can be used for x , so the domain is the set of all real numbers, $(-\infty, \infty)$. See **FIGURE 54**.

NOW TRY
EXERCISE 6

Determine whether each relation defines y as a function of x . Give the domain.

(a) $y = 4x - 3$

(b) $y = \sqrt{2x - 4}$

(c) $y = \frac{1}{x - 2}$

(d) $y < 3x + 1$

(e) $y = \frac{5}{x - 1}$

Given any value of x in the domain, we find y by subtracting 1 and then dividing the result into 5. This process produces exactly one value of y for each value in the domain, so the given equation defines a function.

The domain includes all real numbers except those which make the denominator 0.

$$x - 1 = 0 \quad \text{Set the denominator equal to 0.}$$

$$x = 1 \quad \text{Add 1.}$$

The domain includes all real numbers *except* 1, written $(-\infty, 1) \cup (1, \infty)$. In **FIGURE 55**, the open circle on the graph indicates that 1 is excluded from the domain.




FIGURE 55

NOW TRY

In summary, we give three variations of the definition of a function.

Variations of the Definition of a Function

1. A **function** is a relation in which, for each distinct value of the first component of the ordered pairs, there is exactly one value of the second component.
2. A **function** is a set of distinct ordered pairs in which no first component is repeated.
3. A **function** is a correspondence (mapping) or an equation (rule) that assigns exactly one range value to each distinct domain value.

NOW TRY ANSWERS

6. (a) function; $(-\infty, \infty)$
 (b) function; $[2, \infty)$
 (c) function; $(-\infty, 2) \cup (2, \infty)$
 (d) not a function; $(-\infty, \infty)$

2.5 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Complete each statement. Choices may be used more than once.

function independent variable vertical line test relation
 domain ordered pairs dependent variable range

1. A(n) _____ is any set of _____ $\{(x, y)\}$.
2. A(n) _____ is a relation in which, for each distinct value of the first component of the _____, there is exactly one value of the second component.
3. In a relation $\{(x, y)\}$, the _____ is the set of x -values, and the _____ is the set of y -values.
4. The relation $\{(0, -2), (2, -1), (2, -4), (5, 3)\}$ (*does / does not*) define a function. The set $\{0, 2, 5\}$ is its _____, and the set $\{-2, -1, -4, 3\}$ is its _____.
5. Consider the function $d = 50t$, where d represents distance and t represents time. The value of d depends on the value of t , so the variable t is the _____, and the variable d is the _____.
6. The _____ is used to determine whether a graph is that of a function. It says that any vertical line can intersect the graph of a(n) _____ in no more than (*zero / one / two*) point(s).

Write each relation as a set of ordered pairs. See Example 1.

7.

x	y
2	-2
2	0
2	1

8.

x	y
-1	-1
0	-1
1	-1

9.



Year	Average Movie Ticket Price (in dollars)
1960	0.76
1980	2.69
2000	5.39
2016	8.65

Data from Motion Picture Association of America.

10.

Year	Average ACT Composite Score
2010	21.0
2012	21.1
2014	21.0
2016	20.8

Data from ACT.





Concept Check Express each relation using a different form. (For example, if the given form is a set of ordered pairs, use a graph.) There is more than one correct way to do this. See Objective 1.

13. $\{(0, 2), (2, 4), (4, 6)\}$

14. y is half of x .

15.

x	y
-1	-3
0	-1
1	1
3	3



18. **Concept Check** Which of the relations represented in Exercises 13–17 does *not* define a function? Explain.

Determine whether each relation defines a function, and give the domain and range. See Examples 2–5.

19. $\{(5, 1), (3, 2), (4, 9), (7, 6)\}$

20. $\{(8, 0), (5, 4), (9, 3), (3, 8)\}$

21. $\{(2, 4), (0, 2), (2, 5)\}$

22. $\{(9, -2), (-3, 5), (9, 2)\}$

23. $\{(-3, 1), (4, 1), (-2, 7)\}$

24. $\{(-12, 5), (-10, 3), (8, 3)\}$

25. $\{(1, 1), (1, -1), (0, 0), (2, 4), (2, -4)\}$

26. $\{(2, 5), (3, 7), (4, 9), (5, 11)\}$

27.

x	y
1	5
1	2
1	-1
1	-4

28.



















x	y
-4	-4
-4	0
-4	4
-4	8

29.

x	y
4	-3
2	-3
0	-3
-2	-3

30.

x	y
-3	-6
-1	-6
1	-6
3	-6



Determine whether each relation defines y as a function of x . (Solve for y first if necessary.) Give the domain. See Example 6.

49. $y = -6x$

50. $y = -9x$

51. $y = 2x - 6$

52. $y = 6x + 8$

53. $y = x^2$

54. $y = x^3$

55. $x = y^6$

56. $x = y^4$

57. $x + y < 4$

58. $x - y < 3$

59. $y = \sqrt{x}$

60. $y = -\sqrt{x}$

61. $y = \sqrt{x - 3}$

62. $y = \sqrt{x - 7}$

63. $y = \sqrt{4x + 2}$

64. $y = \sqrt{2x + 9}$

65. $y = \frac{x + 4}{5}$

66. $y = \frac{x - 3}{2}$

67. $y = -\frac{2}{x}$

68. $y = -\frac{6}{x}$

69. $y = \frac{2}{x - 4}$

70. $y = \frac{7}{x - 2}$

71. $xy = 1$

72. $xy = 3$

Solve each problem.

73. The table shows the percentage of students at 4-year colleges who graduated within 5 years.

Year	Percentage
2013	52.8
2014	52.6
2015	52.6
2016	53.2
2017	53.7

Data from ACT.

- Does the table define a function?
- What are the domain and range?
- What is the range element that corresponds to 2015? The domain element that corresponds to 53.7?
- Call this function f . Give two ordered pairs that belong to f .

74. The table shows the percentage of persons age 12 or older who smoked cigarettes.

Year	Percentage
1985	38.7
2000	24.9
2005	24.9
2010	23.0
2015	19.4

Data from National Survey on Drug Use and Health.

- Does the table define a function?
- What are the domain and range?
- What is the range element that corresponds to 2015? The domain element that corresponds to 23.0?
- Call this function g . Give two ordered pairs that belong to g .

2.6 Function Notation and Linear Functions

OBJECTIVES

- 1 Use function notation.
- 2 Graph linear and constant functions.

OBJECTIVE 1 Use function notation.

When a function f is defined with a rule or an equation using x and y for the independent and dependent variables, we say, “ y is a function of x ” to emphasize that y depends on x . We use the notation

$$y = f(x),$$

The parentheses here do *not* indicate multiplication.

called **function notation**, to express this and read $f(x)$ as “ **f of x ,**” or “ **f at x .**” The letter f is a name for this particular function. For example, if $y = 3x - 5$, we can name this function f and write the following.

$$y = \underset{\substack{\text{Name of the function} \\ \text{Value of the function}}}{f(x)} = \overset{\text{Defining expression}}{3x - 5}$$

Value of the independent variable

VOCABULARY

- linear function
- constant function

$f(x)$ is just another name for the dependent variable y .

We evaluate a function at different values of x by substituting x -values from the domain into the function.

NOW TRY EXERCISE 1

Let $f(x) = 4x + 3$. Find the value of function f for each value of x .

- (a) $x = -2$ (b) $x = 0$

EXAMPLE 1 Evaluating a Function

Let $f(x) = 3x - 5$. Find the value of function f for each value of x .

- (a) $x = 2$

$$f(x) = 3x - 5$$

Read $f(2)$ as “ f of 2” or “ f at 2.”

$$f(2) = 3 \cdot 2 - 5 \quad \text{Replace } x \text{ with } 2.$$

$$f(2) = 6 - 5 \quad \text{Multiply.}$$

$$f(2) = 1 \quad \text{Subtract.}$$

For $x = 2$, the corresponding function value (or y -value) is **1**. $f(2) = 1$ symbolizes the statement

“If $x = 2$ in the function f , then $y = 1$ ”

and is represented by the ordered pair $(2, 1)$.

- (b) $x = -1$

$$f(x) = 3x - 5$$

Use parentheses to avoid errors.

$$f(-1) = 3(-1) - 5 \quad \text{Replace } x \text{ with } -1.$$

$$f(-1) = -3 - 5 \quad \text{Multiply.}$$

$$f(-1) = -8 \quad \text{Subtract.}$$

Thus, $f(-1) = -8$ and the ordered pair $(-1, -8)$ belongs to f .

NOW TRY ANSWERS

1. (a) -5 (b) 3

CAUTION The symbol $f(x)$ does *not* indicate “ f times x ,” but represents the y -value associated with the indicated x -value. As shown in **Example 1(a)**, $f(2)$ is the y -value that corresponds to the x -value 2 in f .

NOTE In the function $f(x) = 3x - 5$ in **Example 1**, $f(2) = 1$ and $f(-1) = -8$ correspond to the ordered pairs $(2, 1)$ and $(-1, -8)$. Because the domain of f is $(-\infty, \infty)$ —that is, x can be any real number—this function defines an infinite set of ordered pairs whose graph is a line with slope 3 and y -intercept $(0, -5)$. See **FIGURE 56**. This makes sense because $f(x)$ is another name for y , and

$$f(x) = 3x - 5 \text{ is equivalent to } y = 3x - 5.$$

f is a linear function.




FIGURE 56

NOW TRY EXERCISE 2

Let $f(x) = 2x^2 - 4x + 1$. Find the following.

- (a) $f(-2)$ (b) $f(a)$

EXAMPLE 2 Evaluating a Function

Let $f(x) = -x^2 + 5x - 3$. Find the following.

- (a) $f(4)$

Do *not* read this as “ f times 4.” Read it as “ f of 4,” or “ f at 4.”

$$f(x) = -x^2 + 5x - 3$$

The base in $-x^2$ is x , *not* $(-x)$.

$$f(4) = -4^2 + 5 \cdot 4 - 3$$

Replace x with 4.

$$f(4) = -16 + 20 - 3$$

Apply the exponent. Multiply.

$$f(4) = 1$$

Add and subtract.

Thus, $f(4) = 1$, and the ordered pair $(4, 1)$ belongs to f .

- (b) $f(q)$

$$f(x) = -x^2 + 5x - 3$$

$$f(q) = -q^2 + 5q - 3$$

Replace x with q .

The replacement of one variable with another is important in later courses.

NOW TRY

Sometimes letters other than f , such as g , h , or capital letters F , G , and H are used to name functions.

NOW TRY EXERCISE 3

Let $g(x) = 8x - 5$. Find and simplify $g(a - 2)$.

EXAMPLE 3 Evaluating a Function

Let $g(x) = 2x + 3$. Find and simplify $g(a + 1)$.

$$g(x) = 2x + 3$$

$$g(a + 1) = 2(a + 1) + 3$$

Replace x with $a + 1$.

$$g(a + 1) = 2a + 2 + 3$$

Distributive property

$$g(a + 1) = 2a + 5$$

Add.

NOW TRY ANSWERS

2. (a) 17 (b) $2a^2 - 4a + 1$
3. $8a - 21$

NOW TRY

NOW TRY
EXERCISE 4

For each function, find $f(-1)$.

(a) $f = \{(-5, -1), (-3, 2), (-1, 4)\}$

(b) $f(x) = x^2 - 12$

EXAMPLE 4 Evaluating Functions

For each function, find $f(3)$.

(a) $f(x) = 3x - 7$

$f(3) = 3(3) - 7$ Replace x with 3.

$f(3) = 9 - 7$ Multiply.

$f(3) = 2$ Subtract.


(b)

x	$y = f(x)$
6	-12
3	-6
0	0
-3	6

← Here, $f(3) = -6$.

(c) $f = \{(-3, 5), (0, 3), (3, 1), (6, -1)\}$

We want $f(3)$, the y -value of the ordered pair whose first component is $x = 3$. As indicated by the ordered pair $(3, 1)$, for $x = 3$, $y = 1$. Thus, $f(3) = 1$.



The domain element 3 is paired with 5 in the range, so

$$f(3) = 5.$$

NOW TRY




FIGURE 57

EXAMPLE 5 Finding Function Values from a Graph

Refer to the function graphed in FIGURE 57.

(a) Find $f(3)$.

Locate 3 on the x -axis. See FIGURE 58. Moving up to the graph of f and over to the y -axis gives 4 for the corresponding y -value. Thus, $f(3) = 4$, which corresponds to the ordered pair $(3, 4)$.

(b) Find $f(0)$.

Refer to FIGURE 58 to see that $f(0) = 1$.




FIGURE 58




FIGURE 59

(c) For what value of x is $f(x) = 5$?

Because $f(x) = y$, we want the value of x that corresponds to $y = 5$. See FIGURE 59, and locate 5 on the y -axis. Moving across to the graph of f and down to the x -axis gives $x = 4$. Thus, $f(4) = 5$, which corresponds to the ordered pair $(4, 5)$.

NOW TRY

If a function f is defined by an equation in x and y instead of function notation, use the following steps to find $f(x)$.

Writing an Equation Using Function Notation

Step 1 Solve the equation for y if it is not given in that form.

Step 2 Replace y with $f(x)$.

NOW TRY ANSWERS

4. (a) 4 (b) -11
5. (a) 0 (b) 1


NOW TRY EXERCISE 6

Write the equation using function notation $f(x)$. Then find $f(-3)$.

$$-4x^2 + y = 5$$

EXAMPLE 6 Writing Equations Using Function Notation

Write each equation using function notation $f(x)$. Then find $f(-2)$.

(a) $y = x^2 + 1$ ← This equation is already solved for y . (Step 1)

$$f(x) = x^2 + 1 \quad \text{Replace } y \text{ with } f(x). \quad \text{(Step 2)}$$

To find $f(-2)$, let $x = -2$.

$$f(x) = x^2 + 1$$

$$f(-2) = (-2)^2 + 1 \quad \text{Let } x = -2.$$

$$f(-2) = 4 + 1 \quad (-2)^2 = (-2)(-2)$$

$$f(-2) = 5 \quad \text{Add.}$$

(b) $x - 4y = 5$

Step 1 $-4y = -x + 5$ Subtract x .

$$y = \frac{1}{4}x - \frac{5}{4} \quad \text{Divide by } -4.$$

Step 2 $f(x) = \frac{1}{4}x - \frac{5}{4}$ Replace y with $f(x)$.

Now find $f(-2)$.

$$f(-2) = \frac{1}{4}(-2) - \frac{5}{4} = -\frac{7}{4} \quad \text{Let } x = -2.$$

NOW TRY 

OBJECTIVE 2 Graph linear and constant functions.

Linear equations (except for vertical lines with equations of the form $x = a$) define *linear functions*.

Linear Function

A function f that can be written in the form

$$f(x) = ax + b,$$

where a and b are real numbers, is a **linear function**. The value of a is the slope m of the graph of the function. The domain of a linear function is $(-\infty, \infty)$, unless specified otherwise.

Examples:

$$f(x) = 2x + 4, \quad f(x) = -5x, \quad f(x) = -\frac{1}{2}x - \frac{5}{4}, \quad f(x) = 0.75x + 10$$

A linear function whose graph is a horizontal line has the form

$$f(x) = b \quad \text{Constant function}$$

and is a **constant function**. The range of any nonconstant linear function is $(-\infty, \infty)$, but the range of a constant function $f(x) = b$ is $\{b\}$.

NOW TRY ANSWER

6. $f(x) = 4x^2 + 5$;
 $f(-3) = 41$


NOW TRY EXERCISE 7

Graph each function. Give the domain and range.


(a) $f(x) = \frac{1}{3}x - 2$

(b) $f(x) = -3$

NOW TRY ANSWERS



domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$



domain: $(-\infty, \infty)$;
range: $\{-3\}$

EXAMPLE 7 Graphing Linear and Constant Functions

Graph each function. Give the domain and range.

(a) $f(x) = \frac{1}{4}x - \frac{5}{4}$ (from Example 6(b))
Slope \uparrow $\frac{1}{4}$ \uparrow y-intercept is $(0, -\frac{5}{4})$.

To graph this function, plot the y-intercept $(0, -\frac{5}{4})$. Use the geometric definition of slope as $\frac{\text{rise}}{\text{run}}$ to find a second point on the line. The slope is $\frac{1}{4}$, so we move up 1 unit from $(0, -\frac{5}{4})$ and to the right 4 units to the point $(4, -\frac{1}{4})$. Draw the straight line through these points. See FIGURE 60. The domain and range are both $(-\infty, \infty)$.




FIGURE 60




FIGURE 61

(b) $f(x) = 4$

The graph of this constant function is the horizontal line containing all points with y-coordinate 4. See FIGURE 61. The domain is $(-\infty, \infty)$. The value of $f(x)$ —that is, y —is 4 for every value of x , so the range is $\{4\}$.

NOW TRY

2.6 Exercises

FOR EXTRA HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Time management can be a challenge for students.

Review Study Skill 6, Managing Your Time.

Concept Check Work each problem.

- To emphasize that “ y is a function of x ” for a given function f , we use function notation and write $y = \underline{\hspace{2cm}}$. Here, f is the name of the $\underline{\hspace{2cm}}$, x is a value from the $\underline{\hspace{2cm}}$, and $f(x)$ is the function value (or y -value) that corresponds to $\underline{\hspace{2cm}}$. We read $f(x)$ as “ $\underline{\hspace{2cm}}$.”
- Choose the correct response.
For a function f , the notation $f(3)$ means $\underline{\hspace{2cm}}$.
 - the variable f times 3, or $3f$.
 - the value of the dependent variable when the independent variable is 3.
 - the value of the independent variable when the dependent variable is 3.
 - f equals 3.
- Fill in each blank with the correct response.

The equation $2x + y = 4$ has a straight $\underline{\hspace{2cm}}$ as its graph. One point that lies on the graph is $(3, \underline{\hspace{1cm}})$. If we solve the equation for y and use function notation, we have a(n) $\underline{\hspace{1cm}}$ function $f(x) = \underline{\hspace{2cm}}$. For this function, $f(3) = \underline{\hspace{1cm}}$, meaning that the point $(\underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ lies on the graph of the function.

4. Which of the following defines y as a linear function of x ?



A. $y = \frac{1}{4}x - \frac{5}{4}$ B. $y = \frac{1}{x}$ C. $y = x^2$ D. $y = \sqrt{x}$

Let $f(x) = -3x + 4$ and $g(x) = -x^2 + 4x + 1$. Find the following. See Examples 1–3.

5. $f(0)$ 6. $g(0)$ 7. $f(-3)$ 8. $f(-5)$
9. $g(-2)$ 10. $g(-1)$ 11. $g(3)$ 12. $g(10)$
13. $f(100)$ 14. $f(-100)$ 15. $f\left(\frac{1}{3}\right)$ 16. $f\left(\frac{7}{3}\right)$
17. $g(0.5)$ 18. $g(1.5)$ 19. $f(p)$ 20. $g(k)$
21. $f(-x)$ 22. $g(-x)$ 23. $f(x + 2)$ 24. $f(x - 2)$
25. $f(2t + 1)$ 26. $f(3t - 2)$ 27. $g(\pi)$ 28. $g(t)$
29. $f(x + h)$ 30. $f(a + b)$ 31. $g\left(\frac{p}{3}\right)$ 32. $g\left(\frac{1}{x}\right)$

For each function, find (a) $f(2)$ and (b) $f(-1)$. See Examples 4, 5(a), and 5(b).

33. $f = \{(-2, 2), (-1, -1), (2, -1)\}$ 34. $f = \{(-1, -5), (0, 5), (2, -5)\}$
35. $f = \{(-1, 3), (4, 7), (0, 6), (2, 2)\}$ 36. $f = \{(2, 5), (3, 9), (-1, 11), (5, 3)\}$







39.

x	$y = f(x)$
2	4
1	1
0	0
-1	1
-2	4


40.

x	$y = f(x)$
8	6
5	3
2	0
-1	-3
-4	-6




Refer to the given graph. Find the value of x for each value of $f(x)$. See Example 5(c).

45. (a) $f(x) = 3$
 (b) $f(x) = -1$
 (c) $f(x) = -3$



46. (a) $f(x) = 4$
 (b) $f(x) = -2$
 (c) $f(x) = 0$



An equation that defines y as a function f of x is given. (a) Solve for y in terms of x , and write each equation using function notation $f(x)$. (b) Find $f(3)$. See Example 6.

47. $x + 3y = 12$ 48. $x - 4y = 8$ 49. $y + 2x^2 = 3$
 50. $y - 3x^2 = 2$ 51. $4x - 3y = 8$ 52. $-2x + 5y = 9$

Graph each linear or constant function. Give the domain and range. See Example 7.

53. $f(x) = -2x + 5$ 54. $g(x) = 4x - 1$ 55. $h(x) = \frac{1}{2}x + 2$
 56. $F(x) = -\frac{1}{4}x + 1$ 57. $f(x) = x$ 58. $f(x) = -x$
 59. $H(x) = -3x$ 60. $G(x) = 2x$ 61. $g(x) = -4$
 62. $f(x) = 5$ 63. $f(x) = 0$ 64. $f(x) = -2.5$

65. **Concept Check** What is the name that is usually given to the graph of $f(x) = 0$?
 66. **Concept Check** Can the graph of a linear function have an undefined slope? Explain.

Solve each problem.

67. A package weighing x pounds costs $f(x)$ dollars to mail to a given location, where

$$f(x) = 3.75x.$$


- (a) Evaluate $f(3)$.
 (b) Describe what 3 and the value $f(3)$ mean in part (a), using the terms *independent variable* and *dependent variable*.
 (c) How much would it cost to mail a 5-lb package? Express this situation using function notation.



68. A taxicab driver charges \$2.50 per mile.

- (a) Fill in the table with the correct response for the price $f(x)$ the driver charges for a trip of x miles.
 (b) The linear function that gives a rule for the amount charged is $f(x) =$ _____.
 (c) Graph this function for the domain $\{0, 1, 2, 3\}$ using the set of axes at the right.

x	$f(x)$
0	
1	
2	
3	



69. To print t-shirts, there is a \$100 set-up fee, plus a \$12 charge per t-shirt. Let x represent the number of t-shirts printed and $f(x)$ represent the total charge.
- Write a linear function that models this situation.
 - Find $f(125)$. Interpret the answer in the context of this problem.
 - Find the value of x if $f(x) = 1000$. Express this situation using function notation, and interpret it in the context of this problem.

70. Rental on a car is \$150, plus \$0.50 per mile. Let x represent the number of miles driven and $f(x)$ represent the total cost to rent the car.
- Write a linear function that models this situation.
 - Find $f(250)$. Interpret the answer in the context of this problem.
 - Find the value of x if $f(x) = 400$. Express this situation using function notation, and interpret it in the context of this problem.

71. The table represents a linear function.

x	$y = f(x)$
0	3.5
1	2.3
2	1.1
3	-0.1
4	-1.3


- What is $f(2)$?
- If $f(x) = -1.3$, what is the value of x ?
- What is the slope of the line?
- What is the y -intercept of the line?
- Using the answers from parts (c) and (d), write an equation for $f(x)$.

72. The table represents a linear function.

x	$y = f(x)$
-1	-3.9
0	-2.4
1	-0.9
2	0.6
3	2.1


- What is $f(2)$?
- If $f(x) = 2.1$, what is the value of x ?
- What is the slope of the line?
- What is the y -intercept of the line?
- Using the answers from parts (c) and (d), write an equation for $f(x)$.

73. The graph shows water in a swimming pool over time.



- What numbers are possible values of the independent variable? The dependent variable?
- For how long is the water level increasing? Decreasing?
- How many gallons of water are in the pool after 90 hr?
- Call this function f . What is $f(0)$? What does it mean?
- What is $f(25)$? What does it mean?

74. The graph shows electricity use on a summer day.



- Is this the graph of a function? Why or why not?
- What is the domain?
- Estimate the number of megawatts used at 8 A.M.
- At what time was the most electricity used? The least electricity?
- Call this function f . What is $f(12)$? What does it mean?

75. Forensic scientists use the lengths of certain bones to calculate the height of a person. Two such bones are the tibia (t), the bone from the ankle to the knee, and the femur (r), the bone from the knee to the hip socket. A person's height (h) in centimeters is determined from the lengths of these bones using the following functions.

$$\text{For men: } h(r) = 69.09 + 2.24r \quad \text{or} \quad h(t) = 81.69 + 2.39t$$


$$\text{For women: } h(r) = 61.41 + 2.32r \quad \text{or} \quad h(t) = 72.57 + 2.53t$$

- (a) Find the height of a man with a femur measuring 56 cm.
 (b) Find the height of a man with a tibia measuring 40 cm.
 (c) Find the height of a woman with a femur measuring 50 cm.
 (d) Find the height of a woman with a tibia measuring 36 cm.
76. Based on federal regulations, a pool to house sea otters must have a volume that is “the square of the sea otter’s average adult length (in meters) multiplied by 3.14 and by 0.91 meter.” If x represents the sea otter’s average adult length and $f(x)$ represents the volume (in cubic meters) of the corresponding pool size, this formula can be written as the function

$$f(x) = 0.91(3.14)x^2.$$

Find the volume of the pool for each adult sea otter length (in meters). Round answers to the nearest hundredth.


- (a) 0.8 (b) 1.0 (c) 1.2 (d) 1.5



RELATING CONCEPTS For Individual or Group Work (Exercises 77–85)

Refer to the straight-line graph and work Exercises 77–85 in order.

77. By just looking at the graph, how can we tell whether the slope is *positive*, *negative*, *0*, or *undefined*?
78. Apply the slope formula to find the slope of the line.
79. What is the slope of any line parallel to the line shown? Perpendicular to the line shown?
80. Find the x -intercept of the graph.
81. Find the y -intercept of the graph.
82. Use function notation to write the equation of the graphed line. Use f to designate the function.
83. Find $f(8)$.
84. If $f(x) = -8$, what is the value of x ?
85. From the graph, find $f(1)$. Then confirm this value using the equation written in Exercise 82.



Chapter 2

Summary

STUDY SKILLS REMINDER

How do you best prepare for a test? **Review Study Skill 7, *Reviewing a Chapter*.**

Key Terms

2.1

ordered pair
origin
 x -axis
 y -axis
rectangular (Cartesian)
coordinate system
components
plot
coordinate

quadrant
table of ordered pairs
(table of values)
graph of an equation
first-degree equation
linear equation in two
variables
 x -intercept
 y -intercept

2.2

rise
run
slope

2.3

scatter diagram

2.4

linear inequality in two
variables
boundary line

2.5

relation
function
dependent variable
independent variable
domain
range

2.6

linear function
constant function

New Symbols

(x, y) ordered pair

(x_1, y_1) subscript notation
(read “ x -sub-one,
 y -sub-one”)

Δ Greek letter delta
 m slope

$f(x)$ function notation;
function of x (read
“ f of x ” or “ f at x ”)

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- An **ordered pair** is a pair of numbers written
 - in numerical order within brackets
 - within parentheses or brackets
 - within parentheses in which the order of the numbers is important
 - within parentheses in which order does not matter.
- A **linear equation in two variables** is an equation that can be written in the form
 - $Ax + By < C$
 - $ax + b = 0$
 - $y = x^2$
 - $Ax + By = C$.
- An **intercept** is
 - the point where the x -axis and y -axis intersect
 - a pair of numbers written between parentheses in which order matters
 - one of the regions determined by a coordinate system
 - the point where a graph intersects the x -axis or the y -axis.
- The **slope** of a line is
 - the measure of the run over the rise of the line
 - the distance between two points on the line
 - the ratio of the change in y to the change in x along the line
 - the horizontal change compared to the vertical change of two points on the line.
- A **relation** is
 - a set of ordered pairs
 - the ratio of the change in y to the change in x along a line
 - the set of all possible values of the independent variable
 - all the second components of a set of ordered pairs.
- A **function** is
 - the numbers in an ordered pair
 - a set of ordered pairs in which each x -value corresponds to exactly one y -value
 - a pair of numbers written within parentheses
 - the set of all ordered pairs that satisfy an equation.
- The **domain** of a relation is
 - the set of all possible values of the dependent variable y
 - a set of ordered pairs
 - the difference of the x -values
 - the set of all possible values of the independent variable x .
- The **range** of a relation is
 - the set of all possible values of the dependent variable y
 - a set of ordered pairs
 - the difference of the y -values
 - the set of all possible values of the independent variable x .

ANSWERS

1. C; *Examples:* (0, 3), (3, 8), (4, 0) 2. D; *Examples:* $3x + 2y = 6$, $x = y - 7$ 3. D; *Example:* In the graph of the equation $2x + 3y = 6$, the x -intercept is (3, 0) and the y -intercept is (0, 2). 4. C; *Example:* The line passing through the points (3, 6) and (5, 4) has slope $\frac{4-6}{5-3} = \frac{-2}{2} = -1$. 5. A; *Example:* The set $\{(2, 0), (4, 3), (6, 6)\}$ defines a relation. 6. B; *Example:* The relation given in Answer 5 is a function. 7. D; *Example:* In the relation in Answer 5, the domain is the set of x -values, $\{2, 4, 6\}$. 8. A; *Example:* In the relation in Answer 5, the range is the set of y -values, $\{0, 3, 6\}$.

Quick Review

CONCEPTS

2.1 Linear Equations in Two Variables

Finding Intercepts

To find the x -intercept, let $y = 0$ and solve for x .

To find the y -intercept, let $x = 0$ and solve for y .

Midpoint Formula

The midpoint M of a line segment PQ with endpoints $P(x_1, y_1)$ and $Q(x_2, y_2)$ is found as follows.

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

2.2 The Slope of a Line

The slope m of the line passing through the points (x_1, y_1) and (x_2, y_2) is defined as follows.

$$\text{slope } m = \frac{\text{rise}}{\text{run}} = \frac{\text{change in } y}{\text{change in } x} = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$

(where $x_1 \neq x_2$)

A horizontal line has slope 0.

A vertical line has undefined slope.

Parallel lines have the same slope.

Perpendicular lines, neither of which is vertical, **have slopes that are negative reciprocals** (that is, have a product of -1).

Slope gives the **average rate of change** in y per unit change in x , where the value of y depends on the value of x .

EXAMPLES

Find the intercepts of the graph of $2x + 3y = 12$.

$$\begin{array}{l|l} 2x + 3(0) = 12 & \text{Let } y = 0. \\ 2x = 12 & 2(0) + 3y = 12 \\ x = 6 & 3y = 12 \\ & y = 4 \end{array}$$

The x -intercept is (6, 0).

The y -intercept is (0, 4).

Find the midpoint of the segment with endpoints (4, -7) and (-10 , -13).

$$M = \left(\frac{4 + (-10)}{2}, \frac{-7 + (-13)}{2} \right) = (-3, -10)$$

Find the slope of the graph of $2x + 3y = 12$.

Use the intercepts (6, 0) and (0, 4) and the slope formula.

$$m = \frac{4 - 0}{0 - 6} = \frac{4}{-6} = -\frac{2}{3} \quad \begin{array}{l} (x_1, y_1) = (6, 0) \\ (x_2, y_2) = (0, 4) \end{array}$$

The graph of the horizontal line $y = -5$ has slope $m = 0$.

The graph of the vertical line $x = 3$ has undefined slope.

The lines $y = 2x + 3$ and $4x - 2y = 6$ are **parallel**.

Both have slope 2.

$$\begin{array}{l|l} y = 2x + 3 & 4x - 2y = 6 \leftarrow \text{Solve for } y. \\ & -2y = -4x + 6 \\ & y = 2x - 3 \end{array}$$

The lines $y = 3x - 1$ and $x + 3y = 4$ are **perpendicular**.

Their slopes are negative reciprocals because $3\left(-\frac{1}{3}\right) = -1$.

$$\begin{array}{l|l} y = 3x - 1 & x + 3y = 4 \leftarrow \text{Solve for } y. \\ & 3y = -x + 4 \\ & y = -\frac{1}{3}x + \frac{4}{3} \end{array}$$

The weight of a young child increased from 30 lb to 60 lb between the ages of 3 and 8. What was the child's average change in weight per year over these years?

$$\frac{60 - 30}{8 - 3} = \frac{30}{5} = 6 \text{ lb per yr}$$

CONCEPTS

2.3 Writing Equations of Lines

Slope-Intercept Form

$$y = mx + b$$

Point-Slope Form

$$y - y_1 = m(x - x_1)$$

Standard Form

$Ax + By = C$, where A , B , and C are real numbers and A and B are not both 0. (We give A , B , and C as integers, with greatest common factor 1, and $A \geq 0$.)

Horizontal Line

$$y = b$$

Vertical Line

$$x = a$$

2.4 Linear Inequalities in Two Variables

Graphing a Linear Inequality

Step 1 Draw the graph of the straight line that is the boundary.

- Make the line solid if the inequality involves \leq or \geq .
- Make the line dashed if the inequality involves $<$ or $>$.

Step 2 Choose any point not on the line as a test point. Substitute the coordinates of that point in the inequality.

Step 3 Shade the region that includes the test point if it satisfies the original inequality. Otherwise, shade the region on the other side of the boundary line.

2.5 Introduction to Relations and Functions

A **relation** is any set of ordered pairs. A **function** is a set of ordered pairs in which, for each distinct value of the first component, there is *exactly one* value of the second component.

The set of first components (x -values) is the **domain**.

The set of second components (y -values) is the **range**.

2.6 Function Notation and Linear Functions

To evaluate a function f , where $f(x)$ defines the range value for a given value of x in the domain, substitute the domain value wherever x appears.

To write an equation that defines a function f in function notation, follow these steps.

Step 1 Solve the equation for y if it is not given in that form.

Step 2 Replace y with $f(x)$.

EXAMPLES

$$y = 2x + 3 \quad m = 2; y\text{-intercept is } (0, 3).$$

$$y - 3 = 4(x - 5) \quad m = 4; \text{ The point } (5, 3) \text{ is on the line.}$$

$$2x - 5y = 8 \quad \text{Standard form}$$

$$y = 4 \quad \text{Horizontal line}$$

$$x = -1 \quad \text{Vertical line}$$

Graph $2x - 3y \leq 6$.


Draw the graph of $2x - 3y = 6$ using the intercepts $(3, 0)$ and $(0, -2)$. Draw a solid line because of the inclusion of equality in the symbol \leq .

Choose $(0, 0)$ as a test point.

$$2(0) - 3(0) \stackrel{?}{\leq} 6$$

$$0 < 6 \quad \text{True}$$

Shade the region that includes $(0, 0)$.



The set of ordered pairs $\{(-1, 4), (0, 6), (1, 4)\}$ defines a function.

$$\text{Domain: } \{-1, 0, 1\} \quad \text{Set of } x\text{-values}$$

$$\text{Range: } \{4, 6\} \quad \text{Set of } y\text{-values}$$

The equation $y = x^2$ defines a function.

$$\text{Domain: } (-\infty, \infty) \quad \text{Range: } [0, \infty)$$

Let $f(x) = x^2 - 7x + 12$. Find $f(1)$.

$$f(x) = x^2 - 7x + 12$$

$$f(1) = 1^2 - 7(1) + 12 \quad \text{Let } x = 1.$$

$$f(1) = 6$$

Write $2x + 3y = 12$ using function notation $f(x)$.

$$2x + 3y = 12$$

$$3y = -2x + 12 \quad \text{Subtract } 2x.$$

$$y = -\frac{2}{3}x + 4 \quad \text{Divide by } 3.$$

$$f(x) = -\frac{2}{3}x + 4 \quad \text{Replace } y \text{ with } f(x).$$

Chapter 2

Review Exercises

2.1 Complete the given table of ordered pairs for each equation, and then graph the equation.

1. $3x + 2y = 10$

x	y
0	
	0
2	
	-2

2. $x - y = 8$

x	y
2	
	-3
3	
	-2

Find the x - and y -intercepts. Then graph each equation.

3. $4x - 3y = 12$

4. $5x + 7y = 28$

5. $2x + 5y = 20$

6. $x - 4y = 8$

Find the midpoint of each segment with the given endpoints.

7. $(-8, -12)$ and $(8, 16)$

8. $(0, -5)$ and $(-9, 8)$

2.2 Find the slope of each line. (In Exercises 17 and 18, coordinates of the points shown are integers.)

9. Through $(-1, 2)$ and $(4, -5)$

10. Through $(0, 3)$ and $(-2, 4)$

11. Through $(-1, 5)$ and $(-1, -4)$

12. $y = 2x + 3$


13. $3x - 4y = 5$

14. $y = 4$


15. Perpendicular to $3x - y = 4$

16. Parallel to $3y = 2x + 5$

17.



18.




19.


x	y
-2	6
-1	3
0	0
1	-3

20. Determine whether the slope of the line is *positive*, *negative*, *0*, or *undefined*.


(a)




(b)



(c)



(d)




Determine whether each pair of lines is parallel, perpendicular, or neither.

21. $3x - y = 6$ and $x + 3y = 12$

22. $3x - y = 4$ and $6x + 12 = 2y$

Solve each problem.

23. If the pitch of a roof is $\frac{1}{4}$, how many feet in the horizontal direction correspond to a rise of 3 ft?



24. In 1980, the median family income in the United States was about \$21,000 per year. In 2016, it was about \$59,039 per year. Find the average rate of change in median family income per year to the nearest dollar during this period. (Data from U.S. Census Bureau.)

2.3 Write an equation of the line passing through the given point(s) and satisfying any given condition. Give the equation (a) in slope-intercept form and (b) in standard form.

25. (0, -8); slope $\frac{3}{5}$ 26. (0, 5); slope $-\frac{1}{3}$
 27. (2, -5) and (1, 4) 28. (-3, -1) and (2, 6)
 29. (6, -2); parallel to $4x - y = 3$ 30. (0, 1); perpendicular to $2x - 5y = 7$

Write an equation of the line passing through the given point and satisfying the given condition.

31. (0, 12); slope 0 32. (2, 7); undefined slope
 33. (0.3, 0.6); vertical 34. (-1, 4); horizontal


Solve each problem.

35. A lease on a 2018 Chevrolet Sonic LT hatchback costs \$2829 at signing, plus \$229 per month. Let x represent the number of months since the signing and y represent the amount paid (in dollars). (Data from www.carsdirect.com)

- (a) Write an equation in slope-intercept form that represents this situation.
 (b) How much will be paid over the lease's full term of 39 months?

36. Average annual charges for tuition, fees, room, and board (in dollars) at private nonprofit 4-year universities in the United States are shown in the graph.

Costs at 4-Year Universities



Data from The College Board.




- (a) Use the information given for the years 2013 and 2017, letting $x = 13$ represent 2013 and $x = 17$ represent 2017, and letting y represent cost (in dollars), to write an equation that models the data. Write the equation in slope-intercept form. Interpret the slope.
 (b) Use the equation from part (a) to approximate costs at private 4-year universities in 2015.

2.4 Graph each inequality or compound inequality.

37. $3x - 2y \leq 12$ 38. $5x - y > 6$
 39. $2x + y \leq 1$ and $x \geq 2y$ 40. $x \geq 2$ or $y \geq 2$

2.5 Determine whether each relation defines a function, and give the domain and range.

41. $\{(-4, 2), (-4, -2), (1, 5), (1, -5)\}$



2.5, 2.6 Determine whether each relation defines y as a function of x . Give the domain. Identify any linear functions.

45. $y = 3x - 3$

46. $y < x + 2$

47. $y = |x|$

48. $y = \sqrt{4x + 7}$

49. $x = y^2$

50. $y = \frac{7}{x - 6}$

2.6 Let $f(x) = -2x^2 + 3x - 6$. Find the following.

51. $f(0)$

52. $f(3)$

53. $f\left(-\frac{1}{2}\right)$

54. $f(k)$

Solve each problem.

55. The equation $2x^2 - y = 0$ defines y as a function f of x . Write it using function notation $f(x)$, and find $f(3)$.

56. The linear equation $2x - 5y = 7$ defines y as a function of x . If $y = f(x)$, which of the following defines the same function?

A. $f(x) = -\frac{2}{5}x + \frac{7}{5}$

B. $f(x) = -\frac{2}{5}x - \frac{7}{5}$

C. $f(x) = \frac{2}{5}x - \frac{7}{5}$

D. $f(x) = \frac{2}{5}x + \frac{7}{5}$

57. Describe the graph of a constant function.

58. The table shows life expectancy at birth in the United States for selected years.

(a) Does the table define a function?

(b) What are the domain and range?

(c) Call this function f . Give two ordered pairs that belong to f .

(d) Find $f(1980)$. What does this mean?

(e) If $f(x) = 76.8$, what does x equal?

Year	Life Expectancy at Birth (years)
1960	69.7
1970	70.8
1980	73.7
1990	75.4
2000	76.8
2010	78.7
2015	78.8

Data from National Center for Health Statistics.

Chapter 2

Mixed Review Exercises

Determine whether each pair of lines is parallel, perpendicular, or neither.


1. $4x + 3y = 8$ and $6y = 7 - 8x$

2. $3x + y = 4$ and $3y = x - 6$

The graph shows per capita consumption of potatoes (in pounds) in the United States from 2008 to 2016.

3. Use the given ordered pairs to find and interpret the average rate of change in per capita potato consumption per year to the nearest tenth during this period.

4. Write an equation in slope-intercept form that models per capita consumption of potatoes y (in pounds) in year x , where $x = 0$ represents 2008.




Data from U.S. Department of Agriculture.

Write an equation of a line (in the form specified, if given) passing through the given point(s) and satisfying any given conditions.

5. $(0, 0)$; $m = 3$ (slope-intercept form)
6. $(0, 3)$ and $(-2, 4)$ (standard form)
7. $(2, -3)$; perpendicular to $x = 2$

Solve each problem.

8. Which equations have a graph with just one intercept?
 - A. $x - 6 = 0$
 - B. $x + y = 0$
 - C. $x - y = 4$
 - D. $y = -4$
9. Which inequality has as its graph a dashed boundary line and shading below the line?
 - A. $y \geq 4x + 3$
 - B. $y > 4x + 3$
 - C. $y \leq 4x + 3$
 - D. $y < 4x + 3$
10. Refer to the graph of function f .



- (a) Find $f(-2)$.
- (b) Find $f(0)$.
- (c) For what value of x is $f(x) = -3$?
- (d) Give the domain and range.

STUDY SKILLS REMINDER

Using test-taking strategies can help you improve your test scores. **Review Study Skill 8, Taking Math Tests.**

Chapter 2

Test

FOR EXTRA HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

View the complete solutions to all Chapter Test exercises in MyLab Math.

1. Complete the table of ordered pairs for the equation $2x - 3y = 12$.


x	y
1	
3	
	-4

Find the x - and y -intercepts. Then graph each equation.

2. $4x - 3y = -12$
3. $x = 2$
4. $y = -2x$
5. Find the slope of the line passing through the points $(6, 4)$ and $(-4, -1)$.
6. Describe how the graph of a line with undefined slope is situated in a rectangular coordinate system.

For each line, find the slope and the x - and y -intercepts.

7. $3x - 2y = 13$
8. $y = 5$
9. Which line has positive slope and negative y -coordinate for its y -intercept?



Determine whether each pair of lines is parallel, perpendicular, or neither.

10. $5x - y = 8$ and $5y = -x + 3$
11. $2y = 3x + 12$ and $3y = 2x - 5$

12. In 2000, there were 94,000 farms in Iowa. As of 2016, there were 87,000. Find and interpret the average rate of change in the number of farms per year, to the nearest whole number. (Data from U.S. Department of Agriculture.)

Write an equation of the line passing through the given point(s) and satisfying any given conditions. Give the equation (a) in slope-intercept form and (b) in standard form.

13. $(0, 3)$; slope $-\frac{2}{5}$ 14. $(-2, 3)$ and $(6, -1)$ 15. $(4, -1)$; $m = -5$





Write an equation of the line passing through the given point and satisfying the given condition.

16. $(-3, 14)$; horizontal 17. $(5, -6)$; vertical 18. $(0, 0)$; horizontal
19. Write an equation in slope-intercept form of the line passing through the point $(-7, 2)$ and satisfying the given condition.
- (a) parallel to $3x + 5y = 6$ (b) perpendicular to $y = 2x$
20. A ticket to a rock concert costs \$142.75. An advance parking pass costs \$45. Let x represent the number of tickets purchased and y represent the total cost in dollars.
- (a) Write an equation in slope-intercept form that represents this situation.
- (b) How much does it cost for 6 tickets and a parking pass?

Graph each inequality or compound inequality.

21. $3x - 2y > 6$ 22. $y < 2x - 1$ and $x - y < 3$

23. Which of the following is the graph of a function? Give its domain and range.



24. Which of the following does *not* define a function? Give its domain and range.

A. $\{(0, 1), (-2, 3), (4, 8)\}$

B. $y = 2x - 6$

C. $y = \sqrt{x + 2}$

D.

x	y
0	1
3	2
0	2
6	3

25. For $f(x) = -x^2 + 2x - 1$, find $f(1)$ and $f(a)$.

26. Graph the linear function $f(x) = \frac{2}{3}x - 1$. Give the domain and range.

Chapters R-2 Cumulative Review Exercises

1. Add or subtract as indicated.

(a) $\frac{1}{6} + \frac{3}{4}$

(b) $\frac{11}{12} - \frac{1}{4}$

(c) $7.5 - 2.75$

2. Multiply or divide as indicated.

(a) $6 \div \frac{3}{4}$

(b) 1.5×100

(c) $1.25 \div 10$

3. Complete the table of fraction, decimal, and percent equivalents.

	Fraction in Lowest Terms (or Whole Number)	Decimal	Percent
(a)	$\frac{1}{100}$		
(b)		0.5	
(c)			75%
(d)		1.0	

Determine whether each statement is always true, sometimes true, or never true. If the statement is sometimes true, give examples for which it is true and for which it is false.

- The absolute value of a negative number equals the additive inverse of the number.
- The sum of two negative numbers is positive.
- The sum of a positive number and a negative number is 0.

Perform each operation.

7. $-|-2| - 4 + |-3|$

8. $(-0.8)^2$

9. -3^2

10. $\sqrt{-64}$

Evaluate each expression for $p = -4$, $q = \frac{1}{2}$, and $r = 16$.

11. $-3(2q - 3p)$

12. $\frac{\sqrt{r}}{8p + 2r}$

Simplify each expression.

13. $-(-4x + 3)$

14. $\sqrt{25} - 5(-1)^2$


Solve each equation.

15. $2z - 5 + 3z = 4 - (z + 2)$

16. $\frac{3x - 1}{5} + \frac{x + 2}{2} = -\frac{3}{10}$

Solve each problem.

17. If each side of a square were increased by 4 in., the perimeter would be 8 in. less than twice the perimeter of the original square. Find the length of a side of the original square.



18. Two planes leave the Dallas–Fort Worth airport at the same time. One travels east at 550 mph, and the other travels west at 500 mph. Assuming no wind, how long will it take for the planes to be 2100 mi apart?



Solve.


19. $-4 < 3 - 2k < 9$ 20. $-0.3x + 2.1(x - 4) \leq -6.6$
21. $\frac{1}{2}x > 3$ and $\frac{1}{3}x < \frac{8}{3}$ 22. $-5x + 1 \geq 11$ or $3x + 5 > 26$
23. $|2k - 7| + 4 = 11$ 24. $|3x + 6| \geq 0$

Solve each problem.

25. Find the x - and y -intercepts of the line with equation $3x + 5y = 12$, and graph the line.
26. Consider the points $A(-2, 1)$ and $B(3, -5)$.
- (a) Find the slope of the line AB .
- (b) Find the slope of a line perpendicular to line AB .
27. Graph the inequality $-2x + y < -6$.

Write an equation of the line passing through the given point(s) and satisfying any given conditions. Give the equation (a) in slope-intercept form and (b) in standard form.

28. $(0, -1)$; slope $-\frac{3}{4}$ 29. $(4, -3)$ and $(1, 1)$
30. Give the domain and range of the relation. Does it define a function?



STUDY SKILLS REMINDER

We can learn from the mistakes we make. **Review Study Skill 9, Analyzing Your Test Results.**

SYSTEMS OF LINEAR EQUATIONS

Just as the *intersection* of condensation trails in the sky consists of a region common to both trails, a solution of a *system of linear equations* (represented graphically by lines) is an ordered pair common to both solution sets (both lines).

3.1 Systems of Linear Equations in Two Variables

3.2 Systems of Linear Equations in Three Variables

3.3 Applications of Systems of Linear Equations

3.1 Systems of Linear Equations in Two Variables

OBJECTIVES

- 1 Determine whether an ordered pair is a solution of a linear system.
- 2 Solve linear systems by graphing.
- 3 Solve linear systems (with two equations and two variables) by substitution.
- 4 Solve linear systems (with two equations and two variables) by elimination.
- 5 Solve special systems.

VOCABULARY


- system of equations
- system of linear equations (linear system)
- solution set of a linear system
- consistent system
- independent equations
- inconsistent system
- dependent equations

STUDY SKILLS REMINDER

Be sure to read and work through the section material before working the exercises.

Review Study Skill 2, Reading Your Math Text.

In recent years, the number of Americans who subscribe to Netflix has increased, while the number who subscribe to cable TV has decreased. These trends can be seen in the graph in **FIGURE 1**. The two straight-line graphs intersect at the time when Netflix and cable TV had the *same* number of subscribers.



Data from Netflix, Leichtman Research Group.

FIGURE 1

$$\begin{aligned} -5.61x + y &= 23.9 && \text{Linear system} \\ 0.77x + y &= 52.1 && \text{of equations} \end{aligned}$$

(Here, $x = 0$ represents 2012, $x = 1$ represents 2013, and so on. y represents millions of subscribers.)

As shown beside **FIGURE 1**, we can use a linear equation to model the graph of the number of Netflix subscribers (red graph) and another linear equation to model the graph of the number of cable TV subscribers (blue graph). Such a set of equations is a **system of equations**—in this case, a **system of linear equations**, or a **linear system**.

The point where the graphs in **FIGURE 1** intersect is a solution of each of the individual equations. It is also the solution of the linear system of equations.

OBJECTIVE 1 Determine whether an ordered pair is a solution of a linear system.

The **solution set of a linear system** of equations contains *all* ordered pairs that satisfy *all* the equations of the system *at the same time*.

EXAMPLE 1 Determining Whether an Ordered Pair Is a Solution

Determine whether the given ordered pair is a solution of the given system.

$$\begin{aligned} \text{(a)} \quad x + y &= 6 \\ 4x - y &= 14 \end{aligned} ; (4, 2)$$

Replace x with 4 and y with 2 in each equation of the system.

$$\begin{array}{l|l} \begin{aligned} x + y &= 6 \\ 4 + 2 &\stackrel{?}{=} 6 \\ 6 &= 6 \quad \checkmark \quad \text{True} \end{aligned} & \begin{aligned} 4x - y &= 14 \\ 4(4) - 2 &\stackrel{?}{=} 14 \\ 16 - 2 &\stackrel{?}{=} 14 \\ 14 &= 14 \quad \checkmark \quad \text{True} \end{aligned} \end{array}$$

Because $(4, 2)$ makes *both* equations true, $(4, 2)$ is a solution of the system.

NOW TRY EXERCISE 1

Determine whether the ordered pair $(2, 5)$ is a solution of the given system.

(a) $2x - y = -1$
 $3x + 2y = 16$

(b) $-x + 2y = 8$
 $3x - 2y = 0$

(b) $3x + 2y = 11$; $(-1, 7)$
 $x + 5y = 36$

$$3x + 2y = 11$$

$$3(-1) + 2(7) \stackrel{?}{=} 11$$

$$-3 + 14 \stackrel{?}{=} 11$$

$$11 = 11 \quad \checkmark \quad \text{True}$$

$$x + 5y = 36$$

$$-1 + 5(7) \stackrel{?}{=} 36$$

$$-1 + 35 \stackrel{?}{=} 36$$

$$34 = 36 \quad \text{False}$$

The ordered pair $(-1, 7)$ is not a solution of the system—it does not make *both* equations true.

NOW TRY

OBJECTIVE 2 Solve linear systems by graphing.

One way to find the solution set of a linear system of equations is to graph each equation and find the point where the graphs intersect.

EXAMPLE 2 Solving a System by Graphing

Solve the system of equations by graphing.

$$x + y = 5 \quad (1)$$

$$2x - y = 4 \quad (2)$$

To graph these linear equations, we plot several points for each line.

$$x + y = 5$$

$$2x - y = 4$$

The intercepts are a convenient choice.

x	y
0	5
5	0
2	3

x	y
0	-4
2	0
4	4

Find a third ordered pair as a check.

The graph in **FIGURE 2** suggests that the point of intersection is the ordered pair $(3, 2)$.




FIGURE 2

To confirm that $(3, 2)$ is a solution of *both* equations, we check by substituting 3 for x and 2 for y in each equation.

CHECK $x + y = 5 \quad (1)$

$$3 + 2 \stackrel{?}{=} 5$$

$$5 = 5 \quad \checkmark \quad \text{True}$$

$2x - y = 4 \quad (2)$

$$2(3) - 2 \stackrel{?}{=} 4$$

$$6 - 2 \stackrel{?}{=} 4$$

$$4 = 4 \quad \checkmark \quad \text{True}$$

Because $(3, 2)$ makes both equations true, the solution set of the system is $\{(3, 2)\}$.

NOW TRY

NOW TRY EXERCISE 2


Solve the system of equations by graphing.

$$3x - 2y = 6$$

$$x - y = 1$$

NOW TRY ANSWERS

- (a) solution (b) not a solution
- $\{(4, 3)\}$



Three Cases for Solutions of Linear Systems in Two Variables

Case 1 The two graphs intersect in a single point.

The coordinates of this point give the only solution of the system. Because the system has a solution, it is **consistent**. The equations are *not* equivalent, so they are **independent**. See FIGURE 3(a).

Case 2 The graphs are parallel lines.

There is no solution common to both equations, so the solution set is \emptyset and the system is **inconsistent**. Because the equations are *not* equivalent, they are **independent**. See FIGURE 3(b).

Case 3 The graphs are the same line—that is, they coincide.

Because any solution of one equation of the system is a solution of the other, the solution set is an infinite set of ordered pairs representing the points on the line. This type of system is **consistent** because there is a solution. The equations are equivalent, so they are **dependent**. See FIGURE 3(c).




FIGURE 3

OBJECTIVE 3 Solve linear systems (with two equations and two variables) by substitution.

It can be difficult to read exact coordinates from a graph, especially if they are not integers, so we generally use algebraic methods to solve systems. One such method, the **substitution method**, is well suited for solving linear systems in which one equation is solved or can be easily solved for one variable in terms of the other.

EXAMPLE 3 Solving a System by Substitution (Case 1)

Solve the system.

$$2x - y = 6 \quad (1)$$

$$x = y + 2 \quad (2)$$

Because equation (2) is solved for x , we substitute $y + 2$ for x in equation (1).

$$2x - y = 6 \quad (1) \quad \text{Substitute in the other equation.}$$

$$2(y + 2) - y = 6 \quad \text{Let } x = y + 2.$$

$$2y + 4 - y = 6 \quad \text{Distributive property}$$

$$y + 4 = 6 \quad \text{Combine like terms.}$$

$$y = 2 \quad \text{Subtract 4.}$$

Be sure to use parentheses here.


NOW TRY EXERCISE 3

Solve the system.

$$\begin{aligned}x &= 3 + 2y \\ 4x - 3y &= 32\end{aligned}$$




FIGURE 4

We found y . Now we solve for x by substituting 2 for y in equation (2).

$$x = y + 2 \quad (2)$$

$$x = 2 + 2 \quad \text{Let } y = 2.$$

$$x = 4 \quad \text{Add.}$$

Write the x -value first
in the ordered pair.

Check the ordered pair $(4, 2)$ in both equations of the original system.

CHECK	$2x - y = 6 \quad (1)$		$x = y + 2 \quad (2)$
	$2(4) - 2 \stackrel{?}{=} 6$		$4 \stackrel{?}{=} 2 + 2$
	$8 - 2 \stackrel{?}{=} 6$		$4 = 4 \quad \checkmark \quad \text{True}$
	$6 = 6 \quad \checkmark \quad \text{True}$		

This check and the graph in FIGURE 4 show that the solution set is $\{(4, 2)\}$.
NOW TRY 
Solving a Linear System by Substitution

- Step 1** **Solve one of the equations for either variable.** If one equation has a variable term with coefficient 1 or -1 , choose it because the substitution method is usually easier.
- Step 2** **Substitute** for that variable in the other equation. The result should be an equation with just one variable.
- Step 3** **Solve** the equation from Step 2.
- Step 4** **Find the other value.** Substitute the result from Step 3 into the equation from Step 1 and solve for the value of the other variable.
- Step 5** **Check** the values in *both* of the *original* equations. Then write the solution set as a set containing an ordered pair.

EXAMPLE 4 Solving a System by Substitution (Case 1)

Solve the system.

$$3x + 2y = 13 \quad (1)$$

$$4x - y = -1 \quad (2)$$

Step 1 First solve one of the equations for either x or y . Because the coefficient of y in equation (2) is -1 , it is easiest to solve for y in equation (2).

$$4x - y = -1 \quad (2)$$

$$-y = -1 - 4x \quad \text{Subtract } 4x.$$

$$y = 1 + 4x \quad \text{Multiply by } -1.$$

Step 2 Substitute $1 + 4x$ for y in equation (1).

$$3x + 2y = 13 \quad (1)$$

$$3x + 2(1 + 4x) = 13 \quad \text{Let } y = 1 + 4x.$$

Step 3 Solve.

$$3x + 2 + 8x = 13 \quad \text{Distributive property}$$

$$11x = 11 \quad \text{Combine like terms. Subtract 2.}$$

$$x = 1 \quad \text{Divide by 11.}$$

Substitute in the
other equation.

NOW TRY ANSWER
3. $\{(11, 4)\}$


NOW TRY EXERCISE 4

Solve the system.

$$\begin{aligned} 5x + y &= 7 \\ 3x - 2y &= 25 \end{aligned}$$

Step 4 Now find y using the equation from Step 1.

$$y = 1 + 4x \quad \text{From Step 1}$$

$$y = 1 + 4(1) \quad \text{Let } x = 1.$$

$$y = 5 \quad \text{Multiply, and then add.}$$

Step 5 Check the ordered pair $(1, 5)$ in both equations (1) and (2).

CHECK	$3x + 2y = 13 \quad (1)$		$4x - y = -1 \quad (2)$
	$3(1) + 2(5) \stackrel{?}{=} 13$		$4(1) - 5 \stackrel{?}{=} -1$
	$3 + 10 \stackrel{?}{=} 13$		$4 - 5 \stackrel{?}{=} -1$
	$13 = 13 \quad \checkmark \quad \text{True}$		$-1 = -1 \quad \checkmark \quad \text{True}$

The solution set is $\{(1, 5)\}$.**NOW TRY** **EXAMPLE 5** Solving a System with Fractional Coefficients (Case 1)

Solve the system.

$$\frac{2}{3}x - \frac{1}{2}y = \frac{7}{6} \quad (1)$$

$$3x - y = 6 \quad (2)$$

This system will be easier to solve if we clear the fractions in equation (1).

$$6\left(\frac{2}{3}x - \frac{1}{2}y\right) = 6\left(\frac{7}{6}\right) \quad \text{Multiply (1) by the LCD, 6.}$$

Remember to multiply *each* term by 6.

$$6 \cdot \frac{2}{3}x - 6 \cdot \frac{1}{2}y = 6 \cdot \frac{7}{6} \quad \text{Distributive property}$$

$$4x - 3y = 7 \quad (3)$$

Now the system consists of equations (2) and (3).

This equation is equivalent to equation (1).

$$3x - y = 6 \quad (2)$$

$$4x - 3y = 7 \quad (3)$$

To use the substitution method, we solve equation (2) for y .

$$3x - y = 6 \quad (2)$$

$$-y = 6 - 3x \quad \text{Subtract } 3x.$$

$$y = 3x - 6 \quad \text{Multiply by } -1. \text{ Rewrite.}$$

Substitute $3x - 6$ for y in equation (3).

$$4x - 3y = 7 \quad (3)$$

$$4x - 3(3x - 6) = 7 \quad \text{Let } y = 3x - 6.$$

$$4x - 9x + 18 = 7 \quad \text{Distributive property}$$

Be careful with signs.

$$-5x + 18 = 7 \quad \text{Combine like terms.}$$

$$-5x = -11 \quad \text{Subtract 18.}$$

$$x = \frac{11}{5} \quad \text{Divide by } -5.$$

NOW TRY ANSWER4. $\{(3, -8)\}$


**NOW TRY
EXERCISE 5**

Solve the system.

$$\begin{aligned}\frac{1}{10}x + \frac{3}{5}y &= \frac{2}{5} \\ 2x + y &= 2\end{aligned}$$

Now find y using $y = 3x - 6$ (equation (2) solved for y).

$$y = 3\left(\frac{11}{5}\right) - 6 \quad \text{Let } x = \frac{11}{5}.$$

$$y = \frac{33}{5} - \frac{30}{5} \quad \text{Multiply; } 6 = \frac{30}{5}$$

$$y = \frac{3}{5} \quad \text{Subtract fractions.}$$

A check verifies that the solution set is $\left\{\left(\frac{11}{5}, \frac{3}{5}\right)\right\}$.
NOW TRY 

NOTE If an equation in a system contains decimal coefficients, first clear the decimals by multiplying by 10, 100, or 1000, depending on the number of decimal places. Then solve the system. For example, we multiply *each side* of the equation

$$0.5x + 0.75y = 3.25$$

by 10^2 , or 100, to obtain an equivalent equation

$$50x + 75y = 325.$$

OBJECTIVE 4 Solve linear systems (with two equations and two variables) by elimination.

The **elimination method** involves combining the two equations in a system so that one variable is eliminated. This is done using the following logic.

$$\text{If } a = b \text{ and } c = d, \text{ then } a + c = b + d.$$


**NOW TRY
EXERCISE 6**

Solve the system.

$$\begin{aligned}8x - 2y &= 18 \\ 5x + 2y &= -18\end{aligned}$$

EXAMPLE 6 Solving a System by Elimination (Case 1)

Solve the system.

$$2x + 3y = -6 \quad (1)$$

$$4x - 3y = 6 \quad (2)$$

Notice that adding the equations together will eliminate the variable y .

$$2x + 3y = -6 \quad (1)$$

$$4x - 3y = 6 \quad (2)$$

$$\text{Solve for } x \rightarrow 6x = 0 \quad \text{Add.}$$

$$x = 0 \quad \text{Divide by 6.}$$

To find y , substitute 0 for x in either equation (1) or equation (2).

$$2x + 3y = -6 \quad (1)$$

$$2(0) + 3y = -6 \quad \text{Let } x = 0.$$

$$0 + 3y = -6 \quad \text{Multiply.}$$

$$y = -2 \quad \text{Add, and then divide by 3.}$$

Check by substituting 0 for x and -2 for y in both equations of the original system to verify that $(0, -2)$ satisfies both equations. It does, so the solution set is $\{(0, -2)\}$.

NOW TRY 
NOW TRY ANSWERS

5. $\left\{\left(\frac{8}{11}, \frac{6}{11}\right)\right\}$

6. $\{(0, -9)\}$

From **Example 6**:

$$\begin{array}{rcl} 2x + 3y & = & -6 \quad (1) \\ 4x - 3y & = & 6 \quad (2) \\ \hline 6x & = & 0 \end{array}$$

By adding the equations in **Example 6** (repeated in the margin), we eliminated the variable y because the coefficients of the y -terms were opposites. In many cases the coefficients will *not* be opposites, and we must transform one or both equations so that the coefficients of one pair of variable terms are opposites.

Solving a Linear System by Elimination

Step 1 Write both equations in the form $Ax + By = C$.

Step 2 Transform the equations as needed so that the coefficients of one pair of variable terms are opposites. Multiply one or both equations by appropriate numbers so that the sum of the coefficients of either the x - or y -terms is 0.

Step 3 Add the new equations to eliminate a variable. The sum should be an equation with just one variable.

Step 4 Solve the equation from Step 3 for the remaining variable.

Step 5 Find the other value. Substitute the result from Step 4 into either of the original equations and solve for the other variable.

Step 6 Check the values in *both* of the *original* equations. Then write the solution set as a set containing an ordered pair.

EXAMPLE 7 Solving a System by Elimination (Case 1)

Solve the system.

$$5x - 2y = 4 \quad (1)$$

$$2x + 3y = 13 \quad (2)$$

Step 1 Both equations are in $Ax + By = C$ form.

Step 2 Suppose that we wish to eliminate the variable x . One way to do this is to multiply equation (1) by 2 and equation (2) by -5 .

The goal is to have opposite coefficients.

$$\begin{array}{rcl} 10x - 4y & = & 8 \quad \text{2 times each side of equation (1)} \\ -10x - 15y & = & -65 \quad \text{-5 times each side of equation (2)} \end{array}$$

Step 3 Now add.

$$\begin{array}{rcl} 10x - 4y & = & 8 \\ -10x - 15y & = & -65 \\ \hline -19y & = & -57 \quad \text{Add.} \end{array}$$

Step 4 Solve for y . $y = 3$ Divide by -19 .

Step 5 To find x , substitute 3 for y in either equation (1) or equation (2).

$$\begin{array}{rcl} 2x + 3y & = & 13 \quad (2) \\ 2x + 3(3) & = & 13 \quad \text{Let } y = 3. \\ 2x + 9 & = & 13 \quad \text{Multiply.} \\ 2x & = & 4 \quad \text{Subtract 9.} \\ x & = & 2 \quad \text{Divide by 2.} \end{array}$$

NOW TRY
EXERCISE 7

Solve the system.

$$2x - 5y = 14$$

$$5x + 2y = 6$$

Step 6 To check, substitute 2 for x and 3 for y in both equations (1) and (2).

CHECK	$5x - 2y = 4$ (1)	$2x + 3y = 13$ (2)
	$5(2) - 2(3) \stackrel{?}{=} 4$	$2(2) + 3(3) \stackrel{?}{=} 13$
	$10 - 6 \stackrel{?}{=} 4$	$4 + 9 \stackrel{?}{=} 13$
	$4 = 4$ ✓ True	$13 = 13$ ✓ True

The solution set is $\{(2, 3)\}$.

NOW TRY

NOTE Look again at the system in **Example 7**.

$$5x - 2y = 4 \quad (1)$$

$$2x + 3y = 13 \quad (2)$$

In Step 2, we could eliminate the variable y instead of x .

$$15x - 6y = 12 \quad \text{3 times each side of equation (1)}$$

$$\underline{4x + 6y = 26} \quad \text{2 times each side of equation (2)}$$

$$19x = 38 \quad \text{Add.}$$

$$x = 2 \quad \text{Divide by 19.}$$

If we substitute 2 for x in equation (1) or (2), we find that $y = 3$. The same ordered-pair solution $(2, 3)$ results.

OBJECTIVE 5 Solve special systems.

As we saw in **FIGURES 3(b) and (c)**, some systems of linear equations have no solution or an infinite number of solutions.

NOW TRY
EXERCISE 8

Solve the system.

$$-2x + 5y = 6$$

$$6x - 15y = 4$$

$$x + 3y = 4 \quad (1)$$

$$-2x - 6y = 3 \quad (2)$$

We multiply equation (1) by 2 and then add the result to equation (2).

$$2x + 6y = 8 \quad \text{Equation (1) multiplied by 2}$$

$$\underline{-2x - 6y = 3} \quad (2)$$

$$0 = 11 \quad \text{False (contradiction)}$$

The result of the addition step is a false statement, $0 = 11$, which indicates that the system is inconsistent. As shown in **FIGURE 5**, the graphs of the equations of the system are parallel lines.

There are no ordered pairs that satisfy both equations, so there is no solution for the system. The solution set is \emptyset .




FIGURE 5

NOW TRY

NOW TRY ANSWERS

7. $\{(2, -2)\}$

8. \emptyset


NOW TRY EXERCISE 9

Solve the system.

$$\begin{aligned}x - 3y &= 7 \\ -3x + 9y &= -21\end{aligned}$$




FIGURE 6

EXAMPLE 9 Solving a System of Dependent Equations (Case 3)

Solve the system.

$$2x - y = 3 \quad (1)$$

$$6x - 3y = 9 \quad (2)$$

We multiply equation (1) by -3 and then add the result to equation (2).

$$-6x + 3y = -9 \quad \text{\textit{-3 times each side of equation (1)}}$$

$$\begin{array}{r} 6x - 3y = 9 \quad (2) \\ -6x + 3y = -9 \end{array}$$

$$0 = 0 \quad \text{\textit{True (identity)}}$$

Adding these equations gives the true statement $0 = 0$. In the original system, we could obtain equation (2) from equation (1) by multiplying equation (1) by 3. Because of this, equations (1) and (2) are equivalent and have the same graph, as shown in **FIGURE 6**. The equations are dependent.

The solution set is the set of all points on the line with equation $2x - y = 3$, which is written in set-builder notation as

$$\{(x, y) \mid 2x - y = 3\}$$

and is read “the set of all ordered pairs (x, y) , such that $2x - y = 3$.” Either equation of the system could be used to write the solution set.

We use the equation in $Ax + By = C$ form with coefficients that are integers having greatest common factor 1 and positive coefficient of x .


NOW TRY
Special Cases of Linear Systems

If both variables are eliminated when a system of linear equations is solved, then the solution sets are determined as follows.

- There is no solution if the resulting statement is *false*. (See **Example 8**.)
- There are infinitely many solutions if the resulting statement is *true*. (See **Example 9**.)

EXAMPLE 10 Using Slope-Intercept Form to Determine the Number of Solutions

Refer to **Examples 8 and 9**. Write the equations of each system in slope-intercept form, and use the results to determine how many solutions the system has.

$$(a) \quad x + 3y = 4 \quad (1)$$

$$-2x - 6y = 3 \quad (2)$$

Solve each equation from **Example 8** for y .

$$x + 3y = 4 \quad (1)$$

$$3y = -x + 4 \quad \text{\textit{Add } -x.}$$

$$y = -\frac{1}{3}x + \frac{4}{3} \quad \text{\textit{Divide by 3.}}$$

$$\begin{array}{c} \uparrow \quad \uparrow \\ \text{Slope} \quad y\text{-intercept } \left(0, \frac{4}{3}\right) \end{array}$$

$$-2x - 6y = 3 \quad (2)$$

$$-6y = 2x + 3 \quad \text{\textit{Add } 2x.}$$

$$y = -\frac{1}{3}x - \frac{1}{2} \quad \text{\textit{Divide by } -6; \frac{2}{-6} = -\frac{1}{3}; \frac{3}{-6} = -\frac{1}{2}}$$

$$\begin{array}{c} \uparrow \quad \uparrow \\ \text{Slope} \quad y\text{-intercept } \left(0, -\frac{1}{2}\right) \end{array}$$

NOW TRY ANSWER

9. $\{(x, y) \mid x - 3y = 7\}$

The lines have the *same* slope, but *different* y -intercepts, indicating that they are parallel. Thus, the system has no solution.

NOW TRY
EXERCISE 10

Write the equations of each system in slope-intercept form, and use the results to determine how many solutions the system has.

- (a) $2x - 3y = 3$
 $4x - 6y = -6$
- (b) $5y = -x - 4$
 $-10y = 2x + 8$

(b) $2x - y = 3$ (1)
 $6x - 3y = 9$ (2)

Solve each equation from **Example 9** for y .

$2x - y = 3 \quad (1)$ $-y = -2x + 3 \quad \text{Add } -2x.$ $y = 2x - 3 \quad \text{Multiply by } -1.$ <p style="margin-left: 40px;"> ↑ Slope ↑ y-intercept $(0, -3)$ </p>	$6x - 3y = 9 \quad (2)$ $-3y = -6x + 9 \quad \text{Add } -6x.$ $y = 2x - 3 \quad \text{Divide by } -3.$ <p style="margin-left: 40px;"> ↑ Slope ↑ y-intercept $(0, -3)$ </p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

The lines have the *same* slope and the *same* y -intercept, which means that they coincide. There are infinitely many solutions.

NOW TRY

NOTE In **Example 2**, we solved the system

$$x + y = 5 \quad (1)$$

$$2x - y = 4 \quad (2)$$

by graphing the two lines and finding their point of intersection. We can also do this with a graphing calculator, as shown in **FIGURE 7**. The two lines were graphed by solving each equation for y .

$$y = 5 - x \quad \text{Equation (1) solved for } y$$

$$y = 2x - 4 \quad \text{Equation (2) solved for } y$$




FIGURE 7

The coordinates of their point of intersection are displayed at the bottom of the screen, indicating that the solution set is $\{(3, 2)\}$. (Compare this graph with the one in **FIGURE 2**.)

NOW TRY ANSWERS

10. (a) $y = \frac{2}{3}x - 1$; $y = \frac{2}{3}x + 1$;
no solution
- (b) Both are $y = -\frac{1}{5}x - \frac{4}{5}$;
infinitely many solutions

3.1 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER


Reread your class notes before working the assigned exercises. **Review Study Skill 3, Taking Lecture Notes.**

Concept Check Complete each statement.

- If $(4, -3)$ is a solution of a linear system in two variables, then substituting _____ for x and _____ for y leads to true statements in *both* equations.
- A solution of a system of independent linear equations in two variables is an ordered _____.
- If solving a system leads to a false statement such as $0 = 3$, the solution set is _____.
- If solving a system leads to a true statement such as $0 = 0$, the system has _____ equations.
- If the two lines forming a system have the same slope and different y -intercepts, the system has (*no / one / infinitely many*) solution(s).
- If the two lines forming a system have different slopes, the system has (*no / one / infinitely many*) solution(s).


7. Concept Check Which ordered pair could be a solution of the graphed system of equations? Why?

- A. (4, 4)
- B. (-4, 4)
- C. (-4, -4)
- D. (4, -4)



8. Concept Check Which ordered pair could be a solution of the graphed system of equations? Why?

- A. (4, 0)
- B. (-4, 0)
- C. (0, 4)
- D. (0, -4)







9. Concept Check Match each system with the correct graph.

(a) $x + y = 6$
 $x - y = 0$

(b) $x + y = -6$
 $x - y = 0$

(c) $x + y = 0$
 $x - y = -6$

(d) $x + y = 0$
 $x - y = 6$



10. Concept Check The following systems have infinitely many solutions. Write the solution set of each system using set-builder notation and the method described in **Example 9**.

(a) $5x - 3y = 2$
 $10x - 6y = 4$

(b) $4x - 2y = 8$
 $2x = y + 4$

(c) $4x - 6y = 10$
 $-8x + 12y = -20$

Determine whether the given ordered pair is a solution of the given system. See **Example 1**.

11. $x - y = 17$
 $x + y = -1$; (8, -9)

12. $x + y = 6$
 $x - y = 4$; (5, 1)

13. $3x - 5y = -12$
 $x - y = 1$; (-1, 2)

14. $2x - y = 8$
 $3x + 2y = 20$; (5, 2)

Solve each system by graphing. See **Example 2**.

15. $x + y = -5$
 $-2x + y = 1$

16. $x + y = 4$
 $2x - y = 2$

17. $x - 4y = -4$
 $3x + y = 1$

18. $6x - y = 2$
 $x - 2y = 4$

19. $2x + y = 4$
 $3x - y = 6$

20. $2x + 3y = -6$
 $x - 3y = -3$

Solve each system using the substitution method. If a system is inconsistent or has dependent equations, say so. See **Examples 3–5, 8, and 9**.

21. $4x + y = 6$
 $y = 2x$

22. $2x - y = 6$
 $y = 5x$

23. $-x - 4y = -14$
 $y = 2x - 1$

24. $-3x - 5y = -17$
 $y = 4x + 8$

25. $4x - 5y = -11$
 $x + 2y = 7$

26. $3x - y = 10$
 $2x + 5y = 1$

$$\begin{aligned} 27. \quad & 2x - y = 4 \\ & 5x - 2y = 8 \end{aligned}$$

$$\begin{aligned} 30. \quad & -3x + y = -5 \\ & 3x + 6y = 0 \end{aligned}$$

$$\begin{aligned} 33. \quad & x = 3y + 5 \\ & x = \frac{3}{2}y \end{aligned}$$

$$\begin{aligned} 36. \quad & \frac{1}{4}x - \frac{1}{5}y = 9 \\ & 5x - y = 0 \end{aligned}$$

$$\begin{aligned} 39. \quad & \frac{1}{2}x - \frac{1}{4}y = -5 \\ & \frac{1}{8}x + \frac{1}{4}y = 0 \end{aligned}$$

$$\begin{aligned} 42. \quad & x = 3y \\ & 3x - 9y = 0 \end{aligned}$$

$$\begin{aligned} 28. \quad & 3x + 2y = 6 \\ & 4x + y = 3 \end{aligned}$$

$$\begin{aligned} 31. \quad & 5x - 4y = 9 \\ & 3 - 2y = -x \end{aligned}$$

$$\begin{aligned} 34. \quad & x = 6y - 2 \\ & x = \frac{3}{4}y \end{aligned}$$

$$\begin{aligned} 37. \quad & y = 0.5x \\ & 1.5x - 0.5y = 5.0 \end{aligned}$$

$$\begin{aligned} 40. \quad & -\frac{1}{3}x + \frac{2}{5}y = -6 \\ & -\frac{1}{2}x - \frac{3}{2}y = 12 \end{aligned}$$

$$\begin{aligned} 43. \quad & x = 5y \\ & 5x - 25y = 5 \end{aligned}$$

$$\begin{aligned} 29. \quad & 3x - 4y = -22 \\ & -3x + y = 0 \end{aligned}$$

$$\begin{aligned} 32. \quad & 6x - y = -9 \\ & 4 + 7x = -y \end{aligned}$$

$$\begin{aligned} 35. \quad & \frac{1}{2}x + \frac{1}{3}y = 3 \\ & -3x + y = 0 \end{aligned}$$

$$\begin{aligned} 38. \quad & y = 1.4x \\ & 0.5x + 1.5y = 26.0 \end{aligned}$$

$$\begin{aligned} 41. \quad & y = 2x \\ & 4x - 2y = 0 \end{aligned}$$

$$\begin{aligned} 44. \quad & y = -4x \\ & 8x + 2y = 4 \end{aligned}$$

Solve each system using the elimination method. If a system is inconsistent or has dependent equations, say so. See Examples 6–9.

$$\begin{aligned} 45. \quad & -2x + 3y = -16 \\ & 2x - 5y = 24 \end{aligned}$$

$$\begin{aligned} 48. \quad & -2x + 3y = 1 \\ & -4x + y = -3 \end{aligned}$$

$$\begin{aligned} 51. \quad & 7x + 2y = 6 \\ & -14x - 4y = -12 \end{aligned}$$

$$\begin{aligned} 54. \quad & 8x + 4y = 0 \\ & 4x - 2y = 2 \end{aligned}$$

$$\begin{aligned} 57. \quad & x + y = 0 \\ & 2x - 2y = 0 \end{aligned}$$

$$\begin{aligned} 60. \quad & \frac{3}{2}x + y = 3 \\ & \frac{2}{3}x + \frac{1}{3}y = 1 \end{aligned}$$

$$\begin{aligned} 46. \quad & 6x + 5y = -7 \\ & -6x - 11y = 1 \end{aligned}$$

$$\begin{aligned} 49. \quad & 3x + 4y = -6 \\ & 5x + 3y = 1 \end{aligned}$$

$$\begin{aligned} 52. \quad & x - 4y = 2 \\ & 4x - 16y = 8 \end{aligned}$$

$$\begin{aligned} 55. \quad & 5x - 5y = 3 \\ & x - y = 12 \end{aligned}$$

$$\begin{aligned} 58. \quad & 3x + 3y = 0 \\ & -2x - y = 0 \end{aligned}$$

$$\begin{aligned} 61. \quad & \frac{1}{2}x + \frac{1}{3}y = \frac{49}{18} \\ & \frac{1}{2}x + 2y = \frac{4}{3} \end{aligned}$$

$$\begin{aligned} 47. \quad & 2x - 5y = 11 \\ & 3x + y = 8 \end{aligned}$$

$$\begin{aligned} 50. \quad & 4x + 3y = 1 \\ & 3x + 2y = 2 \end{aligned}$$

$$\begin{aligned} 53. \quad & 3x + 3y = 0 \\ & 4x + 2y = 3 \end{aligned}$$

$$\begin{aligned} 56. \quad & 2x - 3y = 7 \\ & -4x + 6y = 14 \end{aligned}$$

$$\begin{aligned} 59. \quad & x - \frac{1}{2}y = 2 \\ & -x + \frac{2}{5}y = -\frac{8}{5} \end{aligned}$$

$$\begin{aligned} 62. \quad & \frac{1}{5}x + \frac{1}{7}y = \frac{12}{5} \\ & \frac{1}{10}x + \frac{1}{3}y = \frac{5}{6} \end{aligned}$$

63. Concept Check To minimize the amount of work required, determine whether to use the substitution or the elimination method to solve each system. Explain. *Do not actually solve.*

$$\begin{aligned} \text{(a)} \quad & 3x + y = -7 \\ & x - y = -5 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad & 6x - y = 5 \\ & y = 11x \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad & 3x - 2y = 0 \\ & 9x + 8y = 7 \end{aligned}$$

64. Concept Check Explain why *System A* would be more difficult to solve by substitution than *System B*.

$$\begin{array}{l} \text{System A: } 3x + 7y = 19 \\ \quad \quad \quad 8x - 5y = 4 \end{array} \qquad \begin{array}{l} \text{System B: } x + 7y = 8 \\ \quad \quad \quad 3x - 2y = 4 \end{array}$$

Solve each system using the method of your choice. See Examples 3–9.

- | | | |
|------------------------------------------------------------------|--------------------------------------------------------------------------|------------------------------------------------------------------------------------|
| 65. $3x + y = -7$
$x - y = -5$ | 66. $6x - y = 5$
$y = 11x$ | 67. $3x - 2y = 0$
$9x + 8y = 7$ |
| 68. $3x - 5y = 7$
$2x + 3y = 30$ | 69. $2x + 3y = 10$
$-3x + y = 18$ | 70. $x + y = 10$
$2x - y = 5$ |
| 71. $y = -\frac{1}{5}x + 1$


$y = \frac{3}{5}x - 3$ | 72. $7x - 8y = 0$
$2x + 5y = 0$ | 73. $0.3x + 0.2y = 0.4$
$0.5x + 0.4y = 0.7$ |
| 74. $0.2x + 0.5y = 6$
$0.4x + y = 9$ | 75. $\frac{1}{2}x - \frac{1}{8}y = -\frac{1}{4}$
$4x - y = -2$ | 76. $\frac{1}{6}x + \frac{1}{3}y = 8$
$\frac{1}{4}x + \frac{1}{2}y = 12$ |

Write the equations of each system in slope-intercept form, and use the results to determine how many solutions the system has. Do not actually solve. See Example 10.


- | | | | |
|--------------------------------------------|-------------------------------------------|---------------------------------------------|----------------------------------------------|
| 77. $3x + 7y = 4$
$6x + 14y = 3$ | 78. $-x + 2y = 8$
$4x - 8y = 1$ | 79. $2x = -3y + 1$
$6x = -9y + 3$ | 80. $5x = -2y + 1$
$10x = -4y + 2$ |
|--------------------------------------------|-------------------------------------------|---------------------------------------------|----------------------------------------------|

Answer each question using the graphs provided.


- 81.** The figure shows graphs that represent supply and demand for a certain brand of low-fat frozen yogurt at various prices per half-gallon (in dollars).
- At what price does supply equal demand?
 - For how many half-gallons does supply equal demand?
 - What are the supply and demand at a price of \$2 per half-gallon?



- 82.** Sharon compared the monthly payments she would incur for two types of mortgages: fixed rate and variable rate. Her observations led to the following graphs.
- For which years would the monthly payment be more for the fixed-rate mortgage than for the variable-rate mortgage?
 - In what year would the payments be the same? What would those payments be?
 - In year 10, what are the fixed-rate and variable-rate monthly payments?



83. The graph shows college student enrollment (in thousands) for three states from 2011 through 2016.



Data from National Student Clearinghouse Research Center.


- During what years did the number of college students enrolled in Colorado exceed that for Utah?
 - Between what two consecutive years did Utah first have the highest enrollment?
 - In what year were the numbers of college students enrolled in Kentucky and Utah approximately equal? Approximately how many college students were enrolled in each state during that year?
 - Write the answer for part (c) as an ordered pair of the form (year, enrollment).
 - Describe the trends in enrollment in Kentucky and Utah from 2012 through 2014.
84. The graph shows numbers of nuclear weapons possessed by the United States and USSR/Russia from 1950 through 2017.



Data from *The World Almanac and Book of Facts*.

- In what year were the numbers of nuclear weapons approximately equal?
- Approximately how many nuclear weapons did each country possess in that year?
- Express the point of intersection of the graphs for U.S. and Russian nuclear weapons as an ordered pair of the form (year, number of nuclear weapons).
- Over the entire period 1950–2017, which country possessed the greatest number of nuclear weapons at any one particular time? In what year did that maximum occur, and what was the approximate number of nuclear weapons possessed?
- Describe the trend in number of nuclear weapons possessed by Russia from 1990 through 2017. If a straight line were used to approximate its graph, would the line have a slope that is positive, negative, or 0?

Use the graph shown in **FIGURE 1** at the beginning of this section (and repeated here) to work each exercise.



Data from Netflix, Leichtman Research Group.

85. Which type of subscription was more popular in 2017?
86. Between which two years were the numbers of Netflix and cable TV subscribers the same? About how many people subscribed to each video content provider at that point?
87. If $x = 0$ represents 2012 and $x = 5$ represents 2017, the number of subscribers y in millions can be modeled by the linear equations in the following system.

$$\begin{aligned} -5.61x + y &= 23.9 && \text{Netflix} \\ 0.77x + y &= 52.1 && \text{Cable TV} \end{aligned}$$

Solve this system. Express values as decimals rounded to the nearest tenth. Write the solution as an ordered pair of the form (year, number of subscribers).

88. Interpret the answer for **Exercise 87**, rounding down for the year. How does this compare to the estimate from **Exercise 86**?

Extending Skills The following systems can be solved by elimination. One way to do this is to let $p = \frac{1}{x}$ and $q = \frac{1}{y}$. Substitute, solve for p and q , and then find x and y . (For example, in **Exercise 89**, $\frac{3}{x} = 3 \cdot \frac{1}{x} = 3p$.)

Use this method to solve each system.

89.
$$\begin{aligned} \frac{3}{x} + \frac{4}{y} &= \frac{5}{2} \\ \frac{5}{x} - \frac{3}{y} &= \frac{7}{4} \end{aligned}$$

90.
$$\begin{aligned} \frac{2}{x} - \frac{5}{y} &= \frac{3}{2} \\ \frac{4}{x} + \frac{1}{y} &= \frac{4}{5} \end{aligned}$$

91.
$$\begin{aligned} \frac{2}{x} + \frac{3}{y} &= \frac{11}{2} \\ -\frac{1}{x} + \frac{2}{y} &= -1 \end{aligned}$$

92. **Extending Skills** Make up a system of the following form, where $a, b, c, d, e,$ and f are consecutive integers. Solve the system.

$$\begin{aligned} ax + by &= c \\ dx + ey &= f \end{aligned}$$

Extending Skills Solve by any method. Assume that a and b represent nonzero constants.

93.
$$\begin{aligned} ax + by &= c \\ ax - 2by &= c \end{aligned}$$

 94.
$$\begin{aligned} ax + by &= 2 \\ -ax + 2by &= 1 \end{aligned}$$

 95.
$$\begin{aligned} 2ax - y &= 3 \\ y &= 5ax \end{aligned}$$

 96.
$$\begin{aligned} 3ax + 2y &= 1 \\ -ax + y &= 2 \end{aligned}$$

3.2 Systems of Linear Equations in Three Variables

OBJECTIVES

- 1 Understand the geometry of systems of three equations in three variables.
- 2 Solve linear systems (with three equations and three variables) by elimination.
- 3 Solve linear systems (with three equations and three variables) in which some of the equations have missing terms.
- 4 Solve special systems.

A solution of an equation in three variables, such as

$$2x + 3y - z = 4, \quad \text{Linear equation in three variables}$$

is an **ordered triple** and is written (x, y, z) . For example, the ordered triple $(0, 1, -1)$ is a solution of the preceding equation, because

$$2(0) + 3(1) - (-1) = 4 \quad \text{is a true statement.}$$

Verify that another solution of this equation is $(10, -3, 7)$.

We now extend the term *linear equation* to equations of the form

$$Ax + By + Cz + \cdots + Dw = K,$$

where not all the coefficients A, B, C, \dots, D equal 0. For example,

$$2x + 3y - 5z = 7 \quad \text{and} \quad x - 2y - z + 3w = 8$$

are linear equations, the first with three variables and the second with four.

OBJECTIVE 1 Understand the geometry of systems of three equations in three variables.

Consider the solution of a system such as the following.

$$4x + 8y + z = 2$$

$$x + 7y - 3z = -14 \quad \text{System of linear equations in three variables}$$

$$2x - 3y + 2z = 3$$

Theoretically, a system of this type can be solved by graphing. However, the graph of a linear equation with three variables is a *plane*, not a line. Because visualizing a plane requires three-dimensional graphing, the method of graphing is not practical with these systems. However, it does illustrate the number of solutions possible for such systems, as shown in **FIGURE 8**.

VOCABULARY

- ordered triple
- focus variable
- working equation




FIGURE 8

STUDY SKILLS REMINDER

Make study cards to help you learn and remember the material in this chapter.

Review Study Skill 5, Using Study Cards.

FIGURE 8 illustrates the following cases.

Graphs of Linear Systems in Three Variables

Case 1 The three planes may meet at a single, common point.

This point is the solution of the system. See FIGURE 8(a).

Case 2 The three planes may have the points of a line in common.

The infinite set of points that satisfy the equation of the line is the solution of the system. See FIGURES 8(b) and (c).

Case 3 The three planes may coincide.

The solution of the system is the set of all points on a plane. See FIGURE 8(d).

Case 4 The planes may have no points common to all three.

There is no solution of the system. See FIGURES 8(e)–(h).

OBJECTIVE 2 Solve linear systems (with three equations and three variables) by elimination.

Because graphing to find the solution set of a system of three equations in three variables is impractical, these systems are solved with an extension of the elimination method.

In the steps that follow, we use the term **focus variable** to identify the first variable to be eliminated. The focus variable will always be present in the **working equation**, which will be used twice to eliminate this variable.

Solving a Linear System in Three Variables

Step 1 **Select a variable and an equation.** A good choice for the variable, which we call the *focus variable*, is one that has coefficient 1 or -1 . Then select an equation, one that contains the focus variable, as the *working equation*.

Step 2 **Eliminate the focus variable.** Use the working equation and one of the other two equations of the original system. The result is an equation in two variables.

Step 3 **Eliminate the focus variable again.** Use the working equation and the remaining equation of the original system. The result is another equation in two variables.

Step 4 **Write the equations in two variables that result from Steps 2 and 3 as a system, and solve it.** Doing this gives the values of two of the variables.

Step 5 **Find the value of the remaining variable.** Substitute the values of the two variables found in Step 4 into the working equation to obtain the value of the focus variable.

Step 6 **Check** the three values in *each* of the *original* equations of the system. Then write the solution set as a set containing an ordered triple.

EXAMPLE 1 Solving a System in Three Variables

Solve the system.

$$4x + 8y + z = 2 \quad (1)$$

$$x + 7y - 3z = -14 \quad (2)$$

$$2x - 3y + 2z = 3 \quad (3)$$

Step 1 Because z in equation (1) has coefficient 1, we choose z as the focus variable and (1) as the working equation. (Another option would be to choose x as the focus variable—it also has coefficient 1—and use (2) as the working equation.)

$$4x + 8y + \underset{\substack{\downarrow \text{Focus variable} \\ \leftarrow \text{Working equation}}}{z} = 2 \quad (1)$$

Step 2 Multiply working equation (1) by 3 and add the result to equation (2).

$$12x + 24y + 3z = 6 \quad \text{Multiply each side of (1) by 3.}$$

$$x + 7y - 3z = -14 \quad (2)$$

Focus variable z was eliminated.

$$13x + 31y = -8 \quad \text{Add. (4)}$$

Step 3 Multiply working equation (1) by -2 and add the result to remaining equation (3) to again eliminate focus variable z .

$$-8x - 16y - 2z = -4 \quad \text{Multiply each side of (1) by } -2.$$

$$2x - 3y + 2z = 3 \quad (3)$$

Focus variable z was eliminated.

$$-6x - 19y = -1 \quad \text{Add. (5)}$$

Step 4 Write the equations in two variables that result in Steps 2 and 3 as a system.

Make sure these equations have the same two variables.

$$13x + 31y = -8 \quad (4) \quad \text{The result from Step 2}$$

$$-6x - 19y = -1 \quad (5) \quad \text{The result from Step 3}$$

Now solve this system. We choose to eliminate x .

$$78x + 186y = -48 \quad \text{Multiply each side of (4) by 6.}$$

$$-78x - 247y = -13 \quad \text{Multiply each side of (5) by 13.}$$

$$-61y = -61 \quad \text{Add.}$$

$$y = 1 \quad \text{Divide by } -61.$$

Substitute 1 for y in either equation (4) or equation (5) to find x .

$$-6x - 19y = -1 \quad (5)$$

$$-6x - 19(1) = -1 \quad \text{Let } y = 1.$$

$$-6x - 19 = -1 \quad \text{Multiply.}$$

$$-6x = 18 \quad \text{Add 19.}$$

$$x = -3 \quad \text{Divide by } -6.$$

Step 5 Now substitute the two values we found in Step 4 in working equation (1) to find the value of the remaining variable, focus variable z .

$$4x + 8y + z = 2 \quad (1)$$

$$4(-3) + 8(1) + z = 2 \quad \text{Let } x = -3 \text{ and } y = 1.$$


$$-4 + z = 2 \quad \text{Multiply, and then add.}$$

$$z = 6 \quad \text{Add 4.}$$

NOW TRY
EXERCISE 1

Solve the system.

$$\begin{aligned}x - y + 2z &= 1 \\3x + 2y + 7z &= 8 \\-3x - 4y + 9z &= -10\end{aligned}$$



A single solution

FIGURE 8(a) (repeated)

Write the values of x , y , and z in the correct order.

Step 6 It appears that $(-3, 1, 6)$ is the only solution of the system. We must check that this ordered triple satisfies all three original equations of the system. We begin with equation (1).

CHECK

$$\begin{aligned}4x + 8y + z &= 2 && (1) \\4(-3) + 8(1) + 6 &\stackrel{?}{=} 2 && \text{Substitute.} \\-12 + 8 + 6 &\stackrel{?}{=} 2 && \text{Multiply.} \\2 &= 2 && \checkmark \text{ True}\end{aligned}$$

Because $(-3, 1, 6)$ also satisfies equations (2) and (3), the solution set contains a single ordered triple, $\{(-3, 1, 6)\}$. This is Case 1, shown earlier in FIGURE 8(a).

NOW TRY

OBJECTIVE 3 Solve linear systems (with three equations and three variables) in which some of the equations have missing terms.

If a linear system includes an equation that is missing a term or terms, one elimination step can be omitted.

EXAMPLE 2 Solving a System of Equations with Missing Terms

Solve the system.

$$\begin{aligned}6x - 12y &= -5 && (1) \text{ Missing } z \\8y + z &= 0 && (2) \text{ Missing } x \\9x - z &= 12 && (3) \text{ Missing } y\end{aligned}$$

Equation (1) is missing the variable z , so one way to begin is to use z as the focus variable and eliminate z again using equations (2) and (3).

$$\begin{aligned}8y + z &= 0 && (2) \\9x - z &= 12 && (3) \\ \hline 9x + 8y &= 12 && \text{Add. (4)}\end{aligned}$$

Use resulting equation (4) in x and y , together with equation (1), $6x - 12y = -5$, to eliminate x .

$$\begin{aligned}18x - 36y &= -15 && \text{Multiply each side of (1) by 3.} \\-18x - 16y &= -24 && \text{Multiply each side of (4) by } -2. \\ \hline -52y &= -39 && \text{Add.} \\ y &= \frac{-39}{-52} && \text{Divide by } -52. \\ y &= \frac{3}{4} && \text{Write in lowest terms.}\end{aligned}$$

We can find z by substituting this value for y in equation (2).

$$\begin{aligned}8y + z &= 0 && (2) \\8\left(\frac{3}{4}\right) + z &= 0 && \text{Let } y = \frac{3}{4}. \\6 + z &= 0 && \text{Multiply.} \\z &= -6 && \text{Subtract 6.}\end{aligned}$$

NOW TRY ANSWER

1. $\{(2, 1, 0)\}$


NOW TRY EXERCISE 2

Solve the system.

$$3x - z = -10$$

$$4y + 5z = 24$$

$$x - 6y = -8$$

We can find x by substituting -6 for z in equation (3).

$$9x - z = 12 \quad (3)$$

$$9x - (-6) = 12 \quad \text{Let } z = -6.$$

$$9x + 6 = 12 \quad -(-a) = a$$

$$x = \frac{6}{9} \quad \text{Subtract 6. Divide by 9.}$$

$$x = \frac{2}{3} \quad \text{Write in lowest terms.}$$

Check to verify that the solution set is $\left\{\left(\frac{2}{3}, \frac{3}{4}, -6\right)\right\}$. This is also an example of Case 1.**NOW TRY** 

NOTE Another way to solve the system in **Example 2** is to begin by eliminating the variable y from equations (1) and (2). The resulting equation together with equation (3) forms a system of two equations in the variables x and z . Try working **Example 2** this way to see that the same solution results. There are often multiple ways to solve a system of equations. Some ways may involve more work than others.

OBJECTIVE 4 Solve special systems.

NOW TRY EXERCISE 3

Solve the system.

$$x - 3y + 2z = 10$$

$$-2x + 6y - 4z = -20$$

$$\frac{1}{2}x - \frac{3}{2}y + z = 5$$

EXAMPLE 3 Solving a System of Dependent Equations with Three Variables

Solve the system.

$$2x - 3y + 4z = 8 \quad (1)$$

$$-x + \frac{3}{2}y - 2z = -4 \quad (2)$$
 Use as the working equation, with focus variable x .

$$6x - 9y + 12z = 24 \quad (3)$$

Eliminate focus variable x using equations (1) and (2).

$$2x - 3y + 4z = 8 \quad (1)$$

$$\underline{-2x + 3y - 4z = -8} \quad \text{Multiply each side of (2) by 2.}$$

$$0 = 0 \quad \text{True (identity)}$$

Eliminating x from equations (2) and (3) gives the same result.

$$-6x + 9y - 12z = -24 \quad \text{Multiply each side of (2) by 6.}$$

$$\underline{6x - 9y + 12z = 24} \quad (3)$$

$$0 = 0 \quad \text{True (identity)}$$

When solving a system such as this, attempting to eliminate one variable results in elimination of *all* variables. The equations are dependent—that is, they are equivalent forms of the *same* equation—and have the same graph. This is Case 3, as illustrated in **FIGURE 8(d)**. The solution set is written

$$\{(x, y, z) \mid 2x - 3y + 4z = 8\}. \quad \text{Set-builder notation}$$

Although any one of the three equations could be used to write the solution set, we use the equation with integer coefficients having greatest common factor 1 and positive coefficient of x , as in the previous section.

NOW TRY 

All points in common

FIGURE 8(d) (repeated)**NOW TRY ANSWERS**

2. $\{(-2, 1, 4)\}$

3. $\{(x, y, z) \mid x - 3y + 2z = 10\}$

NOW TRY EXERCISE 4

Solve the system.

$$\begin{aligned}x - 5y + 2z &= 4 \\3x + y - z &= 6 \\-2x + 10y - 4z &= 7\end{aligned}$$

EXAMPLE 4 Solving an Inconsistent System with Three Variables

Solve the system.


$$\begin{aligned}2x - 4y + 6z &= 5 & (1) \\-x + 3y - 2z &= -1 & (2) \\x - 2y + 3z &= 1 & (3)\end{aligned}$$

Use as the working equation, with focus variable x .

Eliminate the focus variable, x , using equations (1) and (3).

$$\begin{array}{r} -2x + 4y - 6z = -2 \quad \text{Multiply each side of (3) by } -2. \\ \underline{2x - 4y + 6z = 5} \quad (1) \\ \hline 0 = 3 \quad \text{False (contradiction)} \end{array}$$

The resulting false statement indicates that equations (1) and (3) have no common solution. Thus, the system is inconsistent and the solution set is \emptyset . The graph of this system would show two planes parallel to one another, and a third plane that intersects both, as in **FIGURE 8(f)**. This is Case 4.



No points in common

FIGURE 8(f) (repeated)

NOW TRY

NOTE If a false statement results when adding as in **Example 4**, it is not necessary to go any further with the solution. Because two of the three planes are parallel, it is not possible for the three planes to have any points in common.

NOW TRY EXERCISE 5

Solve the system.

$$\begin{aligned}x - 3y + 2z &= 4 \\ \frac{1}{3}x - y + \frac{2}{3}z &= 7 \\ \frac{1}{2}x - \frac{3}{2}y + z &= 2\end{aligned}$$

EXAMPLE 5 Solving Another Special System

Solve the system.

$$\begin{aligned}2x - y + 3z &= 6 & (1) \\x - \frac{1}{2}y + \frac{3}{2}z &= 3 & (2) \\4x - 2y + 6z &= 1 & (3)\end{aligned}$$


Equations (1) and (2) are equivalent. If we multiply each side of equation (2) by 2, we obtain equation (1). These two equations are dependent and have the same graph.

Equations (1) and (3) are *not* equivalent, however. If we multiply each side of equation (3) by $\frac{1}{2}$, we obtain

$$2x - y + 3z = \frac{1}{2}.$$

This equation has the same coefficients as equation (1), but a different constant term. Therefore, the graphs of equations (1) and (3) have *no* points in common—that is, the planes are parallel.

Thus, this system is inconsistent and the solution set is \emptyset , as illustrated in **FIGURE 8(h)**. This is another example of Case 4.



No points in common

FIGURE 8(h) (repeated)

NOW TRY

NOW TRY ANSWERS

4. \emptyset
5. \emptyset

3.2 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

How are you doing on your homework? **Review Study Skill 4, Completing Your Homework.**

Concept Check Answer each of the following.

- Using your immediate surroundings, give an example of three planes that satisfy the condition.
 - They intersect in a single point.
 - They do not intersect.
 - They intersect in infinitely many points.

- Suppose that a system has infinitely many ordered-triple solutions of the form (x, y, z) such that

$$x + y + 2z = 1.$$

Give three specific ordered triples that are solutions of the system.

- Explain what the following statement means: “The solution set of the following system is $\{(-1, 2, 3)\}$.”

$$2x + y + z = 3$$

$$3x - y + z = -2$$

$$4x - y + 2z = 0$$

- The following two equations have a common solution of $(1, 2, 3)$.

$$x + y + z = 6$$

$$2x - y + z = 3$$

Which equation would complete a system of three linear equations in three variables having solution set $\{(1, 2, 3)\}$?

A. $3x + 2y - z = 1$

B. $3x + 2y - z = 4$

C. $3x + 2y - z = 5$

D. $3x + 2y - z = 6$

- What constant should replace the question mark in this system so that the solution set is $\{(1, 1, 1)\}$?

$$2x - 3y + z = 0$$

$$-5x + 2y - z = -4$$

$$x + y + 2z = ?$$

- Complete the work of **Example 1** and show that the ordered triple $(-3, 1, 6)$ is also a solution of equations (2) and (3).

$$x + 7y - 3z = -14 \quad \text{Equation (2)}$$

$$2x - 3y + 2z = 3 \quad \text{Equation (3)}$$

Solve each system. See **Example 1**.

7. $2x - 5y + 3z = -1$

8. $x + 3y - 6z = 1$

9. $3x + 2y + z = 8$

$$x + 4y - 2z = 9$$

$$2x - y + z = 7$$

$$2x - 3y + 2z = -16$$

$$x - 2y - 4z = -5$$

$$x + 2y + 2z = 14$$

$$x + 4y - z = 20$$

10. $-3x + y - z = -10$

11. $2x + 5y + 2z = 0$

12. $5x - 2y + 3z = -9$

$$-4x + 2y + 3z = -1$$

$$4x - 7y - 3z = 1$$

$$4x + 3y + 5z = 4$$

$$2x + 3y - 2z = -5$$

$$3x - 8y - 2z = -6$$

$$2x + 4y - 2z = 14$$

13. $x + 2y + z = 4$

14. $x - 2y + 5z = -7$

15. $-x + 2y + 6z = 2$

$$2x + y - z = -1$$

$$-2x - 3y + 4z = -14$$

$$3x + 2y + 6z = 6$$

$$x - y - z = -2$$

$$-3x + 5y - z = -7$$

$$x + 4y - 3z = 1$$

$$\begin{array}{lll} 16. & 2x + y + 2z = 1 & 17. & x + y - z = -2 & 18. & x + 2y + 3z = 1 \\ & x + 2y + z = 2 & & 2x - y + z = -5 & & -x - y + 3z = 2 \\ & x - y - z = 0 & & -x + 2y - 3z = -4 & & -6x + y + z = -2 \end{array}$$

$$\begin{array}{ll} 19. & \frac{1}{3}x + \frac{1}{6}y - \frac{2}{3}z = -1 \\ & -\frac{3}{4}x - \frac{1}{3}y - \frac{1}{4}z = 3 \\ & \frac{1}{2}x + \frac{3}{2}y + \frac{3}{4}z = 21 \end{array} \qquad \begin{array}{l} 20. & \frac{2}{3}x - \frac{1}{4}y + \frac{5}{8}z = 0 \\ & \frac{1}{5}x + \frac{2}{3}y - \frac{1}{4}z = -7 \\ & -\frac{3}{5}x + \frac{4}{3}y - \frac{7}{8}z = -5 \end{array}$$

$$\begin{array}{ll} 21. & 5.5x - 2.5y + 1.6z = 11.83 \\ & 2.2x + 5.0y - 0.1z = -5.97 \\ & 3.3x - 7.5y + 3.2z = 21.25 \end{array} \qquad \begin{array}{l} 22. & 6.2x - 1.4y + 2.4z = -1.80 \\ & 3.1x + 2.8y - 0.2z = 5.68 \\ & 9.3x - 8.4y - 4.8z = -34.20 \end{array}$$

Solve each system. See Example 2.

$$\begin{array}{lll} 23. & 2x - 3y + 2z = -1 & 24. & 2x - y + 3z = 6 & 25. & 4x + 2y - 3z = 6 \\ & x + 2y + z = 17 & & x + 2y - z = 8 & & x - 4y + z = -4 \\ & 2y - z = 7 & & 2y + z = 1 & & -x + 2z = 2 \end{array}$$

$$\begin{array}{lll} 26. & 2x + 3y - 4z = 4 & 27. & 2x + y = 6 & 28. & 4x - 8y = -7 \\ & x - 6y + z = -16 & & 3y - 2z = -4 & & 4y + z = 7 \\ & -x + 3z = 8 & & 3x - 5z = -7 & & -8x + z = -4 \end{array}$$

$$\begin{array}{lll} 29. & -5x + 2y + z = 5 & 30. & -4x + 3y - z = 4 & 31. & 7x - 3z = -34 \\ & -3x - 2y - z = 3 & & -5x - 3y + z = -4 & & 2y + 4z = 20 \\ & -x + 6y = 1 & & -2x - 3z = 12 & & \frac{3}{4}x + \frac{1}{6}y = -2 \end{array}$$

$$\begin{array}{lll} 32. & 5x - 2z = 8 & 33. & 4x - z = -6 & 34. & 5x - z = 38 \\ & 4y + 3z = -9 & & \frac{3}{5}y + \frac{1}{2}z = 0 & & \frac{2}{3}y + \frac{1}{4}z = -17 \\ & \frac{1}{2}x + \frac{2}{3}y = -1 & & \frac{1}{3}x + \frac{2}{3}z = -5 & & \frac{1}{5}y + \frac{5}{6}z = 4 \end{array}$$

Solve each system. If the system is inconsistent or has dependent equations, say so. See Examples 1, 3, 4, and 5.

$$\begin{array}{lll} 35. & 2x + 2y - 6z = 5 & 36. & -2x + 5y + z = -3 & 37. & -5x + 5y - 20z = -40 \\ & -3x + y - z = -2 & & 5x + 14y - z = -11 & & x - y + 4z = 8 \\ & -x - y + 3z = 4 & & 7x + 9y - 2z = -5 & & 3x - 3y + 12z = 24 \end{array}$$

$$\begin{array}{lll} 38. & x + 4y - z = 3 & 39. & x + 5y - 2z = -1 & 40. & x + 3y + z = 2 \\ & -2x - 8y + 2z = -6 & & -2x + 8y + z = -4 & & 4x + y + 2z = -4 \\ & 3x + 12y - 3z = 9 & & 3x - y + 5z = 19 & & 5x + 2y + 3z = -2 \end{array}$$

$$\begin{array}{lll} 41. & 2x + y - z = 6 & 42. & 2x - 8y + 2z = -10 & 43. & x + y - 2z = 0 \\ & 4x + 2y - 2z = 12 & & -x + 4y - z = 5 & & 3x - y + z = 0 \\ & -x - \frac{1}{2}y + \frac{1}{2}z = -3 & & \frac{1}{8}x - \frac{1}{2}y + \frac{1}{8}z = -\frac{5}{8} & & 4x + 2y - z = 0 \end{array}$$

$$\begin{array}{lll}
 44. \quad 2x + 3y - z = 0 & 45. \quad x - 2y + \frac{1}{3}z = 4 & 46. \quad 4x + y - 2z = 3 \\
 \quad \quad x - 4y + 2z = 0 & \quad \quad 3x - 6y + z = 12 & \quad \quad x + \frac{1}{4}y - \frac{1}{2}z = \frac{3}{4} \\
 \quad \quad 3x - 5y - z = 0 & \quad \quad -6x + 12y - 2z = -3 & \quad \quad 2x + \frac{1}{2}y - z = 1
 \end{array}$$

Extending Skills Extend the method of this section to solve each system. Express the solution in the form (x, y, z, w) .

$$\begin{array}{ll}
 47. \quad x + y + z - w = 5 & 48. \quad 3x + y - z + 2w = 9 \\
 \quad \quad 2x + y - z + w = 3 & \quad \quad x + y + 2z - w = 10 \\
 \quad \quad x - 2y + 3z + w = 18 & \quad \quad x - y - z + 3w = -2 \\
 \quad \quad -x - y + z + 2w = 8 & \quad \quad -x + y - z + w = -6 \\
 \\
 49. \quad 3x + y - z + w = -3 & 50. \quad x - 3y + 7z + w = 11 \\
 \quad \quad 2x + 4y + z - w = -7 & \quad \quad 2x + 4y + 6z - 3w = -3 \\
 \quad \quad -2x + 3y - 5z + w = 3 & \quad \quad 3x + 2y + z + 2w = 19 \\
 \quad \quad 5x + 4y - 5z + 2w = -7 & \quad \quad 4x + y - 3z + w = 22
 \end{array}$$

RELATING CONCEPTS For Individual or Group Work (Exercises 51–56)


A circle has an equation of the following form.

$$x^2 + y^2 + ax + by + c = 0 \quad \text{Equation of a circle}$$

It is a fact from geometry that given three **non-collinear** points—that is, points that do not all lie on the same straight line—there will be a circle that contains them. For example, the points $(4, 2)$, $(-5, -2)$, and $(0, 3)$ lie on the circle whose equation is shown in the figure.

Work Exercises 51–56 in order, to find an equation of the circle passing through the points

$$(2, 1), (-1, 0), \text{ and } (3, 3).$$



$$x^2 + y^2 - \frac{7}{5}x + \frac{27}{5}y - \frac{126}{5} = 0$$

51. Let $x = 2$ and $y = 1$ in the general equation $x^2 + y^2 + ax + by + c = 0$ to find an equation in a , b , and c .
52. Let $x = -1$ and $y = 0$ to find a second equation in a , b , and c .
53. Let $x = 3$ and $y = 3$ to find a third equation in a , b , and c .
54. Form a system of three equations using the answers from **Exercises 51–53**. Solve the system to find the values of a , b , and c .
55. Use the values of a , b , and c from **Exercise 54** and the form of the equation of a circle given above to write an equation of the circle passing through the given points.
56. Explain why the relation whose graph is a circle is not a function.

3.3 Applications of Systems of Linear Equations

OBJECTIVES

- 1 Solve geometry problems using two variables.
- 2 Solve money problems using two variables.
- 3 Solve mixture problems using two variables.
- 4 Solve distance-rate-time problems using two variables.
- 5 Solve problems with three variables using a system of three equations.

Although some problems with two unknowns can be solved using just one variable, it is often easier to use two variables and a system of equations. The following problem, which can be solved with a system, appeared in a Hindu work that dates back to about A.D. 850. (See Exercise 24.)

The mixed price of 9 citrons (a lemonlike fruit) and 7 fragrant wood apples is 107; again, the mixed price of 7 citrons and 9 fragrant wood apples is 101. O you arithmetician, tell me quickly the price of a citron and the price of a wood apple here, having distinctly separated those prices well.



PROBLEM-SOLVING HINT When solving an applied problem using two variables, it is a good idea to pick letters that correspond to the descriptions of the unknown quantities. In the example above, we could choose

c to represent the number of citrons,
and w to represent the number of wood apples.

The following steps are based on the problem-solving method presented earlier in the text.

Solving an Applied Problem Using a System of Equations

- Step 1** **Read** the problem carefully. *What information is given? What is to be found?*
- Step 2** **Assign variables** to represent the unknown values. Write down what each variable represents. Make a sketch, diagram, or table, as needed.
- Step 3** **Write a system of equations** using all the variables.
- Step 4** **Solve** the system of equations.
- Step 5** **State the answer.** Label it appropriately. *Does it seem reasonable?*
- Step 6** **Check** the answer in the words of the *original* problem.

STUDY SKILLS REMINDER

Are you fully utilizing the features of your text? **Review Study Skill 1, Using Your Math Text.**

OBJECTIVE 1 Solve geometry problems using two variables.

EXAMPLE 1 Finding the Dimensions of a Soccer Field

A rectangular soccer field may have a width between 50 and 100 yd and a length between 100 and 130 yd. One particular soccer field has a perimeter of 320 yd. Its length measures 40 yd more than its width. What are the dimensions of this field? (Data from www.soccer-training-guide.com)

- Step 1** **Read** the problem again. We must find the dimensions of the field.
- Step 2** **Assign variables.** A sketch may be helpful. See **FIGURE 9** on the next page.

Let L = the length and W = the width.


NOW TRY EXERCISE 1

A rectangular parking lot has a length that is 10 ft more than twice its width. The perimeter of the parking lot is 620 ft. What are the dimensions of the parking lot?




FIGURE 9

Step 3 Write a system of equations. Because the perimeter is 320 yd, we find one equation by using the perimeter formula.

$$2L + 2W = 320 \quad 2L + 2W = P$$

For a second equation, use the information given about the length.

$$L = W + 40 \quad \text{The length is 40 yd more than the width.}$$

These two equations form a system of equations.

$$2L + 2W = 320 \quad (1)$$

$$L = W + 40 \quad (2)$$

Step 4 Solve the system. Equation (2) is solved for L , so we use the substitution method, substituting $W + 40$ for L in equation (1), and solving for W .

$$2L + 2W = 320 \quad (1)$$

$$2(W + 40) + 2W = 320 \quad \text{Let } L = W + 40.$$

$$2W + 80 + 2W = 320 \quad \text{Distributive property}$$

$$4W + 80 = 320 \quad \text{Combine like terms.}$$

$$4W = 240 \quad \text{Subtract 80.}$$

$$W = 60 \quad \text{Divide by 4.}$$

Be sure to use parentheses around $W + 40$.

Don't stop here.

Let $W = 60$ in the equation $L = W + 40$ to find L .

$$L = 60 + 40 = 100$$

Step 5 State the answer. The length is 100 yd, and the width is 60 yd. Both dimensions are within the ranges given in the problem.

Step 6 Check using the words of the *original* problem.

$$2(100) + 2(60) = 320 \quad \text{The perimeter is 320 yd, as required}$$

$$100 = 60 + 40 \quad \text{Length is 40 yd more than width, as required.}$$

NOW TRY 

PROBLEM-SOLVING HINT There is often more than one way to write the equations in a system used to solve an application. In **Example 1**, we might write the second equation as

$$W = L - 40. \quad (2)$$

In this case, we would substitute $L - 40$ for W in equation (1) to obtain

$$2L + 2(L - 40) = 320 \quad \text{Let } W = L - 40.$$

and solve for L first (instead of W , as in Step 4 above). The *same* answer results.

NOW TRY ANSWER

1. length: 210 ft; width: 100 ft


**NOW TRY
EXERCISE 2**

Two general admission tickets and three tickets for children to a theme park cost \$239.95. One general admission ticket and four tickets for children cost \$224.95. Determine the ticket prices for general admission and for children.

OBJECTIVE 2 Solve money problems using two variables.

EXAMPLE 2 Solving a Problem about Ticket Prices

For the 2015–2016 National Football League and National Basketball Association seasons, two football tickets and one basketball ticket purchased at their average prices cost \$241.84. One football ticket and two basketball tickets cost \$204.74. What were the average ticket prices for the two sports? (Data from Team Marketing Report.)

Step 1 **Read** the problem again. There are two unknowns.

Step 2 **Assign variables.**

Let f = the average price for a football ticket
and b = the average price for a basketball ticket.

Step 3 **Write a system of equations.** We write one equation using the fact that two football tickets and one basketball ticket cost \$241.84.

$$2f + b = 241.84$$

By similar reasoning, we can write a second equation.

$$f + 2b = 204.74$$

These two equations form a system of equations.

$$2f + b = 241.84 \quad (1)$$

$$f + 2b = 204.74 \quad (2)$$

Step 4 **Solve** the system. Either the substitution or the elimination method can be used, based on personal preference. We choose the elimination method. To eliminate f , multiply equation (2) by -2 and add.

$$2f + b = 241.84 \quad (1)$$

$$\underline{-2f - 4b = -409.48} \quad \text{Multiply each side of (2) by } -2.$$

$$-3b = -167.64 \quad \text{Add.}$$

$$b = 55.88 \quad \text{Divide by } -3.$$

To find the value of f , let $b = 55.88$ in equation (2).

$$f + 2b = 204.74 \quad (2)$$

$$f + 2(55.88) = 204.74 \quad \text{Let } b = 55.88.$$

$$f + 111.76 = 204.74 \quad \text{Multiply.}$$

$$f = 92.98 \quad \text{Subtract } 111.76.$$

Step 5 **State the answer.** The average price for one basketball ticket was \$55.88. For one football ticket, the average price was \$92.98.

Step 6 **Check** that these values satisfy the problem conditions.

$$2(\$92.98) + \$55.88 = \$241.84, \quad \text{as required.}$$

$$\$92.98 + 2(\$55.88) = \$204.74, \quad \text{as required.}$$

NOW TRY 

PROBLEM-SOLVING HINT Rework Step 4 of **Example 2** using the substitution method to confirm that the same answer results. (There are two ways to do this—either solve equation (1) for b or solve equation (2) for f .)

Solving using a different method can be a good strategy for checking your answer.

NOW TRY ANSWER

2. general admission: \$56.99;
children: \$41.99

OBJECTIVE 3 Solve mixture problems using two variables.

Some mixture problems can be solved using one variable. For many mixture problems, we can use more than one variable and a system of equations.

EXAMPLE 3 Solving a Mixture Problem

How many ounces each of 5% hydrochloric acid and 20% hydrochloric acid must be combined to obtain 10 oz of solution that is 12.5% hydrochloric acid?

Step 1 **Read** the problem. Two solutions of different strengths are being mixed to obtain a specific amount of a solution with an “in-between” strength.

Step 2 **Assign variables.**

Let x = the number of ounces of 5% solution

and y = the number of ounces of 20% solution.

Use a table to summarize the information from the problem.

Ounces of Solution	Percent (as a decimal)	Ounces of Pure Acid
x	5% = 0.05	$0.05x$
y	20% = 0.20	$0.20y$
10	12.5% = 0.125	$(0.125)10$

Gives equation (1)
Gives equation (2)

Multiply the amount of each solution (given in the first column) by its concentration of acid (given in the second column) to find the amount of acid in that solution (given in the third column).

FIGURE 10 illustrates what is happening in the problem.




FIGURE 10

Step 3 **Write a system of equations.** When x ounces of 5% solution and y ounces of 20% solution are combined, the total number of ounces is 10.

$$x + y = 10$$

The ounces of acid in the 5% solution ($0.05x$) added to the ounces of acid in the 20% solution ($0.20y$) must equal the total ounces of acid in the mixture, which is $(0.125)10$, or 1.25.

$$0.05x + 0.20y = 1.25$$

Notice that these equations can be quickly determined by reading down the table or using the labels in FIGURE 10.

Multiply the second equation by 100 to clear the decimals and obtain an equivalent system of equations.

$$\begin{array}{rcl}
 x + y = 10 & (1) & \text{This is the system} \\
 5x + 20y = 125 & (2) & \text{to solve.}
 \end{array}$$


NOW TRY EXERCISE 3

How many liters each of a 15% acid solution and a 25% acid solution should be mixed to obtain 30 L of an 18% acid solution?

Step 4 Solve the system. Again, either the substitution or the elimination method can be used. We choose to eliminate x .

$$-5x - 5y = -50 \quad \text{Multiply each side of (1) by } -5.$$

$$5x + 20y = 125 \quad (2)$$

$$15y = 75 \quad \text{Add.}$$

$$\begin{array}{l} \text{Ounces of} \\ \text{20\% solution} \end{array} \rightarrow y = 5 \quad \text{Divide by 15.}$$

Substitute 5 for y in equation (1) to find the value of x .

$$x + y = 10 \quad (1)$$

$$x + 5 = 10 \quad \text{Let } y = 5.$$

$$\begin{array}{l} \text{Ounces of} \\ \text{5\% solution} \end{array} \rightarrow x = 5 \quad \text{Subtract 5.}$$

Step 5 State the answer. The desired mixture will require 5 oz of the 5% solution and 5 oz of the 20% solution.

Step 6 Check.

$$\begin{aligned} \text{Total amount of solution:} \quad x + y &= 5 \text{ oz} + 5 \text{ oz} \\ &= 10 \text{ oz, as required.} \end{aligned}$$

$$\begin{aligned} \text{Total amount of acid:} \quad 5\% \text{ of } 5 \text{ oz} + 20\% \text{ of } 5 \text{ oz} \\ &= 0.05(5) + 0.20(5) \\ &= 1.25 \text{ oz} \end{aligned}$$

Percent of acid in solution:

$$\frac{\text{Total acid} \rightarrow 1.25}{\text{Total solution} \rightarrow 10} = 0.125, \text{ or } 12.5\%, \text{ as required.} \quad \text{NOW TRY } \img alt="Now Try icon" data-bbox="885 518 910 543"/>$$

OBJECTIVE 4 Solve distance-rate-time problems using two variables.

Motion problems require a form of the distance formula $d = rt$, where d is distance, r is rate (or speed), and t is time.

EXAMPLE 4 Solving a Motion Problem

A car travels 250 km in the same time that a truck travels 225 km. If the rate of the car is 8 km per hr faster than the rate of the truck, find both rates.

Step 1 Read the problem again. Given the distances traveled, we need to find the rate of each vehicle.

Step 2 Assign variables.

Let x = the rate of the car, and y = the rate of the truck.

As in **Example 3**, a table helps organize the information. Fill in the distance for each vehicle, and the variables for the unknown rates.

	d	r	t
Car	250	x	$\frac{250}{x}$
Truck	225	y	$\frac{225}{y}$

To find the expressions for time, we solved the distance formula $d = rt$ for t . Thus, $\frac{d}{r} = t$.

NOW TRY ANSWER

3. 25% acid solution: 9 L;
15% acid solution: 21 L


**NOW TRY
EXERCISE 4**

On a bicycle ride, Vann can travel 50 mi in the same amount of time that Ivy can travel 40 mi. Determine each bicyclist's rate, if Vann's rate is 2 mph faster than Ivy's.

Step 3 Write a system of equations. The rate x of the car is 8 km per hr faster than the rate y of the truck.

Be careful writing this equation. *The car is faster.* $x = y + 8$ (1)

Both vehicles travel for the *same* time, so the times must be equal.

$$\text{Time for car} \rightarrow \frac{250}{x} = \frac{225}{y} \leftarrow \text{Time for truck}$$

Multiply both sides by xy to obtain an equivalent equation with no variable denominators.

$$xy \cdot \frac{250}{x} = \frac{225}{y} \cdot xy \quad \text{Multiply by the LCD, } xy.$$

$$\frac{250xy}{x} = \frac{225xy}{y} \quad \text{Multiply; } xy = \frac{xy}{1}$$

$$250y = 225x \quad \text{Divide out the common factors. (2)}$$

We now have a system of linear equations.

$$x = y + 8 \quad (1)$$

$$250y = 225x \quad (2)$$

Step 4 Solve the system by substitution. Replace x with $y + 8$ in equation (2).

$$250y = 225x \quad (2)$$

Be sure to use parentheses around $y + 8$. $250y = 225(y + 8)$ Let $x = y + 8$.

$$250y = 225y + 1800 \quad \text{Distributive property}$$

$$25y = 1800 \quad \text{Subtract } 225y.$$

$$\text{Truck's rate} \rightarrow y = 72 \quad \text{Divide by } 25.$$

Let $y = 72$ in the equation $x = y + 8$ to find x .

$$\text{Car's rate} \rightarrow x = 72 + 8 = 80$$

Step 5 State the answer. The rate of the car is 80 km per hr, and the rate of the truck is 72 km per hr.

Step 6 Check.

$$\begin{array}{l} \text{Car: } t = \frac{d}{r} = \frac{250}{80} = 3.125 \\ \text{Truck: } t = \frac{d}{r} = \frac{225}{72} = 3.125 \end{array} \left\{ \begin{array}{l} \text{Times are equal,} \\ \text{as required.} \end{array} \right.$$

The rate of the car, 80 km per hr, is 8 km per hour greater than that of the truck, 72 km per hr, as required.

NOW TRY 

PROBLEM-SOLVING HINT When one quantity in an application is compared to another (as are the rates of the car and truck in **Example 4**), be careful to translate correctly in terms of the two variables. In Step 3 of **Example 4**,

$$y = x - 8 \text{ is an alternative for equation (1).}$$

NOW TRY ANSWER

4. Vann: 10 mph; Ivy: 8 mph

NOW TRY
EXERCISE 5

In his motorboat, Ed traveled 42 mi upstream at top speed in 2.1 hr. Still at top speed, his return trip to the same spot took only 1.5 hr. Find the rate of Ed's boat in still water and the rate of the current.




FIGURE 11

EXAMPLE 5 Solving a Motion Problem

While kayaking on the Blackledge River, Rebecca traveled 9 mi upstream (against the current) in 2.25 hr. It took her only 1 hr paddling downstream (with the current) back to the spot where she started. Find Rebecca's kayaking rate in still water and the rate of the current.

Step 1 **Read** the problem. We must find two rates—Rebecca's kayaking rate in still water and the rate of the current.

Step 2 **Assign variables.**

Let x = Rebecca's kayaking rate in still water
and y = the rate of the current.

When the kayak is traveling *against* the current, the current slows it down. The rate of the kayak is the *difference* between its rate in still water and the rate of the current, which is $(x - y)$ mph.

When the kayak is traveling *with* the current, the current speeds it up. The rate of the kayak is the *sum* of its rate in still water and the rate of the current, which is $(x + y)$ mph.

Thus, $x - y$ = the rate of the kayak upstream (*against* the current),
and $x + y$ = the rate of the kayak downstream (*with* the current).

See FIGURE 11. Make a table. Use the formula $d = rt$, or $rt = d$.

	r	t	d
Upstream	$x - y$	2.25	$2.25(x - y)$
Downstream	$x + y$	1	$1(x + y)$

← The distance is the same in each direction, 9 mi.

Step 3 **Write a system of equations.**

$$2.25(x - y) = 9 \quad \text{Upstream}$$

$$1(x + y) = 9 \quad \text{Downstream}$$

Clear parentheses in each equation, and then divide each term in the first equation by 2.25 to obtain an equivalent system.

$$x - y = 4 \quad (1)$$

$$x + y = 9 \quad (2)$$

Step 4 **Solve.**

$$2x = 13 \quad \text{Add.}$$

$$\text{Rebecca's kayaking rate} \rightarrow x = 6.5 \quad \text{Divide by 2.}$$

Substitute 6.5 for x in equation (2) and solve for y .

$$x + y = 9 \quad (2)$$

$$6.5 + y = 9 \quad \text{Let } x = 6.5.$$

$$\text{Rate of current} \rightarrow y = 2.5 \quad \text{Subtract 6.5.}$$

Step 5 **State the answer.** Rebecca's kayaking rate in still water was 6.5 mph, and the rate of the current was 2.5 mph.

Step 6 **Check.**

$$\text{Distance upstream: } 2.25(6.5 - 2.5) = 9 \quad \leftarrow \text{True statements}$$

$$\text{Distance downstream: } 1(6.5 + 2.5) = 9 \quad \leftarrow \text{result.}$$

NOW TRY

NOW TRY ANSWER

5. boat: 24 mph; current: 4 mph

OBJECTIVE 5 Solve problems with three variables using a system of three equations.

EXAMPLE 6 Solving a Geometry Problem

The sum of the measures of the angles of any triangle is 180° . The smallest angle of a certain triangle measures 36° less than the middle-sized angle. The largest angle measures 16° more than twice the smallest angle. Find the measure of each angle.

Step 1 Read the problem again. There are three unknowns in this problem.

Step 2 Assign variables. Make a sketch, as in FIGURE 12.

Let x = the measure of the smallest angle,

y = the measure of the middle-sized angle,

and z = the measure of the largest angle.




FIGURE 12

Step 3 Write a system of three equations. The sum of the measures of the angles of any triangle is 180° .

$$x + y + z = 180$$

We can write two more equations from the information given.

Smallest angle	measures	36° less than middle-sized angle.
↓	↓	↓
x	=	$y - 36$

Largest angle	measures	16° more than twice smallest angle.
↓	↓	↓
z	=	$16 + 2x$

These three equations form a system.

$$x + y + z = 180 \quad (1)$$

$$x = y - 36 \quad (2)$$

$$z = 16 + 2x \quad (3)$$

Step 4 Solve the system of equations. If we write equation (2) in terms of y , then equations (2) and (3) will give both y and z in terms of x .

$$x = y - 36 \quad (2)$$

$$y = x + 36 \quad \text{Add 36. Interchange sides.}$$

Now substitute $x + 36$ for y and $16 + 2x$ (from equation (3)) for z in equation (1).

$$x + y + z = 180 \quad (1)$$

$$x + (x + 36) + (16 + 2x) = 180 \quad \text{Let } y = x + 36 \text{ and } z = 16 + 2x.$$

$$4x + 52 = 180 \quad \text{Combine like terms.}$$

$$x = 32 \quad \text{Subtract 52. Divide by 4.}$$

Substitute 32 for x in $y = x + 36$ (equation (2) solved for y) to find y .

$$y = 32 + 36 \quad \text{Let } x = 32.$$

$$y = 68 \quad \text{Add.}$$

Substitute 32 for x in $z = 16 + 2x$ (equation (3)) to find z .

$$z = 16 + 2(32) \quad \text{Let } x = 32.$$

$$z = 80 \quad \text{Multiply, and then add.}$$


NOW TRY EXERCISE 6

The sum of the measures of the angles of any triangle is 180° . The measure of the third angle of a certain triangle is four times that of the first angle. The sum of the measures of the first and second angles is 84° . Find the measure of each angle.

Step 5 State the answer. The three angles measure 32° , 68° , and 80° .

Step 6 Check. The sum of the measures of the three angles is

$$32^\circ + 68^\circ + 80^\circ = 180^\circ, \text{ as required.}$$

Also, each of the following is a true statement.

$$32^\circ = 68^\circ - 36^\circ \quad \text{Smallest measures } 36^\circ \text{ less than middle-sized.}$$

$$80^\circ = 16^\circ + 2(32^\circ) \quad \text{Largest measures } 16^\circ \text{ more than twice smallest.}$$

NOW TRY 

PROBLEM-SOLVING HINT In Step 4 of **Example 6**, we could also have substituted $16 + 2x$ (from equation (3)) for z in equation (1) as follows.

$$x + y + z = 180 \quad (1)$$

$$x + y + (16 + 2x) = 180 \quad \text{Let } z = 16 + 2x.$$

$$3x + y = 164 \quad \text{Combine like terms. Subtract 16.} \quad (4)$$

This gives a system of two equations in x and y .

$$x - y = -36 \quad (2) \text{ in standard form}$$

$$3x + y = 164 \quad (4)$$

Eliminating y gives $x = 32$, which can be used to obtain $y = 68$ and $z = 80$. There is often more than one way to solve applications involving systems of three equations.


EXAMPLE 7 Solving a Problem Involving Prices

At Panera Bread, a loaf of honey wheat bread costs \$3.99, a loaf of tomato basil bread costs \$4.99, and a loaf of French bread costs \$3.19. On a recent day, three times as many loaves of honey wheat were sold as loaves of tomato basil. The number of loaves of French bread sold was 5 less than the number of loaves of honey wheat sold. Total receipts for these breads were \$90.17. How many loaves of each type of bread were sold? (Data from Panera Bread menu.)

Step 1 Read the problem again. There are three unknowns in this problem.

Step 2 Assign variables to represent the three unknowns.

Let x = the number of loaves of honey wheat bread,

y = the number of loaves of tomato basil bread,

and z = the number of loaves of French bread.

Step 3 Write a system of three equations. Three times as many loaves of honey wheat bread were sold as loaves of tomato basil bread.

$$x = 3y, \text{ or } x - 3y = 0 \quad \text{Subtract } 3y. \quad (1)$$

Also, we have the information needed for another equation.

Number of loaves of French	was (equals)	5 less than the number of loaves of honey wheat.
↓	↓	↓
z	=	$x - 5$

$$-x + z = -5 \quad \text{Subtract } x.$$

$$x - z = 5 \quad \text{Multiply by } -1. \quad (2)$$

NOW TRY ANSWER

6. 24° , 60° , 96°


**NOW TRY
EXERCISE 7**

At Panera Bread, a loaf of white bread costs \$3.99, a loaf of cheese bread costs \$4.79, and a loaf of sesame semolina bread costs \$8.19. On a recent day, twice as many loaves of white bread were sold as loaves of cheese bread. The number of loaves of sesame semolina sold was 3 less than the number of loaves of white sold. Total receipts for these breads were \$150.33. How many loaves of each type of bread were sold? (Data from Panera Bread menu.)

Multiplying the cost of a loaf of each kind of bread by the number of loaves of that kind sold and adding gives an equation for the total receipts.

$$\begin{aligned} 3.99x + 4.99y + 3.19z &= 90.17 \\ 399x + 499y + 319z &= 9017 \end{aligned} \quad \begin{array}{l} \text{Multiply each term by} \\ \text{100 to clear decimals. (3)} \end{array}$$

These three equations form a system.

$$x - 3y = 0 \quad (1)$$

$$x - z = 5 \quad (2)$$

$$399x + 499y + 319z = 9017 \quad (3)$$

Step 4 **Solve** the system of equations. Equation (1) is missing the variable z , so one way to begin is to eliminate z again, using equations (2) and (3).

$$\begin{array}{r} 319x \quad \quad - 319z = 1595 \quad \text{Multiply (2) by 319.} \\ \underline{399x + 499y + 319z = 9017} \quad (3) \\ 718x + 499y \quad \quad = 10,612 \quad \text{Add. (4)} \end{array}$$

Use resulting equation (4) in x and y , together with equation (1), $x - 3y = 0$, to eliminate x .

$$\begin{array}{r} -718x + 2154y = 0 \quad \text{Multiply (1) by } -718. \\ \underline{718x + 499y = 10,612} \quad (4) \\ 2653y = 10,612 \quad \text{Add.} \\ y = 4 \quad \text{Divide by 2653.} \end{array}$$

We can find x by substituting this value for y in equation (1).

$$\begin{aligned} x - 3y &= 0 \quad (1) \\ x - 3(4) &= 0 \quad \text{Let } y = 4. \\ x - 12 &= 0 \quad \text{Multiply.} \\ x &= 12 \quad \text{Add 12.} \end{aligned}$$

We can find z by substituting this value for x in equation (2).

$$\begin{aligned} x - z &= 5 \quad (2) \\ 12 - z &= 5 \quad \text{Let } x = 12. \\ z &= 7 \quad \text{Subtract 12. Multiply by } -1. \end{aligned}$$

Step 5 **State the answer.** There were 12 loaves of honey wheat bread, 4 loaves of tomato basil bread, and 7 loaves of French bread sold.

Step 6 **Check.** Each of the following is a true statement.

$$\begin{aligned} 12 &= 3 \cdot 4 \quad \text{Honey wheat is three times tomato basil.} \\ 7 &= 12 - 5 \quad \text{French is 5 less than honey wheat.} \end{aligned}$$

Multiply cost per loaf by number of loaves and add to confirm total receipts.

$$\$3.99(12) + \$4.99(4) + \$3.19(7) = \$90.17, \text{ as required.}$$

NOW TRY ANSWER

7. white bread: 12 loaves;
cheese bread: 6 loaves;
sesame semolina bread: 9 loaves

3.3 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Time management can be a challenge for students.

Review Study Skill 6, Managing Your Time.


Concept Check Answer each question.

- If a container of liquid contains 60 oz of solution, what is the number of ounces of pure acid if the given solution contains the following acid concentrations?
 - 10%
 - 25%
 - 40%
 - 50%
- If \$5000 is invested in an account paying simple annual interest, how much interest will be earned during the first year at the following rates?
 - 2%
 - 3%
 - 4%
 - 3.5%
- If one pound of turkey costs \$1.99, how much will x pounds cost?
- If one movie ticket costs \$13.50, how much will y tickets cost?
- If the rate of a boat in still water is 10 mph and the rate of the current of a river is x mph, what is the rate of the boat in each case?
 - The boat is going upstream (that is, against the current, which slows the boat down).
 - The boat is going downstream (that is, with the current, which speeds the boat up).
- If the rate of a plane in still air is x mph and the rate of a steady wind is 20 mph, what is the rate of the plane in each case?
 - The plane is flying into the wind (that is, into a headwind, which slows the plane down).
 - The plane is flying with the wind (that is, with a tailwind, which speeds the plane up).
- A bear is running at a rate of 44 ft per sec.
 - If the bear runs for 5 sec, what is its distance?
 - If the bear travels d ft, what is its time?
- The swimming rate of a whale is 25 mph.
 - If the whale swims for y hours, what is its distance?
 - If the whale travels 10 mi, what is its time?




Solve each problem. See Example 1.

- Venus and Serena measured a tennis court and found that it was 42 ft longer than it was wide and had a perimeter of 228 ft. What were the length and the width of the tennis court?



- LeBron and Jose measured a basketball court and found that the width of the court was 44 ft less than the length. If the perimeter was 288 ft, what were the length and the width of the basketball court?





11. During the 2017 Major League Baseball season, the Cleveland Indians played 162 games. They won 42 more games than they lost. What was the team's win-loss record that year?
12. During the 2017 Major League Baseball season, the Detroit Tigers played 162 games. They lost 34 more games than they won. What was the team's win-loss record?
13. In 2016, the two American telecommunication companies with the greatest revenues were AT&T and Verizon. The two companies had combined revenues of \$288.9 billion. AT&T's revenue was \$38.7 billion more than that of Verizon. What was the revenue for each company? (Data from Verizon and AT&T Annual Reports.)
14. In 2017, U.S. exports to Canada were \$39.4 billion more than exports to Mexico. Together, exports to these two countries totaled \$525.4 billion. How much were exports to each country? (Data from U.S. Census Bureau.)

Team	W	L
Cleveland	_____	_____
Minnesota	85	77
Kansas City	80	82
Chicago White Sox	67	95
Detroit	_____	_____

Data from Major League Baseball.

Find the measures of angles x and y . Remember that (1) the sum of the measures of the angles of any triangle is 180° , (2) supplementary angles have a sum of 180° , and (3) vertical angles have equal measures.



The Fan Cost Index (FCI) represents the cost of four average-price tickets to a sporting event (two adult, two child), four small soft drinks, two small beers, four hot dogs, parking for one car, two game programs, and two souvenir caps. (Data from Team Marketing Report.)

Use the concept of FCI to solve each problem. See Example 2.

17. For the 2016 Major League Baseball season, the FCI prices for the Cleveland Indians and the Boston Red Sox totaled \$540.10. The Boston FCI was \$181.22 more than that of Cleveland. What were the FCIs for these teams?
18. In 2016, the FCI prices for Major League Baseball and the National Football League totaled \$722.37. The football FCI was \$283.31 more than that of baseball. What were the FCIs for these sports?



Solve each problem. See Example 2.

19. Leanna is a waitress at Bonefish Grill. During one particular day she sold 15 ribeye steak dinners and 20 grilled salmon dinners, totaling \$886.50. Another day she sold 25 ribeye steak dinners and 10 grilled salmon dinners, totaling \$973.50. How much did each type of dinner cost? (Data from Bonefish Grill Menu.)
20. At a business meeting at Panera Bread, the bill for two cappuccinos and three caffe lattes was \$19.75. At another table, the bill for one cappuccino and two caffe lattes was \$11.97. How much did each type of beverage cost? (Data from Panera Bread menu.)



21. Two days at Busch Gardens (Tampa Bay) and 3 days at Universal Studios Florida (Orlando) cost \$510, while 4 days at Busch Gardens and 2 days at Universal Studios cost \$580. (Prices are based on single-day admissions.) What was the cost per day for each park? (Data from Busch Gardens and Universal Studios.)
22. On the basis of average total costs per day for business travel to New York City and Washington, DC (which include a hotel room, car rental, and three meals), 2 days in New York and 3 days in Washington cost \$2484, while 4 days in New York and 2 days in Washington cost \$3120. What was the average cost per day in each city? (Data from *Business Travel News*.)
23. Tickets to a production of *A Midsummer Night's Dream* at Broward College cost \$5 for general admission or \$4 with a student ID. If 184 people paid to see a performance and \$812 was collected, how many of each type of ticket were sold?
24. The mixed price of 9 citrons and 7 fragrant wood apples is 107; again, the mixed price of 7 citrons and 9 fragrant wood apples is 101. O you arithmetician, tell me quickly the price of a citron and the price of a wood apple here, having distinctly separated those prices well. (Source: Hindu work, A.D. 850.)

Solve each problem. See Example 3.

25. How many gallons each of 25% alcohol and 35% alcohol should be mixed to obtain 20 gal of 32% alcohol?
26. How many liters each of 15% acid and 33% acid should be mixed to obtain 120 L of 21% acid?

Gallons of Solution	Percent (as a decimal)	Gallons of Pure Alcohol
x	25% = 0.25	
y	35% = 0.35	
20	32% =	

Liters of Solution	Percent (as a decimal)	Liters of Pure Acid
x	15% = 0.15	
y	33% =	
120	21% =	

27. A party mix is made by adding nuts that sell for \$2.50 per kg to a cereal mixture that sells for \$1 per kg. How much of each should be added to obtain 30 kg of a mix that will sell for \$1.70 per kg?
28. A fruit drink is made by mixing juices. Such a drink with 50% juice is to be mixed with a drink that is 30% juice to obtain 200 L of a drink that is 45% juice. How much of each should be used?

	Number of Kilograms	Price per Kilogram (in dollars)	Value (in dollars)
Nuts	x	2.50	
Cereal	y	1.00	
Mixture		1.70	

	Liters of Drink	Percent (as a decimal)	Liters of Pure Juice
50% Juice	x	0.50	
30% Juice	y	0.30	
Mixture		0.45	

29. A total of \$3000 is invested, part at 2% simple interest and part at 4%. If the total annual return from the two investments is \$100, how much is invested at each rate?
30. An investor will invest a total of \$15,000 in two accounts, one paying 4% annual simple interest and the other 3%. If he wants to earn \$550 annual interest, how much should he invest at each rate?

Principal (in dollars)	Rate (as a decimal)	Interest (in dollars)
x	0.02	0.02x
y	0.04	0.04y
3000	XXXXXXXXXX	100

Principal (in dollars)	Rate (as a decimal)	Interest (in dollars)
x	0.04	
y	0.03	
15,000	XXXXXXXXXX	

31. Pure acid is to be added to a 10% acid solution to obtain 54 L of a 20% acid solution. What amounts of each should be used? (*Hint*: Pure acid is 100% acid.)
32. A truck radiator holds 36 L of fluid. How much pure antifreeze must be added to a mixture that is 4% antifreeze to fill the radiator with a mixture that is 20% antifreeze?
33. How many pounds of candy that sells for \$1.75 per lb must be mixed with candy that sells for \$1.25 per lb to obtain 10 lb of a mixture that should sell for \$1.60 per lb?
34. How many pounds of candy that sells for \$2.50 per lb must be mixed with candy that sells for \$1.75 per lb to obtain 6 lb of a mixture that sells for \$2.10 per lb?



Solve each problem. See Examples 4 and 5.

35. A train travels 150 km in the same time that a plane travels 400 km. If the rate of the plane is 20 km per hr less than three times the rate of the train, find both rates.

	r	t	d
Train	x		150
Plane	y		400


36. A motor scooter travels 20 mi in the same time that a bicycle travels 8 mi. If the rate of the scooter is 5 mph more than twice the rate of the bicycle, find both rates.
37. A freight train and an express train leave towns 390 km apart, traveling toward one another. The freight train travels 30 km per hr slower than the express train. They pass one another 3 hr later. What are their rates?

	r	t	d
Freight Train	x	3	
Express Train	y	3	

38. A car and a truck leave towns 230 mi apart, traveling toward each other. The car travels 15 mph faster than the truck. They pass each other 2 hr later. What are their rates?
39. In his motorboat, Bill travels upstream at top speed to his favorite fishing spot, a distance of 36 mi, in 2 hr. Returning, he finds that the trip downstream, still at top speed, takes only 1.5 hr. Find the rate of Bill's boat and the rate of the current. Let x = the rate of the boat and y = the rate of the current.

	r	t	d
Upstream	$x - y$	2	
Downstream	$x + y$		


40. Traveling for 3 hr into a steady headwind, a plane flies 1650 mi. The pilot determines that flying *with* the same wind for 2 hr, he could make a trip of 1300 mi. Find the rate of the plane and the wind speed.




41. A plane flies 560 mi in 1.75 hr traveling with the wind. The return trip later against the same wind takes the plane 2 hr. Find the rate of the plane and the wind speed.
42. Braving blizzard conditions on the planet Hoth, Luke Skywalker sets out in his snow speeder for a rebel base 4800 mi away. He travels into a steady headwind and makes the trip in 3 hr. Returning, he finds that the trip back, now with a tailwind, takes only 2 hr. Find the rate of Luke's snow speeder and the wind speed.

Solve each problem. See Examples 6 and 7.


43. In the triangle below, $z = x + 10$ and $x + y = 100$. Determine a third equation involving x , y , and z , and then find the measures of the three angles.




44. In the triangle below, x is 10 less than y and 20 less than z . Write a system of equations and find the measures of the three angles.



45. In a certain triangle, the measure of the second angle is 10° greater than three times the first. The third angle measure is equal to the sum of the measures of the other two. Find the measures of the three angles.
46. The measure of the largest angle of a triangle is 12° less than the sum of the measures of the other two. The smallest angle measures 58° less than the largest. Find the measures of the angles.
47. The perimeter of a triangle is 70 cm. The longest side is 4 cm less than the sum of the other two sides. Twice the shortest side is 9 cm less than the longest side. Find the length of each side of the triangle.




48. The perimeter of a triangle is 56 in. The longest side measures 4 in. less than the sum of the other two sides. Three times the shortest side is 4 in. more than the longest side. Find the lengths of the three sides.



49. In the 2016 Summer Olympics, host Brazil earned 1 more gold medal than silver. The number of silver medals that Brazil earned was the same as the number of its bronze medals. Brazil earned a total of 19 medals. How many of each kind of medal did Brazil earn? (Data from *The World Almanac and Book of Facts*.)
50. In 2017, the average Facebook user in the United States spent 693 more minutes per month using Facebook than the average Instagram user spent using Instagram. The average Instagram user spent 36 fewer minutes per month using Instagram than the average Snapchat user spent using Snapchat. The total amount of time that average users of each social network spent per month using their respective networks was 1347 min. How much time did the average user spend on each social network? (Data from Verito Analytics.)



51. Tickets for a Harlem Globetrotters show cost \$30 for upper level, \$51 for center court, or \$76 for floor seats. Nine times as many upper level tickets were sold as floor tickets, and the number of upper level tickets sold was 55 more than the sum of the number of center court tickets and floor tickets. Sales of all three kinds of tickets totaled \$95,215. How many of each kind of ticket were sold? (Data from www.harlemglobetrotters.com)



52. Three kinds of tickets are available for a rock concert: “up close,” “in the middle,” and “far out.” “Up close” tickets cost \$10 more than “in the middle” tickets. “In the middle” tickets cost \$10 more than “far out” tickets. Twice the cost of an “up close” ticket is \$20 more than three times the cost of a “far out” ticket. Find the price of each kind of ticket.
53. A wholesaler supplies college t-shirts to three college bookstores: A, B, and C. The wholesaler recently shipped a total of 800 t-shirts to the three bookstores. Twice as many t-shirts were shipped to bookstore B as to bookstore A, and the number shipped to bookstore C was 40 less than the sum of the numbers shipped to the other two bookstores. How many t-shirts were shipped to each bookstore?
54. An office supply store sells three models of computer desks: A, B, and C. In one month, the store sold a total of 85 computer desks. The number of model B desks was five more than the number of model C desks. The number of model A desks was four more than twice the number of model C desks. How many of each model did the store sell that month?
55. A plant food is to be made from three chemicals. The mix must include 60% of the first and second chemicals. The second and third chemicals must be in the ratio of 4 to 3 by weight. How much of each chemical is needed to make 750 kg of the plant food?
56. How many ounces of 5% hydrochloric acid, 20% hydrochloric acid, and water must be combined to obtain 10 oz of solution that is 8.5% hydrochloric acid if the amount of water used must equal the total amount of the other two solutions?

The National Hockey League uses a point system to determine team standings. A team is awarded 2 points for a win (W), 0 points for a loss in regulation play (L), and 1 point for an overtime loss (OTL). Use this information to solve each problem.


Team	GP	W	L	OTL	Points
Anaheim	82	—	—	—	105
Edmonton	82	47	26	9	103
San Jose	82	46	29	7	99
Calgary	82	45	33	4	94
Los Angeles	82	—	—	—	86

Data from www.nhl.com

57. During the 2016–2017 NHL regular season, the Anaheim Ducks played 82 games. Their wins and overtime losses resulted in a total of 105 points. They had 10 more losses in regulation play than overtimes losses. How many wins, losses, and overtime losses did they have that season?
58. During the same NHL regular season, the Los Angeles Kings also played 82 games. Their wins and overtimes losses resulted in a total of 86 points. They had 4 more total losses (in regulation play and overtime) than wins. How many wins, losses, and overtime losses did they have that season?

The following exercises are based on the “Bait Box Puzzle,” featured in an “Ask Marilyn” column in *Parade Magazine*.

59. There are three kinds of bait boxes—those containing fish, those containing bugs, and those containing worms. Although the bait boxes look alike, each kind has a different weight. Use the information in **FIGURE A** to write a system of three equations, and solve it to determine the weight of each kind of bait box.



60. Write an equation that describes the situation depicted in **FIGURE B**. Use the same three variables used in **Exercise 59**. Then determine the total weight of the bait boxes in **FIGURE B**.

Chapter 3

Summary

STUDY SKILLS REMINDER

Be prepared for your math test on this chapter. **Review Study Skills 7 and 8, Reviewing a Chapter and Taking Math Tests.**

Key Terms

3.1

system of equations
system of linear equations
(linear system)
solution set of a linear
system

consistent system
independent equations
inconsistent system
dependent equations

3.2

ordered triple
focus variable
working equation

New Symbol

(x, y, z) ordered triple

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- A **system of equations** consists of
 - at least two equations with different variables
 - two or more equations that have an infinite number of solutions
 - two or more equations with the same variables
 - two or more inequalities that are to be solved.
- The **solution set of a system of equations** is
 - all ordered pairs that satisfy one equation of the system
 - all ordered pairs that satisfy all the equations of the system
 - any ordered pair that satisfies one or more equations of the system
 - the set of values that make all the equations of the system false.
- A **consistent system** has
 - no solution
 - a solution
 - exactly two solutions
 - exactly three solutions.

4. An **inconsistent system** is a system of equations
- with one solution
 - with no solution
 - with an infinite number of solutions
 - that have the same graph.
5. **Dependent equations**
- have different graphs
 - have no solution
 - have one solution
 - are different forms of the same equation.
6. **Independent equations**
- have different graphs
 - have no solution
 - have graphs that are parallel lines
 - are different forms of the same equation.

ANSWERS

1. C; *Example:* $\begin{cases} 3x - y = 3 \\ 2x + y = 7 \end{cases}$ 2. B; *Example:* The ordered pair (2, 3) satisfies both equations of the system in Answer 1, so $\{(2, 3)\}$ is the solution set of the system. 3. B; *Example:* The system in Answer 1 is a consistent system. 4. B; *Example:* The equations of two parallel lines form an inconsistent system. Their graphs never intersect, so the system has no solution. 5. D; *Example:* The equations $4x - y = 8$ and $8x - 2y = 16$ are dependent because their graphs are the same line. 6. A; *Example:* The equations in the system in Answer 1 are not equivalent. They are independent equations with different graphs.

Quick Review

CONCEPTS	EXAMPLES
<p>3.1 Systems of Linear Equations in Two Variables</p> <p>Solving a Linear System by Substitution</p> <p><i>Step 1</i> Solve one of the equations for either variable.</p> <p><i>Step 2</i> Substitute for that variable in the other equation. The result should be an equation with just one variable.</p> <p><i>Step 3</i> Solve the equation from Step 2.</p> <p><i>Step 4</i> Find the value of the other variable by substituting the result from Step 3 into the equation from Step 1.</p> <p><i>Step 5</i> Check the values in <i>both</i> of the <i>original</i> equations. Then write the solution set as a set containing an ordered pair.</p> <p>Solving a Linear System by Elimination</p> <p><i>Step 1</i> Write both equations in standard form.</p> <p><i>Step 2</i> Transform the equations as needed so that the coefficients of one pair of variable terms are opposites.</p> <p><i>Step 3</i> Add the new equations. The sum should be an equation with just one variable.</p> <p><i>Step 4</i> Solve the equation from Step 3.</p> <p><i>Step 5</i> Find the value of the other variable by substituting the result from Step 4 into either of the original equations.</p> <p><i>Step 6</i> Check the values in <i>both</i> of the <i>original</i> equations. Then write the solution set as a set containing an ordered pair.</p>	<p>Solve by substitution.</p> $\begin{aligned} 4x - y &= 7 & (1) \\ 3x + 2y &= 30 & (2) \end{aligned}$ <p>Solve for y in equation (1).</p> $y = 4x - 7$ <p>Substitute $4x - 7$ for y in equation (2), and solve for x.</p> $\begin{aligned} 3x + 2y &= 30 & (2) \\ 3x + 2(4x - 7) &= 30 & \text{Let } y = 4x - 7. \\ 3x + 8x - 14 &= 30 & \text{Distributive property} \\ 11x - 14 &= 30 & \text{Combine like terms.} \\ 11x &= 44 & \text{Add 14.} \\ x &= 4 & \text{Divide by 11.} \end{aligned}$ <p>Substitute 4 for x in the equation $y = 4x - 7$ to find y.</p> $\begin{aligned} y &= 4(4) - 7 \\ y &= 9 & \text{Multiply. Subtract.} \end{aligned}$ <p>Check to verify that $\{(4, 9)\}$ is the solution set.</p> <p>Solve by elimination.</p> $\begin{aligned} 5x + y &= 2 & (1) \\ 2x - 3y &= 11 & (2) \end{aligned}$ <p>To eliminate y, multiply equation (1) by 3, and add the result to equation (2).</p> $\begin{aligned} 15x + 3y &= 6 & \text{3 times equation (1)} \\ \underline{2x - 3y} &= 11 & (2) \\ 17x &= 17 & \text{Add.} \\ x &= 1 & \text{Divide by 17.} \end{aligned}$

CONCEPTS

Solving Special Systems

If the result of the addition step (Step 3) is a false statement, such as $0 = 4$, the graphs are parallel lines and *there is no solution*. The solution set is \emptyset .

If the result is a true statement, such as $0 = 0$, the graphs are the same line, and *an infinite number of ordered pairs are solutions*. The solution set is written in set-builder notation as

$$\{(x, y) \mid \text{—————}\},$$

where a form of the equation is written in the blank.

3.2 Systems of Linear Equations in Three Variables

Solving a Linear System in Three Variables

- Step 1** Select a focus variable, preferably one with coefficient 1 or -1 , and a working equation.
- Step 2** Eliminate the focus variable, using the working equation and one of the equations of the system.
- Step 3** Eliminate the focus variable again, using the working equation and the remaining equation of the system.
- Step 4** Solve the system of two equations in two variables formed by the equations from Steps 2 and 3.
- Step 5** Find the value of the remaining variable.
- Step 6** Check the values in *each* of the *original* equations of the system. Then write the solution set as a set containing an ordered triple.

EXAMPLES

Let $x = 1$ in equation (1), and solve for y .

$$5x + y = 2 \quad (1)$$

$$5(1) + y = 2 \quad \text{Let } x = 1.$$

$$y = -3 \quad \text{Multiply. Subtract 5.}$$

Check to verify that $\{(1, -3)\}$ is the solution set.

Solve each system.

$$\begin{array}{r} x - 2y = 6 \\ -x + 2y = -2 \\ \hline 0 = 4 \end{array} \quad \begin{array}{l} \text{Inconsistent system} \\ \text{Solution set: } \emptyset \end{array}$$

$$\begin{array}{r} x - 2y = 6 \\ -x + 2y = -6 \\ \hline 0 = 0 \end{array} \quad \begin{array}{l} \text{Dependent equations} \\ \text{Solution set: } \{(x, y) \mid x - 2y = 6\} \end{array}$$

Solve the system.

$$x + 2y - z = 6 \quad (1)$$

$$x + y + z = 6 \quad (2)$$

$$2x + y - z = 7 \quad (3)$$

We choose z as the focus variable and (2) as the working equation.

Add equations (1) and (2).

$$2x + 3y = 12 \quad (4)$$

Add equations (2) and (3).

$$3x + 2y = 13 \quad (5)$$

Use equations (4) and (5) to eliminate x .

$$-6x - 9y = -36 \quad \text{Multiply (4) by } -3.$$

$$6x + 4y = 26 \quad \text{Multiply (5) by 2.}$$

$$-5y = -10 \quad \text{Add.}$$

$$y = 2 \quad \text{Divide by } -5.$$

To find x , substitute 2 for y in equation (4).

$$2x + 3(2) = 12 \quad \text{Let } y = 2 \text{ in (4).}$$

$$2x + 6 = 12 \quad \text{Multiply.}$$

$$2x = 6 \quad \text{Subtract 6.}$$

$$x = 3 \quad \text{Divide by 2.}$$

Substitute 3 for x and 2 for y in working equation (2).

$$x + y + z = 6 \quad (2)$$

$$3 + 2 + z = 6 \quad \text{Substitute.}$$

$$z = 1 \quad \text{Subtract 5.}$$

Check to verify that $\{(3, 2, 1)\}$ is the solution set.

CONCEPTS

3.3 Applications of Systems of Linear Equations

Use the six-step problem-solving method.

Step 1 Read the problem carefully.

Step 2 Assign variables.

Step 3 Write a system of equations.

Step 4 Solve the system.

Step 5 State the answer.

Step 6 Check.

EXAMPLES

The perimeter of a rectangle is 18 ft. The length is 3 ft more than twice the width. What are the dimensions of the rectangle?

Let x = the length and y = the width.

Write a system of equations.

$$2x + 2y = 18 \quad (1) \text{ From the perimeter formula}$$

$$x = 2y + 3 \quad (2) \text{ Length is 3 ft more than twice the width.}$$

Substitute $2y + 3$ for x in equation (1), and solve for y .

$$2x + 2y = 18 \quad (1)$$

$$2(2y + 3) + 2y = 18 \quad \text{Let } x = 2y + 3.$$

$$4y + 6 + 2y = 18 \quad \text{Distributive property}$$

$$6y + 6 = 18 \quad \text{Combine like terms.}$$

$$6y = 12 \quad \text{Subtract 6.}$$

$$y = 2 \quad \text{Divide by 6.}$$

To find x , substitute 2 for y in equation (2).

$$x = 2y + 3 \quad (2)$$

$$x = 2(2) + 3 \quad \text{Let } y = 2.$$

$$x = 7 \quad \text{Multiply, and then add.}$$

The length is 7 ft, and the width is 2 ft. The answer checks because

$$2(7) + 2(2) = 18 \quad \text{and} \quad 2(2) + 3 = 7, \quad \text{as required.}$$

Chapter 3

Review Exercises

3.1 Solve each system by graphing.

1. $x + 3y = 8$
 $2x - y = 2$

2. $2x + y = -8$
 $x - 4y = -4$

3. $x - y = 2$
 $x + 2y = -1$

4. Which ordered pair is *not* a solution of the equation $3x + 2y = 6$?

- A. (2, 0) B. (0, 3) C. (4, -3) D. (3, -2)

Solve each system using the substitution method. If a system is inconsistent or has dependent equations, say so.

5. $3x + y = -4$
 $x = \frac{2}{3}y$

6. $9x - y = -4$
 $y = x + 4$

7. $-5x + 2y = -2$
 $x + 6y = 26$

8. $5x + y = 12$
 $2x - 2y = 0$

Solve each system using the elimination method. If a system is inconsistent or has dependent equations, say so.

9. $5x - y = 14$
 $4x + 2y = 0$

10. $x - 4y = -4$
 $3x + y = 1$

11. $6x + 5y = 4$
 $-4x + 2y = 8$

$$12. \begin{cases} \frac{1}{6}x + \frac{1}{6}y = -\frac{1}{2} \\ x - y = -9 \end{cases}$$

$$13. \begin{cases} -3x + y = 6 \\ y = 6 + 3x \end{cases}$$

$$14. \begin{cases} 5x - 4y = 2 \\ -10x + 8y = 7 \end{cases}$$


3.2 Solve each system. If a system is inconsistent or has dependent equations, say so.

$$15. \begin{cases} 2x + 3y - z = -16 \\ x + 2y + 2z = -3 \\ -3x + y + z = -5 \end{cases} \quad 16. \begin{cases} 3x - y - z = -8 \\ 4x + 2y + 3z = 15 \\ -6x + 2y + 2z = 10 \end{cases} \quad 17. \begin{cases} 4x - y = 2 \\ 3y + z = 9 \\ x + 2z = 7 \end{cases}$$

$$18. \begin{cases} 5x - y = 26 \\ 4y + 3z = -4 \\ x + z = 5 \end{cases} \quad 19. \begin{cases} 3x - 4y + z = 8 \\ -6x + 8y - 2z = -16 \\ \frac{3}{2}x - 2y + \frac{1}{2}z = 4 \end{cases} \quad 20. \begin{cases} 2x - y + 3z = 0 \\ 5x + y - z = 0 \\ -2x + 3y + 4z = 0 \end{cases}$$

3.3 Solve each problem.

21. A regulation National Hockey League ice rink has perimeter 570 ft. The length of the rink is 30 ft longer than twice the width. What are the dimensions of an NHL ice rink? (Data from www.nhl.com)



22. A plane flies 560 mi in 1.75 hr traveling with the wind. The return trip later against the same wind takes the plane 2 hr. Find the rate of the plane and the speed of the wind. Let x = the rate of the plane and y = the speed of the wind.


	r	t	d
With Wind	$x + y$	1.75	
Against Wind		2	

23. In the 2016 Summer Olympics in Rio de Janeiro, Japan earned 9 fewer gold medals than bronze. The number of silver medals earned was 34 less than twice the number of bronze medals. Japan earned a total of 41 medals. How many of each kind of medal did Japan earn? (Data from *The World Almanac and Book of Facts*.)
24. For Valentine's Day, Ms. Sweet will mix some \$6-per-lb nuts with some \$3-per-lb chocolate candy to obtain 100 lb of mix, which she will sell at \$3.90 per lb. How many pounds of each should she use?

	Number of Pounds	Price per Pound (in dollars)	Value (in dollars)
Nuts	x		
Chocolate	y		
Mixture	100		

25. How many liters each of 8% and 20% hydrogen peroxide should be mixed together to obtain 8 L of 12.5% solution?

26. The sum of the measures of the angles of any triangle is 180° . The largest angle measures 10° less than the sum of the other two. The measure of the middle-sized angle is the average of the other two. Find the measures of the three angles.



27. Noemi sells real estate. On three recent sales, she made 10% commission, 6% commission, and 5% commission. Her total commissions on these sales were \$17,000, and she sold property worth \$280,000. If the 5% sale amounted to the sum of the other two, what were the three sales prices?
28. In the great baseball year of 1961, Yankee teammates Mickey Mantle, Roger Maris, and Yogi Berra combined for 137 home runs. Mantle hit 7 fewer than Maris, and Maris hit 39 more than Berra. What were the home run totals for each player? (Data from www.mlb.com)

Chapter 3

Mixed Review Exercises

1. Suppose that two linear equations are graphed on the same set of coordinate axes. Sketch what the graph might look like if the system has the given description.
- (a) The system has a single solution.
- (b) The system has no solution.
- (c) The system has infinitely many solutions.
2. Without doing any algebraic work, but answering on the basis of knowledge of the graphs of the two lines, explain why the following system has \emptyset as its solution set.

$$y = 3x + 2$$

$$y = 3x - 4$$

Solve each system.

3. $-7x + 3y = 12$

$$5x + 2y = 8$$

4. $2x - 5y = 8$

$$3x + 4y = 10$$

5. $x + 2y = 48$

$$\frac{1}{4}x + \frac{1}{2}y = 12$$

6. $x + 4y = 17$

$$-3x + 2y = -9$$

7. $\frac{2}{3}x + \frac{1}{6}y = \frac{19}{2}$

$$\frac{1}{3}x - \frac{2}{9}y = 2$$

8. $x = 7y + 10$

$$2x + 3y = 3$$

9. $x + y - z = 0$

$$2y - z = 1$$

$$2x + 3y - 4z = -4$$

10. $x + 3y - 6z = 7$

$$2x - y + z = 1$$

$$x + 2y + 2z = -1$$

11. $2x + 5y - z = 12$

$$-x + y - 4z = -10$$

$$-8x - 20y + 4z = 31$$

Solve each problem.


12. Which system would be easier to solve using the substitution method? Why?

System A: $5x - 3y = 7$ System B: $7x + 2y = 4$

$$2x + 8y = 3$$

$$y = -3x + 1$$

13. To make a 10% acid solution, Jeffrey wants to mix some 5% solution with 10 L of 20% solution. How many liters of 5% solution should he use?
14. In the 2016 Summer Olympics, China, the United States, and Great Britain won a combined total of 258 medals. China won 51 fewer medals than the United States. Great Britain won 54 fewer medals than the United States. How many medals did each country win? (Data from *The World Almanac and Book of Facts*.)



Chapter 3

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.


► View the complete solutions to all Chapter Test exercises in MyLab Math.

Assuming growth rates continue, the populations of Houston, Phoenix, Dallas, and Philadelphia will follow the trends indicated in the graph. Use the graph to work each problem.

- Which of these cities will experience population growth?
 - Which city will experience population decline?
 - Rank the city populations from least to greatest for the year 2000.
- In which year did the population of Dallas equal that of Philadelphia? About what was this population?
 - Write as an ordered pair (year, population in millions) the point at which Houston and Phoenix will have the same population.
- Solve the system by graphing.

$$\begin{aligned}x + y &= 7 \\x - y &= 5\end{aligned}$$

The Growth Game



Data from U.S. Census Bureau, Chronicle research.

Solve each system. If a system is inconsistent or has dependent equations, say so.

4. $2x - 3y = 24$

$$y = -\frac{2}{3}x$$

5. $3x - y = -8$

$$2x + 6y = 3$$

6. $12x - 5y = 8$

$$3x = \frac{5}{4}y + 2$$

7. $3x + y = 12$

$$2x - y = 3$$

8. $-5x + 2y = -4$

$$6x + 3y = -6$$

9. $3x + 4y = 8$

$$8y = 7 - 6x$$

10. $3x + 5y + 3z = 2$

$$6x + 5y + z = 0$$

$$3x + 10y - 2z = 6$$

11. $4x + y + z = 11$

$$x - y - z = 4$$


$$y + 2z = 0$$

12. $-2x + 4y + 10z = 3$

$$x - 2y - 5z = 1$$


$$x + 2y + z = -1$$

Solve each problem.

- Two top-grossing super hero films, *Captain America: Civil War* and *Deadpool*, earned \$771.2 million together. If *Deadpool* grossed \$45 million less than *Captain America*, how much did each film gross? (Data from comScore, Inc.)
- Two cars start from points 420 mi apart and travel toward each other. They meet after 3.5 hr. Find the average rate of each car if one travels 30 mph slower than the other.
 
- A chemist needs 12 L of a 40% alcohol solution. She must mix a 20% solution and a 50% solution. How many liters of each will be required to obtain what she needs?

Liters of Solution	Percent (as a decimal)	Liters of Pure Alcohol

16. A local electronics store will sell seven AC adaptors and two rechargeable flashlights for \$86, or three AC adaptors and four rechargeable flashlights for \$84. What is the price of a single AC adaptor and a single rechargeable flashlight?
17. The largest angle of a triangle measures 10° less than twice the middle-sized angle. The middle-sized angle measures 5° more than twice the smallest angle. The sum of the measures of the angles of any triangle is 180° . Find the measure of each angle.
18. The owner of a tea shop wants to mix three kinds of tea to make 100 oz of a mixture that will sell for \$0.83 per oz. He uses Orange Pekoe, which sells for \$0.80 per oz, Irish Breakfast, for \$0.85 per oz, and Earl Grey, for \$0.95 per oz. If he wants to use twice as much Orange Pekoe as Irish Breakfast, how much of each kind of tea should he use?



Chapters R-3 Cumulative Review Exercises

Evaluate each expression if possible.

1. $(-3)^4$ 2. -3^4 3. $-(-3)^4$
 4. $\sqrt{0.49}$ 5. $-\sqrt{0.49}$ 6. $\sqrt{-0.49}$

Perform the indicated operations.

7. $-15.42 - (-8.25)$ 8. $-\frac{3}{4} - \left(\frac{2}{3} + \frac{1}{8}\right)$ 9. $\frac{7}{6} \div \left(-\frac{11}{8}\right)$
 10. Simplify $\frac{(-23 + \sqrt{64})(-4^2)}{-6 - 2}$.

Evaluate for $x = -4$, $y = 3$, and $z = 6$.

11. $|2x| + 3y - z^3$ 12. $-5(x^3 - y^3)$

Solve each equation.


13. $7(2x + 3) - 4(2x + 1) = 2(x + 1)$ 14. $|6x - 8| = 4$
 15. $ax + by = d$ for x 16. $0.04x + 0.06(x - 1) = 1.04$

Solve each inequality.

17. $\frac{2}{3}x + \frac{5}{12}x \leq 20$ 18. $|3x + 2| \leq 4$
 19. $|12t + 7| \geq 0$ 20. $2x + 3 > 5$ or $x - 1 \leq 6$

Solve each problem.

21. A jar contains only pennies, nickels, and dimes. The number of dimes is one more than the number of nickels, and the number of pennies is six more than the number of nickels. How many of each denomination are in the jar, if the total value is \$4.80?
22. Two angles of a triangle have the same measure. The measure of the third angle is 4° less than twice the measure of each of the equal angles. Find the measures of the three angles.



23. A survey measured public recognition of some popular contemporary advertising slogans. Complete the results shown in the table if 2500 people were surveyed.

Slogan (product or company)	Percent Recognition (nearest tenth of a percent)	Actual Number Who Recognized Slogan (nearest whole number)
Please Don't Squeeze the . . . (Charmin)	80.4%	
The Breakfast of Champions (Wheaties)	72.5%	
The King of Beers (Budweiser)		1570
Like a Good Neighbor (State Farm)		1430

Data from Department of Integrated Marketing Communications, Northwestern University.

Point A has coordinates $(-2, 6)$ and point B has coordinates $(4, -2)$.

24. What is the equation of the horizontal line through A ?
25. What is the equation of the vertical line through B ?
26. What is the slope of line AB ?
27. What is the slope of a line perpendicular to line AB ?
28. What is the standard form of the equation of line AB ?


Solve each problem.

29. Graph the line having slope $\frac{2}{3}$ and passing through the point $(-1, -3)$.
30. Graph the inequality $-3x - 2y \leq 6$.
31. Given that $f(x) = x^2 + 3x - 6$, find (a) $f(-3)$ and (b) $f(a)$.

Solve each system.

32. $-2x + 3y = -15$
 $4x - y = 15$
33. $x - 3y = 7$
 $2x - 6y = 14$
34. $x + y + z = 10$
 $x - y - z = 0$
 $-x + y - z = -4$

35. The graph shows a company's costs to produce computer parts and the revenue from the sale of computer parts.



STUDY SKILLS REMINDER

It is not too soon to begin preparing for your final exam. **Review Study Skill 10, Preparing for Your Math Final Exam.**

- (a) At what production level does the cost equal the revenue? What is the revenue at that point?
- (b) Profit is revenue less cost. Estimate the profit on the sale of 1100 parts.



EXPONENTS, POLYNOMIALS, AND POLYNOMIAL FUNCTIONS

“Big Data” describes the massive amounts of information collected, analyzed, and applied worldwide today. Large numbers like Big Data numbers can be written using *scientific notation*, one of the topics of this chapter.

- 4.1 Integer Exponents
- 4.2 Scientific Notation
- 4.3 Adding and Subtracting Polynomials

- 4.4 Polynomial Functions, Graphs, and Composition
- 4.5 Multiplying Polynomials
- 4.6 Dividing Polynomials

4.1 Integer Exponents

OBJECTIVES

- 1 Use the product rule for exponents.
- 2 Define 0 and negative exponents.
- 3 Use the quotient rule for exponents.
- 4 Use the power rules for exponents.
- 5 Simplify exponential expressions.

VOCABULARY

- exponent (power)
- base
- exponential expression

STUDY SKILLS REMINDER

Make study cards to help you learn and remember the material in this chapter.

Review Study Skill 5,
Using Study Cards.

NOW TRY EXERCISE 1

Apply the product rule, if possible.

- (a) $8^5 \cdot 8^4$ (b) $4 \cdot 4^2$
 (c) $r^7 \cdot r \cdot r^9$ (d) $p^2 \cdot q^2$
 (e) $(5x^4y^7)(-7xy^3)$

NOW TRY ANSWERS

1. (a) 8^9 (b) 4^3 (c) r^{17}
 (d) The product rule does not apply.
 (e) $-35x^5y^{10}$

Recall that we use exponents to write products of repeated factors.

2^5 is defined as $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$, which equals 32.

In 2^5 , the number 5 is the **exponent** (or **power**). It indicates that the **base** 2 appears as a factor five times. The quantity 2^5 is an **exponential expression**. We read 2^5 as “2 to the fifth power” or “2 to the fifth.”

OBJECTIVE 1 Use the product rule for exponents.

Consider the product $2^5 \cdot 2^3$, which can be simplified as follows.

$$\begin{array}{c} \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ 2^5 \cdot 2^3 = (2 \cdot 2 \cdot 2 \cdot 2 \cdot 2)(2 \cdot 2 \cdot 2) = 2^8 \end{array}$$

$5 + 3 = 8$

This result suggests the **product rule for exponents**.

Product Rule for Exponents

If m and n are natural numbers and a is any real number, then

$$a^m \cdot a^n = a^{m+n}.$$

That is, when multiplying powers of like bases, keep the same base and add the exponents.

To see that the product rule is true, use the definition of an exponent.

$$\begin{array}{cc} a^m = \underbrace{a \cdot a \cdot a \cdot \dots \cdot a}_m & a^n = \underbrace{a \cdot a \cdot a \cdot \dots \cdot a}_n \\ \text{\textit{a appears as a factor } m \text{ times.}} & \text{\textit{a appears as a factor } n \text{ times.}} \end{array}$$

$$\begin{aligned} \text{From this, } a^m \cdot a^n &= \underbrace{a \cdot a \cdot a \cdot \dots \cdot a}_m \cdot \underbrace{a \cdot a \cdot a \cdot \dots \cdot a}_n \\ &= \underbrace{a \cdot a \cdot a \cdot \dots \cdot a}_{(m+n) \text{ factors}} \\ a^m \cdot a^n &= a^{m+n}. \end{aligned}$$

EXAMPLE 1 Using the Product Rule for Exponents

Apply the product rule for exponents, if possible.

(a) $3^4 \cdot 3^7$ (b) $5^3 \cdot 5$ (c) $y^3 \cdot y^8 \cdot y^2$

$$\begin{array}{lll} = 3^{4+7} & = 5^3 \cdot 5^1 & = y^{3+8+2} \\ = 3^{11} & = 5^{3+1} & = y^{13} \end{array}$$

Keep the same base.

$= 5^4$

(d) $x^2 \cdot y^4$ The product rule does not apply because the bases are not the same.

(e) $(5y^2)(-3y^4)$ (f) $(7p^3q)(2p^5q^2)$

$$\begin{array}{lll} = 5(-3)y^2y^4 & \text{Commutative property} & = 7(2)p^3p^5q^1q^2 \\ = -15y^{2+4} & \text{Multiply; product rule} & = 14p^{3+5}q^{1+2} \\ = -15y^6 & & = 14p^8q^3 \end{array}$$

! CAUTION Do not multiply the bases.

$$3^4 \cdot 3^7 = 3^{11}, \text{ not } 9^{11}. \quad \text{See Example 1(a).}$$

Keep the same base and add the exponents.

OBJECTIVE 2 Define 0 and negative exponents.

Consider the following, where the product rule is extended to whole numbers.

$$4^2 \cdot 4^0 = 4^{2+0} = 4^2$$

For the product rule to hold, 4^0 must equal 1, so we define a^0 this way for any nonzero real number a .

Zero Exponent

If a is any nonzero real number, then

$$a^0 = 1.$$

The expression 0^0 is undefined.*

EXAMPLE 2 Using 0 as an Exponent

Evaluate.

(a) $6^0 = 1$

(b) $(-6)^0 = 1$

(c) $-6^0 = -(6)^0 = -1$

Here the base is -6 .

The base is 6, not -6 .

(d) $-(-6)^0 = -1$

(e) $(8k)^0 = 1 \quad (k \neq 0)$

(f) $5^0 + 12^0$

Any nonzero quantity raised to the zero power equals 1.

$$= 1 + 1$$

$$= 2$$

NOW TRY 

To define a negative exponent, we extend the product rule, as follows.

$$8^2 \cdot 8^{-2} = 8^{2+(-2)} = 8^0 = 1$$

Because their product is 1, it appears that 8^{-2} is the *reciprocal* of 8^2 . But we know that $\frac{1}{8^2}$ is the reciprocal of 8^2 , and a number can have only one reciprocal. Therefore, 8^{-2} must equal $\frac{1}{8^2}$.

We can generalize this result.

Negative Exponent

For any natural number n and any nonzero real number a ,

$$a^{-n} = \frac{1}{a^n}.$$

With this definition, the expression a^n is meaningful for any integer exponent n and any nonzero real number a .

NOW TRY EXERCISE 2

Evaluate.

(a) 5^0 (b) -5^0 (c) $(-5)^0$

(d) $(-5x)^0 \quad (x \neq 0)$

(e) $10^0 - 9^0$

NOW TRY ANSWERS

2. (a) 1 (b) -1 (c) 1
(d) 1 (e) 0

*In advanced courses, 0^0 is called an *indeterminate form*.

! CAUTION A negative exponent does not indicate that an expression represents a negative number. Negative exponents lead to reciprocals.

$$3^{-2} = \frac{1}{3^2} = \frac{1}{9} \quad \text{Not negative} \quad \Bigg| \quad -3^{-2} = -\frac{1}{3^2} = -\frac{1}{9} \quad \text{Negative}$$

The base is 3 in both cases. 3^{-2} represents the reciprocal of 3^2 .

NOW TRY EXERCISE 3

Write using only positive exponents.

- (a) 9^{-4} (b) 2^{-1}
 (c) $(3y)^{-6}$ ($y \neq 0$)
 (d) $-4k^{-3}$ ($k \neq 0$)

EXAMPLE 3 Using Negative Exponents

Write using only positive exponents.

- (a) $2^{-3} = \frac{1}{2^3}$ This is the reciprocal of 2^3 .
 (b) $6^{-1} = \frac{1}{6^1} = \frac{1}{6}$ This is the reciprocal of 6^1 , or 6.
 (c) $(5z)^{-3} = \frac{1}{(5z)^3}$ ($z \neq 0$)
↑
The base is $5z$. (Note the parentheses.)
 (d) $5z^{-3} = 5\left(\frac{1}{z^3}\right) = \frac{5}{z^3}$ ($z \neq 0$)
↑
The base is z , not $5z$.
 (e) $-m^{-2} = -\frac{1}{m^2}$ ($m \neq 0$)
(What is the base here?)
 (f) $(-m)^{-2} = \frac{1}{(-m)^2}$, or $\frac{1}{m^2}$ ($m \neq 0$)
(What is the base here?) **NOW TRY**

NOW TRY EXERCISE 4

Evaluate.

- (a) $4^{-1} + 6^{-1}$ (b) $\frac{1}{5^{-3}}$
 (c) $\frac{10^{-2}}{2^{-5}}$

EXAMPLE 4 Using Negative Exponents

Evaluate.

- (a) $3^{-1} + 4^{-1}$

$$= \frac{1}{3} + \frac{1}{4} \quad \text{Definition of negative exponent}$$

$$= \frac{4}{12} + \frac{3}{12} \quad \frac{1}{3} \cdot \frac{4}{4} = \frac{4}{12}; \frac{1}{4} \cdot \frac{3}{3} = \frac{3}{12}$$

$$= \frac{7}{12} \quad \frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$
 (b) $5^{-1} - 2^{-1}$

$$= \frac{1}{5} - \frac{1}{2} \quad \text{Definition of negative exponent}$$

$$= \frac{2}{10} - \frac{5}{10} \quad \text{Find a common denominator.}$$

$$= -\frac{3}{10} \quad \frac{a}{c} - \frac{b}{c} = \frac{a-b}{c}$$
 (c) $\frac{1}{2^{-3}} = \frac{1}{\frac{1}{2^3}} = 1 \div \frac{1}{2^3} = 1 \cdot \frac{2^3}{1} = 2^3 = 8$
Multiply by the reciprocal of the divisor. $\frac{1}{2^{-3}}$ represents the reciprocal of 2^{-3} . Because $2^{-3} = \frac{1}{8}$, the reciprocal is 8, which agrees with the final answer.
 (d) $\frac{2^{-3}}{3^{-2}} = \frac{\frac{1}{2^3}}{\frac{1}{3^2}} = \frac{1}{2^3} \div \frac{1}{3^2} = \frac{1}{2^3} \cdot \frac{3^2}{1} = \frac{3^2}{2^3} = \frac{9}{8}$

NOW TRY

NOW TRY ANSWERS

3. (a) $\frac{1}{9^4}$ (b) $\frac{1}{2}$
 (c) $\frac{1}{(3y)^6}$ (d) $-\frac{4}{k^3}$
 4. (a) $\frac{5}{12}$ (b) 125 (c) $\frac{8}{25}$

! CAUTION In Example 4(a), $3^{-1} + 4^{-1} \neq (3 + 4)^{-1}$. The expression $3^{-1} + 4^{-1}$ simplifies to $\frac{7}{12}$, while the expression $(3 + 4)^{-1}$ simplifies to 7^{-1} , which equals $\frac{1}{7}$. Similar reasoning can be applied in Example 4(b).

Example 4 suggests the following generalizations.

Special Rules for Negative Exponents

If $a \neq 0$ and $b \neq 0$, then

$$\frac{1}{a^{-n}} = a^n \quad \text{and} \quad \frac{a^{-n}}{b^{-m}} = \frac{b^m}{a^n}.$$

OBJECTIVE 3 Use the quotient rule for exponents.

We simplify a quotient, such as $\frac{a^8}{a^3}$, in much the same way as a product. (In all quotients of this type, assume that the denominator is not 0.) Consider this example.

$$\frac{a^8}{a^3} = \frac{a \cdot a \cdot a \cdot a \cdot a \cdot a \cdot a \cdot a}{a \cdot a \cdot a} = a \cdot a \cdot a \cdot a \cdot a = a^5$$

Notice that $8 - 3 = 5$. In the same way, we simplify $\frac{a^3}{a^8}$.

$$\frac{a^3}{a^8} = \frac{a \cdot a \cdot a}{a \cdot a \cdot a \cdot a \cdot a \cdot a \cdot a \cdot a} = \frac{1}{a^5} = a^{-5}$$

Here, $3 - 8 = -5$. These examples suggest the **quotient rule for exponents**.

Quotient Rule for Exponents

If a is any nonzero real number and m and n are integers, then

$$\frac{a^m}{a^n} = a^{m-n}.$$

That is, when dividing powers of like bases, keep the same base and subtract the exponent of the denominator from the exponent of the numerator.

NOW TRY EXERCISE 5

Apply the quotient rule, if possible, and write each result using only positive exponents.

- (a) $\frac{t^8}{t^2}$ ($t \neq 0$) (b) $\frac{4^5}{4^{-2}}$
 (c) $\frac{2^3}{2^7}$ (d) $\frac{m^4}{n^3}$ ($n \neq 0$)

NOW TRY ANSWERS

5. (a) t^6 (b) 4^7 (c) $\frac{1}{2^4}$
 (d) The quotient rule does not apply.

EXAMPLE 5 Using the Quotient Rule for Exponents

Apply the quotient rule for exponents, if possible, and write each result using only positive exponents.

- Numerator exponent**
Denominator exponent
Subtraction symbol
- (a) $\frac{3^7}{3^2} = 3^{7-2} = 3^5$
- (b) $\frac{p^6}{p^2} = p^{6-2} = p^4$ ($p \neq 0$)
- (c) $\frac{k^7}{k^{12}} = k^{7-12} = k^{-5} = \frac{1}{k^5}$ ($k \neq 0$)
- (d) $\frac{2^7}{2^{-3}} = 2^{7-(-3)} = 2^{7+3} = 2^{10}$
Use parentheses to avoid errors.
- (e) $\frac{8^{-2}}{8^5} = 8^{-2-5} = 8^{-7} = \frac{1}{8^7}$
- (f) $\frac{6}{6^{-1}} = \frac{6^1}{6^{-1}} = 6^{1-(-1)} = 6^2$
- (g) $\frac{z^{-5}}{z^{-8}} = z^{-5-(-8)} = z^3$ ($z \neq 0$)
Be careful with signs.
- (h) $\frac{a^3}{b^4}$ ($b \neq 0$)
This expression cannot be simplified further.

The quotient rule does not apply because the bases are different.

OBJECTIVE 4 Use the power rules for exponents.

We can simplify $(3^4)^2$ as follows.

$$(3^4)^2 = 3^4 \cdot 3^4 = 3^{4+4} = 3^8$$

Notice that $4 \cdot 2 = 8$. This example suggests the first **power rule for exponents**. The other two power rules can be demonstrated similarly.

Power Rules for Exponents

If a and b are real numbers and m and n are integers, then

$$(a) \ (a^m)^n = a^{mn}, \quad (b) \ (ab)^m = a^m b^m, \quad \text{and} \quad (c) \ \left(\frac{a}{b}\right)^m = \frac{a^m}{b^m} \quad (b \neq 0).$$

That is,

- (a) To raise a power to a power, multiply exponents.
- (b) To raise a product to a power, raise each factor to that power.
- (c) To raise a quotient to a power, raise the numerator and the denominator to that power.

NOW TRY EXERCISE 6

Simplify using the power rules.

(a) $(x^4)^2$ (b) $(9x)^3$

(c) $(-2m^3)^4$

(d) $\left(\frac{3x^2}{y^3}\right)^3$ ($y \neq 0$)

EXAMPLE 6 Using the Power Rules for Exponents

Simplify using the power rules.

$$(a) \ (p^8)^3 \quad \text{Power rule (a)} \quad = p^{8 \cdot 3} = p^{24}$$

$$(b) \ (3y)^4 \quad \text{Power rule (b)} \quad = 3^4 y^4 = 81y^4$$

$$(c) \ \left(\frac{2}{3}\right)^4 \quad \text{Power rule (c)} \quad = \frac{2^4}{3^4} = \frac{16}{81}$$

$$(d) \ (6p^7)^2 \quad \text{Power rule (b)} \quad = 6^2 (p^7)^2 = 6^2 p^{7 \cdot 2} = 36p^{14} \quad \text{Power rule (a), Square 6, Multiply exponents.}$$

$$(e) \ \left(\frac{-2m^5}{z}\right)^3 \quad \text{Power rule (c)} \quad = \frac{(-2m^5)^3}{z^3} = \frac{(-2)^3 m^{5 \cdot 3}}{z^3} = \frac{-8m^{15}}{z^3} \quad \text{Power rule (b), Simplify.}$$

NOW TRY

The reciprocal of a^n is $\frac{1}{a^n}$, which equals $\left(\frac{1}{a}\right)^n$. Also, a^n and a^{-n} are reciprocals.

$$a^n \cdot a^{-n} = a^n \cdot \frac{1}{a^n} = 1 \quad \text{Reciprocals have product 1.}$$

Thus, because both $\left(\frac{1}{a}\right)^n$ and a^{-n} are reciprocals of a^n , the following is true.

$$a^{-n} = \left(\frac{1}{a}\right)^n \quad a^{-n} \text{ represents the reciprocal of } a^n.$$

NOW TRY ANSWERS

6. (a) x^8 (b) $729x^3$

(c) $16m^{12}$ (d) $\frac{27x^6}{y^9}$

Special Rules for Negative Exponents, Continued

If $a \neq 0$ and $b \neq 0$ and n is an integer, then

$$a^{-n} = \left(\frac{1}{a}\right)^n \quad \text{and} \quad \left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n.$$

That is, any nonzero number raised to the negative n th power is equal to the reciprocal of that number raised to the n th power.


NOW TRY EXERCISE 7

Write using only positive exponents and then evaluate.

(a) $\left(\frac{1}{3}\right)^{-2}$ (b) $\left(\frac{5}{3}\right)^{-3}$

(c) $\left(\frac{3}{2x}\right)^{-4}$ ($x \neq 0$)

EXAMPLE 7 Using Negative Exponents with Fractions

Write using only positive exponents and then evaluate.

<p>(a) $\left(\frac{1}{2}\right)^{-4}$</p> $= 2^4 \quad \left(\frac{1}{a}\right)^n = a^{-n}$ $= 16$	<p>(b) $\left(\frac{3}{7}\right)^{-2}$</p> $= \left(\frac{7}{3}\right)^2 \quad \left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n$ $= \frac{7^2}{3^2} \quad \text{Power rule (c)}$ $= \frac{49}{9} \quad \text{Apply exponents.}$	<p>(c) $\left(\frac{4x}{5}\right)^{-3}$ ($x \neq 0$)</p> $= \left(\frac{5}{4x}\right)^3 \quad \left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n$ $= \frac{5^3}{4^3 x^3} \quad \text{Power rules (c) and (b)}$ $= \frac{125}{64x^3} \quad \text{Apply exponents.}$
----------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Applying the negative exponent involves changing the base to its reciprocal and changing the sign of the exponent.

NOW TRY 

Summary of Definitions and Rules for Exponents

For all integers m and n and all real numbers a and b for which the following are defined, these rules hold true.

		Examples
Product Rule	$a^m \cdot a^n = a^{m+n}$	$5^2 \cdot 5^4 = 5^{2+4} = 5^6$
Quotient Rule	$\frac{a^m}{a^n} = a^{m-n}$	$\frac{4^9}{4^2} = 4^{9-2} = 4^7$
Zero Exponent	$a^0 = 1$	$(-2)^0 = 1$
Negative Exponent	$a^{-n} = \frac{1}{a^n}$	$3^{-4} = \frac{1}{3^4}$
Power Rules	(a) $(a^m)^n = a^{mn}$	$(3^2)^5 = 3^{2 \cdot 5} = 3^{10}$
	(b) $(ab)^m = a^m b^m$	$(3x)^2 = 3^2 x^2$
	(c) $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$	$\left(\frac{2}{3}\right)^5 = \frac{2^5}{3^5}$

Special Rules for Negative Exponents

$$\frac{1}{a^{-n}} = a^n \quad \frac{1}{4^{-3}} = 4^3 \quad \frac{a^{-n}}{b^{-m}} = \frac{b^m}{a^n} \quad \frac{9^{-2}}{3^{-4}} = \frac{3^4}{9^2}$$

$$a^{-n} = \left(\frac{1}{a}\right)^n \quad 6^{-2} = \left(\frac{1}{6}\right)^2 \quad \left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n \quad \left(\frac{4}{7}\right)^{-3} = \left(\frac{7}{4}\right)^3$$

NOW TRY ANSWERS

7. (a) 9 (b) $\frac{27}{125}$ (c) $\frac{16x^4}{81}$


NOW TRY EXERCISE 8

Simplify. Assume that all variables represent nonzero real numbers.

(a) $x^{-8} \cdot x \cdot x^4$

(b) $(5^{-3})^2$

(c) $\frac{p^2q^{-4}}{p^{-2}q^{-1}}$

(d) $\left(\frac{2x^2}{y^2}\right)^3 \left(\frac{-5x^{-2}}{y}\right)^{-2}$

OBJECTIVE 5 Simplify exponential expressions.

EXAMPLE 8 Using the Definitions and Rules for Exponents

Simplify. Assume that all variables represent nonzero real numbers.

(a) $3^2 \cdot 3^{-5}$

$$= 3^{2+(-5)} \quad \text{Product rule}$$

$$= 3^{-3} \quad \text{Add exponents.}$$

$$= \frac{1}{3^3} \quad a^{-n} = \frac{1}{a^n}$$

$$= \frac{1}{27} \quad 3^3 = 27$$

(b) $x^{-3}x^{-4}x^2$

$$= x^{-3+(-4)+2} \quad \text{Product rule}$$

$$= x^{-5} \quad \text{Add exponents.}$$

$$= \frac{1}{x^5} \quad a^{-n} = \frac{1}{a^n}$$

(c) $(4^{-2})^{-5}$

$$= 4^{(-2)(-5)} \quad \text{Power rule (a)}$$

$$= 4^{10} \quad \text{Multiply exponents.}$$

(d) $(x^{-4})^6$

$$= x^{(-4)6} \quad \text{Power rule (a)}$$

$$= x^{-24} \quad \text{Multiply exponents.}$$

$$= \frac{1}{x^{24}} \quad a^{-n} = \frac{1}{a^n}$$

(e) $\frac{x^{-4}y^2}{x^2y^{-5}}$

$$= \frac{x^{-4}}{x^2} \cdot \frac{y^2}{y^{-5}} \quad \frac{ab}{cd} = \frac{a}{c} \cdot \frac{b}{d}$$

$$= x^{-4-2} \cdot y^{2-(-5)} \quad \text{Quotient rule}$$

$$= x^{-6}y^7 \quad \text{Subtract exponents.}$$

$$= \frac{y^7}{x^6} \quad a^{-n} = \frac{1}{a^n}$$

(f) $(2^3x^{-2})^{-2}$

$$= (2^3)^{-2} \cdot (x^{-2})^{-2} \quad \text{Power rule (b)}$$

$$= 2^{-6}x^4 \quad \text{Power rule (a)}$$

$$= \frac{x^4}{2^6} \quad a^{-n} = \frac{1}{a^n}$$

$$= \frac{x^4}{64} \quad 2^6 = 64$$

(g) $\left(\frac{3x^2}{y}\right)^2 \left(\frac{4x^3}{y^{-2}}\right)^{-1}$

$$= \frac{3^2(x^2)^2}{y^2} \cdot \frac{y^{-2}}{4x^3} \quad \text{Combination of rules}$$

$$= \frac{9x^4}{y^2} \cdot \frac{y^{-2}}{4x^3} \quad 3^2 = 9; \quad \text{Power rule (a)}$$

$$= \frac{9}{4}x^{4-3}y^{-2-2} \quad \text{Quotient rule}$$

$$= \frac{9}{4}x^1y^{-4} \quad \text{Subtract exponents.}$$

$$= \frac{9x}{4y^4} \quad a^{-n} = \frac{1}{a^n}$$

(h) $\left(\frac{-4m^5n^4}{24mn^{-7}}\right)^{-2}$

$$= \left(\frac{m^{5-1}n^{4-(-7)}}{-6}\right)^{-2} \quad \text{Quotient rule; Divide coefficients.}$$

$$= \left(\frac{m^4n^{11}}{-6}\right)^{-2} \quad \text{Subtract exponents.}$$

$$= \frac{(m^4)^{-2}(n^{11})^{-2}}{(-6)^{-2}} \quad \text{Power rules (b) and (c)}$$

$$= \frac{m^{-8}n^{-22}}{(-6)^{-2}} \quad \text{Power rule (a)}$$

$$= \frac{(-6)^2}{m^8n^{22}} \quad \frac{a^{-n}}{b^{-m}} = \frac{b^m}{a^n}$$

$$= \frac{36}{m^8n^{22}} \quad (-6)^2 = 36$$

NOW TRY ANSWERS


8. (a) $\frac{1}{x^3}$ (b) $\frac{1}{5^6}$

(c) $\frac{p^4}{q^3}$ (d) $\frac{8x^{10}}{25y^4}$

The sign on -6 does *not* change in this step.

4.1 Exercises

FOR
EXTRA
HELP MyLab Math

 Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

We can learn from the mistakes we make. **Review Study Skill 9, Analyzing Your Test Results.**

Concept Check Determine whether each statement is true or false. If false, correct the right-hand side of the statement.

1. $(ab)^2 = ab^2$

2. $(5x)^3 = 5^3x^3$

3. $\left(\frac{4}{a}\right)^3 = \frac{4^3}{a}$ ($a \neq 0$)

4. $x^3 \cdot x^4 = x^7$

5. $xy^0 = 0$ ($y \neq 0$)

6. $(5^2)^3 = 5^6$

7. $-(-10)^0 = 1$

8. $3^{-1} = -\frac{1}{3}$

9. **Concept Check** A friend incorrectly simplified $4^5 \cdot 4^2$ as 16^7 .

$4^5 \cdot 4^2$ as 16^7 .

10. **Concept Check** A student incorrectly simplified $\frac{6^5}{3^2}$ as 2^3 .

$\frac{6^5}{3^2}$ as 2^3 .

WHAT WENT WRONG? Give the correct answer.

WHAT WENT WRONG? Give the correct answer.

Apply the product rule for exponents, if possible. See Example 1.

11. $13^4 \cdot 13^8$

12. $11^6 \cdot 11^4$

13. $8^9 \cdot 8$

14. $12 \cdot 12^6$

15. $x^3 \cdot x^5 \cdot x^9$

16. $y^4 \cdot y^5 \cdot y^6$

17. $r^2 \cdot s^4$

18. $p^3 \cdot q^2$

19. $(-3w^5)(9w^3)$

20. $(-5x^2)(3x^4)$

21. $(2x^2y^5)(9xy^3)$

22. $(8s^4t)(3s^3t^5)$

Match the expression in Column I with its equivalent expression in Column II. Choices may be used once, more than once, or not at all. See Example 2.

I	II	I	II
23. (a) 9^0	A. 0	24. (a) $2x^0$	A. 0
(b) -9^0	B. 1	(b) $-2x^0$	B. 1
(c) $(-9)^0$	C. -1	(c) $(2x)^0$	C. -1
(d) $-(-9)^0$	D. 9	(d) $(-2x)^0$	D. 2
	E. -9	(Note: $x \neq 0$)	E. -2

Evaluate. Assume that all variables represent nonzero real numbers. See Example 2.

25. 15^0

26. 19^0

27. -8^0

28. -10^0

29. $(-25)^0$

30. $(-30)^0$

31. $-(-1)^0$

32. $-(-2)^0$

33. x^0

34. y^0

35. $3^0 + (-3)^0$

36. $5^0 + (-5)^0$

37. $-3^0 + 3^0$

38. $-5^0 + 5^0$

39. $-4^0 - m^0$

40. $-8^0 - k^0$

Match the expression in Column I with its equivalent expression in Column II. Choices may be used once, more than once, or not at all. See Example 3.

I	II	I	II
41. (a) 5^{-2}	A. 25	42. (a) 4^{-3}	A. 64
(b) -5^{-2}	B. $\frac{1}{25}$	(b) -4^{-3}	B. -64
(c) $(-5)^{-2}$	C. -25	(c) $(-4)^{-3}$	C. $\frac{1}{64}$
(d) $-(-5)^{-2}$	D. $-\frac{1}{25}$	(d) $-(-4)^{-3}$	D. $-\frac{1}{64}$

Write each expression using only positive exponents. Assume that all variables represent nonzero real numbers. See Example 3.

- | | | | |
|-----------------|-----------------|------------------|------------------|
| 43. 5^{-4} | 44. 7^{-2} | 45. 3^{-5} | 46. 2^{-3} |
| 47. 9^{-1} | 48. 4^{-1} | 49. $(4x)^{-2}$ | 50. $(5t)^{-3}$ |
| 51. $4x^{-2}$ | 52. $5t^{-3}$ | 53. $-a^{-3}$ | 54. $-b^{-4}$ |
| 55. $(-a)^{-4}$ | 56. $(-b)^{-6}$ | 57. $(-3x)^{-3}$ | 58. $(-2x)^{-5}$ |

Evaluate each expression. See Example 4.

- | | | | |
|-----------------------------|-----------------------------|-----------------------------|-----------------------------|
| 59. $5^{-1} + 6^{-1}$ | 60. $2^{-1} + 8^{-1}$ | 61. $4^{-1} + 5^{-1}$ | 62. $6^{-1} + 7^{-1}$ |
| 63. $8^{-1} - 3^{-1}$ | 64. $6^{-1} - 4^{-1}$ | 65. $\frac{1}{4^{-2}}$ | 66. $\frac{1}{3^{-3}}$ |
| 67. $\frac{2^{-2}}{3^{-3}}$ | 68. $\frac{3^{-3}}{2^{-2}}$ | 69. $\frac{5^{-2}}{2^{-4}}$ | 70. $\frac{3^{-3}}{7^{-2}}$ |

Apply the quotient rule for exponents, if possible, and write each result using only positive exponents. Assume that all variables represent nonzero real numbers. See Example 5.

- | | | | |
|-----------------------------|-----------------------------|--------------------------|-----------------------------|
| 71. $\frac{4^8}{4^6}$ | 72. $\frac{5^9}{5^7}$ | 73. $\frac{x^{12}}{x^8}$ | 74. $\frac{y^{14}}{y^{10}}$ |
| 75. $\frac{r^7}{r^{10}}$ | 76. $\frac{y^8}{y^{12}}$ | 77. $\frac{6^4}{6^{-2}}$ | 78. $\frac{7^5}{7^{-3}}$ |
| 79. $\frac{6^{-3}}{6^7}$ | 80. $\frac{5^{-4}}{5^2}$ | 81. $\frac{7}{7^{-1}}$ | 82. $\frac{8}{8^{-1}}$ |
| 83. $\frac{r^{-3}}{r^{-6}}$ | 84. $\frac{s^{-4}}{s^{-8}}$ | 85. $\frac{x^3}{y^2}$ | 86. $\frac{y^5}{t^3}$ |

Simplify using the power rules. Assume that all variables represent nonzero real numbers. See Example 6.

- | | | | |
|--------------------------------------|----------------------------------------|---------------------------------------|----------------------------------------|
| 87. $(x^3)^6$ | 88. $(y^5)^4$ | 89. $\left(\frac{3}{5}\right)^3$ | 90. $\left(\frac{4}{3}\right)^2$ |
| 91. $(4t)^3$ | 92. $(5t)^4$ | 93. $(6x^2)^3$ | 94. $(2x^5)^5$ |
| 95. $\left(\frac{-4m^2}{t}\right)^3$ | 96. $\left(\frac{-5n^4}{r^2}\right)^3$ | 97. $\left(\frac{-s^3}{t^5}\right)^4$ | 98. $\left(\frac{-2a^4}{b^5}\right)^6$ |

Match the expression in Column I with its equivalent expression in Column II. Choices may be used once, more than once, or not at all. See Example 7.

- | | I | II | | I | II |
|---------|-----------------------------------|-------------------|----------|-----------------------------------|--------------------|
| 99. (a) | $\left(\frac{1}{3}\right)^{-1}$ | A. $\frac{1}{3}$ | 100. (a) | $\left(\frac{2}{5}\right)^{-2}$ | A. $\frac{25}{4}$ |
| (b) | $\left(-\frac{1}{3}\right)^{-1}$ | B. 3 | (b) | $\left(-\frac{2}{5}\right)^{-2}$ | B. $-\frac{25}{4}$ |
| (c) | $-\left(\frac{1}{3}\right)^{-1}$ | C. $-\frac{1}{3}$ | (c) | $-\left(\frac{2}{5}\right)^{-2}$ | C. $\frac{4}{25}$ |
| (d) | $-\left(-\frac{1}{3}\right)^{-1}$ | D. -3 | (d) | $-\left(-\frac{2}{5}\right)^{-2}$ | D. $-\frac{4}{25}$ |

Write using only positive exponents and then evaluate. Assume that all variables represent nonzero real numbers. See Example 7.

101. $\left(\frac{1}{4}\right)^{-3}$

102. $\left(\frac{1}{5}\right)^{-2}$

103. $\left(\frac{2}{3}\right)^{-3}$

104. $\left(\frac{3}{2}\right)^{-3}$

105. $\left(\frac{4}{5}\right)^{-2}$

106. $\left(\frac{5}{4}\right)^{-2}$

107. $\left(\frac{2t}{3}\right)^{-4}$

108. $\left(\frac{3z}{4}\right)^{-3}$

109. $\left(\frac{1}{4x}\right)^{-2}$

110. $\left(\frac{1}{5x}\right)^{-3}$

111. $\left(\frac{x}{2}\right)^{-5}$

112. $\left(\frac{t}{4}\right)^{-4}$

Simplify each expression. Assume that all variables represent nonzero real numbers. See Examples 1–8.

113. $3^5 \cdot 3^{-6}$

114. $4^4 \cdot 4^{-6}$

115. $a^{-3}a^2a^{-4}$

116. $k^{-5}k^{-3}k^4$

117. $(5^{-3})^{-2}$

118. $(4^{-4})^{-2}$

119. $(x^{-4})^3$

120. $(x^{-3})^6$

121. $(k^2)^{-3}k^4$

122. $(x^3)^{-4}x^5$

123. $-4r^{-2}(r^4)^2$

124. $-2m^{-1}(m^3)^2$

125. $(5a^{-1})^4(a^2)^{-3}$

126. $(3p^{-4})^2(p^3)^{-1}$

127. $(z^{-4}x^3)^{-1}$

128. $(y^{-2}z^4)^{-3}$

129. $\frac{m^2n^{-8}}{m^{-6}n^3}$

130. $\frac{a^{-9}b^6}{a^3b^{-4}}$

131. $\frac{(p^{-2})^3}{5p^4}$

132. $\frac{(m^4)^{-1}}{9m^3}$

133. $\frac{(3pq)q^2}{6p^2q^4}$

134. $\frac{(-8xy)y^3}{4x^5y^4}$

135. $\frac{4a^5(a^{-1})^3}{(a^{-2})^{-2}}$

136. $\frac{12k^{-2}(k^{-3})^{-4}}{6k^5}$

137. $\frac{(-y^{-4})^2}{6(y^{-5})^{-1}}$

138. $\frac{2(-m^{-1})^{-4}}{9(m^{-3})^2}$

139. $\frac{(2k)^2m^{-5}}{(km)^{-3}}$

140. $\frac{(3rs)^{-2}}{3^2r^2s^{-4}}$

141. $\left(\frac{2p}{q^2}\right)^3\left(\frac{3p^4}{q}\right)^{-1}$

142. $\left(\frac{5z^3}{2a^2}\right)^{-3}\left(\frac{8a^{-1}}{15z^{-2}}\right)^{-3}$

143. $\frac{2^2y^4(y^{-3})^{-1}}{2^5y^{-2}}$

144. $\frac{3^{-1}m^4(m^2)^{-1}}{3^2m^{-2}}$

145. $\left(\frac{5m^4n^{-3}}{m^{-5}n^2}\right)^{-2}$

146. $\left(\frac{8x^{-6}y^3}{x^4y^{-4}}\right)^{-2}$

147. $\left(\frac{-3x^4y^6}{15x^{-6}y^7}\right)^{-3}$

148. $\left(\frac{-4a^3b^2}{12a^5b^{-4}}\right)^{-3}$

Extending Skills Simplify each expression. Assume that all variables represent nonzero real numbers.

149. $\frac{(2k)^2k^3}{k^{-1}k^{-5}}(5k^{-2})^{-3}$

150. $\frac{(3r^2)^2r^{-5}}{r^{-2}r^3}(2r^{-6})^2$

151. $\frac{(2m^2p^3)^2(4m^2p)^{-2}}{(-3mp^4)^{-1}(2m^3p^4)^3}$

152. $\frac{(-5y^3z^4)^2(2yz^5)^{-2}}{10(y^4z)^3(3y^3z^2)^{-1}}$

4.2 Scientific Notation

OBJECTIVES

- 1 Write numbers in scientific notation.
- 2 Convert numbers in scientific notation to standard notation.
- 3 Use scientific notation in calculations.

OBJECTIVE 1 Write numbers in scientific notation.

The number of one-celled organisms that will sustain a whale for a few hours is 400,000,000,000,000, and the shortest wavelength of visible light is approximately 0.0000004 m. It is often simpler to write such numbers in *scientific notation*.

Scientific Notation

A number is written in **scientific notation** when it is expressed in the form

$$a \times 10^n, \text{ where } 1 \leq |a| < 10 \text{ and } n \text{ is an integer.}$$

Examples:

$$400,000,000,000,000 = 4 \times 10^{14}$$

$$8500 = 8.5 \times 10^3$$

$$0.039 = 3.9 \times 10^{-2}$$

$$0.0000004 = 4 \times 10^{-7}$$

Scientific notation

$1 \leq a < 10$

↓

It is customary to use \times rather than \cdot for multiplication.

! CAUTION Although a number such as 8500 can be written using exponents in multiple ways, these forms are *not* considered scientific notation.

$$0.85 \times 10^4 \text{ or } 85 \times 10^2, \quad \text{Not in scientific notation}$$

↑ ↑

$a < 1$ $a > 10$

In scientific notation, a number is written with the decimal point after the first nonzero digit and then multiplied by a power of 10.

Converting a Positive Number to Scientific Notation

- Step 1 Position the decimal point.** Place a caret \wedge to the right of the first nonzero digit, where the decimal point will be placed.
- Step 2 Determine the numeral for the exponent.** Count the number of digits from the decimal point to the caret. This number gives the absolute value of the exponent on 10.
- Step 3 Determine the sign for the exponent.** Decide whether multiplying by 10^n should make the result of Step 1 greater or less.
- The exponent should be positive to make the result greater.
 - The exponent should be negative to make the result less.

It is helpful to remember the following.

$$\text{For } n \geq 1, \quad 10^{-n} < 1 \quad \text{and} \quad 10^n \geq 10.$$

To convert a negative number to scientific notation, temporarily ignore the negative sign and go through the steps in the box above. Then attach a negative sign to the result.


NOW TRY EXERCISE 1

Write each number in scientific notation.

- (a) 7,560,000,000
(b) -0.000000245

EXAMPLE 1 Writing Numbers in Scientific Notation

Write each number in scientific notation.

- (a) 820,000

Step 1 Place a caret to the right of the 8 (the first nonzero digit) to mark the new location of the decimal point.

$$8_{\wedge}20,000$$

Step 2 Count from the decimal point, which is understood to be after the last 0, to the caret.

$$8.20,000. \leftarrow \text{Decimal point}$$

Count 5 places.

Step 3 Because 8.2 is to be made greater, the exponent on 10 is positive.

$$820,000 = 8.2 \times 10^5$$

- (b) 0.0000072 0.000007_{\wedge}2 Place a caret to the right of the first nonzero digit.

$$0.000007.2$$

Count from left to right.
6 places

Because 7.2 is to be made less, the exponent on 10 is negative.

$$0.0000072 = 7.2 \times 10^{-6}$$

- (c) $-0.0004_{\wedge}62 = -4.62 \times 10^{-4}$ Remember the negative sign.
Count 4 places.

NOW TRY 

OBJECTIVE 2 Convert numbers in scientific notation to standard notation.

Converting a Positive Number from Scientific Notation

Multiplying a positive number by a positive power of 10 makes the number greater, so move the decimal point to the right if n is positive in 10^n .

Multiplying a positive number by a negative power of 10 makes the number less, so move the decimal point to the left if n is negative in 10^n .

If n is 0, leave the decimal point where it is in 10^n .


NOW TRY EXERCISE 2

Write each number in standard notation.

- (a) 4.45×10^{10}
(b) -5.9×10^{-5}

EXAMPLE 2 Converting from Scientific Notation to Standard Notation

Write each number in standard notation.

- (a) 6.93×10^7 6.9300000 Attach 0's as necessary.
7 places

We moved the decimal point 7 places to the right. (We had to attach five 0's.)

$$6.93 \times 10^7 = 69,300,000$$

Insert commas.
Standard notation

- (b) 4.7×10^{-3} .004.7 Move the decimal point 3 places to the left.
3 places Attach 0's as necessary.

$$4.7 \times 10^{-3} = 0.0047$$

Attach a leading zero.

- (c) $-1.083 \times 10^0 = -1.083 \times 1 = -1.083$

NOW TRY 

NOW TRY ANSWERS

1. (a) 7.56×10^9
(b) -2.45×10^{-7}
2. (a) 44,500,000,000
(b) -0.000059

NOTE When converting from scientific notation to standard notation, use the exponent to determine the number of places and the direction in which to move the decimal point.

OBJECTIVE 3 Use scientific notation in calculations.

NOW TRY EXERCISE 3

Evaluate.

$$\frac{0.00063 \times 400,000}{1400 \times 0.000003}$$

EXAMPLE 3 Using Scientific Notation in Computation

Evaluate.

$$\frac{1,800,000 \times 0.0015}{0.00003 \times 45,000}$$

$$= \frac{1.8 \times 10^6 \times 1.5 \times 10^{-3}}{3 \times 10^{-5} \times 4.5 \times 10^4}$$

Express all numbers in scientific notation.

$$= \frac{1.8 \times 1.5 \times 10^6 \times 10^{-3}}{3 \times 4.5 \times 10^{-5} \times 10^4}$$

Commutative property

$$= \frac{1.8 \times 1.5 \times 10^3}{3 \times 4.5 \times 10^{-1}}$$

Product rule

$$= \frac{1.8 \times 1.5}{3 \times 4.5} \times 10^4$$

Quotient rule

$$= 0.2 \times 10^4$$

Simplify.

Don't stop here.

$$= (2 \times 10^{-1}) \times 10^4$$

Write 0.2 in scientific notation.

$$= 2 \times 10^3$$

Scientific notation

$$= 2000$$

Standard notation

NOW TRY 

NOW TRY EXERCISE 4

The speed of light is approximately 30,000,000,000 cm per sec. How long will it take light to travel 1.2×10^{15} cm?

EXAMPLE 4 Using Scientific Notation to Solve Problems

In 1990, national health care expenditures in the United States were \$724 billion. By 2016, health care expenditures were 4.5 times this amount. (Data from Centers for Medicare & Medicaid Services.)

(a) Write the amount of health care expenditures in 1990 using scientific notation.

724 billion

$$= 724 \times 10^9$$

1 billion = 1,000,000,000 = 10^9

$$= (7.24 \times 10^2) \times 10^9$$

Write 724 in scientific notation.

$$= 7.24 \times 10^{11}$$

Product rule; $10^2 \times 10^9 = 10^{2+9}$

In 1990, health care expenditures were $\$7.24 \times 10^{11}$.

(b) What were health care expenditures in 2016?

$$(7.24 \times 10^{11}) \times 4.5$$

Multiply the result in part (a) by 4.5.

$$= (4.5 \times 7.24) \times 10^{11}$$

Commutative and associative properties

$$= 32.58 \times 10^{11}$$

Multiply.

$$= (3.258 \times 10^1) \times 10^{11}$$

Write 32.58 in scientific notation.

$$= 3.258 \times 10^{12}$$

Product rule; $10^1 \times 10^{11} = 10^{12}$

Expenditures in 2016 were about \$3,258,000,000,000 (over \$3 trillion).

NOW TRY ANSWERS

3. 6×10^4 , or 60,000

4. 4.0×10^4 sec, or 40,000 sec

NOW TRY 

4.2 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Reread your class notes before working the assigned exercises. **Review Study**

Skill 3, Taking Lecture Notes.

1. Concept Check In scientific notation, a number is written with a decimal point (before/after) the first nonzero digit and multiplied by a _____ of 10. A number written in scientific notation is expressed in the form _____ \times _____, where $1 \leq |a| < 10$ and n is an integer.

2. Concept Check Match each number written in scientific notation in (a)–(d) in Column I with the equivalent number written in standard notation in A–F in Column II. Not all choices will be used.

I		II		
(a) 1.5×10^4	(b) 1.5×10^5	A. 150,000	B. 1500	C. 0.00015
(c) 1.5×10^{-4}	(d) 1.5×10^{-5}	D. 15,000	E. 0.0015	F. 0.000015

3. Concept Check When asked to write 9,275,000 in scientific notation, a student incorrectly wrote

$$92.75 \times 10^5.$$

WHAT WENT WRONG? What is the correct answer?

4. Concept Check A student incorrectly claimed that the following number is written in scientific notation.

$$0.2 \times 10^{-2}$$

WHAT WENT WRONG? Write this number in scientific notation.

Concept Check The term “Big Data,” originally coined by NASA researchers, includes all data streaming from sources such as cell phones, tablets, satellites, Amazon, Google, Facebook, and Twitter. Write each “Big Data” number in scientific notation. (Data from Thrivent Magazine.)

- | | |
|-----------------------------------------|-------------------------------------------------|
| 5. 1,000,000 bytes (a megabyte) | 6. 1,000,000,000 bytes (a gigabyte) |
| 7. 1,000,000,000,000 bytes (a terabyte) | 8. 1,000,000,000,000,000,000 bytes (an exabyte) |

Write each number in scientific notation. See Example 1.

- | | | | |
|----------------|-----------|---------------|--------------------|
| 9. 530 | 10. 1600 | 11. 0.830 | 12. 0.0072 |
| 13. 0.00000692 | 14. 0.875 | 15. $-38,500$ | 16. $-976,000,000$ |

Write the boldfaced numbers in each problem in scientific notation. See Example 1.

- 17.** The annual U.S. budget first passed **\$1,000,000,000** in 1917. In 1987, it exceeded **\$1,000,000,000,000** for the first time. The budget request for fiscal-year 2018 was **\$4,094,000,000,000**. If stacked in dollar bills, this amount would stretch **277,843** mi. (Data from Office of Management and Budget.)
- 18.** The largest of the **50** United States is Alaska, with land area of about **365,482,000** acres, while the smallest is Rhode Island, with land area of about **677,000** acres. The total land area of the United States is about **2,271,343,000** acres. (Data from General Services Administration.)

Write each number in standard notation. See Example 2.

- | | | | |
|----------------------------|----------------------------|---------------------------|---------------------------|
| 19. 7.2×10^4 | 20. 8.91×10^2 | 21. 2.54×10^{-3} | 22. 5.42×10^{-4} |
| 23. -6×10^4 | 24. -9×10^3 | 25. 1.2×10^{-5} | 26. 2.7×10^{-6} |
| 27. -2.89×10^{-2} | 28. -7.68×10^{-4} | 29. 8.761×10^0 | 30. 5.92×10^0 |

Evaluate. Express answers in standard notation. See Example 3.

31. $\frac{12 \times 10^4}{2 \times 10^6}$

32. $\frac{16 \times 10^5}{4 \times 10^8}$

33. $\frac{3 \times 10^{-2}}{12 \times 10^3}$

34. $\frac{5 \times 10^{-3}}{25 \times 10^2}$

35. $\frac{18 \times 10^{-3} \times 24 \times 10^4}{8 \times 10^5 \times 2 \times 10^{-9}}$

36. $\frac{15 \times 10^{-4} \times 12 \times 10^5}{4 \times 10^3 \times 5 \times 10^{-8}}$

37. $\frac{0.05 \times 1600}{0.0004}$

38. $\frac{0.003 \times 40,000}{0.00012}$

39. $\frac{15,000 \times 0.25}{0.0075 \times 50,000}$

40. $\frac{400,000 \times 0.05}{0.008 \times 25,000}$

41. $\frac{20,000 \times 0.018}{300 \times 0.0004}$

42. $\frac{840,000 \times 0.03}{0.00021 \times 600}$

Solve each problem. See Example 4.

43. In 2016, the population of the United States was approximately 323.1 million. (Data from U.S. Census Bureau.)

- (a) Write the population using scientific notation.
- (b) Write \$1 trillion—that is, \$1,000,000,000,000—using scientific notation.
- (c) Using the answers from parts (a) and (b), calculate how much each person would have had to contribute in order to make someone a trillionaire. Write this amount in standard notation to the nearest dollar.



44. In 2016, the national debt of the U.S. government was about \$19.32 trillion. The U.S. population at that time was approximately 323.1 million. (Data from U.S. Census Bureau, U.S. Department of the Treasury.)

- (a) Write the population using scientific notation.
- (b) Write the amount of the national debt using scientific notation.
- (c) Using the answers for parts (a) and (b), calculate how much debt this is per American. Write this amount in standard notation to the nearest dollar.



45. In a lottery, a player must choose five numbers from 1 through 50 and one number from 1 through 40. There are about 8.5×10^7 different ways to do this. If 2000 people purchase tickets, at \$1.00 each, for all these numbers, how much will each person pay?

LOTTERY	
Choose 5	Choose 1
1 2 3 4 5 6 7 8 9 10	1 2 3 4 5 6 7 8 9 10
11 12 13 14 15 16 17 18 19 20	11 12 13 14 15 16 17 18 19 20
21 22 23 24 25 26 27 28 29 30	21 22 23 24 25 26 27 28 29 30
31 32 33 34 35 36 37 38 39 40	31 32 33 34 35 36 37 38 39 40
41 42 43 44 45 46 47 48 49 50	



46. In 2017, the population of India was about 1.28×10^9 , which was 40,000 times the population of Monaco. What was the population of Monaco? (Data from *The World Almanac and Book of Facts*.)

47. In 2017, the population of Luxembourg was approximately 5.94×10^5 . The population density was 595 people per square mile. (Data from *The World Almanac and Book of Facts*.)

- (a) Write the population density in scientific notation.
- (b) To the nearest square mile, what is the area of Luxembourg?

48. In 2017, the population of Costa Rica was approximately 4.93×10^6 . The population density was 96.6 people per square kilometer. (Data from *The World Almanac and Book of Facts*.)
- (a) Write the population density in scientific notation.
- (b) To the nearest square kilometer, what is the area of Costa Rica?
49. The speed of light is approximately 3×10^{10} cm per sec. How long will it take light to travel 9×10^{12} cm?
50. The average distance from Earth to the sun is 9.3×10^7 mi. How long would it take a rocket, traveling at 2.9×10^3 mph, to reach the sun? Round to the nearest thousand hours.
51. A *light-year* is the distance that light travels in one year. Find the number of miles in a light-year if light travels 1.86×10^5 mi per sec.
52. Use the information given in the previous two exercises to find the number of minutes necessary for light from the sun to reach Earth.

Calculators can express numbers in scientific notation. The displays often use notation such as

$$2.5\text{E}5 \text{ to represent } 2.5 \times 10^5.$$


Similarly, $2.5\text{E}-3$ represents 2.5×10^{-3} . They can also perform operations with numbers entered in scientific notation.

Predict the result the calculator will give for each screen. (Use the customary scientific notation to write the answers.)

53. 

54. 

55. 

56. 

RELATING CONCEPTS For Individual or Group Work (Exercises 57–60)

The *intensity* of an earthquake is measured relative to the intensity of a standard **zero-level** earthquake of intensity I_0 . The relationship is equivalent to $I = I_0 \times 10^R$, where R is the **Richter scale** measure. For example, if an earthquake has magnitude 5.0 on the Richter scale, then its intensity is calculated as

$$I = I_0 \times 10^{5.0} = I_0 \times 100,000,$$

which is 100,000 times as intense as a zero-level earthquake.

To compare an earthquake that measures 8.1 on the Richter scale to one that measures 5.2, find the ratio of the intensities.

$$\frac{\text{intensity } 8.1}{\text{intensity } 5.2} = \frac{I_0 \times 10^{8.1}}{I_0 \times 10^{5.2}} = \frac{10^{8.1}}{10^{5.2}} = 10^{8.1-5.2} = 10^{2.9} \approx 794 \quad \text{Use a calculator.}$$

Therefore, an earthquake that measures 8.1 on the Richter scale is almost 800 times as intense as one that measures 5.2.



Use the information in the table to work Exercises 57–60 in order.

Earthquake		Richter Scale Measurement
1960	Chile: Valdivia–Puerto Montt area	9.5
2002	Alaska: Slana, Mentasta Lake, Fairbanks	7.9
2005	Montana: Dillon, Silver Star, Twin Bridges	5.6
2010	Haiti: Port-au-Prince	7.0
2018	Mexico: Santiago Jamiltepec	5.9

Data from www.earthquake.usgs.gov

57. Compare the intensity of the 2002 Alaska earthquake to that of the 2018 Mexico earthquake.
58. Compare the intensity of the 1960 Chile earthquake to that of the 2010 Haiti earthquake.
59. Compare the intensity of the 2010 Haiti earthquake to that of the 2005 Montana earthquake.
60. Suppose an earthquake measures a value of x on the Richter scale. How does the intensity of a second earthquake compare if its Richter scale measure is $x + 3.0$? If its Richter scale measure is $x - 1.0$?

4.3 Adding and Subtracting Polynomials

OBJECTIVES

- 1 Define and classify polynomials.
- 2 Add and subtract polynomials.

VOCABULARY

- term
- algebraic expression
- polynomial
- numerical coefficient (coefficient)
- degree of a term
- polynomial in x
- descending powers
- leading term
- leading coefficient
- trinomial
- binomial
- monomial
- degree of a polynomial
- like terms
- negative of a polynomial

OBJECTIVE 1 Define and classify polynomials.

Recall that a **term** is a number (constant), a variable, or the product of a number and one or more variables raised to powers.

Examples: $4x$, $\frac{1}{2}m^5$ (or $\frac{m^5}{2}$), $-7z^9$, $6x^2z$, $\frac{5}{3x^2}$, $3\sqrt{x}$, 9 Terms

A term or a sum of two or more terms is an **algebraic expression**. The simplest kind of algebraic expression is a *polynomial*.

Polynomial

A **polynomial** is a term or a finite sum of terms of the form ax^n , where a is a real number, x is a variable, and the exponent n is a whole number.

Examples: $12x^9$, $3t - 5$, $4m^3 - 5m^2 + 8$ Polynomials in x , t , and m

Even though an expression such as $3t - 5$ involves subtraction, it can be written as a sum of terms, here $3t + (-5)$.

Some algebraic expressions are *not* polynomials.

Examples: $x^{-1} + 3x^{-2}$, $\sqrt{9 - x}$, $\frac{1}{x}$ Not polynomials

The first has negative integer exponents, the second involves a variable under a radical, and the third has a variable in the denominator.

For each term ax^n of a polynomial, the factor a is the **numerical coefficient** of x^n , or just the **coefficient** of x^n , and the exponent n is the **degree of the term**.

▼ Terms and Their Coefficients and Degrees

Term ax^n	Numerical Coefficient	Degree
$12x^9$	12	9
$3x$, or $3x^1$	3	1
-6 , or $-6x^0$	-6	0
$-x^4$, or $-1x^4$	-1	4
$\frac{x^2}{3} = \frac{1x^2}{3} = \frac{1}{3}x^2$	$\frac{1}{3}$	2

Any nonzero constant has degree 0.

NOTE The number 0 has no degree because 0 times a variable to any power is 0.

A polynomial containing only the variable x is a **polynomial in x** . A polynomial in one variable is written in **descending powers** of the variable if the exponents on the variable in the terms decrease from left to right.

$$\underbrace{x^5 - 6x^2 + 12x - 5}_{\text{Descending powers of } x}$$

Think of $12x$ as $12x^1$ and -5 as $-5x^0$.

When a polynomial is written in descending powers of the variable, the greatest-degree term is written first and is the **leading term** of the polynomial. Its coefficient is the **leading coefficient**.

NOW TRY EXERCISE 1

Write the polynomial in descending powers of the variable. Then give the leading term and the leading coefficient.

$$-2x^3 - 2x^5 + 4x^2 + 7 - x$$

EXAMPLE 1 Writing Polynomials in Descending Powers

Write each polynomial in descending powers of the variable. Then give the leading term and the leading coefficient.

(a) $y - 6y^3 + 8y^5 - 9y^4 + 12$ is written as $8y^5 - 9y^4 - 6y^3 + y + 12$.

(b) $-2 + m + 6m^2 - m^3$ is written as $-m^3 + 6m^2 + m - 2$.

Each leading term is shown in color. In part (a), the leading coefficient is 8, and in part (b) it is -1 (because $-m^3 = -1m^3$).

NOW TRY

Some polynomials with a specific number of terms are given special names.

- A polynomial with exactly three terms is a **trinomial**.
- A two-term polynomial is a **binomial**.
- A single-term polynomial is a **monomial**.

Although many polynomials contain only one variable, polynomials may have more than one variable. The degree of a term with more than one variable is the sum of the exponents on the variables. The **degree of a polynomial** is the greatest degree of all of its terms.

▼ Polynomials and Their Degrees

Type of Polynomial	Example	Degree
Monomial	7	0 ($7 = 7x^0$)
	$5x^3y^7$	10 ($3 + 7 = 10$)
Binomial	$6 + 2x^3$	3
	$11y + 8$	1 ($y = y^1$)
Trinomial	$t^2 + 11t + 4$	2
	$-3 + 2k^5 + 9z^4$	5
	$x^3y^9 + 12xy^4 + 7xy$	12 (The terms have degrees 12, 5, and 2, and 12 is the greatest.)

NOW TRY ANSWER

1. $-2x^5 - 2x^3 + 4x^2 - x + 7$;
 $-2x^5$; -2

NOTE If a polynomial in a single variable is written in descending powers of that variable, the degree of the polynomial will be the degree of the leading term.

NOW TRY EXERCISE 2

Identify each polynomial as a *monomial*, *binomial*, *trinomial*, or *none of these*. Give the degree.

- (a) $-4x^3 + 10x^5 + 7$
 (b) $-6m^2n^5$

EXAMPLE 2 Classifying Polynomials

Identify each polynomial as a *monomial*, *binomial*, *trinomial*, or *none of these*. Give the degree.

- (a) $-x^2 + 5x + 1$ This is a trinomial of degree 2.
 (b) $\frac{3}{4}xy^4$ (or $\frac{3}{4}x^1y^4$) This is a monomial of degree 5 (because $1 + 4 = 5$).
 (c) $7m^9 + 18m^{14}$ This is a binomial of degree 14.
 (d) $p^4 - p^2 - 6p - 5$ Polynomials of four terms or more do not have special names, so *none of these* is the answer that applies here. This polynomial has degree 4.

NOW TRY 

OBJECTIVE 2 Add and subtract polynomials.

We use the distributive property to simplify polynomials by combining like terms.

$$\begin{aligned} x^3 + 4x^2 + 5x^2 - 1 \\ &= x^3 + (4 + 5)x^2 - 1 && \text{Distributive property} \\ &= x^3 + 9x^2 - 1 && \text{Add.} \end{aligned}$$

The terms in a polynomial such as $4x + 5x^2$ cannot be combined. **Only terms containing exactly the same variables raised to the same powers may be combined.** Recall that such terms are **like terms**.

CAUTION Only like terms can be combined.

NOW TRY EXERCISE 3

Combine like terms.

- (a) $2x^2 - 8x^2 + x^2$
 (b) $3p^3 - 2q + p^3 - 5q$
 (c) $-x^2t + 4x^2t + 3xt^2 - 7xt^2$

EXAMPLE 3 Combining Like Terms

Combine like terms.

- (a) $-5y^3 + 8y^3 - y^3$
 $= (-5 + 8 - 1)y^3$ $-y^3 = -1y^3$; Distributive property
 $= 2y^3$ Add and subtract.
 (b) $6x + 5y - 9x + 2y$
 $= 6x - 9x + 5y + 2y$ Commutative property
 $= -3x + 7y$ Combine like terms.

Because $-3x$ and $7y$ are unlike terms, no further simplification is possible.

- (c) $5x^2y - 6xy^2 + 9x^2y + 13xy^2$
 $= 5x^2y + 9x^2y - 6xy^2 + 13xy^2$ Commutative property
 $= 14x^2y + 7xy^2$ Combine like terms.

NOW TRY ANSWERS

2. (a) trinomial; 5
 (b) monomial; 7
 3. (a) $-5x^2$ (b) $4p^3 - 7q$
 (c) $3x^2t - 4xt^2$

NOW TRY 

Adding Polynomials

To find the sum of two polynomials, combine like terms.

NOW TRY EXERCISE 4

Add.

$$\begin{aligned} &(7x^2 - 9x + 4) + \\ &(x^3 - 3x^2 - 5) \end{aligned}$$

EXAMPLE 4 Adding Polynomials

Add $(3a^5 - 9a^3 + 4a^2) + (-8a^5 + 8a^3 + 2)$.

$$\begin{aligned} &(3a^5 - 9a^3 + 4a^2) + (-8a^5 + 8a^3 + 2) \\ &= 3a^5 - 8a^5 - 9a^3 + 8a^3 + 4a^2 + 2 \\ &= -5a^5 - a^3 + 4a^2 + 2 \end{aligned}$$

Commutative and
associative properties
Combine like terms.

Alternatively, we can add these two polynomials vertically.

$$\begin{array}{r} 3a^5 - 9a^3 + 4a^2 + 0 \\ -8a^5 + 8a^3 + 0a^2 + 2 \\ \hline -5a^5 - a^3 + 4a^2 + 2 \end{array}$$

Place like terms in columns.
Using placeholders may help.

Add only the
coefficients.

Add in columns.

NOW TRY 

NOTE The vertical addition in **Example 4** could be written as follows.

$$\begin{array}{r} 3a^5 - 9a^3 + 4a^2 + 0 \\ + (-8a^5 + 8a^3 + 0a^2 + 2) \\ \hline -5a^5 - a^3 + 4a^2 + 2 \end{array}$$

Recall that subtraction of real numbers is defined as follows.

$$a - b = a + (-b)$$

That is, we add the first number and the negative (or opposite) of the second. We define the **negative of a polynomial** as that polynomial with the sign of every coefficient changed.

Subtracting Polynomials

To find the difference of two polynomials, add the first polynomial (minuend) and the negative (or opposite) of the *second* polynomial (subtrahend).

EXAMPLE 5 Subtracting Polynomials

Subtract $(-6m^2 - 8m + 5) - (-5m^2 + 7m - 8)$.

Change every sign in the second polynomial (subtrahend) and add.

$$\begin{aligned} &(-6m^2 - 8m + 5) - (-5m^2 + 7m - 8) \\ &= -6m^2 - 8m + 5 + 5m^2 - 7m + 8 \\ &= -6m^2 + 5m^2 - 8m - 7m + 5 + 8 \\ &= -m^2 - 15m + 13 \end{aligned}$$

Definition of subtraction

Rearrange terms.

Combine like terms.

CHECK $(-m^2 - 15m + 13) + (-5m^2 + 7m - 8) = -6m^2 - 8m + 5$ ✓

↑
Difference

↑
Subtrahend

↑
Minuend

NOW TRY ANSWER

4. $x^3 + 4x^2 - 9x - 1$


NOW TRY EXERCISE 5

Subtract.

$$(2y^2 - 7y - 4) - (8y^2 - 2y + 10)$$

Alternatively, we can subtract these two polynomials vertically.

$$\begin{array}{r} -6m^2 - 8m + 5 \\ -5m^2 + 7m - 8 \\ \hline \end{array}$$

Write the subtrahend below the minuend, lining up like terms in columns.

Change all the signs in the subtrahend and add.

$$\begin{array}{r} -6m^2 - 8m + 5 \\ +5m^2 - 7m + 8 \\ \hline -m^2 - 15m + 13 \end{array}$$

Change all signs.

Add in columns.

The difference is the same.

NOW TRY 
NOTE The vertical subtraction in **Example 5** could be written as follows.

$$\begin{array}{r} -6m^2 - 8m + 5 \\ -(-5m^2 + 7m - 8) \\ \hline -m^2 - 15m + 13 \end{array}$$

Change each sign and add.


$$\begin{array}{r} -6m^2 - 8m + 5 \\ +(+5m^2 - 7m + 8) \\ \hline -m^2 - 15m + 13 \end{array}$$

NOW TRY ANSWER
 5. $-6y^2 - 5y - 14$

4.3 Exercises

FOR EXTRA HELP

MyLab Math

 Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Time management can be a challenge for students.

Review Study Skill 6, Managing Your Time.

Concept Check Work each problem involving the vocabulary of polynomials.

1. Match each description in Column I with the correct polynomial in Column II. Choices in Column II may be used once, more than once, or not at all.

I

- (a) Monomial of degree 2
 (b) Trinomial of degree 5
 (c) Polynomial with leading coefficient 1
 (d) Binomial in descending powers
 (e) Term with degree 0

II

- A.** $x^5 + 5x^4 - 5x$
B. 5
C. $5 + x - x^2$
D. $3ab$
E. $-m + 5$

2. Which polynomial is a trinomial in descending powers that has degree 6?
A. $5x^6 - 4x^5 + 12$ **B.** $6x^5 - x^6 + 4$
C. $2x + 4x^2 - x^6$ **D.** $4x^6 - 6x^4 + 9x^2 - 8$
3. Give an example of a polynomial of four terms in the variable x that is of degree 5, is written in descending powers, and lacks a fourth-degree term.
4. Give an example of a trinomial in the variable y that is of degree 3, has leading coefficient -1 , and lacks a term with degree 0.
5. **Concept Check** A student incorrectly gave 2 as the leading coefficient of the following polynomial.

$$2x^2 - 5x^3 - 8x^4$$

WHAT WENT WRONG? Give the correct leading coefficient.

6. **Concept Check** A student incorrectly gave 6 as the degree of the following polynomial.

$$2xy^6 + 3xy^5 + 4xy^4$$

WHAT WENT WRONG? Give the degree of this polynomial.

Give the numerical coefficient and the degree of each term. See **Objective 1**.

7. $7z$ 8. $3r$ 9. $-15p^2$ 10. $-27k^3$ 11. x^4 12. y^6
 13. $\frac{t}{6}$ 14. $\frac{m}{4}$ 15. 8 16. 2 17. $-x^3$ 18. $-y^9$

Write each polynomial in descending powers of the variable. Then give the leading term and the leading coefficient. See Example 1.

19. $2x^3 + x - 3x^2 + 4$ 20. $q^2 + 3q^4 - 2q + 1$ 21. $4p^3 - 8p^5 + p^7$
 22. $3y^2 + y^4 - 2y^3$ 23. $10 - m^3 - 3m^4$ 24. $4 - x - 8x^2$

Identify each polynomial as a monomial, binomial, trinomial, or none of these. Give the degree. See Example 2.

25. 25 26. 15 27. $7m - 22$ 28. $6x + 15$
 29. $-7y^6 + 11y^8$ 30. $12k^2 - 9k^5$ 31. $-mn^5$ 32. $-a^3b$
 33. $-5m^3 + 6m - 9m^2$ 34. $4z^2 - 11z + 2$
 35. $-6p^4q - 3p^3q^2 + 2pq^3 - q^4$ 36. $8s^3t - 4s^2t^2 + 2st^3 + 9$

Combine like terms. See Example 3.

37. $5z^4 + 3z^4$ 38. $8r^5 - 2r^5$ 39. $-m^3 + 2m^3 + 6m^3$
 40. $3p^4 + 5p^4 - 2p^4$ 41. $x + x + x + x + x$ 42. $z - z - z + z$
 43. $m^4 - 3m^2 + m$ 44. $5a^5 + 2a^4 - 9a^3$ 45. $5t + 4s - 6t + 9s$
 46. $8p - 9q - 3p + q$ 47. $2k + 3k^2 + 5k^2 - 7$ 48. $4x^2 + 2x - 6x^2 - 6$
 49. $n^4 - 2n^3 + n^2 - 3n^4 + n^3$ 50. $2q^3 + 3q^2 - 4q - q^3 + 5q^2$
 51. $3ab^2 + 7a^2b - 5ab^2 + 13a^2b$ 52. $6m^2n - 8mn^2 + 3mn^2 - 7m^2n$

53. **Concept Check** A student added two polynomials vertically as follows.

$$\begin{array}{r} 4x^2 - 7x + 4 \\ 2x^2 + 3x - 5 \\ \hline 6x^4 - 4x^2 - 1 \end{array} \quad \text{Incorrect}$$

WHAT WENT WRONG? Give the correct sum.

54. **Concept Check** A student subtracted $(15x^2 + 12) - (7x^2 - 3)$ vertically as follows.

$$\begin{array}{r} 15x^2 + 12 \\ 7x^2 - 3 \\ \hline 8x^2 + 9 \end{array} \quad \text{Incorrect}$$

WHAT WENT WRONG? Give the correct difference.

Add or subtract as indicated. See Examples 4 and 5.

55. $(5x^2 + 7x - 4) + (3x^2 - 6x + 2)$ 56. $(4k^3 + k^2 + k) + (2k^3 - 4k^2 - 3k)$
 57. $(6t^2 - 4t^4 - t) + (3t^4 - 4t^2 + 5)$ 58. $(3p^2 + 2p - 5) + (7p^2 - 4p^3 + 3p)$
 59. $(y^3 + 3y + 2) + (4y^3 - 3y^2 + 2y - 1)$ 60. $(2x^5 - 2x^4 + x^3 - 1) + (x^4 - 3x^3 + 2)$
 61. $(3r + 8) - (2r - 5)$ 62. $(2d + 7) - (3d - 1)$
 63. $(2a^2 + 3a - 1) - (4a^2 + 5a + 6)$ 64. $(q^4 - 2q^2 + 10) - (3q^4 + 5q^2 - 5)$
 65. $(z^5 + 3z^2 + 2z) - (4z^5 + 2z^2 - 5z)$ 66. $(5t^3 - 3t^2 + 2t) - (4t^3 + 2t^2 + 3t)$
 67. Add. 68. Add. 69. Add. 70. Add.

$$\begin{array}{r} 21p - 8 \\ -9p + 4 \\ \hline \end{array}$$

$$\begin{array}{r} 15m - 9 \\ 4m + 12 \\ \hline \end{array}$$

$$\begin{array}{r} -12p^2 + 4p - 1 \\ 3p^2 + 7p - 8 \\ \hline \end{array}$$

$$\begin{array}{r} -6y^3 + 8y + 5 \\ 9y^3 + 4y - 6 \\ \hline \end{array}$$

 71. Subtract. 72. Subtract. 73. Subtract. 74. Subtract.

$$\begin{array}{r} 12a + 15 \\ 7a - 3 \\ \hline \end{array}$$

$$\begin{array}{r} -3b + 6 \\ 2b - 8 \\ \hline \end{array}$$

$$\begin{array}{r} 6m^2 - 11m + 5 \\ -8m^2 + 2m - 1 \\ \hline \end{array}$$

$$\begin{array}{r} -4z^2 + 2z - 1 \\ 3z^2 - 5z + 2 \\ \hline \end{array}$$

75. Add.

$$\begin{array}{r} 12z^2 - 11z + 8 \\ 5z^2 + 16z - 2 \\ \hline -4z^2 + 5z - 9 \end{array}$$

76. Add.

$$\begin{array}{r} -6m^3 + 2m^2 + 5m \\ 8m^3 + 4m^2 - 6m \\ \hline -3m^3 + 2m^2 - 7m \end{array}$$

77. Add.

$$\begin{array}{r} 6y^3 - 9y^2 + 8 \\ 4y^3 + 2y^2 + 5y \\ \hline \end{array}$$

78. Add.

$$\begin{array}{r} -7r^8 + 2r^6 - r^5 \\ 3r^6 + 5 \\ \hline \end{array}$$

79. Subtract.


$$\begin{array}{r} -5a^4 + 8a^2 - 9 \\ 6a^3 - a^2 + 2 \\ \hline \end{array}$$

80. Subtract.


$$\begin{array}{r} -2m^3 + 8m^2 \\ m^4 - m^3 + 2m \\ \hline \end{array}$$

Extending Skills Perform the indicated operations.81. Subtract $4y^2 - 2y + 3$ from $7y^2 - 6y + 5$.82. Subtract $-(-4x + 2z^2 + 3m)$ from $[(2z^2 - 3x + m) + (z^2 - 2m)]$.83. $(-4m^2 + 3n^2 - 5n) - [(3m^2 - 5n^2 + 2n) + (-3m^2) + 4n^2]$ 84. $[-(4m^2 - 8m + 4m^3) - (3m^2 + 2m + 5m^3)] + m^2$ 85. $[-(y^4 - y^2 + 1) - (y^4 + 2y^2 + 1)] + (3y^4 - 3y^2 - 2)$ 86. $[2p - (3p - 6)] - [(5p - (8 - 9p)) + 4p]$ 87. $-[3z^2 + 5z - (2z^2 - 6z)] + [(8z^2 - [5z - z^2]) + 2z^2]$ 88. $5k - (5k - [2k - (4k - 8k)]) + 11k - (9k - 12k)$ Find the perimeter of each figure. Express it as a polynomial in descending powers of the variable x .

89.



90.



4.4

Polynomial Functions, Graphs, and Composition

OBJECTIVES

- 1 Recognize and evaluate polynomial functions.
- 2 Use a polynomial function to model data.
- 3 Add and subtract polynomial functions.
- 4 Graph basic polynomial functions.
- 5 Find the composition of functions.

OBJECTIVE 1 Recognize and evaluate polynomial functions.

We have studied linear (first-degree polynomial) functions $f(x) = ax + b$. Now we consider more general polynomial functions.

Polynomial Function

A **polynomial function of degree n** is defined by

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

for real numbers a_n, a_{n-1}, \dots, a_1 , and a_0 , where $a_n \neq 0$ and n is a whole number.

Examples: $f(x) = 2$, $f(x) = -6x + 1$, $f(x) = 5x^2 - x + 4$,
 $f(x) = x^3 + 2x^2 - 5x - 3$

Another way of describing a polynomial function is to say that it is a function defined by a polynomial in one variable, consisting of one or more terms. It is usually written in descending powers of the variable, and its degree is the degree of the polynomial that defines it.

VOCABULARY

- polynomial function
- identity function
- squaring function
- cubing function
- composite function
(composition of functions)

NOW TRY
EXERCISE 1

Let $f(x) = x^3 - 2x^2 + 7$.
Find $f(-3)$.

We can evaluate a polynomial function $f(x)$ at different values of the variable x .

EXAMPLE 1 Evaluating Polynomial Functions

Let $f(x) = 4x^3 - x^2 + 5$. Find each value.

(a) $f(3)$

Read this as “ f of 3,”
not “ f times 3.”

$$\begin{aligned} f(x) &= 4x^3 - x^2 + 5 && \text{Given function} \\ f(3) &= 4(3)^3 - 3^2 + 5 && \text{Substitute 3 for } x. \\ f(3) &= 4(27) - 9 + 5 && \text{Apply the exponents.} \\ f(3) &= 108 - 9 + 5 && \text{Multiply.} \\ f(3) &= 104 && \text{Subtract, and then add.} \end{aligned}$$

Thus, $f(3) = 104$ and the ordered pair $(3, 104)$ belongs to f .

(b) $f(-4)$

$$\begin{aligned} f(x) &= 4x^3 - x^2 + 5 && \text{Use parentheses.} \\ f(-4) &= 4(-4)^3 - (-4)^2 + 5 && \text{Let } x = -4. \\ f(-4) &= 4(-64) - 16 + 5 && \text{Be careful with signs.} \\ f(-4) &= -256 - 16 + 5 && \text{Multiply.} \\ f(-4) &= -267 && \text{Subtract, and then add.} \end{aligned}$$

So, $f(-4) = -267$. The ordered pair $(-4, -267)$ belongs to f .

NOW TRY

The capital letter P is sometimes used for polynomial functions. The function

$$P(x) = 4x^3 - x^2 + 5$$

yields the same ordered pairs as the function f in Example 1.

OBJECTIVE 2 Use a polynomial function to model data.

NOW TRY
EXERCISE 2

Use the function in Example 2 to approximate the number of public school students in the United States in 2025 (to two decimal places).

EXAMPLE 2 Using a Polynomial Model to Approximate Data

The number of public school students (grades pre-K–12) in the United States from 1990 projected through 2025 can be modeled by the polynomial function

$$P(x) = -0.0002221x^2 + 0.1392x + 46.80,$$

where $x = 0$ corresponds to the year 1990, $x = 1$ corresponds to 1991, and so on, and $P(x)$ is in millions. Use this function to approximate the number of public school students in 2020. (Data from Department of Education.)

Here $x = 30$ corresponds to 2020, so we find $P(30)$.

$$\begin{aligned} P(x) &= -0.0002221x^2 + 0.1392x + 46.80 && \text{Given function} \\ P(30) &= -0.0002221(30)^2 + 0.1392(30) + 46.80 && \text{Let } x = 30. \\ P(30) &\approx 50.78 && \text{Evaluate.} \end{aligned}$$



The model suggests that there will be approximately 50.78 million public school students in 2020.

NOW TRY

NOW TRY ANSWERS

1. -38
2. 51.40 million students


OBJECTIVE 3 Add and subtract polynomial functions.

The graph in **FIGURE 1** shows dollars (in billions) spent for general science and for space/other technologies over a 20-year period.

$G(x)$ represents dollars spent for general science.

$S(x)$ represents dollars spent for space/other technologies.

$T(x)$ represents total expenditures for these two categories.



Data from U.S. Office of Management and Budget.

FIGURE 1

The total expenditures function can be found by *adding* the spending functions for the two individual categories.

$$T(x) = G(x) + S(x)$$

As another example, businesses use the equation “profit equals revenue minus cost,” which can be written using function notation.

$$\begin{array}{ccccc}
 P(x) & = & R(x) & - & C(x) \\
 \uparrow & & \uparrow & & \uparrow \\
 \text{Profit} & & \text{Revenue} & & \text{Cost} \\
 \text{function} & & \text{function} & & \text{function}
 \end{array}$$

x is the number of items produced and sold.

The profit function is found by *subtracting* the cost function from the revenue function.

We define the following **operations on functions**.

Adding and Subtracting Functions

If $f(x)$ and $g(x)$ define functions, then

$$(f + g)(x) = f(x) + g(x) \quad \text{Sum function}$$

and

$$(f - g)(x) = f(x) - g(x) \quad \text{Difference function}$$

In each case, the domain of the new function is the intersection of the domains of $f(x)$ and $g(x)$.

NOW TRY EXERCISE 3

For $f(x) = x^3 - 3x^2 + 4$
and $g(x) = -2x^3 + x^2 - 12$,
find each of the following.

(a) $(f + g)(x)$

(b) $(f - g)(x)$

EXAMPLE 3 Adding and Subtracting Functions

Find each of the following for the polynomial functions f and g as defined.

$$f(x) = x^2 - 3x + 7 \quad \text{and} \quad g(x) = -3x^2 - 7x + 7$$

(a) $(f + g)(x)$ This notation does not indicate the distributive property.

$$= f(x) + g(x) \quad \text{Sum function}$$

$$= (x^2 - 3x + 7) + (-3x^2 - 7x + 7) \quad \text{Substitute.}$$

$$= -2x^2 - 10x + 14 \quad \text{Add the polynomials.}$$

(b) $(f - g)(x)$

$$= f(x) - g(x) \quad \text{Difference function}$$

$$= (x^2 - 3x + 7) - (-3x^2 - 7x + 7) \quad \text{Substitute.}$$

$$= x^2 - 3x + 7 + 3x^2 + 7x - 7 \quad \text{Change subtraction to addition.}$$

$$= 4x^2 + 4x \quad \text{Add the polynomials.}$$

NOW TRY ANSWERS

3. (a) $-x^3 - 2x^2 - 8$

(b) $3x^3 - 4x^2 + 16$

NOW TRY EXERCISE 4

For $f(x) = x^2 - 4$
and $g(x) = -6x^2$,
find each of the following.

- (a) $(f + g)(x)$
(b) $(f - g)(-4)$

EXAMPLE 4 Adding and Subtracting Functions

Find each of the following for the polynomial functions f and g as defined.

$$f(x) = 10x^2 - 2x \quad \text{and} \quad g(x) = 2x$$

(a) $(f + g)(2)$

$$= f(2) + g(2) \quad \text{Sum function}$$

$$= \underbrace{[10(2)^2 - 2(2)]}_{f(x) = 10x^2 - 2x} + \underbrace{2(2)}_{g(x) = 2x}$$

This is a key step.

$$= [40 - 4] + 4$$

Substitute.

$$= 40$$

Order of operations
Subtract, and then add.

Alternative method: $(f + g)(x)$

$$= f(x) + g(x) \quad \text{Sum function}$$

$$= (10x^2 - 2x) + 2x \quad \text{Substitute.}$$

$$= 10x^2 \quad \text{Combine like terms.}$$

Now find $(f + g)(2)$.

$$(f + g)(2) = 10(2)^2$$

$(f + g)(x) = 10x^2$; Substitute.

The result is the same. $\rightarrow = 40$

Apply the exponent. Multiply.

(b) $(f - g)(x)$ and $(f - g)(1)$

$$(f - g)(x)$$

$$= f(x) - g(x) \quad \text{Difference function}$$

$$= (10x^2 - 2x) - 2x \quad \text{Substitute.}$$

$$= 10x^2 - 4x \quad \text{Combine like terms.}$$

Now find $(f - g)(1)$.

$$(f - g)(1) = 10(1)^2 - 4(1)$$

$(f - g)(x) = 10x^2 - 4x$; Substitute.

$$= 6$$


Perform the operations. **NOW TRY**

Confirm that $f(1) - g(1)$ gives the same result.

OBJECTIVE 4 Graph basic polynomial functions.

Recall that each input (or x -value) of a function results in one output (or y -value). The set of input values (for x) defines the domain of the function, and the set of output values (for y) defines the range.

The simplest polynomial function is the **identity function** $f(x) = x$, graphed in **FIGURE 2**. This function pairs each real number with itself.



Identity function

$$f(x) = x$$

Domain: $(-\infty, \infty)$

Range: $(-\infty, \infty)$


FIGURE 2

NOW TRY ANSWERS

4. (a) $-5x^2 - 4$
(b) 108

NOTE The identity function $f(x) = x$ shown in **FIGURE 2** is a *linear function* of the form $f(x) = ax + b$, where the slope a is 1 and the y -value of the y -intercept b is 0.

Another polynomial function, the **squaring function** $f(x) = x^2$, is graphed in **FIGURE 3**. For this function, every real number is paired with its square. The graph of the squaring function is a *parabola*.



Squaring function


$$f(x) = x^2$$

Domain: $(-\infty, \infty)$

Range: $[0, \infty)$

FIGURE 3

The **cubing function** $f(x) = x^3$ is graphed in **FIGURE 4**. This function pairs every real number with its cube.



Cubing function

$$f(x) = x^3$$

Domain: $(-\infty, \infty)$

Range: $(-\infty, \infty)$

FIGURE 4

EXAMPLE 5 Graphing Variations of Polynomial Functions

Graph each function by creating a table of ordered pairs. Give the domain and range of each function by observing its graph.

(a) $f(x) = 2x$

To find each range value, multiply the domain value by 2. Plot the points and join them with a straight line. See **FIGURE 5**. Both the domain and the range are $(-\infty, \infty)$.

x	$f(x) = 2x$
-2	-4
-1	-2
0	0
1	2
2	4




FIGURE 5

NOW TRY EXERCISE 5

Graph $f(x) = x^2 - 4$. Give the domain and range.

(b) $f(x) = -x^2$

For each input x , square it and then take its opposite. Plotting and joining the points gives a parabola that opens down. It is a *reflection* of the graph of the squaring function across the x -axis. See the table and FIGURE 6. The domain is $(-\infty, \infty)$ and the range is $(-\infty, 0]$.




FIGURE 6




FIGURE 7

(c) $f(x) = x^3 - 2$

For this function, cube the input and then subtract 2 from the result. The graph is that of the cubing function *shifted* down 2 units. See the table and FIGURE 7. The domain and range are both $(-\infty, \infty)$.

NOW TRY
OBJECTIVE 5 Find the composition of functions.

The diagram in FIGURE 8 shows a function g that assigns, to each element x of set X , some element y of set Y . Suppose that a function f takes each element of set Y and assigns a value z of set Z . Then f and g together assign an element x in X to an element z in Z .

The result of this process is a new function h that takes an element x in X and assigns it an element z in Z .





FIGURE 8

Function h is the *composition* of functions f and g , written $f \circ g$.

NOW TRY ANSWER


domain: $(-\infty, \infty)$;
range: $[-4, \infty)$

Composition of Functions

The **composite function**, or **composition**, of functions f and g is defined by

$$(f \circ g)(x) = f(g(x)),$$

for all x in the domain of g such that $g(x)$ is in the domain of f .

Read $f \circ g$ as “*f of g*” (or “*f compose g*”).



As a real-life example of how composite functions occur, consider the following retail situation.

A \$40 pair of blue jeans is on sale for 25% off. If we purchase the jeans before noon, the retailer offers an additional 10% off. What is the final sale price of the blue jeans?

We might be tempted to say that the jeans are 35% off and calculate as follows.

$$\begin{aligned} \$40 - 0.35(\$40) & \quad \text{Original price} - \text{Discount} = \text{Sale price} \\ & = \$40 - \$14 \\ & = \$26 \leftarrow \text{This is not correct.} \end{aligned}$$

To find the correct final sale price, we must first find the price after taking 25% off, and then take an additional 10% off *that* price.

$$\begin{array}{ll} \text{Take 25\% off the} & \text{Take an additional} \\ \text{original price.} & \text{10\% off.} \\ \\ \$40 - 0.25(\$40) & \rightarrow \$30 - 0.10(\$30) \\ = \$40 - \$10 & = \$30 - \$3 \\ = \$30 & = \$27 \end{array}$$

This is the idea behind composition of functions.

As another real-life example of composition, suppose an oil well off the coast is leaking, with the leak spreading oil in a circular layer over the surface. See **FIGURE 9**.

At any time t , in minutes, after the beginning of the leak, the radius of the circular oil slick is given by $r(t) = 5t$ feet. Because $\mathcal{A}(r) = \pi r^2$ gives the area of a circle of radius r , the area can be expressed as a function of time.




FIGURE 9

$$\begin{aligned} \mathcal{A}(r) &= \pi r^2 && \text{Area of a circle} \\ \mathcal{A}(r(t)) &= \pi(5t)^2 && \text{Substitute } 5t \text{ for } r. \\ \mathcal{A}(r(t)) &= 25\pi t^2 && (5t)^2 = 5^2 t^2 = 25t^2; \\ &&& \text{Commutative property} \end{aligned}$$

The function $\mathcal{A}(r(t))$ is a composite function of the functions \mathcal{A} and r .

NOW TRY EXERCISE 6

Let $f(x) = 3x + 7$
and $g(x) = x - 2$.
Find $(f \circ g)(7)$.

EXAMPLE 6 Finding a Composite Function

Let $f(x) = x^2$ and $g(x) = x + 3$. Find $(f \circ g)(4)$.

$$\begin{aligned} (f \circ g)(4) & \quad \text{Evaluate the "inside" function value first.} \\ & = f(g(4)) && \text{Definition of composition} \\ & = f(4 + 3) && \text{Use the rule for } g(x); g(4) = 4 + 3 \\ & = f(7) && \text{Add.} \\ & = 7^2 && \text{Use the rule for } f(x); f(7) = 7^2 \\ & = 49 && \text{Square 7.} \end{aligned}$$

Now evaluate the "outside" function.

NOW TRY ANSWER

6. 22

In this composition, g is the innermost "operation" and acts on x (here 4) first. Then the output value of g (here 7) becomes the input (domain) value of f . **NOW TRY**

If we interchange the order of functions f and g , the composition $g \circ f$, read “ g of f ” (or “ g compose f ”), is defined by

$$(g \circ f)(x) = g(f(x)), \quad \text{for all } x \text{ in the domain of } f \text{ such that } f(x) \text{ is in the domain of } g.$$

NOW TRY EXERCISE 7


Let $f(x) = 3x + 7$
and $g(x) = x - 2$
as in **Now Try Exercise 6**.
Find $(g \circ f)(7)$.

EXAMPLE 7 Finding a Composite Function

Let $f(x) = x^2$ and $g(x) = x + 3$ as in **Example 6**. Find $(g \circ f)(4)$.

$$\begin{aligned} (g \circ f)(4) & \quad \text{Evaluate the "inside" function value first.} \\ &= g(f(4)) \quad \text{Definition of composition} \\ &= g(4^2) \quad \text{Use the rule for } f(x); f(4) = 4^2 \\ &= g(16) \quad \text{Square 4.} \\ &= 16 + 3 \quad \text{Use the rule for } g(x); g(16) = 16 + 3 \\ &= 19 \quad \text{Add.} \end{aligned}$$

Now evaluate the "outside" function.

In this composition, f is the innermost “operation” and acts on x (again 4) first. Then the output value of f (here 16) becomes the input (domain) value of g . **NOW TRY** 

We see in **Examples 6 and 7** that $(f \circ g)(4) \neq (g \circ f)(4)$ because $49 \neq 19$. In general,

$$(f \circ g)(x) \neq (g \circ f)(x).$$

NOW TRY EXERCISE 8

Let $f(x) = x - 5$
and $g(x) = -x^2 + 2$.
Find each of the following.
(a) $(g \circ f)(-1)$
(b) $(f \circ g)(x)$

EXAMPLE 8 Finding Composite Functions

Let $f(x) = 4x - 1$ and $g(x) = x^2 + 5$. Find each of the following.

$$\begin{aligned} \text{(a)} \quad (f \circ g)(2) &= f(g(2)) \quad \text{Definition of composition} \\ &= f(2^2 + 5) \quad g(x) = x^2 + 5 \\ &= f(9) \quad \text{Work inside the parentheses.} \\ &= 4(9) - 1 \quad f(x) = 4x - 1 \\ &= 35 \quad \text{Multiply, and then subtract.} \\ \text{(b)} \quad (f \circ g)(x) &= f(g(x)) \quad \text{Use } g(x) \text{ as the input for function } f. \\ &= 4(g(x)) - 1 \quad \text{Use the rule for } f(x); f(x) = 4x - 1 \\ &= 4(x^2 + 5) - 1 \quad g(x) = x^2 + 5 \\ &= 4x^2 + 20 - 1 \quad \text{Distributive property} \\ &= 4x^2 + 19 \quad \text{Combine like terms.} \end{aligned}$$

(c) Find $(f \circ g)(2)$ again, this time using the rule obtained in part (b).

$$\begin{aligned} (f \circ g)(x) &= 4x^2 + 19 \quad \text{From part (b)} \\ (f \circ g)(2) &= 4(2)^2 + 19 \quad \text{Let } x = 2. \\ &= 4(4) + 19 \quad \text{Square 2.} \\ &= 16 + 19 \quad \text{Multiply.} \\ &= 35 \quad \text{Add.} \end{aligned}$$

NOW TRY ANSWERS

7. 26
8. (a) -34 (b) $-x^2 - 3$

4.4 Exercises

FOR
EXTRA
HELP

MyLab Math

▶ Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

How are you doing on your homework? **Review Study Skill 4, Completing Your Homework.**

Concept Check Work each problem involving polynomial functions and function notation.

- A polynomial function is a function defined by a _____ in (*one / two / three*) variable(s), consisting of one or more (*factors / terms*) and usually written in descending _____ of the variable.
- Which of the following are *not* polynomial functions?
 - $P(x) = x^{-2} - 2x$
 - $f(x) = \frac{1}{2}x^2 + x - 1$
 - $g(x) = -4x + 1.5$
 - $p(x) = x^3 - x^2 - \frac{5}{x}$
- For a function f , the notation $f(5)$ means _____.
 - the variable f times 5, or $5f$
 - f equals 5
 - the range value when the domain value is 5
 - the domain value when the range value is 5
- If $f(x) = 2x$, then $f(5) = \underline{\hspace{2cm}}$. Here $f(5) = 10$ is an abbreviation for the statement “If $x = \underline{\hspace{2cm}}$ in the function f , then $y = \underline{\hspace{2cm}}$.” The ordered pair _____ belongs to f .
- For $f(x) = x$, find $f(0)$, $f(1)$, and $f(2)$. Use the results to write three ordered pairs that belong to f .
- For $f(x) = x^2$, find $f(-2)$, $f(0)$, and $f(2)$. Use the results to write three ordered pairs that belong to f .
- Concept Check** For the function $f(x) = -x^2 + 4$, a student claimed that $f(-2) = 8$. This is incorrect. **WHAT WENT WRONG?** Find the correct value of $f(-2)$.
- Concept Check** For the function $f(x) = x^2 - 1$, a student claimed that $f(-1) = -2$. This is incorrect. **WHAT WENT WRONG?** Find the correct value of $f(-1)$.

For each polynomial function, find (a) $f(-1)$, (b) $f(2)$, and (c) $f(0)$. See Example 1.

- $f(x) = 6x - 4$
- $f(x) = x^2 + 5x$
- $f(x) = 2x^2 - 4x + 1$
- $f(x) = 4x^4 + 2x^2 - 1$
- $f(x) = -2x + 5$
- $f(x) = x^2 - 3x + 4$
- $f(x) = 3x^2 + x - 5$
- $f(x) = -x^2 + 2x^3 - 8$
- $f(x) = x^2 - 7x$
- $f(x) = x^2 - 5x - 4$
- $f(x) = 5x^4 - 3x^2 + 6$
- $f(x) = -x^2 - x^3 + 11$

Solve each problem. See Example 2.

21. Imports of Fair Trade Certified™ coffee into the United States during the years 2007 through 2016 can be modeled by the polynomial function

$$P(x) = -2.169x^2 + 60.28x - 256.6,$$

where $x = 7$ represents 2007, $x = 8$ represents 2008, and so on, and $P(x)$ is in millions of pounds. Use this function to approximate the amount of Fair Trade coffee imported into the United States (to the nearest hundredth) in each given year. (Data from Fair Trade USA.)

- (a) 2007 (b) 2014 (c) 2016



22. The percent of the U.S. population that was foreign-born during the years 1930 through 2016 can be modeled by the polynomial function

$$P(x) = 0.0039x^2 - 0.3001x + 11.33,$$

where $x = 0$ represents 1930, $x = 10$ represents 1940, and so on, and $P(x)$ is percent. Use this function to approximate the percent of the U.S. population that was foreign-born (to the nearest hundredth as needed) in each given year. (Data from U.S. Census Bureau.)

- (a) 1930 (b) 1970 (c) 2016

23. The amount spent by international visitors to the United States during the years 1990 through 2016 can be modeled by the polynomial function

$$P(x) = 0.01287x^3 - 0.3514x^2 + 4.979x + 45.24,$$

where $x = 0$ represents 1990, $x = 1$ represents 1991, and so on, and $P(x)$ is in billions of dollars. Use this function to approximate the amount spent by international visitors to the United States (to the nearest tenth) in each given year. (Data from U.S. Travel Association.)

- (a) 1990 (b) 2005 (c) 2016

24. World tourism receipts during the years 2000 through 2016 can be modeled by the polynomial function

$$P(x) = -0.2869x^3 + 6.139x^2 + 22.84x + 465.3,$$

where $x = 0$ represents 2000, $x = 1$ represents 2001, and so on, and $P(x)$ is in billions of dollars. Use this function to approximate world tourism receipts (to the nearest tenth as needed) in each given year. (Data from World Tourism Association.)

- (a) 2000 (b) 2008 (c) 2016



For each pair of functions, find (a) $(f + g)(x)$ and (b) $(f - g)(x)$. See Example 3.

25. $f(x) = 5x - 10$, $g(x) = 3x + 7$

26. $f(x) = -4x + 1$, $g(x) = 6x + 2$

27. $f(x) = 4x^2 + 8x - 3$, $g(x) = -5x^2 + 4x - 9$

28. $f(x) = 3x^2 - 9x + 10$, $g(x) = -4x^2 + 2x + 12$

Concept Check Find two polynomial functions defined by $f(x)$ and $g(x)$ such that each statement is true.

29. $(f + g)(x) = 3x^3 - x + 3$

30. $(f - g)(x) = -x^2 + x - 5$

Let $f(x) = x^2 - 9$, $g(x) = 2x$, and $h(x) = x - 3$. Find each of the following. See Example 4.

31. $(f + g)(x)$ 32. $(f - g)(x)$ 33. $(f + g)(3)$ 34. $(f - g)(-3)$

35. $(f - h)(x)$ 36. $(f + h)(x)$ 37. $(f - h)(-3)$ 38. $(f + h)(-2)$

39. $(g + h)(-10)$ 40. $(g - h)(10)$ 41. $(g - h)(-3)$ 42. $(g + h)(1)$

43. $(g + h)\left(\frac{1}{4}\right)$ 44. $(g + h)\left(\frac{1}{3}\right)$ 45. $(g + h)\left(-\frac{1}{2}\right)$ 46. $(g + h)\left(-\frac{1}{4}\right)$

Solve each problem. See Objective 3.

47. The cost in dollars to produce x t-shirts is $C(x) = 2.5x + 50$. The revenue in dollars from sales of x t-shirts is $R(x) = 10.99x$.

- (a) Write and simplify a function P that gives profit in terms of x .
 (b) Find the profit if 100 t-shirts are produced and sold.

48. The cost in dollars to produce x youth baseball caps is $C(x) = 4.3x + 75$. The revenue in dollars from sales of x caps is $R(x) = 25x$.

- (a) Write and simplify a function P that gives profit in terms of x .
 (b) Find the profit if 50 caps are produced and sold.



Graph each polynomial function. Give the domain and range. See Example 5.

49. $f(x) = 3x$ 50. $f(x) = -4x$ 51. $f(x) = -2x + 1$
 52. $f(x) = 3x + 2$ 53. $f(x) = -3x^2$ 54. $f(x) = \frac{1}{2}x^2$
 55. $f(x) = x^2 - 2$ 56. $f(x) = -x^2 + 2$ 57. $f(x) = x^3 + 1$
 58. $f(x) = -x^3 + 2$ 59. $f(x) = -2x^3 - 1$ 60. $f(x) = \frac{1}{2}x^3 + 3$

Concept Check Let $f(x) = x^2$ and $g(x) = 2x - 1$. Match each expression in Column I with the description of how to evaluate it in Column II.

I

II

61. $(f \circ g)(5)$ A. Square 5. Take the result and square it.
 62. $(g \circ f)(5)$ B. Double 5 and subtract 1. Take the result and square it.
 63. $(f \circ f)(5)$ C. Double 5 and subtract 1. Take the result, double it, and subtract 1.
 64. $(g \circ g)(5)$ D. Square 5. Take the result, double it, and subtract 1.

65. **Concept Check** For $f(x) = x - 8$ and $g(x) = 2$, a student found $(f \circ g)(x)$ incorrectly as follows.

$$\begin{aligned}(f \circ g)(x) &= 2(x - 8) \\ &= 2x - 16\end{aligned}$$

WHAT WENT WRONG? Give the correct composition.

66. **Concept Check** For $f(x) = 2x + 3$ and $g(x) = x + 5$, a student found $(f \circ g)(x)$ correctly as

$$(f \circ g)(x) = 2x + 13.$$

When asked to find $(g \circ f)(x)$, he gave the same result. **WHAT WENT WRONG?** Give the correct composition.

Let $f(x) = x^2 + 4$, $g(x) = 2x + 3$, and $h(x) = x - 5$. Find each of the following. See Examples 6–8.

67. $(h \circ g)(4)$ 68. $(f \circ g)(4)$ 69. $(g \circ f)(6)$ 70. $(h \circ f)(6)$
 71. $(f \circ h)(-2)$ 72. $(h \circ g)(-2)$ 73. $(f \circ g)(0)$ 74. $(f \circ h)(0)$
 75. $(g \circ f)(x)$ 76. $(g \circ h)(x)$ 77. $(h \circ g)(x)$ 78. $(h \circ f)(x)$
 79. $(f \circ h)\left(\frac{1}{2}\right)$ 80. $(h \circ f)\left(\frac{1}{2}\right)$ 81. $(f \circ g)\left(-\frac{1}{2}\right)$ 82. $(g \circ f)\left(-\frac{1}{2}\right)$

Extending Skills The tables give some selected ordered pairs for functions f and g .

x	3	4	6	8	x	2	7	1	9
$f(x)$	1	3	9	2	$g(x)$	3	6	9	12

Tables like these can be used to evaluate composite functions. For example, to evaluate $(g \circ f)(6)$, use the first table to find $f(6) = 9$. Then use the second table to find

$$(g \circ f)(6) = g(f(6)) = g(9) = 12.$$


Find each of the following.

83. $(f \circ g)(2)$ 84. $(f \circ g)(7)$ 85. $(g \circ f)(3)$
 86. $(g \circ f)(8)$ 87. $(f \circ f)(4)$ 88. $(g \circ g)(1)$


Solve each problem. See **Objective 5**.

89. The function $f(x) = 12x$ computes the number of inches in x feet, and the function $g(x) = 5280x$ computes the number of feet in x miles. Find and simplify $(f \circ g)(x)$. What does it compute?
 90. The function $f(x) = 60x$ computes the number of minutes in x hours, and the function $g(x) = 24x$ computes the number of hours in x days. Find and simplify $(f \circ g)(x)$. What does it compute?
 91. The perimeter x of a square with sides of length s is given by the formula $x = 4s$.

- (a) Solve for s in terms of x .
 (b) If y represents the area of this square, write y as a function of the perimeter x .
 (c) Use the composite function of part (b) to find the area of a square with perimeter 6.



92. The perimeter x of an equilateral triangle with sides of length s is given by the formula $x = 3s$.
 (a) Solve for s in terms of x .
 (b) The area y of an equilateral triangle with sides of length s is given by the formula $y = \frac{s^2\sqrt{3}}{4}$. Write y as a function of the perimeter x .
 (c) Use the composite function of part (b) to find the area of an equilateral triangle with perimeter 12.




93. In a sale room at a clothing store, every item is on sale for half its original price, plus \$1.
 (a) Write a function g that finds half of x .
 (b) Write a function f that adds 1 to x .
 (c) Write and simplify the function $(f \circ g)(x)$.
 (d) Use the function from part (c) to find the sale price of a shirt that has an original price of \$60.



94. A pair of shoes is marked 50% off. A customer has a coupon for an additional \$10 off.
 (a) Write a function g that finds 50% of x .
 (b) Write a function f that subtracts 10 from x .
 (c) Write and simplify the function $(f \circ g)(x)$.
 (d) Use the function from part (c) to find the sale price of a pair of shoes that has an original price of \$100.

95. When a thermal inversion layer is over a city (as happens often in Los Angeles), pollutants cannot rise vertically, but are trapped below the layer and must disperse horizontally.

Assume that a factory smokestack begins emitting a pollutant at 8 A.M. and that the pollutant disperses horizontally over a circular area. Suppose that t represents the time, in hours, since the factory began emitting pollutants ($t = 0$ represents 8 A.M.), and assume that the radius of the circle of pollution is $r(t) = 2t$ miles. Let $\mathcal{A}(r) = \pi r^2$ represent the area of a circle of radius r . Find and interpret $(\mathcal{A} \circ r)(t)$.




96. An oil well is leaking, with the leak spreading oil over the surface as a circle. At any time t , in minutes, after the beginning of the leak, the radius of the circular oil slick on the surface is $r(t) = 4t$ feet. Let $\mathcal{A}(r) = \pi r^2$ represent the area of a circle of radius r . Find and interpret $(\mathcal{A} \circ r)(t)$.

RELATING CONCEPTS For Individual or Group Work (Exercises 97–100)


The polynomial function

$$f(x) = \frac{1}{24}x^4 - \frac{1}{4}x^3 + \frac{23}{24}x^2 - \frac{3}{4}x + 1$$


will give the maximum number of interior regions formed in a circle if x points on the circumference are joined by all possible chords. For $x = 1, 2, 3, 4,$ and 5 , see FIGURES A–E.




$x = 1$
FIGURE A




$x = 2$
FIGURE B



$x = 3$
FIGURE C



$x = 4$
FIGURE D



$x = 5$
FIGURE E

For example, in FIGURE A we have 1 point, and because no chords can be drawn, we have only 1 interior region. In FIGURE B there are 2 points, and 2 interior regions are formed.

Apply this information and work Exercises 97–100 in order.

97. Use the given polynomial function to verify that $f(1) = 1$ and $f(2) = 2$.
98. Based on the appropriate figure alone, what is a logical prediction for the value of $f(3)$? $f(4)$? $f(5)$?
99. Verify the answers in Exercise 98 by evaluating $f(3)$, $f(4)$, and $f(5)$.
100. Observe a pattern in the results of Exercises 97–99. Use the pattern to make a logical prediction for the value of $f(6)$. Does the prediction equal $f(6)$?

4.5 Multiplying Polynomials

OBJECTIVES

- 1 Multiply terms.
- 2 Multiply any two polynomials.
- 3 Multiply binomials.
- 4 Find the product of a sum and difference of two terms.
- 5 Find the square of a binomial.
- 6 Multiply polynomial functions.

NOW TRY EXERCISE 1

Find the product.

$$-3s^2t(15s^3t^4)$$

NOW TRY EXERCISE 2

Find each product.

(a) $3k^3(-2k^5 + 3k^2 - 4)$

(b) $5x(2x - 1)(x + 4)$

NOW TRY ANSWERS

1. $-45s^5t^5$
2. (a) $-6k^8 + 9k^5 - 12k^3$
(b) $10x^3 + 35x^2 - 20x$

OBJECTIVE 1 Multiply terms.

EXAMPLE 1 Multiplying Monomials

Find each product.

(a) $3x^4(5x^3)$

$$= 3(5)x^4 \cdot x^3 \quad \text{Commutative and associative properties}$$

$$= 15x^{4+3} \quad \text{Multiply; product rule for exponents}$$

$$= 15x^7 \quad \text{Add the exponents.}$$

(b) $-4a^3(3a^5)$

$$= -4(3)a^3 \cdot a^5$$

$$= -12a^8$$

(c) $2m^2z^4(8m^3z^2)$

$$= 2(8)m^2 \cdot m^3 \cdot z^4 \cdot z^2$$

$$= 16m^5z^6$$

NOW TRY 

OBJECTIVE 2 Multiply any two polynomials.

EXAMPLE 2 Multiplying Polynomials

Find each product.

(a) $-2(8x^3 - 9x^2)$

$$= -2(8x^3) - 2(-9x^2) \quad \text{Distributive property}$$

$$= -16x^3 + 18x^2 \quad \text{Multiply.}$$

Be careful with signs.

(b) $5x^2(-4x^2 + 3x - 2)$

$$= 5x^2(-4x^2) + 5x^2(3x) + 5x^2(-2) \quad \text{Distributive property}$$

$$= -20x^4 + 15x^3 - 10x^2 \quad \text{Multiply.}$$

(c) $(5x + 4)(2x^2 + x)$

$$= 5x(2x^2 + x) + 4(2x^2 + x)$$

$$= 10x^3 + 5x^2 + 8x^2 + 4x \quad \text{Distributive property}$$

$$= 10x^3 + 13x^2 + 4x \quad \text{Combine like terms.}$$

Treat $2x^2 + x$ as a single expression.

(d) $2x^2(x + 1)(x - 3)$

$$= 2x^2[x(x - 3) + 1(x - 3)] \quad \text{Distributive property}$$

$$= 2x^2[x^2 - 3x + x - 3] \quad \text{Distributive property}$$

$$= 2x^2(x^2 - 2x - 3) \quad \text{Combine like terms.}$$

$$= 2x^4 - 4x^3 - 6x^2 \quad \text{Distributive property}$$

NOW TRY 


**NOW TRY
EXERCISE 3**

Find the product.

$$\begin{array}{r} 3t^2 - 5t + 4 \\ \underline{ t - 3} \end{array}$$

EXAMPLE 3 Multiplying Polynomials Vertically

Find each product.

(a) $(5a - 2b)(3a + b)$

$$\begin{array}{r} 5a - 2b \\ 3a + b \\ \hline \end{array}$$

Write the factors vertically.

$$\begin{array}{r} 5a - 2b \\ 3a + b \\ \hline \end{array}$$

$$5ab - 2b^2 \leftarrow \text{Multiply } b(5a - 2b).$$

$$\begin{array}{r} 15a^2 - 6ab \\ \hline \end{array} \leftarrow \text{Multiply } 3a(5a - 2b).$$

$$\begin{array}{r} 15a^2 - ab - 2b^2 \\ \hline \end{array} \leftarrow \text{Combine like terms.}$$

(b) $(3m^3 - 2m^2 + 4)(3m - 5)$

$$\begin{array}{r} 3m^3 - 2m^2 + 4 \\ 3m - 5 \\ \hline \end{array}$$

Be sure to write like terms in columns.

$$\begin{array}{r} 3m^3 - 2m^2 + 4 \\ 3m - 5 \\ \hline \end{array}$$

$$\begin{array}{r} 3m^3 - 2m^2 + 4 \\ 3m - 5 \\ \hline -15m^3 + 10m^2 - 20 \end{array} \leftarrow \text{Multiply } -5(3m^3 - 2m^2 + 4).$$

$$\begin{array}{r} 9m^4 - 6m^3 + 12m \\ \hline \end{array} \leftarrow \text{Multiply } 3m(3m^3 - 2m^2 + 4).$$

$$\begin{array}{r} 9m^4 - 21m^3 + 10m^2 + 12m - 20 \\ \hline \end{array} \leftarrow \text{Combine like terms.}$$

NOW TRY **NOTE** We can use a rectangle to model polynomial multiplication.

$$(5a - 2b)(3a + b) \quad \text{See Example 3(a).}$$

First, we label a rectangle with each term, as shown below on the left. Then we write the product of each pair of monomials in the appropriate box, as shown on the right.

	3a	b
5a		
-2b		

	3a	b
5a	15a ²	5ab
-2b	-6ab	-2b ²

$$(5a - 2b)(3a + b)$$

$$= 15a^2 + 5ab - 6ab - 2b^2 \quad \text{Add the four monomial products.}$$

$$= 15a^2 - ab - 2b^2 \quad \text{The result is the same as in Example 3(a).}$$

OBJECTIVE 3 Multiply binomials.There is a shortcut method for finding the product of two binomials. In **Example 2**, we found such products using the distributive property.

$$(3x - 4)(2x + 3)$$

$$= 3x(2x + 3) - 4(2x + 3) \quad \text{Distributive property}$$

$$= 3x(2x) + 3x(3) - 4(2x) - 4(3) \quad \text{Distributive property again}$$

$$= 6x^2 + 9x - 8x - 12 \quad \text{Multiply.}$$

Before combining like terms to find the simplest form of the answer, we check the origin of each of the four terms in the sum

$$6x^2 + 9x - 8x - 12.$$

NOW TRY ANSWER

3. $3t^3 - 14t^2 + 19t - 12$

From the discussion on the preceding page, multiplying

$$(3x - 4)(2x + 3)$$

using the distributive property gave

$$6x^2 + 9x - 8x - 12.$$

The term $6x^2$ is the product of the two *first* terms of the binomials.

$$(3x - 4)(2x + 3) \quad 3x(2x) = 6x^2 \quad \text{First terms}$$

To obtain $9x$, the *outer* terms are multiplied.

$$(3x - 4)(2x + 3) \quad 3x(3) = 9x \quad \text{Outer terms}$$

The term $-8x$ comes from the *inner* terms.

$$(3x - 4)(2x + 3) \quad -4(2x) = -8x \quad \text{Inner terms}$$

Finally, -12 comes from the *last* terms.

$$(3x - 4)(2x + 3) \quad -4(3) = -12 \quad \text{Last terms}$$

The product is found by adding these four results.

$$\begin{aligned} (3x - 4)(2x + 3) \\ &= 6x^2 + 9x - 8x - 12 \quad \text{FOIL method} \\ &= 6x^2 + x - 12 \quad \text{Combine like terms.} \end{aligned}$$

To keep track of the order of multiplying terms, we use the initials FOIL, where the letters refer to the positions of the terms (**F**irst, **O**uter, **I**nnner, **L**ast). The above steps can be done as follows.

Compact form:

$$\begin{array}{ccc} \text{First} & & \text{Last} \\ (3x - 4)(2x + 3) & & (3x - 4)(2x + 3) \\ & \text{Inner} & \\ & \text{Outer} & \\ & & \begin{array}{c} 6x^2 \quad -12 \\ -8x \\ 9x \\ \hline x \quad \text{Add.} \end{array} \end{array}$$

Try to do as many of these steps as possible mentally.

CAUTION The FOIL method applies only to multiplying two binomials.

EXAMPLE 4 Using the FOIL Method

Use the FOIL method to find each product.

(a) $(4m - 5)(3m + 1)$

First terms $(4m - 5)(3m + 1) \quad 4m(3m) = 12m^2$

Outer terms $(4m - 5)(3m + 1) \quad 4m(1) = 4m$

Inner terms $(4m - 5)(3m + 1) \quad -5(3m) = -15m$

Last terms $(4m - 5)(3m + 1) \quad -5(1) = -5$

$$\begin{aligned} (4m - 5)(3m + 1) \\ &= 12m^2 + 4m - 15m - 5 \quad \text{FOIL method} \\ &= 12m^2 - 11m - 5 \quad \text{Combine like terms.} \end{aligned}$$


NOW TRY EXERCISE 4

Use the FOIL method to find each product.

- (a) $(2x + 1)(7x + 2)$
 (b) $(3p - k)(5p + 4k)$

Compact form: $(4m - 5)(3m + 1)$

$$\begin{array}{r}
 \begin{array}{cc}
 12m^2 & -5 \\
 \hline
 -15m & 4m \\
 \hline
 -11m &
 \end{array} \\
 \text{Add.}
 \end{array}$$

Add these results to obtain

$$12m^2 - 11m - 5.$$

(b) $(5x - 4)(2x - 5)$

$$\begin{array}{cccc}
 \text{First} & \text{Outer} & \text{Inner} & \text{Last} \\
 \downarrow & \downarrow & \downarrow & \downarrow \\
 = 5x(2x) + 5x(-5) - 4(2x) - 4(-5) & \text{FOIL method} \\
 = 10x^2 - 25x - 8x + 20 & \text{Multiply.} \\
 = 10x^2 - 33x + 20 & \text{Combine like terms.}
 \end{array}$$

(c) $(6a + 5b)(3a - 4b)$

$$\begin{array}{cccc}
 \text{First} & \text{Outer} & \text{Inner} & \text{Last} \\
 \downarrow & \downarrow & \downarrow & \downarrow \\
 = 18a^2 - 24ab + 15ab - 20b^2 & \text{FOIL method} \\
 = 18a^2 - 9ab - 20b^2 & \text{Combine like terms.}
 \end{array}$$

(d) $(2k + 3z)(5k + 3z)$

$$\begin{array}{l}
 = 10k^2 + 6kz + 15kz + 9z^2 \quad \text{FOIL method} \\
 = 10k^2 + 21kz + 9z^2 \quad \text{Combine like terms.}
 \end{array}$$

NOW TRY 

OBJECTIVE 4 Find the product of a sum and difference of two terms.

The product of a sum and difference of the same two terms occurs frequently.

$$\begin{array}{l}
 (x + y)(x - y) \\
 = x^2 - xy + xy - y^2 \quad \text{FOIL method} \\
 = x^2 - y^2 \quad \text{Combine like terms.}
 \end{array}$$

Product of a Sum and Difference of Two Terms

The **product of the sum and difference of the two terms x and y** is the difference of the squares of the terms.

$$(x + y)(x - y) = x^2 - y^2$$


NOW TRY EXERCISE 5

Find each product.

- (a) $(x + 8)(x - 8)$
 (b) $(3x - 7y)(3x + 7y)$
 (c) $5k(2k - 3)(2k + 3)$

NOW TRY ANSWERS

4. (a) $14x^2 + 11x + 2$
 (b) $15p^2 + 7kp - 4k^2$
 5. (a) $x^2 - 64$
 (b) $9x^2 - 49y^2$
 (c) $20k^3 - 45k$

EXAMPLE 5 Multiplying a Sum and Difference of Two Terms

Find each product.

(a) $(p + 7)(p - 7)$

$$\begin{array}{l}
 = p^2 - 7^2 \\
 = p^2 - 49
 \end{array}$$

(b) $(2r + 5)(2r - 5)$

$$\begin{array}{l}
 = (2r)^2 - 5^2 \\
 = 4r^2 - 25 \quad (2r)^2 = 2^2r^2
 \end{array}$$

Be careful.
 $(ab)^2 = a^2b^2$

(c) $(6m + 5n)(6m - 5n)$

$$\begin{array}{l}
 = (6m)^2 - (5n)^2 \\
 = 36m^2 - 25n^2
 \end{array}$$

$(6m)^2 = 6^2m^2$;
 $(5n)^2 = 5^2n^2$

(d) $2x^3(x + 3)(x - 3)$

$$\begin{array}{l}
 = 2x^3(x^2 - 9) \\
 = 2x^5 - 18x^3
 \end{array}$$

NOW TRY 

OBJECTIVE 5 Find the square of a binomial.

To find the square of a binomial sum $x + y$ —that is, $(x + y)^2$ —multiply $x + y$ by itself.

$$\begin{aligned}(x + y)^2 &= (x + y)(x + y) && a^2 = a \cdot a \\ &= x^2 + xy + xy + y^2 && \text{FOIL method} \\ &= x^2 + 2xy + y^2 && \text{Combine like terms.}\end{aligned}$$

A similar result is true for the square of a binomial difference.

Square of a Binomial

The **square of a binomial** is the sum of the square of the first term, twice the product of the two terms, and the square of the last term.

$$\begin{aligned}(x + y)^2 &= x^2 + 2xy + y^2 \\ (x - y)^2 &= x^2 - 2xy + y^2\end{aligned}$$

 **NOW TRY EXERCISE 6**

Find each product.

- (a) $(y - 10)^2$
(b) $(4x + 5y)^2$

EXAMPLE 6 Squaring Binomials

Find each product.

(a) $(m + 7)^2$

$$\begin{aligned}&= m^2 + 2 \cdot m \cdot 7 + 7^2 && (x + y)^2 = x^2 + 2xy + y^2 \\ &= m^2 + 14m + 49 && \text{Multiply. Apply the exponent.}\end{aligned}$$

(b) $(p - 4)^2$

$$\begin{aligned}&= p^2 - 2 \cdot p \cdot 4 + 4^2 && (x - y)^2 = x^2 - 2xy + y^2 \\ &= p^2 - 8p + 16 && \text{Multiply. Apply the exponent.}\end{aligned}$$

(c) $(2p + 3v)^2$

$$\begin{aligned}&= (2p)^2 + 2(2p)(3v) + (3v)^2 \\ &= 4p^2 + 12pv + 9v^2\end{aligned}$$

(d) $(3r - 5s)^2$

$$\begin{aligned}&= (3r)^2 - 2(3r)(5s) + (5s)^2 \\ &= 9r^2 - 30rs + 25s^2\end{aligned}$$

NOW TRY 

CAUTION As the products in the formula for the square of a binomial show,

$$(x + y)^2 \neq x^2 + y^2.$$

In general for $n \neq 1$, $(x + y)^n \neq x^n + y^n$.

EXAMPLE 7 Multiplying More Complicated Binomials

Find each product.

(a) $[(3p - 2) + 5q][(3p - 2) - 5q]$

$$\begin{aligned}&= (3p - 2)^2 - (5q)^2 && \text{Product of a sum and difference of terms} \\ &= 9p^2 - 12p + 4 - 25q^2 && \text{Square both quantities.}\end{aligned}$$

(b) $[(2z + r) + 1]^2$

$$\begin{aligned}&= (2z + r)^2 + 2(2z + r)(1) + 1^2 && \text{Square of a binomial} \\ &= 4z^2 + 4zr + r^2 + 4z + 2r + 1 && \text{Square again; distributive property}\end{aligned}$$

NOW TRY ANSWERS

6. (a) $y^2 - 20y + 100$
(b) $16x^2 + 40xy + 25y^2$


NOW TRY EXERCISE 7

Find each product.

(a) $[(4x - y) + 2][(4x - y) - 2]$

(b) $(y - 3)^4$

(c) $(x + y)^3$

This does not equal $x^3 + y^3$.

$$\begin{aligned}
 &= (x + y)(x + y)^2 && a^3 = a \cdot a^2 \\
 &= (x + y)(x^2 + 2xy + y^2) && \text{Square } x + y. \\
 &= x(x^2 + 2xy + y^2) + y(x^2 + 2xy + y^2) && \text{Distributive property} \\
 &= x^3 + 2x^2y + xy^2 + x^2y + 2xy^2 + y^3 && \text{Distributive property} \\
 &= x^3 + 3x^2y + 3xy^2 + y^3 && \text{Combine like terms.}
 \end{aligned}$$

(d) $(2a + b)^4$

$$\begin{aligned}
 &= (2a + b)^2(2a + b)^2 && a^4 = a^2 \cdot a^2 \\
 &= (4a^2 + 4ab + b^2)(4a^2 + 4ab + b^2) && \text{Square } 2a + b \text{ twice.} \\
 &= 4a^2(4a^2 + 4ab + b^2) + 4ab(4a^2 + 4ab + b^2) && \text{Distributive property} \\
 &\quad + b^2(4a^2 + 4ab + b^2) \\
 &= 16a^4 + 16a^3b + 4a^2b^2 + 16a^3b + 16a^2b^2 + 4ab^3 && \text{Distributive property again} \\
 &\quad + 4a^2b^2 + 4ab^3 + b^4 \\
 &= 16a^4 + 32a^3b + 24a^2b^2 + 8ab^3 + b^4 && \text{Combine like terms.}
 \end{aligned}$$

NOW TRY 
OBJECTIVE 6 Multiply polynomial functions.

Previously, we added and subtracted functions. They can also be multiplied.

Multiplying Functions
If $f(x)$ and $g(x)$ define functions, then

$$(fg)(x) = f(x) \cdot g(x). \quad \text{Product function}$$

The domain of the product function is the intersection of the domains of $f(x)$ and $g(x)$.

NOW TRY EXERCISE 8

For $f(x) = 3x^2 - 1$

and $g(x) = 8x + 7$,

find $(fg)(x)$ and $(fg)(-2)$.

EXAMPLE 8 Multiplying Polynomial Functions
For $f(x) = 3x + 4$ and $g(x) = 2x^2 + x$, find $(fg)(x)$ and $(fg)(-1)$.

(f) $(fg)(x)$

This notation indicates function multiplication.

$$\begin{aligned}
 &= f(x) \cdot g(x) && \text{Use the definition.} \\
 &= (3x + 4)(2x^2 + x) && \text{Substitute.} \\
 &= 6x^3 + 3x^2 + 8x^2 + 4x && \text{FOIL method} \\
 &= 6x^3 + 11x^2 + 4x && \text{Combine like terms.}
 \end{aligned}$$

(g) $(fg)(-1)$

$$\begin{aligned}
 &= 6(-1)^3 + 11(-1)^2 + 4(-1) && (fg)(x) = 6x^3 + 11x^2 + 4x \\
 &= -6 + 11 - 4 && \text{Let } x = -1 \text{ in } (fg)(x). \\
 &= 1 && \text{Apply the exponents. Multiply.} \\
 & && \text{Add and subtract.}
 \end{aligned}$$

Be careful with signs.

NOW TRY ANSWERS

7. (a) $16x^2 - 8xy + y^2 - 4$

(b) $y^4 - 12y^3 + 54y^2 - 108y + 81$

8. $24x^3 + 21x^2 - 8x - 7; -99$

An alternative method for finding $(fg)(-1)$ is to find $f(-1)$ and $g(-1)$ and then multiply the results. Verify that $f(-1) \cdot g(-1) = 1$. This follows from the definition.
NOW TRY 

CAUTION Write the product $f(x) \cdot g(x)$ as $(fg)(x)$, **not** $f(g(x))$, which indicates the composition of functions f and g .

4.5 Exercises

FOR
EXTRA
HELP



MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Match each product in Column I with the correct polynomial in Column II.

I	II
1. $(2x - 5)(3x + 4)$	A. $6x^2 + 23x + 20$
2. $(2x + 5)(3x + 4)$	B. $6x^2 + 7x - 20$
3. $(2x - 5)(3x - 4)$	C. $6x^2 - 7x - 20$
4. $(2x + 5)(3x - 4)$	D. $6x^2 - 23x + 20$

Find each product. See Examples 1–3.

5. $-8m^3(3m^2)$	6. $-4p^2(5p^4)$	7. $14x^2y^3(-2x^5y)$
8. $5m^3n^4(-4m^2n^5)$	9. $3x(-2x + 5)$	10. $5y(-6y + 1)$
11. $-q^3(2 + 3q)$	12. $-3a^4(4 + a)$	13. $6k^2(3k^2 + 2k + 1)$
14. $5r^3(2r^2 + 3r + 4)$	15. $(5x + 1)(3x^2 + x)$	16. $(9a + 2)(4a^2 + 3a)$
17. $(2y + 3)(3y - 4)$	18. $(2m + 6)(5m - 3)$	19. $m(m + 5)(m - 8)$
20. $p(p + 4)(p - 6)$	21. $4z(2z + 1)(3z - 4)$	22. $2y(2y + 1)(8y - 3)$
23. $4x^3(x - 3)(x + 2)$	24. $2y^5(y - 8)(y + 2)$	25. $(2t + 3)(3t^2 - 4t - 1)$
26. $(4z + 2)(z^2 - 3z - 5)$	27. $5m - 3n$	28. $2k + 6q$
	<u>$5m + 3n$</u>	<u>$2k - 6q$</u>
29. $-b^2 + 3b + 3$	30. $-r^2 - 4r + 8$	31. $2p^2 + 3p + 6$
<u>$2b + 4$</u>	<u>$3r - 2$</u>	<u>$3p^2 - 4p - 1$</u>
32. $5y^2 - 2y + 4$	33. $2z^3 - 5z^2 + 8z - 1$	34. $3z^4 - 2z^3 + z - 5$
<u>$2y^2 + y + 3$</u>	<u>$4z + 3$</u>	<u>$2z - 5$</u>

Use the FOIL method to find each product. See Example 4.

35. $(m + 5)(m - 8)$	36. $(p + 4)(p - 6)$	37. $(4k + 3)(3k - 2)$
38. $(5w + 2)(2w - 5)$	39. $(3x - 2)(5x - 1)$	40. $(4x - 1)(6x - 7)$
41. $(5x + 2y)(4x + 3y)$	42. $(3x + 4y)(2x + 3y)$	43. $(z - w)(3z + 4w)$
44. $(s - t)(2s + 5t)$	45. $(6c - d)(2c + 3d)$	46. $(2m - n)(3m + 5n)$

47. Concept Check A student multiplied incorrectly as follows.

$$(x + 4)^2 = x^2 + 16$$

WHAT WENT WRONG? Give the correct product.

48. Concept Check A student multiplied incorrectly as follows.

$$(x - 9)(x + 9) = x^2 + 81$$

WHAT WENT WRONG? Give the correct product.

Find each product. See Example 5.

49. $(x + 9)(x - 9)$ 50. $(z + 6)(z - 6)$ 51. $(2p - 3)(2p + 3)$
 52. $(3x - 8)(3x + 8)$ 53. $(5m - 1)(5m + 1)$ 54. $(6y - 3)(6y + 3)$
 55. $(3a + 2c)(3a - 2c)$ 56. $(5r + 4s)(5r - 4s)$ 57. $(4m + 7n^2)(4m - 7n^2)$
 58. $(2k^2 + 6h)(2k^2 - 6h)$ 59. $3y(5y^3 + 2)(5y^3 - 2)$ 60. $4x(3x^3 + 4)(3x^3 - 4)$

Find each product. See Example 6.

61. $(y - 5)^2$ 62. $(a - 3)^2$ 63. $(x + 1)^2$ 64. $(t + 2)^2$
 65. $(2p + 7)^2$ 66. $(3z + 8)^2$ 67. $(4n - 3m)^2$ 68. $(5r - 7s)^2$

Extending Skills The factors in the following exercises involve fractions or decimals. Apply the methods of this section, and find each product.

69. $\left(k - \frac{5}{7}p\right)^2$ 70. $\left(q - \frac{3}{4}r\right)^2$ 71. $(0.2x - 1.4y)^2$
 72. $(0.3x - 1.6y)^2$ 73. $\left(4x - \frac{2}{3}\right)\left(4x + \frac{2}{3}\right)$ 74. $\left(3t - \frac{5}{4}\right)\left(3t + \frac{5}{4}\right)$
 75. $(0.2x + 1.3)(0.5x - 0.1)$ 76. $(0.1y + 2.1)(0.5y - 0.4)$
 77. $\left(3w + \frac{1}{4}z\right)(w - 2z)$ 78. $\left(5r + \frac{2}{3}y\right)(r - 5y)$



Find each product. See Example 7.



79. $[(5x + 1) + 6y]^2$ 80. $[(3m - 2) + p]^2$
 81. $[(2a + b) - 3]^2$ 82. $[(4k + h) - 4]^2$
 83. $[(2a + b) - 3][(2a + b) + 3]$ 84. $[(m + p) - 5][(m + p) + 5]$
 85. $[(2h - k) + j][(2h - k) - j]$ 86. $[(3m - y) + z][(3m - y) - z]$
 87. $(y + 2)^3$ 88. $(z - 3)^3$ 89. $(5r - s)^3$
 90. $(x + 3y)^3$ 91. $(q - 2)^4$ 92. $(r + 3)^4$

Extending Skills Find each product.

93. $(2a + b)(3a^2 + 2ab + b^2)$ 94. $(m - 5p)(m^2 - 2mp + 3p^2)$
 95. $(4z - x)(z^3 - 4z^2x + 2zx^2 - x^3)$ 96. $(3r + 2s)(r^3 + 2r^2s - rs^2 + 2s^3)$
 97. $(m^2 - 2mp + p^2)(m^2 + 2mp - p^2)$ 98. $(3 + x + y)(-3 + x - y)$
 99. $ab(a + b)(a + 2b)(a - 3b)$ 100. $mp(m - p)(m - 2p)(2m + p)$

Find the area of each figure. Express it as a polynomial in descending powers of the variable x . Refer to the formulas at the back of this text if necessary.





For each pair of functions, find $(fg)(x)$. See Example 8.

105. $f(x) = 2x$, $g(x) = 5x - 1$

106. $f(x) = 3x$, $g(x) = 6x - 8$

107. $f(x) = x + 1$, $g(x) = 2x - 3$

108. $f(x) = x - 7$, $g(x) = 4x + 5$

109. $f(x) = 2x - 3$, $g(x) = 4x^2 + 6x + 9$

110. $f(x) = 3x + 4$, $g(x) = 9x^2 - 12x + 16$

Let $f(x) = x^2 - 9$, $g(x) = 2x$, and $h(x) = x - 3$. Find each of the following. See Example 8.

111. $(fg)(x)$

112. $(fh)(x)$

113. $(fg)(2)$

114. $(fh)(1)$

115. $(fh)(-1)$

116. $(gh)(-2)$

117. $(gh)(-3)$

118. $(fg)(-2)$

119. $(fg)(0)$

120. $(fh)(0)$

121. $(fg)\left(-\frac{1}{2}\right)$

122. $(fg)\left(-\frac{1}{3}\right)$

RELATING CONCEPTS For Individual or Group Work (Exercises 123–130)

Consider the figure. Work Exercises 123–130 in order.

123. What is the length of each side of the blue square in terms of a and b ?

124. What is the formula for the area of a square? Use the formula to write an expression, in the form of a product, for the area of the blue square.

125. Each green rectangle has an area of _____. Therefore, the total area in green is represented by the polynomial _____.

126. The yellow square has an area of _____.


127. The area of the entire colored region is represented by _____, because each side of the entire colored region has length _____.

128. The area of the blue square is equal to the area of the entire colored region, minus the total area of the green squares, minus the area of the yellow square. Write this as a simplified polynomial in a and b .

129. (a) What must be true about the expressions for the area of the blue square found in Exercises 124 and 128?

(b) Write an equation based on the answer in part (a).

130. Draw a figure and give a similar proof for $(a + b)^2 = a^2 + 2ab + b^2$.



4.6 Dividing Polynomials

OBJECTIVES

- 1 Divide a polynomial by a monomial.
- 2 Divide a polynomial by a polynomial of two or more terms.
- 3 Divide polynomial functions.

OBJECTIVE 1 Divide a polynomial by a monomial.

Recall that a monomial is a single term, such as 3 , $5m^2$, or x^2y^2 .

Dividing a Polynomial by a Monomial

To divide a polynomial by a monomial, divide each term of the polynomial by the monomial.

$$\frac{a + b}{c} = \frac{a}{c} + \frac{b}{c} \quad (\text{where } c \neq 0)$$

Then write each quotient in lowest terms.

EXAMPLE 1 Dividing Polynomials by Monomials

Divide.

$$\begin{aligned} \text{(a)} \quad & \frac{15x^2 - 12x + 3}{3} && \text{Do not divide out the 3's.} \\ & = \frac{15x^2}{3} - \frac{12x}{3} + \frac{3}{3} && \text{Divide each term by 3.} \\ & = 5x^2 - 4x + 1 && \text{Write in lowest terms.} \end{aligned}$$

$$\begin{array}{c} \text{CHECK} \quad 3(5x^2 - 4x + 1) = 15x^2 - 12x + 3 \quad \checkmark \\ \begin{array}{ccc} \uparrow & \uparrow & \uparrow \\ \text{Divisor} & \text{Quotient} & \text{Original polynomial (Dividend)} \end{array} \end{array}$$

$$\begin{aligned} \text{(b)} \quad & \frac{5m^3 - 9m^2 + 10m}{5m^2} \\ & = \frac{5m^3}{5m^2} - \frac{9m^2}{5m^2} + \frac{10m}{5m^2} && \text{Think: } \frac{10m}{5m^2} = \frac{10}{5} m^{1-2} = 2m^{-1} = \frac{2}{m} \\ & = m - \frac{9}{5} + \frac{2}{m} && \text{Divide each term by } 5m^2. \\ & && \text{Simplify each term.} \\ & && \text{Use the quotient rule for exponents.} \end{aligned}$$

$$\text{CHECK} \quad 5m^2 \left(m - \frac{9}{5} + \frac{2}{m} \right) = 5m^3 - 9m^2 + 10m \quad \checkmark \quad \text{Divisor} \times \text{Quotient} = \text{Original polynomial}$$

The result $m - \frac{9}{5} + \frac{2}{m}$ is not a polynomial because the last term has a variable in its denominator. The quotient of two polynomials need not be a polynomial.

$$\begin{aligned} \text{(c)} \quad & \frac{8xy^2 - 9x^2y + 6x^2y^2}{x^2y^2} \\ & = \frac{8xy^2}{x^2y^2} - \frac{9x^2y}{x^2y^2} + \frac{6x^2y^2}{x^2y^2} && \text{Divide each term by } x^2y^2. \\ & = \frac{8}{x} - \frac{9}{y} + 6 && \frac{a^m}{a^n} = a^{m-n} \end{aligned}$$

NOW TRY EXERCISE 1

Divide.

$$\begin{aligned} \text{(a)} \quad & \frac{16x^3 - 8x^2 - 4}{4} \\ \text{(b)} \quad & \frac{81y^4 - 54y^3 + 18y}{9y^3} \end{aligned}$$

NOW TRY ANSWERS

1. (a) $4x^3 - 2x^2 - 1$
- (b) $9y - 6 + \frac{2}{y^2}$

OBJECTIVE 2 Divide a polynomial by a polynomial of two or more terms.

This process is similar to that for dividing whole numbers.

$$\begin{array}{r} 23 \\ 25 \overline{)581} \\ \underline{-50} \\ 81 \\ \underline{-75} \\ 6 \end{array} \qquad 581 \div 25 = 23 \frac{6}{25} \quad \begin{array}{l} \text{Dividend} \\ \downarrow \end{array}$$

CHECK $25 \times 23 + 6 = 581 \quad \checkmark$

$\begin{array}{ccc} \uparrow & \uparrow & \uparrow \\ \text{Divisor} & \text{Quotient} & \text{Remainder} \end{array}$


NOW TRY EXERCISE 2

Divide.

$$\frac{3x^2 - 4x - 15}{x - 3}$$

EXAMPLE 2 Dividing a Polynomial by a Polynomial

Divide $\frac{2x^2 + x - 10}{x - 2}$.

Make sure each polynomial is written in descending powers of the variables.

$$x - 2 \overline{)2x^2 + x - 10} \quad \begin{array}{l} \text{Write as if dividing} \\ \text{whole numbers.} \end{array}$$

Divide the first term of the dividend $2x^2 + x - 10$ by the first term of the divisor $x - 2$. Here $\frac{2x^2}{x} = 2x$.

$$\begin{array}{r} 2x \\ x - 2 \overline{)2x^2 + x - 10} \end{array} \quad \leftarrow \text{Result of } \frac{2x^2}{x}$$

Multiply $x - 2$ and $2x$, and write the result below $2x^2 + x - 10$.

$$\begin{array}{r} 2x \\ x - 2 \overline{)2x^2 + x - 10} \\ \underline{2x^2 - 4x} \end{array} \quad \leftarrow 2x(x - 2) = 2x^2 - 4x$$

Now subtract by mentally changing the signs on $2x^2 - 4x$ and *adding*.

$$\begin{array}{r} 2x \\ x - 2 \overline{)2x^2 + x - 10} \\ \underline{2x^2 - 4x} \\ 5x \end{array} \quad \leftarrow \text{Subtract. The difference is } 5x.$$

Subtract.
 $\begin{array}{r} 2x^2 + x \\ - (2x^2 - 4x) \end{array}$

Bring down -10 and continue by dividing $5x$ by x .

$$\begin{array}{r} 2x + 5 \\ x - 2 \overline{)2x^2 + x - 10} \\ \underline{2x^2 - 4x} \\ 5x - 10 \\ \underline{5x - 10} \\ 0 \end{array} \quad \begin{array}{l} \leftarrow \frac{5x}{x} = 5 \\ \leftarrow \text{Bring down } -10. \\ \leftarrow 5(x - 2) = 5x - 10 \\ \leftarrow \text{Subtract. The difference is } 0. \end{array}$$

Think:
 $\begin{array}{r} 5x - 10 \\ - (5x - 10) \end{array}$

CHECK Multiply $x - 2$ (the divisor) and $2x + 5$ (the quotient).

$$\begin{aligned} (x - 2)(2x + 5) &= 2x^2 + 5x - 4x - 10 && \text{FOIL method} \\ &= 2x^2 + x - 10 && \text{Combine like terms.} \end{aligned}$$

The result is $2x^2 + x - 10$ (the dividend). \checkmark

NOW TRY ANSWER

2. $3x + 5$

NOW TRY
EXERCISE 3

Divide

$$2x^3 - 12x - 10 \text{ by } x - 4.$$

EXAMPLE 3 Dividing a Polynomial with a Missing Term

Divide $3x^3 - 2x + 5$ by $x - 3$.

Add a term with 0 coefficient as a placeholder for the missing x^2 -term.

$$x - 3 \overline{) 3x^3 + 0x^2 - 2x + 5}$$

Missing term
Descending powers of the variable

Start with $\frac{3x^3}{x} = 3x^2$.

$$x - 3 \overline{) 3x^3 + 0x^2 - 2x + 5}$$

$$\underline{3x^3 - 9x^2}$$

$\frac{3x^3}{x} = 3x^2$
 $3x^2(x - 3)$

Subtract by mentally changing the signs on $3x^3 - 9x^2$ and adding.

$$x - 3 \overline{) 3x^3 + 0x^2 - 2x + 5}$$

$$\underline{3x^3 - 9x^2}$$

$$9x^2$$

Subtract.

Bring down the next term.

$$x - 3 \overline{) 3x^3 + 0x^2 - 2x + 5}$$

$$\underline{3x^3 - 9x^2}$$

$$9x^2 - 2x$$

Bring down $-2x$.

In the next step, $\frac{9x^2}{x} = 9x$.

$$x - 3 \overline{) 3x^3 + 0x^2 - 2x + 5}$$

$$\underline{3x^3 - 9x^2}$$

$$9x^2 - 2x$$

$$\underline{9x^2 - 27x}$$

$$25x + 5$$

$\frac{9x^2}{x} = 9x$
 $9x(x - 3)$
Subtract. Bring down 5.

Finally, $\frac{25x}{x} = 25$.

$$x - 3 \overline{) 3x^3 + 0x^2 - 2x + 5}$$

$$\underline{3x^3 - 9x^2}$$

$$9x^2 - 2x$$

$$\underline{9x^2 - 27x}$$

$$25x + 5$$

$$\underline{25x - 75}$$

$$80$$

$\frac{25x}{x} = 25$
 $25(x - 3)$
Remainder

Write the remainder, 80, as the numerator of a fraction, $\frac{80}{x - 3}$.

$$3x^2 + 9x + 25 + \frac{80}{x - 3}$$

Be sure to add remainder/divisor.
Don't forget the + sign.

NOW TRY ANSWER

3. $2x^2 + 8x + 20 + \frac{70}{x - 4}$

CHECK Multiply $x - 3$ (the divisor) and $3x^2 + 9x + 25$ (the quotient), and then add 80 (the remainder). The result is $3x^3 - 2x + 5$ (the dividend). ✓


NOW TRY EXERCISE 4

Divide

$$2x^4 + 8x^3 + 2x^2 - 5x - 3$$

by $2x^2 - 2$.

EXAMPLE 4 Dividing by a Polynomial with a Missing Term
Divide $6r^4 + 9r^3 + 2r^2 - 8r + 7$ by $3r^2 - 2$.

$$\begin{array}{r}
 2r^2 + 3r + 2 \\
 3r^2 + 0r - 2 \overline{) 6r^4 + 9r^3 + 2r^2 - 8r + 7} \\
 \underline{6r^4 + 0r^3 - 4r^2} \\
 9r^3 + 6r^2 - 8r \\
 \underline{9r^3 + 0r^2 - 6r} \\
 6r^2 - 2r + 7 \\
 \underline{6r^2 + 0r - 4} \\
 -2r + 11 \leftarrow \text{Remainder}
 \end{array}$$

Missing term \longrightarrow \uparrow

Stop when the degree of the remainder is less than the degree of the divisor.

The degree of the remainder, $-2r + 11$, is less than the degree of the divisor, $3r^2 - 2$, so the division process is finished. The result is written as follows.

$$2r^2 + 3r + 2 + \frac{-2r + 11}{3r^2 - 2} \quad \text{Quotient} + \frac{\text{remainder}}{\text{divisor}}$$

CHECK $(3r^2 - 2)(2r^2 + 3r + 2) + (-2r + 11)$

Divisor \times Quotient + Remainder

$$= 6r^4 + 9r^3 + 6r^2 - 4r^2 - 6r - 4 - 2r + 11$$

$$= 6r^4 + 9r^3 + 2r^2 - 8r + 7 \quad \checkmark \quad \text{Original polynomial (Dividend)}$$

NOW TRY 
! CAUTION When dividing a polynomial by a polynomial of two or more terms:

1. Be sure the terms in both polynomials are in descending powers.
2. Write any missing terms with 0 placeholders.


NOW TRY EXERCISE 5

Divide

$$6m^3 - 8m^2 - 5m - 6$$

by $3m - 6$.

EXAMPLE 5 Finding a Quotient with a Fractional Coefficient
Divide $2p^3 + 5p^2 + p - 2$ by $2p + 2$.

$$\begin{array}{r}
 p^2 + \frac{3}{2}p - 1 \\
 2p + 2 \overline{) 2p^3 + 5p^2 + p - 2} \\
 \underline{2p^3 + 2p^2} \\
 3p^2 + p \\
 \underline{3p^2 + 3p} \\
 -2p - 2 \\
 \underline{-2p - 2} \\
 0
 \end{array}$$

$\frac{3p^2}{2p} = \frac{3}{2}p$

The remainder is 0, so the quotient is $p^2 + \frac{3}{2}p - 1$.

NOW TRY 
NOW TRY ANSWERS

4. $x^2 + 4x + 2 + \frac{3x + 1}{2x^2 - 2}$

5. $2m^2 + \frac{4}{3}m + 1$

OBJECTIVE 3 Divide polynomial functions.**Dividing Functions**

If $f(x)$ and $g(x)$ define functions, then

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}. \quad \text{Quotient function}$$

The domain of the quotient function is the intersection of the domains of $f(x)$ and $g(x)$, excluding any values of x for which $g(x) = 0$.

NOW TRY EXERCISE 6

For $f(x) = 8x^2 + 2x - 3$

and $g(x) = 2x - 1$,

find $\left(\frac{f}{g}\right)(x)$ and $\left(\frac{f}{g}\right)(8)$.

EXAMPLE 6 Dividing Polynomial Functions

For $f(x) = 2x^2 + x - 10$ and $g(x) = x - 2$, find $\left(\frac{f}{g}\right)(x)$ and $\left(\frac{f}{g}\right)(-3)$.

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} = \frac{2x^2 + x - 10}{x - 2}$$

This quotient, found in **Example 2**, is $2x + 5$. Thus,

$$\left(\frac{f}{g}\right)(x) = 2x + 5, \quad x \neq 2. \quad \leftarrow \begin{array}{l} 2 \text{ is not in the domain.} \\ \text{It causes denominator} \\ g(x) = x - 2 \text{ to equal } 0. \end{array}$$

$$\left(\frac{f}{g}\right)(-3) = 2(-3) + 5 = -1 \quad \text{Let } x = -3 \text{ in } \left(\frac{f}{g}\right)(x) = 2x + 5.$$

Alternative method: The same result is found by evaluating $\frac{f(-3)}{g(-3)}$.

$$\begin{aligned} \left(\frac{f}{g}\right)(-3) &= \frac{f(-3)}{g(-3)} \\ &= \frac{2(-3)^2 + (-3) - 10}{-3 - 2} \quad \leftarrow \begin{array}{l} f(x) = 2x^2 + x - 10 \\ g(x) = x - 2 \end{array} \\ &= \frac{5}{-5} \\ &= -1 \quad \text{The result is the same.} \end{aligned}$$

NOW TRY ANSWER

6. $4x + 3$, $x \neq \frac{1}{2}$; 35


NOW TRY 

4.6 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Complete each statement.

- We find the quotient of two monomials using the _____ rule for _____.
- When dividing polynomials that are not monomials, first write them in _____ powers.
- If a polynomial in a division problem has a missing term, insert a term with coefficient equal to _____ as a placeholder.

4. To check a division problem, multiply the _____ by the quotient. Then add the _____.

Divide. See Example 1.

$$5. \frac{15x^3 - 10x^2 + 5}{5}$$

$$6. \frac{27m^4 - 18m^3 - 9}{9}$$

$$7. \frac{9y^2 + 12y - 15}{3y}$$

$$8. \frac{80r^2 - 40r + 10}{10r}$$

$$9. \frac{15m^3 + 25m^2 + 30m}{5m^3}$$

$$10. \frac{64x^3 - 72x^2 + 12x}{8x^3}$$

$$11. \frac{4m^2n^2 - 21mn^3 + 18mn^2}{14m^2n^3}$$

$$12. \frac{24h^2k + 56hk^2 - 28hk}{16h^2k^2}$$

$$13. \frac{8wxy^2 + 3wx^2y + 12w^2xy}{4wx^2y}$$

$$14. \frac{12ab^2c + 10a^2bc + 18abc^2}{6a^2bc}$$

Complete the division. See Example 2.

$$15. \begin{array}{r} r^2 \\ 3r - 1 \overline{) 3r^3 - 22r^2 + 25r - 6} \\ \underline{3r^3 - r^2} \\ -21r^2 \end{array}$$

$$16. \begin{array}{r} 3b^2 \\ 2b - 5 \overline{) 6b^3 - 7b^2 - 4b - 40} \\ \underline{6b^3 - 15b^2} \\ 8b^2 \end{array}$$

Divide. See Examples 2–5.

$$17. \frac{y^2 + y - 20}{y + 5}$$

$$18. \frac{y^2 + 3y - 18}{y + 6}$$

$$19. \frac{q^2 + 4q - 32}{q - 4}$$

$$20. \frac{q^2 + 2q - 35}{q - 5}$$

$$21. \frac{3t^2 + 17t + 10}{3t + 2}$$

$$22. \frac{2k^2 - 3k - 20}{2k + 5}$$

$$23. \frac{p^2 + 2p + 20}{p + 6}$$

$$24. \frac{x^2 + 11x + 16}{x + 8}$$

$$25. \frac{3m^3 + 5m^2 - 5m + 1}{3m - 1}$$

$$26. \frac{8z^3 - 6z^2 - 5z + 3}{4z + 3}$$

$$27. \frac{4x^3 + 9x^2 - 10x - 6}{4x + 1}$$

$$28. \frac{10z^3 - 26z^2 + 17z - 13}{5z - 3}$$

$$29. \frac{6x^3 - 19x^2 + 14x - 15}{3x^2 - 2x + 4}$$

$$30. \frac{8m^3 - 18m^2 + 37m - 13}{2m^2 - 3m + 6}$$

$$31. \frac{m^3 - 2m^2 - 9}{m - 3}$$

$$32. \frac{p^3 + 3p^2 - 4}{p + 2}$$

$$33. (x^3 + 2x - 3) \div (x - 1)$$

$$34. (x^3 + 5x^2 - 18) \div (x + 3)$$

$$35. (2x^3 - 11x^2 + 25) \div (x - 5)$$

$$36. (2x^3 + 3x^2 - 5) \div (x - 1)$$

$$37. (3x^3 - x + 4) \div (x - 2)$$

$$38. (3k^3 + 9k - 14) \div (k - 2)$$

$$39. \frac{4k^4 + 6k^3 + 3k - 1}{2k^2 + 1}$$

$$40. \frac{9k^4 + 12k^3 - 4k - 1}{3k^2 - 1}$$

$$41. (2z^3 - 5z^2 + 6z - 15) \div (2z - 5)$$

$$42. (3p^3 + p^2 + 18p + 6) \div (3p + 1)$$

$$43. \frac{6y^4 + 4y^3 + 4y - 6}{3y^2 + 2y - 3}$$

$$44. \frac{8t^4 + 6t^3 + 12t - 32}{4t^2 + 3t - 8}$$

45. $\frac{p^3 - 1}{p - 1}$

47. $(x^4 - 4x^3 + 5x^2 - 3x + 2) \div (x^2 + 3)$

49. $(2p^3 + 7p^2 + 9p + 3) \div (2p + 2)$

51. $(3a^2 - 11a + 17) \div (2a + 6)$

46. $\frac{8a^3 + 1}{2a + 1}$

48. $(3t^4 + 5t^3 - 8t^2 - 13t + 2) \div (t^2 - 5)$

50. $(3x^3 + 4x^2 + 7x + 4) \div (3x + 3)$

52. $(5t^2 + 19t + 7) \div (4t + 12)$

Extending Skills Divide.

53. $\left(2x^2 - \frac{7}{3}x - 1\right) \div (3x + 1)$

55. $\left(3a^2 - \frac{23}{4}a - 5\right) \div (4a + 3)$


54. $\left(m^2 + \frac{7}{2}m + 3\right) \div (2m + 3)$

56. $\left(3q^2 + \frac{19}{5}q - 3\right) \div (5q - 2)$

Solve each problem.


57. Suppose that the volume of a box is

$(2p^3 + 15p^2 + 28p)$ cubic feet.

The height is p feet and the length is $(p + 4)$ feet. Find an expression in p that represents the width.

58. Suppose that a minivan travels

$(2m^3 + 15m^2 + 35m + 36)$ miles

in $(2m + 9)$ hours. Find an expression in m that represents the rate of the van in miles per hour (mph).

- 59.
- Concept Check**
- Let
- $P(x) = 4x^3 - 8x^2 + 13x - 2$
- and
- $D(x) = 2x - 1$
- . Use division to find polynomials
- $Q(x)$
- and
- $R(x)$
- such that

$$P(x) = Q(x) \cdot D(x) + R(x).$$

60. For
- $P(x) = x^3 - 4x^2 + 3x - 5$
- , find
- $P(-1)$
- . Then divide
- $P(x)$
- by
- $D(x) = x + 1$
- . Compare the remainder with
- $P(-1)$
- . What do these results suggest?

For each pair of functions, find $\left(\frac{f}{g}\right)(x)$ and give any x -values that are not in the domain of the quotient function. See Example 6.

61. $f(x) = 10x^2 - 2x$, $g(x) = 2x$

62. $f(x) = 18x^2 - 24x$, $g(x) = 3x$

63. $f(x) = 2x^2 - x - 3$, $g(x) = x + 1$

64. $f(x) = 4x^2 - 23x - 35$, $g(x) = x - 7$

65. $f(x) = 8x^3 - 27$, $g(x) = 2x - 3$

66. $f(x) = 27x^3 + 64$, $g(x) = 3x + 4$

Let $f(x) = x^2 - 9$, $g(x) = 2x$, and $h(x) = x - 3$. Find each of the following. See Example 6.

67. $\left(\frac{f}{g}\right)(x)$

68. $\left(\frac{f}{h}\right)(x)$

69. $\left(\frac{f}{g}\right)(2)$

70. $\left(\frac{f}{h}\right)(1)$

71. $\left(\frac{h}{g}\right)(x)$

72. $\left(\frac{g}{h}\right)(x)$

73. $\left(\frac{h}{g}\right)(3)$

74. $\left(\frac{g}{h}\right)(-1)$

75. $\left(\frac{f}{g}\right)\left(\frac{1}{2}\right)$

76. $\left(\frac{f}{g}\right)\left(\frac{3}{2}\right)$

77. $\left(\frac{h}{g}\right)\left(-\frac{1}{2}\right)$

78. $\left(\frac{h}{g}\right)\left(-\frac{3}{2}\right)$

Chapter 4

Summary

STUDY SKILLS REMINDER

Be prepared for your math test over this chapter. **Review Study Skill 7, Reviewing a Chapter, and Study Skill 8, Taking Math Tests.**

Key Terms**4.1**

exponent (power)
base
exponential expression

4.3

term
algebraic expression

polynomial
numerical coefficient
(coefficient)
degree of a term
polynomial in x
descending powers
leading term
leading coefficient

trinomial
binomial
monomial
degree of a polynomial
like terms
negative of a polynomial

4.4

polynomial function
identity function
squaring function
cubing function
composite function
(composition of functions)

New Symbols

$(f \circ g)(x) = f(g(x))$ composite function

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- A **polynomial** is an algebraic expression made up of
 - a term or a finite product of terms with positive coefficients and exponents
 - the sum of two or more terms with whole number coefficients and exponents
 - the product of two or more terms
 - a term or a finite sum of terms with real number coefficients and whole number exponents.
- A **monomial** is a polynomial with
 - only one term
 - exactly two terms
 - exactly three terms
 - more than three terms.
- A **binomial** is a polynomial with
 - only one term
 - exactly two terms
 - exactly three terms
 - more than three terms.
- A **trinomial** is a polynomial with
 - only one term
 - exactly two terms
 - exactly three terms
 - more than three terms.
- The **FOIL** method is used to
 - add two binomials
 - add two trinomials
 - multiply two binomials
 - multiply two trinomials.

ANSWERS

1. D; Example: $5x^3 + 2x^2 - 7$ 2. A; Examples: $-4, 2x^3, 15a^2b$ 3. B; Example: $3t^3 + 5t$ 4. C; Example: $2a^2 - 3ab + b^2$
5. C; Example: $(m + 4)(m - 3) = m(m) - 3m + 4m + 4(-3) = m^2 + m - 12$


F O I L

Quick Review


CONCEPTS	EXAMPLES
<p>4.1 Integer Exponents</p> <p>Definitions and Rules for Exponents For all integers m and n and all real numbers a and b for which the following are defined, these rules hold true.</p> <p>Product Rule $a^m \cdot a^n = a^{m+n}$</p> <p>Quotient Rule $\frac{a^m}{a^n} = a^{m-n}$</p> <p>Zero Exponent $a^0 = 1$</p> <p>Negative Exponent $a^{-n} = \frac{1}{a^n}$</p> <p>Power Rules (a) $(a^m)^n = a^{mn}$ (b) $(ab)^m = a^m b^m$ (c) $\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}$</p> <p>Special Rules for Negative Exponents</p> $\frac{1}{a^{-n}} = a^n \quad \frac{a^{-n}}{b^{-m}} = \frac{b^m}{a^n}$ $a^{-n} = \left(\frac{1}{a}\right)^n \quad \left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n$	<p>Apply the rules for exponents.</p> $3^4 \cdot 3^2 = 3^6$ $\frac{2^5}{2^3} = 2^2$ $27^0 = 1, \quad (-5)^0 = 1$ $5^{-2} = \frac{1}{5^2}$ $(6^3)^4 = 6^{12}, \quad (5p)^4 = 5^4 p^4$ $\left(\frac{2}{3}\right)^5 = \frac{2^5}{3^5}$ $\frac{1}{x^{-3}} = x^3 \quad \frac{r^{-3}}{t^{-6}} = \frac{t^6}{r^3}$ $4^{-3} = \left(\frac{1}{4}\right)^3 \quad \left(\frac{4}{7}\right)^{-2} = \left(\frac{7}{4}\right)^2$
<p>4.2 Scientific Notation</p> <p>Scientific Notation A number is written in scientific notation when it is expressed in the form</p> $a \times 10^n,$ <p>where $1 \leq a < 10$ and n is an integer.</p>	<p>Write 23,500,000,000 in scientific notation.</p> $23,500,000,000 = 2.35 \times 10^{10}$ <p>Write 4.3×10^{-6} in standard notation.</p> $4.3 \times 10^{-6} = 0.0000043$
<p>4.3 Adding and Subtracting Polynomials</p> <p>Add or subtract polynomials by combining like terms.</p>	<p>Add.</p> $\begin{array}{r} 13x^2 - 3x + 1 \\ -4x^2 + 5x - 7 \\ \hline 9x^2 + 2x - 6 \end{array}$ <p>Subtract.</p> $\begin{aligned} (5x^4 + 3x^2) - (7x^4 + x^2 - x) \\ = 5x^4 + 3x^2 - 7x^4 - x^2 + x \\ = -2x^4 + 2x^2 + x \end{aligned}$
<p>4.4 Polynomial Functions, Graphs, and Composition</p> <p>Adding and Subtracting Functions If $f(x)$ and $g(x)$ define functions, then</p> $(f + g)(x) = f(x) + g(x)$ <p>and</p> $(f - g)(x) = f(x) - g(x).$	<p>Let $f(x) = x^2$ and $g(x) = 2x + 1$. Find $(f + g)(x)$ and $(f - g)(x)$.</p> $\begin{array}{l l} (f + g)(x) & (f - g)(x) \\ = f(x) + g(x) & = f(x) - g(x) \\ = x^2 + 2x + 1 & = x^2 - (2x + 1) \\ & = x^2 - 2x - 1 \end{array}$

CONCEPTS


Graphs of Basic Polynomial Functions



Identity function
 $f(x) = x$
 Domain: $(-\infty, \infty)$
 Range: $(-\infty, \infty)$



Squaring function
 $f(x) = x^2$
 Domain: $(-\infty, \infty)$
 Range: $[0, \infty)$



Cubing function
 $f(x) = x^3$
 Domain: $(-\infty, \infty)$
 Range: $(-\infty, \infty)$

Composition of f and g

$$(f \circ g)(x) = f(g(x))$$

$$(g \circ f)(x) = g(f(x))$$

Let $f(x) = x^2$ and $g(x) = 2x + 1$. Find $(f \circ g)(x)$ and $(g \circ f)(x)$.

$(f \circ g)(x)$ $= f(g(x))$ $= f(2x + 1)$ $= (2x + 1)^2$	$(g \circ f)(x)$ $= g(f(x))$ $= g(x^2)$ $= 2x^2 + 1$
-----------------------------------------------------------	------------------------------------------------------

4.5 Multiplying Polynomials

To multiply two polynomials, multiply each term of one by each term of the other.

To multiply two binomials, use the **FOIL method**. Multiply the **F**irst terms, the **O**uter terms, the **I**nner terms, and the **L**ast terms. Then add these products.

Special Products

$$(x + y)(x - y) = x^2 - y^2$$

$$(x + y)^2 = x^2 + 2xy + y^2$$

$$(x - y)^2 = x^2 - 2xy + y^2$$

Multiplying Functions

If $f(x)$ and $g(x)$ define functions, then

$$(fg)(x) = f(x) \cdot g(x).$$

Multiply.

$$(x^3 + 3x)(4x^2 - 5x + 2)$$

$$= x^3(4x^2 - 5x + 2) + 3x(4x^2 - 5x + 2)$$

$$= 4x^5 - 5x^4 + 2x^3 + 12x^3 - 15x^2 + 6x$$

$$= 4x^5 - 5x^4 + 14x^3 - 15x^2 + 6x$$

$$(2x + 3)(x - 7)$$

$$= 2x(x) + 2x(-7) + 3x + 3(-7) \quad \text{FOIL method}$$

$$= 2x^2 - 14x + 3x - 21 \quad \text{Multiply.}$$

$$= 2x^2 - 11x - 21 \quad \text{Combine like terms.}$$

$$(3m + 8)(3m - 8)$$

$$= 9m^2 - 64$$

$(5a + 3b)^2$ $= 25a^2 + 30ab + 9b^2$	$(2k - 1)^2$ $= 4k^2 - 4k + 1$
---------------------------------------	--------------------------------

Let $f(x) = x^2$ and $g(x) = 2x + 1$. Find $(fg)(x)$.

$$(fg)(x)$$

$$= f(x) \cdot g(x)$$

$$= x^2(2x + 1)$$

$$= 2x^3 + x^2$$

4.6 Dividing Polynomials

Dividing by a Monomial

To divide a polynomial by a monomial, divide each term in the polynomial by the monomial.

$$\frac{a + b}{c} = \frac{a}{c} + \frac{b}{c} \quad (\text{where } c \neq 0)$$

Then write each quotient in lowest terms.

Divide.

$$\frac{2x^3 - 4x^2 + 6x - 8}{2x}$$

$$= \frac{2x^3}{2x} - \frac{4x^2}{2x} + \frac{6x}{2x} - \frac{8}{2x}$$

$$= x^2 - 2x + 3 - \frac{4}{x}$$

CONCEPTS

Dividing by a Polynomial

Use the “long division” process. The process ends when the remainder is 0 or when the degree of the remainder is less than the degree of the divisor.

Dividing Functions

If $f(x)$ and $g(x)$ define functions, then

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}, \quad (\text{where } g(x) \neq 0).$$

EXAMPLES

Divide $m^3 - m^2 + 2m + 5$ by $m + 1$.

$$\begin{array}{r} m^2 - 2m + 4 \\ m + 1 \overline{) m^3 - m^2 + 2m + 5} \\ \underline{m^3 + m^2} \\ -2m^2 + 2m \\ \underline{-2m^2 - 2m} \\ 4m + 5 \\ \underline{4m + 4} \\ 1 \end{array}$$

The answer is

$$m^2 - 2m + 4 + \frac{1}{m + 1}.$$

1 ← Remainder

Let $f(x) = x^2$ and $g(x) = 2x + 1$. Find $\left(\frac{f}{g}\right)(x)$.

$$\begin{aligned} \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \\ &= \frac{x^2}{2x + 1}, \quad x \neq -\frac{1}{2} \end{aligned}$$

Chapter 4

Review Exercises

4.1 Simplify each expression. Assume that all variables represent nonzero real numbers.

- | | | |
|---------------------------------------------------------------------|------------------------------------------------------------------------------|------------------------------------------------------------------------------------|
| 1. 4^3 | 2. $\left(\frac{1}{3}\right)^4$ | 3. $(-5)^3$ |
| 4. $\frac{2}{(-3)^{-2}}$ | 5. $\left(\frac{2}{3}\right)^{-4}$ | 6. $\left(\frac{5}{4}\right)^{-2}$ |
| 7. $5^{-1} - 6^{-1}$ | 8. $2^{-1} + 4^{-1}$ | 9. $-3^0 + 3^0$ |
| 10. $(3^{-4})^2$ | 11. $(x^{-4})^{-2}$ | 12. $(xy^{-3})^{-2}$ |
| 13. $(z^{-3})^3 z^{-6}$ | 14. $(5m^{-3})^2 (m^4)^{-3}$ | 15. $\frac{12k^{-3}(k^{-2})^{-5}}{3k^{-4}}$ |
| 16. $\left(\frac{5z^{-3}}{z^{-1}}\right)\left(\frac{5}{z^2}\right)$ | 17. $\left(\frac{6m^{-4}}{m^{-9}}\right)^{-1}\left(\frac{m^{-2}}{16}\right)$ | 18. $\left(\frac{3r^5}{5r^{-3}}\right)^{-2}\left(\frac{9r^{-1}}{2r^{-5}}\right)^3$ |
| 19. $(-3x^4y^3)(4x^{-2}y^5)$ | 20. $\frac{6m^{-4}n^3}{-3mn^2}$ | 21. $\frac{(5p^{-2}q)(4p^5q^{-3})}{2p^{-5}q^5}$ |

22. Explain the difference between the expressions $(-6)^0$ and -6^0 .

4.2 Write each number in scientific notation.

- | | | |
|------------|------------------|------------|
| 23. 13,450 | 24. 0.0000000765 | 25. -0.138 |
|------------|------------------|------------|

26. The total population of the United States was recently estimated at **321,400,000**. Of this amount, about **72,000** Americans were centenarians, that is, age **100** or older. Write the three boldfaced numbers using scientific notation. (Data from U.S. Census Bureau, Pew Research Center.)



Write each number in standard notation.

27. 1.21×10^6 28. -2.67×10^8 29. 5.8×10^{-3}

Evaluate. Express answers in both scientific notation and standard notation.

30. $\frac{16 \times 10^4}{8 \times 10^8}$ 31. $\frac{6 \times 10^{-2}}{4 \times 10^{-5}}$
 32. $\frac{0.0000000164}{0.0004}$ 33. $\frac{0.0009 \times 12,000,000}{400,000}$

34. The planet Mercury has an average distance from the sun of 3.6×10^7 mi, and the average distance from Venus to the sun is 6.7×10^7 mi.
- (a) How long would it take a spacecraft traveling at 1.55×10^3 mph to travel from Venus to Mercury? Give the answer in hours, in standard notation.
- (b) Use the answer from part (a) to find the number of days it would take the spacecraft to travel from Venus to Mercury, to the nearest whole number.

4.3 Give the numerical coefficient and the degree of each term.

35. $14p^5$ 36. $-z^2$ 37. $\frac{x}{10}$ 38. $504p^3r^5$

For each polynomial, (a) write in descending powers, (b) identify as a monomial, binomial, trinomial, or none of these, and (c) give the degree.


39. $9k + 11k^3 - 3k^2$ 40. $14m^6 + 9m^7$
 41. $-5y^4 + 3y^3 + 7y^2 - 2y$ 42. $-7q^5r^3$

Add or subtract as indicated.

43. Add.

$$\begin{array}{r} 3x^2 - 5x + 6 \\ -4x^2 + 2x - 5 \\ \hline \end{array}$$
44. Subtract.

$$\begin{array}{r} -5y^3 \quad + 8y - 3 \\ \quad \quad \quad 4y^2 + 2y + 9 \\ \hline \end{array}$$
45. $(4a^3 - 9a + 15) - (-2a^3 + 4a^2 + 7a)$ 46. $(3y^2 + 2y - 1) + (5y^2 - 11y + 6)$
47. Find the perimeter of the triangle.



48. Give an example of a polynomial in the variable x that is of degree 5, is written in descending powers, and lacks a third-degree term.

4.4 Work each problem.

49. For the polynomial function $f(x) = -2x^2 + 5x + 7$, find each value.

- (a) $f(-2)$ (b) $f(3)$ (c) $f(0)$

50. The number of twin births in the United States during the years 2005 through 2014 can be modeled by the polynomial function

$$P(x) = 140.33x^3 - 4034.71x^2 + 36,125x + 35,677,$$

where $x = 5$ represents 2005, $x = 6$ represents 2006, and so on. Use this function to approximate the number of twin births (to the nearest whole number as needed) in each given year. (Data from National Center for Health Statistics.)

- (a) 2005 (b) 2010 (c) 2014



51. Let $f(x) = 2x + 3$ and $g(x) = 5x^2 - 3x + 2$. Find each of the following.

- (a) $(f + g)(x)$ (b) $(f - g)(x)$ (c) $(f + g)(-1)$ (d) $(f - g)(-1)$

Graph each polynomial function. Give the domain and range.

52. $f(x) = -2x + 5$ 53. $f(x) = x^2 - 6$ 54. $f(x) = -x^3 + 1$

55. Let $f(x) = 3x^2 + 2x - 1$ and $g(x) = 5x + 7$. Find each of the following.

- (a) $(g \circ f)(3)$ (b) $(f \circ g)(3)$ (c) $(f \circ g)(-2)$
 (d) $(g \circ f)(-2)$ (e) $(f \circ g)(x)$ (f) $(g \circ f)(x)$

4.5 Find each product.

56. $-6k(2k^2 + 7)$ 57. $2x(x + 4)(x + 7)$ 58. $(3m - 2)(5m + 1)$
 59. $(3w - 2t)(2w - 3t)$ 60. $(2p^2 + 6p)(5p^2 - 4)$ 61. $(3q^2 + 2q - 4)(q - 5)$
 62. $(6r^2 - 1)(6r^2 + 1)$ 63. $(4m + 3)^2$ 64. $(3t - 2s)^2$

4.6 Divide.

65. $\frac{4y^3 - 12y^2 + 5y}{4y}$

66. $\frac{x^3 - 9x^2 + 26x - 30}{x - 5}$

67. $\frac{2p^3 + 9p^2 + 27}{2p - 3}$

68. $\frac{5p^4 + 15p^3 - 33p^2 - 9p + 18}{5p^2 - 3}$

4.5, 4.6 Let $f(x) = 12x^2 - 3x$ and $g(x) = 3x$. Find each of the following.

69. (a) $(fg)(x)$ (b) $(fg)(-1)$ 70. (a) $\left(\frac{f}{g}\right)(x)$ (b) $\left(\frac{f}{g}\right)(2)$

Chapter 4

Mixed Review Exercises

Solve each problem.

1. Match each expression (a)–(j) in Column I with its equivalent expression A–J in Column II. Choices may be used once, more than once, or not at all.

I		II	
(a) 4^{-2}	(b) -4^2	A. $\frac{1}{16}$	B. 0
(c) 4^0	(d) $(-4)^0$	C. 1	D. $-\frac{1}{16}$
(e) $(-4)^{-2}$	(f) -4^0	E. -1	F. $\frac{5}{16}$
(g) $-4^0 + 4^0$	(h) $-4^0 - 4^0$	G. -16	H. -2
(i) $4^{-2} + 4^{-1}$	(j) 4^2	I. 16	J. none of these

2. In 2017, the estimated population of New Zealand was 4.510×10^6 . The population density was 44.16 people per square mile. What is the area of New Zealand to the nearest square mile? (Data from *The World Almanac and Book of Facts*.)

Perform the indicated operations and then simplify. Assume that all variables represent non-zero real numbers.

- | | | |
|--------------------------------------------------|------------------------------------------------------|------------------------|
| 3. $\frac{6^{-1}y^3(y^2)^{-2}}{6y^{-4}(y^{-1})}$ | 4. 5^{-3} | 5. $-(-3)^2$ |
| 6. $\frac{(-z^{-2})^3}{5(z^{-3})^{-1}}$ | 7. $7p^5(3p^4 + p^3 + 2p^2)$ | 8. $(2x - 9)^2$ |
| 9. $(y^6)^{-5}(2y^{-3})^{-4}$ | 10. $\left(\frac{3w^{-2}z^4}{-6wz^{-5}}\right)^{-2}$ | 11. $(4x + 1)(2x - 3)$ |
| 12. $\frac{8x^2 - 23x + 2}{x - 3}$ | 13. $[(3m - 5n) + p][(3m - 5n) - p]$ | |
| 14. $\frac{20y^3x^3 + 15y^4x + 25yx^4}{10yx^2}$ | 15. $(2k - 1) - (3k^2 - 2k + 6)$ | |

Chapter 4

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

► View the complete solutions to all Chapter Test exercises in MyLab Math.

1. Match each expression (a)–(j) in Column I with its equivalent expression A–J in Column II. Choices may be used once, more than once, or not at all.

I		II	
(a) 7^{-2}	(b) 7^0	A. 1	B. $\frac{1}{9}$
(c) -7^0	(d) $(-7)^0$	C. $\frac{1}{49}$	D. -1
(e) -7^2	(f) $7^{-1} + 2^{-1}$	E. -49	F. $\frac{9}{14}$
(g) $(7 + 2)^{-1}$	(h) $\frac{7^{-1}}{2^{-1}}$	G. $\frac{2}{7}$	H. 0
(i) 7^2	(j) $(-7)^{-2}$	I. 49	J. none of these

Simplify each expression. Assume that all variables represent nonzero real numbers.

$$2. (3x^{-2}y^3)^{-2}(4x^3y^{-4}) \qquad 3. \frac{36r^{-4}(r^2)^{-3}}{6r^4}$$

$$4. \left(\frac{4p^2}{q^4}\right)^3\left(\frac{6p^8}{q^{-8}}\right)^{-2} \qquad 5. (-2x^4y^{-3})^0(-4x^{-3}y^{-8})^2$$

6. Write 9.1×10^{-7} in standard notation.

7. Evaluate. Express the answer in both scientific notation and standard notation.

$$\frac{2,500,000 \times 0.00003}{0.05 \times 5,000,000}$$

8. Let $f(x) = -2x^2 + 5x - 6$ and $g(x) = 7x - 3$. Find each of the following.

(a) $f(4)$ (b) $(f + g)(x)$ (c) $(f - g)(x)$ (d) $(f - g)(-2)$

Graph each polynomial function. Give the domain and range.

9. $f(x) = -2x^2 + 3$ 10. $f(x) = -x^3 + 3$

11. Let $f(x) = 3x + 5$ and $g(x) = x^2 + 2$. Find each of the following.

(a) $(f \circ g)(-2)$ (b) $(f \circ g)(x)$ (c) $(g \circ f)(x)$

Perform the indicated operations.

12. $(5x - 3)(2x + 1)$ 13. $(2m - 5)(3m^2 + 4m - 5)$

14. $(6x + y)(6x - y)$ 15. $(3k + q)^2$

16. $[2y + (3z - x)][2y - (3z - x)]$ 17. $\frac{16p^3 - 32p^2 + 24p}{4p^2}$

18. $(x^3 + 3x^2 - 4) \div (x - 1)$

19. $(4x^3 - 3x^2 + 2x - 5) - (3x^3 + 11x + 8) + (x^2 - x)$

20. Let $f(x) = x^2 + 3x + 2$ and $g(x) = x + 1$. Find each of the following.

(a) $(fg)(x)$ (b) $(fg)(-2)$ (c) $\left(\frac{f}{g}\right)(x)$ (d) $\left(\frac{f}{g}\right)(-2)$

Chapters R–4

Cumulative Review Exercises

1. Match each number in Column I with the equivalent number(s) in Column II. Choices may be used more than once.

I		II		
(a) $\frac{1}{100}$	(b) $\frac{1}{10}$	A. 0.5	B. 0.25	C. 0.1
(c) 1	(d) $\frac{1}{2}$	D. 0.01	E. 125%	F. 50%
(e) $\frac{1}{4}$	(f) $\frac{5}{4}$	G. 100%	H. 10%	I. 1%

2. Match each number in Column I with the choice or choices of sets of numbers in Column II to which the number belongs.

- | I | | II | |
|-----------------|--------------------|-----------------------|---------------------|
| (a) 34 | (b) 0 | A. Natural numbers | B. Whole numbers |
| (c) 2.16 | (d) $-\sqrt{36}$ | C. Integers | D. Rational numbers |
| (e) $\sqrt{13}$ | (f) $-\frac{4}{5}$ | E. Irrational numbers | F. Real numbers |

Evaluate.

3. $9 \cdot 4 - 16 \div 4$ 4. $\left(\frac{1}{3}\right)^2 - \left(\frac{1}{2}\right)^3$ 5. $-|8 - 13| - |-4| + |-9|$
6. Simplify $0.5(8x - 10) - 0.4(5 + 10x)$.


Solve.

7. $-5(8 - 2z) + 4(7 - z) = 7(8 + z) - 3$ 8. $3(x + 2) - 5(x + 2) = -2x - 4$
9. $A = p + prt$ for t 10. $2(m + 5) - 3m + 1 > 5$
11. $|3x - 1| = 2$ 12. $|3z + 1| \geq 7$
13. A survey polled Internet users age 12 and older about the Internet activities they engaged in daily. Complete the results shown in the table if 5000 such users were surveyed.

Daily Internet Activity	Percent	Actual Number
Browse the web	70%	
Download/listen to music	39%	
Download/watch videos		1800
Play games		1550

Data from the 2017 Center for the Digital Future report.

14. Find the measure of each angle of the triangle.



Find the slope of each line described.

15. Through $(-4, 5)$ and $(2, -3)$ 16. Through $(4, 5)$; horizontal

Write an equation of each line that satisfies the given conditions. Give the equation (a) in slope-intercept form and (b) in standard form.

17. Through $(4, -1)$; $m = -4$ 18. Through $(0, 0)$ and $(1, 4)$


Graph each equation or inequality.

19. $-3x + 4y = 12$ 20. $y \leq 2x - 6$ 21. $3x + 2y < 0$

22. Per capita consumption of whole milk in the United States (in gallons) is shown in the graph, where $x = 0$ represents 1970.

- (a) Use the given ordered pairs to find the average rate of change in per capita consumption of whole milk (in gallons) per year during this period. Interpret the answer.
- (b) Use the answer from part (a) to write an equation of the line in slope-intercept form that models per capita consumption of whole milk y (in gallons).
- (c) Use the equation from part (b) to approximate per capita consumption of whole milk in 2000.

Whole Milk Consumption



Data from U.S. Department of Agriculture.

23. Give the domain and range of the relation

$$\{(-4, -2), (-1, 0), (2, 0), (5, 2)\}.$$

Does this relation define a function?

24. If $g(x) = -x^2 - 2x + 6$, find $g(3)$.

Solve each system.

$$\begin{aligned} 25. \quad 3x - 4y &= 1 \\ 2x + 3y &= 12 \end{aligned}$$

$$\begin{aligned} 26. \quad 3x - 2y &= 4 \\ -6x + 4y &= 7 \end{aligned}$$

$$\begin{aligned} 27. \quad x + 3y - 6z &= 7 \\ 2x - y + z &= 1 \\ x + 2y + 2z &= -1 \end{aligned}$$

Solve each problem.

28. The Star-Spangled Banner that flew over Fort McHenry during the War of 1812 had a perimeter of 144 ft. Its length measured 12 ft more than its width. Find the dimensions of this flag, which is displayed in the Smithsonian Institution's Museum of American History in Washington, DC. (Data from National Park Service brochure.)



29. Agbe needs 9 L of a 20% solution of alcohol. Agbe has a 15% solution on hand, as well as a 30% solution. How many liters of the 15% solution and how many liters of the 30% solution should Agbe mix to obtain the 20% solution needed?

30. Simplify $\frac{x^{-6}y^3z^{-1}}{x^7y^{-4}z}$ ($x, y, z \neq 0$). Write the answer with only positive exponents.

Perform the indicated operations.

$$31. (3x^2 - 8x + 1) - (x^2 - 3x - 9)$$

$$32. (x + 2y)(x^2 - 2xy + 4y^2)$$

$$33. (3x + 2y)(5x - y)$$

$$34. \frac{16x^3y^5 - 8x^2y^2 + 4}{4x^2y}$$

$$35. \frac{m^3 - 3m^2 + 5m - 3}{m - 1}$$

STUDY SKILLS REMINDER

It is not too soon to begin preparing for your final exam. **Review Study Skill 10, Preparing for Your Math Final Exam.**



FACTORING

Determining the dimensions for a garden of fixed area is accomplished by solving a *quadratic equation*, one of the topics covered in this chapter on *factoring*.

- 5.1** Greatest Common Factors and Factoring by Grouping
- 5.2** Factoring Trinomials
- 5.3** Special Factoring
- 5.4** A General Approach to Factoring
- 5.5** Solving Quadratic Equations Using the Zero-Factor Property

5.1 Greatest Common Factors and Factoring by Grouping

OBJECTIVES

- Factor out the greatest common factor.
- Factor by grouping.

VOCABULARY

- factoring
- greatest common factor (GCF)

Writing a polynomial as the product of two or more simpler polynomials is called **factoring** the polynomial. For example, consider the following.

$$3x(5x - 2) = 15x^2 - 6x \quad \text{Multiplying}$$

$$15x^2 - 6x = 3x(5x - 2) \quad \text{Factoring}$$

Notice that both multiplying and factoring use the distributive property, but in opposite directions. **Factoring “undoes,” or reverses, multiplying.**

OBJECTIVE 1 Factor out the greatest common factor.

The first step in factoring a polynomial is to find the *greatest common factor* for the terms of the polynomial. The product of the greatest common numerical factor and each variable factor of least degree common to every term in a polynomial is the **greatest common factor (GCF)** of the terms of the polynomial.

For example, the greatest common factor for $8x + 12$ is 4 because 4 is the greatest factor that *divides into* both $8x$ and 12.

$$\begin{aligned} 8x + 12 &= 4(2x) + 4(3) && \text{Factor 4 from each term.} \\ &= 4(2x + 3) && \text{Distributive property, } ab + ac = a(b + c) \end{aligned}$$

CHECK Multiply $4(2x + 3)$ to obtain $8x + 12$. ✓ Original polynomial

Using the distributive property in this way is called **factoring out the greatest common factor**.

NOW TRY EXERCISE 1

Factor out the greatest common factor.

- $54m - 45$
- $2k - 7$
- $18 + 54r$

EXAMPLE 1 Factoring Out the Greatest Common Factor

Factor out the greatest common factor.

$$\begin{aligned} \text{(a)} \quad 9z - 18 &= 9 \cdot z - 9 \cdot 2 && \text{GCF} = 9; \text{ Factor 9 from each term.} \\ &= 9(z - 2) && \text{Distributive property} \end{aligned}$$

CHECK Multiply $9(z - 2)$ to obtain $9z - 18$. ✓ Original polynomial

$$\begin{aligned} \text{(b)} \quad 56m + 35p &= 7(8m + 5p) && \text{GCF} = 7 \\ \text{(c)} \quad 2y + 5 &&& \text{There is no common factor other than 1.} \end{aligned}$$

$$\begin{aligned} \text{(d)} \quad 12 + 24z &= 12 \cdot 1 + 12 \cdot 2z && \text{Identity property; 12 is the GCF.} \\ &= 12(1 + 2z) && \text{Distributive property} \end{aligned}$$

Remember to write the 1.

$$\begin{aligned} \text{CHECK} \quad 12(1 + 2z) &= 12(1) + 12(2z) && \text{Distributive property} \\ &= 12 + 24z && \text{Original polynomial} \end{aligned}$$

NOW TRY ANSWERS

- $9(6m - 5)$
 - There is no common factor other than 1.
 - $18(1 + 3r)$


NOW TRY EXERCISE 2

Factor out the greatest common factor.

- (a) $16y^4 + 8y^3$
 (b) $14p^2 - 9p^3 + 6p^4$
 (c) $9x^3y^2 - 3x^2y^2 + 6x^2y^3$

EXAMPLE 2 Factoring Out the Greatest Common Factor

Factor out the greatest common factor.

(a) $9x^2 + 12x^3$

The numerical part of the GCF is 3, the greatest integer that divides into both 9 and 12. The least exponent that appears on x is 2. The GCF is $3x^2$.

$$\begin{aligned} 9x^2 + 12x^3 &= 3x^2(3) + 3x^2(4x) && \text{GCF} = 3x^2 \\ &= 3x^2(3 + 4x) && \text{Distributive property} \end{aligned}$$

(b) $32p^4 - 24p^3 + 40p^5$

$$\begin{aligned} &= 8p^3(4p) + 8p^3(-3) + 8p^3(5p^2) && \text{GCF} = 8p^3 \\ &= 8p^3(4p - 3 + 5p^2) && \text{Distributive property} \end{aligned}$$

(c) $3k^4 - 15k^7 + 24k^9$

Remember the 1.

$$= 3k^4(1 - 5k^3 + 8k^5) \quad \text{GCF} = 3k^4$$

(d) $24m^3n^2 - 18m^2n + 6m^4n^3$

$$\begin{aligned} &= 6m^2n(4mn) + 6m^2n(-3) + 6m^2n(m^2n^2) && \text{GCF} = 6m^2n \\ &= 6m^2n(4mn - 3 + m^2n^2) && \text{Distributive property} \end{aligned}$$

(e) $25x^2y^3 + 30y^5 - 15x^4y^7$

$$= 5y^3(5x^2 + 6y^2 - 3x^4y^4)$$

In each case, check the factored form by multiplying.

NOW TRY 


NOW TRY EXERCISE 3

Factor out the greatest common factor.

- (a) $(3x - 2)(x + 1) + (3x - 2)(2x - 5)$
 (b) $z(a - b)^3 - 2w(a - b)^2$

EXAMPLE 3 Factoring Out a Binomial Factor

Factor out the greatest common factor.

(a) $(x - 5)(x + 6) + (x - 5)(2x + 5)$ The greatest common factor is $x - 5$.

$$= (x - 5)[(x + 6) + (2x + 5)]$$
 Factor out $x - 5$.

$$= (x - 5)(3x + 11)$$
 Combine like terms inside the brackets.

(b) $z^2(m + n) + x^2(m + n)$

$$= (m + n)(z^2 + x^2)$$
 Factor out the common factor $(m + n)$.

(c) $p(r + 2s)^2 - q(r + 2s)^3$ $(r + 2s)^3 = (r + 2s)^2(r + 2s)$

$$= (r + 2s)^2[p - q(r + 2s)]$$
 Factor out the common factor $(r + 2s)^2$.

$$= (r + 2s)^2(p - qr - 2qs)$$
 Be careful with signs.

(d) $(p - 5)(p + 2) - (p - 5)(3p + 4)$

$$= (p - 5)[(p + 2) - (3p + 4)]$$
 Factor out the common factor $p - 5$.

$$= (p - 5)[p + 2 - 3p - 4]$$
 Distributive property

$$= (p - 5)[-2p - 2]$$
 Combine like terms.

$$= (p - 5)[-2(p + 1)]$$
 Look for a common factor.

$$= -2(p - 5)(p + 1)$$
 Commutative property

NOW TRY 

NOW TRY ANSWERS

2. (a) $8y^3(2y + 1)$
 (b) $p^2(14 - 9p + 6p^2)$
 (c) $3x^2y^2(3x - 1 + 2y)$
 3. (a) $(3x - 2)(3x - 4)$
 (b) $(a - b)^2(za - zb - 2w)$

Don't stop here.


**NOW TRY
EXERCISE 4**

Factor $-4y^5 - 3y^3 + 8y$
in two ways.

EXAMPLE 4 Factoring Out a Negative Common Factor

Factor $-a^3 + 3a^2 - 5a$ in two ways.

First, a could be used as the common factor.

$$\begin{aligned} & -a^3 + 3a^2 - 5a \\ &= a(-a^2) + a(3a) + a(-5) && \text{Factor out } a. \\ &= a(-a^2 + 3a - 5) && \text{Distributive property} \end{aligned}$$

CHECK Multiply $a(-a^2 + 3a - 5)$ to obtain $-a^3 + 3a^2 - 5a$. ✓

Because of the leading negative sign, $-a$ could also be used as the common factor.

$$\begin{aligned} & -a^3 + 3a^2 - 5a \\ &= -a(a^2) + (-a)(-3a) + (-a)(5) && \text{Factor out } -a. \\ &= -a(a^2 - 3a + 5) && \text{Distributive property} \end{aligned}$$

CHECK Multiply $-a(a^2 - 3a + 5)$ to obtain $-a^3 + 3a^2 - 5a$. ✓

Sometimes there may be a reason to prefer one of these forms, but either is correct.

NOW TRY 

NOTE In cases where the leading coefficient is negative (as in **Example 4**), the answer section will usually give the factored form that has the common factor with a negative coefficient.

OBJECTIVE 2 Factor by grouping.

Sometimes the *individual terms* of a polynomial have greatest common factor of 1, but it still may be possible to factor the polynomial using a process called **factoring by grouping**.

We usually factor by grouping when a polynomial has more than three terms.

EXAMPLE 5 Factoring by Grouping

Factor $ax - ay + bx - by$.

Group the terms in pairs so that each pair has a common factor.

$$\begin{aligned} & ax - ay + bx - by \\ & \begin{array}{cc} \text{Terms with common factor } a & \text{Terms with common factor } b \\ \downarrow \quad \downarrow & \downarrow \quad \downarrow \end{array} \\ &= (ax - ay) + (bx - by) \\ &= a(x - y) + b(x - y) && \text{Factor each group.} \\ &= (x - y)(a + b) && \text{The common factor is } x - y. \end{aligned}$$

By the commutative property,
 $(a + b)(x - y)$
is also correct.

CHECK $(x - y)(a + b)$

$$\begin{aligned} &= xa + xb - ya - yb && \text{Multiply using the FOIL method.} \\ &= ax + bx - ay - by && \text{Commutative property} \\ &= ax - ay + bx - by && \checkmark \text{ Original polynomial} \end{aligned}$$

NOW TRY 


**NOW TRY
EXERCISE 5**

Factor.

$$3m - 3n + xm - xn$$

NOW TRY ANSWERS

4. $y(-4y^4 - 3y^2 + 8)$;
 $-y(4y^4 + 3y^2 - 8)$
5. $(m - n)(3 + x)$


NOW TRY EXERCISE 6

Factor.

$$ab - 7a - 5b + 35$$

EXAMPLE 6 Factoring by Grouping
Factor $3x - 3y - ax + ay$.

$$\begin{aligned} 3x - 3y - ax + ay & \quad \text{Pay close attention here.} \\ &= (3x - 3y) + (-ax + ay) \quad (*) \text{ Group the terms.} \\ &= 3(x - y) + a(-x + y) \quad \text{Factor out 3, and factor out } a. \end{aligned}$$

The factors $(x - y)$ and $(-x + y)$ are opposites. If we factor out $-a$ instead of a in the second group, we obtain the common binomial factor $(x - y)$. So we start over.

$$\begin{aligned} (3x - 3y) + (-ax + ay) & \quad \text{Equation (*) from above} \\ &= 3(x - y) - a(x - y) \quad \text{Be careful with signs.} \\ &= (x - y)(3 - a) \quad \text{Factor out } x - y. \end{aligned}$$

CHECK $(x - y)(3 - a)$

$$\begin{aligned} &= 3x - ax - 3y + ay \quad \text{Multiply using the FOIL method.} \\ &= 3x - 3y - ax + ay \quad \checkmark \text{ Original polynomial} \end{aligned}$$

NOW TRY 

NOTE In **Example 6**, a different grouping would lead to the factored form $(a - 3)(y - x)$. Verify by multiplying that this is also correct.

Factoring by Grouping

Step 1 Group terms. Collect the terms into groups so that each group has a common factor.

Step 2 Factor within the groups. Factor out the common factor in each group.

Step 3 Factor the entire polynomial. If each group now has a common factor, factor it out. If not, try a different grouping.

Always check the factored form by multiplying.


NOW TRY EXERCISE 7

Factor.

$$3ax - 6xy - a + 2y$$

EXAMPLE 7 Factoring by Grouping
Factor $6ax + 12bx + a + 2b$.

$$\begin{aligned} 6ax + 12bx + a + 2b \\ &= (6ax + 12bx) + (a + 2b) \quad \text{Group the terms.} \end{aligned}$$

Now factor $6x$ from the first group, and use the identity property of multiplication to introduce the factor 1 in the second group.

$$\begin{aligned} &= 6x(a + 2b) + 1(a + 2b) \quad \text{Remember to write the 1.} \\ &= (a + 2b)(6x + 1) \quad \text{Factor each group.} \\ & \quad \text{Factor out } a + 2b. \end{aligned}$$

CHECK $(a + 2b)(6x + 1)$

$$\begin{aligned} &= 6ax + a + 12bx + 2b \quad \text{Multiply using the FOIL method.} \\ &= 6ax + 12bx + a + 2b \quad \checkmark \text{ Original polynomial} \end{aligned}$$

NOW TRY 
NOW TRY ANSWERS

6. $(b - 7)(a - 5)$
7. $(a - 2y)(3x - 1)$


NOW TRY EXERCISE 8


Factor.

$$2kp^2 + 6 - 3p^2 - 4k$$

EXAMPLE 8 Rearranging Terms before Factoring by Grouping
Factor $p^2q^2 - 10 - 2q^2 + 5p^2$.

Neither the first two terms nor the last two terms have a common factor except 1. We can rearrange and group the terms as follows.

$$\begin{aligned} p^2q^2 - 10 - 2q^2 + 5p^2 &= (p^2q^2 - 2q^2) + (5p^2 - 10) && \text{Rearrange and group the terms.} \\ &= q^2(p^2 - 2) + 5(p^2 - 2) && \text{Factor out the common factors.} \\ &= (p^2 - 2)(q^2 + 5) && \text{Factor out } p^2 - 2. \text{ Use parentheses.} \end{aligned}$$



CHECK $(p^2 - 2)(q^2 + 5)$

$$\begin{aligned} &= p^2q^2 + 5p^2 - 2q^2 - 10 && \text{Multiply using the FOIL method.} \\ &= p^2q^2 - 10 - 2q^2 + 5p^2 \quad \checkmark && \text{Original polynomial} \end{aligned}$$

NOW TRY 

CAUTION In **Example 8**, do not stop at the step

$$q^2(p^2 - 2) + 5(p^2 - 2).$$

This expression is *not in factored form* because it is a *sum* of two terms, $q^2(p^2 - 2)$ and $5(p^2 - 2)$, not a *product*.


NOW TRY EXERCISE 9

Factor.

$$12wy - 24xy + 4wz - 8xz$$

EXAMPLE 9 Factoring Out a GCF before Factoring by Grouping
Factor $10ax - 5ay + 10bx - 5by$.

Always start by factoring out any greatest common factor from the terms.

$$\begin{aligned} 10ax - 5ay + 10bx - 5by &= 5(2ax - ay + 2bx - by) && \text{Factor out the GCF, 5.} \\ &= 5[(2ax - ay) + (2bx - by)] && \text{Group the terms inside the brackets.} \\ &= 5[a(2x - y) + b(2x - y)] && \text{Factor out the common factors.} \\ &= 5[(2x - y)(a + b)] && \text{Factor out } 2x - y. \\ &= 5(2x - y)(a + b) && \text{Write without the brackets.} \end{aligned}$$

CHECK $5(2x - y)(a + b)$

$$\begin{aligned} &= 5(2ax + 2bx - ay - by) && \text{Use the FOIL method.} \\ &= 10ax + 10bx - 5ay - 5by && \text{Distributive property} \\ &= 10ax - 5ay + 10bx - 5by \quad \checkmark && \text{Original polynomial} \end{aligned}$$

NOW TRY 

NOTE Verify that the result in **Example 9** can also be obtained by grouping the terms of the polynomial

$$2ax - ay + 2bx - by \quad \text{as} \quad (2ax + 2bx) + (-ay - by).$$

NOW TRY ANSWERS

8. $(p^2 - 2)(2k - 3)$
 9. $4(w - 2x)(3y + z)$

5.1 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Choose the letter of the correct response.

- Which choice is an example of a polynomial in factored form?
 - $3x^2y^3 + 6x^2(2x + y)$
 - $5(x + y)^2 - 10(x + y)^3$
 - $(-2 + 3x)(5y^2 + 4y + 3)$
 - $(3x + 4)(5x - y) - (3x + 4)(2x - 1)$
- Which choice is the correct completely factored form of $8 + 24r$?
 - $8(3r)$
 - $-8(-3r)$
 - $8(1 + 3r)$
 - $4(2 + 6r)$

3. Concept Check A student factored the following polynomial as shown.

$$\begin{aligned} 2tw^2 - 16t + 5w^2 - 40 \\ &= (2tw^2 - 16t) + (5w^2 - 40) \\ &= 2t(w^2 - 8) + 5(w^2 - 8) \end{aligned}$$

The teacher did not give the student full credit. **WHAT WENT WRONG?** Give the correct answer.

4. Concept Check A student factored the following polynomial as shown.

$$\begin{aligned} 4x^2y^5 - 8xy^3 \\ &= 2xy^3(2xy^2 - 4) \end{aligned}$$

The teacher did not give the student full credit. **WHAT WENT WRONG?** Give the correct answer.

Concept Check Find the greatest common factor for each list of terms.

- $9m^3, 3m^2, 15m$
- $4a^2, 6a, 2a^3$
- $16xy^3, 24x^2y^2, 8x^2y$
- $10m^2n^2, 25mn^3, 50m^2n$
- $6m(r + t)^2, 3p(r + t)^4$
- $7z^2(m + n)^4, 9z^3(m + n)^5$

Factor out the greatest common factor. See Examples 1–4.

- $12m - 60$
- $15r - 45$
- $8s + 16t$
- $35p + 70q$
- $4 + 20z$
- $9 + 27x$
- $8y - 15$
- $7x - 40$
- $8k^3 + 24k$
- $9z^4 + 81z$
- $-4p^3q^4 - 2p^2q^5$
- $-3z^5w^2 - 18z^3w^4$
- $7x^3 + 35x^4 - 14x^5$
- $6k^3 - 36k^4 - 48k^5$
- $10t^5 - 2t^3 - 4t^4$
- $6p^3 - 3p^2 - 9p^4$
- $15a^2c^3 - 25ac^2 + 5ac$
- $15y^3z^3 - 27y^2z^4 + 3yz^3$
- $16z^2n^6 + 64zn^7 - 32z^3n^3$
- $5r^3s^5 + 10r^2s^2 - 15r^4s^2$
- $14a^3b^2 + 7a^2b - 21a^5b^3 + 42ab^4$
- $12km^3 - 24k^3m^2 + 36k^2m^4 - 60k^4m^3$
- $(m - 4)(m + 2) + (m - 4)(m + 3)$
- $(z - 5)(z + 7) + (z - 5)(z + 10)$
- $(2z - 1)(z + 6) - (2z - 1)(z - 5)$
- $(3x + 2)(x - 4) - (3x + 2)(x + 8)$
- $5(2 - x)^2 - 2(2 - x)^3$
- $2(5 - x)^3 - 3(5 - x)^2$
- $4(3 - x)^2 - (3 - x)^3 + 3(3 - x)$
- $2(t - s) + 4(t - s)^2 - (t - s)^3$
- $15(2z + 1)^3 + 10(2z + 1)^2 - 25(2z + 1)$
- $6(a + 2b)^2 - 4(a + 2b)^3 + 12(a + 2b)^4$
- $5(m + p)^3 - 10(m + p)^2 - 15(m + p)^4$
- $-9a^2(p + q) - 3a^3(p + q)^2 + 6a(p + q)^3$

Factor each polynomial in two ways. First use a common factor with a positive coefficient, and then use a common factor with a negative coefficient. See Example 4.

45. $-r^3 + 3r^2 + 5r$

46. $-t^4 + 8t^3 - 12t$

47. $-12s^5 + 48s^4$

48. $-16y^4 + 64y^3$

49. $-2x^5 + 6x^3 + 4x^2$

50. $-5a^3 + 10a^4 - 15a^5$

Factor by grouping. See Examples 5–9.

51. $mx + qx + my + qy$

52. $2k + 2h + jk + jh$

53. $10m + 2n + 5mk + nk$

54. $3ma + 3mb + 2ab + 2b^2$

55. $4 - 2q - 6p + 3pq$

56. $20 + 5m + 12n + 3mn$

57. $p^2 - 4zq + pq - 4pz$

58. $r^2 - 9tw + 3rw - 3rt$

59. $2xy + 3y + 2x + 3$

60. $7ab + 35bc + a + 5c$

61. $m^3 + 4m^2 - 6m - 24$

62. $2a^3 + a^2 - 14a - 7$

63. $-3a^3 - 3ab^2 + 2a^2b + 2b^3$

64. $-16m^3 + 4m^2p^2 - 4mp + p^3$

65. $4 + xy - 2y - 2x$

66. $10ab - 21 - 6b + 35a$

67. $8 + 9y^4 - 6y^3 - 12y$

68. $x^3y^2 - 3 - 3y^2 + x^3$

69. $1 - a + ab - b$

70. $2ab^2 - 8b^2 + a - 4$

71. $2mx - 6qx + 2my - 6qy$

72. $12 - 6q - 18p + 9pq$

73. $4a^3 - 4a^2b^2 + 8ab - 8b^3$

74. $5x^3 + 15x^2y^2 - 5xy - 15y^3$

75. $2x^3y^2 + x^2y^2 - 14xy^2 - 7y^2$

76. $3m^2n^3 + 15m^2n - 2m^2n^2 - 10m^2$

Extending Skills Factor out the variable that is raised to the lesser exponent. (For example, in Exercise 77, factor out m^{-5} .)

77. $3m^{-5} + m^{-3}$

78. $k^{-2} + 2k^{-4}$

79. $3p^{-3} + 2p^{-2}$

80. $-5q^{-3} + 8q^{-2}$

5.2 Factoring Trinomials

OBJECTIVES

- Factor trinomials when the coefficient of the second-degree term is 1.
- Factor trinomials by grouping when the coefficient of the second-degree term is not 1.
- Factor trinomials using the FOIL method when the coefficient of the second-degree term is not 1.
- Factor using substitution.

VOCABULARY

- prime polynomial

OBJECTIVE 1 Factor trinomials when the coefficient of the second-degree term is 1.

We begin by finding the product of $x + 3$ and $x - 5$.

$$\begin{aligned} (x + 3)(x - 5) &= x^2 - 5x + 3x - 15 && \text{FOIL method} \\ &= x^2 - 2x - 15 && \text{Combine like terms.} \end{aligned}$$

By this result, the factored form of $x^2 - 2x - 15$ is $(x + 3)(x - 5)$.

$$\begin{array}{c} \text{Multiplication} \\ \xrightarrow{\hspace{1.5cm}} \\ \text{Factored form} \rightarrow (x + 3)(x - 5) = x^2 - 2x - 15 \leftarrow \text{Product} \\ \xleftarrow{\hspace{1.5cm}} \\ \text{Factoring} \end{array}$$

Because multiplying and factoring are operations that “undo” each other, factoring trinomials involves using the FOIL method in reverse. As shown here, the x^2 -term comes from multiplying x and x , and -15 comes from multiplying 3 and -5 .

$$\begin{array}{c}
 \text{Product of } x \quad \text{and } x \quad \text{is } x^2. \\
 \downarrow \quad \downarrow \quad \downarrow \\
 (x + 3)(x - 5) = x^2 - 2x - 15 \\
 \uparrow \quad \uparrow \quad \uparrow \\
 \text{Product of } 3 \quad \text{and } -5 \quad \text{is } -15.
 \end{array}$$

We find the term $-2x$ in $x^2 - 2x - 15$ by multiplying the outer terms, multiplying the inner terms, and adding the results.

$$\begin{array}{c}
 \text{Outer terms: } x(-5) = -5x \\
 \downarrow \quad \downarrow \\
 (x + 3)(x - 5) \\
 \uparrow \quad \uparrow \\
 \text{Inner terms: } 3 \cdot x = 3x \\
 \text{Add to obtain } -2x.
 \end{array}$$

STUDY SKILLS REMINDER

Are you getting the most out of your class time?

Review Study Skill 3, Taking Lecture Notes.

Factoring $x^2 + bx + c$ (Coefficient of Second-Degree Term = 1)

- Step 1** Find pairs of integers whose product is c . Write all pairs whose product is c , the third term of the trinomial.
- Step 2** Find the sum of each pair. Choose the pair whose sum is b , the coefficient of the middle term.
- Step 3** Factor the trinomial as the product of two binomials. Use the pair of integers identified in Step 2.

If there is no pair of integers in Step 2 whose sum is b , then the polynomial cannot be factored. A polynomial that cannot be factored with integer coefficients is a **prime polynomial**.

Examples: $x^2 + x + 2$, $x^2 - x - 1$, $2x^2 + x + 7$ Prime polynomials



NOW TRY EXERCISE 1

Factor each trinomial.

- (a) $t^2 - t - 30$
 (b) $w^2 + 12w + 32$

EXAMPLE 1 Factoring Trinomials in $x^2 + bx + c$ Form

Factor each trinomial.

(a) $x^2 + 2x - 35$

Step 1 Find pairs of integers whose product is -35 .

$$\begin{array}{l}
 35(-1) \\
 -35(1) \\
 7(-5) \\
 -7(5)
 \end{array}$$

Step 2 Write sums of those pairs of integers.

$$\begin{array}{l}
 35 + (-1) = 34 \\
 -35 + 1 = -34 \\
 7 + (-5) = 2 \leftarrow \text{Coefficient of the middle term} \\
 -7 + 5 = -2
 \end{array}$$

Step 3 The integers 7 and -5 have the necessary product and sum.

$$x^2 + 2x - 35 \text{ factors as } (x + 7)(x - 5). \quad \text{Multiply to check.}$$

(b) $r^2 + 8r + 12$

Look for two integers with a product of 12 and a sum of 8. Of all pairs having a product of 12, only the pair 6 and 2 has a sum of 8.

$$r^2 + 8r + 12 \text{ factors as } (r + 6)(r + 2). \quad \text{Multiply to check.}$$

By the commutative property, it is equally correct to write $(r + 2)(r + 6)$.

NOW TRY ANSWERS

1. (a) $(t - 6)(t + 5)$
 (b) $(w + 4)(w + 8)$


NOW TRY EXERCISE 2
Factor $m^2 + 12m - 11$.

NOW TRY EXERCISE 3
Factor $a^2 + ab - 20b^2$.

NOW TRY EXERCISE 4
Factor $5w^3 - 40w^2 + 60w$.**EXAMPLE 2** Recognizing a Prime PolynomialFactor $m^2 + 6m + 7$.

Look for two integers whose product is 7 and whose sum is 6. Only 7 and 1 and -7 and -1 give a product of 7. Neither pair has a sum of 6, so $m^2 + 6m + 7$ cannot be factored with integer coefficients and is prime.

NOW TRY **EXAMPLE 3** Factoring a Multivariable TrinomialFactor $x^2 + 6ax - 16a^2$.This trinomial is in the form $x^2 + bx + c$, where $b = 6a$ and $c = -16a^2$.**Step 1** Find pairs of expressions whose product is $-16a^2$.

$$\begin{aligned} &16a(-a) \\ &-16a(a) \\ &8a(-2a) \\ &-8a(2a) \\ &-4a(4a) \end{aligned}$$

Step 2 Write sums of those pairs of expressions.

$$\begin{aligned} 16a + (-a) &= 15a \\ -16a + a &= -15a \\ 8a + (-2a) &= 6a \leftarrow \text{Coefficient of the middle term} \\ -8a + 2a &= -6a \\ -4a + 4a &= 0 \end{aligned}$$

Step 3 The expressions $8a$ and $-2a$ have the necessary product and sum.

$$x^2 + 6ax - 16a^2 \text{ factors as } (x + 8a)(x - 2a).$$

CHECK $(x + 8a)(x - 2a)$

$$= x^2 - 2ax + 8ax - 16a^2 \quad \text{FOIL method}$$

$$= x^2 + 6ax - 16a^2 \quad \checkmark \quad \text{Original polynomial}$$

NOW TRY **EXAMPLE 4** Factoring a Trinomial (Terms Have a Common Factor)Factor $16y^3 - 32y^2 - 48y$.

$$\begin{aligned} &16y^3 - 32y^2 - 48y \\ &= 16y(y^2 - 2y - 3) \quad \text{Factor out the GCF, } 16y. \end{aligned}$$

To factor $y^2 - 2y - 3$, look for two integers whose product is -3 and whose sum is -2 . The necessary integers are -3 and 1 .

Remember to include the GCF, $16y$.

$$= 16y(y - 3)(y + 1) \quad \text{Factor the trinomial.}$$

NOW TRY 

CAUTION When factoring, always look for a common factor first. Remember to write the common factor as part of the answer.

OBJECTIVE 2 Factor trinomials by grouping when the coefficient of the second-degree term is not 1.

We can use a generalization of the method shown in **Objective 1** to factor a trinomial of the form $ax^2 + bx + c$, where $a \neq 1$. To factor $3x^2 + 7x + 2$, for example, we first identify the values a , b , and c .

$$\begin{array}{c} ax^2 + bx + c \\ \downarrow \quad \downarrow \quad \downarrow \\ 3x^2 + 7x + 2, \text{ so } a = 3, \quad b = 7, \quad c = 2 \end{array}$$

NOW TRY ANSWERS

- prime
- $(a + 5b)(a - 4b)$
- $5w(w - 6)(w - 2)$

The product ac is $3 \cdot 2 = 6$, so we must find integers having a product of 6 and a sum of 7 (because the middle term has coefficient 7). The necessary integers are 1 and 6.

$$\begin{aligned} & 3x^2 + 7x + 2 \\ &= 3x^2 + \underbrace{x + 6x} + 2 && \text{Write } 7x \text{ as } 1x + 6x, \text{ or } x + 6x. \\ & \qquad \qquad \qquad \uparrow \\ & \qquad \qquad \qquad x + 6x = 7x \\ &= (3x^2 + x) + (6x + 2) && \text{Group the terms.} \\ &= x(3x + 1) + 2(3x + 1) && \text{Factor by grouping.} \\ &= \underbrace{(3x + 1)}(x + 2) && \text{Factor out the common factor.} \end{aligned}$$

Check by multiplying.

 **NOW TRY EXERCISE 5**

Factor $8y^2 - 10y - 3$.

EXAMPLE 5 Factoring a Trinomial in $ax^2 + bx + c$ Form

Factor $12r^2 - 5r - 2$.

Here $a = 12$, $b = -5$, and $c = -2$, so the product ac is -24 . The two integers whose product is -24 and whose sum b is -5 are 3 and -8 .

$$\begin{aligned} & 12r^2 - 5r - 2 \\ &= 12r^2 + 3r - 8r - 2 && \text{Write } -5r \text{ as } 3r - 8r. \\ &= (12r^2 + 3r) + (-8r - 2) && \text{Group the terms.} \\ &= 3r(4r + 1) - 2(4r + 1) && \text{Factor by grouping.} \\ &= \underbrace{(4r + 1)}(3r - 2) && \text{Factor out the common factor.} \end{aligned}$$

Check by multiplying.

NOW TRY 

OBJECTIVE 3 Factor trinomials using the FOIL method when the coefficient of the second-degree term is not 1.

This method involves trying repeated combinations and using the FOIL method.

EXAMPLE 6 Factoring Trinomials in $ax^2 + bx + c$ Form

Factor each trinomial.

(a) $3x^2 + 7x + 2$

The goal is to find the correct numbers to put in the blanks.

$$(\underline{\hspace{1cm}}x + \underline{\hspace{1cm}})(\underline{\hspace{1cm}}x + \underline{\hspace{1cm}})$$

Addition signs are used because all the signs in the polynomial indicate addition. The first two expressions have a product of $3x^2$, so they must be $3x$ and x .

$$(3x + \underline{\hspace{1cm}})(x + \underline{\hspace{1cm}})$$

The product of the two last terms must be 2, so the numbers must be 2 and 1. There is a choice. The 2 could be placed with the $3x$ or with the x . Only one of these choices will give the correct middle term, $7x$.

$$\begin{array}{c} \text{3x} \\ \text{-----} \\ (3x + 2)(x + 1) \\ \text{-----} \\ \text{2x} \end{array}$$

$$3x + 2x = 5x$$

Wrong middle term

$$\begin{array}{c} \text{6x} \\ \text{-----} \\ (3x + 1)(x + 2) \\ \text{-----} \\ \text{x} \end{array}$$

$$6x + x = 7x$$

Correct middle term

Use the FOIL method to check each factored form.

NOW TRY ANSWER

5. $(2y - 3)(4y + 1)$

Therefore, $3x^2 + 7x + 2$ factors as $(3x + 1)(x + 2)$.


**NOW TRY
EXERCISE 6**
Factor $10r^2 + 19r + 6$.

(b) $12r^2 - 5r - 2$

To reduce the number of trials, we note that the terms of the trinomial have greatest common factor 1. This means that neither of its factors can have a common factor except 1. We try 4 and 3 for the two first terms.

$$(4r \quad)(3r \quad)$$

The factors of -2 are -2 and 1 or -1 and 2 . We try both possibilities.

$$(4r - 2)(3r + 1)$$

Wrong: The terms of $4r - 2$ have a common factor of 2. This cannot be correct because 2 is not a factor of $12r^2 - 5r - 2$.

$$(4r - 1)(3r + 2)$$

$8r - 3r = 5r$
Wrong middle term

The middle term on the right is $5r$, instead of the $-5r$ that is needed. We obtain $-5r$ by interchanging the signs of the second terms in the factors.

$$(4r + 1)(3r - 2)$$

$-8r + 3r = -5r$
Correct middle term

Thus, $12r^2 - 5r - 2$ factors as $(4r + 1)(3r - 2)$. (Compare to **Example 5**.)

CHECK $(4r + 1)(3r - 2)$

$$= 12r^2 - 8r + 3r - 2 \quad \text{FOIL method}$$

$$= 12r^2 - 5r - 2 \quad \checkmark \quad \text{Original polynomial}$$

NOW TRY 

NOTE As shown in **Example 6(b)**, if the terms of a polynomial have greatest common factor 1, then none of the terms of its factors can have a common factor (except 1). Remembering this will eliminate some potential factors.

Factoring $ax^2 + bx + c$ (GCF of $a, b, c = 1$)

Step 1 Find pairs of integers whose product is a . Write all pairs of integer factors of a , the coefficient of the second-degree term.

Step 2 Find pairs of integers whose product is c . Write all pairs of integer factors of c , the last term.

Step 3 Choose inner and outer terms. Use the FOIL method, and try various combinations of the factors from Steps 1 and 2 until the necessary middle term is obtained.

If no such combinations exist, the trinomial is prime.

NOW TRY ANSWER

6. $(2r + 3)(5r + 2)$


**NOW TRY
EXERCISE 7**
Factor $12a^2 - 19ab - 21b^2$.
EXAMPLE 7 Factoring a Multivariable Trinomial
Factor $18m^2 - 19mx - 12x^2$.

Follow the steps in the preceding box to factor the trinomial. There are many possible factors of both 18 and -12 . Try 6 and 3 for 18 and -3 and 4 for -12 .

$$\begin{array}{l|l} (6m - 3x)(3m + 4x) & (6m + 4x)(3m - 3x) \\ \text{Wrong: common factor} & \text{Wrong: common factors} \end{array}$$

Because 6 and 3 do not work as factors of 18, try 9 and 2 instead, with 3 and -4 as factors of -12 .

$$\begin{array}{l|l} (9m + 3x)(2m - 4x) & \begin{array}{l} \text{27mx} \\ \overbrace{(9m - 4x)(2m + 3x)} \\ \text{-8mx} \\ \hline 27mx + (-8mx) = 19mx \\ \text{Wrong middle term} \end{array} \\ \text{Wrong: common factors} & \end{array}$$

The result on the right differs from the correct middle term only in sign, so interchange the signs of the second terms in the factors.

$$18m^2 - 19mx - 12x^2 \text{ factors as } (9m + 4x)(2m - 3x).$$

Check by multiplying.

NOW TRY 

**NOW TRY
EXERCISE 8**
Factor $-8x^2 + 22x - 15$.
EXAMPLE 8 Factoring a Trinomial in $ax^2 + bx + c$ Form ($a < 0$)
Factor $-3x^2 + 16x + 12$.

While we could factor directly, it is helpful to first factor out -1 so that the coefficient of the x^2 -term is positive.

$$\begin{aligned} & -3x^2 + 16x + 12 \\ &= -1(3x^2 - 16x - 12) && \text{Factor out } -1. \\ &= -1(3x + 2)(x - 6) && \text{Factor the trinomial.} \\ &= -(3x + 2)(x - 6) && -1a = -a \end{aligned}$$

NOW TRY 
NOTE The factored form in **Example 8** can be written in other ways, such as

$$(-3x - 2)(x - 6) \quad \text{and} \quad (3x + 2)(-x + 6).$$

Verify that these both give the original trinomial when multiplied.


**NOW TRY
EXERCISE 9**
Factor $16y^3 + 33y^2 - 9y$.
EXAMPLE 9 Factoring a Trinomial (Terms Have a Common Factor)
Factor $16y^3 + 24y^2 - 16y$.

$$16y^3 + 24y^2 - 16y$$

$$\begin{aligned} &= 8y(2y^2 + 3y - 2) && \text{GCF} = 8y \\ &= 8y(2y - 1)(y + 2) && \text{Factor the trinomial.} \end{aligned}$$

Remember the common factor.

NOW TRY 
NOW TRY ANSWERS

7. $(4a + 3b)(3a - 7b)$

8. $-(4x - 5)(2x - 3)$

9. $3y(4y - 1)(y + 3)$

2. Which is the completely factored form of the trinomial $2x^6 - 5x^5 - 3x^4$?
- A. $x^4(2x + 1)(x - 3)$ B. $x^4(2x - 1)(x + 3)$
 C. $(2x^5 + x^4)(x - 3)$ D. $x^3(2x^2 + x)(x - 3)$
3. Which is the completely factored form of the trinomial $4x^2 - 4x - 24$?
- A. $4(x - 2)(x + 3)$ B. $4(x + 2)(x + 3)$
 C. $4(x + 2)(x - 3)$ D. $4(x - 2)(x - 3)$
4. Which is *not* a factored form of the trinomial $-x^2 + 16x - 60$?
- A. $(x - 10)(-x + 6)$ B. $(-x - 10)(x + 6)$
 C. $(-x + 10)(x - 6)$ D. $-1(x - 10)(x - 6)$
5. **Concept Check** When a student was given the polynomial $4x^2 + 2x - 20$ to factor completely on a test, the student lost some credit when her answer was $(4x + 10)(x - 2)$.
6. **Concept Check** When asked to factor $x^2y^2 - 6x^2 + 5y^2 - 30$, a student gave the following incorrect answer:
 $x^2(y^2 - 6) + 5(y^2 - 6)$

WHAT WENT WRONG? Give the correct answer.

WHAT WENT WRONG? What is the correct answer?

Complete each factoring. See Examples 1 and 3–8.

- | | | |
|-----------------------------------------------------------------|----------------------------------------------------------------------|---------------------------------------------------------------------|
| 7. $x^2 + 8x + 15$
= $(x + 5)(\underline{\hspace{2cm}})$ | 8. $y^2 + 11y + 18$
= $(y + 2)(\underline{\hspace{2cm}})$ | 9. $m^2 - 10m + 21$
= $(m - 3)(\underline{\hspace{2cm}})$ |
| 10. $n^2 - 14n + 48$
= $(n - 6)(\underline{\hspace{2cm}})$ | 11. $r^2 - r - 20$
= $(r + 4)(\underline{\hspace{2cm}})$ | 12. $x^2 + 4x - 32$
= $(x - 4)(\underline{\hspace{2cm}})$ |
| 13. $x^2 + ax - 6a^2$
= $(x + 3a)(\underline{\hspace{2cm}})$ | 14. $m^2 - 3mn - 10n^2$
= $(m + 2n)(\underline{\hspace{2cm}})$ | 15. $4x^2 - 4x - 3$
= $(2x + 1)(\underline{\hspace{2cm}})$ |
| 16. $6z^2 - 11z + 4$
= $(3z - 4)(\underline{\hspace{2cm}})$ | 17. $12u^2 + 10uv + 2v^2$
= $2(3u + v)(\underline{\hspace{2cm}})$ | 18. $16p^2 - 4pq - 2q^2$
= $2(2p - q)(\underline{\hspace{2cm}})$ |

Factor each trinomial. See Examples 1–9.

- | | | |
|------------------------------------|------------------------------------|---------------------------|
| 19. $y^2 + 7y - 30$ | 20. $z^2 + 2z - 24$ | 21. $p^2 + 15p + 56$ |
| 22. $k^2 - 11k + 30$ | 23. $m^2 - 11m + 60$ | 24. $p^2 - 12p - 27$ |
| 25. $a^2 - 2ab - 35b^2$ | 26. $z^2 + 8zw + 15w^2$ | 27. $a^2 - 9ab + 18b^2$ |
| 28. $k^2 - 11hk + 28h^2$ | 29. $x^2y^2 + 12xy + 18$ | 30. $p^2q^2 - 5pq - 18$ |
| 31. $-6m^2 - 13m + 15$ | 32. $-15y^2 + 17y + 18$ | 33. $10x^2 + 3x - 18$ |
| 34. $8k^2 + 34k + 35$ | 35. $20k^2 + 47k + 24$ | 36. $27z^2 + 42z - 5$ |
| 37. $15a^2 - 22ab + 8b^2$ | 38. $14c^2 - 17cd - 6d^2$ | 39. $36m^2 - 60m + 25$ |
| 40. $25r^2 - 90r + 81$ | 41. $40x^2 + xy + 6y^2$ | 42. $15p^2 + 24pq + 8q^2$ |
| 43. $6x^2z^2 + 5xz - 4$ | 44. $8m^2n^2 - 10mn + 3$ | 45. $24x^2 + 42x + 15$ |
| 46. $36x^2 + 18x - 4$ | 47. $-15a^2 - 70a + 120$ | 48. $-12a^2 - 10a + 42$ |
| 49. $-11x^3 + 110x^2 - 264x$ | 50. $-9k^3 - 36k^2 + 189k$ | |
| 51. $2x^3y^3 - 48x^2y^4 + 288xy^5$ | 52. $6m^3n^2 - 60m^2n^3 + 150mn^4$ | |
| 53. $6a^3 + 12a^2 - 90a$ | 54. $3m^4 + 6m^3 - 72m^2$ | |
| 55. $13y^3 + 39y^2 - 52y$ | 56. $4p^3 + 24p^2 - 64p$ | |

57. $12p^3 - 12p^2 + 3p$

58. $45t^3 + 60t^2 + 20t$

Factor each trinomial. See Example 10.

59. $12p^6 - 32p^3r + 5r^2$

60. $2y^6 + 7xy^3 + 6x^2$

61. $10(k+1)^2 - 7(k+1) + 1$

62. $4(m-5)^2 - 4(m-5) - 15$

63. $3(m+p)^2 - 7(m+p) - 20$

64. $4(x-y)^2 - 23(x-y) - 6$

Factor each trinomial. See Example 11.

65. $p^4 - 10p^2 + 16$

66. $k^4 + 10k^2 + 9$

67. $2x^4 - 9x^2 - 18$

68. $6z^4 + z^2 - 1$

69. $16x^4 + 16x^2 + 3$

70. $9r^4 + 9r^2 + 2$

Extending Skills Factor each trinomial. (Hint: Factor out the GCF first.)

71. $z^2(z-x) - zx(x-z) - 2x^2(z-x)$

72. $r^2(r-s) - 5rs(s-r) - 6s^2(r-s)$

73. $a^2(a+b)^2 - ab(a+b)^2 - 6b^2(a+b)^2$

74. $m^2(m-p)^2 - mp(m-p)^2 - 12p^2(m-p)^2$

75. $p^2(p+q) + 4pq(p+q) + 3q^2(p+q)$

76. $2k^2(5-y) - 7k(5-y) + 5(5-y)$

RELATING CONCEPTS For Individual or Group Work (Exercises 77–82)

The trinomial $6x^2 + 7x - 20$ may be factored using substitution. To see this, work Exercises 77–82 in order.

77. Multiply each term of $6x^2 + 7x - 20$ by 6 (the leading coefficient). Then compensate by dividing the result by 6 (that is, write it as a fraction with denominator 6).

78. The first term in the numerator of the answer from Exercise 77 is now a perfect square. Let $t = 6x$ (the square root of $36x^2$), and rewrite the expression.

79. Factor the numerator of the expression from Exercise 78 in terms of the variable t .

80. Replace t with $6x$ in the answer from Exercise 79, and simplify to obtain the correct factorization of the trinomial.

81. Factor $3x^2 - 4x - 32$ using the method just described. (Multiply each term and then divide the result by the leading coefficient 3.)

82. Factor $12x^2 + 35x + 8$ using the method just described. (Multiply and divide by 12.)

5.3 Special Factoring

OBJECTIVES

- Factor a difference of squares.
- Factor a perfect square trinomial.
- Factor a difference of cubes.
- Factor a sum of cubes.

OBJECTIVE 1 Factor a difference of squares.

The special products introduced earlier are used in reverse when factoring. Recall that the product of the sum and difference of two terms leads to a **difference of squares**.

Factoring a Difference of Squares

$$x^2 - y^2 = (x + y)(x - y)$$

EXAMPLE 1 Factoring Differences of Squares

Factor each polynomial.

(a) $t^2 - 36$

$$= t^2 - 6^2 \quad 36 = 6^2$$

$$= (t + 6)(t - 6) \quad \text{Factor the difference of squares.}$$

(b) $4a^2 - 64$

$$= 4(a^2 - 16) \quad \text{Factor out the common factor, 4.}$$

$$= 4(a + 4)(a - 4) \quad \text{Factor the difference of squares.}$$

Always check for a common factor first.

VOCABULARY

- difference of squares
- perfect square trinomial
- difference of cubes
- sum of cubes

NOW TRY EXERCISE 1

Factor each polynomial.

(a) $4m^2 - 25n^2$

(b) $9x^2 - 729$

(c) $(a + b)^2 - 25$

(d) $v^4 - 1$

$$(c) \quad \begin{array}{ccccccc} x^2 & - & y^2 & = & (x + y) & (x - y) \\ \downarrow & & \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow \\ (4m)^2 & - & (7p)^2 & = & (4m + 7p) & (4m - 7p) \end{array}$$

$$(d) \quad \begin{array}{ccccccc} x^2 & - & y^2 & = & (x + y) & (x - y) \\ \downarrow & & \downarrow & & \downarrow & \downarrow & \downarrow & \downarrow \\ (9k)^2 & - & (a + 2)^2 & = & (9k + a + 2) & (9k - (a + 2)) \\ & & & & & & & \downarrow \\ & & & & & & & (9k - a - 2) \end{array}$$

$$= (9k + a + 2)(9k - a - 2)$$

We could have used the method of substitution here.

(e) $x^4 - 81$

$$= (x^2 + 9)(x^2 - 9) \quad \text{Factor the difference of squares.}$$

$$= (x^2 + 9)(x + 3)(x - 3) \quad \text{Factor the difference of squares again.}$$

This sum of squares cannot be factored.

Don't stop here.

NOW TRY**Sum of Squares**If x and y have greatest common factor 1, then the sum of squares $x^2 + y^2$ cannot be factored using real numbers and is prime.Examples: $x^2 + 9$, $t^2 + 36$, $4m^2 + 49$ Prime polynomialsA sum of squares can be factored *only* if the terms have a common factor.Example: $4x^2 + 36 = 4(x^2 + 9)$ Factor out the GCF, 4.**OBJECTIVE 2** Factor a perfect square trinomial.

Two other special products lead to the following rules for factoring.

Factoring a Perfect Square Trinomial

$$x^2 + 2xy + y^2 = (x + y)^2$$

$$x^2 - 2xy + y^2 = (x - y)^2$$

NOW TRY ANSWERS

1. (a) $(2m + 5n)(2m - 5n)$
- (b) $9(x + 9)(x - 9)$
- (c) $(a + b + 5)(a + b - 5)$
- (d) $(v^2 + 1)(v + 1)(v - 1)$

Because the trinomial $x^2 + 2xy + y^2$ is the square of $x + y$, it is a **perfect square trinomial**. Both the first and the last terms of the trinomial must be perfect squares.

In the factored form $(x + y)^2$, twice the product of the first and the last terms must give the middle term of the trinomial.

$$4m^2 + 20m + 25$$

$$4m^2 = (2m)^2, 25 = 5^2,$$

$$\text{and } 2(2m)(5) = 20m.$$

Perfect square trinomial

$$p^2 - 8p + 64$$

p^2 and $64 = 8^2$ are perfect squares, but the middle term would have to be $16p$ or $-16p$.
Not a perfect square trinomial

NOW TRY EXERCISE 2

Factor each polynomial.

- (a) $a^2 + 12a + 36$
 (b) $16x^2 - 56xy + 49y^2$
 (c) $y^2 - 16y + 64 - z^2$

EXAMPLE 2 Factoring Perfect Square Trinomials

Factor each polynomial.

(a) $144p^2 - 120p + 25$

Here, $144p^2 = (12p)^2$ and $25 = 5^2$. The sign on the middle term is $-$, so if $144p^2 - 120p + 25$ is a perfect square trinomial, the factored form will have to be

$$(12p - 5)^2.$$

Determine twice the product of the two terms to see whether this is correct.

$$2(12p)(-5) = -120p \leftarrow \text{Desired middle term}$$

This is the middle term of the given trinomial.

$$144p^2 - 120p + 25 \text{ factors as } (12p - 5)^2.$$

(b) $4m^2 + 20mn + 49n^2$

If this is a perfect square trinomial, it will equal $(2m + 7n)^2$. If multiplied out, this squared binomial has a middle term of

$$2(2m)(7n) = 28mn, \text{ which does not equal } 20mn.$$

This trinomial cannot be factored by any other method either. It is prime.

(c) $(r + 5)^2 + 6(r + 5) + 9 \quad 9 = 3^2$
 $= [(r + 5) + 3]^2 \quad 2(r + 5)(3) = 6(r + 5) \text{ is the middle term.}$
 $= (r + 8)^2 \quad \text{Add.}$

(d) $m^2 - 8m + 16 - p^2$ The first three terms here form a perfect square trinomial.
 $= (m^2 - 8m + 16) - p^2$ Group the terms of the perfect square trinomial.
 $= (m - 4)^2 - p^2$ Factor the perfect square trinomial.
 $= (m - 4 + p)(m - 4 - p)$ Factor the difference of squares. **NOW TRY**

NOTE Perfect square trinomials can be factored by the general methods for other trinomials. The patterns given here provide “shortcuts.”

Example:

$$4m^2 + 20m + 25$$

$$= (2m + 5)(2m + 5) \quad 10m + 10m = 20m$$

$$= (2m + 5)^2 \quad a \cdot a = a^2$$

NOW TRY ANSWERS

2. (a) $(a + 6)^2$
 (b) $(4x - 7y)^2$
 (c) $(y - 8 + z)(y - 8 - z)$

OBJECTIVE 3 Factor a difference of cubes.

A **difference of cubes**, $x^3 - y^3$, can be factored as follows.

Factoring a Difference of Cubes

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2)$$

 **NOW TRY EXERCISE 3**

Factor each polynomial.

(a) $t^3 - 1$

(b) $125a^3 - 8b^3$

EXAMPLE 3 Factoring Differences of Cubes

Factor each polynomial.

$$x^3 - y^3 = (x - y)(x^2 + x \cdot y + y^2)$$

$$\begin{aligned} \text{(a) } m^3 - 8 &= m^3 - 2^3 = (m - 2)(m^2 + m \cdot 2 + 2^2) \\ &= (m - 2)(m^2 + 2m + 4) \end{aligned}$$

CHECK $(m - 2)(m^2 + 2m + 4)$

$$= m^3 + 2m^2 + 4m - 2m^2 - 4m - 8 \quad \text{Distributive property}$$

$$= m^3 - 8 \quad \checkmark \quad \text{Combine like terms.}$$

(b) $27x^3 - 8y^3$

$$= (3x)^3 - (2y)^3 \quad \text{Difference of cubes}$$

$$= (3x - 2y)[(3x)^2 + (3x)(2y) + (2y)^2] \quad \text{Factor.}$$

$$= (3x - 2y)(9x^2 + 6xy + 4y^2)$$

$$\begin{aligned} (3x)^2 &= 3^2x^2, \text{ not } 3x^2. \\ (2y)^2 &= 2^2y^2, \text{ not } 2y^2. \end{aligned}$$

(c) $1000k^3 - 27n^3$

$$= (10k)^3 - (3n)^3 \quad \text{Difference of cubes}$$

$$= (10k - 3n)[(10k)^2 + (10k)(3n) + (3n)^2] \quad \text{Factor.}$$

$$= (10k - 3n)(100k^2 + 30kn + 9n^2) \quad \text{Multiply.}$$

NOW TRY **OBJECTIVE 4** Factor a sum of cubes.

Although the binomial $x^2 + y^2$ (a sum of *squares*) cannot be factored with real numbers, a **sum of cubes**, such as $x^3 + y^3$, can be factored as follows.

Factoring a Sum of Cubes

$$x^3 + y^3 = (x + y)(x^2 - xy + y^2)$$

Compare the rules for factoring a *difference* of cubes and a *sum* of cubes.

$$\begin{aligned} x^3 - y^3 &= (x - y)(x^2 + xy + y^2) \\ &\quad \begin{array}{l} \uparrow \text{ Same sign } \uparrow \\ \downarrow \text{ Always positive } \downarrow \\ \uparrow \text{ Opposite sign } \uparrow \end{array} \\ \\ x^3 + y^3 &= (x + y)(x^2 - xy + y^2) \\ &\quad \begin{array}{l} \downarrow \text{ Same sign } \downarrow \\ \downarrow \text{ Opposite sign } \downarrow \\ \uparrow \text{ Always positive } \uparrow \end{array} \end{aligned}$$

NOW TRY ANSWERS

3. (a) $(t - 1)(t^2 + t + 1)$

(b) $(5a - 2b)(25a^2 + 10ab + 4b^2)$


NOW TRY EXERCISE 4

Factor each polynomial.

(a) $1000 + z^3$

(b) $81a^6 + 3b^3$

(c) $(x - 3)^3 + y^3$

EXAMPLE 4 Factoring Sums of Cubes

Factor each polynomial.

(a) $r^3 + 27$

$$= r^3 + 3^3 \quad \text{Sum of cubes}$$

$$= (r + 3)(r^2 - 3r + 3^2) \quad \text{Factor.}$$

$$= (r + 3)(r^2 - 3r + 9) \quad \text{This trinomial cannot be factored further.}$$

(b) $27z^3 + 125$

$$= (3z)^3 + 5^3 \quad \text{Sum of cubes}$$

$$= (3z + 5)[(3z)^2 - (3z)(5) + 5^2] \quad \text{Factor.}$$

$$= (3z + 5)(9z^2 - 15z + 25) \quad \text{Multiply; } (3z)^2 = 3^2z^2$$

(c) $125t^3 + 216s^6$

$$= (5t)^3 + (6s^2)^3 \quad \text{Sum of cubes}$$

$$= (5t + 6s^2)[(5t)^2 - (5t)(6s^2) + (6s^2)^2] \quad \text{Factor.}$$

$$= (5t + 6s^2)(25t^2 - 30ts^2 + 36s^4) \quad \text{Multiply.}$$

(d) $3x^2 + 192$

$$= 3(x^3 + 64) \quad \text{Factor out the common factor.}$$

$$= 3(x^3 + 4^3) \quad \text{Write as a sum of cubes.}$$

$$= 3(x + 4)(x^2 - 4x + 16) \quad \text{Factor.}$$

(e) $(x + 2)^3 + t^3$

$$= [(x + 2) + t][(x + 2)^2 - (x + 2)t + t^2] \quad \text{Sum of cubes}$$

$$= (x + 2 + t)(x^2 + 4x + 4 - xt - 2t + t^2) \quad \text{Multiply.}$$

NOW TRY **STUDY SKILLS REMINDER**

Make study cards to help you learn and remember the material in this chapter.

Review Study Skill 5,
Using Study Cards.**CAUTION** A common error when factoring $x^3 + y^3$ or $x^3 - y^3$ is to think that the xy -term has a coefficient of 2. Because there is no coefficient of 2, expressions of the form $x^2 + xy + y^2$ and $x^2 - xy + y^2$ usually cannot be factored further.**NOW TRY ANSWERS**

4. (a) $(10 + z) \cdot (100 - 10z + z^2)$
 (b) $3(3a^2 + b) \cdot (9a^4 - 3a^2b + b^2)$
 (c) $(x - 3 + y) \cdot (x^2 - 6x + 9 - xy + 3y + y^2)$

Summary of Special Types of Factoring**Difference of Squares**

$$x^2 - y^2 = (x + y)(x - y)$$

Perfect Square Trinomial

$$x^2 + 2xy + y^2 = (x + y)^2$$

$$x^2 - 2xy + y^2 = (x - y)^2$$

Difference of Cubes

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2)$$

Sum of Cubes

$$x^3 + y^3 = (x + y)(x^2 - xy + y^2)$$

5.3 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Choose the letter(s) of the correct response.

- Which of the following binomials are differences of squares?
A. $64 - k^2$ B. $2x^2 - 25$ C. $k^2 + 9$ D. $4z^4 - 49$
- Which of the following binomials are sums or differences of cubes?
A. $64 + r^3$ B. $125 - p^6$ C. $9x^3 + 125$ D. $(x + y)^3 - 1$
- Which of the following trinomials are perfect squares?
A. $x^2 - 8x - 16$ B. $4m^2 + 20m + 25$
C. $9z^4 + 30z^2 + 25$ D. $25p^2 - 45p + 81$
- Which sum of squares can be factored? Factor it.
A. $x^2 + 100$ B. $9m^2 + 25$ C. $t^2 + 16$ D. $4x^2 + 64$
- Concept Check** A student incorrectly factored $x^2 + 4$ as $(x + 2)^2$.
- Concept Check** A student incorrectly factored $x^3 + 8$ as $(x + 2)^3$.

WHAT WENT WRONG? Explain this error.

WHAT WENT WRONG? Explain this error.

Factor each polynomial. See Examples 1–4.

- | | | |
|---------------------------------|--------------------------------|----------------------------|
| 7. $p^2 - 16$ | 8. $k^2 - 9$ | 9. $25x^2 - 4$ |
| 10. $36m^2 - 25$ | 11. $18a^2 - 98b^2$ | 12. $32c^2 - 98d^2$ |
| 13. $64m^4 - 4y^4$ | 14. $243x^4 - 3t^4$ | 15. $(y + z)^2 - 81$ |
| 16. $(h + k)^2 - 9$ | 17. $16 - (x + 3y)^2$ | 18. $64 - (r + 2t)^2$ |
| 19. $p^4 - 256$ | 20. $a^4 - 625$ | 21. $k^2 - 6k + 9$ |
| 22. $x^2 + 10x + 25$ | 23. $4z^2 + 4zw + w^2$ | 24. $9y^2 + 6yz + z^2$ |
| 25. $16m^2 - 8m + 1 - n^2$ | 26. $25c^2 - 20c + 4 - d^2$ | 27. $4r^2 - 12r + 9 - s^2$ |
| 28. $9a^2 - 24a + 16 - b^2$ | 29. $x^2 - y^2 + 2y - 1$ | 30. $-k^2 - h^2 + 2kh + 4$ |
| 31. $98m^2 + 84mn + 18n^2$ | 32. $80z^2 - 40zw + 5w^2$ | |
| 33. $(p + q)^2 + 2(p + q) + 1$ | 34. $(x + y)^2 + 6(x + y) + 9$ | |
| 35. $(a - b)^2 + 8(a - b) + 16$ | 36. $(m - n)^2 + 4(m - n) + 4$ | |
| 37. $x^3 - 27$ | 38. $y^3 - 64$ | 39. $216 - t^3$ |
| 40. $512 - m^3$ | 41. $x^3 + 64$ | 42. $r^3 + 343$ |
| 43. $1000 + y^3$ | 44. $729 + x^3$ | 45. $8x^3 + 1$ |
| 46. $27y^3 + 1$ | 47. $125x^3 - 216$ | 48. $8w^3 - 125$ |
| 49. $x^3 - 8y^3$ | 50. $z^3 - 125p^3$ | 51. $64g^3 - 27h^3$ |
| 52. $27a^3 - 8b^3$ | 53. $343p^3 + 125q^3$ | 54. $512t^3 + 27s^3$ |
| 55. $24n^3 + 81p^3$ | 56. $250x^3 + 16y^3$ | 57. $(y + z)^3 + 64$ |
| 58. $(p - q)^3 + 125$ | 59. $m^6 - 125$ | 60. $x^6 - 216$ |

61. $27 - 1000x^9$

62. $64 - 729p^9$

63. $125y^6 + z^3$

64. $64y^9 + z^6$

65. $125k - 64k^4$

66. $216k^2 + 125k^5$

Extending Skills In some cases, the method of factoring by grouping can be combined with the methods of special factoring discussed in this section. Consider this example.

$$8x^3 + 4x^2 + 27y^3 - 9y^2$$

$$= (8x^3 + 27y^3) + (4x^2 - 9y^2)$$

Associative and commutative properties

$$= (2x + 3y)(4x^2 - 6xy + 9y^2) + (2x + 3y)(2x - 3y)$$

Factor within groups.

$$= (2x + 3y)[(4x^2 - 6xy + 9y^2) + (2x - 3y)]$$

Factor out the greatest common factor, $2x + 3y$.

$$= (2x + 3y)(4x^2 - 6xy + 9y^2 + 2x - 3y)$$

Combine like terms.

In problems such as this, how we choose to group in the first step is essential to factoring correctly. If we reach a “dead end,” then we should group differently and try again.

Use the method just described to factor each polynomial.

67. $125p^3 + 25p^2 + 8q^3 - 4q^2$

68. $27x^3 + 9x^2 + y^3 - y^2$

69. $27a^3 + 15a - 64b^3 - 20b$

70. $1000k^3 + 20k - m^3 - 2m$

71. $8t^4 - 24t^3 + t - 3$

72. $y^4 + y^3 + y + 1$

73. $64m^2 - 512m^3 - 81n^2 + 729n^3$

74. $10x^2 + 5x^3 - 10y^2 + 5y^3$

RELATING CONCEPTS For Individual or Group Work (Exercises 75–80)

The binomial $x^6 - y^6$ may be considered as either a difference of squares or a difference of cubes. **Work Exercises 75–80 in order.**

75. Factor $x^6 - y^6$ by first factoring as a difference of squares. Then factor further by considering one of the factors as a sum of cubes and the other factor as a difference of cubes.

76. Based on the answer in **Exercise 75**, fill in the blank with the correct factors so that $x^6 - y^6$ is factored completely.

$$x^6 - y^6 = (x - y)(x + y) \underline{\hspace{2cm}}$$

77. Factor $x^6 - y^6$ by first factoring as a difference of cubes. Then factor further by considering one of the factors as a difference of squares.

78. Based on the answer in **Exercise 77**, fill in the blank with the correct factor so that $x^6 - y^6$ is factored.

$$x^6 - y^6 = (x - y)(x + y) \underline{\hspace{2cm}}$$

79. Notice that the factor written in the blank in **Exercise 78** is a fourth-degree polynomial, while the two factors written in the blank in **Exercise 76** are both second-degree polynomials. What must be true about the product of the two factors written in the blank in **Exercise 76**? Verify this.

80. If we have a choice of factoring as a difference of squares or a difference of cubes, how should we start to more easily obtain the completely factored form of the polynomial? Base the answer on the results in **Exercises 75–79**.

5.4 A General Approach to Factoring

OBJECTIVE

1 Factor any polynomial.

OBJECTIVE 1 Factor any polynomial.

A polynomial is *completely factored* when it is written as a product of prime polynomials with integer coefficients.

Factoring a Polynomial

Step 1 Factor out any common factor.

Step 2 How many terms are in the polynomial?

If the polynomial is a binomial, check to see whether it is a difference of squares, a difference of cubes, or a sum of cubes.

If the polynomial is a trinomial, check to see whether it is a perfect square trinomial. If it is not, use one of the following methods.

- To factor $x^2 + bx + c$, find two integers whose product is c and whose sum is b , the coefficient of the middle term.
- To factor $ax^2 + bx + c$, find two integers having product ac and sum b . Use these integers to rewrite the middle term, and factor by grouping.

Alternatively, use the FOIL method and try various combinations of the factors until the correct middle term is found.

If the polynomial has more than three terms, try to factor it by grouping.

Step 3 If any of the factors can be factored further, do so.

Step 4 Check the factored form by multiplying.

NOW TRY EXERCISE 1

Factor each polynomial.

- (a) $13x + 39$
 (b) $21x^3y^2 - 27x^2y^4$
 (c) $8y(m - n) - 5(m - n)$
 (d) $(y - 1)(2y + 1) - (y - 1)(y + 4)$

EXAMPLE 1 Factoring Out a Common Factor

Factor each polynomial.

- (a) $9p + 45$
 $= 9(p + 5)$ GCF = 9
- (b) $8m^2p^2 + 4mp$
 $= 4mp(2mp + 1)$ GCF = $4mp$
- (c) $5x(a + b) - y(a + b)$
 $= (a + b)(5x - y)$ Factor out $(a + b)$.
- (d) $(x - 4)(x + 2) + (x - 4)(2x - 1)$
 $= (x - 4)[(x + 2) + (2x - 1)]$ Factor out $x - 4$.
 $= (x - 4)(3x + 1)$ Combine like terms inside the brackets.

NOW TRY 

Factoring a Binomial

For a **binomial** (two terms), check for the following patterns.

Difference of squares $x^2 - y^2 = (x + y)(x - y)$

Difference of cubes $x^3 - y^3 = (x - y)(x^2 + xy + y^2)$

Sum of cubes $x^3 + y^3 = (x + y)(x^2 - xy + y^2)$

NOW TRY ANSWERS

1. (a) $13(x + 3)$
 (b) $3x^2y^2(7x - 9y^2)$
 (c) $(m - n)(8y - 5)$
 (d) $(y - 1)(y - 3)$


**NOW TRY
EXERCISE 2**

Factor each binomial if possible.

(a) $4a^2 - 49b^2$

(b) $9x^2 + 100$

(c) $27v^3 - 1000$

EXAMPLE 2 Factoring Binomials

Factor each binomial if possible.

$$\begin{aligned} \text{(a)} \quad 64m^2 - 9n^2 &= (8m)^2 - (3n)^2 && \text{Difference of squares} \\ &= (8m + 3n)(8m - 3n) && x^2 - y^2 = (x + y)(x - y) \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad 8p^3 - 27 &= (2p)^3 - 3^3 && \text{Difference of cubes} \\ &= (2p - 3)[(2p)^2 + (2p)(3) + 3^2] && x^3 - y^3 = (x - y)(x^2 + xy + y^2) \\ &= (2p - 3)(4p^2 + 6p + 9) && (2p)^2 = 2^2p^2 = 4p^2 \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad 1000m^3 + 1 &= (10m)^3 + 1^3 && \text{Sum of cubes} \\ &= (10m + 1)[(10m)^2 - (10m)(1) + 1^2] && x^3 + y^3 = (x + y)(x^2 - xy + y^2) \\ &= (10m + 1)(100m^2 - 10m + 1) && (10m)^2 = 10^2m^2 = 100m^2 \end{aligned}$$

(d) $25m^2 + 121$

This *sum* of squares is prime. There is no common factor (except 1).



NOTE Although the binomial $25m^2 + 625$ is a sum of squares, it can be factored because the greatest common factor of the terms is *not* 1.

$$25m^2 + 625 = 25(m^2 + 25) \quad \text{Factor out the common factor 25.}$$

This sum of squares cannot be factored further.

Factoring a Trinomial

For a **trinomial** (three terms), decide whether it is a perfect square trinomial of either of these forms.

$$x^2 + 2xy + y^2 = (x + y)^2 \quad \text{or} \quad x^2 - 2xy + y^2 = (x - y)^2$$

If not, use the methods shown in **Examples 3(c)–(f)**.

EXAMPLE 3 Factoring Trinomials

Factor each trinomial.

$$\begin{aligned} \text{(a)} \quad p^2 + 10p + 25 & \text{Perfect square trinomial} && \text{(b)} \quad 49z^2 - 42z + 9 & \text{Perfect square trinomial} \\ &= (p + 5)^2 && &= (7z - 3)^2 \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad y^2 - 5y - 6 &= (y - 6)(y + 1) && \text{The numbers } -6 \text{ and } 1 \text{ have a product of } -6 \text{ and a sum of } -5. \end{aligned}$$

NOW TRY ANSWERS

2. (a) $(2a + 7b)(2a - 7b)$

(b) prime

(c) $(3v - 10)(9v^2 + 30v + 100)$


**NOW TRY
EXERCISE 3**

Factor each trinomial.

(a) $25x^2 - 90x + 81$

(b) $7x^2 - 7xy - 84y^2$

(c) $12m^2 + 5m - 28$

(d) $12x^2 + 28xy - 5y^2$

$$\begin{aligned} \text{(d)} \quad 2k^2 - k - 6 &= (2k + 3)(k - 2) \\ &\quad \begin{array}{c} \text{---}4k\text{---} \\ \text{---}3k\text{---} \\ -4k + 3k = -k \end{array} \end{aligned}$$

Use the FOIL method and try various combinations of the factors until the correct middle term is found.

$$\begin{aligned} \text{(e)} \quad 10x^2 + xy - 3y^2 &= 10x^2 - 5xy + 6xy - 3y^2 \\ &\quad \begin{array}{c} \text{---}5xy\text{---} \\ \text{---}6xy\text{---} \\ xy \end{array} \end{aligned}$$

The integers -5 and 6 have a product of $10(-3) = -30$ and a sum of 1 , the coefficient of the middle term.

$$= (10x^2 - 5xy) + (6xy - 3y^2)$$

Group the terms.

$$= 5x(2x - y) + 3y(2x - y)$$

Factor each group.

$$= (2x - y)(5x + 3y)$$

Factor out the common factor.

(f) $28z^2 + 6z - 10$

Remember the common factor.

$$= 2(14z^2 + 3z - 5)$$

Factor out the common factor.

$$= 2(7z + 5)(2z - 1)$$

Factor the trinomial.

NOW TRY 

If a polynomial has more than three terms, consider factoring by grouping.


**NOW TRY
EXERCISE 4**

Factor each polynomial.

(a) $5a^3 + 5a^2b - ab^2 - b^3$

(b) $9u^2 - 48u + 64 - v^2$

(c) $x^3 - 9y^2 - 27y^3 + x^2$

EXAMPLE 4 Factoring Polynomials with More Than Three Terms

Factor each polynomial.

$$\begin{aligned} \text{(a)} \quad xy^2 - y^3 + x^3 - x^2y &= (xy^2 - y^3) + (x^3 - x^2y) && \text{Group the terms.} \\ &= y^2(x - y) + x^2(x - y) && \text{Factor each group.} \\ &= (x - y)(y^2 + x^2) && x - y \text{ is a common factor.} \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad 20k^3 + 4k^2 - 45k - 9 &= (20k^3 + 4k^2) + (-45k - 9) && \text{Be careful with signs.} \\ &= 4k^2(5k + 1) - 9(5k + 1) && \text{Group the terms.} \\ &= (5k + 1)(4k^2 - 9) && \text{Factor each group.} \\ &= (5k + 1)(2k + 3)(2k - 3) && 5k + 1 \text{ is a common factor.} \\ &&& \text{Factor the difference of squares.} \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad 4a^2 + 4a + 1 - b^2 &= (4a^2 + 4a + 1) - b^2 \\ &= (2a + 1)^2 - b^2 && \text{The first three terms form a perfect square trinomial. } 4a^2 = (2a)^2, 1 = 1^2, \text{ and } 2(2a)(1) = 4a, \text{ as required.} \\ &= (2a + 1 + b)(2a + 1 - b) && \text{Factor the perfect square trinomial.} \\ &&& \text{Factor the difference of squares.} \end{aligned}$$

$$\begin{aligned} \text{(d)} \quad 8m^3 + 4m^2 - n^3 - n^2 &= (8m^3 - n^3) + (4m^2 - n^2) && \text{Rearrange and group the terms.} \\ &= \underbrace{(2m - n)(4m^2 + 2mn + n^2)}_{\text{Difference of cubes}} + \underbrace{(2m - n)(2m + n)}_{\text{Difference of squares}} \\ &= (2m - n)(4m^2 + 2mn + n^2) + (2m - n)(2m + n) \\ &= (2m - n)(4m^2 + 2mn + n^2 + 2m + n) && \text{Factor the difference of cubes and the difference of squares.} \\ &&& \text{Factor out the common factor } 2m - n. \end{aligned}$$

NOW TRY **NOW TRY ANSWERS**

3. (a) $(5x - 9)^2$

(b) $7(x + 3y)(x - 4y)$

(c) $(4m + 7)(3m - 4)$

(d) $(6x - y)(2x + 5y)$

4. (a) $(a + b)(5a^2 - b^2)$

(b) $(3u - 8 + v)(3u - 8 - v)$


(c) $(x - 3y)(x^2 + 3xy + 9y^2 + x + 3y)$

NOTE Observe the final two lines in **Example 4(d)**. Students often ask, “What happened to the other $(2m - n)$?” Think of the variable a in the expression $ax + ay$, which is factored as $a(x + y)$. The binomial $(2m - n)$ is handled just like a in this simpler example.

5.4 Exercises

FOR
EXTRA
HELP

 MyLab Math

 Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

How are you doing on your homework? **Review Study Skill 4, Completing Your Homework.**

Concept Check Match each polynomial in Column I with the method or methods for factoring it in Column II. The choices in Column II may be used once, more than once, or not at all.

I

1. (a) $49x^2 - 81y^2$
- (b) $125z^6 + 1$
- (c) $88r^2 - 55s^2$
- (d) $64a^3 - 8b^9$
- (e) $50x^2 - 128y^4$

I

2. (a) $ab - 5a + 3b - 15$
- (b) $z^2 - 3z + 6$
- (c) $25x^2 + 100$
- (d) $r^2 - 24r + 144$
- (e) $2y^2 + 36y + 162$

II

- A. Factor out the GCF.
- B. Factor a difference of squares.
- C. Factor a difference of cubes.
- D. Factor a sum of cubes.
- E. The polynomial is prime.

II

- A. Factor out the GCF.
- B. Factor a perfect square trinomial.
- C. Factor by grouping.
- D. Factor into two distinct binomials.
- E. The polynomial is prime.

The following exercises are of mixed variety. Factor each polynomial. **See Examples 1–4.**

3. $3p^4 - 3p^3 - 90p^2$
4. $k^4 - 16$
5. $3a^2pq + 3abpq - 90b^2pq$
6. $49z^2 - 16$
7. $225p^2 + 256$
8. $18m^3n + 3m^2n^2 - 6mn^3$
9. $6b^2 - 17b - 3$
10. $k^2 - 6k - 16$
11. $x^3 - 1000$
12. $6t^2 + 19tu - 77u^2$
13. $4(p + 2) + m(p + 2)$
14. $40p - 32r$
15. $9m^2 - 45m + 18m^3$
16. $4k^2 + 28kr + 49r^2$
17. $54m^3 - 2000$
18. $mn - 2n + 5m - 10$
19. $9m^2 - 30mn + 25n^2$
20. $2a^2 - 7a - 4$
21. $kq - 9q + kr - 9r$
22. $56k^3 - 875$
23. $16z^3x^2 - 32z^2x$
24. $9r^2 + 100$
25. $(r + 4)(3r - 1) - (r + 4)(r + 2)$
26. $9 - a^2 + 2ab - b^2$
27. $625 - x^4$
28. $2m^2 - mn - 15n^2$

29. $p^3 + 1$
31. $64m^2 - 625$
33. $12z^3 - 6z^2 + 18z$
35. $256b^2 - 400c^2$
37. $512 + 1000z^3$
39. $10r^2 + 23rs - 5s^2$
41. $24p^3q + 52p^2q^2 + 20pq^3$
43. $48k^4 - 243$
45. $m^3 + m^2 - n^3 - n^2$
47. $x^2 - 4m^2 - 4mn - n^2$
49. $18p^5 - 24p^3 + 12p^6$
51. $2x^2 - 2x - 40$
53. $(2m + n)^2 - (2m - n)^2$
55. $50p^2 - 162$
57. $12m^2rx + 4mnrx + 40n^2rx$
59. $21a^2 - 5ab - 4b^2$
61. $x^2 - y^2 - 4$
63. $(p + 8q)^2 - 10(p + 8q) + 25$
65. $21m^4 - 32m^2 - 5$
67. $(r + 2t)^3 + (r - 3t)^3$
69. $x^5 + 3x^4 - x - 3$
71. $m^2 - 4m + 4 - n^2 + 6n - 9$
30. $(t + 8)(4t + 1) - (t + 8)(3t - 11)$
32. $14z^2 - 3zk - 2k^2$
34. $225k^2 - 36r^2$
36. $z^2 - zp - 20p^2$
38. $64m^2 - 25n^2$
40. $12k^2 - 17kq - 5q^2$
42. $32x^2 + 16x^3 - 24x^5$
44. $14x^2 - 25xq - 25q^2$
46. $64x^3 + y^3 - 16x^2 + y^2$
48. $4r^2 - s^2 - 2st - t^2$
50. $k^2 - 6k + 16$
52. $27x^3 - 3y^3$
54. $(3k + 5)^2 - 4(3k + 5) + 4$
56. $y^2 + 3y - 10$
58. $18p^2 + 53pr - 35r^2$
60. $x^2 - 2xy + y^2 - 4$
62. $(5r + 2s)^2 - 6(5r + 2s) + 9$
64. $z^4 - 9z^2 + 20$
66. $(x - y)^3 - (27 - y)^3$
68. $16x^3 + 32x^2 - 9x - 18$
70. $1 - x^{16}$
72. $x^2 + 4 + x^2y + 4y$

5.5

Solving Quadratic Equations Using the Zero-Factor Property

OBJECTIVES

- 1 Use the zero-factor property.
- 2 Solve applied problems that require the zero-factor property.
- 3 Solve a formula for a specified variable, where factoring is necessary.

We have solved linear, or first-degree, equations. Solving higher-degree polynomial equations requires other methods, one of which involves factoring.

OBJECTIVE 1 Use the zero-factor property.

The **zero-factor property** is a special property of the number 0.

Zero-Factor Property

If two numbers have a product of 0, then at least one of the numbers must be 0.

If $ab = 0$, then either $a = 0$ or $b = 0$ (or both).

To prove the zero-factor property, we first assume that $a \neq 0$. (If $a = 0$, then the property is proved already.) If $a \neq 0$, then $\frac{1}{a}$ exists.

$$ab = 0$$

$$\frac{1}{a} \cdot ab = \frac{1}{a} \cdot 0 \quad \text{Multiply each side by } \frac{1}{a}.$$

$$b = 0 \quad \text{Multiply.}$$

Thus, if $a \neq 0$, then $b = 0$, and the property is proved.

VOCABULARY

- quadratic equation
- double solution

CAUTION If $ab = 0$, then $a = 0$ or $b = 0$. However, if $ab = 6$, for example, it is not necessarily true that $a = 6$ or $b = 6$. In fact, it is very likely that *neither* $a = 6$ *nor* $b = 6$. **The zero-factor property works only for a product equal to 0.**

NOW TRY EXERCISE 1

Solve.

$$(x + 5)(4x - 7) = 0$$

EXAMPLE 1 Using the Zero-Factor Property to Solve an Equation

Solve $(x + 6)(2x - 3) = 0$.

The product of $x + 6$ and $2x - 3$ is 0. The zero-factor property applies.

$$x + 6 = 0 \quad \text{or} \quad 2x - 3 = 0 \quad \text{Zero-factor property}$$

$$x = -6 \quad \text{or} \quad 2x = 3 \quad \text{Solve each of these equations.}$$

$$x = \frac{3}{2}$$

CHECK $(x + 6)(2x - 3) = 0$

$$(-6 + 6)[2(-6) - 3] \stackrel{?}{=} 0$$

Let $x = -6$.

$$0(-15) \stackrel{?}{=} 0$$

$$0 = 0 \quad \checkmark \quad \text{True}$$

$$(x + 6)(2x - 3) = 0$$

$$\left(\frac{3}{2} + 6\right)\left(2 \cdot \frac{3}{2} - 3\right) \stackrel{?}{=} 0$$

Let $x = \frac{3}{2}$.

$$\frac{15}{2}(0) \stackrel{?}{=} 0$$

$$0 = 0 \quad \checkmark \quad \text{True}$$

Both values check, so the solution set is $\left\{-6, \frac{3}{2}\right\}$.

NOW TRY 

Because the product $(x + 6)(2x - 3)$ equals $2x^2 + 9x - 18$, the equation of **Example 1** has a term with a squared variable and is an example of a *quadratic equation*. **A quadratic equation has degree 2.**

Quadratic Equation

A **quadratic equation** (in x here) is an equation that can be written in the form

$$ax^2 + bx + c = 0,$$

where a , b , and c are real numbers, and $a \neq 0$. The given form is called **standard form**.

Examples: $3x^2 - 7x + 2 = 0$, $x^2 - 4 = 0$, $t^2 + 2t = 0$ Quadratic equations

NOW TRY ANSWER

1. $\left\{-5, \frac{7}{4}\right\}$

Solving a Quadratic Equation Using the Zero-Factor Property

- Step 1** Write the equation in standard form—that is, with all terms on one side of the equality symbol in descending powers of the variable and with 0 on the other side.
- Step 2** Factor completely.
- Step 3** Apply the zero-factor property. Set each factor with a variable equal to 0.
- Step 4** Solve the resulting equations.
- Step 5** Check each value in the *original* equation. Write the solution set.


NOW TRY EXERCISE 2
Solve $7x = 3 - 6x^2$.**EXAMPLE 2** Solving a Quadratic EquationSolve $2x^2 + 3x = 2$.

$$2x^2 + 3x = 2$$

Step 1 $2x^2 + 3x - 2 = 0$ Standard form

Step 2 $(2x - 1)(x + 2) = 0$ Factor.

Step 3 $2x - 1 = 0$ or $x + 2 = 0$ Zero-factor property

Step 4 $2x = 1$ or $x = -2$ Solve each equation.

$$x = \frac{1}{2}$$

Step 5 Check each value in the original equation.

CHECK $2x^2 + 3x = 2$

$$2\left(\frac{1}{2}\right)^2 + 3\left(\frac{1}{2}\right) \stackrel{?}{=} 2 \quad \text{Let } x = \frac{1}{2}.$$

$$2\left(\frac{1}{4}\right) + \frac{3}{2} \stackrel{?}{=} 2$$

$$\frac{1}{2} + \frac{3}{2} \stackrel{?}{=} 2$$

$$2 = 2 \quad \checkmark \quad \text{True}$$

$$2x^2 + 3x = 2$$

$$2(-2)^2 + 3(-2) \stackrel{?}{=} 2 \quad \text{Let } x = -2.$$

$$2(4) - 6 \stackrel{?}{=} 2$$

$$8 - 6 \stackrel{?}{=} 2$$

$$2 = 2 \quad \checkmark \quad \text{True}$$

We give solutions from least to greatest.

Because both values check, the solution set is $\left\{-2, \frac{1}{2}\right\}$.

NOW TRY 


NOW TRY EXERCISE 3
Solve $16x^2 = -40x - 25$.**EXAMPLE 3** Solving a Quadratic Equation (Double Solution)Solve $4x^2 = 4x - 1$.

$$4x^2 = 4x - 1$$

$$4x^2 - 4x + 1 = 0 \quad \text{Standard form}$$

$$(2x - 1)^2 = 0 \quad \text{Factor.}$$

$$2x - 1 = 0 \quad \text{Zero-factor property}$$

$$x = \frac{1}{2} \quad \text{Add 1. Divide by 2.}$$

We could factor as $(2x - 1)(2x - 1)$. The same solution results.

NOW TRY ANSWERS

2. $\left\{-\frac{3}{2}, \frac{1}{3}\right\}$

3. $\left\{-\frac{5}{4}\right\}$

There is only one *distinct* solution, called a **double solution**, because the trinomial $4x^2 - 4x + 1$ is a perfect square. The solution set is $\left\{\frac{1}{2}\right\}$.

NOW TRY 


**NOW TRY
EXERCISE 4**
Solve $3x^2 + 12x = 0$.
EXAMPLE 4 Solving a Quadratic Equation (Missing Constant Term)
Solve $4x^2 - 20x = 0$.

This quadratic equation has a missing constant term. Comparing it with the standard form $ax^2 + bx + c = 0$ shows that $c = 0$. The zero-factor property can still be used.

$$4x^2 - 20x = 0$$

$$4x(x - 5) = 0 \quad \text{Factor.}$$

Set each variable factor equal to 0.

$$4x = 0 \quad \text{or} \quad x - 5 = 0 \quad \text{Zero-factor property}$$

$$x = 0 \quad \text{or} \quad x = 5 \quad \text{Solve each equation.}$$

CHECK $4x^2 - 20x = 0$


$4(0)^2 - 20(0) \stackrel{?}{=} 0 \quad \text{Let } x = 0.$

$0 - 0 = 0 \quad \checkmark \quad \text{True}$

$4x^2 - 20x = 0$

$4(5)^2 - 20(5) \stackrel{?}{=} 0 \quad \text{Let } x = 5.$

$100 - 100 = 0 \quad \checkmark \quad \text{True}$

The solution set is $\{0, 5\}$. 

NOW TRY 

**NOW TRY
EXERCISE 5**
Solve $4x^2 - 100 = 0$.
EXAMPLE 5 Solving a Quadratic Equation (Missing Linear Term)
Solve $3x^2 - 108 = 0$.

$$3x^2 - 108 = 0$$

$$3(x^2 - 36) = 0 \quad \text{Factor out 3.}$$

The factor 3 does *not* lead to a solution.

$$3(x + 6)(x - 6) = 0 \quad \text{Factor } x^2 - 36.$$

$$x + 6 = 0 \quad \text{or} \quad x - 6 = 0 \quad \text{Zero-factor property}$$

$$x = -6 \quad \text{or} \quad x = 6 \quad \text{Solve each equation.}$$

Check that the solution set is $\{-6, 6\}$.

NOW TRY 

CAUTION The factor $4x$ in **Example 4** is a **variable factor** and leads to the solution 0. The factor 3 in **Example 5** is **not a variable factor**, so it does *not* lead to a solution of the equation.

EXAMPLE 6 Solving an Equation That Requires Rewriting
Solve $(2x + 1)(x + 1) = 2(1 - x) + 6$.

$$(2x + 1)(x + 1) = 2(1 - x) + 6$$

$$2x^2 + 3x + 1 = 2 - 2x + 6 \quad \text{Multiply on each side.}$$

$$2x^2 + 3x + 1 = 8 - 2x \quad \text{Add on the right.}$$

Write in standard form.

$$2x^2 + 5x - 7 = 0 \quad \text{Add } 2x. \text{ Subtract } 8.$$

$$(2x + 7)(x - 1) = 0 \quad \text{Factor.}$$

$$2x + 7 = 0 \quad \text{or} \quad x - 1 = 0 \quad \text{Zero-factor property}$$

$$x = -\frac{7}{2} \quad \text{or} \quad x = 1 \quad \text{Solve each equation.}$$

NOW TRY ANSWERS
4. $\{0, -4\}$ 5. $\{-5, 5\}$

NOW TRY
EXERCISE 6

Solve.

$$\begin{aligned}(x+3)(2x-1) \\ = 4(x+4) - 4\end{aligned}$$

CHECK

$$\begin{aligned}(2x+1)(x+1) &= 2(1-x) + 6 \\ \left[2\left(-\frac{7}{2}\right) + 1\right]\left(-\frac{7}{2} + 1\right) &\stackrel{?}{=} 2\left[1 - \left(-\frac{7}{2}\right)\right] + 6 && \text{Let } x = -\frac{7}{2}. \\ (-7+1)\left(-\frac{5}{2}\right) &\stackrel{?}{=} 2\left(\frac{9}{2}\right) + 6 && \text{Simplify; } 1 = \frac{2}{2}. \\ (-6)\left(-\frac{5}{2}\right) &\stackrel{?}{=} 9 + 6 \\ 15 &= 15 \quad \checkmark && \text{True}\end{aligned}$$

Check that 1 is also a solution. The solution set is $\left\{-\frac{7}{2}, 1\right\}$.

NOW TRY

The zero-factor property can be extended to solve certain polynomial equations of degree 3 or greater, as shown in the next example.

NOW TRY
EXERCISE 7

Solve $12x = 2x^3 + 5x^2$.**EXAMPLE 7** Solving an Equation of Degree 3Solve $-x^3 + x^2 = -6x$.

$$\begin{aligned}-x^3 + x^2 &= -6x \\ -x^3 + x^2 + 6x &= 0 && \text{Add } 6x \text{ to each side.} \\ x^3 - x^2 - 6x &= 0 && \text{Multiply each side by } -1. \\ x(x^2 - x - 6) &= 0 && \text{Factor out } x. \\ x(x-3)(x+2) &= 0 && \text{Factor the trinomial.} \\ x = 0 \quad \text{or} \quad x-3 = 0 \quad \text{or} \quad x+2 = 0 &&& \text{Extend the zero-factor property} \\ &&& \text{to the three variable factors.} \\ x = 3 \quad \text{or} \quad x = -2\end{aligned}$$

Remember to set x equal to 0.

Check that the solution set is $\{-2, 0, 3\}$.

NOW TRY**OBJECTIVE 2** Solve applied problems that require the zero-factor property.

An application may lead to a quadratic equation.

EXAMPLE 8 Using a Quadratic Equation in an Application


A piece of sheet metal is in the shape of a parallelogram. The longer sides of the parallelogram are each 8 m longer than the distance between them. The area of the piece is 48 m^2 . Find the length of the longer sides and the distance between them.

Step 1 Read the problem again. There will be two answers.

Step 2 Assign a variable.

Let x = the distance between the two longer sides

and $x + 8$ = the length of each of the longer sides. (See **FIGURE 1**.)

**FIGURE 1****NOW TRY ANSWERS**

6. $\left\{-3, \frac{5}{2}\right\}$

7. $\left\{-4, 0, \frac{3}{2}\right\}$

NOW TRY EXERCISE 8

The height of a triangle is 1 ft less than twice the length of the base. The area is 14 ft^2 . What are the measures of the base and the height?

Step 3 Write an equation. The area of a parallelogram is given by $\mathcal{A} = bh$, where b is the length of the longer side and h is the distance between the longer sides. Here, $b = x + 8$ and $h = x$.

$$\mathcal{A} = bh \quad \text{Formula for area of a parallelogram}$$

$$48 = (x + 8)x \quad \text{Let } \mathcal{A} = 48, b = x + 8, h = x.$$

Step 4 Solve. $48 = x^2 + 8x$ Distributive property

$$x^2 + 8x - 48 = 0 \quad \text{Standard form}$$

$$(x + 12)(x - 4) = 0 \quad \text{Factor.}$$

$$x + 12 = 0 \quad \text{or} \quad x - 4 = 0 \quad \text{Zero-factor property}$$

$$x = -12 \quad \text{or} \quad x = 4 \quad \text{Solve each equation.}$$

Step 5 State the answer. *The distance between the longer sides cannot be negative, so reject -12 as an answer.* The only possible answer is 4 , so the required distance is 4 m. The length of the longer sides is $4 + 8 = 12$ m.

Step 6 Check. The length of the longer sides is 8 m more than the distance between them, and the area is

$$4 \cdot 12 = 48 \text{ m}^2, \quad \text{as required.}$$

NOW TRY 

CAUTION When an application leads to a quadratic equation, a solution of the equation may not satisfy the physical requirements of the problem, as in **Example 8**. Reject such solutions as answers.

A function defined by a quadratic polynomial is a **quadratic function**. The next example uses such a function.

NOW TRY EXERCISE 9

Refer to **Example 9**. After how many seconds will the rocket be 192 ft above the ground?

EXAMPLE 9 Using a Quadratic Function in an Application

If a small rocket is launched vertically upward from ground level with an initial velocity of 128 ft per sec, then its height in feet after t seconds (if air resistance is neglected) can be modeled by the function

$$h(t) = -16t^2 + 128t.$$

After how many seconds will the rocket be 220 ft above the ground?

We let $h(t) = 220$ and solve for t .

$$220 = -16t^2 + 128t \quad \text{Let } h(t) = 220.$$

$$16t^2 - 128t + 220 = 0 \quad \text{Standard form}$$

$$4t^2 - 32t + 55 = 0 \quad \text{Divide by 4.}$$

$$(2t - 5)(2t - 11) = 0 \quad \text{Factor.}$$

$$2t - 5 = 0 \quad \text{or} \quad 2t - 11 = 0 \quad \text{Zero-factor property}$$

$$t = 2.5 \quad \text{or} \quad t = 5.5 \quad \text{Solve each equation.}$$


The rocket will reach a height of 220 ft twice: on its way up at 2.5 sec and again on its way down at 5.5 sec.



NOW TRY 

NOW TRY ANSWERS

8. base: 4 ft; height: 7 ft
9. 2 sec and 6 sec



Rectangular solid
 $S = 2HW + 2LW + 2LH$

FIGURE 2

NOW TRY EXERCISE 10

Solve the formula for H .

$$S = 2HW + 2LW + 2LH$$

OBJECTIVE 3 Solve a formula for a specified variable, where factoring is necessary.

A rectangular solid has the shape of a box, but is solid. See FIGURE 2. The surface area of any solid three-dimensional figure is the total area of its surface. For a rectangular solid, the surface area S is

$$S = 2HW + 2LW + 2LH.$$

H , W , and L represent height, width, and length.

EXAMPLE 10 Solving for a Specified Variable

Solve the formula $S = 2HW + 2LW + 2LH$ for L .

To solve for L , treat L as the only variable and treat H and W as constants.

$$S = 2HW + 2LW + 2LH$$

We must isolate the L -terms.

$$S - 2HW = 2LW + 2LH$$

Subtract $2HW$.

This is a key step.

$$S - 2HW = L(2W + 2H)$$

Factor out L .

$$\frac{S - 2HW}{2W + 2H} = L, \text{ or } L = \frac{S - 2HW}{2W + 2H}$$

Divide by $2W + 2H$. **NOW TRY**

CAUTION In Example 10, we must write the expression so that the specified variable is a factor. Then we can divide by its coefficient in the final step.

NOTE Earlier we saw that the graph of $f(x) = x^2$ is a parabola. In general, the graph of $f(x) = ax^2 + bx + c$ (where $a \neq 0$)

is a parabola, and the x -intercepts of its graph give the real number solutions of the equation $ax^2 + bx + c = 0$. A graphing calculator can locate these x -intercepts (called **zeros** of the function).

Consider $f(x) = 2x^2 + 3x - 2$. Notice that this quadratic expression was found on the left side of the equation in Step 1 in **Example 2** earlier in this section, where the equation was written in standard form. The x -intercepts (zeros) given with the graphs in **FIGURE 3** are the same as the solutions found in **Example 2**.




FIGURE 3

NOW TRY ANSWER

10. $H = \frac{S - 2LW}{2W + 2L}$

5.5 Exercises **FOR EXTRA HELP** **MyLab Math**

Video solutions for select problems available in MyLab Math

Concept Check Choose the letter(s) of the correct response.

- Which of the following are quadratic equations?

A. $2x + 3 = 6$	B. $x^2 - 3x + 2 = 0$	C. $5t^2 + 3t = 2$
D. $2x^3 + x^2 - 6x = 0$	E. $x^2 + 16 = 0$	F. $3x^2 - x = 0$

STUDY SKILLS REMINDER

Time management can be a challenge for students.

Review Study Skill 6,
Managing Your Time.

2. Which of the following can be solved using the zero-factor property?

A. $25x^2 + 20x + 4 = 0$

B. $3x^2 = 15x$

C. $2x^2 - 128 = 0$

D. $x(4x - 15) - 30 = 0$

3. **Concept Check** A student solved the following equation incorrectly as shown.

$$\begin{aligned}(x - 3)(x + 2) &= 6 \\ x - 3 &= 6 \quad \text{or} \quad x + 2 = 6 \\ x &= 9 \quad \text{or} \quad x = 4\end{aligned}$$

The student gave the solution set as $\{4, 9\}$.

WHAT WENT WRONG? Give the correct solution set.

5. **Concept Check** A student solved the following equation incorrectly as shown.

$$\begin{aligned}3x^2 + 12x &= 0 \\ 3x(x + 4) &= 0 && \text{Factor.} \\ x + 4 &= 0 && \text{Divide by } 3x. \\ x &= -4\end{aligned}$$

The student gave $\{-4\}$ as the solution set.

WHAT WENT WRONG? Give the correct solution set.

4. **Concept Check** A student solved the following equation incorrectly as shown.

$$\begin{aligned}7(x + 4)(x - 3) &= 0 \\ x = 7 \quad \text{or} \quad x + 4 = 0 \quad \text{or} \quad x - 3 = 0 \\ x = 7 \quad \text{or} \quad x = -4 \quad \text{or} \quad x = 3\end{aligned}$$

The student gave the solution set as $\{-4, 3, 7\}$.

WHAT WENT WRONG? Give the correct solution set.

6. **Concept Check** A student solved the following formula incorrectly for L as shown.

$$\begin{aligned}S &= 2HW + 2LW + 2LH \\ S - 2LW - 2HW &= 2LH \\ \frac{S - 2LW - 2HW}{2H} &= L\end{aligned}$$

WHAT WENT WRONG?

Solve each equation. See Examples 1–6.

7. $(x + 10)(x - 5) = 0$

8. $(x + 7)(x + 3) = 0$

9. $(3k + 8)(2k - 5) = 0$

10. $(3q - 4)(2q + 5) = 0$

11. $x^2 - 3x - 10 = 0$

12. $x^2 + x - 12 = 0$

13. $x^2 + 9x + 18 = 0$

14. $x^2 - 18x + 80 = 0$

15. $2x^2 = 7x + 4$

16. $2x^2 = 3 - x$

17. $15x^2 - 7x = 4$

18. $12x^2 + 4x = 5$

19. $4p^2 + 16p = 0$

20. $2t^2 + 8t = 0$

21. $6x^2 - 36x = 0$

22. $3x^2 - 27x = 0$

23. $4p^2 - 16 = 0$

24. $9z^2 - 81 = 0$

25. $-3x^2 + 27 = 0$

26. $-2x^2 + 8 = 0$

27. $-x^2 = 9 - 6x$

28. $-x^2 - 8x = 16$

29. $9x^2 + 24x + 16 = 0$

30. $4x^2 - 20x + 25 = 0$

31. $(x - 3)(x + 5) = -7$

32. $(x + 8)(x - 2) = -21$

33. $(2x + 1)(x - 3) = 6x + 3$

34. $(3x + 2)(x - 3) = 7x - 1$

35. $2x^2 - 12 - 4x = x^2 - 3x$

36. $3x^2 + 9x + 30 = 2x^2 - 2x$

37. $(5x + 1)(x + 3) = -2(5x + 1)$

38. $(3x + 1)(x - 3) = 2 + 3(x + 5)$

39. $(x + 3)(x - 6) = (2x + 2)(x - 6)$

40. $(2x + 1)(x + 5) = (x + 11)(x + 3)$

Solve each equation. See Example 7.

41. $2x^3 - 9x^2 - 5x = 0$

42. $6x^3 - 13x^2 - 5x = 0$

43. $x^3 - 2x^2 = 3x$

44. $x^3 - 6x^2 = -8x$

45. $9x^3 = 16x$

46. $25x^3 = 64x$


47. $2x^3 + 5x^2 - 2x - 5 = 0$ 48. $2x^3 + x^2 - 98x - 49 = 0$
 49. $x^3 - 6x^2 - 9x + 54 = 0$ 50. $x^3 - 3x^2 - 4x + 12 = 0$

Extending Skills Solve each equation. (Hint: In Exercises 51–54, use the substitution of variable method.)


51. $2(x - 1)^2 - 7(x - 1) - 15 = 0$ 52. $4(2x + 3)^2 - (2x + 3) - 3 = 0$
 53. $5(3x - 1)^2 + 3 = -16(3x - 1)$ 54. $2(x + 3)^2 = 5(x + 3) - 2$
 55. $(2x - 3)^2 = 16x^2$ 56. $9x^2 = (5x + 2)^2$

Solve each problem. See Examples 8 and 9.


57. A garden has an area of 320 ft^2 . Its length is 4 ft more than its width. What are the dimensions of the garden?



58. A square mirror has sides measuring 2 ft less than the sides of a square painting. If the difference between their areas is 32 ft^2 , find the lengths of the sides of the mirror and the painting.




59. The base of a parallelogram is 7 ft more than the height. If the area of the parallelogram is 60 ft^2 , what are the measures of the base and the height?




60. A sign has the shape of a triangle. The length of the base is 3 m less than the height. What are the measures of the base and the height if the area is 44 m^2 ?



61. A farmer has 300 ft of fencing and wants to enclose a rectangular area of 5000 ft^2 . What dimensions should she use?
 62. A rectangular landfill has an area of $30,000 \text{ ft}^2$. Its length is 200 ft more than its width. What are the dimensions of the landfill?
 63. Find two consecutive integers such that the sum of their squares is 61.
 64. Find two consecutive integers such that their product is 72.
 65. A box with no top is to be constructed from a piece of cardboard whose length measures 6 in. more than its width. The box is to be formed by cutting squares that measure 2 in. on each side from the four corners and then folding up the sides. If the volume of the box will be 110 in.^3 , what are the dimensions of the piece of cardboard?



66. The surface area of the box with open top shown in the figure is 161 in.^2 . Find the dimensions of the base. (*Hint:* The surface area of the box is modeled by the function $S(x) = x^2 + 16x$.)



67. If an object is projected upward with an initial velocity of 64 ft per sec from a height of 80 ft, then its height in feet t seconds after it is projected is modeled by the function

$$f(t) = -16t^2 + 64t + 80.$$

How long after it is projected will it hit the ground? (*Hint:* When it hits the ground, its height is 0 ft.)

68. Refer to **Example 9**. After how many seconds will the rocket be

(a) 240 ft above the ground? (b) 112 ft above the ground?

69. If a rock is dropped from a building 576 ft high, then its distance in feet from the ground t seconds later is modeled by the function

$$f(t) = -16t^2 + 576.$$

How long after it is dropped will it hit the ground?

70. If a baseball is dropped from a helicopter 625 ft above the ground, then its distance in feet from the ground t seconds later is modeled by the function

$$f(t) = -16t^2 + 625.$$

How long after it is dropped will it hit the ground?

Solve each equation for the specified variable. *See Example 10.*

71. $k = dF - DF$ for F

72. $Mv = mv - Vm$ for m

73. $A = P + Prt$ for P

74. $P = kR + R - u$ for R

75. $2k + ar = r - 3y$ for r

76. $4s + 7p = tp - 7$ for p

77. $w = \frac{3y - x}{y}$ for y

78. $c = \frac{-2t + 4}{t}$ for t

Chapter 5

Summary

STUDY SKILLS REMINDER

Be prepared for your math test over this chapter. **Review Study Skill 7, Reviewing a Chapter, and Study Skill 8, Taking Math Tests.**

Key Terms

5.1

factoring
greatest common factor
(GCF)

5.2

prime polynomial

5.3

difference of squares
perfect square trinomial
difference of cubes
sum of cubes

5.5

quadratic equation
double solution

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- Factoring** is
 - a method of multiplying polynomials
 - the process of writing a polynomial as a product of prime factors
 - the answer in a multiplication problem
 - a way to add the terms of a polynomial.
- A **difference of squares** is a binomial
 - that can be factored as the difference of two cubes
 - that cannot be factored
- that is squared
 - that can be factored as the product of the sum and difference of two terms.
- A **perfect square trinomial** is a trinomial
 - that can be factored as the square of a binomial
 - that cannot be factored
 - that is multiplied by a binomial
 - where all terms are perfect squares.
- A **quadratic equation** is a polynomial equation of
 - degree 1
 - degree 2
 - degree 3
 - degree 4.
- The **zero-factor property** is used to
 - factor a perfect square trinomial
 - factor by grouping
 - solve a polynomial equation of degree 2 or more
 - solve a linear equation.

ANSWERS

1. B; *Example:* $x^2 - 5x - 14$ factors as $(x - 7)(x + 2)$. 2. D; *Example:* $b^2 - 49$ is the difference of the squares b^2 and 7^2 . It can be factored as $(b + 7)(b - 7)$. 3. A; *Example:* $a^2 + 2a + 1$ is a perfect square trinomial. Its factored form is $(a + 1)^2$. 4. B; *Examples:* $x^2 - 3x + 2 = 0$, $x^2 - 9 = 0$, and $2x^2 = 6x + 8$ 5. C; *Example:* Use the zero-factor property to write $(x + 4)(x - 2) = 0$ as $x + 4 = 0$ or $x - 2 = 0$, and then solve each linear equation to find the solution set $\{-4, 2\}$.

Quick Review

CONCEPTS	EXAMPLES										
<p>5.1 Greatest Common Factors and Factoring by Grouping</p> <p>Factoring Out the Greatest Common Factor Use the distributive property to write the given polynomial as a product of two factors, one of which is the greatest common factor of the terms of the polynomial.</p> <p>Factoring by Grouping</p> <p><i>Step 1</i> Group the terms so that each group has a common factor.</p> <p><i>Step 2</i> Factor out the common factor in each group.</p> <p><i>Step 3</i> If each group now has a common factor, factor it out. If not, try a different grouping.</p> <p><i>Check the factored form by multiplying.</i></p>	<p>Factor $4x^2y - 50xy^2$.</p> $4x^2y - 50xy^2$ $= 2xy(2x - 25y) \quad \text{The greatest common factor is } 2xy.$ <p>Factor by grouping.</p> $5a - 5b - ax + bx$ $= (5a - 5b) + (-ax + bx) \quad \text{Group the terms.}$ $= 5(a - b) - x(a - b) \quad \text{Factor out 5 and } -x.$ $= (a - b)(5 - x) \quad \text{Factor out } a - b.$										
<p>5.2 Factoring Trinomials</p> <p>Factoring $x^2 + bx + c$</p> <p><i>Step 1</i> Find pairs of integers whose product is c.</p> <p><i>Step 2</i> Choose the pair whose sum is b.</p> <p><i>Step 3</i> Factor the trinomial as the product of two binomials using the pair of integers identified in Step 2.</p> <p><i>Check the factored form by multiplying.</i></p>	<p>Factor $x^2 - 6x - 27$.</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="border-right: 1px solid black;">Pairs of integers with product -27</th> <th>Sum of each pair</th> </tr> </thead> <tbody> <tr> <td style="border-right: 1px solid black; text-align: center;">$27(-1)$</td> <td style="text-align: center;">26</td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">$-27(1)$</td> <td style="text-align: center;">-26</td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">$9(-3)$</td> <td style="text-align: center;">6</td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">$-9(3)$</td> <td style="text-align: center;">$-6 \leftarrow$ Coefficient b</td> </tr> </tbody> </table> <p>The integers -9 and 3 have the necessary product and sum.</p> $x^2 - 6x - 27 \text{ factors as } (x - 9)(x + 3).$	Pairs of integers with product -27	Sum of each pair	$27(-1)$	26	$-27(1)$	-26	$9(-3)$	6	$-9(3)$	$-6 \leftarrow$ Coefficient b
Pairs of integers with product -27	Sum of each pair										
$27(-1)$	26										
$-27(1)$	-26										
$9(-3)$	6										
$-9(3)$	$-6 \leftarrow$ Coefficient b										

CONCEPTS

Factoring $ax^2 + bx + c$

Use one of the following methods.

- Identify the values of a , b , and c . Find two integers having product ac and sum b . Use these integers to rewrite the middle term, and factor by grouping.

- Choose factors of the first term and factors of the last term. Write them in a pair of parentheses of this form.

$$\left(\quad \right) \left(\quad \right)$$

Use the FOIL method and try various combinations of the factors until the correct middle term is found.

Check the factored form by multiplying.

EXAMPLES

Factor $15x^2 + 14x - 8$.

$a = 15$, $b = 14$, and $c = -8$, so $ac = 15(-8) = -120$.

We must find two integers with product -120 and sum 14 .

$$20(-6) = -120 \quad \text{and} \quad 20 + (-6) = 14$$

The necessary integers are 20 and -6 .

$$\begin{aligned} 15x^2 + 14x - 8 &= 15x^2 + 20x - 6x - 8 && 20x + (-6x) = 14x \\ &= (15x^2 + 20x) + (-6x - 8) && \text{Group the terms.} \\ &= 5x(3x + 4) - 2(3x + 4) && \text{Factor each group.} \\ &= (3x + 4)(5x - 2) && \text{Factor out } 3x + 4. \end{aligned}$$

Factor $15x^2 + 14x - 8$.

The positive factors of 15 are 5 and 3 , and 15 and 1 .

The factors of -8 are

$$-4 \text{ and } 2, \quad 4 \text{ and } -2, \quad -1 \text{ and } 8, \quad \text{and } 1 \text{ and } -8.$$

Various combinations of these factors lead to the following.

$$\begin{aligned} 15x^2 + 14x - 8 &= (5x - 2)(3x + 4) && 20x - 6x = 14x, \\ & && \text{the correct middle term.} \end{aligned}$$

5.3 Special Factoring**Difference of Squares**

$$x^2 - y^2 = (x + y)(x - y)$$

Perfect Square Trinomials

$$x^2 + 2xy + y^2 = (x + y)^2$$

$$x^2 - 2xy + y^2 = (x - y)^2$$

Difference of Cubes

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2)$$

Sum of Cubes

$$x^3 + y^3 = (x + y)(x^2 - xy + y^2)$$

$$\begin{aligned} \text{Factor.} \quad 4m^2 - 25n^2 &= (2m)^2 - (5n)^2 \\ &= (2m + 5n)(2m - 5n) \end{aligned}$$

$$\begin{array}{l|l} \text{Factor.} \quad 9y^2 + 6y + 1 & 16p^2 - 56p + 49 \\ = (3y + 1)^2 & = (4p - 7)^2 \end{array}$$

$$\begin{aligned} \text{Factor.} \quad 8 - 27a^3 &= (2 - 3a)(4 + 6a + 9a^2) \end{aligned}$$

$$\begin{aligned} 64z^3 + 1 &= (4z + 1)(16z^2 - 4z + 1) \end{aligned}$$

5.4 A General Approach to Factoring

See this section for guidelines and examples.

CONCEPTS

5.5 Solving Quadratic Equations Using the Zero-Factor Property**Zero-Factor Property**

If two numbers have a product of 0, then at least one of the numbers must be 0.

If $ab = 0$, then $a = 0$ or $b = 0$ (or both).

Solving a Quadratic Equation Using the Zero-Factor Property

Step 1 Write the equation in standard form.

Step 2 Factor completely.

Step 3 Apply the zero-factor property.

Step 4 Solve the resulting equations.

Step 5 Check each value in the original equation. Write the solution set.

EXAMPLES

Solve. $2x^2 + 5x = 3$

$$2x^2 + 5x - 3 = 0 \quad \text{Standard form}$$

$$(x + 3)(2x - 1) = 0 \quad \text{Factor.}$$

$$x + 3 = 0 \quad \text{or} \quad 2x - 1 = 0 \quad \text{Zero-factor property}$$

$$x = -3 \quad \text{or} \quad x = \frac{1}{2} \quad \text{Solve each equation.}$$

CHECK $2x^2 + 5x = 3$

$$2(-3)^2 + 5(-3) \stackrel{?}{=} 3 \quad \text{Let } x = -3.$$

$$2(9) - 15 \stackrel{?}{=} 3$$

$$3 = 3 \quad \checkmark \quad \text{True}$$

A check of $\frac{1}{2}$ is done similarly.

Solution set: $\left\{-3, \frac{1}{2}\right\}$

Chapter 5

Review Exercises

5.1 Factor out the greatest common factor.

1. $12p^2 - 6p$

2. $21x^2 + 35x$

3. $12q^2b + 8qb^2 - 20q^3b^2$

4. $6r^3t - 30r^2t^2 + 18rt^3$

5. $(x + 3)(4x - 1) - (x + 3)(3x + 2)$

6. $(z + 1)(z - 4) + (z + 1)(2z + 3)$

Factor by grouping.

7. $4m + nq + mn + 4q$

8. $x^2 + 5y + 5x + xy$

9. $2m + 6 - am - 3a$

10. $x^2 + 3x - 3y - xy$

5.2 Factor completely.

11. $3p^2 - p - 4$

12. $6k^2 + 11k - 10$

13. $12r^2 - 5r - 3$

14. $10m^2 + 37m + 30$

15. $10k^2 - 11kh + 3h^2$

16. $9x^2 + 4xy - 2y^2$

17. $24x - 2x^2 - 2x^3$

18. $6b^3 - 9b^2 - 15b$

19. $y^4 + 2y^2 - 8$

20. $2k^4 - 5k^2 - 3$

21. $p^2(p + 2)^2 + p(p + 2)^2 - 6(p + 2)^2$

22. $3(r + 5)^2 - 11(r + 5) - 4$

23. $24x^8 - 126x^6 + 81x^4$

24. $30y^9 - 155y^6 - 55y^3$

5.3 Factor completely.

25. $16x^2 - 25$

26. $9t^2 - 49$

27. $36m^2 - 25n^2$

28. $x^2 + 14x + 49$

29. $9k^2 - 12k + 4$

30. $r^3 + 27$


31. $125x^3 - 1$ 32. $m^6 - 1$ 33. $x^8 - 1$
 34. $x^2 + 6x + 9 - 25y^2$ 35. $(a + b)^3 - (a - b)^3$ 36. $x^5 - x^3 - 8x^2 + 8$

5.5 Solve each equation.


37. $x^2 - 8x + 16 = 0$ 38. $(5x + 2)(x + 1) = 0$ 39. $x^2 - 5x + 6 = 0$
 40. $x^2 + 2x = 8$ 41. $6x^2 = 5x + 50$ 42. $6x^2 + 7x = 3$
 43. $8x^2 + 14x + 3 = 0$ 44. $-4x^2 + 36 = 0$
 45. $6x^2 + 9x = 0$ 46. $(2x + 1)(x - 2) = -3$
 47. $(x - 1)(x + 3) = (2x - 1)(x - 1)$ 48. $2x^3 - x^2 - 28x = 0$
 49. $-x^3 - 3x^2 + 4x + 12 = 0$ 50. $(x + 2)(5x^2 - 9x - 18) = 0$

Solve each problem.

51. A triangular wall brace has the shape of a right triangle. One of the perpendicular sides is 1 ft longer than twice the other. The area enclosed by the triangle is 10.5 ft^2 . Find the shorter of the perpendicular sides.



52. A rectangular parking lot has a length 20 ft more than its width. Its area is 2400 ft^2 . What are the dimensions of the lot?



A rock is projected directly upward from ground level. After t seconds, its height (if air resistance is neglected) is modeled by the function

$$f(t) = -16t^2 + 256t.$$

53. After how many seconds will the rock return to the ground?
 54. After how many seconds will it be 240 ft above the ground?
 55. After how many seconds does the rock reach its maximum height, 1024 ft?
 56. Why does the question in **Exercise 54** have two answers?

Solve each equation for the specified variable.

57. $3s + bk = k - 2t$ for k 58. $z = \frac{3w + 7}{w}$ for w

Chapter 5 Mixed Review Exercises

Factor completely.

- | | | |
|-------------------------|---------------------|--------------------------|
| 1. $x^2 + 5y + 5x + xy$ | 2. $-x^2 + 3x + 10$ | 3. $16x^2 + 144$ |
| 4. $30a + am - am^2$ | 5. $8 - a^3$ | 6. $81k^2 - 16$ |
| 7. $9x^2 + 13xy - 3y^2$ | 8. $15y^3 + 20y^2$ | 9. $25z^2 - 30zm + 9m^2$ |


10. Which equation is *not* in proper form for using the zero-factor property? Why?

- | | |
|---------------------------|------------------------------|
| A. $(x + 2)(x - 6) = 0$ | B. $x(3x - 7) = 0$ |
| C. $3t(t + 8)(t - 9) = 0$ | D. $y(y - 3) + 6(y - 3) = 0$ |


Solve.

- | | | |
|-----------------------------|-----------------------------------|---------------------|
| 11. $5x^2 - 17x - 12 = 0$ | 12. $x^3 - x = 0$ | 13. $3m^2 - 9m = 0$ |
| 14. $2sA - hb = hB$ for h | 15. $S = 2HW + 2LW + 2LH$ for H | |

16. The rectangular floor area of a typical Native American longhouse was about 2750 ft^2 . The length was 85 ft greater than the width. What were the dimensions of the floor?




17. The length of a rectangular picture frame is 2 in. longer than its width. The area enclosed by the frame is 48 in.^2 . What is the width?



18. An arrow is shot upward from a platform 40 ft high with an initial velocity of 224 ft per sec. Its height h in feet after t seconds is modeled by the function

$$h(t) = -16t^2 + 224t + 40.$$

After how many seconds will the arrow be 760 ft above the ground?



STUDY SKILLS REMINDER

We can learn from the mistakes we make. **Review Study Skill 9, Analyzing Your Test Results.**

Chapter 5

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

► View the complete solutions to all Chapter Test exercises in MyLab Math.

Factor.


1. $11z^2 - 44z$
2. $10x^2y^5 - 5x^2y^3 - 25x^5y^3$
3. $3x + by + bx + 3y$
4. $-2x^2 - x + 36$
5. $6x^2 + 11x - 35$
6. $4p^2 + 3pq - q^2$
7. $16a^2 + 40ab + 25b^2$
8. $x^2 + 2x + 1 - 4z^2$
9. $a^3 + 2a^2 - ab^2 - 2b^2$
10. $9k^2 - 121j^2$
11. $y^3 - 216$
12. $6k^4 - k^2 - 35$
13. $27x^6 + 1$
14. $(t^2 + 3)^2 + 4(t^2 + 3) - 5$
15. $16x^2 + 9$
16. $-x^2 + x + 30$
17. $125 - 216y^6$
18. Which one of the following is *not* a factored form of $-x^2 - x + 12$?
 - A. $(3 - x)(x + 4)$
 - B. $-(x - 3)(x + 4)$
 - C. $(-x + 3)(x + 4)$
 - D. $(x - 3)(-x + 4)$

Solve each equation.

19. $3x^2 + 8x = -4$
20. $3x^2 - 5x = 0$
21. $5m(m - 1) = 2(1 - m)$
22. $ar + 2 = 3r - 6t$ for r

Solve each problem.

23. The area of the rectangle shown is 40 in.^2 . Find the length and the width of the rectangle.

The area is 40 in.^2 .

24. A ball is projected upward from ground level. After t seconds, its height in feet is modeled by the function

$$f(t) = -16t^2 + 96t.$$

After how many seconds will it reach a height of 128 ft?

Chapters R–5

Cumulative Review Exercises

Perform the indicated operations.

1. $\frac{5}{6} + \frac{7}{12} - \frac{1}{3}$
2. $2.5(6.5) - 0.15$

Simplify.

3. $-2(m - 3)$
4. $3x^2 - 4x + 4 + 9x - x^2$

Evaluate for $p = -4$, $q = -2$, and $r = 5$.

5. $\frac{5p + 6r^2}{p^2 + q - 1}$
6. $\frac{\sqrt{r}}{-p + 2q}$


Solve.

7. $2x - 5 + 3x = 4 - (x + 2)$ 8. $\frac{3x - 1}{5} + \frac{x + 2}{2} = -\frac{3}{10}$
9. $3 - 2(x + 3) < 4x$ 10. $2x + 4 < 10$ and $3x - 1 > 5$
11. $2x + 4 > 10$ or $3x - 1 < 5$ 12. $|5x + 3| - 10 = 3$
13. $|x + 2| < 9$ 14. $|2x - 5| \geq 9$

15. Two planes leave the Dallas–Fort Worth airport at the same time. One travels east at 550 mph, and the other travels west at 500 mph. Assuming no wind, how long will it take for the planes to be 2100 mi apart?

Plane	r	t	d
Eastbound	550	x	
Westbound	500	x	
			← Total

16. What is the slope of the line shown here?



17. Find the slope of the line passing through the points $(-4, 8)$ and $(-2, 6)$.
18. Graph $4x + 2y = -8$.

Use the function $f(x) = 2x + 7$ to find each of the following.

19. $f(-4)$ 20. The x -intercept of its graph 21. The y -intercept of its graph

Solve each system.

22. $3x - 2y = -7$ 23. $y = 5x + 6$ 24. $2x + 3y - 6z = 5$
 $2x + 3y = 17$ $-10x = -2y - 12$ $8x - y + 3z = 7$
 $3x + 4y - 3z = 7$

Perform the indicated operations.

25. $(7x + 3y)^2$ 26. $\frac{2x^4 + x^3 + 7x^2 + 2x + 6}{x^2 + 2}$
27. $(3x^3 + 4x^2 - 7) - (2x^3 - 8x^2 + 3x)$
28. (a) Write 0.0004638 using scientific notation.
 (b) Write 5.66×10^5 in standard notation.

Factor.


29. $16w^2 + 50wz - 21z^2$ 30. $4x^2 - 4x + 1 - y^2$
31. $100x^4 - 81$ 32. $8p^3 + 27$

Solve.


33. $(x - 1)(2x + 3)(x + 4) = 0$

34. $9x^2 = 6x - 1$

35. A sign is to have the shape of a triangle with a height 3 ft greater than the length of the base. How long should the base be if the area is to be 14 ft²?



36. A game board has the shape of a rectangle. The longer sides are each 2 in. longer than the distance between them. The area of the board is 288 in.². Find the length of the longer sides and the distance between them.



STUDY SKILLS REMINDER

It is not too soon to begin preparing for your final exam.

Review Study Skill 10,
Preparing for Your Math Final Exam.



RATIONAL EXPRESSIONS AND FUNCTIONS

Ratios and proportions, topics covered in this chapter and used in health care to determine medication dosages, involve rational expressions.

- 6.1** Rational Expressions and Functions; Multiplying and Dividing
- 6.2** Adding and Subtracting Rational Expressions
- 6.3** Complex Fractions
- 6.4** Equations with Rational Expressions and Graphs

SUMMARY EXERCISES Simplifying Rational Expressions vs. Solving Rational Equations

- 6.5** Applications of Rational Expressions
- 6.6** Variation

6.1 Rational Expressions and Functions; Multiplying and Dividing

OBJECTIVES

- 1 Define rational expressions.
- 2 Define rational functions and give their domains.
- 3 Write rational expressions in lowest terms.
- 4 Multiply rational expressions.
- 5 Find reciprocals of rational expressions.
- 6 Divide rational expressions.

OBJECTIVE 1 Define rational expressions.

In arithmetic, a rational number is the quotient of two integers, with the denominator not 0. In algebra, a **rational expression**, or *algebraic fraction*, is the quotient of two polynomials, again with the denominator not 0.

$$\frac{x}{y}, \quad \frac{-a}{4}, \quad \frac{m+4}{m-2}, \quad \frac{8x^2-2x+5}{4x^2+5x}, \quad x^5 \left(\text{or } \frac{x^5}{1} \right) \quad \text{Rational expressions}$$

Rational expressions are elements of the set

$$\left\{ \frac{P}{Q} \mid P \text{ and } Q \text{ are polynomials, where } Q \neq 0 \right\}.$$

OBJECTIVE 2 Define rational functions and give their domains.

Rational Function

A **rational function** is defined by a quotient of polynomials and has the form

$$f(x) = \frac{P(x)}{Q(x)}, \quad \text{where } Q(x) \neq 0.$$

The domain of a rational function includes all real numbers except those that make $Q(x)$ —that is, the denominator—equal to 0.

VOCABULARY

- rational expression
- rational function
- reciprocal

For example, the domain of the rational function

$$f(x) = \frac{2}{x-5} \leftarrow \text{Cannot equal 0}$$

includes all real numbers except 5 because 5 would make the denominator equal to 0. See FIGURE 1. The graph does not exist when $x = 5$. (It does not intersect the dashed vertical line with equation $x = 5$. This line is an *asymptote*.)




FIGURE 1

EXAMPLE 1 Finding Domains of Rational Functions

Give the domain of each rational function.

(a) $f(x) = \frac{3}{7x-14}$

Values of x that make the denominator 0 must be *excluded* from the domain. To find these values, set the denominator equal to 0 and solve.

$$7x - 14 = 0$$

$$7x = 14 \quad \text{Add 14.}$$

$$x = 2 \quad \text{Divide by 7.}$$

The number 2 cannot be used as a replacement for x . Therefore, the domain of f includes all real numbers *except* 2. It can be written using set-builder or interval notation.

Set-builder notation: $\{x \mid x \text{ is a real number, } x \neq 2\}$

Interval notation: $(-\infty, 2) \cup (2, \infty)$


NOW TRY EXERCISE 1

Give the domain of each rational function using set-builder notation and interval notation.

(a) $\frac{1}{x+9}$

(b) $\frac{2x-1}{x^2-4x-5}$

(c) $\frac{15}{2x^2+1}$

(b) $g(x) = \frac{3+x}{x^2-4x+3}$ Values that make the denominator 0 must be excluded.

$$x^2 - 4x + 3 = 0 \quad \text{Set the denominator equal to 0.}$$

$$(x-1)(x-3) = 0 \quad \text{Factor.}$$

$$x-1 = 0 \quad \text{or} \quad x-3 = 0 \quad \text{Zero-factor property}$$

$$x = 1 \quad \text{or} \quad x = 3 \quad \text{Solve each equation.}$$

Domain: $\{x \mid x \text{ is a real number, } x \neq 1, 3\}$ Set-builder notation

$$(-\infty, 1) \cup (1, 3) \cup (3, \infty) \quad \text{Interval notation}$$

(c) $h(x) = \frac{8x+2}{3}$ The denominator, 3, can never be 0, so the domain of h includes all real numbers.

Domain: $\{x \mid x \text{ is a real number}\}$ Set-builder notation

$$(-\infty, \infty) \quad \text{Interval notation}$$

(d) $f(x) = \frac{2}{x^2+4}$

Setting the denominator $x^2 + 4$ equal to 0 leads to $x^2 = -4$. There is no real number whose square is -4 , so any real number can be used as a replacement for x .

Domain: $\{x \mid x \text{ is a real number}\}$ Set-builder notation

$$(-\infty, \infty) \quad \text{Interval notation}$$

NOW TRY 

OBJECTIVE 3 Write rational expressions in lowest terms.

In arithmetic, we use the **fundamental property of rational numbers** to write a fraction in lowest terms by dividing out any common factors in the numerator and denominator.

Fundamental Property of Rational Numbers

If $\frac{a}{b}$ is a rational number and if c is any nonzero real number, then

$$\frac{a}{b} = \frac{ac}{bc}.$$

That is, the numerator and denominator of a rational number may either be multiplied or divided by the same *nonzero number* without changing the value of the rational number.

Example: $\frac{15}{20} = \frac{3 \cdot 5}{4 \cdot 5} = \frac{3}{4} \cdot 1 = \frac{3}{4}$

Because $\frac{c}{c}$ is equivalent to 1, the fundamental property is based on the identity property of multiplication.

We write a rational expression in lowest terms similarly.

Writing a Rational Expression in Lowest Terms

Step 1 Factor both numerator and denominator to find their greatest common factor (GCF).

Step 2 Apply the fundamental property. Divide out common factors.

NOW TRY ANSWERS

1. (a) $\{x \mid x \text{ is a real number, } x \neq -9\};$

$$(-\infty, -9) \cup (-9, \infty)$$

(b) $\{x \mid x \text{ is a real number, } x \neq -1, 5\};$

$$(-\infty, -1) \cup (-1, 5) \cup (5, \infty)$$

(c) $\{x \mid x \text{ is a real number}\};$

$$(-\infty, \infty)$$


NOW TRY EXERCISE 2

Write each rational expression in lowest terms.

(a) $\frac{3a^2 - 7a + 2}{a^2 + 2a - 8}$

(b) $\frac{t^3 + 8}{t + 2}$

(c) $\frac{am - bm + an - bn}{am + bm + an + bn}$

EXAMPLE 2 Writing Rational Expressions in Lowest Terms

Write each rational expression in lowest terms.

(a) $\frac{(x + 5)(x + 2)}{(x + 2)(x - 3)}$

$\frac{x + 2}{x + 2} = 1$

$= \frac{(x + 5)(x + 2)}{(x - 3)(x + 2)}$ Commutative property

$= \frac{x + 5}{x - 3}$ Fundamental property

(b) $\frac{a^2 - a - 6}{a^2 + 5a + 6}$

$= \frac{(a - 3)(a + 2)}{(a + 3)(a + 2)}$ Factor the numerator.
Factor the denominator.

$= \frac{a - 3}{a + 3}$ $\frac{a + 2}{a + 2} = 1$; Fundamental property

(c) $\frac{y^2 - 4}{2y + 4}$

$= \frac{(y + 2)(y - 2)}{2(y + 2)}$ Factor the difference of squares in the numerator.
Factor the denominator.

$= \frac{y - 2}{2}$ $\frac{y + 2}{y + 2} = 1$; Fundamental property

(d) $\frac{x^3 - 27}{x - 3}$

$= \frac{(x - 3)(x^2 + 3x + 9)}{x - 3}$ Factor the difference of cubes in the numerator.

$= x^2 + 3x + 9$ Fundamental property

(e) $\frac{pr + qr + ps + qs}{pr + qr - ps - qs}$

$= \frac{(pr + qr) + (ps + qs)}{(pr + qr) - (ps + qs)}$ Group the terms.
Be careful with signs.

$= \frac{r(p + q) + s(p + q)}{r(p + q) - s(p + q)}$ Factor within the groups.

$= \frac{(p + q)(r + s)}{(p + q)(r - s)}$ Factor by grouping.

$= \frac{r + s}{r - s}$ Fundamental property

(f) $\frac{8 + k}{16}$ Be careful. The numerator cannot be factored.

This expression cannot be simplified further and is in lowest terms.

NOW TRY 

NOW TRY ANSWERS

2. (a) $\frac{3a - 1}{a + 4}$

(b) $t^2 - 2t + 4$

(c) $\frac{a - b}{a + b}$

CAUTION Only common factors may be divided out.

Examples: $\frac{8+k}{16} \neq \frac{k}{2}$ and $\frac{8+k}{16} \neq \frac{1}{2} + k$ The 8 in $8+k$ is *not* a factor of the numerator.

The expression $\frac{8+k}{16}$ (Example 2(f)) indicates that the *entire* numerator is being divided by 16, not just one term. It is already in lowest terms, although it could be written in the equivalent form $\frac{8}{16} + \frac{k}{16}$, or $\frac{1}{2} + \frac{k}{16}$.

Compare this to the expression $\frac{8k}{16}$, which *can* be written in lowest terms.

$$\frac{8k}{16} = \frac{8 \cdot k}{8 \cdot 2} = 1 \cdot \frac{k}{2} = \frac{k}{2} \quad \text{Factor before writing a fraction in lowest terms.}$$

Look again at the rational expression from Example 2(b).

$$\frac{a^2 - a - 6}{a^2 + 5a + 6}, \quad \text{or} \quad \frac{(a-3)(a+2)}{(a+3)(a+2)}$$

In this expression, a can take any value *except* -3 or -2 , because these values make the denominator 0. In the simplified expression $\frac{a-3}{a+3}$, a cannot equal -3 . Thus,

$$\frac{a^2 - a - 6}{a^2 + 5a + 6} = \frac{a-3}{a+3}, \quad \text{for all values of } a \text{ except } -3 \text{ or } -2.$$

From now on, such statements of equality will be made with the understanding that they apply only to those real numbers that make neither denominator equal 0. We will no longer state such restrictions.

NOW TRY EXERCISE 3

Write each rational expression in lowest terms.

(a) $\frac{a-10}{10-a}$ (b) $\frac{81-y^2}{y-9}$

EXAMPLE 3 Writing Rational Expressions in Lowest Terms

Write each rational expression in lowest terms.

(a) $\frac{m-3}{3-m}$  Here, the numerator and denominator are opposites.

To write in lowest terms, we write the denominator as $-1(m-3)$, as shown on the left. Alternatively, we could write the numerator as $-1(3-m)$ and obtain the same result, as shown on the right.

$$\frac{m-3}{3-m} = \frac{m-3}{-1(m-3)} = \frac{1}{-1} = -1 \quad \left| \quad \frac{m-3}{3-m} = \frac{-1(3-m)}{3-m} = \frac{-1}{1} = -1$$

Factor out -1 in the denominator. Factor out -1 in the numerator.

(b) $\frac{r^2 - 16}{4 - r}$

$$= \frac{(r+4)(r-4)}{4-r}$$

Factor the difference of squares in the numerator.

$$= \frac{(r+4)(r-4)}{-1(r-4)}$$

Factor out -1 in the denominator to write $4-r$ as $-1(r-4)$.

Distribute to check.
 $-1(r-4)$
 $= -1 \cdot r - 1(-4)$
 $= -r + 4, \text{ or } 4-r$

$$= \frac{r+4}{-1}$$

$$\frac{r-4}{r-4} = 1; \text{ Fundamental property}$$

$$= -(r+4), \quad \text{or} \quad -r-4 \quad \text{Lowest terms}$$

NOW TRY ANSWERS

3. (a) -1
 (b) $-(y+9)$, or $-y-9$

As shown in **Example 3**, the quotient $\frac{a}{-a}$ (where $a \neq 0$) can be simplified.

$$\frac{a}{-a} = \frac{a}{-1(a)} = \frac{1}{-1} = -1$$

The quotient $\frac{-a}{a}$ (again where $a \neq 0$) can be simplified similarly.

Quotient of Opposites

In general, if the numerator and the denominator of a rational expression are opposites, then the expression equals -1 .

Examples: $\frac{q-7}{7-q} = -1$ and $\frac{-5a+2b}{5a-2b} = -1$ Numerator and denominator in each expression are opposites.

However, the following expression *cannot* be simplified further.

$$\frac{r-2}{r+2} \quad \text{Numerator and denominator are not opposites.}$$

OBJECTIVE 4 Multiply rational expressions.

Recall that if $\frac{a}{b}$ and $\frac{c}{d}$ are rational numbers (where $b \neq 0$ and $d \neq 0$), then

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd} \quad \begin{array}{l} \text{Multiply numerators.} \\ \text{Multiply denominators.} \end{array}$$

Examples: $\frac{2}{3} \cdot \frac{4}{5} = \frac{2 \cdot 4}{3 \cdot 5} = \frac{8}{15}$ and $\frac{7}{10} \cdot \frac{5}{6} = \frac{7}{2 \cdot 5} \cdot \frac{5}{2 \cdot 3} = \frac{7}{12}$

Multiplying Rational Expressions

Step 1 Factor all numerators and denominators as completely as possible.

Step 2 Apply the fundamental property—that is, $\frac{ac}{bc} = \frac{a}{b}$ (where $b, c \neq 0$).

Step 3 Multiply remaining factors in the numerators and remaining factors in the denominators. The final expression can be left in factored form.

Step 4 Check to be sure the product is in lowest terms.

EXAMPLE 4 Multiplying Rational Expressions

Multiply.

$$\begin{aligned} \text{(a)} \quad & \frac{5p-5}{p} \cdot \frac{3p^2}{10p-10} \\ &= \frac{5(p-1)}{p} \cdot \frac{3p \cdot p}{2 \cdot 5(p-1)} && \text{Factor.} \\ &= \frac{5(p-1)}{5(p-1)} \cdot \frac{p}{p} \cdot \frac{3p}{2} && \text{Commutative property} \\ &= 1 \cdot 1 \cdot 1 \cdot \frac{3p}{2} && \text{Fundamental property} \\ &= \frac{3p}{2} && \text{Lowest terms} \end{aligned}$$

In practice, this step is usually done mentally.


NOW TRY EXERCISE 4

Multiply.

$$(a) \frac{8t^2}{t^2 - 4} \cdot \frac{3t + 6}{9t}$$

$$(b) \frac{m^2 + 2m - 15}{m^2 - 5m + 6} \cdot \frac{m^2 - 4}{m^2 + 5m}$$

$$(b) \frac{k^2 + 2k - 15}{k^2 - 4k + 3} \cdot \frac{k^2 - k}{k^2 + k - 20}$$

$$= \frac{(k+5)(k-3)}{(k-3)(k-1)} \cdot \frac{k(k-1)}{(k+5)(k-4)} \quad \text{Factor.}$$

$$= \frac{k}{k-4} \quad \text{Fundamental property; Multiply.}$$

$$(c) (p-4) \cdot \frac{3}{5p-20}$$

$$= \frac{p-4}{1} \cdot \frac{3}{5p-20} \quad \text{Write } p-4 \text{ as } \frac{p-4}{1}.$$

$$= \frac{p-4}{1} \cdot \frac{3}{5(p-4)} \quad \text{Factor.}$$

$$= \frac{3}{5} \quad \text{Fundamental property; Multiply.}$$

$$(d) \frac{x^2 + 2x}{x+1} \cdot \frac{x^2 - 1}{x^3 + x^2}$$

$$= \frac{x(x+2)}{x+1} \cdot \frac{(x+1)(x-1)}{x^2(x+1)} \quad \text{Factor.}$$

$$= \frac{(x+2)(x-1)}{x(x+1)} \quad \text{Fundamental property; Multiply.}$$

The final expression can be left in factored form. It is in lowest terms.

$$(e) \frac{x-6}{x^2-12x+36} \cdot \frac{x^2-3x-18}{x^2+7x+12}$$

$$= \frac{x-6}{(x-6)^2} \cdot \frac{(x+3)(x-6)}{(x+3)(x+4)} \quad \text{Factor.}$$

Remember to include 1 in the numerator when all other factors are eliminated.

$$= \frac{1}{x+4}$$

Fundamental property; Multiply.

NOW TRY 

▼ **Reciprocals of Rational Expressions**

Rational Expression	Reciprocal
3, or $\frac{3}{1}$	$\frac{1}{3}$
$\frac{5}{k}$	$\frac{k}{5}$
$\frac{m^2-9m}{2}$	$\frac{2}{m^2-9m}$
$\frac{0}{4}$	undefined

Reciprocals have a product of 1. Recall that 0 has no reciprocal.

NOW TRY ANSWERS

4. (a) $\frac{8t}{3(t-2)}$ (b) $\frac{m+2}{m}$

OBJECTIVE 5 Find reciprocals of rational expressions.

When dividing rational numbers, we use reciprocals. The rational numbers $\frac{a}{b}$ and $\frac{c}{d}$ are reciprocals of each other if they have a product of 1.

Example: $\frac{3}{4} \cdot \frac{4}{3} = 1$, so $\frac{3}{4}$ and $\frac{4}{3}$ are reciprocals.

The **reciprocal** of a rational expression is defined in the same way. *Two rational expressions are reciprocals of each other if they have a product of 1.*

Finding the Reciprocal

To find the reciprocal of a nonzero rational expression, interchange the numerator and denominator of the expression. (See the table.)

OBJECTIVE 6 Divide rational expressions.

Recall that if $\frac{a}{b}$ and $\frac{c}{d}$ are rational numbers (where $b \neq 0$, $c \neq 0$, and $d \neq 0$), then

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} = \frac{ad}{bc}. \quad \text{To divide by a fraction, multiply by its reciprocal.}$$

Examples: $\frac{3}{4} \div \frac{2}{3} = \frac{3}{4} \cdot \frac{3}{2} = \frac{9}{8}$ and $\frac{6}{7} \div \frac{4}{3} = \frac{6}{7} \cdot \frac{3}{4} = \frac{2 \cdot 3}{7 \cdot 2} \cdot \frac{3}{2} = \frac{9}{14}$

Dividing rational expressions is similar to dividing rational numbers.

Dividing Rational Expressions

To divide two rational expressions, *multiply* the first expression (the *dividend*) by the reciprocal of the second expression (the *divisor*).

NOW TRY EXERCISE 5

Divide.

(a) $\frac{16k^2}{5} \div \frac{3k}{10}$

(b) $\frac{3k^2 + 5k - 2}{9k^2 - 1} \div \frac{4k^2 + 8k}{k^2 - 7k}$

EXAMPLE 5 Dividing Rational Expressions

Divide.

(a) $\frac{2z}{9} \div \frac{5z^2}{18}$

$$= \frac{2z}{9} \cdot \frac{18}{5z^2} \quad \text{Multiply by the reciprocal of the divisor.}$$

$$= \frac{2z}{9} \cdot \frac{2 \cdot 9}{5z \cdot z} \quad \text{Factor.}$$

$$= \frac{4}{5z} \quad \text{Fundamental property; Multiply.}$$

(b) $\frac{8k - 16}{3k} \div \frac{3k - 6}{4k^2}$

$$= \frac{8k - 16}{3k} \cdot \frac{4k^2}{3k - 6} \quad \text{Multiply by the reciprocal.}$$

$$= \frac{8(k - 2)}{3k} \cdot \frac{4k \cdot k}{3(k - 2)} \quad \text{Factor.}$$

$$= \frac{32k}{9} \quad \text{Fundamental property; Multiply.}$$

(c) $\frac{5m^2 + 17m - 12}{3m^2 + 7m - 20} \div \frac{5m^2 + 2m - 3}{15m^2 - 34m + 15}$

$$= \frac{5m^2 + 17m - 12}{3m^2 + 7m - 20} \cdot \frac{15m^2 - 34m + 15}{5m^2 + 2m - 3} \quad \text{Definition of division}$$

$$= \frac{(5m - 3)(m + 4)}{(3m - 5)(m + 4)} \cdot \frac{(5m - 3)(3m - 5)}{(5m - 3)(m + 1)} \quad \text{Factor.}$$

$$= \frac{5m - 3}{m + 1} \quad \text{Fundamental property; Multiply.}$$

NOW TRY ANSWERS

5. (a) $\frac{32k}{3}$ (b) $\frac{k - 7}{4(3k + 1)}$

6.1 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

We can learn from the mistakes we make. **Review Study Skill 9, Analyzing Your Test Results.**

Concept Check Complete each statement.

- A rational number such as $\frac{3}{4}$ is the quotient of two _____, with denominator not _____.
A _____ expression such as $\frac{2x^2}{2x+1}$ is the quotient of two _____, with denominator not _____.
- A function of the form $f(x) = \frac{P(x)}{Q(x)}$, where $P(x)$ and $Q(x)$ are polynomials and $Q(x) \neq$ _____, is a _____ function. The domain includes all _____ except any values of x that make the (*numerator / denominator*) equal to 0.

Give the domain of each rational function using (a) set-builder notation and (b) interval notation. See Example 1.

3. $f(x) = \frac{x}{x-7}$

4. $f(x) = \frac{x}{x+3}$

5. $f(x) = \frac{6x-5}{7x+1}$

6. $f(x) = \frac{8x-3}{2x+7}$

7. $f(x) = \frac{12x+3}{x}$

8. $f(x) = \frac{9x+8}{x}$

9. $f(x) = \frac{3x+1}{2x^2+x-6}$

10. $f(x) = \frac{2x+4}{3x^2+11x-42}$

11. $f(x) = \frac{x+2}{14}$

12. $f(x) = \frac{x-9}{26}$

13. $f(x) = \frac{2x^2-3x+4}{3x^2+8}$

14. $f(x) = \frac{9x^2-8x+3}{4x^2+1}$

Concept Check As review, multiply or divide the rational numbers as indicated. Write answers in lowest terms.

15. $\frac{4}{21} \cdot \frac{7}{10}$

16. $\frac{5}{9} \cdot \frac{12}{25}$

17. $\frac{3}{8} \div \frac{5}{12}$

18. $\frac{5}{6} \div \frac{14}{15}$

19. $\frac{2}{3} \div \frac{8}{9}$

20. $\frac{3}{8} \div \frac{9}{14}$

Concept Check Answer each question.

21. Which rational expressions are equivalent to $-\frac{x}{y}$?

- A. $\frac{-x}{-y}$ B. $\frac{x}{-y}$ C. $\frac{x}{y}$ D. $\frac{-x}{-y}$ E. $\frac{-x}{y}$ F. $\frac{-x}{-y}$

22. Which rational expression can be simplified?

- A. $\frac{x^2+2}{x^2}$ B. $\frac{x^2+2}{2}$ C. $\frac{x^2+y^2}{y^2}$ D. $\frac{x^2-5x}{x}$

23. Which rational expression is *not* equivalent to $\frac{x-3}{4-x}$?

- A. $\frac{3-x}{x-4}$ B. $\frac{x+3}{4+x}$ C. $\frac{3-x}{4-x}$ D. $\frac{x-3}{x-4}$

24. Which two rational expressions are equivalent to -1 ?

- A. $\frac{2x+3}{2x-3}$ B. $\frac{2x-3}{3-2x}$ C. $\frac{2x+3}{3+2x}$ D. $\frac{2x+3}{-2x-3}$

25. Concept Check Consider the expressions $\frac{x-2}{4}$ and $\frac{2x}{4}$. Only one of these expressions is not in lowest terms. Which one is it and why? Write it in lowest terms.

26. Concept Check Identify the two *terms* in the numerator and the two *terms* in the denominator of the rational expression $\frac{x^2 + 4x}{x + 4}$, and write it in lowest terms.

27. Concept Check A student was asked to simplify the rational expression

$$\frac{2x + y}{4x + 2y}$$

and wrote the incorrect answer $\frac{x + y}{4x + y}$.

WHAT WENT WRONG? What is the correct answer?

28. Concept Check A student was asked to simplify the rational expression

$$\frac{a + b}{a - b}$$

and wrote the incorrect answer $\frac{+1}{-1}$.

WHAT WENT WRONG? What is the correct answer?

Write each rational expression in lowest terms. See Example 2.

29. $\frac{x^2(x + 1)}{x(x + 1)}$

30. $\frac{y^3(y - 4)}{y^2(y - 4)}$

31. $\frac{(x + 4)(x - 3)}{(x + 5)(x + 4)}$

32. $\frac{(2x + 7)(x - 1)}{(2x + 3)(2x + 7)}$

33. $\frac{4x(x + 3)}{8x^2(x - 3)}$

34. $\frac{5y^2(y + 8)}{15y(y - 8)}$

35. $\frac{3x + 7}{3}$

36. $\frac{4x - 9}{4}$

37. $\frac{6m + 18}{7m + 21}$

38. $\frac{5r - 20}{3r - 12}$

39. $\frac{3z^2 + z}{18z + 6}$

40. $\frac{2x^2 - 5x}{16x - 40}$

41. $\frac{t^2 - 9}{3t + 9}$

42. $\frac{m^2 - 25}{4m - 20}$

43. $\frac{2t + 6}{t^2 - 9}$

44. $\frac{5s - 25}{s^2 - 25}$

45. $\frac{x^2 + 2x - 15}{x^2 + 6x + 5}$

46. $\frac{y^2 - 5y - 14}{y^2 + y - 2}$

47. $\frac{8x^2 - 10x - 3}{8x^2 - 6x - 9}$

48. $\frac{12x^2 - 4x - 5}{8x^2 - 6x - 5}$

49. $\frac{a^3 + b^3}{a + b}$

50. $\frac{r^3 - s^3}{r - s}$

51. $\frac{2c^2 + 2cd - 60d^2}{2c^2 - 12cd + 10d^2}$

52. $\frac{3s^2 - 9st - 54t^2}{3s^2 - 6st - 72t^2}$

53. $\frac{ac - ad + bc - bd}{ac - ad - bc + bd}$

54. $\frac{2xy + 2xw + y + w}{2xy + y - 2xw - w}$

Write each rational expression in lowest terms. See Example 3.

55. $\frac{7 - b}{b - 7}$

56. $\frac{r - 13}{13 - r}$

57. $\frac{x^2 - y^2}{y - x}$

58. $\frac{m^2 - n^2}{n - m}$

59. $\frac{x^2 - 4}{2 - x}$

60. $\frac{x^2 - 81}{9 - x}$

61. $\frac{(a - 3)(x + y)}{(3 - a)(x - y)}$

62. $\frac{(8 - p)(x + 2)}{(p - 8)(x - 2)}$

63. $\frac{5k - 10}{20 - 10k}$

64. $\frac{7x - 21}{63 - 21x}$

65. $\frac{a^2 - b^2}{a^2 + b^2}$

66. $\frac{p^2 + q^2}{p^2 - q^2}$

Multiply or divide as indicated. See Examples 4 and 5.

67. $\frac{(x+2)(x+1)}{(x+3)(x-2)} \cdot \frac{(x+3)(x+4)}{(x+2)(x+1)}$
68. $\frac{(x+3)(x-6)}{(x-4)(x+2)} \cdot \frac{(x+5)(x-4)}{(x+3)(x-6)}$
69. $\frac{(2x+3)(x-4)}{(x+8)(x-4)} \div \frac{(x-4)(x+2)}{(x-4)(x+8)}$
70. $\frac{(6x+5)(x-3)}{(x-1)(x-3)} \div \frac{(2x+7)(x+9)}{(x-1)(x+9)}$
71. $\frac{4x}{8x+4} \cdot \frac{14x+7}{6}$
72. $\frac{12x-20}{5x} \cdot \frac{6}{9x-15}$
73. $\frac{p^2-25}{4p} \cdot \frac{2}{5-p}$
74. $\frac{a^2-1}{4a} \cdot \frac{2}{1-a}$
75. $(7k+7) \div \frac{4k+4}{5}$
76. $(8y-16) \div \frac{3y-6}{10}$
77. $(z^2-1) \cdot \frac{1}{1-z}$
78. $(y^2-4) \cdot \frac{8}{2-y}$
79. $\frac{4x-20}{5x} \div \frac{2x-10}{7x^3}$
80. $\frac{3x+12}{4x} \div \frac{4x+16}{6x^2}$
81. $\frac{m^2-49}{m+1} \div \frac{7-m}{m}$
82. $\frac{k^2-4}{3k^2} \div \frac{2-k}{11k}$
83. $\frac{12x-10y}{3x+2y} \cdot \frac{6x+4y}{10y-12x}$
84. $\frac{9s-12t}{2s+2t} \cdot \frac{3s+3t}{4t-3s}$
85. $\frac{x^2-25}{x^2+x-20} \cdot \frac{x^2+7x+12}{x^2-2x-15}$
86. $\frac{t^2-49}{t^2+4t-21} \cdot \frac{t^2+8t+15}{t^2-2t-35}$
87. $\frac{a^3-b^3}{a^2-b^2} \div \frac{2a-2b}{2a+2b}$
88. $\frac{x^3+y^3}{2x+2y} \div \frac{x^2-y^2}{2x-2y}$
89. $\frac{8x^3-27}{2x^2-18} \cdot \frac{2x+6}{8x^2+12x+18}$
90. $\frac{64x^3+1}{4x^2-100} \cdot \frac{4x+20}{64x^2-16x+4}$
91. $\frac{a^3-8b^3}{a^2-ab-6b^2} \cdot \frac{a^2+ab-12b^2}{a^2+2ab-8b^2}$
92. $\frac{p^3-27q^3}{p^2+pq-12q^2} \cdot \frac{p^2-2pq-24q^2}{p^2-5pq-6q^2}$
93. $\frac{6x^2+5x-6}{12x^2-11x+2} \div \frac{4x^2-12x+9}{8x^2-14x+3}$
94. $\frac{8a^2-6a-9}{6a^2-5a-6} \div \frac{4a^2+11a+6}{9a^2+12a+4}$

Extending Skills Multiply or divide as indicated.

95. $\frac{(-3mn)^2 \cdot 64(m^2n)^3}{16m^2n^4(mn^2)^3} \div \frac{24(m^2n^2)^4}{(3m^2n^3)^2}$
96. $\frac{(-4a^2b^3)^2 \cdot 9(a^2b^4)^2}{(2a^2b^3)^4 \cdot (3a^3b)^2} \div \frac{(ab)^4}{(a^2b^3)^2}$
97. $\frac{3k^2+17kp+10p^2}{6k^2+13kp-5p^2} \div \frac{6k^2+kp-2p^2}{6k^2-5kp+p^2}$
98. $\frac{16c^2+24cd+9d^2}{16c^2-16cd+3d^2} \div \frac{16c^2-9d^2}{16c^2-24cd+9d^2}$
99. $\left(\frac{6k^2-13k-5}{k^2+7k} \div \frac{2k-5}{k^3+6k^2-7k} \right) \cdot \frac{k^2-5k+6}{3k^2-8k-3}$
100. $\left(\frac{2x^3+3x^2-2x}{3x-15} \div \frac{2x^3-x^2}{x^2-3x-10} \right) \cdot \frac{5x^2-10x}{3x^2+12x+12}$

6.2 Adding and Subtracting Rational Expressions

OBJECTIVES

- 1 Add and subtract rational expressions with the same denominator.
- 2 Find a least common denominator.
- 3 Add and subtract rational expressions with different denominators.

VOCABULARY

- least common denominator (LCD)

OBJECTIVE 1 Add and subtract rational expressions with the same denominator.

Recall that if $\frac{a}{b}$ and $\frac{c}{b}$ are rational numbers (where $b \neq 0$), then

$$\frac{a}{b} + \frac{c}{b} = \frac{a + c}{b} \quad \text{and} \quad \frac{a}{b} - \frac{c}{b} = \frac{a - c}{b}. \quad \begin{array}{l} \text{Add or subtract the numerators.} \\ \text{Keep the same denominator.} \end{array}$$

Examples: $\frac{4}{7} + \frac{1}{7} = \frac{5}{7}$, $\frac{4}{7} - \frac{1}{7} = \frac{3}{7}$, and $\frac{3}{10} - \frac{1}{10} = \frac{2}{10} = \frac{2 \cdot 1}{2 \cdot 5} = \frac{1}{5}$

Adding and subtracting rational expressions is similar.

Adding or Subtracting Rational Expressions

Step 1 If the denominators are the same, add or subtract the numerators. Place the result over the common denominator.

If the denominators are different, find the least common denominator and write all rational expressions with this LCD. Add or subtract the numerators. Place the result over the common denominator.

Step 2 Simplify. Write all answers in lowest terms.

EXAMPLE 1 Adding and Subtracting Rational Expressions (Same Denominators)

Add or subtract as indicated.

$$\begin{aligned} \text{(a)} \quad & \frac{3y}{5} + \frac{x}{5} \\ &= \frac{3y + x}{5} \quad \begin{array}{l} \leftarrow \text{Add the numerators.} \\ \leftarrow \text{Keep the common denominator.} \end{array} \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad & \frac{7}{2r^2} - \frac{11}{2r^2} \\ &= \frac{7 - 11}{2r^2} \quad \begin{array}{l} \text{Subtract the numerators.} \\ \text{Keep the common denominator.} \end{array} \\ &= \frac{-4}{2r^2} \\ &= -\frac{2}{r^2} \quad \frac{-4}{2r^2} = \frac{-2 \cdot 2}{2 \cdot r^2}; \text{ Write in lowest terms.} \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad & \frac{m}{m^2 - p^2} + \frac{p}{m^2 - p^2} \\ &= \frac{m + p}{m^2 - p^2} \quad \begin{array}{l} \text{Add the numerators.} \\ \text{Keep the common denominator.} \end{array} \\ &= \frac{m + p}{(m + p)(m - p)} \quad \text{Factor.} \\ &= \frac{1}{m - p} \quad \text{Fundamental property} \end{aligned}$$

Remember to write 1 in the numerator.


NOW TRY EXERCISE 1

Add or subtract as indicated.

(a) $\frac{5}{3x} + \frac{2}{3x}$

(b) $\frac{x^2}{x-3} - \frac{9}{x-3}$

(c) $\frac{2}{x^2+x-2} + \frac{x}{x^2+x-2}$

(d) $\frac{4}{x^2+2x-8} + \frac{x}{x^2+2x-8}$

$$= \frac{4+x}{x^2+2x-8} \quad \text{Add.}$$

$$= \frac{4+x}{(x-2)(x+4)} \quad \text{Factor.}$$

$$= \frac{1}{x-2} \quad \text{Fundamental property}$$

NOW TRY 
OBJECTIVE 2 Find a least common denominator.

We add or subtract rational numbers with different denominators by first writing them with a common denominator, usually the **least common denominator (LCD)**.

Finding the Least Common Denominator

Step 1 **Factor** each denominator.

Step 2 **Find the least common denominator.** The LCD is the product of all of the different factors from each denominator, with each factor raised to the *greatest* power that occurs in any denominator.

Example: To find the LCD for $\frac{7}{15}$ and $\frac{5}{12}$, we factor each denominator.

$$15 = 3 \cdot 5 \quad \text{and} \quad 12 = 2 \cdot 2 \cdot 3 = 2^2 \cdot 3$$

$$\text{LCD} = 2^2 \cdot 3 \cdot 5 = 60 \quad \text{Find the product of the factors, each raised to its greatest power.}$$

We use a similar procedure to find the LCD of two or more rational expressions.

EXAMPLE 2 Finding Least Common Denominators

Suppose that the given expressions are denominators of rational expressions. Find the LCD for each group of denominators.

(a) $5xy^2, 2x^3y$

$$5xy^2 = 5 \cdot x \cdot y^2 \quad \text{Each denominator is already factored.}$$

$$2x^3y = 2 \cdot x^3 \cdot y$$

$$\begin{aligned} \text{LCD} &= 5 \cdot 2 \cdot x^3 \cdot y^2 \quad \leftarrow \text{Greatest exponent on } y \text{ is } 2. \\ &= 10x^3y^2 \quad \leftarrow \text{Greatest exponent on } x \text{ is } 3. \end{aligned}$$

(b) $k-3, k$ The LCD must be divisible by both $k-3$ and k .

$$\text{LCD} = k(k-3) \quad \leftarrow \text{Don't forget the factor } k.$$

It is usually best to leave a least common denominator in factored form.

(c) $y^2 - 2y - 8, y^2 + 3y + 2$

$$\left. \begin{aligned} y^2 - 2y - 8 &= (y-4)(y+2) \\ y^2 + 3y + 2 &= (y+2)(y+1) \end{aligned} \right\} \text{Factor.}$$

$$\text{LCD} = (y-4)(y+2)(y+1)$$

NOW TRY ANSWERS

1. (a) $\frac{7}{3x}$ (b) $x+3$ (c) $\frac{1}{x-1}$


NOW TRY EXERCISE 2

Find the LCD for each group of denominators.

(a) $15m^3n, 10m^2n$

(b) $t, t - 8$

(c) $3x^2 + 9x - 30, x^2 - 4,$
 $x^2 + 10x + 25$

(d) $8z - 24, 5z^2 - 15z$

$$\left. \begin{aligned} 8z - 24 &= 8(z - 3) \\ 5z^2 - 15z &= 5z(z - 3) \end{aligned} \right\} \text{Factor.}$$

$$\text{LCD} = 8 \cdot 5z \cdot (z - 3)$$

$$= 40z(z - 3)$$

(e) $m^2 + 5m + 6, m^2 + 4m + 4, 2m^2 + 4m - 6$

$$m^2 + 5m + 6 = (m + 3)(m + 2)$$

$$m^2 + 4m + 4 = (m + 2)^2$$

$$2m^2 + 4m - 6 = 2(m + 3)(m - 1)$$

Factor. Notice that $(m + 2)$ is squared in the second denominator.

Use each factor, raised to its *greatest* power in any denominator. This means that $(m + 2)^2$ must be used.

$$\text{LCD} = 2(m + 3)(m + 2)^2(m - 1)$$

NOW TRY 

OBJECTIVE 3 Add and subtract rational expressions with different denominators.

Recall how we add fractions with different denominators.

Example: $\frac{7}{15} + \frac{5}{12}$

The LCD for 15 and 12 is 60.
(See Objective 2.)

$\frac{4}{4}$ and $\frac{5}{5}$ are forms of 1, the identity element of multiplication.

$$= \frac{7 \cdot 4}{15 \cdot 4} + \frac{5 \cdot 5}{12 \cdot 5}$$

Fundamental property

$$= \frac{28}{60} + \frac{25}{60}$$

Write each fraction with the common denominator.

$$= \frac{28 + 25}{60}$$

Add the numerators.

Keep the common denominator.

$$= \frac{53}{60}$$

We add and subtract rational expressions with different denominators similarly.

EXAMPLE 3 Adding and Subtracting Rational Expressions (Different Denominators)

Add or subtract as indicated.

(a) $\frac{5}{2p} + \frac{3}{8p}$

The LCD for $2p$ and $8p$ is $8p$.

$$= \frac{5 \cdot 4}{2p \cdot 4} + \frac{3}{8p}$$

Fundamental property

$$= \frac{20}{8p} + \frac{3}{8p}$$

Write the first fraction with the common denominator.

$$= \frac{23}{8p}$$

Add the numerators.

Keep the common denominator.

NOW TRY ANSWERS

2. (a) $30m^3n$ (b) $t(t - 8)$

(c) $3(x - 2)(x + 2)(x + 5)^2$


NOW TRY EXERCISE 3

Add or subtract as indicated.

(a) $\frac{2}{3x} + \frac{7}{4x}$

(b) $\frac{3}{z} - \frac{6}{z-5}$

(b) $\frac{6}{r} - \frac{5}{r-3}$

The LCD is $r(r-3)$.

$$= \frac{6(r-3)}{r(r-3)} - \frac{r \cdot 5}{r(r-3)}$$

Fundamental property

$$= \frac{6r-18}{r(r-3)} - \frac{5r}{r(r-3)}$$

Distributive and commutative properties

$$= \frac{6r-18-5r}{r(r-3)}$$

Subtract the numerators.

Keep the common denominator.

$$= \frac{r-18}{r(r-3)}$$

Combine like terms in the numerator.

NOW TRY 

! CAUTION Sign errors occur easily when a rational expression with two or more terms in the numerator is being subtracted. *The subtraction sign must be distributed to every term in the numerator of the fraction that follows it.* Study Example 4 carefully.


NOW TRY EXERCISE 4

Subtract.

(a) $\frac{18x+7}{4x+5} - \frac{2x-13}{4x+5}$

(b) $\frac{5}{x-3} - \frac{5}{x+3}$


EXAMPLE 4 Subtracting Rational Expressions

Subtract.

(a) $\frac{7x}{3x+1} - \frac{x-2}{3x+1}$

The denominators are the same. *The subtraction sign must be applied to both terms in the numerator of the second rational expression.*

$$\frac{7x}{3x+1} - \frac{x-2}{3x+1}$$



$$= \frac{7x - (x-2)}{3x+1}$$

Subtract the numerators.

Keep the common denominator.



$$= \frac{7x - x + 2}{3x+1}$$

Distributive property

$$= \frac{6x+2}{3x+1}$$

Combine like terms in the numerator.

$$= \frac{2(3x+1)}{3x+1}$$

Factor the numerator.

$$= 2$$

Fundamental property

(b) $\frac{1}{q-1} - \frac{1}{q+1}$

The LCD is $(q-1)(q+1)$.

$$= \frac{1(q+1)}{(q-1)(q+1)} - \frac{1(q-1)}{(q+1)(q-1)}$$

Fundamental property

$$= \frac{(q+1) - (q-1)}{(q-1)(q+1)}$$

Subtract the numerators.

$$= \frac{q+1 - q+1}{(q-1)(q+1)}$$



Distributive property

$$= \frac{2}{(q-1)(q+1)}$$

Combine like terms in the numerator.

NOW TRY 
NOW TRY ANSWERS

3. (a) $\frac{29}{12x}$ (b) $\frac{-3z-15}{z(z-5)}$

4. (a) 4 (b) $\frac{30}{(x-3)(x+3)}$


NOW TRY EXERCISE 5

Add.

$$\frac{t}{t-9} + \frac{2}{9-t}$$

EXAMPLE 5 Adding Rational Expressions (Denominators Are Opposites)

Add.

$$\frac{y}{y-2} + \frac{8}{2-y} \leftarrow \text{Denominators are opposites.}$$

$$= \frac{y}{y-2} + \frac{8(-1)}{(2-y)(-1)} \quad \text{Multiply the second expression by 1 in the form } \frac{-1}{-1}.$$

$$= \frac{y}{y-2} + \frac{-8}{y-2} \quad \text{The LCD is } y-2.$$

$$= \frac{y-8}{y-2} \quad \text{Add the numerators.}$$

We could use $2-y$ as the common denominator and rewrite the first expression.

$$\frac{y}{y-2} + \frac{8}{2-y}$$

$$= \frac{y(-1)}{(y-2)(-1)} + \frac{8}{2-y} \quad \text{Multiply the first expression by 1 in the form } \frac{-1}{-1}.$$

$$\rightarrow \frac{-y}{2-y} + \frac{8}{2-y} \quad \text{The LCD is } 2-y.$$

$$= \frac{-y+8}{2-y} \quad \text{Add the numerators.}$$

$$= \frac{8-y}{2-y} \quad \text{This is an equivalent form of the answer in red above. Multiply it by } \frac{-1}{-1} \text{ to see this.}$$

NOW TRY 

NOW TRY EXERCISE 6

Add and subtract as indicated.

$$\frac{6}{y} - \frac{1}{y-3} + \frac{3}{y^2-3y}$$

EXAMPLE 6 Adding and Subtracting Three Rational Expressions

Add and subtract as indicated.

$$\frac{3}{x-2} + \frac{5}{x} - \frac{6}{x^2-2x}$$

$$= \frac{3}{x-2} + \frac{5}{x} - \frac{6}{x(x-2)} \quad \text{Factor the third denominator.}$$

$$= \frac{3x}{x(x-2)} + \frac{5(x-2)}{x(x-2)} - \frac{6}{x(x-2)} \quad \text{The LCD is } x(x-2); \text{ fundamental property}$$

$$= \frac{3x + 5(x-2) - 6}{x(x-2)} \quad \text{Add and subtract the numerators.}$$

$$= \frac{3x + 5x - 10 - 6}{x(x-2)} \quad \text{Distributive property}$$

$$= \frac{8x - 16}{x(x-2)} \quad \text{Combine like terms in the numerator.}$$

$$= \frac{8(x-2)}{x(x-2)} \quad \text{Factor the numerator.}$$

$$= \frac{8}{x} \quad \text{Fundamental property}$$

NOW TRY 
NOW TRY ANSWERS

5. $\frac{t-2}{t-9}$, or $\frac{2-t}{9-t}$

6. $\frac{5}{y}$


**NOW TRY
EXERCISE 7**

Subtract.

$$\frac{t-1}{t^2-2t-8} - \frac{2t+3}{t^2+3t+2}$$

EXAMPLE 7 Subtracting Rational Expressions

Subtract.

$$\frac{m+4}{m^2-2m-3} - \frac{2m-3}{m^2-5m+6}$$

$$= \frac{m+4}{(m-3)(m+1)} - \frac{2m-3}{(m-3)(m-2)} \quad \text{Factor each denominator.}$$

$$= \frac{(m+4)(m-2)}{(m-3)(m+1)(m-2)} - \frac{(2m-3)(m+1)}{(m-3)(m-2)(m+1)} \quad \text{Fundamental property}$$

The LCD is $(m-3)(m+1)(m-2)$.

$$= \frac{(m+4)(m-2) - (2m-3)(m+1)}{(m-3)(m+1)(m-2)} \quad \text{Subtract the numerators.}$$

$$= \frac{m^2 + 2m - 8 - (2m^2 - m - 3)}{(m-3)(m+1)(m-2)} \quad \text{Multiply in the numerator.}$$

Note the careful use of parentheses.

$$= \frac{m^2 + 2m - 8 - 2m^2 + m + 3}{(m-3)(m+1)(m-2)} \quad \text{Distributive property}$$

Be careful with signs.

$$= \frac{-m^2 + 3m - 5}{(m-3)(m+1)(m-2)} \quad \text{Combine like terms in the numerator.}$$

If we try to factor the numerator, we find that this expression is in lowest terms.

NOW TRY 

**NOW TRY
EXERCISE 8**

Add.

$$\frac{2}{m^2-6m+9} + \frac{4}{m^2+m-12}$$

EXAMPLE 8 Adding Rational Expressions

Add.

$$\frac{5}{x^2+10x+25} + \frac{2}{x^2+7x+10}$$

$$= \frac{5}{(x+5)^2} + \frac{2}{(x+5)(x+2)} \quad \text{Factor each denominator.}$$

$$= \frac{5(x+2)}{(x+5)^2(x+2)} + \frac{2(x+5)}{(x+5)(x+2)(x+5)} \quad \text{The LCD is } (x+5)^2(x+2); \text{ fundamental property}$$

$$= \frac{5(x+2) + 2(x+5)}{(x+5)^2(x+2)} \quad \text{Add the numerators.}$$

$$= \frac{5x+10+2x+10}{(x+5)^2(x+2)} \quad \text{Distributive property}$$

$$= \frac{7x+20}{(x+5)^2(x+2)} \quad \text{Combine like terms in the numerator.}$$

NOW TRY ANSWERS

7. $\frac{-t^2+5t+11}{(t+2)(t-4)(t+1)}$

8. $\frac{6m-4}{(m-3)^2(m+4)}$

NOW TRY 

6.2 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Make study cards to help you learn and remember the material in this chapter.

Review Study Skill 5,
Using Study Cards.

1. Concept Check The least common denominator of two fractions is the (*sum/product*) of all the different (*terms/factors*) from each denominator, with each factor raised to the (*least/greatest*) power that occurs in any (*numerator/denominator*).

2. Concept Check Find the missing numerator so that the two fractions are equivalent.

$$(a) \frac{4}{5} = \frac{?}{25} \quad (b) \frac{7}{x} = \frac{?}{10x} \quad (c) \frac{3}{2-x} = \frac{?}{x-2}$$

3. Concept Check What is the LCD for the expressions $\frac{1}{x}$ and $-\frac{1}{9x}$?

A. 9 B. x C. $9x$ D. $9x^2$

4. Concept Check What is the LCD for the expressions $\frac{4}{a}$ and $\frac{2}{a+6}$?

A. a B. $a+6$ C. $2a+6$ D. $a(a+6)$

Concept Check As review, add or subtract the rational numbers as indicated. Write answers in lowest terms.

5. $\frac{8}{15} + \frac{4}{15}$

6. $\frac{5}{16} + \frac{9}{16}$

7. $\frac{5}{6} - \frac{8}{9}$

8. $\frac{3}{4} - \frac{5}{6}$

9. $\frac{5}{18} + \frac{7}{12}$

10. $\frac{3}{10} + \frac{7}{15}$

Add or subtract as indicated. See Example 1.

11. $\frac{7}{t} + \frac{2}{t}$

12. $\frac{5}{r} + \frac{9}{r}$

13. $\frac{6x}{7} + \frac{y}{7}$

14. $\frac{12t}{5} + \frac{s}{5}$

15. $\frac{11}{5x} - \frac{1}{5x}$

16. $\frac{7}{4y} - \frac{3}{4y}$

17. $\frac{9}{4x^3} - \frac{17}{4x^3}$

18. $\frac{6}{5y^4} - \frac{21}{5y^4}$

19. $\frac{5x+4}{6x+5} + \frac{x+1}{6x+5}$

20. $\frac{6y+12}{4y+3} + \frac{2y-6}{4y+3}$

21. $\frac{x^2}{x+5} - \frac{25}{x+5}$

22. $\frac{y^2}{y+6} - \frac{36}{y+6}$

23. $\frac{4}{p^2+7p+12} + \frac{p}{p^2+7p+12}$

24. $\frac{5}{x^2+x-20} + \frac{x}{x^2+x-20}$

25. $\frac{-3x+7}{x^2+2x-8} + \frac{8x+13}{x^2+2x-8}$

26. $\frac{5x+6}{x^2+11x+30} + \frac{4-3x}{x^2+11x+30}$

27. $\frac{a^3}{a^2+ab+b^2} - \frac{b^3}{a^2+ab+b^2}$

28. $\frac{p^3}{p^2-pq+q^2} + \frac{q^3}{p^2-pq+q^2}$

Suppose that the given expressions are denominators of rational expressions. Find the least common denominator (LCD) for each group of denominators. See Example 2.

29. $18x^2y^3, 24x^4y^5$

30. $24a^3b^4, 18a^5b^2$

31. $z-2, z$

32. $x+3, x$

33. $2y+8, y+4$

34. $3r-21, r-7$

35. $x^2 - 81$, $x^2 + 18x + 81$

37. $m + n$, $m - n$, $m^2 - n^2$

39. $x^2 - 3x - 4$, $x + x^2$

41. $2t^2 + 7t - 15$, $t^2 + 3t - 10$

43. $2y + 6$, $y^2 - 9$, y

45. $2x - 6$, $x^2 - x - 6$, $(x + 2)^2$

36. $y^2 - 16$, $y^2 - 8y + 16$

38. $r + s$, $r - s$, $r^2 - s^2$

40. $y^2 - 8y + 12$, $y^2 - 6y$

42. $s^2 - 3s - 4$, $3s^2 + s - 2$

44. $9x + 18$, $x^2 - 4$, x

46. $3a - 3b$, $a^2 + ab - 2b^2$, $(a - b)^2$

47. **Concept Check** Consider the following incorrect work.

$$\begin{aligned} \frac{x}{x+2} - \frac{4x-1}{x+2} \\ &= \frac{x-4x-1}{x+2} \\ &= \frac{-3x-1}{x+2} \end{aligned}$$

WHAT WENT WRONG? Give the correct answer.

48. **Concept Check** On a test a student wrote the answer

$$\frac{-1}{6-x}$$

His friend claimed that this is wrong and that the correct answer is

$$\frac{1}{x-6}$$

The friend's assertion is incorrect.

WHAT WENT WRONG?

Add or subtract as indicated. See Examples 3–8.

49. $\frac{8}{t} + \frac{7}{3t}$

50. $\frac{5}{x} + \frac{9}{4x}$

51. $\frac{5}{12x^2y} - \frac{11}{6xy}$

52. $\frac{7}{18a^3b^2} - \frac{2}{9ab}$

53. $\frac{4}{15a^4b^5} + \frac{3}{20a^2b^6}$

54. $\frac{5}{12x^5y^2} + \frac{5}{18x^4y^5}$

55. $\frac{2r}{7p^3q^4} + \frac{3s}{14p^4q}$

56. $\frac{4t}{9a^8b^7} + \frac{5s}{27a^4b^3}$

57. $\frac{1}{a^3b^2} - \frac{2}{a^4b} + \frac{3}{a^5b^7}$

58. $\frac{5}{t^4u^7} - \frac{3}{t^5u^9} + \frac{6}{t^{10}u}$

59. $\frac{1}{x-1} - \frac{1}{x}$

60. $\frac{3}{x-3} - \frac{1}{x}$

61. $\frac{3a}{a+1} + \frac{2a}{a-3}$

62. $\frac{2x}{x+4} + \frac{3x}{x-7}$

63. $\frac{11x-13}{2x-3} - \frac{3x-1}{2x-3}$

64. $\frac{13x-5}{4x-1} - \frac{x-2}{4x-1}$

65. $\frac{17y+3}{9y+7} - \frac{-10y-18}{9y+7}$

66. $\frac{4x+1}{3x+2} - \frac{-2x-3}{3x+2}$

67. $\frac{2}{4-x} + \frac{5}{x-4}$

68. $\frac{3}{2-t} + \frac{1}{t-2}$

69. $\frac{w}{w-z} - \frac{z}{z-w}$

70. $\frac{a}{a-b} - \frac{b}{b-a}$

71. $\frac{1}{x+1} - \frac{1}{x-1}$

72. $\frac{-2}{x-1} + \frac{2}{x+1}$

73. $\frac{4x}{x-1} - \frac{2}{x+1} - \frac{4}{x^2-1}$

74. $\frac{4}{x+3} - \frac{x}{x-3} - \frac{18}{x^2-9}$

75. $\frac{15}{y^2+3y} + \frac{2}{y} + \frac{5}{y+3}$

76. $\frac{7}{t-2} - \frac{6}{t^2-2t} - \frac{3}{t}$

77. $\frac{5}{x-2} + \frac{1}{x} + \frac{2}{x^2-2x}$

78. $\frac{5x}{x-3} + \frac{2}{x} + \frac{6}{x^2-3x}$

79. $\frac{3x}{x+1} + \frac{4}{x-1} - \frac{6}{x^2-1}$

80. $\frac{5x}{x+3} + \frac{x+2}{x} - \frac{6}{x^2+3x}$

81. $\frac{4}{x+1} + \frac{1}{x^2-x+1} - \frac{12}{x^3+1}$

83. $\frac{2x+4}{x+3} + \frac{3}{x} - \frac{6}{x^2+3x}$

85. $\frac{3}{(p-2)^2} - \frac{5}{p-2} + 4$

87. $\frac{3}{x^2-5x+6} - \frac{2}{x^2-4x+4}$

89. $\frac{3}{x^2+4x+4} + \frac{7}{x^2+5x+6}$

91. $\frac{5x}{x^2+xy-2y^2} - \frac{3x}{x^2+5xy-6y^2}$

93. $\frac{5x-y}{x^2+xy-2y^2} - \frac{3x+2y}{x^2+5xy-6y^2}$

95. $\frac{r+s}{3r^2+2rs-s^2} - \frac{s-r}{6r^2-5rs+s^2}$

82. $\frac{5}{x+2} + \frac{2}{x^2-2x+4} - \frac{60}{x^3+8}$

84. $\frac{4x+1}{x+5} - \frac{2}{x} + \frac{10}{x^2+5x}$

86. $\frac{8}{(3r-1)^2} + \frac{2}{3r-1} - 6$

88. $\frac{2}{m^2-4m+4} + \frac{3}{m^2+m-6}$

90. $\frac{5}{x^2+6x+9} - \frac{2}{x^2+4x+3}$

92. $\frac{6x}{6x^2+5xy-4y^2} - \frac{2y}{9x^2-16y^2}$

94. $\frac{6x+5y}{6x^2+5xy-4y^2} - \frac{x+2y}{9x^2-16y^2}$

96. $\frac{3y}{y^2+yz-2z^2} + \frac{4y-1}{y^2-z^2}$

Work each problem.

97. A **Concours d'Elegance** is a competition in which a maximum of 100 points is awarded to a car on the basis of its general attractiveness. The function defined by the rational expression

$$C(x) = \frac{9010}{49(101-x)} - \frac{10}{49}$$

approximates the cost, in thousands of dollars, of restoring a car so that it will win x points.

- (a) Simplify the expression for $C(x)$ by performing the indicated subtraction.
- (b) Use the simplified expression to determine, to two decimal places, how much it would cost to win 95 points.
98. A **cost-benefit model** expresses cost of an undertaking in terms of benefits received. One such model gives the cost, in thousands of dollars, to remove x percent of a pollutant as

$$C(x) = \frac{6.7x}{100-x}$$

Another model produces the relationship

$$C(x) = \frac{6.5x}{102-x}$$

- (a) What is the cost found by averaging the two models? (*Hint:* The average of two quantities is half their sum.)
- (b) Using the two given models and the answer to part (a), find the cost to the nearest thousand dollars to remove 95% ($x = 95$) of the pollutant.
- (c) Average the two costs in part (b) from the given models, to the nearest thousand dollars. Compare this result with the cost obtained using the average of the two models.



6.3 Complex Fractions

OBJECTIVES

- 1 Simplify complex fractions by simplifying the numerator and denominator (Method 1).
- 2 Simplify complex fractions by multiplying by a common denominator (Method 2).
- 3 Compare the two methods of simplifying complex fractions.
- 4 Simplify rational expressions with negative exponents.

A **complex fraction** is a quotient having a fraction in the numerator, denominator, or both.

$$\frac{\frac{2}{3}}{\frac{4}{9}}, \quad 1 + \frac{1}{x}, \quad \frac{\frac{4}{y}}{6 - \frac{3}{y}}, \quad \text{and} \quad \frac{\frac{m^2 - 9}{m + 1}}{\frac{m + 3}{m^2 - 1}} \quad \text{Complex fractions}$$

OBJECTIVE 1 Simplify complex fractions by simplifying the numerator and denominator (Method 1).

Simplifying a Complex Fraction (Method 1)

- Step 1** Simplify the numerator and denominator separately.
- Step 2** Divide by multiplying the numerator by the reciprocal of the denominator.
- Step 3** Simplify the resulting fraction if possible.

In Step 2, we are treating the complex fraction as a quotient of two rational expressions and dividing. *In order for us to perform this step, both the numerator and the denominator must be single fractions.*

VOCABULARY

□ complex fraction

$$\begin{array}{ccc} \frac{q-5}{8} \leftarrow \text{Single fraction} & \left| \quad \frac{1}{x} + x \leftarrow \text{Not a single fraction} \quad \right| & \frac{6 + \frac{3}{x}}{4 + \frac{7}{8}} \leftarrow \text{Not a single fraction} \\ \frac{q+5}{3} \leftarrow \text{Single fraction} & \left| \quad \frac{x^2+1}{8} \leftarrow \text{Single fraction} \quad \right| & \frac{x}{4} + \frac{7}{8} \leftarrow \text{Not a single fraction} \end{array}$$

EXAMPLE 1 Simplifying Complex Fractions (Method 1)

Use Method 1 to simplify each complex fraction.

(a)
$$\frac{\frac{x+1}{x}}{\frac{x-1}{2x}}$$
 Both the numerator and the denominator are single fractions. Each is already simplified. (Step 1)

$$= \frac{x+1}{x} \div \frac{x-1}{2x}$$
 Write as a division problem.
$$= \frac{x+1}{x} \cdot \frac{2x}{x-1}$$
 Multiply by the reciprocal of $\frac{x-1}{2x}$. (Step 2)
$$= \frac{2x(x+1)}{x(x-1)}$$
 Multiply.
$$= \frac{2(x+1)}{x-1}$$
 Simplify. (Step 3)


NOW TRY EXERCISE 1

Use Method 1 to simplify each complex fraction.

(a) $\frac{\frac{t+4}{3t}}{\frac{2t+1}{9t}}$ (b) $\frac{5 - \frac{2}{y}}{4 + \frac{1}{y}}$

(b)
$$\begin{aligned} & \frac{2 + \frac{1}{y}}{3 - \frac{2}{y}} \\ &= \frac{\frac{2y}{y} + \frac{1}{y}}{\frac{3y}{y} - \frac{2}{y}} \\ &= \frac{2y + 1}{3y - 2} \\ &= \frac{2y + 1}{y} \div \frac{3y - 2}{y} \\ &= \frac{2y + 1}{y} \cdot \frac{y}{3y - 2} \\ &= \frac{2y + 1}{3y - 2} \end{aligned}$$

The numerator is a single fraction. So is the denominator.

The numerator and denominator are *not* single fractions. Simplify them separately. (Step 1)

Prepare to write the numerator and denominator as single fractions.

In the numerator, $\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$.

In the denominator, $\frac{a}{c} - \frac{b}{c} = \frac{a-b}{c}$.

Write as a division problem.

Multiply by the reciprocal of $\frac{3y-2}{y}$. (Step 2)

Multiply and simplify. (Step 3)

NOW TRY 

OBJECTIVE 2 Simplify complex fractions by multiplying by a common denominator (Method 2).

This method uses the identity property for multiplication.


Simplifying a Complex Fraction (Method 2)

Step 1 Multiply the numerator and denominator of the complex fraction by the least common denominator of the fractions in the numerator and the fractions in the denominator of the complex fraction.

Step 2 Simplify the resulting fraction if possible.

EXAMPLE 2 Simplifying Complex Fractions (Method 2)

Use Method 2 to simplify each complex fraction.

(a) $\frac{2 + \frac{1}{y}}{3 - \frac{2}{y}}$ 

This is the same fraction as in Example 1(b). Compare the solution methods.

$$= \frac{\left(2 + \frac{1}{y}\right) \cdot y}{\left(3 - \frac{2}{y}\right) \cdot y}$$

The LCD of all the fractions is y . Multiply the numerator and denominator by y because $\frac{y}{y} = 1$. (Step 1)

$$= \frac{2 \cdot y + \frac{1}{y} \cdot y}{3 \cdot y - \frac{2}{y} \cdot y}$$

Distributive property (Step 2)

$$= \frac{2y + 1}{3y - 2}$$

Multiply. We obtain the same answer as in Example 1(b).

NOW TRY ANSWERS

1. (a) $\frac{3(t+4)}{2t+1}$ (b) $\frac{5y-2}{4y+1}$


NOW TRY EXERCISE 2

Use Method 2 to simplify each complex fraction.

$$(a) \frac{5 - \frac{2}{y}}{4 + \frac{1}{y}} \quad (b) \frac{x + \frac{5}{x}}{4x - \frac{1}{x-2}}$$

$$(b) \frac{2p + \frac{5}{p-1}}{3p - \frac{2}{p}}$$

$$= \frac{\left(2p + \frac{5}{p-1}\right) \cdot p(p-1)}{\left(3p - \frac{2}{p}\right) \cdot p(p-1)}$$

$$= \frac{2p[p(p-1)] + \frac{5}{p-1} \cdot p(p-1)}{3p[p(p-1)] - \frac{2}{p} \cdot p(p-1)}$$

$$= \frac{2p[p(p-1)] + 5p}{3p[p(p-1)] - 2(p-1)}$$

$$= \frac{2p[p^2 - p] + 5p}{3p[p^2 - p] - 2p + 2}$$

$$= \frac{2p^3 - 2p^2 + 5p}{3p^3 - 3p^2 - 2p + 2}$$

Multiply the numerator and denominator by the LCD, $p(p-1)$. (Step 1)

Distributive property (Step 2)

Multiply.

Distributive property

Distributive property again

This rational expression is in lowest terms.

NOW TRY 

OBJECTIVE 3 Compare the two methods of simplifying complex fractions.

Some students prefer one method over the other, while other students feel comfortable with both methods and rely on practice with many examples to determine which method they will use in a particular problem.

EXAMPLE 3 Simplifying Complex Fractions (Both Methods)

Use both Method 1 and Method 2 to simplify each complex fraction.

Method 1

$$(a) \frac{\frac{2}{x-3}}{\frac{5}{x^2-9}}$$

$$= \frac{\frac{2}{x-3}}{\frac{5}{(x-3)(x+3)}}$$

$$= \frac{2}{x-3} \div \frac{5}{(x-3)(x+3)}$$

$$= \frac{2}{x-3} \cdot \frac{(x-3)(x+3)}{5}$$

$$= \frac{2(x+3)}{5}$$

Method 2

$$(a) \frac{\frac{2}{x-3}}{\frac{5}{x^2-9}}$$

$$= \frac{\frac{2}{x-3}}{\frac{5}{(x-3)(x+3)}}$$

$$= \frac{\frac{2}{x-3} \cdot (x-3)(x+3)}{\frac{5}{(x-3)(x+3)} \cdot (x-3)(x+3)}$$

$$= \frac{2(x+3)}{5}$$

NOW TRY ANSWERS

2. (a) $\frac{5y-2}{4y+1}$
 (b) $\frac{x^3-2x^2+5x-10}{4x^3-8x^2-x}$

NOW TRY EXERCISE 3

Use both Method 1 and Method 2 to simplify each complex fraction.

(a) $\frac{\frac{1}{p-6}}{\frac{5}{p^2-36}}$ (b) $\frac{\frac{1}{m^2} - \frac{1}{n^2}}{\frac{1}{m} + \frac{1}{n}}$

Method 1

(b)
$$\frac{\frac{1}{x} + \frac{1}{y}}{\frac{1}{x^2} - \frac{1}{y^2}}$$

$$= \frac{\frac{y}{xy} + \frac{x}{xy}}{\frac{y^2}{x^2y^2} - \frac{x^2}{x^2y^2}}$$

$$= \frac{\frac{y+x}{xy}}{\frac{y^2-x^2}{x^2y^2}}$$

$$= \frac{y+x}{xy} \div \frac{y^2-x^2}{x^2y^2}$$

$$= \frac{y+x}{xy} \cdot \frac{x^2y^2}{(y-x)(y+x)}$$

$$= \frac{xy}{y-x}$$

Method 2

(b)
$$\frac{\frac{1}{x} + \frac{1}{y}}{\frac{1}{x^2} - \frac{1}{y^2}}$$

$$= \frac{\left(\frac{1}{x} + \frac{1}{y}\right) \cdot x^2y^2}{\left(\frac{1}{x^2} - \frac{1}{y^2}\right) \cdot x^2y^2}$$

$$= \frac{\left(\frac{1}{x}\right)x^2y^2 + \left(\frac{1}{y}\right)x^2y^2}{\left(\frac{1}{x^2}\right)x^2y^2 - \left(\frac{1}{y^2}\right)x^2y^2}$$

$$= \frac{xy^2 + x^2y}{y^2 - x^2}$$

$$= \frac{xy(y+x)}{(y+x)(y-x)}$$

$$= \frac{xy}{y-x}$$

NOW TRY 

OBJECTIVE 4 Simplify rational expressions with negative exponents.

We begin by rewriting the expressions with only positive exponents.

EXAMPLE 4 Simplifying Rational Expressions with Negative Exponents

Simplify each expression, using only positive exponents in the answer.

(a) $\frac{m^{-1} + p^{-2}}{2m^{-2} - p^{-1}}$

$a^{-n} = \frac{1}{a^n}$

$$= \frac{\frac{1}{m} + \frac{1}{p^2}}{\frac{2}{m^2} - \frac{1}{p}}$$

The base of $2m^{-2}$ is m , not $2m$:
 $2m^{-2} = \frac{2}{m^2}$.

Write with positive exponents.

$2m^{-2} = 2 \cdot m^{-2} = \frac{2}{1} \cdot \frac{1}{m^2} = \frac{2}{m^2}$

$$= \frac{m^2p^2\left(\frac{1}{m} + \frac{1}{p^2}\right)}{m^2p^2\left(\frac{2}{m^2} - \frac{1}{p}\right)}$$

Simplify by Method 2. Multiply the numerator and denominator by the LCD, m^2p^2 .

$$= \frac{m^2p^2 \cdot \frac{1}{m} + m^2p^2 \cdot \frac{1}{p^2}}{m^2p^2 \cdot \frac{2}{m^2} - m^2p^2 \cdot \frac{1}{p}}$$

Distributive property

$$= \frac{mp^2 + m^2}{2p^2 - m^2p}$$

Write in lowest terms.

NOW TRY ANSWERS

3. (a) $\frac{p+6}{5}$ (b) $\frac{n-m}{mn}$

NOW TRY EXERCISE 4

Simplify each expression, using only positive exponents in the answer.

(a) $\frac{r^{-2} - s^{-1}}{4r^{-1} + s^{-2}}$

(b) $\frac{2y^{-1} - 3y^{-2}}{y^{-2} + 3x^{-1}}$

NOW TRY ANSWERS

4. (a) $\frac{s^2 - r^2s}{4rs^2 + r^2}$ (b) $\frac{2xy - 3x}{x + 3y^2}$

(b) $\frac{x^{-2} - 2y^{-1}}{y - 2x^2}$

$= \frac{1}{x^2} - \frac{2}{y}$

The 2 does *not* go in the denominator of this fraction.

Write with positive exponents.

$= \frac{\left(\frac{1}{x^2} - \frac{2}{y}\right)x^2y}{(y - 2x^2)x^2y}$

Use Method 2. Multiply by the LCD, x^2y .

$= \frac{y - 2x^2}{(y - 2x^2)x^2y}$

Use the distributive property in the numerator.

$= \frac{1}{x^2y}$

Remember to write 1 in the numerator.

Write in lowest terms.

NOW TRY

6.3 Exercises

FOR EXTRA HELP

MyLab Math

Video solutions for select problems available in MyLab Math

1. **Concept Check** A(n) _____ fraction is a quotient that has a fraction in the _____, denominator, or _____.

2. **Concept Check** Find the slope of the line that passes through each pair of points. (*Hint:* This will involve simplifying complex fractions. Recall that slope $m = \frac{y_2 - y_1}{x_2 - x_1}$.)

(a) $\left(-\frac{5}{2}, \frac{1}{6}\right)$ and $\left(\frac{5}{3}, \frac{3}{8}\right)$ (b) $\left(-\frac{5}{6}, -\frac{1}{2}\right)$ and $\left(-\frac{1}{3}, -\frac{3}{2}\right)$

Concept Check Simplify.

3. $\frac{\frac{2}{3}}{\frac{3}{4}}$

4. $\frac{\frac{3}{5}}{\frac{5}{6}}$

5. $\frac{\frac{3}{4}}{\frac{5}{12}}$

6. $\frac{\frac{5}{6}}{\frac{4}{9}}$

7. $\frac{\frac{5}{9} - \frac{1}{3}}{\frac{2}{3} + \frac{1}{6}}$

8. $\frac{\frac{7}{8} - \frac{3}{2}}{-\frac{1}{4} - \frac{3}{8}}$

9. $\frac{2 - \frac{1}{4}}{\frac{5}{4} + 3}$

10. $\frac{\frac{4}{3} - 2}{1 - \frac{3}{8}}$

Use either method to simplify each complex fraction. See Examples 1–3.

11. $\frac{\frac{12}{x-1}}{\frac{6}{x}}$

12. $\frac{\frac{24}{t+4}}{\frac{6}{t}}$

13. $\frac{\frac{k+1}{2k}}{\frac{3k-1}{4k}}$

14. $\frac{\frac{1-r}{4r}}{\frac{1+r}{8r}}$

15. $\frac{\frac{4z^2x^4}{9}}{\frac{12x^2z^5}{15}}$

16. $\frac{\frac{3y^2x^3}{8}}{\frac{9y^3x^4}{16}}$

17. $\frac{\frac{1}{x} + 1}{-\frac{1}{x} + 1}$

18. $\frac{\frac{2}{k} - 1}{\frac{2}{k} + 1}$

19. $\frac{6 + \frac{1}{x}}{7 - \frac{3}{x}}$

20. $\frac{4 - \frac{1}{p}}{9 + \frac{5}{p}}$

21. $\frac{\frac{3}{x} + \frac{3}{y}}{\frac{3}{x} - \frac{3}{y}}$

22. $\frac{\frac{4}{t} - \frac{4}{s}}{\frac{4}{t} + \frac{4}{s}}$

23.
$$\frac{\frac{8x - 24y}{10}}{\frac{x - 3y}{5x}}$$

24.
$$\frac{\frac{20x - 10y}{12y}}{\frac{2x - y}{6y^2}}$$

25.
$$\frac{\frac{6}{y - 4}}{\frac{12}{y^2 - 16}}$$

26.
$$\frac{\frac{8}{t + 7}}{\frac{24}{t^2 - 49}}$$

27.
$$\frac{\frac{x^2 - 16y^2}{xy}}{\frac{1}{y} - \frac{4}{x}}$$

28.
$$\frac{\frac{4t^2 - 9s^2}{st}}{\frac{2}{s} - \frac{3}{t}}$$

29.
$$\frac{\frac{1}{b^2} - \frac{1}{a^2}}{\frac{1}{b} - \frac{1}{a}}$$

30.
$$\frac{\frac{1}{x^2} - \frac{1}{y^2}}{\frac{1}{x} + \frac{1}{y}}$$

31.
$$\frac{\frac{x + y}{\frac{1}{y} + \frac{1}{x}}}{\frac{1}{y} + \frac{1}{x}}$$

32.
$$\frac{\frac{s - r}{\frac{1}{r} - \frac{1}{s}}}{\frac{1}{r} - \frac{1}{s}}$$

33.
$$\frac{\frac{y - y - 3}{3}}{\frac{4}{9} + \frac{2}{3y}}$$

34.
$$\frac{p - \frac{p + 2}{4}}{\frac{3}{4} - \frac{5}{2p}}$$

35.
$$\frac{\frac{1}{x^3} - \frac{1}{y^3}}{\frac{1}{x^2} - \frac{1}{y^2}}$$

36.
$$\frac{\frac{1}{y^3} + \frac{1}{x^3}}{\frac{1}{y^2} - \frac{1}{x^2}}$$

37.
$$\frac{\frac{x + 2}{x} + \frac{1}{x + 2}}{\frac{5}{x} + \frac{x}{x + 2}}$$

38.
$$\frac{\frac{y + 3}{y} - \frac{4}{y - 1}}{\frac{y}{y - 1} + \frac{1}{y}}$$

39. **Concept Check** When simplifying the expression

$$\frac{1}{x^{-2} + y^{-2}},$$

a student noted the negative exponents in the denominator and wrote his answer as $x^2 + y^2$. This is incorrect.

WHAT WENT WRONG?

40. **Concept Check** A student simplifying the expression

$$\frac{a^{-2} - 4b^{-2}}{3b - 6a} \text{ wrote } \frac{\frac{1}{a^2} - \frac{1}{4b^2}}{3b - 6a}$$

as her first step. This is a common error.

WHAT WENT WRONG?

Simplify each expression, using only positive exponents in the answer. See Example 4.

41.
$$\frac{1}{x^{-2} + y^{-2}}$$

42.
$$\frac{1}{p^{-2} - q^{-2}}$$

43.
$$\frac{x^{-2} + y^{-2}}{x^{-1} + y^{-1}}$$

44.
$$\frac{x^{-1} - y^{-1}}{x^{-2} - y^{-2}}$$

45.
$$\frac{k^{-1} + p^{-2}}{k^{-1} - 3p^{-2}}$$

46.
$$\frac{x^{-2} - y^{-1}}{x^{-2} + 4y^{-1}}$$

47.
$$\frac{x^{-1} + 2y^{-1}}{2y + 4x}$$

48.
$$\frac{a^{-2} - 4b^{-2}}{3b - 6a}$$

RELATING CONCEPTS For Individual or Group Work (Exercises 49–54)

Simplifying a complex fraction by Method 1 is a good way to review the methods of adding, subtracting, multiplying, and dividing rational expressions. Method 2 gives a good review of the fundamental property of rational numbers.

Refer to the following complex fraction, and work Exercises 49–54 in order.

$$\frac{\frac{4}{m} + \frac{m + 2}{m - 1}}{\frac{m + 2}{m} - \frac{2}{m - 1}}$$

49. Add the fractions in the numerator.

50. Subtract as indicated in the denominator.

51. Divide the answer from Exercise 49 by the answer from Exercise 50.

52. Go back to the original complex fraction and find the LCD of all denominators.

53. Multiply the numerator and denominator of the complex fraction by the answer from Exercise 52.

54. The answers for Exercises 51 and 53 should be the same. Which method do you prefer? Explain why.

6.4

Equations with Rational Expressions and Graphs

OBJECTIVES

- 1 Determine the domain of the variable in a rational equation.
- 2 Solve rational equations.
- 3 Recognize the graph of a rational function.

VOCABULARY

- domain of the variable in a rational equation
- proposed solution
- extraneous solution
- discontinuous
- reciprocal function
- vertical asymptote
- horizontal asymptote


NOW TRY EXERCISE 1

Give the domain of the variable in each equation.

(a) $\frac{1}{3x} - \frac{3}{4x} = \frac{1}{3}$

(b) $\frac{1}{x-7} + \frac{2}{x+7} = \frac{14}{x^2-49}$

NOW TRY ANSWERS

1. (a) $\{x \mid x \text{ is a real number, } x \neq 0\}$
- (b) $\{x \mid x \text{ is a real number, } x \neq \pm 7\}$

OBJECTIVE 1 Determine the domain of the variable in a rational equation.

The **domain of the variable in a rational equation** is the intersection (overlap) of the domains of the rational expressions in the equation. When we solve rational equations, knowing the domain of the variable enables us to determine whether a **proposed solution**—that is, a value of the variable that *appears* to be a solution—must be discarded.

EXAMPLE 1 Determining Domains of Variables

Give the domain of the variable in each equation.

(a) $\frac{2}{x} - \frac{3}{2} = \frac{7}{2x}$

The domains of the expressions $\frac{2}{x}$, $\frac{3}{2}$, and $\frac{7}{2x}$ in the equation are, in order,

$$\{x \mid x \text{ is a real number, } x \neq 0\},$$

$$\{x \mid x \text{ is a real number}\},$$

and

$$\{x \mid x \text{ is a real number, } x \neq 0\}.$$

Set-builder notation
(We could use interval notation.)

The intersection of these three domains is $\{x \mid x \text{ is a real number, } x \neq 0\}$.

(b) $\frac{2}{x-3} - \frac{3}{x+3} = \frac{12}{x^2-9}$

The domains of the three expressions are, respectively,

$$\{x \mid x \text{ is a real number, } x \neq 3\},$$

$$\{x \mid x \text{ is a real number, } x \neq -3\},$$

and

$$\{x \mid x \text{ is a real number, } x \neq \pm 3\}.$$

\pm is read "positive or negative," or "plus or minus."

The domain of the variable is the intersection of the three domains, all real numbers except 3 and -3 .

$$\{x \mid x \text{ is a real number, } x \neq \pm 3\}$$

NOW TRY 

OBJECTIVE 2 Solve rational equations.

To solve rational equations, we usually multiply all terms by the least common denominator to clear the fractions. **We can do this only with an equation** because we want to solve for a variable (not with an expression, where we are carrying out operations).

Solving an Equation with Rational Expressions

Step 1 Determine the domain of the variable.

Step 2 Multiply each side of the equation by the LCD to clear the fractions.

Step 3 Solve the resulting equation.

Step 4 Check that each proposed solution is in the domain, and discard any values that are not. Check the remaining proposed solution(s) in the original equation.


NOW TRY EXERCISE 2

Solve.

$$\frac{1}{3x} - \frac{3}{4x} = \frac{1}{3}$$

EXAMPLE 2 Solving a Rational Equation

Solve $\frac{2}{x} - \frac{3}{2} = \frac{7}{2x}$.

Step 1 The domain, which excludes 0, was found in **Example 1(a)**.

Step 2 $2x\left(\frac{2}{x} - \frac{3}{2}\right) = 2x\left(\frac{7}{2x}\right)$ Multiply by the LCD, $2x$.

Step 3 $2x\left(\frac{2}{x}\right) - 2x\left(\frac{3}{2}\right) = 2x\left(\frac{7}{2x}\right)$ Distributive property

$$4 - 3x = 7$$
 Multiply.

$$-3x = 3$$
 Subtract 4.

This proposed solution is in the domain.

$$x = -1$$
 Divide by -3 .

Step 4 CHECK $\frac{2}{x} - \frac{3}{2} = \frac{7}{2x}$ Original equation

Don't forget this step.

$$\frac{2}{-1} - \frac{3}{2} \stackrel{?}{=} \frac{7}{2(-1)}$$
 Let $x = -1$.

$$-\frac{7}{2} = -\frac{7}{2} \quad \checkmark$$
 True

 A true statement results, so the solution set is $\{-1\}$.

NOW TRY 

CAUTION When each side of an equation is multiplied by a *variable* expression, the resulting “solutions” may not satisfy the original equation. **We must either determine and observe the domain or check all proposed solutions in the original equation. It is wise to do both.**

EXAMPLE 3 Solving a Rational Equation with No Solution

Solve $\frac{2}{x-3} - \frac{3}{x+3} = \frac{12}{x^2-9}$.

Step 1 The domain, which excludes ± 3 , was found in **Example 1(b)**.

Step 2 Factor $x^2 - 9$. Then multiply each side by the LCD, $(x+3)(x-3)$.

$$(x+3)(x-3)\left(\frac{2}{x-3} - \frac{3}{x+3}\right) = (x+3)(x-3)\left[\frac{12}{(x+3)(x-3)}\right]$$

Step 3 $(x+3)(x-3)\left(\frac{2}{x-3}\right) - (x+3)(x-3)\left(\frac{3}{x+3}\right)$

$$= (x+3)(x-3)\left[\frac{12}{(x+3)(x-3)}\right]$$

Distributive property

$$(x+3)(2) - (x-3)(3) = 12$$
 Multiply.

$$2x + 6 - 3x + 9 = 12$$
 Distributive property

$$-x + 15 = 12$$
 Combine like terms.

$$\text{Proposed solution} \rightarrow x = 3$$
 Subtract 15. Multiply by -1 .

NOW TRY ANSWER

2. $\left\{-\frac{5}{4}\right\}$


NOW TRY EXERCISE 3

Solve.

$$\frac{1}{t-7} + \frac{2}{t+7} = \frac{14}{t^2-49}$$

Step 4 The proposed solution, 3, is not in the domain. Such solutions are **extraneous** and must be rejected. Substituting 3 into the original equation shows why.

CHECK $\frac{2}{x-3} - \frac{3}{x+3} = \frac{12}{x^2-9}$ Original equation

$\frac{2}{3-3} - \frac{3}{3+3} \stackrel{?}{=} \frac{12}{3^2-9}$ Let $x = 3$.

$\frac{2}{0} - \frac{3}{6} \stackrel{?}{=} \frac{12}{0}$ Division by 0 is undefined.

The equation has no solution. The solution set is \emptyset .
NOW TRY 

NOW TRY EXERCISE 4

Solve.

$$\frac{2}{t^2-2t-8} - \frac{4}{t^2+6t+8} = \frac{2}{t^2-16}$$

EXAMPLE 4 Solving a Rational Equation

Solve $\frac{3}{p^2+p-2} - \frac{1}{p^2-1} = \frac{7}{2(p^2+3p+2)}$.

Factor each denominator to find the domain and the LCD.

$$\frac{3}{(p-1)(p+2)} - \frac{1}{(p+1)(p-1)} = \frac{7}{2(p+2)(p+1)} \quad \text{Factor the denominators.}$$

The domain excludes 1, -2, and -1. Multiply each side of the equation by the LCD, $2(p-1)(p+2)(p+1)$.

$$2(p-1)(p+2)(p+1) \left[\frac{3}{(p-1)(p+2)} - \frac{1}{(p+1)(p-1)} \right] = 2(p-1)(p+2)(p+1) \left[\frac{7}{2(p+2)(p+1)} \right]$$

$$2(p+1)(3) - 2(p+2) = (p-1)7 \quad \text{Distributive property}$$

$$6p + 6 - 2p - 4 = 7p - 7 \quad 2 \cdot 3 = 6; \text{ Distributive property}$$

$$4p + 2 = 7p - 7 \quad \text{Combine like terms.}$$

$$9 = 3p \quad \text{Subtract } 4p. \text{ Add } 7.$$

$$\text{Proposed solution} \rightarrow 3 = p \quad \text{Divide by } 3.$$

CHECK 3 is in the domain. Substitute it in the original equation to check.

$$\frac{3}{p^2+p-2} - \frac{1}{p^2-1} = \frac{7}{2(p^2+3p+2)} \quad \text{Original equation}$$

$$\frac{3}{3^2+3-2} - \frac{1}{3^2-1} \stackrel{?}{=} \frac{7}{2(3^2+3 \cdot 3+2)} \quad \text{Let } p = 3.$$

$$\frac{\frac{3}{10} - \frac{1}{8}}{\frac{12}{40} - \frac{5}{40}} = \frac{3}{10} - \frac{1}{8} \stackrel{?}{=} \frac{7}{40} \quad \text{Work in the denominators.}$$

$$\frac{7}{40} = \frac{7}{40} \quad \checkmark \quad \text{True}$$

NOW TRY ANSWERS

3. \emptyset
4. $\{5\}$

The solution set is $\{3\}$.
NOW TRY 


NOW TRY EXERCISE 5

Solve.

$$\frac{2x}{x-2} = \frac{-3}{x} + \frac{4}{x-2}$$

EXAMPLE 5 Solving a Rational Equation

Solve $\frac{2}{3x+1} = \frac{1}{x} - \frac{6x}{3x+1}$. We must exclude $-\frac{1}{3}$ and 0 from the domain.

$$x(3x+1)\left(\frac{2}{3x+1}\right) = x(3x+1)\left(\frac{1}{x} - \frac{6x}{3x+1}\right) \quad \text{Multiply by the LCD, } x(3x+1).$$

$$x(3x+1)\left(\frac{2}{3x+1}\right) = x(3x+1)\left(\frac{1}{x}\right) - x(3x+1)\left(\frac{6x}{3x+1}\right)$$

Distributive property

This is a quadratic equation.

$$2x = 3x + 1 - 6x^2$$

Multiply.

$$6x^2 - x - 1 = 0$$

Standard form

$$(3x+1)(2x-1) = 0$$

Factor.

$$3x+1 = 0 \quad \text{or} \quad 2x-1 = 0$$

Zero-factor property

Proposed solutions $\rightarrow x = -\frac{1}{3}$ or $x = \frac{1}{2}$

Solve each equation.

Because $-\frac{1}{3}$ is not in the domain, it is not a solution. Check the proposed solution $\frac{1}{2}$ in the original equation. The solution set is $\left\{\frac{1}{2}\right\}$.

NOW TRY 

NOW TRY EXERCISE 6

Solve.

$$\frac{4}{x} + \frac{3}{x-6} = 2$$

EXAMPLE 6 Solving a Rational Equation

Solve $\frac{5}{x} + \frac{6}{x+2} = 1$. We must exclude 0 and -2 from the domain.

$$x(x+2)\left(\frac{5}{x} + \frac{6}{x+2}\right) = x(x+2)(1) \quad \text{Multiply by the LCD, } x(x+2).$$

$$x(x+2)\left(\frac{5}{x}\right) + x(x+2)\left(\frac{6}{x+2}\right) = x(x+2)(1) \quad \text{Distributive property}$$

$$(x+2)(5) + 6x = x^2 + 2x \quad \text{Multiply.}$$

$$5x + 10 + 6x = x^2 + 2x \quad \text{Distributive property}$$

$$x^2 - 9x - 10 = 0 \quad \text{Standard form}$$

$$(x+1)(x-10) = 0 \quad \text{Factor.}$$

$$x+1 = 0 \quad \text{or} \quad x-10 = 0 \quad \text{Zero-factor property}$$

$$x = -1 \quad \text{or} \quad x = 10 \quad \text{Solve each equation.}$$

Both -1 and 10 are in the domain. Check that the solution set is $\{-1, 10\}$.

NOW TRY 
OBJECTIVE 3 Recognize the graph of a rational function.

A function defined by a quotient of polynomials is a **rational function**. Because one or more values of x may be excluded from the domain of a rational function, its graph is often **discontinuous**—that is, there will be one or more breaks in the graph.

One simple rational function is the **reciprocal function** $f(x) = \frac{1}{x}$. The domain of this function includes all real numbers except 0. Thus, this function pairs every real number except 0 with its reciprocal.

NOW TRY ANSWERS

5. $\left\{-\frac{3}{2}\right\}$

6. $\left\{\frac{3}{2}, 8\right\}$

Some ordered pairs that belong to the function $f(x) = \frac{1}{x}$ are listed in the table.

The closer negative values of x are to 0, the less (“more negative”) y is.

The closer positive values of x are to 0, the greater y is.

x	-3	-2	-1	-0.5	-0.25	-0.1	0.1	0.25	0.5	1	2	3
y	$-\frac{1}{3}$	$-\frac{1}{2}$	-1	-2	-4	-10	10	4	2	1	$\frac{1}{2}$	$\frac{1}{3}$

Plotting the points from the table, we obtain the graph in **FIGURE 2**.




FIGURE 2

Reciprocal function

$$f(x) = \frac{1}{x}$$

Domain: $\{x \mid x \text{ is a real number, } x \neq 0\}$

Range: $\{y \mid y \text{ is a real number, } y \neq 0\}$

Because the domain of $f(x) = \frac{1}{x}$ includes all real numbers except 0, there is no point on the graph where $x = 0$. The vertical line with equation $x = 0$ is a **vertical asymptote**. The horizontal line with equation $y = 0$ is a **horizontal asymptote**. The graph approaches these asymptotes but does not cross them.

NOW TRY EXERCISE 7

Graph the rational function. Give the equations of the vertical and horizontal asymptotes.

$$f(x) = \frac{1}{x+1}$$

EXAMPLE 7 Graphing a Rational Function

Graph the rational function. Give the equations of the vertical and horizontal asymptotes.

$$g(x) = \frac{-2}{x-3}$$

Some ordered pairs that belong to the function are listed in the table.

x	-2	-1	0	1	2	2.5	2.75	3.25	3.5	4	5	6	7
y	$\frac{2}{5}$	$\frac{1}{2}$	$\frac{2}{3}$	1	2	4	8	-8	-4	-2	-1	$-\frac{2}{3}$	$-\frac{1}{2}$

There is no point on the graph, shown in **FIGURE 3**, for $x = 3$, because 3 is excluded from the domain. The dashed line

$x = 3$ represents the vertical asymptote.

It is not part of the graph. The graph gets closer to the vertical asymptote as the x -values get closer to 3. Again,

$y = 0$ is a horizontal asymptote.





FIGURE 3

NOW TRY

NOW TRY ANSWER



vertical asymptote: $x = -1$;
horizontal asymptote: $y = 0$


NOTE In general, the following statements are true of a rational function.

1. If the y -values approach ∞ or $-\infty$ as the x -values approach a real number a , the **vertical line $x = a$ is a vertical asymptote** of the graph.
2. If the y -values approach a real number b as $|x|$ increases without bound, the **horizontal line $y = b$ is a horizontal asymptote** of the graph.

6.4 Exercises

FOR
EXTRA
HELP

 MyLab Math

 Video solutions for select problems available in MyLab Math

1. **Concept Check** Decide whether each of the following is an *expression* or an *equation*.

(a) $\frac{2}{x} = \frac{4}{3x} + \frac{1}{3}$

(b) $\frac{4}{3x} + \frac{1}{3}$

(c) $\frac{5}{x+1} - \frac{2}{x-1}$

(d) $\frac{5}{x+1} - \frac{2}{x-1} = \frac{4}{x^2-1}$

2. **Concept Check** The following instruction is incorrect.

“Solve $\frac{2x+1}{3x-4} + \frac{1}{2x+3}$.”

WHAT WENT WRONG?

3. **CONCEPT CHECK** A student solved

$$\frac{3}{x-3} - \frac{2}{x-2} = \frac{3}{x^2-5x+6}$$

performing all algebraic steps correctly, and obtained $x = 3$ as the final step. She gave the solution set $\{3\}$, but this is incorrect. **WHAT WENT WRONG?** Give the correct solution set.

4. **Concept Check** Consider the following incorrect “solution.”

$$x - \frac{9}{x} = -8 \quad \text{Multiply by the LCD, } x.$$

$$x - 9 = -8x$$

$$9x = 9$$

$$x = 1$$

Solution set: $\{1\}$

WHAT WENT WRONG? Give the correct solution set.

Give the domain of the variable in each equation. See Example 1.

5. $\frac{1}{3x} + \frac{1}{2x} = \frac{x}{3}$

6. $\frac{5}{6x} - \frac{8}{2x} = \frac{x}{4}$

7. $\frac{1}{x+1} - \frac{1}{x-2} = 0$

8. $\frac{3}{x+4} - \frac{2}{x-9} = 0$

9. $\frac{3x+1}{x-4} = \frac{6x+5}{2x-7}$

10. $\frac{4x-1}{2x+3} = \frac{12x-25}{6x-2}$

11. $\frac{6}{4x+7} - \frac{3}{x} = \frac{5}{6x-13}$

12. $\frac{4}{3x-5} + \frac{2}{x} = \frac{9}{4x+13}$

13. $\frac{1}{x^2-16} - \frac{2}{x-4} = \frac{1}{x+4}$

14. $\frac{2}{x^2-25} - \frac{1}{x+5} = \frac{1}{x-5}$

15. $\frac{2}{x^2-x} + \frac{1}{x+3} = \frac{4}{x-2}$

16. $\frac{3}{x^2+x} - \frac{1}{x+5} = \frac{2}{x-7}$

Solve each equation. See Examples 2–6.

17. $\frac{3}{4x} = \frac{5}{2x} - \frac{7}{4}$

18. $\frac{2}{3x} = \frac{6}{5x} + \frac{8}{45}$

19. $x - \frac{24}{x} = -2$

20. $p + \frac{15}{p} = -8$

21. $\frac{x}{4} - \frac{21}{4x} = -1$

22. $\frac{x}{2} - \frac{12}{x} = 1$

23. $\frac{x-4}{x+6} = \frac{2x+3}{2x-1}$

24. $\frac{5x-3}{x+2} = \frac{5x-1}{x+3}$

25. $\frac{3x+1}{x-4} = \frac{6x+5}{2x-7}$

26. $\frac{4x-1}{2x+3} = \frac{12x-25}{6x-2}$

27. $\frac{1}{y-1} + \frac{5}{12} = \frac{-2}{3y-3}$

28. $\frac{4}{m+2} - \frac{11}{9} = \frac{1}{3m+6}$

29. $\frac{-2}{3t-6} - \frac{1}{36} = \frac{-3}{4t-8}$

30. $\frac{6}{t+1} - \frac{34}{15} = \frac{-4}{5t+5}$

31. $\frac{7}{6x+3} - \frac{1}{3} = \frac{2}{2x+1}$

32. $\frac{3}{4m+2} = \frac{17}{2} - \frac{7}{2m+1}$

33. $\frac{9}{x} + \frac{4}{6x-3} = \frac{2}{6x-3}$

34. $\frac{5}{n} + \frac{4}{6-3n} = \frac{2}{6-3n}$

35. $\frac{6}{x-4} + \frac{5}{x} = \frac{-20}{x^2-4x}$

36. $\frac{7}{x-4} + \frac{3}{x} = \frac{-12}{x^2-4x}$

37. $\frac{3}{x+2} - \frac{2}{x^2-4} = \frac{1}{x-2}$

38. $\frac{3}{x-2} + \frac{21}{x^2-4} = \frac{14}{x+2}$

39. $\frac{1}{y+2} + \frac{3}{y+7} = \frac{5}{y^2+9y+14}$

40. $\frac{1}{t+3} + \frac{4}{t+5} = \frac{2}{t^2+8t+15}$

41. $\frac{6}{w+3} + \frac{-7}{w-5} = \frac{-48}{w^2-2w-15}$

42. $\frac{2}{r-5} + \frac{3}{2r+1} = \frac{22}{2r^2-9r-5}$

43. $\frac{2}{x} + \frac{4}{x+9} = 1$

44. $\frac{3}{x} - \frac{9}{x-8} = -2$

45. $\frac{2}{4x+7} + \frac{x}{3} = \frac{6}{12x+21}$

46. $\frac{3}{2x+5} + \frac{x}{2} = \frac{9}{6x+15}$

47. $\frac{1}{x-2} + \frac{1}{4} = \frac{1}{4(x^2-4)}$

48. $\frac{1}{x+4} + \frac{1}{3} = \frac{-10}{3(x^2-16)}$

49. $\frac{x}{x-3} + \frac{4}{x+3} = \frac{18}{x^2-9}$

50. $\frac{2x}{x-3} + \frac{4}{x+3} = \frac{-24}{x^2-9}$

51. $\frac{1}{x+4} + \frac{x}{x-4} = \frac{-8}{x^2-16}$

52. $\frac{5}{x-4} - \frac{3}{x-1} = \frac{x^2-1}{x^2-5x+4}$

53. $\frac{2}{k^2+k-6} + \frac{1}{k^2-k-2} = \frac{4}{k^2+4k+3}$

54. $\frac{5}{p^2+3p+2} - \frac{3}{p^2-4} = \frac{1}{p^2-p-2}$


55. $\frac{5x+14}{x^2-9} = \frac{-2x^2-5x+2}{x^2-9} + \frac{2x+4}{x-3}$

56. $\frac{4x-7}{4x^2-9} = \frac{-2x^2+5x-4}{4x^2-9} + \frac{x+1}{2x+3}$

57. **Concept Check** Match each function in Column I with its type in Column II.

- | I | II |
|---------------------------|-----------------------|
| (a) $f(x) = 3x - 4$ | A. quadratic function |
| (b) $f(x) = \frac{10}{x}$ | B. cubing function |
| (c) $f(x) = x^2 + 2x - 8$ | C. rational function |
| (d) $f(x) = x^3$ | D. linear function |

58. **Concept Check** Which graph has vertical and horizontal asymptotes? What are their equations?



Graph each rational function. Give the equations of the vertical and horizontal asymptotes. See Example 7.

59. $f(x) = \frac{2}{x}$ 60. $f(x) = \frac{3}{x}$ 61. $g(x) = -\frac{1}{x}$ 62. $g(x) = -\frac{2}{x}$
63. $f(x) = \frac{1}{x-2}$ 64. $f(x) = \frac{1}{x+2}$ 65. $g(x) = \frac{-2}{x-1}$ 66. $g(x) = \frac{-1}{x+3}$

Solve each problem.

67. The average number of vehicles waiting in line to enter a parking area is modeled by the function

$$W(x) = \frac{x^2}{2(1-x)},$$

where x is a quantity between 0 and 1 known as the **traffic intensity**. (Data from Mannering, F., and W. Kilareski, *Principles of Highway Engineering and Traffic Control*, John Wiley and Sons.) For each traffic intensity, find the average number of vehicles waiting (to the nearest tenth).

- (a) 0.1 (b) 0.8 (c) 0.9
- (d) What happens to waiting time as traffic intensity increases?



68. The percent of deaths caused by smoking is modeled by the rational function

$$P(x) = \frac{x-1}{x},$$


where x is the number of times more likely a smoker is to die of lung cancer than a non-smoker. This is the **incidence rate**. (Data from Walker, A., *Observation and Inference: An Introduction to the Methods of Epidemiology*, University of Michigan.) For example, $x = 10$ means that a smoker is 10 times more likely than a nonsmoker to die of lung cancer.

- (a) Find $P(x)$ if $x = 10$.
- (b) For what values of x is $P(x) = 80\%$? (Hint: Change 80% to a decimal.)
- (c) Can the incidence rate equal 0? Explain.

69. The force required to keep a 2000-lb car that is going 30 mph from skidding on a curve, where r is the radius of the curve in feet, is given by

$$F(r) = \frac{225,000}{r}.$$


- (a) What radius must a curve have if a force of 450 lb is needed to keep the car from skidding?
 (b) As the radius of the curve is lengthened, how is the force affected?



70. The amount of heating oil produced (in gallons per day) by an oil refinery is modeled by the rational function

$$f(x) = \frac{125,000 - 25x}{125 + 2x},$$

where x is the amount of gasoline produced (in hundreds of gallons per day). Suppose the refinery must produce 300 gal of heating oil per day to meet the needs of its customers.



- (a) How much gasoline will be produced per day?
 (b) A graph of f , for $x > 0$, is shown in the figure. Use it to decide what happens to the amount of gasoline (x) produced as the amount of heating oil (y) produced increases.

SUMMARY EXERCISES Simplifying Rational Expressions vs. Solving Rational Equations

A common student error is to confuse an equation, such as $\frac{x}{2} + \frac{x}{3} = -5$, with an expression involving an operation, such as $\frac{x}{2} + \frac{x}{3}$. **Equations are solved for a numerical answer, while problems involving operations result in simplified expressions.**

Solving an Equation

Solve: $\frac{x}{2} + \frac{x}{3} = -5$ **Look for the equality symbol.**

Multiply each side by the LCD, 6.

$$\begin{aligned} 6\left(\frac{x}{2} + \frac{x}{3}\right) &= 6(-5) \\ 6\left(\frac{x}{2}\right) + 6\left(\frac{x}{3}\right) &= 6(-5) \\ 3x + 2x &= -30 \\ 5x &= -30 \\ x &= -6 \end{aligned}$$

Check that the solution set is $\{-6\}$.

Simplifying an Expression Involving an Operation

Add: $\frac{x}{2} + \frac{x}{3}$ **There is no equality symbol.**

Write both fractions with the LCD, 6.

$$\begin{aligned} \frac{x}{2} + \frac{x}{3} &= \frac{x \cdot 3}{2 \cdot 3} + \frac{x \cdot 2}{3 \cdot 2} \\ &= \frac{3x}{6} + \frac{2x}{6} \\ &= \frac{3x + 2x}{6} \\ &= \frac{5x}{6} \end{aligned}$$

Identify each exercise as an expression or an equation. Then either simplify the expression by performing the indicated operation, or solve the equation, as appropriate.

1. $\frac{x}{2} - \frac{x}{4} = 5$

2. $\frac{4x - 20}{x^2 - 25} \cdot \frac{(x + 5)^2}{10}$

3. $\frac{6}{7x} - \frac{4}{x}$

4. $\frac{\frac{1}{x} + \frac{1}{y}}{\frac{1}{x} - \frac{1}{y}}$

5. $\frac{5}{7t} = \frac{52}{7} - \frac{3}{t}$

6. $\frac{x - 5}{3} + \frac{1}{3} = \frac{x - 2}{5}$

7. $\frac{7}{6x} + \frac{5}{8x}$

8. $\frac{4}{x} - \frac{8}{x + 1} = 0$

9. $\frac{\frac{6}{x + 1} - \frac{1}{x}}{\frac{2}{x} - \frac{4}{x + 1}}$

10. $\frac{8}{r + 2} - \frac{7}{4r + 8}$

11. $\frac{x}{x + y} + \frac{2y}{x - y}$

12. $\frac{3p^2 - 6p}{p + 5} \div \frac{p^2 - 4}{8p + 40}$

13. $\frac{x - 2}{9} \cdot \frac{5}{8 - 4x}$

14. $\frac{a - 4}{3} + \frac{11}{6} = \frac{a + 1}{2}$

15. $\frac{5}{x^2 - 2x} - \frac{3}{x^2 - 4}$

16. $\frac{3}{t - 1} + \frac{1}{t} = \frac{7}{2}$

17. $\frac{\frac{5}{x} - \frac{3}{y}}{\frac{9x^2 - 25y^2}{x^2y}}$

18. $\frac{\frac{t}{4} - \frac{1}{t}}{1 + \frac{t + 4}{t}}$

19. $\frac{3r}{r - 2} = 1 + \frac{6}{r - 2}$

20. $\frac{7}{2x^2 - 8x} + \frac{3}{x^2 - 16}$

21. $\frac{b^2 + b - 6}{b^2 + 2b - 8} \cdot \frac{b^2 + 8b + 16}{3b + 12}$

22. $\frac{10z^2 - 5z}{3z^3 - 6z^2} \div \frac{2z^2 + 5z - 3}{z^2 + z - 6}$

23. $\frac{4y^2 - 13y + 3}{2y^2 - 9y + 9} \div \frac{4y^2 + 11y - 3}{6y^2 - 5y - 6}$

24. $\frac{-2}{a^2 + 2a - 3} - \frac{5}{3 - 3a} = \frac{4}{3a + 9}$

25. $\frac{-1}{3 - x} - \frac{2}{x - 3}$

26. $\frac{8}{3k + 9} - \frac{8}{15} = \frac{2}{5k + 15}$

27. $\frac{2}{y + 1} - \frac{3}{y^2 - y - 2} = \frac{3}{y - 2}$

28. $\frac{6z^2 - 5z - 6}{6z^2 + 5z - 6} \cdot \frac{12z^2 - 17z + 6}{12z^2 - z - 6}$

29. $\frac{3}{y - 3} - \frac{3}{y^2 - 5y + 6} = \frac{2}{y - 2}$

30. $\frac{2k + \frac{5}{k - 1}}{3k - \frac{2}{k}}$

6.5 Applications of Rational Expressions

OBJECTIVES

- 1 Find the value of an unknown variable in a formula.
- 2 Solve a formula for a specified variable.
- 3 Solve applications using proportions.
- 4 Solve applications about distance, rate, and time.
- 5 Solve applications about work rates.

VOCABULARY

- ratio
- proportion
- cross products of a proportion

NOW TRY EXERCISE 1

Use the formula in **Example 1** to find f if $p = 50$ cm and $q = 10$ cm.

NOW TRY EXERCISE 2

Solve for m .

$$\frac{1}{m} - \frac{2}{n} = 5$$

NOW TRY ANSWERS

1. $\frac{25}{3}$ cm
2. $m = \frac{n}{5n+2}$

OBJECTIVE 1 Find the value of an unknown variable in a formula.


Formulas may contain rational expressions, such as $t = \frac{d}{r}$ and $\frac{1}{f} = \frac{1}{p} + \frac{1}{q}$.

EXAMPLE 1 Finding the Value of a Variable in a Formula

In physics, the focal length f of a lens is given by the formula

$$\frac{1}{f} = \frac{1}{p} + \frac{1}{q}$$

In the formula, p is the distance from the object to the lens and q is the distance from the lens to the image. See **FIGURE 4**. Find q if $p = 20$ cm and $f = 10$ cm.



Focal Length of Camera Lens

FIGURE 4

$$\frac{1}{f} = \frac{1}{p} + \frac{1}{q}$$

$$\frac{1}{10} = \frac{1}{20} + \frac{1}{q}$$

Solve this equation for q .

Let $f = 10$, $p = 20$.

$$20q \cdot \frac{1}{10} = 20q \left(\frac{1}{20} + \frac{1}{q} \right)$$

Multiply by the LCD, $20q$.

$$20q \cdot \frac{1}{10} = 20q \left(\frac{1}{20} \right) + 20q \left(\frac{1}{q} \right)$$

Distributive property

$$2q = q + 20$$

Multiply.

$$q = 20$$

Subtract q .

The distance from the lens to the image is 20 cm.

NOW TRY

OBJECTIVE 2 Solve a formula for a specified variable.

We solve for a specified variable by isolating it on one side of the equality symbol.

EXAMPLE 2 Solving a Formula for a Specified Variable

Solve $\frac{1}{f} = \frac{1}{p} + \frac{1}{q}$ for p .

$$\frac{1}{f} = \frac{1}{p} + \frac{1}{q}$$

$$fpq \cdot \frac{1}{f} = fpq \left(\frac{1}{p} + \frac{1}{q} \right)$$

To clear the fractions, multiply by the LCD, fpq .

$$pq = fq + fp$$

Multiply; distributive property

We want the two terms with p on the same side.

$$pq - fp = fq$$

Subtract fp .

This is a key step.

$$p(q - f) = fq$$

Factor out p .

$$p = \frac{fq}{q - f}$$

Divide by $q - f$.

NOW TRY

NOW TRY
EXERCISE 3

Solve for N .

$$\frac{NR}{N-n} = t$$

EXAMPLE 3 Solving a Formula for a Specified Variable

Solve $I = \frac{nE}{R + nr}$ for n .

$$I = \frac{nE}{R + nr}$$

$$(R + nr)I = (R + nr) \frac{nE}{R + nr}$$

To clear the fraction, multiply by $R + nr$.

$$RI + nrI = nE$$

Distributive property on the left

Write the n -terms on the **same** side in preparation for factoring.

$$RI = nE - nrI$$

Subtract nrI .

$$RI = n(E - rI)$$

Factor out n .

$$\frac{RI}{E - rI} = n, \quad \text{or} \quad n = \frac{RI}{E - rI}$$

Divide by $E - rI$.
Interchange sides.

NOW TRY 

CAUTION Refer to the steps in **Examples 2 and 3** that factor out the desired variable. **This variable must be a factor on only one side of the equation.** Then each side can be divided by the remaining factor in the last step.

OBJECTIVE 3 Solve applications using proportions.

A **ratio** is a comparison of two quantities. The ratio of a to b may be written in any of the following ways.

$$a \text{ to } b, \quad a:b, \quad \text{or} \quad \frac{a}{b} \quad \text{Ratio of } a \text{ to } b$$

Ratios are usually written as quotients in algebra. A **proportion** is a statement that two ratios are equal.

$$\frac{a}{b} = \frac{c}{d} \quad \text{Proportion}$$

EXAMPLE 4 Solving a Proportion

In 2016, about 9 of every 100 Americans had no health insurance. The population at that time was about 323 million. How many million Americans had no health insurance? (Data from U.S. Census Bureau.)

Step 1 Read the problem.

Step 2 Assign a variable.

Let x = the number (in millions) who had no health insurance.

Step 3 Write an equation. We set up a proportion. The ratio 9 to 100 should equal the ratio x to 323.

$$\frac{9}{100} = \frac{x}{323} \quad \text{Write a proportion.}$$



NOW TRY ANSWER

3. $N = \frac{nt}{t-R}$


NOW TRY EXERCISE 4

In 2016, 18 of every 100 West Virginians lived in poverty. West Virginia's population was about 1,831,000 at that time. How many West Virginians lived in poverty? (Data from U.S. Census Bureau.)


NOW TRY EXERCISE 5

Clayton's SUV uses 28 gal of gasoline to drive 500 mi. He has 10 gal of gasoline in the SUV, and he still needs to drive 400 mi. If the car continues to use gasoline at the same rate, how many more gallons will he need?

Step 4 Solve. $32,300\left(\frac{9}{100}\right) = 32,300\left(\frac{x}{323}\right)$ *Multiply by a common denominator, $100 \cdot 323 = 32,300$.*

$$2907 = 100x$$

Simplify.

$$29.07 = x$$

Divide by 100.

Step 5 State the answer. 29.07 million Americans had no health insurance.

Step 6 Check that the ratio of 29.07 million to 323 million equals $\frac{9}{100}$.

$$\frac{29.07}{323} = \frac{9}{100}$$

Use a calculator to divide 29.07 by 323. A true statement results.

NOW TRY 

EXAMPLE 5 Solving a Proportion Involving Rates

Jodie's car uses 10 gal of gasoline to travel 210 mi. She has 5 gal of gasoline in the car. She still needs to drive 640 mi. If the car continues to use gasoline at the same rate, how many more gallons will she need?

Step 1 Read the problem.

Step 2 Assign a variable.

Let x = the additional number of gallons of gas needed.

Step 3 Write an equation. We set up a proportion.

$$\begin{array}{ccc} \text{gallons} \rightarrow & \frac{10}{210} & = \frac{5+x}{640} \leftarrow \text{gallons} \\ \text{miles} \rightarrow & & \leftarrow \text{miles} \end{array}$$

Step 4 Solve. We could multiply each side by the LCD $10 \cdot 21 \cdot 64$. Instead we use an alternative method that involves **cross products of a proportion**.

For $\frac{a}{b} = \frac{c}{d}$ to be true, the cross products ad and bc must be equal.

$$\begin{array}{ccc} a \cdot d = b \cdot c \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \end{array}$$

$$10 \cdot 640 = 210(5 + x) \quad \text{If } \frac{a}{b} = \frac{c}{d}, \text{ then } ad = bc.$$

$$6400 = 1050 + 210x \quad \text{Multiply; distributive property}$$

$$5350 = 210x \quad \text{Subtract 1050.}$$

$$25.5 = x \quad \text{Divide by 210. Round to the nearest tenth.}$$

Step 5 State the answer. Jodie will need 25.5 more gallons of gas.

Step 6 Check. 25.5 more gallons plus the 5 that she has equals 30.5 gal. From Step 3, the ratio of 10 to 210 must equal the ratio of 30.5 to 640.

$$\frac{10}{210} \approx 0.048 \quad \text{and} \quad \frac{30.5}{640} \approx 0.048 \quad \text{Use a calculator.}$$

Because the ratios are approximately equal, the answer is correct.

NOW TRY 

OBJECTIVE 4 Solve applications about distance, rate, and time.

When the distance formula $d = rt$ is solved for r or t , rational expressions result.

Rate is the ratio of distance to time, or $r = \frac{d}{t}$.

Time is the ratio of distance to rate, or $t = \frac{d}{r}$.

NOW TRY ANSWERS

4. 329,580 West Virginians
5. 12.4 more gallons


NOW TRY EXERCISE 6

A small fishing boat travels 36 mi against the current in a river in the same time that it travels 44 mi with the current. If the rate of the boat in still water is 20 mph, find the rate of the current.

EXAMPLE 6 Solving a Distance, Rate, Time Problem

A paddle wheeler travels 10 mi against the current in a river in the same time that it travels 15 mi with the current. If the rate of the current is 3 mph, find the rate of the boat in still water.



Step 1 Read the problem.

Step 2 Assign a variable. Let x = the rate of the boat in still water.

Traveling *against* the current slows the boat down, so the rate of the boat is the *difference* between its rate in still water and the rate of the current—that is, $(x - 3)$ mph.

Traveling *with* the current speeds the boat up, so the rate of the boat is the *sum* of its rate in still water and the rate of the current—that is, $(x + 3)$ mph.

Thus, $x - 3$ = the rate of the boat *against* the current,

and $x + 3$ = the rate of the boat *with* the current.

Because the time is the same going against the current as with the current, find time in terms of distance and rate for each situation. Against the current, the distance is 10 mi and the rate is $(x - 3)$ mph.

$$t = \frac{d}{r} = \frac{10}{x - 3} \quad \text{Time against the current}$$

With the current, the distance is 15 mi and the rate is $(x + 3)$ mph.

$$t = \frac{d}{r} = \frac{15}{x + 3} \quad \text{Time with the current}$$

	Distance	Rate	Time
Against Current	10	$x - 3$	$\frac{10}{x - 3}$
With Current	15	$x + 3$	$\frac{15}{x + 3}$

Summarize the information in a table.

Times are equal.

Step 3 Write an equation. Use the fact that the times are equal.

$$\underbrace{\frac{10}{x - 3}}_{\text{Time against current}} = \underbrace{\frac{15}{x + 3}}_{\text{time with current.}}$$

To solve, we can multiply by the LCD or use cross products.

Step 4 Solve.

$$\begin{aligned} (x + 3)(x - 3)\left(\frac{10}{x - 3}\right) &= (x + 3)(x - 3)\left(\frac{15}{x + 3}\right) && \text{Multiply by the LCD, } (x + 3)(x - 3). \\ (x + 3)10 &= (x - 3)15 && \text{Multiply.} \\ 10x + 30 &= 15x - 45 && \text{Distributive property} \\ 75 &= 5x && \text{Subtract } 10x. \text{ Add } 45. \\ 15 &= x && \text{Divide by } 5. \end{aligned}$$

Step 5 State the answer. The rate of the boat in still water is 15 mph.

Step 6 Check the answer: $\frac{10}{15 - 3} = \frac{15}{15 + 3}$ is true.

NOW TRY ANSWER

6. 2 mph

EXAMPLE 7 Solving a Distance, Rate, Time Problem

At O'Hare International Airport in Chicago, Cheryl and Bill are walking to the gate at the same rate to catch their flight to Denver. Bill wants a window seat, so he steps onto the moving sidewalk and continues to walk while Cheryl uses the stationary sidewalk. If the sidewalk moves at 1 m per sec and Bill saves 50 sec covering the 300-m distance, what is their walking rate?

Step 1 Read the problem. We must find their walking rate.

Step 2 Assign a variable.

Let x = their walking rate in meters per second.

Thus, Cheryl travels at a rate of x meters per second and Bill travels at a rate of $(x + 1)$ meters per second. Express their times in terms of the known distances and the variable rates, as in **Example 6**. Cheryl travels 300 m at a rate of x meters per second.

$$t = \frac{d}{r} = \frac{300}{x} \quad \text{Cheryl's time}$$

Bill travels 300 m at a rate of $(x + 1)$ meters per second.

$$t = \frac{d}{r} = \frac{300}{x + 1} \quad \text{Bill's time}$$

	Distance	Rate	Time
Cheryl	300	x	$\frac{300}{x}$
Bill	300	$x + 1$	$\frac{300}{x + 1}$

← Bill's time is 50 sec less than Cheryl's time.

Step 3 Write an equation. Use the times from the table.

$$\underbrace{\frac{300}{x + 1}}_{\text{Bill's time}} \text{ is } \underbrace{\frac{300}{x}}_{\text{Cheryl's time}} \underbrace{- 50}_{\text{less 50 sec}}$$

We *cannot* solve using cross products because there are two terms on the right.

Step 4 Solve.

$$x(x + 1)\left(\frac{300}{x + 1}\right) = x(x + 1)\left(\frac{300}{x} - 50\right) \quad \text{Multiply by the LCD, } x(x + 1).$$

$$x(x + 1)\left(\frac{300}{x + 1}\right) = x(x + 1)\left(\frac{300}{x}\right) - x(x + 1)(50) \quad \text{Distributive property}$$

$$300x = 300(x + 1) - 50x(x + 1) \quad \text{Multiply.}$$

$$300x = 300x + 300 - 50x^2 - 50x \quad \text{Distributive property}$$

$$50x^2 + 50x - 300 = 0 \quad \text{Standard form}$$

$$x^2 + x - 6 = 0 \quad \text{Divide by 50.}$$

$$(x + 3)(x - 2) = 0 \quad \text{Factor.}$$

$$x + 3 = 0 \quad \text{or} \quad x - 2 = 0 \quad \text{Zero-factor property}$$

$$x = -3 \quad \text{or} \quad x = 2 \quad \text{Solve each equation.}$$

Discard the negative solution because rate (speed) cannot be negative.


NOW TRY EXERCISE 7

James and Pat are driving from Atlanta to Jacksonville, a distance of 310 mi. James, whose average rate is 5 mph faster than Pat's, will drive the first 130 mi of the trip, and then Pat will drive the rest of the way to their destination. If the total driving time is 5 hr, determine the average rate of each driver.

Step 5 State the answer. Their walking rate is 2 m per sec.

Step 6 Check the solution in the words of the original problem.



CAUTION We can solve a rational equation using cross products only when there is a single rational expression on each side, such as in the equation in Example 6.

$$\frac{10}{x-3} = \frac{15}{x+3} \quad \text{To solve, we can multiply by the LCD or use cross products of the proportion.}$$

We cannot solve the rational equation in Example 7 using cross products.

$$\frac{300}{x+1} = \frac{300}{x} - 50 \quad \text{There is more than one term on the right. To solve, we must multiply by the LCD.}$$

OBJECTIVE 5 Solve applications about work rates.

PROBLEM-SOLVING HINT If the letters r , t , and A represent the rate at which work is done, the time required, and the amount of work accomplished, respectively, then $A = rt$. Notice the similarity to the distance formula, $d = rt$.

Amount of work can be measured in terms of jobs accomplished. Thus, if 1 job is completed, then $A = 1$, and the formula gives the rate as

$$1 = rt, \quad \text{or} \quad r = \frac{1}{t}.$$

To solve a work problem, we use the following fact to express all rates of work.

Rate of Work

If 1 complete job can be accomplished in t units of time, then the rate of work is

$$\frac{1}{t} \text{ job per unit of time.}$$

EXAMPLE 8 Solving a Work Problem

Letitia and Kareem are working on a neighborhood cleanup. Kareem can clean up all the trash in the area in 7 hr, while Letitia can do the same job in 5 hr. How long will it take them if they work together?

Steps 1 and 2 Read the problem, and assign a variable.

Let x = the number of hours it will take the two people working together.

We use the formula $A = rt$. Because $A = 1$, the rate for each person will be $\frac{1}{t}$, where t is the time it takes the person to complete the job alone. Kareem can clean up all the trash in 7 hr, so his rate is $\frac{1}{7}$ of the job per hour. Letitia's rate is $\frac{1}{5}$ of the job per hour.

	Rate	Time Working Together	Fractional Part of the Job Done
Kareem	$\frac{1}{7}$	x	$\frac{1}{7}x$
Letitia	$\frac{1}{5}$	x	$\frac{1}{5}x$

NOW TRY ANSWER

7. 60 mph; 65 mph




NOW TRY EXERCISE 8

Using her zero-turn riding lawn mower, Tracy can mow her lawn in 2 hr. Using a traditional mower, Matt can mow the same lawn in 3 hr. How long will it take them to mow the lawn working together?

Step 3 Write an equation.

$$\underbrace{\frac{1}{7}x}_{\text{Part done by Kareem}} + \underbrace{\frac{1}{5}x}_{\text{part done by Letitia}} = \underbrace{1}_{\text{1 whole job.}}$$

Together they complete 1 job. The sum of the fractional parts must equal 1.

Step 4 Solve.

$$35\left(\frac{1}{7}x + \frac{1}{5}x\right) = 35 \cdot 1$$

Multiply by the LCD, 35.

$$5x + 7x = 35$$

(*) Distributive property

$$12x = 35$$

Combine like terms.

$$x = \frac{35}{12}$$

Divide by 12.

Step 5 State the answer. Kareem and Letitia can do the entire job working together in $\frac{35}{12}$ hr, or 2 hr, 55 min.

$$\frac{35}{12} = 2\frac{11}{12} = 2\frac{11 \cdot 5}{12 \cdot 5} = 2\frac{55}{60},$$

or 2 hr, 55 min

Step 6 Check this result in the original problem.

NOW TRY 

NOTE There is another way to approach problems about rates of work. In **Example 8**, x represents the number of hours it will take the two people working together to complete the entire job. In 1 hr, $\frac{1}{x}$ of the entire job will be completed, so in 1 hr, Kareem completes $\frac{1}{7}$ of the job and Letitia completes $\frac{1}{5}$ of the job.

$$\frac{1}{7} + \frac{1}{5} = \frac{1}{x} \quad \text{The sum of their rates equals } \frac{1}{x}.$$

Multiplying each side of this equation by the LCD $35x$ gives $5x + 7x = 35$. This is equation (*) in **Example 8**. The same solution results using either approach.

NOW TRY ANSWER


8. $\frac{6}{5}$ hr, or 1 hr, 12 min

6.5 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Each exercise presents a familiar formula. Give the letter of the choice that is an equivalent form of the formula.

1. $p = br$ (percent)

A. $b = \frac{p}{r}$ B. $r = \frac{b}{p}$ C. $b = \frac{r}{p}$ D. $p = \frac{r}{b}$

2. $V = LWH$ (geometry)

A. $H = \frac{LW}{V}$ B. $L = \frac{V}{WH}$ C. $L = \frac{WH}{V}$ D. $W = \frac{H}{VL}$

3. $m = \frac{F}{a}$ (physics)

A. $a = mF$ B. $F = \frac{m}{a}$ C. $F = \frac{a}{m}$ D. $F = ma$

4. $I = \frac{E}{R}$ (electricity)

A. $R = \frac{I}{E}$ B. $R = IE$ C. $E = \frac{I}{R}$ D. $E = RI$

Solve each problem. See Example 1.

5. In work with electric circuits, the formula

$$\frac{1}{a} = \frac{1}{b} + \frac{1}{c}$$

occurs. Find b if $a = 8$ and $c = 12$.

7. A formula from anthropology says that

$$c = \frac{100b}{L}$$

Find L if $c = 80$ and $b = 5$.

9. **Concept Check** To solve the equation

$$m = \frac{ab}{a-b} \text{ for } a,$$

a student multiplied by $a - b$ and divided by b to obtain

$$\frac{m(a-b)}{b} = a.$$

WHAT WENT WRONG? Give the correct answer.

6. A gas law in chemistry says that

$$\frac{PV}{T} = \frac{pv}{t}.$$

Suppose that $T = 300$, $t = 350$, $V = 9$, $P = 50$, and $v = 8$. Find p .

8. The gravitational force between two masses is given by

$$F = \frac{GMm}{d^2}.$$

Find M to the nearest thousandth if $F = 10$, $G = 6.67 \times 10^{-11}$, $m = 1$, and $d = 3 \times 10^{-6}$.

10. **Concept Check** To solve the equation

$$rp - rq = p + q \text{ for } r,$$

a student began by adding rq and then dividing by p to obtain

$$r = \frac{p + q + rq}{p}.$$

WHAT WENT WRONG? Give the correct answer.

Solve each formula for the specified variable. See Examples 2 and 3.

11. $F = \frac{GMm}{d^2}$ for G (physics)

12. $F = \frac{GMm}{d^2}$ for M (physics)

13. $\frac{1}{a} = \frac{1}{b} + \frac{1}{c}$ for a (electricity)

14. $\frac{1}{a} = \frac{1}{b} + \frac{1}{c}$ for b (electricity)

15. $\frac{PV}{T} = \frac{pv}{t}$ for v (chemistry)

16. $\frac{PV}{T} = \frac{pv}{t}$ for T (chemistry)

17. $I = \frac{nE}{R + nr}$ for r (engineering)

18. $a = \frac{V - v}{t}$ for V (physics)

19. $\mathcal{A} = \frac{1}{2}h(b + B)$ for b (mathematics)

20. $S = \frac{n}{2}(a + \ell)d$ for n (mathematics)

21. $\frac{E}{e} = \frac{R + r}{r}$ for r (engineering)

22. $y = \frac{x + z}{a - x}$ for x (mathematics)

23. $D = \frac{R}{1 + RT}$ for R (banking)

24. $R = \frac{D}{1 - DT}$ for D (banking)

Concept Check Use proportions to solve each problem mentally.

25. In a mathematics class, 3 of every 4 students are girls. If there are 28 students in the class, how many are girls? How many are boys?

26. In a certain southern state, sales tax on a purchase of \$1.50 is \$0.12. What is the sales tax on a purchase of \$9.00?

The water content of snow is affected by the temperature, wind speed, and other factors present when the snow is falling. The average snow-to-liquid ratio is 10 in. of snow to 1 in. of liquid precipitation. This means that if 10 in. of snow fell and was melted, it would produce 1 in. of liquid precipitation in a rain gauge. (Data from www.theweatherprediction.com)

Use a proportion to solve each problem.

See Examples 4 and 5.

27. A dry snow might have a snow-to-liquid ratio of 18 to 1. Using this ratio, how much liquid precipitation would be produced by 31.5 in. of snow?
28. A wet, sticky snow good for making a snow man might have a snow-to-liquid ratio of 5 to 1. How many inches of fresh snow would produce 3.25 in. of liquid precipitation using this ratio?



Solve each problem. Give answers to the nearest tenth if an approximation is needed (unless specified otherwise). See Examples 4 and 5.


29. On a U.S. map, the distance between Seattle and Durango is 4.125 in. The two cities are actually 1238 mi apart. On this same map, what would be the distance between Chicago and El Paso, two cities that are actually 1606 mi apart? (Data from Universal Map Atlas.)
30. On a U.S. map, the distance between Reno and Phoenix is 2.5 in. The two cities are actually 768 mi apart. On this same map, what would be the distance between St. Louis and Jacksonville, two cities that are actually 919 mi apart? (Data from Universal Map Atlas.)
31. On a world globe, the distance between New York and Cairo, two cities that are actually 5619 mi apart, is 8.5 in. On this same globe, how far apart are Madrid and Rio de Janeiro, two cities that are actually 5045 mi apart? (Data from *The World Almanac and Book of Facts*.)
32. On a world globe, the distance between San Francisco and Melbourne, two cities that are actually 7856 mi apart, is 11.875 in. On this same globe, how far apart are Mexico City and Singapore, two cities that are actually 10,327 mi apart? (Data from *The World Almanac and Book of Facts*.)
33. In 2016, the average ratio of teachers to students in public elementary and secondary schools was approximately 1 to 16. If a public school had 896 students, how many teachers would be at the school according to this ratio? (Data from U.S. National Center for Education Statistics.)
34. At one point during the 2017–2018 NBA Season, the Indiana Pacers had won 19 of their 38 regular season games. If the team continued to win the same fraction of its games, how many games would the Pacers win for the complete 82-game season? Round the answer to the nearest whole number. (Data from www.nba.com)
35. To estimate the deer population of a forest preserve, wildlife biologists caught, tagged, and then released 42 deer. A month later, they returned and caught a sample of 75 deer and found that 15 of them were tagged. Based on this experiment, how many deer lived in the forest preserve?
36. Suppose that in the experiment in Exercise 35, only 5 of the previously tagged deer were collected in the sample of 75. What would be the estimate of the deer population?




37. Biologists tagged 500 fish in a lake on January 1. On February 1, they returned and collected a random sample of 400 fish, 8 of which had been previously tagged. On the basis of this experiment, how many fish does the lake have?
38. Suppose that in the experiment of **Exercise 37**, 10 of the previously tagged fish were collected on February 1. What would be the estimate of the fish population?
39. Bruce Johnston's Shelby Cobra uses 5 gal of gasoline to drive 156 mi. He has 3 gal of gasoline in the car, and he still needs to drive 300 mi. If the car continues to use gasoline at the same rate, how many more gallons will he need?
40. Mike Love's T-bird uses 6 gal of gasoline to drive 141 mi. He has 4 gal of gasoline in the car, and he still needs to drive 275 mi. If the car continues to use gasoline at the same rate, how many more gallons will he need?



In geometry, two triangles with corresponding angle measures equal, called **similar triangles**, have corresponding sides proportional. For example, in the figure, angle $A =$ angle D , angle $B =$ angle E , and angle $C =$ angle F , so the triangles are similar. Then the following ratios of corresponding sides are equal.



41. Solve for x , using the given proportion to find the lengths of the third sides of the triangles.
42. Suppose the following triangles are similar. Find y and the lengths of the two unknown sides of each triangle.



Nurses use proportions to determine the amount of a drug to administer when the dose is measured in milligrams but the drug is packaged in a diluted form in milliliters. (Data from Hoyles, C., R. Noss, and S. Pozzi, "Proportional Reasoning in Nursing Practice," *Journal for Research in Mathematics Education*.)

For example, to find the number of milliliters of fluid needed to administer 300 mg of a drug that comes packaged as 120 mg in 2 mL of fluid, a nurse sets up the proportion

$$\frac{120 \text{ mg}}{2 \text{ mL}} = \frac{300 \text{ mg}}{x \text{ mL}},$$

where x represents the amount to administer in milliliters. Use this method to find the correct dose for each prescription.

43. 120 mg of amikacin packaged as 100 mg in 2-mL vials
44. 1.5 mg of morphine packaged as 20-mg ampules diluted in 10 mL of fluid



Concept Check Solve each problem mentally.

45. If Marin can mow her yard in 3 hr, what is her rate (in terms of the proportion of the job per hour)?
46. A van traveling from Atlanta to Detroit averages 50 mph and takes 14 hr to make the trip. How far is it from Atlanta to Detroit?
47. **Concept Check** Greg paddles his canoe at a rate of 4 mph. If the rate of a stream is x mph, what is an expression that represents his rate in miles per hour when traveling each direction?

(a) Upstream (b) Downstream

48. **Concept Check** Which equation can be solved using cross products of the proportion? Explain. Then solve each equation.

$$\text{Equation A: } \frac{12}{x} - 1 = \frac{5}{x - 1}$$

$$\text{Equation B: } \frac{12}{x + 3} = \frac{8}{x - 3}$$

Solve each problem. See Examples 6 and 7.

49. Kellen's boat travels 12 mph. Find the rate of the current of the river if she can travel 6 mi upstream in the same amount of time she can go 10 mi downstream. (Let x = the rate of the current.)
50. Kasey can travel 8 mi upstream in the same time it takes her to go 12 mi downstream. Her boat goes 15 mph in still water. What is the rate of the current? (Let x = the rate of the current.)

	Distance	Rate	Time
Downstream	10	$12 + x$	
Upstream	6	$12 - x$	


	Distance	Rate	Time
Downstream			
Upstream		$15 - x$	

51. In his boat, Sheldon can travel 30 mi downstream in the same time that it takes to travel 10 mi upstream. If the rate of the current is 5 mph, find the rate of the boat in still water.
52. In his boat, Leonard can travel 24 mi upstream in the same time that it takes to travel 36 mi downstream. If the rate of the current is 2 mph, find the rate of the boat in still water.
53. On his drive from Montpelier, Vermont, to Columbia, South Carolina, Victor averaged 51 mph. If he had been able to average 60 mph, he would have reached his destination 3 hr earlier. What is the driving distance between Montpelier and Columbia?
54. Kelli lives in an off-campus apartment. When she rides her bike to campus, she arrives 36 min faster than when she walks. If her average walking rate is 3 mph and her average biking rate is 12 mph, how far is it from her apartment to campus?

	Distance	Rate	Time
Actual Trip			
Alternative Trip	x	60	

	Distance	Rate	Time
Bike	x	12	
Walk			

55. A plane averaged 500 mph on a trip going east, but only 350 mph on the return trip. The total flying time in both directions was 8.5 hr. What was the one-way distance?



56. Mary drove from her apartment to her parents' house. She averaged 60 mph because traffic was light. On the return trip, which took 1.5 hr longer, she averaged only 45 mph on the same route. What is the distance between Mary's apartment and her parents' house?

57. On the first part of a trip to Carmel traveling on the freeway, Marge averaged 60 mph. On the rest of the trip, which was 10 mi longer than the first part, she averaged 50 mph. Find the total distance to Carmel if the second part of the trip took 30 min more than the first part.
58. During the first part of a trip on the highway, Jim and Annie averaged 60 mph. In Houston, traffic caused them to average only 30 mph. The distance they drove in Houston was 100 mi less than their distance on the highway. What was their total driving distance if they spent 50 min more on the highway than they did in Houston?

Solve each problem. See Example 8.

59. A chef can finish a catering job in 6 hr. His student can do the same job in 12 hr. How long will it take them to do the job if they work together?
60. A printer can complete a large job in 8 hr using a new machine. Using an older machine requires 24 hr to do the same job. How long would it take to complete the job using both machines?
61. Butch and Peggy want to pick up the mess that their grandson, Grant, has made in his playroom. Butch could do it in 15 min working alone. Peggy, working alone, could clean it in 12 min. How long will it take them if they work together?
62. Lou can groom a customer's dogs in 8 hr, but it takes his business partner, Janet, only 5 hr to groom the same dogs. How long will it take them to groom the dogs if they work together?

	Rate	Time Working Together	Fractional Part of the Job Done
Butch	$\frac{1}{15}$	x	
Peggy	$\frac{1}{12}$	x	

	Rate	Time Working Together	Fractional Part of the Job Done
Lou	$\frac{1}{8}$	x	
Janet	$\frac{1}{5}$	x	

63. Jerry and Kuba are laying a hardwood floor. Working alone, Jerry can do the job in 20 hr. If the two of them work together, they can complete the job in 12 hr. How long would it take Kuba to lay the floor working alone? (Let x = the time it would take Kuba working alone.)
64. Mrs. Disher can grade a set of tests in 5 hr working alone. If her student teacher Mr. Howes helps her, it will take 3 hr to grade the tests. How long would it take Mr. Howes to grade the tests if he worked alone? (Let x = the time it would take Mr. Howes working alone.)

	Rate	Time Working Together	Fractional Part of the Job Done
Jerry		12	
Kuba		12	

	Rate	Time Working Together	Fractional Part of the Job Done
Mrs. Disher		3	
Mr. Howes		3	

65. Dixie can paint a room in 3 hr working alone. Trixie can paint the same room in 6 hr working alone. How long after Dixie starts to paint the room will it be finished if Trixie joins her 1 hr later?
66. Howard can wash a car in 30 min working alone. Raj can do the same job in 45 min working alone. How long after Howard starts to wash the car will it be finished if Raj joins him 5 min later?
67. If a vat of acid can be filled by an inlet pipe in 10 hr and emptied by an outlet pipe in 20 hr, how long will it take to fill the vat if both pipes are open?
68. A winery has a vat to hold Zinfandel. An inlet pipe can fill the vat in 9 hr, and an outlet pipe can empty it in 12 hr. How long will it take to fill the vat if both the outlet and the inlet pipes are open?

69. Suppose that Hortense and Mort can clean their entire house in 7 hr, while their toddler, Mimi, can mess it up in only 2 hr. If Hortense and Mort clean the house while Mimi is at her grandma's and then start cleaning up after Mimi the minute she gets home, how long does it take from the time Mimi gets home until the whole place is a shambles?



70. An inlet pipe can fill an artificial lily pond in 60 min, while an outlet pipe can empty it in 80 min. Through an error, both pipes are left open. How long will it take for the pond to fill?


6.6 Variation

OBJECTIVES

- 1 Write an equation expressing direct variation.
- 2 Find the constant of variation, and solve direct variation problems.
- 3 Solve inverse variation problems.
- 4 Solve joint variation problems.
- 5 Solve combined variation problems.

OBJECTIVE 1 Write an equation expressing direct variation.

The circumference of a circle is given by the formula $C = 2\pi r$, where r is the radius of the circle. See FIGURE 5. The circumference is always a constant multiple of the radius—that is, C is always found by multiplying r by the constant 2π .



$$C = 2\pi r$$

FIGURE 5

As the *radius increases*, the *circumference increases*.

As the *radius decreases*, the *circumference decreases*.

Because of these relationships, the circumference is said to *vary directly* as the radius.

Direct Variation

y **varies directly as** x if there exists a real number k such that

$$y = kx.$$

Stated another way, y is *directly proportional to* x . The number k is the **constant of variation**.

In direct variation, for $k > 0$, as the value of x increases, the value of y increases. Similarly, as x decreases, y decreases.

VOCABULARY

- direct variation
- constant of variation
- inverse variation
- joint variation
- combined variation

OBJECTIVE 2 Find the constant of variation, and solve direct variation problems.

The direct variation equation $y = kx$ defines a linear function, where the constant of variation k is the slope of the line. For example, the following equation describes the cost y to buy x gallons of gasoline.

$$y = 3.50x$$

The cost varies directly as the number of gallons purchased.

As the *number of gallons increases*, the *cost increases*.

As the *number of gallons decreases*, the *cost decreases*.

The constant of variation k is 3.50, the cost of 1 gal of gasoline.

NOW TRY EXERCISE 1

One week Morgan sold 8 dozen eggs for \$20. How much does she charge for one dozen eggs?

EXAMPLE 1 Solving a Direct Variation Problem

Eva is paid an hourly wage. One week she worked 43 hr and was paid \$795.50. How much does she earn per hour?

Let h = the number of hours she works

and P = her corresponding pay.

Write a variation equation.

k represents Eva's hourly wage.

$$P = kh$$

P varies directly as h .

$$795.50 = k \cdot 43$$

Let $P = 795.50$ and $h = 43$.

This is the constant of variation.

$$k = 18.50$$

Use a calculator.

Her hourly wage is \$18.50, and P and h are related by the equation

$$P = 18.50h.$$

We can use this equation to find her pay for any number of hours worked.

NOW TRY 

NOW TRY EXERCISE 2

For a constant height, the area of a parallelogram is directly proportional to its base. If the area is 20 cm^2 when the base is 4 cm, find the area when the base is 7 cm.

EXAMPLE 2 Solving a Direct Variation Problem

Hooke's law for an elastic spring states that the distance a spring stretches is directly proportional to the force applied. If a force of 150 newtons* stretches a certain spring 8 cm, how much will a force of 400 newtons stretch the spring? See FIGURE 6.

Let d = the distance the spring stretches

and f = the force applied.




FIGURE 6

Then $d = kf$ for some constant k . A force of 150 newtons stretches the spring 8 cm, so we use these values to find k .

$$d = kf$$

Variation equation

Solve for k .

$$8 = k \cdot 150$$

Let $d = 8$ and $f = 150$.

$$k = \frac{8}{150}$$

Solve for k .

$$k = \frac{4}{75}$$

Write in lowest terms.


Now we rewrite the variation equation $d = kf$ using $\frac{4}{75}$ for k .

$$d = \frac{4}{75}f \quad \text{Let } k = \frac{4}{75}.$$

For a force of 400 newtons, substitute 400 for f .

$$d = \frac{4}{75}(400) = \frac{64}{3} \quad \text{Let } f = 400.$$

The spring will stretch $\frac{64}{3}$ cm, or $21\frac{1}{3}$ cm, if a force of 400 newtons is applied.

NOW TRY 

NOW TRY ANSWERS

- \$2.50
- 35 cm^2

*A newton is a unit of measure of force used in physics.

Solving a Variation Problem


- Step 1** Write a variation equation.
- Step 2** Substitute the initial values and solve for k .
- Step 3** Rewrite the variation equation with the value of k from Step 2.
- Step 4** Substitute the remaining values, solve for the unknown, and find the required answer.

One variable can be directly proportional to a power of another variable.

Direct Variation as a Power

y varies directly as the n th power of x if there exists a real number k such that

$$y = kx^n.$$



$$A = \pi r^2$$

FIGURE 7

An example of direct variation as a power is the formula for the area of a circle, $A = \pi r^2$. See FIGURE 7. Here, π is the constant of variation, and the area A varies directly as the *square* of the radius r .


NOW TRY EXERCISE 3

Suppose y varies directly as the square of x , and $y = 200$ when $x = 5$. Find y when $x = 7$.

EXAMPLE 3 Solving a Direct Variation Problem

The distance a body falls from rest varies directly as the square of the time it falls (disregarding air resistance). If a skydiver falls 64 ft in 2 sec, how far will she fall in 8 sec?

- Step 1** Let d = the distance the skydiver falls
and t = the time it takes to fall.

Then d is a function of t for some constant k .

$$d = kt^2 \quad d \text{ varies directly as the square of } t.$$

- Step 2** To find the value of k , use the fact that the skydiver falls 64 ft in 2 sec.

$$\begin{aligned} d &= kt^2 && \text{Variation equation} \\ 64 &= k(2)^2 && \text{Let } d = 64 \text{ and } t = 2. \\ k &= 16 && \text{Find } k. \end{aligned}$$

- Step 3** Now we rewrite the variation equation $d = kt^2$ using 16 for k .


$$d = 16t^2 \quad \text{Let } k = 16.$$

- Step 4** Let $t = 8$ to find the number of feet the skydiver will fall in 8 sec.

$$d = 16(8)^2 = 1024 \quad \text{Let } t = 8.$$

The skydiver will fall 1024 ft in 8 sec.

NOW TRY 



As pressure on trash **increases**, volume of trash **decreases**.

FIGURE 8

OBJECTIVE 3 Solve inverse variation problems.

Another type of variation is *inverse variation*. **With inverse variation, for $k > 0$, as one variable increases, the other variable decreases.**

For example, in a closed space, volume decreases as pressure increases, which can be illustrated by a trash compactor. See FIGURE 8. As the compactor presses down, the pressure on the trash increases, and in turn, the trash occupies a smaller space.

NOW TRY ANSWER

3. 392

Inverse Variation

y varies inversely as x if there exists a real number k such that

$$y = \frac{k}{x}.$$

y varies inversely as the n th power of x if there exists a real number k such that

$$y = \frac{k}{x^n}.$$

The inverse variation equation $y = \frac{k}{x}$ defines a rational function.

Another example of inverse variation comes from the distance formula.

$$d = rt \quad \text{Distance formula}$$

$$t = \frac{d}{r} \quad \text{Divide each side by } r.$$

In the form $t = \frac{d}{r}$, t (time) varies inversely as r (rate or speed), with d (distance) serving as the constant of variation. For example, if the distance between Chicago and Des Moines is 300 mi, then

$$t = \frac{300}{r}.$$

The values of r and t might be any of the following.

$$\left. \begin{array}{l} r = 50, t = 6 \\ r = 60, t = 5 \\ r = 75, t = 4 \end{array} \right\} \begin{array}{l} \text{As } r \text{ increases,} \\ t \text{ decreases.} \end{array} \quad \left. \begin{array}{l} r = 30, t = 10 \\ r = 25, t = 12 \\ r = 20, t = 15 \end{array} \right\} \begin{array}{l} \text{As } r \text{ decreases,} \\ t \text{ increases.} \end{array}$$

If we *increase* the rate (speed) at which we drive, time *decreases*. If we *decrease* the rate (speed) at which we drive, time *increases*.

NOW TRY EXERCISE 4

For a constant area, the height of a triangle varies inversely as the base. If the height is 7 cm when the base is 8 cm, find the height when the base is 14 cm.

EXAMPLE 4 Solving an Inverse Variation Problem

In the manufacture of a phone-charging device, the cost of producing the device varies inversely as the number produced. If 10,000 units are produced, the cost is \$2 per unit. Find the cost per unit to produce 25,000 units.

Let x = the number of units produced

and c = the cost per unit.

Here, as production increases, cost decreases, and as production decreases, cost increases. We write a variation equation using the variables c and x and the constant k .

$$c = \frac{k}{x} \quad c \text{ varies inversely as } x.$$

$$2 = \frac{k}{10,000} \quad \text{Let } c = 2 \text{ and } x = 10,000.$$

$$20,000 = k \quad \text{Multiply by } 10,000.$$

Thus, the variation equation $c = \frac{k}{x}$ becomes $c = \frac{20,000}{x}$.

$$c = \frac{20,000}{25,000} = 0.80 \quad \text{Let } x = 25,000.$$

The cost per unit to make 25,000 units is \$0.80.

NOW TRY ANSWER

4. 4 cm

NOW TRY EXERCISE 5

The weight of an object above Earth varies inversely as the square of its distance from the center of Earth. If an object weighs 150 lb on the surface of Earth, and the radius of Earth is about 3960 mi, how much does it weigh when it is 1000 mi above Earth's surface? Round to the nearest pound.

EXAMPLE 5 Solving an Inverse Variation Problem

The weight of an object above Earth varies inversely as the square of its distance from the center of Earth. A space shuttle in an elliptical orbit has a maximum distance from the center of Earth (**apogee**) of 6700 mi. Its minimum distance from the center of Earth (**perigee**) is 4090 mi. See **FIGURE 9**. If an astronaut in the shuttle weighs 57 lb at its apogee, what does the astronaut weigh at its perigee?




FIGURE 9

Let w = the astronaut's weight and d = the distance from the center of Earth, for some constant k . We write a variation equation using these variables.

$$w = \frac{k}{d^2} \quad w \text{ varies inversely as the square of } d.$$

At the apogee, the astronaut weighs 57 lb, and the distance from the center of Earth is 6700 mi. Use these values to find k .

$$57 = \frac{k}{(6700)^2} \quad \text{Let } w = 57 \text{ and } d = 6700.$$

$$k = 57(6700)^2 \quad \text{Solve for } k.$$


Substitute $k = 57(6700)^2$ and $d = 4090$ in the variation equation $w = \frac{k}{d^2}$ to find the astronaut's weight at the perigee.

$$w = \frac{57(6700)^2}{(4090)^2} = 153 \text{ lb} \quad \begin{array}{l} \text{Use a calculator.} \\ \text{Round to the nearest pound.} \end{array}$$

NOW TRY

OBJECTIVE 4 Solve joint variation problems.

If one variable varies directly as the *product* of several other variables (perhaps raised to powers), the first variable is said to *vary jointly* as the others.



B = area of the base

$$V = \frac{1}{3}Bh$$

FIGURE 10

Joint Variation

y varies jointly as x and z if there exists a real number k such that

$$y = kxz.$$

An example of joint variation is the formula for the volume of a right pyramid, $V = \frac{1}{3}Bh$. See **FIGURE 10**. Here, $\frac{1}{3}$ is the constant of variation, and the volume V varies jointly as the area of the base B and the height h .

CAUTION Note that *and* in the expression “ y varies jointly as x and z ” translates as a product in the variation equation $y = kxz$. The word *and* does **not** indicate addition here.

NOW TRY ANSWER

5. 96 lb

NOW TRY EXERCISE 6

The volume of a right pyramid varies jointly as the height and the area of the base. If the volume is 100 ft^3 when the area of the base is 30 ft^2 and the height is 10 ft, find the volume when the area of the base is 90 ft^2 and the height is 20 ft.

EXAMPLE 6 Solving a Joint Variation Problem

The interest on a loan or an investment is given by the formula $I = prt$. Here, for a given principal p , the interest earned I varies jointly as the interest rate r and the time t the principal is left earning interest. If an investment earns \$100 interest at 5% for 2 yr, how much interest will the same principal earn at 4.5% for 3 yr?

We use the formula $I = prt$, where p is the constant of variation because it is the same for both investments.

$$\begin{array}{ll} I = prt & \text{Here, } p \text{ is the constant of variation.} \\ \text{Solve for } p. \quad 100 = p(0.05)(2) & \text{Let } I = 100, r = 5\% = 0.05, \text{ and } t = 2. \\ 100 = 0.1p & \text{Multiply.} \\ p = 1000 & \text{Divide by 0.1. Rewrite.} \end{array}$$

Now we find I when $p = 1000$, $r = 4.5\% = 0.045$, and $t = 3$.

$$I = 1000(0.045)(3) = 135 \quad \text{Let } p = 1000, r = 4.5\% = 0.045, \text{ and } t = 3.$$

The interest will be \$135.

NOW TRY 

OBJECTIVE 5 Solve combined variation problems.

There are combinations of direct and inverse variation, called **combined variation**.

NOW TRY EXERCISE 7

In statistics, the sample size used to estimate a population mean varies directly as the variance and inversely as the square of the maximum error of the estimate. If the sample size is 200 when the variance is 25 m^2 and the maximum error of the estimate is 0.5 m, find the sample size when the variance is 25 m^2 and the maximum error of the estimate is 0.1 m.

EXAMPLE 7 Solving a Combined Variation Problem

Body mass index (BMI) is used to assess whether a person's weight is healthy. A BMI from 19 through 25 is considered desirable. BMI varies directly as an individual's weight in pounds and inversely as the square of his or her height in inches.

A person who weighs 118 lb and is 64 in. tall has a BMI of 20. (BMI is rounded to the nearest whole number.) Find the BMI of a man who weighs 165 lb and is 70 in. tall. (Data from *Washington Post*.)

Let $B = \text{BMI}$, $w = \text{weight}$, and $h = \text{height}$. Write a variation equation.

$$B = \frac{kw}{h^2} \quad \begin{array}{l} \leftarrow \text{BMI varies directly as the weight.} \\ \leftarrow \text{BMI varies inversely as the square} \\ \quad \text{of the height.} \end{array}$$

To find k , let $B = 20$, $w = 118$, and $h = 64$.

$$\begin{array}{ll} 20 = \frac{k(118)}{64^2} & B = \frac{kw}{h^2} \\ k = \frac{20(64^2)}{118} & \text{Multiply by } 64^2. \\ k = 694 & \text{Divide by } 118. \\ & \text{Use a calculator. Round to} \\ & \text{the nearest whole number.} \end{array}$$

Now find B when $k = 694$, $w = 165$, and $h = 70$.

$$B = \frac{694(165)}{70^2} = 23 \quad \text{Round to the nearest whole number.}$$

The man's BMI is 23.




NOW TRY ANSWERS

6. 600 ft^3
7. 5000

NOW TRY 

6.6 Exercises

FOR
EXTRA
HELP
 MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Fill in each blank with the correct response.

- For $k > 0$, if y varies directly as x , then when x increases, y _____, and when x decreases, y _____.
- For $k > 0$, if y varies inversely as x , then when x increases, y _____, and when x decreases, y _____.

Concept Check Use personal experience or intuition to determine whether the situation suggests direct or inverse variation.

- The number of movie tickets purchased and the total price for the tickets
- The rate and the distance traveled by a pickup truck in 3 hr
- The amount of pressure put on the accelerator of a car and the speed of the car
- The percentage off an item that is on sale and the price of the item
- Your age and the probability that you believe in the tooth fairy
- The surface area of a balloon and its diameter
- The demand for an item and the price of the item
- The number of hours worked by an hourly worker and the amount of money earned

Concept Check Determine whether each equation represents direct, inverse, joint, or combined variation.

- | | | | |
|-----------------------|-----------------------|-------------------------|-------------------------|
| 11. $y = \frac{3}{x}$ | 12. $y = \frac{8}{x}$ | 13. $y = 10x^2$ | 14. $y = 2x^3$ |
| 15. $y = 3xz^4$ | 16. $y = 6x^3z^2$ | 17. $y = \frac{4x}{wz}$ | 18. $y = \frac{6x}{st}$ |

Concept Check Write each formula using the “language” of variation. For example, the formula for the circumference of a circle, $C = 2\pi r$, can be written as

“The circumference of a circle varies directly as the length of its radius.”

- $P = 4s$, where P is the perimeter of a square with side of length s
- $d = 2r$, where d is the diameter of a circle with radius r
- $S = 4\pi r^2$, where S is the surface area of a sphere with radius r
- $V = \frac{4}{3}\pi r^3$, where V is the volume of a sphere with radius r
- $\mathcal{A} = \frac{1}{2}bh$, where \mathcal{A} is the area of a triangle with base b and height h
- $V = \frac{1}{3}\pi r^2h$, where V is the volume of a cone with radius r and height h
- Concept Check** What is the constant of variation in each of the variation equations in Exercises 19–24?
- Concept Check** What is meant by the constant of variation in a direct variation problem? If we were to graph the linear equation $y = kx$ for some nonnegative constant k , what role would k play in the graph?

Write a variation equation for each situation. Use k as the constant of variation. See Examples 1–6.

- A varies directly as b .
- W varies directly as f .



29. h varies inversely as t .
 30. p varies inversely as s .
 31. M varies directly as the square of d .
 32. P varies inversely as the cube of x .
 33. I varies jointly as g and h .
 34. C varies jointly as a and the square of b .

Solve each problem. See Examples 1–7.


35. If x varies directly as y , and $x = 9$ when $y = 3$, find x when $y = 12$.
 36. If x varies directly as y , and $x = 10$ when $y = 7$, find y when $x = 50$.
 37. If a varies directly as the square of b , and $a = 4$ when $b = 3$, find a when $b = 2$.
 38. If h varies directly as the square of m , and $h = 15$ when $m = 5$, find h when $m = 7$.
 39. If z varies inversely as w , and $z = 10$ when $w = 0.5$, find z when $w = 8$.
 40. If t varies inversely as s , and $t = 3$ when $s = 5$, find s when $t = 5$.
 41. If m varies inversely as the square of p , and $m = 20$ when $p = 2$, find m when $p = 5$.
 42. If a varies inversely as the square of b , and $a = 48$ when $b = 4$, find a when $b = 7$.
 43. p varies jointly as q and the square of r , and $p = 200$ when $q = 2$ and $r = 3$. Find p when $q = 5$ and $r = 2$.
 44. f varies jointly as h and the square of g , and $f = 50$ when $h = 2$ and $g = 4$. Find f when $h = 6$ and $g = 3$.

Solve each problem. See Examples 1–7.

45. Ben bought 8.5 gal of gasoline and paid \$33.32. What is the price of gasoline per gallon?
 46. Sara gives horseback rides at Shadow Mountain Ranch. A 2.5-hr ride costs \$50.00. What is the price per hour?
 47. The weight of an object on Earth is directly proportional to the weight of that same object on the moon. A 200-lb astronaut would weigh 32 lb on the moon. How much would a 50-lb dog weigh on the moon?
 48. The pressure exerted by a certain liquid at a given point is directly proportional to the depth of the point beneath the surface of the liquid. The pressure at 30 m is 80 newtons. What pressure is exerted at 50 m?
 49. The volume of a can of tomatoes is directly proportional to the height of the can. If the volume of the can is 300 cm^3 when its height is 10.62 cm, find the volume to the nearest whole number of a can with height 15.92 cm.





51. For a body falling freely from rest (disregarding air resistance), the distance the body falls varies directly as the square of the time. If an object is dropped from the top of a tower 576 ft high and hits the ground in 6 sec, how far did it fall in the first 4 sec?



52. The amount of water emptied by a pipe varies directly as the square of the diameter of the pipe. For a certain constant water flow, a pipe emptying into a canal will allow 200 gal of water to escape in an hour. The diameter of the pipe is 6 in. How much water would a 12-in. pipe empty into the canal in an hour, assuming the same water flow?




53. Over a specified distance, rate varies inversely as time. If a Dodge Viper on a test track goes a certain distance in one-half minute at 160 mph, what rate is needed to go the same distance in three-fourths minute?
54. For a constant area, the length of a rectangle varies inversely as the width. The length of a rectangle is 27 ft when the width is 10 ft. Find the width of a rectangle with the same area if the length is 18 ft.
55. The frequency of a vibrating string varies inversely as its length. That is, a longer string vibrates fewer times in a second than a shorter string. Suppose a piano string 2 ft long vibrates at 250 cycles per sec. What frequency would a string 5 ft long have?
56. The current in a simple electrical circuit varies inversely as the resistance. If the current is 20 amps when the resistance is 5 ohms, find the current when the resistance is 7.5 ohms.
57. The amount of light (measured in foot-candles) produced by a light source varies inversely as the square of the distance from the source. If the illumination produced 1 m from a light source is 768 foot-candles, find the illumination produced 6 m from the same source.
58. The force with which Earth attracts an object above Earth's surface varies inversely as the square of the distance of the object from the center of Earth. If an object 4000 mi from the center of Earth is attracted with a force of 160 lb, find the force of attraction if the object were 6000 mi from the center of Earth.



59. For a given interest rate, simple interest varies jointly as principal and time. If \$2000 left in an account for 4 yr earned interest of \$280, how much interest would be earned in 6 yr?
60. The collision impact of an automobile varies jointly as its mass and the square of its speed. Suppose a 2000-lb car traveling at 55 mph has a collision impact of 6.1. What is the collision impact (to the nearest tenth) of the same car at 65 mph?
61. The weight of a bass varies jointly as its girth and the square of its length. (**Girth** is the distance around the body of the fish.) A prize-winning bass weighed in at 22.7 lb and measured 36 in. long with a 21-in. girth. How much (to the nearest tenth of a pound) would a bass 28 in. long with an 18-in. girth weigh?
62. The weight of a trout varies jointly as its length and the square of its girth. One angler caught a trout that weighed 10.5 lb and measured 26 in. long with an 18-in. girth. Find the weight (to the nearest tenth of a pound) of a trout that is 22 in. long with a 15-in. girth.
63. The force needed to keep a car from skidding on a curve varies inversely as the radius of the curve and jointly as the weight of the car and the square of the speed. If 242 lb of force keeps a 2000-lb car from skidding on a curve of radius 500 ft at 30 mph, what force (to the nearest tenth of a pound) would keep the same car from skidding on a curve of radius 750 ft at 50 mph?



64. The maximum load that a cylindrical column with a circular cross section can hold varies directly as the fourth power of the diameter of the cross section and inversely as the square of the height. A 9-m column 1 m in diameter will support 8 metric tons. How many metric tons can be supported by a column 12 m high and $\frac{2}{3}$ m in diameter?



65. The number of long-distance phone calls between two cities during a certain period varies jointly as the populations of the cities, p_1 and p_2 , and inversely as the distance between them, in miles. If 80,000 calls are made between two cities 400 mi apart, with populations of 70,000 and 100,000, how many calls (to the nearest hundred) are made between cities with populations of 50,000 and 75,000 that are 250 mi apart?
66. The volume of gas varies inversely as the pressure and directly as the temperature. (Temperature must be measured in *kelvins* (K), a unit of measurement used in physics.) If a certain gas occupies a volume of 1.3 L at 300 K and a pressure of 18 newtons, find the volume at 340 K and a pressure of 24 newtons.
67. A body mass index from 27 through 29 carries a slight risk of weight-related health problems, while a BMI of 30 or more indicates a great increase in risk. Use your own height and weight and the information in **Example 7** to determine your BMI and whether you are at risk.
68. The maximum load of a horizontal beam that is supported at both ends varies directly as the width and the square of the height and inversely as the length between the supports. A beam 6 m long, 0.1 m wide, and 0.06 m high supports a load of 360 kg. What is the maximum load supported by a beam 16 m long, 0.2 m wide, and 0.08 m high?

RELATING CONCEPTS For Individual or Group Work (Exercises 69–74)

A routine activity such as pumping gasoline can be related to many of the concepts studied in this text. Suppose that premium unleaded costs \$3.75 per gal. **Work Exercises 69–74 in order.**



69. 0 gal of gasoline cost \$0.00, while 1 gal costs \$3.75. Represent these two pieces of information as ordered pairs of the form (gallons, price).
70. Use the information from **Exercise 69** to find the slope of the line on which the two points lie.
71. Write the slope-intercept form of the equation of the line on which the two points lie.
72. Using function notation, if $f(x) = ax + b$ represents the line from **Exercise 71**, what are the values of a and b ?
73. How is the value of a from **Exercise 72** related to gasoline in this situation? With relationship to the line, what do we call this number?
74. Why does the equation from **Exercise 72** satisfy the conditions for direct variation? In the context of variation, what do we call the value of a ?

Chapter 6

Summary

Key Terms

6.1

rational expression
rational function
reciprocal

6.2

least common denominator
(LCD)

6.3

complex fraction

6.4

domain of the variable in a
rational equation
proposed solution
extraneous solution
discontinuous

reciprocal function
vertical asymptote
horizontal asymptote

6.5

ratio
proportion
cross products of a
proportion

6.6

direct variation
constant of variation
inverse variation
joint variation
combined variation

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- A **rational expression** is
 - an algebraic expression made up of a term or the sum of a finite number of terms with real coefficients and integer exponents
 - a polynomial equation of degree 2
 - a quotient with one or more fractions in the numerator, denominator, or both
 - a quotient of two polynomials with denominator not 0.
- In a given set of fractions, the **least common denominator** is
 - the smallest denominator of all the denominators
 - the product of the factors from each denominator, with each factor raised to the greatest power that occurs
- the largest integer that evenly divides the numerator and denominator of all the fractions
 - the sum of the factors raised to the greatest power that occurs in each denominator.
- A **complex fraction** is
 - an algebraic expression made up of a term or the sum of a finite number of terms with real coefficients and integer exponents
 - a polynomial equation of degree 2
 - a quotient with one or more fractions in the numerator, denominator, or both
 - a quotient of two polynomials with denominator not 0.
- A **ratio**
 - compares two quantities using a quotient
 - says that two quotients are equal
 - is a product of two quantities
 - is a difference of two quantities.
- A **proportion**
 - compares two quantities using a quotient
 - says that two quotients are equal
 - is a product of two quantities
 - is a difference of two quantities.

ANSWERS

- D; Examples: $-\frac{3}{4y^2}$, $\frac{5x^3}{x+2}$, $\frac{a+3}{a^2-4a-5}$
- B; Example: The LCD of $\frac{1}{x}$, $\frac{2}{3}$, and $\frac{5}{x+1}$ is $3x(x+1)$.
- C; Examples: $\frac{\frac{2}{3}}{\frac{4}{7}}$, $\frac{x-\frac{1}{x}}{x+\frac{1}{y}}$, $\frac{\frac{2}{a+1}}{a^2-1}$
- A; Example: $\frac{7 \text{ in.}}{12 \text{ in.}}$ compares two quantities.
- B; Example: The proportion $\frac{2}{3} = \frac{8}{12}$ states that the two ratios are equal.

Quick Review

CONCEPTS

6.1 Rational Expressions and Functions;
Multiplying and Dividing**Rational Function**

A rational function is defined by a quotient of polynomials and has the form

$$f(x) = \frac{P(x)}{Q(x)}, \quad \text{where } Q(x) \neq 0.$$

Its domain includes all real numbers except those that make $Q(x)$ equal to 0.

Fundamental Property of Rational Numbers

If $\frac{a}{b}$ is a rational number and if c is any nonzero real number, then

$$\frac{a}{b} = \frac{ac}{bc}.$$

Writing a Rational Expression in Lowest Terms

Step 1 Factor both numerator and denominator to find their greatest common factor (GCF).

Step 2 Apply the fundamental property. Divide out common factors.

Multiplying Rational Expressions

Step 1 Factor numerators and denominators.

Step 2 Apply the fundamental property.

Step 3 Multiply remaining factors in the numerators and in the denominators. The final expression can be left in factored form.

Step 4 Check that the product is in lowest terms.

Dividing Rational Expressions

Multiply the first rational expression (the dividend) by the reciprocal of the second expression (the divisor).

EXAMPLES

Give the domain of the rational function.

$$f(x) = \frac{2x + 1}{3x + 6}$$

Solve $3x + 6 = 0$ to find $x = -2$. This value must be excluded from the domain, which can be written

$$\{x \mid x \text{ is a real number, } x \neq -2\} \quad \text{or} \quad (-\infty, -2) \cup (-2, \infty).$$

Write in lowest terms.

$$\frac{15}{20} = \frac{3 \cdot 5}{4 \cdot 5} = \frac{3}{4} \quad \frac{15}{20} \text{ and } \frac{3}{4} \text{ are equivalent.}$$

$$\frac{2x + 8}{x^2 - 16}$$

$$= \frac{2(x + 4)}{(x - 4)(x + 4)} \quad \text{Factor.}$$

$$= \frac{2}{x - 4} \quad \text{Fundamental property}$$

Multiply or divide.

$$\frac{x^2 + 2x + 1}{x^2 - 1} \cdot \frac{5}{3x + 3}$$

$$= \frac{(x + 1)^2}{(x - 1)(x + 1)} \cdot \frac{5}{3(x + 1)} \quad \text{Factor.}$$

$$= \frac{5}{3(x - 1)} \quad \text{Fundamental property; Multiply.}$$

$$\frac{2x + 5}{x - 3} \div \frac{2x^2 + 3x - 5}{x^2 - 9}$$

$$= \frac{2x + 5}{x - 3} \cdot \frac{x^2 - 9}{2x^2 + 3x - 5} \quad \text{Multiply by the reciprocal.}$$

$$= \frac{2x + 5}{x - 3} \cdot \frac{(x + 3)(x - 3)}{(2x + 5)(x - 1)} \quad \text{Factor.}$$

$$= \frac{x + 3}{x - 1} \quad \text{Fundamental property; Multiply.}$$

CONCEPTS

6.2 Adding and Subtracting Rational Expressions

Adding or Subtracting Rational Expressions

Step 1 If the denominators are the same, add or subtract the numerators. Place the result over the common denominator.

If the denominators are different, write all rational expressions with the LCD. Then add or subtract the numerators, and place the result over the common denominator.

Step 2 Make sure that the answer is in lowest terms.

EXAMPLES

Subtract.

$$\begin{aligned} \frac{1}{x+6} - \frac{3}{x+2} & \quad \text{The LCD is } (x+6)(x+2). \\ &= \frac{x+2}{(x+6)(x+2)} - \frac{3(x+6)}{(x+6)(x+2)} \\ &= \frac{x+2-3(x+6)}{(x+6)(x+2)} \quad \text{Subtract the numerators.} \\ &= \frac{x+2-3x-18}{(x+6)(x+2)} \quad \text{Distributive property} \\ &= \frac{-2x-16}{(x+6)(x+2)} \quad \text{Combine like terms.} \end{aligned}$$

The numerator factors as $-2(x+8)$, so there is no common factor in the numerator and denominator. The answer is in lowest terms.

6.3 Complex Fractions

Simplifying a Complex Fraction

Method 1

Step 1 Simplify the numerator and denominator separately.

Step 2 Divide by multiplying the numerator by the reciprocal of the denominator.

Step 3 Simplify the resulting fraction if possible.

Method 2

Step 1 Multiply the numerator and denominator of the complex fraction by the least common denominator of all the fractions appearing in the complex fraction.

Step 2 Simplify the resulting fraction if possible.

Simplify using both methods.

<i>Method 1</i>	<i>Method 2</i>
$\begin{aligned} \frac{\frac{1}{x^2} - \frac{1}{y^2}}{\frac{1}{x} + \frac{1}{y}} &= \frac{\frac{y^2}{x^2y^2} - \frac{x^2}{x^2y^2}}{\frac{y}{xy} + \frac{x}{xy}} \\ &= \frac{\frac{y^2 - x^2}{x^2y^2}}{\frac{y+x}{xy}} \\ &= \frac{y^2 - x^2}{x^2y^2} \div \frac{y+x}{xy} \\ &= \frac{(y+x)(y-x)}{x^2y^2} \cdot \frac{xy}{y+x} \\ &= \frac{y-x}{xy} \end{aligned}$	$\begin{aligned} \frac{\frac{1}{x^2} - \frac{1}{y^2}}{\frac{1}{x} + \frac{1}{y}} &= \frac{x^2y^2\left(\frac{1}{x^2} - \frac{1}{y^2}\right)}{x^2y^2\left(\frac{1}{x} + \frac{1}{y}\right)} \\ &= \frac{y^2 - x^2}{xy^2 + x^2y} \\ &= \frac{(y-x)(y+x)}{xy(y+x)} \\ &= \frac{y-x}{xy} \end{aligned}$

CONCEPTS

6.4 Equations with Rational Expressions and Graphs**Solving an Equation with Rational Expressions**

- Step 1** Determine the domain of the variable.
- Step 2** Multiply each side of the equation by the LCD to clear the fractions.
- Step 3** Solve the resulting equation.
- Step 4** Check that each proposed solution is in the domain, and discard any values that are not. Check the remaining proposed solution(s) in the original equation.

Graphing a Rational Function

The graph of a rational function (written in lowest terms) may have one or more breaks. At such points, the graph will approach an asymptote.

6.5 Applications of Rational Expressions**Solving a Distance, Rate, Time Problem**

Use the distance formula

$$d = rt$$

or one of its equivalents,

$$t = \frac{d}{r} \quad \text{or} \quad r = \frac{d}{t}$$

Solving a Work Problem

Use the fact that if 1 complete job is done in t units of time, then the rate of work is $\frac{1}{t}$ job per unit of time.

$$1 = rt, \quad \text{or} \quad r = \frac{1}{t}$$

EXAMPLES

Solve.

$$\frac{3x+2}{x-2} + \frac{2}{x(x-2)} = \frac{-1}{x}$$

$$x(3x+2) + 2 = -(x-2)$$

$$3x^2 + 2x + 2 = -x + 2$$

$$3x^2 + 3x = 0$$

$$3x(x+1) = 0$$

$$3x = 0 \quad \text{or} \quad x + 1 = 0$$

$$x = 0 \quad \text{or} \quad x = -1$$

Note that 0 and 2 are excluded from the domain.

Multiply by the LCD, $x(x-2)$.

Distributive property

Add x . Subtract 2.


Factor.

Zero-factor property

Solve each equation.

Of the two proposed solutions, 0 must be discarded because it is not in the domain. A check confirms that the solution set is $\{-1\}$.

Graph $f(x) = \frac{1}{x+2}$.



Solve.

A canal has a current of 2 mph. Find the rate of Amy's boat in still water if it travels 11 mi downstream in the same time that it travels 8 mi upstream.

Let x = the rate of the boat in still water.

	Distance	Rate	Time
Downstream	11	$x + 2$	$\frac{11}{x+2}$
Upstream	8	$x - 2$	$\frac{8}{x-2}$

The times are equal.

$$\frac{11}{x+2} = \frac{8}{x-2}$$

Write an equation. Use $t = \frac{d}{r}$.

$$11(x-2) = 8(x+2)$$

If $\frac{a}{b} = \frac{c}{d}$, then $ad = bc$.

$$11x - 22 = 8x + 16$$

Distributive property

$$3x = 38$$

Subtract $8x$. Add 22.

$$x = \frac{38}{3}, \quad \text{or} \quad 12\frac{2}{3}$$

Divide by 3.

The rate in still water is $12\frac{2}{3}$ mph.

CONCEPTS

6.6 Variation

Let k be a real number.

If $y = kx$, then y varies directly as x .

If $y = kx^n$, then y varies directly as x^n .

If $y = \frac{k}{x}$, then y varies inversely as x .

If $y = kxz$, then y varies jointly as x and z .

EXAMPLES

The diameter of a circle varies directly as the radius.

$$d = kr \quad \text{Here, } k = 2.$$

The area of a circle varies directly as the square of the radius.

$$\mathcal{A} = kr^2 \quad \text{Here, } k = \pi.$$

Pressure varies inversely as volume.

$$p = \frac{k}{V}$$

For a given principal, interest varies jointly as interest rate and time.

$$I = krt \quad k \text{ is the given principal.}$$

Chapter 6

Review Exercises

6.1 Give the domain of each rational function using (a) set-builder notation and (b) interval notation.

$$1. f(x) = \frac{-7}{3x + 18}$$

$$2. f(x) = \frac{5x + 17}{x^2 - 7x + 10}$$

$$3. f(x) = \frac{9}{x^2 - 18x + 81}$$

Write each rational expression in lowest terms.

$$4. \frac{12x^2 + 6x}{24x + 12}$$

$$5. \frac{25m^2 - n^2}{25m^2 - 10mn + n^2}$$

$$6. \frac{r - 2}{4 - r^2}$$

Multiply or divide as indicated.

$$7. \frac{(2y + 3)^2}{5y} \cdot \frac{15y^3}{4y^2 - 9}$$

$$8. \frac{w^2 - 16}{w} \cdot \frac{3}{4 - w}$$

$$9. \frac{x^2 - 4}{x - 2} \div \frac{x + 2}{x - 4}$$

$$10. \frac{z^2 - z - 6}{z - 6} \cdot \frac{z^2 - 6z}{z^2 + 2z - 15}$$

$$11. \frac{8 - x}{8 + x} \cdot \frac{16 + 2x}{x^2 - 64}$$

$$12. \frac{m^3 - n^3}{m^2 - n^2} \div \frac{m^2 + mn + n^2}{m + n}$$

6.2 Suppose that the given expressions are denominators of rational expressions. Find the least common denominator (LCD) for each group of denominators.

$$13. 32b^3, 24b^5$$

$$14. 9r^2, 3r + 1, 9$$

$$15. 6x^2 + 13x - 5, 9x^2 + 9x - 4$$

$$16. 3x - 12, x^2 - 2x - 8, x^2 - 8x + 16$$

Add or subtract as indicated.

$$17. \frac{8}{z} - \frac{3}{2z^2}$$

$$18. \frac{5y + 13}{y + 1} - \frac{1 - 7y}{y + 1}$$

$$19. \frac{6}{5a + 10} + \frac{7}{6a + 12}$$

$$20. \frac{3r}{10r^2 - 3rs - s^2} + \frac{2r}{2r^2 + rs - s^2}$$

6.3 Simplify each complex fraction.

21.
$$\frac{\frac{3}{t} + 2}{\frac{4}{t} - 7}$$

22.
$$\frac{\frac{2}{m-3n}}{\frac{1}{3n-m}}$$

23.
$$\frac{\frac{\frac{3}{p} - \frac{2}{q}}{9q^2 - 4p^2}}{qp}$$

24.
$$\frac{x^{-2} - y^{-2}}{x^{-1} - y^{-1}}$$

6.4 Solve each equation.

25.
$$\frac{1}{t+4} + \frac{1}{2} = \frac{3}{2t+8}$$

26.
$$\frac{-5m}{m+1} + \frac{m}{3m+3} = \frac{56}{6m+6}$$

27.
$$\frac{4x+3}{5x+5} = \frac{8x+2}{10x+5}$$

28.
$$\frac{2}{k-1} - \frac{4k+1}{k^2-1} = \frac{-1}{k+1}$$

29.
$$\frac{r}{r-2} - \frac{8}{r^2-4} = \frac{-1}{r+2}$$

30.
$$\frac{5}{x+2} + \frac{3}{x+3} = \frac{x}{x^2+5x+6}$$

31. Decide whether each of the following is an *expression* or an *equation*. Simplify the one that is an expression, and solve the one that is an equation.

(a)
$$\frac{4}{x} + \frac{1}{2} = \frac{1}{3}$$

(b)
$$\frac{4}{x} + \frac{1}{2} - \frac{1}{3}$$

32. Graph the rational function $f(x) = \frac{2}{x+1}$. Give the equations of the vertical and horizontal asymptotes.

6.5 Solve each problem.

33. According to a law from physics,

$$\frac{1}{A} = \frac{1}{B} + \frac{1}{C}.$$

Find A if $B = 30$ and $C = 10$.

34. In banking and finance, the formula

$$P = \frac{M}{1 + RT}$$

gives present value at simple interest. If $P = 600$, $M = 750$, and $R = 0.125$, find T .

Solve each formula for the specified variable.

35.
$$F = \frac{GMm}{d^2}$$
 for m (physics)

36.
$$\mu = \frac{Mv}{M+m}$$
 for M (electronics)

Solve each problem.

37. An article in *Scientific American* predicted that in the year 2050, about 23,200 of the 58,000 passenger-km per day in North America will be provided by high-speed trains. If the traffic volume in a typical region of North America is 15,000, how many passenger-kilometers per day will high-speed trains provide there? (Data from Schafer, A., and D. Victor, "The Past and Future of Global Mobility," *Scientific American*.)

38. Johnny's car uses 15 gal of gasoline to travel 495 mi. He has 6 gal of gasoline in the car. He still needs to drive 600 mi. If the car continues to use gasoline at the same rate, how many more gallons will he need (to the nearest tenth)?

39. A river has a current of 4 km per hr. Find the rate of Herby's boat in still water if it travels 40 km downstream in the same time that it takes to travel 24 km upstream.
40. A sink can be filled by a cold-water tap in 8 min and by a hot-water tap in 12 min. How long would it take to fill the sink with both taps open?

	d	r	t
Upstream	24	$x - 4$	
Downstream	40		

	Rate	Time Working Together	Fractional Part of the Job Done
Cold		x	
Hot		x	

41. Jane and Jessica need to sort a pile of bottles at the recycling center. Working alone, Jane could do the entire job in 9 hr, while Jessica could do the entire job in 6 hr. How long will it take them if they work together?


6.6

42. In which one of the following does y vary inversely as x ?

A. $y = 2x$ B. $y = \frac{x}{3}$ C. $y = \frac{3}{x}$ D. $y = x^2$

Solve each problem.

43. For a particular camera, the viewing distance varies directly as the amount of enlargement. A picture that is taken with this camera and enlarged 5 times should be viewed from a distance of 250 mm. Suppose a print 8.6 times the size of the negative is made. From what distance should it be viewed?
44. The frequency (number of vibrations per second) of a vibrating guitar string varies inversely as its length. That is, a longer string vibrates fewer times in a second than a shorter string. Suppose a guitar string 0.65 m long vibrates 4.3 times per sec. What frequency would a string 0.5 m long have?
45. The volume of a rectangular box of a given height is jointly proportional to its width and length. A box with width 2 ft and length 4 ft has volume 12 ft³. Find the volume of a box with the same height that is 3 ft wide and 5 ft long.



Write each rational expression in lowest terms.

1. $\frac{x + 2y}{x^2 - 4y^2}$

2. $\frac{x^2 + 2x - 15}{x^2 - x - 6}$

Perform the indicated operations.

3. $\frac{2}{m} + \frac{5}{3m^2}$

4. $\frac{9}{3 - x} - \frac{2}{x - 3}$

5. $\frac{-\frac{3}{x} + \frac{x}{2}}{1 + \frac{x+1}{x}}$

6. $\frac{\frac{3}{x} - 5}{6 + \frac{1}{x}}$

7. $\frac{4y + 16}{30} \div \frac{2y + 8}{5}$

8. $\frac{t^{-2} + s^{-2}}{t^{-1} - s^{-1}}$

9. $\frac{k^2 - 6k + 9}{1 - 216k^3} \cdot \frac{6k^2 + 17k - 3}{9 - k^2}$

10. $\frac{9x^2 + 46x + 5}{3x^2 - 2x - 1} \div \frac{x^2 + 11x + 30}{x^3 + 5x^2 - 6x}$

11. $\frac{4a}{a^2 - ab - 2b^2} - \frac{6b - a}{a^2 + 4ab + 3b^2}$

12. $\frac{a}{b} + \frac{b}{c} + \frac{c}{d}$

Solve each equation.

13. $\frac{x + 3}{x^2 - 5x + 4} - \frac{1}{x} = \frac{2}{x^2 - 4x}$

14. $\frac{3x}{x - 4} + \frac{2}{x} = \frac{48}{x^2 - 4x}$

15. $1 - \frac{5}{r} = \frac{-4}{r^2}$

16. $A = \frac{Rr}{R + r}$ for r

Solve each problem.


17. The strength of a contact lens is given in units called diopters and also in millimeters of arc. As the diopters increase, the millimeters of arc decrease. The rational function

$$f(x) = \frac{337}{x}$$

relates the arc measurement $f(x)$ to the diopter measurement x . (Data from Bausch and Lomb.)

- (a) To the nearest hundredth, what arc measurement will correspond to 40.5-diopter lenses?
- (b) A lens with an arc measurement of 7.51 mm will provide what diopter strength, to the nearest hundredth?
18. At a certain gasoline station, 3 gal of unleaded gasoline cost \$12.27. How much would 13 gal of the same gasoline cost?
19. If Dr. Dawson rides his bike to his office, he averages 12 mph. If he drives his car, he averages 36 mph. His time driving is $\frac{1}{4}$ hr less than his time riding his bike. How far is his office from home?

20. The area of a triangle varies jointly as the lengths of the base and height. A triangle with base 10 ft and height 4 ft has area 20 ft
- ²
- . Find the area of a triangle with base 3 ft and height 8 ft.



Chapter 6

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

► View the complete solutions to all Chapter Test exercises in MyLabMath.

1. Give the domain of the following rational function using (a) set-builder notation and (b) interval notation.

$$f(x) = \frac{x + 3}{3x^2 + 2x - 8}$$

2. Write
- $\frac{6x^2 - 13x - 5}{9x^3 - x}$
- in lowest terms.

Multiply or divide as indicated.

3. $\frac{(x + 3)^2}{4} \cdot \frac{6}{2x + 6}$

4. $\frac{y^2 - 16}{y^2 - 25} \cdot \frac{y^2 + 2y - 15}{y^2 - 7y + 12}$

5. $\frac{3 - t}{5} \div \frac{t - 3}{10}$

6. $\frac{x^2 - 9}{x^3 + 3x^2} \div \frac{x^2 + x - 12}{x^3 + 9x^2 + 20x}$

7. Find the least common denominator for the group of denominators.

$$t^2 + t - 6, \quad t^2 + 3t, \quad t^2$$

Add or subtract as indicated.

8. $\frac{7}{6t^2} - \frac{1}{3t}$

9. $\frac{9}{x-7} + \frac{4}{x+7}$

10. $\frac{9}{x^2 - 6x + 9} + \frac{2}{x^2 - 9}$

11. $\frac{6}{x+4} + \frac{1}{x+2} - \frac{3x}{x^2 + 6x + 8}$

Simplify each expression.

12. $\frac{\frac{12}{r+4}}{\frac{11}{6r+24}}$

13. $\frac{\frac{1}{a} - \frac{1}{b}}{\frac{a}{b} - \frac{b}{a}}$

14. $\frac{2x^{-2} + y^{-2}}{x^{-1} - y^{-1}}$

15. Decide whether each of the following is an *expression* or an *equation*. Simplify the one that is an expression, and solve the one that is an equation.

(a) $\frac{2x}{3} + \frac{x}{4} - \frac{11}{2}$

(b) $\frac{2x}{3} + \frac{x}{4} = \frac{11}{2}$

Solve each equation.

16. $\frac{1}{x} - \frac{4}{3x} = \frac{1}{x-2}$

17. $\frac{y}{y+2} - \frac{1}{y-2} = \frac{8}{y^2-4}$

18. Solve for the variable ℓ in this formula from mathematics: $S = \frac{n}{2}(a + \ell)$.

19. Graph the rational function $f(x) = \frac{-2}{x+1}$. Give the equations of the vertical and horizontal asymptotes.

Solve each problem.

20. Chris can do a job in 9 hr, while Dana can do the same job in 5 hr. How long would it take them to do the job if they worked together?
21. The rate of the current in a stream is 3 mph. Nana's boat can travel 36 mi downstream in the same time that it takes to travel 24 mi upstream. Find the rate of her boat in still water.
22. Biologists collected a sample of 600 fish from West Okoboji Lake on May 1 and tagged each of them. When they returned on June 1, a new sample of 800 fish was collected and 10 of these had been previously tagged. Use this experiment to determine the approximate fish population of West Okoboji Lake.
23. In biology, the rational function

$$g(x) = \frac{5x}{2+x}$$

gives the growth rate g of a population for x units of available food. (Data from Smith, J. Maynard, *Models in Ecology*, Cambridge University Press.)

- (a) What amount of food (in appropriate units) would produce a growth rate of 3 units of growth per unit of food?
- (b) What is the growth rate if no food is available?

24. The current in a simple electrical circuit is inversely proportional to the resistance. If the current is 80 amps when the resistance is 30 ohms, find the current when the resistance is 12 ohms.
25. The force of the wind blowing on a vertical surface varies jointly as the area of the surface and the square of the velocity. If a wind blowing at 40 mph exerts a force of 50 lb on a surface of 500 ft², how much force will a wind of 80 mph exert on a surface of 2 ft²?

Chapters R–6 Cumulative Review Exercises

Perform the indicated operations.

1. $-\frac{2}{3} - \frac{1}{4}$ 2. $-\frac{3}{4} \left(-\frac{5}{6}\right)$ 3. $-\frac{3}{5} \div \frac{1}{2}$
 4. $-8.75 + 3.08$ 5. $5.25 \div 100$ 6. 43.6×100

Evaluate.


7. (a) 5^2 (b) -5^2 (c) $(-5)^2$ (d) $\sqrt{25}$ (e) $-\sqrt{25}$ (f) $\sqrt{-25}$
 8. Evaluate $|2x| + 3y - z^3$ for $x = -4$, $y = 3$, and $z = 6$.

Solve each equation or inequality.

9. $7(2x + 3) - 4(2x + 1) = 2(x + 1)$ 10. $|6x - 8| - 4 = 0$
 11. $\frac{2}{3}y + \frac{5}{12}y \leq 20$ 12. $|3x + 2| \geq 4$

Solve each problem.

13. Abushieba invested some money at 4% interest and twice as much at 3% interest. His interest for the first year was \$400. How much did he invest at each rate?
14. A triangle has area 42 m². The base is 14 m long. Find the height of the triangle.



15. Graph $-4x + 2y = 8$ and give the intercepts.

Find the slope of each line described.

16. Passing through $(-5, 8)$ and $(-1, 2)$ 17. Perpendicular to $4x - 3y = 12$
 18. Write an equation of the line in **Exercise 16**. Give the equation in the form $y = mx + b$.

Graph each inequality or compound inequality.

19. $2x + 5y > 10$ 20. $x - y \geq 3$ and $3x + 4y \leq 12$
 21. Consider the equation $5x - 3y = 8$.
 (a) Write y as a function f of x , using function notation $f(x)$.
 (b) Find $f(1)$.
 22. Consider the relation $y = -\sqrt{x + 2}$.
 (a) Does it define a function?
 (b) Give its domain and range.

Solve each system.

23. $4x - y = -7$
 $5x + 2y = 1$

24. $x + y - 2z = -1$
 $2x - y + z = -6$
 $3x + 2y - 3z = -3$

25. $x + 2y + z = 5$
 $x - y + z = 3$
 $2x + 4y + 2z = 11$

26. Simplify $\left(\frac{m^{-4}n^2}{m^2n^{-3}}\right) \cdot \left(\frac{m^5n^{-1}}{m^{-2}n^5}\right)$. Write the answer with only positive exponents. Assume that all variables represent nonzero real numbers.

Perform the indicated operations.

27. $(3y^2 - 2y + 6) - (-y^2 + 5y + 12)$

28. $(4f + 3)(3f - 1)$

29. $\left(\frac{1}{4}x + 5\right)^2$

30. $(3x^3 + 13x^2 - 17x - 7) \div (3x + 1)$

31. Find each of the following for the polynomial functions

$$f(x) = x^2 + 2x - 3, \quad g(x) = 2x^3 - 3x^2 + 4x - 1, \quad \text{and} \quad h(x) = x^2.$$

(a) $(f + g)(x)$ (b) $(g - f)(x)$ (c) $(f + g)(-1)$ (d) $(f \circ h)(x)$

Factor each polynomial completely.

32. $2x^2 - 13x - 45$

33. $100r^4 - 25$

34. $8p^3 + 125$

Perform the indicated operations. Express answers in lowest terms.

35. $\frac{2a^2}{a+b} \cdot \frac{a-b}{4a}$

36. $\frac{x^2 - 9}{2x + 4} \div \frac{x^3 - 27}{4}$

37. $\frac{x+4}{x-2} + \frac{2x-10}{x-2}$

Solve.

38. $3x^2 + 4x = 7$

39. $\frac{-3x}{x+1} + \frac{4x+1}{x} = \frac{-3}{x^2+x}$

40. Machine A can complete a certain job in 2 hr. To speed up the work, Machine B, which can complete the job alone in 3 hr, is brought in to help. How long will it take the two machines to complete the job working together?

STUDY SKILLS REMINDER

Have you begun to prepare for your final exam? **Review Study Skill 10, Preparing for Your Math Final Exam.**

This page intentionally left blank



ROOTS, RADICALS, AND ROOT FUNCTIONS

The formula for calculating the distance one can see to the horizon from the top of a tall building involves a *square root radical*, one of the topics covered in this chapter.

- 7.1** Radical Expressions and Graphs
- 7.2** Rational Exponents
- 7.3** Simplifying Radicals, the Distance Formula, and Circles
- 7.4** Adding and Subtracting Radical Expressions

- 7.5** Multiplying and Dividing Radical Expressions

SUMMARY EXERCISES Performing Operations with Radicals and Rational Exponents

- 7.6** Solving Equations with Radicals
- 7.7** Complex Numbers

7.1 Radical Expressions and Graphs

OBJECTIVES

- 1 Find roots of numbers.
- 2 Find principal roots.
- 3 Graph functions defined by radical expressions.
- 4 Find n th roots of n th powers.
- 5 Use a calculator to find roots.

OBJECTIVE 1 Find roots of numbers.

Recall that $6^2 = 36$. We say that “6 squared equals 36.” The opposite (or inverse) of *squaring* a number is taking its *square root*.

It is customary to write $\sqrt{\quad}$ rather than $\sqrt[2]{\quad}$. $\sqrt{36} = 6$, because $6^2 = 36$.


We extend this discussion to *cube roots* $\sqrt[3]{\quad}$, *fourth roots* $\sqrt[4]{\quad}$, and higher roots.

Meaning of $\sqrt[n]{a}$

The n th root of a , written $\sqrt[n]{a}$, is a number whose n th power equals a . That is,

$$\sqrt[n]{a} = b \text{ means } b^n = a.$$

The number a is the **radicand**, n is the **index**, or **order**, and the expression $\sqrt[n]{a}$ is a **radical**.



VOCABULARY

- radicand
- index (order)
- radical
- principal root
- radical expression
- square root function
- cube root function

NOW TRY EXERCISE 1

Find each root.

- (a) $\sqrt[3]{1000}$ (b) $\sqrt[4]{625}$
 (c) $\sqrt[4]{\frac{1}{256}}$ (d) $\sqrt[3]{0.027}$

EXAMPLE 1 Simplifying Higher Roots

Find each root.

- (a) $\sqrt[3]{64} = 4$, because $4^3 = 64$. (b) $\sqrt[3]{125} = 5$, because $5^3 = 125$.
 (c) $\sqrt[4]{16} = 2$, because $2^4 = 16$. (d) $\sqrt[5]{32} = 2$, because $2^5 = 32$.
 (e) $\sqrt[3]{\frac{8}{27}} = \frac{2}{3}$, because $\left(\frac{2}{3}\right)^3 = \frac{8}{27}$. (f) $\sqrt[4]{0.0016} = 0.2$, because $(0.2)^4 = 0.0016$.

NOW TRY

OBJECTIVE 2 Find principal roots.

If n is even, positive numbers have two n th roots. For example, both 4 and -4 are square roots of 16, and 2 and -2 are fourth roots of 16. For $a > 0$, $\sqrt[n]{a}$ represents the positive root, called the **principal root**, and $-\sqrt[n]{a}$ represents the negative root. For all n , $\sqrt[n]{0} = 0$.

n th Root

Case 1 If n is *even* and a is *positive or 0*, then

$\sqrt[n]{a}$ represents the **principal n th root** of a ,

and $-\sqrt[n]{a}$ represents the **negative n th root** of a .

Case 2 If n is *even* and a is *negative*, then $\sqrt[n]{a}$ is not a real number.

Case 3 If n is *odd*, then there is exactly one real n th root of a , written $\sqrt[n]{a}$.

NOW TRY ANSWERS

1. (a) 10 (b) 5 (c) $\frac{1}{4}$ (d) 0.3

If n is even (Case 1), then the two n th roots of a are often written together as $\pm\sqrt[n]{a}$, with \pm read “*positive or negative*” or “*plus or minus*.”

NOW TRY
EXERCISE 2

Find each root.

- (a) $-\sqrt{25}$ (b) $\sqrt[4]{-625}$
(c) $\sqrt[5]{-32}$ (d) $-\sqrt[3]{64}$

EXAMPLE 2 Finding Roots

Find each root.

(a) $\sqrt{100} = 10$ (Case 1)

Because the radicand, 100, is *positive*, there are two square roots: 10 and -10 . We want the principal square root, which is 10.

(b) $-\sqrt{100} = -10$ (Case 1)

Here, we want the negative square root, -10 .

(c) $\sqrt[4]{81} = 3$ Principal 4th root (Case 1)

(d) $-\sqrt[4]{81} = -3$ Negative 4th root (Case 1)

(e) $\sqrt[4]{-81}$ (Case 2)

The index is *even* and the radicand is *negative*. Any real number raised to an *even* power is *positive*—here $3^4 = 81$ and $(-3)^4 = 81$. Thus,

$$\sqrt[4]{-81} \text{ is not a real number.}$$

(f) $\sqrt[3]{8} = 2$, because $2^3 = 8$. (Case 3)

(g) $\sqrt[3]{-8} = -2$, because $(-2)^3 = -8$. (Case 3)

In parts (f) and (g), the index is *odd*. Each radical represents exactly one *n*th root (regardless of whether the radicand is positive, negative, or 0).

NOW TRY

OBJECTIVE 3 Graph functions defined by radical expressions.

A **radical expression** is an algebraic expression that contains radicals.


$$3 - \sqrt{x}, \quad \sqrt[3]{x}, \quad \sqrt{2x - 1} \quad \text{Radical expressions}$$

In earlier chapters, we graphed functions defined by polynomial and rational expressions. Now we graph functions defined by basic radical expressions, such as $f(x) = \sqrt{x}$ and $f(x) = \sqrt[3]{x}$.

FIGURE 1 shows the graph of the **square root function**,

$$f(x) = \sqrt{x},$$

together with a table of selected points. Only nonnegative values can be used for x , so the domain is $[0, \infty)$. Because \sqrt{x} is the principal square root of x , it always has a nonnegative value, so the range is also $[0, \infty)$.



Square root function

$$f(x) = \sqrt{x}$$

Domain: $[0, \infty)$

Range: $[0, \infty)$

NOW TRY ANSWERS


2. (a) -5
(b) It is not a real number.
(c) -2 (d) -4

FIGURE 1

FIGURE 2 shows the graph of the **cube root function**

$$f(x) = \sqrt[3]{x}.$$

Any real number (positive, negative, or 0) can be used for x in the cube root function, so $\sqrt[3]{x}$ can be positive, negative, or 0. Thus, both the domain and the range of the cube root function are $(-\infty, \infty)$.



Cube root function

$$f(x) = \sqrt[3]{x}$$

Domain: $(-\infty, \infty)$

Range: $(-\infty, \infty)$

FIGURE 2

NOW TRY EXERCISE 3

Graph each function, and give its domain and range.

(a) $f(x) = \sqrt{x+1}$

(b) $f(x) = \sqrt[3]{x} - 1$

EXAMPLE 3 Graphing Functions Defined with Radicals

Graph each function, and give its domain and range.


(a) $f(x) = \sqrt{x-3}$

Create a table of values as given with the graph in FIGURE 3. The x -values were chosen in such a way that the function values are all integers. For the radicand to be nonnegative, we must have

$$x - 3 \geq 0, \text{ or } x \geq 3.$$

Therefore, the domain of this function is $[3, \infty)$. Function values are positive or 0, so the range is $[0, \infty)$.


x	$f(x) = \sqrt{x-3}$
3	$\sqrt{3-3} = 0$
4	$\sqrt{4-3} = 1$
7	$\sqrt{7-3} = 2$




This graph is shifted to the right 3 units compared to the graph of $y = \sqrt{x}$.

FIGURE 3

NOW TRY ANSWERS



domain: $[-1, \infty)$
range: $[0, \infty)$




domain: $(-\infty, \infty)$
range: $(-\infty, \infty)$

(b) $f(x) = \sqrt[3]{x} + 2$

See FIGURE 4. Both the domain and the range are $(-\infty, \infty)$.

x	$f(x) = \sqrt[3]{x} + 2$
-8	$\sqrt[3]{-8} + 2 = 0$
-1	$\sqrt[3]{-1} + 2 = 1$
0	$\sqrt[3]{0} + 2 = 2$
1	$\sqrt[3]{1} + 2 = 3$
8	$\sqrt[3]{8} + 2 = 4$



This graph is shifted up 2 units compared to the graph of $y = \sqrt[3]{x}$.

FIGURE 4

OBJECTIVE 4 Find n th roots of n th powers.

Consider the expression $\sqrt{a^2}$. At first glance, we might think that it is equivalent to a . However, this is not necessarily true. For example, consider the following.

$$\text{If } a = 6, \quad \text{then } \sqrt{a^2} = \sqrt{6^2} = \sqrt{36} = 6.$$

$$\text{If } a = -6, \quad \text{then } \sqrt{a^2} = \sqrt{(-6)^2} = \sqrt{36} = 6. \quad \leftarrow \text{Instead of } -6, \text{ we get } 6, \text{ the absolute value of } -6.$$

The symbol $\sqrt{a^2}$ represents the *nonnegative* square root, so we express $\sqrt{a^2}$ with absolute value bars as $|a|$ because a may be a negative number.

Meaning of $\sqrt{a^2}$

$$\text{For any real number } a, \quad \sqrt{a^2} = |a|.$$

That is, the principal square root of a^2 is the absolute value of a .

NOW TRY EXERCISE 4

Find each square root.

(a) $\sqrt{11^2}$ (b) $\sqrt{(-11)^2}$
 (c) $\sqrt{z^2}$ (d) $\sqrt{(-z)^2}$

EXAMPLE 4 Simplifying Square Roots Using Absolute Value

Find each square root.

(a) $\sqrt{7^2} = |7| = 7$ (b) $\sqrt{(-7)^2} = |-7| = 7$

(c) $\sqrt{k^2} = |k|$ (d) $\sqrt{(-k)^2} = |-k| = |k|$

NOW TRY

We can generalize this idea to any n th root.

Meaning of $\sqrt[n]{a^n}$

If n is an *even* positive integer, then $\sqrt[n]{a^n} = |a|$.

If n is an *odd* positive integer, then $\sqrt[n]{a^n} = a$.

That is, use the absolute value symbol when n is even.

NOW TRY EXERCISE 5

Simplify each root.

(a) $\sqrt[8]{(-2)^8}$ (b) $\sqrt[3]{(-9)^3}$
 (c) $-\sqrt[4]{(-10)^4}$ (d) $-\sqrt{m^8}$
 (e) $\sqrt[3]{x^{18}}$ (f) $\sqrt[4]{t^{20}}$

EXAMPLE 5 Simplifying Higher Roots Using Absolute Value

Simplify each root.

(a) $\sqrt[6]{(-3)^6} = |-3| = 3$ n is even. Use absolute value.

(b) $\sqrt[5]{(-4)^5} = -4$ n is odd. Absolute value is not necessary.

(c) $-\sqrt[4]{(-9)^4} = -|-9| = -9$ n is even. Use absolute value.

(d) $-\sqrt{m^4} = -|m^2| = -m^2$ For all m , $|m^2| = m^2$.

No absolute value bars are needed here, because m^2 is nonnegative for any real number value of m .


(e) $\sqrt[3]{a^{12}} = a^4$, because $a^{12} = (a^4)^3$.

(f) $\sqrt[4]{x^{12}} = |x^3|$


We use absolute value to guarantee that the result is not negative (because x^3 is negative when x is negative). If desired, $|x^3|$ can be written as $x^2 \cdot |x|$.

NOW TRY**NOW TRY ANSWERS**

4. (a) 11 (b) 11 (c) $|z|$
 (d) $|z|$
 5. (a) 2 (b) -9 (c) -10
 (d) $-m^4$ (e) x^6 (f) $|t^5|$



(a)



(b)

FIGURE 5

OBJECTIVE 5 Use a calculator to find roots.

While numbers such as $\sqrt{9}$ and $\sqrt[3]{-8}$ are rational, radicals are often irrational numbers. To find approximations of such radicals, we usually use a calculator. For example,

$$\sqrt{15} \approx 3.872983346, \quad \sqrt[3]{10} \approx 2.15443469, \quad \text{and} \quad \sqrt[4]{2} \approx 1.189207115,$$

where the symbol \approx means “is approximately equal to.” In this text, we often show approximations rounded to three decimal places. Thus,

$$\sqrt{15} \approx 3.873, \quad \sqrt[3]{10} \approx 2.154, \quad \text{and} \quad \sqrt[4]{2} \approx 1.189.$$

FIGURE 5 shows how the preceding approximations are displayed on a TI-83/84 Plus graphing calculator. In FIGURE 5(a), eight or nine decimal places are shown, while in FIGURE 5(b), the number of decimal places is fixed at three.

There is a simple way to check that a calculator approximation is “in the ballpark.” For example, because 16 is a little larger than 15, $\sqrt{16} = 4$ should be a little larger than $\sqrt{15}$. Thus, 3.873 is reasonable as an approximation for $\sqrt{15}$.

NOTE Methods for finding approximations differ among makes and models of calculators. **Consult your owner’s manual for keystroke instructions.** Be aware that graphing calculators often differ from scientific calculators in the order in which keystrokes are made.

NOW TRY EXERCISE 6

Use a calculator to approximate each radical to three decimal places.

- (a) $\sqrt{73}$ (b) $-\sqrt{92}$
 (c) $\sqrt[4]{92}$ (d) $\sqrt[5]{33}$

NOW TRY EXERCISE 7

Use the formula in Example 7 to approximate f to the nearest thousand if

$$L = 7 \times 10^{-5}$$

and $C = 3 \times 10^{-9}$.

NOW TRY ANSWERS

6. (a) 8.544 (b) -9.592
 (c) 3.097 (d) 2.012
 7. 347,000 cycles per sec

EXAMPLE 6 Finding Approximations for Roots

Use a calculator to approximate each radical to three decimal places.

- (a) $\sqrt{39} \approx 6.245$ (b) $-\sqrt{72} \approx -8.485$
 (c) $\sqrt[3]{93} \approx 4.531$ (d) $\sqrt[4]{39} \approx 2.499$

NOW TRY

EXAMPLE 7 Using Roots to Calculate Resonant Frequency

In electronics, the resonant frequency f of a circuit may be found using the formula

$$f = \frac{1}{2\pi\sqrt{LC}}, \quad \text{Electronics formula}$$

where f is in cycles per second, L is in henrys, and C is in farads. (Henrys and farads are units of measure in electronics.) Find the resonant frequency f if $L = 5 \times 10^{-4}$ henry and $C = 3 \times 10^{-10}$ farad. Give the answer to the nearest thousand.

Find the value of f when $L = 5 \times 10^{-4}$ and $C = 3 \times 10^{-10}$.

$$f = \frac{1}{2\pi\sqrt{LC}} \quad \text{Given formula}$$

$$f = \frac{1}{2\pi\sqrt{(5 \times 10^{-4})(3 \times 10^{-10})}} \quad \text{Substitute for } L \text{ and } C.$$

$$f \approx 411,000 \quad \text{Use a calculator.}$$

The resonant frequency f is approximately 411,000 cycles per sec.

NOW TRY

NOTE The expression in the second-to-last line of **Example 7** can be difficult to compute on a calculator. There are several ways to approach it, but here is one way that works nicely. Use a calculator to verify the following steps.

Step 1 Calculate the product under the radical in the denominator. The result is

$$1.5 \times 10^{-13}.$$

Step 2 Use the square root function (\sqrt{x}) to find the square root of this number. The result is approximately

$$3.872983346 \times 10^{-7}.$$

Step 3 Multiply by 2π , using the π function of the calculator. The result is approximately

$$2.433467206 \times 10^{-6}.$$

Step 4 Now use the reciprocal function (labeled $\frac{1}{x}$ or x^{-1}) of the calculator. The result should be 410936.296, which, rounded to the nearest thousand, is 411,000.

If the numerator had not been 1, and perhaps an expression to evaluate, one approach would be to save the result found in Step 3 in the calculator memory, evaluate the numerator, and then divide by the number saved in the memory.

7.1 Exercises

FOR
EXTRA
HELP



MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Make study cards to help you learn and remember the material in this chapter.

Review Study Skill 5,
Using Study Cards.

Concept Check Find each square root.

1. (a) $\sqrt{25}$ (b) $\sqrt{81}$ (c) $\sqrt{144}$ 2. (a) $\sqrt{0.25}$ (b) $\sqrt{0.81}$ (c) $\sqrt{1.44}$

3. **Concept Check** A student incorrectly claimed that $\sqrt{16} = 8$. **WHAT WENT WRONG?** Evaluate $\sqrt{16}$ correctly.

4. **Concept Check** When approximating $\sqrt{19}$ to three decimal places, a student incorrectly rounded as follows.

$$\sqrt{19} \approx 4.358898944$$

$$\sqrt{19} \approx 4.358$$

WHAT WENT WRONG? Give the correct approximation.

Concept Check Match each expression in Column I with the equivalent choice in Column II. Answers may be used once, more than once, or not at all.

I

II

5. $-\sqrt{16}$

6. $\sqrt{-16}$

A. 3

B. -2

7. $\sqrt[3]{-27}$

8. $\sqrt[5]{-32}$

C. 2

D. -3

9. $\sqrt[4]{16}$

10. $-\sqrt[3]{64}$

E. -4

F. Not a real number

Concept Check Choose the closest approximation of each square root. Do not use a calculator.

11. $\sqrt{123}$

12. $\sqrt{67}$

A. 9

B. 10

C. 11

D. 12


A. 7

B. 8

C. 9

D. 10

Concept Check Refer to the figure to answer each question.



13. Which one of the following is the best estimate of its area?
 A. 50 B. 100 C. 250 D. 2500
14. Which one of the following is the best estimate of its perimeter?
 A. 15 B. 30 C. 100 D. 250
15. **Concept Check** Consider the expression $-\sqrt{-a}$. Decide whether it is *positive, negative, 0, or not a real number* in each case.
 (a) $a > 0$ (b) $a < 0$ (c) $a = 0$
16. **Concept Check** If n is odd, under what conditions is $\sqrt[n]{a}$ the following?
 (a) positive (b) negative (c) 0

Find each root. See Examples 1 and 2.

- | | | | |
|-------------------------------|-------------------------------|-------------------------------|-------------------------------|
| 17. $-\sqrt{81}$ | 18. $-\sqrt{121}$ | 19. $\sqrt[3]{216}$ | 20. $\sqrt[3]{343}$ |
| 21. $\sqrt[3]{-64}$ | 22. $\sqrt[3]{-125}$ | 23. $-\sqrt[3]{512}$ | 24. $-\sqrt[3]{1000}$ |
| 25. $\sqrt[4]{1296}$ | 26. $\sqrt[4]{625}$ | 27. $-\sqrt[4]{16}$ | 28. $-\sqrt[4]{256}$ |
| 29. $\sqrt[4]{-625}$ | 30. $\sqrt[4]{-256}$ | 31. $\sqrt[6]{64}$ | 32. $\sqrt[6]{729}$ |
| 33. $\sqrt[6]{-32}$ | 34. $\sqrt[8]{-1}$ | 35. $\sqrt{\frac{64}{81}}$ | 36. $\sqrt{\frac{100}{9}}$ |
| 37. $\sqrt[3]{\frac{64}{27}}$ | 38. $\sqrt[4]{\frac{81}{16}}$ | 39. $-\sqrt[6]{\frac{1}{64}}$ | 40. $-\sqrt[5]{\frac{1}{32}}$ |
| 41. $-\sqrt[3]{-27}$ | 42. $-\sqrt[3]{-64}$ | 43. $\sqrt{0.25}$ | 44. $\sqrt{0.36}$ |
| 45. $-\sqrt{0.49}$ | 46. $-\sqrt{0.81}$ | 47. $\sqrt[3]{0.001}$ | 48. $\sqrt[3]{0.125}$ |

Graph each function, and give its domain and range. See Example 3.

- | | | |
|----------------------------|----------------------------|----------------------------|
| 49. $f(x) = \sqrt{x+3}$ | 50. $f(x) = \sqrt{x-5}$ | 51. $f(x) = \sqrt{x-2}$ |
| 52. $f(x) = \sqrt{x+4}$ | 53. $f(x) = \sqrt[3]{x-3}$ | 54. $f(x) = \sqrt[3]{x+1}$ |
| 55. $f(x) = \sqrt[3]{x-3}$ | 56. $f(x) = \sqrt[3]{x+1}$ | |

Simplify each root. See Examples 4 and 5.

- | | | | |
|-------------------------|-------------------------|------------------------|------------------------|
| 57. $\sqrt{12^2}$ | 58. $\sqrt{19^2}$ | 59. $\sqrt{(-10)^2}$ | 60. $\sqrt{(-13)^2}$ |
| 61. $\sqrt[6]{(-2)^6}$ | 62. $\sqrt[6]{(-4)^6}$ | 63. $\sqrt[5]{(-9)^5}$ | 64. $\sqrt[5]{(-8)^5}$ |
| 65. $-\sqrt[6]{(-5)^6}$ | 66. $-\sqrt[6]{(-7)^6}$ | 67. $\sqrt{x^2}$ | 68. $-\sqrt{x^2}$ |
| 69. $\sqrt{(-z)^2}$ | 70. $\sqrt{(-q)^2}$ | 71. $\sqrt[3]{x^3}$ | 72. $-\sqrt[3]{x^3}$ |
| 73. $\sqrt[3]{x^{15}}$ | 74. $\sqrt[3]{m^9}$ | 75. $\sqrt[6]{x^{30}}$ | 76. $\sqrt[4]{k^{20}}$ |

Use a calculator to approximate each radical to three decimal places. See Example 6.

77. $\sqrt{9483}$ 78. $\sqrt{6825}$ 79. $\sqrt{284.361}$ 80. $\sqrt{846.104}$
 81. $-\sqrt{82}$ 82. $-\sqrt{91}$ 83. $\sqrt[3]{423}$ 84. $\sqrt[3]{555}$
 85. $\sqrt[4]{100}$ 86. $\sqrt[4]{250}$ 87. $\sqrt[5]{23.8}$ 88. $\sqrt[5]{98.4}$

Solve each problem. See Example 7.

89. Use the electronics formula

$$f = \frac{1}{2\pi\sqrt{LC}}$$

to calculate the resonant frequency f of a circuit, in cycles per second, to the nearest thousand for the following values of L and C .

- (a) $L = 7.237 \times 10^{-5}$ and $C = 2.5 \times 10^{-10}$
 (b) $L = 5.582 \times 10^{-4}$ and $C = 3.245 \times 10^{-9}$
90. The threshold weight T for a person is the weight above which the risk of death increases greatly. The threshold weight in pounds for men aged 40–49 is related to height h in inches by the formula

$$h = 12.3\sqrt[3]{T}.$$

What height corresponds to a threshold weight of 216 lb for a 43-year-old man? Round the answer to the nearest inch and then to the nearest tenth of a foot.

91. According to an article in *The World Scanner Report*, the distance D , in miles, to the horizon from an observer's point of view over water or "flat" earth is given by

$$D = \sqrt{2H},$$




where H is the height of the point of view, in feet. If a person whose eyes are 6 ft above ground level is standing at the top of a hill 44 ft above "flat" earth, approximately how far to the horizon will she be able to see?

92. The time for one complete swing of a simple pendulum is given by

$$t = 2\pi\sqrt{\frac{L}{g}},$$


where t is time in seconds, L is the length of the pendulum in feet, and g , the force due to gravity, is about 32 ft per sec². Find the time of a complete swing of a 2-ft pendulum to the nearest tenth of a second.



The coefficient of self-induction L (in henrys), the energy P stored in an electronic circuit (in joules), and the current I (in amps) are related by the formula

$$I = \sqrt{\frac{2P}{L}}.$$

93. Find I if $P = 120$ and $L = 80$.
 94. Find I if $P = 100$ and $L = 40$.



Heron's formula gives a method of finding the area of a triangle if the lengths of its sides are known. Suppose that a , b , and c are the lengths of the sides. Let s denote one-half of the perimeter of the triangle (called the **semiperimeter**)—that is,


$$s = \frac{1}{2}(a + b + c).$$

Then the area of the triangle is given by


$$\mathcal{A} = \sqrt{s(s - a)(s - b)(s - c)}.$$

Use Heron's formula to solve each problem.

95. Find the area of the Bermuda Triangle, to the nearest thousand square miles, if the “sides” of this triangle measure approximately 960 mi, 1030 mi, and 1030 mi.



96. The Vietnam Veterans Memorial in Washington, DC, is in the shape of an unenclosed isosceles triangle with equal sides of length 246.75 ft. If the triangle were enclosed, the third side would have length 438.14 ft. Find the area of this enclosure to the nearest hundred square feet. (Data from information pamphlet obtained at the Vietnam Veterans Memorial.)



97. Find the area of a triangle with sides of lengths $a = 11$ m, $b = 60$ m, and $c = 61$ m.
98. Find the area of a triangle with sides of lengths $a = 20$ ft, $b = 34$ ft, and $c = 42$ ft.

7.2 Rational Exponents

OBJECTIVES

- 1 Use exponential notation for n th roots.
- 2 Define and use expressions of the form $a^{m/n}$.
- 3 Convert between radicals and rational exponents.
- 4 Use the rules for exponents with rational exponents.

OBJECTIVE 1 Use exponential notation for n th roots.

Consider the expression $(3^{1/2})^2$. We can simplify it as follows.

$$\begin{aligned} (3^{1/2})^2 &= 3^{1/2} \cdot 3^{1/2} & a^2 = a \cdot a \\ &= 3^{1/2+1/2} & \text{Product rule: } a^m \cdot a^n = a^{m+n} \\ &= 3^1 & \text{Add exponents.} \\ &= 3 & a^1 = a \end{aligned}$$

Also, by definition,

$$(\sqrt{3})^2 = \sqrt{3} \cdot \sqrt{3} = 3.$$

Because both $(3^{1/2})^2$ and $(\sqrt{3})^2$ are equal to 3, it seems reasonable to define

$$3^{1/2} = \sqrt{3}.$$

This suggests the following generalization.

Meaning of $a^{1/n}$

If $\sqrt[n]{a}$ is a real number, then $a^{1/n} = \sqrt[n]{a}$.

Examples: $4^{1/2} = \sqrt{4}$, $8^{1/3} = \sqrt[3]{8}$, $16^{1/4} = \sqrt[4]{16}$

Notice that the denominator of the rational exponent is the index of the radical.

NOW TRY EXERCISE 1

Evaluate each exponential.

- (a) $81^{1/2}$ (b) $125^{1/3}$
 (c) $-625^{1/4}$ (d) $(-625)^{1/4}$
 (e) $(-125)^{1/3}$ (f) $\left(\frac{1}{16}\right)^{1/4}$

EXAMPLE 1 Evaluating Exponentials of the Form $a^{1/n}$

Evaluate each exponential.

The denominator is the index, or root.

(a) $64^{1/3} = \sqrt[3]{64} = 4$

(c) $-256^{1/4} = -\sqrt[4]{256} = -4$

(d) $(-256)^{1/4} = \sqrt[4]{-256}$ is not a real number because the radicand, -256 , is negative and the index is even.

(e) $(-32)^{1/5} = \sqrt[5]{-32} = -2$

The denominator is the index, or root. $\sqrt{\quad}$ means $\sqrt[2]{\quad}$.

(b) $100^{1/2} = \sqrt{100} = 10$

(f) $\left(\frac{1}{8}\right)^{1/3} = \sqrt[3]{\frac{1}{8}} = \frac{1}{2}$

NOW TRY 

CAUTION Notice the distinction between **Examples 1(c) and 1(d)**. The radical in part (c) is the **negative fourth root of a positive number**, while the radical in part (d) is the **principal fourth root of a negative number, which is not a real number**.

OBJECTIVE 2 Define and use expressions of the form $a^{m/n}$.

We know that $8^{1/3} = \sqrt[3]{8}$. Now we can define a number like $8^{2/3}$, where the numerator of the exponent is not 1. For past rules of exponents to be valid,

$$8^{2/3} = 8^{(1/3)2} = (8^{1/3})^2.$$

Because $8^{1/3} = \sqrt[3]{8}$,

$$8^{2/3} = (\sqrt[3]{8})^2 = 2^2 = 4.$$

Generalizing from this example, we define $a^{m/n}$ as follows.

Meaning of $a^{m/n}$

If m and n are positive integers with m/n in lowest terms, then

$$a^{m/n} = (a^{1/n})^m,$$

provided that $a^{1/n}$ is a real number. If $a^{1/n}$ is not a real number, then $a^{m/n}$ is not a real number.

EXAMPLE 2 Evaluating Exponentials of the Form $a^{m/n}$

Evaluate each exponential.

Think:

$$36^{1/2} = \sqrt{36} = 6$$

(a) $36^{3/2} = (36^{1/2})^3 = 6^3 = 216$

Think:

$$125^{1/3} = \sqrt[3]{125} = 5$$

(b) $125^{2/3} = (125^{1/3})^2 = 5^2 = 25$

NOW TRY ANSWERS

1. (a) 9 (b) 5 (c) -5
 (d) It is not a real number.
 (e) -5 (f) $\frac{1}{2}$


NOW TRY EXERCISE 2

Evaluate each exponential.

- (a) $32^{2/5}$ (b) $8^{5/3}$
 (c) $-100^{3/2}$ (d) $(-125)^{4/3}$
 (e) $(-121)^{3/2}$

Be careful.
The base is 4.

$$(c) -4^{5/2} = -(4^{5/2}) = -(4^{1/2})^5 = -(2)^5 = -32$$

Because the base is 4, the negative sign is *not* affected by the exponent.

$$(d) (-27)^{2/3} = [(-27)^{1/3}]^2 = (-3)^2 = 9$$

Notice that in part (c) we first evaluate the exponential and then find its negative. In part (d), the $-$ sign is part of the base, -27 .

$$(e) (-100)^{3/2} = [(-100)^{1/2}]^3, \text{ which is not a real number, because}$$

$$(-100)^{1/2}, \text{ or } \sqrt{-100}, \text{ is not a real number. } \quad \text{NOW TRY } \img alt="Now Try icon" data-bbox="885 258 910 278"/>$$

Recall that for any natural number n ,

$$a^{-n} = \frac{1}{a^n} \quad (\text{where } a \neq 0). \quad \text{Definition of negative exponent}$$

When a rational exponent is negative, this earlier interpretation is applied.

Meaning of $a^{-m/n}$

If $a^{m/n}$ is a real number, then

$$a^{-m/n} = \frac{1}{a^{m/n}} \quad (\text{where } a \neq 0).$$


NOW TRY EXERCISE 3

Evaluate each exponential.

- (a) $243^{-3/5}$ (b) $4^{-5/2}$
 (c) $\left(\frac{216}{125}\right)^{-2/3}$

EXAMPLE 3 Evaluating Exponentials of the Form $a^{-m/n}$

Evaluate each exponential.

$$(a) 16^{-3/4} = \frac{1}{16^{3/4}} = \frac{1}{(16^{1/4})^3} = \frac{1}{(\sqrt[4]{16})^3} = \frac{1}{2^3} = \frac{1}{8}$$

The denominator of $3/4$ is the index and the numerator is the exponent.

$$(b) 25^{-3/2} = \frac{1}{25^{3/2}} = \frac{1}{(25^{1/2})^3} = \frac{1}{(\sqrt{25})^3} = \frac{1}{5^3} = \frac{1}{125}$$

$$(c) \left(\frac{8}{27}\right)^{-2/3} = \frac{1}{\left(\frac{8}{27}\right)^{2/3}} = \frac{1}{\left(\sqrt[3]{\frac{8}{27}}\right)^2} = \frac{1}{\left(\frac{2}{3}\right)^2} = \frac{1}{\frac{4}{9}} = \frac{9}{4}$$

$$\frac{1}{\frac{4}{9}} = 1 \div \frac{4}{9} = 1 \cdot \frac{9}{4}$$

We can also use the rule $\left(\frac{b}{a}\right)^{-m} = \left(\frac{a}{b}\right)^m$ here, as follows.

$$\left(\frac{8}{27}\right)^{-2/3} = \left(\frac{27}{8}\right)^{2/3} = \left(\sqrt[3]{\frac{27}{8}}\right)^2 = \left(\frac{3}{2}\right)^2 = \frac{9}{4}$$

The result is the same.

NOW TRY ANSWERS

2. (a) 4 (b) 32 (c) -1000
 (d) 625
 (e) It is not a real number.
 3. (a) $\frac{1}{27}$ (b) $\frac{1}{32}$ (c) $\frac{25}{36}$

Take the reciprocal of only the base, *not* the exponent.

! CAUTION Be careful to distinguish between exponential expressions like the following.

$$16^{-1/4}, \text{ which equals } \frac{1}{2}, \quad -16^{1/4}, \text{ which equals } -2, \quad \text{and} \quad -16^{-1/4}, \text{ which equals } -\frac{1}{2}$$

A negative exponent does not necessarily lead to a negative result. Negative exponents lead to reciprocals, which may be positive.

We obtain an alternative definition of $a^{m/n}$ by applying the power rule for exponents differently than in the earlier definition.

Alternative Meaning of $a^{m/n}$

If all indicated roots are real numbers, then

$$a^{m/n} = (a^{1/n})^m = (a^m)^{1/n}.$$

As a result, we can evaluate an expression such as $27^{2/3}$ in two ways.

$$27^{2/3} = (27^{1/3})^2 = 3^2 = 9$$

or

$$27^{2/3} = (27^2)^{1/3} = 729^{1/3} = 9$$

The result is the same.

In most cases, it is easier to use $(a^{1/n})^m$.

Radical Form of $a^{m/n}$

If all indicated roots are real numbers, then

$$a^{m/n} = \sqrt[n]{a^m} = (\sqrt[n]{a})^m.$$

That is, raise a to the m th power and then take the n th root, or take the n th root of a and then raise it to the m th power.

For example, $8^{2/3} = \sqrt[3]{8^2} = \sqrt[3]{64} = 4$, and $8^{2/3} = (\sqrt[3]{8})^2 = 2^2 = 4$,

$$\text{so} \quad 8^{2/3} = \sqrt[3]{8^2} = (\sqrt[3]{8})^2.$$

NOW TRY EXERCISE 4

Write each exponential as a radical. Assume that all variables represent positive real numbers. Use the definition that takes the root first.

- (a) $21^{1/2}$ (b) $17^{5/4}$
 (c) $4t^{3/5} + (4t)^{2/3}$
 (d) $w^{-2/5}$ (e) $(a^2 - b^2)^{1/4}$

NOW TRY ANSWERS

4. (a) $\sqrt{21}$ (b) $(\sqrt[4]{17})^5$
 (c) $4(\sqrt[5]{t})^3 + (\sqrt[3]{4t})^2$
 (d) $\frac{1}{(\sqrt[5]{w})^2}$ (e) $\sqrt[4]{a^2 - b^2}$

OBJECTIVE 3 Convert between radicals and rational exponents.

Using the definition of rational exponents, we can simplify many problems involving radicals by converting the radicals to numbers with rational exponents. After simplifying, we can convert the answer back to radical form if required.

EXAMPLE 4 Converting Exponentials to Radicals

Write each exponential as a radical. Assume that all variables represent positive real numbers. Use the definition that takes the root first.

(a) $13^{1/2} = \sqrt{13}$ (b) $6^{3/4} = (\sqrt[4]{6})^3$ (c) $9m^{5/8} = 9(\sqrt[8]{m})^5$

(d) $6x^{2/3} - (4x)^{3/5} = 6(\sqrt[3]{x})^2 - (\sqrt[5]{4x})^3$

(e) $r^{-2/3} = \frac{1}{r^{2/3}} = \frac{1}{(\sqrt[3]{r})^2}$

(f) $(a^2 + b^2)^{1/2} = \sqrt{a^2 + b^2}$ ⚠ $\sqrt{a^2 + b^2} \neq a + b$


NOW TRY EXERCISE 5

Write each radical as an exponential and simplify. Assume that all variables represent positive real numbers.

(a) $\sqrt[3]{15}$ (b) $\sqrt[4]{4^2}$
 (c) $\sqrt[4]{x^4}$

EXAMPLE 5 Simplifying Radicals Using Rational Exponents

Write each radical as an exponential and simplify. Assume that all variables represent positive real numbers.

(a) $\sqrt{10} = 10^{1/2}$
 (b) $\sqrt{5^4} = 5^{4/2} = 5^2 = 25$
 (c) $\sqrt[4]{3^8} = 3^{8/4} = 3^2 = 9$
 (d) $\sqrt[6]{z^6} = z^{6/6} = z^1 = z$, because z is positive.

NOW TRY 

NOTE In **Example 5(d)**, it is not necessary to use absolute value bars because the directions specifically state that the variable represents a positive real number. The absolute value of the positive real number z is z itself, so the answer is simply z .

OBJECTIVE 4 Use the rules for exponents with rational exponents.

The definition of rational exponents allows us to apply the rules for exponents.

Rules for Rational Exponents

Let r and s be rational numbers. For all real numbers a and b for which the following are defined, these rules hold true.

Rules	Examples	Rules	Examples
$a^r \cdot a^s = a^{r+s}$	$5^{2/3} \cdot 5^{5/3} = 5^{2/3+5/3} = 5^{7/3}$	$\left(\frac{a}{b}\right)^r = \frac{a^r}{b^r}$	$\left(\frac{5}{7}\right)^{2/3} = \frac{5^{2/3}}{7^{2/3}}$
$\frac{a^r}{a^s} = a^{r-s}$	$\frac{4^{5/3}}{4^{4/3}} = 4^{5/3-4/3} = 4^{1/3}$	$a^{-r} = \frac{1}{a^r}$	$4^{-3/7} = \frac{1}{4^{3/7}}$
$(a^r)^s = a^{rs}$	$(5^{1/2})^{2/5} = 5^{1/2 \cdot 2/5} = 5^{1/5}$	$\frac{a^{-r}}{b^{-s}} = \frac{b^s}{a^r}$	$\frac{5^{-1/2}}{3^{-3/4}} = \frac{3^{3/4}}{5^{1/2}}$
$(ab)^r = a^r b^r$	$(3x)^{1/2} = 3^{1/2} x^{1/2}$		

EXAMPLE 6 Applying Rules for Rational Exponents

Simplify each expression. Assume that all variables represent positive real numbers.

(a) $2^{1/2} \cdot 2^{1/4}$
 $= 2^{1/2+1/4}$ **Product rule**
 $= 2^{2/4+1/4}$ **Write exponents with a common denominator.**
 $= 2^{3/4}$ **Add exponents.**

(b) $\frac{5^{2/3}}{5^{7/3}}$
 $= 5^{2/3-7/3}$ **Quotient rule**
 $= 5^{-5/3}$ **Subtract exponents.**
 $= \frac{1}{5^{5/3}}$ $a^{-r} = \frac{1}{a^r}$

NOW TRY ANSWERS

5. (a) $15^{1/3}$ (b) 2 (c) x


NOW TRY EXERCISE 6

Simplify each expression. Assume that all variables represent positive real numbers.

(a) $5^{1/4} \cdot 5^{2/3}$ (b) $\frac{9^{3/5}}{9^{7/5}}$

(c) $\frac{(r^{2/3}t^{1/4})^8}{t}$

(d) $\left(\frac{2x^{1/2}y^{-2/3}}{x^{-3/5}y^{-1/5}}\right)^{-3}$

(e) $y^{2/3}(y^{1/3} + y^{5/3})$

$$\begin{aligned} \text{(c)} \quad & \frac{(x^{1/2}y^{2/3})^4}{y} \\ &= \frac{(x^{1/2})^4(y^{2/3})^4}{y} && \text{Power rule} \\ &= \frac{x^2y^{8/3}}{y^1} && \text{Power rule; } y = y^1 \\ &= x^2y^{8/3-1} && \text{Quotient rule} \\ &= x^2y^{5/3} && \frac{8}{3} - 1 = \frac{8}{3} - \frac{3}{3} = \frac{5}{3} \end{aligned}$$

$$\begin{aligned} \text{(d)} \quad & \left(\frac{x^4y^{-6}}{x^{-2}y^{1/3}}\right)^{-2/3} \\ &= \frac{(x^4)^{-2/3}(y^{-6})^{-2/3}}{(x^{-2})^{-2/3}(y^{1/3})^{-2/3}} && \text{Power rule} \\ &= \frac{x^{-8/3}y^4}{x^{4/3}y^{-2/9}} && \text{Power rule} \\ &= x^{-8/3-4/3}y^{4-(-2/9)} && \text{Quotient rule} \\ &= x^{-4}y^{38/9} && 4 - \left(-\frac{2}{9}\right) = \frac{36}{9} + \frac{2}{9} = \frac{38}{9} \\ &= \frac{y^{38/9}}{x^4} && \text{Definition of negative exponent} \end{aligned}$$

Use parentheses to avoid errors.

Alternative method: We can simplify within the parentheses first, as follows.

$$\begin{aligned} & \left(\frac{x^4y^{-6}}{x^{-2}y^{1/3}}\right)^{-2/3} \\ &= (x^{4-(-2)}y^{-6-1/3})^{-2/3} && \text{Quotient rule} \\ &= (x^6y^{-19/3})^{-2/3} && -6 - \frac{1}{3} = -\frac{18}{3} - \frac{1}{3} = -\frac{19}{3} \\ &= (x^6)^{-2/3}(y^{-19/3})^{-2/3} && \text{Power rule} \\ &= x^{-4}y^{38/9} && \text{Power rule} \\ &= \frac{y^{38/9}}{x^4} && \text{Definition of negative exponent; The result is the same.} \end{aligned}$$

$$\begin{aligned} \text{(e)} \quad & m^{3/4}(m^{5/4} - m^{1/4}) \\ &= m^{3/4}(m^{5/4}) - m^{3/4}(m^{1/4}) && \text{Distributive property} \\ &= m^{3/4+5/4} - m^{3/4+1/4} && \text{Product rule} \\ &= m^{8/4} - m^{4/4} && \text{Add exponents.} \\ &= m^2 - m && \text{Write the exponents in lowest terms.} \end{aligned}$$

Do not make the common mistake of multiplying exponents in the first step.

NOW TRY ANSWERS

6. (a) $5^{11/12}$ (b) $\frac{1}{9^{4/5}}$
 (c) $r^{16/3}t$ (d) $\frac{y^{7/5}}{8x^{33/10}}$
 (e) $y + y^{7/3}$

CAUTION Use the rules of exponents in problems like those in **Example 6**. Do not convert the expressions to radical form.


NOW TRY EXERCISE 7

Write each radical as an exponential and simplify. Leave answers in exponential form. Assume that all variables represent positive real numbers.

(a) $\sqrt[5]{y^3} \cdot \sqrt[3]{y}$ (b) $\frac{\sqrt[4]{y^3}}{\sqrt[5]{y^5}}$
 (c) $\sqrt{\sqrt[3]{y}}$

NOW TRY ANSWERS

7. (a) $y^{14/15}$ (b) $\frac{1}{y^{7/4}}$ (c) $y^{1/6}$

EXAMPLE 7 Applying Rules for Rational Exponents

Write each radical as an exponential and simplify. Leave answers in exponential form. Assume that all variables represent positive real numbers.

(a) $\sqrt[3]{x^2} \cdot \sqrt[4]{x}$
 $= x^{2/3} \cdot x^{1/4}$ Convert to rational exponents.
 $= x^{2/3+1/4}$ Product rule
 $= x^{8/12+3/12}$ Write exponents with a common denominator.
 $= x^{11/12}$ Add exponents.

(b) $\frac{\sqrt{x^3}}{\sqrt[3]{x^2}}$
 $= \frac{x^{3/2}}{x^{2/3}}$ Convert to rational exponents.
 $= x^{3/2-2/3}$ Quotient rule
 $= x^{9/6-4/6}$ Write exponents with a common denominator.
 $= x^{5/6}$ Subtract exponents.


(c) $\sqrt{\sqrt[4]{z}}$
 $= \sqrt{z^{1/4}}$ Convert the inside radical to a rational exponent.
 $= (z^{1/4})^{1/2}$ Convert the square root to a rational exponent.
 $= z^{1/8}$ Power rule

NOW TRY 

7.2 Exercises

FOR
EXTRA
HELP


MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Match each expression in Column I with the equivalent choice in Column II.

I		II	
1. $3^{1/2}$	2. $(-27)^{1/3}$	A. -4	B. 8
3. $-16^{1/2}$	4. $(-25)^{1/2}$	C. $\sqrt{3}$	D. $-\sqrt{6}$
5. $(-32)^{1/5}$	6. $(-32)^{2/5}$	E. -3	F. $\sqrt{6}$
7. $4^{3/2}$	8. $6^{2/4}$	G. 4	H. -2
9. $-6^{2/4}$	10. $36^{0.5}$	I. 6	J. Not a real number

11. **Concept Check** A student incorrectly evaluated $27^{1/3}$ as 9. **WHAT WENT WRONG?** Evaluate $27^{1/3}$ correctly.

12. **Concept Check** A student evaluating $16^{-3/2}$ incorrectly suggested that the result is a negative number because the exponent is negative. **WHAT WENT WRONG?** Evaluate $16^{-3/2}$ correctly.

Evaluate each exponential. See Examples 1–3.

- | | | | |
|-------------------|-------------------|----------------------------------------|---------------------------------------|
| 13. $169^{1/2}$ | 14. $121^{1/2}$ | 15. $729^{1/3}$ | 16. $512^{1/3}$ |
| 17. $16^{1/4}$ | 18. $625^{1/4}$ | 19. $\left(\frac{64}{81}\right)^{1/2}$ | 20. $\left(\frac{8}{27}\right)^{1/3}$ |
| 21. $(-27)^{1/3}$ | 22. $(-32)^{1/5}$ | 23. $(-144)^{1/2}$ | 24. $(-36)^{1/2}$ |
| 25. $100^{3/2}$ | 26. $64^{3/2}$ | 27. $81^{3/4}$ | 28. $216^{2/3}$ |
| 29. $-16^{5/2}$ | 30. $-32^{3/5}$ | 31. $(-8)^{4/3}$ | 32. $(-243)^{2/5}$ |

$$\begin{array}{llll}
 33. 32^{-3/5} & 34. 27^{-4/3} & 35. 64^{-3/2} & 36. 81^{-3/2} \\
 37. \left(\frac{125}{27}\right)^{-2/3} & 38. \left(\frac{64}{125}\right)^{-2/3} & 39. \left(\frac{16}{81}\right)^{-3/4} & 40. \left(\frac{729}{64}\right)^{-5/6}
 \end{array}$$

Write each exponential as a radical. Assume that all variables represent positive real numbers. Use the definition that takes the root first. **See Example 4.**

$$\begin{array}{lll}
 41. 10^{1/2} & 42. 3^{1/2} & 43. 8^{3/4} \\
 44. 7^{2/3} & 45. 5x^{2/3} & 46. 8x^{3/4} \\
 47. 9q^{5/8} - (2x)^{2/3} & 48. (3p)^{3/4} + 4x^{1/3} & 49. x^{-3/5} \\
 50. z^{-4/9} & 51. (2y + x)^{2/3} & 52. (r + 2z)^{3/2}
 \end{array}$$

Write each radical as an exponential and simplify. Assume that all variables represent positive real numbers. **See Example 5.**

$$\begin{array}{lllll}
 53. \sqrt{15} & 54. \sqrt{26} & 55. \sqrt{2^{12}} & 56. \sqrt{5^{10}} & 57. \sqrt[3]{4^9} \\
 58. \sqrt[4]{6^8} & 59. \sqrt[8]{x^8} & 60. \sqrt{t^2} & 61. \sqrt{x^{20}} & 62. \sqrt{r^{50}}
 \end{array}$$

Simplify each expression. Assume that all variables represent positive real numbers. **See Example 6.**

$$\begin{array}{lll}
 63. 3^{1/2} \cdot 3^{3/2} & 64. 6^{4/3} \cdot 6^{2/3} & 65. \frac{64^{5/3}}{64^{4/3}} \\
 66. \frac{125^{7/3}}{125^{5/3}} & 67. y^{7/3} \cdot y^{-4/3} & 68. r^{-8/9} \cdot r^{17/9} \\
 69. x^{2/3} \cdot x^{-1/4} & 70. x^{2/5} \cdot x^{-1/3} & 71. \frac{k^{1/3}}{k^{2/3} \cdot k^{-1}} \\
 72. \frac{z^{3/4}}{z^{5/4} \cdot z^{-2}} & 73. \frac{(x^{1/4}y^{2/5})^{20}}{x^2} & 74. \frac{(r^{1/5}s^{2/3})^{15}}{r^2} \\
 75. \frac{(x^{2/3})^2}{(x^2)^{7/3}} & 76. \frac{(p^3)^{1/4}}{(p^{5/4})^2} & 77. \frac{m^{3/4}n^{-1/4}}{(m^2n)^{1/2}} \\
 78. \frac{a^{-1/2}b^{-5/4}}{(a^{-3}b^2)^{1/6}} & 79. \left(\frac{x^8y^{-8}}{x^{-4}y^{1/2}}\right)^{-2/3} & 80. \left(\frac{x^{2/3}y^{-6}}{x^{-1/12}y^{1/4}}\right)^{-4/5} \\
 81. \left(\frac{b^{-3/2}}{c^{-5/3}}\right)^2 (b^{-1/4}c^{-1/3})^{-1} & 82. \left(\frac{m^{-2/3}}{a^{-3/4}}\right)^4 (m^{-3/8}a^{1/4})^{-2} & 83. \left(\frac{p^{-1/4}q^{-3/2}}{3^{-1}p^{-2}q^{-2/3}}\right)^{-2} \\
 84. \left(\frac{2^{-2}w^{-3/4}x^{-5/8}}{w^{3/4}x^{-1/2}}\right)^{-3} & 85. p^{2/3}(p^{1/3} + 2p^{4/3}) & 86. z^{5/8}(3z^{5/8} + 5z^{11/8}) \\
 87. k^{1/4}(k^{3/2} - k^{1/2}) & 88. r^{3/5}(r^{1/2} + r^{3/4}) & 89. 6a^{7/4}(a^{-7/4} + 3a^{-3/4}) \\
 90. 4m^{5/3}(m^{-2/3} - 4m^{-5/3}) & 91. -5x^{7/6}(x^{5/6} - x^{-1/6}) & 92. -8y^{11/7}(y^{3/7} - y^{-4/7})
 \end{array}$$

Write each radical as an exponential and simplify. Leave answers in exponential form. Assume that all variables represent positive numbers. **See Example 7.**

$$\begin{array}{llll}
 93. \sqrt[5]{x^3} \cdot \sqrt[4]{x} & 94. \sqrt[6]{y^5} \cdot \sqrt[3]{y^2} & 95. \frac{\sqrt[3]{t^4}}{\sqrt[5]{t^4}} & 96. \frac{\sqrt[4]{w^3}}{\sqrt[6]{w}} \\
 97. \frac{\sqrt{x^5}}{\sqrt{x^8}} & 98. \frac{\sqrt[3]{k^5}}{\sqrt[3]{k^7}} & 99. \sqrt{y} \cdot \sqrt[3]{yz} & 100. \sqrt[3]{xz} \cdot \sqrt{z}
 \end{array}$$

101. $\sqrt[4]{\sqrt[3]{m}}$

102. $\sqrt[3]{\sqrt{k}}$

103. $\sqrt{\sqrt[3]{\sqrt[4]{x}}}$

104. $\sqrt[3]{\sqrt[5]{\sqrt{y}}}$

Concept Check Work each problem.

105. Replace a with 3 and b with 4 to show that, in general,

$$\sqrt{a^2 + b^2} \neq a + b.$$

106. Suppose someone claims that $\sqrt[n]{a^n + b^n}$ must equal $a + b$, because when $a = 1$ and $b = 0$, a true statement results:

$$\sqrt[n]{a^n + b^n} = \sqrt[n]{1^n + 0^n} = \sqrt[n]{1^n} = 1 = 1 + 0 = a + b.$$

Explain why this is faulty reasoning.

Solve each problem.

107. Meteorologists can determine the duration of a storm using the function

$$T(d) = 0.07d^{3/2},$$

where d is the diameter of the storm in miles and T is the time in hours. Find the duration of a storm with a diameter of 16 mi. Round the answer to the nearest tenth of an hour.

108. The threshold weight t , in pounds, for a person is the weight above which the risk of death increases greatly. The threshold weight in pounds for men aged 40–49 is related to height H in inches by the function

$$H(t) = (1860.867t)^{1/3}.$$

What height corresponds to a threshold weight of 200 lb for a 46-yr-old man? Round the answer to the nearest inch and then to the nearest tenth of a foot.

The **windchill factor** is a measure of the cooling effect that the wind has on a person's skin. It calculates the equivalent cooling temperature if there were no wind. The National Weather Service uses the formula

$$\text{Windchill temperature} = 35.74 + 0.6215T - 35.75V^{4/25} + 0.4275TV^{4/25},$$

where T is the temperature in $^{\circ}\text{F}$ and V is the wind speed in miles per hour, to calculate windchill. The table gives the windchill factor for various wind speeds and temperatures at which frostbite is a risk, and how quickly it may occur.



		Temperature ($^{\circ}\text{F}$)									
		Calm	40	30	20	10	0	-10	-20	-30	-40
Wind speed (mph)	5	36	25	13	1	-11	-22	-34	-46	-57	
	10	34	21	9	-4	-16	-28	-41	-53	-66	
	15	32	19	6	-7	-19	-32	-45	-58	-71	
	20	30	17	4	-9	-22	-35	-48	-61	-74	
	25	29	16	3	-11	-24	-37	-51	-64	-78	
	30	28	15	1	-12	-26	-39	-53	-67	-80	
	35	28	14	0	-14	-27	-41	-55	-69	-82	
40	27	13	-1	-15	-29	-43	-57	-71	-84		

Frostbites times: ■ 30 minutes ■ 10 minutes ■ 5 minutes

Data from National Oceanic and Atmospheric Administration, National Weather Service.

Use the formula to determine the windchill temperature to the nearest tenth of a degree, given the following conditions. Compare answers with the appropriate entries in the table.

109. 30°F , 15-mph wind

110. 10°F , 30-mph wind

111. 20°F , 20-mph wind

112. 40°F , 10-mph wind

7.3

Simplifying Radicals, the Distance Formula, and Circles

OBJECTIVES

- 1 Use the product rule for radicals.
- 2 Use the quotient rule for radicals.
- 3 Simplify radicals.
- 4 Simplify products and quotients of radicals with different indexes.
- 5 Use the Pythagorean theorem.
- 6 Use the distance formula.
- 7 Find an equation of a circle given its center and radius.

OBJECTIVE 1 Use the product rule for radicals.

Consider the expressions $\sqrt{36 \cdot 4}$ and $\sqrt{36} \cdot \sqrt{4}$.

$$\begin{aligned}\sqrt{36 \cdot 4} &= \sqrt{144} = 12 \\ \sqrt{36} \cdot \sqrt{4} &= 6 \cdot 2 = 12\end{aligned}$$

The result is the same.

This is an example of the **product rule for radicals**.

Product Rule for Radicals

If n is a natural number and $\sqrt[n]{a}$ and $\sqrt[n]{b}$ are real numbers, then the following holds true.

$$\sqrt[n]{a} \cdot \sqrt[n]{b} = \sqrt[n]{ab}$$

That is, the product of two n th roots is the n th root of the product.

We justify the product rule using the rules for rational exponents. Because $\sqrt[n]{a} = a^{1/n}$ and $\sqrt[n]{b} = b^{1/n}$,

$$\sqrt[n]{a} \cdot \sqrt[n]{b} = a^{1/n} \cdot b^{1/n} = (ab)^{1/n} = \sqrt[n]{ab}.$$

CAUTION Use the product rule only when the radicals have the same index.

EXAMPLE 1 Using the Product Rule

Multiply. Assume that all variables represent positive real numbers.

$$\begin{array}{lll} \text{(a)} \sqrt{5} \cdot \sqrt{7} & \text{(b)} \sqrt{11} \cdot \sqrt{p} & \text{(c)} \sqrt{7} \cdot \sqrt{11xyz} \\ = \sqrt{5 \cdot 7} & = \sqrt{11p} & = \sqrt{77xyz} \\ = \sqrt{35} & & \end{array}$$

NOW TRY 

EXAMPLE 2 Using the Product Rule

Multiply. Assume that all variables represent positive real numbers.

$$\begin{array}{lll} \text{(a)} \sqrt[3]{3} \cdot \sqrt[3]{12} & \text{(b)} \sqrt[4]{8y} \cdot \sqrt[4]{3r^2} & \text{(c)} \sqrt[6]{10m^4} \cdot \sqrt[6]{5m} \\ = \sqrt[3]{3 \cdot 12} & = \sqrt[4]{24yr^2} & = \sqrt[6]{50m^5} \\ = \sqrt[3]{36} & & \end{array}$$

Remember to write the index.

(d) $\sqrt[4]{2} \cdot \sqrt[5]{2}$ This product cannot be simplified using the product rule for radicals because the indexes (4 and 5) are different.

NOW TRY 

 NOW TRY EXERCISE 1

Multiply. Assume that all variables represent positive real numbers.

$$\begin{array}{l} \text{(a)} \sqrt{7} \cdot \sqrt{11} \\ \text{(b)} \sqrt{2mn} \cdot \sqrt{15} \end{array}$$

 NOW TRY EXERCISE 2

Multiply. Assume that all variables represent positive real numbers.

$$\begin{array}{ll} \text{(a)} \sqrt[3]{4} \cdot \sqrt[3]{5} & \text{(b)} \sqrt[4]{5t} \cdot \sqrt[4]{6r^3} \\ \text{(c)} \sqrt[7]{20x} \cdot \sqrt[7]{3xy^3} \\ \text{(d)} \sqrt[3]{5} \cdot \sqrt[4]{9} \end{array}$$

NOW TRY ANSWERS

1. (a) $\sqrt{77}$ (b) $\sqrt{30mn}$
2. (a) $\sqrt[3]{20}$ (b) $\sqrt[4]{30tr^3}$
- (c) $\sqrt[7]{60x^2y^3}$
- (d) This expression cannot be simplified using the product rule.

VOCABULARY

- hypotenuse
- legs (of a right triangle)
- circle
- center
- radius


NOW TRY EXERCISE 3

Simplify. Assume that all variables represent positive real numbers.

(a) $\sqrt{\frac{49}{36}}$ (b) $\sqrt{\frac{5}{144}}$

(c) $\sqrt[3]{-\frac{27}{1000}}$ (d) $\sqrt[4]{\frac{t}{16}}$

(e) $-\sqrt[5]{\frac{m^{15}}{243}}$

OBJECTIVE 2 Use the quotient rule for radicals.

The **quotient rule for radicals** is similar to the product rule.

Quotient Rule for Radicals

If n is a natural number and $\sqrt[n]{a}$ and $\sqrt[n]{b}$ are real numbers, then the following holds true.

$$\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}} \quad (\text{where } b \neq 0)$$

That is, the n th root of a quotient is the quotient of the n th roots.

EXAMPLE 3 Using the Quotient Rule

Simplify. Assume that all variables represent positive real numbers.

(a) $\sqrt{\frac{16}{25}} = \frac{\sqrt{16}}{\sqrt{25}} = \frac{4}{5}$

(b) $\sqrt{\frac{7}{36}} = \frac{\sqrt{7}}{\sqrt{36}} = \frac{\sqrt{7}}{6}$

(c) $\sqrt[3]{-\frac{8}{125}} = \sqrt[3]{\frac{-8}{125}} = \frac{\sqrt[3]{-8}}{\sqrt[3]{125}} = \frac{-2}{5} = -\frac{2}{5} \quad \frac{-a}{b} = -\frac{a}{b}$

(d) $\sqrt[3]{\frac{7}{216}} = \frac{\sqrt[3]{7}}{\sqrt[3]{216}} = \frac{\sqrt[3]{7}}{6}$

(e) $\sqrt[5]{\frac{x}{32}} = \frac{\sqrt[5]{x}}{\sqrt[5]{32}} = \frac{\sqrt[5]{x}}{2}$

(f) $-\sqrt[3]{\frac{m^6}{125}} = -\frac{\sqrt[3]{m^6}}{\sqrt[3]{125}} = -\frac{m^2}{5}$ Think: $\sqrt[3]{m^6} = m^{6/3} = m^2$

NOW TRY 

OBJECTIVE 3 Simplify radicals.

We use the product and quotient rules to simplify radicals. A radical is **simplified** if the following four conditions are met.

Conditions for a Simplified Radical

1. The radicand has no factor raised to a power greater than or equal to the index.
2. The radicand has no fractions.
3. No denominator contains a radical.
4. Exponents in the radicand and the index of the radical have greatest common factor 1.

Examples: $\sqrt{22}$, $\sqrt{15xy}$, $\sqrt[3]{18}$, $\frac{\sqrt[4]{m^3}}{m}$

These radicals are simplified.

$\sqrt{28}$, $\sqrt[3]{\frac{3}{5}}$, $\frac{7}{\sqrt{7}}$, $\sqrt[3]{r^{12}}$

These radicals are not simplified. Each violates one of the above conditions.

NOW TRY ANSWERS

3. (a) $\frac{7}{6}$ (b) $\frac{\sqrt{5}}{12}$ (c) $-\frac{3}{10}$
 (d) $\frac{\sqrt[4]{t}}{2}$ (e) $-\frac{m^3}{3}$


NOW TRY EXERCISE 4

Simplify.

- (a) $\sqrt{50}$ (b) $\sqrt{192}$
 (c) $\sqrt{42}$ (d) $\sqrt[3]{108}$
 (e) $-\sqrt[4]{80}$

EXAMPLE 4 Simplifying Roots of Numbers

Simplify.

(a) $\sqrt{24}$

Check to see whether 24 is divisible by a perfect square (the square of a natural number) such as 4, 9, 16, The greatest perfect square that divides into 24 is 4.

$$\begin{aligned}\sqrt{24} &= \sqrt{4 \cdot 6} && \text{Factor. 4 is a perfect square.} \\ &= \sqrt{4} \cdot \sqrt{6} && \text{Product rule} \\ &= 2\sqrt{6} && \sqrt{4} = 2\end{aligned}$$

(b) $\sqrt{108}$

As shown on the left, the number 108 is divisible by the perfect square 36. If this perfect square is not immediately clear, try factoring 108 into its prime factors, as shown on the right.

$\begin{aligned}\sqrt{108} &= \sqrt{36 \cdot 3} && \text{Factor.} \\ &= \sqrt{36} \cdot \sqrt{3} && \text{Product rule} \\ &= 6\sqrt{3} && \sqrt{36} = 6\end{aligned}$	$\begin{aligned}\sqrt{108} &= \sqrt{2^2 \cdot 3^3} && \text{Factor into prime factors.} \\ &= \sqrt{2^2 \cdot 3^2 \cdot 3} && a^3 = a^2 \cdot a \\ &= \sqrt{2^2} \cdot \sqrt{3^2} \cdot \sqrt{3} && \text{Product rule} \\ &= 2 \cdot 3 \cdot \sqrt{3} && \sqrt{2^2} = 2, \sqrt{3^2} = 3 \\ &= 6\sqrt{3} && \text{Multiply.}\end{aligned}$
------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

(c) $\sqrt{10}$ No perfect square (other than 1) divides into 10, so $\sqrt{10}$ cannot be simplified further.

(d) $\sqrt[3]{16}$

Remember to write the index.

$$\begin{aligned}\sqrt[3]{16} &= \sqrt[3]{8 \cdot 2} \\ &= \sqrt[3]{8} \cdot \sqrt[3]{2} \\ &= 2\sqrt[3]{2}\end{aligned}$$

Look for the greatest perfect cube that divides into 16.

Factor. 8 is a perfect cube.

Product rule

$\sqrt[3]{8} = 2$

(e)

Remember the negative sign in each line.

$$\begin{aligned}-\sqrt[4]{162} &= -\sqrt[4]{81 \cdot 2} \\ &= -\sqrt[4]{81} \cdot \sqrt[4]{2} \\ &= -3\sqrt[4]{2}\end{aligned}$$

81 is a perfect fourth power.

Product rule

$\sqrt[4]{81} = 3$

NOW TRY 
NOW TRY ANSWERS

4. (a) $5\sqrt{2}$ (b) $8\sqrt{3}$
 (c) $\sqrt{42}$ cannot be simplified further.
 (d) $3\sqrt[3]{4}$ (e) $-2\sqrt[4]{5}$

! CAUTION Be careful with which factors belong outside the radical symbol and which belong inside. In Example 4(b), the $2 \cdot 3$ is written *outside* because $\sqrt{2^2} = 2$ and $\sqrt{3^2} = 3$. The remaining 3 is left inside the radical.


NOW TRY EXERCISE 5

Simplify. Assume that all variables represent positive real numbers.

- (a) $\sqrt{36x^5}$ (b) $\sqrt{32m^5n^4}$
 (c) $\sqrt[3]{-125k^3p^7}$
 (d) $-\sqrt[4]{162x^7y^8}$

EXAMPLE 5 Simplifying Radicals Involving Variables

Simplify. Assume that all variables represent positive real numbers.

$$\begin{aligned} \text{(a)} \quad \sqrt{16m^3} &= \sqrt{16m^2 \cdot m} && \text{Factor.} \\ &= \sqrt{16m^2} \cdot \sqrt{m} && \text{Product rule} \\ &= 4m\sqrt{m} && \text{Take the square root.} \end{aligned}$$

Absolute value bars are not needed around the m in color because all the variables represent *positive* real numbers.

$$\begin{aligned} \text{(b)} \quad \sqrt{200k^7q^8} &= \sqrt{10^2 \cdot 2 \cdot (k^3)^2 \cdot k \cdot (q^4)^2} && \text{Factor into perfect squares.} \\ &= 10k^3q^4\sqrt{2k} && \text{Take the square root.} \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad \sqrt[3]{-8x^4y^5} &= \sqrt[3]{(-8x^3y^3)(xy^2)} && \text{Choose } -8x^3y^3 \text{ as the perfect cube that} \\ &= \sqrt[3]{-8x^3y^3} \cdot \sqrt[3]{xy^2} && \text{divides into } -8x^4y^5. \\ &= -2xy\sqrt[3]{xy^2} && \text{Product rule} \\ &&& \text{Take the cube root.} \end{aligned}$$

$$\begin{aligned} \text{(d)} \quad -\sqrt[4]{32y^9} &= -\sqrt[4]{(16y^8)(2y)} && 16y^8 \text{ is the greatest fourth power that} \\ &= -\sqrt[4]{16y^8} \cdot \sqrt[4]{2y} && \text{divides into } 32y^9. \\ &= -2y^2\sqrt[4]{2y} && \text{Product rule} \\ &&& \text{Take the fourth root.} \end{aligned}$$

NOW TRY 

NOTE From **Example 5**, we see that if a variable is raised to a power with an exponent divisible by 2, it is a perfect square. If it is raised to a power with an exponent divisible by 3, it is a perfect cube. *In general, if it is raised to a power with an exponent divisible by n , it is a perfect n th power.*


NOW TRY EXERCISE 6

Simplify. Assume that all variables represent positive real numbers.

- (a) $\sqrt[6]{7^2}$ (b) $\sqrt[6]{y^4}$
 (c) $\sqrt[4]{x^{10}}$

NOW TRY ANSWERS

5. (a) $6x^2\sqrt{x}$ (b) $4m^2n^2\sqrt{2m}$
 (c) $-5kp^2\sqrt[3]{p}$ (d) $-3xy^2\sqrt[4]{2x^3}$
 6. (a) $\sqrt[6]{7}$ (b) $\sqrt[6]{y^2}$ (c) $x^2\sqrt{x}$

EXAMPLE 6 Simplifying Radicals Using Lesser Indexes

Simplify. Assume that all variables represent positive real numbers.

(a) $\sqrt[9]{5^6}$ Exponents in the radicand and the index must have GCF 1.
 We write this radical using rational exponents and then write the exponent in lowest terms. We then express the answer as a radical.

$$\sqrt[9]{5^6} = (5^6)^{1/9} = 5^{6/9} = 5^{2/3} = \sqrt[3]{5^2} = \sqrt[3]{25}$$

(b) $\sqrt[4]{p^2} = (p^2)^{1/4} = p^{2/4} = p^{1/2} = \sqrt{p}$ (Recall the assumption that $p > 0$.)

(c) $\sqrt[4]{x^{18}} = (x^{18})^{1/4} = x^{18/4} = x^{9/2} = \sqrt{x^9} = \sqrt{x^8 \cdot x} = \sqrt{x^8} \cdot \sqrt{x} = x^2\sqrt{x}$

NOW TRY 

These examples suggest the following rule.

Meaning of $\sqrt[kn]{a^{km}}$

If m is an integer, n and k are natural numbers, and all indicated roots exist, then the following holds true.

$$\sqrt[kn]{a^{km}} = \sqrt[n]{a^m}$$

OBJECTIVE 4 Simplify products and quotients of radicals with different indexes.

We multiply and divide radicals with different indexes using rational exponents.

EXAMPLE 7 Multiplying Radicals with Different Indexes

Simplify $\sqrt{7} \cdot \sqrt[3]{2}$.

The indexes, 2 and 3, have a least common multiple of 6, so we use rational exponents to write each radical as a **sixth** root.

$$\sqrt{7} = 7^{1/2} = 7^{3/6} = \sqrt[6]{7^3} = \sqrt[6]{343}$$

$$\sqrt[3]{2} = 2^{1/3} = 2^{2/6} = \sqrt[6]{2^2} = \sqrt[6]{4}$$

Now we can multiply.

$$\begin{aligned} \sqrt{7} \cdot \sqrt[3]{2} &= \sqrt[6]{343} \cdot \sqrt[6]{4} && \text{Substitute; } \sqrt{7} = \sqrt[6]{343}, \sqrt[3]{2} = \sqrt[6]{4} \\ &= \sqrt[6]{1372} && \text{Product rule} \end{aligned}$$

NOW TRY 

Results such as the one in **Example 7** can be supported with a calculator, as shown in **FIGURE 6**. Notice that the calculator gives the same *approximation* for the initial product and the final radical that we obtained.




FIGURE 6

CAUTION The computation in **FIGURE 6** is not *proof* that the two expressions are equal. The algebra in **Example 7**, however, is valid proof of their equality.


OBJECTIVE 5 Use the Pythagorean theorem.

The **Pythagorean theorem** provides an equation that relates the lengths of the three sides of a right triangle.

Pythagorean Theorem

If a and b are the lengths of the shorter sides of a right triangle and c is the length of the longest side, then the following holds true.

$$a^2 + b^2 = c^2$$



The two shorter sides are the **legs** of the triangle, and the longest side is the **hypotenuse**. The hypotenuse is the side opposite the right angle.

$$\text{leg}^2 + \text{leg}^2 = \text{hypotenuse}^2$$



NOW TRY ANSWER

7. $\sqrt[6]{1944}$

Later we will see that an equation such as $x^2 = 7$ has two solutions: $\sqrt{7}$ (the principal, or positive, square root of 7) and $-\sqrt{7}$. Similarly, $c^2 = 15$ has two solutions: $\pm\sqrt{15}$. In applications we often choose only the principal (positive) square root.

NOW TRY EXERCISE 8

Find the length of the unknown side in each right triangle.



EXAMPLE 8 Using the Pythagorean Theorem

Find the length of the unknown side in the right triangle in FIGURE 7.

Substitute carefully.

$$a^2 + b^2 = c^2$$

$$4^2 + 6^2 = c^2$$

$$16 + 36 = c^2$$

$$c^2 = 52$$

$$c = \sqrt{52}$$

$$c = \sqrt{4 \cdot 13}$$

$$c = \sqrt{4} \cdot \sqrt{13}$$

$$c = 2\sqrt{13}$$

Pythagorean theorem

Let $a = 4$ and $b = 6$.

Apply the exponents.

Add. Interchange sides.

Choose the principal root.

Factor.

Product rule

Simplify.




FIGURE 7

The length of the hypotenuse is $2\sqrt{13}$.

NOW TRY 

CAUTION In the equation $a^2 + b^2 = c^2$, be sure that the length of the hypotenuse is substituted for c and that the lengths of the legs are substituted for a and b .

OBJECTIVE 6 Use the distance formula.

The *distance formula* enables us to find the distance between two points in the coordinate plane, or the length of the line segment joining those two points.

FIGURE 8 shows the points $(3, -4)$ and $(-5, 3)$. The vertical line through $(-5, 3)$ and the horizontal line through $(3, -4)$ intersect at the point $(-5, -4)$. Thus, the point $(-5, -4)$ becomes the vertex of the right angle in a right triangle.

By the Pythagorean theorem, the sum of the squares of the lengths of the two legs a and b of the right triangle in FIGURE 8 is equal to the square of the length of the hypotenuse, c .

$$a^2 + b^2 = c^2, \quad \text{or} \quad c^2 = a^2 + b^2.$$

The length a is the difference of the y -coordinates of the endpoints. The x -coordinate of both points in FIGURE 8 is -5 , so the side is vertical, and we can find a by finding the difference of the y -coordinates. We subtract -4 from 3 to obtain a positive value for a .

$$a = 3 - (-4) = 7$$

Similarly, we find b by subtracting -5 from 3 .

$$b = 3 - (-5) = 8$$




FIGURE 8

NOW TRY ANSWERS

8. (a) $\sqrt{89}$ (b) $6\sqrt{3}$

Now substitute these values into the equation.

$$c^2 = a^2 + b^2 \quad \text{Pythagorean theorem}$$

$$c^2 = 7^2 + 8^2 \quad \text{Let } a = 7 \text{ and } b = 8.$$

$$c^2 = 49 + 64 \quad \text{Apply the exponents.}$$

$$c^2 = 113 \quad \text{Add.}$$

$$c = \sqrt{113} \quad \text{Choose the principal root.}$$

We choose the principal root because the distance cannot be negative. Therefore, the distance between $(-5, 3)$ and $(3, -4)$ is $\sqrt{113}$.

NOTE It is customary to leave the distance in simplified radical form. Do not use a calculator to find an approximation unless specifically directed to do so.




FIGURE 9

This work can be generalized. **FIGURE 9** shows the two points (x_1, y_1) and (x_2, y_2) . The distance a between (x_1, y_1) and (x_2, y_1) is given by

$$a = |x_2 - x_1|,$$

and the distance b between (x_2, y_2) and (x_2, y_1) is given by

$$b = |y_2 - y_1|.$$

From the Pythagorean theorem, we obtain the following.

$$c^2 = a^2 + b^2$$

$$c^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2$$

For all real numbers a ,
 $|a|^2 = a^2$.

Choosing the principal square root gives the **distance formula**. In this formula, we use d (to denote distance) rather than c .

Distance Formula

The distance d between the points (x_1, y_1) and (x_2, y_2) is given by the following.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

NOW TRY EXERCISE 9

Find the distance between the points $(-4, -3)$ and $(-8, 6)$.

EXAMPLE 9 Using the Distance Formula

Find the distance between the points $(-3, 5)$ and $(6, 4)$.

We arbitrarily choose to let $(x_1, y_1) = (-3, 5)$ and $(x_2, y_2) = (6, 4)$.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad \text{Distance formula}$$

$$d = \sqrt{[6 - (-3)]^2 + (4 - 5)^2} \quad \text{Let } x_2 = 6, y_2 = 4, x_1 = -3, y_1 = 5.$$

$$d = \sqrt{9^2 + (-1)^2} \quad \text{Substitute carefully.}$$

$$d = \sqrt{82} \quad \begin{array}{l} \text{Leave in radical form.} \\ \text{The distance is } \sqrt{82}. \end{array}$$

NOW TRY ANSWER

9. $\sqrt{97}$

NOW TRY EXERCISE 10

Find an equation of the circle with center $(0, 0)$ and radius 6, and graph it.

OBJECTIVE 7 Find an equation of a circle given its center and radius.

A **circle** is the set of all points in a plane that lie a fixed distance from a fixed point. The fixed point is the **center**, and the fixed distance is the **radius**. We use the distance formula to find an equation of a circle.

EXAMPLE 10 Finding an Equation of a Circle and Graphing It

Find an equation of the circle with center $(0, 0)$ and radius 3, and graph it.

If the point (x, y) is on the circle, then the distance from (x, y) to the center $(0, 0)$ is the radius 3.

$$\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = d \quad \text{Distance formula}$$

$$\sqrt{(x - 0)^2 + (y - 0)^2} = 3 \quad \text{Let } x_1 = 0, y_1 = 0, \text{ and } d = 3.$$

$$\sqrt{x^2 + y^2} = 3 \quad \text{Simplify.}$$

$$(\sqrt{x^2 + y^2})^2 = 3^2 \quad \text{Square each side.}$$

$$x^2 + y^2 = 9 \quad (\sqrt{a})^2 = a; 3^2 = 9$$




FIGURE 10

An equation of this circle is $x^2 + y^2 = 9$. The graph is shown in FIGURE 10.

NOW TRY

A circle may not be centered at the origin, as seen in the next example.

NOW TRY EXERCISE 11

Find an equation of the circle with center $(-2, 2)$ and radius 3, and graph it.

EXAMPLE 11 Finding an Equation of a Circle and Graphing It

Find an equation of the circle with center $(4, -3)$ and radius 5, and graph it.

$$\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = d \quad \text{Distance formula}$$

$$\sqrt{(x - 4)^2 + [y - (-3)]^2} = 5 \quad \text{Let } x_1 = 4, y_1 = -3, \text{ and } d = 5.$$

$$(x - 4)^2 + (y + 3)^2 = 25 \quad \text{Square each side.}$$

To graph the circle

$$(x - 4)^2 + (y + 3)^2 = 25,$$

plot the center $(4, -3)$, and then, because the radius is 5, move right, left, up, and down 5 units from the center, plotting the points

$$(9, -3), \quad (-1, -3), \quad (4, 2), \quad \text{and} \quad (4, -8).$$

Draw a smooth curve through these four points. When graphing by hand, it is helpful to sketch one quarter of the circle at a time. See FIGURE 11. (Note that the center is not part of the actual graph, but provides help in drawing a more accurate graph.)





FIGURE 11


NOW TRY

NOW TRY ANSWERS

10. $x^2 + y^2 = 36$



11. $(x + 2)^2 + (y - 2)^2 = 9$



Examples 10 and 11 suggest the form of an equation of a circle with radius r and center (h, k) . If (x, y) is a point on the circle, then the distance from the center (h, k) to the point (x, y) is r . See **FIGURE 12**. By the distance formula,

$$\sqrt{(x - h)^2 + (y - k)^2} = r.$$

Squaring each side gives the **center-radius form** of the equation of a circle.




FIGURE 12

Equation of a Circle (Center-Radius Form)

A circle with center (h, k) and radius $r > 0$ has an equation of the form

$$(x - h)^2 + (y - k)^2 = r^2.$$

As a special case, a circle with center at the origin $(0, 0)$ and radius $r > 0$ has the following equation.

$$x^2 + y^2 = r^2$$

NOW TRY EXERCISE 12

Find an equation of the circle with center $(-5, 4)$ and radius $\sqrt{6}$.

NOW TRY ANSWER

12. $(x + 5)^2 + (y - 4)^2 = 6$

EXAMPLE 12 Using the Center-Radius Form of the Equation of a Circle

Find an equation of the circle with center $(-1, 2)$ and radius $\sqrt{7}$.

$$(x - h)^2 + (y - k)^2 = r^2 \quad \text{Center-radius form}$$

$$[x - (-1)]^2 + (y - 2)^2 = (\sqrt{7})^2 \quad \text{Let } h = -1, k = 2, \text{ and } r = \sqrt{7}.$$

Pay attention to signs here.

$$(x + 1)^2 + (y - 2)^2 = 7 \quad \text{Simplify; } (\sqrt{a})^2 = a$$

NOW TRY

7.3 Exercises

FOR EXTRA HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Choose the correct response.

- Which is the greatest perfect square factor of 128?

A. 12 B. 16 C. 32 D. 64
- Which is the greatest perfect cube factor of $81a^7$?

A. $8a^3$ B. $27a^3$ C. $81a^6$ D. $27a^6$
- Which radical can be simplified?

A. $\sqrt{21}$ B. $\sqrt{48}$ C. $\sqrt[3]{12}$ D. $\sqrt[4]{10}$
- Which radical *cannot* be simplified?

A. $\sqrt[3]{30}$ B. $\sqrt[3]{27a^2b}$ C. $\sqrt{\frac{25}{81}}$ D. $\frac{2}{\sqrt{7}}$
- Which one of the following is *not* equal to $\sqrt{\frac{1}{2}}$? (Do not use calculator approximations.)

A. $\sqrt{0.5}$ B. $\sqrt{\frac{2}{4}}$ C. $\sqrt{\frac{3}{6}}$ D. $\frac{\sqrt{4}}{\sqrt{16}}$

6. Which one of the following is *not* equal to $\sqrt[3]{\frac{2}{5}}$? (Do not use calculator approximations.)

- A. $\sqrt[3]{\frac{6}{15}}$ B. $\frac{\sqrt[3]{50}}{5}$ C. $\frac{\sqrt[3]{10}}{\sqrt[3]{25}}$ D. $\frac{\sqrt[3]{10}}{5}$

7. **Concept Check** A student multiplied incorrectly as follows.

$$\begin{aligned}\sqrt[3]{13} \cdot \sqrt[3]{5} \\ &= \sqrt{13 \cdot 5} && \text{Product rule} \\ &= \sqrt{65} && \text{Multiply.}\end{aligned}$$

WHAT WENT WRONG? Give the correct product.

8. **Concept Check** A student multiplied incorrectly as follows.

$$\sqrt[3]{x} \cdot \sqrt[3]{x} = x$$

WHAT WENT WRONG? Give the correct product.

Multiply. Assume that all variables represent positive real numbers. See Examples 1 and 2.

- | | | | |
|-----------------------------------------|------------------------------------------|--------------------------------------|--------------------------------------|
| 9. $\sqrt{3} \cdot \sqrt{3}$ | 10. $\sqrt{5} \cdot \sqrt{5}$ | 11. $\sqrt{18} \cdot \sqrt{2}$ | 12. $\sqrt{12} \cdot \sqrt{3}$ |
| 13. $\sqrt{5} \cdot \sqrt{6}$ | 14. $\sqrt{10} \cdot \sqrt{3}$ | 15. $\sqrt{14} \cdot \sqrt{x}$ | 16. $\sqrt{23} \cdot \sqrt{t}$ |
| 17. $\sqrt{14} \cdot \sqrt{3pqr}$ | 18. $\sqrt{7} \cdot \sqrt{5xt}$ | 19. $\sqrt[3]{2} \cdot \sqrt[3]{5}$ | 20. $\sqrt[3]{3} \cdot \sqrt[3]{6}$ |
| 21. $\sqrt[3]{7x} \cdot \sqrt[3]{2y}$ | 22. $\sqrt[3]{9x} \cdot \sqrt[3]{4y}$ | 23. $\sqrt[4]{11} \cdot \sqrt[4]{3}$ | 24. $\sqrt[4]{6} \cdot \sqrt[4]{9}$ |
| 25. $\sqrt[4]{2x} \cdot \sqrt[4]{3y^2}$ | 26. $\sqrt[4]{3y^2} \cdot \sqrt[4]{6yz}$ | 27. $\sqrt[3]{7} \cdot \sqrt[4]{3}$ | 28. $\sqrt[5]{8} \cdot \sqrt[6]{12}$ |

Simplify. Assume that all variables represent positive real numbers. See Example 3.

- | | | | |
|---------------------------------|----------------------------------|----------------------------------|-----------------------------------|
| 29. $\sqrt{\frac{64}{121}}$ | 30. $\sqrt{\frac{16}{49}}$ | 31. $\sqrt{\frac{3}{25}}$ | 32. $\sqrt{\frac{13}{49}}$ |
| 33. $\sqrt{\frac{x}{25}}$ | 34. $\sqrt{\frac{k}{100}}$ | 35. $\sqrt{\frac{p^6}{81}}$ | 36. $\sqrt{\frac{w^{10}}{36}}$ |
| 37. $\sqrt[3]{-\frac{27}{64}}$ | 38. $\sqrt[3]{-\frac{216}{125}}$ | 39. $\sqrt[3]{\frac{r^2}{8}}$ | 40. $\sqrt[3]{\frac{t}{125}}$ |
| 41. $-\sqrt[4]{\frac{81}{x^4}}$ | 42. $-\sqrt[4]{\frac{625}{y^4}}$ | 43. $\sqrt[5]{\frac{1}{x^{15}}}$ | 44. $\sqrt[5]{\frac{32}{y^{20}}}$ |

45. **Concept Check** A student simplified $\sqrt{48}$ as follows and did not receive credit for his answer.

$$\begin{aligned}\sqrt{48} \\ &= \sqrt{4 \cdot 12} \\ &= \sqrt{4} \cdot \sqrt{12} \\ &= 2\sqrt{12}\end{aligned}$$

WHAT WENT WRONG? Give the correct simplified form.

46. **Concept Check** A student incorrectly claimed that the following radical is in simplified form.

$$\sqrt[3]{k^4}$$

WHAT WENT WRONG? Give the correct simplified form.

Simplify. See Example 4.

47. $\sqrt{12}$

48. $\sqrt{18}$

49. $\sqrt{288}$

50. $\sqrt{72}$

51. $-\sqrt{32}$

52. $-\sqrt{48}$

53. $-\sqrt{28}$

54. $-\sqrt{24}$

55. $\sqrt{30}$

56. $\sqrt{46}$

57. $\sqrt[3]{128}$

58. $\sqrt[3]{24}$

59. $\sqrt[3]{40}$

60. $\sqrt[3]{375}$

61. $\sqrt[3]{-16}$

62. $\sqrt[3]{-250}$

63. $-\sqrt[4]{512}$

64. $-\sqrt[4]{1250}$

65. $\sqrt[5]{64}$

66. $\sqrt[5]{128}$

67. $-\sqrt[5]{486}$

68. $-\sqrt[5]{2048}$

69. $\sqrt[6]{128}$

70. $\sqrt[6]{1458}$

Simplify. Assume that all variables represent positive real numbers. See Example 5.

71. $\sqrt{72k^2}$

72. $\sqrt{18m^2}$

73. $\sqrt{144x^3y^9}$

74. $\sqrt{169s^5t^{10}}$

75. $\sqrt{121x^6}$

76. $\sqrt{256z^{12}}$

77. $-\sqrt[3]{27t^{12}}$

78. $-\sqrt[3]{64y^{18}}$

79. $-\sqrt{100m^8z^4}$

80. $-\sqrt{25t^6s^{20}}$

81. $-\sqrt[3]{-125a^6b^9c^{12}}$

82. $-\sqrt[3]{-216y^{15}x^6z^3}$

83. $\sqrt[4]{\frac{1}{16}r^8t^{20}}$

84. $\sqrt[4]{\frac{81}{256}t^{12}u^8}$

85. $\sqrt{50x^3}$

86. $\sqrt{300z^3}$

87. $-\sqrt{500r^{11}}$

88. $-\sqrt{200p^{13}}$

89. $\sqrt{13x^7y^8}$

90. $\sqrt{23k^9p^{14}}$

91. $\sqrt[3]{8z^6w^9}$

92. $\sqrt[3]{64a^{15}b^{12}}$

93. $\sqrt[3]{-16z^5t^7}$

94. $\sqrt[3]{-81m^4n^{10}}$

95. $\sqrt[4]{81x^{12}y^{16}}$

96. $\sqrt[4]{81t^8u^{28}}$

97. $-\sqrt[4]{162r^{15}s^9}$

98. $-\sqrt[4]{32k^5m^{11}}$

99. $\sqrt{\frac{y^{11}}{36}}$

100. $\sqrt{\frac{v^{13}}{49}}$

101. $\sqrt[3]{\frac{x^{16}}{27}}$

102. $\sqrt[3]{\frac{y^{17}}{125}}$

Simplify. Assume that $x \geq 0$. See Example 6.

103. $\sqrt[4]{48^2}$

104. $\sqrt[4]{50^2}$

105. $\sqrt[4]{25}$

106. $\sqrt[6]{8}$

107. $\sqrt[10]{x^{25}}$

108. $\sqrt[12]{x^{44}}$

109. $\sqrt[10]{x^{16}}$

110. $\sqrt[12]{x^{38}}$

Simplify. Assume that all variables represent positive real numbers. See Example 7.

111. $\sqrt[3]{4} \cdot \sqrt{3}$

112. $\sqrt[3]{5} \cdot \sqrt{6}$



113. $\sqrt[4]{3} \cdot \sqrt[3]{4}$





114. $\sqrt[3]{2} \cdot \sqrt[5]{3}$

115. $\sqrt{x} \cdot \sqrt[3]{x}$

116. $\sqrt[3]{y} \cdot \sqrt[4]{y}$

Find the length of the unknown side in each right triangle. Simplify answers if possible. See Example 8.





Find the distance between each pair of points. See Example 9.

123. (6, 13) and (1, 1)

124. (8, 13) and (2, 5)

125. (-6, 5) and (3, -4)

126. (-1, 5) and (-7, 7)

127. (-8, 2) and (-4, 1)

128. (-1, 2) and (5, 3)

129. (4.7, 2.3) and (1.7, -1.7)

130. (-2.9, 18.2) and (2.1, 6.2)

131. $(\sqrt{2}, \sqrt{6})$ and $(-2\sqrt{2}, 4\sqrt{6})$

132. $(\sqrt{7}, 9\sqrt{3})$ and $(-\sqrt{7}, 4\sqrt{3})$



133. $(x + y, y)$ and $(x - y, x)$

134. $(c, c - d)$ and $(d, c + d)$

Concept Check Work each problem.



135. Match each equation with the correct graph.

(a) $(x - 3)^2 + (y - 2)^2 = 25$




(b) $(x - 3)^2 + (y + 2)^2 = 25$

(c) $(x + 3)^2 + (y - 2)^2 = 25$



(d) $(x + 3)^2 + (y + 2)^2 = 25$

136. A circle can be drawn on a piece of posterboard by fastening one end of a string with a thumbtack, pulling the string taut with a pencil, and tracing a curve, as shown in the figure. Explain why this method works.



Find the equation of a circle satisfying the given conditions. See Examples 10–12.

137. Center: $(0, 0)$; radius: 12

138. Center: $(0, 0)$; radius: 9

139. Center: $(-4, 3)$; radius: 2

140. Center: $(5, -2)$; radius: 4

141. Center: $(-8, -5)$; radius: $\sqrt{5}$

142. Center: $(-12, 13)$; radius: $\sqrt{7}$

Graph each circle. Identify the center and the radius. See Examples 10–12.

143. $x^2 + y^2 = 9$

144. $x^2 + y^2 = 4$

145. $x^2 + y^2 = 16$

146. $x^2 + y^2 = 25$

147. $(x + 3)^2 + (y - 2)^2 = 9$


148. $(x - 1)^2 + (y + 3)^2 = 16$

149. $(x - 2)^2 + (y - 3)^2 = 4$

150. $(x + 4)^2 + (y + 1)^2 = 25$

Solve each problem.


151. A Panasonic Smart Viera E50 LCD HDTV has a rectangular screen with a 36.5-in. width. Its height is 20.8 in. What is the length of the diagonal of the screen to the nearest tenth of an inch? (Data from measurements of the author's television.)



152. The length of the diagonal of a box is given by

$$D = \sqrt{L^2 + W^2 + H^2},$$

where L , W , and H are, respectively, the length, width, and height of the box. Find the length of the diagonal D of a box that is 4 ft long, 2 ft wide, and 3 ft high. Give the exact value, and then round to the nearest tenth of a foot.



153. In the study of sound, one version of the law of tensions is

$$f_1 = f_2 \sqrt{\frac{F_1}{F_2}}.$$

If $F_1 = 300$, $F_2 = 60$, and $f_2 = 260$, find f_1 to the nearest unit.

154. The illumination I , in foot-candles, produced by a light source is related to the distance d , in feet, from the light source by the equation

$$d = \sqrt{\frac{k}{I}},$$

where k is a constant. If $k = 640$, how far from the light source will the illumination be 2 foot-candles? Give the exact value, and then round to the nearest tenth of a foot.

Tom owns a condominium in a high rise building on the shore of Lake Michigan, and has a beautiful view of the lake from his window. He discovered that he can find the number of miles to the horizon by multiplying 1.224 by the square root of his eye level in feet from the ground.



- 155.** Use Tom's discovery to do the following.
- Write a formula that could be used to calculate the distance d in miles to the horizon from a height h in feet from the ground.
 - Tom lives on the 14th floor, which is 150 ft above the ground. His eyes are 6 ft above his floor. Use the formula from part (a) to calculate the distance, to the nearest tenth of a mile, that Tom can see to the horizon from his condominium window.
- 156.** Tom's neighbor Sheri lives on a floor that is 100 ft above the ground. Assuming that her eyes are 5 ft above the ground, to the nearest tenth of a mile, how far can she see to the horizon?


RELATING CONCEPTS For Individual or Group Work (Exercises 157–162)

Heron's formula for finding the area \mathcal{A} of a triangle with sides a , b , and c is

$$\mathcal{A} = \sqrt{s(s-a)(s-b)(s-c)},$$

where s is the semiperimeter, and $s = \frac{1}{2}(a + b + c)$.

Consider the triangle, and **work Exercises 157–162** in order.



- The lengths of the sides of the entire triangle shown are 7, 7, and 12. Find the semiperimeter s .
- Use Heron's formula to find the area of the entire triangle. Write it as a simplified radical.
- Find the height h of the triangle using the Pythagorean theorem.
- Find the area of each congruent right triangle forming the entire triangle, using the common area formula, $\mathcal{A} = \frac{1}{2}bh$.
- Double the result from **Exercise 160** to determine the area of the entire triangle.
- How do the answers in **Exercises 158** and **161** compare?

7.4

Adding and Subtracting Radical Expressions

OBJECTIVE

- 1 Simplify radical expressions involving addition and subtraction.


NOW TRY EXERCISE 1

Add or subtract to simplify each radical expression.

- (a) $3\sqrt{5} + 7\sqrt{5}$
 (b) $2\sqrt{7} - 3\sqrt{7}$
 (c) $\sqrt{12} + \sqrt{75}$
 (d) $-\sqrt{63t} + 3\sqrt{28t}$, $t \geq 0$
 (e) $6\sqrt{7} - 2\sqrt{3}$

OBJECTIVE 1 Simplify radical expressions involving addition and subtraction.

We do so using the distributive property,

$$ac + bc = (a + b)c.$$

EXAMPLE 1 Adding and Subtracting Radicals

Add or subtract to simplify each radical expression.

$$\begin{aligned} \text{(a)} \quad & 4\sqrt{2} + 3\sqrt{2} \\ &= (4 + 3)\sqrt{2} \\ &= 7\sqrt{2} \end{aligned}$$

This is similar to simplifying $4x + 3x$ as $7x$.

$$\begin{aligned} \text{(b)} \quad & 2\sqrt{3} - 5\sqrt{3} \\ &= (2 - 5)\sqrt{3} \\ &= -3\sqrt{3} \end{aligned}$$

This is similar to simplifying $2x - 5x$ as $-3x$.

$$\begin{aligned} \text{(c)} \quad & 3\sqrt{24} + \sqrt{54} \\ &= 3\sqrt{4 \cdot 6} + \sqrt{9 \cdot 6} \\ &= 3\sqrt{4} \cdot \sqrt{6} + \sqrt{9} \cdot \sqrt{6} \\ &= 3 \cdot 2\sqrt{6} + 3\sqrt{6} \\ &= 6\sqrt{6} + 3\sqrt{6} \\ &= (6 + 3)\sqrt{6} \\ &= 9\sqrt{6} \end{aligned}$$

Simplify each individual radical.

Factor the radicands so that one factor is a perfect square.

Product rule

Find the square roots.

Multiply.

Distributive property

Add.

$$\text{(d)} \quad 2\sqrt{20x} - \sqrt{45x}, \quad x \geq 0$$

$$= 2\sqrt{4} \cdot \sqrt{5x} - \sqrt{9} \cdot \sqrt{5x} \quad \text{Product rule}$$

$$= 2 \cdot 2\sqrt{5x} - 3\sqrt{5x} \quad \text{Find the square roots.}$$

$$= 4\sqrt{5x} - 3\sqrt{5x} \quad \text{Multiply.}$$

$$= (4 - 3)\sqrt{5x} \quad \text{Distributive property}$$

$$= \sqrt{5x} \quad \text{Subtract.}$$

$$1\sqrt{5x} = \sqrt{5x}$$

$$\text{(e)} \quad 2\sqrt{3} - 4\sqrt{5}$$

The radicands differ and are already simplified, so this expression cannot be simplified further.

NOW TRY 

NOW TRY ANSWERS

1. (a) $10\sqrt{5}$ (b) $-\sqrt{7}$
 (c) $7\sqrt{3}$ (d) $3\sqrt{7t}$
 (e) The expression cannot be simplified further.

CAUTION Only radical expressions with the same index and the same radicand may be combined.


NOW TRY EXERCISE 2

Add or subtract to simplify each radical expression. Assume that all variables represent positive real numbers.

(a) $3\sqrt[3]{2000} - 4\sqrt[3]{128}$

(b) $5\sqrt[4]{a^5b^3} + \sqrt[4]{81ab^7}$

(c) $\sqrt[3]{128t^4} - 2\sqrt[3]{72t^3}$

EXAMPLE 2 Adding and Subtracting Radicals with Higher Indexes

Add or subtract to simplify each radical expression. Assume that all variables represent positive real numbers.

(a) $2\sqrt[3]{16} - 5\sqrt[3]{54}$ Remember to write the index with each radical.

$$= 2\sqrt[3]{8 \cdot 2} - 5\sqrt[3]{27 \cdot 2} \quad \text{Factor.}$$

$$= 2\sqrt[3]{8} \cdot \sqrt[3]{2} - 5\sqrt[3]{27} \cdot \sqrt[3]{2} \quad \text{Product rule}$$

$$= 2 \cdot 2 \cdot \sqrt[3]{2} - 5 \cdot 3 \cdot \sqrt[3]{2} \quad \text{Find the cube roots.}$$

$$= 4\sqrt[3]{2} - 15\sqrt[3]{2} \quad \text{Multiply.}$$

$$= (4 - 15)\sqrt[3]{2} \quad \text{Distributive property}$$

$$= -11\sqrt[3]{2} \quad \text{Combine like terms.}$$

In practice, the step indicating $(4 - 15)\sqrt[3]{2}$ can be done mentally, giving the final answer $-11\sqrt[3]{2}$ directly.

(b) $2\sqrt[3]{x^2y} + \sqrt[3]{8x^5y^4}$

$$= 2\sqrt[3]{x^2y} + \sqrt[3]{8x^3y^3 \cdot x^2y} \quad \text{Factor.}$$

$$= 2\sqrt[3]{x^2y} + \sqrt[3]{8x^3y^3} \cdot \sqrt[3]{x^2y} \quad \text{Product rule}$$

$$= 2\sqrt[3]{x^2y} + 2xy\sqrt[3]{x^2y} \quad \text{Find the cube root.}$$

This result cannot be simplified further.

$$= (2 + 2xy)\sqrt[3]{x^2y} \quad \text{Distributive property}$$

Although we were able to use the distributive property in the last step, 2 and $2xy$ are not like terms and cannot be combined into a single term.

(c) $5\sqrt{4x^3} + 3\sqrt[3]{64x^4}$ Be careful. The indexes are different.

$$= 5\sqrt{4x^2 \cdot x} + 3\sqrt[3]{64x^3 \cdot x} \quad \text{Factor.}$$

$$= 5\sqrt{4x^2} \cdot \sqrt{x} + 3\sqrt[3]{64x^3} \cdot \sqrt[3]{x} \quad \text{Product rule}$$

$$= 5 \cdot 2x\sqrt{x} + 3 \cdot 4x\sqrt[3]{x} \quad \text{Keep track of the indexes.}$$

$$= 10x\sqrt{x} + 12x\sqrt[3]{x} \quad \text{Multiply.}$$

The two terms in the final expression cannot be combined into a single term—the indexes are different.

NOW TRY 

NOW TRY ANSWERS

2. (a) $14\sqrt[3]{2}$

(b) $(5a + 3b)\sqrt[4]{ab^3}$

(c) $4t\sqrt[3]{2t} - 12t\sqrt{2t}$

CAUTION *The root of a sum does not equal the sum of the roots.*

Example: $\sqrt{9 + 16} \neq \sqrt{9} + \sqrt{16}$

because $\sqrt{9 + 16} = \sqrt{25} = 5$, but $\sqrt{9} + \sqrt{16} = 3 + 4 = 7$.


NOW TRY EXERCISE 3

Perform the indicated operations. Assume that all variables represent positive real numbers.

(a) $5\sqrt{\frac{\sqrt{5}}{45}} - 4\sqrt{\frac{28}{9}}$

(b) $6\sqrt[3]{\frac{16}{x^{12}}} + 7\sqrt[3]{\frac{9}{x^9}}$

EXAMPLE 3 Adding and Subtracting Radicals with Fractions

Perform the indicated operations. Assume that all variables represent positive real numbers.

$$\begin{aligned} \text{(a)} \quad & 2\sqrt{\frac{75}{16}} + 4\sqrt{\frac{\sqrt{8}}{32}} \\ &= 2\frac{\sqrt{25 \cdot 3}}{\sqrt{16}} + 4\frac{\sqrt{4 \cdot 2}}{\sqrt{16 \cdot 2}} \\ &= 2\left(\frac{5\sqrt{3}}{4}\right) + 4\left(\frac{2\sqrt{2}}{4\sqrt{2}}\right) \\ &= \frac{5\sqrt{3}}{2} + 2 \\ &= \frac{5\sqrt{3}}{2} + \frac{4}{2} \\ &= \frac{5\sqrt{3} + 4}{2} \end{aligned}$$

Quotient rule; Factor.

Product rule; Find the square roots.

Multiply; $\frac{\sqrt{2}}{\sqrt{2}} = 1$

Write with a common denominator; $2 = \frac{4}{2}$

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

$$\begin{aligned} \text{(b)} \quad & 10\sqrt[3]{\frac{5}{x^6}} - 3\sqrt[3]{\frac{4}{x^9}} \\ &= 10\frac{\sqrt[3]{5}}{\sqrt[3]{x^6}} - 3\frac{\sqrt[3]{4}}{\sqrt[3]{x^9}} \\ &= \frac{10\sqrt[3]{5}}{x^2} - \frac{3\sqrt[3]{4}}{x^3} \\ &= \frac{10\sqrt[3]{5} \cdot x}{x^2 \cdot x} - \frac{3\sqrt[3]{4}}{x^3} \\ &= \frac{10x\sqrt[3]{5} - 3\sqrt[3]{4}}{x^3} \end{aligned}$$

Quotient rule

Simplify denominators.

Write with a common denominator.

$$\frac{a}{c} - \frac{b}{c} = \frac{a-b}{c}$$

NOW TRY ANSWERS

3. (a) $\frac{5 - 8\sqrt{7}}{3}$
 (b) $\frac{12\sqrt[3]{2} + 7x\sqrt[3]{9}}{x^4}$

This equals x^3 , so there is a common denominator.


NOW TRY 

7.4 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Choose the correct response.

- Which sum can be simplified without first simplifying the individual radical expressions?
 - $\sqrt{50} + \sqrt{32}$
 - $3\sqrt{6} + 9\sqrt{6}$
 - $\sqrt[3]{32} - \sqrt[3]{108}$
 - $\sqrt[5]{6} - \sqrt[5]{192}$
- Which difference can be simplified without first simplifying the individual radical expressions?
 - $\sqrt{81} - \sqrt{18}$
 - $\sqrt[3]{8} - \sqrt[3]{16}$
 - $4\sqrt[3]{7} - 9\sqrt[3]{7}$
 - $\sqrt{75} - \sqrt{12}$

3. Concept Check A student incorrectly simplified

$$(3 + 3xy)\sqrt[3]{xy^2} \text{ as } 6xy\sqrt[3]{xy^2}.$$

His teacher did not give him any credit for this answer. **WHAT WENT WRONG?**

4. Concept Check A student incorrectly gave the difference

$$28 - 4\sqrt{2} \text{ as } 24\sqrt{2}.$$

Her teacher did not give her any credit for this answer. **WHAT WENT WRONG?**

Add or subtract to simplify each radical expression. Assume that all variables represent positive real numbers. See Examples 1 and 2.

5. $\sqrt{36} - \sqrt{100}$

6. $\sqrt{25} - \sqrt{81}$

7. $6\sqrt{10} + 2\sqrt{10}$

8. $5\sqrt{6} + 4\sqrt{6}$

9. $6\sqrt{5} - 7\sqrt{5}$

10. $3\sqrt{2} - 4\sqrt{2}$

11. $-2\sqrt{48} + 3\sqrt{75}$

12. $4\sqrt{32} - 2\sqrt{8}$

13. $5\sqrt{6} + 2\sqrt{10}$

14. $3\sqrt{11} - 5\sqrt{13}$

15. $\sqrt[3]{16} + 4\sqrt[3]{54}$

16. $3\sqrt[3]{24} - 2\sqrt[3]{192}$

17. $6\sqrt{18} - \sqrt{32} + 2\sqrt{50}$

18. $5\sqrt{8} + 3\sqrt{72} - 3\sqrt{50}$

19. $2\sqrt{5} + 3\sqrt{20} + 4\sqrt{45}$

20. $5\sqrt{54} - 2\sqrt{24} - 2\sqrt{96}$

21. $\sqrt{72x} - \sqrt{8x}$

22. $\sqrt{18k} - \sqrt{72k}$

23. $3\sqrt{72m^2} - 5\sqrt{32m^2} - 3\sqrt{18m^2}$

24. $9\sqrt{27p^2} - 14\sqrt{108p^2} + 2\sqrt{48p^2}$

25. $\sqrt[4]{32} + 3\sqrt[4]{2}$

26. $\sqrt[4]{405} - 2\sqrt[4]{5}$

27. $2\sqrt[3]{16} + \sqrt[3]{54}$

28. $15\sqrt[3]{81} + 4\sqrt[3]{24}$

29. $2\sqrt[3]{27x} - 2\sqrt[3]{8x}$

30. $6\sqrt[3]{128m} - 3\sqrt[3]{16m}$

31. $3\sqrt[3]{x^2y} - 5\sqrt[3]{8x^2y}$

32. $3\sqrt[3]{x^2y^2} - 2\sqrt[3]{64x^2y^2}$

33. $3x\sqrt[3]{xy^2} - 2\sqrt[3]{8x^4y^2}$

34. $6q^2\sqrt[3]{5q} - 2q\sqrt[3]{40q^4}$

35. $5\sqrt[4]{32} + 3\sqrt[4]{162}$

36. $2\sqrt[4]{512} + 4\sqrt[4]{32}$

37. $3\sqrt[4]{x^5y} - 2x\sqrt[4]{xy}$

38. $2\sqrt[4]{m^9p^6} - 3m^2p\sqrt[4]{mp^2}$

39. $2\sqrt[4]{32a^3} + 5\sqrt[4]{2a^3}$

40. $5\sqrt[4]{243x^3} + 2\sqrt[4]{3x^3}$

41. $\sqrt[3]{64xy^2} + \sqrt[3]{27x^4y^5}$

42. $\sqrt[4]{625s^3t} + \sqrt[4]{81s^7t^5}$

43. $\sqrt[3]{192st^4} - \sqrt[3]{27s^3t}$

44. $\sqrt{125a^5b^5} + \sqrt[3]{125a^4b^4}$

45. $2\sqrt[3]{8x^4} + 3\sqrt[4]{16x^5}$

46. $3\sqrt[3]{64m^4} + 5\sqrt[4]{81m^5}$

Perform the indicated operations. Assume that all variables represent positive real numbers. See Example 3.

47. $\sqrt{8} - \frac{\sqrt{64}}{\sqrt{16}}$

48. $\sqrt{48} - \frac{\sqrt{81}}{\sqrt{9}}$

49. $\frac{2\sqrt{5}}{3} + \frac{\sqrt{5}}{6}$

50. $\frac{4\sqrt{3}}{3} + \frac{2\sqrt{3}}{9}$

51. $\sqrt{\frac{8}{9}} + \sqrt{\frac{18}{36}}$

52. $\sqrt{\frac{12}{16}} + \sqrt{\frac{48}{64}}$

53. $\frac{\sqrt{32}}{3} + \frac{2\sqrt{2}}{3} - \frac{\sqrt{2}}{\sqrt{9}}$

54. $\frac{\sqrt{27}}{2} - \frac{3\sqrt{3}}{2} + \frac{\sqrt{3}}{\sqrt{4}}$

55. $3\sqrt{\frac{50}{9}} + 8\frac{\sqrt{2}}{\sqrt{8}}$

56. $5\sqrt{\frac{288}{25}} + 21\frac{\sqrt{2}}{\sqrt{18}}$

57. $\sqrt{\frac{25}{x^8}} + \sqrt{\frac{9}{x^6}}$

58. $\sqrt{\frac{100}{y^4}} + \sqrt{\frac{81}{y^{10}}}$

59. $3\sqrt{\frac{50}{49}} - \frac{\sqrt{27}}{\sqrt{12}}$

60. $9\sqrt{\frac{48}{25}} - 2\frac{\sqrt{2}}{\sqrt{98}}$

61. $3\sqrt[3]{\frac{m^5}{27}} - 2m\sqrt[3]{\frac{m^2}{64}}$

62. $2a\sqrt[4]{\frac{a}{16}} - 5a\sqrt[4]{\frac{a}{81}}$

63. $3\sqrt[3]{\frac{2}{x^6}} - 4\sqrt[3]{\frac{5}{x^9}}$

64. $-4\sqrt[3]{\frac{4}{t^9}} + 3\sqrt[3]{\frac{9}{t^{12}}}$

65. **Concept Check** Consider the expression

$$\sqrt{63} + \sqrt{112} - \sqrt{252}.$$

- (a) Simplify this expression using the methods of this section.
 (b) Use a calculator to approximate the given expression.
 (c) Use a calculator to approximate the simplified expression in part (a).
 (d) Complete the following: Assuming the work in part (a) is correct, the approximations in parts (b) and (c) should be (*equal/unequal*).

66. **Concept Check** Let $a = 1$ and let $b = 64$.

- (a) Evaluate $\sqrt{a} + \sqrt{b}$. Then find $\sqrt{a+b}$. Are they equal?
 (b) Evaluate $\sqrt[3]{a} + \sqrt[3]{b}$. Then find $\sqrt[3]{a+b}$. Are they equal?
 (c) Complete the following: In general,

$$\sqrt[n]{a} + \sqrt[n]{b} \neq \underline{\hspace{2cm}},$$

based on the observations in parts (a) and (b) of this exercise.

Solve each problem.




67. A rectangular yard has a length of $\sqrt{192}$ m and a width of $\sqrt{48}$ m. Choose the best estimate of its dimensions. Then estimate the perimeter.

- A. 14 m by 7 m B. 5 m by 7 m C. 14 m by 8 m D. 15 m by 8 m


68. If the sides of a triangle are $\sqrt{65}$ in., $\sqrt{35}$ in., and $\sqrt{26}$ in., which one of the following is the best estimate of its perimeter?

- A. 19 in. B. 20 in. C. 24 in. D. 26 in.



Find the perimeter of each figure. Give answers as simplified radical expressions.



72. Find the area of the trapezoid. Give the answer as a simplified radical.



Extending Skills Find the perimeter of each triangle.



7.5 Multiplying and Dividing Radical Expressions

OBJECTIVES

- 1 Multiply radical expressions.
- 2 Rationalize denominators with one radical term.
- 3 Rationalize denominators with binomials involving radicals.
- 4 Write radical quotients in lowest terms.

VOCABULARY

□ conjugates

NOW TRY EXERCISE 1

Multiply.

- (a) $\sqrt{10}(4 + \sqrt{7})$
- (b) $3(\sqrt{20} - \sqrt{45})$

NOW TRY ANSWERS

1. (a) $4\sqrt{10} + \sqrt{70}$
- (b) $-3\sqrt{5}$

OBJECTIVE 1 Multiply radical expressions.

The distributive property may be used when multiplying radical expressions.

EXAMPLE 1 Using the Distributive Property with Radicals

Multiply.

$$\begin{aligned} \text{(a)} \quad & \sqrt{5}(2 + \sqrt{6}) \\ &= \sqrt{5} \cdot 2 + \sqrt{5} \cdot \sqrt{6} && \text{Distributive property: } a(b + c) = ab + ac \\ &= 2\sqrt{5} + \sqrt{30} && \text{Commutative property; product rule} \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad & 4(\sqrt{12} - \sqrt{27}) \\ &= 4\sqrt{12} - 4\sqrt{27} && \text{Distributive property} \\ &= 4\sqrt{4 \cdot 3} - 4\sqrt{9 \cdot 3} && \text{Factor the radicands so that} \\ & && \text{one factor is a perfect square.} \\ &= 4 \cdot 2\sqrt{3} - 4 \cdot 3\sqrt{3} && \sqrt{4} = 2; \sqrt{9} = 3 \\ &= 8\sqrt{3} - 12\sqrt{3} && \text{Multiply.} \\ &= -4\sqrt{3} && 8\sqrt{3} - 12\sqrt{3} = (8 - 12)\sqrt{3} \end{aligned}$$

NOW TRY

We multiply binomial expressions involving radicals using the FOIL method. Recall that the acronym **FOIL** refers to the positions of the terms. We multiply the **F**irst terms, **O**uter terms, **I**nner terms, and **L**ast terms of the binomials.

EXAMPLE 2 Multiplying Binomials Involving Radical Expressions

Multiply, using the FOIL method.

$$\begin{aligned} \text{(a)} \quad & (\sqrt{5} + 3)(\sqrt{6} + 1) \\ & \begin{array}{cccc} \text{First} & \text{Outer} & \text{Inner} & \text{Last} \\ \hline & & & \\ & & & \\ & & & \end{array} \\ &= \sqrt{5} \cdot \sqrt{6} + \sqrt{5} \cdot 1 + 3 \cdot \sqrt{6} + 3 \cdot 1 && \text{FOIL method} \\ &= \sqrt{30} + \sqrt{5} + 3\sqrt{6} + 3 && \text{This result cannot be simplified further.} \end{aligned}$$


NOW TRY EXERCISE 2

Multiply, using the FOIL method.

(a) $(8 - \sqrt{5})(9 - \sqrt{2})$

(b) $(\sqrt{7} + \sqrt{5})(\sqrt{7} - \sqrt{5})$

(c) $(\sqrt{15} - 4)^2$

(d) $(8 + \sqrt[3]{5})(8 - \sqrt[3]{5})$

(e) $(\sqrt{m} - \sqrt{n})(\sqrt{m} + \sqrt{n})$,
 $m \geq 0$ and $n \geq 0$

(b) $(7 - \sqrt{3})(\sqrt{5} + \sqrt{2})$

$$\begin{array}{ccccccc} & \text{F} & \text{O} & \text{I} & \text{L} & & \\ = & 7\sqrt{5} & + & 7\sqrt{2} & - & \sqrt{3} \cdot \sqrt{5} & - & \sqrt{3} \cdot \sqrt{2} & \text{FOIL method} \end{array}$$

$$= 7\sqrt{5} + 7\sqrt{2} - \sqrt{15} - \sqrt{6} \quad \text{Product rule}$$

(c) $(\sqrt{10} + \sqrt{3})(\sqrt{10} - \sqrt{3})$

$$= \sqrt{10} \cdot \sqrt{10} - \sqrt{10} \cdot \sqrt{3} + \sqrt{3} \cdot \sqrt{10} - \sqrt{3} \cdot \sqrt{3} \quad \text{FOIL method}$$

$$= 10 - 3 \quad \text{Product rule; } -\sqrt{30} + \sqrt{30} = 0$$

$$= 7 \quad \text{Subtract.}$$

A product such as $(\sqrt{10} + \sqrt{3})(\sqrt{10} - \sqrt{3}) = (\sqrt{10})^2 - (\sqrt{3})^2$ is a difference of squares.

$$(x + y)(x - y) = x^2 - y^2 \quad \text{Here, } x = \sqrt{10} \text{ and } y = \sqrt{3}.$$

(d) $(\sqrt{7} - 3)^2$

$$= (\sqrt{7} - 3)(\sqrt{7} - 3) \quad a^2 = a \cdot a$$

$$= \sqrt{7} \cdot \sqrt{7} - 3\sqrt{7} - 3\sqrt{7} + 3 \cdot 3 \quad \text{FOIL method}$$

$$= 7 - 6\sqrt{7} + 9 \quad \text{Multiply. Combine like terms.}$$

$$= 16 - 6\sqrt{7} \quad \text{Add.}$$

Be careful. These terms cannot be combined.

(e) $(5 - \sqrt[3]{3})(5 + \sqrt[3]{3})$

$$= 5 \cdot 5 + 5\sqrt[3]{3} - 5\sqrt[3]{3} - \sqrt[3]{3} \cdot \sqrt[3]{3} \quad \text{FOIL method}$$

$$= 25 - \sqrt[3]{3^2} \quad \text{Multiply. Combine like terms.}$$

$$= 25 - \sqrt[3]{9} \quad \text{Apply the exponent.}$$

Remember to write the index 3 in each radical.

(f) $(\sqrt{k} + \sqrt{y})(\sqrt{k} - \sqrt{y})$, $k \geq 0$ and $y \geq 0$

$$= (\sqrt{k})^2 - (\sqrt{y})^2 \quad \text{Difference of squares}$$

$$= k - y$$

NOW TRY 

NOTE In **Example 2(d)**, we could have used the formula for the square of a binomial to obtain the same result.

$$(\sqrt{7} - 3)^2$$

$$= (\sqrt{7})^2 - 2(\sqrt{7})(3) + 3^2 \quad (x - y)^2 = x^2 - 2xy + y^2$$

$$= 7 - 6\sqrt{7} + 9 \quad \text{Apply the exponents. Multiply.}$$

$$= 16 - 6\sqrt{7} \quad \text{Add.}$$

NOW TRY ANSWERS

2. (a) $72 - 8\sqrt{2} - 9\sqrt{5} + \sqrt{10}$

(b) 2 (c) $31 - 8\sqrt{15}$

(d) $64 - \sqrt[3]{25}$ (e) $m - n$

OBJECTIVE 2 Rationalize denominators with one radical term.

A simplified radical expression has no radical in the denominator. The origin of this agreement no doubt occurred before the days of high-speed calculation, when computation was a tedious process performed by hand.

Consider the expression $\frac{1}{\sqrt{2}}$. To find a decimal approximation by hand, it is necessary to divide 1 by a decimal approximation for $\sqrt{2}$, such as 1.414. It is much easier if the divisor is a whole number. This can be accomplished by multiplying $\frac{1}{\sqrt{2}}$ by 1 in the form $\frac{\sqrt{2}}{\sqrt{2}}$. **Multiplying by 1 in any form does not change the value of the original expression.**

$$\frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2} \quad \text{Multiply by 1; } \frac{\sqrt{2}}{\sqrt{2}} = 1$$

Now the computation requires dividing 1.414 by 2 to obtain 0.707, which is easier.

With current technology, either form $\frac{1}{\sqrt{2}}$ or $\frac{\sqrt{2}}{2}$ can be approximated with the same number of keystrokes. See **FIGURE 13**, which shows that a calculator gives the same approximation for both forms of the expression.




FIGURE 13

Rationalizing the Denominator

The process of removing radicals from a denominator so that the denominator contains only rational numbers is called **rationalizing the denominator**. This is done by multiplying by a form of 1.

EXAMPLE 3 Rationalizing Denominators with Square Roots

Rationalize each denominator.

(a) $\frac{3}{\sqrt{7}}$

Multiply by $\frac{\sqrt{7}}{\sqrt{7}}$. This is an application of the identity property of multiplication—in effect, we are multiplying by 1.

$$\frac{3}{\sqrt{7}} = \frac{3 \cdot \sqrt{7}}{\sqrt{7} \cdot \sqrt{7}} = \frac{3\sqrt{7}}{7} \quad \begin{array}{l} \text{In the denominator,} \\ \sqrt{7} \cdot \sqrt{7} = \sqrt{7 \cdot 7} = \sqrt{49} = 7. \\ \text{The final denominator is now a rational number.} \end{array}$$

(b) $\frac{5\sqrt{2}}{\sqrt{5}} = \frac{5\sqrt{2} \cdot \sqrt{5}}{\sqrt{5} \cdot \sqrt{5}} = \frac{5\sqrt{10}}{5} = \sqrt{10}$ Divide out the common factor, 5.


NOW TRY EXERCISE 3

Rationalize each denominator.

(a) $\frac{8}{\sqrt{13}}$ (b) $\frac{9\sqrt{7}}{\sqrt{3}}$

(c) $\frac{-10}{\sqrt{20}}$

(c) $\frac{-6}{\sqrt{12}}$

Less work is involved if the radical in the denominator is simplified first.

$$\frac{-6}{\sqrt{12}} = \frac{-6}{\sqrt{4 \cdot 3}} = \frac{-6}{2\sqrt{3}} = \frac{-3}{\sqrt{3}}$$

Now we rationalize the denominator.

$$\frac{-3}{\sqrt{3}} = \frac{-3 \cdot \sqrt{3}}{\sqrt{3} \cdot \sqrt{3}} = \frac{-3\sqrt{3}}{3} = -\sqrt{3}$$

NOW TRY 

NOW TRY EXERCISE 4

Simplify.

(a) $-\sqrt{\frac{27}{80}}$

(b) $\sqrt{\frac{48x^8}{y^3}}, y > 0$

EXAMPLE 4 Rationalizing Denominators in Roots of Fractions

Simplify.

(a) $-\sqrt{\frac{18}{125}}$

$$= -\frac{\sqrt{18}}{\sqrt{125}}$$

Quotient rule

$$= -\frac{\sqrt{9 \cdot 2}}{\sqrt{25 \cdot 5}}$$

Factor.

$$= -\frac{3\sqrt{2}}{5\sqrt{5}}$$

Product rule

$$= -\frac{3\sqrt{2} \cdot \sqrt{5}}{5\sqrt{5} \cdot \sqrt{5}}$$

Multiply by $\frac{\sqrt{5}}{\sqrt{5}}$.

$$= -\frac{3\sqrt{10}}{5 \cdot 5}$$

Product rule

$$= -\frac{3\sqrt{10}}{25}$$

Multiply.

(b) $\sqrt{\frac{50m^4}{p^5}}, p > 0$

$$= \frac{\sqrt{50m^4}}{\sqrt{p^5}}$$

Quotient rule

$$= \frac{\sqrt{25m^4 \cdot 2}}{\sqrt{p^4 \cdot p}}$$

Factor.

$$= \frac{5m^2\sqrt{2}}{p^2\sqrt{p}}$$

Product rule

$$= \frac{5m^2\sqrt{2} \cdot \sqrt{p}}{p^2\sqrt{p} \cdot \sqrt{p}}$$

Multiply by $\frac{\sqrt{p}}{\sqrt{p}}$.

$$= \frac{5m^2\sqrt{2p}}{p^2 \cdot p}$$

Product rule

$$= \frac{5m^2\sqrt{2p}}{p^3}$$

Multiply.

NOW TRY 
NOW TRY ANSWERS

3. (a) $\frac{8\sqrt{13}}{13}$ (b) $3\sqrt{21}$

(c) $-\sqrt{5}$

4. (a) $-\frac{3\sqrt{15}}{20}$ (b) $\frac{4x^4\sqrt{3y}}{y^2}$

EXAMPLE 5 Rationalizing Denominators with Higher Roots

Simplify.

(a) $\sqrt[3]{\frac{27}{16}}$

Use the quotient rule, and simplify the numerator and denominator.

$$\sqrt[3]{\frac{27}{16}} = \frac{\sqrt[3]{27}}{\sqrt[3]{16}} = \frac{3}{\sqrt[3]{8 \cdot 2}} = \frac{3}{2\sqrt[3]{2}}$$

NOW TRY EXERCISE 5

Simplify.

(a) $\sqrt[3]{\frac{8}{81}}$

(b) $\sqrt[4]{\frac{7x}{y}}$, $x \geq 0, y > 0$

Because $2 \cdot 4 = 8$ is a perfect cube, multiply the numerator and denominator of the simplified expression by $\sqrt[3]{4}$.

$$\frac{3}{2\sqrt[3]{2}}$$

$\sqrt[3]{\frac{27}{16}} = \frac{3}{2\sqrt[3]{2}}$ from above

$$= \frac{3 \cdot \sqrt[3]{4}}{2\sqrt[3]{2} \cdot \sqrt[3]{4}}$$

Multiply by $\sqrt[3]{4}$ in the numerator and denominator. This will give $\sqrt[3]{8} = 2$ in the denominator.

$$= \frac{3\sqrt[3]{4}}{2\sqrt[3]{8}}$$

Multiply.

$$= \frac{3\sqrt[3]{4}}{2 \cdot 2}$$

$\sqrt[3]{8} = 2$

$$= \frac{3\sqrt[3]{4}}{4}$$

Multiply.

(b) $\sqrt[4]{\frac{5x}{z}}$, $x \geq 0, z > 0$

$$= \frac{\sqrt[4]{5x}}{\sqrt[4]{z}}$$

Quotient rule

$$= \frac{\sqrt[4]{5x} \cdot \sqrt[4]{z^3}}{\sqrt[4]{z} \cdot \sqrt[4]{z^3}}$$

Multiply by $\sqrt[4]{z^3}$ in the numerator and denominator.

$\sqrt[4]{z} \cdot \sqrt[4]{z^3}$
will give $\sqrt[4]{z^4}$.

$$= \frac{\sqrt[4]{5xz^3}}{\sqrt[4]{z^4}}$$

Product rule

$$= \frac{\sqrt[4]{5xz^3}}{z}$$

NOW TRY

CAUTION In Example 5(a), a typical error is to multiply the numerator and denominator by $\sqrt[3]{2}$, forgetting that

$$\sqrt[3]{2} \cdot \sqrt[3]{2} = \sqrt[3]{2^2}, \text{ which does not equal } 2.$$

We need **three** factors of 2 to obtain 2^3 under the radical.

$$\sqrt[3]{2} \cdot \sqrt[3]{2} \cdot \sqrt[3]{2} = \sqrt[3]{2^3}, \text{ which does equal } 2.$$

OBJECTIVE 3 Rationalize denominators with binomials involving radicals.

To rationalize a denominator that contains a binomial expression (one that contains exactly two terms) involving radicals, such as

$$\frac{3}{1 + \sqrt{2}},$$

we use the special product $(x + y)(x - y) = x^2 - y^2$ and the concept of *conjugates*. The conjugate of $1 + \sqrt{2}$ is $1 - \sqrt{2}$.

NOW TRY ANSWERS

5. (a) $\frac{2\sqrt[3]{9}}{9}$ (b) $\frac{\sqrt[4]{7xy^3}}{y}$

In general, $x + y$ and $x - y$ are **conjugates**. Specifically, if x and y represent non-negative rational numbers, the product

$$(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y}) \text{ produces the rational number } x - y.$$

Rationalizing a Binomial Denominator

Whenever a radical expression has a sum or difference with square root radicals in the denominator, rationalize the denominator by multiplying both the numerator and denominator by the conjugate of the denominator.

EXAMPLE 6 Rationalizing Binomial Denominators

Rationalize each denominator.

(a)
$$\frac{3}{1 + \sqrt{2}}$$

Again, we are multiplying by a form of 1.

$$= \frac{3(1 - \sqrt{2})}{(1 + \sqrt{2})(1 - \sqrt{2})}$$

Multiply the numerator and denominator by $1 - \sqrt{2}$, the conjugate of the denominator.

$$(1 + \sqrt{2})(1 - \sqrt{2}) = 1^2 - (\sqrt{2})^2 = 1 - 2 = -1$$

The denominator is now a rational number.

$$= \frac{3(1 - \sqrt{2})}{-1}$$

$$= \frac{3}{-1}(1 - \sqrt{2})$$

$$\frac{a \cdot b}{c} = \frac{a}{c} \cdot b$$

$$= -3(1 - \sqrt{2})$$

$$\frac{a}{-1} = -a$$

$$= -3 + 3\sqrt{2}$$

Distributive property

Either of the forms in the last two lines can be given as the answer.

(b)
$$\frac{5}{4 - \sqrt{3}}$$

$$= \frac{5(4 + \sqrt{3})}{(4 - \sqrt{3})(4 + \sqrt{3})}$$

Multiply the numerator and denominator by $4 + \sqrt{3}$.

$$= \frac{5(4 + \sqrt{3})}{16 - 3}$$

Multiply in the denominator.

$$= \frac{5(4 + \sqrt{3})}{13}$$

Subtract in the denominator.

We leave the numerator in factored form. This makes it easier to determine whether the expression is written in lowest terms.


NOW TRY EXERCISE 6

Rationalize each denominator.

(a) $\frac{4}{1 + \sqrt{3}}$ (b) $\frac{4}{5 - \sqrt{7}}$

(c) $\frac{\sqrt{3} + \sqrt{7}}{\sqrt{5} - \sqrt{2}}$

(d) $\frac{8}{\sqrt{3x} + \sqrt{y}}$,
 $3x \neq y, x > 0, y > 0$

(c) $\frac{\sqrt{2} - \sqrt{3}}{\sqrt{5} + \sqrt{3}}$

$$= \frac{(\sqrt{2} - \sqrt{3})(\sqrt{5} - \sqrt{3})}{(\sqrt{5} + \sqrt{3})(\sqrt{5} - \sqrt{3})}$$

Multiply the numerator and denominator by $\sqrt{5} - \sqrt{3}$.

$$= \frac{\sqrt{10} - \sqrt{6} - \sqrt{15} + 3}{5 - 3}$$

Multiply.

$$= \frac{\sqrt{10} - \sqrt{6} - \sqrt{15} + 3}{2}$$

Subtract in the denominator.

(d) $\frac{3}{\sqrt{5m} - \sqrt{p}}$, $5m \neq p, m > 0, p > 0$

$$= \frac{3(\sqrt{5m} + \sqrt{p})}{(\sqrt{5m} - \sqrt{p})(\sqrt{5m} + \sqrt{p})}$$

Multiply the numerator and denominator by $\sqrt{5m} + \sqrt{p}$.

$$= \frac{3(\sqrt{5m} + \sqrt{p})}{5m - p}$$

Multiply in the denominator.

NOW TRY **OBJECTIVE 4** Write radical quotients in lowest terms.

NOW TRY EXERCISE 7

Write each quotient in lowest terms.

(a) $\frac{15 - 6\sqrt{2}}{18}$

(b) $\frac{15k + \sqrt{50k^2}}{20k}$, $k > 0$

(a) $\frac{6 + 2\sqrt{5}}{4}$

$$= \frac{2(3 + \sqrt{5})}{2 \cdot 2}$$

This is a key step.

Factor the numerator and denominator.

$$= \frac{3 + \sqrt{5}}{2}$$

Divide out the common factor.

Alternative method: $\frac{6 + 2\sqrt{5}}{4} = \frac{6}{4} + \frac{2\sqrt{5}}{4} = \frac{3}{2} + \frac{\sqrt{5}}{2} = \frac{3 + \sqrt{5}}{2}$

(b) $\frac{5y - \sqrt{8y^2}}{6y}$, $y > 0$

$$= \frac{5y - 2y\sqrt{2}}{6y}$$

$$\sqrt{8y^2} = \sqrt{4y^2 \cdot 2} = 2y\sqrt{2}$$

$$= \frac{y(5 - 2\sqrt{2})}{6y}$$

Factor the numerator.

$$= \frac{5 - 2\sqrt{2}}{6}$$

Divide out the common factor.

NOW TRY **NOW TRY ANSWERS**

6. (a) $-2 + 2\sqrt{3}$

(b) $\frac{2(5 + \sqrt{7})}{9}$

(c) $\frac{\sqrt{15} + \sqrt{6} + \sqrt{35} + \sqrt{14}}{3}$

(d) $\frac{8(\sqrt{3x} - \sqrt{y})}{3x - y}$

7. (a) $\frac{5 - 2\sqrt{2}}{6}$ (b) $\frac{3 + \sqrt{2}}{4}$

CAUTION Be careful to factor before writing a quotient in lowest terms.

7.5 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Match each part of a rule for a special product in Column I with the part it equals in Column II. Assume that A and B represent positive real numbers.

I

1. $(x + \sqrt{y})(x - \sqrt{y})$

2. $(\sqrt{x} + y)(\sqrt{x} - y)$

3. $(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y})$

4. $(\sqrt{x} + \sqrt{y})^2$

5. $(\sqrt{x} - \sqrt{y})^2$

6. $(\sqrt{x} + y)^2$

II

A. $x - y$

B. $x + 2y\sqrt{x} + y^2$

C. $x - y^2$

D. $x - 2\sqrt{xy} + y$

E. $x^2 - y$

F. $x + 2\sqrt{xy} + y$

Multiply, and then simplify each product. Assume that all variables represent positive real numbers. See Examples 1 and 2.

7. $\sqrt{6}(3 + \sqrt{2})$

8. $\sqrt{10}(5 + \sqrt{3})$

9. $\sqrt{3}(\sqrt{12} - 4)$

10. $\sqrt{5}(\sqrt{125} - 6)$

11. $5(\sqrt{72} - \sqrt{8})$

12. $7(\sqrt{50} - \sqrt{18})$

13. $\sqrt{2}(\sqrt{18} - \sqrt{3})$

14. $\sqrt{5}(\sqrt{15} + \sqrt{5})$

15. $(\sqrt{2} + 1)(\sqrt{3} + 1)$

16. $(\sqrt{3} + 3)(\sqrt{5} + 2)$

17. $(\sqrt{2} - \sqrt{3})(\sqrt{2} + \sqrt{3})$

18. $(\sqrt{7} + \sqrt{14})(\sqrt{7} - \sqrt{14})$

19. $(\sqrt{8} - \sqrt{2})(\sqrt{8} + \sqrt{2})$

20. $(\sqrt{20} - \sqrt{5})(\sqrt{20} + \sqrt{5})$

21. $(\sqrt{11} - \sqrt{7})(\sqrt{2} + \sqrt{5})$

22. $(\sqrt{13} - \sqrt{7})(\sqrt{3} + \sqrt{11})$

23. $(2\sqrt{3} + \sqrt{5})(3\sqrt{3} - 2\sqrt{5})$

24. $(\sqrt{7} - \sqrt{11})(2\sqrt{7} + 3\sqrt{11})$

25. $(\sqrt{5} + 2)^2$

26. $(\sqrt{11} - 1)^2$

27. $(\sqrt{21} - \sqrt{5})^2$

28. $(\sqrt{6} - \sqrt{2})^2$

29. $(2 + \sqrt[3]{6})(2 - \sqrt[3]{6})$

30. $(\sqrt[3]{3} + 6)(\sqrt[3]{3} - 6)$

31. $(2 + \sqrt[3]{2})(4 - 2\sqrt[3]{2} + \sqrt[3]{4})$

32. $(\sqrt[3]{3} - 1)(\sqrt[3]{9} + \sqrt[3]{3} + 1)$

33. $(3\sqrt{x} - \sqrt{5})(2\sqrt{x} + 1)$

34. $(4\sqrt{p} + \sqrt{7})(\sqrt{p} - 9)$

35. $(3\sqrt{r} - \sqrt{s})(3\sqrt{r} + \sqrt{s})$

36. $(\sqrt{k} + 4\sqrt{m})(\sqrt{k} - 4\sqrt{m})$

37. $(\sqrt[3]{2y} - 5)(4\sqrt[3]{2y} + 1)$

38. $(\sqrt[3]{9z} - 2)(5\sqrt[3]{9z} + 7)$

39. $(\sqrt{3x} + 2)(\sqrt{3x} - 2)$

40. $(\sqrt{6y} - 4)(\sqrt{6y} + 4)$

41. Concept Check A student incorrectly simplified the radical expression

$$6 - 4\sqrt{3} \text{ as } 2\sqrt{3}.$$

WHAT WENT WRONG?

42. Concept Check A student rationalized the following denominator as shown.

$$\frac{5}{\sqrt[3]{2}} = \frac{5 \cdot \sqrt[3]{2}}{\sqrt[3]{2} \cdot \sqrt[3]{2}} = \frac{5\sqrt[3]{2}}{2} \quad \text{Incorrect}$$

WHAT WENT WRONG? Give the correct answer.

Rationalize each denominator. Assume that all variables represent positive real numbers. See Examples 3 and 4.

43. $\frac{7}{\sqrt{7}}$

44. $\frac{11}{\sqrt{11}}$

45. $\frac{15}{\sqrt{3}}$

46. $\frac{12}{\sqrt{6}}$

47. $\frac{\sqrt{3}}{\sqrt{2}}$

48. $\frac{\sqrt{7}}{\sqrt{6}}$

49. $\frac{9\sqrt{3}}{\sqrt{5}}$

50. $\frac{3\sqrt{2}}{\sqrt{11}}$

51. $\frac{-7}{\sqrt{48}}$

52. $\frac{-5}{\sqrt{24}}$

53. $\sqrt{\frac{7}{2}}$

54. $\sqrt{\frac{10}{3}}$

55. $-\sqrt{\frac{7}{50}}$

56. $-\sqrt{\frac{13}{75}}$

57. $\sqrt{\frac{24}{x}}$

58. $\sqrt{\frac{52}{y}}$

59. $\frac{-8\sqrt{3}}{\sqrt{k}}$

60. $\frac{-4\sqrt{13}}{\sqrt{m}}$

61. $-\sqrt{\frac{150m^5}{n^3}}$

62. $-\sqrt{\frac{98r^3}{s^5}}$

63. $\sqrt{\frac{288x^7}{y^9}}$

64. $\sqrt{\frac{242t^9}{u^{11}}}$

65. $\frac{5\sqrt{2m}}{\sqrt{y^3}}$

66. $\frac{2\sqrt{5r}}{\sqrt{m^3}}$

67. $-\sqrt{\frac{48k^2}{z}}$

68. $-\sqrt{\frac{75m^3}{p}}$

Simplify. Assume that all variables represent positive real numbers. See Example 5.

69. $\sqrt[3]{\frac{2}{3}}$

70. $\sqrt[3]{\frac{4}{5}}$

71. $\sqrt[3]{\frac{4}{9}}$

72. $\sqrt[3]{\frac{5}{16}}$

73. $\sqrt[3]{\frac{9}{32}}$

74. $\sqrt[3]{\frac{10}{9}}$

75. $-\sqrt[3]{\frac{2p}{r^2}}$

76. $-\sqrt[3]{\frac{6x}{y^2}}$

77. $\sqrt[3]{\frac{x^6}{y}}$

78. $\sqrt[3]{\frac{m^9}{q}}$

79. $\sqrt[4]{\frac{16}{x}}$

80. $\sqrt[4]{\frac{81}{y}}$

81. $\sqrt[4]{\frac{2y}{z}}$

82. $\sqrt[4]{\frac{7t}{s^2}}$

Rationalize each denominator. Assume that all variables represent positive real numbers and that no denominators are 0. See Example 6.

83. $\frac{3}{4 + \sqrt{5}}$

84. $\frac{4}{5 + \sqrt{6}}$

85. $\frac{6}{\sqrt{5} + \sqrt{3}}$

86. $\frac{12}{\sqrt{6} + \sqrt{3}}$

87. $\frac{\sqrt{8}}{3 - \sqrt{2}}$

88. $\frac{\sqrt{27}}{3 - \sqrt{3}}$

89. $\frac{2}{3\sqrt{5} + 2\sqrt{3}}$

90. $\frac{-1}{3\sqrt{2} - 2\sqrt{7}}$

91. $\frac{\sqrt{2} - \sqrt{3}}{\sqrt{6} - \sqrt{5}}$

92. $\frac{\sqrt{5} + \sqrt{6}}{\sqrt{3} - \sqrt{2}}$

93. $\frac{m - 4}{\sqrt{m} + 2}$

94. $\frac{r - 9}{\sqrt{r} - 3}$

95. $\frac{4}{\sqrt{x} - 2\sqrt{y}}$

96. $\frac{5}{3\sqrt{r} + \sqrt{s}}$

97. $\frac{\sqrt{x} - \sqrt{y}}{\sqrt{x} + \sqrt{y}}$

98. $\frac{\sqrt{a} + \sqrt{b}}{\sqrt{a} - \sqrt{b}}$

99. $\frac{5\sqrt{k}}{2\sqrt{k} + \sqrt{q}}$

100. $\frac{3\sqrt{x}}{\sqrt{x} - 2\sqrt{y}}$

Write each quotient in lowest terms. Assume that all variables represent positive real numbers. See Example 7.

101. $\frac{30 + 20\sqrt{6}}{10}$

102. $\frac{24 + 12\sqrt{5}}{12}$

103. $\frac{3 - 3\sqrt{5}}{3}$

104. $\frac{-5 + 5\sqrt{2}}{5}$

105. $\frac{16 - 4\sqrt{8}}{12}$

106. $\frac{12 - 9\sqrt{72}}{18}$

107. $\frac{6p + \sqrt{24p^3}}{3p}$

108. $\frac{11y - \sqrt{242y^5}}{22y}$

Extending Skills Rationalize each denominator. Assume that all radicals represent real numbers and that no denominators are 0.

109. $\frac{3}{\sqrt{x+y}}$

110. $\frac{5}{\sqrt{m-n}}$

111. $\frac{p}{\sqrt{p+2}}$

112. $\frac{q}{\sqrt{5+q}}$

Solve each problem.

113. The following expression occurs in a standard problem in trigonometry.

$$\frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \cdot \frac{1}{2}$$

Show that it simplifies to $\frac{\sqrt{6} - \sqrt{2}}{4}$. Then verify, using a calculator approximation.

114. The following expression occurs in a standard problem in trigonometry.

$$\frac{\sqrt{3} + 1}{1 - \sqrt{3}}$$

Show that it simplifies to $-2 - \sqrt{3}$. Then verify, using a calculator approximation.

RELATING CONCEPTS For Individual or Group Work (Exercises 115–118)

In calculus, it is sometimes desirable to rationalize the numerator. To rationalize a numerator, we multiply the numerator and the denominator by the conjugate of the numerator. For example,

$$\frac{6 - \sqrt{2}}{4} = \frac{(6 - \sqrt{2})(6 + \sqrt{2})}{4(6 + \sqrt{2})} = \frac{36 - 2}{4(6 + \sqrt{2})} = \frac{34}{4(6 + \sqrt{2})} = \frac{17}{2(6 + \sqrt{2})}$$

Rationalize each numerator. Assume that all variables represent positive real numbers.

115. $\frac{6 - \sqrt{3}}{8}$

116. $\frac{2\sqrt{5} - 3}{2}$

117. $\frac{2\sqrt{x} - \sqrt{y}}{3x}$

118. $\frac{\sqrt{p} - 3\sqrt{q}}{4q}$

SUMMARY EXERCISES Performing Operations with Radicals and Rational Exponents

Conditions for a Simplified Radical

1. The radicand has no factor raised to a power greater than or equal to the index.
2. The radicand has no fractions.
3. No denominator contains a radical.
4. Exponents in the radicand and the index of the radical have greatest common factor 1.

Concept Check Give the reason why each radical is not simplified.

1. $\sqrt{\frac{2}{5}}$

2. $\sqrt[15]{x^5}$

3. $\frac{5}{\sqrt[3]{10}}$

4. $\sqrt[3]{x^5y^6}$

Perform the indicated operations, and express each answer in simplest form. Assume that all variables represent positive real numbers.

5. $6\sqrt{10} - 12\sqrt{10}$

6. $\sqrt{7}(\sqrt{7} - \sqrt{2})$

7. $(1 - \sqrt{3})(2 + \sqrt{6})$

8. $\sqrt{50} - \sqrt{98} + \sqrt{72}$

9. $(3\sqrt{5} + 2\sqrt{7})^2$

10. $\frac{-3}{\sqrt{6}}$

11. $\frac{8}{\sqrt{7} + \sqrt{5}}$

12. $\frac{1 - \sqrt{2}}{1 + \sqrt{2}}$

13. $(\sqrt{5} + 7)(\sqrt{5} - 7)$

14. $\frac{1}{\sqrt{x} - \sqrt{5}}, x \neq 5$

15. $\sqrt[3]{8a^3b^5c^9}$

16. $\frac{15}{\sqrt[3]{9}}$

17. $\frac{3}{\sqrt{5} + 2}$

18. $\sqrt{\frac{3}{5x}}$

19. $\frac{16\sqrt{3}}{5\sqrt{12}}$

20. $\frac{2\sqrt{25}}{8\sqrt{50}}$

21. $\frac{-10}{\sqrt[3]{10}}$

22. $\frac{\sqrt{6} + \sqrt{5}}{\sqrt{6} - \sqrt{5}}$

23. $\sqrt{12x} - \sqrt{75x}$

24. $(5 - 3\sqrt{3})^2$

25. $\sqrt[3]{\frac{13}{81}}$

26. $\frac{\sqrt{3} + \sqrt{7}}{\sqrt{6} - \sqrt{5}}$

27. $\frac{6}{\sqrt[4]{3}}$

28. $\sqrt[3]{\frac{x^2y}{x^{-3}y^4}}$

29. $\sqrt{12} - \sqrt{108} - \sqrt[3]{27}$

30. $\frac{4^{1/2} + 3^{1/2}}{4^{1/2} - 3^{1/2}}$

31. $\sqrt[3]{16x^2} - \sqrt[3]{54x^2} + \sqrt[3]{128x^2}$

32. $(1 - \sqrt[3]{3})(1 + \sqrt[3]{3} + \sqrt[3]{9})$

Simplify each expression. Write answers with positive exponents. Assume that all variables represent positive real numbers.

33. $3^{1/2} \cdot 3^{1/3}$

34. $\left(\frac{x^2y}{x^{-3}y^4}\right)^{1/3}$

35. $\frac{x^{-2/3}y^{4/5}}{x^{-5/3}y^{-2/5}}$

36. $\left(\frac{x^{3/4}y^{2/3}}{x^{1/3}y^{5/8}}\right)^{24}$

37. $(125x^3)^{-2/3}$

38. $(3x^{-2/3}y^{1/2})(-2x^{5/8}y^{-1/3})$

7.6 Solving Equations with Radicals

OBJECTIVES

- 1 Solve radical equations using the power rule.
- 2 Solve radical equations that require additional steps.
- 3 Solve radical equations with indexes greater than 2.
- 4 Use the power rule to solve a formula for a specified variable.

VOCABULARY

- radical equation
- proposed solution
- extraneous solution

OBJECTIVE 1 Solve radical equations using the power rule.

An equation that includes one or more radical expressions with a variable in a radicand is a **radical equation**.

$$\sqrt{x-4} = 8, \quad \sqrt{5x+12} = 3\sqrt{2x-1}, \quad \sqrt[3]{6+x} = 27 \quad \text{Radical equations}$$

Solving radical equations involves a process that we have not yet seen, and it requires careful application. Notice that the equation $x = 1$ has only one solution. Its solution set is $\{1\}$. If we square both sides of this equation, we obtain $x^2 = 1$. This new equation has *two* solutions: -1 and 1 . The solution of the original equation is also a solution of the “squared” equation. However, that equation has another solution, -1 , that is *not* a solution of the original equation.

When solving equations with radicals, we use this idea of raising both sides to a power, which is an application of the **power rule**.

Power Rule for Solving a Radical Equation

If both sides of an equation are raised to the same power, all solutions of the original equation are also solutions of the new equation.

The power rule does not say that all solutions of the new equation are solutions of the original equation. They may or may not be. A value of the variable that appears to be a solution is a **proposed solution**. Such values that do not satisfy the original equation are **extraneous solutions**. They must be rejected.

CAUTION When the power rule is used to solve an equation, every solution of the new equation must be checked in the original equation.

NOW TRY EXERCISE 1

Solve $\sqrt{9x+7} = 5$.

EXAMPLE 1 Using the Power Rule

Solve $\sqrt{3x+4} = 8$.

$$(\sqrt{a})^2 = \sqrt{a} \cdot \sqrt{a} = a$$

for $a \geq 0$.

$$(\sqrt{3x+4})^2 = 8^2$$

Use the power rule and square each side.

$$3x + 4 = 64$$

Apply the exponents.

$$3x = 60$$

Subtract 4.

$$x = 20$$

Divide by 3.

CHECK

$$\sqrt{3x+4} = 8$$

Original equation

$$\sqrt{3 \cdot 20 + 4} \stackrel{?}{=} 8$$

Let $x = 20$.

$$\sqrt{64} \stackrel{?}{=} 8$$

Simplify.

$$8 = 8 \quad \checkmark \quad \text{True}$$

Because 20 satisfies the *original* equation, the solution set is $\{20\}$.

NOW TRY ANSWER

1. $\{2\}$

Solving an Equation with Radicals

- Step 1 Isolate the radical.** Make sure that one radical term is alone on one side of the equation.
- Step 2 Apply the power rule.** Raise each side of the equation to a power that is the same as the index of the radical.
- Step 3 Solve** the resulting equation. If it still contains a radical, repeat Steps 1 and 2.
- Step 4 Check** all proposed solutions in the *original* equation. Discard any values that are not solutions of the original equation.

CAUTION Remember to check all proposed solutions (Step 4) or an incorrect solution set may result.

NOW TRY EXERCISE 2

Solve $\sqrt{3x + 4} + 5 = 0$.

EXAMPLE 2 Using the Power Rule

Solve $\sqrt{5x - 1} + 3 = 0$.

- | | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|----------------------------------------------------------------|
| Step 1 | $\sqrt{5x - 1} = -3$ | To isolate the radical on one side, subtract 3 from each side. |
| Step 2 | $(\sqrt{5x - 1})^2 = (-3)^2$ | Square each side. |
| Step 3 | $5x - 1 = 9$ | Apply the exponents. |
| | $5x = 10$ | Add 1. |
| | $x = 2$ | Divide by 5. |
| Step 4 CHECK | $\sqrt{5x - 1} + 3 = 0$ | Original equation |
| <div style="border: 1px solid #ccc; border-radius: 5px; padding: 2px; width: fit-content; margin-left: 20px;">Be sure to check the proposed solution.</div> | $\sqrt{5 \cdot 2 - 1} + 3 \stackrel{?}{=} 0$ | Let $x = 2$. |
| | $\sqrt{9} + 3 \stackrel{?}{=} 0$ | Evaluate the radicand. |
| | $3 + 3 \stackrel{?}{=} 0$ | Take the square root. |
| | $6 = 0$ | False |

This false result shows that the *proposed* solution 2 is *not* a solution of the original equation. It is extraneous. The solution set is \emptyset .

NOW TRY

NOTE We could have determined after Step 1 that the equation in **Example 2** has no solution because the expression on the left cannot equal a negative number.

$$\sqrt{5x - 1} = -3 \quad \text{A square root radical cannot be negative.}$$

OBJECTIVE 2 Solve radical equations that require additional steps.

Recall the following rule for squaring a binomial.

$$(x + y)^2 = x^2 + 2xy + y^2$$

NOW TRY ANSWER

2. \emptyset


**NOW TRY
EXERCISE 3**
Solve $\sqrt{16 - x} = x + 4$.
EXAMPLE 3 Using the Power Rule (Squaring a Binomial)
Solve $\sqrt{4 - x} = x + 2$.**Step 1** The radical is isolated on the left side of the equation.**Step 2** Square each side. On the right, $(x + 2)^2 = x^2 + 2(x)(2) + 2^2$.

$$(\sqrt{4 - x})^2 = (x + 2)^2$$

Remember the middle term.

$$4 - x = x^2 + 4x + 4$$

 Twice the product of 2 and x
Step 3 The new equation is quadratic, so write it in standard form.

$$x^2 + 5x = 0$$

Subtract 4, add x , and interchange sides.

$$x(x + 5) = 0$$

Factor.

Set each factor equal to 0.

$$x = 0 \quad \text{or} \quad x + 5 = 0$$

Zero-factor property

$$x = -5$$

Solve.

Step 4 Check each proposed solution in the original equation.

CHECK $\sqrt{4 - x} = x + 2$

$$\sqrt{4 - 0} \stackrel{?}{=} 0 + 2 \quad \text{Let } x = 0.$$

$$\sqrt{4} \stackrel{?}{=} 2$$

$$2 = 2 \quad \checkmark \quad \text{True}$$

$\sqrt{4 - x} = x + 2$

$$\sqrt{4 - (-5)} \stackrel{?}{=} -5 + 2 \quad \text{Let } x = -5.$$

$$\sqrt{9} \stackrel{?}{=} -3$$

$$3 = -3 \quad \text{False}$$

The solution set is $\{0\}$. The proposed solution -5 is extraneous.**NOW TRY** 

**NOW TRY
EXERCISE 4**

Solve

$\sqrt{x^2 - 3x + 18} = x + 3$.

EXAMPLE 4 Using the Power Rule (Squaring a Binomial)
Solve $\sqrt{x^2 - 4x + 9} = x - 1$.Square each side. On the right, $(x - 1)^2 = x^2 - 2(x)(1) + 1^2$.

$$(\sqrt{x^2 - 4x + 9})^2 = (x - 1)^2$$

Remember the middle term.

$$x^2 - 4x + 9 = x^2 - 2x + 1$$

 Twice the product of x and -1

$$-2x = -8$$

Subtract x^2 and 9. Add $2x$.

$$x = 4$$

Divide by -2 .

CHECK $\sqrt{x^2 - 4x + 9} = x - 1$ Original equation

$$\sqrt{4^2 - 4 \cdot 4 + 9} \stackrel{?}{=} 4 - 1 \quad \text{Let } x = 4.$$

$$\sqrt{9} \stackrel{?}{=} 4 - 1 \quad \text{Evaluate the radicand.}$$

$$3 = 3 \quad \checkmark \quad \text{True}$$

The solution set is $\{4\}$.**NOW TRY** **NOW TRY ANSWERS**3. $\{0\}$ 4. $\{1\}$


**NOW TRY
EXERCISE 5**

Solve

$$\sqrt{3x+1} - \sqrt{x+4} = 1.$$

EXAMPLE 5 Using the Power Rule (Squaring Twice)
Solve $\sqrt{5x+6} + \sqrt{3x+4} = 2$.Isolate one radical on one side by subtracting $\sqrt{3x+4}$ from each side.

$$\sqrt{5x+6} = 2 - \sqrt{3x+4} \quad \text{Subtract } \sqrt{3x+4}.$$

$$(\sqrt{5x+6})^2 = (2 - \sqrt{3x+4})^2 \quad \text{Square each side.}$$

$$5x + 6 = 4 - 4\sqrt{3x+4} + (3x+4) \quad \text{Be careful here.}$$

Remember the middle term.

Twice the product of 2 and $-\sqrt{3x+4}$

The equation still contains a radical, so isolate the radical term on the right and square both sides again.

$$5x + 6 = 8 - 4\sqrt{3x+4} + 3x \quad \text{Combine like terms.}$$

$$2x - 2 = -4\sqrt{3x+4} \quad \text{Subtract 8 and 3x.}$$

Divide each term by 2.

$$x - 1 = -2\sqrt{3x+4} \quad \text{Divide each term by 2 to obtain smaller numbers.}$$

$$(x - 1)^2 = (-2\sqrt{3x+4})^2 \quad \text{Square each side again.}$$

$$x^2 - 2x + 1 = (-2)^2(\sqrt{3x+4})^2 \quad \text{On the right, } (ab)^2 = a^2b^2.$$

$$x^2 - 2x + 1 = 4(3x+4) \quad \text{Apply the exponents.}$$

$$x^2 - 2x + 1 = 12x + 16 \quad \text{Distributive property}$$

$$x^2 - 14x - 15 = 0 \quad \text{Standard form}$$

$$(x - 15)(x + 1) = 0 \quad \text{Factor.}$$

$$x - 15 = 0 \quad \text{or} \quad x + 1 = 0 \quad \text{Zero-factor property}$$

$$x = 15 \quad \text{or} \quad x = -1 \quad \text{Solve each equation.}$$

CHECK First check the proposed solution 15. Then check -1 .

$$\sqrt{5x+6} + \sqrt{3x+4} = 2 \quad \text{Original equation}$$

$$\sqrt{5(15)+6} + \sqrt{3(15)+4} \stackrel{?}{=} 2 \quad \text{Let } x = 15.$$

$$\sqrt{81} + \sqrt{49} \stackrel{?}{=} 2 \quad \text{Evaluate the radicands.}$$

$$9 + 7 \stackrel{?}{=} 2 \quad \text{Take square roots.}$$

$$16 = 2 \quad \text{False}$$


$$\sqrt{5x+6} + \sqrt{3x+4} = 2 \quad \text{Original equation}$$

$$\sqrt{5(-1)+6} + \sqrt{3(-1)+4} \stackrel{?}{=} 2 \quad \text{Let } x = -1.$$

$$\sqrt{1} + \sqrt{1} \stackrel{?}{=} 2 \quad \text{Evaluate the radicands.}$$

$$1 + 1 \stackrel{?}{=} 2 \quad \text{Take square roots.}$$

$$2 = 2 \quad \checkmark \quad \text{True}$$

NOW TRY ANSWER
5. $\{5\}$ The proposed solution 15 is extraneous and must be rejected, but -1 is valid. Thus, the solution set is $\{-1\}$.**NOW TRY** 


**NOW TRY
EXERCISE 6**

Solve $\sqrt[3]{4x-5} = \sqrt[3]{3x+2}$.

OBJECTIVE 3 Solve radical equations with indexes greater than 2.**EXAMPLE 6** Using the Power Rule for a Power Greater Than 2

Solve $\sqrt[3]{z+5} = \sqrt[3]{2z-6}$.

$$\left(\sqrt[3]{z+5}\right)^3 = \left(\sqrt[3]{2z-6}\right)^3 \quad \text{Cube each side.}$$

$$z+5 = 2z-6 \quad \left(\sqrt[3]{a}\right)^3 = a$$

$$11 = z \quad \text{Subtract } z. \text{ Add } 6.$$

CHECK

$$\sqrt[3]{z+5} = \sqrt[3]{2z-6} \quad \text{Original equation}$$

$$\sqrt[3]{11+5} \stackrel{?}{=} \sqrt[3]{2 \cdot 11 - 6} \quad \text{Let } z = 11.$$

$$\sqrt[3]{16} = \sqrt[3]{16} \quad \checkmark \quad \text{True}$$

The solution set is $\{11\}$.**NOW TRY** **OBJECTIVE 4** Use the power rule to solve a formula for a specified variable.**EXAMPLE 7** Solving a Formula from Electronics for a Variable

An important property of a radio-frequency transmission line is its **characteristic impedance**, represented by Z and measured in ohms. If L and C are the inductance and capacitance, respectively, per unit of length of the line, then these quantities are related by the following formula. Solve this formula for C .

$$Z = \sqrt{\frac{L}{C}}$$

Our goal is to isolate C on one side of the equality symbol.

$$Z^2 = \left(\sqrt{\frac{L}{C}}\right)^2 \quad \text{Square each side.}$$

$$Z^2 = \frac{L}{C} \quad \left(\sqrt{a}\right)^2 = a$$

$$CZ^2 = L \quad \text{Multiply by } C.$$

$$C = \frac{L}{Z^2} \quad \text{Divide by } Z^2.$$

NOW TRY 

**NOW TRY
EXERCISE 7**
Solve the formula for a .


$$x = \sqrt{\frac{y+2}{a}}$$

NOW TRY ANSWERS

6. $\{7\}$

7. $a = \frac{y+2}{x^2}$

7.6 Exercises**FOR
EXTRA
HELP****MyLab Math**

 Video solutions for select problems available in MyLab Math

Concept Check Check each equation to see if the given value for x is a solution.

1. $\sqrt{3x+18} - x = 0$
(a) 6 (b) -3

2. $\sqrt{3x-3} - x + 1 = 0$
(a) 1 (b) 4

3. $\sqrt{x+2} - \sqrt{9x-2} = -2\sqrt{x-1}$
(a) 2 (b) 7

4. $\sqrt{8x-3} = 2x$
(a) $\frac{3}{2}$ (b) $\frac{1}{2}$

5. **Concept Check** A student solved the following equation and claimed that 9 is a solution of the equation.

$$\sqrt{x} = -3$$

He received no credit for his answer.

WHAT WENT WRONG? Give the correct solution set.

6. **Concept Check** A student solved the following equation and obtained the proposed solutions $x = -3$ and $x = 6$.

$$\sqrt{3x + 18} = x$$

She gave $\{-3, 6\}$ as the solution set.

WHAT WENT WRONG? Give the correct solution set.

Solve each equation. See Examples 1–4.

7. $\sqrt{x - 2} = 3$

9. $\sqrt{6k - 1} = 1$

11. $\sqrt{4r + 3} + 1 = 0$

13. $\sqrt{3x + 1} - 4 = 0$

15. $4 - \sqrt{x - 2} = 0$

17. $\sqrt{9x - 4} = \sqrt{8x + 1}$

19. $2\sqrt{x} = \sqrt{3x + 4}$

21. $3\sqrt{x - 1} = 2\sqrt{2x + 2}$

23. $x = \sqrt{x^2 + 4x - 20}$

25. $x = \sqrt{x^2 + 3x + 9}$

27. $\sqrt{9 - x} = x + 3$

29. $\sqrt{5 - x} = x + 1$

31. $\sqrt{6x + 7} = x + 2$

33. $\sqrt{k^2 + 2k + 9} = k + 3$

35. $\sqrt{z^2 + 12z - 4} + 4 - z = 0$

37. $\sqrt{r^2 + 9r + 15} - r - 4 = 0$

39. **Concept Check** When solving the equation $\sqrt{3x + 4} = 8 - x$, a student wrote the following as her first step.

$$3x + 4 = 64 + x^2$$

WHAT WENT WRONG? Solve the given equation correctly.

Solve each equation. See Example 5.

41. $\sqrt{k + 2} - \sqrt{k - 3} = 1$

43. $\sqrt{2r + 11} - \sqrt{5r + 1} = -1$

45. $\sqrt{3p + 4} - \sqrt{2p - 4} = 2$

47. $\sqrt{3 - 3p} - 3 = \sqrt{3p + 2}$

49. $\sqrt{2\sqrt{x + 11}} = \sqrt{4x + 2}$

8. $\sqrt{x + 1} = 7$

10. $\sqrt{7x - 3} = 6$

12. $\sqrt{5k - 3} + 2 = 0$

14. $\sqrt{5x + 1} - 11 = 0$

16. $9 - \sqrt{4x + 1} = 0$

18. $\sqrt{4x - 2} = \sqrt{3x + 5}$

20. $2\sqrt{x} = \sqrt{5x - 16}$

22. $5\sqrt{4x + 1} = 3\sqrt{10x + 25}$

24. $x = \sqrt{x^2 - 3x + 18}$

26. $x = \sqrt{x^2 - 4x - 8}$

28. $\sqrt{36 - x} = x + 6$

30. $\sqrt{3 - x} = x + 3$

32. $\sqrt{4x + 13} = x + 4$

34. $\sqrt{x^2 - 3x + 3} = x - 1$

36. $\sqrt{x^2 - 15x + 15} + 5 - x = 0$

38. $\sqrt{m^2 + 3m + 12} - m - 2 = 0$

40. **Concept Check** When solving the equation $\sqrt{5x + 6} - \sqrt{x + 3} = 3$, a student wrote the following as his first step.

$$(5x + 6) + (x + 3) = 9$$

WHAT WENT WRONG? Solve the given equation correctly.

42. $\sqrt{r + 6} - \sqrt{r - 2} = 2$

44. $\sqrt{3x - 2} - \sqrt{x + 3} = 1$

46. $\sqrt{4x + 5} - \sqrt{2x + 2} = 1$

48. $\sqrt{4x + 7} - 4 = \sqrt{4x - 1}$

50. $\sqrt{1 + \sqrt{24 - 10x}} = \sqrt{3x + 5}$

Solve each equation. See Example 6.

51. $\sqrt[3]{p-1} = 2$

52. $\sqrt[3]{x+8} = 3$

53. $\sqrt[3]{2x+5} = \sqrt[3]{6x+1}$

54. $\sqrt[3]{p+5} = \sqrt[3]{2p-4}$

55. $\sqrt[3]{2m-1} = \sqrt[3]{m+13}$

56. $\sqrt[3]{2k-11} = \sqrt[3]{5k+1}$

57. $\sqrt[3]{x^2+5x+1} = \sqrt[3]{x^2+4x}$

58. $\sqrt[3]{r^2+2r+8} = \sqrt[3]{r^2+3r+12}$

59. $\sqrt[4]{x+12} = \sqrt[4]{3x-4}$

60. $\sqrt[4]{z+11} = \sqrt[4]{2z+6}$

61. $\sqrt[3]{x-8} + 2 = 0$

62. $\sqrt[3]{r+1} + 1 = 0$

63. $\sqrt[4]{2k-5} + 4 = 0$

64. $\sqrt[4]{8z-3} + 2 = 0$

65. $\sqrt[3]{r^2+2r+8} = \sqrt[3]{r^2+3r+12}$

66. $\sqrt[3]{x^2+7x+2} = \sqrt[3]{x^2+6x+1}$

Extending Skills For each equation, write the expressions with rational exponents as radical expressions, and then solve, using the procedures explained in this section.

67. $(2x-9)^{1/2} = 2 + (x-8)^{1/2}$

68. $(3w+7)^{1/2} = 1 + (w+2)^{1/2}$

69. $(2w-1)^{2/3} - w^{1/3} = 0$

70. $(x^2-2x)^{1/3} - x^{1/3} = 0$

Solve each formula for the specified variable. See Example 7. (Data from Cooke, N., and J. Orleans, *Mathematics Essential to Electricity and Radio*, McGraw-Hill.)

71. $Z = \sqrt{\frac{L}{C}}$ for L

72. $r = \sqrt{\frac{\mathcal{A}}{\pi}}$ for \mathcal{A}

73. $V = \sqrt{\frac{2K}{m}}$ for K

74. $V = \sqrt{\frac{2K}{m}}$ for m


75. $r = \sqrt{\frac{Mm}{F}}$ for M

76. $r = \sqrt{\frac{Mm}{F}}$ for F

To find the rotational rate N of a space station, the formula

$$N = \frac{1}{2\pi} \sqrt{\frac{a}{r}}$$

can be used. Here, a is the acceleration and r represents the radius of the space station in meters. To find the value of r that will make N simulate the effect of gravity on Earth, the equation must be solved for r , using the required value of N . (Data from Kastner, B., *Space Mathematics*, NASA.)



77. Solve the equation for the indicated variable.

- (a) for r (b) for a

78. If $a = 9.8$ m per sec², find the value of r (to the nearest tenth) using each value of N .

- (a) $N = 0.063$ rotation per sec (b) $N = 0.04$ rotation per sec

7.7 Complex Numbers

OBJECTIVES

- 1 Simplify numbers of the form $\sqrt{-b}$, where $b > 0$.
- 2 Identify subsets of the complex numbers.
- 3 Add and subtract complex numbers.
- 4 Multiply complex numbers.
- 5 Divide complex numbers.
- 6 Simplify powers of i .

VOCABULARY

- complex number
- real part
- imaginary part
- nonreal complex number
- pure imaginary number
- complex conjugates

Recall that the set of real numbers includes many other number sets—the rational numbers, integers, and natural numbers, for example. In this section, we introduce a new set of numbers that includes the set of real numbers, as well as numbers that are even roots of negative numbers, like $\sqrt{-2}$.

OBJECTIVE 1 Simplify numbers of the form $\sqrt{-b}$, where $b > 0$.

The equation $x^2 + 1 = 0$ has no real number solution because any solution must be a number whose square is -1 . In the set of real numbers, all squares are *nonnegative* numbers because the product of two positive numbers or two negative numbers is positive and $0^2 = 0$. To provide a solution of the equation

$$x^2 + 1 = 0,$$

we introduce a new number i .

Imaginary Unit i

The **imaginary unit** i is defined as follows.

$$i = \sqrt{-1}, \text{ and thus } i^2 = -1$$

That is, i is the principal square root of -1 .

We can use this definition to define any square root of a negative real number.

Meaning of $\sqrt{-b}$

For any positive real number b , $\sqrt{-b} = i\sqrt{b}$.

NOW TRY EXERCISE 1

Write each number as a product of a real number and i .

- (a) $\sqrt{-49}$ (b) $-\sqrt{-121}$
 (c) $\sqrt{-3}$ (d) $\sqrt{-32}$

EXAMPLE 1 Simplifying Square Roots of Negative Numbers

Write each number as a product of a real number and i .

- (a) $\sqrt{-100} = i\sqrt{100} = 10i$ (b) $-\sqrt{-36} = -i\sqrt{36} = -6i$
 (c) $\sqrt{-2} = i\sqrt{2}$ (d) $\sqrt{-54} = i\sqrt{54} = i\sqrt{9 \cdot 6} = 3i\sqrt{6}$

NOW TRY 

CAUTION It is easy to mistake $\sqrt{2}i$ for $\sqrt{2i}$ with the i under the radical. For this reason, we usually write $\sqrt{2}i$ as $i\sqrt{2}$, as in the definition of $\sqrt{-b}$.

When finding a product such as $\sqrt{-4} \cdot \sqrt{-9}$, we cannot use the product rule for radicals because it applies only to *nonnegative* radicands.

For this reason, we change $\sqrt{-b}$ to the form $i\sqrt{b}$ before performing any multiplications or divisions.

NOW TRY ANSWERS

1. (a) $7i$ (b) $-11i$
 (c) $i\sqrt{3}$ (d) $4i\sqrt{2}$


NOW TRY EXERCISE 2

Multiply.

- (a) $\sqrt{-4} \cdot \sqrt{-16}$
 (b) $\sqrt{-5} \cdot \sqrt{-11}$
 (c) $\sqrt{-3} \cdot \sqrt{-12}$
 (d) $\sqrt{13} \cdot \sqrt{-2}$

EXAMPLE 2 Multiplying Square Roots of Negative Numbers

Multiply.

(a) $\sqrt{-4} \cdot \sqrt{-9}$

First write all square roots in terms of i .

$$= i\sqrt{4} \cdot i\sqrt{9} \quad \sqrt{-b} = i\sqrt{b}$$

$$= i \cdot 2 \cdot i \cdot 3 \quad \text{Take square roots.}$$

$$= 6i^2 \quad \text{Multiply.}$$

$$= 6(-1) \quad \text{Substitute } -1 \text{ for } i^2.$$

$$= -6$$

(b) $\sqrt{-3} \cdot \sqrt{-7}$

First write all square roots in terms of i .

$$= i\sqrt{3} \cdot i\sqrt{7} \quad \sqrt{-b} = i\sqrt{b}$$

$$= i^2\sqrt{3 \cdot 7} \quad \text{Product rule}$$

$$= (-1)\sqrt{21} \quad \text{Substitute } -1 \text{ for } i^2.$$

$$= -\sqrt{21} \quad (-1)a = -a$$

(c) $\sqrt{-2} \cdot \sqrt{-8}$

$$= i\sqrt{2} \cdot i\sqrt{8} \quad \sqrt{-b} = i\sqrt{b}$$

$$= i^2\sqrt{2 \cdot 8} \quad \text{Product rule}$$

$$= (-1)\sqrt{16} \quad i^2 = -1$$

$$= -4 \quad \text{Take the square root.}$$

(d) $\sqrt{-5} \cdot \sqrt{6}$

$$= i\sqrt{5} \cdot \sqrt{6}$$

$$= i\sqrt{30}$$

NOW TRY 

! CAUTION Using the product rule for radicals *before* using the definition of $\sqrt{-b}$ gives an *incorrect* answer. **Example 2(a)** shows that

$$\sqrt{-4} \cdot \sqrt{-9} = -6, \quad \text{Correct (Example 2(a))}$$

but $\sqrt{-4(-9)} = \sqrt{36} = 6. \quad \text{Incorrect}$

Thus, $\sqrt{-4} \cdot \sqrt{-9} \neq \sqrt{-4(-9)}.$


NOW TRY EXERCISE 3

Divide.

(a) $\frac{\sqrt{-72}}{\sqrt{-8}}$ (b) $\frac{\sqrt{-48}}{\sqrt{3}}$

EXAMPLE 3 Dividing Square Roots of Negative Numbers

Divide.

(a) $\frac{\sqrt{-75}}{\sqrt{-3}}$

First write all square roots in terms of i .

$$= \frac{i\sqrt{75}}{i\sqrt{3}}$$

$$= \sqrt{\frac{75}{3}} \quad \frac{i}{i} = 1; \text{ Quotient rule}$$

$$= \sqrt{25} \quad \text{Divide.}$$

$$= 5$$

(b) $\frac{\sqrt{-32}}{\sqrt{8}}$

$$= \frac{i\sqrt{32}}{\sqrt{8}} \quad \sqrt{-32} = i\sqrt{32}$$

$$= i\sqrt{\frac{32}{8}} \quad \text{Quotient rule}$$

$$= i\sqrt{4} \quad \text{Divide.}$$

$$= 2i$$

NOW TRY ANSWERS

2. (a) -8 (b) $-\sqrt{55}$
 (c) -6 (d) $i\sqrt{26}$
 3. (a) 3 (b) $4i$

NOW TRY 

OBJECTIVE 2 Identify subsets of the complex numbers.

A new set of numbers, the *complex numbers*, is defined as follows.

Complex Number

If a and b are real numbers, then any number of the form

$$a + bi$$

\uparrow \uparrow
 Real part Imaginary part

is a **complex number**. In the complex number $a + bi$, the number a is the **real part** and b is the **imaginary part**.*

The following important concepts apply to a complex number $a + bi$.

1. If $b = 0$, then $a + bi = a$, which is a real number.

Thus, the set of real numbers is a subset of the set of complex numbers. See FIGURE 14.

2. If $b \neq 0$, then $a + bi$ is a **nonreal complex number**.

Examples: $7 + 2i$, $-1 - i$

3. If $a = 0$ and $b \neq 0$, then the nonreal complex number is a **pure imaginary number**.

Examples: $3i$, $-16i$

A complex number written in the form $a + bi$ is in **standard form**. In this section, most answers will be given in standard form, but if $a = 0$ or $b = 0$, we consider answers such as a or bi to be in standard form.

The relationships among the various sets of numbers are shown in FIGURE 14.




FIGURE 14

*Some texts define bi as the imaginary part of the complex number $a + bi$.

OBJECTIVE 3 Add and subtract complex numbers.

The commutative, associative, and distributive properties for real numbers are also valid for complex numbers.

To add complex numbers, we add their real parts and add their imaginary parts.

 **NOW TRY EXERCISE 4**

Add.

(a) $(-3 + 2i) + (4 + 7i)$

(b) $(5 - i) + (-3 + 3i) + (6 - 4i)$

EXAMPLE 4 Adding Complex Numbers

Add.

(a) $(2 + 3i) + (6 + 4i)$

$$= (2 + 6) + (3 + 4)i$$

Commutative, associative, and distributive properties

$$= 8 + 7i$$

Add real parts. Add imaginary parts.

(b) $(4 + 2i) + (3 - i) + (-6 + 3i)$

$$= [4 + 3 + (-6)] + [2 + (-1) + 3]i$$

Commutative, associative, and distributive properties

$$= 1 + 4i$$

Add real parts. Add imaginary parts.

NOW TRY 

To subtract complex numbers, we subtract their real parts and subtract their imaginary parts.

 **NOW TRY EXERCISE 5**

Subtract.

(a) $(7 + 10i) - (3 + 5i)$

(b) $(5 - 2i) - (9 - 7i)$

(c) $(-1 + 12i) - (-1 - i)$

EXAMPLE 5 Subtracting Complex Numbers

Subtract.

(a) $(6 + 5i) - (3 + 2i)$

$$= (6 - 3) + (5 - 2)i$$

Commutative, associative, and distributive properties

$$= 3 + 3i$$

Subtract real parts. Subtract imaginary parts.

(b) $(7 - 3i) - (8 - 6i)$

$$= (7 - 8) + [-3 - (-6)]i$$

$$= -1 + 3i$$

(c) $(-9 + 4i) - (-9 + 8i)$

$$= (-9 + 9) + (4 - 8)i$$

$$= 0 - 4i$$

Be careful.

$$= -4i$$

NOW TRY **OBJECTIVE 4** Multiply complex numbers.

We multiply complex numbers in the same way that we multiply polynomials.

EXAMPLE 6 Multiplying Complex Numbers

Multiply.

(a) $4i(2 + 3i)$

$$= 4i(2) + 4i(3i)$$

Distributive property

$$= 8i + 12i^2$$

Multiply.

$$= 8i + 12(-1)$$

Substitute -1 for i^2 .

$$= -12 + 8i$$

Standard form

NOW TRY ANSWERS

4. (a) $1 + 9i$ (b) $8 - 2i$

5. (a) $4 + 5i$ (b) $-4 + 5i$

(c) $13i$


NOW TRY EXERCISE 6

Multiply.

(a) $8i(3 - 5i)$

(b) $(7 - 2i)(4 + 3i)$

(b) $(3 + 5i)(4 - 2i)$

$$= \underbrace{3(4)}_{\text{First}} + \underbrace{3(-2i)}_{\text{Outer}} + \underbrace{5i(4)}_{\text{Inner}} + \underbrace{5i(-2i)}_{\text{Last}}$$

Use the FOIL method.

$$= 12 - 6i + 20i - 10i^2$$

Multiply.

$$= 12 + 14i - 10(-1)$$

Add imaginary parts; $i^2 = -1$

$$= 12 + 14i + 10$$

Multiply.

$$= 22 + 14i$$

Add real parts.

(c) $(2 + 3i)(1 - 5i)$

$$= 2(1) + 2(-5i) + 3i(1) + 3i(-5i)$$

FOIL method

$$= 2 - 10i + 3i - 15i^2$$

Multiply.

$$= 2 - 7i - 15(-1)$$

Add imaginary parts; $i^2 = -1$

$$= 2 - 7i + 15$$

Multiply.

$$= 17 - 7i$$

Add real parts.

NOW TRY 

The two complex numbers $a + bi$ and $a - bi$ are **complex conjugates**, or simply *conjugates*, of each other. **The product of a complex number and its conjugate is always a real number**, as shown here.

$$(a + bi)(a - bi)$$

$$= a^2 - abi + abi - b^2i^2$$

FOIL method

$$= a^2 - b^2(-1)$$

Combine like terms; $i^2 = -1$

$$= a^2 + b^2$$

The product eliminates i .

$$\text{Example: } (3 + 7i)(3 - 7i) = 3^2 + 7^2 = 9 + 49 = 58$$

OBJECTIVE 5 Divide complex numbers.**EXAMPLE 7** Dividing Complex Numbers

Divide.

(a) $\frac{8 + 9i}{5 + 2i}$

$$\frac{5 - 2i}{5 - 2i} = 1$$

$$= \frac{(8 + 9i)(5 - 2i)}{(5 + 2i)(5 - 2i)}$$

Multiply numerator and denominator by $5 - 2i$, the conjugate of the denominator.

$$= \frac{40 - 16i + 45i - 18i^2}{5^2 + 2^2}$$

In the denominator, $(a + bi)(a - bi) = a^2 + b^2$.

$$= \frac{40 + 29i - 18(-1)}{25 + 4}$$

In the numerator, add imaginary parts; $i^2 = -1$

$$= \frac{58 + 29i}{29}$$

Multiply. Add real parts.

Add in the denominator.

$$= \frac{29(2 + i)}{29}$$

Factor the numerator.

$$= 2 + i$$

Lowest terms

NOW TRY ANSWERS6. (a) $40 + 24i$ (b) $34 + 13i$

Factor first. Then divide out the common factor.


NOW TRY EXERCISE 7

Find each quotient.

(a) $\frac{4 + 2i}{1 + 3i}$ (b) $\frac{5 - 4i}{i}$

$$\begin{aligned}
 \text{(b)} \quad & \frac{1 + i}{i} \\
 &= \frac{(1 + i)(-i)}{i(-i)} && \text{Multiply numerator and denominator by } -i, \\
 & && \text{the conjugate of } i. \\
 &= \frac{-i - i^2}{-i^2} && \text{Use the distributive property in the numerator.} \\
 & && \text{Multiply in the denominator.} \\
 &= \frac{-i - (-1)}{-(-1)} && \text{Substitute } -1 \text{ for } i^2. \\
 & && \text{Use parentheses to avoid errors.} \\
 &= \frac{-i + 1}{1} \\
 &= 1 - i && \frac{a}{1} = a
 \end{aligned}$$

NOW TRY 
OBJECTIVE 6 Simplify powers of i .
Powers of i can be simplified using the facts

$$i^2 = -1 \quad \text{and} \quad i^4 = (i^2)^2 = (-1)^2 = 1.$$

Consider the following powers of i .

$$\begin{aligned}
 i^1 &= i && i^5 = i^4 \cdot i = 1 \cdot i = i \\
 i^2 &= -1 && i^6 = i^4 \cdot i^2 = 1(-1) = -1 \\
 i^3 &= i^2 \cdot i = (-1) \cdot i = -i && i^7 = i^4 \cdot i^3 = 1 \cdot (-i) = -i \\
 i^4 &= i^2 \cdot i^2 = (-1)(-1) = 1 && i^8 = i^4 \cdot i^4 = 1 \cdot 1 = 1, \quad \text{and so on.}
 \end{aligned}$$

Powers of i cycle through the same four outcomes

$$i, \quad -1, \quad -i, \quad \text{and} \quad 1$$

because i^4 has the same multiplicative property as 1. Also, any power of i with an exponent that is a multiple of 4 has value 1. As with real numbers, $i^0 = 1$.

NOW TRY EXERCISE 8
Find each power of i .

(a) i^{16} (b) i^{21}
 (c) i^{-6} (d) i^{-13}

EXAMPLE 8 Simplifying Powers of i
Find each power of i .

$$\begin{aligned}
 \text{(a)} \quad & i^{12} = (i^4)^3 = 1^3 = 1 && i^4 = 1 \\
 \text{(b)} \quad & i^{39} = i^{36} \cdot i^3 = (i^4)^9 \cdot i^3 = 1^9 \cdot (-i) = 1 \cdot (-i) = -i \\
 \text{(c)} \quad & i^{-2} = \frac{1}{i^2} = \frac{1}{-1} = -1 \\
 \text{(d)} \quad & i^{-15} = \frac{1}{i^{15}} = \frac{1}{(i^4)^3 \cdot i^3} = \frac{1}{1^3 \cdot (-i)} = \frac{1}{-i}
 \end{aligned}$$

We divide by multiplying the numerator and denominator by i , the conjugate of $-i$.

$$\frac{1}{-i} = \frac{1(i)}{-i(i)} = \frac{i}{-i^2} = \frac{i}{-(-1)} = \frac{i}{1} = i$$

NOW TRY 
NOW TRY ANSWERS

7. (a) $1 - i$ (b) $-4 - 5i$
 8. (a) 1 (b) i (c) -1 (d) $-i$

NOTE In **Example 8(d)**, we could also multiply by a power of i that is a multiple of 4 (because $i^4 = 1$).

$$i^{-15} = i^{-15} \cdot i^{16} = i^1 = i \quad i^{16} = 1$$

7.7 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check List all of the following sets to which each number belongs. A number may belong to more than one set.

real numbers pure imaginary numbers nonreal complex numbers complex numbers

1. $3 + 5i$

2. $-7i$

3. $\sqrt{2}$

4. $\frac{13}{3}$

5. $\sqrt{-49}$

6. $-\sqrt{-8}$

Concept Check Decide whether each expression is equal to 1, -1 , i , or $-i$.

7. $\sqrt{-1}$

8. $-\sqrt{-1}$

9. i^2

10. $-i^2$

11. $\frac{1}{i}$

12. $(-i)^2$

Write each number as a product of a real number and i . Simplify all radical expressions. See Example 1.

13. $\sqrt{-169}$

14. $\sqrt{-225}$

15. $-\sqrt{-144}$

16. $-\sqrt{-196}$

17. $\sqrt{-5}$

18. $\sqrt{-21}$

19. $\sqrt{-48}$

20. $\sqrt{-96}$

21. **Concept Check** A student incorrectly multiplied as follows.

$$\sqrt{-9} \cdot \sqrt{-25} = \sqrt{225} = 15$$

WHAT WENT WRONG? Find the product correctly.

22. **Concept Check** A student incorrectly simplified $-\sqrt{-16}$ as follows.

$$-\sqrt{-16} = -(-4) = 4$$

WHAT WENT WRONG? Simplify the expression correctly.

Multiply or divide as indicated. See Examples 2 and 3.

23. $\sqrt{-15} \cdot \sqrt{-15}$

24. $\sqrt{-19} \cdot \sqrt{-19}$

25. $\sqrt{-7} \cdot \sqrt{-15}$

26. $\sqrt{-3} \cdot \sqrt{-19}$

27. $\sqrt{-4} \cdot \sqrt{-25}$

28. $\sqrt{-9} \cdot \sqrt{-81}$

29. $\sqrt{-3} \cdot \sqrt{11}$

30. $\sqrt{-5} \cdot \sqrt{13}$

31. $\sqrt{5} \cdot \sqrt{-30}$

32. $\sqrt{-10} \cdot \sqrt{2}$

33. $\frac{\sqrt{-300}}{\sqrt{-100}}$

34. $\frac{\sqrt{-40}}{\sqrt{-10}}$

35. $\frac{\sqrt{-75}}{\sqrt{3}}$

36. $\frac{\sqrt{-160}}{\sqrt{10}}$

37. $\frac{-\sqrt{-64}}{\sqrt{-16}}$

38. $\frac{-\sqrt{-100}}{\sqrt{-25}}$

Add or subtract as indicated. Give answers in standard form. See Examples 4 and 5.

39. $(3 + 2i) + (-4 + 5i)$

40. $(7 + 15i) + (-11 + 14i)$

41. $(5 - i) + (-5 + i)$

42. $(-2 + 6i) + (2 - 6i)$

43. $(4 + i) - (-3 - 2i)$

44. $(9 + i) - (3 + 2i)$

45. $(-3 - 4i) - (-1 - 4i)$

46. $(-2 - 3i) - (-5 - 3i)$

47. $(-4 + 11i) + (-2 - 4i) + (7 + 6i)$

48. $(-1 + i) + (2 + 5i) + (3 + 2i)$

49. $[(7 + 3i) - (4 - 2i)] + (3 + i)$

50. $[(7 + 2i) + (-4 - i)] - (2 + 5i)$

Concept Check Fill in the blank with the correct response.

51. Because $(4 + 2i) - (3 + i) = 1 + i$, using the definition of subtraction we can check this to find that

$$(1 + i) + (3 + i) = \underline{\hspace{2cm}}.$$

52. Because $\frac{-5}{2-i} = -2 - i$, using the definition of division we can check this to find that

$$(-2 - i)(2 - i) = \underline{\hspace{2cm}}.$$

Multiply. Give answers in standard form. See Example 6.

- | | | |
|--------------------------|--------------------------|--------------------------|
| 53. $(3i)(27i)$ | 54. $(5i)(125i)$ | 55. $(-8i)(-2i)$ |
| 56. $(-32i)(-2i)$ | 57. $5i(-6 + 2i)$ | 58. $3i(4 + 9i)$ |
| 59. $(4 + 3i)(1 - 2i)$ | 60. $(7 - 2i)(3 + i)$ | 61. $(4 + 5i)^2$ |
| 62. $(3 + 2i)^2$ | 63. $2i(-4 - i)^2$ | 64. $3i(-3 - i)^2$ |
| 65. $(12 + 3i)(12 - 3i)$ | 66. $(6 + 7i)(6 - 7i)$ | 67. $(4 + 9i)(4 - 9i)$ |
| 68. $(7 + 2i)(7 - 2i)$ | 69. $(1 + i)^2(1 - i)^2$ | 70. $(2 - i)^2(2 + i)^2$ |

Concept Check Answer each of the following.

71. Let a and b represent real numbers.
- What is the conjugate of $a + bi$?
 - If we multiply $a + bi$ by its conjugate, we obtain $\underline{\hspace{2cm}} + \underline{\hspace{2cm}}$, which is always a real number.
72. By what complex number should we multiply the numerator and denominator of $\frac{2 + i\sqrt{2}}{2 - i\sqrt{2}}$ to write the quotient in standard form?

- A. $\sqrt{2}$ B. $i\sqrt{2}$ C. $2 + i\sqrt{2}$ D. $2 - i\sqrt{2}$

Divide. Give answers in standard form. See Example 7.

- | | | | |
|--------------------------|---------------------------|-------------------------|--------------------------|
| 73. $\frac{2}{1-i}$ | 74. $\frac{2}{1+i}$ | 75. $\frac{8i}{2+2i}$ | 76. $\frac{-8i}{1+i}$ |
| 77. $\frac{-7+4i}{3+2i}$ | 78. $\frac{-38-8i}{7+3i}$ | 79. $\frac{2-3i}{2+3i}$ | 80. $\frac{-1+5i}{3+2i}$ |
| 81. $\frac{3+i}{i}$ | 82. $\frac{5-i}{i}$ | 83. $\frac{3-i}{-i}$ | 84. $\frac{5+i}{-i}$ |

Find each power of i . See Example 8.

- | | | | |
|--------------|---------------|---------------|---------------|
| 85. i^{18} | 86. i^{26} | 87. i^{89} | 88. i^{48} |
| 89. i^{38} | 90. i^{102} | 91. i^{43} | 92. i^{83} |
| 93. i^{-5} | 94. i^{-17} | 95. i^{-20} | 96. i^{-27} |

Concept Check Provide a short response for each problem.

97. A student simplified i^{-18} as shown. Provide justification for this correct work.

$$i^{-18} = i^{-18} \cdot i^{20} = i^{-18+20} = i^2 = -1$$

98. Explain why the following two expressions must be equal. (Do not actually perform the computation.)

$$(46 + 25i)(3 - 6i) \quad \text{and} \quad (46 + 25i)(3 - 6i)i^{12}$$

Ohm's law for the current I in a circuit with voltage E , resistance R , capacitive reactance X_c , and inductive reactance X_L is

$$I = \frac{E}{R + (X_L - X_c)i}$$

Use this law to work each exercise.

99. Find I if $E = 2 + 3i$, $R = 5$, $X_L = 4$, and $X_c = 3$.
 100. Find E if $I = 1 - i$, $R = 2$, $X_L = 3$, and $X_c = 1$.

RELATING CONCEPTS For Individual or Group Work (Exercises 101–105)

Some equations have nonreal complex solutions. **Work Exercises 101–105 in order**, to see how these nonreal complex solutions are related.

101. Show that $1 + 5i$ is a solution of $x^2 - 2x + 26 = 0$.
 102. Show that $1 - 5i$ is a solution of $x^2 - 2x + 26 = 0$.
 103. From **Exercises 101 and 102**, the nonreal complex solutions of the equation

$$x^2 - 2x + 26 = 0$$
 are $1 + 5i$ and $1 - 5i$. What do we call two complex numbers $a + bi$ and $a - bi$?
 104. Show that $3 + 2i$ is a solution of the equation $x^2 - 6x + 13 = 0$.
 105. Using the results of **Exercises 101–104**, make a conjecture about another nonreal complex solution of the equation $x^2 - 6x + 13 = 0$, and verify it.

Chapter 7

Summary

Key Terms

7.1
 radicand
 index (order)
 radical
 principal root
 radical expression
 square root function
 cube root function

7.3
 hypotenuse
 legs (of a right triangle)
 circle
 center
 radius

7.5
 conjugates
7.6
 radical equation
 proposed solution
 extraneous solution

7.7
 complex number
 real part
 imaginary part
 nonreal complex number
 pure imaginary number
 complex conjugates

New Symbols

$\sqrt{\quad}$ radical symbol

\pm “positive or negative” or
 “plus or minus”

$a^{1/n}$ a to the power $\frac{1}{n}$

i imaginary unit

$\sqrt[n]{a}$ radical; principal n th
 root of a

\approx is approximately equal to

$a^{m/n}$ a to the power $\frac{m}{n}$

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- A **radicand** is
 - the index of a radical
 - the number or expression under the radical symbol
 - the positive root of a number
 - the radical symbol.
- The **Pythagorean theorem** states that, in a right triangle,
 - the sum of the measures of the angles is 180°
 - the sum of the lengths of the two shorter sides equals the length of the longest side
 - the longest side is opposite the right angle
 - the square of the length of the longest side equals the sum of the squares of the lengths of the two shorter sides.
- A **hypotenuse** is
 - either of the two shorter sides of a triangle
 - the shortest side of a triangle
 - the side opposite the right angle in a triangle
 - the longest side in any triangle.
- Rationalizing the denominator** is the process of
 - eliminating fractions from a radical expression
 - changing the denominator of a fraction from a radical expression to a rational number
 - clearing a radical expression of radicals
 - multiplying radical expressions.
- An **extraneous solution** is a value
 - that does not satisfy the original equation
 - that makes an equation true
 - that makes an expression equal 0
 - that checks in the original equation.
- A **complex number** is
 - a real number that includes a complex fraction
 - a zero multiple of i
 - a number of the form $a + bi$, where a and b are real numbers
 - the square root of -1 .

ANSWERS

- B; *Example:* In $\sqrt{3xy}$, $3xy$ is the radicand. 2. D; *Example:* In a right triangle where $a = 6$, $b = 8$, and $c = 10$, $6^2 + 8^2 = 10^2$.
- C; *Example:* In a right triangle where the sides measure 9, 12, and 15 units, the hypotenuse is the side opposite the right angle, with measure 15 units.
- B; *Example:* To rationalize the denominator of $\frac{5}{\sqrt{3}+1}$, multiply both numerator and denominator by $\sqrt{3}-1$ to obtain $\frac{5(\sqrt{3}-1)}{2}$.
- A; *Example:* The proposed solution 2 is extraneous in $\sqrt{5x-1} + 3 = 0$. 6. C; *Examples:* -5 (or $-5 + 0i$), $7i$ (or $0 + 7i$), $\sqrt{2} - 4i$

Quick Review

CONCEPTS

7.1 Radical Expressions and Graphs

$$\sqrt[n]{a} = b \text{ means } b^n = a.$$

$\sqrt[n]{a}$ is the **principal n th root** of a .

$$\sqrt[n]{a^n} = |a| \text{ if } n \text{ is even. } \sqrt[n]{a^n} = a \text{ if } n \text{ is odd.}$$

Functions Defined by Radical Expressions

The square root function is

$$f(x) = \sqrt{x}.$$

The cube root function is

$$f(x) = \sqrt[3]{x}.$$


EXAMPLES

Find each root.

$$\sqrt{64} = 8 \quad \text{Principal square root}$$


$$-\sqrt{64} = -8 \quad \sqrt{-64} \text{ is not a real number.}$$

$$\sqrt[4]{(-2)^4} = |-2| = 2 \quad \sqrt[3]{-27} = -3$$



Square root function

Domain and range: $[0, \infty)$



Cube root function

Domain and range: $(-\infty, \infty)$

CONCEPTS

7.2 Rational Exponents

$$a^{1/n} = \sqrt[n]{a} \text{ whenever } \sqrt[n]{a} \text{ exists.}$$

If m and n are positive integers with m/n in lowest terms, then

$$a^{m/n} = (a^{1/n})^m, \text{ provided that } a^{1/n} \text{ is a real number.}$$

All of the usual definitions and rules for exponents are valid for rational exponents.

7.3 Simplifying Radicals, the Distance Formula, and Circles

Product and Quotient Rules for Radicals

If n is a natural number and $\sqrt[n]{a}$ and $\sqrt[n]{b}$ are real numbers, then the following hold true.

$$\sqrt[n]{a} \cdot \sqrt[n]{b} = \sqrt[n]{ab} \quad \text{and} \quad \sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}} \quad (\text{where } b \neq 0)$$

Conditions for a Simplified Radical

1. The radicand has no factor raised to a power greater than or equal to the index.
2. The radicand has no fractions.
3. No denominator contains a radical.
4. Exponents in the radicand and the index of the radical have greatest common factor 1.

Pythagorean Theorem

If a and b are the lengths of the shorter sides of a right triangle and c is the length of the longest side, then the following holds.

$$a^2 + b^2 = c^2$$

Distance Formula

The distance d between the points (x_1, y_1) and (x_2, y_2) is given by the following.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

EXAMPLES

Apply the rules for rational exponents.

$$81^{1/2} = \sqrt{81} = 9 \quad -64^{1/3} = -\sqrt[3]{64} = -4$$

$$8^{5/3} = (8^{1/3})^5 = 2^5 = 32 \quad (y^{2/5})^{10} = y^4$$

Write with positive exponents.

$$\begin{aligned} 5^{-1/2} \cdot 5^{1/4} &= 5^{-1/2+1/4} \\ &= 5^{-1/4} \\ &= \frac{1}{5^{1/4}} \end{aligned} \quad \left| \quad \begin{aligned} \frac{x^{-1/3}}{x^{-1/2}} &= x^{-1/3-(-1/2)} \\ &= x^{-1/3+1/2} \\ &= x^{1/6}, \quad x > 0 \end{aligned}$$

Simplify.

$$\sqrt{3} \cdot \sqrt{7} = \sqrt{21} \quad \sqrt[5]{x^3y} \cdot \sqrt[5]{xy^2} = \sqrt[5]{x^4y^3}$$

$$\frac{\sqrt{x^5}}{\sqrt{x^4}} = \sqrt{\frac{x^5}{x^4}} = \sqrt{x}, \quad x > 0$$

$$\sqrt{18} = \sqrt{9 \cdot 2} = \sqrt{9} \cdot \sqrt{2} = 3\sqrt{2}$$

$$\sqrt[3]{54x^5y^3} = \sqrt[3]{27x^3y^3 \cdot 2x^2} = \sqrt[3]{27x^3y^3} \cdot \sqrt[3]{2x^2} = 3xy\sqrt[3]{2x^2}$$

$$\sqrt{\frac{7}{4}} = \frac{\sqrt{7}}{\sqrt{4}} = \frac{\sqrt{7}}{2}$$

$$\sqrt[9]{x^3} = x^{3/9} = x^{1/3}, \quad \text{or} \quad \sqrt[3]{x}$$

Find the length of the unknown side in the right triangle.


$$10^2 + b^2 = (2\sqrt{61})^2$$

$$100 + b^2 = 4(61)$$

$$100 + b^2 = 244$$

$$b^2 = 144$$

$$b = 12$$



Find the distance between $(3, -2)$ and $(-1, 1)$.

$$d = \sqrt{(-1 - 3)^2 + [1 - (-2)]^2} \quad \text{Substitute.}$$

$$d = \sqrt{(-4)^2 + 3^2} \quad \text{Subtract.}$$

$$d = \sqrt{16 + 9} \quad \text{Square.}$$

$$d = \sqrt{25} \quad \text{Add.}$$

$$d = 5 \quad \text{Take the square root.}$$

CONCEPTS

Equation of a Circle

A circle with center (h, k) and radius $r > 0$ has an equation of the form

$$(x - h)^2 + (y - k)^2 = r^2.$$

A circle with center at the origin $(0, 0)$ and radius $r > 0$ has the following equation.

$$x^2 + y^2 = r^2$$


EXAMPLES

Graph $(x + 2)^2 + (y - 3)^2 = 25$.

This equation, which can be written

$$[x - (-2)]^2 + (y - 3)^2 = 5^2,$$

is a circle with center $(-2, 3)$ and radius 5.



7.4 Adding and Subtracting Radical Expressions

Add or subtract radical expressions using the distributive property.

$$ac + bc = (a + b)c$$

Only radical expressions with the same index and the same radicand may be combined.

Simplify.

$$\begin{aligned} 2\sqrt{28} - 3\sqrt{63} + 8\sqrt{112} &= 2\sqrt{4 \cdot 7} - 3\sqrt{9 \cdot 7} + 8\sqrt{16 \cdot 7} \\ &= 2 \cdot 2\sqrt{7} - 3 \cdot 3\sqrt{7} + 8 \cdot 4\sqrt{7} \\ &= 4\sqrt{7} - 9\sqrt{7} + 32\sqrt{7} \\ &= (4 - 9 + 32)\sqrt{7} \\ &= 27\sqrt{7} \end{aligned}$$

$$\left. \begin{array}{l} \sqrt{15} + \sqrt{30} \\ \sqrt{3} + \sqrt[3]{9} \end{array} \right\} \text{Cannot be simplified further}$$

7.5 Multiplying and Dividing Radical Expressions

Multiply binomial radical expressions using the FOIL method. Special product rules may apply.

$$\begin{aligned} (x + y)(x - y) &= x^2 - y^2 \\ (x + y)^2 &= x^2 + 2xy + y^2 \\ (x - y)^2 &= x^2 - 2xy + y^2 \end{aligned}$$

Rationalizing a Denominator

Rationalize the denominator by multiplying both the numerator and the denominator by the same expression, one that will yield a rational number in the final denominator.

To write a radical quotient in lowest terms, factor the numerator and denominator. Then divide out any common factor(s).

Perform the operations and simplify.

$$\begin{aligned} (\sqrt{2} + \sqrt{7})(\sqrt{3} - \sqrt{6}) &= \sqrt{6} - 2\sqrt{3} + \sqrt{21} - \sqrt{42} \quad \sqrt{12} = 2\sqrt{3} \\ (\sqrt{5} - \sqrt{10})(\sqrt{5} + \sqrt{10}) &= 5 - 10 \\ &= -5 \end{aligned} \quad \left| \quad \begin{aligned} (\sqrt{3} - \sqrt{2})^2 &= 3 - 2\sqrt{3} \cdot \sqrt{2} + 2 \\ &= 5 - 2\sqrt{6} \end{aligned}$$

$$\frac{\sqrt{7}}{\sqrt{5}} = \frac{\sqrt{7} \cdot \sqrt{5}}{\sqrt{5} \cdot \sqrt{5}} = \frac{\sqrt{35}}{5}$$

$$\begin{aligned} \frac{4}{\sqrt{5} - \sqrt{2}} &= \frac{4(\sqrt{5} + \sqrt{2})}{(\sqrt{5} - \sqrt{2})(\sqrt{5} + \sqrt{2})} \\ &= \frac{4(\sqrt{5} + \sqrt{2})}{5 - 2} = \frac{4(\sqrt{5} + \sqrt{2})}{3} \end{aligned}$$

$$\frac{5 + 15\sqrt{6}}{10} = \frac{5(1 + 3\sqrt{6})}{5 \cdot 2} = \frac{1 + 3\sqrt{6}}{2} \quad \text{Factor first, then divide.}$$

CONCEPTS

7.6 Solving Equations with Radicals**Solving a Radical Equation**

- Step 1** Isolate one radical on one side of the equation.
- Step 2** Raise each side of the equation to a power that is the same as the index of the radical.
- Step 3** Solve the resulting equation. If it still contains a radical, repeat Steps 1 and 2.
- Step 4** Check all proposed solutions in the *original* equation. Discard any values that are not solutions of the original equation.

7.7 Complex Numbers

$i = \sqrt{-1}$, and thus $i^2 = -1$.

For any positive real number b , $\sqrt{-b} = i\sqrt{b}$.

To multiply radicals with negative radicands, first change each factor to the form $i\sqrt{b}$ and then multiply. The same procedure applies to quotients.

Adding and Subtracting Complex Numbers

Add (or subtract) the real parts and add (or subtract) the imaginary parts.

Multiplying Complex Numbers

Multiply complex numbers using the FOIL method.

Dividing Complex Numbers

Divide complex numbers by multiplying the numerator and the denominator by the conjugate of the denominator.

EXAMPLES

Solve. $\sqrt{2x+3} - x = 0$

$$\sqrt{2x+3} = x \quad \text{Add } x.$$

$$(\sqrt{2x+3})^2 = x^2 \quad \text{Square each side.}$$

$$2x+3 = x^2 \quad \text{Apply the exponents.}$$

$$x^2 - 2x - 3 = 0 \quad \text{Standard form}$$

$$(x-3)(x+1) = 0 \quad \text{Factor.}$$

$$x-3 = 0 \quad \text{or} \quad x+1 = 0 \quad \text{Zero-factor property}$$

$$x = 3 \quad \text{or} \quad x = -1 \quad \text{Solve each equation.}$$

A check shows that 3 is a solution, but -1 is extraneous (because it leads to $2 = 0$, a false statement). The solution set is $\{3\}$.

Simplify.

$$\sqrt{-25} = i\sqrt{25} = 5i$$

$$\sqrt{-3} \cdot \sqrt{-27}$$

$$= i\sqrt{3} \cdot i\sqrt{27} \quad \sqrt{-b} = i\sqrt{b}$$

$$= i^2\sqrt{81} \quad \text{Product rule}$$

$$= -1 \cdot 9 \quad i^2 = -1; \text{ Take the square root.}$$

$$= -9 \quad \text{Multiply.}$$

$$\frac{\sqrt{-18}}{\sqrt{-2}} = \frac{i\sqrt{18}}{i\sqrt{2}} = \sqrt{\frac{18}{2}} = \sqrt{9} = 3$$

Perform the operations.

$$(5+3i) + (8-7i) \quad | \quad (5+3i) - (8-7i)$$

$$= 13-4i \quad | \quad = -3+10i$$

Multiply. $(2+i)(5-3i)$

$$= 10 - 6i + 5i - 3i^2 \quad \text{Use the FOIL method.}$$

$$= 10 - i - 3(-1) \quad \text{Add imaginary parts; } i^2 = -1$$

$$= 13 - i \quad \text{Multiply. Add real parts.}$$

Divide. $\frac{20}{3+i}$

$$= \frac{20(3-i)}{(3+i)(3-i)} \quad \text{Multiply both numerator and denominator by the conjugate of the denominator.}$$

$$= \frac{20(3-i)}{9-i^2} \quad (x+y)(x-y) = x^2 - y^2$$

$$= \frac{20(3-i)}{10} \quad i^2 = -1 \text{ and } 9 - (-1) = 10.$$

$$= 2(3-i) \quad \text{Divide out the common factor, 10.}$$

$$= 6-2i \quad \text{Distributive property}$$

Chapter 7

Review Exercises

7.1 Find each root.

1. $\sqrt{1764}$

2. $-\sqrt{289}$

3. $\sqrt[3]{216}$

4. $\sqrt[3]{-125}$

5. $-\sqrt[3]{27}$

6. $\sqrt[5]{-32}$

7. $\sqrt{x^2}$

8. $\sqrt[3]{x^3}$

9. $\sqrt[4]{x^{20}}$

10. Under what conditions is $\sqrt[n]{a}$ not a real number?

Graph each function, and give its domain and range.

11. $f(x) = \sqrt{x-1}$

12. $f(x) = \sqrt[3]{x} + 4$

Use a calculator to approximate each radical to three decimal places.

13. $-\sqrt{47}$

14. $\sqrt[3]{-129}$

15. $\sqrt[4]{605}$

16. $\sqrt[4]{500}$

17. $-\sqrt[3]{500}$

18. $-\sqrt{28}$

Solve each problem.

19. The time t in seconds for one complete swing of a pendulum is given by

$$t = 2\pi\sqrt{\frac{L}{g}},$$

where L is the length of the pendulum in feet, and g , the force due to gravity, is about 32 ft per sec². Find the time of a complete swing of a 3-ft pendulum to the nearest tenth of a second.

20. Use Heron's formula

$$A = \sqrt{s(s-a)(s-b)(s-c)},$$

where $s = \frac{1}{2}(a+b+c)$, to find the area of a triangle with sides of lengths 11 in., 13 in., and 20 in.

7.2 Evaluate each exponential.

21. $49^{1/2}$

22. $-121^{1/2}$

23. $16^{5/4}$

24. $-8^{2/3}$

25. $-\left(\frac{36}{25}\right)^{3/2}$

26. $\left(-\frac{1}{8}\right)^{-5/3}$

27. $\left(\frac{81}{10,000}\right)^{-3/4}$

28. $(-16)^{3/4}$

Write each exponential as a radical. Assume that all variables represent positive real numbers. Use the definition that takes the root first.

29. $10x^{1/2}$

30. $(3a+b)^{-5/3}$

Write each radical as an exponential and simplify. Assume that all variables represent positive real numbers.

31. $\sqrt{7^9}$

32. $\sqrt{11^6}$

33. $\sqrt[3]{r^{12}}$

34. $\sqrt[4]{2^4}$

Simplify each expression. Assume that all variables represent positive real numbers.

$$35. 5^{1/4} \cdot 5^{7/4} \qquad 36. \frac{96^{2/3}}{96^{-1/3}} \qquad 37. \frac{(a^{1/3})^4}{a^{2/3}}$$

$$38. \frac{y^{-1/3} \cdot y^{5/6}}{y} \qquad 39. \left(\frac{z^{-1}x^{-3/5}}{2^{-2}z^{-1/2}x} \right)^{-1} \qquad 40. r^{-1/2}(r + r^{3/2})$$

Write each radical as an exponential and simplify. Leave answers in exponential form. Assume that all variables represent positive real numbers.

$$41. \sqrt[5]{y} \cdot \sqrt[3]{y} \qquad 42. \frac{\sqrt{x^3}}{\sqrt{x^4}} \qquad 43. \frac{\sqrt{p^5}}{p^2}$$

$$44. \sqrt[4]{k^3} \cdot \sqrt{k^3} \qquad 45. \sqrt[3]{m^5} \cdot \sqrt[3]{m^8} \qquad 46. \frac{\sqrt[3]{t^5}}{\sqrt[6]{t}}$$

$$47. \sqrt{\sqrt{\sqrt{x}}} \qquad 48. \sqrt[3]{\sqrt[5]{x}} \qquad 49. \sqrt{\sqrt[6]{\sqrt[3]{x}}}$$

50. The product rule does not apply to $3^{1/4} \cdot 2^{1/5}$. Why?

7.3 Simplify. Assume that all variables represent positive real numbers.

$$51. \sqrt{6} \cdot \sqrt{11} \qquad 52. \sqrt{5} \cdot \sqrt{r} \qquad 53. \sqrt[3]{6} \cdot \sqrt[3]{5} \qquad 54. \sqrt[4]{7} \cdot \sqrt[4]{3}$$


$$55. \sqrt{20} \qquad 56. \sqrt{75} \qquad 57. -\sqrt{125} \qquad 58. \sqrt[3]{-108}$$

$$59. \sqrt{100y^7} \qquad 60. \sqrt[3]{64p^4q^6} \qquad 61. \sqrt[3]{108a^8b^5} \qquad 62. \sqrt[3]{632r^8t^4}$$

$$63. \sqrt{\frac{y^3}{144}} \qquad 64. \sqrt[3]{\frac{m^{15}}{27}} \qquad 65. \sqrt[3]{\frac{r^2}{8}} \qquad 66. \sqrt[4]{\frac{a^9}{81}}$$

$$67. \sqrt[6]{15^3} \qquad 68. \sqrt[4]{p^6} \qquad 69. \sqrt[3]{2} \cdot \sqrt[4]{5} \qquad 70. \sqrt{x} \cdot \sqrt[5]{x}$$

71. Find the length c of the hypotenuse of the right triangle shown.



72. Find the distance between the points $(-4, 7)$ and $(10, 6)$.

Find the equation of a circle satisfying the given conditions.

$$73. \text{Center } (0, 0), r = 11 \qquad 74. \text{Center } (-2, 4), r = 3$$

$$75. \text{Center } (-1, -3), r = 5 \qquad 76. \text{Center } (4, 2), r = 6$$

Graph each circle. Identify the center and the radius.

$$77. x^2 + y^2 = 25 \qquad 78. (x + 3)^2 + (y - 3)^2 = 9 \qquad 79. (x - 2)^2 + (y + 5)^2 = 9$$

80. Why does the equation $x^2 + y^2 = -1$ have no points on its graph?

7.4 Perform the indicated operations. Assume that all variables represent positive real numbers.

81. $2\sqrt{8} - 3\sqrt{50}$

82. $8\sqrt{80} - 3\sqrt{45}$

83. $-\sqrt{27y} + 2\sqrt{75y}$

84. $2\sqrt{54m^3} + 5\sqrt{96m^3}$

85. $3\sqrt[3]{54} + 5\sqrt[3]{16}$

86. $-6\sqrt[4]{32} + \sqrt[4]{512}$

7.5 Multiply, and then simplify.

87. $(\sqrt{3} + 1)(\sqrt{3} - 2)$

88. $(\sqrt{7} + \sqrt{5})(\sqrt{7} - \sqrt{5})$

89. $(3\sqrt{2} + 1)(2\sqrt{2} - 3)$

90. $(\sqrt{13} - \sqrt{2})^2$

91. $(\sqrt[3]{2} + 3)(\sqrt[3]{4} - 3\sqrt[3]{2} + 9)$

92. $(\sqrt[3]{4y} - 1)(\sqrt[3]{4y} + 3)$

Rationalize each denominator. Assume that all variables represent positive real numbers.

93. $\frac{\sqrt{6}}{\sqrt{5}}$

94. $\frac{-6\sqrt{3}}{\sqrt{2}}$

95. $\frac{3\sqrt{7p}}{\sqrt{y}}$

96. $\sqrt{\frac{11}{8}}$

97. $-\sqrt[3]{\frac{9}{25}}$

98. $\sqrt[3]{\frac{108m^3}{n^5}}$

99. $\frac{1}{\sqrt{2} + \sqrt{7}}$

100. $\frac{-5}{\sqrt{6} - 3}$

Write each quotient in lowest terms.

101. $\frac{2 - 2\sqrt{5}}{8}$

102. $\frac{-18 + \sqrt{27}}{6}$

7.6 Solve each equation.

103. $\sqrt{8x + 9} = 5$

104. $\sqrt{2x - 3} - 3 = 0$

105. $\sqrt{7x + 1} = x + 1$

106. $3\sqrt{x} = \sqrt{10x - 9}$

107. $\sqrt{x^2 + 3x + 7} = x + 2$

108. $\sqrt{x + 2} - \sqrt{x - 3} = 1$

109. $\sqrt[3]{5x - 1} = \sqrt[3]{3x - 2}$

110. $\sqrt[3]{1 - 2x} - \sqrt[3]{-x - 13} = 0$

111. $\sqrt[4]{x - 1} + 2 = 0$


112. $\sqrt[4]{x + 7} = \sqrt[4]{2x}$

Carpenters stabilize wall frames with a diagonal brace, as shown in the figure. The length of the brace is given by

$$L = \sqrt{H^2 + W^2}.$$

113. Solve this formula for H .

114. If the bottom of the brace is attached 9 ft from the corner and the brace is 12 ft long, how far up the corner post should it be nailed? Give the answer to the nearest tenth of a foot.



7.7 Write each number as a product of a real number and i . Simplify.

115. $\sqrt{-16}$

116. $\sqrt{-200}$

Perform the indicated operations. Give answers in standard form.

117. $(-2 + 5i) + (-8 - 7i)$

118. $(5 + 4i) - (-9 - 3i)$

119. $\sqrt{-5} \cdot \sqrt{-7}$

120. $\sqrt{-25} \cdot \sqrt{-81}$

121. $\frac{\sqrt{-72}}{\sqrt{-8}}$

122. $(2 + 3i)(1 - i)$

123. $(6 - 2i)^2$

124. $\frac{3 - i}{2 + i}$

Find each power of i .

125. i^{11}

126. i^{36}

127. i^{-10}

128. i^{-8}

Chapter 7

Mixed Review Exercises

Simplify. Assume that all variables represent positive real numbers.

1. $-\sqrt{169a^2b^4}$

2. $1000^{-2/3}$

3. $\frac{z^{-1/5} \cdot z^{3/10}}{z^{7/10}}$

4. $\sqrt[3]{54z^9t^8}$

5. $\sqrt{-49}$

6. $\frac{-1}{\sqrt{12}}$

7. $\sqrt[3]{\frac{12}{25}}$

8. i^{-1000}

9. $-5\sqrt{18} + 12\sqrt{72}$

10. $(4 - 9i) - (-1 + 2i)$

11. $\frac{\sqrt{50}}{\sqrt{-2}}$

12. $\frac{3 + \sqrt{54}}{6}$

13. $(3 + 2i)^2$

14. $8\sqrt[3]{x^3y^2} - 2x\sqrt[3]{y^2}$

15. $5i(3 - 7i)$

16. $\sqrt[3]{2} \cdot \sqrt[4]{5}$

17. $\frac{2\sqrt{z}}{\sqrt{z-2}}, z \neq 4$

18. Find the perimeter of a triangular electronic highway road sign having the dimensions shown in the figure.



Solve each equation.

19. $\sqrt{x+4} = x-2$

20. $\sqrt[3]{2x-9} = \sqrt[3]{5x+3}$

21. $\sqrt{6+2x}-1 = \sqrt{7-2x}$

22. $\sqrt{11+2x}+1 = \sqrt{5x+1}$

Chapter 7

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

► View the complete solutions to all Chapter Test exercises in MyLab Math.

Evaluate.

1. $-\sqrt{841}$

2. $\sqrt[3]{-512}$

3. $125^{1/3}$

Use a calculator to approximate each radical to three decimal places.

4. $\sqrt{478}$

5. $\sqrt[3]{-832}$

6. Graph the function $f(x) = \sqrt{x+6}$, and give its domain and range.

Simplify each expression. Leave answers in exponential form. Assume that all variables represent positive real numbers.

7. $\left(\frac{16}{25}\right)^{-3/2}$


8. $(-64)^{-4/3}$

9. $\frac{3^{2/5}x^{-1/4}y^{2/5}}{3^{-8/5}x^{7/4}y^{1/10}}$

10. $\left(\frac{x^{-4}y^{-6}}{x^{-2}y^3}\right)^{-2/3}$

11. $7^{3/4} \cdot 7^{-1/4}$

12. $\sqrt[3]{a^4} \cdot \sqrt[3]{a^7}$

13. Find the exact length of side b in the figure.14. Find the distance between the points $(-4, 2)$ and $(2, 10)$.15. Write an equation of a circle with center $(-4, 6)$ and radius 5.16. Graph the circle $(x-2)^2 + (y+3)^2 = 16$. Identify the center and the radius.

Simplify. Assume that all variables represent positive real numbers.

17. $\sqrt{54x^5y^6}$

18. $\sqrt[4]{32a^7b^{13}}$

19. $\sqrt{2} \cdot \sqrt[3]{5}$

20. $3\sqrt{20} - 5\sqrt{80} + 4\sqrt{500}$

21. $\sqrt[3]{16t^3s^5} - \sqrt[3]{54t^6s^2}$

22. $(7\sqrt{5} + 4)(2\sqrt{5} - 1)$

23. $(\sqrt{3} - 2\sqrt{5})^2$

24. $\frac{-5}{\sqrt{40}}$

25. $\frac{2}{\sqrt[3]{5}}$

26. $\frac{-4}{\sqrt{7} + \sqrt{5}}$

27. Write $\frac{6 + \sqrt{24}}{2}$ in lowest terms.28. The following formula from physics relates the velocity V of sound to the temperature T .

$$V = \frac{V_0}{\sqrt{1 - kT}}$$

(a) Approximate V to the nearest tenth if $V_0 = 50$, $k = 0.01$, and $T = 30$.(b) Solve the formula for T .

Solve each equation.

29. $\sqrt[3]{5x} = \sqrt[3]{2x-3}$

30. $\sqrt{x+6} = 9 - 2x$

31. $\sqrt{x+4} - \sqrt{1-x} = -1$

Perform the indicated operations. Give the answers in standard form.

$$32. (-2 + 5i) - (3 + 6i) - 7i \quad 33. (1 + 5i)(3 + i) \quad 34. \frac{7 + i}{1 - i}$$

35. Simplify i^{37} .

36. Answer *true* or *false* to each of the following.

$$(a) i^2 = -1 \quad (b) i = \sqrt{-1} \quad (c) i = -1 \quad (d) \sqrt{-3} = i\sqrt{3}$$

Chapters R–7 Cumulative Review Exercises

Perform the indicated operations.

$$1. -4|2 - 7| - (-3)(-9) \quad 2. \frac{8(-4) - 3^2 \cdot 6}{-4 \cdot 2^3 + 12} \quad 3. \left(-\frac{5}{8} + \frac{1}{6}\right) - \left(-\frac{2}{3}\right)$$

Solve each equation or inequality.

$$4. 3(x + 2) - 4(2x + 3) = -3x + 2 \quad 5. \frac{1}{3}x + \frac{1}{4}(x + 8) = x + 7$$

$$6. 0.04x + 0.06(100 - x) = 5.88 \quad 7. |6x + 7| = 13$$

$$8. -5 - 3(x - 2) < 11 - 2(x + 2) \quad 9. -2 < 1 - 3x < 7$$

Solve each problem.

10. A piggy bank has 100 coins, all of which are nickels and quarters. The total value of the money is \$17.80. How many of each denomination are there in the bank?
11. How many liters of pure alcohol must be mixed with 40 L of 18% alcohol to obtain a 22% alcohol solution?
12. Graph the equation $4x - 3y = 12$.
13. Find the slope of the line passing through the points $(-4, 6)$ and $(2, -3)$. Then find the equation of the line and write it in the form $y = mx + b$.
14. If $f(x) = 3x - 7$, find $f(-10)$.

Solve each system.

$$15. \begin{cases} 3x - y = 23 \\ 2x + 3y = 8 \end{cases} \quad 16. \begin{cases} x + y + z = 1 \\ x - y - z = -3 \\ x + y - z = -1 \end{cases}$$

17. In 2017, sending five 2-oz letters and three 3-oz letters by first-class mail would have cost \$6.23. Sending three 2-oz letters and five 3-oz letters would have cost \$6.65. What was the rate for one 2-oz letter and one 3-oz letter? (Data from U.S. Postal Service.)



Perform the indicated operations.

18. $(3k^3 - 5k^2 + 8k - 2) - (4k^3 + 11k + 7) + (2k^2 - 5k)$

19. $(8x - 7)(x + 3)$

20. $\frac{6y^4 - 3y^3 + 5y^2 + 6y - 9}{2y + 1}$

Factor each polynomial completely.

21. $2p^2 - 5pq + 3q^2$

22. $3k^4 + k^2 - 4$

23. $x^3 + 512$

24. What is the domain of $f(x) = \frac{4}{x^2 - 9}$? Express it using set-builder notation.

Perform each operation, and express the answer in lowest terms.

25. $\frac{y^2 + y - 12}{y^3 + 9y^2 + 20y} \div \frac{y^2 - 9}{y^3 + 3y^2}$

26. $\frac{1}{x + y} + \frac{3}{x - y}$

Simplify.

27. $\frac{\frac{-6}{x - 2}}{\frac{8}{3x - 6}}$

28. $\frac{\frac{1}{a} - \frac{1}{b}}{\frac{a}{b} - \frac{b}{a}}$

29. $\frac{x^{-1}}{y - x^{-1}}$

Solve.

30. $2x^2 + 11x + 15 = 0$

31. $5x(x - 1) = 2(1 - x)$

32. $\frac{x + 1}{x - 3} = \frac{4}{x - 3} + 6$

33. $\sqrt{3x - 8} = x - 2$

34. Danielle can ride her bike 4 mph faster than her husband, Richard. If Danielle can ride 48 mi in the same time that Richard can ride 24 mi, what are their rates?

Simplify. Assume that all variables represent positive real numbers.

35. $27^{-2/3}$

36. $\sqrt[3]{16x^2y} \cdot \sqrt[3]{3x^3y}$

37. $\sqrt{50} + \sqrt{8}$

38. $\frac{1}{\sqrt{10} - \sqrt{8}}$

39. Find the distance between the points $(-4, 4)$ and $(-2, 9)$.

40. Express $\frac{6 - 2i}{1 - i}$ in standard form.

STUDY SKILLS REMINDER

Have you begun to prepare for your final exam? **Review Study Skill 10, Preparing for Your Math Final Exam.**

This page intentionally left blank

QUADRATIC EQUATIONS, INEQUALITIES, AND FUNCTIONS

Quadratic functions, one of the topics of this chapter, have graphs that are *parabolas*. Cross sections of telescopes, satellite dishes, and automobile headlights form parabolas, as do the cables that support suspension bridges.

8.1 The Square Root Property and Completing the Square

8.2 The Quadratic Formula

8.3 Equations That Lead to Quadratic Methods

SUMMARY EXERCISES Applying Methods for Solving Quadratic Equations

8.4 Formulas and Further Applications

8.5 Graphs of Quadratic Functions

8.6 More about Parabolas and Their Applications

8.7 Polynomial and Rational Inequalities

8.1 The Square Root Property and Completing the Square

OBJECTIVES

- 1 Review the zero-factor property.
- 2 Learn the square root property.
- 3 Solve quadratic equations of the form $(ax + b)^2 = c$ by extending the square root property.
- 4 Solve quadratic equations by completing the square.
- 5 Solve quadratic equations with nonreal complex solutions.

Recall that a *quadratic equation* is defined as follows.

Quadratic Equation

A **quadratic equation** (in x here) is an equation that can be written in the form

$$ax^2 + bx + c = 0,$$

where a , b , and c are real numbers and $a \neq 0$. The given form is called **standard form**.

Examples: $4x^2 + 4x - 5 = 0$, $3x^2 = 4x - 8$

Quadratic equations
(The first equation is
in standard form.)

A quadratic equation is a **second-degree equation**—that is, an equation with a squared variable term and no terms of greater degree.

OBJECTIVE 1 Review the zero-factor property.

We have used the zero-factor property to solve quadratic equations.

Zero-Factor Property

If two numbers have a product of 0, then at least one of the numbers must be 0. **That is, if $ab = 0$, then $a = 0$ or $b = 0$.**

EXAMPLE 1 Using the Zero-Factor Property

Solve $3x^2 - 5x - 28 = 0$.

This trinomial can
be factored.

$$3x^2 - 5x - 28 = 0$$

The quadratic equation
must be in standard form.

$$(3x + 7)(x - 4) = 0$$

Factor.

$$3x + 7 = 0 \quad \text{or} \quad x - 4 = 0$$

Zero-factor property

$$3x = -7 \quad \text{or} \quad x = 4$$

Solve each equation.

$$x = -\frac{7}{3}$$

To check, substitute each value in the original equation. The solution set is $\left\{-\frac{7}{3}, 4\right\}$.

NOW TRY 

VOCABULARY

- quadratic equation
- second-degree equation

NOW TRY EXERCISE 1

Solve $2x^2 + 5x - 12 = 0$.

OBJECTIVE 2 Learn the square root property.

Not every quadratic equation can be solved using the zero-factor property. Other methods of solving quadratic equations are based on the following property.

Square Root Property

If x and k are complex numbers and $x^2 = k$, then

$$x = \sqrt{k} \quad \text{or} \quad x = -\sqrt{k}.$$

NOW TRY ANSWER

1. $\left\{-4, \frac{3}{2}\right\}$

These steps justify the square root property.

$$\begin{aligned}
 x^2 &= k \\
 x^2 - k &= 0 && \text{Subtract } k. \\
 (x - \sqrt{k})(x + \sqrt{k}) &= 0 && \text{Factor.} \\
 x - \sqrt{k} = 0 & \quad \text{or} \quad x + \sqrt{k} = 0 && \text{Zero-factor property} \\
 x = \sqrt{k} & \quad \text{or} \quad x = -\sqrt{k} && \text{Solve each equation.}
 \end{aligned}$$

CAUTION If $k \neq 0$, then using the square root property always produces two square roots, one positive and one negative.

NOW TRY EXERCISE 2

Solve each equation.

- (a) $t^2 = 10$
 (b) $2x^2 - 90 = 0$

EXAMPLE 2 Using the Square Root Property

Solve each equation.

(a) $x^2 = 5$

By the square root property, if $x^2 = 5$, then

$$x = \sqrt{5} \quad \text{or} \quad x = -\sqrt{5}.$$

Don't forget the negative solution.

These solutions are irrational numbers. The solution set is $\{\sqrt{5}, -\sqrt{5}\}$.

(b) $4x^2 - 48 = 0$

$$4x^2 = 48 \quad \text{Add 48.}$$

$$x^2 = 12 \quad \text{Divide by 4.}$$

Don't stop here. Simplify the radicals.

$$x = \sqrt{12} \quad \text{or} \quad x = -\sqrt{12} \quad \text{Square root property}$$

$$x = 2\sqrt{3} \quad \text{or} \quad x = -2\sqrt{3} \quad \sqrt{12} = \sqrt{4} \cdot \sqrt{3} = 2\sqrt{3}$$

CHECK $4x^2 - 48 = 0$ Original equation

$4(2\sqrt{3})^2 - 48 \stackrel{?}{=} 0$ $4(12) - 48 \stackrel{?}{=} 0$ $48 - 48 \stackrel{?}{=} 0$ $0 = 0 \quad \checkmark \quad \text{True}$	Let $x = 2\sqrt{3}$.		$4(-2\sqrt{3})^2 - 48 \stackrel{?}{=} 0$ $4(12) - 48 \stackrel{?}{=} 0$ $48 - 48 \stackrel{?}{=} 0$ $0 = 0 \quad \checkmark \quad \text{True}$	Let $x = -2\sqrt{3}$.
-----------------------------------------------------------------------------------------------------------------------------------------------	-----------------------	--	------------------------------------------------------------------------------------------------------------------------------------------------	------------------------

$(2\sqrt{3})^2 = 2^2 \cdot (\sqrt{3})^2$

The solution set is $\{2\sqrt{3}, -2\sqrt{3}\}$.

(c) $3x^2 + 5 = 11$

$$3x^2 = 6 \quad \text{Subtract 5.}$$

$$x^2 = 2 \quad \text{Divide by 3.}$$

$$x = \sqrt{2} \quad \text{or} \quad x = -\sqrt{2} \quad \text{Square root property}$$

The solution set is $\{\sqrt{2}, -\sqrt{2}\}$.

NOW TRY

NOW TRY ANSWERS

2. (a) $\{\sqrt{10}, -\sqrt{10}\}$
 (b) $\{3\sqrt{5}, -3\sqrt{5}\}$

NOTE Using the symbol \pm (read “positive or negative,” or “plus or minus”), the solutions in **Example 2** could be written $\pm\sqrt{5}$, $\pm 2\sqrt{3}$, and $\pm\sqrt{2}$.

NOW TRY
EXERCISE 3

Tim is dropping roofing nails from the top of a roof 25 ft high into a large bucket on the ground. Use the formula in **Example 3** to determine how long it will take a nail dropped from 25 ft to hit the bottom of the bucket.


EXAMPLE 3 Using the Square Root Property in an Application

Galileo Galilei developed a formula for freely falling objects described by

$$d = 16t^2,$$

where d is the distance in feet that an object falls (disregarding air resistance) in t seconds, regardless of weight.

If the Leaning Tower of Pisa is about 180 ft tall, use Galileo's formula to determine how long it would take an object dropped from the top of the tower to fall to the ground. (Data from www.brittanica.com)



Galileo Galilei (1564–1642)

$$d = 16t^2 \quad \text{Galileo's formula}$$

$$180 = 16t^2 \quad \text{Let } d = 180.$$

$$11.25 = t^2 \quad \text{Divide by 16.}$$

$$t = \sqrt{11.25} \quad \text{or} \quad t = -\sqrt{11.25} \quad \text{Square root property}$$

Time cannot be negative, so we discard $-\sqrt{11.25}$. Using a calculator,

$$\sqrt{11.25} \approx 3.4 \quad \text{so} \quad t \approx 3.4. \quad \text{Round to the nearest tenth.}$$

The object would fall to the ground in 3.4 sec.

NOW TRY

CAUTION When solving applications, a solution of a quadratic equation may not satisfy the physical requirements, as in **Example 3**. Reject such solutions as answers.

OBJECTIVE 3 Solve quadratic equations of the form $(ax + b)^2 = c$ by extending the square root property.

EXAMPLE 4 Extending the Square Root Property

Solve $(x - 5)^2 = 36$.

$$\begin{array}{ccc} x^2 & = & k \\ \downarrow & & \downarrow \\ (x - 5)^2 & = & 36 \end{array} \quad \begin{array}{l} \text{Substitute } (x - 5)^2 \text{ for } x^2 \text{ and } 36 \text{ for } k \\ \text{in the square root property.} \end{array}$$

$$x - 5 = \sqrt{36} \quad \text{or} \quad x - 5 = -\sqrt{36} \quad \text{Square root property}$$

$$x - 5 = 6 \quad \text{or} \quad x - 5 = -6 \quad \text{Take square roots.}$$

$$x = 11 \quad \text{or} \quad x = -1 \quad \text{Add 5.}$$

CHECK $(x - 5)^2 = 36$ Original equation

$$(11 - 5)^2 \stackrel{?}{=} 36 \quad \text{Let } x = 11. \quad \left| \quad (-1 - 5)^2 \stackrel{?}{=} 36 \quad \text{Let } x = -1.$$

$$6^2 \stackrel{?}{=} 36 \quad \left| \quad (-6)^2 \stackrel{?}{=} 36$$

$$36 = 36 \quad \checkmark \quad \text{True} \quad \left| \quad 36 = 36 \quad \checkmark \quad \text{True}$$

Both values satisfy the original equation. The solution set is $\{-1, 11\}$.

NOW TRY

NOW TRY
EXERCISE 4

Solve $(x + 3)^2 = 25$.

NOW TRY ANSWERS

3. 1.25 sec

4. $\{-8, 2\}$


NOW TRY EXERCISE 5
Solve $(5x - 4)^2 = 27$.**EXAMPLE 5** Extending the Square Root PropertySolve $(2x - 3)^2 = 18$.

$$2x - 3 = \sqrt{18} \quad \text{or} \quad 2x - 3 = -\sqrt{18} \quad \text{Square root property}$$

$$2x = 3 + \sqrt{18} \quad \text{or} \quad 2x = 3 - \sqrt{18} \quad \text{Add 3.}$$

$$x = \frac{3 + \sqrt{18}}{2} \quad \text{or} \quad x = \frac{3 - \sqrt{18}}{2} \quad \text{Divide by 2.}$$

$$x = \frac{3 + 3\sqrt{2}}{2} \quad \text{or} \quad x = \frac{3 - 3\sqrt{2}}{2} \quad \sqrt{18} = \sqrt{9 \cdot 2} = 3\sqrt{2}$$

We show the check for the first value. The check for the other value is similar.

CHECK $(2x - 3)^2 = 18$ Original equation

$$\left[2\left(\frac{3 + 3\sqrt{2}}{2}\right) - 3 \right]^2 \stackrel{?}{=} 18 \quad \text{Let } x = \frac{3 + 3\sqrt{2}}{2}.$$

$$(3 + 3\sqrt{2} - 3)^2 \stackrel{?}{=} 18 \quad \text{Multiply.}$$

$$(3\sqrt{2})^2 \stackrel{?}{=} 18 \quad \text{Simplify.}$$

$$18 = 18 \quad \checkmark \quad \text{True}$$

The \pm symbol denotes two solutions.

The solution set is $\left\{ \frac{3 + 3\sqrt{2}}{2}, \frac{3 - 3\sqrt{2}}{2} \right\}$, abbreviated $\left\{ \frac{3 \pm 3\sqrt{2}}{2} \right\}$.

NOW TRY **OBJECTIVE 4** Solve quadratic equations by completing the square.

We can use the square root property to solve *any* quadratic equation by writing it in the form

$$\text{Square of a binomial} \rightarrow (x + k)^2 = n. \leftarrow \text{Constant}$$

That is, we must write the left side of the equation as a perfect square trinomial that can be factored as $(x + k)^2$, the square of a binomial, and the right side must be a constant. This process is called **completing the square**.

Recall that the perfect square trinomial

$$x^2 + 10x + 25 \quad \text{can be factored as} \quad (x + 5)^2.$$

In the trinomial, the coefficient of x (the first-degree term) is 10 and the constant term is 25. If we take half of 10 and square it, we obtain the constant term, 25.

$$\begin{array}{ccc} \text{Coefficient of } x & & \text{Constant} \\ \downarrow & & \downarrow \\ \left[\frac{1}{2}(10) \right]^2 = 5^2 = 25 \end{array}$$

Similarly, in $x^2 + 12x + 36$, $\left[\frac{1}{2}(12) \right]^2 = 6^2 = 36$,

and in $m^2 - 6m + 9$, $\left[\frac{1}{2}(-6) \right]^2 = (-3)^2 = 9$.

NOW TRY ANSWER

5. $\left\{ \frac{4 \pm 3\sqrt{3}}{5} \right\}$

This relationship is true in general and is the idea behind completing the square.


NOW TRY EXERCISE 6
Solve $x^2 + 6x - 2 = 0$.
EXAMPLE 6 Solving a Quadratic Equation by Completing the Square ($a = 1$)
Solve $x^2 + 8x + 10 = 0$.

The trinomial on the left is nonfactorable, so this quadratic equation cannot be solved using the zero-factor property. It is not in the correct form to solve using the square root property. We can solve it by completing the square.

$$\begin{array}{ll}
 x^2 + 8x + 10 = 0 & \text{Original equation} \\
 \text{Only terms with variables remain of the left side.} \rightarrow x^2 + 8x = -10 & \text{Subtract 10.} \\
 x^2 + 8x + \underline{\quad ? \quad} = -10 & \text{We must add a constant.} \\
 \underbrace{\hspace{1.5cm}} & \text{Needs to be a perfect square trinomial}
 \end{array}$$

Take half the coefficient of the first-degree term, $8x$, and square the result.

$$\left[\frac{1}{2}(8)\right]^2 = 4^2 = 16 \leftarrow \text{Desired constant}$$

We add this constant, 16, to *each* side of the equation and continue solving as shown.

$$\begin{array}{ll}
 x^2 + 8x + 16 = -10 + 16 & \text{Add 16 to each side.} \\
 \text{This is a key step.} \rightarrow (x + 4)^2 = 6 & \text{Factor on the left.} \\
 & \text{Add on the right.} \\
 x + 4 = \sqrt{6} \quad \text{or} \quad x + 4 = -\sqrt{6} & \text{Square root property} \\
 x = -4 + \sqrt{6} \quad \text{or} \quad x = -4 - \sqrt{6} & \text{Add } -4.
 \end{array}$$

CHECK

$$\begin{array}{ll}
 x^2 + 8x + 10 = 0 & \text{Original equation} \\
 \text{Remember the middle term when squaring } (-4 + \sqrt{6})^2 + 8(-4 + \sqrt{6}) + 10 \stackrel{?}{=} 0 & \text{Let } x = -4 + \sqrt{6}. \\
 -4 + \sqrt{6}. \rightarrow 16 - 8\sqrt{6} + 6 - 32 + 8\sqrt{6} + 10 \stackrel{?}{=} 0 & \text{Multiply.} \\
 0 = 0 \quad \checkmark & \text{True}
 \end{array}$$

The check of $-4 - \sqrt{6}$ is similar. The solution set is

$$\{-4 + \sqrt{6}, -4 - \sqrt{6}\}, \quad \text{or} \quad \{-4 \pm \sqrt{6}\}.$$

NOW TRY 
STUDY SKILLS REMINDER

Make study cards to help you learn and remember the material in this chapter.

Review Study Skill 5, Using Study Cards.

Completing the Square to Solve $ax^2 + bx + c = 0$ (Where $a \neq 0$)
Step 1 Be sure the second-degree term has coefficient 1.

- If the coefficient of the second-degree term is 1, go to Step 2.
- If it is not 1, but some other nonzero number a , divide each side of the equation by a .

Step 2 Write the equation in correct form. Make sure that all variable terms are on one side of the equality symbol and the constant term is on the other side.

Step 3 Complete the square.

- Take half the coefficient of the first-degree term, and square it.
- Add the square to each side of the equation.
- Factor the variable side, which should be a perfect square trinomial, as the square of a binomial. Combine terms on the other side.

Step 4 Solve the equation using the square root property.

NOW TRY ANSWER6. $\{-3 \pm \sqrt{11}\}$


NOW TRY EXERCISE 7
Solve $x^2 + x - 3 = 0$.**EXAMPLE 7** Solving a Quadratic Equation by Completing the Square ($a = 1$)Solve $x^2 + 5x - 1 = 0$.

The coefficient of the second-degree term is 1, so we begin with Step 2.

Step 2 $x^2 + 5x = 1$ Add 1 to each side.

Step 3 $\left[\frac{1}{2}(5)\right]^2 = \left(\frac{5}{2}\right)^2 = \frac{25}{4}$ Take half the coefficient of the first-degree term and square the result.

$$x^2 + 5x + \frac{25}{4} = 1 + \frac{25}{4}$$

Add the square to each side of the equation.

$$\left(x + \frac{5}{2}\right)^2 = \frac{29}{4}$$

Factor on the left. Add on the right.

Step 4 $x + \frac{5}{2} = \sqrt{\frac{29}{4}}$ or $x + \frac{5}{2} = -\sqrt{\frac{29}{4}}$ Square root property

$$x + \frac{5}{2} = \frac{\sqrt{29}}{2}$$
 or $x + \frac{5}{2} = -\frac{\sqrt{29}}{2}$ $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$

$$x = -\frac{5}{2} + \frac{\sqrt{29}}{2}$$
 or $x = -\frac{5}{2} - \frac{\sqrt{29}}{2}$ Add $-\frac{5}{2}$.

$$x = \frac{-5 + \sqrt{29}}{2}$$
 or $x = \frac{-5 - \sqrt{29}}{2}$ $\frac{a}{c} \pm \frac{b}{c} = \frac{a \pm b}{c}$

The solution set is $\left\{\frac{-5 \pm \sqrt{29}}{2}\right\}$.**NOW TRY** **EXAMPLE 8** Solving a Quadratic Equation by Completing the Square ($a \neq 1$)Solve $2x^2 - 4x - 5 = 0$.

Divide each side by 2 to obtain 1 as the coefficient of the second-degree term.

$$x^2 - 2x - \frac{5}{2} = 0$$

Divide by 2. (Step 1)

$$x^2 - 2x = \frac{5}{2}$$

Add $\frac{5}{2}$. (Step 2)

$$\left[\frac{1}{2}(-2)\right]^2 = (-1)^2 = 1$$

Complete the square. (Step 3)

$$x^2 - 2x + 1 = \frac{5}{2} + 1$$

Add 1 to each side.

$$(x - 1)^2 = \frac{7}{2}$$

Factor on the left. Add on the right.

$$x - 1 = \sqrt{\frac{7}{2}}$$
 or $x - 1 = -\sqrt{\frac{7}{2}}$

Square root property (Step 4)

$$x = 1 + \sqrt{\frac{7}{2}}$$
 or $x = 1 - \sqrt{\frac{7}{2}}$

Add 1.

$$x = 1 + \frac{\sqrt{14}}{2}$$
 or $x = 1 - \frac{\sqrt{14}}{2}$

$\sqrt{\frac{7}{2}} = \frac{\sqrt{7}}{\sqrt{2}} = \frac{\sqrt{7}}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{14}}{2}$

NOW TRY ANSWER

7. $\left\{\frac{-1 \pm \sqrt{13}}{2}\right\}$


NOW TRY EXERCISE 8
Solve $3x^2 + 12x - 5 = 0$.

Add the two terms in each solution as follows.

$$1 + \frac{\sqrt{14}}{2} = \frac{2}{2} + \frac{\sqrt{14}}{2} = \frac{2 + \sqrt{14}}{2}$$

$$1 - \frac{\sqrt{14}}{2} = \frac{2}{2} - \frac{\sqrt{14}}{2} = \frac{2 - \sqrt{14}}{2}$$

$1 = \frac{2}{2}$

The solution set is $\left\{\frac{2 \pm \sqrt{14}}{2}\right\}$.
NOW TRY 
OBJECTIVE 5 Solve quadratic equations with nonreal complex solutions.
If $k < 0$ in the equation $x^2 = k$, then there will be two nonreal complex solutions.

NOW TRY EXERCISE 9

Solve each equation.

- (a) $t^2 = -24$
 (b) $(x + 4)^2 = -36$
 (c) $x^2 + 8x + 21 = 0$

EXAMPLE 9 Solving Quadratic Equations (Nonreal Complex Solutions)

Solve each equation.

(a) $x^2 = -15$

$$x = \sqrt{-15} \quad \text{or} \quad x = -\sqrt{-15} \quad \text{Square root property}$$

$$x = i\sqrt{15} \quad \text{or} \quad x = -i\sqrt{15} \quad \sqrt{-a} = i\sqrt{a}$$

The solution set is $\{i\sqrt{15}, -i\sqrt{15}\}$, or $\{\pm i\sqrt{15}\}$.

(b) $(x + 2)^2 = -16$

$$x + 2 = \sqrt{-16} \quad \text{or} \quad x + 2 = -\sqrt{-16} \quad \text{Square root property}$$

$$x + 2 = 4i \quad \text{or} \quad x + 2 = -4i \quad \sqrt{-16} = 4i$$

$$x = -2 + 4i \quad \text{or} \quad x = -2 - 4i \quad \text{Add } -2.$$

The solution set is $\{-2 + 4i, -2 - 4i\}$, or $\{-2 \pm 4i\}$.

(c) $x^2 + 2x + 7 = 0$

$$x^2 + 2x = -7 \quad \text{Subtract 7.}$$

$$x^2 + 2x + 1 = -7 + 1 \quad \left[\frac{1}{2}(2)\right]^2 = 1; \text{ Add 1 to each side.}$$

$$(x + 1)^2 = -6 \quad \text{Factor on the left. Add on the right.}$$

$$x + 1 = \sqrt{-6} \quad \text{or} \quad x + 1 = -\sqrt{-6} \quad \text{Square root property}$$

$$x + 1 = i\sqrt{6} \quad \text{or} \quad x + 1 = -i\sqrt{6} \quad \sqrt{-6} = i\sqrt{6}$$

$$x = -1 + i\sqrt{6} \quad \text{or} \quad x = -1 - i\sqrt{6} \quad \text{Add } -1.$$

The solution set is $\{-1 + i\sqrt{6}, -1 - i\sqrt{6}\}$, or $\{-1 \pm i\sqrt{6}\}$.
NOW TRY 
NOW TRY ANSWERS

8. $\left\{\frac{-6 \pm \sqrt{51}}{3}\right\}$
 9. (a) $\{\pm 2i\sqrt{6}\}$
 (b) $\{-4 \pm 6i\}$
 (c) $\{-4 \pm i\sqrt{5}\}$

8.1 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Fill in each blank with the correct response.

- An equation in the form $ax^2 + bx + c = 0$, where a , b , and c are real numbers and $a \neq 0$, is a(n) _____ equation, also called a(n) _____ degree equation. The greatest degree of the variable is _____.
- Which of the following are quadratic equations?
 - $x + y = 0$
 - $x^2 - 4x + 4 = 0$
 - $x^3 + x^2 + 8 = 0$
 - $2t^2 - 7t = -4$
- Concept Check** A student incorrectly solved the following equation as shown.

$$x^2 - x - 2 = 4$$

$$(x - 2)(x + 1) = 4 \quad \text{Factor.}$$

$$x - 2 = 4 \quad \text{or} \quad x + 1 = 4 \quad \text{Zero-factor property}$$

$$x = 6 \quad \text{or} \quad x = 3 \quad \text{Solve each equation.}$$

WHAT WENT WRONG? Solve correctly and give the solution set.

- Concept Check** A student was asked to solve the quadratic equation

$$x^2 = 16$$

and did not get full credit for the solution set $\{4\}$. **WHAT WENT WRONG?** Give the correct solution set.

Use the zero-factor property to solve each equation. (Hint: In Exercises 11–14, write the equation in standard form first.) See Example 1.

5. $x^2 + 3x + 2 = 0$

6. $x^2 + 8x + 15 = 0$

7. $3x^2 + 8x - 3 = 0$

8. $2x^2 + x - 6 = 0$

9. $5x^2 - 7x - 6 = 0$

10. $2x^2 - 5x - 3 = 0$

11. $x^2 - 6x = -8$

12. $x^2 - 10x = -24$

13. $2x^2 = 9x - 4$

14. $5x^2 = 11x - 2$

15. $8x^2 + 14x + 3 = 0$

16. $12x^2 + 19x + 5 = 0$

Use the square root property to solve each equation. See Examples 2, 4, and 5.

17. $x^2 = 81$

18. $x^2 = 225$

19. $x^2 = 17$

20. $x^2 = 19$

21. $x^2 = 32$

22. $x^2 = 54$

23. $x^2 - 20 = 0$

24. $p^2 - 50 = 0$

25. $3x^2 - 72 = 0$

26. $5z^2 - 200 = 0$

27. $2x^2 + 7 = 61$

28. $3x^2 + 8 = 80$

29. $3x^2 - 10 = 86$

30. $4x^2 - 7 = 65$

31. $(x + 2)^2 = 25$

32. $(t + 8)^2 = 9$

33. $(x - 6)^2 = 49$

34. $(x - 4)^2 = 64$

35. $(x - 4)^2 = 3$

36. $(x + 3)^2 = 11$

37. $(t + 5)^2 = 48$

38. $(m - 6)^2 = 27$

39. $(3x - 1)^2 = 7$

40. $(2x - 5)^2 = 10$

41. $(4p + 1)^2 = 24$

42. $(5t + 2)^2 = 12$

43. $(2 - 5t)^2 = 12$

44. $(1 - 4p)^2 = 24$

Use Galileo's formula to solve each problem. Round answers to the nearest tenth. See Example 3.

45. The sculpture of American presidents at Mount Rushmore National Memorial is 500 ft above the valley floor. How long would it take a rock dropped from the top of the sculpture to fall to the ground? (Data from www.travelsd.com)
46. The Gateway Arch in St. Louis, Missouri, is 630 ft tall. How long would it take an object dropped from the top of the arch to fall to the ground? (Data from www.gatewayarch.com)



47. **Concept Check** Which one of the two equations

$$(2x + 1)^2 = 5 \quad \text{and} \quad x^2 + 4x = 12$$

is more suitable for solving by the square root property? By completing the square?

48. **Concept Check** According to the procedure described in this section, what would be the first step in solving $2x^2 + 8x = 9$ by completing the square?

Concept Check Find the constant that must be added to make each expression a perfect square trinomial. Then factor the trinomial.

- | | | |
|------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|
| 49. $x^2 + 6x + \underline{\hspace{1cm}}$
It factors as $\underline{\hspace{1cm}}$. | 50. $x^2 + 14x + \underline{\hspace{1cm}}$
It factors as $\underline{\hspace{1cm}}$. | 51. $p^2 - 12p + \underline{\hspace{1cm}}$
It factors as $\underline{\hspace{1cm}}$. |
| 52. $x^2 - 20x + \underline{\hspace{1cm}}$
It factors as $\underline{\hspace{1cm}}$. | 53. $q^2 + 9q + \underline{\hspace{1cm}}$
It factors as $\underline{\hspace{1cm}}$. | 54. $t^2 + 3t + \underline{\hspace{1cm}}$
It factors as $\underline{\hspace{1cm}}$. |

Solve each equation by completing the square. See Examples 6–8.

- | | | |
|--------------------------------------------------------------------|--------------------------------------------------------------------|-------------------------------|
| 55. $x^2 - 2x - 24 = 0$ | 56. $m^2 - 4m - 32 = 0$ | 57. $x^2 + 4x - 2 = 0$ |
| 58. $t^2 + 2t - 1 = 0$ | 59. $x^2 + 10x + 18 = 0$ | 60. $x^2 + 8x + 11 = 0$ |
| 61. $3w^2 - w = 24$ | 62. $4z^2 - z = 39$ | 63. $x^2 + 7x - 1 = 0$ |
| 64. $x^2 + 13x - 3 = 0$ | 65. $2k^2 + 5k - 2 = 0$ | 66. $3r^2 + 2r - 2 = 0$ |
| 67. $5x^2 - 10x + 2 = 0$ | 68. $2x^2 - 16x + 25 = 0$ | 69. $9x^2 - 24x = -13$ |
| 70. $25n^2 - 20n = 1$ | 71. $z^2 - \frac{4}{3}z = -\frac{1}{9}$ | 72. $p^2 - \frac{8}{3}p = -1$ |
| 73. $0.1x^2 - 0.2x - 0.1 = 0$
(Hint: First clear the decimals.) | 74. $0.1p^2 - 0.4p + 0.1 = 0$
(Hint: First clear the decimals.) | |

Solve each equation. (All solutions for these equations are nonreal complex numbers.) See Example 9.

75. $x^2 = -100$

76. $x^2 = -64$

77. $x^2 = -12$

78. $x^2 = -18$

79. $(x + 3)^2 = -4$

80. $(x - 5)^2 = -36$

81. $(r - 5)^2 = -3$

82. $(t + 6)^2 = -5$

83. $(6k - 1)^2 = -8$

84. $(4m - 7)^2 = -27$

85. $m^2 + 4m + 13 = 0$

86. $t^2 + 6t + 10 = 0$

87. $m^2 + 6m + 12 = 0$

88. $x^2 + 10x + 27 = 0$

89. $3r^2 + 4r + 4 = 0$

90. $4x^2 + 5x + 5 = 0$

91. $-k^2 - 5k - 10 = 0$

92. $-x^2 - 3x - 8 = 0$

Extending Skills Solve for x . Assume that a and b represent positive real numbers.

93. $x^2 - b = 0$

94. $x^2 = 4b$

95. $4x^2 = b^2 + 16$


96. $9x^2 - 25a = 0$

97. $(5x - 2b)^2 = 3a$

98. $x^2 - a^2 - 36 = 0$

RELATING CONCEPTS For Individual or Group Work (Exercises 99–104)

The Greeks had a method of completing the square geometrically in which they literally changed a figure into a square. For example, to complete the square for $x^2 + 6x$, we begin with a square of side x , as in the figure on the top. We add three rectangles of width 1 to the right side and the bottom to get a region with area $x^2 + 6x$. To fill in the corner (complete the square), we must add nine 1-by-1 squares as shown.



Work Exercises 99–104 in order.

99. What is the area of the original square?


100. What is the area of each strip?

101. What is the total area of the six strips?

102. What is the area of each small square in the corner of the second figure?

103. What is the total area of the small squares?

104. What is the area of the new “complete” square?



8.2

The Quadratic Formula

OBJECTIVES

- 1 Derive the quadratic formula.
- 2 Solve quadratic equations using the quadratic formula.
- 3 Use the discriminant to determine number and type of solutions.

In this section, we complete the square to solve the general quadratic equation

$$ax^2 + bx + c = 0,$$

where a , b , and c are complex numbers and $a \neq 0$. The solution of this general equation gives a formula for finding the solution of any specific quadratic equation.

OBJECTIVE 1 Derive the quadratic formula.

We solve $ax^2 + bx + c = 0$ by completing the square (where $a > 0$) as follows.

$$ax^2 + bx + c = 0$$

$$x^2 + \frac{b}{a}x + \frac{c}{a} = 0 \quad \text{Divide by } a. \text{ (Step 1)}$$

$$x^2 + \frac{b}{a}x = -\frac{c}{a} \quad \text{Subtract } \frac{c}{a}. \text{ (Step 2)}$$

$$\left[\frac{1}{2}\left(\frac{b}{a}\right)\right]^2 = \left(\frac{b}{2a}\right)^2 = \frac{b^2}{4a^2} \quad \text{Complete the square. (Step 3)}$$

$$x^2 + \frac{b}{a}x + \frac{b^2}{4a^2} = -\frac{c}{a} + \frac{b^2}{4a^2} \quad \text{Add } \frac{b^2}{4a^2} \text{ to each side.}$$

$$\left(x + \frac{b}{2a}\right)^2 = \frac{b^2}{4a^2} + \frac{-c}{a} \quad \begin{array}{l} \text{Factor on the left.} \\ \text{Rearrange the terms on the right.} \end{array}$$

$$\left(x + \frac{b}{2a}\right)^2 = \frac{b^2}{4a^2} + \frac{-4ac}{4a^2} \quad \text{Write with a common denominator.}$$

$$\left(x + \frac{b}{2a}\right)^2 = \frac{b^2 - 4ac}{4a^2} \quad \text{Add fractions.}$$

$$x + \frac{b}{2a} = \sqrt{\frac{b^2 - 4ac}{4a^2}} \quad \text{or} \quad x + \frac{b}{2a} = -\sqrt{\frac{b^2 - 4ac}{4a^2}} \quad \begin{array}{l} \text{Square root property} \\ \text{(Step 4)} \end{array}$$

We can simplify $\sqrt{\frac{b^2 - 4ac}{4a^2}}$ as $\frac{\sqrt{b^2 - 4ac}}{\sqrt{4a^2}}$, or $\frac{\sqrt{b^2 - 4ac}}{2a}$.

The right side of each equation can be expressed as follows.

$$x + \frac{b}{2a} = \frac{\sqrt{b^2 - 4ac}}{2a} \quad \text{or} \quad x + \frac{b}{2a} = \frac{-\sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-b}{2a} + \frac{\sqrt{b^2 - 4ac}}{2a} \quad \text{or} \quad x = \frac{-b}{2a} - \frac{\sqrt{b^2 - 4ac}}{2a}$$

If $a < 0$, the same two solutions are obtained.

$$x = \frac{-b + \sqrt{b^2 - 4ac}}{2a} \quad \text{or} \quad x = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

This result is the **quadratic formula**, which is abbreviated as follows.

Quadratic Formula

The solutions of the equation $ax^2 + bx + c = 0$ (where $a \neq 0$) are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

! CAUTION In the quadratic formula, the square root is added to or subtracted from the value of $-b$ before dividing by $2a$.


**NOW TRY
EXERCISE 1**
Solve $2x^2 + 3x - 20 = 0$.**OBJECTIVE 2** Solve quadratic equations using the quadratic formula.**EXAMPLE 1** Using the Quadratic Formula (Two Rational Solutions)Solve $6x^2 - 5x - 4 = 0$.

This equation is in standard form, so we identify the values of a , b , and c . Here a , the coefficient of the second-degree term, is 6 , and b , the coefficient of the first-degree term, is -5 . The constant c is -4 . Now substitute into the quadratic formula.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Quadratic formula

$$x = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(6)(-4)}}{2(6)}$$

 $a = 6, b = -5, c = -4$

Use parentheses and substitute carefully to avoid errors.

$$x = \frac{5 \pm \sqrt{25 + 96}}{12}$$

$$x = \frac{5 \pm \sqrt{121}}{12}$$

Add under the radical.

$$x = \frac{5 \pm 11}{12}$$

Take the square root.

There are two values represented, one from the $+$ sign and one from the $-$ sign.

$$x = \frac{5 + 11}{12} = \frac{16}{12} = \frac{4}{3} \quad \text{or} \quad x = \frac{5 - 11}{12} = \frac{-6}{12} = -\frac{1}{2}$$

Check each value in the original equation. The solution set is $\left\{-\frac{1}{2}, \frac{4}{3}\right\}$.

NOW TRY 

NOTE We could have factored the trinomial and then used the zero-factor property to solve the equation in **Example 1**.

$$6x^2 - 5x - 4 = 0$$

See Example 1.

$$(3x - 4)(2x + 1) = 0$$

Factor.

$$3x - 4 = 0 \quad \text{or} \quad 2x + 1 = 0$$

Zero-factor property

$$3x = 4 \quad \text{or} \quad 2x = -1$$

Solve each equation.

$$x = \frac{4}{3} \quad \text{or} \quad x = -\frac{1}{2}$$

Same solutions as in Example 1

When solving a quadratic equation, it is a good idea to try to factor the quadratic expression first. If it can be factored, then apply the zero-factor property. If it cannot be factored or if factoring is difficult, then use the quadratic formula.

EXAMPLE 2 Using the Quadratic Formula (One Rational Solution)Solve $9x^2 + 12x + 4 = 0$.

The trinomial on the left side of the equality symbol can be factored, so this equation could be solved using the zero-factor property. However, if we did not recognize this and solved using the quadratic formula, our work might look similar to that shown on the next page.

NOW TRY ANSWER1. $\left\{-4, \frac{5}{2}\right\}$


**NOW TRY
EXERCISE 2**
Solve $4x^2 - 20x + 25 = 0$.The equation $9x^2 + 12x + 4 = 0$ is in standard form, so $a = 9$, $b = 12$, and $c = 4$.


$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{Quadratic formula}$$

$$x = \frac{-12 \pm \sqrt{12^2 - 4(9)(4)}}{2(9)} \quad a = 9, b = 12, c = 4$$

$$x = \frac{-12 \pm \sqrt{144 - 144}}{18} \quad \text{Simplify.}$$

$$x = \frac{-12 \pm 0}{18} \quad \sqrt{0} = 0$$

$$x = -\frac{2}{3} \quad \text{Write } \frac{-12}{18} \text{ in lowest terms.}$$

In this case, $b^2 - 4ac = 0$ because $9x^2 + 12x + 4$ is a perfect square trinomial. There is one *distinct* solution, $-\frac{2}{3}$. The solution set is $\left\{-\frac{2}{3}\right\}$. **NOW TRY** 

NOTE Solve the equation in **Example 2** by factoring $9x^2 + 12x + 4$ and using the zero-factor property to confirm that the same solution, $-\frac{2}{3}$, results.


**NOW TRY
EXERCISE 3**
Solve $3x^2 + 1 = -5x$.**EXAMPLE 3** Using the Quadratic Formula (Two Irrational Solutions)Solve $4x^2 = 8x - 1$.Write the equation in standard form as $4x^2 - 8x + 1 = 0$. 

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{Quadratic formula}$$

$$x = \frac{-(-8) \pm \sqrt{(-8)^2 - 4(4)(1)}}{2(4)} \quad a = 4, b = -8, c = 1$$

$$x = \frac{8 \pm \sqrt{64 - 16}}{8} \quad \text{Simplify in the numerator and denominator.}$$

$$x = \frac{8 \pm \sqrt{48}}{8} \quad \text{Subtract under the radical.}$$

$$x = \frac{8 \pm 4\sqrt{3}}{8} \quad \sqrt{48} = \sqrt{16} \cdot \sqrt{3} = 4\sqrt{3}$$

$$x = \frac{4(2 \pm \sqrt{3})}{4(2)} \quad \text{Factor.}$$

$$x = \frac{2 \pm \sqrt{3}}{2} \quad \text{Divide out the common factor 4 to write in lowest terms.}$$

NOW TRY ANSWERS

2. $\left\{\frac{3}{2}\right\}$

3. $\left\{\frac{-5 \pm \sqrt{13}}{6}\right\}$

The solution set is $\left\{\frac{2 \pm \sqrt{3}}{2}\right\}$.**NOW TRY** 

! CAUTION

1. Before solving, every quadratic equation must be expressed in standard form $ax^2 + bx + c = 0$, whether we use the zero-factor property or the quadratic formula.
2. When writing solutions in lowest terms, be sure to factor first. Then divide out the common factor. See the last two steps in Example 3.

NOW TRY EXERCISE 4

Solve $(x + 5)(x - 1) = -18$.

EXAMPLE 4 Using the Quadratic Formula (Two Nonreal Complex Solutions)

Solve $(9x + 3)(x - 1) = -8$.

This is a quadratic equation—when the first terms $9x$ and x are multiplied, we obtain a second-degree term, $9x^2$. We must write the equation in standard form.

$$(9x + 3)(x - 1) = -8$$

$$9x^2 - 6x - 3 = -8 \quad \text{Multiply.}$$

$$\text{Standard form} \rightarrow 9x^2 - 6x + 5 = 0 \quad \text{Add 8.}$$

From the equation $9x^2 - 6x + 5 = 0$, we identify $a = 9$, $b = -6$, and $c = 5$.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{Quadratic formula}$$

$$x = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(9)(5)}}{2(9)} \quad \text{Substitute.}$$

$$x = \frac{6 \pm \sqrt{-144}}{18} \quad \text{Simplify.}$$

$$x = \frac{6 \pm 12i}{18} \quad \sqrt{-144} = 12i$$

$$x = \frac{6(1 \pm 2i)}{6(3)} \quad \text{Factor first. Then divide out the common factor.} \quad \text{Factor.}$$

$$x = \frac{1 \pm 2i}{3} \quad \text{Divide out the common factor 6 to write in lowest terms.}$$

$$x = \frac{1}{3} \pm \frac{2}{3}i \quad \text{Standard form } a + bi \text{ for a complex number}$$

The solution set is $\left\{\frac{1}{3} \pm \frac{2}{3}i\right\}$.

NOW TRY **OBJECTIVE 3** Use the discriminant to determine number and type of solutions.

The solutions of the quadratic equation $ax^2 + bx + c = 0$ are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \leftarrow \text{Discriminant}$$

The expression under the radical symbol, $b^2 - 4ac$, is called the **discriminant** because it distinguishes among the number of solutions—one or two—and the type of solutions—rational, irrational, or nonreal complex—of a quadratic equation.

NOW TRY ANSWER

4. $\{-2 \pm 3i\}$

Using the Discriminant

If a , b , and c are integers in a quadratic equation $ax^2 + bx + c = 0$, then the discriminant $b^2 - 4ac$ can be used to determine the number and type of solutions of the equation as follows.

Discriminant $b^2 - 4ac$	Number and Type of Solutions
Positive, and the square of an integer	Two rational solutions
Positive, but not the square of an integer	Two irrational solutions
Zero	One rational solution
Negative	Two nonreal complex solutions

We can also use the discriminant to help decide how to solve a quadratic equation.

If a , b , and c are integers and the discriminant is a perfect square (including 0), then the equation can be solved using the zero-factor property. Otherwise, the quadratic formula should be used.

EXAMPLE 5 Using the Discriminant

Find the discriminant. Use it to predict the number and type of solutions for each equation. Tell whether the equation can be solved using the zero-factor property, or if the quadratic formula should be used instead.

(a) $6x^2 - x - 15 = 0$

First identify the values of a , b , and c . Because $-x = -1x$, the value of b is -1 . We find the discriminant by evaluating $b^2 - 4ac$.

$$\begin{aligned}
 b^2 - 4ac &= (-1)^2 - 4(6)(-15) && \text{Use parentheses and substitute carefully.} \\
 &= 1 + 360 && a = 6, b = -1, c = -15 \text{ (all integers)} \\
 &= 361 && \text{Apply the exponent. Multiply.} \\
 &= 19^2, \text{ which is a perfect square.} && \text{Add.}
 \end{aligned}$$

The discriminant 361 is a perfect square, so referring to the table we see that there will be two rational solutions. We can solve using the zero-factor property.

(b) $3x^2 - 4x = 5$

Write in standard form as $3x^2 - 4x - 5 = 0$.

$$\begin{aligned}
 b^2 - 4ac &= (-4)^2 - 4(3)(-5) && \text{Discriminant} \\
 &= 16 + 60 && a = 3, b = -4, c = -5 \text{ (all integers)} \\
 &= 76 && \text{Apply the exponent. Multiply.} \\
 &&& \text{Add.}
 \end{aligned}$$

Because 76 is positive but *not* the square of an integer, the equation will have two irrational solutions. We solve using the quadratic formula.


NOW TRY EXERCISE 5

Find the discriminant. Use it to predict the number and type of solutions for each equation. Tell whether the equation can be solved using the zero-factor property, or if the quadratic formula should be used instead.

- (a) $8x^2 - 6x - 5 = 0$
 (b) $9x^2 = 24x - 16$
 (c) $3x^2 + 2x = -1$

NOW TRY ANSWERS

5. (a) 196; two rational solutions; zero-factor property
 (b) 0; one rational solution; zero-factor property
 (c) -8; two nonreal complex solutions; quadratic formula

(c) $4x^2 + x + 1 = 0$

$x = 1x$, so $b = 1$.

$$b^2 - 4ac$$

$$= 1^2 - 4(4)(1)$$

$$= 1 - 16$$

$$= -15$$

Discriminant

$a = 4, b = 1, c = 1$ (all integers)

Apply the exponent. Multiply.

Subtract.

Because the discriminant is negative, there will be two nonreal complex solutions. We solve using the quadratic formula.

(d) $4x^2 + 9 = 12x$ Write in standard form as $4x^2 - 12x + 9 = 0$.

$$b^2 - 4ac$$

$$= (-12)^2 - 4(4)(9)$$

$$= 144 - 144$$

$$= 0$$

Discriminant

$a = 4, b = -12, c = 9$ (all integers)

Apply the exponent. Multiply.

Subtract.

The discriminant is 0, so there is only one rational solution. We solve using the zero-factor property.


NOW TRY 

8.2 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Answer each question.

1. The documentation for an early version of Microsoft *Word* for Windows used the following for the quadratic formula. Was this correct? If not, correct it.

$$x = -b \pm \frac{\sqrt{b^2 - 4ac}}{2a}$$

Correct or incorrect?

2. One patron wrote the quadratic formula, as shown here, on a wall at the Cadillac Bar in Houston, Texas. Was this correct? If not, correct it.

$$x = \frac{-b\sqrt{b^2 - 4ac}}{2a}$$

Correct or incorrect?

3. A student solved $5x^2 - 5x + 1 = 0$ incorrectly as follows.

$$x = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(5)(1)}}{2(5)}$$

$a = 5, b = -5, c = 1$

$$x = \frac{5 \pm \sqrt{5}}{10}$$

Simplify.

$$x = \frac{1}{2} \pm \sqrt{5}$$

Write in lowest terms.

WHAT WENT WRONG? Give the correct solution set.

4. A student incorrectly claimed that the equation $2x^2 - 5 = 0$ cannot be solved using the quadratic formula because there is no first-degree x -term. **WHAT WENT WRONG?** Give the values of a , b , and c for this equation.

Use the quadratic formula to solve each equation. (All solutions for these equations are real numbers.) See Examples 1–3.

- | | | |
|---------------------------------------|---------------------------------------|-------------------------------------|
| 5. $x^2 - 8x + 15 = 0$ | 6. $x^2 + 3x - 28 = 0$ | 7. $6x^2 + 11x - 10 = 0$ |
| 8. $8x^2 + 10x - 3 = 0$ | 9. $4x^2 + 12x + 9 = 0$ | 10. $16x^2 + 40x + 25 = 0$ |
| 11. $36x^2 - 12x + 1 = 0$ | 12. $9x^2 - 6x + 1 = 0$ | 13. $2x^2 + 4x + 1 = 0$ |
| 14. $2x^2 + 3x - 1 = 0$ | 15. $2x^2 - 2x = 1$ | 16. $9x^2 + 6x = 1$ |
| 17. $x^2 + 18 = 10x$ | 18. $x^2 - 4 = 2x$ | 19. $4x^2 + 4x - 1 = 0$ |
| 20. $4r^2 - 4r - 19 = 0$ | 21. $2 - 2x = 3x^2$ | 22. $26r - 2 = 3r^2$ |
| 23. $\frac{x^2}{4} - \frac{x}{2} = 1$ | 24. $p^2 + \frac{p}{3} = \frac{1}{6}$ | 25. $-2t(t + 2) = -3$ |
| 26. $-3x(x + 2) = -4$ | 27. $(r - 3)(r + 5) = 2$ | 28. $(x + 1)(x - 7) = 1$ |
| 29. $(x + 2)(x - 3) = 1$ | 30. $(x - 5)(x + 2) = 6$ | 31. $p = \frac{5(5 - p)}{3(p + 1)}$ |
| 32. $x = \frac{2(x + 3)}{x + 5}$ | 33. $(2x + 1)^2 = x + 4$ | 34. $(2x - 1)^2 = x + 2$ |

Use the quadratic formula to solve each equation. (All solutions for these equations are nonreal complex numbers.) See Example 4.

- | | | |
|----------------------------------------|----------------------------------------|-------------------------|
| 35. $x^2 - 3x + 6 = 0$ | 36. $x^2 - 5x + 20 = 0$ | 37. $r^2 - 6r + 14 = 0$ |
| 38. $t^2 + 4t + 11 = 0$ | 39. $4x^2 - 4x = -7$ | 40. $9x^2 - 6x = -7$ |
| 41. $x(3x + 4) = -2$ | 42. $z(2z + 3) = -2$ | |
| 43. $(2x - 1)(8x - 4) = -1$ | 44. $(x - 1)(9x - 3) = -2$ | |
| 45. $(6x + 1)(x - 2) = (x + 2)(x - 5)$ | 46. $(4x + 3)(x - 1) = (x + 3)(x - 6)$ | |

Find the discriminant. Use it to determine whether the solutions for each equation are

- A. two rational numbers B. one rational number
C. two irrational numbers D. two nonreal complex numbers.

Tell whether the equation can be solved using the zero-factor property, or if the quadratic formula should be used instead. Do not actually solve. See Example 5.

- | | | |
|----------------------------|----------------------------|------------------------|
| 47. $25x^2 + 70x + 49 = 0$ | 48. $4x^2 - 28x + 49 = 0$ | 49. $x^2 + 4x + 2 = 0$ |
| 50. $9x^2 - 12x - 1 = 0$ | 51. $3x^2 = 5x + 2$ | 52. $4x^2 = 4x + 3$ |
| 53. $3m^2 - 10m + 15 = 0$ | 54. $18x^2 + 60x + 82 = 0$ | |
55. Find the discriminant for each quadratic equation. Use it to tell whether the equation can be solved using the zero-factor property, or the quadratic formula should be used instead. Then solve each equation.
- (a) $3x^2 + 13x = -12$ (b) $2x^2 + 19 = 14x$
56. Refer to the answers in Exercises 47–54, and solve the equation given in each exercise.
- (a) $25x^2 + 70x + 49 = 0$ (Exercise 47) (b) $4x^2 - 28x + 49 = 0$ (Exercise 48)
(c) $3x^2 = 5x + 2$ (Exercise 51) (d) $4x^2 = 4x + 3$ (Exercise 52)

Extending Skills Find the value of a , b , or c so that each equation will have exactly one rational solution. (Hint: The discriminant must equal 0 for an equation to have one rational solution.)

57. $p^2 + bp + 25 = 0$

58. $r^2 - br + 49 = 0$

59. $am^2 + 8m + 1 = 0$

60. $at^2 + 24t + 16 = 0$

61. $9x^2 - 30x + c = 0$

62. $4m^2 + 12m + c = 0$

63. One solution of $4x^2 + bx - 3 = 0$ is $-\frac{5}{2}$. Find b and the other solution.64. One solution of $3x^2 - 7x + c = 0$ is $\frac{1}{3}$. Find c and the other solution.

8.3

Equations That Lead to Quadratic Methods

OBJECTIVES

- 1 Solve rational equations that lead to quadratic equations.
- 2 Solve applied problems involving quadratic equations.
- 3 Solve radical equations that lead to quadratic equations.
- 4 Solve equations that are quadratic in form.

VOCABULARY

□ quadratic in form


NOW TRY EXERCISE 1

Solve $\frac{2}{x} + \frac{3}{x+2} = 1$.

OBJECTIVE 1 Solve rational equations that lead to quadratic equations.

A variety of nonquadratic equations can be written in the form of a quadratic equation and solved using the methods of this chapter.

EXAMPLE 1 Solving a Rational Equation That Leads to a Quadratic Equation

Solve $\frac{1}{x} + \frac{1}{x-1} = \frac{7}{12}$.

Clear fractions by multiplying each side by the least common denominator, $12x(x-1)$. (The domain is $\{x \mid x \text{ is a real number, } x \neq 0, 1\}$.)

$$12x(x-1)\left(\frac{1}{x} + \frac{1}{x-1}\right) = 12x(x-1)\left(\frac{7}{12}\right) \quad \text{Multiply by the LCD.}$$

$$12x(x-1)\frac{1}{x} + 12x(x-1)\frac{1}{x-1} = 12x(x-1)\frac{7}{12} \quad \text{Distributive property}$$

$$12(x-1) + 12x = x(x-1) \cdot 7 \quad \text{Multiply.}$$

$$12x - 12 + 12x = 7x^2 - 7x \quad \text{Distributive property}$$

$$24x - 12 = 7x^2 - 7x \quad \text{Combine like terms.}$$

This trinomial is factorable. $7x^2 - 31x + 12 = 0$

$$(7x-3)(x-4) = 0 \quad \text{Standard form}$$

$$7x-3=0 \quad \text{or} \quad x-4=0 \quad \text{Factor.}$$

$$x = \frac{3}{7} \quad \text{or} \quad x = 4 \quad \text{Zero-factor property}$$

Solve each equation.

These values are in the domain. Check them in the original equation. The solution set is $\left\{\frac{3}{7}, 4\right\}$.

NOW TRY 

OBJECTIVE 2 Solve applied problems involving quadratic equations.


Some distance-rate-time (or motion) problems lead to quadratic equations. We use a form of the distance formula $d = rt$ to solve them.

NOW TRY ANSWER

1. $\{-1, 4\}$


NOW TRY EXERCISE 2

A small fishing boat averages 18 mph in still water. It takes the boat $\frac{9}{10}$ hr to travel 8 mi upstream and return. Find the rate of the current.



Riverboat traveling upstream—the current slows it down.

FIGURE 1

EXAMPLE 2 Solving a Motion Problem

A riverboat for tourists averages 12 mph in still water. It takes the boat 1 hr, 4 min to travel 6 mi upstream and return. Find the rate of the current.

Step 1 Read the problem carefully.

Step 2 Assign a variable. Let x = the rate of the current.

The current slows down the boat as it travels upstream, so the rate of the boat traveling upstream is its rate in still water *less* the rate of the current, or $(12 - x)$ mph. See **FIGURE 1**.

Similarly, the current speeds up the boat as it travels downstream, so its rate downstream is $(12 + x)$ mph. Thus,

$$12 - x = \text{the rate upstream in miles per hour,}$$

and $12 + x = \text{the rate downstream in miles per hour.}$

	d	r	t
Upstream	6	$12 - x$	$\frac{6}{12 - x}$
Downstream	6	$12 + x$	$\frac{6}{12 + x}$

Complete a table. Use the distance formula, $d = rt$, solved for time t , $t = \frac{d}{r}$, to write expressions for t .

Step 3 Write an equation. We use the total time, 1 hr, 4 min, written as a fraction.

$$1 + \frac{4}{60} = 1 + \frac{1}{15} = \frac{16}{15} \text{ hr} \quad \text{Total time}$$

The time upstream plus the time downstream equals $\frac{16}{15}$ hr.

$$\begin{array}{rcc} \text{Time upstream} & + & \text{Time downstream} & = & \text{Total time} \\ \downarrow & & \downarrow & & \downarrow \\ \frac{6}{12 - x} & + & \frac{6}{12 + x} & = & \frac{16}{15} \end{array}$$

Step 4 Solve the equation. The LCD is $15(12 - x)(12 + x)$.

$$\begin{aligned} 15(12 - x)(12 + x) \left(\frac{6}{12 - x} + \frac{6}{12 + x} \right) \\ = 15(12 - x)(12 + x) \left(\frac{16}{15} \right) \end{aligned}$$

Multiply by the LCD.

$$15(12 + x) \cdot 6 + 15(12 - x) \cdot 6 = (12 - x)(12 + x)16$$

Distributive property; Multiply.

$$90(12 + x) + 90(12 - x) = 16(144 - x^2)$$

Multiply.

$$1080 + 90x + 1080 - 90x = 2304 - 16x^2$$

Distributive property

$$2160 = 2304 - 16x^2$$

Combine like terms.

$$16x^2 = 144$$

Add $16x^2$. Subtract 2160.

$$x^2 = 9$$

Divide by 16.

$$x = 3 \quad \text{or} \quad x = -3$$

Square root property

Step 5 State the answer. The current rate cannot be -3 , so the answer is 3 mph.

Step 6 Check that this value satisfies the original problem.

NOW TRY 

NOW TRY ANSWER

2. 2 mph

PROBLEM-SOLVING HINT Recall from earlier work that a person's work rate is $\frac{1}{t}$ part of the job per hour, where t is the time in hours required to do the complete job. Thus, the part of the job the person will do in x hours is $\frac{1}{t}x$.



EXAMPLE 3 Solving a Work Problem

It takes two carpet layers 4 hr to carpet a room. If each worked alone, one of them could do the job in 1 hr less time than the other. How long would it take each carpet layer to complete the job alone?

Step 1 Read the problem again. There will be two answers.

Step 2 Assign a variable.

Let x = the number of hours for the slower carpet layer to complete the job.

Then $x - 1$ = the number of hours for the faster carpet layer to complete the job.

The slower worker's rate is $\frac{1}{x}$, and the faster worker's rate is $\frac{1}{x-1}$. Together they can do the job in 4 hr. Complete a table as shown.

	Rate	Time Working Together	Fractional Part of the Job Done
Slower Worker	$\frac{1}{x}$	4	$\frac{1}{x}(4)$
Faster Worker	$\frac{1}{x-1}$	4	$\frac{1}{x-1}(4)$

Sum is 1 whole job.

Step 3 Write an equation.

$$\begin{array}{rcccl} \text{Part done by slower worker} & + & \text{Part done by faster worker} & = & \text{1 whole job} \\ \downarrow & & \downarrow & & \downarrow \\ \frac{4}{x} & + & \frac{4}{x-1} & = & 1 \end{array}$$

Step 4 Solve the equation. The LCD is $x(x-1)$.

$$\begin{aligned} x(x-1)\left(\frac{4}{x} + \frac{4}{x-1}\right) &= x(x-1)(1) && \text{Multiply by the LCD.} \\ (x-1) \cdot 4 + 4x &= x(x-1) && \text{Distributive property} \\ 4x - 4 + 4x &= x^2 - x && \text{Distributive property} \\ x^2 - 9x + 4 &= 0 && \text{Standard form} \end{aligned}$$

The trinomial on the left cannot be factored, so the equation cannot be solved using the zero-factor property. We use the quadratic formula.

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} && \text{Quadratic formula} \\ x &= \frac{-(-9) \pm \sqrt{(-9)^2 - 4(1)(4)}}{2(1)} && a = 1, b = -9, c = 4 \\ x &= \frac{9 \pm \sqrt{65}}{2} && \text{Simplify.} \\ x &= \frac{9 + \sqrt{65}}{2} = 8.5 \quad \text{or} \quad x = \frac{9 - \sqrt{65}}{2} = 0.5 && \text{Use a calculator. Round to the nearest tenth.} \end{aligned}$$


NOW TRY EXERCISE 3

Two electricians are running wire to finish a basement. One electrician could finish the job in 2 hr less time than the other. Together, they complete the job in 6 hr. How long (to the nearest tenth) would it take the slower electrician to complete the job alone?


NOW TRY EXERCISE 4

Solve each equation.

(a) $x = \sqrt{9x - 20}$

(b) $x + \sqrt{x} = 20$

Step 5 State the answer. Only the solution 8.5 makes sense in the original problem.

$$\text{If } x = 0.5, \text{ then } x - 1 = 0.5 - 1 = -0.5, \quad \text{Time cannot be negative.}$$

which cannot represent the time for the faster worker. The slower worker could do the job in 8.5 hr and the faster in $8.5 - 1 = 7.5$ hr.

Step 6 Check that these results satisfy the original problem.

NOW TRY 

OBJECTIVE 3 Solve radical equations that lead to quadratic equations.

EXAMPLE 4 Solving Radical Equations That Lead to Quadratic Equations

Solve each equation.

(a) $x = \sqrt{6x - 8}$

This equation is not quadratic. However, squaring each side of the equation gives a quadratic equation that can be solved using the zero-factor property.

$$x^2 = (\sqrt{6x - 8})^2 \quad \text{Square each side.}$$

$$x^2 = 6x - 8 \quad (\sqrt{a})^2 = a$$

Standard form

This trinomial is factorable. $x^2 - 6x + 8 = 0$

$$(x - 4)(x - 2) = 0 \quad \text{Factor.}$$

$$x - 4 = 0 \quad \text{or} \quad x - 2 = 0 \quad \text{Zero-factor property}$$

$$x = 4 \quad \text{or} \quad x = 2 \quad \text{Proposed solutions}$$

Squaring each side of a radical equation can introduce extraneous solutions. **All proposed solutions must be checked in the original (not the squared) equation.**

CHECK $x = \sqrt{6x - 8}$	$x = \sqrt{6x - 8}$
$4 \stackrel{?}{=} \sqrt{6(4) - 8}$ Let $x = 4$.	$2 \stackrel{?}{=} \sqrt{6(2) - 8}$ Let $x = 2$.
$4 \stackrel{?}{=} \sqrt{16}$	$2 \stackrel{?}{=} \sqrt{4}$
$4 = 4$ ✓ True	$2 = 2$ ✓ True

Both proposed solutions check, so the solution set is $\{2, 4\}$.

(b) $x + \sqrt{x} = 6$

$$\sqrt{x} = 6 - x \quad \text{Isolate the radical on one side.}$$

$$(\sqrt{x})^2 = (6 - x)^2 \quad \text{Square each side.}$$

$$x = 36 - 12x + x^2 \quad (x - y)^2 = x^2 - 2xy + y^2$$

$$x^2 - 13x + 36 = 0 \quad \text{Write in standard form.}$$

$$(x - 4)(x - 9) = 0 \quad \text{Factor.}$$

$$x - 4 = 0 \quad \text{or} \quad x - 9 = 0 \quad \text{Zero-factor property}$$

$$x = 4 \quad \text{or} \quad x = 9 \quad \text{Proposed solutions}$$

CHECK $x + \sqrt{x} = 6$	$x + \sqrt{x} = 6$
$4 + \sqrt{4} \stackrel{?}{=} 6$ Let $x = 4$.	$9 + \sqrt{9} \stackrel{?}{=} 6$ Let $x = 9$.
$6 = 6$ ✓ True	$12 = 6$ False

Only the proposed solution 4 checks, so the solution set is $\{4\}$.

NOW TRY ANSWERS

3. 13.1 hr

4. (a) $\{4, 5\}$ (b) $\{16\}$

NOW TRY 

OBJECTIVE 4 Solve equations that are quadratic in form.

A nonquadratic equation that can be written in the form

$$au^2 + bu + c = 0,$$

for $a \neq 0$ and an algebraic expression u , is **quadratic in form**.

Many equations that are quadratic in form can be solved more easily by defining and substituting a “temporary” variable u for an expression involving the variable in the original equation.

 **NOW TRY EXERCISE 5**

Define a variable u in terms of x , and write each equation in the quadratic form

$$au^2 + bu + c = 0.$$

(a) $x^4 - 10x^2 + 9 = 0$

(b) $6(x + 2)^2 - 11(x + 2) + 4 = 0$

EXAMPLE 5 Defining Substitution Variables

Define a variable u in terms of x , and write each equation in the quadratic form $au^2 + bu + c = 0$.

(a) $x^4 - 13x^2 + 36 = 0$

Look at the two terms involving the variable x , ignoring their coefficients. Try to find one variable expression that is the square of the other. Here $x^4 = (x^2)^2$, so we can define $u = x^2$, and rewrite the original equation as a quadratic equation in u .

$$u^2 - 13u + 36 = 0 \quad \text{Here, } u = x^2.$$

(b) $2(4x - 3)^2 + 7(4x - 3) + 5 = 0$

Because this equation involves both $(4x - 3)^2$ and $(4x - 3)$, we let $u = 4x - 3$.

$$2u^2 + 7u + 5 = 0 \quad \text{Here, } u = 4x - 3.$$

(c) $2x^{2/3} - 11x^{1/3} + 12 = 0$

We apply a power rule for exponents, $(a^m)^n = a^{mn}$. Because $(x^{1/3})^2 = x^{2/3}$, we define $u = x^{1/3}$ and write the original equation as follows.

$$2u^2 - 11u + 12 = 0 \quad \text{Here, } u = x^{1/3}.$$

NOW TRY 

EXAMPLE 6 Solving Equations That Are Quadratic in Form

Solve each equation.

(a) $x^4 - 13x^2 + 36 = 0$ See Example 5(a).

$$(x^2)^2 - 13x^2 + 36 = 0 \quad x^4 = (x^2)^2$$

Quadratic in form $\rightarrow u^2 - 13u + 36 = 0$ Let $u = x^2$.

$$(u - 4)(u - 9) = 0 \quad \text{Factor.}$$

$$u - 4 = 0 \quad \text{or} \quad u - 9 = 0 \quad \text{Zero-factor property}$$

Don't stop here. $\rightarrow u = 4$ or $u = 9$ Solve.

$$x^2 = 4 \quad \text{or} \quad x^2 = 9 \quad \text{Substitute } x^2 \text{ for } u.$$

$$x = \pm 2 \quad \text{or} \quad x = \pm 3 \quad \text{Square root property}$$

Each value can be verified by substituting it into the original equation for x . The equation $x^4 - 13x^2 + 36 = 0$ is a fourth-degree equation and has four solutions, $-3, -2, 2, 3$.* The solution set is abbreviated $\{\pm 2, \pm 3\}$.

NOW TRY ANSWERS

5. (a) $u = x^2$; $u^2 - 10u + 9 = 0$

(b) $u = x + 2$;
 $6u^2 - 11u + 4 = 0$

*In general, an equation in which an n th-degree polynomial equals 0 has n complex solutions, although some of them may be repeated.


NOW TRY EXERCISE 6

Solve each equation.

(a) $x^4 - 17x^2 + 16 = 0$

(b) $x^4 + 4 = 8x^2$

(b) $4x^4 + 1 = 5x^2$

$4x^4 - 5x^2 + 1 = 0$ Standard form

$4(x^2)^2 - 5x^2 + 1 = 0$ $x^4 = (x^2)^2$

$4u^2 - 5u + 1 = 0$ Let $u = x^2$.

$(4u - 1)(u - 1) = 0$ Factor.

$4u - 1 = 0$ or $u - 1 = 0$ Zero-factor property

$u = \frac{1}{4}$ or $u = 1$ Solve.


 $x^2 = \frac{1}{4}$ or $x^2 = 1$ Substitute x^2 for u .

$x = \pm \frac{1}{2}$ or $x = \pm 1$ Square root property

Check that the solution set is $\left\{\pm \frac{1}{2}, \pm 1\right\}$.

(c) $x^4 = 6x^2 - 3$

$x^4 - 6x^2 + 3 = 0$ Standard form

$(x^2)^2 - 6x^2 + 3 = 0$ $x^4 = (x^2)^2$

$u^2 - 6u + 3 = 0$ Let $u = x^2$.

The trinomial on the left is nonfactorable, so we cannot solve the equation using the zero-factor property. To solve, we use the quadratic formula.

$$u = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(1)(3)}}{2(1)}$$
 $a = 1, b = -6, c = 3$

$$u = \frac{6 \pm \sqrt{24}}{2}$$
Simplify.

$$u = \frac{6 \pm 2\sqrt{6}}{2}$$
 $\sqrt{24} = \sqrt{4} \cdot \sqrt{6} = 2\sqrt{6}$

$$u = \frac{2(3 \pm \sqrt{6})}{2}$$
Factor.

$$u = 3 \pm \sqrt{6}$$
Divide out the common factor 2.


 $x^2 = 3 + \sqrt{6}$ or $x^2 = 3 - \sqrt{6}$
Substitute x^2 for u .

$x = \pm \sqrt{3 + \sqrt{6}}$ or $x = \pm \sqrt{3 - \sqrt{6}}$
Square root property

The solution set contains four numbers and is written as follows.

NOW TRY ANSWERS

6. (a) $\{\pm 1, \pm 4\}$

(b) $\left\{\pm \sqrt{4 + 2\sqrt{3}}, \pm \sqrt{4 - 2\sqrt{3}}\right\}$

$\left\{\pm \sqrt{3 + \sqrt{6}}, \pm \sqrt{3 - \sqrt{6}}\right\}$

NOW TRY 

NOTE Expressions in equations like those in **Examples 6(a) and (b)** can be factored directly.

$$x^4 - 13x^2 + 36 = 0 \quad \text{Example 6(a) equation}$$

$$(x^2 - 9)(x^2 - 4) = 0 \quad \text{Factor.}$$

$$(x + 3)(x - 3)(x + 2)(x - 2) = 0 \quad \text{Factor again.}$$

Using the zero-factor property gives the same solutions that we obtained in **Example 6(a)**. Equations that include nonfactorable quadratic expressions (as in **Example 6(c)**) must be solved using substitution and the quadratic formula.

Solving an Equation That Is Quadratic in Form by Substitution

Step 1 Define a temporary variable u , based on the relationship between the variable expressions in the given equation. Substitute u in the original equation and rewrite the equation in the form

$$au^2 + bu + c = 0.$$

Step 2 Solve the quadratic equation obtained in **Step 1** either by factoring the trinomial and applying the zero-factor property or by using the quadratic formula.

Step 3 Replace u with the expression it defined in **Step 1**.

Step 4 Solve the resulting equations for the original variable.

Step 5 Check all solutions by substituting them in the original equation. Write the solution set.

EXAMPLE 7 Solving Equations That Are Quadratic in Form

Solve each equation.

(a) $2(4x - 3)^2 + 7(4x - 3) + 5 = 0$

Step 1 Because of the repeated quantity $4x - 3$, substitute u for $4x - 3$.

$$2(4x - 3)^2 + 7(4x - 3) + 5 = 0 \quad \text{See Example 5(b).}$$

$$2u^2 + 7u + 5 = 0 \quad \text{Let } u = 4x - 3.$$

Step 2 $(2u + 5)(u + 1) = 0$ Factor.

$$2u + 5 = 0 \quad \text{or} \quad u + 1 = 0 \quad \text{Zero-factor property}$$

Don't stop here. $u = -\frac{5}{2}$ or $u = -1$ Solve for u .

Step 3 $4x - 3 = -\frac{5}{2}$ or $4x - 3 = -1$ Substitute $4x - 3$ for u .

Step 4 $4x = \frac{1}{2}$ or $4x = 2$ Solve for x .

$$x = \frac{1}{8} \quad \text{or} \quad x = \frac{1}{2}$$

Step 5 Check that the solution set of the original equation is $\left\{\frac{1}{8}, \frac{1}{2}\right\}$.


NOW TRY EXERCISE 7

Solve each equation.

(a) $6(x - 4)^2 + 11(x - 4) - 10 = 0$

(b) $2x^{2/3} - 7x^{1/3} + 3 = 0$

(b) $2x^{2/3} - 11x^{1/3} + 12 = 0$

Step 1 Because $x^{2/3} = (x^{1/3})^2$, we substitute u for $x^{1/3}$.

$$2x^{2/3} - 11x^{1/3} + 12 = 0$$

See Example 5(c).

$$2(x^{1/3})^2 - 11x^{1/3} + 12 = 0$$

$$x^{2/3} = (x^{1/3})^2$$

$$2u^2 - 11u + 12 = 0$$

Let $u = x^{1/3}$.

Step 2

$$(2u - 3)(u - 4) = 0$$

Factor.

$$2u - 3 = 0 \quad \text{or} \quad u - 4 = 0$$

Zero-factor property

$$u = \frac{3}{2} \quad \text{or} \quad u = 4$$

Solve for u .**Step 3**

$$x^{1/3} = \frac{3}{2} \quad \text{or} \quad x^{1/3} = 4$$

Substitute $x^{1/3}$ for u .**Step 4**

$$(x^{1/3})^3 = \left(\frac{3}{2}\right)^3 \quad \text{or} \quad (x^{1/3})^3 = 4^3$$

Cube each side.

$$x = \frac{27}{8} \quad \text{or} \quad x = 64$$

Apply the exponents.

Step 5 Because the original equation involves variables with rational exponents, check that neither of these solutions is extraneous. The solution set is $\left\{\frac{27}{8}, 64\right\}$.


NOW TRY

CAUTION A common error when solving problems like those in Examples 6 and 7 is to stop too soon. *Once we have solved for u , we must remember to substitute and solve for the values of the original variable.*


NOW TRY ANSWERS

7. (a) $\left\{\frac{3}{2}, \frac{14}{3}\right\}$ (b) $\left\{\frac{1}{8}, 27\right\}$

8.3 Exercises

FOR
EXTRA
HELP

MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Based on the discussion and examples of this section, give the first step to solve each equation. Do not actually solve.

1. $\frac{14}{x} = x - 5$

2. $\sqrt{1+x} + x = 5$

3. $(x^2 + x)^2 - 8(x^2 + x) + 12 = 0$

4. $3x = \sqrt{16 - 10x}$

5. **Concept Check** Study this incorrect “solution.”

6. **Concept Check** Study this incorrect “solution.”

$$x = \sqrt{3x + 4} \quad \text{Square each side.}$$

$$x^2 = 3x + 4$$

$$2(x - 1)^2 - 3(x - 1) + 1 = 0 \quad \text{Let } u = x - 1.$$

$$2u^2 - 3u + 1 = 0$$

$$x^2 - 3x - 4 = 0$$

$$(2u - 1)(u - 1) = 0$$

$$(x - 4)(x + 1) = 0$$

$$2u - 1 = 0 \quad \text{or} \quad u - 1 = 0$$

$$x - 4 = 0 \quad \text{or} \quad x + 1 = 0$$

$$u = \frac{1}{2} \quad \text{or} \quad u = 1$$

$$x = 4 \quad \text{or} \quad x = -1$$

Solution set: $\{4, -1\}$ Solution set: $\left\{\frac{1}{2}, 1\right\}$

WHAT WENT WRONG? Give the correct solution set.

WHAT WENT WRONG? Give the correct solution set.

Solve each equation. Check the solutions. See Example 1.

7. $\frac{14}{x} = x - 5$

8. $\frac{-12}{x} = x + 8$

9. $1 - \frac{3}{x} - \frac{28}{x^2} = 0$

10. $4 - \frac{7}{r} - \frac{2}{r^2} = 0$

11. $3 - \frac{1}{t} = \frac{2}{t^2}$

12. $1 + \frac{2}{x} = \frac{3}{x^2}$

13. $\frac{1}{x} + \frac{2}{x+2} = \frac{17}{35}$

14. $\frac{2}{m} + \frac{3}{m+9} = \frac{11}{4}$

15. $\frac{2}{x+1} + \frac{3}{x+2} = \frac{7}{2}$

16. $\frac{4}{3-p} + \frac{2}{5-p} = \frac{26}{15}$

17. $\frac{3}{2x} - \frac{1}{2(x+2)} = 1$

18. $\frac{4}{3x} - \frac{1}{2(x+1)} = 1$

19. $3 = \frac{1}{t+2} + \frac{2}{(t+2)^2}$

20. $1 + \frac{2}{3z+2} = \frac{15}{(3z+2)^2}$

21. $\frac{6}{p} = 2 + \frac{p}{p+1}$

22. $\frac{x}{2-x} + \frac{2}{x} = 5$

23. $1 - \frac{1}{2x+1} - \frac{1}{(2x+1)^2} = 0$

24. $1 - \frac{1}{3x-2} - \frac{1}{(3x-2)^2} = 0$

Concept Check Answer each question.

25. A boat travels 20 mph in still water, and the rate of the current is t mph.

- (a) What is the rate of the boat when it travels upstream?
 (b) What is the rate of the boat when it travels downstream?

26. It takes m hours to grade a set of papers.

- (a) What is the grader's rate (in job per hour)?
 (b) How much of the job will the grader do in 2 hr?

Solve each problem. Round answers to the nearest tenth as needed. See Examples 2 and 3.

27. In 4 hr, Kerrie can travel 15 mi upriver and come back. The rate of the current is 5 mph. Find the rate of her boat in still water.

Let $x =$ _____.

The rate traveling upriver (*against* the current) is _____ mph.

The rate traveling back downriver (*with* the current) is _____ mph.

Complete the table.

	d	r	t
Up			
Down			

Write an equation, and complete the solution.

28. Carlos can complete a certain lab test in 2 hr less time than Jaime can. If they can finish the job together in 2 hr, how long would it take each of them working alone?

Let $x =$ Jaime's time alone (in hours).

Then _____ = Carlos' time alone (in hours).

Complete the table.

	Rate	Time Working Together	Fractional Part of the Job Done
Carlos			
Jaime			

Write an equation, and complete the solution.

29. On a windy day William found that he could travel 16 mi downstream and then 4 mi back upstream at top speed in a total of 48 min. What was the top speed of William’s boat if the rate of the current was 15 mph? (Let x represent the rate of the boat in still water.)

	d	r	t
Upstream	4	$x - 15$	
Downstream	16		

30. Vera flew for 6 hr at a constant rate. She traveled 810 mi with the wind, then turned around and traveled 720 mi against the wind. The wind speed was a constant 15 mph. Find the rate of the plane.

	d	r	t
With Wind	810		
Against Wind	720		

31. The distance from Jackson to Lodi is about 40 mi, as is the distance from Lodi to Manteca. Adrian drove from Jackson to Lodi, stopped in Lodi for a high-energy drink, and then drove on to Manteca at 10 mph faster. Driving time for the entire trip was 88 min. Find her rate from Jackson to Lodi. (Data from *State Farm Road Atlas*.)
32. Medicine Hat and Cranbrook are 300 km apart. Steve rides his Harley 20 km per hr faster than Mohammad rides his Yamaha. Find Steve’s average rate if he travels from Cranbrook to Medicine Hat in $1\frac{1}{4}$ hr less time than Mohammad. (Data from *State Farm Road Atlas*.)



33. Working together, two people can cut a large lawn in 2 hr. One person can do the job alone in 1 hr less time than the other. How long would it take the faster worker to do the job? (Let x represent the time of the faster worker.)
34. Working together, two people can clean an office building in 5 hr. One person takes 2 hr longer than the other to clean the building alone. How long would it take the slower worker to clean the building alone? (Let x represent the time of the slower worker.)

	Rate	Time Working Together	Fractional Part of the Job Done
Faster Worker	$\frac{1}{x}$	2	
Slower Worker		2	

	Rate	Time Working Together	Fractional Part of the Job Done
Faster Worker			
Slower Worker	$\frac{1}{x}$		

35. Rusty and Nancy are planting flowers. Working alone, Rusty would take 2 hr longer than Nancy to plant the flowers. Working together, they do the job in 12 hr. How long would it have taken each person working alone?
36. Joel can work through a stack of invoices in 1 hr less time than Noel can. Working together they take $1\frac{1}{2}$ hr. How long would it take each person working alone?

37. A washing machine can be filled in 6 min if both the hot water and the cold water taps are fully opened. Filling the washer with hot water alone takes 9 min longer than filling it with cold water alone. How long does it take to fill the washer with cold water?
38. Two pipes together can fill a tank in 2 hr. One of the pipes, used alone, takes 3 hr longer than the other to fill the tank. How long would each pipe take to fill the tank alone?

Solve each equation. Check the solutions. See Example 4.

39. $x = \sqrt{7x - 10}$	40. $z = \sqrt{5z - 4}$	41. $2x = \sqrt{11x + 3}$
42. $4x = \sqrt{6x + 1}$	43. $3x = \sqrt{16 - 10x}$	44. $4t = \sqrt{8t + 3}$
45. $t + \sqrt{t} = 12$	46. $p - 2\sqrt{p} = 8$	47. $x = \sqrt{\frac{6 - 13x}{5}}$
48. $r = \sqrt{\frac{20 - 19r}{6}}$	49. $-x = \sqrt{\frac{8 - 2x}{3}}$	50. $-x = \sqrt{\frac{3x + 7}{4}}$

Solve each equation. Check the solutions. See Examples 5–7.

51. $x^4 - 29x^2 + 100 = 0$	52. $x^4 - 37x^2 + 36 = 0$
53. $4q^4 - 13q^2 + 9 = 0$	54. $9x^4 - 25x^2 + 16 = 0$
55. $x^4 + 48 = 16x^2$	56. $z^4 + 72 = 17z^2$
57. $(x + 3)^2 + 5(x + 3) + 6 = 0$	58. $(x - 4)^2 + (x - 4) - 20 = 0$
59. $3(m + 4)^2 - 8 = 2(m + 4)$	60. $(t + 5)^2 + 6 = 7(t + 5)$
61. $x^{2/3} + x^{1/3} - 2 = 0$	62. $x^{2/3} - 2x^{1/3} - 3 = 0$
63. $r^{2/3} + r^{1/3} - 12 = 0$	64. $3x^{2/3} - x^{1/3} - 24 = 0$
65. $4x^{4/3} - 13x^{2/3} + 9 = 0$	66. $9t^{4/3} - 25t^{2/3} + 16 = 0$
67. $2 + \frac{5}{3x - 1} = \frac{-2}{(3x - 1)^2}$	68. $3 - \frac{7}{2p + 2} = \frac{6}{(2p + 2)^2}$
69. $2 - 6(z - 1)^{-2} = (z - 1)^{-1}$	70. $3 - 2(x - 1)^{-1} = (x - 1)^{-2}$

The following exercises are not grouped by type. Solve each equation. (Exercises 83 and 84 require knowledge of complex numbers.) See Examples 1 and 4–7.

71. $12x^4 - 11x^2 + 2 = 0$	72. $\left(x - \frac{1}{2}\right)^2 + 5\left(x - \frac{1}{2}\right) - 4 = 0$
73. $\sqrt{2x + 3} = 2 + \sqrt{x - 2}$	74. $\sqrt{m + 1} = -1 + \sqrt{2m}$
75. $2(1 + \sqrt{r})^2 = 13(1 + \sqrt{r}) - 6$	76. $(x^2 + x)^2 + 12 = 8(x^2 + x)$
77. $2m^6 + 11m^3 + 5 = 0$	78. $8x^6 + 513x^3 + 64 = 0$
79. $6 = 7(2w - 3)^{-1} + 3(2w - 3)^{-2}$	80. $x^6 - 10x^3 = -9$
81. $2x^4 - 9x^2 = -2$	82. $8x^4 + 1 = 11x^2$
83. $2x^4 + x^2 - 3 = 0$	84. $4x^4 + 5x^2 + 1 = 0$

SUMMARY EXERCISES Applying Methods for Solving Quadratic Equations

We have introduced four methods for solving quadratic equations written in standard form $ax^2 + bx + c = 0$.

Method	Advantages	Disadvantages
Zero-factor property	This is usually the fastest method.	Not all polynomials are factorable. Some factorable polynomials are difficult to factor.
Square root property	This is the simplest method for solving equations of the form $(ax + b)^2 = c$.	Few equations are given in this form.
Completing the square	This method can always be used.	It requires more steps than other methods.
Quadratic formula	This method can always be used.	Sign errors may occur when evaluating $\sqrt{b^2 - 4ac}$.

Concept Check Decide whether the zero-factor property, the square root property, or the quadratic formula is most appropriate for solving each quadratic equation. Do not actually solve.

- | | | |
|---------------------|--------------------|-----------------------|
| 1. $(2x + 3)^2 = 4$ | 2. $4x^2 - 3x = 1$ | 3. $x^2 + 5x - 8 = 0$ |
| 4. $2x^2 + 3x = 1$ | 5. $3x^2 = 2 - 5x$ | 6. $x^2 = 5$ |

Solve each quadratic equation by the method of your choice.

- | | | |
|----------------------------------------|----------------------------------------|---------------------------------------------------|
| 7. $p^2 = 7$ | 8. $6x^2 - x - 15 = 0$ | 9. $n^2 + 6n + 4 = 0$ |
| 10. $(x - 3)^2 = 25$ | 11. $\frac{5}{x} + \frac{12}{x^2} = 2$ | 12. $3x^2 = 3 - 8x$ |
| 13. $2r^2 - 4r + 1 = 0$ | *14. $x^2 = -12$ | 15. $x\sqrt{2} = \sqrt{5x - 2}$ |
| 16. $x^4 - 10x^2 + 9 = 0$ | 17. $(2x + 3)^2 = 8$ | 18. $\frac{2}{x} + \frac{1}{x - 2} = \frac{5}{3}$ |
| 19. $t^4 + 14 = 9t^2$ | 20. $8x^2 - 4x = 2$ | *21. $z^2 + z + 1 = 0$ |
| 22. $5x^6 + 2x^3 - 7 = 0$ | 23. $4t^2 - 12t + 9 = 0$ | 24. $x\sqrt{3} = \sqrt{2 - x}$ |
| 25. $r^2 - 72 = 0$ | 26. $-3x^2 + 4x = -4$ | 27. $x^2 - 5x - 36 = 0$ |
| 28. $w^2 = 169$ | *29. $3p^2 = 6p - 4$ | 30. $z = \sqrt{\frac{5z + 3}{2}}$ |
| *31. $\frac{4}{r^2} + 3 = \frac{1}{r}$ | 32. $2(3x - 1)^2 + 5(3x - 1) = -2$ | |

*This exercise requires knowledge of complex numbers.

8.4

Formulas and Further Applications

OBJECTIVES

- 1 Solve formulas involving squares and square roots for specified variables.
- 2 Solve applied problems using the Pythagorean theorem.
- 3 Solve applied problems using area formulas.
- 4 Solve applied problems using quadratic functions as models.


NOW TRY EXERCISE 1

Solve each formula for the specified variable. Keep \pm in the answer in part (a).

(a) $n = \frac{ab}{E^2}$ for E

(b) $S = \sqrt{\frac{pq}{n}}$ for p

NOW TRY ANSWERS

1. (a) $E = \frac{\pm \sqrt{abn}}{n}$ (b) $p = \frac{nS^2}{q}$

OBJECTIVE 1 Solve formulas involving squares and square roots for specified variables.

EXAMPLE 1 Solving for Specified Variables

Solve each formula for the specified variable. Keep \pm in the answer in part (a).

(a) $w = \frac{kFr}{v^2}$ for v The goal is to isolate v on one side.

$$v^2 w = kFr \quad \text{Multiply by } v^2.$$

$$v^2 = \frac{kFr}{w} \quad \text{Divide by } w.$$

$$v = \pm \sqrt{\frac{kFr}{w}} \quad \text{Square root property}$$

Include both positive and negative roots.

$$v = \frac{\pm \sqrt{kFr}}{\sqrt{w}} \cdot \frac{\sqrt{w}}{\sqrt{w}} \quad \text{Rationalize the denominator.}$$

$$v = \frac{\pm \sqrt{kFrw}}{w} \quad \begin{array}{l} \sqrt{a} \cdot \sqrt{b} = \sqrt{ab}; \\ \sqrt{a} \cdot \sqrt{a} = a \end{array}$$

(b) $d = \sqrt{\frac{4\mathcal{A}}{\pi}}$ for \mathcal{A} The goal is to isolate \mathcal{A} on one side.

$$d^2 = \frac{4\mathcal{A}}{\pi} \quad \text{Square each side.}$$

$$\pi d^2 = 4\mathcal{A} \quad \text{Multiply by } \pi.$$

$$\frac{\pi d^2}{4} = \mathcal{A}, \quad \text{or} \quad \mathcal{A} = \frac{\pi d^2}{4} \quad \begin{array}{l} \text{Divide by 4.} \\ \text{Interchange sides.} \end{array}$$

NOW TRY 

EXAMPLE 2 Solving for a Specified Variable

Solve $s = 2t^2 + kt$ for t .

Because the given equation has terms with t^2 and t , write it in standard form $ax^2 + bx + c = 0$, with t as the variable instead of x .

$$s = 2t^2 + kt$$

$$0 = 2t^2 + kt - s \quad \text{Subtract } s.$$

$$2t^2 + kt - s = 0 \quad \text{Standard form}$$

We can solve this equation using the quadratic formula.

NOW TRY
EXERCISE 2

Solve for r .

$$r^2 + 9r = -c$$

In the equation $2t^2 + kt - s = 0$, we have $a = 2$, $b = k$, and $c = -s$.

$$t = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{Quadratic formula}$$

$$t = \frac{-k \pm \sqrt{k^2 - 4(2)(-s)}}{2(2)} \quad \text{In } 2t^2 + kt - s = 0, \text{ } a = 2, b = k, \text{ and } c = -s.$$

$$t = \frac{-k \pm \sqrt{k^2 + 8s}}{4} \quad \text{Simplify.}$$

The solutions are $t = \frac{-k + \sqrt{k^2 + 8s}}{4}$ and $t = \frac{-k - \sqrt{k^2 + 8s}}{4}$.

NOW TRY

OBJECTIVE 2 Solve applied problems using the Pythagorean theorem.

The Pythagorean theorem is represented by the equation

$$a^2 + b^2 = c^2. \quad \text{See FIGURE 2.}$$




FIGURE 2

NOW TRY
EXERCISE 3

Matt is building a new barn, with length 10 ft more than width. While determining the footprint of the barn, he measured the diagonal as 50 ft. What will be the dimensions of the barn?

EXAMPLE 3 Using the Pythagorean Theorem

Two cars left an intersection at the same time, one heading due north, the other due west. Some time later, they were exactly 100 mi apart. The car headed north had gone 20 mi farther than the car headed west. How far had each car traveled?

Step 1 Read the problem carefully.

Step 2 Assign a variable.

Let x = the distance traveled by the car headed west.

Then $x + 20$ = the distance traveled by the car headed north.

See FIGURE 3. The cars are 100 mi apart, so the hypotenuse of the right triangle equals 100.




FIGURE 3

Step 3 Write an equation. Use the Pythagorean theorem.

$$a^2 + b^2 = c^2 \quad \text{Pythagorean theorem}$$

$$x^2 + (x + 20)^2 = 100^2 \quad \text{See FIGURE 3.}$$

$$(x + y)^2 = x^2 + 2xy + y^2$$

Step 4 Solve. $x^2 + x^2 + 40x + 400 = 10,000$ Square the binomial.

$$2x^2 + 40x - 9600 = 0 \quad \text{Standard form}$$

$$x^2 + 20x - 4800 = 0 \quad \text{Divide by 2.}$$

$$(x + 80)(x - 60) = 0 \quad \text{Factor.}$$

$$x + 80 = 0 \quad \text{or} \quad x - 60 = 0 \quad \text{Zero-factor property}$$

$$x = -80 \quad \text{or} \quad x = 60 \quad \text{Solve for } x.$$

Step 5 State the answer. Distance cannot be negative, so discard the negative solution. The required distances are 60 mi and $60 + 20 = 80$ mi.

Step 6 Check. Here $60^2 + 80^2 = 100^2$, as required.

NOW TRY

NOW TRY ANSWERS

2. $r = \frac{-9 \pm \sqrt{81 - 4c}}{2}$

3. 30 ft by 40 ft

NOW TRY EXERCISE 4

A football practice field is 30 yd wide and 40 yd long. A strip of grass sod of uniform width is to be placed around the perimeter of the practice field. There is enough money budgeted for 296 sq yd of sod. How wide will the strip be?

OBJECTIVE 3 Solve applied problems using area formulas.

EXAMPLE 4 Solving an Area Problem

A rectangular reflecting pool in a park is 20 ft wide and 30 ft long. The gardener wants to plant a strip of grass of uniform width around the edge of the pool. She has enough seed to cover 336 ft^2 . How wide will the strip be?




FIGURE 4

Step 1 Read the problem carefully.

Step 2 Assign a variable. The pool is shown in FIGURE 4.

Let x = the unknown width of the grass strip.

Then $20 + 2x$ = the width of the large rectangle (the width of the pool plus two grass strips),

and $30 + 2x$ = the length of the large rectangle.

Step 3 Write an equation. Refer to FIGURE 4.

$$(30 + 2x)(20 + 2x) \quad \text{Area of large rectangle (length} \cdot \text{width)}$$

$$30 \cdot 20, \text{ or } 600 \quad \text{Area of pool (in square feet)}$$

The area of the large rectangle minus the area of the pool should equal 336 ft^2 , the area of the grass strip.

$$\begin{array}{rcc} \text{Area} & & \text{area} & = & \text{area} \\ \text{of large} & - & \text{of} & & \text{of} \\ \text{rectangle} & & \text{pool} & & \text{grass.} \\ \downarrow & & \downarrow & & \downarrow \\ (30 + 2x)(20 + 2x) - 600 = 336 \end{array}$$

Step 4 Solve. $600 + 60x + 40x + 4x^2 - 600 = 336$ Multiply.

$$4x^2 + 100x - 336 = 0 \quad \text{Standard form}$$

$$x^2 + 25x - 84 = 0 \quad \text{Divide by 4.}$$

$$(x + 28)(x - 3) = 0 \quad \text{Factor.}$$

$$x + 28 = 0 \quad \text{or} \quad x - 3 = 0 \quad \text{Zero-factor property}$$

$$x = -28 \quad \text{or} \quad x = 3 \quad \text{Solve for } x.$$

Step 5 State the answer. The width cannot be -28 ft, so the grass strip should be 3 ft wide.

Step 6 Check. If $x = 3$, we can find the area of the large rectangle (which includes the grass strip).

$$(30 + 2 \cdot 3)(20 + 2 \cdot 3) = 36 \cdot 26 = 936 \text{ ft}^2 \quad \text{Area of pool and strip}$$

The area of the pool is $30 \cdot 20 = 600 \text{ ft}^2$, so the area of the grass strip is

$$936 - 600 = 336 \text{ ft}^2, \text{ as required.}$$

NOW TRY

NOW TRY ANSWER

4. 2 yd

NOW TRY
EXERCISE 5

If an object is projected upward from the top of a 120-ft building at 60 ft per second, its position (in feet above the ground) is given by

$$s(t) = -16t^2 + 60t + 120,$$

where t is time in seconds after it was projected. When does it hit the ground (to the nearest tenth)?

OBJECTIVE 4 Solve applied problems using quadratic functions as models.

Some applied problems can be modeled by *quadratic functions*, which for real numbers a , b , and c , can be written in the form

$$f(x) = ax^2 + bx + c \quad (\text{where } a \neq 0).$$

EXAMPLE 5 Solving an Applied Problem Using a Quadratic Function

If an object is projected upward from the top of a 144-ft building at 112 ft per second, its position (in feet above the ground) is given by

$$s(t) = -16t^2 + 112t + 144,$$

where t is time in seconds after it was projected. When does it hit the ground?

When the object hits the ground, its distance above the ground is 0. We must find the value of t that makes $s(t) = 0$.

$$s(t) = -16t^2 + 112t + 144$$

Given model

$$0 = -16t^2 + 112t + 144$$

Let $s(t) = 0$.

$$0 = t^2 - 7t - 9$$

Divide by -16 .

$$t = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Quadratic formula

$$t = \frac{-(-7) \pm \sqrt{(-7)^2 - 4(1)(-9)}}{2(1)}$$

Let $a = 1$, $b = -7$,
and $c = -9$.

$$t = \frac{7 \pm \sqrt{85}}{2}$$

Simplify.

$$t \approx \frac{7 \pm 9.2}{2}$$

Approximate the square root to the nearest tenth.

$$t \approx 8.1 \quad \text{or} \quad t \approx -1.1$$

Time cannot be negative, so discard -1.1 .

Find the two solutions.

The object will hit the ground about 8.1 sec after it is projected.

NOW TRY

EXAMPLE 6 Using a Quadratic Function to Model the CPI

The Consumer Price Index (CPI) is used to measure trends in prices for a “basket” of goods purchased by typical American families. This index uses a base period of 1982–1984, which means that the index number for that period is 100. The quadratic function

$$f(x) = -0.00833x^2 + 4.745x + 83.10$$

approximates the CPI for the years 1980–2016, where x is the number of years that have elapsed since 1980. (Data from Bureau of Labor Statistics.)

(a) Use the model to approximate the CPI for 2016.

For 2016, $x = 2016 - 1980 = 36$, so we find $f(36)$.

$$f(x) = -0.00833x^2 + 4.745x + 83.10$$

Given model

$$f(36) = -0.00833(36)^2 + 4.745(36) + 83.10$$

Let $x = 36$.

$$f(36) = 243$$

Nearest whole number

According to the model, the CPI for 2016 was 243.



NOW TRY ANSWER

5. 5.2 sec after it is projected


NOW TRY EXERCISE 6
Refer to **Example 6**.

- (a) Use the model to approximate the CPI for 2010, to the nearest whole number.
- (b) In what year did the CPI reach 175? (Round down for the year.)

- (b) In what year did the CPI reach 200?
Find the value of x that makes $f(x) = 200$.

$$f(x) = -0.00833x^2 + 4.745x + 83.10$$

$$200 = -0.00833x^2 + 4.745x + 83.10$$

$$0 = -0.00833x^2 + 4.745x - 116.9$$

$$x = \frac{-4.745 \pm \sqrt{4.745^2 - 4(-0.00833)(-116.9)}}{2(-0.00833)}$$

$$x = 25.8 \quad \text{or} \quad x = 543.8$$

Given model

Let $f(x) = 200$.

Subtract 200.

Use $a = -0.00833$, $b = 4.745$, and $c = -116.9$ in the quadratic formula.

Use a calculator. Round to the nearest tenth.


Rounding the first solution 25.8 down, the CPI first reached 200 in

$$1980 + 25 = 2005.$$

(Reject the solution $x = 543.8$, as this corresponds to a totally unreasonable year.)**NOW TRY ANSWERS**



6. (a) 218 (b) 2000

NOW TRY **8.4 Exercises**FOR
EXTRA
HELP**MyLab Math**

 Video solutions for select problems available in MyLab Math

Concept Check Answer each question.

- In solving a formula that has the specified variable in the denominator, what is the first step?
- What is the first step in solving a formula like $gw^2 = 2r$ for w ?
- What is the first step in solving a formula like $gw^2 = kw + 24$ for w ?
- Why is it particularly important to check all proposed solutions to an applied problem against the information in the original problem?

For each triangle, solve for m in terms of the other variables (where $m > 0$).Solve each formula for the specified variable. (Leave \pm in the answers as needed.) See **Examples 1 and 2**.

7. $d = kt^2$ for t

8. $S = 6e^2$ for e

9. $S = 4\pi r^2$ for r

10. $s = kwd^2$ for d

11. $I = \frac{ks}{d^2}$ for d

12. $R = \frac{k}{d^2}$ for d

13. $F = \frac{kA}{v^2}$ for v

14. $L = \frac{kd^4}{h^2}$ for h

15. $V = \frac{1}{3}\pi r^2 h$ for r

16. $V = \pi r^2 h$ for r

17. $At^2 + Bt = -C$ for t

18. $S = 2\pi rh + \pi r^2$ for r

19. $D = \sqrt{kh}$ for h

20. $F = \frac{k}{\sqrt{d}}$ for d

21. $p = \sqrt{\frac{k\ell}{g}}$ for ℓ

22. $p = \sqrt{\frac{k\ell}{g}}$ for g

Extending Skills Solve each equation for the specified variable. (Leave \pm in the answers.)

23. $p = \frac{E^2R}{(r+R)^2}$ for R (where $E > 0$)

24. $S(6S - t) = t^2$ for S

25. $10p^2c^2 + 7pcr = 12r^2$ for r


26. $S = vt + \frac{1}{2}gt^2$ for t

27. $LI^2 + RI + \frac{1}{c} = 0$ for I


28. $P = EI - RI^2$ for I

Solve each problem. When appropriate, round answers to the nearest tenth. See Example 3.


29. Find the lengths of the sides of the triangle.




30. Find the lengths of the sides of the triangle.



31. Two ships leave port at the same time, one heading due south and the other heading due east. Several hours later, they are 170 mi apart. If the ship traveling south traveled 70 mi farther than the other ship, how many miles did they each travel?




32. Deborah is flying a kite that is 30 ft farther above her hand than its horizontal distance from her. The string from her hand to the kite is 150 ft long. How high is the kite?



33. A game board is in the shape of a right triangle. The hypotenuse is 2 in. longer than the longer leg, and the longer leg is 1 in. less than twice as long as the shorter leg. How long is each side of the game board?
34. Manuel is planting a vegetable garden in the shape of a right triangle. The longer leg is 3 ft longer than the shorter leg, and the hypotenuse is 3 ft longer than the longer leg. Find the lengths of the three sides of the garden.
35. The diagonal of a rectangular rug measures 26 ft, and the length is 4 ft more than twice the width. Find the length and width of the rug.




36. A 13-ft ladder is leaning against a house. The distance from the bottom of the ladder to the house is 7 ft less than the distance from the top of the ladder to the ground. How far is the bottom of the ladder from the house?




Solve each problem. See Example 4.


37. A club swimming pool is 30 ft wide and 40 ft long. The club members want an exposed aggregate border in a strip of uniform width around the pool. They have enough material for 296 ft². How wide can the strip be?




38. Lyudmila wants to buy a rug for a room that is 20 ft long and 15 ft wide. She wants to leave an even strip of flooring uncovered around the edges of the room. How wide a strip will she have if she buys a rug with an area of 234 ft²?



39. A rectangle has a length 2 m less than twice its width. When 5 m are added to the width, the resulting figure is a square with an area of 144 m². Find the dimensions of the original rectangle.
40. Mariana's backyard measures 20 m by 30 m. She wants to put a flower garden in the middle of the yard, leaving a strip of grass of uniform width around the flower garden. Mariana must have 184 m² of grass. Under these conditions, what will the length and width of the garden be?
41. A rectangular piece of sheet metal has a length that is 4 in. less than twice the width. A square piece 2 in. on a side is cut from each corner. The sides are then turned up to form an uncovered box of volume 256 in.³. Find the length and width of the original piece of metal.



42. A rectangular piece of cardboard is 2 in. longer than it is wide. A square piece 3 in. on a side is cut from each corner. The sides are then turned up to form an uncovered box of volume 765 in.³. Find the dimensions of the original piece of cardboard.




Solve each problem. When appropriate, round answers to the nearest tenth. See Example 5.

43. An object is projected directly upward from the ground. After t seconds its distance in feet above the ground is

$$s(t) = 144t - 16t^2.$$

After how many seconds will the object be 128 ft above the ground? (*Hint:* Look for a common factor before solving the equation.)




44. When does the object in Exercise 43 strike the ground?

45. A ball is projected upward from the ground. Its distance in feet from the ground in t seconds is given by

$$s(t) = -16t^2 + 128t.$$


At what times will the ball be 213 ft from the ground?



46. A toy rocket is launched from ground level. Its distance in feet from the ground in t seconds is given by

$$s(t) = -16t^2 + 208t.$$


At what times will the rocket be 550 ft from the ground?



47. The following function gives the distance in feet a car going approximately 68 mph will skid in t seconds.

$$D(t) = 13t^2 - 100t$$

Find the time it would take for the car to skid 180 ft.



48. Refer to the function in **Exercise 47**. Find the time it would take for the car to skid 500 ft.

A ball is projected upward from ground level, and its distance in feet from the ground in t seconds is given by

$$s(t) = -16t^2 + 160t.$$

49. After how many seconds does the ball reach a height of 400 ft? Describe in words its position at this height.
50. After how many seconds does the ball reach a height of 425 ft? Interpret the mathematical result here.

Extending Skills Solve each problem using a quadratic equation.

51. A certain bakery has found that the daily demand for blueberry muffins is $\frac{6000}{p}$, where p is the price of a muffin in cents. The daily supply is $3p - 410$. Find the price at which supply and demand are equal.
52. In one area the demand for Blu-ray discs is $\frac{1900}{P}$ per day, where P is the price in dollars per disc. The supply is $5P - 1$ per day. At what price, to the nearest cent, does supply equal demand?
53. The formula $A = P(1 + r)^2$ gives the amount A in dollars that P dollars will grow to in 2 yr at interest rate r (where r is given as a decimal), using compound interest. What interest rate will cause \$2000 to grow to \$2142.45 in 2 yr?
54. Use the formula $A = P(1 + r)^2$ to find the interest rate r at which a principal P of \$10,000 will increase to \$10,920.25 in 2 yr.

William Froude was a 19th century naval architect who used the following expression, known as the **Froude number**, in shipbuilding.

$$\frac{v^2}{g\ell}$$

This expression was also used by R. McNeill Alexander in his research on dinosaurs. (Data from “How Dinosaurs Ran,” Scientific American.)

Use this expression to find the value of v (in meters per second), given $g = 9.8$ m per sec². (Round to the nearest tenth.)





55. Rhinoceros: $\ell = 1.2$;
Froude number = 2.57


56. Triceratops: $\ell = 2.8$;
Froude number = 0.16

Recall that corresponding sides of similar triangles are proportional. Use this fact to find the lengths of the indicated sides of each pair of similar triangles. Check all possible solutions in both triangles. Sides of a triangle cannot be negative (and are not drawn to scale here).

57. Side AC




58. Side RQ



Total spending (in billions of dollars) in the United States from all sources on physician and clinical services for the years 2000–2016 are shown in the bar graph. One model for the data is the quadratic function

$$f(x) = -0.2901x^2 + 25.90x + 291.6.$$

Here, $x = 0$ represents 2000, $x = 2$ represents 2002, and so on. Use the graph and the model to work each of the following exercises. See Example 6.



Data from Centers for Medicare and Medicaid Services.

59. Approximate spending on physician and clinical services in 2014 to the nearest \$10 billion using (a) the graph and (b) the model. How do the two approximations compare?
60. Repeat Exercise 59 for the year 2008.

61. According to the model, in what year did spending on physician and clinical services first exceed \$500 billion? (Round down for the year.)
62. Repeat **Exercise 61** for \$400 billion.

RELATING CONCEPTS For Individual or Group Work (Exercises 63–66)

In the 1939 classic movie *The Wizard of Oz*, Ray Bolger's character, the Scarecrow, wants a brain. When the Wizard grants him his "Th.D." (Doctor of Thinkology), the Scarecrow replies with the following statement.

Scarecrow: The sum of the square roots of any two sides of an isosceles triangle is equal to the square root of the remaining side.

His statement sounds like the formula for the Pythagorean theorem. To see why it is incorrect, **work Exercises 63–66 in order.**

63. To what kind of triangle does the Scarecrow refer in his statement? To what kind of triangle does the Pythagorean theorem actually refer?
64. In the Scarecrow's statement, he refers to square roots. In applying the formula for the Pythagorean theorem, do we find square roots of the sides? If not, what do we find?
65. An isosceles triangle has two sides of equal length. Draw an isosceles triangle with two sides of length 9 units and remaining side of length 4 units. Show that this triangle does not satisfy the Scarecrow's statement. (This is a *counterexample* and is sufficient to show that his statement is false in general.)
66. Use wording similar to that of the Scarecrow, but state the Pythagorean theorem correctly.



8.5 Graphs of Quadratic Functions

OBJECTIVES

- 1 Graph a quadratic function.
- 2 Graph parabolas with horizontal and vertical shifts.
- 3 Use the coefficient of x^2 to predict the shape and direction in which a parabola opens.
- 4 Find a quadratic function to model data.

OBJECTIVE 1 Graph a quadratic function.

FIGURE 5 gives a graph of the simplest *quadratic function* $y = x^2$. This graph is a **parabola**. The point $(0, 0)$, the lowest point on the curve, is the **vertex** of the parabola. The vertical line through the vertex is the **axis of symmetry**, or simply the **axis**, of the parabola. Here its equation is $x = 0$. A parabola is **symmetric about its axis**—that is, if the graph were folded along the axis, the two portions of the curve would coincide.

As **FIGURE 5** suggests, x can be any real number, so the domain of the function $y = x^2$, written in interval notation, is $(-\infty, \infty)$. Values of y are always nonnegative, so the range is $[0, \infty)$.




FIGURE 5

Quadratic Function

A function that can be written in the form

$$f(x) = ax^2 + bx + c$$

for real numbers a , b , and c , where $a \neq 0$, is a **quadratic function**.

Examples: $f(x) = x^2$, $f(x) = x^2 - 4x + 4$, $f(x) = -\frac{1}{2}x^2 + 3$


VOCABULARY

- parabola
- vertex
- axis of symmetry (axis)
- quadratic function

The graph of any quadratic function is a parabola with a vertical axis.

NOTE We use the variable y and function notation $f(x)$ interchangeably. Although we use the letter f most often to name quadratic functions, other letters can be used. We use the capital letter F to distinguish between different parabolas graphed on the same coordinate axes.

Parabolas have a special reflecting property that makes them useful in the design of telescopes, radar equipment, solar furnaces, and automobile headlights. (See FIGURE 6.)



Headlight

FIGURE 6


OBJECTIVE 2 Graph parabolas with horizontal and vertical shifts.

Parabolas need not have their vertices at the origin, as does the graph of $f(x) = x^2$.

EXAMPLE 1 Graphing a Parabola (Vertical Shift)

Graph $F(x) = x^2 - 2$.

If $x = 0$, then $F(x) = -2$, which gives the vertex $(0, -2)$. The graph of $F(x) = x^2 - 2$ has the same shape as that of $f(x) = x^2$ but is *shifted*, or *translated*, down 2 units. Every function value is 2 less than the corresponding function value of $f(x) = x^2$. Plotting points on both sides of the vertex gives the graph in FIGURE 7.



$$F(x) = x^2 - 2$$

Vertex: $(0, -2)$

Axis of symmetry: $x = 0$

Domain: $(-\infty, \infty)$

Range: $[-2, \infty)$

The graph of $f(x) = x^2$ is shown for comparison.

FIGURE 7


This parabola is symmetric about its axis $x = 0$, so the plotted points are “mirror images” of each other. Because x can be any real number, the domain is $(-\infty, \infty)$. The value of y (or $F(x)$) is always greater than or equal to -2 , so the range is $[-2, \infty)$.

NOW TRY

NOW TRY EXERCISE 1

Graph $f(x) = x^2 - 3$. Give the vertex, axis, domain, and range.

NOW TRY ANSWER



vertex: $(0, -3)$; axis: $x = 0$;

domain: $(-\infty, \infty)$; range: $[-3, \infty)$

Parabola with a Vertical Shift

The graph of $F(x) = x^2 + k$ is a parabola.

- The graph has the same shape as the graph of $f(x) = x^2$.
- The parabola is shifted up k units if $k > 0$, and down $|k|$ units if $k < 0$.
- The vertex of the parabola is $(0, k)$.

NOW TRY EXERCISE 2

Graph $f(x) = (x + 1)^2$.
Give the vertex, axis,
domain, and range.

EXAMPLE 2 Graphing a Parabola (Horizontal Shift)

Graph $F(x) = (x - 2)^2$.

If $x = 2$, then $F(x) = 0$, giving the vertex $(2, 0)$. The graph of


$$F(x) = (x - 2)^2$$

has the same shape as that of $f(x) = x^2$ but is shifted *to the right* 2 units. We plot several points on one side of the vertex and then use symmetry about the axis $x = 2$ to find corresponding points on the other side of the vertex.

If $x = 0$, for example, then

$$F(0) = (0 - 2)^2 = 4,$$

and the point $(0, 4)$ lies on the graph. The corresponding point two units on the “other” side of the axis $x = 2$ is the point $(4, 4)$, which also lies on the graph. See **FIGURE 8**.



$$F(x) = (x - 2)^2$$

Vertex: $(2, 0)$

Axis of symmetry: $x = 2$

Domain: $(-\infty, \infty)$

Range: $[0, \infty)$

FIGURE 8

NOW TRY

Parabola with a Horizontal Shift

The graph of $F(x) = (x - h)^2$ is a parabola.

- The graph has the same shape as the graph of $f(x) = x^2$.
- The parabola is shifted to the right h units if $h > 0$, and to the left $|h|$ units if $h < 0$.
- The vertex of the parabola is $(h, 0)$.

CAUTION *Errors frequently occur when horizontal shifts are involved.* To determine the direction and magnitude of a horizontal shift, find the value that causes the expression $x - h$ to equal 0, as shown below.

$$F(x) = (x - 5)^2$$

Because $+5$ causes $x - 5$ to equal 0, the graph of $F(x)$ illustrates a shift


to the right 5 units.

$$F(x) = (x + 5)^2$$

Because -5 causes $x + 5$ to equal 0, the graph of $F(x)$ illustrates a shift

to the left 5 units.

NOW TRY ANSWER



vertex: $(-1, 0)$; axis: $x = -1$;
domain: $(-\infty, \infty)$; range: $[0, \infty)$


NOW TRY EXERCISE 3

Graph $f(x) = (x + 1)^2 - 2$.
Give the vertex, axis, domain, and range.

EXAMPLE 3 Graphing a Parabola (Horizontal and Vertical Shifts)

Graph $F(x) = (x + 3)^2 - 2$.

This graph has the same shape as that of $f(x) = x^2$, but is shifted *to the left* 3 units (because $x + 3 = 0$ if $x = -3$) and *down* 2 units (because of the negative sign in -2). This gives the vertex $(-3, -2)$. Find and plot several ordered pairs, using symmetry as needed, to obtain the graph in **FIGURE 9**.



$F(x) = (x + 3)^2 - 2$
Vertex: $(-3, -2)$
Axis of symmetry: $x = -3$
Domain: $(-\infty, \infty)$
Range: $[-2, \infty)$

FIGURE 9

NOW TRY

Parabola with Horizontal and Vertical Shifts

The graph of $F(x) = (x - h)^2 + k$ is a parabola.

- The graph has the same shape as the graph of $f(x) = x^2$.
- The vertex of the parabola is (h, k) .
- The axis of symmetry is the vertical line $x = h$.

OBJECTIVE 3 Use the coefficient of x^2 to predict the shape and direction in which a parabola opens.

Not all parabolas open up, and not all parabolas have the same shape as the graph of $f(x) = x^2$.

NOW TRY EXERCISE 4

Graph $f(x) = -3x^2$.
Give the vertex, axis, domain, and range.

EXAMPLE 4 Graphing a Parabola That Opens Down


Graph $f(x) = -\frac{1}{2}x^2$.

This parabola is shown in **FIGURE 10**. The coefficient of x^2 , $-\frac{1}{2}$, affects the shape of the graph—the $\frac{1}{2}$ makes the parabola wider (because the values of $\frac{1}{2}x^2$ increase more slowly than those of x^2), and the negative sign makes the parabola open down.

The graph is not shifted in any direction, so the vertex is $(0, 0)$. Unlike the parabolas graphed in **Examples 1–3**, the vertex $(0, 0)$ has the *greatest* function value of any point on the graph.


NOW TRY ANSWERS

3.



vertex: $(-1, -2)$;
axis: $x = -1$;
domain: $(-\infty, \infty)$;
range: $[-2, \infty)$


4.



vertex: $(0, 0)$;
axis: $x = 0$;
domain: $(-\infty, \infty)$;
range: $(-\infty, 0]$

$f(x) = -\frac{1}{2}x^2$

x	f(x)
-2	-2
-1	$-\frac{1}{2}$
0	0
1	$-\frac{1}{2}$
2	-2



$f(x) = -\frac{1}{2}x^2$
Vertex: $(0, 0)$
Axis of symmetry: $x = 0$
Domain: $(-\infty, \infty)$
Range: $(-\infty, 0]$

FIGURE 10

NOW TRY

General Characteristics of the Graph of a Vertical Parabola

The graph of the quadratic function $F(x) = a(x - h)^2 + k$ (where $a \neq 0$) is a parabola.

- The vertex of the parabola is (h, k) .
- The axis of symmetry is the vertical line $x = h$.
- The graph opens up if $a > 0$ and down if $a < 0$.
- The graph is wider than that of $f(x) = x^2$ if $0 < |a| < 1$.
The graph is narrower than that of $f(x) = x^2$ if $|a| > 1$.

NOW TRY EXERCISE 5

Graph $f(x) = 2(x - 1)^2 + 2$.
Give the vertex, axis, domain, and range.

EXAMPLE 5 Using the General Characteristics to Graph a Parabola


Graph $F(x) = -2(x + 3)^2 + 4$.

The parabola opens down (because $a = -2$ and $-2 < 0$) and is narrower than the graph of $f(x) = x^2$ (because $|-2| = 2$ and $2 > 1$). This causes values of $F(x)$ to decrease more quickly than those of $f(x) = -x^2$.

The graph is shifted *to the left* 3 units (because $x + 3 = 0$ when $x = -3$) and *up* 4 units (because of the $+4$), which gives the vertex $(-3, 4)$. To complete the graph, we plotted the ordered pairs $(-4, 2)$ and, by symmetry, $(-2, 2)$. Symmetry can be used to find additional ordered pairs that satisfy the equation. See FIGURE 11.

$$F(x) = -2(x + 3)^2 + 4$$

x	$F(x)$
-5	-4
-4	2
-3	4
-2	2
-1	-4



$$F(x) = -2(x + 3)^2 + 4$$

Vertex: $(-3, 4)$

Axis of symmetry: $x = -3$

Domain: $(-\infty, \infty)$

Range: $(-\infty, 4]$

FIGURE 11

NOW TRY

OBJECTIVE 4 Find a quadratic function to model data.

EXAMPLE 6 Modeling the Number of Multiple Births


The number of higher-order multiple births (triplets or more) in the United States is shown in the table. Here, x represents the number of years since 1996, and y represents the number of higher-order multiple births (to the nearest hundred).

Year	x	y
1996	0	5900
2000	4	7300
2002	6	7400
2004	8	7300
2006	10	6500
2008	12	6300
2010	14	5500
2012	16	4900
2014	18	4500



Data from National Center for Health Statistics.

NOW TRY ANSWER



$$f(x) = 2(x - 1)^2 + 2$$

vertex: $(1, 2)$; axis: $x = 1$;

domain: $(-\infty, \infty)$; range: $[2, \infty)$


NOW TRY EXERCISE 6

Using the points $(0, 5900)$, $(4, 7300)$, and $(12, 6300)$, find another quadratic model for the data on higher-order multiple births in **Example 6**. (Round values of a and b to the nearest tenth.)

Find a quadratic function that models the data in the table on the previous page.

A scatter diagram of the ordered pairs (x, y) is shown in **FIGURE 12**. The general shape suggested by the scatter diagram indicates that a parabola should approximate these points, as shown by the dashed curve in **FIGURE 13**. The equation for such a parabola would have a negative coefficient for x^2 because the graph opens down.




FIGURE 12




FIGURE 13

To find a quadratic function of the form

$$y = ax^2 + bx + c$$

that models, or *fits*, these data, we choose three representative ordered pairs from the table and use them to write a system of three equations.

$$(0, 5900), \quad (6, 7400), \quad \text{and} \quad (12, 6300) \quad \text{Three ordered pairs } (x, y)$$

We substitute the x - and y -values from each ordered pair into the quadratic form $y = ax^2 + bx + c$ to obtain three equations.

$$a(0)^2 + b(0) + c = 5900 \quad \xrightarrow{\text{Simplify.}} \quad c = 5900 \quad (1)$$

$$a(6)^2 + b(6) + c = 7400 \quad \longrightarrow \quad 36a + 6b + c = 7400 \quad (2)$$

$$a(12)^2 + b(12) + c = 6300 \quad \longrightarrow \quad 144a + 12b + c = 6300 \quad (3)$$

To find the values of a , b , and c , we solve this system of three equations in three variables. From equation (1), $c = 5900$, so we substitute 5900 for c in equations (2) and (3) to obtain two equations in two variables.

$$36a + 6b + 5900 = 7400 \quad \longrightarrow \quad 36a + 6b = 1500 \quad (4)$$

$$144a + 12b + 5900 = 6300 \quad \xrightarrow{\text{Subtract 5900.}} \quad 144a + 12b = 400 \quad (5)$$

We eliminate b from this system of equations in two variables by multiplying equation (4) by -2 and adding the results to equation (5).

$$-72a - 12b = -3000 \quad \text{Multiply equation (4) by } -2.$$

$$144a + 12b = 400 \quad (5)$$

$$72a = -2600 \quad \text{Add.}$$

$$a \approx -36.1 \quad \text{Use a calculator. Round to one decimal place.}$$

We substitute -36.1 for a in equation (4) to find that $b \approx 466.6$. (Substituting in equation (5) will give $b \approx 466.5$ due to rounding procedures.) Using the values we found for a , b , and c , the model is

$$y = \underset{\substack{a \\ \downarrow}}{-36.1}x^2 + \underset{\substack{b \\ \downarrow}}{466.6}x + \underset{\substack{c \\ \downarrow}}{5900}.$$

NOW TRY ANSWER

6. $y = -39.6x^2 + 508.4x + 5900$
(Answers may vary slightly due to rounding.)

NOTE If we had chosen three different ordered pairs of data in **Example 6**, a slightly different, though similar, model would have resulted. (See **Now Try Exercise 6**.)

The *quadratic regression* feature on a graphing calculator can also be used to generate the quadratic model that best fits given data. See your owner's manual for details.

8.5 Exercises

FOR
EXTRA
HELP


MyLab Math

▶ Video solutions for select problems available in MyLab Math


Concept Check Match each quadratic function in parts (a)–(d) with its graph from choices A–D.

1. (a) $f(x) = (x + 2)^2 - 1$

A.




B.




(b) $f(x) = (x + 2)^2 + 1$

(c) $f(x) = (x - 2)^2 - 1$

C.




D.




(d) $f(x) = (x - 2)^2 + 1$

2. (a) $f(x) = -x^2 + 2$

A.




B.




(b) $f(x) = -x^2 - 2$

(c) $f(x) = -(x + 2)^2$

C.



D.



(d) $f(x) = -(x - 2)^2$

3. **CONCEPT CHECK** Match each quadratic function in Column I with the description of the parabola that is its graph in Column II.

I

- (a) $f(x) = (x - 4)^2 - 2$
 (b) $f(x) = (x - 2)^2 - 4$
 (c) $f(x) = (x + 4)^2 + 2$
 (d) $f(x) = -(x - 4)^2 - 2$
 (e) $f(x) = -(x - 2)^2 - 4$
 (f) $f(x) = -(x - 4)^2 + 2$

II

- A. Vertex $(2, -4)$, opens down
 B. Vertex $(2, -4)$, opens up
 C. Vertex $(4, -2)$, opens down
 D. Vertex $(4, -2)$, opens up
 E. Vertex $(4, 2)$, opens down
 F. Vertex $(-4, 2)$, opens up

4. Concept Check For the quadratic function $f(x) = a(x - h)^2 + k$, in what quadrant is the vertex if the values of h and k are as follows?

- (a) $h > 0, k > 0$ (b) $h > 0, k < 0$
 (c) $h < 0, k > 0$ (d) $h < 0, k < 0$

Consider the value of a , and make the correct choice.

- (e) If $a > 0$, then the graph opens (*up / down*).
 (f) If $a < 0$, then the graph opens (*up / down*).
 (g) If $|a| > 1$, then the graph is (*narrower / wider*) than the graph of $f(x) = x^2$.
 (h) If $0 < |a| < 1$, then the graph is (*narrower / wider*) than the graph of $f(x) = x^2$.

Identify the vertex of each parabola. See Examples 1–4.

5. $f(x) = -3x^2$ 6. $f(x) = -4x^2$ 7. $f(x) = \frac{1}{3}x^2$
 8. $f(x) = \frac{1}{2}x^2$ 9. $f(x) = x^2 + 4$ 10. $f(x) = x^2 - 4$
 11. $f(x) = (x - 1)^2$ 12. $f(x) = (x + 3)^2$ 13. $f(x) = (x + 3)^2 - 4$
 14. $f(x) = (x + 5)^2 - 8$ 15. $f(x) = -(x - 5)^2 + 6$ 16. $f(x) = -(x - 2)^2 + 1$

For each quadratic function, tell whether the graph opens up or down and whether the graph is wider, narrower, or the same shape as the graph of $f(x) = x^2$. See Examples 4 and 5.


17. $f(x) = -\frac{2}{5}x^2$ 18. $f(x) = -2x^2$
 19. $f(x) = 3x^2 + 1$ 20. $f(x) = \frac{2}{3}x^2 - 4$
 21. $f(x) = -4(x + 2)^2 + 5$ 22. $f(x) = -\frac{1}{3}(x + 6)^2 + 3$

Graph each parabola. Give the vertex, axis of symmetry, domain, and range. See Examples 1–5.

23. $f(x) = 3x^2$ 24. $f(x) = \frac{1}{2}x^2$
 25. $f(x) = -2x^2$ 26. $f(x) = -\frac{1}{3}x^2$
 27. $f(x) = x^2 - 1$ 28. $f(x) = x^2 + 3$
 29. $f(x) = -x^2 + 2$ 30. $f(x) = -x^2 - 2$
 31. $f(x) = (x - 4)^2$ 32. $f(x) = (x + 1)^2$
 33. $f(x) = (x + 2)^2 - 1$ 34. $f(x) = (x - 1)^2 + 2$
 35. $f(x) = (x - 1)^2 - 3$ 36. $f(x) = (x + 1)^2 + 1$
 37. $f(x) = 2(x - 2)^2 - 4$ 38. $f(x) = 3(x - 2)^2 + 1$
 39. $f(x) = \frac{1}{2}(x - 2)^2 - 3$ 40. $f(x) = \frac{4}{3}(x - 3)^2 - 2$
 41. $f(x) = -2(x + 3)^2 + 4$ 42. $f(x) = -2(x - 2)^2 - 3$
 43. $f(x) = -\frac{1}{2}(x + 1)^2 + 2$ 44. $f(x) = -\frac{2}{3}(x + 2)^2 + 1$


Determine whether a linear function or a quadratic function would be a more appropriate model for each set of graphed data. If linear, tell whether the slope should be positive or negative. If quadratic, tell whether the coefficient of x^2 should be positive or negative. See Example 6.

45. Time Spent Playing Video Games




Data from www.statista.com

46. Average Daily Volume of First-Class Mail




Data from USPS.

47. Food Assistance Spending in Iowa




Data from Iowa DHS.

48. U.S. Foreign-Born Population




Data from U.S. Census Bureau.

49. High School Students Who Smoke



Data from www.cdc.gov

50. Social Security Assets



Data from SSA.

Solve each problem. See Example 6.

- 51.** The number of U.S. households (in millions) with cable television service is shown in the table, where x represents the number of years since 2007 and y represents households.

Year	x	y
2007	0	95
2009	2	103
2011	4	105
2013	6	103
2015	8	100
2017	10	96

Data from Nielsen Media Research.



- Use the ordered pairs (x, y) to make a scatter diagram of the data.
- Would a linear or a quadratic function better model the data?
- Should the coefficient a of x^2 in a quadratic model $y = ax^2 + bx + c$ be positive or negative?
- Use the ordered pairs $(0, 95)$, $(4, 105)$, and $(10, 96)$ to find a quadratic function that models the data.
- Use the model from part (d) to approximate the number of U.S. households (in millions) with cable television service in 2009 and 2015. How well does the model approximate the actual data from the table?

52. The number (in thousands) of new, privately owned housing units completed in the United States is shown in the table. Here x represents the number of years since 2006, and y represents total housing units completed.

Year	x	y
2006	0	1980
2008	2	1120
2010	4	650
2012	6	650
2014	8	880
2016	10	1060

Data from U.S. Census Bureau.



- Use the ordered pairs (x, y) to make a scatter diagram of the data.
- Would a linear or a quadratic function better model the data?
- Should the coefficient a of x^2 in a quadratic model $y = ax^2 + bx + c$ be positive or negative?
- Use the ordered pairs $(0, 1980)$, $(4, 650)$, and $(8, 880)$ to find a quadratic function that models the data.
- Use the model from part (d) to approximate the number of such housing units (in thousands) completed in 2008 and 2012. How well does the model approximate the actual data from the table?

8.6

More about Parabolas and Their Applications

OBJECTIVES

- Find the vertex of a vertical parabola.
- Graph a quadratic function.
- Use the discriminant to find the number of x -intercepts of a parabola with a vertical axis.
- Use quadratic functions to solve problems involving maximum or minimum value.
- Graph parabolas with horizontal axes.



NOW TRY EXERCISE 1

Find the vertex of the graph of

$$f(x) = x^2 + 2x - 8.$$

NOW TRY ANSWER

- $(-1, -9)$

OBJECTIVE 1 Find the vertex of a vertical parabola.

When the equation of a parabola is given in the form $f(x) = ax^2 + bx + c$, there are two ways to locate the vertex.

- Complete the square. (See **Examples 1 and 2**.)
- Use a formula derived by completing the square. (See **Example 3**.)

EXAMPLE 1 Completing the Square to Find the Vertex ($a = 1$)

Find the vertex of the graph of $f(x) = x^2 - 4x + 5$.

We can express $x^2 - 4x + 5$ in the form $(x - h)^2 + k$ by completing the square on $x^2 - 4x$, as in the first section of this chapter. The process is slightly different here because we want to keep $f(x)$ alone on one side of the equation. Instead of adding the appropriate number to each side, we *add and subtract* it on the right.

$$f(x) = x^2 - 4x + 5$$

$$f(x) = (x^2 - 4x \quad \quad) + 5$$

This is equivalent to adding 0. $\rightarrow \left[\frac{1}{2}(-4)\right]^2 = (-2)^2 = 4$

$$f(x) = (x^2 - 4x + 4 - 4) + 5$$

$$f(x) = (x^2 - 4x + 4) - 4 + 5$$

$$f(x) = (x - 2)^2 + 1$$

Group the variable terms.

Square half the coefficient of the first-degree term.

Add and subtract 4.

Bring -4 outside the parentheses.

Factor. Combine like terms.

The vertex of this parabola is $(2, 1)$.


**NOW TRY
EXERCISE 2**

Find the vertex of the graph of

$$f(x) = -4x^2 + 16x - 10.$$

EXAMPLE 2 Completing the Square to Find the Vertex ($a \neq 1$)
Find the vertex of the graph of $f(x) = -3x^2 + 6x - 1$.Because the x^2 -term has a coefficient other than 1, we factor that coefficient out of the first two terms before completing the square.

$$f(x) = -3x^2 + 6x - 1$$

$$f(x) = (-3x^2 + 6x) - 1$$

Group the variable terms.

$$f(x) = -3(x^2 - 2x) - 1$$

Factor out -3 .

$$f(x) = -3(x^2 - 2x \quad \quad) - 1$$

Prepare to complete the square.

$$\rightarrow \left[\frac{1}{2}(-2)\right]^2 = (-1)^2 = 1$$

Square half the coefficient of the first-degree term.

$$f(x) = -3(x^2 - 2x + 1 - 1) - 1$$

Add and subtract 1.

Now bring -1 outside the parentheses. Be sure to multiply it by -3 .

This is a key step.

$$f(x) = -3(x^2 - 2x + 1) + (-3)(-1) - 1 \quad \text{Distributive property}$$

$$f(x) = -3(x^2 - 2x + 1) + 3 - 1$$

Multiply.

$$f(x) = -3(x - 1)^2 + 2$$

Factor. Combine like terms.

The vertex is $(1, 2)$.
NOW TRY 
We can complete the square to derive a formula for the vertex of the graph of the quadratic function $f(x) = ax^2 + bx + c$ (where $a \neq 0$).

$$f(x) = ax^2 + bx + c$$

Standard form

$$f(x) = (ax^2 + bx) + c$$

Group the terms with x .

$$f(x) = a\left(x^2 + \frac{b}{a}x\right) + c$$

Factor a from the first two terms.

$$\rightarrow \left[\frac{1}{2}\left(\frac{b}{a}\right)\right]^2 = \left(\frac{b}{2a}\right)^2 = \frac{b^2}{4a^2}$$

Square half the coefficient of the first-degree term.

$$f(x) = a\left(x^2 + \frac{b}{a}x + \frac{b^2}{4a^2} - \frac{b^2}{4a^2}\right) + c$$

Add and subtract $\frac{b^2}{4a^2}$.

$$f(x) = a\left(x^2 + \frac{b}{a}x + \frac{b^2}{4a^2}\right) + a\left(-\frac{b^2}{4a^2}\right) + c$$

Distributive property

$$f(x) = a\left(x^2 + \frac{b}{a}x + \frac{b^2}{4a^2}\right) - \frac{b^2}{4a} + c$$

$$-\frac{ab^2}{4a^2} = -\frac{b^2}{4a}$$

$$f(x) = a\left(x + \frac{b}{2a}\right)^2 + \frac{4ac - b^2}{4a}$$

Factor. Rewrite terms with a common denominator.

$$f(x) = a\left[\underbrace{x - \left(\frac{-b}{2a}\right)}_h\right]^2 + \underbrace{\frac{4ac - b^2}{4a}}_k$$

 $f(x) = a(x - h)^2 + k$;
The vertex (h, k) can be expressed in terms of $a, b,$ and c .

NOW TRY ANSWER
2. $(2, 6)$

The expression for k can be found by replacing x with $\frac{-b}{2a}$. Using function notation, if $y = f(x)$, then the y -value of the vertex is $f\left(\frac{-b}{2a}\right)$.

Vertex Formula

The graph of the quadratic function $f(x) = ax^2 + bx + c$ (where $a \neq 0$) has vertex

$$\left(\frac{-b}{2a}, f\left(\frac{-b}{2a}\right) \right).$$

The axis of symmetry of the parabola is the line having equation

$$x = \frac{-b}{2a}.$$

 **NOW TRY EXERCISE 3**

Use the vertex formula to find the vertex of the graph of

$$f(x) = 3x^2 - 2x + 8.$$

EXAMPLE 3 Using the Formula to Find the Vertex

Use the vertex formula to find the vertex of the graph of $f(x) = x^2 - x - 6$.

The x -coordinate of the vertex of the parabola is given by $\frac{-b}{2a}$.

$$\frac{-b}{2a} = \frac{-(-1)}{2(1)} = \frac{1}{2} \leftarrow \begin{array}{l} a = 1, b = -1, \text{ and } c = -6 \\ x\text{-coordinate of vertex} \end{array}$$

The y -coordinate of the vertex of $f(x) = x^2 - x - 6$ is $f\left(\frac{-b}{2a}\right) = f\left(\frac{1}{2}\right)$.

$$f\left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)^2 - \frac{1}{2} - 6 = \frac{1}{4} - \frac{1}{2} - 6 = -\frac{25}{4} \leftarrow \begin{array}{l} \frac{1}{4} - \frac{1}{2} - 6 = \frac{1}{4} - \frac{2}{4} - \frac{24}{4} \\ y\text{-coordinate of vertex} \end{array}$$

The vertex is $\left(\frac{1}{2}, -\frac{25}{4}\right)$.

NOW TRY 

OBJECTIVE 2 Graph a quadratic function.Graphing a Quadratic Function $y = f(x)$

Step 1 Determine whether the graph opens up or down.

- If $a > 0$, then the parabola opens up.
- If $a < 0$, then it opens down.

Step 2 Find the vertex. Use the vertex formula or complete the square.

Step 3 Find any intercepts.

- To find the x -intercepts (if any), solve $f(x) = 0$.
- To find the y -intercept, evaluate $f(0)$.

Step 4 Complete the graph. Plot the points found so far. Find and plot additional points as needed, using symmetry about the axis.

EXAMPLE 4 Graphing a Quadratic Function

Graph the quadratic function $f(x) = x^2 - x - 6$.

Step 1 From the equation, $a = 1$, so the graph of the function opens up.

Step 2 The vertex, $\left(\frac{1}{2}, -\frac{25}{4}\right)$, was found in **Example 3** using the vertex formula.

NOW TRY ANSWER

3. $\left(\frac{1}{3}, \frac{23}{3}\right)$

NOW TRY
EXERCISE 4

Graph the quadratic function

$$f(x) = x^2 + 2x - 3.$$

Give the vertex, axis, domain, and range.

Step 3 Find any intercepts. The vertex, $(\frac{1}{2}, -\frac{25}{4})$, is in quadrant IV and the graph opens up, so there will be two x -intercepts. Let $f(x) = 0$ and solve to find them.

$$f(x) = x^2 - x - 6$$

$$0 = x^2 - x - 6$$

Let $f(x) = 0$.

$$0 = (x - 3)(x + 2)$$

Factor.

$$x - 3 = 0 \quad \text{or} \quad x + 2 = 0$$

Zero-factor property

$$x = 3 \quad \text{or} \quad x = -2$$

Solve each equation.

The x -intercepts are $(3, 0)$ and $(-2, 0)$.

Find the y -intercept by evaluating $f(0)$.

$$f(x) = x^2 - x - 6$$

$$f(0) = 0^2 - 0 - 6$$

Let $x = 0$.

$$f(0) = -6$$

Apply the exponent. Subtract.

The y -intercept is $(0, -6)$.

Step 4 Plot the points found so far and additional points as needed using symmetry about the axis, $x = \frac{1}{2}$. The graph is shown in **FIGURE 14**.




FIGURE 14

$$f(x) = x^2 - x - 6$$

$$\text{Vertex: } \left(\frac{1}{2}, -\frac{25}{4}\right)$$

$$\text{Axis of symmetry: } x = \frac{1}{2}$$

$$\text{Domain: } (-\infty, \infty)$$

$$\text{Range: } \left[-\frac{25}{4}, \infty\right)$$

NOW TRY

OBJECTIVE 3 Use the discriminant to find the number of x -intercepts of a parabola with a vertical axis.

Recall that the expression under the radical in $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, that is,


$$b^2 - 4ac, \quad \text{Discriminant}$$

is the *discriminant* of the quadratic equation $ax^2 + bx + c = 0$ and that we can use it to determine the number of real solutions of a quadratic equation.

In a similar way, we can use the discriminant of a quadratic function to determine the number of x -intercepts of its graph. The three possibilities are shown in **FIGURE 15** on the next page.

1. If the discriminant is positive, the parabola will have two x -intercepts.
2. If the discriminant is 0, there will be only one x -intercept, and it will be the vertex of the parabola.
3. If the discriminant is negative, the graph will have no x -intercepts.

NOW TRY ANSWER



vertex: $(-1, -4)$; axis: $x = -1$;
domain: $(-\infty, \infty)$; range: $[-4, \infty)$




FIGURE 15

NOW TRY EXERCISE 5

Find the discriminant and use it to determine the number of x -intercepts of the graph of each quadratic function.

(a) $f(x) = -2x^2 + 3x - 2$

(b) $f(x) = 3x^2 + 2x - 1$

(c) $f(x) = 4x^2 - 12x + 9$

EXAMPLE 5 Using the Discriminant to Determine Number of x -Intercepts

Find the discriminant and use it to determine the number of x -intercepts of the graph of each quadratic function.

(a) $f(x) = 2x^2 + 3x - 5$

$$b^2 - 4ac$$

$$= 3^2 - 4(2)(-5)$$

$$= 9 - (-40)$$

$$= 49$$

Discriminant

$$a = 2, b = 3, c = -5$$

Apply the exponent. Multiply.

Subtract.

Because the discriminant is positive, the parabola has two x -intercepts.

(b) $f(x) = -3x^2 - 1$ (which can be written as $f(x) = -3x^2 + 0x - 1$)

$$b^2 - 4ac$$

$$= 0^2 - 4(-3)(-1)$$

$$= 0 - 12$$

$$= -12$$

$$a = -3, b = 0, c = -1$$

Apply the exponent. Multiply.

Subtract.

The discriminant is negative, so the graph has no x -intercepts.

(c) $f(x) = 9x^2 + 6x + 1$

$$b^2 - 4ac$$

$$= 6^2 - 4(9)(1)$$

$$= 36 - 36$$

$$= 0$$

$$a = 9, b = 6, c = 1$$

Apply the exponent. Multiply.

Subtract.

Because the discriminant is 0, the parabola has only one x -intercept (its vertex).

NOW TRY

OBJECTIVE 4 Use quadratic functions to solve problems involving maximum or minimum value.

The vertex of the graph of a quadratic function is either the highest or the lowest point on the parabola. It provides the following information.

1. The y -value of the vertex gives the maximum or minimum value of y .
2. The x -value tells where the maximum or minimum occurs.

NOW TRY ANSWERS

5. (a) -7 ; none (b) 16 ; two
(c) 0 ; one

NOW TRY
EXERCISE 6

Solve the problem in **Example 6** if the farmer has only 80 ft of fencing.

EXAMPLE 6 Finding Maximum Area

A farmer has 120 ft of fencing to enclose a rectangular area next to a building. (See **FIGURE 16**.) Find the maximum area he can enclose and the dimensions of the field when the area is maximized.

Let x = the width of the rectangle.

$$x + x + \text{length} = 120$$

Sum of the sides is 120 ft.

$$2x + \text{length} = 120$$

Combine like terms.

$$\text{length} = 120 - 2x \quad \text{Subtract } 2x.$$




FIGURE 16

The area $\mathcal{A}(x)$ is given by the product of the length and width.

$$\mathcal{A}(x) = (120 - 2x)x \quad \text{Area} = \text{length} \cdot \text{width}$$

$$\mathcal{A}(x) = 120x - 2x^2 \quad \text{Distributive property}$$

$$\mathcal{A}(x) = -2x^2 + 120x \quad \text{Standard form}$$

The graph of $\mathcal{A}(x) = -2x^2 + 120x$ is a parabola that opens down. To determine the maximum area, use the vertex formula to find the vertex of the parabola.

$$x = \frac{-b}{2a} = \frac{-120}{2(-2)} = \frac{-120}{-4} = 30 \quad \text{Vertex formula; } a = -2, b = 120, c = 0$$

$$\mathcal{A}(30) = -2(30)^2 + 120(30) = -2(900) + 3600 = 1800$$

The vertex is $(30, 1800)$. The maximum area will be 1800 ft² when x , the width, is 30 ft and the length is

$$120 - 2(30) = 60 \text{ ft.}$$

NOW TRY

! CAUTION *Be careful when interpreting the meanings of the coordinates of the vertex.* The first coordinate, x , gives the value for which the *function value*, y or $f(x)$, is a maximum or a minimum.

Read a problem carefully to determine whether to find the value of the independent variable, the function value, or both.

EXAMPLE 7 Finding Maximum Height

If air resistance is neglected, a projectile on Earth shot straight upward with an initial velocity of 40 m per sec will be at a height s in meters given by

$$s(t) = -4.9t^2 + 40t,$$

where t is the number of seconds elapsed after projection. After how many seconds will it reach its maximum height, and what is this maximum height?

For this function, $a = -4.9$, $b = 40$, and $c = 0$. Use the vertex formula.

$$t = \frac{-b}{2a} = \frac{-40}{2(-4.9)} \approx 4.1 \quad \text{Use a calculator. Round to the nearest tenth.}$$

NOW TRY ANSWER

6. The field should be 20 ft by 40 ft with maximum area 800 ft².

This indicates that the maximum height is attained at 4.1 sec.

NOW TRY EXERCISE 7

A stomp rocket is launched from the ground with an initial velocity of 48 ft per sec so that its distance in feet above the ground after t seconds is

$$s(t) = -16t^2 + 48t.$$

Find the maximum height attained by the rocket and the number of seconds it takes to reach that height.

To find the maximum height at 4.1 sec, calculate $s(4.1)$.

$$\begin{aligned} s(t) &= -4.9t^2 + 40t \\ s(4.1) &= -4.9(4.1)^2 + 40(4.1) && \text{Let } t = 4.1. \\ s(4.1) &\approx 81.6 && \text{Use a calculator. Round to the nearest tenth.} \end{aligned}$$

The projectile will attain a maximum height of 81.6 m at 4.1 sec.

NOW TRY

PROBLEM-SOLVING HINT Many applied problems ask for the greatest or least value of some quantity. When we can express that quantity in terms of a quadratic function, the value of k in the vertex (h, k) gives that optimum value.

OBJECTIVE 5 Graph parabolas with horizontal axes.

If x and y are interchanged in the equation

$$y = ax^2 + bx + c, \quad \text{the equation becomes} \quad x = ay^2 + by + c.$$

Because of the interchange of the roles of x and y , these parabolas are horizontal (with horizontal lines as axes of symmetry).

General Characteristics of the Graph of a Horizontal Parabola

The graph of an equation of the form

$$x = ay^2 + by + c \quad \text{or} \quad x = a(y - k)^2 + h$$

is a horizontal parabola.

- The vertex of the parabola is (h, k) .
- The axis of symmetry is the horizontal line $y = k$.
- The graph opens to the right if $a > 0$ and to the left if $a < 0$.

NOW TRY EXERCISE 8

Graph $x = (y + 2)^2 - 1$.
Give the vertex, axis, domain, and range.

EXAMPLE 8 Graphing a Horizontal Parabola ($a = 1$)

Graph $x = (y - 2)^2 - 3$. Give the vertex, axis, domain, and range.


This graph has its vertex at $(-3, 2)$ because the roles of x and y are interchanged. It opens to the right (the positive x -direction) because $a = 1$ and $1 > 0$, and has the same shape as $y = x^2$ (but situated horizontally).

To find additional points to plot, it is easiest to substitute a value for y and find the corresponding value for x . For example, let $y = 3$. Then

$$x = (3 - 2)^2 - 3 = -2, \quad \text{giving the point } (-2, 3).$$

 Write the x -value first.

Using symmetry, we can locate the point $(-2, 1)$. See FIGURE 17.




$$\begin{aligned} x &= (y - 2)^2 - 3 \\ \text{Vertex: } &(-3, 2) \\ \text{Axis of symmetry: } &y = 2 \\ \text{Domain: } &[-3, \infty) \\ \text{Range: } &(-\infty, \infty) \end{aligned}$$

FIGURE 17

NOW TRY
NOW TRY ANSWERS

7. 36 ft; 1.5 sec



vertex: $(-1, -2)$; axis: $y = -2$;
domain: $[-1, \infty)$; range: $(-\infty, \infty)$

NOW TRY EXERCISE 9

Graph $x = -3y^2 - 6y - 5$.
Give the vertex, axis, domain, and range.

EXAMPLE 9 Graphing a Horizontal Parabola ($a \neq 1$)

Graph $x = -2y^2 + 4y - 3$. Give the vertex, axis, domain, and range.

$$x = -2y^2 + 4y - 3$$

$$x = (-2y^2 + 4y) - 3$$

Group the variable terms.

$$x = -2(y^2 - 2y) - 3$$

Factor out -2 .

$$x = -2(y^2 - 2y \quad) - 3$$

Prepare to complete the square.

$$\left[\frac{1}{2}(-2) \right]^2 = (-1)^2 = 1$$

Square half the coefficient of the first-degree term.

$$x = -2(y^2 - 2y + 1 - 1) - 3$$

Complete the square within the parentheses. Add and subtract 1.

$$x = -2(y^2 - 2y + 1) + (-2)(-1) - 3$$


Distributive property

Be careful here.

$$x = -2(y - 1)^2 - 1$$

Factor. Simplify.

The vertex is $(-1, 1)$. Because of the negative coefficient -2 in $x = -2(y - 1)^2 - 1$, the graph opens to the left (the negative x -direction). The graph is narrower than the graph of $y = x^2$ because $|-2| = 2$, and $2 > 1$. See **FIGURE 18**.



$$x = -2y^2 + 4y - 3$$

Vertex: $(-1, 1)$

Axis of symmetry: $y = 1$

Domain: $(-\infty, -1]$

Range: $(-\infty, \infty)$

FIGURE 18


NOW TRY

CAUTION Only quadratic equations that are solved for y (whose graphs are vertical parabolas) are functions. The horizontal parabolas in **Examples 8 and 9** are *not* graphs of functions because they do not satisfy the conditions of the vertical line test.

Summary of Graphs of Parabolas

Equation	Graph
$y = ax^2 + bx + c$ or $y = a(x - h)^2 + k$	
$x = ay^2 + by + c$ or $x = a(y - k)^2 + h$	

NOW TRY ANSWER 9.



vertex: $(-2, -1)$; axis: $y = -1$;
domain: $(-\infty, -2]$; range: $(-\infty, \infty)$

8.6 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Answer each question.

- How can we determine just by looking at the equation of a parabola whether it has a vertical or a horizontal axis?
- Why can't the graph of a quadratic function be a parabola with a horizontal axis?
- How can we determine the number of x -intercepts of the graph of a quadratic function without graphing the function?
- If the vertex of the graph of a quadratic function is $(1, -3)$, and the graph opens down, how many x -intercepts does the graph have?
- Which equations have a graph that is a vertical parabola? A horizontal parabola?
 - $y = -x^2 + 20x + 80$
 - $x = 2y^2 + 6y + 5$
 - $x + 1 = (y + 2)^2$
 - $f(x) = (x - 4)^2$
- Which of the equations in **Exercise 5** represent functions?

Find the vertex of each parabola. See **Examples 1–3**.







- $f(x) = x^2 + 8x + 10$
- $f(x) = x^2 + 10x + 23$
- $f(x) = -2x^2 + 4x - 5$
- $f(x) = -3x^2 + 12x - 8$
- $f(x) = x^2 + x - 7$
- $f(x) = x^2 - x + 5$

Find the vertex of each parabola. For each equation, decide whether the graph opens up, down, to the left, or to the right, and whether it is wider, narrower, or the same shape as the graph of $y = x^2$. If it is a parabola with a vertical axis of symmetry, find the discriminant and use it to determine the number of x -intercepts. See **Examples 1–3, 5, 8, and 9**.

- $f(x) = 2x^2 + 4x + 5$
- $f(x) = 3x^2 - 6x + 4$
- $f(x) = -x^2 + 5x + 3$
- $f(x) = -x^2 + 7x + 2$
- $x = \frac{1}{3}y^2 + 6y + 24$
- $x = \frac{1}{2}y^2 + 10y - 5$

Concept Check Match each equation with its graph in choices A–F.

- $y = 2x^2 + 4x - 3$
- $y = -x^2 + 3x + 5$
- $y = -\frac{1}{2}x^2 - x + 1$
- $x = y^2 + 6y + 3$
- $x = -y^2 - 2y + 4$
- $x = 3y^2 + 6y + 5$



Graph each parabola. Give the vertex, axis of symmetry, domain, and range. See Examples 4, 8, and 9.

25. $f(x) = x^2 + 8x + 10$

26. $f(x) = x^2 + 10x + 23$

27. $f(x) = x^2 + 4x + 3$

28. $f(x) = x^2 + 2x - 2$

29. $f(x) = -2x^2 + 4x - 5$

30. $f(x) = -3x^2 + 12x - 8$

31. $f(x) = -3x^2 - 6x + 2$

32. $f(x) = -2x^2 + 12x - 13$

33. $x = (y + 2)^2 + 1$

34. $x = (y + 3)^2 - 2$

35. $x = -(y - 3)^2 - 1$

36. $x = -(y - 2)^2 + 4$

37. $x = -\frac{1}{5}y^2 + 2y - 4$



38. $x = -\frac{1}{2}y^2 - 4y - 6$

39. $x = 3y^2 + 12y + 5$

40. $x = 4y^2 + 16y + 11$

Solve each problem. See Examples 6 and 7.

41. Find the pair of numbers whose sum is 40 and whose product is a maximum. (*Hint:* Let x and $40 - x$ represent the two numbers.)
42. Find the pair of numbers whose sum is 60 and whose product is a maximum.
43. Polk Community College wants to construct a rectangular parking lot on land bordered on one side by a highway. It has 280 ft of fencing that is to be used to fence off the other three sides. What should be the dimensions of the lot if the enclosed area is to be a maximum? What is the maximum area?
44. Bonnie has 100 ft of fencing material to enclose a rectangular exercise run for her dog. One side of the run will border her house, so she will only need to fence three sides. What dimensions will give the enclosure the maximum area? What is the maximum area?



45. Two physics students from American River College find that when a bottle of California sparkling wine is shaken several times, held upright, and uncorked, its cork travels according to the function

$$s(t) = -16t^2 + 64t + 1,$$

where s is its height in feet above the ground t seconds after being released. After how many seconds will it reach its maximum height? What is the maximum height?



46. Professor Barbu has found that the number of students attending his intermediate algebra class is approximated by

$$S(x) = -x^2 + 20x + 80,$$

where x is the number of hours that the Campus Center is open daily. Find the number of hours that the center should be open so that the number of students attending class is a maximum. What is this maximum number of students?

47. Klaus has a taco stand. He has found that his daily costs are approximated by

$$C(x) = x^2 - 40x + 610,$$

where $C(x)$ is the cost, in dollars, to sell x units of tacos. Find the number of units of tacos he should sell to minimize his costs. What is the minimum cost?

48. Mohammad has a frozen yogurt cart. His daily costs are approximated by

$$C(x) = x^2 - 70x + 1500,$$

where $C(x)$ is the cost, in dollars, to sell x units of frozen yogurt. Find the number of units of frozen yogurt he must sell to minimize his costs. What is the minimum cost?

49. If an object on Earth is projected upward with an initial velocity of 32 ft per sec, then its height after t seconds is given by

$$s(t) = -16t^2 + 32t.$$

Find the maximum height attained by the object and the number of seconds it takes to hit the ground.

50. A projectile on Earth is fired straight upward so that its distance (in feet) above the ground t seconds after firing is given by

$$s(t) = -16t^2 + 400t.$$


Find the maximum height it reaches and the number of seconds it takes to reach that height.

51. The graph shows how the average retail price (in dollars per gallon) of regular unleaded gasoline in the United States changed between 2009 and 2016. The graph suggests that a quadratic function would be a good fit to the data. The data are approximated by the function

$$f(x) = -0.1171x^2 + 0.7768x + 2.296.$$

In the model, $x = 0$ represents 2009, $x = 1$ represents 2010, and so on.

- (a) How could we have predicted that this quadratic model would have a negative coefficient for x^2 , based only on the graph shown?
- (b) Algebraically determine the vertex of the graph, with coordinates rounded to the nearest hundredth. Interpret the answer as it applies to this application.



Data from International Energy Agency.


52. The graph shows how Social Security trust fund assets are expected to change and suggests that a quadratic function would be a good fit to the data. The data are approximated by the function

$$f(x) = -20.57x^2 + 758.9x - 3140.$$

In the model, $x = 10$ represents 2010, $x = 15$ represents 2015, and so on, and $f(x)$ is in billions of dollars.

- (a) How could we have predicted that this quadratic model would have a negative coefficient for x^2 , based only on the graph shown?
- (b) Algebraically determine the vertex of the graph. Express the x -coordinate to the nearest hundredth and the y -coordinate to the nearest whole number. Interpret the answer as it applies to this application.

Social Security Assets*



*Projected

Data from Social Security Administration.

Extending Skills In each problem, find the following.

- (a) A function $R(x)$ that describes the total revenue received
- (b) The graph of the function from part (a)
- (c) The number of unsold seats that will produce the maximum revenue
- (d) The maximum revenue
53. A charter flight charges a fare of \$200 per person, plus \$4 per person for each unsold seat on the plane. The plane holds 100 passengers. Let x represent the number of unsold seats. (*Hint:* To find $R(x)$, multiply the number of people flying, $100 - x$, by the price per ticket, $200 + 4x$.)
54. A charter bus charges a fare of \$48 per person, plus \$2 per person for each unsold seat on the bus. The bus has 42 seats. Let x represent the number of unsold seats. (*Hint:* To find $R(x)$, multiply the number riding, $42 - x$, by the price per ticket, $48 + 2x$.)

8.7

Polynomial and Rational Inequalities

OBJECTIVES

- 1 Solve quadratic inequalities.
- 2 Solve polynomial inequalities of degree 3 or greater.
- 3 Solve rational inequalities.

OBJECTIVE 1 Solve quadratic inequalities.

We can combine methods of solving linear inequalities and methods of solving quadratic equations to solve *quadratic inequalities*.

Quadratic Inequality

A **quadratic inequality** (in x here) is an inequality that can be written in the form

$$ax^2 + bx + c < 0, \quad ax^2 + bx + c > 0,$$

$$ax^2 + bx + c \leq 0, \quad \text{or} \quad ax^2 + bx + c \geq 0,$$

where a , b , and c are real numbers and $a \neq 0$.


Examples: $x^2 - 3x - 10 < 0$, $6x^2 + x \geq 1$ [Quadratic inequalities](#)

VOCABULARY

- quadratic inequality
- rational inequality


NOW TRY EXERCISE 1

Use the graph to solve each quadratic inequality.



- (a) $x^2 - 3x - 4 > 0$
 (b) $x^2 - 3x - 4 < 0$

One way to solve a quadratic inequality involves graphing the related quadratic function. This method is justified because the graph is *continuous*—that is, it has no breaks.

EXAMPLE 1 Solving Quadratic Inequalities by Graphing

Solve each inequality.

(a) $x^2 - x - 12 > 0$

We graph the related quadratic function $f(x) = x^2 - x - 12$. We are particularly interested in the x -intercepts, which are found by letting $f(x) = 0$ and solving the following quadratic equation.

$$x^2 - x - 12 = 0 \quad \text{Let } f(x) = 0.$$


$$(x - 4)(x + 3) = 0 \quad \text{Factor.}$$

$$x - 4 = 0 \quad \text{or} \quad x + 3 = 0 \quad \text{Zero-factor property}$$

$$x = 4 \quad \text{or} \quad x = -3 \leftarrow \text{The } x\text{-intercepts are } (4, 0) \text{ and } (-3, 0).$$


The graph opens up because the coefficient of x^2 is positive. See **FIGURE 19(a)**. Notice that x -values less than -3 or greater than 4 result in y -values *greater than* 0 . Thus, the solution set of $x^2 - x - 12 > 0$, written in interval notation, is

$$(-\infty, -3) \cup (4, \infty).$$



The graph is above the x -axis for $(-\infty, -3) \cup (4, \infty)$.

(a)



The graph is below the x -axis for $(-3, 4)$.

(b)

FIGURE 19

(b) $x^2 - x - 12 < 0$

We want values of y that are *less than* 0 . See **FIGURE 19(b)**. Notice from the graph that x -values between -3 and 4 result in y -values less than 0 . Thus, the solution set of $x^2 - x - 12 < 0$, written in interval notation, is $(-3, 4)$.

NOW TRY 

NOTE If the inequalities in **Example 1** had used \geq and \leq , the solution sets would have included the x -values of the intercepts, which make the quadratic expression equal to 0 . They would have been written in interval notation as

$$(-\infty, -3] \cup [4, \infty) \quad \text{and} \quad [-3, 4].$$

Square brackets would indicate that the endpoints -3 and 4 are *included* in the solution sets.

NOW TRY ANSWERS

1. (a) $(-\infty, -3) \cup (4, \infty)$
 (b) $(-3, 4)$

Another method for solving a quadratic inequality uses the basic ideas of **Example 1** without actually graphing the related quadratic function.

EXAMPLE 2 Solving a Quadratic Inequality Using Test Values

Solve and graph the solution set of $x^2 - x - 12 > 0$.

Solve the quadratic equation $x^2 - x - 12 = 0$. (See **Example 1(a)**.)

$$x^2 - x - 12 = 0 \quad \text{Let } f(x) = 0.$$

$$(x - 4)(x + 3) = 0 \quad \text{Factor.}$$

$$x - 4 = 0 \quad \text{or} \quad x + 3 = 0 \quad \text{Zero-factor property}$$

$$x = 4 \quad \text{or} \quad x = -3 \quad \text{Solve each equation.}$$

The numbers 4 and -3 divide a number line into Intervals A, B, and C, as shown in **FIGURE 20**. *Be careful to put the lesser number on the left.*




FIGURE 20

Notice the similarity between **FIGURE 20** and the x -axis with intercepts $(-3, 0)$ and $(4, 0)$ in **FIGURE 19(a)**.

The numbers 4 and -3 are the only values that make the quadratic expression $x^2 - x - 12$ equal to 0. All other numbers make the expression either positive or negative. The sign of the expression can change from positive to negative or from negative to positive only at a number that makes it 0.

Therefore, if one number in an interval satisfies the inequality, then all numbers in that interval will satisfy the inequality.

To see if the numbers in Interval A satisfy the inequality, choose any number from Interval A in **FIGURE 20** (that is, any number less than -3). We choose -5 . Substitute this test value for x in the original inequality $x^2 - x - 12 > 0$.

$$x^2 - x - 12 > 0 \quad \text{Original inequality}$$

$$(-5)^2 - (-5) - 12 \stackrel{?}{>} 0 \quad \text{Let } x = -5.$$

$$25 + 5 - 12 \stackrel{?}{>} 0 \quad \text{Simplify.}$$

$$18 > 0 \quad \text{True}$$

Use parentheses to avoid sign errors.

Because -5 satisfies the inequality, *all* numbers from Interval A are solutions, indicated by **T** in **FIGURE 20**.

Now try 0 from Interval B.

$$x^2 - x - 12 > 0 \quad \text{Original inequality}$$

$$0^2 - 0 - 12 \stackrel{?}{>} 0 \quad \text{Let } x = 0.$$

$$-12 > 0 \quad \text{False}$$

The numbers in Interval B are *not* solutions, indicated by **F** in **FIGURE 20**.

Now try 5 from Interval C.

$$x^2 - x - 12 > 0 \quad \text{Original inequality}$$

$$5^2 - 5 - 12 \stackrel{?}{>} 0 \quad \text{Let } x = 5.$$

$$8 > 0 \quad \text{True}$$

All numbers from Interval C are solutions, indicated by **T** in **FIGURE 20**.

NOW TRY EXERCISE 2

Solve and graph the solution set.

$$x^2 + 2x - 8 > 0$$

Based on these results (shown by the colored letters in **FIGURE 20**), the solution set includes all numbers in Intervals A and C, as shown in **FIGURE 21**.




FIGURE 21

The solution set is written in interval notation as

$$(-\infty, -3) \cup (4, \infty).$$


-3 and 4 are *not* included because the symbol $>$ does not include equality.

This agrees with the solution set found in **Example 1(a)**.

NOW TRY

Solving a Quadratic Inequality

- Step 1** Write the inequality as an equation and solve it.
- Step 2** Use the solutions from Step 1 to determine intervals. Graph the values found in Step 1 on a number line. These values divide the number line into intervals.
- Step 3** Find the intervals that satisfy the inequality. Substitute a test value from each interval into the original inequality to determine the intervals that satisfy the inequality. All numbers in those intervals are in the solution set. A graph of the solution set will usually look like one of these.



- Step 4** Consider the endpoints separately. The values from Step 1 are included in the solution set if the inequality symbol is \leq or \geq . They are not included if it is $<$ or $>$.

EXAMPLE 3 Solving a Quadratic Inequality

- Step 1** Solve and graph the solution set of $2x^2 + 5x \leq 12$.

$$2x^2 + 5x = 12 \quad \text{Related quadratic equation}$$

$$2x^2 + 5x - 12 = 0 \quad \text{Standard form}$$

$$(2x - 3)(x + 4) = 0 \quad \text{Factor.}$$

$$2x - 3 = 0 \quad \text{or} \quad x + 4 = 0 \quad \text{Zero-factor property}$$

$$x = \frac{3}{2} \quad \text{or} \quad x = -4 \quad \text{Solve each equation.}$$

- Step 2** The numbers $\frac{3}{2}$ and -4 divide a number line into three intervals. See **FIGURE 22**.




FIGURE 22

NOW TRY ANSWER

2.
 $(-\infty, -4) \cup (2, \infty)$

NOW TRY EXERCISE 3

Solve and graph the solution set.

$$3x^2 - 11x \leq 4$$

Steps 3 and 4 Substitute a test value from each interval in the *original* inequality $2x^2 + 5x \leq 12$ to determine which intervals satisfy the inequality.

Interval	Test Value	Test of Inequality	True or False?
A	-5	$25 \leq 12$	F
B	0	$0 \leq 12$	T
C	2	$18 \leq 12$	F

We use a table to organize this information. (Verify it.)

The numbers in Interval B are solutions. See **FIGURE 23**.




FIGURE 23

The solution set is the interval $\left[-4, \frac{3}{2}\right]$.

-4 and $\frac{3}{2}$ are included because the symbol \leq includes equality.

NOW TRY

NOW TRY EXERCISE 4

Solve each inequality.

- (a) $(4x - 1)^2 > -3$
- (b) $(4x - 1)^2 < -3$

EXAMPLE 4 Solving Special Cases

Solve each inequality.

(a) $(2x - 3)^2 > -1$

Because $(2x - 3)^2$ is never negative, it is always greater than -1 . Thus, the solution set for $(2x - 3)^2 > -1$ is the set of all real numbers, $(-\infty, \infty)$.

(b) $(2x - 3)^2 < -1$

Using similar reasoning as in part (a), there is no solution for this inequality. The solution set is \emptyset .

NOW TRY

OBJECTIVE 2 Solve polynomial inequalities of degree 3 or greater.

EXAMPLE 5 Solving a Third-Degree Polynomial Inequality

Solve and graph the solution set of $(x - 1)(x + 2)(x - 4) \leq 0$.

This *cubic* (third-degree) inequality can be solved by extending the zero-factor property to more than two factors. (Step 1)

$$(x - 1)(x + 2)(x - 4) = 0$$

Set the factored polynomial equal to 0.

$$x - 1 = 0 \quad \text{or} \quad x + 2 = 0 \quad \text{or} \quad x - 4 = 0$$

Zero-factor property

$$x = 1 \quad \text{or} \quad x = -2 \quad \text{or} \quad x = 4$$

Solve each equation.

Locate the numbers -2 , 1 , and 4 on a number line, as in **FIGURE 24**, to determine the Intervals A, B, C, and D. (Step 2)





FIGURE 24

NOW TRY ANSWERS



$$\left[-\frac{1}{3}, 4\right]$$

4. (a) $(-\infty, \infty)$ (b) \emptyset


NOW TRY EXERCISE 5

Solve and graph the solution set.

$$(x + 4)(x - 3)(2x + 1) \leq 0$$

Substitute a test value from each interval in the *original* inequality to determine which intervals satisfy $(x - 1)(x + 2)(x - 4) \leq 0$. (Step 3)

Interval	Test Value	Test of Inequality	True or False?
A	-3	$-28 \leq 0$	T
B	0	$8 \leq 0$	F
C	2	$-8 \leq 0$	T
D	5	$28 \leq 0$	F

The numbers in Intervals A and C are in the solution set. The three endpoints are included because the inequality symbol \leq includes equality. (Step 4) See FIGURE 25.



FIGURE 25

The solution set is the interval $(-\infty, -2] \cup [1, 4]$.

NOW TRY 

OBJECTIVE 3 Solve rational inequalities.

Rational inequalities involve rational expressions and are solved similarly.

Solving a Rational Inequality

- Step 1** Write the inequality so that 0 is on one side and there is a single fraction on the other side.
- Step 2** Determine the values that make the numerator or denominator equal to 0.
- Step 3** Divide a number line into intervals. Use the values from Step 2.
- Step 4** Find the intervals that satisfy the inequality. Test a value from each interval by substituting it into the *original* inequality.
- Step 5** Consider the endpoints separately. Exclude any values that make the denominator 0.

EXAMPLE 6 Solving a Rational Inequality

Solve and graph the solution set of $\frac{-1}{x - 3} > 1$.

Write the inequality so that 0 is on one side. (Step 1)

$$\frac{-1}{x - 3} - 1 > 0 \quad \text{Subtract 1.}$$

$$\frac{-1}{x - 3} - \frac{x - 3}{x - 3} > 0 \quad \text{Use } x - 3 \text{ as the common denominator.}$$

$$\frac{-1 - (x - 3)}{x - 3} > 0 \quad \text{Write the left side as a single fraction.}$$

Be careful with signs.

$$\frac{-1 - x + 3}{x - 3} > 0 \quad \text{Distributive property}$$

$$\frac{-x + 2}{x - 3} > 0 \quad \text{Combine like terms in the numerator.}$$

NOW TRY ANSWER

5.

$$(-\infty, -4] \cup \left[-\frac{1}{2}, 3\right]$$


NOW TRY EXERCISE 6

Solve and graph the solution set.

$$\frac{3}{x+1} > 4$$

The sign of $\frac{-x+2}{x-3}$ will change from positive to negative or negative to positive only at those values that make the numerator or denominator 0. The number 2 makes the numerator 0, and 3 makes the denominator 0. (Step 2) These two numbers, 2 and 3, divide a number line into three intervals. See FIGURE 26. (Step 3)




FIGURE 26

Substituting a test value from each interval in the *original inequality*, $\frac{-1}{x-3} > 1$, gives the results shown in the table. (Step 4)

Interval	Test Value	Test of Inequality	True or False?
A	0	$\frac{1}{3} > 1$	F
B	2.5	$2 > 1$	T
C	4	$-1 > 1$	F

The numbers in Interval B are solutions. This interval does not include 3 because it makes the denominator in the original inequality 0. The number 2 is not included either because the inequality symbol $>$ does not include equality. (Step 5) See FIGURE 27.



FIGURE 27

The solution set is the interval $(2, 3)$.

NOW TRY

CAUTION When solving a rational inequality, any number that makes the denominator 0 must be excluded from the solution set.

EXAMPLE 7 Solving a Rational Inequality
Solve and graph the solution set of $\frac{x-2}{x+2} \leq 2$.

Write the inequality so that 0 is on one side. (Step 1)

$$\frac{x-2}{x+2} - 2 \leq 0 \quad \text{Subtract 2.}$$

$$\frac{x-2}{x+2} - \frac{2(x+2)}{x+2} \leq 0 \quad \text{Use } x+2 \text{ as the common denominator.}$$


$$\frac{x-2-2(x+2)}{x+2} \leq 0 \quad \text{Write as a single fraction.}$$

Be careful with signs.

$$\frac{x-2-2x-4}{x+2} \leq 0 \quad \text{Distributive property}$$

$$\frac{-x-6}{x+2} \leq 0 \quad \text{Combine like terms in the numerator.}$$

NOW TRY ANSWER

6. 
 $(-1, -\frac{1}{4})$

NOW TRY EXERCISE 7

Solve and graph the solution set.

$$\frac{x - 3}{x + 3} \leq 2$$

In the inequality $\frac{-x - 6}{x + 2} \leq 0$, the number -6 makes the numerator 0, and -2 makes the denominator 0. (Step 2) These two numbers determine three intervals on a number line. See FIGURE 28. (Step 3)




FIGURE 28

Substitute a test value from each interval in the *original* inequality $\frac{x - 2}{x + 2} \leq 2$. (Step 4)

Interval	Test Value	Test of Inequality	True or False?
A	-8	$\frac{5}{3} \leq 2$	T
B	-4	$3 \leq 2$	F
C	0	$-1 \leq 2$	T

The numbers in Intervals A and C are solutions. The number -6 satisfies the original inequality, but -2 does not because it makes the denominator 0. (Step 5) See FIGURE 29.




FIGURE 29

NOW TRY ANSWER

7. $(-\infty, -9] \cup (-3, \infty)$

The solution set is the interval $(-\infty, -6] \cup (-2, \infty)$.

NOW TRY

8.7 Exercises

FOR EXTRA HELP

MyLab Math

Video solutions for select problems available in MyLab Math



1. **Concept Check** The solution set of the inequality $x^2 + x - 12 < 0$ is the interval $(-4, 3)$. Without actually performing any work, give the solution set of the inequality

$$x^2 + x - 12 \geq 0.$$

2. Explain how to determine whether to include or to exclude endpoints when solving a quadratic or higher-degree inequality.

In each exercise, the graph of a quadratic function f is given. Use the graph to find the solution set of each equation or inequality. See Example 1.


3. (a) $x^2 - 4x + 3 = 0$
 (b) $x^2 - 4x + 3 > 0$
 (c) $x^2 - 4x + 3 < 0$
4. (a) $3x^2 + 10x - 8 = 0$
 (b) $3x^2 + 10x - 8 \geq 0$
 (c) $3x^2 + 10x - 8 < 0$



5. (a) $-x^2 + 3x + 10 = 0$

(b) $-x^2 + 3x + 10 \geq 0$


(c) $-x^2 + 3x + 10 \leq 0$



6. (a) $-2x^2 - x + 15 = 0$

(b) $-2x^2 - x + 15 \geq 0$

(c) $-2x^2 - x + 15 \leq 0$



Solve each inequality, and graph the solution set. See Examples 2 and 3. (Hint: In Exercises 23 and 24, use the quadratic formula.)

7. $(x + 1)(x - 5) > 0$

8. $(x + 6)(x - 2) > 0$

9. $(x + 4)(x - 6) < 0$

10. $(x + 4)(x - 8) < 0$

11. $x^2 - 4x + 3 \geq 0$

12. $x^2 - 3x - 10 \geq 0$

13. $10x^2 + 9x \geq 9$

14. $3x^2 + 10x \geq 8$

15. $4x^2 - 9 \leq 0$

16. $9x^2 - 25 \leq 0$

17. $6x^2 + x \geq 1$

18. $4x^2 + 7x \geq -3$

19. $z^2 - 4z \geq 0$

20. $x^2 + 2x < 0$

21. $3x^2 - 5x \leq 0$

22. $2z^2 + 3z > 0$

23. $x^2 - 6x + 6 \geq 0$

24. $3x^2 - 6x + 2 \leq 0$

Solve each inequality. See Example 4.

25. $(4 - 3x)^2 \geq -2$

26. $(7 - 6x)^2 \geq -1$

27. $(3x + 5)^2 \leq -4$

28. $(8x + 5)^2 \leq -5$

29. $(2x + 5)^2 < 0$

30. $(3x - 7)^2 < 0$

31. $(5x - 1)^2 \geq 0$

32. $(4x + 1)^2 \geq 0$

Solve each inequality, and graph the solution set. See Example 5.

33. $(x - 1)(x - 2)(x - 4) < 0$

34. $(2x + 1)(3x - 2)(4x + 7) < 0$

35. $(x - 4)(2x + 3)(3x - 1) \geq 0$

36. $(x + 2)(4x - 3)(2x + 7) \geq 0$

Solve each inequality, and graph the solution set. See Examples 6 and 7.

37. $\frac{x - 1}{x - 4} > 0$

38. $\frac{x + 1}{x - 5} > 0$

39. $\frac{2x + 3}{x - 5} \leq 0$

40. $\frac{3x + 7}{x - 3} \leq 0$

41. $\frac{8}{x - 2} \geq 2$

42. $\frac{20}{x - 1} \geq 1$

43. $\frac{3}{2x - 1} < 2$

44. $\frac{6}{x - 1} < 1$

45. $\frac{x - 3}{x + 2} \geq 2$

46. $\frac{m + 4}{m + 5} \geq 2$

47. $\frac{x - 8}{x - 4} < 3$

48. $\frac{2t - 3}{t + 1} > 4$

49. $\frac{4k}{2k-1} < k$

50. $\frac{r}{r+2} < 2r$

51. $\frac{2x-3}{x^2+1} \geq 0$

52. $\frac{9x-8}{4x^2+25} < 0$

53. $\frac{(3x-5)^2}{x+2} > 0$

54. $\frac{(5x-3)^2}{2x+1} \leq 0$

RELATING CONCEPTS For Individual or Group Work (Exercises 55–58)

A model rocket is projected vertically upward from the ground. Its distance s in feet above the ground after t seconds is given by the quadratic function

$$s(t) = -16t^2 + 256t.$$

Work Exercises 55–58 in order, to see how quadratic equations and inequalities are related.

55. At what times will the rocket be 624 ft above the ground? (*Hint*: Let $s(t) = 624$ and solve the quadratic equation.)
56. At what times will the rocket be more than 624 ft above the ground? (*Hint*: Let $s(t) > 624$ and solve the quadratic inequality.)
57. At what times will the rocket be at ground level? (*Hint*: Let $s(t) = 0$ and solve the quadratic equation.)
58. At what times will the rocket be less than 624 ft above the ground? (*Hint*: Let $s(t) < 624$, solve the quadratic inequality, and observe the solutions in **Exercises 56 and 57** to determine the least and greatest possible values of t .)



Chapter 8

Summary

Key Terms

8.1

quadratic equation
second-degree equation

8.2

quadratic formula
discriminant

8.5

parabola
vertex
axis of symmetry (axis)
quadratic function

8.7

quadratic inequality
rational inequality

8.3

quadratic in form

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- The **quadratic formula** is
 - a formula to find the number of solutions of a quadratic equation
 - a formula to find the type of solutions of a quadratic equation
 - the standard form of a quadratic equation
 - a general formula for solving any quadratic equation.
- A **quadratic function** is a function that can be written in the form
 - $f(x) = mx + b$, for real numbers m and b
 - $f(x) = \frac{P(x)}{Q(x)}$, where $Q(x) \neq 0$
 - $f(x) = ax^2 + bx + c$, for real numbers a , b , and c ($a \neq 0$)
 - $f(x) = \sqrt{x}$, for $x \geq 0$.
- A **parabola** is the graph of
 - any equation in two variables
 - a linear equation
 - an equation of degree 3
 - a quadratic equation in two variables.

4. The **vertex** of a parabola is
- the point where the graph intersects the y -axis
 - the point where the graph intersects the x -axis
 - the lowest point on a parabola that opens up or the highest point on a parabola that opens down
 - the origin.
5. The **axis of symmetry** of a parabola is
- either the x -axis or the y -axis
 - the vertical line (of a vertical parabola) or the horizontal line (of a horizontal parabola) through the vertex
 - the lowest or highest point on a graph of a parabola
 - a line through the origin.
6. A parabola is **symmetric about its axis** because its graph
- is near the axis
 - is identical on each side of the axis
 - looks different on each side of the axis
 - intersects the axis.

ANSWERS

1. D; *Example:* The solutions of $ax^2 + bx + c = 0$ ($a \neq 0$) are given by $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$. 2. C; *Examples:* $f(x) = x^2 - 2$, $f(x) = (x + 4)^2 + 1$, $f(x) = x^2 - 4x + 5$ 3. D; *Example:* The quadratic equation $y = x^2$ has a parabola as its graph. 4. C; *Example:* The graph of $y = (x + 3)^2$ has vertex $(-3, 0)$, which is the lowest point on the graph. 5. B; *Example:* The axis of $y = (x + 3)^2$ is the vertical line $x = -3$. 6. B; *Example:* Because the graph of $y = (x + 3)^2$ is symmetric about its axis $x = -3$, the points $(-2, 1)$ and $(-4, 1)$ are on the graph.

Quick Review

CONCEPTS

8.1 The Square Root Property and Completing the Square

Square Root Property

If x and k are complex numbers and $x^2 = k$, then

$$x = \sqrt{k} \quad \text{or} \quad x = -\sqrt{k}.$$

Completing the Square

To solve $ax^2 + bx + c = 0$ (where $a \neq 0$), follow these steps.

- Step 1** If $a \neq 1$, divide each side by a .
- Step 2** Write the equation with the variable terms on one side and the constant on the other.
- Step 3** Complete the square.
- Take half the coefficient of x and square it.
 - Add the square to each side.
 - Factor the perfect square trinomial, and write it as the square of a binomial. Combine terms on the other side.
- Step 4** Use the square root property to solve.

EXAMPLES

Solve $(x - 1)^2 = 8$.

$$x - 1 = \sqrt{8} \quad \text{or} \quad x - 1 = -\sqrt{8}$$

$$x = 1 + 2\sqrt{2} \quad \text{or} \quad x = 1 - 2\sqrt{2}$$

Solution set: $\{1 + 2\sqrt{2}, 1 - 2\sqrt{2}\}$, or $\{1 \pm 2\sqrt{2}\}$

Solve $2x^2 - 4x - 18 = 0$.

$$x^2 - 2x - 9 = 0 \quad \text{Divide by 2.}$$

$$x^2 - 2x = 9 \quad \text{Add 9.}$$

$$\left[\frac{1}{2}(-2)\right]^2 = (-1)^2 = 1$$

$$x^2 - 2x + 1 = 9 + 1 \quad \text{Add 1.}$$

$$(x - 1)^2 = 10 \quad \text{Factor. Add.}$$

$$x - 1 = \sqrt{10} \quad \text{or} \quad x - 1 = -\sqrt{10} \quad \text{Square root property}$$

$$x = 1 + \sqrt{10} \quad \text{or} \quad x = 1 - \sqrt{10}$$

Solution set: $\{1 + \sqrt{10}, 1 - \sqrt{10}\}$, or $\{1 \pm \sqrt{10}\}$

8.2 The Quadratic Formula

Quadratic Formula

The solutions of $ax^2 + bx + c = 0$ (where $a \neq 0$) are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Solve $3x^2 + 5x + 2 = 0$.

$$x = \frac{-5 \pm \sqrt{5^2 - 4(3)(2)}}{2(3)} \quad a = 3, b = 5, c = 2$$

$$x = \frac{-5 \pm 1}{6} \quad \text{Simplify.}$$

$$x = -\frac{2}{3} \quad \text{or} \quad x = -1 \quad \text{Two solutions, one from + and one from -}$$

Solution set: $\{-1, -\frac{2}{3}\}$

CONCEPTS

Using the Discriminant

The discriminant $b^2 - 4ac$ of $ax^2 + bx + c = 0$ (where a , b , and c are integers) can be used to determine the number and type of solutions.

Discriminant $b^2 - 4ac$	Number and Type of Solutions
Positive, and the square of an integer	Two rational solutions
Positive, but not the square of an integer	Two irrational solutions
Zero	One rational solution
Negative	Two nonreal complex solutions

8.3 Equations That Lead to Quadratic Methods

A nonquadratic equation that can be written in the form

$$au^2 + bu + c = 0,$$

for $a \neq 0$ and an algebraic expression u , is quadratic in form.

Solving an Equation Quadratic in Form by Substitution

Step 1 Define a temporary variable u .

Step 2 Solve the quadratic equation obtained in Step 1.

Step 3 Replace u with the expression it defined.

Step 4 Solve the resulting equations for the original variable.

Step 5 Check all solutions in the original equation.

8.4 Formulas and Further Applications

Solving a Formula for a Squared Variable

Case 1 If the variable appears only to the second power: Isolate the second-degree variable on one side of the equation, and then use the square root property.

Case 2 If the variable appears to the first and second powers: Write the equation in standard form, and then use the quadratic formula.

EXAMPLES

For $x^2 + 3x - 10 = 0$, the discriminant is as follows.

$$\begin{aligned} b^2 - 4ac &= 3^2 - 4(1)(-10) && a = 1, b = 3, c = -10 \\ &= 49 && \text{There are two rational} \\ &= 7^2 && \text{solutions.} \\ &&& \text{Simplify.} \end{aligned}$$

Because the discriminant is a perfect square, the quadratic equation can be solved using the zero-factor property.

Solve $3(x + 5)^2 + 7(x + 5) + 2 = 0$.

$$\begin{aligned} 3u^2 + 7u + 2 &= 0 && \text{Let } u = x + 5. \\ (3u + 1)(u + 2) &= 0 && \text{Factor.} \\ 3u + 1 = 0 &\quad \text{or} \quad u + 2 = 0 && \text{Zero-factor} \\ &&& \text{property} \\ u = -\frac{1}{3} &\quad \text{or} \quad u = -2 && \text{Solve for } u. \\ x + 5 = -\frac{1}{3} &\quad \text{or} \quad x + 5 = -2 && \text{Replace } u \text{ with} \\ &&& \text{ } x + 5. \\ x = -\frac{16}{3} &\quad \text{or} \quad x = -7 && \text{Subtract 5.} \end{aligned}$$

Solution set: $\left\{-7, -\frac{16}{3}\right\}$

Solve $A = \frac{2mp}{r^2}$ for r . (Case 1)

$$\begin{aligned} r^2A &= 2mp && \text{Multiply by } r^2. \\ r^2 &= \frac{2mp}{A} && \text{Divide by } A. \\ r &= \pm \sqrt{\frac{2mp}{A}} && \text{Square root property} \\ r &= \frac{\pm \sqrt{2mpA}}{A} && \text{Rationalize denominator.} \end{aligned}$$

Solve $x^2 + rx = t$ for x . (Case 2)

$$\begin{aligned} x^2 + rx - t &= 0 && \text{Standard form} \\ x &= \frac{-r \pm \sqrt{r^2 - 4(1)(-t)}}{2(1)} && a = 1, b = r, \\ &&& c = -t \\ x &= \frac{-r \pm \sqrt{r^2 + 4t}}{2} && \text{Simplify.} \end{aligned}$$

CONCEPTS

8.5 Graphs of Quadratic Functions

The graph of the quadratic function $F(x) = a(x - h)^2 + k$ (where $a \neq 0$) is a parabola.

- The vertex of the parabola is (h, k) .
- The axis of symmetry is the vertical line $x = h$.
- The graph opens up if $a > 0$ and down if $a < 0$.
- The graph is wider than the graph of $f(x) = x^2$ if $0 < |a| < 1$ and narrower if $|a| > 1$.

8.6 More about Parabolas and Their Applications

The vertex of the graph of $f(x) = ax^2 + bx + c$ (where $a \neq 0$) may be found by completing the square or by using the vertex formula $\left(\frac{-b}{2a}, f\left(\frac{-b}{2a}\right)\right)$.

Graphing a Quadratic Function

Step 1 Determine whether the graph opens up or down.

Step 2 Find the vertex.

Step 3 Find any intercepts.

Step 4 Find and plot additional points as needed.

Horizontal Parabolas

The graph of an equation of the form

$$x = ay^2 + by + c \quad \text{or} \quad x = a(y - k)^2 + h$$

is a horizontal parabola.

- The vertex of the parabola is (h, k) .
- The axis of symmetry is the horizontal line $y = k$.
- The graph opens to the right if $a > 0$ and to the left if $a < 0$.

Horizontal parabolas do not represent functions.

8.7 Polynomial and Rational Inequalities

Solving a Quadratic (or Higher-Degree Polynomial) Inequality

Step 1 Write the inequality as an equation and solve it.


Step 2 Use the values found in Step 1 to divide a number line into intervals.

Step 3 Substitute a test value from each interval into the *original* inequality to determine the intervals that belong to the solution set.

Step 4 Consider the endpoints separately.

EXAMPLES

Graph $f(x) = -(x + 3)^2 + 1$.



The graph opens down because $a < 0$.


Vertex: $(-3, 1)$

Axis of symmetry: $x = -3$

Domain: $(-\infty, \infty)$

Range: $(-\infty, 1]$

Graph $f(x) = x^2 + 4x + 3$.



The graph opens up because $a > 0$.

Vertex: $(-2, -1)$

The solutions of $x^2 + 4x + 3 = 0$ are -1 and -3 , so the x -intercepts are $(-1, 0)$ and $(-3, 0)$.


$f(0) = 3$, so the y -intercept is $(0, 3)$.

Axis of symmetry: $x = -2$

Domain: $(-\infty, \infty)$

Range: $[-1, \infty)$

Graph $x = 2y^2 + 6y + 5$.



The graph opens to the right because $a > 0$.

Vertex: $\left(\frac{1}{2}, -\frac{3}{2}\right)$

Axis of symmetry: $y = -\frac{3}{2}$

Domain: $\left[\frac{1}{2}, \infty\right)$

Range: $(-\infty, \infty)$


Solve $2x^2 + 5x + 2 < 0$.

$$2x^2 + 5x + 2 = 0 \quad \text{Related equation}$$

$$(2x + 1)(x + 2) = 0 \quad \text{Factor.}$$

$$2x + 1 = 0 \quad \text{or} \quad x + 2 = 0 \quad \text{Zero-factor property}$$

$$x = -\frac{1}{2} \quad \text{or} \quad x = -2 \quad \text{Solve each equation.}$$



Test values: -3 (Interval A), -1 (Interval B), 0 (Interval C); $x = -3$ makes the original inequality false, $x = -1$ makes it true, and $x = 0$ makes it false.

Solution set: $\left(-2, -\frac{1}{2}\right)$ (Endpoints are not included because $<$ does not include equality.)

CONCEPTS

Solving a Rational Inequality

- Step 1** Write the inequality so that 0 is on one side and there is a single fraction on the other side.
- Step 2** Determine the values that make the numerator or denominator 0.
- Step 3** Use the values from Step 2 to divide a number line into intervals.
- Step 4** Substitute a test value from each interval into the *original* inequality to determine the intervals that belong to the solution set.
- Step 5** Consider the endpoints separately. Exclude any values that make the denominator 0.

EXAMPLES

Solve $\frac{x}{x+2} \geq 4$.

$$\frac{x}{x+2} - 4 \geq 0 \quad \text{Subtract 4.}$$

$$\frac{x}{x+2} - \frac{4(x+2)}{x+2} \geq 0 \quad \text{Write with a common denominator.}$$

$$\frac{x - 4x - 8}{x+2} \geq 0 \quad \text{Write as a single fraction; distributive property}$$

$$\frac{-3x - 8}{x+2} \geq 0 \quad \text{Subtract fractions.}$$

$-\frac{8}{3}$ makes the numerator 0, and -2 makes the denominator 0.

Interval A	Interval B	Interval C	→	Intervals:
F	$-\frac{8}{3}$	T	-2	F
				$(-\infty, -\frac{8}{3}),$ $(-\frac{8}{3}, -2), (-2, \infty)$

Test values: -4 from Interval A makes the original inequality false, $-\frac{7}{3}$ from Interval B makes it true, and 0 from Interval C makes it false.

Solution set: $[-\frac{8}{3}, -2)$ (The endpoint -2 is not included because it makes the denominator 0.)

Chapter 8

Review Exercises

8.1 Solve each equation using the square root property or completing the square.

1. $t^2 = 121$

2. $p^2 = 3$

3. $(2x + 5)^2 = 100$

*4. $(3x - 2)^2 = -25$

5. $x^2 + 4x = 15$

6. $2x^2 - 3x = -1$

Solve each problem. Round answers to the nearest tenth of a second.

7. The height of the Eiffel Tower is 984 ft. Use Galileo's formula

$$d = 16t^2 \quad (\text{where } d \text{ is in feet})$$

to find how long it would take a phone dropped from the top of the Eiffel Tower to reach the ground. (Data from www.wonderopolis.org)

8. The High Roller observation wheel in Las Vegas has a height of about 168 m. Use the metric version of Galileo's formula,

$$d = 4.9t^2 \quad (\text{where } d \text{ is in meters}),$$

to find how long it would take a wallet dropped from the top of the High Roller to reach the ground. (Data from www.caesars.com)

*This exercise requires knowledge of complex numbers.

8.2 Solve each equation using the quadratic formula.

9. $2x^2 + x - 21 = 0$

10. $x^2 + 5x = 7$

11. $(t + 3)(t - 4) = -2$

*12. $2x^2 + 3x + 4 = 0$

*13. $3p^2 = 2(2p - 1)$

14. $x(2x - 7) = 3x^2 + 3$

Find the discriminant and use it to predict whether the solutions to each equation are

A. two rational numbers

B. one rational number

C. two irrational numbers

D. two nonreal complex numbers.

Tell whether each equation can be solved using the zero-factor property, or the quadratic formula should be used instead. Do not actually solve.

15. $x^2 + 5x + 2 = 0$

16. $4t^2 = 3 - 4t$

17. $4x^2 = 6x - 8$

18. $9z^2 + 30z + 25 = 0$

8.3 Solve each equation. Check the solutions.

19. $\frac{15}{x} = 2x - 1$

20. $\frac{1}{n} + \frac{2}{n+1} = 2$

21. $-2r = \sqrt{\frac{48 - 20r}{2}}$

22. $8(3x + 5)^2 + 2(3x + 5) - 1 = 0$

23. $2x^{2/3} - x^{1/3} - 28 = 0$

24. $p^4 - 10p^2 + 9 = 0$

Solve each problem. Round answers to the nearest tenth, as necessary.

25. Bahaa paddled a canoe 20 mi upstream, then paddled back. If the rate of the current was 3 mph and the total trip took 7 hr, what was Bahaa's rate?

26. Carol Ann drove 8 mi to pick up a friend, and then drove 11 mi to a mall at a rate 15 mph faster. If Carol Ann's total travel time was 24 min, what was her rate on the trip to pick up her friend?

27. An old machine processes a batch of checks in 1 hr more time than a new one. How long would it take the old machine to process a batch of checks that the two machines together process in 2 hr?

28. Zoran can process a stack of invoices 1 hr faster than Claude can. Working together, they take 1.5 hr. How long would the job take each person working alone?

	Rate	Time Working Together	Fractional Part of the Job Done
Zoran			
Claude			

8.4 Solve each formula for the specified variable. (Leave \pm in the answers as needed.)


29. $k = \frac{rF}{wv^2}$ for v

30. $p = \sqrt{\frac{yz}{6}}$ for y


31. $mt^2 = 3mt + 6$ for t

Solve each problem.


32. A large machine requires a part in the shape of a right triangle with a hypotenuse 9 ft less than twice the length of the longer leg. The shorter leg must be $\frac{3}{4}$ the length of the longer leg. Find the lengths of the three sides of the part.



33. A square has an area of 256 cm^2 . If the same amount is removed from one dimension and added to the other, the resulting rectangle has an area 16 cm^2 less. Find the dimensions of the rectangle.



34. Allen wants to buy a mat for a photograph that measures 14 in. by 20 in. He wants to have an even border around the picture when it is mounted on the mat. If the area of the mat he chooses is 352 in.^2 , how wide will the border be?



35. If a square piece of cardboard has 3-in. squares cut from its corners and then has the flaps folded up to form an open-top box, the volume of the box is given by the formula


$$V = 3(x - 6)^2,$$

where x is the length of each side of the original piece of cardboard in inches. What original length would yield a box with volume 432 in.^3 ?

36. A searchlight moves horizontally back and forth along a wall. The distance of the light from a starting point at t minutes is given by the quadratic function

$$f(t) = 100t^2 - 300t.$$

How long will it take before the light returns to the starting point?



8.5, 8.6 Identify the vertex of each parabola.

37. $f(x) = -(x - 1)^2$

38. $f(x) = (x - 3)^2 + 7$

39. $x = (y - 3)^2 - 4$

40. $y = -3x^2 + 4x - 2$

Graph each parabola. Give the vertex, axis of symmetry, domain, and range.

41. $y = 2(x - 2)^2 - 3$

42. $f(x) = -2x^2 + 8x - 5$

43. $x = 2(y + 3)^2 - 4$

44. $x = -\frac{1}{2}y^2 + 6y - 14$

Solve each problem.

45. The height (in feet) of a projectile t seconds after being fired from Earth into the air is given by

$$f(t) = -16t^2 + 160t.$$

Find the number of seconds required for the projectile to reach maximum height. What is the maximum height?

46. Find the length and width of a rectangle having a perimeter of 200 m if the area is to be a maximum. What is the maximum area?

8.7 Solve each inequality, and graph the solution set.

47. $(x - 4)(2x + 3) > 0$

48. $x^2 + x \leq 12$

49. $(x + 2)(x - 3)(x + 5) \leq 0$

50. $(4x + 3)^2 \leq -4$

51. $\frac{6}{2z - 1} < 2$

52. $\frac{3t + 4}{t - 2} \leq 1$

Chapter 8

Mixed Review Exercises

Solve each equation or inequality.

1. $V = r^2 + R^2h$ for R

*2. $3t^2 - 6t = -4$

3. $(3x + 11)^2 = 7$

4. $S = \frac{Id^2}{k}$ for d

5. $(8x - 7)^2 \geq -1$

6. $2x - \sqrt{x} = 6$

7. $x^4 - 8x^2 = -1$

8. $\frac{-2}{x + 5} \leq -5$

9. $6 + \frac{15}{s^2} = -\frac{19}{s}$

10. $(x^2 - 2x)^2 = 11(x^2 - 2x) - 24$

11. $(r - 1)(2r + 3)(r + 6) < 0$

Work each problem.

12. Match each equation in parts (a) – (f) with the figure that most closely resembles its graph in choices A–F.

(a) $g(x) = x^2 - 5$







(b) $h(x) = -x^2 + 4$

(c) $F(x) = (x - 1)^2$

(d) $G(x) = (x + 1)^2$

(e) $H(x) = (x - 1)^2 + 1$

(f) $K(x) = (x + 1)^2 + 1$



13. Graph $f(x) = 4x^2 + 4x - 2$. Give the vertex, axis of symmetry, domain, and range.

14. In 4 hr, Rajeed can travel 15 mi upriver and come back. The rate of the current is 5 mph. Find the rate of the boat in still water.

15. Two pieces of a large wooden puzzle fit together to form a rectangle with length 1 cm less than twice the width. The diagonal, where the two pieces meet, is 2.5 cm in length. Find the length and width of the rectangle.

*This exercise requires knowledge of complex numbers.

16. The percent of the U.S. population that was foreign-born during the years 1930–2010 can be modeled by the quadratic function

$$f(x) = 0.0043x^2 - 0.3245x + 11.53,$$

where $x = 0$ represents 1930, $x = 10$ represents 1940, and so on. (Data from U.S. Census Bureau.)

- (a) The coefficient of x^2 in the model is positive, so the graph of this quadratic function is a parabola that opens up. Will the y -value of the vertex of this graph be a maximum or a minimum?
- (b) According to the model, in what year during this period was the percent of foreign-born population a minimum? (Round down for the year.) Use the actual x -value of the vertex, to the nearest tenth, to find this percent, also to the nearest tenth.

Chapter 8

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

► View the complete solutions to all Chapter Test exercises in MyLab Math.

1. Solve $5x^2 + 13x = 6$ using the zero-factor property.

Solve each equation using the square root property or completing the square.

2. $t^2 = 54$

3. $(7x + 3)^2 = 25$

4. $x^2 + 2x = 4$

Solve each equation using the quadratic formula.

5. $2x^2 - 3x - 1 = 0$

*6. $3t^2 - 4t = -5$

7. $3x = \sqrt{\frac{9x + 2}{2}}$

- *8. If k is a negative number, then which one of the following equations will have two nonreal complex solutions?

A. $x^2 = 4k$

B. $x^2 = -4k$

C. $(x + 2)^2 = -k$

D. $x^2 + k = 0$

9. What is the discriminant for $2x^2 - 8x - 3 = 0$? How many and what type of solutions does this equation have? (Do not actually solve.)

Solve each equation by any method.

10. $3 - \frac{16}{x} - \frac{12}{x^2} = 0$

11. $4x^2 + 7x - 3 = 0$

12. $9x^4 + 4 = 37x^2$

13. $12 = (2n + 1)^2 + (2n + 1)$


14. Solve $S = 4\pi r^2$ for r . (Leave \pm in your answer.)

Solve each problem.


15. Terry and Callie do word processing. For a certain prospectus, Callie can prepare it 2 hr faster than Terry can. If they work together, they can do the entire prospectus in 5 hr. How long will it take each of them working alone to prepare the prospectus? Round answers to the nearest tenth of an hour.
16. Qihong paddled a canoe 10 mi upstream and then paddled back to the starting point. If the rate of the current was 3 mph and the entire trip took $3\frac{1}{2}$ hr, what was Qihong's rate?

*This exercise requires knowledge of complex numbers.

17. Endre has a pool 24 ft long and 10 ft wide. He wants to construct a concrete walk around the pool. If he plans for the walk to be of uniform width and cover 152 ft^2 , what will the width of the walk be?







18. At a point 30 m from the base of a tower, the distance to the top of the tower is 2 m more than twice the height of the tower. Find the height of the tower.



19. Which one of the following figures most closely resembles the graph of

$$f(x) = a(x - h)^2 + k \quad \text{if } a < 0, h > 0, \text{ and } k < 0?$$



Graph each parabola. Identify the vertex, axis of symmetry, domain, and range.

20. $f(x) = \frac{1}{2}x^2 - 2$

21. $f(x) = -x^2 + 4x - 1$

22. $x = -(y - 2)^2 + 2$

23. Houston Community College is planning to construct a rectangular parking lot on land bordered on one side by a highway. The plan is to use 640 ft of fencing to fence off the other three sides. What should the dimensions of the lot be if the enclosed area is to be a maximum?

Solve each inequality, and graph the solution set.

24. $2x^2 + 7x > 15$

25. $\frac{5}{t - 4} \leq 1$

Chapters R-8 Cumulative Review Exercises

Perform the indicated operations.

1. $-8.84 - (-3.46)$
2. $\frac{7}{5} - \left(\frac{9}{10} - \frac{3}{2}\right)$
3. $|-12| + |13|$
4. Find 6% of 12.
5. Simplify $3 - 6(4^2 - 8)$.
6. Let $S = \left\{-\frac{7}{3}, -2, -\sqrt{3}, 0, 0.7, \sqrt{12}, \sqrt{-8}, 7, \frac{32}{3}\right\}$. List the elements of S that are elements of each set.
 - (a) Integers
 - (b) Rational numbers
 - (c) Real numbers
 - (d) Complex numbers

Solve each equation or inequality.

7. $7 - (4 + 3t) + 2t = -6(t - 2) - 5$
8. $|6x - 9| = |-4x + 2|$
9. $2x = \sqrt{\frac{5x + 2}{3}}$
10. $\frac{3}{x - 3} - \frac{2}{x - 2} = \frac{3}{x^2 - 5x + 6}$
11. $(r - 5)(2r + 3) = 1$
12. $x^4 - 5x^2 + 4 = 0$
13. $-2x + 4 \leq -x + 3$
14. $|3x - 7| \leq 1$
15. $x^2 - 4x + 3 < 0$
16. $\frac{3}{p + 2} > 1$

Graph each relation. Decide whether or not y can be expressed as a function f of x , and if so, give its domain and range, and write using function notation.

17. $4x - 5y = 15$
18. $4x - 5y < 15$
19. $y = -2(x - 1)^2 + 3$
20. Find the slope and intercepts of the line with equation $-2x + 7y = 16$.

Write an equation for the specified line. Express each equation in slope-intercept form.

21. Passing through $(2, -3)$ and parallel to the line with equation $5x + 2y = 6$
22. Passing through $(-4, 1)$ and perpendicular to the line with equation $5x + 2y = 6$

Solve each system of equations.

23. $2x - 4y = 10$
 $9x + 3y = 3$
24. $x + 2y = 5$
 $4y = -2x + 8$
25. $x + y + 2z = 3$
 $-x + y + z = -5$
 $2x + 3y - z = -8$

26. The two top-grossing films of 2017 were *Star Wars: The Last Jedi* and *Beauty and the Beast*. The two films together grossed \$1049 million. *Beauty and the Beast* grossed \$41 million less than *Star Wars: The Last Jedi*. How much did each film gross? (Data from Box Office Mojo.)

Write with positive exponents only. Assume that variables represent positive real numbers.

27. $\left(\frac{x^{-3}y^2}{x^5y^{-2}}\right)^{-1}$
28. $\frac{(4x^{-2})^2(2y^3)}{8x^{-3}y^5}$

Perform the indicated operations.

$$29. \left(\frac{2}{3}t + 9\right)^2$$

$$30. \text{ Divide } 4x^3 + 2x^2 - x + 26 \text{ by } x + 2.$$

Factor completely.

$$31. 24m^2 + 2m - 15$$

$$32. 4t^2 - 100$$

$$33. 8x^3 + 27y^3$$

$$34. 9x^2 - 30xy + 25y^2$$


Simplify. Express each answer in lowest terms. Assume denominators are nonzero.

$$35. \frac{5x + 2}{-6} \div \frac{15x + 6}{5}$$

$$36. \frac{3}{2-x} - \frac{5}{x} + \frac{6}{x^2 - 2x}$$

$$37. \frac{\frac{r}{s} - \frac{s}{r}}{\frac{r}{s} + 1}$$

38. Two cars left an intersection at the same time, one heading due south and the other due east. Later they were exactly 95 mi apart. The car heading east had gone 38 mi less than twice as far as the car heading south. How far had each car traveled?



Simplify each radical expression.

$$39. \sqrt[3]{\frac{27}{16}}$$

$$40. \frac{2}{\sqrt{7} - \sqrt{5}}$$

41. Graph $f(x) = 3x^2 + 12x + 6$. Give the vertex, axis of symmetry, domain, and range.

42. Solve $\frac{4 - 2t}{6 + t} > 0$. Give the solution set using interval notation.

STUDY SKILLS REMINDER

Have you begun to prepare for your final exam? **Review Study Skill 10, Preparing for Your Math Final Exam.**



INVERSE, EXPONENTIAL, AND LOGARITHMIC FUNCTIONS

Compound interest earned on money, intensities of sounds, and population growth and decay are some examples of applications of *exponential* and *logarithmic functions*.

- 9.1** Inverse Functions
- 9.2** Exponential Functions
- 9.3** Logarithmic Functions
- 9.4** Properties of Logarithms

- 9.5** Common and Natural Logarithms
- 9.6** Exponential and Logarithmic Equations; Further Applications

9.1 Inverse Functions

OBJECTIVES

- 1 Decide whether a function is one-to-one and, if it is, find its inverse.
- 2 Use the horizontal line test to determine whether a function is one-to-one.
- 3 Find the equation of the inverse of a function.
- 4 Graph f^{-1} , given the graph of f .

In this chapter we study two important types of functions, *exponential* and *logarithmic*. These functions are related: They are *inverses* of one another.

OBJECTIVE 1 Decide whether a function is one-to-one and, if it is, find its inverse.

Suppose we define the function

$$G = \{(-2, 2), (-1, 1), (0, 0), (1, 3), (2, 5)\}.$$

We can form another set of ordered pairs from G by interchanging the x - and y -values of each pair in G . We can call this set F , so

$$F = \{(2, -2), (1, -1), (0, 0), (3, 1), (5, 2)\}.$$

To show that these two sets are related as just described, F is called the *inverse* of G . For a function f to have an inverse, f must be a *one-to-one function*.

One-to-One Function

In a **one-to-one function**, each x -value corresponds to only one y -value, and each y -value corresponds to only one x -value.

VOCABULARY

- one-to-one function
- inverse of a function

STUDY SKILLS REMINDER

Make study cards to help you learn and remember the material in this chapter.

Review Study Skill 5,
Using Study Cards.

The function in **FIGURE 1(a)** is one-to-one. The function shown in **FIGURE 1(b)** is not one-to-one because the y -value 7 corresponds to *two* x -values, 2 and 3. That is, the ordered pairs $(2, 7)$ and $(3, 7)$ both belong to the function.




FIGURE 1

The *inverse* of any one-to-one function f is found by interchanging the components of the ordered pairs of f . The inverse of f is written f^{-1} . Read f^{-1} as “*the inverse of f* ” or “ *f -inverse*.”

CAUTION The symbol $f^{-1}(x)$ does **not** represent $\frac{1}{f(x)}$.

Inverse of a Function

The **inverse** of a one-to-one function f , written f^{-1} , is the set of all ordered pairs of the form (y, x) , where (x, y) belongs to f . **The inverse is formed by interchanging x and y , so the domain of f becomes the range of f^{-1} and the range of f becomes the domain of f^{-1} .**

For inverses f and f^{-1} , it follows that for all x in their domains,

$$(f \circ f^{-1})(x) = x \quad \text{and} \quad (f^{-1} \circ f)(x) = x.$$

NOW TRY EXERCISE 1

Determine whether each function is one-to-one. If it is, find the inverse.

- (a) $F = \{(0, 0), (1, 1), (4, 2), (9, 3)\}$
- (b) $G = \{(-1, -2), (0, 0), (1, -2), (2, -8)\}$
- (c) A Norwegian physiologist has developed a rule for predicting running times based on the time to run 5 km (5K). An example for one runner is shown here.

Distance	Time
1.5K	4:22
3K	9:18
5K	16:00
10K	33:40

Data from Stephen Seiler, Agder College, Kristiansand, Norway.

EXAMPLE 1 Finding Inverses of One-to-One Functions

Determine whether each function is one-to-one. If it is, find the inverse.

(a) $F = \{(3, 1), (0, 2), (2, 3), (4, 0)\}$

Every x -value in F corresponds to only one y -value, and every y -value corresponds to only one x -value, so F is a one-to-one function. The inverse function is found by interchanging the x - and y -values in each ordered pair.

$$F^{-1} = \{(1, 3), (2, 0), (3, 2), (0, 4)\}$$

The domain and range of F become the range and domain, respectively, of F^{-1} .

(b) $G = \{(-2, 1), (-1, 0), (0, 1), (1, 2), (2, 2)\}$

Each x -value in G corresponds to just one y -value. However, the y -value 1 corresponds to two x -values, -2 and 0 . Also, the y -value 2 corresponds to both 1 and 2. Because some y -values correspond to more than one x -value, G is not one-to-one and does not have an inverse.

(c) The table shows the number of days in which the air in the Los Angeles–Long Beach–Anaheim metropolitan area exceeded air-quality standards in recent years.

Year	Number of Days Exceeding Standards
2010	4
2011	14
2012	37
2013	25
2014	28
2015	38
2016	25

Data from U.S. Environmental Protection Agency.



Let f be the function defined in the table, with the years forming the domain and the number of days exceeding air-quality standards forming the range. Then f is not one-to-one because in two different years (2013 and 2016) the number of days exceeding air-quality standards was the same, 25.

NOW TRY

OBJECTIVE 2 Use the horizontal line test to determine whether a function is one-to-one.

By graphing a function and observing the graph, we can use the *horizontal line test* to determine whether the function is one-to-one.

Horizontal Line Test

A function is one-to-one if every horizontal line intersects the graph of the function at most once.

The horizontal line test follows from the definition of a one-to-one function. Any two points that lie on the same horizontal line have the same y -coordinate. No two ordered pairs that belong to a one-to-one function may have the same y -coordinate. Therefore, no horizontal line will intersect the graph of a one-to-one function more than once.



NOW TRY ANSWERS

1. (a) one-to-one;
 $F^{-1} = \{(0, 0), (1, 1), (2, 4), (3, 9)\}$
- (b) not one-to-one
- (c) one-to-one

Time	Distance
4:22	1.5K
9:18	3K
16:00	5K
33:40	10K

NOW TRY EXERCISE 2

Use the horizontal line test to determine whether each graph is the graph of a one-to-one function.



EXAMPLE 2 Using the Horizontal Line Test

Use the horizontal line test to determine whether each graph is the graph of a one-to-one function.




FIGURE 2

A horizontal line intersects the graph in **FIGURE 2** in more than one point. This function is not one-to-one.




FIGURE 3

Every horizontal line will intersect the graph in **FIGURE 3** in exactly one point. This function is one-to-one. **NOW TRY**

OBJECTIVE 3 Find the equation of the inverse of a function.

The inverse of a one-to-one function is found by interchanging the x - and y -values of each of its ordered pairs. The equation of the inverse of a function $y = f(x)$ is found in the same way.

Finding the Equation of the Inverse of $y = f(x)$

For a one-to-one function f defined by an equation $y = f(x)$, find the defining equation of the inverse function f^{-1} as follows.

Step 1 Interchange x and y .

Step 2 Solve for y .

Step 3 Replace y with $f^{-1}(x)$.

EXAMPLE 3 Finding Equations of Inverses

Determine whether each equation defines a one-to-one function. If so, find the equation that defines the inverse.

(a) $f(x) = 2x + 5$

The graph of $y = 2x + 5$ is a nonvertical line, so by the horizontal line test, f is a one-to-one function. Find the inverse as follows.

$$f(x) = 2x + 5$$

$$y = 2x + 5 \quad \text{Let } y = f(x).$$

Step 1 $x = 2y + 5$ Interchange x and y .

Step 2 $2y = x - 5$ Subtract 5.

$$y = \frac{x - 5}{2} \quad \text{Interchange sides. } \left. \begin{array}{l} \text{Solve for } y. \\ \text{Divide by 2.} \end{array} \right\}$$

Step 3 $f^{-1}(x) = \frac{x - 5}{2}$ Replace y with $f^{-1}(x)$.

These are equivalent forms of the same equation.

$$f^{-1}(x) = \frac{x}{2} - \frac{5}{2}, \quad \text{or} \quad f^{-1}(x) = \frac{1}{2}x - \frac{5}{2} \quad \frac{a-b}{c} = \frac{a}{c} - \frac{b}{c}$$

NOW TRY ANSWERS

- 2. (a) one-to-one function
- (b) not a one-to-one function

NOW TRY EXERCISE 3

Determine whether each equation defines a one-to-one function. If so, find the equation that defines the inverse.

- (a) $f(x) = 5x - 7$
- (b) $f(x) = (x + 1)^2$
- (c) $f(x) = x^3 - 4$

Thus, f^{-1} is a linear function. In the given function

$$f(x) = 2x + 5,$$

we start with a value of x , *multiply* by 2, and *add* 5. In the equation for the inverse

$$f^{-1}(x) = \frac{x - 5}{2}, \quad \text{One form of } f^{-1}(x)$$

we *subtract* 5, and then *divide* by 2. This shows how an inverse is used to “undo” what a function does to the variable x .

(b) $y = x^2 + 2$

This equation has a vertical parabola as its graph, so some horizontal lines will intersect the graph at two points. For example, both $x = 1$ and $x = -1$ correspond to $y = 3$. See **FIGURE 4**.




FIGURE 4

Because of the x^2 -term in $y = x^2 + 2$, there are many pairs of x -values that correspond to the same y -value. The function is not one-to-one and does not have an inverse. If we try to find the equation of an inverse, we obtain the following.

$$y = x^2 + 2$$

$$x = y^2 + 2 \quad \text{Interchange } x \text{ and } y.$$

$$y^2 = x - 2 \quad \text{Solve for } y.$$

$$y = \pm \sqrt{x - 2} \quad \text{Square root property}$$

Remember both roots.

The last step shows that there are two y -values for each choice of x in $(2, \infty)$ —that is, for $x > 2$ —so the function is not one-to-one. It does not have an inverse.

(c) $f(x) = (x - 2)^3$

A cubing function like this is one-to-one. See the graph in **FIGURE 5**.




FIGURE 5

NOW TRY ANSWERS

3. (a) one-to-one function;

$$f^{-1}(x) = \frac{x + 7}{5}, \text{ or}$$

$$f^{-1}(x) = \frac{1}{5}x + \frac{7}{5}$$

(b) not a one-to-one function

(c) one-to-one function;

$$f^{-1}(x) = \sqrt[3]{x + 4}$$

Step 1

$$x = (y - 2)^3 \quad \text{Interchange } x \text{ and } y.$$

Step 2

$$\sqrt[3]{x} = \sqrt[3]{(y - 2)^3} \quad \text{Take the cube root on each side.}$$

$$\sqrt[3]{x} = y - 2 \quad \sqrt[3]{a^3} = a \quad \left. \begin{array}{l} \text{Solve for } y. \end{array} \right\}$$

$$y = \sqrt[3]{x} + 2 \quad \text{Add 2. Interchange sides.}$$

Step 3

$$f^{-1}(x) = \sqrt[3]{x} + 2 \quad \text{Replace } y \text{ with } f^{-1}(x).$$

NOW TRY
EXERCISE 4

Find $f^{-1}(x)$ for

$$f(x) = \frac{x+3}{x-4}, \quad x \neq 4.$$

EXAMPLE 4 Using Factoring to Find an Inverse

It is shown in standard college algebra texts that the following function is one-to-one.

$$f(x) = \frac{x+1}{x-2}, \quad x \neq 2$$

Find $f^{-1}(x)$.

$$f(x) = \frac{x+1}{x-2}, \quad x \neq 2 \quad \text{Given equation}$$

$$y = \frac{x+1}{x-2}, \quad x \neq 2 \quad \text{Replace } f(x) \text{ with } y.$$

Step 1 $x = \frac{y+1}{y-2}, \quad y \neq 2 \quad \text{Interchange } x \text{ and } y.$

Step 2

$x(y-2) = y+1$	Multiply by $y-2$.	}	Solve for y .
$xy - 2x = y + 1$	Distributive property		
$xy - y = 2x + 1$	Subtract y . Add $2x$.		
$y(x-1) = 2x + 1$	Factor out y .		
$y = \frac{2x+1}{x-1}$	Divide by $x-1$.		

Step 3 $f^{-1}(x) = \frac{2x+1}{x-1}, \quad x \neq 1 \quad \text{Replace } y \text{ with } f^{-1}(x). \text{ Note the restriction.}$

NOW TRY

OBJECTIVE 4 Graph f^{-1} , given the graph of f .

One way to graph the inverse of a function f whose equation is given follows.

Graphing the Inverse

- Step 1** Find several ordered pairs that belong to f .
- Step 2** Interchange x and y to obtain ordered pairs that belong to f^{-1} .
- Step 3** Plot those points, and sketch the graph of f^{-1} through them.

We can also select points on the graph of f and use *symmetry* to find corresponding points on the graph of f^{-1} .

For example, suppose the point (a, b) shown in **FIGURE 6** belongs to a one-to-one function f . Then the point (b, a) belongs to f^{-1} . The line segment connecting the points (a, b) and (b, a) is perpendicular to, and cut in half by, the line $y = x$. The points (a, b) and (b, a) are “mirror images” of each other with respect to $y = x$.




FIGURE 6


We can find the graph of f^{-1} from the graph of f by locating the mirror image of each point in f with respect to the line $y = x$.

NOW TRY ANSWER

4. $f^{-1}(x) = \frac{4x+3}{x-1}, \quad x \neq 1$

NOW TRY
EXERCISE 5

Graph the inverse of the function labeled f in the figure.



EXAMPLE 5 Graphing Inverses of Functions

Graph the inverse of each function labeled f in the figures.

(a) **FIGURE 7(a)** shows the graph of a one-to-one function f . The points

$$\left(-1, \frac{1}{2}\right), (0, 1), (1, 2), \text{ and } (2, 4) \quad \text{Points on } f$$

lie on its graph. Interchange x and y to obtain ordered pairs that belong to f^{-1} .

$$\left(\frac{1}{2}, -1\right), (1, 0), (2, 1), \text{ and } (4, 2) \quad \text{Points on } f^{-1}$$

Plot these points, and sketch the graph of f^{-1} through them. See **FIGURE 7(b)**.




FIGURE 7

(b) Each function f (shown in blue) in **FIGURES 8** and **9** is a one-to-one function.

Each inverse f^{-1} is shown in red. In both cases, the graph of f^{-1} is a reflection of the graph of f across the line $y = x$.




FIGURE 8





FIGURE 9

NOW TRY

NOW TRY ANSWER



9.1 Exercises

FOR
EXTRA
HELP




MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Choose the correct response.

- If a function is made up of ordered pairs in such a way that the same y -value appears in a correspondence with two different x -values, then
 - the function is one-to-one
 - the function is not one-to-one
 - its graph does not pass the vertical line test
 - it has an inverse function associated with it.
- Which equation defines a one-to-one function? Explain why the others do not, using specific examples.
 - $f(x) = x$
 - $f(x) = x^2$
 - $f(x) = |x|$
 - $f(x) = -x^2 + 2x - 1$

3. Only one of the graphs illustrates a one-to-one function. Which one is it? (See Example 2.)



4. If a function f is one-to-one and the point (p, q) lies on the graph of f , then which point *must* lie on the graph of f^{-1} ?

- A. $(-p, q)$ B. $(-q, -p)$ C. $(p, -q)$ D. (q, p)

Answer each question.

5. The table shows fat content of various menu items at McDonald's. If the set of menu items is the domain and the set of fat contents is the range of a function, is it one-to-one? Why or why not?

Menu Item	Fat Content (in grams)
Chicken McNuggets (20 pieces)	60
Quarter Pounder with Cheese	43
Big Breakfast with Egg Whites	41
Bacon Clubhouse Burger	41
McFlurry with Reese's	37
Sausage Biscuit with Egg	33

Data from www.thebalancesmb.com

6. The table shows the most-visited social networking web sites, by number of visitors in millions, in June 2017. If the set of web sites is the domain and the set of numbers of visitors is the range of a function, is it one-to-one? Why or why not?

Social Networking Web Site	Number of Visitors (in millions)
Facebook	202
Instagram	121
Twitter	110
LinkedIn	103
Snapchat	95
Pinterest	86

Data from comScore, Inc.

7. The road mileage between Denver, Colorado, and several selected U.S. cities is shown in the table. If we consider this a function that pairs each city with a distance, is it one-to-one? How could we change the answer to this question by adding 1 mile to one of the distances shown?

City	Distance to Denver (in miles)
Atlanta	1398
Dallas	781
Indianapolis	1058
Kansas City, MO	600
Los Angeles	1059

8. The table lists caffeine amounts in several popular 12-oz sodas. If the set of sodas is the domain and the set of caffeine amounts is the range of a function, is it one-to-one? Why or why not?

Soda	Caffeine (in mg)
Mountain Dew	54
Diet Coke	46
Sunkist Orange Soda	41
Diet Pepsi-Cola	34
Coca-Cola Classic	34

Data from www.caffeineinformer.com

Determine whether each function is one-to-one. If it is, find the inverse. See Examples 1 and 3.

9. $\{(3, 6), (2, 10), (5, 12)\}$ 10. $\left\{(-1, 3), (0, 5), (5, 0), \left(7, -\frac{1}{2}\right)\right\}$

11. $\{(-1, 3), (2, 7), (4, 3), (5, 8)\}$ 12. $\{(-8, 6), (-4, 3), (0, 6), (5, 10)\}$

13. $\{(0, 4.5), (2, 8.6), (4, 12.7)\}$ 14. $\{(1, 5.8), (2, 8.8), (3, 8.5)\}$

15. $f(x) = x + 3$ 16. $f(x) = x + 8$ 17. $f(x) = -\frac{1}{2}x - 2$

18. $f(x) = -\frac{1}{4}x - 8$ 19. $f(x) = 2x + 4$ 20. $f(x) = 3x + 1$

21. $g(x) = -4x + 3$ 22. $g(x) = -6x - 8$ 23. $f(x) = 5$

24. $f(x) = -7$ 25. $f(x) = \sqrt{x-3}, x \geq 3$ 26. $f(x) = \sqrt{x+2}, x \geq -2$

27. $f(x) = \sqrt{x+6}, x \geq -6$ 28. $f(x) = \sqrt{x-4}, x \geq 4$

29. $f(x) = 3x^2 + 2$ 30. $f(x) = 4x^2 - 1$ 31. $g(x) = (x+1)^3$

32. $g(x) = (x-4)^3$ 33. $f(x) = x^3 - 4$ 34. $f(x) = x^3 + 5$

Each function is one-to-one. Find its inverse. See Example 4.

35. $f(x) = \frac{x+4}{x+2}, x \neq -2$ 36. $f(x) = \frac{x+3}{x+5}, x \neq -5$

37. $f(x) = \frac{4x-2}{x+5}, x \neq -5$ 38. $f(x) = \frac{5x-10}{x+4}, x \neq -4$

39. $f(x) = \frac{-2x+1}{2x-5}, x \neq \frac{5}{2}$ 40. $f(x) = \frac{-3x+2}{3x-4}, x \neq \frac{4}{3}$




Concept Check Let $f(x) = 2^x$. This function is one-to-one. Find each value.




41. (a) $f(3)$ 42. (a) $f(4)$ 43. (a) $f(0)$ 44. (a) $f(-2)$
 (b) $f^{-1}(8)$ (b) $f^{-1}(16)$ (b) $f^{-1}(1)$ (b) $f^{-1}\left(\frac{1}{4}\right)$

Graphs of selected functions are given in the following exercises.

(a) Use the horizontal line test to determine whether each function graphed is one-to-one. See Example 2.

(b) If the function is one-to-one, graph its inverse. See Example 5.





Each of the following functions is one-to-one. Graph the function as a solid line (or curve), and then graph its inverse on the same set of axes as a dashed line (or curve). Complete any tables to help graph the functions. See Example 5.

51. $f(x) = 2x - 1$ 52. $f(x) = 2x + 3$ 53. $f(x) = -4x$ 54. $f(x) = -2x$

55. $f(x) = \sqrt{x}$, 56. $f(x) = -\sqrt{x}$, 57. $f(x) = x^3 - 2$ 58. $f(x) = x^3 + 3$

$x \geq 0$

x	f(x)
0	
1	
4	

$x \geq 0$

x	f(x)
0	
1	
4	

x	f(x)
-1	
0	
1	
2	

x	f(x)
-2	
-1	
0	
1	

RELATING CONCEPTS For Individual or Group Work (Exercises 59–62)

Inverse functions can be used to send and receive coded information. A simple example might use the function

$f(x) = 2x + 5$. (Note that it is one-to-one.)

Suppose that each letter of the alphabet is assigned a numerical value according to its position, as follows.

A	1	G	7	L	12	Q	17	V	22
B	2	H	8	M	13	R	18	W	23
C	3	I	9	N	14	S	19	X	24
D	4	J	10	O	15	T	20	Y	25
E	5	K	11	P	16	U	21	Z	26
F	6								



This is an Enigma machine used by the Germans in World War II to send coded messages.

Using the function, the word ALGEBRA would be encoded as

7 29 19 15 9 41 7,

because

$f(A) = f(1) = 2(1) + 5 = 7$, $f(L) = f(12) = 2(12) + 5 = 29$, and so on.

The message would then be decoded using the inverse of f , which is $f^{-1}(x) = \frac{x-5}{2}$.

$f^{-1}(7) = \frac{7-5}{2} = 1 = A$, $f^{-1}(29) = \frac{29-5}{2} = 12 = L$, and so on.

Work Exercises 59–62 in order.

59. Suppose that you are an agent for a detective agency. Today's encoding function is $f(x) = 4x - 5$. Find the rule for f^{-1} algebraically.

60. You receive the following coded message today. (Read across from left to right.)

47 95 7 -1 43 7 79 43 -1 75 55 67 31 71 75 27
 15 23 67 15 -1 75 15 71 75 75 27 31 51 23 71
 31 51 7 15 71 43 31 7 15 11 3 67 15 -1 11

Use the letter/number assignment described on the previous page to decode the message.

61. Why is a one-to-one function essential in this encoding/decoding process?

62. Use $f(x) = x^3 + 4$ to encode your name, using the above letter/number assignment.

9.2 Exponential Functions

OBJECTIVES

- 1 Evaluate exponential expressions using a calculator.
- 2 Define and graph exponential functions.
- 3 Solve exponential equations of the form $a^x = a^k$ for x .
- 4 Use exponential functions in applications involving growth or decay.

VOCABULARY

- exponential function with base a
- asymptote
- exponential equation

NOW TRY EXERCISE 1

Use a calculator to approximate each exponential expression to three decimal places.

- (a) $2^{3.1}$ (b) $2^{-1.7}$ (c) $2^{1/4}$

NOW TRY ANSWERS

1. (a) 8.574 (b) 0.308 (c) 1.189

OBJECTIVE 1 Evaluate exponential expressions using a calculator.

Consider the exponential expression 2^x for rational values of x .

$$2^3 = 8, \quad 2^{-1} = \frac{1}{2}, \quad 2^{1/2} = \sqrt{2}, \quad 2^{3/4} = \sqrt[4]{2^3} = \sqrt[4]{8} \quad \text{Examples of } 2^x \text{ for rational } x$$

In more advanced courses it is shown that 2^x exists for all real number values of x , both rational and irrational. We can use a calculator to find approximations of exponential expressions that are not easily determined.

EXAMPLE 1 Evaluating Exponential Expressions

Use a calculator to approximate each exponential expression to three decimal places.

- (a) $2^{1.6}$ (b) $2^{-1.3}$ (c) $2^{1/3}$

FIGURE 10 shows how a TI-84 Plus calculator approximates these values. The display shows more decimal places than we usually need, so we round to three decimal places as directed.

$$2^{1.6} \approx 3.031, \quad 2^{-1.3} \approx 0.406, \quad 2^{1/3} \approx 1.260$$




FIGURE 10

NOW TRY

OBJECTIVE 2 Define and graph exponential functions.

The definition of an exponential function assumes that a^x exists for all real numbers x .

Exponential Function

For $a > 0$, $a \neq 1$, and all real numbers x ,

$$f(x) = a^x$$

defines the **exponential function with base a** .

When graphing an exponential function of the form $f(x) = a^x$, pay particular attention to whether $a > 1$ or $0 < a < 1$.

NOW TRY EXERCISE 2
Graph $f(x) = 4^x$.

EXAMPLE 2 Graphing an Exponential Function ($a > 1$)

Graph $f(x) = 2^x$. Then compare it to the graph of $F(x) = 5^x$.

Choose some values of x , and find the corresponding values of $f(x) = 2^x$. Plotting these points and drawing a smooth curve through them gives the darker graph shown in **FIGURE 11**. This graph is typical of the graph of an exponential function of the form $f(x) = a^x$, where $a > 1$.

The larger the value of a , the faster the graph rises.

To see this, compare the graph of $F(x) = 5^x$ with the graph of $f(x) = 2^x$ in **FIGURE 11**. When graphing such functions, be sure to plot a sufficient number of points to see how rapidly the graph rises.




FIGURE 11

Exponential function with base $a > 1$

Domain: $(-\infty, \infty)$

Range: $(0, \infty)$

y-intercept: $(0, 1)$

The function is one-to-one, and its graph rises from left to right.

The vertical line test assures us that the graphs in **FIGURE 11** represent functions. These graphs show an important characteristic of exponential functions where $a > 1$.

As x gets larger, y increases at a faster and faster rate.

NOW TRY

NOW TRY EXERCISE 3

Graph $g(x) = \left(\frac{1}{10}\right)^x$.

EXAMPLE 3 Graphing an Exponential Function ($0 < a < 1$)

Graph $g(x) = \left(\frac{1}{2}\right)^x$.

Find and plot some points on the graph. The graph in **FIGURE 12** is similar to that of $f(x) = 2^x$ (**FIGURE 11**) with the same domain and range, except that here *as x gets larger, y decreases*. This graph is typical of the graph of an exponential function of the form $f(x) = a^x$, where $0 < a < 1$.




FIGURE 12

Exponential function with base $0 < a < 1$



Domain: $(-\infty, \infty)$

Range: $(0, \infty)$

y-intercept: $(0, 1)$

The function is one-to-one, and its graph falls from left to right.

NOW TRY ANSWERS



NOW TRY

! CAUTION The graph of an exponential function of the form

$$f(x) = a^x$$

approaches the x -axis, but does **not** touch it. Recall that such a line is called an **asymptote**.

Characteristics of the Graph of $f(x) = a^x$

1. The graph contains the point $(0, 1)$, which is its y -intercept.
2. The function is one-to-one.
 - When $a > 1$, the graph *rises* from left to right. (See FIGURE 11.)
 - When $0 < a < 1$, the graph *falls* from left to right. (See FIGURE 12.)

In both cases, the graph goes from the second quadrant to the first.
3. The graph approaches the x -axis but never touches it—that is, the x -axis is an asymptote.
4. The domain is $(-\infty, \infty)$, and the range is $(0, \infty)$.

NOW TRY
EXERCISE 4
Graph $f(x) = 4^{2x-1}$.


EXAMPLE 4 Graphing a More Complicated Exponential Function

Graph $f(x) = 3^{2x-4}$.

Find some ordered pairs. We let $x = 0$ and $x = 2$ and find values of $f(x)$, or y .

$f(x) = 3^{2x-4}$	$f(x) = 3^{2x} - 4$
$y = 3^{2(0)-4}$ Let $x = 0$.	$y = 3^{2(2)-4}$ Let $x = 2$.
$y = 3^{-4}$	$y = 3^0$
$y = \frac{1}{81}$ $a^{-n} = \frac{1}{a^n}$	$y = 1$ $a^0 = 1$

These ordered pairs, $(0, \frac{1}{81})$ and $(2, 1)$, along with the other ordered pairs shown in the table, lead to the graph in FIGURE 13.



The graph of $f(x) = 3^{2x-4}$ is similar to the graph of $f(x) = 3^x$ except that it is shifted to the right and rises more rapidly.

NOW TRY ANSWER




FIGURE 13

NOTE The two restrictions on the value of a in the definition of an exponential function $f(x) = a^x$ are important.

1. The restriction $a > 0$ is necessary so that the function can be defined for *all* real numbers x . Letting a be negative, such as $a = -2$, and letting $x = \frac{1}{2}$ gives the expression $(-2)^{1/2}$, which is not a real number.
2. The restriction $a \neq 1$ is necessary because 1 raised to *any* power is equal to 1, resulting in the linear function $f(x) = 1$.

OBJECTIVE 3 Solve exponential equations of the form $a^x = a^k$ for x .

Up to this point, we have solved only equations that had the variable as a base, like $x^2 = 8$. In these equations, all exponents have been constants. An **exponential equation** is an equation that has a variable in an exponent, such as

$$9^x = 27.$$

We can use the following property to solve many exponential equations.

Property for Solving an Exponential Equation

For $a > 0$ and $a \neq 1$, if $a^x = a^y$ then $x = y$.

This property would not necessarily be true if $a = 1$.

Solving an Exponential Equation

- Step 1 Both sides must have the same base.** If the two sides of the equation do not have the same base, express each as a power of the same base if possible.
- Step 2 Simplify exponents** if necessary, using the rules of exponents.
- Step 3 Set exponents equal** using the property given in this section.
- Step 4 Solve** the equation obtained in Step 3.

EXAMPLE 5 Solving an Exponential Equation

Solve $9^x = 27$.

$$9^x = 27$$

Step 1

$$(3^2)^x = 3^3$$

Write with the same base;
 $9 = 3^2$ and $27 = 3^3$.

Step 2

$$3^{2x} = 3^3$$

Power rule for exponents

Step 3

$$2x = 3$$

If $a^x = a^y$, then $x = y$.

Step 4

$$x = \frac{3}{2}$$

Solve for x .

CHECK Substitute $\frac{3}{2}$ for x : $9^x = 9^{3/2} = (9^{1/2})^3 = 3^3 = 27$. ✓ True

The solution set is $\left\{\frac{3}{2}\right\}$.

NOW TRY EXERCISE 5

Solve $8^x = 16$.

NOW TRY ANSWER

5. $\left\{\frac{4}{3}\right\}$


**NOW TRY
EXERCISE 6**

Solve each equation.

(a) $3^{2x-1} = 27^{x+4}$

(b) $5^x = \frac{1}{625}$

(c) $\left(\frac{2}{7}\right)^x = \frac{343}{8}$

EXAMPLE 6 Solving Exponential Equations

Solve each equation.

(a) $4^{3x-1} = 16^{x+2}$

Be careful multiplying the exponents.

$4^{3x-1} = (4^2)^{x+2}$ Write with the same base; $16 = 4^2$ (Step 1)

$4^{3x-1} = 4^{2x+4}$ Power rule for exponents (Step 2)

$3x - 1 = 2x + 4$ Set the exponents equal. (Step 3)

$x = 5$ Subtract $2x$. Add 1. (Step 4)

CHECK $4^{3x-1} = 16^{x+2}$

$4^{3(5)-1} \stackrel{?}{=} 16^{5+2}$ Substitute. Let $x = 5$.

$4^{14} \stackrel{?}{=} 16^7$ Perform the operations in the exponents.

$4^{14} \stackrel{?}{=} (4^2)^7$ $16 = 4^2$

$4^{14} = 4^{14}$ ✓ True

The solution set is $\{5\}$.

(b) $6^x = \frac{1}{216}$

$6^x = \frac{1}{6^3}$ $216 = 6^3$

$6^x = 6^{-3}$ Write with the same base; $\frac{1}{6^3} = 6^{-3}$.

$x = -3$ Set exponents equal.

CHECK Substitute -3 for x .

$6^x = 6^{-3} = \frac{1}{6^3} = \frac{1}{216}$ ✓ True

The solution set is $\{-3\}$.

(c) $\left(\frac{2}{3}\right)^x = \frac{9}{4}$

$\left(\frac{2}{3}\right)^x = \left(\frac{4}{9}\right)^{-1}$ $\frac{9}{4} = \left(\frac{4}{9}\right)^{-1}$

$\left(\frac{2}{3}\right)^x = \left[\left(\frac{2}{3}\right)^2\right]^{-1}$ Write with the same base.

$\left(\frac{2}{3}\right)^x = \left(\frac{2}{3}\right)^{-2}$ Power rule for exponents

$x = -2$ Set exponents equal.

Check that the solution set is $\{-2\}$.**NOW TRY ANSWERS**

6. (a) $\{-13\}$ (b) $\{-4\}$

(c) $\{-3\}$


NOW TRY
EXERCISE 7

Use the function in **Example 7** to approximate average annual carbon dioxide concentration in 2010, to the nearest unit.

OBJECTIVE 4 Use exponential functions in applications involving growth or decay.

EXAMPLE 7 Applying an Exponential Growth Function

The graph in **FIGURE 14** shows average annual concentration of carbon dioxide (in parts per million) in the air. This concentration is increasing exponentially.



Data from National Oceanic and Atmospheric Administration.

FIGURE 14

The data in **FIGURE 14** are approximated by the exponential function

$$f(x) = 275.8 + 1.5172(1.0161)^x,$$

where x is number of years since 1750. Use this function to approximate average annual concentration of carbon dioxide in parts per million, to the nearest unit, for each year.

(a) 1900

Because x represents number of years since 1750, $x = 1900 - 1750 = 150$.

$$f(x) = 275.8 + 1.5172(1.0161)^x \quad \text{Given function}$$

$$f(150) = 275.8 + 1.5172(1.0161)^{150} \quad \text{Let } x = 150.$$

$$f(150) \approx 292 \quad \text{Evaluate with a calculator.}$$

The concentration in 1900 was approximately 292 parts per million.

(b) 2000

$$f(x) = 275.8 + 1.5172(1.0161)^x \quad \text{Given function}$$

$$f(250) = 275.8 + 1.5172(1.0161)^{250} \quad x = 2000 - 1750 = 250$$

$$f(250) \approx 358 \quad \text{Evaluate with a calculator.}$$

The concentration in 2000 was approximately 358 parts per million.

NOW TRY

EXAMPLE 8 Applying an Exponential Decay Function

Atmospheric pressure (in millibars) at a given altitude x , in meters, can be approximated by the exponential function

$$f(x) = 1038(1.000134)^{-x}, \quad \text{for values of } x \text{ between 0 and 10,000.}$$

Because the base is greater than 1 and the coefficient of x in the exponent is negative, function values decrease as x increases. This means that as altitude increases, atmospheric pressure decreases. (Data from Miller, A. and J. Thompson, *Elements of Meteorology*, Fourth Edition, Charles E. Merrill Publishing Company.)

NOW TRY ANSWER
7. 372 parts per million


NOW TRY EXERCISE 8

Use the function in **Example 8** to approximate the pressure at 6000 m, to the nearest unit.

(a) According to this function, what is the pressure at ground level?

At ground level, $x = 0$.	$f(x) = 1038(1.000134)^{-x}$	Given function
	$f(0) = 1038(1.000134)^{-0}$	Let $x = 0$.
	$f(0) = 1038(1)$	$a^0 = 1$
	$f(0) = 1038$	Identity property

The pressure is 1038 millibars.

(b) What is the pressure at 5000 m, to the nearest unit?

$f(x) = 1038(1.000134)^{-x}$	Given function
$f(5000) = 1038(1.000134)^{-5000}$	Let $x = 5000$.
$f(5000) \approx 531$	Evaluate with a calculator.

The pressure is approximately 531 millibars.

NOW TRY 

NOTE The function in **Example 8** is equivalent to

$$f(x) = 1038 \left(\frac{1}{1.000134} \right)^x.$$

In this form, the base a satisfies the condition $0 < a < 1$.

NOW TRY ANSWER


8. 465 millibars

9.2 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Choose the correct response.

- For an exponential function $f(x) = a^x$, if $a > 1$, then the graph (*rises / falls*) from left to right.
- For an exponential function $f(x) = a^x$, if $0 < a < 1$, then the graph (*rises / falls*) from left to right.
- Which point lies on the graph of $f(x) = 3^x$?
 A. $(1, 0)$ B. $(3, 1)$ C. $(0, 1)$ D. $\left(\sqrt{3}, \frac{1}{3}\right)$
- The asymptote of the graph of $f(x) = a^x$
 A. is the x -axis B. is the y -axis
 C. has equation $x = 1$ D. has equation $y = 1$.
- Which statement is true?
 A. The point $\left(\frac{1}{2}, \sqrt{5}\right)$ lies on the graph of $f(x) = 5^x$.
 B. $f(x) = 5^x$ is not a one-to-one function.
 C. The y -intercept of the graph of $f(x) = 5^x$ is $(0, 5)$.
 D. The graph of $y = 5^x$ rises at a faster rate than the graph of $y = 10^x$.

6. Which statement is false?

A. The domain of the function $f(x) = \left(\frac{1}{4}\right)^x$ is $(-\infty, \infty)$.

B. The graph of the function $f(x) = \left(\frac{1}{4}\right)^x$ has one x -intercept.

C. The range of the function $f(x) = \left(\frac{1}{4}\right)^x$ is $(0, \infty)$.

D. The point $(-2, 16)$ lies on the graph of $f(x) = \left(\frac{1}{4}\right)^x$.

Use a calculator to approximate each exponential expression to three decimal places. See Example 1.

7. $2^{1.9}$

8. $2^{2.7}$

9. $2^{-1.54}$

10. $2^{-1.88}$

11. $10^{0.3}$

12. $10^{0.5}$

13. $4^{1/3}$

14. $6^{1/5}$

15. $\left(\frac{1}{3}\right)^{1.5}$

16. $\left(\frac{1}{3}\right)^{2.4}$

17. $\left(\frac{1}{4}\right)^{-3.1}$

18. $\left(\frac{1}{4}\right)^{-1.4}$

Graph each exponential function. See Examples 2–4.

19. $f(x) = 3^x$

20. $f(x) = 5^x$

21. $g(x) = \left(\frac{1}{3}\right)^x$

22. $g(x) = \left(\frac{1}{5}\right)^x$

23. $f(x) = 4^{-x}$

24. $f(x) = 6^{-x}$

25. $f(x) = 2^{2x-2}$

26. $f(x) = 2^{2x+1}$

27. **Concept Check** A student incorrectly solved the following equation as shown.

$$2^x = 32 \quad \text{Given equation}$$

$$\frac{2^x}{2} = \frac{32}{2} \quad \text{Divide by 2.}$$

$$x = 16$$

WHAT WENT WRONG? Give the correct solution set.

28. **Concept Check** A student incorrectly solved the following equation as shown.

$$3^x = 81 \quad \text{Given equation}$$

$$3^x - 3 = 81 - 3 \quad \text{Subtract 3.}$$

$$x = 78$$

WHAT WENT WRONG? Give the correct solution set.

Solve each equation. See Examples 5 and 6.

29. $6^x = 36$

30. $8^x = 64$

31. $100^x = 1000$

32. $8^x = 4$

33. $16^x = 64$

34. $8^x = 32$

35. $4^{x-5} = 64^{2x}$

36. $125^{3x} = 5^{2x-7}$

37. $16^{2x+1} = 64^{x+3}$

38. $9^{2x-8} = 27^{x-4}$

39. $5^x = \frac{1}{125}$

40. $3^x = \frac{1}{81}$

41. $9^x = \frac{1}{27}$

42. $8^x = \frac{1}{32}$

43. $5^x = 0.2$

44. $10^x = 0.1$

45. $\left(\frac{3}{2}\right)^x = \frac{8}{27}$

46. $\left(\frac{4}{3}\right)^x = \frac{27}{64}$

47. $\left(\frac{5}{4}\right)^x = \frac{16}{25}$

48. $\left(\frac{3}{2}\right)^x = \frac{16}{81}$

59. A small business estimates that the value $V(t)$ of a copy machine is decreasing according to the exponential function

$$V(t) = 5000(2)^{-0.15t},$$

where t is the number of years that have elapsed since the machine was purchased, and $V(t)$ is in dollars.

- What was the original value of the machine?
- What is the value of the machine 5 yr after purchase, to the nearest dollar?
- What is the value of the machine 10 yr after purchase, to the nearest dollar?
- Graph the function.



60. Refer to the exponential function in **Exercise 59**.

- When will the value of the machine be \$2500? (*Hint*: Let $V(t) = 2500$, divide both sides by 5000, and use the method of **Example 5**.)
- When will the value of the machine be \$1250?

9.3 Logarithmic Functions

OBJECTIVES

- Define a logarithm.
- Convert between exponential and logarithmic forms, and evaluate logarithms.
- Solve logarithmic equations of the form $\log_a b = k$ for a , b , or k .
- Use the definition of logarithm to simplify logarithmic expressions.
- Define and graph logarithmic functions.
- Use logarithmic functions in applications involving growth or decay.

OBJECTIVE 1 Define a logarithm.

The graph of $y = 2^x$ is the blue curve in **FIGURE 15**. Because $y = 2^x$ defines a one-to-one function, it has an inverse. Interchanging x and y gives

$$x = 2^y, \quad \text{the inverse of } y = 2^x. \quad \text{Roles of } x \text{ and } y \text{ are interchanged.}$$

The graph of the inverse is found by reflecting the graph of $y = 2^x$ across the line $y = x$. The graph of the inverse $x = 2^y$ is the red curve in **FIGURE 15**.




FIGURE 15

We can also write the equation of the red curve using a new notation that involves the concept of *logarithm*.

VOCABULARY


- logarithm
- logarithmic equation
- logarithmic function with base a

Logarithm

For all positive numbers a , where $a \neq 1$, and all positive real numbers x ,

$$y = \log_a x \quad \text{is equivalent to} \quad x = a^y.$$

The abbreviation **log** is used for the word **logarithm**. Read $\log_a x$ as “*the logarithm of x with base a*” or “*the base a logarithm of x.*” To remember the location of the base and the exponent in each form, refer to the following diagrams.



Meaning of $\log_a x$

A logarithm is an exponent. *The expression $\log_a x$ represents the exponent to which the base a must be raised to obtain x .*

OBJECTIVE 2 Convert between exponential and logarithmic forms, and evaluate logarithms.

We can use the definition of logarithm to carry out these conversions.

NOW TRY EXERCISE 1

- (a) Write $6^3 = 216$ in logarithmic form.
 (b) Write $\log_{64} 4 = \frac{1}{3}$ in exponential form.

EXAMPLE 1 Converting between Exponential and Logarithmic Forms

The table shows several pairs of equivalent forms.

Exponential Form	Logarithmic Form
$3^2 = 9$	$\log_3 9 = 2$
$\left(\frac{1}{5}\right)^{-2} = 25$	$\log_{1/5} 25 = -2$
$10^5 = 100,000$	$\log_{10} 100,000 = 5$
$4^{-3} = \frac{1}{64}$	$\log_4 \frac{1}{64} = -3$

$y = \log_a x$
is equivalent to
 $x = a^y.$

NOW TRY

NOW TRY EXERCISE 2

Use a calculator to approximate each logarithm to four decimal places.

- (a) $\log_2 7$ (b) $\log_5 8$
 (c) $\log_{1/3} 12$ (d) $\log_{10} 18$

EXAMPLE 2 Evaluating Logarithms

Use a calculator to approximate each logarithm to four decimal places.

- (a) $\log_2 5$ (b) $\log_3 12$ (c) $\log_{1/2} 12$ (d) $\log_{10} 20$

FIGURE 16 shows how a TI-84 Plus calculator approximates the logarithms in parts (a)–(c).




FIGURE 16




FIGURE 17

FIGURE 17 shows the approximation for the expression $\log_{10} 20$ in part (d). Notice that the second display, which indicates $\log 20$ (with no base shown), gives the same result. We shall see in a later section that when no base is indicated, the base is understood to be 10. A base 10 logarithm is a *common logarithm*.

NOW TRY

NOW TRY ANSWERS

1. (a) $\log_6 216 = 3$
 (b) $64^{1/3} = 4$
 2. (a) 2.8074 (b) 1.2920
 (c) -2.2619 (d) 1.2553


NOW TRY EXERCISE 3

Solve each equation.

- (a) $\log_2 x = -5$
 (b) $\log_{3/2} (2x - 1) = 3$
 (c) $\log_x 10 = 2$
 (d) $\log_{125} \sqrt[3]{5} = x$

(c) $\log_x 3 = 2$

$x^2 = 3$

Write in exponential form.

Be careful here.
 $-\sqrt{3}$ is extraneous.

$x = \pm \sqrt{3}$

Take square roots.

Only the *principal* square root $\sqrt{3}$ satisfies the equation because the base must be a positive number.

CHECK $\log_x 3 = 2$

$\log_{\sqrt{3}} 3 \stackrel{?}{=} 2$ Let $x = \sqrt{3}$.

$(\sqrt{3})^2 \stackrel{?}{=} 3$ Write in exponential form.

$3 = 3$ ✓ True

The solution set is $\{\sqrt{3}\}$.

(d) $\log_{49} \sqrt[3]{7} = x$

$49^x = \sqrt[3]{7}$ Write in exponential form.

$(7^2)^x = 7^{1/3}$ Write with the same base.

$7^{2x} = 7^{1/3}$ Power rule for exponents

$2x = \frac{1}{3}$ Set the exponents equal.

$x = \frac{1}{6}$ Divide by 2 (which is the same as multiplying by $\frac{1}{2}$).

Check to verify that the solution set is $\{\frac{1}{6}\}$.**NOW TRY** **OBJECTIVE 4** Use the definition of logarithm to simplify logarithmic expressions.

The definition of logarithm enables us to state several special properties.

Special Properties of LogarithmsFor any positive real number b , where $b \neq 1$, the following hold true.

$\log_b b = 1$

$\log_b 1 = 0$

$\log_b b^r = r$ (r is real.)

$b^{\log_b r} = r$ ($r > 0$)

To prove the last statement, let $x = \log_b r$.

$x = \log_b r$

$b^x = r$ Write in exponential form.

$b^{\log_b r} = r$ Replace x with $\log_b r$.

This is the statement to be proved.

NOW TRY ANSWERS

3. (a) $\{\frac{1}{32}\}$ (b) $\{\frac{35}{16}\}$

(c) $\{\sqrt{10}\}$ (d) $\{\frac{1}{9}\}$

NOW TRY EXERCISE 4

Use the special properties to evaluate each expression.

- (a) $\log_{10} 10$ (b) $\log_8 1$
 (c) $\log_{0.1} 1$ (d) $\log_3 3^9$
 (e) $5^{\log_5 3}$ (f) $\log_3 81$

EXAMPLE 4 Using Special Properties of Logarithms

Use the special properties to evaluate.

- (a) $\log_7 7 = 1$ $\log_b b = 1$ (b) $\log_{\sqrt{2}} \sqrt{2} = 1$ $\log_b b = 1$
 (c) $\log_9 1 = 0$ $\log_b 1 = 0$ (d) $\log_{0.2} 1 = 0$ $\log_b 1 = 0$
 (e) $\log_2 2^6 = 6$ $\log_b b^r = r$ (f) $\log_3 3^{-2.5} = -2.5$ $\log_b b^r = r$
 (g) $4^{\log_4 9} = 9$ $b^{\log_b r} = r$ (h) $10^{\log_{10} 13} = 13$ $b^{\log_b r} = r$
 (i) $\log_2 32$ (j) $\log_3 \frac{1}{3}$
 $= \log_2 2^5$ $32 = 2^5$ $= \log_3 3^{-1}$ Definition of
 $= 5$ $\log_b b^r = r$ $= -1$ negative exponent
 $\log_b b^r = r$

NOW TRY

OBJECTIVE 5 Define and graph logarithmic functions.

Logarithmic Function

If a and x are positive real numbers, where $a \neq 1$, then

$$g(x) = \log_a x$$

defines the **logarithmic function with base a** .

NOW TRY EXERCISE 5

Graph $f(x) = \log_6 x$.

EXAMPLE 5 Graphing a Logarithmic Function ($a > 1$)

Graph $f(x) = \log_2 x$.

By writing $y = f(x) = \log_2 x$ in exponential form as $x = 2^y$, we can identify ordered pairs that satisfy the equation. It is easier to choose values for y and find the corresponding values of x . Plotting the points in the table of ordered pairs and connecting them with a smooth curve gives the graph in FIGURE 18. This graph is typical of logarithmic functions with base $a > 1$.




FIGURE 18

Logarithmic function with base $a > 1$

Domain: $(0, \infty)$

Range: $(-\infty, \infty)$

x -intercept: $(1, 0)$

The function is one-to-one, and its graph rises from left to right.

NOW TRY

EXAMPLE 6 Graphing a Logarithmic Function ($0 < a < 1$)

Graph $g(x) = \log_{1/2} x$.


We write $y = g(x) = \log_{1/2} x$ in exponential form as

$$x = \left(\frac{1}{2}\right)^y,$$

and then choose values for y and find the corresponding values of x . Plotting these points and connecting them with a smooth curve gives the graph in FIGURE 19. This graph is typical of logarithmic functions with base $0 < a < 1$.

NOW TRY ANSWERS

4. (a) 1 (b) 0 (c) 0
 (d) 9 (e) 3 (f) 4



NOW TRY EXERCISE 6
Graph $g(x) = \log_{1/4} x$.




FIGURE 19

Logarithmic function with base $0 < a < 1$

Domain: $(0, \infty)$

Range: $(-\infty, \infty)$

x-intercept: $(1, 0)$

The function is one-to-one, and its graph falls from left to right.

NOW TRY

Characteristics of the Graph of $g(x) = \log_a x$

1. The graph contains the point $(1, 0)$, which is its x -intercept.
2. The function is one-to-one.
 - When $a > 1$, the graph *rises* from left to right, from the fourth quadrant to the first. (See FIGURE 18.)
 - When $0 < a < 1$, the graph *falls* from left to right, from the first quadrant to the fourth. (See FIGURE 19.)
3. The graph approaches the y -axis, but never touches it—that is, the y -axis is an asymptote.
4. The domain is $(0, \infty)$, and the range is $(-\infty, \infty)$.

OBJECTIVE 6 Use logarithmic functions in applications involving growth or decay.

NOW TRY EXERCISE 7

Suppose the gross national product (GNP) of a small country (in millions of dollars) is approximated by the logarithmic function

$$G(t) = 15.0 + 2.00 \log_{10} t,$$

where t is time in years since 2013. Approximate to the nearest tenth the GNP for each value of t .

- (a) $t = 1$ (b) $t = 10$

EXAMPLE 7 Applying a Logarithmic Function

Barometric pressure in inches of mercury at a distance of x miles from the eye of a typical hurricane can be approximated by the logarithmic function

$$f(x) = 27 + 1.105 \log_{10} (x + 1).$$

(Data from Miller, A. and R. Anthes, *Meteorology*, Fifth Edition, Charles E. Merrill Publishing Company.)

Approximate the pressure 9 mi from the eye of the hurricane.



$$f(x) = 27 + 1.105 \log_{10} (x + 1)$$

$$f(9) = 27 + 1.105 \log_{10} (9 + 1) \quad \text{Let } x = 9.$$

$$f(9) = 27 + 1.105 \log_{10} 10 \quad \text{Add inside parentheses.}$$


$$f(9) = 27 + 1.105(1) \quad \log_{10} 10 = 1$$

$$f(9) = 28.105 \quad \text{Add.}$$

The pressure 9 mi from the eye of the hurricane is 28.105 in.

NOW TRY

NOW TRY ANSWERS



6.
 7. (a) \$15.0 million
 (b) \$17.0 million

9.3 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

1. Concept Check Match each logarithmic equation in Column I with the corresponding exponential equation in Column II.

I	II
(a) $\log_{1/3} 3 = -1$	A. $8^{1/3} = \sqrt[3]{8}$
(b) $\log_5 1 = 0$	B. $\left(\frac{1}{3}\right)^{-1} = 3$
(c) $\log_2 \sqrt{2} = \frac{1}{2}$	C. $4^1 = 4$
(d) $\log_{10} 1000 = 3$	D. $2^{1/2} = \sqrt{2}$
(e) $\log_8 \sqrt[3]{8} = \frac{1}{3}$	E. $5^0 = 1$
(f) $\log_4 4 = 1$	F. $10^3 = 1000$

2. Concept Check Match each logarithm in Column I with its corresponding value in Column II.

I	II
(a) $\log_4 16$	A. -2
(b) $\log_3 81$	B. -1
(c) $\log_3 \left(\frac{1}{3}\right)$	C. 2
(d) $\log_{10} 0.01$	D. 0
(e) $\log_5 \sqrt{5}$	E. $\frac{1}{2}$
(f) $\log_{13} 1$	F. 4

3. Concept Check The domain of $f(x) = a^x$ is $(-\infty, \infty)$, while the range is $(0, \infty)$. Therefore because $g(x) = \log_a x$ is the inverse of f , the domain of g is _____, while the range of g is _____.

4. Concept Check The graphs of both $f(x) = 3^x$ and $g(x) = \log_3 x$ rise from left to right. Which one rises at a faster rate as x gets large?

Write in logarithmic form. See Example 1.

- | | | | |
|------------------------------|-------------------------------|----------------------------------------|------------------------------------------|
| 5. $4^5 = 1024$ | 6. $3^6 = 729$ | 7. $\left(\frac{1}{2}\right)^{-3} = 8$ | 8. $\left(\frac{1}{6}\right)^{-3} = 216$ |
| 9. $10^{-3} = 0.001$ | 10. $36^{1/2} = 6$ | 11. $\sqrt[4]{625} = 5$ | 12. $\sqrt[3]{343} = 7$ |
| 13. $8^{-2/3} = \frac{1}{4}$ | 14. $16^{-3/4} = \frac{1}{8}$ | 15. $5^0 = 1$ | 16. $7^0 = 1$ |

Write in exponential form. See Example 1.

- | | | |
|--------------------------------------------|---------------------------------|--------------------------------------------|
| 17. $\log_4 64 = 3$ | 18. $\log_2 512 = 9$ | 19. $\log_{12} 12 = 1$ |
| 20. $\log_{100} 100 = 1$ | 21. $\log_6 1 = 0$ | 22. $\log_\pi 1 = 0$ |
| 23. $\log_9 3 = \frac{1}{2}$ | 24. $\log_{64} 2 = \frac{1}{6}$ | 25. $\log_{1/4} \frac{1}{2} = \frac{1}{2}$ |
| 26. $\log_{1/8} \frac{1}{2} = \frac{1}{3}$ | 27. $\log_5 5^{-1} = -1$ | 28. $\log_{10} 10^{-2} = -2$ |

29. Concept Check Match each logarithm in Column I with its value in Column II.

I	II
(a) $\log_8 8$	A. -1
(b) $\log_{16} 1$	B. 0
(c) $\log_{0.3} 1$	C. 1
(d) $\log_{\sqrt{7}} \sqrt{7}$	D. 0.1

30. Concept Check When a student asked his teacher to explain how to evaluate

$$\log_9 3$$

without showing any work, his teacher told him to “Think radically.” Explain what the teacher meant by this hint.

Use a calculator to approximate each logarithm to four decimal places. *See Example 2.*

- | | | | |
|---------------------|---------------------|--------------------------|--------------------------|
| 31. $\log_2 9$ | 32. $\log_2 15$ | 33. $\log_5 18$ | 34. $\log_5 26$ |
| 35. $\log_{1/4} 12$ | 36. $\log_{1/5} 27$ | 37. $\log_2 \frac{1}{3}$ | 38. $\log_2 \frac{1}{7}$ |
| 39. $\log_{10} 84$ | 40. $\log_{10} 126$ | 41. $\log 50$ | 42. $\log 90$ |

Solve each equation. *See Example 3.*

- | | | |
|--------------------------------|--------------------------------|----------------------------------------|
| 43. $x = \log_{27} 3$ | 44. $x = \log_{125} 5$ | 45. $\log_5 x = -3$ |
| 46. $\log_{10} x = -2$ | 47. $\log_x 9 = \frac{1}{2}$ | 48. $\log_x 5 = \frac{1}{2}$ |
| 49. $\log_x 125 = -3$ | 50. $\log_x 64 = -6$ | 51. $\log_{12} x = 0$ |
| 52. $\log_4 x = 0$ | 53. $\log_x x = 1$ | 54. $\log_x 1 = 0$ |
| 55. $\log_x \frac{1}{25} = -2$ | 56. $\log_x \frac{1}{10} = -1$ | 57. $\log_8 32 = x$ |
| 58. $\log_{81} 27 = x$ | 59. $\log_\pi \pi^4 = x$ | 60. $\log_{\sqrt{2}} (\sqrt{2})^9 = x$ |
| 61. $\log_6 \sqrt{216} = x$ | 62. $\log_4 \sqrt{64} = x$ | 63. $\log_4 (2x + 4) = 3$ |
| 64. $\log_3 (2x + 7) = 4$ | 65. $\log_{1/3} (x - 4) = 2$ | 66. $\log_{1/2} (2x - 1) = 3$ |

Use the special properties of logarithms to evaluate each expression. *See Example 4.*

- | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|
| 67. $\log_3 3$ | 68. $\log_8 8$ | 69. $\log_5 1$ | 70. $\log_{12} 1$ |
| 71. $\log_4 4^9$ | 72. $\log_5 5^6$ | 73. $\log_2 2^{-1}$ | 74. $\log_4 4^{-6}$ |
| 75. $6^{\log_6 9}$ | 76. $12^{\log_{12} 3}$ | 77. $8^{\log_8 5}$ | 78. $5^{\log_5 11}$ |
| 79. $\log_2 64$ | 80. $\log_2 128$ | 81. $\log_3 81$ | 82. $\log_3 27$ |
| 83. $\log_4 \frac{1}{4}$ | 84. $\log_6 \frac{1}{6}$ | 85. $\log_6 \sqrt[3]{6}$ | 86. $\log_9 \sqrt[3]{9}$ |

Graph each logarithmic function. *See Examples 5 and 6.*


- | | | | |
|---------------------------|---------------------------|---------------------------|---------------------------|
| 87. $g(x) = \log_3 x$ | 88. $g(x) = \log_5 x$ | 89. $f(x) = \log_4 x$ | 90. $f(x) = \log_6 x$ |
| 91. $f(x) = \log_{1/3} x$ | 92. $f(x) = \log_{1/5} x$ | 93. $g(x) = \log_{1/4} x$ | 94. $g(x) = \log_{1/6} x$ |

Use the graph at the right to predict the value of $f(t)$ for the given value of t .

- | | | |
|-------------|--------------|--------------|
| 95. $t = 0$ | 96. $t = 10$ | 97. $t = 60$ |
|-------------|--------------|--------------|

98. Show that the points determined in Exercises 95–97 lie on the graph of

$$f(t) = 8 \log_5 (2t + 5).$$



Concept Check Answer each question.

99. Why is 1 not allowed as a base for a logarithmic function?
100. Why is a negative number not allowed as a base for a logarithmic function?
101. Why is $\log_a 1 = 0$ true for any value of a that is allowed as the base of a logarithm? Use a rule of exponents introduced earlier in the explanation.
102. Why is $\log_a a = 1$ true for any value of a that is allowed as the base of a logarithm?

Solve each problem. See Example 7.

103. Sales (in thousands of units) of a new product are approximated by the logarithmic function

$$S(t) = 100 + 30 \log_3 (2t + 1),$$

where t is the number of years after the product is introduced.

- (a) What were the sales, to the nearest unit, after 1 yr?
 (b) What were the sales, to the nearest unit, after 13 yr?
 (c) Graph $y = S(t)$.
104. A study showed that the number of mice in an old abandoned house was approximated by the logarithmic function

$$M(t) = 6 \log_4 (2t + 4),$$

where t is measured in months and $t = 0$ corresponds to January 2018. Find the number of mice in the house for each month.

- (a) January 2018 (b) July 2018 (c) July 2020 (d) Graph $y = M(t)$.
105. An online sales company finds that its sales (in millions of dollars) are approximated by the logarithmic function

$$S(x) = \log_2 (3x + 1),$$

where x is the number of advertisements placed on a popular website. How many advertisements must be placed to earn sales of \$4 million?

106. The population of deer (in thousands) in a certain area is approximated by the logarithmic function

$$f(x) = \log_5 (100x - 75),$$


where x is the number of years since 2017. During what year is the population expected to be 4 thousand deer?




RELATING CONCEPTS For Individual or Group Work (Exercises 107–110)

To see how exponential and logarithmic functions are related, work Exercises 107–110 in order.

107. Complete the table of values, and sketch the graph of $y = 10^x$. Give the domain and range of the function.



108. Complete the table of values, and sketch the graph of $y = \log_{10} x$. Give the domain and range of the function.



109. Describe the symmetry between the graphs in Exercises 107 and 108.

110. What can we conclude about the functions

$$y = f(x) = 10^x \quad \text{and} \quad y = g(x) = \log_{10} x?$$

9.4 Properties of Logarithms

OBJECTIVES

- 1 Use the product rule for logarithms.
- 2 Use the quotient rule for logarithms.
- 3 Use the power rule for logarithms.
- 4 Use properties to write alternative forms of logarithmic expressions.

Logarithms were used as an aid to numerical calculation for several hundred years. Today the widespread use of calculators has made the use of logarithms for calculation obsolete. However, logarithms are still very important in applications and in further work in mathematics.

OBJECTIVE 1 Use the product rule for logarithms.

One way in which logarithms simplify problems is by changing a problem of multiplication into one of addition. For example, we know that

$$\log_2 4 = 2, \quad \log_2 8 = 3, \quad \text{and} \quad \log_2 32 = 5.$$

Therefore, we can make the following statements.

$$\log_2 32 = \log_2 4 + \log_2 8 \quad 5 = 2 + 3$$

$$\log_2 (4 \cdot 8) = \log_2 4 + \log_2 8 \quad 32 = 4 \cdot 8$$

This is an example of the product rule for logarithms.

Product Rule for Logarithms

If x , y , and b are positive real numbers, where $b \neq 1$, then the following holds true.

$$\log_b xy = \log_b x + \log_b y$$

That is, the logarithm of a product is the sum of the logarithms of the factors.

Examples: $\log_3 (4 \cdot 7) = \log_3 4 + \log_3 7$, $\log_{10} 8 + \log_{10} 9 = \log_{10} (8 \cdot 9)$

To prove this rule, let $m = \log_b x$ and $n = \log_b y$, and recall that

$\log_b x = m$ is equivalent to $b^m = x$ and $\log_b y = n$ is equivalent to $b^n = y$.

Now consider the product xy .

$$xy = b^m \cdot b^n \quad \text{Substitute.}$$

$$xy = b^{m+n} \quad \text{Product rule for exponents}$$

$$\log_b xy = m + n \quad \text{Write in logarithmic form.}$$

$$\log_b xy = \log_b x + \log_b y \quad \text{Substitute for } m \text{ and } n.$$

The last statement is the result we wished to prove.

NOTE The word statement of the product rule can be restated by replacing the word “logarithm” with the word “exponent.” The rule then becomes the familiar rule for multiplying exponential expressions:

The exponent of a product is equal to the sum of the exponents of the factors.


NOW TRY EXERCISE 1

Use the product rule to rewrite each logarithm.

- (a) $\log_{10}(7 \cdot 9)$
 (b) $\log_5 11 + \log_5 8$
 (c) $\log_5(5x)$, $x > 0$
 (d) $\log_2 t^3$, $t > 0$

EXAMPLE 1 Using the Product Rule

Use the product rule to rewrite each logarithm. Assume $x > 0$.

- (a) $\log_5(6 \cdot 9)$
 $= \log_5 6 + \log_5 9$ **Product rule**
- (b) $\log_7 8 + \log_7 12$
 $= \log_7(8 \cdot 12)$ **Product rule**
 $= \log_7 96$ **Multiply.**
- (c) $\log_3(3x)$
 $= \log_3 3 + \log_3 x$ **Product rule**
 $= 1 + \log_3 x$ **$\log_b b = 1$**
- (d) $\log_4 x^3$
 $= \log_4(x \cdot x \cdot x)$ **$x^3 = x \cdot x \cdot x$**
 $= \log_4 x + \log_4 x + \log_4 x$
Product rule
 $= 3 \log_4 x$ **Combine like terms.**

NOW TRY 

OBJECTIVE 2 Use the quotient rule for logarithms.

The rule for division is similar to the rule for multiplication.

Quotient Rule for Logarithms

If x , y , and b are positive real numbers, where $b \neq 1$, then the following holds true.

$$\log_b \frac{x}{y} = \log_b x - \log_b y$$

That is, the logarithm of a quotient is the difference of the logarithm of the numerator and the logarithm of the denominator.

Examples: $\log_5 \frac{2}{3} = \log_5 2 - \log_5 3$, $\log_7 3 - \log_7 5 = \log_7 \frac{3}{5}$

The proof of this rule is similar to the proof of the product rule.

EXAMPLE 2 Using the Quotient Rule

Use the quotient rule to rewrite each logarithm. Assume $x > 0$.

- (a) $\log_4 \frac{7}{9}$
 $= \log_4 7 - \log_4 9$ **Quotient rule**
- (b) $\log_5 6 - \log_5 x$
 $= \log_5 \frac{6}{x}$ **Quotient rule**
- (c) $\log_3 \frac{27}{5}$
 $= \log_3 27 - \log_3 5$ **Quotient rule**
 $= 3 - \log_3 5$ **$\log_3 27 = 3$**
- (d) $\log_6 28 - \log_6 7$
 $= \log_6 \frac{28}{7}$ **Quotient rule**
 $= \log_6 4$ **$\frac{28}{7} = 4$**

NOW TRY 


NOW TRY EXERCISE 2

Use the quotient rule to rewrite each logarithm.

- (a) $\log_{10} \frac{7}{9}$
 (b) $\log_4 x - \log_4 12$, $x > 0$
 (c) $\log_5 \frac{25}{27}$
 (d) $\log_2 15 - \log_2 3$

NOW TRY ANSWERS

1. (a) $\log_{10} 7 + \log_{10} 9$
 (b) $\log_5 88$
 (c) $1 + \log_5 x$
 (d) $3 \log_2 t$
2. (a) $\log_{10} 7 - \log_{10} 9$
 (b) $\log_4 \frac{x}{12}$
 (c) $2 - \log_5 27$
 (d) $\log_2 5$

! CAUTION There is no property of logarithms to rewrite the logarithm of a sum.

$$\log_b (x + y) \neq \log_b x + \log_b y$$

Also, $\log_b x \cdot \log_b y \neq \log_b xy$, and $\frac{\log_b x}{\log_b y} \neq \log_b \frac{x}{y}$.

OBJECTIVE 3 Use the power rule for logarithms.

Consider the exponential expression 2^3 .

$$2^3 \text{ means } 2 \cdot 2 \cdot 2. \quad \text{The base 2 is used as a factor 3 times.}$$

Similarly, the product rule can be extended to rewrite the logarithm of a power as the product of the exponent and the logarithm of the base.

$$\begin{array}{l|l} \log_5 2^3 & \log_2 7^4 \\ = \log_5 (2 \cdot 2 \cdot 2) & = \log_2 (7 \cdot 7 \cdot 7 \cdot 7) \\ = \log_5 2 + \log_5 2 + \log_5 2 & = \log_2 7 + \log_2 7 + \log_2 7 + \log_2 7 \\ = 3 \log_5 2 & = 4 \log_2 7 \end{array}$$

Furthermore, we saw in **Example 1(d)** that $\log_4 x^3 = 3 \log_4 x$. These examples suggest the following rule.

Power Rule for Logarithms

If x and b are positive real numbers, where $b \neq 1$, and if r is any real number, then the following holds true.

$$\log_b x^r = r \log_b x$$

That is, the logarithm of a number to a power equals the exponent times the logarithm of the number.

Examples: $\log_b m^5 = 5 \log_b m$, $\log_3 5^4 = 4 \log_3 5$

To prove the power rule, let $\log_b x = m$.

$$b^m = x \quad \text{Write in exponential form.}$$

$$(b^m)^r = x^r \quad \text{Raise each side to the power } r.$$

$$b^{mr} = x^r \quad \text{Power rule for exponents}$$

$$\log_b x^r = mr \quad \text{Write in logarithmic form.}$$

$$\log_b x^r = rm \quad \text{Commutative property}$$

$$\log_b x^r = r \log_b x \quad m = \log_b x \text{ from above}$$

This is the statement to be proved.

As a special case of the power rule, let $r = \frac{1}{p}$, so

$$\log_b \sqrt[p]{x} = \log_b x^{1/p} = \frac{1}{p} \log_b x.$$

Examples: $\log_b \sqrt[5]{x} = \log_b x^{1/5} = \frac{1}{5} \log_b x$, $\log_b \sqrt[3]{x^4} = \log_b x^{4/3} = \frac{4}{3} \log_b x$ ($x > 0$)

Another special case is

$$\log_b \frac{1}{x} = \log_b x^{-1} = -\log_b x. \quad -a = -1 \cdot a$$

Example: $\log_9 \frac{1}{5} = \log_9 5^{-1} = -\log_9 5$

NOW TRY EXERCISE 3

Use the power rule to rewrite each logarithm. Assume $a > 0$, $x > 0$, and $a \neq 1$.

(a) $\log_7 5^3$ (b) $\log_a \sqrt{10}$

(c) $\log_3 \sqrt[4]{x^3}$ (d) $\log_4 \frac{1}{x^5}$

EXAMPLE 3 Using the Power Rule

Use the power rule to rewrite each logarithm. Assume $b > 0$, $x > 0$, and $b \neq 1$.

(a) $\log_5 4^2$
 $= 2 \log_5 4$
 Power rule

(b) $\log_b x^5$
 $= 5 \log_b x$
 Power rule

(c) $\log_b \sqrt{7}$
 $= \log_b 7^{1/2}$ $\sqrt{x} = x^{1/2}$
 $= \frac{1}{2} \log_b 7$ Power rule

(d) $\log_2 \sqrt[5]{x^2}$
 $= \log_2 x^{2/5}$ $\sqrt[5]{x^2} = x^{2/5}$
 $= \frac{2}{5} \log_2 x$ Power rule

(e) $\log_3 \frac{1}{x^4}$
 $= \log_3 x^{-4}$ Definition of negative exponent
 $= -4 \log_3 x$ Power rule

NOW TRY 

We summarize the properties of logarithms from the previous section and this one.

Properties of Logarithms

If x , y , and b are positive real numbers, where $b \neq 1$, and r is any real number, then the following hold true.

Special Properties	$\log_b b = 1$	$\log_b 1 = 0$
	$\log_b b^r = r$	$b^{\log_b r} = r \quad (r > 0)$
Product Rule	$\log_b xy = \log_b x + \log_b y$	
Quotient Rule	$\log_b \frac{x}{y} = \log_b x - \log_b y$	
Power Rule	$\log_b x^r = r \log_b x$	

OBJECTIVE 4 Use properties to write alternative forms of logarithmic expressions.

EXAMPLE 4 Writing Logarithms in Alternative Forms

Use the properties of logarithms to rewrite each expression if possible. Assume that all variables represent positive real numbers.

(a) $\log_4 4x^3$
 $= \log_4 4 + \log_4 x^3$ Product rule
 $= 1 + 3 \log_4 x$ $\log_b b = 1$; Power rule

NOW TRY ANSWERS

3. (a) $3 \log_7 5$ (b) $\frac{1}{2} \log_a 10$
 (c) $\frac{3}{4} \log_3 x$ (d) $-5 \log_4 x$


NOW TRY EXERCISE 4

Use the properties of logarithms to rewrite each expression if possible. Assume that all variables represent positive real numbers.

(a) $\log_3 9z^4$

(b) $\log_6 \sqrt{\frac{n}{3m}}$

(c) $\log_2 x + 3 \log_2 y - \log_2 z$

(d) $\log_5 (x + 10)$
 $+ \log_5 (x - 10)$
 $- \frac{3}{5} \log_5 x, \quad x > 10$

(e) $\log_7 (49 + 2x)$

(b) $\log_7 \sqrt{\frac{m}{n}}$
 $= \log_7 \left(\frac{m}{n}\right)^{1/2}$ Write the radical expression with a rational exponent.

$$= \frac{1}{2} \log_7 \frac{m}{n}$$
 Power rule

$$= \frac{1}{2} (\log_7 m - \log_7 n)$$
 Quotient rule

(c) $\log_5 \frac{a^2}{bc}$
 $= \log_5 a^2 - \log_5 bc$ Quotient rule

$$= 2 \log_5 a - \log_5 bc$$
 Power rule

$$= 2 \log_5 a - (\log_5 b + \log_5 c)$$
 Product rule

$$= 2 \log_5 a - \log_5 b - \log_5 c$$

Parenttheses are necessary here.

(d) $4 \log_b m - \log_b n, \quad b \neq 1$

$$= \log_b m^4 - \log_b n$$
 Power rule

$$= \log_b \frac{m^4}{n}$$
 Quotient rule

(e) $\log_b (x + 1) + \log_b (2x + 1) - \frac{2}{3} \log_b x, \quad b \neq 1$

$$= \log_b (x + 1) + \log_b (2x + 1) - \log_b x^{2/3}$$
 Power rule

$$= \log_b \frac{(x + 1)(2x + 1)}{x^{2/3}}$$
 Product and quotient rules

$$= \log_b \frac{2x^2 + 3x + 1}{x^{2/3}}$$
 Multiply in the numerator.

(f) $\log_8 (2p + 3r)$ cannot be rewritten using the properties of logarithms. **There is no property of logarithms to rewrite the logarithm of a sum.**

NOW TRY 

In the next example, we use numerical values for $\log_2 5$ and $\log_2 3$. While we use the equality symbol to give these values, they are actually approximations because most logarithms of this type are irrational numbers. **We use = with the understanding that the values are correct to four decimal places.**

EXAMPLE 5 Using the Properties of Logarithms with Numerical Values

Given that $\log_2 5 = 2.3219$ and $\log_2 3 = 1.5850$, use these values and the properties of logarithms to evaluate each expression.

(a) $\log_2 15$

$$= \log_2 (3 \cdot 5)$$
 Factor 15.

$$= \log_2 3 + \log_2 5$$
 Product rule

$$= 1.5850 + 2.3219$$
 Substitute the given values.

$$= 3.9069$$
 Add.

NOW TRY ANSWERS

4. (a) $2 + 4 \log_3 z$

(b) $\frac{1}{2} (\log_6 n - \log_6 3 - \log_6 m)$

(c) $\log_2 \frac{xy^3}{z}$ (d) $\log_5 \frac{x^2 - 100}{x^{3/5}}$

(e) cannot be rewritten


NOW TRY EXERCISE 5

Given that $\log_2 7 = 2.8074$ and $\log_2 10 = 3.3219$, use these values and the properties of logarithms to evaluate each expression.

- (a) $\log_2 70$ (b) $\log_2 0.7$
 (c) $\log_2 49$

$$\begin{aligned} \text{(b)} \log_2 0.6 &= \log_2 \frac{3}{5} && 0.6 = \frac{6}{10} = \frac{3}{5} \\ &= \log_2 3 - \log_2 5 && \text{Quotient rule} \\ &= 1.5850 - 2.3219 && \text{Substitute the given values.} \\ &= -0.7369 && \text{Subtract.} \end{aligned}$$

$$\begin{aligned} \text{(c)} \log_2 27 &= \log_2 3^3 && \text{Write 27 as a power of 3.} \\ &= 3 \log_2 3 && \text{Power rule} \\ &= 3(1.5850) && \text{Substitute the given value.} \\ &= 4.7550 && \text{Multiply.} \end{aligned}$$

NOW TRY 


NOW TRY EXERCISE 6

Determine whether each statement is *true* or *false*.

- (a) $\log_2 16 + \log_2 16 = \log_2 32$
 (b) $(\log_2 4)(\log_3 9) = \log_6 6^4$

EXAMPLE 6 Determining Whether Statements about Logarithms Are True

Determine whether each statement is *true* or *false*.

(a) $\log_2 8 - \log_2 4 = \log_2 4$

Evaluate each side.

$\begin{aligned} \log_2 8 - \log_2 4 & \text{Left side} \\ &= \log_2 2^3 - \log_2 2^2 && \text{Write 8 and 4 as powers of 2.} \\ &= 3 - 2 && \log_b b^r = r \\ &= 1 && \text{Subtract.} \end{aligned}$		$\begin{aligned} \log_2 4 & \text{Right side} \\ &= \log_2 2^2 && \text{Write 4 as a power of 2.} \\ &= 2 && \log_b b^r = r \end{aligned}$
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--------------------------------------------------------------------------------------------------------------------------------------------

The statement is false because $1 \neq 2$.

(b) $\log_3 (\log_2 8) = \frac{\log_7 49}{\log_8 64}$

$\begin{aligned} \log_3 (\log_2 8) & \text{Left side} \\ &= \log_3 (\log_2 2^3) && \text{Write 8 as a power of 2.} \\ &= \log_3 3 && \log_b b^r = r \\ &= 1 && \log_b b = 1 \end{aligned}$		$\begin{aligned} \frac{\log_7 49}{\log_8 64} & \text{Right side} \\ &= \frac{\log_7 7^2}{\log_8 8^2} && \text{Write 49 and 64 using exponents.} \\ &= \frac{2}{2} && \log_b b^r = r \\ &= 1 && \text{Simplify.} \end{aligned}$
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

The statement is true because $1 = 1$.

NOW TRY 

NOW TRY ANSWERS


5. (a) 6.1293 (b) -0.5145
 (c) 5.6148
 6. (a) false (b) true

9.4 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Determine whether each statement of a logarithmic property is true or false. If it is false, correct it by changing the right side of the equation.

1. $\log_b x + \log_b y = \log_b (x + y)$
2. $\log_b \frac{x}{y} = \log_b x - \log_b y$
3. $\log_b xy = \log_b x + \log_b y$
4. $b^{\log_b r} = 1$
5. $\log_b b^r = r$
6. $\log_b x^r = \log_b rx$

7. Concept Check A student erroneously wrote $\log_a(x + y) = \log_a x + \log_a y$. When his teacher explained that this was indeed wrong, the student claimed that he had used the distributive property. **WHAT WENT WRONG?**

8. Concept Check Consider the following “proof” that $\log_2 16$ does not exist.

$$\begin{aligned}\log_2 16 \\ &= \log_2 (-4)(-4) \\ &= \log_2 (-4) + \log_2 (-4)\end{aligned}$$

The logarithm of a negative number is not defined, so the final step cannot be evaluated. Thus $\log_2 16$ does not exist. **WHAT WENT WRONG?**

Concept Check Use the indicated rule of logarithms to complete each equation.

9. $\log_{10}(7 \cdot 8) = \underline{\hspace{2cm}}$ (product rule)

10. $\log_{10} \frac{7}{8} = \underline{\hspace{2cm}}$ (quotient rule)

11. $3^{\log_3 4} = \underline{\hspace{2cm}}$ (special property)

12. $\log_{10} 3^6 = \underline{\hspace{2cm}}$ (power rule)

13. $\log_3 3^9 = \underline{\hspace{2cm}}$ (special property)

14. $\log_3 9^2 = \underline{\hspace{2cm}}$ (special property)

Use the properties of logarithms to express each logarithm as a sum or difference of logarithms, or as a single logarithm if possible. Assume that all variables represent positive real numbers. See Examples 1–4.

15. $\log_7(4 \cdot 5)$ 16. $\log_8(9 \cdot 11)$ 17. $\log_5 \frac{8}{3}$

18. $\log_3 \frac{7}{5}$ 19. $\log_4 6^2$ 20. $\log_5 7^4$

21. $\log_3 \frac{\sqrt[3]{4}}{x^2 y}$ 22. $\log_7 \frac{\sqrt[3]{13}}{pq^2}$ 23. $\log_3 \sqrt{\frac{xy}{5}}$

24. $\log_6 \sqrt{\frac{pq}{7}}$ 25. $\log_2 \frac{\sqrt[3]{x} \cdot \sqrt[5]{y}}{r^2}$ 26. $\log_4 \frac{\sqrt[4]{z} \cdot \sqrt[5]{w}}{s^2}$

Use the properties of logarithms to write each expression as a single logarithm. Assume that all variables are defined in such a way that the variable expressions are positive, and bases are positive numbers not equal to 1. See Examples 1–4.

27. $\log_b x + \log_b y$ 28. $\log_b w + \log_b z$

29. $\log_a m - \log_a n$ 30. $\log_b x - \log_b y$

31. $(\log_a r - \log_a s) + 3 \log_a t$ 32. $(\log_a p - \log_a q) + 2 \log_a r$

33. $3 \log_a 5 - 4 \log_a 3$ 34. $3 \log_a 5 - \frac{1}{2} \log_a 9$

35. $\log_{10}(x + 3) + \log_{10}(x - 3)$ 36. $\log_{10}(y + 4) + \log_{10}(y - 4)$

37. $\log_{10}(x + 3) + \log_{10}(x + 5)$ 38. $\log_{10}(x + 4) + \log_{10}(x + 6)$

39. $3 \log_p x + \frac{1}{2} \log_p y - \frac{3}{2} \log_p z - 3 \log_p a$ 40. $\frac{1}{3} \log_b x + \frac{2}{3} \log_b y - \frac{3}{4} \log_b s - \frac{2}{3} \log_b t$

To four decimal places, the values of $\log_{10} 2$ and $\log_{10} 9$ are

$$\log_{10} 2 = 0.3010 \quad \text{and} \quad \log_{10} 9 = 0.9542.$$

Use these values and the properties of logarithms to evaluate each expression. **DO NOT USE A CALCULATOR. See Example 5.**

- | | | |
|-----------------------------|-----------------------------|-----------------------------|
| 41. $\log_{10} 18$ | 42. $\log_{10} 4$ | 43. $\log_{10} \frac{2}{9}$ |
| 44. $\log_{10} \frac{9}{2}$ | 45. $\log_{10} 36$ | 46. $\log_{10} 162$ |
| 47. $\log_{10} \sqrt[4]{9}$ | 48. $\log_{10} \sqrt[5]{2}$ | 49. $\log_{10} 3$ |
| 50. $\log_{10} \frac{1}{9}$ | 51. $\log_{10} 9^5$ | 52. $\log_{10} 2^{19}$ |

Determine whether each statement is true or false. **See Example 6.**

- | | |
|------------------------------------------------------|---------------------------------------------------------|
| 53. $\log_2 (8 + 32) = \log_2 8 + \log_2 32$ | 54. $\log_2 (64 - 16) = \log_2 64 - \log_2 16$ |
| 55. $\log_3 7 + \log_3 7^{-1} = 0$ | 56. $\log_3 49 + \log_3 49^{-1} = 0$ |
| 57. $\log_6 60 - \log_6 10 = 1$ | 58. $\log_3 8 + \log_3 \frac{1}{8} = 0$ |
| 59. $\frac{\log_{10} 7}{\log_{10} 14} = \frac{1}{2}$ | 60. $\frac{\log_{10} 10}{\log_{10} 100} = \frac{1}{10}$ |

9.5 Common and Natural Logarithms

OBJECTIVES

- 1 Evaluate common logarithms using a calculator.
- 2 Use common logarithms in applications.
- 3 Evaluate natural logarithms using a calculator.
- 4 Use natural logarithms in applications.
- 5 Use the change-of-base rule.

Logarithms are important in many applications in biology, engineering, economics, and social science. In this section we find numerical approximations for logarithms. Traditionally, base 10 logarithms were used most often because our number system is base 10. Logarithms to base 10 are **common logarithms**, and

$\log_{10} x$ is abbreviated as **log x**,

where the base is understood to be 10.

OBJECTIVE 1 Evaluate common logarithms using a calculator.

We evaluate common logarithms using a calculator with a **LOG** key.

EXAMPLE 1 Evaluating Common Logarithms

Evaluate each logarithm to four decimal places using a calculator as needed.

- | | |
|-----------------------------------|----------------------------------------------------------------------|
| (a) $\log 327.1 \approx 2.5147$ | (b) $\log 437,000 \approx 5.6405$ |
| (c) $\log 0.0615 \approx -1.2111$ | (d) $\log 10^{6.1988} = 6.1988$
Special property $\log_b b^r = r$ |

VOCABULARY

- common logarithm
- natural logarithm

NOW TRY EXERCISE 1

Evaluate each logarithm to four decimal places using a calculator as needed.

- (a) $\log 115$
 (b) $\log 539,000$
 (c) $\log 0.023$
 (d) $\log 10^{12.2139}$

In part (c), $\log 0.0615 \approx -1.2111$, which is a negative result. **The common logarithm of a number between 0 and 1 is always negative** because the logarithm is the exponent on 10 that produces the number. In this case, we have

$$10^{-1.2111} \approx 0.0615.$$

If the exponent (the logarithm) were positive, the result would be greater than 1 because $10^0 = 1$. The graph in FIGURE 20 illustrates these concepts.




FIGURE 20

NOW TRY

OBJECTIVE 2 Use common logarithms in applications.

In chemistry, pH is a measure of the acidity or alkalinity of a solution. Water, for example, has pH 7. In general, acids have pH numbers less than 7, and alkaline solutions have pH values greater than 7. FIGURE 21 illustrates the pH scale.




FIGURE 21 pH Scale

The **pH** of a solution is defined as

$$\text{pH} = -\log [\text{H}_3\text{O}^+],$$

where $[\text{H}_3\text{O}^+]$ is the hydronium ion concentration in moles per liter. It is customary to round pH values to the nearest tenth.

EXAMPLE 2 Using pH in an Application

Wetlands are classified as *bogs*, *fens*, *marshes*, and *swamps*, on the basis of pH values. A pH value between 6.0 and 7.5, such as that of Summerby Swamp in Michigan's Hiawatha National Forest, indicates that the wetland is a "rich fen." When the pH is between 3.0 and 6.0, the wetland is a "poor fen," and if the pH falls to 3.0 or less, it is a "bog." (Data from Mohlenbrock, R., "Summerby Swamp, Michigan," *Natural History*.)



Suppose that the hydronium ion concentration of a sample of water from a wetland is 6.3×10^{-3} mole per liter. How would this wetland be classified?

$\text{pH} = -\log [\text{H}_3\text{O}^+]$	Definition of pH
$\text{pH} = -\log (6.3 \times 10^{-3})$	Let $[\text{H}_3\text{O}^+] = 6.3 \times 10^{-3}$.
$\text{pH} = -(\log 6.3 + \log 10^{-3})$	Product rule
$\text{pH} \approx -[0.7993 - 3]$	Use a calculator to approximate $\log 6.3$; $\log_b b^r = r$
$\text{pH} = -0.7993 + 3$	Distributive property
$\text{pH} \approx 2.2$	Add.

The pH is less than 3.0, so the wetland is a bog.

NOW TRY

NOW TRY EXERCISE 2

Water taken from a wetland has a hydronium ion concentration of

$$3.4 \times 10^{-5} \text{ mole per liter.}$$

Find the pH value for the water and classify the wetland as a rich fen, a poor fen, or a bog.

NOW TRY ANSWERS

1. (a) 2.0607 (b) 5.7316
 (c) -1.6383 (d) 12.2139
 2. 4.5; poor fen


NOW TRY EXERCISE 3

Find the hydronium ion concentration of a solution with pH 2.6.

EXAMPLE 3 Finding Hydronium Ion Concentration

Find the hydronium ion concentration of drinking water with pH 6.5.

$$\text{pH} = -\log [\text{H}_3\text{O}^+] \quad \text{Definition of pH}$$

$$6.5 = -\log [\text{H}_3\text{O}^+] \quad \text{Let pH} = 6.5.$$

$$\log [\text{H}_3\text{O}^+] = -6.5 \quad \text{Multiply by } -1. \text{ Interchange sides.}$$

$$[\text{H}_3\text{O}^+] = 10^{-6.5} \quad \text{Write in exponential form, base 10.}$$

$$[\text{H}_3\text{O}^+] \approx 3.2 \times 10^{-7} \quad \text{Evaluate with a calculator.}$$

The hydronium ion concentration of drinking water is approximately 3.2×10^{-7} mole per liter.

NOW TRY 

The loudness of sound is measured in a unit called a **decibel**, abbreviated **dB**. To measure with this unit, we first assign an intensity of I_0 to a very faint sound, called the **threshold sound**. If a particular sound has intensity I , then the decibel level of this louder sound is

$$D = 10 \log \left(\frac{I}{I_0} \right).$$

The loudness of some common sounds are shown in the table. Any sound over 85 dB exceeds what hearing experts consider safe. Permanent hearing damage can be suffered at levels above 150 dB.

▼ Loudness of Common Sounds

Decibel Level	Example
60	Normal conversation
90	Rush hour traffic, lawn mower
100	Garbage truck, chain saw, pneumatic drill
120	Rock concert, thunderclap
140	Gunshot blast, jet engine
180	Rocket launching pad

Data from Deafness Research Foundation.


NOW TRY EXERCISE 4

Find the decibel level to the nearest whole number of the sound from a jet engine with intensity I of

$$(6.312 \times 10^{13})I_0.$$

EXAMPLE 4 Measuring the Loudness of Sound

If music delivered through Bluetooth headphones has intensity I of $(3.162 \times 10^9)I_0$, find the average decibel level.

$$D = 10 \log \left(\frac{I}{I_0} \right)$$

$$D = 10 \log \left(\frac{(3.162 \times 10^9)I_0}{I_0} \right) \quad \text{Substitute the given value for } I.$$

$$D = 10 \log (3.162 \times 10^9)$$

$$D \approx 95 \text{ dB}$$

Evaluate with a calculator.

NOW TRY 

NOW TRY ANSWERS

3. 2.5×10^{-3}

4. 138 dB



Leonhard Euler (1707–1783)

The number e is named after Euler.

OBJECTIVE 3 Evaluate natural logarithms using a calculator.

Logarithms used in applications are often **natural logarithms**, which have as base the number e . The letter e was chosen to honor Leonhard Euler, who published extensive results on the number in 1748. It is an irrational number, so its decimal expansion never terminates and never repeats.

One way to see how e appears in an exponential situation involves calculating the values of $\left(1 + \frac{1}{x}\right)^x$ as x gets larger without bound. See the table.

x	$\left(1 + \frac{1}{x}\right)^x$
1	2
10	2.59374246
100	2.704813829
1000	2.716923932
10,000	2.718145927
100,000	2.718268237

These approximations are found using a calculator.

It appears that as x gets larger without bound, $\left(1 + \frac{1}{x}\right)^x$ approaches some number. This number is e .

Approximation for e

$$e \approx 2.718281828459$$

A calculator with an (e^x) key can approximate powers of e .

$$e^2 \approx 7.389056099, \quad e^3 \approx 20.08553692, \quad \text{and} \quad e^{0.6} \approx 1.8221188$$

Logarithms with base e are called natural logarithms because they occur in natural situations that involve growth or decay.

The base e logarithm of x is written $\ln x$ (read “el en x ”).

A graph of $y = \ln x$ is given in FIGURE 22.




FIGURE 22

A calculator key labeled (LN) is used to evaluate natural logarithms.

NOW TRY EXERCISE 5

Evaluate each logarithm to four decimal places using a calculator as needed.

- (a) $\ln 0.26$ (b) $\ln 12$
 (c) $\ln 150$ (d) $\ln e^{5.8321}$

NOW TRY ANSWERS

5. (a) -1.3471 (b) 2.4849
 (c) 5.0106 (d) 5.8321

EXAMPLE 5 Evaluating Natural Logarithms

Evaluate each logarithm to four decimal places using a calculator as needed.

- (a) $\ln 0.5841 \approx -0.5377$ (b) $\ln 0.9215 \approx -0.0818$

As with common logarithms, a number between 0 and 1 has a negative natural logarithm.

- (c) $\ln 192.7 \approx 5.2611$ (d) $\ln e^{4.6832} = 4.6832$

Special property $\log_b b^r = r$


**NOW TRY
EXERCISE 6**

Use the natural logarithmic function in **Example 6** to approximate the altitude when atmospheric pressure is 600 millibars. Round to the nearest hundred.

OBJECTIVE 4 Use natural logarithms in applications.

EXAMPLE 6 Applying a Natural Logarithmic Function

Altitude in meters that corresponds to an atmospheric pressure of x millibars can be approximated by the natural logarithmic function

$$f(x) = 51,600 - 7457 \ln x.$$

(Data from Miller, A. and J. Thompson, *Elements of Meteorology*, Fourth Edition, Charles E. Merrill Publishing Company.)

Use this function to find the altitude when atmospheric pressure is 400 millibars. Round to the nearest hundred.

$$f(x) = 51,600 - 7457 \ln x$$

$$f(400) = 51,600 - 7457 \ln 400 \quad \text{Let } x = 400.$$

$$f(400) \approx 6900 \quad \text{Evaluate with a calculator.}$$

Atmospheric pressure is 400 millibars at 6900 m.



NOTE In **Example 6**, the final answer was obtained using a calculator *without* rounding the intermediate values. In general, it is best to wait until the final step to round the answer. Otherwise, a buildup of round-off error may cause the final answer to have an incorrect final decimal place digit or digits.

OBJECTIVE 5 Use the change-of-base rule.

The change-of-base rule enables us to convert logarithms from one base to another.

Change-of-Base Rule

If $a > 0$, $a \neq 1$, $b > 0$, $b \neq 1$, and $x > 0$, then the following holds true.

$$\log_a x = \frac{\log_b x}{\log_b a}$$

To derive the change-of-base rule, let $\log_a x = m$.

$$\log_a x = m$$

$$a^m = x \quad \text{Write in exponential form.}$$

$$\log_b a^m = \log_b x \quad \text{Take the logarithm on each side.}$$

$$m \log_b a = \log_b x \quad \text{Power rule}$$

$$(\log_a x)(\log_b a) = \log_b x \quad \text{Substitute for } m.$$

$$\log_a x = \frac{\log_b x}{\log_b a} \quad \text{Divide by } \log_b a.$$

NOW TRY ANSWER

6. 3900 m

This last statement is the change-of-base rule.

NOTE Any positive number other than 1 can be used for base b in the change-of-base rule. Usually the only practical bases are e and 10, because calculators give logarithms for these two bases.

NOW TRY EXERCISE 7

Use the change-of-base rule to approximate each logarithm to four decimal places.

(a) $\log_2 7$ (b) $\log_{1/3} 12$

NOW TRY ANSWERS

7. (a) 2.8074 (b) -2.2619

EXAMPLE 7 Using the Change-of-Base Rule

Use the change-of-base rule to approximate $\log_5 12$ to four decimal places.

$$\log_5 12 = \frac{\log 12}{\log 5} \approx 1.5440$$

$$\log_5 12 = \frac{\ln 12}{\ln 5} \approx 1.5440$$

Either common or natural logarithms can be used.


NOW TRY 

9.5 Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Choose the correct response.

- What is the base in the expression $\log x$?
A. x B. 1 C. 10 D. e
- What is the base in the expression $\ln x$?
A. x B. 1 C. 10 D. e
- Given that $10^0 = 1$ and $10^1 = 10$, between what two consecutive integers is the value of $\log 6.3$?
A. -1 and 0 B. 0 and 1 C. 6 and 7 D. 10 and 11
- Given that $e^1 \approx 2.718$ and $e^2 \approx 7.389$, between what two consecutive integers is the value of $\ln 6.3$?
A. 0 and 1 B. 1 and 2 C. 2 and 3 D. 6 and 7

Concept Check Without using a calculator, give the value of each expression.

- | | | |
|-----------------------|-----------------------|-------------------------|
| 5. $\log 10^{9.6421}$ | 6. $\ln e^{\sqrt{2}}$ | 7. $10^{\log \sqrt{3}}$ |
| 8. $e^{\ln 75.2}$ | 9. $\ln e^{-11.4007}$ | 10. $10 \ln e^4$ |

Use a calculator for most of the remaining exercises in this set.

Evaluate each logarithm to four decimal places. See Examples 1 and 5.

- | | | |
|-------------------------------|-------------------------------|-----------------------------|
| 11. $\log 43$ | 12. $\log 98$ | 13. $\log 328.4$ |
| 14. $\log 457.2$ | 15. $\log 0.0326$ | 16. $\log 0.1741$ |
| 17. $\log (4.76 \times 10^9)$ | 18. $\log (2.13 \times 10^4)$ | 19. $\ln 7.84$ |
| 20. $\ln 8.32$ | 21. $\ln 0.0556$ | 22. $\ln 0.0217$ |
| 23. $\ln 388.1$ | 24. $\ln 942.6$ | 25. $\ln (8.59 \times e^2)$ |
| 26. $\ln (7.46 \times e^3)$ | 27. $\ln 10$ | 28. $\log e$ |

29. Concept Check Use a calculator to find approximations of each logarithm.

- (a) $\log 356.8$ (b) $\log 35.68$ (c) $\log 3.568$
 (d) Observe the answers and make a conjecture concerning the decimal values of the common logarithms of numbers greater than 1 that have the same digits.

- 30. Concept Check** Let k represent the number of letters in your last name.
- Use a calculator to find $\log k$.
 - Raise 10 to the power indicated by the number found in part (a). What is the result?
 - Explain why we obtained the answer found in part (b). Would it matter what number we used for k to observe the same result?

Suppose that water from a wetland area is sampled and found to have the given hydronium ion concentration. Is the wetland a rich fen, a poor fen, or a bog? **See Example 2.**

31. 3.1×10^{-5} 32. 2.5×10^{-5} 33. 2.5×10^{-2}
 34. 3.6×10^{-2} 35. 2.7×10^{-7} 36. 2.5×10^{-7}

Find the pH (to the nearest tenth) of the substance with the given hydronium ion concentration. **See Example 2.**

37. Ammonia, 2.5×10^{-12} 38. Egg white, 1.6×10^{-8}
 39. Sodium bicarbonate, 4.0×10^{-9} 40. Tuna, 1.3×10^{-6}
 41. Grapes, 5.0×10^{-5} 42. Grapefruit, 6.3×10^{-4}

Find the hydronium ion concentration of the substance with the given pH. **See Example 3.**

43. Human blood plasma, 7.4 44. Milk, 6.4
 45. Human gastric contents, 2.0 46. Spinach, 5.4
 47. Bananas, 4.6 48. Milk of magnesia, 10.5

Solve each problem. **See Examples 4 and 6.**

- 49.** Managements of sports stadiums and arenas often encourage fans to make as much noise as possible. Find the average decibel level

$$D = 10 \log \left(\frac{I}{I_0} \right)$$

for each venue with the given intensity I .

- (a) NFL fans, Kansas City Chiefs at Arrowhead Stadium: $I = (1.58 \times 10^{14})I_0$
 (b) NBA fans, Sacramento Kings at Sleep Train Arena: $I = (3.9 \times 10^{12})I_0$
 (c) MLB fans, Baltimore Orioles at Camden Yards: $I = (1.1 \times 10^{12})I_0$
 (Data from www.baltimoresportsreport.com, www.guinnessworldrecords.com)



- 50.** Find the decibel level of each sound.

- (a) noisy restaurant: $I = 10^8 I_0$
 (b) farm tractor: $I = (5.340 \times 10^9) I_0$
 (c) snowmobile: $I = 31,622,776,600 I_0$
 (Data from The Canadian Society of Otolaryngology.)



51. The number of years, $N(x)$, since two independently evolving languages split off from a common ancestral language is approximated by

$$N(x) = -5000 \ln x,$$

where x is the percent of words (in decimal form) from the ancestral language common to both languages now. Find the number of years (to the nearest hundred years) since the split for each percent of common words.

- (a) 85% (or 0.85) (b) 35% (or 0.35) (c) 10% (or 0.10)

52. The time t in years for an amount of money invested at an interest rate r (in decimal form) to double is given by

$$t(r) = \frac{\ln 2}{\ln(1+r)}.$$

This is the **doubling time**. Find the doubling time to the nearest tenth for an investment at each interest rate.

- (a) 2% (or 0.02) (b) 5% (or 0.05) (c) 8% (or 0.08)

53. The number of monthly active Twitter users (in millions) worldwide during the third quarter of each year from 2010 to 2017 is approximated by

$$f(x) = 25.6829 + 149.1368 \ln x,$$

where $x = 1$ represents 2010, $x = 2$ represents 2011, and so on. (Data from Twitter.)

- (a) What does this model give for the number of monthly active Twitter users in 2012?
 (b) According to this model, when did the number of monthly active Twitter users reach 300 million? (*Hint*: Substitute for $f(x)$, and then write the equation in exponential form to solve it.)
54. In the central Sierra Nevada of California, the percent of moisture that falls as snow rather than rain is approximated by

$$f(x) = 86.3 \ln x - 680,$$

where x is the altitude in feet.

- (a) What percent of the moisture at 5000 ft falls as snow?
 (b) What percent at 7500 ft falls as snow?
55. The approximate tax $T(x)$, in dollars per ton, that would result in an $x\%$ (in decimal form) reduction in carbon dioxide emissions is approximated by the **cost-benefit equation**

$$T(x) = -0.642 - 189 \ln(1-x).$$

- (a) What tax will reduce emissions 25%?
 (b) Explain why the equation is not valid for $x = 0$ or $x = 1$.
56. The age in years of a female blue whale of length x in feet is approximated by

$$f(x) = -2.57 \ln \left(\frac{87-x}{63} \right).$$

- (a) How old is a female blue whale that measures 80 ft?
 (b) The equation that defines this function has domain $24 < x < 87$. Explain why.



Use the change-of-base rule (with either common or natural logarithms) to approximate each logarithm to four decimal places. See Example 7.

- | | | | |
|------------------------|--------------------------|--------------------|--------------------|
| 57. $\log_3 12$ | 58. $\log_4 18$ | 59. $\log_5 3$ | 60. $\log_7 4$ |
| 61. $\log_{21} 0.7496$ | 62. $\log_{19} 0.8325$ | 63. $\log_{1/2} 5$ | 64. $\log_{1/3} 7$ |
| 65. $\log_3 \sqrt{2}$ | 66. $\log_6 \sqrt[3]{5}$ | 67. $\log_\pi e$ | 68. $\log_\pi 10$ |

9.6

Exponential and Logarithmic Equations; Further Applications

OBJECTIVES

- 1 Solve equations involving variables in the exponents.
- 2 Solve equations involving logarithms.
- 3 Solve applications of compound interest.
- 4 Solve applications involving base e exponential growth and decay.

VOCABULARY

- compound interest
- continuous compounding


NOW TRY EXERCISE 1

Solve $5^x = 20$.
Approximate the solution to three decimal places.

General methods for solving these equations depend on the following properties.

Properties for Solving Exponential and Logarithmic Equations

For all real numbers $b > 0$, $b \neq 1$, and any real numbers x and y , the following hold true.

1. If $x = y$, then $b^x = b^y$.
2. If $b^x = b^y$, then $x = y$.
3. If $x = y$, and $x > 0$, $y > 0$, then $\log_b x = \log_b y$.
4. If $x > 0$, $y > 0$, and $\log_b x = \log_b y$, then $x = y$.

OBJECTIVE 1 Solve equations involving variables in the exponents.

EXAMPLE 1 Solving an Exponential Equation

Solve $3^x = 12$. Approximate the solution to three decimal places.

$$3^x = 12$$

$$\log 3^x = \log 12$$

$$x \log 3 = \log 12$$

$$\text{Exact solution} \longrightarrow x = \frac{\log 12}{\log 3}$$

$$\text{Decimal approximation} \longrightarrow x \approx 2.262$$

Property 3 (common logarithms):
If $x = y$, then $\log_b x = \log_b y$.

Power rule

Divide by $\log 3$.

Evaluate with a calculator.

CHECK $3^x = 3^{2.262} \approx 12$ ✓ True

The solution set is $\{2.262\}$.

NOW TRY 

CAUTION Be careful: $\frac{\log 12}{\log 3}$ is **not** equal to $\log 4$.

$$\frac{\log 12}{\log 3} \approx 2.262, \quad \text{but} \quad \log 4 \approx 0.6021.$$

NOTE In Example 1, we could have used Property 3 with natural logarithms.

$$\frac{\ln 12}{\ln 3} = \frac{\log 12}{\log 3} \approx 2.262 \quad (\text{Verify this.})$$

NOW TRY ANSWER

1. $\{1.861\}$


NOW TRY EXERCISE 2
Solve $e^{0.12x} = 10$.

Approximate the solution to three decimal places.

EXAMPLE 2 Solving an Exponential Equation (Base e)
Solve $e^{0.003x} = 40$. Approximate the solution to three decimal places.

$$e^{0.003x} = 40$$

$$\ln e^{0.003x} = \ln 40$$

$$0.003x \ln e = \ln 40$$

$$0.003x = \ln 40$$

$$x = \frac{\ln 40}{0.003}$$

$$x \approx 1229.626$$

Property 3 (natural logarithms):
If $x = y$, then $\ln x = \ln y$.

Power rule

$$\ln e = \ln e^1 = 1$$

Divide by 0.003.

Evaluate with a calculator.

CHECK $e^{0.003x} = e^{0.003(1229.626)} \approx 40$ ✓ TrueThe solution set is $\{1229.626\}$.**NOW TRY** 

NOW TRY EXERCISE 3
Solve $3^{-x+5} = 8$. Approximate the solution to three decimal places.
EXAMPLE 3 Solving an Exponential Equation
Solve $7^{-x+4} = 17$. Approximate the solution to three decimal places.

$$7^{-x+4} = 17$$

$$\log 7^{-x+4} = \log 17$$

$$(-x + 4) \log 7 = \log 17$$

$$-x \log 7 + 4 \log 7 = \log 17$$

$$-x \log 7 = \log 17 - 4 \log 7$$

$$x = \frac{\log 17 - 4 \log 7}{-\log 7}$$

$$x \approx 2.544$$

Property 3 (common logarithms):
If $x = y$, then $\log_b x = \log_b y$.

Power rule

Distributive property

Subtract $4 \log 7$.Divide by $-\log 7$.

Evaluate with a calculator.

CHECK $7^{-x+4} = 7^{-2.544+4} \approx 17$ ✓ TrueThe solution set is $\{2.544\}$.**NOW TRY** 
General Method for Solving an Exponential Equation
Take logarithms having the same base on both sides and then use the power rule of logarithms or the special property $\log_b b^x = x$. (**See Examples 1–3.**)

As a special case, if both sides can be written as exponentials with the same base, do so, and set the exponents equal.

OBJECTIVE 2 Solve equations involving logarithms.
EXAMPLE 4 Solving a Logarithmic Equation
Solve $\log_3(4x + 1) = 4$.

$$\log_3(4x + 1) = 4$$

$$4x + 1 = 3^4$$
 Write in exponential form.

$$4x + 1 = 81$$
 $3^4 = 81$

$$4x = 80$$
 Subtract 1.

$$x = 20$$
 Divide by 4.

NOW TRY ANSWERS2. $\{19.188\}$ 3. $\{3.107\}$

NOW TRY EXERCISE 4Solve $\log_6(2x + 4) = 2$.**CHECK**

$$\log_3(4x + 1) = 4$$

$$\log_3[4(20) + 1] \stackrel{?}{=} 4$$

$$\log_3 81 \stackrel{?}{=} 4$$

$$4 = 4 \quad \checkmark$$

Let $x = 20$.

Work inside the brackets.

True; $\log_3 81 = 4$ A true statement results, so the solution set is $\{20\}$.**NOW TRY** **NOW TRY EXERCISE 5**Solve $\log_5(x - 1)^3 = 2$.

Give the exact solution.

EXAMPLE 5 Solving a Logarithmic EquationSolve $\log_2(x + 5)^3 = 4$. Give the exact solution.

$$\log_2(x + 5)^3 = 4$$

$$(x + 5)^3 = 2^4$$

$$(x + 5)^3 = 16$$

$$x + 5 = \sqrt[3]{16}$$

$$x = -5 + \sqrt[3]{16}$$

$$x = -5 + 2\sqrt[3]{2}$$

Write in exponential form.

$$2^4 = 16$$

Take the cube root on each side.

Add -5 .

$$\sqrt[3]{16} = \sqrt[3]{8 \cdot 2} = \sqrt[3]{8} \cdot \sqrt[3]{2} = 2\sqrt[3]{2}$$

CHECK

$$\log_2(x + 5)^3 = 4$$

$$\log_2(-5 + 2\sqrt[3]{2} + 5)^3 \stackrel{?}{=} 4$$

$$\log_2(2\sqrt[3]{2})^3 \stackrel{?}{=} 4$$

$$\log_2 16 \stackrel{?}{=} 4$$

$$4 = 4 \quad \checkmark$$

Original equation

Let $x = -5 + 2\sqrt[3]{2}$.

Work inside the parentheses.

$$(2\sqrt[3]{2})^3 = 2^3(\sqrt[3]{2})^3 = 8 \cdot 2 = 16$$

True; $\log_2 16 = 4$ A true statement results, so the solution set is $\{-5 + 2\sqrt[3]{2}\}$.**NOW TRY** **! CAUTION** Recall that the domain of $f(x) = \log_b x$ is $(0, \infty)$. *For this reason, always check that each proposed solution of an equation with logarithms yields only logarithms of positive numbers in the original equation.***EXAMPLE 6 Solving a Logarithmic Equation**Solve $\log_2(x + 1) - \log_2 x = \log_2 7$.

$$\log_2(x + 1) - \log_2 x = \log_2 7$$

Transform the left side to an expression with only *one* logarithm.

$$\log_2 \frac{x + 1}{x} = \log_2 7$$

Quotient rule

$$\frac{x + 1}{x} = 7$$

Property 4: If $\log_b x = \log_b y$, then $x = y$.

$$x + 1 = 7x$$

Multiply by x .

$$1 = 6x$$

Subtract x .

This proposed solution must be checked.

$$\frac{1}{6} = x$$

Divide by 6.

NOW TRY ANSWERS4. $\{16\}$ 5. $\{1 + \sqrt[3]{25}\}$ We cannot take the logarithm of a *nonpositive* number, so both $x + 1$ and x must be positive here. If $x = \frac{1}{6}$, then this condition is satisfied.


NOW TRY EXERCISE 6

Solve.

$$\log_4(2x + 13) - \log_4(x + 1) = \log_4 10$$

CHECK $\log_2(x + 1) - \log_2 x = \log_2 7$ Original equation

$$\log_2\left(\frac{1}{6} + 1\right) - \log_2 \frac{1}{6} \stackrel{?}{=} \log_2 7$$
 Let $x = \frac{1}{6}$.

$$\log_2 \frac{7}{6} - \log_2 \frac{1}{6} \stackrel{?}{=} \log_2 7$$
 Add.

$$\log_2 \frac{7}{\frac{1}{6}} \stackrel{?}{=} \log_2 7$$
 Quotient rule

$$\frac{\frac{7}{6}}{\frac{1}{6}} = \frac{7}{6} \div \frac{1}{6} = \frac{7}{6} \cdot \frac{6}{1} = 7$$

$$\log_2 7 = \log_2 7 \quad \checkmark$$
 True

A true statement results, so the solution set is $\left\{\frac{1}{6}\right\}$.

NOW TRY 

NOW TRY EXERCISE 7

Solve.

$$\log_4(x + 2) + \log_4 2x = 2$$

EXAMPLE 7 Solving a Logarithmic Equation
Solve $\log x + \log(x - 21) = 2$.

$$\log x + \log(x - 21) = 2$$

$$\log x(x - 21) = 2$$
 Product rule

$$x(x - 21) = 10^2$$
 Write in exponential form.

$$x^2 - 21x = 100$$
 Distributive property; $10^2 = 100$

$$x^2 - 21x - 100 = 0$$
 Standard form

$$(x - 25)(x + 4) = 0$$
 Factor.

$$x - 25 = 0 \quad \text{or} \quad x + 4 = 0$$
 Zero-factor property

$$x = 25 \quad \text{or} \quad x = -4$$
 Solve each equation.

The value -4 must be rejected as a solution because it leads to the logarithm of a negative number in the original equation.

$$\log(-4) + \log(-4 - 21) = 2$$
 The left side is undefined.

Check that the only solution is 25, so the solution set is $\{25\}$.

NOW TRY 

CAUTION Do not reject a potential solution just because it is nonpositive. Reject any value that leads to the logarithm of a nonpositive number.

Solving a Logarithmic Equation

Step 1 Transform the equation so that a single logarithm appears on one side. Use the product rule or quotient rule of logarithms to do this.

Step 2 Do one of the following.

(a) Use Property 4.

If $\log_b x = \log_b y$, then $x = y$. (See Example 6.)

(b) Write the equation in exponential form.

If $\log_b x = k$, then $x = b^k$. (See Examples 4, 5, and 7.)

NOW TRY ANSWERS

6. $\left\{\frac{3}{8}\right\}$

7. $\{2\}$

OBJECTIVE 3 Solve applications of compound interest.

We have solved simple interest problems using the formula $I = prt$. In most cases, interest paid or charged is **compound interest** (interest paid on both principal and interest). The formula for compound interest is an application of exponential functions. *In this text, monetary amounts are given to the nearest cent.*

Compound Interest Formula (for a Finite Number of Periods)

If a principal of P dollars is deposited at an annual rate of interest r compounded (paid) n times per year, then the account will contain

$$A = P \left(1 + \frac{r}{n} \right)^{nt}$$

dollars after t years. (In this formula, r is expressed as a decimal.)

NOW TRY EXERCISE 8

How much money will there be in an account at the end of 10 yr if \$10,000 is deposited at 2.5% compounded monthly?

EXAMPLE 8 Solving a Compound Interest Problem for A

How much money will there be in an account at the end of 5 yr if \$1000 is deposited at 3% compounded quarterly? (Assume no withdrawals are made.)

Because interest is compounded quarterly, $n = 4$.

$$A = P \left(1 + \frac{r}{n} \right)^{nt} \quad \text{Compound interest formula}$$

$$A = 1000 \left(1 + \frac{0.03}{4} \right)^{4 \cdot 5} \quad \text{Substitute } P = 1000, r = 0.03 \text{ (because } 3\% = 0.03), n = 4, \text{ and } t = 5.$$

$$A = 1000(1.0075)^{20} \quad \text{Simplify.}$$

$$A = 1161.18 \quad \text{Evaluate with a calculator.}$$

The account will contain \$1161.18.

NOW TRY **NOW TRY EXERCISE 9**

Find the number of years, to the nearest hundredth, it will take for money deposited in an account paying 2% interest compounded quarterly to double.

EXAMPLE 9 Solving a Compound Interest Problem for t

Suppose inflation is averaging 3% per year. To the nearest hundredth of a year, how long will it take for prices to double? (This is the **doubling time** of the money.)

We want to find the number of years t for P dollars to grow to $2P$ dollars.

$$A = P \left(1 + \frac{r}{n} \right)^{nt} \quad \text{Compound interest formula}$$

$$2P = P \left(1 + \frac{0.03}{1} \right)^{1t} \quad \text{Let } A = 2P, r = 0.03, \text{ and } n = 1.$$

$$2 = (1.03)^t \quad \text{Divide by } P. \text{ Simplify.}$$

$$\log 2 = \log (1.03)^t \quad \text{Property 3}$$

$$\log 2 = t \log (1.03) \quad \text{Power rule}$$

$$t = \frac{\log 2}{\log 1.03} \quad \text{Interchange sides. Divide by } \log 1.03.$$

$$t \approx 23.45 \quad \text{Evaluate with a calculator.}$$

Prices will double in 23.45 yr. To check, verify that $1.03^{23.45} \approx 2$.

NOW TRY **NOW TRY ANSWERS**

8. \$12,836.92

9. 34.74 yr

Interest can be compounded over various time periods per year, including annually, semiannually, quarterly, daily, and so on.

If the value of n increases without bound, we have an example of **continuous compounding**.

Continuous Compound Interest Formula

If a principal of P dollars is deposited at an annual rate of interest r compounded continuously for t years, the final amount A on deposit is given by

$$A = Pe^{rt}.$$



NOW TRY EXERCISE 10

Suppose that \$4000 is invested at 3% interest for 2 yr.

- (a) How much will the investment grow to if it is compounded continuously?
- (b) How long, to the nearest tenth of a year, would it take for the original investment to double?

EXAMPLE 10 Solving a Continuous Compound Interest Problem

In **Example 8** we found that \$1000 invested for 5 yr at 3% interest compounded quarterly would grow to \$1161.18.

- (a) How much would this investment grow to if it is compounded continuously?

$$A = Pe^{rt} \quad \text{Continuous compounding formula}$$

$$A = 1000e^{0.03(5)} \quad \text{Let } P = 1000, r = 0.03, \text{ and } t = 5.$$

$$A = 1000e^{0.15} \quad \text{Multiply in the exponent.}$$

$$A = 1161.83 \quad \text{Evaluate with a calculator.}$$

The investment will grow to \$1161.83 (which is \$0.65 more than the amount in **Example 8** when interest was compounded quarterly).

- (b) How long, to the nearest tenth of a year, would it take for the initial investment to triple?

We must find the value of t that will cause A to be $3(\$1000) = \3000 .

$$A = Pe^{rt} \quad \text{Continuous compounding formula}$$

$$3000 = 1000e^{0.03t} \quad \text{Let } A = 3P = 3000, P = 1000, \text{ and } r = 0.03.$$

$$3 = e^{0.03t} \quad \text{Divide by 1000.}$$

$$\ln 3 = \ln e^{0.03t} \quad \text{Take natural logarithms.}$$

$$\ln 3 = 0.03t \quad \text{In } e^k = k$$

$$t = \frac{\ln 3}{0.03} \quad \text{Divide by 0.03. Interchange sides.}$$

$$t \approx 36.6 \quad \text{Evaluate with a calculator.}$$

It would take 36.6 yr for the original investment to triple.



OBJECTIVE 4 Solve applications involving base e exponential growth and decay.

When situations involve growth or decay of a population, the amount or number of some quantity present at time t can be approximated by

$$f(t) = y_0 e^{kt}.$$

In this equation, y_0 is the amount or number present at time $t = 0$ and k is a constant.

NOW TRY ANSWERS

10. (a) \$4247.35
(b) 23.1 yr


NOW TRY EXERCISE 11

Radium-226 decays according to the function

$$f(t) = y_0 e^{-0.00043t},$$

where t is time in years.

- (a) If an initial sample contains $y_0 = 4.5$ g of radium-226, how many grams, to the nearest tenth, will be present after 150 yr?
- (b) What is the half-life of radium-226? Round to the nearest year.

The continuous compounding of money is an example of exponential growth. In **Example 11**, we investigate exponential decay.

EXAMPLE 11 Applying an Exponential Decay Function

After a plant or animal dies, the amount of radioactive carbon-14 that is present disintegrates according to the natural logarithmic function

$$f(t) = y_0 e^{-0.000121t},$$

where t is time in years, $f(t)$ is the amount of the sample at time t , and y_0 is the initial amount present at $t = 0$.

- (a) If an initial sample contains $y_0 = 10$ g of carbon-14, how many grams, to the nearest hundredth, will be present after 3000 yr?

$$f(3000) = 10e^{-0.000121(3000)} \approx 6.96 \text{ g} \quad \text{Let } y_0 = 10 \text{ and } t = 3000 \text{ in the formula.}$$

- (b) How long would it take to the nearest year for the initial sample to decay to half of its original amount? (This is the **half-life** of the sample.)

$$f(t) = y_0 e^{-0.000121t} \quad \text{Exponential decay formula}$$

$$5 = 10e^{-0.000121t} \quad \text{Let } f(t) = \frac{1}{2}(10) = 5 \text{ and } y_0 = 10.$$

$$\frac{1}{2} = e^{-0.000121t} \quad \text{Divide by 10.}$$

$$\ln \frac{1}{2} = -0.000121t \quad \text{Take natural logarithms; } \ln e^k = k$$

$$t = \frac{\ln \frac{1}{2}}{-0.000121} \quad \text{Divide by } -0.000121. \text{ Interchange sides.}$$

$$t \approx 5728 \quad \text{Evaluate with a calculator.}$$

The half-life is 5728 yr.

NOW TRY ANSWERS


11. (a) 4.2 g (b) 1612 yr

NOW TRY 

9.6 Exercises

FOR EXTRA HELP

MyLab Math

 Video solutions for select problems available in MyLab Math

Many of the problems in these exercises require a calculator.

Concept Check Determine whether common logarithms or natural logarithms would be a better choice to use for solving each equation. Do not actually solve.

1. $10^{0.0025x} = 75$

2. $10^{3x+1} = 13$

3. $e^{x-2} = 24$

4. $e^{-0.28x} = 30$

Solve each equation. Approximate solutions to three decimal places. See Examples 1 and 3.

5. $7^x = 5$

6. $4^x = 3$

7. $9^{-x+2} = 13$

8. $6^{-x+1} = 22$

9. $3^{2x} = 14$

10. $5^{3x} = 11$

11. $2^{x+3} = 5^x$

12. $6^{x+3} = 4^x$

13. $2^{x+3} = 3^{x-4}$

14. $4^{x-2} = 5^{3x+2}$

15. $4^{2x+3} = 6^{x-1}$

16. $3^{2x+1} = 5^{x-1}$

Solve each equation. Use natural logarithms. Approximate solutions to three decimal places when appropriate. See Example 2.

- | | | |
|--------------------------------|---------------------------------|------------------------------------|
| 17. $e^{0.012x} = 23$ | 18. $e^{0.006x} = 30$ | 19. $e^{-0.205x} = 9$ |
| 20. $e^{-0.103x} = 7$ | 21. $\ln e^{3x} = 9$ | 22. $\ln e^{2x} = 4$ |
| 23. $\ln e^{0.45x} = \sqrt{7}$ | 24. $\ln e^{0.04x} = \sqrt{3}$ | 25. $\ln e^{-x} = \pi$ |
| 26. $\ln e^{2x} = \pi$ | 27. $e^{\ln 2x} = e^{\ln(x+1)}$ | 28. $e^{\ln(6-x)} = e^{\ln(4+2x)}$ |

Solve each equation. Give exact solutions. See Examples 4 and 5.

- | | |
|---------------------------|----------------------------|
| 29. $\log_4(2x + 8) = 2$ | 30. $\log_5(5x + 10) = 3$ |
| 31. $\log_3(6x + 5) = 2$ | 32. $\log_5(12x - 8) = 3$ |
| 33. $\log_2(2x - 1) = 5$ | 34. $\log_6(4x + 2) = 2$ |
| 35. $\log_7(x + 1)^3 = 2$ | 36. $\log_4(x - 3)^3 = 4$ |
| 37. $\log_2(x^2 + 7) = 4$ | 38. $\log_6(x^2 + 11) = 2$ |

39. **Concept Check** Suppose that in solving a logarithmic equation having the term $\log(x - 3)$, we obtain the proposed solution 2. We know that our algebraic work is correct, so we give $\{2\}$ as the solution set. **WHAT WENT WRONG?**
40. **Concept Check** Suppose that in solving a logarithmic equation having the term $\log(3 - x)$, we obtain the proposed solution -4 . We know that our algebraic work is correct, so we reject -4 and give \emptyset as the solution set. **WHAT WENT WRONG?**

Solve each equation. Give exact solutions. See Examples 6 and 7.

- | | |
|--------------------------------------------|------------------------------------------------|
| 41. $\log(6x + 1) = \log 3$ | 42. $\log(7 - 2x) = \log 4$ |
| 43. $\log_5(3t + 2) - \log_5 t = \log_5 4$ | 44. $\log_2(t + 5) - \log_2(t - 1) = \log_2 3$ |
| 45. $\log 4x - \log(x - 3) = \log 2$ | 46. $\log(-x) + \log 3 = \log(2x - 15)$ |
| 47. $\log_2 x + \log_2(x - 7) = 3$ | 48. $\log(2x - 1) + \log 10x = \log 10$ |
| 49. $\log 5x - \log(2x - 1) = \log 4$ | 50. $\log_3 x + \log_3(2x + 5) = 1$ |
| 51. $\log_2 x + \log_2(x - 6) = 4$ | 52. $\log_2 x + \log_2(x + 4) = 5$ |

Solve each problem. See Examples 8–10.

53. Suppose that \$2000 is deposited at 4% compounded quarterly.
- How much money will be in the account at the end of 6 yr? (Assume no withdrawals are made.)
 - To one decimal place, how long will it take for the account to grow to \$3000?
54. Suppose that \$3000 is deposited at 3.5% compounded quarterly.
- How much money will be in the account at the end of 7 yr? (Assume no withdrawals are made.)
 - To one decimal place, how long will it take for the account to grow to \$5000?
55. What will be the amount A in an account with initial principal \$4000 if interest is compounded continuously at an annual rate of 3.5% for 6 yr?
56. What will be the amount A in an account with initial principal \$10,000 if interest is compounded continuously at an annual rate of 2.5% for 5 yr?



57. How long, to the nearest hundredth of a year, would it take an initial principal P to double if it were invested at 2.5% compounded continuously?
58. How long, to the nearest hundredth of a year, would it take \$4000 to double at 3.25% compounded continuously?
59. Find the amount of money in an account after 12 yr if \$5000 is deposited at 7% annual interest compounded as follows.
 (a) Annually (b) Semiannually (c) Quarterly
 (d) Daily (Use $n = 365$.) (e) Continuously
60. Find the amount of money in an account after 8 yr if \$4500 is deposited at 6% annual interest compounded as follows.
 (a) Annually (b) Semiannually (c) Quarterly
 (d) Daily (Use $n = 365$.) (e) Continuously
61. How much money must be deposited today to amount to \$1850 in 40 yr at 6.5% compounded continuously?
62. How much money must be deposited today to amount to \$1000 in 10 yr at 5% compounded continuously?

Solve each problem. See Example 11.

63. Revenues of software publishers in the United States for the years 2004–2016 can be modeled by the function

$$S(x) = 91.412e^{0.05195x},$$

where $x = 4$ represents 2004, $x = 5$ represents 2005, and so on, and $S(x)$ is in billions of dollars. Approximate, to the nearest unit, revenue for 2016. (Data from U.S. Census Bureau.)

64. Based on selected figures obtained during the years 1970–2015, the total number of bachelor's degrees earned in the United States can be modeled by the function

$$D(x) = 792,377e^{0.01798x},$$

where $x = 0$ corresponds to 1970, $x = 5$ corresponds to 1975, and so on. Approximate, to the nearest unit, the number of bachelor's degrees earned in 2015. (Data from U.S. National Center for Education Statistics.)



65. Suppose that the amount, in grams, of plutonium-241 present in a given sample is determined by the function

$$A(t) = 2.00e^{-0.053t},$$

where t is measured in years. Approximate the amount present, to the nearest hundredth, in the sample after the given number of years.

- (a) 4 (b) 10 (c) 20 (d) What was the initial amount present?
66. Suppose that the amount, in grams, of radium-226 present in a given sample is determined by the function

$$A(t) = 3.25e^{-0.00043t},$$

where t is measured in years. Approximate the amount present, to the nearest hundredth, in the sample after the given number of years.

- (a) 20 (b) 100 (c) 500 (d) What was the initial amount present?

67. A sample of 400 g of lead-210 decays to polonium-210 according to the function

$$A(t) = 400e^{-0.032t},$$

where t is time in years. Approximate answers to the nearest hundredth.

- (a) How much lead will be left in the sample after 25 yr?
 (b) How long will it take the initial sample to decay to half of its original amount?
68. The concentration of a drug in a person's system decreases according to the function

$$C(t) = 2e^{-0.125t},$$

where $C(t)$ is in appropriate units, and t is in hours. Approximate answers to the nearest hundredth.

- (a) How much of the drug will be in the system after 1 hr?
 (b) How long will it take for the concentration to be half of its original amount?

RELATING CONCEPTS For Individual or Group Work (Exercises 69–72)

Previously, we solved an equation such as $5^x = 125$ as follows.

$$5^x = 125 \quad \text{Original equation}$$

$$5^x = 5^3 \quad 125 = 5^3$$

$$x = 3 \quad \text{Set exponents equal.}$$

Solution set: $\{3\}$

The method described in this section can also be used to solve this equation. **Work Exercises 69–72 in order, to see how this is done.**

69. Take common logarithms on both sides, and write this equation.
 70. Apply the power rule for logarithms on the left.
 71. Write the equation so that x is alone on the left.
 72. Use a calculator to find the decimal form of the solution. What is the solution set?

Chapter 9

Summary

Key Terms

9.1

one-to-one function
inverse of a function

9.2

exponential function
with base a
asymptote
exponential equation

9.3

logarithm
logarithmic equation
logarithmic function
with base a

9.5

common logarithm
natural logarithm

9.6

compound interest
continuous compounding

New Symbols

$f^{-1}(x)$ inverse of $f(x)$
 $\log_a x$ logarithm of x
with base a

$\log x$ common (base 10)
logarithm of x

$\ln x$ natural (base e)
logarithm of x

e a constant,
approximately
2.718281828459

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- In a **one-to-one function**
 - each x -value corresponds to only one y -value
 - each x -value corresponds to one or more y -values
 - each x -value is the same as each y -value
 - each x -value corresponds to only one y -value and each y -value corresponds to only one x -value.
- If f is a one-to-one function, then the **inverse** of f is
 - the set of all solutions of f
 - the set of all ordered pairs formed by interchanging the coordinates of the ordered pairs of f
 - the set of all ordered pairs that are the opposite (negative) of the coordinates of the ordered pairs of f
 - an equation involving an exponential expression.
- An **exponential function** is a function defined by an expression of the form
 - $f(x) = ax^2 + bx + c$ for real numbers a, b, c ($a \neq 0$)
 - $f(x) = \log_a x$ for positive numbers a and x ($a \neq 1$)
 - $f(x) = a^x$ for all real numbers x ($a > 0, a \neq 1$)
 - $f(x) = \sqrt{x}$ for $x \geq 0$.
- A **logarithmic function** is a function defined by an expression of the form
 - $f(x) = ax^2 + bx + c$ for real numbers a, b, c ($a \neq 0$)
 - $f(x) = \log_a x$ for positive numbers a and x ($a \neq 1$)
 - $f(x) = a^x$ for all real numbers x ($a > 0, a \neq 1$)
 - $f(x) = \sqrt{x}$ for $x \geq 0$.
- A **logarithm** is
 - an exponent
 - a base
 - an equation
 - a polynomial.
- An **asymptote** is
 - a line that a graph intersects just once
 - a line that the graph of a function more and more closely approaches as the x -values increase or decrease without bound
 - the x -axis or y -axis
 - a line about which a graph is symmetric.

ANSWERS

1. D; *Example:* The function $f = \{(0, 2), (1, -1), (3, 5), (-2, 3)\}$ is one-to-one. 2. B; *Example:* The inverse of the one-to-one function f defined in Answer 1 is $f^{-1} = \{(2, 0), (-1, 1), (5, 3), (3, -2)\}$. 3. C; *Examples:* $f(x) = 4^x, g(x) = \left(\frac{1}{2}\right)^x$ 4. B; *Example:* The graph of $f(x) = 2^x$ has the x -axis ($y = 0$) as an asymptote. 5. B; *Examples:* $f(x) = \log_3 x, g(x) = \log_{1/3} x$ 6. A; *Example:* $\log_a x$ is the exponent to which a must be raised to obtain x ; $\log_3 9 = 2$ because $3^2 = 9$.

Quick Review

CONCEPTS

9.1 Inverse Functions

Horizontal Line Test

A function is one-to-one if every horizontal line intersects the graph of the function at most once.

Inverse Functions

For a one-to-one function f defined by an equation $y = f(x)$, find the defining equation of the inverse function f^{-1} as follows.

Step 1 Interchange x and y .

Step 2 Solve for y .

Step 3 Replace y with $f^{-1}(x)$.

EXAMPLES

Find f^{-1} if $f(x) = 2x - 3$.

The graph of f is a nonhorizontal (slanted) straight line, so f is one-to-one by the horizontal line test and has an inverse.

$$f(x) = 2x - 3$$

$$y = 2x - 3$$

Let $y = f(x)$.

$$x = 2y - 3$$

Interchange x and y .

$$x + 3 = 2y$$

Add 3.

$$y = \frac{x + 3}{2}$$

Divide by 2.
Interchange sides.

Solve for y .

$$f^{-1}(x) = \frac{1}{2}x + \frac{3}{2}$$


Replace y with $f^{-1}(x)$; $\frac{a+b}{c} = \frac{a}{c} + \frac{b}{c}$

CONCEPTS

In general, the graph of f^{-1} is the mirror image of the graph of f with respect to the line $y = x$. If the point (a, b) lies on the graph of f , then the point (b, a) lies on the graph of f^{-1} .

EXAMPLES

The graphs of a function f and its inverse f^{-1} are shown here.



9.2 Exponential Functions

For $a > 0$, $a \neq 1$, and all real numbers x ,

$$f(x) = a^x$$

defines the **exponential function with base a** .


Graph of $f(x) = a^x$

1. The graph contains the point $(0, 1)$, which is its y -intercept.
2. When $a > 1$, the graph rises from left to right.
When $0 < a < 1$, the graph falls from left to right.
3. The x -axis is an asymptote.
4. The domain is $(-\infty, \infty)$, and the range is $(0, \infty)$.

$f(x) = 2^x$ defines the exponential function with base 2.

$F(x) = \left(\frac{1}{2}\right)^x$ defines the exponential function with base $\frac{1}{2}$.

x	$f(x) = 2^x$	$F(x) = \left(\frac{1}{2}\right)^x$
-2	$\frac{1}{4}$	4
-1	$\frac{1}{2}$	2
0	1	1
1	2	$\frac{1}{2}$
2	4	$\frac{1}{4}$



9.3 Logarithmic Functions

For all positive real numbers a , where $a \neq 1$, and all positive real numbers x ,

$$y = \log_a x \text{ is equivalent to } x = a^y.$$

Special Properties of Logarithms

For $b > 0$, where $b \neq 1$, the following hold true.

$$\log_b b = 1$$

$$\log_b 1 = 0$$

$$\log_b b^r = r \text{ (} r \text{ is real.)} \quad b^{\log_b r} = r \text{ (} r > 0 \text{)}$$

If a and x are positive real numbers, where $a \neq 1$, then

$$g(x) = \log_a x$$

defines the **logarithmic function with base a** .

Graph of $g(x) = \log_a x$

1. The graph contains the point $(1, 0)$, which is its x -intercept.
2. When $a > 1$, the graph rises from left to right.
When $0 < a < 1$, the graph falls from left to right.
3. The y -axis is an asymptote.
4. The domain is $(0, \infty)$, and the range is $(-\infty, \infty)$.

$$y = \log_2 x \text{ is equivalent to } x = 2^y.$$

Evaluate.


$$\log_3 3 = 1 \quad \log_5 1 = 0$$

$$\log_5 5^6 = 6 \quad 4^{\log_4 6} = 6$$

$g(x) = \log_2 x$ defines the logarithmic function with base 2.

$G(x) = \log_{1/2} x$ defines the logarithmic function with base $\frac{1}{2}$.

x	$g(x) = \log_2 x$	$G(x) = \log_{1/2} x$
$\frac{1}{4}$	-2	2
$\frac{1}{2}$	-1	1
1	0	0
2	1	-1
4	2	-2



CONCEPTS	EXAMPLES
<p>9.4 Properties of Logarithms</p> <p>If x, y, and b are positive real numbers, where $b \neq 1$, and r is any real number, then the following hold true.</p> <p>Product Rule $\log_b xy = \log_b x + \log_b y$</p> <p>Quotient Rule $\log_b \frac{x}{y} = \log_b x - \log_b y$</p> <p>Power Rule $\log_b x^r = r \log_b x$</p>	<p>Rewrite each logarithm.</p> <p>$\log_2 3m = \log_2 3 + \log_2 m$ Product rule</p> <p>$\log_5 \frac{9}{4} = \log_5 9 - \log_5 4$ Quotient rule</p> <p>$\log_{10} 2^3 = 3 \log_{10} 2$ Power rule</p>
<p>9.5 Common and Natural Logarithms</p> <p>Common logarithms (base 10) are used in applications such as pH and loudness of sound. Use the (LOG) key of a calculator to evaluate common logarithms.</p> <p>Natural logarithms (base e) are most often used in applications of growth and decay, such as continuous compounding of money, decay of chemical compounds, and biological growth. Use the (LN) key of a calculator to evaluate natural logarithms.</p> <p>Change-of-Base Rule If $a > 0$, $a \neq 1$, $b > 0$, $b \neq 1$, and $x > 0$, then the following holds true.</p> $\log_a x = \frac{\log_b x}{\log_b a}$	<p>Use the formula pH = $-\log [\text{H}_3\text{O}^+]$ to find the pH of grapes with hydronium ion concentration 5.0×10^{-5}.</p> <p>$\text{pH} = -\log (5.0 \times 10^{-5})$ Substitute.</p> <p>$\text{pH} = -(\log 5.0 + \log 10^{-5})$ Property of logarithms</p> <p>$\text{pH} \approx 4.3$ Evaluate with a calculator.</p> <p>Use the following formula to find doubling time, to the nearest hundredth of a year, for an amount of money invested at 4%.</p> $t(r) = \frac{\ln 2}{\ln (1 + r)}$ Formula for doubling time (in years) $t(0.04) = \frac{\ln 2}{\ln (1 + 0.04)}$ Let $r = 0.04$. $t(0.04) \approx 17.67$ Evaluate with a calculator. <p>The doubling time is 17.67 yr.</p> <p>Use the change-of-base rule to approximate $\log_3 17$.</p> $\log_3 17 = \frac{\ln 17}{\ln 3} = \frac{\log 17}{\log 3} \approx 2.5789$
<p>9.6 Exponential and Logarithmic Equations; Further Applications</p> <p>To solve exponential equations, use these properties (where $b > 0$, $b \neq 1$).</p> <ol style="list-style-type: none"> If $b^x = b^y$, then $x = y$. If $x = y$ and $x > 0$, $y > 0$, then $\log_b x = \log_b y$. 	<p>Solve each equation.</p> $2^{3x} = 2^5$ $3x = 5$ Set the exponents equal. $x = \frac{5}{3}$ Divide by 3. <p>Solution set: $\left\{\frac{5}{3}\right\}$</p> $5^x = 8$ $\log 5^x = \log 8$ Take common logarithms. $x \log 5 = \log 8$ Power rule $x = \frac{\log 8}{\log 5}$ Divide by $\log 5$. $x \approx 1.292$ Evaluate with a calculator. <p>Solution set: $\{1.292\}$</p>

CONCEPTS

To solve logarithmic equations, use these properties (where $b > 0$, $b \neq 1$, $x > 0$, and $y > 0$). First use the product rule, quotient rule, power rule, or special properties as necessary to write the equation in the proper form.

1. If $\log_b x = \log_b y$, then $x = y$.

2. If $\log_b x = k$, then $x = b^k$.

Always check that each proposed solution of a logarithmic equation yields only logarithms of positive numbers in the original equation.

EXAMPLES

Solve each equation.

$$\log_3 2x = \log_3 (x + 1)$$

$$2x = x + 1 \quad \text{Property 1}$$

$$x = 1 \quad \text{Subtract } x.$$

A true statement results when 1 is substituted for x .

Solution set: $\{1\}$

$$\log_2 (3x - 1) = 4$$

$$3x - 1 = 2^4 \quad \text{Write in exponential form.}$$

$$3x - 1 = 16 \quad \text{Apply the exponent.}$$

$$3x = 17 \quad \text{Add 1.}$$

$$x = \frac{17}{3} \quad \text{Divide by 3.}$$



A true statement results when $\frac{17}{3}$ is substituted for x .

Solution set: $\left\{\frac{17}{3}\right\}$

Chapter 9

Review Exercises

9.1 Determine whether each graph is the graph of a one-to-one function.



Determine whether each function is one-to-one. If it is, find its inverse.

3. $\{(-2, 4), (-1, 1), (0, 0), (1, 1), (2, 4)\}$

4. $\{(-2, -8), (-1, -1), (0, 0), (1, 1), (2, 8)\}$

5. $f(x) = -3x + 7$

6. $f(x) = \sqrt[3]{6x - 4}$



7. $f(x) = -x^2 + 3$

8. The table lists basic minimum wages in several states. If the set of states is the domain and the set of wage amounts is the range of a function, is it one-to-one? Why or why not?

State	Minimum Wage (in dollars)
Washington	11.50
Massachusetts	11.00
California	10.50
Arizona	10.50
Ohio	8.30
Iowa	7.25

Data from U.S. Department of Labor.

Each function graphed is one-to-one. Graph its inverse.



9.2 Use a calculator to approximate each exponential expression to three decimal places.

11. $5^{3.2}$

12. $\left(\frac{1}{5}\right)^{2.1}$

13. $8.3^{-1.2}$

Graph each exponential function.

14. $f(x) = 3^x$

15. $f(x) = \left(\frac{1}{3}\right)^x$

16. $f(x) = 2^{2x+3}$

Solve each equation.

17. $5^{2x+1} = 25$

18. $4^{3x} = 8^{x+4}$

19. $\left(\frac{1}{27}\right)^{x-1} = 9^{2x}$

20. The U.S. Hispanic population (in millions) can be approximated by

$$f(x) = 46.9 \cdot 2^{0.035811x},$$

where x represents the number of years since 2008. Use this function to approximate, to the nearest tenth, the Hispanic population in each year. (Data from U.S. Census Bureau.)

(a) 2015

(b) 2030



9.3 Work each problem.

21. Convert each equation to the indicated form.

(a) Write in exponential form: $\log_5 625 = 4$.

(b) Write in logarithmic form: $5^{-2} = 0.04$.

22. Fill in each blank with the correct response: The value of $\log_3 81$ is _____. This means that if we raise _____ to the _____ power, the result is _____.

Use a calculator to approximate each logarithm to four decimal places.

23. $\log_2 6$

24. $\log_7 3$

25. $\log_{10} 55$

Graph each logarithmic function.

26. $g(x) = \log_3 x$

27. $g(x) = \log_{1/3} x$

28. Use the special properties of logarithms to evaluate each expression.

(a) $4^{\log_4 12}$

(b) $\log_9 9^{13}$

(c) $\log_5 625$

Solve each equation.

$$\begin{array}{lll} 29. \log_8 64 = x & 30. \log_2 \sqrt{8} = x & 31. \log_x \left(\frac{1}{49} \right) = -2 \\ 32. \log_4 x = \frac{3}{2} & 33. \log_k 4 = 1 & 34. \log_6 x = -2 \end{array}$$

A company has found that total sales, in thousands of dollars, are given by the function

$$S(x) = 100 \log_2(x + 2),$$

where x is the number of weeks after a major advertising campaign was introduced.

35. What were total sales 6 weeks after the campaign was introduced?
36. Graph the function.

9.4 Use the properties of logarithms to express each logarithm as a sum or difference of logarithms. Assume that all variables represent positive real numbers.

$$\begin{array}{ll} 37. \log_4 3x^2 & 38. \log_5 \frac{a^3 b^2}{c^4} \\ 39. \log_4 \frac{\sqrt{x} \cdot w^2}{z} & 40. \log_2 \frac{p^2 r}{\sqrt{z}} \end{array}$$

Use the properties of logarithms to rewrite each expression as a single logarithm. Assume that all variables are defined in such a way that the variable expressions are positive, and bases are positive numbers not equal to 1.

$$\begin{array}{ll} 41. 2 \log_a 7 - 4 \log_a 2 & 42. 3 \log_a 5 + \frac{1}{3} \log_a 8 \\ 43. \log_b 3 + \log_b x - 2 \log_b y & 44. \log_3(x + 7) - \log_3(4x + 6) \end{array}$$

9.5 Evaluate each logarithm to four decimal places.

$$45. \log 28.9 \quad 46. \log 0.257 \quad 47. \ln 28.9 \quad 48. \ln 0.257$$

Use the change-of-base rule (with either common or natural logarithms) to approximate each logarithm to four decimal places.

$$49. \log_{16} 13 \quad 50. \log_4 12 \quad 51. \log_{1/4} 17 \quad 52. \log_\pi \sqrt{2}$$

Find the pH (to the nearest tenth) of the substance with the given hydronium ion concentration.

$$53. \text{Milk, } 4.0 \times 10^{-7} \quad 54. \text{Crackers, } 3.8 \times 10^{-9}$$

Solve each problem.

55. If orange juice has pH 4.6, what is its hydronium ion concentration?
56. The magnitude of a star is given by the equation

$$M = 6 - 2.5 \log \frac{I}{I_0},$$

where I_0 is the measure of the faintest star and I is the actual intensity of the star being measured. The dimmest stars are of magnitude 6, and the brightest are of magnitude 1. Determine the ratio of intensities between stars of magnitude 1 and 3.

57. The concentration of a drug injected into the bloodstream decreases with time. The intervals of time T when the drug should be administered are given by

$$T = \frac{1}{k} \ln \frac{C_2}{C_1},$$

where k is a constant determined by the drug in use, C_2 is the concentration at which the drug is harmful, and C_1 is the concentration below which the drug is ineffective. (Data from Horelick, B. and S. Koont, "Applications of Calculus to Medicine: Prescribing Safe and Effective Dosage," *UMAP Module 202*.) Thus, if $T = 4$, the drug should be administered every 4 hr.

For a certain drug, $k = \frac{1}{3}$, $C_2 = 5$, and $C_1 = 2$. How often should the drug be administered? (*Hint*: Round down.)

9.6 Solve each equation. Approximate solutions to three decimal places.

58. $3^x = 9.42$ 59. $2^{x-1} = 15$ 60. $e^{0.06x} = 3$

Solve each equation. Give exact solutions.

61. $\log_3(9x + 8) = 2$ 62. $\log_5(x + 6)^3 = 2$
 63. $\log_3(x + 2) - \log_3 x = \log_3 2$ 64. $\log(2x + 3) = 1 + \log x$
 65. $\log_4 x + \log_4(8 - x) = 2$ 66. $\log_2 x + \log_2(x + 15) = \log_2 16$

Solve each problem.

67. Suppose \$20,000 is deposited at 4% annual interest compounded quarterly. How much will be in the account at the end of 5 yr? (Assume no withdrawals are made.)
 68. How much will \$10,000 compounded continuously at 3.75% annual interest amount to in 3 yr?
 69. Which plan is better? How much more would it pay?

Plan A: Invest \$1000 at 4% compounded quarterly for 3 yr

Plan B: Invest \$1000 at 3.9% compounded monthly for 3 yr



70. Find the half-life of a radioactive substance that decays according to the function

$$Q(t) = A_0 e^{-0.05t}, \quad \text{where } t \text{ is in days, to the nearest tenth.}$$

Evaluate.

1. $\log_2 128$ 2. $\log_{12} 1$ 3. $\log_{2/3} \frac{27}{8}$ 4. $5^{\log_5 36}$
 5. $e^{\ln 4}$ 6. $10^{\log e}$ 7. $\log_3 3^{-5}$ 8. $\ln e^{5.4}$

Solve.

9. $\log_3(x + 9) = 4$ 10. $\log_2 32 = x$ 11. $\log_x \frac{1}{81} = 2$
 12. $27^x = 81$ 13. $2^{2x-3} = 8$ 14. $5^{x+2} = 25^{2x+1}$
 15. $\log_3(x + 1) - \log_3 x = 2$ 16. $\log(3x - 1) = \log 10$
 17. $\log_4(x + 2) - \log_4 x = 3$ 18. $\ln(x^2 + 3x + 4) = \ln 2$

19. A machine purchased for business use **depreciates**, or loses value, over a period of years. The value of the machine at the end of its useful life is its **scrap value**. By one method of depreciation, the scrap value, S , is given by

$$S = C(1 - r)^n,$$

where C is original cost, n is useful life in years, and r is the constant percent of depreciation.

- (a) Find the scrap value of a machine costing \$30,000, having a useful life of 12 yr and a constant annual rate of depreciation of 15%. Round to the nearest dollar.
 (b) A machine has a “half-life” of 6 yr. Find the constant annual rate of depreciation, to the nearest percent.
20. One measure of the diversity of the species in an ecological community is the **index of diversity**, given by the logarithmic expression

$$-(p_1 \ln p_1 + p_2 \ln p_2 + \cdots + p_n \ln p_n),$$

where p_1, p_2, \dots, p_n are the proportions of a sample belonging to each of n species in the sample. (Data from Ludwig, J. and J. Reynolds, *Statistical Ecology: A Primer on Methods and Computing*, New York, John Wiley and Sons.)



Approximate the index of diversity to the nearest thousandth if a sample of 100 from a community produces the following numbers.

- (a) 90 of one species, 10 of another (b) 60 of one species, 40 of another
21. To solve the equation $5^x = 7$, we must find the exponent to which 5 must be raised in order to obtain 7. This is $\log_5 7$.
- (a) Use the change-of-base rule and a calculator to find $\log_5 7$.
 (b) Raise 5 to the number found in part (a). What is the result?
 (c) Using as many decimal places as the calculator gives, write the solution set of $5^x = 7$.
22. Let m be the number of letters in your first name, and let n be the number of letters in your last name.
- (a) Explain what $\log_m n$ means. (b) Use a calculator to find $\log_m n$.
 (c) Raise m to the power indicated by the number found in part (b). What is the result?

Chapter 9

Test

FOR
EXTRA
HELP


Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

► View the complete solutions to all Chapter Test exercises in MyLab Math.

1. Decide whether each function is one-to-one.

(a) $f(x) = x^2 + 9$


(b)



2. Find $f^{-1}(x)$ for the one-to-one function

$$f(x) = \sqrt[3]{x+7}.$$

3. Graph the inverse given the graph of $y = f(x)$.



Graph each function.

4. $f(x) = 6^x$

5. $g(x) = \log_6 x$

6. Explain how the graph of the function in **Exercise 5** can be obtained from the graph of the function in **Exercise 4**.

Use the special properties of logarithms to evaluate each expression.

7. $7^{\log_7 9}$

8. $\log_3 3^6$

9. $\log_5 1$

Solve each equation. Give exact solutions.

10. $5^x = \frac{1}{625}$

11. $2^{3x-7} = 8^{2x+2}$

12. The atmospheric pressure (in millibars) at a given altitude x (in meters) is approximated by

$$f(x) = 1013e^{-0.0001341x}.$$

Use this function to approximate the atmospheric pressure, to the nearest unit, at each altitude.

- (a) 2000 m (b) 10,000 m

13. Write in logarithmic form: $4^{-2} = 0.0625$.

14. Write in exponential form: $\log_7 49 = 2$.

Solve each equation.

15. $\log_{1/2} x = -5$

16. $x = \log_9 3$

17. $\log_x 16 = 4$

18. Fill in each blank with the correct response: The value of $\log_2 32$ is _____. This means that if we raise _____ to the _____ power, the result is _____.

Use the properties of logarithms to write each expression as a sum or difference of logarithms. Assume that variables represent positive real numbers.

19. $\log_3 x^2 y$

20. $\log_5 \left(\frac{\sqrt{x}}{yz} \right)$

Use the properties of logarithms to rewrite each expression as a single logarithm. Assume that all variables represent positive real numbers, and that bases are positive numbers not equal to 1.

21. $3 \log_b s - \log_b t$

22. $\frac{1}{4} \log_b r + 2 \log_b s - \frac{2}{3} \log_b t$

Evaluate each logarithm to four decimal places.

23. $\log 23.1$

24. $\ln 0.82$

25. $\log_6 45$

Work each problem.

26. Use the change-of-base rule to express $\log_3 19$ as described.

(a) in terms of common logarithms (b) in terms of natural logarithms

(c) approximated to four decimal places

27. Solve $3^x = 78$, giving the solution to three decimal places.

28. Solve $\log_8(x + 5) + \log_8(x - 2) = 1$.

29. Suppose that \$10,000 is invested at 3.5% annual interest, compounded quarterly.

(a) How much will be in the account in 5 yr if no money is withdrawn?

(b) How long, to the nearest tenth of a year, will it take for the initial principal to double?

30. Suppose that \$15,000 is invested at 3% annual interest, compounded continuously.

(a) How much will be in the account in 5 yr if no money is withdrawn?

(b) How long, to the nearest tenth of a year, will it take for the initial principal to double?

Chapters R-9 Cumulative Review Exercises

1. Write each fraction as a decimal and a percent.

(a) $\frac{1}{20}$ (b) $\frac{5}{4}$

2. Perform the indicated operations. Give answers in lowest terms.

(a) $-\frac{7}{10} - \frac{1}{2}$ (b) $-\frac{8}{15} \div \frac{2}{3}$

3. Multiply or divide as indicated.

(a) 37.5×100 (b) $37.5 \div 1000$

4. Let $S = \left\{ -\frac{9}{4}, -2, -\sqrt{2}, 0, 0.6, \sqrt{11}, \sqrt{-8}, 6, \frac{30}{3} \right\}$. List the elements of S that are members of each set.

(a) Integers (b) Rational numbers (c) Irrational numbers

Simplify each expression.

5. $|-8| + 6 - |-2| - (-6 + 2)$

6. $2(-5) + (-8)(4) - (-3)$

Solve each equation or inequality.

7. $7 - (3 + 4x) + 2x = -5(x - 1) - 3$

8. $2x + 2 \leq 5x - 1$

9. $|2x - 5| = 9$

10. $|4x + 2| > 10$


Graph.

11. $5x + 2y = 10$

12. $-4x + y \leq 5$

13. The graph projects that the number of international travelers to the United States will increase from 51.2 million in 2000 to 90.3 million in 2020.

- (a) Is this the graph of a function?
 (b) What is the slope of the line in the graph? Interpret the slope in the context of international travelers to the United States.



Data from U.S. Department of Commerce.

14. Find an equation of the line passing through $(5, -1)$ and parallel to the line with equation $3x - 4y = 12$. Write the equation in slope-intercept form.

Solve each system.

15. $5x - 3y = 14$
 $2x + 5y = 18$

16. $3x - 2y = 3$
 $x = \frac{2}{3}y + 1$

17. $x + 2y + 3z = 11$
 $3x - y + z = 8$
 $2x + 2y - 3z = -12$

18. Candy worth \$1.00 per lb is to be mixed with 10 lb of candy worth \$1.96 per lb to obtain a mixture that will be sold for \$1.60 per lb. How many pounds of the \$1.00 candy should be used?

Number of Pounds	Price per Pound (in dollars)	Value (in dollars)
x	1.00	$1x$
	1.60	

Perform the indicated operations.

19. $(2p + 3)(3p - 1)$

20. $(4k - 3)^2$

21. $(3m^3 + 2m^2 - 5m) - (8m^3 + 2m - 4)$

22. Divide $15x^3 - x^2 + 22x + 8$ by $3x + 1$.

Factor.

23. $8x + x^3$

24. $24y^2 - 7y - 6$

25. $5z^3 - 19z^2 - 4z$

26. $16a^2 - 25b^4$

27. $8c^3 + d^3$

28. $16r^2 + 56rq + 49q^2$

Perform the indicated operations.

29. $\frac{(5p^3)^4(-3p^7)}{2p^2(4p^4)}$

30. $\frac{x^2 - 9}{x^2 + 7x + 12} \div \frac{x - 3}{x + 5}$

31. $\frac{2}{k + 3} - \frac{5}{k - 2}$

Simplify.

32. $\sqrt{288}$

33. $2\sqrt{32} - 5\sqrt{98}$

34. $(5 + 4i)(5 - 4i)$

Solve each equation or inequality.

35. $\sqrt{2x + 1} - \sqrt{x} = 1$

36. $3x^2 - x - 1 = 0$

37. $x^2 + 2x - 8 > 0$

38. $x^4 - 5x^2 + 4 = 0$

39. $5^{x+3} = \left(\frac{1}{25}\right)^{3x+2}$

40. $\log_5 x + \log_5(x + 4) = 1$

Graph.

41. $f(x) = \frac{1}{3}(x - 1)^2 + 2$ 42. $f(x) = 2^x$ 43. $f(x) = \log_3 x$

44. Use properties of logarithms to write the following as a sum or difference of logarithms. Assume that variables represent positive real numbers.

$$\log \frac{x^3 \sqrt{y}}{z}$$

45. Let the number of bacteria present in a certain culture be given by

$$B(t) = 25,000e^{0.2t},$$

where t is time measured in hours, and $t = 0$ corresponds to noon. Approximate, to the nearest hundred, the number of bacteria present at each time.

- (a) noon (b) 1 P.M. (c) 2 P.M.
 (d) When will the population double?

STUDY SKILLS REMINDER

Have you begun to prepare for your final exam? **Review Study Skill 10, Preparing for Your Math Final Exam.**

This page intentionally left blank



NONLINEAR FUNCTIONS, CONIC SECTIONS, AND NONLINEAR SYSTEMS

An *ellipse*, one of a group of curves known as *conic sections*, has a special reflecting property responsible for “whispering galleries” like that in the Old House Chamber of the U.S. Capitol. We investigate ellipses in this chapter.

10.1 Additional Graphs of Functions

10.2 Circles Revisited and Ellipses

10.3 Hyperbolas and Functions Defined by Radicals

10.4 Nonlinear Systems of Equations

10.5 Second-Degree Inequalities and Systems of Inequalities


10.1 Additional Graphs of Functions

OBJECTIVES

- 1 Recognize graphs of the absolute value, reciprocal, and square root functions, and graph their translations.
- 2 Recognize and graph step functions.

OBJECTIVE 1 Recognize graphs of the absolute value, reciprocal, and square root functions, and graph their translations.

The elementary function $f(x) = |x|$ is the **absolute value function**. This function pairs each real number with its absolute value. Its graph is shown in **FIGURE 1**.



Absolute value function

$$f(x) = |x|$$

Domain: $(-\infty, \infty)$


Range: $[0, \infty)$

FIGURE 1

VOCABULARY

- absolute value function
- reciprocal function
- asymptote
- square root function
- greatest integer function
- step function

Recall that the **reciprocal function** $f(x) = \frac{1}{x}$ pairs every real number except 0 with its reciprocal. Its graph is shown in **FIGURE 2**. Because x can never equal 0, as x gets closer and closer to 0, $\frac{1}{x}$ approaches either ∞ or $-\infty$. Also, $\frac{1}{x}$ can never equal 0, and as x approaches ∞ or $-\infty$, $\frac{1}{x}$ approaches 0. The axes are the **asymptotes** for the function.



Reciprocal function


$$f(x) = \frac{1}{x}$$

Domain: $(-\infty, 0) \cup (0, \infty)$

Range: $(-\infty, 0) \cup (0, \infty)$

FIGURE 2

The **square root function** $f(x) = \sqrt{x}$, also introduced earlier, pairs every non-negative real number with its principal square root. See **FIGURE 3**.



Square root function

$$f(x) = \sqrt{x}$$

Domain: $[0, \infty)$

Range: $[0, \infty)$

FIGURE 3

The graphs of these elementary functions can be shifted, or translated.

NOW TRY EXERCISE 1

Graph $f(x) = \frac{1}{x+3}$. Give the domain and range.

EXAMPLE 1 Applying a Horizontal Shift

Graph $f(x) = |x - 2|$. Give the domain and range.

If $x = 2$, then $f(x) = 0$, which gives the lowest point on the graph $(2, 0)$. The graph of $f(x) = |x - 2|$ has the same shape as that of $f(x) = |x|$ but is *shifted*, or *translated*, to the right 2 units, as shown in **FIGURE 4**.




FIGURE 4

NOW TRY

As seen in **Example 1**, the graph of

$$y = f(x + h) \text{ is a horizontal translation of the graph of } y = f(x).$$

In **Example 2**, we use the fact that the graph of

$$y = f(x) + k \text{ is a vertical translation of the graph of } y = f(x).$$

NOW TRY EXERCISE 2

Graph $f(x) = \sqrt{x + 2}$. Give the domain and range.

EXAMPLE 2 Applying a Vertical Shift

Graph $f(x) = \frac{1}{x} + 3$. Give the domain and range.

The graph is found by shifting the graph of $y = \frac{1}{x}$ up 3 units. See **FIGURE 5**.






FIGURE 5

NOW TRY

NOW TRY ANSWERS



domain: $(-\infty, -3) \cup (-3, \infty)$;
range: $(-\infty, 0) \cup (0, \infty)$



domain: $[0, \infty)$; range: $[2, \infty)$

EXAMPLE 3 Applying Both Horizontal and Vertical Shifts

Graph $f(x) = \sqrt{x + 1} - 4$. Give the domain and range.

This graph has the same shape as that of $f(x) = \sqrt{x}$ but is shifted to the left 1 unit (because $x + 1 = 0$ if $x = -1$) and down 4 units (because of the negative sign in -4). See **FIGURE 6** on the next page.


NOW TRY EXERCISE 3

Graph $f(x) = |x + 1| - 3$.
Give the domain and range.

$$f(x) = \sqrt{x + 1} - 4$$

x	y
-1	-4
0	-3
3	-2

Compare this table of values to that with FIGURE 3.



Domain: $[-1, \infty)$
Range: $[-4, \infty)$

FIGURE 6

NOW TRY

OBJECTIVE 2 Recognize and graph step functions.

The greatest integer function is defined as follows.

$$f(x) = \llbracket x \rrbracket$$

The **greatest integer function**

$$f(x) = \llbracket x \rrbracket$$

pairs every real number x with the greatest integer less than or equal to x .

NOW TRY EXERCISE 4

Evaluate.

- (a) $\llbracket 5 \rrbracket$
- (b) $\llbracket -6 \rrbracket$
- (c) $\llbracket 3.5 \rrbracket$
- (d) $\llbracket -4.1 \rrbracket$


EXAMPLE 4 Finding the Greatest Integer

Evaluate.

- (a) $\llbracket 8 \rrbracket = 8$
 - (b) $\llbracket -1 \rrbracket = -1$
 - (c) $\llbracket 0 \rrbracket = 0$
- If x is an integer, then $\llbracket x \rrbracket = x$.

(d) $\llbracket 7.45 \rrbracket = 7$ The greatest integer less than or equal to 7.45 is 7. This is like “rounding down.”

(e) $\llbracket -2.6 \rrbracket = -3$ Think of a number line with -2.6 graphed on it. Because -3 is to the left of (and therefore less than) -2.6 , the greatest integer less than or equal to -2.6 is -3 , **not** -2 .



NOW TRY


EXAMPLE 5 Graphing the Greatest Integer Function

Graph $f(x) = \llbracket x \rrbracket$. Give the domain and range.

- For $\llbracket x \rrbracket$, if $-1 \leq x < 0$, then $\llbracket x \rrbracket = -1$;
- if $0 \leq x < 1$, then $\llbracket x \rrbracket = 0$;
- if $1 \leq x < 2$, then $\llbracket x \rrbracket = 1$;
- if $2 \leq x < 3$, then $\llbracket x \rrbracket = 2$;
- if $3 \leq x < 4$, then $\llbracket x \rrbracket = 3$, and so on.

The graph, shown in FIGURE 7 on the next page, consists of a series of horizontal line segments. In each segment, the left endpoint is included and the right endpoint is excluded. These segments continue infinitely following this pattern to the left and right. The appearance of the graph is the reason why this function is called a **step function**.

NOW TRY ANSWERS



domain: $(-\infty, \infty)$; range: $[-3, \infty)$

4. (a) 5 (b) -6 (c) 3 (d) -5

NOW TRY EXERCISE 5

Graph $f(x) = \llbracket x - 1 \rrbracket$. Give the domain and range.




FIGURE 7

Greatest integer function

$$f(x) = \llbracket x \rrbracket$$

Domain: $(-\infty, \infty)$

Range: $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
(the set of integers)

The ellipsis points indicate that the graph continues indefinitely in the same pattern.

The graph of a step function also may be shifted. For example, the graph of

$$h(x) = \llbracket x - 2 \rrbracket \text{ is the graph of } f(x) = \llbracket x \rrbracket \text{ shifted to the right 2 units.}$$

Similarly, the graph of

$$g(x) = \llbracket x \rrbracket + 2 \text{ is the graph of } f(x) \text{ shifted up 2 units. } \text{ NOW TRY } \curvearrowright$$

NOW TRY EXERCISE 6

The cost of parking a car at an airport hourly parking lot is \$4 for the first hour and \$2 for each additional hour or fraction thereof. Let $y = f(x)$ represent the cost of parking a car for x hours. Graph $f(x)$ for x in the interval $(0, 5]$.

EXAMPLE 6 Applying a Greatest Integer Function

An overnight delivery service charges \$25 for a package weighing up to 2 lb. For each additional pound or fraction of a pound, there is an additional charge of \$3. Let $y = D(x)$ represent the cost to send a package weighing x pounds. Graph $D(x)$ for x in the interval $(0, 6]$.

$$\text{For } x \text{ in the interval } (0, 2], \quad y = 25.$$

$$\text{For } x \text{ in the interval } (2, 3], \quad y = 25 + 3 = 28.$$

$$\text{For } x \text{ in the interval } (3, 4], \quad y = 28 + 3 = 31.$$

$$\text{For } x \text{ in the interval } (4, 5], \quad y = 31 + 3 = 34.$$

$$\text{For } x \text{ in the interval } (5, 6], \quad y = 34 + 3 = 37.$$

The graph, which is that of a step function, is shown in FIGURE 8.

NOW TRY ANSWERS

5.
domain: $(-\infty, \infty)$;
range: $\{\dots, -2, -1, 0, 1, 2, \dots\}$

6.
Dollars
Hours




FIGURE 8









10.1 Exercises

FOR
EXTRA
HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Refer to the basic graphs in A–F to answer each of the following.




- Which is the graph of $f(x) = |x|$? The lowest point on its graph has coordinates (____, ____).
- Which is the graph of $f(x) = x^2$? Give the domain and range.
- Which is the graph of $f(x) = [x]$? Give the domain and range.
- Which is the graph of $f(x) = \sqrt{x}$? Give the domain and range.
- Which is not the graph of a function? Why?
- Which is the graph of $f(x) = \frac{1}{x}$? The lines with equations $x = 0$ and $y = 0$ are called its _____.

Concept Check Without actually plotting points, match each function defined by the absolute value expression with its graph.


7. $f(x) = |x - 2| + 2$

A.




8. $f(x) = |x + 2| + 2$

B.




9. $f(x) = |x - 2| - 2$

C.



10. $f(x) = |x + 2| - 2$

D.



- Concept Check** How is the graph of $f(x) = \frac{1}{x-3} + 2$ obtained from the graph of $g(x) = \frac{1}{x}$?
- Concept Check** How is the graph of $f(x) = \frac{1}{x+5} - 3$ obtained from the graph of $g(x) = \frac{1}{x}$?

Graph each function. Give the domain and range. See Examples 1–3.

13. $f(x) = |x + 1|$

14. $f(x) = |x - 1|$

15. $f(x) = \frac{1}{x} + 1$

16. $f(x) = \frac{1}{x} - 1$

17. $f(x) = \sqrt{x - 2}$

18. $f(x) = \sqrt{x + 5}$

19. $f(x) = \frac{1}{x - 2}$

20. $f(x) = \frac{1}{x + 2}$

21. $f(x) = \sqrt{x + 3} - 3$

22. $f(x) = \sqrt{x - 2} + 2$

23. $f(x) = |x - 3| + 1$

24. $f(x) = |x + 1| - 4$

25. **Concept Check** A student incorrectly evaluated $\lceil -5.1 \rceil$ as -5 . **WHAT WENT WRONG?** Evaluate correctly.

26. **Concept Check** A student incorrectly evaluated $\lceil 3.75 \rceil$ as 4 . **WHAT WENT WRONG?** Evaluate correctly.

Evaluate each expression. See Example 4.

27. $\lceil 3 \rceil$

28. $\lceil 18 \rceil$

29. $\lceil 4.5 \rceil$

30. $\lceil 8.7 \rceil$

31. $\left\lfloor \frac{1}{2} \right\rfloor$

32. $\left\lfloor \frac{3}{4} \right\rfloor$

33. $\left\lfloor \frac{8}{3} \right\rfloor$

34. $\left\lfloor \frac{5}{2} \right\rfloor$

35. $\lceil -14 \rceil$

36. $\lceil -5 \rceil$

37. $\lceil -10.1 \rceil$

38. $\lceil -6.9 \rceil$

Graph each step function. See Examples 5 and 6.

39. $f(x) = \lceil x - 3 \rceil$

40. $f(x) = \lceil x + 2 \rceil$

41. $f(x) = \lceil x \rceil - 1$


42. $f(x) = \lceil x \rceil + 1$

Solve each problem. See Example 6.

43. Suppose that postage rates are \$0.55 for the first ounce, plus \$0.24 for each additional ounce, and that each letter carries one \$0.55 stamp and as many \$0.24 stamps as necessary. Graph the function

$$y = p(x) = \text{the number of stamps}$$


on a letter weighing x ounces. Use the interval $(0, 5]$.



44. The cost of parking a car at an airport hourly parking lot is \$3 for the first half-hour and \$2 for each additional half-hour or fraction thereof. Graph the function

$$y = f(x) = \text{the cost of parking a car}$$

for x hours. Use the interval $(0, 2]$.



45. A certain long-distance carrier provides service between Podunk and Nowhereville. If x represents the number of minutes for the call, where $x > 0$, then the function f defined by

$$f(x) = 0.40\lceil x \rceil + 0.75$$

gives the total cost of the call in dollars. Find the cost of a 5.5-minute call.

46. Total rental cost in dollars for a power washer, where x represents the number of hours with $x > 0$, can be represented by the function

$$f(x) = 12\lceil x \rceil + 25.$$

Find the cost of a $7\frac{1}{2}$ -hr rental.

10.2 Circles Revisited and Ellipses

OBJECTIVES

- 1 Graph circles.
- 2 Write an equation of a circle given its center and radius.
- 3 Determine the center and radius of a circle given its equation.
- 4 Recognize the equation of an ellipse.
- 5 Graph ellipses.

When an infinite cone is intersected by a plane, the resulting figure is called a **conic section**. The parabola is one example of a conic section. Circles, ellipses, and hyperbolas may also result. See FIGURE 9.




FIGURE 9

VOCABULARY

- conic section
- circle
- center (of a circle)
- radius
- ellipse
- foci (singular *focus*)
- center (of an ellipse)

OBJECTIVE 1 Graph circles.

Recall from our earlier work that a **circle** is a set of all points in a plane that lie a fixed distance from a fixed point. This fixed point is the **center** of the circle. The fixed distance is its **radius**.

Suppose that the point (h, k) is the center of a circle with radius r . We let (x, y) represent any point on the circle. See FIGURE 10. By definition, the distance between (h, k) and (x, y) must be r .

$$\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = r \quad \text{Distance formula}$$

$$\sqrt{(x - h)^2 + (y - k)^2} = r \quad \text{Let } (h, k) = (x_1, y_1) \text{ and } (x, y) = (x_2, y_2).$$

$$(x - h)^2 + (y - k)^2 = r^2 \quad \text{Square each side.}$$




FIGURE 10

This result is the *center-radius form* of the equation of a circle.

Equation of a Circle (Center-Radius Form)

A circle with center (h, k) and radius r has an equation that can be written in the form

$$(x - h)^2 + (y - k)^2 = r^2 \quad (\text{where } r > 0).$$

If a circle has its center at the origin $(0, 0)$, then its equation becomes

$$x^2 + y^2 = r^2. \quad \text{Here, } h = 0 \text{ and } k = 0.$$

NOW TRY EXERCISE 1

Find the center and radius of each circle. Then graph the circle.

- (a) $(x - 2)^2 + (y - 3)^2 = 9$
- (b) $x^2 + y^2 = 49$

EXAMPLE 1 Graphing Circles

Find the center and radius of each circle. Then graph the circle.

(a) $(x - 3)^2 + (y + 1)^2 = 16$ The radius r is the positive square root of 16.

$$(x - 3)^2 + [y - (-1)]^2 = 4^2 \quad (x - h)^2 + (y - k)^2 = r^2$$

\uparrow
 \uparrow
 \uparrow
 h
 k
 r

The center (h, k) is $(3, -1)$, and the radius r is 4.

To graph the circle, plot the center $(3, -1)$, and then move right, left, up, and down 4 units from the center to plot four points on the circle. Draw a smooth curve through the four points. See **FIGURE 11**. (The center is *not* part of the graph.)




FIGURE 11




FIGURE 12

(b) $x^2 + y^2 = 64$


$$(x - 0)^2 + (y - 0)^2 = 8^2 \quad \text{Here, } h = 0 \text{ and } k = 0.$$

The center is $(0, 0)$ and the radius is 8. The graph, which uses a scale of 2 on the axes, is shown in **FIGURE 12**.


NOW TRY

NOW TRY ANSWERS

1. (a) center: $(2, 3)$; radius: 3



- (b) center: $(0, 0)$; radius: 7



NOTE The x - and y -coordinates of the center of a circle are the values of x and y that make each squared binomial in its center-radius form equal 0. Consider the equation

$$(x - 3)^2 + (y + 1)^2 = 16. \quad \text{See Example 1(a).}$$

If we let $x = 3$, then $(x - 3)^2$ equals 0. If we let $y = -1$, then $(y + 1)^2$ equals 0. Thus, the center of the circle is $(3, -1)$.

NOW TRY EXERCISE 2

Write the center-radius form of each circle described. Then graph the circle.

- (a) Center: $(1, -2)$; radius: 3
- (b) Center: $(0, 0)$; radius: $\sqrt{7}$

OBJECTIVE 2 Write an equation of a circle given its center and radius.

EXAMPLE 2 Writing Equations of Circles and Graphing Them

Write the center-radius form of each circle described. Then graph the circle.

- (a) Center: $(-3, 4)$; radius: 2

Here, $(h, k) = (-3, 4)$ and $r = 2$.

$$(x - h)^2 + (y - k)^2 = r^2 \quad \text{Center-radius form}$$

$$[x - (-3)]^2 + (y - 4)^2 = 2^2 \quad \text{Substitute for } h, k, \text{ and } r.$$

Watch signs here. $(x + 3)^2 + (y - 4)^2 = 4$ Simplify.

See FIGURE 13.




FIGURE 13




FIGURE 14

- (b) Center: $(2, 0)$; radius: $\sqrt{10}$

$$(x - h)^2 + (y - k)^2 = r^2 \quad \text{Center-radius form}$$

$$(x - 2)^2 + (y - 0)^2 = (\sqrt{10})^2 \quad \text{Let } h = 2, k = 0, \text{ and } r = \sqrt{10}.$$

$$(x - 2)^2 + y^2 = 10 \quad \text{Simplify; } (\sqrt{a})^2 = a.$$


The graph is shown in FIGURE 14. For the radius,

$$r = \sqrt{10} \approx 3.16. \quad \text{Evaluate with a calculator.}$$


NOW TRY

NOW TRY ANSWERS

- 2. (a) $(x - 1)^2 + (y + 2)^2 = 9$



- (b) $x^2 + y^2 = 7$



OBJECTIVE 3 Determine the center and radius of a circle given its equation.

If we begin with the center-radius form of the equation of a circle and rewrite it so that the binomials are expanded and the right side equals 0, we obtain another form of the equation. Consider the following.

$$(x + 3)^2 + (y - 4)^2 = 4 \quad \text{See Example 2(a).}$$

Remember the middle term when squaring each binomial. $x^2 + 6x + 9 + y^2 - 8y + 16 = 4$ Square each binomial.

$$x^2 + y^2 + 6x - 8y + 21 = 0 \quad \text{Subtract 4. Combine and rearrange terms.}$$

This result is a different, yet equivalent, form of the equation of the circle.

Equation of a Circle (General Form)

For some real numbers c , d , and e , an equation of the form

$$x^2 + y^2 + cx + dy + e = 0$$

may represent a circle.

In the above general form, both x^2 - and y^2 -terms have equal coefficients, here 1. If the coefficients are equal but *not* 1, the above form can be obtained by dividing through by that coefficient.


NOW TRY EXERCISE 3

Write the center-radius form of the circle with equation

$$x^2 + y^2 - 8x + 10y - 8 = 0.$$

Give the center and radius.




FIGURE 15

EXAMPLE 3 Completing the Square to Write the Center-Radius Form

Write the center-radius form of the circle with equation

$$x^2 + y^2 + 2x + 6y - 15 = 0.$$

Give the center and radius, and graph the circle.

To write the center-radius form, we complete the squares on x and y .

$$x^2 + y^2 + 2x + 6y - 15 = 0$$

$$x^2 + y^2 + 2x + 6y = 15$$

$$(x^2 + 2x \quad) + (y^2 + 6y \quad) = 15$$

$$\left[\frac{1}{2}(2)\right]^2 = 1 \quad \left[\frac{1}{2}(6)\right]^2 = 9$$

$$(x^2 + 2x + 1) + (y^2 + 6y + 9) = 15 + 1 + 9$$

Add 1 and 9
on both sides
of the equation.

$$(x + 1)^2 + (y + 3)^2 = 25$$

$$[x - (-1)]^2 + [y - (-3)]^2 = 5^2$$

Transform so that the constant is on the right.

Write in anticipation of completing the square.

Square half the coefficient of each middle term.

Complete the squares on both x and y .

Factor. Add. Center-radius form

Rewrite to identify the center and radius.

The circle has center $(-1, -3)$ and radius 5. See FIGURE 15.

NOW TRY 

NOTE There are three possibilities for the graph of an equation of the form

$$(x - h)^2 + (y - k)^2 = m, \quad \text{where } m \text{ is a constant.}$$

1. If $m > 0$, then $r^2 = m$, and the graph of the equation is a circle with radius \sqrt{m} .
2. If $m = 0$, then the graph of the equation is the single point (h, k) .
3. If $m < 0$, then no points satisfy the equation and the graph is nonexistent.

OBJECTIVE 4 Recognize the equation of an ellipse.

An **ellipse** is the set of all points in a plane the *sum* of whose distances from two fixed points is constant. These fixed points are the **foci** (singular: *focus*). The ellipse in FIGURE 16 has foci $(c, 0)$ and $(-c, 0)$, with x -intercepts $(a, 0)$ and $(-a, 0)$ and y -intercepts $(0, b)$ and $(0, -b)$. It is shown in more advanced courses that $c^2 = a^2 - b^2$ for an ellipse of this type. The origin is the **center** of the ellipse.




FIGURE 16

NOW TRY ANSWER

3. $(x - 4)^2 + (y + 5)^2 = 49$;
center: $(4, -5)$; radius: 7


An ellipse has the following equation.

Equation of an Ellipse

An ellipse with x -intercepts $(a, 0)$ and $(-a, 0)$ and y -intercepts $(0, b)$ and $(0, -b)$ has an equation that can be written in the form

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

NOTE A circle is a special case of an ellipse, where $a^2 = b^2$.



Reflecting property of an ellipse

FIGURE 17

When a ray of light or sound emanating from one focus of an ellipse bounces off the ellipse, it passes through the other focus. See **FIGURE 17**. This reflecting property is responsible for whispering galleries. In the Old House Chamber of the U.S. Capitol, John Quincy Adams was able to listen in on his opponents' conversations—his desk was positioned at one of the foci beneath the ellipsoidal ceiling, and his opponents were located across the room at the other focus. (See the chapter opener.)

Elliptical bicycle gears are designed to respond to the legs' natural strengths and weaknesses. At the top and bottom of the powerstroke, where the legs have the least leverage, the gear offers little resistance, but as the gear rotates, the resistance increases. This allows the legs to apply more power where it is most naturally available. See **FIGURE 18**.




FIGURE 18

OBJECTIVE 5 Graph ellipses.

To graph an ellipse centered at the origin, we plot the four intercepts and sketch the ellipse through those points.

EXAMPLE 4 Graphing Ellipses

Graph each ellipse.

(a) $\frac{x^2}{49} + \frac{y^2}{36} = 1$

Here, $a^2 = 49$, so $a = 7$, and the x -intercepts are $(7, 0)$ and $(-7, 0)$. Similarly, $b^2 = 36$, so $b = 6$, and the y -intercepts are $(0, 6)$ and $(0, -6)$. Plotting the intercepts and sketching the ellipse through them gives the graph in **FIGURE 19**.




FIGURE 19

NOW TRY EXERCISE 4

Graph $\frac{x^2}{16} + \frac{y^2}{25} = 1$.

(b) $\frac{x^2}{36} + \frac{y^2}{121} = 1$

The x -intercepts are $(6, 0)$ and $(-6, 0)$, and the y -intercepts are $(0, 11)$ and $(0, -11)$. Join these intercepts with the smooth curve of an ellipse. See **FIGURE 20**. (Note the scale of 2 on the axes.)




FIGURE 20

NOW TRY

CAUTION Hand-drawn graphs of ellipses are smooth curves that show symmetry with respect to the center. Again, the center is not part of the actual graph. It just provides help in drawing a more accurate graph.

NOW TRY EXERCISE 5

Graph $\frac{(x-3)^2}{36} + \frac{(y-4)^2}{4} = 1$.

EXAMPLE 5 Graphing an Ellipse Shifted Horizontally and Vertically

Graph $\frac{(x-2)^2}{25} + \frac{(y+3)^2}{49} = 1$.

Just as $(x-2)^2$ and $(y+3)^2$ would indicate that the center of a circle would be $(2, -3)$, so it is with this ellipse. **FIGURE 21** shows that the graph goes through the four points

$(2, 4)$, $(7, -3)$, $(2, -10)$, and $(-3, -3)$.






FIGURE 21

The x -values of these points are found by adding $\pm a = \pm 5$ to 2 (the x -value of the center). The y -values are found by adding $\pm b = \pm 7$ to -3 (the y -value of the center).

NOW TRY ANSWERS



NOW TRY

Graphs of circles and ellipses are not graphs of functions—they fail the conditions of the vertical line test. The only conic section whose graph represents a function is the vertical parabola with equation $f(x) = ax^2 + bx + c$.

NOTE A graphing calculator in function mode cannot directly graph a circle or an ellipse because they do not represent functions. We must first solve the equation for y to obtain two functions y_1 and y_2 . The union of these two graphs is the graph of the entire figure. Consider the following equation of a circle.

$$(x + 3)^2 + (y + 2)^2 = 25 \quad \text{Solve the equation for } y.$$

$$(y + 2)^2 = 25 - (x + 3)^2 \quad \text{Subtract } (x + 3)^2.$$


$$y + 2 = \pm \sqrt{25 - (x + 3)^2} \quad \text{Take square roots.}$$

$$y = -2 \pm \sqrt{25 - (x + 3)^2} \quad \text{Add } -2.$$

The two functions to be graphed are

$$y_1 = -2 + \sqrt{25 - (x + 3)^2} \quad \text{and} \quad y_2 = -2 - \sqrt{25 - (x + 3)^2}.$$

To get an undistorted screen, a **square viewing window** must be used. See **FIGURE 22**. The two semicircles seem to be disconnected because the calculator cannot show a true picture of the behavior at these points.



Square Viewing Window

FIGURE 22

10.2 Exercises

FOR
EXTRA
HELP



MyLab Math

▶ Video solutions for select problems available in MyLab Math



Concept Check Complete each statement. Choices may be used once, more than once, or not at all.

circle	parabolas	radius	intercepts
foci	center	ellipse	conic sections

- When a plane intersects an infinite cone at different angles, _____ such as _____, circles, ellipses, and hyperbolas result.
- A set of all points in a plane that lie a fixed distance from a fixed point is a(n) _____. The fixed distance is the _____ and the fixed point is the _____.
- A set of all points in a plane the sum of whose distances from two fixed points is constant is a(n) _____. The fixed points are the _____.
- The equation $(x + 1)^2 + (y - 4)^2 = 4$ represents a(n) _____ with center $[(1, -4)/(1, 4)/(-1, 4)]$ and _____ equal to 2.
- Concept Check** Consider the circle whose equation is $x^2 + y^2 = 25$.
 - What are the coordinates of its center?
 - What is its radius?
 - Sketch its graph.
- Concept Check** Complete the following: The graph of a circle (*is/is not*) the graph of a function because it fails the conditions of the _____. The graph of an ellipse (*is/is not*) the graph of a function.



Concept Check Match each equation with the correct graph.

7. $(x - 3)^2 + (y - 1)^2 = 25$



8. $(x + 3)^2 + (y - 1)^2 = 25$

9. $(x - 3)^2 + (y + 1)^2 = 25$



10. $(x + 3)^2 + (y + 1)^2 = 25$

Find the center and radius of each circle. Then graph the circle. See Example 1.

11. $x^2 + y^2 = 4$

12. $x^2 + y^2 = 1$

13. $x^2 + y^2 = 81$

14. $x^2 + y^2 = 36$

15. $(x - 5)^2 + (y + 4)^2 = 49$

16. $(x + 2)^2 + (y - 5)^2 = 16$

17. $(x + 1)^2 + (y + 3)^2 = 25$

18. $(x - 3)^2 + (y - 2)^2 = 4$

Write the center-radius form of each circle described. Then graph the circle. See Examples 1 and 2.

19. Center: $(0, 0)$; radius: 4

20. Center: $(0, 0)$; radius: 3

21. Center: $(-3, 2)$; radius: 3

22. Center: $(1, -3)$; radius: 4

23. Center: $(4, 3)$; radius: 5

24. Center: $(-3, -2)$; radius: 6

25. Center: $(-2, 0)$; radius: $\sqrt{5}$

26. Center: $(3, 0)$; radius: $\sqrt{13}$

27. Center: $(0, -3)$; radius: 7

28. Center: $(0, 4)$; radius: 4

Write the center-radius form of the circle with the given equation. Give the center and radius. (Hint: Divide each side by a common factor as needed.) See Example 3.

29. $x^2 + y^2 + 4x + 6y + 9 = 0$

30. $x^2 + y^2 - 8x - 12y + 3 = 0$

31. $x^2 + y^2 + 10x - 14y - 7 = 0$

32. $x^2 + y^2 - 2x + 4y - 4 = 0$

33. $3x^2 + 3y^2 - 12x - 24y + 12 = 0$

34. $2x^2 + 2y^2 + 20x + 16y + 10 = 0$

Write the center-radius form of the circle with the given equation. Give the center and radius, and graph the circle. See Example 3.

35. $x^2 + y^2 - 4x - 6y + 9 = 0$

36. $x^2 + y^2 + 8x + 2y - 8 = 0$

37. $x^2 + y^2 + 6x - 6y + 9 = 0$


38. $x^2 + y^2 - 4x + 10y + 20 = 0$

Concept Check Answer each question, and give a short explanation. See the Note following Example 3.


39. How many points are there on the graph of $(x - 4)^2 + (y - 1)^2 = 0$?

40. How many points are there on the graph of $(x - 4)^2 + (y - 1)^2 = -1$?

- 41. Concept Check** A circle can be drawn on a piece of posterboard by fastening one end of a string with a thumbtack, pulling the string taut with a pencil, and tracing a curve, as shown in the figure. Why does this method work?



- 42. Concept Check** An ellipse can be drawn on a piece of posterboard by fastening two ends of a length of string with thumbtacks, pulling the string taut with a pencil, and tracing a curve, as shown in the figure. Why does this method work?



Graph each ellipse. See Examples 4 and 5.

43. $\frac{x^2}{9} + \frac{y^2}{25} = 1$

44. $\frac{x^2}{9} + \frac{y^2}{16} = 1$

45. $\frac{x^2}{36} + \frac{y^2}{16} = 1$

46. $\frac{x^2}{9} + \frac{y^2}{4} = 1$

47. $\frac{x^2}{16} + \frac{y^2}{4} = 1$

48. $\frac{x^2}{49} + \frac{y^2}{81} = 1$

49. $\frac{y^2}{25} = 1 - \frac{x^2}{49}$

50. $\frac{y^2}{9} = 1 - \frac{x^2}{16}$

51. $\frac{(x+1)^2}{64} + \frac{(y-2)^2}{49} = 1$

52. $\frac{(x-4)^2}{9} + \frac{(y+2)^2}{4} = 1$

53. $\frac{(x-2)^2}{16} + \frac{(y-1)^2}{9} = 1$

54. $\frac{(x+3)^2}{25} + \frac{(y+2)^2}{36} = 1$

Extending Skills A *lithotripter* is a machine used to crush kidney stones using shock waves. The patient is in an elliptical tub with the kidney stone at one focus of the ellipse. Each beam is projected from the other focus to the tub so that it reflects to hit the kidney stone. See the figure.

55. Suppose a lithotripter is based on an ellipse with equation


$$\frac{x^2}{36} + \frac{y^2}{9} = 1.$$

How far from the center of the ellipse must the kidney stone and the source of the beam be placed? (*Hint*: Use the fact that $c^2 = a^2 - b^2$ because $a > b$ here.)

56. Rework Exercise 55 if the equation of the ellipse is

$$9x^2 + 4y^2 = 36.$$

(*Hint*: Write the equation in fractional form by dividing each term by 36, and use $c^2 = b^2 - a^2$ because $b > a$ here.)




The top of an ellipse is illustrated in this depiction of how a lithotripter crushes a kidney stone.

Extending Skills Solve each problem.

57. An arch has the shape of half an ellipse. The equation of the ellipse, where x and y are in meters, is


$$100x^2 + 324y^2 = 32,400.$$

- (a) How high is the center of the arch?
 (b) How wide is the arch across the bottom?



Not to scale


58. A one-way street passes under an overpass, which is in the form of the top half of an ellipse, as shown in the figure. Suppose that a truck 12 ft wide passes directly under the overpass. What is the maximum possible height of this truck?



Not to scale

59. Work each of the following.

- (a) The Roman Colosseum is an ellipse with $a = 310$ ft and $b = \frac{513}{2}$ ft. Find the distance, to the nearest tenth, between the foci of this ellipse.
 (b) The approximate perimeter of an ellipse is given by




$$P \approx 2\pi \sqrt{\frac{a^2 + b^2}{2}},$$

where a and b are the lengths given in part (a). Use this formula to find the approximate perimeter, to the nearest tenth, of the Roman Colosseum.

60. A satellite is in an elliptical orbit around Earth with least distance (**perigee**) of 160 km and greatest distance (**apogee**) of 16,000 km. Earth is located at a focus of the elliptical orbit. See the figure. (Data from *Space Mathematics*, Kastner, B., NASA.)

Find the equation of the ellipse. (*Hint*: Use the fact that $c^2 = a^2 - b^2$ here.)




Not to scale

Extending Skills Solve each problem. See **FIGURE 16** and use the fact that $c^2 = a^2 - b^2$, where $a^2 > b^2$. Round answers to the nearest tenth. (Data from *Astronomy!*, Kaler, J. B., Addison-Wesley.)

61. The orbit of Mars is an ellipse with the sun at one focus. For x and y in millions of miles, the equation of the orbit is

$$\frac{x^2}{141.7^2} + \frac{y^2}{141.1^2} = 1.$$

- (a) Find the greatest distance (apogee) from Mars to the sun.
 (b) Find the least distance (perigee) from Mars to the sun.



$c^2 = a^2 - b^2$ Not to scale

62. The orbit of Venus around the sun (one of the foci) is an ellipse with equation

$$\frac{x^2}{5013} + \frac{y^2}{4970} = 1,$$

where x and y are measured in millions of miles.

- (a) Find the greatest distance (apogee) between Venus and the sun.
 (b) Find the least distance (perigee) between Venus and the sun.

10.3 Hyperbolas and Functions Defined by Radicals

OBJECTIVES

- 1 Recognize the equation of a hyperbola.
- 2 Graph hyperbolas using asymptotes.
- 3 Identify conic sections using their equations.
- 4 Graph generalized square root functions.

VOCABULARY

- hyperbola
- transverse axis
- fundamental rectangle
- asymptotes (of a hyperbola)
- generalized square root function



OBJECTIVE 1 Recognize the equation of a hyperbola.

A **hyperbola** is the set of all points in a plane such that the absolute value of the *difference* of the distances from two fixed points (the *foci*) is constant. The graph of a hyperbola has two parts, or *branches*, and two intercepts (or *vertices*) that lie on its axis, called the **transverse axis**.

The hyperbola in **FIGURE 23** has a horizontal transverse axis, with foci $(c, 0)$ and $(-c, 0)$ and x -intercepts $(a, 0)$ and $(-a, 0)$. (A hyperbola with vertical transverse axis would have its intercepts on the y -axis.)

A hyperbola centered at the origin has one of the following equations. (It is shown in more advanced courses that for a hyperbola, $c^2 = a^2 + b^2$.)




FIGURE 23

Equations of Hyperbolas

A hyperbola with x -intercepts $(a, 0)$ and $(-a, 0)$ has an equation that can be written in the form

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1. \quad \text{Transverse axis on } x\text{-axis}$$

A hyperbola with y -intercepts $(0, b)$ and $(0, -b)$ has an equation that can be written in the form

$$\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1. \quad \text{Transverse axis on } y\text{-axis}$$

If we were to throw two stones into a pond, the ensuing concentric ripples would be shaped like a hyperbola. A cross-section of the cooling towers for a nuclear power plant is hyperbolic, as shown in the photo.

OBJECTIVE 2 Graph hyperbolas using asymptotes.

The two branches of the graph of a hyperbola approach a pair of intersecting straight lines, which are its *asymptotes*. (See **FIGURES 24** and **25** on the next page.) The asymptotes are useful for sketching the graph of the hyperbola.

Asymptotes of Hyperbolas

The extended diagonals of a rectangle, called the **fundamental rectangle**, with vertices (corners) at the points (a, b) , $(-a, b)$, $(-a, -b)$, and $(a, -b)$ are the **asymptotes** of the hyperbolas

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \quad \text{and} \quad \frac{y^2}{b^2} - \frac{x^2}{a^2} = 1.$$

Using previous methods, we could show that the equations of these asymptotes are

$$y = \frac{b}{a}x \quad \text{and} \quad y = -\frac{b}{a}x. \quad \text{Equations of the asymptotes of a hyperbola}$$

Graphing a Hyperbola

Step 1 Find and locate the intercepts.

- At $(a, 0)$ and $(-a, 0)$ if the x^2 -term has a positive coefficient
- At $(0, b)$ and $(0, -b)$ if the y^2 -term has a positive coefficient

Step 2 Find the fundamental rectangle. Its vertices will be located at the points (a, b) , $(-a, b)$, $(-a, -b)$, and $(a, -b)$.

Step 3 Sketch the asymptotes. The extended diagonals of the fundamental rectangle are the asymptotes of the hyperbola. They have equations $y = \pm \frac{b}{a}x$.

Step 4 Draw the graph. Sketch each branch of the hyperbola through an intercept, approaching (but not touching) the asymptotes.

NOW TRY EXERCISE 1

Graph $\frac{x^2}{25} - \frac{y^2}{9} = 1$.

EXAMPLE 1 Graphing a Horizontal Hyperbola


Graph $\frac{x^2}{16} - \frac{y^2}{25} = 1$.

Step 1 Here $a = 4$ and $b = 5$. The x -intercepts are $(4, 0)$ and $(-4, 0)$.

Step 2 The vertices of the fundamental rectangle are the four points

$$\begin{array}{cccc} (a, b) & (-a, b) & (-a, -b) & (a, -b) \\ \downarrow \downarrow & \downarrow \downarrow & \downarrow \downarrow & \downarrow \downarrow \\ (4, 5), & (-4, 5), & (-4, -5), & \text{and } (4, -5). \end{array}$$

Steps 3 and 4 The equations of the asymptotes are $y = \pm \frac{b}{a}x$, or $y = \pm \frac{5}{4}x$. The hyperbola approaches these lines as x and y get larger in absolute value. See **FIGURE 24**.




When graphing a hyperbola, the fundamental rectangle and the asymptotes are not part of the actual graph. They provide help in drawing a more accurate graph.

Be sure that the branches do not touch the asymptotes.

FIGURE 24

NOW TRY ANSWER



NOW TRY
EXERCISE 2

Graph $\frac{y^2}{9} - \frac{x^2}{16} = 1$.

EXAMPLE 2 Graphing a Vertical Hyperbola

Graph $\frac{y^2}{49} - \frac{x^2}{16} = 1$.

This hyperbola has y -intercepts $(0, 7)$ and $(0, -7)$. The asymptotes are the extended diagonals of the fundamental rectangle with vertices at

$$(4, 7), (-4, 7), (-4, -7), \text{ and } (4, -7).$$

Their equations are $y = \pm \frac{7}{4}x$. See FIGURE 25.




FIGURE 25

NOW TRY

OBJECTIVE 3 Identify conic sections using their equations.

Rewriting a second-degree equation in one of the forms given for ellipses, hyperbolas, circles, or parabolas makes it possible to identify the graph of the equation.

NOW TRY
EXERCISE 3

Identify the graph of each equation.

- (a) $y^2 - 10 = -x^2$
 (b) $y - 2x^2 = 8$
 (c) $3x^2 + y^2 = 4$

EXAMPLE 3 Identifying the Graphs of Equations

Identify the graph of each equation.

(a) $9x^2 = 108 + 12y^2$

Both variables are squared, so the graph is either an ellipse or a hyperbola. (This situation also occurs for a circle, which is a special case of an ellipse.) To see which conic section it is, rewrite the equation so that the x^2 - and y^2 -terms are on one side of the equation and 1 is on the other side.

$$9x^2 - 12y^2 = 108 \quad \text{Subtract } 12y^2.$$

$$\frac{x^2}{12} - \frac{y^2}{9} = 1 \quad \text{Divide by } 108.$$

The subtraction symbol indicates that the graph of this equation is a hyperbola.

(b) $x^2 = y - 3$

Only one of the two variables, x , is squared, so this is the vertical parabola $y = x^2 + 3$.


(c) $x^2 = 9 - y^2$

Write the variable terms on the same side of the equation.

$$x^2 + y^2 = 9 \quad \text{Add } y^2.$$

The graph of this equation is a circle with center at the origin and radius 3.







NOW TRY ANSWERS



3. (a) circle (b) parabola
 (c) ellipse

NOW TRY

▼ SUMMARY OF CONIC SECTIONS

Equation	Graph	Description	Identification
$y = ax^2 + bx + c$ or $y = a(x - h)^2 + k$	 <p style="text-align: center;">Parabola</p>	It opens up if $a > 0$, down if $a < 0$. The vertex is (h, k) .	It has an x^2 -term. y is not squared.
$x = ay^2 + by + c$ or $x = a(y - k)^2 + h$	 <p style="text-align: center;">Parabola</p>	It opens to the right if $a > 0$, to the left if $a < 0$. The vertex is (h, k) .	It has a y^2 -term. x is not squared.
$(x - h)^2 + (y - k)^2 = r^2$	 <p style="text-align: center;">Circle</p>	The center is (h, k) , and the radius is r .	x^2 - and y^2 -terms have the same positive coefficient.
$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$	 <p style="text-align: center;">Ellipse</p>	The x -intercepts are $(a, 0)$ and $(-a, 0)$. The y -intercepts are $(0, b)$ and $(0, -b)$.	x^2 - and y^2 -terms have different positive coefficients.
$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$	 <p style="text-align: center;">Hyperbola</p>	The x -intercepts are $(a, 0)$ and $(-a, 0)$. The asymptotes are found from (a, b) , $(a, -b)$, $(-a, -b)$, and $(-a, b)$.	x^2 has a positive coefficient. y^2 has a negative coefficient.
$\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$	 <p style="text-align: center;">Hyperbola</p>	The y -intercepts are $(0, b)$ and $(0, -b)$. The asymptotes are found from (a, b) , $(a, -b)$, $(-a, -b)$, and $(-a, b)$.	y^2 has a positive coefficient. x^2 has a negative coefficient.

OBJECTIVE 4 Graph generalized square root functions.

Because they do not satisfy the conditions of the vertical line test, the graphs of horizontal parabolas and all circles, ellipses, and hyperbolas with horizontal or vertical axes do not satisfy the conditions of a function. However, by considering only a part of the graph of each of these, we have the graph of a function, as seen in **FIGURE 26**.




FIGURE 26

In parts (a)–(d) of **FIGURE 26**, the top portion of a conic section is shown (parabola, circle, ellipse, and hyperbola, respectively). In part (e), the top two portions of a hyperbola are shown. In each case, the graph is that of a function because the graph satisfies the conditions of the vertical line test.

Earlier in this chapter, we worked with the square root function $f(x) = \sqrt{x}$. To find equations for the types of graphs shown in **FIGURE 26**, we extend its definition.

Generalized Square Root Function

For an algebraic expression in x defined by u , where $u \geq 0$, a function of the form

$$f(x) = \sqrt{u}$$

is a **generalized square root function**.

NOW TRY EXERCISE 4

Graph $f(x) = \sqrt{64 - x^2}$.
Give the domain and range.

EXAMPLE 4 Graphing a Semicircle

Graph $f(x) = \sqrt{25 - x^2}$. Give the domain and range.

$$f(x) = \sqrt{25 - x^2} \quad \text{Given function}$$

$$y = \sqrt{25 - x^2} \quad \text{Replace } f(x) \text{ with } y.$$

$$y^2 = (\sqrt{25 - x^2})^2 \quad \text{Square each side.}$$

$$y^2 = 25 - x^2 \quad (\sqrt{a})^2 = a$$

$$x^2 + y^2 = 25 \quad \text{Add } x^2.$$

This is the equation of a circle with center at $(0, 0)$ and radius 5.

Because $f(x)$, or y , represents a principal square root in the original equation, $f(x)$ must be nonnegative. This restricts the graph to the upper half of the circle.

See **FIGURE 27**. Use the graph and the vertical line test to verify that it is indeed a function. The domain is $[-5, 5]$, and the range is $[0, 5]$.





FIGURE 27


NOW TRY ANSWER

domain: $[-8, 8]$; range: $[0, 8]$

**NOW TRY
EXERCISE 5**

Graph $\frac{y}{4} = -\sqrt{1 - \frac{x^2}{9}}$.

Give the domain and range.

NOW TRY ANSWER

 domain: $[-3, 3]$; range: $[-4, 0]$
EXAMPLE 5 Graphing a Portion of an Ellipse

 Graph $\frac{y}{6} = -\sqrt{1 - \frac{x^2}{16}}$. Give the domain and range.

$$\frac{y}{6} = -\sqrt{1 - \frac{x^2}{16}} \quad \text{Given equation}$$


$$\left(\frac{y}{6}\right)^2 = \left(-\sqrt{1 - \frac{x^2}{16}}\right)^2 \quad \text{Square each side.}$$

$$\frac{y^2}{36} = 1 - \frac{x^2}{16} \quad \text{Apply the exponents.}$$

$$\frac{x^2}{16} + \frac{y^2}{36} = 1 \quad \text{Add } \frac{x^2}{16}.$$

This is the equation of an ellipse with center at $(0, 0)$. The x -intercepts are $(4, 0)$ and $(-4, 0)$. The y -intercepts are $(0, 6)$ and $(0, -6)$.

Because $\frac{y}{6}$ equals a negative square root in the original equation, y must be nonpositive, restricting the graph to the lower half of the ellipse.


FIGURE 28

See **FIGURE 28**. The graph is that of a function. The domain is $[-4, 4]$, and the range is $[-6, 0]$.

NOW TRY



10.3 Exercises

 FOR
EXTRA
HELP



Video solutions for select problems available in MyLab Math

Concept Check Based on the discussions of ellipses in the previous section and of hyperbolas in this section, match each equation with its graph.

1. $\frac{x^2}{25} + \frac{y^2}{9} = 1$



2. $\frac{x^2}{9} + \frac{y^2}{25} = 1$



3. $\frac{x^2}{9} - \frac{y^2}{25} = 1$

4. $\frac{x^2}{25} - \frac{y^2}{9} = 1$

5. **Concept Check** A student incorrectly described the graph of the equation

$$\frac{x^2}{9} - \frac{y^2}{16} = 1$$

as a vertical hyperbola with y -intercepts $(0, -4)$ and $(0, 4)$. **WHAT WENT WRONG?** Give the correct description of the graph.

6. **Concept Check** A student incorrectly described the graph of the function

$$f(x) = \sqrt{49 - x^2}$$

as a circle centered at the origin with radius 7. **WHAT WENT WRONG?** Give the correct description of the graph.

Graph each hyperbola. See Examples 1 and 2.

7. $\frac{x^2}{16} - \frac{y^2}{9} = 1$

8. $\frac{x^2}{25} - \frac{y^2}{9} = 1$

9. $\frac{y^2}{4} - \frac{x^2}{25} = 1$

10. $\frac{y^2}{9} - \frac{x^2}{4} = 1$

11. $\frac{x^2}{25} - \frac{y^2}{36} = 1$

12. $\frac{x^2}{49} - \frac{y^2}{16} = 1$

13. $\frac{y^2}{9} - \frac{x^2}{9} = 1$

14. $\frac{y^2}{16} - \frac{x^2}{16} = 1$

Identify the graph of each equation as a parabola, circle, ellipse, or hyperbola, and then sketch the graph. See Example 3.

15. $x^2 - y^2 = 16$

16. $x^2 + y^2 = 16$

17. $4x^2 + y^2 = 16$

18. $9x^2 = 144 + 16y^2$

19. $y^2 = 36 - x^2$

20. $9x^2 + 25y^2 = 225$

21. $x^2 + 9y^2 = 9$

22. $x^2 - 2y = 0$

23. $x^2 = 4y - 8$

24. $y^2 = 4 + x^2$

Graph each generalized square root function. Give the domain and range. See Examples 4 and 5.

25. $f(x) = \sqrt{16 - x^2}$

26. $f(x) = \sqrt{9 - x^2}$

27. $f(x) = -\sqrt{36 - x^2}$

28. $f(x) = -\sqrt{25 - x^2}$

29. $\frac{y}{3} = \sqrt{1 + \frac{x^2}{9}}$

30. $\frac{y}{2} = \sqrt{1 + \frac{x^2}{4}}$

31. $y = -2\sqrt{1 - \frac{x^2}{9}}$

32. $y = -3\sqrt{1 - \frac{x^2}{25}}$


Extending Skills Solve each problem.

33. Two buildings in a sports complex are shaped and positioned like a portion of the branches of the hyperbola with equation


$$400x^2 - 625y^2 = 250,000,$$

where x and y are in meters.

- (a) How far apart are the buildings at their closest point?
 (b) Find the distance d in the figure to the nearest tenth of a meter.



34. Using LORAN, a location-finding system, a radio transmitter at M sends out a series of pulses. When each pulse is received at transmitter S , it then sends out a pulse. A ship at P receives pulses from both M and S . A receiver on the ship measures the difference in the arrival times of the pulses. A special map gives hyperbolas that correspond to the differences in arrival times (which give the distances d_1 and d_2 in the figure). The ship can then be located as lying on a branch of a particular hyperbola.



Suppose $d_1 = 80$ mi and $d_2 = 30$ mi, and the distance between transmitters M and S is 100 mi. Use the definition to find an equation of the hyperbola on which the ship is located.

Extending Skills In rugby, after a try (similar to a touchdown in American football) the scoring team attempts a kick for extra points. The ball must be kicked from directly behind the point where the try was scored. The kicker can choose the distance but cannot move the ball sideways. It can be shown that the kicker's best choice is on the hyperbola with equation

$$\frac{x^2}{g^2} - \frac{y^2}{g^2} = 1,$$



where $2g$ is the distance between the goal posts. Since the hyperbola approaches its asymptotes, it is easier for the kicker to estimate points on the asymptotes instead of on the hyperbola. (Data from Isaksen, Daniel C., "How to Kick a Field Goal." The College Mathematics Journal.)

35. What are the asymptotes of this hyperbola?
36. Why is it relatively easy to estimate the asymptotes?

RELATING CONCEPTS For Individual or Group Work (Exercises 37–40)

We have seen that the center of an ellipse may be shifted away from the origin. The same process applies to hyperbolas. For example, the hyperbola


$$\frac{(x + 5)^2}{4} - \frac{(y - 2)^2}{9} = 1,$$

has the same graph as

$$\frac{x^2}{4} - \frac{y^2}{9} = 1,$$

but it is centered at $(-5, 2)$, as shown at the right.

Graph each hyperbola with center shifted away from the origin.



$$\frac{(x + 5)^2}{4} - \frac{(y - 2)^2}{9} = 1$$

37. $\frac{(x - 2)^2}{4} - \frac{(y + 1)^2}{9} = 1$

38. $\frac{(x + 3)^2}{16} - \frac{(y - 2)^2}{25} = 1$

39. $\frac{y^2}{36} - \frac{(x - 2)^2}{49} = 1$

40. $\frac{(y - 5)^2}{9} - \frac{x^2}{25} = 1$

10.4 Nonlinear Systems of Equations

OBJECTIVES

- 1 Solve a nonlinear system using substitution.
- 2 Solve a nonlinear system with two second-degree equations using elimination.
- 3 Solve a nonlinear system that requires a combination of methods.

An equation in which some terms have more than one variable or a variable of degree 2 or greater is called a **nonlinear equation**. A **nonlinear system of equations** includes at least one nonlinear equation.

When solving a nonlinear system, it helps to visualize the types of graphs of the equations of the system to determine the possible number of points of intersection. For example, if a system includes two equations where the graph of one is a circle and the graph of the other is a line, then there may be zero, one, or two points of intersection, as illustrated in **FIGURE 29**.




FIGURE 29

VOCABULARY

- nonlinear equation
- nonlinear system of equations

OBJECTIVE 1 Solve a nonlinear system using substitution.

We can usually solve a nonlinear system using the substitution method if one equation is linear.

EXAMPLE 1 Solving a Nonlinear System Using Substitution

Solve the system.

$$x^2 + y^2 = 9 \quad (1)$$

$$2x - y = 3 \quad (2)$$

The graph of equation (1) is a circle and the graph of equation (2) is a line, so the graphs could intersect in zero, one, or two points, as in **FIGURE 29**. We begin by solving the linear equation (2) for one of the two variables.

$$2x - y = 3 \quad (2)$$

$$y = 2x - 3 \quad \text{Solve for } y. \quad (3)$$

Then we substitute $2x - 3$ for y in the nonlinear equation (1).

$$x^2 + y^2 = 9 \quad (1)$$

$$x^2 + (2x - 3)^2 = 9 \quad \text{Let } y = 2x - 3.$$

$$x^2 + 4x^2 - 12x + 9 = 9 \quad \text{Square } 2x - 3.$$

$$5x^2 - 12x = 0 \quad \text{Combine like terms. Subtract 9.}$$

$$x(5x - 12) = 0 \quad \text{Factor. The GCF is } x.$$

$$x = 0 \quad \text{or} \quad 5x - 12 = 0 \quad \text{Zero-factor property}$$

Set both factors equal to 0.

$$x = \frac{12}{5} \quad \text{Solve for } x.$$

NOW TRY
EXERCISE 1

Solve the system.

$$4x^2 + y^2 = 36$$

$$x - y = 3$$

Let $x = 0$ in equation (3), $y = 2x - 3$, to obtain $y = -3$. Then let $x = \frac{12}{5}$ in equation (3).

$$y = 2\left(\frac{12}{5}\right) - 3 \quad \text{Let } x = \frac{12}{5}.$$

$$y = \frac{24}{5} - \frac{15}{5} \quad \text{Multiply; } 3 = \frac{15}{5}$$

$$y = \frac{9}{5} \quad \frac{a}{c} - \frac{b}{c} = \frac{a-b}{c}$$

The solution set of the system is $\left\{(0, -3), \left(\frac{12}{5}, \frac{9}{5}\right)\right\}$. The graph in **FIGURE 30** confirms the two points of intersection.




FIGURE 30

NOW TRY

EXAMPLE 2 Solving a Nonlinear System Using Substitution

Solve the system.

$$6x - y = 5 \quad (1)$$

$$xy = 4 \quad (2)$$

The graph of (1) is a line. It can be shown by plotting points that the graph of (2) is a hyperbola. Visualizing a line and a hyperbola indicates that there may be zero, one, or two points of intersection.

Since neither equation has a squared term, we can solve either equation for one of the variables and then substitute the result into the other equation. Solving $xy = 4$ for x gives $x = \frac{4}{y}$. We substitute $\frac{4}{y}$ for x in equation (1).

$$6x - y = 5 \quad (1)$$

$$6\left(\frac{4}{y}\right) - y = 5 \quad \text{Let } x = \frac{4}{y}.$$

$$\frac{24}{y} - y = 5 \quad \text{Multiply.}$$

$$24 - y^2 = 5y \quad \text{Multiply by } y, y \neq 0.$$

$$y^2 + 5y - 24 = 0 \quad \text{Standard form}$$

$$(y - 3)(y + 8) = 0 \quad \text{Factor.}$$

$$y - 3 = 0 \quad \text{or} \quad y + 8 = 0 \quad \text{Zero-factor property}$$

$$y = 3 \quad \text{or} \quad y = -8 \quad \text{Solve each equation.}$$

NOW TRY ANSWER

1. $\left\{(3, 0), \left(-\frac{9}{5}, -\frac{24}{5}\right)\right\}$

NOW TRY
EXERCISE 2

Solve the system.

$$xy = 2$$

$$x - 3y = 1$$

We substitute these values of y into $x = \frac{4}{y}$ to obtain the corresponding values of x .

$$\text{If } y = 3, \quad \text{then } x = \frac{4}{3}.$$

$$\text{If } y = -8, \quad \text{then } x = -\frac{1}{2}.$$

The solution set of the system is

$$\left\{ \left(\frac{4}{3}, 3 \right), \left(-\frac{1}{2}, -8 \right) \right\}.$$

Write the x -coordinates first in the ordered-pair solutions.




FIGURE 31


NOW TRY

NOTE In **Example 2**, we could solve the *linear* equation for one of its variables and substitute for this variable in the *nonlinear* equation. There is often more than one way to solve a nonlinear system of equations.

OBJECTIVE 2 Solve a nonlinear system with two second-degree equations using elimination.

If a system consists of two second-degree equations, then there may be zero, one, two, three, or four solutions. **FIGURE 32** shows a case where a system consisting of a circle and a parabola has four solutions, all made up of ordered pairs of real numbers.

The elimination method is often used when both equations of a nonlinear system are second degree.



This system has four solutions because there are four points of intersection.

FIGURE 32

EXAMPLE 3 Solving a Nonlinear System Using Elimination

Solve the system.

$$x^2 + y^2 = 9 \quad (1)$$

$$2x^2 - y^2 = -6 \quad (2)$$

The graph of (1) is a circle, while the graph of (2) is a hyperbola. By analyzing the possibilities, we conclude that there may be zero, one, two, three, or four points of intersection. Adding the two equations will eliminate y .

$$x^2 + y^2 = 9 \quad (1)$$

$$2x^2 - y^2 = -6 \quad (2)$$

$$3x^2 = 3 \quad \text{Add.}$$

$$x^2 = 1 \quad \text{Divide by 3.}$$

$$x = 1 \quad \text{or} \quad x = -1 \quad \text{Square root property}$$

NOW TRY ANSWER

2. $\{(-2, -1), (3, \frac{2}{3})\}$

Each value of x gives corresponding values for y when substituted into one of the original equations. Using equation (1) is easier because the coefficients of the x^2 - and y^2 -terms are 1.

NOW TRY
EXERCISE 3

Solve the system.

$$\begin{aligned}x^2 + y^2 &= 16 \\4x^2 + 13y^2 &= 100\end{aligned}$$




FIGURE 33

$$\begin{aligned}x^2 + y^2 &= 9 && (1) \\1^2 + y^2 &= 9 && \text{Let } x = 1. \\y^2 &= 8 \\y &= \sqrt{8} \quad \text{or} \quad y = -\sqrt{8} \\y &= 2\sqrt{2} \quad \text{or} \quad y = -2\sqrt{2}\end{aligned}$$

$$\begin{aligned}x^2 + y^2 &= 9 && (1) \\(-1)^2 + y^2 &= 9 && \text{Let } x = -1. \\y^2 &= 8 \\y &= \sqrt{8} \quad \text{or} \quad y = -\sqrt{8} \\y &= 2\sqrt{2} \quad \text{or} \quad y = -2\sqrt{2}\end{aligned}$$

The solution set is

$$\{(1, 2\sqrt{2}), (1, -2\sqrt{2}), (-1, 2\sqrt{2}), (-1, -2\sqrt{2})\}.$$

FIGURE 33 shows the four points of intersection.

NOW TRY

OBJECTIVE 3 Solve a nonlinear system that requires a combination of methods.

EXAMPLE 4 Solving a Nonlinear System Using a Combination of Methods

Solve the system.

$$x^2 + 2xy - y^2 = 7 \quad (1)$$

$$x^2 - y^2 = 3 \quad (2)$$

While we have not graphed equations like (1), its graph is a hyperbola. The graph of (2) is also a hyperbola. Two hyperbolas may have zero, one, two, three, or four points of intersection. We use the elimination method here in combination with the substitution method.

$$\begin{aligned}x^2 + 2xy - y^2 &= 7 && (1) \\-x^2 &+ y^2 = -3 && \text{Multiply (2) by } -1. \\ \hline 2xy &= 4 && \text{Add.}\end{aligned}$$

The x^2 - and y^2 -terms were eliminated.

Next, we solve $2xy = 4$ for one of the variables. We choose y .

$$\begin{aligned}2xy &= 4 \\y &= \frac{2}{x} && \text{Divide by } 2x. \quad (3)\end{aligned}$$

Now, we substitute $y = \frac{2}{x}$ into one of the original equations.

$$\begin{aligned}x^2 - y^2 &= 3 && \text{The substitution is easier in (2).} \\x^2 - \left(\frac{2}{x}\right)^2 &= 3 && \text{Let } y = \frac{2}{x}. \\x^2 - \frac{4}{x^2} &= 3 && \text{Square } \frac{2}{x}. \\x^4 - 4 &= 3x^2 && \text{Multiply by } x^2, x \neq 0. \\x^4 - 3x^2 - 4 &= 0 && \text{Subtract } 3x^2. \\(x^2 - 4)(x^2 + 1) &= 0 && \text{Factor.}\end{aligned}$$

$$x^2 - 4 = 0 \quad \text{or} \quad x^2 + 1 = 0 \quad \text{Zero-factor property}$$

$$x^2 = 4 \quad \text{or} \quad x^2 = -1 \quad \text{Solve each equation.}$$

$$x = 2 \quad \text{or} \quad x = -2 \quad \text{or} \quad x = i \quad \text{or} \quad x = -i$$

NOW TRY ANSWER

3. $\{(2\sqrt{3}, 2), (2\sqrt{3}, -2), (-2\sqrt{3}, 2), (-2\sqrt{3}, -2)\}$

NOW TRY EXERCISE 4

Solve the system.

$$x^2 + 3xy - y^2 = 23$$

$$x^2 - y^2 = 5$$

Substituting these four values of x into $y = \frac{2}{x}$ (equation (3)) gives the corresponding values for y .

If $x = 2$, then $y = \frac{2}{2} = 1$.

If $x = -2$, then $y = \frac{2}{-2} = -1$.

If $x = i$, then $y = \frac{2}{i} = \frac{2}{i} \cdot \frac{-i}{-i} = -2i$.

If $x = -i$, then $y = \frac{2}{-i} = \frac{2}{-i} \cdot \frac{i}{i} = 2i$.

Multiply by the complex conjugate of the denominator.
 $i(-i) = 1$

If we substitute the x -values we found into equation (1) or (2) instead of into equation (3), we get extraneous solutions. **It is always wise to check all solutions in both of the given equations.** We show a check for the ordered pair $(i, -2i)$.

CHECK Let $x = i$ and $y = -2i$ in both equations (1) and (2).

$x^2 + 2xy - y^2 = 7$ (1)		$x^2 - y^2 = 3$ (2)
$i^2 + 2(i)(-2i) - (-2i)^2 \stackrel{?}{=} 7$		$i^2 - (-2i)^2 \stackrel{?}{=} 3$
$i^2 - 4i^2 - 4i^2 \stackrel{?}{=} 7$		$i^2 - 4i^2 \stackrel{?}{=} 3$
$-1 - 4(-1) - 4(-1) \stackrel{?}{=} 7$	$i^2 = -1$	$-1 - 4(-1) \stackrel{?}{=} 3$
$7 = 7$ ✓ True		$3 = 3$ ✓ True




FIGURE 34

The other ordered pairs would be checked similarly. There are four ordered pairs in the solution set, two with real values and two with pure imaginary values. The solution set is

$$\{(2, 1), (-2, -1), (i, -2i), (-i, 2i)\}.$$

The graph of the system, shown in FIGURE 34, shows only the two real intersection points because the graph is in the real number plane. In general, if solutions contain nonreal complex numbers as components, they do not appear on the graph.

NOW TRY

NOTE It is not essential to visualize the number of points of intersection of the graphs in order to solve a nonlinear system. Sometimes we are unfamiliar with the graphs or, as in **Example 4**, there are nonreal complex solutions that do not appear as points of intersection in the real plane. Visualizing the graphs is only an aid to solving these systems.

NOW TRY ANSWER

4. $\{(3, 2), (-3, -2), (2i, -3i), (-2i, 3i)\}$


10.4 Exercises

FOR EXTRA HELP

MyLab Math

Video solutions for select problems available in MyLab Math

Concept Check Each sketch represents the graphs of a pair of equations in a system. How many ordered pairs of real numbers are in each solution set?



Concept Check Suppose that a nonlinear system is composed of equations whose graphs are those described, and the number of points of intersection of the two graphs is as given. Make a sketch satisfying these conditions. (There may be more than one way to do this.)

- | | |
|--------------------------------------------|--------------------------------------------|
| 5. A line and a circle; no points | 6. A line and a circle; one point |
| 7. A line and a hyperbola; one point | 8. A line and an ellipse; no points |
| 9. A circle and an ellipse; four points | 10. A parabola and an ellipse; one point |
| 11. A parabola and an ellipse; four points | 12. A parabola and a hyperbola; two points |

Solve each system using the substitution method. See Examples 1 and 2.

- | | |
|------------------------------------------|-------------------------------------------|
| 13. $y = 4x^2 - x$
$y = x$ | 14. $y = x^2 + 6x$
$3y = 12x$ |
| 15. $y = x^2 + 6x + 9$
$x + y = 3$ | 16. $y = x^2 + 8x + 16$
$x - y = -4$ |
| 17. $x^2 + y^2 = 2$
$2x + y = 1$ | 18. $2x^2 + 4y^2 = 4$
$x = 4y$ |
| 19. $xy = 4$
$3x + 2y = -10$ | 20. $xy = -5$
$2x + y = 3$ |
| 21. $xy = -3$
$x + y = -2$ | 22. $xy = 12$
$x + y = 8$ |
| 23. $y = 3x^2 + 6x$
$y = x^2 - x - 6$ | 24. $y = 2x^2 + 1$
$y = 5x^2 + 2x - 7$ |
| 25. $2x^2 - y^2 = 6$
$y = x^2 - 3$ | 26. $x^2 + y^2 = 4$
$y = x^2 - 2$ |
| 27. $x^2 - xy + y^2 = 0$
$x - 2y = 1$ | 28. $x^2 - 3x + y^2 = 4$
$2x - y = 3$ |

Solve each system using the elimination method or a combination of the elimination and substitution methods. See Examples 3 and 4.

- | | |
|-----------------------------------------------|--------------------------------------------|
| 29. $3x^2 + 2y^2 = 12$
$x^2 + 2y^2 = 4$ | 30. $x^2 + 6y^2 = 9$
$4x^2 + 3y^2 = 36$ |
| 31. $5x^2 - 2y^2 = -13$
$3x^2 + 4y^2 = 39$ | 32. $x^2 + y^2 = 41$
$x^2 - y^2 = 9$ |
| 33. $2x^2 + 3y^2 = 6$
$x^2 + 3y^2 = 3$ | 34. $3x^2 + y^2 = 15$
$x^2 + y^2 = 5$ |
| 35. $2x^2 + y^2 = 28$
$4x^2 - 5y^2 = 28$ | 36. $x^2 + 3y^2 = 40$
$4x^2 - y^2 = 4$ |
| 37. $xy = 6$
$3x^2 - y^2 = 12$ | 38. $xy = 5$
$2y^2 - x^2 = 5$ |

$$39. \begin{aligned} 2x^2 &= 8 - 2y^2 \\ 3x^2 &= 24 - 4y^2 \end{aligned}$$

$$41. \begin{aligned} x^2 + xy + y^2 &= 15 \\ x^2 + y^2 &= 10 \end{aligned}$$

$$43. \begin{aligned} x^2 + xy - y^2 &= 11 \\ x^2 - y^2 &= 8 \end{aligned}$$

$$45. \begin{aligned} 3x^2 + 2xy - 3y^2 &= 5 \\ -x^2 - 3xy + y^2 &= 3 \end{aligned}$$

$$40. \begin{aligned} 5x^2 &= 20 - 5y^2 \\ 2y^2 &= 2 - x^2 \end{aligned}$$


$$42. \begin{aligned} 2x^2 + 3xy + 2y^2 &= 21 \\ x^2 + y^2 &= 6 \end{aligned}$$

$$44. \begin{aligned} x^2 + xy - y^2 &= 29 \\ x^2 - y^2 &= 24 \end{aligned}$$


$$46. \begin{aligned} -2x^2 + 7xy - 3y^2 &= 4 \\ 2x^2 - 3xy + 3y^2 &= 4 \end{aligned}$$

Extending Skills Solve each problem using a nonlinear system.

47. The area of a rectangular rug is 84 ft^2 and its perimeter is 38 ft . Find the length and width of the rug.



48. Find the length and width of a rectangular room whose perimeter is 50 m and whose area is 100 m^2 .



49. A company has found that the price p (in dollars) of its scientific calculator is related to the supply x (in thousands) by the equation

$$px = 16.$$

The price is related to the demand x (in thousands) for the calculator by the equation

$$p = 10x + 12.$$

The **equilibrium price** is the value of p where demand equals supply. Find the equilibrium price and the supply/demand at that price. (*Hint:* Demand, price, and supply must all be positive.)

50. A company has determined that the cost y to make x (thousand) computer tablets is

$$y = 4x^2 + 36x + 20,$$

and that the revenue y from the sale of x (thousand) tablets is

$$36x^2 - 3y = 0.$$

Find the **break-even point**, where cost equals revenue.

10.5

Second-Degree Inequalities and Systems of Inequalities

OBJECTIVES

- 1 Graph second-degree inequalities.
- 2 Graph the solution set of a system of inequalities.

OBJECTIVE 1 Graph second-degree inequalities.

A **second-degree inequality** is an inequality with at least one variable of degree 2 and no variable of degree greater than 2.

Examples: $x^2 + y^2 > 9$, $y \leq 2x^2 - 4$, $x \geq y^2$

EXAMPLE 1 Graphing a Second-Degree Inequality

Graph $x^2 + y^2 \leq 36$.

The boundary of the inequality $x^2 + y^2 \leq 36$ is the graph of the equation $x^2 + y^2 = 36$, a circle with radius 6 and center at the origin, as shown in **FIGURE 35**.

The inequality $x^2 + y^2 \leq 36$ includes the points of the boundary (because the symbol \leq includes equality) and either the points “outside” the circle or the points “inside” the circle. To decide which region to shade, we substitute any test point not on the circle.




FIGURE 35

NOW TRY

VOCABULARY

- second-degree inequality
- system of inequalities

NOW TRY EXERCISE 1

Graph $x^2 + y^2 \geq 9$.

$$\begin{aligned}
 x^2 + y^2 &< 36 && \text{We are testing the region.} \\
 0^2 + 0^2 &\stackrel{?}{<} 36 && \text{Use } (0, 0) \text{ as a test point.} \\
 0 &< 36 && \text{True}
 \end{aligned}$$

Because a true statement results, the original inequality includes the points *inside* the circle, the shaded region in **FIGURE 35**, and the boundary.

NOTE Because the substitution is easy, the origin is the test point of choice unless the graph actually passes through $(0, 0)$.

EXAMPLE 2 Graphing a Second-Degree Inequality

Graph $y < -2(x - 4)^2 - 3$.

The boundary, $y = -2(x - 4)^2 - 3$, is a parabola that opens down with vertex $(4, -3)$. We must decide whether to shade the region “inside” or “outside” the parabola.

$$\begin{aligned}
 y &< -2(x - 4)^2 - 3 && \text{Original inequality} \\
 0 &\stackrel{?}{<} -2(0 - 4)^2 - 3 && \text{Use } (0, 0) \text{ as a test point.} \\
 0 &\stackrel{?}{<} -32 - 3 && \text{Simplify.} \\
 0 &< -35 && \text{False}
 \end{aligned}$$

Because the final inequality is a false statement, the points in the region containing $(0, 0)$ do not satisfy the inequality. As a result, the region inside (or below) the parabola is shaded in **FIGURE 36**. The parabola is drawn as a dashed curve since the points of the parabola itself do not satisfy the inequality.




FIGURE 36

CHECK As additional confirmation, select a test point in the shaded region, such as $(4, -7)$, and substitute it into the original inequality.

$$\begin{aligned}
 y &< -2(x - 4)^2 - 3 \\
 -7 &\stackrel{?}{<} -2(4 - 4)^2 - 3 && \text{Test } (4, -7). \\
 -7 &< -3 && \checkmark \quad \text{True}
 \end{aligned}$$

A true statement results, so the correct region is shaded.

NOW TRY

NOW TRY EXERCISE 2

Graph $y \geq -(x + 2)^2 + 1$.

NOW TRY ANSWERS

- 1.
- 2.

NOW TRY
EXERCISE 3

Graph $25x^2 - 16y^2 > 400$.




FIGURE 37

EXAMPLE 3 Graphing a Second-Degree Inequality

Graph $16y^2 \leq 144 + 9x^2$.

$$\begin{aligned} 16y^2 &\leq 144 + 9x^2 \\ 16y^2 - 9x^2 &\leq 144 && \text{Subtract } 9x^2. \\ \frac{y^2}{9} - \frac{x^2}{16} &\leq 1 && \text{Divide by } 144. \end{aligned}$$

This form shows that the boundary is the hyperbola given by

$$\frac{y^2}{9} - \frac{x^2}{16} = 1.$$

Because the graph is a vertical hyperbola, the desired region will be either the region “between” the branches or the regions “above” the top branch and “below” the bottom branch. We choose $(0, 0)$ as a test point.

$$\begin{aligned} 16y^2 &< 144 + 9x^2 \\ 16(0)^2 &\stackrel{?}{<} 144 + 9(0)^2 && \text{Test } (0, 0). \\ 0 &< 144 && \text{True} \end{aligned}$$

Because a true statements results, we shade the region between the branches containing $(0, 0)$. See FIGURE 37.

NOW TRY

OBJECTIVE 2 Graph the solution set of a system of inequalities.

If two or more inequalities are considered at the same time, we have a **system of inequalities**. To find the solution set of the system, we find the intersection of the graphs (solution sets) of the inequalities in the system.

NOW TRY
EXERCISE 4

Graph the solution set of the system.

$$\begin{aligned} 2x + 3y &\geq 6 \\ x - 5y &\geq 5 \end{aligned}$$

EXAMPLE 4 Graphing a System of Two Inequalities

Graph the solution set of the system.

$$\begin{aligned} x - y &\leq 4 \\ x + 2y &< 2 \end{aligned}$$

Both inequalities in the system are linear. We begin by graphing the solution set of $x - y \leq 4$. The inequality includes the points of the boundary line $x - y = 4$ because the symbol \leq includes equality. The test point $(0, 0)$ leads to a true statement in $x - y < 4$, so we shade the region above the line. See FIGURE 38.

The graph of the solution set of $x + 2y < 2$ does not include the boundary line $x + 2y = 2$. Testing the point $(0, 0)$ leads to a true statement, indicating that we should shade the region below the dashed boundary line. See FIGURE 39.




FIGURE 38




FIGURE 39






FIGURE 40

NOW TRY ANSWERS

3.



4.



The solution set is the intersection of the graphs of the two inequalities. See the overlapping region in FIGURE 40, which includes one boundary line.

NOW TRY

NOW TRY EXERCISE 5

Graph the solution set of the system.

$$\begin{aligned} x^2 + y^2 &\leq 25 \\ x + y &\leq 3 \end{aligned}$$

EXAMPLE 5 Graphing a System of Two Inequalities

Graph the solution set of the system.

$$\begin{aligned} 2x + 3y &> 6 \\ x^2 + y^2 &< 16 \end{aligned}$$

We begin by graphing the solution set of the linear inequality $2x + 3y > 6$. The boundary line is the graph of $2x + 3y = 6$ and is a dashed line because the symbol $>$ does not include equality. The test point $(0, 0)$ leads to a false statement in $2x + 3y > 6$, so we shade the region above the line, as shown in **FIGURE 41**.

The graph of $x^2 + y^2 < 16$ is the region inside of a dashed circle centered at the origin with radius 4. This is shown in **FIGURE 42**.




FIGURE 41

FIGURE 42

FIGURE 43

The graph of the solution set of the system is the intersection of the graphs of the two inequalities. The overlapping region in **FIGURE 43** is the solution set.

CHECK As additional confirmation, select a test point in the shaded region, such as $(3, 1)$, and substitute it into *both* inequalities.

$2x + 3y > 6$	$x^2 + y^2 < 16$
$2(3) + 3(1) \stackrel{?}{>} 6$ Test $(3, 1)$.	$3^2 + 1^2 \stackrel{?}{<} 16$ Test $(3, 1)$.
$9 > 6$ ✓ True	$10 < 16$ ✓ True

True statements result, so the correct region is shaded.

NOW TRY

NOW TRY EXERCISE 6

Graph the solution set of the system.

$$\begin{aligned} 3x + 2y &> 6 \\ y &\geq \frac{1}{2}x - 2 \\ x &\geq 0 \end{aligned}$$

EXAMPLE 6 Graphing a System of Three Inequalities

Graph the solution set of the system.

$$\begin{aligned} x + y &< 1 \\ y &\leq 2x + 3 \\ y &\geq -2 \end{aligned}$$

We graph each linear inequality on the same axes.

- The graph of $x + y < 1$ is the region that lies below the dashed line $x + y = 1$.
- The graph of $y \leq 2x + 3$ is the region that lies below the solid line $y = 2x + 3$.
- The graph of $y \geq -2$ is the region above the solid horizontal line $y = -2$.




FIGURE 44

The test point $(0, 0)$ satisfies all three inequalities. The graph of the system, the intersection of these three graphs, is the triangular region enclosed by the three boundary lines in **FIGURE 44**, including two of its boundary lines.

NOW TRY

NOW TRY ANSWERS


- 5.
- 6.

NOW TRY EXERCISE 7

Graph the solution set of the system.

$$\begin{aligned} \frac{x^2}{4} + \frac{y^2}{16} &\leq 1 \\ y &\leq x^2 - 2 \\ y + 3 &> 0 \end{aligned}$$

NOW TRY ANSWER



EXAMPLE 7 Graphing a System of Three Inequalities

Graph the solution set of the system.

$$\begin{aligned} y &\geq x^2 - 2x + 1 \\ 2x^2 + y^2 &> 4 \\ y &< 4 \end{aligned}$$

We graph each inequality on the same axes.

- The graph of $y = x^2 - 2x + 1$ is a parabola with vertex at $(1, 0)$. Those points inside (or above) the parabola satisfy the condition $y > x^2 - 2x + 1$. Thus, the solution set of $y \geq x^2 - 2x + 1$ includes points on or inside the parabola.
- The graph of the equation $2x^2 + y^2 = 4$ is an ellipse. We draw it as a dashed curve. To satisfy the inequality $2x^2 + y^2 > 4$, a point must lie outside the ellipse.
- The graph of $y < 4$ includes all points below the dashed line $y = 4$.

The graph of the system is the shaded region in **FIGURE 45**, which lies outside the ellipse, inside or on the boundary of the parabola, and below the line $y = 4$.




FIGURE 45

NOW TRY

10.5 Exercises





FOR EXTRA HELP



Video solutions for select problems available in MyLab Math

1. Concept Check Match each inequality in parts (a)–(d) with its graph in choices A–D.

- (a) $y \geq x^2 + 4$ (b) $y \leq x^2 + 4$ (c) $y < x^2 + 4$ (d) $y > x^2 + 4$







2. **Concept Check** Match each system of inequalities in parts (a)–(d) with its graph in choices A–D.

(a) $x > -2$
 $y \leq 2$

(b) $x < -2$
 $y \geq 2$

(c) $x \geq 2$
 $y < -2$

(d) $x \leq 2$
 $y > -2$



Concept Check Write a system of inequalities for the indicated region.

3. Quadrant I, including the x - and y -axes
4. Quadrant IV, not including the x - and y -axes
5. Quadrant III, including the x -axis but not including the y -axis
6. Quadrant II, not including the x -axis but including the y -axis

7. **Concept Check** Which one of the following is a description of the graph of the solution set of the following system?

$$x^2 + y^2 < 25$$

$$y > -2$$

- A. All points outside the circle $x^2 + y^2 = 25$ and above the line $y = -2$
- B. All points outside the circle $x^2 + y^2 = 25$ and below the line $y = -2$
- C. All points inside the circle $x^2 + y^2 = 25$ and above the line $y = -2$
- D. All points inside the circle $x^2 + y^2 = 25$ and below the line $y = -2$

8. **Concept Check** Fill in each blank with the appropriate response. The graph of the solution set of the system

$$y > x^2 + 1$$

$$\frac{x^2}{9} + \frac{y^2}{4} > 1$$

$$y < 5$$

consists of all points _____ the parabola $y = x^2 + 1$, _____ the
(above / below) (inside / outside)

ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$, and _____ the line $y = 5$.
(above / below)

Graph each inequality. See Examples 1–3.

- | | |
|-----------------------------|----------------------------|
| 9. $y \geq x^2 - 2$ | 10. $y > x^2 - 1$ |
| 11. $2y^2 \geq 8 - x^2$ | 12. $y^2 \leq 4 - 2x^2$ |
| 13. $x^2 > 4 - y^2$ | 14. $x^2 \leq 16 - y^2$ |
| 15. $9x^2 > 16y^2 + 144$ | 16. $x^2 \leq 16 + 4y^2$ |
| 17. $9x^2 < 16y^2 - 144$ | 18. $y^2 > 4 + x^2$ |
| 19. $x^2 - 4 \geq -4y^2$ | 20. $4y^2 \leq 36 - 9x^2$ |
| 21. $x \leq -y^2 + 6y - 7$ | 22. $x \geq y^2 - 8y + 14$ |
| 23. $25x^2 \leq 9y^2 + 225$ | 24. $y^2 - 16x^2 \leq 16$ |


Graph each system of inequalities. See Examples 4–7.

- | | | |
|----------------------------------------------------------------------|-------------------------------------------------------------------|------------------------------------------|
| 25. $2x + 5y < 10$
$x - 2y < 4$ | 26. $3x - y > -6$
$4x + 3y > 12$ | 27. $5x - 3y \leq 15$
$4x + y \geq 4$ |
| 28. $4x - 3y \leq 0$
$x + y \leq 5$ | 29. $x \leq 5$
$y \leq 4$ | 30. $x \geq -2$
$y \leq 4$ |
| 31. $x^2 + y^2 > 9$
$y > x^2 - 1$ | 32. $x^2 - y^2 \geq 9$
$\frac{x^2}{16} + \frac{y^2}{9} \leq 1$ | |
| 33. $y > -x^2 - 4$
$y < -x^2 + 3$ | 34. $y \leq -x^2 + 5$
$y \leq x^2 - 3$ | |
| 35. $x + y > 1$
$y \geq 2x - 2$
$y \leq 4$ | 36. $3x - 4y \geq 12$
$x + 3y > 6$
$y \leq 2$ | |
| 37. $x^2 + y^2 \geq 4$
$x + y \leq 5$
$x \geq 0$
$y \geq 0$ | 38. $y^2 - x^2 \geq 4$
$-5 \leq y \leq 5$ | |
| 39. $y \leq -x^2$
$y \geq x - 3$
$y \leq -1$
$x < 1$ | 40. $y < x^2$
$y > -2$
$x + y < 3$
$3x - 2y > -6$ | |

Extending Skills The nonlinear inequality

$$x^2 + y^2 \leq 4, \quad x \geq 0$$

is graphed in the figure. Only the right half of the interior of the circle and its boundary is shaded because of the restriction that x must be nonnegative. Graph each nonlinear inequality with the given restrictions.



- | | |
|---------------------------------------------------|-----------------------------------------------------|
| 41. $x^2 + y^2 > 36, \quad x \geq 0$ | 42. $4x^2 + 25y^2 < 100, \quad y < 0$ |
| 43. $x < y^2 - 3, \quad x < 0$ | 44. $x^2 - y^2 < 4, \quad x < 0$ |
| 45. $4x^2 - y^2 > 16, \quad x < 0$ | 46. $x^2 + y^2 > 4, \quad y < 0$ |
| 47. $x^2 + 4y^2 \geq 1, \quad x \geq 0, y \geq 0$ | 48. $2x^2 - 32y^2 \leq 8, \quad x \leq 0, y \geq 0$ |

Chapter 10 Summary

Key Terms

10.1 absolute value function reciprocal function asymptote square root function greatest integer function step function	10.2 conic section circle center (of a circle) radius ellipse foci (singular <i>focus</i>) center (of an ellipse)	10.3 hyperbola transverse axis fundamental rectangle asymptotes (of a hyperbola) generalized square root function	10.4 nonlinear equation nonlinear system of equations 10.5 second-degree inequality system of inequalities
--------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------

New Symbols

$\lfloor x \rfloor$ greatest integer less than or equal to x

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- Conic sections are
 - graphs of first-degree equations
 - the result of two or more intersecting planes
 - graphs of first-degree inequalities
 - figures that result from the intersection of an infinite cone with a plane.
- A **circle** is the set of all points in a plane
 - such that the absolute value of the difference of the distances from two fixed points is constant
 - that lie a fixed distance from a fixed point
 - the sum of whose distances from two fixed points is constant
 - that make up the graph of any second-degree equation.
- An **ellipse** is the set of all points in a plane
 - such that the absolute value of the difference of the distances from two fixed points is constant
 - that lie a fixed distance from a fixed point
 - the sum of whose distances from two fixed points is constant
 - that make up the graph of any second-degree equation.
- A **hyperbola** is the set of all points in a plane
 - such that the absolute value of the difference of the distances from two fixed points is constant
 - that lie a fixed distance from a fixed point
 - the sum of whose distances from two fixed points is constant
 - that make up the graph of any second-degree equation.
- A **nonlinear equation** is an equation
 - in which some terms have more than one variable or a variable of degree 2 or greater
 - in which the terms have only one variable
 - of degree 1
 - of a linear function.
- A **nonlinear system of equations** is a system
 - with at least one linear equation
 - with two or more inequalities
 - with at least one nonlinear equation
 - with at least two linear equations.

ANSWERS

1. D; *Example:* Parabolas, circles, ellipses, and hyperbolas are conic sections. 2. B; *Example:* The graph of $x^2 + y^2 = 9$ is a circle centered at the origin with radius 3. 3. C; *Example:* The graph of $\frac{x^2}{49} + \frac{y^2}{36} = 1$ is an ellipse centered at the origin with x -intercepts $(7, 0)$ and $(-7, 0)$ and y -intercepts $(0, 6)$ and $(0, -6)$. 4. A; *Example:* The graph of $\frac{x^2}{16} - \frac{y^2}{25} = 1$ is a horizontal parabola centered at the origin with vertices $(4, 0)$ and $(-4, 0)$. 5. A; *Examples:* $y = x^2 + 8x + 16$, $xy = 5$, $2x^2 - y^2 = 6$ 6. C; *Example:* $x^2 + y^2 = 2$
 $2x + y = 1$

Quick Review


CONCEPTS

10.1 Additional Graphs of Functions

Other Elementary Functions

- Absolute value function $f(x) = |x|$
- Reciprocal function $f(x) = \frac{1}{x}$
- Square root function $f(x) = \sqrt{x}$
- Greatest integer function $f(x) = \llbracket x \rrbracket$ (a step function)

Their graphs can be translated, as shown in the first three examples at the right.



10.2 Circles Revisited and Ellipses

Circle

A circle with center (h, k) , and radius $r > 0$ has an equation that can be written in the form

$$(x - h)^2 + (y - k)^2 = r^2 \quad \text{Center-radius form}$$

If its center is $(0, 0)$, then this equation becomes

$$x^2 + y^2 = r^2.$$

For some real numbers c , d , and e , an equation of the form

$$x^2 + y^2 + cx + dy + e = 0 \quad \text{General form}$$

may represent a circle. The x^2 - and y^2 -terms have equal coefficients, here 1.

Ellipse

An ellipse with x -intercepts $(a, 0)$ and $(-a, 0)$ and y -intercepts $(0, b)$ and $(0, -b)$ has an equation that can be written in the form

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

10.3 Hyperbolas and Functions Defined by Radicals

Hyperbola

A hyperbola centered at the origin has an equation that can be written in one of the following forms.

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \quad \text{or} \quad \frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$$

x -intercepts $(a, 0)$ and $(-a, 0)$ y -intercepts $(0, b)$ and $(0, -b)$

The extended diagonals of the fundamental rectangle with vertices at the points (a, b) , $(-a, b)$, $(-a, -b)$, and $(a, -b)$ are the asymptotes of these hyperbolas.


EXAMPLES

Graph $(x + 2)^2 + (y - 3)^2 = 25$.

This equation, which can be written

$$[x - (-2)]^2 + (y - 3)^2 = 5^2,$$

represents a circle with center $(-2, 3)$ and radius 5.



The general form of the above equation is found as follows.

$$(x + 2)^2 + (y - 3)^2 = 25 \quad \text{Center-radius form}$$


$$x^2 + 4x + 4 + y^2 - 6y + 9 = 25 \quad \text{Square each binomial.}$$

$$x^2 + y^2 + 4x - 6y - 12 = 0 \quad \text{General form}$$

Graph $\frac{x^2}{9} + \frac{y^2}{4} = 1$.

x -intercepts: $(3, 0)$ and $(-3, 0)$


y -intercepts: $(0, 2)$ and $(0, -2)$



Graph $\frac{x^2}{4} - \frac{y^2}{4} = 1$.

x -intercepts: $(2, 0)$ and $(-2, 0)$

Vertices of the fundamental rectangle:
 $(2, 2)$, $(-2, 2)$, $(-2, -2)$, and $(2, -2)$



CONCEPTS

Generalized Square Root Function

For an algebraic expression in x defined by u , where $u \geq 0$, a function of the form

$$f(x) = \sqrt{u}$$

is a generalized square root function.


EXAMPLES

Graph $y = -\sqrt{4 - x^2}$.

Square each side and rearrange terms.

$$x^2 + y^2 = 4$$


This equation has a circle as its graph. However, graph only the lower half of the circle because the original equation indicates that y cannot be positive.

**10.4 Nonlinear Systems of Equations****Solving a Nonlinear System**

A nonlinear system can be solved by the substitution method, the elimination method, or a combination of the two.

Geometric Interpretation

If, for example, a nonlinear system includes two second-degree equations, then there may be zero, one, two, three, or four solutions of the system.



This system has four solutions because there are four points of intersection.

Solve the system.

$$x^2 + 2xy - y^2 = 14 \quad (1)$$

$$x^2 - y^2 = -16 \quad (2)$$

Multiply equation (2) by -1 and use the elimination method.

$$x^2 + 2xy - y^2 = 14$$

$$-x^2 \quad + y^2 = 16$$

$$\hline 2xy = 30$$

$$xy = 15$$

Solve $xy = 15$ for y to obtain $y = \frac{15}{x}$, and substitute into equation (2).

$$x^2 - y^2 = -16 \quad (2)$$

$$x^2 - \left(\frac{15}{x}\right)^2 = -16 \quad \text{Let } y = \frac{15}{x}.$$

$$x^2 - \frac{225}{x^2} = -16 \quad \text{Apply the exponent.}$$

$$x^4 + 16x^2 - 225 = 0 \quad \text{Multiply by } x^2. \text{ Add } 16x^2.$$

$$(x^2 - 9)(x^2 + 25) = 0 \quad \text{Factor.}$$

$$x^2 - 9 = 0 \quad \text{or} \quad x^2 + 25 = 0 \quad \text{Zero-factor property}$$

$$x = \pm 3 \quad \text{or} \quad x = \pm 5i \quad \text{Solve each equation}$$

Substitute these values of x into $y = \frac{15}{x}$ to obtain the corresponding values of y .

$$\text{If } x = 3, \quad \text{then } y = \frac{15}{3} = 5.$$

$$\text{If } x = -3, \quad \text{then } y = \frac{15}{-3} = -5.$$

$$\text{If } x = 5i, \quad \text{then } y = \frac{15}{5i} = \frac{3}{i} \cdot \frac{-i}{-i} = -3i.$$

$$\text{If } x = -5i, \quad \text{then } y = \frac{15}{-5i} = \frac{3}{-i} \cdot \frac{i}{i} = 3i.$$

Solution set: $\{(3, 5), (-3, -5), (5i, -3i), (-5i, 3i)\}$

CONCEPTS

10.5 Second-Degree Inequalities and Systems of Inequalities**Graphing a Second-Degree Inequality**

To graph a second-degree inequality, graph the corresponding equation as a boundary, and use test points to determine which region(s) form the solution set. Shade the appropriate region(s).

Graphing a System of Inequalities


The solution set of a system of inequalities is the intersection of the graphs (solution sets) of the inequalities in the system.

EXAMPLES

Graph $y \geq x^2 - 2x + 3$.

The boundary is a parabola that opens up with vertex $(1, 2)$. Use $(0, 0)$ as a test point. Substituting into the inequality $y > x^2 - 2x + 3$ gives a false statement,

$$0 > 3. \quad \text{False}$$




Shade the region inside (or above) the parabola.

Graph the solution set of the system.

$$3x - 5y > -15$$

$$x^2 + y^2 \leq 25$$

Graph each inequality separately, on the same axes. The graph of the system is the region that lies below the dashed line *and* inside the solid circle.



Chapter 10 Review Exercises

10.1 Graph each function. Give the domain and range.

1. $f(x) = |x + 4|$

2. $f(x) = \sqrt{x} + 3$

3. $f(x) = \frac{1}{x - 4}$

4. $f(x) = \lceil -x \rceil$

Evaluate each expression.

5. $\lceil 12 \rceil$

6. $\left\lceil 2\frac{3}{4} \right\rceil$

7. $\lceil -21 \rceil$

8. $\lceil -4.75 \rceil$

10.2 Write the center-radius form of each circle described.

9. Center: $(0, 0)$; radius: 7

10. Center: $(-2, 4)$; radius: 3

11. Center: $(4, 2)$; radius: 6

12. Center: $(-5, 0)$; radius: $\sqrt{2}$

Write the center-radius form of the circle with the given equation. Give the center and radius.

13. $x^2 + y^2 + 6x - 4y - 3 = 0$

14. $x^2 + y^2 - 8x - 2y + 13 = 0$

15. $2x^2 + 2y^2 + 4x + 20y = -34$

16. $4x^2 + 4y^2 - 24x + 16y = 48$

17. Identify the graph of each equation as a *circle* or an *ellipse*.

(a) $x^2 + y^2 = 121$

(b) $x^2 + 4y^2 = 4$

(c) $\frac{x^2}{16} + \frac{y^2}{25} = 1$

(d) $3x^2 + 3y^2 = 300$

$$39. \begin{aligned} x^2 + y^2 &= 6 \\ x^2 - 2y^2 &= -6 \end{aligned}$$

$$40. \begin{aligned} 3x^2 - 2y^2 &= 12 \\ x^2 + 4y^2 &= 18 \end{aligned}$$

10.5 Graph each inequality.

$$41. 9x^2 \geq 16y^2 + 144$$

$$42. 4x^2 + y^2 \geq 16$$

$$43. y < -(x + 2)^2 + 1$$

44. Which one of the following is a description of the graph of $x^2 + y^2 > 4$?

- A. The region inside a circle with radius 4
- B. The region outside a circle with radius 4
- C. The region inside a circle with radius 2
- D. The region outside a circle with radius 2

Graph each system of inequalities.

$$45. \begin{aligned} 2x + 5y &\leq 10 \\ 3x - y &\leq 6 \end{aligned}$$

$$46. \begin{aligned} 9x^2 &\leq 4y^2 + 36 \\ x^2 + y^2 &\leq 16 \end{aligned}$$

$$47. \begin{aligned} x + y &\geq -2 \\ y &< x + 4 \\ x &< 1 \end{aligned}$$

48. Match each system of inequalities in Column I with the quadrant that represents its graph in Column II.

I

- (a) $x < 0$
 $y > 0$
- (b) $x > 0$
 $y < 0$
- (c) $x < 0$
 $y < 0$
- (d) $x > 0$
 $y > 0$

II

- A. Quadrant I
- B. Quadrant II
- C. Quadrant III
- D. Quadrant IV

Chapter 10 Mixed Review Exercises

1. Find the center and radius of the circle with equation

$$x^2 + y^2 + 4x - 10y - 7 = 0.$$

2. *True or false:* The graph of $y = -\sqrt{1 - x^2}$ is the graph of a function.

Graph.

$$3. x^2 + 9y^2 = 9$$

$$4. x^2 - 9y^2 = 9$$

$$5. f(x) = \sqrt{4 - x}$$

$$6. x^2 + y^2 = 25$$

$$7. \frac{y^2}{4} - 1 \leq \frac{x^2}{9}$$

$$8. \begin{aligned} 4y &> 3x - 12 \\ x^2 &< 16 - y^2 \end{aligned}$$

Solve each system.

$$9. \begin{aligned} y &= x^2 - 84 \\ 4y &= 20x \end{aligned}$$

$$10. \begin{aligned} x^2 - xy - y^2 &= -5 \\ x^2 - y^2 &= -3 \end{aligned}$$

Chapter 10

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.


View the complete solutions to all Chapter Test exercises in MyLab Math.

Fill in each blank with the correct response.


- For the reciprocal function defined by $f(x) = \frac{1}{x}$, _____ is the only real number not in the domain.
- The range of the square root function $f(x) = \sqrt{x}$ is _____.
- The range of $f(x) = \llbracket x \rrbracket$, the greatest integer function, is _____.
- Match each function in parts (a)–(d) with its graph from choices A–D.

(a) $f(x) = \sqrt{x - 2}$

A.




B.




(b) $f(x) = \sqrt{x + 2}$

(c) $f(x) = \sqrt{x} + 2$

C.



D.



(d) $f(x) = \sqrt{x} - 2$

- Graph $f(x) = |x - 3| + 4$. Give the domain and range.
- Find the center and radius of the circle with equation $(x - 2)^2 + (y + 3)^2 = 16$. Sketch the graph.
- Which one of the following equations is represented by the graph of a circle?
 - $x^2 + y^2 = 0$
 - $x^2 + y^2 = -1$
 - $x^2 + y^2 = x^2 - y^2$
 - $x^2 + y^2 = 1$

For the equation that is represented by a circle, what are the coordinates of the center? What is the radius?

- Find the center and radius of the circle with equation $x^2 + y^2 + 8x - 2y = 8$.

Graph.

9. $f(x) = \sqrt{9 - x^2}$

10. $4x^2 + 9y^2 = 36$

11. $16y^2 - 4x^2 = 64$

12. $\frac{y}{2} = -\sqrt{1 - \frac{x^2}{9}}$

Identify the graph of each equation as a parabola, hyperbola, ellipse, or circle.

13. $6x^2 + 4y^2 = 12$

14. $16x^2 = 144 + 9y^2$

15. $y^2 = 20 - x^2$

16. $4y^2 + 4x = 9$

Solve each system.

$$\begin{aligned} 17. \quad & 2x - y = 9 \\ & xy = 5 \end{aligned}$$

$$\begin{aligned} 18. \quad & x - 4 = 3y \\ & x^2 + y^2 = 8 \end{aligned}$$

$$\begin{aligned} 19. \quad & x^2 + y^2 = 25 \\ & x^2 - 2y^2 = 16 \end{aligned}$$

Graph.

$$20. \quad y < x^2 - 2$$

$$\begin{aligned} 21. \quad & x - y \leq 2 \\ & x + 3y > 6 \end{aligned}$$

$$\begin{aligned} 22. \quad & x^2 + 25y^2 \leq 25 \\ & x^2 + y^2 \leq 9 \end{aligned}$$

Chapters R–10 Cumulative Review Exercises

1. Write $\frac{1}{100}$ as a decimal and as a percent.

2. Simplify $-|-1| - 5 + |4 - 10|$.

3. Match each number in Column I with the set (or sets) of numbers in Column II to which the number belongs.

I		II	
(a) 1	(b) 0	A. Natural numbers	B. Whole numbers
(c) $-\frac{2}{3}$	(d) π	C. Integers	D. Rational numbers
(e) $-\sqrt{49}$	(f) 2.75	E. Irrational numbers	F. Real numbers

4. Evaluate.

$$(a) 6^2 \quad (b) -6^2 \quad (c) (-6)^2 \quad (d) \sqrt{36} \quad (e) -\sqrt{36} \quad (f) \sqrt{-36}$$

Solve.

$$5. \quad 4 - (2x + 3) + x = 5x - 3$$

$$6. \quad -4x + 7 \geq 6x + 1$$

$$7. \quad |5x| - 6 = 14$$

$$8. \quad |2p - 5| > 15$$

9. Find the slope of the line passing through the points (2, 5) and (-4, 1).

10. Find the equation of the line passing through the point (-3, -2) and perpendicular to the graph of $2x - 3y = 7$.

Solve each system.

$$\begin{aligned} 11. \quad & 3x - y = 12 \\ & 2x + 3y = -3 \end{aligned}$$

$$\begin{aligned} 12. \quad & x + y - 2z = 9 \\ & 2x + y + z = 7 \\ & 3x - y - z = 13 \end{aligned}$$

$$\begin{aligned} 13. \quad & xy = -5 \\ & 2x + y = 3 \end{aligned}$$

14. Al and Bev traveled from their apartment to a picnic 20 mi away. Al traveled on his bike while Bev, who left later, took her car. Al's average rate was half of Bev's average rate. The trip took Al $\frac{1}{2}$ hr longer than Bev. What was Bev's average rate?

Perform the indicated operations.

$$15. \quad (5y - 3)^2$$

$$16. \quad \frac{8x^4 - 4x^3 + 2x^2 + 13x + 8}{2x + 1}$$

17. $\frac{y^2 - 4}{y^2 - y - 6} \div \frac{y^2 - 2y}{y - 1}$

18. $\frac{5}{c + 5} - \frac{2}{c + 3}$

19. $\frac{p}{p^2 + p} + \frac{1}{p^2 + p}$

20. $\frac{1}{\frac{1}{x} - \frac{1}{y}}$

21. $4\sqrt[3]{16} - 2\sqrt[3]{54}$

22. $\frac{5 + 3i}{2 - i}$

Factor.

23. $12x^2 - 7x - 10$

24. $z^4 - 1$

25. $a^3 - 27b^3$

26. Henry and Lawrence want to clean their office. Henry can do the job alone in 3 hr, and Lawrence can do it alone in 2 hr. How long will it take them if they work together?

	Rate	Time Working Together	Fractional Part of the Job Done
Henry	$\frac{1}{3}$		
Lawrence			

Simplify. Assume all variables represent positive real numbers.

27. $\frac{(2a)^{-2}a^4}{a^{-3}}$

28. $\frac{3\sqrt{5x}}{\sqrt{2x}}$

Solve.

29. $2\sqrt{x} = \sqrt{5x + 3}$

30. $10q^2 + 13q = 3$

31. $3x^2 - 3x - 2 = 0$

32. $2(x^2 - 3)^2 - 5(x^2 - 3) = 12$

33. $\log(x + 2) + \log(x - 1) = 1$

34. $F = \frac{kvw^2}{r}$ for v

35. If $f(x) = x^2 + 2x - 4$ and $g(x) = 3x + 2$, find the following.

(a) $(g \circ f)(1)$

(b) $(f \circ g)(x)$

36. If $f(x) = x^3 + 4$, find $f^{-1}(x)$.

37. Evaluate. (a) $3^{\log_3 4}$ (b) $e^{\ln 7}$

38. Use properties of logarithms to write $2 \log(3x + 7) - \log 4$ as a single logarithm.

39. Give the domain and range of the function defined by $f(x) = |x - 3|$.

Graph.

40. $f(x) = -3x + 5$

41. $f(x) = -2(x - 1)^2 + 3$

42. $\frac{x^2}{25} + \frac{y^2}{16} \leq 1$

43. $f(x) = \sqrt{x - 2}$

44. $\frac{x^2}{4} - \frac{y^2}{16} = 1$

45. $f(x) = 3^x$

STUDY SKILLS REMINDER

Have you begun to prepare for your final exam? **Review Study Skill 10, Preparing for Your Math Final Exam.**

This page intentionally left blank


FURTHER TOPICS IN ALGEBRA

The number of ancestors in the family tree of a male honeybee forms the *Fibonacci sequence*

$$1, 1, 2, 3, 5, 8, \dots,$$

where each term after the second is found by adding the two previous terms. This chapter covers *sequences, series,* and other topics in algebra.

- 11.1** Sequences and Series
- 11.2** Arithmetic Sequences
- 11.3** Geometric Sequences
- 11.4** The Binomial Theorem



11.1 Sequences and Series

OBJECTIVES

- 1 Define infinite and finite sequences.
- 2 Find the terms of a sequence, given the general term.
- 3 Find the general term of a sequence.
- 4 Use sequences to solve applied problems.
- 5 Use summation notation to evaluate a series.
- 6 Write a series using summation notation.
- 7 Find the arithmetic mean (average) of a group of numbers.

VOCABULARY

- infinite sequence
- finite sequence
- terms of a sequence
- general term
- series
- summation notation
- index of summation
- arithmetic mean (average)

NOW TRY EXERCISE 1

Given an infinite sequence with $a_n = 5 - 3n$, find a_3 .

NOW TRY ANSWER

1. $a_3 = -4$

OBJECTIVE 1 Define infinite and finite sequences.

In the Palace of the Alhambra, residence of the Moorish rulers of Granada, Spain, the Sultana's quarters feature an interesting architectural pattern.

There are 2 matched marble slabs inlaid in the floor, 4 walls, an octagon (8-sided) ceiling, 16 windows, 32 arches, and so on.

If this pattern is continued indefinitely, the set of numbers forms an *infinite sequence* whose *terms* are powers of 2.



Sequence

An **infinite sequence** is a function with the set of positive integers (natural numbers)* as domain.

A **finite sequence** is a function with domain of the form $\{1, 2, 3, \dots, n\}$, where n is a positive integer.

For any positive integer n , the function value of a sequence is written as a_n (read “*a sub-n*”). The function values

$$a_1, a_2, a_3, \dots,$$

written in order, are the **terms** of the sequence, with a_1 the first term, a_2 the second term, and so on. The expression a_n , which defines the sequence, is its **general term**.

In the Palace of the Alhambra example, the first five terms of the sequence are

$$a_1 = 2, \quad a_2 = 4, \quad a_3 = 8, \quad a_4 = 16, \quad \text{and} \quad a_5 = 32.$$

The general term for this sequence is $a_n = 2^n$.

OBJECTIVE 2 Find the terms of a sequence, given the general term.

EXAMPLE 1 Writing the Terms of Sequences from the General Term

Given an infinite sequence with $a_n = n + \frac{1}{n}$, find the following.

- (a) The second term of the sequence

$$a_2 = 2 + \frac{1}{2} = \frac{5}{2} \quad \text{Replace } n \text{ in } a_n = n + \frac{1}{n} \text{ with } 2.$$


$$(b) \quad a_{10} = 10 + \frac{1}{10} = \frac{101}{10} \quad 10 + \frac{1}{10} = \frac{10}{1} + \frac{1}{10} = \frac{100}{10} + \frac{1}{10} = \frac{101}{10}$$

$$(c) \quad a_{12} = 12 + \frac{1}{12} = \frac{145}{12} \quad 12 + \frac{1}{12} = \frac{12}{1} + \frac{1}{12} = \frac{144}{12} + \frac{1}{12} = \frac{145}{12}$$

NOW TRY

*In this chapter we will use the terms *positive integers* and *natural numbers* interchangeably.

Graphing calculators can generate and graph sequences. See **FIGURE 1**. The calculator must be in dot mode so that the discrete points on the graph are not connected. **Remember that the domain of a sequence consists only of positive integers.**



The first five terms of the sequence $a_n = 2^n$

The first five terms of $a_n = 2^n$ are graphed here. The display indicates that the fourth term is 16; that is, $a_4 = 2^4 = 16$.

FIGURE 1

OBJECTIVE 3 Find the general term of a sequence.

EXAMPLE 2 Finding the General Term of a Sequence

Determine an expression for the general term a_n of the sequence.

$$5, 10, 15, 20, 25, \dots$$

Notice that the terms are all multiples of 5. The **first** term is $5(1)$, the **second** is $5(2)$, and so on. The general term that will produce the given first five terms is

$$a_n = 5n.$$

NOW TRY

OBJECTIVE 4 Use sequences to solve applied problems.

Applied problems may involve *finite sequences*.

EXAMPLE 3 Using a Sequence in an Application

Saad borrows \$5000 and agrees to pay \$500 monthly, plus interest of 1% on the unpaid balance from the beginning of the first month. Find the payments for the first four months and the remaining debt at the end of that period.

The payments and remaining balances are calculated as follows.

<i>First month</i>	Payment: $\$500 + 0.01(\$5000) = \$550$ Balance: $\$5000 - \$500 = \$4500$
<i>Second month</i>	Payment: $\$500 + 0.01(\$4500) = \$545$ Balance: $\$5000 - 2 \cdot \$500 = \$4000$
<i>Third month</i>	Payment: $\$500 + 0.01(\$4000) = \$540$ Balance: $\$5000 - 3 \cdot \$500 = \$3500$
<i>Fourth month</i>	Payment: $\$500 + 0.01(\$3500) = \$535$ Balance: $\$5000 - 4 \cdot \$500 = \$3000$

The payments for the first four months are

$$\$550, \$545, \$540, \$535$$

and the remaining debt at the end of the period is $\$3000$.

NOW TRY

NOW TRY EXERCISE 2

Determine an expression for the general term a_n of the sequence.

$$-3, 9, -27, 81, \dots$$

NOW TRY EXERCISE 3

Chase borrows \$8000 and agrees to pay \$400 monthly, plus interest of 2% on the unpaid balance from the beginning of the first month. Find the payments for the first four months and the remaining debt at the end of that period.

NOW TRY ANSWERS

- $a_n = (-3)^n$
- payments: \$560, \$552, \$544, \$536; balance: \$6400

OBJECTIVE 5 Use summation notation to evaluate a series.

By adding the terms of a sequence, we obtain a *series*.

Series

A **series** is the indicated sum of the terms of a sequence.

For example, if we consider the sum of the payments listed in **Example 3**, namely,

$$550 + 545 + 540 + 535,$$

we have a series that represents the total payments for the first four months. Because a sequence can be finite or infinite, there are both finite and infinite series.

We use a compact notation, called **summation notation**, to write a series from the general term of the corresponding sequence. In mathematics, the Greek letter Σ (**sigma**) is used to denote summation. For example, the sum of the first six terms of the sequence with general term

$$a_n = 3n + 2$$

is written as

$$\sum_{i=1}^6 (3i + 2).$$

The letter i is the **index of summation**. We read this as “the sum from $i = 1$ to 6 of $3i + 2$.” To find this sum, we replace the letter i in $3i + 2$ with 1, 2, 3, 4, 5, and 6, and add the resulting terms.

CAUTION This use of i as the index of summation has no connection with the complex number i .

EXAMPLE 4 Evaluating Series Written Using Summation Notation

Write each series as a sum of terms and then find the sum.

- (a) $\sum_{i=1}^6 (3i + 2)$
- Multiply and then add.
- $$= (3 \cdot 1 + 2) + (3 \cdot 2 + 2) + (3 \cdot 3 + 2)$$
- $$+ (3 \cdot 4 + 2) + (3 \cdot 5 + 2) + (3 \cdot 6 + 2)$$
- Replace i with 1, 2, 3, 4, 5, 6.
- $$= 5 + 8 + 11 + 14 + 17 + 20$$
- Simplify inside the parentheses.
- $$= 75$$
- Add.
- (b) $\sum_{i=1}^5 (i - 4)$
- $$= (1 - 4) + (2 - 4) + (3 - 4) + (4 - 4) + (5 - 4)$$
- $i = 1, 2, 3, 4, 5$
- $$= -3 - 2 - 1 + 0 + 1$$
- Subtract.
- $$= -5$$
- Simplify.


NOW TRY EXERCISE 4

Write the series as a sum of terms and then find the sum.

$$\sum_{i=1}^5 (i^2 - 4)$$

$$(c) \sum_{i=3}^7 3i^2$$

$$= 3(3)^2 + 3(4)^2 + 3(5)^2 + 3(6)^2 + 3(7)^2 \quad i = 3, 4, 5, 6, 7$$

$$= 27 + 48 + 75 + 108 + 147$$

$$= 405$$

Square, and then multiply.

Add.

NOW TRY 

OBJECTIVE 6 Write a series using summation notation.

In **Example 4**, we started with summation notation and wrote each series using + signs. Given a series, we can write it using summation notation by observing a pattern in the terms and writing the general term accordingly.


NOW TRY EXERCISE 5

Write each series using summation notation.

(a) $3 + 5 + 7 + 9 + 11$

(b) $-1 - 4 - 9 - 16 - 25$

EXAMPLE 5 Writing Series Using Summation Notation

Write each series using summation notation.

(a) $2 + 5 + 8 + 11$

First, find a general term a_n that will give these four terms for $a_1, a_2, a_3,$ and $a_4,$ respectively. Each term is one less than a multiple of 3, so try $3i - 1$.

$$3(1) - 1 = 2 \quad i = 1$$

$$3(2) - 1 = 5 \quad i = 2$$

$$3(3) - 1 = 8 \quad i = 3$$

$$3(4) - 1 = 11 \quad i = 4$$

(There may be other expressions that also work.) Because i ranges from 1 to 4,

$$2 + 5 + 8 + 11 = \sum_{i=1}^4 (3i - 1).$$

(b) $8 + 27 + 64 + 125 + 216$

These numbers are the cubes of 2, 3, 4, 5, and 6, so the general term is i^3 .

$$8 + 27 + 64 + 125 + 216 = \sum_{i=2}^6 i^3$$

NOW TRY 

OBJECTIVE 7 Find the arithmetic mean (average) of a group of numbers.

Arithmetic Mean or Average

The **arithmetic mean**, or **average**, of a set of values is symbolized \bar{x} and is found by dividing their sum by the number of values.

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n}$$

The values of x_i represent the individual numbers in the group, and n represents the number of values.

NOW TRY ANSWERS

4. $-3 + 0 + 5 + 12 + 21 = 35$

5. (a) $\sum_{i=1}^5 (2i + 1)$ (b) $\sum_{i=1}^5 -i^2$

NOW TRY EXERCISE 6

The table shows the top five American Quarter Horse States in 2016 based on the total number of registered Quarter Horses. To the nearest whole number, what is the average number of Quarter Horses registered per state in these top five states?

State	Number of Registered Quarter Horses
Texas	418,249
Oklahoma	164,265
California	114,623
Missouri	95,877
Montana	87,418

Data from American Quarter Horse Association.

NOW TRY ANSWER

6. 176,086 quarter horses

EXAMPLE 6 Finding the Arithmetic Mean (Average)

The table shows the number of FDIC-insured financial institutions for each year during the period from 2011 through 2017. What was the average number of institutions per year for this 7-yr period?

Year	Number of Institutions
2011	7523
2012	7255
2013	6950
2014	6669
2015	6358
2016	6068
2017	5797

Data from U.S. Federal Deposit Insurance Corporation.



$$\bar{x} = \frac{\sum_{i=1}^7 x_i}{7}$$

Let $x_1 = 7523$, $x_2 = 7255$, and so on. There are 7 values in the group, so $n = 7$.

$$\bar{x} = \frac{7523 + 7255 + 6950 + 6669 + 6358 + 6068 + 5797}{7}$$

$$\bar{x} = 6660$$

The average number of institutions per year for this 7-yr period was 6660.

NOW TRY

11.1 Exercises

FOR EXTRA HELP



MyLab Math

Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Make study cards to help you learn and remember the material in this chapter.

Review Study Skill 5,
Using Study Cards.

Concept Check Fill in each blank with the correct response.

- The domain of an infinite sequence is _____.
- In the sequence 3, 6, 9, 12, the term $a_3 =$ _____.
- If $a_n = 2n$, then $a_{40} =$ _____.
- If $a_n = (-1)^n$, then $a_{115} =$ _____.
- The value of the sum $\sum_{i=1}^3 (i + 2)$ is _____.
- The arithmetic mean of -4 , -2 , 0 , 2 , and 4 is _____.

Write the first five terms of each sequence. See Example 1.

7. $a_n = n + 1$

8. $a_n = n + 4$

9. $a_n = \frac{n + 3}{n}$

10. $a_n = \frac{n + 2}{n}$

11. $a_n = 3^n$

12. $a_n = 2^n$

13. $a_n = -\frac{1}{n^2}$

14. $a_n = -\frac{2}{n^2}$

15. $a_n = 5(-1)^{n-1}$

16. $a_n = 6(-1)^{n+1}$

17. $a_n = n - \frac{1}{n}$

18. $a_n = n + \frac{4}{n}$

Find the indicated term for each sequence. See Example 1.

19. $a_n = -9n + 2$; a_8

20. $a_n = -3n + 7$; a_{12}

21. $a_n = \frac{3n + 7}{2n - 5}$; a_{14}

22. $a_n = \frac{5n - 9}{3n + 8}$; a_{16}

23. $a_n = (n + 1)(2n + 3)$; a_8

24. $a_n = (5n - 2)(3n + 1)$; a_{10}

Determine an expression for the general term a_n of each sequence. See Example 2.

25. 4, 8, 12, 16, ...

26. 7, 14, 21, 28, ...

27. -8, -16, -24, -32, ...

28. -10, -20, -30, -40, ...

29. $\frac{1}{3}, \frac{1}{9}, \frac{1}{27}, \frac{1}{81}, \dots$

30. $\frac{2}{5}, \frac{2}{25}, \frac{2}{125}, \frac{2}{625}, \dots$

31. $\frac{2}{5}, \frac{3}{6}, \frac{4}{7}, \frac{5}{8}, \dots$

32. $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \dots$

Solve each applied problem by writing the first few terms of a sequence. See Example 3.

33. Horacio borrows \$1000 and agrees to pay \$100 plus interest of 1% on the unpaid balance each month. Find the payments for the first six months and the remaining debt at the end of that period.

34. Leslie is offered a new job with a salary of $20,000 + 2500n$ dollars per year at the end of the n th year. Write a sequence showing her salary at the end of each of the first 5 yr. If she continues in this way, what will her salary be at the end of the tenth year?

35. Suppose that an automobile loses $\frac{1}{5}$ of its value each year; that is, at the end of any given year, the value is $\frac{4}{5}$ of the value at the beginning of that year. If a car costs \$20,000 new, what is its value at the end of 5 yr, to the nearest dollar?

36. A certain car loses $\frac{1}{2}$ of its value each year. If this car cost \$40,000 new, what is its value at the end of 6 yr?

Write each series as a sum of terms and then find the sum. See Example 4.

37. $\sum_{i=1}^5 (i + 3)$

38. $\sum_{i=1}^6 (i + 9)$

39. $\sum_{i=1}^3 (i^2 + 2)$

40. $\sum_{i=1}^4 (i^3 + 3)$

41. $\sum_{i=1}^6 (-1)^i \cdot 2$

42. $\sum_{i=1}^5 (-1)^i \cdot i$

43. $\sum_{i=3}^7 (i - 3)(i + 2)$

44. $\sum_{i=2}^6 (i + 3)(i - 4)$

Write each series using summation notation. See Example 5.

45. $3 + 4 + 5 + 6 + 7$

46. $7 + 8 + 9 + 10 + 11$

47. $-2 + 4 - 8 + 16 - 32$

48. $-1 + 2 - 3 + 4 - 5 + 6$

49. $1 + 4 + 9 + 16$

50. $1 + 16 + 81 + 256$

51. **Concept Check** When asked to write the series $-1 - 4 - 9 - 16 - 25$ using summation notation, a student incorrectly wrote the following.

$$\sum_{i=1}^5 (-i)^2$$

WHAT WENT WRONG? Give the correct summation notation.

52. **Concept Check** When asked to write the first five terms of the sequence defined by $a_n = 5n - 1$, a student incorrectly wrote the following.

$$4 + 9 + 14 + 19 + 24$$

WHAT WENT WRONG? Give the correct answer.

Find the arithmetic mean for each set of values. See Example 6.

53. 8, 11, 14, 9, 7, 6, 8

54. 10, 12, 8, 19, 23, 12

55. 5, 9, 8, 2, 4, 7, 3, 2, 0

56. 2, 1, 4, 8, 3, 7, 10, 8, 0

Solve each problem. See Example 6.

57. The number of mutual funds operating in the United States each year during the period 2012 through 2016 is given in the table. To the nearest whole number, what was the average number of mutual funds operating per year during the given period?

Year	Number of Mutual Funds
2012	8744
2013	8972
2014	9258
2015	9517
2016	9511

Data from Investment Company Institute.

58. The total assets of mutual funds operating in the United States, in billions of dollars, for each year during the period 2012 through 2016 are shown in the table. What were the average assets per year during this period?

Year	Assets (in billions of dollars)
2012	13,054
2013	15,049
2014	15,873
2015	15,650
2016	16,344

Data from Investment Company Institute.

11.2 Arithmetic Sequences

OBJECTIVES

- 1 Find the common difference of an arithmetic sequence.
- 2 Find the general term of an arithmetic sequence.
- 3 Use an arithmetic sequence in an application.
- 4 Find any specified term or the number of terms of an arithmetic sequence.
- 5 Find the sum of a specified number of terms of an arithmetic sequence.

OBJECTIVE 1 Find the common difference of an arithmetic sequence.

In this section, we introduce *arithmetic sequences*.

Arithmetic Sequence

An **arithmetic sequence**, or **arithmetic progression**, is a sequence in which each term after the first is found by adding a constant number to the preceding term.

For example, $6, 11, 16, 21, 26, \dots$ Arithmetic sequence

is an arithmetic sequence in which the difference between any two adjacent terms is always 5. The number 5 is the **common difference** of the arithmetic sequence. The common difference d is found by subtracting a_n from a_{n+1} in any such pair of terms.

$$d = a_{n+1} - a_n \quad \text{Common difference}$$

VOCABULARY

- arithmetic sequence
(arithmetic progression)
- common difference

NOW TRY EXERCISE 1

Determine the common difference d for the arithmetic sequence.

$$-4, -13, -22, -31, -40, \dots$$

NOW TRY EXERCISE 2

Write the first five terms of the arithmetic sequence with first term 10 and common difference -8 .

EXAMPLE 1 Finding the Common Difference

Determine the common difference d for the arithmetic sequence.

$$-11, -4, 3, 10, 17, 24, \dots$$

The sequence is arithmetic, so d is the difference between any two adjacent terms. We arbitrarily choose the terms 10 and 17.

$$d = a_{n+1} - a_n$$

$$d = 17 - 10 \quad a_5 - a_4$$

$$d = 7 \quad \text{Common difference}$$

Verify that *any* two adjacent terms would give the same result.

NOW TRY **EXAMPLE 2** Writing the Terms of an Arithmetic Sequence

Write the first five terms of the arithmetic sequence with first term 3 and common difference -2 .

$$a_1 = 3 \quad \text{First term}$$

$$a_2 = 3 + (-2) = 1 \quad \text{Add } d = -2.$$

$$a_3 = 1 + (-2) = -1 \quad \text{Add } -2.$$

$$a_4 = -1 + (-2) = -3 \quad \text{Add } -2.$$

$$a_5 = -3 + (-2) = -5 \quad \text{Add } -2.$$

The first five terms are 3, 1, -1 , -3 , -5 .

NOW TRY **OBJECTIVE 2** Find the general term of an arithmetic sequence.

Generalizing from **Example 2**, if we know the first term a_1 and the common difference d of an arithmetic sequence, then the sequence is completely defined as

$$a_1, \quad a_2 = a_1 + d, \quad a_3 = a_1 + 2d, \quad a_4 = a_1 + 3d, \dots$$

Writing the terms of the sequence in this way suggests the following formula for a_n .

General Term of an Arithmetic Sequence

The general term a_n of an arithmetic sequence with first term a_1 and common difference d is given by the following.

$$a_n = a_1 + (n - 1)d$$

Because $a_n = a_1 + (n - 1)d$ Formula for a_n

$$a_n = dn + (a_1 - d) \quad \text{Properties of real numbers}$$

is a linear function in n , any linear expression of the form $dn + c$, where d and c are real numbers, defines an arithmetic sequence.

NOW TRY ANSWERS

1. $d = -9$
2. 10, 2, -6 , -14 , -22


**NOW TRY
EXERCISE 3**

Determine an expression for the general term of the arithmetic sequence. Then find a_{20} .

$$-5, 0, 5, 10, 15, \dots$$

EXAMPLE 3 Finding the General Term of an Arithmetic Sequence

Determine an expression for the general term of the arithmetic sequence. Then find a_{20} .

$$-9, -6, -3, 0, 3, 6, \dots$$

The first term is $a_1 = -9$. Begin by finding d .

$$d = -6 - (-9) \quad \text{Let } d = a_2 - a_1.$$

$$d = 3 \quad \text{Subtract.}$$

Now find a_n .

$$a_n = a_1 + (n - 1)d \quad \text{Formula for } a_n$$

$$a_n = -9 + (n - 1)(3) \quad \text{Let } a_1 = -9, \text{ and } d = 3.$$

$$a_n = -9 + 3n - 3 \quad \text{Distributive property}$$

$$a_n = 3n - 12 \quad \text{Combine like terms.}$$

The general term is $a_n = 3n - 12$. Now find a_{20} .

$$a_{20} = 3(20) - 12 \quad \text{Let } n = 20.$$

$$a_{20} = 60 - 12 \quad \text{Multiply.}$$

$$a_{20} = 48 \quad \text{Subtract.}$$

NOW TRY 

OBJECTIVE 3 Use an arithmetic sequence in an application.


**NOW TRY
EXERCISE 4**

Ginny is saving money for her son's college education. She makes an initial contribution of \$1000 and deposits an additional \$120 each month for the next 96 months. Disregarding interest, how much money will be in the account after 96 months?

EXAMPLE 4 Applying an Arithmetic Sequence

Leonid's uncle decides to start a fund for Leonid's education. He makes an initial contribution of \$3000 and deposits an additional \$500 each month. How much will be in the fund after 24 months? (Disregard any interest.)

After n months, the fund will contain

$$a_n = 3000 + 500n \text{ dollars.} \quad \text{Use an arithmetic sequence.}$$

To find the amount in the fund after 24 months, find a_{24} .

$$a_{24} = 3000 + 500(24) \quad \text{Let } n = 24.$$

$$a_{24} = 3000 + 12,000 \quad \text{Multiply.}$$

$$a_{24} = 15,000 \quad \text{Add.}$$

The fund will contain \$15,000 (disregarding interest) after 24 months. **NOW TRY** 

OBJECTIVE 4 Find any specified term or the number of terms of an arithmetic sequence.

The formula for the general term of an arithmetic sequence has four variables:

$$a_n, a_1, n, \text{ and } d.$$

If we know the values of any three of these variables, we can use the formula to find the value of the fourth variable.

NOW TRY ANSWERS

3. $a_n = 5n - 10$; $a_{20} = 90$

4. \$12,520


**NOW TRY
EXERCISE 5**

Find the indicated term for each arithmetic sequence.

(a) $a_1 = 21, d = -3; a_{22}$

(b) $a_7 = 25, a_{12} = 40; a_{19}$

EXAMPLE 5 Finding Specified Terms in Arithmetic Sequences

Find the indicated term for each arithmetic sequence.

(a) $a_1 = -6, d = 12; a_{15}$

$$a_n = a_1 + (n - 1)d \quad \text{Formula for } a_n$$

$$a_{15} = a_1 + (15 - 1)d \quad \text{Let } n = 15.$$

$$a_{15} = -6 + 14(12) \quad \text{Let } a_1 = -6, d = 12.$$

$$a_{15} = 162 \quad \text{Multiply, and then add.}$$

(b) $a_5 = 2, a_{11} = -10; a_{17}$

Any term can be found if a_1 and d are known. Use the formula for a_n .

$$a_n = a_1 + (n - 1)d$$

$$a_5 = a_1 + (5 - 1)d \quad \text{Let } n = 5.$$

$$a_5 = a_1 + 4d \quad \text{Simplify.}$$

$$2 = a_1 + 4d \quad a_5 = 2$$

$$a_n = a_1 + (n - 1)d$$

$$a_{11} = a_1 + (11 - 1)d \quad \text{Let } n = 11.$$

$$a_{11} = a_1 + 10d \quad \text{Simplify.}$$

$$-10 = a_1 + 10d \quad a_{11} = -10$$

This gives a system of two equations in two variables, a_1 and d .

$$a_1 + 4d = 2 \quad (1)$$

$$a_1 + 10d = -10 \quad (2)$$

Multiply equation (2) by -1 and add to equation (1) to eliminate a_1 .

$$a_1 + 4d = 2 \quad (1)$$

$$-a_1 - 10d = 10 \quad \text{Multiply (2) by } -1.$$

$$-6d = 12 \quad \text{Add.}$$

$$d = -2 \quad \text{Divide by } -6.$$

Now find a_1 by substituting -2 for d into either equation.

$$a_1 + 10d = -10 \quad (2)$$

$$a_1 + 10(-2) = -10 \quad \text{Let } d = -2 \text{ in (2).}$$

$$a_1 - 20 = -10 \quad \text{Multiply.}$$

$$a_1 = 10 \quad \text{Add 20.}$$

Use the formula for a_n to find a_{17} .

$$a_n = a_1 + (n - 1)d \quad \text{Formula for } a_n$$

$$a_{17} = a_1 + (17 - 1)d \quad \text{Let } n = 17.$$

$$a_{17} = a_1 + 16d \quad \text{Subtract.}$$

$$a_{17} = 10 + 16(-2) \quad \text{Let } a_1 = 10, d = -2.$$

$$a_{17} = -22 \quad \text{Multiply, and then add.}$$

NOW TRY ANSWERS

5. (a) -42 (b) 61


NOW TRY EXERCISE 6

Find the number of terms in the arithmetic sequence.

$$1, \frac{4}{3}, \frac{5}{3}, 2, \dots, 11$$

EXAMPLE 6 Finding the Number of Terms in an Arithmetic Sequence

Find the number of terms in the arithmetic sequence.

$$-8, -2, 4, 10, \dots, 52$$

Let n represent the number of terms in the sequence. Here $a_n = 52$, $a_1 = -8$, and $d = -2 - (-8) = 6$, so we use the formula for a_n to find n .

$$a_n = a_1 + (n - 1)d \quad \text{Formula for } a_n$$

$$52 = -8 + (n - 1)(6) \quad \text{Let } a_n = 52, a_1 = -8, d = 6.$$

$$52 = -8 + 6n - 6 \quad \text{Distributive property}$$

$$52 = 6n - 14 \quad \text{Simplify.}$$

$$66 = 6n \quad \text{Add 14.}$$

$$n = 11 \quad \text{Divide by 6.}$$

The sequence has 11 terms.

NOW TRY 

OBJECTIVE 5 Find the sum of a specified number of terms of an arithmetic sequence.

To find a formula for the sum S_n of the first n terms of a given arithmetic sequence, we can write out the terms in two ways. We start with the first term, and then with the last term. Then we add the terms in columns.

$$S_n = a_1 + (a_1 + d) + (a_1 + 2d) + \cdots + [a_1 + (n - 1)d]$$

$$S_n = a_n + (a_n - d) + (a_n - 2d) + \cdots + [a_n - (n - 1)d]$$

$$S_n + S_n = (a_1 + a_n) + (a_1 + a_n) + (a_1 + a_n) + \cdots + (a_1 + a_n)$$

$$2S_n = n(a_1 + a_n) \quad \text{There are } n \text{ terms of } a_1 + a_n \text{ on the right.}$$

Formula for S_n

$$S_n = \frac{n}{2}(a_1 + a_n) \quad \text{Divide by 2.}$$


NOW TRY EXERCISE 7

Evaluate the sum of the first seven terms of the arithmetic sequence in which $a_n = 5n - 7$.

EXAMPLE 7 Finding the Sum of the First n Terms of an Arithmetic Sequence

Evaluate the sum of the first five terms of the arithmetic sequence in which $a_n = 2n - 5$.

Begin by evaluating a_1 and a_5 .

$$\begin{array}{l|l} a_1 = 2(1) - 5 & a_5 = 2(5) - 5 \\ a_1 = -3 & a_5 = 5 \end{array}$$

Now evaluate the sum using $a_1 = -3$, $a_5 = 5$, and $n = 5$.

$$S_n = \frac{n}{2}(a_1 + a_n) \quad \text{Formula for } S_n$$

$$S_5 = \frac{5}{2}(-3 + 5) \quad \text{Substitute.}$$

$$S_5 = 5 \quad \text{Add, and then multiply.}$$

NOW TRY 

NOW TRY ANSWERS

6. 31
7. 91

It is possible to express the sum S_n of an arithmetic sequence in terms of a_1 and d , the quantities that define the sequence. We have established that

$$S_n = \frac{n}{2}(a_1 + a_n) \quad \text{and} \quad a_n = a_1 + (n - 1)d.$$

We substitute the expression for a_n into the expression for S_n .

$$S_n = \frac{n}{2}(a_1 + [a_1 + (n - 1)d]) \quad \text{Substitute for } a_n.$$

$$S_n = \frac{n}{2}[2a_1 + (n - 1)d] \quad \text{Combine like terms.}$$

Sum of the First n Terms of an Arithmetic Sequence

The sum S_n of the first n terms of the arithmetic sequence with first term a_1 , n th term a_n , and common difference d is given by either of the following.

$$S_n = \frac{n}{2}(a_1 + a_n) \quad \text{or} \quad S_n = \frac{n}{2}[2a_1 + (n - 1)d]$$

The first formula is used when the first and last terms are known; otherwise, the second formula is used.

NOW TRY EXERCISE 8

Evaluate the sum of the first nine terms of the arithmetic sequence having first term -8 and common difference -5 .

EXAMPLE 8 Finding the Sum of the First n Terms of an Arithmetic Sequence

Evaluate the sum of the first eight terms of the arithmetic sequence having first term 3 and common difference -2 .

The known values $a_1 = 3$, $d = -2$, and $n = 8$ appear in the second formula for S_n , so we use it.

$$S_n = \frac{n}{2}[2a_1 + (n - 1)d] \quad \text{Second formula for } S_n$$

$$S_8 = \frac{8}{2}[2(3) + (8 - 1)(-2)] \quad \text{Let } a_1 = 3, d = -2, n = 8.$$

$$S_8 = 4[6 - 14] \quad \text{Work inside the brackets.}$$

$$S_8 = -32 \quad \text{Subtract, and then multiply.}$$

NOW TRY 

As mentioned earlier, a linear expression of the form

$$dn + c, \quad \text{where } d \text{ and } c \text{ are real numbers,}$$

defines an arithmetic sequence. For example, the sequences defined by $a_n = 2n + 5$ and $a_n = n - 3$ are arithmetic sequences. For this reason,

$$\sum_{i=1}^n (di + c)$$

represents the sum of the first n terms of an arithmetic sequence having first term $a_1 = d(1) + c = d + c$ and general term $a_n = d(n) + c = dn + c$. We can find this sum with the first formula for S_n , as shown in the next example.

NOW TRY ANSWER

8. -252


NOW TRY EXERCISE 9

Evaluate $\sum_{i=1}^{11} (5i - 7)$.

EXAMPLE 9 Using S_n to Evaluate a Summation

Evaluate $\sum_{i=1}^{12} (2i - 1)$.

This is the sum of the first 12 terms of the arithmetic sequence having $a_n = 2n - 1$. We find this sum, S_{12} , using the first formula for S_n .

$$S_n = \frac{n}{2}(a_1 + a_n) \quad \text{First formula for } S_n$$

$$S_{12} = \frac{12}{2} [(2(1) - 1) + (2(12) - 1)] \quad \text{Let } n = 12, a_1 = 2(1) - 1, \text{ and } a_{12} = 2(12) - 1.$$

$$S_{12} = 6(1 + 23) \quad \text{Evaluate } a_1 \text{ and } a_{12}.$$

$$S_{12} = 6(24) \quad \text{Add.}$$

$$S_{12} = 144 \quad \text{Multiply.}$$

NOW TRY 


NOW TRY ANSWER

9. 253

11.2 Exercises

FOR
EXTRA
HELP


MyLab Math

 Video solutions for select problems available in MyLab Math

STUDY SKILLS REMINDER

Have you begun to prepare for your final exam? **Review Study Skill 10, Preparing for Your Math Final Exam.**

Concept Check Fill in each blank with the correct response.

- In an arithmetic sequence, if any term is subtracted from the term that follows it, the result is the common _____ of the sequence.
- For the arithmetic sequence having $a_n = 2n + 4$, the term $a_3 =$ _____.
- The sum of the first five terms of the arithmetic sequence 1, 6, 11, ... is _____.
- The number of terms in the arithmetic sequence 2, 4, 6, ..., 100 is _____.

Concept Check Fill in the blanks to complete the terms of each arithmetic sequence.

- $1, \frac{3}{2}, 2, \underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}}$
- $\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, 1, 5, 9$
- $11, \underline{\hspace{1cm}}, 21, \underline{\hspace{1cm}}, 31$
- $7, \underline{\hspace{1cm}}, \underline{\hspace{1cm}}, -17, \underline{\hspace{1cm}}$

If the given sequence is arithmetic, find the common difference d . If the sequence is not arithmetic, say so. **See Example 1.**

- 1, 2, 3, 4, 5, ...
- 2, 5, 8, 11, ...
- 2, -4, 6, -8, 10, -12, ...
- 1, 2, 4, 7, 11, 16, ...
- 10, 5, 0, -5, -10, ...
- 6, -10, -14, -18, ...

Write the first five terms of each arithmetic sequence. **See Example 2.**

- $a_1 = 5, d = 4$
- $a_1 = 6, d = 7$
- $a_1 = -2, d = -4$
- $a_1 = -3, d = -5$
- $a_1 = \frac{1}{3}, d = \frac{2}{3}$
- $a_1 = 0.25, d = 0.55$

Determine an expression for the general term of each arithmetic sequence. Then find a_{25} . See Example 3.

21. $a_1 = 2, d = 5$

22. $a_1 = 5, d = 3$

23. $3, \frac{15}{4}, \frac{9}{2}, \frac{21}{4}, \dots$

24. $1, \frac{5}{3}, \frac{7}{3}, 3, \dots$

25. $-3, 0, 3, \dots$

26. $-10, -5, 0, \dots$

Find the indicated term for each arithmetic sequence. See Example 5.

27. $a_1 = 4, d = 3; a_{25}$

28. $a_1 = 1, d = -3; a_{12}$

29. $2, 4, 6, \dots; a_{24}$

30. $1, 5, 9, \dots; a_{50}$

31. $a_{12} = -45, a_{10} = -37; a_1$

32. $a_{10} = -2, a_{15} = 13; a_3$

Find the number of terms in each arithmetic sequence. See Example 6.

33. $3, 5, 7, \dots, 33$

34. $4, 8, 12, \dots, 204$

35. $-7, -13, -19, \dots, -157$

36. $4, 1, -2, \dots, -32$

37. $\frac{3}{4}, 3, \frac{21}{4}, \dots, 12$

38. $2, \frac{3}{2}, 1, \frac{1}{2}, \dots, -5$

39. **Concept Check** A student incorrectly claimed that the common difference for the arithmetic sequence

$$-15, -10, -5, 0, 5, \dots$$

is -5 . **WHAT WENT WRONG?** Find the correct common difference.

40. **Concept Check** A student incorrectly claimed that there are 100 terms in the arithmetic sequence

$$2, 4, 6, 8, \dots, 100.$$

WHAT WENT WRONG? How many terms are there?

Evaluate S_6 for each arithmetic sequence. See Examples 7 and 8.

41. $a_n = 3n - 8$

42. $a_n = 5n - 12$

43. $a_n = 4 + 3n$

44. $a_n = 9 + 5n$

45. $a_n = -\frac{1}{2}n + 7$

46. $a_n = -\frac{2}{3}n + 6$

47. $a_1 = 6, d = 3$

48. $a_1 = 5, d = 4$

49. $a_1 = 7, d = -3$

50. $a_1 = -5, d = -4$

51. $a = 0.9, d = -0.5$

52. $a_1 = 2.4, d = 0.7$

Use a formula for S_n to evaluate each series. See Example 9.

53. $\sum_{i=1}^{10} (8i - 5)$

54. $\sum_{i=1}^{17} (3i - 1)$

55. $\sum_{i=1}^{20} \left(\frac{3}{2}i + 4 \right)$

56. $\sum_{i=1}^{11} \left(\frac{1}{2}i - 1 \right)$

57. $\sum_{i=1}^{250} i$

58. $\sum_{i=1}^{2000} i$

Solve each problem. See Examples 4, 7, 8, and 9.

59. Nancy's aunt has promised to deposit \$1 in her account on the first day of her birthday month, \$2 on the second day, \$3 on the third day, and so on for 30 days. How much will this amount to over the entire month?



60. Billy's aunt deposits \$2 in his savings account on the first day of February, \$4 on the second day, \$6 on the third day, and so on for the entire month. How much will this amount to? (Assume that it is a leap year.)
61. Suppose that Cherian is offered a job at \$1600 per month with a guaranteed increase of \$50 every six months for 5 yr. What will Cherian's salary be at the end of that time?
62. Malik is offered a job with a starting salary of \$2000 per month with a guaranteed increase of \$100 every four months for 3 yr. What will Malik's salary be at the end of that time?
63. A seating section in a theater-in-the-round has 20 seats in the first row, 22 in the second row, 24 in the third row, and so on for 25 rows. How many seats are there in the last row? How many seats are there in the section?
64. Constantin has started on a fitness program. He plans to jog 10 min per day for the first week and then to add 10 min per day each week until he is jogging an hour each day. In which week will this occur? What is the total number of minutes he will run during the first four weeks?
65. A child builds with blocks, placing 35 blocks in the first row, 31 in the second row, 27 in the third row, and so on. Continuing this pattern, can he end with a row containing exactly 1 block? If not, how many blocks will the last row contain? How many rows can he build this way?



66. A stack of firewood has 28 pieces on the bottom, 24 on top of those, then 20, and so on. If there are 108 pieces of wood, how many rows are there?

11.3 Geometric Sequences

OBJECTIVES

- 1 Find the common ratio of a geometric sequence.
- 2 Find the general term of a geometric sequence.
- 3 Find any specified term of a geometric sequence.
- 4 Find the sum of a specified number of terms of a geometric sequence.
- 5 Apply the formula for the future value of an ordinary annuity.
- 6 Find the sum of an infinite number of terms of a geometric sequence.

OBJECTIVE 1 Find the common ratio of a geometric sequence.

In an arithmetic sequence, each term after the first is found by *adding* a fixed number to the previous term. A *geometric sequence* is defined using multiplication.

Geometric Sequence

A **geometric sequence**, or **geometric progression**, is a sequence in which each term after the first is found by multiplying the preceding term by a nonzero constant.

We find the constant multiplier r , called the **common ratio**, by dividing any term a_{n+1} by the preceding term, a_n .

$$r = \frac{a_{n+1}}{a_n} \quad \text{Common ratio}$$

For example, $2, 6, 18, 54, 162, \dots$ Geometric sequence

is a geometric sequence in which the first term is 2 and the common ratio is 3.

$$r = \frac{6}{2} = \frac{18}{6} = \frac{54}{18} = \frac{162}{54} = 3 \leftarrow \frac{a_{n+1}}{a_n} = 3 \text{ for all } n.$$

VOCABULARY

- geometric sequence (geometric progression)
- common ratio
- annuity
- ordinary annuity
- payment period
- term of an annuity
- future value of an annuity

**NOW TRY EXERCISE 1**

Determine the common ratio r for the geometric sequence.

$$\frac{1}{4}, -1, 4, -16, 64, \dots$$

EXAMPLE 1 Finding the Common Ratio

Determine the common ratio r for the geometric sequence.

$$15, \frac{15}{2}, \frac{15}{4}, \frac{15}{8}, \dots$$

To find r , choose any two successive terms and divide the second one by the first. We choose the second and third terms of the sequence.

$$r = \frac{a_3}{a_2} \quad \text{Common ratio } r = \frac{a_{n+1}}{a_n}$$

$$r = \frac{\frac{15}{4}}{\frac{15}{2}} \quad \text{Substitute.}$$

$$r = \frac{15}{4} \div \frac{15}{2} \quad \text{Write as a division problem.}$$

$$r = \frac{15}{4} \cdot \frac{2}{15} \quad \text{Definition of division}$$

$$r = \frac{1}{2} \quad \text{Multiply. Write in lowest terms.}$$

Any other two successive terms could have been used to find r . Additional terms of the sequence can be found by multiplying each successive term by $\frac{1}{2}$. **NOW TRY**

OBJECTIVE 2 Find the general term of a geometric sequence.

The general term a_n of a geometric sequence a_1, a_2, a_3, \dots is expressed in terms of a_1 and r by writing the first few terms.

$$a_1, \quad a_2 = a_1r, \quad a_3 = a_1r^2, \quad a_4 = a_1r^3, \dots, \quad a_n = a_1r^{n-1}.$$

General Term of a Geometric Sequence

The general term a_n of the geometric sequence with first term a_1 and common ratio r is given by the following.

$$a_n = a_1r^{n-1}$$

**NOW TRY EXERCISE 2**

Determine an expression for the general term of the geometric sequence.

$$\frac{1}{4}, -1, 4, -16, 64, \dots$$

EXAMPLE 2 Finding the General Term of a Geometric Sequence

Determine an expression for the general term of the geometric sequence.

$$15, \frac{15}{2}, \frac{15}{4}, \frac{15}{8}, \dots \quad \text{See Example 1.}$$

The first term is $a_1 = 15$ and the common ratio is $r = \frac{1}{2}$.

$$a_n = a_1r^{n-1} \quad \text{Formula for the general term } a_n$$

$$a_n = 15\left(\frac{1}{2}\right)^{n-1} \quad \text{Let } a_1 = 15 \text{ and } r = \frac{1}{2}.$$

It is not possible to simplify further because the exponent must be applied before the multiplication can be done. **NOW TRY**

NOW TRY ANSWERS

1. $r = -4$
2. $a_n = \frac{1}{4}(-4)^{n-1}$


NOW TRY EXERCISE 3

Find the indicated term for each geometric sequence.

(a) $a_1 = 3, r = -2; a_8$

(b) $10, 2, \frac{2}{5}, \frac{2}{25}, \dots; a_7$

OBJECTIVE 3 Find any specified term of a geometric sequence.

EXAMPLE 3 Finding Specified Terms in Geometric Sequences

Find the indicated term for each geometric sequence.

(a) $a_1 = 4, r = -3; a_6$

$$a_n = a_1 r^{n-1} \quad \text{Formula for the general term } a_n$$

$$a_6 = a_1 \cdot r^{6-1} \quad \text{Let } n = 6.$$

$$a_6 = 4(-3)^5 \quad \text{Let } a_1 = 4, r = -3.$$

$$a_6 = -972 \quad \text{Evaluate } (-3)^5 \text{ and then multiply.}$$

(b) $\frac{3}{4}, \frac{3}{8}, \frac{3}{16}, \dots; a_7$

$$a_7 = \frac{3}{4} \left(\frac{1}{2} \right)^{7-1} \quad \text{Let } a_1 = \frac{3}{4}, r = \frac{1}{2}, n = 7.$$

$$a_7 = \frac{3}{4} \left(\frac{1}{64} \right) \quad \text{Apply the exponent.}$$

$$a_7 = \frac{3}{256} \quad \text{Multiply.}$$

NOW TRY 


NOW TRY EXERCISE 4

Write the first five terms of the geometric sequence having first term 25 and common ratio $-\frac{1}{5}$.

EXAMPLE 4 Writing the Terms of a Geometric Sequence

Write the first five terms of the geometric sequence having first term 5 and common ratio $\frac{1}{2}$.

$$a_1 = 5, \quad a_2 = 5 \left(\frac{1}{2} \right) = \frac{5}{2}, \quad a_3 = 5 \left(\frac{1}{2} \right)^2 = \frac{5}{4}, \quad \text{Use } a_n = a_1 r^{n-1}, \text{ with}$$

$$a_4 = 5 \left(\frac{1}{2} \right)^3 = \frac{5}{8}, \quad a_5 = 5 \left(\frac{1}{2} \right)^4 = \frac{5}{16} \quad \text{with } a_1 = 5, r = \frac{1}{2}, \text{ and}$$

$$n = 1, 2, 3, 4, 5.$$

NOW TRY 

OBJECTIVE 4 Find the sum of a specified number of terms of a geometric sequence.

It is convenient to have a formula for the sum S_n of the first n terms of a geometric sequence. We can develop such a formula by first writing out S_n .

$$S_n = a_1 + a_1 r + a_1 r^2 + a_1 r^3 + \dots + a_1 r^{n-1}$$

Next, we multiply both sides by $-r$.

$$-rS_n = -a_1 r - a_1 r^2 - a_1 r^3 - a_1 r^4 - \dots - a_1 r^n$$

Now add.

$$S_n = a_1 + a_1 r + a_1 r^2 + a_1 r^3 + \dots + a_1 r^{n-1}$$

$$-rS_n = -a_1 r - a_1 r^2 - a_1 r^3 - \dots - a_1 r^{n-1} - a_1 r^n$$

$$S_n - rS_n = a_1 \qquad \qquad \qquad -a_1 r^n$$

$$S_n(1 - r) = a_1 - a_1 r^n \quad \text{Factor on the left.}$$

$$S_n = \frac{a_1(1 - r^n)}{1 - r} \quad \text{Factor on the right. Divide each side by } 1 - r.$$

NOW TRY ANSWERS

3. (a) -384 (b) $\frac{2}{3125}$

4. $25, -5, 1, -\frac{1}{5}, \frac{1}{25}$

Sum of the First n Terms of a Geometric Sequence

The sum S_n of the first n terms of the geometric sequence having first term a_1 and common ratio r is given by the following.

$$S_n = \frac{a_1(1 - r^n)}{1 - r} \quad (\text{where } r \neq 1)$$

If $r = 1$, then $S_n = a_1 + a_1 + a_1 + \cdots + a_1 = na_1$.

Multiplying the formula for S_n by $\frac{-1}{-1}$ gives an alternative form.

$$S_n = \frac{a_1(1 - r^n)}{1 - r} \cdot \frac{-1}{-1} = \frac{a_1(r^n - 1)}{r - 1} \quad \text{Alternative form}$$

NOW TRY EXERCISE 5

Evaluate the sum of the first six terms of the geometric sequence with first term 4 and common ratio 2.

EXAMPLE 5 Finding the Sum of the First n Terms of a Geometric Sequence

Evaluate the sum of the first six terms of the geometric sequence having first term -2 and common ratio 3.

$$S_n = \frac{a_1(1 - r^n)}{1 - r} \quad \text{Formula for } S_n$$

$$S_6 = \frac{-2(1 - 3^6)}{1 - 3} \quad \text{Let } n = 6, a_1 = -2, r = 3.$$

$$S_6 = \frac{-2(1 - 729)}{-2} \quad \text{Evaluate } 3^6. \text{ Subtract in the denominator.}$$

$$S_6 = -728 \quad \text{Simplify.}$$

NOW TRY 

A series of the form

$$\sum_{i=1}^n a \cdot b^i$$

represents the sum of the first n terms of a geometric sequence having first term $a_1 = a \cdot b^1 = ab$ and common ratio b . The next example illustrates this form.

NOW TRY EXERCISE 6

Evaluate $\sum_{i=1}^5 8\left(\frac{1}{2}\right)^i$.

EXAMPLE 6 Using the Formula for S_n to Evaluate a Summation

Evaluate $\sum_{i=1}^4 3 \cdot 2^i$.

Because the series is in the form $\sum_{i=1}^n a \cdot b^i$, it represents the sum of the first n terms of the geometric sequence with $a_1 = a \cdot b^1 = 3 \cdot 2^1 = 6$ and $r = b = 2$.

$$S_n = \frac{a_1(1 - r^n)}{1 - r} \quad \text{Formula for } S_n$$

$$S_4 = \frac{6(1 - 2^4)}{1 - 2} \quad \text{Let } n = 4, a_1 = 6, r = 2.$$

$$S_4 = \frac{6(1 - 16)}{-1} \quad \text{Evaluate } 2^4. \text{ Subtract in the denominator.}$$

$$S_4 = 90 \quad \text{Simplify.}$$

NOW TRY 

NOW TRY ANSWERS

5. 252

6. $\frac{31}{4}$, or 7.75




FIGURE 2

FIGURE 2 shows how a graphing calculator can store the terms in a list and then find the sum of these terms. The figure supports the result of **Example 6**.

OBJECTIVE 5 Apply the formula for the future value of an ordinary annuity.

A sequence of equal payments made over equal periods is an **annuity**. If the payments are made at the end of the period, and if the frequency of payments is the same as the frequency of compounding, the annuity is an **ordinary annuity**. The time between payments is the **payment period**, and the time from the beginning of the first payment period to the end of the last is the **term of the annuity**. The **future value of the annuity**, the final sum on deposit, is defined as the sum of the compound amounts of all the payments, compounded to the end of the term.

We state the following formula without proof.

Future Value of an Ordinary Annuity

The future value of an ordinary annuity is given by

$$S = R \left[\frac{(1 + i)^n - 1}{i} \right],$$

where S is the future value,
 R is the payment at the end of each period,
 i is the interest rate per period,
 and n is the number of periods.

NOW TRY EXERCISE 7

Work each problem. Give answers to the nearest cent.

- (a) Billy deposits \$600 at the end of each year into an account paying 2% per yr, compounded annually. Find the total amount on deposit after 18 yr.
- (b) How much will be in Billy's account after 18 yr if he deposits \$100 at the end of each month at 1.5% interest compounded monthly?

NOW TRY ANSWERS

7. (a) \$12,847.39
 (b) \$24,779.49

EXAMPLE 7 Applying the Formula for Future Value of an Annuity

Work each problem. Give answers to the nearest cent.

- (a) Igor is an athlete who believes that his playing career will last 7 yr. He deposits \$22,000 at the end of each year for 7 yr in an account paying 2% compounded annually. How much will he have on deposit after 7 yr?
 Igor's payments form an ordinary annuity with $R = 22,000$, $n = 7$, and $i = 0.02$.

$$S = 22,000 \left[\frac{(1 + 0.02)^7 - 1}{0.02} \right] \quad \text{Substitute into the formula for } S.$$

$$S = 163,554.23 \quad \text{Evaluate with a calculator.}$$

The future value of this annuity is \$163,554.23.

- (b) Amy has decided to deposit \$200 at the end of each month in an account that pays interest of 2.8% compounded monthly for retirement in 20 yr. How much will be in the account at that time?
 Because the interest is compounded monthly, $i = \frac{0.028}{12}$. Also, $R = 200$ and $n = 12(20)$. Substitute into the formula for S to find the future value.

$$S = 200 \left[\frac{\left(1 + \frac{0.028}{12}\right)^{12(20)} - 1}{\frac{0.028}{12}} \right] = 64,245.50, \quad \text{or} \quad \$64,245.50$$

OBJECTIVE 6 Find the sum of an infinite number of terms of a geometric sequence.

Consider an infinite geometric sequence such as

$$\frac{1}{3}, \frac{1}{6}, \frac{1}{12}, \frac{1}{24}, \frac{1}{48}, \dots, \quad \text{with first term } \frac{1}{3} \text{ and common ratio } \frac{1}{2}.$$

The sum of the first two terms is

$$S_2 = \frac{1}{3} + \frac{1}{6} = \frac{1}{2}.$$

In a similar manner, we can find additional “partial sums.”

$$S_3 = S_2 + \frac{1}{12} = \frac{1}{2} + \frac{1}{12} = \frac{7}{12} \approx 0.583$$

$$S_4 = S_3 + \frac{1}{24} = \frac{7}{12} + \frac{1}{24} = \frac{15}{24} = 0.625$$

$$S_5 = \frac{31}{48} \approx 0.64583$$

$$S_6 = \frac{21}{32} = 0.65625$$

$$S_7 = \frac{127}{192} \approx 0.6614583, \quad \text{and so on.}$$

Each term of the geometric sequence is less than the preceding one, so each additional term is contributing less and less to the partial sum. In decimal form (to the nearest thousandth), the first 7 terms and the 10th term are given in the table.

Term	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_{10}
Value	0.333	0.167	0.083	0.042	0.021	0.010	0.005	0.001

As the table suggests, the value of a term gets closer and closer to 0 as the number of the term increases. To express this idea, we say that as n increases without bound (written $n \rightarrow \infty$), the limit of the term a_n is 0, which is written

$$\lim_{n \rightarrow \infty} a_n = 0.$$

A number that can be defined as the sum of an infinite number of terms of a geometric sequence is found by starting with the expression for the sum of a finite number of terms.

$$S_n = \frac{a_1(1 - r^n)}{1 - r}$$

If $|r| < 1$, then as n increases without bound, the value of r^n gets closer and closer to 0. As r^n approaches 0, $1 - r^n$ approaches $1 - 0 = 1$. S_n approaches the quotient $\frac{a_1}{1 - r}$.

$$\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \frac{a_1(1 - r^n)}{1 - r} = \frac{a_1(1 - 0)}{1 - r} = \frac{a_1}{1 - r}$$

This limit is defined to be the *sum* of the terms of the infinite geometric sequence.

$$a_1 + a_1r + a_1r^2 + a_1r^3 + \dots = \frac{a_1}{1 - r}, \quad \text{if } |r| < 1$$

Sum of the Terms of an Infinite Geometric Sequence

The sum S of the terms of an infinite geometric sequence having first term a_1 and common ratio r , where $|r| < 1$, is given by the following.

$$S = \frac{a_1}{1 - r}$$

If $|r| \geq 1$, then the sum does not exist.

Now consider $|r| > 1$. For example, suppose the sequence is

$$6, 12, 24, \dots, 3(2)^n, \dots \quad \text{Here, } a_1 = 6 \text{ and } r = 2.$$

In this kind of sequence, as n increases, the value of r^n also increases and so does the sum S_n . Each new term adds a greater and greater amount to the sum, so there is no limit to the value of S_n . The sum S does not exist. Recall that if $r = 1$, then

$$S_n = a_1 + a_1 + a_1 + \dots + a_1 = na_1.$$

NOW TRY EXERCISE 8

Evaluate the sum of the terms of the infinite geometric sequence having $a_1 = -4$ and $r = \frac{2}{3}$.

EXAMPLE 8 Finding the Sum of the Terms of an Infinite Geometric Sequence

Evaluate the sum of the terms of the infinite geometric sequence having $a_1 = 3$ and $r = -\frac{1}{3}$.

$$S = \frac{a_1}{1 - r} \quad \text{Infinite sum formula}$$

$$S = \frac{3}{1 - \left(-\frac{1}{3}\right)} \quad \text{Let } a_1 = 3, r = -\frac{1}{3}.$$

$$S = \frac{3}{\frac{4}{3}} \quad \text{Subtract in the denominator; } 1 - \left(-\frac{1}{3}\right) = 1 + \frac{1}{3} = \frac{3}{3} + \frac{1}{3} = \frac{4}{3}.$$

$$S = 3 \div \frac{4}{3} \quad \text{Write as a division problem.}$$

$$S = 3 \cdot \frac{3}{4} \quad \text{Definition of division}$$

$$S = \frac{9}{4} \quad \text{Multiply.}$$

NOW TRY 

In summation notation, the sum of an infinite geometric sequence is written as

$$\sum_{i=1}^{\infty} a_i.$$

For instance, the sum in **Example 8** would be written

$$\sum_{i=1}^{\infty} 3\left(-\frac{1}{3}\right)^{i-1}.$$

NOW TRY ANSWER

8. -12


NOW TRY EXERCISE 9

Evaluate $\sum_{i=1}^{\infty} \left(\frac{5}{8}\right)\left(\frac{3}{4}\right)^i$.

EXAMPLE 9 Evaluating an Infinite Geometric Series

Evaluate $\sum_{i=1}^{\infty} \left(\frac{2}{3}\right)\left(\frac{1}{2}\right)^i$.

This is the infinite geometric series

$$\frac{1}{3} + \frac{1}{6} + \frac{1}{12} + \cdots,$$

 with $a_1 = \frac{1}{3}$ and $r = \frac{1}{2}$. Because $|r| < 1$, we find the sum as follows.

$$S = \frac{a_1}{1 - r} \quad \text{Infinite sum formula}$$

$$S = \frac{\frac{1}{3}}{1 - \frac{1}{2}} \quad \text{Let } a_1 = \frac{1}{3}, r = \frac{1}{2}.$$

$$S = \frac{\frac{1}{3}}{\frac{1}{2}} \quad \text{Subtract in the denominator; } 1 - \frac{1}{2} = \frac{2}{2} - \frac{1}{2} = \frac{1}{2}.$$

$$S = \frac{2}{3} \quad \text{Divide; } \frac{\frac{1}{3}}{\frac{1}{2}} = \frac{1}{3} \cdot \frac{2}{1} = \frac{2}{3}.$$


NOW TRY 
NOW TRY ANSWER

9. $\frac{15}{8}$

11.3 Exercises

 FOR
EXTRA
HELP

MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Fill in each blank with the correct response.

- In a geometric sequence, if any term after the first is divided by the term that precedes it, the result is the common _____ of the sequence.
- For the geometric sequence having $a_n = (-2)^n$, the term $a_5 =$ _____.
- The sum of the first five terms of the geometric sequence 1, 2, 4, ... is _____.
- The number of terms in the geometric sequence 1, 2, 4, ..., 2048 is _____.

Concept Check Fill in the blanks to complete the terms of each geometric sequence.

- $\frac{1}{3}, -\frac{1}{9}, \frac{1}{27},$ _____, _____, _____
- _____, _____, 1, 5, 25
- 7, _____, 28, _____, 112
- 2, _____, _____, -54, _____

If the given sequence is geometric, find the common ratio r . If the sequence is not geometric, say so. See Example 1.

- 4, 8, 16, 32, ...
- 5, 15, 45, 135, ...
- $\frac{1}{3}, \frac{2}{3}, \frac{3}{3}, \frac{4}{3}, \dots$
- $\frac{5}{7}, \frac{8}{7}, \frac{11}{7}, 2, \dots$
- 1, -3, 9, -27, 81, ...
- 2, -8, 32, -128, ...
- 1, $-\frac{1}{2}, \frac{1}{4}, -\frac{1}{8}, \dots$
- $\frac{2}{3}, -\frac{2}{15}, \frac{2}{75}, -\frac{2}{375}, \dots$

Determine an expression for the general term of each geometric sequence. See Example 2.

17. $-5, -10, -20, \dots$ 18. $-2, -6, -18, \dots$ 19. $-2, \frac{2}{3}, -\frac{2}{9}, \dots$
 20. $-3, \frac{3}{2}, -\frac{3}{4}, \dots$ 21. $10, -2, \frac{2}{5}, \dots$ 22. $8, -2, \frac{1}{2}, \dots$

Find the indicated term for each geometric sequence. See Example 3.

23. $a_1 = 2, r = 5; a_{10}$ 24. $a_1 = 1, r = 3; a_{15}$
 25. $\frac{1}{2}, \frac{1}{6}, \frac{1}{18}, \dots; a_{12}$ 26. $\frac{2}{3}, \frac{1}{3}, \frac{1}{6}, \dots; a_{18}$
 27. $a_2 = 18, a_5 = -486; a_7$ 28. $a_5 = 48, a_8 = -384; a_{10}$

Write the first five terms of each geometric sequence. See Example 4.

29. $a_1 = 2, r = 3$ 30. $a_1 = 4, r = 2$ 31. $a_1 = 5, r = -\frac{1}{5}$
 32. $a_1 = 6, r = -\frac{1}{3}$ 33. $a_1 = -4, r = 0.5$ 34. $a_1 = -40, r = 0.25$

Evaluate the sum of the terms of each geometric sequence. In Exercises 39–44, give answers to the nearest thousandth. See Examples 5 and 6.

35. $a_1 = -3, r = 4; \text{ Find } S_{10}.$ 36. $a_1 = -5, r = 7; \text{ Find } S_9.$
 37. $\frac{1}{3}, \frac{1}{9}, \frac{1}{27}, \frac{1}{81}, \frac{1}{243}$ 38. $\frac{1}{4}, \frac{1}{16}, \frac{1}{64}, \frac{1}{256}, \frac{1}{1024}$
 39. $-\frac{4}{3}, -\frac{4}{9}, -\frac{4}{27}, -\frac{4}{81}, -\frac{4}{243}, -\frac{4}{729}$ 40. $\frac{5}{16}, -\frac{5}{32}, \frac{5}{64}, -\frac{5}{128}, \frac{5}{256}$
 41. $\sum_{i=1}^7 4\left(\frac{2}{5}\right)^i$ 42. $\sum_{i=1}^8 5\left(\frac{2}{3}\right)^i$
 43. $\sum_{i=1}^{10} (-2)\left(\frac{3}{5}\right)^i$ 44. $\sum_{i=1}^6 (-2)\left(-\frac{1}{2}\right)^i$

Solve each problem. Round answers to the nearest cent. See Example 7.

45. A father opened a savings account for his daughter on her first birthday, depositing \$1000. Each year on her birthday he deposits another \$1000, making the last deposit on her 21st birthday. If the account pays 1.5% interest compounded annually, how much is in the account at the end of the day on the daughter's 21st birthday?



46. A teacher puts \$1000 in a retirement account at the end of each quarter ($\frac{1}{4}$ of a year) for 15 yr. If the account pays 2.2% annual interest compounded quarterly, how much will be in the account at that time?

47. At the end of each quarter, a 50-year-old woman puts \$1200 in a retirement account that pays 2% interest compounded quarterly. When she reaches age 60, she withdraws the entire amount and places it in a fund that pays 1% annual interest compounded monthly. From then on, she deposits \$300 in the fund at the end of each month. How much is in the account when she reaches age 65?
48. At the end of each quarter, a 45-year-old man puts \$1500 in a retirement account that pays 1.5% interest compounded quarterly. When he reaches age 55, he withdraws the entire amount and places it in a fund that pays 0.75% annual interest compounded monthly. From then on, he deposits \$400 in the fund at the end of each month. How much is in the account when he reaches age 60?

Find the sum, if it exists, of the terms of each infinite geometric sequence. See Examples 8 and 9.

49. $a_1 = 6, r = \frac{1}{3}$

50. $a_1 = 10, r = \frac{1}{5}$

51. $a_1 = 1000, r = -\frac{1}{10}$

52. $a_1 = 8800, r = -\frac{3}{5}$

53. $\sum_{i=1}^{\infty} \frac{9}{8} \left(-\frac{2}{3}\right)^i$


54. $\sum_{i=1}^{\infty} \frac{3}{5} \left(\frac{5}{6}\right)^i$

55. $\sum_{i=1}^{\infty} \frac{12}{5} \left(\frac{5}{4}\right)^i$


56. $\sum_{i=1}^{\infty} \left(-\frac{16}{3}\right) \left(-\frac{9}{8}\right)^i$

Extending Skills Solve each application.

57. When dropped from a certain height, a ball rebounds to $\frac{3}{5}$ of the original height. How high will the ball rebound after the fourth bounce if it was dropped from a height of 10 ft? Round to the nearest tenth.



58. A fully wound yo-yo has a string 40 in. long. It is allowed to drop, and on its first rebound it returns to a height 15 in. lower than its original height. Assuming that this “rebound ratio” remains constant until the yo-yo comes to rest, how far does it travel on its third trip up the string? Round to the nearest tenth.



59. A ball is dropped from a height of 20 m, and on each bounce it returns to $\frac{3}{4}$ of its previous height. How far will the ball travel before it comes to rest? (*Hint:* Consider the sum of two sequences.)
60. A fully wound yo-yo is dropped the length of its 30-in. string. Each time it drops, it returns to $\frac{1}{2}$ of its original height. How far does it travel before it comes to rest? (*Hint:* Consider the sum of two sequences.)
61. A particular substance decays in such a way that it loses half its weight each day. In how many days will 256 g of the substance be reduced to 32 g? How much of the substance is left after 10 days?
62. A tracer dye is injected into a system with an ingestion and an excretion. After 1 hr, $\frac{2}{3}$ of the dye is left. At the end of the second hour, $\frac{2}{3}$ of the remaining dye is left, and so on. If one unit of the dye is injected, how much is left after 6 hr?

- 63.** In a certain community, the consumption of electricity has increased about 6% per yr.
- (a) If the community uses 1.1 billion units of electricity now, how much will it use 5 yr from now? Round to the nearest tenth.
- (b) Find the number of years (to the nearest year) it will take for the consumption to double.
- 64.** A growing community increases its consumption of electricity 2% per yr.
- (a) If the community uses 1.1 billion units of electricity now, how much will it use 5 yr from now? Round to the nearest tenth.
- (b) Find the number of years (to the nearest year) it will take for the consumption to double.
- 65.** A machine depreciates by $\frac{1}{4}$ of its value each year. If it cost \$50,000 new, what is its value after 8 yr?
- 66.** A vehicle depreciates by 20% of its value each year. If it cost \$35,000 new, what is its value after 6 yr?
- 67.** The repeating decimal $0.99999 \dots$ can be written as the sum of the terms of a geometric sequence with $a_1 = 0.9$ and $r = 0.1$.
- $$0.99999 \dots = 0.9 + 0.9(0.1) + 0.9(0.1)^2 + 0.9(0.1)^3 + 0.9(0.1)^4 + 0.9(0.1)^5 + \dots$$
- Because $|0.1| < 1$, this sum can be found from the formula $S = \frac{a_1}{1-r}$. Use this formula to find a more common way of writing the decimal $0.99999 \dots$.
- 68.** If the result of **Exercise 67** seems hard to believe, look at it this way: Use long division to find the repeating decimal representations of $\frac{1}{3}$ and $\frac{2}{3}$. Then line up the decimals vertically and add them. The result must equal

$$\frac{1}{3} + \frac{2}{3} = 1.$$

11.4 The Binomial Theorem

OBJECTIVES

- Expand a binomial raised to a power.
- Find any specified term of the expansion of a binomial.

VOCABULARY

- Pascal's triangle

OBJECTIVE 1 Expand a binomial raised to a power.

Observe the expansion of the expression $(x + y)^n$ for the first six nonnegative integer values of n .

$$(x + y)^0 = 1$$

$$(x + y)^1 = x + y$$

$$(x + y)^2 = x^2 + 2xy + y^2$$

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

$$(x + y)^5 = x^5 + 5x^4y + 10x^3y^2 + 10x^2y^3 + 5xy^4 + y^5$$

Expansions of $(x + y)^n$



By identifying patterns, we can write a general expansion for $(x + y)^n$.

First, if n is a positive integer, each expansion after $(x + y)^0$ begins with x raised to the same power to which the binomial is raised. That is, the expansion of $(x + y)^1$ has a first term of x^1 , the expansion of $(x + y)^2$ has a first term of x^2 , and so on. Also, the last term in each expansion is y to this same power, so the expansion of

$$(x + y)^n$$

should begin with the term x^n and end with the term y^n .

The exponents on x decrease by 1 in each term after the first, while the exponents on y , beginning with y in the second term, increase by 1 in each succeeding term. Thus, the *variables* in the expansion of $(x + y)^n$ have the following pattern.

$$x^n, x^{n-1}y, x^{n-2}y^2, x^{n-3}y^3, \dots, xy^{n-1}, y^n$$

This pattern suggests that the sum of the exponents on x and y in each term is n . For example, in the third term shown, the variable part is $x^{n-2}y^2$ and the sum of the exponents is $n - 2 + 2 = n$.

Now examine the pattern for the *coefficients* of the terms of the preceding expansions. Writing the coefficients alone in a triangular pattern gives **Pascal's triangle**, named in honor of the 17th-century mathematician Blaise Pascal.



Blaise Pascal (1623–1662)

Pascal's Triangle

						Row
1						0
1		1				1
1	2	1				2
1	3	3	1			3
1	4	6	4	1		4
1	5	10	10	5	1	5

In this triangle, the first and last terms of each row are 1. Each number in the interior of the triangle is the sum of the two numbers just above it (one to the right and one to the left). For example, in row 4 of the triangle, 4 is the sum of 1 and 3, 6 is the sum of 3 and 3, and so on.

To obtain the coefficients for $(x + y)^6$, we need to attach row 6 to the table by starting and ending with 1, and adding pairs of numbers from row 5.

$$1 \quad 6 \quad 15 \quad 20 \quad 15 \quad 6 \quad 1 \quad \text{Row 6}$$

We then use these coefficients to expand $(x + y)^6$.

$$(x + y)^6 = x^6 + 6x^5y + 15x^4y^2 + 20x^3y^3 + 15x^2y^4 + 6xy^5 + y^6$$

Although it is possible to use Pascal's triangle to find the coefficients in $(x + y)^n$ for any positive integer value of n , it is impractical for large values of n . A more efficient way to determine these coefficients uses the symbol $n!$ (read "*n factorial*").

n Factorial ($n!$)

For any positive integer n ,

$$n! = n(n - 1)(n - 2)(n - 3) \cdots (3)(2)(1).$$

By definition, $0! = 1$.

Mode	Value
3!	6
5!	120
0!	1

FIGURE 3

NOW TRY
EXERCISE 1
Evaluate $7!$.

NOW TRY ANSWER
1. 5040

EXAMPLE 1 Evaluating Factorials

Evaluate each factorial.

(a) $3! = 3 \cdot 2 \cdot 1 = 6$ (b) $5! = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 120$ (c) $0! = 1$

FIGURE 3 shows how a graphing calculator computes factorials.

NOW TRY


NOW TRY EXERCISE 2

Find the value of each expression.

- (a) $\frac{8!}{6!2!}$ (b) $\frac{8!}{5!3!}$
 (c) $\frac{6!}{6!0!}$ (d) $\frac{6!}{5!1!}$

EXAMPLE 2 Evaluating Expressions Involving Factorials

Find the value of each expression.

- (a) $\frac{5!}{4!1!} = \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{(4 \cdot 3 \cdot 2 \cdot 1)(1)} = 5$
 (b) $\frac{5!}{3!2!} = \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{(3 \cdot 2 \cdot 1)(2 \cdot 1)} = \frac{5 \cdot 4}{2 \cdot 1} = 10$
 (c) $\frac{6!}{3!3!} = \frac{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{(3 \cdot 2 \cdot 1)(3 \cdot 2 \cdot 1)} = \frac{6 \cdot 5 \cdot 4}{3 \cdot 2 \cdot 1} = 20$
 (d) $\frac{4!}{4!0!} = \frac{4 \cdot 3 \cdot 2 \cdot 1}{(4 \cdot 3 \cdot 2 \cdot 1)(1)} = 1$

NOW TRY 

Now look again at the coefficients of the expansion

$$(x + y)^5 = x^5 + 5x^4y + 10x^3y^2 + 10x^2y^3 + 5xy^4 + y^5.$$

The coefficient of the second term is **5**, and the exponents on the variables in that term are 4 and 1. From **Example 2(a)**, $\frac{5!}{4!1!} = 5$. The coefficient of the third term is **10**, and the exponents are 3 and 2. From **Example 2(b)**, $\frac{5!}{3!2!} = 10$. Similar results are true for the remaining terms. The first term can be written as $1x^5y^0$, and the last term can be written as $1x^0y^5$. Then the coefficient of the first term should be $\frac{5!}{5!0!} = 1$, and the coefficient of the last term would be $\frac{5!}{0!5!} = 1$. (This is why $0!$ is *defined* to be 1.)

The coefficient of a term in $(x + y)^n$ in which the variable part is $x^r y^{n-r}$ is

$$\frac{n!}{r!(n-r)!} \cdot \quad \text{This is the binomial coefficient.}$$

The **binomial coefficient** $\frac{n!}{r!(n-r)!}$ is often represented by the symbol ${}_n C_r$. This notation comes from the fact that if we choose *combinations* of n things taken r at a time, the result is given by that expression. We read ${}_n C_r$ as “*combinations of n things taken r at a time.*” Another common representation is $\binom{n}{r}$.

Formula for the Binomial Coefficient ${}_n C_r$

For nonnegative integers n and r , where $r \leq n$, ${}_n C_r$ is defined as follows.

$${}_n C_r = \frac{n!}{r!(n-r)!}$$

EXAMPLE 3 Evaluating Binomial Coefficients

Evaluate each binomial coefficient.

- (a) ${}_5 C_4 = \frac{5!}{4!(5-4)!}$ Let $n = 5, r = 4$.
 $= \frac{5!}{4!1!}$ Subtract.
 $= \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{4 \cdot 3 \cdot 2 \cdot 1 \cdot 1}$ Definition of n factorial
 $= 5$ Lowest terms

NOW TRY ANSWERS

2. (a) 28 (b) 56 (c) 1 (d) 6

NOW TRY EXERCISE 3
Evaluate ${}_7C_2$.

NORMAL FLOAT AUTO REAL RADIAN HP	
${}_5C_4$	5
${}_5C_3$	10
${}_6C_3$	20

FIGURE 4

$$(b) {}_5C_3 = \frac{5!}{3!(5-3)!} = \frac{5!}{3!2!} = \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{3 \cdot 2 \cdot 1 \cdot 2 \cdot 1} = 10$$

$$(c) {}_6C_3 = \frac{6!}{3!(6-3)!} = \frac{6!}{3!3!} = \frac{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{3 \cdot 2 \cdot 1 \cdot 3 \cdot 2 \cdot 1} = 20$$

Binomial coefficients will always be whole numbers.

FIGURE 4 displays the binomial coefficients computed here.

NOW TRY 

Our observations about the expansion of $(x + y)^n$ are summarized as follows.

- There are $n + 1$ terms in the expansion.
- The first term is x^n , and the last term is y^n .
- In each succeeding term, the exponent on x decreases by 1 and the exponent on y increases by 1.
- The sum of the exponents on x and y in any term is n .
- The coefficient of the term with $x^r y^{n-r}$ or $x^{n-r} y^r$ is ${}_n C_r$.

We now state the **binomial theorem**, or the **general binomial expansion**.

Binomial Theorem

For any positive integer n , $(x + y)^n$ is expanded as follows.

$$\begin{aligned} (x + y)^n &= x^n + \frac{n!}{1!(n-1)!} x^{n-1}y + \frac{n!}{2!(n-2)!} x^{n-2}y^2 \\ &\quad + \frac{n!}{3!(n-3)!} x^{n-3}y^3 + \cdots + \frac{n!}{(n-1)!1!} xy^{n-1} + y^n \end{aligned}$$

The binomial theorem can be written using summation notation as follows.

$$(x + y)^n = \sum_{r=0}^n \frac{n!}{r!(n-r)!} x^{n-r} y^r$$

NOW TRY EXERCISE 4
Expand $(a + 3b)^5$.

EXAMPLE 4 Using the Binomial Theorem

Expand $(2m + 3)^4$.

$$\begin{aligned} &(2m + 3)^4 \\ &= (2m)^4 + \frac{4!}{1!3!} (2m)^3(3) + \frac{4!}{2!2!} (2m)^2(3)^2 + \frac{4!}{3!1!} (2m)(3)^3 + 3^4 \\ &= 16m^4 + 4(8m^3)(3) + 6(4m^2)(9) + 4(2m)(27) + 81 \\ &= 16m^4 + 96m^3 + 216m^2 + 216m + 81 \end{aligned}$$

Remember:
 $(ab)^m = a^m b^m$

NOW TRY 

NOTE The binomial coefficients for the first and last terms in **Example 4** are both 1. These values can be taken from Pascal's triangle or computed as follows.

$$\frac{4!}{0!4!} = 1 \quad \text{Binomial coefficient of the first term}$$

$$\frac{4!}{4!0!} = 1 \quad \text{Binomial coefficient of the last term}$$

NOW TRY ANSWERS

3. 21

4. $a^5 + 15a^4b + 90a^3b^2 + 270a^2b^3 + 405ab^4 + 243b^5$


NOW TRY EXERCISE 5
Expand $\left(\frac{x}{3} - 2y\right)^4$.**EXAMPLE 5** Using the Binomial TheoremExpand $\left(a - \frac{b}{2}\right)^5$.

$$\begin{aligned} & \left(a - \frac{b}{2}\right)^5 \\ &= a^5 + \frac{5!}{1!4!}a^4\left(-\frac{b}{2}\right) + \frac{5!}{2!3!}a^3\left(-\frac{b}{2}\right)^2 + \frac{5!}{3!2!}a^2\left(-\frac{b}{2}\right)^3 \\ & \quad + \frac{5!}{4!1!}a\left(-\frac{b}{2}\right)^4 + \left(-\frac{b}{2}\right)^5 \\ &= a^5 + 5a^4\left(-\frac{b}{2}\right) + 10a^3\left(\frac{b^2}{4}\right) + 10a^2\left(-\frac{b^3}{8}\right) \\ & \quad + 5a\left(\frac{b^4}{16}\right) + \left(-\frac{b^5}{32}\right) \end{aligned}$$

Notice that signs alternate positive and negative.

$$= a^5 - \frac{5}{2}a^4b + \frac{5}{2}a^3b^2 - \frac{5}{4}a^2b^3 + \frac{5}{16}ab^4 - \frac{1}{32}b^5$$

NOW TRY 

CAUTION When the binomial is a *difference* of two terms, as in **Example 5**, the signs of the terms in the expansion will alternate.

- Those terms with odd exponents on the second variable expression $\left(-\frac{b}{2}\right)$ in **Example 5** will be negative.
- Those terms with even exponents on the second variable expression will be positive.

OBJECTIVE 2 Find any specified term of the expansion of a binomial.

Any single term of a binomial expansion can be determined without writing out the whole expansion. For example, if $n \geq 10$, then the 10th term of $(x + y)^n$ has y raised to the ninth power (because y has the power of 1 in the second term, the power of 2 in the third term, and so on).

The exponents on x and y in any term must have a sum of n , so the exponent on x in the 10th term is $n - 9$. The quantities 9 and $n - 9$ determine the factorials in the denominator of the coefficient. Thus, the 10th term of $(x + y)^n$ is

$$\frac{n!}{9!(n-9)!}x^{n-9}y^9.$$

kth Term of the Binomial Expansion

If $n \geq k - 1$, then the k th term of the expansion of $(x + y)^n$ is given by the following.

$$\frac{n!}{(k-1)![n-(k-1)]!}x^{n-(k-1)}y^{k-1}$$

NOW TRY ANSWER

5. $\frac{x^4}{81} - \frac{8x^3y}{27} + \frac{8x^2y^2}{3} - \frac{32xy^3}{3} + 16y^4$


NOW TRY EXERCISE 6

Find the sixth term of the expansion of $(2m - n^2)^8$.

EXAMPLE 6 Finding a Single Term of a Binomial Expansion

Find the fourth term of the expansion of $(a + 2b)^{10}$.

In the fourth term, $2b$ has an exponent of $4 - 1 = 3$, and a has an exponent of $10 - 3 = 7$. The fourth term is determined as follows.

$$\begin{aligned} \frac{10!}{3!7!}(a^7)(2b)^3 & \quad \text{Parentheses MUST be used with } 2b. \quad \text{Let } n = 10, x = a, y = 2b, \text{ and } k = 4. \\ &= 120(a^7)(2^3b^3) \quad \text{Simplify the factorials; } (ab)^m = a^mb^m \\ &= 120a^7(8b^3) \quad \text{Simplify.} \\ &= 960a^7b^3 \quad \text{Multiply.} \end{aligned}$$

NOW TRY ANSWER


6. $-448m^3n^{10}$

NOW TRY 

11.4 Exercises

FOR EXTRA HELP

MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Fill in each blank with the correct response.

- In each row of Pascal's triangle, the first and last terms are _____, and each number in the interior of the triangle is the _____ of the two numbers just above it (one to the right and one to the left).
- The complete row of Pascal's triangle that begins with the terms 1, 4, is _____.
- The first term in the expansion of $(x + y)^3$ is _____.
- The last term in the expansion of $(x - y)^4$ is _____.
- The value of $3!$ is _____.
- The value of $0!$ is _____.
- For any nonnegative integer n , the binomial coefficient ${}_nC_0$ is equal to _____.
- For any nonnegative integer n , the binomial coefficient ${}_nC_n$ is equal to _____.

Evaluate each expression. See Examples 1–3.

- | | | | |
|-----------------------|-----------------------|-----------------------|-----------------------|
| 9. $6!$ | 10. $4!$ | 11. $8!$ | 12. $9!$ |
| 13. $\frac{6!}{4!2!}$ | 14. $\frac{7!}{3!4!}$ | 15. $\frac{4!}{0!4!}$ | 16. $\frac{5!}{5!0!}$ |
| 17. $4! \cdot 5$ | 18. $6! \cdot 7$ | 19. ${}_6C_2$ | 20. ${}_7C_4$ |
| 21. ${}_{13}C_{11}$ | 22. ${}_{13}C_2$ | 23. ${}_{10}C_7$ | 24. ${}_{10}C_3$ |

Use the binomial theorem to expand each binomial. See Examples 4 and 5.

- | | | |
|--------------------------------------|---------------------------------------|----------------------|
| 25. $(m + n)^4$ | 26. $(x + r)^5$ | 27. $(a - b)^5$ |
| 28. $(p - q)^4$ | 29. $(2x + 3)^3$ | 30. $(4x + 2)^3$ |
| 31. $\left(\frac{x}{2} - y\right)^4$ | 32. $\left(\frac{x}{3} - 2y\right)^5$ | 33. $(x^2 + 1)^4$ |
| 34. $(y^3 + 2)^4$ | 35. $(3x^2 - y^2)^3$ | 36. $(2p^2 - q^2)^3$ |

Write the first four terms of each binomial expansion. See Examples 4 and 5.

37. $(r + 2s)^{12}$ 38. $(m + 3n)^{20}$ 39. $(3x - y)^{14}$
 40. $(2p - 3q)^{11}$ 41. $(t^2 + u^2)^{10}$ 42. $(x^2 + y^2)^{15}$

Find the indicated term of each binomial expansion. See Example 6.

43. $(2m + n)^{10}$; fourth term 44. $(a + 3b)^{12}$; fifth term
 45. $\left(x + \frac{y}{2}\right)^8$; seventh term 46. $\left(a + \frac{b}{3}\right)^{15}$; eighth term
 47. $(k - 1)^9$; third term 48. $(r - 4)^{11}$; fourth term
 49. The middle term of $(x^2 + 2y)^6$ 50. The middle term of $(m^3 + 2y)^8$
 51. The term with x^9y^4 in $(3x^3 - 4y^2)^5$ 52. The term with x^8y^2 in $(2x^2 + 3y)^6$

Chapter 11 Summary

Key Terms

11.1 infinite sequence finite sequence terms of a sequence general term series summation notation	index of summation arithmetic mean (average)	11.3 geometric sequence (geometric progression) common ratio annuity ordinary annuity payment period	term of an annuity future value of an annuity
	11.2 arithmetic sequence (arithmetic progression) common difference		11.4 Pascal's triangle

New Symbols

a_n n th term of a sequence	S_n sum of first n terms of a sequence	$\sum_{i=1}^{\infty} a_i$ sum of an infinite number of terms	${}_nC_r$ binomial coefficient (combinations of n things taken r at a time)
$\sum_{i=1}^n a_i$ summation notation with index i	$\lim_{n \rightarrow \infty} a_n$ limit of a_n as n increases without bound	$n!$ n factorial	

Test Your Word Power

See how well you have learned the vocabulary in this chapter.

- An **infinite sequence** is
 - the values of a function
 - a function whose domain is the set of positive integers
 - the sum of the terms of a function
 - the average of a group of numbers.
- A **series** is
 - the sum of the terms of a sequence
 - the product of the terms of a sequence
 - the average of the terms of a sequence
 - the function values of a sequence.
- An **arithmetic sequence** is a sequence in which
 - each term after the first is a constant multiple of the preceding term
 - the numbers are written in a triangular array
 - the terms are added
 - each term after the first differs from the preceding term by a common amount.

4. A **geometric sequence** is a sequence in which
- A. each term after the first is a constant multiple of the preceding term
 - B. the numbers are written in a triangular array
 - C. the terms are multiplied
 - D. each term after the first differs from the preceding term by a common amount.
5. The **common difference** is
- A. the average of the terms in a sequence
 - B. the constant multiplier in a geometric sequence
 - C. the difference between any two adjacent terms in an arithmetic sequence
 - D. the sum of the terms of an arithmetic sequence.
6. The **common ratio** is
- A. the average of the terms in a sequence
 - B. the constant multiplier in a geometric sequence
 - C. the difference between any two adjacent terms in an arithmetic sequence
 - D. the product of the terms of a geometric sequence.

ANSWERS

1. B; *Example:* The ordered list of numbers 3, 6, 9, 12, 15, . . . is an infinite sequence. 2. A; *Example:* $3 + 6 + 9 + 12 + 15$, written in summation notation as $\sum_{i=1}^5 3i$, is a series. 3. D; *Example:* The sequence $-3, 2, 7, 12, 17, \dots$ is arithmetic. 4. A; *Example:* The sequence 1, 4, 16, 64, 256, . . . is geometric. 5. C; *Example:* The common difference of the arithmetic sequence $-3, 2, 7, 12, 17, \dots$ is 5, since $2 - (-3) = 5, 7 - 2 = 5$, and so on. 6. B; *Example:* The common ratio of the geometric sequence 1, 4, 16, 64, 256, . . . is 4, since $\frac{4}{1} = \frac{16}{4} = \frac{64}{16} = \frac{256}{64} = 4$.

Quick Review

CONCEPTS	EXAMPLES
<p>11.1 Sequences and Series</p> <p>An infinite sequence is a function with the set of positive integers as domain.</p> <p>A finite sequence is a function with domain of the form $\{1, 2, 3, \dots, n\}$, where n is a positive integer.</p> <p>A series is the indicated sum of the terms of a sequence.</p>	<p>The infinite sequence $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots, \frac{1}{n}$ has general term $a_n = \frac{1}{n}$.</p> <p>The corresponding series is the <i>sum</i></p> $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n}.$
<p>11.2 Arithmetic Sequences</p> <p>Assume that a_1 is the first term, a_n is the nth term, and d is the common difference in an arithmetic sequence.</p> <p>Common Difference</p> $d = a_{n+1} - a_n$ <p>nth Term</p> $a_n = a_1 + (n - 1)d$ <p>Sum of the First n Terms</p> $S_n = \frac{n}{2}(a_1 + a_n)$ <p>or</p> $S_n = \frac{n}{2}[2a_1 + (n - 1)d]$	<p>Consider the following arithmetic sequence.</p> $2, 5, 8, 11, \dots$ <p>$a_1 = 2$ a_1 is the first term.</p> <p>$d = 3$ $d = a_2 - a_1$</p> <p>(Any two successive terms could have been used.)</p> <p>The tenth term is found as follows.</p> $a_{10} = 2 + (10 - 1)(3) \quad \text{Let } a_1 = 2, d = 3, \text{ and } n = 10.$ $a_{10} = 2 + 9 \cdot 3 \quad \text{Subtract inside the parentheses.}$ $a_{10} = 29 \quad \text{Multiply, and then add.}$ <p>The sum of the first ten terms can be found in either of two ways.</p> $S_{10} = \frac{10}{2}(2 + a_{10}) \quad \left \quad S_{10} = \frac{10}{2}[2(2) + (10 - 1)(3)]$ $S_{10} = 5(2 + 29) \quad \left \quad S_{10} = 5(4 + 9 \cdot 3)$ $S_{10} = 5(31) \quad \left \quad S_{10} = 5(4 + 27)$ $S_{10} = 155 \quad \left \quad S_{10} = 5(31)$ $S_{10} = 155 \quad \left \quad S_{10} = 155$

CONCEPTS

11.3 Geometric Sequences

Assume that a_1 is the first term, a_n is the n th term, and r is the common ratio in a geometric sequence.

Common Ratio

$$r = \frac{a_{n+1}}{a_n}$$

 n th Term

$$a_n = a_1 r^{n-1}$$

Sum of the First n Terms

$$S_n = \frac{a_1(1 - r^n)}{1 - r} \quad (\text{where } r \neq 1)$$

Future Value of an Ordinary Annuity

$$S = R \left[\frac{(1 + i)^n - 1}{i} \right],$$

where S is the future value, R is the payment at the end of each period, i is the interest rate per period, and n is the number of periods.

Sum of the Terms of an Infinite Geometric Sequence

$$S = \frac{a_1}{1 - r} \quad (\text{where } |r| < 1)$$

11.4 The Binomial Theorem **n Factorial ($n!$)**

For any positive integer n ,

$$n! = n(n - 1)(n - 2)(n - 3) \cdots (3)(2)(1).$$

By definition, $0! = 1$.

Binomial Coefficient

$${}_n C_r = \frac{n!}{r!(n - r)!} \quad (\text{where } r \leq n)$$

EXAMPLES

Consider the following geometric sequence.

$$1, 2, 4, 8, \dots$$

$$a_1 = 1 \quad a_1 \text{ is the first term.}$$

$$r = 2 \quad r = \frac{a_4}{a_3}$$

(Any two successive terms could have been used.)

The sixth term is found as follows.

$$a_6 = 1(2)^{6-1} \quad \text{Let } a_1 = 1, r = 2, \text{ and } n = 6.$$

$$a_6 = 1(2)^5 \quad \text{Subtract in the exponent.}$$

$$a_6 = 32 \quad \text{Apply the exponent and multiply.}$$

The sum of the first six terms is found as follows.

$$S_6 = \frac{1(1 - 2^6)}{1 - 2} \quad \text{Substitute for } a_1, r, \text{ and } n.$$

$$S_6 = \frac{1 - 64}{-1} \quad \text{Evaluate } 2^6. \text{ Subtract in the denominator.}$$

$$S_6 = 63 \quad \text{Simplify.}$$

If \$5800 is deposited into an ordinary annuity at the end of each quarter for 4 yr and interest is earned at 2.4% compounded quarterly, how much will be in the account at that time?

$$R = \$5800, \quad i = \frac{0.024}{4} = 0.006, \quad n = 4(4) = 16$$

$$S = 5800 \left[\frac{(1 + 0.006)^{16} - 1}{0.006} \right] = \$97,095.24 \quad \text{Nearest cent}$$

Evaluate the sum S of the terms of an infinite geometric sequence with $a_1 = 1$ and $r = \frac{1}{2}$.

$$S = \frac{1}{1 - \frac{1}{2}} = \frac{1}{\frac{1}{2}} = 1 \div \frac{1}{2} = 1 \cdot \frac{2}{1} = 2$$

Evaluate $4!$.

$$4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24$$

Evaluate ${}_5 C_3$.

$${}_5 C_3 = \frac{5!}{3!(5 - 3)!} = \frac{5!}{3!2!} = \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{3 \cdot 2 \cdot 1 \cdot 2 \cdot 1} = 10$$

CONCEPTS
Binomial Theorem

For any positive integer n , $(x + y)^n$ is expanded as follows.

$$\begin{aligned} (x + y)^n &= x^n + \frac{n!}{1!(n-1)!}x^{n-1}y + \frac{n!}{2!(n-2)!}x^{n-2}y^2 \\ &+ \frac{n!}{3!(n-3)!}x^{n-3}y^3 + \cdots + \frac{n!}{(n-1)!1!}xy^{n-1} \\ &+ y^n \end{aligned}$$

 k th Term of the Binomial Expansion of $(x + y)^n$

If $n \geq k - 1$, then the k th term of the expansion of $(x + y)^n$ is given by the following.

$$\frac{n!}{(k-1)![n-(k-1)]!}x^{n-(k-1)}y^{k-1}$$

EXAMPLES

Expand $(2m + 3)^4$.

$$\begin{aligned} (2m + 3)^4 &= (2m)^4 + \frac{4!}{1!3!}(2m)^3(3) + \frac{4!}{2!2!}(2m)^2(3)^2 \\ &+ \frac{4!}{3!1!}(2m)(3)^3 + 3^4 \\ &= 2^4m^4 + 4(2)^3m^3(3) + 6(2)^2m^2(9) + 4(2m)(27) + 81 \\ &= 16m^4 + 12(8)m^3 + 54(4)m^2 + 216m + 81 \\ &= 16m^4 + 96m^3 + 216m^2 + 216m + 81 \end{aligned}$$

Find the eighth term of the expansion of $(a - 2b)^{10}$.

$$\begin{aligned} \frac{10!}{7!3!}(a^3)(-2b)^7 & \quad \text{Let } n = 10, x = a, y = -2b, \\ & \quad \text{and } k = 8. \\ &= 120(a^3)(-2)^7b^7 \quad \text{Simplify the factorials;} \\ & \quad (ab)^m = a^mb^m \\ &= 120a^3(-128b^7) \quad \text{Simplify.} \\ &= -15,360a^3b^7 \quad \text{Multiply.} \end{aligned}$$

Chapter 11

Review Exercises

11.1 Write the first four terms of each sequence.

- $a_n = 2n - 3$
- $a_n = \frac{n-1}{n}$
- $a_n = n^2$
- $a_n = \left(\frac{1}{2}\right)^n$
- $a_n = (n+1)(n-1)$
- $a_n = n(-1)^{n-1}$

Write each series as a sum of terms and then find the sum.

- $\sum_{i=1}^5 i^2$
- $\sum_{i=1}^6 (i+1)$
- $\sum_{i=3}^6 (5i-4)$
- $\sum_{i=1}^4 (i+2)^2$
- $\sum_{i=1}^6 2^i$
- $\sum_{i=4}^7 \frac{i}{i+1}$

- 13.** The table shows the worldwide number of electric vehicles, in thousands, in use from 2012 to 2017. To the nearest tenth, what was the average number of electric vehicles for this period?

Year	Electric Vehicles (in thousands)
2012	110
2013	220
2014	409
2015	727
2016	1186
2017	1928

Data from www.statista.com

11.2, 11.3 Determine whether each sequence is arithmetic, geometric, or neither. If the sequence is arithmetic, find the common difference d . If it is geometric, find the common ratio r .

14. 2, 5, 8, 11, ... 15. -6, -2, 2, 6, 10, ... 16. $\frac{2}{3}, -\frac{1}{3}, \frac{1}{6}, -\frac{1}{12}, \dots$

17. -1, 1, -1, 1, -1, ... 18. 64, 32, 8, $\frac{1}{2}, \dots$ 19. 64, 32, 8, 1, ...

20. The *Fibonacci sequence* begins 1, 1, 2, 3, 5, 8, What is the eleventh term of this sequence?

11.2 Find the indicated term for each arithmetic sequence.

21. $a_1 = -2, d = 5; a_{16}$ 22. $a_6 = 12, a_8 = 18; a_{25}$

Determine an expression for the general term of each arithmetic sequence.

23. $a_1 = -4, d = -5$ 24. 6, 3, 0, -3, ...

Find the number of terms in each arithmetic sequence.

25. 7, 10, 13, ..., 49 26. 5, 1, -3, ..., -79

Evaluate S_8 for each arithmetic sequence.

27. $a_1 = -2, d = 6$ 28. $a_n = -2 + 5n$

11.3 Determine an expression for the general term of each geometric sequence.

29. -1, -4, -16, ... 30. $\frac{2}{3}, \frac{2}{15}, \frac{2}{75}, \dots$

Find the indicated term for each geometric sequence.

31. 2, -6, 18, ...; a_{11} 32. $a_3 = 20, a_5 = 80; a_{10}$

Evaluate each sum if it exists.

33. $\sum_{i=1}^5 \left(\frac{1}{4}\right)^i$ 34. $\sum_{i=1}^8 \frac{3}{4}(-1)^i$ 35. $\sum_{i=1}^{\infty} 4\left(\frac{1}{5}\right)^i$ 36. $\sum_{i=1}^{\infty} 2(3)^i$

11.4 Use the binomial theorem to expand each binomial.

37. $(2p - q)^5$ 38. $(x^2 + 3y)^4$ 39. $(3t^3 - s^2)^4$

40. Write the fourth term of the expansion of $(3a + 2b)^{19}$.

Chapter 11 Mixed Review Exercises

Find the indicated term and evaluate S_{10} for each sequence.

1. a_{10} ; geometric; -3, 6, -12, ... 2. a_{40} ; arithmetic; 1, 7, 13, ...
 3. a_{15} ; arithmetic; $a_1 = -4, d = 3$ 4. a_9 ; geometric; $a_1 = 1, r = -3$

Determine an expression for the general term of each arithmetic or geometric sequence.

5. 2, 8, 32, ... 6. 2, 7, 12, ... 7. 12, 9, 6, ... 8. 27, 9, 3, ...

Solve each problem.

9. When Faith's sled goes down the hill near her home, she covers 3 ft in the first second. Then, for each second after that, she goes 4 ft more than in the preceding second. If the distance she covers going down is 210 ft, how long does it take her to reach the bottom?
10. An ordinary annuity is set up so that \$672 is deposited at the end of each quarter for 7 yr. The money earns 4.5% annual interest compounded quarterly. What is the future value of the annuity?
11. The school population in Middleton has been dropping 3% per yr. The current population is 50,000. If this trend continues, what will the population be in 6 yr?
12. A pump removes $\frac{1}{2}$ of the liquid in a container with each stroke. What fraction of the liquid is left in the container after seven strokes?
13. Consider the repeating decimal number 0.55555
- Write it as the sum of the terms of an infinite geometric sequence.
 - What is r for this sequence?
 - Find this infinite sum if it exists, and write it as a fraction in lowest terms.
14. Can the sum of the terms of the infinite geometric sequence defined by $a_n = 5(2)^n$ be found? Explain.

Chapter 11

Test

FOR
EXTRA
HELP

Step-by-step test solutions are found on the Chapter Test Prep Videos available in MyLab Math.

► View the complete solutions to all Chapter Test exercises in MyLab Math.

Write the first five terms of each sequence described.

1. $a_n = (-1)^n + 1$ 2. arithmetic, with $a_1 = 4$ and $d = 2$
 3. geometric, with $a_4 = 6$ and $r = \frac{1}{2}$

Determine a_4 for each sequence described.

4. arithmetic, with $a_1 = 6$ and $d = -2$ 5. geometric, with $a_5 = 16$ and $a_7 = 9$

Evaluate S_5 for each sequence described.

6. arithmetic, with $a_2 = 12$ and $a_3 = 15$ 7. geometric, with $a_5 = 4$ and $a_7 = 1$

8. The first eight time intervals, in minutes, between eruptions of the Old Faithful geyser in Yellowstone National Park on July 19, 2018, are given here. Calculate the average number of minutes between eruptions to the nearest tenth. (Data from www.geysertimes.org)

89, 94, 99, 85, 101, 89, 97, 86



9. If \$4000 is deposited in an ordinary annuity at the end of each quarter for 7 yr and earns 6% interest compounded quarterly, how much will be in the account at the end of this term?
10. Under what conditions does an infinite geometric series have a sum?

Evaluate each sum if it exists.

$$11. \sum_{i=1}^5 (2i + 8) \qquad 12. \sum_{i=1}^6 (3i - 5) \qquad 13. \sum_{i=1}^{500} i$$

$$14. \sum_{i=1}^3 \frac{1}{2}(4^i) \qquad 15. \sum_{i=1}^{\infty} \left(\frac{1}{4}\right)^i \qquad 16. \sum_{i=1}^{\infty} 6\left(\frac{3}{2}\right)^i$$

Evaluate each expression.

$$17. 8! \qquad 18. 0! \qquad 19. \frac{6!}{4!2!} \qquad 20. {}_{12}C_{10}$$

21. Expand $(3k - 5)^4$.

22. Write the fifth term of the expansion of $\left(2x - \frac{y}{3}\right)^{12}$.

Solve each problem.

23. Christian bought a new bicycle for \$300. He agreed to pay \$20 per month for 15 months, plus interest of 1% each month, on the unpaid balance. Find the total cost of the bicycle.
24. During the summer months, the population of a certain insect colony triples each week. If there are 20 insects in the colony at the end of the first week in July, how many are present by the end of September? (Assume exactly four weeks in a month.)

Chapters R–11 Cumulative Review Exercises

Simplify each expression.

$$1. |-7| + 6 - |-10| - (-8 + 3) \qquad 2. -\frac{7}{30} + \frac{11}{45} - \frac{3}{10}$$

Let $S = \left\{-\frac{8}{3}, 10, 0, \sqrt{13}, -\sqrt{3}, \frac{45}{15}, \sqrt{-7}, 0.82, -3\right\}$. List the elements of S that are members of each set.

3. Rational numbers

4. Irrational numbers

Solve each equation or inequality.

$$5. 9 - (5 + 3x) + 5x = -4(x - 3) - 7 \qquad 6. \frac{x + 3}{12} - \frac{x - 3}{6} = 0$$

7. $7x + 18 \leq 9x - 2$

8. $2x > 8$ or $-3x > 9$

9. $|4x - 3| = 21$

10. $|2x - 5| \geq 11$

11. $2x^2 + x = 10$

12. $6x^2 + 5x = 8$

$$13. \frac{4}{x-3} - \frac{6}{x+3} = \frac{24}{x^2-9}$$

$$14. 3^{2x-1} = 81$$

$$15. \log_8 x + \log_8 (x+2) = 1$$

Solve each problem.

$$16. \text{ Find the slope of the line passing through } (4, -5) \text{ and } (-12, -17).$$

$$17. \text{ Find the standard form of the equation of the line passing through } (-2, 10) \text{ and parallel to the line with equation } 3x + y = 7.$$

$$18. \text{ Write the equation of a circle with center } (-5, 12) \text{ and radius } 9.$$

Solve each system of equations.

$$19. \begin{cases} y = 5x + 3 \\ 2x + 3y = -8 \end{cases}$$

$$20. \begin{cases} x + 2y + z = 8 \\ 2x - y + 3z = 15 \\ -x + 3y - 3z = -11 \end{cases}$$

$$21. \begin{cases} xy = -5 \\ 2x + y = 3 \end{cases}$$

$$22. \text{ Nuts worth } \$3 \text{ per lb are to be mixed with } 8 \text{ lb of nuts worth } \$4.25 \text{ per lb to obtain a mixture that will be sold for } \$4 \text{ per lb. How many pounds of the } \$3 \text{ nuts should be used?}$$



Simplify. Assume that all variables represent nonzero real numbers.

$$23. \left(\frac{2}{3}\right)^{-2}$$

$$24. \frac{(3p^2)^3(-2p^6)}{4p^3(5p^7)}$$

Perform the indicated operations.

$$25. (4p + 2)(5p - 3)$$

$$26. (2m^3 - 3m^2 + 8m) - (7m^3 + 5m - 8)$$

$$27. (6t^4 + 5t^3 - 18t^2 + 14t - 1) \div (3t - 2)$$

$$28. (8 + 3i)(8 - 3i)$$

Factor.

$$29. 6z^3 + 5z^2 - 4z$$

$$30. 49a^4 - 9b^2$$

$$31. c^3 + 27d^3$$

Simplify.

$$32. \frac{x^2 - 16}{x^2 + 2x - 8} \div \frac{x - 4}{x + 7}$$

$$33. \frac{5}{p^2 + 3p} - \frac{2}{p^2 - 4p}$$

$$34. 5\sqrt{72} - 4\sqrt{50}$$

Graph.

$$35. x - 3y = 6$$

$$36. 4x - y < 4$$

37. $f(x) = 2(x - 2)^2 - 3$

38. $g(x) = \left(\frac{1}{3}\right)^x$

39. $y = \log_{1/3} x$

40. $f(x) = \frac{1}{x - 3}$

41. $\frac{x^2}{9} + \frac{y^2}{25} = 1$

42. $x^2 - y^2 = 9$

Solve each problem.

43. Find $f^{-1}(x)$ if $f(x) = 9x + 5$.

44. Write the first five terms of the sequence with general term $a_n = 5n - 12$.

45. Find the sum of the first six terms of the arithmetic sequence with $a_1 = 8$ and $d = 2$.

46. Find the sum of the geometric series $15 - 6 + \frac{12}{5} - \frac{24}{25} + \dots$.

47. Find the sum $\sum_{i=1}^4 3i$.

48. Evaluate. (a) $9!$ (b) $0!$ (c) ${}_{10}C_4$

49. Use the binomial theorem to expand $(2a - 1)^5$.

50. Find the fourth term in the expansion of $\left(3x^4 - \frac{1}{2}y^2\right)^5$.

Synthetic Division

APPENDIX

OBJECTIVES

- 1 Use synthetic division to divide by a polynomial of the form $x - k$.
- 2 Use the remainder theorem to evaluate a polynomial.
- 3 Use the remainder theorem to decide whether a given number is a solution of an equation.

OBJECTIVE 1 Use synthetic division to divide by a polynomial of the form $x - k$.

If a polynomial in x is divided by a binomial of the form $x - k$, a shortcut method called **synthetic division** can be used. Consider the following.

Polynomial Division	Synthetic Division
$\begin{array}{r} 3x^2 + 9x + 25 \\ x - 3 \overline{) 3x^3 + 0x^2 - 2x + 5} \\ \underline{3x^3 - 9x^2} \\ 9x^2 - 2x \\ \underline{9x^2 - 27x} \\ 25x + 5 \\ \underline{25x - 75} \\ 80 \end{array}$	$\begin{array}{r} 3 \quad 9 \quad 25 \\ 1 - 3 \overline{) 3 \quad 0 \quad -2 \quad 5} \\ \underline{3 \quad -9} \\ 9 \quad -2 \\ \underline{9 \quad -27} \\ 25 \quad 5 \\ \underline{25 \quad -75} \\ 80 \end{array}$

On the right above, exactly the same division is shown written without the variables. This is why it is *essential* to use 0 as a placeholder in synthetic division. All the numbers in color on the right are repetitions of the numbers directly above them, so we omit them to condense our work, as shown on the left below.

$\begin{array}{r} 3 \quad 9 \quad 25 \\ 1 - 3 \overline{) 3 \quad 0 \quad -2 \quad 5} \\ \underline{-9} \\ 9 \quad -2 \\ \underline{-27} \\ 25 \quad 5 \\ \underline{-75} \\ 80 \end{array}$	$\begin{array}{r} 3 \quad 9 \quad 25 \\ 1 - 3 \overline{) 3 \quad 0 \quad -2 \quad 5} \\ \underline{-9} \\ 9 \\ \underline{-27} \\ 25 \\ \underline{-75} \\ 80 \end{array}$
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

The numbers in color on the left are again repetitions of the numbers directly above them. They too are omitted, as shown on the right above. If we bring the 3 in the dividend down to the beginning of the bottom row, the top row can be omitted because it duplicates the bottom row.

$$\begin{array}{r} 1 - 3 \overline{) 3 \quad 0 \quad -2 \quad 5} \\ \underline{-9 \quad -27 \quad -75} \\ 3 \quad 9 \quad 25 \quad 80 \end{array}$$

We omit the 1 at the upper left—it represents $1x$, which will always be the first term in the divisor. To simplify the arithmetic, we replace subtraction in the second row by addition. To compensate, we change the -3 at the upper left to its additive inverse, 3.

$$\begin{array}{r}
 \text{Additive inverse of } -3 \rightarrow 3 \overline{)3 \quad 0 \quad -2 \quad 5} \\
 \underline{ \quad 9 \quad 27 \quad 75} \leftarrow \text{Signs changed} \\
 3 \quad 9 \quad 25 \quad 80 \leftarrow \text{Remainder} \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 3x^2 + 9x + 25 + \frac{80}{x-3}
 \end{array}$$

The quotient is read from the bottom row.

Remember to add $\frac{\text{remainder}}{\text{divisor}}$.

The first three numbers in the bottom row are the coefficients of the quotient polynomial with degree one less than the degree of the dividend. The last number gives the remainder.

Synthetic Division

Synthetic division is a shortcut procedure for polynomial division that eliminates writing the variable factors. It is used only when dividing a polynomial by a binomial of the form $x - k$.

NOW TRY EXERCISE 1

Use synthetic division to divide.

$$\frac{4x^3 + 18x^2 + 19x + 7}{x + 3}$$

EXAMPLE 1 Using Synthetic Division

Use synthetic division to divide.

$$\frac{5x^2 + 16x + 15}{x + 2}$$

Remember that a fraction bar means division.

We change $x + 2$ into the form $x - k$ by writing it as

$$x + 2 = x - (-2), \quad \text{where } k = -2.$$

Now we write the coefficients of $5x^2 + 16x + 15$, placing -2 to the left.

$$\begin{array}{r}
 x + 2 \text{ leads to } -2. \rightarrow -2 \overline{)5 \quad 16 \quad 15} \leftarrow \text{Coefficients} \\
 \underline{-2 \overline{)5 \quad 16 \quad 15}} \\
 \downarrow \quad -10 \\
 5 \quad \quad \\
 \underline{-2 \overline{)5 \quad 16 \quad 15}} \\
 \quad -10 \quad -12 \\
 \quad \quad \\
 \underline{-2 \overline{)5 \quad 16 \quad 15}} \\
 \quad -10 \quad -12 \\
 \quad \quad 3 \leftarrow \text{Remainder} \\
 \quad \quad
 \end{array}$$

Bring down the 5, and multiply: $-2 \cdot 5 = -10$.

Add 16 and -10 , obtaining 6, and multiply 6 and -2 to obtain -12 .

Add 15 and -12 , obtaining 3.

Read the result from the bottom row.

$$\frac{5x^2 + 16x + 15}{x + 2} = 5x + 6 + \frac{3}{x + 2}$$

Add $\frac{\text{remainder}}{\text{divisor}}$.

CHECK We can check this result as with polynomial division.

$$\begin{aligned}
 (x + 2)(5x + 6) + 3 & \quad \text{Divisor} \times \text{Quotient} + \text{Remainder} \\
 = 5x^2 + 6x + 10x + 12 + 3 & \quad \text{FOIL method} \\
 = 5x^2 + 16x + 15 \quad \checkmark & \quad \text{Dividend}
 \end{aligned}$$

NOW TRY ANSWER

1. $4x^2 + 6x + 1 + \frac{4}{x+3}$

NOW TRY EXERCISE 2

Use synthetic division to divide.

$$\frac{-3x^4 + 13x^3 - 6x^2 + 31}{x - 4}$$

EXAMPLE 2 Using Synthetic Division with a Missing Term

Use synthetic division to divide.

$$\frac{-4x^5 + x^4 + 6x^3 + 2x^2 + 50}{x - 2}$$

This quotient could also be written as

$$(-4x^5 + x^4 + 6x^3 + 2x^2 + 50) \div (x - 2).$$

In long division form, the procedure is set up as follows.

$$x - 2 \overline{) -4x^5 + x^4 + 6x^3 + 2x^2 + 50}$$

Now use synthetic division.

$$\begin{array}{r|rrrrrr} 2 & -4 & 1 & 6 & 2 & 0 & 50 \\ & & -8 & -14 & -16 & -28 & -56 \\ \hline & -4 & -7 & -8 & -14 & -28 & -6 \end{array}$$

Use the steps given previously, first inserting a 0 for the missing x -term.

Read the result from the bottom row.

$$\frac{-4x^5 + x^4 + 6x^3 + 2x^2 + 50}{x - 2} = -4x^4 - 7x^3 - 8x^2 - 14x - 28 + \frac{-6}{x - 2}$$

NOW TRY

OBJECTIVE 2 Use the remainder theorem to evaluate a polynomial.

We can use synthetic division to evaluate polynomials. In the synthetic division of **Example 2**, where the polynomial was divided by $x - 2$, the remainder was -6 .

Replacing x in the polynomial with 2 gives the following.

$$\begin{aligned} -4x^5 + x^4 + 6x^3 + 2x^2 + 50 & \quad \text{Dividend in Example 2} \\ = -4 \cdot 2^5 + 2^4 + 6 \cdot 2^3 + 2 \cdot 2^2 + 50 & \quad \text{Replace } x \text{ with } 2. \\ = -4 \cdot 32 + 16 + 6 \cdot 8 + 2 \cdot 4 + 50 & \quad \text{Evaluate the powers.} \\ = -128 + 16 + 48 + 8 + 50 & \quad \text{Multiply.} \\ = -6 & \quad \text{Add.} \end{aligned}$$

This number, -6 , is the same number as the remainder. Dividing by $x - 2$ produced a remainder equal to the result when x is replaced with 2. This always happens, as the **remainder theorem** states. This result is proved in more advanced courses.

Remainder Theorem

If the polynomial $f(x)$ is divided by $x - k$, then the remainder is equal to $f(k)$.

NOW TRY EXERCISE 3

Let $f(x) = 3x^3 - 2x^2 + 5x + 30$. Use the remainder theorem to evaluate $f(-2)$.

EXAMPLE 3 Using the Remainder Theorem

Let $f(x) = 2x^3 - 5x^2 - 3x + 11$. Use the remainder theorem to evaluate $f(-2)$.

Divide $f(x)$ by $x - (-2)$, using synthetic division.

$$\begin{array}{r|rrrr} \text{Value of } k \rightarrow -2 & 2 & -5 & -3 & 11 \\ & & -4 & 18 & -30 \\ \hline & 2 & -9 & 15 & -19 \end{array}$$

Thus, $f(-2) = -19$.

$-19 \leftarrow$ Remainder

NOW TRY

NOW TRY ANSWERS

2. $-3x^3 + x^2 - 2x - 8 + \frac{-1}{x-4}$
3. -12


NOW TRY EXERCISE 4

Use the remainder theorem to decide whether -4 is a solution of the equation.

$$5x^3 + 19x^2 - 2x + 8 = 0$$

OBJECTIVE 3 Use the remainder theorem to decide whether a given number is a solution of an equation.

EXAMPLE 4 Using the Remainder Theorem

Use the remainder theorem to decide whether -5 is a solution of the equation.

$$2x^4 + 12x^3 + 6x^2 - 5x + 75 = 0$$

If synthetic division gives a remainder of 0, then -5 is a solution. Otherwise, it is not.

$$\begin{array}{r|rrrrr} \text{Proposed solution} \rightarrow -5 & 2 & 12 & 6 & -5 & 75 \\ & & -10 & -10 & 20 & -75 \\ \hline & 2 & 2 & -4 & 15 & 0 \leftarrow \text{Remainder} \end{array}$$

Because the remainder is 0, the polynomial has value 0 when $k = -5$. So -5 is a solution of the given equation.

NOW TRY 

The synthetic division in **Example 4** shows that $x - (-5)$ divides the polynomial with 0 remainder. Thus $x - (-5) = x + 5$ is a *factor* of the polynomial.

$$2x^4 + 12x^3 + 6x^2 - 5x + 75 \text{ factors as } (x + 5)(2x^3 + 2x^2 - 4x + 15).$$

The second factor is the quotient polynomial in the last row of the synthetic division.

NOW TRY ANSWER


4. yes

Exercises

FOR
EXTRA
HELP



MyLab Math

 Video solutions for select problems available in MyLab Math

Concept Check Choose the letter of the correct setup to perform synthetic division on the indicated quotient.

1. $\frac{x^2 + 3x - 6}{x - 2}$

A. $-2 \overline{) 1 \quad 3 \quad -6}$

B. $-2 \overline{) -1 \quad -3 \quad 6}$

C. $2 \overline{) 1 \quad 3 \quad -6}$

D. $2 \overline{) -1 \quad -3 \quad 6}$

2. $\frac{x^3 - 3x^2 + 2}{x - 1}$

A. $1 \overline{) 1 \quad -3 \quad 2}$

B. $-1 \overline{) 1 \quad -3 \quad 2}$

C. $1 \overline{) 1 \quad -3 \quad 0 \quad 2}$

D. $1 \overline{) -1 \quad 3 \quad 0 \quad -2}$

Concept Check Fill in each blank with the appropriate response.

3. In arithmetic, the result of the division

$$\begin{array}{r} 3 \\ 5 \overline{) 19} \\ \underline{15} \\ 4 \end{array}$$

can be written

$$19 = 5 \cdot \underline{\hspace{1cm}} + \underline{\hspace{1cm}}.$$

4. In algebra, the result of the division

$$\begin{array}{r} x + 3 \\ x - 1 \overline{) x^2 + 2x + 3} \\ \underline{x^2 - x} \\ 3x + 3 \\ \underline{3x - 3} \\ 6 \end{array}$$

can be written $x^2 + 2x + 3 =$

$$(x - 1)(\underline{\hspace{1cm}}) + \underline{\hspace{1cm}}.$$

5. To perform the division $x - 3 \overline{) x^3 + 6x^2 + 2x}$ using synthetic division, we begin by writing the following.

$$\underline{\hspace{1cm}} \overline{) 1 \quad \underline{\hspace{1cm}} \quad 2 \quad \underline{\hspace{1cm}}}$$

6. Consider the following function.

$$f(x) = 2x^4 + 6x^3 - 5x^2 + 3x + 8$$

$$f(x) = (x - 2)(2x^3 + 10x^2 + 15x + 33) + 74$$

By inspection, we can state that $f(2) = \underline{\hspace{2cm}}$.

7. **Concept Check** A student attempted to divide

$$4x^3 + 2x^2 + 6 \text{ by } x + 2$$

synthetically by setting up the division as follows.

$$\begin{array}{r} -2 \overline{)4 \ 2 \ 6} \end{array}$$

This is incorrect. **WHAT WENT WRONG?** Give the correct setup and the answer.

8. **Concept Check** A student attempted to divide

$$4x^3 + 2x^2 + 6x \text{ by } x + 2$$

synthetically by setting up the division as follows.

$$\begin{array}{r} -2 \overline{)4 \ 2 \ 6} \end{array}$$

This is incorrect. **WHAT WENT WRONG?** Give the correct setup and the answer.

Use synthetic division to divide. See Examples 1 and 2.

9. $\frac{x^2 - 6x + 5}{x - 1}$

10. $\frac{x^2 + 4x - 21}{x - 3}$

11. $\frac{4x^2 + 19x - 5}{x + 5}$

12. $\frac{3x^2 + 5x - 12}{x + 3}$

13. $\frac{2x^2 + 8x + 13}{x + 2}$

14. $\frac{4x^2 - 5x - 20}{x - 4}$

15. $\frac{x^2 - 3x + 5}{x + 1}$

16. $\frac{x^2 + 4x - 6}{x - 5}$

17. $\frac{4x^3 - 3x^2 + 2x - 3}{x - 1}$

18. $\frac{5x^3 - 6x^2 + 3x + 14}{x + 1}$

19. $\frac{x^3 + 2x^2 - 4x + 3}{x - 4}$

20. $\frac{x^3 - 3x^2 + 5x - 1}{x - 5}$

21. $\frac{2x^5 - 2x^3 + 3x^2 - 24x - 2}{x - 2}$

22. $\frac{3x^5 + x^4 - 84x^2 - 12x + 3}{x - 3}$

23. $\frac{-3x^5 - 3x^4 + 5x^3 - 6x^2 + 3}{x + 1}$

24. $\frac{-3x^5 + 2x^4 - 5x^3 - 6x^2 - 1}{x + 2}$

25. $\frac{x^5 + x^4 + x^3 + x^2 + x + 3}{x + 1}$

26. $\frac{x^5 - x^4 + x^3 - x^2 + x - 2}{x - 1}$

Use the remainder theorem to evaluate $f(k)$. See Example 3.

27. $f(x) = 2x^3 - 4x^2 + 5x - 3; \quad k = 2$

28. $f(x) = x^3 + 3x^2 - x + 5; \quad k = -1$

29. $f(x) = -x^3 - 5x^2 - 4x - 2; \quad k = -4$

30. $f(x) = -x^3 + 5x^2 - 3x + 4; \quad k = 3$

31. $f(x) = 2x^3 - 4x^2 + 5x - 33; \quad k = 3$

32. $f(x) = x^3 - 3x^2 + 4x - 4; \quad k = 2$

Use the remainder theorem to decide whether the given number is a solution of the equation. See Example 4.

33. $x^3 - 2x^2 - 3x + 10 = 0; \quad x = -2$

34. $x^3 - 3x^2 - x + 10 = 0; \quad x = -2$

35. $3x^3 + 2x^2 - 2x + 11 = 0; \quad x = -2$

36. $3x^3 + 10x^2 + 3x - 9 = 0; \quad x = -2$

37. $2x^3 - x^2 - 13x + 24 = 0$; $x = -3$

38. $5x^3 + 22x^2 + x - 28 = 0$; $x = -4$

39. $x^4 + 2x^3 - 3x^2 + 8x = 8$; $x = -2$

40. $x^4 - x^3 - 6x^2 + 5x = -10$; $x = -2$

RELATING CONCEPTS For Individual or Group Work (Exercises 41–46)

We can show a connection between dividing one polynomial by another and factoring the first polynomial. Let $f(x) = 2x^2 + 5x - 12$. **Work Exercises 41–46 in order.**

41. Factor $f(x)$.

42. Solve $f(x) = 0$.

43. Evaluate $f(-4)$.

44. Evaluate $f\left(\frac{3}{2}\right)$.

45. Complete the sentence: If $f(a) = 0$, then $x - \underline{\hspace{2cm}}$ is a factor of $f(x)$.

46. Use the conclusion reached in **Exercise 45** to decide whether $x - 3$ is a factor of $g(x) = 3x^3 - 4x^2 - 17x + 6$. If so, factor $g(x)$ completely.

ANSWERS TO SELECTED EXERCISES

In this section, we provide the answers that we think most students will obtain when they work the exercises using the methods explained in the text. If your answer does not look exactly like the one given here, it is not necessarily wrong. In many cases, there are equivalent forms of the answer that are correct. For example, if the answer section shows $\frac{3}{4}$ and your answer is 0.75, you have obtained the right answer, but written it in a different (yet equivalent) form. Unless the directions specify otherwise, 0.75 is just as valid an answer as $\frac{3}{4}$.

In general, if your answer does not agree with the one given in the text, see whether it can be transformed into the other form. If it can, then it is the correct answer. If you still have doubts, talk with your instructor.

R REVIEW OF THE REAL NUMBER SYSTEM

Section R.1


1. true 3. false; This is an improper fraction. Its value is 1. 5. C
 7. $\frac{1}{2}$ 9. $\frac{5}{6}$ 11. $\frac{3}{5}$ 13. $\frac{1}{5}$ 15. $\frac{6}{5}$ 17. $1\frac{5}{7}$ 19. $6\frac{5}{12}$ 21. $\frac{13}{5}$ 23. $\frac{38}{3}$
 25. $\frac{24}{35}$ 27. $\frac{1}{20}$ 29. $\frac{6}{25}$ 31. $\frac{6}{5}$, or $1\frac{1}{5}$ 33. 9 35. $\frac{65}{12}$, or $5\frac{5}{12}$ 37. $\frac{14}{27}$
 39. $\frac{10}{3}$, or $3\frac{1}{3}$ 41. 12 43. $\frac{1}{16}$ 45. 10 47. 18 49. $\frac{35}{24}$, or $1\frac{11}{24}$
 51. $\frac{84}{47}$, or $1\frac{37}{47}$ 53. $\frac{11}{15}$ 55. $\frac{2}{3}$ 57. $\frac{8}{9}$ 59. $\frac{29}{24}$, or $1\frac{5}{24}$ 61. $\frac{107}{144}$
 63. $\frac{43}{8}$, or $5\frac{3}{8}$ 65. $\frac{5}{9}$ 67. $\frac{2}{3}$ 69. $\frac{1}{4}$ 71. $\frac{17}{36}$ 73. $\frac{67}{20}$, or $3\frac{7}{20}$
 75. $\frac{32}{9}$, or $3\frac{5}{9}$ 77. (a) $\frac{1}{2}$ (b) $\frac{1}{4}$ (c) $\frac{1}{3}$ (d) $\frac{1}{6}$ 79. (a) 6 (b) 9
 (c) 1 (d) 7 (e) 4 81. (a) 46.25 (b) 46.2 (c) 46 (d) 50
 83. $\frac{4}{10}$ 85. $\frac{64}{100}$ 87. $\frac{138}{1000}$ 89. $\frac{43}{1000}$ 91. $\frac{3805}{1000}$ 93. 143.094
 95. 25.61 97. 15.33 99. 21.77 101. 81.716 103. 15.211
 105. 116.48 107. 739.53 109. 0.006 111. 7.15 113. 2.8
 115. 2.05 117. 1232.6 119. 5711.6 121. 94 123. 0.162
 125. 1.2403 127. 0.02329 129. 1% 131. $\frac{1}{20}$ 133. $12\frac{1}{2}\%$, or
 12.5% 135. 0.25; 25% 137. $\frac{1}{2}$; 0.5 139. $\frac{3}{4}$; 75% 141. 4.2
 143. 2.25 145. 0.375 147. $0.\overline{5}$; 0.556 149. $0.1\overline{6}$; 0.167 151. 0.54
 153. 0.07 155. 1.17 157. 0.024 159. 0.0625 161. 79% 163. 2%
 165. 0.4% 167. 128% 169. 40% 171. $\frac{51}{100}$ 173. $\frac{3}{20}$ 175. $\frac{1}{50}$
 177. $\frac{7}{5}$, or $1\frac{2}{5}$ 179. $\frac{3}{40}$ 181. 80% 183. 14% 185. $18.\overline{18}\%$
 187. 225% 189. $216.\overline{6}\%$

Section R.2

1. yes 3. {1, 2, 3, 4, 5} 5. {5, 6, 7, 8, ...}
 7. {..., -1, 0, 1, 2, 3, 4} 9. {10, 12, 14, 16, ...}
 11. {-4, 4} 13. \emptyset

In Exercises 15 and 17, we give one possible answer.

15. { x | x is an even natural number less than or equal to 8}
 17. { x | x is a positive multiple of 4}



19. $-\frac{6}{5}, -\frac{1}{4}, \frac{5}{6}, \frac{13}{4}, 5.2, \frac{11}{2}$
 23. (a) 8, 13, $\frac{75}{5}$ (or 15) (b) 0, 8, 13, $\frac{75}{5}$ (c) -9, 0, 8, 13, $\frac{75}{5}$
 (d) -9, -0.7, 0, $\frac{6}{7}$, $4.\overline{6}$, 8, $\frac{21}{2}$, 13, $\frac{75}{5}$ (e) $-\sqrt{6}$, $\sqrt{7}$ (f) All are real numbers.
 25. false; Some are whole numbers, but negative integers are not. 27. false; No irrational number is an integer. 29. true 31. true
 33. true 35. (a) A (b) A (c) B (d) B 37. (a) -6 (b) 6
 39. (a) 12 (b) 12 41. (a) $-\frac{6}{5}$ (b) $\frac{6}{5}$ 43. 8 45. $\frac{3}{2}$ 47. -5
 49. -2 51. -4.5 53. 5 55. -1 57. 6 59. 0
 61. (a) New Orleans; The population increased by 13.9%.
 (b) Toledo; The population decreased by 3.0%. 63. Pacific Ocean, Indian Ocean, Caribbean Sea, South China Sea, Gulf of California 65. true
 67. true 69. false 71. true 73. false 75. $2 < 6$ 77. $4 > -9$
 79. $-10 < -5$ 81. $7 > -1$ 83. $5 \geq 5$ 85. $13 - 3 \leq 10$
 87. $5 + 0 \neq 0$ 89. false 91. true 93. $-6 < 10$; true
 95. $10 \geq 10$; true 97. $-3 \geq -3$; true 99. $-8 > -6$; false
 101. Iowa (IA), Ohio (OH), Pennsylvania (PA) 103. $x < y$

Section R.3

1. positive; $18 + 6 = 24$ 3. greater; $-14 + 9 = -5$ 5. the number with greater absolute value is subtracted from the one with lesser absolute value; $5 - 12 = -7$ 7. positive; $-2(-8) = 16$
 9. undefined; 0; $\frac{-17}{0}$ is undefined; $\frac{0}{42} = 0$ 11. -19 13. -9 15. 9
 17. 5 19. $-\frac{19}{12}$ 21. 1.7 23. -5 25. -11 27. 21 29. -13
 31. -10.18 33. $\frac{67}{30}$ 35. 14 37. -5 39. -6 41. -11
 43. 0 45. 12 47. -4 49. 3 51. $-\frac{7}{4}$ 53. $-\frac{7}{8}$ 55. 2 57. 1
 59. 6 61. $\frac{13}{2}$, or $6\frac{1}{2}$ 63. It is true for multiplication (and division). It is false for addition and false for subtraction when the number to be subtracted has the lesser absolute value. A more precise statement is "The product or quotient of two negative numbers is positive."
 65. 40 67. -35 69. 0 71. 0 73. -12 75. 2 77. $\frac{6}{5}$
 79. 1 81. 0.4 83. -0.024 85. -7 87. 6 89. -4 91. 0
 93. undefined 95. $\frac{25}{102}$ 97. $-\frac{9}{13}$ 99. -9 101. 3.2 103. $\frac{17}{18}$
 105. $\frac{17}{36}$ 107. $-\frac{19}{24}$ 109. $-\frac{22}{45}$ 111. $-\frac{2}{15}$ 113. $\frac{3}{5}$ 115. $-\frac{20}{7}$
 117. $-\frac{1}{3}$ 119. $-\frac{4}{3}$ 121. -32.351 123. -3 125. -4.14
 127. 4800 129. 10,000 131. -4.39 133. 180°F 135. \$30.13
 137. (a) \$466.02 (b) \$190.68 139. (a) 31.33% (b) -36.87%
 (c) 58.64% 141. 2001: \$128 billion; 2006: -\$248 billion; 2011: -\$1300 billion; 2016: -\$585 billion

Section R.4


1. false; $-7^6 = -(7^6)$ 3. true 5. true 7. true 9. false; The base is 2. 11. 10^4 13. $(\frac{3}{4})^5$ 15. $(-9)^3$ 17. $(0.8)^2$ 19. z^7

21. (a) 64 (b) -64 (c) 64 (d) -64 23. 16 25. 0.027
 27. $\frac{1}{125}$ 29. $\frac{256}{625}$ 31. -125 33. 256 35. -729 37. -4096
 39. (a) negative (b) positive (c) negative (d) negative
 (e) negative (f) positive 41. 9 43. 13 45. -20 47. $\frac{10}{11}$
 49. -0.7 51. not a real number 53. not a real number
 55. (a) The grandson's answer was correct. (b) The reasoning was incorrect. The operation of division must be done first, and then the addition follows. The grandson's "Order of Process rule" is **not correct**. It happened coincidentally, in this problem, that he obtained the correct answer the wrong way. 57. 24 59. 15 61. 55 63. -91 65. -8
 67. -48 69. -2 71. -97 73. 11 75. -10 77. 2 79. 1
 81. undefined 83. -1 85. 17 87. -96 89. 180 91. $\frac{5}{9}$
 93. $-\frac{15}{238}$ 95. 8 97. $-\frac{5}{16}$ 99. $-\frac{11}{4}$, or -2.75 101. \$2434
 103. (a) $48 \times 3.2 \times 0.075 \div 190 - 2 \times 0.015$ (b) 0.031
 105. (a) 0.024; 0.023; Increased weight results in lower BACs.
 (b) Decreased weight will result in higher BACs; 0.040; 0.053
 107. (a) \$31.5 billion (b) \$48.9 billion (c) \$63.9 billion
 (d) The amount spent on pets roughly doubled from 2003 to 2016.

Section R.5

1. B 3. A 5. $ab + ac$ 7. grouping 9. like 11. $2m + 2p$
 13. $-12x + 12y$ 15. $8k$ 17. $-2r$ 19. cannot be simplified
 21. $8a$ 23. $2x$ 25. $-2d + f$ 27. $x + y$ 29. $2x - 6y + 4z$
 31. $-3y$ 33. $p + 11$ 35. $-2k + 15$ 37. $-4y + 2$ 39. $30xy$
 41. $-56wz$ 43. $m - 14$ 45. -1 47. $2p + 7$ 49. $-6z - 39$
 51. $(5 + 8)x = 13x$ 53. $(5 \cdot 9)r = 45r$ 55. $9y + 5x$ 57. 7
 59. 0 61. $8(-4) + 8x = -32 + 8x$ 63. 0 65. Answers will vary. One example is washing your face and brushing your teeth. 67. 1900
 69. 75 71. 87.5 73. associative property 74. associative property
 75. commutative property 76. associative property 77. distributive property 78. arithmetic facts

Chapter R Test

- [R.1] 1. $\frac{11}{12}$ 2. $\frac{2}{3}$ 3. 6.833 4. 0.028 5. $\frac{1}{25}$; 0.04 6. $0.8\bar{3}$; $83\frac{1}{3}\%$, or $83.\bar{3}\%$ 7. $\frac{3}{2}$, or $1\frac{1}{2}$; 150% [R.2] 8. 
 9. 0, 3, $\sqrt{25}$ (or 5), $\frac{24}{2}$ (or 12) 10. -1, 0, 3, $\sqrt{25}$ (or 5), $\frac{24}{2}$ (or 12)
 11. -1, -0.5, 0, 3, $\sqrt{25}$ (or 5), 7.5, $\frac{24}{2}$ (or 12) 12. All are real numbers except $\sqrt{-4}$. [R.3, R.4] 13. 0 14. $-\frac{17}{63}$ 15. -26
 16. 19 17. $\frac{16}{7}$ 18. undefined [R.4] 19. 14 20. -15
 21. not a real number 22. $-\frac{6}{23}$ [R.5] 23. $10k - 10$ 24. $7r + 2$
 [R.3, R.5] 25. (a) B (b) D (c) A (d) F (e) C (f) C (g) E

1 LINEAR EQUATIONS, INEQUALITIES, AND APPLICATIONS

Section 1.1

1. algebraic expression; does; is not 3. true; solution; solution set
 5. identity; all real numbers 7. A, C 9. equation 11. expression

13. equation 15. A sign error was made when the distributive property was applied. The left side of the second line should be $8x - 4x + 6$. The correct solution is 1. 17. (a) B (b) A (c) C 19. $\{-1\}$
 21. $\{-4\}$ 23. $\{-7\}$ 25. $\{0\}$ 27. $\{4\}$ 29. $\{-\frac{7}{8}\}$
 31. \emptyset ; contradiction 33. $\{-9\}$ 35. $\{-\frac{1}{2}\}$ 37. $\{2\}$ 39. $\{-2\}$
 41. {all real numbers}; identity 43. $\{-1\}$ 45. $\{7\}$ 47. $\{2\}$
 49. {all real numbers}; identity 51. 12 53. (a) 10^2 , or 100
 (b) 10^3 , or 1000 55. $\{-\frac{18}{5}\}$ 57. $\{-\frac{5}{6}\}$ 59. $\{6\}$ 61. $\{4\}$
 63. $\{3\}$ 65. $\{3\}$ 67. $\{0\}$ 69. $\{35\}$ 71. $\{-10\}$ 73. $\{2000\}$
 75. $\{25\}$ 77. $\{40\}$ 79. $\{3\}$

Section 1.2

1. formula 3. (a) 35% (b) 18% (c) 2% (d) 7.5% (e) 150%
 There may be other acceptable forms of the answers in Exercises 5–35.
 5. $r = \frac{I}{pt}$ 7. (a) $W = \frac{\mathcal{A}}{L}$ (b) $L = \frac{\mathcal{A}}{W}$ 9. $L = \frac{P - 2W}{2}$,
 or $L = \frac{P}{2} - W$ 11. (a) $W = \frac{V}{LH}$ (b) $H = \frac{V}{LW}$ 13. $r = \frac{C}{2\pi}$
 15. (a) $b = \frac{2\mathcal{A} - hb}{h}$, or $b = \frac{2\mathcal{A}}{h} - B$ (b) $B = \frac{2\mathcal{A} - hb}{h}$,
 or $B = \frac{2\mathcal{A}}{h} - b$ 17. $C = \frac{5}{9}(F - 32)$ 19. (a) $x = -\frac{b}{a}$ (b) $a = -\frac{b}{x}$
 21. $t = \frac{A - P}{Pr}$ 23. D 25. $y = -4x + 1$ 27. $y = \frac{1}{2}x + 3$
 29. $y = -\frac{4}{9}x + \frac{11}{9}$ 31. $y = \frac{3}{2}x + \frac{5}{2}$ 33. $y = \frac{6}{5}x - \frac{7}{5}$ 35. $y = \frac{3}{2}x - 3$
 37. 3.492 hr 39. 52 mph 41. 108°F 43. 230 m 45. (a) 240 in.
 (b) 480 in. 47. 2 in. 49. 25% 51. 20% 53. 75% 55. 75% water;
 25% alcohol 57. 3% 59. (a) .494 (b) .481 (c) .481 (d) .463
 61. 19% 63. 6.5 million 65. \$72,324 67. \$14,465 69. 12%
 71. 5.1% 73. 10.1% 75. (a) 0.79 m^2 (b) 0.96 m^2 77. (a) 116 mg
 (b) 141 mg 79. (a) $7x + 8 = 36$ (b) $ax + k = tc$
 80. (a) $7x + 8 - 8 = 36 - 8$ (b) $ax + k - k = tc - k$ 81. (a) $7x = 28$
 (b) $ax = tc - k$ 82. (a) $x = 4$; $\{4\}$ (b) $x = \frac{tc - k}{a}$ 83. $a \neq 0$;
 If $a = 0$, the denominator is 0. 84. $\{-5\}$ 85. $a = \frac{2S - n\ell}{n}$, or
 $a = \frac{2S}{n} - \ell$; $n \neq 0$

Section 1.3

1. (a) $x + 15$ (b) $15 > x$ 3. (a) $x - 8$ (b) $8 < x$ 5. Because the unknown number comes first in the verbal phrase, it must come first in the mathematical expression. The correct expression is $x - 7$.
 7. $2x - 13$ 9. $12 + 4x$ 11. $8(x - 16)$ 13. $\frac{3x}{10}$ 15. $x + 6 = -31$; -37
 17. $x - (-4x) = x + 9$; $\frac{9}{4}$ 19. $12 - \frac{2}{3}x = 10$; 3 21. expression;
 $-11x + 63$ 23. equation; $\{\frac{51}{11}\}$ 25. expression; $\frac{1}{3}x - \frac{13}{2}$
 27. Step 1: the number of patents each corporation secured;
 Step 2: patents that Samsung secured; Step 3: x ; $x - 2250$;
 Step 4: 7309; Step 5: 7309; 5059; Step 6: 2250; IBM patents; 5059; 12,368

29. width: 165 ft; length: 265 ft 31. 24.34 in. by 29.88 in.
 33. 850 mi; 925 mi; 1300 mi 35. Walmart: \$486.9 billion; Berkshire Hathaway: \$223.6 billion 37. Eiffel Tower: 984 ft; Leaning Tower: 180 ft
 39. National Geographic: 83.4 million followers; Nike: 75.4 million followers 41. 19.0% 43. 1300 million users 45. \$49.86 47. \$225
 49. We cannot expect the final mixture to be worth more than the more expensive of the two ingredients. 51. \$4000 at 3%; \$8000 at 4%
 53. \$10,000 at 4.5%; \$19,000 at 3% 55. \$24,000 57. 5 L 59. 4 L
 61. 1 gal 63. 150 lb 65. (a) $800 - x$ (b) $800 - y$
 66. (a) $0.03x$; $0.06(800 - x)$ (b) $0.03y$; $0.06(800 - y)$
 67. (a) $0.03x + 0.06(800 - x) = 800(0.0525)$
 (b) $0.03y + 0.06(800 - y) = 800(0.0525)$
 68. (a) \$200 at 3%; \$600 at 6% (b) 200 L of 3% acid; 600 L of 6% acid (c) The processes are the same. The amounts of money in Problem A correspond to the amounts of solution in Problem B.

Section 1.4
















1. \$5.70 3. 30 mph 5. 81 in.² 7. The right side of the equation should be 180, not 90. The correct equation is $x + (x + 1) + (x + 2) = 180$. The angles measure 59°, 60°, and 61°. 9. 17 pennies; 17 dimes; 10 quarters 11. 23 loonies; 14 toonies 13. 28 \$10 coins; 13 \$20 coins
 15. 872 adult tickets 17. 8.01 m per sec 19. 10.11 m per sec
 21. 2½ hr 23. 7:50 P.M. 25. 45 mph 27. ½ hr 29. 60°, 60°, 60°
 31. 40°, 45°, 95° 33. 41°, 52°, 87° 35. Both measure 122°.
 37. 64°, 26° 39. 65°, 115° 41. 19, 20, 21 43. 27, 28, 29, 30
 45. 61 yr old 47. 28, 30, 32 49. 21, 23, 25 51. 40°, 80° 52. 120°
 53. The sum is equal to the measure of the angle found in **Exercise 52**.
 54. The sum of the measures of angles ① and ② is equal to the measure of angle ③.

SUMMARY EXERCISES Applying Problem-Solving Techniques









1. length: 8 in.; width: 5 in. 2. length: 60 m; width: 30 m 3. \$279.98
 4. \$425 5. \$800 at 4%; \$1600 at 5% 6. \$12,000 at 3%; \$14,000 at 4%
 7. Westbrook: 2558 points; Harden: 2356 points 8. 2016: 718 films; 2009: 557 films 9. 5 hr 10. 1½ cm 11. 13⅓ L 12. 53⅓ kg
 13. 84 fives; 42 tens 14. 1650 tickets at \$9; 810 tickets at \$7
 15. 20°, 30°, 130° 16. 107°, 73° 17. 31, 32, 33 18. 9, 11
 19. 6 in., 12 in., 16 in. 20. 23 in.

Section 1.5





1. D 3. B 5. F 7. (a) $x < 100$ (b) $100 \leq x \leq 129$
 (c) $130 \leq x \leq 159$ (d) $160 \leq x \leq 189$ (e) $x \geq 190$
 9. The student divided by 4, a *positive* number. Reverse the symbol only when multiplying or dividing by a *negative* number. The solution set is $[-16, \infty)$.



75. at least 78 77. 26 months 79. more than 4.4 in. 81. (a) 140 to 184 lb (b) 107 to 141 lb (c) Answers will vary. 83. 26 DVDs




45. \emptyset





65. (0, 1) 67. (-2, 2)

69. $[3, \infty)$ 71. $[-9, \infty)$

73. at least 80






















89. $\{-3\}$; $(-3, \infty)$; $(-\infty, -3)$ 90. It is the set of all real numbers; $(-\infty, \infty)$

Section 1.6

1. true 3. false; The union is $(-\infty, 7) \cup (7, \infty)$. 5. $\{4\}$, or D
7. $\{1, 3, 5\}$, or B 9. \emptyset 11. $\{1, 2, 3, 4, 5, 6\}$, or A 13. $\{1, 3, 5, 6\}$























47. $[-4, -1]$ 49. $[-9, -6]$
51. $(-\infty, 3)$ 53. $[3, 9)$



63. {Tuition and fees} 65. {Tuition and fees, Board rates, Dormitory charges} 67. 160 ft; 150 ft; 220 ft; 120 ft 69. Maria, Joe 71. none of them 73. Maria, Joe 75. (a) A: $185 + x \geq 585$; B: $185 + x \geq 520$; C: $185 + x \geq 455$ (b) A: $x \geq 400$; 89%; B: $x \geq 335$; 75%; C: $x \geq 270$; 60% 76. $520 \leq 185 + x \leq 584$; $335 \leq x \leq 399$; $75\% \leq \text{average} \leq 89\%$ 77. A: $105 + x \geq 585$; $x \geq 480$; impossible; B: $105 + x \geq 520$; $x \geq 415$; 93%; C: $105 + x \geq 455$; $x \geq 350$; 78% 78. $455 \leq 105 + x \leq 519$; $350 \leq x \leq 414$; $78\% \leq \text{average} \leq 92\%$

Section 1.7

1. E; C; D; B; A 3. (a) one (b) two (c) none 5. $\{-12, 12\}$
7. $\{-5, 5\}$ 9. $\{-6, 12\}$ 11. $\{-5, 6\}$ 13. $\{-3, \frac{11}{2}\}$ 15. $\{-\frac{19}{2}, \frac{9}{2}\}$
17. $\{\frac{7}{3}, 3\}$ 19. $\{12, 36\}$ 21. $\{-12, 12\}$ 23. $\{-10, -2\}$
25. $\{-\frac{32}{3}, 8\}$ 27. $\{-75, 175\}$








67. $(-\infty, \frac{1}{2}] \cup [\frac{7}{6}, \infty)$ 69. $\{-\frac{5}{3}, \frac{11}{3}\}$ 71. $(-\infty, -20) \cup (40, \infty)$
73. $\{-5, 1\}$ 75. $\{3, 9\}$ 77. $\{0, 20\}$ 79. $\{-5, 5\}$ 81. $\{-5, -3\}$
83. $(-\infty, -3) \cup (2, \infty)$ 85. $[-10, 0]$ 87. $(-\infty, 20] \cup [30, \infty)$
89. $\{-\frac{5}{3}, \frac{1}{3}\}$ 91. $\{-1, 3\}$ 93. $\{-3, \frac{5}{3}\}$ 95. $\{-\frac{1}{3}, -\frac{1}{15}\}$ 97. $\{-\frac{5}{4}\}$
99. $(-\infty, \infty)$ 101. \emptyset 103. $\{-\frac{1}{4}\}$ 105. \emptyset 107. $(-\infty, \infty)$
109. $\{-\frac{3}{7}\}$ 111. $\{\frac{2}{5}\}$ 113. $(-\infty, \infty)$ 115. \emptyset 117. between 60.8 and 67.2 oz, inclusive 119. between 31.36 and 32.64 oz, inclusive
121. $(-0.05, 0.05)$ 123. $(2.74975, 2.75025)$ 125. between 6.8 and 9.8 lb 127. $|x - 1000| \leq 100$; $900 \leq x \leq 1100$
129. 814.8 ft 130. Bank of America Center

131. Williams Tower, Bank of America Center, Texaco Heritage Plaza, 609 Main at Texas, Enterprise Plaza, Centerpoint Energy Plaza, 1600 Smith St., Fulbright Tower 132. (a) $|x - 814.8| \geq 95$ (b) $x \geq 909.8$ or $x \leq 719.8$ (c) JPMorgan Chase Tower, Wells Fargo Plaza (d) It makes sense because it includes all buildings *not* listed in the answer to Exercise 131.

SUMMARY EXERCISES Solving Linear and Absolute Value Equations and Inequalities

1. $\{12\}$ 2. $\{-5, 7\}$ 3. $\{7\}$ 4. $\{-\frac{2}{5}\}$ 5. \emptyset 6. $(-\infty, -1]$
 7. $[-\frac{2}{3}, \infty)$ 8. $\{-1\}$ 9. $\{-3\}$ 10. $\{1, \frac{11}{3}\}$ 11. $(-\infty, 5]$
 12. $(-\infty, \infty)$ 13. $\{2\}$ 14. $(-\infty, -8] \cup [8, \infty)$ 15. \emptyset 16. $(-\infty, \infty)$
 17. $(-5.5, 5.5)$ 18. $\{\frac{13}{3}\}$ 19. $\{-\frac{96}{5}\}$ 20. $(-\infty, 32]$ 21. $(-\infty, -24)$
 22. $[\frac{3}{4}, \frac{15}{8}]$ 23. $[\frac{7}{2}]$ 24. $\{60\}$ 25. $\{\text{all real numbers}\}$ 26. $(-\infty, 5)$
 27. $[-\frac{9}{2}, \frac{15}{2}]$ 28. $\{24\}$ 29. $\{-\frac{1}{5}\}$ 30. $(-\infty, -\frac{5}{2}]$ 31. $[-\frac{1}{3}, 3]$
 32. $[1, 7]$ 33. $\{-\frac{1}{6}, 2\}$ 34. $\{-3\}$ 35. $(-\infty, -1] \cup [\frac{5}{3}, \infty)$
 36. $\{\frac{3}{8}\}$ 37. $\{-\frac{5}{2}\}$ 38. $(-6, 8)$ 39. $(-\infty, -4) \cup (7, \infty)$
 40. $(1, 9)$ 41. $(-\infty, \infty)$ 42. $\{\frac{1}{3}, 9\}$ 43. $\{\text{all real numbers}\}$
 44. $[-\frac{10}{9}]$ 45. $\{-2\}$ 46. \emptyset 47. $(-\infty, -1) \cup (2, \infty)$ 48. $[-3, -2]$

Chapter 1 Review Exercises





1. $\{-\frac{9}{5}\}$ 2. $\{16\}$ 3. $\{-\frac{7}{5}\}$ 4. \emptyset 5. $\{\text{all real numbers}\}$;
 identity 6. \emptyset ; contradiction 7. $\{0\}$; conditional equation
 8. $\{\text{all real numbers}\}$; identity 9. $L = \frac{V}{HW}$ 10. $b = \frac{2sA - Bh}{h}$, or
 $b = \frac{2sA}{h} - B$ 11. $c = P - a - b - B$ 12. $y = -\frac{4}{7}x + \frac{9}{7}$ 13. 6 ft
 14. 14.4% 15. 3.25% 16. 25° 17. 138.6 million subscriptions
 18. 67.3 million subscriptions 19. $9 - \frac{1}{3}x$ 20. $\frac{4x}{x+9}$ 21. length: 13 m;
 width: 8 m 22. 17 in., 17 in., 19 in. 23. 12 kg 24. 30 L 25. 10 L
 26. \$10,000 at 6%; \$6000 at 4% 27. 15 dimes; 8 quarters 28. 7 nickels;
 12 dimes 29. C 30. 530 mi 31. 328 mi 32. 2.2 hr 33. 50 km
 per hr; 65 km per hr 34. 46 mph 35. $40^\circ, 45^\circ, 95^\circ$ 36. $150^\circ, 30^\circ$
 37. 48, 49, 50 38. 38, 40, 42 39. $(-9, \infty)$ 40. $(-\infty, -3]$
 41. $(\frac{3}{2}, \infty)$ 42. $[-3, \infty)$ 43. $[3, 5)$ 44. $(\frac{59}{31}, \infty)$ 45. (a) $x < 6.75$
 (b) $6.75 \leq x \leq 7.25$ (c) $x > 7.25$ 46. 38 m or less 47. 31 tickets or
 less (but at least 15) 48. any score greater than or equal to 61
 49. $\{a, c\}$ 50. $\{a\}$ 51. $\{a, c, e, f, g\}$ 52. $\{a, b, c, d, e, f, g\}$
 53.  (6, 9) 54.  (8, 14)
 55.  $(-\infty, -3] \cup (5, \infty)$ 56.  $(-\infty, \infty)$
 57. \emptyset
 58.  $(-\infty, -2] \cup [7, \infty)$ 59. $(-3, 4]$ 60. $(-\infty, 2]$
 61. $(4, \infty)$ 62. $(1, \infty)$
 63. $\{-7, 7\}$ 64. $\{-11, 7\}$ 65. $\{-\frac{1}{3}, 5\}$ 66. \emptyset 67. $\{0, 7\}$

68. $\{-\frac{3}{2}, \frac{1}{2}\}$ 69. $\{-\frac{3}{4}, \frac{1}{2}\}$ 70. $\{-\frac{1}{2}\}$ 71. $(-14, 14)$ 72. $[-1, 13]$
 73. $[-3, -2]$ 74. $(-\infty, \infty)$ 75. \emptyset 76. $(-\infty, \infty)$ 77. between
 46.56 and 49.44 oz, inclusive 78. between 31.68 and 32.32 oz, inclusive





Chapter 1 Mixed Review Exercises

1. $(-2, \infty)$ 2. $k = \frac{6r - bt}{a}$ 3. $[-2, 3)$ 4. $\{0\}$ 5. $(-\infty, \infty)$
 6. $(-\infty, 2]$ 7. $\{-\frac{7}{3}, 1\}$ 8. $\{300\}$ 9. $[-16, 10]$ 10. $(-\infty, \frac{14}{17})$
 11. $(-3, \frac{7}{2})$ 12. $(-\infty, 1]$ 13. $(-\infty, -\frac{13}{5}) \cup (3, \infty)$ 14. $(-\infty, \infty)$
 15. $\{-4, -\frac{2}{3}\}$ 16. $\{30\}$ 17. $\{1, \frac{11}{3}\}$ 18. \emptyset 19. $(6, 8)$
 20. $(-\infty, -2] \cup [7, \infty)$ 21. 10 ft 22. 46, 47, 48 23. any amount
 greater than or equal to \$1100 24. 5 L

Chapter 1 Test

- [1.1] 1. $\{-19\}$ 2. $\{5\}$ 3. (a) \emptyset ; contradiction
 (b) $\{\text{all real numbers}\}$; identity (c) $\{0\}$; conditional equation
 (d) $\{\text{all real numbers}\}$; identity [1.2] 4. $h = \frac{3V}{b}$ 5. $y = \frac{3}{2}x + 3$
 [1.3, 1.4] 6. 3.497 hr 7. 1.25% 8. (a) .562 (b) .494 (c) .469
 (d) .463 9. \$8000 at 1.5%; \$20,000 at 2.5% 10. 13.3% 11. 80 L
 12. faster car: 60 mph; slower car: 45 mph 13. $40^\circ, 40^\circ, 100^\circ$
 [1.5]
 14.  $[1, \infty)$ 15.  $(-\infty, 28]$
 16.  $(1, 2)$ 17.  $[-3, 3]$
 18. C 19. 82 or more [1.6] 20. (a) $\{1, 5\}$ (b) $\{1, 2, 5, 7, 9, 12\}$
 21. $[2, 9)$ 22. $(-\infty, 3) \cup [6, \infty)$ [1.7] 23. $[-\frac{5}{2}, 1]$
 24. $(-\infty, -\frac{7}{6}) \cup (\frac{17}{6}, \infty)$ 25. $\{1, 5\}$ 26. $(\frac{1}{3}, \frac{7}{3})$ 27. \emptyset 28. $\{-\frac{5}{3}, 3\}$
 29. $\{-\frac{5}{7}, \frac{11}{3}\}$ 30. (a) \emptyset (b) $(-\infty, \infty)$ (c) \emptyset

Chapters R and 1 Cumulative Review Exercises


- [R.1] 1. (a) 0.35 (b) 35% 2. (a) 66% (b) $\frac{33}{50}$ 3. $\frac{37}{60}$
 4. -7.865 [R.3, R.4] 5. -8 6. 10 [R.4] 7. -125 8. $\frac{81}{16}$
 9. -34 10. $\frac{3}{16}$ [R.5] 11. $-20r + 17$ 12. $13k + 42$
 [1.1] 13. $\{-12\}$ 14. $\{26\}$ [1.7] 15. $\{\frac{3}{4}, \frac{7}{2}\}$ [1.2] 16. $n = \frac{A - P}{iP}$
 [1.5-1.7]
 17.  $[-14, \infty)$ 18.  $[\frac{5}{3}, 3)$
 19.  $(-\infty, 0) \cup (2, \infty)$ 20.  $(-\infty, -\frac{1}{7}] \cup [1, \infty)$
 [1.3] 21. \$5000 [1.5] 22. $6\frac{1}{3}$ g 23. 74 or greater
 [R.3] 24. 11,331 ft [1.3] 25. 4 cm; 9 cm; 27 cm

2 LINEAR EQUATIONS, GRAPHS, AND FUNCTIONS


Section 2.1

1. origin 3. y ; x ; x ; y 5. two; 2; 6 7. (a) x represents the year; y represents personal spending on medical care in billions of dollars.
 (b) about \$2830 billion (c) (2016, 2830) (d) 2010
 9. (a) I (b) III (c) II (d) IV (e) none (f) none
 11. (a) I or III (b) II or IV (c) II or IV (d) I or III
 13. The student interchanged the x - and y -coordinates. To plot this point correctly, move from 0 to the left 4 units on the x -axis and then up 2 units parallel to the y -axis.


15.-23.




25. -4 ; -3 ; -2 ; -1 ; 0




27. -3 ; 3 ; 2 ; -1



29. $\frac{5}{2}$; 5 ; $\frac{3}{2}$; 1




31. -4 ; 5 ; $-\frac{12}{5}$; $\frac{5}{4}$




33. (a) C (b) D (c) B (d) A


35. $(6, 0)$; $(0, 4)$




37. $(6, 0)$; $(0, -2)$




39. $(-2, 0)$; $(0, -\frac{5}{3})$




41. $(\frac{21}{2}, 0)$; $(0, -\frac{7}{3})$




43. $(0, 0)$; $(0, 0)$




45. $(0, 0)$; $(0, 0)$




47. $(0, 0)$; $(0, 0)$




49. $(0, 0)$; $(0, 0)$




51. none; $(0, 5)$




53. $(2, 0)$; none





55. $(-4, 0)$; none



57. none; $(0, -2)$



59. (a) $(-2, 0)$; $(0, 3)$ (b) B 61. (a) none; $(0, -1)$ (b) A




63. $(-5, -1)$ 65. $(\frac{9}{2}, -\frac{3}{2})$ 67. $(0, \frac{11}{2})$ 69. $(2.1, 0.9)$ 71. $(1, 1)$
 73. $(-\frac{5}{12}, \frac{5}{28})$ 75. $Q(11, -4)$ 77. $Q(4.5, 0.75)$


Section 2.2

1. A, B, D, F 3. 5 ft 5. (a) C (b) A (c) D (d) B 7. 2
 9. undefined 11. 1 13. -1 15. (a) B (b) C (c) A (d) D 17. The x -values in the denominator should be subtracted in the same order as the y -values in the numerator. The denominator should be $2 - (-4)$. The correct slope is $\frac{2}{6} = \frac{1}{3}$. 19. 2 21. 0
 23. undefined 25. (a) 8 (b) rises 27. (a) $\frac{5}{6}$ (b) rises
 29. (a) 0 (b) horizontal 31. (a) $-\frac{1}{2}$ (b) falls 33. (a) undefined (b) vertical 35. (a) -1 (b) falls 37. 6 39. -3 41. -2
 43. $\frac{4}{3}$ 45. $-\frac{5}{2}$ 47. undefined
 In part (a) of Exercises 49-53, we used the intercepts. Other points can be used.
 49. (a) $(4, 0)$ and $(0, -8)$; 2 (b) $y = 2x - 8$; 2 (c) $A = 2, B = -1$; 2
 51. (a) $(4, 0)$ and $(0, 3)$; $-\frac{3}{4}$ (b) $y = -\frac{3}{4}x + 3$; $-\frac{3}{4}$
 (c) $A = 3, B = 4$; $-\frac{3}{4}$ 53. (a) $(-3, 0)$ and $(0, -3)$; -1
 (b) $y = -x - 3$; -1 (c) $A = 1, B = 1$; -1


55. $-\frac{1}{2}$




57. $\frac{5}{2}$




59. 4




61. undefined




63. 0




65. 0




67. $(-2, 3)$; $(-4, 2)$




69.




71.




73. $(0, 0)$; $(5, 1)$



75.



77.




79. $-\frac{4}{9}$; $\frac{9}{4}$ 81. parallel 83. perpendicular 85. neither
 87. parallel 89. neither 91. perpendicular 93. perpendicular
 95. $\frac{7}{10}$ 97. $-\$4000$ per year; The value of the machine is decreasing $\$4000$ per year during these years. 99. 0% per year (or no change); The percent of pay raise is not changing—it is 3% per year during these years. 101. (a) In 2016, there were 396 million wireless subscriber connections in the U.S.

101. (b) 16 (c) The number of subscribers increased by an average of 16 million per year from 2011 to 2016. 103. (a) -9 theaters per yr (b) The negative slope means that the number of drive-in theaters decreased by an average of 9 per year from 2010 to 2017. 105. \$0.04 per year; The price of a gallon of gasoline increased by an average of \$0.04 per year from 2000 to 2016. 107. -16 million digital cameras per year; The number of digital cameras sold decreased by an average of 16 million per year from 2010 to 2016. 109. Because the slopes of both pairs of opposite sides are equal, the figure is a parallelogram. 111. $\frac{1}{3}$ 112. $\frac{1}{3}$ 113. $\frac{1}{3}$ 114. $\frac{1}{3} = \frac{1}{3} = \frac{1}{3}$ is true. 115. collinear 116. not collinear


Section 2.3

1. A 3. A 5. $3x + y = 10$ 7. A 9. C 11. H 13. B
 15. $y = 5x + 15$ 17. $y = -\frac{2}{3}x + \frac{4}{5}$ 19. $y = x - 1$
 21. $y = \frac{2}{5}x + 5$ 23. $y = \frac{2}{3}x + 1$ 25. $y = -x - 2$ 27. $y = 2x - 4$
 29. $y = -\frac{3}{5}x + 3$


31. (a) $y = x + 4$ (b) 1
 (c) (0, 4)




33. (a) $y = -\frac{6}{5}x + 6$ (b) $-\frac{6}{5}$
 (c) (0, 6)



35. (a) $y = \frac{4}{5}x - 4$ (b) $\frac{4}{5}$
 (c) (0, -4)



37. (a) $y = -\frac{1}{2}x - 2$ (b) $-\frac{1}{2}$
 (c) (0, -2)



39. (a) $y = -2x + 18$ (b) $2x + y = 18$ 41. (a) $y = -\frac{3}{4}x + \frac{5}{2}$
 (b) $3x + 4y = 10$ 43. (a) $y = \frac{1}{2}x + \frac{13}{2}$ (b) $x - 2y = -13$
 45. (a) $y = 4x - 12$ (b) $4x - y = 12$ 47. (a) $y = 1.4x + 4$
 (b) $7x - 5y = -20$ 49. $2x - y = 2$ 51. $x + 2y = 8$ 53. $y = 5$
 55. $x = 7$ 57. $y = -3$ 59. $2x - 13y = -6$ 61. $y = 5$
 63. $x = 9$ 65. $y = -\frac{3}{2}$ 67. $y = 8$ 69. $x = 0.5$ 71. $y = 0$
 73. (a) $y = 3x - 19$ (b) $3x - y = 19$ 75. (a) $y = \frac{1}{2}x - 1$
 (b) $x - 2y = 2$ 77. (a) $y = -\frac{1}{2}x + 9$ (b) $x + 2y = 18$
 79. (a) $y = 7$ (b) $y = 7$ 81. $y = 45x$; (0, 0), (5, 225), (10, 450)
 83. $y = 3.75x$; (0, 0), (5, 18.75), (10, 37.50) 85. $y = 150x$; (0, 0), (5, 750), (10, 1500) 87. (a) $y = 140x + 18.50$ (b) (5, 718.50);
 The cost for 5 tickets and the delivery fee is \$718.50. (c) \$298.50
 89. (a) $y = 41x + 99$ (b) (5, 304); The cost for a 5-month membership is \$304. (c) \$591 91. (a) $y = 90x + 25$ (b) (5, 475);
 The cost of the plan for 5 months is \$475. (c) \$2185
 93. (a) $y = 6x + 30$ (b) (5, 60); It costs \$60 to rent the saw for 5 days.
 (c) 18 days 95. (a) $y = -79.7x + 716$; Sales of e-readers in the United States decreased by \$79.7 million per year from 2013 to 2016.

- (b) \$636.3 million 97. (a) The slope would be positive because spending on home health care is increasing over these years.
 (b) $y = 3.5x + 36$ (c) \$88.5 billion; It is very close to the actual value. 99. 32; 212 100. (a) (0, 32) and (100, 212)
 (b) $\frac{9}{5}$ 101. $F = \frac{9}{5}C + 32$ 102. $C = \frac{5}{9}(F - 32)$; -40°
 103. 60° 104. 59° ; They differ by 1° . 105. 90° ; 86° ; They differ by 4° . 106. Because $\frac{9}{5}$ is a little less than 2, and 32 is a little more than 30, $\frac{9}{5}C + 32 \approx 2C + 30$.




SUMMARY EXERCISES Finding Slopes and Equations of Lines

1. $-\frac{3}{5}$ 2. 0 3. 1 4. $\frac{3}{7}$ 5. undefined 6. $-\frac{4}{7}$ 7. (a) B
 (b) F (c) A (d) C (e) E (f) D 8. C is in standard form;
 A: $4x + y = -7$; B: $3x - 4y = -12$; D: $x + 2y = 0$;
 E: $3x - y = 5$; F: $5x - 3y = 15$ 9. (a) $y = -3x + 10$
 (b) $3x + y = 10$ 10. (a) $y = \frac{2}{3}x + 8$ (b) $2x - 3y = -24$
 11. (a) $y = -\frac{5}{6}x + \frac{13}{3}$ (b) $5x + 6y = 26$ 12. (a) $y = -\frac{5}{2}x + 2$
 (b) $5x + 2y = 4$ 13. (a) $y = 3x + 11$ (b) $3x - y = -11$
 14. (a) $y = -\frac{5}{2}x$ (b) $5x + 2y = 0$ 15. (a) $y = -8$
 (b) $y = -8$ 16. (a) $y = -\frac{7}{9}$ (b) $9y = -7$
 17. (a) $y = 2x - 10$ (b) $2x - y = 10$ 18. (a) $y = \frac{2}{3}x + \frac{14}{3}$
 (b) $2x - 3y = -14$ 19. (a) $y = \frac{1}{5}x - \frac{7}{5}$ (b) $x - 5y = 7$
 20. (a) $y = -\frac{3}{4}x - 6$ (b) $3x + 4y = -24$


Section 2.4

1. (a) yes (b) yes (c) no (d) yes 3. (a) no (b) no (c) no
 (d) yes 5. solid; below 7. dashed; above 9. \leq 11. $>$
 13. 15. 17. 19. 21. 23. 25. 27. 29. 31. 33. 35.





37. 2; (0, -4); $2x - 4$; solid; above; \geq ; $\geq 2x - 4$




45. $-3 < x < 3$



47. $-2 < x + 1 < 2$
 $-3 < x < 1$



55. $x \leq 200$,
 $x \geq 100$,
 $y \geq 3000$




57. $C = 50x + 100y$ 58. Some examples are (100, 5000), (150, 3000), and (150, 5000). The corner points are (100, 3000) and (200, 3000). 59. The least value occurs when $x = 100$ and $y = 3000$. 60. The company

should use 100 workers and manufacture 3000 units to achieve the least possible cost.

Section 2.5

1. relation; ordered pairs 3. domain; range 5. independent variable; dependent variable 7. $\{(2, -2), (2, 0), (2, 1)\}$ 9. $\{(1960, 0.76), (1980, 2.69), (2000, 5.39), (2016, 8.65)\}$ 11. $\{(A, 4), (B, 3), (C, 2), (D, 1), (F, 0)\}$

In Exercises 13–17, answers will vary.




15. $\{(-1, -3), (0, -1), (1, 1), (3, 3)\}$
17. $y = 2$ 19. function; domain: $\{5, 3, 4, 7\}$;
range: $\{1, 2, 9, 6\}$ 21. not a function; domain:
 $\{2, 0\}$; range: $\{4, 2, 5\}$

23. function; domain: $\{-3, 4, -2\}$; range: $\{1, 7\}$
25. not a function; domain: $\{1, 0, 2\}$; range: $\{1, -1, 0, 4, -4\}$
27. not a function; domain: $\{1\}$; range: $\{5, 2, -1, -4\}$ 29. function;
domain: $\{4, 2, 0, -2\}$; range: $\{-3\}$ 31. function; domain:
 $\{2, 5, 11, 17, 3\}$; range: $\{1, 7, 20\}$ 33. function; domain: $\{-2, 0, 3\}$;
range: $\{2, 3\}$ 35. function; domain: $(-\infty, \infty)$; range: $(-\infty, \infty)$
37. not a function; domain: $\{-2\}$; range: $(-\infty, \infty)$ 39. function;
domain: $(-\infty, \infty)$; range: $\{-2\}$ 41. not a function; domain: $(-\infty, 0]$;
range: $(-\infty, \infty)$ 43. function; domain: $(-\infty, \infty)$; range: $(-\infty, 4]$
45. not a function; domain: $[-4, 4]$; range: $[-3, 3]$ 47. not a
function; domain: $(-\infty, \infty)$; range: $[2, \infty)$ 49. function; $(-\infty, \infty)$
51. function; $(-\infty, \infty)$ 53. function; $(-\infty, \infty)$ 55. not a function;
 $[0, \infty)$ 57. not a function; $(-\infty, \infty)$ 59. function; $[0, \infty)$
61. function; $[3, \infty)$ 63. function; $[-\frac{1}{2}, \infty)$ 65. function; $(-\infty, \infty)$
67. function; $(-\infty, 0) \cup (0, \infty)$ 69. function; $(-\infty, 4) \cup (4, \infty)$


71. function; $(-\infty, 0) \cup (0, \infty)$ 73. (a) yes (b) domain:
 $\{2013, 2014, 2015, 2016, 2017\}$; range: $\{52.8, 52.6, 53.2, 53.7\}$
(c) 52.6; 2017 (d) Answers will vary. Two possible answers are
(2014, 52.6) and (2017, 53.7).

Section 2.6


1. $f(x)$; function; domain; x ; f of x (or “ f at x ”) 3. line; -2 ; linear;
 $-2x + 4$; -2 ; 3; -2 5. 4 7. 13 9. -11 11. 4
13. -296 15. 3 17. 2.75 19. $-3p + 4$ 21. $3x + 4$
23. $-3x - 2$ 25. $-6t + 1$ 27. $-\pi^2 + 4\pi + 1$
29. $-3x - 3h + 4$ 31. $-\frac{p^2}{9} + \frac{4p}{3} + 1$ 33. (a) -1 (b) -1
35. (a) 2 (b) 3 37. (a) 15 (b) 10 39. (a) 4 (b) 1
41. (a) 3 (b) -3 43. (a) -3 (b) 2 45. (a) 2
(b) 0 (c) -1 47. (a) $f(x) = -\frac{1}{3}x + 4$ (b) 3
49. (a) $f(x) = 3 - 2x^2$ (b) -15 51. (a) $f(x) = \frac{4}{3}x - \frac{8}{3}$ (b) $\frac{4}{3}$




domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$




domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$




domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$



domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$



domain: $(-\infty, \infty)$;
range: $\{-4\}$





domain: $(-\infty, \infty)$;
range: $\{0\}$


65. x -axis 67. (a) 11.25 (dollars) (b) 3 is the value of the independent
variable, which represents a package weight of 3 lb. $f(3)$ is the value of the
dependent variable, which represents the cost to mail a 3-lb package.
(c) \$18.75; $f(5) = 18.75$ 69. (a) $f(x) = 12x + 100$ (b) 1600;
The cost to print 125 t-shirts is \$1600. (c) 75; $f(75) = 1000$; The cost
to print 75 t-shirts is \$1000. 71. (a) 1.1 (b) 4 (c) -1.2
(d) (0, 3.5) (e) $f(x) = -1.2x + 3.5$ 73. (a) $[0, 100]$; $[0, 3000]$
(b) 25 hr; 25 hr (c) 2000 gal (d) $f(0) = 0$; The pool is empty at
time 0. (e) $f(25) = 3000$; After 25 hr, there are 3000 gal of water
in the pool. 75. (a) 194.53 cm (b) 177.29 cm (c) 177.41 cm
(d) 163.65 cm 77. Because it falls from left to right, the slope
is negative. 78. $-\frac{3}{2}$ 79. $-\frac{3}{2}; \frac{2}{3}$ 80. $(\frac{7}{3}, 0)$ 81. $(0, \frac{7}{2})$
82. $f(x) = -\frac{3}{2}x + \frac{7}{2}$ 83. $-\frac{17}{2}$ 84. $\frac{23}{3}$ 85. 2; $f(1) = -\frac{3}{2}(1) + \frac{7}{2}$,
which simplifies on the right to $f(1) = -\frac{3}{2} + \frac{7}{2}$. This gives $f(1) = 2$.

Chapter 2 Review Exercises


1. $5; \frac{10}{3}; 2; \frac{14}{3}$ 2. $-6; 5; -5; 6$




3. (3, 0); (0, -4)




4. $(\frac{28}{5}, 0)$; (0, 4)



5. (10, 0); (0, 4)



6. (8, 0); (0, -2)



7. (0, 2) 8. $(-\frac{9}{2}, \frac{3}{2})$ 9. $-\frac{7}{5}$ 10. $-\frac{1}{2}$ 11. undefined 12. 2

13. $\frac{3}{4}$ 14. 0 15. $-\frac{1}{3}$ 16. $\frac{2}{3}$ 17. $-\frac{1}{3}$ 18. -1 19. -3

20. (a) positive (b) negative (c) undefined (d) 0

21. perpendicular 22. parallel 23. 12 ft 24. \$1057 per year

25. (a) $y = \frac{3}{5}x - 8$ (b) $3x - 5y = 40$ 26. (a) $y = -\frac{1}{3}x + 5$

(b) $x + 3y = 15$ 27. (a) $y = -9x + 13$ (b) $9x + y = 13$




28. (a) $y = \frac{7}{5}x + \frac{16}{5}$ (b) $7x - 5y = -16$ 29. (a) $y = 4x - 26$

(b) $4x - y = 26$ 30. (a) $y = -\frac{5}{2}x + 1$ (b) $5x + 2y = 2$

31. $y = 12$ 32. $x = 2$ 33. $x = 0.3$ 34. $y = 4$

35. (a) $y = 229x + 2829$ (b) \$11,760 36. (a) $y = 1007.5x + 29,822.5$;

The cost at private 4-year universities increased by an average of \$1007.50 per year from 2013 to 2017. (b) \$44,935



40.

41. not a function; domain: $\{-4, 1\}$; range:

$\{2, -2, 5, -5\}$ 42. function; domain:

$\{9, 11, 4, 17, 25\}$; range: $\{32, 47, 69, 14\}$

43. function; domain: $[-4, 4]$; range: $[0, 2]$

44. not a function; domain: $(-\infty, 0]$; range: $(-\infty, \infty)$ 45. function; domain: $(-\infty, \infty)$; linear function 46. not a function; domain: $(-\infty, \infty)$

47. function; domain: $(-\infty, \infty)$ 48. function; domain: $[-\frac{7}{4}, \infty)$

49. not a function; domain: $[0, \infty)$ 50. function; domain:

$(-\infty, 6) \cup (6, \infty)$ 51. -6 52. -15 53. -8 54. $-2k^2 + 3k - 6$

55. $f(x) = 2x^2; 18$ 56. C 57. It is a horizontal line. 58. (a) yes

(b) domain: $\{1960, 1970, 1980, 1990, 2000, 2010, 2015\}$; range:

$\{69.7, 70.8, 73.7, 75.4, 76.8, 78.7, 78.8\}$ (c) Answers will vary.

Two possible answers are (1960, 69.7) and (2010, 78.7). (d) 73.7;

In 1980, life expectancy at birth was 73.7 yr. (e) 2000

Chapter 2 Mixed Review Exercises

1. parallel 2. perpendicular 3. -0.575 lb per year; Per capita consumption of potatoes decreased by an average of 0.575 lb per year from 2008 to 2016. 4. $y = -0.575x + 37.8$ 5. $y = 3x$


6. $x + 2y = 6$ 7. $y = -3$ 8. A, B, D 9. D 10. (a) -1

(b) -2 (c) 2 (d) $(-\infty, \infty); (-\infty, \infty)$


Chapter 2 Test

[2.1] 1. $-\frac{10}{3}; -2; 0$


2. $(-3, 0); (0, 4)$



3. (2, 0); none



4. (0, 0); (0, 0)



[2.2] 5. $\frac{1}{2}$ 6. It is a vertical line. [2.1, 2.2] 7. $\frac{3}{2}; (\frac{13}{3}, 0); (0, -\frac{13}{2})$

8. 0; none; (0, 5) [2.2] 9. B 10. perpendicular 11. neither

12. -438 farms per yr; The number of farms decreased an average of



438 farms per year from 2000 to 2016. [2.3] 13. (a) $y = -\frac{2}{5}x + 3$

(b) $2x + 5y = 15$ 14. (a) $y = -\frac{1}{2}x + 2$ (b) $x + 2y = 4$

15. (a) $y = -5x + 19$ (b) $5x + y = 19$ 16. $y = 14$ 17. $x = 5$

18. $y = 0$ 19. (a) $y = -\frac{3}{5}x - \frac{11}{5}$ (b) $y = -\frac{1}{2}x - \frac{3}{2}$

20. (a) $y = 142.75x + 45$ (b) \$901.50



[2.5] 23. D; domain: $(-\infty, \infty)$; range: $[0, \infty)$

24. D; domain: $\{0, 3, 6\}$; range: $\{1, 2, 3\}$ [2.6] 25. 0; $-a^2 + 2a - 1$

26. domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$

Chapters R-2 Cumulative Review Exercises

[R.1] 1. (a) $\frac{11}{12}$ (b) $\frac{2}{3}$ (c) 4.75 2. (a) 8 (b) 150 (c) 0.125

3. (a) 0.01; 1% (b) $\frac{1}{2}$; 50% (c) $\frac{3}{4}$; 0.75 (d) 1; 100%

[R.2] 4. always true 5. never true 6. sometimes true; For example, $3 + (-3) = 0$, but $3 + (-1) = 2$ and $2 \neq 0$. [R.3] 7. -3

[R.4] 8. 0.64 9. -9 10. not a real number [R.4, R.5] 11. -39


12. undefined 13. $4x - 3$ 14. 0 [1.1] 15. $\{\frac{7}{6}\}$ 16. $\{-1\}$

[1.4] 17. 6 in. 18. 2 hr [1.5] 19. $(-3, \frac{7}{2})$ 20. $(-\infty, 1]$


[1.6] 21. (6, 8) 22. $(-\infty, -2] \cup (7, \infty)$ [1.7] 23. $\{0, 7\}$

24. $(-\infty, \infty)$

[2.1] 25. $(4, 0); (0, \frac{12}{5})$



[2.2] 26. (a) $-\frac{6}{5}$ (b) $\frac{5}{6}$



[2.3] 28. (a) $y = -\frac{3}{4}x - 1$ (b) $3x + 4y = -4$


29. (a) $y = -\frac{4}{3}x + \frac{7}{3}$ (b) $4x + 3y = 7$

[2.5] 30. domain: $\{14, 91, 75, 23\}$; range: $\{9, 70, 56, 5\}$; not a function

3 SYSTEMS OF LINEAR EQUATIONS

Section 3.1

1. 4; -3 3. \emptyset 5. no 7. D; The ordered-pair solution must be in quadrant IV because that is where the graphs of the equations intersect.
 9. (a) B (b) C (c) A (d) D 11. solution 13. not a solution
 15. $\{(-2, -3)\}$ 17. $\{(0, 1)\}$ 19. $\{(2, 0)\}$



21. $\{(1, 2)\}$ 23. $\{(2, 3)\}$ 25. $\{(1, 3)\}$ 27. $\{(0, -4)\}$
 29. $\left\{\left(\frac{22}{9}, \frac{22}{3}\right)\right\}$ 31. $\{(5, 4)\}$ 33. $\left\{\left(-5, -\frac{10}{3}\right)\right\}$ 35. $\{(2, 6)\}$
 37. $\{(4, 2)\}$ 39. $\{(-8, 4)\}$ 41. $\{(x, y) \mid 2x - y = 0\}$; dependent equations
 43. \emptyset ; inconsistent system 45. $\{(2, -4)\}$ 47. $\{(3, -1)\}$
 49. $\{(2, -3)\}$ 51. $\{(x, y) \mid 7x + 2y = 6\}$; dependent equations
 53. $\left\{\left(\frac{3}{2}, -\frac{3}{2}\right)\right\}$ 55. \emptyset ; inconsistent system 57. $\{(0, 0)\}$
 59. $\{(0, -4)\}$ 61. $\left\{\left(6, -\frac{5}{6}\right)\right\}$ 63. (a) Use elimination because the coefficients of the y-terms are opposites. (b) Use substitution because the second equation is solved for y. (c) Use elimination because the equations are in standard form with no coefficients of 1 or -1. Solving by substitution would involve fractions. 65. $\{(-3, 2)\}$
 67. $\left\{\left(\frac{1}{3}, \frac{1}{2}\right)\right\}$ 69. $\{(-4, 6)\}$ 71. $\{(5, 0)\}$ 73. $\left\{\left(1, \frac{1}{2}\right)\right\}$
 75. $\{(x, y) \mid 4x - y = -2\}$ 77. $y = -\frac{3}{7}x + \frac{4}{7}; y = -\frac{3}{7}x + \frac{3}{14}$; no solution 79. Both are $y = -\frac{2}{3}x + \frac{1}{3}$; infinitely many solutions
 81. (a) \$4 (b) 300 half-gallons (c) supply: 200 half-gallons; demand: 400 half-gallons 83. (a) 2011–2015 (b) 2015 and 2016 (c) 2013; 260 thousand students (d) (2013, 260) (e) Enrollment in Kentucky was decreasing, and enrollment in Utah remained constant. 85. Netflix
 87. (4.4, 48.7) 89. $\{(2, 4)\}$ 91. $\left\{\left(\frac{1}{2}, 2\right)\right\}$ 93. $\left\{\left(\frac{c}{a}, 0\right)\right\}$
 95. $\left\{\left(-\frac{1}{a}, -5\right)\right\}$

Section 3.2

1. Answers will vary. Some possible answers are (a) two perpendicular walls and the ceiling in a normal room, (b) the floors of three different levels of an office building, and (c) three pages of a book (because they intersect in the spine). 3. The statement means that when -1 is substituted for x, 2 is substituted for y, and 3 is substituted for z in the three equations, the resulting three statements are true. 5. 4 7. $\{(3, 2, 1)\}$
 9. $\{(1, 4, -3)\}$ 11. $\{(0, 2, -5)\}$ 13. $\{(1, 0, 3)\}$ 15. $\left\{\left(1, \frac{3}{10}, \frac{2}{5}\right)\right\}$
 17. $\left\{\left(-\frac{7}{3}, \frac{22}{3}, 7\right)\right\}$ 19. $\{(-12, 18, 0)\}$ 21. $\{(0.8, -1.5, 2.3)\}$
 23. $\{(4, 5, 3)\}$ 25. $\{(2, 2, 2)\}$ 27. $\left\{\left(\frac{8}{3}, \frac{2}{3}, 3\right)\right\}$ 29. $\{(-1, 0, 0)\}$
 31. $\{(-4, 6, 2)\}$ 33. $\{(-3, 5, -6)\}$ 35. \emptyset ; inconsistent system


37. $\{(x, y, z) \mid x - y + 4z = 8\}$; dependent equations 39. $\{(3, 0, 2)\}$
 41. $\{(x, y, z) \mid 2x + y - z = 6\}$; dependent equations 43. $\{(0, 0, 0)\}$
 45. \emptyset ; inconsistent system 47. $\{(2, 1, 5, 3)\}$ 49. $\{(-2, 0, 1, 4)\}$
 51. $2a + b + c = -5$ 52. $a - c = 1$ 53. $3a + 3b + c = -18$
 54. $a = 1, b = -7, c = 0$ 55. $x^2 + y^2 + x - 7y = 0$ 56. The relation is not a function because it fails the vertical line test—that is, a vertical line can intersect its graph more than once.

Section 3.3

1. (a) 6 oz (b) 15 oz (c) 24 oz (d) 30 oz 3. \$1.99x
 5. (a) $(10 - x)$ mph (b) $(10 + x)$ mph 7. (a) 220 ft (b) $\frac{d}{44}$ sec
 9. length: 78 ft; width: 36 ft 11. wins: 102; losses: 60 13. AT&T: \$163.8 billion; Verizon: \$125.1 billion 15. $x = 40$ and $y = 50$, so the angles measure 40° and 50° . 17. Red Sox: \$360.66; Indians: \$179.44 19. ribeye: \$30.30; salmon: \$21.60 21. Busch Gardens: \$90; Universal Studios: \$110 23. general admission: 76; with student ID: 108 25. 25% alcohol: 6 gal; 35% alcohol: 14 gal 27. nuts: 14 kg; cereal: 16 kg 29. 2%: \$1000; 4%: \$2000 31. pure acid: 6 L; 10% acid: 48 L 33. \$1.75-per-lb candy: 7 lb; \$1.25-per-lb candy: 3 lb
 35. train: 60 km per hr; plane: 160 km per hr 37. freight train: 50 km per hr; express train: 80 km per hr 39. boat: 21 mph; current: 3 mph
 41. plane: 300 mph; wind: 20 mph 43. $x + y + z = 180; 70^\circ, 30^\circ, 80^\circ$ 45. $20^\circ, 70^\circ, 90^\circ$ 47. 12 cm, 25 cm, 33 cm 49. gold: 7; silver: 6; bronze: 6 51. upper level: 1170; center court: 985; floor: 130 53. bookstore A: 140; bookstore B: 280; bookstore C: 380
 55. first chemical: 50 kg; second chemical: 400 kg; third chemical: 300 kg
 57. wins: 46; losses: 23; overtime losses: 13 59. box of fish: 8 oz; box of bugs: 2 oz; box of worms: 5 oz

Chapter 3 Review Exercises


1. $\{(2, 2)\}$ 2. $\{(-4, 0)\}$ 3. $\{(1, -1)\}$



4. D 5. $\left\{\left(-\frac{8}{9}, -\frac{4}{3}\right)\right\}$ 6. $\{(0, 4)\}$ 7. $\{(2, 4)\}$ 8. $\{(2, 2)\}$
 9. $\{(2, -4)\}$ 10. $\{(0, 1)\}$ 11. $\{(-1, 2)\}$ 12. $\{(-6, 3)\}$
 13. $\{(x, y) \mid 3x - y = -6\}$; dependent equations 14. \emptyset ; inconsistent system
 15. $\{(1, -5, 3)\}$ 16. \emptyset ; inconsistent system 17. $\{(1, 2, 3)\}$
 18. $\{(5, -1, 0)\}$ 19. $\{(x, y, z) \mid 3x - 4y + z = 8\}$; dependent equations
 20. $\{(0, 0, 0)\}$ 21. length: 200 ft; width: 85 ft 22. plane: 300 mph; wind: 20 mph 23. gold: 12; silver: 8; bronze: 21 24. \$6-per-lb nuts: 30 lb; \$3-per-lb candy: 70 lb 25. 8% solution: 5 L; 20% solution: 3 L
 26. $85^\circ, 35^\circ, 60^\circ$ 27. 10%: \$40,000; 6%: \$100,000; 5%: \$140,000
 28. Mantle: 54; Maris: 61; Berra: 22

Chapter 3 Mixed Review Exercises

1. Answers will vary.




2. The two lines have the same slope, 3, but the y-intercepts, (0, 2) and (0, -4), are different. Therefore, the lines are parallel, do not intersect, and have no common solution. 3. $\{(0, 4)\}$ 4. $\left\{\left(\frac{82}{23}, -\frac{4}{23}\right)\right\}$
 5. $\{(x, y) \mid x + 2y = 48\}$ 6. $\{(5, 3)\}$ 7. $\{(12, 9)\}$ 8. $\{(3, -1)\}$
 9. $\{(1, 2, 3)\}$ 10. $\{(1, 0, -1)\}$ 11. \emptyset 12. B; The second equation is already solved for y. 13. 20 L 14. United States: 121; China: 70; Great Britain: 67

Chapter 3 Test

- [3.1] 1. (a) Houston, Phoenix, Dallas (b) Philadelphia (c) Dallas, Phoenix, Philadelphia, Houston 2. (a) 2010; 1.45 million (b) (2025, 2.8)



3. $\{(6, 1)\}$ 4. $\{(6, -4)\}$ 5. $\left\{\left(-\frac{9}{4}, \frac{5}{4}\right)\right\}$
 6. $\{(x, y) \mid 12x - 5y = 8\}$; dependent equations
 7. $\{(3, 3)\}$ 8. $\{(0, -2)\}$
 9. \emptyset ; inconsistent system [3.2] 10. $\left\{\left(-\frac{2}{3}, \frac{4}{5}, 0\right)\right\}$
 11. $\{(3, -2, 1)\}$ 12. \emptyset [3.3] 13. Captain



- America: \$408.1 million; Deadpool: \$363.1 million 14. slower car: 45 mph; faster car: 75 mph 15. 20% solution: 4 L; 50% solution: 8 L
 16. AC adaptor: \$8; rechargeable flashlight: \$15 17. 25°, 55°, 100°
 18. Orange Pekoe: 60 oz; Irish Breakfast: 30 oz; Earl Grey: 10 oz

Chapters R-3 Cumulative Review Exercises

- [R.4] 1. 81 2. -81 3. -81 4. 0.7 5. -0.7 6. It is not a real number. [R.3] 7. -7.17 8. $-\frac{37}{24}$, or $-1\frac{13}{24}$ 9. $-\frac{28}{33}$ [R.4] 10. -30
 11. -199 12. 455 [1.1] 13. $\left\{-\frac{15}{4}\right\}$ [1.7] 14. $\left\{\frac{2}{3}, 2\right\}$
 [1.2] 15. $x = \frac{d - by}{a}$ [1.1] 16. $\{11\}$ [1.5] 17. $\left(-\infty, \frac{240}{13}\right)$
 [1.7] 18. $\left[-2, \frac{2}{3}\right]$ 19. $(-\infty, \infty)$ [1.6] 20. $(-\infty, \infty)$
 [1.3, 1.4] 21. pennies: 35; nickels: 29; dimes: 30 22. 46°, 46°, 88°
 [R.1, 1.2] 23. 2010; 1813; 62.8%; 57.2% [2.1] 24. $y = 6$
 25. $x = 4$ [2.2] 26. $-\frac{4}{3}$ 27. $\frac{3}{4}$ [2.3] 28. $4x + 3y = 10$



- [2.6] 31. (a) -6 (b) $a^2 + 3a - 6$ [3.1] 32. $\{(3, -3)\}$

33. $\{(x, y) \mid x - 3y = 7\}$ [3.2] 34. $\{(5, 3, 2)\}$
 [3.1] 35. (a) $x = 8$, or 800 items; \$3000 (b) about \$400

4 EXPONENTS, POLYNOMIALS, AND POLYNOMIAL FUNCTIONS

Section 4.1

1. false; $(ab)^2 = a^2b^2$ 3. false; $\left(\frac{4}{a}\right)^3 = \frac{4^3}{a^3}$ 5. false; $xy^0 = x \cdot 1 = x$
 7. false; $-(-10)^0 = -(-1) = 1$ 9. The bases should not be multiplied; $4^5 \cdot 4^2 = 4^7$ 11. 13^{12} 13. 8^{10} 15. x^{17}
 17. The product rule does not apply. 19. $-27w^8$ 21. $18x^3y^8$
 23. (a) B (b) C (c) B (d) C 25. 1 27. -1 29. 1 31. -1
 33. 1 35. 2 37. 0 39. -2 41. (a) B (b) D (c) B (d) D
 43. $\frac{1}{5^4}$ 45. $\frac{1}{3^5}$ 47. $\frac{1}{9}$ 49. $\frac{1}{(4x)^2}$ 51. $\frac{4}{x^2}$ 53. $-\frac{1}{a^3}$ 55. $\frac{1}{(-a)^4}$, or $\frac{1}{a^4}$
 57. $\frac{1}{(-3x)^3}$ 59. $\frac{11}{30}$ 61. $\frac{9}{20}$ 63. $-\frac{5}{24}$ 65. 16 67. $\frac{27}{4}$ 69. $\frac{16}{25}$
 71. 4^2 73. x^4 75. $\frac{1}{r^3}$ 77. 6^6 79. $\frac{1}{6^{10}}$ 81. 7^2 83. r^3
 85. The quotient rule does not apply. 87. x^{18} 89. $\frac{27}{125}$ 91. $64r^3$
 93. $216x^6$ 95. $-\frac{64m^6}{t^3}$ 97. $\frac{s^{12}}{t^{20}}$ 99. (a) B (b) D (c) D (d) B
 101. 64 103. $\frac{27}{8}$ 105. $\frac{25}{16}$ 107. $\frac{81}{16t^4}$ 109. $16x^2$ 111. $\frac{32}{x^5}$ 113. $\frac{1}{3}$
 115. $\frac{1}{a^5}$ 117. 5^6 119. $\frac{1}{x^{12}}$ 121. $\frac{1}{k^2}$ 123. $-4r^6$ 125. $\frac{625}{a^{10}}$
 127. $\frac{z^4}{x^3}$ 129. $\frac{m^8}{n^{11}}$ 131. $\frac{1}{5p^{10}}$ 133. $\frac{1}{2pq}$ 135. $\frac{4}{a^2}$ 137. $\frac{1}{6y^{13}}$
 139. $\frac{4k^5}{m^2}$ 141. $\frac{8}{3pq^{10}}$ 143. $\frac{y^9}{8}$ 145. $\frac{n^{10}}{25m^{18}}$ 147. $-\frac{125y^3}{x^{30}}$
 149. $\frac{4k^{17}}{125}$ 151. $-\frac{3}{32m^8p^4}$

Section 4.2


1. after; power; a ; 10^a 3. A number in scientific notation should have only one nonzero digit to the left of the decimal point. The correct answer is 9.275×10^6 . 5. 1×10^6 7. 1×10^{12}
 9. 5.3×10^2 11. 8.3×10^{-1} 13. 6.92×10^{-6} 15. -3.85×10^4
 17. $\$1 \times 10^9$ (or $\$10^9$); $\$1 \times 10^{12}$ (or $\$10^{12}$); $\$4.094 \times 10^{12}$; 2.77843×10^5 19. 72,000 21. 0.00254 23. -60,000
 25. 0.000012 27. -0.0289 29. 8.761 31. 0.06 33. 0.0000025
 35. 2,700,000 37. 200,000 39. 10 41. 3000 43. (a) 3.231×10^8
 (b) $\$1 \times 10^{12}$ (or $\$10^{12}$) (c) \$3095 45. \$42,500
 47. (a) 5.95×10^2 (b) 998 mi² 49. 300 sec 51. approximately 5.87×10^{12} mi
 53. 7.5×10^9 55. 4×10^{17} 57. The 2002 quake was 100 times as intense as the 2018 quake. 58. The 1960 quake was 316 times as intense as the 2010 quake. 59. The 2010 quake was 25 times as intense as the 2005 quake. 60. $10^3 = 1000$ times as intense; $10^{-1} = \frac{1}{10}$ (one-tenth) as intense.

Section 4.3


1. (a) D (b) A (c) A (d) E (e) B 3. Answers will vary. One example is $7x^5 + 2x^3 - 6x^2 + 9x$. 5. To identify the leading coefficient, it is necessary to write the polynomial in descending powers of the variable as $-8x^4 - 5x^3 + 2x^2$. The leading term is $-8x^4$, so the leading coefficient is -8 . 7. 7; 1 9. -15 ; 2 11. 1; 4 13. $\frac{1}{6}$; 1 15. 8; 0 17. -1 ; 3 19. $2x^3 - 3x^2 + x + 4$; $2x^3$; 2 21. $p^7 - 8p^5 + 4p^3$; p^7 ; 1 23. $-3m^4 - m^3 + 10$; $-3m^4$; -3 25. monomial; 0 27. binomial; 1 29. binomial; 8 31. monomial; 6 33. trinomial; 3 35. none of these; 5 37. $8z^4$ 39. $7m^3$ 41. $5x$ 43. already simplified 45. $-t + 13x$ 47. $8k^2 + 2k - 7$ 49. $-2n^4 - n^3 + n^2$ 51. $-2ab^2 + 20a^2b$ 53. Only the coefficients of the like terms should be added, *not* the exponents. The correct sum is $6x^2 - 4x - 1$. 55. $8x^2 + x - 2$ 57. $-t^4 + 2t^2 - t + 5$ 59. $5y^3 - 3y^2 + 5y + 1$ 61. $r + 13$ 63. $-2a^2 - 2a - 7$ 65. $-3z^5 + z^2 + 7z$ 67. $12p - 4$ 69. $-9p^2 + 11p - 9$ 71. $5a + 18$ 73. $14m^2 - 13m + 6$ 75. $13z^2 + 10z - 3$ 77. $10y^3 - 7y^2 + 5y + 3$ 79. $-5a^4 - 6a^3 + 9a^2 - 11$ 81. $3y^2 - 4y + 2$ 83. $-4m^2 + 4n^2 - 7n$ 85. $y^4 - 4y^2 - 4$ 87. $10z^2 - 16z$ 89. $12x^2 + 8x + 5$

Section 4.4


1. polynomial; one; terms; powers 3. C 5. 0; 1; 2; (0, 0), (1, 1), (2, 2) 7. The student either did not substitute correctly or did not apply the exponent correctly. Here $f(-2) = -(-2)^2 + 4 = -(4) + 4 = 0$, so $f(-2) = 0$. 9. (a) -10 (b) 8 (c) -4 11. (a) 8 (b) -10 (c) 0 13. (a) 8 (b) 2 (c) 4 15. (a) 7 (b) 1 (c) 1 17. (a) 8 (b) 74 (c) 6 19. (a) -11 (b) 4 (c) -8 21. (a) 59.08 million lb (b) 162.20 million lb (c) 152.62 million lb 23. (a) \$45.2 billion (b) \$84.3 billion (c) \$163.4 billion 25. (a) $8x - 3$ (b) $2x - 17$ 27. (a) $-x^2 + 12x - 12$ (b) $9x^2 + 4x + 6$ 29. $f(x)$ and $g(x)$ can be any two polynomials that have a sum of $3x^3 - x + 3$, such as $f(x) = 3x^3 + 1$ and $g(x) = -x + 2$. 31. $x^2 + 2x - 9$ 33. 6 35. $x^2 - x - 6$ 37. 6 39. -33 41. 0 43. $-\frac{9}{4}$ 45. $-\frac{9}{2}$ 47. (a) $P(x) = 8.49x - 50$ (b) \$799




domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$




domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$




domain: $(-\infty, \infty)$;
range: $(-\infty, 0]$



domain: $(-\infty, \infty)$;
range: $[-2, \infty)$




domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$



domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$

61. B 63. A 65. The student multiplied the functions instead of composing them. The correct answer is $(f \circ g)(x) = -6$. 67. 6 69. 83 71. 53 73. 13 75. $2x^2 + 11$ 77. $2x - 2$ 79. $\frac{97}{4}$ 81. 8 83. 1 85. 9 87. 1 89. $(f \circ g)(x) = 63,360x$; It computes the number of inches in x miles. 91. (a) $s = \frac{x}{4}$ (b) $y = \frac{x^2}{16}$ (c) 2.25 93. (a) $g(x) = \frac{1}{2}x$ (b) $f(x) = x + 1$ (c) $(f \circ g)(x) = \frac{1}{2}x + 1$ (d) $(f \circ g)(60) = 31$; The sale price is \$31. 95. $(\mathcal{A} \circ r)(t) = 4\pi t^2$; This is the area of the circular layer as a function of time. 97. $f(1) = 1$; $f(2) = 2$ 98. $f(3) = 4$; $f(4) = 8$; $f(5) = 16$ 99. $f(3) = 4$; $f(4) = 8$; $f(5) = 16$ 100. The pattern 1, 2, 4, 8, 16 emerges, so most students predict 32 because the terms are doubling each time. However, $f(6) = 31$ (not 32). See the figure.



Section 4.5

1. C 3. D 5. $-24m^5$ 7. $-28x^7y^4$ 9. $-6x^2 + 15x$ 11. $-2q^3 - 3q^4$ 13. $18k^4 + 12k^3 + 6k^2$ 15. $15x^3 + 8x^2 + x$ 17. $6y^2 + y - 12$ 19. $m^3 - 3m^2 - 40m$ 21. $24z^3 - 20z^2 - 16z$ 23. $4x^5 - 4x^4 - 24x^3$ 25. $6t^3 + t^2 - 14t - 3$ 27. $25m^2 - 9n^2$ 29. $-2b^3 + 2b^2 + 18b + 12$ 31. $6p^4 + p^3 + 4p^2 - 27p - 6$ 33. $8z^4 - 14z^3 + 17z^2 + 20z - 3$ 35. $m^2 - 3m - 40$ 37. $12k^2 + k - 6$ 39. $15x^2 - 13x + 2$ 41. $20x^2 + 23xy + 6y^2$ 43. $3z^2 + zw - 4w^2$ 45. $12c^2 + 16cd - 3d^2$ 47. The student squared each term instead of multiplying the binomial times itself. The correct product is $x^2 + 8x + 16$. 49. $x^2 - 81$ 51. $4p^2 - 9$ 53. $25m^2 - 1$ 55. $9a^2 - 4c^2$ 57. $16m^2 - 49n^4$ 59. $75y^7 - 12y$ 61. $y^2 - 10y + 25$ 63. $x^2 + 2x + 1$ 65. $4p^2 + 28p + 49$ 67. $16n^2 - 24nm + 9m^2$ 69. $k^2 - \frac{10}{7}kp + \frac{25}{49}p^2$ 71. $0.04x^2 - 0.56xy + 1.96y^2$ 73. $16x^2 - \frac{4}{9}$ 75. $0.1x^2 + 0.63x - 0.13$ 77. $3w^2 - \frac{23}{4}wz - \frac{1}{2}z^2$ 79. $25x^2 + 10x + 1 + 60xy + 12y + 36y^2$ 81. $4a^2 + 4ab + b^2 - 12a - 6b + 9$ 83. $4a^2 + 4ab + b^2 - 9$ 85. $4h^2 - 4hk + k^2 - j^2$ 87. $y^3 + 6y^2 + 12y + 8$ 89. $125r^3 - 75r^2s + 15rs^2 - s^3$ 91. $q^4 - 8q^3 + 24q^2 - 32q + 16$ 93. $6a^3 + 7a^2b + 4ab^2 + b^3$ 95. $4z^4 - 17z^3x + 12z^2x^2 - 6zx^3 + x^4$ 97. $m^4 - 4m^2p^2 + 4mp^3 - p^4$ 99. $a^4b - 7a^2b^3 - 6ab^4$ 101. $\frac{9}{2}x^2 - 2y^2$ 103. $15x^2 - 2x - 24$ 105. $10x^2 - 2x$ 107. $2x^2 - x - 3$ 109. $8x^3 - 27$ 111. $2x^3 - 18x$ 113. -20 115. 32 117. 36 119. 0 121. $\frac{35}{4}$ 123. $a - b$ 124. $\mathcal{A} = s^2$; $(a - b)^2$ 125. $(a - b)b$, or $ab - b^2$; $2ab - 2b^2$ 126. b^2 127. a^2 ; a 128. $a^2 - (2ab - 2b^2) - b^2 = a^2 - 2ab + b^2$ 129. (a) They must be equal. (b) $(a - b)^2 = a^2 - 2ab + b^2$ 130.

| | | |
|---|----------------|----------------|
| a | Area:
a^2 | Area:
ab |
| b | Area:
ab | Area:
b^2 |




The large square is made up of two smaller squares and two congruent rectangles. The sum of the areas is $a^2 + 2ab + b^2$. Because $(a + b)^2$ must represent the same quantity, they must be equal. Thus, $(a + b)^2 = a^2 + 2ab + b^2$.

Section 4.6

1. quotient; exponents 3. 0 5. $3x^3 - 2x^2 + 1$ 7. $3y + 4 - \frac{5}{y}$
 9. $3 + \frac{5}{m} + \frac{6}{m^2}$ 11. $\frac{2}{7n} - \frac{3}{2m} + \frac{9}{7mn}$ 13. $\frac{2y}{x} + \frac{3}{4} + \frac{3w}{x}$
 15. $r^2 - 7r + 6$ 17. $y - 4$ 19. $q + 8$ 21. $t + 5$
 23. $p - 4 + \frac{44}{p + 6}$ 25. $m^2 + 2m - 1$ 27. $x^2 + 2x - 3 + \frac{-3}{4x + 1}$
 29. $2x - 5 + \frac{-4x + 5}{3x^2 - 2x + 4}$ 31. $m^2 + m + 3$ 33. $x^2 + x + 3$
 35. $2x^2 - x - 5$ 37. $3x^2 + 6x + 11 + \frac{26}{x - 2}$ 39. $2k^2 + 3k - 1$
 41. $z^2 + 3$ 43. $2y^2 + 2$ 45. $p^2 + p + 1$ 47. $x^2 - 4x + 2 + \frac{9x - 4}{x^2 + 3}$
 49. $p^2 + \frac{5}{2}p + 2 + \frac{-1}{2p + 2}$ 51. $\frac{3}{2}a - 10 + \frac{77}{2a + 6}$ 53. $\frac{2}{3}x - 1$
 55. $\frac{3}{4}a - 2 + \frac{1}{4a + 3}$ 57. $(2p + 7)$ feet 59. $Q(x) = 2x^2 - 3x + 5$;
 $R(x) = 3$ 61. $5x - 1$; 0 63. $2x - 3$; -1 65. $4x^2 + 6x + 9$; $\frac{3}{2}$
 67. $\frac{x^2 - 9}{2x}$, $x \neq 0$ 69. $-\frac{5}{4}$ 71. $\frac{x - 3}{2x}$, $x \neq 0$ 73. 0
 75. $-\frac{35}{4}$ 77. $\frac{7}{2}$

Chapter 4 Review Exercises

1. 64 2. $\frac{1}{81}$ 3. -125 4. 18 5. $\frac{81}{16}$ 6. $\frac{16}{25}$ 7. $\frac{1}{30}$ 8. $\frac{3}{4}$ 9. 0
 10. $\frac{1}{3^8}$ 11. x^8 12. $\frac{y^6}{x^2}$ 13. $\frac{1}{z^{15}}$ 14. $\frac{25}{m^{18}}$ 15. $4k^{11}$ 16. $\frac{25}{z^4}$
 17. $\frac{1}{96m^7}$ 18. $\frac{2025}{8r^4}$ 19. $-12x^2y^8$ 20. $-\frac{2n}{m^5}$ 21. $\frac{10p^8}{q^7}$
 22. In $(-6)^0$, the base is -6 and the expression simplifies to 1.
 In -6^0 , the base is 6 and the expression simplifies to -1 .
 23. 1.345×10^4 24. 7.65×10^{-8} 25. -1.38×10^{-1} 26. 3.214×10^8 ;
 7.2×10^4 ; 1×10^2 (or 10^2) 27. 1,210,000 28. $-267,000,000$
 29. 0.0058 30. 2×10^{-4} ; 0.0002 31. 1.5×10^3 ; 1500
 32. 4.1×10^{-5} ; 0.000041 33. 2.7×10^{-2} ; 0.027
 34. (a) 20,000 hr (b) 833 days 35. 14; 5 36. -1 ; 2
 37. $\frac{1}{10}$; 1 38. 504; 8 39. (a) $11k^3 - 3k^2 + 9k$ (b) trinomial
 (c) 3 40. (a) $9m^7 + 14m^6$ (b) binomial (c) 7
 41. (a) $-5y^4 + 3y^3 + 7y^2 - 2y$ (b) none of these (c) 4
 42. (a) $-7q^5r^3$ (b) monomial (c) 8 43. $-x^2 - 3x + 1$
 44. $-5y^3 - 4y^2 + 6y - 12$ 45. $6a^3 - 4a^2 - 16a + 15$
 46. $8y^2 - 9y + 5$ 47. $20x^2 - 6x + 6$ 48. One example is
 $x^5 + 2x^4 - x^2 + x + 2$. 49. (a) -11 (b) 4 (c) 7
 50. (a) 132,976 twin births (b) 133,786 twin births (c) 135,689 twin
 births 51. (a) $5x^2 - x + 5$ (b) $-5x^2 + 5x + 1$ (c) 11 (d) -9



55. (a) 167 (b) 1495 (c) 20 (d) 42 (e) $75x^2 + 220x + 160$
 (f) $15x^2 + 10x + 2$ 56. $-12k^3 - 42k$ 57. $2x^3 + 22x^2 + 56x$
 58. $15m^2 - 7m - 2$ 59. $6w^2 - 13wt + 6t^2$
 60. $10p^4 + 30p^3 - 8p^2 - 24p$ 61. $3q^3 - 13q^2 - 14q + 20$
 62. $36r^4 - 1$ 63. $16m^2 + 24m + 9$ 64. $9t^2 - 12ts + 4s^2$
 65. $y^2 - 3y + \frac{5}{4}$ 66. $x^2 - 4x + 6$ 67. $p^2 + 6p + 9 + \frac{54}{2p - 3}$
 68. $p^2 + 3p - 6$ 69. (a) $36x^3 - 9x^2$ (b) -45
 70. (a) $4x - 1$, $x \neq 0$ (b) 7

Chapter 4 Mixed Review Exercises

1. (a) A (b) G (c) C (d) C (e) A (f) E (g) B
 (h) H (i) F (j) I 2. 102,129 mi² 3. $\frac{y^4}{36}$ 4. $\frac{1}{125}$ 5. -9
 6. $-\frac{1}{52^9}$ 7. $21p^9 + 7p^8 + 14p^7$ 8. $4x^2 - 36x + 81$ 9. $\frac{1}{16y^{18}}$
 10. $\frac{4w^6}{z^{18}}$ 11. $8x^2 - 10x - 3$ 12. $8x + 1 + \frac{5}{x - 3}$
 13. $9m^2 - 30mn + 25n^2 - p^2$ 14. $2y^2x + \frac{3y^3}{2x} + \frac{5x^2}{2}$
 15. $-3k^2 + 4k - 7$

Chapter 4 Test

- [4.1] 1. (a) C (b) A (c) D (d) A (e) E (f) F (g) B
 (h) G (i) I (j) C 2. $\frac{4x^7}{9y^{10}}$ 3. $\frac{6}{r^{14}}$ 4. $\frac{16}{9p^{10}q^{28}}$ 5. $\frac{16}{x^6y^{16}}$
 [4.2] 6. 0.00000091 7. 3×10^{-4} ; 0.0003 [4.4] 8. (a) -18
 (b) $-2x^2 + 12x - 9$ (c) $-2x^2 - 2x - 3$ (d) -7
 9.
 domain: $(-\infty, \infty)$;
 range: $(-\infty, 3]$
 10.
 domain: $(-\infty, \infty)$;
 range: $(-\infty, \infty)$
 11. (a) 23 (b) $3x^2 + 11$ (c) $9x^2 + 30x + 27$
 [4.5] 12. $10x^2 - x - 3$ 13. $6m^3 - 7m^2 - 30m + 25$ 14. $36x^2 - y^2$
 15. $9k^2 + 6kq + q^2$ 16. $4y^2 - 9z^2 + 6zx - x^2$
 [4.6] 17. $4p - 8 + \frac{6}{p}$ 18. $x^2 + 4x + 4$ [4.3] 19. $x^3 - 2x^2 - 10x - 13$
 [4.5, 4.6] 20. (a) $x^3 + 4x^2 + 5x + 2$ (b) 0 (c) $x + 2$, $x \neq -1$
 (d) 0


Chapters R-4 Cumulative Review Exercises

- [R.1] 1. (a) D, I (b) C, H (c) G (d) A, F (e) B (f) E
 [R.2] 2. (a) A, B, C, D, F (b) B, C, D, F (c) D, F (d) C, D, F
 (e) E, F (f) D, F [R.4] 3. 32 4. $-\frac{1}{72}$ 5. 0 [R.5] 6. -7
 [1.1] 7. $\{-65\}$ 8. {all real numbers} [1.2] 9. $t = \frac{A - p}{pr}$
 [1.5] 10. $(-\infty, 6)$ [1.7] 11. $\{-\frac{1}{3}, 1\}$ 12. $(-\infty, -\frac{8}{3}] \cup [2, \infty)$

[R.1, 1.2] 13. 3500; 1950; 36%; 31% [1.4] 14. $15^\circ, 35^\circ, 130^\circ$

[2.2] 15. $-\frac{4}{3}$ 16. 0 [2.3] 17. (a) $y = -4x + 15$

(b) $4x + y = 15$ 18. (a) $y = 4x$ (b) $4x - y = 0$



[2.2, 2.3] 22. (a) -0.44 gal per yr; Per capita consumption of whole milk decreased by an average of 0.44 gal per year from 1970 to 2015. (b) $y = -0.44x + 25.3$ (c) 12.1 gal

[2.5] 23. domain: $\{-4, -1, 2, 5\}$; range: $\{-2, 0, 2\}$; function

[2.6] 24. -9 [3.1] 25. $\{(3, 2)\}$ 26. \emptyset [3.2] 27. $\{(1, 0, -1)\}$

[3.3] 28. length: 42 ft; width: 30 ft 29. 15% solution: 6 L;

30% solution: 3 L [4.1] 30. $\frac{y^7}{x^{13}z^2}$ [4.3] 31. $2x^2 - 5x + 10$

[4.5] 32. $x^3 + 8y^3$ 33. $15x^2 + 7xy - 2y^2$

[4.6] 34. $4xy^4 - 2y + \frac{1}{x^2y}$ 35. $m^2 - 2m + 3$

5 FACTORING

Section 5.1

1. C 3. The student stopped too soon. This is a sum of two terms, not a product. $w^2 - 8$ must be factored out to obtain $(w^2 - 8)(2t + 5)$. 5. $3m$
 7. $8xy$ 9. $3(r + t)^2$ 11. $12(m - 5)$ 13. $8(s + 2t)$ 15. $4(1 + 5z)$
 17. cannot be factored 19. $8k(k^2 + 3)$ 21. $-2p^2q^4(2p + q)$
 23. $7x^3(1 + 5x - 2x^2)$ 25. $2t^3(5t^2 - 1 - 2t)$ 27. $5ac(3ac^2 - 5c + 1)$
 29. $16zn^3(zn^3 + 4n^4 - 2z^2)$ 31. $7ab(2a^2b + a - 3a^4b^2 + 6b^3)$
 33. $(m - 4)(2m + 5)$ 35. $11(2z - 1)$ 37. $(2 - x)^2(1 + 2x)$
 39. $(3 - x)(6 + 2x - x^2)$ 41. $20z(2z + 1)(3z + 4)$
 43. $5(m + p)^2(m + p - 2 - 3m^2 - 6mp - 3p^2)$ 45. $r(-r^2 + 3r + 5); -r(r^2 - 3r - 5)$ 47. $12s^4(-s + 4); -12s^4(s - 4)$
 49. $2x^2(-x^3 + 3x + 2); -2x^2(x^3 - 3x - 2)$ 51. $(m + q)(x + y)$
 53. $(5m + n)(2 + k)$ 55. $(2 - q)(2 - 3p)$ 57. $(p + q)(p - 4z)$
 59. $(2x + 3)(y + 1)$ 61. $(m + 4)(m^2 - 6)$ 63. $(a^2 + b^2)(-3a + 2b)$
 65. $(y - 2)(x - 2)$ 67. $(3y - 2)(3y^3 - 4)$ 69. $(1 - a)(1 - b)$
 71. $2(m - 3q)(x + y)$ 73. $4(a^2 + 2b)(a - b^2)$ 75. $y^2(2x + 1)(x^2 - 7)$
 77. $m^{-5}(3 + m^2)$ 79. $p^{-3}(3 + 2p)$

Section 5.2

1. D 3. C 5. The factor $(4x + 10)$ can be factored further into $2(2x + 5)$, giving the completely factored form as $2(2x + 5)(x - 2)$.
 7. $x + 3$ 9. $m - 7$ 11. $r - 5$ 13. $x - 2a$ 15. $2x - 3$ 17. $2u + v$
 19. $(y - 3)(y + 10)$ 21. $(p + 8)(p + 7)$ 23. prime
 25. $(a + 5b)(a - 7b)$ 27. $(a - 6b)(a - 3b)$ 29. prime
 31. $-(6m - 5)(m + 3)$ 33. $(5x - 6)(2x + 3)$ 35. $(4k + 3)(5k + 8)$
 37. $(3a - 2b)(5a - 4b)$ 39. $(6m - 5)^2$ 41. prime
 43. $(2xz - 1)(3xz + 4)$ 45. $3(4x + 5)(2x + 1)$

47. $-5(a + 6)(3a - 4)$ 49. $-11x(x - 6)(x - 4)$ 51. $2xy^3(x - 12y)^2$
 53. $6a(a - 3)(a + 5)$ 55. $13y(y + 4)(y - 1)$ 57. $3p(2p - 1)^2$
 59. $(6p^3 - r)(2p^3 - 5r)$ 61. $(5k + 4)(2k + 1)$
 63. $(3m + 3p + 5)(m + p - 4)$ 65. $(p^2 - 8)(p^2 - 2)$
 67. $(2x^2 + 3)(x^2 - 6)$ 69. $(4x^2 + 3)(4x^2 + 1)$ 71. $(z - x)^2(z + 2x)$
 73. $(a + b)^2(a - 3b)(a + 2b)$ 75. $(p + q)^2(p + 3q)$
 77. $\frac{36x^2 + 42x - 120}{6}$ 78. $\frac{t^2 + 7t - 120}{6}$ 79. $\frac{(t + 15)(t - 8)}{6}$
 80. $\frac{(6x + 15)(6x - 8)}{6} = \frac{3(2x + 5) \cdot 2(3x - 4)}{6} = (2x + 5)(3x - 4)$
 81. $(x - 4)(3x + 8)$ 82. $(4x + 1)(3x + 8)$

Section 5.3

1. A, D 3. B, C 5. $x^2 + 4$ is a sum of squares with GCF = 1. It cannot be factored and is prime. Also, $(x + 2)^2 = x^2 + 4x + 4$, not $x^2 + 4$. In general, $x^2 + y^2 \neq (x + y)^2$. 7. $(p + 4)(p - 4)$
 9. $(5x + 2)(5x - 2)$ 11. $2(3a + 7b)(3a - 7b)$
 13. $4(4m^2 + y^2)(2m + y)(2m - y)$ 15. $(y + z + 9)(y + z - 9)$
 17. $(4 + x + 3y)(4 - x - 3y)$ 19. $(p^2 + 16)(p + 4)(p - 4)$
 21. $(k - 3)^2$ 23. $(2z + w)^2$ 25. $(4m - 1 + n)(4m - 1 - n)$
 27. $(2r - 3 + s)(2r - 3 - s)$ 29. $(x + y - 1)(x - y + 1)$
 31. $2(7m + 3n)^2$ 33. $(p + q + 1)^2$ 35. $(a - b + 4)^2$
 37. $(x - 3)(x^2 + 3x + 9)$ 39. $(6 - t)(36 + 6t + t^2)$
 41. $(x + 4)(x^2 - 4x + 16)$ 43. $(10 + y)(100 - 10y + y^2)$
 45. $(2x + 1)(4x^2 - 2x + 1)$ 47. $(5x - 6)(25x^2 + 30x + 36)$
 49. $(x - 2y)(x^2 + 2xy + 4y^2)$ 51. $(4g - 3h)(16g^2 + 12gh + 9h^2)$
 53. $(7p + 5q)(49p^2 - 35pq + 25q^2)$ 55. $3(2n + 3p)(4n^2 - 6np + 9p^2)$
 57. $(y + z + 4)(y^2 + 2yz + z^2 - 4y - 4z + 16)$
 59. $(m^2 - 5)(m^4 + 5m^2 + 25)$ 61. $(3 - 10x^3)(9 + 30x^3 + 100x^6)$
 63. $(5y^2 + z)(25y^4 - 5y^2z + z^2)$ 65. $k(5 - 4k)(25 + 20k + 16k^2)$
 67. $(5p + 2q)(25p^2 - 10pq + 4q^2 + 5p - 2q)$
 69. $(3a - 4b)(9a^2 + 12ab + 16b^2 + 5)$
 71. $(t - 3)(2t + 1)(4t^2 - 2t + 1)$
 73. $(8m - 9n)(8m + 9n - 64m^2 - 72mn - 81n^2)$
 75. $(x^3 - y^3)(x^3 + y^3); (x - y)(x^2 + xy + y^2)(x + y)(x^2 - xy + y^2)$
 76. $(x^2 + xy + y^2)(x^2 - xy + y^2)$ 77. $(x^2 - y^2)(x^4 + x^2y^2 + y^4); (x - y)(x + y)(x^4 + x^2y^2 + y^4)$ 78. $x^4 + x^2y^2 + y^4$ 79. The product must equal $x^4 + x^2y^2 + y^4$. Multiply $(x^2 + xy + y^2)(x^2 - xy + y^2)$ to verify this. 80. Start by factoring as a difference of squares.

Section 5.4

1. (a) B (b) D (c) A (d) A, C (e) A, B 3. $3p^2(p - 6)(p + 5)$
 5. $3pq(a + 6b)(a - 5b)$ 7. prime 9. $(6b + 1)(b - 3)$
 11. $(x - 10)(x^2 + 10x + 100)$ 13. $(p + 2)(4 + m)$
 15. $9m(m - 5 + 2m^2)$ 17. $2(3m - 10)(9m^2 + 30m + 100)$
 19. $(3m - 5n)^2$ 21. $(k - 9)(q + r)$ 23. $16z^2x(zx - 2)$
 25. $(r + 4)(2r - 3)$ 27. $(25 + x^2)(5 - x)(5 + x)$
 29. $(p + 1)(p^2 - p + 1)$ 31. $(8m + 25)(8m - 25)$ 33. $6z(2z^2 - z + 3)$
 35. $16(4b + 5c)(4b - 5c)$ 37. $8(4 + 5z)(16 - 20z + 25z^2)$

39. $(5r - s)(2r + 5s)$ 41. $4pq(2p + q)(3p + 5q)$
 43. $3(4k^2 + 9)(2k + 3)(2k - 3)$ 45. $(m - n)(m^2 + mn + n^2 + m + n)$
 47. $(x - 2m - n)(x + 2m + n)$ 49. $6p^3(3p^2 - 4 + 2p^3)$
 51. $2(x + 4)(x - 5)$ 53. $8mn$ 55. $2(5p + 9)(5p - 9)$
 57. $4rx(3m^2 + mn + 10n^2)$ 59. $(7a - 4b)(3a + b)$ 61. prime
 63. $(p + 8q - 5)^2$ 65. $(7m^2 + 1)(3m^2 - 5)$
 67. $(2r - t)(r^2 - rt + 19t^2)$ 69. $(x + 3)(x^2 + 1)(x + 1)(x - 1)$
 71. $(m + n - 5)(m - n + 1)$

Section 5.5

1. B, C, E, F 3. The zero-factor property can be applied only with a product that equals 0. Multiply on the left, write the equation in standard form $x^2 - x - 12 = 0$, factor, and then apply the zero-factor property. The solution set is $\{-3, 4\}$. 5. By dividing each side by a variable expression, here $3x$, the student “lost” the solution 0. On the left, set $3x = 0$ when the zero-factor property is applied. The solution set is $\{-4, 0\}$.

7. $\{-10, 5\}$ 9. $\{-\frac{8}{3}, \frac{5}{2}\}$ 11. $\{-2, 5\}$ 13. $\{-6, -3\}$
 15. $\{-\frac{1}{2}, 4\}$ 17. $\{-\frac{1}{3}, \frac{4}{5}\}$ 19. $\{-4, 0\}$ 21. $\{0, 6\}$ 23. $\{-2, 2\}$
 25. $\{-3, 3\}$ 27. $\{3\}$ 29. $\{-\frac{4}{3}\}$ 31. $\{-4, 2\}$ 33. $\{-\frac{1}{2}, 6\}$
 35. $\{-3, 4\}$ 37. $\{-5, -\frac{1}{5}\}$ 39. $\{1, 6\}$ 41. $\{-\frac{1}{2}, 0, 5\}$
 43. $\{-1, 0, 3\}$ 45. $\{-\frac{4}{3}, 0, \frac{4}{3}\}$ 47. $\{-\frac{5}{2}, -1, 1\}$ 49. $\{-3, 3, 6\}$
 51. $\{-\frac{1}{2}, 6\}$ 53. $\{-\frac{2}{3}, \frac{4}{15}\}$ 55. $\{-\frac{3}{2}, \frac{1}{2}\}$ 57. width: 16 ft; length: 20 ft
 59. base: 12 ft; height: 5 ft 61. 50 ft by 100 ft 63. -6 and -5 or 5 and 6
 65. length: 15 in.; width: 9 in. 67. 5 sec 69. 6 sec 71. $F = \frac{k}{d - D}$
 73. $P = \frac{A}{1 + rt}$ 75. $r = \frac{-2k - 3y}{a - 1}$, or $r = \frac{2k + 3y}{1 - a}$ 77. $y = \frac{-x}{w - 3}$,
 or $y = \frac{x}{3 - w}$

Chapter 5 Review Exercises

1. $6p(2p - 1)$ 2. $7x(3x + 5)$ 3. $4qb(3q + 2b - 5q^2b)$
 4. $6rt(r^2 - 5rt + 3t^2)$ 5. $(x + 3)(x - 3)$ 6. $(z + 1)(3z - 1)$
 7. $(m + q)(4 + n)$ 8. $(x + y)(x + 5)$ 9. $(m + 3)(2 - a)$
 10. $(x + 3)(x - y)$ 11. $(3p - 4)(p + 1)$ 12. $(3k - 2)(2k + 5)$
 13. $(3r + 1)(4r - 3)$ 14. $(2m + 5)(5m + 6)$ 15. $(2k - h)(5k - 3h)$
 16. prime 17. $2x(4 + x)(3 - x)$ 18. $3b(2b - 5)(b + 1)$
 19. $(y^2 + 4)(y^2 - 2)$ 20. $(2k^2 + 1)(k^2 - 3)$
 21. $(p + 2)^2(p + 3)(p - 2)$ 22. $(3r + 16)(r + 1)$
 23. $3x^4(2x^2 - 9)(4x^2 - 3)$ 24. $5y^3(3y^3 + 1)(2y^3 - 11)$
 25. $(4x + 5)(4x - 5)$ 26. $(3t + 7)(3t - 7)$
 27. $(6m - 5n)(6m + 5n)$ 28. $(x + 7)^2$ 29. $(3k - 2)^2$
 30. $(r + 3)(r^2 - 3r + 9)$ 31. $(5x - 1)(25x^2 + 5x + 1)$
 32. $(m + 1)(m^2 - m + 1)(m - 1)(m^2 + m + 1)$
 33. $(x^4 + 1)(x^2 + 1)(x + 1)(x - 1)$ 34. $(x + 3 + 5y)(x + 3 - 5y)$
 35. $2b(3a^2 + b^2)$ 36. $(x + 1)(x - 1)(x - 2)(x^2 + 2x + 4)$
 37. $\{4\}$ 38. $\{-1, -\frac{2}{5}\}$ 39. $\{2, 3\}$ 40. $\{-4, 2\}$ 41. $\{-\frac{5}{2}, \frac{10}{3}\}$
 42. $\{-\frac{3}{2}, \frac{1}{3}\}$ 43. $\{-\frac{3}{2}, -\frac{1}{4}\}$ 44. $\{-3, 3\}$ 45. $\{-\frac{3}{2}, 0\}$

46. $\{\frac{1}{2}, 1\}$ 47. $\{1, 4\}$ 48. $\{-\frac{7}{2}, 0, 4\}$ 49. $\{-3, -2, 2\}$
 50. $\{-2, -\frac{6}{5}, 3\}$ 51. 3 ft 52. length: 60 ft; width: 40 ft 53. 16 sec
 54. 1 sec and 15 sec 55. 8 sec 56. The rock reaches a height of 240 ft once on its way up and once on its way down. 57. $k = \frac{-3s - 2t}{b - 1}$,
 or $k = \frac{3s + 2t}{1 - b}$ 58. $w = \frac{7}{z - 3}$, or $w = \frac{-7}{3 - z}$


Chapter 5 Mixed Review Exercises

1. $(x + y)(x + 5)$ 2. $-(x + 2)(x - 5)$ 3. $16(x^2 + 9)$
 4. $a(6 - m)(5 + m)$ 5. $(2 - a)(4 + 2a + a^2)$ 6. $(9k + 4)(9k - 4)$
 7. prime 8. $5y^2(3y + 4)$ 9. $(5z - 3m)^2$ 10. D; The polynomial is not factored. 11. $\{-\frac{3}{5}, 4\}$ 12. $\{-1, 0, 1\}$ 13. $\{0, 3\}$
 14. $h = \frac{2sA}{b + B}$ 15. $H = \frac{S - 2LW}{2W + 2L}$ 16. width: 25 ft; length: 110 ft
 17. 6 in. 18. 5 sec and 9 sec

Chapter 5 Test

- [5.1–5.4] 1. $11z(z - 4)$ 2. $5x^2y^3(2y^2 - 1 - 5x^3)$
 3. $(x + y)(3 + b)$ 4. $-(2x + 9)(x - 4)$ 5. $(3x - 5)(2x + 7)$
 6. $(4p - q)(p + q)$ 7. $(4a + 5b)^2$ 8. $(x + 1 + 2z)(x + 1 - 2z)$
 9. $(a + b)(a - b)(a + 2)$ 10. $(3k + 11j)(3k - 11j)$
 11. $(y - 6)(y^2 + 6y + 36)$ 12. $(2k^2 - 5)(3k^2 + 7)$
 13. $(3x^2 + 1)(9x^4 - 3x^2 + 1)$ 14. $(t^2 + 8)(t^2 + 2)$
 15. prime 16. $-(x + 5)(x - 6)$ 17. $(5 - 6y^2)(25 + 30y^2 + 36y^4)$
 18. D [5.5] 19. $\{-2, -\frac{2}{3}\}$ 20. $\{0, \frac{5}{3}\}$ 21. $\{-\frac{2}{5}, 1\}$
 22. $r = \frac{-2 - 6t}{a - 3}$, or $r = \frac{2 + 6t}{3 - a}$ 23. length: 8 in.; width: 5 in.
 24. 2 sec and 4 sec

Chapters R–5 Cumulative Review Exercises

- [R.1] 1. $\frac{13}{12}$ 2. 16.1 [R.5] 3. $-2m + 6$ 4. $2x^2 + 5x + 4$
 [R.4] 5. 10 6. undefined [1.1] 7. $\{\frac{7}{6}\}$ 8. $\{-1\}$
 [1.5] 9. $(-\frac{1}{2}, \infty)$ [1.6] 10. $(2, 3)$ 11. $(-\infty, 2) \cup (3, \infty)$
 [1.7] 12. $(-\frac{16}{5}, 2)$ 13. $(-11, 7)$ 14. $(-\infty, -2] \cup [7, \infty)$
 [1.4] 15. 2 hr [2.2] 16. 0 17. -1
 [2.1] 18.  19. -1 20. $(-\frac{7}{2}, 0)$ 21. $(0, 7)$
 [3.1] 22. $\{(1, 5)\}$ 23. \emptyset
 [3.2] 24. $\{(1, 1, 0)\}$
 [4.5] 25. $49x^2 + 42xy + 9y^2$ [4.6] 26. $2x^2 + x + 3$
 [4.3] 27. $x^3 + 12x^2 - 3x - 7$ [4.2] 28. (a) 4.638×10^{-4} (b) 566,000
 [5.1–5.4] 29. $(2w + 7z)(8w - 3z)$ 30. $(2x - 1 + y)(2x - 1 - y)$
 31. $(10x^2 + 9)(10x^2 - 9)$ 32. $(2p + 3)(4p^2 - 6p + 9)$
 [5.5] 33. $\{-4, -\frac{3}{2}, 1\}$ 34. $\{\frac{1}{3}\}$ 35. 4 ft 36. longer sides: 18 in.;
 distance between: 16 in.

6 RATIONAL EXPRESSIONS AND FUNCTIONS

Section 6.1

1. integers; 0; rational; polynomials; 0

3. (a) $\{x|x \text{ is a real number, } x \neq 7\}$ (b) $(-\infty, 7) \cup (7, \infty)$ 5. (a) $\{x|x \text{ is a real number, } x \neq -\frac{1}{7}\}$ (b) $(-\infty, -\frac{1}{7}) \cup (-\frac{1}{7}, \infty)$ 7. (a) $\{x|x \text{ is a real number, } x \neq 0\}$ (b) $(-\infty, 0) \cup (0, \infty)$ 9. (a) $\{x|x \text{ is a real number, } x \neq -2, \frac{3}{2}\}$ (b) $(-\infty, -2) \cup (-2, \frac{3}{2}) \cup (\frac{3}{2}, \infty)$ 11. (a) $\{x|x \text{ is a real number}\}$ (b) $(-\infty, \infty)$ 13. (a) $\{x|x \text{ is a real number}\}$ (b) $(-\infty, \infty)$ 15. $\frac{2}{15}$ 17. $\frac{9}{10}$ 19. $\frac{3}{4}$ 21. B, E, F 23. B 25. The expression $\frac{2x}{4}$ has 2 as a factor of both the numerator and the denominator. It can be written in lowest terms as $\frac{x}{2}$. 27. The student tried to simplify beforefactoring the numerator and denominator. The correct answer is $\frac{1}{2}$.29. x 31. $\frac{x-3}{x+5}$ 33. $\frac{x+3}{2x(x-3)}$ 35. It is already in lowest terms.37. $\frac{6}{7}$ 39. $\frac{z}{6}$ 41. $\frac{t-3}{3}$ 43. $\frac{2}{t-3}$ 45. $\frac{x-3}{x+1}$ 47. $\frac{4x+1}{4x+3}$ 49. $a^2 - ab + b^2$ 51. $\frac{c+6d}{c-d}$ 53. $\frac{a+b}{a-b}$ 55. -1 57. $-(x+y)$, or $-x-y$ 59. $-(x+2)$, or $-x-2$ 61. $-\frac{x+y}{x-y}$ (There are other answers.) 63. $-\frac{1}{2}$ 65. It is already in lowestterms. 67. $\frac{x+4}{x-2}$ 69. $\frac{2x+3}{x+2}$ 71. $\frac{7x}{6}$ 73. $-\frac{p+5}{2p}$ (There areother answers.) 75. $\frac{35}{4}$ 77. $-(z+1)$, or $-z-1$ 79. $\frac{14x^2}{5}$ 81. $\frac{-m(m+7)}{m+1}$ (There are other answers.) 83. -2 85. $\frac{x+4}{x-4}$ 87. $\frac{a^2+ab+b^2}{a-b}$ 89. $\frac{2x-3}{2(x-3)}$ 91. $\frac{a^2+2ab+4b^2}{a+2b}$ 93. $\frac{2x+3}{2x-3}$ 95. $\frac{27}{2mn^7}$ 97. $\frac{k+5p}{2k+5p}$ 99. $(k-1)(k-2)$

Section 6.2

1. product; factors; greatest; denominator 3. C 5. $\frac{4}{5}$ 7. $-\frac{1}{18}$ 9. $\frac{31}{36}$ 11. $\frac{9}{t}$ 13. $\frac{6x+y}{7}$ 15. $\frac{2}{x}$ 17. $-\frac{2}{x^3}$ 19. 1 21. $x-5$ 23. $\frac{1}{p+3}$ 25. $\frac{5}{x-2}$ 27. $a-b$ 29. $72x^4y^5$ 31. $z(z-2)$ 33. $2(y+4)$ 35. $(x+9)^2(x-9)$ 37. $(m+n)(m-n)$ 39. $x(x-4)(x+1)$ 41. $(t+5)(t-2)(2t-3)$ 43. $2y(y+3)(y-3)$ 45. $2(x+2)^2(x-3)$ 47. The expression $\frac{x-4x-1}{x+2}$ is incorrect. The third term in the numerator shouldbe $+1$ because the $-$ sign should be distributed over both $4x$ and -1 .The correct answer is $\frac{-3x+1}{x+2}$. 49. $\frac{31}{3t}$ 51. $\frac{5-22x}{12x^2y}$ 53. $\frac{16b+9a^2}{60a^4b^6}$ 55. $\frac{4pr+3sq^3}{14p^4q^4}$ 57. $\frac{a^2b^5-2ab^6+3}{a^5b^7}$ 59. $\frac{1}{x(x-1)}$ 61. $\frac{5a^2-7a}{(a+1)(a-3)}$ 63. 4 65. 367. $\frac{3}{x-4}$, or $\frac{-3}{4-x}$ 69. $\frac{w+z}{w-z}$, or $\frac{-w-z}{z-w}$ 71. $\frac{-2}{(x+1)(x-1)}$ 73. $\frac{2(2x-1)}{x-1}$ 75. $\frac{7}{y}$ 77. $\frac{6}{x-2}$ 79. $\frac{3x-2}{x-1}$ 81. $\frac{4x-7}{x^2-x+1}$ 83. $\frac{2x+1}{x}$ 85. $\frac{4p^2-21p+29}{(p-2)^2}$ 87. $\frac{x}{(x-2)^2(x-3)}$ 89. $\frac{10x+23}{(x+2)^2(x+3)}$ 91. $\frac{2x^2+24xy}{(x+2y)(x-y)(x+6y)}$ 93. $\frac{2x^2+21xy-10y^2}{(x+2y)(x-y)(x+6y)}$ 95. $\frac{3r-2s}{(2r-s)(3r-s)}$ 97. (a) $C(x) = \frac{8000+10x}{49(101-x)}$

(b) 30.44 thousand dollars

Section 6.3

1. complex; numerator; both 3. $\frac{1}{6}$ 5. $\frac{9}{5}$ 7. $\frac{4}{15}$ 9. $\frac{7}{17}$ 11. $\frac{2x}{x-1}$ 13. $\frac{2(k+1)}{3k-1}$ 15. $\frac{5x^2}{9z^3}$ 17. $\frac{1+x}{-1+x}$ 19. $\frac{6x+1}{7x-3}$ 21. $\frac{y+x}{y-x}$ 23. $4x$ 25. $\frac{y+4}{2}$ 27. $x+4y$ 29. $\frac{a+b}{ab}$ 31. xy 33. $\frac{3y}{2}$ 35. $\frac{y^2+yx+x^2}{xy(y+x)}$ 37. $\frac{x^2+5x+4}{x^2+5x+10}$

39. The negative exponents are on terms, not factors. Terms with negative exponents cannot be simply moved across a fraction bar.

41. $\frac{x^2y^2}{y^2+x^2}$ 43. $\frac{y^2+x^2}{xy^2+x^2y}$, or $\frac{y^2+x^2}{xy(y+x)}$ 45. $\frac{p^2+k}{p^2-3k}$ 47. $\frac{1}{2xy}$ 49. $\frac{m^2+6m-4}{m(m-1)}$ 50. $\frac{m^2-m-2}{m(m-1)}$ 51. $\frac{m^2+6m-4}{m^2-m-2}$ 52. $m(m-1)$ 53. $\frac{m^2+6m-4}{m^2-m-2}$ 54. Answers will vary.

Section 6.4


1. (a) equation (b) expression (c) expression (d) equation

3. The proposed solution 3 must be rejected because it causes a denominator to equal 0. The solution set is \emptyset .


In Exercises 5–15, we give the domains using set-builder notation.

5. $\{x|x \text{ is a real number, } x \neq 0\}$ 7. $\{x|x \text{ is a real number, } x \neq -1, 2\}$ 9. $\{x|x \text{ is a real number, } x \neq 4, \frac{7}{2}\}$ 11. $\{x|x \text{ is a real number, } x \neq -\frac{7}{4}, 0, \frac{13}{6}\}$ 13. $\{x|x \text{ is a real number, } x \neq \pm 4\}$ 15. $\{x|x \text{ is a real number, } x \neq 0, 1, -3, 2\}$ 17. $\{1\}$ 19. $\{-6, 4\}$ 21. $\{-7, 3\}$ 23. $\{-\frac{7}{12}\}$ 25. \emptyset 27. $\{-3\}$ 29. $\{5\}$ 31. $\{0\}$ 33. $\{\frac{27}{56}\}$ 35. \emptyset 37. $\{5\}$ 39. \emptyset 41. \emptyset 43. $\{-6, 3\}$ 45. $\{0\}$ 47. $\{-3, -1\}$ 49. $\{-10\}$ 51. $\{-1\}$ 53. $\{13\}$ 55. $\{x|x \text{ is a real number, } x \neq \pm 3\}$ 57. (a) D (b) C


(c) A (d) B




$x = 0; y = 0$



$x = 0; y = 0$



$x = 2; y = 0$



$x = 1; y = 0$

67. (a) 0 (b) 1.6 (c) 4.1 (d) The waiting time also increases.
 69. (a) 500 ft (b) It decreases.

SUMMARY EXERCISES Simplifying Rational Expressions vs. Solving Rational Equations

1. equation; {20} 2. expression; $\frac{2(x+5)}{5}$ 3. expression; $-\frac{22}{7x}$
 4. expression; $\frac{y+x}{y-x}$ 5. equation; $\left\{\frac{1}{2}\right\}$ 6. equation; {7}
 7. expression; $\frac{43}{24x}$ 8. equation; {1} 9. expression; $\frac{5x-1}{-2x+2}$, or $\frac{5x-1}{-2(x-1)}$ 10. expression; $\frac{25}{4(r+2)}$ 11. expression; $\frac{x^2+xy+2y^2}{(x+y)(x-y)}$
 12. expression; $\frac{24p}{p+2}$ 13. expression; $-\frac{5}{36}$ 14. equation; {0}
 15. expression; $\frac{2x+10}{x(x-2)(x+2)}$ 16. equation; $\left\{\frac{1}{7}, 2\right\}$
 17. expression; $\frac{-x}{3x+5y}$ 18. expression; $\frac{t-2}{8}$ 19. equation; \emptyset
 20. expression; $\frac{13x+28}{2x(x+4)(x-4)}$ 21. expression; $\frac{b+3}{3}$
 22. expression; $\frac{5}{3z}$ 23. expression; $\frac{3y+2}{y+3}$ 24. equation; {-13}
 25. expression; $\frac{-1}{x-3}$, or $\frac{1}{3-x}$ 26. equation; $\left\{\frac{5}{4}\right\}$
 27. equation; {-10} 28. expression; $\frac{2z-3}{2z+3}$ 29. equation; \emptyset
 30. expression; $\frac{k(2k^2-2k+5)}{(k-1)(3k^2-2)}$

Section 6.5

1. A 3. D 5. 24 7. $\frac{25}{4}$ 9. The variable a cannot appear on both sides of the final answer. The correct answer is
 $a = \frac{mb}{m-b}$ 11. $G = \frac{Fd^2}{Mm}$ 13. $a = \frac{bc}{c+b}$ 15. $v = \frac{PVt}{pT}$
 17. $r = \frac{nE-IR}{In}$, or $r = \frac{IR-nE}{-In}$ 19. $b = \frac{2sA}{h} - B$, or $b = \frac{2sA-hB}{h}$
 21. $r = \frac{eR}{E-e}$ 23. $R = \frac{D}{1-DT}$, or $R = \frac{-D}{DT-1}$ 25. 21 girls, 7 boys
 27. 1.75 in. 29. 5.4 in. 31. 7.6 in. 33. 56 teachers 35. 210 deer

37. 25,000 fish 39. 6.6 more gallons 41. $x = \frac{7}{2}; AC = 8; DF = 12$
 43. 2.4 mL 45. $\frac{1}{3}$ job per hr 47. (a) $4-x$ (b) $4+x$
 49. 3 mph 51. 10 mph 53. 1020 mi 55. 1750 mi 57. 190 mi
 59. 4 hr 61. $6\frac{2}{3}$ min, or 6 min, 40 sec 63. 30 hr 65. $2\frac{1}{3}$ hr, or 2 hr, 20 min 67. 20 hr 69. $2\frac{4}{5}$ hr, or 2 hr, 48 min

Section 6.6

1. increases; decreases 3. direct 5. direct 7. inverse 9. inverse
 11. inverse 13. direct 15. joint 17. combined 19. The perimeter of a square varies directly as the length of its side. 21. The surface area of a sphere varies directly as the square of its radius. 23. The area of a triangle varies jointly as the lengths of its base and height.
 25. 4; 2; $4\pi; \frac{4}{3}\pi; \frac{1}{2}; \frac{1}{3}\pi$ 27. $A = kb$ 29. $h = \frac{k}{t}$ 31. $M = kd^2$
 33. $I = kgh$ 35. 36 37. $\frac{16}{9}$ 39. 0.625 41. $\frac{16}{5}$ 43. $222\frac{2}{9}$
 45. \$3.92 47. 8 lb 49. 450 cm³ 51. 256 ft 53. $106\frac{2}{3}$ mph
 55. 100 cycles per sec 57. $21\frac{1}{3}$ foot-candles 59. \$420 61. 11.8 lb
 63. 448.1 lb 65. 68,600 calls 67. Answers will vary. 69. (0, 0), (1, 3.75) 70. 3.75 71. $y = 3.75x + 0$, or $y = 3.75x$
 72. $a = 3.75, b = 0$ 73. It is the price per gallon and the slope of the line. 74. It can be written in the form $y = kx$ (where $k = a$). The value of a is the constant of variation.


Chapter 6 Review Exercises

1. (a) $\{x|x \text{ is a real number, } x \neq -6\}$ (b) $(-\infty, -6) \cup (-6, \infty)$
 2. (a) $\{x|x \text{ is a real number, } x \neq 2, 5\}$ (b) $(-\infty, 2) \cup (2, 5) \cup (5, \infty)$
 3. (a) $\{x|x \text{ is a real number, } x \neq 9\}$ (b) $(-\infty, 9) \cup (9, \infty)$ 4. $\frac{x}{2}$
 5. $\frac{5m+n}{5m-n}$ 6. $\frac{-1}{2+r}$ 7. $\frac{3y^2(2y+3)}{2y-3}$ 8. $\frac{-3(w+4)}{w}$ 9. $x-4$
 10. $\frac{z(z+2)}{z+5}$ 11. $\frac{-2}{x+8}$ 12. 1 13. $96b^5$ 14. $9r^2(3r+1)$
 15. $(3x-1)(2x+5)(3x+4)$ 16. $3(x-4)^2(x+2)$
 17. $\frac{16z-3}{2z^2}$ 18. 12 19. $\frac{71}{30(a+2)}$ 20. $\frac{13r^2+5rs}{(5r+s)(2r-s)(r+s)}$
 21. $\frac{3+2t}{4-7t}$ 22. -2 23. $\frac{1}{3q+2p}$ 24. $\frac{y+x}{xy}$ 25. {-3}
 26. {-2} 27. \emptyset 28. {0} 29. {-5} 30. \emptyset
 31. (a) equation; {-24} (b) expression; $\frac{24+x}{6x}$
 32.
 $x = -1; y = 0$
 33. $\frac{15}{2}$ 34. 2 35. $m = \frac{Fd^2}{GM}$ 36. $M = \frac{m\mu}{v-\mu}$
 37. 6000 passenger-km per day 38. 12.2 more gallons 39. 16 km per hr
 40. $4\frac{4}{5}$ min, or 4 min, 48 sec 41. $3\frac{3}{5}$ hr, or 3 hr, 36 min 42. C
 43. 430 mm 44. 5.59 vibrations per sec 45. 22.5 ft³


Chapter 6 Mixed Review Exercises


1. $\frac{1}{x-2y}$ 2. $\frac{x+5}{x+2}$ 3. $\frac{6m+5}{3m^2}$ 4. $\frac{11}{3-x}$, or $\frac{-11}{x-3}$ 5. $\frac{x^2-6}{2(2x+1)}$
 6. $\frac{3-5x}{6x+1}$ 7. $\frac{1}{3}$ 8. $\frac{s^2+t^2}{st(s-t)}$ 9. $\frac{k-3}{36k^2+6k+1}$ 10. $\frac{x(9x+1)}{3x+1}$
 11. $\frac{5a^2+4ab+12b^2}{(a+3b)(a-2b)(a+b)}$ 12. $\frac{acd+b^2d+bc^2}{bcd}$ 13. $\left\{\frac{1}{3}\right\}$
 14. $\left\{-\frac{14}{3}\right\}$ 15. $\{1, 4\}$ 16. $r = \frac{AR}{R-A}$, or $r = \frac{-AR}{A-R}$
 17. (a) 8.32 mm (b) 44.87 diopters 18. \$53.17 19. $4\frac{1}{2}$ mi
 20. 12 ft²

Chapter 6 Test

- [6.1] 1. (a) $\{x \mid x \text{ is a real number, } x \neq -2, \frac{4}{3}\}$
 (b) $(-\infty, -2) \cup (-2, \frac{4}{3}) \cup (\frac{4}{3}, \infty)$ 2. $\frac{2x-5}{x(3x-1)}$ 3. $\frac{3(x+3)}{4}$
 4. $\frac{y+4}{y-5}$ 5. -2 6. $\frac{x+5}{x}$ [6.2] 7. $t^2(t+3)(t-2)$
 8. $\frac{7-2t}{6t^2}$ 9. $\frac{13x+35}{(x-7)(x+7)}$ 10. $\frac{11x+21}{(x-3)^2(x+3)}$ 11. $\frac{4}{x+2}$
 [6.3] 12. $\frac{72}{11}$ 13. $-\frac{1}{a+b}$ 14. $\frac{2y^2+x^2}{xy(y-x)}$ [6.4] 15. (a) expression;
 $\frac{11(x-6)}{12}$ (b) equation; $\{6\}$ 16. $\left\{\frac{1}{2}\right\}$ 17. $\{5\}$
 18. $\ell = \frac{2S}{n} - a$, or $\ell = \frac{2S-na}{n}$
 19. 
 $x = -1; y = 0$ [6.5] 20. $3\frac{3}{14}$ hr 21. 15 mph
 22. 48,000 fish 23. (a) 3 units (b) 0
 [6.6] 24. 200 amps 25. 0.8 lb

Chapters R-6 Cumulative Review Exercises

- [R.1, R.3] 1. $-\frac{11}{12}$ 2. $\frac{5}{8}$ 3. $-\frac{6}{5}$ 4. -5.67 5. 0.0525 6. 4360
 [R.4] 7. (a) 25 (b) -25 (c) 25 (d) 5 (e) -5 (f) not a real number
 8. -199 [1.1] 9. $\left\{-\frac{15}{4}\right\}$ [1.7] 10. $\left\{\frac{2}{3}, 2\right\}$
 [1.5] 11. $(-\infty, \frac{240}{13})$ [1.7] 12. $(-\infty, -2] \cup [\frac{2}{3}, \infty)$
 [1.3] 13. \$4000 at 4%; \$8000 at 3% 14. 6 m
 [2.1] 15. 
 x-intercept: (-2, 0); y-intercept: (0, 4) [2.2] 16. $-\frac{3}{2}$ 17. $-\frac{3}{4}$
 [2.3] 18. $y = -\frac{3}{2}x + \frac{1}{2}$







- [2.6] 21. (a) $f(x) = \frac{5}{3}x - \frac{8}{3}$ (b) -1 [2.5] 22. (a) yes
 (b) domain: $[-2, \infty)$; range: $(-\infty, 0]$ [3.1] 23. $\{(-1, 3)\}$
 [3.2] 24. $\{(-2, 3, 1)\}$ 25. \emptyset [4.1] 26. $\frac{m}{n}$
 [4.3] 27. $4y^2 - 7y - 6$ [4.5] 28. $12f^2 + 5f - 3$
 29. $\frac{1}{16}x^2 + \frac{5}{2}x + 25$ [4.6] 30. $x^2 + 4x - 7$
 [4.4] 31. (a) $2x^3 - 2x^2 + 6x - 4$ (b) $2x^3 - 4x^2 + 2x + 2$
 (c) -14 (d) $x^4 + 2x^2 - 3$ [5.2] 32. $(2x+5)(x-9)$
 [5.3] 33. $25(2t^2+1)(2t^2-1)$ 34. $(2p+5)(4p^2-10p+25)$
 [6.1] 35. $\frac{a(a-b)}{2(a+b)}$ 36. $\frac{2(x+3)}{(x+2)(x^2+3x+9)}$ [6.2] 37. 3
 [5.5] 38. $\left\{-\frac{7}{3}, 1\right\}$ [6.4] 39. $\{-4\}$ [6.5] 40. $1\frac{1}{5}$ hr, or 1 hr, 12 min

7 ROOTS, RADICALS, AND ROOT FUNCTIONS

Section 7.1

1. (a) 5 (b) 9 (c) 12 3. $\sqrt{16}$ is a number whose square equals 16, not a number that gives 16 when doubled. Thus, $\sqrt{16} = 4$. 5. E 7. D
 9. C 11. C 13. A 15. (a) not a real number (b) negative
 (c) 0 17. -9 19. 6 21. -4 23. -8 25. 6 27. -2 29. It is not a real number. 31. 2 33. It is not a real number. 35. $\frac{8}{9}$ 37. $\frac{4}{3}$
 39. $-\frac{1}{2}$ 41. 3 43. 0.5 45. -0.7 47. 0.1

In Exercises 49-55, we give the domain and then the range.

49. 
 $[-3, \infty); [0, \infty)$ 51. 
 $[0, \infty); [-2, \infty)$ 53. 
 $(-\infty, \infty); (-\infty, \infty)$ 55. 
 $(-\infty, \infty); (-\infty, \infty)$

57. 12 59. 10 61. 2 63. -9 65. -5 67. $|x|$ 69. $|z|$ 71. x
 73. x^5 75. $|x|^5$ (or $|x^5|$) 77. 97.381 79. 16.863 81. -9.055
 83. 7.507 85. 3.162 87. 1.885 89. (a) 1,183,000 cycles per sec
 (b) 118,000 cycles per sec 91. 10 mi 93. 1.732 amps
 95. 437,000 mi² 97. 330 m²





Section 7.2

1. C 3. A 5. H 7. B 9. D 11. In $27^{1/3}$, the base and exponent should not be multiplied. The denominator of the rational exponent is the index of the radical, $\sqrt[3]{27}$. The correct answer is 3. 13. 13 15. 9

17. 2 19. $\frac{8}{9}$ 21. -3 23. It is not a real number. 25. 1000
 27. 27 29. -1024 31. 16 33. $\frac{1}{8}$ 35. $\frac{1}{512}$ 37. $\frac{9}{25}$ 39. $\frac{27}{8}$
 41. $\sqrt{10}$ 43. $(\sqrt[4]{8})^3$ 45. $5(\sqrt[3]{x})^2$ 47. $9(\sqrt[8]{q})^5 - (\sqrt[3]{2x})^2$
 49. $\frac{1}{(\sqrt[5]{x})^3}$ 51. $(\sqrt[3]{2y+x})^2$ 53. $15^{1/2}$ 55. 64 57. 64
 59. x 61. x^{10} 63. 9 65. 4 67. y 69. $x^{5/12}$ 71. $k^{2/3}$
 73. x^3y^8 75. $\frac{1}{x^{10/3}}$ 77. $\frac{1}{m^{1/4}n^{3/4}}$ 79. $\frac{y^{17/3}}{x^8}$ 81. $\frac{c^{11/3}}{b^{11/4}}$ 83. $\frac{q^{5/3}}{9p^{7/2}}$
 85. $p + 2p^2$ 87. $k^{7/4} - k^{3/4}$ 89. $6 + 18a$ 91. $-5x^2 + 5x$
 93. $x^{17/20}$ 95. $t^{8/15}$ 97. $\frac{1}{x^{3/2}}$ 99. $y^{5/6}z^{1/3}$ 101. $m^{1/12}$ 103. $x^{1/24}$
 105. $\sqrt{a^2 + b^2} = \sqrt{3^2 + 4^2} = 5$; $a + b = 3 + 4 = 7$; $5 \neq 7$
 107. 4.5 hr 109. 19.0° ; The table gives 19° . 111. 4.2° ; The table gives 4° .

Section 7.3

1. D 3. B 5. D 7. The student “dropped” the index, 3. The correct product is $\sqrt[3]{65}$. 9. $\sqrt{9}$, or 3 11. $\sqrt{36}$, or 6 13. $\sqrt{30}$ 15. $\sqrt{14x}$
 17. $\sqrt{42pqr}$ 19. $\sqrt[3]{10}$ 21. $\sqrt[3]{14xy}$ 23. $\sqrt[4]{33}$ 25. $\sqrt[4]{6xy^2}$
 27. This expression cannot be simplified by the product rule. 29. $\frac{8}{11}$
 31. $\frac{\sqrt{3}}{5}$ 33. $\frac{\sqrt{x}}{5}$ 35. $\frac{p^3}{9}$ 37. $-\frac{3}{4}$ 39. $\frac{\sqrt[3]{r^2}}{2}$ 41. $-\frac{3}{x}$ 43. $\frac{1}{x^3}$
 45. $\sqrt{12}$ can be simplified further. The *greatest* perfect square factor that divides into 48 is 16, not 4. Thus, $\sqrt{48} = \sqrt{16 \cdot 3} = \sqrt{16} \cdot \sqrt{3} = 4\sqrt{3}$.
 47. $2\sqrt{3}$ 49. $12\sqrt{2}$ 51. $-4\sqrt{2}$ 53. $-2\sqrt{7}$ 55. This radical cannot be simplified further. 57. $4\sqrt[3]{2}$ 59. $2\sqrt[3]{5}$ 61. $-2\sqrt[3]{2}$
 63. $-4\sqrt[4]{2}$ 65. $2\sqrt[5]{2}$ 67. $-3\sqrt[5]{2}$ 69. $2\sqrt[6]{2}$ 71. $6k\sqrt{2}$
 73. $12xy^4\sqrt{xy}$ 75. $11x^3$ 77. $-3t^4$ 79. $-10m^4z^2$ 81. $5a^2b^3c^4$
 83. $\frac{1}{2}r^2t^5$ 85. $5x\sqrt{2x}$ 87. $-10r^5\sqrt{5r}$ 89. $x^3y^4\sqrt{13x}$ 91. $2z^2w^3$
 93. $-2zt^2\sqrt[3]{2z^2t}$ 95. $3x^3y^4$ 97. $-3r^3s^2\sqrt[4]{2r^3s}$ 99. $\frac{y^5\sqrt{y}}{6}$
 101. $\frac{x^5\sqrt[3]{x}}{3}$ 103. $4\sqrt{3}$ 105. $\sqrt{5}$ 107. $x^2\sqrt{x}$ 109. $x\sqrt[5]{x^3}$
 111. $\sqrt[6]{432}$ 113. $\sqrt[12]{6912}$ 115. $\sqrt[6]{x^5}$ 117. 5 119. $8\sqrt{2}$
 121. $2\sqrt{14}$ 123. 13 125. $9\sqrt{2}$ 127. $\sqrt{17}$ 129. 5
 131. $6\sqrt{2}$ 133. $\sqrt{5y^2 - 2xy + x^2}$ 135. (a) B (b) C
 (c) D (d) A 137. $x^2 + y^2 = 144$ 139. $(x + 4)^2 + (y - 3)^2 = 4$
 141. $(x + 8)^2 + (y + 5)^2 = 5$



151. 42.0 in. 153. 581
 155. (a) $d = 1.224\sqrt{h}$ (b) 15.3 mi
 157. $s = 13$ units 158. $6\sqrt{13}$ sq. units
 159. $h = \sqrt{13}$ units 160. $3\sqrt{13}$ sq. units
 161. $6\sqrt{13}$ sq. units 162. The answers are equal, both $6\sqrt{13}$ sq. units, as expected.

Section 7.4

1. B 3. The terms 3 and $3xy$ are not like terms and cannot be combined. The expression $(3 + 3xy)\sqrt[3]{xy^2}$ cannot be simplified further. 5. -4 7. $8\sqrt{10}$ 9. $-\sqrt{5}$ 11. $7\sqrt{3}$
 13. The expression cannot be simplified further. 15. $14\sqrt[3]{2}$
 17. $24\sqrt{2}$ 19. $20\sqrt{5}$ 21. $4\sqrt{2x}$ 23. $-11m\sqrt{2}$ 25. $5\sqrt[4]{2}$
 27. $7\sqrt[3]{2}$ 29. $2\sqrt[3]{x}$ 31. $-7\sqrt[3]{x^2y}$ 33. $-x\sqrt[3]{xy^2}$ 35. $19\sqrt[4]{2}$
 37. $x\sqrt[4]{xy}$ 39. $9\sqrt[4]{2a^3}$ 41. $(4 + 3xy)\sqrt[3]{xy^2}$ 43. $4t\sqrt[3]{3st} - 3s\sqrt[3]{3st}$
 45. $4x\sqrt[3]{x} + 6x\sqrt[4]{x}$ 47. $2\sqrt{2} - 2$ 49. $\frac{5\sqrt{5}}{6}$ 51. $\frac{7\sqrt{2}}{6}$
 53. $\frac{5\sqrt{2}}{3}$ 55. $5\sqrt{2} + 4$ 57. $\frac{5 + 3x}{x^4}$ 59. $\frac{30\sqrt{2} - 21}{14}$
 61. $\frac{m\sqrt[3]{m^2}}{2}$ 63. $\frac{3x\sqrt[3]{2} - 4\sqrt[3]{5}}{x^3}$ 65. (a) $\sqrt{7}$ (b) 2.645751311
 (c) 2.645751311 (d) equal 67. A; 42 m 69. $(12\sqrt{5} + 5\sqrt{3})$ in.
 71. $(24\sqrt{2} + 12\sqrt{3})$ in. 73. $2\sqrt{106} + 4\sqrt{2}$

Section 7.5

1. E 3. A 5. D 7. $3\sqrt{6} + 2\sqrt{3}$ 9. $6 - 4\sqrt{3}$
 11. $20\sqrt{2}$ 13. $6 - \sqrt{6}$ 15. $\sqrt{6} + \sqrt{2} + \sqrt{3} + 1$
 17. -1 19. 6 21. $\sqrt{22} + \sqrt{55} - \sqrt{14} - \sqrt{35}$ 23. $8 - \sqrt{15}$
 25. $9 + 4\sqrt{5}$ 27. $26 - 2\sqrt{105}$ 29. $4 - \sqrt[3]{36}$
 31. 10 33. $6x + 3\sqrt{x} - 2\sqrt{5x} - \sqrt{5}$ 35. $9r - s$
 37. $4\sqrt[3]{4y^2} - 19\sqrt[3]{2y} - 5$ 39. $3x - 4$ 41. Because 6 and $4\sqrt{3}$ are not like terms, they cannot be combined. The expression $6 - 4\sqrt{3}$ cannot be simplified further. 43. $\sqrt{7}$ 45. $5\sqrt{3}$ 47. $\frac{\sqrt{6}}{2}$
 49. $\frac{9\sqrt{15}}{5}$ 51. $-\frac{7\sqrt{3}}{12}$ 53. $\frac{\sqrt{14}}{2}$ 55. $-\frac{\sqrt{14}}{10}$ 57. $\frac{2\sqrt{6x}}{x}$
 59. $-\frac{8\sqrt{3k}}{k}$ 61. $-\frac{5m^2\sqrt{6mn}}{n^2}$ 63. $\frac{12x^3\sqrt{2xy}}{y^5}$ 65. $\frac{5\sqrt{2my}}{y^2}$
 67. $-\frac{4k\sqrt{3z}}{z}$ 69. $\frac{\sqrt[3]{18}}{3}$ 71. $\frac{\sqrt[3]{12}}{3}$ 73. $\frac{\sqrt[3]{18}}{4}$ 75. $-\frac{\sqrt[3]{2pr}}{r}$
 77. $\frac{x^2\sqrt[3]{y^2}}{y}$ 79. $\frac{2\sqrt[4]{x^3}}{x}$ 81. $\frac{\sqrt[4]{2yz^3}}{z}$ 83. $\frac{3(4 - \sqrt{5})}{11}$
 85. $3(\sqrt{5} - \sqrt{3})$ 87. $\frac{6\sqrt{2} + 4}{7}$ 89. $\frac{2(3\sqrt{5} - 2\sqrt{3})}{33}$
 91. $2\sqrt{3} + \sqrt{10} - 3\sqrt{2} - \sqrt{15}$ 93. $\sqrt{m} - 2$

95. $\frac{4(\sqrt{x} + 2\sqrt{y})}{x - 4y}$ 97. $\frac{x - 2\sqrt{xy} + y}{x - y}$ 99. $\frac{5\sqrt{k}(2\sqrt{k} - \sqrt{q})}{4k - q}$
 101. $3 + 2\sqrt{6}$ 103. $1 - \sqrt{5}$ 105. $\frac{4 - 2\sqrt{2}}{3}$ 107. $\frac{6 + 2\sqrt{6p}}{3}$
 109. $\frac{3\sqrt{x+y}}{x+y}$ 111. $\frac{p\sqrt{p+2}}{p+2}$ 113. Each expression is approximately equal to 0.2588190451.
 115. $\frac{33}{8(6 + \sqrt{3})}$ 116. $\frac{11}{2(2\sqrt{5} + 3)}$
 117. $\frac{4x - y}{3x(2\sqrt{x} + \sqrt{y})}$ 118. $\frac{p - 9q}{4q(\sqrt{p} + 3\sqrt{q})}$

SUMMARY EXERCISES Performing Operations with Radicals and Rational Exponents

- The radicand is a fraction, $\frac{2}{5}$.
- The exponent in the radicand and the index of the radical have greatest common factor 5.
- The denominator contains a radical, $\sqrt[3]{10}$.
- The radicand has two factors, x and y , that are raised to powers greater than the index, 3.
- $-6\sqrt{10}$
- $7 - \sqrt{14}$
- $2 + \sqrt{6} - 2\sqrt{3} - 3\sqrt{2}$
- $4\sqrt{2}$
- $73 + 12\sqrt{35}$
- $\frac{-\sqrt{6}}{2}$
- $4(\sqrt{7} - \sqrt{5})$
- $-3 + 2\sqrt{2}$
- -44
- $\frac{\sqrt{x} + \sqrt{5}}{x - 5}$
- $2abc^3\sqrt[3]{b^2}$
- $5\sqrt[3]{3}$
- $3(\sqrt{5} - 2)$
- $\frac{\sqrt{15x}}{5x}$
- $\frac{8}{5}$
- $\frac{\sqrt{2}}{8}$
- $-\sqrt[3]{100}$
- $11 + 2\sqrt{30}$
- $-3\sqrt{3x}$
- $52 - 30\sqrt{3}$
- $\frac{\sqrt[3]{117}}{9}$
- $3\sqrt{2} + \sqrt{15} + \sqrt{42} + \sqrt{35}$
- $2\sqrt[4]{27}$
- $\frac{x\sqrt[3]{x^2}}{y}$
- $-4\sqrt{3} - 3$
- $7 + 4 \cdot 3^{1/2}$, or $7 + 4\sqrt{3}$
- $3\sqrt[3]{2x^2}$
- -2
- $3^{5/6}$
- $\frac{x^{5/3}}{y}$
- $xy^{6/5}$
- $x^{10}y$
- $\frac{1}{25x^2}$
- $\frac{-6y^{1/6}}{x^{1/24}}$

Section 7.6


- (a) yes (b) no
- (a) yes (b) no
- There is no solution. The radical expression, which is nonnegative, cannot equal a negative number. The solution set is \emptyset .
- $\{11\}$
- $\{\frac{1}{3}\}$
- \emptyset
- $\{5\}$
- $\{18\}$
- $\{5\}$
- $\{4\}$
- $\{17\}$
- $\{5\}$
- \emptyset
- $\{0\}$
- $\{1\}$
- $\{-1, 3\}$
- $\{0\}$
- \emptyset
- $\{1\}$
- We cannot just square each term. The right side should be $(8 - x)^2 = 64 - 16x + x^2$. The correct first step is $3x + 4 = 64 - 16x + x^2$. The solution set is $\{4\}$.
- $\{7\}$
- $\{7\}$
- $\{4, 20\}$
- \emptyset
- $\{\frac{5}{4}\}$
- $\{9\}$
- $\{1\}$
- $\{14\}$
- $\{-1\}$
- $\{8\}$
- $\{0\}$
- \emptyset
- $\{-4\}$
- $\{9, 17\}$
- $\{\frac{1}{4}, 1\}$
- $L = CZ^2$
- $K = \frac{V^2m}{2}$
- $M = \frac{r^2F}{m}$
- (a) $r = \frac{a}{4\pi^2N^2}$ (b) $a = 4\pi^2N^2r$

Section 7.7


- nonreal complex, complex
- real, complex
- pure imaginary, nonreal complex, complex
- i
- -1
- $-i$
- $13i$
- $-12i$
- $i\sqrt{5}$
- $4i\sqrt{3}$
- It is incorrect to use the product rule for radicals before using the definition of $\sqrt{-b}$. The correct product is -15 .
- -15
- $-\sqrt{105}$
- -10
- $i\sqrt{33}$
- $5i\sqrt{6}$
- $\sqrt{3}$
- $5i$
- -2
- $-1 + 7i$
- 0
- $7 + 3i$
- -2
- $1 + 13i$
- $6 + 6i$
- $4 + 2i$
- -81
- -16
- $-10 - 30i$
- $10 - 5i$
- $-9 + 40i$
- $-16 + 30i$
- 153
- 97
- 4
- (a) $a - bi$ (b) $a^2; b^2$
- $1 + i$
- $2 + 2i$
- $-1 + 2i$
- $-\frac{5}{13} - \frac{12}{13}i$
- $1 - 3i$
- $1 + 3i$
- -1
- i
- -1
- $-i$
- $-i$
- 1
- Because $i^{20} = (i^4)^5 = 1^5 = 1$, the student multiplied by 1, which is justified by the identity property for multiplication.
- $\frac{1}{2} + \frac{1}{2}i$
- Substitute $1 + 5i$ for x in the equation. A true statement results—that is, $(1 + 5i)^2 - 2(1 + 5i) + 26$ will simplify to 0 when the operations are applied. Thus, $1 + 5i$ is a solution.
- Substituting $1 - 5i$ for x in the equation results in a true statement, indicating that $1 - 5i$ is a solution.
- They are complex conjugates.
- Substituting $3 + 2i$ for x in the equation results in a true statement, indicating that $3 + 2i$ is a solution.
- $3 - 2i$; Substituting $3 - 2i$ for x in the equation results in a true statement, indicating that $3 - 2i$ is a solution.

Chapter 7 Review Exercises

- 42
- -17
- 6
- -5
- -3
- -2
- $|x|$
- x
- $|x|^5$ (or $|x^5|$)
- $\sqrt[n]{a}$ is not a real number if n is even and a is negative.



domain: $[1, \infty)$;
range: $[0, \infty)$



domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$


- -6.856
- -5.053
- 4.960
- 4.729
- -7.937
- -5.292
- 1.9 sec
- 66 in.²
- 7
- -11
- 32
- -4
- $-\frac{216}{125}$
- -32
- $\frac{1000}{27}$
- It is not a real number.
- $10\sqrt{x}$
- $\frac{1}{(\sqrt[3]{3a+b})^5}$
- $7^{9/2}$
- 1331
- r^4
- z
- 25
- 96
- $a^{2/3}$
- $\frac{1}{y^{1/2}}$
- $\frac{z^{1/2}x^{8/5}}{4}$
- $r^{1/2} + r$
- $y^{8/15}$
- $\frac{1}{x^{1/2}}$
- $p^{1/2}$
- $k^{9/4}$
- $m^{13/3}$
- $t^{3/2}$
- $x^{1/8}$
- $x^{1/15}$
- $x^{1/36}$
- The product rule for exponents applies only if the bases are the same.
- $\sqrt{66}$
- $\sqrt{5r}$
- $\sqrt[3]{30}$
- $\sqrt[4]{21}$
- $2\sqrt{5}$
- $5\sqrt{3}$
- $-5\sqrt{5}$
- $-3\sqrt[3]{4}$
- $10y^3\sqrt{y}$
- $4pq^2\sqrt[3]{p}$
- $3a^2b\sqrt[3]{4a^2b^2}$

62. $2r^2t\sqrt[3]{79r^2t}$ 63. $\frac{y\sqrt{y}}{12}$ 64. $\frac{m^5}{3}$ 65. $\frac{\sqrt[3]{r^2}}{2}$ 66. $\frac{a^2\sqrt[4]{a}}{3}$


67. $\sqrt{15}$ 68. $p\sqrt{p}$ 69. $\sqrt[12]{2000}$ 70. $\sqrt[10]{x^7}$ 71. 10

72. $\sqrt{197}$ 73. $x^2 + y^2 = 121$ 74. $(x + 2)^2 + (y - 4)^2 = 9$


75. $(x + 1)^2 + (y + 3)^2 = 25$ 76. $(x - 4)^2 + (y - 2)^2 = 36$



center: (0, 0);
radius: 5



center: (-3, 3);
radius: 3



center: (2, -5);
radius: 3

80. It is impossible for the sum of the squares of two real numbers to be negative. 81. $-11\sqrt{2}$ 82. $23\sqrt{5}$ 83. $7\sqrt{3y}$ 84. $26m\sqrt{6m}$

85. $19\sqrt[3]{2}$ 86. $-8\sqrt[4]{2}$ 87. $1 - \sqrt{3}$ 88. 2 89. $9 - 7\sqrt{2}$

90. $15 - 2\sqrt{26}$ 91. 29 92. $2\sqrt[3]{2y^2} + 2\sqrt[3]{4y} - 3$ 93. $\frac{\sqrt{30}}{5}$

94. $-3\sqrt{6}$ 95. $\frac{3\sqrt[3]{7py}}{y}$ 96. $\frac{\sqrt{22}}{4}$ 97. $-\frac{\sqrt[3]{45}}{5}$ 98. $\frac{3m\sqrt[3]{4n}}{n^2}$

99. $\frac{\sqrt{2} - \sqrt{7}}{-5}$ 100. $\frac{5(\sqrt{6} + 3)}{3}$ 101. $\frac{1 - \sqrt{5}}{4}$ 102. $\frac{-6 + \sqrt{3}}{2}$

103. {2} 104. {6} 105. {0, 5} 106. {9} 107. {3} 108. {7}

109. $\{-\frac{1}{2}\}$ 110. {14} 111. \emptyset 112. {7} 113. $H = \sqrt{L^2 - W^2}$

114. 7.9 ft 115. 4i 116. $10i\sqrt{2}$ 117. $-10 - 2i$ 118. $14 + 7i$

119. $-\sqrt{35}$ 120. -45 121. 3 122. $5 + i$ 123. $32 - 24i$

124. $1 - i$ 125. $-i$ 126. 1 127. -1 128. 1

Chapter 7 Mixed Review Exercises

1. $-13ab^2$ 2. $\frac{1}{100}$ 3. $\frac{1}{z^{3/5}}$ 4. $3z^3t^2\sqrt[3]{2t^2}$ 5. 7i 6. $-\frac{\sqrt{3}}{6}$

7. $\frac{\sqrt[3]{60}}{5}$ 8. 1 9. $57\sqrt{2}$ 10. $5 - 11i$ 11. $-5i$ 12. $\frac{1 + \sqrt{6}}{2}$


13. $5 + 12i$ 14. $6x\sqrt[3]{y^2}$ 15. $35 + 15i$ 16. $\sqrt[12]{2000}$

17. $\frac{2\sqrt{z}(\sqrt{z} + 2)}{z - 4}$ 18. $(12\sqrt{3} + 5\sqrt{2})$ ft 19. {5}

20. {-4} 21. $\{\frac{3}{2}\}$ 22. {7}

Chapter 7 Test


[7.1] 1. -29 2. -8 [7.2] 3. 5 [7.1] 4. 21.863 5. -9.405

6.  [7.2] 7. $\frac{125}{64}$ 8. $\frac{1}{256}$ 9. $\frac{9y^{3/10}}{x^2}$ 10. $x^{4/3}y^6$

domain: $[-6, \infty)$;
range: $[0, \infty)$

11. $7^{1/2}$ 12. $a^{11/3}$ [7.3] 13. $\sqrt{145}$

14. 10 15. $(x + 4)^2 + (y - 6)^2 = 25$



$(x - 2)^2 + (y + 3)^2 = 16$
center: (2, -3);
radius: 4

17. $3x^2y^3\sqrt{6x}$ 18. $2ab^3\sqrt[4]{2a^3b}$ 19. $\sqrt[6]{200}$

[7.4] 20. $26\sqrt{5}$ 21. $(2ts - 3t^2)\sqrt[3]{2s^2}$

[7.5] 22. $66 + \sqrt{5}$ 23. $23 - 4\sqrt{15}$

24. $-\frac{\sqrt{10}}{4}$ 25. $\frac{2\sqrt[3]{25}}{5}$

26. $-2(\sqrt{7} - \sqrt{5})$ 27. $3 + \sqrt{6}$ [7.6] 28. (a) 59.8

(b) $T = \frac{V_0^2 - V^2}{-V^2k}$, or $T = \frac{V^2 - V_0^2}{V^2k}$ 29. {-1} 30. {3}

31. {-3} [7.7] 32. $-5 - 8i$ 33. $-2 + 16i$ 34. $3 + 4i$

35. i 36. (a) true (b) true (c) false (d) true

Chapters R-7 Cumulative Review Exercises

[R.4] 1. -47 2. $\frac{43}{10}$ [R.3] 3. $\frac{5}{24}$ [1.1] 4. {-4} 5. {-12}

6. {6} [1.7] 7. $\{-\frac{10}{3}, 1\}$ [1.5] 8. $(-6, \infty)$ 9. (-2, 1)

[1.3] 10. 36 nickels; 64 quarters 11. $2\frac{2}{39}$ L

[2.1] 12.  [2.2, 2.3] 13. $-\frac{3}{2}$; $y = -\frac{3}{2}x$

[2.6] 14. -37 [3.1] 15. {(7, -2)}

[3.2] 16. {(-1, 1, 1)}

[3.3] 17. 2-oz letter: \$0.70; 3-oz letter: \$0.91

[4.3] 18. $-k^3 - 3k^2 - 8k - 9$ [4.5] 19. $8x^2 + 17x - 21$

[4.6] 20. $3y^3 - 3y^2 + 4y + 1 + \frac{-10}{2y + 1}$

[5.2, 5.3] 21. $(2p - 3q)(p - q)$ 22. $(3k^2 + 4)(k - 1)(k + 1)$

23. $(x + 8)(x^2 - 8x + 64)$ [6.1] 24. $\{x | x \text{ is a real number, } x \neq \pm 3\}$

25. $\frac{y}{y + 5}$ [6.2] 26. $\frac{4x + 2y}{(x + y)(x - y)}$ [6.3] 27. $-\frac{9}{4}$ 28. $\frac{-1}{a + b}$

29. $\frac{1}{xy - 1}$ [5.5] 30. $\{-3, -\frac{5}{2}\}$ 31. $\{-\frac{2}{5}, 1\}$ [6.4] 32. \emptyset

[7.6] 33. {3, 4} [6.5] 34. Danielle: 8 mph; Richard: 4 mph

[7.2] 35. $\frac{1}{9}$ [7.3] 36. $2x\sqrt[3]{6x^2y^2}$ [7.4] 37. $7\sqrt{2}$

[7.5] 38. $\frac{\sqrt{10} + 2\sqrt{2}}{2}$ [7.3] 39. $\sqrt{29}$ [7.7] 40. $4 + 2i$

8 QUADRATIC EQUATIONS, INEQUALITIES, AND FUNCTIONS

Section 8.1

1. quadratic; second; two 3. The zero-factor property requires a product equal to 0. The first step should have been to write the equation with 0 on one side. The solution set is $\{-2, 3\}$. 5. $\{-2, -1\}$

7. $\{-3, \frac{1}{3}\}$ 9. $\{-\frac{3}{5}, 2\}$ 11. {2, 4} 13. $\{\frac{1}{2}, 4\}$ 15. $\{-\frac{3}{2}, -\frac{1}{4}\}$

17. $\{\pm 9\}$ 19. $\{\pm\sqrt{17}\}$ 21. $\{\pm 4\sqrt{2}\}$ 23. $\{\pm 2\sqrt{5}\}$

25. $\{\pm 2\sqrt{6}\}$ 27. $\{\pm 3\sqrt{3}\}$ 29. $\{\pm 4\sqrt{2}\}$ 31. $\{-7, 3\}$

33. $\{-1, 13\}$ 35. $\{4 \pm \sqrt{3}\}$ 37. $\{-5 \pm 4\sqrt{3}\}$
 39. $\left\{\frac{1 \pm \sqrt{7}}{3}\right\}$ 41. $\left\{\frac{-1 \pm 2\sqrt{6}}{4}\right\}$ 43. $\left\{\frac{2 \pm 2\sqrt{3}}{5}\right\}$
 45. 5.6 sec 47. Solve $(2x + 1)^2 = 5$ by the square root property. Solve $x^2 + 4x = 12$ by completing the square. 49. 9; $(x + 3)^2$
 51. 36; $(p - 6)^2$ 53. $\frac{81}{4}; \left(q + \frac{9}{2}\right)^2$ 55. $\{-4, 6\}$
 57. $\{-2 \pm \sqrt{6}\}$ 59. $\{-5 \pm \sqrt{7}\}$ 61. $\left\{-\frac{8}{3}, 3\right\}$
 63. $\left\{\frac{-7 \pm \sqrt{53}}{2}\right\}$ 65. $\left\{\frac{-5 \pm \sqrt{41}}{4}\right\}$ 67. $\left\{\frac{5 \pm \sqrt{15}}{5}\right\}$
 69. $\left\{\frac{4 \pm \sqrt{3}}{3}\right\}$ 71. $\left\{\frac{2 \pm \sqrt{3}}{3}\right\}$ 73. $\{1 \pm \sqrt{2}\}$
 75. $\{\pm 10i\}$ 77. $\{\pm 2i\sqrt{3}\}$ 79. $\{-3 \pm 2i\}$
 81. $\{5 \pm i\sqrt{3}\}$ 83. $\left\{\frac{1}{6} \pm \frac{\sqrt{2}}{3}i\right\}$ 85. $\{-2 \pm 3i\}$
 87. $\{-3 \pm i\sqrt{3}\}$ 89. $\left\{-\frac{2}{3} \pm \frac{2\sqrt{2}}{3}i\right\}$ 91. $\left\{-\frac{5}{2} \pm \frac{\sqrt{15}}{2}i\right\}$
 93. $\{\pm \sqrt{b}\}$ 95. $\left\{\pm \frac{\sqrt{b^2 + 16}}{2}\right\}$ 97. $\left\{\frac{2b \pm \sqrt{3a}}{5}\right\}$
 99. x^2 100. x 101. $6x$ 102. 1 103. 9
 104. $(x + 3)^2$, or $x^2 + 6x + 9$

Section 8.2

1. No. The fraction bar should extend under the term $-b$. The correct formula is $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$. 3. The last step is wrong. Because 5 is not a common factor in the numerator, the fraction cannot be simplified. The solution set is $\left\{\frac{5 \pm \sqrt{5}}{10}\right\}$. 5. $\{3, 5\}$ 7. $\left\{-\frac{5}{2}, \frac{2}{3}\right\}$
 9. $\left\{-\frac{3}{2}\right\}$ 11. $\left\{\frac{1}{6}\right\}$ 13. $\left\{\frac{-2 \pm \sqrt{2}}{2}\right\}$ 15. $\left\{\frac{1 \pm \sqrt{3}}{2}\right\}$
 17. $\{5 \pm \sqrt{7}\}$ 19. $\left\{\frac{-1 \pm \sqrt{2}}{2}\right\}$ 21. $\left\{\frac{-1 \pm \sqrt{7}}{3}\right\}$
 23. $\{1 \pm \sqrt{5}\}$ 25. $\left\{\frac{-2 \pm \sqrt{10}}{2}\right\}$ 27. $\{-1 \pm 3\sqrt{2}\}$
 29. $\left\{\frac{1 \pm \sqrt{29}}{2}\right\}$ 31. $\left\{\frac{-4 \pm \sqrt{91}}{3}\right\}$ 33. $\left\{\frac{-3 \pm \sqrt{57}}{8}\right\}$
 35. $\left\{\frac{3}{2} \pm \frac{\sqrt{15}}{2}i\right\}$ 37. $\{3 \pm i\sqrt{5}\}$ 39. $\left\{\frac{1}{2} \pm \frac{\sqrt{6}}{2}i\right\}$
 41. $\left\{-\frac{2}{3} \pm \frac{\sqrt{2}}{3}i\right\}$ 43. $\left\{\frac{1}{2} \pm \frac{1}{4}i\right\}$ 45. $\left\{\frac{4}{5} \pm \frac{2\sqrt{6}}{5}i\right\}$
 47. 0; B; zero-factor property 49. 8; C; quadratic formula
 51. 49; A; zero-factor property 53. -80 ; D; quadratic formula

55. (a) 25; zero-factor property; $\left\{-3, -\frac{4}{3}\right\}$ (b) 44; quadratic formula;
 $\left\{\frac{7 \pm \sqrt{11}}{2}\right\}$ 57. -10 or 10 59. 16 61. 25 63. $b = \frac{44}{5}; \frac{3}{10}$

Section 8.3

1. Multiply by the LCD, x . 3. Substitute a variable for $x^2 + x$.
 5. The proposed solution -1 does not check. The solution set is $\{4\}$.
 7. $\{-2, 7\}$ 9. $\{-4, 7\}$ 11. $\left\{-\frac{2}{3}, 1\right\}$ 13. $\left\{-\frac{14}{17}, 5\right\}$
 15. $\left\{-\frac{11}{7}, 0\right\}$ 17. $\left\{\frac{-1 \pm \sqrt{13}}{2}\right\}$ 19. $\left\{-\frac{8}{3}, -1\right\}$
 21. $\left\{\frac{2 \pm \sqrt{22}}{3}\right\}$ 23. $\left\{\frac{-1 \pm \sqrt{5}}{4}\right\}$ 25. (a) $(20 - t)$ mph
 (b) $(20 + t)$ mph 27. the rate of her boat in still water; $x - 5$; $x + 5$;
 row 1 of table: $15, x - 5, \frac{15}{x - 5}$; row 2 of table: $15, x + 5, \frac{15}{x + 5}$;
 $\frac{15}{x - 5} + \frac{15}{x + 5} = 4$; 10 mph 29. 25 mph 31. 50 mph
 33. 3.6 hr 35. Rusty: 25.0 hr; Nancy: 23.0 hr 37. 9 min
 39. $\{2, 5\}$ 41. $\{3\}$ 43. $\left\{\frac{8}{9}\right\}$ 45. $\{9\}$ 47. $\left\{\frac{2}{5}\right\}$ 49. $\{-2\}$
 51. $\{\pm 2, \pm 5\}$ 53. $\left\{\pm 1, \pm \frac{3}{2}\right\}$ 55. $\{\pm 2, \pm 2\sqrt{3}\}$
 57. $\{-6, -5\}$ 59. $\left\{-\frac{16}{3}, -2\right\}$ 61. $\{-8, 1\}$ 63. $\{-64, 27\}$
 65. $\left\{\pm 1, \pm \frac{27}{8}\right\}$ 67. $\left\{-\frac{1}{3}, \frac{1}{6}\right\}$ 69. $\left\{-\frac{1}{2}, 3\right\}$ 71. $\left\{\pm \frac{\sqrt{6}}{3}, \pm \frac{1}{2}\right\}$
 73. $\{3, 11\}$ 75. $\{25\}$ 77. $\left\{-\sqrt[3]{5}, -\frac{\sqrt[3]{4}}{2}\right\}$ 79. $\left\{\frac{4}{3}, \frac{9}{4}\right\}$
 81. $\left\{\pm \frac{\sqrt{9 + \sqrt{65}}}{2}, \pm \frac{\sqrt{9 - \sqrt{65}}}{2}\right\}$ 83. $\left\{\pm 1, \pm \frac{\sqrt{6}}{2}i\right\}$

SUMMARY EXERCISES Applying Methods for Solving Quadratic Equations

1. square root property 2. zero-factor property 3. quadratic formula
 4. quadratic formula 5. zero-factor property 6. square root property
 7. $\{\pm \sqrt{7}\}$ 8. $\left\{-\frac{3}{2}, \frac{5}{3}\right\}$ 9. $\{-3 \pm \sqrt{5}\}$ 10. $\{-2, 8\}$
 11. $\left\{-\frac{3}{2}, 4\right\}$ 12. $\left\{-3, \frac{1}{3}\right\}$ 13. $\left\{\frac{2 \pm \sqrt{2}}{2}\right\}$ 14. $\{\pm 2i\sqrt{3}\}$
 15. $\left\{\frac{1}{2}, 2\right\}$ 16. $\{\pm 1, \pm 3\}$ 17. $\left\{\frac{-3 \pm 2\sqrt{2}}{2}\right\}$ 18. $\left\{\frac{4}{5}, 3\right\}$
 19. $\{\pm \sqrt{2}, \pm \sqrt{7}\}$ 20. $\left\{\frac{1 \pm \sqrt{5}}{4}\right\}$ 21. $\left\{-\frac{1}{2} \pm \frac{\sqrt{3}}{2}i\right\}$
 22. $\left\{-\frac{\sqrt[3]{175}}{5}, 1\right\}$ 23. $\left\{\frac{3}{2}\right\}$ 24. $\left\{\frac{2}{3}\right\}$ 25. $\{\pm 6\sqrt{2}\}$
 26. $\left\{-\frac{2}{3}, 2\right\}$ 27. $\{-4, 9\}$ 28. $\{\pm 13\}$ 29. $\left\{1 \pm \frac{\sqrt{3}}{3}i\right\}$
 30. $\{3\}$ 31. $\left\{\frac{1}{6} \pm \frac{\sqrt{47}}{6}i\right\}$ 32. $\left\{-\frac{1}{3}, \frac{1}{6}\right\}$

Section 8.4

1. Find a common denominator, and then multiply both sides by the common denominator. 3. Write it in standard form (with 0 on one side,

in decreasing powers of w). 5. $m = \sqrt{p^2 - n^2}$ 7. $t = \frac{\pm \sqrt{dk}}{k}$

9. $r = \frac{\pm \sqrt{S\pi}}{2\pi}$ 11. $d = \frac{\pm \sqrt{skI}}{I}$ 13. $v = \frac{\pm \sqrt{kAF}}{F}$

15. $r = \frac{\pm \sqrt{3\pi Vh}}{\pi h}$ 17. $t = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}$ 19. $h = \frac{D^2}{k}$

21. $\ell = \frac{p^2 g}{k}$ 23. $R = \frac{E^2 - 2pr \pm E\sqrt{E^2 - 4pr}}{2p}$

25. $r = \frac{5pc}{4}$ or $r = -\frac{2pc}{3}$ 27. $I = \frac{-cR \pm \sqrt{c^2 R^2 - 4cL}}{2cL}$

29. 7.9, 8.9, 11.9 31. eastbound ship: 80 mi; southbound ship: 150 mi

33. 8 in., 15 in., 17 in. 35. length: 24 ft; width: 10 ft 37. 2 ft


39. 7 m by 12 m 41. 20 in. by 12 in. 43. 1 sec and 8 sec

45. 2.4 sec and 5.6 sec 47. 9.2 sec 49. It reaches its *maximum* height at 5 sec because this is the only time it reaches 400 ft.

51. \$1.50 53. 0.035, or 3.5% 55. 5.5 m per sec 57. 5 or 14

59. (a) \$600 billion (b) \$597 billion; They are about the same.

61. 2008 63. isosceles triangle; right triangle 64. no; The formula for the Pythagorean theorem involves the *squares* of the sides, not the square roots.

65.  $\sqrt{9} + \sqrt{9} = 3 + 3 = 6;$
 $\sqrt{4} = 2;$
 $6 \neq 2$

66. The sum of the squares of the two shorter sides (legs) of a right triangle is equal to the square of the longest side (hypotenuse).


Section 8.5

1. (a) B (b) C (c) A (d) D 3. (a) D (b) B (c) F (d) C


(e) A (f) E 5. (0, 0) 7. (0, 0) 9. (0, 4) 11. (1, 0)

13. (-3, -4) 15. (5, 6) 17. down; wider 19. up; narrower


21. down; narrower




vertex: (0, 0);
axis: $x = 0$;
domain: $(-\infty, \infty)$;
range: $[0, \infty)$




vertex: (0, 0);
axis: $x = 0$;
domain: $(-\infty, \infty)$;
range: $(-\infty, 0]$




vertex: (0, -1);
axis: $x = 0$;
domain: $(-\infty, \infty)$;
range: $[-1, \infty)$




vertex: (0, 2);
axis: $x = 0$;
domain: $(-\infty, \infty)$;
range: $(-\infty, 2]$




vertex: (4, 0);
axis: $x = 4$;
domain: $(-\infty, \infty)$;
range: $[0, \infty)$




vertex: (-2, -1);
axis: $x = -2$;
domain: $(-\infty, \infty)$;
range: $[-1, \infty)$




vertex: (1, -3);
axis: $x = 1$;
domain: $(-\infty, \infty)$;
range: $[-3, \infty)$




vertex: (2, -4);
axis: $x = 2$;
domain: $(-\infty, \infty)$;
range: $[-4, \infty)$



vertex: (2, -3);
axis: $x = 2$;
domain: $(-\infty, \infty)$;
range: $[-3, \infty)$



vertex: (-3, 4);
axis: $x = -3$;
domain: $(-\infty, \infty)$;
range: $(-\infty, 4]$




vertex: (-1, 2);
axis: $x = -1$;
domain: $(-\infty, \infty)$;
range: $(-\infty, 2]$

45. linear function; positive

47. quadratic function; positive

49. quadratic function; negative



(b) quadratic function

(c) negative

(d) $y = -0.4x^2 + 4.1x + 95$


(e) 2009: 101.6 million households; 2015: 102.2 million households; The model approximates the data fairly well.

Section 8.6


1. If x is squared, it has a vertical axis. If y is squared, it has a horizontal axis. 3. Find the discriminant of the function. If it is positive, there are two x -intercepts. If it is 0, there is one x -intercept (at the vertex), and if it is negative, there is no x -intercept. 5. A, D are vertical parabolas. B, C are horizontal parabolas. 7. (-4, -6) 9. (1, -3)

11. $(-\frac{1}{2}, -\frac{29}{4})$ 13. (-1, 3); up; narrower; no x -intercepts


15. $(\frac{5}{2}, \frac{37}{4})$; down; same shape; two x -intercepts 17. (-3, -9); to the right; wider 19. F 21. C 23. D




vertex: $(-4, -6)$;
axis: $x = -4$;
domain: $(-\infty, \infty)$;
range: $[-6, \infty)$




vertex: $(-2, -1)$;
axis: $x = -2$;
domain: $(-\infty, \infty)$;
range: $[-1, \infty)$




vertex: $(1, -3)$;
axis: $x = 1$;
domain: $(-\infty, \infty)$;
range: $(-\infty, -3]$




vertex: $(-1, 5)$;
axis: $x = -1$;
domain: $(-\infty, \infty)$;
range: $(-\infty, 5]$




vertex: $(1, -2)$;
axis: $y = -2$;
domain: $[1, \infty)$;
range: $(-\infty, \infty)$



vertex: $(-1, 3)$;
axis: $y = 3$;
domain: $(-\infty, -1]$;
range: $(-\infty, \infty)$




vertex: $(1, 5)$;
axis: $y = 5$;
domain: $(-\infty, 1]$;
range: $(-\infty, \infty)$



vertex: $(-7, -2)$;
axis: $y = -2$;
domain: $[-7, \infty)$;
range: $(-\infty, \infty)$

41. 20 and 20 43. 140 ft by 70 ft; 9800 ft² 45. 2 sec; 65 ft
47. 20 units; \$210 49. 16 ft; 2 sec 51. (a) The coefficient of x^2 is negative because a parabola that models the data must open down. (b) (3.32, 3.58); In 2012, the average retail price of regular unleaded gasoline reached a maximum value of \$3.58 per gallon.
















53. (a) $R(x) = (100 - x)(200 + 4x)$
 $= 20,000 + 200x - 4x^2$








(c) 25 (d) \$22,500

Section 8.7

1. $(-\infty, -4] \cup [3, \infty)$ 3. (a) $\{1, 3\}$ (b) $(-\infty, 1) \cup (3, \infty)$
(c) $(1, 3)$ 5. (a) $\{-2, 5\}$ (b) $[-2, 5]$ (c) $(-\infty, -2] \cup [5, \infty)$



25. $(-\infty, \infty)$ 27. \emptyset
29. \emptyset 31. $(-\infty, \infty)$



55. 3 sec and 13 sec 56. between 3 sec and 13 sec 57. at 0 sec (the time when it is initially projected) and at 16 sec (the time when it hits the ground) 58. between 0 and 3 sec and between 13 and 16 sec

Chapter 8 Review Exercises

1. $\{\pm 11\}$ 2. $\{\pm \sqrt{3}\}$ 3. $\{-\frac{15}{2}, \frac{5}{2}\}$ 4. $\{\frac{2}{3} \pm \frac{5}{3}i\}$
5. $\{-2 \pm \sqrt{19}\}$ 6. $\{\frac{1}{2}, 1\}$ 7. 7.8 sec 8. 5.9 sec 9. $\{-\frac{7}{2}, 3\}$
10. $\{\frac{-5 \pm \sqrt{53}}{2}\}$ 11. $\{\frac{1 \pm \sqrt{41}}{2}\}$ 12. $\{-\frac{3}{4} \pm \frac{\sqrt{23}}{4}i\}$

13. $\left\{\frac{2}{3} \pm \frac{\sqrt{2}}{3}i\right\}$ 14. $\left\{\frac{-7 \pm \sqrt{37}}{2}\right\}$ 15. 17; C; quadratic formula

16. 64; A; zero-factor property 17. -92; D; quadratic formula

18. 0; B; zero-factor property 19. $\left\{-\frac{5}{2}, 3\right\}$ 20. $\left\{-\frac{1}{2}, 1\right\}$

21. $\{-4\}$ 22. $\left\{-\frac{11}{6}, -\frac{19}{12}\right\}$ 23. $\left\{-\frac{343}{8}, 64\right\}$ 24. $\{\pm 1, \pm 3\}$


25. 7 mph 26. 40 mph 27. 4.6 hr 28. Zoran: 2.6 hr; Claude: 3.6 hr

29. $v = \frac{\pm \sqrt{rFkw}}{kw}$ 30. $y = \frac{6p^2}{z}$ 31. $t = \frac{3m \pm \sqrt{9m^2 + 24m}}{2m}$

32. 9 ft, 12 ft, 15 ft 33. 12 cm by 20 cm 34. 1 in. 35. 18 in.


36. 3 min 37. (1, 0) 38. (3, 7) 39. (-4, 3) 40. $\left(\frac{2}{3}, -\frac{2}{3}\right)$

41.




vertex: (2, -3);
axis: $x = 2$;
domain: $(-\infty, \infty)$;
range: $[-3, \infty)$

42.




vertex: (2, 3);
axis: $x = 2$;
domain: $(-\infty, \infty)$;
range: $(-\infty, 3]$

43.



vertex: (-4, -3);
axis: $y = -3$;
domain: $[-4, \infty)$;
range: $(-\infty, \infty)$


44.




vertex: (4, 6);
axis: $y = 6$;
domain: $(-\infty, 4]$;
range: $(-\infty, \infty)$

45. 5 sec; 400 ft


46. length: 50 m; width: 50 m;
maximum area: 2500 m²



$(-\infty, -\frac{3}{2}) \cup (4, \infty)$




$[-4, 3]$




$(-\infty, -5] \cup [-2, 3]$

50. \emptyset



$(-\infty, \frac{1}{2}) \cup (2, \infty)$



$[-3, 2]$

Chapter 8 Mixed Review Exercises

1. $R = \frac{\pm \sqrt{Vh - r^2h}}{h}$ 2. $\left\{1 \pm \frac{\sqrt{3}}{3}i\right\}$


3. $\left\{\frac{-11 \pm \sqrt{7}}{3}\right\}$ 4. $d = \frac{\pm \sqrt{SkI}}{I}$ 5. $(-\infty, \infty)$

6. $\{4\}$ 7. $\{\pm \sqrt{4 + \sqrt{15}}, \pm \sqrt{4 - \sqrt{15}}\}$

8. $\left(-5, -\frac{23}{5}\right]$ 9. $\left\{-\frac{5}{3}, -\frac{3}{2}\right\}$ 10. $\{-2, -1, 3, 4\}$

11. $(-\infty, -6) \cup \left(-\frac{3}{2}, 1\right)$ 12. (a) F (b) B (c) C (d) A

(e) E (f) D



vertex: $\left(-\frac{1}{2}, -3\right)$;
axis: $x = -\frac{1}{2}$;
domain: $(-\infty, \infty)$;
range: $[-3, \infty)$

14. 10 mph 15. length: 2 cm;
width: 1.5 cm

16. (a) minimum
(b) 1967; 5.4%

Chapter 8 Test

[8.1] 1. $\left\{-3, \frac{2}{5}\right\}$ 2. $\{\pm 3\sqrt{6}\}$ 3. $\left\{-\frac{8}{7}, \frac{2}{7}\right\}$ 4. $\{-1 \pm \sqrt{5}\}$

[8.2] 5. $\left\{\frac{3 \pm \sqrt{17}}{4}\right\}$ 6. $\left\{\frac{2}{3} \pm \frac{\sqrt{11}}{3}i\right\}$ [8.3] 7. $\left\{\frac{2}{3}\right\}$


[8.1] 8. A [8.2] 9. discriminant: 88; There are two irrational solutions.

[8.1-8.3] 10. $\left\{-\frac{2}{3}, 6\right\}$ 11. $\left\{\frac{-7 \pm \sqrt{97}}{8}\right\}$ 12. $\left\{\pm \frac{1}{3}, \pm 2\right\}$

13. $\left\{-\frac{5}{2}, 1\right\}$ [8.4] 14. $r = \frac{\pm \sqrt{\pi S}}{2\pi}$ [8.3] 15. Terry: 11.1 hr;


Callie: 9.1 hr 16. 7 mph [8.4] 17. 2 ft 18. 16 m [8.5, 8.6] 19. A

20.




vertex: (0, -2);
axis: $x = 0$;
domain: $(-\infty, \infty)$;
range: $[-2, \infty)$

21. $f(x) = -x^2 + 4x - 1$




vertex: (2, 3);
axis: $x = 2$;
domain: $(-\infty, \infty)$;
range: $(-\infty, 3]$

22.




vertex: (2, 2);
axis: $y = 2$;
domain: $(-\infty, 2]$;
range: $(-\infty, \infty)$

23. 160 ft by 320 ft



$(-\infty, -5) \cup \left(\frac{3}{2}, \infty\right)$



$(-\infty, 4) \cup [9, \infty)$

Chapters R-8 Cumulative Review Exercises

[R.1-R.3] 1. -5.38 2. 2 3. 25 4. 0.72 [R.4] 5. -45


[R.2, 7.7] 6. (a) -2, 0, 7 (b) $-\frac{7}{3}, -2, 0, 0.7, 7, \frac{32}{3}$

(c) All are real except $\sqrt{-8}$. (d) All are complex numbers.


[1.1] 7. $\left\{\frac{4}{5}\right\}$ [1.7] 8. $\left\{\frac{11}{10}, \frac{7}{2}\right\}$ [7.6] 9. $\left\{\frac{2}{3}\right\}$ [6.4] 10. \emptyset

[8.1, 8.2] 11. $\left\{\frac{7 \pm \sqrt{177}}{4}\right\}$ [8.3] 12. $\{\pm 1, \pm 2\}$


[1.5] 13. $[1, \infty)$ [1.7] 14. $\left[2, \frac{8}{3}\right]$ [8.7] 15. (1, 3) 16. (-2, 1)



function;
domain: $(-\infty, \infty)$;
range: $(-\infty, \infty)$;
 $f(x) = \frac{4}{5}x - 3$



not a function



function;
domain: $(-\infty, \infty)$;
range: $(-\infty, 3]$;
 $f(x) = -2(x - 1)^2 + 3$

[2.2] 20. $m = \frac{2}{7}$; x-intercept: $(-8, 0)$; y-intercept: $(0, \frac{16}{7})$

[2.3] 21. $y = -\frac{5}{2}x + 2$ 22. $y = \frac{2}{5}x + \frac{13}{5}$ [3.1] 23. $\{(1, -2)\}$

24. \emptyset [3.2] 25. $\{(3, -4, 2)\}$ [3.3] 26. *Star Wars: The Last Jedi*: \$545 million; *Beauty and the Beast*: \$504 million

[4.1] 27. $\frac{x^8}{y^4}$ 28. $\frac{4}{xy^2}$ [4.5] 29. $\frac{4}{9}t^2 + 12t + 81$

[4.6] 30. $4x^2 - 6x + 11 + \frac{4}{x+2}$ [5.1-5.3] 31. $(4m - 3)(6m + 5)$

32. $4(t+5)(t-5)$ 33. $(2x+3y)(4x^2-6xy+9y^2)$ 34. $(3x-5y)^2$

[6.1] 35. $-\frac{5}{18}$ [6.2] 36. $-\frac{8}{x}$ [6.3] 37. $\frac{r-s}{r}$

[8.4] 38. southbound car: 57 mi; eastbound car: 76 mi

[7.3] 39. $\frac{3\sqrt[3]{4}}{4}$ [7.5] 40. $\sqrt{7} + \sqrt{5}$

[8.5, 8.6] 41.  [8.7] 42. $(-6, 2)$

vertex: $(-2, -6)$;
axis: $x = -2$;
domain: $(-\infty, \infty)$;
range: $[-6, \infty)$

$f(x) = 3x^2 + 12x + 6$

21. $g^{-1}(x) = \frac{-x+3}{4}$, or $g^{-1}(x) = -\frac{1}{4}x + \frac{3}{4}$ 23. not one-to-one





25. $f^{-1}(x) = x^2 + 3, x \geq 0$ 27. $f^{-1}(x) = x^2 - 6, x \geq 0$

29. not one-to-one 31. $g^{-1}(x) = \sqrt[3]{x} - 1$ 33. $f^{-1}(x) = \sqrt[3]{x+4}$

35. $f^{-1}(x) = \frac{-2x+4}{x-1}, x \neq 1$ 37. $f^{-1}(x) = \frac{-5x-2}{x-4}, x \neq 4$


39. $f^{-1}(x) = \frac{5x+1}{2x+2}, x \neq -1$ 41. (a) 8 (b) 3 43. (a) 1 (b) 0

45. (a) one-to-one 47. (a) not one-to-one 49. (a) one-to-one




55.

| x | f(x) |
|---|------|
| 0 | 0 |
| 1 | 1 |
| 4 | 2 |



57.

| x | f(x) |
|----|------|
| -1 | -3 |
| 0 | -2 |
| 1 | -1 |
| 2 | 6 |






59. $f^{-1}(x) = \frac{x+5}{4}$, or
 $f^{-1}(x) = \frac{1}{4}x + \frac{5}{4}$


60. MY CALCULATOR IS THE GREATEST THING SINCE SLICED BREAD. 61. If the function were not one-to-one, there would be ambiguity in some of the characters, as they could represent more than one letter. 62. Answers will vary. For example, Jane Doe is 1004 5 2748 129 68 3379 129.

Section 9.2

1. rises 3. C 5. A 7. 3.732 9. 0.344 11. 1.995 13. 1.587
15. 0.192 17. 73.517



$g(x) = (\frac{1}{3})^x$



27. The division in the second step does not lead to x on the left side. By expressing each side using the same base, $2^x = 2^5$, we obtain the correct solution set, $\{5\}$. 29. $\{2\}$ 31. $\{\frac{3}{2}\}$ 33. $\{\frac{3}{2}\}$

35. $\{-1\}$ 37. $\{7\}$ 39. $\{-3\}$ 41. $\{-\frac{3}{2}\}$ 43. $\{-1\}$

45. $\{-3\}$ 47. $\{-2\}$ 49. 100 g 51. 0.30 g 53. (a) 0.6°C

(b) 0.3°C 55. (a) 1.4°C (b) 0.5°C 57. (a) 80.599 million users
(b) 165.912 million users (c) 341.530 million users

9 INVERSE, EXPONENTIAL, AND LOGARITHMIC FUNCTIONS

Section 9.1


1. B 3. A 5. It is not one-to-one. The Big Breakfast with Egg Whites and the Bacon Clubhouse Burger are paired with the same fat content, 41 g. 7. Yes, it is one-to-one. Adding 1 to 1058 would make two distances be the same, so the function would not be one-to-one.

9. $\{(6, 3), (10, 2), (12, 5)\}$ 11. not one-to-one

13. $\{(4.5, 0), (8.6, 2), (12.7, 4)\}$ 15. $f^{-1}(x) = x - 3$




17. $f^{-1}(x) = -2x - 4$ 19. $f^{-1}(x) = \frac{x-4}{2}$, or $f^{-1}(x) = \frac{1}{2}x - 2$

59. (a) \$5000 (b) \$2973 (c) \$1768



Section 9.3


1. (a) B (b) E (c) D (d) F (e) A (f) C 3. $(0, \infty); (-\infty, \infty)$
 5. $\log_4 1024 = 5$ 7. $\log_{1/2} 8 = -3$ 9. $\log_{10} 0.001 = -3$
 11. $\log_{625} 5 = \frac{1}{4}$ 13. $\log_8 \frac{1}{4} = -\frac{2}{3}$ 15. $\log_5 1 = 0$ 17. $4^3 = 64$
 19. $12^1 = 12$ 21. $6^0 = 1$ 23. $9^{1/2} = 3$ 25. $(\frac{1}{4})^{1/2} = \frac{1}{2}$
 27. $5^{-1} = 5^{-1}$ 29. (a) C (b) B (c) B (d) C 31. 3.1699
 33. 1.7959 35. -1.7925 37. -1.5850 39. 1.9243 41. 1.6990
 43. $\{\frac{1}{3}\}$ 45. $\{\frac{1}{125}\}$ 47. $\{81\}$ 49. $\{\frac{1}{5}\}$ 51. $\{1\}$
 53. $\{x | x > 0, x \neq 1\}$ 55. $\{5\}$ 57. $\{\frac{5}{3}\}$ 59. $\{4\}$ 61. $\{\frac{3}{2}\}$
 63. $\{30\}$ 65. $\{\frac{37}{9}\}$ 67. 1 69. 0 71. 9 73. -1 75. 9
 77. 5 79. 6 81. 4 83. -1 85. $\frac{1}{3}$



93. 95. 8 97. 24 99. Because every real number power of 1 equals 1, if $y = \log_1 x$, then $x = 1^y$ and so $x = 1$ for every y . This contradicts the definition of a function.

101. $x = \log_a 1$ is equivalent to $a^x = 1$. The only value of x that makes $a^x = 1$ is 0. (Recall that $a \neq 1$.)


103. (a) 130 thousand units (b) 190 thousand units




105. 5 advertisements

107. $\frac{1}{100}, \frac{1}{10}, 1; 10; 100$

108. -2; -1; 0; 1; 2



domain: $(-\infty, \infty)$;
range: $(0, \infty)$



domain: $(0, \infty)$;
range: $(-\infty, \infty)$

109. The graphs have symmetry across the line $y = x$.

110. They are inverses.

Section 9.4

1. false; $\log_b x + \log_b y = \log_b xy$ 3. true 5. true 7. In the notation $\log_a (x + y)$, the parentheses do not indicate multiplication. They indicate that $x + y$ is the result of raising a to some power.
 9. $\log_{10} 7 + \log_{10} 8$ 11. 4 13. 9 15. $\log_7 4 + \log_7 5$
 17. $\log_5 8 - \log_5 3$ 19. $2 \log_4 6$ 21. $\frac{1}{3} \log_3 4 - 2 \log_3 x - \log_3 y$
 23. $\frac{1}{2} \log_3 x + \frac{1}{2} \log_3 y - \frac{1}{2} \log_3 5$ 25. $\frac{1}{3} \log_2 x + \frac{1}{5} \log_2 y - 2 \log_2 r$
 27. $\log_b xy$ 29. $\log_a \frac{m}{n}$ 31. $\log_a \frac{r^3}{s}$ 33. $\log_a \frac{125}{81}$
 35. $\log_{10} (x^2 - 9)$ 37. $\log_{10} (x^2 + 8x + 15)$ 39. $\log_p \frac{x^3 y^{1/2}}{z^{3/2} a^3}$
 41. 1.2552 43. -0.6532 45. 1.5562 47. 0.2386 49. 0.4771
 51. 4.7710 53. false 55. true 57. true 59. false

Section 9.5

1. C 3. B 5. 9.6421 7. $\sqrt{3}$ 9. -11.4007 11. 1.6335
 13. 2.5164 15. -1.4868 17. 9.6776 19. 2.0592 21. -2.8896
 23. 5.9613 25. 4.1506 27. 2.3026 29. (a) 2.552424846
 (b) 1.552424846 (c) 0.552424846 (d) The whole number parts will vary, but the decimal parts are the same. 31. poor fen 33. bog
 35. rich fen 37. 11.6 39. 8.4 41. 4.3 43. 4.0×10^{-8}
 45. 1.0×10^{-2} 47. 2.5×10^{-5} 49. (a) 142 dB (b) 126 dB
 (c) 120 dB 51. (a) 800 yr (b) 5200 yr (c) 11,500 yr
 53. (a) 189.53 million monthly active Twitter users (b) 2015
 55. (a) \$54 per ton (b) If $x = 0$, then $\ln(1 - x) = \ln 1 = 0$, so $T(x)$ would be negative. If $x = 1$, then $\ln(1 - x) = \ln 0$, but the domain of $\ln x$ is $(0, \infty)$. 57. 2.2619 59. 0.6826 61. -0.0947
 63. -2.3219 65. 0.3155 67. 0.8736



Section 9.6

1. common logarithms 3. natural logarithms 5. $\{0.827\}$
 7. $\{0.833\}$ 9. $\{1.201\}$ 11. $\{2.269\}$ 13. $\{15.967\}$
 15. $\{-6.067\}$ 17. $\{261.291\}$ 19. $\{-10.718\}$ 21. $\{3\}$
 23. $\{5.879\}$ 25. $\{-\pi\}$, or $\{-3.142\}$ 27. $\{1\}$ 29. $\{4\}$
 31. $\{\frac{2}{3}\}$ 33. $\{\frac{33}{2}\}$ 35. $\{-1 + \sqrt[3]{49}\}$ 37. $\{\pm 3\}$
 39. 2 cannot be a solution because $\log(2 - 3) = \log(-1)$, and -1 is not in the domain of $\log x$. 41. $\{\frac{1}{3}\}$ 43. $\{2\}$ 45. \emptyset
 47. $\{8\}$ 49. $\{\frac{4}{3}\}$ 51. $\{8\}$ 53. (a) \$2539.47 (b) 10.2 yr
 55. \$4934.71 57. 27.73 yr 59. (a) \$11,260.96 (b) \$11,416.64
 (c) \$11,497.99 (d) \$11,580.90 (e) \$11,581.83 61. \$137.41
 63. \$210 billion 65. (a) 1.62 g (b) 1.18 g (c) 0.69 g (d) 2.00 g
 67. (a) 179.73 g (b) 21.66 yr 69. $\log 5^x = \log 125$
 70. $x \log 5 = \log 125$ 71. $x = \frac{\log 125}{\log 5}$ 72. $\frac{\log 125}{\log 5} = 3; \{3\}$



Chapter 9 Review Exercises

1. not one-to-one 2. one-to-one 3. not one-to-one
 4. $\{(-8, -2), (-1, -1), (0, 0), (1, 1), (8, 2)\}$ 5. $f^{-1}(x) = \frac{x-7}{-3}$, or $f^{-1}(x) = -\frac{1}{3}x + \frac{7}{3}$ 6. $f^{-1}(x) = \frac{x^3+4}{6}$ 7. not one-to-one

8. This function is not one-to-one because two states in the list have minimum wage \$10.50.



17. $\{\frac{1}{2}\}$ 18. $\{4\}$ 19. $\{\frac{3}{7}\}$ 20. (a) 55.8 million people
 (b) 81.0 million people 21. (a) $5^4 = 625$ (b) $\log_5 0.04 = -2$
 22. 4; 3; fourth; 81 23. 2.5850 24. 0.5646 25. 1.7404



40. $2 \log_2 p + \log_2 r - \frac{1}{2} \log_2 z$ 41. $\log_a \frac{49}{16}$ 42. $\log_a 250$
 43. $\log_b \frac{3x}{y^2}$ 44. $\log_3 \frac{x+7}{4x+6}$ 45. 1.4609 46. -0.5901 47. 3.3638
 48. -1.3587 49. 0.9251 50. 1.7925 51. -2.0437 52. 0.3028
 53. 6.4 54. 8.4 55. 2.5×10^{-5} 56. Magnitude 1 is about 6.3 times as intense as magnitude 3. 57. every 2 hr 58. $\{2.042\}$
 59. $\{4.907\}$ 60. $\{18.310\}$ 61. $\{\frac{1}{9}\}$ 62. $\{-6 + \sqrt[3]{25}\}$
 63. $\{2\}$ 64. $\{\frac{3}{8}\}$ 65. $\{4\}$ 66. $\{1\}$ 67. \$24,403.80
 68. \$11,190.72 69. Plan A is better because it would pay \$2.92 more.
 70. 13.9 days


Chapter 9 Mixed Review Exercises

1. 7 2. 0 3. -3 4. 36 5. 4 6. e 7. -5 8. 5.4
 9. $\{72\}$ 10. $\{5\}$ 11. $\{\frac{1}{9}\}$ 12. $\{\frac{4}{3}\}$ 13. $\{3\}$ 14. $\{0\}$
 15. $\{\frac{1}{8}\}$ 16. $\{\frac{11}{3}\}$ 17. $\{\frac{2}{63}\}$ 18. $\{-2, -1\}$ 19. (a) \$4267 (b) 11%

20. (a) 0.325 (b) 0.673 21. (a) 1.209061955
 (b) 7 (c) $\{1.209061955\}$ 22. Answers will vary. Suppose the name is Jeffery Cole, with $m = 7$ and $n = 4$. (a) $\log_7 4$ is the exponent to which 7 must be raised to obtain 4.
 (b) 0.7124143742 (c) 4

Chapter 9 Test

- [9.1] 1. (a) not one-to-one (b) one-to-one 2. $f^{-1}(x) = x^3 - 7$



6. Interchange the x - and y -values of the ordered pairs, because the functions are inverses. 7. 9 8. 6 9. 0 [9.2] 10. $\{-4\}$ 11. $\{-\frac{13}{3}\}$
 12. (a) 775 millibars (b) 265 millibars [9.3] 13. $\log_4 0.0625 = -2$
 14. $7^2 = 49$ 15. $\{32\}$ 16. $\{\frac{1}{2}\}$ 17. $\{2\}$ 18. 5; 2; fifth; 32

- [9.4] 19. $2 \log_3 x + \log_3 y$ 20. $\frac{1}{2} \log_5 x - \log_5 y - \log_5 z$

21. $\log_b \frac{s^3}{t}$ 22. $\log_b \frac{r^{1/4} s^2}{t^{2/3}}$ [9.5] 23. 1.3636 24. -0.1985

25. 2.1245 26. (a) $\frac{\log 19}{\log 3}$ (b) $\frac{\ln 19}{\ln 3}$ (c) 2.6801


- [9.6] 27. $\{3.966\}$ 28. $\{3\}$ 29. (a) \$11,903.40 (b) 19.9 yr
 30. (a) \$17,427.51 (b) 23.1 yr

Chapters R-9 Cumulative Review Exercises

- [R.1] 1. (a) 0.05; 5% (b) 1.25; 125% [R.1, R.3] 2. (a) $-\frac{6}{5}$
 (b) $-\frac{4}{5}$ 3. (a) 3750 (b) 0.0375 [R.2] 4. (a) -2, 0, 6, $\frac{30}{3}$ (or 10)
 (b) $-\frac{9}{4}$, -2, 0, 0.6, 6, $\frac{30}{3}$ (or 10) (c) $-\sqrt{2}$, $\sqrt{11}$

- [R.3, R.4] 5. 16 6. -39 [I.1] 7. $\{-\frac{2}{3}\}$ [I.5] 8. $[1, \infty)$

- [I.7] 9. $\{-2, 7\}$ 10. $(-\infty, -3) \cup (2, \infty)$



- [2.2, 2.5] 13. (a) yes (b) 1.955; The number of travelers shows an increase of an average of 1.955 million per year during these years.

- [2.3] 14. $y = \frac{3}{4}x - \frac{19}{4}$ [3.1, 3.2] 15. $\{(4, 2)\}$

16. $\{(x, y) | 3x - 2y = 3\}$ 17. $\{(1, -1, 4)\}$ [3.3] 18. 6 lb

- [4.5] 19. $6p^2 + 7p - 3$ 20. $16k^2 - 24k + 9$

- [4.3] 21. $-5m^3 + 2m^2 - 7m + 4$ [4.6] 22. $5x^2 - 2x + 8$

- [5.1] 23. $x(8 + x^2)$ [5.2] 24. $(3y - 2)(8y + 3)$ 25. $z(5z + 1)(z - 4)$

[5.3] 26. $(4a + 5b^2)(4a - 5b^2)$ 27. $(2c + d)(4c^2 - 2cd + d^2)$
 28. $(4r + 7q)^2$ [4.1] 29. $-\frac{1875p^{13}}{8}$ [6.1] 30. $\frac{x + 5}{x + 4}$



[6.2] 31. $\frac{-3k - 19}{(k + 3)(k - 2)}$ [7.3] 32. $12\sqrt{2}$ [7.4] 33. $-27\sqrt{2}$

[7.7] 34. 41 [7.6] 35. $\{0, 4\}$ [8.1, 8.2] 36. $\left\{\frac{1 \pm \sqrt{13}}{6}\right\}$


[8.7] 37. $(-\infty, -4) \cup (2, \infty)$ [8.3] 38. $\{\pm 1, \pm 2\}$

[9.2] 39. $\{-1\}$ [9.6] 40. $\{1\}$

[8.5] 41. $f(x) = \frac{1}{3}(x - 1)^2 + 2$ [9.2] 42.



[9.3] 43. $f(x) = \log_3 x$ [9.4] 44. $3 \log x + \frac{1}{2} \log y - \log z$




- [9.6] 45. (a) 25,000 bacteria
 (b) 30,500 bacteria (c) 37,300 bacteria
 (d) in 3.5 hr, or at 3:30 P.M.

10 NONLINEAR FUNCTIONS, CONIC SECTIONS, AND NONLINEAR SYSTEMS


Section 10.1

1. E; 0; 0 3. A; $(-\infty, \infty)$; $\{\dots, -2, -1, 0, 1, 2, \dots\}$ 5. B; It does not satisfy the conditions of the vertical line test. 7. B 9. A


11. Shift the graph of $g(x) = \frac{1}{x}$ to the right 3 units and up 2 units.




domain: $(-\infty, \infty)$;
 range: $[0, \infty)$




domain: $(-\infty, 0) \cup (0, \infty)$;
 range: $(-\infty, 1) \cup (1, \infty)$




domain: $[2, \infty)$;
 range: $[0, \infty)$



domain: $(-\infty, 2) \cup (2, \infty)$;
 range: $(-\infty, 0) \cup (0, \infty)$






domain: $[-3, \infty)$;
 range: $[-3, \infty)$



domain: $(-\infty, \infty)$;
 range: $[1, \infty)$

25. $\llbracket x \rrbracket$ means the greatest integer less than or equal to x . Because -6 is less than -5.1 , the greatest integer less than -5.1 is -6 . 27. 3

29. 4 31. 0 33. 2 35. -14 37. -11




45. \$2.75

Section 10.2


1. conic sections; parabolas 3. ellipse; foci

5. (a) $(0, 0)$ (b) 5 (c)




7. B 9. C


11. $(0, 0)$; $r = 2$




13. $(0, 0)$; $r = 9$




15. $(5, -4)$; $r = 7$




17. $(-1, -3)$; $r = 5$




19. $x^2 + y^2 = 16$




21. $(x + 3)^2 + (y - 2)^2 = 9$




23. $(x - 4)^2 + (y - 3)^2 = 25$



25. $(x + 2)^2 + y^2 = 5$



27. $x^2 + (y + 3)^2 = 49$




29. $(x + 2)^2 + (y + 3)^2 = 4$; $(-2, -3)$; $r = 2$


31. $(x + 5)^2 + (y - 7)^2 = 81$; $(-5, 7)$; $r = 9$

33. $(x - 2)^2 + (y - 4)^2 = 16$; $(2, 4)$; $r = 4$

35. $(x - 2)^2 + (y - 3)^2 = 4$;
 $(2, 3)$; $r = 2$









37. $(x + 3)^2 + (y - 3)^2 = 9$;
 $(-3, 3)$; $r = 3$



39. one point; The only ordered pair that satisfies the equation is $(4, 1)$.

41. The thumbtack acts as the center, and the length of the string acts as the radius.






















55. $3\sqrt{3}$ units 57. (a) 10 m (b) 36 m 59. (a) 348.2 ft
(b) 1787.6 ft 61. (a) 154.7 million mi (b) 128.7 million mi

Section 10.3

1. C 3. D 5. Because the coefficient of the x^2 -term is positive, this is a horizontal hyperbola with x -intercepts $(-3, 0)$ and $(3, 0)$.







33. (a) 50 m
(b) 69.3 m
35. $y = \pm x$



Section 10.4













1. one 3. none









13. $\{(0, 0), (\frac{1}{2}, \frac{1}{2})\}$ 15. $\{(-6, 9), (-1, 4)\}$
17. $\{(-\frac{1}{5}, \frac{7}{5}), (1, -1)\}$ 19. $\{(-2, -2), (-\frac{4}{3}, -3)\}$
21. $\{(-3, 1), (1, -3)\}$ 23. $\{(-\frac{3}{2}, -\frac{9}{4}), (-2, 0)\}$
25. $\{(-\sqrt{3}, 0), (\sqrt{3}, 0), (-\sqrt{5}, 2), (\sqrt{5}, 2)\}$
27. $\{(\frac{\sqrt{3}}{3}i, -\frac{1}{2} + \frac{\sqrt{3}}{6}i), (-\frac{\sqrt{3}}{3}i, -\frac{1}{2} - \frac{\sqrt{3}}{6}i)\}$
29. $\{(-2, 0), (2, 0)\}$ 31. $\{(1, 3), (1, -3), (-1, 3), (-1, -3)\}$
33. $\{(-\sqrt{3}, 0), (\sqrt{3}, 0)\}$
35. $\{(-2\sqrt{3}, -2), (-2\sqrt{3}, 2), (2\sqrt{3}, -2), (2\sqrt{3}, 2)\}$
37. $\{(i\sqrt{2}, -3i\sqrt{2}), (-i\sqrt{2}, 3i\sqrt{2}), (-\sqrt{6}, -\sqrt{6}), (\sqrt{6}, \sqrt{6})\}$
39. $\{(-2i\sqrt{2}, -2\sqrt{3}), (-2i\sqrt{2}, 2\sqrt{3}), (2i\sqrt{2}, -2\sqrt{3}), (2i\sqrt{2}, 2\sqrt{3})\}$
41. $\{(-\sqrt{5}, -\sqrt{5}), (\sqrt{5}, \sqrt{5})\}$
43. $\{(-3, -1), (3, 1), (-i, 3i), (i, -3i)\}$
45. $\{(i, 2i), (-i, -2i), (2, -1), (-2, 1)\}$ 47. length: 12 ft; width: 7 ft
49. $\$20; \frac{4}{5}$ thousand or 800 calculators

Section 10.5

1. (a) B (b) D (c) A (d) C 3. $x \geq 0$ 5. $x < 0$ 7. C
 $y \geq 0$ $y \leq 0$










35. $\{(6, -9), (-2, -5)\}$ 36. $\{(1, 2), (-5, 14)\}$




37. $\{(4, 2), (-1, -3)\}$ 38. $\{(-2, -4), (8, 1)\}$

39. $\{(-\sqrt{2}, 2), (-\sqrt{2}, -2), (\sqrt{2}, -2), (\sqrt{2}, 2)\}$


40. $\{(-\sqrt{6}, -\sqrt{3}), (-\sqrt{6}, \sqrt{3}), (\sqrt{6}, -\sqrt{3}), (\sqrt{6}, \sqrt{3})\}$




44. D




Chapter 10 Review Exercises




domain: $(-\infty, \infty)$;
range: $[0, \infty)$



domain: $[0, \infty)$;
range: $[3, \infty)$









domain: $(-\infty, 4) \cup (4, \infty)$;
range: $(-\infty, 0) \cup (0, \infty)$



domain: $(-\infty, \infty)$;
range: $\{\dots, -2, -1, 0, 1, 2, \dots\}$

5. 12 6. 2 7. -21 8. -5 9. $x^2 + y^2 = 49$
10. $(x + 2)^2 + (y - 4)^2 = 9$ 11. $(x - 4)^2 + (y - 2)^2 = 36$
12. $(x + 5)^2 + y^2 = 2$ 13. $(x + 3)^2 + (y - 2)^2 = 16$; $(-3, 2)$; $r = 4$
14. $(x - 4)^2 + (y - 1)^2 = 4$; $(4, 1)$; $r = 2$
15. $(x + 1)^2 + (y + 5)^2 = 9$; $(-1, -5)$; $r = 3$
16. $(x - 3)^2 + (y + 2)^2 = 25$; $(3, -2)$; $r = 5$
17. (a) circle (b) ellipse (c) ellipse (d) circle
18. (a) A (b) C (c) D (d) B



25. $f(x) = -\sqrt{16 - x^2}$
26. (a) C (b) A (c) D (d) B
27. circle 28. parabola 29. hyperbola
30. ellipse 31. parabola 32. hyperbola
33. 0, 1, or 2 34. 0, 1, 2, 3, or 4

48. (a) B (b) D (c) C (d) A


Chapter 10 Mixed Review Exercises

1. $(-2, 5)$; $r = 6$ 2. true
3. $x^2 + 9y^2 = 9$
4. $x^2 - 9y^2 = 9$
5. $f(x) = \sqrt{4 - x}$
6. $x^2 + y^2 = 25$
7. $\frac{y^2}{4} - 1 \leq \frac{x^2}{9}$
8. $4y > 3x - 12$
 $x^2 < 16 - y^2$
9. $\{(-7, -35), (12, 60)\}$ 10. $\{(-1, -2), (1, 2), (-2i, i), (2i, -i)\}$


Chapter 10 Test

- [10.1] 1. 0 2. $[0, \infty)$ 3. $\{\dots, -2, -1, 0, 1, 2, \dots\}$

4. (a) C (b) A (c) D (d) B



domain: $(-\infty, \infty)$;
range: $[4, \infty)$








7. D; $(0, 0)$; $r = 1$ 8. $(-4, 1)$; $r = 5$

[10.3]


[10.2]

[10.3]





[10.5]



Chapters R-10 Cumulative Review Exercises

[R.1] 1. 0.01; 1% [R.3] 2. 0 [R.2] 3. (a) A, B, C, D, F
 (b) B, C, D, F (c) D, F (d) E, F (e) C, D, F (f) D, F

[R.4] 4. (a) 36 (b) -36 (c) 36 (d) 6 (e) -6

(f) not a real number [1.1] 5. $\left\{ \frac{2}{3} \right\}$ [1.5] 6. $\left(-\infty, \frac{3}{5} \right)$

[1.7] 7. $\{-4, 4\}$ 8. $(-\infty, -5) \cup (10, \infty)$ [2.2] 9. $\frac{2}{3}$

[2.3] 10. $3x + 2y = -13$ [3.1] 11. $\{(3, -3)\}$ [3.2] 12. $\{(4, 1, -2)\}$

[10.4] 13. $\left\{ (-1, 5), \left(\frac{5}{2}, -2\right) \right\}$ [3.3] 14. 40 mph

[4.5] 15. $25y^2 - 30y + 9$ [4.6] 16. $4x^3 - 4x^2 + 3x + 5 + \frac{3}{2x + 1}$

[6.1] 17. $\frac{y - 1}{y(y - 3)}$ [6.2] 18. $\frac{3c + 5}{(c + 5)(c + 3)}$ 19. $\frac{1}{p}$

[6.3] 20. $\frac{xy}{y - x}$ [7.4] 21. $2\sqrt[3]{2}$ [7.7] 22. $\frac{7}{5} + \frac{11}{5}i$

[5.2] 23. $(3x + 2)(4x - 5)$ [5.3] 24. $(z^2 + 1)(z + 1)(z - 1)$

25. $(a - 3b)(a^2 + 3ab + 9b^2)$ [6.5] 26. $1\frac{1}{5}$ hr [4.1] 27. $\frac{a^5}{4}$

[7.5] 28. $\frac{3\sqrt{10}}{2}$ [7.6] 29. \emptyset [5.5] 30. $\left\{ \frac{1}{5}, -\frac{3}{2} \right\}$

[8.1, 8.2] 31. $\left\{ \frac{3 \pm \sqrt{33}}{6} \right\}$ [8.3] 32. $\left\{ \pm \frac{\sqrt{6}}{2}, \pm \sqrt{7} \right\}$


[9.6] 33. $\{3\}$ [8.4] 34. $v = \frac{\pm \sqrt{rFkw}}{kw}$ [4.4] 35. (a) -1

(b) $9x^2 + 18x + 4$ [9.1] 36. $f^{-1}(x) = \sqrt[3]{x - 4}$


[9.4, 9.5] 37. (a) 4 (b) 7 [9.4] 38. $\log \frac{(3x + 7)^2}{4}$

[10.1] 39. domain: $(-\infty, \infty)$; range: $[0, \infty)$


[2.6]




[8.5]




[10.5]




[10.1]



[10.3]



[9.2]



11 FURTHER TOPICS IN ALGEBRA

Section 11.1

1. the set of all positive integers (natural numbers) 3. 80
 5. 12 7. 2, 3, 4, 5, 6 9. $4, \frac{5}{2}, 2, \frac{7}{4}, \frac{8}{5}$ 11. 3, 9, 27, 81, 243
 13. $-1, -\frac{1}{4}, -\frac{1}{9}, -\frac{1}{16}, -\frac{1}{25}$ 15. 5, -5, 5, -5, 5 17. $0, \frac{3}{2}, \frac{8}{3}, \frac{15}{4}, \frac{24}{5}$
 19. -70 21. $\frac{49}{23}$ 23. 171 25. $4n$ 27. $-8n$ 29. $\frac{1}{3^n}$ 31. $\frac{n + 1}{n + 4}$
 33. \$110, \$109, \$108, \$107, \$106, \$105; \$400 35. \$6554
 37. $4 + 5 + 6 + 7 + 8 = 30$ 39. $3 + 6 + 11 = 20$
 41. $-2 + 2 - 2 + 2 - 2 + 2 = 0$ 43. $0 + 6 + 14 + 24 + 36 = 80$

Answers may vary for Exercises 45-49.

45. $\sum_{i=1}^5 (i + 2)$ 47. $\sum_{i=1}^5 2^i(-1)^i$ 49. $\sum_{i=1}^4 i^2$

51. In summation notation, each replacement for i should be squared first before the negative symbol is applied. The correct answer is $\sum_{i=1}^5 -i^2$.

53. 9 55. $\frac{40}{9}$ 57. 9200 mutual funds

Section 11.2

1. difference 3. 55 5. $\frac{5}{2}; 3; \frac{7}{2}$ 7. 16; 26 9. $d = 1$
 11. not arithmetic 13. $d = -5$ 15. 5, 9, 13, 17, 21
 17. -2, -6, -10, -14, -18 19. $\frac{1}{3}, 1, \frac{5}{3}, \frac{7}{3}, 3$ 21. $a_n = 5n - 3; 122$
 23. $a_n = \frac{3}{4}n + \frac{9}{4}; 21$ 25. $a_n = 3n - 6; 69$ 27. 76 29. 48
 31. -1 33. 16 35. 26 37. 6 39. The student subtracted terms in the wrong order. Subtracting the first term from the second term, we find that $d = -10 - (-15) = -10 + 15 = 5$. 41. 15 43. 87 45. $\frac{63}{2}$
 47. 81 49. -3 51. -2.1 53. 390 55. 395 57. 31,375
 59. \$465 61. \$2100 per month 63. 68 seats; 1100 seats
 65. no; 3 blocks; 9 rows

Section 11.3

1. ratio 3. 31 5. $-\frac{1}{81}; \frac{1}{243}; -\frac{1}{729}$ 7. 14; 56 9. $r = 2$
 11. not geometric 13. $r = -3$ 15. $r = -\frac{1}{2}$

There are alternative forms of the answers in Exercises 17-21.

17. $a_n = -5(2)^{n-1}$ 19. $a_n = -2\left(-\frac{1}{3}\right)^{n-1}$ 21. $a_n = 10\left(-\frac{1}{5}\right)^{n-1}$
 23. 3,906,250 25. $\frac{1}{354,294}$ 27. -4374 29. 2, 6, 18, 54, 162

31. $5, -1, \frac{1}{5}, -\frac{1}{25}, \frac{1}{125}$ 33. $-4, -2, -1, -0.5, -0.25$ 35. $-1,048,575$
 37. $\frac{121}{243}$ 39. -1.997 41. 2.662 43. -2.982 45. $\$24,470.52$
 47. $\$74,156.06$ 49. 9 51. $\frac{10,000}{11}$ 53. $-\frac{9}{20}$
 55. The sum does not exist. 57. 1.3 ft 59. 140 m 61. 3 days; $\frac{1}{4}$ g
 63. (a) 1.5 billion units (b) 12 yr 65. $\$5005.65$
 67. $\frac{a_1}{1-r} = \frac{0.9}{1-0.1} = \frac{0.9}{0.9} = 1$; Therefore, $0.99999 \dots = 1$.

Section 11.4

1. 1; sum 3. x^3 5. 6 7. 1 9. 720 11. $40,320$
 13. 15 15. 1 17. 120 19. 15 21. 78 23. 120
 25. $m^4 + 4m^3n + 6m^2n^2 + 4mn^3 + n^4$
 27. $a^5 - 5a^4b + 10a^3b^2 - 10a^2b^3 + 5ab^4 - b^5$
 29. $8x^3 + 36x^2 + 54x + 27$ 31. $\frac{x^4}{16} - \frac{x^3y}{2} + \frac{3x^2y^2}{2} - 2xy^3 + y^4$
 33. $x^8 + 4x^6 + 6x^4 + 4x^2 + 1$ 35. $27x^6 - 27x^4y^2 + 9x^2y^4 - y^6$
 37. $r^{12} + 24r^{11}s + 264r^{10}s^2 + 1760r^9s^3$
 39. $3^{14}x^{14} - 14(3^{13})x^{13}y + 91(3^{12})x^{12}y^2 - 364(3^{11})x^{11}y^3$
 41. $t^{20} + 10t^{18}u^2 + 45t^{16}u^4 + 120t^{14}u^6$ 43. $120(2^7)m^7n^3$
 45. $\frac{7x^2y^6}{16}$ 47. $36k^7$ 49. $160x^6y^3$ 51. $4320x^9y^4$

Chapter 11 Review Exercises

1. $-1, 1, 3, 5$ 2. $0, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}$ 3. $1, 4, 9, 16$ 4. $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}$
 5. $0, 3, 8, 15$ 6. $1, -2, 3, -4$ 7. $1 + 4 + 9 + 16 + 25 = 55$
 8. $2 + 3 + 4 + 5 + 6 + 7 = 27$ 9. $11 + 16 + 21 + 26 = 74$
 10. $9 + 16 + 25 + 36 = 86$ 11. $2 + 4 + 8 + 16 + 32 + 64 = 126$
 12. $\frac{4}{5} + \frac{5}{6} + \frac{6}{7} + \frac{7}{8} = \frac{2827}{840}$ 13. 763.3 thousand electric vehicles
 14. arithmetic; $d = 3$ 15. arithmetic; $d = 4$ 16. geometric; $r = -\frac{1}{2}$
 17. geometric; $r = -1$ 18. neither 19. neither 20. 89 21. 73
 22. 69 23. $a_n = -5n + 1$ 24. $a_n = -3n + 9$ 25. 15 26. 22
 27. 152 28. 164 29. $a_n = -1(4)^{n-1}$ 30. $a_n = \frac{2}{3}(\frac{1}{5})^{n-1}$
 31. $118,098$ 32. 2560 or -2560 33. $\frac{341}{1024}$ 34. 0 35. 1
 36. The sum does not exist. 37. $32p^5 - 80p^4q + 80p^3q^2 - 40p^2q^3 + 10pq^4 - q^5$ 38. $x^8 + 12x^6y + 54x^4y^2 + 108x^2y^3 + 81y^4$
 39. $81t^{12} - 108t^9s^2 + 54t^6s^4 - 12t^3s^6 + s^8$ 40. $7752(3)^{16}a^{16}b^3$

Chapter 11 Mixed Review Exercises

1. $a_{10} = 1536; S_{10} = 1023$ 2. $a_{40} = 235; S_{10} = 280$
 3. $a_{15} = 38; S_{10} = 95$ 4. $a_9 = 6561; S_{10} = -14,762$ 5. $a_n = 2(4)^{n-1}$
 6. $a_n = 5n - 3$ 7. $a_n = -3n + 15$ 8. $a_n = 27(\frac{1}{3})^{n-1}$
 9. 10 sec 10. $\$21,973.00$ 11. approximately $42,000$ people





12. $\frac{1}{128}$ 13. (a) $0.5 + 0.05 + 0.005 + 0.0005 + \dots$ (b) 0.1 (c) $\frac{5}{9}$
 14. No, the sum cannot be found, because $r = 2$. This value of r does not satisfy $|r| < 1$.

Chapter 11 Test


- [11.1] 1. $0, 2, 0, 2, 0$ [11.2] 2. $4, 6, 8, 10, 12$ [11.3] 3. $48, 24, 12, 6, 3$
 [11.2] 4. 0 [11.3] 5. $\frac{64}{3}$ or $-\frac{64}{3}$ [11.2] 6. 75 [11.3] 7. 124 or 44
 [11.1] 8. 92.5 min [11.3] 9. $\$137,925.91$ 10. It has a sum if $|r| < 1$.
 [11.2] 11. 70 12. 33 13. $125,250$ [11.3] 14. 42 15. $\frac{1}{3}$
 16. The sum does not exist. [11.4] 17. $40,320$ 18. 1 19. 15
 20. 66 21. $81k^4 - 540k^3 + 1350k^2 - 1500k + 625$ 22. $\frac{14,080x^8y^4}{9}$
 [11.1] 23. $\$324$ [11.3] 24. $3,542,940$

Chapters R-11 Cumulative Review Exercises


- [R.2, R.3] 1. 8 2. $-\frac{13}{45}$ 3. $-\frac{8}{3}, 10, 0, \frac{45}{15}$ (or 3), $0.82, -3$
 4. $\sqrt{13}, -\sqrt{3}$ [1.1] 5. $\{\frac{1}{6}\}$ 6. $\{9\}$ [1.5] 7. $[10, \infty)$
 [1.6] 8. $(-\infty, -3) \cup (4, \infty)$ [1.7] 9. $\{-\frac{9}{2}, 6\}$
 10. $(-\infty, -3) \cup [8, \infty)$ [5.5] 11. $\{-\frac{5}{2}, 2\}$
 [8.2] 12. $\{\frac{-5 \pm \sqrt{217}}{12}\}$ [6.4] 13. \emptyset [9.2] 14. $\{\frac{5}{2}\}$
 [9.6] 15. $\{2\}$ [2.2] 16. $\frac{3}{4}$ [2.3] 17. $3x + y = 4$
 [7.3, 10.2] 18. $(x + 5)^2 + (y - 12)^2 = 81$
 [3.1] 19. $\{(-1, -2)\}$ [3.2] 20. $\{(2, 1, 4)\}$
 [10.4] 21. $\{(-1, 5), (\frac{5}{2}, -2)\}$ [3.3] 22. 2 lb [4.1] 23. $\frac{9}{4}$
 24. $-\frac{27p^2}{10}$ [4.5] 25. $20p^2 - 2p - 6$ [4.3] 26. $-5m^3 - 3m^2 + 3m + 8$
 [4.6] 27. $2t^3 + 3t^2 - 4t + 2 + \frac{3}{3t-2}$ [7.7] 28. 73
 [5.2] 29. $z(3z + 4)(2z - 1)$ [5.3] 30. $(7a^2 + 3b)(7a^2 - 3b)$
 31. $(c + 3d)(c^2 - 3cd + 9d^2)$ [6.1] 32. $\frac{x+7}{x-2}$
 [6.2] 33. $\frac{3p-26}{p(p+3)(p-4)}$ [7.4] 34. $10\sqrt{2}$




[9.3] 39.




[10.1] 40.



[10.2] 41.



[10.3] 42.



[9.1] 43. $f^{-1}(x) = \frac{x-5}{9}$, or $f^{-1}(x) = \frac{1}{9}x - \frac{5}{9}$

[11.1] 44. $-7, -2, 3, 8, 13$ [11.2] 45. 78 [11.3] 46. $\frac{75}{7}$

[11.1] 47. 30 [13.4] 48. (a) $362,880$ (b) 1 (c) 210

49. $32a^5 - 80a^4 + 80a^3 - 40a^2 + 10a - 1$ 50. $-\frac{45x^8y^6}{4}$

APPENDIX SYNTHETIC DIVISION

1. C 3. $3; 4$ 5. $3; 6; 0$ 7. The student neglected to include 0 as the coefficient of the linear term. The correct setup is $-2 \overline{)4 \ 2 \ 0 \ 6}$ and

the correct answer is $4x^2 - 6x + 12 + \frac{-18}{x+2}$. 9. $x - 5$

11. $4x - 1$ 13. $2x + 4 + \frac{5}{x+2}$ 15. $x - 4 + \frac{9}{x+1}$

17. $4x^2 + x + 3$ 19. $x^2 + 6x + 20 + \frac{83}{x-4}$

21. $2x^4 + 4x^3 + 6x^2 + 15x + 6 + \frac{10}{x-2}$

23. $-3x^4 + 5x^2 - 11x + 11 + \frac{-8}{x+1}$ 25. $x^4 + x^2 + 1 + \frac{2}{x+1}$

27. 7 29. -2 31. 0 33. yes 35. no 37. yes 39. no

41. $(2x - 3)(x + 4)$ 42. $\{-4, \frac{3}{2}\}$ 43. 0 44. 0 45. a

46. Yes, $x - 3$ is a factor; $g(x) = (x - 3)(3x - 1)(x + 2)$

STUDY SKILLS

p. S-1 Borchee/E+/Getty Images; **p. S-2** Origenoo stock/123RF; **p. S-4** G-stockstudio/Shutterstock; **p. S-6** Rawpixel/123RF; **p. S-8** Studio DMM Photography, Designs & Art/Shutterstock; **p. S-10** Doyeol (David) Ahn/Alamy Stock Photo

CHAPTER R

p. 17 Terry McGinnis; **p. 25** Callie Daniels; **p. 35** Piotr Adamowicz/Shutterstock; **p. 44** (top, bottom) Terry McGinnis; **p. 45** Maximilian Laschon/Shutterstock; **p. 47** Tatyana Tomsickova/123RF

CHAPTER 1

p. 55 Callie Daniels; **p. 70** John Hornsby; **p. 75** Anne Richard/Shutterstock; **p. 76** Dinodia/123RF; **p. 82** Pedro Monteiro/Shutterstock; **p. 89** (top) Michele Perbellini/Shutterstock, (bottom) Hornsby stamp; **p. 90** Rob/Fotolia; **p. 91** Spotmatik Ltd/Shutterstock, **p. 97** Pearson Education, Inc.; **p. 98** (top) Josefauer/Shutterstock, (bottom) Pete Niesen/Shutterstock; **p. 102** Oles Ishchuk/123RF; **p. 114** (top) Fredrick Kippe/Alamy Stock Photo, (bottom left) Duplass/Shutterstock, (bottom right) John Hornsby; **p. 121** Jeayesy/123RF; **p. 123** Michaeljung/Shutterstock; **p. 124** Chad McDermott/Shutterstock; **p. 131** Leonid Shecheglov/Shutterstock; **p. 135** Katherine Martin/123RF; **p. 144** John Hornsby

CHAPTER 2

p. 149 Michelangelo/Shutterstock; **p. 151** Library of Congress; **p. 182** Kazzland Inc. /Shutterstock; **p. 184** Podfoto/Shutterstock; **p. 187** (top) Siraphol/123RF, (bottom) Karen roach/123RF; **p. 188** (top) Wavebreakmedia/Shutterstock, (bottom) Goran Bogicevic/123RF; **p. 190** Barry Blackburn/Shutterstock; **p. 198** Derek Meijer/Alamy Stock Photo; **p. 216** Rido/Shutterstock; **p. 218** Neelsky/Shutterstock

CHAPTER 3

p. 229 Graham Moore/123RF; **p. 254** Pavol Kmeto/Shutterstock; **p. 255** Blickwinkel/Alamy Stock Photo; **p. 262** Marie C Fields/Shutterstock; **p. 264** AndreAnita/Shutterstock; **p. 265** (top) Chris Whitehead/Cultura Creative (RF)/Alamy Stock Photo, (bottom) Thomas M Perkins/Shutterstock; **p. 267** Callie Daniels; **p. 268** Dpa picture alliance/Alamy Stock Photo; **p. 269** Callie Daniels

CHAPTER 4

p. 279 Rawpixel/123RF; **p. 294** (top) Vovan/Shutterstock, (bottom) Terry McGinnis; **p. 295** Doug McLean/Shutterstock; **p. 303** Gareth Boden/Pearson Education Ltd; **p. 308** Gorbelabda/Shutterstock; **p. 310** Fair Trade USA; **p. 311** Topseller/Shutterstock; **p. 312** Coprid/Shutterstock; **p. 313** Terry McGinnis; **p. 335** Lucky Dragon/Fotolia; **p. 336** Lopolo/Shutterstock; **p. 340** National Atomic Museum Foundation

CHAPTER 5

p. 341 Don Fink/Shutterstock; **p. 372** Konstantin Shaklein/123RF

CHAPTER 6

p. 385 Ballemans/Shutterstock; **p. 404** Terry McGinnis; **p. 418** Lars Christensen/Shutterstock; **p. 422** Disability Images/Alamy Stock Photo; **p. 424** Lucio pepi/Fotolia; **p. 426** Dmitry Kalinovsky/Shutterstock; **p. 429** (top) John Hornsby, (center) Hurst Photo/Shutterstock, (bottom) CCImage/Shutterstock; **p. 430** (top) Terry McGinnis, (bottom) Scott Rothstein/Shutterstock; **p. 438** Dana Bartekoske/123RF; **p. 441** Dmitry Deshevykh/Alamy Stock Photo; **p. 442** Cathy Yeulet/123RF

CHAPTER 7

p. 455 Giulio Meinardi/Fotolia; **p. 463** Stoupa/Shutterstock; **p. 472** Maria Moroz/Fotolia; **p. 485** (dog) Anneka/Shutterstock, (screen) Cobalt/Fotolia; **p. 486** Kevin Kipper/Alamy Stock Photo; **p. 509** Johnson Space Center/NASA; **p. 528** Scol22/Fotolia

CHAPTER 8

p. 531 Sarot Chamnankit/123RF; **p. 534** Library of Congress; **p. 540** (left) Piotrek Jastrzebski/Shutterstock, (right) Rudi1976/Alamy Stock Photo; **p. 551** Phovoir/Shutterstock; **p. 564** Micro10x/Shutterstock; **p. 569** Look Die Bildagentur der Fotografen GmbH/Alamy Stock Photo; **p. 570** Steven J. Everts/123RF; **p. 574** Zurijeta/Shutterstock; **p. 578** Andriy Popov/123RF; **p. 579** Alexwhite/Shutterstock; **p. 588** Kaband/Shutterstock; **p. 589** Bonniej/E+/Getty Images; **p. 599** Hartrockets/iStock/Getty Images

CHAPTER 9

p. 611 Chien321/Shutterstock; **p. 613** Frontpage/Shutterstock; **p. 620** Mark Higgins/Shutterstock;
p. 629 Bernhard Staehli/Shutterstock; **p. 630** Maksym Bondarchuk/123RF; **p. 635** Per Tillmann/Fotolia;
p. 638 Prisma by Dukas Presseagentur GmbH/Alamy Stock Photo; **p. 647** Romrodphoto/Shutterstock;
p. 649 Georgios Kollidas/Fotolia; **p. 652** (top) Jim West/Alamy Stock Photo, (bottom) Silvano Rebai/
Fotolia; **p. 653** WaterFrame/Alamy Stock Photo; **p. 661** Sergey Nivens/Shutterstock; **p. 662** Armadillo
Stock/Shutterstock; **p. 668** Monkey Business Images/Shutterstock; **p. 670** Syda Productions/
Shutterstock; **p. 671** IndustryAndTravel/Alamy Stock Photo

CHAPTER 10

p. 677 Terry McGinnis; **p. 681** Samuel Acosta/Shutterstock; **p. 694** Petr Student/Shutterstock;
p. 701 Faiz Azizan/Shutterstock

CHAPTER 11

p. 725 Callie Daniels; **p. 726** Madeinitaly4k/Shutterstock; **p. 730** Frank11/Shutterstock;
p. 739 Blend Images/Shutterstock; **p. 740** Creativa Images/Shutterstock; **p. 748** Paul Hakimata
Photography/Shutterstock; **p. 751** Atlaspix/Alamy Stock Photo; **p. 761** Jtbaskinphoto/Shutterstock;
p. 763 Baloncici/123RF

- A**
- Absolute value
 - definition of, 19
 - distance definition of, 125–126
 - explanation of, 125
 - simplifying square roots using, 459
 - solving equations involving, 125–131
 - solving inequalities involving, 125–131
 - Absolute value equations, 125–131
 - Absolute value functions, 678
 - Absolute value inequalities, 125–131
 - Addition
 - associative property of, 48–50
 - commutative property of, 48–50
 - of complex numbers, 513
 - of decimals, 6
 - of fractions, 4–5
 - identity element for, 46
 - identity property for, 46
 - inverse property for, 47–48
 - of polynomial functions, 304–305
 - of polynomials, 298–299
 - of radical expressions, 487–489
 - of rational expressions, 396–401
 - of real numbers, 26–27
 - Addition property
 - of equality, 57
 - of inequality, 105–106
 - Additive identity, 46–47
 - Additive inverse of a real number, 18–19, 47–48
 - Agreement on domain, 205
 - Algebraic expressions
 - definition of, 41, 296
 - evaluating, 41
 - explanation of, 56
 - Algebraic fraction, 386
 - Angles
 - complementary, 100
 - steps to solve measures of, 95–96
 - supplementary, 100, 265
 - vertical, 100, 265
 - Annuity
 - definition of, 744
 - ordinary, 744
 - terms of, 744
 - Apogee, 437, 693
 - Applied problems, steps for solving, 81
 - Approximately equal symbol, 460
 - Area problem, 563
 - Arithmetic mean, 729–730
 - Arithmetic progression, 732
 - Arithmetic sequence
 - application of, 734
 - common difference of, 732–733
 - definition of, 732
 - general term of, 733
 - specified term of, 734–735
 - sum of terms of, 736–737
 - Associative properties, 48–50
 - Asymptotes
 - explanation of, 386
 - horizontal, 415
 - of a hyperbola, 694–695
 - of reciprocal function, 678
 - vertical, 415
 - Average, 729–730
 - Average rate, 68
 - Average rate of change, 168–169
 - Axis
 - of a coordinate system, 151
 - of a hyperbola, 694
 - of a parabola, 570, 573
- B**
- Base
 - of an exponent, 37, 280
 - of an exponential expression, 37
 - Binomial coefficient formula, 752
 - Binomial expansion
 - general, 753
 - specified term of, 754–755
 - Binomial theorem, 753–754
 - Binomials
 - conjugates of, 497–498
 - definition of, 297
 - factoring, 363–364
 - multiplication of, 316–318, 492–493
 - raising to a power, 750–754
 - square of, 319
 - Body surface area, 71–72
 - Boundary line, 192
 - Braces, 14
- C**
- Calculator graphing method. *See* Graphing calculators
 - Cartesian coordinate system, 151
 - Celsius-Fahrenheit relationship, 190
 - Center
 - of circle, 480, 684–687
 - of ellipse, 687
 - Center-radius form of a circle, 481, 685, 687
 - Change-of-base rule, 650–651
 - Circle(s)
 - center of, 480, 684–687
 - center-radius form of, 481, 685, 687
 - definition of, 684
 - equation of, 253, 480–481, 686–687
 - explanation of, 684
 - graphing, 480–481
 - graphs of, 480, 684–686, 697
 - radius of, 684–687
 - Closed interval, 104
 - Coefficient(s)
 - binomial, 752
 - decimal, 235
 - definition of, 48, 296
 - fractional, 234–235, 327
 - leading, 297
 - numerical, 48, 296
 - Collinear points, 175
 - Combinations, definition of, 752
 - Combined variation, 438
 - Combining like terms, 48, 298
 - Common denominators, 4–5
 - Common difference of an arithmetic sequence, 732–733
 - Common factors, 1–2
 - Common logarithms
 - applications of, 647–648
 - definition of, 646
 - evaluating, 646–647
 - Common ratio of a geometric sequence, 740–741
 - Commutative properties, 48–50
 - Complementary angles, 100
 - Completing the square, 535–536, 560, 579–580
 - Complex conjugates, 514
 - Complex fraction
 - comparing methods of simplifying, 407–408
 - definition of, 405
 - simplifying with negative exponents, 408–409
 - steps to simplify, 405–406
 - Complex numbers
 - addition of, 513
 - conjugates of, 514
 - definition of, 512
 - division of, 514–515
 - imaginary part of, 512
 - multiplication of, 513–514
 - nonreal, 512
 - real part of, 512
 - standard form of, 512
 - subtraction of, 513
 - Components, 151
 - Composite function
 - definition of, 307
 - finding, 307–309

- Composition of functions, 307–309
- Compound inequalities
with *and*, 116–118
definition of, 116
with *or*, 119–121
- Compound interest
continuous, 659
formula for, 568
formulas for, 658–659
solving investment problems, 84
- Concours d'Elegance, 404
- Conditional equation, 61–62
- Conic sections
definition of, 684
examples of, 684
explanation of, 684
summary of, 697
- Conjugate
of a binomial, 497–498
of a complex number, 514
- Consecutive integers, 96
- Consistent system, 232
- Constant function, 213
- Constant of variation, 433
- Consumer Price Index (CPI), 564
- Continuous compounding
definition of, 659
formula for, 659
- Contradiction, 61–62
- Coordinate(s)
on a line, 16
of a point, 16
of points in a plane, 151
- Coordinate system
Cartesian, 151
rectangular, 151
- Cost-benefit equation, 653
- Cost-benefit model, 404
- Counting numbers, 14, 17
- Cross products of a proportion, 423
- Cube(s)
difference of, 359, 360, 363–364
of a number, 37
sum of, 359–360, 363–364
- Cube root function
definition of, 458
graph of, 458
- Cubing function, 306
- D**
- Data modeling, 182–185
- Decay
applications of, 626–627
exponential, 626–627, 660
- Decibel, 648
- Decimal(s)
addition of, 6
division of, 7–8
equivalents, 8–9
explanation of, 5
linear equations with, 59–61
multiplication of, 7–8
operations with, 6–8
place value in, 5
repeating, 8
rounding of, 7
subtraction of, 6
terminating, 8
writing as percents, 8–9
writing fractions as, 8
writing percents as, 8–9
written as fractions, 6
- Decimal coefficients, 235
- Decimal places, 7–8
- Degree
of a polynomial, 297
of a term, 296
- Denominator(s)
addition of fractions with same, 4
common, 4
explanation of, 1, 2
least common, 4–5, 397–398
rationalizing, 494–498
- Dependent equations
definition of, 232
solving a system of, 238
- Dependent variable, 201
- Depreciation, 671
- Descartes, René, 150–151
- Descending powers, 297
- Difference, 27
- Difference of cubes
definition of, 359
factoring, 359, 360, 363–364
factoring of, 359, 360
- Difference of squares, 356–357, 360, 363–364
- Direct variation
definition of, 433
as a power, 435
solving problems involving, 433–435
- Discontinuous graphs, 414
- Discriminant, 545–546, 582
- Disjoint interval, 104
- Distance, rate, time relationship, 93–95,
258–260, 423–426
- Distance between points, 29
- Distance formula(s)
distance between points, 478–479
distance to the horizon, 463, 486
- Distributive property, 45–46, 58–59, 342
- Dividend, 31
- Division
of complex numbers, 514–515
of decimals, 7–8
of fractions, 3–4
of polynomial functions, 328
of polynomials, 324–327
by powers of ten, 7–8
of rational expressions, 392
of real numbers, 31–32
synthetic, 765–768
by zero, 31
- Divisor, 31
- Domain
agreement on, 205
of a function, 202–203
of a rational function, 386–387, 414–415
of a relation, 202–203
of the variable in a rational equation, 411
- Double solution, 369
- Doubling time, 653, 658
- Downward opening parabola, 573
- E**
- e*, 649
- Earthquakes, intensity of, 295
- Elements of a set
definition of, 14
symbol for, 14
- Elimination method
for solving nonlinear systems, 704–705
for solving systems, 235–237
- Ellipse(s)
center of, 687
equation of, 688
explanation of, 684
foci of, 687
graphs of, 687–690, 697, 699
horizontal shift of, 689
intercepts of, 687–688
vertical shift of, 689
- Empty set
definition of, 14
notation for, 14
- Equality
addition property of, 57
multiplication property of, 57
- Equality symbol, 56, 58
- Equation(s)
absolute value, 125–131
of an ellipse, 688
of a circle, 253, 480–481, 686–687
conditional, 61–62
contradiction, 61–62
definition of, 21, 56
dependent, 232, 238
distinguishing from expressions, 80
equivalent, 57
exponential, 624–625, 654–655,
658–659
first-degree, 56, 153
graphs of, 152–153, 696
of a horizontal line, 155–156, 180
of hyperbola, 694
identity, 61–62
independent, 232
of an inverse function, 614–615
linear in one variable. *See* Linear
equation in one variable
linear in three variables, 245–250
linear in two variables. *See* Linear
equations in two variables
linear system of, 230, 245

- of lines, 176–185
- literal, 66
- logarithmic, 632, 654–657
- with no solution, 412–413
- nonlinear, 56, 702
- nonlinear systems of, 702–706
- power rule for, 503–504
- quadratic. *See* Quadratic equations
- quadratic in form, 549, 553–556
- radical, 503–507
- rational, 411–414, 549
- with rational expressions, 411–416
- second-degree, 532, 696, 704–705
- solution of, 56
- solution set of, 57
- translating words into, 80
- of a vertical line, 155–156, 180
- working, 246
- Equivalent equations, 57
- Equivalent forms of fractions, 32
- Equivalent inequalities, 105
- Euclid, 95
- Euclidean geometry, 95
- Euler, Leonhard, 649
- Exponential decay, 626–627, 660
- Exponential equations
 - applications of, 658–659
 - definition of, 624
 - general method for solving, 655
 - properties for solving, 624, 654
 - steps to solve, 624
- Exponential expressions
 - base of, 37
 - definition of, 37, 280
 - evaluating, 37–38
 - simplifying, 286
- Exponential functions
 - applications of, 626–627
 - characteristics of graph of, 623
 - converting to logarithmic form, 631
 - explanation of, 621
 - graphs of, 621–623
- Exponential growth, 626, 659–660
- Exponential notation, 37, 464–465
- Exponents
 - base of, 37, 280
 - definition of, 36–37, 280
 - integer, 280–285
 - negative, 281–283, 285, 408–409
 - power rules for, 284–285
 - product rule for, 280–281, 285
 - quotient rule for, 283, 285
 - rational, 464–470
 - summary of rules for, 285
 - zero, 281, 285
- Expressions
 - algebraic, 41, 56, 296
 - distinguishing from equations, 80
 - exponential, 37–38, 280, 286
 - radical. *See* Radical expressions
 - rational. *See* Rational expressions
 - translating from words to, 78–79
- Extraneous solutions, 413, 503
- F**
- Factor(s)
 - common, 1–2
 - greatest common, 1–2, 342–344
 - of numbers, 36
- Factorial notation, 751
- Factoring
 - binomials, 363–364
 - definition of, 342
 - difference of cubes, 359, 360, 363–364
 - difference of squares, 356–357, 360, 363–364
 - by grouping, 344–346
 - perfect square trinomials, 357–358, 360, 364
 - polynomials, 342, 363, 365–366
 - solving quadratic equations by, 369–373
 - sum of cubes, 359–360, 363–364
 - summary of special types of, 360
 - trinomials, 348–354, 364–365
 - using FOIL, 348–349, 351–353
- Fahrenheit-Celsius relationship, 190
- Farads, 460
- Fibonacci sequence, 725
- Finite sequence, 726
- Finite set, 14
- First-degree equations. *See also* Linear equations
 - definition of, 56, 153
 - explanation of, 153
 - graph of, 153
- Foci (*sing*: Focus)
 - of ellipse, 687
 - of hyperbola, 694
- Focus variable, 246
- FOIL method, 317–318, 348–349, 351–353, 492–493, 514
- Formula(s)
 - binomial coefficient, 752
 - for compound interest, 658–659
 - definition of, 65
 - distance, 463, 478–479, 486
 - Galileo's, 534, 603
 - Heron's, 464, 486
 - involving parentheses, 67
 - midpoint, 156–157
 - perimeter, 255
 - of the Pythagorean theorem, 477, 562
 - quadratic, 542–545, 560
 - with rational expressions, 421–422
 - resonant frequency, 460
 - slope, 161
 - solving for a specified variable of, 66–68, 373, 421–422, 561–562
 - with square roots, 561
 - vertex, 581
- Fraction(s)
 - addition of, 4–5
 - algebraic, 386
 - complex, 405–409
 - denominator of, 1, 2
 - division of, 3–4
 - equivalent forms of, 32
 - equivalents, 8–9
 - explanation of, 1
 - improper, 1, 2
 - least common denominator of, 4–5
 - linear equations with, 59–60
 - linear inequalities with, 109
 - lowest terms of, 1–2
 - mixed numbers, 2
 - multiplication of, 3
 - numerator of, 1, 2
 - operations on, 3–5
 - proper, 1
 - reciprocals of, 3
 - simplifying the, 1
 - subtraction of, 4–5
 - writing as percents, 9–10
 - writing decimals as, 6
 - writing percents as, 9–10
 - written as decimals, 8
- Fraction bar, 1, 2
- Fractional coefficients, 234–235, 327
- Froude, William, 569
- Froude number, 569
- Function(s)
 - absolute value, 678
 - coding information using, 620
 - composition, 307–309
 - composition of, 307–309
 - constant, 213
 - cube root, 458
 - cubing, 306
 - definition of, 199–200
 - definitions, variations of, 206
 - division of, 328
 - domain of, 202–203
 - equation of the inverse of, 614–615
 - evaluating, 210–212
 - exponential, 621–627, 631
 - greatest integer, 680–681
 - identity, 305
 - as Input-output machine, 202
 - inverse of, 612–617
 - linear, 176, 213, 306
 - logarithmic, 631, 634–635
 - multiplication of, 320–321
 - one-to-one, 612–614
 - operations on, 304–305
 - polynomial, 302–309, 314, 320–321
 - quadratic. *See* Quadratic functions
 - range of, 202–203
 - rational, 386–387, 414–416
 - reciprocal, 414–415, 678
 - square root, 457, 678, 698–699
 - squaring, 306

Function(s) (*continued*)
 step, 680
 translation of, 678–680
 vertical line test for, 203–204
 zeros of, 373

Function notation, 210–213

Fundamental property of rational numbers, 387

Fundamental rectangle of hyperbola, 695

Future value of an ordinary annuity, 744

$f(x)$ notation, 210–213

G

Galilei, Galileo, 534

Galileo's formula, 534, 603

General binomial expansion, 753

General term
 of an arithmetic sequence, 733
 of a geometric sequence, 741
 of a sequence, 726–727

Generalized square root function, 698–699

Geometric progression, 740

Geometric sequence
 common ratio of, 740–741
 definition of, 740
 general term of, 741
 specified term of, 742
 sum of terms of, 742–744

Grade, 161

Graph(s)
 of absolute value function, 678
 of boundary passing through origin, 194
 of circles, 480
 of cube root functions, 458
 definition of, 16
 discontinuous, 414
 of elementary functions, 678–680
 of ellipses, 687–690, 697
 of equations, 152–153, 696
 of exponential functions, 621–623
 of first-degree equations, 153
 of greatest integer functions, 681
 of horizontal lines, 155–156
 of hyperbolas, 694–697
 of intervals on number line, 103–104
 of inverses, 616–617
 of linear equations, 152–153
 of linear inequalities, 192–194
 of linear systems, 231–232, 245, 246
 of lines passing through origin, 154–155
 of logarithmic functions, 634–635
 of numbers, 16–17
 of ordered pairs of numbers, 150–151
 of parabolas, 570, 585–586, 697
 of polynomial functions, 305–307
 of quadratic functions, 570–576, 581–582
 of quadratic inequalities, 591
 of radical expressions, 457–458
 of rational functions, 414–416
 of reciprocal functions, 678

of second-degree inequalities, 708–712
 of semicircles, 698
 of square root functions, 457, 678, 698–699
 of systems of inequalities, 710–712
 of vertical lines, 155–156

Graphing calculators
 for approximation of roots, 460–461
 for displaying binomial coefficients, 753
 for generating quadratic models, 576
 for generating sequences, 727
 for graphing a parabola, 373
 for graphing circles or ellipses, 690
 for solving linear systems, 239
 square viewing window in, 690
 standard viewing window in, 157

(is) Greater than symbol, 21–22, 103

Greatest common factor (GCF)
 definition of, 342
 explanation of, 1–2
 factoring out, 342–344

Greatest integer functions
 graphs of, 681
 method for applying, 680–681

Grouping
 factoring by, 344–346
 steps to factor by, 345

Growth
 applications of, 626
 exponential, 626, 659–660

H

Half-closed interval, 104

Half-life, 660

Half-open interval, 104

Henrys, 460

Heron's formula, 464, 486

Horizontal asymptote, 415

Horizontal hyperbola, 695

Horizontal line
 equation of, 155–156, 180, 182
 graph of, 155–156
 slope of, 163–164

Horizontal line test for a one-to-one function, 613–614

Horizontal parabola, 585–586

Horizontal shift
 of an ellipse, 689
 method for applying, 679–680
 of a parabola, 572–573

Hyperbola(s)
 asymptotes of, 694–695
 equations of, 694
 explanation of, 684
 foci of, 694
 fundamental rectangle of, 695
 graphs of, 694–697
 intercepts of, 694
 transverse axis of, 694

Hypotenuse of a right triangle, 477

I

i
 definition of, 510
 powers of, 515

Identity elements, 46

Identity equations, 61–62

Identity function, 305

Identity properties, 46–47

Imaginary part of a complex number, 512

Imaginary unit
 definition of, 510
 powers of, 515

Improper fractions
 converting between mixed numbers and, 2
 explanation of, 1

Incidence rate, 418

Inconsistent system
 definition of, 232
 solving, 237, 250

Independent equations, 232

Independent variable, 201

Index of a radical, 456

Index of summation, 728

Inequality(ies)
 absolute value, 125–131
 addition property of, 105–106
 compound, 116–121
 definition of, 103
 equivalent, 105
 explanation of, 21–22
 linear in one variable. *See* Linear inequalities in one variable
 linear in two variables, 192–196
 multiplication property of, 106–109
 polynomial, 594–595
 quadratic, 590–594
 rational, 595–597
 second-degree, 708–712
 summary of symbols, 22
 symbols for, 21–22, 103
 systems of, 710–712
 three-part, 110–111

Infinite geometric sequence
 explanation of, 745
 sum of terms of, 745–747

Infinite interval, 104

Infinite sequence
 definition of, 726
 terms of, 726

Infinite set, 14

Infinity symbol, 103

Input-output machine, 202

Integer exponents, 280–285

Integers
 consecutive, 96
 definition of, 16, 17

Intensity of earthquakes, 295

Intercepts
 of ellipse, 687–688
 of hyperbola, 694

- of a parabola, 582–583
 - x -, 153–155
 - y -, 153–155, 177
 - Interest
 - compound, 84, 568, 658
 - simple, 84, 658
 - Intersection
 - of linear inequalities, 195
 - of sets, 116
 - Interval notation, 103–104
 - Inverse
 - additive, 18–19, 47–48
 - multiplicative, 30–31, 47–48
 - Inverse of a function
 - definition of, 612
 - equation of, 614–615
 - graph of, 616–617
 - steps to find the equation of, 614
 - symbol for, 612
 - Inverse properties, 47–48
 - Inverse variation
 - explanation of, 435–436
 - as a power, 436
 - Irrational numbers, 16, 17
- J**
- Joint variation, 437–438
- L**
- Leading coefficient, 297
 - Leading term, 297
 - Least common denominator (LCD)
 - definition of, 397
 - explanation of, 4
 - finding, 4–5
 - steps to find, 397
 - Legs of a right triangle, 477
 - (is) Less than symbol, 21–22, 103
 - Light-year, 295
 - Like terms
 - combining, 48, 298
 - definition of, 48, 298
 - Limit notation, 745
 - Line(s)
 - equations of, 176–185
 - horizontal, 155–156, 180, 182
 - number, 16–17
 - slope of, 161–169
 - vertical, 155–156, 180, 182
 - Line graph, 150
 - Line segment, midpoint of, 156–157
 - Linear equation in one variable
 - applications of, 92–96
 - with decimals, 59–61
 - definition of, 56
 - with fractions, 59–60
 - identifying, 56–57
 - solution of, 56
 - solution set of, 57, 111
 - solving using distributive property, 58–59
 - solving using properties of equality, 57–58
 - steps to solve, 58
 - types of, 61–62
 - Linear equations in three variables
 - explanation of, 245
 - graphs of, 246
 - system of, 245–250
 - Linear equations in two variables
 - definition of, 153
 - explanation of, 153
 - graph of, 152–153
 - point-slope form of, 178–179, 182
 - slope-intercept form of, 176, 182
 - standard form of, 153, 179, 182
 - summary of forms of, 182
 - system of, 230–239
 - x -intercept of, 153–155
 - y -intercept of, 153–155, 177
 - Linear functions, 176, 213, 306
 - Linear inequalities in one variable
 - applications of, 111–112
 - definition of, 104
 - with fractions, 109
 - solution sets of, 111
 - solving using addition property, 105–106
 - solving using multiplication property, 106–109
 - steps to solve, 108
 - three-part, 110–111
 - Linear inequalities in two variables
 - boundary line of graph, 192
 - explanation of, 192
 - graph of, 192–194
 - intersection of, 195
 - region of solution, 192
 - union of, 195–196
 - Linear models, creating, 182–185
 - Linear programming, 198
 - Linear system of equations. *See* System of linear equations
 - Literal equation, 66
 - Lithotripter, 692
 - Logarithm(s)
 - alternative forms of, 642–643
 - change-of-base rule for, 650–651
 - common, 646–648
 - definition of, 630–631
 - evaluating, 631–632, 646–647
 - exponential form of, 631
 - natural, 649–650
 - power rule for, 641–642
 - product rule for, 639–640
 - properties of, 633, 639–644, 642
 - quotient rule for, 640–641
 - Logarithmic equations
 - definition of, 632
 - properties for solving, 654
 - solving, 655–657
 - steps to solve, 657
 - Logarithmic functions
 - applications of, 635
 - with base a , 634–635
 - characteristics of graph of, 635
 - converting to exponential form, 631
 - graphs of, 634–635
 - Lowest terms
 - of a fraction, 1–2
 - of a rational expression, 387–390
 - writing radical quotients in, 498
- M**
- Mapping of sets, 201
 - Mathematical expressions from words, 78–79
 - Mathematical models, 65
 - Maximum value of a quadratic function, 583–585
 - Mean, arithmetic, 729–730
 - Members of a set, 14
 - Midpoint of a line segment formula, 156–157
 - Minimum value of a quadratic function, 583–585
 - Mixed numbers
 - converting between improper fractions and, 2
 - explanation of, 2
 - Mixture problems, 85–86, 257–258
 - Model(s)
 - to approximate data, 303
 - data, 182–185
 - mathematical, 65
 - quadratic functions as, 564–565, 574–576
 - Money problems, 92–93, 256
 - Monomials
 - definition of, 297
 - multiplication of, 315
 - Motion problems, 93–95, 258–260, 423–426, 550
 - Multiplication
 - associative property of, 48–50
 - of binomials, 316–318, 492–493
 - commutative property of, 48–50
 - of complex numbers, 513–514
 - of decimals, 7–8
 - distributive property of, 45–46
 - FOIL method of, 317–318, 492–493
 - of fractions, 3
 - identity element for, 46
 - identity property of, 46–47
 - inverse property of, 47–48
 - of monomials, 315
 - of polynomial functions, 320–321
 - of polynomials, 315–320
 - by powers of ten, 7–8
 - of radical expressions, 492–493
 - of radicals, 473
 - of radicals with different indexes, 477
 - of rational expressions, 390–391
 - of real numbers, 29–30

- Multiplication (*continued*)
 - of sum and difference of two terms, 318
 - using logarithms, 639–640
 - by zero, 30
 - Multiplication property
 - of equality, 57
 - of inequality, 106–109
 - Multiplicative identity, 46–47
 - Multiplicative inverse of a real number, 30–31, 47–48
- N**
- Natural logarithms
 - applications of, 650
 - definition of, 649
 - evaluating, 649–650
 - Natural numbers, 14, 17
 - Negative exponents
 - definition of, 281–283
 - in rational expressions, 408–409
 - rules for, 283, 285
 - Negative n th root, 456
 - Negative of a number, 18–19
 - Negative of a polynomial, 299
 - Negative reciprocals, 167
 - Negative root, 456
 - Negative slope, 166
 - Negative square root, 38
 - Newtons, 434
 - n factorial, 751
 - Noncollinear points, 253
 - Nonlinear equations, 56, 702
 - Nonlinear systems of equations
 - explanation of, 702
 - solving, 702–706
 - Nonreal complex numbers, 512
 - (is) Not equal to
 - explanation of, 21–22
 - symbol for, 21–22
 - Notation
 - exponential, 37, 464–466
 - factorial, 751
 - function, 210–213
 - interval, 103–104
 - limit, 745
 - scientific, 290–292
 - set, 14
 - set-builder, 15
 - sigma, 728
 - square root, 38, 456
 - subscript, 156
 - summation, 728–729
 - n th root(s)
 - explanation of, 456
 - exponential notation for, 464–466
 - finding for n th powers, 459
 - Null set
 - definition of, 14
 - notation for, 14
 - Number(s)
 - absolute value of, 19–20, 459
 - additive inverse of, 18–19, 47–48
 - complex, 512–515
 - counting, 14, 17
 - cubes of, 37
 - factors of, 36
 - fractions, 1
 - graphs of, 16–17. *See* Graph(s)
 - of numbers
 - imaginary, 512
 - integers, 16, 17
 - irrational, 16, 17
 - mixed, 2
 - natural, 14, 17
 - negative of, 18–19
 - nonreal complex, 512
 - opposite of, 18–19
 - ordered pair of, 150–152
 - pure imaginary, 512
 - rational, 16–17, 387
 - real. *See* Real numbers
 - reciprocal of, 3, 30–31
 - roots of, 456
 - sets of, 17–18
 - signed, 19
 - square roots of. *See* Square roots of a number
 - squares of, 37
 - whole, 14, 17
 - Number line(s)
 - coordinate of a point on, 16
 - definition of, 16
 - distance between points, 29
 - graph of a point, 16
 - graphing intervals on, 103–104
 - using, 16–17
 - Numerator(s), 1, 2
 - Numerical coefficient, 48, 296
- O**
- Ohm's law, 518
 - One-to-one function
 - definition of, 612
 - horizontal line test for, 613–614
 - inverse of, 612–613
 - Open interval, 104
 - Operations
 - with decimals, 6–8
 - on fractions, 3–5
 - on functions, 304–305, 320–321, 328
 - order of, 39–40
 - on rational expressions, 390–392, 396–401
 - on real numbers, 26–32
 - on sets, 116, 118
 - Opposite of a number, 18–19
 - Opposites, quotient of, 390
 - Order
 - of operations, 39–40
 - of a radical, 456
 - Ordered pairs
 - definition of, 150
 - graph of, 150–151
 - as solution of a linear system, 230–231
 - table of, 152
 - Ordered triple, 245
 - Ordinary annuity
 - definition of, 744
 - future value of, 744
 - payment period of, 744
 - Origin
 - definition of, 150
 - graphing line passing through, 154–155
- P**
- Pairs, ordered, 150–152
 - Parabola(s)
 - applications of, 583–585
 - axis of, 570, 573, 585
 - definition of, 570
 - explanation of, 684
 - graph of, 570, 585
 - graphing, 306
 - graphing by calculator, 373
 - graphs of, 697
 - horizontal, 585–586
 - horizontal shift of, 572–573
 - intercepts of, 582–583
 - summary of graphs of, 586
 - symmetry of, 570
 - vertex formula for, 581
 - vertex of, 570, 573, 579–581, 585
 - vertical, 571–573
 - vertical shift of, 571–573
 - Parallel lines, slope of, 166, 180–182
 - Parentheses
 - in interval notation, 103
 - solving a formula with, 67
 - Pascal, Blaise, 751
 - Pascal's triangle, 751
 - Payment period of an ordinary annuity, 744
 - Percents and percentages
 - applications of, 71–72
 - equivalents, 8–9
 - explanation of, 8
 - interpreting from graphs, 70
 - involving percent increase or decrease, 70–71
 - solving, 68–69, 83
 - writing as decimals, 8–9
 - writing as fractions, 9–10
 - writing decimals as, 8–9
 - writing fractions as, 9–10
 - Perfect square trinomial
 - definition of, 357
 - factoring of, 357–358, 360, 364
 - Perigee, 437, 693
 - Perimeter formula, 255
 - Perpendicular lines, slopes of, 167–168, 180–181

- pH
 application of, 647–648
 definition of, 647
- Pi (π), 17
- Place value in decimals, 5
- Plane
 coordinates of points in, 151
 orientation of a line in the, 166
 plotting points in, 151
 in three-dimensional
 graphing, 245
- Point(s)
 coordinate on a line, 16
 coordinates in a plane, 151
 distance between, 29
 noncollinear, 253
- Point-slope form, 178–179, 182
- Polynomial(s)
 addition of, 298–299
 definition of, 296
 degree of, 297
 in descending powers, 297
 division of, 324–327
 factoring, 342, 363, 365–366
 factoring by substitution, 354
 greatest common factor of, 342
 multiplication of, 315–320
 negative of, 299
 prime, 349
 steps to factor, 363
 subtraction of, 299–300
 term of, 296
 in x , 297
- Polynomial function(s)
 addition of, 304–305
 cubing, 306
 definition of, 302, 314
 of degree n , 302
 division of, 328
 domain of, 305
 evaluating, 302–303
 graphs of, 305–307
 identity, 305
 modeling data using, 303
 multiplication of, 320–321
 range of, 305
 squaring, 306
 subtraction of, 304–305
- Polynomial inequality
 solving, 594–595
 third-degree, 594–595
- Positive root, 456
- Positive slope, 166
- Positive square root, 38
- Power rule
 for exponents, 284–285
 for logarithms, 641–642
 for radical equations, 503–504
- Powers
 definition of, 37, 280
 descending, 297
- Powers of i
 explanation of, 515
 simplifying, 515
- Powers of ten
 division by, 7–8
 explanation of, 5
 multiplication by, 7–8
- Pricing problem, 262–263
- Prime polynomial, 349
- Principal n th root, 456
- Principal square root, 38, 456
- Product
 definition of, 29
 of sum and difference of two terms, 318
- Product rule
 for exponents, 280–281, 285
 for logarithms, 639–640, 642
 for radicals, 473
- Progression
 arithmetic, 732
 geometric, 740
- Proper fractions, 1
- Proportion
 cross products of a , 423
 definition of, 422
 solving, 422–423
- Proportional, 433
- Proposed solution, 503
- Pure imaginary number, 512
- Pythagorean theorem, 477–478, 562
- Q**
- Quadrants, 151
- Quadratic equations
 applications of, 371–372, 549–552,
 562–565
 completing the square method for
 solving, 535–538, 560
 with complex solutions, 538
 definition of, 368, 532
 discriminant of, 545–547, 582
 double solution for, 369
 factoring method for solving, 369
 with nonreal complex solutions, 538
 quadratic formula for solving,
 542–545, 560
 solving radical equations that lead
 to, 552
 square root property for solving,
 532–535, 560
 standard form of, 368, 532
 steps to solve by completing the
 square, 536
 steps to solve by factoring, 369
 substitution method for solving, 553–556
 summary of methods for solving, 560
 types of solutions, 545–547
 zero-factor property for solving,
 369–370, 532, 560
- Quadratic formula
 definition of, 542
- derivation of, 542
 solving quadratic equations using,
 542–545, 560
- Quadratic functions
 applications using, 564–565, 574–576,
 583–585
 definition of, 372, 571
 general characteristics of, 574
 graphs of, 570–576, 581–582
 maximum value of, 583–585
 minimum value of, 583–585
 solving applied problems using, 564–565
 steps to graph, 581
 zeros of, 373
- Quadratic in form equations, 549, 553–556
- Quadratic inequalities
 definition of, 590
 graphs of, 591
 steps to solve, 593
- Quotient, 31
- Quotient of opposites, 390
- Quotient rule
 for exponents, 283, 285
 for logarithms, 640–641, 642
 for radicals, 474
- R**
- Radical equations
 definition of, 503
 extraneous solutions of, 503
 power rule for solving, 503–504
 solving with additional steps, 504–506
 solving with indexes greater than 2, 507
 steps for solving, 504
- Radical expressions
 addition of, 487–489
 definition of, 457
 graphs of, 457–458
 multiplication of, 492–493
 rationalizing the denominator of,
 494–498
 subtraction of, 487–489
- Radical symbol, 38, 456
- Radicals
 conditions for simplified form, 474, 502
 converting between rational exponents
 and, 467–468
 definition of, 456
 equations with, 503–507
 index of, 456
 multiplication of, 473
 order of, 456
 product rule for, 473
 quotient rule for, 474
 simplifying, 474–477
- Radicand, 456
- Radius, 684–687
- Radius of a circle, 480–481
- Range
 of a function, 202–203
 of a relation, 202–203

- Rate in motion problems, 93–95, 258–260, 423–426
 - Rate of change
 - average, 168–169
 - comparing, 20
 - Rate of work, 426–427
 - Ratio, 422
 - Rational equations
 - definition of, 411
 - domain of, 411
 - with no solutions, 412–413
 - solving, 411–414
 - solving in quadratic form, 549
 - Rational exponents
 - converting between radicals and, 467–468
 - evaluating terms with, 464–467
 - explanation of, 465
 - radical form of, 467
 - rules for, 468–470
 - Rational expressions
 - addition of, 396–397, 400–401
 - addition with different denominators, 398–399
 - addition with opposite denominators, 400
 - applications of, 421–427
 - definition of, 386
 - division of, 392
 - equations with, 411–416
 - formulas with, 421–422
 - lowest terms of, 387–390
 - multiplication of, 390–391
 - operations on, 390–392, 396–401
 - quotient of opposites in, 390
 - reciprocals of, 391
 - simplifying with negative exponents, 408–409
 - steps to multiply, 390
 - subtraction of, 396, 399–401
 - subtraction with different denominators, 398–399
 - Rational functions
 - definition of, 386, 414
 - discontinuous, 414
 - domains of, 386–387, 414–415
 - graphs of, 414–416
 - inverse variation and, 436
 - Rational inequality
 - definition of, 595
 - steps to solve, 595
 - Rational numbers
 - definition of, 16–17
 - as exponents, 465
 - fundamental property of, 387
 - Rationalizing
 - a binomial denominator, 497–498
 - the denominator, 494–498
 - the numerator, 501
 - Real numbers
 - addition of, 26–27
 - additive inverse of, 19, 47–48
 - definition of, 17
 - division of, 31–32
 - multiplication of, 29–30
 - multiplicative inverse of, 30–31, 47–48
 - operations on, 26–32
 - properties of, 45–50
 - reciprocals of, 30–31
 - subtraction of, 27–29
 - Real part of a complex number, 512
 - Reciprocal(s)
 - explanation of, 3
 - of fractions, 3
 - negative, 167
 - negative exponents leading to, 282
 - of a rational expression, 391
 - of a real number, 30–31
 - Reciprocal function
 - definition of, 414, 678
 - graph of, 414–415
 - Rectangular coordinate system
 - definition of, 151
 - plotting points in, 151
 - quadrants of, 151
 - Regions in the real number plane, 192
 - Relation
 - definitions of, 199–202
 - domain of, 202–203
 - range of, 202–203
 - Relative error, 131
 - Remainder theorem, 767–768
 - Repeating decimals, 8
 - Resonant frequency formula, 460
 - Richter scale, 295
 - Right triangle
 - hypotenuse of, 477
 - legs of, 477
 - Pythagorean theorem on, 477
 - Rise, 161
 - Roots
 - calculator approximation of, 460–461
 - cube, 456
 - fourth, 456
 - negative, 456
 - n th, 456
 - of numbers, 456
 - positive, 456
 - principal, 456
 - simplifying, 456
 - square, 38–39, 456
 - Rounding, of decimals, 7
 - Run, 161
- S**
- Scale, 157
 - Scatter diagram, 183, 575
 - Scientific notation
 - application of, 292
 - converting positive numbers from scientific notation, 291–292
 - converting positive numbers to, 290–291
 - definition of, 290
 - Scrap value, 671
 - Second-degree equations, 532, 696, 704–705
 - Second-degree inequalities
 - explanation of, 708
 - graphs of, 708–712
 - Semicircles, 698
 - Semiperimeter, 464
 - Sequence(s)
 - applications of, 727
 - arithmetic, 732–738
 - definition of, 726
 - Fibonacci, 725
 - finite, 726
 - general term of, 726–727
 - geometric, 740–747
 - infinite, 726
 - terms of, 726
 - Series
 - definition of, 728
 - finite, 728
 - infinite, 728
 - Set(s)
 - definition of, 14
 - elements of, 14
 - empty, 14
 - finite, 14
 - infinite, 14
 - intersection of, 116
 - mapping of, 201
 - members of, 14
 - null, 14
 - operations on, 116, 118
 - solution. *See* Solution set(s)
 - union of, 116, 118
 - Set braces, 14
 - Set-builder notation, 15
 - Set operations, 116, 118
 - Sigma notation, 728
 - Signed numbers, 19
 - Similar triangles, 430
 - Simple interest, 66, 84, 658
 - Simplified form of a radical, 474, 502
 - Simplifying the fraction, 1
 - Slope(s)
 - definition of, 161
 - formula for, 161
 - of a horizontal line, 163–164
 - of a line, 161–169
 - negative, 166
 - of parallel lines, 166, 180–181
 - of perpendicular lines, 167–168, 180–181
 - positive, 166
 - undefined, 164, 166, 167
 - of a vertical line, 163–164
 - Slope-intercept form, 176, 182
 - Solution of an equation, 56
 - Solution set(s)
 - definition of, 57
 - of equations and inequalities, 111

- of a linear system, 230
 - of system of inequalities, 710–712
 - Solving a literal equation, 66
 - Solving for a specified variable
 - definition of, 66
 - explanation of, 421–422
 - with power rule, 507
 - with square roots, 561
 - steps to solve, 66
 - where factoring is necessary, 373
 - Speed in motion problems, 93–95, 258–260, 423–426
 - Square(s)
 - of a binomial, 319
 - difference of, 356–357, 360, 363–364
 - of a number, 37
 - sum of, 357, 364
 - Square bracket in interval notation, 103
 - Square root function
 - definition of, 457
 - generalized, 698–699
 - graph of, 457
 - graphs of, 678, 698–699
 - Square root notation, 456
 - Square root property, 532–533, 560
 - Square roots of a number
 - definition of, 38, 456
 - finding, 38–39
 - for a negative real number, 510
 - principal, 38, 456
 - simplifying, 459
 - solving formulas involving, 561–562
 - symbol for, 38, 456
 - Square viewing window, 690
 - Squaring function, 306
 - Standard form
 - of a complex number, 512
 - of a linear equation, 153, 179, 182
 - of a quadratic equation, 368, 532
 - Standard viewing window, 157
 - Step functions, 680
 - Study skills
 - analyzing test results, S-9
 - completing homework, S-4
 - managing time, S-6
 - preparing for math final exam, S-10
 - reading math textbook, S-2
 - reviewing a chapter, S-7
 - taking lecture notes, S-3
 - taking math tests, S-8
 - using math textbook, S-1
 - using study cards, S-5
 - Subscript notation, 156
 - Substitution method
 - for factoring a polynomial, 354
 - to solve nonlinear systems, 702–704
 - for solving quadratic equations, 553–556
 - for solving systems, 232–235
 - Subtraction
 - of complex numbers, 513
 - of decimals, 6
 - of fractions, 4–5
 - of polynomial functions, 304–305
 - of polynomials, 299–300
 - of radical expressions, 487–489
 - of rational expressions, 396–401
 - of real numbers, 27–29
 - Sum, definition of, 26
 - Sum and difference of two terms, product of, 318
 - Sum of cubes
 - definition of, 359
 - factoring, 359–360, 363–364
 - Sum of measures of angles of a triangle, 265
 - Sum of squares, 357, 364
 - Sum of terms of a geometric sequence, 742–744, 745–747
 - Sum of terms of an arithmetic sequence, 736–738
 - Summation notation, 728–729
 - Supplementary angles, 100, 265
 - Symmetry about an axis, 570
 - Synthetic division, 765–768
 - System of linear equations in three variables
 - applications of, 261–263
 - explanation of, 245
 - geometry of, 245–246, 261–262
 - graphs of, 245–246
 - inconsistent, 250
 - special cases of, 249–250
 - steps to solve, 246
 - System of linear equations in two variables
 - applications of, 254–263
 - calculator graphing method for solving, 239
 - consistent, 232
 - with dependent equations, 232
 - elimination method for solving, 235–237
 - explanation of, 230
 - with fractional coefficients, 234–235
 - graphs of, 231–232
 - inconsistent, 232
 - with independent equations, 232
 - solution set of, 230
 - special cases of, 237–239
 - steps to solve applications of, 254
 - steps to solve by elimination, 236
 - steps to solve by substitution, 233
 - substitution method for solving, 232–235
 - Systems of inequalities
 - explanation of, 710
 - graphs of, 710–712
 - solution set of, 710–712
 - Systems of nonlinear equations, 702–706
- T**
- Table of ordered pairs, 152
 - Term(s)
 - of an annuity, 744
 - of a binomial expansion, 754–755
 - coefficient of, 48, 297
 - combining, 48, 298
 - definition of, 48
 - degree of, 296
 - leading, 297
 - like, 48, 298
 - numerical coefficient of, 48, 296
 - of a polynomial, 296
 - of a sequence, 726
 - unlike, 48
 - Terminating decimals, 8
 - Third-degree polynomial inequalities, 594–595
 - Three-part inequalities
 - definition of, 110
 - solving, 110–111
 - Threshold sound, 648
 - Threshold weight, 472
 - Time in motion problems, 258–260, 423–426
 - Tolerance, 131
 - Traffic intensity, 418
 - Translating words into equations, 80
 - Translating words into mathematical expressions, 78–79
 - Translations of functions, 678–680
 - Transverse axis of hyperbola, 694
 - Triangle(s)
 - Pascal's, 751
 - right, 477
 - similar, 430
 - sum of angles of, 265
 - Trinomials
 - definition of, 297
 - factoring of, 348–354, 364–365
 - perfect square, 357–358, 360, 364
 - steps to factor, 349, 352
 - Triple, ordered, 245
- U**
- Undefined slope, 164, 166, 167
 - Union
 - of linear inequalities, 195–196
 - of sets, 116, 118
 - Unlike terms, 48
- V**
- Variable
 - definition of, 15
 - dependent, 201
 - independent, 201
 - solving for a specified. *See* Solving for a specified variable
 - Variation
 - combined, 438
 - constant of, 433
 - direct, 433
 - explanation of, 433
 - inverse, 435–437
 - joint, 437–438
 - steps to solve problems with, 435
 - Verbal expressions into mathematical expressions, 78–79
 - Verbal sentences into equations, 80

Vertex of a parabola
 definition of, 570
 explanation of, 573, 585
 finding, 579–581
 formula for, 581

Vertical angles, 100, 265

Vertical asymptote, 415

Vertical hyperbola, 696

Vertical line
 equation of, 155–156, 180, 182
 graph of, 155–156
 slope of, 163–164

Vertical line test for a function, 203–204

Vertical parabola
 graphing, 571–573
 vertex of, 571, 579–581
 x -intercepts of, 582–583

Vertical shift
 of ellipse, 689
 method for applying, 679–680

Vertical shift of a parabola, 571–573

W

Whole numbers, 14, 17

Windchill factor, 472

Work problems, 426–427,
 551–552

Working equation, 246

X

x -axis, 151

x -intercept
 definition of, 153
 of a line, 153–154
 of a parabola, 582–583

Y

y -axis, 151

y -intercept
 definition of, 153
 of a line, 153–154, 177

Z

Zero(s)
 division by, 31
 of a function, 373
 multiplication by, 30
 of a quadratic function,
 373

Zero exponent
 definition of, 281
 summary of rules for, 285


Zero-factor property
 definition of, 367
 explanation of, 532, 560
 solving a formula for a specified
 variable with, 373
 solving an equation with,
 367–371, 532
 solving applied problems with,
 371–372

Zero-level earthquakes, 295

Triangles and Angles

Right Triangle

Triangle has one 90° (right) angle.




Pythagorean Theorem (for right triangles)

$$a^2 + b^2 = c^2$$

Right Angle


Measure is 90° .



Isosceles Triangle


Two sides are equal.

$$AB = BC$$



Straight Angle


Measure is 180° .



Equilateral Triangle


All sides are equal.

$$AB = BC = CA$$




Complementary Angles

The sum of the measures of two complementary angles is 90° .




Sum of the Angles of Any Triangle

$$A + B + C = 180^\circ$$



Supplementary Angles

The sum of the measures of two supplementary angles is 180° .




Similar Triangles

Corresponding angles are equal. Corresponding sides are proportional.


$$A = D, B = E, C = F$$

$$\frac{AB}{DE} = \frac{AC}{DF} = \frac{BC}{EF}$$



Vertical Angles

Vertical angles have equal measures.



Geometry Formulas

Figure


Formulas

Illustration

Square

Perimeter: $P = 4s$

Area: $\mathcal{A} = s^2$



Rectangle

Perimeter: $P = 2L + 2W$


Area: $\mathcal{A} = LW$



Triangle

Perimeter: $P = a + b + c$


Area: $\mathcal{A} = \frac{1}{2}bh$



Parallelogram

Perimeter: $P = 2a + 2b$


Area: $\mathcal{A} = bh$



Trapezoid

Perimeter: $P = a + b + c + B$

Area: $\mathcal{A} = \frac{1}{2}h(b + B)$




Circle







Diameter: $d = 2r$

Circumference: $C = 2\pi r$
 $C = \pi d$

Area: $\mathcal{A} = \pi r^2$



Geometry Formulas

| Figure | Formulas | Illustration |
|--------------------------------|-----------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| Cube | Volume: $V = e^3$
Surface area: $S = 6e^2$ |  |
| Rectangular Solid | Volume: $V = LWH$
Surface area:
$S = 2HW + 2LW + 2LH$ |  |
| Right Circular Cylinder | Volume: $V = \pi r^2 h$
Surface area: $S = 2\pi r h + 2\pi r^2$
(Includes both circular bases) |  |
| Cone | Volume: $V = \frac{1}{3}\pi r^2 h$
Surface area: $S = \pi r\sqrt{r^2 + h^2} + \pi r^2$
(Includes circular base) |  |
| Right Pyramid | Volume: $V = \frac{1}{3}Bh$
$B =$ area of the base |  |
| Sphere | Volume: $V = \frac{4}{3}\pi r^3$
Surface area: $S = 4\pi r^2$ |  |

Other Formulas

Distance: $d = rt$ (r = rate or speed, t = time)

Percent: $p = br$ (p = percentage, b = base, r = rate)

Temperature: $F = \frac{9}{5}C + 32$ $C = \frac{5}{9}(F - 32)$

Simple Interest: $I = prt$ (p = principal or amount invested, r = rate or percent, t = time in years)