

Protocol 2000

Protocol 2000

For All Cancers, Stage IV Cancer, and Other Immediately Life-Threatening Diseases

Depending on how bad you feel, start with 1, 2 or 3 drops of MMS every hour for at least 8 hours a day. The worse you feel, the fewer drops you should take.

This is an intense program that often works on cancer at any stage, and has worked for Stage IV cancers. It is the best I know at this time for advanced cancers. I have used it quite a few times on people here in Africa and also for two years in Mexico. It could be used with many other diseases as well. We have successfully treated most known life-threatening diseases in the last stages with this protocol. Hepatitis A, B, and C are examples and the final result was a negative reading. Please read about Protocol 1000 first for the basic technical data you will need.

Never give up on a person. Even if they have 2 hours left to live, get them started. Get some MMS1 into them. Up to this point, MMS is the opposite of drugs. Hundreds of thousands of people die taking drugs and everybody thinks that it is normal. "Well," the doctor will say, "we were trying."

Think of this – over the past ten years, there has only been one death reported of someone taking MMS and that person was just not in good enough shape to be helped, even by MMS. Now millions of people have at least tried MMS and most of the millions were sick from something, else there would be no point in their trying it.

- No deaths have been reported.

In the 100,000 malaria cases treated by MMS, there were no deaths. Normally there would have been from 400 to 500 deaths in that number of malaria cases.

Step One of Two

Begin by taking a one-drop dose of MMS. Do this each hour for several hours until it appears that one drop is tolerated OK. Just add one drop of MMS to an empty dry glass and then a drop of 50% citric acid. Be sure to wait for at least 20 seconds before adding juice or water and drinking it.

Protocol 2000

You can use up to a full glass of water or juice to overcome most of the bad taste. Do this every hour for 10 hours straight each day.

However, one drop is not enough. That is just to get you started. Once you determine that you can tolerate that one drop, go to two drops each hour. You don't have to wait a whole day to go to two drops. If there is no nausea in the first two hours, go ahead and try two drops an hour. Two drops are activated with 10 drops of lemon or 10% citric or two drops of 50% citric.

- The one juice you must not use is orange juice. And forego use of vitamin C while on the 2000 protocol. Both orange juice and vitamin C prevent the action of MMS. If you add juice to improve the flavor, use only juices with no added Vitamin C.
- You can wait until two hours after completing the protocol for the day and then take your vitamin C.
- If you become sicker than you already are, stop taking MMS until the increased portion of the sickness is gone and start back with a smaller dose.

You can increase to 8 or even 10 drops of MMS1 each hour. However that is a lot, and most people will become nauseous and not tolerate that much until the cancer or other disease is completely gone. If you do go this high, place the activated mix in at least three quarters of a glass of water or juice or a full glass (8 ounces). That way you may be able to tolerate the taste a little more than if you used less water or juice and thus had a more concentrated mix.

Step Two of Two

Now the next step is a little hard for some people, but if you want a good result, it is needed. You must take MMS2 capsules at least twice a day at first and then increase to 5 times a day.

Start slowly, as with everything. MMS2 is hypochlorous acid, the same acid that the human immune system uses to kill things. It has the smell of chlorine, but it is not chlorine -- it is just made with chlorine like table salt. Although it tastes a little worse than table salt, it has approximately the same poison index. So you could take a tremendous amount more than Protocol 2000 calls for before it became dangerous.

At the time of this writing, 1/15/2011 send an email to sales@min-eral-solutions.net and he will ship a 60-capsule bottle to you. The cost is about \$60 USD but that includes the shipping fee as well. I feel safe in telling you

Protocol 2000

about him because he is in Thailand. He ships them right out and you usually get them in less than a week.

The immune system simply cannot make enough of this chemical to kill a cancer, but you can supplement it with these capsules. It's the very same thing your immune system makes. This acid is generated by the chemical called calcium hypochlorite when you buy it.

You could make your own MMS 2 capsules by scooping this chemical into a size zero gel capsule. Cap it and take a capsule 4 times a day with 2 hours separating each capsule, but go slowly. Start with capsules only one quarter filled and if there is little or no nausea or diarrhea, fill the capsules more until they are full each time. This data is explained on my website www.jimhumble.biz under the MMS2 menu item.

If you doubt my word about this acid being used by the immune system, go to Google and search for "hypochlorous acid and the immune system." You will get dozens or hundreds of sites telling you about it being used by the immune system.

However, I am the only one who suggests that you take it by mouth. I have been taking it for 4 years, and hundreds of people I sent it to in the beginning have taken it as well and now thousands have bought it from a supplier or made their own. (No, I never did charge for it. I mention this because so many critics keep saying I am making millions.)

Actually it is perfectly logical to take it. Your system uses it extensively throughout the body to kill pathogens and the human body has been using it for hundreds of thousands of years. It would seem perfectly logical to assume that the system might need some if a disease is not being destroyed. Only people who sell drugs would want you to not take it, because it will heal you and you won't need their ineffective drugs.

Managing Your Protocol 2000 Doses

OK, so now you have the two things that you must take to kill your cancer. You must saturate your body with MMS1 and MMS2 to the point of killing the cancer quickly, but not so quickly that it makes you feel sicker than you already are. Killing the cancer too quickly can kill you too, so go slowly. You must take as much as you can without developing nausea, vomiting, diarrhea, or severe fatigue.

Protocol 2000

It is a balancing act. Increase your doses until you notice some slight indication that nausea, vomiting, diarrhea, severe fatigue, or all four are beginning to show up. If you know nausea is coming on, just reduce the next dose by about half. If nausea still hits, then stop the intake of MMS1 and 2 until the nausea passes, and start again immediately, but reduce the dose of both MMS1 and 2.

You should notice the cancer getting better (smaller or less pain) very soon, or maybe you'll just notice it not getting worse. However, do not remain taking a reduced dosage. After several hours of taking a reduced dose that does not make you feel worse, begin slowly increasing again. Soon you will get the hang of the balancing act of taking as much as possible but not letting it make you feel sicker. Find your body's balancing point so that you do not feel any of the negative symptoms I have mentioned. Overdosing is never productive.

You may find it necessary to also use Protocol 3000. If the cancer is not going away, many people find that Protocol 3000 helps overcome it. A few people have been successful in using Protocol 3000 on their cancer, and nothing else, as it is a very powerful protocol and it can kill many diseases. But normally one will need both Protocols 2000 and 3000.

However, the same rules apply.

- Do not make yourself feel worse and if you do, reduce your intake of MMS 1 and 2.

The minute you feel that nausea is starting, or something else is feeling bad, reduce your dosage or at least do not increase. This goes for both MMS1 and 2. With MMS2 you can reduce the dosage by opening the capsule and dumping part of the powder out. Generally, once you are taking a full capsule, you would increase the dose merely by just taking more capsules in a day. Maintain the two-hour separation, but you can take it more often if it doesn't make you feel sicker. Don't take more than one capsule at one time. You can also reduce the amount of MMS used with Protocol 3000 by using smaller areas of the skin.