











□ APPENDIXES


A Numbers, Inequalities, and Absolute Values

1. $|5 - 23| = |-18| = 18$
2. $|5| - |-23| = 5 - 23 = -18$
3. $|-π| = π$ because $π > 0$.
4. $|π - 2| = π - 2$ because $π - 2 > 0$.
5. $|\sqrt{5} - 5| = -(\sqrt{5} - 5) = 5 - \sqrt{5}$ because $\sqrt{5} - 5 < 0$.
6. $||-2| - |-3|| = |2 - 3| = |-1| = 1$
7. For $x < 2$, $x - 2 < 0$, so $|x - 2| = -(x - 2) = 2 - x$.
8. For $x > 2$, $x - 2 > 0$, so $|x - 2| = x - 2$.
9. $|x + 1| = \begin{cases} x + 1 & \text{for } x + 1 \geq 0 \Leftrightarrow x \geq -1 \\ -(x + 1) & \text{for } x + 1 < 0 \Leftrightarrow x < -1 \end{cases}$
10. $|2x - 1| = \begin{cases} 2x - 1 & \text{for } 2x - 1 \geq 0 \Leftrightarrow x \geq \frac{1}{2} \\ 1 - 2x & \text{for } 2x - 1 < 0 \Leftrightarrow x < \frac{1}{2} \end{cases}$
11. $|x^2 + 1| = x^2 + 1$ (since $x^2 + 1 \geq 0$ for all x).
12. Determine when $1 - 2x^2 < 0 \Leftrightarrow 1 < 2x^2 \Leftrightarrow x^2 > \frac{1}{2} \Leftrightarrow \sqrt{x^2} > \sqrt{\frac{1}{2}} \Leftrightarrow |x| > \sqrt{\frac{1}{2}} \Leftrightarrow x < -\frac{1}{\sqrt{2}}$ or $x > \frac{1}{\sqrt{2}}$. Thus, $|1 - 2x^2| = \begin{cases} 1 - 2x^2 & \text{if } -\frac{1}{\sqrt{2}} \leq x \leq \frac{1}{\sqrt{2}} \\ 2x^2 - 1 & \text{if } x < -\frac{1}{\sqrt{2}} \text{ or } x > \frac{1}{\sqrt{2}} \end{cases}$
13. $2x + 7 > 3 \Leftrightarrow 2x > -4 \Leftrightarrow x > -2$, so $x \in (-2, \infty)$. 
14. $3x - 11 < 4 \Leftrightarrow 3x < 15 \Leftrightarrow x < 5$, so $x \in (-\infty, 5)$. 
15. $1 - x \leq 2 \Leftrightarrow -x \leq 1 \Leftrightarrow x \geq -1$, so $x \in [-1, \infty)$. 
16. $4 - 3x \geq 6 \Leftrightarrow -3x \geq 2 \Leftrightarrow x \leq -\frac{2}{3}$, so $x \in (-\infty, -\frac{2}{3}]$. 
17. $2x + 1 < 5x - 8 \Leftrightarrow 9 < 3x \Leftrightarrow 3 < x$, so $x \in (3, \infty)$. 
18. $1 + 5x > 5 - 3x \Leftrightarrow 8x > 4 \Leftrightarrow x > \frac{1}{2}$, so $x \in (\frac{1}{2}, \infty)$. 
19. $-1 < 2x - 5 < 7 \Leftrightarrow 4 < 2x < 12 \Leftrightarrow 2 < x < 6$, so $x \in (2, 6)$. 
20. $1 < 3x + 4 \leq 16 \Leftrightarrow -3 < 3x \leq 12 \Leftrightarrow -1 < x \leq 4$, so $x \in (-1, 4]$. 
21. $0 \leq 1 - x < 1 \Leftrightarrow -1 \leq -x < 0 \Leftrightarrow 1 \geq x > 0$, so $x \in (0, 1]$. 


22. $-5 \leq 3 - 2x \leq 9 \Leftrightarrow -8 \leq -2x \leq 6 \Leftrightarrow 4 \geq x \geq -3$, so $x \in [-3, 4]$.



23. $4x < 2x + 1 \leq 3x + 2$. So $4x < 2x + 1 \Leftrightarrow 2x < 1 \Leftrightarrow x < \frac{1}{2}$, and $2x + 1 \leq 3x + 2 \Leftrightarrow -1 \leq x$. Thus, $x \in [-1, \frac{1}{2})$.



24. $2x - 3 < x + 4 < 3x - 2$. So $2x - 3 < x + 4 \Leftrightarrow x < 7$, and $x + 4 < 3x - 2 \Leftrightarrow 6 < 2x \Leftrightarrow 3 < x$, so $x \in (3, 7)$.




25. $(x - 1)(x - 2) > 0$. Case 1: (both factors are positive, so their product is positive)

$x - 1 > 0 \Leftrightarrow x > 1$, and $x - 2 > 0 \Leftrightarrow x > 2$, so $x \in (2, \infty)$.

Case 2: (both factors are negative, so their product is positive)


$x - 1 < 0 \Leftrightarrow x < 1$, and $x - 2 < 0 \Leftrightarrow x < 2$, so $x \in (-\infty, 1)$.

Thus, the solution set is $(-\infty, 1) \cup (2, \infty)$.




26. $(2x + 3)(x - 1) \geq 0$. Case 1: $2x + 3 \geq 0 \Leftrightarrow x \geq -\frac{3}{2}$, and $x - 1 \geq 0 \Leftrightarrow x \geq 1$, so $x \in [1, \infty)$.


Case 2: $2x + 3 \leq 0 \Leftrightarrow x \leq -\frac{3}{2}$, and $x - 1 \leq 0 \Leftrightarrow x \leq 1$, so $x \in (-\infty, -\frac{3}{2}]$. Thus, the solution set is $(-\infty, -\frac{3}{2}] \cup [1, \infty)$.




27. $2x^2 + x \leq 1 \Leftrightarrow 2x^2 + x - 1 \leq 0 \Leftrightarrow (2x - 1)(x + 1) \leq 0$. Case 1: $2x - 1 \geq 0 \Leftrightarrow x \geq \frac{1}{2}$, and $x + 1 \leq 0 \Leftrightarrow x \leq -1$, which is an impossible combination. Case 2: $2x - 1 \leq 0 \Leftrightarrow x \leq \frac{1}{2}$, and $x + 1 \geq 0 \Leftrightarrow x \geq -1$, so $x \in [-1, \frac{1}{2}]$. Thus, the solution set is $[-1, \frac{1}{2}]$.



28. $x^2 < 2x + 8 \Leftrightarrow x^2 - 2x - 8 < 0 \Leftrightarrow (x - 4)(x + 2) < 0$. Case 1: $x > 4$ and $x < -2$, which is impossible. Case 2: $x < 4$ and $x > -2$. Thus, the solution set is $(-2, 4)$.




29. $x^2 + x + 1 > 0 \Leftrightarrow x^2 + x + \frac{1}{4} + \frac{3}{4} > 0 \Leftrightarrow (x + \frac{1}{2})^2 + \frac{3}{4} > 0$. But since $(x + \frac{1}{2})^2 \geq 0$ for every real x , the original inequality will be true for all real x as well. Thus, the solution set is $(-\infty, \infty)$.




30. $x^2 + x > 1 \Leftrightarrow x^2 + x - 1 > 0$. Using the quadratic formula, we obtain


$x^2 + x - 1 = (x - \frac{-1-\sqrt{5}}{2})(x - \frac{-1+\sqrt{5}}{2}) > 0$. Case 1: $x - \frac{-1-\sqrt{5}}{2} > 0$ and $x - \frac{-1+\sqrt{5}}{2} > 0$, so that $x > \frac{-1+\sqrt{5}}{2}$. Case 2: $x - \frac{-1-\sqrt{5}}{2} < 0$ and $x - \frac{-1+\sqrt{5}}{2} < 0$, so that $x < \frac{-1-\sqrt{5}}{2}$. Thus, the solution set is $(-\infty, \frac{-1-\sqrt{5}}{2}) \cup (\frac{-1+\sqrt{5}}{2}, \infty)$.




31. $x^2 < 3 \Leftrightarrow x^2 - 3 < 0 \Leftrightarrow (x - \sqrt{3})(x + \sqrt{3}) < 0$. *Case 1:* $x > \sqrt{3}$ and $x < -\sqrt{3}$, which is impossible.
Case 2: $x < \sqrt{3}$ and $x > -\sqrt{3}$. Thus, the solution set is $(-\sqrt{3}, \sqrt{3})$.
 Another method: $x^2 < 3 \Leftrightarrow |x| < \sqrt{3} \Leftrightarrow -\sqrt{3} < x < \sqrt{3}$.



32. $x^2 \geq 5 \Leftrightarrow x^2 - 5 \geq 0 \Leftrightarrow (x - \sqrt{5})(x + \sqrt{5}) \geq 0$. *Case 1:* $x \geq \sqrt{5}$ and $x \geq -\sqrt{5}$, so $x \in [\sqrt{5}, \infty)$.
Case 2: $x \leq \sqrt{5}$ and $x \leq -\sqrt{5}$, so $x \in (-\infty, -\sqrt{5}]$. Thus, the solution set is $(-\infty, -\sqrt{5}] \cup [\sqrt{5}, \infty)$.
 Another method: $x^2 \geq 5 \Leftrightarrow |x| \geq \sqrt{5} \Leftrightarrow x \geq \sqrt{5}$ or $x \leq -\sqrt{5}$.




33. $x^3 - x^2 \leq 0 \Leftrightarrow x^2(x - 1) \leq 0$. Since $x^2 \geq 0$ for all x , the inequality is satisfied when $x - 1 \leq 0 \Leftrightarrow x \leq 1$. Thus, the solution set is $(-\infty, 1]$.



34. $(x + 1)(x - 2)(x + 3) = 0 \Leftrightarrow x = -1, 2, \text{ or } -3$. Constructing a table:

Interval	$x + 1$	$x - 2$	$x + 3$	$(x + 1)(x - 2)(x + 3)$
$x < -3$	-	-	-	-
$-3 < x < -1$	-	-	+	+
$-1 < x < 2$	+	-	+	-
$x > 2$	+	+	+	+


Thus, $(x + 1)(x - 2)(x + 3) \geq 0$ on $[-3, -1]$ and $[2, \infty)$, and the solution set is $[-3, -1] \cup [2, \infty)$.



35. $x^3 > x \Leftrightarrow x^3 - x > 0 \Leftrightarrow x(x^2 - 1) > 0 \Leftrightarrow x(x - 1)(x + 1) > 0$. Constructing a table:

Interval	x	$x - 1$	$x + 1$	$x(x - 1)(x + 1)$
$x < -1$	-	-	-	-
$-1 < x < 0$	-	-	+	+
$0 < x < 1$	+	-	+	-
$x > 1$	+	+	+	+


Since $x^3 > x$ when the last column is positive, the solution set is $(-1, 0) \cup (1, \infty)$.




$$36. x^3 + 3x < 4x^2 \Leftrightarrow x^3 - 4x^2 + 3x < 0 \Leftrightarrow x(x^2 - 4x + 3) < 0 \Leftrightarrow x(x-1)(x-3) < 0.$$

Interval	x	$x-1$	$x-3$	$x(x-1)(x-3)$
$x < 0$	-	-	-	-
$0 < x < 1$	+	-	-	+
$1 < x < 3$	+	+	-	-
$x > 3$	+	+	+	+


Thus, the solution set is $(-\infty, 0) \cup (1, 3)$.



37. $1/x < 4$. This is clearly true for $x < 0$. So suppose $x > 0$. then $1/x < 4 \Leftrightarrow 1 < 4x \Leftrightarrow \frac{1}{4} < x$. Thus, the solution set is $(-\infty, 0) \cup (\frac{1}{4}, \infty)$.



38. $-3 < 1/x \leq 1$. We solve the two inequalities separately and take the intersection of the solution sets. First, $-3 < 1/x$ is clearly true for $x > 0$. So suppose $x < 0$. Then $-3 < 1/x \Leftrightarrow -3x > 1 \Leftrightarrow x < -\frac{1}{3}$, so for this inequality, the solution set is $(-\infty, -\frac{1}{3}) \cup (0, \infty)$. Now $1/x \leq 1$ is clearly true if $x < 0$. So suppose $x > 0$. Then $1/x \leq 1 \Leftrightarrow 1 \leq x$, and the solution set here is $(-\infty, 0) \cup [1, \infty)$. Taking the intersection of the two solution sets gives the final solution set: $(-\infty, -\frac{1}{3}) \cup [1, \infty)$.



39. $C = \frac{5}{9}(F - 32) \Rightarrow F = \frac{9}{5}C + 32$. So $50 \leq F \leq 95 \Rightarrow 50 \leq \frac{9}{5}C + 32 \leq 95 \Rightarrow 18 \leq \frac{9}{5}C \leq 63 \Rightarrow 10 \leq C \leq 35$. So the interval is $[10, 35]$.

40. Since $20 \leq C \leq 30$ and $C = \frac{5}{9}(F - 32)$, we have $20 \leq \frac{5}{9}(F - 32) \leq 30 \Rightarrow 36 \leq F - 32 \leq 54 \Rightarrow 68 \leq F \leq 86$. So the interval is $[68, 86]$.

41. (a) Let T represent the temperature in degrees Celsius and h the height in km. $T = 20$ when $h = 0$ and T decreases by 10°C for every km (1°C for each 100-m rise). Thus, $T = 20 - 10h$ when $0 \leq h \leq 12$.

(b) From part (a), $T = 20 - 10h \Rightarrow 10h = 20 - T \Rightarrow h = 2 - T/10$. So $0 \leq h \leq 5 \Rightarrow 0 \leq 2 - T/10 \leq 5 \Rightarrow -2 \leq -T/10 \leq 3 \Rightarrow -20 \leq -T \leq 30 \Rightarrow 20 \geq T \geq -30 \Rightarrow -30 \leq T \leq 20$. Thus, the range of temperatures (in $^\circ\text{C}$) to be expected is $[-30, 20]$.

42. The ball will be at least 32 ft above the ground if $h \geq 32 \Leftrightarrow 128 + 16t - 16t^2 \geq 32 \Leftrightarrow 16t^2 - 16t - 96 \leq 0 \Leftrightarrow 16(t-3)(t+2) \leq 0$. $t = 3$ and $t = -2$ are endpoints of the interval we're looking for, and constructing a table gives $-2 \leq t \leq 3$. But $t \geq 0$, so the ball will be at least 32 ft above the ground in the time interval $[0, 3]$.

43. $|2x| = 3 \Leftrightarrow$ either $2x = 3$ or $2x = -3 \Leftrightarrow x = \frac{3}{2}$ or $x = -\frac{3}{2}$.

44. $|3x + 5| = 1 \Leftrightarrow$ either $3x + 5 = 1$ or -1 . In the first case, $3x = -4 \Leftrightarrow x = -\frac{4}{3}$, and in the second case, $3x = -6 \Leftrightarrow x = -2$. So the solutions are -2 and $-\frac{4}{3}$.

45. $|x + 3| = |2x + 1| \Leftrightarrow$ either $x + 3 = 2x + 1$ or $x + 3 = -(2x + 1)$. In the first case, $x = 2$, and in the second case, $x + 3 = -2x - 1 \Leftrightarrow 3x = -4 \Leftrightarrow x = -\frac{4}{3}$. So the solutions are $-\frac{4}{3}$ and 2 .

46. $\left| \frac{2x-1}{x+1} \right| = 3 \Leftrightarrow$ either $\frac{2x-1}{x+1} = 3$ or $\frac{2x-1}{x+1} = -3$. In the first case, $2x-1 = 3x+3 \Leftrightarrow x = -4$, and in the second case, $2x-1 = -3x-3 \Leftrightarrow x = -\frac{2}{5}$.
47. By Property 5 of absolute values, $|x| < 3 \Leftrightarrow -3 < x < 3$, so $x \in (-3, 3)$.
48. By Properties 4 and 6 of absolute values, $|x| \geq 3 \Leftrightarrow x \leq -3$ or $x \geq 3$, so $x \in (-\infty, -3] \cup [3, \infty)$.
49. $|x-4| < 1 \Leftrightarrow -1 < x-4 < 1 \Leftrightarrow 3 < x < 5$, so $x \in (3, 5)$.
50. $|x-6| < 0.1 \Leftrightarrow -0.1 < x-6 < 0.1 \Leftrightarrow 5.9 < x < 6.1$, so $x \in (5.9, 6.1)$.
51. $|x+5| \geq 2 \Leftrightarrow x+5 \geq 2$ or $x+5 \leq -2 \Leftrightarrow x \geq -3$ or $x \leq -7$, so $x \in (-\infty, -7] \cup [-3, \infty)$.
52. $|x+1| \geq 3 \Leftrightarrow x+1 \geq 3$ or $x+1 \leq -3 \Leftrightarrow x \geq 2$ or $x \leq -4$, so $x \in (-\infty, -4] \cup [2, \infty)$.
53. $|2x-3| \leq 0.4 \Leftrightarrow -0.4 \leq 2x-3 \leq 0.4 \Leftrightarrow 2.6 \leq 2x \leq 3.4 \Leftrightarrow 1.3 \leq x \leq 1.7$, so $x \in [1.3, 1.7]$.
54. $|5x-2| < 6 \Leftrightarrow -6 < 5x-2 < 6 \Leftrightarrow -4 < 5x < 8 \Leftrightarrow -\frac{4}{5} < x < \frac{8}{5}$, so $x \in (-\frac{4}{5}, \frac{8}{5})$.
55. $1 \leq |x| \leq 4$. So either $1 \leq x \leq 4$ or $1 \leq -x \leq 4 \Leftrightarrow -1 \geq x \geq -4$. Thus, $x \in [-4, -1] \cup [1, 4]$.
56. $0 < |x-5| < \frac{1}{2}$. Clearly $0 < |x-5|$ for $x \neq 5$. Now $|x-5| < \frac{1}{2} \Leftrightarrow -\frac{1}{2} < x-5 < \frac{1}{2} \Leftrightarrow 4.5 < x < 5.5$. So the solution set is $(4.5, 5) \cup (5, 5.5)$.
57. $a(bx-c) \geq bc \Leftrightarrow bx-c \geq \frac{bc}{a} \Leftrightarrow bx \geq \frac{bc}{a} + c = \frac{bc+ac}{a} \Leftrightarrow x \geq \frac{bc+ac}{ab}$
58. $a \leq bx+c < 2a \Leftrightarrow a-c \leq bx < 2a-c \Leftrightarrow \frac{a-c}{b} \leq x < \frac{2a-c}{b}$ (since $b > 0$)
59. $ax+b < c \Leftrightarrow ax < c-b \Leftrightarrow x > \frac{c-b}{a}$ (since $a < 0$)
60. $\frac{ax+b}{c} \leq b \Leftrightarrow ax+b \geq bc$ (since $c < 0$) $\Leftrightarrow ax \geq bc-b \Leftrightarrow x \leq \frac{b(c-1)}{a}$ (since $a < 0$)
61. $|(x+y)-5| = |(x-2)+(y-3)| \leq |x-2| + |y-3| < 0.01 + 0.04 = 0.05$
62. Use the Triangle Inequality: $|x+3| < \frac{1}{2} \Rightarrow$
 $|4x+13| = |4(x+3)+1| \leq |4(x+3)| + |1| = 4|x+3| + 1 < 4(\frac{1}{2}) + 1 = 3$
Another method: $|x+3| < \frac{1}{2} \Rightarrow -\frac{1}{2} < x+3 < \frac{1}{2} \Rightarrow -2 < 4x+12 < 2 \Rightarrow -1 < 4x+13 < 3 \Rightarrow$
 $|4x+13| < 3$
63. If $a < b$ then $a+a < a+b$ and $a+b < b+b$. So $2a < a+b < 2b$. Dividing by 2, we get $a < \frac{1}{2}(a+b) < b$.
64. If $0 < a < b$, then $\frac{1}{ab} > 0$. So $a < b \Rightarrow \frac{1}{ab} \cdot a < \frac{1}{ab} \cdot b \Leftrightarrow \frac{1}{b} < \frac{1}{a}$.
65. $|ab| = \sqrt{(ab)^2} = \sqrt{a^2b^2} = \sqrt{a^2} \sqrt{b^2} = |a| |b|$
66. $\left| \frac{a}{b} \right| |b| = \left| \frac{a}{b} \cdot b \right| = |a|$ (using the result of Exercise 65). Dividing the equation through by $|b|$ gives $\left| \frac{a}{b} \right| = \frac{|a|}{|b|}$.
67. If $0 < a < b$, then $a \cdot a < a \cdot b$ and $a \cdot b < b \cdot b$ [using Rule 3 of Inequalities]. So $a^2 < ab < b^2$ and hence $a^2 < b^2$.
68. Following the hint, the Triangle Inequality becomes $|(x-y)+y| \leq |x-y| + |y| \Leftrightarrow |x| \leq |x-y| + |y| \Leftrightarrow$
 $|x-y| \geq |x| - |y|$.

69. Observe that the sum, difference and product of two integers is always an integer. Let the rational numbers be represented by $r = m/n$ and $s = p/q$ (where m, n, p and q are integers with $n \neq 0, q \neq 0$). Now

$$r + s = \frac{m}{n} + \frac{p}{q} = \frac{mq + pn}{nq}, \text{ but } mq + pn \text{ and } nq \text{ are both integers, so } \frac{mq + pn}{nq} = r + s \text{ is a rational number by}$$

definition. Similarly, $r - s = \frac{m}{n} - \frac{p}{q} = \frac{mq - pn}{nq}$ is a rational number. Finally, $r \cdot s = \frac{m}{n} \cdot \frac{p}{q} = \frac{mp}{nq}$ but mp and

nq are both integers, so $\frac{mp}{nq} = r \cdot s$ is a rational number by definition.

70. (a) No. Consider the case of $\sqrt{2}$ and $-\sqrt{2}$. Both are irrational numbers, yet $\sqrt{2} + (-\sqrt{2}) = 0$ and 0, being an integer, is not irrational.
- (b) No. Consider the case of $\sqrt{2}$ and $\sqrt{2}$. Both are irrational numbers, yet $\sqrt{2} \cdot \sqrt{2} = 2$ is not irrational.

B Coordinate Geometry and Lines

1. From the Distance Formula with $x_1 = 1, x_2 = 4, y_1 = 1, y_2 = 5$, we find the distance from $(1, 1)$ to $(4, 5)$ to be

$$\sqrt{(4-1)^2 + (5-1)^2} = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$$

2. The distance from $(1, -3)$ to $(5, 7)$ is $\sqrt{(5-1)^2 + [7-(-3)]^2} = \sqrt{4^2 + 10^2} = \sqrt{116} = 2\sqrt{29}$.

3. $\sqrt{(-1-6)^2 + [3-(-2)]^2} = \sqrt{(-7)^2 + 5^2} = \sqrt{74}$

4. $\sqrt{(-1-1)^2 + [-3-(-6)]^2} = \sqrt{(-2)^2 + 3^2} = \sqrt{13}$

5. $\sqrt{(4-2)^2 + (-7-5)^2} = \sqrt{2^2 + (-12)^2} = \sqrt{148} = 2\sqrt{37}$

6. $\sqrt{(b-a)^2 + (a-b)^2} = \sqrt{(a-b)^2 + (a-b)^2} = \sqrt{2(a-b)^2} = \sqrt{2}|a-b|$

7. From (2), the slope is $\frac{11-5}{4-1} = \frac{6}{3} = 2$.

8. $m = \frac{-3-6}{4-(-1)} = -\frac{9}{5}$

9. With $P(-3, 3)$ and $Q(-1, -6)$, the slope m of the line through P and Q is $m = \frac{-6-3}{-1-(-3)} = -\frac{9}{2}$.

10. $m = \frac{0-(-4)}{6-(-1)} = \frac{4}{7}$

11. Since $|AC| = \sqrt{(-4-0)^2 + (3-2)^2} = \sqrt{(-4)^2 + 1^2} = \sqrt{17}$ and

$|BC| = \sqrt{[-4-(-3)]^2 + [3-(-1)]^2} = \sqrt{(-1)^2 + 4^2} = \sqrt{17}$, the triangle has two sides of equal length, and so is isosceles.

12. (a) $|AB| = \sqrt{(11-6)^2 + [-3-(-7)]^2} = \sqrt{5^2 + 4^2} = \sqrt{41}$,

$|AC| = \sqrt{(2-6)^2 + [-2-(-7)]^2} = \sqrt{(-4)^2 + 5^2} = \sqrt{41}$, and

$|BC| = \sqrt{(2-11)^2 + [-2-(-3)]^2} = \sqrt{(-9)^2 + 1^2} = \sqrt{82}$, so

$|AB|^2 + |AC|^2 = 41 + 41 = 82 = |BC|^2$, and so $\triangle ABC$ is a right triangle.

(b) $m_{AB} = \frac{-3 - (-7)}{11 - 6} = \frac{4}{5}$ and $m_{AC} = \frac{-2 - (-7)}{2 - 6} = -\frac{5}{4}$. Thus $m_{AB} \cdot m_{AC} = -1$ and so AB is perpendicular to AC and $\triangle ABC$ must be a right triangle.

(c) Taking lengths from part (a), the base is $\sqrt{41}$ and the height is $\sqrt{41}$. Thus the area is $\frac{1}{2}bh = \frac{1}{2}\sqrt{41}\sqrt{41} = \frac{41}{2}$.

13. Using $A(-2, 9)$, $B(4, 6)$, $C(1, 0)$, and $D(-5, 3)$, we have

$$|AB| = \sqrt{[4 - (-2)]^2 + (6 - 9)^2} = \sqrt{6^2 + (-3)^2} = \sqrt{45} = \sqrt{9} \sqrt{5} = 3\sqrt{5},$$

$$|BC| = \sqrt{(1 - 4)^2 + (0 - 6)^2} = \sqrt{(-3)^2 + (-6)^2} = \sqrt{45} = \sqrt{9} \sqrt{5} = 3\sqrt{5},$$

$$|CD| = \sqrt{(-5 - 1)^2 + (3 - 0)^2} = \sqrt{(-6)^2 + 3^2} = \sqrt{45} = \sqrt{9} \sqrt{5} = 3\sqrt{5}, \text{ and}$$

$$|DA| = \sqrt{[-2 - (-5)]^2 + (9 - 3)^2} = \sqrt{3^2 + 6^2} = \sqrt{45} = \sqrt{9} \sqrt{5} = 3\sqrt{5}. \text{ So all sides are of equal length and}$$

we have a rhombus. Moreover, $m_{AB} = \frac{6 - 9}{4 - (-2)} = -\frac{1}{2}$, $m_{BC} = \frac{0 - 6}{1 - 4} = 2$, $m_{CD} = \frac{3 - 0}{-5 - 1} = -\frac{1}{2}$, and

$m_{DA} = \frac{9 - 3}{-2 - (-5)} = 2$, so the sides are perpendicular. Thus, A , B , C , and D are vertices of a square.

14. (a) Using $A(-1, 3)$, $B(3, 11)$, and $C(5, 15)$, we have

$$|AB| = \sqrt{[3 - (-1)]^2 + (11 - 3)^2} = \sqrt{4^2 + 8^2} = \sqrt{80} = 4\sqrt{5},$$

$$|BC| = \sqrt{(5 - 3)^2 + (15 - 11)^2} = \sqrt{2^2 + 4^2} = \sqrt{20} = 2\sqrt{5}, \text{ and}$$

$$|AC| = \sqrt{[5 - (-1)]^2 + (15 - 3)^2} = \sqrt{6^2 + 12^2} = \sqrt{180} = 6\sqrt{5}. \text{ Thus, } |AC| = |AB| + |BC|.$$


(b) $m_{AB} = \frac{11 - 3}{3 - (-1)} = \frac{8}{4} = 2$ and $m_{AC} = \frac{15 - 3}{5 - (-1)} = \frac{12}{6} = 2$. Since the segments AB and AC have the same slope, A , B and C must be collinear.

15. The slope of the line segment AB is $\frac{4 - 1}{7 - 1} = \frac{1}{2}$, the slope of CD is $\frac{7 - 10}{-1 - 5} = \frac{1}{2}$, the slope of BC is $\frac{10 - 4}{5 - 7} = -3$, and the slope of DA is $\frac{1 - 7}{1 - (-1)} = -3$. So AB is parallel to CD and BC is parallel to DA .


Hence $ABCD$ is a parallelogram.

16. The slopes of the four sides are $m_{AB} = \frac{3 - 1}{11 - 1} = \frac{1}{5}$, $m_{BC} = \frac{8 - 3}{10 - 11} = -5$, $m_{CD} = \frac{6 - 8}{0 - 10} = \frac{1}{5}$, and $m_{DA} = \frac{1 - 6}{1 - 0} = -5$. Hence $AB \parallel CD$, $BC \parallel DA$, $AB \perp BC$, $BC \perp CD$, $CD \perp DA$, and $DA \perp AB$, and so $ABCD$ is a rectangle.


17. $x = 3$




18. $y = -2$



19. $xy = 0 \Leftrightarrow x = 0$ or $y = 0$. The graph consists of the coordinate axes.




20. $|y| = 1 \Leftrightarrow y = 1$ or $y = -1$




21. By the point-slope form of the equation of a line, an equation of the line through $(2, -3)$ with slope 6 is $y - (-3) = 6(x - 2)$ or $y = 6x - 15$.
22. $y - 4 = -3[x - (-1)]$ or $y = -3x + 1$
23. $y - 7 = \frac{2}{3}(x - 1)$ or $y = \frac{2}{3}x + \frac{19}{3}$
24. $y - (-5) = -\frac{7}{2}[x - (-3)]$ or $y = -\frac{7}{2}x - \frac{31}{2}$
25. The slope of the line through $(2, 1)$ and $(1, 6)$ is $m = \frac{6 - 1}{1 - 2} = -5$, so an equation of the line is $y - 1 = -5(x - 2)$ or $y = -5x + 11$.
26. For $(-1, -2)$ and $(4, 3)$, $m = \frac{3 - (-2)}{4 - (-1)} = 1$. So $y - 3 = 1(x - 4)$ or $y = x - 1$.
27. By the slope-intercept form of the equation of a line, an equation of the line is $y = 3x - 2$.
28. By the slope-intercept form of the equation of a line, an equation of the line is $y = \frac{2}{5}x + 4$.
29. Since the line passes through $(1, 0)$ and $(0, -3)$, its slope is $m = \frac{-3 - 0}{0 - 1} = 3$, so an equation is $y = 3x - 3$.
Another method: From Exercise 61, $\frac{x}{1} + \frac{y}{-3} = 1 \Rightarrow -3x + y = -3 \Rightarrow y = 3x - 3$.
30. For $(-8, 0)$ and $(0, 6)$, $m = \frac{6 - 0}{0 - (-8)} = \frac{3}{4}$. So an equation is $y = \frac{3}{4}x + 6$.
Another method: From Exercise 61, $\frac{x}{-8} + \frac{y}{6} = 1 \Rightarrow -3x + 4y = 24 \Rightarrow y = \frac{3}{4}x + 6$.
31. Since $m = 0$, $y - 5 = 0(x - 4)$ or $y = 5$.
32. Since m is undefined, we have the vertical line $x = 4$.
33. Putting the line $x + 2y = 6$ into its slope-intercept form gives us $y = -\frac{1}{2}x + 3$, so we see that this line has slope $-\frac{1}{2}$. Thus, we want the line of slope $-\frac{1}{2}$ that passes through the point $(1, -6)$: $y - (-6) = -\frac{1}{2}(x - 1) \Leftrightarrow y = -\frac{1}{2}x - \frac{11}{2}$.
34. $2x + 3y + 4 = 0 \Leftrightarrow y = -\frac{2}{3}x - \frac{4}{3}$, so $m = -\frac{2}{3}$ and the required line is $y = -\frac{2}{3}x + 6$.
35. $2x + 5y + 8 = 0 \Leftrightarrow y = -\frac{2}{5}x - \frac{8}{5}$. Since this line has slope $-\frac{2}{5}$, a line perpendicular to it would have slope $\frac{5}{2}$, so the required line is $y - (-2) = \frac{5}{2}[x - (-1)] \Leftrightarrow y = \frac{5}{2}x + \frac{1}{2}$.
36. $4x - 8y = 1 \Leftrightarrow y = \frac{1}{2}x - \frac{1}{8}$. Since this line has slope $\frac{1}{2}$, a line perpendicular to it would have slope -2 , so the required line is $y - (-\frac{2}{3}) = -2(x - \frac{1}{2}) \Leftrightarrow y = -2x + \frac{1}{3}$.


37. $x + 3y = 0 \Leftrightarrow y = -\frac{1}{3}x$,
so the slope is $-\frac{1}{3}$ and the
 y -intercept is 0.




38. $2x - 5y = 0 \Leftrightarrow y = \frac{2}{5}x$, so
the slope is $\frac{2}{5}$ and the y -intercept
is 0.




39. $y = -2$ is a horizontal line with
slope 0 and y -intercept -2 .




40. $2x - 3y + 6 = 0 \Leftrightarrow$
 $y = \frac{2}{3}x + 2$, so the slope is $\frac{2}{3}$
and the y -intercept is 2.




41. $3x - 4y = 12 \Leftrightarrow$
 $y = \frac{3}{4}x - 3$, so the slope is $\frac{3}{4}$
and the y -intercept is -3 .




42. $4x + 5y = 10 \Leftrightarrow$
 $y = -\frac{4}{5}x + 2$, so the slope is
 $-\frac{4}{5}$ and the y -intercept is 2.




43. $\{(x, y) \mid x < 0\}$




44. $\{(x, y) \mid y > 0\}$



45. $\{(x, y) \mid xy < 0\} =$
 $\{(x, y) \mid x < 0 \text{ and } y > 0\}$
 $\cup \{(x, y) \mid x > 0 \text{ and } y < 0\}$




46. $\{(x, y) \mid x \geq 1 \text{ and } y < 3\}$




47. $\{(x, y) \mid |x| \leq 2\} =$


$\{(x, y) \mid -2 \leq x \leq 2\}$




48. $\{(x, y) \mid |x| < 3 \text{ and } |y| < 2\}$




49. $\{(x, y) \mid 0 \leq y \leq 4, x \leq 2\}$




50. $\{(x, y) \mid y > 2x - 1\}$



51. $\{(x, y) \mid 1 + x \leq y \leq 1 - 2x\}$



52. $\{(x, y) \mid -x \leq y < \frac{x+3}{2}\}$



53. Let $P(0, y)$ be a point on the y -axis. The distance from P to $(5, -5)$

is $\sqrt{(5-0)^2 + (-5-y)^2} = \sqrt{5^2 + (y+5)^2}$. The distance from P

to $(1, 1)$ is $\sqrt{(1-0)^2 + (1-y)^2} = \sqrt{1^2 + (y-1)^2}$. We want

these distances to be equal: $\sqrt{5^2 + (y+5)^2} = \sqrt{1^2 + (y-1)^2} \Leftrightarrow$

$$5^2 + (y+5)^2 = 1^2 + (y-1)^2 \Leftrightarrow$$

$$25 + (y^2 + 10y + 25) = 1 + (y^2 - 2y + 1) \Leftrightarrow 12y = -48 \Leftrightarrow$$

$y = -4$. So the desired point is $(0, -4)$.

54. Let M be the point $(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2})$. Then

$$|MP_1|^2 = \left(x_1 - \frac{x_1 + x_2}{2}\right)^2 + \left(y_1 - \frac{y_1 + y_2}{2}\right)^2 = \left(\frac{x_1 - x_2}{2}\right)^2 + \left(\frac{y_1 - y_2}{2}\right)^2 \text{ and}$$

$|MP_2|^2 = \left(x_2 - \frac{x_1 + x_2}{2}\right)^2 + \left(y_2 - \frac{y_1 + y_2}{2}\right)^2 = \left(\frac{x_2 - x_1}{2}\right)^2 + \left(\frac{y_2 - y_1}{2}\right)^2$. Hence, $|MP_1| = |MP_2|$; that is, M is equidistant from P_1 and P_2 .

55. (a) Using the midpoint formula from Exercise 54 with $(1, 3)$ and $(7, 15)$, we get $(\frac{1+7}{2}, \frac{3+15}{2}) = (4, 9)$.

(b) Using the same formula, we get $(\frac{-1+8}{2}, \frac{6-12}{2}) = (\frac{7}{2}, -3)$.


56. The midpoint M_1 of AB is $(\frac{1+3}{2}, \frac{0+6}{2}) = (2, 3)$, the midpoint M_2 of BC is $(\frac{3+8}{2}, \frac{6+2}{2}) = (\frac{11}{2}, 4)$, and the midpoint M_3 of CA is $(\frac{8+1}{2}, \frac{2+0}{2}) = (\frac{9}{2}, 1)$. The lengths of the medians are $|AM_2| = \sqrt{(\frac{11}{2} - 1)^2 + (4 - 0)^2} = \sqrt{(\frac{9}{2})^2 + 4^2} = \sqrt{\frac{145}{4}} = \frac{\sqrt{145}}{2}$,
 $|BM_3| = \sqrt{(\frac{9}{2} - 3)^2 + (1 - 6)^2} = \sqrt{(\frac{3}{2})^2 + (-5)^2} = \sqrt{\frac{109}{4}} = \frac{\sqrt{109}}{2}$, and
 $|CM_1| = \sqrt{(2 - 8)^2 + (3 - 2)^2} = \sqrt{(-6)^2 + 1^2} = \sqrt{37}$.
57. $2x - y = 4 \Leftrightarrow y = 2x - 4 \Rightarrow m_1 = 2$ and $6x - 2y = 10 \Leftrightarrow 2y = 6x - 10 \Leftrightarrow y = 3x - 5 \Rightarrow m_2 = 3$. Since $m_1 \neq m_2$, the two lines are not parallel. To find the point of intersection: $2x - 4 = 3x - 5 \Leftrightarrow x = 1 \Rightarrow y = -2$. Thus, the point of intersection is $(1, -2)$.
58. $3x - 5y + 19 = 0 \Leftrightarrow 5y = 3x + 19 \Leftrightarrow y = \frac{3}{5}x + \frac{19}{5} \Rightarrow m_1 = \frac{3}{5}$ and $10x + 6y - 50 = 0 \Leftrightarrow 6y = -10x + 50 \Leftrightarrow y = -\frac{5}{3}x + \frac{25}{3} \Rightarrow m_2 = -\frac{5}{3}$. Since $m_1 m_2 = \frac{3}{5}(-\frac{5}{3}) = -1$, the two lines are perpendicular. To find the point of intersection: $\frac{3}{5}x + \frac{19}{5} = -\frac{5}{3}x + \frac{25}{3} \Leftrightarrow 9x + 57 = -25x + 125 \Leftrightarrow 34x = 68 \Leftrightarrow x = 2 \Rightarrow y = \frac{3}{5} \cdot 2 + \frac{19}{5} = \frac{25}{5} = 5$. Thus, the point of intersection is $(2, 5)$.
59. With $A(1, 4)$ and $B(7, -2)$, the slope of segment AB is $\frac{-2-4}{7-1} = -1$, so its perpendicular bisector has slope 1. The midpoint of AB is $(\frac{1+7}{2}, \frac{4+(-2)}{2}) = (4, 1)$, so an equation of the perpendicular bisector is $y - 1 = 1(x - 4)$ or $y = x - 3$.
60. (a) Side PQ has slope $\frac{4-0}{3-1} = 2$, so its equation is $y - 0 = 2(x - 1) \Leftrightarrow y = 2x - 2$. Side QR has slope $\frac{6-4}{-1-3} = -\frac{1}{2}$, so its equation is $y - 4 = -\frac{1}{2}(x - 3) \Leftrightarrow y = -\frac{1}{2}x + \frac{11}{2}$. Side RP has slope $\frac{0-6}{1-(-1)} = -3$, so its equation is $y - 0 = -3(x - 1) \Leftrightarrow y = -3x + 3$.
- (b) M_1 (the midpoint of PQ) has coordinates $(\frac{1+3}{2}, \frac{0+4}{2}) = (2, 2)$. M_2 (the midpoint of QR) has coordinates $(\frac{3-1}{2}, \frac{4+6}{2}) = (1, 5)$. M_3 (the midpoint of RP) has coordinates $(\frac{1-1}{2}, \frac{0+6}{2}) = (0, 3)$. RM_1 has slope $\frac{2-6}{2-(-1)} = -\frac{4}{3}$ and hence equation $y - 2 = -\frac{4}{3}(x - 2) \Leftrightarrow y = -\frac{4}{3}x + \frac{14}{3}$. PM_2 is a vertical line with equation $x = 1$. QM_3 has slope $\frac{3-4}{0-3} = \frac{1}{3}$ and hence equation $y - 3 = \frac{1}{3}(x - 0) \Leftrightarrow y = \frac{1}{3}x + 3$. PM_2 and RM_1 intersect where $x = 1$ and $y = -\frac{4}{3}(1) + \frac{14}{3} = \frac{10}{3}$, or at $(1, \frac{10}{3})$. PM_2 and QM_3 intersect where $x = 1$ and $y = \frac{1}{3}(1) + 3 = \frac{10}{3}$, or at $(1, \frac{10}{3})$, so this is the point where all three medians intersect.
61. (a) Since the x -intercept is a , the point $(a, 0)$ is on the line, and similarly since the y -intercept is b , $(0, b)$ is on the line. Hence, the slope of the line is $m = \frac{b-0}{0-a} = -\frac{b}{a}$. Substituting into $y = mx + b$ gives $y = -\frac{b}{a}x + b \Leftrightarrow \frac{b}{a}x + y = b \Leftrightarrow \frac{x}{a} + \frac{y}{b} = 1$.
- (b) Letting $a = 6$ and $b = -8$ gives $\frac{x}{6} + \frac{y}{-8} = 1 \Leftrightarrow -8x + 6y = -48$ [multiply by -48] $\Leftrightarrow 6y = 8x - 48 \Leftrightarrow 3y = 4x - 24 \Leftrightarrow y = \frac{4}{3}x - 8$.

62. (a) Let
- d
- = distance traveled (in miles) and
- t
- = time elapsed (in hours). (b)

At $t = 0$, $d = 0$ and at $t = 50$ minutes $= 50 \cdot \frac{1}{60} = \frac{5}{6}$ h, $d = 40$.

Thus, we have two points: $(0, 0)$ and $(\frac{5}{6}, 40)$, so $m = \frac{40 - 0}{5/6 - 0} = 48$

and $d = 48t$.




- (c) The slope is 48 and represents the car's speed in mi/h.


C Graphs of Second-Degree Equations

- An equation of the circle with center $(3, -1)$ and radius 5 is $(x - 3)^2 + (y + 1)^2 = 5^2 = 25$.
- From (1), the equation is $(x + 2)^2 + (y + 8)^2 = 100$.
- The equation has the form $x^2 + y^2 = r^2$. Since $(4, 7)$ lies on the circle, we have $4^2 + 7^2 = r^2 \Rightarrow r^2 = 65$. So the required equation is $x^2 + y^2 = 65$.
- The equation has the form $(x + 1)^2 + (y - 5)^2 = r^2$. Since $(-4, -6)$ lies on the circle, we have $r^2 = (-4 + 1)^2 + (-6 - 5)^2 = 130$. So an equation is $(x + 1)^2 + (y - 5)^2 = 130$.
- $x^2 + y^2 - 4x + 10y + 13 = 0 \Leftrightarrow x^2 - 4x + y^2 + 10y = -13 \Leftrightarrow (x^2 - 4x + 4) + (y^2 + 10y + 25) = -13 + 4 + 25 = 16 \Leftrightarrow (x - 2)^2 + (y + 5)^2 = 4^2$. Thus, we have a circle with center $(2, -5)$ and radius 4.
- $x^2 + y^2 + 6y + 2 = 0 \Leftrightarrow x^2 + (y^2 + 6y + 9) = -2 + 9 \Leftrightarrow x^2 + (y + 3)^2 = 7$. Thus, we have a circle with center $(0, -3)$ and radius $\sqrt{7}$.
- $x^2 + y^2 + x = 0 \Leftrightarrow (x^2 + x + \frac{1}{4}) + y^2 = \frac{1}{4} \Leftrightarrow (x + \frac{1}{2})^2 + y^2 = (\frac{1}{2})^2$. Thus, we have a circle with center $(-\frac{1}{2}, 0)$ and radius $\frac{1}{2}$.
- $16x^2 + 16y^2 + 8x + 32y + 1 = 0 \Leftrightarrow 16(x^2 + \frac{1}{2}x + \frac{1}{16}) + 16(y^2 + 2y + 1) = -1 + 1 + 16 \Leftrightarrow 16(x + \frac{1}{4})^2 + 16(y + 1)^2 = 16 \Leftrightarrow (x + \frac{1}{4})^2 + (y + 1)^2 = 1$. Thus, we have a circle with center $(-\frac{1}{4}, -1)$ and radius 1.
- $2x^2 + 2y^2 - x + y = 1 \Leftrightarrow 2(x^2 - \frac{1}{2}x + \frac{1}{16}) + 2(y^2 + \frac{1}{2}y + \frac{1}{16}) = 1 + \frac{1}{8} + \frac{1}{8} \Leftrightarrow 2(x - \frac{1}{4})^2 + 2(y + \frac{1}{4})^2 = \frac{5}{4} \Leftrightarrow (x - \frac{1}{4})^2 + (y + \frac{1}{4})^2 = \frac{5}{8}$. Thus, we have a circle with center $(\frac{1}{4}, -\frac{1}{4})$ and radius $\frac{\sqrt{5}}{2\sqrt{2}} = \frac{\sqrt{10}}{4}$.
- $x^2 + y^2 + ax + by + c = 0 \Leftrightarrow (x^2 + ax + \frac{1}{4}a^2) + (y^2 + by + \frac{1}{4}b^2) = -c + \frac{1}{4}a^2 + \frac{1}{4}b^2 \Leftrightarrow (x + \frac{1}{2}a)^2 + (y + \frac{1}{2}b)^2 = \frac{1}{4}(a^2 + b^2 - 4c)$. For this to represent a nondegenerate circle, $\frac{1}{4}(a^2 + b^2 - 4c) > 0$ or $a^2 + b^2 > 4c$. If this condition is satisfied, the circle has center $(-\frac{1}{2}a, -\frac{1}{2}b)$ and radius $\frac{1}{2}\sqrt{a^2 + b^2 - 4c}$.


11. $y = -x^2$. Parabola




12. $y^2 - x^2 = 1$. Hyperbola



13. $x^2 + 4y^2 = 16 \Leftrightarrow \frac{x^2}{16} + \frac{y^2}{4} = 1$. Ellipse




14. $x = -2y^2$. Parabola




15. $16x^2 - 25y^2 = 400 \Leftrightarrow \frac{x^2}{25} - \frac{y^2}{16} = 1$.


Hyperbola




16. $25x^2 + 4y^2 = 100 \Leftrightarrow \frac{x^2}{4} + \frac{y^2}{25} = 1$. Ellipse




17. $4x^2 + y^2 = 1 \Leftrightarrow \frac{x^2}{1/4} + y^2 = 1$. Ellipse



18. $y = x^2 + 2$. Parabola with vertex at (0, 2)




19. $x = y^2 - 1$. Parabola with vertex at $(-1, 0)$




20. $9x^2 - 25y^2 = 225 \Leftrightarrow \frac{x^2}{25} - \frac{y^2}{9} = 1$.


Hyperbola




21. $9y^2 - x^2 = 9 \Leftrightarrow y^2 - \frac{x^2}{9} = 1$. Hyperbola




22. $2x^2 + 5y^2 = 10 \Leftrightarrow \frac{x^2}{5} + \frac{y^2}{2} = 1$. Ellipse



23. $xy = 4$. Hyperbola




24. $y = x^2 + 2x = (x^2 + 2x + 1) - 1 = (x + 1)^2 - 1$.

 Parabola with vertex at $(-1, -1)$


25. $9(x - 1)^2 + 4(y - 2)^2 = 36 \Leftrightarrow$


$$\frac{(x - 1)^2}{4} + \frac{(y - 2)^2}{9} = 1$$
. Ellipse centered at $(1, 2)$




26. $16x^2 + 9y^2 - 36y = 108 \Leftrightarrow$

$$16x^2 + 9(y^2 - 4y + 4) = 108 + 36 = 144 \Leftrightarrow$$


$$\frac{x^2}{9} + \frac{(y - 2)^2}{16} = 1$$
. Ellipse centered at $(0, 2)$




27. $y = x^2 - 6x + 13 = (x^2 - 6x + 9) + 4 = (x - 3)^2 + 4$. Parabola with vertex at $(3, 4)$



29. $x = -y^2 + 4$. Parabola with vertex at $(4, 0)$




31. $x^2 + 4y^2 - 6x + 5 = 0 \Leftrightarrow (x^2 - 6x + 9) + 4y^2 = -5 + 9 = 4 \Leftrightarrow \frac{(x - 3)^2}{4} + y^2 = 1$. Ellipse centered at $(3, 0)$




33. $y = 3x$ and $y = x^2$ intersect where $3x = x^2 \Leftrightarrow 0 = x^2 - 3x = x(x - 3)$, that is, at $(0, 0)$ and $(3, 9)$.



28. $x^2 - y^2 - 4x + 3 = 0 \Leftrightarrow (x^2 - 4x + 4) - y^2 = -3 + 4 = 1 \Leftrightarrow (x - 2)^2 - y^2 = 1$. Hyperbola centered at $(2, 0)$




30. $y^2 - 2x + 6y + 5 = 0 \Leftrightarrow y^2 + 6y + 9 = 2x + 4 \Leftrightarrow (y + 3)^2 = 2(x + 2)$. Parabola with vertex $(-2, -3)$



32. $4x^2 + 9y^2 - 16x + 54y + 61 = 0 \Leftrightarrow 4(x^2 - 4x + 4) + 9(y^2 + 6y + 9) = -61 + 16 + 81 = 36 \Leftrightarrow \frac{(x - 2)^2}{9} + \frac{(y + 3)^2}{4} = 1$. Ellipse centered at $(2, -3)$



34.



$y = 4 - x^2$, $x - 2y = 2$. Substitute y from the first equation into the second: $x - 2(4 - x^2) = 2 \Leftrightarrow 2x^2 + x - 10 = 0$
 $\Leftrightarrow (2x + 5)(x - 2) = 0 \Leftrightarrow x = -\frac{5}{2}$ or 2 . So the points of intersection are $(-\frac{5}{2}, -\frac{9}{4})$ and $(2, 0)$.

35. The parabola must have an equation of the form $y = a(x - 1)^2 - 1$. Substituting $x = 3$ and $y = 3$ into the equation gives $3 = a(3 - 1)^2 - 1$, so $a = 1$, and the equation is $y = (x - 1)^2 - 1 = x^2 - 2x$. Note that using the other point $(-1, 3)$ would have given the same value for a , and hence the same equation.

36. The ellipse has an equation of the form $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$. Substituting $x = 1$ and $y = -\frac{10\sqrt{2}}{3}$ gives


$$\frac{1^2}{a^2} + \frac{(-10\sqrt{2}/3)^2}{b^2} = \frac{1}{a^2} + \frac{200}{9b^2} = 1. \text{ Substituting } x = -2 \text{ and } y = \frac{5\sqrt{5}}{3} \text{ gives}$$

$$\frac{(-2)^2}{a^2} + \frac{(5\sqrt{5}/3)^2}{b^2} = \frac{4}{a^2} + \frac{125}{9b^2} = 1. \text{ From the first equation, } \frac{1}{a^2} = 1 - \frac{200}{9b^2}. \text{ Putting this into the second}$$


$$\text{equation gives } 4\left(1 - \frac{200}{9b^2}\right) + \frac{125}{9b^2} = 1 \Leftrightarrow 3 = \frac{675}{9b^2} \Leftrightarrow b^2 = \frac{675}{27} = 25, \text{ so } b = 5. \text{ Hence}$$

$$\frac{1}{a^2} = 1 - \frac{200}{9(5)^2} = \frac{1}{9} \text{ and so } a = 3. \text{ The equation of the ellipse is } \frac{x^2}{9} + \frac{y^2}{25} = 1.$$


37. $\{(x, y) \mid x^2 + y^2 \leq 1\}$




38. $\{(x, y) \mid x^2 + y^2 > 4\}$



39. $\{(x, y) \mid y \geq x^2 - 1\}$




40. $\{(x, y) \mid x^2 + 4y^2 \leq 4\}$




D Trigonometry

1. $210^\circ = 210\left(\frac{\pi}{180}\right) = \frac{7\pi}{6}$ rad
2. $300^\circ = 300\left(\frac{\pi}{180}\right) = \frac{5\pi}{3}$ rad
3. $9^\circ = 9\left(\frac{\pi}{180}\right) = \frac{\pi}{20}$ rad
4. $-315^\circ = -315\left(\frac{\pi}{180}\right) = -\frac{7\pi}{4}$ rad
5. $900^\circ = 900\left(\frac{\pi}{180}\right) = 5\pi$ rad
6. $36^\circ = 36\left(\frac{\pi}{180}\right) = \frac{\pi}{5}$ rad
7. 4π rad $= 4\pi\left(\frac{180}{\pi}\right) = 720^\circ$
8. $-\frac{7\pi}{2}$ rad $= -\frac{7\pi}{2}\left(\frac{180}{\pi}\right) = -630^\circ$
9. $\frac{5\pi}{12}$ rad $= \frac{5\pi}{12}\left(\frac{180}{\pi}\right) = 75^\circ$
10. $\frac{8\pi}{3}$ rad $= \frac{8\pi}{3}\left(\frac{180}{\pi}\right) = 480^\circ$
11. $-\frac{3\pi}{8}$ rad $= -\frac{3\pi}{8}\left(\frac{180}{\pi}\right) = -67.5^\circ$
12. 5 rad $= 5\left(\frac{180}{\pi}\right) = \left(\frac{900}{\pi}\right)^\circ$
13. Using Formula 3, $a = r\theta = 36 \cdot \frac{\pi}{12} = 3\pi$ cm.
14. Using Formula 3, $a = r\theta = 10 \cdot 72\left(\frac{\pi}{180}\right) = 4\pi$ cm.
15. Using Formula 3, $\theta = a/r = \frac{1}{1.5} = \frac{2}{3}$ rad $= \frac{2}{3}\left(\frac{180}{\pi}\right) = \left(\frac{120}{\pi}\right)^\circ \approx 38.2^\circ$.
16. $a = r\theta \Rightarrow r = \frac{a}{\theta} = \frac{6}{3\pi/4} = \frac{8}{\pi}$ cm


17.




18.




19.




20.




21.



22.



23.




From the diagram we see that a point on the terminal side is $P(-1, 1)$.

Therefore, taking $x = -1$, $y = 1$, $r = \sqrt{2}$ in the definitions of the trigonometric ratios, we have $\sin \frac{3\pi}{4} = \frac{1}{\sqrt{2}}$, $\cos \frac{3\pi}{4} = -\frac{1}{\sqrt{2}}$,

$\tan \frac{3\pi}{4} = -1$, $\csc \frac{3\pi}{4} = \sqrt{2}$, $\sec \frac{3\pi}{4} = -\sqrt{2}$, and $\cot \frac{3\pi}{4} = -1$.

24.




From the diagram and Figure 8, we see that a point on the terminal side is $P(-1, -\sqrt{3})$. Therefore, taking $x = -1$, $y = -\sqrt{3}$, $r = 2$ in the

definitions of the trigonometric ratios, we have $\sin \frac{4\pi}{3} = -\frac{\sqrt{3}}{2}$,

$\cos \frac{4\pi}{3} = -\frac{1}{2}$, $\tan \frac{4\pi}{3} = \sqrt{3}$, $\csc \frac{4\pi}{3} = -\frac{2}{\sqrt{3}}$, $\sec \frac{4\pi}{3} = -2$, and

$\cot \frac{4\pi}{3} = \frac{1}{\sqrt{3}}$.


25.



From the diagram we see that a point on the terminal line is $P(0, 1)$.

Therefore taking $x = 0$, $y = 1$, $r = 1$ in the definitions of the trigonometric ratios, we have $\sin \frac{9\pi}{2} = 1$, $\cos \frac{9\pi}{2} = 0$, $\tan \frac{9\pi}{2} = y/x$ is undefined since $x = 0$, $\csc \frac{9\pi}{2} = 1$, $\sec \frac{9\pi}{2} = r/x$ is undefined since $x = 0$, and $\cot \frac{9\pi}{2} = 0$.

26.




From the diagram, we see that a point on the terminal line is $P(-1, 0)$.

Therefore taking $x = -1$, $y = 0$, $r = 1$ in the definitions of the trigonometric ratios we have $\sin(-5\pi) = 0$, $\cos(-5\pi) = -1$,

$\tan(-5\pi) = 0$, $\csc(-5\pi)$ is undefined, $\sec(-5\pi) = -1$, and $\cot(-5\pi)$ is undefined.

27.




Using Figure 8 we see that a point on the terminal line is $P(-\sqrt{3}, 1)$.

Therefore taking $x = -\sqrt{3}$, $y = 1$, $r = 2$ in the definitions of the trigonometric ratios, we have $\sin \frac{5\pi}{6} = \frac{1}{2}$, $\cos \frac{5\pi}{6} = -\frac{\sqrt{3}}{2}$,

$\tan \frac{5\pi}{6} = -\frac{1}{\sqrt{3}}$, $\csc \frac{5\pi}{6} = 2$, $\sec \frac{5\pi}{6} = -\frac{2}{\sqrt{3}}$, and $\cot \frac{5\pi}{6} = -\sqrt{3}$.

28.



From the diagram, we see that a point on the terminal line is $P(-1, 1)$.

Therefore taking $x = -1$, $y = 1$, $r = \sqrt{2}$ in the definitions of the trigonometric ratios we have $\sin \frac{11\pi}{4} = \frac{1}{\sqrt{2}}$, $\cos \frac{11\pi}{4} = -\frac{1}{\sqrt{2}}$,

$\tan \frac{11\pi}{4} = -1$, $\csc \frac{11\pi}{4} = \sqrt{2}$, $\sec \frac{11\pi}{4} = -\sqrt{2}$, and $\cot \frac{11\pi}{4} = -1$.

29. $\sin \theta = y/r = \frac{3}{5} \Rightarrow y = 3, r = 5$, and $x = \sqrt{r^2 - y^2} = 4$ (since $0 < \theta < \frac{\pi}{2}$). Therefore taking $x = 4, y = 3, r = 5$ in the definitions of the trigonometric ratios, we have $\cos \theta = \frac{4}{5}, \tan \theta = \frac{3}{4}, \csc \theta = \frac{5}{3}, \sec \theta = \frac{5}{4}$, and $\cot \theta = \frac{4}{3}$.

30. Since $0 < \alpha < \frac{\pi}{2}$, α is in the first quadrant where x and y are both positive. Therefore, $\tan \alpha = y/x = \frac{2}{1} \Rightarrow y = 2, x = 1$, and $r = \sqrt{x^2 + y^2} = \sqrt{5}$. Taking $x = 1, y = 2, r = \sqrt{5}$ in the definitions of the trigonometric ratios, we have $\sin \alpha = \frac{2}{\sqrt{5}}, \cos \alpha = \frac{1}{\sqrt{5}}, \csc \alpha = \frac{\sqrt{5}}{2}, \sec \alpha = \sqrt{5}$, and $\cot \alpha = \frac{1}{2}$.

31. $\frac{\pi}{2} < \phi < \pi \Rightarrow \phi$ is in the second quadrant, where x is negative and y is positive. Therefore $\sec \phi = r/x = -1.5 = -\frac{3}{2} \Rightarrow r = 3, x = -2$, and $y = \sqrt{r^2 - x^2} = \sqrt{5}$. Taking $x = -2, y = \sqrt{5}$, and $r = 3$ in the definitions of the trigonometric ratios, we have $\sin \phi = \frac{\sqrt{5}}{3}, \cos \phi = -\frac{2}{3}, \tan \phi = -\frac{\sqrt{5}}{2}, \csc \phi = \frac{3}{\sqrt{5}}$, and $\cot \theta = -\frac{2}{\sqrt{5}}$.

32. Since $\pi < x < \frac{3\pi}{2}$, x is in the third quadrant where x and y are both negative. Therefore $\cos x = x/r = -\frac{1}{3} \Rightarrow x = -1, r = 3$, and $y = -\sqrt{r^2 - x^2} = -\sqrt{8} = -2\sqrt{2}$. Taking $x = -1, r = 3, y = -2\sqrt{2}$ in the definitions of the trigonometric ratios, we have $\sin x = -\frac{2\sqrt{2}}{3}, \tan x = 2\sqrt{2}, \csc x = -\frac{3}{2\sqrt{2}}, \sec x = -3$, and $\cot x = \frac{1}{2\sqrt{2}}$.

33. $\pi < \beta < 2\pi$ means that β is in the third or fourth quadrant where y is negative. Also since $\cot \beta = x/y = 3$ which is positive, x must also be negative. Therefore $\cot \beta = x/y = \frac{3}{1} \Rightarrow x = -3, y = -1$, and $r = \sqrt{x^2 + y^2} = \sqrt{10}$. Taking $x = -3, y = -1$ and $r = \sqrt{10}$ in the definitions of the trigonometric ratios, we have $\sin \beta = -\frac{1}{\sqrt{10}}, \cos \beta = -\frac{3}{\sqrt{10}}, \tan \beta = \frac{1}{3}, \csc \beta = -\sqrt{10}$, and $\sec \beta = -\frac{\sqrt{10}}{3}$.

34. Since $\frac{3\pi}{2} < \theta < 2\pi$, θ is in the fourth quadrant where x is positive and y is negative. Therefore $\csc \theta = r/y = -\frac{4}{3} \Rightarrow r = 4, y = -3$, and $x = \sqrt{r^2 - y^2} = \sqrt{7}$. Taking $x = \sqrt{7}, y = -3$, and $r = 4$ in the definitions of the trigonometric ratios, we have $\sin \theta = -\frac{3}{4}, \cos \theta = \frac{\sqrt{7}}{4}, \tan \theta = -\frac{3}{\sqrt{7}}, \sec \theta = \frac{4}{\sqrt{7}}$, and $\cot \theta = -\frac{\sqrt{7}}{3}$.


35. $\sin 35^\circ = \frac{x}{10} \Rightarrow x = 10 \sin 35^\circ \approx 5.73576$ cm

36. $\cos 40^\circ = \frac{x}{25} \Rightarrow x = 25 \cos 40^\circ \approx 19.15111$ cm

37. $\tan \frac{2\pi}{5} = \frac{x}{8} \Rightarrow x = 8 \tan \frac{2\pi}{5} \approx 24.62147$ cm

38. $\cos \frac{3\pi}{8} = \frac{22}{x} \Rightarrow x = \frac{22}{\cos \frac{3\pi}{8}} \approx 57.48877$ cm

39.



(a) From the diagram we see that $\sin \theta = \frac{y}{r} = \frac{a}{c}$, and

$$\sin(-\theta) = \frac{-a}{c} = -\frac{a}{c} = -\sin \theta.$$

(b) Again from the diagram we see that $\cos \theta = \frac{x}{r} = \frac{b}{c} = \cos(-\theta)$.

40. (a) Using (12a) and (12b), we have

$$\tan(x+y) = \frac{\sin(x+y)}{\cos(x+y)} = \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y - \sin x \sin y} = \frac{\frac{\sin x \cos y}{\cos x \cos y} + \frac{\cos x \sin y}{\cos x \cos y}}{\frac{\cos x \cos y}{\cos x \cos y} - \frac{\sin x \sin y}{\cos x \cos y}} = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

(b) From (10a) and (10b), we have $\tan(-\theta) = -\tan \theta$, so (14a) implies that

$$\tan(x-y) = \tan(x+(-y)) = \frac{\tan x + \tan(-y)}{1 - \tan x \tan(-y)} = \frac{\tan x - \tan y}{1 + \tan x \tan y}$$

41. (a) Using (12a) and (13a), we have

$$\begin{aligned} \frac{1}{2} [\sin(x+y) + \sin(x-y)] &= \frac{1}{2} [\sin x \cos y + \cos x \sin y + \sin x \cos y - \cos x \sin y] \\ &= \frac{1}{2} (2 \sin x \cos y) = \sin x \cos y \end{aligned}$$

(b) This time, using (12b) and (13b), we have

$$\begin{aligned} \frac{1}{2} [\cos(x+y) + \cos(x-y)] &= \frac{1}{2} [\cos x \cos y - \sin x \sin y + \cos x \cos y + \sin x \sin y] \\ &= \frac{1}{2} (2 \cos x \cos y) = \cos x \cos y \end{aligned}$$

(c) Again using (12b) and (13b), we have

$$\begin{aligned} \frac{1}{2} [\cos(x-y) - \cos(x+y)] &= \frac{1}{2} [\cos x \cos y + \sin x \sin y - \cos x \cos y + \sin x \sin y] \\ &= \frac{1}{2} (2 \sin x \sin y) = \sin x \sin y \end{aligned}$$

42. Using (13b), $\cos\left(\frac{\pi}{2} - x\right) = \cos\frac{\pi}{2} \cos x + \sin\frac{\pi}{2} \sin x = 0 \cdot \cos x + 1 \cdot \sin x = \sin x$.

43. Using (12a), we have $\sin\left(\frac{\pi}{2} + x\right) = \sin\frac{\pi}{2} \cos x + \cos\frac{\pi}{2} \sin x = 1 \cdot \cos x + 0 \cdot \sin x = \cos x$.

44. Using (13a), we have $\sin(\pi - x) = \sin \pi \cos x - \cos \pi \sin x = 0 \cdot \cos x - (-1) \sin x = \sin x$.

45. Using (6), we have $\sin \theta \cot \theta = \sin \theta \cdot \frac{\cos \theta}{\sin \theta} = \cos \theta$.

46. $(\sin x + \cos x)^2 = \sin^2 x + 2 \sin x \cos x + \cos^2 x = (\sin^2 x + \cos^2 x) + \sin 2x$ [by (15a)]
 $= 1 + \sin 2x$ [by (7)]

47. $\sec y - \cos y = \frac{1}{\cos y} - \cos y$ [by (6)] $= \frac{1 - \cos^2 y}{\cos y} = \frac{\sin^2 y}{\cos y}$ [by (7)] $= \frac{\sin y}{\cos y} \sin y = \tan y \sin y$ [by (6)]

48. $\tan^2 \alpha - \sin^2 \alpha = \frac{\sin^2 \alpha}{\cos^2 \alpha} - \sin^2 \alpha = \frac{\sin^2 \alpha - \sin^2 \alpha \cos^2 \alpha}{\cos^2 \alpha} = \frac{\sin^2 \alpha (1 - \cos^2 \alpha)}{\cos^2 \alpha} = \tan^2 \alpha \sin^2 \alpha$ [by (6), (7)]

49. $\cot^2 \theta + \sec^2 \theta = \frac{\cos^2 \theta}{\sin^2 \theta} + \frac{1}{\cos^2 \theta}$ [by (6)] $= \frac{\cos^2 \theta \cos^2 \theta + \sin^2 \theta}{\sin^2 \theta \cos^2 \theta}$
 $= \frac{(1 - \sin^2 \theta)(1 - \sin^2 \theta) + \sin^2 \theta}{\sin^2 \theta \cos^2 \theta}$ [by (7)] $= \frac{1 - \sin^2 \theta + \sin^4 \theta}{\sin^2 \theta \cos^2 \theta}$
 $= \frac{\cos^2 \theta + \sin^4 \theta}{\sin^2 \theta \cos^2 \theta}$ [by (7)] $= \frac{1}{\sin^2 \theta} + \frac{\sin^2 \theta}{\cos^2 \theta} = \csc^2 \theta + \tan^2 \theta$ [by (6)]

50. $2 \csc 2t = \frac{2}{\sin 2t} = \frac{2}{2 \sin t \cos t}$ [by (15a)] $= \frac{1}{\sin t \cos t} = \sec t \csc t$

51. Using (14a), we have $\tan 2\theta = \tan(\theta + \theta) = \frac{\tan \theta + \tan \theta}{1 - \tan \theta \tan \theta} = \frac{2 \tan \theta}{1 - \tan^2 \theta}$.

52. $\frac{1}{1 - \sin \theta} + \frac{1}{1 + \sin \theta} = \frac{1 + \sin \theta + 1 - \sin \theta}{(1 - \sin \theta)(1 + \sin \theta)} = \frac{2}{1 - \sin^2 \theta} = \frac{2}{\cos^2 \theta}$ [by (7)] $= 2 \sec^2 \theta$

53. Using (15a) and (16a),

$$\begin{aligned} \sin x \sin 2x + \cos x \cos 2x &= \sin x (2 \sin x \cos x) + \cos x (2 \cos^2 x - 1) = 2 \sin^2 x \cos x + 2 \cos^3 x - \cos x \\ &= 2 (1 - \cos^2 x) \cos x + 2 \cos^3 x - \cos x \text{ [by (7)]} \\ &= 2 \cos x - 2 \cos^3 x + 2 \cos^3 x - \cos x = \cos x \end{aligned}$$

Or: $\sin x \sin 2x + \cos x \cos 2x = \cos(2x - x)$ [by 13(b)] $= \cos x$

54. Working backward, we start with equations (12a) and (13a):

$$\begin{aligned} \sin(x + y) \sin(x - y) &= (\sin x \cos y + \cos x \sin y) (\sin x \cos y - \cos x \sin y) \\ &= \sin^2 x \cos^2 y - \sin x \cos y \cos x \sin y + \cos x \sin y \sin x \cos y - \cos^2 x \sin^2 y \\ &= \sin^2 x (1 - \sin^2 y) - (1 - \sin^2 x) \sin^2 y \text{ [by (7)]} \\ &= \sin^2 x - \sin^2 x \sin^2 y - \sin^2 y + \sin^2 x \sin^2 y = \sin^2 x - \sin^2 y \end{aligned}$$

55. $\frac{\sin \phi}{1 - \cos \phi} = \frac{\sin \phi}{1 - \cos \phi} \cdot \frac{1 + \cos \phi}{1 + \cos \phi} = \frac{\sin \phi (1 + \cos \phi)}{1 - \cos^2 \phi} = \frac{\sin \phi (1 + \cos \phi)}{\sin^2 \phi}$ [by (7)]
 $= \frac{1 + \cos \phi}{\sin \phi} = \frac{1}{\sin \phi} + \frac{\cos \phi}{\sin \phi} = \csc \phi + \cot \phi$ [by (6)]

56. $\tan x + \tan y = \frac{\sin x}{\cos x} + \frac{\sin y}{\cos y} = \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y} = \frac{\sin(x + y)}{\cos x \cos y}$ [by (12a)]


57. Using (12a),

$$\begin{aligned} \sin 3\theta + \sin \theta &= \sin(2\theta + \theta) + \sin \theta = \sin 2\theta \cos \theta + \cos 2\theta \sin \theta + \sin \theta \\ &= \sin 2\theta \cos \theta + (2 \cos^2 \theta - 1) \sin \theta + \sin \theta \text{ [by (16a)]} \\ &= \sin 2\theta \cos \theta + 2 \cos^2 \theta \sin \theta - \sin \theta + \sin \theta = \sin 2\theta \cos \theta + \sin 2\theta \cos \theta \text{ [by (15a)]} \\ &= 2 \sin 2\theta \cos \theta \end{aligned}$$

58. We use (12b) with $x = 2\theta$, $y = \theta$ to get

$$\begin{aligned} \cos 3\theta &= \cos(2\theta + \theta) = \cos 2\theta \cos \theta - \sin 2\theta \sin \theta \\ &= (2 \cos^2 \theta - 1) \cos \theta - 2 \sin^2 \theta \cos \theta \text{ [by (16a) and (15a)]} \\ &= (2 \cos^2 \theta - 1) \cos \theta - 2(1 - \cos^2 \theta) \cos \theta \text{ [by (7)]} \\ &= 2 \cos^3 \theta - \cos \theta - 2 \cos \theta + 2 \cos^3 \theta = 4 \cos^3 \theta - 3 \cos \theta \end{aligned}$$

59. Since $\sin x = \frac{1}{3}$ we can label the opposite side as having length 1, the hypotenuse as having length 3, and use the Pythagorean Theorem to get that the adjacent side has length $\sqrt{8}$. Then, from the diagram, $\cos x = \frac{\sqrt{8}}{3}$. Similarly we have that $\sin y = \frac{3}{5}$. Now use (12a):



$$\sin(x + y) = \sin x \cos y + \cos x \sin y = \frac{1}{3} \cdot \frac{4}{5} + \frac{\sqrt{8}}{3} \cdot \frac{3}{5} = \frac{4}{15} + \frac{3\sqrt{8}}{15} = \frac{4 + 6\sqrt{2}}{15}.$$

60. Use (12b) and the values for $\sin y$ and $\cos x$ obtained in Exercise 59 to get

$$\cos(x + y) = \cos x \cos y - \sin x \sin y = \frac{\sqrt{8}}{3} \cdot \frac{4}{5} - \frac{1}{3} \cdot \frac{3}{5} = \frac{8\sqrt{2} - 3}{15}$$

61. Using (13b) and the values for $\cos x$ and $\sin y$ obtained in Exercise 59, we have

$$\cos(x - y) = \cos x \cos y + \sin x \sin y = \frac{\sqrt{8}}{3} \cdot \frac{4}{5} + \frac{1}{3} \cdot \frac{3}{5} = \frac{8\sqrt{2} + 3}{15}$$

62. Using (13a) and the values for $\sin y$ and $\cos x$ obtained in Exercise 59, we get

$$\sin(x - y) = \sin x \cos y - \cos x \sin y = \frac{1}{3} \cdot \frac{4}{5} - \frac{\sqrt{8}}{3} \cdot \frac{3}{5} = \frac{4 - 6\sqrt{2}}{15}$$

63. Using (15a) and the values for $\sin y$ and $\cos y$ obtained in Exercise 59, we have

$$\sin 2y = 2 \sin y \cos y = 2 \cdot \frac{3}{5} \cdot \frac{4}{5} = \frac{24}{25}$$

64. Using (16a) with $\cos y = \frac{4}{5}$, we have $\cos 2y = 2 \cos^2 y - 1 = 2\left(\frac{4}{5}\right)^2 - 1 = \frac{32}{25} - 1 = \frac{7}{25}$.

65. $2 \cos x - 1 = 0 \Leftrightarrow \cos x = \frac{1}{2} \Rightarrow x = \frac{\pi}{3}, \frac{5\pi}{3}$ for $x \in [0, 2\pi]$.

66. $3 \cot^2 x = 1 \Leftrightarrow 3 = 1/\cot^2 x \Leftrightarrow \tan^2 x = 3 \Leftrightarrow \tan x = \pm\sqrt{3} \Rightarrow x = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3},$ and $\frac{5\pi}{3}$.

67. $2 \sin^2 x = 1 \Leftrightarrow \sin^2 x = \frac{1}{2} \Leftrightarrow \sin x = \pm\frac{1}{\sqrt{2}} \Rightarrow x = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$.

68. $|\tan x| = 1 \Leftrightarrow \tan x = -1$ or $\tan x = 1 \Leftrightarrow x = \frac{3\pi}{4}, \frac{7\pi}{4}$ or $x = \frac{\pi}{4}, \frac{5\pi}{4}$.


69. Using (15a), we have $\sin 2x = \cos x \Leftrightarrow 2 \sin x \cos x - \cos x = 0 \Leftrightarrow \cos x(2 \sin x - 1) = 0 \Leftrightarrow \cos x = 0$ or $2 \sin x - 1 = 0 \Rightarrow x = \frac{\pi}{2}, \frac{3\pi}{2}$ or $\sin x = \frac{1}{2} \Rightarrow x = \frac{\pi}{6}$ or $\frac{5\pi}{6}$. Therefore, the solutions are $x = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}, \frac{3\pi}{2}$.

70. By (15a), $2 \cos x + \sin 2x = 0 \Leftrightarrow 2 \cos x + 2 \sin x \cos x = 0 \Leftrightarrow 2 \cos x(1 + \sin x) = 0 \Leftrightarrow \cos x = 0$ or $1 + \sin x = 0 \Leftrightarrow x = \frac{\pi}{2}, \frac{3\pi}{2}$ or $\sin x = -1 \Rightarrow x = \frac{3}{2}\pi$. So the solutions are $x = \frac{\pi}{2}, \frac{3\pi}{2}$.


71. $\sin x = \tan x \Leftrightarrow \sin x - \tan x = 0 \Leftrightarrow \sin x - \frac{\sin x}{\cos x} = 0 \Leftrightarrow \sin x \left(1 - \frac{1}{\cos x}\right) = 0 \Leftrightarrow \sin x = 0$ or $1 - \frac{1}{\cos x} = 0 \Rightarrow x = 0, \pi, 2\pi$ or $1 = \frac{1}{\cos x} \Rightarrow \cos x = 1 \Rightarrow x = 0, 2\pi$. Therefore the solutions are $x = 0, \pi, 2\pi$.

72. By (16a), $2 + \cos 2x = 3 \cos x \Leftrightarrow 2 + 2 \cos^2 x - 1 = 3 \cos x \Leftrightarrow 2 \cos^2 x - 3 \cos x + 1 = 0 \Leftrightarrow (2 \cos x - 1)(\cos x - 1) = 0 \Leftrightarrow \cos x = 1$ or $\cos x = \frac{1}{2} \Rightarrow x = 0, 2\pi$ or $x = \frac{\pi}{3}, \frac{5\pi}{3}$.


73. We know that $\sin x = \frac{1}{2}$ when $x = \frac{\pi}{6}$ or $\frac{5\pi}{6}$, and from Figure 13(a), we see that $\sin x \leq \frac{1}{2} \Rightarrow 0 \leq x \leq \frac{\pi}{6}$ or $\frac{5\pi}{6} \leq x \leq 2\pi$ for $x \in [0, 2\pi]$.
74. $2 \cos x + 1 > 0 \Rightarrow 2 \cos x > -1 \Rightarrow \cos x > -\frac{1}{2}$. $\cos x = -\frac{1}{2}$ when $x = \frac{2\pi}{3}, \frac{4\pi}{3}$ and from Figure 13(b), we see that $\cos x > -\frac{1}{2}$ when $0 \leq x < \frac{2\pi}{3}, \frac{4\pi}{3} < x \leq 2\pi$.
75. $\tan x = -1$ when $x = \frac{3\pi}{4}, \frac{7\pi}{4}$, and $\tan x = 1$ when $x = \frac{\pi}{4}$ or $\frac{5\pi}{4}$. From Figure 14 we see that $-1 < \tan x < 1 \Rightarrow 0 \leq x < \frac{\pi}{4}, \frac{3\pi}{4} < x < \frac{5\pi}{4}$, and $\frac{7\pi}{4} < x \leq 2\pi$.
76. We know that $\sin x = \cos x$ when $x = \frac{\pi}{4}, \frac{5\pi}{4}$, and from the diagram we see that $\sin x > \cos x$ when $\frac{\pi}{4} < x < \frac{5\pi}{4}$.




77. $y = \cos(x - \frac{\pi}{3})$. We start with the graph of $y = \cos x$ and shift it $\frac{\pi}{3}$ units to the right.




78. $y = \tan 2x$. Start with the graph of $y = \tan x$ with period π and compress it to a period of $\frac{\pi}{2}$.




79. $y = \frac{1}{3} \tan(x - \frac{\pi}{2})$. We start with the graph of $y = \tan x$, shift it $\frac{\pi}{2}$ units to the right and compress it to $\frac{1}{3}$ of its original vertical size.




80. $y = 1 + \sec x$. Start with the graph of $y = \sec x$ and raise it by one unit.



81. $y = |\sin x|$. We start with the graph of $y = \sin x$ and reflect the parts below the x -axis about the x -axis.



82. $y = 2 + \sin(x + \frac{\pi}{4})$. Start with the graph of $y = \sin x$, and shift it $\frac{\pi}{4}$ units to the left and 2 units up.



83. From the figure in the text, we see that $x = b \cos \theta$, $y = b \sin \theta$, and from the distance formula we have that the distance c from (x, y) to $(a, 0)$ is $c = \sqrt{(x-a)^2 + (y-0)^2} \Rightarrow$

$$\begin{aligned} c^2 &= (b \cos \theta - a)^2 + (b \sin \theta)^2 = b^2 \cos^2 \theta - 2ab \cos \theta + a^2 + b^2 \sin^2 \theta \\ &= a^2 + b^2 (\cos^2 \theta + \sin^2 \theta) - 2ab \cos \theta = a^2 + b^2 - 2ab \cos \theta \quad [\text{by (7)}] \end{aligned}$$


84. $|AB|^2 = |AC|^2 + |BC|^2 - 2|AC||BC| \cos \angle C = (820)^2 + (910)^2 - 2(820)(910) \cos 103^\circ$
 $\approx 1,836,217 \Rightarrow |AB| \approx 1355 \text{ m}$

85. Using the Law of Cosines, we have $c^2 = 1^2 + 1^2 - 2(1)(1) \cos(\alpha - \beta) = 2[1 - \cos(\alpha - \beta)]$. Now, using the distance formula, $c^2 = |AB|^2 = (\cos \alpha - \cos \beta)^2 + (\sin \alpha - \sin \beta)^2$. Equating these two expressions for c^2 , we get $2[1 - \cos(\alpha - \beta)] = \cos^2 \alpha + \sin^2 \alpha + \cos^2 \beta + \sin^2 \beta - 2 \cos \alpha \cos \beta - 2 \sin \alpha \sin \beta \Rightarrow$
 $1 - \cos(\alpha - \beta) = 1 - \cos \alpha \cos \beta - \sin \alpha \sin \beta \Rightarrow \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$.

86. $\cos(x + y) = \cos(x - (-y)) = \cos x \cos(-y) + \sin x \sin(-y)$
 $= \cos x \cos y - \sin x \sin y$ [using Equations 10a and 10b]

87. In Exercise 86 we used the subtraction formula for cosine to prove the addition formula for cosine. Using that formula with $x = \frac{\pi}{2} - \alpha$, $y = \beta$, we get $\cos[(\frac{\pi}{2} - \alpha) + \beta] = \cos(\frac{\pi}{2} - \alpha) \cos \beta - \sin(\frac{\pi}{2} - \alpha) \sin \beta \Rightarrow$
 $\cos[\frac{\pi}{2} - (\alpha - \beta)] = \cos(\frac{\pi}{2} - \alpha) \cos \beta - \sin(\frac{\pi}{2} - \alpha) \sin \beta$. Now we use the identities given in the problem, $\cos(\frac{\pi}{2} - \theta) = \sin \theta$ and $\sin(\frac{\pi}{2} - \theta) = \cos \theta$, to get $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$.

88. If $0 < \theta < \frac{\pi}{2}$, we have the case depicted in the first diagram. In this case, we see that the height of the triangle is $h = a \sin \theta$. If $\frac{\pi}{2} \leq \theta < \pi$, we have the case depicted in the second diagram. In this case, the height of the triangle is $h = a \sin(\pi - \theta) = a \sin \theta$ (by the identity proved in Exercise 78). So in either case, the area of the triangle is $\frac{1}{2}bh = \frac{1}{2}ab \sin \theta$.



89. Using the formula from part (a), the area of the triangle is $\frac{1}{2}(10)(3) \sin 107^\circ \approx 14.34457 \text{ cm}^2$.

$$26. \sum_{i=1}^{100} 4 = \underbrace{4 + 4 + 4 + \cdots + 4}_{(100 \text{ summands})} = 100 \cdot 4 = 400$$

$$27. \sum_{i=0}^4 (2^i + i^2) = (1 + 0) + (2 + 1) + (4 + 4) + (8 + 9) + (16 + 16) = 61$$

$$28. \sum_{i=-2}^4 2^{3-i} = 2^5 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 + 2^{-1} = 63.5$$

$$29. \sum_{i=1}^n 2i = 2 \sum_{i=1}^n i = 2 \cdot \frac{n(n+1)}{2} \text{ [by Theorem 3(c)]} = n(n+1)$$

$$30. \sum_{i=1}^n (2 - 5i) = \sum_{i=1}^n 2 - \sum_{i=1}^n 5i = 2n - 5 \sum_{i=1}^n i = 2n - \frac{5n(n+1)}{2} = \frac{4n}{2} - \frac{5n^2 + 5n}{2} = -\frac{n(5n+1)}{2}$$

$$\begin{aligned} 31. \sum_{i=1}^n (i^2 + 3i + 4) &= \sum_{i=1}^n i^2 + 3 \sum_{i=1}^n i + \sum_{i=1}^n 4 = \frac{n(n+1)(2n+1)}{6} + \frac{3n(n+1)}{2} + 4n \\ &= \frac{1}{6} [(2n^3 + 3n^2 + n) + (9n^2 + 9n) + 24n] = \frac{1}{6} (2n^3 + 12n^2 + 34n) \\ &= \frac{1}{3} n(n^2 + 6n + 17) \end{aligned}$$

$$\begin{aligned} 32. \sum_{i=1}^n (3 + 2i)^2 &= \sum_{i=1}^n (9 + 12i + 4i^2) = \sum_{i=1}^n 9 + 12 \sum_{i=1}^n i + 4 \sum_{i=1}^n i^2 \\ &= 9n + 6n(n+1) + \frac{2n(n+1)(2n+1)}{3} = \frac{27n + 18n^2 + 18n + 4n^3 + 6n^2 + 2n}{3} \\ &= \frac{1}{3} (4n^3 + 24n^2 + 47n) = \frac{1}{3} n(4n^2 + 24n + 47) \end{aligned}$$

$$\begin{aligned} 33. \sum_{i=1}^n (i+1)(i+2) &= \sum_{i=1}^n (i^2 + 3i + 2) = \sum_{i=1}^n i^2 + 3 \sum_{i=1}^n i + \sum_{i=1}^n 2 \\ &= \frac{n(n+1)(2n+1)}{6} + \frac{3n(n+1)}{2} + 2n = \frac{n(n+1)}{6} [(2n+1) + 9] + 2n \\ &= \frac{n(n+1)}{3} (n+5) + 2n = \frac{n}{3} [(n+1)(n+5) + 6] = \frac{n}{3} (n^2 + 6n + 11) \end{aligned}$$

$$\begin{aligned} 34. \sum_{i=1}^n i(i+1)(i+2) &= \sum_{i=1}^n (i^3 + 3i^2 + 2i) = \sum_{i=1}^n i^3 + 3 \sum_{i=1}^n i^2 + 2 \sum_{i=1}^n i \\ &= \left[\frac{n(n+1)}{2} \right]^2 + \frac{3n(n+1)(2n+1)}{6} + \frac{2n(n+1)}{2} \\ &= n(n+1) \left[\frac{n(n+1)}{4} + \frac{2n+1}{2} + 1 \right] = \frac{n(n+1)}{4} (n^2 + n + 4n + 2 + 4) \\ &= \frac{n(n+1)}{4} (n^2 + 5n + 6) = \frac{n(n+1)(n+2)(n+3)}{4} \end{aligned}$$

$$\begin{aligned} 35. \sum_{i=1}^n (i^3 - i - 2) &= \sum_{i=1}^n i^3 - \sum_{i=1}^n i - \sum_{i=1}^n 2 = \left[\frac{n(n+1)}{2} \right]^2 - \frac{n(n+1)}{2} - 2n \\ &= \frac{1}{4} n(n+1) [n(n+1) - 2] - 2n = \frac{1}{4} n(n+1)(n+2)(n-1) - 2n \\ &= \frac{1}{4} n [(n+1)(n-1)(n+2) - 8] = \frac{1}{4} n [(n^2 - 1)(n+2) - 8] = \frac{1}{4} n (n^3 + 2n^2 - n - 10) \end{aligned}$$

36. By Theorem 3(c) we have that $\sum_{i=1}^n i = \frac{n(n+1)}{2} = 78 \Leftrightarrow n(n+1) = 156 \Leftrightarrow n^2 + n - 156 = 0 \Leftrightarrow (n+13)(n-12) = 0 \Leftrightarrow n = 12$ or -13 . But $n = -13$ produces a negative answer for the sum, so $n = 12$.

37. By Theorem 2(a) and Example 3, $\sum_{i=1}^n c = c \sum_{i=1}^n 1 = cn$.

38. Let S_n be the statement that $\sum_{i=1}^n i^3 = \left[\frac{n(n+1)}{2} \right]^2$.

1. S_1 is true because $1^3 = \left(\frac{1 \cdot 2}{2} \right)^2$.

2. Assume S_k is true. Then $\sum_{i=1}^k i^3 = \left[\frac{k(k+1)}{2} \right]^2$, so

$$\begin{aligned} \sum_{i=1}^{k+1} i^3 &= \left[\frac{k(k+1)}{2} \right]^2 + (k+1)^3 = \frac{(k+1)^2}{4} [k^2 + 4(k+1)] = \frac{(k+1)^2}{4} (k+2)^2 \\ &= \left(\frac{(k+1)[(k+1)+1]}{2} \right)^2 \end{aligned}$$

showing that S_{k+1} is true.

Therefore, S_n is true for all n by mathematical induction.

39.
$$\begin{aligned} \sum_{i=1}^n [(i+1)^4 - i^4] &= (2^4 - 1^4) + (3^4 - 2^4) + (4^4 - 3^4) + \cdots + [(n+1)^4 - n^4] \\ &= (n+1)^4 - 1^4 = n^4 + 4n^3 + 6n^2 + 4n \end{aligned}$$

On the other hand,

$$\begin{aligned} \sum_{i=1}^n [(i+1)^4 - i^4] &= \sum_{i=1}^n (4i^3 + 6i^2 + 4i + 1) = 4 \sum_{i=1}^n i^3 + 6 \sum_{i=1}^n i^2 + 4 \sum_{i=1}^n i + \sum_{i=1}^n 1 \\ &= 4S + n(n+1)(2n+1) + 2n(n+1) + n \left[\text{where } S = \sum_{i=1}^n i^3 \right] \\ &= 4S + 2n^3 + 3n^2 + n + 2n^2 + 2n + n = 4S + 2n^3 + 5n^2 + 4n \end{aligned}$$

Thus, $n^4 + 4n^3 + 6n^2 + 4n = 4S + 2n^3 + 5n^2 + 4n$, from which it follows that

$$4S = n^4 + 2n^3 + n^2 = n^2(n^2 + 2n + 1) = n^2(n+1)^2 \text{ and } S = \left[\frac{n(n+1)}{2} \right]^2.$$

40. The area of G_i is

$$\begin{aligned} \left(\sum_{k=1}^i k \right)^2 - \left(\sum_{k=1}^{i-1} k \right)^2 &= \left[\frac{i(i+1)}{2} \right]^2 - \left[\frac{(i-1)i}{2} \right]^2 = \frac{i^2}{4} [(i+1)^2 - (i-1)^2] \\ &= \frac{i^2}{4} [(i^2 + 2i + 1) - (i^2 - 2i + 1)] = \frac{i^2}{4} (4i) = i^3 \end{aligned}$$

Thus, the area of $ABCD$ is $\sum_{i=1}^n i^3 = \left[\frac{n(n+1)}{2} \right]^2$.

$$41. (a) \sum_{i=1}^n [i^4 - (i-1)^4] = (1^4 - 0^4) + (2^4 - 1^4) + (3^4 - 2^4) + \cdots + [n^4 - (n-1)^4] = n^4 - 0 = n^4$$

$$(b) \sum_{i=1}^{100} (5^i - 5^{i-1}) = (5^1 - 5^0) + (5^2 - 5^1) + (5^3 - 5^2) + \cdots + (5^{100} - 5^{99}) = 5^{100} - 5^0 = 5^{100} - 1$$

$$(c) \sum_{i=3}^{99} \left(\frac{1}{i} - \frac{1}{i+1} \right) = \left(\frac{1}{3} - \frac{1}{4} \right) + \left(\frac{1}{4} - \frac{1}{5} \right) + \left(\frac{1}{5} - \frac{1}{6} \right) + \cdots + \left(\frac{1}{99} - \frac{1}{100} \right) = \frac{1}{3} - \frac{1}{100} = \frac{97}{300}$$

$$(d) \sum_{i=1}^n (a_i - a_{i-1}) = (a_1 - a_0) + (a_2 - a_1) + (a_3 - a_2) + \cdots + (a_n - a_{n-1}) = a_n - a_0$$

42. Summing the inequalities $-|a_i| \leq a_i \leq |a_i|$ for $i = 1, 2, \dots, n$, we get $-\sum_{i=1}^n |a_i| \leq \sum_{i=1}^n a_i \leq \sum_{i=1}^n |a_i|$. Since

$$|x| \leq c \Leftrightarrow -c \leq x \leq c, \text{ we have } \left| \sum_{i=1}^n a_i \right| \leq \sum_{i=1}^n |a_i|. \text{ Another method: Use mathematical induction.}$$

$$43. \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} \left(\frac{i}{n} \right)^2 = \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{i=1}^n i^2 = \lim_{n \rightarrow \infty} \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6} = \lim_{n \rightarrow \infty} \frac{1}{6} \left(1 + \frac{1}{n} \right) \left(2 + \frac{1}{n} \right) \\ = \frac{1}{6}(1)(2) = \frac{1}{3}$$

$$44. \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} \left[\left(\frac{i}{n} \right)^3 + 1 \right] = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[\frac{i^3}{n^4} + \frac{1}{n} \right] = \lim_{n \rightarrow \infty} \left[\frac{1}{n^4} \sum_{i=1}^n i^3 + \frac{1}{n} \sum_{i=1}^n 1 \right] \\ = \lim_{n \rightarrow \infty} \left[\frac{1}{n^4} \left(\frac{n(n+1)}{2} \right)^2 + \frac{1}{n}(n) \right] = \lim_{n \rightarrow \infty} \frac{1}{4} \left(1 + \frac{1}{n} \right)^2 + 1 = \frac{1}{4} + 1 = \frac{5}{4}$$

$$45. \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{2}{n} \left[\left(\frac{2i}{n} \right)^3 + 5 \left(\frac{2i}{n} \right) \right] = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[\frac{16}{n^4} i^3 + \frac{20}{n^2} i \right] = \lim_{n \rightarrow \infty} \left[\frac{16}{n^4} \sum_{i=1}^n i^3 + \frac{20}{n^2} \sum_{i=1}^n i \right] \\ = \lim_{n \rightarrow \infty} \left[\frac{16}{n^4} \frac{n^2(n+1)^2}{4} + \frac{20}{n^2} \frac{n(n+1)}{2} \right] = \lim_{n \rightarrow \infty} \left[\frac{4(n+1)^2}{n^2} + \frac{10n(n+1)}{n^2} \right] \\ = \lim_{n \rightarrow \infty} \left[4 \left(1 + \frac{1}{n} \right)^2 + 10 \left(1 + \frac{1}{n} \right) \right] = 4 \cdot 1 + 10 \cdot 1 = 14$$

$$46. \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{3}{n} \left[\left(1 + \frac{3i}{n} \right)^3 - 2 \left(1 + \frac{3i}{n} \right) \right] = \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{3}{n} \left[1 + \frac{9i}{n} + \frac{27i^2}{n^2} + \frac{27i^3}{n^3} - 2 - \frac{6i}{n} \right] \\ = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[\frac{81}{n^4} i^3 + \frac{81}{n^3} i^2 + \frac{9}{n^2} i - \frac{3}{n} \right] \\ = \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \frac{n^2(n+1)^2}{4} + \frac{81}{n^3} \frac{n(n+1)(2n+1)}{6} + \frac{9}{n^2} \frac{n(n+1)}{2} - \frac{3}{n} n \right] \\ = \lim_{n \rightarrow \infty} \left[\frac{81}{4} \left(1 + \frac{1}{n} \right)^2 + \frac{27}{2} \left(1 + \frac{1}{n} \right) \left(2 + \frac{1}{n} \right) + \frac{9}{2} \left(1 + \frac{1}{n} \right) - 3 \right] = \frac{81}{4} + \frac{54}{2} + \frac{9}{2} - 3 = \frac{195}{4}$$

47. Let $S = \sum_{i=1}^n ar^{i-1} = a + ar + ar^2 + \cdots + ar^{n-1}$. Multiplying both sides by r gives us

$rS = ar + ar^2 + \cdots + ar^{n-1} + ar^n$. Subtracting the first equation from the second, we find

$$(r-1)S = ar^n - a = a(r^n - 1), \text{ so } S = \frac{a(r^n - 1)}{r-1} \text{ (since } r \neq 1\text{).}$$

48. $\sum_{i=1}^n \frac{3}{2^{i-1}} = 3 \sum_{i=1}^n \left(\frac{1}{2}\right)^{i-1} = \frac{3\left[\left(\frac{1}{2}\right)^n - 1\right]}{\frac{1}{2} - 1}$ [using Exercise 47 with $a = 3$ and $r = \frac{1}{2}$] $= 6\left[1 - \left(\frac{1}{2}\right)^n\right]$

49. $\sum_{i=1}^n (2i + 2^i) = 2 \sum_{i=1}^n i + \sum_{i=1}^n 2 \cdot 2^{i-1} = 2 \frac{n(n+1)}{2} + \frac{2(2^n - 1)}{2 - 1} = 2^{n+1} + n^2 + n - 2$.

For the first sum we have used Theorem 3(c), and for the second, Exercise 47 with $a = r = 2$.

50. $\sum_{i=1}^m \left[\sum_{j=1}^n (i+j) \right] = \sum_{i=1}^m \left[\sum_{j=1}^n i + \sum_{j=1}^n j \right]$ [Theorem 2(b)] $= \sum_{i=1}^m \left[ni + \frac{n(n+1)}{2} \right]$ [Theorem 3(b) and (c)]

$$= \sum_{i=1}^m ni + \sum_{i=1}^m \frac{n(n+1)}{2} = \frac{nm(m+1)}{2} + \frac{nm(n+1)}{2} = \frac{nm}{2}(m+n+2)$$

G Complex Numbers

- $(5 - 6i) + (3 + 2i) = (5 + 3) + (-6 + 2)i = 8 + (-4)i = 8 - 4i$
- $(4 - \frac{1}{2}i) - (9 + \frac{5}{2}i) = (4 - 9) + (-\frac{1}{2} - \frac{5}{2})i = -5 + (-3)i = -5 - 3i$
- $(2 + 5i)(4 - i) = 2(4) + 2(-i) + (5i)(4) + (5i)(-i) = 8 - 2i + 20i - 5i^2$
 $= 8 + 18i - 5(-1) = 8 + 18i + 5 = 13 + 18i$
- $(1 - 2i)(8 - 3i) = 8 - 3i - 16i + 6(-1) = 2 - 19i$
- $\overline{12 + 7i} = 12 - 7i$
- $2i(\frac{1}{2} - i) = i - 2(-1) = 2 + i \Rightarrow \overline{2i(\frac{1}{2} - i)} = \overline{2 + i} = 2 - i$
- $\frac{1 + 4i}{3 + 2i} = \frac{1 + 4i}{3 + 2i} \cdot \frac{3 - 2i}{3 - 2i} = \frac{3 - 2i + 12i - 8(-1)}{3^2 + 2^2} = \frac{11 + 10i}{13} = \frac{11}{13} + \frac{10}{13}i$
- $\frac{3 + 2i}{1 - 4i} = \frac{3 + 2i}{1 - 4i} \cdot \frac{1 + 4i}{1 + 4i} = \frac{3 + 12i + 2i + 8(-1)}{1^2 + 4^2} = \frac{-5 + 14i}{17} = -\frac{5}{17} + \frac{14}{17}i$
- $\frac{1}{1 + i} = \frac{1}{1 + i} \cdot \frac{1 - i}{1 - i} = \frac{1 - i}{1 - (-1)} = \frac{1 - i}{2} = \frac{1}{2} - \frac{1}{2}i$
- $\frac{3}{4 - 3i} = \frac{3}{4 - 3i} \cdot \frac{4 + 3i}{4 + 3i} = \frac{12 + 9i}{16 - 9(-1)} = \frac{12}{25} + \frac{9}{25}i$
- $i^3 = i^2 \cdot i = (-1)i = -i$
- $i^{100} = (i^2)^{50} = (-1)^{50} = 1$
- $\sqrt{-25} = \sqrt{25}i = 5i$
- $\sqrt{-3}\sqrt{-12} = \sqrt{3}i\sqrt{12}i = \sqrt{3 \cdot 12}i^2 = \sqrt{36}(-1) = -6$

$$15. \overline{12 - 5i} = 12 + 15i \text{ and } |12 - 15i| = \sqrt{12^2 + (-5)^2} = \sqrt{144 + 25} = \sqrt{169} = 13$$

$$16. \overline{-1 + 2\sqrt{2}i} = -1 - 2\sqrt{2}i \text{ and } |-1 + 2\sqrt{2}i| = \sqrt{(-1)^2 + (2\sqrt{2})^2} = \sqrt{1 + 8} = \sqrt{9} = 3$$

$$17. \overline{-4i} = \overline{0 - 4i} = 0 + 4i = 4i \text{ and } |-4i| = \sqrt{0^2 + (-4)^2} = \sqrt{16} = 4$$

18. Let $z = a + bi$, $w = c + di$.

$$(a) \overline{z + w} = \overline{(a + bi) + (c + di)} = \overline{(a + c) + (b + d)i} \\ = (a + c) - (b + d)i = (a - bi) + (c - di) = \bar{z} + \bar{w}$$

$$(b) \overline{zw} = \overline{(a + bi)(c + di)} = \overline{(ac - bd) + (ad + bc)i} = (ac - bd) - (ad + bc)i. \\ \text{On the other hand, } \bar{z}\bar{w} = (a - bi)(c - di) = (ac - bd) - (ad + bc)i = \overline{zw}.$$

(c) Use mathematical induction and part (b): Let S_n be the statement that $\overline{z^n} = \bar{z}^n$.

S_1 is true because $\overline{z^1} = \bar{z} = \bar{z}^1$. Assume S_k is true, that is $\overline{z^k} = \bar{z}^k$. Then

$\overline{z^{k+1}} = \overline{z^{1+k}} = \overline{z z^k} = \overline{z} \overline{z^k}$ [part (b) with $w = z^k$] $= \bar{z}^1 \bar{z}^k = \bar{z}^{1+k} = \bar{z}^{k+1}$, which shows that S_{k+1} is true.

Therefore, by mathematical induction, $\overline{z^n} = \bar{z}^n$ for every positive integer n .

Another proof: Use part (b) with $w = z$, and mathematical induction.

$$19. 4x^2 + 9 = 0 \Leftrightarrow 4x^2 = -9 \Leftrightarrow x^2 = -\frac{9}{4} \Leftrightarrow x = \pm\sqrt{-\frac{9}{4}} = \pm\sqrt{\frac{9}{4}}i = \pm\frac{3}{2}i.$$

$$20. x^4 = 1 \Leftrightarrow x^4 - 1 = 0 \Leftrightarrow (x^2 - 1)(x^2 + 1) = 0 \Leftrightarrow x^2 - 1 = 0 \text{ or } x^2 + 1 = 0 \Leftrightarrow \\ x = \pm 1 \text{ or } x = \pm i.$$

$$21. x^2 + 2x + 5 = 0 \Leftrightarrow x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 1 \cdot 5}}{2 \cdot 1} = \frac{-2 \pm \sqrt{-16}}{2} = \frac{-2 \pm 4i}{2} = -1 \pm 2i$$

$$22. 2x^2 - 2x + 1 = 0 \Leftrightarrow x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \cdot 2 \cdot 1}}{2 \cdot 2} = \frac{2 \pm \sqrt{-4}}{4} = \frac{2 \pm 2i}{4} = \frac{1}{2} \pm \frac{1}{2}i$$

$$23. \text{By the quadratic formula, } z^2 + z + 2 = 0 \Leftrightarrow z = \frac{-1 \pm \sqrt{1^2 - 4(1)(2)}}{2(1)} = \frac{-1 \pm \sqrt{-7}}{2} = -\frac{1}{2} \pm \frac{\sqrt{7}}{2}i.$$

$$24. z^2 + \frac{1}{2}z + \frac{1}{4} = 0 \Leftrightarrow 4z^2 + 2z + 1 = 0 \Leftrightarrow$$

$$z = \frac{-2 \pm \sqrt{2^2 - 4(4)(1)}}{2(4)} = \frac{-2 \pm \sqrt{-12}}{8} = \frac{-2 \pm 2\sqrt{3}i}{8} = -\frac{1}{4} \pm \frac{\sqrt{3}}{4}i$$

25. For $z = -3 + 3i$, $r = \sqrt{(-3)^2 + 3^2} = 3\sqrt{2}$ and $\tan \theta = \frac{3}{-3} = -1 \Rightarrow \theta = \frac{3\pi}{4}$ (since z lies in the second quadrant). Therefore, $-3 + 3i = 3\sqrt{2}(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4})$.

26. For $z = 1 - \sqrt{3}i$, $r = \sqrt{1^2 + (-\sqrt{3})^2} = 2$ and $\tan \theta = \frac{-\sqrt{3}}{1} = -\sqrt{3} \Rightarrow \theta = \frac{5\pi}{3}$ (since z lies in the fourth quadrant). Therefore, $1 - \sqrt{3}i = 2(\cos \frac{5\pi}{3} + i \sin \frac{5\pi}{3})$.

27. For $z = 3 + 4i$, $r = \sqrt{3^2 + 4^2} = 5$ and $\tan \theta = \frac{4}{3} \Rightarrow \theta = \tan^{-1}(\frac{4}{3})$ (since z lies in the first quadrant).
Therefore, $3 + 4i = 5[\cos(\tan^{-1} \frac{4}{3}) + i \sin(\tan^{-1} \frac{4}{3})]$.

28. For $z = 8i$, $r = \sqrt{0^2 + 8^2} = 8$ and $\tan \theta = \frac{8}{0}$ is undefined, so $\theta = \frac{\pi}{2}$ (since z lies on the positive imaginary axis).
Therefore, $8i = 8(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2})$.

29. For $z = \sqrt{3} + i$, $r = \sqrt{(\sqrt{3})^2 + 1^2} = 2$ and $\tan \theta = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \Rightarrow z = 2(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$.

For $w = 1 + \sqrt{3}i$, $r = 2$ and $\tan \theta = \sqrt{3} \Rightarrow \theta = \frac{\pi}{3} \Rightarrow w = 2(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3})$.

Therefore, $zw = 2 \cdot 2[\cos(\frac{\pi}{6} + \frac{\pi}{3}) + i \sin(\frac{\pi}{6} + \frac{\pi}{3})] = 4(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2})$,

$z/w = \frac{2}{2}[\cos(\frac{\pi}{6} - \frac{\pi}{3}) + i \sin(\frac{\pi}{6} - \frac{\pi}{3})] = \cos(-\frac{\pi}{6}) + i \sin(-\frac{\pi}{6})$, and $1 = 1 + 0i = 1(\cos 0 + i \sin 0) \Rightarrow$

$1/z = \frac{1}{2}[\cos(0 - \frac{\pi}{6}) + i \sin(0 - \frac{\pi}{6})] = \frac{1}{2}[\cos(-\frac{\pi}{6}) + i \sin(-\frac{\pi}{6})]$. For $1/z$, we could also use the formula that precedes Example 5 to obtain $1/z = \frac{1}{2}(\cos \frac{\pi}{6} - i \sin \frac{\pi}{6})$.

30. For $z = 4\sqrt{3} - 4i$, $r = \sqrt{(4\sqrt{3})^2 + (-4)^2} = \sqrt{64} = 8$ and $\tan \theta = \frac{-4}{4\sqrt{3}} = -\frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{11\pi}{6} \Rightarrow$

$z = 8(\cos \frac{11\pi}{6} + i \sin \frac{11\pi}{6})$. For $w = 8i$, $r = \sqrt{0^2 + 8^2} = 8$ and $\tan \theta = \frac{8}{0}$ is undefined, so $\theta = \frac{\pi}{2} \Rightarrow$

$w = 8(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2})$. Therefore, $zw = 8 \cdot 8[\cos(\frac{11\pi}{6} + \frac{\pi}{2}) + i \sin(\frac{11\pi}{6} + \frac{\pi}{2})] = 64(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3})$,

$z/w = \frac{8}{8}[\cos(\frac{11\pi}{6} - \frac{\pi}{2}) + i \sin(\frac{11\pi}{6} - \frac{\pi}{2})] = \cos \frac{4\pi}{3} + i \sin \frac{4\pi}{3}$, and

$1 = 1 + 0i = 1(\cos 0 + i \sin 0) \Rightarrow 1/z = \frac{1}{8}[\cos(0 - \frac{11\pi}{6}) + i \sin(0 - \frac{11\pi}{6})] = \frac{1}{8}[\cos(\frac{\pi}{6}) + i \sin(\frac{\pi}{6})]$.

For $1/z$, we could also use the formula that precedes Example 5 to obtain $1/z = \frac{1}{8}(\cos \frac{11\pi}{6} - i \sin \frac{11\pi}{6})$.

31. For $z = 2\sqrt{3} - 2i$, $r = \sqrt{(2\sqrt{3})^2 + (-2)^2} = 4$ and $\tan \theta = \frac{-2}{2\sqrt{3}} = -\frac{1}{\sqrt{3}}$

$\Rightarrow \theta = -\frac{\pi}{6} \Rightarrow z = 4[\cos(-\frac{\pi}{6}) + i \sin(-\frac{\pi}{6})]$. For $w = -1 + i$, $r = \sqrt{2}$,

$\tan \theta = \frac{1}{-1} = -1 \Rightarrow \theta = \frac{3\pi}{4} \Rightarrow w = \sqrt{2}(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4})$. Therefore,

$zw = 4\sqrt{2}[\cos(-\frac{\pi}{6} + \frac{3\pi}{4}) + i \sin(-\frac{\pi}{6} + \frac{3\pi}{4})] = 4\sqrt{2}(\cos \frac{7\pi}{12} + i \sin \frac{7\pi}{12})$,

$z/w = \frac{4}{\sqrt{2}}[\cos(-\frac{\pi}{6} - \frac{3\pi}{4}) + i \sin(-\frac{\pi}{6} - \frac{3\pi}{4})] = \frac{4}{\sqrt{2}}[\cos(-\frac{11\pi}{12}) + i \sin(-\frac{11\pi}{12})]$

$= 2\sqrt{2}(\cos \frac{13\pi}{12} + i \sin \frac{13\pi}{12})$, and

$1/z = \frac{1}{4}[\cos(-\frac{\pi}{6}) - i \sin(-\frac{\pi}{6})] = \frac{1}{4}(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$.

32. For $z = 4(\sqrt{3} + i) = 4\sqrt{3} + 4i$, $r = \sqrt{(4\sqrt{3})^2 + 4^2} = \sqrt{64} = 8$ and $\tan \theta = \frac{4}{4\sqrt{3}} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \Rightarrow$

$z = 8(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$. For $w = -3 - 3i$, $r = \sqrt{(-3)^2 + (-3)^2} = \sqrt{18} = 3\sqrt{2}$ and

$\tan \theta = \frac{-3}{-3} = 1 \Rightarrow \theta = \frac{5\pi}{4} \Rightarrow w = 3\sqrt{2}(\cos \frac{5\pi}{4} + i \sin \frac{5\pi}{4})$. Therefore,

$zw = 8 \cdot 3\sqrt{2}[\cos(\frac{\pi}{6} + \frac{5\pi}{4}) + i \sin(\frac{\pi}{6} + \frac{5\pi}{4})] = 24\sqrt{2}(\cos \frac{17\pi}{12} + i \sin \frac{17\pi}{12})$,

$z/w = \frac{8}{3\sqrt{2}}[\cos(\frac{\pi}{6} - \frac{5\pi}{4}) + i \sin(\frac{\pi}{6} - \frac{5\pi}{4})] = \frac{4\sqrt{2}}{3}[\cos(-\frac{13\pi}{12}) + i \sin(-\frac{13\pi}{12})]$, and

$1/z = \frac{1}{8}(\cos \frac{\pi}{6} - i \sin \frac{\pi}{6})$.

33. For $z = 1 + i$, $r = \sqrt{2}$ and $\tan \theta = \frac{1}{1} = 1 \Rightarrow \theta = \frac{\pi}{4} \Rightarrow z = \sqrt{2}(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4})$. So by

De Moivre's Theorem,

$$\begin{aligned} (1+i)^{20} &= \left[\sqrt{2}(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4}) \right]^{20} = (2^{1/2})^{20} (\cos \frac{20 \cdot \pi}{4} + i \sin \frac{20 \cdot \pi}{4}) \\ &= 2^{10}(\cos 5\pi + i \sin 5\pi) = 2^{10}[-1 + i(0)] = -2^{10} = -1024 \end{aligned}$$

34. For $z = 1 - \sqrt{3}i$, $r = \sqrt{1^2 + (-\sqrt{3})^2} = 2$ and $\tan \theta = \frac{-\sqrt{3}}{1} = -\sqrt{3} \Rightarrow \theta = \frac{5\pi}{3} \Rightarrow z = 2 \left(\cos \frac{5\pi}{3} + i \sin \frac{5\pi}{3} \right)$. So by De Moivre's Theorem,

$$\begin{aligned} (1 - \sqrt{3}i)^5 &= [2(\cos \frac{5\pi}{3} + i \sin \frac{5\pi}{3})]^5 = 2^5 (\cos \frac{5 \cdot 5\pi}{3} + i \sin \frac{5 \cdot 5\pi}{3}) \\ &= 2^5 (\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}) = 32 \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i \right) = 16 + 16\sqrt{3}i \end{aligned}$$

35. For $z = 2\sqrt{3} + 2i$, $r = \sqrt{(2\sqrt{3})^2 + 2^2} = \sqrt{16} = 4$ and $\tan \theta = \frac{2}{2\sqrt{3}} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \Rightarrow z = 4(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})$. So by De Moivre's Theorem,

$$(2\sqrt{3} + 2i)^5 = [4(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})]^5 = 4^5 (\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6}) = 1024 \left[-\frac{\sqrt{3}}{2} + \frac{1}{2}i \right] = -512\sqrt{3} + 512i$$

36. For $z = 1 - i$, $r = \sqrt{2}$ and $\tan \theta = \frac{-1}{1} = -1 \Rightarrow \theta = \frac{7\pi}{4} \Rightarrow z = \sqrt{2}(\cos \frac{7\pi}{4} + i \sin \frac{7\pi}{4}) \Rightarrow$

$$\begin{aligned} (1 - i)^8 &= [\sqrt{2}(\cos \frac{7\pi}{4} + i \sin \frac{7\pi}{4})]^8 = 2^4 (\cos \frac{8 \cdot 7\pi}{4} + i \sin \frac{8 \cdot 7\pi}{4}) \\ &= 16(\cos 14\pi + i \sin 14\pi) = 16(1 + 0i) = 16 \end{aligned}$$

37. $1 = 1 + 0i = 1(\cos 0 + i \sin 0)$. Using Equation 3 with $r = 1$, $n = 8$, and $\theta = 0$, we have


$$w_k = 1^{1/8} \left[\cos \left(\frac{0 + 2k\pi}{8} \right) + i \sin \left(\frac{0 + 2k\pi}{8} \right) \right] = \cos \frac{k\pi}{4} + i \sin \frac{k\pi}{4}, \text{ where } k = 0, 1, 2, \dots, 7.$$

$$w_0 = 1(\cos 0 + i \sin 0) = 1, w_1 = 1(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4}) = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i,$$

$$w_2 = 1(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}) = i, w_3 = 1(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4}) = -\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i,$$

$$w_4 = 1(\cos \pi + i \sin \pi) = -1, w_5 = 1(\cos \frac{5\pi}{4} + i \sin \frac{5\pi}{4}) = -\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i,$$

$$w_6 = 1(\cos \frac{3\pi}{2} + i \sin \frac{3\pi}{2}) = -i, w_7 = 1(\cos \frac{7\pi}{4} + i \sin \frac{7\pi}{4}) = \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i$$



38. $32 = 32 + 0i = 32(\cos 0 + i \sin 0)$. Using Equation 3 with $r = 32$, $n = 5$, and $\theta = 0$, we have

$$w_k = 32^{1/5} \left[\cos \left(\frac{0 + 2k\pi}{5} \right) + i \sin \left(\frac{0 + 2k\pi}{5} \right) \right] = 2(\cos \frac{2}{5}\pi k + i \sin \frac{2}{5}\pi k), \text{ where } k = 0, 1, 2, 3, 4.$$


$$w_0 = 2(\cos 0 + i \sin 0) = 2$$

$$w_1 = 2(\cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5})$$

$$w_2 = 2(\cos \frac{4\pi}{5} + i \sin \frac{4\pi}{5})$$

$$w_3 = 2(\cos \frac{6\pi}{5} + i \sin \frac{6\pi}{5})$$

$$w_4 = 2(\cos \frac{8\pi}{5} + i \sin \frac{8\pi}{5})$$




39. $i = 0 + i = 1\left(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}\right)$. Using Equation 3 with $r = 1$, $n = 3$, and $\theta = \frac{\pi}{2}$, we have

$$w_k = 1^{1/3} \left[\cos \left(\frac{\frac{\pi}{2} + 2k\pi}{3} \right) + i \sin \left(\frac{\frac{\pi}{2} + 2k\pi}{3} \right) \right], \text{ where } k = 0, 1, 2.$$

$$w_0 = \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right) = \frac{\sqrt{3}}{2} + \frac{1}{2}i$$

$$w_1 = \left(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6} \right) = -\frac{\sqrt{3}}{2} + \frac{1}{2}i$$

$$w_2 = \left(\cos \frac{9\pi}{6} + i \sin \frac{9\pi}{6} \right) = -i$$




40. $1 + i = \sqrt{2}\left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4}\right)$. Using Equation 3 with $r = \sqrt{2}$, $n = 3$, and $\theta = \frac{\pi}{4}$, we have

$$w_k = (\sqrt{2})^{1/3} \left[\cos \left(\frac{\frac{\pi}{4} + 2k\pi}{3} \right) + i \sin \left(\frac{\frac{\pi}{4} + 2k\pi}{3} \right) \right], \text{ where } k = 0, 1, 2.$$

$$w_0 = 2^{1/6} \left(\cos \frac{\pi}{12} + i \sin \frac{\pi}{12} \right)$$

$$w_1 = 2^{1/6} \left(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4} \right) = 2^{1/6} \left(-\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i \right) = -2^{-1/3} + 2^{-1/3}i$$

$$w_2 = 2^{1/6} \left(\cos \frac{17\pi}{12} + i \sin \frac{17\pi}{12} \right)$$



41. Using Euler's formula (6) with $y = \frac{\pi}{2}$, we have $e^{i\pi/2} = \cos \frac{\pi}{2} + i \sin \frac{\pi}{2} = 0 + 1i = i$.

42. Using Euler's formula (6) with $y = 2\pi$, we have $e^{2\pi i} = \cos 2\pi + i \sin 2\pi = 1$.

43. Using Euler's formula (6) with $y = \frac{\pi}{3}$, we have $e^{i\pi/3} = \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$.

44. Using Euler's formula (6) with $y = -\pi$, we have $e^{-i\pi} = \cos(-\pi) + i \sin(-\pi) = -1$.

45. Using Equation 7 with $x = 2$ and $y = \pi$, we have $e^{2+i\pi} = e^2 e^{i\pi} = e^2(\cos \pi + i \sin \pi) = e^2(-1 + 0) = -e^2$.

46. Using Equation 7 with $x = \pi$ and $y = 1$, we have $e^{\pi+i} = e^\pi \cdot e^{1i} = e^\pi(\cos 1 + i \sin 1) = e^\pi \cos 1 + (e^\pi \sin 1)i$.

47. $F(x) = e^{rx} = e^{(a+bi)x} = e^{ax+bx i} = e^{ax}(\cos bx + i \sin bx) = e^{ax} \cos bx + i(e^{ax} \sin bx) \Rightarrow$

$$\begin{aligned} F'(x) &= (e^{ax} \cos bx)' + i(e^{ax} \sin bx)' = (ae^{ax} \cos bx - be^{ax} \sin bx) + i(ae^{ax} \sin bx + be^{ax} \cos bx) \\ &= a[e^{ax}(\cos bx + i \sin bx)] + b[e^{ax}(-\sin bx + i \cos bx)] = ae^{rx} + b[e^{ax}(i^2 \sin bx + i \cos bx)] \\ &= ae^{rx} + bi[e^{ax}(\cos bx + i \sin bx)] = ae^{rx} + bie^{rx} = (a + bi)e^{rx} = re^{rx} \end{aligned}$$

48. (a) From Exercise 47, $F(x) = e^{(1+i)x} \Rightarrow F'(x) = (1+i)e^{(1+i)x}$. So

$$\int e^{(1+i)x} dx = \frac{1}{1+i} \int F'(x) dx = \frac{1}{1+i} F(x) + C = \frac{1-i}{2} F(x) + C = \frac{1-i}{2} e^{(1+i)x} + C$$

$$(b) \int e^{(1+i)x} dx = \int e^x e^{ix} dx = \int e^x (\cos x + i \sin x) dx = \int e^x \cos x dx + i \int e^x \sin x dx \quad (1).$$

Also,

$$\begin{aligned} \frac{1-i}{2} e^{(1+i)x} &= \frac{1}{2} e^{(1+i)x} - \frac{1}{2} i e^{(1+i)x} = \frac{1}{2} e^{x+ix} - \frac{1}{2} i e^{x+ix} \\ &= \frac{1}{2} e^x (\cos x + i \sin x) - \frac{1}{2} i e^x (\cos x + i \sin x) \\ &= \frac{1}{2} e^x \cos x + \frac{1}{2} e^x \sin x + \frac{1}{2} i e^x \sin x - \frac{1}{2} i e^x \cos x \\ &= \frac{1}{2} e^x (\cos x + \sin x) + i \left[\frac{1}{2} e^x (\sin x - \cos x) \right] \quad (2) \end{aligned}$$

Equating the real and imaginary parts in (1) and (2), we see that $\int e^x \cos x dx = \frac{1}{2} e^x (\cos x + \sin x) + C$ and $\int e^x \sin x dx = \frac{1}{2} e^x (\sin x - \cos x) + C$.